MARROW ED8

Previous Year Question Papers

Comprehensive Question Bank

Contents

Chapter	Title	Page
1	Anatomy AIIMS 2017	16
2	Anatomy AIIMS 2018	45
3	Anatomy AIIMS 2019 (May & Nov)	67
4	Anatomy AIIMS 2020 (May and Nov INI-CET)	95
5	Anatomy NEET 2018	126
6	Anatomy NEET 2019	139
7	Anatomy NEET 2020	156
8	Anatomy INI-CET 2021	175
9	Anatomy NEET 2021	218
10	Anatomy INI-CET 2022	233
11	Anatomy NEET 2022	298
12	Anatomy NEET 2023	311
13	Anatomy INI-CET 2023	323
14	Anatomy INI-CET 2024	366
15	Anatomy NEET 2024	401
16	Physiology AIIMS 2017	417
17	Physiology AIIMS 2018	439
18	Physiology AIIMS 2019 (May & Nov)	459
19	Physiology AIIMS 2020 (May and Nov INI-CET)	471
20	Physiology NEET 2018	493

Chapter	Title	Page
21	Physiology NEET 2019	503
22	Physiology NEET 2020	517
23	Physiology INI-CET 2021	529
24	Physiology NEET 2021	554
25	Physiology INI-CET 2022	559
26	Physiology NEET 2022	592
27	Physiology NEET 2023	602
28	Physiology INI-CET 2023	615
29	Physiology INI-CET 2024	642
30	Physiology NEET 2024	678
31	Biochemistry AIIMS 2017	692
32	Biochemistry AIIMS 2018	707
33	Biochemistry AIIMS 2019 (May & Nov)	723
34	Biochemistry AIIMS 2020 (May and Nov INI-CET)	739
35	Biochemistry NEET 2018	758
36	Biochemistry NEET 2019	770
37	Biochemistry NEET 2020	778
38	Biochemistry INI-CET 2021	791
39	Biochemistry NEET 2021	822
40	Biochemistry INI-CET 2022	845
41	Biochemistry NEET 2022	872
42	Biochemistry NEET 2023	885

Chapter	Title	Page
43	Biochemistry INI-CET 2023	903
44	Biochemistry INI-CET 2024	924
45	Biochemistry NEET 2024	946
46	Pharmacology AIIMS 2017	977
47	Pharmacology AIIMS 2018	999
48	Pharmacology AIIMS 2019 (May & Nov)	1025
49	Pharmacology AIIMS 2020 (May and Nov INI-CET)	1038
50	Pharmacology NEET 2018	1066
51	Pharmacology NEET 2019	1080
52	Pharmacology NEET 2020	1095
53	Pharmacology INI-CET 2021	1110
54	Pharmacology NEET 2021	1145
55	Pharmacology INI-CET 2022	1161
56	Pharmacology NEET 2022	1217
57	Pharmacology NEET 2023	1235
58	Pharmacology INI-CET 2023	1247
59	Pharmacology INI-CET 2024	1287
60	Pharmacology NEET 2024	1310
61	Microbiology AIIMS 2017	1334
62	Microbiology AIIMS 2018	1360
63	Microbiology AIIMS 2019 (May & Nov)	1385
64	Microbiology AIIMS 2020 (May and Nov INI-CET)	1402

Chapter	Title	Page
65	Microbiology NEET 2018	1443
66	Microbiology NEET 2019	1456
67	Microbiology NEET 2020	1479
68	Microbiology INI-CET 2021	1499
69	Microbiology NEET 2021	1548
70	Microbiology INI-CET 2022	1570
71	Microbiology NEET 2022	1629
72	Microbiology NEET 2023	1649
73	Microbiology INI-CET 2023	1671
74	Microbiology INI-CET 2024	1718
75	Microbiology NEET 2024	1768
76	Pathology AIIMS 2017	1807
77	Pathology AIIMS 2018	1844
78	Pathology AIIMS 2019 (May & Nov)	1865
79	Pathology AIIMS 2020 (May and Nov INI-CET)	1896
80	Pathology NEET 2018	1931
81	Pathology NEET 2019	1947
82	Pathology NEET 2020	1966
83	Pathology INI-CET 2021	1996
84	Pathology NEET 2021	2060
85	Pathology INI-CET 2022	2086
86	Pathology NEET 2022	2177

Chapter	Title	Page
87	Pathology NEET 2023	2196
88	Pathology INI-CET 2023	2218
89	Pathology INI-CET 2024	2268
90	Pathology NEET 2024	2303
91	Community Medicine AIIMS 2017	2365
92	Community Medicine AIIMS 2018	2383
93	Community Medicine AIIMS 2019 (May & Nov)	2406
94	Community Medicine AIIMS 2020 (May and Nov INI-...	2428
95	Community Medicine NEET 2018	2442
96	Community Medicine NEET 2019	2456
97	Community Medicine NEET 2020	2472
98	Community Medicine INI-CET 2021	2487
99	Community Medicine NEET 2021	2514
100	Community Medicine INI-CET 2022	2533
101	Community medicine NEET 2022	2565
102	Community Medicine NEET 2023	2578
103	Community Medicine INI-CET 2023	2594
104	Community Medicine INI-CET 2024	2619
105	Community Medicine NEET 2024	2633
106	Forensic Medicine AIIMS 2017	2665
107	Forensic Medicine AIIMS 2018	2678
108	Forensic Medicine AIIMS 2019 (May & Nov)	2693

Chapter	Title	Page
109	Forensic Medicine AIIMS 2020 (May and Nov INI-CET)	2698
110	Forensic Medicine NEET 2018	2717
111	Forensic Medicine NEET 2019	2726
112	Forensic Medicine NEET 2020	2735
113	Forensic Medicine INI-CET 2021	2742
114	Forensic Medicine NEET 2021	2770
115	Forensic Medicine INI-CET 2022	2787
116	Forensic Medicine NEET 2022	2824
117	Forensic Medicine NEET 2023	2835
118	Forensic Medicine INI-CET 2023	2843
119	Forensic Medicine INI-CET 2024	2872
120	Forensic Medicine NEET 2024	2885
121	Ophthalmology AIIMS 2017	2904
122	Ophthalmology AIIMS 2018	2931
123	Ophthalmology AIIMS 2019 (May & Nov)	2968
124	Ophthalmology AIIMS 2020 (May and Nov INI-CET)	2987
125	Ophthalmology NEET 2018	2997
126	Ophthalmology NEET 2019	3015
127	Ophthalmology NEET 2020	3030
128	Ophthalmology INI-CET 2021	3039
129	Ophthalmology NEET 2021	3055
130	Ophthalmology INICET 2022	3072

Chapter	Title	Page
131	Ophthalmology NEET 2022	3096
132	Ophthalmology NEET 2023	3111
133	Ophthalmology INI-CET 2023	3122
134	Ophthalmology INI-CET 2024	3142
135	Ophthalmology NEET 2024	3164
136	ENT AIIMS 2017	3188
137	ENT AIIMS 2018	3202
138	ENT AIIMS 2019 (May & Nov)	3212
139	ENT AIIMS 2020 (May and Nov INI-CET)	3218
140	ENT NEET 2018	3240
141	ENT NEET 2019	3247
142	ENT NEET 2020	3256
143	ENT INICET 2021	3266
144	ENT NEET 2021	3286
145	ENT INICET 2022	3297
146	ENT NEET 2022	3318
147	ENT NEET 2023	3334
148	ENT INICET 2023	3344
149	ENT INI-CET 2024	3365
150	ENT NEET 2024	3384
151	Anesthesia AIIMS 2017	3407
152	Anesthesia AIIMS 2018	3412

Chapter	Title	Page
153	Anesthesia AIIMS 2019 (May & Nov)	3415
154	Anesthesia AIIMS 2020 (May and Nov INI-CET)	3421
155	Anesthesia NEET 2018	3430
156	Anesthesia NEET 2019	3436
157	Anesthesia NEET 2020	3441
158	Anesthesia INI-CET 2021	3445
159	Anesthesia NEET 2021	3464
160	Anesthesia INI-CET 2022	3473
161	Anesthesia NEET 2022	3477
162	Anesthesia NEET 2023	3482
163	Anesthesia INI-CET 2023	3487
164	Anesthesia INI-CET 2024	3492
165	Anesthesia NEET 2024	3505
166	Dermatology AIIMS 2017	3524
167	Dermatology AIIMS 2018	3539
168	Dermatology AIIMS 2019 (May & Nov)	3550
169	Dermatology AIIMS 2020 (May and Nov INI-CET)	3563
170	Dermatology NEET 2018	3577
171	Dermatology NEET 2019	3587
172	Dermatology NEET 2020	3602
173	Dermatology INI-CET 2021	3611
174	Dermatology NEET 2021	3639

Chapter	Title	Page
175	Dermatology INI-CET 2022	3653
176	Dermatology NEET 2022	3678
177	Dermatology NEET 2023	3695
178	Dermatology INI-CET 2023	3701
179	Dermatology INI-CET 2024	3724
180	Dermatology NEET 2024	3734
181	Psychiatry AIIMS 2017	3756
182	Psychiatry AIIMS 2018	3766
183	Psychiatry AIIMS 2019 (May & Nov)	3773
184	Psychiatry AIIMS 2020 (May and Nov INI-CET)	3780
185	Psychiatry NEET 2018	3792
186	Psychiatry NEET 2019	3799
187	Psychiatry NEET 2020	3805
188	Psychiatry INI-CET 2021	3812
189	Psychiatry NEET 2021	3821
190	Psychiatry INI-CET 2022	3827
191	Psychiatry NEET 2022	3839
192	Psychiatry NEET 2023	3842
193	Psychiatry INI-CET 2023	3848
194	Psychiatry INI-CET 2024	3855
195	Psychiatry NEET 2024	3860
196	Radiology AIIMS 2017	3870

Chapter	Title	Page
197	Radiology AIIMS 2018	3880
198	Radiology AIIMS 2019 (May & Nov)	3892
199	Radiology AIIMS 2020 (May and Nov INI-CET)	3897
200	Radiology NEET 2018	3905
201	Radiology NEET 2019	3923
202	Radiology NEET 2020	3935
203	Radiology INI-CET 2021	3945
204	Radiology NEET 2021	3967
205	Radiology INI-CET 2022	3975
206	Radiology NEET 2022	4007
207	Radiology NEET 2023	4012
208	Radiology INI-CET 2023	4020
209	Radiology INI-CET 2024	4034
210	Radiology NEET 2024	4044
211	Medicine AIIMS 2017	4071
212	Medicine AIIMS 2018	4114
213	Medicine AIIMS 2019 (May & Nov)	4150
214	Medicine AIIMS 2020 (May and Nov INI-CET)	4173
215	Medicine NEET 2018	4206
216	Medicine NEET 2019	4234
217	Medicine NEET 2020	4261
218	Medicine INI-CET 2021	4286

Chapter	Title	Page
219	Medicine NEET 2021	4318
220	Medicine INI-CET 2022	4352
221	Medicine NEET 2022	4422
222	Medicine NEET 2023	4457
223	Medicine INI-CET 2023	4488
224	Medicine INI-CET 2024	4532
225	Medicine NEET 2024	4576
226	Surgery AIIMS 2017	4639
227	Surgery AIIMS 2018	4674
228	Surgery AIIMS 2019 (May & Nov)	4696
229	Surgery AIIMS 2020 (May and Nov INI-CET)	4730
230	Surgery NEET 2018	4779
231	Surgery NEET 2019	4810
232	Surgery NEET 2020	4835
233	Surgery INI-CET 2021	4865
234	Surgery NEET 2021	4907
235	Surgery INI-CET 2022	4937
236	Surgery NEET 2022	4994
237	Surgery NEET 2023	5034
238	Surgery INI-CET 2023	5073
239	Surgery INI-CET 2024	5111
240	Surgery NEET 2024	5146

Chapter	Title	Page
241	Orthopedics AIIMS 2017	5215
242	Orthopedics AIIMS 2018	5240
243	Orthopedics AIIMS 2019 (May & Nov)	5256
244	Orthopedics AIIMS 2020 (May and Nov INI-CET)	5277
245	Orthopedics NEET 2018	5297
246	Orthopedics NEET 2019	5309
247	Orthopedics NEET 2020	5329
248	Orthopedics INI-CET 2021	5341
249	Orthopedics NEET 2021	5365
250	Orthopedics INI-CET 2022	5380
251	Orthopedics NEET 2022	5410
252	Orthopedics NEET 2023	5427
253	Orthopedics INI-CET 2023	5441
254	Orthopedics INI-CET 2024	5457
255	Orthopedics NEET 2024	5469
256	Paediatrics AIIMS 2017	5498
257	Paediatrics AIIMS 2018	5515
258	Paediatrics AIIMS 2019 (May & Nov)	5535
259	Paediatrics AIIMS 2020 (May and Nov INI-CET)	5553
260	Paediatrics NEET 2018	5567
261	Paediatrics NEET 2019	5583
262	Paediatrics NEET 2020	5602

Chapter	Title	Page
263	Paediatrics INI-CET 2021	5613
264	Paediatrics NEET 2021	5629
265	Paediatrics INI-CET 2022	5645
266	Paediatrics NEET 2022	5672
267	Paediatrics NEET 2023	5700
268	Paediatrics INI-CET 2023	5714
269	Paediatrics INI-CET 2024	5735
270	Paediatrics NEET 2024	5760
271	Obstetrics and Gynecology AIIMS 2017	5776
272	Obstetrics and Gynecology AIIMS 2018	5814
273	Obstetrics and Gynecology AIIMS 2019 (May & Nov)	5852
274	Obstetrics and Gynecology AIIMS 2020 (May and N...	5875
275	Obstetrics and Gynecology NEET 2018	5900
276	Obstetrics and Gynecology NEET 2019	5927
277	Obstetrics and Gynecology NEET 2020	5952
278	Obstetrics and Gynecology INICET 2021	5983
279	Obstetrics & Gynecology NEET 2021	6047
280	Obstetrics and Gynecology INICET 2022	6073
281	Obstetrics and Gynecology NEET 2022	6133
282	Obstetrics and Gynecology NEET 2023	6166
283	Obstetrics and Gynecology INI-CET 2023	6191
284	Obstetrics and Gynecology INI-CET 2024	6225

Chapter	Title	Page
285	Obstetrics and Gynecology NEET 2024	6262

Anatomy AIIMS 2017

Question 1:

Nucleus pulposus is derived from _____.



a) 1

b) 2

c) 3

d) 4

Question 2:

What is the origin of the fibers marked by the arrow?



a) Hippocampus

b) Habenular

c) Mamillary body

d) Amygdala

Question 3:

Among the following, what is the function of the marked muscles?



a) Depression

b) Protrusion

c) Elevation

d) Retraction

Question 4:

Which of the marked foramen does the mandibular division of the trigeminal nerve pass through?



a) A

b) B

c) C

d) D

Question 5:

The structure marked with the asterisk in the electron micrograph indicates _____.



a) Mitochondria

b) Golgi bodies

c) Secretory vesicles

d) Ribosomes

Question 6:

In the image given below, identify the muscle responsible for the opening of the mouth.



a) A

b) B

c) C

d) D

Question 7:

Identify the structure passing through the foramen marked.



a) Abducens nerve

b) Spinal accessory nerve

c) Ophthalmic artery

d) Middle meningeal artery

Question 8:

The talocalcaneonavicular joint is a _____type of joint.

a) Ball and Socket

b) Saddle type

c) Plane

d) Ellipsoid

Question 9:

Which of the following is connected to the structure marked as A in the image below?



a) Orbital cortex

b) Hippocampus

c) Striate cortex

d) Optic chiasma

Question 10:

The structure passing through the central tendon of the diaphragm is:

a) Oesophagus

b) Aorta

c) Inferior vena cava

d) Sympathetic Chain

Question 11:

Shape of Trapezius muscle is _____

a) Triangular

b) Quadrilateral

c) Pentagonal

d) Linear

Question 12:

All of the following are true about grey rami communicantes except:

a) Unmyelinated

b) Connected to spinal nerve

c) Preganglionic

d) Present medial to white rami communicantes

Question 13:

Which of the following represents a holocrine gland?



a) A

b) B

c) C

d) D

Question 14:

Which of the following structures marked in the below image gives rise to the heart in an adult?



a) A

b) C

c) B

d) D

Question 15:

A 58-year-old woman was undergoing cholecystectomy. During the procedure, the resident accidentally advanced the scissors along the superior border of the epiploic foramen causing injury. Which of the following structures is likely to be injured?

a) Right lobe of liver

b) Quadrate lobe of liver

c) Caudate lobe of liver

d) Fundus of gall bladder

Question 16:

Which of the following structures does the cell pointed in arrow inhibit?



a) Vestibular nuclei

b) Anterior horn of spinal

c) Basal ganglia

d) Deep nuclei of cerebellum

Question 17:

The image below shows the insertion of transversus abdominis muscle. Identify the conjoint tendon:



a) A

b) B

c) C

d) D

Question 18:

What represents the marked area in the given histology of lymph node?



a) Marginal zone

b) Paracortical zone

c) Germinal center

d) Mantle zone

Question 19:

A young patient presented with hypoparathyroidism, absent thymus and tetany. Which of the following marked areas in the picture is defective in this condition?



a) 1

b) 2

c) 3

d) 4

Question 20:

A patient came with complaints of inability to move his 4th and 5th digit, cannot hold a pen and he was not able to hold a paper between fingers. Which of the following sites given below is the probable cause of injury to the nerve in question?



a) A

b) B

c) C

d) D

Question 21:

Which of the following structures of the scapula is palpable in the infraclavicular fossa?



a) A

b) B

c) C

d) D

Question 22:

Identify the pelvic diaphragm in the picture given below.



a) A

b) B

c) C

d) D

Question 23:

What is the root value of cremasteric reflex?

a) L1,L2

b) L3,L4

c) S1,S2

d) T8,T9

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	a
5	c

6	b
7	a
8	a
9	c
10	c
11	a
12	c
13	b
14	b
15	c
16	d
17	c
18	d
19	c
20	c
21	a
22	a
23	a

Detailed Explanations

Solution to Question 1:

Nucleus pulposus is derived from the notochord (marked as 1).

The other labelled structures are

2 - Ectoderm

3 - Endoderm

4 - Intraembryonic mesoderm

Solution to Question 2:

The arrow indicates the fornix. It arises from the hippocampus and ends in the mamillary body of the hypothalamus.

Fornix is an example of association fiber. It is supplied by the anterior cerebral artery.

Solution to Question 3:

The muscle marked in the above image is the lateral pterygoid muscle and its primary function is depression of the mandible (opening of the mouth).

Bilateral contraction of the lateral pterygoid alone causes depression of the mandible (making it the primary function) whereas, protrusion is the result of bilateral contraction of both the medial and lateral pterygoid muscles.

The lateral pterygoid has the following actions:

• Opening of the jaw (depression): Lateral pterygoid has a specialized role in mandibular opening mediated by its horizontal fibres. When left and right lateral pterygoid muscles contract together, the mandibular condyle is pulled forwards and slightly downwards. This protrusive movement alone has little or no function except in assisting the opening of the jaw (Gray's Anatomy 41st edition).

• Side-to-side movement: When the lateral pterygoid contracts unilaterally along with the adjacent medial pterygoid the jaw is pulled medially. This is used for grinding food between teeth of the same side. This action is mediated by the inferior head of the lateral pterygoid.

Muscles of mastication	Nerve supply	Action (On mandible)
Temporalis	Mandibular nerve (CN V3)	ElevationRetraction (posterior fibers)
Masseter	Mandibular nerve (CN V3)	ElevationProtrusion (superficial fibers)
Lateral pterygoid	Mandibular nerve (CN V3)	DepressionSwinging of Jaw(unilaterally)
Medial pterygoid	Mandibular nerve (CN V3)	ElevationProtrusion
Suprahyoid muscles		
Digastric	Facial nerve (CN VII),Mandibular nerve (CN V3)	Depression of mandible and elevation of hyoid bone
Stylohyoid	Facial nerve (CN VII)	Elevation of hyoid bone
Mylohyoid	Mandibular nerve (CN V3)	Elevation of hyoid bone
Geniohyoid	Nerve to geniohyoid	Depression of mandible and elevation of hyoid bone
Infrahyoid muscles		
Omohyoid	Ansa cervicalis (C1-C3)	Fixes or depresses hyoid bone
Sternohyoid	Ansa cervicalis (C1-C3)	Fixes or depresses hyoid bone
Sternothyroid	Ansa cervicalis (C1-C3)	Draws thyroid cartilage downwards
Thyrohyoid	C1 via hypoglossal nerve (CN XII)	Fixes or depresses hyoid bone

Solution to Question 4:

The mandibular nerve passes through the foramen ovale which is marked as A in the above image.

Other foramina marked in the above image:

B - Foramen spinosum

C - Internal acoustic meatus

D - Jugular foramen

Solution to Question 5:

The structure marked with the asterisk in the electron micrograph indicates presynaptic secretory vesicles. The structure marked with the arrowhead is postsynaptic density.

The secretory vesicles are synthesized by the neurons and are involved in electrochemical signalling in the brain and neuromuscular junctions.



On electron microscopy:

• Mitochondria appear as round or elliptical bodies of 0.5-2.0 micron long on an electron microscope, consisting of an outer and an inner mitochondrial membrane, cristae, and mitochondrial matrix.



• Golgi bodies (or Golgi apparatus) appear as stacks of flatenned membranous cisternae surrounded by clusters of vesicles.



• Ribosomes may be attached to the membrane of the rough endoplasmic reticulum (as shown in the image below), or else, they may exist as monosomes or as a chain of granular structures called polyribosomes associated with mRNA.



Solution to Question 6:

In the given image the structure marked B represents the lateral pterygoid muscle which is responsible for the opening of the mouth.

Lateral pterygoid:

• Origin -

• Upper head - Infratemporal surface and infratemporal crest of the greater wing of the sphenoid bone.

• Lower head - Lateral surface of the lateral pterygoid plate.

• Insertion - It is inserted into the pterygoid fovea, capsule of the temporomandibular joint, and the articular disc.

• Nerve supply - Lateral pterygoid nerve which is a branch of the trigeminal nerve.

Other structures marked:

A - Temporalis

C - Masseter

D - Medial pterygoid



Solution to Question 7:

The marked foramen is the superior orbital fissure and the abducens nerve passes through it.

The abducens nerve is the 6th cranial nerve and arises from the brainstem at the pontomedullary junction. It runs through the Dorello's canal which is present at the medial most end of the petrous ridge and then enters the cavernous sinus where it lies lateral to the internal carotid artery. It enters the orbit through the superior orbital fissure within the common tendinous ring and supplies the lateral rectus.

Other options:

Spinal accessory nerve passes through the foramen magnum along with the medulla oblongata and the verterbral arteries

Ophthalmic artery passes through the optic canal along with the optic nerve.

Middle meningeal artery and vein pass through the foramen spinosum along with the meningeal branch of mandibular nerve.

Solution to Question 8:

The talocalcaneonavicular joint is a ball and socket joint.

• The ball part is formed by the rounded head of the talus.

• The socket is formed by 3 structures:

• Posterior surface of the navicular

• Anterior articular surface of the calcaneus

• Upper surface of the plantar calcaneonavicular ligament (spring ligament).



Solution to Question 9:

The structure marked A is the splenium of the corpus callosum and it is connected to the striate cortex.

The body of the corpus callosum arches posteriorly and ends as the thickened posterior portion called the splenium. The fibers from the splenium form the forceps major and arch backward into the occipital lobe (striate cortex).

The hippocampus is not connected directly to the splenium, but situated inferior to the splenium.

The image given below shows the parts of the corpus callosum.



Solution to Question 10:

The inferior vena cava passes through the central tendon of the diaphragm.



Solution to Question 11:

The trapezius is a broad, flat and triangular muscle

Trapezius muscle shape:

• considering left or right side - triangular shape

• considering both sides together - diamond shaped, hence named “trapezius”



Solution to Question 12:

The grey rami communicantes are not preganglionic fibers. They are postganglionic fibers.

The sympathetic trunks on either side of the vertebral columns consist of a series of ganglia. They are connected to the spinal nerves by white and grey rami communicantes. The cell bodies of the preganglionic sympathetic neurons are located in the lateral horn in the spinal grey matter. They leave the cord in the ventral nerve roots to form spinal nerves. These fibers enter the sympathetic trunk in white rami communicantes. They synapse with the postganglionic sympathetic neurons in the sympathetic ganglia. The axons of the postganglionic neurons return to the spinal nerve through the grey rami communicantes.

Fibers	White rami communicantes	Grey rami communicantes
Location	Preganglionic	Postganglionic
Myelination	Myelinated	Unmyelinated
Principal neurotransmitter	Acetylcholine	Noradrenaline



Solution to Question 13:

The histopathological image marked ‘B’ represents the sebaceous gland, which is a type of holocrine gland.

Holocrine secretions are produced in the cytoplasm of the cell and released by the rupture of the plasma membrane, which destroys the cell and results in the secretion of the product into the lumen.

Examples of holocrine glands are

• Sebaceous glands of the skin and

• Meibomian glands of the eyelid

Image A: Sweat glands - Eccrine/merocrine glands; glands remain intact

Image C: Mammary gland - Apocrine gland; apical part sheds to secrete

Image D: Ceruminous gland - Apocrine glands

Solution to Question 14:

The structure marked 'C' in the above image depicts the heart bulge.



Solution to Question 15:

The superior border of the epiploic foramen (foramen of Winslow) is formed by the caudate lobe of the liver. It is likely to be injured in this case.

Epiploic foramen is a vertical slit of about 3 cm behind the free right border of the lesser omentum. It connects the lesser sac to the greater sac.

Clinical note: Pringle maneuver (as shown below) is a technique where the free margin of the lesser omentum is compressed to obtain rapid control of bleeding from the hepatic artery and portal vein in liver trauma and surgery.



Solution to Question 16:

The image shows a histopathological section of the cerebellum. The cells marked with the arrow are Purkinje cells and they inhibit deep nuclei of the cerebellum.

Solution to Question 17:

The structure marked C is the conjoint tendon.

The conjoint tendon is formed from the lower fibres of internal oblique and the lower part of the aponeurosis of transversus abdominis. It is attached to the pubic crest and extends to a variable extent along the pectineal line.

Transversus Abdominis muscle

Origin: internal surface of 7th -12th costal cartilages, thoracolumbar fascia, iliac crest, connective tissue deep to lateral third of inguinal ligament.

Insertion: Linea Alba with aponeurosis of internal oblique, pubic crest, and pecten pubis via the conjoint tendon.

A: Lumbar fascia

B: Transversus Abdominis

D: Inguinal Ligament

Solution to Question 18:

The area marked in the image is the mantle zone of a lymph node.

Lymph node has primary follicle composed of small, quiescent lymphocytes, and a secondary follicle with germinal centre composed mainly of antigen-stimulated B cells.

The mantle zone is produced as surrounding cells are marginalised by the rapidly growing germinal centre. It is populated by mainly quiescent B cells with condensed heterochromatic nuclei and little cytoplasm (hence the deeply basophilic staining of this region in routine preparations), a few helper T cells, Follicular Dendritic cells and macrophages.

In peripheral lymph nodes, the marginal zone are often inapparent because they are mixed with the mantle zone cells. And often not well developed in lymph nodes.

The paracortex lies between the cortical follicles and the medulla and is populated mainly by T cells, which are not organised into follicles



Solution to Question 19:

The given clinical scenario of hypoparathyroidism, absent thymus, and tetany is suggestive of DiGeorge syndrome. It is caused by defective development of the third and fourth pharyngeal pouch. In the image of pharyngeal pouches, "3" points to the thymus (derivative of 3rd pharyngeal pouch). Since in the above question, the thymus is absent, option 3 will be the most appropriate answer.



DiGeorge syndrome is a complex embryopathy that is caused by the hemizygous interstitial deletion of chromosome 22. It is characterized by multiple developmental defects, including conotruncal defects, hypoparathyroidism, and dysmorphic syndrome. It is also known as velocardiofacial syndrome (palate, facial, and cardiac defects).

It has variable presentations including Cleft palate, Abnormal facies, Thymic aplasia leading to T-cell deficiency, Cardiac defects, and Hypocalcemia secondary to parathyroid aplasia (CATCH-22).

Solution to Question 20:

The given clinical situation suggests an ulnar nerve injury. The most likely site of injury would thus be medial epicondyle (marked as C)

Tests for ulnar nerve lesions are as follows:

• Card test positive

• Claw hand deformity

• Froment's sign positive

• Wartenberg sign

• Palmaris brevis sign

Positive card test is due to weakness of palmar interrosei, causing loss of adduction. As a result, the patient is unable to hold the card firmly between his or her fingers.

Option A: It is the radial groove where radial nerve runs along with profunda brachii artery.

Option D: It is the neck of humerus where axillary nerve runs around.

Option B: It is the lateral epicondyle. Fracture of this part may lead to tardy ulnar nerve palsy, but incidence is less.

Solution to Question 21:

The part of the scapula marked as "A" in this image is palpable and it is the coracoid process.

The apex of the coracoid process lies approximately 2.5cm below the clavicle, immediately lateral to the infraclavicular fossa (deltopectoral triangle). Hence, the coracoid process can be palpated below the clavicle.



Solution to Question 22:

The structure marked "A" is the levator ani muscle which is part of the pelvic diaphragm.

Other structures marked in the above image are

B: External anal sphincter

C: Internal anal sphincter

D: Pudendal canal

Levator ani and ischiococcygeus form the pelvic diaphragm and delineate the lower limit of the true pelvis.



Levator ani is formed by:

• Pubococcygeus

• Iliococcygeus

• Puborectalis.



Solution to Question 23:

Cremasteric reflex is mediated by L1 and L2. In this reflex, there is ipsilateral elevation of the testicle following stroking of the medial thigh.

The cremasteric reflex's afferent and efferent branches:

• Afferent - Femoral branch of genitofemoral nerve

• Efferent - Genital branch of genitofemoral nerve

Given below are the various reflexes with their root values.

Deep reflexes	Superficial reflexes
Biceps -C5, C6	Corneal - CN V(A) - CN VII(E) - Pons
Brachioradialis - C5, C6	Conjunctival - CN V (A) - CN VII(E) - Pons
Triceps - C6, C7	Scapular - C5, T1
Finger flexion - C8	Abdominal - T7-T12
Knee jerk - L3,L4	Anal - S3, S4
Ankle -S1, S2	Cremastric -L1, L2
Jaw jerk - CN V (Pons)	Plantar - L5S1

Anatomy AIIMS 2018

Question 1:

The structure marked with the asterisk in the electron micrograph indicates _____.



a) Mitochondria

b) Golgi bodies

c) Secretory vesicles

d) Ribosomes

Question 2:

Inferior thyroid artery is the branch of which of the following?

a) External carotid artery

b) Internal carotid artery

c) Thyrocervical trunk

d) Lingual artery

Question 3:

Neurovascular bundle is absent in which compartment of the leg?

a) Lateral compartment

b) Posterior superficial compartment

c) Posterior deep compartment

d) Anterior compartment

Question 4:

A patient presents to you with inability to form words but can make noises. He also knows what he wants to say. What area of the brain has been damaged in this patient?

a) Wernicke's area

b) Broca's area

c) Heschl's gyrus

d) Striate area

Question 5:

Identify the wrongly marked structure in the diagram



a) A-Corpus Callosum

b) C-Hypothalamus

c) B-Fornix

d) D-Cerebellum

Question 6:

Loss of flexion at the interphalangeal joint of thumb after supracondylar fracture is due to the involvement of?

a) Deep branch of ulnar nerve

b) Median Nerve

c) Posterior interosseous nerve

d) Anterior interosseous nerve

Question 7:

Purkinje cells of cerebellum relay into _____

a) Caudate nucleus

b) Lentiform nucleus

c) Dentate nucleus

d) Red nucleus

Question 8:

Which of the following sensations is not perceived by the spinal nucleus of the trigeminal nerve?

a) Temperature

b) Pain

c) Touch

d) Proprioception

Question 9:

The structure marked in the given picture arises from the _____.



a) Mamillary body

b) Hippocampus

c) Thalamus

d) Amygdala

Question 10:

Which of the following arteries does not supply the corresponding region of the dura mater?

a) Middle meningeal artery – Anterior cranial fossa

b) Accessory meningeal branch of maxillary artery – Middle cranial fossa

c) Meningeal branch of internal carotid artery – Posterior cranial fossa

d) Meningeal branch of anterior and posterior ethmoid artery – Anterior cranial fossa

Question 11:

All of the following take part in the formation of the wrist joint, except:

a) Scaphoid

b) Triquetrum

c) Distal end of radius

d) Distal end of ulna

Question 12:

Which of the following nerves is not involved in olfaction?

a) Trigeminal nerve

b) Vagus nerve

c) Glossopharyngeal nerve

d) Hypoglossal nerve

Question 13:

Which of the following is the most conspicuous sign of breast cancer?

a) Peau d'orange

b) Nipple retraction

c) Puckering

d) Both nipple retraction and puckering

Question 14:

Identify the cranial nerve that passes through the marked foramen marked in the image below.



a) Oculomotor nerve

b) Olfactory nerve

c) Optic nerve

d) Facial nerve

Question 15:

Which of the following veins drain into brachiocephalic vein?

a) Hemiazygous vein

b) Superior thyroid veins

c) Right superior intercostal vein

d) Left superior intercostal vein

Question 16:

What is the main action of the muscle marked below?



a) Elevation

b) Depression

c) Protrusion

d) Retraction

Question 17:

Ligamentum flavum has high levels of:

a) Type 1 collagen

b) Type 2 collagen

c) Reticular fibres

d) Elastin fibres

Question 18:

Which muscle is not involved in the inversion of foot?

a) Tibialis anterior

b) Tibialis posterior

c) Peroneus longus

d) Extensor hallucis longus

Question 19:

Which of the following is normally seen in lumbar spine?

a) Scoliosis

b) Kyphosis

c) Lordosis

d) Recurvatum

Question 20:

Type I collagen is not seen in _____.

a) Bone

b) Ligament

c) Aponeurosis

d) Elastic cartilage

Question 21:

The correct pair of lymph node level in the neck and anatomical boundaries amongst the following are:

a) Level 3 - Cricoid to clavicle

b) Level 4 - Hyoid to cricoid

c) Level 5 - Base of skull to cricoid

d) Level 6 - Hyoid to clavicle

Question 22:

Meissner corpuscles are present in:

a) Papillary dermis

b) Reticular dermis

c) Basal epidermis

d) Stratum lucidum

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	b
5	b
6	d
7	c
8	d
9	b
10	c
11	d
12	d
13	a
14	a
15	d
16	a
17	d
18	c

19	c
20	d
21	d
22	a

Detailed Explanations

Solution to Question 1:

The structure marked with the asterisk in the electron micrograph indicates presynaptic secretory vesicles. The structure marked with the arrowhead is postsynaptic density.

The secretory vesicles are synthesized by the neurons and are involved in electrochemical signalling in the brain and neuromuscular junctions.



On electron microscopy:

• Mitochondria appear as round or elliptical bodies of 0.5-2.0 micron long on an electron microscope, consisting of an outer and an inner mitochondrial membrane, cristae, and mitochondrial matrix.



• Golgi bodies (or Golgi apparatus) appear as stacks of flatenned membranous cisternae surrounded by clusters of vesicles.



• Ribosomes may be attached to the membrane of the rough endoplasmic reticulum (as shown in the image below), or else, they may exist as monosomes or as a chain of granular structures called polyribosomes associated with mRNA.



Solution to Question 2:

Inferior thyroid artery is a branch of thyrocervical trunk, which in turn, is a branch of the first part of the subclavian artery. The inferior thyroid artery loops upwards anterior to the medial border of the scalenus anterior, turns medially just below the sixth cervical transverse process, and then descends on longus colli to the lower border of the thyroid gland.

Relations between the terminal branches and recurrent laryngeal nerve are very variable and of considerable surgical importance.



Solution to Question 3:

Posterior superficial compartment of the leg does not contain any neurovascular bundle.

The posterior compartment of the leg is divided into the deep and superficial parts by the transverse intermuscular septum.

The contents of the posterior superficial compartment are gastrocnemius, plantaris, and soleus muscle.

Neurovascular components in various compartments of leg are as follows:

• Anterior compartment: Anterior tibial artery and Deep fibular/peroneal nerve

• Posterior deep compartment: Tibial nerve and posterior tibial vessels in the deep plane.

• Lateral Compartment: Superficial peroneal/fibular nerve and peroneal/fibular vessel. Peroneal artery is a branch of posterior tibial artery.

Note: MRI is the best imaging modality for evaluating the soft tissues of the leg.

Solution to Question 4:

The given clinical scenario is suggestive of Broca's aphasia in which the Broca's area will be affected.

In Broca's aphasia/ Motor aphasia / Non-fluent aphasia, areas 44 and 45 in the prefrontal and premotor facial regions of the cerebral cortex are affected. The affected individuals will have difficulty forming verbal or written words. The comprehension aspect of speech is preserved but the person cannot repeat or name the words and lacks fluency.

Patients in whom the area is severely affected may have very few words to express all of their emotions.

Solution to Question 5:

The image shows the midsagittal section of the brain. The wrongly marked structure is C-hypothalamus. The structure marked as C corresponds to the pituitary gland.



Solution to Question 6:

Loss of flexion at the interphalangeal (IP) joint of the thumb, is due to the involvement of the anterior interosseous nerve (AIN).

Flexion of thumb occurs at 2 joints:

• IP joint: Flexor pollicis longus (FPL) supplied by the anterior interosseous nerve (a branch of the median nerve, C7 and C8).

• MCP joint: Flexor pollicis brevis (FPB) supplied by the recurrent motor branch of the median nerve.

Other options:

• Deep branch of ulnar nerve > ulnar claw hand

Solution to Question 7:

Purkinje cells of the cerebellum relay into the dentate nucleus which is one of the deep cerebellar nuclei.

The input to the Purkinje cells come from the climbing fibres. The output of Purkinje cells is inhibitory to the deep cerebellar nuclei.

Options A and B: Caudate and lentiform nuclei are part of the basal ganglia.

Option D: Red nucleus receives afferents from the cerebellum and plays a role in modulating signals to the upper motor neurons of the cerebral cortex.

Solution to Question 8:

Proprioception is perceived by the mesencephalic nucleus of the trigeminal nerve, not the spinal nucleus.

The trigeminal nerve consists of 4 nuclei: 3 sensory (Mesencephalic, spinal, and principal sensory nucleus) and 1 motor.

Nucleus	Location	Function
Mesencephalic Nucleus	Composed of a column ofuni polar nervecells situated in th e midbrain and extends inferi orly into the pons as far as th e main sensory nucleus	Receives the sensation ofpro prioceptionfrom the muscles of mastication and from the f acial and extraocular muscles
Principal sensory nucleu s	Posterior part of the pons lat eral to the motor nucleus	Receives the sensations oftou ch and pressurefrom the face
Spinal nucleus	Continuous superiorly with t he main sensory nucleus and extends inferiorly the whole of the medulla oblongatatill t he 2ndCervical Segment	Receives the sensation ofpain and temperaturefrom the fac e and some fibers oftouch
Motor nucleus	The pons medial to the main sensory nucleus	Supplies the muscles of masti cation, the tensor tympani, t he tensor veli palitini and the mylohyoid and the anterior belly of digastric muscle



Solution to Question 9:

This is a coronal section of the brain and the marked structure is the fornix. It arises from the hippocampus and ends in the mammillary body of the hypothalamus.



Solution to Question 10:

The meningeal branch of the internal carotid artery (ICA) does not supply the posterior cranial fossa. It supplies the anterior cranial fossa.

The dura mater is supplied by meningeal arteries that pass along the inner periosteum (endosteum) of the skull over the endosteal layer of the dura and supply the following regions.

Region	Arteries
Anterior cranial fossa	Meningeal branches of anterior and posterior ethmoid arteriesMeningeal branchesofinternal carotid arteriesA branch of themiddle meningealartery
Middle cranial fossa	Middle meningeal artery and accessory meningeal branches of the maxillary arteryA branch of the ascending pharyngeal arteryBranches of the internal carotid arteryA recurrent branch of the lacrimal artery
Posterior cranial fossa	Meningeal branches of the occipital arteryPosterior meningeal branches of the vertebral arteryDorsal meningeal and tentorial arteries arising from the meningohypophyseal trunk

Solution to Question 11:

The distal end of the ulna does not take part in the formation of the radiocarpal joint.

The radiocarpal joint (Wrist joint) is a synovial, biaxial, and ellipsoid joint formed by the articulation of:

• The distal end of the radius

• Articular disc of the triangular fibrocartilage

• Scaphoid, lunate and triquetrum

In the neutral position of the wrist, only the scaphoid and lunate are in contact with the radius and articular disc; the triquetrum comes into apposition with the disc only in full adduction of the wrist joint.



Solution to Question 12:

The hypoglossal nerve is purely motor cranial nerve and is not involved in olfaction.

The other three cranial nerves—trigeminal, glossopharyngeal, and vagus nerves are mixed cranial nerves and contribute to olfaction.

Vagal nerve stimulation modifies the electrical activity of the olfactory bulb.

Solution to Question 13:

The most conspicuous sign of breast cancer is Peau d' orange appearance.

Peau d'orange (orange peel like texture), is the edematous swollen and pitted breast skin due to obstruction of the subcutaneous lymphatics.

Solution to Question 14:

The marked structure is the superior orbital fissure and the oculomotor nerve passes through this fissure.

Superior orbital fissure is divided into three compartments by the common tendinous ring. The oculomotor nerve passes through its intermediate compartment.

Solution to Question 15:

The left superior intercostal vein drains into the left brachiocephalic vein.

Tributaries of the brachiocephalic vein:

• Left and right internal thoracic veins

• Left and right inferior thyroid veins

• Left superior intercostal vein drains into the left brachiocephalic vein.

The right superior intercostal vein drains into the azygos vein.

Option A - Hemiazygous vein drains into the azygous vein.

Option B - Superior thyroid veins drain directly into internal jugular veins.

Option C - Right superior intercostal vein drains into the azygous vein.



Solution to Question 16:

The muscle marked in the image is the temporalis and its main action is elevation of the mandible.

Other actions are:

• It helps in the side-to-side grinding movements of the mandible.

• The posterior fibres which are horizontal, help in the retraction of the mandible.

Solution to Question 17:

Ligamentum flavum has high levels of elastin fibres.

The ligamenta flava are a series of ligaments that connect the ventral parts of the laminae of adjacent vertebrae. They predominantly contain elastic tissue and form part of the posterior surface of the vertebral canal. The high content of elastin allows it to be stretched by 80% without damage. The ligamentum flava resist separation of the laminae in flexion and assist in extension back to the anatomical position.



Solution to Question 18:

Peroneus longus muscle is not involved in the inversion of the foot. It causes eversion and plantar flexion.

Inversion of the foot is a movement in which the medial border of the foot is elevated so that the sole faces medially.

• Tibialis anterior - Causes dorsiflexion and inversion of the foot

• Tibialis posterior - Causes plantar flexion and inversion of the foot

• Extensor hallucis longus - Causes extension of the great toe, dorsiflexion of the foot, and inversion of the foot (minimal action)

Eversion of the foot is a movement where the lateral border of the foot is elevated so that the sole faces laterally. Peroneus longus, peroneus brevis, and peroneus tertius are the muscles involved.

These movements take place, mainly at the talocalcaneal and the talocalcaneonavicular joints, and partly at the transverse tarsal joint.



Solution to Question 19:

Lordosis is normally seen in lumbar spine.

In adults,

• The cervical curve is a lordosis (convex forwards),

• The thoracic curve is a kyphosis (convex dorsally),

• The lumbar curve is also a lordosis,

• The pelvic curve is concave anteroinferiorly.



Solution to Question 20:

Cartilage is made up of type II collagen.

Type I collagen (fibrocartilage), Type II collagen (hyaline and elastic).

There are nearly 28 types of collagen. The important ones are given below with mnemonic:

Bone, Ligament, Tendon, Aponeurosis	Type I Collagen [B-ONE]
Hyaline and Elastic Cartilage	Type II Collagen [Car-Two-Lage]
Reticular Tissue	Type III Collagen [Re-Three-Cular]
Basement Membrane	Type IV Collagen [Floor-Four]

Solution to Question 21:

The correct pair of lymph nodes and levels among the following options are level 6- Hyoid to clavicle.



Levels of lymph nodes

Level	Lymph node
1a	Submental
1b	Submandibular
2	Upper jugular
3	Middle jugular
4	Lower jugular
5	Posterior triangle
6	Anterior group
7	Mediastinal

Solution to Question 22:

Meissner corpuscles are present in the papillary dermis. These corpuscles are rapidly adapting mechanoreceptors particularly sensitive to the movement of objects over the skin's surface and low-frequency vibrations. Their acute sensitivity plays a crucial role in tasks such as reading Braille text.

It is an elongated encapsulated nerve ending of a large myelinated sensory nerve fiber (type Aβ). Each corpuscle consists of a connective tissue capsule and a central core composed of stacked modified Schwann cells. Meissner's corpuscles are found in nonhairy areas of the skin, especially in fingertips, lips, and other regions where spatial touch perception is highly developed.



Anatomy AIIMS 2019 (May & Nov)

Question 1:

The region marked corresponds to the surface marking for which of the following structures:



a) Thoracic duct

b) Right atrium

c) Inferomedial bronchopulmonary segments

d) Hilum

Question 2:

The action of the muscle attached to the marked area?

a) Flexion

b) Adduction

c) Abduction

d) Extension

Question 3:

Which of the following finger has two dorsal interossei muscles?

a) Index finger

b) Little finger

c) Middle finger

d) Ring finger

Question 4:

The structure that divides the liver in Couinaud's classification system is:

a) Portal vein

b) Hepatic artery

c) Bile duct

d) Cystic duct

Question 5:

Which of the following muscle is not supplied by the structure marked A?



a) Superior rectus

b) Superior oblique

c) Medial rectus

d) Levator palpebrae superioris (LPS)

Question 6:

The wrong statement regarding the levator ani muscle is:

a) Supports pelvic viscera

b) Converge downwards and medially

c) Consists of iliococcygeus and pubococcygeus

d) Attached to pelvic brim

Question 7:

In the given section of the heart, match the marked (A, B, C & D) structures.



a) A-5, B-2, C-1, D-4

b) A-1, B-2, C-5, D-3

c) A-5, B-3, C-2, D-1

d) A-1, B-2, C-4, D-5

Question 8:

A lesion of the structure marked in the image below causes?



a) Contralateral hemiplegia

b) Ataxia

c) Tinnitus

d) Ipsilateral facial palsy

Question 9:

Identify the structure marked in the image given below:



a) Ulnar nerve

b) Ulnar artery

c) Superficial radial artery

d) Median nerve

Question 10:

True regarding the lesion of the structure marked by the arrow is:



a) Ape thumb deformity

b) Claw hand

c) Claudication

d) Wrist drop

Question 11:

Which of the following structures given below is related to the structure highlighted (red) in the image?



a) Pudendal nerve

b) Sciatic nerve

c) Obturator nerve

d) Femoral nerve

Question 12:

All of the following form the marked structure except:



a) 2nd Dorsal interossei

b) Lumbricals

c) Extensor digitorum

d) 2nd Palmar interossei

Question 13:

Which of the following structures of brachial plexus do not give branches?

a) Roots

b) Cords

c) Trunks

d) Divisions

Question 14:

Lesion of the structure marked 'A' causes:



a) Loss of sensation of the nail of index finger

b) Hypothenar muscles wasting

c) Loss of sensation at the tip of radial tubercle

d) Loss of sensation of the 4th interdigital cleft

Question 15:

Which of the following is related to the marked area in the image?



a) Stomach

b) Spleen

c) 2nd part of duodenum

d) Pancreas

Question 16:

The blood supply to the structure marked in the image is by:



a) Vertebral artery

b) Internal carotid artery

c) Anterior cerebral artery

d) Middle cerebral artery

Question 17:

Marked structure is derived from the_____



a) Endocardial cushions

b) Septum primum

c) Left venous valve

d) Bulbous venosus

Question 18:

All of the following supply the marked structure except :

a) Right coronary artery

b) Anterior interventricular artery

c) Marginal artery

d) Diagonal artery

Question 19:

What is the nerve supply of the structure marked in the image below?



a) Oculomotor nerve

b) Trochlear nerve

c) Trigeminal nerve

d) Abducens nerve

Answer Key

Question No.	Correct Option
1	b
2	c
3	c
4	a
5	b
6	d
7	c

8	a
9	b
10	d
11	a
12	d
13	d
14	a
15	a
16	c
17	a
18	d
19	b

Detailed Explanations

Solution to Question 1:

The region marked corresponds to the surface marking for the right atrium.

The costal margin, xiphoid process, and the mid-clavicular line are seen in this image. The examiner draws a laterally convex line that extends from a level below the 2nd costal cartilage (as clued in by the sternal angle) i.e., 3rd costal cartilage to about 3 spaces below i.e., the 6th costal cartilage. This line is about 1.2 cms lateral to the sternal edge. These suggest that the region marked corresponds to the surface marking for the right atrium.

The projection of the cardiac borders, of an average adult, onto the anterior thoracic wall forms a trapezoid. This can be mapped out using the percussion method by looking for cardiac dullness.

The varying heart borders and the corresponding surface markings are as follows:

• Upper border: Slopes gently from the second left costal cartilage to the third right costal cartilage. It corresponds to the left atrium.

• Right border: Convex to the right and runs from the third to the sixth right costal cartilages, 1–2 cm lateral to the sternal edge. It corresponds to the right atrium.

• Inferior border or acute border: Runs leftwards from the 6th right costal cartilage to the cardiac apex which is located approximately 9 cm lateral to the midline, often in the left 5th intercostal space or level with the 5th or 6th rib. It corresponds to the right ventricle.

• Left border or obtuse border: Convex laterally on the left. It extends superomedially from the cardiac apex to meet the second left costal cartilage about 1 cm from the left sternal edge. It corresponds to the left ventricle.





Solution to Question 2:

The image shows the greater trochanter and the muscle fibers of the gluteus medius and the gluteus minimus attach(insert) here and their function is to abduct the thigh and their anterior fibers rotate the thigh medially.

Both are innervated by the superior gluteal nerve (L4, L5, S1)

Greater and Lesser trochanters:

• Greater trochanter

• Large and quadrangular

• Junction of the neck and shaft

• Lesser trochanter

• Conical projection

• Posteromedial to shaft

• Posteroinferior to junction with the neck.

• Surface marking: Greater trochanter lies approximately a hand's breadth below the iliac tuberculum, the surface is palpable, especially when the muscles are relaxed, Lesser trochanter is not palpable.



Solution to Question 3:

The Middle Finger has two Dorsal interossei as seen in the image below.

Dorsal Interossei causes Abduction of fingers from the Middle/3rd finger:

Muscle	Palmar Interossei	Dorsal Interossei
Origin	First arises from the base of 1 st metacarpal; Other three fr om the anterior surface of sh afts of 2nd, 4th, and 5th met acarpals	Contiguous sides of shafts of metacarpal bones
Insertion	Proximal phalanges of thumb and index, ring, and little fin gers and dorsal extensor exp ansion of each finger	Proximal phalanges of index, middle, and ring fingers and dorsal extensor expansion
Nerve sup ply	Deep branch of the ulnar ner ve	
Nerve root s	C8; T1	

Muscle	Palmar Interossei	Dorsal Interossei
Action	Palmar interosseiadductfingerstoward the centerof the third finger & flexes the metacarpophalangeal joints and extend interphalangeal joints	Dorsal interosseiabductfingersfrom the centerof the third finger flexes metacarpophalangeal joints and extend interphalangeal joints



Solution to Question 4:

The structure that divides the liver in Couinaud's classification system is the portal vein.

Solution to Question 5:

The structure marked in the image is the oculomotor nerve and it does not supply the superior oblique muscle. Superior oblique muscle is supplied by the trochlear nerve.

All the extraocular muscles are innervated by the oculomotor (III) nerve except:

- Lateral rectus which is supplied by abducens (VI) nerve
- Superior oblique which is supplied by trochlear (IV) nerve

Optic and Oculomotor nerves



Origin of cranial nerves at the base of the brain



Solution to Question 6:

Levator Ani Muscle is not attached to the pelvic brim. It takes its origin from the body of the pubis, the arcus tendinous of the levator ani (a thickened part of the obturator fascia), and the ischial spine.

• Insertion :- The coccyx and the anococcygeal raphe or ligament.

• Nerve supply :- The branches of the anterior rami of sacral nerves S3 and S4 and the perineal branch of the pudendal nerve.

• Supports and raises the pelvic floor.

• Consists of the puborectalis, pubococcygeus, and iliococcygeus.



Pelvic floor

©MARROW

Solution to Question 7:

The image shows a cut and reflected section of the heart.

The ventricles can be identified based upon the wall thickness. Left ventricle has thicker walls. The correct match is:



©MARROW

Solution to Question 8:

The structure marked in the image is the pyramid containing the pyramidal tracts. A lesion here leads to contralateral hemiplegia.





Solution to Question 9:

The structure marked in the image is the ulnar artery, seen contributing to the superficial palmar arch.

It passes deep to the Guyon's canal in the wrist and contributes to the superficial and deep palmar arch.





Solution to Question 10:

The structure marked in the image is the radial nerve demonstrated in the cubital fossa. A lesion of the radial nerve leads to a wrist drop.





Ape thumb deformity is seen in a lesion of the median nerve.

Claw hand deformity is seen in a lesion of the ulnar nerve.

Solution to Question 11:

The structure marked in red is the ischial spine and the pudendal nerve is related to it.

The pudendal nerve originates from the sacral plexus S2-S4. It exits out of the pelvis by passing through the greater sciatic foramen. It then enters the perineum below the pelvic floor by passing around the ischial spine. The ischial spine is used as a landmark for pudendal nerve block.



Solution to Question 12:

The structure marked in the image is the dorsal digital expansion of the middle finger. Palmar interossei do not attach to the middle finger and therefore it does not contribute to the dorsal digital expansion of the middle finger.

Palmar interossei are unipennate muscles originating from their own metacarpal bone. Their function is adduction. The middle finger does not need adduction and hence does not have palmar interossei.



Dorsal interossei are bipennate muscles originating from the adjacent metacarpal bone. They attach to the middle three fingers on both sides and do not attach on the thumb and little finger. Their function is abduction. The thumb and little finger have other muscles for abduction.

Interossei muscles of left hand (dorsal view)



The lateral 2 lumbricals are unipennate and medial 2 lumbricals are bipennate. Their function is flexion at metacarpophalangeal joints.

Superficial muscles of the hand (palmar view)



Solution to Question 13:

The divisions in the brachial plexus do not give any branches.

However, the roots, trunks, and cords of the brachial plexus give out direct branches.

Solution to Question 14:

The structure marked 'A' in the diagram is median nerve in the cubital fossa. A lesion of the median nerve causes loss of sensation of the nail of the index finger.



Median nerve cutaneous supply:

• Lateral 2/3rd of the palm by superficial cutaneous branch

• Lateral 3½ fingers (palmar aspect and dorsal aspect of nailbeds) by palmar digital nerves



Contents of cubital fossa: (From medial to lateral) - “MBBR”

• Median nerve

• Brachial artery

• Biceps brachii tendon

• Radial nerve

Solution to Question 15:

The marked area in the image of the left kidney is related to the stomach.

The anterior relations of the right kidney:

The anterior relations of the left kidney:

The posterior relations of the kidneys:

• One vessel - subcostal vessels

• Two ligaments - medial and lateral arcuate ligaments

• Three nerves - subcostal, iliohypogastric, and ilioinguinal nerves

• Four muscles - diaphragm, psoas major, quadratus lumborum, and transverse abdominis.

• The right kidney is related to the 12th rib and the left kidney is related to the 11th and 12th ribs

Intraperitoneal	Right lobe of liverSmall intestine
Retroperitoneal	Right suprarenal gland2ndpart of duoednumHepatic flexure of colon

Intraperitoneal	StomachJejunumSpleen
Retroperitoneal	Left suprarenal glandPancreasSplenic vesselsSplenic flexure and beginning of descending colon





Solution to Question 16:

The marked structure is fornix which is supplied by the anterior cerebral artery.

The anteromedial central branches from the anterior cerebral artery supply the fornix along with optic chiasma, lamina terminalis, hypothalamus, para-olfactory areas, and cingulate gyrus.



Solution to Question 17:

The structure marked is the membranous part of the interventricular septum. It is derived from endocardial cushions.

Endocardial cushions are soft tissue masses that protrude in the atrioventricular and conotruncal regions.

• Atrioventricular cushions are derived from overlying endocardial cells.

• The conotruncal cushions are derived from the neural crest cells.

Endocardial cushions contribute to the formation of:

• Atrial septa

• Ventricular septa (membranous portion)

• Atrioventricular canals

• Atrioventricular valves

• Aortic and pulmonary channels

Defects in endocardial cushion formation can result in cardiac malformations, including ASDs, VSDs, and defects involving the great vessels (transposition of the great vessels, common truncus arteriosus, and tetralogy of Fallot).

Solution to Question 18:

The structure marked is the right ventricle. It is not supplied by the diagonal artery.

The diagonal artery is a branch of the anterior interventricular artery supplying the left ventricle. This is present in about 30-50% of the population.

The right ventricle is mostly supplied by the right coronary artery and its branches such as the marginal artery.

The anterior interventricular artery gives off 1-2 anterior right ventricular branches.



Solution to Question 19:

The structure marked in the image is the superior oblique muscle. It is innervated by the trochlear nerve.



There are seven extraocular muscles present in the eye. They are:

• Levator palpebrae superioris - Which elevates the upper eyelids

• Four recti muscles - Superior, inferior, medial, and lateral recti

• Two obliques - Superior and inferior oblique

The muscles originate from the back of the orbit in a common tendinous ring called the annulus of Zinn and are inserted into the sclera.

All the extraocular muscles are innervated by the oculomotor (III) nerve except:

• Lateral rectus which is supplied by abducens (VI) nerve

• Superior oblique which is supplied by trochlear (IV) nerve

Origin of extraocular muscles in orbit



Anatomy AIIMS 2020 (May and Nov INI-CET)

Question 1:

Given is a model of embryological development of the heart. Which structure develops from the area marked as "A" ?



a) Atria

b) Left Ventricle

c) Infundibulum

d) Interventricular septum

Question 2:

A muscle is attached to the area marked in the given image. What is the action of this muscle on the hip joint?

a) Flexion and medial rotation

b) Adduction and lateral rotation

c) Abduction and medial rotation

d) Extension and lateral rotation

Question 3:

Neurotransmitter secreted by the area marked in the given image is?



a) Acetylcholine

b) Dopamine

c) Glutamine

d) Noradrenaline

Question 4:

Which of the following arteries passes in close proximity to the area marked by the arrow in the given image?



a) Internal carotid artery

b) Middle meningeal artery

c) Anterior cerebral artery

d) Deep temporal artery

Question 5:

Where are the Ito cells in the liver located?

a) Space of disse

b) Space of moll

c) Sinusoids

d) Bile canaliculi

Question 6:

Which of the following features differentiates the exocrine pancreas from the parotid gland?

a) Basal lamina lining the serous acini

b) Absence of striated duct

c) Presence of serous acini with acidophilic tip

d) Presence of apical acinar villi

Question 7:

Medial lemniscus is a continuation of which of the following structures?

a) Spinothalamic tract

b) Spinocerebellar tract

c) Spinotectal tract

d) Fasciculus gracilis

Question 8:

The type of fibers present in the fornix are:

a) Association and commisural fibers

b) Association and projection fibers

c) Commissural and projection fibers

d) Association fibers

Question 9:

Decussation of the superior cerebellar peduncle occurs at which level?

a) Pons

b) Midbrain

c) Diencephalon

d) Medulla

Question 10:

The pretracheal layer of deep cervical fascia encloses all of the following except:

a) Oesophagus

b) Sternocleidomastoid

c) Sternohyoid

d) Thyroid gland

Question 11:

All of the following are the contents of the carotid sheath except?

a) Sympathetic chain

b) Carotid artery

c) Internal jugular vein

d) 10th cranial nerve

Question 12:

A patient presented with paradoxical respiration. Which of the following nerves is most likely to be affected?



a) A

b) B

c) C

d) D

Question 13:

Which of the following muscles is attached to the area marked in the image below?



a) Abductor pollicis longus

b) Palmar interossei

c) Opponens pollicis

d) Flexor pollicis brevis

Question 14:

Which of the following marked structures does not form part of the laryngeal skeleton?



a) A

b) B

c) C

d) D

Question 15:

Identify the foramen marked in the image below.



a) Foramen rotundum

b) Foramen ovale

c) Vidian canal

d) Foramen spinosum

Question 16:

Reticular fibres are not present in which of the following structures?

a) Thymus

b) Lymph node

c) Bone marrow

d) Spleen

Question 17:

Which of the following muscles is not involved in the overhead abduction of the arm?

a) Serratus anterior

b) Deltoid

c) Trapezius

d) Pectoralis major

Question 18:

The structures marked in the image below are found in the epithelial lining of all of the following sites, except:



a) Oviduct

b) Epididymis

c) Bronchi

d) Ependyma

Question 19:

Which of the following bony structures is present behind the marked area?



a) Post clivus

b) Lesser wing of sphenoid

c) Sella turcica

d) Carotid canal

Question 20:

All of the following nerves are involved in the movement of the eyeball except:



a) A

b) B

c) C

d) D

Question 21:

The middle radioulnar joint is an example of a:

a) Plane synovial

b) Primary cartilaginous

c) Secondary cartilaginous

d) Fibrous joint

Question 22:

What is the action of the muscle at the hip, which is attached to the marked area in the image below.

a) Abduction and external rotation

b) Extension and lateral rotation

c) Extension, adduction, and lateral rotation

d) Flexion and abduction

Question 23:

Identify the level of the cross-section shown in the image below.



a) T_1 - T_2

b) T_2 - T_3

c) T_3 - T_4

d) T_4 - T_5

Question 24:

A patient is unable to abduct his arm. Which of the following root values is likely to be affected?

a) C_4

b) C_5

c) C_6

d) C_7

Answer Key

Question No.	Correct Option
1	c
2	d

3	b
4	b
5	a
6	b
7	d
8	a
9	b
10	b
11	a
12	d
13	c
14	a
15	a
16	a
17	d
18	b
19	b
20	a
21	d
22	b
23	c
24	b

Detailed Explanations

Solution to Question 1:

The structure marked is conus cordis (middle part of bulbus cordis). The infundibulum of the heart develops from this structure.

The model shows the embryologic stage of the heart at which both the heart tubes have fused and are undergoing dextro-looping or bending. This happens around day 23 of intrauterine life.

Heart tube and its derivatives	
Part of the heart tube	Derivative structure in the he art

Heart tube and its derivatives	
Bulbus cordisProximal 1/3Middle 1/3 (conus cordis)Distal 1/3 (truncus arteriosus)	Trabeculated part of Rt.ventricleOutflow tract of Rt. and Lt.ventriclesRoot and proximal portions of aorta and pulmonary artery
Primitive ventricle	Trabeculated part of the Lt.ventricle
Primitive atrium	The common atrium that splits to Rt. and Lt.atrium



Solution to Question 2:

The area marked in the given image is the gluteal tuberosity. The muscle that is inserted here is the gluteus maximus which causes extension and lateral rotation of the hip joint.

Gluteus maximus inserts onto iliotibial tract (3/4th part) and deep fibers (1/4th) attach on gluteal tuberosity (marked in image). It is supplied by the inferior gluteal nerve. It is the chief extensor of the lower limb at the hip joint. It additionally also causes lateral rotation, abduction.



Solution to Question 3:

The area marked in the given image is substantia nigra (meaning- dark substance). The neurotransmitter secreted by this area is dopamine.

Substantia nigra is a part of the nigrostriatal pathway. Degeneration of dopaminergic neurons is implicated in Parkinson's disease.

Other dopaminergic pathways include:

- Mesocortical
- Mesolimbic
- Tuberoinfundibular







Solution to Question 4:

The area marked by an arrow in the given image is pterion and the artery that passes in close proximity to this area is the middle meningeal artery.

The pterion is an H shaped sutural junction between four bones of the skull:

• Frontal bone

• Squamous part of the temporal bone

• Greater wing of the sphenoid bone

• Parietal bone.

It is an important structure as it overlies the anterior branch of the middle meningeal artery and lateral fissure of the cerebral hemisphere. The bone in this area is thin and any injury to this area

may lead to rupture of the middle meningeal artery leading to extradural hematoma.



Solution to Question 5:

Ito cells/Hepatic stellate cells are located in the space of Disse. They are mesenchymal in origin.

The space of Disse or the perisinusoidal space (or space of Disse) is a location in the liver between hepatocytes and the liver sinusoid.

Significance of Ito cells:

Ito cells are in a dormant state for the majority of the duration. They store the fat-soluble vitamin A in their lipid droplets and are a signi■cant source of growth factors involved in liver homeostasis and regeneration.

When there is liver injury, Ito cells get activated. They replace damaged hepatocytes with collagenous scar tissue and secrete intralobular matrix components, including collagen type III (reticular) ■bres. This leads to hepatic ■brosis.

Note: A narrow space which separates limiting plate (a sleeve of modified hepatocytes) from the connective tissue of the portal area is known as space of moll.



Solution to Question 6:

Absence of striated duct in exocrine pancreas differentiates it from the parotid gland.

The exocrine pancreas is comprised entirely of serous acini. Hence, its general structure is similar to that of the parotid gland, but the two are easily distinguished by:

• The parotid gland has striated and intercalated intralobular ducts, whereas the pancreas has only the intercalated type.

• The excretory ducts of the parotid gland are striated columnar, whereas the excretory duct of the pancreas is a simple columnar.

Solution to Question 7:

Medial lemniscus is a continuation of fasciculus gracilis.

Posterior column–medial lemniscal pathway:

First-order axons carrying discriminative touch and vibration sensations enter the dorsal horn and continue traveling upward in the posterior white column as the fasciculus gracilis and fasciculus cuneatus.

Fibers of these fasciculi synapse on the second-order neurons of the ipsilateral nuclei gracilis and cuneatus in the medulla oblongata, respectively.

Axons of these second-order neurons then cross over as the internal arcuate fibers to form the medial lemniscus in the contralateral medulla.

These second-order axons ascend through the brainstem to reach the third-order neuronal cell bodies in the ventral posterior lateral nucleus of the thalamus. Axons from the third-order neurons then project through the posterior limb of the internal capsule to reach the primary

somatosensory cortex.



Solution to Question 8:

The fibers present in the fornix are association fibers except for the hippocampal commissure of fornix which has commissural fibers. Hence the most appropriate answer is option A. i.e both association and commissural fibers.

Association fibers connect various cortical regions within the same hemisphere whereas commissural fibers connect corresponding regions of two hemispheres.

Fornix arises from the hippocampus and ends in the mamillary body of the hypothalamus. It is supplied by the anterior cerebral artery.

Note: The function of the commissure of the fornix is to connect the hippocampal formations of the two sides.

Solution to Question 9:

The decussation of the superior cerebellar peduncle occurs in the midbrain, at the level of the inferior colliculus.

The midbrain has four decussations:

• At the level of the superior colliculus:

• Ventral tegmental decussation of the rubrospinal tract

• Dorsal tegmental decussation of the tectospinal tract

• At the level of the inferior colliculus:

• Decussation of superior cerebellar peduncles

• Decussation of trochlear nerve

Solution to Question 10:

All the above structures are enclosed by the pretracheal layer of deep cervical fascia except sternocleidomastoid. It is enclosed by the investing layer of deep cervical fascia.

The deep cervical fascia of the neck is organized into the following layers:

• The investing layer, which surrounds all structures in the neck

• The prevertebral layer, which surrounds the vertebral column and the deep muscles associated with the back.

• The pretracheal layer, which encloses the viscera of the neck

• The carotid sheath, which receives a contribution from the other three fascial layers

In the neck, the pretracheal fascia covers the strap muscles i.e sternohyoid, omohyoid, sternothyroid, thyrohyoid. It splits to enclose the thyroid gland. It also surrounds the trachea and the esophagus including the recurrent laryngeal nerve present in the groove between them.

Solution to Question 11:

The sympathetic chain is not the content of the carotid sheath. It lies posterior to the carotid sheath and anterior to the transverse processes of the cervical vertebrae in the neck.

The carotid sheath is a layer of loose connective tissue formed by adjacent portions of the investing layer of deep cervical fascia, the pretracheal fascia, and the prevertebral fascia.

Contents of carotid sheath:

• Common and internal carotid arteries,

• Internal jugular vein,

• Vagus nerve

• Ansa cervicalis

Note: The glossopharyngeal, accessory, and hypoglossal nerves lie within the upper part of the carotid sheath.

Solution to Question 12:

Paradoxical respiration occurs due to the damage of the phrenic nerve, which is marked as D, in the given image.

The structures marked in the image are:

A. Right vagus nerve

B. Left vagus nerve

C. Recurrent laryngeal nerve

D. Right phrenic nerve



Paradoxical movements in the chest are as follows:

• Paradoxical respiration - the abnormal movement of the chest wall or abdominal wall, where they move in during inspiration instead of moving out. It is seen in flail chest.

• Paradoxical movement of the diaphragm - normally, the diaphragm moves down during inspiration. However, in cases of phrenic nerve palsy, the affected side of the diaphragm moves upward (paradoxical respiration) due to diaphragmatic paralysis. The gold standard investigation to detect this is the electrical or magnetic stimulation of the phrenic nerve with a recording of the compound muscle action potential and/or the transdiaphragmatic pressure. This can be observed on ultrasound or fluoroscopy.

Solution to Question 13:

The area marked in the image is the palmar aspect of the first metacarpal. The opponens pollicis muscle is inserted at this site.

Opponens pollicis:

Origin: Tubercle of the trapezium and flexor retinaculum

Insertion: Whole length of the lateral border and adjoining lateral half of the palmar aspect of the first metacarpal.

Action: Flexes the metacarpal bone of the thumb.

The image below shows various intrinsic muscles of the hand.

Superficial muscles of the hand (palmar view)



Deep muscles of the hand (palmar view)



Interossei muscles of left hand (palmar view)





Solution to Question 14:

The structure marked as 'A', is the hyoid bone, which is not a part of the laryngeal skeleton.

The structures marked in the image are:

• Hyoid bone

• Thyroid cartilage

• Epiglottis

• Cricoid cartilage

The laryngeal skeleton is formed by a series of cartilages that are connected by ligaments and fibrous membranes:

• The thyroid, cricoid, and the greater part of the arytenoid cartilages consist of hyaline cartilage.

• The corniculate, cuneiform, tritiate, and epiglottic cartilages and the apices of the arytenoid cartilage consist of elastic cartilage.

Solution to Question 15:

The image shows the sphenoid bone. The foramen marked is the foramen rotundum. Maxillary branch of trigeminal nerve passes through it.



Solution to Question 16:

Reticular fibres are not present in the thymus.

Reticular fibres are fine, anastomosing fibres composed of mainly type III collagen. They differ from type I collagen fibres in that, they are finer and are arranged in the form of a network (reticulum). The fibres appear black in colour on staining with silver impregnation. Due to this affinity to silver salts, they are also known as argentophil fibres.



Reticular fibres are mainly found in:

• All lymphoid organs like spleen, lymph node, and bone marrow (except thymus).

• Glands, liver and kidneys.

• The basement membrane of all epithelia.

• Smooth muscle and nerve cells.

Solution to Question 17:

Pectoralis major is mainly involved in adduction and medial rotation of the arm and is not involved in the abduction of the arm.

Glenohumeral abduction is about 120° or more. The glenohumeral arthrodesis restricts the abduction at about 45°. Further abduction occurs at the sterno and acromioclavicular articulations, and contralateral vertebral flexion.

Muscles involved in the abduction of the shoulder:

• The first 15° of abduction is initiated by the supraspinatus muscle.

• 15° to 90° of abduction is carried out mainly by the middle fibers of the deltoid muscle.

• Abduction of the arm, beyond 90°, is carried out by the movement of the shoulder girdle caused by the action of trapezius and serratus anterior.

Solution to Question 18:

The image shows an arrow mark pointing towards the cilia. All the structures given in the options have a ciliated lining epithelium except epididymis, which is lined by pseudostratified columnar epithelium with stereocilia.

The given image is a histological section of the epididymis showing the pseudostratified columnar epithelium with stereocilia.



Cilia are minute hair-like projections, while stereocilia are very long, thick microvilli measuring about 5–10 μm in length.

Ciliated epithelium:

• The epithelium bears cilia, which moves in a coordinated fashion, and help in the movement of the mucus, or propel the movement of any other structures such as ova or spermatozoa.

• The ciliated columnar epithelium is found in most of the respiratory tract, the uterus, and the uterine tube (oviduct), efferent ductules of the testis, parts of the middle ear and auditory tube; the ependyma lining the central canal of the spinal cord and the ventricles of the brain.

• The pseudostratified ciliated columnar epithelium is seen in the trachea and in large bronchi.

Solution to Question 19:

The structure marked in the image is pterion, and the bony structure that is present behind it is the lesser wing of the sphenoid.

The pterion is an H-shaped sutural junction between four bones of the skull:

• Frontal bone

• Squamous part of the temporal bone

• Greater wing of the sphenoid bone

• Parietal bone



The pterion overlies the anterior branch of the middle meningeal artery and lateral fissure of the cerebral hemisphere. The bone in this area is thin and any injury to this area may lead to rupture of the middle meningeal artery leading to extradural hematoma.

The pterion also corresponds to the site of the anterolateral (sphenoidal) fontanelle of the neonatal skull, which closes in the third month after birth.



Solution to Question 20:

The nerve marked as 'A' is the optic nerve (II CN), and is not involved in the movement of the eyeball.

The nerves marked in the image are:

• Optic nerve (II CN)

• Oculomotor nerve (III CN)

• Abducens nerve (VI CN)

• Trochlear nerve (IV CN)



Extraocular muscles involved in the movement of the eyeball:

• Superior rectus, inferior rectus, medial rectus, inferior oblique, and levator palpebrae superioris muscles - supplied by the oculomotor (III) nerve.

• Lateral rectus - supplied by abducens (VI) nerve.

• Superior oblique - supplied by trochlear (IV) nerve.

Solution to Question 21:

The middle radioulnar joint is an example of a fibrous joint (syndesmosis).

The radius and ulna are united with each other by three joints:

• Proximal (superior) radioulnar joint

• Middle radioulnar joint

• Distal (inferior) radioulnar joint

Proximal and distal radioulnar joints are uniaxial pivot joints (synovial) and are involved in pronation and supination movements of the forearm.

The middle radioulnar joint is a fibrous joint (syndesmosis). The radius and ulna, in the middle, are connected by two syndesmoses namely:

• The oblique cord - extends from the lateral side of the ulnar tuberosity to the radius, distal to the radial tuberosity.

• Interosseous membrane - a broad, thin collagenous structure, which runs perpendicular to the oblique cord. It extends distomedially between the interosseous borders of the radius and ulna.

Based on the structure, joints are classified into three types:

• Fibrous joints

• Sutures

• Syndesmosis

• Gomphosis

• Cartilaginous joints

• Primary cartilaginous joints or synchondrosis

• Secondary cartilaginous joints or symphysis

• Synovial joints

• Ball-and-socket or spheroidal joints

• Sellar or saddle joints

• Condylar or bicondylar joints

• Ellipsoid joints

• Hinge joints

• Pivot or trochoid joints

• Plane joints





Solution to Question 22:

The image shows the posterior surface of the femur, and the structure marked is gluteal tuberosity, to which the gluteus maximus muscle is attached. It causes extension and lateral rotation at the hip joint.

The femur can be divided into:

• The proximal femur - Consists of the head, neck, greater and lesser trochanters. The intertrochanteric line is present on the anterior surface at the junction of the neck and shaft of the femur while the intertrochanteric crest is present on the posterior surface.

• The shaft of the femur - Consists of linea aspera (bony crest present on the posterior surface), gluteal tuberosity, and pectineal line

• The distal femur - Consists of the medial condyle, lateral condyle, intercondylar fossa, and patellar surface





The image below shows attachments of various muscles over the posterior surface of the right femur.



Gluteus maximus muscle:

Origin:

• Posterior aspect of the dorsal ilium, posterior to the posterior gluteal line

• Posterior superior iliac spine

• Posterior aspect of sacrum and coccyx

• Sacrotuberous ligament.

Insertion - It is inserted mainly in fascia lata at the iliotibial band and also into the gluteal tuberosity on the posterior femoral surface.

Action:

• Major extensor of the hip joint

• Lateral rotation of the hip joint

• Active in powerful abduction of the thigh

Arterial Supply - Inferior and superior gluteal arteries and the first perforating branch of the profunda femoris artery.

Nerve Supply- Inferior gluteal nerve (L5, S1 and S2).

Solution to Question 23:

The cross-section shown in the image is at the level of T3-T4.

The cross-section shows a large vascular opening of the arch of the aorta, along with three small openings corresponding to the brachiocephalic trunk, left common carotid artery and left subclavian artery. This suggests that the section was taken at the level between T3 and T4.

Options A, B: In cross-sections at the levels of T1-T2 and T2-T3, only the small vascular openings of the branches arising from the arch of aorta would be visible.

Option D: In cross-section at the level of T4-T5, where, the arch of the aorta begins and ends, bifurcation of the pulmonary trunk is visible.



Solution to Question 24:

The inability to abduct the arm is most likely due to a lesion of the C5 root.

The deltoid muscle is the main muscle involved in the abduction of the arm. It is innervated by the C5 root of the axillary nerve. So, a lesion involving the C5 causes inability to abduct the arm.

The table below shows the movements and muscles tested to determine the location of the lesion in the upper limb.

Arm movement	Muscle	Root	Nerve
Shoulder abduction	Deltoid	C5	Axillary
Elbow extension	Triceps	C7	Radial
Elbow flexion	BicepsBrachioradialis	C5/6 C6	MusculocutaneousRadial
Radial wrist extensor	Extensor carpi radialis longus	C6	Radial
Finger extensors	Extensor digitorum	C7	Posterior interosseus
Finger flexors	Flexor pollicis longus and flexor digitorum profundus, index Flexor digitorum profundus, ring and little	C8C8	Anterior interosseousUlnar
Finger abduction	First dorsal interosseusAbductor pollicis brevis	T1T1	UlnarMedian

Anatomy NEET 2018

Question 1:

Identify the thickened nerve marked in the arrow in the image shown below:



a) Great auricular nerve

b) Lesser occipital nerve

c) Facial nerve

d) Auriculotemporal nerve

Question 2:

Broca's area is situated in the _____.

a) Inferior frontal gyrus

b) Superior temporal gyrus

c) Angular gyrus

d) Postcentral gyrus

Question 3:

Which of the following cranial nerve nuclei is not a part of the special visceral afferent (SVA) column?

a) Facial nerve

b) Trochlear nerve

c) Vagus nerve

d) Glossopharyngeal nerve

Question 4:

The joint between the attachment of the 8th and 9th rib to the 7th rib is a:

a) Fibrocartilaginous joint

b) Symphyseal joint

c) Synovial joint

d) Fibrous joint

Question 5:

Which is the last fontanelle to close?

a) Anterior

b) Posterior

c) Occipital

d) Mastoid

Question 6:

Which among the following is incorrect regarding Sibsons's fascia?

a) Attached to the inner border of 2nd rib

b) Covers apical part of lung

c) Part of scalenus minimus muscle

d) Vessels pass above the fascia

Question 7:

The structure not passing through the aortic opening of the diaphragm is:

a) Aorta

b) Azygous vein

c) Thoracic duct

d) Vagal trunk

Question 8:

The hard palate contains _____

a) Keratinised epithelium, submucosa, minor salivary glands

b) Keratinised epithelium, absent submucosa, no salivary glands

c) Non-keratinised epithelium, submucosa, minor salivary glands

d) Non-keratinised epithelium , absent submucosa, minor salivary glands

Question 9:

What is the function of the muscle marked as A in the image below, at the metacarpophalangeal joint?



a) Flexion

b) Extension

c) Adduction

d) Abduction

Question 10:

Which among the following is a tensor of vocal cords?

a) Cricothyroid

b) Lateral cricoarytenoid

c) Thyroarytenoids

d) Posterior cricoarytenoids

Question 11:

The nerve of the second branchial arch is:

a) Glossopharyngeal nerve

b) Trigeminal nerve

c) Facial nerve

d) Vagal nerve

Question 12:

Which of the following statements is true about the iliopubic tract?

a) It is a thickening of the lumbar fascia

b) It lies anterior to the Poupart's ligament

c) It inserts into the pubic tubercle in front of the conjoint tendon

d) Its fibers fan out with fascia iliaca laterally

Question 13:

Ureteric bud develops from:

a) Mesonephric duct

b) Metanephros

c) Pronephros

d) Genital sinus

Question 14:

The heart starts to beat in the _____week of development.

a) 2nd week

b) 4th week

c) 6th week

d) 10th week

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	c
5	a
6	a
7	d
8	a
9	a
10	a
11	c
12	d
13	a
14	b

Detailed Explanations

Solution to Question 1:

The thickened nerve marked with the arrow in the above image is the great auricular nerve and the thickened nerve visible below the great auricular nerve is the transverse cervical nerve.

Nerve thickening in this case is likely to be due to leprosy.



Great auricular nerve:

It is the largest ascending branch of cervical plexus and arises from the second and the third cervical spinal nerves.

It winds over the posterior border of the sternocleidomastoid muscle and ascends upwards. On reaching the parotid gland it divides into the anterior and posterior branches:

• Anterior branch - supplies the facial skin over the parotid gland and joins the facial nerve in the gland.

• Posterior branch - supplies the skin over the mastoid process, posteroinferior part of the auricle, lobule, and concha.

Clinical considerations:

• Leprosy - Nerve thickening is present. The nerve is palpable over the upper part of the sternocleidomastoid muscle. (as seen in the above image)

• Frey's syndrome - An aberrant connection between the auriculotemporal nerve and great auricular nerve causes gustatory sweating.

Solution to Question 2:

Broca’s area is situated in the inferior frontal gyrus.

Broca’s motor speech area (44,45) is located in the inferior frontal gyrus.

Wernicke’s area (22) is located in the superior temporal gyrus.



Solution to Question 3:

The trochlear nerve nucleus is a part of the general somatic efferent (GSE) column, not the special visceral afferent column.

The trochlear nerve is a motor nerve that innervates superior oblique muscle in the orbit.

The neurons of the general somatic efferent (GSE) column innervate the extrinsic ocular muscles and muscles of the tongue (III, IV, VI, and XII cranial nerves).



Solution to Question 4:

The interchondral joint between the attachment of the 8th and 9th rib with the 7th rib is a synovial joint.

Joints in the thoracic cavity are of the following types:

• Synarthrosis - structures are connected by a fibrous band of tissue. E.g, 1st sternocostal joint

• Synchondrosis - primary cartilaginous joints. E.g., costochondral joints

• Symphysis - secondary cartilaginous joints. E.g. manubriosternal and xiphisternal joints

• Synovial joints - the presence of these help the thoracic cavity expand during inspiration. E.g.:

• Interchondral joint

• Sternocostal joint

• Costovertebral joint

• Costotransverse joint

Note:

The interchondral joint between the 9th and 10th cartilages is never synovial and sometimes absent.



Solution to Question 5:

The anterior fontanelle is the last fontanelle to close. It closes at around 18-24 months of age.

Fontanelles of the skull:

Fontanelle	Location	Closes at
Anterior	Between frontal and parietal bonesJunction marked by br egma	18-24 months

Fontanelle	Location	Closes at
Posterior	Sagittal and lambdoid sutures	2-3 months
Sphenoid	Pterion	2-3 months
Mastoid/Posterolateral	Intersection of temporal, parietal, and occipital bones.	6-18 months (average 12 months)



Solution to Question 6:

Sibson’s fascia is attached to the inner border of the 1st rib, not the 2nd rib.

Sibson's fascia (suprapleural membrane) serves as a plane of separation between the lower neck and the thorax and covers the apical part of the lung. It is attached to the inner border of the 1st rib and posteriorly to the anterior border of the transverse process of the C7 vertebra. It is strengthened by a few spreading muscular fibers derived from the scalene muscles. The subclavian vessels lie on top of it.



Solution to Question 7:

The vagal trunk does not pass through the aortic opening of the diaphragm.

It passes through the esophageal opening at the level of T10 vertebra.

Solution to Question 8:

The hard palate contains keratinised stratified squamous epithelium, has a submucosa, and contains minor salivary gland. This keratinised stratified squamous epithelium has regional variations and may be orthokeratinised or parakeratinised(found in hard palate and gingiva).

The bony structure of the hard palate is covered by a firmly attached mucosa in the central part. In the lateral parts, the hard palate also has a submucosal layer containing blood vessels. There are minor mucous type salivary glands in the submucosa in the posterior part of the hard palate.

(Note: In the orthokeratinized epithelium the cell nuclei disappear in the keratinized layer, whereas in the parakeratinized epithelium flattened, highly condensed nuclei remain in the cell cytoplasm of the keratinized layer until exfoliation)

Solution to Question 9:

The muscle marked 'A' is lumbricals and they cause flexion at the metacarpophalangeal joints.

Lumbricals are intrinsic muscles of the hand that flex the metacarpophalangeal joints (MCP) and extend the interphalangeal joints (PIP & DIP).

They originate from flexor digitorum profundus.

• The lateral two are unipennate, supplied by the median nerve.

• The medial two are bipinnate, supplied by the ulnar nerve.

Superficial muscles of the hand



Superficial muscles of the hand (palmar view)



Solution to Question 10:

Cricothyroid is a tensor of vocal cords.

Solution to Question 11:

The facial nerve is the nerve of the 2nd pharyngeal arch.

Each pharyngeal arch is characterized by its own muscular components. The muscular components of each arch have their own cranial nerve. Wherever the muscle cells migrate, they carry their nerve component with them.

The neurons of the 5th, 7th, 9th, and 10th cranial sensory ganglia that supply the pharyngeal arch structures are derived from ectodermal placodes (epipharyngeal placodes) and from the neural crest cells.

Solution to Question 12:

The fibers of the iliopubic tract fan out with fascia iliaca laterally.

The iliopubic tract is a thickening of the transversalis fascia lying immediately posterior to the inguinal (Poupart's) ligament. It extends from the anterior superior iliac spine (ASIS) superolaterally and arches across the front of the femoral sheath.

Attachments:

• Medially, its fibres insert by a broad attachment into the pubic tubercle and pectineal line, behind the conjoint tendon.

• Laterally, its fibres fan out within the fascia transversalis and fascia iliaca without bony attachment to the iliac spines.

Clinical note: It is considered as an important landmark in laparoscopic hernia surgeries.



Solution to Question 13:

Ureteric bud develops from Mesonephric duct.

• Intermediate mesoderm is a type of mesoderm that is located between the paraxial mesoderm and lateral plate.

• It develops into a part of the urogenital system as well as the reproductive system.

• Intermediate mesoderm develops into Genital ridge and Nephrogenic cord.

• Nephrogenic cord further develops into Pronephros, Mesonephros and Metanephros.

• Ureteric bud develops from mesonephros which later fuses with Mesonephric duct.

• Metanephros proliferates around the ureteric bud which forms metanephric blastema (definitive kidneys).

Structures arising from metanephric blastema and ureteric bud are as follows -

Metanephric blastema	Ureteric bud
Bowmans capsule	Collecting tubules
Proximal convoluted tubule	Renal papillae
Loop of Henle	Calyces
Distal convoluted tubule	Renal pelvis
	Ureter



Solution to Question 14:

The heart starts to beat in the 4th week of development.

Anatomy NEET 2019

Question 1:

The movements at the _____ joint permit a person to look to the right and left.

a) Atlanto-occipital

b) Atlanto-axial

c) C2-C3

d) C3-C4

Question 2:

All of the following are true about the right coronary artery except _____

a) It's diameter is less than left coronary artery

b) It arises from the anterior aortic sinus

c) It gives rise to circumflex coronary branch

d) Right conal artery is it's first branch

Question 3:

What is the basis for the formation of omphalocele?

a) Excessive length of the intestine

b) Physiological hernia failing to go back

c) Herniation of liver

d) Herniation of umbilicus

Question 4:

Which fibre is marked by the arrow in the image given below?



a) Short association

b) Long association

c) Projection

d) Commissural

Question 5:

Which nerve passes through the foramen marked in the image below?



a) Lingual nerve

b) Mandibular nerve

c) Chorda tympani nerve

d) Inferior alveolar nerve

Question 6:

Boundaries of Calot triangle are formed by all except:

a) Common hepatic duct

b) Cystic duct

c) Cystic artery

d) Gall bladder

Question 7:

Area marked in the given image is supplied by dermatome number?



a) T8

b) T9

c) T10

d) T11

Question 8:

The arrow in the image denotes which of the following muscles?



a) Extensor carpi radialis longus

b) Brachioradialis

c) Supinator

d) Flexor carpi radialis brevis

Question 9:

What is the nerve supply of the structure marked in the image below?



a) Posterior interosseus nerve

b) Median nerve

c) Anterior interosseus nerve

d) Ulnar nerve

Question 10:

Nerve supply of the area marked by arrow in the image is:



a) Radial nerve

b) Posterior interosseus nerve

c) Median nerve

d) Ulnar nerve

Question 11:

Cholecysto-vena caval line separates which of the following?

a) Gallbladder and inferior vena cava

b) Porta hepatis and inferior vena cava

c) Caudate lobe and quadrangular lobe

d) Right and left lobe of liver

Question 12:

Which of the following is false about skin?

a) Dermis is derived from ectoderm

b) Skin accounts for 15% of total body weight

c) Epidermis is derived from ectoderm

d) Dermis is made up of type 1 and type 3 collagen

Question 13:

All of the following structures are derived from the aponeurosis of the external oblique muscle except:

a) Pectineal ligament

b) Inguinal ligament

c) Lacunar ligament

d) Linea semilunaris

Question 14:

Which of the following lobes of the prostate forms the uvula vesicae?

a) Anterior

b) Posterior

c) Median

d) Lateral

Question 15:

Foot drop is caused by damage to:

a) Common peroneal nerve

b) Femoral nerve

c) Tibial nerve

d) Sciatic nerve

Question 16:

Which of the following is true regarding De Quervain's tenosynovitis?

a) Fingers are held in mild extension

b) It affects APL and EPB

c) Most common involvement is index finger

d) Treatment is surgery

Question 17:

Identify the structure marked B in the image given below :



a) Pineal gland

b) Falx cerebri

c) Fornix

d) Pituitary gland

Question 18:

1st pharyngeal arch gives rise to all except:

a) Medial pterygoids

b) Levator veli palatini

c) Lateral pterygoids

d) Tensor veli palatini

Answer Key

Question No.	Correct Option
1	b
2	c
3	b
4	c
5	d
6	d
7	c
8	a
9	b
10	a
11	d
12	a
13	d
14	c
15	a
16	b
17	c
18	b

Detailed Explanations

Solution to Question 1:

The movements at the atlanto-axial joint allow a person to look left and right.

The atlanto-axial joints include 3 synovial joints:

• Median atlanto-axial joint- The central pivot joint between the dens of the axis and a ring formed by the anterior arch and transverse atlantal ligament

• Lateral atlanto-axial joints- Plain synovial joints between lateral masses of atlas and upper facets of the body of axis

Movements at the atlanto-axial joint almost exclusively is rotation around the axis, which help side-to-side rotation of the head. The normal range of atlanto-axial rotation is about 40°.



Superior view of atlas

Solution to Question 2:

The right coronary artery does not give rise to the circumflex coronary artery.

The right coronary artery arises from the anterior aortic sinus (right coronary sinus) of the ascending aorta. It passes anteriorly and descends vertically in the coronary sulcus between the right atrium and the right ventricle. On reaching the inferior margin, it turns posteriorly and continues in the sulcus onto the diaphragmatic surface and the base of the heart.

The diameter of the right coronary artery is less than that of the left coronary artery.

The right conal artery is, usually the first branch of the right coronary artery.

The circumflex coronary artery is a branch of the left coronary artery and is the exclusive supplier to the right bundle branch.



Solution to Question 3:

Omphalocele is protrusion of abdominal contents through the umbilical ring. It is due to failure of physiological hernia to reduce into the abdominal cavity, which results in non-closure of the umbilical ring. .

An omphalocele may contain stomach, liver, and intestines. And these contents are enclosed in a sac.

Omphalocele is usually associated with other congenital anomalies (e.g., trisomy 13, trisomy 18, or Beckwith-Wiedemann syndrome) and with increased levels of α-fetoprotein.

Note: In gastroschisis, there is an absence of a membranous sac and is paraumbilical.



Solution to Question 4:

The marked structure is the internal capsule, which is a form of projection fiber.



Solution to Question 5:

The marked structure is mandibular foramen and inferior alveolar nerve passes through it.

The inferior alveolar nerve (IAN), a branch of the mandibular nerve (CN V3), it descends behind the lateral pterygoid, at the lower border of the muscle, the nerve passes between the sphenomandibular ligament and the mandibular ramus and enters the mandibular canal via the mandibular foramen

The mandibular foramen is located above the mandibular angle, the opening through which the IAN and the vessels of the mandible pass.

The IAN continues its course through the mandibular canal and then branches off to supply the lower teeth, gums, and lower lip.



Solution to Question 6:

The gall bladder does not form a boundary of the Calot's triangle.

Calot's triangle is bounded by 3C's:

• Medially – Common hepatic duct

• Inferiorly – Cystic duct

• Superior – Cystic artery

Solution to Question 7:

The marked dermatome which includes the umbilicus is supplied by T10.

The dermatomal distribution of the abdominal region is as follows:

• T7, T8, T9 - supplies skin from the xiphoid process to just above the umbilicus.

• T10 - supplies skin around the umbilicus.

• T11, T12 - supplies skin below the umbilicus upto the suprapubic region.

• L1 - supplies skin around the genitalia and upper medial thigh.



Solution to Question 8:

The arrow shows extensor carpi radialis longus.

It originates from the distal third of the lateral supracondylar ridge of the humerus and inserts at the second metacarpal (radial side). It is innervated by the radial nerve, C6, and C7.

The vascular supply is by radial recurrent artery and it is responsible for the extension and abduction of the wrist joint.



Solution to Question 9:

The marked structure is second lumbrical which is supplied by the median nerve.

The lateral two lumbricals are supplied by the median nerve and the medial two are supplied by the ulnar nerve. Function of these muscles is flexion at the metacarpophalangeal joints and extension at both the interphalangeal joints.



Solution to Question 10:

The area marked in the image is supplied by radial nerve.

Solution to Question 11:

Cholecysto-vena caval line (Cantlie line) separates right and left lobe of the liver.

Cantlie line is an imaginary line that that divides the liver into 2 planes. It extends from the midpoint of the gallbladder fossa at the inferior margin of the liver back to the midpoint of the inferior vena cava (IVC) and contains the middle hepatic vein.

Solution to Question 12:

Dermis is derived from mesoderm and epidermis is derived from ectoderm.

Solution to Question 13:

Linea semilunaris is derived from the aponeurosis of the internal oblique muscle.

Pectineal ligament, inguinal ligament and lacunar ligament are derived from the external oblique aponeurosis.

Solution to Question 14:

The median lobe of the prostate forms the uvula vesicae.

Uvula vesicae is an elevation in the mucous membrane of the urinary bladder, behind the internal urethral orifice. It is often enlarged in benign prostatic hypertrophy.



Solution to Question 15:

Foot drop is caused by damage to the common peroneal nerve.

The common peroneal nerve is a component of the sciatic nerve. In the posterior thigh, it supplies the short head of the biceps femoris. It curves around the neck of the fibula deep to peroneus longus and divides into superficial and deep peroneal nerves.

• The superficial peroneal nerve innervates peroneus longus and peroneus brevis which are the evertors of the foot and to the skin over the dorsal area of the foot except for the first webspace and lateral side of the little toe.

• The deep peroneal nerve innervates all muscles of the anterior compartment which are the dorsiflexors of the foot, extensor digitorum brevis, first two dorsal interossei muscles, and skin between great and second toes.

Hence, injury to the common peroneal nerve will lead to:

• Weakness of dorsiflexion of foot or foot drop

• Loss of eversion

• Loss of extension of toes

• Sensory loss

• Anterior and lateral side of the leg

• Dorsum of the foot including the medial side of the great toe

Th image given below shows the course of common peroneal nerve.



Solution to Question 16:

De Quervain's disease is a stenosing tenosynovitis of the extensor tendons of the first dorsal compartment, namely extensor pollicis brevis (EPB) and abductor pollicis longus (APL).

It is prominent among females, especially pregnant women. Most commonly thumb is involved.

Finkelstein's test: On grasping the patient's thumb and quickly abducting the hand ulnar ward, the pain over the radial styloid tip is excruciating. It's the most pathognomonic objective sign but is also seen in other conditions like radial nerve entrapment and arthritis.

Treatment:

• Majority of the patients are treated medically

• Only few require the release of the 1st dorsal compartment which houses the APL and EPB tendons.

Solution to Question 17:

The image given shows a midsagittal section of the brain. The structure marked B is the Fornix.



Solution to Question 18:

Levator veli palatini is not derived from first pharyngeal arch.

All soft palate muscles including levator veli palatini are derived from the 4th pharyngeal arch and are supplied by vagal plexus.

First pharyngeal arch gives rise to:

• Muscles of mastication (temporalis, masseter, lateral and medial pterygoids)

• Mylohyoid

• Anterior belly of digastric

- Tensor tympani
- Tensor veli palatini and
- Anterior 2⁄3 of tongue.

Anatomy NEET 2020

Question 1:

Which of the following forms the lateral boundary of the anatomical snuff box?

a) Extensor pollicis brevis and abductor pollicis longus

b) Extensor pollicis longus and abductor pollicis brevis

c) Extensor pollicis longus and extensor pollicis brevis

d) Abductor pollicis longus and abductor pollicis brevis

Question 2:

Identify the cell marked in the histological image of the cerebellum shown below :



a) Purkinje cells

b) Basket cells

c) Golgi cells

d) Granule cell

Question 3:

Which of the following is not used for the identification of the facial nerve trunk?

a) Retrograde dissection from distal branch

b) Tragal pointer

c) Inferior belly of omohyoid muscle

d) Posterior belly of digastric

Question 4:

The false statement about fetal circulation is:

a) Anatomical closure of foramen ovale is by 3 to 5 days

b) Anatomical closure of ductus venosus is by 3 to 7 days

c) Right umbilical vein is absent

d) Anatomical closure of ductus arteriosus is by 2 to 3 weeks

Question 5:

A tumor of the uncinate process of the pancreas affects which of the following vessels?

a) Superior mesenteric artery

b) Portal vein

c) Common hepatic artery

d) Inferior mesenteric artery

Question 6:

A patient presented with clinical features of ataxia and incoordination. This is due to thrombosis of which artery?

a) Posterior cerebral

b) Middle cerebral

c) Internal carotid

d) Superior cerebellar

Question 7:

Which of the following structures prevents the upward extension of thyroid swelling through its attachment to the thyroid cartilage?

a) Pretracheal fascia

b) Sternothyroid

c) Thyrohyoid membrane

d) Ligament of berry

Question 8:

A patient presented with the extension of MCP joint and flexion of IP joints. Which muscle is affected?

a) Extensor digitorum

b) Interossei and lumbricals

c) Abductor pollicis

d) Abductor pollicis brevis

Question 9:

Meiosis occurs in:

a) Adrenal

b) Adult ovary

c) Prepubertal testis

d) Hypothalamus

Question 10:

The long left recurrent laryngeal nerve is due to the persistence of which arch artery?

a) 3rd arch

b) 4th arch

c) 5th arch

d) 6th arch

Question 11:

The muscle marked by the arrow in the image below is innervated by the_____.



a) Dorsal scapular nerve

b) Suprascapular nerve

c) From the dorsal rami of C1

d) Subscapular nerve

Question 12:

A patient presented with weakness of the right side of the face with the loss of pain and temperature. Pain and temperature sensation of the left leg is also lost. The lesion is most likely located at?

a) Medial medulla

b) Lateral pons

c) Medial pons

d) Lateral medulla

Question 13:

Ligation of common hepatic artery will impair blood supply in:

a) Right gastric and right gastroepiploic artery

b) Right gastric and left gastric artery

c) Right gastroepiploic and short gastric vessels

d) Right gastric and short gastric vessels

Question 14:

The type of joint marked in the image below is _____:



a) Syndesmosis

b) Synarthrosis

c) Synchondrosis

d) Synovial

Question 15:

Identify the sign shown in the image below:

a) Popeye sign

b) Hill-Sachs sign

c) Griesinger sign

d) Rising sun sign

Question 16:

A patient presented with an enlarged inguinal lymph node. The most likely primary cause is

a) Testicular cancer

b) Anal cancer

c) Sigmoid colon cancer

d) Prostate cancer

Question 17:

Identify the type of cartilage shown in the image below:



a) Hyaline cartilage

b) Elastic cartilage

c) Articular cartilage

d) Fibrocartilage

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	a
5	a
6	d
7	b
8	b
9	b
10	d
11	a
12	b
13	a
14	d

15	a
16	b
17	b

Detailed Explanations

Solution to Question 1:

The lateral boundary of the anatomical snuff box is formed by extensor pollicis brevis and abductor pollicis longus.

Boundaries of the anatomical snuffbox:

• Lateral wall:

• Abductor pollicis longus (APL)

• Extensor pollicis brevis (EPB)

• Medial wall:

• Extensor pollicis longus (EPL)

• Roof:

• Cephalic vein

• Superficial branch of the radial nerve

• Floor:

• Radial styloid, scaphoid, trapezium, and 1st metacarpal

Only content: Radial artery



Solution to Question 2:

The cells marked in the above histological image are Purkinje cells.

Characteristics of Purkinje cells:

• Purkinje cells are the flask-shaped cells present in the Purkinje cell layer of the cerebellum.

• The cells are evenly spaced and are arranged in a single layer.

• A dendrite arises from the neck of each flask-shaped cell, which passes upward to enter the molecular layer, which then branches extensively to form a dendritic tree.

• The axon of each cell passes downwards, crosses the granular cell layer to enter the white matter.

• Axons of Purkinje cells are the only efferents of the cerebellar cortex.

• The initial segment of the axon receives synaptic contacts from basket cell axons.

• The main axon terminates in one of the cerebellar or vestibular nuclei.

• Purkinje cells are inhibitory and use GABA as their neurotransmitter.

Pathological significance:

• Negri bodies in rabies, are usually found as eosinophilic, cytoplasmic inclusions in the neurons of the hippocampus and Purkinje cells.

• Loss of Purkinje cells and granule cells is seen in ataxia-telangiectasia.

• In patients with subacute cerebellar degeneration, there is the destruction of Purkinje cells. Some patients show circulating PCA1 antibody (antiYo) against Purkinje cells.

• Purkinje cells and neurons in pyramidal cell layers of hippocampus and cerebral cortex are the most sensitive neurons to hypoxia, and are likely to be damaged in cases of global cerebral ischemia.

Solution to Question 3:

The inferior belly of the omohyoid muscle is not used as an anatomical landmark for the identification of the facial nerve trunk.

There are two methods for localization of the facial nerve trunk:

1. Anterograde method: The facial nerve trunk is identified based on the anatomical landmarks after it exits from the stylomastoid foramen. This includes:

• Tragal pointer/Conley's pointer - It is the inferior portion of the cartilaginous canal. The facial nerve trunk lies 1 cm deep and inferior to its tip.

• Posterior belly of digastric - The facial nerve lies superior to the upper border of the muscle.

• Squamotympanic fissure - It is a fissure in the temporal bone, which runs from TMJ to the tympanic cavity. The facial nerve runs through this fissure.

• Styloid process - The facial nerve lies superficial to it.

• Mastoid process - Upon drilling it, the nerve is identified proximally.

2. Retrograde method: The main branches of the facial nerve trunk are identified and are then traced proximally until the main tunk is identified. Some branches used for tracing the facial nerve trunk include:

• Buccal branch - It lies in relation to the parotid gland.

• Marginal mandibular branch - It lies in relation to the facial vessels.



Solution to Question 4:

The anatomical closure of the foramen ovale takes 1 month to a year.

Given below are the fetal circulatory adjustments after birth:

	Functional closure	Anatomical closu re
Ductus venosus	After removal of the placenta	7 days
Foramen ovale	At birth - after breathing is in itiated	Months to years
Ductus arterios us	10–15 hours	10–21 days

Solution to Question 5:

A tumor of the uncinate process of pancreas affects the superior mesenteric artery.

The superior mesenteric artery arises from the abdominal aorta. It lies first behind the body of the pancreas and then in front of the uncinate process.



Relations of the other parts of pancreas:

Head of pancreas:

• Anterior:

• 1st part of the duodenum

• Origin of the transverse mesocolon

• Posterior:

• Right renal vein

• Right renal artery

• Inferior vena cava

• Common bile duct (which is embedded in the head)

• Right crus of the diaphragm

• Termination of the right gonadal vein

Neck of the pancreas:

• Anterior:

• Lesser sac

• pylorus

• The anterior superior pancreaticoduodenal branch of the gastroduodenal artery

• Posterior:

• Portal vein

• Superior mesenteric vein

• Inferior mesenteric vein (in 1/3rd individuals)



Solution to Question 6:

Thrombosis of the superior cerebellar artery causes cerebellar lesion which presents with ataxia and incoordination.

Blood supply of cerebellum:

- Superior cerebellar artery
- Posterior inferior cerebellar artery
- Anterior inferior cerebellar artery

Solution to Question 7:

Sternothyroid and sternohyoid prevent the upward extension of the thyroid swelling through its attachment to the thyroid cartilage.

Pretracheal fascia encloses the thyroid gland and is attached to the larynx. This attachment is responsible for the movement of the enlarged thyroid gland on deglutition.

The thyrohyoid membrane is a fibro-elastic ligament, which arises from the superior margin of the thyroid cartilage, and gets attached to the hyoid bone. It facilitates the upward movement of the larynx during swallowing.

The ligament of Berry, also called the suspensory ligament extends from the cricoid cartilage to the thyroid. Its attachment prevents the thyroid gland from sinking down into the mediastinum.



Solution to Question 8:

The lumbricals and interossei muscles are affected.

The lumbricals and interossei muscles through their attachments to the dorsal digital expansions, cause:

• Flexion of the metacarpophalangeal (MCP) joints

• Extension of the interphalangeal (IP) joints

In cases of ulnar nerve injury, these muscles may get atrophied leading to the extension of MCP joints and flexion of IP joints.

Note: The ulnar nerve innervates all the intrinsic muscles of the hand, except for

• First and second lumbrical

• Abductor pollicis brevis

• Opponens pollicis

• Part of flexor pollicis brevis

Solution to Question 9:

Meiosis occurs in the adult ovary.

Meiosis takes place only in germ cells and results in the formation of haploid gametes. Meiosis requires two cell divisions, meiosis I and meiosis II, to reduce the number of chromosomes to the haploid number of 23.

Option C: Spermatogenesis begins at puberty. Primordial germ cells (PGCs) remain dormant in the testis until puberty. At puberty, PGCs differentiate into spermatogonial stem cells. These

spermatogonial stem cells then give rise to spermatogonia which marks the beginning of spermatogenesis.

Options A and D: These are somatic cells that only undergo mitosis and not meiosis.

Solution to Question 10:

The long left recurrent laryngeal nerve is due to the persistence of the left 6th arch artery.

During the development, the laryngeal nerves initially arise below the 6th arch artery and cross here to innervate the intrinsic muscles of the larynx. As the larynx moves cranially relative to the aortic arch arteries, the recurrent laryngeal nerves on the right and left sides are caught under the 6th arch arteries.

On the right side, the distal 6th arch artery disappears. Since there is no 5th arch artery, the right recurrent laryngeal nerve is caught under the 4th arch artery which becomes the future right subclavian artery.

On the left side, the 6th arch artery persists, which develops into ductus arteriosus, which later forms ligamentum arteriosum. Due to the persistence of the 6th arch artery, the left recurrent laryngeal nerve is caught under it and therefore, has a relatively long course.

Solution to Question 11:

The muscle marked in the above image is levator scapulae, which is innervated by the dorsal scapular nerve.

The appendicular group of back muscles and their innervations include:

Option B: The suprascapular nerve innervates the supraspinatus and infraspinatus muscles.

Option D: The subscapular nerves supply the teres major and subscapularis muscles.

Muscle	Nerve Supply
Trapezius	Accessory nerve
Latissimus dorsi	Thoracodorsal nerve
Levator scapulae	Dorsal scapular nerve
Rhomboid major and minor	Dorsal scapular nerve



Solution to Question 12:

The above clinical scenario is suggestive of lateral pontine syndrome which occurs due to a lesion located at lateral pons.

Lateral pontine syndrome occurs due to the involvement of the anterior inferior cerebellar artery.

• Ipsilateral manifestations include:

• CN VII: Facial paralysis

• CN V: loss of pain and temperature sensation over the face.

• Vestibular nerve or nucleus: Horizontal and vertical nystagmus, vertigo, nausea, vomiting, oscillopsia.

• Auditory nerve or cochlear nucleus: Deafness, tinnitus.

• Middle cerebellar peduncle: Ataxia of limb and gait.

• Contralateral manifestations include:

• Spinothalamic tract: Loss of pain and temperature sensation over the body.

Medial pontine syndrome occurs due to the involvement of the paramedian branch of the basilar artery.

• Ipsilateral manifestation

• CN VI: Diplopia(lateral rectus muscle paralysis - the affected eye looks downwards and inwards)

• Vestibular nucleus: Nystagmus.

• Middle cerebellar peduncle: Ataxia of limb and gait.

• Contralateral manifestation

• Corticobulbar and corticospinal tract: Paralysis of face, arm, and leg.

• Medial lemniscus: loss of touch and proprioception from the body.

Solution to Question 13:

Ligation of the common hepatic artery will cause impairment in the blood supply through its branches i.e right gastric artery and right gastroepiploic artery.

Blood supply of the stomach:

The arterial supply of the stomach is contributed exclusively by the branches of the celiac trunk. They are:

• Left gastric artery - Supplies the lesser curvature of the stomach

• Splenic artery- This further divides into:

• Short gastric arteries- Supplies the fundus of the stomach

• Left gastroepiploic artery- Supplies the greater curvature of the stomach

• Common hepatic artery, which gives the following branches

• Right gastric artery- Supplies the lesser curvature of the stomach

• Gastroduodenal artery- Right gastroepiploic artery arises from this and supplies the greater curvature of the stomach.



Solution to Question 14:

The joint marked in the above image is the costotransverse joint, which is a synovial type of joint.

The various types of joints in the thoracic wall include:

Sternocostal joints: These are the joints between the costal cartilages of the first seven ribs and the sternum.

• The joint between the 1st costal cartilage and manubrium is a fibrocartilaginous joint (synarthrosis).

• The joints between the 2nd to 7th costal cartilages and the sternum are synovial joints.

The interchondral joint between the costal cartilages of the 9th and 10th ribs is never synovial and is usually absent.

Type of Joint	Examples
Synovial joint	Costovertebral jointCostotransverse jointInterchondral joints
Primary cartilaginous joint/Synchondrosis	Costochondral joint
Secondary cartilaginous/Symphyses joint	Manubriosternal and xiphisternal joints

Solution to Question 15:

The sign shown in the above image is the popeye sign, which occurs due to the rupture of the tendon of the long head of the biceps brachii.

On flexing the elbow, the characteristic deformity appears due to the bulging of the muscle belly caused by the contraction of the unrestrained muscles.

Option B: Hill-Sachs sign refers to a defect in the postero-lateral surface of the humeral head due to repeated impact against anterior glenoid rim. It is a common complication of Anterior shoulder dislocation.

Option C: Griesinger sign refers to oedema of the postauricular soft tissues overlying the mastoid process as a result of thrombosis of the mastoid emissary vein. It is a complication of acute

mastoiditis.



Option D: Rising sun sign , also known as Schwartze sign, is the otoscopic sign of active otospongiosis characterised by reddish discolouration of the cochlear promontory.

Solution to Question 16:

The most likely primary cause of an enlarged inguinal lymph node is anal cancer.

Lymphatics from the anal canal distal to the dentate line (anal margin) drains into the inguinal lymph nodes. Hence, anal cancer arising distal to dentate line can cause enlarged inguinal lymph nodes. Lymphatics proximal to the dentate line (anal canal) drains into internal iliac lymph nodes through superior rectal lymphatics.

Option A: In testicular cancer, lymphatic spread occurs to the paraaortic lymph nodes.

Option C: In sigmoid colon cancer, lymphatic spread occurs to the regional lymph node and then to the central lymph node.

Option D: In prostate cancer, lymphatic spread occurs to the:

• Lymph nodes beside the internal iliac vein and in the hollow of the sacrum

• External iliac lymph nodes.

Solution to Question 17:

The above image shows an elastic cartilage.

The elastic cartilage, like the hyaline cartilage, contains individual or small groups of chondrocytes that are surrounded by type II collagen fibrils. However, it differs from hyaline cartilage in that, much of the interterritorial matrix consists of fine yellow elastic

fibres containing elastin protein.

On microscopy, elastic cartilage consists of:

• Chondrocytes within lacunae

• Elastic fibres in the matrix

• Perichondrium with outer fibrous layer and the inner cellular layer

Anatomy INI-CET 2021

Question 1:

A 6-month-old boy was brought to the casualty with seizures. The pediatrician tries to do CSF sampling. What are the structures punctured by the pediatrician while piercing through the marked structure?



a) Scalp, dura, arachnoid

b) Scalp, epicranium, endocranium and dura

c) Scalp, synchondral membrane, dura, arachnoid

d) Pericranium, dura, arachnoid

Question 2:

A 12-year-old presented with fever and difficulty swallowing. He had swelling in the marked region and was advised to undergo surgery. Post-surgery the gauze continued to soak with blood. Which of the following vessels must have been injured?

a) Tonsillar branch of facial artery

b) Ascending pharyngeal artery

c) Paratonsillar vein

d) Retromandibular vein

Question 3:

At which level does the nerve supply for the marked structure arise?

a) Red nucleus

b) Sub thalamic nuclei

c) Decussation of pyramidal tract

d) Olivary nucleus

Question 4:

Which of the following are branches of the anterior division of the internal iliac artery?

a) 1, 2, 3, 5

b) 2, 3, 4, 6

c) 2, 3, 4, 5, 6

d) 1, 4, 5, 6

Question 5:

Fibers from the marked structure terminate at which of the following?



a) Fastigial nucleus

b) Inferior olivary nucleus

c) Red nucleus

d) Subthalamus

Question 6:

A woman with right-sided loss of sensations of both the upper and lower limb complains of shooting pain from her fingers to the right shoulder and a burning sensation when touching cold water. Motor functions are normal. Which of the following structures is likely to be involved?



a) A

b) B

c) C

d) D

Question 7:

A 25-year-old male presented with symptoms of inability to flex the distal interphalangeal joint of the 4th and 5th digits. He was also not able to hold a piece of paper between his fingers. What is the likely site of injury?



a) A

b) B

c) C

d) D

Question 8:

A young child with absent thymus is brought to the OPD with features of hypoparathyroidism and tetany. Which of the following marked area in the picture is defective in this condition?



a) 1

b) 2

c) 3

d) 4

Question 9:

A patient who has taken the first COVID vaccine comes for the second dose. An astute nurse noticed that the shoulder was flabby, flat, and was asymmetrical. There was an associated loss of contour of the shoulder joint. Injury to which of the structures might have resulted and was avoidable?



a) Rotator cuff

b) Posterior circumflex artery

c) Lateral cutaneous nerve of arm

d) Axillary nerve

Question 10:

The Para-sagittal section of the abdomen is given below. What is the superior border of the marked structure?



a) Caudate lobe of liver

b) Duodenum

c) Lesser omentum

d) Inferior vena cava

Question 11:

Through which of the following marked sites does SARS-COVID 19 spread to the brain?



a) D

b) C

c) B

d) A

Question 12:

A boy with a history of a fall from a tree presents with a lurching gait and dragging of his feet. Which of the following structures are involved?



a) 1

b) 2

c) 3

d) 4

Question 13:

A patient after a road traffic accident presents to the emergency room with difficulty in swallowing and slurred speech. Investigations reveal fractures in the occipitotemporal region. Which of the following areas should be tested in order to find the nerve which is involved?



a) 1

b) 2

c) 3

d) 4

Question 14:

Fibers from the marked structure end in which of the following areas of the thalamus?



a) Posterior nucleus

b) Reticular nucleus

c) Anterior nucleus

d) Ventral Lateral nucleus

Question 15:

The cell marked in the below slide is inhibitory to



a) Deep nuclei of cerebellum

b) Vestibular nuclei

c) Inferior olivary nucleus

d) Basal ganglia

Question 16:

The marked structure develops from which of the following structures?



a) Common cardinal vein

b) Anterior cardinal vein

c) Supracardinal vein

d) Subcardinal vein

Question 17:

Inferior thyroid artery supplies which of the following structures?

a) 1, 2, 3, and 4

b) 1, 2, and 3 only

c) 1 and 2 only

d) 1, 2 and 4 only

Question 18:

Which part of this bone is palpable in the infraclavicular fossa?



a) 1

b) 2

c) 3

d) 4

Question 19:

Identify the pelvic diaphragm in the picture given below.



a) 1

b) 2

c) 3

d) 4

Question 20:

Identify the incorrect match.



a) 1- Cilia

b) 4- Lamina propria

c) 2- Goblet cells

d) 3- Nuclei of epithelial cell

Question 21:

Third part of vertebral artery is related to which of the following ?



a) 4

b) 3

c) 2

d) 1

Question 22:

What is the most common complication of the central line insertion below the marked bone?



a) Arterial puncture

b) Thrombophlebitis

c) Empyema

d) Chylothorax

Question 23:

Identify the cranial nerve passing through the given structure



a) III

b) V

c) IV

d) VI

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	c
5	c
6	b
7	c

8	c
9	d
10	a
11	d
12	b
13	b
14	c
15	a
16	c
17	a
18	a
19	a
20	b
21	c
22	a
23	c

Detailed Explanations

Solution to Question 1:

The given image shows the skull of an infant and the arrow points towards the anterior fontanelle. The procedure of collecting CSF from the anterior fontanelle is called ventricular tapping. The various structures pierced are the scalp, dura, and arachnoid.

Fontanelle tap/ Percutaneous ventricular puncture:

A fontanelle tap or percutaneous ventricular puncture is performed in infants without access to a VP shunt, who require urgent ventricular decompression and have an open anterior fontanelle. The infants' open fontanelle is directly punctured with a butterfly needle to enter the lateral ventricle to collect CSF samples or release the pressure or place a drain/catheter.

A 25-gauge butterfly needle is inserted through the anterolateral aspect of the anterior fontanelle (away from the midline, avoiding the superior sagittal sinus).

The image below shows the various fontanelles.



The scalp is a soft tissue that covers the calvaria of the skull. It consists of five layers (Mnemonic: SCALP). Below the scalp and the calvaria lie the meningeal layers, namely, pachymeninges [arachnoid + dura mater (P-AD)] and leptomeninges [arachnoid + pia mater (L-AP)]. One must reach the subarachnoid space below the arachnoid layer to collect or tap CSF. So, the layers pierced in the order are as follows:

• Skin

• Connective tissue (superficial fascia) with blood vessels and nerves

• Aponeurosis (galea aponeurotica) with occipitofrontalis muscle

• Loose areolar (subaponeurotic) tissue

• Pericranium

• Dura mater

• Arachnoid membrane

The given images show the various layers that are pierced.





Solution to Question 2:

The given image is a cadaveric specimen with the arrow pointing towards, the tonsillar fossa, and the given clinical vignette is suggestive of tonsillitis. Post tonsillectomy hemorrhage is commonly venous in origin, from the paratonsillar/ external palatine vein.

Torrential hemorrhage can be due to injury to an aberrantly large or aberrantly placed tonsillar arterial branch of the facial artery.

Blood supply of the tonsils:

Arterial supply:

• The tonsillar branch of the facial artery is the main supply.

• Ascending palatine branch of the facial artery.

• Ascending pharyngeal branch of the external carotid artery.

• Dorsal lingual branches of the lingual artery.

• Descending palatine branches of the maxillary artery.

Venous supply:

• Veins from the tonsils drain into paratonsillar vein which joins the common facial vein and pharyngeal venous plexus.

Types of Hemorrhage seen during/post tonsillectomy:

• Primary hemorrhage:

• Occurs during surgery due to injury to blood vessels.

• Can be controlled by pressure, ligation, or electrocoagulation of the bleeding vessels.

• Reactionary hemorrhage:

• It occurs within 24 hours of surgery.

• It may be due to dislodgement of the clot, slippage of ligature, or return of blood pressure to normal.

• It is managed by removal of clot and application of pressure or vasoconstrictor.

• If not controlled the patient is anesthetized and hemorrhage is managed by re-exploration.

• Secondary hemorrhage:

• It occurs between 24 hours to 10 days (most common between 5th to 8th postoperative days).

• It is due to sepsis and premature separation of the membrane.

• The patient is started on IV antibiotics and if not controlled, general anesthesia is given and the bleeding vessel is electrocoagulated or ligated.

The images below summarize the blood supply to the tonsils and the relations of the tonsillar bed respectively.





Solution to Question 3:

The muscle marked in the given CT image is that of the medial rectus. The medial rectus is supplied by the oculomotor nerve, the nucleus of which is situated at the level of the superior colliculus close to the red nucleus.

The nuclear complex from which the oculomotor nerve arises consists of:

• Several groups of large motor neurons are collectively called the oculomotor nucleus. The motor neuron groups of the oculomotor nucleus innervate, in dorsoventral order, the ipsilateral inferior rectus, inferior oblique, and medial rectus. A medially placed column of cells innervates the contralateral superior rectus.

• Smaller preganglionic parasympathetic neurons called the Edinger–Westphal preganglionic nucleus

Red Nucleus/ Nucleus Ruber

It is a rounded mass of grey matter located dorsomedial to the substantial nigra. It has a reddish hue in fresh specimens due to its extensive vascularity and the presence of ferric iron pigments in multipolar neurons.

Clinical correlate:

Injury to the red nucleus can cause Benedikt's syndrome characterized by ipsilateral oculomotor nerve palsy and contralateral tremors, chorea, and athetosis.

The given image shows the midbrain section at the level of superior colliculus. We can appreciate the two nuclei being close to each other.

Afferent connections	Efferent connections
The cerebral cortex by the co rticospinal fibers	The spinal cord by the rubros pinal tract

Afferent connections	Efferent connections
The cerebellum through the s uperior cerebellar peduncle	The reticular formation throu gh the rubroreticular tract
The lentiform nucleus, hypot halamic and subthalamic nuc lei, substantial nigra, and spi nal cord.	The substantia nigra and the thalamus



The image attached below summarizes the course of the oculomotor nerve.



Solution to Question 4:

The branches from the anterior division include 2, 3, 4, 5, 6.

The internal iliac artery is the major artery of the pelvis. At the superior border of the greater sciatic foramen in the pelvis, it divides into anterior and posterior trunks.



Branches of the anterior trunk of the internal iliac artery:

- Obturator artery
- Superior vesical artery
- Inferior vesical artery
- Inferior gluteal artery
- Middle rectal artery
- Vaginal artery (in females)
- Uterine artery (in females)
- Internal pudendal artery

Branches of the posterior trunk of the internal iliac artery:

- Iliolumbar artery
- Lateral sacral arteries
- Superior gluteal artery



Solution to Question 5:

The peduncle highlighted in the image is the superior cerebellar peduncle. The superior cerebellar peduncle includes fibers from deep cerebellar nuclei that project to the brainstem, red nucleus, and thalamus. It also contains the ventral spinocerebellar tract fibers.

The superior cerebellar peduncle contains all the efferent fibers from the dentate, emboliform, and the globose nuclei, and a small fascicle from the fastigial nucleus. Its fibers decussate in the caudal mesencephalon and are destined to synapse in the contralateral red nucleus and thalamus. The middle cerebellar peduncle contains only afferent fibers from the contralateral pontine nuclei. The inferior cerebellar peduncle is a mixture of afferent fibers from the brainstem and spinal cord and efferent fibers to the vestibular nuclei.

The image below summarizes the various tracts in the cerebellar unit.



The cerebellum consists of two hemispheres (right and left) joined by a median vermis. It is divided into three lobes:-

• Anterior lobe - lies superiorly above the primary fissure

• Middle lobe - lies inferiorly below the primary fissure. It is the largest lobe. It is also called the posterior lobe.

• Flocculonodular lobe - lies posterior to the uvulonodular fissure

It has two surfaces - superior and inferior separated by a horizontal fissure. It is connected to the other parts of the central nervous system via three peduncles

• Superior cerebellar peduncle -connecting to the midbrain

• Middle cerebellar peduncle - connecting to the pons

• Inferior cerebellar peduncle - connecting to the medulla oblongata

Solution to Question 6:

The given clinical stem of a hemi-sensory loss and severe excruciating pain of the affected areas is suggestive of thalamic pain syndrome also known as Dejerine Roussy syndrome. Hence, thalamus marked as B is most likely to be involved.

Occlusion of these thalamic and thalamogeniculate branches results in the thalamic pain syndrome. It is characterized by C/L hemisensory loss which starts as numbness and progresses to cause severe burning pain in the affected regions. As the thalamus is the site for the relay of sensory information between the periphery and cortex, a lesion in the thalamus causes a miscommunication between the afferent pathway and cortex of the brain. Pain occurs during the recovery from the thalamic infarct. This pain responds poorly to analgesics. Anticonvulsants such as carbamazepine or gabapentin or tricyclic antidepressants may be beneficial.

Other markings:

A: Caudate nucleus

C: Substantia nigra

D: Pons

The image below shows the coronal section of the brain with different parts of the basal ganglia:





Solution to Question 7:

The given clinical situation with a positive card test and an inability to flex the distal interphalangeal joint of the 4th and 5th digits, suggests an ulnar nerve injury. The most likely site of injury would thus be medial epicondyle: site C

Injury at the medial epicondyle damages the ulnar nerve, leading to paralysis of flexor carpi ulnaris (loss of ulnar deviation) and medial half of flexor digitorum profundus (loss of distal interphalangeal joint flexion). It leads to hypothenar paralysis (lead to atrophy) and sensory loss. Also, flexor digitorum superficialis (supplied by median nerve) and extensor digitorum (supplied by the radial nerve) are still functional. Hence complete paralysis of the 3rd and 4th digits is not seen. The positive card test is due to the weakness of palmar interossei leading to loss of adduction.

Ulnar nerve branches and supply:

Superficial branch supplies palmaris brevis, skin on the palmar surface of medial 1½ fingers. The deep branch runs within the concavity of deep palmar arch and supplies all intrinsic muscles of the hand (except first 2 lumbricals and thenar muscles).

Injury to the Ulnar nerve:

Other tests seen in ulnar nerve lesions are as follows:

• Claw hand deformity

• Froment's sign positive

• Wartenberg sign

• Palmaris brevis sign

	High lesion(Elbow)	Low lesion(Wrist)
Site of injury	Fracture/dislocation of theme dial epicondyleCubital tunnel syndromeCompression betw een the two heads of FCUTar dy ulnar nerve palsy (Valgus deformity)	Wrist laceration/ slashingGuy on's canalsyndrome
Motor loss	Claw-handdeformityHypothe nar & interosseous wastingLo ss of:• Abduction & adduction of fingers(Card test- Positive) • Adduction of thumb(Frome nt sign-Positive)• Flexion of MCP jointsWeakness of wrist flexion (radial deviation)	Claw-handdeformity(more pr onounced)Hypothenar & inte rosseous wastingLoss of:• Ab duction & adduction of finger s(Card test- Positive)• Adduct ion of thumb(Froment sign-P ositive)
Sensory loss	Over palmar & dorsal aspect of themedial thirdof thehand Over palmar & dorsal aspect of themedial 11/2fingers	No sensory lossover palmar & dorsal aspect of the medial third of the handOver palmar & dorsal aspect of themedial 11/2fingers

The image below shows a positive card test.



Positive Card Test
In the weakness of palmar interossei, there is a weakness of finger adduction. So the patient is unable to hold card firmly between fingers.

Wartenberg's sign: (Weakness of palmar interossei), Sensory loss Palmar and dorsal surface of the medial third of hand and medial 1 and a half fingers.



Other options:

• Option A: Radial groove - Radial nerve

• Option D: Neck of humerus - Axillary nerve

• Option B: Lateral epicondyle – May lead to tardy ulnar nerve palsy after fracture but the incidence is less.

Solution to Question 8:

The image shows pharyngeal pouches and based on the clinical history, the patient most likely has DiGeorge syndrome(thymic and parathyroid defects). It may be caused by defective development of the third and fourth pharyngeal pouch. In the given stem, it is clearly mentioned that the thymus is absent. Thus, the better of the two options is 3.

DiGeorge syndrome is a complex embryopathy that is caused by hemizygous interstitial deletion of chromosome 22, leading to multiple developmental defects, including conotruncal defects, hypoparathyroidism, and dysmorphic syndrome. Microdeletion at chromosome 22q11 can lead to variable presentations including

• Cleft palate

• Abnormal facies

• Thymic aplasia leading to T-cell deficiency

• Cardiac defects

• Hypocalcemia 2° to parathyroid aplasia. (Mnemonic: CATCH-22)

Velocardiofacial syndrome - palate, facial, and cardiac defects.



In the diagram, I-IV are the branchial arches, 1-4 are pharyngeal pouches (inside), and/or pharyngeal grooves (outside)

• Tuberculum lateral

• Tuberculum impar

• Foramen cecum

• Ductus thyroglossus

• Sinus cervicalis

• First pharyngeal pouch, middle ear cavity, and epithelial lining of the auditory tube.

• Second pharyngeal pouch, palatine tonsils

• Points towards third pharyngeal pouch which gives rise to inferior parathyroid gland and thymus.

• Fourth pharyngeal pouch, Superior parathyroid gland, and ultimobranchial body.

Solution to Question 9:

The given clinical vignette is suggestive of injury to axillary nerve which leads to deltoid muscle palsy.

The axillary nerve supplies the deltoid muscle. Iatrogenic injuries of the axillary nerve can occur due to incorrectly administered IM injection or while performing transaxillary operations on the shoulder with lateral deltoid-splitting incisions. But, note that this nerve is commonly injured during shoulder dislocation (most commonly anterior type) or fractures of the humeral neck as both the nerve and the posterior circumflex vessels (Option-B) pass horizontally and posterior to the neck of the humerus.

Injury to the axillary nerve can lead to:

• Weakness of abduction and lateral rotation of the arm

• Retropulsion (extension of the shoulder with the arm abducted to 90 degrees) is impossible.

• Atrophy of the Deltoid muscle leading to the loss of muscle contours of the shoulder

• Loss of sensation in the Regimental badge area

Note:

Upper (superior) lateral cutaneous nerve of the arm (Option-C) that innervates the skin over the lower deltoid (regimental badge area). Thus, this is not the best of the given four options.

The images below show the Seargent/regimental badge sign and the various anatomical structures that can be appreciated in the transverse section of the shoulder joint respectively.



Vaccine administration in the deltoid can sometimes injure the axillary and the radial nerve.



Solution to Question 10:

The structure marked in the given image is epiploic foramen. The superior border of the epiploic foramen (foramen of Winslow) is formed by the caudate lobe of the liver.

Epiploic foramen is a vertical slit of about 3 cm behind the free right border of the lesser omentum. It connects the lesser sac to the greater sac. The pearl summarizes the various relations of the foramen.

The image attached shows the various peritoneal reflections.



Clinical note: Pringle maneuver (as shown below) is a technique where the free margin of the lesser omentum is compressed to obtain rapid control of the hepatic artery and portal vein in liver trauma and surgery.



Solution to Question 11:

The given image shows various skull foramina's. Infection through the cribriform plate and olfactory bulb (A) and dissemination through trans-synaptic transfer are some of the mechanisms proposed for the SARS-COV2 virus to reach the brain.

Encephalitis, demyelination, neuropathy, and stroke are the neurological sequelae to severe COVID-19 infection. The neurological invasion of the virus on the medullary cardiorespiratory center can lead to refractory respiratory failure in patients infected with COVID-19.

The pathogenesis implicated is explained in the image attached below.



Solution to Question 12:

The structure involved in causing lurching gait in the child is gluteus medius (marked as 2).

The gluteus medius originates from the outer surface of the ilium between the anterior and posterior gluteal lines. Insertion of the muscle is at the lateral surface of the greater trochanter. It is supplied by the superior gluteal nerve.

Gluteus medius and minimus are powerful abductors of the hip joint. Mechanism of actions of gluteus medius and minimus:

• When the leg on one side is lifted off the ground, the gluteus medius and minimus muscles, on the opposite supported side, exert traction on the hip bone by acting from the femur. This action results in a slight raising of the pelvis on the unsupported side, which helps maintain the trunk in the upright position.

• The actions of these muscles are affected when there is an injury to the superior gluteal nerve, neck of the femur, or loss of normal angulation of the shaft of the femur. This occurs in conditions like paralysis of the glutei, developmental dysplasia of the hip, etc.

• When the gluteus medius and minimus are paralyzed, the pelvis sinks down on the unsupported (normal) side, when the patient tries to stand on the affected limb. This leads to a positive Trendelenburg's sign with a characteristic lurching gait - the patient lurches towards the affected side with contralateral pelvis drop.



Other options

1 - Gluteus maximus - involved in the extension of flexed thigh, abduction (only upper fibres), and lateral rotation at the hip joint. Insertion is at the iliotibial tract.

3 - Piriformis - involved in lateral rotation and assists abduction at the hip joint. It is inserted into the medial side of greater trochanter of the femur. Entrapment of the sciatic nerve in the piriformis can lead to piriformis syndrome.

4 - Sciatic nerve - The nerve passes over the obturator internus, gemellus superior, gemellus inferior and quadratus femoris. It supplies knee flexors and muscles below the knee. Paralysis of this nerve results in flail foot and severe difficulty in walking.



Solution to Question 13:

The given clinical scenario is suggestive of the involvement of 9th and 10th cranial nerves due to occipitotemporal fracture and they are examined by testing the uvula which is marked as 2 in the given image.

The jugular foramen is located between the lower border of the petrous part of the temporal bone and the condylar part of the occipital bone. As the investigations reveal fractures in the occipitotemporal region, along with the history of difficulty in swallowing and slurred speech, are suggestive of injury to the glossopharyngeal and vagus nerve passing through the jugular foramen.

Unilateral lesions of IX and X cranial nerves are commonly caused by skull-base fractures, strokes, or tumors.

Glossopharyngeal and Vagus nerve examination include:

• Observing the position and symmetry of the palate and uvula(marked as 2 in the given image) at rest and with phonation (“aah”).

• The pharyngeal (“gag”) reflex is evaluated by stimulating the posterior pharyngeal wall on each side with a sterile, blunt object (e.g., tongue blade)

Patients with a proximal unilateral lesion of the vagus nerve complain of dysphagia. There will be a weakness of the soft palate muscles on the affected side. Attempts to voluntarily elevate the soft palate, like by asking the patient to say “aah” will reveal a weakness in elevation of the palate on the affected side. Along with this, there will be deviation of the uvula to the normal side, due to the unopposed action of palatal muscles on that side.

The below image shows the base of the skull and its foramina:



Solution to Question 14:

The structure marked in the above image is mamillary body and the fibers from the mamillary body end in the anterior nucleus of the thalamus.

The mammillary bodies are two small hemispherical, pea-sized eminences, lying side by side, forming the floor of the third ventricle. The fibres from medial mammillary nucleus project to the

anterior nucleus of the thalamus.

The anterior thalamic nuclei have 3 subdivisions—anteroventral nucleus, anteromedial nucleus, and anterodorsal nucleus.

• Major input: Mamillothalamic tract from the mammillary nuclei of the hypothalamus.

• Major cortical output: Anterior limbic area, cingulate, and the parahippocampal gyrus

• Major function:

• Regulation of alertness and attention

• Acquisition of memory

Solution to Question 15:

The structure marked in the image is a flask-shaped neuron in the cerebellar cortex, which is a Purkinje cell. It is inhibitory to deep cerebellar nuclei.

Purkinje cells are the flask-shaped cells present in the Purkinje cell layer of the cerebellum. They are evenly spaced and arranged in a single layer between molecular and granular cell layers. The dendrite from each cell passes upward to enter the molecular layer and the axon of each cell passes downwards, crosses the granular cell layer, and terminates on the deep cerebellar nuclei. They are inhibitory and use GABA as their neurotransmitter.

Axons of Purkinje cells are the only efferents of the cerebellar cortex. They receive excitatory input from the climbing fibers from the inferior olivary nucleus and mossy fibers.

Pathological significance of Purkinje cells:

• Negri bodies in rabies are usually found as eosinophilic, cytoplasmic inclusions in the neurons of the hippocampus and Purkinje cells of the cerebellum.

• In paraneoplastic cerebellar degeneration, there is the destruction of Purkinje cells due to antibodies directed against them.

• Purkinje cells and neurons in pyramidal cell layers of the hippocampus and cerebral cortex are the most sensitive neurons to hypoxia and are likely to be damaged in cases of global cerebral ischemia.

Solution to Question 16:

The structure marked in the given image is the azygos vein, which develops from the supracardinal vein.

The azygos system of veins forms a connecting channel between the superior and inferior venae cavae. It occupies the upper part of the posterior abdominal wall and the posterior mediastinum. It also connects the portal, caval system, and vertebral venous systems.It drains the thoracic wall and the upper lumbar region.

Development: The azygos venous system arises from the supra cardinal veins embryologically. The right supra cardinal vein becomes the azygos vein, while the left supra cardinal vein becomes

the hemiazygos vein.

• The 4th to 11th right intercostal veins empty into the right supracardinal vein, which together with a portion of the posterior cardinal vein forms the azygos vein.

• On the left, the 4th to 7th intercostal veins enter into the left supracardinal vein, and the left supracardinal vein, then known as the hemiazygos vein, which also empties into the azygos vein.

Other options: Cardinal veins form the main venous drainage system of the embryo. It consists of,

• Anterior cardinal veins, which drain the cephalic part of the embryo, and

• Posterior cardinal veins, which drain the rest of the embryo.

During the fifth to the seventh weeks, a number of additional veins are formed. It includes,

• Subcardinal veins, which mainly drain the kidneys;

• Sacrocardinal veins, which drain the lower extremities;

• Supracardinal veins, which drain the body wall by way of the intercostal veins.

Below image shows cardinal veins development.



Solution to Question 17:

All the given 4 structures are supplied by the Inferior thyroid artery.

Inferior thyroid artery is a branch of the thyrocervical trunk, which in turn, is a branch of the first part of the subclavian artery. The inferior thyroid artery loops upwards anterior to the medial border of the scalenus anterior, turns medially just below the sixth cervical transverse process, and then descends on longus colli to the lower border of the thyroid gland.

Branches:

• Pharyngeal branches: These supply the lower part of the pharynx.

• Tracheal branches supply the trachea and anastomose with the bronchial arteries;

• Oesophageal branches supply the oesophagus and anastomose with the oesophageal branches of the thoracic aorta; and

• Inferior and ascending glandular branches supply the posterior and inferior regions of the thyroid gland, and anastomose with the contralateral inferior and ipsilateral superior thyroid arteries.

• The ascending branch also supplies the parathyroid glands.

• Ascending cervical artery

• Inferior laryngeal

• Muscular branches

Note: Thymic branches originating from the internal thoracic, inferior thyroid artery, and sometimes superior thyroid arteries supply the thymus and the surrounding mediastinal connective tissue.

Solution to Question 18:

The part of the scapula marked as 1 is the coracoid process and it is palpable in the infraclavicular fossa. The tip of the coracoid process lies approximately 2.5 cm below the clavicle, immediately lateral to the infraclavicular fossa (deltopectoral triangle). Hence, the coracoid process can be palpated below the clavicle.

The coracoid process arises from the upper border of the neck of the scapula and it projects forwards and slightly laterally. Attachments at the coracoid process include

• The conoid portion of the coracoclavicular ligament

• Pectoralis minor

• Medial end of coraco-acromial ligament

• Coracohumeral ligament

• Glenocorocoid ligament

• Coracobrachialis muscle

Other options

Option B - Acromian process continues from the spine of the scapula. The acromion is subcutaneous over its dorsal surface, The attachments at the acromion are the coracoacromial ligament, the articular capsule of the acromioclavicular joint, and horizontal fibers of the trapezius.

Option C - The glenoid fossa is present at the lateral angle of the scapula. It is articular with the humerus to form the shoulder joint.

Option D - The inferior angle lies over the seventh rib. Dorsally, it is related to latissimus dorsi. The angle is visible when the arm is raised over the head.

Parts of the scapula are given in the image below.



Solution to Question 19:

The structure marked "1" is the levator ani muscle which is part of the pelvic diaphragm.

Other structures marked in the above image are

2: External anal sphincter

3: Internal anal sphincter

4: Pudendal canal

Pelvic diaphragm

The muscular part of the pelvic floor is the pelvic diaphragm. It is attached to the pelvic wall on either side between the greater and lesser sciatic foramen. Levator ani and coccygeus muscles form the pelvic diaphragm and delineate the lower limit of the true pelvis.

Levator ani is formed by pubococcygeus, iliococcygeus, and puborectalis. The muscles blend together posterior to the vagina in women and around the anus in both sexes. Posterior to the anal aperture, the form the anococcygeal body attaches to the coccyx. Urogenital hiatus is present anteriorly separating the muscles. This allows the urethra and vagina to pass through. Levator ani reinforces the external anal sphincter. In women, it functions as the vaginal sphincter.

The coccygeus muscles, one on each side are triangular in shape and form the posterior part of the pelvic diaphragm. The apices are attached to the tips of ischial spines. The bases are attached to the coccyx and sacrum.

Solution to Question 20:

Among the given options, Option B represents an incorrect match, as the structure labeled as 4 is showing the basement membrane and not lamina propria.

The given image is of pseudostratified ciliated columnar epithelium, as it shows irregular cells, which are attached to the basement membrane(marked as 4 in the image). It appears to be multi-layered. However, there is actually only one layer of cells. The multi-layered appearance is due to the fact that the nuclei of epithelial cell(marked as 3 in the image) lie at different levels.

The epithelium shows hair-like projections called cilia (marked as 1 in the image) and goblet cells (marked as 2 in the image). Goblet cells secrete mucous which helps to trap dust entering the passages. The cilia are capable of movement and help in the clearance of mucous.

Example include epithelium lining the trachea and in large bronchi.



Solution to Question 21:

The third part of the vertebral artery (V3) is related to the area marked 2, which is the posterior arch of the first cervical vertebra or the atlas.

The given image shows the first cervical vertebra - atlas.

• The area marked 1 is the posterior tubercle.

• The area marked 2 is the posterior arch. The superior surface bears a groove for the of vertebral artery and venous plexus immediately behind.

• The area marked 3 is the superior articular facet.

• The area marked 4 is the posterior surface of the anterior arch. It gives attachment to the apex of dens from the second cervical vertebra.

The vertebral artery arises from the first part of the subclavian artery. It passes through the foramina of transverse processes of C6 to C1 cervical vertebrae and enters the cranium through the foramen magnum. The vertebral arteries of both sides join to form the basilar artery. In its course, it is divided into four parts:

The first part, V1, begins at its origin from the subclavian artery and ends at its entry into the transverse process of the C6 cervical vertebra. This part is crossed by the inferior thyroid artery on

both sides, and thoracic duct on the left side, and the right lymphatic duct on the right side.

The second part, V2 is that part of the artery ascending through the foramina in the transverse process of the cervical vertebrae.

The third part, V3 is curved backward behind the atlas. In this position, it lies in a groove on the posterior arch of the atlas. This part of the artery lies in the suboccipital triangle.

The fourth part, V4 pierces the dura and arachnoid mater and unites with the contralateral vertebral artery to form the midline basilar artery at the lower border of pons.



Solution to Question 22:

The bone marked in the above image is the left clavicle. The most common complication of a central line insertion below this bone is an arterial puncture.

Complications of central vein catheterization are

• Arterial puncture- this is the most common complication.

• Cardiac tamponade is the most important life-threatening complication. It occurs as a result of puncture of intrapericardial superior vena cava, right atrium, or right ventricle.

• Pneumothorax- injury to the underlying pleura may lead to pneumothorax. This is more common in subclavian vein catheterization.

• Chylothorax can occur particularly in left subclavian vein catheterisation. But it is not the most common.

• Injury to the associated nerves like brachial plexus, stellate ganglion, phrenic nerve, or vocal cords.

• Thromboembolism leading to venous thrombosis, pulmonary embolism is more common with femoral veins.

• Infections

Central venous catheterization is the insertion of the catheter into a large central vein. The veins preferred largely depend on the underlying disease in the patient, the clinical setting, and the expertise of the doctor.

The internal jugular vein is preferred when the patient has:

• Bleeding diathesis- Bleeding from this vein can be easily detected and can be controlled with pressure.

• Pneumothorax- The lesser risk of injury to the lung

• Needs cardiac pacing- right internal jugular vein is used.

Femoral or subclavian veins can be used in patients when the neck is immobilized using a cervical collar.

The below image shows central vein catheter



The below image shows subclavian vein catheterization.



The below image shows catheterisation into jugular vein



Below image shows radiograph of subclavian vein catherisation



Solution to Question 23:

In the given image, the dorsal aspect of the brainstem is marked and the cranial nerve that is passing there is the trochlear nerve (CN IV).

The trochlear nerve is the fourth cranial nerve that originates from the trochlear nuclei within the midbrain at the level of the inferior colliculus. It is the only cranial nerve that emerges from the dorsal surface of the brainstem and has the longest intracranial course. It passes on to the lateral surface of the crus of cerebellar peduncle and lateral wall of the cavernous sinus. The nerve then crosses the oculomotor nerve to enter the orbit through the superior orbital fissure to exclusively

supply the contralateral superior oblique muscle.

The main actions of superior oblique are

• Depression in adducted position- The muscle elevates the posterior of the eye causing the front of the eye to be depressed. Patients with trochlear nerve palsy have difficulty in looking downwards when eyes are in the adducted position.

• Abduction

• Intortion of the eyeball.

Other options- Oculomotor (CN III), trigeminal (CN V) and abducens (CN VI) nerves exit ventrally from the brainstem.



Anatomy NEET 2021

Question 1:

A patient presented with vision loss. On radiological investigation, an aneurysm causing damage to the optic chiasma was noted. Which of the following arteries is most likely to be the artery that is causing the damage?

a) Anterior communicating artery

b) Anterior choroidal artery

c) Middle cerebral artery

d) Anterior cerebral artery

Question 2:

A 34-year-old man presented with fever and massive splenomegaly. On palpation, the spleen was 15 cm and directed obliquely when measured from the right iliac fossa. Which of the following structures prevent the vertical and downward descent of the spleen?

a) Lienorenal ligament

b) Gastrosplenic ligament

c) Phrenicocolic ligament

d) Lienophrenic ligament

Question 3:

Identify the papillae from the given histological section.



a) Fungiform

b) Circumvallate

c) Filiform

d) Foliate

Question 4:

A 7-year-old boy was brought to the hospital with multiple fractures of the humerus secondary to a fall from height. On examination, there is difficulty in flexion of the elbow and supination of the forearm, and associated loss of sensation over the lateral aspect of the forearm. Which is the nerve most likely to be injured?

a) Median nerve

b) Radial nerve

c) Musculocutaneous nerve

d) Ulnar nerve

Question 5:

A patient presented with jaw swelling and weight loss. Suspecting a malignancy, the submandibular gland was resected. Which is the most likely nerve to be damaged during submandibular gland resection?

a) Nerve to mylohyoid

b) Inferior alveolar nerve

c) Lingual nerve

d) Hypoglossal nerve

Question 6:

A patient underwent a coronary bypass graft with the great saphenous vein. The patient now complains of paresthesia and sensory loss over the medial aspect of the leg and foot. Which of the following nerves is most likely to be involved?

a) Superficial peroneal nerve

b) Saphenous nerve

c) Femoral nerve

d) Sural nerve

Question 7:

During the clinical examination of the nervous system, which of the following nerves can be tested by touching the highlighted structure?



a) IX

b) X

c) XI

d) XII

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	c
5	c
6	b
7	a

Detailed Explanations

Solution to Question 1:

An aneurysm of the anterior communicating artery is most likely to be causing damage to the optic chiasma, leading to vision loss in this patient.

Circle of Willis is a large arterial anastomosis that unites the internal carotid and vertebrobasilar systems. It lies in the subarachnoid space within the basal cisterns that surround the optic chiasma and infundibulum.

Anterior communicating artery aneurysm:

Anterior communicating artery (Acom) links the right and left anterior cerebral arteries and is directed anterior and superior to the optic chiasma. The aneurysm will protrude in the posterior direction and will impinge upon the optic chiasma from above, leading to loss of vision.

Compression of the optic chiasma (central lesion) by an ACom aneurysm, will lead to bitemporal hemianopia i.e, loss of vision in the left/temporal halves of both visual fields. Nasal fibers of each retina cross through the chiasma at this level and hence, these fibers will be compromised.

In the case of a bilateral peripheral lesion of the optic chiasma, binasal hemianopia will be seen.

Other causes of chiasmal lesions are

• Pituitary adenoma (most common)

• Craniopharyngioma

• Meningioma

• Glioma

Blood supply of optic chiasma: The left and right anterior cerebral arteries, along with the superior hypophyseal artery (branch of the internal carotid artery) form the blood supply of optic

chiasma.



The given cadaveric image below is of the vascular structures (Circle of Willis) at the base of the brain.



Investigation: Digital subtraction angiogram is the definitive imaging modality for aneurysms arising from the circle of Willis. Other investigations are CT and MRI.

Other options:

Aneurysms of the anterior choroidal artery (option B) and middle cerebral artery (option C) aneurysms will not compress the optic chiasma as they are anatomically not in close proximity to be able to exert pressure on the chiasma and cause loss of vision.

Despite the anatomical proximity of the anterior cerebral artery (option D) to the optic chiasma, anterior communicating artery (Acom) aneurysm is the commonest intracranial aneurysm.

Solution to Question 2:

The vertical and downward descent of the spleen is prevented by the phrenicocolic ligament.

The phrenicocolic ligament is not attached to the spleen. It attaches the splenic flexure of the colon to the diaphragm.

When the spleen gets enlarged, the phrenicocolic ligament along with the splenic flexure of the colon restricts its downward displacement. The enlargement of the spleen occurs towards the right iliac fossa and hence the spleen is directed obliquely.

Splenic ligaments and their attachments:

Splenic ligament	Attachments	Contents
Phrenicocolic ligament	Diaphragm to the splenic flexure of the colon	-
Phrenicosplenic ligament	The undersurface of the diaphragm to the spleen	-
Splenorenal (lienorenal) ligament	Hilum of the spleen to the anterior surface of the left kidney	Tail of pancreasSplenic artery and vein
Gastrosplenic ligament	The greater curvature of the stomach to the upper pole of the spleen	Short gastric vesselsLeft gastroepiploic artery



Cross-sectional view of the spleen and its ligaments



©MARROW

The spleen becomes palpable when it is enlarged 2-3 times more than its normal size. A characteristic splenic notch can be felt in the medial border in an enlarged spleen. Also, fingers cannot be insinuated between the left costal margin and the enlarged spleen.

In the classical method of palpation, the right hand (with fingers pointed towards the left costal margin) is placed on the right iliac fossa. Palpation begins from the right iliac fossa towards the left hypochondrium in an upwards and lateral direction. Between each breath of the patient, the edge of the spleen can be felt.

In the hooking method, the hooked fingers of the right hand are placed under the left costal margin, and with the left hand, pressure is exerted over the posterolateral aspect of the lower thorax. Spleen is felt at the deep inspiratory phase.

Solution to Question 3:

The given histological image is a section of the tongue showing circumvallate papillae, the largest papillae of the tongue.

Circumvallate papillae are large cylindrical structures that are around 8-12 in number, covered by non-keratinized stratified squamous epithelium.

They are arranged in a V-shaped row immediately in front of the sulcus terminalis (not on the sides of the tongue) on the dorsal surface of the tongue. It is narrower at its base than its apex.

Numerous taste buds (around 250 taste buds for each of the 8-12 papillae) are present within the papillary walls and they contain chemosensory cells that synapse with the gustatory nerves. Serous glands of von Ebner empty via the ducts into the base of the sulcus.

Identification: They can be identified by their characteristic dome-shaped structure lined by stratified squamous epithelium.

Other options:

Option A - Fungiform papillae are large, rounded, and red in color and are mainly present on the lingual margins and also on the dorsal surface of the tongue. They are covered by non-keratinized stratified squamous epithelium. It is supplied by the lingual nerve.

Option C- Filiform papillae are minute cylindrical or conical projections that are mainly present on the dorsal part of the anterior 2/3rd of the tongue. All except filiform papillae contain taste buds. They are covered by keratinized stratified squamous epithelium. It is supplied by the lingual nerve.

Option D - Foliate papillae are red, leaf-like mucosal ridges that are present bilaterally in two zones at the sides of the tongue near the sulcus terminalis. They are covered by non-keratinized stratified squamous epithelium.



Solution to Question 4:

The given clinical vignette in a patient with difficulty in flexion of the elbow and supination of the forearm, with associated loss of sensation over the lateral aspect of the forearm, is suggestive of injury to the musculocutaneous nerve.

Fall from a height on an outstretched hand is one of the main causes of humerus fractures.

Course:

The musculocutaneous nerve (C5, C6, C7) is a branch of the lateral cord of the brachial plexus. It emerges at the lower border of the pectoralis minor and pierces coracobrachialis. It then descends between biceps brachii and brachialis where it supplies a branch to the shaft of the humerus. Then it enters the elbow and continues as the lateral cutaneous nerve of the forearm.

It innervates all the muscles of the anterior compartment of the arm i.e.,

• Biceps brachii - involved in flexion of elbow and supination of the forearm

• Brachialis - involved in the flexion of the elbow

• Coracobrachialis - flexes the arm at the glenohumeral joint

Cutaneous innervation of the lateral aspect of the forearm is by a branch of the musculocutaneous nerve, namely, the lateral cutaneous nerve of the forearm (C5, C6).

Hence, injury to the musculocutaneous nerve results in weakness of supination of forearm (biceps brachii), weakness in flexion of the elbow (biceps brachii, brachialis), and loss of sensation over the lateral aspect of the forearm (lateral cutaneous nerve of the forearm).

Muscles of the anterior compartment of the arm:

The below image shows the anterior compartment of the arm

Muscles	Origin	Insertion	Innervation
Coracobrachialis	Coracoid Process of scapula	The medial side of the midshaft of the humerus	Musculocutaneous nerve
Brachialis	Distal half of anterior aspect of the humerus	Ulnar tuberosity	Musculocutaneous and radial nerve
Biceps Brachii	Short head: Coracoid process Long head: Supraglenoid tubercle	Radial tuberosity	Musculocutaneous nerve



Other options:

Option A:

Median nerve injury is characterized by sensory loss over the radial 3 1/2 fingers and loss of thumb abduction. In high lesions, the long flexors of the thumb, index, and middle fingers, the radial wrist flexors, and the forearm pronator muscles are also paralyzed. Wasting of the thenar eminence is seen in long-standing cases.

Option B:

Radial nerve injury can present with wrist drop, thumb drop, and finger drop. Extension of the elbow will be spared in lesions below the elbow. The extension of the IP joints is intact due to the action of the intrinsic muscles of the hand. Loss of sensation over the first dorsal web space is observed.

Option D:

Ulnar nerve injury is characterized by ulnar claw hand, wasting of the hypothenar muscles and intrinsic muscles of the hand, and sensory loss over palmar & dorsal aspect of the medial third of the hand and medial 1 1/2 fingers.





Solution to Question 5:

The most likely nerve to be damaged during submandibular gland resection is the lingual nerve.

It loops around the submandibular duct and hence is susceptible to injury during submandibular gland resection.

Course:

The lingual nerve arises from the posterior trunk of the mandibular nerve (V3 branch of the trigeminal nerve). The lingual nerve descends between the tensor veli palatini muscle and the lateral pterygoid muscle, where it is joined by the chorda tympani nerve. It then descends across the medial pterygoid muscle to enter the oral cavity. As the lingual nerve enters the floor of the oral cavity, it lies on the deep portion of the submandibular gland. It then passes into the tongue on the hyoglossus muscle where it is attached to the submandibular ganglion.

The submandibular duct lies between the lingual and hypoglossal nerves, superficial to the hyoglossus muscle. As the duct reaches the anterior border of the hyoglossus, it is crossed by the lingual nerve and lies in close proximity to the nerve. Hence lingual nerve injury is the most common.

Injury:

The lingual nerve is at risk to be injured during submandibular gland resection because dissection of the submandibular duct from the nerve is required. Also, it is connected to the submandibular ganglion which will lead to the nerve being exposed within the operating field.

The manifestations of lingual nerve injury depend on whether the lesion is proximal or distal to where it is joined by the chorda tympani nerve.

• Proximal injury: There is loss of sensation from the anterior two-thirds of the tongue, gingivae, oral mucosa, lower lip, and chin.

• Distal injury: In addition to features of proximal injury, secretion from the salivary glands below the oral fissure and taste from the anterior two-thirds of the tongue is also lost.

Note:

The cranial nerves that are at risk during submandibular gland resection include:

• The marginal mandibular branch of the facial nerve

• Lingual nerve

• Hypoglossal nerve







Other options:

Option A: Nerve to mylohyoid is a branch of the inferior alveolar nerve and lies in the mylohyoid groove inferior to the mandibular foramen, hence the risk of injury is less.

Option B: Inferior alveolar nerve enters the mandibular canal via the mandibular foramen, hence is unlikely to be injured during submandibular gland excision.

Option D: Hypoglossal nerve injury is less common than lingual nerve injury as it only lies below and parallels to the submandibular duct.

Solution to Question 6:

The given clinical vignette in a patient with paresthesia and sensory loss over the medial aspect of the leg and foot is suggestive of injury to the saphenous nerve.

Injury to the saphenous nerve is common following venous cut down for access to the great saphenous vein, in this case for coronary bypass graft.

Saphenous nerve:

It is the longest and largest cutaneous branch of the femoral nerve and is the longest nerve in the body. It arises from the posterior division of the femoral nerve. It then enters the adductor canal and exits it by piercing the fascia lata between the gracilis and sartorius and supplies the skin over the patella. The saphenous nerve then descends along the medial border of the tibia along with the great saphenous vein (anterior to the vein) and supplies the skin on the medial side of the knee, leg, and foot as far as up to the first metatarsophalangeal joint.

Saphenous nerve injury:

When the great saphenous vein is being harvested for coronary bypass graft, injury to the saphenous nerve is common as it closely travels with the great saphenous vein. Sensory loss over the medial side of the knee, leg, and foot is observed in saphenous nerve injury.





Other options:

Option A: Superficial peroneal nerve injury causes sensory loss over the lateral aspect of the leg and the dorsum of the foot.

Option C: Femoral nerve gives rise to anterior cutaneous branches, the saphenous nerve, and motor supply to the quadriceps and sartorius. Anterior cutaneous branches supply the skin on the anterior thigh and knee. If the saphenous nerve was not an option in the question, the femoral nerve would have been the next best answer.

Option D: Sural nerve injury causes sensory loss over the posterior and lateral aspects of the distal third of the leg and lateral side of the foot and fifth toe. It is often used as an autologous peripheral nerve graft as it is readily identified and easy to harvest.

Solution to Question 7:

The structure marked in the given sagittal section of the head and neck is the uvula. The sensory supply to the uvula is by the glossopharyngeal nerve and it is tested in this case.

Nerve supply by glossopharyngeal nerve:

• Sensory fibers to the uvula, tympanic cavity, pharyngotympanic tube, tonsils, fauces, nasopharynx, and posterior third of the tongue.

• Motor fibers to the stylopharyngeus muscle

• Parasympathetic secretomotor fibers to the parotid gland

• Gustatory fibers for the postsulcal part of the tongue.

Pharyngeal plexus:

The nerve supply of the pharynx is mainly from the pharyngeal plexus of nerves. It is formed by the pharyngeal branches of the glossopharyngeal nerve and vagus nerve. It also has contributions from the superior cervical ganglion. The vagus nerve provides the motor supply and the glossopharyngeal nerve provides the sensory supply.

Nerve supply by the pharyngeal branch of the vagus nerve:

• All the muscles of the soft palate(except tensor veli palatini - supplied by mandibular division of the trigeminal nerve)

• All the muscles of the pharynx(except stylopharyngeus - supplied by the glossopharyngeal nerve)

Note:

Previously cranial root of the spinal accessory nerve was thought to contribute to the motor functions of the pharyngeal plexus, but recent evidence suggests that it is not a major part of the motor drive to the pharyngeal plexus.

Innervation of the soft palate:

General sensation - It is mostly carried by the lesser palatine nerve which is a branch of maxillary nerve. The sensation from posterior part is carried by the pharyngeal branches of the glossopharyngeal nerve and from the plexus around the tonsil which is formed by the tonsillar branches of the glossopharyngeal and lesser palatine nerves.

Taste sensation - There are taste buds in the oral surface of soft palate and sensation from them is carried by the lesser palatine nerve.

Secretomotor supply - It is by the lesser palatine nerve via postganglionic fibres from the pterygopalatine ganglion.

Motor supply - It is by the pharyngeal branch of the vagus nerve except for tensor veli palatini which is supplied by mandibular division of the trigeminal nerve.

Anatomy INI-CET 2022

Question 1:

Arrange in sequence the structures involved in the direct pathway:

a) 1,2,3,4,5

b) 5,4,2,3,1

c) 5,1,2,3,4

d) 1,3,4,2,5

Question 2:

What is the volume of the prostate in normal adult males?

a) 20cc

b) 50cc

c) 75cc

d) 100cc

Question 3:

Which ligament connects medial cuneiform to the base of the 2nd metatarsal?

a) Chopart

b) Spring

c) Lisfranc

d) Deltoid

Question 4:

Fibres of the given structure originate from ?



a) Amygdala

b) Mammillary body

c) Caudate nucleus

d) Hippocampus

Question 5:

Which structure is supplied by the nerve causing this elevation?



a) Lateral rectus

b) Superior oblique

c) Risorius

d) Levator palpebrae superioris

Question 6:

Arrange lung hilar structures from anterior to posterior:

a) 1,2,3,4

b) 2,3,4,1

c) 3,4,1,2

d) 4,3,2,1

Question 7:

Which is not a branch of facial artery in the face?

a) Lateral nasal

b) Mental

c) Superior labial

d) Inferior labial

Question 8:

Which of the following structures passing via jugular foramen has a separate canal?

a) Hypoglossal nerve

b) Glossopharyngeal nerve

c) Vagus nerve

d) Accessory nerve

Question 9:

What is the nerve supply of the structure marked in the image?



a) Radial nerve

b) Median nerve

c) Ulnar nerve

d) Musculocutaneous nerve

Question 10:

A child lacks thymus and inferior parathyroid glands. Defective development of which of the following structures is likely to be the cause?

a) 1

b) 2

c) 3

d) 4

Question 11:

Identify which of the following structure is a merocrine gland:



a) 1

b) 2

c) 3

d) 4

Question 12:

What is the blood supply of the structure marked in the image below?



a) Anterior cerebral artery and middle cerebral artery

b) Anterior cerebral artery and posterior cerebral artery

c) Anterior cerebral artery and anterior communicating artery

d) Middle cerebral artery and posterior cerebral artery

Question 13:

Superior gluteal nerve supplies all of the following structures except:

a) Gluteus medius

b) Gluteus minimus

c) Piriformis

d) Tensor fasciae latae

Question 14:

Arrange the structures of the cubital fossa in sequence from medial to lateral.

a) 1,2,3,4

b) 1,3,4,2

c) 1,4,3,2

d) 4,3,2,1

Question 15:

Arrange the cells according to their positions from the basal layer towards the lumen in the seminiferous tubules.

a) 1,5,4,2,3

b) 1,2,3,4,5

c) 5,4,3,2,1

d) 5,1,2,3,4

Question 16:

Which skull foramen is formed by 3 bones?

a) Foramen lacerum

b) Foramen magnum

c) Foramen rotundum

d) Foramen jugulare

Question 17:

Which marked structure causes lateral rotation, flexion, and abduction at the hip joint, along with flexion at the knee joint?



a) 1

b) 2

c) 3

d) 4

Question 18:

Which among the following statements is true regarding the course of the mandibular nerve?

a) 2,3

b) 1,3

c) 3,4

d) 1,4

Question 19:

Identify the incorrectly matched pair(s) of pharyngeal arch and its artery.

a) iv, v

b) iii

c) v

d) ii, iii

Question 20:

The rupture of penile urethra does not cause extravasation of blood into which of the following spaces?

a) Superficial perineal pouch

b) Deep perineal pouch

c) Ischiorectal fossa

d) Skin around the scrotum

Question 21:

Which nerve is blocked when local anesthesia is given below the inguinal ligament during femoral vein catheterization?

a) Femoral nerve

b) Genitofemoral nerve

c) Lateral cutaneous nerve of thigh

d) Ilioinguinal nerve

Question 22:

Identify the nerve passing around the marked structure.



a) Common peroneal nerve

b) Sural nerve

c) Tibial nerve

d) Femoral nerve

Question 23:

Which of the following cells are present in the molecular layer of the cerebellum?

a) 1, 2

b) 1, 3

c) 2, 4

d) 2, 3

Question 24:

Bleeding as shown in the image is due to which of the following vessels?

a) Bridging veins

b) Middle meningeal artery

c) Lenticulostriate artery

d) Vertebral artery

Question 25:

Which of the following statements are true about the internal capsule?

a) 1, 3, 4

b) 2, 3, 4

c) 1, 2, 4

d) 1, 2, 3

Question 26:

Which of the following is not a branch of posterior division of internal iliac artery ?

a) Iliolumbar artery

b) Lateral sacral artery

c) Inferior gluteal artery

d) Superior gluteal artery

Question 27:

Identify the incorrect statement regarding the marked structure.



a) It is the meeting point of frontal, parietal, temporal and sphenoid bones.

b) Blow to the lateral side of the skull injures the marked structure.

c) Accessory middle meningeal artery lies under this structure.

d) It corresponds to site of anterolateral fontanelle of fetal skull.

Question 28:

What are the derivatives of the marked cells?



a) 1, 3, 5, 6

b) 2, 3, 4, 6

c) 1, 2, 3, 6

d) 1, 2, 3, 4

Question 29:

Which of the following muscles form the pelvic diaphragm?

a) A, B

b) A, B, C, D

c) A, B, D

d) A,D

Question 30:

Which of the following arteries passes in close proximity to the area marked by the arrow in the given image?



a) Anterior division of middle cerebral artery

b) Anterior division of the middle meningeal artery

c) Posterior division of middle cerebral artery

d) Stem of middle meningeal artery

Question 31:

Fracture of the jugular canal can present as

a) Loss of taste sensation in anterioir 2/3rd of tongue

b) Loss of taste sensation in posterior 1/3rd of tongue

c) Loss of general sensation in anterior 2/3rd of tongue

d) Loss of motor sensation in anterior 2/3rd of tongue

Question 32:

Injury of the lower brachial plexus leads to Klumpke's paralysis. What is the characteristic clinical finding seen in this lesion?

a) Policeman tip hand

b) Wrist drop

c) Claw hand

d) Horner's syndrome

Question 33:

Identify the liver segment marked in the image below:

a) V

b) IVa

c) VII

d) III

Question 34:

Which of the following lobes of the lung is affected when there is aspiration in supine position?

a) Lingula

b) Upper part of right inferior lobe

c) Right superior lobe

d) Right middle lobe

Question 35:

Identify the cranial nerve that arises from the marked part.



a) Occulomotor nerve

b) Glossopharyngeal nerve

c) Abducens nerve

d) Facial nerve

Question 36:

Which of the following are a part of the limbic system?

a) 1, 2

b) 1, 3

c) 2, 3

d) 1, 2, 3

Question 37:

A 50-year-old male presented with a positive Trendelenburg test on the right side. Identify the nerve injured.

a) Right inferior gluteal nerve

b) Right superior gluteal nerve

c) Left superior gluteal nerve

d) Left inferior gluteal nerve

Answer Key

Question No.	Correct Option
1	c
2	a
3	c
4	d
5	c
6	c
7	b
8	b
9	b
10	c
11	b
12	c
13	c
14	c
15	d
16	a
17	b
18	a
19	c
20	c
21	b
22	a
23	a
24	a
25	c
26	c
27	c
28	c
29	a
30	b
31	b
32	c

33	a
34	b
35	c
36	d
37	b

Detailed Explanations

Solution to Question 1:

The sequence of structures involved in the direct pathway is cortex > striatum > globus pallidus > substantia nigra > thalamus.

There are five parallel circuits that connect the frontal cortex to the striatum, pallidum, substantia nigra, and thalamus. These circuits have both direct and indirect pathways. The direct pathway stimulates the cortex to produce movement, whereas the indirect pathway inhibit it.

The cerebral cortex projects into the striatum. The striatum is seen projected into the pallidum and substantia nigra pars reticularis. Fibres originate from different striatal cells according to their endings in the external and internal pallidum. In the direct pathway, striatal neurones project directly to the internal pallidum. Whereas some other fibres that arise from the neurons project to the external pallidum. This is called the indirect pathway.

The sequence of structures involved in the indirect pathway is cortex > striatum > globus pallidus externa > subthalamic nucleus > globus pallidus interna and substantia nigra >Thalamus

The striatal activity in the direct pathway is mediated by D1 dopamine receptors and in the indirect pathway by D2 dopamine receptors.

Putamen lies lateral and the internal capsule lies medial to the pallidum. It receives fibres from the striatum and subthalamic nuclei. It acts as the main output of the basal ganglia to the thalamus and superior colliculus. The substantia nigra is seen in each cerebral peduncle of the midbrain located deep to the crus cerebri. The levels of dopamine in the substantia nigra and striata are seen reduced in parkinsonism disease.





In parkinsonism disease, due to the death of neurons in substantia nigra activity of the direct pathway is reduced and the indirect pathway is increased. This results in bradykinesia in parkinsonism disease. Whereas in Huntington's disease, the activity of the indirect pathway is affected resulting in choreic involuntary movements.



Solution to Question 2:

The volume of the prostate in normal adult males is 20cc.

Prostate is an accessory glandular tissue embedded in the fibromuscular stroma. It usually weighs 8 grams in the young adults and its weight ranges from 40-150 grams in men aged >50 years.

The prostate gland is situated in the lesser pelvis and surrounds the prostatic urethra which extends from bladder base to the membranous urethra. Pubic symphysis and pubic arch lie anteriorly to the prostate whereas rectourethralis and rectal ampulla lie posteriorly to it.

The prostate has three distinct zones as follows:

Zone	Structures enclosed	Clinical significance
Transitional zone	Urethra	Most common site for benign prostatic hyperplasia (BPH)
Central zone	Ejaculatory ducts	Atrophies secondary to BPH
Peripheral zone	Central and transitional zones Pre-prostatic urethra	Most common site for prostatic carcinoma



The prostate is composed of 5 lobes. They are anterior, posterior, two lateral and one medial or middle lobe.

The median lobe of the prostate forms the uvula vesicae. It is an elevation found in the lower part of the bladder trigone. It is the most common site of adenoma.



Microstructure of the prostate gland:

The glandular tissue of the prostate gland is composed of many follicles with frequent internal papillae. The follicular epithelium is either a single-layered or pseudostratified columnar epithelum. These follicles open into canals, which join to form the main ducts. Small eosinophilic colloid amyloid bodies called corpora amylacea are frequently seen within a follicle.

The prostatic ducts open into the prostatic sinus in the prostatic urethra. The ducts of the prostate follicles are lined by bilayered epithelium (basal cuboidal and apical columnar).

The secretions of the prostate are acidic and contain acid phosphatase, amylase, prostate-specific antigen (PSA), fibrin, and zinc.

The blood supply to the prostate is by the inferior vesical, internal pudendal and middle rectal arteries. It recieves nerve supply from the inferior hypogastric plexus.

Solution to Question 3:

The Lisfranc ligament connects the medial cuneiform to the base of the 2nd metatarsal.

Lisfranc ligament is a cuneometatarsal interosseous ligament. It is the strongest ligament among the three cuneometatarsal interosseous ligaments. It passes from the lateral surface of the medial cuneiform to the adjacent angle of the second metatarsal bone.

Deformity of this ligament can lead to instability and deformity of the tarsometatarsal joint.



Other options:

Option A: Chopart's ligament is related to both calcaneocuboid and calcaneonavicular joints. It is a strong Y-shaped band, having a proximal stem and 2 distal limbs.

• The proximal limb is attached by a stem to the anterior part of the upper calcaneal surface.

• The distal limb divides into calcaneocuboid and calcaneonavicular parts.



Option B: Spring ligament also known as the plantar calcaneonavicular ligament connects the sustentaculum tali to the navicular. It ties the calcaneus to the navicular and maintains the medial longitudinal arch of the foot. It is made up of the superomedial calcaneonavicular portion and the inferior calcaneonavicular portion.



Option D: Deltoid ligament also called the medial collateral ligament is attached to the apex, anterior and posterior borders of the medial malleolus. It has 2 layers, superficial and deep:

• The superficial layer consists of tibionavicular fibres anteriorly, tibiocalcaneal fibres in the middle, and superficial posterior tibiotalar fibres posteriorly.

• Deep layer consists of anterior tibiotalar fibres, tibiocalcaneal in the middle and deep posterior tibiotalar posteriorly.



Solution to Question 4:

The given structure is the fornix and its fibres arise from the hippocampus.

The fornix is a part of the connecting pathways of the limbic system. It arises from the hippocampus which wraps around the thalamus and ends in the mammillary body. It connects the hippocampus with distal brain sites.

The fornix has two types of fibres:

• Association fibres: They connect the various cortical regions within the same hemisphere

• Commissural fibres: They connect the corresponding regions of two hemispheres. It is seen in the hippocampal commissure of the fornix.

The blood supply of fornix is by the anterior communicating artery which arises from the anterior cerebral artery.

The limbic system is a group of structures seen between the cerebral cortex and the hypothalamus. It is involved in emotional and motivation activity, learning, and memory.





Other options:

Option A: Amygdala is a part of the limbic system that consists of lateral, central and basal nuclei. It can Influence the body's response to enviromnental changes. In a fearful situation it can change the heart rate, blood pressure, respiratory rate etc.

Option C: Caudate nucleus is a C-shaped mass of grey matter. It is divided into a head, body, and tail. Degeneration of the caudate nucleus is seen in Huntington's disease.

Solution to Question 5:

Risorius is supplied by the facial nerve which causes the elevation marked. The image shows the floor of the fourth ventricle and the facial colliculus is marked.

The fourth ventricle is located between the brainstem and the cerebellum. The rhomboid fossa is a diamond-shaped depression seen on the floor of the fourth ventricle. The superior and the inferior part of the floor is triangular in shape. It is divided by a longitudinal median sulcus into two halves. Each half is again divided into the medial and lateral regions.

The facial colliculus is present in the superior part, which presents as a small elevation produced by the root of the facial nerve winding around the nucleus of the abducens nerve. Columns are formed by groups of facial neurones, which innervate individual muscles and give branches of the facial nerve. Dorsal neurones innervate muscles in the scalp and upper face and ventral neurons supply the lower facial musculature.

Risorius falls under the buccolabial muscle group and act as the elevator, retractor and evertor of the upper lip. Some of its peripheral attachments are the zygomatic arch, parotid fascia and fascia over the mastoid process. It is involved in actions like grinning and laughing and is innervated by the buccal branches of the facial nerve.





Other options:

Option A: Lateral rectus is an extraocular muscle that arises from the lateral part of the common tendinous ring. It is innervated by the abducens nerve.

Option B: Superior oblique is an extraocular muscle that arises from the body of the sphenoid. It is innervated by the trochlear nerve.

Option D: Levator palpebrae superiors is an extraocular muscle that arises from the lesser wing of sphenoid. It is innervated by the oculomotor nerve.









Solution to Question 6:

Lung hilar structures are arranged from the anterior to posterior in the order:

- Pulmonary vein
- Pulmonary artery
- Primary bronchus
- Bronchial artery

The medial surface of the lung is connected to the heart and trachea by a group of structures that enter or leave the hilum. It is called the pulmonary root. It is anteriorly related to the phrenic nerve, pericardiacophrenic vessels , and anterior pulmonary plexus. The vagus nerve and posterior pulmonary plexus form the posterior relation and the pulmonary ligament the inferior

relation.

The superior vena cava and right atrium are seen in front of the right root and terminal part of the azygos vein superiorly. The hilar structures are arranged from the superior to inferior as:

• Superior lobar bronchus

• Pulmonary artery

• Principal bronchus

• Inferior pulmonary vein

Structures related to right lung



The aortic arch is seen superiorly and descending thoracic aorta posteriorly of the left root. The hilar structures are vertically arranged as:

• Pulmonary artery

• Principal bronchus

• Inferior pulmonary vein

Structures related to Left lung



Solution to Question 7:

Mental is not a branch of the facial artery but the branch of the maxillary artery in the face.

The first part of the maxillary artery gives rise to the mental artery as a terminal branch of the inferior alveolar artery.

The facial artery is a branch of the external carotid artery which arises in the carotid triangle, above the origin of the lingual artery. It runs upwards in the neck and curves around the anteroinferior border of the masseter to enter the face.

The premasseteric artery passes along the anterior border of the masseter and the surrounding tissues are supplied by it. The inferior labial artery mainly supplies the inferior labial glands. It then anastomoses with its controlateral fellow and with the mental branch of the inferior alveolar artery. The superior labial artery supplies the upper lip whereas the lateral nasal artery supplies the dorsum and alae of the nose. Both these arteries anastomose with its controlateral fellow.

The image shows the branches of facial artery:

Branches of facial artery in th e neck	Branches of facial artery in th e face
Ascending palatine artery	Premasseteric artery
Tonsillar branch	Superior and inferior labial a rteries
Submental artery	Lateral nasal artery
Branches to submandibular g land	Angular artery



Solution to Question 8:

The glossopharyngeal nerve passing through the jugular foramen has a separate canal.

The jugular foramen is formed by the occipital bone and the petrous part of the temporal bone. The spinal accessory nerve and vagus nerve also pass through this foramen.

The glossopharyngeal nerve leaves the skull through the anteromedial part of the jugular foramen in a separate dural sheath.

The branches of the nerve are tympanic, carotid, pharyngeal, muscular, tonsillar and lingual branches.

Infection, malignancy and postoperative inflammation of the tonsil and tonsillar fossa will cause referred pain in the ear. This is conducted through the tympanic branch of the glossopharyngeal nerve.

Pharyngeal branches of the glossopharyngeal nerve and vagus nerve form the pharyngeal plexus of nerves. These plexes mainly supply the pharynx. It also has contributions from the superior cervical ganglion.

Nerve supply by the pharyngeal branch of the vagus nerve are:

• All the muscles of the soft palate(except tensor veli palatini - supplied by mandibular division of the trigeminal nerve)

• All the muscles of the pharynx(except stylopharyngeus - supplied by the glossopharyngeal nerve)

Motor fibres to the stylopharyngeus and parasympathetic secretomotor fibres to the parotid gland are also supplied by the glossopharyngeal nerve.

Sensory supply includes uvula, posterior one-third of the tongue, tympanic cavity etc.

The glossopharyngeal nerve is rarely damaged alone. Lesions will result in loss of sensation of the posterior third of the tongue, reduced gag reflex, loss of sensation over the ipsilateral soft palate etc.

The main functions of IX and X can be tested together by assessing swallowing, phonation/articulation and sensation from the pharynx/larynx.

Glossopharyngeal and vagus nerve examinations include:

• Observing the position and symmetry of the palate and uvula at rest and with phonation (“aah”).

• By assessing the strength of the cough

The image below shows the base of the skull and foramina:





Solution to Question 9:

The median nerve supplies the 1st lumbrical which is marked.

There are four lumbrical muscles. They originate from the tendons of the flexor digitorum profundus. It passes to the radial side of the corresponding fingers and is inserted into the dorsal digital expansion of the extensor digitorum.

The action of lumbricals is flexion at the metacarpophalangeal joint and extension of interphalangeal joints.

Lumbricals can be tested only along with interosseous muscle. The examiner will hold the metacarpophalangeal joint of the index finger in hyperextension. Then the patient is asked to extend the proximal interphalangeal joint against resistance. The test is repeated on other fingers as well.

The median nerve innervates the first and second lumbricals. The deep terminal branch of the ulnar nerve innervates the third and the fourth.





The median nerve is formed by the union of the lateral root (C6, 7) from the lateral cord, and the medial root (C8, T1) from the medial cord.

On the hand, it is seen proximal to the flexor retinaculum and lateral to the tendons of flexor digitorum superficialis. It also lies between the tendons of flexor carpi radialis and palmaris longus.

Its cutaneous supply in hand is as follows:

• Lateral 2/3rd of the palm by superficial cutaneous branch

• Lateral 3½ fingers (palmar aspect and dorsal aspect of nailbeds) by palmar digital nerves



The muscles supplied by the median nerve in the hand are flexor pollicis brevis, abductor pollicis brevis, and opponens pollicis

The median nerve sometimes gets compressed under the fibro-osseous tunnel beneath the flexor retinaculum leading to carpal tunnel syndrome.

Pen test is used to test the median nerve in the hand. It tests the abductor pollicis brevis muscle. The hand must remain flat with the palm facing upwards. A pen is held above the palm and the patient is asked to touch the pen with his thumb.

Solution to Question 10:

The given clinical scenario is suggestive of DiGeorge syndrome. Defective development of the third pharyngeal pouch marked as 3 causes this condition.

A child with DiGeorge syndrome lacks thymus and inferior parathyroid glands due to defective development of the 3rd pharyngeal pouch. The child can also lack superior parathyroid glands, due to defective development of the 4th pharyngeal pouch. It is caused by chromosome 22q deletion and is characterized by the following (Mnemonic: CATCH-22)

• Cleft palate

• Abnormal facies

• Thymic aplasia leading to T-cell deficiency and recurrent infections

• Cardiac defects

• Hypocalcemia, secondary to parathyroid aplasia.

Derivatives of the Pharyngeal Pouches

1st Pouch

• Tympanic cavity (middle ear cavity)

• Auditory tube (eustachian tube)

2nd Pouch

• Palatine tonsils

• Tonsillar fossa

3rd Pouch

• Inferior parathyroid glands

• Thymus

4th Pouch

• Superior parathyroid glands

• Ultimobranchial body - gives rise to parafollicular (C) cells of the thyroid gland



The medulla of the thymus has Hassall's corpuscles. They are small rounded structures formed from whorls of medullary epithelial cells with a central partially keratinized core appearing as an eosinophilic-staining mass.

Histological image of Hassall's corpuscles is given below:



Solution to Question 11:

Sweat gland marked 2 is a merocrine gland.

Eccrine and apocrine are the two types of sweat glands. Secretions of the sweat glands are by exocytosis hence they are also known as merocrine sweat glands.

A histopathological image of the sweat gland is shown below:





The other structures marked are:

1- Hair follicle

3- Sebaceous gland

4- Hair shaft



Glands are divided into holocrine, apocrine, and merocrine on the basis of the manner in which their secretions are poured out of the cells.

In holocrine secretions, the secretory cells themselves are disintegrated as part of their terminal differentiation. Afterwhich, the terminally differentiated cells filled with lipid products are released. The sebaceous gland and hair follicles are examples.

A histopathological image of the sebaceous gland is shown below:





Merocrine uses exocytosis for secretions. Most of the exocrine glands are merocrine.

In apocrine, product-filled areas of cells are released, the mammary gland is an example.

Histopathological image of mammary gland is shown below:





Solution to Question 12:

Blood is supplied by the anterior cerebral artery and anterior communicating artery as the marked structure is the anterior part of the corpus callosum.

Corpus callosum is the largest fibre pathway of the brain. The parts of the corpus callosum are the rostrum, genu, trunk or body, and splenium.

The image below shows the parts of the corpus callosum:



Anterior cerebral artery is a branch of the internal carotid artery. The frontal branch of the anterior cerebral artery supplies the corpus callosum and the central branch which supplies the rostrum.

Central or perforating arteries arises from the circle of willis. They are again divided into the anteromedial, anterolateral, posteromedial and posterolateral groups. The anterior cerebral and anterior communicating artery give rise to anteromedial group which supplies the rostrum of the corpus callosum.

The image below shows the mid sagittal section of the brain:



Solution to Question 13:

All of the structures except pyriformis are supplied by the superior gluteal nerve. Pyriformis is supplied by the nerve to the piriformis.

The dorsal branches of the L4, L5, and S1 rami give rise to the superior gluteal nerve. It is accompanied by superior gluteal vessels. This nerve leaves the pelvis via the greater sciatic foramen above piriformis and divides into superior and inferior branches. It supplies gluteus medius, gluteus minimus, and tensor fasciae latae.

The dorsal branches of the S1 and S2 rami give rise to the nerve to the piriformis.

Gluteus medius and gluteus minimus act as the abductors of the hip joint. Stabilizing the pelvis on the femur during walking is done by these muscles. It is demonstrated by asking the subject to stand on one limb. If the ipsilateral abductors are normal, the pelvis remains in a horizontal position. If the ipsilateral abductors are weak, then the pelvis tilts towards the opposite (unsupported) side. This is called positive Trendelenburg's sign.



Gluteal region



Solution to Question 14:

The structures are arranged in the cubital fossa from medial to lateral as follows:

• Median nerve

• Brachial artery

• Tendon of biceps brachii

• Radial nerve

The cubital fossa is a triangular depression in the anterior aspect of the upper forearm. It is related to the epicondylar line proximally, lateral border of pronator teres medially, and by the medial surface of brachioradialis laterally.

Its roof is formed by the deep fascia of the forearm. The floor of the fossa is formed by the brachialis and supinator. Veins like basilic, median cubital, and cephalic veins are arranged in the superficial fascia from medial to lateral side.



Solution to Question 15:

From the basal layer towards the lumen in the seminiferous tubules, the cells are arranged in the order of:

• Myoepithelial cells

• Spermatogonia

• Primary spermatocyte

• Spermatid

• Spermatozoa

Spermatogenesis is the process in which spermatogonia are transformed into spermatozoa. It begins at puberty. During puberty, the sex cords acquire lumen to become seminiferous tubules.

The primordial germ cells give rise to spermatogonial stem cells. The transformation of spermatogonial stem cells to spermatozoa is given in the flowchart below.



The process by which spermatids are transformed into spermatozoa is called spermiogenesis.



Solution to Question 16:

Foramen lacerum is formed by three bones.

The three bones involved in the formation of foramen lacerum are:

- Petrous part of the temporal bone
- Greater wing of the sphenoid bone
- Basilar part of the occipital bone

The internal carotid artery enters the skull through this foramen. It ascends through the carotid canal and from the upper end of the foramen lacerum.

It is accompanied by venous and sympathetic nerve plexuses. It is also transversed by the meningeal branches of the ascending pharyngeal artery and emissary veins from the cavernous sinus.

On the anterior margin of the foramen lacerum pterygoid canal can be seen which leads to the pterygopalatine fossa. The nerve of the pterygoid canal and accompanying vessels are transmitted by them.

The image below shows the base of the skull and foraminas:



Other options:

Option B: Foramen magnum is formed by the occipital bone.

Option C: Foramen rotundum is a hole seen in the greater wing of the sphenoid bone.



Option D: Foramen jugulare is formed by the petrous part of the temporal bone and occipital bone.

The image below shows superior and inferior views of the skull base:



Solution to Question 17:

Sartorius is involved in lateral rotation, flexion, and abduction at the hip joint, and flexion at the knee. The structure marked '2' is sartorius.

The sartorius muscle is known as the tailor muscle and is also the longest muscle in the body. The word "sartorius" is derived from the Latin word sartor, which translates to tailor, due to the way the individual will position one's leg while working.

Origin: Anterior superior iliac spine and the upper half of the notch below it.

Insertion: Into the superior aspect of the medial surface of the tibial shaft near the tibial tuberosity (joins aponeurosis of gracilis and semitendinosus to form pes anserinus).

Action: Flexion, abduction, and lateral rotation of the thigh; and flexion of the leg.

Innervation: By femoral nerve (L2 and L3)



Other options:

Option A: Rectus femoris is involved in flexion at the hip and extension at the knee.

Option C: Gracilis is involved in adduction at the hip and flexion and medial rotation at the knee.

Option D: Adductor longus is involved in adduction, flexion, and medial rotation at the hip.

Solution to Question 18:

The true statements are 2 and 3.

The lingual nerve is a branch from the posterior division of the mandibular nerve, and it arcs around the submandibular duct.

The buccal branch is a branch from the anterior division of the mandibular nerve, and it emerges from behind the mandibular ramus.

The other statements:

1. The auriculotemporal nerve emerges from behind the temporomandibular joint and is a branch from the posterior division of the mandibular nerve.

4. The ophthalmic nerve arises in the middle cranial fossa and is a division of the trigeminal nerve, along with the maxillary and mandibular nerves.

Solution to Question 19:

All the given pairs of pharyngeal arteries and their arteries are correctly matched except for 'v' (option C).

The 6th pharyngeal arch has its blood supply on the left side from the left pulmonary artery and ductus arteriosus, not from the aorta.

Solution to Question 20:

Rupture of the penile urethra causes extravasation of blood into the superficial perineal pouch, scrotum, penis, and then up to the lower anterior abdominal wall.

The superficial perineal pouch is a compartment of the perineum enclosed by the membranous layer of the perineal fascia that surrounds the scrotum or labia and penis or clitoris.

Anteriorly, the membranous layer of perineal fascia is continuous over the pubic symphysis and pubic bones. It is later continuous with the membranous layer of the anterior abdominal wall. Thus, the fluid that accumulates in the perineal pouch extends to the lower anterior abdomen.

In the lower lateral abdominal wall, the abdominal fascia is attached to the deep fascia of the thigh, inferior to the inguinal ligament. Thus, on trauma at the penile urethra, urine does not extravasate into the anterior thigh.

The superficial perineal pouch is closed posteriorly by attaching the membranous layer of superficial fascia to the posterior margin of the urogenital diaphragm. Thus, commonly there is no extravasation into the deep perineal pouch and the ischiorectal fossa.

Also, the membranous layer of superficial fascia attaches to the ischiopubic rami and forms the lateral borders of the urogenital triangle. Thus, the urine does not extravasate to the thigh.

But, the ischiorectal fossa would be a better answer because of the following reasons:

• Bulbourethral glands that lie in the deep perineal space open their ducts into the spongy urethra, which is related to the superficial perineal space. Also, the muscles surrounding these glands were recently found to be a continuation of the deep transverse perinei and tail of the external urethral sphincter, both of which are components of the deep perineal space. Injury to the penile urethra can damage bulbourethral glands' ducts, which can lead to damage to the above-mentioned structures and may cause extravasation of urine into the deep perineal space.

• Previously, the perineal membrane was thought to be a triangular membrane. However, recent studies show that it is contiguous anteriorly with the structures that support the urethra. Posteriorly, it is related to the support of the perineal body and lateral vaginal wall. So damage to the urethra might damage the anterior part of the perineal membrane which can make way for extravasation of blood into the deep perineal space.

• Deep perineal pouch: It was believed initially that the urogenital diaphragm was formed by 3 layers: An inferior fascia (also called the perineal membrane), deep perineal space, and a superior fascia. The superior fascia formed the inferior border of the ischioanal fossa. The Colles fascia also separates the perineal membrane from the ischioanal fossa. So, extravasation of blood into the deep perineal space is possible but extravasation into the ischioanal fossa due to penile urethral injury is highly unlikely. Recent studies have shown that this might be a slightly more complex structure.





Solution to Question 21:

Genitofemoral nerve is blocked when the local anesthetic is given below the inguinal ligament during femoral vein catheterization.

The genitofemoral nerve originates from the L1 and L2 ventral rami. It divides into the genital and femoral branches after crossing behind the ureter.

The genital branch enters the inguinal canal through the deep ring and accompanies the spermatic cord or round ligament. It exits the superficial inguinal ring and supplies the cremaster muscle and skin of the external genitalia.

The femoral branch descends lateral to the external iliac artery to pass behind the inguinal ligament and enter the femoral sheath lateral to the femoral artery. It pierces the anterior wall of the femoral sheath and fascia lata and supplies them.

The genitofemoral nerve gives sensory supply to the anteromedial thigh via the femoral branch. The genital branch supplies skin of the scrotum in males and mons pubis and labium majus in females. The genitofemoral nerve also innervates the cremaster muscle via the genital branch and is responsible for the cremasteric reflex.

The image given below shows nerves related to the inguinal region:



Other options:

Option A: Femoral nerve arises from the posterior division of ventral rami of L2-L4 spinal segments on the posterior abdominal wall. It enters the femoral triangle by passing under the inguinal ligament. In the femoral triangle, it lies lateral to the femoral artery and outside the femoral sheath. Later, it is divided into anterior and posterior branches by the lateral circumflex femoral artery.

Option C: Lateral cutaneous nerve (a.k.a lateral femoral cutaneous nerve) of the thigh is most commonly injured at the site where it leaves the pelvis. An injury to the lateral cutaneous nerve of the thigh leads to pain and paresthesia over the anterolateral region of the thigh, a condition called meralgia paresthetica.

Option D: Ilioinguinal nerve is most commonly injured within the inguinal canal in open hernia repair due to traction or thermal injury. Injury causes loss of sensation in the root of the penis and adjacent part of the scrotum.

Solution to Question 22:

The structure marked in the given image is the neck of fibula. The nerve passing around the neck of fibula is the common peroneal nerve (common fibular nerve).

The common peroneal nerve is derived from the dorsal branches of the ventral rami of L4, L5, S1, and S2. The sciatic nerve divides into the tibial nerve and the common fibular nerve.

The common fibular nerve runs obliquely along the lateral side of the popliteal fossa to the fibular head. It lies between the biceps femoris and the lateral head of the gastrocnemius. It then curves

laterally around the neck of the fibula and gets divided into superficial and deep fibular nerves.

Superficial fibular nerve:

Motor innervation:

• Peroneus longus

• Peroneus brevis

Sensory innervation:

• Lateral aspect of the lower leg

• Dorsum of the toes except for the lateral side of the 5th toe and the first interdigital webspace

Deep fibular nerve:

Motor innervation:

• Extensor digitorum longus

• Extensor hallucis longus

• Tibialis anterior

• Peroneus tertius

Sensory innervation:

• The first interdigital space







The common peroneal nerve is more prone to injury as it lies superficial at the level of the lateral aspect of the neck of the fibula. Injury results in foot drop (which causes high-stepping gait), loss of eversion of the foot, and sensory loss on the dorsum of the foot.

Solution to Question 23:

The molecular layer of the cerebellum contains stellate and basket cells.

The molecular layer is the outermost layer that contains stellate cells and basket cells, and dendrites of Purkinje cells. They receive excitatory input from the parallel fibers and form inhibitory synapses on the Purkinje cells. This layer contains many myelinated and unmyelinated nerve fibers and appears pale on H&E staining.



Solution to Question 24:

The image showing a bleed over the arachnoid layer but under the dura is suggestive of a subdural hemorrhage. Most commonly, it results from the shearing of cortical bridging veins.

Subdural hemorrhage (SDH) is a collection of blood between the dura and the arachnoid layers of the meninges. The causes of SDH include

• trauma

• anticoagulants, antiplatelets

• alcohol abuse

• battered baby syndrome.

Non-contrast CT is the investigation of choice.

Other options:

Middle meningeal artery rupture would result in an extradural hemorrhage.

Lenticulostriate artery would lead to an intraparenchymal hemorrhage.

Vertebral artery dissection leads to subarachnoid hemorrhage or sometimes intraparenchymal hemorrhage.



The given image shows types of brain hemorrhage:



Solution to Question 25:

Statements 1, 2, and 4 are true about the internal capsule.

The charcot's artery of hemorrhage is a larger lenticulostriate or lateral striate artery. Lateral striate arteries supply the anterior limb, genu and the posterior limb of the internal capsule. These arteries are also commonly involved in ischemic and hemorrhagic strokes.

The retrolentiform part of the internal capsule carries fibers mainly from optic radiation. A lesion here produces homonymous hemianopia, superior quadrantanopia, or inferior quadrantanopia.

The anterior limb of the internal capsule carries fronto-pontine fibers. A lesion here produces confusion, impaired attention, agitation, and dysarthria.

The posterior limb of the internal capsule carries cortico-spinal motor fibers in the anterior part and the lesion here produces contralateral hemiparesis. The posterior part of the posterior limb of the internal capsule carries sensory fibers and lesions here may produce contralateral hemi-sensory deficits.

The internal capsule predominantly contains projection fibers, not commissural fibers. Projection fibers are the fibers that project from higher to lower centers or vice versa in CNS.



Solution to Question 26:

The inferior gluteal artery is a branch of the anterior division of the internal iliac artery.



Branches of the anterior trunk of the internal iliac artery:

• Obturator artery

• Superior vesical artery

• Inferior vesical artery

• Inferior gluteal artery

• Middle rectal artery

• Vaginal artery (in females)

• Uterine artery (in females)

• Internal pudendal artery

Branches of the posterior trunk of the internal iliac artery:

• Iliolumbar artery

• Lateral sacral arteries

• Superior gluteal artery



Solution to Question 27:

The structure marked in the image is the pterion. The incorrect statement is statement 3, which is the accessory middle meningeal artery lies under this structure.

The pterion is an H-shaped sutural junction between four bones of the skull:

• Frontal bone

• Squamous part of the temporal bone

• Greater wing of the sphenoid bone

• Parietal bone

It is located approximately 4 cm above the midpoint of the zygomatic arch.



The pterion overlies the anterior branch of the middle meningeal artery and the lateral fissure of the cerebral hemisphere. The bone in this area is thin, and any injury to this area may lead to rupture of the middle meningeal artery, leading to extradural hematoma.

The pterion also corresponds to the site of the anterolateral (sphenoidal) fontanelle of the neonatal skull, which closes in the third month after birth.



The accessory meningeal artery may be a branch of the middle meningeal artery or might arise directly from the maxillary artery. It reaches the cranium via the foramen ovale and distributes principally in the extracranial region. It is distributed over the medial and lateral pterygoids, tensor veli palatini, greater wing and pterygoid processes of the sphenoid bone, mandibular nerve, and the otic ganglion. So, it supplies the trigeminal ganglion, dura mater and bone. But in some people, it may be replaced by small separate arteries.

Solution to Question 28:

The cells shown in the image are neural crest cells (NCC) present adjacent to the neural tube (NT). NCCs contribute to many organs and tissues, and the neural crest is sometimes referred to as the fourth germ layer:

• In the trunk region, they form:

• Melanocytes

• Sensory ganglia

• Sympathetic and enteric neurons

• Schwann cells

• Adrenal medulla cells

• In the cranial region, they contribute to the:

• Craniofacial skeleton

• Neurons for cranial ganglia

• Satellite glial cells

• Melanocytes

• Some of the other neural crest derivatives are

• C cells of the thyroid gland

• Conotruncal septum in the heart

• Odontoblasts

• Smooth muscle cells to blood vessels to the face and forebrain.

The intermediate mesoderm is a longitudinal strip of intraembryonic mesoderm that connects the paraxial with the lateral plate mesoderm. The intermediate mesoderm develops into the genital ridge and nephrogenic cord.

The majority of the heart, including the right ventricle and outflow tract, is derived from the bulbus cordis, which is further derived from the pharyngeal mesoderm.



Solution to Question 29:

Iliococcygeus and pubococcygeus are parts of the pelvic diaphragm, which is also known as the levator ani.

The levator ani muscle is a complex unit that consists of 3 muscle components:

• Pubococcygeus

• Pubovaginalis - elevates the urethra and hence contributes to urinary continence

• Puboperinealis - draws the perineal body towards the pubic symphysis

• Puboanalis - elevates the anus and keeps the urogenital hiatus narrowed

• Puborectalis

• Iliococcygeus

Levator ani muscles are the most important muscles in the pelvic floor and maintain a constant state of contraction. They provide a stable floor and support the weight of the abdominopelvic contents against intra-abdominal forces. Weakening of levator ani and pelvic fascia from stretching or tearing during childbirth, may decrease support for the vagina, bladder, uterus, or rectum or alter the position of the neck of the bladder and urethra.

Other options:

The obturator internus forms the lateral wall of the ischioanal fossa and is not a part of the levator ani.

The deep transverse perineal muscle is the content of the deep perineal space and contributes to the formation of the perineal body.



Solution to Question 30:

The area marked by an arrow in the given image is pterion and the artery that passes in close proximity to this area is the middle meningeal artery.

The pterion is an H shaped sutural junction between four bones of the skull:

• Frontal bone

• Squamous part of the temporal bone

• Greater wing of the sphenoid bone

• Parietal bone.

It is an important structure as it overlies the anterior branch of the middle meningeal artery and lateral fissure of the cerebral hemisphere. The bone in this area is thin and any injury to this area may lead to rupture of the middle meningeal artery leading to extradural hematoma.



Solution to Question 31:

Fracture of the jugular canal affects glossopharyngeal nerve and presents as loss of taste sensation in the posterior 1/3rd of the tongue.

The lingual nerve which is a branch of the mandibular division of the trigeminal nerve gives general sensation, chorda tympani branch of facial nerve gives taste sensation for the anterior 2/3rd of tongue. Both taste and general sensation of the posterior 1/3rd is by the glossopharyngeal nerve. The muscles of the tongue, with the exception of palatoglossus, are supplied by the hypoglossal nerve. Palatoglossus is supplied via the pharyngeal plexus.

The other nerves passing through the jugular foramen are glossopharyngeal(cranial nerve IX), vagus(cranial nerve X), and accessory nerve(cranial nerve XI).

The following is an image of the base of skull showing jugular foramen:



Solution to Question 32:

The characteristic clinical finding in Klumpke's paralysis is a complete claw hand.

Klumpke's paralysis is caused by injury of the lower trunk of the brachial plexus-mainly T1 and partially C8, due to traction injury of the lower trunk that may be seen in undue abduction of the arm (outstretched arm grabbing an object during a fall from height) or birth injury (traction of the extended arm during delivery)

This leads to paralysis of the intrinsic muscles of the hand (T1) that are responsible for flexion at metacarpophalangeal joints (MCP) and extension at proximal and distal interphalangeal joints(PIP and DIP). The unopposed action of extensor digitorum at MCP causes hyperextension. The unopposed action of flexor digitorum superficialis and profundus at PIP and DIP causes flexion. Thus giving rise to the characteristic claw hand.

Other changes include anesthesia and analgesia along the ulnar border of the forearm, vasomotor changes, absence of sweating, and trophic changes in long-standing cases. If the C8 is also damaged, the extent of anesthesia will be greater and will involve the medial side of the forearm, hand, and medial two fingers.

Horner's syndrome (option D) may occur in one-third of the cases due to injury of sympathetic fibers to the head and neck that leave the spinal cord through nerve T1.

The following image shows a complete claw hand in Klumpke's paralysis:

Other options:

Option A: Policeman tip deformity, also known as waiter's tip hand or porter's tip hand, is a finding of Erb's palsy due to injury of the upper trunk of the brachial plexus involving C5 and C6. The hand is adducted, medially rotated, and hangs by the side with the elbow extended and pronated.

The following image shows policeman tip deformity in Erb's palsy:



Option B: Wrist drop is a feature of radial nerve injury in the axilla, radial groove, anterolateral side of the lower arm, or injury of the deep branch in the cubital fossa.

Option D: Horner's syndrome is caused due to the disruption of sympathetic supply to the face. It causes ptosis, miosis, anhydrosis, enophthalmos, and loss of ciliospinal reflex of the affected eye. It may occur in some cases of Klumpke's paralysis and is not a characteristic finding.

The following image shows the ocular manifestation of Horner syndrome:

Solution to Question 33:

The liver segment highlighted in green in this CT section is segment V. The section of the CT given is taken below the portal vein as the cross-section of the portal vein before its bifurcation can be viewed.

A horizontal plane known as the portal plane divides the liver where the portal vein bifurcates and becomes horizontal. This divides each section of the liver into superior and inferior segments:

- Left lateral section: segment II above and segment III below the portal plane
- Left medial section: segment IVa above and segment IVb below the portal plane
- Right anterior (or medial) section: segment VIII above and segment V below the portal plane
- Right posterior (or lateral) section: segment VII above and segment VI below the portal plane

Shown below is a section from below the portal plane.



Couinaud classification is the commonly used classification of the liver. It divides the liver is divided into eight segments. It is based on the distribution of portal venous branches in the liver parenchyma. Each segment has its own vascular supply, biliary drainage, and lymphatic drainage.

The Couinaud's liver segments are as follows:



Solution to Question 34:

The upper part of the right inferior lobe is prone to aspiration in the supine position. It is commonly seen in alcoholics.

The posterior segment of the right superior lobe and the superior segment of the inferior lobes are the most commonly involved lung sites when aspiration occurs in a recumbent patient.

The location of aspiration depends on the position of the patient since it is gravity-dependent. Aspiration while standing leads to infiltration of bilateral inferior lobes because of larger and more vertical corresponding bronchi, the middle lobes and the lingula.

Aspiration of contents into foreign substances into the bronchial tree, usually oral or gastric contents, produces aspiration pneumonia. Chemical (gastric acid) injury or bacterial infection complicates the condition.

Solution to Question 35:

The structure marked in the image is the abducens nerve that arises from the pontomedullary junction.

In the groove between the medulla and pons, abducens, facial, and vestibulocochlear nerves emerge arranged medially to laterally.

The abducens nerve is the 6th cranial nerve that arises from the brainstem at the pontomedullary junction, passes through Dorello's canal, and enters the cavernous sinus where it lies lateral to the internal carotid artery. It then enters the orbit through the common tendinous ring in the superior orbital fissure to supply the lateral rectus.

Solution to Question 36:

The amygdala, hippocampus, and thalamus are all a part of the limbic system.

The limbic system refers to the interconnected neuronal circuit in the basal part of the brain that controls emotional behavior and other vegetative functions such as body temperature, the osmolality of the body fluids, and the drives to eat and drink and to control body weight.

The limbic system includes the following:

• Subcallosal gyrus

• Cingulate gyrus

• Parahippocampal gyrus

• Hippocampal formation

• Amygdaloid nucleus

• Mammillary bodies

• Anterior thalamic nucleus

• Hypothalamus

The connecting pathways for this system comprise the alveus, the fimbria, the fornix, the mammillothalamic tract, and the stria terminalis.



Solution to Question 37:

Positive Trendelenburg test on the right side indicates a right superior gluteal nerve injury.

When the patient stands on his right foot, there is drooping of the left pelvis, this is a Positive Trendelenburg test on the right side.

The gluteus medius and minimus are powerful abductors of the hip supplied by the superior gluteal nerve.

Normally, when the leg is lifted off the ground, the gluteus medius and minimus on the opposite (unsupported) side exert traction on the hip bone. This results in a slight raising of the pelvis on the unsupported side, which helps maintain the trunk upright position.

If the gluteus medius and minimus are paralyzed, the pelvis drops on the contralateral side. This is known as a positive Trendelenburg sign. It is positive in the paralysis of the hip abductors, and in developmental dysplasia of the hip. A characteristic lurching gait is seen.

The image below shows the Trendelenburg test.


Trendelenberg test
Normal
Pelvic stabilisation
Paralysis of gluteus medius and minimus muscle
No pelvic stabilisation on affected limb
Trendelenberg
Sign positive when made to stand on affected limb
Pelvic sags on healthy side
Iliac crest
Bony landmark to diagnose this problem in pelvic stabilisation
MARROW

Anatomy NEET 2022

Question 1:

The structure marked A in the image below gives rise to which of the following structures?



a) Gastrosplenic ligament

b) Lienorenal ligament

c) Falciform ligament

d) Gastrophrenic ligament

Question 2:

A patient underwent surgery for the varicose vein. He now complains of sensory loss over the medial aspect of the leg and foot. Which of the following nerves is most likely to be injured?

a) Sural nerve

b) Superficial peroneal nerve

c) Deep peroneal nerve

d) Saphenous nerve

Question 3:

Identify the cartilage given below:



a) Non articular hyaline cartilage

b) Articular hyaline cartilage

c) Yellow cartilage

d) White fibrocartilage

Question 4:

A patient presents with a unilateral throbbing headache, photophobia, and excessive lacrimation. He also complains of hemifacial pain on clenching of teeth. On examination, pupillary reaction, light reflex, and accommodation reflex are normal. Which of the following marked nerves is involved in the above scenario?



a) 1

b) 2

c) 3

d) 4

Question 5:

Identify the given structure:



a) Lymph node

b) Glomerulus

c) Spleen

d) Pancreatic islet cells

Answer Key

Question No.	Correct Option
1	c
2	d
3	d
4	c
5	d

Detailed Explanations

Solution to Question 1:

The structure marked A is ventral mesentery. The falciform ligament is a derivative of the ventral mesentery.



Mesenteries are double layers of the peritoneum that enclose an organ and connect it to the body wall. They provide pathways for vessels, nerves, and lymphatics to and from the abdominal viscera.

• Ventral mesentery: Only the foregut has a ventral mesentery. It is attached to the terminal part of the esophagus, the stomach, and the upper part of the duodenum. The ventral mesentery is derived from the septum transversum. When the liver grows into the mesenchyme of the septum transversum, it divides the ventral mesentery into 2 parts:

• Lesser omentum- extends from the lower portion of the esophagus, the stomach, and the upper portion of the duodenum to the liver

• Ligaments of the liver- all ligaments of the liver (falciform ligament, right and left triangular ligaments, and coronary ligaments), except ligamentum teres

• Dorsal mesentery: The foregut, midgut, and hindgut have a dorsal mesentery. This connects the organs to the posterior abdominal wall. It runs from the lower end of the esophagus to the cloacal region of the hindgut. It has the following special names in different regions:

• Dorsal mesogastrium (or greater omentum)- stomach

• Dorsal mesoduodenum- duodenum

• Mesentery proper- jejunum and ileum

• Mesoappendix- appendix

• Transverse mesocolon- transverse colon

• Sigmoid mesocolon- sigmoid colon

• Gastrosplenic ligament (stomach to the spleen), lienorenal ligament (spleen to the left kidney), and gastrophrenic ligament (stomach to diaphragm) are derived from the dorsal mesentery.





Solution to Question 2:

The given clinical scenario in a patient with sensory loss over the medial aspect of the leg and foot is suggestive of injury to the saphenous nerve. Injury to the saphenous nerve is common following varicose vein surgery.

The saphenous nerve is the longest and largest cutaneous branch of the femoral nerve and is the longest nerve in the body. It arises from the posterior division of the femoral nerve. It then enters the adductor canal and exits it by piercing the fascia lata between the gracilis and sartorius and supplies the skin over the patella. The saphenous nerve then descends along the medial border of the tibia along with the great saphenous vein (anterior to the vein) and supplies the skin on the medial side of the knee, leg, and foot up to the first metatarsophalangeal joint.

During stripping of veins in varicose surgery, injury to the saphenous nerve is common as it closely travels with the great saphenous vein. Sensory loss over the medial side of the knee, leg, and foot is observed in saphenous nerve injury.





Other options:

Option A: Sural nerve injury causes sensory loss over the posterior and lateral aspects of the distal third of the leg and lateral side of the foot and fifth toe. It is often used as an autologous peripheral nerve graft as it is readily identified and easy to harvest.

Option B: Superficial peroneal nerve injury causes sensory loss over the lateral aspect of the leg and the dorsum of the foot. It also causes weakness of foot eversion.

Option C: Deep peroneal nerve injury causes weakness of ankle dorsiflexion and toe extension but normal eversion of the foot.

Solution to Question 3:

The given image shows prominent collagen fibers arranged in bundles with intervening rows of chondrocytes. This is suggestive of white fibrocartilage.

Microscopy of fibrocartilage shows the following:

• Chondrocytes- round in shape (in tendons they are flattened and elongated)

• Ground tissue with collagen fibers- type I collagen fibers

• Absence of perichondrium



Fibrocartilage is found in:

• Articular discs (temporomandibular joint and sternoclavicular joint)

• Intervertebral discs

• Menisci

• Glenoid and acetabular labrum

• Pubic symphysis

• Manubriosternal joint

• The lining of the bony grooves for tendons

The image below describes the histology of different types of catilages.



Solution to Question 4:

The above patient presents with features suggesting an involvement of the trigeminal nerve. The structure marked 3 is the trigeminal nerve.

Other marked structures:

• 1- Optic nerve. The normal visual field and light reflex rules out the involvement of the optic nerve.

• 2- Oculomotor nerve. Normal accommodation and pupillary reflex rule out the involvement of the oculomotor nerve.

• 4- Facial nerve. Absence of motor involvement of the face rule out the involvement of the facial nerve.



The trigeminal nerve leaves the anterior aspect of the pons as a small motor root and a large sensory root. The large sensory root expands to form the crescent-shaped trigeminal ganglion, which lies within a pouch of dura mater called the trigeminal or Meckel's cave. The ophthalmic, maxillary, and mandibular nerves arise from the anterior border of the ganglion.

Branches of the trigeminal nerve:

• The ophthalmic nerve (V1)

• Leaves the skull through the superior orbital fissure to enter the orbital cavity.

• Contains only sensory fibers

• Cornea (photophobia)

• The skin of the forehead, scalp, eyelids, and nose

• The mucous membrane of paranasal sinuses and nasal cavity

• The maxillary nerve (V2)

• Leaves the skull through the foramen rotundum.

• Contains only sensory fibers from

• Teeth of the upper jaw

• The skin of the face over the maxilla

• The mucous membrane of the nose, the maxillary sinus, and the palate

• The mandibular nerve (V3)

• Leaves the skull through the foramen ovale.

• Contains both sensory and motor fibers.

• Sensory from:

• The skin of the cheek, over mandible, and side of the head

• Teeth of the lower jaw and temporomandibular joint (hemifacial pain on clenching of teeth)

• The mucous membrane of the mouth and anterior part of the tongue

• Motor to:

• Muscles of mastication (pterygoids, masseter, and temporalis)

• Tensor tympani

• Anterior belly of digastric

• Mylohyoid

Solution to Question 5:

The above image shows pancreatic islet cells. Islet of Langerhans makes up the endocrine part of the pancreas.

Islets are characterized by secretory cells supported by reticulin fibers, and containing numerous fenestrated capillaries. There is a delicate capsule around each islet. They are paler than the surrounding exocrine cells, as they have less rough endoplasmic reticulum.

The image given below shows the histology of the pancreas.



The islets of Langerhans contain four major and two minor cell types:

• Four main types are β, α, δ, and PP (pancreatic polypeptide) cells:

• α or A cells- located peripherally and secrete glucagon

• β or B cells- located centrally and produce insulin

• δ or D cells- scattered between beta cells and secrete somatostatin

• PP cells- are present in islets and also in the exocrine pancreas. They secrete a pancreatic polypeptide, which exerts several gastrointestinal effects, such as stimulation of secretion of gastric and intestinal enzymes and inhibition of intestinal motility.

• Minor cell types:

• D1 cells secrete vasoactive intestinal peptide (VIP), a hormone that induces glycogenolysis and hyperglycemia. It also stimulates gastrointestinal fluid secretion and causes secretory diarrhea.

• Enterochromaffin cells synthesize serotonin and are the source of pancreatic tumors that cause carcinoid syndrome.

Other options:

Option A: Lymph nodes are secondary lymphoid organs. The structure of lymph nodes consists of an outer cortex and inner medulla. The cortex contains an aggregation of B-lymphocytes (primary follicles) and quiescent secondary follicles with a pale central germinal center.

The inner medulla consists of medullary cords (lymphocytes and plasma cells) and medullary sinuses. Between the cortex and medulla, there is an ill-defined zone called the paracortex, which contains T lymphocytes.

The image given below shows the histology of the lymph node.



Option B: Glomerulus is a rounded tuft of anastomosing capillaries. The glomerular capillary wall is the filtering membrane and consists of fenestrated endothelial cells, basement membrane, and visceral epithelial cells called podocytes, which contain foot processes.

The glomerular tuft is supported by the mesangium containing mesangial cells and a non-cellular mesangial matrix.

The image given below shows the histology of the glomerulus.



Option C: Spleen is a secondary lymphoid organ. It contains round white pulp surrounded by red pulp. The white pulp is made up of aggregations of lymphocytes that surround a small artery or arterioles. Lymphocytes surrounding the arteriole are called peri-arterial lymphatic sheath (PALS) and contain T cells. Follicles in white pulp contain B cells.

The red pulp is a modified lymphoid tissue infiltrated with all the cells of circulating blood. It is like a sponge permeated by spaces (splenic sinusoids) lined by reticular cells. The zone of red pulp immediately surrounding white pulp is the marginal zone. This zone has a rich network of

sinusoids.

The image given below shows the histology of the spleen.



Anatomy NEET 2023

Question 1:

A defect in which of the following forms the structure marked below?



a) Internal oblique

b) External oblique

c) Parietal peritoneum

d) Fascia transversalis

Question 2:

A student had his jaw locked while yawning. Which of the following muscles is attached to the articular disc of the temporomandibular joint?

a) Lateral pterygoid

b) Temporalis

c) Medial pterygoid

d) Masseter

Question 3:

Identify the structure(s) in the following histological image.



a) Leydig cells of testis

b) Pancreatic islet cells

c) Hassall's corpuscles

d) Glomerulus

Question 4:

Injury at which of the following marked sites causes failure of dorsiflexion?



a) 3

b) 2

c) 1

d) 4

Question 5:

A defect in which of the following aortic arches causes the defect shown in the image?



a) 4th right

b) 4th left

c) 6th right

d) 6th left

Question 6:

Which of the following structures develops from the structure marked A?



a) Medial umbilical ligament

b) Median umbilical ligament

c) Meckel's diverticulum

d) Ligamentum teres

Question 7:

A patient comes with abdominal pain, jaundice, and portal hypertension. Anastomosis between which of the following veins is seen?

a) Left colic vein and middle colic veins

b) Esophageal veins and left gastric veins

c) Superior rectal and phrenic veins

d) Sigmoid and superior rectal veins

Question 8:

A patient presents with swelling under the left ear lobule and complains of pain. Which of the following structures is the likely cause of the pain?

a) Facial nerve

b) Greater auricular nerve

c) Auriculotemporal nerve

d) Investing layer of deep cervical fascia

Answer Key

Question No.	Correct Option
1	d
2	a
3	d
4	a
5	d
6	b
7	b
8	b

Detailed Explanations

Solution to Question 1:

The structure marked is the deep inguinal ring, which is formed by a defect in fascia transversalis.

The deep inguinal ring is an opening in the transversalis fascia present 1 cm above the inguinal ligament, midway between the anterior superior iliac spine and the pubic symphysis. An indirect inguinal hernia arises through the deep inguinal ring lateral to the inferior epigastric vessel.



Solution to Question 2:

The muscle that is attached to the articular disc of the temporomandibular joint is the lateral pterygoid.

Lateral pterygoid muscle:

• Origin - Upper head arises from the infratemporal surface and infratemporal crest of the greater wing of the sphenoid bone. The lower head arises from the lateral surface of the lateral pterygoid plate.

• Insertion - It is inserted into the pterygoid fovea, the capsule of the temporomandibular joint, and the articular disc.

The action of lateral pterygoid muscle is depression, protrusion, and side-to-side movement of the jaw.

Solution to Question 3:

The above histology image shows the glomerulus.

Glomerulus is a rounded tuft of anastomosing capillaries lined by fenestrated endothelium. Blood enters the tuft through an afferent arteriole and leaves it through an efferent arteriole. It is supported by the mesangium containing mesangial cells and a non-cellular mesangial matrix.

The glomerular capillary wall is the filtering membrane and consists of fenestrated endothelial cells, basement membrane, and visceral epithelial cells called podocytes, which contain foot processes.

The image given below shows the histology of the glomerulus.



Option A: Leydig cells, also known as interstitial cells of Leydig, are present in the interstitial connective between seminiferous tubules.

The testis contains numerous seminiferous tubules separated by connective tissue in between. Seminiferous tubules are lined by several spermatogonia cells, which produce spermatozoa, and Sertoli (sustentacular) cells, which have a supportive function.

The connective tissue in between the tubules is made up of Leydig cells, collagen fibers, fibroblasts, macrophages, mast cells, blood vessels, and lymphatics. Leydig cells are large, round, or polyhedral cells with eccentric nuclei. They are lightly stained and often have a foamy appearance. They secrete testosterone upon stimulation by luteinizing hormone from the anterior pituitary.

The image below shows seminiferous tubules and Leydig cells.



Option B: Islets of Langerhans makes up the endocrine part of the pancreas.

Islets are characterized by secretory cells supported by reticulin fibers and contain numerous fenestrated capillaries. There is a delicate capsule around each islet. The islet cells are paler than the surrounding exocrine cells, as they have less rough endoplasmic reticulum.

The islet of Langerhans is mainly composed of three types of cells:

• α or A cells - located peripherally and secrete glucagon

• β or B cells - located centrally and produce insulin

• δ or D cells - scattered between beta cells and secrete somatostatin

The image given below shows the islets of Langerhans.



Option C: Hassall's corpuscles are small rounded structures present in the medulla of the thymus. Each corpuscle has a central core of degenerated epithelial cells and macrophages surrounded by concentrically arranged epithelial cells. The degenerated cells ultimately give pink staining hyaline mass. The function of the corpuscles of Hassall is still not known.

The image given below shows Hassall's corpuscle.



Solution to Question 4:

The site marked as '3' in the x-ray given above is the neck of the fibula. Injury at this site causes failure of dorsiflexion (foot drop) due to injury to the common fibular nerve (also know as common peroneal nerve)

The common fibular nerve is derived from the dorsal branches of the 4th and 5th lumbar and 1st and 2nd sacral ventral rami. It runs obliquely along the lateral side of the popliteal fossa to the fibular head. It lies between the biceps femoris and the lateral head of the gastrocnemius. It then curves laterally around the neck of the fibula and gets divided into superficial and deep fibular nerves.

	Superficial fibular nerve	Deep fibular nerve
Motor innervation	Peroneus longusPeroneus brevis	Extensor digitorum longusExtensor hallucis longusTibialis anteriorPeroneus tertius
Sensory innervation	Lateral aspect of the lower legDorsum of the toes except for the lateral side of the 5th toe and the first interdigital web-space	The first interdigital space







Injury to the common peroneal nerve (common fibular nerve) at the level of the neck of the fibula causes foot drop. Ankle dorsiflexion, toe extension, and eversion of the foot are lost, while the inversion and plantar flexion are usually preserved. Sensory innervation to the dorsum of the foot, including the first web space, is lost. The ankle reflex is usually preserved.

Solution to Question 5:

The image above shows a patent ductus arteriosus due to a defect in the left side of the 6th aortic arch.

Usually, in term infants, the ductus arteriosus closes within 72 hours of birth. Failure of the ductus arteriosus to close results in patent ductus arteriosus (PDA).

PDA usually presents with a hyperdynamic precordium, bounding peripheral pulses, wide pulse pressure, a machine-like continuous or systolic murmur, cardiomegaly, increased pulmonary vascular markings, hepatomegaly, increasing oxygen dependence, carbon dioxide retention, and renal failure.

In a majority of cases, PDA closes spontaneously. If spontaneous closure fails to occur, COX inhibitors like indomethacin or ibuprofen can be used. Surgical ligation is indicated in case of contraindications to medical management or failure of the same.

Other options

Option A: The 4th right aortic arch gives rise to the proximal part of the right subclavian artery.

Option B: The 4th left aortic arch gives rise to the arch of the aorta. Defects involving the 4th left aortic arch could cause a variety of aortic arch anomalies, such as interruption of the aortic arch, but not PDA.

Option C: The 6th right aortic arch forms the proximal part of the right pulmonary artery.

Solution to Question 6:

The structure marked A is the allantois. Its distal portion forms the urachus, which in turn gives rise to the median umbilical ligament.

The given image shows an embryo at the end of the third week.



Meckel's diverticulum is a congenital diverticulum arising from anti-mesenteric border of ileum. It is a remnant of the proximal part of the vitelline duct, which is made up of pluripotent cell lining. Meckel's diverticulum may harbor abnormal tissues, containing embryonic remnants of other tissue types, mostly gastric mucosa.

Solution to Question 7:

In the given scenario of a patient with portal hypertension, anastomosis between esophageal veins and left gastric veins can be seen.

In portal hypertension, increased pressure within the portal venous system may result in the dilation of portal venous tributaries and reversed flow at sites of portosystemic anastomoses.

Common sites of porto-systemic anastomoses and their clinical presentations are as follows:

Portal	Systemic	Clinical presentation
Left gastric vein	Distal esophageal veins	Esophageal and gastric varices
Superior rectal vein	Middle and inferior rectal veins	Rectal varices
Persistent tributaries of the left branch of the portal vein in ligamentum teres	Periumbilical branches ofepigastric and intercostal veins	Caput medusae

Portal	Systemic	Clinical presentation
Right branch of portal vein	Retroperitoneal veins draining into azygos, hemiazygos, lumbar, intercostal, and phrenic veins	Dilated retroperitoneal veins at risk during surgery or interventions
Omental and colonic veins	Retroperitoneal veins near hepatic and splenic ■exures	At risk during surgery

Solution to Question 8:

The cause of the pain in this patient complaining of pain in the parotid region is likely to be the greater auricular nerve.

The greater auricular nerve provides sensory innervation to the skin overlying the parotid gland and the ear lobule. Pain associated with swelling and inflammation in the parotid region could be transmitted through the greater auricular nerve. The parotid gland is covered by a parotid capsule that is derived from the investing layer of deep cervical fascia. Any acute inflammation of the gland such as acute sialadenitis results in pain in the preauricular region. The pain is mainly due to stretching of the capsule resulting in stimulation of the greater auricular nerve.

Other options:

Option A: The facial nerve is not the primary cause of pain in the parotid region due to swelling. The Hitzelberger sign(numbness in the posterosuperior meatal wall of the external auditory canal), associated with facial nerve involvement, is seen in acoustic neuroma and is not the cause of pain in this scenario.

Option D: Investing layer of deep cervical fascia is not the correct answer because the investing layer of deep cervical fascia itself does not transmit pain sensation. The investing layer of deep cervical fascia is a layer of connective tissue that envelops the neck and extends to cover the parotid gland, forming the parotid capsule. While inflammation or swelling in the parotid gland can stretch the capsule and cause pain, the actual transmission of pain sensation is mediated by sensory nerves, such as the greater auricular nerve, which innervates the skin over the parotid gland and the ear lobule.

The greater auricular nerve (Option B) is a more appropriate answer in this context, as it is responsible for carrying the pain sensation from the area of swelling and inflammation in the parotid region.

Option C: The auriculotemporal nerve can cause referred pain due to caries tooth, ulcerative lesions of the oral cavity, temporomandibular joint disorders, and sphenopalatine neuralgia. However, it is not the primary cause of pain in the parotid region due to swelling.

Anatomy INI-CET 2023

Question 1:

Which of the following vessels is involved in the condition affecting the region marked in red in the image shown below?



a) Superior Cerebellar Artery

b) Posterior Inferior Cerebellar Artery

c) Posterior Superior Cerebellar Artery

d) Cerebral artery

Question 2:

Which of the statements given below regarding the muscle shown in the image are correct?



a) 1, 2, 3

b) 1, 2, 4

c) Only 1

d) 2, 3, 4

Question 3:

Match the following:

a) 1-D, 2-C, 3-A, 4-B

b) 1-C, 2-A, 3-B, 4-D

c) 1-A, 2-B, 3-C, 4-D

d) 1-B, 2-A, 3-C, 4-D

Question 4:

Which of the following muscles are supplied by Superior Gluteal Nerve?

a) 1, 2, 4

b) 2, 3, 4

c) 1, 2, 3

d) 1, 3, 4

Question 5:

A 25-year-old male presented with symptoms of inability to flex the distal interphalangeal joint of the 4th and 5th digits. He was also not able to hold a piece of paper between his fingers. What is the likely site of injury?



a) A

b) B

c) C

d) D

Question 6:

Which of the following muscles form the pelvic diaphragm?

a) A, B

b) A, B, C, D

c) A, B, D

d) A,D

Question 7:

Anterior relations of the third part of the duodenum are all except:

a) Superior mesenteric vessels

b) Loops of jejunum

c) Root of mesentery

d) Fundus and body of Gall bladder

Question 8:

Arrange the following afferent columns in the floor of the fourth ventricle from medial to lateral.

a) 3, 1, 4, 2

b) 2, 4, 1, 3

c) 1, 2, 3, 4

d) 4, 3, 2, 1

Question 9:

Match the following muscles with their respective actions at the shoulder joint.

a) A-3, B-4, C-1, D-2

b) A-1, B-3, C-2, D-4

c) A-2, B-4, C-3, D-1

d) A-2, B-1, C-3, D-4

Question 10:

Given below is the image of a 33-hour chick embryo. Which of the following is derived from the marked structure?



a) Kidney

b) Dorsal root ganglion

c) Ribs

d) Parietal pleura

Question 11:

Which of the following is false regarding the anatomy of the adrenal glands?

a) The right adrenal gland is semi lunar in shape

b) The left adrenal gland is closely related with tail of pancreas and spleen

c) The adrenals are yellow on gross appearance

d) The adrenals are retroperitoneal in location

Question 12:

The sulcus located deep to the marked structure is?



a) Central sulcus

b) Stem of the lateral sulcus

c) Inferior frontal sulcus

d) Superior temporal sulcus

Question 13:

Given below is an illustration of a transverse section of the heart. Identify the marked structure.



a) Sub-aortic curtain

b) Posterior leaflet of mitral valve

c) Aortic valve leaflet

d) Anterior aortic sinus

Question 14:

Identify the structure marked as A.



a) Intra-pulmonary bronchus

b) Terminal bronchiole

c) Respiratory bronchiole

d) Alveolar sac

Question 15:

The fibres of the internal oblique muscle and transverse abdominus muscle contribute to which boundaries of the inguinal canal walls:

a) ii and iv

b) i, ii and iv

c) i and iii

d) iii and iv

Question 16:

Which of the following is a function of the marked layer?



a) Gluconeogenesis

b) Sodium and water retention

c) Increased sex hormones

d) Immune cell interaction

Question 17:

A patient's right hip drops when he takes the right foot off the ground. Which of the following would not be involved in the patient?



a) Superior gluteal nerve

b) Gluteus medius

c) Gluteus minimus

d) Sciatic nerve

Question 18:

Which of the following statements is incorrect about the anomaly shown below?

a) It occurs due to the persistance of prechordal plate

b) It occurs due to the defective migration of primordial cells to the genital ridge

c) It contains skin, cartilage and bone

d) It is a benign tumor of coccyx

Question 19:

Identify the arrow-marked structure.

a) Carina

b) Superior Vena Cava

c) Pulmonary Trunk

d) Arch of Aorta

Question 20:

What is the role of the marked structure in the given image?



a) Motor function

b) Reward-punishment

c) Judgement

d) Memory

Question 21:

Which of the following muscles are supplied by the lateral plantar nerve?

a) 2 and 5

b) 3 and 4

c) 1 and 3

d) 2 and 4

Question 22:

The marked structure sends efferents to which nucleus of the thalamus?



a) Pulvinar

b) Ventro-postero-lateral nucleus

c) Anterior nucleus

d) Lateral nucleus

Question 23:

On neurologic examination, a patient is not able to identify an object but is able to recognize the object from its texture, colour and pattern. Which of the following is likely to have been affected?

a) Occipital lobe

b) Lateral parietal lobe

c) Medial parietal lobe

d) Inferior temporal lobe

Question 24:

Apocrine glands are present in which of the following?



a) Face

b) Scalp

c) Palms and soles

d) Axilla

Question 25:

Which of the following is a tributary of the inferior vena cava?

a) Left inferior phrenic vein

b) Left supra-renal vein

c) Left gonadal vein

d) Right ascending lumbar vein

Answer Key

Question No.	Correct Option
1	b
2	a
3	d
4	c
5	c

6	a
7	d
8	b
9	c
10	c
11	a
12	b
13	a
14	b
15	a
16	b
17	d
18	a
19	c
20	a
21	a
22	c
23	d
24	d
25	a

Detailed Explanations

Solution to Question 1:

The given image is suggestive of lateral medullary syndrome/Wallenberg syndrome. This may result from a stroke affecting the Posterior Inferior Cerebellar Artery.

The clinical findings correspond to the neuroanatomical structures affected in the lateral portion of the medulla:

• Descending tract and nucleus of CN V (trigeminal)- loss of pain and sensation over the ipsilateral half of the face.

• Inferior cerebellar peduncle – ipsilateral ataxia (gait and limb)

• Lateral spinothalamic tract – loss of pain and temperature sensation over the contralateral half of the body.

• IX and X CN - ipsilateral paralysis of palate, vocal cord, diminished gag reflex giving rise to dysphagia, dysarthria.

• Vestibular nucleus - Nystagmus, diplopia, vertigo, nausea

• Descending sympathetic tract – ipsilateral Horner's syndrome (miosis, ptosis, and anhidrosis).



Solution to Question 2:

The muscles marked in the above image are rhomboids minor and major. These muscles cause retraction and elevation of the scapula and they rotate the scapula tilting the glenoid cavity inferiorly.

Rhomboids minor and major are innervated by the dorsal scapular nerve. They are supplied by the dorsal scapular artery or the deep branch of the transverse cervical artery.

Muscles involved in movements of the scapula are:

• Protraction (forward movement): Serratus anterior and pectoralis minor are prime movers. Rhomboid muscles control the rate and range of movements. The upper part of the latissimus dorsi acts like a strap across the inferior scapular angle in protraction.

• Retraction (backward movement): Middle trapezius and rhomboids.

• Elevation: Trapezius, Levator scapulae and rhomboids.

• Depression: Usually gravity alone is sufficient for the depression of the scapula. When required, a part of the serratus anterior, pectoralis minor, and trapezius are used.

Appendicular group of back muscles and their innervations:

Muscle	Nerve Supply	Action
Trapezius	Accessory nerve	Elevation of the scapula, rota tion of scapula during abduct ion of humerus above horizo ntal; middle fibers retract sca pula; lower fibers depress the scapula

Muscle	Nerve Supply	Action
Latissimus dorsi	Thoracodorsal nerve	Adduction, medial rotation, a nd extension of the arm at th e glenohumeral joint
Levator scapulae	Dorsal scapular nerveBranch es directly from anterior rami of C3 and C4	Elevation of the scapula
Rhomboid major and m inor	Dorsal scapular nerve	Elevation and retraction of th e scapula





Solution to Question 3:

The correct match is as follows:

Chorda tympani nerve which is a branch of facial nerve carries taste from anterior 2/3rd of the tongue.

The trapezius is innervated by the spinal accessory nerve (CN XI). It causes retraction of scapula, shrugging of the shoulder and overhead abduction (along with serratus anterior). Paralysis of trapezius causes drooping of the shoulder and slight winging of the scapula (the superior angle of scapula becomes more prominent).

The glossopharyngeal nerve carries taste sensation from the posterior 1/3rd of the tongue.

The mandibular nerve (V3) is responsible for chewing movements as it gives motor supply to muscles of mastication. It contains both sensory and motor fibers.

• Sensory from

• Skin of cheek, over mandible and side of head

• Teeth of lower jaw and temporomandibular joint

• Mucous membrane of mouth and anterior part of tongue

• Motor to

• Muscles of mastication (pterygoids, masster and temporalis)

• Tensor tympani

• Anterior belly of digastric

• Mylohyoid

1. Facial nerve	B. Taste from anterior two th ird of the tongue
2. Spinal accessory nerve	A. Shrugging of shoulder
3. Glossopharyngeal nerv e	C. Taste from posterior one-t hird of the tongue
4. Mandibular nerve	D. Chewing

Solution to Question 4:

Gluteus Medius, Gluteus Minimus, and Tensor fascia lata are supplied by Superior Gluteal Nerve. Piriformis is supplied by the nerve to the piriformis.

The dorsal branches of the L4, L5, and S1 rami give rise to the superior gluteal nerve. It is accompanied by superior gluteal vessels. This nerve leaves the pelvis via the greater sciatic foramen above piriformis and divides into superior and inferior branches. It supplies gluteus medius, gluteus minimus, and tensor fasciae latae.

The dorsal divisions of the S1 and S2 rami give rise to the nerve to the piriformis.

Gluteus medius and gluteus minimus act as the abductors of the hip joint. Stabilizing the pelvis on the femur during walking is done by these muscles. It is demonstrated by asking the subject to stand on one limb. If the ipsilateral abductors are normal, the pelvis remains in a horizontal

position. If the ipsilateral abductors are weak, then the pelvis tilts towards the opposite (unsupported) side. This is called positive Trendelenburg's sign.





Solution to Question 5:

The given clinical situation with a positive card test and an inability to flex the distal interphalangeal joint of the 4th and 5th digits, suggests an ulnar nerve injury. The most likely site of injury would thus be medial epicondyle: site C

Injury at the medial epicondyle damages the ulnar nerve, leading to paralysis of flexor carpi ulnaris (loss of ulnar deviation) and medial half of flexor digitorum profundus (loss of distal interphalangeal joint flexion). It leads to hypothenar paralysis (lead to atrophy) and sensory loss. Also, flexor digitorum superficialis (supplied by median nerve) and extensor digitorum (supplied by the radial nerve) are still functional. Hence complete paralysis of the 3rd and 4th digits

is not seen. The positive card test is due to the weakness of palmar interossei leading to loss of adduction.

Ulnar nerve branches and supply:

Superficial branch supplies palmaris brevis, skin on the palmar surface of medial 1½ fingers. The deep branch runs within the concavity of deep palmar arch and supplies all intrinsic muscles of the hand (except first 2 lumbricals and thenar muscles).

Injury to the Ulnar nerve:

Other tests seen in ulnar nerve lesions are as follows:

• Claw hand deformity

• Froment's sign positive

• Wartenberg sign

• Palmaris brevis sign

	High lesion(Elbow)	Low lesion(Wrist)
Site of injury	Fracture/dislocation of theme dial epicondyleCubital tunnel syndromeCompression betw een the two heads of FCUTar dy ulnar nerve palsy (Valgus deformity)	Wrist laceration/ slashingGuy on's canalsyndrome
Motor loss	Claw-handdeformityHypothe nar & interosseous wastingLo ss of:• Abduction & adduction of fingers(Card test- Positive) • Adduction of thumb(Frome nt sign-Positive)• Flexion of MCP jointsWeakness of wrist flexion (radial deviation)	Claw-handdeformity(more pr onounced)Hypothenar & inte rosseous wastingLoss of:• Ab duction & adduction of finger s(Card test- Positive)• Adduct ion of thumb(Froment sign-P ositive)
Sensory loss	Over palmar & dorsal aspect of themedial thirdof thehand Over palmar & dorsal aspect of themedial 11/2fingers	No sensory lossover palmar & dorsal aspect of the medial third of the handOver palmar & dorsal aspect of themedial 11/2fingers

The image below shows a positive card test.



Positive Card Test
In the weakness of palmar interossei, there is a weakness of finger adduction. So the patient is unable to hold card firmly between fingers.

Wartenberg's sign: (Weakness of palmar interossei), Sensory loss Palmar and dorsal surface of the medial third of hand and medial 1 and a half fingers.



Other options:

• Option A: Radial groove - Radial nerve

• Option D: Neck of humerus - Axillary nerve

• Option B: Lateral epicondyle – May lead to tardy ulnar nerve palsy after fracture but the incidence is less.

Solution to Question 6:

Iliococcygeus and pubococcygeus are parts of the pelvic diaphragm, which is also known as the levator ani.

The levator ani muscle is a complex unit that consists of 3 muscle components:

• Pubococcygeus

• Pubovaginalis - elevates the urethra and hence contributes to urinary continence

• Puboperinealis - draws the perineal body towards the pubic symphysis

• Puboanalis - elevates the anus and keeps the urogenital hiatus narrowed

• Puborectalis

• Iliococcygeus

Levator ani muscles are the most important muscles in the pelvic floor and maintain a constant state of contraction. They provide a stable floor and support the weight of the abdominopelvic contents against intra-abdominal forces. Weakening of levator ani and pelvic fascia from stretching or tearing during childbirth, may decrease support for the vagina, bladder, uterus, or rectum or alter the position of the neck of the bladder and urethra.

Other options:

The obturator internus forms the lateral wall of the ischioanal fossa and is not a part of the levator ani.

The deep transverse perineal muscle is the content of the deep perineal space and contributes to the formation of the perineal body.



Solution to Question 7:

Anterior relations of the third part of the duodenum include the Superior mesenteric vessels, the Root of the mesentery, and Loops of the jejunum.

Relations of the 3rd part of duodenum:

• Posterior relations of 3rd part of duodenum include right ureter, right psoas major, right gonadal vessels, inferior vena cava and abdominal aorta (at the origin of the inferior mesenteric artery).

• Anterior relations: Loops of jejunum, Root of mesentery and Superior mesenteric vessels.

• Superiorly related to head of the pancreas and Inferiorly to loops of jejunum.

Note: Root of mesentery (containing Superior mesenteric vessels) lies anterior to the 3rd part of Duodenum. Hence, posterior relations of 3rd part of duodenum are the same as structures crossed by root of mesentery.

Solution to Question 8:

From medial to lateral, the afferent columns in the floor of the fourth ventricle are general visceral, special visceral, general somatic, special somatic.

Embryologically, the continuous addition of neuroblasts to the mantle layer of the neural tube produces thickenings ventrally and dorsally - producing the basal and alar plates respectively. A longitudinal groove, called sulcus limitans, marks the boundary between the two.

Basal Plate	Alar plate
Ventralthickening	Dorsalthickenings
Give rise tomotor nuclei-general visceral efferent nuclei, special visceral efferent of CN IX, X, and XII; and somatic efferent of XII	Give rise tosensory nucleiCN V, VII, VIII, IX, X, inferior olivary nucleus, and mesencephalic nucleus of CN V
Forms the motor part of the spinal cord (ventral horns)	Forms the sensory part of the spinal cord (dorsal horns).The rhombic lip develops into the cerebellum and its nuclei.





Solution to Question 9:

Muscles	Actions
Supraspinatus	Steadies the head of the humerus during movements,abduction of shoulder joint from 0 -15°
Deltoid	Abduction, anterior fibers - flexors and medial rotators, posterior fibers - extensors and lateral rotators
Infraspinatus	Lateral rotatorof the arm
Teres minor	Lateral rotator of the arm
Teres major	Adduction and medial rotation of the arm
Latissimus Dorsi	Adduction, medial rotation, andextension of the arm
The long head of the triceps	Extension of elbow joint
Pectoralis major	Adduction, medial rotation, andflexion of the arm



Solution to Question 10:

The structure marked in the given image is the para-axial mesoderm. The ribs are derived from the para-axial mesoderm.

During the third week of embryological development, the process of gastrulation occurs, which refers to the formation of the three germ layers: ectoderm, mesoderm and endoderm. This is followed by the formation of the notochord, which is the axial structure, which induces the formation of the neural tube above it. During the formation of the neural tube, some cells at the lateral border dissociate and form the neural crest cells, which is sometimes referred to as the fourth germ layer.





After the formation of the notochord, which is the axial structure, the mesoderm on either side can be seen to be organized into three layers as follows:

1. Para-axial mesoderm - Thickened plate of the mesoderm close to the midline, on either side of the notochord. This forms the somitomeres and somites, which have the following derivatives:

• Dermatome: Forms the segmental dermis of the skin

• Myotome: Forms the skeletal muscle

• Sclerotome: Forms the vertebral column and ribs

2. Intermediate mesoderm - Longitudinal strip of mesoderm that connects the para-axial with the lateral plate mesoderm. The urinary (Option A) and genital systems are derived from the intermediate mesoderm.

3. Lateral plate mesoderm - A thinner layer of mesoderm that lies laterally. This layer develops a large cavity called the intraembryonic coelom that gives rise to pericardial, pleural (Option C), and peritoneal cavities.

The neural crest cells also contribute to the formation of structures as follows:

In the trunk region, they form melanocytes, sensory ganglia (Option B), sympathetic and enteric neurons, Schwann cells, and adrenal medulla cells.

In the cranial region, they contribute to the craniofacial skeleton, neurons for cranial ganglia, satellite glial cells, and melanocytes.

Solution to Question 11:

Option A is false as the right adrenal gland is pyramidal in shape while the left adrenal gland is semi-lunar/crescent shaped.

The adrenal glands, also called the supra-renal glands, are endocrine glands which are located at the upper pole of each kidney. They are grossly yellow in appearance and are retroperitoneal in

location. Both glands are surrounded by areolar tissue and perirenal fascia. The following are the key differences between the right and the left adrenal glands and their relations:

Each gland is supplied by:

• Superior suprarenal artery - branch of the inferior phrenic artery.

• .Middle suprarenal artery - branch of the abdominal aorta.

• Inferior suprarenal artery - branch of the renal artery.

Each gland has one draining vein.

• The right suprarenal vein drains into the IVC.

• The left suprarenal vein drains into the left renal vein.

	Right Adrenal Gland	Left Adrenal Gland
Shape	Pyramidal or triangular	Semi-lunar or crescenteric
Parts	Apex and base	Upper end and lower end
Superior relation	Bare area of liver	Spleen
Inferior relation	Right kidney	Left kidney and left supra-re nal vein
Anterior relation	IVC, bare area of liver	Stomach and tail of pancreas with splenic artery
Posterior relation	Right crus of diaphragm, righ t kidney	Left crus of diaphragm, left k idney
Medial relation	Coeliac ganglion	Coeliac ganglion
Lateral relation	Liver	Stomach



Solution to Question 12:

The area marked by an arrow in the given image is pterion and the stem of the lateral sulcus lies under it.

The pterion is an H-shaped sutural junction between four bones of the skull:

• Frontal bone

• Squamous part of the temporal bone

• Greater wing of the sphenoid bone

• Parietal bone.

The surface of the cerebrum consists of folds called gyri separated by fissures called sulci. The lateral sulcus, also called the Sylvian Fissure, is present on the infero-lateral surface of each cerebral hemisphere and separates the frontal lobe from the temporal lobe. The lateral sulcus consists of a stem which divides into three rami, which are, the anterior horizontal, anterior ascending, and the posterior rami.

The structures lying deep to the pterion are:

• The anterior branch of the middle meningeal artery

• The middle meningeal vein

• The stem of the lateral sulcus (Sylvian fissure)

The bone in this area is thin and an injury to this area may lead to rupture of the middle meningeal artery leading to extradural hematoma.





Solution to Question 13:

The marked structure is the sub-aortic curtain. It is also called the trigonum fibrosum sinistrum.

The fibrous skeleton of the heart is formed by fibrous tissue that is continuous between the atrio-ventricular and arterial orifices. They exist to keep the valves competent, allow attachment of cardiac musculature, and act as an insulating layer for the conducting system of the heart. The fibrous rings of importance are:

• The atrioventricular fibrous rings - existing in a figure of 8 configuration, connecting the atria, ventricles and the membranous part of the interventricular septum.

• The trigonum fibrosum dextrum - between the atrioventricular rings and the aortic ring.

• The trigonum fibrosum sinistrum - between the mitral valve and the aortic valve.



Solution to Question 14:

The arrow-marked structure in the given image is the terminal bronchiole. It can be identified by the absence of cartilage, which would be present in the bronchus, and the presence of cilia and stratified columnar epithelium, which would not be seen in respiratory bronchioles and alveolar sac.

The following is a summary of the intrapulmonary passages:

Trachea → Main Bronchi → Lobar Bronchi → Segmental Bronchi → Terminal Bronchiole → Respiratory Bronchiole → Alveolar Duct → Alveolar Sac → Alveoli

Bronchi and bronchioles can be distinguished by the presence of cartilage and glands in the bronchi which are not present in the bronchioles.

Terminal bronchioles represent the most distal parts of the conducting passage. Each terminal bronchiole ends by dividing into respiratory bronchioles. Terminal bronchioles contain cilia and stratified columnar epithelium, whereas respiratory bronchioles contain few or no cilia, and simple cuboidal epithelium. Another differentiating point between the two is that terminal bronchioles would not communicate with the alveoli but respiratory bronchioles would have a communication with the alveoli.

Each respiratory bronchiole ends by dividing into a few alveolar ducts. Each alveolar duct ends in a passage, the atrium, which leads into a number of rounded alveolar sacs. Each alveolar sac is studded with a number of air sacs or alveoli.



A: Terminal Bronchiole, B: Respiratory Bronchiole, C: Alveolar Sac, D: Alveolar Duct

The coloured strips in the following illustration represent the location of different cells/structures in the trachea and bronchial tree.



Note - The image in the question has been updated for better clarity

Solution to Question 15:

The fibres of the internal oblique muscle and transverse abdominus muscle contribute to the roof and posterior wall of the inguinal canal walls.

The boundaries of the inguinal canal walls are as follows:

• Anterior wall: Formed by the external oblique aponeurosis.

• Roof: Formed by the arched fibres of the internal oblique and transversus abdominis muscles.

• Floor: Formed by the grooved upper surface of the Abdomen and Pelvis inguinal ligament.

• Posterior wall: Consists of the transversalis fascia and conjoint tendon (the lowest fibres of the transverses abdominis muscle fuse with the lowest fibres of the internal oblique to form the conjoint tendon).

Solution to Question 16:

The image shows a histological section of the adrenal gland and the layer marked is the outermost layer, ie, the zona glomerulosa, which secretes the mineralocorticoid - aldosterone, responsible for sodium and water retention.

Based on structure and function, the adrenal cortex has three zones, which from outer to inner, are:

• Zona glomerulosa: Secretes mineralocorticoids (aldosterone) - which acts in response to the activation of the renin-angiotensin-aldosterone-system (RAAS) and acts at the distal convoluted tubule and the collecting ducts favouring sodium and water retention and potassium excretion.

• Zona fasiculata: Secretes glucocorticoids (cortisol and corticosterone) - which has multiple functions that include metabolic effects which induce hyperglycemia inducing gluconeogenesis (Option A) and mediates anti-inflammatory and immunosuppresive effects by interacting with the cells of the immune system (Option D).

• Zona reticularis: Secretes sex steroids (dehydroepiandrosterone, androstenedione, estrogens, and glucocorticoids) (Option C)



Solution to Question 17:

A patient with dropping of the right hip on lifting the right foot off the ground, has a positive Trendelenburg's sign. The sciatic nerve would not be involved.

A positive Trendelenburg's sign is seen in the paralysis of the gluteus medius and gluteus minimus muscles, both of which are supplied by the superior gluteal nerve. In the scenario mentioned, since the patient has dropping of the hip on the right side, the structures on the left side would be affected.

Mechanism of actions of gluteus medius and minimus:

• When the leg on one side is lifted off the ground, making it the unsupported side, gluteus medius and minimus muscles on the opposite side, which is the supported side, exert traction on the hip bone by acting from the femur. This action results in slight raising of the pelvis on the unsupported side, which helps maintain the trunk in the upright position.

• When the gluteus medius and minimus are paralyzed, the pelvis on the unsupported side sinks when the patient lifts the foot off the ground. This causes the patients to have a characteristic lurching gait.



Solution to Question 18:

The image shows a sacrococcygeal teratoma, which occurs due to the persistence of pluripotent cells of the primitive streak in the sacrococcygeal region.

During the third week of embryological development, the process of gastrulation occurs, which refers to the formation of the three germ layers. These germ layers are formed from the primordial germ cells of the epiblast, which are pluripotent. The process begins with the formation of the primitive streak on the epiblast, which develops into the primitive node in the cranial region, and a primitive pit. The cells from the epiblast, migrate towards the primitive streak and invaginate into the primitive pit and organizes itself into cells arranged in three layers, which eventually develop into ectoderm, mesoderm, and endoderm. The prechordal plate simply refers to the cells that have invaginated in the cranial region.



Sacrococcygeal teratomas occur due to remnants of pluripotent cells from the primitive streak due to defective migration of the cells (Option B). As these cells are from the epiblast, teratomas contain tissues from all germ layers which may include skin, hair, cartilage and bone (Option C). These tumors are benign in nature (Option D), but may rarely become malignant. If they occur in the cranial end of the primitive streak, they form oro-pharyngeal teratomas (as shown below), whereas if they occur in the caudal end of the primitive streak in the coccygeal region, they are called sacro-coccygeal teratomas.

Solution to Question 19:

The arrow marks the bifurcation of the pulmonary trunk.

The image shows a cross section of a cadaver at the T5 vertebral level. This corresponds anteriorly to the level of the sternal angle, and is also the level of division between the superior and inferior mediastinum.

At this level, the other structures visible are:

• The origin and termination of the arch of aorta. The descending aorta can be identified to the left of the oesophagus. The arch of the aorta (Option D) would be visible at the T3-T4 cross sectional level.

• The superior vena cava (Option B), just lateral to the origin of the aorta, which can visually differentiated from the aorta based on the wall thickness. At this level, it penetrates the right atrium.

• The bifurcation of the pulmonary trunk.

• The left and right main bronchi. The carina (Option A) is not visible as the trachea bifurcates more proximally at the T4 level.



Solution to Question 20:

The marked structure is the subthalamic nucleus, which is a part of the basal ganglia and is concerned with motor function.

The basal ganglia refers to the mass of grey matter located within the cerebral white matter and is responsible for a number of coordinated functions including the control of extra-pyramidal pathways for motor coordination. There are two major pathways involved which are, the direct and indirect pathways.

The direct pathway stimulates the cortex to produce movement, whereas the indirect pathway inhibits it. The striatal activity in the direct pathway is mediated by D1 dopamine receptors and in the indirect pathway by D2 dopamine receptors. The subthalamic nucleus is a part of the indirect pathway. A lesion at subthalamic nuclei leads to loss of inhibitory effect of the indirect pathway resulting in dopaminergic overactivity responsible for features of hemiballismus.



Other Options:

Option B: Reward-punishment is associated with the prefrontal area and the dopaminergic connections linked to it from the ventral tegmental area.

Option C: Judgement is controlled by the frontal lobe.

Option D: Memory is controlled by the hippocampus.



Solution to Question 21:

From the given list, the muscles supplied by the lateral plantar nerve are the flexor digitorum accesorius and the first interossei.

The sole of the foot is supplied by the medial and lateral plantar nerves which are both branches of the tibial nerve.

All the muscles of the sole of the foot are supplied by the lateral plantar nerve, except for the following four muscles which are supplied by the medial plantar nerve:

- Abductor hallucis
- Flexor digitorum brevis
- Flexor hallucis brevis
- First lumbrical

The sensory supply of the sole of the foot is as shown in the illustration below.



The muscles of the sole of the foot are arranged in 4 layers as shown below:

1st layer: It consists of the abductor hallucis, flexor digitorum brevis and abductor digiti minimi.



2nd layer: It consists of flexor accessorius and the four lumbrical muscles,the tendons of flexor hallucis longus and flexor digitorum longus run in the same plane as the muscles of the second layer.



3rd layer: It consists of the shorter intrinsic muscles of the toes, which are flexor hallucis brevis, adductor hallucis and flexor digiti minimi brevis.



4th layer: It consists of plantar and dorsal interossei and the tendons of tibialis posterior and fibularis longus.



Solution to Question 22:

The marked structure in the given coronal section of the brain is the fornix. The fornix carries associations from the hippocampus to the anterior nucleus of the thalamus via the mamillary bodies.

The Papez circuit is a part of the limbic system concerned with the control of emotional behaviour and memory.

It consists of the following structures, which can be remembered using the given mnemonic:

• Hippocampus (He)

• Mamillary body (Man)

• Anterior thalamus (Ate)

• Cingulate gyrus (Cat)

(Mnemonic-He Man Ate Cat)



Solution to Question 23:

The clinical scenario suggests that the patient has visual agnosia and the likely site of involvement would be the inferior temporal lobe.

• Agnosia refers to the inability to visually identify objects and is also called optic aphasia. According to De Jong, visual agnosia is further classified into:

• Apperceptive visual agnosia - the visual image appears distorted to the patient and hence they are unable to recognise it. This occurs due lesions in the parieto-occipital region

• Associative visual agnosia - inability to visually identify objects in the absence of visual impairment, with ability to identify the object using other sensory modalities (as suggested in the scenario). This occurs due to lesions of occipito-temporal junction, which is present in the inferior temporal lobe. Prosopagnosia is a type of associatetive visual agnosia which refers to the inability to visually identify faces.

The occipital lobe acts as the primary visual area and a lesion in this region would result in cortical blindness and achromatopsia, which is the inability to visualise colour. In the given scenario, as the patient is also able to identify colour, it indicates that the occipital lobe (Option A) is unaffected and (hence inferior temporal lobe is the better answer).

Stereognosis is the ability to identify an object by a perception of the form and nature of an object without looking at it. In the given scenario, the patient is able to recognize the object from its texture and pattern which indicates that stereognosis is intact. Loss of this ability is known as astereognosis which occurs in lesions of the superior parietal lobule (Options B and C) and dorsal column pathways.

Solution to Question 24:

Apocrine glands are present in the axilla.

Exocrine glands can be classified into three subtypes, based on the manner in which their secretions are poured out of the cells:

• Apocrine

• Merocrine

• Holocrine

In apocrine glands, the apical parts of the cells are shed off to discharge the secretion. Examples of apocrine glands are :

• Secretion of milk fat by mammary glands

• Atypical sweat glands in the axilla, perianal region, periumbilical skin, mons pubis.

• Ceruminous glands of the external auditory meatus are modified apocrine glands

Apocrine glands are large glands found in the dermis or subcutaneous fat. They are a type of sweat gland. It consists of a basal secretory coil followed by a straight duct that opens into the pilary canal above the duct of the sebaceous gland. In hairless skin, it directly opens onto the skin surface.

Apocrine activity is minimal before puberty, after which it is androgen-dependent and responsive to emotional stimuli. It is controlled by adrenergic nerves and is sensitive to adrenaline and noradrenaline. The secretion is initially sterile and odorless, but it undergoes bacterial decomposition to generate odorous and musky compounds. Obstruction of apocrine sweat ducts and associated upper hair follicles in the axillae, breast areolae, and pubic region, mainly in women, is thought to cause Fox–Fordyce disease. It is also implicated in the causation of a painful, occasionally debilitating condition called hidradenitis suppurativa.

The given image shows an illustration of apocrine secretion.







Solution to Question 25:

The left inferior phrenic vein is a tributary of the inferior vena cava (IVC).

From inferior to superior, the following vessels are direct tributaries of the IVC:

- Common iliac veins (left and right common iliac veins merge to form the IVC)
- Median sacral vein
- Four pairs of lumbar veins
- Right gonadal vein
- Renal veins (both left and right)
- Right supra-renal vein

• Hepatic veins

• Inferior phrenic veins (both left and right)

Other Options:

Option B: The left supra-renal vein drains into the left renal vein.

Option C: The left gonadal vein drains into the left renal vein.

Option D: The right ascending lumbar vein joins the subcostal vein on the right side to form the azygos vein. The left ascending lumbar vein joins the subcostal vein on the left side to form the hemi-azygos vein.



Anatomy INI-CET 2024

Question 1:

Which of the following joints is not formed by the structure shown in the picture?



a) Synchondrosis

b) Syndesmosis

c) Symphysis

d) Plane synovial joint

Question 2:

The spinal sensory nucleus of the trigeminal nerve has 2nd order neurons to carry which sensation?

a) Proprioception

b) Two point discrimination

c) Pain

d) Vibration

Question 3:

Which of the following is not seen in the floor of the inferior horn of the lateral ventricle?

a) Fimbria

b) Collateral eminence

c) Hippocampus

d) Tail of caudate nucleus

Question 4:

Which of the following marked areas in the histopathological specimen shows calcification?



a) 1

b) 2

c) 3

d) 4

Question 5:

What are the structures passing between the muscles marked as 1 and 2?



©MARROW

a) Recurrent laryngeal nerve and inferior thyroid vessels

b) Internal laryngeal nerve and superior laryngeal artery

c) Auditory tube and levator palitini

d) Glossopharyngeal nerve and stylopharyngeus muscle

Question 6:

At what vertebral level does the part marked in the image lie in a normal anatomical position?



©MARROW

a) S3 spine

b) Upper border of L5

c) Tip of coccyx

d) Disc space between L4 and L5

Question 7:

The nerve related to the area marked in the picture has its root value from?



a) Ventral horn of C4

b) Intermediate Horn of T2-T6

c) Open part of Medulla Oblongata

d) Dorsal root ganglia of T1 and T2

Question 8:

The area marked by the arrow is supplied by which of the following arteries?



©MARROW

a) Diagonal artery

b) Right Marginal artery

c) Anterior interventricular artery

d) Left circumflex artery

Question 9:

What level of the vertebral column does the given cross-sectional image correspond to?



a) T12

b) T9

c) T10

d) T11

Question 10:

The structure marked as 'X' in image A has a direct connection with which of the following structures marked in image B?



a) 1

b) 2

c) 3

d) 4

Question 11:

Which of the following statements is true concerning the anatomy of the breast?

a) Inner quadrants contain most breast tissue

b) Ligaments of cooper are found only in lower quadrants

c) Tail of Spence extends across anterior axillary fold

d) Breast is made of 10 lobes

Question 12:

A professor underwent a left parotidectomy. Post surgery she complains of left lower lip paralysis. Which structure is involved?

a) Facial nerve - Main trunk

b) Facial nerve- Cervical branch

c) Facial nerve-Temporal branch

d) Parotid duct

Question 13:

Identify the normal anatomical variation given in the following CT image.

a) Tracheal diverticulum

b) Esophageal duplication

c) Aberrant right subclavian artery

d) Right aortic arch

Question 14:

Identify the marked structure

a) Superficial temporal artery

b) Posterior auricular artery

c) Middle meningeal artery

d) Occipital artery

Question 15:

The marked structure serves as a surface landmark for which anatomical structure?



a) Pineal gland

b) Wenicke's area

c) Stem of the lateral sulcus

d) Internal carotid artery siphon

Question 16:

A 60-year-old male recovering from facial nerve palsy complains of food getting stuck between the cheek and teeth while eating. What is the muscle affected in this case?

a) Buccinator

b) Levator labii superioris

c) Zygomaticus major

d) Orbicularis oris

Question 17:

Which of the following is not a derivative of neural crest cells?

a) Ciliary muscles

b) Retinal pigment epithelium

c) Ciliary ganglion

d) Melanocytes

Question 18:

The cranial component of the neural crest cells leads to the formation of?

a) SA node

b) AV cushion

c) Conotruncal septum

d) Chordae tendinae

Question 19:

A 15-year-old boy underwent splenectomy for immune thrombocytopenic purpura. During the procedure, the surgeon noticed bleeding while separating the spleen from the stomach. Which vessels are most likely implicated in this bleeding?

a) Left gastric vessels

b) Short gastric vessels

c) Right gastro-epiploic vessels

d) Splenic vessels

Question 20:

Identify the marked structure

a) Primitive streak

b) Notochord

c) Neural groove

d) Secondary neural plate

Question 21:

Which of the following is the first to be formed by the migration of the epiblast along the primitive streak?

a) Endoderm

b) Cardiac mesoderm

c) Epidermis

d) Neural groove

Question 22:

During walking, the hip bone on the side of the suspended leg is raised by the action of which muscle present in the supporting leg?

a) Gluteus Medius

b) Gluteus Maximus

c) Obdurator Internus

d) Quadratus Femoris

Question 23:

Identify the marked structure?



a) Oculomotor nerve

b) Abducens nerve

c) Trochlear nerve

d) Ophthalmic division

Question 24:

Injury to the posterior interosseous nerve leads to?

a) Loss of extension of digits

b) Loss of extension of the wrist

c) Loss of supination

d) Wrist drop

Answer Key

Question No.	Correct Option
1	a
2	c
3	d
4	c
5	d
6	c
7	a
8	c
9	c
10	a
11	c
12	b
13	c
14	a
15	c
16	a
17	b
18	c
19	b
20	a
21	a
22	a
23	a
24	a

Detailed Explanations

Solution to Question 1:

The provided image depicts a thoracic vertebra, which does not participate in forming a synchondrosis-type joint. The example for a synchondrosis is the costochondral joint formed between the ribs and the costal cartilage.

A vertebral bone participates in the following joints :

• Intervertebral joint - secondary cartilaginous joint (symphysis type)

• Costovertebral joint - plane synovial joint

• costotransverse joint - plane synovial joint

• Facet joints or zygapophysial joints - synovial joint

A syndesmosis is a type of complex fibrous joint between two bones connected by ligaments and a strong membrane[1] with slight movement allowed.The vertebra is connected by a anterior and posterior ligament where some degree of syndesmosis type of joint is seen.

Structurally, joints are classified as fibrous joints, cartilaginous joints, and synovial joints.

Type	Subtypes
Fibrous joints	SuturesSyndesmosisGomphosis
Cartilaginous joints	Primary cartilaginous (synchondrosis)Secondary cartilaginous (symphysis)
Synovial joints	Ball-and-socket or spheroidal jointsSellar or saddle jointsCondylar or bicondylar jointsEllipsoid jointsHinge jointsPivot or trochoid jointsPlane joints

Solution to Question 2:

The trigeminal nerve (cranial nerve V) is responsible for sensory input from the face, as well as motor functions such as chewing. Its sensory pathway involves three branches: the ophthalmic (V1), maxillary (V2), and mandibular (V3) divisions. The first-order neurons for each division originate from different sensory ganglia:

1.Ophthalmic Division (V1):

First-order neurons: Cell bodies are located in the trigeminal ganglion (also called the semilunar or Gasserian ganglion).

2. Maxillary Division (V2):

First-order neurons: Cell bodies are located in the trigeminal ganglion.

3. Mandibular Division (V3):

First-order neurons: Cell bodies are located in the trigeminal ganglion.

These first-order neurons carry sensory information from the face to the brainstem. Upon entering the brainstem, the fibers synapse with second-order neurons in specific nuclei:

1. Principal Sensory Nucleus (Main Sensory Nucleus):
   a. Receives input from all three divisions of the trigeminal nerve.
   b. Located in the pons.
   c. Second-order neurons here carry discriminative touch and proprioceptive information.
2. Spinal Trigeminal Nucleus (Spinal Nucleus of Trigeminal):
   a. Receives input mainly from the maxillary and mandibular divisions.
   b. Located in the medulla.
   c. Second-order neurons here carry pain and temperature sensations.

After synapsing in these nuclei, the sensory information is further relayed to higher brain centers for processing and perception.

Solution to Question 3:

The tail of the caudate nucleus is not seen on the floor of the inferior horn of the lateral ventricle. It is present on the roof of the inferior horn of the lateral ventricle.

The inferior horn is the largest compartment of the lateral ventricle and extends up to 2.5 cm from the temporal pole. The relations are:

Roof - Tapetum of corpus callosum laterally and tail of caudate nucleus medially.

Floor (lateral to medial) - Collateral eminence produced by collateral fissure, hippocampus.

The floor of the temporal horn consists of the hippocampus medially and the collateral eminence, formed by the infolding of the collateral sulcus, laterally. The inferior part of the choroidal fissure, located between the fimbria of the fornix and the inferolateral aspect of the thalamus, lies in the medial wall.

The lateral ventricle is divided into the body and 3 horns. The locations of the parts are:

- Body - parietal lobe
- Anterior horn - frontal lobe
- Posterior horn - occipital lobe
- Inferior horn - temporal lobe.



The anterior horn extends from the genu of the corpus callosum up to the interventricular foramen. The relations of the anterior horn are:

• Anteriorly - Genu and rostrum of the corpus callosum

• Roof - Anterior part of the body of the corpus callosum

• Floor and lateral wall - Caudate nucleus

• Medial wall - Septum pellucidum and anterior column of fornix

The body or central part extends from the interventricular foramen to the splenium of the corpus callosum. The relations of the body are:

• Roof - Body of the corpus callosum

• Medial wall - Septum pellucidum

• Floor (from lateral to medial) - Caudate nucleus, stria terminalis, thalamostriate vein, thalamus, choroid plexus, and body of the fornix

The posterior horn extends from the splenium to the occipital pole. The relations are:

• Roof and lateral wall - Tapetum of corpus callosum

• Medial wall - Forceps major and calcar avis (calcarine sulcus

Note: There is no lateral wall for the lateral ventricle.



Solution to Question 4:

The given specimen is that of an epiphyseal cartilage, where zone 3 corresponds to the zone of calcification.

Epiphyseal cartilage is divided into five zones :

• The resting zone consists of hyaline cartilage without morphological changes in the cells (option 1)

• In the proliferative zone, chondrocytes divide rapidly and form columns of stacked cells parallel to the long axis of the bone (option 2)

• The hypertrophic cartilage zone contains large chondrocytes whose cytoplasm has accumulated glycogen. The resorbed matrix is reduced to thin septa between the chondrocytes.

• Simultaneous with the death of chondrocytes in the calcified cartilage zone, the thin septa of cartilage matrix become calcified by the deposit of hydroxyapatite.

• In the ossification zone, endochondral bone tissue appears (option 4). Blood capillaries and osteoprogenitor cells formed by mitosis of cells originating from the periosteum invade the cavities left by the chondrocytes. The osteoprogenitor cells form osteoblasts, which are distributed in a discontinuous layer over the septa of the calcified cartilage matrix. Ultimately, the osteoblasts deposit bone matrix over the three-dimensional calcified cartilage matrix.

The epiphyseal cartilage is responsible for the growth in the length of bone and it disappears in adults, which is why bone growth ceases in adulthood.





Solution to Question 5:

The structures passing in between the superior constrictor muscle (structure 1) and middle constrictor muscle (structure 2) are glossopharyngeal nerve and stylopharyngeus muscle.

The structures passing in between pharyngeal muscles are as follows:

1. Sinus of Morgagni - It is a semilunar space that exists between the upper concave border of the superior constrictor and the skull base. It is closed off by the upper sturdy portion of the pharyngobasilar fascia. Structures passing through this space include:

a. The auditory tube

b. The levator veli palatini muscle (Option C)

c. The ascending palatine artery

d. Palatine branch of the ascending pharyngeal artery.

2. Between the superior and middle constrictors - the stylopharyngeus muscle and the glossopharyngeal nerve.

3. Between the middle and inferior constrictors -the internal laryngeal nerve and the superior laryngeal vessels penetrate the thyrohyoid membrane (Option B).

4. Between the lower border of the inferior constrictor and the esophagus - The recurrent laryngeal nerve and the inferior laryngeal vessels.

Solution to Question 6:

In the right pelvic bone given above the marked structure is the ischial spine which lies approximately at the level of the fifth sacral vertebrae in a normal anatomical position. As the tip of the coccyx is the closest option relating to the spine , it can be the most appropriate answer.

The ischial spine is a bony prominence found on the ischium, one of the three bones that make up the hip bone. The ischial spine forms a part of the plane of least pelvic dimensions or narrow pelvic plane which comprises of :

• lower border of the symphysis pubis

• the tip of ischial spines

• fifth sacral vertebra

The interspinous distance is the distance between the tip of two ischial spines. (Bispinous diameter = 10.5 cm or 4.5")

The important events occurring at the narrowest plane in the pelvis :

• It corresponds roughly to the origin of levator ani muscles.

• The internal rotation of the fetal head occurs during labour at this plane.

• It marks the beginning of the forward curve of the pelvic axis

• This level of ischial spines indicates station ‘O’

• When the fetal head station is at ‘O’, the head is considered to be engaged.

• Deep transverse arrest occurs at this plane.

• The ischial spine can serve as a landmark in pudendal anaesthesia, as the pudendal nerve lies close to the ischial spine

The image given below highlights the interspinous distance seen through the pelvic inlet.





Solution to Question 7:

The above is an image of the left lung with the arrow pointing at the nerve passing anterior to the left hilum. The phrenic nerve passes in front of the left hilum and has its root value from the ventral horn of C4.

In the given image, the left lung is identified by the vertical sequence of structures at the left hilum - pulmonary artery, principal (hyparterial) bronchus, and inferior pulmonary vein. The image also shows an oblique fissure dividing the lung into superior and inferior lobes.

The phrenic nerve, originating from the cervical nerves C3–C5, is a mixed nerve that carries motor fibres to the diaphragm and sensory fibres from the mediastinal pleura, pericardium, and a portion of the peritoneum.

The course of the left phrenic nerve:

Cervical region - both phrenic nerves pass, Anterior to scalenus anterior muscle

Posterior to sternocleidomastoid and inferior belly of omohyoid

Posterior to internal jugular vein, transverse cervical artery, and suprascapular artery

Posterior to the thoracic duct on the left

Root of the neck - It lies medial to the trunks of the brachial plexus in the neck. At the root of the neck, it runs anterior to the second part of the subclavian artery and posterior to the subclavian vein.

Thorax - The phrenic nerve enters the thorax by crossing medially in front of the internal thoracic artery. Within the thorax, the phrenic nerve descends anterior to the pulmonary hilum between the fibrous pericardium and mediastinal pleura, accompanied by the pericardiacophrenic vessels.



Solution to Question 8:

The marked structure represents the interventricular septum, which is supplied by the anterior interventricular artery.

Blood Supply to the ventricular septum is as follows :

• Septal branch from posterior interventricular artery - Posterior one-third of the ventricular septum

• Septal branch from anterior interventricular artery - Anterior two-thirds of ventricular septum

Blood Supply of Heart :

Other options :

Option A - Diagonal artery supplies the left ventricle.

Option B - Right marginal artery supplies the right ventricle and the apex of the heart.

Option D - Left circumflex artery supplies the left atrium and the left ventricle.

	Right coronary artery	Left coronary artery
Origin	Anterior aortic sinus of ascending aorta, diameter less than left coronary artery	Left posterior aortic sinus of ascending aorta. (Right posterior aortic sinus – noncoronary sinus)
Branches	Right conus artery (third coronary artery)Right anterior ventricular artery – supplies sternocostal surface of right ventricle.Atrial rami – supplies right atrium.Posterior interventricular artery – supplies diaphragmatic surface of right and left ventricle.Right posterior atrial and ventricular rami	Anterior interventricular arteryAnterior ventricular ramiCircumflex arterySA nodal branch arise from this in 35% of subjectsSingle Left marginal (obtuse)arteryAtrial branch
Areas of distribution	The whole of right atriumMost of the right ventricle except a strip along the anterior interventricular groovePart left ventricle adjoining posterior interventricular groove	Most of the left atriumMost of the left ventricle except area adjoining the posterior interventricular grooveA small part of right ventricle along anterior interventricular groove



Solution to Question 9:

The above cross-sectional picture is denoted to be T10 because ,

At the level of the T10 vertebra, we observe:

• The esophagus enters the abdomen through the diaphragm (at the level of T10), connecting shortly afterward to the stomach at the cardiac orifice. In this image, the esophagus is positioned slightly left to the midline and is surrounded by the crura of the diaphragm.

• Adipose tissue around the stomach representing greater omentum

• Nearby structures include the spleen posteriorly, and the right and anterior left lobes of the liver.

• The inferior vena cava (IVC), slightly right of midline, is mostly surrounded by the liver.

• The descending aorta, azygos vein, and thoracic duct are within the thorax.

• A small portion of the left and right lung are still visible.

• Small cross-sectional area of the sternum and the costal cartilages of the 7th ribs. The 7th ribs are the last true ribs with costal cartilages articulating with the sternum

• Predominance of thoracic and upper limb musculature.

• Diaphragm encircling the abdominal cavity, still positioned deep to the thoracic wall.

Solution to Question 10:

Image A shows a section of the cerebellum and image B shows a section of the midbrain. The structure labeled as 'X' represents the dentate nucleus of the cerebellum, which has a direct connection with the structure marked as '1', which is the red nucleus.

The dentate nucleus is the largest of the cerebellar nuclei. It has the shape of a crumpled bag with the opening facing medially. The interior of the bag is filled with white matter made up of efferent fibers that leave the nucleus through the opening to form a large part of the superior cerebellar peduncle.

The efferent pathways from the dentate nucleus connect to the thalamus and the red nucleus, via the dentato- rubro -thalamic pathway.

Other structures in the image B :

2 - Represents superior colliculus, which receives afferents from the retina and the occipital cortex.

3 - Represents substantia nigra, which receives afferents mainly from the caudate nucleus

4 - Represents the crus cerebri, which contains fibers from the cerebral cortex.

Solution to Question 11:

The tail of Spence extends across the anterior axillary fold and is a small extension of the upper lateral quadrant of the breast.

The breast lies in the superficial fascia of the pectoral region. It is divided into four quadrants, i.e. upper medial, upper lateral, lower medial and lower lateral.

Vertically, it extends from the second to the sixth ribs. Horizontally, it extends from the lateral border of the sternum to the midaxillary line. A small extension of the upper lateral quadrant, called the axillary tail of Spence, passes through an opening in the deep fascia and lies in the axilla. The opening is called the foramen of Langer.

The breast lies on the deep pectoral fascia covering the pectoralis major and serratus anterior superiorly, the external abdominal oblique and its aponeurosis inferiorly, and the clavipectoral fascia laterally. Fibrous strands or sheets consisting of condensations of connective tissue extend between the posterior lamina of the superficial fascia and the dermis. These suspensory ligaments (retinacula or Cooper's ligaments) are often well-developed in the superior parts of the breast and support the breast tissue, helping to minimize drooping.

The breasts normally comprise up to 20 discrete lobes that contain ductal-glandular tissue consisting of branching ducts and terminal secretory lobules in a connective tissue stroma. Each lobe is derived by branching from one central lactiferous duct in the nipple. The terminal duct lobular unit (TDLU or lobule) is the functional milk secretory component of the breast.

The upper outer quadrant of the breast contains a greater volume of tissue than the other quadrants. This is the reason for the majority of the carcinomas arising in the upper outer quadrant.





Solution to Question 12:

Lower lip paralysis is caused by damage to the cervical branch of the facial nerve.

The cervical branch of the facial nerve is responsible for innervating the platysma muscle, which is one of the muscles involved in the movements of the lower lip. Damage to this nerve can potentially lead to paralysis of the lower lip.

The facial nerve is the motor nerve of the face. Its five terminal branches emerge from the parotid gland and diverge to supply the various facial muscles as follows :

• Temporal — frontalis, auricular muscles, orbicularis oculi

• Zygomatic — orbicularis oculi (lower eyelid part)

• Buccal — muscles of the cheek and upper lip

• Marginal mandibular — muscles of the lower lip

• Cervical — platysma



Solution to Question 13:

The CT finding in the image is suggestive of an aberrant right subclavian artery.

An aberrant right subclavian artery occurs when the right fourth aortic arch and right dorsal aorta involute cranial to the seventh intersegmental artery.

This artery arises distal to the left subclavian artery and crosses the midline posterior to the esophagus. Because of the absence of a normal right subclavian-carotid junction to pull down the right recurrent laryngeal nerve (RLN) during embryologic development, the right RLN follows a straight path from the vagus to the larynx and is called a non-recurrent right laryngeal nerve.

Most patients with this anomaly are asymptomatic, and the aberrant subclavian artery is of no clinical consequence. Dysphagia caused by compression of the esophagus by an abnormal subclavian artery is called dysphagia lusoria.



Solution to Question 14:

The marked structure is superficial temporal artery.

The superficial temporal artery is one of the terminal branches of the external carotid artery, along with the maxillary artery. It ascends in front of the ear, passing over the zygomatic arch and travels superficially, supplying the scalp and areas around the temple.



Other Options:

Option B: The posterior auricular artery is smaller and located posterior to the ear.

Option C: The middle meningeal artery is deep within the cranial cavity and supplies the meninges.

Option D: The occipital artery lies posteriorly and supplies the scalp at the back of the head.

Solution to Question 15:

The marked structure indicates the pterion and it serves as a landmark for the stem of the lateral sulcus.

The frontal, parietal, temporal, and sphenoid bones meet at the pterion. It is a landmark for the stem of the lateral sulcus (Sylvian fissure), the most prominent sulcus of the cerebral hemisphere. It separates the frontal lobe from the temporal lobe. Deep to the lateral sulcus lies the insular cortex.



Pterion marks as a surface landmark for the following structures:

• Middle meningeal artery and vein: fractures in this area can rupture the middle meningeal artery, leading to an epidural hematoma.

• Broca's area: Located in the dominant hemisphere (usually left) in the inferior frontal gyrus, it lies in proximity to the pterion

• The insular cortex and parts of the basal ganglia, such as the putamen, are located deep to the pterion, beneath the lateral sulcus.

• Stem of the lateral sulcus, also called the Sylvian fissure. It houses the middle cerebral artery and its branches, playing a crucial role in brain vascularization and cortical organization.

During surgeries, the pterion is a common site for trephining (making a hole in the skull).

Other Options:

Option A: The pineal gland is located in the posterior aspect of the cranial fossa in the brain. It is located deep in the brain, near the midline.

Option B: Wenicke's area is in the superior temporal gyrus in the dominant cerebral hemisphere. It is found in the posterior part of the superior temporal gyrus.

Option D: The internal carotid artery siphon is a part of the internal carotid artery's course. Characterized by its S-shaped curvature as it traverses the cavernous sinus and enters the cranial cavity. This configuration helps dissipate pulsatile energy, reducing stress on cerebral vessels, and allows adaptability to cranial movements while maintaining cerebral perfusion.

Solution to Question 16:

This patient, recovering from facial nerve palsy, getting food stuck between the cheek and teeth while eating, is because of the buccinator being affected.

The buccinator is a muscle of facial expression. It is innervated by the facial nerve. It is known as the whistling muscle.

The buccinator muscle is primarily responsible for keeping food between the teeth during mastication by pressing the cheek against the teeth (involved in flattening the cheeks against the gums and teeth).

In facial nerve palsy, paralysis of the buccinator muscle can lead to food accumulating in the cheek pouch, as the muscle is unable to contract properly.

Other options:

Option B: Levator Labii Superioris: This muscle elevates the upper lip and contributes to facial expressions like showing the upper teeth and dilates the nostrils. It is not involved in keeping food in place during chewing.

Option C : Zygomaticus Major: Responsible for pulling the corner of the mouth upward and outward, contributing to smiling. It does not influence food placement during mastication.

Option D : Orbicularis Oris: Encircles the mouth and helps in closing the lips. While it is involved in mastication and speech, its paralysis would not cause food to accumulate in the cheek.

Solution to Question 17:

Retinal pigment epithelium is not a derivative of neural crest cells.

Retinal pigment epithelium (RPE) is derived from the neuroectoderm. The RPE develops from the outer layer of the optic cup, a structure that arises from the neural tube (neuroectodermal origin), not from neural crest cells.

Neural crest cells are a group of cells that migrate during embryonic development and give rise to various structures, including parts of the nervous system, face, and heart.

Neural crest cells (NCC) are present adjacent to the neural tube and are sometimes referred to as the fourth germ layer. It contributes to the formation of many organs and tissues:

In the trunk region, they form:

• Melanocytes (Option D)

• Sensory ganglia

• Sympathetic and enteric neurons

• Schwann cells

• Adrenal medulla cells

In the cranial region, they contribute to the:

• Craniofacial skeleton

• Neurons for cranial ganglia-Ciliary ganglion (Option C)

• Ciliary muscles (Option A)

• Satellite glial cells

• Melanocytes

Some of the other neural crest derivatives are:

• C cells of the thyroid gland

• Conotruncal septum in the heart

• Odontoblasts

• Smooth muscle cells to blood vessels to the face and forebrain.

Solution to Question 18:

The cranial component of neural crest cells (NCC) leads to the formation of Conotruncal septum.

The conotruncal septum is also known as the aortopulmonary septum. The formation of the aortopulmonary septum in the truncus arteriosus divides the outflow channel of the heart into the ventral aorta and the pulmonary trunk. Persistent truncus arteriosus results when the conotruncal ridges fail to form.

The development of the heart begins in the third week of embryogenesis with the formation of the cardiogenic mesoderm. The heart tube forms through fusion of paired endocardial tubes, which then undergo looping and septation to establish the four chambers. By the end of the eighth week, the heart achieves its definitive structure, with separate atria, ventricles, and major vessels.

The heart develops from mesodermal origins, with different parts deriving from specific embryonic structures:

• First Heart Field (Primary Heart Field): Forms the left ventricle and part of the atria.

• Second Heart Field: Contributes to the right ventricle, outflow tract (conus arteriosus and truncus arteriosus), and part of the atria.

• Neural Crest Cells: Contribute to the septation of the outflow tract and the formation of the aorticopulmonary septum.

• Proepicardium (Epicardial Cells): Forms the epicardium and contributes to coronary vessels.

• Sinus Venosus: Forms part of the right atrium (sinus venarum) and the coronary sinus.

• Endocardial Cushions: Form the atrioventricular valves and contribute to septation.

Other options:

Option A: SA node is derived from the primary heart field - sinus venosus.

Option B: AV cushions are derived from adjacent endocardial cells. The endocardial cells along with the epicardium and mesenchyme, form the AV cushions.

Option D: Chordae tendineae are formed from mesenchymal tissue within the heart.



Solution to Question 19:

This patient, undergoing splenectomy, while separating the spleen from the stomach, is likely bleeding from the short gastric vessels.

In an open splenectomy, the gastrosplenic ligament is opened and short gastric vessels are ligated and divided.

The spleen is attached to the stomach through the gastrosplenic ligament. The gastrosplenic ligament attaches the greater curvature of stomach to the upper pole of the spleen.



Contents of the gastrosplenic ligament include:

• Short gastric vessels

• Left gastroepiploic vessels

Hence, while separating the spleen from the stomach, there are maximum chances of bleeding if there is injury to the short gastric vessels.

Other Options:

Option A: Left gastric vessels: Left gastric vessels run along the lesser curvature of the stomach. It is unlikely to be the source of bleeding in this case.

Option C: Right gastro-epiploic vessels: The right gastroepiploic vessels run along the pylorus and inferior region of the greater curvature of the stomach. It is not a part of the gastrosplenic ligament.

Option D: Splenic vessels: The splenic artery and vein run along the superior border of the pancreas. They are not directly involved in the separation of the spleen from the stomach.

Solution to Question 20:

The marked structure is the primitive streak.

The primitive streak appears in the early stages of gastrulation, a process that reorganizes the epiblast into three germ layers: ectoderm, mesoderm, and endoderm. It appears as a linear thickening on the dorsal surface of the epiblast in the bilaminar embryonic disc. It is found in the caudal (tail-end) region of the developing embryo.



Functions of the primitive streak:

• Serves as the site where cells migrate inward to form the mesoderm and endoderm.

• Marks the beginning of gastrulation.

• Defines the body axis (head-to-tail, left-to-right).

Other Options:

Option B: Notochord: A rod-like structure derived from mesoderm that forms after the primitive streak and is crucial for the development of the neural tube. Located cranially relative to the primitive streak.



Option C: Neural Groove: A depression in the ectoderm that forms during neurulation, leading to the development of the neural tube. It appears after the primitive streak.

Option D: Secondary Neural Plate: Part of the neural plate involved in the later stages of neural tube development. It is not directly related to the primitive streak.

Solution to Question 21:

Endoderm is the first to be formed by the migration of the epiblast along the primitive streak.

Gastrulation is when the single-layered blastula reorganizes into a multi-layered structure called the gastrula. This process establishes the three primary germ layers that give rise to all tissues and organs of the body. During gastrulation, epiblast cells migrate through the primitive streak to form the three germ layers: endoderm, mesoderm, and ectoderm.

Cells from the epiblast migrate inward (invagination) through the primitive streak.

The first group of migrating epiblast cells replaces the hypoblast to form the definitive endoderm. The endoderm gives rise to structures like the gastrointestinal tract, liver, and lungs.

Subsequent migrating cells spread between the epiblast and the newly formed endoderm to create the mesoderm, which forms structures like muscles, bones, and the cardiovascular system.

Remaining epiblast cells that do not migrate through the primitive streak form the ectoderm, which gives rise to the epidermis and the nervous system.



Other Options:

Option B: Cardiac Mesoderm: The mesoderm forms after the endoderm and differentiates later into subtypes, including the cardiac mesoderm.

Option C: Epidermis: The epidermis arises from the ectoderm, which forms after both the endoderm and mesoderm.

Option D: Neural Groove: The neural groove is part of the ectoderm and forms during neurulation, which occurs much later than the formation of the endoderm.

Solution to Question 22:

Gluteus Medius of the supporting leg is responsible for raising the hip bone on the side of the suspended leg.

Actions of Gluteus Medius:

- Hip abduction
- Stabilizes the pelvis during gait.
- Assists the internal rotation of the thigh.

Gluteus medius and gluteus minimus act as the abductors of the hip joint. Stabilizing the pelvis on the femur during walking is done by these muscles. It is demonstrated by asking the subject to stand on one limb.

Trendelenburg Sign:

Trendelenburg sign is a clinical sign used to assess the function of the gluteus medius.

It is positive when the pelvis drops on the opposite side of the stance leg, indicating weakness or dysfunction of the gluteus medius on the supporting leg.

In a healthy individual, the gluteus medius contracts to prevent the pelvis from tilting when one leg is lifted. If the gluteus medius is weak or paralyzed, the pelvis tilts toward the side of the suspended leg.



Option B: Gluteus Maximus: Extends and externally rotates the hip; key for climbing stairs and rising from sitting. Weakness causes difficulty in hip extension, affecting activities like standing up or climbing.

Option C: Obturator Internus: Externally rotates and stabilizes the hip joint; abducts the thigh when the hip is flexed. Dysfunction can impair hip rotation, commonly affected in pelvic injuries.

Option D: Quadratus Femoris: Externally rotates and adducts the hip. Injury or entrapment of the nerve to quadratus femoris can cause deep hip pain and weakness in rotation.

Solution to Question 23:

The structure shown in the above image is the oculomotor nerve. The oculomotor nerve (CN III) passes through the superior orbital fissure, which lies within the region of the cavernous sinus.



Option B: The abducens nerve (CN VI) also passes through the cavernous sinus and superior orbital fissure. However, it is located more medially in the sinus and the fissure compared to the oculomotor nerve.

Option C: The trochlear nerve (CN IV) passes through the superior orbital fissure as well, but it is the smallest cranial nerve and follows a higher and more lateral path than the oculomotor nerve.

Option D: The ophthalmic division of the trigeminal nerve enters the lateral wall of the cavernous sinus, below the oculomotor nerve and trochlear nerve.



Solution to Question 24:

Injury to the posterior interosseous nerve leads to loss of extension of digits.

The posterior interosseous nerve (PIN) is a continuation of the deep branch of the radial nerve. It originates from the radial nerve in the cubital fossa, near the lateral epicondyle of the humerus. After passing through the supinator muscle, it transitions into the PIN.

The PIN is purely a motor nerve and supplies the posterior compartment of the forearm.

Clinical correlation based on location of lesion:

Monteggia fracture involves a fracture of the proximal ulna with a dislocation of the radial head. The radial nerve, especially its posterior interosseous branch, is at risk due to its proximity to the fracture site and leads to loss of extension of digits.

Level of lesion	Nerve affected	Cause	Clinical feature
Low	Posterior interosseous nerve and Superficial Cutaneous	Fracture or dislocations of the elbow.- Commonly seen in Monteggia fracture	Cannot extend the MCP joint (digits)(Option A).Weakness in the extension of the thumb. Radial deviation of the wrist.
High	Radial nerve in spiral groove	Humerus fracture.	Wrist drop (Option D)"Saturday night palsy" due to compression of the radial nerveThumb drop, finger drop
Very High	Radial nerve proper	Lesions around the shoulder joint & axilla and crutch palsy	Loss of extension of the wrist and hand.(Option B)Inability to extend elbow

Anatomy NEET 2024

Question 1:

An absent gag reflex can result from injury to which of the following nerves?

a) ■ and ■

b) ■ and ■

c) ■ and ■

d) ■ and ■

Question 2:

If a tumor compresses the structures passing through the opening marked in the given image, which of the following sensations would be affected?



a) Sensation of the nasopharynx

b) Mucosa of the nasal cavity

c) Sensation in the anterior scalp and dorsum of the nose

d) Sensation of the chin

Question 3:

A chronic tobacco consumer went to the dental clinic with bleeding lesions on his tongue and was diagnosed with carcinoma of the tongue. Which of the following papillae, which do not have taste buds, is least likely to be involved in such a cancer?

a) Circumvallate

b) Filiform

c) Fungiform

d) Foliate

Question 4:

Which artery is used for arteriovenous shunting with the great saphenous vein for hemodialysis, and is located behind the medial malleolus and 2.5 cm in front of the tendocalcaneus?

a) Anterior tibial artery

b) Posterior tibial artery

c) Peroneal artery

d) Circumflex fibular artery

Question 5:

What is the outer covering of the condition shown in the image?

a) Chorion

b) Amnion

c) Endoderm

d) Ectoderm

Question 6:

Identify the marked structure.



a) Thoracodorsal nerve

b) Medial pectoral nerve

c) Subscapular nerve

d) Long thoracic nerve

Question 7:

A patient presents with sciatica and tenderness in the hamstring muscles. Which nerve supplies the hybrid muscle that is likely to be spared in this condition?

a) Common peroneal nerve

b) Tibial nerve

c) Femoral nerve

d) Obturator nerve

Question 8:

Where will you find the epithelium shown in the image?



a) Gall bladder

b) Trachea

c) Ureter

d) Duodenum

Question 9:

During the examination of a patient, a professor asks the patient to invert his foot and points to a prominent tendon located behind the medial malleolus. Which tendon is being highlighted?

a) Tibialis anterior

b) Extensor digitorum

c) Tibialis posterior

d) Extensor hallucis longus

Question 10:

An elderly woman reports difficulty in buttoning her shirt and increased clumsiness during daily activities. On examination, she is unable to make an 'O' shape by opposing her thumb and index finger. Which of the following muscles is most likely involved?

a) Opponens pollicis

b) Palmar interossei

c) Flexor pollicis brevis

d) Abductor pollicis brevis

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	b
5	b
6	b
7	d
8	c
9	c
10	d

Detailed Explanations

Solution to Question 1:

An absent gag reflex can result from injury to the glossopharyngeal nerve (IX) and the vagus nerve (X).

The afferent nerve for the gag reflex is the glossopharyngeal nerve, while the efferent nerve consists of both the glossopharyngeal (to the stylopharyngeus muscle) and vagus. Both these nerves pass through the jugular foramen and damage to them may be the cause for loss of gag reflex.

Pharyngeal branches of the glossopharyngeal nerve and vagus nerve form the pharyngeal plexus of nerves. These plexes mainly supply the pharynx. It also has contributions from the superior cervical ganglion.

Nerve supply by the pharyngeal branch of the vagus nerve are:

• All the muscles of the soft palate(except tensor veli palatini - supplied by mandibular division of the trigeminal nerve)

• All the muscles of the pharynx(except the stylopharyngeus - supplied by the glossopharyngeal nerve).

The glossopharyngeal nerve is rarely damaged alone. Lesions will result in loss of sensation of the posterior third of the tongue, reduced gag reflex, loss of sensation over the ipsilateral soft palate, etc. Motor fibres to the stylopharyngeus and parasympathetic secretomotor fibres to the parotid gland are also supplied by the glossopharyngeal nerve. Sensory supply includes uvula, posterior one-third of the tongue, tympanic cavity etc.

The main functions of IX and X can be tested together by assessing swallowing, phonation/articulation and sensation from the pharynx/larynx.

Glossopharyngeal and vagus nerve examinations include:

• Observing the position and symmetry of the palate and uvula at rest and with phonation (“aah”).

• By assessing the strength of the cough

Solution to Question 2:

The marked foramen is the superior orbital fissure. If a tumour compresses the structures passing through this foramen, it would impair sensation in the anterior scalp and dorsum of the nose, as this would affect the frontal nerve and nasociliary nerve.

The superior orbital fissure is an important anatomical opening in the skull that allows several critical structures to pass through. Among these structures, the frontal nerve and nasociliary nerve are branches of the ophthalmic nerve, the first division of the trigeminal nerve (CN V1), which is responsible for sensory innervation to parts of the face.

The frontal nerve provides sensation to the forehead, the anterior scalp, and the dorsum of the nose, while the nasociliary nerve supplies sensation to parts of the nasal cavity. Therefore, if a tumour or lesion compresses the structures passing through the superior orbital fissure, it would likely affect the sensation in the anterior scalp and dorsum of the nose because these areas are innervated by the branches of the trigeminal nerve that pass through this fissure.

Other options:

Option A: Sensation of the nasopharynx is innervated by other branches of the trigeminal and glossopharyngeal nerve, not by structures passing through the superior orbital fissure.

Option B: While some sensation from the nasal cavity comes from the trigeminal nerve, the specific branches affected by a lesion in the superior orbital fissure are more focused on the anterior areas of the face.

Option D: Sensation of the chin is primarily innervated by the mandibular branch (CN V3) of the trigeminal nerve, which passes through the foramen ovale, not the superior orbital fissure.

Solution to Question 3:

Among the different types of papillae on the tongue, the one that does not have taste buds is the filiform papillae.

Filiform papillae are minute cylindrical or conical projections that are mainly present on the dorsal part of the anterior 2/3rd of the tongue. The filiform papillae do not contain taste buds. They are covered by keratinized stratified squamous epithelium.

Other papillae of the tongue are:

Circumvallate papillae are 8-12 in number which are large cylindrical structures. They form a V-shaped row just in front of the sulcus terminalis. Usually, the entire papillae are covered by non-keratinized stratified squamous epithelium. Taste buds are present in the lateral wall of these papillae which surrounds the papillae circumferentially.

Fungiform papillae are large, rounded, and red and are mainly present on the lingual margins and also on the dorsal surface of the tongue. They are covered by non-keratinized stratified squamous epithelium. They are supplied by the facial nerve.

Foliate papillae are red, leaf-like mucosal ridges that are present bilaterally in two zones at the sides of the tongue near the sulcus terminalis. They are covered by non-keratinized stratified squamous epithelium.

A diagrammatic representation of the various papillae of the tongue is given below:



Below is the image showing the various parts of the tongue.



The given image depicts the histology image of filiform papillae:



The given image depicts the histology image of circumvallate papillae:



The given image depicts the histology image of foliate papillae:

The given image depicts the histology image of fungiform papillae:



Solution to Question 4:

The posterior tibial artery, located behind the medial malleolus and 2.5 cm in front of the tendocalcaneus, that is used for arteriovenous (AV) shunting with the great saphenous vein for hemodialysis.

The posterior tibial artery is situated behind the medial malleolus and approximately 2.5 cm in front of the tendocalcaneus, making it a suitable site for creating an AV fistula with the great saphenous vein. This location allows for optimal access and ensures that the shunt can handle the necessary blood flow for effective dialysis with the great saphenous vein. However, they are not commonly used for AV shunting in common practices.

Arteries used for AV shunting include the radial artery and the brachial artery in the upper arm, which are frequently employed for hemodialysis access. In the lower leg, the anterior tibial artery and the peroneal (fibular) artery are less commonly used for this procedure.

Other options:

Option A: In comparison, the anterior tibial artery is located on the anterior aspect of the leg and runs down the front of the tibia, not behind the medial malleolus.

Option C: The peroneal (fibular) artery is situated on the lateral side of the leg, travelling alongside the fibula, and is also not located behind the medial malleolus.

Option D: The fibular circumflex artery is a small artery and is not commonly used for hemodialysis access. It is not a standard choice and does not fit the anatomical location described, as it is a branch of the anterior tibial artery.

Solution to Question 5:

The diagnosis based on the given image is omphalocele and is covered by the amnion.

Omphalocele is a midline abdominal wall defect where the bowel and other solid viscera protrude through the umbilicus. In this condition, the herniated organs are covered by both peritoneum and an amnion. The umbilical cord inserts directly into this sac, which helps in identifying the condition. The defect occurs due to the failure of the physiological hernia to return to the body cavity by the 11th week of gestation.

The size of an omphalocele can vary significantly. It may be relatively small, containing only the gut, or it can be classified as a giant omphalocele if it measures 4 cm or more in diameter, encompassing the gut, liver, and other abdominal organs. The condition is often associated with other congenital anomalies such as trisomies 13, 15, 18, or 21, Beckwith-Wiedemann syndrome, exstrophy of the bladder or cloaca, and pentalogy of Cantrell.

Treatment for omphalocele involves covering the exposed organs to prevent fluid loss and protect them from infection. Prophylactic antibiotics may be administered in case of rupture. Surgical correction is usually performed once the baby is stable. In omphalocele, since the umbilicus is covered by a reflection of the amnion, this membrane also covers the omphalocele itself, differentiating it from other types of abdominal wall defects.



Note : Gastroschisis is a condition where the intestinal loops herniate through the defect in the abdominal wall. The defect usually lies on one side of the umbilicus (right side commonly). The loops of the bowel are not covered by amnion. It is associated with elevated alpha-fetoprotein levels.

Solution to Question 6:

The marked structure in the image is the medial pectoral nerve.

The given image shows the medial pectoral nerve arising from the medial cord of the brachial plexus. It also shows the axillary artery traversing inferior to the median pectoral nerve and the thoracoacromial artery. The medial pectoral nerve and the thoracoacromial artery both contribute to the neurovascular supply of the pectoral muscles.

After originating from the medial cord, the medial pectoral nerve passes anterior to the axillary artery. This nerve is specifically known for its anatomical path of piercing the pectoralis minor muscle before branching out to supply both the pectoralis minor and the pectoralis major muscles.



Solution to Question 7:

The adductor magnus is the hybrid muscle that is likely to be spared in this condition.

In sciatica, the sciatic nerve is typically affected, which branches into the tibial nerve and the common peroneal nerve. The adductor magnus muscle is an example of a hybrid muscle with dual innervation:

• The adductor part of the adductor magnus is innervated by the obturator nerve.

• The hamstring part of the adductor magnus is innervated by the tibial nerve.

During sciatica, the tibial nerve is often affected, which would impact the hamstring part of the adductor magnus. However, because the adductor part of the adductor magnus is innervated by the obturator nerve, it remains functional and is likely to be spared.

The obturator nerve arises from the anterior division of the ventral rami of L2-L4. Near the obturator foramen, it divides into anterior and posterior branches.

• Anterior division supplies pectineus, adductor longus, adductor brevis, and gracilis

• Posterior division supplies adductor magnus, obturator externus. It also supplies adductor brevis if the muscle is not supplied by the anterior division

The image given below shows the course of the obturator nerve.



Other options:

Option A: The common peroneal nerve is a branch of the sciatic nerve and can also be affected in sciatica. It is responsible for innervating muscles in the anterior and lateral compartments of the leg.

Option B: The tibial nerve is a branch of the sciatic nerve and is often affected in sciatica. It supplies the hamstring part of the adductor magnus, which would be impacted during sciatica.

Option C: The femoral nerve is not involved in sciatica. It supplies the quadriceps and other muscles in the anterior compartment of the thigh, and does not innervate the adductor magnus.

Solution to Question 8:

The epithelium shown in the above image can be found in the ureter.

The ureter is lined by transitional epithelium (also called urothelium), similar to the urinary bladder. This specialized epithelium allows stretching, which is crucial as the urine passes through. The presence of transitional epithelium is characteristic of structures in the urinary system, such as the ureters, bladder, and part of the urethra.

Given below is a cross-section of the ureter showing the transitional epithelium and the layer of smooth muscle.



Other options:

Option A: Gall bladder: The gall bladder is lined by simple columnar epithelium, which facilitates the absorption of water and electrolytes from bile.

Option B: Trachea: The trachea is lined by pseudostratified columnar ciliated epithelium with goblet cells, which helps trap and remove particles from the respiratory tract.

Option D: Duodenum: The duodenum is lined by simple columnar epithelium with goblet cells, aiding in digestion and nutrient absorption.

Solution to Question 9:

The tendon being highlighted here is of the tibialis posterior.

It passes behind the medial malleolus and is a key muscle responsible for foot inversion. The tendon of this muscle becomes prominent in this area during foot inversion. Dysfunction of this tendon can lead to flatfoot deformity.

The given image shows the tarsal tunnel and position of the tendon of tibialis posterior:



Other options:

Option A: Tibialis anterior lies anterior to the medial malleolus and aids in dorsiflexion and inversion of the foot but is not located behind the medial malleolus.

Option B: Extensor digitorum is involved in extending the toes and runs anteriorly to the foot, not behind the medial malleolus.

Option D: Extensor hallucis longus extends the big toe and runs along the anterior portion of the foot, also not behind the medial malleolus.

Solution to Question 10:

The difficulty in buttoning the shirt and the inability to make the "O" shape with fingers suggest the involvement of the abductor pollicis brevis muscle.

This is a case of anterior interosseous nerve (AIN) syndrome. The AIN is a purely motor branch of the median nerve that affects specific muscles in the forearm, the flexor pollicis longus and flexor digitorum profundus leading to difficulty in performing tasks requiring fine motor control, such as the "OK" sign. Among the options provided, the abductor pollicis brevis is uniquely and consistently innervated by the recurrent motor branch of the median nerve, making it susceptible to dysfunction in conditions affecting the median nerve.

Other options:

Option A: The opponens pollicis is innervated primarily by the lateral terminal branch of the median nerve, with some contributions from a branch of the deep terminal branch of the ulnar nerve. While it plays a role in thumb opposition, it is not consistently affected in anterior interosseous nerve (AIN) syndrome, which primarily impacts deeper flexor muscles like the flexor pollicis longus. Therefore, it is less likely to be the cause of the patient's inability to perform the "OK" sign.

Option B: The palmar interossei muscles are innervated by the deep branch of the ulnar nerve. Because they do not receive any innervation from the median nerve, they are not implicated in

AIN syndrome. Their primary function involves the adduction of the fingers.

Option C: The flexor pollicis brevis has dual innervation: the superficial head is supplied by the recurrent branch of the median nerve, while the deep head is innervated by the deep branch of the ulnar nerve. This dual supply makes it less likely to be solely affected in AIN syndrome.

Physiology AIIMS 2017

Question 1:

Given below is the demonstration of tetanization in the gastrocnemius muscle of a frog. Identify the tetanising frequency.

a) 25-20 Hz

b) 30-35 Hz

c) 15-20 Hz

d) 10-15 Hz

Question 2:

The oxygen carrying capacity of a 18 year old boy with haemoglobin of 14g/dl is

a) 22

b) 16

c) 14

d) 18

Question 3:

A 30-year-old patient with respiratory volume of 500 mL. The intraplueral pressure measured -4 before inspiration and -8 at the end of inspiration. The pulmonary compliance of the patient is

a) 125 ml/cm H2O

b) 150 ml/cm H2O

c) 50 ml/cm H2O

d) 62.5 ml/cm H2O

Question 4:

The function of the Golgi tendon organ is to _____

a) Sense changes in muscle direction

b) Sense changes in muscle tension

c) Sense changes in position of the muscle

d) Sense changes in length of the muscle

Question 5:

Difference in trajectory between inspiratory loop and the expiratory loop in the curve is due to:



a) Difference in alveolar pressure during inspiration and expiration

b) Difference in conc. of surfactant during inspiration and expiration

c) Difference in airway resistance during inspiration and expiration

d) Inspiration is active and expiration is passive

Question 6:

Reflex responsible for tachycardia during right atrial distension is:

a) Bezold-Jarisch reflex

b) Bainbridge reflex

c) Cushing reflex

d) J reflex

Question 7:

Identify the hormone from the picture



a) Growth hormone

b) Cortisol

c) Estrogen

d) Insulin

Question 8:

Feed forward control system is employed during the regulation of

a) Blood volume

b) pH

c) Temperature

d) Blood pressure

Question 9:

Cellular junctions present in a cardiac muscle are all except:

a) Zonula occludens

b) Fascia adherens

c) Gap junctions

d) Macula adherens

Question 10:

Efferent arteriolar constriction causes all except:

a) Decrease in GFR

b) Decreases renal blood flow

c) Decreases oncotic pressure in glomerular capillaries

d) Increases hydrostatic pressure in glomerular capillaries

Question 11:

Absolute refractory period is due to:

a) Opening of calcium channels

b) Closure of potassium channels

c) Closure of active gates of sodium channel

d) Closure of inactive gates of sodium channel

Question 12:

X, Y, Z are three permeable ions. X = -50 and Y = -30. At RMP, if there is no net electrogenic transfer, what is the value of Z?

a) +20

b) -20

c) +80

d) -80

Question 13:

Slow waves are generated by:

a) Myenteric neurons

b) Smooth muscle cells

c) Interstitial cells of Cajal

d) Parasympathetic neurons

Question 14:

A patient has a ■blood glucose level of 200 mg/dL and GFR of 90 ml/min. What is the amount of glucose excreted if the transport maximum of the patient is as shown below?



a) 80 mg/min

b) 50 mg/min

c) 40 mg/min

d) 30 mg/min

Question 15:

Which of the following can be responsible for Bezold-Jarisch reflex?

a) Serotonin

b) Histamine

c) Prostaglandin

d) Angiotensin

Question 16:

Which of the following is the constituent of the marked area in the electron microscopic picture of muscle?



a) Alpha actinin

b) Actin

c) Titin

d) Tropomyosin

Question 17:

The fourth heart sound is caused by:

a) Closure of aortic and pulmonary valves

b) Vibrations in the ventricular wall during systole

c) Ventricular filling

d) Closure of mitral and tricuspid valves

Question 18:

"a" wave on JVP is for

a) Atrial contraction

b) Atrial filling

c) Ventricular relaxation

d) Ventricular filling

Question 19:

In the 2 vessels shown below, assume the pressure along both the vessels is the same and both follow linear flow pattern. What will be the amount of blood flow in A compared to B



©MARROW

a) 4 times

b) 32 times

c) 8 times

d) 16 times

Question 20:

A 13-year-old boy presents with involuntary dancing movements of his right hand. He was diagnosed with rheumatic fever. What constitutes the ganglia involved in the presenting complication?

a) Caudate nucleus and globus pallidus

b) Globus pallidus and putamen

c) Globus pallidus and lentiform nucleus

d) Caudate nucleus and putamen

Answer Key

Question No.	Correct Option
1	b
2	d
3	a
4	b
5	b
6	b
7	a
8	c
9	a
10	c
11	d
12	c
13	c
14	d
15	a
16	a
17	c
18	a
19	c
20	d

Detailed Explanations

Solution to Question 1:

Tetanizing frequency demonstrated by the muscle is 33Hz. In the above graph, we know,

• The speed of the paper is 1000mm/sec.

• The y-axis represents millivolts, the x-axis represents time.

• The scale on the x-axis is 10mm.

• The area under the graph represents the muscle contraction.

Contraction period (latent period to maximum force of contraction) includes 3 scales = 30 mm.

As paper speed is 1000mm/sec,

Contraction period = 30mm = (30/1000) = 30 msec.

Tetanizing frequency = 1/contraction period = 1000/30 = 33Hz.

Tetanization

The features of muscle contraction can be demonstrated by eliciting single muscle twitches. The adding together of individual twitch contractions to increase the intensity of overall muscle contraction is Summation. Summation occurs in two ways:

• by increasing the number of motor units contracting simultaneously, which is multiple ■ber summations.

• by increasing the frequency of contraction, which is called frequency summation and can lead to tetanization.

When the frequency of a contraction reaches a critical level, the successive contractions eventually fuse together, and the whole muscle contraction appears to be completely smooth and continuous, this is called tetanization.

Tetanization occurs in two ways:

• Complete tetanus occurs when no relaxation occurs between stimuli.

• Incomplete tetanus when periods of incomplete relaxation take place between the summated stimuli.



Solution to Question 2:

The oxygen carrying capacity of a boy with haemoglobin 14g/dl is 18.

When blood is saturated with 100% O2, the haemoglobin is also 100% saturated and each gram of normal haemoglobin contains 1.39 mL of O2.

However, blood normally contains some inactive haemoglobin derivatives, and the measured value in vivo is thus slightly lower hence, each gram of haemoglobin in vivo contains 1.31 mL of O2 (Hufner's constant).

According to the above question, when Hb=14g/dl, O2= 1.31*14= 18.34

The oxygen-hemoglobin dissociation curve

• Demonstrates a progressive increase in the percentage of haemoglobin bound with oxygen as blood Po2 increases, which is called the percent saturation of haemoglobin.

• Blood leaving the lungs and entering the systemic arteries usually has a Po2 of about 95 mm Hg, the usual oxygen saturation of systemic arterial blood averages 97 per cent.

• Conversely, in normal venous blood returning from the peripheral tissues, the Po2 is about 40 mm Hg, and the saturation of haemoglobin averages 75 per cent.



Solution to Question 3:

Pulmonary compliance of this patient is 125 mL/cm of H2O.

Compliance is developed due to the tendency for tissue to resume its original position after an applied force has been removed.

After an expiration during quiet breathing, the lungs have a tendency to collapse and the chest wall has a tendency to expand. The interaction between the recoil of the lungs and recoil of the chest can be measured through a spirometer and a curve of airway pressure plotted against volume is the pressure-volume curve of the total respiratory system (Ptr).

Compliance of the lung and chest wall is measured as the slope of the Ptr curve, or, as a change in lung volume per unit change in airway pressure ($\Delta V/\Delta P$).

Compliance = Change in lung volume / Change in pressure

Compliance = 500/4

Compliance = 125 ml/cm H20

Solution to Question 4:

Golgi tendon organ sense changes in muscle tension.

Golgi tendon reflex (Inverse myotatic/ Inverse stretch reflex) is disynaptic, is the opposite of the stretch reflex.

When the tension in a contracted muscle is significantly elevated, the inhibitory effect from the Golgi tendon organ leads to instantaneous relaxation of the entire muscle. This effect is called the lengthening reaction.

It is a protective mechanism to prevent tearing of the muscle or avulsion of the tendon from its attachments to the bone.

Note: Muscle spindles senses changes in muscle length (stretch).





Solution to Question 5:

The difference between the inflation and the deflation paths is called hysteresis.

It exists because a greater transpulmonary pressure is required to open a previously closed airway, due to a deficit of surfactant at the air-water interface, than to keep an open airway from closing, as then it has abundant surfactant.

Solution to Question 6:

Reflex responsible for tachycardia during right atrial distension is Bainbridge Reflex.

The stretch receptors of the atria that elicit the Bainbridge reflex transmit their afferent signals through the vagus nerves to the medulla of the brain. Then efferent signals are transmitted back through vagal and sympathetic nerves to increase heart rate and strength of heart contraction. Thus, this reflex helps prevent damming of blood in the veins, atria, and pulmonary circulation.

Other options:

Option A: Bezold–Jarisch Reflex: Activation of chemosensitive vagal C fibers in the cardiopulmonary region causes profound bradycardia, hypotension and a brief period of apnoea followed by rapid shallow breathing.

Option C: Cushing Reflex: When intracranial pressure is increased, the blood supply to Rostral Ventro-Lateral Medulla (RVLM) neurons is compromised. The local hypoxia and hypercapnia cause an increase in discharge from the neurons, producing hypertension, bradycardia, and irregular respiration.

Option D: J Reflex: The stimulation of the J-receptors (C receptors/juxta-pulmonary capillary receptors) causes apnea followed by rapid breathing, bradycardia, and hypotension.

Solution to Question 7:

The image depicts the fluctuations of secretion of growth hormone throughout the day.

• Growth hormone is found at relatively low levels during the day unless specific triggers such as strenuous exercise are present.

• During sleep, on the other hand, large pulsatile bursts of growth hormone secretion occur.



Stimuli that Affect Growth Hormone Secretion in Humans

Note:

The early morning surge of Growth Hormone during sleep, as seen in the image, plays a role in the Dawn Phenomenon

Stimuli that Increase Secretion	Stimuli that Decrease Secretion
ExerciseGoing to sleepFasting/HypoglycemiaStressful stimuli (including various psychological stresses)High Protein MealGlucagonEstrogens and androgensPyrogen	GlucoseCortisolFFAMedroxyprogesteroneGrowth hormone and IGF-I

Solution to Question 8:

Feed forward control system is employed during regulation of temperature.

Feed forward control

There is a control system in our body when no stimulus is required but still the system anticipates and makes corrective changes.

Such a system is called Feed forward or Anticipatory or Adaptive Control.

Examples include

• Temperature control – Thermoregulatory responses are initiated by hypothalamus BEFORE the changes in environmental temperature have succeeded in changing the body's core body temperature

• Cephalic phase of gastric secretion- Just thinking about food increases gastric acid production

• Thinking about exercise itself increases heart rate & respiratory rate

• Role of cerebellum in motor coordination

Solution to Question 9:

Zonula occludens is the other name for the tight junction. It is absent in a cardiac muscle.

Cardiac muscle cells are connected by Fascia adherens (a structural analog of zonula adherens), Gap junctions and Macula adherens.

Cardiac muscle fibers are made up of many individual cells connected in series and parallel to one another. The areas where the end of one muscle fiber abuts on another are called intercalated disks. At each intercalated disc, the plasma membrane fuses with one another forming gap junction which provides low resistance bridges for the spread of excitation from one fiber to another permitting the cardiac muscle to function as a syncytium.

Solution to Question 10:

Efferent arteriolar constriction increases oncotic pressure in glomerular capillaries.

Efferent arteriolar constriction has a biphasic response on GFR depending on the severity of resistance.

Moderate efferent arteriolar constriction causes increased glomerular hydrostatic pressure and slightly increased GFR.

However, in cases of severe efferent arteriolar constriction (more than threefold increase in efferent arteriolar resistance), the renal blood flow is significantly reduced. This results in increased plasma protein concentration, thereby increasing the glomerular colloid oncotic pressure. As the increase in colloid oncotic pressure exceeds the increase in hydrostatic pressure, the net force for filtration decreases, resulting in decreased GFR.



Solution to Question 11:

Absolute refractory period is due to closure of inactive gates of sodium channel.

Voltage-gated sodium channels have 3 states that you can find them in. They can either be closed, open, or inactive.

When the stimulus causes activation of the open gates, sodium rushes into the cell causing the membrane potential to become more positive. After 0.5 to 1 millisecond, the inactive gate automatically closes and no stimulus, no matter how strong, can excite the nerve. This is called the absolute refractory period.



Solution to Question 12:

No net electro genic transfer at RMP means X + Y + Z = 0

Since X = -50 and Y = -30, then Z must be +80 since (-80 +80 = 0)

Solution to Question 13:

The interstitial cells of Cajal undergo cyclic changes in membrane potential due to unique ion channels that periodically open and produce inward (pacemaker) currents that generate slow-wave activity.

Interstitial cells of Cajal are the pacemaker cells of the GIT that initiate the basic electrical rhythm (BER). These are stellate mesenchymal pacemaker cells with smooth muscle-like features.

In the stomach and the small intestine, these cells are located in the circular smooth muscle layer near the myenteric plexus. In the colon, they are at the submucosal border of the circular muscle layer.

The slow waves usually do not by themselves cause muscle contraction in most parts of the gastrointestinal tract, except perhaps in the stomach. Instead, they mainly excite the appearance of intermittent spike potentials, and the spike potentials in turn actually excite the muscle contraction.

Note: Gastrointestinal stromal tumors arise from the interstitial cells of Cajal.

Solution to Question 14:

The amount of glucose excreted in the above scenario is 30 mg/min.

Calculation:

Step 1: Calculate total filtered glucose:

In this question,

• GFR = 90 ml/min

• Plasma glucose concentration = 200 mg/dL= 2 mg/ml (1 dL=100 ml).

• Total filtered glucose = GFR X Plasma glucose concentration.

= 90 ml/min X 2 mg/ml

= 180 mg/min.

Step 2: Check transport maximum:

In the above question transport maximum is 150 mg/min.

Step 3: Calculate the excretion:

Excretion = Total filtered glucose - Total reabsorbed glucose (Transport Maximum)

= 180-150 mg/min

= 30 mg/min

Solution to Question 15:

Serotonin can be responsible for Bezold–Jarisch reflex.

Activation of chemosensitive vagal C fibers located in the cardiopulmonary region (ex: ventricles, atria, great veins, pulmonary artery, juxtacapillary region of alveoli) causes profound bradycardia, hypotension, and apnea followed by rapid shallow breathing. This response is called Bezold-Jarisch reflex.

Other substances that can cause Bezold–Jarisch reflex are capsaicin, phenyl biguanide, and veratridine.

Solution to Question 16:

Alpha actinin is the constituent of the marked area.

The image depicts sarcomere and the arrow points towards Z-line. Alpha actinin binds actin to Z-lines.



The orderly arrangement of actin, myosin, and related proteins that produces this pattern. The area between two Z-lines is called sarcomere.

The thick filaments, which are about twice the diameter of the thin filaments, are made up of myosin, which are lined to form A bands; the thin filaments are made up of actin, tropomyosin, and troponin. The array of thin filaments extends out of the A band and into the less dense staining I bands.

The lighter H bands in the center of the A bands are the regions where, when the muscle is relaxed, the thin filaments do not overlap the thick filaments. The Z lines allow for anchoring of

the thin filaments.

Actinin, titin and desmin are additional structural proteins.Titin connects Z-lines to the M line and provides the scaffolding to the sarcomere. Desmin adds structure to Z-lines by connecting Z-line to plasma membrane.

Note: Some sources say that even titin is a component of Z-disc so if alpha actinin is not given in the options titin is the second best answer.



Solution to Question 17:

The fourth heart sound is caused by ventricular filling.

A fourth sound is heard immediately before the first sound when atrial pressure is high or the ventricle is stiff in conditions such as ventricular hypertrophy. It is due to ventricular filling and is rarely heard in normal adults.



Solution to Question 18:

"a" wave in JVP represents atrial contraction.

Waves of JVP :

• a- atrial systole pushing the blood to the ventricles.

• c- isovolumetric contraction of ventricles causing the tricuspid valve to bulge.

• x'- fall in atrial pressure during ventricular systole due to pull of valve leaflets.

• v- venous filling which creates an increase in atrial pressure.

• y- passive filling of ventricles by a passive flow of the blood from the right atrium.



Solution to Question 19:

The relationship between the flow in a long narrow tube, the viscosity of the fluid, and the radius of the tube is expressed mathematically in the formula called as Poiseuille's law :



Blood flow (F) inside the vessel depends on:

• Pressure difference: directly proportional

• Viscosity: inversely proportional

• Radius: directly proportional to the fourth power

• Length of the vessel: inversely proportional

Here, the radius is doubled so the flow increases by 2 to the power of 4 = 16.

the length of the vessel is also doubled, so the flow decreases by half (÷2) = 8

Solution to Question 20:

The given clinical scenario is suggestive of Sydenham's chorea, a complication of rheumatic fever. The choreic movements are caused by the destruction of D2 neurons of the striatum. The neostriatum consists of the caudate nucleus and putamen.

The image given below shows Sydenham's chorea in a patient.



Nomenclature of basal ganglia:

• Corpus striatum: Caudate nucleus + putamen+ globus pallidus

• Lentiform nucleus: Putamen + globus pallidus

• Neostriatum: Caudate nucleus + putamen



Coronal section of human brain



Physiology AIIMS 2018

Question 1:

Which of the following is the diluting segment of the kidney?

a) PCT

b) Collecting duct

c) Ascending thick loop of Henle

d) Descending loop of Henle

Question 2:

Identify the ions A and B based on conductance graph of a neuron shown below:



a) Na, K

b) K, Na

c) Na, Ca

d) Ca, Na

Question 3:

How is an electrical synapse different from a chemical synapse?

a) No cytoplasmic continuity between presynaptic and postsynaptic membrane

b) Direction of impulse is unidirectional

c) Connexons are involved in electrical synapse

d) Electrical synapse has more latent period than chemical synapse

Question 4:

Hormone degradation as an important mechanism for regulating local concentrations is seen in _____.

a) 11β-hydroxysteroid dehydrogenase inactivation of glucocorticoids

b) Thyroid hormone deiodinases inactivating T3

c) Degradation of retinoic acid by Cyp26b1 in primordial germ cells of males

d) All of the above

Question 5:

The factors that can cause a person to drink water due to stimulation of thirst mechanism is/are _____.

a) Dehydration

b) Hemorrhage

c) Prandial Drinking

d) All of the above

Question 6:

On performing a histamine test on an individual with brachial plexus trunk injury, you can expect _____.

a) Central capillary dilatation → wheal → Surrounding flare

b) Central capillary dilatation → wheal without surrounding flare

c) Central capillary dilatation → wheal with exaggerated surrounding flare

d) Absent triple response

Question 7:

GFR is increased by all except

a) Increased renal blood flow

b) Efferent arteriolar constriction

c) Ureteric stone

d) Decreased oncotic pressure

Question 8:

Baroreceptor provides _____

a) Negative feedback

b) Positive feedback

c) Both negative and positive feedback

d) Feedforward control

Question 9:

The diagram below shows the pressure-volume curves for 3 different lungs. Which of the following is the best representation of compliance between the three lungs?



a) A<B<C

b) B>A>C

c) B>C>A

d) C>A>B

Question 10:

Identify the site which is responsible for relieving pain when pressure is applied to the painful area.



©MARROW

a) A

b) B

c) C

d) D

Question 11:

Inhibitory control by the hypothalamus is exerted on:

a) Prolactin only

b) Prolactin and growth hormone

c) Growth hormone only

d) Neither

Question 12:

Which of the following hormones play a permissive role in the attainment of puberty?

a) GnRH

b) Leptin

c) Insulin

d) GH

Question 13:

Weber-Fechner law relates magnitude of sensation to ____.

a) Strength of muscle contraction

b) Log of Intensity of stimulus

c) Amplitude of action potential

d) Duration of stimulus

Question 14:

The following Darrow Yannet diagram is suggestive of which condition? Note : The solid line is the original fluid compartment and the colour shaded compartment within the dotted line denotes the change caused by the condition.



a) Excessive sweating

b) Adrenal Insufficiency

c) Diarrhoea

d) SIADH

Question 15:

The mechanism by which glucose is absorbed in the intestine is:

a) Primary active transport

b) Secondary active transport

c) Facilitated diffusion

d) Any of the above

Question 16:

The earliest stage at which erythropoietin is effective in red blood cell formation is?

a) Burst forming units

b) Colony forming units

c) Late erythroblast stage

d) Normoblast stage

Question 17:

Integrin connects with:

a) Cadherin

b) Vitronectin

c) Fibronectin

d) Selectin

Question 18:

Which of the following organelles is absent in the area of the cardiac muscle marked:



a) Mitochondria

b) Lysosome

c) Golgi bodies

d) Endoplasmic reticulum

Question 19:

Which of the following fibres reach the Purkinje cell directly?

a) Inferior olivary nucleus

b) Raphe nucleus

c) Vestibular nucleus

d) Locus coeruleus

Question 20:

Calculate the net filtration pressure across the GBM for a patient with Glomerular ultrafiltration coefficient = 6 Glomerular hydrostatic pressure =45 Glomerular capillary oncotic pressure =20 Bowman's capsule hydrostatic pressure =10 Bowman's capsule oncotic pressure =0

a) 90 mmHg

b) 60 mmHg

c) 30 mmHg

d) 15 mmHg

Question 21:

In a 100-m dash, most of the energy consumed by skeletal muscles to replenish ATP is derived from _____

a) Phosphocreatine

b) Aerobic glycolysis

c) Oxidation of Fatty acids

d) None of the above

Question 22:

In the homunculus, which of the following has maximum representation?

a) Low control and skilled movements

b) More muscle fibres are present

c) Increased preciseness of movements

d) None of the above

Question 23:

What is the usual position of the Golgi apparatus in serous acinar cells?

a) Supranuclear

b) Basal

c) Adjacent to nucleus

d) Apical

Answer Key

Question No.	Correct Option
1	c
2	a

3	c
4	d
5	d
6	b
7	c
8	a
9	b
10	a
11	b
12	b
13	b
14	b
15	b
16	a
17	c
18	b
19	a
20	d
21	a
22	c
23	a

Detailed Explanations

Solution to Question 1:

The ascending thick limb of the loop of Henle is called the diluting segment.

In the ascending limb, there is the movement of Na+ and Cl– out of the tubular lumen, and the tubular fluid becomes more dilute.

When the fluid reaches the top of the ascending limb, it becomes hypotonic to plasma.

Solution to Question 2:

Channel A: Na+, Channel B: K+

Na+ channels are fast activating and are responsible for depolarization phase of action potential.

K+ channels are slow activating and open for prolonged duration than the sodium channels. They are responsible for repolarization phase of action potential. Also, the postassium channels return slowly to their closed state, which is responsible for after-hyperpolarization.



The sequence of events during the action potential occur in 7 steps as described below.

1. Resting state - Both channels are closed

2. Na+ channels open in response to depolarizing stimulus. Sodium enters into the cell. The membrane reaches threshold potential.

3. The sodium influx causes opening of more sodium channels through positive feedback mechanism. Rapid depolarization occurs.

4. Sodium conductance decreases.

5. Na+ channels close and K+ channels open. Repolarization occurs.

6. Sodium channels are reset, and are ready to respond to another depolarizing stimulus (relative refractory period). During this phase, the potassium channels remain open and slowly return to closed state causing after-hyperpolarization.

7. Both channels are closed. The membrane returns to RMP.



Solution to Question 3:

In electrical synapses, the cytoplasm of adjacent cells is directly connected by gap junctions (connexons) that allow free movement of ions from the interior of one cell to the interior of the next cell.

The transmission in electrical synapses is bidirectional and there is no synaptic delay. Unlike chemical synapses, electrical synapse has cytoplasmic continuity.

Note:

Most of the synapses used for signal transmission in the central nervous system are chemical synapses.

Solution to Question 4:

Hormone degradation as an important mechanism for regulating local concentrations is seen in all of the above.

11β-hydroxysteroid dehydrogenase inactivates glucocorticoids in renal tubular cells, preventing actions through the mineralocorticoid receptor.

During development, degradation of retinoic acid by Cyp26b1 prevents the primordial germ cells in the male from entering meiosis, which occurs in the female ovary.

The thyroid hormone deiodinases convert T4 to T3 and can inactivate T3. All the three are hence examples of local hormone degradation being utilized as a method for regulation of hormone concentration locally.

Solution to Question 5:

Dehydration, hemorrhage, and prandial drinking: All three can cause increased thirst resulting in water drinking.

Increased osmolality acts via osmoreceptors in the anterior hypothalamus.

The decreased plasma volume, as in hemorrhage, acts via

1. Renin-angiotensin system

2. Baroreceptors in heart and blood vessels.

The intake of liquids during a meal is known as prandial drinking. The increase is known as learned or habit response but is not understood completely.

The following factors are thought to play a role

1. Increase in plasma osmolality due to food absorption

2. Action on hypothalamus by one or more gastrointestinal hormones

Solution to Question 6:

Brachial plexus trunk injury leads to anesthetized skin over the region. On performing a histamine test on such an individual, one can expect central capillary dilatation → wheal without surrounding flare. The incomplete triple response over the anesthetic skin is characteristic of postganglionic lesions.

Intradermal injection of histamine usually causes a triple response in the surrounding skin i.e, red reaction→ flare → wheal

• Red reaction is independent of innervation and is due to histamine-mediated relaxation of the precapillary sphincters

• Flare reaction (Flush, Axon reflex) is due to Substance P mediated arteriolar dilatation (antidromic conduction of afferent impulses via collaterals of the sensory nerve fiber that lead to the release of vasoactive mediators)

• Wheal (skin edema) is due to histamine-mediated increased capillary permeability

• The triple response persists after total sympathectomy

• Flare is absent in locally anesthetized skin

Solution to Question 7:

Ureteric stone causes a reduction in glomerular filtration rate (GFR).

Following ureteral obstruction, there is a progressive fall in glomerular filtration rate (GFR) due to a reduction in a single-nephron glomerular filtration rate (SNGFR) and a reduced number of filtering nephrons.

All the other factors/options increase GFR.

Solution to Question 8:

Baroreceptors provide negative feedback in response to an increase in blood pressure.

Increased baroreceptor discharge inhibits the tonic discharge of sympathetic nerves and excites the vagal innervation of the heart.

Baroreceptor reflex

• Increased BP → stimulation of baroreceptors → stimulation of NTS → inhibition of RVLM → inhibition of sympathetic system

• Stimulation of NTS also stimulates CVC → parasympathetic stimulation

• Net effect = sympathetic inhibition and parasympathetic stimulation → decreased HR and BP

NTS - Nucleus tractus solitarius

CVC - Cardiovagal center

RVLM - Rostral ventrolateral medulla

HR- Heart rate

BP - Blood pressure

Solution to Question 9:

• The pressure-volume relationship is the fundamental relation between the work of breathing and lung mechanics.

• The diagram depicts the change in volume with respect to the change in intrapleural pressure, which is compliance. Here, the change is more (steep curve) with B and least (flat curve) with C.

• The “stiffer” lung displays a more horizontal curve (C), meaning that it takes a greater pressure to achieve the same inflation volume compared with a normal lung (A). These lungs are called less compliant.

• Curve (B) is more compliant as it takes lesser pressure to achieve the same inflation volume compared to normal (A).

Applied aspect:

Compliance is increased in emphysema. The mechanism is shown in the flow chart below.



Compliance is decreased in case of pulmonary edema or fibrosis.



Solution to Question 10:

It is a cross section of brain stem at pons.

• A is a periaqueductal gray matter which also consists of raphe nucleus.

• Descending pathways from the mesencephalic periaqueductal gray matter inhibit transmission in nociceptive pathways. This descending pathway includes a synapse in the ventromedial medulla (Raphe nucleus) and the release of endogenous opiates. So periaqueductal gray matter releases endogenous opiates.

Other marked structures are

B: Corticospinal tract

C: Medial lemniscus

D: Medial longitudinal fasciculus

Solution to Question 11:

Inhibitory control by the hypothalamus is exerted mainly on prolactin and growth hormone.

The major hypothalamic releasing hormones are:

• Thyrotropin releasing hormone (TRH)

• Corticotropin releasing hormone (CRH)

• Growth hormone releasing hormone (GHRH)

• Gonadotropin releasing hormone (GnRH)

The major hypothalamic inhibitory hormones are:

• Prolactin inhibitory hormone or dopamine

• Growth hormone inhibitory hormone (GHIH), also called Somatostatin.

Solution to Question 12:

The hormone that plays a permissive role in the attainment of puberty is leptin.

It now appears that leptin may be the link between body weight and puberty. Leptin is a satiety-producing hormone secreted by adipocytes. High leptin levels reduce appetite and increase energy expenditure.

Evidence for which is given by:

• Obese ob/ob mice that cannot make leptin are infertile, and their fertility is restored by injections of leptin.

• Leptin treatment also induces precocious puberty in immature female mice.

• However, the way that leptin fits into the overall control of puberty remains to be determined.

Leptin also accelerates GnRH pulsatility in neurons, fastening puberty.

Solution to Question 13:

Weber-Fechner Law states that the magnitude of sensation felt is proportionate to the log of the intensity of the stimulus.

Other Laws in Physiology:

• Doctrine of specific nerve energies - When the nerve pathways from a particular sense organ are stimulated, the sensation evoked is that for which the receptor is specialized no matter how or

where along the pathway the activity is initiated.

• Law of projection - No matter where a particular sensory pathway is stimulated along its course to the cortex, the conscious sensation produced is referred to the location of the receptor. This principle is called the law of projection.

Solution to Question 14:

In adrenal insufficiency there is hypo-osmotic volume contraction. Aldosterone deficiency leads to increased sodium loss (loss of hyperosmotic fluid).

• The osmolarity of ECF decreases.

• ECF volume decreases because of the loss of volume.

• Water shifts into ICF; as a result of the shift, ICF osmolarity decreases until it is equal to ECF osmolarity, and ICF volume increases.

• So, loss of hypertonic fluid causes hypotonic volume contraction.

Please note that in the diagram, the horizontal axis represents volume and vertical axis represents osmolarity.





The Darrow Yannet diagrams of various conditions are shown above.

Solution to Question 15:

Glucose is absorbed in the intestine by secondary active transport.

On the basolateral side, there is active transport of sodium ions out of the cell into the interstitial fluid. This depletes sodium inside the cell. Because of this Na+ concentration difference, more Na+ binds to the SGLT on the luminal side. This increases the affinity of SGLT for glucose and glucose binds to it.

Then, both the Na+ and glucose are transported together to the interior of the cell. Thus, glucose is absorbed in the intestine by secondary active transport. The energy for glucose transport is provided indirectly, by the active transport of Na+ out of the cell. The glucose moves out of the cell into the blood across the basolateral membrane by facilitated diffusion mediated via GLUT.



Solution to Question 16:

BFU is the earliest stage of hematopoiesis where erythropoietin (EPO) acts

Under the influence of EPO, both BFU and CFU grow in semisolid culture media and give rise to colonies of well-hemoglobinized erythroblasts.

Erythroid burst-forming unit (BFU-E) is the most primitive single lineage committed erythroid progenitor. In addition to EPO, they form the sub-population of Erythroid Colony Forming Units (CFU-E).

CFU is derived from burst-forming units.

Note: While CFUs have the highest number of receptors for EPO, the earliest stage at which it acts is on the BFU.

Solution to Question 17:

Integrin connects with fibronectin.

Integrins are transmembrane glycoproteins (integral proteins) that allow cells to attach to extracellular matrix constituents such as laminin, fibronectin and collagen. It helps in linking the intracellular cytoskeleton functionally and structurally with the outside world.

Solution to Question 18:

The area of the cardiac muscle marked is devoid of nuclei. Lysosomes are commonly observed in perinuclear areas. Other organelles are present in abundance throughout the cell.

Solution to Question 19:

The fibres of inferior olivary nucleus (climbing fibres) reach the purkinje cell directly.

Note:

The input from the Purkinje cells to the deep nuclei is inhibitory.

However, the deep nuclei output is always excitatory.

Solution to Question 20:

Net filtration pressure = Hydrostatic pressure in the glomerular capillaries - Pressure in Bowman's capsule - Oncotic pressure in the glomerular capillaries.

Net filtration pressure= [PG -PB- πG+ πB]

Therefore,

Net filtration pressure = (45-10-20+0)

= 25-10

= 15 mmHg

Note:

Here, just the net filtration pressure is asked and not the GFR.

GFR= Kf [PG -PB- πG+ πB]

Solution to Question 21:

In a 100-m dash that takes 10 s, 85% of the energy consumed is derived anaerobically by phosphocreatine, which is hydrolyzed to creatine and phosphate groups with the release of considerable energy.

The ATP concentration in a muscle is low enough (4 millimolar) to sustain contractions for 1-2 seconds after which ATP is dephosphorylated into ADP. The ADP is re-phosphorylated by phosphocreatine for the next 5 to 8 seconds.

ADP is re-phosphorylated by 3 sources.

• Phosphocreatine after 2 seconds upto 10 seconds.

• Glycolysis (anaerobic) after 10 seconds upto a minute

• Oxidative metabolism - after a minute upto 2-4 hours.

Solution to Question 22:

In a homunculus, the more the representation of an area, the more precise the movements are.

• The relative sizes of the cortical receiving areas of the sensory homunculus are proportionate to the use of the part.

• The sizes of these areas are directly proportional to the number of specialized sensory receptors in each respective peripheral area of the body.

• Some body parts are represented by large areas in the somatic cortex—the lips the greatest of all, followed by the face and thumb, whereas the trunk and lower part of the body are represented by relatively small areas.



Solution to Question 23:

Golgi complex is located in supranuclear cytoplasm in serous acinar cells. Serous acinar cells are generally found in pancreas and the submandibular salivary glands.

Physiology AIIMS 2019 (May & Nov)

Question 1:

The function of the Golgi tendon organ is to _____

a) Sense changes in muscle direction

b) Sense changes in muscle tension

c) Sense changes in position of the muscle

d) Sense changes in length of the muscle

Question 2:

Which is the first-order neuron in the optic pathway?

a) Rods and cones

b) Ganglion cells

c) Cells in lateral geniculate body

d) Bipolar cells

Question 3:

Pacemaker generating the rhythm for breathing is _____

a) Pneumotaxic centre

b) Dorsal group of neurons in the medulla

c) Pre-B■tzinger complex

d) Apneustic centre

Question 4:

Which of the following is not an example of feed forward control mechanism?

a) Temperature Regulation

b) Cephalic phase of gastric secretion

c) Cerebellar action in motor coordination

d) Exercise resulting in heart rate increase

Question 5:

Single breath nitrogen method is used for measuring ____

a) Dead space volume

b) FRC

c) Tidal volume

d) Diffusion capacity

Question 6:

I-125 labelled albumin is used for measurement of:

a) Plasma volume

b) Interstitial fluid volume

c) ICF

d) ECF

Question 7:

Which of the following hormones is exclusively secreted by adrenal gland but not by the ovaries of females?

a) Progesterone

b) DHEA

c) Estrogen

d) DHEA-S

Question 8:

Which hormone crosses the plasma membrane for its action?

a) Epinephrine

b) Thyroxine

c) Glucagon

d) Insulin

Question 9:

Afferent fibers from baroreceptors for blood pressure control goes to?

a) Nucleus ambiguous

b) Rostral-ventro lateral medulla

c) Nucleus Tractus Solitarius

d) Caudal ventrolateral medulla

Question 10:

If the lung is allowed to undergo elastic recoil without the chest wall, the resultant volume of the lung would be?

a) Minimum volume

b) Zero

c) Functional residual capacity

d) Residual volume

Question 11:

Myosin head detaches from actin by which of the following mechanism?

a) Binding of ATP

b) Entry of calcium into sarcoplasmic reticulum

c) Change in troponin C configuration

d) Release of ADP and Pi

Question 12:

You are an intern and want to determine the quantity of IV fluids needed for a patient. How much amount of transient loss/insensible loss per day would you account for in your calculations?

a) 150-200 ml

b) 500-600 ml

c) 200-350 ml

d) 350-400 ml

Question 13:

A soldier posted in Siachen is brought to you in the ER. In order to rule out hypothermia, how do you best measure his temperature?

a) Oral

b) Rectal

c) Infrared tympanic thermometer

d) Axillary

Question 14:

"Vasopressin escape" in SIADH results due to:

a) Downregulation of aquaporin 2 production

b) Increase in urine osmolarity

c) Decrease in ADH secretion

d) Upregulation of aquaporin 2 production

Question 15:

Which of the following hormones do not act on the pancreas?

a) Secretin

b) Gastrin

c) Cholecystokinin

d) GIP

Answer Key

Question No.	Correct Option
1	b

2	a
3	c
4	d
5	a
6	a
7	d
8	b
9	c
10	a
11	a
12	b
13	b
14	a
15	b

Detailed Explanations

Solution to Question 1:

Golgi tendon organ sense changes in muscle tension.

Golgi tendon reflex (Inverse myotatic/ Inverse stretch reflex) is disynaptic, is the opposite of the stretch reflex.

When the tension in a contracted muscle is significantly elevated, the inhibitory effect from the Golgi tendon organ leads to instantaneous relaxation of the entire muscle. This effect is called the lengthening reaction.

It is a protective mechanism to prevent tearing of the muscle or avulsion of the tendon from its attachments to the bone.

Note: Muscle spindles senses changes in muscle length (stretch).





Solution to Question 2:

Photoreceptors - rods and cones are the first-order neurons in the visual pathway.

Solution to Question 3:

Pre-Botzinger complex generates the rhythm for automated breathing.

- These are also known as pacemaker cells located in the medulla.
- These are the first set of neurons that initiate inspiration.

• These neurons discharge rhythmically, and they produce rhythmic discharges in the phrenic motor neurons.

Breathing is under the control of both the following:

• Voluntary control: Located in the cerebral cortex (via corticospinal tract)

• Automatic control: Driven by pacemaker cells in the medulla (Pre-Botzinger complex). These activate motor neurons via cervical (Phrenic nerve innervating diaphragm) and thoracic spinal cord (external intercostal muscle).

There is a reciprocal innervation to expiratory and inspiratory muscles in that motor neurons supplying expiratory muscles are inactive when motor neurons supplying inspiratory muscles are active, and vice versa.

Inspiration	
Initiation	Pre-Botzinger complex (Pace maker cells) in Medulla
Maintenance	Apneustic center in lower Pons stimulates Dorsal group of neurons in the medulla
Termination	Apneustic center inhibited by Vagus and Pneumotaxic center

Regulation of expiration: During quiet breathing, expiration is a passive process; there is no firing of neurons. Ventral respiratory group of neurons are involved in forceful expiration like during exercise.

Mnemonic DIVE: Dorsal respiratory group for Inspiration and Ventral respiratory group for Expiration.

Solution to Question 4:

Exercise resulting in heart rate increase is not an example of feedforward control mechanism. However, thinking about exercise and it resulting in a heart rate increase would be an example of feedforward control mechanism.

Feedforward control is that which makes an anticipatory or adaptive response without a specific stimulus.

Temperature control - Thermoregulatory responses are initiated by the hypothalamus before the changes in environmental temperature changes the core body temperature. Thus, shivering in response to cold environment would be an example of feedforward control.

In the cephalic phase of gastric secretion, the thought of food itself and not food per se produces a gastric secretion.

The cerebellum has a feed-forward inhibition that limits the duration of the excitation produced by any given afferent volley.

Solution to Question 5:

Anatomical dead space can be measured by the single-breath N2 curve (Fowler's method).

• Anatomical dead space is the respiratory system volume exclusive of alveoli.

• Alveolar dead space is the volume of gas not equilibrating with blood i.e. wasted ventilation.

• Physiological/total dead space is the sum of anatomical and alveolar dead space.

• The equation used for the calculation of total/physiological dead space is the Bohr's equation.



The patient takes a deep breath of O2 From mid-inspiration, then exhales steadily. The changes in the N2 concentration of expired gas during expiration are plotted.

Region I - Dead Space (DS)

From I–II is a mixture of DS and alveolar gas.

The transition from III-IV is the closing volume (CV).

End of IV - Residual Volume (RV).

Factors affecting dead space:

Option B: Functional Residual Capacity (FRC) is measured using body plethysmography.

Option C: Tidal volume (TV) is measured using spirometry.

Option D: Diffusion Capacity is measured using carbon monoxide diffusion capacity (DLCO).

Factor	Effect
Posture:UprightSupine	IncreaseDecrease
Position of airway:Neck ExtensionNeck Flexion	IncreaseDecrease

Factor	Effect
Age	Increase
Artificial airway	Decrease
Positive pressure ventilation	Increase
Drug anti-cholinergic	Increase
Pulmonary perfusion:Pulmonary embolismHypotension	IncreaseIncrease
Pulmonary disease:Emphysema	Increase

Solution to Question 6:

I-125 labelled albumin is the most commonly used substance for the measurement of plasma volume.

Solution to Question 7:

DHEAS is exclusively secreted from the adrenals but not by ovaries because ovaries lack the enzyme dehydroepiandrosterone-sulphotransferase needed to form DHEAS.

90% of DHEA is synthesised by the adrenals and 10% comes from the ovaries.

Solution to Question 8:

Thyroxine crosses the plasma membrane for its action.

Thyroid hormones enter the nucleus and bind to thyroid hormone receptors. The hormone-receptor complex in the nucleus binds to DNA via zinc fingers. This increases the expression of a variety of different genes that code for proteins regulating cell function. Although thyroxine is an AA derivative, there are transporters to move it into the cell, hence they are intracellular receptors acting on the nucleus.

Epinephrine, glucagon, and insulin bind to cell surface receptors.

Solution to Question 9:

Afferent fiber for blood pressure control goes to the Nucleus Tractus Solitarius (NTS).

Increased BP stimulates the baroreceptors which provide negative feedback in response to the increase in blood pressure.

Baroreceptor reflex:

• Increased BP → stimulation of baroreceptors → stimulation of NTS → excitation of CVLM → inhibition of RVLM → inhibition of sympathetic system

• Stimulation of NTS also stimulates CVC → parasympathetic stimulation

• Net effect = sympathetic inhibition and parasympathetic stimulation → decreased HR and BP

NTS - Nucleus tractus solitarius

CVC - Cardiovagal centre

CVLM - Caudal ventrolateral medulla

RVLM - Rostral ventrolateral medulla

HR - Heart rate

BP - Blood pressure

Solution to Question 10:

If lung is allowed to undergo elastic recoil without the chest wall, the resultant volume of the lung would be minimum volume as the opposing force by the chest wall would be non existant.



The yellow line in the above graph denotes the compliance of lungs alone. According to the graph, when the pressure (x axis) inside the lung is zero, there is some volume of air lesser than FRC still present (y axis) inside the lungs which is less than the functional residual capacity and residual volume which can be called as minimum volume.

Solution to Question 11:

Binding of ATP on the free site of myosin leads to the detachment of myosin head from the thin filament.



Solution to Question 12:

500-600 ml of transient loss/insensible loss per day would have to be accounted for in the calculation.

Although water is lost through sweating especially in hot climates or during physical exertion, we tend to lose water at rest as well. Physiologically, water evaporates from the skin and lungs at a rate of about 600 to 700 ml/day. This is known as insensible water loss.

Solution to Question 13:

In the given clinical scenario rectal temperature would be the preferred site to measure the core body temperature.

During low or high temperatures like in Siachen, the axillary and the oral temperatures are highly misleading and should not be used to measure the core body temperatures

Infrared tympanic thermometer measures the temperature of the eardrum which is representative of core body temperature. However, more often the thermometer is not inserted adequately or for safety reasons, they are manufactured in such a way that they do not reach deep into the ear canal, or the ear canal contains cerumen which can disturb the measurement. Hence the infrared thermometer more commonly delivers the temperature of ear canal cerumen and hence not a reliable method of measuring the core body temperature.

Solution to Question 14:

Vasopressin escape in SIADH (Syndrome of inappropriate ADH secretion) results due to downregulation of aquaporin 2 production.

Prolonged exposure to elevated levels of vasopressin (ADH) leads eventually to downregulation of aquaporin-2 receptors. This prevents excessive reabsorption of water, despite high serum ADH levels and hence maintains euvolemia. This is known as vasopressin escape. This helps prevent excessive dilutional hyponatremia in SIADH.

Solution to Question 15:

Gastrin does not act upon the pancreas. Gastrin stimulates gastric acid secretion and gastric mucosal growth.

GI hormones acting on the pancreas include:

GIP is also known as Glucose-dependent insulinotropic peptide.

	CCK(Cholecystokinin)	Secretin	GIP(Gastric inhibitory peptide)
Stimulates	Pancreatic enzyme secretion Pancreatic bicarbonate secretionGallbladder contractionGrowth of exocrine pancreas	Pepsin secretionPancreatic bicarbonate secretionBiliary bicarbonate secretionGrowth of exocrine pancreas	Insulin release
Inhibits	Gastric emptying	Gastric acid secretion	Gastric acid secretion

Physiology AIIMS 2020 (May and Nov INI-CET)

Question 1:

A 65-year old patient visits his family practitioner for a yearly examination. On examination, his BP was 190/100 mm Hg. His pulse pressure was 90 mm Hg. What would have decreased resulting in this increase in pulse pressure?

a) Myocardial contractility

b) Stroke volume

c) Arterial wall compliance

d) Cardiac output

Question 2:

Which of the following curves represents gamma-globin chain production?



a) Green Curve

b) Black Curve

c) Blue Curve

d) Orange curve

Question 3:

Transcutaneous electric nerve stimulation is used to reduce pain. What is the mechanism behind this?

a) Central inhibition

b) Desensitization

c) Gate control theory

d) Descending inhibition

Question 4:

Which of the following is not required for calculation of creatinine clearance?

a) Urine creatinine

b) Serum creatinine

c) 24 hour urine output

d) Volume of serum passing through kidney in 24 hours

Question 5:

A neuron is receiving 400 inputs from 150 sensory neurons. This is an example of?

a) Divergence

b) Convergence

c) Feedback inhibition

d) Feedforward excitation

Question 6:

Calculate the ECF volume in a volunteer in whom 10 gm of mannitol was injected in venous blood. At equilibrium urinary excretion of mannitol was found to be 10% and the plasma concentration of mannitol was 65 mg/100 ml.

a) 14L

b) 10L

c) 28L

d) 20L

Question 7:

Gibbs-Donnan equilibrium is mainly due to?

a) Phosphate

b) Bicarbonate in the intracellular compartment

c) Calcium in the extracellular compartment

d) Intracellular proteins

Question 8:

Which of these hormones have intracellular receptors?

a) Thyroxine

b) Epinephrine

c) Glucagon

d) Parathormone

Question 9:

Intercellular communication in eukaryotes is by?

a) Lysosome

b) Peroxisome

c) Exosome

d) Genome

Question 10:

In skeletal muscle, dihydropyridine and ryanodine receptors are coupled by?

a) Chemically

b) Electrically

c) Chemically and then electrically

d) Mechanically

Question 11:

In negative feedback, feedback gain of infinity is seen in which of the following?

a) Blood volume control by kidney

b) Temperature control in hypothalamus

c) BP control by barorecptors

d) Infinite gain is not possible

Question 12:

The image below shows simple muscle twitch along with the time trace, recorded in a frog gastrocnemius muscle. Calculate the minimum tetanizing frequency. (The time tracer was made with a 100Hz tuning fork).



a) 25 per second

b) 15 per second

c) 17 per second

d) 20 per second

Question 13:

Least frequency of slow waves is seen in:

a) Duodenum

b) Ileum

c) Jejunum

d) Stomach

Question 14:

Macula densa is formed by which part of the nephron?

a) PCT

b) DCT

c) Afferent arteriole

d) Bowman's capsule

Question 15:

As filtrate flows through PCT, the concentration of all of the following decreases except:

a) Aminoacid

b) Glucose

c) Chloride

d) Bicarbonate

Question 16:

This is the graph depicting the filterability of dextran through the kidney, 1 representing complete filtration and 0 representing no filtration. Choose the substance A, B, C respectively.



©MARROW

a) Polycationic, Neutral, Polyanionic

b) Polyanionic, Neutral, Polycationic

c) Neutral, Polycationic, Polyanionic

d) Polyanionic, Polycationic, Neutral

Question 17:

Calculate the respiratory quotient (RQ) in a patient with 50kg body weight, if the CO2 exhaled is 200ml/min and O2 consumed is 250 ml/min.

a) 0.8

b) 1

c) 1.25

d) 1.5

Question 18:

Which of the following types of amino acids are present in the transmembranous part of the membrane receptor?

a) Amphipathic

b) Hydrophobic

c) Hydrophilic

d) None of the above

Question 19:

Which of the following is not a cause of secondary hyperparathyroidism?

a) Parathyroid adenoma

b) Lithium intake

c) Vitamin D deficiency

d) Chronic kidney disease

Question 20:

Which of the following events occur immediately when a person changes his/her posture from supine to standing?

a) Decrease in firing from baroreceptor

b) Vagal output to cardiac fibres increase

c) Mesenteric Vasoconstriction

d) Increase cardiac contractility

Question 21:

Which of the following statements is true about skeletal muscle contraction?

a) 2, 3, 4

b) 3 only

c) 1, 2, 4

d) 1, 3, 4

Question 22:

Which of the following statements are correct regarding mechanisms of depolarisation, repolarisation, and hyperpolarisation in a skeletal muscle action potential?

a) 2 and 4 are correct

b) 1, 3, and 5 are correct

c) 1, 4, and 5 are correct

d) 2, 4, and 5 are correct

Question 23:

Arrange the following events in the sequence in which they occur, after 'p wave' in ECG:

a) 1-2-3-4

b) 1-2-4-3

c) 3-1-2-4

d) 1-3-2-4

Question 24:

Which of the following increases in COPD?

a) 1 only

b) 1 and 4

c) 1 and 3

d) 1 and 2

Answer Key

Question No.	Correct Option
1	c
2	d
3	c
4	d
5	b
6	a
7	d
8	a
9	c
10	d
11	a
12	d
13	d
14	b
15	c
16	a
17	a
18	b
19	a
20	a
21	b
22	b
23	b
24	d

Detailed Explanations

Solution to Question 1:

Pulse pressure increases with age due to decreased arterial wall compliance.

With advancing age, there is an increase in systolic blood pressure. Whilst diastolic blood pressure falls after 60 years of age due to increased arterial stiffness (decreased compliance). As a result of these two changes, pulse pressure increases (↑PP = ↑SBP - ↓DBP).

Solution to Question 2:

In the given image, the Orange curve represents the gamma-globin chain production. Gamma globin chain is produced in the fetal period and drops to undetectable levels 6 months postpartum which is the pattern represented by the Orange curve.

Hemoglobin in blood has different variants throughout life, in different proportions. The type of hemoglobin essentially depends on the combination of globin chains. Adult hemoglobin is made of 2 alpha and 2 beta chains, whereas fetal hemoglobin is made of 2 alpha and 2 gamma-globin chains. Other combinations of chains can be found in embryonic hemoglobin.

Here,

• Gamma chain - Orange curve

• Alpha chain - Red curve

• Beta chain - Green curve

• Epsilon chain - Black curve

• Delta chain - Blue curve

Solution to Question 3:

Transcutaneous electrical nerve stimulation (TENS) used for pain relief is based on the gate theory of pain. This method uses electrodes to activate Aα and Aβ fibers near the injury to relieve pain.

Gate control theory is based on the inhibition of pain pathway in the spinal cord. Various sensory inputs from a particular dermatome reach the dorsal horn of the spinal cord. The pain is transmitted via A-delta and C fibers, whereas, A-beta fibers carry tactile sensations (touch, pressure) and A-alpha fibers carry proprioception.

When these large-diameter A-beta/A-alpha fibers are stimulated, they inhibit the pain pathway in the spinal cord. This inhibition is thought to result from the local lateral inhibition of interneurons in the spinal cord.

Gate control theory operates at the substantia gelatinosa of the dorsal horn of the spinal cord.

The given image depicts the gate control theory of pain:



Solution to Question 4:

The volume of plasma passing through the kidney in 24 hours is not used in the calculation of creatinine clearance.

The parameters used to calculate creatinine clearance (CCr) are:

• Urine creatinine

• Serum creatinine

• 24hr urine output

The formula used to calculate creatinine clearance (CCr) is:

$UCr \times V / PCr$ where,

UCr - Urine creatinine(mg/dl)

PCr - Serum creatinine(mg/dl)

V - Urine volume(ml/24 hrs)

Creatinine is a metabolite of muscle metabolism and is cleared from the body mostly by glomerular filtration. Therefore, the clearance of creatinine is used to assess GFR.

Urine collection for the measurement of creatinine clearance may be subject to practical constraints. Therefore, approximate changes in GFR can be obtained by measuring the serum creatinine concentration, which is inversely proportional to GFR.

Solution to Question 5:

A single neuron receiving 400 inputs from 150 sensory neurons is an example of convergence.

Convergence means signals from multiple inputs uniting to excite a single neuron. It can be of 2 types:

• Convergence from a single source

• Convergence from multiple sources

The importance of convergence is that neurons are almost never excited by an action potential from a single input terminal. However, action potentials converging on the neuron from multiple terminals provide enough spatial summation to bring the neuron to the threshold required for discharge.

Solution to Question 6:

ECF volume of the above volunteer is calculated to be 14 L.

The ECF volume can be calculated using the formula:

Where,

• V = ECF volume

• Amount injected = 10 gm

• Amount of mannitol excreted = 10% of 10 gm = 1gm

• Plasma concentration of mannitol = 65 mg/100mL = 0.65 gm/L

Calculation:

ECF Volume = (10-1)/0.65

= 13.8 L = approximately 14 L

Mannitol cannot cross cell membranes, hence, it will be distributed in ECF but not in ICF. Thus mannitol is a marker for ECF volume and therefore calculating the volume of distribution of mannitol gives the value of ECF volume.

Solution to Question 7:

Gibbs-Donnan equilibrium is mainly due to intracellular proteins.

Donnan and Gibbs showed that "When an ion on one side of a membrane cannot diffuse through the membrane, the distribution of other ions to which the membrane is permeable is affected in a predictable way".

According to this, in the presence of intracellular proteins, which are negatively charged non-diffusable ions, the distribution of other ions to which the membrane is permeable is affected in a predictable way (Donnan Effect) so that at equilibrium their concentration ratios are equal.

Solution to Question 8:

Thyroxine hormone act by binding to intracellular nuclear receptors.

All other hormones mentioned in the options bind to receptors on the cell membrane.

Solution to Question 9:

Intercellular communication in eukaryotes is by Exosome.

In the last decade, a class of small, heterogeneous, secreted vesicles, broadly termed as extracellular vesicles have been identified.

These extracellular vesicles have been implicated as a new and important mediator of cell-cell communication that likely contributes importantly to both normal and pathologic physiology.

These vesicles are enclosed by a lipid bilayer.

They are heterogeneous in size (30-2000 nm diameter), and are generated by at least two distinct mechanisms:

• Microvesicles are generated by budding from the plasma membrane of a source cell.

• Exosomes are generated from the multivesicular body (MVB).

Note: The terms microvesicle and exosome are often lumped together as simply exosomes.

Solution to Question 10:

In skeletal muscle, dihydropyridine and ryanodine receptors show mechanical coupling involving structural interaction between the two.

Dihydropyridine receptors (DHPRs) are voltage-gated Ca^{2+} channels in the T tubule membrane.

The ryanodine receptor (RyR) is a ligand-gated Ca^{2+} channel with Ca^{2+} as its natural ligand.

Skeletal muscle excitation-contraction (E-C) coupling is thought to occur by a "mechanical coupling" mechanism involving protein-protein interactions that lead to activation of the sarcoplasmic reticulum ryanodine receptor by the voltage-sensing transverse (T-) tubule dihydropyridine receptor.

Solution to Question 11:

In negative feedback, feedback gain of infinity could be seen in blood volume control by the kidney.

When the blood volume falls below the equilibrium, the renal–body fluid mechanism causes the blood volume returns back to the equilibrium point resulting in zero error and hence the gain of kidney in its ability to regulate blood volume is infinity (∞).

The degree of effectiveness with which a regulatory system maintains constant conditions is determined by the gain of the negative feedback system.

Gain= Correction/Error

Example: If an external or internal factor led to a decrease in mean blood pressure from 100 mm Hg to 80 mm Hg. Activation of a negative feedback mechanism i.e. baroreflex leads to a correction of blood pressure to 96 mm Hg. So in this case,

• Correction = 16 mm Hg

• Error = 4 mm Hg

• Gain = Correction/Error = 16/4 = 4

Note: A gain of 0 would imply a total failure of the regulatory system and a gain of infinity (∞) would indicate complete restoration of the regulated variable.

Solution to Question 12:

The tetanizing frequency of the muscle, based on the given diagram, is 20 per second.

Tetanizing frequency = 1/contraction period. It is the frequency at which individual muscle fibres twitch that fuse and cause sustained muscle contraction, where the contraction period is the time taken from latent period to maximum force of contraction.

In this case, the time tracer was taken using a 100Hz tuning fork. Which means the tracer recorded 100 waves per second.

Therefore, the duration of each wave in time tracer = 0.01 seconds.

In the image given in the question, between the point B (end of latent period) to point C (maximum force of contraction), there are about 5 waves.

Hence, the duration of contraction = 5 x 0.01 = 0.05 seconds.

Substituting the value in the formula, we get:

Tetanizing frequency = 1/0.05 sec = 100/5 = 20/sec.



Tetanization:

The features of muscle contraction can be demonstrated by eliciting single muscle twitches. The adding together of individual twitch contractions to increase the intensity of overall muscle contraction is Summation. Summation occurs in two ways:

• by increasing the number of motor units contracting simultaneously, which is multiple ■ber summations.

• by increasing the frequency of contraction, which is called frequency summation and can lead to tetanization.

When the frequency of a contraction reaches a critical level, the successive contractions eventually fuse together, and the whole muscle contraction appears to be completely smooth and continuous, this is called tetanization. Beyond this point, a further increase in the frequency, will not result in increased force of contraction.

Tetanization occurs in two ways:

• Complete tetanus occurs when no relaxation occurs between stimuli.

• Incomplete tetanus when periods of incomplete relaxation take place between the summated stimuli.

Solution to Question 13:

Among the given options, the least frequency of slow waves is seen in the stomach (4/min).

The Basal electrical rhythm (BER) also called slow-wave, is a rhythmic fluctuation of membrane potential in the gastrointestinal smooth muscle. It is initiated by the interstitial cells of Cajal. The membrane potential fluctuates between -65mV and -45mV.

The BER is seen in all sites of gastrointestinal smooth muscle except in the esophagus and proximal stomach.

The rate of BER in various sites is as follows:

• Stomach: 4/min

• Duodenum: 12/min

• Distal ileum: 8/min

• Cecum: 2/min

• Sigmoid colon: 6/min

The BER does not cause contraction of the smooth muscle. However, when the BER is superimposed by the spike potentials, it increases muscle tension.

Acetylcholine causes an increase in the number of spike potentials and muscle tension. Epinephrine causes a decrease in the number of spike potentials and muscle tension.



Muscle contraction can occur only when the motor stimulus is provided during the depolarisation part of the BER.

Thus, BER does not cause muscle contraction by itself, but controls the rate and organization sites of the muscle contractions, as in coordination of the peristaltic activity and setting the rhythm of segmentation.

Solution to Question 14:

Macula densa is formed by specialized tubular epithelial cells of the distal convoluted tubule.



The juxtaglomerular apparatus consists of the following parts:

• Macula densa: At the end of the thick ascending limb of Henle, and at the beginning of the distal tubule, the tubular epithelial cells are specialized to form the macula densa. They detect sodium

concentration of the fluid in the tubule (chemoreceptor). In response to elevated sodium, the macula densa cells trigger the contraction of the afferent arteriole, reducing the flow of blood to the glomerulus and the glomerular filtration rate. This happens as follows: When there is increased sodium, there is increased sodium-potassium ATPase activity which leads to increased ATP hydrolysis causing more adenosine formation. The adenosine receptors on the macula densa cells stimulate it to release more calcium, so that there is vasoconstriction of the afferent arteriole, thereby decreasing the GFR.

• Lacis cells: Extraglomerular mesangial cells situated between afferent and efferent arteriole.

• Granular cells/JG cells: Modified smooth muscle cells of the afferent arteriole, they secrete renin. Fall in pressure in afferent arteriole stimulates renin release (mechanoreceptor), which, in turn, increases blood pressure.

Solution to Question 15:

As the filtrate flows through the proximal convoluted tubule, the concentration of the chloride increases.

• The concentration of chloride in the early part of the pCT is ≈105 mEq/L.

• As the sodium is reabsorbed, it preferentially carries glucose, amino acids, and bicarbonate.

• This leaves behind the solute, that has a higher concentration of the chloride, which as it reaches the second half of the PCT is ≈140 mEq/L.

Solution to Question 16:

In the given graph, the order of filterability is A>B>C, which likely indicates that the substances are Polycationic, Neutral, Polyanionic.

The glomerular filtration barrier is negatively charged, which will repel most of the negative charged (anionic substances). Whereas, the cationic substances are easily filtered.

Hence, the order of filterability would be: Polycationic > Neutral > Polyanionic, represented by A, B, and C in the graph shown.

The glomerular filtration barrier consists of:

• Endothelium of the capillary

• Basement membrane:

• Lamina densa - electron-dense layer. It contains type IV collagen

• Lamina rara interna - contains glycosaminoglycan heparan sulfate which has a negative charge

• Lamina rara externa - contains glycosaminoglycan heparan sulfate which has a negative charge

• Layer of epithelial cells (podocytes) surrounding the outer surface of the capillary basement membrane

The capillary endothelium is perforated by thousands of small holes called fenestrae, which measure from 4 nm up to 8nm. Therefore, neutral substances with molecular diameters of less than 4 nm are freely filtered, and filtration of those with diameters of more than 8 nm approaches zero.

However, in addition to the size of the particles, the filtration also depends mainly on the charge of the particle being filtered. The glomerular capillary wall consists of highly negatively charged proteins called sialoproteins, which repel the negatively charged substances (anions). The property can be studied by using charged dextran.

When the filtration of the negatively and positively charged dextrans was studied, the results indicated that the negatively charged dextrans in the blood are repelled, with the result that filtration of anionic substances 4 nm in diameter is less than half that of neutral substances of the same size.

This explains, why albumin despite having an effective molecular diameter of approximately 7 nm, has a very low filtration rate.

Solution to Question 17:

The respiratory quotient is 0.8.

The respiratory quotient is the ratio of carbon dioxide production to oxygen utilization.

In the given question, the CO_2 production is 200mL/min, while the O_2 consumption is 250mL/min.

Therefore,

Respiratory quotient = 200/250 = 0.8

The respiratory quotient (RQ) is used for determining the metabolic utilization of various foods.

• When carbohydrates are metabolized with one molecule of oxygen, one carbon dioxide molecule is released. Therefore, the RQ of carbohydrates = 1.

• When fats are oxidized, for every 100 molecules of oxygen consumed, 70 molecules of CO_2 are released. Therefore, the RQ of fats is 0.70.

• Similarly, the respiratory quotient for proteins is 0.80.

Solution to Question 18:

The transmembranous part of the membrane receptor consists of hydrophobic amino acids.

A membrane protein can be either integral protein (spans the thickness of the cell membrane) or peripheral protein (attached to the peripheral side of the membrane). The membrane-spanning segment of integral proteins are usually hydrophobic alpha-helices.

The central core of the cell membrane is hydrophobic. Integral proteins have their hydrophobic portions usually in the interior of the membrane and the hydrophilic portions towards the surfaces. The presence of these hydrophobic amino acids in the interior is a special adaptation

that allows the protein to remain stable in the hydrophobic milieu of the cell membrane.



Solution to Question 19:

Parathyroid adenoma is a benign tumor of the parathyroid gland, which causes primary hyperparathyroidism (not secondary).

Secondary hyperparathyroidism is characterized by high levels of parathyroid hormone, to compensate for the low levels of serum calcium, due to some other underlying condition.

Causes of secondary hyperparathyroidism, include:

• Chronic kidney disease: The damaged kidneys are unable to produce sufficient amounts of the active form of vitamin D, 1,25-dihydroxycholecalciferol. This further causes hypocalcemia and triggers PTH secretion.

• Vitamin D deficiency: Causes hypocalcemia, which triggers the increased secretion of PTH.

• Lithium: May cause an increased set-point of calcium for parathyroid hormone suppression, which leads to parathyroid hyperplasia.

Solution to Question 20:

There is a decreased firing from the baroreceptors.

• When a person stands up from a lying down position, it causes pooling of the blood in the legs, due to gravity. This reduces the venous return and causes a decrease in cardiac output.

• Reduced cardiac output, leads to a fall in blood pressure, which in turn causes reduced firing from baroreceptors.

• As a result, the vagal tone is suppressed leading to increased heart rate.

Solution to Question 21:

During the contraction of the skeletal muscle, the Ca2+ is released from the sarcoplasmic reticulum. Only statement 3 is correct.

Excitation contraction coupling refers to the process by which the membrane depolarization of the muscle fibre (excitation) leads to the contraction of the muscle.

The steps of excitation-contraction coupling in skeletal muscle are as follows:

• Motor-neuron discharge causes the release of acetylcholine at the motor endplate, which binds to the nicotinic acetylcholine receptors.

• This causes increased Na+ and K+ conductance in the endplate membrane, leading to the generation of end-plate potential, which is a local potential.

• When the local endplate potential reaches a threshold for the muscle fibre, and an action potential is generated.

• The generated action potential passes inward through the T tubules, leading to the spread of depolarisation.

• The depolarization of the muscle fibre causes the release of Ca2+ from terminal cisterns of sarcoplasmic reticulum and diffusion to thick and thin filaments.

• The calcium released, binds to troponin C, uncovering myosin-binding sites on actin.

• The formation of cross-linkages between actin and myosin and sliding of thin on thick filaments causes contraction of the skeletal muscle.

Steps in relaxation:

• The cytoplasmic calcium is pumped back into the sarcoplasmic reticulum, by the action of SERCA (sarcoplasmic endoplasmic reticulum calcium ATPase).

• Once the calcium concentration is lowered, the interaction between actin and myosin ceases, and the muscle relaxes.

Other options:

Statement 1: Incorrect - Extracellular calcium levels play an important role in muscle contraction. When there is a decrease in extracellular calcium as in hypocalcemia, the voltage-gated sodium channels are activated, which causes sodium influx. The increased intracellular sodium concentration causes sustained depolarization, which may lead to tetany.

Statement 2: Incorrect - Ca2+ is not released from mitochondria. It is released from the sarcoplasmic reticulum.

Statement 4: Voltage-gated calcium channels on T-tubules (part of sarcolemma) are activated by a depolarizing stimulus, which causes the release of calcium from the sarcoplasmic reticulum. The increased calcium concentration will eventually lead to muscle contraction.

Solution to Question 22:

Statements 1, 3, and 5 are correct.

Action potential in skeletal muscle:

• Motor-neuron discharge causes end-plate potential, which when reaches to a threshold value, causes action potential in the muscle fibre.

• The depolarization of the muscle fibre is caused by the activation of the voltage-gated sodium channels, which causes the influx of sodium.

• During repolarization, the sodium channels are closed, and voltage-gated potassium channels open, which causes efflux of potassium.

• The voltage-gated potassium channels close slowly. As a result, the potassium efflux continues, leading to hyperpolarization of the membrane.



The steps in an action potential in muscle fibre and the neuron, are similar except for the quantitative differences, such as:

• RMP is -80 to -90 mV in skeletal muscle fibres, while it is about -70mV in neurons.

• The duration of the action potential in skeletal muscle is 1 to 5 msec, which is about 5 times longer than in the largest myelinated nerve fibre.

• The velocity of conduction is about 3-5m/s, which is about 1/13th of the velocity of the nerve fibre.

Solution to Question 23:

The correct sequence of events that occur after the p-wave is 1-2-4-3, ie:-

a wave in JVP → First heart sound → T wave in ECG → Rapid filling of ventricles.

The sequence of the events in this question can be better understood with the following diagram:



• The p wave in the ECG represents atrial depolarization, which causes contraction of the atria, represented by the 'a' wave in the JVP. Atrial contraction causes final filling of the ventricles (fourth phase of ventricular diastole)

• Atrial contraction is followed by the closure of the mitral valve, which causes the first-heart sound.

• This is followed by the isovolumetric contraction of the ventricle, which causes c wave in JVP, due to bulging of the tricuspid valve. This is followed by the rapid ejection phase of the ventricle.

• The T wave of ECG occurs during ventricular repolarization, which is followed by the opening of the mitral valve, followed by rapid filling phase of the ventricle (second phase of ventricular diastole).

Note: Passive rapid filling occurs as the mitral valve opens just before atrial contraction occurs.

Solution to Question 24:

In COPD, the functional residual capacity increases. Total lung capacity may be increased or normal. Hence, both 1 and 2 are correct.

• In obstructive lung diseases, both FEV1 and FVC reduces, where the decrease in FEV1 is more than the reduction in FVC. As a result, the ratio of FEV1/FVC also decreases.

• The total lung capacity is usually normal in chronic bronchitis, while increased in emphysema.

• Functional Residual Capacity (FRC) is the volume of air present in the lungs at the end of passive expiration. In COPD, FRC is increased.

Lung Capacities:

The total lung capacity can be broken down into alternative capacities that help to define functioning lungs.

• The vital lung capacity refers to the maximum amount of air expired from the fully inflated lung or maximum inspiratory level.

• The inspiratory capacity is the maximum amount of air inspired from the end-expiratory level.

• The functional residual capacity represents the volume of the air remaining in the lungs after expiration of a normal breath. Functional Residual Capacity (FRC) is expiratory reserve volume + residual volume.

Physiology NEET 2018

Question 1:

The predominant organ producing insulin-like growth factor I is

a) Liver

b) Pituitary gland

c) Pancreas

d) Adrenal glands

Question 2:

What is Prosopagnosia?

a) Impairment of consciousness

b) Being unaware of one's problems

c) Difficulty in identifying known faces

d) Failure to identify objects

Question 3:

All of the following are components responsible for the countercurrent mechanism in the kidney except _____.

a) Sodium outflow in thick ascending limb

b) Water outflow in thin descending limb

c) Sodium outflow in thin ascending limb

d) Flow of tubular fluid from PCT to DCT

Question 4:

Glucose is absorbed in intestine by?

a) Primary active transport

b) Facilitated diffusion

c) Simple diffusion

d) Secondary active transport

Question 5:

In which of the following conditions is oxygen therapy not useful?

a) Asthma

b) Acute myocardial infarction

c) Pulmonary edema

d) Central respiratory depression

Question 6:

Va/Q = infinity means _____.

a) Partial pressure of O2 becomes zero

b) No exchange of O2 & CO2

c) CO2 alone equilibrates with the venous blood

d) Partial pressure of both CO2 and O2 remain normal

Question 7:

'c' wave in JVP is seen in

a) Iso-volumetric contraction

b) Slow filling at end of diastole

c) End of systole

d) Start of diastole

Question 8:

All of the following are true regarding the actions of endothelin-1 except _____

a) Bronchodilation

b) Vasoconstriction

c) Decreased GFR

d) Has inotropic effect

Question 9:

Aldosterone secretion is maximally stimulated by_____

a) Hyperkalemia

b) ACTH

c) Hypernatremia

d) Exogenous steroid

Question 10:

cGMP is the second messenger of

a) Somatostatin

b) Angiotensin II

c) ADH

d) NO (nitric oxide)

Question 11:

Transport molecule for Fe^{2+} into the enterocytes is

a) Ferroportin 1

b) Hephaestin

c) Divalent Metal Transporter 1

d) Divalent Metal Transporter 2

Question 12:

Nuclear signalling pathways are all except

a) Caveolins

b) NF-kB

c) Importins

d) Rat proteins

Question 13:

Which of the following statements is false about Cuthbertson's metabolic ebb and flow responses to trauma?

a) The "ebb" phase is associated with a decline in body temperature and oxygen consumption.

b) The "flow" phase occurs after resuscitation from a state of shock.

c) The "ebb" phase is aimed at reducing posttraumatic energy depletion.

d) The "ebb" phase leads to activation of the innate immune system and induction of the hepatic acute-phase response.

Question 14:

Kidney responds to shock by _____.

a) Increasing renal blood flow

b) Increasing afferent arteriole resistance

c) Increasing GFR

d) Both B and C

Answer Key

Question No.	Correct Option
1	a
2	c
3	c
4	d
5	d
6	b
7	a
8	a
9	a
10	d
11	c

12	d
13	d
14	b

Detailed Explanations

Solution to Question 1:

Growth hormone acts on the liver to form small proteins called “somatomedins”

• Somatomedins increases bone growth in all aspect

• Effects are similar to insulin, hence referred to as insulin-like growth factor (IGF).

• Four types available:

• Most important is somatomedin C, specifically referred to as “insulin-like growth factor-1 (IGF-1)

• Binds to carrier protein in blood

• Hence, longer duration of action than growth hormone.

• Half-life - about 20 Hrs (compared to growth hormone - 20 mins)

• Blood concentration of IGF-1, follows the levels of growth hormone.

• Growth effects of GH are mostly attributed to somatomedin (rather than its direct effect on bones & peripheral tissues)

Solution to Question 2:

Prosopagnosia is difficulty in identifying known faces.

In humans, storage and recognition of faces is more strongly represented in the right inferior temporal lobe in right-handed individuals, though the left lobe is also active. Damage to this area can cause prosopagnosia.

Patients with this abnormality can recognize forms and reproduce them. They can recognize people by their voices, and many of them show autonomic responses when they see familiar as opposed to unfamiliar faces. However, they cannot identify the familiar faces they see. The left hemisphere is also involved, but the role of the right hemisphere is primary.

The presence of autonomic response to a familiar face in the absence of recognition has been explained by postulating the existence of a separate dorsal pathway for processing information about faces that leads to recognition at only a subconscious level.

Note:

• Autotopagnosia - inability to recognize or to orient any part of one's own body

• Simultagnosia - inability to recognize multiple elements in a visual presentation, one object or some elements of a scene can be appreciated but not the display as a whole

• Somatagnosia - inability to identify any part of the body, either one's own or another's body

• Tactile agnosia (astereognosis) - inability to recognize objects by touch, in the presence of intact cutaneous and proprioceptive hand sensation; caused by a lesion in the contralateral parietal lobe

Solution to Question 3:

Thin ascending limb is not a component of the countercurrent mechanism.

The loop of Henle function as countercurrent multipliers and vasa recta as countercurrent exchangers.

The functioning of the loop of Henle depends upon:

• High permeability of the thin descending limb to water (via aquaporin-1)

• Active transport of Na+ and Cl- from the thick ascending limb

• Inflow of tubular fluid from proximal tubule and outflow to the distal tubule.

Note: The thin ascending limbs of juxtamedullary nephrons are permeable to Na+ and Cl- and hence do contribute passively to countercurrent multiplication. However, among the options provided, Option C is the best answer as the other options are directly responsible for the countercurrent mechanism.

Solution to Question 4:

Glucose is absorbed in the intestine by secondary active transport (sodium co-transport).

The energy for glucose transport is provided indirectly, by the active transport of Na+ out of the cell. This maintains the concentration gradient across the luminal border of the cell so that more Na+ and consequently more glucose enter.

Though both secondary active transport and facilitated diffusion play a role in glucose absorption, it is the initial active transport of sodium through the basolateral membranes of the intestinal epithelial cells that provide the eventual force for moving glucose through the membranes as well. So, secondary active transport is the single best answer for this question.

Solution to Question 5:

Oxygen therapy is not beneficial in central respiratory depression.

High dose oxygen reduces the output of peripheral chemoreceptors. So, if given to patients who already have central respiratory depression, it will indirectly reduce the hypoxic drive to breathe even further, leading to a reduction in respiratory frequency and minute ventilation.

Note: In acute severe asthma, a high concentration of oxygen should be given by face mask to achieve an oxygen saturation of >90%.

Solution to Question 6:

Va /Q = infinity means No exchange of O2 & CO2.

• Va represents the ventilation in alveoli.

• Q represents the blood flow through the alveolus.

• The ratio of Va and Q explains the respiratory exchange when there is an imbalance between alveolar ventilation and alveolar blood flow.

• Ventilation-perfusion ratio is considered normal if there is an equal amount of blood flow & ventilation through alveolus.

• When Va /Q ratio is infinity,

• There is adequate ventilation but no perfusion.

• No exchange of gases through the respiratory membranes of corresponding alveoli.

• The alveolar air equilibrates the humidified inspired air

• No exchange of oxygen and carbon-di-oxide occurs.

• Partial pressures of the O2 and CO2 are 149 mmHg & 0 mm Hg respectively

Solution to Question 7:

'c' waves are produced by bulging of the tricuspid valve into the right atrium during the Iso-volumetric contraction of the right ventricle.



Solution to Question 8:

Endothelin-1

• Potent vasoconstrictor agent.

• Produced by endothelial cells.

• They are secreted locally and act in a paracrine fashion.

Actions of Endothelin-1 (ET-1)

• Potent vasoconstrictor.

• Platelet activation.

• Inotropic effect.

• Closure of ductus arteriosus.

• Have a role in the pathogenesis of hypertension, myocardial infarction, and congestive heart failure.

• In the brain, regulate the transport of substances across blood brain barrier (BBB).

• In the kidneys, they are involved in tubuloglomerular feedback→ ↓ GFR.

• In the lungs, it causes bronchoconstriction (mediated via ET-B receptors).

• In GIT, they are involved in the migration of myenteric plexus to distal colon (failure leads to Hirschsprung's/megacolon).

• Deletion of both alleles of ET-1 gene→craniofacial abnormalities and respiratory failure at birth.

Solution to Question 9:

Hyperkalemia and activation of renin-angiotensin system are the most potent stimulators of aldosterone secretion.



Solution to Question 10:

Vasodilatory actions of NO are mediated by cGMP.

NO is a hormone derived from the amino acid arginine. It activates guanylyl cyclases in vascular smooth muscle cells resulting in the conversion of cGTP to cGMP (2nd messenger). cGMP activates cGMP dependent protein kinase, which causes blood vessels to relax.

cGMP is the second messenger for nitric oxide and atrial natriuretic factor.

Solution to Question 11:

Transport of Fe2+ into the enterocytes occurs via divalent metal transporter 1 (DMT1), through an active transport process.

Protein DMT-1 (Divalent Metal Transporter-1) is expressed on the apical surface of enterocytes in the initial part of the duodenum. DMT-1 is not specific to iron & transports other substances like zinc, copper, cobalt & manganese.



Solution to Question 12:

Rat proteins are not nuclear signalling pathways. Recombinant rat IFN- is a bioactive protein intended for use in cell culture applications.

Importin:

• Transporting protein molecules into nucleus by binding to specific recognition sequences, called Nuclear Localization Sequences (NLS).

• 2 subunits, importin α and importin β.

Caveolins:

• Family of integral membrane proteins that are principal components of caveolae membranes

• Involved in receptor-independent endocytosis.

• Acts as scaffolding proteins within caveolar membranes by compartmentalizing and concentrating signaling molecules.

NF-kB

• Part of a prototypical proinflammatory signaling pathway

• based on the role of NF-κB in the expression of proinflammatory genes including cytokines, chemokines, and adhesion molecules

Solution to Question 13:

The "flow" phase (not "ebb") leads to activation of the innate immune system and induction of the hepatic acute-phase response.

Cuthbertson's ebb and flow response to trauma:

The "ebb" phase lasts for a brief duration and is associated with a decline in body temperature and oxygen consumption, aimed at reducing post traumatic energy depletion.

The "flow" phase occurs after resuscitation from a state of shock, which leads to an increased metabolic turnover, activation of the innate immune system and induction of the hepatic acute-phase response. This hypercatabolic condition leads to a significantly increased oxygen consumption and energy expenditure.

Solution to Question 14:

Kidney responds to shock by increasing afferent arteriolar resistance.

During shock, both afferent and efferent arterioles in kidney are constricted, however, the efferent vessels are constricted much more.

The glomerular filtration rate and the plasma renal flow is decreased during shock. As plasma renal flow is decreased to a greater extent, the filtration fraction increases. (Option 1 & 3)

Physiology NEET 2019

Question 1:

According to myogenic hypothesis of renal autoregulation the afferent arterioles contract in response to stretch-induced by_____

a) NO

b) Noradrenaline

c) Opening of calcium channels

d) Adenosine

Question 2:

According to Herrington classification, the decerebrate rigidity is characterized by all except _____

a) Rigidity occurs in all the muscles of the body

b) Increase in the rate of discharge of the ■ efferent neuron

c) Increased excitability of the motor neuron pool

d) Decerebration produces no phenomenon akin to spinal shock

Question 3:

Which of the following is true for decorticate rigidity?

a) It is produced by the removal of cerebral cortex and basal ganglia

b) Flexion of lower limbs and extension of upper limbs occurs

c) Rigidity is pronounced

d) It is characterized by flexion of upper limbs and extension of lower limbs

Question 4:

Acid Citrate Dextrose (ACD), used to store blood, preserves the RBCs for a period of 21 days. The storage period when adding phosphate alone and when adding adenine and phosphate together to ACD would respectively be?

a) 35 days and 42 days

b) 21 days and 35 days

c) 28 days and 45 days

d) 35 days and 28 days

Question 5:

Functional residual capacity is:

a) Volume expired after normal inspiration

b) Volume that remains after forced expiration

c) Expiratory reserve volume + residual volume

d) Tidal volume + inspiratory reserve volume

Question 6:

Which of the following is not a dietary fiber?

a) Cellulose

b) Inulin

c) Pectin

d) Gums

Question 7:

Lesion in the medial lemniscus at the level of pons causes which of the following?

a) Pain and temperature loss in the same side

b) Pain and temperature loss in the opposite side

c) Loss of tactile and proprioception on the same side

d) Loss of tactile and proprioception on the opposite side

Question 8:

Which neurotransmitter is responsible for anterograde dilatation of the small intestine?

a) VIP

b) Acetylcholine

c) Substance P

d) Serotonin

Question 9:

Reflex responsible for tachycardia during right atrial distension is:

a) J reflex

b) Cushing reflex

c) Bezold jarisch reflex

d) Bainbridge reflex

Question 10:

Testosterone production is mainly contributed by _____

a) Leydig cells

b) Sertoli cells

c) Seminiferous tubules

d) Epidydmis

Question 11:

What is the function of tubuloglomerular feedback?

a) Regulate GFR

b) Decrease plasma sodium

c) Increase plasma volume

d) Decrease tubular secretion

Question 12:

Based on the agglutination test shown below, what is the blood group?



a) B positive

b) A positive

c) O positive

d) AB positive

Question 13:

Which of the following neurons predominantly fire during forceful expiration?

a) Dorsal group of neurons

b) Ventral group of neurons

c) Pneumotaxic center

d) Chemoreceptors

Question 14:

ADH acts through which aquaporin channel?

a) 1

b) 2

c) 3

d) 4

Question 15:

Which of the following has the least conduction velocity?

a) Postganglionic sympathetic fibers

b) Somatic motor fibers

c) Pre-ganglionic autonomic fibers

d) Touch and pressure fibers

Question 16:

Which part of the liver stores vitamin A?

a) Ito cells

b) Hepatocytes

c) Endothelial cells

d) Kupffer cells

Question 17:

Liver synthesizes all of the following except:

a) Albumin

b) Coagulation factors

c) Acute phase proteins

d) Immunoglobulins

Question 18:

Which of the following ions is involved in peripheral oxygen sensing chemoreceptors?

a) Potassium

b) Calcium

c) Sodium

d) Chlorine

Question 19:

Brown sequard syndrome is characterized by _____

a) Fine touch lost on the opposite side

b) Pain and temperature lost on the same side

c) Pain and temperature loss on the opposite side

d) Only pain lost on the same side

Answer Key

Question No.	Correct Option
1	c
2	a
3	d
4	b
5	c
6	d
7	d
8	a
9	d
10	a
11	a
12	c
13	b
14	b
15	a
16	a
17	d
18	a
19	c

Detailed Explanations

Solution to Question 1:

The afferent arterioles contract in response to stretch induced by the opening of calcium channels. The mechanism is called myogenic autoregulation of renal blood flow and GFR.

The stretch of the vascular wall allows increased movement of calcium ions from the extracellular fluid into the cells, causing them to contract. This contraction prevents excessive stretch of the vessel. This prevents excessive increase of renal blood flow and GFR by raising vascular resistance.

This mechanism may be more important in protecting the kidney from hypertension-induced injury.

Autoregulation has two main mechanisms:

• Metabolic mechanism: Tissues produce vasodilatory metabolites that regulate flow e.g. prostaglandins, NO, CO_2, H^+ and K^+

• Myogenic mechanism



Solution to Question 2:

In decerebrate rigidity, there is rigidity in extensor muscles of all the four extremities (not all muscles of the body).

Decerebration

• Complete transection of the brain stem between the superior and inferior colliculi permits the brain stem pathways to function independently of their input from higher brain structures. This is called a mid-collicular decerebration.

• This lesion interrupts all input from the cortex (corticospinal and corticobulbar tracts) and red nucleus (rubrospinal tract), primarily to distal muscles of the extremities.

• The excitatory and inhibitory reticulospinal pathways (primarily to postural extensor muscles) remain intact.

• The dominance of drive from ascending sensory pathways to the excitatory reticulospinal pathway leads to hyperactivity in extensor muscles in all four extremities which are called decerebrate rigidity.

• The excitatory input from the reticulospinal pathway activates γ-motor neurons which indirectly activate α-motor neurons (via Ia spindle afferent activity). This is called the gamma loop.

Option 4. Spinal shock is different from decebrate rigidity as it is a transient state that can occur immediately after a spinal cord injury. It is characterized by a temporary loss or depression of spinal reflexes below the level of the injury.

Solution to Question 3:

• Removal of the cerebral cortex (decortication) produces decorticate rigidity.

• Characterized by flexion of the upper extremities at the elbow and extensor hyperactivity in the lower extremities.

• The flexion can be explained by active rubrospinal excitation of flexor muscles in the upper extremities; the hyperextension of lower extremities is due to the same changes that occur after midcollicular decerebration.



Solution to Question 4:

The storage period when adding phosphate alone to ACD would be 21 days and when adding adenine and phosphate together would be 35 days.

When adding phosphate to ACD, Citrate Phosphate Dextrose (CPD) is formed – It has a storage period of 21 days. There is no change in the shelf life as phosphate only increases the 2,3 DPG levels of the stored blood.

When adding both phosphate and adenine to ACD, Citrate Phosphate Dextrose Adenine (CPDA-1) is formed. It has a storage period of 35 days. It increases the shelf life by improving ATP synthesis of RBCs.

Using additive solutions containing saline, adenine, and glucose, with or without mannitol (SAGM) decreases RBC hemolysis, thus increasing the storage period to 42 days.

Solution to Question 5:

Functional Residual Capacity (FRC) is expiratory reserve volume + residual volume.

From the picture given below, (FRC = ERV + RV).



Lung Capacities:

The total lung capacity can be broken down into alternative capacities that help to define functioning lungs.

• The vital lung capacity refers to the maximum amount of air expired from the fully inflated lung or maximum inspiratory level.

• The inspiratory capacity is the maximum amount of air inspired from the end-expiratory level.

• The functional residual capacity represents the volume of the air remaining in the lungs after expiration of a normal breath.

Solution to Question 6:

Gum is the best answer among the given options. Even though all of the given options are dietary fibers, the consumption of gums through diet is very low and hence, it is the best answer.

Dietary fibers consist of both soluble and insoluble forms and include lignin, pectin, inulin, cellulose, hemicellulose, etc.

Advantages of fibers:

• Forms bulk of the stool and reduces stool transit time

• Reduces incidence of carcinoma colon, DM, CAD, diverticulitis

Solution to Question 7:

Lesion in the medial lemniscus at the level of pons causes loss of tactile and proprioception on the opposite side.

Fibers of medial lemniscus ascend ipsilaterally in the dorsal columns of the spinal cord to the medulla, where they synapse in the gracilus and cuneate nuclei.

The second-order neurons from these nuclei cross the midline and ascend in the medial lemniscus to end in the contralateral ventral posterior lateral (VPL) nucleus and related specific sensory relay nuclei of the thalamus.

Solution to Question 8:

VIP is responsible for anterograde dilatation of the small intestine.

Peristalsis involves the contraction of a segment of intestine behind the bolus of food and relaxation of a segment of the intestine in front of the bolus of food.

Cholinergic neurons passing in anterograde direction produce NO and VIP which cause relaxation of a segment of the intestine in front of the bolus.

Cholinergic neurons passing in retrograde direction produce substance P and acetylcholine which cause contraction of a segment of the intestine behind the bolus.



Note: In achalasia cardia, there is decreased release of NO and VIP in the distal part of the esophagus.

Solution to Question 9:

Reflex responsible for tachycardia during right atrial distension is Bainbridge reflex.

Atrial Reflex Control of Heart Rate: The Bainbridge Reflex

• The stretch receptors of the atria that elicit the Bainbridge reflex transmit their afferent signals through the vagus nerves to the medulla of the brain.

• Then efferent signals are transmitted back through vagal and sympathetic nerves to increase heart rate and strength of heart contraction.

Bezold-Jarisch Reflex:

• Activation of chemosensitive vagal C fibres in the cardiopulmonary region (e.g. Juxta capillary region of alveoli, ventricles, atria, great veins, and pulmonary artery) causes profound brady-cardia, hypotension and a brief period of apnea followed by rapid shallow breathing.

Cushing Reflex:

• When ICP is increased, the blood supply to RVLM neurons is compromised.

• The local hypoxia and hypercapnia causes an increase in discharge from the neurons, producing hypertension, bradycardia and irregular respiration.

• The resultant rise in systemic arterial pressure tends to restore the blood flow to the medulla.

Solution to Question 10:

Testosterone production is mainly contributed by Leydig cells.

Testosterone is synthesized from cholesterol in the Leydig cells and also from androstenedione secreted by the adrenal cortex.

The diagnosis of testosterone deficiency should be made only on the basis of clinical signs and symptoms consistent with androgen deficiency, and consistently subnormal serum testosterone concentrations at 8 to 10 AM on three occasions.

Testosterone has been used to treat primary or secondary hypogonadism in adult men.

Solution to Question 11:

Tubuloglomerular feedback regulates glomerular filtration rate (GFR).

As the rate of ■ow through the ascending limb of the loop of Henle and first part of the distal tubule increases, glomerular filtration in the same nephron decreases, and, conversely, a decrease

in ■ow increases the GFR.

Solution to Question 12:

Based on the agglutination test shown in the image, the blood group is O positive.

As agglutination is present only in 'Rh' and not with 'A' and 'B' hence the blood group is O Rh positive.



Solution to Question 13:

Ventral groups of neurons fire, when forceful expiration is needed like during heavy exercise.

Dorsal group of neurons control the normal inspiration and expiration.

Pneumotaxic center limits the duration of inspiration.

Solution to Question 14:

The antidiuretic hormone acts through aquaporin channel 2.

ADH binds to V2 receptors in the late distal tubules, collecting tubules, and collecting ducts, and stimulates the movement of an intracellular protein, called aquaporin-2 (AQP-2), to the luminal side of the cell membranes.

Solution to Question 15:

Postganglionic sympathetic (type C) has the least conduction velocity among the given options.

Types of nerve fibers:

Fiber type	Function	Fiber diameter (μm)	Conduction velocity (m/s)
Aα	Proprioception, somatic motor	12-20	70-120
Aβ	Touch, pressure	5-12	30-70
Aγ	Motor to muscle spindles	3-6	15-30
Aδ	Pain, temperature	2-5	12-30
B	Preganglionic autonomic	<3	3-15
C, Dorsal root	Pain, temperature	0.4-1.2	0.5-2
C, Sympathetic	Post ganglionic sympathetic	0.3-1.3	0.7-2.3

Solution to Question 16:

Vitamin A is stored in the liver perisinusoidal stellate (Ito) cells as retinyl ester (retinol palmitate).

Kupffer cells, also known as stellate macrophages and Kupffer-Borowicz cells; are specialized macrophages located in the liver that form part of the mononuclear phagocyte system. These are present, lining the walls of the sinusoids, not the perisinusoidal space (Space of Disse).

Solution to Question 17:

Immunoglobulins are not synthesized by the liver. They are produced by plasma cells.

Plasma proteins synthesized by the liver are:

• Albumin

• Coagulation/clotting factors

• Acute-phase proteins

• Steroid binding and other hormone-binding proteins

Solution to Question 18:

Potassium is involved in peripheral oxygen sensing chemoreceptors.

Type I glomus cells (carotid and aortic body) have O2-sensitive K+ channels, whose conductance is reduced in proportion to the degree of hypoxia.

This reduces the K+ efflux, depolarizing the cell and causing Ca^{2+} in■ux.

The Ca^{2+} influx triggers action potentials and consequent excitation of the afferent nerve endings.

Solution to Question 19:

Hemisection of the spinal cord causes damage to ascending sensory (dorsal column pathway, ventrolateral spinothalamic tract) and descending motor (corticospinal tract) pathways, which is called the Brown- Séquard syndrome.

The lesion to fasciculus gracilus or fasciculus cuneatus leads to ipsilateral loss of discriminative touch, vibration, and proprioception below the level of the lesion.(fibers from the sacral cord are positioned most medially and those from the cervical cord are positioned most laterally.)

The loss of the spinothalamic tract leads to contralateral loss of pain and temperature sensation beginning one or two segments below the lesion.

Damage to the corticospinal tract produces weakness and spasticity in certain muscle groups on the same side of the body.

Physiology NEET 2020

Question 1:

A person develops headache and breathlessness on a trekking expedition following rapid ascent to over 3000 metres above the sea level. Which of the following is not used in the treatment of this condition?

a) IV digoxin

b) Descent to lower altitude

c) Administration of oxygen

d) Tablet acetazolamide

Question 2:

V/Q = infinity holds true in which of the following conditions?

a) Foreign body obstruction in the bronchus

b) Equal to dead space

c) When O_2 and CO_2 ratio is equal

d) The PO_2 of alveolar air is 159 mmHg and PCO_2 is 40 mmHg

Question 3:

Prolactin level is highest during :

a) 24 hours after ovulation

b) 24 hours after delivery

c) REM Sleep

d) After running for 1 hour

Question 4:

C wave of JVP represents:

a) Tricuspid bulge into right atrium during isovolumetric ventricular contraction

b) Contraction of right atrium

c) Passive filling of right atrium

d) Passive emptying of right atrium

Question 5:

The protein in the glomerular basement membrane responsible for charge-dependent filtration is?

a) Fibronectin

b) Heparan sulfate

c) Collagen Type IV

d) Entactin

Question 6:

Which of the following clotting factors would have decreased gamma carboxyglutamate residue in a patient on warfarin therapy?

a) Factor 2

b) Factor 11

c) Tissue factor

d) Factor 5

Question 7:

Which of the following conditions would have the highest levels of ACTH and cortisol?

a) Early morning

b) Normal individuals with dexamethasone administration

c) Addison's disease

d) Cushing's disease

Question 8:

Following complete ileal and partial jejunal resection, the patient is most likely to have

a) Folic acid deficiency

b) Vitamin B12 deficiency

c) Gastric ulcer

d) Constipation

Question 9:

Respiratory distress syndrome is due to defective

a) Type 1 pneumocytes

b) Type 2 pneumocytes

c) Alveolar capillary endothelial cells

d) Bronchial mucosal epithelial cells

Question 10:

Which of the following hormones use receptor tyrosine kinase signalling?

a) Insulin

b) TSH

c) TRH

d) MSH

Question 11:

What is the site of lesion in Wernicke's hemianopic pupil?

a) Optic nerve

b) Optic tract

c) Optic radiation

d) Optic chiasma

Question 12:

In which of the following reflexes, gastric emptying is delayed when acidic content enters the duodenum?

a) Gastrocolic

b) Enterocolic

c) Gastroileal

d) Enterogastric

Question 13:

A lady presents with vascular injury to the inferior frontal gyrus. Which functional area is likely to be affected?

a) Motor speech area

b) Wernicke's area

c) Auditory area

d) Visual area

Question 14:

A man falls into a deep sleep with one arm under his head. He finds he is not able to move this arm when he awakens. However, it tingles and pain sensation in the arm is still intact. What is the reason for the loss of motor function without loss of pain sensation is that in the nerves to his arm?

a) B fibers are more susceptible to pressure than A fibers

b) A fibers are more sensitive to pressure than C fibers

c) C fibers are more sensitive to pressure than A fibers

d) Motor nerves are more affected by sleep than sensory nerves

Question 15:

Which of the following has the highest oxygen concentration in fetal circulation?

a) Inferior vena cava

b) Right atrium

c) Superior vena cava

d) Ascending aorta

Question 16:

Respiratory distress syndrome is due to the deficiency of

a) Dipalmitoyl Phosphatidylcholine

b) Phosphatidylserine

c) Phosphatidylethanolamine

d) Phosphatidylinositol

Question 17:

What makes CPDA superior than ACD for blood storage:

a) The fall in 2, 3 DPG is less

b) Decreased release of O2

c) It has less P50

d) It is less acidic

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	a
5	b
6	a
7	d
8	b
9	b
10	a
11	b
12	d
13	a
14	b
15	a
16	a

17	a

Detailed Explanations

Solution to Question 1:

The symptoms of headache and breathlessness following rapid ascent indicate acute mountain/altitude sickness and IV digoxin is not used to treat this condition.

On ascent to high altitudes, the low partial pressure of oxygen leads to hypoxia and systemic arteriolar dilatation. Consequently, the capillary pressure rises with increased transudation of fluid into the brain tissue and the following symptoms are seen:

• Headache

• Irritation

• Insomnia

• Breathlessness

• Nausea and vomiting

Treatment involves administration of oxygen, descent to a lower altitude and administration of acetazolamide. Acetazolamide counteracts hypoxia by stimulating respiration and also reduces the formation of cerebrospinal fluid, thereby reducing intracranial tension.

Portable hyperbaric chambers are increasingly being used in the treatment of high-altitude illnesses when rapid descent is not possible. These chambers are available in armed forces hospitals in high altitude areas

Solution to Question 2:

VA /Q= infinity means that there is no exchange of O2 & CO2, which means the volume of the lung equals to dead space and the physiological effort of breathing is wasted.

VA represents the ventilation in alveoli and Q represents the blood flow through the alveolus. The ratio of VA and Q explains the respiratory exchange when there is an imbalance between alveolar ventilation and alveolar blood flow. The ventilation-perfusion ratio is considered normal if there is an equal amount of blood flow & ventilation through the alveolus.

VA /Q ratio is infinity when:

• There is adequate ventilation but no perfusion.

• No exchange of gases (O2 and CO2)through the respiratory membranes of corresponding alveoli.

• The alveolar air equilibrates the humidified inspired air.

• Partial pressures of the O2 and CO2 are 149 mmHg & 0 mm Hg respectively.

Solution to Question 3:

Among the given options, the prolactin level is highest 24 hours after delivery.

The maximum level of prolactin is seen during pregnancy and in the immediate postpartum period. During pregnancy, the secretion is increased reaching a peak at the time of delivery. After delivery, the plasma concentration falls to nonpregnant levels in a few weeks. Suckling produces surges of prolactin release.

Solution to Question 4:

'C' wave in JVP represents tricuspid bulge into right atrium during isovolumetric ventricular contraction.

Characteristics of JVP :

• a wave corresponds to atrial systole which causes regurgitation of some blood into the great veins.

• c wave occurs due to the increase in atrial pressure caused by bulging of the tricuspid valve into the right atrium during isovolumetric ventricular contraction.

• x descent occurs due to the fall in atrial pressure during ventricular systole due to pulling of valve leaflets.

• v wave is seen at the end of ventricular contraction and occurs due to the slow venous filling which causes an increase in atrial pressure.

• y descent occurs due to the passive emptying of the right atrium into the right ventricle.



Solution to Question 5:

The protein in the glomerular basement membrane responsible for charge-dependent filtration is heparan sulfate. It prevents filtration by repelling negatively charged proteins.

Collagen type IV, fibronectin, entactin are structural components of basement membrane.

Solution to Question 6:

Factor 2 would have decreased gamma carboxyglutamate residue in a patient on warfarin therapy.

Activation of coagulation factors II, VII, IX, X is done by carboxylation of their glutamate residues. This is done by the enzyme γ-glutamyl carboxylase with the reduced form of vitamin K as a co-factor.

This reduced form of vitamin K needs to be constantly recycled back from its oxidized form for the reaction to continue. This recycling is done by the enzyme vitamin K epoxide reductase. This enzyme is inhibited by warfarin.

Hence, warfarin indirectly inhibits the γ-carboxylation of vitamin K dependent clotting factors - Factors II, VII, IX, X by directly inhibiting vitamin K epoxide reductase. Thus leading to decreased gamma carboxyglutamate residue in these factors.



Solution to Question 7:

Among the given conditions, the levels of ACTH and cortisol will be highest in Cushing's disease.

Cushing's disease is caused by an ACTH secreting pituitary adenoma. The increased ACTH causes increased cortisol production.

Note: In normal physiologic state, ACTH levels are higher in the early morning but not as high as in Cushing's disease.



Solution to Question 8:

After complete ileal and partial jejunal resection, the patient is likely to have vitamin B12 deficiency.

Vitamin B12 is absorbed in the ileum. Vitamin B12 binds to intrinsic factor, a protein secreted by the parietal cells of the stomach, and the complex is absorbed across the ileal mucosa. Hence after ileal resection, there is impaired absorption of vitamin B12.

Solution to Question 9:

Respiratory distress syndrome is due to defective type 2 pneumocytes which produce lung surfactants. The surfactants reduce the surface tension in the alveoli. A defect in type 2 pneumocytes or surfactant deficiency can lead to respiratory distress syndrome.

The primary alveoli develop and the production of the surfactant begins by week 20-24. However, it does not reach the lungs until later. The surfactant appears in the amniotic fluid between 28 and 32 weeks of gestation. Mature levels of the pulmonary surfactant are present usually after 35 weeks of gestation.

Antenatal corticosteroids are given to increase production of surfactant.

Solution to Question 10:

Insulin uses the receptor tyrosine kinase signalling to exert its functions.

Other hormones acting in a similar fashion includes

• Growth hormone

• Growth factors like VEGF, PDGF, FGF, NGF

• Prolactin

• Leptin

Solution to Question 11:

The site of lesion in Wernicke's hemianopic pupil is optic tract.

Wernicke hemianopic pupil is the light reflex, where, illumination of the blind half of the retina shows no pupillary reaction while stimulating the normal side of the retina shows a pupillary reaction. It is seen in lesions involving the optic tract. It is also seen in cases such as TB, syphilis and tumours of the optic thalamus.

Homonymous hemianopia (HH) involves loss of vision on the same side of the visual field in both eyes. HH is characteristically seen in lesions involving the visual pathway posterior to the optic chiasma.

Solution to Question 12:

In the enterogastric reflex, gastric emptying is delayed when acidic content enters the duodenum.

When the food enters the duodenum, multiple reflexes are initiated from the duodenal wall which passes back to the stomach. These reflexes slow or stop the stomach emptying if the volume of the chyme in the duodenum becomes too much. These reflexes are called enterogastric reflexes.

This reflex can be triggered due to the following reasons:

• Presence of irritants and acids in duodenal chyme (most sensitive factor)

• Distension of duodenum due to the presence of an excessive volume of chyme

• Rapid entry of hypertonic or hypotonic fluids into the duodenum

• Presence of breakdown products of protein digestion and sometimes fat

Solution to Question 13:

Broca's motor speech area is likely to be affected in the vascular injury of the inferior frontal gyrus

A lesion in this region leads to motor/Broca's aphasia. Speech is non-fluent and interrupted by many word-finding pauses. The speech is impoverished in function words that contribute to the syntax but is enriched in meaningful nouns and verbs. However, comprehension of speech is intact and the patient knows what he wants to say.

For example, a patient with Broca's aphasia may say "walk dog" instead of "I want to take the dog for a walk".

Solution to Question 14:

The reason for the loss of motor function without loss of pain sensation is that in the nerves to his arm, A-fibers are more sensitive to pressure than C fibers.

Sleeping for a long period with the arms under the head puts pressure on a nerve. It leads to the loss of conduction in large-diameter motor, touch, and pressure fibers, while pain sensation remains relatively intact. A-fibers (includes the motor fibers) are affected.

The susceptibility of the nerve fibers to pressure is A>B>C.

This is responsible for Saturday night or Sunday morning paralysis.

Solution to Question 15:

Inferior vena cava has the highest oxygen concentration in fetal circulation.

The blood gets oxygenated in the placenta and this passes to the umbilical vein. From the umbilical vein, 50% of the blood enters the hepatic circulation and 50% bypasses the liver to enter the inferior vena cava. From here, the blood enters the heart via the right atrium. The right atrium also receives deoxygenated blood from the superior vena cava. This mixed blood is distributed to the body via aorta.

Therefore, inferior vena cava has the highest concentration of oxygen in the fetal circulation.



Solution to Question 16:

Respiratory distress syndrome is due to the deficiency of dipalmitoyl-phosphatidyl-choline(DPPC) /dipalmitoyl lecithin. It is the major pulmonary surfactant.

Clinical relevance- Test to assess fetal lung maturity is L/S ratio - lecithin/sphingomyelin ratio. A ratio of ≥ 2 is considered normal.

Solution to Question 17:

The fall in 2, 3 DPG is less when the blood is stored in CPDA compared with ACD.

2, 3 DPG (2,3-diphosphoglycerate) binds to deoxygenated haemoglobin. Therefore, an increase in 2,3 DPG causes increased release of O2 to the tissues. The equilibrium between Hb and O2 can be understood from the following equation:

$HbO_2 + 2,3\text{-DPG} \blacksquare Hb–2,3\text{-DPG} + O_2$

In stored blood, the 2,3 DPG levels usually fall. As a result, stored blood releases less oxygen to the tissues than the normal circulating blood, thus limiting its use in hypoxic patients (Eg: anaemia).

However, such a fall in 2,3 DPG is less when it is stored in a phosphate-containing solution like CPDA (Citrate, phosphate, dextrose, adenine), rather than ACD (Acid citrate dextrose). Therefore, the blood stored in CPDA, in comparison with blood stored in ACD has:

- Higher 2,3 DPG concentration
- More amount of deoxygenated blood bound to 2,3 DPG
- Decreased affinity of Hb to the oxygen.
- Increased release of oxygen to the tissues.
- Raised P50 (PO_2 at which Hb is half saturated)
- Shift in O_2-Hb dissociation curve to the right

Physiology INI-CET 2021

Question 1:

Which of the following is not involved in renal auto-regulation?

a) Tubuloglomerular feedback

b) Myogenic reflex

c) Renin-angiotensin system

d) Renal sympathetic system fibres

Question 2:

The formula for urea clearance is U*V/P, where U is

a) Urine volume

b) Serum urea in mg /100ml

c) Urine urea in g/24h

d) Urine urea in mg /100ml

Question 3:

The relaxation of the intestinal segment distal to the segment with the bolus of food during peristalsis is because of

a) VIP

b) Substance P

c) Dopamine from adrenergic fibres

d) Acetylcholine from myenteric fibres

Question 4:

A patient presented with severe watery diarrhea resembling that of rice gruel stool. Damage of which structure is likely to have resulted in this diarrhea?

a) Zona occludens

b) Gap junction

c) Adherent junction

d) Hemidesmosomes

Question 5:

Beta hCG is structurally similar to which biochemical moiety?

a) LH

b) FSH

c) ACTH

d) Oxytocin

Question 6:

What is the duration of short term memory?

a) 3-6 Hours

b) 12-18 hours

c) 30-300 seconds

d) 3-4 seconds

Question 7:

The stroke volume depends on end-diastolic volume. This is best represented as

a) Length-force relationship

b) Length- velocity relationship

c) Length-tension relationship

d) Length-viscosity relationship

Question 8:

Which of the following ions in the thick ascending limb of the loop of Henle is responsible for the resorption of divalent ions?

a) H^+

b) HCO_3^-

c) K+

d) Na+

Question 9:

Which of the following sensory receptors are responsible for tactile discrimination?

a) Merkel disc

b) Ruffini end organ

c) Pacinian corpuscle

d) Meissner's corpuscle

Question 10:

All of the following increases gastric acid secretion except:

a) Gastrin

b) Histamine

c) Acetylcholine

d) Serotonin

Question 11:

Pacinian corpuscle is stimulated by which of the following?

a) Pain

b) Temperature

c) Touch

d) Pressure

Question 12:

Which of the following jugular venous pressure (JVP) waves corresponds to the isovolumetric contraction phase of the cardiac cycle?

a) a wave

b) c wave

c) x descent

d) y descent

Question 13:

The extracellular concentration of chloride is 100mmol/L and the intracellular concentration is 10 mmol/L. What is the equilibrium potential of chloride?

a) + 10 mV

b) - 10 mV

c) - 60 mV

d) + 60 mV

Question 14:

After reaching an equilibrium state during exercising, it was found that the body was consuming oxygen at the rate of 1500 ml/min. Calculate the MET score if the baseline oxygen consumption at rest is 250 ml/min.

a) 5

b) 6

c) 1/5

d) 1/6

Question 15:

All of the following cause a right shift of the oxygen-hemoglobin dissociation curve, except ___.

a) Increased HbF

b) Raised 2,3 BPG

c) Acidosis

d) Increased CO_2

Question 16:

The latent period, contraction period, and relaxation period of a skeletal muscle are 10 ms, 40 ms, and 50 ms respectively. Calculate the tetanizing frequency.

a) 100 per second

b) 50 per second

c) 20 per second

d) 25 per second

Question 17:

An individual is in an environment of a temperature of 47°C. What is the mechanism of heat loss?

a) Radiation

b) Conduction

c) Convection

d) Sweating

Question 18:

Which among the following has the highest airway resistance?

a) Alveolar duct

b) Bronchi

c) Respiratory bronchioles

d) Small bronchioles

Question 19:

If the contractility of the heart is decreased, which of the following is seen

a) End diastolic volume and pressure curve shifts to left without change in slope

b) Slope changes in ESV pressure curve

c) End systolic volume and pressure curve shifts to left without change in slope

d) Slope changes in EDV and pressure curve

Question 20:

A lesion in which lobe will lead to impairment of hand-eye coordination?

a) Parietal lobe

b) Frontal lobe

c) Temporal lobe

d) Occipital lobe

Question 21:

All of the following ions have decreased concentration on the luminal side of proximal convoluted tubule except :

a) Amino acids

b) Glucose

c) Chloride

d) Bicarbonate

Question 22:

Sodium iodide symporter is present in all of the following except:

a) Thyroid gland

b) Placenta

c) Salivary gland

d) Pituitary gland

Answer Key

Question No.	Correct Option
1	d
2	d
3	a
4	a
5	a
6	c

7	c
8	c
9	d
10	d
11	d
12	b
13	c
14	b
15	a
16	d
17	d
18	b
19	b
20	a
21	c
22	d

Detailed Explanations

Solution to Question 1:

Renal sympathetic system fibres is not involved in the autoregulation of renal blood flow, because a denervated kidney also maintains the auto regulation of renal blood flow as that of a normal kidney.

Tubulo glomerular feedback, myogenic reflex, and renin-angiotensin system are all involved in the autoregulation of renal blood flow.

Tubulo glomerular feedback:

• When the renal artery pressure increases, the renal blood flow and glomerular filtration rate (GFR) increases

• This increased GFR results in increased delivery of solute and water to the early distal tubule, particularly the macula densa

• The macula dense, then secretes a vasoactive substance that constricts afferent arteriole via a paracrine mechanism

• This vasoconstriction then reduces renal blood flow and GFR back to normal

Myogenic reflex:

• The increase in renal arterial pressure stretches the walls of the afferent arterioles, which then contracts

• This contraction increases the afferent arteriolar resistance

• This increase in resistance then compensates for the increase in blood pressure, thereby keeping the renal blood flow constant.

Renin-angiotensin system:

• Angiotensin II is a potent vasoconstrictor of both afferent and efferent arterioles

• However efferent arterioles are more sensitive to angiotensin II than afferent arteriole

• Thus low levels of angiotensin II produce an increase in GFR by constricting efferent arterioles

• High levels of angiotensin II decreases GFR by constricting afferent and efferent arterioles, thereby decreasing the renal blood flow

Solution to Question 2:

'U' in the urea clearance stands for urine concentration of urea in mg /100ml.

The formula for renal clearance of a substance is,

Clearance (C) = U × V / P,

where

U = Urine concentration of the substance in mg / ml

V = Urine flow rate in ml/min

P = Plasma concentration of substance in mg /ml.

Renal clearance of a substance is the volume of plasma that is completely cleared of the substance by the kidneys per unit of time.

Solution to Question 3:

The relaxation of the intestinal segment distal to the segment of a bolus of food during peristalsis is due to Vasoactive Intestinal Peptide (VIP).

Peristaltic contractions in the small intestine are designed to propel the food towards the large intestine.

The food bolus in the small intestine is sensed by enterochromaffin cells of the intestinal mucosa, which release 5-hydroxytryptamine (5-HT)(serotonin). The 5-HT then binds to the receptors on intrinsic primary afferent neurons (IPANs), which when activated, initiate the peristaltic reflex in that segment of the small intestine. Behind the bolus, excitatory transmitters like acetylcholine, Substance P, neuropeptide Y are released in the circular muscle, while these pathways are inhibited in the longitudinal muscle; thus this segment of small intestine (behind the bolus) contracts and lengthens.

(Recall that contraction of circular muscle decreases the diameter of that small intestinal segment whereas contraction of the longitudinal muscle decreases the length of that segment. To prevent the conflict that would occur if circular and longitudinal are contracted at the same time, they are

reciprocally innervated)

In front of the bolus, inhibitory pathways like vasoactive intestinal peptide (VIP) and nitric oxide (NO) are activated in the circular muscle, while excitatory pathways are activated in the longitudinal muscle;thereby relaxing and shortening this segment of small intestine.

Solution to Question 4:

The given clinical picture of a patient having rice gruel stools is suggestive of the diagnosis of 'cholera'. Vibrio cholerae in addition to the cholera toxin (CT) produces some additional toxins such as,

• Zonula occludens toxin, which alters the intestinal permeability by acting on intestinal epithelial cell tight junctions (which are also called zonula occludens)

• Accessory cholera enterotoxin.

Zonula occludens:

• Also known as tight junctions

• Characteristically surround the apical margins of the cells in epithelia such as the intestinal mucosa, the walls of the renal tubules, and the choroid plexus.

• Also important to the endothelial barrier function and endothelium-dependent vasodilation.

• Made up of ridges- half from one cell and a half from another cell- which adhere so strongly that there is no space between the cells.

• Three families of transmembrane proteins are involved:

• Occludin

• Junctional adhesion molecule

• Claudins

• Paracellular pathway - these tight junctions permit the passage of ions, solutes, and intracellular signaling molecules in between adjacent cells.



Other options:

B) Gap junctions:

• The gap between the two cells in these types of junctions is about 4nm.

• These junctions are formed by units called connexons, which are present in the membrane of each cell.

• Each connexon is made up of six protein subunits called connexins.

• They permit substances to pass between the cells, without entering the ECF.

• The pore diameter is around 0.8 to 1.4 nm, which permits the passage of ions, sugars, amino acids, and other solutes which weigh up to 1000 Da.

• They also permit rapid propagation of electrical activity from cell to cell, as well as the exchange of various chemical messengers and intracellular signaling molecules.

C) Adherent junctions:

• Adherent junctions or Zonula adherens is a continuous structure on the basal side of the zonula occludens.

• It contains cadherins.

D) Hemidesmosomes:

• They look like half-desmosomes.

• They attach cells to the underlying basal lamina and are connected intracellularly to intermediate filaments.

• They contain integrins.

Solution to Question 5:

Beta hCG (human chorionic gonadotropin) has biological activity similar to that of luteinizing hormone (LH).

hCG is a glycoprotein produced by syncytiotrophoblasts. It is made up of α and β subunits. Though the α subunit of hCG is similar to the α subunits of LH, FSH, and TSH, the β subunit has structural similarity to LH only. Hence, LH is a better answer than FSH.

hCG is primarily luteinizing and luteotrophic. It appears to act on the same receptor as that of LH. The function of hCG is to prevent regression of the corpus luteum. It can be detected in blood as early as 6 days after conception and forms the basis of various tests used to identify pregnancy.

Its level peaks at gestational week 9 and then declines gradually but is present in lower concentration till the end of pregnancy. But hCG is not specific for pregnancy as it is secreted in a variety of gastrointestinal and genital tumors in both males and females.

Solution to Question 6:

Short-term memory lasts for seconds to minutes, hence of the given 4 options, 30-300 seconds is the most appropriate answer.

Memory is usually divided into two forms: Explicit or declarative memory which is associated with consciousness or awareness. Implicit or nondeclarative memory does not involve awareness and its retention does not usually involve processing in the hippocampus.

Explicit and many forms of implicit memory involve

• Short-term memory and

• Long-term memory.

Short-term memory:

It lasts seconds to minutes; the encoding process for short-term explicit memory involves the hippocampus. It lasts only till the person continues to think about the number or facts.

Working memory is a form of short-term memory that keeps information available, usually for very short periods of time, until the individual plans action based on it.

Long-term memory:

It stores memories that can be recollected after years or even for a lifetime. These memories are stored in various parts of the neocortex. The various parts of the memories such as, visual, auditory, olfactory, etc., are localised to the cortical regions concerned with these functions. All these parts of the memories are brought together by long-term changes in the strength of transmission at relevant synaptic junctions so that all the components are brought to consciousness when that particular memory is recalled.

Neural basis of memory:

The basis of memory is the alteration in the strength of the selected synaptic connections. Second messenger systems contribute to the changes in the neural circuits which are required for learning and memory. Alterations in the cellular membrane channels also contribute to it.

Solution to Question 7:

The relationship between end-diastolic volume and stroke volume is best represented by length-tension relationship.

This is explained by the 'Frank-Starling law of heart' which states that ' the volume of blood ejected by the ventricle depends on the volume present in the ventricle at the end of the diastole'. The length of the myocytes depend on the volume (and hence tension) present at the end of the diastole, which in turn, depends on the volume returned to the heart, or the venous return.

• The Frank-Starling law governs the normal ventricular function and ensures that the cardiac output equals venous return, within physiological limits.

• There is a curvilinear relationship between stroke volume or cardiac output and the ventricular end-diastolic volume.

• Agents that increase the contractility of the cardiac muscle have a positive inotropic effect and produce increased stroke volume and thereby increased cardiac output for a given end-diastolic volume. (Eg: Digoxin)

• Agents that decrease the contractility of the cardiac muscle have a negative inotropic effect and produce decreased stroke volume (and cardiac output) for a given end-diastolic volume.

Solution to Question 8:

Potassium (K+) in the thick ascending limb of loop of Henle is responsible for the resorption of divalent ions such as, calcium, magnesium, etc.

This resorption is driven by the lumen-positive potential difference across the cells of the thick ascending limb, created by the potassium leak into the lumen.

Mechanism of resorption in thick ascending limb:

• The luminal membrane contains a Na+-K+-2Cl- cotransporter.

• There is net resorption of Na+, K+ and Cl- in the thick ascending limb as follows:

• All three ions are transported into the cell on the cotransporter.

• Na+ is extruded from the cell by the Na+-K+ ATPase in the basolateral membrane.

• K+ and Cl- diffuse through channels in the basolateral membrane, down their respective electrochemical gradients.

• Most of the K+ leaves the cell across the basolateral membrane; the remaining portion of K+ however diffuses back into the lumen.

• This K+ leak into the lumen generates a positive electrochemical potential difference in the lumen.

• This lumen-positive potential difference drives the resorption of ions such as sodium(Na+), magnesium (Mg2+), calcium (Ca+),etc.,

Solution to Question 9:

Meissner's Corpuscles are responsible for discriminative touch.

Meissner's corpuscles are encapsulated receptors found in the dermis of the non-hairy skin, most commonly on the fingertips, lips, and other locations where tactile discrimination is very good. Their field of reception is very small and hence used for two-point discrimination. These are rapidly adapting receptors that encode

• Point discrimination

• Precise location

• Tapping

• Flutter

Solution to Question 10:

All of the above increase gastric acid secretion except serotonin.

Serotonin is localized in the enterochromaffin cells of the gastrointestinal (GI) mucosa and within neurons of the GI tract. Serotonin inhibits gastric acid secretion. It stimulates the production and release of gastric and colonic mucus. Serotonin acts as an enteric neurotransmitter and increases intestinal motility.

The main factors that stimulate gastric secretion are acetylcholine, gastrin, and histamine. Parasympathetic stimulation releases Ach which stimulates the secretion of

• pepsinogen by peptic cells

• hydrochloric acid by parietal cells

• mucus by mucous cells.

Both gastrin and histamine strongly stimulate acid secretion by parietal cells but have little effect on other cells.

The stomach mucosa has two important types of tubular glands:

• The oxyntic (gastric) glands secrete hydrochloric acid, pepsinogen, intrinsic factor, and mucus.

• The pyloric glands secrete mainly mucus for protection of the pyloric mucosa from the stomach acid. They also secrete the hormone gastrin.

Note- Serotonin released from the small intestine triggers vomiting by stimulation of 5HT3 receptors. Therefore, 5HT3 antagonists like ondansetron are used as antiemetics.

Solution to Question 11:

Pacinian corpuscle is stimulated by pressure.

Pacinian corpuscles are the largest cutaneous mechanoreceptors. They are present in the dermis of glabrous and hairy skin. They are composed of a nerve ending which is encapsulated by

concentric layers of connective tissue that give it an onion-like appearance. These receptors respond to fast vibration and deep pressure and adapt very rapidly. The sensory nerves from cutaneous mechanoreceptors are myelinated Aα and Aβ fibers.

Image shows histology of Pacinian corpuscle.



Other mechanoreceptors:

• Meissner's corpuscles are present mainly in the dermis of non-hairy skin particularly in the fingertips, lips, and other locations where tactile discrimination is well developed. They have small receptive fields and can be used for two point discrimination. Meissner corpuscles are rapidly adapting receptors that sense point discrimination, precise location, tapping, and flutter. These are encapsulated nerve endings of a large myelinated sensory nerve (Aβ type).

The image shows the histology of Meissner's corpuscle in the dermis.



• Merkel discs are seen in the epidermis of glabrous skin. They are slowly adapting receptors and have very small receptive fields. These receptors detect vertical indentations of the skin. They

are expanded dendritic endings.

• Ruffini corpuscles are capsulated, enlarged, dendritic endings. They are located in the dermis of glabrous and hairy skin, and in joint capsules. They are slowly adapting receptors. Ruffini corpuscles detect stretch and joint rotation.

Pain and temperature sensations arise from receptors located on unmyelinated sensory neurons present throughout the glabrous skin, hairy skin, and deep tissue. Impulses from nociceptors are transmitted via thinly myelinated Aδ fibers and unmyelinated C fibers.

Activation of Aδ fibers is responsible for first pain (also called fast pain). It is a rapid response and helps in the discrimination of pain or the ability to localize the site and intensity of the noxious stimulus. It is mediated by the release of glutamate. An example is pinprick.

Activation of C fibers is responsible for delayed second pain(also called slow pain). It is a dull, intense, diffuse, and unpleasant feeling associated with a noxious stimulus. . Second pain is mediated by the release of a combination of glutamate and substance P. Example is visceral pain.

Types of nociceptors :

• Mechanical nociceptors respond to strong pressure (eg, from a sharp object).

• Thermal nociceptors are activated by skin temperatures above 45°C or by severe cold (<20°C).

• Chemically sensitive nociceptors respond to chemicals such as bradykinin, histamine, high acidity, and environmental irritants.

• Polymodal nociceptors respond to combinations of these stimuli.

Solution to Question 12:

The c wave of jugular venous pressure wave corresponds to the isovolumetric contraction phase in the cardiac cycle.

During isovolumetric contraction, the ventricles start contracting with valves closed. This leads to bulging of the tricuspid valve. This bulging increases pressure in the right atrium which is transmitted to the right internal jugular vein. This appears as a c wave in jugular venous pressure (JVP) tracing.

JVP measured in the internal jugular vein (IJV) is a reflection of the right atrial pressure. IJV is preferred because it is not valved and is in direct line with superior vena cava and right atrium.



The waveforms seen in a normal JVP are:

• a wave corresponds to atrial contraction and occurs just after the P wave (on ECG) and just before the first heart sound.

• c wave represents the bulging of the tricuspid valve into the right atrium during ventricular systole.

• x descent corresponds to the atrial relaxation during atrial diastole.

• v wave represents atrial filling during ventricular systole and peaks at the second heart sound.

• y descent represents passive emptying of the right atrium into the right ventricle after tricuspid valve opening.

The Wiggers diagram given below correlates JVP with other aspects of cardiac cycle.



Solution to Question 13:

The equilibrium potential of chloride in the given question is -60 mV.

Solution to Question 14:

The MET score in the given scenario is 6.

One metabolic equivalent (MET) is defined as the amount of oxygen consumed while sitting at rest.

Normally, one MET= 3.5 ml O2 /kg body weight/min

In the above question, the baseline O2 consumption at rest is given as 250 ml/min. According to the definition of one MET, we can deduce that the given 250ml/min is one MET.

O2 consumption at reaching equilibrium during exercise= 1500ml/min

MET score= O2 consumption during exercise/ one MET

= 1500/250

= 6

The MET concept represents a simple, practical, and easily understandable procedure for expressing the energy cost of physical activities as a multiple of the resting metabolic rate. If the MET score is low, it means that the activity requires less oxygen compared to an activity that has a higher MET score. Eg. Gardening requires around 3-7 METS compared to that jogging that may need around 9-11 METS. We can classify activities as light, moderate or heavy, depending on MET scores. It can also be used to identify the functional capacity of a patient.

Solution to Question 15:

Increased HbF (fetal hemoglobin) causes a left shift of the oxygen-hemoglobin dissociation curve.

Oxygen-hemoglobin dissociation curve:

This is a plot of the partial pressure of O2 and the percentage saturation of Hb with O2. It is normally sigmoid in shape. If the oxygen-hemoglobin dissociation curve is shifted to the right, it means that the affinity of Hb for O2 has become less, which favors O2 delivery. This means a higher P50, which implies that a higher PO2 is required to maintain 50% saturation (R - Right shift Releases oxygen). If the curve is shifted to the left, it means that affinity has increased for O2 (which favors O2 binding) i.e. lower P50 value (L-Left shift Leashes oxygen).



Causes of the right shift of the oxygen-hemoglobin dissociation curve:

• Hypoxia and Hypercarbia - metabolically active tissues utilize O2 and tissues produce CO2

• ↓ pH - Secondary to increased CO2 production

• ↑ in temperature

• ↑ in 2,3-DPG - increased anaerobic glycolysis

• Growth hormone, thyroid hormone, androgen - by increasing 2,3 DPG

• Sickle cell anemia - reduced affinity for O2

• Exercise

Causes of the left shift of oxygen-hemoglobin dissociation curve:

• ↑ pH (alkalosis)

• ↓ PCO2

• ↓ 2,3-DPG

• ↓ Temperature

• HbF

• Carboxyhemoglobin

Bohr's effect- The decrease in the O2 affinity of hemoglobin when the pH of the blood falls

Solution to Question 16:

The tetanizing frequency of the given skeletal muscle is 25 per second.

In the question, it is given that the contraction period is 40 msec (40 msec is 0.04 seconds)

Tetanizing frequency = 1/contraction period

Therefore, tetanizing frequency = 1/40 millisecond = 1/0.04 per second = 1000/40 per second = 25 per second.

When there is a repeated stimulus to a muscle fiber before relaxation, it produces an additional contraction of the muscle. The second contraction adds to the first contraction and the total strength of contraction increases. This phenomenon is known as a summation of contractions.

With repeated stimulation, contraction occurs repeatedly before any relaxation has occurred. This causes the individual responses to fuse into one continuous contraction. Such a response is called tetanus (tetanic contraction). ■■■

Tetanizing frequency is the minimum frequency required for producing tetanus.

Tetanus is of two types.

• Complete tetanus- when no relaxation occurs between stimuli

• Incomplete tetanus- when periods of incomplete relaxation occur between the summated stimuli.



Solution to Question 17:

The mechanism of heat loss in the given case where the surrounding temperatures are higher than the body temperature is sweating.

Different methods of heat loss:

• Radiation is heat loss in the form of infrared rays.

• Conduction is heat loss that occurs by direct contact with an object.

• Convection is heat loss resulting from air movement.

When the temperature of the surroundings is more than that of the skin, the body gains heat by both radiation and conduction. Under these conditions, the heat can be lost from the body only by evaporation or sweating.

When the body temperature is greater than the temperature of the surroundings, heat can be lost by radiation and conduction.

Solution to Question 18:

Among the given options, bronchi (medium-sized airway) offer maximum resistance.

The best answer here is bronchi, as all the other options suggest smaller airways, while bronchi are the only medium-sized airway.

Airflow can be derived from Poiseuille-Hagen formula:

$Q = \pi \Delta P r^4 / 8\eta L$

Where

• Q is the flow rate

• π is pie symbol

• ΔP is the pressure difference

• r is radius

• η is viscosity

• L is length

Also,

Flow rate (Q) = pressure difference (ΔP)/ resistance (R).

Therefore, substituting this in Poiseuille- Hagen formula,

$\Delta P / R = \pi \Delta P r^4 / 8\eta l$

$1/R = \pi r^4 / 8\eta l$

$R = 8\eta l / \pi r^4$

Therefore it can be derived that airway resistance is inversely proportional to the radius by the power of four.

With Poiseuille- Hagen formula we might feel that the smallest airways are the major site of resistance. However, the contribution by the smallest airways is very little to the overall total resistance. This is because, the airway branches in each generation exist in parallel rather than in series. As the parallel branching increases, the resistance decreases. (Total cross-sectional diameter of small airways put together is high). Thus the overall resistance of the small airways is very small. Also, airflow velocity decreases substantially as the effective cross-sectional area increases (i.e., the flow becomes laminar).

Other factors affecting airway resistance :

• Lung volume, as lung volume increases the resistance decreases and vice versa. Lung volume increases the transairway pressure thus increasing the caliber of the airway.

• Airway mucus, edema, and contraction of bronchial smooth muscle increase the airway resistance. Its clinical application is seen in asthma, bronchitis, and emphysema

• The density and viscosity of inspired gas also contribute to airway resistance. A gas with more density imparts more resistance.

Solution to Question 19:

If the contractility of the heart is decreased, slope changes in the ESV pressure curve are seen.

To answer this question, we will first understand the volume-pressure diagram and the end-systolic pressure-volume relationship (ESPVR) during cardiac cycle.

Volume-pressure diagram during the cardiac cycle

The volume pressure diagram plots the changes in the pressure and volume in the left ventricle that occurs during the cardiac cycle. The x-axis shows the changes in ventricular volume, while the y-axis shows the changes in ventricular pressure.



It has the following phases.

Phase I- This phase is called the period of filling and extends between points A and B. Point A is at the end of systole when the ventricular volume is about 50 mL. This volume is called end-systolic volume. The pressure at this point is about 2 to 3 mm Hg. These volume and pressure go on increasing as the blood from the atrium flows to the ventricle during diastole. At the end of diastole, marked by point B, the ventricle will have a volume of around 120 mL. This volume is called end-diastolic volume. The pressure is slightly raised to 5-7mm Hg.

Phase II- It is the phase of isovolumetric contraction, and extends between points B and C. At the end of diastole (point B), a slight increase in the ventricular pressure closes the mitral valve, and contraction starts. The ventricle contracts while both mitral and aortic valves are closed. Therefore, the pressure within the ventricle slowly rises without any changes in the volume.

Phase III- It is the phase of ejection and extends between point C and point D. Point C is the place where the aortic valve opens. The pressure within the left ventricle rises slowly during isovolumetric contraction until it is sufficient to open the aortic valve. Once the aortic valve opens, blood within the left ventricle is ejected into the aorta. This decreases the volume of the left

ventricle, while increasing the pressure within the ventricle.

Phase IV- It is the period of isovolumetric relaxation. It extends between points D and A. At the end of the ejection (Point D), the aortic valve closes. However, the ventricular relaxation continues despite closed aortic and mitral valves. Therefore, the ventricular pressure continues to fall while the volume remains the same. The ventricle returns to its starting point, with about 50 ml of blood left in the ventricle and pressure of about 2 to 3 mm Hg

End-systolic pressure-volume relationship (ESPVR)

ESPVR is the maximum pressure that can be developed by the ventricle at the end-systolic volume. It is determined by the line intersecting the upper left corners of the loops. This line can be extended down to the x-axis. The slope of this line is called as ESPVR slope.



Now coming to the question, when the contractility of the heart is decreased, the force of ejection decreases leading to decreased stroke volume, and increased end-systolic volume. Therefore, the volume-pressure diagram shifts to the right. If the diagram shifts to the right, then ESPVR also changes, leading to a change in the slope of the ESPVR curve.



Other options:

Options A and D- End diastolic volume and pressure curve (EDPVR) mainly assesses the compliance of the ventricle, rather than contractility. The slope of this curve increases when the compliance decreases, and vice versa.

Option C- ESPVR shifts to right with change in slope.

Solution to Question 20:

A lesion in the parietal lobe will lead to impairment of hand-eye coordination.

The somatic sensory area and related parts of the posterior parietal lobe project to the premotor cortex. Some of these neurons are concerned with aiming and manipulating the hands at an object, whereas other neurons are concerned with hand-eye coordination. Neurons in this posterior parietal cortex contribute to the descending pathways involved in motor control.

There are two parietal lobes, the dominant side, and the non-dominant side.

• The functions of the dominant side of the parietal lobe are calculation, language, planned movement, appreciation of size, shape, weight, and texture. A lesion here causes dyscalculia, dysphasia, dyslexia, apraxia, agnosia, homonymous hemianopia.

• The function of the non-dominant side is spatial orientation and constructional skills. Its lesion causes spatial disorientation, constructional apraxia, neglect of the non-dominant side, dressing apraxia, and homonymous hemianopia.

Other options:

Option B: The frontal lobe is responsible for personality, emotional response, social behavior. Any lesion in the frontal lobe causes disinhibition, antisocial behavior, impaired memory, incontinence, anosmia.

Option C: The temporal lobe has two lobes and is divided as dominant and non-dominant.

• The dominant temporal lobe has important functions like auditory perception, speech, language, verbal memory, smell. Lesions lead to dysphasia, dyslexia, poor memory, complex hallucinations (smell, sound, vision), impaired face recognition (Prosopagnosia) and homonymous hemianopia.

• The nondominant temporal lobe auditory perception, music, tone sequences, non-verbal memory (faces, shapes, music). Lesions here leads to poor non-verbal memory, loss of musical skills, complex hallucinations, and homonymous hemianopia.

Option D: The occipital lobe is responsible for the analysis of vision. A lesion in occipital lobe mainly affects vision leading to homonymous hemianopia, visual agnosia, impaired face recognition (prosopagnosia), visual hallucinations (lights, lines and zigzags).

Solution to Question 21:

The concentration of chloride on the luminal side is more when compared to sodium, glucose, and amino acids.

In the first half of the proximal tubule, sodium is reabsorbed by co-transport along with glucose, amino acids, and other solutes. As sodium is reabsorbed in the proximal tubule, it preferentially carries glucose, amino acids, bicarbonate along with it. Thus, their concentrations decrease markedly along the length of the proximal tubule, and a solution having a higher concentration of chloride is left behind. The concentration of chloride here is around 105 mEq/L.

In the second half of the proximal tubule, sodium gets re-absorbed along with chloride since there is only a little glucose and amino acids yet to be re-absorbed. The second half of the proximal tubule has a concentration of chloride ≈140 mEq/L. This higher chloride concentration favors its own diffusion from the lumen into the renal interstitial fluid.

Although the amount of sodium in the tubular fluid decreases markedly along the proximal tubule, the concentration of sodium and the total osmolarity remain relatively constant. This is because the proximal tubule is highly permeable to water and water reabsorption increases to keep pace with the sodium reabsorption.



Solution to Question 22:

Sodium iodide symporter is present in all of the options given above except the pituitary gland.

Sodium iodide symporter is expressed mainly in the thyroid gland and is responsible for iodine uptake in the follicular cells of the thyroid gland.

Sodium iodide symporter (NIS)

The NIS is mainly present in the basolateral membranes of the thyrocytes. The NIS transports two Na+ ions and one I- ion into the cell against the concentration gradient. The energy for this transport is derived from the Na+-K+ ATPase pump. Na+-K+ ATPase pumps sodium outside the cell and causes low intracellular sodium concentration. This causes sodium to enter the cell by facilitated diffusion. Along with sodium, iodide ions also enter the cell. Since the energy to

transport sodium and iodide ions in NIS is derived from Na+-K+ ATPase pump, this is called secondary active transport.

The NIS is also expressed in:

• Salivary glands

• Gastric mucosa

• Placenta

• Ciliary body of the eye

• Choroid plexus

• Mammary glands

• Certain cancers are derived from these tissues.

The significance of these extrathyroidal NIS is not known.

Physiology NEET 2021

Question 1:

A 30-year-old man weighing 70 kg had a sodium level of 120 mEq/L. Calculate the sodium deficit.

a) 280 mEq

b) 480 mEq

c) 840 mEq

d) 1400 mEq

Question 2:

Beta 2 receptor action leading to hyperglycemia/ hypoglycemia is mediated by which of the following enzymes?

a) Adenylyl cyclase

b) Guanylyl cyclase

c) Phospholipase C

d) Phospholipase A2

Question 3:

A body fluid sample is being studied and has Na: 10 mEq/L, K: 140 mEq/L, CI: 4 mEq/L. Identify the compartment from which the sample has been obtained.

a) Interstitial

b) Intracellular fluid

c) Extracellular fluid

d) Plasma

Question 4:

A middle-aged female patient presented with slowly progressive motor weakness, nuchal rigidity, and difficulty in swallowing and carrying out day-to-day activities. A provisional

diagnosis of multiple sclerosis is suspected. Which of the following cells are affected in multiple sclerosis?

a) Microglia

b) Oligodendrocytes

c) Astrocytes

d) Ependymal cells

Question 5:

A child with loose stools for one day presented to your OPD. On examination, signs of dehydration were present and the child was given ORS. It facilitates glucose absorption through which of the following receptors?

a) SGLT1

b) SGLT2

c) GLUT4

d) Na- Ca transporter

Answer Key

Question No.	Correct Option
1	c
2	a
3	b
4	b
5	a

Detailed Explanations

Solution to Question 1:

The sodium deficit in the given scenario is 840 mEq.

Sodium deficit is calculated as 0.6 x body weight x (target plasma sodium concentration - starting plasma sodium concentration).

Substituting the given values in the formula :

Sodium deficit = 0.6 x 70 x (140 - 120) = 840 mEq.

Solution to Question 2:

Beta 2 receptor action leading to hypoglycemia/ hyperglycemia is mediated by adenylyl cyclase. Catecholamines act through beta 2 receptors and they use adenylyl cyclase - cAMP second messenger system to stimulate the target tissues.

Other hormones that use adenylyl cyclase - cAMP second messenger system are :

- Adrenocorticotropic hormone
- Angiotensin II
- Corticotropin-releasing hormone
- Calcitonin
- Follicle-stimulating hormone
- Glucagon
- Growth hormone-releasing hormone
- Human chorionic gonadotropin
- Luteinizing hormone
- Parathyroid hormone
- Secretin
- Somatostatin
- Thyroid-stimulating hormone
- Vasopressin

Hormones exert their actions by stimulating cAMP production within the cells. When the hormone binds to the membrane receptor, the receptor is coupled to the G protein. This G protein can either stimulate (Gs protein) or inhibit (Gi protein) adenylyl cyclase that controls the conversion of cytoplasmic adenosine triphosphate to adenosine monophosphate (cAMP). The protein kinases dependent on cAMP are thus activated. This leads to a series of biochemical reactions enabling the hormone to exert its actions.

Guanylyl cyclase is the enzyme that catalyzes the formation of cGMP.

Solution to Question 3:

The sample obtained is from the intracellular fluid (ICF) compartment. The ICF contains large quantities of potassium (K+) ions - 140 mEq/L . The ICF also has lesser amounts of sodium (Na+) and Chloride (Cl-) ions.

The extracellular fluid (ECF) compartment includes interstitial fluids and plasma. In contrast to the ICF, the ECF has an abundance of sodium, chloride, calcium and bicarbonate ions, and lesser amounts of potassium, phosphate and magnesium ions.

The image below depicts the proportion of the important anions and cations present in the intracellular and extracellular fluids.



Solution to Question 4:

Oligodendrocytes are affected in multiple sclerosis.

In multiple sclerosis, there is patchy destruction of myelin in the central nervous system and the axons are demyelinated. This causes loss of conduction or delayed conduction in the demyelinated axons. Oligodendrocytes and Schwann cells are involved in the formation of myelin. Hence, they are affected by multiple sclerosis.

Other options

Option A: Microglia are a part of the immune system. Microglia are scavenger cells that aid in the removal of cellular debris formed due to various diseases. They originate from macrophages outside of the nervous system and are unrelated to the neural cell types.

Option C: There are three types of macroglia - astrocytes, oligodendrocytes, and Schwann cells. Astrocytes are found throughout the brain and they induce capillaries to form the blood-brain barrier. The astrocytes may be fibrous or protoplasmic.

Option D: Ependymal cells line the ventricular system. They are in contact with the cerebrospinal fluid and play a role in fluid homeostasis.

Solution to Question 5:

ORS facilitates glucose absorption through SGLT 1 (Sodium-dependent glucose co-transporter - 1). In a child with diarrhea, when oral rehydration solution is given, there is coupled absorption of sodium and glucose through SGLT -1.

Glucose transport may be active or passive.

Active transport of glucose takes place through Sodium dependent glucose transporter (SGLT 1 and 2). SGLT 1 helps in glucose absorption in the small intestine and renal tubules. SGLT 2 primarily facilitates glucose absorption in the renal tubules.

Passive absorption of glucose occurs through facilitated diffusion. This is carried out by Glucose transporters (GLUT) 1 -7.

Physiology INI-CET 2022

Question 1:

Which physiological adaptation does not happen at high altitudes?

a) Pulmonary vasoconstriction

b) Polycythemia

c) Respiratory acidosis

d) Hypoxia

Question 2:

Countercurrent mechanism is present in which of the following parts of the body?

a) 1 and 2 are correct

b) 1,2,3 are correct

c) 1,2,3,4,5 are correct

d) 2,3,4,5 are correct

Question 3:

Match the following molecules with their carriers in plasma:

a) 1-b, 2-c, 3-d, 4-a

b) 1-a, 2-b, 3-c 4-d

c) 1-a, 2-d, 3-b 4-c

d) 1-b, 2-c, 3-a, 4-d

Question 4:

Myosin ceases to be in contact with actin filaments as a result of:

a) Dissociation of ADP and Pi

b) Cessation of calcium release from sarcoplasm

c) Conformational change in Troponin T

d) Attachment of ATP to myosin head

Question 5:

Transport of iron into enterocytes is done by which of the following proteins:

a) Transferrin

b) Ferroportin

c) Hepcidin

d) Divalent metal transporter - 1

Question 6:

Vessel A has a length of L and diameter of D and vessel B has 2L and 2D. Considering the flow is laminar through both the cylinders, the flow through B is how many times as much as the flow through A?



a) 4 times

b) 8 times

c) 16 times

d) 32 times

Question 7:

If the pH is reduced from 7.4 to 7.3 with the PO2 being constant at 100 mmHg, which line would PCO2 follow?



a) 1

b) 2

c) 3

d) 4

Question 8:

Substances and the type of transport across a membrane are given below, choose the correct combination:

a) 1 and 2 are correct

b) 1, 2, and 4 are correct

c) 1, 2, 3, 4, and 5 are correct

d) 1, 2, and 5 are correct

Question 9:

A and B in the graph below represent which of the following ions, in terms of permeability for action potential generation?



a) A is Na+, B is K+

b) A is Na+, B is Ca2+

c) A is K+, B is Ca2+

d) A is Ca2+, B is K+

Question 10:

Transcutaneous electrical nerve stimulation (TENS) is used to relieve pain. What is the mechanism behind it?

a) Gate control theory

b) Conditioned pain modulation

c) Descending inhibition

d) Central inhibition

Question 11:

The parents notice that their child is tired and that he has shortness of breath on climbing stairs and while lying down. What is the first investigation that you would do?

a) Echocardiogram

b) Chest x-ray

c) Spirometry

d) MRI to rule out enlarged heart

Question 12:

Match the following molecules with their carriers in plasma:

a) 1-b, 2-c, 3-d, 4-a

b) 1-a, 2-b, 3-c 4-d

c) 1-a, 2-d, 3-b 4-c

d) 1-b, 2-c, 3-a, 4-d

Question 13:

Arrange the following steps of ventricular action potential in the correct order.

a) B, A, C, D

b) D, C, B, A

c) C, D, A, B

d) A, B, C, D

Question 14:

What does the zero pressure indicate in the pressure-volume curve?

a) Residual volume

b) Functional residual capacity

c) Tidal volume

d) Inspiratory reserve volume

Question 15:

How is cerebral perfusion pressure calculated?

a) Systolic blood pressure + Intracranial pressure

b) Systolic blood pressure - intracranial pressure

c) Mean arterial pressure/intracranial pressure

d) Mean arterial pressure - intracranial pressure

Question 16:

Fluid leaving the proximal convoluted tubule in the absence of antidiuretic hormone is:

a) Isotonic

b) Hypotonic

c) Hypertonic

d) Depends on fluid intake

Question 17:

2,3-BPG causes the unloading of oxygen to tissues by _____.

a) Increasing ATP production

b) Binding to beta subunit of Hb

c) Inhibiting taut structure

d) Decreasing ATP production

Question 18:

The graph below shows oxygen dissociation curves. What does the curve marked as 'A' indicate?



a) Myoglobin

b) Carboxyhemoglobinemia

c) Methemoglobin

d) Fetal hemoglobin

Question 19:

Which among the following organs has the least arteriovenous oxygen difference?

a) Brain

b) Kidney

c) Skin

d) Liver

Question 20:

The image shows changes in membrane potential during an action potential. The region marked between 4 to 5 is due to _____.



a) Influx of Na+

b) Efflux of Na+

c) Influx of K+

d) Efflux of K+

Question 21:

What is the value of normal tissue oncotic pressure?

a) 12 mmHg

b) 25 mmHg

c) 32 mmHg

d) 54 mmHg

Question 22:

Where is the signaling sequence in a type 2 membrane protein located?

a) Inside the membrane at the N-terminal end

b) Inside the membrane at the C-terminal end

c) Outside the membrane at the N-terminal end

d) Outside the membrane at the C-terminal end

Question 23:

Which of the following primary second messengers is involved in the contraction of ciliary muscles while focusing on near objects?

a) cAMP

b) cGMP

c) Na+ inflow through a channel

d) IP3

Question 24:

Which of the following tissues are dependent on insulin for glucose uptake?

a) Brain

b) Adipose tissue

c) Pancreas

d) Muscle

Question 25:

The second phase of saliva secretion involves:

a) Active Na+ reabsorption, passive Cl- reabsorption, passive K+ secretion, active secretion of HCO_3^-

b) Passive Na+ reabsorption, active Cl- reabsorption, passive K+ secretion, active secretion of HCO_3^-

c) Passive Na+ secretion, active Cl- reabsorption, active K+ secretion, passive secretion of HCO_3^-

d) Active Na+ reabsorption, passive Cl- reabsorption, active K+ secretion, passive secretion of HCO_3^-

Question 26:

Macula densa is made of specialized cells located at:

a) Junction of thick descending limb and DCT

b) Junction of thick descending limb and PCT

c) Junction of thick ascending limb and DCT

d) Junction of DCT and collecting duct

Question 27:

Which of the following can be used to calculate the cardiac output?

a) A, B, C, D

b) A, D

c) B, C, D

d) A, C, D

Question 28:

Match the given cellular components with their respective marker enzymes.

a) 1-D, 2-A, 3-C, 4-B

b) 1-D,2-A,3-B,4-C

c) 1-C, 2-B, 3-A, 4-D

d) 1-B, 2-C, 3-D, 4-A

Question 29:

Consider the following statements about sympathetic, parasympathetic, and motor nerves. Which of the following statements are true?

a) 1,3

b) 1,4

c) 2,3

d) 2,4

Question 30:

What is the normal insensible water loss?

a) 100 mL/hour

b) 150 mL/hour

c) 50 mL/hour

d) 200 mL/hour

Question 31:

A woman has vomiting whenever she eats spicy food. Arrange the sequence of events during vomiting.

a) 6, 4, 2, 5, 1, 3

b) 6, 4, 2, 3, 1, 5

c) 4, 6, 2, 1, 3, 5

d) 4, 6, 2, 5, 3, 1

Answer Key

Question No.	Correct Option
1	c
2	d
3	a
4	d

5	d
6	b
7	a
8	c
9	a
10	a
11	b
12	a
13	c
14	b
15	d
16	a
17	b
18	a
19	b
20	d
21	a
22	a
23	d
24	b
25	d
26	c
27	d
28	b
29	b
30	c
31	b

Detailed Explanations

Solution to Question 1:

The physiological adaptation that does not happen at high altitudes is respiratory acidosis. Respiratory alkalosis is the physiological adaptation seen at high altitudes due to hyperventilation. It is a method of acclimatization. Compensatory mechanisms that are carried out to increase altitude tolerance are called acclimatization.

Following are the physiological changes that occur during acclimatization:

• Hyperventilation leads to respiratory alkalosis and shift of the oxygen–hemoglobin dissociation curve to the left (and shifts to right after aclimitization).

• Increased erythropoietin secretion, leading to an increase in circulating red blood cells

• Increased number of mitochondria

• Increased myoglobin to facilitate the movement of O2 into the tissues.

• Increased tissue cytochrome oxidase

• Increased sympathetic tone

• Renal suppression of carbonic anhydrase and excretion of dilute alkaline urine brings the pH of the blood to normal.

• Decreased maximal exercise capacity (by ~1% for every 100 meters gained above 1500 meters)

Other options:

Option A- In response to the high altitude hypoxia, pulmonary vasoconstriction occurs due to which blood flow is distributed where it is most effective, leading to distribution of blood flow to the pulmonary areas in proportion to their alveolar O2 pressures.

Option B- Polycythemia - Erythropoietin secretion increases on the ascent to high altitude which results in polycythemia

Option D- Hypoxia - Total barometric pressure decreases as altitude increases, resulting in decreased Po2 (partial pressure of oxygen) leading to hypoxic hypoxia.

Solution to Question 2:

The countercurrent mechanism is present in the testis, kidney, gut, and lungs. When the inflow runs parallel to, counter to, and in close proximity to the outflow for some distance it is called as the counter current mechanism.

Spermatogenesis requires a temperature considerably lower than that of the interior of the body. The testes are kept cool by heat exchange in a countercurrent manner between the spermatic arteries and veins.

The concentration of the urine by the kidney requires an osmolality gradient. This gradient is produced by the loop of Henle as counter-current multipliers and maintained by vasa recta as counter-current exchangers.

The functioning of the loop of Henle depends upon:

• The high permeability of the thin descending limb to water (via aquaporin-1)

• Active transport of Na+ and Cl- from the thick ascending limb, thereby diluting the tubular fluid.

• The inflow of tubular fluid from proximal tubule and outflow to the distal tubule.



The countercurrent mechanism is used in the villi to decrease the oxygen availability for local metabolic reactions of the gut. Hence this will increase the blood flow in the intestine during gastrointestinal activity.

Countercurrent exchange of oxygen is seen in the lungs as well.

Solution to Question 3:

Plasma contains various components like ions, inorganic and organic molecules which help in the transport of substances across various parts of the body. Albumin, globulin, and fibrinogen fractions are important plasma proteins. Some of the plasma proteins act as antibodies and are also concerned with blood clotting. Where some others are non-specific carriers of hormones, drugs, and other solutes. Majority of plasma proteins are synthesized in the liver. The above table shows some of the proteins synthesized by the liver and the molecules they carry.

Thyroxine is carried by thyroid-binding pre-albumin which is also known as transthyretin.

Albumin is a carrier protein as well as an osmotic regulator which carries hormones, amino acids, vitamins, and steroids too.

Hemopexin binds with heme, particularly for heme recycling.

Haptoglobin binds to free hemoglobin and prevents its excretion in urine.Hence, haptoglobin levels decrease in the presence of free hemoglobin as seen in hemolytic anemia.

Molecules	Carriers
Thyroxine	Prealbumin
Free fatty acid	Albumin

Molecules	Carriers
Hemoglobin	Haptoglobin
Heme	Hemopexin

Solution to Question 4:

Myosin ceases to be in contact with actin filaments as a result of the attachment of ATP to the myosin head.

Muscle contraction occurs by sliding of thin filament over thick filament. The sequence of events are as follows:

• Troponin I is bound to actin and tropomyosin and covers the sites where actin interacts with myosin heads in a resting muscle. The myosin head contains tightly bound adenosine phosphate (ADP) and inorganic phosphate (Pi).

• Following an action potential, cytosolic Ca^{2+} is increased due to the Ca^{2+} released from the sarcoplasmic reticulum. Simultaneously Ca^{2+} influx happens through voltage-gated Ca^{2+} channels too.

• The free Ca^{2+} binds with troponin C. This binding will weaken the troponin I interaction with actin and exposes the actin-binding site for myosin.

• This results in the formation of myosin-actin cross-bridges.

• The cross-bridge between the head of the myosin and the active site of the actin causes a conformational change in the head and the power stroke occurs, moving the thin filament relative to the thick filament.

• Once the head of the cross-bridge tilts, the ADP and phosphate ion that were previously attached to the head are released and a new ATP molecule can bind in that place.

• The binding of the new ATP molecule to the free site on the myosin causes detachment of the head of the myosin from the actin.

• ATP is hydrolyzed and Pi released, causing a re-cocking of the myosin head and completing the cycle.



Solution to Question 5:

Transport of iron into enterocytes is done by divalent metal transporter-1 protein.

Iron absorption occurs in the duodenum. Iron in our gut exists in 2 forms, they are nonheme and heme iron.

Luminal nonheme iron is mostly in the Fe3+ (ferric) state and is reduced to Fe2+ (ferrous) iron by ferrireductases. Fe2+ iron is then transported across the apical membrane by divalent metal transporter 1 (DMT1).

Given below is an image of iron absorption that shows divalent metal transporter-1 protein in intestinal mucosal cells.



Heme iron is moved across the apical membrane into the cytoplasm through heme transporters.

In the duodenal cells, Fe^{2+} iron can follow one of two pathways: transport to the blood or storage as mucosal iron. The nonstorage form of Fe^{2+} iron is transported from the cytoplasm into circulation by the protein ferroportin (Option B).

Once in circulation, Fe^{2+} is again oxidized back to Fe^{3+} iron by hephaestin and ceruloplasmin. The newly absorbed Fe^{3+} iron binds rapidly to transferrin (Option A), which delivers iron to red cell progenitors in the marrow.

Hepcidin (Option C), is a protein secreted from the liver that plays a major role in the regulation of iron absorption. When the body's iron stores are adequate, it downregulates the expression of ferroportin and thereby decreases iron absorption.

Solution to Question 6:

Flow through B will be 8 times more than the flow through A.

According to Poiseuille Hagen formula,

$F = \Delta P \times \pi/8 \times 1/n \times r4/L$

F = flow

ΔP = pressure difference between two ends of the tube

η = viscosity

r = radius of the tube

L = length of the tube

Hence flow directly varies to the 4th power of radius and inversely varies to length.

Flow through B = $(2r)4/2L$ = 8 times

Solution to Question 7:

If the pH is reduced from 7.4 to 7.3 with the alveolar PO_2 being constant at 100 mmHg, PCO_2 would follow line 1.

This concept can be understood in three steps.

Step 1- Effect of CO_2 on ventilation, when the alveolar PO_2 is kept constant.

When the alveolar PO_2 (PaO_2) is kept constant, an increased amount of inspired CO_2 leads to increase in the ventilation. The relation is a straight line.

The below graph shows the effect of CO_2 on ventilation when alveolar PaO_2 is kept constant at 100mm Hg.



Step 2. The shift of the above graph when PO2 changes.

When PO2 changes, the relation between the amount of CO2 inhaled and ventilation still remains in a straight line. However, the slopes of the lines change. The slope increases when alveolar PO2 decreases and vice versa. In other words, the lines move towards the left when PO2 decreases. However, the point of intersection of all these lines remains the same.

The below graph shows the relation between inspired CO2 and ventilation when alveolar PO2 is maintained at 40mm Hg, 55mm Hg, and 100 mm Hg. As the alveolar PO2 has decreased, the lines have shifted to the left. But they are all intersecting at the same point on the x-axis.



Step 3. The shift of the above graph when pH changes.

When pH decreases, as in acidosis, the CO2 response curve is similar to the above. The only difference is they are shifted to the left and have different points on the intersection on x- axis.

The below graphs show CO2 response curves when pH is 7.4, 7.3, and PO2 is 40mm Hg, 55mm Hg, and 100mm Hg. The lines have shifted to the left and have a different point of intersection.



In the question image, only line 1 has been shifted to the left and has a different point of intersection on x-axis compared to other lines. Therefore, line 1 is the answer.

Solution to Question 8:

All the given combinations 1, 2, 3, 4, and 5 are correct.

1) Oxygen is a non-polar molecule and hence can diffuse across the lipid membrane of the cells.

2) Facilitated diffusion - Transporting the molecule from one side of the cell membrane to the other is done with the help of transport proteins called carriers. Molecules move from areas of high concentration to areas of low concentration. No energy input is required when carrier proteins move substances in the direction of their gradients. Glucose transport by the glucose transporter is an example, which moves glucose down its concentration gradient from the ECF to the cytoplasm of the cell.



3) Voltage-gated channels are simple ion channels acting as transport proteins, gated by membrane potential alterations. Sodium transported via voltage-gated channels is an example.

4) Transporting more than one substance together using a single transport protein against the gradient is known as secondary active transport. An example is SGLT 1, the cotransporter of sodium and glucose from the intestinal lumen into mucosal cells.



5)Active transport - Transporting of substances against the gradients, and requires energy. Ca2+ ATPase that pumps out calcium out of the cells is an example.

Solution to Question 9:

In the given graph, showing action potential generation in terms of ion permeability, wave A represents Na+ and B represents K+.

The sequence of events during the action potential occurs in 7 steps as described below.

1. Resting-state - Both channels are closed

2. Na+ channels open in response to depolarizing stimulus. Sodium enters the cell. The membrane reaches threshold potential.

3. The sodium influx causes the opening of more Na+ channels through a positive feedback mechanism. Rapid depolarization occurs.

4. Sodium conductance decreases.

5. Na+ channels close and K+ channels open. Repolarization occurs.

6. Sodium channels are reset, and are ready to respond to another depolarizing stimulus (relative refractory period). During this phase, the potassium channels remain open and slowly return to closed state causing after-hyperpolarization.

7. Both channels are closed. The membrane returns to RMP.



Solution to Question 10:

Gate theory of pain is the mechanism behind transcutaneous electrical nerve stimulation (TENS). This method uses electrodes to activate Aα and Aβ fibres near the injury to relieve pain.

At the site of sensory afferent termination, transmission in nociceptive pathways can be interrupted. This is done by the actions within the dorsal horn of the spinal cord. Pain subsiding when the site of injury is rubbed or shook is an example. Afferents of innocuous cutaneous mechanoreceptors emit collaterals that terminate in the dorsal horn. The responsiveness of dorsal horn neurons to their input from nociceptive afferent terminals is reduced by these afferents This is called the gate-control mechanism of pain modulation.

The given image depicts the gate control theory of pain:



Solution to Question 11:

The first investigation to be done for a child who feels shortness of breath on climbing stairs and in the supine position is a chest x-ray.

The child having shortness of breath on climbing stairs is known as dyspnea on exertion. The most common cause is congestive heart failure. The child feeling shortness of breath in the supine position is orthopnea. It is relieved on sitting or standing.

Respiratory causes include asthma, chronic obstructive pulmonary disorder (COPD), pneumonia, and pulmonary embolism. Cardiovascular causes include congestive heart failure(CHF), pulmonary edema, valvular heart defect, and pulmonary hypertension.

Evaluation begins with assessing the airway, breathing, and circulation of the patient. Oxygen saturation can appear normal at rest, and hence oxygen saturation during physical activity should

be obtained. A chest x-ray is the first diagnostic test which can show pulmonary pathology or pumonary edema and other signs of CHF that could help in identifying the cause and further evaluation.

Other options:

Option A: An echocardiogram can be done if the chest X-ray is abnormal. It is done to evaluate cardiac function, valvular function, and pericardial space.

Option C: Spirometry is done when the chest x-ray is normal. It is done to determine lung function. Abnormal spirometry can indicate either an obstructive pathology such as asthma, or COPD or a restrictive disease such as interstitial fibrosis.

Option D: Cardiac MRI (Magnetic resonance imaging) is an emerging modality done to look for cardiomegaly. It is for the accurate evaluation of left and right ventricular mass, size, and function. It is also used for localizing ischemic and non-ischemic causes such as myocarditis.

Solution to Question 12:

The correct pairs are as follows:

1. Thyroxine - B. Pre-albumin

2. Vitamin B12 - C. Transcobalamin

3. Hemoglobin - D. Haptoglobin

4. Heme - A. Hemopexin

Thyroxine is carried by thyroid-binding prealbumin, which is also known as transthyretin.

Vitamin B12 binds to transcobalamin II in the ileal cells and is secreted into the plasma. Transcobalamin II delivers vitamin B12 to the liver and other cells of the body, including rapidly proliferating cells in the bone marrow and the gastrointestinal tract.

Hemopexin binds to porphyrins, particularly heme, for heme recycling.

Haptoglobin binds to free hemoglobin and prevents its excretion in urine.

Solution to Question 13:

The correct sequence for ventricular action potential is as follows:

C. Opening of voltage-gated sodium channels

D. Transient outward flow of potassium

A. Opening of voltage-gated calcium channels

B. Repolarizing K+ currents

The events of action potential occur in the following sequence:

• Phase 0: Voltage-gated Na+ channels open in response to depolarizing stimulus (rapid depolarization phase)

• Phase 1: Inactivation of voltage-gated Na+ channels and transient outward flow of K+ causes rapid repolarization (rapid repolarization phase)

• Phase 2: Opening of slow voltage-gated Ca²■ channels causes the influx of calcium into the cells and results in the plateau phase (plateau phase)

• Phase 3: Opening of various K+ channels results in the efflux of K+ from the cells and results in slow repolarization (slow repolarization phase).

• Phase 4: Return to resting membrane potential



Solution to Question 14:

The zero pressure in the pressure-volume curve indicates the functional residual capacity.

The pressure-volume relationship is basic to the understanding of the fundamental relationship between the compliance of the lungs and the chest wall. It is measured by a spirometer.

The functional residual capacity is the volume of the air that remains in the lungs after normal expiration. The normal value is about 2.5 liters.

Other options:

Option A: Residual volume is the air left in the lungs after a maximal expiratory effort.

Option C: Tidal volume is the amount of air that moves in during each inspiration or the amount of air that moves out with each expiration, during quiet breathing. The normal tidal volume is 500–750 mL.

Option D: Inspiratory reserve volume is the air inspired with a maximal inspiratory effort in addition to the tidal volume.

Solution to Question 15:

Cerebral perfusion pressure (CPP) is calculated as mean arterial pressure (MAP) - intracranial pressure (ICP).

However, ICP values are not readily available. In this case, CPP may be calculated as MAP - central venous pressure (CVP), and if CVP > ICP.

The brain depends on continuous perfusion for oxygen and glucose delivery, thus its survival. The blood flow to the brain is normally about 55 mL per 100 g of brain tissue. If the cerebral flow reduces to less than 20 mL per minute, the brain undergoes ischemia.

Blood flow depends on cerebral perfusion pressure (CPP). Normally the CPP is 75-105 mmHg, MAP is 90-110 mmHg and ICP is 5-15 mmHg.

According to Monro Kellie doctrine, the cranium is a rigid box containing a nearly incompressible brain. In cases of head trauma, there is an expansion of cranial content, in the form of hematomas and brain swelling. This increase in cranial pressure is accommodated initially by the exclusion of fluid components, venous blood, and cerebrospinal fluid. Further increase in the cranial pressure leads to an uncontrolled rise in ICP resulting in a compromise in cerebral perfusion and eventually cerebral herniation.

Acute rise in ICP causes Cushing reflex, a triad of widened pulse pressure (increasing systolic, decreasing diastolic) bradycardia, and irregular respirations.

Thus, in cases of head injury with intracranial bleeding, cranial decompression is performed to avoid further complications due to the raised ICP.

Solution to Question 16:

Fluid leaving the proximal convoluted tubule (PCT) in the absence of antidiuretic hormone (ADH) is isotonic.

The PCT is the site of maximum water reabsorption under normal physiological conditions. It reabsorbs around 60% of the filtered NaCl and water and other nutrients such as glucose and amino acids. Isosmotic reabsorption occurs in the PCT, so the amount of solute and water reabsorbed are coupled and proportional to each other. Here, sodium is reabsorbed along with the water. The large amounts of sodium and water reabsorption are essential in maintaining the ECF volume.

In addition to the isosmotic reabsorption, the water reabsorption in the PCT also occurs through aquaporin-1 which are active water channels. Reabsorption of sodium and water in PCT is not affected by ADH levels. Thus even in the absence of ADH, fluid is isotonic in PCT.

Solution to Question 17:

2,3-Bisphosphoglycerate (BPG) causes the unloading of oxygen to tissues by binding to the beta subunit of Hb.

2,3 BPG is formed from glycolysis. It is found in high amounts in red blood cells and in trace amounts in most other cells. 2,3 BPG facilitates increased oxygen delivery by reducing the affinity of hemoglobin for oxygen.

Hemoglobin exists in 2 forms in the blood. The deoxyhemoglobin form is called the taut or T form. This hemoglobin has a low affinity for oxygen. The oxyhemoglobin form is called the relaxed or R form. This hemoglobin has a high affinity for oxygen.

2,3 BPG binds more selectively to deoxyhemoglobin and stabilizes the taut conformation of hemoglobin (Option C).

Other options:

Options A and D: The process of unloading oxygen by 2,3 BPG is independent of ATP production.

Solution to Question 18:

The curve 'A' shows a hyperbolic curve that is characteristic of oxygen dissociation in myoglobin.

Myoglobin is an iron-containing component of the skeletal muscle. Myoglobin can bind only to one oxygen molecule and hence lacks co-operative binding. It has a higher affinity to oxygen than hemoglobin and hence releases oxygen only at low PO_2 values. This leads to a leftward shift in the myoglobin oxygen dissociation curve.

Myoglobin is present in large quantities in muscles that are commonly used for sustained contraction. During such contractions, muscle blood supply is decreased due to compression. Myoglobin facilitates oxygen supply in these conditions.

Other options:

Option B: Carbon monoxide (CO) reacts with hemoglobin to form carbon monoxyhemoglobin (carboxyhemoglobin or COHb). The affinity of hemoglobin for CO is 210 times its affinity for O_2, and COHb liberates CO very slowly. An additional difficulty is that when COHb is present, the dissociation curve of the remaining HbO_2 shifts to the left, decreasing the amount of O_2 released.

Option C: Hemoglobin normally has ferrous ion. When it is oxidized to ferric iron, it is called methemoglobin. It causes dark-coloured blood.

Option D: Fetal hemoglobin (HbF) is structurally similar to hemoglobin A except for the gamma chains. It binds less avidly to 2,3 bisphosphoglycerate and hence has higher affinity for oxygen leading to left ward shift in the dissociation curve.

Solution to Question 19:

The organ with the least arteriovenous oxygen difference is the kidney.

The arteriovenous oxygen difference is the amount of oxygen the tissues take up from the blood. If the amount of oxygen utilized by the tissues is more, the greater the arteriovenous oxygen difference.

The renal cortical blood flow is comparatively greater as the main function is to filter large volumes of blood through the glomeruli. Here, the oxygen extracted from the blood is relatively less. The arteriovenous oxygen difference for the kidney is only 14 mL/L of blood, which is very less when compared to the brain and heart.

Organs	Arteriovenous Oxygen Difference (mL/L)
Brain	62
Kidney	14
Skin	25
Liver	34
Heart muscles	114

Solution to Question 20:

The region marked between 4 and 5 in the action potential curve indicates repolarization due to the efflux of K+ ions across the cell membrane.



The sequence of events during the action potential occurs in the following sequence.

1. Resting state: Both channels are closed.

2. Na+ channels open in response to depolarizing stimulus. Sodium enters the cell. The membrane reaches threshold potential.

3. The sodium influx causes the opening of more Na+ channels through a positive feedback mechanism. Rapid depolarization occurs.

4. Sodium conductance decreases.

5. Na+ channels close and K+ channels open. Due to K+ efflux, repolarization occurs.

6. Na+ channels are reset and are ready to respond to another depolarizing stimulus (relative refractory period). During this phase, the K+ channels remain open and slowly return to the closed state causing after-hyperpolarization.

7. Both channels are closed. The membrane returns to RMP.

Solution to Question 21:

The value of normal tissue oncotic pressure is 12 mmHg. It refers to the interstitial osmotic pressure.

The interstitial oncotic pressure, while very small, is the pressure exerted by a small amount of protein present in the interstitium that tends to pull fluids out of the capillaries. It is different for each organ based on the level of protein components. It is mostly negligible, but sometimes can be as high as 6 mmHg in the brain. The values are also positive in the liver and kidneys.

The colloid osmotic pressure or oncotic pressure is the osmotic pressure exerted due to the presence of colloidal proteins like albumin and globulin within the plasma. The colloidal proteins are impermeable to the semi-permeable cell membrane and cannot cross the membrane. Thus, it exerts osmotic pressure, which pulls fluid into the capillaries.

The lowest number in the given options is 12 mmHg and, hence, is the best possible answer to this question.

Solution to Question 22:

The signaling sequence in a type 2 membrane protein is located inside the membrane (transmembrane region).

A membrane protein can be either integral protein (spans the thickness of the cell membrane) or peripheral protein (attached to the peripheral side of the membrane).

Integral membrane proteins or transmembrane proteins are embedded in the cell membrane, spanning through the entire thickness. They are anchored by hydrophobic interactions and require the use of detergent for solubility. Examples are ligand-binding receptors for hormones or neurotransmitters, transport proteins like Na+-K+ ATPase, pores, ion channels, cell adhesion molecules, and GTP-binding proteins (G proteins).

Peripheral proteins are attached only to one surface of the cell membrane and do not penetrate all the way through. They are often attached to integral proteins. Examples are cell surface receptors, enzymes, ankyrin.



Signaling sequences are 16 to 30 amino acid residues in length and located at the N-terminal extension of the membrane proteins. After an appropriate trigger, they are either cleaved (and bind to MHC class I of the endoplasmic reticulum) or act as a membrane anchor (transporting proteins across the membrane).

Integral proteins can be divided into:

• Type I integral proteins have a cleavable N-terminal signal sequence (towards the cytosol). Examples include histocompatibility antigens, glycophorin, and membrane immunoglobulin.

• Type II integral proteins have a non-cleavable N terminal combined signal/anchor sequence. Examples include transferrin receptors and glycosyl transferases.

Solution to Question 23:

IP3 is the primary second messenger involved in the contraction of ciliary muscles to focus on a near object.

On focusing on a near object, stimulation of the parasympathetic nerve fibers leads to the contraction of the bilateral ciliary muscle, via short ciliary nerves, which relaxes the lens ligaments. The lens becomes thicker and the increase in the refractive index allows the eye to focus on the near object. This is known as the accommodation reflex.

In the ciliary muscles of the eye, contraction is mediated via M3 > M2 receptors, while relaxation is mediated via the β2 receptor.

M3 receptors act via the formation of inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG), leading to an increase in intracellular Ca2+ and activation of protein kinase C, respectively. The increase in intracellular Ca2+ sets up a chain reaction which eventually leads to contraction of the ciliary smooth muscle of the eye.

Thus, in the contraction of the ciliary muscle during focusing on a near object, the second messenger involved is the IP3-DAG pathway.

Solution to Question 24:

Adipose tissue is dependent on insulin for glucose uptake.

The uptake of glucose in skeletal muscles is also dependent on insulin but may be mediated via non-insulin-dependent mechanisms as well (such as muscle contractions). Therefore the best answer out of the given options is adipose tissue.

Glucose uptake by skeletal muscle, cardiac muscle, and adipose tissues is via GLUT-4 which is insulin-dependent. In the presence of insulin, GLUT-4 receptors translocate to the cell membrane and hence, glucose uptake by the cell occurs.

Most of the other GLUT transporters are not insulin sensitive and are constitutively expressed in the cell membrane. Ex: GLUT-1 in RBCs, brain and kidneys. (GLUT-3 is also present in the brain and kidneys)

Solution to Question 25:

During the second phase of salivary secretion, Na+ is actively reabsorbed, Cl- is reabsorbed passively, K+ is secreted actively and HCO3- is secreted passively (part of the secretion may be active as well, but among the given options, option D is the most accurate with respect to all the ions).

Salivary secretion occurs in two stages:

The first stage involves acini, which secrete ptyalin and/or mucin together with some ions. The composition of the ions is similar to that of the extra-cellular fluid.

The second stage involves the ducts. As the primary salivary secretion flows through the ducts, two active transport processes occur:

• Na+ is actively re-absorbed and K+ is actively secreted, in exchange for Na+.

• HCO3- is secreted by the ductal epithelium into the lumen of the salivary ducts. Some of this secretion occurs by passive bicarbonate exchange for Cl- and partly active secretion.

As a result, the Na+ and Cl- concentrations are lower than that of plasma, while K+ and HCO3- concentrations are higher than that of plasma.

Composition of ions:

• Under resting conditions, the Na+ and Cl- concentrations in saliva are about 15 mEq/L each, which is 1/7th to 1/10th of the plasma concentrations.

• K+ concentration is about 30 mEq/L, which is seven times higher than that of plasma.

• HCO3- concentration is 50 to 70 mEq/L, which is two to three times higher than that of the plasma.

Solution to Question 26:

Macula densa are the specialized cells present at the distal end of the thick ascending loop of Henle, where it reaches the glomerulus. Distal convoluted tubule starts from macula densa.

Macula forms the part of the juxtaglomerular apparatus.



Signals from the renal tubules affect filtration in its glomerulus. As the rate of flow through the ascending limb of the loop of Henle and the first part of the distal tubule increases, glomerular filtration in the same nephron decreases, and vice versa. This is known as tubuloglomerular feedback which maintains the constancy of the load delivered to the distal tubule.

The sensor is the macula densa. The Na+ and Cl– in the fluid enter the macula densa cells via theNa+–K–2Cl– cotransporter in their apical membranes. The increased Na+ increases the Na+-K+ ATPase activity causing ATP hydrolysis and the formation of adenosine. This acts on adenosine A1 receptors on the macula densa cells, causing the release of Ca2+ into the vascular smooth muscle in the afferent arterioles, leading to vasoconstriction and decreased GFR.



Solution to Question 27:

Fick principle, thermodilution, and ECHO can be used to measure cardiac output.

Cardiac output is the amount of blood pumped by the heart into the aorta every minute. The average cardiac output is 5L/min and can be measured by various indirect methods.

Fick's principle uses the total amount of oxygen absorbed from the air by the lungs and the arteriovenous oxygen difference of blood to calculate the cardiac output using the following formula:

Cardiac output (L/min) = (O_2 absorbed per minute by the lungs (mL/min)) / (Arterio-venous O_2 difference(mL/L of blood))

Thermodilution method employs cold saline which is infused into the right atrium through a double-lumen Swan–Ganz thermodilution catheter and the temperature change is measured at the pulmonary artery using a thermistor at the end of the catheter. The mean decrease in the temperature of blood is inversely proportional to the cardiac output.

An ECHO uses ultrasound waves from a transducer placed on the chest wall to measure the velocity of blood flowing from the left ventricle into the aorta. Stroke volume can be calculated from this. Cardiac Output = Heart Rate X Stroke Volume.

The other methods that can be used to measure cardiac output include impedance cardiography a.k.a. thoracic electrical bioimpedance, electromagnetic or ultrasonic flowmeter,and indicator dilution method.

Ventilation-perfusion scan is not a method of cardiac output measurement. It is used to assess pulmonary circulation.

Solution to Question 28:

The correctly matched answer is 1-D,2-A,3-B,4-C.

Each organelle has some specific enzymes are called marker enzymes. After centrifugation, the separated organelles are identified by the detection of marker enzymes in the sample.

Cellular components	Marker enzymes
1. Mitochondria	D. ATP synthase
2. Peroxisomes	A. Catalase
3. Golgi complex	B. Galactosyl transferase
4. Endoplasmic reticulum	C. Glucose-6-phosphatase

Solution to Question 29:

The true statements are 1,4.

Sympathetic efferents are distributed extensively all over the body whereas parasympathetic distribution is limited to head, neck, and trunk regions.

Sympathetic fibers travel short distances to the sympathetic ganglionic chain, before finally synapsing and then have a long course to supply the target organs. Parasympathetic fibers travel a longer distance to their respective ganglia which are located close to the target organs.

Motor neurons are located in the ventral horn of the spinal cord from where the axons come out and divide into several branches, each of which supplies an individual muscle fiber. A motor unit comprises a motor neuron and a group of skeletal muscle fibers it innervates.

Solution to Question 30:

The normal insensible water loss is 50 mL/hour.

In human beings, there is a loss of water through vaporization from the respiratory tract (300–400 mL) and diffusion from the skin (300–400 mL). This is called insensible water loss. It is about 50 mL/hour. This insensible water loss via the skin is independent of sweating and occurs in people without sweat glands as well.

Solution to Question 31:

The correct sequence of vomiting is 6, 4, 2, 3, 1, 5.

6. Reverse peristalsis in the small intestine.

4. Relaxation of the pyloric sphincter.

2. Strong contractions in the stomach.

3. Inspiration against a closed glottis.

1. LES is open and UES is closed.

5. LES opens and UES opens.

The sensory signals that initiate vomiting are located in the pharynx, esophagus, stomach, and upper portions of the small intestines. Excessive distention or irritation of the duodenum provides a strong stimulus for vomiting. Area postrema constitutes the vomiting centre.

In the early stage, reverse peristalsis begins often minutes before vomiting. It begins far down in the ileum and travels up the intestine. This process pushes a large share of lower small intestine contents to the duodenum and stomach. Relaxation of the pyloric sphincter occurs during this process.

At the onset of vomiting, strong intrinsic contractions are seen in the duodenum and stomach. Once the vomiting centre has been stimulated, inspiration occurs against the closed glottis. It prevents the flow of vomitus into the lungs. The lower esophageal sphincter (LES) is open and

the upper esophageal sphincter (UES) is closed at this stage.

The muscles of the abdominal wall contract, building intra-gastric pressure. Finally LES relaxes completely and UES also relaxes allowing the expulsion of the gastric contents.

Note: The esophageal smooth muscle functions as LES at the gastro-esophageal junction. It normally remains tonically constricted. Receptive relaxation of LES occurs when the peristaltic wave passes down the esophagus.

Physiology NEET 2022

Question 1:

A person after sleeping overnight with the arm under his head now experiences paresis but no numbness in the morning. Which of the following is the best explanation for it?

a) C fibers are more sensitive to pressure than A fibers.

b) A fibers are more sensitive to hypoxia than B fibers.

c) A fibers are more susceptible to pressure changes than C fibers.

d) A fibers are more susceptible to hypoxia than C fibers.

Question 2:

Which of the following mechanisms is seen in the baroreceptor reflex?

a) Feedforward

b) Positive feedback

c) Negative feedback

d) Adaptive control regulation

Question 3:

The image below depicts which of the following types of cell-to-cell signaling?



a) Paracrine

b) Autocrine

c) Endocrine

d) Merocrine

Question 4:

Which of the following statements is true regarding the given cystometrogram?



a) Segment Ia is due to residual urine

b) Segment Ib is due to Laplace law

c) Micturition fails to happen in segment II

d) The dotted line represents that micturition has occurred

Question 5:

In a patient, the hypothalamic thermostat was reset from point A to point C as shown below. Which of the following happens in stage A compared to stage B?



a) Shivering

b) Sweating

c) Increased blood flow to skin

d) Inhibition of chemical thermogenesis

Question 6:

A 65-year-old man suffered from a stroke 2 days ago. He now presents with involuntary, violent, and flinging movements of the limbs on one side. What is the likely site of lesion in this patient?

a) Subthalamic nuclei

b) Globus pallidus

c) Putamen

d) Caudate nucleus

Answer Key

Question No.	Correct Option
1	c
2	c
3	a

4	b
5	a
6	a

Detailed Explanations

Solution to Question 1:

In the above scenario, the patient has developed paresis (partial paralysis) but the pain sensation is intact (no numbness). This is because A fibers that carry motor, touch, and pressure sensations are more susceptible to pressure changes/compression than C fibers that carry pain sensations.

The compression of a large diameter nerve fiber leads to a reversible physiologic conduction block or neuropraxia.

The relative susceptibility of nerve fibers to conduction blocks produced by various agents:

Susceptibility	Most susceptible	Intermediate	Least susceptible
Local anesthetic	C	B	A
Hypoxia	B	A	C
Pressure	A	B	C

Solution to Question 2:

The negative feedback mechanism is seen in the baroreceptor reflex. If the activity of a particular system is increased, the controller system initiates the negative feedback mechanism, which prevents overactivity and ensures normal homeostasis.

Baroreceptors are stretch receptors located in the heart and the adventitia of the blood vessels. They are stimulated by the stretching of the walls of the blood vessels when there is an increased volume or blood pressure in the vessel. They are important in the immediate management of blood pressure. They are of two types: carotid sinus and aortic arch receptors (high-pressure regions) and cardiopulmonary receptors (low-pressure regions).

An increase in blood pressure stimulates the baroreceptors. These signals are transmitted through small Hering's nerves to the glossopharyngeal nerves and then to the nucleus tractus solitarius (NTS) in the medulla. These signals release glutamate and activate the caudal ventrolateral medulla (CVLM). This in turn inhibits the rostral ventrolateral medulla (RVLM) considered the vasoconstrictor center for the brain and the heart.

Simultaneously, this pathway also excites the vagal parasympathetic center. The stimulation of the vagus results in a slowing of impulse generation in the SA node and

a slowing of conduction velocity in the AV node.

These neural changes produce vasodilation, venodilation, hypotension, bradycardia, and a decrease in cardiac output.





Other options:

Option A: The feedforward mechanism responds to the anticipation of a change in a regulated variable. For example, when a meal is in the digestive tract, a feedforward mechanism increases the secretion of insulin that promotes the cellular uptake and storage of ingested nutrients.

Option B: Positive feedback mechanism refers to the enhancement or amplification of the output so that it moves in the direction of the initial change. An example is seen in the luteinizing hormone (LH) surge where the estrogen positively regulates the further secretion of LH.

Option D: The adaptive control mechanism refers to the delayed type of negative feedback mechanism. When the feedforward mechanism has an error, the correction is made by a retrospective feedback mechanism called the adaptive control mechanism.

Solution to Question 3:

The given image shows the paracrine cell-to-cell signaling mechanism. In paracrine signaling, the signaling molecules released by a cell only affect target cells in close proximity to it.

An example of paracrine signaling is the release of endothelin-1, a potent vasoconstrictor released by the endothelial cells.

Synaptic signaling is a special case of paracrine signaling (for chemical synapses) between neurons and target cells.

Other options:

Option B: In the autocrine signaling mechanism, the cells respond to the signaling molecules that are released by themselves, which is the site of release and action within the same cell. Note that the signaling molecules are first secreted out of the cell and then act on the same cell through receptors. An example is the proliferation of activated T-cells by IL-2 released by the same cell after it interacts with antigen-presenting cells.

Option C: In endocrine signaling, a signaling molecule, a hormone (for example insulin, FSH, LH), is usually carried by the blood from its site of release to act on its target at a distance.

The below image shows the different types of extracellular cell-to-cell communication mechanisms.



Option D: Merocrine is a type of glandular secretion and not an extracellular signaling mechanism. In this, the secretions are poured out by exocytosis and the cell remains intact. An example of the merocrine gland is the goblet cell and eccrine sweat gland.

An illustration depicting this mechanism is given below.



Solution to Question 4:

The true statement regarding the given cystometrogram is that segment Ib is due to Laplace law.

Cystometrogram is a plot of intravesical pressure against the volume of fluid in the bladder. The curve shows three segments:

• Ia- initial slight rise in pressure when the first increments in the volume are produced

• Ib- long, nearly flat segment as further increments are produced

• II- sudden, sharp rise in pressure as the micturition reflex is triggered (not failure).

• The dotted line- pressure-volume relation if the micturition had not occurred.

The flatness of segment Ib is a demonstration of the law of Laplace. The law of Laplace states that the pressure (P) in a spherical viscus is equal to twice the wall tension (T) divided by the radius (R). This is expressed as P= 2T/R. In the bladder, the tension increases as the bladder get filled. Since the radius also increases there is only a slight increase in pressure until the bladder is relatively full.

The first urge to void is felt at a bladder volume of about 150 mL. A marked sense of fullness is felt at a bladder volume of 400ml.

Micturition reflex is the result of sensory signals from the stretch receptors in the bladder wall, which are conducted to the sacral segments of the cord through the pelvic nerves and then efferent signals conducts back again to the bladder through the parasympathetic nerve fibers.

The micturition reflex pathway is given below.



Solution to Question 5:

In the above scenario, the patient's hypothalamic thermostat has been reset from point A to a higher temperature (point C). The body responds to this change by using certain measures in stage A such as shivering.

In stage A the body activates various heat gain mechanisms to match the hypothalamic temperature setpoint. These include:

• Shivering- the shivering center is located in the posterior hypothalamus. It is normally inhibited by signals from the heat center in the anterior hypothalamic-preoptic area but is excited by cold signals from the skin and spinal cord. ■A sudden increase in heat setpoint results in the activation of this center. It then transmits non-rhythmic signals through bilateral tracts through the lateral columns of the spinal cord, and finally to the anterior motor neurons. They increase the tone of the skeletal muscles throughout the body. As the tone rises above a certain critical level, shivering begins. During maximum shivering, body heat production can rise to four to five times normal.

• Vasoconstriction (decreased blood flow to skin) is mediated by increased sympathetic drive by the activation of the posterior hypothalamic sympathetic center.

• An increase in chemical thermogenesis in which heat production takes place in the brown fat by the mechanism of uncoupling of oxidative phosphorylation. This is the predominant mechanism seen in neonates.

In stage B, the hypothalamic temperature setpoint is lower than the normal body temperature. Hence body activates various heat loss mechanisms to match the hypothalamic temperature setpoint. It includes vasodilatation of the skin blood vessel, sweating, and decreased thermogenesis.

The temperature of the body is regulated majorly by a nervous feedback mechanism that operates through temperature-regulating centers located in the hypothalamus. Minor regulation is done by the thermal receptors in the skin and the deep thermal receptors present in the spinal

cord and abdominal viscera.



Solution to Question 6:

The above clinical scenario is suggestive of hemiballismus. A lesion in subthalamic nuclei is responsible for this condition.

A lesion in the subthalamic nucleus causes sudden, violent, involuntary, and flinging movements confined to one side of the body, commonly affecting the proximal muscles of the limbs, a condition called hemiballismus. The movements are seen contralateral to the side of the lesion, i.e., involuntary movements of the right arm and leg are seen in the lesion of the left subthalamic nucleus and vice versa.



Basal ganglia circuits have both direct and indirect pathways. The direct pathway stimulates the cortex to produce movement, whereas the indirect pathway inhibits it. The striatal activity in the direct pathway is mediated by D1 dopamine receptors and in the indirect pathway by D2 dopamine receptors.

The subthalamic nucleus is a part of the indirect pathway. Hence, a lesion at subthalamic nuclei leads to loss of inhibitory effect of the indirect pathway resulting in dopaminergic overactivity. This is responsible for features of hemiballismus.

This movement disorder is usually self-limiting and tends to resolve spontaneously after weeks or months. Dopamine antagonistic agents can be given to manage it as well.

Other options:

Option B: A lesion in the globus pallidus can give rise to athetosis. This involves spontaneous and continuous writhing movements of the hand, arm, neck, or face.

Options C and D: A lesion in the putamen and caudate nucleus can give rise to chorea. Chorea is defined as rapid, semi-purposeful, graceful, dance-like nonpatterned involuntary movements involving distal or proximal muscle groups.

Physiology NEET 2023

Question 1:

Identify the marked region in the given spirometry.



a) Functional residual capacity

b) Tidal volume

c) Expiratory reserve volume

d) Expiratory capacity

Question 2:

A man was brought to the emergency after he suddenly became unconscious while working in the field. On examination, his temperature was 105 degrees Fahrenheit and his skin turgor was decreased. Which of the following would not be seen in the patient?

a) Tachypnea

b) Hypotension

c) Sweating

d) Red hot skin

Question 3:

Identify the correctly matched pair of substances with their renal clearance from the graph given below.



a) A - Glucose, B - PAH, C - Bicarbonate and D - Inulin

b) A - Glucose, B - Bicarbonate, C - Inulin and D - PAH

c) A - PAH, B - Inulin, C - Glucose and D - Bicarbonate

d) A - Inulin, B - Glucose, C - Bicarbonate and D - PAH

Question 4:

Identify the true statement regarding the given nerve action potential curve.



a) Threshold point is at A

b) At point E, the nerve is more excitable

c) Point C to D is due to opening of Na+ and closure of K+ channels

d) Point B to D is refractory period

Question 5:

Identify the type of transport across the cell membrane given below.



a) Simple diffusion

b) Facilitated diffusion

c) Primary active transport

d) Secondary active transport

Question 6:

A preterm baby who was delivered at 28 weeks developed respiratory distress syndrome. Which of the following is true about surface tension and compliance in this baby?

a) Surface tension - decreased; Compliance - increased

b) Surface tension - increased; Compliance - decreased

c) Both surface tension and compliance decreased

d) Both surface tension and compliance increased

Question 7:

A patient came to the hospital after a road traffic accident. He had severe muscle injury and his serum K+ level was found to be 5.5 mEq/L. What will happen to the resting membrane potential in this patient?

a) No change

b) Becomes more negative

c) Becomes more positive

d) First becomes more positive then negative

Question 8:

A patient met with a road traffic accident and developed a cervical spine injury. The fracture fragment had pierced the lateral aspect of the dorsal column tract. Which of the following findings is seen in this patient?

a) Absence of ipsilateral lower limb proprioception

b) Absence of fine motor movement of fingers

c) Absence of ipsilateral arm proprioception

d) Absence of contralateral lower limb proprioception

Question 9:

Identify the true statement regarding the point marked on the myocardial action potential curve.



a) Due to opening of Na+ and closure of fast K+ channels

b) Due to slow but prolonged opening of Ca2+ channels

c) Due to closure of Na+ and opening of slow K+ channel

d) Due to Na+/K+ ATPase

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	d
5	b
6	b
7	c
8	c
9	b

Detailed Explanations

Solution to Question 1:

The marked area in the spirometry shows the total amount of air present in the lung after normal expiration. This is called functional residual capacity.

The various volumes and capacities in a spirometry are shown below.



Solution to Question 2:

The given clinical vignette in a patient with signs of dehydration and a body temperature of 105°F is highly suggestive of heatstroke. Sweating would not be seen in a patient with heatstroke.

The body has various mechanisms to maintain body temperature within a range of 36–37°C. When the body temperature is elevated by exposure to high environmental temperatures or physical exertion, the central control area in the medial preoptic and anterior hypothalamic nuclei is activated. This engages the autonomic pathways that promote sweating and cutaneous vasodilation, causing the heat to dissipate and bring the body to normal temperature.

Heatstroke occurs due to the failure of the heat regulation center of the hypothalamus. The person fails to lose heat by cutaneous circulation and sweating. It is likely to develop when the body temperature rises to a range of 105°F to 108°F. The predisposing factors include high temperature, increased humidity, minor infections, old age, and physical activity.

It is a medical emergency that leads to homeostatic imbalance. The symptoms include dizziness, abdominal distress, vomiting, delirium, and eventually leading to loss of consciousness. On examination, the skin is dry, hot, and flushed, with a complete absence of sweating (anhidrosis). Other signs seen in heat stroke are tachycardia, Kussmaul-type rapid deep breathing (tachypnea), hypotension, cardiac failure, dizziness, muscle cramping, confusion, and seizures.

Heatstroke can cause damage to multiple organ systems, including the brain, kidneys, and liver, which can lead to life-threatening complications. It is essential to initiate rapid cooling and supportive care measures to prevent permanent injury or death. The cooling methods used to treat heatstroke include:

• Cold water immersion: The patient is immersed in a tub of cold water, which is the most effective way to lower body temperature.

• Evaporative cooling: The patient's body is sprayed with water while fans blow air over the body to help facilitate evaporation and cooling.

• Applying cold packs: Applying ice packs to the neck, armpits, and groin can help cool the body by targeting areas where large blood vessels are located close to the skin surface.

• Misting and fanning: Spraying the patient with cool water and using a fan to circulate air can help promote evaporative cooling.

Preventive measures for heatstroke include staying hydrated, taking regular breaks in shaded areas, and avoiding strenuous activities during the hottest part of the day.

Other Options

• Option A: Tachypnea - Rapid, deep breathing (Kussmaul breathing) can be seen in heatstroke as the body tries to eliminate excess heat through increased respiratory rate.

• Option B: Hypotension - Dehydration and vasodilation in an attempt to dissipate heat can result in hypotension in heatstroke patients.

• Option D: Red hot skin - Heatstroke patients often present with hot, flushed skin as a result of increased blood flow to the skin surface and the body's inability to dissipate heat effectively.

Solution to Question 3:

The correctly matched pair of substances and their renal clearance is A - Glucose, B - Bicarbonate, C - Inulin and D - PAH.

The concentration of the substance changes as it passes through the tubules. The tubular cells may remove the substance from the filtrate (tubular reabsorption) or add more of the substance to the filtrate (tubular secretion) or both. The amount of the substance finally excreted equals the amount filtered plus the net amount transferred out by the tubules.



PAH is filtered by the glomeruli and secreted by the tubular cells. Its concentration increases on moving through the length of the nephron due to secretion into tubules. It is represented by curve D.

Inulin is freely filtered by glomeruli and does not undergo any reabsorption or secretion. Hence, its concentration does not change throughout the length of the nephron and its excretion is constant and equal to the glomerular filtration rate. It is represented by curve C.

The majority of bicarbonate reabsorption occurs in the proximal tubule and a small amount of it flows into the distal tubules and collecting ducts. Its reabsorption occurs in all parts of the tubules, except the descending and ascending thin limbs of the loop of Henle. It is represented by curve B.

Substances like amino acids or glucose are reabsorbed along with sodium in the early portion of the proximal tubule. They have a graph that is sloping down as shown in curve A.

Solution to Question 4:

The correct statement regarding the given nerve action potential curve is point B to D is the refractory period.

The sequence of events occurring during an action potential is shown below



• Point A represents the resting state where both channels are closed.

• Na+ channels open in response to depolarizing stimulus and sodium enters the cell. The initiation of action potential occurs only after a certain potential is reached, which is called the threshold potential represented by point B. (Option A)

• After this, the sodium influx causes the opening of more Na+ channels through a positive feedback mechanism. Rapid depolarization occurs to reach point C.

• At point C, Na+ channels close and K+ channels open, causing repolarization due to K+ efflux. (Option C)

• Point D represents continued repolarization due to potassium efflux.

• At point E, Na+ channels are reset and are ready to respond to another depolarizing stimulus. During this phase, the K+ channels remain open and slowly return to the closed state, causing after-hyperpolarization. As a stronger stimulus is required to reach the threshold potential, the excitability of nerve fiber is reduced at point E. (Option B)

• As both channels close, the membrane returns to resting membrane potential back to point A.

Point B-D is a part of the refractory period. During this period, the application of even a stronger stimulus will not elicit an action potential.

The refractory period is divided into an absolute refractory period and a relative refractory period. During the absolute refractory period, the nerve will not be excited no matter how strong a stimulus is given. However, during the relative refractory period, stronger than normal stimuli can cause excitation.

Solution to Question 5:

The given image shows the transport of a molecule across a membrane with the help of a carrier protein. This form of transport is called facilitated diffusion.

When carrier proteins help in the movement of substances in the direction of their chemical or electrical gradients (downhill movement), no energy input is required and the process is called facilitated diffusion.

Glucose transporters or GLUTs are examples of transport proteins in the cell membrane responsible for the transfer of glucose into the cells by facilitated diffusion. Facilitated diffusion involves the passage of molecules or ions inside the cell membrane by binding to a carrier/ transport protein.



Other options -

Option A. Simple diffusion - It is along the concentration gradient but is not carrier-mediated. In the cell, examples of molecules that can use simple diffusion to travel in and out of the cell membrane are water, oxygen, carbon dioxide, and nitrogen. They pass directly through the cell membrane along the concentration gradient, without the use of energy.

Option C. Primary active transport - It is a form of cellular transport requiring energy in the form of hydrolysis of ATP to ship and transfer molecules against gradient differences. The important examples of primary active transport include sodium-potassium ATPase, proton pump, and calcium ATPase.

Option D. Secondary active transport - In this form of transport, two or more substances are transported together across the membrane with help of a carrier molecule. As one of the substances (e.g., sodium) diffuses down its electrochemical gradient, the energy released is used to drive another substance (e.g., glucose) against the gradient. It does not require energy directly from ATP.

Solution to Question 6:

This preterm baby who developed respiratory distress syndrome has surfactant deficiency. In case of surfactant deficiency, the surface tension increases and compliance decreases.

Surfactant is secreted by the type 2 pneumocytes in the alveoli. Dipalmitoyl lecithin/dipalmitoylphosphatidylcholine (DPPC) is a major component of lung surfactant. It reduces surface tension inside the alveoli, as shown below.



Compliance is defined as the extent of lung expansion for each unit increase in pressure. An important factor that affects lung compliance is surface tension. As surface tension increases, the pressure required to keep the alveoli open also increases. As more pressure is required to expand the alveoli, lung compliance will be reduced.

Solution to Question 7:

In the given clinical scenario, the patient has hyperkalemia. It would result in the resting membrane potential becoming more positive.

The resting membrane potential represents an equilibrium situation where the concentration gradient of ions across the membrane is equal and opposite to the electrical gradients. At a normal potassium concentration of 3.5–5.0 mEq/L, the equilibrium potential for potassium is –90 mV. This is close to the resting membrane potential of neurons, which is about –70 mV.

A decrease in the extracellular sodium concentration reduces the size of the action potential but has little effect on the resting membrane potential. However, the intracellular and extracellular potassium (K+) concentrations are the major determinants of the RMP.

This is because

• At rest, there are more open K+ channels compared to Na+ channels; therefore, the membrane permeability of K+ is greater.

• The RMP is close to the equilibrium potential of K+.

In hyperkalemia, the extracellular level of K+ is increased. Thus, the RMP moves closer to the threshold to elicit an action potential, leading to a more excitable nerve. This means the RMP has become more positive.

In hypokalemia, the extracellular level of K+ is reduced. Thus, the RMP is reduced or becomes more negative and the nerve is hyperpolarized.

Hyperkalemia can lead to various clinical manifestations depending on the severity of the condition. Mild to moderate hyperkalemia may be asymptomatic or cause nonspecific symptoms such as fatigue, muscle weakness, and paresthesias. Severe hyperkalemia can lead to life-threatening complications such as cardiac arrhythmias and paralysis. The management of hyperkalemia includes treating the underlying cause, administering medications to stabilize the cardiac membrane (e.g., calcium gluconate), promoting the movement of potassium into cells (e.g., insulin and glucose, beta-agonists), and enhancing the removal of potassium from the body (e.g., diuretics, ion-exchange resins, or hemodialysis).

Solution to Question 8:

The above clinical scenario of transection of the spinal cord in the lateral aspect of the dorsal column would affect the fasciculus cuneatus. This lesion would result in absence of ipsilateral proprioception of the arm.

The fibers in the dorsal column are organized in such a way that the sacral segments are positioned most medially and those from the cervical segments are positioned most laterally.

These are called

• Fasciculus gracilis - carrying sensations from the ipsilateral lower limb

• Fasciculus cuneatus - carrying sensations from the ipsilateral upper limb.

This patient with an injury to the lateral part of the cervical cord's dorsal column tract will have damaged fasciculus cuneatus. This would result in loss of touch, vibration, and proprioception in

the arm of the same side of the lesion.

The extent of neurological deficits resulting from a spinal cord injury depends on the level of the injury and the severity of the damage. Injuries may be classified as complete or incomplete, depending on whether there is a total or partial loss of sensory and motor function below the level of injury. The key to solving this question lies in recognizing that the question mentions that the damage is to the lateral aspect of the dorsal column tract.

Other options:

Option A: Absence of ipsilateral lower limb proprioception - This would be seen if the fasciculus gracilis was affected, which is responsible for carrying sensations from the ipsilateral lower limb. However, in this case, the injury is to the lateral part of the dorsal column, affecting the fasciculus cuneatus.

Option B: Absence of fine motor movement of fingers - This would be a result of damage to the corticospinal tract, which carries motor information. The question is specifically focused on the dorsal column tract, which is responsible for carrying sensory information.

Option D: Absence of contralateral lower limb proprioception - Proprioception is carried by the dorsal column pathway, and it is affected ipsilaterally. Damage to the lateral part of the dorsal column would not result in contralateral loss of proprioception.

Solution to Question 9:

The myocardial action potential curve marked above represents the plateau phase, which is due to the slower and prolonged opening of calcium channels.

Two types of voltage-gated Ca2+ channels (T- and L-types) are present in myocardial cells. The opening of the slower L-type Ca2+ channels is mostly responsible for changes in the action potential. The action potential curve of cardiac muscles consists of 5 phases, as shown below.



Phase 0: Depolarization - Opening of voltage-gated sodium channels (fast sodium channels) leads to sodium influx.

Phase 1: Initial repolarization - Closure of Na+ channels and opening up of K+ channels leads to potassium efflux.

Phase 2: Plateau - Slower but prolonged opening of voltage-gated Ca2+ channels, leading to calcium influx and reduced permeability of K+ channels, causes the graph to plateau

Phase 3: Rapid Repolarization - The closure of Ca2+ channels and opening of K+ at the end of the plateau phase causes repolarization.

Phase 4: Resting membrane potential - Is reached at the end of phase 3

Physiology INI-CET 2023

Question 1:

Which of the following hormone acts via this pathway?



a) Leptin

b) Calcitonin

c) Vasopressin

d) Aldosterone

Question 2:

Which of the following options is false regarding the measurement of temperature in a febrile patient?

a) Rectal temperature is ideal for measuring the core body temperature

b) Rectal temperature is lower than axillary temperature

c) Tympanic membrane temperature is lower than rectal temperature

d) Rectal temperature is higher than oral temperature.

Question 3:

Which of the following statements is/are true about skeletal muscle?

a) Only 2

b) Only 3

c) 3, 4

d) 1, 4

Question 4:

Cells with prolific smooth endoplasmic reticulum secrete_____

a) Eicosanoid hormones

b) Steroid hormones

c) Amine hormones

d) Peptide hormones

Question 5:

At which volume or capacity, pressure measured at the upper airway becomes zero?

a) Residual volume

b) Functional residual capacity

c) Total lung capacity

d) Inspiratory capacity

Question 6:

Arrange the following bands in the sarcomere from the periphery to the center.

a) 1,2,3,4

b) 2,3,4,1

c) 1,3,4,2

d) 1,4,2,3

Question 7:

Which of the following are incorrectly paired?

a) Gastrin - G cells, Antrum

b) Secretin - S cells, Duodenum

c) CCK - I cells, Duodenum and jejunum

d) Motilin - M cells, Duodenum and Jejunum

Question 8:

Which of the following is responsible for fetal heart acceleration?

a) Carotid body chemoreceptors

b) Aortic baroreceptors

c) Autonomic brain stem regulation

d) ANP

Question 9:

Our body was built to identify toxic substance by the recognition of bitter taste. Which of the following substances is responsible for the bitter taste?

a) Amino acids

b) Alkalaoids

c) Hydrogen ions

d) Aldehydes

Question 10:

The type of hypoxia based on the graph is_____ (Consider the plot in red as normal)



a) Stagnant hypoxia

b) Histotoxic hypoxia

c) Hypoxic hypoxia

d) Anemic hypoxia

Question 11:

Assertion: RMP is dependent on proteins and phosphate ions. Reason: Diffusion potential can be calculated using Nernst Equation.

a) The assertion is independently a true statement, but the reasons is independently a false statement.

b) The assertion is independently a false statement, but the reasons is independently a true statement.

c) Both assertion and reason are independently true and the reason is not the correct explanation for the assertion

d) Both assertion and reason are independently true and the reason is the correct explanation for the assertion

Question 12:

Windkessel effect is shown by all except _____

a) Radial artery

b) Renal Artery

c) Aorta

d) Carotid artery

Question 13:

Increase in blood pressure at the beginning of Valsalva maneuver is due to_____



a) Increase in intrathoracic pressure adding to left ventricular pressure

b) Increase in intrathoracic pressure adding to aortic pressure

c) Increase in left atrial pressure

d) Increase in left ventricular volume

Question 14:

Which of the following statements is false about male sexual development?

a) At eight weeks the embryo is bipotent.

b) Wolffian duct develops into epidymis and vas deferens on either side of the body.

c) Mullerian inhibiting substance inhibits mullerian duct.

d) Dihydrotestosterone leads to the development of secondary sexual characters and external genitalia.

Question 15:

The upstroke of action potential (phase 0) in the sinoatrial node is due to _____ ?

a) Sodium influx via voltage gated sodium channels

b) Calcium currents only

c) Potassium efflux

d) Influx of both sodium and calcium

Question 16:

Which of the following types of pain stimulus is associated with the Carrageenan-induced experimental pain model?

a) Inflammatory stimulus

b) Acute noxious stimulus

c) Deep boring stimulus

d) Neuropathic stimulus

Question 17:

Insulin-independent glucose transporter is present in all of the following except?

a) Brain

b) Adipose tissue

c) RBC

d) Pancreas

Question 18:

Which of the following is responsible for the phase marked C to D in the given nerve action potential?



a) Na+ influx

b) K+ influx

c) Na+ efflux

d) K+ efflux

Question 19:

The given graph shows the effect of PTH on blood calcium and phosphate levels in an experimental animal. How does PTH increase plasma calcium levels?



a) By increasing osetoclastic activity

b) By decreasing renal excretion of calcium

c) By increasing osteoblastic activity

d) By increasing osteoclastic activity and decreasing renal excretion of calcium

Question 20:

In the given cross-section of a muscle spindle, the structure marked below is innervated by which of the following motor nerve fibres?



a) Gamma

b) Alpha

c) Beta

d) Delta

Question 21:

Which of the following is responsible for pneumoperitoneal bradycardia following CO_2 insufflation during laparoscopic surgery?

a) Decreased afterload

b) Vascular injury

c) Air embolism

d) Peritoneal stretching

Question 22:

Which of the following sodium transporters are present maximally in the early part of the PCT?

a) 2,3,4

b) 2,4

c) 1, 2, 3

d) 1,4

Question 23:

Which of the following statements is incorrect with respect to the baroreceptors present in the carotid sinus?

a) They need the support of electromechanical transducers

b) They are located in the deep smooth muscle layer

c) Afferent fibres pass via cranial nerve IX

d) They express several non-selective cation channels

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	b
5	b
6	c
7	d
8	a
9	b
10	d
11	c
12	a
13	b

14	a
15	b
16	a
17	b
18	d
19	d
20	a
21	d
22	c
23	b

Detailed Explanations

Solution to Question 1:

The given image shows the JAK-STAT signaling pathway, which is associated with tyrosine kinase receptors. Among the given hormones leptin acts via this pathway.

Hormones that use receptor tyrosine kinase/ JAK-STAT signaling include the following:

• Growth hormone

• Fibroblast growth factor

• Hepatocyte growth factor

• Insulin

• Insulin-like growth factor-1

• Leptin

• Prolactin

• Vascular endothelial growth factor

Leptin is a hormone that is secreted by fat cells. The physiological effects of leptin are regulating appetite and energy balance. The leptin receptor acts via tyrosine kinase of the Janus kinase (JAK) family, JAK2. Leptin binds to the extracellular part of the leptin receptor, which causes phosphorylation (P) and activation of the intracellular JAK2. This causes phosphorylation of signal transducer and activation of transcription (STAT) proteins, which results in the transcription of target genes and the synthesis of proteins.

Phosphorylation of JAK2 also results in the activation of other intracellular enzyme pathways such as mitogen-activated protein kinases (MAPK) and phosphatidylinositol 3-kinase (PI3K).

Other options:

Hormones such as adrenal (Aldosterone)and gonadal steroid hormones, thyroid hormones, retinoid hormones, and vitamin D, bind with protein receptors inside the cell rather than on the

cell surface. These hormones are lipid soluble and they readily cross the cell membrane and interact with receptors in the cytoplasm or nucleus.

Vasopressin in the epithelial cells acts via V2 receptor. They use adenylyl cyclase as their second messenger system. Vasopressin in the vascular smooth muscles acts via V1 receptor that uses phospholipase C as the second messenger system.

Solution to Question 2:

With regards to the measurement of temperature in a febrile patient, rectal temperature is higher than axillary temperature.

For humans, the conventional normal value for oral temperature is 37°C (98.6°F).

Rectal temperature represents the core body temperature and is less affected by changes in environmental temperature. It remains relatively constant. In comparison, oral temperature is typically 0.5°C lower than rectal temperature. However, the oral temperature can be influenced by various factors such as consuming hot or cold fluids, chewing gum, smoking, and breathing through the mouth.

Tympanic membrane thermometers measure radiant heat emitted by the tympanic membrane and the nearby ear canal, providing an absolute value of temperature. Tympanic membrane readings are generally 0.8°C (1.6°F) lower than rectal temperatures. The axillary temperature measured is lower than the rectal temperature.

Solution to Question 3:

The true statement about skeletal muscle is that the calcium is released from sarcoplasmic reticulum.

Excitation-contraction coupling refers to the process by which the membrane depolarization of the muscle fiber (excitation) leads to the contraction of the muscle.

For the maximum contraction of the skeletal muscle, the current has to penetrate deeply into the muscle fiber to the vicinity of the separate myofibrils. This penetration of current is achieved by transmission of action potentials along transverse tubules (T tubules). The T tubules penetrate through the muscle fiber, from one side of the fiber to the other.

Sarcoplasmic reticulum contains calcium ions in high concentration within its vesicular tubules. These ions are released from each vesicle when an action potential occurs in the adjacent T tubule. (Statement 3)

This action potential when it reaches the T tubule, it is sensed by the dihydropyridine receptors linked to calcium release channels. Actvation of this receptor leads to the opening of calcium release channels that releases the calcium ions into the sarcoplasm surrounding the myofibrils and causing contraction. (Statement 4)

In skeletal muscles, extracellular calcium supply partially contributes for the maintenance of skeletal muscle contraction.(Statement 1)

Solution to Question 4:

Cells with prolific smooth endoplasmic reticulum secrete steroid hormones.

The endoplasmic reticulum contains a complex series of tubules in the cytoplasm of the cell.

• Ribosomes are attached to the cytoplasmic side of the membrane in the rough(granular) endoplasmic reticulum. Protein synthesis and the initial folding of polypeptide chains with the formation of disulfide bonds are the functions of rough ER.

• Ribosomes are absent in the smooth (agranular) endoplasmic reticulum. Free ribosomes are found in the cytoplasm. It is the site of steroid synthesis in steroid-secreting cells and the site of detoxification processes in other cells.

The sarcoplasmic reticulum is a modified endoplasmic reticulum, that functions as an intracellular calcium store, which releases calcium into the cytoplasm stimulating cell contraction, proliferation, and migration.

Solution to Question 5:

Pressure measured at the upper airway becomes zero at functional residual capacity.

The curve of airway pressure obtained is plotted against the volume, which is the pressure-volume curve of the total respiratory system. This curve is used in understanding the fundamental relationship between the compliance of the lungs and the chest wall. It is measured by a spirometer.

The functional residual capacity is the volume of the air that remains in the lungs after normal expiration. The normal value is about 2.5 liters. At the lung volume of functional residual capacity (FRC), which is the relaxed state of the lungs, the pressure inside the lungs is effectively zero.

Other options:

Option A: Residual volume is the air left in the lungs after a maximal expiratory effort.

Option C: Total lung capacity is the sum of tidal volume, inspiratory reserve volume, expiratory reserve volume, and residual volume.

Option D: Inspiratory capacity is the maximum amount of air inspired from the end-expiratory level. It is the sum of tidal volume and inspiratory reserve volume.

Solution to Question 6:

The correct sequence of bands in the sarcomere from the periphery to the centre is: Z line→ A band→ H zone→ M line.

The sarcomere is the contractile unit of the skeletal muscle fibre.

• Z line provides attachment to actin filaments. The area between two adjacent Z lines is called the sarcomere. It is composed of thick and thin filaments:

• Thick filament: Myosin

• Thin filaments: Actin, tropomyosin and troponin

• I band is isotropic to light. It contains only actin filaments.

• A band is anisotropic to light. It contains myosin plus overlapped actin.

• H bands are the lighter areas in the centre of the A band. These are the areas where the the thick filaments do not overlap the thin filaments, when the muscle is relaxed.

• M line is present in the centre of H band. There are slender cross-connections that hold the thick filaments in proper array.



Solution to Question 7:

Motilin is secreted by the Mo cells(not M cells) in the stomach, small intestine, and colon

Motilin is a gastrointestinal hormones that is secreted by enterochromaffin cells and Mo cells in the stomach, small intestine, and colon. It mediates its action via G-protein-coupled receptors on enteric neurons in the duodenum and colon. It produces contraction of smooth muscle in the stomach and intestines in the period between meals.

M cells are microfold cells are located in the epithelium covering mucosa-associated lymphoid tissues of the small intestine such as the Peyer's patches (PPs). Their function is to transport luminal antigens to the underlying lymphoid follicles to initiate an immune response.

Gastrin is secreted by G cells of the antrum, duodenum, and jejunum. It stimulates gastric acid secretion and mucosal growth.

Secretin is secreted by the S cells of the duodenum. Its primary function is to promote the secretion of bicarbonate-rich pancreatic juice by the ductal cells of the pancreas and duodenum. This secretion is poor in enzymes.

Cholecystokinin (CCK) is secreted by I cells of the duodenum, jejunum, and ileum. It stimulates pancreatic enzyme secretion, pancreatic bicarbonate secretion, gallbladder contraction, and growth of the exocrine pancreas. It also inhibits gastric emptying.

Solution to Question 8:

Amongst the given options, the best answer for the site responsible for Fetal heart acceleration is Carotid body chemoreceptors>Aortic baroreceptors>Autonomic brain stem regulation.

Though the brainstem is critical to mediating this reflex, the primary stimulus for initiating the changes in the heart rate is initiated by the carotid chemo and baro receptors, especially the chemoreceptor which is highly sensitive to the changes in O2 levels in the blood. It plays a permissive role on fetal heart rate acceleration.

Arterial chemoreceptors in the carotid arteries>aortic arch, respond to low pH levels and decreased oxygen saturation. These chemoreceptors detect instances of significant fall in blood oxygen levels, setting off the peripheral chemoreflex and heightening both parasympathetic and sympathetic neural activities. The activation of the parasympathetic branch leads to a swift decline in Fetal Heart Rate (FHR), a clinical manifestation identified as intrapartum decelerations. Concurrently, the sympathetic branch prompts significant peripheral vasoconstriction across a range of vascular territories, which results in the elevation of arterial blood pressure, ensuring continuous perfusion to essential organs.

The permissive action on the FHR acceleration comes in instances of alleviated hypoxemia, detected by the chemoreceptors in the carotid body and aortic arch, suppressing the peripheral chemoreflex, leading to a reduction in parasympathetic and sympathetic neural activities. Deactivation of the parasympathetic branch results in a swift increase in Fetal Heart Rate (FHR), typically seen clinically as an increase in intrapartum accelerations.

Solution to Question 9:

Alkaloids cause an intensely bitter taste, that causes the rejection of the food.

Bitter taste sensations are caused by two particular classes of substances:

• Long-chain organic substances that contain nitrogen.

• Alkaloids

Alkaloids cause an intensely bitter taste, leading to the rejection of certain foods. They are found in many medicinal drugs such as quinine, caffeine, strychnine, and nicotine. The bitter taste sensation in many deadly toxins found in poisonous plants is often caused by alkaloids. This helps the person or animal to reject the food.

Sweet taste is caused by chemicals such as sugars, glycols, alcohols, aldehydes, ketones, amides, esters, some amino acids, some small proteins, sulfonic acids, halogenated acids, and inorganic salts of lead and beryllium.

Sour taste is caused by acids (Hydrogen ion concentration). The sour taste is directly proportional to the strength of the acid.

Solution to Question 10:

The graph represents the oxygen-hemoglobin dissociation curve that would be seen in anemic hypoxia and normal. The X-axis represents the partial pressure of oxygen(PO_2). The Y-axis represents the percentage saturation between the hemoglobin and the oxygen.

The graph indicates two lines: a red line representing normal hemoglobin and a purple line representing reduced hemoglobin levels. The red line corresponds to an arterial blood oxygen content of approximately 19 mL/dL, and venous blood oxygen content of 14 mL/dL. On the other hand, the purple line represents an arterial blood oxygen content of 9 mL/dL and venous blood oxygen content of 4 mL/dL.

This scenario demonstrates a 50% decrease in oxygen saturation of hemoglobin, while the partial pressure of oxygen (PO_2) remains within the normal range. This condition is characterized as anemic hypoxia.

In anemic hypoxia, the blood oxygen content is decreased. The patients might not experience any difficulty during rest because 2,3-DPG increases in the red blood cells. However, they may have difficulty during exercise because of a limited ability to increase O_2 delivery to the active tissues.

Conditions that cause a right shift in the oxygen hemoglobin curve are: (Conditions with the increased requirement of O_2 - metabolically active tissues)

- Hypoxia
- Hypercarbia
- Decreased pH
- Increased temperature
- Increased 2,3-biphosphoglycerate

Conditions that cause a left shift in the oxygen hemoglobin curve

- Increased oxygen
- Decreased carbon dioxide
- Increased pH
- Decreased temperature
- Decreased 2,3-biphosphoglycerate
- Fetal hemoglobin.

Solution to Question 11:

Both assertion and reason are independently true and the reason is not the correct explanation for the assertion.

Resting membrane potential (RMP) is the difference in the electrical potentials between the cytoplasm of the cell and the extracellular space. It represents an equilibrium situation where

the concentration gradient of ions across the membrane is equal and opposite to the electrical gradients.

Resting membrane potential is primarily influenced by sodium (Na+) and potassium (K+) ions, along with negatively charged intracellular proteins and organic phosphates. In neurons, the concentration of potassium is higher inside the cell, while sodium ions exhibit the opposite distribution. The establishment of this concentration difference is facilitated by the Na, K ATPase pump. Potassium ions passively move out of the cell, while sodium ions move into the cell.

The Goldman-Hodgkin-Katz equation is utilized to calculate the resting membrane potential since multiple ions are involved. Among these ions, potassium, which has the greatest conductance across the membrane at rest, becomes the primary contributor to the resting membrane potential.

Goldman-Hodgkin-Katz equation

$$EMF\,(millivolts) = -61 \times \log \frac{C_{Na_i^+} P_{Na^+} + C_{K_i^+} P_{K^+} + C_{Cl_o^-} P_{Cl^-}}{C_{Na_o^+} P_{Na^+} + C_{K_o^+} P_{K^+} + C_{Cl_i^-} P_{Cl^-}}$$

The Nernst equation establishes a connection between the diffusion potential and the ion concentration difference across a membrane. The magnitude of the Nernst potential is determined by the ratio of the ion concentrations on the two sides of the membrane.

Nernst equation :

EMF(millivolts) = [61/z] X log [Co / Ci]

• EMF = electromotive force (Nernst potential)

• z = charge of ion

• Co = concentration outside

• Ci = concentration inside

Solution to Question 12:

Windkessel effect is shown by all except the radial artery.

The windkessel effect is seen in the large elastic arteries like the aorta, carotid, and renal artery. The elasticity or windkessel effect is the exclusive property of large and elastic vessels. These vessels when stretched undergo elastic recoil.

The aorta is stretched during systole. This expansion causes temporary storage of blood, allowing more blood to enter the aorta. This buffer stock of blood is released during diastole, to maintain organ perfusion even during diastole.

The image given below shows the windkessel effect seen in the aorta.



Several methods can be used to evaluate the elastic properties of the aortic Windkessel in living humans. One simple approach is to measure the pulse wave velocity along the aorta, where a higher velocity indicates increased stiffness and elastic resistance. Alternative methods involve assessing the ratio between pulse pressure and changes in aortic volume (delta P/delta V). These methods can be noninvasively performed using ultrasound or tomographic techniques.

Solution to Question 13:

Increase in blood pressure at the beginning of the Valsalva maneuver is due to increase in intrathoracic pressure adding to aortic pressure.

Valsalva maneuver involves the performance of forced expiration against a closed glottis. It occurs during coughing, defecation, and heavy lifting. At the onset of straining, the blood pressure is raised because the increase in intrathoracic pressure is added to the pressure of the blood in the aorta. The pressure then falls as the high intrathoracic pressure compresses the veins and decreases the cardiac output.

When the glottis is opened and intrathoracic pressure returns to its normal state, cardiac output is regained, but there is constriction in the peripheral vessels. As a result, blood pressure elevates beyond the normal range, which triggers the baroreceptors, leading to bradycardia and a subsequent decrease in pressure back to normal levels.

Solution to Question 14:

The embryo is bipotent till seven weeks of gestation, not eight.

At the seventh week, the embryo contains both male and female primordial genital ducts. In the case of a female fetus, uterine tubes and the uterus develops from the Mullerian duct. In case of a male fetus, the wolffian duct system on each side develops into the epididymis and vas deferens.

Testosterone is secreted by the Leydig cells of the fetal testis and MIS (Mullerian inhibiting substance) is secreted by the Sertoli cells. MIS is an amino acid that belongs to the family of transforming growth factor β (TGF-β). MIS causes apoptosis of the Mullerian ducts on the side on which it is secreted.

Dihydrotestosterone, a metabolite of testosterone is responsible for the formation of male external genitalia and male secondary sex characteristics. Development of the vas deferens, epididymis, and other structures from the wolffian ducts is due to the action of testosterone.



Solution to Question 15:

The upstroke of action potential (phase 0) in the sinoatrial node is due to calcium currents only. It occurs due to the influx of Ca2+ ions by the opening of voltage-gated L-type channels.

The action potential of the SA node differs from that of the atria, ventricles, and Purkinje fibres in the following ways: (1) In the absence of neural input, the SA node demonstrates automaticity and the ability to produce its own action potentials. (2) It has an unstable resting membrane potential. (3) It has no sustained plateau.

The resting membrane potential (RMP) of the SA node is -55 to -60 millivolts. Phases of this potential are:

• Phase 0 (Upstroke wave)- This is due to the influx of Ca2+ ions by the opening of voltage-gated L-type channels.

• Phase 3 (Downstroke wave)- This is due to the efflux of K+ leading to repolarization. (Option C)

• Phase 4 (Prepotential)- Because of ongoing K+ efflux, hyperpolarization sets in. This makes the channel more permeable to Na+ ions and since these Na+ channels are activated during hyperpolarization they are referred to as funny currents (Ih/ If). They form the first half of the pre-potential. The second half of the prepotential is completed by the opening of the voltage-gated T-(transient) type Ca2+ channel. The slope of Phase 4 in the SA node determines the heart rate.



Solution to Question 16:

The Carrageenan-induced pain model also called the Carrageenan-Induced Paw Edema model is a well-known model of acute inflammation.

It is employed to evaluate the anti-inflammatory properties of a variety of synthetic and natural substances. A non-antigenic phlogistic polysaccharide derived from red seaweed, carrageenan is frequently added to processed foods as a thickening, stabilising, gelling, and emulsifying agent. The activation of the complement system and inflammatory mediators is caused by the sulphated sugars found in carrageenan.

Carrageenan is injected subcutaneously into the plantar surface of the rat paw to cause inflammation. This process results in temporary inflammation and acute swelling, which peaks 3 to 5 hours after injection and subsides in 24 hours.

Postcapillary venules dilate due to the action of carrageenan, causing inflammatory cells and fluid to exude. During this process, a number of proinflammatory mediators are released. The cyclooxygenase pathway's activation is commonly associated with the carrageenan model. A biphasic curve represents the edema caused by carrageenan.

Part of the initial phase of inflammation caused by carrageenan is attributed to injection trauma and the release of acute phase mediators, particularly bradykinin, histamine, and serotonin. The second phase of carrageenan-induced inflammation, which happens about three hours after carrageenan is added, is mostly caused by prostaglandins.

Solution to Question 17:

Insulin-dependent (not independent) glucose transporter is found in the adipose tissue.

Glucose uptake by skeletal muscle, cardiac muscle and adipose tissues is via GLUT-4 which is insulin-dependent.

In the cytoplasm of insulin-sensitive cells, the vesicles contain a pool of GLUT-4 molecules. When the insulin receptors of these cells are activated, the vesicles move rapidly to the cell membrane and fuse with it, inserting the transporters into the cell membrane.

The membrane patches containing transporters are endocytosed when insulin action stops, preparing the vesicles for subsequent insulin exposure. Phosphoinositol 3-kinase is activated by the insulin receptor, which causes the vesicles to migrate to the cell membrane

Most of the other GLUT transporters are not insulin-sensitive and are constitutively expressed in the cell membrane. Ex: GLUT-1 in RBC's and brain, GLUT-2 in the beta cells of the pancreas and GLUT-3 present in the brain and kidneys.

Solution to Question 18:

The phase C to D in the given nerve action potential is caused by K+ efflux as a result of the opening of the K+ channels, resulting in repolarization.

The sequence of events occurring during an action potential is shown below



• Point A represents the resting state where both channels are closed.

• Na+ channels open in response to depolarizing stimulus resulting in Na+ influx (option A). The initiation of action potential occurs only after a certain potential is reached, which is called the threshold potential represented by point B.

• After this, the sodium influx causes the opening of more Na+ channels through a positive feedback mechanism. Rapid depolarization occurs to reach point C.

• At point C, Na+ channels close and K+ channels open, causing repolarization due to K+ efflux.

• Point D represents continued depolarization due to K+ efflux.

• At point E, Na+ channels are reset and are ready to respond to another depolarizing stimulus. During this phase, the K+ channels remain open and slowly return to the closed state, causing after-hyperpolarization. As a stronger stimulus is required to reach the threshold potential, the excitability of nerve fibre is reduced at point E.

• As both channels close, the membrane returns to resting membrane potential back to point A.

Point B-D is a part of the refractory period. During this period, the application of even a stronger stimulus will not elicit an action potential.

The refractory period is divided into an absolute refractory period and a relative refractory period. During the absolute refractory period, the nerve will not be excited no matter how strong a stimulus is given. However, during the relative refractory period, stronger than normal stimuli can cause excitation.

Solution to Question 19:

The parathyroid hormone increases the plasma calcium levels by increasing osteoclastic activity and decreasing renal excretion of calcium.

By regulating intestinal reabsorption, renal excretion, and the exchange of these ions between the extracellular fluid and bone, PTH offers a potent mechanism for controlling the concentrations of calcium and phosphate in extracellular fluid.

The parathormone (PTH) increases Ca+2 concentration by:

• Increased reabsorption of Ca+2 by renal tubules, decreasing the urinary Ca+2 excretion

• Increasing bone resorption, leading to Ca+2 release into the ECF

• Decreased phosphate reabsorption by the renal tubules, increasing the urinary excretion of phosphate

• Conversion of 25-hydroxycholecalciferol to 1,25-dihydroxycholecalciferol which increases intestinal Ca2+ absorption

Activation of the osteoclasts occurs due to activated osteoblasts and osteocytes that send secondary signals to the osteoclasts. Preosteoclast cells become mature osteoclasts when their receptors are activated by RANKL, a significant secondary signal. Activation of the osteoclastic system occurs in two stages: (1) immediate activation of the osteoclasts that are already formed and (2) formation of new osteoclasts.

PTH administration decreases proximal tubular reabsorption of phosphate ions, which results in a rapid loss of phosphate in the urine. At the same time that PTH reduces phosphate reabsorption, it also increases calcium reabsorption through renal tubules. This ultimately leads to a decrease in the renal excretion of calcium.

Furthermore, it increases the reabsorption of hydrogen and magnesium ions while decreasing the reabsorption of ions from amino acids, potassium, and sodium in a manner similar to how it affects phosphate. Distal tubules and the thick ascending loop of the Henle are the primary sites

of the increased calcium reabsorption.



Various steps in formation and hydroxylation of Vitamin D3

©MARROW

Solution to Question 20:

In the given cross-section of a muscle spindle, the structure marked shows intrafusal muscle fibres, which are innervated by gamma(γ) motor neurons.

Each muscle spindle is 3 to 10 mm long and consists of 3 to 12 tiny intrafusal muscle fibers that are pointed at their ends and attached to the glycocalyx of the surrounding large extrafusal muscle fibers. There are two kinds of sensory endings in each spindle, a single primary (group Ia) ending and up to eight secondary (group II) endings.

Intrafusal fibers are of 2 types: (1) nuclear bag fibers - subdivided as dynamic and static nuclear bag fibers and (2) nuclear chain fibers.

Each intrafusal muscle fiber is a tiny skeletal muscle fiber. Since there are either few or no actin and myosin filaments in the centre of each of these fibres, the centre does not contract when the ends do. It performs the role of a sensory receptor instead. The end portions that do contract are excited by small gamma(γ) motor nerve fibers which help control basic muscle tone.

There are two types of γ motor neurons- (a) dynamic that supplies dynamic nuclear bag fibers and (b) static that supply static nuclear bag fibers and nuclear chain fibers.

Other options:

Option B: Alpha (α) motor fibers supply the extrafusal muscle fibers which are the regular contractile units of the muscle. They are on an average 14 micrometers in diameter and these fibers branch many times after they enter the muscle and innervate the large skeletal muscle fibers at the motor endplate. A single alpha nerve fibre can stimulate anywhere between three to several hundred skeletal muscle fibres, which are collectively called the motor unit.

Option C: Beta (β) motor fibers are for touch and pressure.

Option D: Delta (δ) motor fibers are for pain and temperature.





Solution to Question 21:

Vagal stimulation due to peritoneal stretching is responsible for bradycardia following CO_2 insufflation during laparoscopic surgery.

Carbon dioxide (CO_2) and nitrous oxide (N_2O) are used in pneumoperitoneum, a technique that enhances laparoscopic surgery by distending the abdominal cavity and splitting up its contents.

The physiologic effects of CO_2 pneumoperitoneum can be divided into local and systemic effects:

CO_2 pneumoperitoneum is associated with the risk of pathophysiological cardiovascular changes such as severe bradycardia, arrhythmia, and cardiac arrest requiring cardiopulmonary resuscitation. These changes are known to be associated with vagal-mediated cardiovascular reflexes triggered by rapid peritoneum distension due to insufflation.

This can be managed by desufflation of the abdomen, the use of vagolytic medications (such as atropine), and sufficient volume replacement.

Most people can tolerate the hemodynamic and metabolic effects of pneumoperitoneum for a brief duration, and most healthy people can tolerate them for a longer period of time. Long laparoscopic procedures can be challenging for patients whose cardiovascular systems are compromised.

In such cases, alternative gases like helium, neon and argon can be used. They do not cause any metabolic effects, but if they get direct access to the venous system, they can cause gas emboli because they are not as soluble in blood as CO_2 and N_2O.

Gas embolism is treated by positioning the patient in a left lateral decubitus position, head down, to trap the gas in the right ventricle's apex. The gas can then be aspirated out of the right ventricle using a rapidly placed central venous catheter.

Limiting intra-abdominal pressure (IAP) below 12-15 mmHg during insufflation is also effective in preventing pathophysiological changes of pneumoperitoneum.

Local Effects	Systemic effects
Peritoneal distention	Hypercarbia
Vagal reaction	Acidosis
Elevated diaphragm	Increased afterload
Altered venous return	Increased catecholamines

Solution to Question 22:

The sodium transporters present in the early part of PCT are Na-K-ATPase, Na+-glucose cotransporter (SGLT) and Na+-H+ antiporter.

Reabsorption of sodium in different parts of the nephron are:

• Proximal convoluted tubule (PCT):

• Nearly 60% of the filtered sodium is reabsorbed in the PCT. Solutes mainly absorbed in the proximal tubule are sodium, chloride, bicarbonate, phosphate, lactate, glucose, amino acids etc.

• Na^+ is actively pumped from these cells into the interstitial space by Na-K ATPase in the basolateral membrane after travelling by cotransport or exchange via the tubular lumen into the tubular epithelial cells along its concentration and electrical gradients. It extrudes three Na^+ in exchange for two K^+ that are pumped into the cell.

• In early PCT, the main sodium channels used for reabsorption are Na+-glucose cotransporter (SGLT), Na+-H+ antiporter, Na+-amino acid cotransporter, Na+-phosphate cotransporter, etc.

• In distal PCT, due to high chloride concentration in the lumen, it favours NaCl reabsorption, driven by Cl^- gradient.

• The total solute and water reabsorption are proportional to each other. Thus, the entire proximal tubule also reabsorbs nearly 60% of the filtered water (isosmotic reabsorption).

• Loop of Henle:

• The thin descending limb has no active transport of Na+ outside the lumen but the loop is permeable to water, making the tubular fluid hypertonic.

• A thick ascending limb reabsorbs nearly 30% of the Na+ by Na–2Cl–K cotransporter, an example of secondary active transport. This loop is impermeable to water. Since some of the solutes are reabsorbed and water is not reabsorbed, it dilutes the tubular fluid.

• Distal convoluted tubule: Approximately 7% of the filtered load of sodium chloride is reabsorbed in the early distal tubule. This occurs due to the presence of sodium chloride co-transporter. Even this segment dilutes the tubular fluid as it is impermeable to water.

• Cortical collecting tubule: The remaining 3% of the filtered Na+ is absorbed via ENaCs (epithelial sodium channels) in this duct. This portion is regulated by aldosterone to maintain homeostasis in Na+ balance.



Solution to Question 23:

Baroreceptors are not located in the deep smooth muscle layer. They are located in the adventitia layer.

Baroreceptors are mechanoreceptors which respond to stretch stimuli. They are located in the heart and adventitia of the blood vessels. There are two types of baroreceptors: high-pressure arterial baroreceptors and low-pressure volume receptors. Arterial baroreceptors are located within the carotid sinuses and the aortic arch.

The carotid sinus is a small dilation of the internal carotid artery just above the bifurcation of the common carotid into external and internal carotid branches. The afferent nerve fibres from the carotid sinus form a distinct branch of the glossopharyngeal nerve (CN IX), the carotid sinus nerve.

Afferent signals from both the aortic and carotid sinus baroreceptors are sent to the nucleus tractus solitarius (NTS) and they create a negative feedback loop that keeps arterial blood pressure within normal limits.

By stimulating these arterial baroreceptors in response to vascular distension, elevated arterial blood pressure can be reflexively brought back to normal levels by decreasing peripheral sympathetic outflow and activating peripheral parasympathetic outflow.

Three subsequent functional processes must take place in order to convert a mechanical change in the arterial vasculature into an excitatory signal:

• Vascular distension and the deformation of arterial baroreceptor terminals

• Mechanoelectrical transduction converts the mechanical deformation of arterial baroreceptor nerve terminals into baroreceptor membrane depolarization. Mechanosensitive ion channels such as the epithelial Na+ channels (ENaCs), acid-sensing ion channels (ASICs) and transient receptor potential (TRP) ion channels are responsible for mechanoelectrical transduction in arterial baroreceptor nerve terminals.

• Translation of membrane depolarization into action potentials via voltage-gated ion channels (Na+, Ca2+ and K+).




Baroreceptor reflex Pathway
Stimulation of baroreceptors
Glossopharyngeal & Vagus nerves
⊕ Glutamate
Nucleus Tractus Solitarius (NTS)
⊕ Glutamate
⊕ Glutamate
Caudal venterolateral medulla (CVM)
Vagal motor neurons in Nucleus Ambiguus
⊖ GABA
Rostral ventrolateral medulla (RVLM)
Vagal inhibition of heart
Inhibition of tonic sympathetic discharge to vasculature
1. Decreased heart rate and strength of conduction
2. Vasodilation of arterioles and veins
3. Decreased blood pressure
©MARROW

Physiology INI-CET 2024

Question 1:

Which of the following are properties of ventricular cardiac muscle?

a) 1,2

b) 1,2,3

c) 1,2,4

d) 1,2,3,4

Question 2:

Which of the following may lead to a decrease in GFR?

a) Increase in renal blood flow

b) Decrease in glomerular oncotic pressure

c) Increase in glomerular hydrostatic pressure

d) Decrease in bowman capsule oncotic pressure

Question 3:

Which is the most potent naturally occurring estrogen?

a) Estradiol

b) Estriol

c) Estrone

d) Conjugated Estrogen

Question 4:

Which hormones are produced from the zones labelled in the image given?



a) A-cortisol, B-aldosterone, C-androgens, D-epinephrine

b) A-epinephrine, B-cortisol, C-aldosterone, D-androgens

c) A-aldosterone, B-epinephrine, C-androgens, D-cortisol

d) A-aldosterone, B-cortisol, C-androgens, D-epinephrine

Question 5:

Erythropoietin is produced by:

a) 1 only

b) 2 only

c) 1,2

d) 1,2,3,4

Question 6:

EOG, EEG, and EMG of a patient are given respectively from above downwards. Which stage of sleep does the marked area represent?



a) NREM I sleep

b) NREM II sleep

c) REM sleep

d) NREM III sleep

Question 7:

Which of the following statements is incorrect regarding ABO blood grouping?

a) Antibodies are naturally present since birth

b) The antigens are carbohydrate in nature

c) The antibodies are predominantly IgM

d) Antibodies are present only if antigens are absent

Question 8:

Identify the hormones marked in the image given



a) A-Somatostatin, B-GHRH, C-GH, D-IGF-1

b) A-GHRH, B-GH, C-IGF-1, D-Somatostatin

c) A-GHRH, B-Somatostatin, C-GH, D-IGF-1

d) A-GH, B-GHRH, C-IGF-1, D-Somatostatin

Question 9:

Identify the correct sequence.

a) Protein kinase A » Adenylyl cyclase » cAMP

b) cAMP » Adenylyl cyclase » Protein kinase A

c) Adenylyl cyclase » cAMP » Protein kinase A

d) Protein kinase A » cAMP » Adenylyl cyclase

Question 10:

Which among the following instruments is not used in the Sahli's method of hemoglobin estimation?



a) A

b) B

c) C

d) D

Question 11:

Which neurotransmitter is secreted by the SA node in response to increased blood pressure?

a) Acetylcholine

b) Epinephrine

c) Norepinephrine

d) Dopamine

Question 12:

Match the following hormones with their receptors.

a) A-1, B-2, C-3, D-4

b) A-1, B-3, C-2, D-4

c) A-3, B-4, C-1, D-2

d) A-4, B-2, C-3, D-1

Question 13:

The O2 saturation with respect to PO2 is shown by the curve marked as 5. What happens to the oxygen Hb dissociation curve on transfusion of stored RBCs?



a) 2

b) 3

c) 4

d) 1

Question 14:

Not seen in decompression sickness

a) Vision loss

b) Hearing loss

c) Hemodilution

d) Coagulopathy

Question 15:

Which of the following ovarian structures is responsible for the following finding?



a) Secondary follicle

b) Theca interna

c) Granulosa cell

d) Corpus luteum

Question 16:

Which of the following does not form the inner component of photoreceptors?

a) Mitochondria

b) Membranous disc

c) Golgi apparatus

d) Endoplasmic reticulum

Question 17:

Which of the following does not stimulate catecholamine synthesis?

a) Hyperglycemia

b) Smoking

c) Standing posture

d) Exercise

Question 18:

While collecting a blood sample for serum calcium, which is easily alterable, which of the following should be prevented to measure it accurately?

a) If a tourniquet is needed, tie and release as soon as the sample is collected.

b) Collecting after exercise.

c) Making a fist is not recommended.

d) Not collecting samples from patients immobilized for long periods of time

Question 19:

Which receptor contributes to static 2 point discrimination?

a) Merkel cell

b) Meissner corpuscle

c) Pacinian corpuscle

d) All of them

Question 20:

Mitochondrial sheaths are found in which part of the sperm?

a) Tail

b) Middle piece

c) Principal piece

d) Connecting piece

Question 21:

Identify the correct option



a) 1-Z line, 2-A band, 3-H zone, 4-I band

b) 1-A band, 2-Z line, 3-H zone, 4-I band

c) 1-T tubule, 2-I band, 3-H zone, 4-A band

d) 1-T tubule, 2- A band, 3-I band, 4-H zone

Question 22:

Which of these drug transport systems do not require ATP?



a) 1,2,3

b) 2,3,4

c) 3,4,5

d) 1,2,5

Question 23:

Choose the correct options from the given statements of the enteric nervous system-

a) 1,2,3

b) 1,4

c) 1,3

d) 2

Question 24:

What causes decrease in GFR?

a) Decrease in glomerular coefficient

b) Decrease in glomerular oncotic pressure

c) Increase in glomerular hydrostatic pressure

d) Decrease of Bowman's hydrostatic pressure

Question 25:

A 70-year-old man with a BP of 170/100mmHg. What is the reason for the increased pulse pressure in this patient?

a) Decreased arterial compliance

b) Increased cardiac output

c) Increased sympathetic outflow

d) Increased myocardial contractility

Question 26:

Physiological dead space :

a) Can be measured by spirometry

b) Is lesser than anatomical dead space

c) Represents inadequately ventilated alveoli

d) Represents inadequately perfused alveoli

Question 27:

What contributes to the latent period of muscle twitch?

a) 1,2,4

b) 1,2,3

c) 3,4

d) 2,3

Question 28:

Mechanical nociceptors and their response to a probe with a blunt object, pinprick, and toothed forceps with serrations are recorded as an output of a force transducer coupled to the stimulator. What could be inferred from this?



a) There are no responses from the nociceptors when stimulated by blunt forceps

b) Pinching with toothed forceps causes stimulation of mechano-nociceptors, but no pain

c) Pricking with a needle does not stimulate nociceptors

d) Sharp pain sensation is carried by C fibres

Question 29:

Peripheral chemoreceptors are most sensitive to?

a) Hypercapnia

b) Hypoxia

c) Acidosis

d) Alkalosis

Question 30:

Identify the marked structure.

a) Intercalated disc

b) Connective tissue

c) Cytoskeletal condensation

d) Nerve fibre

Question 31:

Which blood component is not separated by centrifugation in the blood bank?

a) Red blood cells

b) Platelets

c) Cryoprecipitate

d) Albumin

Answer Key

Question No.	Correct Option
1	b
2	d
3	a
4	d
5	c
6	c
7	a
8	a
9	c
10	d
11	a
12	c
13	a
14	c
15	d
16	b
17	a
18	b
19	a
20	b
21	a
22	a
23	d
24	a
25	a
26	d
27	a
28	a
29	b
30	a
31	d

Detailed Explanations

Solution to Question 1:

All or none phenomenon, length tension relationship and long refractory period applies to cardiac muscles. Cardiac muscle cannot be tetanized due to long refractory period.

In cardiac myocytes, depolarization proceeds rapidly and an overshoot of the zero potential is present. This is followed by a plateau before the membrane potential reaches the RMP (-90mV). This plateau phase lasts for 200 msec or more. Repolarization is therefore not complete until contraction is 50% completed.

Phases of cardiac action potential

• 0→ Depolarisation and overshoot-Due to opening of voltage-gated Na+ channels

• 1→ Initial rapid repolarization-Due to closure of Na+ channels and opening up of K+ channels

• 2→ Plateau phase - Due to slow but prolonged opening of Ca2+ channels

• 3→ Late rapid repolarization

• 4→ Baseline (RMP)

Phases 3 and 4 are due closure of Ca2+ channels and a slow, delayed increase of K+ efflux through K+ channels.

During phases 0 to 2 and about half of phase 3 of cardiac action potential, cardiac muscle cannot be excited again, that is, it is in its absolute refractory period. It remains relatively refractory until phase 4. Thus, cardiac muscle cannot be tetanized.



Solution to Question 2:

A decrease in bowman's capsule oncotic pressure would lead to a decrease in GFR.

Glomerular filtration rate (GFR) is the volume of fluid filtered from the renal glomerular capillaries into the Bowman's capsule per unit time.

The GFR is determined by sum of the hydrostatic and colloid osmotic forces across the glomerular membrane, which gives the net filtration pressure. It is dependant on Glomerular Kf (glomerular ultrafiltration coefficient) also.

Mathematically, it can be expressed as follows:

GFR= Kf [PG + πB -PB- πG]

Forces favoring Filtration

• PG: Glomerular hydrostatic pressure

• πB: Bowman's capsule oncotic pressure

Forces opposing Filtration

• PB: Bowman's capsule hydrostatic pressure

• πG: Glomerular capillary oncotic pressure

Factors that can increase the GFR are as follows:

• Increase in glomerular capillary filtration coefficient Kf: This coefficient depends on the hydraulic conductivity and surface area of the glomerular capillaries.

• Decrease in Bowman's capsule hydrostatic pressure: Conditions with urinary tract obstruction can lead to an increase in this pressure and a fall in GFR.

• Decrease in glomerular colloid osmotic pressure: Accumulation of plasma proteins that are not filtered from the blood contributes to this pressure and an increase reduces the further filtration and GFR. It is directly proportional to the following factors:Arterial plasma colloid osmotic pressure

• Filtration fraction

Solution to Question 3:

Estradiol is the most potent naturally occurring estrogen.

Classic estrogens are C18 steroid compounds containing a phenolic ring. Only three estrogens are present in significant quantities in the plasma of the human female—β-estradiol, estrone, and estriol.

The principal estrogen secreted by the ovaries is β-estradiol.

Small amounts of estrone are also secreted, but most of this is formed in peripheral tissues from androgens secreted by the adrenal cortices and by ovarian thecal cells.

Estriol is a weak estrogen; it is an oxidative product derived from both estradiol and estrone, with the conversion occurring mainly in the liver.

The estrogenic potency of β-estradiol is 12 times that of estrone and 80 times that of estriol. Hence, estradiol is the most potent natural form of estrogen.

In the liver, estradiol, estrone, and estriol are converted to inactive glucuronide and sulfate conjugates. All these compounds, along with other metabolites, are excreted in the urine.



Solution to Question 4:

Option D is the correct option. This is a section through an adrenal gland showing both the medulla and the zones of the cortex. Following are the zones of the cortex and medulla, and the hormones they secrete.

A – Zona Glomerulosa – secretes mineralocorticoids (aldosterone)

B – Zona Fasciculata – secretes glucocorticoids (cortisol, corticosterone)

C – Zona Reticularis – secretes androgens (dehydroepiandrosterone)

D – Adrenal Medulla – Produces Catecholamines: Norepinephrine, Epinephrine, and Dopamine

On histological examination, the different layers are identified as:

• Outer zona glomerulosa - small, compact cells

• Central zona fasciculata - larger lipid-rich cells arranged in radial columns

• Inner zona reticularis - pigmented, compact cells

• Adrenal medulla - thin layer of large chromaffin cells

Solution to Question 5:

Erythropoietin is produced mainly by kidney and liver.

In adults, approximately 85% of erythropoietin is produced by the kidneys, while 15% is synthesized by the liver. Both of these organs possess the mRNA necessary for erythropoietin production. Although erythropoietin can be found in the spleen and salivary glands, these tissues

lack the mRNA required for its synthesis, indicating that they do not produce the hormone.

Erythropoietin is secreted by interstitial cells in the peri-tubular capillary bed of the kidneys and by perivenous hepatocytes in the liver. It is also generated in the brain, where it provides a safeguarding effect against excitotoxic harm caused by hypoxia; and in the uterus and oviducts, where it is stimulated by estrogen and seems to facilitate estrogen-driven angiogenesis.

Solution to Question 6:

REM sleep is represented by the marked area in the given polysomnography.

The three zones in the given image represent EOG, EEG, and EMG, respectively. In the marked area,

• EOG shows Positive activity

• EEG shows Mixed waves

• EMG shows Minimal activity

This is characteristic of REM sleep.





Solution to Question 7:

Antibodies are not present since birth. At 2 to 8 months after birth, an infant begins to produce agglutinins (antibodies).

In the absence of type A agglutinogen(antigen) in an individual's RBCs, antibodies called anti-A agglutinins form in the plasma. Similarly, when type B agglutinogen is lacking in the RBCs, antibodies referred to as anti-B agglutinins develop in the plasma. Type A blood carries type A agglutinogens and anti-B agglutinins, while type B blood contains type B agglutinogens and anti-A agglutinins. Type AB blood possesses both A and B agglutinogens but no agglutinins. On the other hand, type O blood, which lacks agglutinogens, does have both anti-A and anti-B agglutinins.

The A and B antigens are complex oligosaccharides that differ in their terminal sugar. The agglutinins are gamma globulins produced by the bone marrow and lymph gland cells. Most of them are IgM and IgG immunoglobulin molecules.

Solution to Question 8:

In the given image, A: Somatostatin, B: GHRH, C: GH, D: IGF-1

• Growth hormone-releasing hormone (GHRH) secreted from the hypothalamus acts on the anterior pituitary to release growth hormone.

• Growth hormone (GH) increases the circulating levels of IGF -1 which exerts its effect on the tissues.

• Ghrelin promotes the secretion of growth hormone from the anterior pituitary.

• Somatostatin secreted from the hypothalamus is an inhibitory effect on the GH secretion from the anterior pituitary.

• Similarly, IGF -1 exerts negative feedback on GH secretion and positive feedback on somatostatin secretion, which in turn inhibits GH secretion.

Solution to Question 9:

The correct sequence in Gs is adenylyl cyclase » cAMP » Protein kinase A.

G protein-coupled receptors (GPCRs) are cell membrane receptors that span across the membrane 7 times. They also serve as receptors for various peptide hormones, neurotransmitters and amino acids. Subtypes of GPCRs are distinguished by their alpha subunits. The important subtypes of GPCRs are Gs, Gi, and Gq.

When hormones bind to the receptor, it enables the receptor to connect with a G protein. If the G protein activates the adenylyl cyclase-cAMP system, it is referred to as a Gs protein, indicating a stimulatory G protein. The Gs protein stimulates adenylyl cyclase, an enzyme located in the cell membrane, which then converts a small amount of cytoplasmic adenosine triphosphate into cAMP. This cAMP activates cAMP-dependent protein kinase, which phosphorylates specific proteins within the cell, initiating biochemical reactions that ultimately result in the cell's response to the hormone.

Solution to Question 10:

Option D is RBC pipette, it is not used in Sahli's method of hemoglobin estimation. The instruments used are comparator with brown glass standard(Option A), graduated hemoglobin tube(Option B), Sahli's pipette (Option C)and stirrer.

Sahli's method of estimation of hemoglobin is based on colorimetric measurement of the intensity of color developed on the addition of dilute hydrochloric acid (N/10 HCl) reagent to the blood.

In Sahli's method, hemoglobin is converted to acid hematin with the action of dilute hydrochloric acid (N/10 HCl). The brown color of the acid hematin is matched to its intensity with a standard brown glass comparator in a visual colorimeter called Sahli's colorimeter. It can be used to estimate oxyHb and reduced Hb.

Sahli's method is a simple bedside test and is cost-effective. However, it has some disadvantages, which include visual errors, the inability to measure other forms of hemoglobin like carboxyhemoglobin, methemoglobin, and sulfhemoglobin, fading of the comparator over time, and the color of acid hematin fading quickly.

The visual error can be reduced by taking the average of three readings; first, when the color is slightly darker than the standard; second, when the color exactly matches the standard; and third, when the color is slightly lighter than the standard.

Solution to Question 11:

Acetylcholine is released in response to increased blood pressure.

In response to raised BP, there is inhibition of sympathetic system and activation of parasympathetic system. Acetylcholine is the universal neurotransmitter of PNS.

Baroreceptors are stretch receptors located in the heart and the adventitia of the blood vessels. They are stimulated by the stretching of the walls of the blood vessels when there is an increased volume or blood pressure in the vessel. They are important in the immediate management of blood pressure. They are of two types: carotid sinus and aortic arch receptors (high-pressure regions) and cardiopulmonary receptors (low-pressure regions).

An increase in blood pressure stimulates the baroreceptors. These signals are transmitted through small Hering's nerves to the glossopharyngeal nerves and then to the nucleus tractus solitarius (NTS) in the medulla. These signals release glutamate and activate the caudal ventrolateral medulla (CVLM). This in turn inhibits the rostral ventrolateral medulla (RVLM) considered the vasoconstrictor center for the brain and the heart.

Simultaneously, this pathway also excites the vagal parasympathetic center. The stimulation of the vagus results in a slowing of impulse generation in the SA node and a slowing of conduction velocity in the AV node.

These neural changes produce vasodilation, venodilation, hypotension, bradycardia, and a decrease in cardiac output.



- NTS: Nucleus Tractus Solitarius
- CVLM: Caudal Ventrolateral Medulla
- RVLM : Rostral Ventrolateral Medulla (Vasomotor center)
- CVC : Cardiovagal center (Cardioinhibitory center)
- IML: intermediolateral nucleus

Solution to Question 12:

The correct answer is A-3, B-4, C-1, D-2.

Insulin binds to a receptor that has intrinsic tyrosine kinase activity. Upon activation, the receptor phosphorylates itself and downstream signaling proteins. (A-3)

Acetylcholine acts on nicotinic receptors, which are ligand-gated ion channels. Binding of acetylcholine opens the channel, allowing ion flow (e.g., Na■ or K■). (B-4)

Growth hormone binds to its receptor, which lacks intrinsic kinase activity but recruits Janus Kinase (JAK) to phosphorylate downstream proteins and activate the STAT (Signal Transducer and Activator of Transcription) pathway. (C-1)

Thyroxine (and its active form, T3) binds to intracellular nuclear receptors. These hormone-receptor complexes act as transcription factors, regulating gene expression directly in the nucleus. (D-2)

Hormone	Receptor
A.Insulin	3. Tyrosine Kinase
B.Acetylcholine	4. Ligand gated

Hormone	Receptor
C.Growth Hormone	1. JAK/STAT
D.Thyroxine	2. Nuclear Receptor

Solution to Question 13:

When stored RBCs are transfused, the oxygen-hemoglobin dissociation curve typically shifts to the left but still retains its sigmoid shape as represented by the curve marked as 2.



This is due to reduced levels of 2,3-bisphosphoglycerate (2,3-BPG) in stored red blood cells. 2,3-BPG is a metabolite in RBCs that binds to hemoglobin and reduces its affinity for oxygen.

During storage, metabolic processes in RBCs slow down, leading to a depletion of 2,3-BPG. Lower 2,3-BPG levels make hemoglobin hold onto oxygen more tightly, impairing oxygen delivery to tissues. Hence, this increases hemoglobin's affinity for oxygen, shifting the curve to the left.

Other Options:

Option B and C: show a right shift to the oxygen-hemoglobin dissociation curve.↑ in 2,3-BPG seen in increased anaerobic glycolysis leads to the right shift of the curve. Storage of blood causes depletion of 2,3-BPG, hence leading to left shift instead of right.

Solution to Question 14:

Hemodilution is not seen in decompression sickness. Instead, hemoconcentration is seen in decompression sickness.

Hemodilution refers to the dilution of blood, usually due to excessive fluid intake or other medical conditions, but it is not a feature of decompression sickness.

Decompression sickness (DCS) is a condition that occurs when a person ascends too quickly after being exposed to high-pressure environments (such as deep-sea diving), causing nitrogen bubbles to form in the bloodstream and tissues.

Other options

Option A: Vision changes or loss can occur in DCS due to gas bubbles affecting the optic nerve or retinal circulation, leading to temporary or permanent vision impairment.

Option B: Gas bubbles in DCS can also affect the auditory system, leading to hearing loss or tinnitus, a common feature of the condition.

Option D: Decompression sickness can lead to coagulopathy, which refers to abnormal blood clotting due to the formation of nitrogen bubbles that disrupt the vascular system, potentially causing bleeding or clotting disorders.

Solution to Question 15:

The ovarian structure responsible for the following finding is the corpus luteum.

The histological image shows a corkscrew appearance of the glands, and this is characteristic of the secretory phase of the endometrium in the menstrual cycle. This phase is stimulated by progesterone, which is secreted by the corpus luteum.

Endometrial Biopsy of the various stages of the ovarian cycle:

1. Proliferative phase: The glands are simple, straight, and tubular glands. The lining of the glands shows pseudostratified columnar epithelium with mitotic figures, indicating active cell proliferation. The stroma is dense and compact, with numerous cells undergoing mitosis.



2. Early secretory phase: The initially straight glands begin to coil as the endometrium transitions into the secretory phase. Subnuclear Vacuolation is seen in the epithelial cells of the endometrial

glands, caused by the accumulation of glycogen at the base of the cells.

3. Late secretory phase: The glands become maximally tortuous and assume a distinctive "saw-tooth" appearance due to extreme coiling. They are filled with secretions rich in glycogen and glycoproteins, essential for supporting a potential embryo. The stroma is edematous.



Other Options:

Option A: Secondary follicle

The secondary follicle is a more advanced stage of follicular development following the primary follicle. It contains a fluid-filled cavity (antrum) surrounded by two layers: theca interna and theca externa.



Option B and C: Theca interna and Granulosa cell

The theca interna is a layer of stromal cells just outside the basal lamina of the granulosa cells. They are composed of cuboidal endocrine cells with lipid droplets and abundant smooth

endoplasmic reticulum, reflecting its role in steroidogenesis.

Granulosa cells line the interior of the follicle. Cells are organized in layers, initially single-layered in primary follicles and multilayered in secondary follicles.



Solution to Question 16:

Among the given options, the membranous disc does not form the inner component of photoreceptors but instead forms the outer component.

The other three options i.e., mitochondria, Golgi apparatus, and endoplasmic reticulum form the inner component of photoreceptors.

Photoreceptors (rods and cones) are specialized neurons in the retina with distinct structural components:

• Outer segment: Contains membranous discs that have photopigments (like rhodopsin in rods and photopsins in cones) essential for light detection.

• Inner segment: Contains cellular organelles responsible for metabolism and protein synthesis, such as mitochondria, Golgi apparatus, and endoplasmic reticulum.



Solution to Question 17:

Hyperglycemia does not stimulate catecholamine synthesis. Instead, hypoglycemia stimulates catecholamine synthesis.

Catecholamine synthesis is primarily stimulated by various physiological stressors and factors that activate the sympathetic nervous system, such as physical exertion(exercise), changes in posture, or certain external stimuli like smoking.

Other options:

Option B: Smoking stimulates catecholamine release through the activation of the sympathetic nervous system, which increases catecholamines like adrenaline and norepinephrine.

Option C: When a person stands up, it can lead to orthostatic hypotension, which triggers the release of catecholamines to help increase blood pressure and maintain adequate perfusion.

Option D: Exercise stimulates the sympathetic nervous system and increases catecholamine production to support increased heart rate, blood pressure, and metabolic demands.

Solution to Question 18:

Blood collection after exercise should be prevented to measure serum calcium accurately.

Option A: A prolonged tourniquet is usually avoided while drawing blood for calcium measurements since hemoconcentration also leads to an increase in serum calcium levels. Hence, the tourniquet should be released as soon as the sample is collected.

Option C: Fist-clenching or forearm exercise can lead to falsely elevated ionized (free) calcium levels. Hence, making a fist while blood collection is not recommended.

Option D: Staying immobilised for long periods of time increases bone resorption and increases calcium levels. Hence, do not collect blood samples for serum calcium estimation in patients who are immobilised for long periods.

Solution to Question 19:

Merkel cell contributes to static 2 point discrimination.

Merkel cells are responsible for static two-point discrimination. These cells are specialized for detecting pressure and texture and are involved in the detection of fine touch. They provide information about the spatial resolution of tactile stimuli, which is essential for static two-point discrimination, where the ability to differentiate between two distinct points of touch is tested.

Merkel cells are expanded dendritic endings in the epidermis of glabrous (non-hairy) skin. They respond to sustained pressure and touch. They are slowly adapting receptors.

Static vs. Dynamic Two-Point Discrimination

Static: Measures the ability to distinguish stationary points; mediated by Merkel cells (slow-adapting receptors).

Dynamic: Measures the ability to distinguish moving points; mediated by Meissner corpuscles (fast-adapting receptors).



Other options:

Option B: Meissner corpuscle: Meissner corpuscles are dendrites encapsulated in the connective tissue beneath the epidermis of glabrous (non-hairy) skin. They respond to tap and flutter (low-frequency vibration - 5-40Hz). They are rapidly adapting receptors.

Option C: Pacinian corpuscle: Pacinian corpuscles are the largest cutaneous mechanoreceptor. They are present in the dermis of glabrous (non-hairy) and hairy skin. A Pacinian corpuscle consists of a nerve ending encapsulated by concentric layers of connective tissue, giving it an onion-like appearance. They respond to fast vibration (60-500Hz) and deep pressure. They are

rapidly adapting receptors.

Solution to Question 20:

Mitochondrial sheaths are found in the middle piece of the sperm.

The middle piece is the segment of the sperm located just behind the head and neck.

It contains a spiral arrangement of mitochondria, forming the mitochondrial sheath around the central axoneme.

These mitochondria generate ATP, which provides the energy required for the flagellar movement of the sperm, enabling motility.



Other options:

Option A: Tail: Comprises the middle piece, principal piece, and end piece and is primarily responsible for motility.

Option C: Principle piece: Longest part of the tail which contains fibrous sheaths around the axoneme but lacks mitochondria.

Option D: Connecting piece: Links the head of the sperm to the tail (neck region). Does not contain mitochondrial sheaths.

Solution to Question 21:

The correct marking is 1-Z line, 2-A band, 3-H zone, 4-I band.



H zone or H band denotes the less refractile region in the centre of A band.

• A band – Anisotropic / Dark band: Formed by thick filaments - Myosin

• I band – Isotropic / Light band: Formed by thin filaments – Actin, Troponin, Tropomyosin

• M line – Seen in the centre of H band: Due to central bulge of A band

• Z line – Seen in the centre of each I band: Narrow line with high refractile material appearing dark.



Solution to Question 22:

Paracellular transport, Simple Diffusion and Facilitated diffusion do not require ATP as they are all passive transport mechanisms.

1: Simple diffusion : Simple diffusion is a passive transport mechanism where molecules (water, carbon dioxide, nitrogen) move from an area of high concentration to an area of low concentration and is not carrier mediated.

2: Paracellular : Paracellular transport involves movement of molecules through tight junctions between cells without the need for energy (ATP).

3: Facilitated diffusion : This also occurs along a concentration gradient, but involves carrier proteins to help move the drug across the membrane. No ATP is used because the process is still passive (driven by the concentration gradient).

4: Antiport (Na/K ATPase): Antiport transport system actively moves ions (Na+, K+) against their concentration gradient using ATP in the opposite direction. It is a form of primary active transport.

5: Symport : Two substances are transported in the same direction across a membrane. ATP is required to move substances against their gradients, often via a secondary active transport mechanism.

Solution to Question 23:

The enteric nervous system functions independently of the brain and spinal Cord. (Statement 2)

Statement 2: Functions independent of brain and spinal cord: The enteric nervous system is often referred to as the "second brain" because it can function independently of the central nervous system (CNS). It has its own reflex circuits and can control gut motility, secretion, and blood flow without input from the brain or spinal cord, although it can be influenced by them.

Statement 1: The vagus is the only afferent from the gut: While the vagus nerve does provide significant afferent (sensory) input from the gut, other nerves, such as those from the pelvic splanchnic nerves, also contribute to afferent pathways. Therefore, the vagus is not the only afferent nerve.

Statement 3: ■Myenteric plexus in the lamina propria: The myenteric plexus is located between the circular and longitudinal muscle layers of the gut, not in the lamina propria (which is part of the mucosal layer of the gut).

Statement 4: Exclusively under the influence of nerves: The enteric nervous system is influenced by nerves, but it also responds to hormonal and immune system signals. Therefore, it is not exclusively under neural influence.

Statement 5: Only parasympathetic connects enteric to CNS: Both parasympathetic and sympathetic connect the enteric nervous system to the central nervous system. Not just the parasympathetic.

The enteric nervous system consists of submucosal and myenteric plexi. This Plexi can function independently and is often referred to as the ‘little brain’. The enteric nervous system regulates gastrointestinal (GI) function and is connected to the central nervous system by parasympathetic and sympathetic fibers.

The myenteric (Auerbach’s) plexus is present between the longitudinal and circular muscle layers. It innervates both these layers along the entire length of the GI tract. It primarily controls GI motility. On stimulation, it has the following effects:

• Increased tone of the GI wall

• Increased intensity and rate of rhythmic contractions

• Increased velocity of excitatory wave conduction

Some of the myenteric plexus neurons primarily release vasoactive inhibitory peptide (VIP), which causes the relaxation of the GI sphincters.

The submucosal (Meissner's) plexus is found between the mucosa and the circular muscle layers. It innervates the GI glands, endocrine cells, and submucosal blood vessels. It primarily controls GI secretions, absorption, and local blood flow.

The parasympathetic innervation of the plexi is from the vagus and pelvic nerves. They release either acetylcholine or substance P. It generally increases GI motility and secretion.

The sympathetic innervation is from the celiac, superior mesenteric, inferior mesenteric, and hypogastric sympathetic ganglia. It generally inhibits GI motility and stimulates sphincteric contraction.

Solution to Question 24:

Decrease in glomerular coefficient causes decrease in GFR.

The glomerular filtration coefficient (Kf) is a measure of how efficiently the glomerulus filters blood into the kidney tubules. It depends on two key factors:

• Surface Area for Filtration: Greater surface area means higher filtration.

• Permeability of the Glomerular Membrane: Higher permeability leads to greater filtration.

Glomerular filtration rate (GFR) is the volume of fluid filtered from the renal glomerular capillaries into the Bowman's capsule per unit time.

The GFR is determined by the sum of the hydrostatic and colloid osmotic forces across the glomerular membrane, which gives the net filtration pressure. It is dependent on Glomerular Kf (glomerular ultrafiltration coefficient) also.

Mathematically, it can be expressed as follows:

$GFR = Kf [PG + \pi B - PB - \pi G]$

Forces favoring Filtration

• PG: Glomerular hydrostatic pressure (Option C)

• πB: Bowman's capsule oncotic pressure

Forces opposing Filtration

• PB: Bowman's capsule hydrostatic pressure (Option D)

• πG: Glomerular capillary oncotic pressure (Option B)

The formula for glomerular filtration rate (GFR) incorporates Kf:

• A high Kf means efficient filtration (e.g., in healthy kidneys)-indicates high GFR

• A low Kf {occurs when the surface area is reduced (e.g., due to glomerular damage) or the membrane becomes less permeable (e.g., in diabetic nephropathy)}- indicates low GFR.

Kf is essential for maintaining proper kidney function, as it directly affects how well the kidneys filter blood to form urine.

Solution to Question 25:

Pulse pressure increases with age due to decreased arterial wall compliance.

Diastolic blood pressure falls after 60 years of age due to increased arterial stiffness and decreased compliance, often associated with arteriosclerosis. As the left ventricle contracts against stiffer, less compliant arteries, systolic blood pressure also increases.

As a result of these two changes, pulse pressure increases (↑PP = ↑SBP - ↓DBP)

The factors that affect the pulse pressure:

• the stroke volume output of the heart

• the compliance or total distensibility of the arterial walls

• the character of ejection from the heart during systole (less important)

Pulse pressure = Stroke volume/ arterial compliance

Widened pulse pressure (> 40 mm Hg) occurs with several diseases such as:

• Aortic regurgitation

• Aortic sclerosis

• Severe iron deficiency anaemia

• Arteriosclerosis

• Hyperthyroidism

In the majority of these cases, systolic pressures increase while diastolic pressures remain near normal.

Solution to Question 26:

Physiological dead space represents inadequately perfused alveoli.

Physiological dead space is the part of the respiratory system where no gas exchange occurs. It includes:

• Anatomic dead space: Air in the conducting airways (nose, trachea, bronchi) that doesn't reach the alveoli.

• Alveolar dead space: Air in alveoli that are ventilated but not perfused by blood.

Together, it represents the total volume of air that doesn't participate in oxygen and carbon dioxide exchange.(Option B)

Anatomical dead space refers to the volume of all the space of the respiratory system other than the alveoli that do not participate in gas exchange. The anatomical dead space is not altered with a change in ventilation.

In healthy individuals:

• Anatomical dead space = 150 mL.

• Alveolar dead space is usually zero (increased in diseased conditions).

Alveolar Dead space represents inadequately perfused alveoli (wasted ventilation). (Option C)

In normal individuals, physiological dead space remains unaltered and is equal to the anatomical dead space (volume of all the space of the respiratory system other than the alveoli that do not participate in gas exchange).

Dead space cannot be measured using spirometry. (Option A)

Solution to Question 27:

Neuromuscular transmission, Excitation contraction coupling and transmission impulse from nerve coupling contribute to latent period of muscle twitch.

The latent period of a muscle twitch is the brief delay between the application of a stimulus (such as a nerve impulse) and the onset of muscle contraction. During this period, several physiological processes occur, but neuromuscular transmission and excitation-contraction coupling primarily contribute to this delay.

Neuromuscular transmission: The action potential travels down the motor neuron to the neuromuscular junction. Acetylcholine (ACh) is released into the synaptic cleft and binds to receptors on the muscle fiber's sarcolemma, triggering an action potential in the muscle fiber. (Statement 1)

Excitation-contraction coupling: The action potential propagates along the sarcolemma and into the transverse (T) tubules. This triggers the release of calcium ions from the sarcoplasmic reticulum into the cytosol. Calcium binds to troponin, initiating the interaction of actin and myosin filaments. (Statement 2)

Reuptake of calcium back into the sarcoplasmic reticulum: This occurs during the relaxation phase, not the latent period. (Statement 3)

Transmission of impulse from nerve: This is a part of neuromuscular transmission and is involved in the latent period. (Statement 4)

Solution to Question 28:

There are no responses from the nociceptors when stimulated by blunt forceps.

Mechanical nociceptors typically respond to noxious mechanical stimuli, such as sharp pressure or intense stretch. A blunt object, such as blunt forceps, may not provide a strong enough stimulus to activate these nociceptors, leading to no response.

The sharp painful sensation felt immediately upon injury is known as fast pain. The poorly localized dull aching pain is characteristic of slow pain. The nerve fibers carrying fast pain and slow pain are Aδ and C respectively.

Nociceptors are innervated by both Aδ and C fibers. Aδ fibers are myelinated and larger in diameter with a faster conduction velocity. C fibers are unmyelinated and conduct impulses at a slower rate. When nociceptors are activated by a noxious stimulus, activated Aδ fibers release glutamate and mediate fast pain. It is a rapid response that is responsible for localizing the noxious stimulus.

Other Options:

Option B: Pinching with toothed forceps causes stimulation of mechanonociceptors, but no pain: Pinching with toothed forceps typically applies more localized pressure, which can activate mechanical nociceptors. However, it may still cause a painful sensation, not just a mechanical response. It is more likely that pain is perceived, even if the response is primarily mechanical.

Option C: Pricking with a needle does not stimulate nociceptors: A needle prick is a sharp stimulus that activates nociceptors, particularly mechanical nociceptors in response to tissue damage.

Option D: Sharp pain sensation is carried by C fibers: Sharp pain (due to a needle prick) is typically carried by A-delta fibers, which transmit fast, localized pain sensations. C fibers are responsible for carrying dull, burning pain that occurs later after the initial sharp pain.

Solution to Question 29:

Peripheral chemoreceptors are most sensitive to hypoxia.

Peripheral Chemoreceptors are located in the carotid bodies (at the bifurcation of the common carotid arteries) and aortic bodies (near the aortic arch). They are sensitive to changes in pO2, pCO2, and pH of the blood. In response to changes in these parameters, they send impulses to the medullary respiratory centers and modify the duration and frequency of respiratory cycles.

Peripheral chemoreceptors are most sensitive to a drop in partial pressure of oxygen (PaO■) in arterial blood.

Mechanism:

Type I glomus cells in the carotid and aortic bodies have O2-sensitive K+ channels, which are inactivated in response to hypoxia. This reduces the K+ efflux, depolarizing the cell and causing Ca2+ in■ux, via L-type voltage-gated calcium channels. The Ca2+ influx triggers action potentials and consequent excitation of the afferent nerve endings in the peripheral chemoreceptors.

Hypoxia (PaO■ < 60 mmHg) strongly stimulates these receptors, leading to an increase in respiratory rate and depth to improve oxygenation.

Other Options:

Peripheral chemoreceptors also respond to hypercapnia (Option A) and acidosis (Option C). However, these are primarily detected by central chemoreceptors, located in the medulla, which monitor CO■ levels in cerebrospinal fluid.

Alkalosis (Option D) causes a weaker response from peripheral chemoreceptors compared to hypoxia.

Solution to Question 30:

The given section shows cardiac muscle and the marked structure is an intercalated disc.

Intercalated discs are specialized structures unique to cardiac muscle tissue. They are critical for the coordinated contraction of cardiac muscle cells. They are found at the junctions between adjacent cardiac muscle cells. They appear as dark, zigzagging lines under a microscope. They facilitate the rhythmic and coordinated contraction of the heart by ensuring efficient electrical and mechanical connections.



On microscopy, cardiac muscle can be identified by:

• Presence of striations similar to the skeletal muscle.

• Presence of single and centrally placed nucleus. A perinuclear halo may be seen surrounding the nucleus.

• The junctions between adjoining cardiac myocytes are seen as dark staining transverse lines running across the muscle fibre. These lines are called intercalated discs.

• Cells within one fibre branch and join the cells in adjacent fibres.

Other Options:

Option B: Connective Tissue: Found surrounding the cardiac muscle cells, providing structural support, but not at the cell junctions.

Option C: Cytoskeletal condensation: Refers to cytoskeletal organization within a cell and is not visible as distinct structures in cardiac muscle histology.

Option D: Nerve Fibre: Present in cardiac tissue to innervate the heart but appears as separate structures and not within cardiac muscle fibers.

Solution to Question 31:

Albumin is not separated by centrifugation in the blood bank.

In blood banks, centrifugation is used to separate the main components of blood into distinct layers for storage and transfusion purposes. These components include:

• Red Blood Cells (RBCs): Separated by centrifugation and stored for patients with anemia or significant blood loss. (Option A)

• Platelets: Isolated through centrifugation of plasma-rich blood for use in patients with bleeding disorders. (Option B)

• Cryoprecipitate: Derived from plasma by controlled freezing and thawing, which allows the collection of clotting factors (e.g., Factor VIII, fibrinogen, von Willebrand factor). (Option C)

Albumin is a plasma protein that is extracted through a process called fractionation, not simple centrifugation. This involves the use of chemical, temperature, and pH adjustments to isolate albumin from plasma.



Physiology NEET 2024

Question 1:

What are the effects of posterior lesions in a Brodmann's area 22 in a stroke patient?

a) Auditory function

b) Receptive Speech Function

c) Expressive Speech Function

d) Visual functions

Question 2:

A 30-year-old female went on vacation and fell asleep while sitting on a beach. After some time she woke with sunburn. Later when she went back home, she experienced pain while taking a warm bath (40°C). Which of the following receptors is responsible for the pain?

a) Innocuos thermal receptor - allodynia

b) Innocuos thermal receptor - hyperalgesia

c) Thermal receptor - allodynia

d) Thermal receptor - hyperalgesia

Question 3:

To prevent polyspermy, which reaction occurs in the zona pellucida of the oocyte after fertilization?

a) Acrosomal reaction

b) Increased membrane potential of the oocyte

c) Cortical reaction

d) Decreased membrane potential of the oocyte

Question 4:

A 60-year-old patient came with a puncture wound on the left chest wall. Radiological investigations confirmed pneumothorax. Which of the following is correct regarding this

condition?

a) Lung volume increases, thoracic volume increases

b) Lung volume decreases, thoracic volume increases

c) Lung volume increases, thoracic volume decreases

d) lung volume decreases, thoracic volume decreases

Question 5:

Sour taste is mediated by which of the following receptors?

a) T1R1 GPCR

b) TRPV1 channel

c) T1R3 GPCR

d) Metabolic receptor

Question 6:

A patient who underwent right limb amputation 2 years back, now presents with severe pain in the limb (Phantom limb pain). On radiological investigations, What is the finding seen?

a) Projection of adjacent fibres to overlap the right somatosensory cortex

b) Expansion of left somatosensory cortex

c) Projection of adjacent fibres to overlap the left somatosensory cortex

d) Expansion of right somatosensory cortex

Question 7:

In a type 2 diabetes mellitus patient, serum insulin levels are found to be raised. He is advised to exercise to improve insulin resistance. Which of the following GLUT will help to improve insulin resistance after exercise ?

a) GLUT 1

b) GLUT 2

c) GLUT 5

d) GLUT 4

Question 8:

Which of the following is incorrect about iron absorption?

a) Ferrireductase convert Fe2+ to Fe 3+

b) RBC protophyrin is increased in deficiency

c) Iron excretion is only through skin, gut and urothelial cells

d) Body cannot compensate if more than 10 RBC/HPF per day is lost in urine

Question 9:

A 50-year-old patient presents with a chronic tingling sensation in the forearm. A neurological examination is done and appears to be normal. Which of the following receptors are correctly matched?

a) Merkel cells-slow vibration

b) Meissner's corpuscles-stretch

c) Pacinian corpuscles-fast vibration

d) Ruffini corpuscles-sustained pressure

Question 10:

The patient presented with alcoholic gait, and abnormal eye movements (nystagmus) after RTA. Which lobe of the cerebellum is likely affected?

a) Flocculonodular Lobe

b) Dentate nucleus

c) Anterior lobe

d) Interposed nucleus

Question 11:

Renal compensation in metabolic acidosis takes time because:

a) Initially the pH of tubular fluid increases

b) Low pH normally decreases bicarbonate excretion

c) It takes hours to days for kidney to synthesize enzymes that produce NH_4^+

d) Other ions in the tubule interfere with buffer action

Question 12:

The functioning of the thyroid involves the transport of iodine from the blood to the thyroid follicle. Iodine is transported from the follicle to the colloid. Which pump or channel is for transport of iodine from the follicle to the colloid?

a) Sodium iodine symporter

b) Calmodulin

c) Claudin

d) Pendrin

Question 13:

The resting membrane potential(RMP) of the nerve is -70mV. Increased permeability of which ion is responsible for the RMP?

a) Calcium

b) Sodium

c) Potassium

d) Chloride

Question 14:

Arrange the sequence of steps involved at neuromuscular junction endplate potential.

a) Acetylcholine release - Calcium entry- Na+ entry

b) Calcium entry- Acetylcholine release- Na+ entry

c) Na+ entry - Calcium entry - Acetylcholine release

d) Na+ entry - Acetylcholine release - Calcium entry

Answer Key

Question No.	Correct Option
1	b

2	a
3	c
4	b
5	b
6	c
7	d
8	a
9	c
10	a
11	c
12	d
13	c
14	b

Detailed Explanations

Solution to Question 1:

A posterior lesion in Brodmann's area 22, which is located in the superior temporal gyrus, typically affects receptive speech function. It is the sensory speech area.

The patient will most likely have Wernicke’s aphasia due to the damage to Wernicke’s area - area 22.

In Wernicke's aphasia, there is difficulty in finding words leading to poorly understood content for the listener. These patients also find it difficult to repeat sentences and name things properly. It is a type of fluent aphasia in which the speech is easy and fluent but there are difficulties related to the output of language like paraphasia (substitution of similar-sounding, non-words, or incorrect words). The speech is full of jargon and neologisms (non-existent words) that make little sense.

Other options:

Option A: Auditory Function is primarily mediated by Brodmann area 41, which is located in the primary auditory cortex within the superior temporal gyrus.

Option C: Expressive Speech Function is associated with Brodmann areas 44 and 45 (Broca's area), which are involved in language production and speech articulation. Lesions here can cause difficulty in producing coherent speech (Broca's aphasia).

Option D: Visual function primarily associated with Brodmann areas 17, 18, and 19 (the primary and secondary visual cortices).

Solution to Question 2:

The receptors responsible for this are innocuous thermal receptors which have perceived ordinarily non-noxious stimulus (i.e., warm bath) as severe pain also known as allodynia.

In this case, the warm bath at 40°C (typically not painful) causes pain due to sunburn, which sensitizes the skin. After sunburn, the skin becomes more sensitive, and thermal receptors that usually detect warmth now trigger pain signals in response to non-painful thermal stimuli, leading to allodynia.

Innocuous thermoreceptors, which detect non-harmful temperature changes(cold and warm), are specialized nerve endings found on the skin. Cold receptors are located on the dendritic endings of Aδ and C fibers, while warm receptors are situated on C fibers. The specific thermosensitive molecular channels mediating innocuous heat stimuli is transient receptor potential (TRP) cation channel TRPV1 and TRPM8 cation channel for innocuous cold stimuli.

Thermal hyperalgesia(increased response to noxious stimulation) is mediated by activation of NMDA receptors. After tissue or nerve injury, C-fibers discharges release glutamate and neurokinins from primary afferent terminals, leading to the release of peptides (e.g., substance P; calcitonin gene-related peptide, CGRP). As a result, a voltage-dependent magnesium block on NMDA receptors is removed allowing for an influx of calcium that triggers a cascade of intracellular events leading to the development and maintenance of hyperalgesia.(Option D)

Solution to Question 3:

To prevent polyspermy, the reaction in the zona pellucida of the oocyte after fertilisation is the cortical reaction.

The chronological steps of fertilisation:

Acrosomal reaction: After a sperm reaches the oocyte and binds to the zona pellucida, the acrosome reaction occurs. The acrosome, an enzyme-rich organelle in the anterior of the sperm head, fuses at many points with the plasma membrane of the oocyte. Then those fused membranes are sloughed off from the head, exposing the acrosomal enzymes. The enzymes digest a path through the zona pellucida, enabling the sperm to advance.

Eventually, the sperm meets and fuses with the secondary oocyte plasma membrane.

This contact results in weak membrane depolarisation(decreased membrane potential) and a calcium wave.

Cortical reaction: The calcium wave triggers the release of lysosomal enzymes from the cortical granules lining the oolemma. These enzymes hydrolyse the ZP3 receptor on the zona pellucida. Digestion of ZP3 and alteration of zona pellucida (zona reaction) prevents other sperms from binding, thus creating a block to polyspermy.

Solution to Question 4:

In pneumothorax, lung volume is decreased and thoracic volume is increased.

Both lungs and the chest wall (thoracic cage) are elastic structures. Under normal conditions, the elastic tendency of the lungs to collapse is exactly balanced by the elastic tendency of the chest

wall to expand.

In pneumothorax, air is introduced into the pleural space, so the pleural pressure, which is normally subatmospheric, becomes equal to atmospheric pressure. The chest wall springs outward and the lungs collapse. This increases the thoracic volume and decreases the lung volume.

Solution to Question 5:

Sour taste is mediated by the TRPV1 channel. The TRPV1 (Transient Receptor Potential Vanilloid 1) channel is involved in detecting the acidity of foods, which contributes to the perception of sour taste.

It is a cation channel that allows the influx of various cations, including sodium (Na^{+}), calcium (Ca^{2+}), and magnesium (Mg^{2+}), in response to changes in temperature and the presence of acidic conditions or capsaicin

Taste Receptors

• Humans have five established basic tastes: sweet, salt, sour, bitter and umami.

• The taste receptors for these five basic tastes are of two major types:

• Ligand-gated channels (ionotropic receptors): for salt and sour taste

• G-Protein Coupled Receptors (metabotropic receptors): for sweet, bitter and umami

• Salt and sour taste are mediated through epithelial sodium channels (ENaC)

• Na+ movement through ENaC → salt taste

• H+ movement through ENaC → sour taste

• Sweet, bitter, and umami are sensed via GPCRs

• T1R2 and T1R3 families→ sweet taste

• T2R family → bitter taste

• T1R1, T1R3, and Metabotropic glutamate receptor (mGluR4) → Umami (Option A and C)

Solution to Question 6:

In the given scenario of phantom limb pain, on the radiological investigation of the brain projection of adjacent fibres to overlap the left somatosensory cortex can be seen. This pain is based on cortical plasticity.

Cortical structures are capable of reorganization after the loss of a body part. The cortical representation of the neighbouring parts spreads into the cortical area that was formerly occupied by the representation of the amputated part. This is called cortical plasticity leads to the phenomenon of phantom limb pain. The sensory cortical area occupied by the right limb is the left somatosensory cortex.

Cortical plasticity / Phantom limb: It is the sensation that an amputated or missing limb is still attached. Plasticity is the ability of the brain to adapt to such a condition by reorganization of thalamic and cortical structures.

The converse also holds true: if the cortical area representing a limb is removed, the somatosensory map of the limb moves to the surrounding cortex.

Diagnosis and Management :

• PET scan is used to document these plastic changes in cortex.

• Use of epidural analgesia during amputation surgery decreases incidence of phantom pain.

• Spinal cord stimulation also helps in reducing phantom pain. (interferes with ascending impulses to brain)

Solution to Question 7:

There is an increase in the number of GLUT 4 receptors in response to exercise. This increases the insulin dependent glucose uptake into the cells, contributing to a decrease in insulin resistance.

GLUT-4 is an insulin-dependent GLUT. Other GLUTs are not insulin-sensitive and are constitutively expressed in the cell membrane.

Insulin resistance is a condition in which the cells in muscles, fat, and liver are less responsive to insulin and have difficulty absorbing glucose from the blood. The pancreas then produces more insulin to facilitate the entry of glucose into the cells.

An arbitrary but clinically useful benchmark considers patients insulin resistant if they require greater than 1 unit/kilogram/day of exogenous insulin to maintain glycemic control. Patients requiring greater than 200 units of exogenous insulin per day are considered severely insulin resistant.

Other options

Option A: GLUT 1 receptors are present in the red blood cell, brain, kidney, colon, retina, and placenta. It is responsible for glucose uptake.

Option B: GLUT-2 is found in the liver, pancreatic β cell, small intestine, and kidney. Its function is the rapid uptake or release of glucose.

Option C: GLUT-5 is present in the small intestine and is involved in the absorption of fructose.

Solution to Question 8:

Ferrireductase is an enzyme that converts Fe^{3+} (ferric iron) to Fe^{2+} (ferrous iron), not the other way around. Ferrous is the form of iron that is more readily absorbed in the gut.

Almost all the iron absorption occurs in the duodenum.

Most of the iron in the diet is in the ferric (Fe^{3+}) form, whereas it is the ferrous (Fe^{2+}) form that is absorbed.

Body iron stores are controlled by intestinal absorption as there is no exit to iron from the body. Divalent metal transporter 1 (DMT1) transports non-heme iron into the enterocyte. Iron absorption in the duodenum is regulated by hepcidin release from the liver in response to increases in intrahepatic iron levels. It binds to ferroportin leading to endocytosis and degradation within the enterocyte. This prevents the transfer of iron from enterocyte to plasma resulting in iron trapping within duodenal cells in the form of mucosal ferritin. The trapped iron is lost as these cells are sloughed. The iron transported by ferroportin into the plasma is transferred to other parts with the help of transferrin and stored as serum ferritin and hemosiderin. The image depicting iron absorption and storage is given below:



Other options:

Option B: RBC protoporphyrin is increased in deficiency, increased protoporphyrin levels in red blood cells can indicate iron deficiency or disorders of heme synthesis.

Option C: Iron is primarily excreted through the gut, skin, and urinary tract, but the body generally has very limited mechanisms for iron excretion compared to iron absorption and recycling.

Option D: The body's iron homeostasis is finely tuned, and a significant loss of red blood cells, such as more than 10 ml per day, can overwhelm the body's ability to compensate and maintain iron balance.

Solution to Question 9:

Pacinian corpuscles are responsible for sensing fast vibration. The chronic tingling sensation is mediated by pressure-sensitive Aβ fibers and low-conduction velocity C fibers.

Pacinian corpuscles are the most sensitive mechanoreceptors in the somatosensory system. They are responsive to high-frequency (30–500 Hz) vibratory stimuli and deep pressure. They are also examples of rapidly adapting receptors.

Chronic tingling causing conditions include Diabetes, carpal tunnel syndrome, ulnar nerve palsy, radial nerve palsy, tarsal tunnel syndrome, sciatica, vitamin B1,B6,B12 deficiencies, and alcoholism.

Other options

Option A: The Merkel cells are expanded dendritic endings in the epidermis of glabrous skin and respond to sustained pressure and touch.

Option B: Meissner's corpuscle senses tap and flutter. They are present in dermal papillae. They are rapidly adapting receptors.

Option D: Ruffini's corpuscles are enlarged dendritic endings with elongated capsules in the dermis of glabrous and hairy skin. They respond to stretch and fluttering vibrations. They are slowly adapting receptors.



Solution to Question 10:

The flocculonodular lobe is involved in this patient with an alcoholic gait and abnormal eye movements (nystagmus) following an RTA.

The cerebellum has three parts based on function.

• Vestibulocerebellum (or flocculonodular lobe): The oldest part of the cerebellum which contains nodulus in the vermis and the flanking flocculus. It has vestibular connections and is concerned with equilibrium and gait. Loss of it can cause disturbance of equilibrium and postural movements, especially during rapid movements.

• Spinocerebellum: The region that receives proprioceptive input from the body as well as a copy of the “motor plan” from the motor cortex. By comparing the plan with performance, it smooths and coordinates movements that are ongoing. The superior vermis is a part of the spinocerebellum. Loss of it can cause problems with coordination.

• Cerebrocerebellum: They are the newest from a phylogenetic point of view, reaching their greatest development in humans. They interact with the motor cortex in planning and

programming movements.

Other options:

Option B: The Dentate nucleus(largest) is one of the 4 deep cerebellar nuclei others being, Emoliform, Globose, and Fastigial nuclei. These receive input from climbing fibers and are the only motor output from the cerebellar cortex.

Option C: Anterior lobe is a part of the Paleo/Spinocerebellum responsible for the coordination of movement and regulation of muscle tone whose lesion results in hypotonia.

Option D: Interposed nuclei / Nucleus interpositus is the term used to describe the cerebellar nuclei, Emboliform, and Globose.

Solution to Question 11:

Compensation in metabolic acidosis happens late in the kidney because it takes hours to days for it to synthesize enzymes that produce NH4+(ammonium ions).

Acid-base balance equation: CO2 + H2O -----> H2CO3 -----> H+(acid) and HCO3-(base)

• First line of defence (in seconds): Buffers (phosphate and ammonium)

• Second line of defence (in Minutes): Lungs

• Third line of defence (in hours - days): Kidney

The kidneys regulate the acid-base balance by two major mechanisms: the reabsorption of bicarbonate and the generation of new bicarbonate via ammonia production and distal nephron acid secretion. Urinary NH4+ is produced primarily in the proximal tubule cells by converting glutamine to glutamate.

Renal failure leads to a buildup of acids due to loss of excretory function of the kidney. The decreased glomerular filtration rate also leads to diminished excretion of the renal buffers phosphate and ammonium. Thus, bicarbonate reabsorption decreases, resulting in metabolic acidosis.

Solution to Question 12:

The channel needed for the transport of iodine from the follicle to the colloid is Pendrin.

Steps in synthesis and secretion of thyroid hormone:

• Iodide uptake: The basolateral sodium-iodide symporter (NIS) actively takes up I-(iodide) from the circulation into the follicular cells. It is then transported into the colloid through a pendrin (chloride-iodide) transporter

• Thyroglobulin synthesis: Thyroglobulin (Tg), is a large glycoprotein rich in tyrosine residues. It is synthesized by follicular cells.

• Iodination of Tg: This is regulated by the thyroid peroxidase (TPO) enzyme. TPO has three functions:

• Oxidation: TPO oxidizes I- to I2 (iodine) in the presence of H2O2

• Organification: TPO links tyrosine residues of thyroglobulin with iodine (I2) to form monoiodotyrosine (MIT), diiodotyrosine (DIT)

• Coupling reactions: TPO combines iodinated tyrosine residues to make MIT + DIT = T3 (triiodothyronine), DIT + DIT = T4 (tetraiodothyronine)

• Storage: Thyroid hormones are bound to thyroglobulin and stored in the follicular lumen as colloids. The colloid has hormone reserves for 2-3 months

• Release: Iodinated thyroglobulin is taken up by the thyroid follicles via endocytosis and fuses with lysosomes. It is cleaved by lysosomal proteolytic enzymes into MIT, DIT, T3, and T4. T4 (80%) and T3 (20%) are released into the blood

• Peripheral T3 conversion: T4 undergoes deiodination in extrathyroidal tissue to form T3, which is the active form of the hormone.

Pendred syndrome: Pendrin is present in the thyroid gland and inner ear. The mutation causes goiter and sensorineural hearing loss.



Other options

Option A: Sodium-iodide symporter (NIS) is responsible for transporting iodine from the bloodstream into the thyroid follicular cell, not into the colloid. It works by bringing in iodine along with sodium from the bloodstream into the follicle.

Option B: Calmodulin is structurally and functionally related to troponin. In place of troponin, smooth muscle cells contain a large amount of the regulatory protein calmodulin; it is different in the manner in which it initiates contraction. Calmodulin initiates contraction by activating the myosin light chain kinase.

Option C: Claudin is a transmembrane protein involved in the formation of tight junctions.

Solution to Question 13:

Increased permeability of potassium ions(K+) is responsible for the RMP of a nerve as there are more K+ channels open at rest, facilitating the flow of potassium ions across the membrane.

Resting membrane potential (RMP) is the difference in the electrical potentials between the cytoplasm of the cell and the extracellular space.

At rest, there are more open K+ channels compared to Na+ channels, thus the membrane permeability of K+ is greater. As a result of this, the intracellular and extracellular K+ concentrations are the major determinants of the RMP. Furthermore, the RMP is close to the equilibrium potential of K+ in general.



Clinical Significance: In hyperkalemia, the extracellular level of K+ is increased. Thus the RMP moves closer to the threshold to elicit an action potential leading to a more excitable nerve. The excitable nerve, stimulates the muscle repeatedly causing fasciculations, exhaustion of muscle, and weakness. In hypokalemia, the extracellular level of K+ is reduced. Thus the RMP is reduced and the neuron is hyperpolarized.

Solution to Question 14:

The sequence of steps involved at neuromuscular junction endplate potential is Calcium entry-Acetylcholine release- Na+ entry.

Events occurring at the Chemical Synapse:

The action potential arriving at the end of the motor neuron increases the permeability of its endings to Ca2+

↓

Ca2+ enters the endings and triggers a marked increase in exocytosis of the acetylcholine-containing synaptic vesicles

↓

The binding of acetylcholine to nicotinic receptors increases the Na+ and K+ conductance

↓

The resultant influx of Na+ produces a local depolarizing potential, the endplate potential

↓

The current created by this local potential depolarizes the adjacent muscle membrane to its firing level

↓

Action potentials are generated on either side of the endplate and are conducted away from the endplate in both directions along the muscle fiber. This muscle action potential, in turn, initiates muscle contraction.

Acetylcholine is then removed from the synaptic cleft by acetylcholinesterase, which is present in high concentration at the neuromuscular junction.



Biochemistry AIIMS 2017

Question 1:

Identify the anaplerotic reaction among the following?

a) Conversion of pyruvate to Acetyl CoA

b) Conversion of pyruvate to Oxaloacetate

c) Conversion of pyruvate to Acetaldehyde

d) Conversion of pyruvate to Lactic acid

Question 2:

True regarding mitochondrial DNA is:

a) One set from each parents

b) It has a higher mutation rate than nuclear DNA

c) It codes for more than 25% of oxidative phosphorylation proteins

d) It has 3 X 10^9 base pairs

Question 3:

Which of the following is Thermogenic food

a) High fat diet

b) It does not depend on the macronutrients

c) High protein diet

d) High carbohydrate diet

Question 4:

Klenow fragment lacks the activity of

a) 3'→5' Exonuclease

b) 5'→3' Exonuclease

c) 5'→3' DNA polymerase

d) 3'→5' DNA polymerase

Question 5:

In SCID, what type of DNA repair is defective:

a) Base excision repair

b) Nucleotide excision repair

c) Mismatch Repair

d) Non-homologous end joining repair

Question 6:

A 10-year-old boy rapidly develops hypoglycemia after moderate activity. Blood examination reveals raised levels of ketone bodies, lactic acid, and triglycerides. On examination, the liver and kidneys are found to be enlarged. Histopathology of the liver shows deposits of glycogen in an excess amount. What is the diagnosis?

a) von Gierke's disease

b) Pompe's disease

c) McArdle's disease

d) Cori's disease

Question 7:

Which one of the following is not involved in iron metabolism:

a) Ceruloplasmin

b) Hepcidin

c) Transthyretin

d) Transferrin

Question 8:

Which of the following amino acids is not formed after post-translational modification?

a) Selenocysteine

b) Triiodothyronine

c) Hydroxy proline

d) Hydroxylysine

Question 9:

Which test will be positive in starvation state in urine?

a) Benedict test and heat coagulation test

b) Heat coagulation test

c) Rothera's test

d) Heat coagulation test and Rothera's test

Question 10:

Identify the marked bond:



a) Ionic bond

b) Covalent bond

c) Hydrogen bond

d) Phosphodiester Bond

Question 11:

Which among the following serves as cofactor for Glycogen phosphorylase:

a) Thiamine

b) Niacin

c) Pyridoxal Phosphate

d) Biotin

Question 12:

Glucosamines are used in which of the following conditions:

a) Arthritis

b) Niemann Pick disease

c) Alzheimer's disease

d) Cancer

Question 13:

Which of the following site does the restriction enzyme act when RFLP is used in-order to identify the five different species of staphylococci in a surgical ICU?

a) ATGGAC-TACGTG

b) GATATC- CTATAG

c) TAGATA-ATCTCT

d) AATATA-TATAAT

Question 14:

Thiamine deficiency results in decreased■ energy production, because TPP _____

a) Interferes with alcohol metabolism

b) Interferes with transketolase activity

c) Is coenzyme for pyruvate and alpha ketoglutarate dehydrogenase

d) Interferes with energy production from amino acids

Question 15:

A young patient came with complaints of acute abdominal pain on and off and tingling sensation of limbs and at times weakness of limbs. She had a history of eating paint from the

wall of a newly built house. Which of the following enzyme defect will be the cause of her condition?

a) ALA synthase

b) ALA dehydratase

c) Heme synthase

d) Coproporphyrinogen oxidase

Question 16:

Under anaerobic metabolism which of the following substrates yields 3 molecules of ATP?

a) Glycogen

b) Amino acid

c) Glucose

d) Galactose

Question 17:

Which of the following is the reason for using phenylbutyrate in urea cycle disorders?

a) It maintains renal output

b) It scavenges nitrogen

c) It maintains energy production

d) It activates enzymes

Question 18:

Which of the following can alter the gene expression by methylation and acetylation but not change the gene sequence?

a) Mutation

b) Epigenetics

c) Translocation

d) Inversion

Question 19:

Which of the following is not seen in low insulin-glucagon ratio?

a) Glycogen breakdown

b) Ketogenesis

c) Glycogen storage

d) Gluconeogenesis

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	b
5	d
6	a
7	c
8	a
9	c
10	c
11	c
12	a
13	b
14	c
15	b
16	a
17	b
18	b
19	c

Detailed Explanations

Solution to Question 1:

The conversion of pyruvate to oxaloacetate is an anaplerotic reaction. It is catalyzed by pyruvate carboxylase.

Anaplerotic reactions are chemical reactions that form intermediates of a metabolic pathway. The above reaction is important in maintaining an adequate concentration of oxaloacetate for the condensation reaction with acetyl-CoA. If acetyl-CoA accumulates, it acts as an activator of pyruvate carboxylase and an inhibitor of pyruvate dehydrogenase, thereby ensuring an adequate supply of oxaloacetate.

Other anaplerotic reactions are:

• Compounds that are metabolized to yield propionyl CoA, which can be then carboxylated and isomerized to succinyl CoA.

• Glutamate and glutamine are important anaplerotic substrates because they yield α-ketoglutarate as a result of the reactions catalyzed by glutaminase and glutamate dehydrogenase.

• Transamination of aspartate leads directly to the formation of oxaloacetate.

Solution to Question 2:

The true statement is that mitochondrial DNA has a high mutation rate.

Mitochondrial DNA is circular and double-stranded. It has 16,569 bp (that makes up approximately 1% of total cellular DNA) while nuclear DNA has 3.3 billion bp. It has 37 genes that encode 13 protein subunits of the respiratory chain (out of a total of 67 proteins) i.e. about 20%. It mutates at a rate 5-10 times that of nuclear DNA. Mitochondrial DNA is transmitted by maternal non-mendelian inheritance.



Solution to Question 3:

A high intake of protein results in a higher rate of protein turnover, which explains the increased diet-induced thermogenesis seen in people consuming a high protein diet.

The state of protein nutrition can be determined by measuring the difference between intake and output of nitrogenous compounds which are known as nitrogen balance.

• In a healthy adult, nitrogen balance is in equilibrium, when intake equals output, and there is no change in the total body content of protein.

• In a growing child, a pregnant woman, the excretion of nitrogenous compounds is less than the dietary intake of nitrogen in the body as protein— positive nitrogen balance.

• In response to trauma or infection, or if the intake of protein is inadequate to meet requirements, there is the net loss of protein nitrogen from the body—negative nitrogen balance.

A high intake of protein does not lead to positive nitrogen balance; although it increases the rate of protein synthesis, it also increases the rate of protein catabolism, so that nitrogen equilibrium is maintained with a higher rate of protein turnover.

Both protein synthesis and catabolism are ATP expensive, hence the increased diet-induced thermogenesis seen in people consuming a high protein diet.

Solution to Question 4:

Klenow fragment lacks 5’ → 3’ exonuclease activity.

DNA polymerase I is an enzyme involved in DNA replication. It has polymerase activity along with both 3' → 5' and 5' → 3' exonuclease activity. When the 5' → 3' exonuclease domain is removed, the remaining fragment is called Klenow fragment. It retains the polymerization and proofreading activities of DNA polymerase I.

Solution to Question 5:

SCID is due to defective non-homologous end joining repair.

Defects in DNA Repair:

Non-Homologous End Joining Repair	Severe combined immunodeficiency disease (SCID)
Homologous Repair	Bloom syndrome Werner syndrome Breast cancer susceptibility
Nucleotide Excision Repair	Xeroderma pigmentosum
Mismatch Repair	Hereditary nonpolyposis colorectal cancer (HNPCC)
Base Excision Repair	MUTYH-associated polyposis (MAP)

Solution to Question 6:

The given features are consistent with von Gierke's disease. It is the most common glycogen storage disease.

Enzyme deficient in this condition is glucose-6-phosphatase (common to both glycogenolysis and gluconeogenesis). Inheritance is autosomal recessive.

Von Gierke disease (type I glycogen storage disorder) occurs due to the deficiency of glucose-6-phosphatase enzyme in the liver, kidney, and intestinal mucosa. Patients present commonly by 3-4 months of age with hypoglycemia (leading to seizures), lactic acidosis, and hepatomegaly. Affected children often have a doll-like face with fat cheeks, relatively thin extremities, short stature, and a protuberant abdomen due to massive hepatomegaly. The kidneys are also enlarged, whereas the spleen and heart are not involved. Laboratory findings include hypoglycemia, lactic acidosis, hyperuricemia, and hyperlipidemia.

A definitive diagnosis is made using a liver biopsy. Fructose and galactose restriction in the diet is advised.

Solution to Question 7:

• Hepcidin, ceruloplasmin, and transferrin are involved in iron metabolism.

• Transthyretin transports thyroxine and retinol-binding protein and has no role in iron metabolism

Proteins involved in iron metabolism:

• Ferritin and hemosiderin: Iron storage

• Transferrin: Iron transport

• Ceruloplasmin: a copper-containing enzyme with ferroxidase activity

• Hepcidin: Chief regulator of systemic iron homeostasis

Solution to Question 8:

Selenocysteine, the 21st amino acid, is formed by co-translational modification. It is coded by the UGA codon, which is normally a stop codon. The protein synthetic apparatus can identify a selenocysteine-encoding UGA by the presence of the selenocysteine insertion sequence (SECIS) in the untranslated region of mRNA.

Selenocysteine is an analog of cysteine and is synthesized from serine.

Selenocysteine is present at the active site of several human enzymes that catalyze redox reactions such as thioredoxin reductase, glutathione peroxidase, and deiodinase. Since Glutathione peroxidase contains selenocysteine and hence it is an antioxidant.

Impairments in human selenoproteins have been implicated in tumorigenesis and atherosclerosis and are associated with selenium deficiency cardiomyopathy (Keshan disease).

Solution to Question 9:

Detectable levels of ketone bodies in urine (ketonuria) are seen in conditions characterized by increased ketogenesis such as starvation. The urine test done for ketone bodies is the Rothera's test in which formation of a purple ring indicates presence of ketone bodies. Gerhardt's test is also done to detect ketone bodies in urine.

Benedict' test done for reducing sugars.

Heat coagulation test done for proteins.

Solution to Question 10:

The marked bond represents a hydrogen bond between thymine and adenine.

The complementary nitrogenous bases of the 2 strands of the DNA double helix are linked by hydrogen bonds. Adenine and thymine are linked by two hydrogen bonds (A=T) whereas three hydrogen bonds are present between cytidine and guanine (C≡G). Van der Waals and hydrophobic interactions are present between the stacked adjacent base pairs.

A β-N-glycosidic bond is present between the nitrogenous base and deoxyribose sugar. C1 of deoxyribose/ribose sugar is joined covalently to N1 of pyrimidines and N9 of purines by β-N-glycosidic bond. An ester bond is present between the sugar and phosphate group of the same nucleotide. An acid anhydride bond is present between two phosphate groups.

A polymer of nucleotides is joined by 3'-5' phosphodiester bond. The bond is present between the sugar of one nucleotide and the phosphate group of another.



Solution to Question 11:

• Pyridoxal phosphate is a cofactor of glycogen phosphorylase, where the phosphate group is catalytically important.

• PLP also acts as the coenzyme for many enzymes involved in amino acid metabolism, especially transamination and decarboxylation.

• Glycogen phosphorylase is a debranching enzyme which cleaves alpha-1,4 bonds of the glycogen chain to produce glucose-1-phosphate in glycogenolysis.



Solution to Question 12:

The combined use of glucosamine sulfate and chondroitin sulfates in the treatment of degenerative joint disease has become an extremely popular supplementation protocol in arthritic conditions of the joints.

• Glucosamines are amino sugars which form the major component of glycosaminoglycans. (GAG's are tandem repeats of amino sugar and uronic acid)

• GAG's combine with proteins to form proteoglycans

• One such proteoglycan is aggrecan containing several GAGs (hyaluronic acid, chondroitin sulphate, and keratan sulphate) and both link and core proteins

• Aggrecan is the major proteoglycan in cartilage

• Hence, from this we can deduce that glucosamines being a major part of GAG's and proteoglycans such as aggrecan are protective against arthritis.

Solution to Question 13:

Restriction endonuclease enzymes act at specific palindromic sequences that create DNA fragments of different lengths. By analyzing the fragments, specific species of bacteria can be identified. Option B is the only palindromic sequence here, hence it is the answer.

RFLP is a restriction fragment length polymorphism (RFLP) analysis technique that involves cutting a particular region of DNA with known variability, with restriction enzymes, then separating the DNA fragments by agarose gel electrophoresis and determining the number of fragments and relative sizes.

Cleavage of a sample of DNA with a restriction enzyme produces a characteristic set of smaller DNA fragments. Deviations in the normal product pattern, produce RFLPs that occur if a mutation renders a restriction site unrecognizable and alternatively, generates a new recognition site.

Solution to Question 14:

Thiamine deficiency results in decreased energy production because thiamine pyrophosphate (TPP) is a coenzyme of pyruvate and α-ketoglutarate dehydrogenase, which are required for the production of ATP using NADH.

Thiamine pyrophosphate is an important coenzyme in three multienzyme complexes that catalyze oxidative decarboxylation reactions:

• Pyruvate dehydrogenase in carbohydrate metabolism.

• α-ketoglutarate dehydrogenase in the citric acid cycle.

• Branched-chain ketoacid dehydrogenase involved in the metabolism of leucine, isoleucine and valine.

NAD is a co-factor in pyruvate and α-ketoglutarate dehydrogenase. Hence in thiamine deficiency, there is decreased ATP production owing to reduced NADH entering into the electron transport chain.

Option A: Thiamine does not interfere in alcohol metabolism but alcohol interferes directly with the absorption of thiamine and with the synthesis of thiamine pyrophosphate.

Option B: Thiamine is also the coenzyme for transketolase in the pentose phosphate pathway (HMP shunt) but there is no ATP synthesized in this pathway.

Solution to Question 15:

A child with acute abdominal pain, tingling sensation and weakness of limbs with history of eating paint likely has lead poisoning. It is characterized by impaired ALA dehydratase enzyme activity.

Lead is a potent inhibitor of both ALA dehydratase and ferrochelatase/ heme synthase (option C). However, it has the greatest inhibitory action on ALA dehydratase. Lead poisonining thus leads to the buildup of protoporphyrin and ALA. Lead also substitutes for calcium in many fundamental

cellular processes. It can cross red blood cell membranes as well as the blood-brain barrier and enter the neuroglial cells which support brain function. The fatal dose of lead is 20g of lead acetate and the fatal period is 1 to 2 days.



Other options:

Option A: ALA synthase 2 defect causes X-linked protoporphyria.

Option C: Heme synthase, or ferrochelatase, defect causes erythropoeitic protoporphyria.

Option D: Coproporphyrinogen oxidase defect causes hereditary corproporphyria.

Solution to Question 16:

Breakdown of glycogen under anaerobic conditions yields 3 ATPs.

Breakdown of glycogen yields glucose-6-phosphate which can enter the glycolysis pathway. Since it bypasses the first step, one ATP is saved and anaerobic glycolysis yields 3 ATPs.

The first step in glycolysis involves the conversion of glucose to glucose-6-phosphate which requires one ATP. Glycogen breakdown leads to formation of glucose-6-phosphate which can enter glycolysis hence bypassing the first step and saving one ATP.

One ATP is used for conversion of fructose-6-phosphate to fructose 1,6 biphosphate hence net ATP formed from anaerobic glycolysis of glycogen is 3ATP

REACTION	ENZYME	METHOD OF ATP FORMATION	NET ATP PRODUCED
1,3-bisphosphoglycerate to 3 phosphoglycerate	Phosphoglycerate kinase	Substrate level phosphorylation	2 ATP
Phosphoenol pyruvate to Pyruvate	Pyruvate kinase	Substrate level phosphorylation	2 ATP
		TOTAL PRODUCED	4 ATP



Solution to Question 17:

Phenylbutyrate scavenges nitrogen by combining with amino acids to form a more water-soluble product, which can be easily excreted by the kidneys.

Phenylbutyrate combines with glutamate to form pheynylbutylglutamine.

Other substances that can be used as nitrogen scavengers are:

• Combination of benzoate (forms hippurate on combining with endogenous glycine)

• Phenylacetate (forms phenylacetylglutamine on combining with glutamate) available as ammonia.

• Arginine is also useful.

Solution to Question 18:

Epigenetics is defined as heritable changes in gene expression that are not caused by alteration in gene sequence.

It means “above genetics” and refers to the fact that gene regulatory mechanisms do not change the underlying regulated DNA sequence, but rather simply the expression patterns of the DNA.

Epigenetic mechanisms play a key role in establishment, maintenance, and reversibility of transcriptional states. A key feature is that the controlled transcriptional on/off states can be maintained through multiple rounds of cell division. This observation indicates there must be robust mechanisms to maintain and stably propagate these epigenetic states.

Histone Methylation Mostly Makes DNA Mute.

Histone Acetylation: Relaxes DNA coiling, allowing for transcription. Histone Acetylation makes DNA Active.

Solution to Question 19:

A low insulin/glucagon ratio indicates low insulin and high glucagon levels, which are observed in the fasting state. In the fasting state, all of the above processes (i.e., glycogen breakdown, ketogenesis, and gluconeogenesis) occur, but not glycogen storage.

Metabolic process	Fasting state/ low insulin-glucagon ratio	Fed state/ high insulin-glucagon ratio
Gluconeogenesis	Increased	Decreased
Glycogenolysis	Increased	Decreased
Ketogenesis	Increased	Decreased
Glycogen storage	Decreased	Increased

Biochemistry AIIMS 2018

Question 1:

An adolescent male patient presented with pain in the calf muscles on exercise. On biopsy, an excessive amount of glycogen was found to be present in the muscle. What is the most likely enzyme deficiency?

a) Muscle-debranching enzyme

b) Phosphofructokinase 1

c) Glucose 6 phosphatase

d) Muscle glycogen phosphorylase

Question 2:

Alkali disease of livestock is caused by an excess of which of the following minerals?

a) Selenium

b) Manganese

c) Molybdenum

d) Iron

Question 3:

Which of the following types of collagens is involved in wound healing?

a) Type II

b) Type III

c) Type IV

d) Type V

Question 4:

An infant is brought with a history of vomiting and poor feeding. A musty odor is noted in the baby. Guthrie test was done and it was found to be positive. All are true regarding this disease except

a) Phenylalanine hydroxylase enzyme defect

b) White patch of hair due to tryptophan deficiency

c) Phenylacetate positive in urine

d) Mental retardation is present

Question 5:

In a reaction, the substrate is available in a concentration that is 1000 times the Km value of the enzyme. After 9 minutes of reaction, 1% of substrate is converted to product (12 microgram/ml). If the concentration of the enzyme is changed to 1/3 and concentration of substrate is doubled, what is the time taken to convert the substrate into the same amount of product that i.e. 12 microgram/ml?

a) 9 minutes

b) 4.5 minutes

c) 27 minutes

d) 13.5 minutes

Question 6:

The best method to measure HbA1C is _____

a) Isoelectric focusing

b) Affinity chromatography

c) Ion exchange chromatography

d) Electrophoresis

Question 7:

Which of the following is the best screening investigation for metabolic disorders?

a) Western blot

b) Tandem mass spectrometry

c) PCR

d) Gel electrophoresis

Question 8:

Which of the following enzyme activity decreases in fasting?

a) Hormone-sensitive lipase

b) Glycogen phosphorylase

c) Pyruvate carboxylase

d) Phosphofructokinase 1

Question 9:

Fluoroacetate is a component of pesticides. Which of the following metabolic pathways is inhibited by this compound?

a) TCA cycle

b) Glycolytic pathway

c) Oxidative phosphorylation

d) ETC

Question 10:

Which of the following does not require 5' capping?

a) tRNA of Alanine

b) U6 snRNA

c) mRNA for Histone

d) siRNA

Question 11:

Vitamin K in its coenzyme form is regenerated by the enzyme ____

a) Glutathione reductase

b) Quinone reductase

c) Vitamin K reductase

d) Epoxide reductase

Question 12:

Identify the phenomenon shown below



a) Translation

b) Transformation

c) Complementation

d) Conjugation

Question 13:

A middle-aged woman presents with fissured tongue and peripheral neuropathy. Investigations showed reduced glutathione reductase activity. Which vitamin deficiency is the likely cause of this?

a) Vitamin B1

b) Vitamin B2

c) Vitamin B6

d) Vitamin B12

Question 14:

Which of the following enhances iron absorption?

a) Vitamin A

b) Vitamin C

c) Thiamine

d) Riboflavin

Question 15:

Which of the following does not favor euchromatin formation due to changes occurring at cytosine residues at CpG islands in DNA?

a) Methylation

b) Phosphorylation

c) Alkylation

d) Sumoylation

Question 16:

Which of the following techniques is based on RNA?

a) RT PCR

b) Sanger's technique

c) Next generation sequencing

d) Western blot

Question 17:

Vitamin C cannot be produced in humans due to lack of

a) L- gulonolactone oxidase

b) Xylitol reductase

c) Pyruvate dehydrogenase

d) UDP glucose dehydrogenase

Question 18:

Klenow fragment lacks the activity of

a) 3'→5' Exonuclease

b) 5'→3' Exonuclease

c) 5'→3' DNA polymerase

d) 3'→5' DNA polymerase

Question 19:

Child presents with hypotonia and seizures. It was confirmed to be Zellweger syndrome. Which among the following accumulates in brain in this condition?

a) Glucose

b) Lactic acid

c) Long chain fatty acid

d) Triglycerides

Question 20:

A 12-year-old girl suffers from an inadequately treated type 1 diabetes mellitus. She is currently hyperglycemic. Which of the following is most likely to occur?

a) Glycogenesis in muscle

b) Increased protein synthesis

c) Increased conversion of fatty acids to acetyl-CoA

d) Decreased cholesterol synthesis

Question 21:

What would be the respiratory quotient if someone is on an exclusive carbohydrate diet?

a) 0.7

b) 0.8

c) 1

d) 1.2

Answer Key

Question No.	Correct Option
1	d

2	a
3	b
4	b
5	c
6	c
7	b
8	d
9	a
10	a
11	d
12	c
13	b
14	b
15	a
16	a
17	a
18	b
19	c
20	c
21	c

Detailed Explanations

Solution to Question 1:

The given clinical features are seen in McArdle's disease. This patient has muscle glycogen phosphorylase deficiency.

The lack of muscle glycogen phosphorylase results in excess glycogen build-up in muscles. The image given below shows the vacuolar myopathy seen in McArdle's disease.



Clinical features include poor exercise tolerance, abnormally high muscle glycogen (2.5-4%), and a very low blood lactate after exercise

The images given below show myophosphorylase enzyme histochemistry showing (A) Normal muscle and (B) McArdle's disease.



Solution to Question 2:

Excess of selenium causes alkali disease in livestock.

Clinical types of selenium intoxication

• Acute selenosis

• Subacute selenosis (i.e., blind staggers type)

• Chronic selenosis (i.e., alkali disease type)

Alkali disease occurs when animals consume moderate levels of selenium for a period of weeks to months.

The common clinical signs of chronic selenosis in horses, cattle, and swine are loss of hair (horses and cattle lose long hair from the mane and tails), emaciation, hoof lesions, weight reduction, and lameness.

Solution to Question 3:

Type III collagen is involved in wound healing.

Collagen types I and III predominate in the skin and aponeurotic layers. In the early stages of wound healing, type III collagen is synthesized. It is replaced by type I collagen as wound healing progresses.

The tensile strength of the healing wound is directly proportional to the amount and type of collagen present. During healing, as disorganized collagen is degraded and reformed, covalent cross-links are formed that enhance tensile strength. The maximum strength of the healed wound depends upon the interconnection of collagen subunits. Approximately 80 percent of the original strength of the tissue is obtained by six weeks after surgery. Wounds slowly continue to get stronger but may never achieve 100 percent of their previous strength.

Solution to Question 4:

The given scenario of an infant with a history of vomiting and poor feeding with a musty odour is suggestive of phenylketonuria (PKU). Tryptophan deficiency is not a feature of this disorder. The deficiency of the enzyme phenylalanine hydroxylase causes PKU.

Characteristic clinical findings in Phenylketonuria:

• Mental retardation

• Failure to walk or talk

• Seizures

• Hyperactivity

• Tremor

• Microcephaly

• Failure to grow

• Deficiency of pigmentation (fair hair, light skin color, and blue eyes)

The Guthrie bacterial inhibition test is a screening test for phenylketonuria. This test uses a phenylalanine requiring strain of Bacillus subtilis. The ferric chloride test is also used as a screening test for PKU. Tandem mass spectrometry is the gold standard investigation.

Solution to Question 5:

The time taken for the formation of the same amount of product is 27 minutes.

In this reaction, the substrate concentration becomes 1000 times i.e much greater than the Km value. Thus, the rate of this reaction is equal to Vmax and is thus independent of the substrate concentration i.e. it follows zero-order kinetics. Thus, doubling the substrate concentration does not change the velocity of the reaction i.e. it remains as Vmax.

No matter the substrate concentration, the rate of the reaction is directly proportional to the concentration of the enzyme. In this situation the enzyme concentration becomes 1/3rd. Thus, the velocity of the reaction (Vmax) also becomes 1/3rd. This means that the time taken for the formation of the same amount of product will become tripled i.e 9 × 3 = 27 minutes.



Solution to Question 6:

The best method to measure HBA1C is ion-exchange chromatography.

In ion-exchange chromatography:

• Proteins interact with the stationary phase by charge-charge interactions.

• Proteins with a net positive charge at a given pH will tightly adhere to beads with negatively charged functional groups such as carboxylates or sulfates (cation exchangers).

• Proteins with a net negative charge adhere to beads with positively charged functional groups, typically tertiary or quaternary amines (anion exchangers).

• Non-adherent proteins flow through the matrix and are washed away.

• Proteins elute in inverse order of the strength of their interactions with the stationary phase.

Solution to Question 7:

Tandem mass spectrometry (MS) is the technique of choice for screening neonates for inherited metabolic diseases.

Tandem MS employs the equivalent of two mass spectrometers linked in series (MS-MS or MS2). It can be used to screen blood samples from newborns for the presence and concentrations of amino acids, fatty acids, and other metabolites.

Abnormalities in metabolite levels indicate inherited metabolic diseases, such as phenylketonuria, ethylmalonic encephalopathy, and glutaric acidemia type 1.

Solution to Question 8:

During fasting, there is low insulin, and hence, the activity of phosphofructokinase-1 gets inhibited.

Phosphofructokinase 1 is an enzyme of glycolysis. It is induced by insulin and a well-fed state.

Other options:

Option A: Hormone-sensitive lipase mobilizes stored fats in a fasting state.

Option B: Glycogen phosphorylase is active in fasting state so as to produce glucose from glycogen.

Option C: Pyruvate carboxylase is an enzyme of gluconeogenesis. During the fasting state, the liver is flooded with fatty acids mobilized from adipose tissue. The resulting elevated hepatic acetyl CoA activates pyruvate carboxylase. Other enzymes of gluconeogenesis that are elevated in fasting include Phosphoenolpyruvate carboxykinase, Fructose-1,6-bisphosphatase, and Glucose-6-phosphatase.

Solution to Question 9:

Fluoroacetate inhibits the TCA cycle enzyme aconitase.

Fluoroacetyl CoA condenses with oxaloacetate to form fluorocitrate which inhibits aconitase, causing citrate to accumulate. This way it inhibits TCA cycle enzyme non-competitively.

Sources of fluoroacetate exposure:

• Fluoroacetate in some plants

• Fluorinated compounds (anticancer agents, industrial chemicals, and pesticides) are metabolized to fluoroacetate.

Solution to Question 10:

tRNA of Alanine does not require 5’ capping.

Capping process protects the 5′ end of mRNA from the action of exonucleases and polyadenylation protects the 3′ end of the mRNA. All mRNA, snRNA, and siRNA undergo this

process, but tRNA are comparatively stable and do not require a capping at the 5’ end.

Note: rRNAs also do not require 5' capping.

Solution to Question 11:

The coenzyme form of Vitamin K is regenerated by the enzyme epoxide reductase.

Epoxide reductase reduces Vitamin K epoxide to its coenzyme form hydroquinone.



Note: In addition to vitamin K reductase, epoxide reductase can also catalyse the reduction of quinone to hydroquinone. Thus, it can regenerate hydroquinone by itself.

Solution to Question 12:

The image shows a schematic of the blue-white assay used in the screening of recombinant vectors for inserted DNA fragments. It uses the phenomenon of complementation (α-complementation of the β-galactosidase gene).

Blue–white assay:

The DNA of interest is ligated into a vector. The vector is then inserted into a competent host cell viable for transformation, which is then grown in the presence of X-gal (X-gal detects active β-galactosidase). Cells transformed with vectors containing recombinant DNA will produce white colonies; cells transformed with non-recombinant plasmids (i.e., only the vector) grow into blue colonies.

The image below shows the result of a blue–white assay.



Solution to Question 13:

This patient is most likely suffering from riboflavin/vitamin B2 deficiency.

Deficiency of riboflavin is characterized by cheilosis, desquamation, and inflammation of the tongue, seborrheic dermatitis, corneal vascularization, anemia, and personality changes.

Riboflavin nutritional status is assessed by measurement of the activation of erythrocyte glutathione reductase by FAD added in vitro.

Solution to Question 14:

Iron absorption is enhanced by vitamin C.

Vitamin C reduces the ferric iron to its ferrous form, thereby helping in the process of absorption of dietary iron from the intestine.

Vitamin C deficiency causes

• Decreased absorption of iron leading to microcytic-hypochromic anemia

• Decreased hydroxylation of proline and lysine, which results in defective collagen synthesis and bleeding manifestations.

Conversely, vitamin C toxicity (>2000 mg/ day) can lead to iron overloaded states.

Note: Alcohol and fructose also enhance iron absorption. However, the absorption of both inorganic and heme iron is impaired by calcium. So, a glass of milk with a meal significantly reduces iron availability.

Solution to Question 15:

Methylation does not favour euchromatin formation due to changes cytosine residues at CpG islands in DNA.

The transcriptional regulatory regions of active genes often contain a high frequency of CpG dinucleotides (referred to as CpG islands) and they are normally unmethylated. Therefore, methylation is an epigenetic modification that inactivates DNA, and hinders the formation of euchromatin (transcriptionally active chromatin). Epigenetic changes include alterations of chromatin structure by:

• Methylation of cytosine residues in CpG dinucleotides

• Modification of histones by acetylation or methylation

• Changes in higher-order chromosome structure

Sumoylation of histones (SUMO - Small Ubiquitin-related MOdifier) is associated with transcription repression. Other possible roles of posttranslationally modified histones:

• Acetylation of core histones is associated with chromosomal assembly during DNA replication.

• Phosphorylation of histone H1 is associated with the condensation of chromosomes during the replication cycle.

• ADP-ribosylation of histones is associated with DNA repair.

• Methylation of histones is correlated with activation and repression of gene transcription.

• Acetylation of histones H3 and H4 is associated with the activation or inactivation of gene transcription.

• Monoubiquitylation is associated with gene activation, repression, and heterochromatic gene silencing.

Solution to Question 16:

RT PCR is a RNA-based technique.

RT PCR is a method used to quantitate mRNA levels that relies upon the first step of cDNA copying of mRNAs catalyzed by reverse transcriptase prior to PCR amplification and quantitation.

Other techniques:

• Sanger's technique and Southern blot: DNA

• Western blot: protein

• Northern blot: RNA

• Next-generation DNA sequencing technologies include Illumina (Solexa) sequencing, Roche 454 sequencing, Ion torrent: Proton/PGM sequencing, SOLID sequencing.

Solution to Question 17:

Ascorbic acid (vitamin C) cannot be synthesized in human beings because of the absence of L-gulonolactone oxidase.

Uronic acid pathway catalyzes the conversion of glucose to glucuronic acid, ascorbic acid, and pentoses. Glucuronate is reduced to L-gulonate, the direct precursor of ascorbate in the animals capable of synthesizing this vitamin.

Solution to Question 18:

Klenow fragment lacks 5' → 3' exonuclease activity.

DNA polymerase I is an enzyme involved in DNA replication. It has polymerase activity along with both 3' → 5' and 5' → 3' exonuclease activity. When the 5' → 3' exonuclease domain is removed, the remaining fragment is called Klenow fragment. It retains the polymerization and proofreading activities of DNA polymerase I.

Solution to Question 19:

Elevated levels of very long chain polyenoic acids have been found in the brains of patients with Zellweger syndrome.

Zellweger syndrome:

• Autosomal recessive

• Cerebro-hepato-renal syndrome: Impaired neuronal migration and myelination, hepatomegaly, renal cysts

• Reduction or absence of functional peroxisomes in tissues.

• Accumulation of very long chain polyenoic acids in the brain.

• Severe neurological symptoms.

• Most patients die within the 1st year of life

Solution to Question 20:

With the background of inadequately treated type I diabetes mellitus, an insulin-deficient state in the body is expected which will lead to increased conversion of fatty acids to acetyl-CoA.

Type 1 diabetes mellitus is commonly found in the adolescent population. In response to certain precipitating factors, such patients may develop a severe insulin-deficient state. Since insulin is an inhibitor of hormone-sensitive lipase (HSL), an insulin-deficient state will cause HSL overactivity in adipocytes and the breakdown of lipid into fatty acids (lipolysis). Free fatty acids will undergo beta-oxidation and produce a large amount of acetyl CoA, which gets converted to ketone bodies (acetoacetate, acetone, beta-hydroxybutyrate, etc.)

Solution to Question 21:

The respiratory quotient, if someone is on an exclusive carbohydrate diet, is 1.

The respiratory quotient is the amount of CO_2 produced /O_2 consumed.

The respiratory quotient for various substances:

Substance	RQ
Carbohydrate	1.00
Protein	0.81
Fat	0.71
Alcohol	0.66

Biochemistry AIIMS 2019 (May & Nov)

Question 1:

Child presents with hypotonia and seizures. It was confirmed to be Zellweger syndrome. Which among the following accumulates in brain in this condition?

a) Glucose

b) Lactic acid

c) Long chain fatty acid

d) Triglycerides

Question 2:

Which of the following does not happen in 5' to 3' direction?

a) Transcription

b) DNA replication

c) DNA repair

d) RNA editing

Question 3:

Chimeric DNA is associated with:

a) Paternity test

b) Maternity test

c) Personal identification

d) Organ transplantation

Question 4:

A bodybuilder starts eating raw eggs for protein. He starts to develop fatigue on moderate exercise. The doctor prescribes a vitamin. Which enzyme is deficient in him?

a) Glucose 6 Phosphate

b) Pyruvate Carboxylase

c) PEPCK

d) Glycogen Phosphorylase

Question 5:

Which of the following shows functional assessment of B1 deficiency?

a) RBC Transketolase

b) RBC Glutathione reductase

c) Serum Thiamine levels

d) RBC Glutathione Peroxidase

Question 6:

Which modification will lead to a change in the biological activity of the hormone given in the picture?



a) Hydrolysis of disulphide bonds

b) Change in 5-6 in A chain

c) Change in 29-30 in B chain

d) Change in 14-16 in A chain

Question 7:

The graph shown below is the titration curve of a biochemical compound. Which of the following statement is true?



a) The maximum buffering capacity of the compound is represented by points A and B

b) The points A and B represent the range of maximum ionization of the amine and carboxyl group

c) The compound has three ionisable side chains

d) The compound has one ionisable group

Question 8:

Which of the following TCA cycle intermediate is a part of heme metabolism?

a) Alpha ketoglutarate

b) Fumarate

c) Succinyl CoA

d) Malate

Question 9:

The catalytic efficiency of enzyme is best expressed by which of the following kinetic constants:

a) Kcat/Km

b) Km/Kcat

c) Km/Ka

d) Ka/Km

Question 10:

Type of mutation seen in sickle cell anemia:

a) Insertion

b) Deletion

c) Point mutation

d) Frameshift mutations

Question 11:

By which mechanism does gene repair occur when mediated by CRISPR-Cas9?

a) Base excision

b) Non-homologous end joining

c) Homologous end joining

d) Mismatch repair

Question 12:

Which of the following is deficient in a patient with features as shown below:

a) Phenylalanine

b) Tryptophan

c) Glutamine

d) Arginine

Question 13:

The attachment of Eukaryotic mRNA to the ribosome is mediated through _____

a) Poly A tail

b) tRNA

c) Guanyl Cap

d) Shine-Dalgarno Sequence

Question 14:

Which of the following would have no effect on the function of the protein product?

a) Glutamine replaced by asparagine

b) Glutamine replaced by alanine

c) Glutamine replaced by glutamate

d) Glutamine replaced by arginine

Question 15:

Which of the following is not true about of proteoglycans?

a) Carries negative charge

b) Chondroitin sulfate is a proteoglylcan

c) Made of amino acids and sugars

d) Holds less water

Question 16:

Which of the following has protective as well as lubricating property?

a) Collagen

b) Mucin

c) Albumin

d) Immunoglobulin

Question 17:

In urea cycle disorders, which of the following can be used to reduce the levels of ammonia?

a) Glutamate

b) Phenylbutyrate

c) Isoleucine

d) L-carnitine

Question 18:

Restriction endonuclease will act on which of the following:

a) AAGAAG

b) GAGAGG

c) AAGCTT

d) TACGAG

Answer Key

Question No.	Correct Option
1	c
2	d
3	d
4	b
5	a
6	c
7	a
8	c
9	a
10	c

11	b
12	b
13	c
14	a
15	d
16	b
17	b
18	c

Detailed Explanations

Solution to Question 1:

Elevated levels of very long chain polyenoic acids have been found in the brains of patients with Zellweger syndrome.

Zellweger syndrome:

• Autosomal recessive

• Cerebro-hepato-renal syndrome: Impaired neuronal migration and myelination, hepatomegaly, renal cysts

• Reduction or absence of functional peroxisomes in tissues.

• Accumulation of very long chain polyenoic acids in the brain.

• Severe neurological symptoms.

• Most patients die within the 1st year of life

Solution to Question 2:

RNA editing occurs in 3’ to 5’ direction. It is the process by which the coding sequence is changed at the mRNA level before translation.

RNA editing occurs by insertion of additional Us at the 3’ end of the mRNA and proceeds in a 3’ to 5’ direction. The guide RNA (gRNA) acts as a template for the editing process. An example of RNA editing is apoB gene and mRNA shown in the image below:



Solution to Question 3:

Chimeric DNA is associated with organ transplantation, especially bone-marrow transplantation.

In organ transplantation, chimeric DNA refers to the mixture of donor and recipient genetic material that occurs when donor cells begin functioning within the recipient's body. It is used for long-term monitoring of bone marrow transplant recipients. Detecting changes in the levels of donor-derived cells can provide early warning signs of graft failure, rejection, or relapse of underlying disease.

Note: The practical applications of recombinant DNA technology include:

• Gene mapping using techniques such as fluorescence in situ hybridization (FISH).

• Producing proteins such as:

• Interferons, tPA (tissue plasminogen activator)

• Insulin, growth hormone

• Hepatitis B vaccine

• Molecular analysis of disease, to detect gene variations and mutations.

• Restriction Fragment Length polymorphism (RFLP) and single nucleotide polymorphism (SNP)

• Microsatellite DNA polymorphisms

• Gene therapy and bone marrow transplantation

Solution to Question 4:

The patient developed pyruvate carboxylase enzyme deficiency.

• Biotin serves as co-enzyme for the synthesis of several carboxylase enzymes such as:

• Acetyl-CoA carboxylase

• Pyruvate carboxylase

• Propionyl CoA carboxylase

• Methylcrotonyl-CoA carboxylase

• Individuals who eat an excess of uncooked egg-white, develops biotin deficiency and as a result deficiency of the carboxylase enzymes. This is because, a protein called, avidin present in the egg-white bind to biotin in the diet, and makes biotin unavailable for intestinal absorption.

Solution to Question 5:

Erythrocyte (RBC) transketolase test is the most widely used method for the functional assessment of thiamine deficiency.

Thiaminediphosphate is a co-enzyme for transketolase. Therefore, in thiamine (Vitamin B1) deficiency, the RBC transketolase activity is reduced. When thiamine diphosphate is added in-vitro to erythrocyte lysate, activation of apo-transketolase is suggestive of B1 deficiency.

Solution to Question 6:

The hormone shown in the above image is insulin. Among the given options, changes in 29-30 amino acids in the B chain cause the generation of short and long-acting insulins depending on the modification.

Structure:

The primary structure of insulin consists of 2 polypeptide chains: A-chain(21 Amino acids) and B-chain(30 amino acids).

It has two interchain disulfide bonds,

• Between 7th amino acid of A chain and 7th amino acid of B chain

• Between 20th amino acid of A chain and 19th amino acid of B chain

One intrachain disulfide bond is present between the 6th amino acid of A-chain and 11th amino acid in A-chain.

Modification:

Various short and long-acting insulin analogs have been developed by modifying these amino acid residues.

• Insulin lispro: Proline at position 28 and lysine at position 29 in the B-chain are interchanged.

• Insulin Aspart: Proline at position 28 in the B-chain is replaced by aspartic acid.

• Insulin glulisine: Asparagine at position 3 and lysine at position 29 in the B-chain are replaced by lysine and glutamic acid.

• Insulin glargine: Asparagine at position 21 in the A-chain is replaced by glycine and the B-chain contains two extra amino acids(two arginine residues) at positions 31 and 32.

• Insulin detemir: Threonine at position 30 of the B-chain is removed and a 14-carbon fatty acid chain is added to position 29 of the B-chain.



Solution to Question 7:

The maximum buffering capacity of the compound is represented by points A and B, which corresponds to the pK1 and pK2 of the compound.

Other options:

• The points A and B represent the pKa of carboxyl and amino groups respectively.

• The amino acid has no ionizable side chains as the graph shows only two pKas.

• The amino acid has two ionizable groups: an amino group and a carboxyl group.

The above graph resembles the titration curve of the amino acid glycine.

Titration curve of Glycine:



The titration is completed at a pH of 12, where the glycine predominantly exists in the form of H2N - CH2 - COO-.

pK1 and pK2 correspond to the point where the buffering capacity is maximum.

Characteristics of the titration curve:

• Buffering capacity is referred to as the ability of the aminoacid to resist the change in pH due to the addition of acid (H+) or base (OH- equivalents), by accepting or losing the protons respectively.

• An ionizable group refers to the group which can either lose or accept a proton and becomes negatively charged or positively charged ion respectively.

• During the titration, the buffering capacity of the compound reaches its maximum, at points where the pKa of the ionizable groups is reached.

• Each of these pKa are labeled as pK1, pK2, pK3, etc.

• The presence of two pKas indicates that the compound has two ionizable groups. Ex. Glycine.

• The presence of three or more pKas indicates the presence of side chains on the amino acid, with additional ionizable groups.

• Ex. histidine has three ionizable groups, an amino group, an imidazole group (on the side chain), and a carboxyl group. The pKas of these groups respectively correspond to the pK1, pK2, pK3 of the titration curve of histidine.

Solution to Question 8:

Succinyl Co-A, an essential part for heme biosynthesis, is an intermediate of the TCA cycle.

The synthesis of heme begins by condensation of succinyl Co-A and glycine to form α-amino-β-ketoadipate, which is then converted to δ-aminolevulinate. These steps are mediated by the enzyme ALA synthase in the mitochrondria.

Various steps involved in the synthesis of heme is illustrated in the image given below:



Solution to Question 9:

The catalytic efficiency of an enzyme is expressed as Kcat/Km.

Kcat is the rate constant, which is the limiting rate of an enzyme-catalyzed reaction at saturation. It is given by Vmax divided by number of active sites of the enzyme. Km is the Michaelis-Menten constant, which is defined as the substrate concentration at which the initial velocity(Vi) is half the maximum velocity (Vmax/2).

For some enzymes, when a substrate (S) binds to the active site of the enzyme (E), and forms an enzyme-substrate complex (ES), the product is instantaneously formed. For such enzymes, the reaction of E+S→ES becomes the rate-limiting step. Its rate is determined on the rate of diffusion of the molecules in a solution. Hence, these enzymes are called diffusion-limited or catalytically perfect.

Examples of the catalytically perfect enzymes, where the catalytic efficiency reaches the diffusion limit of 108-109 M−1s−1:

• Triosephosphate isomerase

• Carbonic anhydrase

• Acetylcholinesterase

• Adenosine deaminase

Solution to Question 10:

Point mutation is seen in sickle cell anemia.

• Sickle cell anemia is caused by the mutation of a single base in the β-globin gene, where adenine is replaced with thymine in the sixth codon of the DNA sequence.

• This leads to change from GAG in the normal gene to GTG in the sickle cell gene, leading to a change in mRNA sequence from GAG to GUG.

• As a result, the mutant gene now codes for valine instead of glutamic acid.



Solution to Question 11:

Non-homologous end joining (NHEJ) is the major mechanism used by CRISPR-Cas9 for gene repair.

The CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats) are clustered palindromic sequences found in many bacteria. It is a form of acquired immunity to infection by bacteriophages which complements the system of restriction endonucleases and methylases. CRISPR uses RNA-based targeting to bring the Cas9 endonuclease to foreign (or any complementary) DNA. Within bacteria this CRISPR-RNA-Cas9 complex then degrades and inactivates the targeted DNA.

The CRISPR system has been adapted for use in eukaryotic cells for gene deletion, gene editing and even modulation of gene transcription.



Solution to Question 12:

The image provided is characteristic of Casal's necklace, seen in pellagra, and indicates a deficiency of Tryptophan.

Pellagra can occur due to:

• Vitamin B3 (niacin)

• Niacin is derived from tryptophan and needs B2 and B6 for its synthesis.

• Hartnup's disease (tryptophan malabsorption)

• Malignant carcinoid syndrome (↑ tryptophan metabolism)

• Isoniazid (↓ vitamin B6)

In Hartnup's disease, defective membrane transport of tryptophan across PCT and enterocytes leads to malabsorption of tryptophan. This causes a deficiency of tryptophan.

As a result, the production of niacin from tryptophan is reduced and this manifests as pellagra with dermatitis, diarrhea and dementia (3D's) and glossitis. A characteristic skin rash called the Casal’s necklace may also be seen.

Solution to Question 13:

The attachment of mRNA to the ribosome is mediated through the guanyl cap at the 5' end. Eukaryotic mRNAs have 7-methylguanosine caps at the 5' end and poly-A tails at the 3' end.

The 7-methylguanosine caps (Guanyl caps) are necessary for:

• Processing of primary transcripts to mRNA.

• Mediating the attachment of mRNA to the ribosome for translation.

• Protection of mRNA against nucleolytic attack by 5'exonucleases.

Other options:

Option A: AAUAAA is present at the 3’ end, which is the polyadenylation sequence (poly-A tail). They help stabilize the mRNA, facilitate its exit from the nucleus and aid in translation. After the mRNA enters the cytosol, the poly-A tail is gradually shortened.

Option B: tRNA molecules serve as adapters for the translation of the information in the sequence of nucleotides of the mRNA into specific amino acids.

Option D: Shine-Dalgarno sequence initiates translation in prokaryotes.

Note: The guanyl cap and poly-A tail aid in attaching the mRNA to the ribosome for translation, however, it is initiated and mediated first through the Guanyl cap. The poly(A) tail stimulates recruitment of the 40S ribosomal subunit to the mRNA, after which the ribosomal subunit binds to the guanyl cap at 5'end of mRNA.

Solution to Question 14:

When glutamine is replaced by asparagine, there is no change in the function of the protein product.

This is due to the concept of conservative or homologous replacement of amino acids. Asparagine and glutamine are both uncharged polar (hydrophilic) amino acids. Thus, when glutamine is replaced with asparagine there is no change in the polarity or the charge of the peptide/protein.

Solution to Question 15:

Proteoglycans can hold large amounts of water, and not less.

Proteoglycans are made of a central protein core with O-linked glycosaminoglycan side-chains. They hold large quantities of water and occupy space.

They provide the ground substance of connective tissue.

Chondroitin sulfate is a glycosaminoglycan that is bound to a protein to form chondroitin sulfate proteoglycan. It is present in the extracellular matrix. Hyaluronic acid and heparin are examples of glycosaminoglycans.

All of the other statements regarding proteoglycans are true.

Solution to Question 16:

Among the given options, mucin has both a protective as well as a lubricating properties.

Solution to Question 17:

Phenylbutyrate can be used to reduce ammonia levels in patients with urea cycle disorders.

Phenylbutyrate diverts urea into an alternate pathway and reduces the amount of ammonia. Hence, acting as an ammonia scavenger. Other drugs that can be used to achieve the same outcome (ammonia scavengers) include phenylacetate and benzoate.

Note: Arginine is used as first-line treatment as it provides ornithine; a positive regulator of the urea cycle. Arginine also activates NAG synthase which increases NAG; a positive regulator of the urea cycle

Solution to Question 18:

Restriction endonucleases can act on AAGCTT.

Restriction endonucleases act and cleave at the site of a palindromic sequence. A palindromic sequence is a sequence that reads the same backward and forward. An example is listed below.

5' --AAGCTT-- 3'

3' --TTCGAA-- 5'

Biochemistry AIIMS 2020 (May and Nov INI-CET)

Question 1:

Dicoumarol inhibits the action of vitamin K by?

a) Suicide inhibition

b) Competitive inhibition

c) Non competitive inhibition

d) Uncompetitive inhibition

Question 2:

A TPN bag consists of 100g glucose, 30g amino acids, and 40g lipids. What is the amount of calories delivered?

a) 840 KCal

b) 880 KCal

c) 640 KCal

d) 680 KCal

Question 3:

What is true about the Hayflick limit?

a) Human cells can divide a finite number of times only

b) Free radical oxidation level is unrestricted

c) Telomerase activity is unrestricted

d) RNA polymerase can function for. a finite number of times only

Question 4:

Ammonia in the brain is detoxified to form _____

a) Urea

b) Glutamine

c) Ammonium ions

d) Alanine

Question 5:

The sequence of vitamin D synthesis is?

a) Liver- Skin- Kidney

b) Skin- Liver- Kidney

c) Skin- Kidney- Liver

d) Liver- Kidney- Skin

Question 6:

Glucose is stored in the form of glycogen because _____

a) It is compact

b) It has multiple reducing ends

c) It can be stored at multiple organs

d) It can provide glucose for about 1 week when needed

Question 7:

DNA methylation is not related to?

a) Mismatch repair

b) Chromatin Remodelling

c) DNA replication

d) Gene silencing

Question 8:

The enzyme that is not used in cholesterol synthesis?

a) Thiolase

b) HMG CoA reductase

c) HMG CoA synthase

d) HMG CoA lyase

Question 9:

The enzyme that is common for both glycolysis and gluconeogenesis is?

a) Pyruvate kinase

b) Phosphofructokinase

c) Phosphoenolpyruvate carboxykinase

d) Phosphoglycerate kinase

Question 10:

The given chemical structure is of:



a) Keratan sulfate

b) Chondroitin sulfate

c) Heparin

d) Dermatan sulfate

Question 11:

The correct order of steps in Polymerase Chain Reaction (PCR) is:

a) Hybridisation, Annealing, Elongation, Denaturation

b) Elongation, Denaturation, Hybridisation, Annealing

c) Denaturation, Annealing, Extension

d) Annealing, Extension, Denaturation

Question 12:

Which of the following is not a true statement regarding telomerase?

a) Has reverse transcriptase activity

b) Present only in eukaryotes

c) Maintain the chromosome length

d) Involved in DNA repair

Question 13:

The illustration below shows alternative splicing of a gene which has intron segments 1, 2, 3, 4, and 5. It yields two products A and B, by the action of different promoters. You are provided with the promoters w, x, y, and z, each of which binds to a different segment as shown below and acts in the marked direction. Which of the following promoters will you use to get product A?



a) W,X,Y

b) W,Z

c) W,X

d) W, Y

Question 14:

Gibbs free energy curve of an enzyme-catalyzed reaction is given by



©MARROW

a) B

b) A

c) C

d) D

Question 15:

The maximum thermic effect is seen with a meal which has

a) Protein

b) Fat

c) Carbohydrates

d) Mixed meal having all 3 in equal amount

Question 16:

DNA methylation is not involved in?

a) 1 and 2

b) 2 and 3

c) 1 and 4

d) 2 and 4

Question 17:

Which of the following enzymes has the highest catalytic efficiency?

a) Km of an enzyme = 10 micromole and Kcat = 20 per sec

b) Km of an enzyme = 2000 nanomole and Kcat = 50 per sec

c) Km of an enzyme = 2 micromole and Kcat = 200 per sec

d) Km of an enzyme = 4 micromole and Kcat = 200 per sec

Question 18:

Which of the following are vitamin-B12 dependent enzymes?

a) 1, 2 , 3 and 4

b) 1 and 4 only

c) 1 and 3 only

d) 1, 2, and 3

Question 19:

Which of the following enzymes are involved in gluconeogenesis? A. Pyruvate carboxylase B. Phosphoenol pyruvate kinase C. Hexokinase D. Pyruvate kinase E. Glucose 6 phosphatase

a) A,B,and E only

b) B and C only

c) A and E only

d) A, B and C only

Question 20:

Ketosis is common in type 1 DM due to:

a) 2,3 and 4

b) 1,2,3 and 4

c) 4 only

d) 1 and 2 only

Answer Key

Question No.	Correct Option
1	b
2	b
3	a
4	b
5	b
6	a
7	c
8	d
9	d
10	b
11	c
12	d
13	d
14	b
15	a
16	d
17	c
18	b
19	c
20	a

Detailed Explanations

Solution to Question 1:

Dicoumarol inhibits the action of vitamin K by competitive inhibition.

Dicoumarol and warfarin competitively inhibit the gamma-carboxylation system due to structural similarity with vitamin K. Therefore, they can act as anticoagulants used for therapeutic purposes.

Vitamin K epoxide is reduced to the quinone form by epoxide reductase, which is sensitive to inhibition by warfarin and dicoumarol.



Solution to Question 2:

The amount of calories delivered by the given TPN (Total Parenteral Nutrition) bag is 880 KCal.

The caloric value of different food components are:

• Carbohydrate (glucose) - 4KCal/g

• Proteins (amino acids) - 4.2KCal/g = ~4KCal/g

• Fat (lipids) - 9Kcal/g

Calculation of the number of calories:

• 100 gm of glucose = 100 x 4KCal=400 KCal

• 30 gm of amino acids = 30 x 4 KCal= 120 KCal

• 40 gm of lipids = 40 x 9 KCal= 360 KCal

Total amount of calories delivered= 400+120+360=880 Kcal

Note: The caloric value of alcohol is 7KCal/g.

Solution to Question 3:

The Hayflick limit says that human cells can divide for a finite number of times.

The Hayflick limit is the number of divisions a cell population can go through in culture before division stops and enters a state known as senescence. With the exception of stem cells, all the other human cells have only a limited capacity for cell division. This biological clock is of great help in the study of the normal ageing process.

The number of cell divisions of the cultured cell is roughly related to:

• The age of the cell donors

• The longevity of the species

For example, fibroblasts from the human embryos, when sustained in tissue culture, divide about 50 times before they die. Those taken from person after birth divide only 20 to 30 times.

Solution to Question 4:

Ammonia in the brain is detoxified to form glutamine.

Glutamine synthetase combines ammonia with glutamate to form glutamine. It requires the hydrolysis of ATP.

Deficiency of glutamine synthetase in neonates results in severe brain damage, multiorgan failure, and death.

Note: Glutamine is the transport form of ammonia from the brain and intestine to the liver; while alanine is the transport form from the muscle.

Solution to Question 5:

The sequence of vitamin D synthesis is skin-liver-kidney.

The sequence of vitamin D synthesis:

Step 1: In the skin: 7-Dehydrocholesterol (an intermediate in cholesterol synthesis that accumulates in the skin) undergoes a nonenzymic reaction on exposure to ultraviolet light, yielding cholecalciferol, which is absorbed into the bloodstream.

Step 2: In the liver: Cholecalciferol is hydroxylated to 25-hydroxycholecalciferol (calcidiol). This is released into the circulation bound to a vitamin D–binding globulin, which is the main storage form of the vitamin.

Step 3: In the kidney: 25-hydroxycholecalciferol is converted to either 1,25 dihydroxycholecalciferol (calcitriol), the biologically active form by the enzyme 1α-hydroxylase.



Solution to Question 6:

Among the given options, the most appropriate answer would be option A i.e Glucose is stored in the form of glycogen because it is compact.

Glycogen is the major storage carbohydrate in animals, corresponding to starch in plants. It is a branched polymer of α-d-glucose.

Other options:

The major sites of storage are liver and muscle, that is, it can't be stored at multiple sites.

It does not have multiple reducing ends. It instead has multiple non-reducing ends.

Liver glycogen functions as a reserve to maintain the blood glucose concentration in the fasting state. After about 18 hours of fasting, most of the liver glycogen is depleted and hence it can not provide glucose for up to 1 week.

Solution to Question 7:

DNA methylation is not related to DNA replication as DNA methylation is stable during replication and cell cycle arrest.

Chromatin remodeling by methylation/histone acetylation is an important aspect of transcription regulation.

Methylation of genes also plays an important part in gene silencing. Ex:- Promoter hypermethylation in tumor suppressor genes contributes to the development of cancers.

The process of mismatch repair relies upon a DNA-methylation excision signal, especially in E. coli.

Solution to Question 8:

Among the given options, the enzyme HMG CoA lyase is not used in cholesterol synthesis. It is a mitochondrial enzyme involved in the breakdown of proteins and fats.

Cholesterol is a 27-carbon compound consisting of four rings and a side chain. It is synthesized from acetyl-CoA.

The major sites of synthesis of cholesterol are the liver, adrenal cortex, testes, ovaries, and intestine. All nucleated cells can synthesize cholesterol, including arterial walls. The enzymes involved in the synthesis of cholesterol are partly located in the endoplasmic reticulum and partly in the cytoplasm.

Initially, two molecules of acetyl-CoA condense to form acetoacetyl-CoA catalyzed by cytosolic thiolase. This acetoacetyl- CoA enters the cholesterol synthesis pathway along with a molecule of acetyl-CoA.

The image below shows the enzymes involved in cholesterol synthesis:



Solution to Question 9:

The enzyme phosphoglycerate kinase is common for both glycolysis and gluconeogenesis. Phosphoglycerate kinase catalyzes the conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, which is a reversible step involved in both the processes.

Glycolysis and gluconeogenesis are processes that are reverse of each other. All the reversible steps of these two processes are catalyzed by the same enzymes, therefore are common to both.

The irreversible steps of glycolysis are catalyzed by the following enzymes:

• Hexokinase

• Phosphofructokinase

• Pyruvate kinase

The irreversible steps of gluconeogenesis are catalyzed by the following enzymes:

• Pyruvate carboxylase

• Phosphoenolpyruvate carboxykinase

• Fructose-1,6-bisphosphatase

• Glucose-6-phosphatase



Solution to Question 10:

The given chemical structure is of chondroitin sulfate.

Chondroitin sulfate is composed of repeating units of N-acetyl galactosamine with sulfate and glucuronic acid. Some important features of chondroitin sulfate are as follows:

• The most abundant GAG in the body.

• Location: Cartilage, CNS, bone

• They are located at sites of calcification in endochondral bone and are a major component of cartilage.

• They are found in high amounts in the ECM of the central nervous system.

Solution to Question 11:

The correct sequence of events for PCR among the given options is denaturation, annealing, and extension.

The process of PCR is as follows:

The DNA sample is first heat-denatured at >90°C (90-96 degrees Celsius for 3 minutes) to separate the two strands of the template DNA containing the target sequence; the primers (which attach to the 3' end of the flanking sequence) are added in excess, are allowed to anneal to the DNA (typically at 50-75°C) in order to generate the required template-primer hybridization complex. Under suitable conditions and presence of deoxynucleotides, elongation of the DNA stand occurs.

The number of copies produced in the presence of an unlimited supply of deoxynucleotides and enzymes is given by - 2 to the power n (where n is the number of PCR cycles)

• 20 cycles provide an amplification of 220 = 106 or 1 million copies

• 30 cycles provides an amplification of 230 =109 or 1 billion copies

Below is an image showing the various ingredients for carrying out PCR.



Solution to Question 12:

The false statement is that telomerase is involved in DNA repair.

Telomeres are the end of chromosomal DNA. Since prokaryotic DNA is circular, it does not have free ends and hence no telomeres. Thus, telomeres are present only in eukaryotes Telomeres have TG- rich repeats and usually do not code for any proteins. With every DNA replication during cell division, telomere shortening occurs. Once telomeres are almost negligible or non-existent in length, the DNA is unable to replicate and the cell dies. Hence, there is a limit to the number of times a cell can divide.

In stem cells and cancer cells, an enzyme telomerase having reverse transcriptase activity (RNA dependent DNA polymerase) adds telomeres to the chromosome ends, thereby preserving chromosomal length. It allows the stem cells and cancer cells to multiply a lot more number of times than a normal adult cell.

Solution to Question 13:

Among the given options, using primers W and Y will yield the desired product A.

Promoters or DNA regulatory elements usually precedes the desired exon to be transcribed. It helps redirect transcription of only the desired exons of a gene to yield a product. Alternative splicing utilizes different promoters to code slightly different products from the same gene.

In this question, the desired product A has segments 1,3,4,5 whilst the other product B has 1,2,4,5. The difference between both is the presence of seg 3 in product A and the presence of seg 2 in product B. As the product in question is A, only seg 3 is required while excluding seg 2 and ensuring seg 1,4 and 5 intact.

• Seg 1- Needs to be transcribed and so W is essential

• Seg 2- Does not need to be transcribed.

• Seg 3- Needs to be transcribed and so Y is essential

• Seg 4- Needs to be transcribed but Z is not essential as Y continues the transcription in the direction shown by the arrow.

The answer is hence promoter W and Y.

Solution to Question 14:

Among the given graphs, the Gibbs free energy change of an enzyme-catalyzed reaction is shown by graph A (Option B).

Graph B (Option A) cannot represent the Gibbs free energy of an enzyme-catalyzed reaction, as the energy state of an intermediate cannot be lower than the product. A lower energy state of the intermediate implies that the enzyme-intermediate complex is stable and such a complex will usually not further form a product.

During any chemical reaction, the product is formed by the breaking of substrate bonds and the conformational change of the substrate into a highly unstable transition state. This transition state is denoted by the peak of the graph. The Gibbs free energy curve of an enzyme-catalyzed reaction (with no intermediate state) is shown in the image given below:



The below graph shows the normal intermediate formation with an energy state still higher than that of the product:



All the reactions have a transition state, but not all reactions have an intermediate state. Every intermediate state is again always followed by another transition state.

Activation energy is the energy difference between the free energy of the substrates and that of the transition state. Because of the high activation energy, the rates of uncatalyzed chemical reactions are often slow.



In an enzyme-catalyzed reaction, the activation energy is reduced by forming a relatively stable intermediate-enzyme complex. Enzymes provide attachment sites for the substrate and speed up the rate of reaction. They don't change the energy state of either the substrate or the product. The Gibbs free energy change remains the same.

Graph C: Free energy change cannot be a straight line. Every reaction passes through a high-energy intermediate transition state.

Graph D: The activation energy of any reaction is never lower than the free energy of the product.

Solution to Question 15:

Among the given options, maximum thermic effect is seen with proteins

Specific dynamic allowance (previously called as thermic effect of food): This refers to the increased heat production or increased metabolic rate following the intake of food (diet-induced thermogenesis).

The values of SDA are:

• 30% for proteins: This means that out of every 100 grams of proteins consumed, the energy available for doing useful work is 30% less than the calculated value.

• 15% for lipids, and

• 5% for carbohydrates.

Solution to Question 16:

DNA methylation is not involved in the base pair excision repair mechanism (statement 4) and RNA splicing (statement 2).

DNA methylation is an epigenetic modification used as a gene expression regulation mechanism. It causes gene silencing by inactivating DNA and hindering the formation of euchromatin (transcriptionally active chromatin). Thus, it also aids in chromatin remodelling.

It however is not involved in the base pair repair mechanism or RNA splicing.



Solution to Question 17:

The highest catalytic efficiency is seen when Km= 2 micromole and Kcat = 200 per sec.

Catalytic efficiency is calculated as Kcat/Km, where Km is inversely proportional to the binding affinity of the enzyme and Kcat is the catalytic constant (Vmax divided by the total number of active sites of the enzyme present). Higher values of Kcat and lower values of Km gives better catalytic efficiency. Catalytic efficiency can be used to compare the performance of 2 different enzymes in catalyzing a reaction.

Step 1: Km of each option after converting respective units to moles will be 10, 2, 2 and 4 (each multiplied by 10-6) moles.

Step 2: The catalytic efficiency is calculated as Kcat/Km. On substituting the values in various options after converting to moles, the values come out to be 2, 25, 100 and 50 (each multiplied by 106) M-1 s-1.

Some enzymes can be so efficient that they convert the substrate to product instantaneously. They are limited only by the rate at which substrate diffuses towards the enzyme in a solution. These are called diffusion-limited enzymes and their catalytic efficiency is as high as 108 M-1 s-1.

Solution to Question 18:

Among the given enzymes, methyl malonyl CoA isomerase and methionine synthase are vitamin B12 dependent enzymes.

There are two vitamin B12-dependent enzymes:

• Methionine synthase (homocysteine methyltransferase): Catalyzes the conversion of homocysteine to methionine

• Methylmalonyl CoA mutase: methylmalonyl CoA, formed as an intermediate in the catabolism of valine, isoleucine, and propionyl CoA is converted to succinyl CoA

The below figure shows B12 serving as a cofactor of methionine synthase. Its deficiency may cause a folate trap.



Solution to Question 19:

Among the given options, A and E are correct. Pyruvate carboxylase and Glucose-6-phosphatase are enzymes used in gluconeogenesis.

Glycolysis and gluconeogenesis are 2 metabolic pathways in opposite directions (glucose being the substrate and product respectively). But a few reactions are not reversible in glycolysis. So a different pathway is used or a different enzyme is used by the body to make the reactions feasible.



Solution to Question 20:

2, 3 and 4 statements correctly explain the reason behind increased ketogenesis as seen in type1 DM.

Type 1 DM is characterized by a deficiency in insulin production. Since insulin is deficient, GLUT4 (insulin-dependent uptake) mediated glucose uptake is not possible in the heart or skeletal muscles. So adipose tissue mobilizes fat to be used as an energy source. These FFA (free fatty acids) are oxidized to acetoacetyl CoA which serves as a precursor for ketone body synthesis.

As the level of serum FFA is raised, proportionately more of the acetyl-CoA produced from their breakdown is converted to ketone bodies and less is oxidized via the citric acid cycle to CO_2. A fall in the concentration of oxaloacetate, particularly within the mitochondria, can impair the ability of the citric acid cycle to metabolize acetyl-CoA and divert fatty acid oxidation toward ketogenesis. Such a fall may occur because of an increase in the (NADH)/(NAD+) ratio caused when increased β-oxidation alters the equilibrium between oxaloacetate and malate so that the concentration of oxaloacetate is decreased.

Beta-hydroxy butyrate dehydrogenase is one of the enzymes used in ketone body synthesis. Increased rate of ketone body synthesis in insulin-deficient state implies an increased activity of beta-hydroxy butyrate dehydrogenase enzyme.

Biochemistry NEET 2018

Question 1:

Which of the following is responsible for the negative charge in fibrinopeptide A?

a) Glutamate and Valine

b) Histidine and Lysine

c) Aspartate and glutamate

d) Serine and Threonine

Question 2:

Serotonin is also known as?

a) 5 hydroxytryptamine

b) 5 hydroxytryptophan

c) 5 carboxytryptamine

d) 5 carboxytryptophan

Question 3:

Which of the following is true about DNA polymerase I?

a) It is involved in the creation of okazaki fragments.

b) It is found in prokaryotes.

c) It synthesises RNA primers.

d) It is the primary enzyme of DNA synthesis.

Question 4:

A patient presents with recurrent kidney stones. Microscopic examination finding of the urine specimen is shown below. Which of the following is not seen in the urine of this patient?

a) Ornithine

b) Lysine

c) Arginine

d) Histidine

Question 5:

In which condition is excessive urinary excretion of 5- hydroxy indole acetic acid seen?

a) Alkaptonuria

b) Albinism

c) Carcinoid syndrome

d) Phenylketonuria

Question 6:

Which of the following should be avoided by a patient of fish odour syndrome?

a) Biotin

b) Choline

c) Niacin

d) Pantothenic acid

Question 7:

Which of the following is true regarding an allosteric modifier?

a) Desaturates the enzyme

b) Causes the enzyme to work faster only.

c) Binds to the catalytic site

d) Participates in feedback regulation

Question 8:

Which of the following is the most common end-product of fatty acid synthesis pathway?

a) Arachidonic acid

b) Oleic acid

c) Palmitic acid

d) Acetyl CoA

Question 9:

Vanillyl mandelic acid is excreted in urine in which of the following conditions?

a) Pheochromocytoma

b) Alkaptonuria

c) Carcinoid syndrome

d) Diabetic ketoacidosis

Question 10:

Which of the following statements is true about vitamin K?

a) Deficiency of vitamin K leads to hypercoagulability.

b) It is a water soluble vitamin.

c) Prolonged use of antibiotics leads to vitamin K deficiency.

d) Menaquinone is the dietary source of vitamin K found in green vegetables.

Question 11:

What do chaperones assist in?

a) Ubiquitinization of proteins

b) Protein modification

c) Protein folding

d) Protein cleavage

Question 12:

Defective fumarylacetoacetate hydrolase enzyme is associated with _____.

a) Type 1 Tyrosinemia

b) Type 2 Tyrosinemia

c) Type 3 Tyrosinemia

d) Type 4 Tyrosinemia

Question 13:

A newborn baby refuses breast milk since the 2nd day of birth but accepts glucose water. He develops vomiting and severe jaundice by the 5th day. Benedict's test was positive for urine and blood glucose was low. The most likely cause is due to the deficiency of

a) Aldose reductase

b) Galactokinase

c) Galactose-1-phosphate uridyl transferase

d) UDP galactose-4-epimerase

Question 14:

A 45-year-old woman with hypertriglyceridemia is considered for niacin therapy. She is a known diabetic managed adequately with sugars well under control. Why should niacin be used cautiously in this patient?

a) It can cause hyperglycemia.

b) It can cause scleroderma which makes injecting insulin difficult.

c) It can increase the metabolism of oral hypoglyemic drugs.

d) It can cause hypoglycemia.

Question 15:

Which of the following can alter the gene expression by methylation and acetylation but not change the gene sequence?

a) Mutation

b) Epigenetics

c) Translocation

d) Inversion

Question 16:

The flipped LDH ratio implies _____.

a) LDH 1 > LDH 2

b) LDH2 > LDH1

c) LDH 2 > LDH 3

d) LDH 3 > LDH 2

Answer Key

Question No.	Correct Option
1	c
2	a
3	b
4	d
5	c
6	b
7	d
8	c
9	a
10	c
11	c
12	a
13	c
14	a

15	b
16	a

Detailed Explanations

Solution to Question 1:

Aspartate and glutamate are responsible for the negative charge in fibrinopeptide A and B of fibrinogen.

The negative charges contribute to the solubility of fibrinogen in plasma and importantly also serve to prevent aggregation by causing electrostatic repulsion between fibrinogen molecules.

Solution to Question 2:

Serotonin is also known as 5 hydroxy-tryptamine, and it is a monoamine neurotransmitter. Serotonin is produced by hydroxylation of tryptophan and subsequent decarboxylation. It is a potent vasoconstrictor and stimulator of smooth muscle contraction.

Clinical correlation:

In carcinoid syndrome, excessive levels of serotonin is produced which causes flushing, diarrhea, wheezing, damage to the tricuspid valve, etc. The levels of 5-OH indole acetic acid (serotonin metabolite) is elevated in the urine.

Solution to Question 3:

The true statement is that DNA polymerase I (pol I) is found in prokaryotes.

The prokaryotic polymerases are DNA polymerases I, II, and III. DNA polymerase I recognize a “nick” or break in the phosphate backbone, and then removes each RNA primer and fills the gaps with DNA. It does this through its 5'→3' exonuclease activity. It has DNA repair activity through 3'→5' exonuclease activity. DNA polymerase II carries out DNA proofreading and repair. DNA polymerase III is the primary enzyme for the synthesis of the leading strand of DNA.

Option A: DNA polymerase δ (eukaryotic polymerase) synthesizes Okazaki fragments which are short, newly synthesized DNA fragments that are formed on the lagging template strand during DNA replication. They are complementary to the lagging template strand.

Option C: Primase synthesizes RNA primers to start the replication.



Solution to Question 4:

The image shows hexagonal, translucent and white crystals suggestive of cystinuria. Histidine is not seen in the urine of this patient.

Cystinuria is an autosomal-recessive disease due to a defect in renal PCT & intestinal amino acid transporter that is essential for reabsorptive transport of Cystine and the dibasic amino acids Ornithine, Lysine, and Arginine (COLA).

Diagnosis of cystinuria is by the cyanide–nitroprusside test. Initial treatment is with adequate hydration, alkalization of the urine with citrate supplementation or acetazolamide, and dietary modification to reduce salt and protein intake. If this fails then patients are usually started on chelation therapy with penicillamine.

Solution to Question 5:

Excessive urinary excretion of 5-hydroxy indole acetic acid (5-HIAA) is seen in carcinoid syndrome.

Carcinoid syndrome is marked by symptoms of bronchospasm, cutaneous flushing, abdominal cramps, and diarrhea. It occurs due to excessive production of serotonin, a specialized product of tryptophan metabolism by tumors commonly in the small intestine, liver and lungs. It is then metabolized and excreted in urine as 5-HIAA.

Solution to Question 6:

A fishy odor is seen in trimethylaminuria. Consumption of food products with choline should be avoided in fish odor syndrome.

Fish odor syndrome (trimethylaminuria) is an autosomal recessive disease where TMA(trimethylamine) is not converted to the non-odorous TMAO (TMA-N-oxide) in the liver. This is due to the deficiency of flavin-containing mono oxygenase 3 (FMO3). TMA then builds up in the body and is released in sweat, urine, and breath, giving off a strong odor of rotting fish from the body.

Choline, lecithin, and TMAO are precursors of trimethylamine which are found in food products. These food products are eggs, liver, peanuts, meat, poultry, fish, dairy foods, pasta, and rice.

Solution to Question 7:

An allosteric modifier of an enzyme usually participates in feedback regulation.

Allosteric regulations are a natural example of control loops, such as feedback from downstream products or feedforward from upstream substrates. Allosteric regulation is also particularly important in the cell's ability to adjust enzyme activity.

An allosteric modifier or effector binds to the allosteric site to induce a conformational change in the catalytic or active site. Those modifiers or effectors that enhance the protein's activity are referred to as allosteric activators, whereas those that decrease the protein's activity are called allosteric inhibitors.

Some examples of allosteric modifiers		
Enzyme	Allosteric inhibitor	Allosteric activator
1. ALA synthase	Heme	-
2. Aspartate transcarbamoylase	CTP	ATP
3. HMG CoA reductase	Cholesterol	-
4.Phosphofructokinase	Citrate, ATP	AMP, F2,6P
5. Acetyl CoA carboxylase	Acyl CoA	Citrate
6. Citrate synthase	ATP	-
7. Carbamoyl phosphate synthase-I	-	NAG
8. Carbamoyl phosphate synthase-II	ATP	-



Solution to Question 8:

The most common end product of fatty acid synthesis is palmitic acid.

Palmitic acid is a 16 carbon saturated long chain fatty acid (LCFA). The substrate for its synthesis is acetyl CoA. Acetyl CoA is also the end product of fatty acid beta-oxidation.

Solution to Question 9:

• Vanillyl Mandelic Acid is the end product of catabolism of catecholamines and is excreted in urine in Pheochromocytoma.

• VMA is the urinary product of both epinephrine and norepinephrine.

• In pheochromocytoma and neuroblastoma, there is an excessive synthesis of catecholamines which causes enhanced synthesis of VMA and its excretion in the urine.

• It is a good screening test for pheochromocytoma and is also used to diagnose and follow up neuroblastoma and ganglioneuroma.

Solution to Question 10:

Prolonged use of antibiotics leads to vitamin K deficiency. This is due to the loss of normal bowel flora that synthesizes menaquinones (vitamin K2).

Vitamin K is a fat-soluble vitamin. The following are the vitamers of vitamin K:

• Phylloquinone (Vitamin K1) is the normal dietary source of Vitamin K that is found in green vegetables.

• Menaquinones (Vitamin K2) are synthesized by intestinal bacteria.

• Menadione (Vitamin K3) is a synthetic compound that is metabolized to phylloquinone.

Solution to Question 11:

Chaperones assist in protein folding.

Chaperones are heat shock proteins. They bind to the short sequences of hydrophobic amino acids of an extended polypeptide. Thus, they shield the new polypeptide being synthesized from aggregating. This provides the polypeptide an opportunity to form an appropriate secondary structure.

When a protein is thermodynamically trapped in a misfolded dead-end form, chaperones can rescue the protein by unfolding their hydrophobic ends. Thus, they provide the protein a second chance to fold productively.

Solution to Question 12:

Defective fumarylacetoacetate hydrolase enzyme is associated with type I tyrosinemia (tyrosinosis). It is the most common type of tyrosinemia.

There are three types of tyrosinemias:

• Type 1 (Hepatorenal tyrosinemia): Fumaryl acetoacetate hydrolase deficiency

• Type 2 (Oculocutaneous tyrosinemia/Richner-Hanhart syndrome): Tyrosine transaminase deficiency

• Type 3 (Neonatal tyrosinemia)

Solution to Question 13:

The above clinical features on ingestion of breastmilk at birth suggest classical galactosemia, a disorder of galactose metabolism. Galactose-1-phosphate uridyl transferase deficiency leads to classical galactosemia.

It is an autosomal recessive disorder. Clinical features include:

• Accumulation of galactose-1-phosphate in the liver causes liver damage & leads to hepatomegaly, vomiting, convulsions.

• Accumulation of galactose-1-phosphate depletes the liver of inorganic phosphate leading to diminished ATP synthesis which affects gluconeogenesis resulting in hypoglycemia.

• Accumulation of galactose-1-phosphate in the nerves causes intellectual disability.

• Accumulation of galactitol in the lens causes an oil-drop cataract.

Urinary reducing sugar (Benedict's test) and the Mucic acid test are positive.

Treatment is a galactose-free diet.

The flowchart given below depicts galactose metabolism.



Solution to Question 14:

Niacin should be used cautiously in diabetics as it can cause hyperglycemia.

Niacin induces insulin resistance and also decreases glucose-stimulated insulin secretion, which may result in severe hyperglycemia. When niacin is prescribed for patients with known or suspected diabetes, blood glucose levels should be monitored at least once a week.

Solution to Question 15:

Epigenetics is defined as heritable changes in gene expression that are not caused by alteration in gene sequence.

It means “above genetics” and refers to the fact that gene regulatory mechanisms do not change the underlying regulated DNA sequence, but rather simply the expression patterns of the DNA.

Epigenetic mechanisms play a key role in establishment, maintenance, and reversibility of transcriptional states. A key feature is that the controlled transcriptional on/off states can be maintained through multiple rounds of cell division. This observation indicates there must be robust mechanisms to maintain and stably propagate these epigenetic states.

Histone Methylation Mostly Makes DNA Mute.

Histone Acetylation: Relaxes DNA coiling, allowing for transcription. Histone Acetylation makes DNA Active.

Solution to Question 16:

The flipped lactate dehydrogenase (LDH) ratio is LDH 1 > LDH 2.

Under normal conditions, LDH 2 is predominantly present in the blood and LDH 1 is predominantly present in the heart. When cardiac tissue is damaged LDH 1 is released into the bloodstream.

LDH 1 > LDH 2 (flipped pattern) in the blood is suggestive of myocardial infarction.

LDH is a tetrameric enzyme with 4 subunits. It has four subunits with two isoforms: H isoform (heart) and M isoform (muscle)

LDH Type	Composition	Location
LDH-1	HHHH	Heart and RBC
LDH-2	HHHM	Heart and RBC
LDH-3	HHMM	Lungs, brain, lymphatic tissues, and platelets
LDH-4	HMMM	Liver and skeletal muscle
LDH-5	MMMM	Liver and skeletal muscle

Biochemistry NEET 2019

Question 1:

Which of the following diseases have autosomal recessive inheritance?

a) Albinism

b) G6PD deficiency

c) Marfan's syndrome

d) Lesch-Nyhan syndrome

Question 2:

Deficiency of which of the following causes increased glutamine in blood?

a) Arginosuccinate lyase

b) Alpha-galactosidase-A

c) Ornithine transcarbamoylase

d) Arginase

Question 3:

Which of the following amino acid is necessary for the conversion of nor-epinephrine to epinephrine

a) Tyrosine

b) Tryptophan

c) Phenyl alanine

d) Methionine

Question 4:

An infant presents with hypotonia and seizures. It was confirmed to be cerebrohepatorenal syndrome. Which of the following is accumulated in the brain in cerebrohepatorenal syndrome?

a) Glucose

b) Long chain fatty acid

c) Lactic acid

d) Triglycerides

Question 5:

Excess ammonia reduces which of the following substrates of the Krebs cycle?

a) Malate

b) Oxaloacetate

c) Fumarate

d) Alpha ketoglutarate

Question 6:

Which of the following is autosomal dominant?

a) Cystic fibrosis

b) Duchene muscular dystrophy

c) Sickle cell anemia

d) Achondroplasia

Question 7:

In Menke's disease, which of the following enzymes would be affected?

a) Lysyl hydroxylase

b) Lysyl oxidase

c) Prolyl oxidase

d) Prolyl hydroxylase

Question 8:

Common precursor amino acid for urea, creatine and nitric oxide is____.

a) Glycine

b) Arginine

c) Aspartate

d) Alanine

Question 9:

True regarding mitochondrial genome:

a) Maternal inheritance

b) Low rate of mutation

c) Single stranded DNA

d) Paternal inheritance

Question 10:

In a patient with lipoprotein lipase deficiency, which of the following is increased following a fatty meal?

a) Chylomicron

b) LDL

c) HDL

d) Apo A

Question 11:

In a patient with Wilson's disease which of the following is decreased in urine?

a) Tyrosine

b) Phosphorous

c) Serine

d) 3-methylhistidine

Question 12:

NADPH is used in _____.

a) Fatty acid synthesis

b) Ketone synthesis

c) Gluconeogenesis

d) Glycolysis

Answer Key

Question No.	Correct Option
1	a
2	c
3	d
4	b
5	d
6	d
7	b
8	b
9	a
10	a
11	d
12	a

Detailed Explanations

Solution to Question 1:

Albinism is inherited in an autosomal recessive pattern.

Solution to Question 2:

Deficiency ornithine transcarbamoylase causes increased glutamine in blood

Ornithine transcarbamoylase catalyzes the condensation of carbamoyl phosphate with ornithine to form citrulline in the urea cycle. Its deficiency leads to the accumulation of carbamoyl phosphate. The carbamoyl phosphate is then shunted to pyrimidine biosynthesis, resulting in high ammonia and orotic acid levels in the blood. This condition is thus called hyperammonemia type 2.

As a compensatory mechanism, ammonia is then combined with glutamate to form glutamine (by the enzyme glutamine synthetase). The levels of glutamine are elevated in the blood, cerebrospinal

fluid, and urine, probably as a result of this dysregulated glutamine synthesis.

Other options:

Option A: Deficiency of argininosuccinate lyase leads to argininosuccinic aciduria

Option C: Deficiency of alpha-galactosidase-A leads to Fabry's disease and an accumulation of globotriaosylceramide.

Option D: Deficiency of arginase leads to hyperargininemia

Solution to Question 3:

Methionine is necessary for the conversion of nor-epinephrine to epinephrine.

Solution to Question 4:

Cerebrohepatorenal syndrome or Zellweger's syndrome is characterized by the accumulation of very-long-chain fatty acid, abnormalities of synthesis of bile acids and a marked reduction of plasmalogens.

It occurs in individuals with a rare inherited absence of peroxisomes. It is due to mutations in genes encoding certain proteins called peroxins (involved in various steps of peroxisome biogenesis) in all tissues. They accumulate C26-C38 polyenoic acids in brain tissues and also exhibit a generalized loss of peroxisomal functions. It causes severe neurological symptoms and most patients die in the first year of life.

Solution to Question 5:

Excess ammonia results in depletion of levels of alpha-ketoglutarate.

This happens because excess ammonia reacts with alpha-ketoglutarate to form glutamate, mainly in the CNS.

Solution to Question 6:

Achondroplasia is an autosomal dominant disorder.

It is caused by a mutation of the fibroblast growth factor receptor-3 (FGFR-3) gene.

The primary defect is abnormal chondrocyte proliferation at the growth plate that causes development of short and thick bones.

Clinical features:

• Short limbs

• Normal trunk

• Large head

• Saddle nose

• Exaggerated lumbar lordosis

• Severe spinal deformity may lead to cord compression

Solution to Question 7:

Menke's disease is due to defective lysyl oxidase.

It is an X■linked recessive disorder caused by a mutation in ATP7A, which encodes for a copper-transporting ATPase■α polypeptide.

Solution to Question 8:

Arginine is the common precursor for urea, creatine and nitric oxide.

Solution to Question 9:

Mitochondrial genome is maternally inherited.

Mitochondrial inheritance:

• Inherited exclusively from the affected mother (ovum) as an individual's mitochondrial genome is entirely derived from the mother because sperms contain few mitochondria, which are typically shed upon fertilization.

• Both male and female children born to an affected mother will inherit the disease.

Pedigree shows mitochondrial inheritance.



Solution to Question 10:

After a fatty meal, chylomicrons are elevated in lipoprotein lipase deficiency.

Chylomicrons transport dietary lipids from the intestine to the peripheral tissues. They are synthesized in the intestine by assembling apo B-48 with lipids (mainly triacylglycerols). This is called a nascent chylomicron, which upon receiving apo C-II and apo E from HDL is remodelled into mature chylomicron.

Apo C-II activates lipoprotein lipase present on the endothelium. Lipoprotein lipase hydrolyzes the triacylglycerols into fatty acids and glycerol which are either used for storage or energy. Therefore, patients who are deficient in lipoprotein lipase or apo C-II (type I hyperlipoproteinemia) show increased levels of chylomicrons in the plasma after a fatty meal.

Solution to Question 11:

Urinary levels of 3-methylhistidine are decreased in patients with Wilson disease.

Amino acids in urine exist either in conjugated form (most) or in free form. In Wilson's disease the amount of conjugated amino acids far exceeds the amount of free amino acids. Methyl Histidine largely exists in free form and does not increase in amount on hydrolysis of acids in urine. Hence low amounts is seen in Wilson's disease.

Wilson disease is an autosomal-recessive disorder caused by mutation of the ATP7B gene located on chromosome 13. The ATP7B gene, encodes a transmembrane copper-transporting ATPase, expressed on the hepatocyte canalicular membrane.

Deficiency in the ATP7B protein causes a decrease in copper excretion into the bile, impairs its incorporation into ceruloplasmin, and inhibits ceruloplasmin secretion into the blood, leading to the accumulation of toxic levels of copper in the liver, brain, and eye.

Solution to Question 12:

NADPH is the source of reducing equivalent for fatty acid synthesis.

This NADPH is derived from 3 sources:

- HMP shunt pathway (main source)
- Extramitochondrial isocitrate dehydrogenase (minor source)
- Malic enzyme

Biochemistry NEET 2020

Question 1:

What is the function of the proteasome?

a) Protein folding

b) Post-translational modification

c) Protein degradation

d) Protein sorting

Question 2:

Replacing alanine by which amino acid, will increase UV absorbance of protein at 280nm wavelength?

a) Leucine

b) Proline

c) Arginine

d) Tryptophan

Question 3:

Vitamin B12 is found maximum in:

a) Animal products

b) Green leafy vegetables

c) Roots and tubers

d) Vegetarian diet

Question 4:

The process of apolipoprotein B48 synthesis involves which of the following?

a) RNA alternate splicing

b) RNA editing

c) DNA editing

d) RNA interference

Question 5:

A patient presents with a clinical scenario of obstructive jaundice. Which of the following results can be seen on Van den bergh's reaction?

a) Direct positive

b) Indirect positive

c) Both positive

d) Coproporphyrin

Question 6:

If the percentage of thymine residues in DNA is 28%. What is the percentage of cytosine?

a) 28%

b) 44%

c) 36%

d) 22%

Question 7:

Enzyme activated by a decrease in insulin: glucagon ratio is:

a) Glucokinase

b) Hexokinase

c) Phosphofructokinase

d) Glucose 6-phosphatase

Question 8:

A patient presented with symptoms of peripheral neuropathy and macrocytic anemia. She was given folate 5mg and the blood picture improved. However, the neurologic manifestations were aggravated. What is the most likely cause?

a) Malabsorption of folate

b) Treatment with folate unmasking pyridoxine deficiency

c) Deficiency of folate reductase in CNS

d) Folate therapy using the B12 stores

Question 9:

Which of the following is true regarding non-competitive inhibition?

a) Constant Km and increased Vmax

b) Constant Km and decreased Vmax

c) Decreased Km and decreased Vmax

d) Decreased Km and increased Vmax

Question 10:

A patient came to the hospital with complaint of abdominal pain. Routine investigations were unremarkable. On observation, the urine turned black on standing. Which of the following enzymes is likely to be defective in this patient?

a) Homogentisate oxidase

b) Xanthine oxidase

c) Phenylalanine hydroxylase

d) Dihydro oratate dehydrogenase

Question 11:

A 34-year-old male was on maize as a staple diet for a prolonged period. He presented with diarrhoea and lesion over his neck as shown in the image below. The condition occurs due to deficiency of:

a) Niacin

b) Zinc

c) Thiamine

d) Iron

Question 12:

Which vitamin, when infused in supraphysiological doses, causes macular edema and macular cysts?

a) Vitamin A

b) Vitamin D

c) Vitamin E

d) Vitamin B3

Question 13:

Which of the following is true about the structural organization of proteins?

a) Tertiary structure is three dimensional

b) Primary, secondary and tertiary structures are destroyed by denaturation

c) Secondary structure is stabilised by disulphide bonds

d) Secondary and tertiary structures depend upon amino acid sequence

Question 14:

Nitric oxide is synthesized from:

a) L - arginine

b) L - citrulline

c) Glycine

d) Lysine

Answer Key

Question No.	Correct Option
1	c
2	d
3	a
4	b
5	a
6	d
7	d
8	d
9	b
10	a
11	a
12	d
13	a
14	a

Detailed Explanations

Solution to Question 1:

The proteasome takes part in protein degradation.

The proteasome is a cylindrical structure in the cytosol that degrades polyubiquitinated proteins. It has a central core with proteolytic enzymes. Proteins that are tagged by ubiquitin to be degraded enter the core through regulatory rings on either side of the cylinder.



Clinical relevance: It plays an important role in antigen presentation to T lymphocytes as it degrades viruses and other molecules to small peptides.

Solution to Question 2:

Replacing alanine with tryptophan will increase UV absorbance of protein at 280nm wavelength

The amino acids which absorb 250-290nm (maximum at 280nm) UV light are Tryptophan (Maximum Absorption), Tyrosine, Phenylalanine.

Solution to Question 3:

Vitamin B12 is found maximum in animal products.

Vitamin B12 or cobalamin is synthesized only by microorganisms. Thus, the only source of cobalamin for humans is animal products. It is not found in plant sources. Therefore, individuals who are strict vegetarians are usually at the risk of developing vitamin B12 deficiency.

Solution to Question 4:

The process of apolipoprotein B48 synthesis involves RNA editing.

RNA editing is the process by which the coding sequence of the mRNA is changed before translation. As a result, the mRNA template differs from the DNA from which it was transcribed. This leads to the formation of a different protein. This occurs by insertion of additional uracils at the 3' end of the mRNA and proceeds in 3' to 5' direction.

RNA editing in apoB synthesis:

• In the liver, the apoB gene is transcribed to mRNA which is then translated to apoB 100.

• In the intestine, the same apoB gene is transcribed to the same mRNA. But now, the mRNA undergoes editing by an enzyme cytidine deaminase which converts the CAA codon in the mRNA to UAA at a single specific site. This edited mRNA translates to apoB 48.



Solution to Question 5:

In a patient with obstructive jaundice , van der bergh reaction gives direct positive result due to conjugated hyperbilirubinemia .

Van den Bergh reaction is used to determine the serum bilirubin level.

Van den Bergh reaction is a colorimetric method to measure serum bilirubin. It is based on the formation of red azodipyrroles when diazotised sulfanilic acid reacts with bilirubin.

Interpretation:

• In the absence of methanol, the water-soluble conjugated bilirubin reacts rapidly with the reagent and is called direct bilirubin.

• In the presence of methanol, both conjugated and unconjugated bilirubin react, providing the total bilirubin.

• Unconjugated (Indirect) bilirubin = Total-direct bilirubin

Solution to Question 6:

The percentage of cytosine in the DNA is 22%.

As per the Chargaff's rule, in the DNA molecules, the concentration of deoxyadenosine (A) equals that of thymidine (T), and the concentration of deoxyguanosine (G) equals that of deoxycytidine (C). Therefore, A=T and G=C.

In the question, the concentration of thymine is given as 28%, therefore the concentration of adenine is also 28%, both of which add up to 56%. The remaining 44% (100-56) of the DNA consists of G-C base pairs. Of which the guanine constitutes 22% and the cytosine constitutes another 22%.

Solution to Question 7:

Enzyme activated by a decrease in insulin: glucagon ratio is glucose 6-phosphatase.

A decrease in insulin: glucagon ratio is seen in fasting state, in which there is the activation of gluconeogenesis and inhibition of glycolysis.

Among the given options, only glucose 6-phosphatase is an enzyme of gluconeogenesis and therefore is activated. The enzymes mentioned as other options are irreversible enzymes of glycolysis and are inhibited in the fasting state.



Solution to Question 8:

The most likely cause is folate therapy which used up the B12 stores, aggravating the symptoms.

Administration of folic acid for patients with macrocytic anemia can partially reverse the hematologic abnormalities associated with vitamin B12 deficiency. However, the neurologic manifestations of vitamin B12 deficiency are not treated by folic acid.

Administration of folic acid to an individual with vitamin B12 deficiency can potentially mask untreated vitamin B12 deficiency or even worsen the neurologic complications. Hence, testing for and/or treatment of vitamin B12 deficiency is required in all patients being treated with folic acid.



Solution to Question 9:

Noncompetitive inhibition has constant Km and decreased Vmax.

Noncompetitive inhibition occurs when the inhibitor binds at a site different from the substrate-binding site. The structure of the inhibitor does not resemble that of the substrate and the inhibitor does not affect the binding of the substrate.



Solution to Question 10:

Urine turning black on exposure to air is suggestive of alkaptonuria which occurs due to the defect in homogentisate oxidase enzyme.

Alkaptonuria is a disorder of tyrosine metabolism, that develops due to the defect in homogentisate oxidase, which oxidizes homogentisate to maleylacetoacetate. The defect in this enzyme leads to the accumulation of homogentisate, which is then excreted in the urine. The excreted homogentisate gets oxidized when exposed to the air and causes the urine to turn black.

In some cases, the excessive homogentisate may get oxidized to benzoquinone acetate, which binds to the connective tissue. This leads to arthritis and pigmentation of the connective tissue, a condition called ochronosis.

Other options:

Option B: Xanthine oxidase deficiency results in hypouricemia, xanthinuria, and xanthine lithiasis.

Option C: Defects in phenylalanine hydroxylase can lead to phenylketonuria characterised by hyperreflexia, seizures, hypopigmentation, low IQ, hyperactivity, and tremors.

Option D: Defects in dihydroorotate dehydrogenase are rarely seen in clinical practice. Miller syndrome is associated with this defect.

Solution to Question 11:

The above history and clinical features suggest niacin/vitamin B3 deficiency.

Excessive consumption of maize-based diets can cause pellagra. Corn/Maize has low levels of niacin and tryptophan. Jowar has high leucine content which inhibits QPRTase, an enzyme in the synthesis of Niacin.

Niacin deficiency leads to a disease called pellagra which is characterized by

- Diarrhea
- Dermatitis
- Dementia

The lesion in the neck is characteristic Casal's necklace pattern, seen in pellagra. The Casal collar or Casal necklace is a clinical sign in which there is an erythematous pigmented skin rash in the distribution of a broad collar (dermatomes C3 and C4).

Solution to Question 12:

Macular cysts and edema may develop in patients due to the infusion of supraphysiological doses of vitamin B3 (niacin toxicity).

The toxic effects of niacin include:

• Flushing associated with skin dryness, itching, paresthesia, and headache is seen. These effects may be seen with even a low dose of 30mg/day. Pre-medication with aspirin alleviates these symptoms as these symptoms are mediated by prostaglandins.

• Macular edema and macular cysts

• Nausea, vomiting, and abdominal pain.

• Hepatic toxicity presents as jaundice with elevated AST and ALT levels.

• Fulminant hepatitis requiring liver transplantation.

• Glucose intolerance

• Hyperuricemia

Solution to Question 13:

The tertiary structure is three dimensional is the right answer among the given options.

Proteins are structurally organised into four levels, as follows:

• Primary: The linear sequence of amino acids in the polypeptide chain

• Secondary: Folded and geometrically ordered units of short segments of polypeptides. Eg: alpha helix, beta-pleated sheets, loops, and bends

• Tertiary: Secondary structures arranged into larger functional units called domains

• Quaternary: Different types of polypeptides arranged in a spatial manner

Other options:

Option B: Protein denaturation results in the unfolding and disorganization of the secondary and tertiary structures without the hydrolysis of peptide bonds, which form the primary structure.

Option C: Hydrogen bond and Van der Waal interactions are involved in the stabilization of secondary structure(helix). Disulfide bonds stabilize the tertiary structure of proteins.

Option D: Primary structure depends upon the amino acid sequence, not secondary or tertiary.



Solution to Question 14:

Nitric oxide is synthesized from arginine by a calcium-dependent reaction catalyzed by NO synthase.

The co-enzymes of NO synthase include FMN, FAD, Heme, and Tetrahydrobiopterin (BH4).


NADPH
NADP+
O2
Ca2+
NO synthase
+ NO
Arginine
Citrulline

Biochemistry INI-CET 2021

Question 1:

What is the effect of RNAi on gene expression?

a) Knock out

b) Knock down

c) Knock up

d) Knock in

Question 2:

Methyl-tetrahydro-folic acid gets accumulated in deficiency of which of the following?

a) B2

b) B12

c) B1

d) B6

Question 3:

Cytochrome oxidase is inhibited in all of the following poisonings except:

a) Hydrogen sulphide

b) Carbon monoxide

c) Methane

d) Cyanide

Question 4:

Which of the following molecular analysis methods is used to detect RNA ?

a) Sanger's method

b) G banding

c) Western blot

d) RT-PCR

Question 5:

Insulin inhibits which of the following lipase enzymes?

a) Lipoprotein lipase

b) Pancreatic lipase

c) Gastric lipase

d) Hormone sensitive lipase

Question 6:

Glycated hemoglobin (HbA1c) is best measured using?

a) Ion exchange chromatography

b) Affinity chromatography

c) Isoelectric focussing

d) Electrophoresis

Question 7:

What is the main substrate and metabolic fuel used by RBCs during fasting?

a) Ketone bodies

b) Glucose

c) Free fatty acids

d) Amino acids

Question 8:

miRNA binds to which part of the mRNA to inhibit translation?

a) Gene body

b) Gene promoter

c) 3' UTR

d) 5' UTR

Question 9:

The enzyme kinetics depicted in the Graph is suggestive of-



a) Allosteric Activation

b) Cooperative Binding

c) Michaelis Menten kinetics

d) Sigmoid Curve

Question 10:

Which of the following methods cannot be used to precipitate proteins?

a) Add alcohol and acetone

b) Using heavy metal ions

c) Moving pH away from isoelectric pH

d) Adding trichloroacetic acid

Question 11:

Which of the following diagnostic markers is/are incorrectly paired?

a) 1 only

b) 1 and 2

c) 1, 2 and 4

d) 2, 3 and 4

Question 12:

Identify the type of marked chromosome in the given karyotype.



a) Acrocentric

b) Metacentric

c) Submetacentric

d) Telocentric

Question 13:

Which of the following consumes the highest energy during assimilation?

a) Proteins

b) Carbohydrates

c) Mono-unsaturated fatty acids

d) Poly-unsaturated fatty acids

Question 14:

Which of the following is the best method to detect HbA1c?

a) Ion-exchange chromatography

b) Electrophoresis

c) Immunoassay

d) Affinity chromatography

Question 15:

Which of the following options is false in a patient with advanced liver disease?

a) Hyperammonemia

b) Decreased ATP synthesis

c) Oxidative decarboxylation of alpha-keto glutarate

d) Increased synthesis of glutamine and glutamate from alpha-ketoglutarate

Question 16:

Which of the following is associated with a defect in collagen maturity?

a) Scurvy

b) Rickets

c) Beri-beri

d) Night blindness

Question 17:

Which of the following is not a substrate for glucose formation?

a) Acetyl coenzyme A

b) Glycerol

c) Alanine

d) Lactate

Question 18:

Which of the following is the best test for quantitative analysis of an RNA sample?

a) Real time PCR

b) Reverse transcriptase PCR

c) PCR - Restriction Fragment Length Polymorphism

d) RTPCR followed by Real time PCR

Question 19:

Wobble hypothesis explains which of the following concepts of the genetic code?

a) Universal

b) Unambiguous

c) Degeneracy

d) Non-punctuated

Question 20:

Which of the following statements are true regarding mitochondrial DNA?

a) 1 and 3

b) 1 and 2

c) 2 and 4

d) 2 and 3

Question 21:

What is the repair mechanism associated with CRISPR-Cas9?

a) Non-homologous end joining

b) Base excision repair

c) Nucleotide excision repair

d) Mismatch repair

Question 22:

A patient with thiamine deficiency develops lactic acidosis. Which of the following enzymes is affected?

a) Pyruvate carboxylase

b) Pyruvate dehydrogenase

c) Phosphoenolpyruvate carboxykinase

d) Transketolase

Question 23:

Which of the following is not amphipathic?

a) Sphingolipids

b) Phosphoglycerol

c) Triglycerides

d) Glycolipids

Question 24:

Which of the following tests will be positive in the urine of a patient who has undergone prolonged starvation?



a) 1 and 3

b) Only 3

c) 1, 2 and 3

d) 1 and 2

Question 25:

The action of inhibition of acetylcholinesterase enzyme by organophosphorus compounds is through which of the following mechanisms?

a) Non-competitive inhibition

b) Reversible competitive inhibition

c) Irreversible competitive inhibition

d) Suicidal inhibition

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	d
5	d
6	a
7	b
8	c
9	c
10	c
11	a
12	a
13	a
14	a
15	c
16	a
17	a
18	d
19	c
20	b
21	a
22	b
23	c

24	b
25	c

Detailed Explanations

Solution to Question 1:

RNA interference(RNAi) is a biological process in which miRNA/siRNA molecules inhibit gene expression ("knockdown") by neutralizing the targeted mRNAs.

• miRNA regulates the expression of the target gene by post-transcriptional silencing.

• Small interfering RNAs (siRNAs) are short RNA sequences that can be introduced experimentally into cells and then serve as substrates for the Dicer enzyme and interact with the RNA-induced silencing complex (RISC) in a manner similar to that observed for endogenous miRNAs.

• Synthetic siRNAs targeted against specific mRNA species are useful in the study of gene function and this technique is referred to as knockdown technology.

• siRNAs are also being developed as possible therapeutic agents to silence pathogenic genes like oncogenes involved in neoplastic transformation.

Knock out: Mice generated through genetic manipulation of embryonic stem (ES) cells such that a specific genetic locus is targeted and rendered non-functional ("knocked-out"), either by insertion of an irrelevant DNA sequence or by deletion of the endogenous DNA sequence.

Knock-in: Mice generated through genetic manipulation of embryonic stem (ES) cells such that a specific genetic locus is altered either by one-for-one substitution of DNA sequence or by addition of sequence not found in the endogenous genetic locus.

Solution to Question 2:

Accumulation of methyl tetrahydro folic acid is seen in vitamin B12 deficiency. Vitamin B12 acts as a cofactor of methionine synthase, an enzyme involved in the regeneration of tetrahydrofolate from methyl-tetrahydrofolate. Hence, in vitamin B12 deficiency methyl-tetrahydrofolate accumulates leading to functional folate deficiency known as the folate trap.



The conversion of 5,10-methylene-THF into 5-methyl-THF (methyltetrahydrofolate) is an irreversible process. The major source of tetrahydrofolate for tissues is 5-methyl-THF. The only way to make further use of 5-methyl-THF and to maintain the folate cycle is by the vitamin B12-dependent remethylation of homocysteine to methionine by the methionine synthase enzyme.

Decreased activity of methionine synthase in vitamin B12 deficiency results in the accumulation of methyltetrahydrofolate, which cannot be used, resulting in a folate trap. The folate cycle acts as the link between the functions of folate and vitamin B12, and deficiency of vitamin B12 can cause a functional deficiency of folate.

Note: Since vitamin B12 is also linked to the regeneration of methionine from homocysteine, its deficiency can result in hyperhomocysteinemia as well. This in turn increases the risk of atherosclerosis, hypertension, and thrombosis.

Solution to Question 3:

Cytochrome oxidase is a hemoprotein, which is the terminal component of the respiratory chain carriers (complex IV) in mitochondria. The action of cytochrome oxidase is inhibited in carbon monoxide, cyanide, and hydrogen sulfide poisoning, resulting in prevention of cellular respiration.

Hemoproteins are metalloproteins that have a typical heme prosthetic group. They are present in myoglobin, hemoglobin, and other cytochromes. Cytochrome oxidase contains two molecules of heme, each having one Fe atom that oscillates between Fe^{3+} and Fe^{2+} during oxidation and reduction. It also contains two atoms of copper, each associated with a heme unit.

Complexes of ETC :

Complex I: NADH-CoQ oxidoreductase

Contains - FeS, FMN, Coenzyme Q

Inhibitors – Rotenone

Complex II: Succinate-CoQ reductase

Contains - FeS, Coenzyme Q

Inhibitor - Carboxin

Complex III: CoQH2-cytochrome c oxidoreductase

Contains - cytochrome c, b, C1, and FeS

Inhibitor - Antimycin A

Complex IV: cytochrome oxidase

Contains - Cytochrome a, a3, and Copper

Inhibitor - Cyanide, Azide, Carbon monoxide

Complex V (informally): ATP Synthase

Inhibitor - Oligomycin

Note: MELAS (mitochondrial encephalopathy, lactic acidosis, and stroke) is an inherited condition due to NADH-Q oxidoreductase (Complex I) or cytochrome oxidase (Complex IV) deficiency.

Solution to Question 4:

RT PCR(Reverse Transcriptase PCR) is a molecular analysis method used for the detection of RNA. It can be used to quantitate mRNA levels by generating complementary DNA (cDNA) copies of the mRNA using the reverse transcriptase enzyme.

The northern blot technique can also be used to detect RNAs. Here, the mixture of mRNA molecules is separated by electrophoresis and then transferred to a nitrocellulose membrane and hybridized with radiolabeled probes.

Sanger's technique:

• This was the first technique used to determine the sequence of proteins.

• Sanger's reagent is 1- Flouro 2,4 Dinitro benzene.

• The first protein to be sequenced by this method was insulin.

• Only dipeptides and tripeptides can be sequenced by this method.

G banding (Giemsa banding):

It is the most common technique used in cytogenetics to produce a visible karyotype by staining condensed chromosomes. It is useful for identifying genetic diseases through the photographic representation of the entire chromosome complement. The metaphase chromosomes are treated with trypsin (to partially digest the chromosome) and stained with Giemsa stain.

Heterochromatic regions, which tend to be rich with adenine and thymine (AT-rich) DNA and relatively gene-poor, stain darker and appear as dark bands in G-banding. In contrast, less condensed chromatin, which tends to be rich with guanine and cytosine (GC-rich) and more transcriptionally active incorporates less Giemsa stain, and these regions appear as light bands in G-banding. The image below shows a G-banding karyogram of a normal human male.

Western blot, also called immunoblots, is a technique used to study protein molecules. The proteins undergo electrophoresis and then are transferred to a membrane. These proteins are then tagged with the specific antibodies and visualized.

Solution to Question 5:

Insulin has an inhibitory effect on lipolytic hormones like hormone sensitive lipase (HSL). In adipose tissues, the stored triglycerides are broken down (lipolysis) by hormone sensitive lipase (HSL) into free fatty acids (FFAs). Insulin inhibits HSL and thus reduces the release of FFAs and also glycerol.

In diabetic patients, low insulin level corresponds to an increase in HSL activity, promoting the breakdown of lipids and release of FFAs from adipose tissue. Hormone-sensitive lipase activity is accelerated by hormones like epinephrine, norepinephrine, glucagon, ACTH, TSH, GH, ADH, and Melanocyte stimulating hormones (MSH).



Insulin induces the activity of lipoprotein lipase (LPL) which is present on the walls of capillaries. LPL promotes hydrolysis of TGs and degradation of VLDL and chylomicrons. In diabetes, low insulin leads to decreased LPL activity resulting in increased VLDL and TGs levels.

Effects of insulin on various tissues:

Well-fed state (Insulin)	During fasting (Glucagon)
Lipogenesis increased	Lipogenesis inhibited
Lipolysis inhibited	Lipolysis increased
Insulin inhibits HS-lipase	Glucagon activates HS-lipase
Lipoprotein lipase active	Lipoprotein lipase inactive

Tissue	Increases	Decreases
Adipose tissue	Increased glucose entryIncreased fatty acid synthesisIncreased glycerol phosphate synthesisIncreased triglycerides depositionActivation of lipoprotein lipaseIncreased K+ uptake	Inhibition of hormone-sensitive lipase
Muscle	Increased glucose entryIncreased glycogen synthesisIncreased amino acid uptakeIncreased protein synthesis in ribosomeIncrease ketone uptakeIncrease K+ uptake	Decrease protein catabolism Decrease release of glucogenic amino acids
Liver	Increased protein synthesisIncreased lipid synthesisIncreased glycogen synthesis & glycolysis	Decrease ketogenesisDecreased glucose output due to decreased gluconeogenesis

Tissue	Increases	Decreases
General	Increased cell growth	

Solution to Question 6:

Glycated hemoglobin (HbA1c) is best measured using ion-exchange chromatography.

HbA1c is formed by non-enzymatic glycation of hemoglobin A present in RBCs. Normally this occurs to a small extent but is increased in diabetic patients with poor glycemic control. Since the half-life of RBCs is typically 60 days, HbA1c can be used to monitor mean blood glucose concentration over the past 6 to 8 weeks.

In ion-exchange chromatography:

• Proteins interact with the stationary phase by charge-charge interactions.

• Proteins with a net positive charge at a given pH will tightly adhere to beads with negatively charged functional groups such as carboxylates or sulfates (cation exchangers).

• Proteins with a net negative charge adhere to beads with positively charged functional groups, typically tertiary or quaternary amines (anion exchangers).

• Non-adherent proteins flow through the matrix and are washed away.

• Proteins elute in inverse order of the strength of their interactions with the stationary phase.

Solution to Question 7:

Glucose is the main substrate and metabolic fuel used by RBCs during fasting. RBCs lack mitochondria and are dependent fully on glycolysis (anaerobic) and pentose phosphate pathway during fasting state.

Glucose is always required by brain and RBCs. Brain can meet only about 20 percent of energy requirements by using ketone bodies and the remaining is supplied by glucose. Metabolic changes occuring during fasting state and starvation therefore serve to preserve glucose and glycogen reserves for brain and RBCs.

Solution to Question 8:

Micro-RNA (miRNA) is an endogenous non-coding RNA that typically binds to the 3'-untranslated regions (UTRs) of target messenger-RNAs (mRNAs). The primary function of miRNA is to modulate the translation of target mRNAs into their corresponding proteins. After the target RNA is transcribed from its gene (post-transcriptional), the miRNA may either induce mRNA cleavage or repress its translation. This is referred to as post-transcriptional gene silencing or RNA interference.

RNA interference(RNAi) is a biological process in which miRNA or siRNA molecules inhibit gene expression by neutralizing targeted mRNAs.The microRNA (miRNA) belongs to a group of non-coding genes. The miRNA genes are transcribed but not translated into proteins. They play a role in the regulation of the expression of the target gene by post-transcriptional silencing.

Mechanism:

• The miRNA gene present in the nucleus is transcribed to primary miRNA, which is further converted to pre-miRNA by an enzyme DROSHA intranuclearly and exported to the cytoplasm by an export protein called Exportin 5.

• The pre-miRNA undergoes trimming, by an enzyme called DICER, into mature single-stranded miRNA.

• The mature miRNA is associated with a multiprotein aggregate called RNA-induced silencing complex (RISC).

• The target mRNAs contain seed sequences in their 3′ untranslated region (UTR) that determines the specificity of miRNA binding and gene silencing.

Depending on the base pairing between the miRNA strand and its target mRNA, the RISC causes gene silencing either by mRNA cleavage (degradation or deadenylation) or by repressing its translation.

siRNA is similar in function to miRNA and is used in RNA interference.

miRNA expression has been found to be dysregulated in many cancers. Such misregulation is thought to have a causal role in the pathogenesis of such cancers. Some types of miRNAs are oncogenic (referred to as oncomiRs) and are overexpressed in cancerous tissue. On the other hand, tumor-suppressive miRNAs that counteract oncogenic characteristics are found to be underexpressed in cancers.



Solution to Question 9:

The graph shows a hyperbola which is suggestive of Michaelis - Menten Kinetics of the enzyme.

For enzymes showing michaelis menten kinetics, there is an increase in Rate of reaction, V, with increase in substrate concentration till a point Vmax is reached beyond which rate of reaction shows saturation.

Cooperative Binding would result in a sigmoid curve and is suggestive of Non Michaelis Menten Kinetics.

Solution to Question 10:

Precipitation of proteins is maximum at isoelectric pH and decreases as pH moves away from it. Hence, moving pH away from isoelectric pH cannot be used for precipitating proteins.

The isoelectric point or isoelectric pH (pI) is the pH at which the molecule carries no net charge.

Significance of Iso-electric pH (pI):

• The number of anions and cations present on the protein molecule will be equal and the net charge is zero.

• The proteins will not migrate in an electrical field

• Buffering capacity is minimum (note: maximum buffering action of an amino acid is seen when pH=pKa, and not at isoelectric pH)

• Viscosity will be minimum

• Precipitability will be maximum and solubility is minimum

Various precipitation reactions of proteins are:

• Salting out: higher the molecular weight of protein, the lesser the salt required. Globulins would require only half-saturated salt whereas albumin would need full saturation.

• Isoelectric precipitation: proteins are least soluble at their isoelectric pH and get precipitated immediately.

• Precipitation by organic solvents like alcohol and acetone.

• Precipitation by heavy metal ions: positively charged metal ions precipitate the negatively charged proteins in an alkaline medium. Salts of heavy metals can precipitate normal proteins of the GI tract in poisoning.

• Precipitation by alkaloidal reagents like tungstic acid, trichloroacetic acid, and picric acid.

• Heat coagulation is an irreversible precipitation process.

Solution to Question 11:

Beta glucocerebrosidase deficiency is seen in Gaucher's disease. It is an inborn error of metabolism due to the deficiency of the enzyme β-glucocerebrosidase (β-glucosidase) that leads to the accumulation of glucocerebroside in the lysosomes of cells.

• It is the most common lysosomal storage disorder. It is inherited in an autosomal recessive manner.

• The clinical features result from the accumulation of lipid-laden macrophages in the spleen, liver, bone marrow, bone, and other tissues.

• There are 3 clinical subtypes, delineated by the difference of neurologic manifestations: type 1, or the adult, non-neuronopathic form; type 2, the infantile or acute neuronopathic form; and type 3, the juvenile or subacute neuronopathic form

• Typical clinical features include delayed puberty, painless hepatomegaly and splenomegaly, liver cirrhosis, hypersplenism and pancytopenia, bleeding tendencies, bone pain due to avascular necrosis, osteoporosis, and pathological fractures.

• Neurological symptoms such as developmental delay, hypertonia, strabismus, and supranuclear gaze palsy (seen in type 3).

• Bone marrow examination usually reveals the presence of Gaucher cells having a characteristic wrinkled-paper appearance.

• Enzyme replacement therapy for Gaucher's disease includes imiglucerase, velaglucerase alfa, taliglucerase alfa, and eliglustat.

Von Gierke disease occurs due to deficiency of glucose-6-phosphatase enzyme in the liver, kidney, and intestinal mucosa. Patients present commonly by 3-4 months of age with hypoglycemia (leading to seizures), lactic acidosis, and hepatomegaly. Affected children often have a doll-like face with fat cheeks, relatively thin extremities, short stature, and a protuberant abdomen due to massive hepatomegaly. The kidneys are also enlarged, whereas spleen and heart are not involved. Laboratory findings include hypoglycemia, lactic acidosis, hyperuricemia, and hyperlipidemia.

Elevated serum lipase and serum amylase levels are associated with acute pancreatitis. Serum amylase levels do not correlate with the severity of pancreatitis and lack specificity whereas serum lipase is a more specific marker.

Low serum ceruloplasmin level is suggestive of hepatolenticular degeneration seen in Wilson's disease. However, it should not be used for definitive diagnosis as levels are normal in 10% of the affected patients and reduced in 20% of carriers. The diagnosis of Wilson's disease is based on the following:

• Increased 24-hour urinary excretion of copper (screening test)

• Decrease in serum ceruloplasmin

• Kayser–Fleischer rings under slit lamp

• Genetic testing for mutations in ATP7B gene

• Liver biopsy with quantitative copper assay (>200 μg/g dry weight of liver) is the gold standard investigation.

A higher level of LDH1 than LDH 2 in the blood is suggestive of myocardial infarction. Under normal conditions, LDH 2 is predominantly present in the blood and LDH 1 is predominantly present in the heart. When cardiac tissue is damaged LDH 1 is released into the bloodstream resulting in a high LDH1:LDH2 ratio known as the flipped LDH ratio (LDH1 > LDH2). LDH is a tetrameric enzyme with 4 subunits. It has four subunits with two isoforms: H isoform (Heart) and M isoform (Muscle)

• LDH-1: Heart

• LDH-2: Blood

• LDH-3: Brain, kidney, lungs, spleen, pancreas

• LDH-4: Liver and skeletal muscle

• LDH-5: Liver and skeletal muscle

Solution to Question 12:

The marked chromosome is chromosome 13 and the type is acrocentric.

Each chromatid has a long arm q and a short arm p. The two arms are separated by a centromere. In this image, the centromere is not midway between the two ends of the chromatids. It is seen towards one end with a marked difference in length between the p and q arms, hence it is an acrocentric chromosome.

Chromosomes are classified based on the position of the centromere as follows:

•

Metacentric: Centromere is centrally placed and the two arms are equal in length

•

Submetacentric: Centromere is slightly away from the center and the two arms are only slightly different in length

•

Acrocentric: Centromere is nearer to one end and the difference in length of arms is marked

•

Telocentric: Centromere lies at one end.

Group	Pairs of chromosomes	Features
A	1–3	Long, metacentric
B	4, 5	Long, submetacentric
C	6–12, X-chromosome	Medium, Submetacentric
D	13–15	Medium, acrocentric Satellite bodies +
E	16–18	Short, metacentric (16) or submetacentric (17 and 18)
F	19, 20	Short, metacentric
G	21, 22 + Y-chromosome	Short, acrocentricSatellite bodies +



Solution to Question 13:

Proteins consume the highest energy during assimilation.

The energy wasted in the assimilation of food is represented as diet-induced thermogenesis or specific dynamic action(SDA). Diet-induced thermogenesis (DIT) can be defined as the increase in energy expenditure above the basal metabolic rate and is expressed as a percentage.

The values for diet-induced thermogenesis are relatively higher protein and alcohol consumption and lower at a high-fat consumption, the hierarchy for it, being in the sequence of alcohol> protein> fat> carbohydrate. The value of DIT for protein is 30%, the lipid is 15% and carbohydrate is 5%. Hence, taking into account the loss of energy as DIT, additional 10% calories should be provided in a mixed diet.

Solution to Question 14:

Ion exchange chromatography is the best method to detect HbA1c.

Ion-exchange chromatography allows the separation of ionizable molecules according to differences in charge. It involves the use of a column which contains a matrix consisting of resin-bound charged groups.

If the charged group is anionic or negatively charged, it is called a cation exchanger and is used in cation exchange chromatography where proteins with a net positive charge migrate through the matrix more slowly than those with a net negative charge due to retardation of the migration of the former which is attracted to the negative charges bound to the resin. Cation exchange chromatography is used for negatively charged HbA1c estimation and is used as a diagnostic tool for diabetes mellitus.

Glycated hemoglobin, reported as HbA1c, has been used to monitor glycemic control in patients with diabetes for many years. HbA1c results from the attachment of glucose to the N-terminal

valine of the hemoglobin β-chain.

The concentration of HbA1c depends on both the concentration of glucose in the blood and the lifespan of the erythrocyte. Since the half-life of an RBC is 60 days, the level of glycated haemoglobin reflects the mean blood glucose concentration over the preceding 6 to 8 weeks. Normal glucose tolerance is characterized by an HbA1c < 5.7%; any value of HbA1c equal to or exceeding 6.5 % is diagnostic of the patient being diabetic. Also, ion exchange chromatography has the advantage of separating hemoglobin variants into peaks with the least interference by other hemoglobin variants; its disadvantage is that it is temperature dependant.

Other options:

Option B: Electrophoresis is an electrochemical process where macromolecules or colloidal particles depending on the electric charge migrate in a solution under the influence of an electric current. It can also be used for HbA1c estimation, its advantage being able to identify different variants of haemoglobin and disadvantage being interference of the process by fetal hemoglobin(HbF).

Option C: Immunoassays are used to quantify molecules based on the specificity and selectivity of antibody reagents generated. It can provide quantitative estimates of HbA1c but HbF produces falsely low results and hemoglobin variants such as HbS and HbC may cause a false increase in HbA1c measurements.

Option D: Affinity chromatography is a type of liquid chromatography. It involves the passing of a matrix through a solution that consists of the molecules of interest. These molecules would have a specific affinity for ligands present in the matrix and bind with them and end up getting separated from the solution. It can be used to measure HbA1c without being affected by temperature. High levels of HbF seen in thalassemia may interfere with its result.

Solution to Question 15:

Oxidative decarboxylation of alpha-ketoglutarate is not seen in a patient with advanced liver disease. Option C is false.

Normally, ammonia, a toxic by-product of protein catabolism, is excreted as urea upon being metabolised in the urea cycle, which takes place in the liver. In advanced liver disease, impaired ammonia metabolism leads to hyperammonemia. Ammonia withdraws α-ketoglutarate to form glutamate and glutamine. This leads to lowered concentrations of all citric acid cycle intermediates, and hence reduced generation of ATP.

The Krebs cycle involves a sequence of reactions occurring in mitochondria that oxidize the acetyl moiety of acetyl-CoA to CO_2 and reduces the coenzymes that are reoxidized through the electron transport chain. It is the final common pathway for the oxidation of carbohydrates, lipids, and protein also has a central role in gluconeogenesis, lipogenesis, and interconversion of amino acids. Although processes occur in many tissues, the liver is the only tissue in which all occur to a significant extent.

The reactions that occur in Kreb's cycle are shown in the image below.



One of the steps of the Krebs cycle involves α-ketoglutarate undergoing oxidative decarboxylation in a reaction catalyzed by α-ketoglutarate dehydrogenase, resulting in the formation of succinyl-CoA. High concentrations of ammonia inhibit α-ketoglutarate dehydrogenase, thereby preventing its oxidative decarboxylation.

Solution to Question 16:

Scurvy is associated with a defect in collagen maturity.

Scurvy results from ascorbic acid or vitamin C deficiency. Vitamin C is an essential cofactor for the enzyme prolyl hydroxylase, required for the hydroxylation of prolyl residues in collagen. This enables each collagen fibril to form a uniform and flexible triple helix configuration in the body's connective tissues.

Its deficiency results in a defective prolyl hydroxylase activity, thereby resulting in the impaired formation of mature connective tissue; the clinical manifestations of this deficiency include petechiae, ecchymosis, coiled hairs, inflamed and bleeding gums, joint effusion, tooth loss, bone fracture, poor wound healing, and fatigue which can be attributed to impaired collagen synthesis all of which are typical of scurvy.

Risk factors include chronic alcoholism, diseases such as celiac disease and Crohn's disease, associated with malabsorption and poor nutrition.

It is treated with vitamin C supplementation.

Other options:

Option B- Rickets is caused due to deficiency of vitamin D in children which is characterized by the continued formation of the collagen matrix of bone but its incomplete mineralization or demineralization. Children have soft, pliable bones.

Option C- Beriberi is caused due to deficiency of thiamine. It can either be dry beriberi characterized by peripheral neuropathy or wet beriberi characterized by edema due to dilated cardiomyopathy.

Option D- Night blindness is caused by a deficiency of vitamin A. Deficiency can also lead to pancreatic insufficiency, repeated gastrointestinal, pulmonary, and urinary infections, and xeropthalmia.

Solution to Question 17:

Acetyl coenzyme A (CoA) is not a substrate for glucose formation.

Acetyl CoA and compounds that give rise to acetyl CoA alone like acetoacetate, lysine, and leucine cannot give rise to a net synthesis of glucose. This is because of the irreversible nature of the pyruvate dehydrogenase complex, which converts pyruvate to acetyl CoA. These compounds give rise to ketone bodies instead.

The synthesis of glucose in the body, in the absence of dietary carbohydrates, occurs in two ways:

• From liver glycogen, by a process known as glycogenolysis in the absence of dietary carbohydrate, for up to 24 hours.

• De novo, from non-carbohydrate sources, by a process known as gluconeogenesis, when glycogen stores have depleted. This occurs primarily in the liver and to a lesser extent in the kidneys while fasting overnight or primarily in the kidneys during prolonged fasting of 48 hours or longer.

The substrate for glycogenolysis is glycogen and during this process, glycogen is first converted into glucose-1-phosphate by the enzyme glycogen phosphorylase. This is a rate-limiting step in the process. Glucose-1-phosphate is then converted to glucose-6-phosphate by the enzyme phosphoglucomutase and eventually, glucose is obtained from glucose-6-phosphate by the enzyme phosphatase. There is no involvement of acetyl CoA.

The steps with enzymes involved in glycogenolysis are shown in the image below.



In gluconeogenesis, the substrates can be:

• Oxaloacetate (from its conjugate acid, oxaloacetic acid) is a direct precursor of phosphoenolpyruvate

• Pyruvate, an α-keto acid that can be converted to glucose. Alanine is a glucogenic amino acid that is transaminated to form pyruvate by the enzyme alanine amino-transferase in the liver; the pyruvate formed, can be used for the synthesis of glucose.

• Lactate, obtained from anaerobic glycolysis, released from skeletal muscles and RBCs into the blood, to be taken up by the liver in the Cori cycle to be oxidised to pyruvate, to eventually synthesize glucose

• Glycerol from adipose tissue is converted to dihydroxyacetone phosphate in the liver, which itself is an intermediate of glycolysis and gluconeogenesis.

The steps and enzymes involved in gluconeogenesis are shown in blue in the flowchart below.



Solution to Question 18:

RTPCR followed by real-time PCR is the best test for quantitative analysis of an RNA sample.

RTPCR (reverse transcriptase PCR) involves the generation of complementary DNA (cDNA) copies of the mRNA using the reverse transcriptase enzyme followed by amplification of the cDNA copies by PCR technique.

Quantification of DNA involves the assessment of the amount/concentration of DNA that is present in a sample when carrying out a technique like PCR. Real-time PCR or quantitative PCR/qPCR involves the quantification of DNA after each cycle of amplification, or, in "real-time" than at the end of the procedure; therefore real-time PCR helps in quantitative analysis of DNA.

When reverse transcriptase PCR and qPCR are combined, they can be used to determine the relative concentrations of a particular mRNA molecule in a cell.

Other options:

Option A: Real-time PCR alone can only be used for the quantification of DNA only.

Option B: RTPCR can convert the RNA into cDNA by reverse transcriptase enzyme and then amplify the cDNA copies by means of PCR but cannot quantify the DNA/RNA.

Option C: PCR-Restriction fragment length polymorphism (PCR-RFLP) is also called cleaved amplified polymorphic sequence (CAPS) assay. Polymorphisms are inherited differences found among individuals in more than 1% of the normal population. PCR-RFLP is a technique in which organisms may be differentiated by variations in their DNA sequence by restriction endonucleases. Typically, gel electrophoresis is used to visualize RFLPs. Thus, DNA but not RNA molecules can be quantified in this process.

Solution to Question 19:

The wobble hypothesis explains the degeneracy of the genetic code.

The wobble hypothesis states that codon-anticodon pairing follows the traditional Watson–Crick rules (G pairs with C and A pairs with U) for the first two bases of the codon but can be less stringent for the third/last base. It has been observed that a tRNA can recognize more than one codon for a specific amino acid, especially in the last nucleotide of the codon triplet. This is referred to as degeneracy and its mechanism can be explained by the wobble hypothesis.

The first base of the anticodon moves to permit nontraditional pairing with the last base of the codon. This movement is called wobble and is possible because the first base of the anticodon is not as spatially defined as the other two bases. Due to the wobbling, 61 tRNA species are not required to read the 61 codons.

For example, three codons for glycine—GGU, GGC, and GGA—can form a base pair from one anticodon, 3′ CCI 5′ ie, I, inosine, can base pair with U, C, and A, as shown in the diagram below.

Other options:

Option A: The genetic code is universal because it is the same for all living organisms.

Option B: The genetic code is unambiguous because each codon is specific for only one amino acid. Example: GGA only codes for Glycine.

Option D: The genetic code is non-punctuated. This means that once the reading commences at a specific start codon, it is continuous and is read continuously as triplets until a stop codon is reached without stopping.



Solution to Question 20:

Statements 1 and 2 are true regarding mitochondrial DNA; it has a higher rate of mutations as compared to the nuclear genome and is a circular double-stranded DNA with about 16,000 base pairs.

Mitochondrial DNA is circular, double-stranded DNA composed of heavy and light chains and comprises about 1% of total cellular DNA. It contains about 16,569 base pairs and encodes about 13 protein subunits of the electron transport chain, of a total of about 67 protein subunits.

Its genetic code/ nucleotide sequence largely differs from that of the standard cellular code. For example, the standard stop codon UGA is read as tryptophan (Trp); in addition, few untranslated sequences exist and a very high mutation rate, about 5-10 times compared to that of nuclear DNA of the cell is observed.

As all mitochondria are contributed by the ovum during zygote formation, human mitochondrial DNA is transmitted by maternal non-mendelian inheritance. This is clinically significant because in diseases resulting from mutations of mitochondrial DNA, except in a few cases, an affected mother would pass the disease to all of her children but only her daughters would transmit the trait.

The below image shows the mitochondrial inheritance



Solution to Question 21:

The repair mechanism associated with CRISPR-Cas9 is non-homologous end-joining.

Non-homologous end-joining (NHEJ) is a mechanism by which a group of proteins mediates the recognition, processing, and direct ligation of the ends of two DNA fragments where a double-strand break in DNA has occurred. This is in contrast to homologous recombination (HR), another mechanism to repair a double-strand break, where DNA repair needs an undamaged homologous DNA template to replace an adjacent damaged one.

Clustered regularly interspaced short palindromic repeats-associated gene 9 or CRISPR-Cas9 is a DNA editing/gene regulatory system. It involves two essential components, Cas9, an endonuclease that causes a double-stranded DNA break, allowing modifications to the genomic DNA to be made; and a synthetic single guide RNA or sgRNA responsible for guiding Cas9 to its target.

Once the sgRNA has directed the Cas9 to the section of the genomic DNA to be edited, the 3' end of the sgRNA comes to lie across a specific sequence of DNA adjacent to the editing side called the protospacer adjacent motif or PAM. The Cas9 now causes a double-stranded break in the DNA at the PAM site which is then repaired by a non-homologous end-joining.

The following image shows CRISPR-Cas9 system



Solution to Question 22:

Pyruvate dehydrogenase is the enzyme affected in a patient with thiamine deficiency who develops lactic acidosis.

Pyruvate is transported from the cytosol into the mitochondria, where undergoes oxidative decarboxylation to form acetyl CoA in the presence of the multi-enzyme pyruvate dehydrogenase complex (PDHC) in the mitochondrial matrix.

The PDHC primarily consists of three enzymes: Pyruvate decarboxylase (also called Pyruvate dehydrogenase), dihydrolipoyl transacetylase, and dihydrolipoyl dehydrogenase. It also requires the following co-enzymes:

• Thiamine pyrophosphate, essential for pyruvate decarboxylase and is tightly bound to it

• Lipoic acid and coenzyme A, essential for dihydrolipoyl transacetylase

• FAD and NAD+, essential for dihydrolipoyl dehydrogenase.

In a patient with thiamine deficiency, due to the inactive thiamine pyrophosphate, the pyruvate dehydrogenase complex is unable to act on pyruvate, which begins to accumulate on the body and is then shunted towards lactic acid formation by the enzyme lactate dehydrogenase.

This results in lactic acid accumulation and subsequent lactic acidosis in the patient.

Wernicke–Korsakoff, an encephalopathy-psychosis syndrome due to thiamine deficiency, may be seen in chronic alcoholics. It occurs due to insufficient production of ATP by the brain cells via the citric acid cycle as the PDHC is inactive.

Other options:

Option A: Pyruvate carboxylase facilitates the conversion of pyruvate to oxaloacetate by carboxylation.

Option C: Phosphoenolpyruvate carboxykinase is involved in the decarboxylation and phosphorylation of oxaloacetate to form phosphoenolpyruvate seen in gluconeogenesis.

Option D: Transketolase with coenzyme thiamine pyrophosphate is an enzyme involved in the non-oxidative phase of the pentose phosphate pathway.



Solution to Question 23:

Triglycerides are not amphipathic.

Amphipathic substances are those which have both a hydrophilic or polar region and a hydrophobic or nonpolar region.

Phosphoglycerols or glycerol-based phospholipids have a hydrophilic head which is a phosphate group with alcohol attached to it and a hydrophobic tail containing fatty acids or fatty acid-based hydrocarbons. Therefore, they are amphipathic.

Sphingolipids, like all membrane-based lipids, are amphipathic molecules having hydrophobic and hydrophilic properties.

Glycolipids are amphipathic molecules consisting of a hydrophilic oligosaccharide chain attached to a hydrophobic lipid that anchors the glycolipid in cell membranes.

But lipids like triacylglycerol and cholesteryl esters are termed neutral lipids and thus are not amphipathic.

Cholesterol is amphipathic.

Solution to Question 24:

Only test 3, which is Rothera's test for ketone bodies in the urine will be positive in the urine of a patient with prolonged starving.

Urine tests

Test 1 - Benedict's test for reducing sugars

Test 2 - Heat and acetic acid test for presence of albumin

Test 3 - Rothera's test for ketone bodies

In case of prolonged starvation,

• After a meal till about 2 hours later, maintenance of blood glucose is primarily from the diet. During this time, hepatic glycogen stores are getting replenished.

• As dietary blood glucose levels start to decline, glucagon secretion increases and insulin secretion decreases, resulting in rapid mobilisation of hepatic glycogen stores.

• If there is fasting for a longer period of time, hepatic glycogen stores cannot maintain blood glucose levels. 12 hours after the last meal, gluconeogenesis becomes the primary source of blood glucose, since most of the hepatic glycogen would have depleted. If an individual continues fasting, gluconeogenesis continues to be the main source.

• However, over the first few days of fasting, ketogenesis happens. This is when the concentration of acetyl CoA from fatty acid oxidation exceeds the oxidative capacity of the TCA cycle. Also, ketogenesis releases CoA, ensuring its availability for continued fatty acid oxidation.
Water-soluble ketone bodies are used as fuel by most tissues, including the brain during fasting. It also reduces the need for gluconeogenesis from amino acid carbon skeletons, thus preserving essential protein in the body.

Therefore, during prolonged fasting, the levels of ketone bodies in the body rise and subsequently begin to be eliminated in urine. Hence, in starvation ketosis, Rothera's test will be positive, but Benedict's test will be negative.

Both Benedict's and Rothera's tests will be positive in diabetic ketoacidosis. (Option 1)

• Benedict's test (Test 1) for reducing sugars is a semiquantitative method for the estimation of sugar in the urine.

• Rothera's test (Test 3) is performed by saturating urine with solid ammonium sulfate. On adding sodium nitroprusside and liquor ammonia along the sides of the test tube, a purple ring is formed at the junction of the two liquids if there is acetone or acetoacetic acid in urine.
Beta-hydroxybutyrate does not give a purple ring.

• Heat and acetic acid test (Test 2) for proteinuria is based on the fact that when heated to an isoelectric point, proteins get coagulated.

Color	Concentration
Green	0.5 g%
Yellow	1 g%
Orange	1.5 g%
Red	2 g%

Solution to Question 25:

The action of inhibition on acetylcholinesterase enzyme by organophosphorus compounds is through irreversible competitive inhibition.

Competitive inhibition is a type of inhibition that occurs when the inhibitor binds reversibly to the same site that the substrate would normally occupy. It, therefore, competes with the substrate for binding to the enzyme active site. The effect of competitive inhibition can be overcome by increasing the concentration of the substrate. Km is increased, but Vmax is unaltered in this type of inhibition.

Competitive inhibition is of two types:

• Irreversible (non-equilibrium type) competitive inhibition: The inhibitor forms strong covalent bonds or has such a high affinity for the enzyme that the normal substrate is not able to displace the inhibitor. For example, organophosphorus poisoning inhibits acetylcholinesterase(AChE). Once an organophosphate binds to AChE, endogenous hydrolysis of the phosphorylated enzyme by esterases further leads to irreversible binding and permanent enzyme inactivation which is termed aging.

• Reversible (equilibrium type) competitive inhibition: Substrate concentration is sufficiently increased, it can displace the inhibitor. Statin drug inhibits HMG CoA reductase.

Noncompetitive inhibitors bind to enzymes at remote sites from the active site. Therefore, noncompetitive inhibition cannot be overcome by increasing the concentration of the substrate. Thus, noncompetitive inhibitors decrease the Vmax of the reaction. They do not interfere with the binding of the substrate to the enzyme. Thus, the enzyme shows the same Km in the presence or absence of noncompetitive inhibitors. For eg, acetazolamide inhibits carbonic anhydrase, digoxin inhibits Na+ K+ ATPase

Graph showing the changes in Vmax and Km noncompetitive and competitive inhibition.





Suicidal inhibition is a special type of irreversible inhibition. It is also known as mechanism-based inactivation. The inhibitor is a specialised substrate analog that can be transformed by the enzyme it is inhibiting. After binding to the active site of the enzyme, a highly reactive group is generated. This covalently binds to and inhibits the enzyme. An example is allopurinol, which is oxidised by xanthine oxidase to alloxanthine. Alloxanthine in turn is a strong inhibitor of xanthine oxidase.

Biochemistry NEET 2021

Question 1:

An alcoholic patient presented with itchy lesions as shown in the image, He also has a red-colored beefy tongue. Which of the following histories would you like to additionally ask/ elicit in this patient?

a) Ask for diarrhea and history of memory loss

b) Ask for diarrhea, thiamine deficiency features, and detailed dietary history

c) History of allergy

d) Ask for diarrhea and conjunctivitis

Question 2:

A patient presented with symptoms of delayed adaptation to vision in dark places, gritty sensation in the eyes, dry eyes, and softening of the cornea. What is the likely deficiency in this case?

a) Retinoic acid

b) Riboflavin

c) Niacin

d) Pantothenic acid

Question 3:

You are conducting an experiment on mitochondrial respiration. You add malate/pyruvate and respiration is normal. You add succinate and respiration is normal. When you add another substance in the presence of pyruvate/succinate and malate, respiration is blocked. Which of the following substances is most likely added?

a) Rotenone

b) Antimycin A

c) Oligomycin

d) 2,4-dinitrophenol

Question 4:

A patient presented with tendon xanthomas. His lipid profile showed serum cholesterol levels to be 283 mg/dL and serum LDL to be 202 mg/dL. He was started on lovastatin. What is the likely diagnosis?

a) Familial hypercholesterolemia

b) Tangier's disease

c) Lipoprotein lipase deficiency

d) Familial hyperchylomicronemia

Question 5:

Which of the following glycosaminoglycans (GAGs) are responsible for corneal transparency?

a) Keratan sulfate

b) Chondroitin sulfate

c) Heparan sulfate

d) Hyaluronic acid

Question 6:

A person who had dinner at 8 pm the previous night, tested his blood glucose levels at 7 am the next morning. What is the source of blood glucose in this case?

a) Hepatic glycogenolysis

b) Hepatic gluconeogenesis

c) Muscle glycogenolysis

d) Dietary glucose

Question 7:

A patient who has been diagnosed with multiple gastrointestinal polyps and carcinoma, with a positive family history/history of hereditary non-polyposis colon carcinoma has a defect in which of the following?

a) Mismatch repair

b) Nucleotide excision repair

c) Base excision repair

d) Point mutation

Question 8:

The enzymes that convert pyruvate to acetyl CoA are given in the diagram. What are the enzymes marked as E1 and E2 respectively?



a) Pyruvate dehydrogenase and dihydrolipoyl transacetylase

b) Pyruvate carboxylase and dihydrolipoyl transacetylase

c) Pyruvate dehydrogenase and dihydrolipoyl dehydrogenase

d) Pyruvate carboxylase and dihydrolipoyl dehydrogenase

Question 9:

A young girl presented to the OPD for examination. She is tall and thin with slender fingers, and dislocated lenses. Workup revealed cystathionine beta-synthase deficiency. Which amino acid supplement is required in this case?

a) Cysteine

b) Methionine

c) Serine

d) Tyrosine

Question 10:

A patient presented with joint pain and on evaluation, urine turns black on standing. What is the likely enzyme deficiency?

a) Homogentisic acid 1, 2 - dioxygenase

b) Tyrosine transaminase

c) Fumarylacetoacetate hydrolase

d) 4-hydroxy-phenyl pyruvate dioxygenase

Question 11:

A 5-year-old boy presented with easy fatigability, irritability and inability to concentrate. Labs revealed the following findings a and b (light and oil immersion respectively) on the bone marrow aspiration. Which of the following is the most likely enzyme deficient in this condition?



a) Hexosaminidase

b) Glucocerebrosidase

c) Sphingomyelinase

d) N-acetylglucosaminidase

Question 12:

A farmer presented with loose stools and rashes over the sun-exposed areas. Careful diet history reveals a diet rich in maize. Which of the following deficiencies would you suspect?

a) Niacin

b) Pyridoxine

c) Thiamine

d) Biotin

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	a

5	a
6	a
7	a
8	a
9	a
10	a
11	b
12	a

Detailed Explanations

Solution to Question 1:

An alcoholic patient with a red-colored beefy tongue, itching and the characteristic Casal's necklace, shown in the image, is suggestive of pellagra. It is caused by niacin/vitamin B3 deficiency. Hence, one should ask for a history of diarrhea and memory loss in this case.

Pellagra presents with 4 Ds: Diarrhea, Dermatitis, Dementia, and if untreated, Death. Early symptoms of pellagra are loss of appetite, generalized weakness and irritability, abdominal pain, and vomiting. Photosensitive dermatitis is seen over sun-exposed areas. Diarrhea occurs due to proctitis or malabsorption. Bright red glossitis (beefy tongue) and Casal's necklace (a pigmented, scaly skin rash that forms a ring around the neck) follow. Depressive psychosis and dementia also commonly occur.

Corn/ maize/ jowar based diets are the main causes of niacin deficiency. Corn/maize lacks niacin and tryptophan. The excess leucine in jowar and maize disrupts the conversion of tryptophan to niacin. Niacin deficiency is also seen in alcoholism. Hence, additional histories of diarrhea and memory loss will aid in supporting the diagnosis of pellagra (along with dietary history).

Treatment includes oral supplementation with 100–200 mg of nicotinamide or nicotinic acid three times daily for 5 days.

Option B: Thiamine deficiency (vitamin B1) features include CVS manifestations like heart failure and edema (wet beriberi), peripheral neuritis (dry beriberi), and Wernicke encephalopathy (in alcoholics).

Option C: Dermatitis seen in pellagra is due to dietary deficiency of niacin and so allergy history is not relevant.

Option D: Conjunctivitis is not a feature of niacin deficiency.

Solution to Question 2:

The clinical picture of delayed adaptation to vision in dark places, gritty sensation in the eyes, dry eyes, and softening of the cornea is suggestive of vitamin A deficiency in the patient. Hence

retinoic acid (the active form of vitamin A) is likely to be deficient.

Vitamin A is a fat-soluble vitamin that can occur as carotenoids (most common - beta carotene from plant sources) or retinoids (from animal sources).

3 forms of retinoids are:

• Retinal is required mainly for normal vision (present in the visual pigment rhodopsin).

• Retinol is important for the normal functioning of the reproductive system.

• Retinoic acid has a role in the regulation of gene expression and tissue differentiation.

Ocular manifestations of vitamin A deficiency include loss of sensitivity to green light (earliest to appear), nyctalopia (night blindness), xerophthalmia.

Xerophthalmia includes conjunctival xerosis (dryness) with Bitot's spots (white patches of keratinized epithelium appearing on the sclera), corneal ulcers, and keratomalacia (softening of the cornea) that leads to corneal scarring and ultimately, blindness.

Other manifestations: General impairment of immunity, keratinization of skin (phrynoderma/toad skin).

Treatment: The WHO recommended schedule for all patients above the age of 1 year (except women of reproductive age): 200,000 IU of vitamin A orally or 100,000 IU by intramuscular injection should be given immediately on diagnosis (day 0) and repeated the following day (day 1) and 14 days later.

Dietary sources of vitamin A are liver, fish, eggs, dark green, and deeply colored fruits and vegetables.

Bitot's spots



The below image shows phrynoderma with multiple keratotic papules

Other options:

Option B: Ocular manifestations of riboflavin deficiency (vitamin B2) are photophobia, burning sensation, and vascularization of the cornea. Other features include cheilosis, magenta-colored tongue, glossitis, and seborrheic dermatitis.

Corneal vascularization. Seen in riboflavin (Vitamin B2) deficiency.



Option C: Niacin deficiency (vitamin B3) leads to pellagra. Dermatitis, diarrhea, and dementia are its common features.

Casal's necklace finding in niacin deficiency



Option D: Features of pantothenic acid deficiency (vitamin B5) are burning foot syndrome ("nutritional melalgia"), dermatitis, enteritis, alopecia, ataxia.

Solution to Question 3:

In the experiment conducted on mitochondrial respiration, respiration will be blocked by the addition of oligomycin, an antibiotic, when added in the presence of pyruvate/ succinate and malate. Oligomycin inhibits oxidative phosphorylation in the mitochondria and hence blocks respiration.

Electron transport chain (ETC)/respiratory chain is the final pathway by which all the electrons derived from the oxidation of carbohydrates, fatty acids, and amino acids are directed towards the final reaction with oxygen, to form water and adenosine triphosphate (ATP).

The chemiosmotic theory states that the flow of electrons through the respiratory chain generates a proton motive force. This force drives the formation of ATP by the process of oxidative phosphorylation.

In the given experiment, malate and pyruvate are electron donors (NADH) at complex I of the ETC, and succinate is the electron donor ($FADH_2$) at complex II. Normal respiration is observed on the addition of substances at both complexes I and II i.e., both oxygen consumption and ATP synthesis from ADP (adenosine diphosphate) take place.

Oligomycin is a competitive inhibitor of the Fo (oligomycin-sensitive) complex of ATP synthase (complex V) and blocks oxidation and phosphorylation by stopping the flow of protons through the ATP synthase enzyme complex. Hence the addition of oligomycin will inhibit complex V and stop respiration.

Other options:

Options A and B: Rotenone (option A) if added, will inhibit complex I. Electron transfer before complex I will be blocked but not after. Hence, normal respiration will occur. Similarly, antimycin A (option B) inhibits complex III, and electron transfer prior to complex III will be blocked and

not after. Hence, respiration will not be completely hindered.

Option C: 2,4-dinitrophenol (DNP) is an uncoupler that dissociates oxidation in the respiratory chain from phosphorylation. Hence, respiration continues without ATP synthesis.

Complexes of ETC	Inhibitors
Complex I: NADH-CoQ oxido reductase	Piericidin, amobarbital, roten one
Complex II: Succinate-CoQ r eductase	TTFA (thenoyltrifluoroaceton e), carboxin, malonate
Complex III: CoQH2-cytochr ome c oxidoreductase	BAL(British anti-lewisite), an timycin A
Complex IV: cytochrome oxid ase	Hydrogen sulfide, carbon mo noxide, cyanide
Complex V (informally): ATP Synthase	Atractyloside, oligomycin

Solution to Question 4:

The clinical presentation with tendon xanthomas and elevated serum LDL (low-density lipoprotein) and elevated serum cholesterol points to a diagnosis of familial hypercholesterolemia (FH).

Familial hypercholesterolemia (type IIa hyperlipoproteinemia) is an inherited disorder due to defects in LDL receptors. This defect causes a decrease in the uptake of LDL and cholesterol. As a result, the blood levels of both LDL and cholesterol are elevated. Such patients are at increased risk to develop atherosclerosis and coronary artery disease.

Features of FH:

• Painful polyarthritis and tendinitis are common.

• Physical findings seen are corneal arcus and tendon xanthomas over Achilles tendon and the dorsum of hands. Xanthomas are tumorous masses produced by foamy cell clusters (lipid-laden macrophages) found in the connective tissue of the skin and in tendons.

Diagnosis is done clinically based on hypercholesterolemia with

• LDL > 190 mg/dl,

• Family history of hypercholesterolemia and/or

• Premature coronary artery disease.

Treatment includes lifestyle changes, restriction of a fatty diet, and pharmacotherapy with statins in heterozygous FH patients.

Recent drugs for specific treatment of homozygous FH are:

• Lomitapide, a microsomal triglyceride transfer protein (MTP) inhibitor, and

• Mipomersen, an antisense oligonucleotide against apoB.

The image below shows Achilles tendon xanthoma and a plain lateral radiograph of both ankles showing bilateral symmetrical swellings (arrows) superior to the calcaneum respectively.



Other options

Option B: Tangier's disease is a hypolipoproteinemia caused by a mutation of the ABCA1 gene. Enlarged orange tonsils are pathognomonic. Low HDL levels are seen. There is an increased risk for atherosclerosis.

The image below shows enlarged yellow-orange colored tonsils in Tangier's disease.



Option C: Lipoprotein lipase (LPL) deficiency causes type I hyperlipoproteinemia due to defects in LPL or apo C-II. Eruptive xanthomas are seen. Chylomicrons and VLDL are elevated. LDL and HDL are low. There is no risk for coronary disease.

Option D: Familial hyperchylomicronemia syndrome is caused due to either LPL deficiency, ApoC-II deficiency, ApoA-V deficiency, or GPIHBP1 mutation. Clinically, patients present with recurrent episodes of severe abdominal pain due to acute pancreatitis. The retinal blood vessels may be opalescent and are called lipemia retinalis. Increased levels of triglycerides (mostly >1000 mg/dL) is seen. The risk of premature coronary heart disease is low.

Solution to Question 5:

The glycosaminoglycan (GAG) responsible for corneal transparency is keratan sulfate.

Glycosaminoglycans or mucopolysaccharides are complex carbohydrates made up of amino sugars and uronic acids. They may be attached to a protein molecule to form a proteoglycan.

Keratan sulfate is a mucopolysaccharide that does not contain uronic acid. It is constituted by repeating disaccharide units of N-acetyl glucosamine and galactose.

Keratan sulfate is present between the collagen fibrils in the cornea and plays a major role in corneal transparency.

2 types of keratan sulfate are seen:

• Keratan sulfate I is present in the cornea.

• Keratan sulfate II is mainly present in the cartilage.

Keratin sulfate I and II differ in their linkage to the core proteins. Keratan sulfate I links to a core protein L-asparagine via N-acetylglucosamine and Keratan sulfate II to L-Serine (or L-Threonine) via N-acetylgalactosamine.

The chemical structure of Keratan sulfate is shown in the image below.



Other options:

Option B: In chondroitin sulfates (chondroitin 4-Sulfate & chondroitin 6-Sulfate), the repeating disaccharide units are glucuronic acid and N-acetylgalactosamine. They maintain the structure of the extracellular matrix (ECM) and are located in endochondral bone and cartilage. They are also found in the ECM of the central nervous system. They act as signaling molecules in the prevention of the repair of nerve endings after injury.

Option C: Heparan sulfate is found on many extracellular cell surfaces and contains glucosamine with fewer N-sulfates than heparin. The predominant uronic acid is d-glucuronic acid. In the plasma membrane of cells, they act as receptors and mediate cell growth and cell-cell communication. It is also found in the basement membrane of the kidney along with type IV collagen and laminin, where it determines the charge selectiveness of glomerular filtration.

Option D: Hyaluronic acid is present in synovial fluid. It is also found in skin, bone, cartilage, vitreous humor, and embryonic tissues. It contains repeating units of N Acetyl Glucosamine and glucuronic acid. It plays a role in morphogenesis and wound repair. The high concentrations of hyaluronic acid along with chondroitin sulfates in cartilage contribute to its compressibility.

Solution to Question 6:

The source of blood glucose in a person fasting overnight for 11 hours (8 pm to 7 am) is hepatic glycogenolysis.

First stage of fasting: The fasting state begins two to four hours after the absorptive/well-fed state. In early fasting i.e; 4-18 hours after the well-fed state, the blood glucose level is maintained by hepatic glycogenolysis. The glycogen stores are sufficient for around 18 hours after a meal following which they start to deplete. So the source of blood glucose in a person fasting for 11 hours will be liver glycogenolysis.

Second stage of fasting: Gluconeogenesis begins 4–6 hours after the last meal but becomes fully active only after stores of liver glycogen are depleted (after 18 hours). They can serve as a source

of glucose for up to 2 days.

Substrates for gluconeogenesis are:

• Glucogenic amino acids (mainly alanine) from the muscle and

• Glycerol from adipose tissue

During fasting, the glucogenic amino acids are transported to the liver, transaminated into pyruvate, and converted to glucose.

Third stage of fasting:

• Lipolysis occurs in the adipose tissue due to the activity of hormone-sensitive lipase, stimulated by increased glucagon:insulin ratio.

• Free fatty acids are the source of energy in this stage and not glucose. Triacylglycerols will be hydrolyzed to fatty acids and glycerol.

• Beta oxidation of fatty acids takes place which provides acetyl CoA.

• In prolonged starvation (>3 days), acetyl CoA is converted to ketone bodies in the liver which will act as the main fuel source for skeletal and heart muscle and can meet up to 20% of the brain's energy needs.

Final stage of fasting: After several weeks of starvation, ketone body synthesis will not take place due to depleted triacylglycerol stores. Proteins remain the only source of fuel. Protein degradation increases, and death results from a loss of heart, liver, or kidney function.

Role of insulin and glucagon:

When blood glucose falls during fasting, there is increased secretion of glucagon and decreased release of insulin. Hence glucagon:insulin ratio is increased.

Effects of increased glucagon:insulin ratio on enzymes of glycogenolysis and gluconeogenesis:

• It activates glycogen phosphorylase, the key enzyme of glycogenolysis, and deactivates glycogen synthase. This causes a rapid mobilization of liver glycogen stores to be utilized as fuel.

• It favors the synthesis of glucose-6-phosphatase which converts glucose-6-phosphate to glucose. The resulting glucose is used by red blood cells (lack mitochondria) and the brain as they are largely dependant on glucose in the fasting state.

• Phosphoenol pyruvate carboxykinase and fructose-1, 6-bisphosphatase are induced, leading to increased gluconeogenesis.

Metabolic process	Fasting state	Fed state
Glucagon: insulin	Increased	Decreased
Gluconeogenesis	Increased	Decreased
Glycogenolysis	Increased	Decreased
Ketogenesis	Increased	Decreased

Metabolic process	Fasting state	Fed state
Glycogen storage	Decreased	Increased



Other options:

Option C: Muscle glycogen cannot contribute directly to plasma glucose, since muscle lacks glucose-6-phosphatase. So glycogen cannot be converted to glucose to be used as fuel during starvation.

Option D: In the well-fed state/ absorptive state of two to four hours after a meal, dietary glucose is the main source of fuel.

The below image shows steps of glycolysis and gluconeogenesis.



Solution to Question 7:

Mismatch repair will be defective in a patient with multiple gastrointestinal polyps and carcinoma, with a positive family history/history of hereditary non-polyposis colon carcinoma (HNPCC)

Mismatch repair is a DNA repair mechanism that corrects single-nucleotide mismatches, insertions, and deletions introduced by the DNA polymerase during DNA replication.

HNPCC or Lynch syndrome is an autosomal dominant genetic condition. It occurs due to defects in DNA mismatch repair genes like MLH1 and MSH2. The genomes of these patients show microsatellite instability. HNPCC is associated with a high risk of colon cancer. It may also be associated with a risk of developing carcinomas of the endometrium, stomach, small bowel, ovary, pancreas, and transitional cell carcinoma of the ureter and renal pelvis.

The diagnostic criteria for HNPCC are referred to as the Amsterdam criteria or the 3-2-1-0 rule. It includes:

• Three or more relatives with HNPCC-associated cancer (colorectal cancer, endometrial cancer, cancer of the small bowel, ureter, or renal pelvis), one of whom is a first-degree relative of the other two.

• At least two successive generations affected

• At least one case diagnosed before age 50 years

• Familial adenomatous polyposis excluded

• Tumors verified by pathologic examination.

Other options:

Option B: Patients with xeroderma pigmentosum have a defect in the nucleotide excision repair pathway.

Option C: Defect in base excision repair pathway is seen in patients with MUTYH-associated polyposis.

Option D: Sickle cell disease and beta-thalassemia are examples of diseases seen in patients with point mutations.

Solution to Question 8:

E1 is pyruvate dehydrogenase and E2 is dihydrolipoyl transacetylase in the image showing oxidative decarboxylation of pyruvate. This is catalyzed by the pyruvate dehydrogenase (PDH) complex where E1, E2, and E3 are the three enzyme components of the complex.

The pyruvate dehydrogenase (PDH) complex is a multienzyme complex that is associated with the inner mitochondrial membrane. It catalyzes the oxidative decarboxylation of pyruvate to acetyl-CoA which enters the citric acid/Kreb's cycle.

Steps of pyruvate decarboxylation:

• Pyruvate is transported into the mitochondria by a proton symporter.

• Pyruvate is decarboxylated by pyruvate dehydrogenase (E1). TPP is required for this step. The decarboxylated pyruvate remains bound to the enzyme, as hydroxyethyl thiamine pyrophosphate.

• The hydroxyethyl group is oxidized to form an acetyl group. It is then transferred from TPP to lipoamide to form acetyl lipoamide. This step is catalyzed by dihydrolipoyl transacetylase (E2).

• Acetyl lipoamide reacts with coenzyme A to form acetyl-CoA and reduced lipoamide.

• The reduced lipoamide is reoxidized by FAD-dependent dihydrolipoyl dehydrogenase (E3).

The overall reaction is : Pyruvate + NAD+ + CoA → Acetyl-CoA + NADH + H+ + CO2

Enzyme components of PDH complex	Co-factors
Pyruvate dehydrogenase (E1)	Thiamine pyrophosphate (TPP)
Dihydrolipoyl transacetylase (E2)	Lipoic acidCoenzyme A
Dihydrolipoyl dehydrogenase (E3)	Flavin adenine dinucleotide (FAD)Nicotinamide adenine dinucleotide (NAD+)



Clinical significance:

• Dietary deficiency of thiamine inhibits PDH. Chronic alcoholics may develop potentially fatal pyruvic and lactic acidosis. Lactic acidosis is a common presentation in patients with inherited pyruvate dehydrogenase deficiency too.

• In arsenic and mercury poisoning, the heavy metal ions react with -SH groups of lipoic acid and inhibit pyruvate dehydrogenase.

Other multienzyme complexes include alpha-ketoglutarate dehydrogenase and branched-chain ketoacid dehydrogenase. All the five coenzymes of PDH are common to these two complexes too.

Solution to Question 9:

The clinical picture of a tall and thin young girl with slender fingers, dislocated lenses, and cystathionine beta-synthase deficiency is suggestive of classic homocystinuria. The amino acid supplement required in this scenario is cysteine.

Classic homocystinuria is a metabolic disorder of methionine metabolism, due to cystathionine beta-synthase deficiency. It is an autosomal recessive condition.

Cystathionine beta-synthase is a pyridoxal phosphate-dependent enzyme that condenses homocysteine with serine to form cystathionine. The absence of this enzyme will cause cysteine deficiency, hence cysteine needs to be supplemented in the diet.

Clinical features:

• Affected children are initially asymptomatic.

• Ectopia lentis (subluxation of the lens) is seen first, after 3 years of age.

• Intellectual disability and psychiatric and behavioral disorders.

• Thromboembolic episodes.

• Skeletal abnormalities resembling those of Marfan syndrome: They are tall and thin with elongated limbs and arachnodactyly (slender fingers).

Elevations of both methionine and homocysteine in body fluids are the diagnostic laboratory findings. The diagnosis is established by molecular analysis of cystathionine β-synthase (CBS) or by assay of the enzyme in cultured fibroblasts, phytohemagglutinin-stimulated lymphocytes, or liver biopsy specimens. Treatment includes restriction of methionine intake and supplementation with cysteine, betaine, vitamins B6, B12, and folate.



The image below shows arachnodactyly.



Methionine supplementation is required in non-classical homocystinuria. It is due to impaired re-methylation of homocysteine to methionine. This can be caused by defective methionine synthase or reduced availability of two essential cofactors, 5-methyltetrahydrofolate, and methylcobalamin.

Elevated homocysteine levels with low levels of methionine will be seen. Treatment includes methionine supplementation and administration of methyl folate, hydroxocobalamin, and betaine.

Solution to Question 10:

The urine turning black in color, on exposure to air, along with joint pain is suggestive of alkaptonuria. It occurs due to the defect in the homogentisic acid 1, 2 - dioxygenase enzyme.

Alkaptonuria is an autosomal recessive disorder of tyrosine metabolism, that develops due to the defect in homogentisate oxidase/homogentisic acid 1, 2 - dioxygenase, which oxidizes homogentisate to maleylacetoacetate. The defect in this enzyme leads to the accumulation of homogentisate, which is then excreted in the urine.

Alkaptonuria is characterized by homogentisic aciduria, ochronosis, and arthropathy (joint pain).

Homogentisic aciduria is the black discoloration of urine on prolonged exposure to air caused due to oxidation of the excreted homogentisate.

Ochronosis is the accumulation of oxidized homogentisic acid pigment (alkaptone bodies) in connective tissues leading to bluish-black discoloration of tissues mostly involving ear cartilage, ocular tissue, the skin of the eyelids, forehead, cheeks, and hands, nail beds, and intervertebral discs. The compound that accumulates in the tissues in ochronosis is benzoquinone acetate.

Painful arthritis (arthropathy) may be seen from the fourth or fifth decade.

Diagnosis: Blackish discoloration of urine on standing is diagnostic of alkaptonuria (shown below). The ferric chloride test will be positive.

The image further below shows Ochronosis.



Treatment: No specific treatment is required. Symptomatic treatment with analgesics, spinal surgery, and arthroplasty along with ascorbic acid and protein restriction can be considered. Nitisinone (p-hydroxyphenylpyruvic acid (pHPPA) oxidase inhibitor) reduces urinary excretion of homogentisic acid.

Other options:

Option B: Tyrosine transaminase deficiency causes type II tyrosinemia (Richner-Hanhart syndrome). Patients present with painful corneal erosions with photophobia, learning disability, and palmoplantar keratosis.

Option C: Fumarylacetoacetate hydrolase deficiency causes type I tyrosinemia (Tyrosinosis). Patients generally present with failure to thrive, neuropathy, boiled-cabbage urine, and liver failure secondary to cirrhosis.

Option D: 4-hydroxy-phenyl pyruvate dioxygenase deficiency causes type III tyrosinemia (neonatal tyrosinemia). Patients might rarely have a mental disability. The liver function is generally normal.

The steps of tyrosine metabolism are shown below.



Solution to Question 11:

The given clinical scenario of a child with easy fatiguability, irritability, and inability to concentrate with the images showing a crumpled tissue paper appearance(Gaucher's cell) is suggestive of Gaucher's disease. The deficient enzyme is glucocerebrosidase (also known as beta-glucosidase). I

Gaucher's disease is an inborn error of metabolism caused due to a deficiency of beta glucocerebrosidase. It is the most common lysosomal storage disorder and the mode of inheritance is autosomal recessive. There is an accumulation of lipid-laden macrophages in the spleen, liver, and other tissues.

There are three clinical subtypes - type I (chronic non-neuronopathic), type II (acute neuronopathic), and type III (intermediate). Clinical features include anemia, thrombocytopenia, hepatosplenomegaly, and neurological features depending on the subtype.

Gaucher's cells - These are formed as a result of the accumulation of massive amounts of glucocerebrosides within the phagocytes. The cells have a fibrillary cytoplasm and give a crumpled tissue paper appearance.

Option A: Deficiency of beta-hexosaminidase A is seen in Tay-Sachs disease. The deficiency of beta-hexosaminidase A and B causes Sandhoff's disease. Clinical features are seizures, macrocephaly, and hyperacusis.

Option C: Deficiency of sphingomyelinase leads to Niemann-Pick disease. It is characterized by the presence of foam cells in the bone marrow. There may be pulmonary infiltrates leading to lung failure.

Option D: Deficiency of N-acetylglucosaminidase leads to Sanfillipo's disease. It is a type of mucopolysaccharidosis (MPS - III). There is an accumulation of heparan sulfate leading to severe cognitive degeneration.

Solution to Question 12:

History of a maize-rich diet along with loose stools and rashes over the sun-exposed areas is suggestive of niacin/vitamin B3 deficiency.

Niacin deficiency causes pellagra, found mostly among people eating corn/maize-based diets, which is low in niacin and tryptophan. Excess leucine in jowar and maize disrupts the conversion of tryptophan to niacin.

Niacin can be synthesized in the body from tryptophan. Two compounds- nicotinic acid, and nicotinamide have the biological activity of niacin. The nicotinamide ring of the coenzymes NAD (nicotinamide adenine dinucleotide) and NADP (nicotinamide adenine dinucleotide phosphate) participate in oxidation/reduction reactions.

The initial symptoms of pellagra include loss of appetite, generalized weakness, irritability, abdominal pain, and vomiting. This is followed by bright red glossitis and a characteristic pigmented, scaly skin rash in sun-exposed areas. This rash is known as Casal’s necklace when it forms a ring around the neck.

The 4Ds of pellagra:

• Diarrhea

• Dermatitis

• Dementia, leading to

• Death

It is diagnosed by measuring urinary levels of 2-methyl nicotinamide and 2-pyridone.

It is treated by oral supplementation of 100–200 mg of nicotinamide or nicotinic acid three times daily for 5 days.

Casal’s necklace finding in niacin deficiency

Other options:

Option B: Dietary pyridoxine deficiency (vitamin B6) is rare but has been observed in alcoholics, newborn infants formulas low in vitamin B6, and women taking oral contraceptives. Isoniazid, an anti-TB drug, leads to pyridoxine deficiency resulting in peripheral neuropathy. Hence pyridoxine supplementation is included in ATT regimens.

Option C: Thiamine deficiency (vitamin B1) can manifest as wet beriberi (CVS features like heart failure and edema) or Shoshin beriberi, an acute and fulminant form of wet beriberi, dry beriberi (peripheral neuritis), or as Wernicke encephalopathy with Korsakoff psychosis (in alcoholics). Dietary deficiency of thiamine inhibits pyruvate dehydrogenase. Chronic alcoholics may develop potentially fatal pyruvic and lactic acidosis.

Option D: Dietary biotin deficiency is unknown but rarely, dermatitis, glossitis, loss of appetite, and nausea may occur after consumption of large amounts of uncooked egg white, which contains the protein avidin. It binds biotin and makes it unavailable for absorption.

Note: A similar image-based question on vitamin B3 deficiency was previously asked in NEET PG 2020. Vitamin deficiencies are a high-yield topic and are repeatedly tested every year.

Biochemistry INI-CET 2022

Question 1:

Which is the correct sequence of steps in the isolation of desirable protein using recombinant DNA technology?

a) 2, 1, 3, 5, 4

b) 1, 2, 4, 3, 5

c) 1, 5, 2, 4, 3

d) 2, 4, 5, 3, 1

Question 2:

Which of the following techniques is used to detect DNA-protein interaction?

a) DNA footprinting

b) DNA fingerprinting

c) ELISA

d) Northern blotting

Question 3:

A patient was diagnosed with Ewing sarcoma with EWSR1 gene mutation. The technique shown in the image was performed to confirm the diagnosis. Identify the technique shown in the image.



a) Fluorescent in situ hybridisation

b) Karyotyping

c) Immunohistochemistry

d) Sanger's technique

Question 4:

In a patient with maple syrup urine disease, all of the following amino acids should be restricted in diet except?

a) Methionine

b) Valine

c) Leucine

d) Isoleucine

Question 5:

Which of the following statements about the human genome project is false?

a) Total 20,500 genes were identified

b) The project spanned over 13 years

c) It required international collaboration

d) Next generation sequencing was used

Question 6:

Banding technique used for dicentric chromosomes is:

a) G banding

b) C banding

c) NOR

d) R banding

Question 7:

Affinity maturation of antibodies is because of?

a) Somatic hypermutation

b) Differential mRNA processing

c) CD40

d) Gene rearrangements

Question 8:

Best method for detection of mutations with low allele frequency is

a) Droplet digital PCR

b) FISH

c) Nested PCR

d) Sanger Sequencing

Question 9:

In Sanger sequencing, ddNTP is used because:

a) It is used to increase DNA polymerase activity

b) It can be made fluorescent

c) They exhibit higher affinity for DNA polymerases

d) It doesn't allow further elongation of DNA

Question 10:

Misincorporation of uracil into DNA is seen in deficiency of:

a) Vitamin B12

b) Folate

c) Vitamin B6

d) Vitamin E

Question 11:

Deamination of methylated cytosine forms which of the following?

a) Uracil

b) Cytosine

c) Guanine

d) Thymine

Question 12:

Which of the following types of inhibition is depicted in the given Lineweaver-Burk plot?



©MARROW

a) Competitive inhibition

b) Non-competitive inhibition

c) Uncompetitive inhibition

d) Suicidal inhibition

Question 13:

Glycogen phosphorylase is activated by:

a) Calcium

b) Glucose-6-phosphate

c) Insulin

d) Protein phosphatase

Question 14:

Mutation of the gene of cystic fibrosis occurs with:

a) Tryptophan at 708 position, chloride channel

b) Phenylalanine at 708 position, calcium channel

c) Phenylalanine at 508 position, chloride channel

d) Tryptophan at 508 position, calcium channel

Question 15:

A baby is brought to the OPD with complaints of lethargy and poor feeding. On examination, it is noted that the urine had a smell of burnt sugar. A defect in which of the following enzymes causes this disease?

a) Branched-chain ketoacid dehydrogenase

b) Branched-chain amino acid dehydrogenase

c) Branched-chain ketoacid oxidase

d) Branched-chain amino acid oxidase

Question 16:

All are true about the hexose monophosphate shunt, except:

a) Source of ribose-5-phosphate

b) Source of Acetyl CoA

c) Source of NADPH

d) Utilises ribose

Question 17:

Which of the following enzymes use selenocysteine?

a) 1, 3, 4

b) 1, 2, 3

c) 2, 3, 4

d) 1, 2, 4

Question 18:

A patient presented with reduced levels of high-density lipoprotein, and ABCA1 mutation. On examination, tonsils appeared as shown in the image. What is the diagnosis?



a) Tangier's disease

b) Abetalipoprotenemia

c) Hyperlipoproteinemia

d) Wolman's disease

Question 19:

A baby is being evaluated for delayed developmental milestones. On examination, the child has hepatosplenomegaly. A microscopic image of the bone marrow evaluation is shown below. What is the diagnosis?



a) Gaucher disease

b) Niemann-Pick disease

c) Hunter disease

d) Hurler disease

Question 20:

Refsum disease is due to:

a) Absence of alpha oxidation in peroxisomes

b) Absence of beta oxidation in peroxisomes

c) Absence of alpha oxidation in lysosomes

d) Absence of beta oxidation in lysosomes

Question 21:

Choose the right sequence of steps in the conversion of cholesterol to testosterone.

a) b, d, a, c

b) a, c, d, b

c) b, a, c, d

d) d, a, b. c

Question 22:

Which of the following describes the relationship between the melting temperature (Tm) of DNA and the concentration of G-C and A-T?

a) No relation of GC with Tm

b) Tm increases with more A-T

c) Tm decreases with more G-C

d) Tm increases with more G-C

Question 23:

An elderly patient on a vegetarian diet presenting with easy fatigability and pallor was investigated. Peripheral smear showed findings of macrocytic anemia. What is the probable cause of anemia in this patient? Other lab findings are as follows:

a) Folate deficiency

b) Alcohol-induced macrocytosis

c) Vitamin B12 deficiency

d) Thiamine deficiency

Question 24:

Zellweger syndrome is due to a defect in the biogenesis of:

a) Peroxisomes

b) Lysosomes

c) Mitochondria

d) Nucleus

Question 25:

True statement about the action of enhancers of gene expression?

a) Work only when in close conjugation with promoter region

b) Work only when in upstream of promoter region

c) They are very specific to the gene

d) Recruit chromatin and modify regulatory mechanisms

Question 26:

Increased H+ ions in the intermembrane space of mitochondria are due to?

a) Decreased oxidative phosphorylation

b) Increased ATP production

c) Reduced chemiosmosis

d) Water retention in inner mitochondrial space

Question 27:

Fructosamine is obtained by glycation of:

a) Frustose

b) Albumin

c) Urea

d) Glycosaminoglycans

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	a
5	d
6	b
7	a
8	a
9	d
10	b
11	d
12	a
13	a

14	c
15	a
16	d
17	a
18	a
19	a
20	a
21	a
22	d
23	a
24	a
25	d
26	a
27	b

Detailed Explanations

Solution to Question 1:

The correct sequence of isolation of desirable protein in recombinant DNA technology is 2, 1, 3, 5, 4.

I) Incorporation of genes into bacteria: The gene needed to produce the desired proteins is to be incorporated into the bacteria. This is done using cloning vectors like bacterial plasmids, phages, or cosmids. The DNA of interest is ligated into the vector DNA and is called recombinant DNA. This vector is then inserted into a competent host cell (eg: E.coli bacteria) where it is replicated.

II) Expression of protein and lysis of bacterial cell: After incorporation of recombinant DNA, bacteria express the inserted gene and produce the desired protein on a large scale along with other indigenous bacterial proteins. To release these proteins bacterial cell is lysed.

III) SDS-PAGE (sodium dodecyl sulfate-polyacrylamide gel electrophoresis): It is the most widely used method for determining the purity of a protein. After the proteins are released, to separate the desirable protein, we subject it to SDS-PAGE. It separates proteins based on molecular weight.

IV) Hybridization and blotting: It is done to confirm that the isolated protein is the desired protein. It is done using western blotting techniques on nitrocellulose paper.

V) Protein elution - extraction of the produced protein.

The practical applications of recombinant DNA technology include:

• Gene mapping using techniques such as fluorescence in situ hybridization (FISH).

• Producing proteins such as:

• Interferons, tPA (tissue plasminogen activator)

• Insulin, growth hormone

• Hepatitis B vaccine

• Molecular analysis of disease, to detect gene variations and mutations.

• Restriction Fragment Length polymorphism (RFLP) and single nucleotide polymorphism (SNP)

• Microsatellite DNA polymorphisms

• Gene therapy and bone marrow transplantation

Solution to Question 2:

The technique used to detect DNA-protein interaction is DNA footprinting.

The parts of DNA that are protein-bound (Eg: RNA polymerase bound to the promoter region of DNA) are resistant to digestion by nuclease enzymes. Since the nucleases are unable to cleave the protein-bound regions of DNA, when the cleaved DNA fragments are run on polyacrylamide gel electrophoresis, there are missing bands indicating, protected sites of protein-bound DNA. These missing areas represent the footprint of the protected DNA-protein complexes. Hence DNA footprinting technique helps to identify the specific binding site of protein on a DNA sequence.

The image below illustrates the DNA footprinting technique.



Other options:

Option B: DNA fingerprinting uses RFLPs (restriction fragment length polymorphism) or repeat sequence DNA to identify a person's unique pattern of DNA. Polymorphisms are inherited differences found among individuals. with this technique, individuals can be differentiated by variations in their DNA sequence, using restriction endonucleases followed by gel electrophoresis, to visualize RFLPs. DNA fingerprinting is used for forensic analysis of crime scenes, and for paternity testing.

Option C: ELISA uses antibodies linked to certain enzymes to detect certain antigens present in the sample.

Option D: Northern blotting technique can be used to detect RNA. Here, the mixture of mRNA molecules is separated by electrophoresis and then transferred to a nitrocellulose membrane and hybridized with radiolabeled probes.

Solution to Question 3:

The technique shown in the image is FISH demonstrating a positive EWSR1 gene rearrangement, characteristic of Ewing sarcoma. The fused region denoted by Yellow corresponds to a normal intact gene, while a split signal pattern showing separate red and green signals is indicative of EWSR1 gene rearrangement.

Fluorescent In Situ Hybridisation (FISH) is a simple technique for the detection of specific genetic information on morphologically intact tissue using fluorescent probes. FISH can be performed on prenatal samples, peripheral blood cells, touch preparations from cancer biopsies, and even fixed archival tissue sections. It is used to detect

• numeric abnormalities of chromosomes (aneuploidy)

• subtle microdeletions and rearrangements.

• complex translocations that are not demonstrable by routine karyotyping

• gene amplification (e.g., HER2 in breast cancer or NMYC amplification in neuroblastomas)

Other options:

Option B: Karyotyping is the process by which chromosomes are isolated, stained, and examined to find chromosomal anomalies. It is used in prenatal diagnosis, investigation of repeat pregnancy loss, investigation of children and adults suspected of having a chromosomal disorder, and subtyping of lymphomas and leukemias for the purpose of exact diagnosis, giving a prognosis, and treatment.

Option C: IHC (Immunohistochemistry) is a staining method for histopathological specimens. It helps in the diagnosis and classification of tumors based on their distinct protein expression patterns. Eg: The expression of estrogen receptor (ER), progesterone receptor (PR), and HER2/neu in breast cancer cells is mainly assessed by immunohistochemistry.

Option D: Sanger's technique: This was the first technique used to determine the sequence of proteins. The first protein to be sequenced by this method was insulin. Sanger's reagent is 1-Flouro 2,4 Dinitro benzene. Only dipeptides and tripeptides can be sequenced by this method.

Solution to Question 4:

In maple syrup urine disease (MSUD), only branched-chain amino acids are restricted, hence, methionine does not need any restriction in diet.

Branched-chain amino acids are:

• Leucine (L)

• Isoleucine (I)

• Valine (V)

MSUD is an autosomal recessive disorder in which there is a partial or complete deficiency of branched-chain ketoacid dehydrogenase complex (BCKD). This impairs the metabolism of branched-chain amino acids. These amino acids, particularly leucine and their metabolite accumulate in the blood, causing toxic effects on the brain. Features include feeding problems, vomiting, ketosis, and a characteristic burnt sugar odor /maple syrup-like odor found in urine, sweat, and cerumen.



Laboratory investigations will show increased levels of branched-chain amino acids in the plasma and urine. Leucine levels are usually higher than isoleucine, and valine.

Treatment involves restricted intake of branched-chain amino acids using commercially available formulas and judicious supplementation of isoleucine and valine. Very high levels of leucine are removed by hemodialysis.

Solution to Question 5:

The human genome project did not use next-generation sequencing.

The Human Genome Project (HGP) was an international, collaborative research program, which involved agencies such as UNESCO, HUGO, the National Institute of Health/Department of Energy (USA), the Medical Research Council (UK), Genethon (France), and the European Union. The primary goal of the Human Genome Project was to determine the sequence of all nucleotide base pairs, that make up human DNA and to create a single linear map of all genes from the human genome. The project was active from 1990 to 2003, taking around 13 years to complete. A total of 20500 genes were identified and mapped. Scientists used various methods of first-generation sequencing technologies including sanger's technique, SGS technique, etc. The HGP has produced a curated and accurate reference sequence for the majority of human

chromosomes. The reference sequence has provided the foundation for the use of second-generation sequencing technologies such as array techniques.

Next-generation sequencing (NGS) is a new technology used for sequencing DNA and RNA and mutation detection. NGS can sequence the genome faster and in a shorter period of time than first-generation sequencing technologies. NGS allows for the detection of a spectrum of mutations, including single nucleotide substitutions, small insertions and deletions, large genomic duplications and deletions, and rare variations. Next-generation DNA sequencing technologies include

• Illumina sequencing

• Roche 454 sequencing

• Ion torrent: Proton/PGM sequencing

• SOLID sequencing.

Solution to Question 6:

The banding technique used for dicentric chromosomes is C banding.

Heterochromatin includes non-coding regions of DNA, such as centromeric regions. The staining of these constitutive heterochromatins, is done by C-banding using silver salt staining. A dicentric chromosome having two centromeres can hence be identified by C banding.

Various techniques are used in karyotyping to study chromosomes. Staining with various compounds produces various results and helps us to understand the chromosomes better.

Common techniques used are:

• Q banding - Fluorescence staining with quinacrine mustard.

• G banding - Giemsa staining after partial denaturation of chromatin. It is the most common technique used in cytogenetics to produce a visible karyotype by staining condensed chromosomes.

• R banding - Reverse Giemsa staining. In this, the light and dark areas are reversed.

• C banding - staining of heterochromatin with silver salts.

• T banding - staining the telomeres of chromosomes.

Solution to Question 7:

Affinity maturation of antibodies is because of somatic hypermutation.

Affinity maturation is the process by which antibodies gain
increased affinity, avidity, and anti-pathogen activity. It is a result of somatic hypermutation of immunoglobulin genes in B cells, coupled with selection for antigen binding.

Somatic hypermutation is a cellular response by which the immune system adapts to the new foreign elements that confront it. They do so by adding nucleotides in V gene regions of heavy and

light chains, resulting in a better fit of antigen with the antibodies. This hence increases the affinity of antibodies to antigens. Due to this, the subset of plasma cells with these improved hypervariable regions are more strongly

selected by the antigen and therefore constitute a larger part of the population of antibody-producing cells. This process is called affinity maturation.

This process occurs in the germinal centers and proceeds for weeks after acute infection or vaccination, or for many cycles during chronic infection. The resulting antibodies can be highly mutated from their germline-encoded counterparts, with an increase in several orders of magnitude in affinity for antigen compared to the corresponding naïve B cell receptors (BCRs).

Solution to Question 8:

Best method for detection of mutation with low allele frequency is Droplet digital PCR (ddPCR)

It is an advanced and more precise type of quantitative PCR which gives more consistent and reproducible results. It can accurately quantify the target nucleic acid sequence within a sample. It is useful for determining somatic mutation with low allele frequency. Mutant alleles of frequency as low as 0.25% can also be detected using this technique.

Polymerase chain reaction (PCR) is a rapid automated method for the amplification of specific DNA sequences (or genes). Reverse transcriptase PCR- The enzyme reverse transcriptase (RT) is used to synthesize a complimentary copy of DNA (cDNA) from RNA and this cDNA is then used as a template for PCR.

Other special PCR types:

Nested PCR- It involves the sequential use of two primer sets. The first primer set is used for amplification and the target sequence for the second primer is present only on the amplicon with the target sequence. Nested PCR is intended to reduce non-specific binding due to the amplification of unexpected primer binding sites.



Multiplex PCR- Multiple primers are used to amplify several DNA sequences simultaneously. As this process targets multiple genes at once, a single test run gives additional information that would otherwise require several times the reagents and technician time to perform.

The Sanger sequencing technique is used for DNA sequencing. It is based on controlled termination of synthesis of DNA by using chain terminating agents like dideoxynucleotides.

Fluorescent In Situ Hybridisation (FISH) is a simple technique for the detection of specific genetic information on morphologically intact tissue using fluorescent probes. FISH can be performed on prenatal samples, peripheral blood cells, touch preparations from cancer biopsies, and even fixed archival tissue sections.

Solution to Question 9:

In Sanger sequencing, dideoxynucleotides triphosphates (ddNTP) are used because it doesn't allow further elongation of DNA.

The Sanger method is used for DNA sequencing. This method uses the mechanism of synthesis of DNA by DNA polymerases. In the process of synthesis of DNA, the 3'-hydroxyl group of primer reacts with an incoming dNTP to form a new phosphodiester bond. This reaction is catalyzed by DNA polymerase.

In the Sanger sequencing method, ddNTP is used because it halts the DNA synthesis. ddNTPs can bind to the template strand but lack the 3'-hydroxyl group which is required to add the next nucleotide. The products of this reaction are a mixture of DNA strands of varying lengths. This mixture is then separated using polyacrylamide gel electrophoresis and autoradiography. This produces a pattern of bands from which the DNA sequence can be read.

The image given below shows the steps involved in Sanger's method.



Sanger's technique was the first technique used to determine the sequence of proteins. The first protein to be sequenced by this method was insulin. Sanger's reagent is 1- fluoro 2,4 dinitro benzene.

Solution to Question 10:

Misincorporation of uracil into DNA is seen in the deficiency of folate.

Folate deficiency leads to decreased thymidylate synthesis from deoxyuridylate due to deficient methylation of deoxyuridine monophosphate (dUMP) to thymidine monophosphate (TMP). The methylation of dUMP to TMP is catalyzed by thymidylate synthase and requires tetrahydrofolate (THFA) as a methyl group donor. This decreased TMP synthesis can result in an imbalance of deoxyribonucleotides and hence leading to excessive uracil misincorporation into DNA during replication and repair.

Folic acid (vitamin B9) is a water-soluble vitamin, and its active form is tetrahydrofolate (THFA). It acts as a carrier of one-carbon groups. Tetrahydrofolate receives one-carbon fragments from donors such as serine, glycine, and histidine and transfers them to intermediates in the synthesis of amino acids, purines, and pyrimidines.



Folate deficiency results in:

- Megaloblastic anemia
- Homocystenemia
- Neural tube defects (spina bifida, Anencephaly) in the fetus.

Solution to Question 11:

Deamination of methylated cytosine forms thymine.

Deamination is the removal of an amino group(NH2) from a chemical compound. Enzymes that catalyze this reaction are called deaminases. Deamination of 5-methylcytosine forms thymine.

Other significant deaminations are given below:

• Deamination of adenosine forms inosine. This is a metabolic step involved in uric acid formation by purine catabolism. It is mediated by the enzyme adenosine deaminase (ADA). ADA deficiency leads to sparse or defective T cells and B cells and patients suffer from Severe combined immunodeficiency (SCID).

• Deamination of cytosine leads to uracil formation - The production of apolipoprotein B-48 (apoB-48) in the intestine is the result of RNA editing in the intestine. This involves post-transcriptional deamination of cytosine to uracil by enzyme cytidine deaminase. This converts sense codon (CAA) in the mRNA to nonsense codon (UAA)at a single specific site. This edited mRNA translates to apoB-48.

• Deamination of guanine forms xanthine. This is a metabolic step involved in uric acid formation by purine catabolism.

Solution to Question 12:

The given graph shows competitive inhibition, in which the value of Vmax remains constant and the Km value increases.

• Km is the substrate concentration at which velocity is half maximum.

• Vmax is the maximum velocity of a reaction.

Competitive inhibition is when the inhibitor binds reversibly to the substrate-binding site and competes with the substrate for the active enzyme-binding site. This effect of competitive inhibition can be overcome by increasing the substrate concentration by which maximum velocity can still be achieved. Therefore, Vmax is unaltered in this type of inhibition. However, the concentration of the substrate required to achieve half of the maximum velocity is increased. Therefore, Km is increased in this condition.

The graph shows the changes in Vmax and Km in competitive inhibition.

Inhibition	Km	Vmax
Competitive	Increases	Same
Non Competitive	Same	Decreases
Uncompetitive	Decreases	Decreases



Other options:

Option B: Non-competitive inhibition is by binding inhibitors to enzymes in locations remote from the active site. Increasing the concentration of the substrate does not overcome non-competitive inhibition. Thus, noncompetitive inhibitors decrease the Vmax of the reaction. They do not interfere with the binding of the substrate to the enzyme. Thus, the enzyme shows the same Km in the presence or absence of noncompetitive inhibitors.

The graph shows the changes in Vmax and Km in non-competitive inhibition.



Option C: In uncompetitive inhibition, inhibitors bind with the enzyme-substrate intermediate complex, and inhibit it. This leads to decrease in both Vmax and Km. An example is the inhibition of placental alkaline phosphate (ALP) by phenylalanine.

Option D: Suicidal inhibition is a special type of irreversible inhibition. The inhibitor here makes use of the enzyme's own reaction mechanism to inactivate it and is hence known as mechanism-based inactivation. An example is allopurinol, which is oxidized by xanthine oxidase

to alloxanthine. Alloxanthine in turn is a more potent inhibitor of xanthine oxidase.

Solution to Question 13:

Glycogen phosphorylase is activated by calcium in the muscle.

Glycogen phosphorylase is the rate-limiting enzyme of glycogenolysis and is decreased in the fed state.

In the muscle, the increased cytosolic calcium ion concentration initiates both contraction and glycogenolysis. Muscle phosphorylase kinase has four different subunits, α, β, γ, and δ. The α and β subunits are phosphorylated by cAMP-dependent protein kinase. The δ subunit is identical to the calcium-binding protein calmodulin and binds four calcium ions. Calcium coordinates the activation of Glycogen Phosphorylase with the contraction of the muscle.

Glycogen phosphorylase catalyzes the cleavage of $\alpha 1 \rightarrow 4$ linkages of glycogen to single units of glucose 1-phosphate. It uses pyridoxal phosphate as the coenzyme. Different isoenzymes of glycogen phosphorylase are seen in the liver, muscles, and brain.

Other options:

Option B: During glycogenolysis, glucose 6-phosphate allosterically inhibits glycogen phosphorylase.

Option C: Insulin entirely suppresses glycogenolysis in the skeletal muscle and liver.

Option D: In glycogenolysis protein phosphatase is inactivated by inhibitor proteins in response to cAMP, to maintain the glycogen phosphorylase in an active state.

Solution to Question 14:

The mutant gene of cystic fibrosis (CF gene) produces a mutant CFTR protein with the deletion of a phenylalanine residue at 508 (F508del), characterised by impaired chloride secretion.

Cystic fibrosis is a condition with an autosomal recessive mode of inheritance. It results due to a defective mutation in the gene encoding for cystic fibrosis transmembrane conductance regulator (CF gene). It plays an important role in mucociliary transport along the respiratory mucosa. It also helps to regulate the amount of exocrine gland secretion.

The defect in the ion channel results in the production of hyper-viscous, adherent secretions which can obstruct the airways. Organisms such as Pseudomonas aeruginosa and Staphylococcus aureus can grow in the sputum. The exocrine pancreas can undergo fibrotic scarring, cyst formation, and acinar tissue loss. The viscid secretions can lead to pancreatic duct obstruction. The flow of digestive enzymes to the duodenum is hence impaired.

Symptoms include sinusitis, steatorrhea, and abdominal pain. Distal intestinal obstruction is a complication due to difficulty in excreting intestinal contents. In neonates, it can present as meconium ileus.

Detection of high levels of trypsinogen is used in newborn screening. Chloride level >60mEq/L in the sweat of affected patients is diagnostic. Hyponatremic hypochloremic alkalosis is seen.

Postural drainage, short-term antibiotics, and inhaled bronchodilators can be used for medical management. The definitive treatment is lung transplantation.

The image below shows the features of the disease.



Solution to Question 15:

The diagnosis, in this case, is that of maple syrup urine disease (MSUD). This disease occurs due to a defect of the enzyme branched-chain keto acid dehydrogenase.

MSUD is an autosomal recessive disorder that occurs due to a deficiency of branched-chain ketoacid dehydrogenase complex. This impairs the metabolism of branched-chain amino acids valine, leucine, isoleucine, and alloisoleucine. These amino acids, particularly leucine and its metabolite accumulate in the blood, causing toxic effects on the brain.

The affected child develops poor feeding and vomiting a few days after birth. There is a period of hypertonicity with a period of flaccidity in between. These appear as repetitive boxing and bicycling movements. The babies may also have hypoglycemia, but this is not corrected by the administration of glucose. The urine, sweat, and cerumen of the baby have a sweetish maple syrup-like odor.

Laboratory investigations will show increased levels of branched-chain amino acids in the plasma and urine. Leucine levels are usually higher than isoleucine, valine, and alloisoleucine. Since the patient excretes ketone bodies in the urine, the urine will be positive for the dinitrophenylhydrazine test.

Treatment involves controlled intake of branched-chain amino acids using commercially available formulas and judicious supplementation of isoleucine and valine. Very high levels of leucine are removed by hemodialysis.

Solution to Question 16:

The hexose monophosphate (HMP) shunt produces ribose and does not utilize it.

The pentose phosphate pathway, also known as the HMP shunt pathway, is an alternative route for the metabolism of glucose. It contributes in two major ways:

• Formation of NADPH for the synthesis of fatty acids and steroids and maintenance of reduced glutathione for antioxidant activity.

• Formation of ribose for the synthesis of nucleic acid and nucleotides

The enzymes for this pathway are found in the cytosol. There are two phases:

• Irreversible oxidative phase: Glucose-6-phosphate is dehydrogenated and decarboxylated to form ribulose-5-phosphate which in turn gets converted to ribose-5-phosphate.

• Reversible non-oxidative phase: Ribulose-5-phosphate is converted back to glucose-6-phosphate.



Solution to Question 17:

The enzymes that use selenocysteine are glutathione peroxidase, deiodinase, and thioredoxin reductase.

Selenocysteine is the 21st amino acid coded by the UGA codon, which is normally a stop codon. It is formed by co-translational modification. The protein synthetic apparatus can identify a selenocysteine-encoding UGA by the presence of the selenocysteine insertion sequence (SECIS) in the untranslated region of mRNA.

Enzymes such as thioredoxin reductase, glutathione peroxidase, and deiodinase use selenocysteine to catalyze redox reactions. Defects in human selenoproteins are associated with tumorigenesis and atherosclerosis. It can cause cardiomyopathy due to selenium deficiency, which is known as Keshan disease.

Glutathione synthase is the second enzyme in the synthesis of glutathione. Glutathione plays an important role in oxidative stress, redox signalling, and regulates cell proliferation and apoptosis.

Solution to Question 18:

The diagnosis in this case with reduced levels of high-density lipoprotein (HDL), ABCA1 mutation, and orange tonsils on examination is Tangier's disease. It is caused by a defect in ABCA1 receptors.

ABCA1 plays a role in the efflux of cholesterol from peripheral cells of HDL. LCAT (lecithin cholesterol acyltransferase) converts cholesterol to cholesterol esters, which accumulate in the core of HDL. Thus, defective ABCA1 will lead to low levels of HDL in the blood.

Other options:

Option B: Abetalipoproteinemia is a hypolipoproteinemia caused by a defect in the MTP gene. Here, loading apolipoprotein B with lipids is defective. Hence, few very low density-lipoproteins (VLDLs) or chylomicrons are formed, leading to the accumulation of triacylglycerols in the liver and intestine.

Option C: Hyperlipoproteinemia is a disorder that causes the body to metabolize lipids incorrectly. This results in hyperlipidemia and can lead to the development of multiple small xanthomas in the skin.

Option D: Acid lipase deficiency can lead to Wolman's disease which is a lysosomal storage disease. The pathognomic feature of this condition is adrenal calcification. So patients generally present with adrenal insufficiency and are neonates.

Solution to Question 19:

The microscopic image of the bone marrow biopsy shows Gaucher cells and the diagnosis is Gaucher disease.

Gaucher disease is a lysosomal storage disease with autosomal recessive inheritance. The condition results due to the deficiency of acid β-glucosidase. This results in the accumulation of glucosylceramide (a glycolipid substrate) in the reticuloendothelial cells. There is progressive infiltration of the liver, spleen, and bone marrow.

There are three clinical types:

• Type 1: Adult (non-neuronopathic form)

• Type 2: Infantile (acute neuronopathic form)

• Type 3: Juvenile (subacute neuronopathic form)

Type 1 is the most common. It presents with a myriad of symptoms, such as bruising secondary to thrombocytopenia, fatigue due to anemia, hepatosplenomegaly, and bone pain. Lytic lesions and osteosclerosis of the femur, ribs, and pelvis may also be seen. A few patients may have pulmonary involvement. Growth retardation may be observed in cases presenting in the first decade.

Type 2 is rare, and presents with hypertonia, strabismus, and organomegaly. Neurodegeneration is the characteristic feature. Stridor and failure to thrive can also be the presenting features.

Ultimately, death can occur due to respiratory arrest.

Type 3 presents with intermediate symptoms, as seen in the other two types. By the age of 10-15 years, the child succumbs to death.

Gaucher cells are reticuloendothelial cells with glucosylceramide, and have a wrinkled-paper appearance due to intracytoplasmic substrate inclusions. These cells stain strongly with periodic acid-Schiff (PAS) stain. This is characteristic of Gaucher's disease.

Other options:

Option B - Acid sphingomyelinase deficiency results in Niemann-Pick disease types A and B. Both are autosomal recessive conditions. Type A has a neurodegenerative course, while type B is non-neuronopathic. The former presents with hepatosplenomegaly, lymphadenopathy, and psychomotor retardation. Type B is associated with hepatosplenomegaly and mild pulmonary involvement.

Option-C - Hunter disease is an X-linked inherited disorder due to a deficiency of the enzyme iduronate sulfatase. It leads to the accumulation of dermatan and heparan sulfate. Clinical features involve coarse facies, cardiovascular involvement, and distinctive skin lesions.

Option-D - Hurler/Scheie disease is a mucopolysaccharidosis that has childhood and adult onset. It presents with cognitive degeneration.

Solution to Question 20:

Refsum disease is due to the absence of alpha oxidation in peroxisomes.

Refsum's disease is a rare autosomal recessive disorder. The defective enzyme is phytanoyl CoA hydroxylase, which causes alpha oxidation of fatty acids and leads to elevated levels of phytanic acid in the blood.

Clinical features are dysmorphic facies and severe neurological symptoms such as polyneuropathy, retinitis pigmentosa, nerve deafness, ichthyosis, and cerebellar ataxia.

Restricted dietary intake of phytanic acid reduces the symptoms. Management includes avoiding sources of phytanic acid such as dairy products, meat, nuts, and green leafy vegetables.

Solution to Question 21:

The correct sequence of steps in the conversion of cholesterol to testosterone is:

b. 20, 22 desmolase

d. 3 beta-hydroxy steroid dehydrogenase

a. 17 alpha-hydroxylase

c. 17, 20 lyase

The step of synthesis of sex steroids is illustrated in the diagram below.



Solution to Question 22:

The correct statement about the melting temperature of DNA (Tm) is that it increases with more G-C concentration.

On heating the DNA, the temperature at which one-half of the helical structure loses its structure defines the melting temperature Tm. The two strands of the double helix structure of the DNA get separated when the hydrogen bonds between the base pairs are disrupted. Thus, the Tm is influenced by the concentrationbase composition of the DNA.

Since G-C pairing has 3 hydrogen bonds and A-T pairing has 2 hydrogen bonds, DNA rich in G-C base pairs requires higher temperatures to denature in comparison to DNA rich in A-T base pairs.

Solution to Question 23:

The probable cause of anemia in this patient of folate deficiency.

The given clinical scenario of macrocytic anemia in a vegetarian with elevated homocysteine levels is due to megaloblastic anemia, which can be caused by the deficiency of vitamin B12 or folic acid. Vitamin B12 deficiency is associated with raised serum methylmalonic acid (MMA) levels, which are normal in this patient; hence, folic acid deficiency is the cause of megaloblastic anemia in this patient.

Folic acid deficiency is associated with the following clinical and laboratory findings:

• Raised MCV (normal range: 80–100 fL) - macrocytosis

• Normal MCHC (normal range: 33–36 g/dL)

• Increased MCH

• Elevated RDW

• Decreased reticulocyte count - due to destruction of abnormal erythroid precursors

• Elevated homocysteine level

• Normal MMA level

where MCV, mean corpuscular volume; MCHC, mean corpuscular hemoglobin concentration; mean corpuscular hemoglobin, MCH; RDW, red cell distribution width

In vitamin B12 deficiency, neurological abnormalities, elevated homocysteine, and elevated MMA levels are seen.

Solution to Question 24:

Zellweger syndrome is due to a defect in the biogenesis of peroxisomes.

Zellweger syndrome is also known as cerebrohepatorenal syndrome. It is caused by mutations in the PEX family of genes. This results in defective beta-oxidation of very long-chain fatty acids.

The child presents with

• Dysmorphic facial features.

• Weakness, hypotonia.

• Neonatal seizures

• Eye abnormalities.

Biochemical findings will show :

• Accumulation of very-long-chain fatty acids, phytanic acid

• Abnormalities in the synthesis of bile acids

• Marked reduction of plasmalogens

There is no appropriate treatment for Zellweger syndrome. It is very severe and infants rarely survive beyond a few months.

Solution to Question 25:

The correct statement about the action of enhancers is that they recruit chromatin and modify regulatory mechanisms.

A promoter is a sequence on DNA where RNA polymerase binds and acts as the start signal for transcription. They are present upstream or downstream but close to the gene that is undergoing transcription.

Enhancers or repressors are proteins that regulate the expression of a gene
by recruiting chromatin(DNA bound with histones) and modifying regulatory
mechanisms and making it more or less accessible to transcription by RNA polymerase. They may be located upstream or downstream and can be present at any distance from the promoter region. These are not specific to any single gene and interact with many other proteins.

Solution to Question 26:

Increased H+ ions in the intermembrane space of mitochondria are due to decreased oxidative phosphorylation.

ATP is synthesized in mitochondria by a process called oxidative phosphorylation. The oxidation of substrates like carbohydrates, amino acids, or fats produces reducing equivalents (NADH, FADH2). These equivalents enter the electron transport chain and transfer a proton (H+) to the inter-membrane space. The inner mitochondrial membrane is not permeable to protons leading to the accumulation of H+ in the intermembrane space. The flow of these protons through the Fo releases the energy that phosphorylates ADP to form ATP.

Drugs that inhibit oxidative phosphorylation like oligomycin inhibit the flow of protons through Fo causing an increase in H+ ions in the intermembrane space.

Solution to Question 27:

Fructosamine is obtained by the glycation of albumin.

Fructosamine is a measure of non-enzymatic glycation of circulating proteins including albumin, globulins, and lipoproteins. It is now used as a reasonable alternative to HbA1c measurement when it is not reliable. Since albumin is the most abundant of the serum proteins, fructosamine is predominantly a measure of glycated albumin.

Fructosamine is used as a short-term marker of glucose control. It reflects the average glycemia over the preceding 2 to 3 weeks. HbA1c reflects glucose control over a period of the preceding 8 to 12 weeks.

Biochemistry NEET 2022

Question 1:

A patient from a Mediterranean country visits Africa, where he develops malaria. He is treated with primaquine and later develops hemolytic anemia. Deficiency of an enzyme involved in which of the following pathways could be the cause?

a) Glycolysis

b) Gluconeogenesis

c) Hexose monophosphate (HMP) pathway

d) Luebering–Rapoport pathway

Question 2:

Which of the following disorders follows autosomal recessive inheritance pattern?

a) Huntington's disease

b) Treacher Collins syndrome

c) Cystic fibrosis

d) Achondroplasia

Question 3:

Which of the following helps in the transport of fatty acids across the inner mitochondrial membrane?

a) Acyl carrier protein

b) Carnitine

c) Lecithin-cholesterol acyltransferase

d) Carnitine and albumin

Question 4:

A single mutation in a nucleotide base pair resulting in a termination codon is known as?

a) Missense mutation

b) Nonsense mutation

c) Stop mutation

d) Silent mutation

Question 5:

DNA packing is done by which of the following ?

a) Histone

b) Glycoprotein

c) Nucleic acid

d) Adenine

Question 6:

An adolescent male patient presents to you with exercise intolerance. He gives a history of developing cramps on exertion. Which of the following enzyme deficiencies could be the cause?

a) Myophosphorylase

b) Hexokinase

c) Glucose-6-phosphatase

d) Hepatic glycogen phosphorylase

Question 7:

Which of the following enzyme activities can be estimated in red blood cells to diagnose vitamin B2 deficiency?

a) Transketolase

b) Glutathione reductase

c) Kynureninase

d) Pyruvate dehydrogenase

Question 8:

The following is due to the deficiency of?

a) Vitamin C

b) Vitamin A

c) Vitamin B

d) Vitamin E

Question 9:

A child presents to you with anemia, thrombocytopenia, and bony pain. On examination, hepatosplenomegaly was noticed. If light microscopy of bone marrow aspirate reveals "crumpled tissue paper" appearance, what is the enzyme defect?

a) Glucocerebrosidase

b) Sphingomyelinase

c) Hexosaminidase

d) Glucose 6-phosphatase

Question 10:

The electron transport chain is a series of redox reactions that result in ATP synthesis. Which of the following is a cytochrome complex IV inhibitor?

a) Cyanide

b) Carbon dioxide

c) Oligomycin

d) Ouabain

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	b
5	a
6	a
7	b
8	b
9	a
10	a

Detailed Explanations

Solution to Question 1:

The deficiency of glucose-6-phosphate dehydrogenase (G6PD) enzyme, which is involved in the hexose monophosphate (HMP) pathway, could be the cause behind this presentation.

G6PD is an enzyme that plays a critical role in redox reactions that occur in all aerobic cells. It is the only source of NADPH in erythrocytes, which helps in the formation of reduced glutathione. This reduced form protects erythrocytes against oxidative stress.

G6PD deficiency is an X-linked inheritance disorder. Most patients with this condition are asymptomatic. But when exposed to oxidative agents, they are prone to develop acute hemolytic anemia. The various oxidative triggers include:

• Fava beans

• Infections

• Drugs such as primaquine, dapsone, and nitrofurantoin

A hemolytic attack can present with weakness and abdominal/ lumbar pain. Later, the patient can develop jaundice, hemoglobinemia, and hemoglobinuria. Investigations will show elevated levels of lactate dehydrogenase (LDH) and low levels of plasma haptoglobin. Anisocytosis and spherocytes can be seen on peripheral smear. Acute renal failure is a complication seen in adults.

Other options:

Option A - Glycolysis, the main pathway of glucose, can occur either aerobically or anaerobically. A deficiency of the enzyme pyruvate kinase can present as hemolytic anemia. Fatigue can be the presentation when there is phosphofructokinase. Both these are enzymes of glycolysis.

Option B - Gluconeogenesis is the process by which non-carbohydrate precursors are used to synthesize glucose. Pyruvate carboxylase, phosphoenolpyruvate carboxykinase, and glucose-6-phosphatase are the enzymes in the pathway. Glucose-6-phosphatase deficiency leads to von Gierke disease, which is a glycogen storage disorder. Defective gluconeogenesis results in hypoglycemia.

Option D - Luebering–Rapoport pathway is a shunt mechanism that generates and dephosphorylates 2,3-bisphosphoglycerate (2,3-BPG). 2,3-BPG mutase is the enzyme involved. A deficiency of this enzyme can cause erythrocytosis.

Solution to Question 2:

Cystic fibrosis is a condition with an autosomal recessive mode of inheritance.

It results due to a defective mutation in the gene encoding for cystic fibrosis transmembrane conductance regulator (CFTR gene). It functions as an epithelial anion channel. It plays an important role in mucociliary transport along the respiratory mucosa. It also helps regulate the amount of exocrine gland secretion.

The defect in the ion channel results in the production of hyper viscous, adherent secretions which can obstruct the airways. Organisms such as Pseudomonas aeruginosa and Staphylococcus aureus can grow in the sputum. The exocrine pancreas can undergo fibrotic scarring, cyst formation, and acinar tissue loss. The viscid secretions can lead to pancreatic duct obstruction. The flow of digestive enzymes to the duodenum is hence impaired.

Symptoms include sinusitis, steatorrhea, and abdominal pain. Distal intestinal obstruction is a complication due to difficulty in excreting intestinal contents. In neonates, it can present as meconium ileus.

Detection of high levels of trypsinogen is used in newborn screening. Chloride level >60mEq/L in the sweat of affected patients is diagnostic. Hyponatremic hypochloremic alkalosis and hypokalemia may also be seen.

Postural drainage, short-term antibiotics, and inhaled bronchodilators can be used for medical management. The definitive treatment is lung transplantation.

In autosomal recessive inheritance, multiple affected members are seen in the same generation. Other generations are not affected. The following pedigree chart depicts an autosomal recessive pattern of inheritance.



Other options:

Option A - Huntington's disease is a trinucleotide (CAG) repeat disorder with an autosomal dominant mode of inheritance. It is characterized by involuntary movements known as chorea.

Option B - Treacher Collins syndrome is an autosomal dominant disorder that presents as mandibular and malar hypoplasia.

Option D - Achondroplasia has an autosomal dominant inheritance pattern and presents with short limbs, saddle nose, and spinal deformity.

Solution to Question 3:

Carnitine is responsible for the transport of fatty acids or long-chain acyl-CoA across the inner mitochondrial membrane.

Fatty acids are oxidized in the mitochondria to produce acetyl-CoA to produce energy. They must first be converted to acyl-CoA (active fatty acid) before they can be catabolized.

Carnitine palmitoyltransferase - I (CPT- I) transports acyl-CoA across the inner mitochondrial membrane only in the presence of carnitine. The acyl group of acyl-CoA is transferred to carnitine. The acylcarnitine so formed can penetrate the inner mitochondrial membrane via the carnitine-acylcarnitine translocase. This transporter transfers the acyl group back to CoA, restoring acyl-CoA and liberating carnitine, which is moved back to the intermembrane space. Carnitine palmitoyltransferase -II catalyzes this reaction. The deficiency of this enzyme is a common cause of recurrent myoglobinuria.

Carnitine deficiency can lead to hypoglycemia and restrictive or dilated cardiomyopathy. It can also cause hepatic steatosis.



Other options:

Option A - Acyl carrier protein is a component of the fatty acid synthase complex involved in the biosynthesis of fatty acids. It has vitamin pantothenic acid (vitamin B5).

Option C - Lecithin-cholesterol acyltransferase (LCAT) is present in the plasma and plays an important role in the esterification of cholesterol. Deficiency leads to low levels of lysolecithin and cholesteryl esters.

Option D - The long-chain fatty acids are bound to albumin in the plasma. But albumin does not transport fatty acids across the inner mitochondrial membrane. Albumin is a negative acute-phase reactant synthesized by the liver. It is a carrier for acidic drugs.

Solution to Question 4:

A single mutation in a nucleotide base pair resulting in a termination codon is known as a nonsense mutation.

A codon is a sequence of three bases present on the mRNA (messenger RNA), and it codes for amino acids. Out of the 64 codons possible, only 61 code for the 20 amino acids. Three codons, UAA, UAG, and UGA do not code for any of the 20 amino acids. These are known as nonsense/punctator/termination/stop codons and function as translation termination signals. The UGA codon is traditionally considered a stop codon, but it codes for selenocysteine (21st amino acid).

The tRNA (transfer RNA) contains the anticodon which recognizes specific codons present on the mRNA. The sequence of the anticodon is complementary to the codon. The pairing between the codon and anticodon can wobble at the third base. For example, GGU, GGC, and GGA all pair with the anticodon CCI. The wobble phenomenon along with degeneracy reduces the effect of mutations.

A mutation involving a single nucleotide base is known as a point mutation. A nonsense mutation is a type of point mutation where a codon previously coding for an amino acid is changed to a stop

codon. This results in premature termination of translation and the formation of a truncated protein. mRNA containing premature stops are removed by the nonsense-mediated degradation pathway.

Other options:

Option A: A missense mutation is a type of point mutation where the new codon, formed by the changed base, codes for a different amino acid.

Option C: Stop mutation is a term that can be used interchangeably with nonsense mutation. Here, nonsense mutation is the best answer.

Option D: Silent mutations are point mutations where the new codon formed still codes for the same amino acid as the codon before the mutation.

Solution to Question 5:

The proteins involved in DNA packing are histones.

Chromatin contains double-stranded DNA molecules and small basic proteins known as histones. In addition, nonhistone proteins are present that are involved in DNA replication and repair, and RNA synthesis. Histones play a role in condensing the DNA.

Nucleosomes are the organizational unit of chromatin and contain an octamer of histones around which 150 base pairs of DNA are packed. There are various types of histone proteins. H2A, H2B, H3, and H4 groups are components of nucleosomes and are known as core histones. They undergo post-translational modifications such as acetylation and methylation. This plays a critical part in chromatin structure and function.

The DNA in the nucleosome is arranged in a left-handed helical configuration over the histone octamer.



Other options:

Option B - Glycoproteins are proteins with oligosaccharide chains (glycans) bound to amino acids. Certain plasma proteins, peptide hormones, and cell membrane proteins are examples. Structural changes in the glycoproteins present in cancer cells may play a role in metastasis.

Option C - Nucleic acids can be either DNA or RNA. DNA is thought to be arranged in a double-stranded helix model. RNA on the other hand exists as a single strand. There are different types of RNA such as mRNA (messenger RNA), tRNA (transfer RNA), and rRNA (ribosomal RNA).

Option D - Adenine is a purine base that when linked to ribose forms adenosine, a nucleoside. When a nucleoside is esterified with a phosphate group a nucleotide is formed. Guanine is the other purine base. Cytosine is the pyrimidine base common to DNA and RNA. Thymine is present only in DNA and Uracil in RNA.

Solution to Question 6:

The clinical scenario points to a diagnosis of McArdle disease, which results due to a deficiency of the enzyme myophosphorylase (muscle glycogen phosphorylase).

McArdle disease, also known as type V glycogen storage disorder, is an autosomal recessive condition. It typically presents during late childhood or in the second decade of life. The muscle glycogen phosphorylase enzyme deficiency interferes with ATP generation by glycogenolysis in the muscles. This causes muscle glycogen accumulation.

Symptoms are exercise intolerance associated with muscle cramps. A second wind phenomenon, where the symptoms subside after a short period of rest, can also be seen. Myoglobinuria secondary to rhabdomyolysis leads to the passage of burgundy-colored urine in some patients.

Rarely, it can present during infancy as hypotonia, muscle weakness, and respiratory complications. It is generally fatal.

Muscle biopsy to measure glycogen content is the standard diagnosis. Avoiding strenuous exercise prevents symptoms. Glucose administration before exercise can improve tolerance. Vitamin B6 is also recommended.

Other options:

Option B - Hexokinase is a glycolytic enzyme. Its deficiency leads to hemolytic anemia. Splenomegaly may be another presenting feature.

Option C - Glucose-6-phosphatase is an enzyme involved in gluconeogenesis. Its deficiency leads to von Gierke disease also known as type I glycogen storage disease.

Option D - Hepatic glycogen phosphorylase deficiency results in type VI glycogen storage disease, also known as Hers disease.

Solution to Question 7:

Glutathione reductase activity in red blood cells can be estimated to diagnose vitamin B2 deficiency.

Vitamin B2, also known as riboflavin, is a water-soluble vitamin. Its active forms are FMN (flavin mononucleotide) and FAD (flavin adenine dinucleotide). Glutathione reductase is a FAD-containing enzyme (flavoenzyme). Hence, riboflavin deficiency can be detected by measuring erythrocyte glutathione reductase activity. Red blood cell and urinary concentrations of riboflavin can also be estimated to detect a deficiency.

Deficiency of riboflavin can present as anemia, corneal vascularization, mucocutaneous lesions, and personality changes. Milk and other dairy products are important dietary sources.

Other options:

Option A - Transketolase contains thiamine diphosphate as a coenzyme. Estimation of erythrocyte apotransketolase (enzyme protein) after in vitro addition of thiamine diphosphate, can be used to assess thiamine nutritional status. Thiamine deficiency can result in peripheral neuritis, beriberi, and Wernicke encephalopathy with Korsakoff psychosis.

Option C - Kynureninase is an enzyme that requires pyridoxal phosphate as a coenzyme. It converts 3-hydroxykynurenine, a tryptophan metabolite, to 3-hydroxyanthranilate. In vitamin B6 deficiency, this is impaired and results in the formation and subsequent urinary excretion of xanthurenate. This can be used for diagnosis.

Option D - Pyruvate is decarboxylated by pyruvate dehydrogenase for which thiamine diphosphate acts as the cofactor. Therefore thiamine deficiency results in the accumulation of pyruvate. This leads to pyruvic and lactic acidosis, commonly seen in alcoholics.



Solution to Question 8:

The given image depicts Bitot's spots characteristically seen in deficiency of vitamin A. Bitot's spots are seen as white patches of keratinized epithelium on the sclera. It is more commonly bilateral and on the temporal side.

Vitamin A deficiency arises mostly due to the inadequate amounts of vitamin A in diet. It can present as night blindness, conjunctival xerosis, Bitot's spots, and keratomalacia. In later stages, it

can lead to xerophthalmia. It also increases the risk of developing measles, diarrhea, and dysentery. Skin manifestations include xerosis, scaling, and dermomalacia. Phrynoderma may also be seen as keratotic follicular papules affecting the thighs and upper arms. It is shown in the image below.

In patients above the age of 1 year, vitamin A deficiency is treated with 200,000 IU of vitamin A orally or 100,000 IU by intramuscular route. It should be repeated the next day and two weeks later.

Vitamin A is a fat-soluble vitamin that plays an essential role in vision, reproduction, and maintenance of epithelial integrity. The various forms of vitamin A include retinol (storage form), retinal, retinoic acid, and beta-carotene. It is stored in the liver perisinusoidal stellate (Ito) cells as retinyl ester (retinyl palmitate).

Other options:

Option A - Vitamin C is a water-soluble vitamin that aids in iron absorption and collagen maturation. It cannot be synthesized in humans because of the absence of L-gulonolactone oxidase. Vitamin C deficiency results in scurvy.

Option C - Vitamin B deficiency has clinical manifestations pertaining to each type.

Option D - Vitamin E deficiency can present as posterior column and spinocerebellar symptoms. Peripheral neuropathy, ophthalmoplegia, and pigmented retinopathy are other features seen.

Solution to Question 9:

The clinical scenario points to a diagnosis of Gaucher's disease, where there is a defect in the enzyme glucocerebrosidase.

Gaucher's disease is a lysosomal storage disease with autosomal recessive inheritance. The condition results due to the deficiency of beta-glucocerebrosidase. This results in the accumulation of glucosylceramide (glycolipid substrate) in the reticuloendothelial cells. There is progressive infiltration of the liver, spleen, and bone marrow.

There are three clinical types:

• Type 1: Adult (non-neuronopathic) form

• Type 2: Infantile (acute neuronopathic) form

• Type 3: Juvenile (subacute neuronopathic) form

Type 1 is the most common. It presents with a myriad of symptoms, such as bruising secondary to thrombocytopenia, fatigue due to anemia, hepatosplenomegaly, and bone pain. Pathologic fractures and lytic lesions of the femur, ribs, and pelvis may also be seen. A few patients may have pulmonary involvement. Growth retardation may be observed in cases presenting in the first decade.

Type 2 is rare, and presents with hypertonia, strabismus, and organomegaly. Neurodegeneration is the characteristic feature. Stridor and failure to thrive can also be the presenting features. Ultimately, death can occur due to respiratory arrest.

Type 3 presents with intermediate symptoms, as seen in the other two types. By the age of 10–15 years, the child succumbs to death.

The Gaucher cells, which are the reticuloendothelial cells with glucosylceramide, have a wrinkled-paper appearance due to intracytoplasmic substrate inclusions. These cells stain strongly with periodic acid-Schiff (PAS) stain. This is characteristic of Gaucher's disease.

Other options:

Option B - Acid sphingomyelinase deficiency results in Niemann-Pick disease types A and B. Both are autosomal recessive conditions. Type A has a neurodegenerative course, while type B is non-neuronopathic. The former presents with hepatosplenomegaly, lymphadenopathy, and psychomotor retardation. Death occurs by the age of 3 years. Type B is associated with hepatosplenomegaly and mild pulmonary involvement.

Option C - Hexosaminidase deficiency causes Sandhoff disease. It has an autosomal recessive pattern of inheritance. Clinical features include ataxia, dysarthria, and mental retardation.

Option D - Glucose-6-phosphatase deficiency leads to von Gierke disease, which is a glycogen storage disorder. It is an autosomal recessive condition. The affected children have characteristic doll-like facies. Hyperuricemia, lactic acidosis, and triglyceridemia can be seen. Hepatomegaly and renal enlargement is also associated.

Solution to Question 10:

Cyanide is an inhibitor of cytochrome complex IV or cytochrome a+a3 (cytochrome c oxidase). It is a component of the electron transport chain.

The electron transport chain (ETC) is a specialized set of electron carriers that can accept electrons. Energy-rich molecules like glucose are oxidized to produce water and carbon dioxide. The intermediate substrates in these reactions can pass on electrons to certain coenzymes such as nicotinamide adenine dinucleotide (NAD+) and flavin adenine dinucleotide ($FADH_2$). The reduced forms can then donate electrons to the ETC by which they lose most of their free energy which is used to move H^+ through the inner mitochondrial membrane. The H^+ gradient thus produced results in the production of ATP. ATP synthesis together with electron transport is

known as oxidative phosphorylation.

The reactions in oxidative phosphorylation and the arrangement of complexes that form the ETC are shown below.



These complexes can accept or donate electrons to coenzyme Q (CoQ) and cytochrome c, which are mobile electron carriers. Certain site-specific inhibitors can inhibit ATP synthesis.

Other options:

Option B - Carbon dioxide along with water is produced by the oxidation of energy-rich molecules in oxidative phosphorylation.

Option C - Oligomycin, an antibiotic, blocks the flow of protons through ATP synthase. This inhibits oxidative phosphorylation.

Option D - Ouabain is a cardiac drug that inhibits the Na+ - K+ ATPase of cell membranes by acting on the extracellular surface.

Complexes of ETC	Inhibitors
Complex I	Piericidin A, amobarbital, rot enone
Complex II	TTFA (thenoyltrifluoroaceton e), carboxin, malonate
Complex III	BAL(British anti-lewisite), an timycin A
Complex IV	Hydrogen sulfide, carbon mo noxide,cyanide

Biochemistry NEET 2023

Question 1:

Which of the following zymogens requires gamma-glutamyl carboxylase?

a) Factors II, VIII, IX, X

b) Factors II, VII, IX, and X

c) Factors II, VII, IX, XI

d) Factors II, VIII, X, XI

Question 2:

A patient presented with diarrhea, dermatitis, and dementia. Which of the following vitamin deficiencies would you suspect in this patient?

a) Vitamin B3 deficiency

b) Vitamin B1 deficiency

c) Vitamin B6 deficiency

d) Vitamin B12 deficiency

Question 3:

A young boy presents to the OPD with hypoglycemia and muscle cramps, on exertion or while playing. Then he becomes normal after resting for a while. These episodes are recurrent after a period of activity. He has decreased serum lactate and glucose levels. Which of the following diseases is he most likely to be suffering from?

a) McArdle disease

b) Hers disease

c) Cori's disease

d) Andersen disease

Question 4:

A family consumes only polished rice. Which of the following combination of vitamin deficiency and enzymatic defect will be present in this family?

a) Riboflavin-glutathione reductase

b) Thiamine-transketolase

c) Thiamine-transaminase

d) Riboflavin-transketolase

Question 5:

A chronic alcoholic patient presented to the casualty with confusion, ataxia, and painful eye movements. What could be the vitamin deficiency seen in this patient?

a) Vitamin B deficiency

b) Vitamin A deficiency

c) Vitamin D deficiency

d) Vitamin C deficiency

Question 6:

A 4-year-old child presents with easy fatiguability. The mother also complains that the child has increased hunger between meals, which is relieved after food. Liver examination revealed no glycogen. The enzyme most likely deficient is _____.

a) Glycogen phosphorylase

b) Glycogen synthase

c) Debranching enzyme

d) Glucose 6 phosphatase

Question 7:

A child was brought with pedal edema and cheilosis. Cardiomegaly was present. What is the vitamin deficiency associated with this clinical presentation?

a) Riboflavin

b) Thiamine

c) Pyridoxine

d) Niacin

Question 8:

A chronic alcoholic patient is diagnosed with gouty arthritis. What biochemical changes will be seen in this patient?

a) Decreased NADH/NAD+ ratio

b) Increased urea and urate levels

c) Increased urate and lactate levels

d) Increased level of alkaline phosphatase

Question 9:

A child presents with erythematous scaly patches in the perioral region, mucosal ulcers, and impaired epithelial wound healing. The most likely mineral deficiency associated with this condition is _____.

a) Iron deficiency

b) Zinc deficiency

c) Calcium deficiency

d) Copper deficiency

Question 10:

A patient presenting with bleeding gums and easy bruisability was diagnosed with scurvy. This is due to _____.

a) Low calcium

b) Defective collagen formation

c) Inhibition of clotting factors

d) Increased keratinization of epithelium

Question 11:

Which of the following is seen in low insulin:glucagon ratio?

a) Activation of lipoprotein lipase

b) Activation of glycogen synthase

c) Activation of phosphofructokinase 1

d) Activation of hormone-sensitive lipase

Question 12:

An infant is brought by his parents with complaints that his urine turns black on standing. Which of the following metabolic disorders is likely?

a) Phenylketonuria

b) Alkaptonuria

c) Homocystinuria

d) Maple syrup urine disease

Question 13:

A child presents with bone pain and hepatosplenomegaly. A trephine biopsy and aspirate show the following finding. Which of the following is the most likely enzyme deficient in this condition?



a) Hexosaminidase

b) Glucocerebrosidase

c) Sphingomyelinase

d) Alpha 1,4-glucosidase

Question 14:

A child presents with thinning of hair and scaling of the scalp. On taking further history, it is found that she consumes raw eggs in her diet regularly. The given manifestations are due to the deficiency of which of the following vitamins?

a) Niacin

b) Pyridoxine

c) Biotin

d) Thiamine

Question 15:

A child presents with mental retardation, bone pain, and inability to walk. On funduscopy examination, a cherry red spot is seen. There is no organomegaly. What is the most likely diagnosis?

a) Tay-Sachs disease

b) Niemann-Pick disease

c) Hurler syndrome

d) Gaucher disease

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	a
6	b
7	b
8	c
9	b
10	b
11	d

12	b
13	b
14	c
15	a

Detailed Explanations

Solution to Question 1:

Clotting factors II, VII, IX, and X are zymogens of serine proteases that contain glutamic acid residues. These residues undergo vitamin-K-dependent carboxylation via gamma-glutamyl carboxylase to form gamma-carboxyglutamic acid residues. These residues chelate calcium ions, allowing the binding of clotting factors to the platelet surface.

Zymogens are the proprotein form of enzymes (proenzymes) that require proteolytic activation to carry out their function.

The reduced form of vitamin K acts as a cofactor in the post-synthetic modification of these zymogens and their activation. This reduced form of vitamin K is recycled back from its oxidized form by the enzyme vitamin K epoxide reductase, which is inhibited by warfarin.

The absence of vitamin K or the presence of warfarin inhibits carboxylation of the glutamate residues, leading to the release of abnormal precursors of clotting factors in the blood.



Solution to Question 2:

The given clinical scenario of a patient with diarrhea, dermatitis, and dementia is suggestive of niacin/vitamin B3 deficiency.

Niacin deficiency causes pellagra, found mostly among people eating corn/maize-based diets, which are low in niacin and tryptophan. Excess leucine in jowar and maize disrupts the conversion of tryptophan to niacin.

Niacin can be synthesized in the body from tryptophan. Two compounds, nicotinic acid and nicotinamide, have the biological activity of niacin. The nicotinamide ring of the coenzymes nicotinamide adenine dinucleotide (NAD) and nicotinamide adenine dinucleotide phosphate (NADP) participate in oxidation/reduction reactions.

The initial symptoms of pellagra include loss of appetite, generalized weakness, irritability, abdominal pain, and vomiting. This is followed by bright red glossitis and a characteristic pigmented, scaly skin rash in sun-exposed areas. This rash is known as Casal's necklace when it forms a ring around the neck.

The 4Ds of pellagra:

• Diarrhea

• Dermatitis

• Dementia, leading to

• Death

It is diagnosed by measuring urinary levels of 2-methyl nicotinamide and 2-pyridone.

It is treated by oral supplementation of 100–200 mg of nicotinamide or nicotinic acid three times daily for 5 days.

Casal's necklace finding in niacin deficiency:

Other options:

Option B: Vitamin B1/Thiamine deficiency can manifest as wet beriberi (CVS features like heart failure and edema) or Shoshin beriberi, an acute and fulminant form of wet beriberi, dry beriberi (peripheral neuritis), or as Wernicke encephalopathy with Korsakoff psychosis (in alcoholics). Dietary deficiency of thiamine inhibits pyruvate dehydrogenase. Chronic alcoholics may develop potentially fatal pyruvic and lactic acidosis.

Option C: Vitamin B6/Pyridoxine deficiency is rare but has been observed in alcoholics, newborns, formulas low in vitamin B6, and women taking oral contraceptives. Isoniazid, an anti-TB drug, leads to pyridoxine deficiency resulting in peripheral neuropathy. Hence pyridoxine supplementation is included in ATT regimens.

Option D: Vitamin B12/Cobalamin deficiency causes megaloblastic anemia, dementia, spinal cord degeneration, homocysteinemia, etc.

Solution to Question 3:

A clinical scenario of a patient with muscle cramps, pain, weakness with exertion, myoglobinuria, and the muscle biopsy specimen showing vacuolar myopathy are suggestive of McArdle's disease, which is due to the deficiency of muscle glycogen phosphorylase.

McArdle disease, also known as type V glycogen storage disorder, is an autosomal recessive condition. It typically presents during late childhood or in the second decade of life. The muscle glycogen phosphorylase enzyme deficiency interferes with ATP generation by glycogenolysis in the muscles. This causes muscle glycogen accumulation.

Symptoms are exercise intolerance associated with muscle cramps. Due to high-intensity exercise, sudden persistent muscle contraction occurs. This leads to severe muscle damage resulting in a massive release of muscle proteins and myoglobin in the blood. Myoglobinuria secondary to rhabdomyolysis leads to the passage of burgundy-colored urine in some patients.

A unique feature associated with McArdle is the improvement in symptoms after rest for a few minutes. This is termed as the second wind phenomenon. This occurs due to the increased blood flow to the muscles and increased utilization of free fatty acids (FFAs) for energy production after resting for a few minutes. This compensates for the impaired ATP generation from glycogenolysis. Patients with McArdle disease can benefit from engaging in regular aerobic exercises, such as walking or cycling, which can help increase their exercise tolerance and cardiovascular fitness. However, they should avoid high-intensity exercises or activities that can trigger muscle cramps and rhabdomyolysis. Glucose administration before exercise can improve tolerance. Vitamin B6 is also recommended.

Standard diagnosis includes muscle biopsy, which shows periodic acid Schiff (PAS)-positive vacuoles (Shown in the image below) of high glycogen content due to lack of myophosphorylase and genetic testing (PYGM gene sequencing).



Other options:

Option B: Hers disease (type VI glycogen storage disorder) - It is caused by the deficiency of liver glycogen phosphorylase. Patients with Hers disease exhibit growth retardation, hepatomegaly, and mild hypoglycemia. They do not typically present with muscle cramps and myoglobinuria.

Option C: Cori's disease (type III glycogen storage disorder) - It is caused by a deficiency of glycogen debranching enzyme. Patients with Cori's disease present with hepatomegaly, growth retardation, and hypoglycemia. Muscle weakness may be present but not muscle cramps or myoglobinuria.

Option D: Andersen disease (type IV glycogen storage disorder) - It is caused by the deficiency of glycogen branching enzyme. This disease presents with hepatomegaly, cirrhosis, and liver failure. It does not cause muscle cramps or myoglobinuria.

Solution to Question 4:

The consumption of polished rice leads to thiamine deficiency which leads to defective transketolase activity.

Polished rice is rice that has had its husk, bran, and germ removed, leaving only the endosperm. This processing method results in a loss of nutrients, including thiamine.

Thiamine pyrophosphate (Vit B1) is the cofactor of transketolase. This reaction requires two cofactors:

• Mg^{2+}

• Thiamine diphosphate (vitamin B1)

Transketolase and transaldolase are involved in the non-oxidative phase of the HMP pathway.

Thiamine deficiency can lead to beriberi, a condition characterized by peripheral neuropathy, weakness, and cardiovascular symptoms. In severe cases, it can also lead to Wernicke-Korsakoff syndrome, which is characterized by confusion, ataxia, and vision disturbances. Thiamine

deficiency can be prevented by consuming a balanced diet that includes unprocessed grains, legumes, and other sources of thiamine. Diagnosis of thiamine deficiency can be established by erythrocyte transketolase activity assessment.

Solution to Question 5:

The above clinical history is suggestive of Wernicke encephalopathy, which is caused by vitamin B1 (Thiamine) deficiency.

Vitamin B1/thiamine deficiency leads to following conditions:

• Wernicke encephalopathy:

• An acute, neuropsychiatric syndrome characterized by nystagmus and ophthalmoplegia, mental status changes (confusion), and unsteadiness of stance and gait.

• Wernicke's encephalopathy is a medical emergency, and thiamine should be initiated immediately, either intravenously or intramuscularly.

• The current standard of treatment for patients with symptoms of Wernicke's encephalopathy is to give thiamine 100 mg intravenously before administering glucose-containing IV fluids.

• Korsakoff psychosis:

• Korsakoff syndrome is characterized by a pronounced anterograde and retrograde amnesia along with potential impairment in visuospatial, abstract, and other types of learning.

• Symptoms are similar to that of Wernicke encephalopathy, but also include memory loss and confabulation.

• Permanent in at least a partial form in about 50% of the people affected.

• Beriberi: It can present with:

• Dry beriberi: sensorimotor peripheral neuropathy, muscle weakness

• Wet beriberi: tachycardia, cardiomegaly, high-output cardiac failure

Solution to Question 6:

The most likely enzyme deficiency in a child with fatigue, excessive hunger between meals, and absent liver glycogen is glycogen synthase deficiency.

Glycogen is a storage carbohydrate synthesized from glucose (glycogenesis) and is stored in the liver, muscles, and brain. Glycogen synthase is a regulating enzyme for glycogenesis.

Glycogen synthase exists in both phosphorylated (inactive) and dephosphorylated (active) forms. Phosphorylation is stimulated by epinephrine, norepinephrine, and glucagon via cAMP. Dephosphorylation is stimulated by insulin via phosphodiesterase.

A deficiency of glycogen synthase leads to glycogen storage disorder and presents in infancy with early morning drowsiness and fatigue, fasting hypoglycemia, pallor, emesis, and hypoglycemic convulsions. It leads to a marked decrease in liver glycogen stores. Hepatomegaly is absent.

Laboratory investigations reveal prolonged hyperglycemia, glycosuria, ketosis, lactic acidosis, low alanine levels, and normal insulin levels after glucose administration. A liver biopsy is diagnostic.

Treatment involves protein-rich frequent meals, and nighttime supplementation with cornstarch to prevent hypoglycemia and hyperketonemia.

Other options:

Option A: Glycogen phosphorylase deficiency, also known as McArdle's disease or glycogen storage disease type V, primarily affects skeletal muscle glycogen metabolism. Affected individuals experience muscle weakness, cramps, and exercise intolerance, but it doesn't typically cause the symptoms mentioned in the question.

Option C: Debranching enzyme deficiency, also known as Cori's disease or glycogen storage disease type III, presents with hepatomegaly, hypoglycemia, and growth retardation. The liver would still contain glycogen, albeit with an abnormal structure.

Option D: Glucose-6-phosphatase deficiency, known as von Gierke's disease or glycogen storage disease type I, is characterized by severe fasting hypoglycemia, lactic acidosis, hyperuricemia, and hepatomegaly. In this case, glycogen is present in the liver but cannot be broken down efficiently into glucose.

Solution to Question 7:

The given clinical presentation with pedal edema and cardiomegaly is suggestive of wet beriberi due to thiamine (vitamin B1) deficiency.

Early symptoms of thiamine deficiency include peripheral neuropathy, irritability, loss of deep tendon reflexes and proprioception. As the disease progresses, increased intracranial pressure, meningismus and coma develop.

Wet beriberi results due to cardiac and renal dysfunction which leads to fluid accumulation. It is associated with tachycardia, cardiomegaly, cardiac failure, and edema. It is commonly seen in the refugee camps where people consume polished rice-based monotonous diets.

Dry beriberi is associated with neurological deficits such as peripheral neuropathy, optic atrophy, ataxia, and irritability. Complications include lactic acidosis and Wernicke encephalopathy in adults.

Diagnosis of thiamine deficiency can be established by erythrocyte transketolase activity assessment.

Other options:

Option A: Riboflavin deficiency can present with cheilosis, but it does not typically cause pedal edema and cardiomegaly.

Option C: Pyridoxine (vitamin B6) deficiency can cause neurological symptoms, anemia, and dermatitis but is not associated with pedal edema, cardiomegaly, and cheilosis.

Option D: Niacin (vitamin B3) deficiency leads to pellagra, characterized by the triad of dermatitis, diarrhea, and dementia. It is not associated with pedal edema, cardiomegaly, and cheilosis.

Solution to Question 8:

This patient with chronic alcoholism and gouty arthritis will most likely have elevated levels of lactate and urate.

The oxidation of ethanol to acetate by alcohol dehydrogenase is accompanied by the reduction of nicotinamide adenine dinucleotide (NAD+) to NADH. This results in a high concentration of NADH, leading to an increased NADH/NAD+ ratio (Option A).

Due to this, the following metabolic reactions that require NAD+ are inhibited:

• Glycolysis

• Citric acid cycle (ketogenesis favored)

• Pyruvate dehydrogenase

• Fatty acid oxidation

Decreased fatty acid oxidation leads to increased esterification and production of triacylglycerol, which causes fatty liver.

Also, there is an increase in two reactions:

• Pyruvate to lactate

• Oxaloacetate to malate

Increased lactate leads to hyper-lactic-acidemia, which leads to increased renal reabsorption of urate and decreased excretion of uric acid. This results in aggravation of gout. Alcoholics are deficient in thiamine which inhibits pyruvate dehydrogenase and leads to lactic acidosis.

A decrease in pyruvate and oxaloacetate levels, which are substrates for gluconeogenesis, leads to a decrease in blood glucose levels, causing symptoms of agitation, impaired judgment, and combativeness.

Other options:

Increased urea and urate levels are not specifically associated with alcoholism and gouty arthritis. While urate levels may be increased due to gout, urea levels are not directly related to the condition (Option B).

Increased levels of alkaline phosphatase can be seen in alcohol-induced hepatitis due to intrahepatic cholestasis but is not directly related to gouty arthritis in a chronic alcoholic patient. Therefore, this option is less relevant compared to increased urate and lactate levels (Option D).

Solution to Question 9:

The clinical features of periorificial dermatitis and stomatitis in a young child are suggestive of acrodermatitis enteropathica, seen in zinc deficiency.

Acrodermatitis enteropathica is a genetic disorder that results in the inability to absorb dietary zinc.

The clinical features develop after weaning and the infant presents with:

• Failure to thrive

• Skin manifestations such as symmetrical vesiculobullous dermatitis in the perioral, perianal, and acral regions

• Anorexia, diarrhea

• Recurrent infections due to T-cell immunodeficiency

• Alopecia with a reddish tint

• Impaired wound healing

• Male hypogonadism, growth retardation

• Neurological manifestations like tremors, nystagmus, photophobia, hypogeusia, irritability, and night blindness

Oral zinc supplementation is the treatment of choice.

Other options:

Option A- Iron deficiency is the most common micronutrient deficiency globally. It presents with iron-deficiency anemia and neurocognitive impairment. It presents clinically with pallor, fatigue, atrophic glossitis, and pica.

Option C- Calcium deficiency leads to rickets as a result of early weaning. It presents with typical symptoms of rickets, including failure to thrive; protruding abdomen; muscle weakness; fractures; craniotabes; delayed dentition; caries; enlargement of wrists and ankles; and hypocalcemic symptoms of tetany, stridor, and seizures.

Option D- Copper deficiency leads to anemia, growth retardation, defective keratinization and pigmentation of hair, hypothermia, degenerative changes in aortic elastin, osteopenia, and mental retardation.

Solution to Question 10:

Bleeding gums and easy bruisability seen in scurvy are due to defective collagen formation.

Scurvy results from the deficiency of vitamin C, which is required for the hydroxylation of prolyl and lysyl residues of collagen reactions. Hence, deficiency of vitamin C results in a defective prolyl hydroxylase activity, leading to defective collagen formation and resulting in generalized bleeding tendencies.

Hemorrhagic manifestations of scurvy include petechiae, purpura, ecchymoses at pressure points, epistaxis, gum bleeding, and characteristic perifollicular hemorrhages. Other manifestations are poor wound and fracture healing, hyperkeratosis of hair follicles, arthralgia, and muscle weakness. It also results in defective osteoid formation. Costo-chondral separation leads to the formation of tender nodules that are sharply angulated and are also known as the scorbutic rosary.

The cutaneous manifestations of scurvy include

• Phrynoderma over posterolateral arms which can then generalize over the back and thighs and legs.

• Corkscrew hairs - fractured and coiled hairs due to impaired keratin cross■links by disulfide bonds

• Perifollicular hemorrhage and purpura

• Lower limb edema

• Splinter hemorrhages

It is treated with vitamin C supplementation.

Other options:

Option A: Low calcium levels do not cause scurvy; they may contribute to other clinical conditions like osteoporosis or muscle cramps.

Option C: Inhibition of clotting factors is not responsible for scurvy. Vitamin K deficiency, however, can lead to the inhibition of clotting factors and cause bleeding tendencies.

Option D: Increased keratinization of epithelium is not the cause of scurvy. Vitamin A deficiency, on the other hand, can result in hyperkeratoses and epithelial changes, such as xerosis and Bitot's spots.

Solution to Question 11:

Low insulin:glucagon ratio occurs in the fasting state. Since insulin has an inhibitory effect on hormone-sensitive lipase (lipolytic hormone), low insulin:glucagon ratio results in the activation of hormone-sensitive lipase.

In adipose tissues, decreased insulin results in the inhibition of lipogenesis and activation of intracellular hormone-sensitive lipase. The stored triglycerides are broken down (lipolysis) by hormone-sensitive lipase (HSL) into free fatty acids (FFAs) and glycerol. In the fasting state, glucose is spared for the brain and glycerol is used as a substrate for gluconeogenesis.

In diabetic patients, low insulin level corresponds to an increase in HSL activity, promoting the breakdown of lipids and the release of FFAs from adipose tissue. HSL activity is accelerated by hormones like epinephrine, norepinephrine, glucagon, ACTH, TSH, GH, ADH, and melanocyte-stimulating hormones (MSH).



Other options:

Option A: Activation of lipoprotein lipase (LPL) is promoted by insulin, not low insulin: glucagon ratio. LPL is involved in the breakdown of triglycerides and is active in the well-fed state. In low insulin: glucagon ratio, lipoprotein lipase activity is reduced, leading to decreased hydrolysis of triglycerides.

Option B: Activation of glycogen synthase is promoted by insulin. In low insulin: glucagon ratio, glycogen synthase activity is reduced, leading to decreased glycogenesis. Instead, glycogenolysis and gluconeogenesis are promoted to maintain blood glucose levels.

Option C: Activation of phosphofructokinase 1 (PFK1) is promoted by insulin. PFK1 is a rate-regulating enzyme in glycolysis. In low insulin: glucagon ratio, glycolysis is downregulated, and gluconeogenesis is upregulated to maintain blood glucose levels.

Well-fed state (Insulin)	During fasting (Glucagon)
Lipogenesis increased	Lipogenesis inhibited
Lipolysis inhibited	Lipolysis increased
Insulin inhibits HS-lipase	Glucagon activates HS-lipase
Lipoprotein lipase active	Lipoprotein lipase inactive

Solution to Question 12:

Blackening of urine on standing is characteristic of alkaptonuria.

Alkaptonuria is an autosomal recessive disorder of tyrosine metabolism, that develops due to the defect in homogentisate oxidase/homogentisic acid 1, 2 - dioxygenase, which oxidizes homogentisate to maleylacetoacetate. The defect in this enzyme leads to the accumulation of homogentisate, which is then excreted in the urine.

Alkaptonuria is characterized by homogentisic aciduria, ochronosis, and arthropathy (joint pain).

Homogentisic aciduria is the black discoloration of urine on prolonged exposure to air caused due to oxidation of the excreted homogentisate.

Ochronosis, which is bluish-black pigmentation in connective tissue, generally occurs after the third decade of life, and it occurs due to the accumulation of homogentisic acid and its oxidation products like benzoquinone acetic acid in connective tissue.

Arthritis generally begins in the third decade. Arthritis often begins in the spine and resembles ankylosing spondylitis in its large-joint distribution. Radiographs of the spine showing flattened and calcified intervertebral discs are pathognomonic. Other findings include degeneration of the intervertebral discs, followed by disc calcification and eventually fusion of the vertebral bodies. Osteophyte formation and calcification of the intervertebral ligaments also occur.

Diagnosis: Blackish discoloration of urine on standing is diagnostic of alkaptonuria (shown below). The ferric chloride test will be positive. The blackening of urine occurs due to the excretion of homogentisic acid in the urine.

Image A - Normal urine

Image B - Alkaptonuria



Other options:

Option A: Phenylketonuria (PKU) is a metabolic disorder resulting from a deficiency of the enzyme phenylalanine hydroxylase. It does not lead to black urine. Patients with PKU have a musty or mousy odor due to phenylacetic acid accumulation.

Option C: Homocystinuria is an inborn error of methionine metabolism caused by a deficiency of the enzyme cystathionine β-synthase. It is characterized by tall stature, long limbs, dislocation of

the lens of the eye, and predisposition to thromboembolism, but it does not lead to black urine.

Option D: Maple syrup urine disease (MSUD) is an inherited metabolic disorder caused by a deficiency of branched-chain α-ketoacid dehydrogenase. It does not cause black urine. Patients with MSUD have urine with a characteristic sweet odor, resembling maple syrup.

Solution to Question 13:

The given clinical scenario of a child with bone pain, hepatosplenomegaly, and biopsy showing Gaucher's cell with a characteristic crumpled tissue paper appearance is suggestive of Gaucher's disease. The deficient enzyme is glucocerebrosidase (also known as beta-glucosidase).

Gaucher's disease is an inborn error of metabolism caused due to a deficiency of beta-glucocerebrosidase. It is the most common lysosomal storage disorder and the mode of inheritance is autosomal recessive. There is an accumulation of lipid-laden macrophages in the spleen, liver, and other tissues.

There are three clinical subtypes - type I (chronic non-neuronopathic), type II (acute neuronopathic), and type III (intermediate). Clinical features include anemia, thrombocytopenia, hepatosplenomegaly, and neurological features depending on the subtype.

Gaucher's cells - These are formed as a result of the accumulation of massive amounts of glucocerebrosides within the phagocytes. The cells have a fibrillary cytoplasm and give a crumpled tissue paper appearance.

Option A: Deficiency of hexosaminidase A is seen in Tay-Sachs disease. The deficiency of beta-hexosaminidase A and B causes Sandhoff's disease. Clinical features are seizures, macrocephaly, and hyperacusis.

Option C: Deficiency of sphingomyelinase leads to Niemann-Pick disease. It is characterized by the presence of foam cells in the bone marrow. There may be pulmonary infiltrates leading to lung failure.

Option D: Pompe's disease (type 2 glycogen storage disease) occurs due to alpha 1,4-glucosidase (acid maltase) deficiency. It is the only glycogen storage disease that is a lysosomal storage disease and causes lysosomal glycogen accumulation in cardiac, skeletal, and smooth muscle cells. The clinical symptoms of Pompe disease are hypotonia (floppy infant), skeletal muscle weakness, hepatomegaly, hypertrophic cardiomyopathy, and macroglossia.

Solution to Question 14:

The given clinical scenario of scaling and alopecia and the history of raw egg consumption are suggestive of biotin deficiency.

Raw egg white contains a glycoprotein known as avidin, which tightly binds to biotin and renders it unabsorbable from the intestine. Hence, there can be a biotin deficiency state.

Biotin deficiency may cause:

• Dermatitis, scaling, seborrheic, and erythematous rash

• Mental status changes (depression, hallucinations)

• Hyperesthesia

• Alopecia

• Atrophic glossitis

In infants, biotin deficiency presents as hypotonia, lethargy, and apathy. In addition, infants may develop alopecia and a characteristic rash that includes the ears.

Solution to Question 15:

The given clinical scenario, along with the fundoscopy image showing a macular cherry-red spot, indicates that the most likely diagnosis is Tay-Sachs disease.

Tay-Sachs disease is a lysosomal storage disease caused by the deficiency of β-hexosaminidase. It can present during infancy with macrocephaly, increased startle reaction, gradual loss of motor skills, and a macular cherry-red spot. Affected children usually die by the age of 10–15 years.

It is of two types—juvenile-onset and adult-onset. The symptoms of the juvenile-onset form include ataxia and dementia. Dysarthria, lower motor neuron signs, and psychiatric disorders are common in the adult-onset type.

Other options:

Options B - Niemann-Pick disease is an autosomal recessive inborn error of lipid metabolism. It results due to the deficiency of the sphingomyelinase enzyme. This results in the accumulation of sphingomyelin in the brain, lungs, liver, and spleen. The clinical features include a combination of hepatosplenomegaly, developmental delay, failure to thrive, severe psychomotor regression, interstitial lung disease, and a cherry-red spot in the macula.

Option C - Hurler syndrome (mucopolysaccharidosis I) is an autosomal recessive condition that affects multiple organs. Presentation is usually at 6–24 months with corneal clouding, coarse facial features, joint stiffness, and skeletal dysplasia. Glaucoma, retinal degeneration, and nasal discharge are the other features. The condition is generally fatal.

Option D - Gaucher disease is a glycolipid storage disorder that can present with hepatosplenomegaly, mental retardation, and erosion of long bones. It is caused by the deficiency of β-glucosidase. The Gaucher cell has a wrinkled-paper appearance due to intracytoplasmic inclusions, which is a characteristic feature.

Biochemistry INI-CET 2023

Question 1:

Xanthurenic aciduria is seen in which of the following vitamin deficiency?

a) Vitamin B2

b) Vitamin B1

c) Vitamin B12

d) Vitamin B6

Question 2:

A 12-year-old boy with type 1 diabetes mellitus presents with unconsciousness. His urine is subjected to a test and the results are shown below. What is the test done?

a) Seliwanoff's test- Fructose

b) Benedict's test- Reducing sugars

c) Fouchet's test- Bilirubin

d) Rothera's test- Ketone bodies

Question 3:

The template DNA strand is 5'CATTCAATATCGATC3'. The resulting mRNA sequence after transcription is?

a) 3'GAUCGAUAUUGAAUA5'

b) 5'GATCGATATTGAATUT3'

c) 5'GAUCGAUAUUGAAUG3'

d) 3'GATCGATATTGAATU5'

Question 4:

Arrange the following steps of Karyotyping in the correct order :

a) 1, 2, 3, 4, 5

b) 1, 3, 2, 4, 5

c) 1, 2 , 3, 5, 4

d) 2, 1, 3, 4, 5

Question 5:

A patient came with complaints of muscle weakness to the OPD. The doctor found that he has cardiomyopathy and diagnosed him to be a case of Barth Syndrome, where the patient has mitochondrial dysfunction. What is the defect in this patient?

a) Cephalin

b) Cardiolipin

c) Lecithin

d) Lysolecithin

Question 6:

In Pompe's disease, the enzyme deficient is?

a) Lactic acid dehydrogenase

b) Phosphofructokinase

c) Muscle glycogen phosphorylase

d) Lysosomal alpha glucosidase

Question 7:

The thermic effect is highest in _____.

a) Carbohydrates

b) Proteins

c) Lipids

d) Independent of components

Question 8:

Which of the following statements are false about collagen?

a) Most common fibrous protein

b) Prolysyl and lysyl hydroxylase are involved in post translational modification

c) Phenylalanine is the most abundant AA in its structure

d) Triple helix structure is important for its function

Question 9:

Ketone bodies are not utilized by the liver because it lacks:

a) Thiokinase

b) Thiophorase

c) Thiolase

d) Thiotransferase

Question 10:

Which of the following methods can detect the alterations in DNA sequence?

a) 1 and 3

b) 1, 2, and 3

c) 2, 3, and 4

d) 1, 2, and 4

Question 11:

Which anticoagulant present in the light blue vacutainer is used for plasma glucose estimation?

a) Sodium fluoride

b) Heparin

c) Sodium Citrate

d) Potassium oxalate

Question 12:

Which of the following is not involved in the repair of DNA following double-stranded breaks?

a) ATM

b) BRCA1

c) MSH2

d) Ku70

Question 13:

2-3 BPG binds to?

a) Heme part of myoglobin

b) Globin part of Hemoglobin

c) Globin part of myoglobin

d) Heme part of Hemoglobin

Question 14:

Which of the following is complementary to the active site of the enzyme?

a) Product

b) Substrate

c) Allosteric activator

d) Transition site

Question 15:

Cytochrome P450 monooxygenase utilises which of the following cofactors during the phase 1 reaction of detoxification of xenobiotics?

a) FMN

b) NADPH

c) FAD

d) NAD

Question 16:

Which biochemical cycle/process is used by monoclonal antibodies in hybridoma technology?

a) TCA cycle

b) Glycolytic pathway

c) Pentose phosphate pathway

d) Purine salvage pathway

Question 17:

Glycogen phosphorylase requires which of the following for its activation?

a) Insulin

b) Glucose 6 phosphate

c) Ca2+

d) cAMP independent protein kinase

Answer Key

Question No.	Correct Option
1	d
2	d
3	c
4	a
5	b
6	d

7	b
8	c
9	b
10	a
11	c
12	c
13	b
14	b
15	b
16	d
17	c

Detailed Explanations

Solution to Question 1:

Xanthurenic aciduria is seen in deficiency of vitamin B6 (pyridoxal phosphate), a coenzyme required by kynureninase enzyme to convert 3-OH kynurenine to 3-OH anthranilic acid during tryptophan metabolism.

Other options-

Option A: Riboflavin or Vitamin B2 deficiency manifests as angular stomatitis, magenta tongue, seborrhea, corneal vascularization. It is seen in alcoholism, and individuals with a poor diet and decreased intake of dairy products.

Option B: A patient with thiamine or vitamin B1 deficiency presents either dry Beriberi with predominantly neurological symptoms like neuropathy, muscle weakness, and wasting or wet Beriberi with predominant cardiac symptoms like cardiomegaly, tachycardia, high output congestive cardiac failure.

Option C: Vitamin B12 or cobalamin deficiency presents as anemia, fatigue, and pallor symptoms. Other presenting complaints may include peripheral neuropathy, glossitis, diarrhea, headaches, neuropsychiatric disturbances like dementia, and subacute combined degeneration of the spinal cord (SACD) leading to ataxia. A peripheral smear shows macrocytic anemia and hypersegmented neutrophils.

Solution to Question 2:

The given clinical scenario points towards a diagnosis of diabetic ketoacidosis. Rothera's test as shown in the image is a specific test used for the determination of ketone bodies in the urine.

Type I diabetes mellitus is a low insulin state. There is an increased conversion of fatty acids to acetyl-CoA which in turn leads to increased ketone body production. The presence of

ketosis can be established by the detection of ketone bodies in urine by Rothera's test.

The test involves the saturation of 5 mL of urine with solid ammonium sulfate. This is followed by the addition of a few drops of freshly prepared sodium nitroprusside followed by 2 mL of liquor ammonia along the sides of the test tube. The development of a purple ring indicates the presence of ketone bodies in urine. Strip tests based on the same principle are also available.

Other options-

Option A: Seliwanoff's test will distinguish glucose and fructose. The reagent is resorcinol in HCl. 3 ml of the reagent and 0.5 ml of urine are heated for 30 seconds up to a boiling point. A red color indicates the presence of fructose.

Option B: Benedict's test is a test commonly used in the follow-up of diabetics for the detection of glucose in urine (glucosuria). It uses Benedict's reagent containing sodium carbonate, copper sulfate, and sodium citrate. Reducing sugars (containing a free aldehyde/ keto group) will reduce Benedict's reagent. Examples include glucose, fructose, and lactose. The test is a semi-quantitative test. The blue color indicates no reducing sugar.

Option C: Fouchet's test is used for detecting the presence of bile pigments in urine. Here, urine is heated with barium sulfate, which adsorbs bilirubin. Ferric chloride is added, which oxidizes the adsorbed bilirubin to biliverdin which produces a green color. Principle for all tests to detect bile pigments: Oxidizing agents readily oxidize bilirubin to biliverdin to form a green color.

Solution to Question 3:

5'GAUCGAUAUUGAAUG3' will be the sequence of mRNA produced from this fragment of template DNA.

The non-template strand or coding strand or sense/plus strand in the DNA will have the same sequence of the RNA that is produced, except that the non-template or coding strand has T is in place of U.

As all sequences are written in the 5′ to 3′ direction, unless otherwise specified, an RNA sequence of GAUCGAUAUUGAAUG would correspond to a DNA sequence, on the non-template strand, of CATTCAATATCGATC. Both the non-template strand and RNA produced will be a complementary sequence to the template strand.

Note:

Template strand - known as minus or antisense or non-coding strand.

Coding strand- known as plus or sense or non-template strand.



Solution to Question 4:

The correct order of steps of Karyotyping is as follows:

• Collect cells by peripheral vein using a heparin syringe.

• Culture in phytohaemoagglutinin.

• Incubation at 37 degrees C for 3 days.

• Harvest with colchicine.

• Routine staining with Giemsa

Karyotyping is the process by which chromosomes are isolated, stained, and examined to find chromosomal anomalies. It is used in prenatal diagnosis, investigation of repeat pregnancy loss, investigation of children and adults suspected of having a chromosomal disorder, and subtyping of lymphomas and leukemias for the purpose of exact diagnosis, giving a prognosis, and treatment.

Various techniques are used in karyotyping to study chromosomes. Staining with various compounds produces various results and helps us to understand the chromosomes better.

Common techniques used are:

• Q banding - fluorescence staining with quinacrine mustard

• G banding - Giemsa staining after partial denaturation of chromatin

• R banding - reverse Giemsa staining. In this, the light and dark areas are reversed

• C banding - staining of heterochromatin with silver salts

• T banding - staining the telomeres of chromosomes

The karyotyping of a normal human male is given below.



Solution to Question 5:

Decreased cardiolipin or alteration in its metabolism results in mitochondrial dysfunction seen in Barth syndrome (cardio skeletal myopathy) which is characterized by an enlarged and weakened heart (dilated cardiomyopathy), weakness in muscles used for movement (skeletal myopathy), recurrent infections due to small numbers of white blood cells (neutropenia), and short stature.

Diphosphatidyl glycerol, also called cardiolipin, is the phospholipid found only in the mitochondria. Cardiolipin, which is a phospholipid with antigenic properties, is responsible for the false-positive VDRL test.

It is an important component of the inner mitochondrial membrane, where it is essential for the following:

• Optimal function of numerous mitochondrial enzymes that are involved in mitochondrial energy metabolism.

• Maintenance of the structure of the mitochondrial membrane.

Other options-

Option A: Phosphatidylethanolamine, also called cephalin, is found in the cell membranes of nervous tissue, particularly in the brain.

Option C: Defect in the synthesis of dipalmitoyl lecithin leads to Respiratory distress syndrome (RDS), also known as hyaline membrane disease.

option D: Lysolecithin is obtained by removing one fatty acid from lecithin. Oxidized lysolecithin is associated with atherosclerosis.

Solution to Question 6:

Pompe's disease results from a deficiency in the enzyme lysosomal alpha-glucosidase.

Pompe's disease is a type II glycogen storage disorder that results from a deficiency in the enzyme acid maltase/ lysosomal acid alpha-glucosidase. It is a rapidly progressive disorder that results in hypotonia and hepatosplenomegaly. Myocardial involvement helps to distinguish it from other similar glycogen storage disorders.

Hepatomegaly results from the ballooning of the lysosomes with glycogen. This creates a characteristic lacy cytoplasmic pattern. Cardiac and skeletal muscles will show enlargement with clear spaces in the cells, representing glycogen storage within the lysosomes. This disease classically has a poor prognosis, but enzyme replacement has helped to prolong survival in recent times.

A muscle biopsy with enlarged glycogen-containing vacuoles can be seen in the image below:



Other options:

Option A: Lactic acid dehydrogenase deficiency is not associated with any disorder.

Option B: Tarui's disease is characterized by a deficiency of muscle and erythrocyte phosphofructokinase 1. This results in abnormal glycogen metabolism and hence muscle cannot generate energy after exercise using glycogen reserves which leads to exercise intolerance.

Option C: Muscle glycogen phosphorylase deficiency is seen in McArdle's disease (type V glycogen storage disorder). Patients present with exercise-induced muscle cramping without elevated blood lactate levels which would normally be expected.

Solution to Question 7:

The thermic effect is highest in proteins.

Specific dynamic allowance (previously called the thermic effect of food): This refers to the increased heat production or increased metabolic rate following the intake of food (diet-induced

thermogenesis).

The values of SDA are

• 30% for proteins: This means that out of every 100 grams of proteins consumed, the energy available for doing useful work is 30% less than the calculated value - i.e 30% of the energy content in protein is used in its own metabolism.

• 15% for lipids, and

• 5% for carbohydrates.

Importance: It is a common experience that during hot weather following the consumption of a protein-rich meal, one feels hot and humid for a while. On the other hand, in cold weather, the same would provide a comfortable feeling.

The energy generated by the metabolism of the macronutrients is used for three energy-requiring processes that occur in the body:

• Resting metabolic rate

• Thermic effect of food (formerly termed specific dynamic action)

• Physical activity

Solution to Question 8:

Glycine is the most abundant amino acid(AA) in collagen and not phenylalanine.

Collagen is the most abundant protein in the animal world. It has a triple helical structure, forming a right-handed superhelix. A striking characteristic of collagen is the occurrence of glycine residues at every third position.

Proline and hydroxyproline confer rigidity on the collagen molecule. Hydroxyproline and hydroxylysine are formed by the post-translational hydroxylation of proline and lysine residues catalyzed by the enzyme prolyl and lysyl hydroxylase, whose cofactors are ascorbic acid (vitamin C) and α-ketoglutarate. Collagen is arranged in a quarter-staggered arrangement and further stabilized by covalent cross-links.

Solution to Question 9:

Ketone bodies are not utilized in the liver because it lacks the enzyme thiophorase.

Ketone bodies are synthesized in the liver mitochondria (by ketogenesis) and transported in the blood to the peripheral tissues. Here they are utilized by the following steps:

• β-hydroxybutyrate (the most predominant ketone body) is oxidized to acetoacetate by β-hydroxybutyrate dehydrogenase producing NADH.

• Acetoacetate is then activated to acetoacetyl CoA by transferring a CoA molecule from succinyl CoA by succinyl CoA-acetoacetyl CoA transferase (thiophorase).

• Acetoacetyl CoA is cleaved into two acetyl CoA molecules by thiolase and oxidized into CO_2 by the TCA cycle.

Thiokinase or Acyl CoA synthetase catalyzes the only energy-requiring step in fatty acid oxidation. In the presence of ATP and coenzyme A, thiokinase catalyzes the conversion of a fatty acid to acyl-CoA. Acyl-CoA is the active form of fatty acid. This step uses one high-energy phosphate and forms AMP and PPi.

When higher than normal ketone bodies are present in the blood or urine, it is called ketonemia or ketonuria, respectively. The overall condition is called ketosis. The metabolism of free fatty acids to produce ketone bodies is exaggerated in pathologic states such as starvation and diabetes mellitus, thus causing ketosis. The continued buffering and excretion of ketone bodies deplete the alkali reserve, causing ketoacidosis.

Solution to Question 10:

The alterations in DNA sequence can be detected by restriction fragment length analysis and pyrosequencing.

Restriction fragment length polymorphism (RFLP) is also called cleaved amplified polymorphic sequence (CAPS) assay. RFLP is a technique in which organisms may be differentiated by variations in their DNA sequence by restriction endonucleases. This technique involves cutting a particular region of DNA with known variability, with restriction enzymes, then separating the DNA fragments by agarose gel electrophoresis and determining the number of fragments and relative sizes.

Pyrosequencing is a synthesis type of approach of DNA sequencing. This technique employs four enzymatic reactions in a single tube to monitor DNA synthesis. Nucleotides are incorporated into the reaction and pyrophosphate (PPi) is released. PPi triggers a series of reactions resulting in the production of light, which is proportional to the amount of DNA and number of incorporated nucleotides. The generated light is detected and recorded by a detector system in the form of a peak signal, which reflects the activity of all four enzymes in the reaction.

The practical applications of recombinant DNA technology include:

• Gene mapping using techniques such as fluorescence in situ hybridization (FISH).

• Producing proteins such as:

• Interferons, tPA (tissue plasminogen activator)

• Insulin, growth hormone

• Hepatitis B vaccine

• Molecular analysis of disease, to detect gene variations and mutations.

• Restriction Fragment Length polymorphism (RFLP) and single nucleotide polymorphism (SNP)

• Microsatellite DNA polymorphisms

• Gene therapy and bone marrow transplantation

Flow cytometry is a laser-based method that analyzes the cells and cell components based on the presence of certain cell surface molecules which are named as per cluster of differentiation (CD)

nomenclature. It is very effective for the study of the immune system and its response to infectious diseases and cancer. It cannot detect the alterations in DNA sequence.

Fluorescent in situ hybridization (FISH) is a technique where cells are stained with fluorescent stains. It uses DNA probes that recognize sequences specific to particular chromosomal regions. It cannot detect the alterations in DNA sequence. FISH can be performed on prenatal samples, peripheral blood cells, touch preparations from cancer biopsies, and even fixed archival tissue sections. Uses include :

• Numerical abnormalities of chromosomes e.g., aneuploidy.

• Subtle microdeletions and rearrangements.

• Complex translocations that are not demonstrable by routine karyotyping.

• Gene amplification of HER2 in breast cancer or NMYC amplification in neuroblastomas.

Solution to Question 11:

Light blue vacutainers contain the anticoagulant sodium citrate.

Citrated plasma generated from whole blood is commonly utilized in laboratory testing for hemostasis. The recommended procedure involves the collection of specimens in tubes that are filled with a solution containing 3.2% sodium citrate (109 mM). The appropriate ratio for this collection is 9 parts blood to 1 part anticoagulant. The primary objective of citrate is to effectively eliminate calcium ions, which play a crucial role in the process of blood coagulation.

The following image shows color coding of vacutainers with the anticoagulant used and their practical uses:

Vacutainer	Additive	Uses
Lightgrey	Sodiumfluorideand potassium oxalate	Blood glucose estimation
Light green	Lithium Heparin	Plasma testing
Light blue	SodiumCitrate	Coagulation tests (APTT and prothrombin time)



Solution to Question 12:

MSH2 is involved in mismatch repair and is not involved in the repair of double-stranded DNA breaks as the repair involves the correction of a single mismatch base pair e.g. C to A, instead of T to A.

DNA mismatch repair proteins function as spell checkers during DNA replication by correcting the incorrect pairing. In HNPCC, the proofreading function of the mismatch repair systems (majorly MLH1 and MSH2) is lost, leading to the accumulation of errors throughout the genome. The genomes of these patients show microsatellite instability, which is characterized by changes in the length of short repeats.

Double-strand breaks (DSBs) in DNA pose a significant concern. Genetic recombination events can give rise to chromosomal translocation, chromosomal breakage, and ultimately cellular demise. Double-strand breaks (DSBs) have the potential to undergo repair through two distinct mechanisms: homologous recombination and non-homologous end joining. Homologous recombination is observed in yeasts, whereas in mammals, non-homologous end joining predominates.



Non-homologous end-joining (NHEJ) is a mechanism by which a group of proteins mediates the recognition, processing, and direct ligation of the ends of two DNA fragments where a double-strand break in DNA has occurred. This is in contrast to homologous repair (HR), another mechanism, where an undamaged homologous DNA template is required to replace an adjacent damaged one.

The type of repair depends on the phase of cell cycles:

Other options:

Option A: The protein known as ataxia telangiectasia mutated (ATM) serves as a crucial kinase that triggers a wide range of cellular stress responses through its signalling activity. The activation of ATM is induced by DNA double-strand breaks (DSBs) and oxidative stress, resulting in the phosphorylation of a wide range of target proteins. Recent advancements in molecular biology have brought to light the participation of the ataxia telangiectasia mutated (ATM) protein in a range of cellular mechanisms triggered by DNA damage. These mechanisms encompass the repair of double-strand breaks (DSBs), the halting of cell cycle progression through checkpoint activation, the initiation of programmed cell death (apoptosis), and the cessation of transcriptional activity. The compromised precision of repair resulting from the combinational failure of stress responses leads to heightened sensitivity to ionizing radiation (IR) in ataxia-telangiectasia (A-T) cells.

Option B: CtIP which is a DNA repair protein involved in Homologous Repair whose function is dependent on phosphorylation at serine 327 as well as the recruitment of BRCA 1.

Option D: The Ku heterodimer, consisting of Ku70 and Ku80 subunits, plays a pivotal role in the nonhomologous end-joining (NHEJ) pathway, which is responsible for the repair of DNA double-strand breaks (DSBs). Ku protein mediates the binding of the damaged DNA fragment and orchestrates the recruitment of additional proteins to enable the subsequent processing and ligation of the broken DNA end.

Phaseofthecellcycle	Repairmechanism
Go /G1 Phase	Nonhomologous End Joining Repair
S, G2, M Phase	Homologous Repair

Solution to Question 13:

2,3 bisphosphoglycerate binds to the globin chains of the hemoglobin while oxygen binds to the heme part of the hemoglobin.

A single unit of 2,3-BPG forms a complex with one tetrameric unit of deoxyhemoglobin within the central cavity of its four subunits. The centre pocket of this structure contains two -globin chains that are positively charged, such as histidine and lysine. Ionic bonds, also known as salt bridges, are established through the interaction between amino acids that carry a positive charge, specifically those found in globins, and phosphate groups that possess a negative charge, specifically those present in 2,3-BPG. The stabilization of deoxygenated haemoglobin (in its T-form) is achieved through the cross-linking of the α-chains by the binding of 2,3-BPG. Upon oxygenation of haemoglobin, the molecule 2,3-BPG is released from its binding site, resulting in the transformation of oxyhemoglobin into the R-form conformation.

A relaxed (R) configuration, exposes more O2 binding sites. The quaternary structure of haemoglobin determines its affinity for O2. In deoxyhemoglobin, the globin units are tightly bound in a tense (T) configuration, which reduces the affinity of the molecule for O2. When O2 is first bound, the bonds holding the globin units are released, producing a relaxed (R) configuration, which exposes more O2 binding sites. The net result is a 500-fold increase in O2. In tissues, these reactions are reversed, resulting in O2 dissociation.



Solution to Question 14:

The substrate is complementary to the active site of the enzyme.

The Lock and Key concept was put forth by Emil Fischer. This model is the initial proposition aimed at elucidating the mechanism of an enzymatic catalytic process. Based on this concept, it can be inferred that the enzyme's structure or conformation exhibits a state of rigidity. The substrate effectively conforms to the binding site, which is now referred to as the active site, in a manner analogous to a key fitting precisely into its corresponding lock or a hand fitting snugly into a well-fitting glove. The active site of an enzyme can be described as a stiff and pre-configured template that selectively accommodates a certain substrate for binding.

The induced fit or Koshland theory states that the active site does not possess a firm and pre-determined conformation. The emerging active site exhibits the fundamental characteristics of the substrate binding site. The substrate's contact with the enzyme elicits a conformational change in the enzyme, leading to the establishment of a robust binding site for the substrate. Moreover, as a result of induced fit, the enzyme undergoes conformational changes that lead to the repositioning of the specific amino acids involved in the formation of the active site, hence facilitating the catalytic process. The induced fit model has garnered substantial empirical support through X-ray diffraction studies. Koshland's model additionally elucidates the mechanisms by which allosteric modulators and competitive inhibition exert their effects on enzymes.

The substrate strain theory states that the substrate experiences strain as a result of the conformational alteration generated by the enzyme. It is plausible that upon the binding of a substrate to the preexisting active site, the enzyme elicits a strain on the substrate. The presence of a stressed or strained substrate facilitates the generation of the desired product. The enzymatic action is believed to involve a combination of the induced fit model and substrate strain.



Other options:

Option A: The complementary active site of the enzyme binds to the substrate to form the product.

Option C: The active site of the enzyme is not complementary to the allosteric activator. Allosteric sites are separate and specific sites on the enzyme that are modulated with allosteric effect

modifiers and hence are complementary to them.

Option D: The active site of the enzyme binds to the substrate which form a transitional state which is unstable and later dissociates to form a product.

Solution to Question 15:

The microsomal cytochrome P450 system, located in the liver, facilitates the process of drug and foreign substance detoxification through hydroxylation processes that rely on the availability of NADPH.

Phase 1 reactions include oxidation, reduction, or hydrolytic reaction. CYPs- Cytochrome P family of enzymes play a major role in phase 1 reactions. Reactions carried out by phase 1 enzymes usually lead to the inactivation of a drug. The addition of functional groups does little to increase the water solubility of the drug but alters its biological properties.

The importance of NADPH is enumerated as follows:

• NADPH is needed for the reductive biosynthesis of fatty acids and steroids, hence HMP shunt is more active in lipogenesis tissues like adipose tissue and the liver.

• Along with Glutamate dehydrogenase helps in the synthesis of various amino acids.

• Maintaining reduced glutathione is critical for RBC membrane integrity. NADPH keeps haemoglobin's ferrous iron (Fe^{2+}) lowered to prevent methemoglobin (Fe^{3+}) buildup.

• High concentrations of NADPH are also required to maintain the transparency of the lens.

The cytochromes are a class of proteins that possess a conjugated heme group. The latter is comprised of a porphyrin ring containing an iron atom. The heme group present in cytochromes exhibits distinct characteristics when compared to the heme group observed in the structures of haemoglobin and myoglobin. The heme iron within cytochromes undergoes a cyclic process of oxidation (Fe^{3+}) and reduction (Fe^{2+}), playing a crucial role in facilitating electron movement throughout the electron transport chain (ETC). This stands in opposition to the heme iron found in haemoglobin and myoglobin, which remains in the ferrous (Fe^{2+}) form. The cytochromes were initially assigned the designations of a, b, and c based on the specific heme type they contained and their corresponding absorption spectra. Subsequent discoveries led to the identification of additional cytochromes, including c1, b1, b2, a3, among others.

Other options

Option A: Flavin adenine mononucleotide (FMN) is a phosphorylated form of riboflavin which functions as a cofactor for various enzymes. Some of these include as a cofactor for blue-light photoreceptors and oxidoreductases.

Option C: Flavin adenine dinucleotide (FAD) is nothing but phosphorylated riboflavin and is used in the electron transport chain and also is a cofactor for various enzymes. Some of these include oxidative phosphorylation, DNA repair, beta-oxidation of fatty acids as well as amino acid catabolism.

Option D: Nicotinamide adenine dinucleotide (NAD) consists of the vitamin B3 (niacin) and also serves as a coenzyme for various enzymatic reactions in the body. Some of these include oxidative phosphorylation (ATP production), DNA repair and intracellular calcium signalling.

Solution to Question 16:

It is the purine salvage pathway that is utilized by monoclonal antibodies in hybridoma technology.

Monoclonal antibodies refer to the antibodies produced by a group of lymphocytes that produce antibodies against a single antigen. These antibodies are produced by hybridoma technology.

In this technology, the B-lymphocytes of mice spleen are immunized with an antigen to produce a specific antibody. These specific antibody-producing B lymphocytes are then fused with myeloma cells that can multiply indefinitely.

The myeloma cells are deficient in the hypoxanthine-guanine-phosphoribosyltransferase (HGPRT) gene and do not produce immunoglobulins. Therefore, the resulting hybrid cell is producing antibodies against a single antigen indefinitely. These hybrid cells are cultured in a HAT medium (containing hypoxanthine, aminopterin, and thymidine) and the monoclonal antibodies are obtained. It is well-established that nucleotides present in DNA and RNA consist of purine and pyrimidine bases. The synthesis of these nucleotides can occur through either the De novo pathway or the salvage pathway. Consequently, the inclusion of aminopterin in the HAT medium effectively inhibits the De novo pathway, thereby rendering the synthesis of purine dependent on the salvage pathway. However, it should be noted that the salvage pathway exclusively takes place in the presence of the enzymes HGPRT and thymidine kinase. The hybridoma cell is uniquely capable of proliferating on the HAT medium due to its distinct composition. Specifically, the myeloma component of the hybridoma possesses the capacity for repetitive cellular division, while the plasma cell component contains the enzyme HGPRT. Consequently, the hybridoma cells can sustain growth and viability over an extended period when cultivated on the HAT medium.

Hybridoma technology is a method for producing large numbers of identical antibodies (monoclonal antibodies).

Hybridomas are somatic cell hybrids produced by fusing the antibody-forming spleen cells with myeloma cells. The resultant hybrid thus has both the antibody-producing ability of the B-cell and the ability to multiply indefinitely of the myeloma.

The following are the steps involved in hybridoma technology:

• Mice are immunized with an antigen.

• After several weeks of immunization, blood samples are obtained from mice for measurement of serum antibodies.

• When the antibody titer is high enough, mice are commonly boosted by injecting antigen without adjuvant, intraperitoneally or intravenously.

• Then the mice are euthanized and their spleens removed for in vitro hybridoma cell production.

• Fusing antibody-producing spleen cells, which have a limited life span, with myeloma cells (derived from an immortal tumour of lymphocytes) results in a hybridoma that is capable of unlimited growth.

• The cells are cultured in a medium containing hypoxanthine, aminopterin, and thymidine (HAT medium) which allows only the fused cells to survive in culture.

• Monoclonal antibodies are then harvested.

Applications of hybridoma technology

• Listing lymphocyte subpopulations

• Specific cell quantitative preparation

• As an immunosuppressant in therapeutic settings

• Blood component nephelometric assays

• ELISA testing

• Quantitative purification of antigens.

• Common therapeutic applications for monoclonal antibodies



Solution to Question 17:

Glycogen phosphorylase requires the presence of calcium to attain its fully activated state.

Glycogen phosphorylase is an enzyme required in glycogenolysis.

In the muscle: During muscle contraction → Ca++ is released → Ca++ binds to calmodulin forming Ca++-Calmodulin complex → activates phosphorylase kinase from b form (inactive form) without phosphorylation → significant Ca++ is required to activate glycogen phosphorylase.

Hormonal regulation of glycogenolysis:

Glucagon (in liver) and epinephrine (in liver and muscle) acts via GPCR to activate adenylyl cyclase to catalyse synthesis of cAMP. cAMP dependent protein kinase activates phosphorylase kinase from inactive(b) to active(a) form, which in turn activates glycogen phosphorylase to active(a) form, hence leading to increased glycogenolysis. (Option D)

Option A: Insulin is a known inhibitor of glycogen phosphorylase which dephosphorylates the glycogen phosphorylase enzyme thus rendering it inactive.

Option B: Glucose 6 phosphate is also an inhibitor of glycogen phosphorylase which dephosphorylates it thus rendering it inactive.

The regulation of glycogen phosphorylase is shown in the image below:



Biochemistry INI-CET 2024

Question 1:

The following Lineweaver-Burk plots depict enzymatic action in the presence of enzyme inhibitors. The enzyme action without inhibitor is represented by a blue line. What type of inhibitors are depicted in images 1 and 2?



a) 1- Uncompetitive inhibition 2- Competitive inhibition

b) 1- Noncompetitive inhibition 2- Competitive inhibition

c) 1- Competitive inhibition 2- Uncompetitive inhibition

d) 1- Competitive inhibition 2- Noncompetitive inhibition

Question 2:

FISH technique helps to identify all except?

a) Gene amplification

b) Deletion

c) Point mutation

d) Translocation

Question 3:

Which of the following micronutrients are associated with decreased sexual maturation and hypogonadism?

a) Zinc

b) Copper

c) Selenium

d) Iron

Question 4:

Which of the following is not a heme breakdown product?

a) Carbon Monoxide

b) Ferric / Fe 3+

c) ALA

d) Biliverdin

Question 5:

Which type of mutation results in the reversal to wild type of phenotype when the mutant gene is suppressed?

a) Deletion of mutated gene

b) Frameshift of mutated gene

c) Addition of another normal gene

d) Mutations in tRNA

Question 6:

Which of the following is not a post-transcriptional modification?

a) 5 prime capping

b) Methylation

c) Glycosylation

d) Endonuclease cleavage

Question 7:

Match the following conditions :

a) 1 - b, 2 -a, 3 -d, 4 -c

b) 1 -c, 2 -d, 3 -b, 4 -a

c) 1 -b, 2 -d, 3 -a, 4 -c

d) 1 -c, 2 - b, 3 -a, 4 -d

Question 8:

All are true statements regarding folate deficiency except

a) Increased levels of homocysteine and normal methylmalonic acid

b) Folate supplementation on diagnosis of pregnancy prevents neural tube defect

c) It is important to treat co-existing vitamin B12 deficiency before initiating folate therapy

d) Folate deficiency is commonly seen in children with hemolytic anemia

Question 9:

Congenital lactic acidosis is seen due to a defect in which of the following enzymes?

a) Pyruvate dehydrogenase

b) Branched chain ketoacid dehydrogenase

c) Transketolase

d) Isocitrate dehydrogenase

Question 10:

Which of the following is not associated with G6PD deficiency?

a) Equal incidence in males and females

b) Defective NADPH synthesis

c) Predisposes the patient to oxidative stress

d) Patient presents with hemolysis

Question 11:

Which of the following is not a neurotransmitter derived from aminoacids ?

a) Gamma amino butyric acid

b) Dopamine

c) 5 hydroxytryptamine

d) Creatine

Question 12:

Which of the following methods is used to test the blood taken from a neonate for metabolic disease screening?

a) Complete blood count

b) Tandem Mass Spectrometry

c) Next Generation Sequencing

d) ELISA

Question 13:

Familial hypercholesterolemia is a result of a defect in which of the following?

a) LDL receptor

b) Apo E

c) Apo CII

d) Apo B48

Question 14:

In a patient with Diabetes Mellitus, the liver produces more TG and VLDL. What is the cause of the same?

a) Decreased action of LPL with increased action of HSL

b) Decreased action of LPL with Decreased action of HSL

c) Increased action of LPL with increased action of HSL

d) Increased action of LPL with decreased action of HSL

Question 15:

Nutritional deficiency of Copper, required by lysyl oxidase which is an importent key step for formation for covalent cross linking in collagen synthesis, gives rise to which of the following condition?

a) Menke's disease

b) Scurvy

c) Ehlers Danlos syndrome

d) Scoliosis

Question 16:

Match the following with appropriate precursors

a) i-b, ii-d, iii-a, iv-c

b) i- c, ii-d, iii-a, iv-b

c) i- a, ii-d, iii-b, iv-c

d) i- b, ii-a, iii-d, iv-c

Question 17:

In the current scenario treatment for urea cycle disorder is available. Administration of amino acid phenylbutyrate can decrease ammonia formation. Which amino acid provides phenylbutyrate?

a) Arginine

b) Glutamate

c) Aspartate

d) Glycine

Question 18:

Which among the following are appropriate techniques for chromosomal aneuploidy?

a) i,ii,iii,iv

b) i,ii

c) ii,iii

d) iii,iv

Question 19:

Enzyme used in recombinant DNA technology are?

a) 1,2,3,4

b) 1,2,3

c) 2,3,4

d) 1,3,4

Answer Key

Question No.	Correct Option
1	b
2	c
3	a
4	c
5	d
6	c
7	b
8	d
9	a
10	a
11	d
12	b
13	a
14	a
15	a
16	a
17	b
18	a
19	c

Detailed Explanations

Solution to Question 1:

The Lineweaver-Burk plot in image 1 depicts non-competitive inhibition, and that in image 2 depicts competitive inhibition.

Important parameters in a reaction include:

• The affinity of the enzyme for its substrate, given by the Michaelis constant or Km. A low Km reflects a high affinity of the enzyme for substrate and vice versa. The value of Km is given by the substrate concentration at which the reaction velocity is equal to one-half Vmax. It does not change with changes in enzyme concentration.

• Vmax is the maximum velocity of a reaction.

Competitive inhibition is when the inhibitor binds reversibly to the substrate-binding site and competes with the substrate for the active enzyme-binding site. This effect can be overcome by increasing the substrate concentration by which maximum velocity can still be achieved. Therefore, Vmax is unaltered in this type of inhibition. However, the concentration of the substrate required to achieve half of the maximum velocity is increased. Therefore, Km is increased in this condition.

In a Lineweaver-Burk plot, the addition of a competitive inhibitor causes plots with the same y-intercept (Vmax is unaffected) but different x-intercepts (due to increased Km).



Noncompetitive inhibition is by binding of inhibitors to enzymes in locations remote from the active site. Increasing the concentration of the substrate does not overcome non-competitive inhibition. Thus, noncompetitive inhibitors decrease the Vmax of the reaction. They do not interfere with the binding of the substrate to the enzyme. Thus, the enzyme shows the same Km in the presence or absence of noncompetitive inhibitors.

In a Lineweaver-Burk plot, the addition of a non-competitive inhibitor causes plots with the same x-intercept (Km is unaffected) but different slopes and y-intercepts (due to decreased Vmax).



In uncompetitive inhibition, inhibitors bind with the enzyme-substrate intermediate complex and inhibit it. This leads to a decrease in both Vmax and Km.



Solution to Question 2:

Fluorescent in situ hybridization (FISH) is not useful in identifying point mutation. Point mutations are identified by other means like sequencing studies and restriction fragment length polymorphism.

FISH is a method of molecular analysis of genetic lesions involving large deletions, duplications, or more complex rearrangements that are not readily amenable to detection using polymerase chain reaction or DNA sequencing approaches.

FISH uses DNA probes that recognize sequences specific to particular chromosomal regions. To perform FISH, large fragments of cloned genomic DNA spanning up to 200 kb are labeled with fluorescent dyes and applied to metaphase chromosome preparations or interphase nuclei. The probe hybridizes to its homologous genomic sequence and thus labels a specific chromosomal region that can be visualized under a fluorescent microscope.

FISH can be performed on prenatal samples, peripheral blood cells, touch preparations from cancer biopsies, and even fixed archival tissue sections. It is used to detect numeric abnormalities of chromosomes (aneuploidy), subtle microdeletions and complex translocations that are not demonstrable by routine karyotyping, and gene amplification (e.g., HER2 in breast cancer or NMYC amplification in neuroblastomas).



Solution to Question 3:

Zinc deficiency is associated with decreased sexual maturation and hypogonadism.

Zinc is the second most common trace element in the body after iron. As a component of zinc finger proteins, zinc regulates gene transcription and participates in nucleic acid metabolism, protein synthesis and thereby cell growth. Zinc is a major antioxidant being part of the enzyme superoxide dismutase.

Zinc is derived mainly from animal protein. Diets based on cereals, plants and legumes are associated with zinc deficiency due to dietary phytic acid decreasing the bioavailability of zinc.

Zinc is absorbed throughout the small intestine by a process of facilitated diffusion. Almost 90% of the total body zinc is localized in bone and skeletal muscle.

Deficiency -

• Poor physical growth is an important feature of zinc depletion in preschool and school-age children. Delayed sexual maturation and hypogonadism are prominent features of zinc deficiency in adolescents. Other features include anemia, anorexia, diarrhea, alopecia, impaired immune function and poor wound healing.

• Acrodermatitis enteropathica is an autosomal recessive disorder of severe zinc deficiency, caused by impaired intestinal absorption due to a defect in intestinal zinc transporter protein. It presents in early infancy with vesiculobullous, dry, scaly, or eczematous skin lesions chiefly in the periorificial and acral areas.

Normal requirements for children range between 3.5 and 5.0 mg/day. Acquired zinc deficiency stats can be treated with 0.5 to 1.0 mg/kg/day for several weeks. Malnourished children have much higher requirements of 2-4mg/kg/day.

Other options :

Option B - Copper deficiency results in microcytic, hypochromic anemia unresponsive to iron therapy, depigmentation of hair, neutropenia, neurological problems and osteoporosis. Disruption in copper transport is linked to Wilson's disease and Menke's disease.

Option C - Selenium deficiency is a cause of Keshan disease, a form of cardiomyopathy in young children seen in some regions of China. In combination with iodine deficiency, lack of selenium can result in myxedematous endemic cretinism.

Option D - Iron deficiency causes anemia, irritability, anorexia, pica, angular stomatitis, leg cramps and congestive heart failure.

Solution to Question 4:

Aminolevulinic acid (ALA) is not a product of heme breakdown; rather it is an intermediate in the synthesis of heme molecules.

The process of heme breakdown occurs over a series of reactions. In the reticuloendothelial system of the liver, spleen and bone marrow, heme is degraded by microsomal heme oxygenase. It requires NADPH and cytochrome c. In this process, the alpha methenyl bridge of the heme molecule is liberated as carbon monoxide. The Fe+ liberated is oxidized to Fe3+++ and taken up by transferrin.

The linear tetrapyrrole formed is biliverdin (green pigment), which is further reduced to bilirubin (red-yellow pigment) by an NADPH-dependent biliverdin reductase. The bilirubin is transported in the plasma bound to albumin and is taken up by the liver, where it is conjugated with glucuronic acid to make it water-soluble, which is finally excreted in the bile.

The intestinal bacteria deconjugate the bilirubin, which is reduced to urobilinogen. It is further reduced to stercobilinogen, which is excreted through feces.



Solution to Question 5:

Mutations in tRNA are known to suppress the mutant gene and cause reversal to wild phenotype. Such mutations are called suppressor mutations.

Suppressor mutations are secondary mutations that alter the phenotypes caused by an existing mutation. This is not to be confused with reversion mutations in which the original mutation has been lost. Instead, suppressor mutations return the phenotype caused by the original mutation to a more wild-type phenotype despite the presence of the original mutation. The suppressor mutation's effect on the phenotype can vary from partial restoration to full restoration of the wild-type phenotype.

In prokaryotic and lower eukaryotic organisms, abnormally functioning tRNA molecules have been discovered that are themselves the results of mutations. Some of these abnormal tRNA molecules are capable of binding to and decoding altered codons, thereby suppressing the effects of mutations in the structural genes.

These suppressor tRNA molecules, usually formed as a result of alterations in their anticodon regions, are capable of suppressing certain missense mutations, nonsense mutations, and frameshift mutations. An example of suppressor mutation is illustrated below.

A primary mutation occurs changing the original codon sequence from UGG (Trp) to UAG (Stop). A secondary mutation occurs in the gene for a glutamine tRNA leading to the anticodon switch from a GUC to AUC. This mutant tRNA can recognize the UAG stop codon and insert glutamine in place of the stop, allowing translation to continue past the stop and suppressing the original mutation.



In the laboratory context, such suppressor tRNAs, coupled with mutated variants of aminoacyl-tRNA synthetases, can be utilized to incorporate unnatural amino acids into defined locations within altered genes that carry engineered nonsense mutations. The resulting labeled proteins can be used for in vivo and in vitro cross-linking and biophysical studies.

Solution to Question 6:

Glycosylation is not a post-transcriptional modification; it is a post-translational modification.

A primary transcript is the initial, linear, RNA copy of a transcription unit. The collection of all the primary transcripts synthesized in the nucleus by RNA polymerase II is known as heterogeneous nuclear RNA (hnRNA). The pre-mRNA components of hnRNA undergo extensive co- and posttranscriptional modification in the nucleus and become mature mRNA. These modifications usually include the following :

• Addition of a 5′ cap - This is the first of the processing reactions for pre-mRNA. The cap is a 7-methylguanosine attached to the 5′ terminal end of the mRNA and is resistant to most nucleases. Creation of the cap requires the removal of the γ-phosphoryl group from the 5′ triphosphate of the pre-mRNA, followed by the addition of guanosine monophosphate. Methylation of this terminal G occurs in the cytosol and is catalyzed by guanine-7-methyltransferase. Additional methylation steps may occur (options A and B).

• Addition of a 3′-poly-A tail - Most eukaryotic mRNA have a chain of 40 to 250 adenylates (adenosine monophosphates) attached to the 3′ end. This poly-A tail is added by the nuclear enzyme, polyadenylate polymerase. The poly-A tail terminates eukaryotic transcription. In addition, it helps stabilize the mRNA, facilitates its exit from the nucleus, and aids in translation.

• Splicing - Maturation of eukaryotic mRNA usually involves removal from the primary transcript of RNA sequences (introns) that do not code for protein (endonuclease cleavage). The remaining coding (expressed) sequences, the exons, are joined together to form the mature mRNA. The process of removing introns and joining exons is called splicing. The molecular complex that accomplishes these tasks is known as the spliceosome (option D).

Solution to Question 7:

The correct answer is option B: 1 -c, 2 -d, 3 -b, 4 -a

Von Gierke disease (type I glycogen storage disorder) occurs due to the deficiency of glucose-6-phosphatase enzyme in the liver, kidney, and intestinal mucosa. Patients present commonly by 3-4 months of age with hypoglycemia (leading to seizures), lactic acidosis, and hepatomegaly. Affected children often have a doll-like face with fat cheeks, relatively thin extremities, short stature, and a protuberant abdomen due to massive hepatomegaly. The kidneys are also enlarged, whereas spleen and heart are not involved. Laboratory findings include hypoglycemia, lactic acidosis, hyperuricemia, and hyperlipidemia.

Pompe's disease (type II glycogen storage disorder) occurs due to a mutation in alpha-1,4-glucosidase (also known as acid maltase), a lysosomal enzyme. A defect in this enzyme leads to the accumulation of glycogen in the lysosomes of peripheral, voluntary muscles and progresses to involve the cardiac muscle and diaphragm. Clinical features include hypotonia (floppy baby syndrome), cardiomyopathy, hepatomegaly, and failure to thrive. Myozyme can prevent worsening or can reverse cardiac and skeletal abnormalities.

Cori disease (type III glycogen storage disorder) occurs due to a deficiency of the glycogen debranching enzyme. The features of Cori disease include hypoglycemia, hepatomegaly, and hyperlipidemia but with normal blood lactate and uric acid levels. Muscle weakness and atrophy can also occur in these patients. Liver biopsy shows the accumulation of glycogen in the form of limit-dextrins (branched structures).

McArdle disease (type V glycogen storage disorder) typically presents during late childhood or in the second decade of life. The muscle glycogen phosphorylase enzyme deficiency interferes with ATP generation by glycogenolysis in the muscles. This causes muscle glycogen accumulation. Symptoms are exercise intolerance associated with muscle cramps. This leads to severe muscle damage resulting in a massive release of myoglobin in the blood.

1) Glycogen accumulation in liver and renal cells and lactic acidemia, associated with hypoglycemia	c) Von Gierk disease
2) Glycogen accumulation in lysosomes	d)Pompe disease
3) Exercise intolerance	b)McArdle disease
4) Accumulation of branched structures	a)Cori disease

Solution to Question 8:

Folate deficiency is seen in children with macrocytic anemia and not with hemolytic anemia.

Folic acid (vitamin B9) is the synthetic form of folate used in supplements and fortification of foods. Folic acid plays a key role in one-carbon metabolism, essential for the biosynthesis of several compounds. Tetrahydrofolate (THF), the reduced, coenzyme form of folate, receives one-carbon fragments from donors such as serine, glycine, and histidine and transfers them to intermediates in the synthesis of amino acids, purine nucleotides, and thymidine monophosphate (TMP), a pyrimidine nucleotide incorporated into DNA.

Leafy vegetables such as spinach, turnip, lettuce and fortified cereal products are rich sources.

Folate deficiency is caused by increased demand (e.g., pregnancy and lactation), poor absorption caused by pathology of the small intestine, alcoholism, or treatment with drugs (e.g., methotrexate) that are dihydrofolate reductase inhibitors. Children with hemolytic anemia have high folate requirements due to increased erythropoiesis predisposing them to deficiency (option B). Its manifestations are :

• Anemia - A primary result of folic acid deficiency is megaloblastic anemia, caused by diminished synthesis of purine nucleotides and TMP, which leads to an inability of cells to make DNA and, therefore, an inability to divide.

• Neural tube defects - Maternal deficiency of folate, during pregnancy, is implicated in neural tube defects. Periconceptional folate supplements, begun 1 month before conception and continued for 3 months after, are recommended to reduce the risk.

• Hyperhomocysteinemia - Folate, along with vitamin B12, is involved in converting homocysteine to methionine, as depicted below. Hence, folate deficiency results in increased levels of homocysteine.



However, levels of methylmalonic acid are normal. Methylmalonic acid levels are elevated in deficiency of vitamin B12, which is required as a coenzyme in the conversion of methylmalonyl CoA to succinyl CoA (option A).



Treatment: In patients with megaloblastic anemia, it is imperative to exclude and treat vitamin B12 deficiency before treating with folate. Otherwise the neurological signs of vitamin B12 deficiency may develop and progress irreversibly (option C). Deficiency is corrected using folic acid at a dose of 0.5-1mg/day orally for 3-4 weeks.

Solution to Question 9:

Congenital lactic acidosis is seen due to a defect in pyruvate dehydrogenase.

The major source of acetyl CoA for the tricarboxylic acid (TCA)cycle is the oxidative decarboxylation of pyruvate by the multienzyme pyruvate dehydrogenase complex (PDHC). The PDHC is a protein aggregate of multiple copies of three enzymes, pyruvate decarboxylase (E1), dihydrolipoyl transacetylase (E2), and dihydrolipoyl dehydrogenase (E3). Each catalyzes a part of the overall reaction.

A deficiency of the E1 component of the PDHC is the most common biochemical cause of congenital lactic acidosis. The deficiency results in a decreased ability to convert pyruvate to acetyl CoA, causing pyruvate to be shunted to lactate via lactate dehydrogenase. This creates particular problems for the brain, which relies on the TCA cycle for most of its energy and is particularly sensitive to acidosis. Symptoms include neurodegeneration, muscle spasticity, and, in the neonatal-onset form, early death. The inheritance pattern is X- linked dominant, with both males and females affected. Although there is no proven treatment for PDHC deficiency, dietary restriction of carbohydrates and supplementation with thiamine may reduce symptoms in select patients.

Other options :

Option B - Branched-chain ketoacid dehydrogenase deficiency results in maple syrup urine disease. There is defective oxidative decarboxylation of leucine, isoleucine and valine. The disease is characterized by feeding problems, ketoacidosis, changes in muscle tone, and coma.

Option C - Defective transketolase is a rare disorder of pentose phosphate metabolism with symptoms apparent in infancy including developmental delay and intellectual disability, delayed or absent speech, short stature, and congenital heart defects.

Option D - Isocitrate dehydrogenase is an enzyme of the TCA cycle which converts isocitrate to alpha-ketoglutarate. Defects in this enzyme are linked to carcinogenesis.

Solution to Question 10:

G6PD Deficiency does not have an equal incidence in males and females. It is an X-linked recessive (XLR) disorder, hence males are more affected than females.

Glucose-6-phosphate dehydrogenase (G6PD) deficiency causes congenital hemolytic anemia due to enzyme deficiency. G6PD is required to generate NADPH while oxidizing glucose-6-phosphate in the hexose monophosphate pathway (HMP). G6PD deficiency leads to reduced NADPH levels (option B) which then reduces the level of reduced glutathione. Reduced glutathione is required to protect the cells against oxidative stress to RBCs and hemolysis.

Clinically, most patients are asymptomatic. When exposed to triggers like fava beans, infections, drugs such as primaquine, chloroquine, sulfonamides/sulfones, dapsone, and nitrofurantoin patient develops:

• Episodic hemolysis, both intravascular and extravascular

• Hemolytic anaemia

• Hemoglobinemia

• Hemoglobinuria

Increased oxidative damage to RBCs leads to the formation of denatured precipitates known as Heinz bodies. These bodies can damage the RBC membrane leading to intravascular hemolysis. Additionally, when such RBCs pass through the splenic cords, macrophages pluck out these precipitates, giving them the characteristic appearance of bite cells.

The images below show the presence of Heinz bodies and bite cells.



©MARROW



Solution to Question 11:

Creatine is not a neurotransmitter derived from amino acids.

Creatine is an intermediate product in the process of formation of creatinine. The steps involved in the formation are as follows:

• The amidino group of arginine is transferred to glycine to form guanidoacetic acid. This step occurs in the kidneys and pancreas.

• Guanido acetic acid is methylated by S-adenosyl methionine to form creatine. This step occurs in the liver.

Creatine is phosphorylated to creatine phosphate by the enzyme creatine kinase, which is present in the muscle, brain and liver. The reaction needs hydrolysis of ATP. The stored creatine phosphate in the muscle serves as an immediate store of energy in the muscle. During muscle contraction, the energy is first derived from ATP hydrolysis. Thereafter, the ATP is regenerated by the hydrolysis of creatine phosphate. This is called the Lohmann's reaction.

The creatine phosphate may be converted to its anhydride, creatinine, which is excreted in the urine. Creatinine level in the blood is an indicator of renal function.

Other options :

Option A - Gamma amino butyric acid (GABA) functions in brain tissue as an inhibitory neurotransmitter. It is formed by decarboxylation of glutamate by L-glutamate decarboxylase.

Option B - Dopa is a precursor in the formation of the neurotransmitters dopamine, norepinephrine and epinephrine. It is formed by the hydroxylation of tyrosine by tyrosine hydroxylase.

Option C - 5 Hydroxytryptamine is an important neurotransmitter in the brain, where it functions as an antidepressant. It is derived from the amino acid tryptophan, in a process of hydroxylation followed by decarboxylation.

Solution to Question 12:

Tandem mass spectrometry is used to screen a blood sample for metabolic diseases.

Mass spectrometry is a very sensitive quantitative analytical technique used to measure a wide range of clinically relevant analytes. It identifies molecules based on their mass or molecular size. A sample is vaporized such that a positive charge is produced on the molecule. An electric field is applied so that particles move. At the same time, a magnetic field is applied, so the cations are deflected and hit on the detector. For molecules of identical charge, the time required for the molecule to reach the detector is inversely proportional to the mass of the molecule.

In tandem mass spectrometry, two mass spectrometers (MS) are arranged sequentially. The first MS separates big peptides, based on their mass. From this observation, the parent ion can be directed into the second MS. Here the ions collide with argon gas molecules and are broken into smaller ions. These daughter ions are detected and quantified by their mass spectrum in the second instrument.

Tandem MS is used to screen blood samples from newborns for the presence and concentrations of amino acids, fatty acids and other metabolites. Abnormalities in metabolite levels can serve as diagnostic indicators for a variety of genetic disorders like phenylketonuria.

Solution to Question 13:

Familial hypercholesterolemia occurs due to a defect in the LDL (low-density lipoprotein) receptor.

The primary function of LDL particles is to provide cholesterol to the peripheral tissues (or return it to the liver). They do so by binding to plasma membrane LDL receptors that recognize apo B-100.

After binding, the LDL–receptor complex is endocytosed. The pH of the endosome falls, which allows the separation of the LDL from its receptor. The receptors can be recycled, whereas the lipoprotein remnants in the vesicle are transferred to lysosomes and degraded by lysosomal acid hydrolases, releasing free cholesterol, amino acids, fatty acids, and phospholipids. These compounds can be reutilized by the cell.

Defects in the synthesis of functional LDL receptors cause a significant elevation in plasma LDL-C. Patients with such deficiencies have type IIa hyperlipidemia /familial hypercholesterolemia (FH). This autosomal dominant condition predisposes individuals to premature atherosclerosis and coronary artery disease.

Patients with homozygous FH must be treated aggressively to delay the onset and progression of CVD. Two drugs that reduce the hepatic production of VLDL and thus LDL, a small-molecule inhibitor of the microsomal TG transfer protein (MTP) and an antisense oligonucleotide to apoB, and an antibody that inhibits ANGPLT3 are approved and should be considered in patients who have insufficient response to statins and PCSK9 inhibitors. LDL apheresis should be considered in patients who have persistently elevated LDL-C levels despite drug therapy.

Other options :

Option B - Defects in apo-E lead to type III hyperlipoproteinemia / familial dysbetalipoproteinemia. It is characterized by defective clearance of chylomicron remnants and intermediate-density lipoproteins by the liver. This leads to hypercholesterolemia, xanthomas, and atherosclerosis.

Option C - Defective apo-C II results in type I hyperlipoproteinemia/ familial lipoprotein lipase deficiency. The resulting inactive lipoprotein lipase leads to slow clearance of chylomicrons and VLDL, and low levels of LDL and HDL. Patients present with eruptive xanthomas and acute pancreatitis.

Option D - Apo-B 48 is involved in the assembly of chylomicrons. Defects in this protein are not associated with familial hypercholesterolemia.

Solution to Question 14:

In a patient with Diabetes Mellitus, the liver produces more TG and VLDL, due to decreased action of LPL with increased action of HSL.

• Lipoprotein lipase (LPL): Enzyme responsible for hydrolyzing triglycerides (TG) in VLDL and chylomicrons into free fatty acids (FFAs) for storage or energy use in tissues.

• Hormone-sensitive lipase (HSL): Enzyme that breaks down stored triglycerides in adipocytes into free fatty acids and glycerol.

Insulin has an inhibitory effect on hormone-sensitive lipase (HSL). In diabetes, low insulin corresponds to an increase in HSL activity. This promotes the breakdown of lipids and a release of TG and free fatty acids from the adipose tissue.

Insulin induces the activity of lipoprotein lipase (LPL) which is present on the capillary walls. LPL promotes hydrolysis of TG and the degradation of the transport proteins, VLDL and chylomicrons. In diabetes, low insulin leads to decreased LPL activity which corresponds to an increase in VLDL and TG.

This results in increased production of triglycerides and VLDL in the liver.

Solution to Question 15:

Nutritional deficiency of copper, required by lysyl oxidase for formation for covalent cross linking in collagen synthesis, gives rise to Menke's disease.

Menke's disease is an X-linked recessive condition caused due to a mutation in the ATP7A gene on chromosome X. This causes impaired transport of dietary copper from the small intestine to the liver leading to low levels of copper-containing enzymes.

The final step of collagen synthesis involves the formation of covalent cross-links. This is done by the enzyme lysyl oxidase, a copper-containing protein. This is essential for the strength and rigidity of collagen. As a result of deficiency of copper in Menkes disease, this step of collagen synthesis is affected.

The patients presents with depigmented kinky hair, failure to thrive, neurodegenerative symptoms (loss of developmental milestones, truncal hypotonia, epilepsy), multiple fractures, anaemia due to ceruloplasmin deficiency, neutropenia, and recurrent infections.

Other Options:

Option B: Scurvy: Caused by vitamin C deficiency, leading to defective collagen synthesis due to impaired hydroxylation of proline and lysine.

Option C: Ehlers-Danlos syndrome: A group of genetic disorders affecting collagen structure or production.

Option D: Scoliosis: A structural deformity of the spine.

Solution to Question 16:

The correct matched sequence is : i-b, ii-d, iii-a, iv-c.

i. Melatonin – b. Tryptophan

Tryptophan is an essential amino acid and the precursor for the synthesis of melatonin, produced by the pineal gland. The process begins with tryptophan being converted into serotonin, and then serotonin is further converted into melatonin.

ii. Histamine – d. Histidine

Histamine is derived from the amino acid histidine. The conversion of histidine into histamine is catalyzed by the enzyme histidine decarboxylase. Histamine is involved in local immune responses, acts as a neurotransmitter, and plays a role in regulating gastric acid secretion.

iii. Norepinephrine – a. Tyrosine

Norepinephrine (noradrenaline) is synthesized from tyrosine. Tyrosine is first converted to L-DOPA by the enzyme tyrosine hydroxylase, and then L-DOPA is converted into dopamine. Finally, dopamine is hydroxylated to form norepinephrine. Norepinephrine is a neurotransmitter involved in the sympathetic nervous system's fight-or-flight response.

iv. Glutathione – c. Cystine

Glutathione is a tripeptide composed of glutamate, cystine, and glycine. The cystine component is crucial because it contains a thiol (-SH) group, which is responsible for the antioxidant properties of glutathione.

i. Melatonin	b. Tryptophan
ii. Histamine	d. Histidine
iii. Norepinephrine	a. Tyrosine
iv. Glutathione	c. Cystine

Solution to Question 17:

The amino acid that provides phenylbutyrate in the treatment of urea cycle disorders is glutamate.

Ammonia to Urea cycle: Ammonia gets converted to glutamate (non toxic form) through transamination. Glutamate is transported to the liver as glutamine (by glutamine synthase). In the liver, ammonia is released from glutamine, converting it to alpha keto glutatamte. The released ammonia enters the urea cycle.

In acetylation therapy, Phenylbutyrate (a prodrug) which is metabolized to phenylacetate. Phenylacetate combines with glutamine (a nitrogen-rich amino acid) to form phenylacetylglutamine. Phenylacetylglutamine is water-soluble and is excreted in the urine. This process helps trap excess nitrogen, bypassing the urea cycle.

Other drugs that can be used to achieve the same outcome (ammonia scavengers) include:

• Phenylacetate

• Benzoate

Other Options:

Option A and C: Arginine and Aspartate do not play a role in this specific reaction.

Option D: Glycine: The combination of benzoate and glycine forms hippuric acid, a non-toxic compound. When Benzoate is administered to the body, it acts as a nitrogen-scavenging agent. Hippuric acid is water-soluble and is excreted in the urine, effectively removing one nitrogen molecule from the body per molecule of hippuric acid formed. This mechanism is similar to how phenylacetate acts as a scavenger for glutamate.

Solution to Question 18:

In situ hybridization, Conventional cytogenetics, Sanger sequencing and Polymerase chain reaction are appropriate techniques for chromosomal aneuploidy.

Techniques to detect aneuploidy:

• In situ hybridization: Fluorescent In Situ Hybridisation (FISH) helps in the detection of specific genetic information using fluorescent probes. FISH is used to detect chromosomal aneuploidy, subtle microdeletions and rearrangements, complex translocations, and gene amplification.

• Conventional cytogenetics: This technique involves the study of chromosomes under a microscope, typically using methods like karyotyping, which arranges chromosomes by size, shape, and number. The analysis allows the identification of large chromosomal abnormalities, such as aneuploidy or structural changes.

• Sanger sequencing: DNA sequencing that uses chain-termination to identify the exact sequence of nucleotides in a DNA molecule. Simple Senger sequencing can detect aneuploidy only with massive parallel sequencing.

• Polymerase chain reaction: Polymerase chain reaction (PCR) is a rapid automated method for the amplification of specific DNA sequences (or genes). It can accurately quantify the target nucleic acid sequence within a sample. It is useful for determining somatic mutation with low allele frequency. Quantitative Fluorescence (QF) PCR is a method of rapid aneuploidy detection. It

is less expensive and allows simultaneous processing of large number of samples of FISH.

• Array Comparitive Genomic Hybridization: microarray based technique which can reveal imbalances across the genome.

• Multiplex Ligation Probe Amplification: to detect gene dosage.

Solution to Question 19:

Phosphatase, Terminal transferase and Crispr Cas9 are used in recombinant DNA technology.

Recombinant DNA technology is a technique where DNA from different sources is combined to create new genetic material. Applications of recombinant DNA technology include producing insulin, growth hormones, vaccines, gene therapy, and studying gene functions and disease mechanisms.

Phosphatase dephosphorylates 5′ ends of RNA and DNA. It is primarily used for the removal of phosphate groups pror to kinase labelling and it is also used to prevent self ligation.

Terminal transferase catalyzes the addition of nucleotides to the 3' terminus of a DNA molecule. It is used in homopolymer tailing.

CRISPR-Cas9 acts as an RNA-directed nuclease. It is a DNA editing/gene regulatory system. It involves two essential components: Cas9, an endonuclease that causes a double-stranded DNA break, allowing modifications to the genomic DNA to be made; and a synthetic single-guide RNA, or sgRNA, responsible for guiding Cas9 to its target.

Isomerase is a class of enzymes that catalyze the structural rearrangement of molecules, converting one isomer into another. Isomerase is not commonly used directly in recombinant DNA technology, it has broader applications. They are used in metabolic pathways and as biocatalysts in industrial processes.

Biochemistry NEET 2024

Question 1:

A patient underwent gastrectomy for gastric adenocarcinoma. Which of the following vitamins needs to be supplemented in this patient?

a) Vitamin B12

b) Vitamin B1

c) Vitamin A

d) Vitamin D

Question 2:

A 10-year-old child from a southern Indian state, hailing from a poor family, presents with rough and scaly skin. The child's diet lacks vegetables and fruits. After treatment with linseed oil, Vitamin A, and Vitamin B, the symptoms subsided. What is the most likely diagnosis?

a) Lichen planus

b) Erythroderma

c) Phrynoderma

d) Keratosis pilaris

Question 3:

A patient presents to the emergency department with severe abdominal pain, nausea, and vomiting. The laboratory results show serum triglyceride levels > 2000 mg/dl. Which deficiency could be contributing to the patient's condition?

a) Apolipoprotein C II

b) Apolipoprotein B48

c) Apolipoprotein B100

d) Apolipoprotein A I

Question 4:

After the age of 12, the boy had an intellectual disability. On examination, fingernail abrasions, excessive biting of lips, and increased levels of serum uric acid were observed. What is the likely diagnosis?

a) Hurler syndrome

b) Lesch-Nyhan disease

c) Hunter syndrome

d) Von Gierke syndrome

Question 5:

A neonate was brought with features of lethargy, and poor oral intake. On examination, a cataract was seen. Urine Benedict's test was positive but negative for Glucose. What will be the enzyme deficiency in this condition?

a) Hexokinase

b) Galactokinase

c) Galactose-1-phosphate uridyl transferase

d) Phosphofructokinase

Question 6:

A patient diagnosed with a rare homozygous recessive condition presented with ectopia lentis. In the peripheral smear, there was no megaloblastic anemia. Later, he was found to have elevated homocysteine levels in the blood. Which of the following supplementations will help the patient?

a) Thiamine

b) Pyridoxine

c) Biotin

d) Riboflavin

Question 7:

You received a text message from your friend JOHN DOE about occupational exposure after he visited his doctor as shown below. What is the test used to diagnose?



©MARROW

a) Anthracosis

b) Lead poisoning

c) Silicosis

d) Asbestosis

Question 8:

A 3-year-old child is brought to the OPD with complaints of persistent delayed developmental milestones, and hyperpigmented macular lesions on the body which are more pronounced on exposure to sunlight. What is the error in DNA multiplication?

a) Base excision repair

b) Nucleotide excision repair

c) DNA mismatch repair

d) Homologous recombination

Question 9:

A patient with a family history of coronary heart disease presented with tendon xanthomas. His lipid profile showed serum LDL to be 280 mg/dL and triglyceride levels was 398 mg/dl. What is the likely defect in this patient?

a) LDL receptor defect

b) LDL metabolism

c) VLDL metabolism

d) Lipoprotein lipase deficiency

Question 10:

Which food item among these is the primary source of thiamine in Indians?

a) 1

b) 2

c) 3

d) 4

Question 11:

A young female comes to the emergency department after ingesting 10 tablets of Phenobarbital. Which of the following components of the ETC is inhibited by this drug?

a) Succinate - Fe

b) Cytochrome I-II

c) Complex I to Q

d) Complex IV

Question 12:

A child presents to you with bone pain, hepatosplenomegaly, and pancytopenia. A trephine biopsy aspirate shows the following finding. Which of the following is the most likely enzyme deficient in this patient?



a) Galactocerebosidase

b) Glucocerebrosidase

c) Sphingomyelinase

d) Alpha 1,4-glucosidase

Question 13:

A 32-year-old lady comes with complaints of erythematous scaly rash, loose stools, and cognitive dysfunction. The rash involved her forearm, neck, and face. She seems to have difficulty remembering things from the recent past. Her electrolytes are within normal range. She admits to having shifted to a corn-based diet due to economic reasons. Which of the following would improve her symptoms?

a) Oral Nicotinamide supplementation

b) High dose of IV thiamine

c) High Dose Vitamin C

d) Stop corn diet

Question 14:

A 3-month-old child was brought to the clinic with a history of difficulty in feeding and failure to thrive. On examination, the child is floppy, 48 cm in length and weighs 2.8 kg. There is generalized hypotonia, macroglossia and hepatomegaly. The chest x-ray of the child is given below. What is the most likely diagnosis?



a) Ebstein's anomaly

b) Pompe's disease

c) Transposition of great arteries

d) Von- Gierke's disease

Question 15:

A patient presented with multiple yellowish raised plaques on eyelids, xanthomas on the achilles tendon. What is the likely cause for the patient's presentation?

a) Familial hypercholesterolemia

b) Tangier's disease

c) Xanthomatosis

d) Fabry's disease

Question 16:

A child presented to the OPD with complaints of fatigue and muscle weakness while exercising. Laboratory investigations found that LCFA oxidation was affected. A muscle biopsy was done which revealed the presence of fat vacuoles. What is the probable diagnosis?

a) LDL defect

b) LPL defect

c) Fatty acid synthase defect

d) Carnitine deficiency

Question 17:

A mother came back home to see her child suffering from difficulty breathing. The child had contact with rotenone while trying to bathe their pet. The child was taken to the emergency department and she later passed away. Inhibition of which of the following enzymes was the likely cause of her death?

a) Cytochrome c reductase

b) Succinate dehydrogenase

c) NADH-Q oxidoreductase

d) Cytochrome oxidase

Question 18:

A child suffering from phenylketonuria was started on necessary dietary modifications but the neurological symptoms persisted. Phenylalanine hydroxylase levels in the serum are normal. Which of the following enzymes is likely to be deficient in this case?

a) Tyrosine hydroxylase

b) Homogentisate oxidase

c) Tetrahydrobiopterin synthase

d) Dopa decarboxylase

Question 19:

Which micronutrient is most likely deficient in a patient with the radiographic findings given below along with anemia, neutropenia, depigmented hair and skin?

a) Iron

b) Zinc

c) Copper

d) Fluoride

Question 20:

A child presented with recurrent abdominal pain, vomiting, and diarrhea following ingestion of milk-based products. Which enzyme is likely to be deficient?

a) Lipase

b) Maltase

c) Amylase

d) Lactase

Question 21:

In the given Lineweaver-Burk plot, the blue line represents the action of an enzyme, and the red line represents the action of the same enzyme in the presence of an inhibitor. What is the type of inhibition shown here?



©MARROW

a) Competitive inhibition

b) Non-competitive inhibition

c) Uncompetitive inhibition

d) Allosteric inhibition

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	b
5	c
6	b
7	b
8	b
9	a
10	d
11	c
12	b
13	a
14	b

15	a
16	d
17	c
18	c
19	c
20	d
21	a

Detailed Explanations

Solution to Question 1:

A patient who has undergone gastrectomy for gastric adenocarcinoma needs to be supplemented with vitamin B12.

Vitamin B12 absorption is a complex process that relies on several factors. Intrinsic factor, a glycoprotein secreted by the parietal cells in the stomach, is essential for this process. Gastric acid and pepsin release vitamin B12 from dietary proteins, allowing it to bind with cobalophilin in the saliva. In the jejunum, cobalophilin is broken down, freeing vitamin B12 to bind with intrinsic factor. This complex is then absorbed in the distal ileum through specific receptors. Following gastrectomy, the production of intrinsic factor is impaired due to the removal of gastric tissue, making vitamin B12 absorption difficult. Therefore, vitamin B12 supplementation is necessary after gastrectomy to prevent deficiency.

Other complications of gastrectomy include:

• Nutritional deficiencies- Iron deficiency anemia (most common), Vitamin B12 deficiency, and osteoporosis

• Dumping syndrome: early and late

• Diarrhea

• Duodenal stump leak/blow out

• Bile reflux gastritis and esophagitis

• Gall stones

• Roux syndrome

Solution to Question 2:

The cutaneous lesions with rough and scaly skin, with dietary history stating lack of vegetables and fruits is suggestive of phrynoderma.

Phrynoderma are keratotic follicular papules commonly present over the elbows and knees. They can also extend to the shoulders, neck, back, thighs, buttocks, and abdomen. It is also known

as toad skin, due to the rough cone-shaped elevated papules. It can caused by deficiency of vitamins A, B complex, C, and E and essential fatty acids.

When it occurs due to vitamin A deficiency, it is associated with ophthalmic symptoms like xerophthalmia, night blindness, and Bitot's spots and cutaneous symptoms like xerosis, scaling, and acne.

Management includes nutritional improvement through diet, multivitamin supplementation, and treatment of the underlying cause.

Note: Linseed oil is rich in alpha-linolenic acid, which is an essential fatty acid.

The toad skin appearance of a child with phrynoderma is given below:

Other options:

Option A: Lichen planus is a papulosquamous lesion presenting as pruritic, purple, polygonal, planar, papules and plaques. These papules are often highlighted by white dots or lines called Wickham striae, which are created by areas of hypergranulosis. The characteristic lesions in lichen planus are shown below:

Option B: Erythroderma is a serious skin condition characterized by widespread redness (erythema) and scaling affecting more than 90% of the body's surface area. It is also known as exfoliative dermatitis and often results from underlying skin disorders, systemic diseases, or drug reactions.

Option D: Keratosis pilaris is an inherited abnormality of keratinization. It affects the follicular orifices leading to keratotic follicular plugging, perifollicular erythema, and follicular atrophy. It usually occurs during puberty. They present as small, keratotic papules on the extensor aspects of the limbs, particularly the arms which can lead to ingrowing of the hair. Treatment is purely reserved for cosmetic purposes.

Solution to Question 3:

The elevated triglyceride levels and symptoms of acute pancreatitis are likely related to a deficiency in apolipoprotein C II.

Apolipoprotein C2 is a lipoprotein lipase cofactor that catalyzes cleavage of chylomicrons. Defective apo C II results in type I hyperlipoproteinemia/ familial lipoprotein lipase deficiency. The resulting inactive lipoprotein lipase leads to slow clearance of chylomicrons and VLDL, and low levels of LDL and HDL. Patients present with eruptive xanthomas and acute pancreatitis.

Lipoprotein lipase is an extracellular enzyme that is present in the capillary walls of most tissues. It is chiefly synthesized by the adipose tissue, and cardiac and skeletal muscles. A deficiency of lipoprotein lipase or apo C-II leads to type 1 hyperlipoproteinemia or familial chylomicronemia which leads to an increased risk of acute pancreatitis.

Other options:

Option B: Apo B48 is involved in the assembly of chylomicrons. Defects in this protein are not associated with familial hypercholesterolemia.

Option C: Apo B100 binds low-density lipoprotein receptors. For the uptake of LDL, apo B100 acts as a ligand for the low-density lipoprotein receptor.

Option D: Apo A1 activates lecithin cholesterol acyltransferase (LCAT). LCAT is the enzyme present in HDL, which is activated by apo A1. LCAT converts free cholesterol (amphipathic) in HDL to cholesterol ester(hydrophobic). The initial discoidal shape now becomes spherical. This facilitates reverse cholesterol transport.

Solution to Question 4:

The above clinical scenario with findings showing self-mutilation and laboratory values of hyperuricemia is most likely suggestive of Lesch-Nyhan disease.

Lesch-Nyhan syndrome is an X-linked recessive disorder due to a complete deficiency of the hypoxanthine-guanine phosphoribosyltransferase (HGPRT). This deficiency results in an inability to salvage hypoxanthine or guanine leading to increased levels of phospho-ribosyl pyrophosphate (PRPP). Increased PRPP leads to increased purine synthesis, which is later degraded to uric acid. Increased hypoxanthine is also converted to uric acid.

The patients have intellectual disability, behaviors of self-injury like lip and finger biting, developmental delay, muscle dystonia, and extrapyramidal symptoms.

Diagnosis is made by the elevated levels of uric acid and by HGPRT enzyme assay.

The image given below shows the metabolism of hypoxanthine and guanine by HGPRT.



Orange diaper syndrome can be an early indicator of Lesch-Nyhan syndrome. At birth, individuals with Lesch-Nyhan syndrome typically appear asymptomatic, with normal prenatal growth and development. The condition is often first detected through the presence of orange-coloured crystals in the diaper, resulting from hyperuricemia. Since hyperuricemia is a common feature across all variants of HGPRT deficiency, which causes Lesch-Nyhan syndrome, predicting associated neurodevelopmental abnormalities requires enzyme activity testing.

Solution to Question 5:

The given clinical scenario of a neonate presenting with lethargy, and poor oral intake, with examination revealing a cataract and urine benedict's test was positive but negative for glucose, is suggestive of classical galactosemia and enzyme deficient is galactose-1-phosphate uridyl transferase.

Galactose-1-phosphate uridyl transferase deficiency leads to classical galactosemia. It is an autosomal recessive disorder. Normally, newborns receive up to 40% of their intake as lactose (glucose + galactose). The deficiency of this enzyme leads to an inability to convert galactose-1-phosphate to uridine diphosphate galactose. This leads to the accumulation of galactose 1-phosphate and causes injury to the kidney, liver, and brain.

Clinical features:

• Accumulation of galactose-1-phosphate in the liver causes liver damage & leads to hepatomegaly, vomiting, and convulsions.

• Accumulation of galactose-1-phosphate depletes the liver of inorganic phosphate leading to diminished ATP synthesis which affects gluconeogenesis resulting in hypoglycemia.

• Accumulation of galactose-1-phosphate in the nerves causes intellectual disability.

• Accumulation of galactitol in the lens causes oil-drop cataracts.

• Patients with galactosemia are at an increased risk of developing Escherichia coli neonatal sepsis.

In uridyl transferase deficiency, the epimerase is present in adequate amounts, so that the galactosemic individual can still form UDPGal from glucose. This explains how normal growth and development of affected children can occur despite the galactose-free diets.

Urinary reducing sugar (Benedict's test) and mucic acid test are positive. Direct enzyme assay using erythrocytes establishes the diagnosis.

Treatment is a galactose-free diet.



Other options:

Option A: Hexokinase is a glycolytic enzyme, involved in maintaining intracellular glucose concentration by catalysing the first step in glucose metabolism, i.e., phosphorylation of intracellular glucose. Its deficiency leads to hemolytic anaemia. Splenomegaly may be another presenting feature.

Option B: Galactokinase is the enzyme responsible for the conversion of galactose to galactose 1-phosphate. Only cataract is seen in the deficiency of this enzyme because galactose-1-phosphate is not formed. Hence no toxic effects of this compound are manifested.

Option D: Phosphofructokinase is a key regulatory enzyme of glycolysis and is necessary for the ATP-dependent conversion of fructose-6-phosphate to fructose-1,6-diphosphate. Tauri's disease is type VII glycogen storage disorder that results due to the deficiency of phosphofructokinase.

Solution to Question 6:

The vitamin that can be used as supplementation in this patient with homocystinuria is pyridoxine (vitamin B6).

Classical homocystinuria is an autosomal recessive disease most commonly caused by a deficiency of the cystathionine β-synthase. The deficiency of this enzyme leads to an accumulation of homocysteine in the blood and a clinical syndrome of intellectual disability, subluxated ocular lenses, long and slender extremities, arachnodactyly, and arterial and venous thrombi. The skeletal features of homocystinuria often mimic Marfan syndrome. Folic acid and vitamin B12

deficiency can also lead to homocystinuria with megaloblastic anemia.



Demonstration of elevated total homocysteine in the blood confirms the diagnosis. 50% of the cases of homocystinuria are termed pyridoxine-responsive because large doses of pyridoxine can enhance the activity of deficient enzymes in these patients. Administration of large doses of pyridoxine (vitamin B6) restores the activity of the deficient enzyme in half of all cases of homocystinuria.

Folate supplementation is also added to overcome any folate deficiency that might occur due to increased conversion of homocysteine to methionine. For the other 50% of the patients, the treatment consists of methionine restriction and cysteine supplementation. In addition, betaine (methyl group donor), folate and vitamin B12 may be added.

Solution to Question 7:

The test for coproporphyrinogen levels is used as a screening test to diagnose lead poisoning.

Lead inhibits the enzymes delta-aminolevulinic acid (ALA) dehydratase and ferrochelatase, which are involved in heme synthesis. ALA is converted to porphobilinogen by ALA dehydratase, inhibited by lead. So, serum porphobilinogen levels are decreased in lead poisoning.

Laboratory findings include the following:

• An elevated serum lead level

• An elevated level of protoporphyrin in blood and elevated ALA and coproporphyrinogen in urine

• Microcytic anemia with basophilic stippling in peripheral smear due to ineffective heme synthesis

Clinical features of chronic lead poisoning include facial pallor, anemia, basophilic stippling of red cells, blue lines in the gums, and retinal stippling. In the later phase, the findings are colic abdominal pain, constipation, nerve palsy, encephalopathy, and disturbance of the genito-urinary and cardiovascular systems.

Diagnosis of lead poisoning:

Laboratory parameter	Value	Remarks
Coproporphyrin in urine	>150 mcg/L	Screening test- indicates exposure
Aminolevulinic acid (ALA) in urine	>5 mg/L	Indicates lead absorption
Lead in blood	>70 mcg/dL	Clinical symptoms appear
Lead in urine	0.8 mg/L	Indicates lead absorption
Basophilic stippling of RBCs		Sensitive indicator

Solution to Question 8:

The history points to a diagnosis of xeroderma pigmentosum. It is characterized by defective nucleotide excision repair.

UV radiation leads to pyrimidine dimerization in DNA strands, preventing further replication by DNA polymerase. The nucleotide excision repair mechanism corrects the dimer formation in DNA. Xeroderma pigmentosum (XP) is an autosomal recessive genetic disorder characterized by an inability to repair DNA pyrimidine dimers caused by UV exposure. This leads to clinical and cellular hypersensitivity to ultraviolet (UV) radiation.

In XP the skin is normal at birth and has a three-stage progression:

• Diffuse erythema, scaling, and freckle-like areas over the sun-exposed regions of the body at around 6 months of age.

• The second stage is characterized by poikiloderma. This results in skin atrophy, telangiectasias with mottled hyperpigmentation, and hyperpigmentation.

• In the third stage appearance of cutaneous malignancies like squamous cell carcinomas, malignant melanoma, basal cell carcinoma, and fibrosarcoma.

Most of the cases with XP have ocular manifestations. Initially, it appears as photophobia, conjunctivitis, ectropion, and symblepharon with ulceration. Accumulation of pyrimidine dimer causes corneal opacification and neovascularization. Ocular malignancies are common in the lids, conjunctiva, and cornea as they are exposed to as much as facial skin. XP is ultimately fatal, and early detection of the skin lesions and their management with prevention of UV ray exposures prolong life.

Solution to Question 9:

The clinical presentation with tendon xanthomas, elevated serum LDL and elevated serum triglycerides points to a diagnosis of familial hypercholesterolemia and is due to a defect in LDL

receptors.

Familial hypercholesterolemia (type IIa hyperlipoproteinemia) is an inherited disorder due to defects in LDL receptors. This defect causes a decrease in the uptake of LDL and cholesterol. As a result, the blood levels of both LDL and cholesterol are elevated. Such patients are at increased risk of developing atherosclerosis and coronary artery disease.

Features of FH:

• Painful polyarthritis and tendinitis are common.

• Physical findings seen are corneal arcus and tendon xanthomas over the Achilles tendon and the dorsum of the hands. Xanthomas are tumorous masses produced by foamy cell clusters (lipid-laden macrophages) found in the connective tissue of the skin and in tendons.

Diagnosis is done clinically based on hypercholesterolemia with

• LDL > 190 mg/dl,

• Family history of hypercholesterolemia and/or

• Premature coronary artery disease.

Treatment includes lifestyle changes, restriction of a fatty diet, and pharmacotherapy with statins in heterozygous FH patients.

Recent drugs for specific treatment of homozygous FH are:

• Lomitapide, a microsomal triglyceride transfer protein (MTP) inhibitor, and

• Mipomersen, an antisense oligonucleotide against apoB.

The image below shows Achilles tendon xanthoma and a plain lateral radiograph of both ankles showing bilateral symmetrical swellings (arrows) superior to the calcaneum respectively.

Note: Lipoprotein lipase (LPL) deficiency causes type I hyperlipoproteinemia due to defects in LPL or apo C-II. Eruptive xanthomas are seen. Chylomicrons and VLDL are elevated. LDL and HDL are low. There is no risk for coronary disease.

Solution to Question 10:

Among the various food items given on the plate, the primary source of thiamine (vitamin B1) in the Indian diet is option 4 i.e. chapati and rice.

The thiamine content in various foods in decreasing order are as follows:

• Gingelly seeds

• Groundnut

• Bengal gram

• Whole wheat

• Rice

Gingelly (sesame) seeds have the highest content of thiamine.

Thiamine is lost from rice during the process of milling. As it is a water-soluble vitamin, washing, and cooking rice cause further losses. Hence, it is advisable to avoid highly polished rice and eat parboiled or under-milled rice.

Thiamine is essential for energy metabolism, particularly in carbohydrate processing. Its active form, thiamine diphosphate, acts as a coenzyme for three critical enzyme complexes: pyruvate dehydrogenase, α-ketoglutarate dehydrogenase, and branched-chain keto acid dehydrogenase. This coenzyme facilitates oxidative decarboxylation by forming a reactive carbanion that attaches to carbonyl groups like those in pyruvate, leading to the release of CO_2. Thiamine diphosphate is also vital for transketolase in the pentose phosphate pathway.

Thiamine triphosphate, another form of thiamin, helps regulate nerve function by activating chloride channels in nerve membranes.

A thiamine deficiency can cause several conditions such as beriberi, characterized by peripheral neuritis, which can also involve heart failure and edema. Additionally, thiamine deficiency can lead to Wernicke's encephalopathy and Korsakoff's psychosis, commonly associated with alcohol or drug abuse. The lack of thiamine impairs the conversion of pyruvate to acetyl-CoA, leading to increased levels of lactate and pyruvate in the blood, which can result in dangerous lactic acidosis, especially in those consuming high-carbohydrate diets.



Solution to Question 11:

Phenobarbital is a barbiturate that inhibits complex I to Q of the electron transport chain.

Complex I of the electron transport chain is NADH-Q oxidoreductase. It oxidizes NADH and reduces coenzyme Q. Phenobarbital inhibits complex I to Q so that the electrons will not be able to be transferred from NADH to coenzyme Q, which is the first step in the electron transport chain. As a result, the rate of oxygen consumption will decrease.

The electron transport chain (ETC) represents the final pathway through which electrons, originating from the oxidation of carbohydrates, fatty acids, and amino acids, are channeled toward a final reaction with oxygen to produce water and ATP. The ETC complexes are embedded in the inner mitochondrial membrane.

The oxidation of compounds by ETC complexes leads to the release of free energy. This is used to pump H+ to the intermembrane space. This results in the formation of a proton motive force that is used for the phosphorylation of ADP to ATP. Thus, oxidation is coupled with phosphorylation to generate ATP. NADH generates 2.5 ATP and FADH2 generates 1.5 ATP.

The table below summarizes the inhibitors of the electron transport chain.

Complexes of ETC	Inhibitors
Complex I:NADH-Q oxidoreductase	Piericidin A,phenobarbital, rotenone
Complex II:Succinate-Q reductase	TTFA (thenoyltrifluoroacetone), carboxin, malonate
Complex III:Q- Cyt c oxidoreductase	BAL (British anti-lewisite), antimycin A
Complex IV:Cyt c oxidase	Hydrogen sulfide, carbon monoxide, cyanide
Complex V:ATP Synthase	Atractyloside, oligomycin

Solution to Question 12:

The given clinical scenario of a child with bone pain, hepatosplenomegaly, and trephine biopsy showing Gaucher's cell with a characteristic crumpled tissue paper appearance is suggestive of Gaucher's disease. The deficient enzyme is glucocerebrosidase (also known as beta-glucosidase).

Gaucher's disease is an inborn error of metabolism caused due to a deficiency of beta-glucocerebrosidase. It is the most common lysosomal storage disorder and the mode of inheritance is autosomal recessive. There is an accumulation of lipid-laden macrophages in the spleen, liver, and other tissues.

There are three clinical subtypes - type I (chronic non-neuronopathic), type II (acute neuronopathic), and type III (intermediate). Clinical features include anemia, thrombocytopenia, hepatosplenomegaly, and neurological features depending on the subtype.

Gaucher's cells are formed as a result of the accumulation of massive amounts of glucocerebrosides within the phagocytes. The cells have a fibrillary cytoplasm and give a crumpled tissue paper appearance.

Solution to Question 13:

The given clinical features suggests the diagnosis of pellagra, due to her corn-based diet. Oral nicotinamide supplements can improve her symtpoms.

Niacin can be synthesized in the body from tryptophan. Two compounds, nicotinic acid and nicotinamide, have the biological activity of niacin. The nicotinamide ring of the coenzymes nicotinamide adenine dinucleotide (NAD) and nicotinamide adenine dinucleotide phosphate (NADP) participate in oxidation/reduction reactions.

The initial symptoms of pellagra include loss of appetite, generalized weakness, irritability, abdominal pain, and vomiting. This is followed by bright red glossitis and a characteristic pigmented, scaly skin rash in sun-exposed areas. This rash is known as Casal’s necklace when it forms a ring around the neck.

The 4Ds of pellagra:

• Diarrhea

• Dermatitis

• Dementia, leading to

• Death

It is diagnosed by measuring urinary levels of 2-methyl nicotinamide and 2-pyridone.

Other options:

Option B: Thiamine (Vitamin B1) deficiency leads to conditions like beriberi and Wernicke-Korsakoff syndrome, which are characterized by symptoms such as muscle weakness, peripheral neuropathy, cardiovascular issues, and confusion. While cognitive dysfunction can result from thiamine deficiency, rash and loose stools do not fit with thiamine deficiency. Therefore, thiamine would not address the full spectrum of this patient's symptoms.

Option C: Vitamin C deficiency causes scurvy, which is associated with symptoms like bleeding gums, bruising, joint pain, and fatigue. Scurvy is not associated with a rash like the one described, nor does it typically cause cognitive dysfunction or diarrhea. Hence, high-dose vitamin C would not improve this patient's symptoms.

Option D: Corn-based diets are low in niacin and tryptophan, leading to pellagra over time, especially if the diet is not balanced with other niacin-rich foods. While stopping the corn-based diet could prevent further niacin deficiency, it will not immediately address the existing deficiency and symptoms like rash, diarrhea, and cognitive dysfunction. The patient would still need niacin supplementation to resolve the current symptoms.

Solution to Question 14:

The child with a clinical history of difficulty feeding and failure to thrive with generalised hypotonia, macroglossia, hepatomegaly and chest x-ray findings of massive cardiomegaly suggest a diagnosis of Pompe's disease.

Pompe's disease is a glycogen storage disorder(type II) caused by a deficiency of lysosomal acid maltase (alpha-glucosidase) which leads to the accumulation of glycogen in the lysosomes of the liver (hepatomegaly), skeletal muscles in the form of large vacuoles (hypotonia and weakness), cardiac muscles (hypertrophic cardiomyopathy). The child generally dies by the age of 2 years due to cardiac failure. The adult form presents with muscle dystrophies.

Laboratory findings show elevated levels of serum creatine kinase (CK), aspartate transaminase (AST), alanine transaminase (ALT), and lactate dehydrogenase (LDH). Urine glucose tetrasaccharide, a glycogen breakdown metabolite, is a reliable biomarker for disease severity and treatment response. Treatment includes enzyme replacement therapy with recombinant enzyme a-glucosidase alfa (Myozyme).

Given below is the image of muscle biopsy showing glycogen-positive vacuoles.



Other options:

Option A- Ebstein's anomaly is the teratogenic effect of maternal lithium consumption which causes a cyanotic heart disease in the newborn with atrialization of the right ventricle and conduction defects. The chest X-ray will show a box-shaped enlargement of the right atrium and pulmonary oligemia.

Option C- Transposition of great vessels is cyanotic congenital heart disease where there is a switch in the position of two main arteries i.e. pulmonary artery and aorta. X-ray shows of egg-on-string appearance with pulmonary plethora.

Option D- Von Gierke disease occurs due to a deficiency of glucose-6-phosphatase. Patients present commonly in 3-4 months of age with hypoglycemia, lactic acidosis, hepatomegaly, growth retardation, hyperuricemia, and hyperlipidemia.

Solution to Question 15:

The clinical presentation with yellowish, raised plaques on eyelids, and tendon xanthomas points to a diagnosis of familial hypercholesterolemia (FH).

Familial hypercholesterolemia (type IIa hyperlipoproteinemia) is an inherited disorder due to defects in LDL receptors and Apo B100. This decreases the uptake of Low-density lipoprotein and cholesterol leading to their increased levels in the blood. The increased LDL accumulates in the blood vessels causing atherosclerosis (premature coronary artery disease)and in the Achilles tendon causing tendon xanthomas. Xanthomas are tumorous masses produced by foamy cell clusters (lipid-laden macrophages) found in the connective tissue of the skin and tendons. Patients can also have yellowish plaques around the eyes (Xanthelasma) and corneal arcus.

Treatment includes lifestyle changes, restriction of a fatty diet, and pharmacotherapy with statins in heterozygous FH patients.

The images below show Achilles tendon xanthoma and xanthelasma respectively:



Other options:

Option B: Tangier's disease is a hypolipoproteinemia caused by a mutation of the ABCA1 gene. Enlarged orange tonsils are pathognomonic. Low HDL levels are seen. There is an increased risk for atherosclerosis.

The image below shows enlarged yellow-orange coloured tonsils in Tangier's disease.



Option C: Xanthomatosis is the deposition of lipids in various parts of the body like tendon xanthomas, xanthelasma, eruptive and tuberous xanthomas. This is a sign of underlying lipid metabolism disorders.

Option D: Fabry's disease is a lysosomal storage disorder with a defect in alpha-galactosidase. It causes angiokeratomas, hypohidrosis and characteristic Maltese cross deposits in urine.

Solution to Question 16:

The clinical scenario of a child with fatigue and muscle weakness on exercise with laboratory findings of LCFA oxidation deficiency and fat vacuoles on muscle biopsy suggests a diagnosis of carnitine deficiency.

Carnitine deficiency is a defect in fatty acid oxidation that leads to hypoketotic hypoglycemia and hyperammonemia. Carnitine transports activated LCFA (long chain fatty acyl CoA) from the cytoplasm to the mitochondria for beta-oxidation (as they cannot cross the mitochondrial membrane), carnitine deficiency impairs this transport causing accumulation of lipids (skeletal muscle weakness). This excess fatty acid inhibits the urea cycle causing hyperammonemia. Since beta-oxidation cannot occur, energy production is reduced, and there is hypoglycemia without a corresponding increase in ketone bodies (hypoketotic) during times of glucose depletion such as exercise.

The role of carnitine in the metabolism of fatty acids is demonstrated in the image below:



Clinical features of carnitine deficiency begin in childhood and include poor feeding, lethargy, liver dysfunction and hepatomegaly, hypotonia, and dilated cardiomyopathy. Treatment involves oral supplementation of carnitine.

Other options:

Option A: An increase in LDL can cause hyperlipoproteinemia, and reduced LDL can cause defective absorption of fat-soluble vitamins.

Option B: LPL is involved in hydrolysis of lipoproteins in the blood to free fatty acid and glycerol which are then taken up by the tissues. A defective LPL can cause type 1 familial chylomicronemia syndrome presenting as pancreatitis.

Option C: Fatty acid synthase is a multienzyme complex involved in fatty acid synthesis in the cytosol which when defective may impair energy production, essential fatty acids, and affect metabolic pathways. There is no accumulation of fat vacuoles since fatty acids are not synthesised.

Solution to Question 17:

In the given clinical vignette, inhibition of NADH-Q oxidoreductase by rotenone was the likely cause of breathlessness, followed by death in this child.

Rotenone inhibits complex I, thereby inhibiting the respiratory chain, so electrons cannot be transferred to oxygen. This halts ATP production, leading to an energy deficit in the cells, eventually causing cell death. The ETC constitutes four enzyme complexes:

• Complex I: NADH-Q oxidoreductase

• Complex II: Succinate-Q reductase

• Complex III: Q-cytochrome c oxidoreductase

• Complex IV: cytochrome c oxidase

Components of complex I are FMN and Fe-S complex. NADH-Q oxidoreductase catalyses the transfer of electrons from NADH to coenzyme Q, along with the transfer of four H+ across the membrane.

Solution to Question 18:

This child with phenylketonuria with neurological symptoms not improving despite dietary modifications and normal serum phenylalanine hydroxylase level is suggestive of tetrahydrobiopterin (BH4) synthase deficiency.

A diet low in phenylalanine (eg: tapioca) is the mainstay of treatment for classic phenylketonuria but may not be sufficient in BH4 deficiency where supplementation with BH4 and deficient neurotransmitter precursors (L-dopa) is required to overcome the block.

Phenylketonuria (PKU) is an inborn error of metabolism seen due to a defect in the enzyme phenylalanine hydroxylase (classic phenylketonuria) and less commonly, due to a defect in BH4 synthase (non-classical). These defects result in the impaired conversion of phenylalanine to tyrosine. The patient develops clinical features due to elevated levels of phenylalanine metabolites and decreased levels of tyrosine. The excess phenylalanine in the blood is metabolized to phenylpyruvate (positive ferric chloride) and phenylacetate (mousy odour) which are excreted in the urine. Infants appear normal at birth and gradually develop symptoms of hyperreflexia, seizures, hypopigmentation, low IQ, hyperactivity, and tremors.

Tandem mass spectroscopy is the investigation of choice. Guthrie test is a screening test that shows with absence of growth of Bacillus subtilis. The ferric chloride test is positive with the development of a transient blue-green colour.

Other options -

Option A: Tyrosine hydroxylase is the rate-limiting enzyme in the synthesis of epinephrine, converts tyrosine to DOPA, its deficiency does not cause phenylketonuria

Option B: Homogentisic acid oxidase converts homogentisic acid to maleylacetoacetic acid in the tyrosine degradation pathway and its deficiency causes alkaptonuria with a dark colour of urine on standing, bluish-black pigmentation in connective tissue (ochronosis) and arthritis.

Option D: Dopa decarboxylase is another enzyme involved in the synthesis of epinephrine. It converts DOPA to dopamine in the presence of vitamin B6. While enzymes involved in epinephrine synthesis cause neurological dysfunction, they do not cause phenylketonuria as such.

Solution to Question 19:

The given clinical features of anaemia and neutropenia, along with depigmented hair and skin and the radiological findings suggestive of pelkan spurs, trummerfeld zone, Frenkel's line, and Wimberger's ring sign point to the diagnosis of Menke's disease. Patients with this condition have low serum levels of copper.

Menke's disease or Menke-Kinky hair syndrome is an X-linked recessive condition caused due to a mutation in the ATP7A gene on chromosome X. This causes impaired transport of dietary copper

from the small intestine to the liver leading to low levels of copper-containing enzymes.

The final step of collagen synthesis involves the formation of covalent cross-links. This is done by the enzyme lysyl oxidase, a copper-containing protein. This is essential for the strength and rigidity of collagen. As a result of deficiency of copper in Menkes disease, this step of collagen synthesis is affected.

The patient presents with depigmented kinky hair, failure to thrive, neurodegenerative symptoms (loss of developmental milestones, truncal hypotonia, epilepsy), multiple fractures, anaemia due to ceruloplasmin and hephaestin deficiency (converts ferrous to ferric to be bound to transferrin), neutropenia, and recurrent infections.

Lab investigations show low serum copper (<65 µg/dL) and low ceruloplasmin(<20 mg/dl). The disease is fatal and has no curative treatment.

A variety of distinct anomalies that are representative of scurvy and acquired copper insufficiency can be seen on radiographs of people with menke's disease.

These include



Early administration of L-histidine and copper chloride injections may be beneficial.

Other options:

Option A: Iron deficiency causes iron deficiency anemia, and hair loss (telogen effluvium. It does not cause neutropenia, depigmentation of hair and skin, or bony abnormalities.

Option B: Zinc deficiency causes acrodermatitis enteropathica characterized by periorificial dermatitis, diarrhea, and alopecia.

Option D: Fluoride is essential for bone and teeth mineralization and its deficiency causes decreased bone-mineral density, osteoporosis, and tooth decay.

Solution to Question 20:

The given scenario indicates intolerance of milk-based products in this child, hence, the enzyme likely to be deficient is lactase.

Lactase is the enzyme that is responsible for breaking down lactose, a disaccharide found in milk and dairy products. When this enzyme is deficient, it can lead to the accumulation of lactose in the intestinal lumen where it is fermented by intestinal bacteria to lactate which produces symptoms of lactose intolerance, including recurrent abdominal pain, vomiting, and diarrhea after consuming lactose-rich food items. Lactase and maltase are present in the intestinal brush border.

Other options:

Option A- Lipases are involved in the hydrolysis of dietary triglycerides to free fatty acids and glycerol. It is deficient in exocrine pancreatic deficiency states such as chronic pancreatitis, cystic fibrosis, and Crohn's. Although it might also present with abdominal pain and diarrhea, the stools are foul-smelling(due to gut bacteria) and greasy due to the presence of fat (steatorrhea) and are also associated with fat-soluble vitamin deficiency.

Option B- Maltase breaks down maltose which is a disaccharide intermediate of starch breakdown.

Option C- Amylase hydrolyses starch which is a glucose homopolymer and is present in germinating cereals, potatoes, legumes, and malt.

Solution to Question 21:

The given graph shows competitive inhibition, in which the value of Vmax remains constant and the Km value increases. The intercept on the y-axis remains unchanged while the intercept on the x-axis shifts close to zero forming a steeper slope.

Competitive inhibition is when the inhibitor binds reversibly to the substrate-binding site and competes with the substrate for the active enzyme-binding site. This effect of competitive inhibition can be overcome by increasing the substrate concentration by which maximum velocity can still be achieved. Therefore, Vmax (maximum velocity of the reaction) is unaltered in this type of inhibition. However, the concentration of the substrate required to achieve half of the maximum velocity is increased. Therefore, Km is increased in this condition.

The graph below shows the changes in Vmax and Km in competitive inhibition.



Other options:

Option B: Non-competitive inhibition is by binding inhibitors to enzymes in locations remote from the active site. Increasing the concentration of the substrate does not overcome non-competitive inhibition. Thus, noncompetitive inhibitors decrease the Vmax of the reaction. They do not interfere with the binding of the substrate to the enzyme. Thus, the enzyme shows the same Km in the presence or absence of noncompetitive inhibitors.

The graph shows the changes in Vmax and Km in non-competitive inhibition.



Option C: In uncompetitive inhibition, inhibitors bind with the enzyme-substrate intermediate complex, and inhibit it. This leads to a decrease in both Vmax and Km. An example is the inhibition of placental alkaline phosphate (ALP) by phenylalanine.

Option D: In Allosteric inhibition, inhibitors bind to a site other than the active site i.e. allosteric site, and induce a conformational change (change in the shape of the active site unlike in non-competitive inhibition) thus, inhibiting the enzyme. ATP is an allosteric inhibitor of

Phosphofructokinase in glycolysis.

Pharmacology AIIMS 2017

Question 1:

How much drug will remain at the end of 6 hours when a patient is given 200 mg drug, 75 mg excreted in 90 min and if the drug follows first-order kinetics?

a) 50 mg

b) 30.5 mg

c) 12.5 mg

d) 6.25 mg

Question 2:

What is the most bactericidal anti leprotic drug amongst the given drugs?

a) Amoxicillin

b) Ciprofloxacin

c) Ofloxacin

d) Erythromycin

Question 3:

Which of the following is not true about P-glycoprotein?

a) Pgp stands for permeability glycoprotein

b) Also known as CD243

c) Also known as ABCB1

d) Also known as MDR2

Question 4:

For which of the following drugs, is absorption increased after a fatty meal?

a) Griseofulvin

b) Amphotericin B

c) Ampicillin

d) Aspirin

Question 5:

A patient with Prinzmetal angina is administered isosorbide mononitrate and is noticed to have symptomatic relief. What is the mechanism of action of nitrate in this case?

a) Endothelium independent coronary vasodilation

b) Reduced cardiac contractility

c) Increased left ventricular end diastolic volume

d) Decreased diastolic perfusion pressure

Question 6:

The drug of choice in anaphylactic shock is?

a) Adrenaline 0.5 mg 1/1000 IM

b) Adrenaline 1 mg 1/10000 IV

c) Atropine 3 mg

d) Adenosine 6 mg followed by 12 mg IV

Question 7:

What instruction do you give to a mother who is lactating for drug use?

a) No advice as most of drug are not present in breast milk

b) Give longer half-life drugs

c) Tell the mother to feed when it is least efficacious

d) To feed just before the next dose, when the plasma concentration of the drug would be the least

Question 8:

In a case of acute exacerbation of asthma, salbutamol was given, however, no improvement was noticed. Intravenous corticosteroids and aminophylline were given and the condition improved. What is the probable mechanism by which the corticosteroids would have improved the condition?

a) Corticosteroids increase bronchial responsiveness to salbutamol

b) Corticosteroids cause direct bronchodilatation when used with xanthines

c) Corticosteroids indirectly increase the effect of xanthines on adenosine receptor

d) Corticosteroids increase mucociliary clearance

Question 9:

Which of the following opioids is a partial agonist at mu and a full agonist at kappa receptors?

a) Pentazocine

b) Buprenorphine

c) Tramadol

d) Fentanyl

Question 10:

The drug of choice for insomnia in blind people is

a) Latanoprostene

b) Zerviate

c) Tasimelteon

d) Ramelteon

Question 11:

Dofetilide belongs to which class of anti-arrhythmic drugs?

a) Class I

b) Class II

c) Class III

d) Class IV

Question 12:

What is the mechanism of action of vancomycin?

a) Cell wall synthesis inhibition

b) Cell membrane inhibition

c) Peptide synthesis inhibition

d) 50s ribosome inhibition

Question 13:

The following table gives BP values before and after administration of drug 'X'. What would be the probable mechanism of action of drug 'X'?

		DRUG X
SBP	120	150
DBP	80	68
HR	72	86

©MARROW

a) α1 agonist and β1 agonist

b) M2 agonist and M3 agonist

c) α1 antagonist and β2 agonist

d) β1 agonist and β2 agonist

Question 14:

A patient requires Ceftriaxone 180 mg. You have a 2 ml syringe with 10 divisions per ml. The vial contains 500 mg/5ml of ceftriaxone. How many divisions in the 2 ml syringe will you fill to give 180 mg ceftriaxone?

a) 18

b) 1.8

c) 20

d) 2

Question 15:

The drug that decreases the size of prostate is:

a) Tamsulosin

b) Sildenafil

c) Finasteride

d) Flutamide

Question 16:

Which one of the following is a placebo?

a) Herbal medication with no effect known

b) Physiotherapy

c) Sham surgery

d) Cognitive behavioural therapy

Question 17:

A young male developed 5 episodes of loose stools 2 hours after eating in a restaurant. He is afebrile and has mild dehydration. What should be the treatment?

a) Ciprofloxacin and tinidazole

b) Only ciprofloxacin

c) Only ORS

d) Ciprofloxacin, tinidazole and ORS

Question 18:

Atropine is indicated in all of the following poisonings except _____.

a) Baygon

b) TIK 20

c) Parathion

d) Endrin

Question 19:

27-year-old female came for treatment of infertility to OPD. Bromocriptine was prescribed. What could be the possible reason?

a) Hyperprolactinemia

b) PCOD

c) Hypogonadotrophic hypogonadism

d) PID

Question 20:

Which is a mood stabiliser with an anti-suicidal role?

a) Carbamazepine

b) Lamotrigine

c) Lithium

d) Valproate

Question 21:

Which of the following is not an idiosyncratic reaction of carbamazepine?

a) Blurred vision

b) Agranulocytosis

c) Rash

d) Steven Johnson Syndrome

Question 22:

In a middle-aged diabetic patient with chronic kidney disease stage 5, which of the following drugs can be given safely without altering the dose?

a) Vildagliptin

b) Linagliptin

c) Saxagliptin

d) Sitagliptin

Question 23:

Which of the following drugs can be used as an antidote for fibrinolytic therapy?

a) Protamine

b) Streptokinase

c) Heparin

d) ε-Aminocaproic acid

Question 24:

A known epileptic patient on levetiracetam has been seizure-free for 2 years. But he develops agitation and anger issues interfering with his day-to-day activities as a result of the drug intake. Which of the following measures is to be taken next?

a) Slowly taper the drug over next 6 months

b) Continue levetriacetam since a 5 year seizure free interval is needed

c) Stop levetiracetam and start on a different anti-epileptic

d) Discontinue the drug as he is seizure free

Question 25:

A drug was given IV at a rate of 1.6mg/min. The clearance of the drug is 640ml/min. The half life of the drug is 1.8 hour. Calculate the plasma concentration of the drug at steady state.

a) 2.88mg/ml

b) 0.004mg/ml

c) 3.55mg/ml

d) 0.002mg/ml

Question 26:

Which of the following disease-modifying anti rheumatoid drug increases extracellular adenosine?

a) Methotrexate

b) Lefunamide

c) Sulphasalazine

d) Hydroxychloroquine

Question 27:

Composition of Lente insulin is:

a) 70% amorphous 30% crystalline

b) 70% crystalline 30% amorphous

c) Only 70% amorphous

d) Other name of NPH insulin

Question 28:

Mechanism of action of Oseltamivir as an antiviral agent is:

a) Inhibition of M2 receptor

b) Neuraminidase inhibition

c) Inhibition of RNA dependent DNA polymerase

d) Apoptosis of infected cells

Question 29:

A male patient who is suffering from a psychotic disorder is on antipsychotic, antidepressant and antihypertensive treatment. He suddenly developed tachycardia and chest tightness. On evaluation he had metabolic acidosis. ECG showed widened QRS complex and prolonged QT interval. Which of the following drugs is responsible for this?

a) Amitriptyline

b) Amlodipine

c) Chlorthalidone

d) Enalapril

Question 30:

You and your friend plan for a mountain trip for the day after. What medication would you advise him to help with his motion sickness?

a) Ranitidine tablet before start of trip

b) Dimenhydrinate before start of trip

c) Scopolamine patch the night before trip

d) Omeprazole the night before and before start of trip

Question 31:

A patient was treated for septicemia with norepinephrine infusion. Which of the following best explains the benefit of this drug?

a) Bradycardia

b) Tachycardia

c) Increases urine output

d) Increases mean arterial blood pressure

Question 32:

Which of the following suggests storage of drugs in tissue?

a) Excretion in saliva

b) Excretion in urine

c) Large volume of distribution

d) Small volume of distribution

Answer Key

Question No.	Correct Option
1	b
2	c
3	d
4	a
5	a

6	a
7	d
8	a
9	a
10	c
11	c
12	a
13	d
14	a
15	c
16	c
17	c
18	d
19	a
20	c
21	a
22	b
23	d
24	a
25	d
26	a
27	b
28	b
29	a
30	c
31	d
32	c

Detailed Explanations

Solution to Question 1:

In first order kinetics, a constant fraction of the drug is excreted in a given time.

Since 75 mg of 200 mg is excreted in 90 minutes, the fraction of drug excreted in each 90 mins is 75/200, which is 3/8. Thus, the fraction of the drug that remains in the body is 5/8.

There will be 4 cycles of excretion of the drug (6 hours = 360 mins, and 360/90 = 4). Thus, with a starting dose of 200mg, the drug remaining in the body after 4 hours will be:

200 x 5/8 x 5/8 x 5/8 x 5/8 = 30.5mg

Alternate way to calculate:

75 mg of 200 mg is excreted in 90 minutes, thus 75/200 = 0.375 is eliminated

After 90 minutes; 200 mg-75 mg = 125 mg remains.

The second 90 minutes (after 180 minutes) = 125 x 0.375 = 46.875 mg is eliminated, and 125-46.875 = 78.125 mg remains

The third 90 minutes (after 270 minutes) = 78.125 x 0.375 = 29.297 mg is eliminated, and 78.125-29.297 = 48.828 remains

The fourth 90 minutes (after 360 minutes) = 48.828 x 0.375 = 18.310 mg is eliminated, 48.828-18.310 = 30.518 mg remains after 6 hours.

Solution to Question 2:

Ofloxacin is the most bactericidal anti leprotic drug amongst the given drugs.

It exhibits high activity against M. leprae, and it is being used in alternative multidrug therapy (MDT) regimens in rifampicin-resistant cases. As a component of MDT, it has been found to hasten the bacteriological and clinical response.

Among the macrolides, clarithromycin has activity against leprosy bacilli and is used for alternative MDT in cases resistant to both rifampicin and fluoroquinolones. Azithromycin shows activity against Mycobacterium avium complex but not lepra bacilli.

Note: Rifampicin is the most potent bactericidal anti leprotic drug.

Solution to Question 3:

P-glycoprotein is also known as multidrug resistance protein 1 or MDR1 (not MDR2). It is also denoted by cluster of differentiation 243 (CD243).

P-glycoprotein (P-gp) is one of the first members of the ATP-binding cassette (ABC) transporter, which acts as an efflux membrane transporter. It is responsible for limiting cellular uptake and the distribution of xenobiotics and toxic substances. It is associated with multidrug resistance (MDR).

• Increased P-gp expression: Reduces the absorption of drugs that are substrates for P-gp

• Decreased P-gp expression: Leads to supratherapeutic plasma concentrations and drug toxicity

Solution to Question 4:

The absorption of griseofulvin is increased after a fatty meal.

Note: Fatty meals increase the rate of absorption of efavirenz to such an extent as to cause toxicity. Hence, efavirenz is recommended to be taken on an empty stomach.

Drug absorption reduced /delayed by food	Drug absorption increased by food
AmpicillinAspirinAtenololAzithromycinCaptoprilIndinavirIsoniazidRifampinSucralfateTetracyclineDoxycycline	AtovaquoneCarbamazepineChlorothiazideCefuroximeDiazepamGanciclovirLithiumMetoprololRitonavirHydralazineLumefantrine

Solution to Question 5:

The mechanism of action of nitrates in prinzmetal angina is endothelium-independent coronary vasodilatation.

Prinzmetal angina occurs due to vasospasm of the coronary arteries. Nitrates relax the smooth muscles of epicardial coronary arteries and relieve coronary artery spasm.

In the case of stable angina, the primary mechanism of action of nitrates is preferential dilatation of veins (by increasing cGMP). Venous dilatation leads to a decrease in the venous return to the heart (preload). This decreases the ventricular end-diastolic volume and thereby a decrease in myocardial oxygen consumption. The diastolic perfusion pressure (aortic blood pressure in diastole - left ventricular pressure in diastole) is increased.

Solution to Question 6:

Adrenaline 0.5 mg 1/1000 by intramuscular route (IM) is the drug of choice for treatment of anaphylactic shock. Hence it is the answer here.

In case the patient develops life threatening condition, adrenaline 0.25 mg IV is given.

Note : This is a controversial question. The following inference from journal may help to clarify the concept.

• As vasodilatation is the main pathological change early in anaphylaxis, this enables the subcutaneous or intramuscular absorption of adrenaline to be rapid and effective.

• Thus, when the disease is treated early and is progressing slowly, or venous access is difficult or the patient is unmonitored, intramuscular adrenaline has advantages in terms of safety and is usually effective.

• Later, when intravascular volume is depleted and shock occurs with a life threatening situation, or there is severe dyspnoea or airway compromise, the intravenous route is necessary to achieve optimal absorption.

Solution to Question 7:

Ideally, the mother should feed just before the next dose, when the plasma concentration of the drug would be the least.

• The drugs are present in breast milk in variable amount but lesser than in plasma of mother.

• Breastfeeding is advised at least 3-4 hours after drug intake so that plasma concentration is lesser and hence the concentration in the breast milk is also less.

Solution to Question 8:

Corticosteroids improved the condition by increasing the airway responsiveness to salbutamol.

Mechanism of action of corticosteroids in an acute asthma attack:

• Increases responsiveness to β-2 agonists

• Decreases mucous secretion

• Decreases inflammatory cell response.

Steroids increase the transcription of β-2 receptor genes and hence increases responsiveness to β-2 agonists.

β-2 agonists increase the effect of steroids by enhancing the action of glucocorticoid receptors.

Theophylline increases the effects of steroids by activating the enzyme histone deacetylase which is the target for steroids as well.

Note: As per recent GINA 2021 guidelines, aminophyllines are no longer preferred in the management of acute severe asthma due to their low efficacy and poor safety profile.

Solution to Question 9:

Pentazocine is a partial agonist at mu and a full agonist at kappa receptors.

Analgesia produced by pentazocine is predominantly mediated by the κ receptor.

Pentazocine is a mixed agonist-antagonist opioid agent. It is a full agonist at the κ receptor and a partial agonist or weak antagonist at the μ receptor

Note: Buprenorphine is a partial agonist at mu receptors but an antagonist at kappa receptors

Solution to Question 10:

The drug of choice for insomnia in blind people is tasimelteon.

Option A - Latanoprostene is used in primary open-angle glaucoma.

Option B - Zerviate is a cetirizine ophthalmic solution used in allergic conjunctivitis.

Option D - Ramelteon is a synthetic tricyclic analog of melatonin used in sleep onset difficulty insomnia.

Solution to Question 11:

Dofetilide belongs to class III of anti-arrhythmic drugs.

Solution to Question 12:

The mechanism of action of vancomycin is cell wall synthesis inhibition.

The bacterial cell wall has long chains of peptidoglycan and is cross-linked to each other by the action of the enzyme transglycosylase. Vancomycin is a glycopeptide group of antibiotics that binds to the D-alanyl-D-alanine terminus of the synthesized peptidoglycan. This inhibits the transglycosylase enzyme and prevents elongation and cross-linking of peptidoglycan. This leads to the weakening of the cell wall causing lysis.

It exhibits a time-dependant or concentration-independent killing of bacteria. It primarily acts on Gram +ve organisms which are resistant to other antibiotics such as methicillin-resistant staphylococcus and ampicillin resistant enterococci.

Lipoglycopeptides are a new generation of glycopeptide congeners which include telavancin, dalbavancin, and oritavancin. Dalbavancin and Oritovancin are administered as a single intravenous dose for skin and soft-tissue infections.



Solution to Question 13:

The probable mechanism of action of drug 'X' is β1 and β2 agonism.

According to the given table, systolic BP and heart rate is increased due to β1 agonistic effect, and diastolic BP is decreased by vasodilatation due to β2 action.

Option A: Action on α1 causes ↑ DBP

Option B: Action on M2 causes ↓ HR and action on M3 causes vasodilatation and reflex increase in HR. Hence there is no net effect on heart rate

Option C: α1 antagonism causes vasodilatation and ↓ DBP. β2 action causes vasodilatation and DBP ↓

Solution to Question 14:

• 500 mg of drug is in 5 ml and hence 100 mg in 1ml. Thus 180 mg in 1.8 ml

• 10 divisions for each ml

• 1 ml = 10 divisions

• Therefore, 1.8 ml = 10*1.8 = 18 Divisions

Solution to Question 15:

Finasteride reduces prostatic size.

Finasteride:

• It is a competitive inhibitor of 5α-reductase enzyme, thus inhibiting the conversion of testosterone into its active form DHT(Dihydrotestosterone) which is responsible for androgenic action in Prostate gland.

• It produces effect on long term use (nearly 6 months) by reducing prostatic epithelial cell proliferation thereby reducing the size of prostate and improves urine flow.

Tamsulosin is an α 1 Selective Antagonist which dilates prostatic urethra and hence produces immediate symptom relief in BPH. It has no effect on reducing size of the prostate.

Sildenafil acts by inhibiting phosphodiesterase-5 (PDE-5) and is used in the treatment of erectile dysfunction.

Flutamide is a Nonsteroidal Androgen Receptor antagonist with no hormonal activity. They are used for treatment of metastatic prostatic cancer in conjunction with GnRH agonists.

Solution to Question 16:

Sham surgery (placebo surgery) is a faked surgical intervention that omits the step thought to be therapeutically necessary.

Note: While option A may be considered a placebo due to there being no known effects, it is not certain as to whether all possible effects (known and unknown) have been considered. Hence,

weighing options A and C, the better option among the two would be C as sham surgery is a widely used placebo.

Solution to Question 17:

It is a case of acute diarrhea in an afebrile, stable patient which can be managed with ORS alone.

Management of diarrhea:

• Mild-moderate dehydration - ORS.

• Severe dehydration- IV fluids are given.

• Afebrile patient - No antibiotics are given.

• Febrile patient - empirical antibiotics. If only bacterial etiology is suspected, ciprofloxacin is given. If giardiasis is suspected, metronidazole or tinidazole are recommended.

Solution to Question 18:

Endrin is an organochloride pesticide that causes neurotoxicity. There is no antidote available and only symptomatic treatment is done.

Parathion and TIK20 are organophosphates and baygon is a combination of organophosphate and carbamate. For all of these, atropine is the drug of choice.

Solution to Question 19:

The patient is likely to be suffering from hyperprolactinemia.

Prolactin inhibits the secretion of GnRH, which leads to the decreased secretion of LH and FSH resulting in infertility. Bromocriptine acts on Dopamine D2 receptors to reduce prolactin secretion.

Solution to Question 20:

Lithium is a mood stabilizer with an antisuicidal role.

Maintenance treatment with lithium has shown to decrease the rate of suicide in mood disorders. Lithium was found to be better than other mood stabilizers such as carbamazepine, lamotrigine, and valproate.

Lithium is also used in recurrent depression with a cyclic pattern is controlled by either lithium or imipramine. It is also used to augment standard antidepressant response in acute major depression in those patients who have had an inadequate response to monotherapy.

Solution to Question 21:

Blurred vision is not an idiosyncratic reaction of carbamazepine.

Idiosyncratic blood dyscrasias with carbamazepine, include fatal cases of aplastic anaemia and agranulocytosis. The most common idiosyncratic reaction is an erythematous skin rash. It can rarely cause Steven Johnson Syndrome.

Solution to Question 22:

Linagliptin can be given safely without altering the dose for this patient.

Linagliptin is a dipeptidyl-peptidase-4 (DPP-4) inhibitor and is eliminated by the hepatobiliary route. Hence no dosage adjustment is needed in renal failure.

It is used in type 2 diabetes as monotherapy or as combination therapy with other drugs when metformin is not tolerated or contraindicated.

Other DPP-4 inhibitors like sitagliptin, vildagliptin, saxagliptin, alogliptin show renal elimination.

Solution to Question 23:

ε-Aminocaproic acid can be used as an antidote for fibrinolytic therapy.

It is a lysine analog that binds competitively to plasminogen, blocking plasminogen from binding to fibrin and the subsequent conversion to plasmin. This activity subsequently results in the inhibition of fibrinolysis.

It should not be given unless hemorrhage is life-threatening because it inhibits intrinsic fibrinolytic activity and can precipitate DIC. Fresh frozen plasma and cryoprecipitate may also be used to replenish fibrin and clotting factors.



Solution to Question 24:

The mood changes observed in the above patient must be due to an adverse effect of levetiracetam. Abrupt stoppage of the drug can result in rebound seizures. So the best management will be to slowly taper the drug over the next 6 months.

Adverse effects of levetiracetam:

- Somnolence
- Asthenia
- Ataxia
- Dizziness.
- Mood and behavioral changes
- Psychotic reactions (rare)

Solution to Question 25:

Plasma steady-state concentration is achieved when the amount of drug being administered equals the amount of drug being excreted from the body.

The amount of drug being administered = 1.6mg/min

Amount of drug being excreted = Clearance (CL) x Plasma concentration of the drug (C)

Clearance = 640ml/min

Therefore,

1.6mg/min = 640ml /min (clearance) x Plasma concentration of the drug

Hence,

Plasma concentration of the drug = 1.6 mg/min / 640 ml/min = 0.002 mg/ml.

Solution to Question 26:

Methotrexate is the disease-modifying anti rheumatoid drug that increases extracellular adenosine. Methotrexate's principal mechanism of action relates to inhibition of amino-imidazole carboxamide ribonucleotide (AICAR) transformylase and thymidylate synthetase.

AICAR accumulates intracellularly, competitively inhibits AMP deaminase, leading to an accumulation of AMP. The AMP is released and converted extracellularly to adenosine, which is a potent inhibitor of inflammation. As a result, the inflammatory functions of neutrophils, macrophages, dendritic cells, and lymphocytes are suppressed.



Solution to Question 27:

Lente insulin is an intermediate-acting insulin and is a combination of Ultralente and Semilente in the ratio of 7:3

Two types of insulin zinc suspensions have been produced. The one with large particles is crystalline and practically insoluble in water (Ultralente). It is long-acting. The other has smaller particles and is amorphous (Semilente), is short-acting. Their 7:3 ratio mixture is called 'Lente insulin' and is intermediate-acting.

Solution to Question 28:

Oseltamivir and zanamivir are neuraminidase inhibitors.

They competitively and reversibly interact with the enzyme sites to inhibit viral neuraminidase activity. This interferes with the release of the progeny from infected host cells, halting the spread of infection. Early administration is crucial.

Oseltamivir is given at a dose of 75 mg, twice a day for 5 days for the treatment of Influenza A and B. For prophylaxis, it is given at a dose of 75mg, once per day and continued for the duration of exposure.

Solution to Question 29:

The given clinical scenario is suggestive of torsades de pointes. Only amitriptyline causes this condition, among the above drugs.

Torsades de pointes (French: twisting of points) is a life-threatening form of polymorphic ventricular tachycardia with rapid asynchronous complexes and an undulating baseline on ECG.



Solution to Question 30:

In the above case, a scopolamine patch the night before the trip should be advised.

Treatment for motion sickness :

• Scopolamine patch- preferred, most effective.

• Do not give to children <10 years of age.

• Place patch behind the ears at least 4 hours before travel and replace every 72 hours as needed.

• Transdermal patch – Most effective

• Oral dose - 0.4–0.6 mg one hour before travel and then TID may be recommended.

• Cinnarizine – 30mg, 2 hours prior to travel, and 15mg TID as required. Not as effective as scopolamine. Moderately effective.

• Promethazine

• Other drugs (minimally effective):

• Cyclizine

• Dimenhydrinate

• Diphenhydramine

• Meclizine

PPI's / ranitidine / traditional ginger – all are ineffective.

Solution to Question 31:

The benefit of using noradrenaline in septicemia is that it increases the mean arterial blood pressure. It acts as a powerful peripheral vasoconstrictor and inotrope.

Noradrenaline causes a rise in both systolic BP and diastolic BP, resulting in a rise in the mean arterial blood pressure. This increases the organ perfusion in patients in septic shock or cardiogenic shock.

Another advantage of noradrenaline is that it does so without causing tachycardia, which is often a troublesome event with other inotropes in critical care.

Options A and B: The rise in mean arterial pressure causes reflex bradycardia which is not a major contributor to the treatment of septic shock.

Option C: Low-dose dopamine infusions and furosemide are used to increase urine output in critically ill patients.

Solution to Question 32:

Storage of drug in tissue suggests large volume distribution.

Drugs with small volume of distribution are confined to vascular component.

Drugs with large volume of distribution are in compartments like adipose tissue, muscle, bone.

Volume of distribution depends on:

• Lipid solubility

• pKa value of a drug

• Degree of plasma protein binding

• presence of tissue-specific transporters

• differences in regional blood flow.

Volume (V) = Dose administered I.V. / Plasma concentration

e.g., when 1000mg of drug is injected I.V. and plasma concentration of drug is 50mg/L,

- V=1000/50
- V=20L

Pharmacology AIIMS 2018

Question 1:

Which of the following is not a side effect of salbutamol?

a) Hypoglycemia

b) Hypokalemia

c) Tremors

d) Tachycardia

Question 2:

Which of the following drugs should be sold only on prescription from a registered medical practitioner?

a) Schedule M drugs

b) Schedule H drugs

c) Schedule J drugs

d) Schedule X drugs

Question 3:

The loading dose of a drug is governed primarily by:

a) Clearance

b) Volume of distribution

c) Plasma half life

d) Bioavailability

Question 4:

Therapeutic index is used for assessing which of the following factors?

a) Safety

b) Efficacy

c) Potency

d) Toxicity

Question 5:

Which of the following drugs causes flagellate pigmentation?

a) Bleomycin

b) Doxorubicin

c) Vincristine

d) Minocycline

Question 6:

Which is the anticoagulant of choice for the management of cancer-associated thromboembolism?

a) Unfractionated Heparin

b) Warfarin

c) Low molecular weight heparin

d) Dabigatran

Question 7:

Which of the following new drugs has been approved for the treatment of amyotrophic lateral sclerosis?

a) Edaravone

b) Piracetam

c) Dabrafenib

d) Trametinib

Question 8:

What is the drug of choice for scrub typhus during pregnancy and otherwise respectively?

a) Chloramphenicol, doxycycline

b) Doxycycline, doxycycline

c) Doxycycline, azithromycin

d) Azithromycin, doxycycline

Question 9:

Which of the following is an example of a GPCR?

a) Muscarinic

b) Nicotinic

c) GABA-A

d) Insulin

Question 10:

The package description of a drug states that 500mg of paracetamol is present in each unit of the tablet formulation, but in reality, each unit contains only 200mg. What is the terminology used to refer to these kinds of drugs?

a) Misbranded drug

b) Adulterated drugs

c) Spurious drugs

d) Counterfeit drugs

Question 11:

A patient presented to the casualty with signs of dehydration due to severe vomiting. The intern gave the patient a parenteral medication, following which the patient developed grimacing and abnormal posturing. What could have been the inciting drug for this reaction?

a) Atropine

b) Ranitidine

c) Metoclopramide

d) Domperidone

Question 12:

Identify the wrongly matched pair of antifungals with its mechanism of action:

a) Ketoconazole : Lanosterol-14-demethylase inhibition

b) Flucytosine : Inhibits microtubule synthesis and thus hinders mitosis

c) Terbinafine : Squalene epoxidase inhibition

d) Amphotericin B : Sequesters ergosterol

Question 13:

Which of the following parameters is used to assess the extent of drug absorption in a plasma concentration vs. time graph?

a) Tmax

b) Cmax

c) T1/2

d) Area under the curve

Question 14:

Which of the following routes of administration will not bypass hepatic first-pass metabolism?

a) Sublingual

b) Oral

c) Intravenous

d) Transdermal

Question 15:

For the action of heparin as an anticoagulant, availability of which of the following chemical substances is mandatory?

a) Thrombin

b) Plasmin

c) Antithrombin-III

d) Thrombomodulin

Question 16:

What is the bioavailability of a drug administered intravenously?

a) 10%

b) 100%

c) 50%

d) Variable

Question 17:

A drug solution containing 500 mg/5 ml of the drug is given to you. You are provided with a 2 ml syringe (having 10 divisions for each ml). How many divisions of the 2 ml syringe need to be administered to deliver 180 mg of the drug to a patient?

a) 1.8

b) 10

c) 14

d) 18

Question 18:

Which of the following is gametocidal to all the species causing malaria?

a) Primaquine

b) Chloroquine

c) Quinine

d) None

Question 19:

A drug label suggests storage in a 'cool' place. What is the temperature at which this drug is to be stored?

a) - 2 degree Celsius

b) 0 degree Celsius

c) 2-6 degree Celsius

d) 8-15 degree Celsius

Question 20:

A patient with AIDS was treated with zidovudine, lamivudine, indinavir, ketoconazole and cotrimoxazole. He developed nephrolithiasis, central obesity, hyperlipidemia and insulin resistance. Which of the drugs could have been the cause?

a) Ketoconazole

b) Indinavir

c) Lamivudine

d) Zidovudine

Question 21:

If carbamazepine is added to ongoing valproate therapy, what adverse effect is risked?

a) Pancreatitis

b) Thrombocytopenia

c) Hepatotoxicity

d) Hyperammonemia

Question 22:

What is the mechanism of action of dapsone, considering that it is used in the treatment of leprosy, fungal infections and, also for immunomodulatory actions?

a) Cell wall synthesis inhibitor

b) Folic acid synthesis inhibitor

c) Protein synthesis inhibitor

d) Cell membrane inhibitor

Question 23:

What is the mechanism of action of sodium nitroprusside?

a) Increased cAMP

b) Stimulation of guanylyl cyclase

c) Calcium channel blockage

d) K+ channel opener

Question 24:

Levothyroxine/liothyronine acts by binding to thyroid receptors. Thyroid hormone receptors belong to which of the following class of receptors?

a) Transmembrane receptor linked to tyrosine kinase

b) Nuclear receptor

c) Ion channel receptor

d) G protein coupled receptor

Question 25:

The Nobel Prize in Physiology or Medicine, 2018 was awarded to James P. Allison and Tasuku Honjo for which of the following discoveries?

a) Discovery of mechanisms of autophagy

b) Discoveries of molecular mechanisms controlling the circadian rhythm

c) Discovery of cancer therapy by inhibition of negative immune regulation

d) Discovery that mature cells can be reprogrammed to become pluripotent

Question 26:

Which of the following drugs does not require dose reduction in patients with renal failure?

a) Gentamicin

b) Vancomycin

c) Amphotericin B

d) Doxycycline

Question 27:

Which of the following drugs does not increase the risk of bleeding in patients undergoing warfarin therapy?

a) Cimetidine

b) Carbamazepine

c) Isoniazid

d) Amiodarone

Question 28:

Which of the following is used in acute mountain sickness?

a) Dexamathasone

b) Digoxin

c) Diltiazem

d) Acetazolamide

Question 29:

Which of the following anti-tubercular drugs is associated with ophthalmic toxicity?

a) Ethambutol

b) Isoniazid

c) Kanamycin

d) Rifampin

Question 30:

A patient with diabetic macular edema developed glaucoma. Which of the following drugs would you consider giving last?

a) Beta blocker

b) Alpha agonist

c) Pilocarpine

d) Prostaglandin analogues

Question 31:

What is the mechanism by which acetylcholine causes a decrease in the heart rate?

a) Delayed diastolic depolarization

b) Increase in plateau

c) Decrease in preload

d) Increase in afterload

Question 32:

Diamox is given to a patient with glaucoma. It is a non-competitive inhibitor of the carbonic anhydrase enzyme. Which of the following would be the effect of this drug?

a) Decrease in Vmax

b) Decrease in Km

c) Decrease in both Km and Vmax

d) No change in Vmax

Question 33:

Low apparent volume of distribution of drug indicates

a) Low t 1/2

b) Low bioavailability

c) Low efficacy

d) Not extensively distributed to tissues

Question 34:

An infant had a serum bilirubin level of 33 mg/dL. The mother might have taken which of the following drugs during the third trimester for this to occur?

a) Ampicillin

b) Co-trimoxazole

c) Azithromycin

d) Chloramphenicol

Question 35:

Lithium usage in pregnancy can lead to which of the following teratogenic effects?

a) Cardiac defects

b) Facial defects

c) CNS defects

d) Urogenital defects

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	a
5	a
6	c
7	a
8	d
9	a
10	a
11	c
12	b
13	d
14	b
15	c
16	b
17	d
18	a
19	d
20	b
21	c
22	b
23	b
24	b
25	c
26	d
27	b
28	d

29	a
30	d
31	a
32	a
33	d
34	b
35	a

Detailed Explanations

Solution to Question 1:

Hypoglycemia is not a side effect of salbutamol. It causes hyperglycemia.

The side effects of salbutamol are due to the stimulation of extrapulmonary β2 receptors. They are more common after oral or intravenous administration. The side effects seen are:

• Muscle tremor

• Tachycardia

• Hypokalemia

• Restlessness

• Hypoxemia

• Increase in free fatty acid, insulin, glucose, pyruvate, and lactate

Solution to Question 2:

Schedule H drugs should be sold only if the prescription is from a registered medical practitioner (RMP).

Important schedules under the “drugs and cosmetics act”:

• C and C1: Biological and special products

• F and F1: Bacterial vaccines

• G: Drugs that must be labelled with the word “caution” should not be taken without medical supervision

• J: List of diseases and ailments which a drug may not claim to prevent or cure

• M: Good Manufacturing Practices (GMP)

• P: Expiry period of drug formulations

• W: Drugs marketed under generic name only

• X: Psychotropic drugs requiring a special license for manufacture and sale

• Y: Requirements and guidelines on clinical trials, import, and manufacture of new drugs.

Solution to Question 3:

The volume of distribution of the drug determines the loading dose of the drug.

Loading Dose = Vol. of distribution x Desired steady state plasma concentration.

When the time to reach a steady-state is long, as it is for drugs with long half-lives, it may be desirable to administer a loading dose that promptly raises the concentration of drug in plasma to the target concentration.

Option A: Clearance is used to determine the maintenance dose.

Maintenance dose = Clearance x Desired Steady-state plasma concentration.

Options C and D: Plasma half-life primarily determines the frequency of dosing, while bioavailability determines the route of administration. Though bioavailability may affect the dose necessary, it will not be the primary factor considered during the calculation of the loading dose.

Solution to Question 4:

Therapeutic index is used to assess the safety of the drug administered.

Therapeutic index = TD50/ED50

• TD50 (Median toxic dose) = Dose required to produce a toxic effect in 50% of animals tested.

• ED50 (Median effect dose) = Dose required to produce a therapeutic effect in 50% of the individuals.

Option B: Efficacy is defined as the maximum effect produced by the drug. It expresses the effect produced by the receptors when bound by a certain drug.

Option C: Potency refers to the concentration (EC50) or dose (ED50) required to produce 50% of maximum effect where EC50 is the concentration of agonist resulting in a half-maximal response.

Option D: Median toxic dose is the dose required to produce a toxic effect in 50% of animals tested. Lethal dose is the dose required to kill 50% of animals tested.

Solution to Question 5:

Flagellate hyperpigmentation/flagellate erythema is a characteristic cutaneous reaction to treatment with bleomycin.

Clinical features include multiple linear streaks (erythematous or hyperpigmented) at sites of trauma and generalized pruritus.

Solution to Question 6:

The anticoagulant of choice for the management of cancer-associated thromboembolism is low-molecular-weight heparin (LMWH).

It is the preferred agent because of more bioavailability, lack of need for anticoagulant monitoring, and a long t1/2.

Venous thromboembolism (including deep vein thrombosis, pulmonary embolism, and splanchnic vein thrombosis), is a frequent complication of cancer and is the second-leading cause of death in cancer patients receiving chemotherapy.

Based on the results from the Comparison of Low-molecular-weight heparin vs. Oral anticoagulants for long-term anticoagulation in cancer patients with venous Thromboembolism (CLOT) Trial, all recent guidelines now advise LMWH for cancer-associated thromboembolism.

However, unfractionated heparin (UFH) can be used in patients with renal failure, and fondaparinux can be considered in patients with heparin-induced thrombocytopenia (HIT).

Solution to Question 7:

Edaravone was approved by the FDA for the treatment of amyotrophic lateral sclerosis (ALS).

It has free radical scavenging properties that may reduce oxidative stress. The exact mechanism of action is unknown. It is to be administered as an IV infusion with an initial treatment cycle of daily dosing for 14 days, followed by a 14-day drug-free period. Subsequent treatment cycles consist of dosing on 10 of 14 days, followed by a 14 day drug-free period.

The major adverse effect of edaravone is hypersensitivity because the infusion contains sodium bisulfite. Other adverse effects include bruising, gait disorder, and headache.

Tafinlar (dabrafenib) and Mekinist (trametinib), are both indicated as monotherapy in BRAF V600E/K (BRAF V600E mutation-positive) melanoma. They are approved for combined administration, for the treatment of anaplastic thyroid cancer (ATC) that cannot be removed by surgery or that has spread to other parts of the body (metastatic).

Piracetam is a nootropic agent (a drug that enhances cognitive function) that is used off-label to enhance memory. It is also used to treat myoclonus.

Solution to Question 8:

The drug of choice for a case of scrub typhus during pregnancy is azithromycin. For other patients the drug of choice is doxycycline.

An alternative treatment option in pregnant women and patients allergic to doxycycline is chloramphenicol.

Solution to Question 9:

Muscarinic receptors are G-protein-coupled receptors among the given options.

M1, M2, M3, M4, and M5 are the muscarinic Ach receptors.

• M1 and M3 act through Gq protein and activate IP3/DAG causing increased cytosolic calcium.

• M2 and M4 act through Gi protein and inhibit adenylyl cyclase causing increased potassium conductance.



Nicotinic receptors and GABA-A are examples of ion channel coupled receptors.



Insulin receptor is a tyrosine kinase receptor.



Solution to Question 10:

A drug stating that 500mg of paracetamol is present in each tablet, but contains only 200mg is a misbranded drug, according to the CDSCO (Central drugs standard control organization).

The drug and cosmetics act initially passed in 1940 and amended in 2016 provides a definition of misbranded, spurious, and adulterated drugs.

A drug is considered misbranded if it fulfills any of the following three conditions:

• If the appearance of the drug has been altered to conceal physical damage or make it appear of greater therapeutic value than it actually is (or)

• If it is not labeled in the prescribed manner (or)

• If the label or container of the drug makes a false or misleading claim about the drug. In this case, the packet states that there is 500 mg of paracetamol present as opposed to the 200 mg that is actually present.

A drug is considered spurious if it fulfills the following conditions:

• If the drug is imported under a name that belongs to another drug (or)

• If the drug is an imitation of another drug with the intent to deceive (or)

• If the label depicts a manufacturer that is either fictitious or a real company with no ties to the product (or)

• If it has been substituted wholly or in part by another drug or substance.

A drug is considered adulterated if it fulfills the following conditions:

• If it consists wholly or in part of any filthy, putrid, poisonous, or decomposed substance (or)

• If it has been manufactured under unsanitary conditions (or)

• If it bears a coloring other than the one prescribed (or)

• If it has been mixed with any substance that may reduce its quality or strength.

*Note: The term counterfeit drugs is a broad term and is used by the WHO. This may also be conceived to include misbranded, spurious, or adulterated drugs. However, the term misbranded drugs would be a more appropriate answer here since it is likely that the reference used is from CDSCO.

Solution to Question 11:

The given clinical scenario describes the adverse effects characteristic of metoclopramide. The most common adverse effects of metoclopramide involve the central nervous system because it crosses the blood-brain barrier (BBB) and blocks the central D2 receptors.

The adverse effects include:

• Restlessness

• Drowsiness

• Insomnia

• Anxiety

• Agitation

• Extrapyramidal effects - acute reaction with high doses and long-term therapy. (due to central dopamine receptor blockade)

• Dystonias

• Akathisia

• Parkinsonian features

• Tardive dyskinesia

Adverse events are more common among the elderly. Long-term use should be avoided unless absolutely necessary.

Solution to Question 12:

Flucytosine acts by inhibition of fungal DNA synthesis.

It is a pyrimidine antimetabolite. After uptake into fungal cells, fungal cytosine deaminase activates it to 5-fluorouracil, which is an anticancer drug. This is further converted into metabolites that inhibit thymidylate synthase which is necessary for DNA synthesis. Thus fungal DNA synthesis is inhibited. Mammalian cells are spared as they lack cytosine deaminase enzyme.

Flucytosine has clinically useful activity against (3C):

- Cryptococcus
- Candida species
- Agents causing chromoblastomycosis

Solution to Question 13:

The area under the curve of plasma concentration vs. time curve gives the extent of absorption, which is calculated by the trapezoidal rule.

The following image shows a plasma concentration vs. time graph:



Two drugs may have the same rate of absorption but a different extent of absorption or bioavailability.

The factors upon which the extent of drug absorption depends are:

• Lipophilicity - If too lipophilic (e.g., acyclovir), the drug is not soluble enough to cross the water layer adjacent to the cell.

• Hydrophilicity - If too hydrophilic (e.g., atenolol), the drug cannot cross the lipid cell membrane.

Solution to Question 14:

Hepatic first-pass metabolism occurs in drugs administered via the oral route.

When drugs are given orally, they are absorbed in the intestine and reach the liver through the portal vein. During this process, the drug is acted on by enzymes of the liver and intestine. Additionally, the liver may excrete part of the drug in the bile, further reducing the bioavailability. This is called the first-pass metabolism.

In all the other routes, the drug directly enters systemic circulation without reaching the liver.

Note: Bioavailability is the fraction of an administered dose of unchanged drug that reaches the systemic circulation. Hence, the bioavailability of a drug administered intravenously (I.V) is 100%. It is neither subjected to the first-pass metabolism as in the oral route nor does it depend on local factors as in subcutaneous, intramuscular, or transdermal.

Solution to Question 15:

Antithrombin-III is mandatory for the action of heparin as an anticoagulant. Reduced antithrombin levels can lead to resistance.

Heparin resistance refers to failure to achieve a therapeutic aPTT with daily doses of heparin of 35,000 units or more.

In true heparin resistance, there is an actual blunting of heparin's anticoagulant effect. This occurs due to:

• Low ATIII levels (most common cause)

• Increased heparin-binding plasma proteins (acute phase reactants)

• Increased heparin clearance (e.g. due to splenomegaly in liver disease)

Low ATIII levels could be due to decreased synthesis (hereditary, hepatic failure), increased consumption (DIC, MAHA, or in a bypass circuit) or increased loss (nephrotic syndrome).

The dose of heparin has to be increased to achieve therapeutic aPTT. Severe antithrombin deficiency may require antithrombin concentrate as well.



Solution to Question 16:

The bioavailability of a drug administered intravenously (I.V) is 100%.

It is neither subjected to the first-pass metabolism as in oral route nor does it depend on local factors as in subcutaneous, intramuscular, or transdermal. In IV route, the drug is directly delivered to the systemic circulation and hence has the maximum availability for action at the intended site.

Bioavailability is the fraction of an administered dose of unchanged drug that reaches the systemic circulation. Hence, by definition, when a medication is administered intravenously, its bioavailability is 100%.

Intrathecal administration does not have a bioavailability of 100%.

Solution to Question 17:

Concentration of drug solution given = 500/5 ml or 100 mg/ml

-> 100 mg of drug in 1ml

i.e 180 mg of drug in "X" ml

-> X = 180/100 = 1.8 ml

So, 1.8 ml of the drug with 180 mg of the drug is to be given to the patient.

In the 2 ml syringe provided, each ml has 10 divisions i.e. each division corresponds to 0.1 ml.

So, for delivering 1.8 ml, 1.8/0.1= 18 divisions need to be administered.

Solution to Question 18:

Primaquine is gametocidal to all species causing malaria. It inhibits transmission of malaria.

The drugs with clinically important gametocidal activity (against all species) are:

• High activity:

• Artemisinins

• Primaquine

• Low to moderate activity:

• Chloroquine

• Quinine

• Atovaquone/proguanil

• No activity:

• Mefloquine

• Pyrimethamine

• Sulfadoxine

• Tetracycline



Solution to Question 19:

Storing at a cool place means to store at 8-15 degrees C.

Examples for drugs to be stored under cool temperature include insulin, nicorandil.

Label instructions	Temperature
Store frozen	-20oC
Keep cool	8-15oC
Store at room temperature	15-25oC

Solution to Question 20:

Indinavir, a protease inhibitor causes the above given adverse effects.

Patients being treated with protease inhibitors (e.g., indinavir) develop a syndrome involving the derangement of lipid metabolism. This causes changes in body fat distribution. Type 2 diabetes mellitus has also been reported. This is called HIV/HAART-associated lipodystrophy.

It is typically seen after 6–12 months of protease inhibitor therapy.

Solution to Question 21:

If carbamazepine is added to ongoing valproate therapy, it can result in hepatotoxicity.

A combination of valproate and carbamazepine increases hepatotoxicity. Valproic acid may inhibit carbamazepine clearance and increase steady-state carbamazepine blood levels.

Solution to Question 22:

The mechanism of action of dapsone is inhibition of folic acid synthesis.

Dapsone [4,4■diaminodiphenylsulphone (DDS)] is a structural analog of para-aminobenzoic acid (PABA) and a competitive inhibitor of dihydropteroate synthase in the folate pathway. It is weakly bactericidal and can cause mild hemolysis, and rarely anemia or psychosis.

DDS syndrome is a rare side effect of dapsone therapy in leprosy patients which starts after 5-6 weeks of starting dapsone and leads to exfoliative dermatitis, hepatosplenomegaly, lymphadenopathy, fever and hepatitis and may be fatal. It is also called 'five-week dermatitis'.

Solution to Question 23:

Sodium nitroprusside acts by stimulation of guanylyl cyclase.

Sodium nitroprusside releases nitric oxide which in turn stimulates guanyl cyclase. This activates cGMP secondary messenger which results in the relaxation of the smooth muscle.

Solution to Question 24:

Thyroid hormone receptors belong to the class of nuclear receptors.

Nuclear receptors are a family of ligand-regulated transcription factors that are activated by various lipid-soluble signals, including retinoic acid, thyroid hormone, and steroid hormones. They primarily function as transcription factors and directly interact with the genomic DNA to regulate gene expression.

Nuclear receptors can either remain anchored in the cytoplasm via chaperone proteins until freed by ligand-binding (e.g., steroid and vitamin D receptors) or can reside in the nucleus bound to specific DNA response elements even in the absence of ligand (e.g., thyroid hormone, PPARγ, and retinoic acid receptors).

Solution to Question 25:

The Nobel prize in Physiology or Medicine, 2018 was awarded to James P. Allison and Tasuku Honjo for the discovery of cancer therapy by inhibition of negative immune regulation.

Immune checkpoint inhibitors inhibit proteins that keep immune responses to cancer cells in check. When these proteins are blocked, the brake on the immune system is released and T cells are able to kill cancer cells more efficiently.

The Nobel prize 2018 in Physiology or Medicine was awarded to James P. Allison and Tasuku Honjo for their breakthrough work on unleashing the body's immune system to attack cancer, that has led to an entirely new class of these drugs and brought lasting remissions to many patients who had run out of options.

Solution to Question 26:

Doxycycline does not require dose reduction in renal failure. Doxycycline, minocycline, and glycylcyclines like tigecycline have fewer renal side effects.

Antibiotics causing acute kidney injury are as follows:

• Acute tubular necrosis:

• Amphotericin B

• Vancomycin

• Acyclovir

• Foscarnet

• Tenofovir

• Acute interstitial nephritis:

• Penicillin

- Cephalosporins
- Quinolones
- Rifampicin

Solution to Question 27:

Carbamazepine does not increase the risk of bleeding in patients undergoing warfarin therapy.

Warfarin is mainly metabolized by the CYP2C9 enzyme. Drugs that inhibit CYP2C9 enzyme reduce the metabolism of warfarin and thus, enhance the anticoagulant effect of warfarin and increase the risk of bleeding.

Drugs causing CYP2C9 inhibition:

- Amiodarone
- Azole antifungals
- Cimetidine
- Clopidogrel
- Cotrimoxazole
- Disulfiram
- Isoniazid
- Metronidazole

Solution to Question 28:

Acetazolamide is used in the treatment of acute mountain sickness for symptomatic relief. It can also be used for prophylaxis.

Solution to Question 29:

Ethambutol is associated with ophthalmic toxicity.

Optic neuritis resulting in decreased visual acuity and loss of ability to differentiate red from green is the most serious adverse effect of ethambutol. It is a dose-dependent adverse effect that can be unilateral or bilateral.

Tests of visual acuity and red-green discrimination before the start of therapy and periodically thereafter are recommended.

Solution to Question 30:

Prostaglandin analogs are the last drugs to be considered in this patient since macular edema is one of their side effects.

Adverse effects of PGF 2alpha analogs are:

• Blurring of vision

• Macular edema (especially in aphakic patients)

• Eyelash hypertrichosis

• Iris pigmentation (Heterochromia iridis)

• Peri-orbital fat depletion

Solution to Question 31:

The mechanism by which acetylcholine causes a decrease in the heart rate is by decreasing the diastolic depolarization.

The acetylcholine released from the vagal nerve acts on the M2 receptor (βγ subunit of G protein) on the SA node. This opens a special set of potassium channels causing an efflux of potassium (IKAch) which hyperpolarizes the SA node (Ih). The resultant current (Ih) slows the diastolic depolarization thereby causing bradycardia.

Solution to Question 32:

Non-competitive inhibitors cause a decrease in Vmax.

In non-competitive inhibition:

• The inhibitor binds to the adjacent site and not the catalytic site of the enzyme

• The catalytic property of the enzyme is reduced and hence Vmax is decreased

• The affinity of the enzyme is not altered and hence Km is unaltered



Solution to Question 33:

Low apparent volume of distribution of a drug indicates that it is not extensively distributed in the tissues.

Apparent Volume of distribution (Vd) = dose administered I.V/ plasma concentration E.g: If a 500 mg of drug A is given i.v (100% bioavailability) and the plasma concentration reached is 10 mg/ liter then the apparent volume of distribution (Vd) is 500/10= 50 liters.

But the normal blood volume is about 5 liters. A high Vd is due to accumulation of drug A in extravascular tissues resulting in low plasma concentration. Therefore, the apparent volume of distribution is an imaginary volume throughout which the drug is distributed if the plasma concentration remains the same throughout.

Option A: t1/2 relationship with volume of distribution (Vd) and clearance (Cl) is given by the formula t 1/2 = 0.693 x Vd /Cl. t1/2 is directly proportional to Vd but it is also inversely proportional to clearance. If a drug has a low volume of distribution but clearance is very low, t1/2 can still be high. So low Vd doesn't always imply low t1/2. So the best answer among the given options is D.

Option B and C: Low Vd does not indicate low bioavailability or low efficacy.

Solution to Question 34:

Cotrimoxazole is a combination drug consisting of trimethoprim and sulfamethoxazole, of which sulfamethoxazole is a sulfonamide. Sulphonamides displace bilirubin from plasma albumin, leading to elevation of serum bilirubin levels.

These drugs can cross the placenta and are secreted in milk and can therefore cause kernicterus in newborn infants if they are given to pregnant women near term or during lactation.

Solution to Question 35:

Lithium use in pregnancy can lead to cardiac defects.

Ebstein's anomaly is the specific cardiac defect with lithium use in pregnancy.

In Ebstein's anomaly, the radiological sign seen is a box-shaped heart.



Pharmacology AIIMS 2019 (May & Nov)

Question 1:

Which of the following is not an adverse effect of salbutamol?

a) Tachycardia

b) Tolerance

c) Hypokalemia

d) Hypoglycemia

Question 2:

Propranolol is drug of choice for:

a) Ulcerated infantile hemangioma

b) Lymphangioma circumscriptum

c) Capillary malformation

d) Pyogenic granuloma

Question 3:

CIostridium difficile associated diarrhea occurs maximally due to:

a) Lincosamides

b) Tetracyclines

c) Carbapenems

d) Macrolide

Question 4:

A 53 year old man is found to have stage 2 hypertension without any co morbidities. Which drug will you start?

a) Furosemide

b) Chlorthalidone

c) Spironolactone

d) Triamterene

Question 5:

Which of the following is a schedule X drug?

a) Halothane

b) Thalidomide

c) Ketamine

d) Colistin

Question 6:

LMWH is given by which route?

a) Intravenous

b) Subcutaneous

c) Intramuscular

d) Intradermal

Question 7:

A patient presents to the emergency department with diarrhea, sweating, excessive lacrimation, and salivation. His pulse rate is 55/min and his plasma cholinesterase level was 50U/mL. Which of the following can be used to treat this patient?

a) Neostigmine

b) Atropine

c) Epinephrine

d) Benztropine

Question 8:

What is pegloticase used for?

a) Chronic gout

b) Rheumatoid arthritis

c) Psoriatric arthritis

d) Paralytic ileus

Question 9:

Which of the following GPCRs does not act via the K+ channel?

a) Angiotensin 1 receptor

b) Muscarinic M2 receptor

c) Dopamine D2 receptor

d) Serotonin 5 HT1 receptor

Question 10:

Which of the following is not used in heart failure?

a) Trimetazidine

b) Sacubitril

c) Metoprolol

d) Nesiritide

Question 11:

Which of the following causes an increase in ACh release from the presynaptic neuron?

a) Potassium channel blocker

b) Sodium channel blocker

c) Calcium channel blocker

d) Chlorine channel blocker

Question 12:

Apart from mu-opioid receptors, on which other receptor does tramadol act?

a) Anticholinergic

b) 5HT1a inhibition

c) Serotonergic and norepinephrine reuptake inhibition

d) Antihistaminic

Question 13:

In which of the following conditions is non-pulsatile dosing of GnRH agonist not used?

a) Male infertility

b) Endometriosis

c) Precocious puberty

d) Prostate cancer

Question 14:

Which of the following cannot be used to decrease intraocular pressure in glaucoma?

a) Dexamethasone

b) Acetazolamide

c) Apraclonidine

d) Mannitol

Question 15:

Prucalopride is a _____.

a) 5HT2a agonist

b) 5HT2b agonist

c) 5HT4 antagonist

d) 5HT4 agonist

Question 16:

Z track technique is used for?

a) Administration of nicotine patches

b) Monitoring of carbamazepine therapy

c) Administration of long acting depot antipsychotics

d) Monitoring of lithium therapy

Question 17:

Tafenoquine is used in which of the following conditions?

a) Radical cure of plasmodium vivax

b) Prophylaxis of malaria in pregnancy

c) Treatment of severe falciparum malaria

d) Treatment of malaria in children <2 years

Question 18:

Which is not given in Wilson's disease?

a) Zinc

b) Penicillamine

c) Calcium citrate

d) Trientine

Question 19:

What dose of epinephrine is used in anaphylactic shock?

a) 1:10000 I.V. - 10 ml

b) 1:1000 I.M. - 0.5 ml

c) 1:1000 I.M. - 1 ml

d) 1:1000 I.V. - 0.5 ml

Question 20:

Which of the following is true about tedizolid?

a) Active against gram positive organisms

b) Poor oral bioavailability

c) Active against anaerobes

d) Causes peripheral neuropathy

Answer Key

Question No.	Correct Option
1	d
2	a
3	a
4	b
5	c
6	b
7	b
8	a
9	a
10	a
11	a
12	c
13	a
14	a
15	d
16	c
17	a
18	c
19	b
20	a

Detailed Explanations

Solution to Question 1:

Salbutamol in large doses can result in hyperglycemia, not hypoglycemia.

Option A: Tachycardia due to reflex sympathetic stimulation due to vasodilator action on β2 receptors and direct effect on cardiac β1 receptors (β2 selectivity is lost at high concentrations)

Option B: Tolerance occurs due to desensitization on long-term use

Option C: Hypokalemia due to direct effect on skeletal muscle uptake of K+ ions

Solution to Question 2:

Propranolol is the drug of choice in the management of infantile hemangioma complicated by ulceration or in symptomatic patients.

It acts by causing vasoconstriction, inducing apoptosis and inhibition of angiogenesis.

Solution to Question 3:

Clostridium difficile-associated diarrhea is mostly due to lincosamides such as clindamycin

Antibiotic-associated colitis is caused by Clostridium difficile infection due to the destruction of the normal gut microbial flora by the use of antibiotics. The antibiotics most frequently implicated include clindamycin, fluoroquinolones and broad-spectrum penicillins, and cephalosporins

Clostridium difficile infection occurs frequently in hospitals and nursing homes where antimicrobial use is high and the environment is contaminated by C. difficile spores.

Although all antibiotics can cause the condition, the most commonly implicated are:

• Clindamycin

• Third-generation cephalosporins (cefotaxime, ceftriaxone, cefuroxime, and ceftazidime)

• Fluoroquinolones (ciprofloxacin, levofloxacin, and moxifloxacin)

• Broad-spectrum penicillins

Solution to Question 4:

Chlorthalidone is one of the first choice drugs in the treatment of hypertension without any comorbidities.

First-choice drugs used in the treatment of hypertension in patients without any comorbidities include:

• Thiazide diuretics: Chlorthalidone is usually preferred to hydrochlorothiazide.

• Calcium channel blockers: Extended-release, long-acting dihydropyridines are preferred.

• ACE inhibitors: Captopril is preferred for the initiation of therapy. Enalapril and ramipril twice daily may also be used.

In the presence of any associated conditions or comorbidities, the first choice drugs for the treatment of hypertension would be as follows:

Conditions associated with hypertension	First choice drugs for hypertension
DiabetesChronic kidney disease with proteinuria	Inhibitors of renin-angiotensin system.
Angina	Beta-blockers or calcium channel blockers
Heart failure	ACE inhibitors and alpha+beta blockers such as carvedilol
Benign prostatic hyperplasia	α1Blockers
Pregnancy	Labetalol
Atrial fibrillation	Diltiazem and verapamil

Solution to Question 5:

Ketamine is a schedule X drug.

According to The Drugs and Cosmetics Act-1945,

Schedule X drugs are those that have high abuse liability and are to be sold with a valid prescription only.

A prescription in duplicate, separate license requirement, meticulous storage, and dispensing records must be maintained by the chemist.

Option B: Thalidomide is a schedule H drug.

Option D: Colistin is a schedule P drug.

Solution to Question 6:

LMWH is given by the subcutaneous route.

Heparin, low molecular weight heparin (LMWH), and fondaparinux are not absorbed through GI mucosa and must be given parenterally.

LMWH and fondaparinux are given subcutaneously.

Heparin can be given through intravenous infusion and subcutaneous routes.

Solution to Question 7:

The clinical features are suggestive of organophosphate poisoning and IV atropine is the treatment of choice.

The organophosphates commonly used as pesticides are anticholinesterases. Toxicity occurs by excessive inhalation of vapors, ingestion, or percutaneous absorption. It inhibits the

cholinesterase activity which results in an increase of acetylcholine, leading to cholinergic symptoms.

The predominant initial signs are of muscarinic excess which includes: (DUMBBELSS)

• Diarrhea

• Urination

• Miosis

• Bronchospasm

• Bradycardia

• Emesis

• Lacrimation

• Salivation

• Sweating

This may be followed by cognitive disturbances, convulsions, and neuromuscular blockade.

Treatment:

• Maintenance of vital signs particularly respiration

• Removal of clothing and washing of the skin

• IV atropine 2- 5 mg. If IV administration is not possible, then 2 mg IM every 5-10 minutes (till muscarinic symptoms disappear)

• Pralidoxime, a cholinesterase reactivator can also be used for treatment

Solution to Question 8:

Pegloticase is used in the treatment of chronic gout.

Pegloticase is a pegylated uricase (urate oxidase). It catalyzes the enzymatic oxidation of uric acid into allantoin, which is a more soluble and inactive metabolite.

It is used in the treatment of severe, refractory chronic gout or in cases where other urate-lowering therapies are contraindicated.

Solution to Question 9:

Angiotensin 1 receptor acts via Ca2+ channels. It does not act through second messengers via the K+ channel.

G-protein coupled receptors activate inwardly rectifying potassium channels and increase the permeability of the potassium ions causing hyperpolarisation of the cell. They are called G protein-coupled inwardly-rectifying potassium channels (GIRKs). These receptors include:

• M2 muscarinic

- A1 adenosine
- α2 adrenergic
- D2 dopamine
- 5HT1 serotonin
- Somatostatin receptors.

Solution to Question 10:

Trimetazidine is not used in heart failure.

Trimetazidine is a newer antianginal drug that is a pFOX inhibitor which partially inhibits the fatty acid oxidation pathway in the myocardium. This decreases the oxygen requirement of the heart.

Nesiritide (natriuretic peptide), sacubitril (neprilysin inhibitor) and metoprolol (beta-blocker) are used in the treatment of heart failure.

Solution to Question 11:

Potassium channel blockers cause an increase in ACh release from the presynaptic neuron.

Presynaptic potassium channel block promotes the activation of voltage-gated calcium channels. This leads to a release of acetylcholine in the neuromuscular junction.

Aminopyridines (K+ channel blockers) are used in the treatment of Lambert-Eaton myasthenic syndrome which helps to improve muscle strength.

Solution to Question 12:

Tramadol inhibits the reuptake of serotonin and norepinephrine, apart from its action on the mu-opioid receptor.

It is a synthetic codeine analog that is a mu-opioid receptor agonist.

Tramadol is a centrally acting analgesic used in the treatment of mild to moderate pain.

Solution to Question 13:

Pulsatile dose of GnRH agonist is used in the treatment of male infertility.

Pulsatile administration of GnRH agonists results in increased FSH and LH secretion and is used to treat male and female infertility.

Non-pulsatile or continuous dose of GnRH agonist results in suppression and decreased concentrations of gonadotropins LH and FSH. They are used in:

• Endometriosis

• Uterine fibroids

• Central precocious puberty

• Ca prostate

• Ca ovary and breast

• Controlled ovary stimulation

Solution to Question 14:

Dexamethasone is a long-acting glucocorticoid and is not used to decrease the intraocular pressure in glaucoma.

Acetazolamide, apraclonidine, and mannitol are used in lowering intraocular pressure in glaucoma.

Solution to Question 15:

Prucalopride is a specific 5HT4 receptor agonist which increases bowel movements.

It is approved for use in women with chronic constipation where other laxatives have failed to provide relief.

Solution to Question 16:

The Z-track technique is used for the intramuscular administration of long-acting depot antipsychotics.

Advantages:

• The 'Z-track' prevents leakage of the medication into the subcutaneous tissue and skin staining through the tract to the surface of the skin

• Pain intensity is also reduced



Solution to Question 17:

Tafenoquine is used in the radical cure of Plasmodium vivax. Total eradication of plasmodium species which includes clinical curative and hypnozoites phase are called radical cure.

Primaquine and tafenoquine (derivative of primaquine) have very high activity against the latent tissue forms (hypnozoites) of the species. It is primarily used for the chemoprophylaxis and radical cure of P. vivax and P. ovale infections.

Solution to Question 18:

Calcium citrate is not used in the treatment of Wilson's disease.

The drugs used in the management of Wilson's disease are:

Chelating agents: These drugs bind to copper to form complexes that get excreted in the urine. D-penicillamine and trientine are first-line drugs in symptomatic patients with hepatic diseases.

• D-penicillamine: It is the recommended drug due to its availability and cost-effectiveness as compared to trientine. Since it is an antagonist to pyridoxine, supplementation of pyridoxine is required. It can precipitate various adverse effects including aplastic anemia, myasthenia gravis, and membranous nephropathy. It can also worsen neurological symptoms

• Trientine: It is administered in those who cannot tolerate D-penicillamine

• Tetrathiomolybdate: It has been identified to be effective in patients with neuropsychiatric manifestation

Elemental zinc:

• Zinc blocks the intestinal absorption of copper by inducing metallothionein (endogenous chelator in enterocytes)

• It is commonly used as a maintenance therapy along with chelating agents. It can also be used as first-line therapy in pregnancy

Solution to Question 19:

Epinephrine (1:1000 solution) 0.5 ml via intramuscular route is preferred in the treatment of anaphylactic shock.

For intravenous use, 1:1000 adrenaline must be further diluted to 1:10,000 and should be used very cautiously as slow bolus.

There is a much greater risk of causing harmful side effects by either inappropriate dosage or misdiagnosis of anaphylaxis when using IV adrenaline. Hence, IM injection is preferred over intravenous (IV) bolus because it is faster in many situations and safer (lower risk of cardiovascular complications, such as severe hypertension and ventricular arrhythmias).

For intramuscular, subcutaneous, and inhalational (nebulization) routes of administration, 1:1000 dilution can be used.

In cardiac arrest, the 1:1000 adrenaline must be diluted to 1:10,000 for IV use. The recommended dose of epinephrine is 1mg = 10 mL of 1:10,000 solution IV administered every 3 to 5 minutes during resuscitation as per the CPR guidelines.

Epinephrine is also used in the treatment of hypotension, to induce local vasoconstriction to reduce the diffusion of local anaesthetics and to induce bronchodilation in asthma and COPD.

Solution to Question 20:

Tedizolid is active against gram-positive organisms.

Tedizolid belongs to the class of oxazolidinones. It has poor activity against most gram-negative bacteria and anaerobes. It inhibits protein synthesis by binding to the P site of the 50S ribosomal subunit.

It is used for the treatment of skin and soft tissue infections, respiratory tract infections caused by streptococci, and S. aureus including MRSA.

Common adverse effects include myelosuppression leading to anemia, leukopenia, thrombocytopenia, and pancytopenia.

Pharmacology AIIMS 2020 (May and Nov INI-CET)

Question 1:

All of the following are the uses of enteric-coated drugs except:

a) Protect the acid labile drug from gastric acidity

b) Protect the gastric mucosa from irritant effects of certain drugs

c) Prolongs the half life of the drug

d) Deliver drugs distal to stomach

Question 2:

A patient who was on antiepileptics for the management of seizure disorder started having visual disturbance. Evaluation reveals the contraction of the visual field. The drug responsible is likely to be?

a) Phenobarbitone

b) Vigabatrin

c) Ethosuximide

d) Levetiracetam

Question 3:

If a patient who is on metoprolol is started on verapamil, which of the following side effects would be seen?

a) Bradycardia with AV block

b) Torsades de pointes

c) Tachycardia

d) Ventricular fibrillation

Question 4:

The response on stimulation of muscarinic receptor is:

a) Erection

b) Ejaculation

c) Bronchodilation

d) Increased myocardial contractility

Question 5:

Rosuvastatin should not be combined with which of the following drugs?

a) Oxycodon

b) Clarithromycin

c) Adalimumab

d) Rivoraxaban

Question 6:

Which of the following is not a prodrug?

a) Proguanil

b) Sulindac

c) Primidone

d) Flurouracil

Question 7:

Which of the following topical antifungal has both anti-inflammatory effects and anti-pruritic action?

a) Ketoconazole

b) Sertaconazole

c) Luliconazole

d) Terbinafine

Question 8:

A patient is started on lithium therapy. Which of the following side effects should the patient be asked to watch out for to monitor for toxicity?

a) Malaise and sore throat

b) Vertigo and Diarrhoea

c) Fever

d) Breathlessness

Question 9:

A psychiatric patient was started on lithium for the management of bipolar disorder. To monitor the blood lithium levels, at what time interval should the blood sample be collected after the administration of lithium tablets?

a) After 24 hours

b) After 12 hours

c) After 6 hours

d) Immediately after taking tablets

Question 10:

Which of the following is a second line anti-tubercular drug?

a) Streptomycin

b) Ethionamide

c) Acyclovir

d) Vancomycin

Question 11:

Which of the following anti-tubercular drug is not bactericidal?

a) Isoniazid

b) Rifapentin

c) Thioacetazone

d) Kanamycin

Question 12:

Which of the following drugs do not need dose adjustment in renal failure?

a) Sitagliptin

b) Saxagliptin

c) Vildagliptin

d) Linagliptin

Question 13:

The antidote for fibrinolytic therapy is?

a) Urokinase

b) Dalteparin

c) Tranexamic acid

d) Alteptase

Question 14:

Which of the following is least likely to be useful in a patient with severe COVID-19?

a) Low molecular weight heparin

b) Oxygen

c) Favipiravir

d) Steroids

Question 15:

If codiene is coming to the market in combination with dextromethorphan as an anti-tussive, this combination _____

a) Should not be used due to risk of addiction

b) Can be used as both have different mechanism of action

c) Should not be used as it is irrational

d) Can be used to treat cough effectively

Question 16:

The normality of 2M HCl is ___ as 1M H_2SO_4.

a) Half

b) Twice

c) Same

d) 4 times

Question 17:

An elderly woman on treatment for migraine presents with chest pain. She has a previous history of angina/MI. Which of the following mechanism of the prescribed drug for migraine might have caused this presentation?

a) 5-HT antagonist

b) 5-HT agonist

c) Calcium Channel Blocker action

d) Nitric oxide mediated vasodilation

Question 18:

A patient presented with pelvic fracture and head trauma after a road traffic accident. He was given IV morphine for pain relief following which he developed respiratory difficulty. What could be the most likely cause?

a) Pelvic Fracture

b) CO_2 retention due to opioid use

c) Cerebral Edema

d) Pulmonary embolism

Question 19:

Which of the following antidepressants causes the least sexual dysfunction?

a) Fluoxetine

b) Venlafaxine

c) Mirtazapine

d) Imipramine

Question 20:

Which of the following statements is true about disulfiram?

a) When used with alcohol, the patients feels drowsy, with chest discomfort and respiratory depression

b) It is an anticraving drug

c) It can be initiated even in minimally motivated person

d) Its action persists up to 2-3 days after stopping the drug

Question 21:

Which of the following drugs promotes bone formation?

a) Calcitonin

b) Raloxifene

c) Bisphosphonates

d) Teriparatide

Question 22:

A known asthmatic patient with fever and sore throat was prescribed an antibiotic. Two days later he presents with a maculopapular rash. Which of the following drugs might have caused this?

a) Prednisolone

b) Salbutamol

c) Fexofenadine

d) Amoxicillin

Question 23:

What is the most common drawback of the given device?

a) Bacterial contamination

b) Does not provide multiple doses

c) Drug deposition in the oropharynx

d) Not portable

Question 24:

Which of the following statements is correct for a patient on high dose of penicillin and gentamicin?

a) TDM is done with penicillin because of narrow therapeutic index

b) TDM is done with gentamicin because of narrow therapeutic index

c) TDM is done with both because of dangerous side effects

d) Both drugs do not need monitoring

Question 25:

A boy presents with pallor and lymphadenopathy, the X-ray of his skull is given below. Which of the following drug is not used for the management of the suspected diagnosis?



a) Cyclosporine

b) Methotrexate

c) Vinblastine

d) Vincristine

Question 26:

A patient was treated with TCA for a major depressive episode in the hospital for 4 weeks. What is the main concern while considering his discharge from the hospital?

a) Need for drug monitoring

b) ECG monitoring for arrhythmias

c) Suicidal risk

d) Sedation being the side effect

Question 27:

Which of the following drugs exacerbates myasthenia and is usually avoided in a patient with myasthenia gravis? 1. Lidocaine 2. Digene 3. Meropenem 4. Phenytoin

a) 1 and 2

b) 2 and 3

c) 1 and 3

d) 1 only

Question 28:

Which of the following drugs are used for absence seizures? A. Carbamazepine B. Valproate C. Ethosuximide D. Gabapentin

a) A, B and C

b) B and C only

c) B only

d) A, B and D

Question 29:

Match the following dilution of adrenaline and its use: A. 1:1000 B. 1:10000 C. 1:100000 1. Cardiac arrest 2. Anaphylaxis 3. Local anesthesia

a) A-1 B-2 C-3

b) A-2 B-1 C-3

c) A-3 B-2 C-1

d) A-1 B-3 C-2

Question 30:

Match the following bacteria with the drug of choice used for each.

a) A-1, B-3, C-4, D-2

b) A-2, B-1, C-4, D-3

c) A-3, B-4, C-1, D-2

d) A-4, B-1, C-2, D-3

Question 31:

Which of the following drug reduces mortality in COVID-19 patients? A. Dexamethasone B. Remdesivir C. Favipravir

a) B and C

b) A and B

c) Only A

d) A, B , C

Question 32:

Which of the following statements are true?

a) A and C

b) A, B and C

c) A and D

d) B, C and D

Question 33:

Which of the following is correct about the drugs used in Gout?

a) A, B & C

b) A, B, C & D

c) A & B

d) C & D

Answer Key

Question No.	Correct Option
1	c
2	b
3	a
4	a
5	b
6	c
7	b
8	b
9	b
10	b
11	c
12	d

13	c
14	c
15	c
16	c
17	b
18	b
19	c
20	a
21	d
22	d
23	c
24	b
25	d
26	c
27	a
28	b
29	b
30	b
31	c
32	b
33	a

Detailed Explanations

Solution to Question 1:

Enteric coating of the drug does not prolong the half-life of the drug.

The half-life of a drug depends on the volume of distribution and rate of clearance from the blood. It is given by t1/2 = 0.693x Volume of distribution / Clearance.

The enteric coating can be done for the following reasons:

• Protection of active pharmaceutical ingredients, from the acidic environment of the stomach (e.g. enzymes and certain antibiotics).

• To prevent gastric distress or nausea from a drug due to irritation (e.g. sodium salicylate).

• For the delivery of drugs that are optimally absorbed in the small intestine to their primary absorption site in their most concentrated form.

• To provide a delayed-release component for repeat action.

• Required for minimizing the first-pass metabolism of the drug

Solution to Question 2:

The given patient has developed a contraction of the visual field which is a side effect of vigabatrin.

Solution to Question 3:

The patient will be prone to developing bradycardia and AV block.

Both ß-blockers and non-dihydropyridine calcium blockers have a propensity to slow down SA Node function and AV conduction. A combination of these drugs increases the risk further and is contraindicated. In contrast, propranolol has safely been used with the dihydropyridine group (nifedipine) of calcium channel blockers.

Solution to Question 4:

Erection is mediated by muscarinic receptors.

Erection results from parasympathetic action which is mediated via muscarinic/nicotinic receptors, whereas sympathetic action is mediated by adrenergic receptors.

Solution to Question 5:

Rosuvastatin should not be combined with clarithromycin.

Clarithromycin is a macrolide antibiotic and it can cause drug interactions because of its ability to inhibit organic anion-transporting polypeptides 1B1 and 1B3 (OATP1B1 and OATP1B3) which are involved in the excretion of statins like rosuvastatin and pravastatin.

Individuals taking rosuvastatin, pravastatin, or fluvastatin with clarithromycin had an increased risk of hospitalization for acute kidney injury, an increased risk of hyperkalemia and an increased risk of all cause mortality.

Solution to Question 6:

Among the given options primidone is not a prodrug.

Prodrugs are drugs that are in their pharmacologically inactive form. They are converted to their active forms by metabolism.

The prodrug may offer advantages over the active form in being more stable, having better bioavailability, or fewer side effects and toxicity. Some prodrugs are activated selectively at the

site of action. The given options are examples of prodrugs:

• Sulindac is a sulfoxide prodrug used in inflammatory diseases including osteoarthritis, rheumatoid arthritis, acute gouty arthritis, ankylosing spondylitis.

• Proguanil is a prodrug and its active metabolite is cycloguanil. It is used in the treatment and prevention of malaria.

• Fluro-uracil is a prodrug having the active metabolite as fluorouridine monophosphate

Solution to Question 7:

Sertaconazole has both anti-inflammatory effects and anti-pruritic action.

Sertaconazole inhibits the release of pro-inflammatory cytokines from activated immune cells.

Solution to Question 8:

Vertigo and diarrhoea are known side-effects of lithium therapy. Lithium therapy can cause side effects at both therapeutic and toxic levels.

Known significant side effects of lithium therapy include

• Polyuria-polydipsia

• Dry mouth

• Leukocytosis Headache

• Hand tremors

• Confusion

• Decreased memory

• Muscle weakness

• Hyperreflexia

• Muscle twitch

• Vertigo

• ECG changes

• Nausea & vomiting

• Diarrhoea

Other options:

No evidence of significant association with sore throat, fever or breathlessness has been documented with lithium therapy

Solution to Question 9:

The blood sample for serum lithium estimation is to be collected after a time interval of 10-12 hours from the previous dose of lithium.

Lithium is used in the management of bipolar disorder, mania, and has anti-suicidal properties. It has a narrow therapeutic index, and blood levels in excess of this may cause symptoms of toxicity. Therapeutic drug monitoring is done by measuring trough lithium levels 12 hours after the previous dose, at least 5 days after the initiation of the treatment with lithium.

Solution to Question 10:

Ethionamide is a second line anti-tubercular drug.

Ethionamide is structurally related to isoniazid and acts by inhibiting the mycolic acid synthesis. It is effective against bacilli resistant to other drugs and has been proved effective in infections due to atypical mycobacteria. It is effective against both intracellular as well as extracellular organisms.

The first line anti-tubercular drugs include: Isoniazid, Rifampicin, Pyrazinamide, Ethambutol

The second line drugs include fluoroquinolones, ethionamide, capreomycin, kanamycin, amikacin, cycloserine, thioacetazone.

Note: This is a recall question when ethionamide is a second-line among the given options, and streptomycin was a first line injectable drug.

Solution to Question 11:

Thioacetazone is not a bactericidal drug, it is bacteriostatic.

Thioacetazone is never used alone. It is only used as a combination with first-line TB medications, such as isoniazid and rifampicin. It is used primarily to prevent the development of bacteria that are resistant to first-line drugs and to treat patients infected with drug-resistant TB.

It should never be used in HIV patients as it can cause severe and fatal skin reactions.
Side-effects include gastrointestinal disturbances, blurring of vision, hemolytic anemia and urticaria.

Isoniazid is a first-line bactericidal drug that is active against both intracellular and extracellular bacilli.

Rifapentin is a rifampin congener having a similar mechanism of action. It is a bactericidal drug having a longer half-life than rifampin.

Kanamycin is bactericidal and is active against bacilli resistant to streptomycin, INH, and cycloserine.

Solution to Question 12:

Linagliptin does not require dose adjustment in patients with renal failure.

Linagliptin is a dipeptidyl-peptidase-4 (DPP-4) inhibitor. It is used in type 2 diabetes as monotherapy or in combination therapy when metformin is not tolerated or contraindicated. It binds extensively to plasma proteins and is cleared primarily by the hepatobiliary system. Hence, no dosage adjustment is needed in renal failure.

Other DPP-4 inhibitors (sitagliptin, vildagliptin, saxagliptin, alogliptin) show renal elimination and therefore require dose reduction in renal failure.

Solution to Question 13:

The antidote for fibrinolytic therapy is tranexamic acid.

Tranexamic acid is a lysine analogue that competitively bind to lysine binding sites on plasminogen and plasmin, and block their interactions with fibrin and thereby inhibiting fibrinolysis. It is given intravenously in trauma resuscitation, massive hemorrhage, and in women with postpartum hemorrhage.



Other options:

Option A (Urokinase) and D (Alteplase) are fibrinolytic drugs.

Option B: Dalteparin is low molecular weight heparin.

Solution to Question 14:

Favipiravir (inhibits viral RNA polymerase) was the first oral drug approved in India for restricted emergency use for the treatment of mild to moderate COVID-19 patients. It was not approved for

severe COVID-19.

Note: This is a question from INICET 2020, during which time Favipiravir had gained approval from DCGI for mild to moderate COVID. The current clinical management protocol for COVID-19 does not recommend Favipiravir.

Solution to Question 15:

It is an irrational drug combination and should not be given.

Codeine is an opioid analgesic having specific antitussive action while dextromethorphan is a centrally active NMDA receptor antagonist that may be effective as an antitussive. Both are centrally acting antitussives, and do not provide any added advantage when combined. Hence it is an irrational drug combination.

Rational fixed-dose combination drugs for the treatment of cough usually contain:

• A centrally acting anti-tussive (either codeine or dextromethorphan; not both), with

• Antihistamines, decongestants and expectorants.

Option A - Both of these drugs carry a risk of addiction, but it is not the reason to not give the drug. The main reason is that the combination is irrational.

Options B & D - Both codeine and dextromethorphan have different mechanisms of action and can treat cough effectively. However, the combination of two centrally acting antitussives does not provide any added advantage compared to using one drug alone. Hence, the combination is irrational.

Solution to Question 16:

The normality of 2M HCl is the same as 1M H_2SO_4.

Normality of a solution can be calculated from its molarity by multiplying it by its valence. Valence is the number of protons a salt releases in aqueous solution.

Here Valence of HCl is 1 while H_2SO_4 is 2.

Hence Normality of HCl = 2M x 1 = 2 N

similarly Normality of H_2SO_4 = 1M x 2 = 2N

Normality is defined as the number of gram equivalents dissolved per liter of solution. Where gram equivalents are the number of moles of a solute multiplied by its valence

Molarity is defined as the number of moles of a solute dissolved per liter of solution.

Solution to Question 17:

The patient may have been given sumatriptan, a 5-HT1b/1d receptor agonist which causes vasoconstriction and used in migraine. Thus, sumatriptan will worsen angina by causing coronary vasoconstriction.

Angina will be worsened if the O2 demand of the heart is increased or there is reduced blood flow to the already ischemic areas.

Solution to Question 18:

The most likely cause of respiratory difficulty in the given clinical scenario is CO2 retention due to opioid use, as the patient was administered morphine after head trauma.

Head injury is considered as a contraindication to morphine use because:

• Morphine retains CO2 that increases intracranial pressure (ICP) and adds to the already increased ICP.

• Therapeutic doses can result in marked respiratory depression.

• Interferes with the assessment of progress by causing miosis, vomiting, and altered mentation.

Solution to Question 19:

Among the given options mirtazapine causes the least sexual dysfunction.

Mirtazapine and mianserin block histamine H1 receptors and also have high antagonistic activity against 5HT2A, 5HT2C, and 5HT3 receptors. Sexual dysfunction is not a known side-effect of mirtazapine.

Due to its sedative effect, mirtazapine is used in depression with insomnia.

Option A: Fluoxetine is a selective serotonin reuptake inhibitor (SSRI) that can cause side effects like GI disturbances, sexual dysfunctions, may increase the risk of suicidal thoughts or behavior, risk of serotonin syndrome with MAOIs, and some CYP interactions. Vilazodone is an SSRI not associated with sexual dysfunction or weight gain.

Option B: Serotonin-norepinephrine reuptake inhibitors like venlafaxine have side effects that include nausea and dizziness, may increase the risk of suicidal thoughts or behavior, and can also cause sexual dysfunctions.

Option D: Tricyclic antidepressants like imipramine have side effects like orthostatic hypotension, weight gain, GI disturbances, sexual dysfunction, seizures, and irregular heartbeats.

The various drugs classes used in depression and anxiety are listed below

Class of drugs	Examples
Selective serotonin reuptake inhibitors (SSRI)	FluoxetineFluvoxaminParoxetineSertralineCitalopram
Serotonin-norepinephrine reuptake inhibitors	Venlafaxine, desvenlafaxineDuloxetineMilnacipran

Class of drugs	Examples
Tricyclic antidepressants	AmitriptylineImipramineNortriptylineDesipramineAmoxapine
Atypical antipsychotics	AripiprazoleOlanzapineQuetiapineRisperidone
Monoamine oxidase inhibitors (MAO)	IsocarboxazidSelegilinePhenelzine
Atypical antidperessants	BupropionTrazodoneMirtazapineMianserin

Solution to Question 20:

Disulfiram is used in behavioural modification technique for alcohol de-addiction. The principle of action is aversive conditioning.

Disulfiram is an irreversible inhibitor of both cytosolic and mitochondrial forms of aldehyde dehydrogenase. Ethanol consumption during disulfiram therapy can lead to increased concentrations of acetaldehyde. Increased acetaldehyde levels can cause flushing, pulsating headache, throbbing pain in head and neck, nausea, vomiting, rashes, respiratory difficulties, chest pain, autonomic manifestations and other aversive reactions.

Disulfiram can only be given to highly motivated individuals. Disulfiram should not be administered until the person has abstained from alcohol for at least 12 hours. The disulfiram-alcohol reaction may occur as long as 14 days after the last dose of disulfiram.

Disulfiram like reaction with alcohol ingestion is also seen with the following drugs:

• Metronidazole

• Sulfonylureas like chlorpropamide

• Cephalosporins like cefoperazone, cefamandole

Other modalities of treatment:

• Pharmacotherapy: Typically 5-10 days, treatment for detoxification: Benzodiazepine and chlordiazepoxide are the drugs of choice.

• Rehabilitation programs: Typically 28-day programs, learn about relapse prevention and identification of triggers.

• Maintenance phase: Baclofen, Acamprosate, Naltrexone, Topiramate, SSRI, Ondansetron

• Psychotherapy: Group therapy such as Alcoholics Anonymous

Solution to Question 21:

Teriparatide increases bone formation especially when given in low doses and intermittently.

Teriparatide is a recombinant preparation of parathyroid hormone.

The pharmacologic activity of teriparatide, which is similar to the physiologic activity of PTH, includes stimulating osteoblast function, increasing gastrointestinal calcium absorption, and increasing renal tubular reabsorption of calcium.

Although it may be more effective than bisphosphonates and estrogen, it has to be administered subcutaneously every day. Hence it is indicated mostly in severe osteoporosis.

Teriparatide increases the risk of osteosarcoma and it is contraindicated in conditions that cause high risk of osteosarcoma such as:

• Paget disease

• Prior h/o of radiotherapy

• Young adults with open epiphysis

• Unexplained elevated ALP

• Hypercalcemia

Option A: Calcitonin decreases bone resorption.

Option B: Raloxifene is a selective estrogen receptor modulator, which decreases bone resorption. It is used to prevent and treat osteoporosis in postmenopausal women.

Option C: Bisphosphonates like alendronate and etidronate decrease bone resorption.

Solution to Question 22:

Skin rashes of all types may be caused by allergy to penicillin.

Amoxicillin and ampicillin are aminopenicillins. The incidence of skin rashes appears to be highest following the use of ampicillin, at about 9%.

Amoxicillin is a close congener of ampicillin with the following features:

• Better oral absorption; food does not interfere with the absorption

• Incidence of diarrhea is less

• More active against penicillin-resistant pneumococci

Solution to Question 23:

The given image shows a metered-dose inhaler (MDI). The most common side effect of this device is drug deposition in the oropharynx.

Most material inhaled using this device will be swallowed and absorbed, entering systemic circulation after undergoing the first-pass effect in the liver. Some drug will also be absorbed into the systemic circulation from the lungs.

The use of a large-volume spacer will reduce the amount of drug deposited in the oropharynx, thereby reducing the amount swallowed and absorbed from the GI tract, thus limiting the

systemic effects of the drug.

The given image shows the use of spacer with MDI:



Spacers reduce the velocity and size of particles entering the upper airways. This reduces the amount of drug that impinges on the oropharynx and increases the proportion of the drug inhaled into the lower airways.

Spacer devices are useful in delivering inhaled drugs to small children, as young as 3 years of age, who are not able to use a pMDI (pressurized MDI), as seen in the image below:

Pressurized metered-dose inhalers: Drugs are propelled from a canister in the pMDI with the aid of a propellant. These devices are convenient, portable, and typically deliver 50–200 doses of the drug.

Dry powder inhalers: Drugs may also be delivered as a dry powder using devices that scatter a fine powder dispersed by air turbulence on inhalation. Children less than 7 years of age find it difficult to use a DPI. DPIs have been developed to deliver peptides and proteins, such as insulin,

systemically but have proved to be problematic because of the consistency of dosing.

The given image shows various types of dry powder inhalers:

Nebulizers: Two types of nebulizers are available. Jet nebulizers are driven by a stream of gas (air or oxygen), whereas ultrasonic nebulizers use a rapidly vibrating piezoelectric crystal and thus do not require a source of compressed gas.

The nebulized drug may be inspired during tidal breathing, and it is possible to deliver much higher doses of the drug as compared to pMDI. Nebulizers are therefore useful in treating acute exacerbations of asthma and COPD, for delivering drugs when airway obstruction is extreme (e.g., in severe COPD), for delivering inhaled drugs to infants and small children who cannot use the other inhalation devices, and for giving drugs such as antibiotics when relatively high doses must be delivered.



Using a Nebulizer

Solution to Question 24:

Gentamicin is an aminoglycoside antibiotic that requires therapeutic drug monitoring (TDM) as it has a narrow therapeutic index. It is nephrotoxic and ototoxic (predominantly vestibular effects).

The threshold for toxicity for gentamicin is the plasma concentration of 2 μg/mL, the maximum recommended for prolonged exposure.

Aminoglycoside antibiotics act by binding to the 30s ribosome and inhibiting protein synthesis. Their ability to cause misreading of DNA templates and the formation of aberrant proteins that get incorporated into bacterial cell membranes makes them bactericidal agents.

Common ototoxic drugs are:

• Antibiotics - aminoglycosides (gentamicin > tobramycin > amikacin > neomycin), erythromycin, tetracyclines.

• Chemotherapeutic agents - cisplatin, bleomycin, nitrogen mustards.

• High dose aspirin, acetaminophen, NSAID's

• Antimalarials - quinine, quinidine

• Phosphodiesterase-5 inhibitors - sildenafil

Solution to Question 25:

The given clinical scenario and the X-ray of the skull are suggestive of Langerhans Cell Histiocytosis. Vincristine is not used in the management of this condition.

Following are skeletal and radiological manifestations of Langerhan's cell histiocytosis:

Skull:

• Solitary or multiple punched-out lytic lesion without sclerotic rim.

• Double contour or beveled edge appearance (hole within a hole sign).

• Button sequestra representing residual bone.

• Geographic skull.

Mandible:

• Floating tooth: loss of lamina dura.

Spine:

• Vertebra Plana: Most common cause of vertebra plana in children; mostly seen in thoracic spine.

Skull in LCH, Vetebra plana:





Solution to Question 26:

Suicidal thoughts or behavior are a side effect of TCAs. Therefore, the attendants of the patient should be warned regarding this when a patient gets discharged after being treated with TCAs for a major depressive episode.

Tricyclic antidepressants (TCAs) are drugs that act by inhibiting the reuptake of noradrenaline and serotonin from the synaptic clefts.

The side effects of TCAs are due to blockade of other receptors:

• Muscarinic M receptors- constipation, dry mouth, arrhythmia

• Histamine H1 receptors- sedation

• 5HT receptors - weight gain due to increased appetite

• Alpha1 receptors - postural hypotension.

The various classes of drugs used in depression and anxiety are listed below:

Class of drugs	Examples
Selective serotonin reuptake inhibitors (SSRI)	FluoxetineFluvoxaminParoxetineSertralineCitalopram
Serotonin-norepinephrine reuptake inhibitors	VenlafaxineDesvenlafaxineDuloxetineMilnacipran
Tricyclic antidepressants	AmitriptylineImipramineNortriptylineDesipramineAmoxapine
Atypical antipsychotics	AripiprazoleOlanzapineQuetiapineRisperidone
Monoamine oxidase inhibitors (MAO)	IsocarboxazidSelegilinePhenelzine
Atypical antidepressants	BupropionTrazodoneMirtazapineMianserin

Solution to Question 27:

Lidocaine can exacerbate myasthenia gravis and should be avoided in patients with this disease. Also, digene is an antacid, that contains magnesium hydroxide as one of its active ingredients, which may exacerbate myasthenia.

Drugs that may aggravate myasthenia gravis are

• Antibiotics like aminoglycosides, quinolones, and macrolides

• Beta-blockers

• Botulinum toxin

• Local anaesthetics

• Magnesium

• Non–depolarising muscle relaxants

• Penicillamine

• Quinine derivatives

Aminoglycosides have to be avoided in a patient with myasthenia gravis. It blocks Ca^{2+} mediated exocytosis of acetylcholine from the presynaptic membrane in the neuromuscular junction. The decrease in ACh levels aggravates myasthenia gravis. Aminoglycosides can also have a direct inhibitory effect on post-synaptic acetylcholine receptors.



Solution to Question 28:

Ethosuximide and Valproate are effective for the treatment of absence seizures. However, between these 2 drugs, Ethosuximide would be a better first-line treatment choice for uncomplicated typical absence seizures followed by Valproate. Atypical absence seizures respond well to valproate.

Typical absence seizures are characterized by sudden, brief lapses of consciousness without loss of postural control. There is no postictal confusion. The following subtle motor signs may accompany the episodes

• Rapid blinking of the eyelids

• Chewing movements

• Small-amplitude clonic movements of the hands

Hyperventilation has a tendency to provoke the absence seizures.

NOTE: In a typical absence seizure, Ethosuximide is the first-line therapy for it and acts by blocking the calcium channel.

Mechanism of action of some common antiepileptics:

Mechanism of action	Antiepileptics
Sodium-channel blockers	PhenytoinCarbamazepine, OxcarbamazepineEslicarbazepineLamotrigineTopiramateFelbamateValproateRufinamideLevetiracetam
Calcium-channel blockers	ValproateLamotrigineEthosuximide

Mechanism of action	Antiepileptics
GABAAreceptor modulators	BenzodiazepinesPhenobarbitalFelbamatePrimidoneTopiramateCarbamazepine, OxcarbamazepineStiripentolClobazam
GABA uptake inhibitors	TiagabineVigabatrin
NMDA receptor antagonist	Felbamate
AMPA/Kainate receptor antagonist	PhenobarbitalTopiramatePerampanel
HCN channel activity enhancers	Lamotrigine
Inhibitors of brain carbonic anhydrase	AcetazolamideZonisamideTopiramate
Potassium-channel openers	RetigabineTopiramate

Solution to Question 29:

A-2 B-1 C-3

Epinephrine (1:1000 solution) 0.5 ml I.M. is preferred in the treatment of anaphylactic shock.

In cardiac arrest, the 1:1000 adrenaline must be diluted to 1:10,000 for IV use.

Adrenaline in a concentration of 1: 50,000 to 1: 200,000 is added to local anesthetics for:

• Reducing the systemic absorption

• Prolonging the duration of action

• Providing a more bloodless field for surgery

1:1000 dilution can be used for intramuscular, subcutaneous and inhalational (nebulization) route of administration. For intravenous use, 1:1000 adrenaline must be further diluted to 1:10,000 and used cautiously.

IM injection is preferred over intravenous (IV) bolus because it is faster in many situations and safer (ie, lower risk of cardiovascular complications, such as severe hypertension and ventricular arrhythmias). There is a much greater risk of causing harmful side effects by inappropriate dosage or misdiagnosis of anaphylaxis when using IV adrenaline.

Epinephrine is also used in the treatment of hypotension, to induce local vasoconstriction to reduce the diffusion of local anaesthetics, to induce bronchodilation in asthma and COPD.

Note: . The recommended dose of epinephrine is 1mg = 10 mL of a 1:10,000 solution administered IV every 3 to 5 minutes during resuscitation as per the CPR guidelines. For nebulization, 1:1000 adrenaline is commonly used.

Solution to Question 30:

A-2, B-1, C-4, D-3

Vancomycin is Drug Of Choice for Methicillin-Resistant Staphylococcus aureus (MRSA) (should be used only if other drugs are resistant).

Metronidazole is the drug of choice for anaerobic infections caused by Bacteroides.

Uses of metronidazole include:

• Giardiasis

• Trichomoniasis

• Amoebiasis

• Bacterial vaginosis

• treatment of pseudomembranous colitis.

Meropenem is used in infections caused by ESBL producing bacteria like Klebsiella.

Carbapenems are not hydrolyzed by ESBL (extended-spectrum beta-lactamases). Therefore, carbapenems like imipenem, meropenem, ertapenem are resistant to hydrolysis by extended-spectrum beta-lactamases and are indicated in the treatment of infections caused by ESBL producing bacteria. ESBL is most prevalent among gram-negative organisms like Klebsiella and E. coli.

Penicillins like amoxicillin, ampicillin can be used in H. influenzae infections.

Solution to Question 31:

Among the given options, only dexamethasone is shown to reduce mortality in severe covid19 cases.

Patients with severe COVID-19 can develop a systemic inflammatory response that can lead to lung injury and multisystem organ dysfunction. It has been proposed that the potent anti-inflammatory effects of corticosteroids might prevent or mitigate these deleterious effects and help reduce mortality.

In severe cases, Dexamethasone is administered at doses of 6mg IV/PO once daily for 10 days or until hospital discharge. If dexamethasone is not available, then, equivalent doses of other corticosteroids, like prednisone, methylprednisone, or hydrocortisone may be used.

Other options:

• There is no evidence that favipiravir reduces mortality. It reduces duration of hospitalisation in mild to moderate disease.

• In a recent update, WHO recommended against the use of remdesivir in hospitalized patients, regardless of disease severity, as there is no evidence that it improves survival.

Solution to Question 32:

Statements A, B and C are true.

A. Morphine used in pulmonary edema

B. Pethidine used in biliary colic

C. Indomethacin is used in headache

Aspirin can also cause damage to the stomach and intestinal lining leading to the development of erosions or ulcers. Erosions may cause further complications like bleeding and perforations.

Note: Unlike Morphine, Pethidine/Mepiridine can be used for pain management in biliary colic as it has no effect on the tone of sphincter of Oddi.

Solution to Question 33:

Gout is a metabolic disease associated with hyperuricemia.

The drug of choice is an acute attack of gout is NSAIDs (Non-steroidal anti-inflammatory drugs) like indomethacin or the drug colchicine, but they are poorly tolerated and should not be used in elderly patients with chronic kidney disease.

Indomethacin is commonly used in the initial treatment of gout. In addition to inhibiting prostaglandin synthase, indomethacin and other NSAIDs also inhibit urate crystal phagocytosis

Intraarticular steroids like triamcinolone or methylprednisolone can also be given.

Aspirin is not used because it causes renal retention of uric acid at low doses. It is uricosuric only at higher doses that are poorly tolerated.

Pharmacology NEET 2018

Question 1:

A colonoscopy of an elderly patient revealed brownish pigmented lesions as shown below. He admitted to chronic laxative abuse. Which of the following laxatives is most likely to cause this condition?



a) Bran

b) Psyllium

c) Senna

d) Methylcellulose

Question 2:

A patient who is on treatment for angina, presents to you with erectile dysfunction. You decide against prescribing sildenafil because the patient is being treated with _____

a) Calcium channel blockers

b) Beta blockers

c) Nitrates

d) ACE inhibitors

Question 3:

An example of a bacteriostatic drug is _____

a) Aminoglycoside

b) Vancomycin

c) Metronidazole

d) Linezolid

Question 4:

The antiviral drug having dual antiviral activity against HIV and HBV is:

a) Enfuviritide

b) Abacavir

c) Emtricitabine

d) Ritonavir

Question 5:

The replacement dose of hydrocortisone for chronic adrenal insufficiency is (in mg/day):

a) 10

b) 12

c) 20

d) 5

Question 6:

Which of the following is the drug of choice for scrub typhus?

a) Doxycycline

b) Azithromycin

c) Chloramphenicol

d) Ciprofloxacin

Question 7:

Which of the following drugs is associated with QT prolongation?

a) Lignocaine

b) Quinidine

c) Magnesium sulfate

d) All of the above

Question 8:

Sacubitril is a new drug approved for chronic heart failure. The mechanism of action of sacubitril is:

a) Angiotensin II inhibitor

b) ACE inhibitor

c) Renin inhibitor

d) Neutral endopeptidase inhibitor

Question 9:

Coagulation profile monitoring is needed in therapy with which of the following drugs?

a) Fondaparinux

b) Enoxaparin

c) Dabigatran

d) Lepirudin

Question 10:

Which of the following antihypertensives are not used in pregnancy?

a) Alpha methyldopa

b) Hydralazine

c) Propranolol

d) Labetalol

Question 11:

Which of the following atypical antipsychotics has the highest risk of galactorrhea?

a) Iloperidone

b) Risperidone

c) Clozapine

d) Aripiprazole

Question 12:

When the pH = pKa of a drug, which of the following statements is true?

a) Concentration of the drug is 50% ionic and 50 % non-ionic

b) Concentration of the drug is 90% ionic and 10% non-ionic

c) Concentration of the drug is 75% ionic and 25 % non-ionic

d) Concentration of the drug is 25% ionic and 75 % non-ionic

Question 13:

Corticosteroid indicated for the stimulation of fetal lung maturation in preterm labor is?

a) Hydrocortisone

b) Triamcinolone

c) Methylprednisolone

d) Betamethasone

Question 14:

What is the site of action of amphotericin B in fungi?

a) Nucleic acid

b) Ribosome

c) Cell membrane

d) Cytoplasm

Question 15:

Which of the following acts by causing both decreased bone resorption and increased bone formation?

a) Strontium ranelate

b) Ibandronate

c) Teriparatide

d) Calcitonin

Question 16:

Which is the drug used for refractory intractable rheumatic chorea?

a) Probenecid

b) Diazepam

c) Haloperidol

d) Sodium valproate

Question 17:

What is basiliximab?

a) IL-1 receptor antagonist

b) Anti-CD3 antibody

c) IL-2 receptor antagonist

d) TNF inhibitor

Question 18:

Which of the following conditions can be treated with pirenzepine?

a) Asthma

b) Peptic ulcer disease

c) Glaucoma

d) Hypertension

Question 19:

Which of the following antimicrobials is effective against an organism producing extended-spectrum beta-lactamases?

a) Amoxicillin - clavulanic acid

b) Penicillin

c) Piperacillin - tazobactam

d) Ceftriaxone

Question 20:

Which of the following inhibits the synthesis of glucocorticoid?

a) Mifepristone

b) Flutamide

c) Finasteride

d) Metyrapone

Question 21:

Isosorbide dinitrate is indicated for congestive heart failure in combination with which of the following vasodilators?

a) Minoxidil

b) Hydralazine

c) Nimodipine

d) Nitrendipine

Question 22:

What is the mechanism of action of colchicine?

a) Inhibits synthesis of uric acid

b) Promotes excretion of uric acid

c) Inhibits xanthine oxidase enzyme

d) Binds to tubulin and inhibits granulocyte migration

Question 23:

Which of the following statements is true about prasugrel?

a) It acts as an P2 Y12 agonist

b) It is 10 times less potent than clopidogrel

c) It is a pro drug

d) Omeprazole interferes in its activation to an active metabolite

Question 24:

Apixaban belongs to which of the following class of drugs?

a) Antifibrinolytic

b) Factor Xa inhibitor

c) Oral direct thrombin inhibitor

d) Parenteral direct thrombin inhibitor

Answer Key

Question No.	Correct Option
1	c
2	c
3	d
4	c
5	c
6	a
7	b
8	d
9	d
10	c
11	b
12	a
13	d
14	c
15	a
16	d

17	c
18	b
19	c
20	d
21	b
22	d
23	c
24	b

Detailed Explanations

Solution to Question 1:

The above colonoscopy image depicts a brownish moiré pattern that is indicative of melanosis coli. Chronic use of anthraquinone laxatives like senna and cascara sagrada can cause melanosis coli.

Bisacodyl is also implicated in causing melanosis coli.

Solution to Question 2:

The patient would be on nitrates for angina and thus should not be prescribed sildenafil as sildenafil causes dangerous potentiation of nitrate action resulting in severe hypotension.

The combination of the two drugs has a synergistic action on the intracellular cGMP levels that can result in uninhibited vasodilation and a precipitous fall in blood pressure. Hence, the use of sildenafil is a contraindication when on nitrates.

Solution to Question 3:

Linezolid is an example of a bacteriostatic drug.

Primarily bacteriostatic drugs	Primarily bactericidal drugs
SulfonamidesErythromycinTetracyclinesClindamycinChloramphenicolLinezolidEthambutol	Penicillins,CephalosporinsAminoglycosidesVancomycinPolypeptidesCiprofloxacinRifampinMetronidazoleIsoniazidCotrimoxazolePyrazinamide

Solution to Question 4:

Antiviral drug having dual antiviral activity against HIV and HBV is emtricitabine.

It is a nucleoside reverse transcriptase inhibitor similar to lamivudine. It has activity against HIV-1, HIV-2, and HBV.

Solution to Question 5:

The replacement dose of hydrocortisone for chronic adrenal insufficiency is 20 mg/day.

The treatment of chronic adrenal insufficiency (not in crisis) requires glucocorticoid replacement at a dose that replaces the physiologic daily cortisol production. This is achieved by administering an oral replacement dose of 15-25mg Hydrocortisone.

A midpoint dose of 20mg is given in 2-3 divided doses: 10mg + 5mg + 5mg.

Half the daily dose should be administered to all patients in the morning. This is to follow the normal circadian rhythm.

Solution to Question 6:

Doxycycline is the drug of choice for scrub typhus.

Azithromycin, chloramphenicol, or rifampin can be administered as alternatives in patients allergic to doxycycline and in pregnancy.

Solution to Question 7:

QT prolongation is seen with quinidine.

It causes torsades de pointes, a life-threatening form of polymorphic ventricular tachycardia.



Solution to Question 8:

Sacubitril is a neutral endopeptidase (neprilysin) inhibitor.

Sacubitril is a prodrug that is activated to sacubitrilat. It can inhibit the enzyme neprilysin (neutral endopeptidase), which is responsible for the degradation of atrial and brain natriuretic peptides (reduction of blood volume). Therefore, it increases the levels of atrial and brain natriuretic peptides and lowers blood pressure.

In 2015, the FDA approved the use of sacubitril in combination with valsartan, at a ratio of 1:1, for the treatment of chronic heart failure.

Solution to Question 9:

Coagulation profile monitoring is needed for lepirudin.

Lepirudin is a direct irreversible thrombin inhibitor and its most adverse reaction is bleeding. The risk of bleeding can be reduced by implementing an appropriate monitoring and dose adjustment strategy. Thus, coagulation profile monitoring is required for lepirudin.

However, enoxaparin, fondaparinux, and dabigatran do not bind to endothelium or many plasma proteins and thus, the response to these drugs at the usual doses is consistent from patient to patient. Hence, monitoring is not required.

Solution to Question 10:

Propranolol is a non-selective beta-blocker and is not safe for use during pregnancy.

Antihypertensives to be avoided in pregnancy:

• ACE inhibitors, ARBS

• Diuretics

• Nonselective beta-blockers

• Nitroprusside

Antihypertensives found to be safe in pregnancy:

• Hydralazine

• Methyldopa

• Dihydropyridine CCBs: should be stopped before labor as they weaken the uterine contractions

• Cardioselective beta-blockers and those with intrinsic sympathomimetic activity: may be used as a last choice

Solution to Question 11:

Among atypical antipsychotics, risperidone and paliperidone have the highest risk of galactorrhea.

The adverse effect occurs due to their higher D2 blocking potency.

Solution to Question 12:

When the pH = pKa of a drug, the concentration of the drug is 50% ionic and 50% non-ionic.

pKa is defined as the pH at which a drug exists as 50% ionized (hydrophilic) and 50% unionized (lipophilic).

For e.g., phenobarbitone is an acidic drug with a pKa of 7.2. At a pH of 7.2, 50% will be unionized. In a more basic plasma (pH=7.4), a lesser fraction (around 40%) will be unionized.

Small changes in pH near pKa of a drug may cause large variations in the degree of ionization.

Solution to Question 13:

Betamethasone is indicated for the stimulation of fetal lung maturation in preterm labor.

As compared to cortisol, betamethasone binds to maternal proteins less and has less placental metabolism. This allows increased transfer to the fetus. Additionally, infants exposed to betamethasone are less likely to require postnatal resuscitation. 2 doses of 12 mg each are administered 18-24 hours apart when the delivery is anticipated before 34 weeks of gestation.

Dexamethasone, another long-acting glucocorticoid is also indicated. It has benefits such as less risk of intraventricular hemorrhage and a shorter length of stay in the NICU.

Solution to Question 14:

The site of action of amphotericin B in fungi is the cell membrane.

Amphotericin B binds to the ergosterol in the cell membrane of fungi and forms pores or channels. They increase the permeability of the membrane to ions and cause leakage of cytosolic molecules. This leads to a loss of integrity of the cell membrane and causes fungicidal effects.

Recent evidence suggests that the amphotericin B sequesters ergosterol from lipid layers, much like a sponge, resulting in fungal cell death.

Solution to Question 15:

Strontium ranelate acts by both decreased bone resorption and increased bone formation.

It is generally reserved for severe osteoporosis. Due to an increased incidence of myocardial infarction and other cardiovascular events, the use is restricted in all the approved regions.

Option B- Ibandronate which is a bisphosphonate, decreases bone resorption only.

Option C- The pharmacologic activity of teriparatide, which is similar to the physiologic activity of PTH, includes stimulating osteoblast function, increasing gastrointestinal calcium absorption, and increasing renal tubular reabsorption of calcium.

Option D- Calcitonin also decreases bone resorption.

Solution to Question 16:

The drug used for refractory intractable rheumatic chorea is sodium valproate, as it doesn't have extrapyramidal side effects.

Sydenham's chorea is characterized by involuntary, purposeless, jerky movements which mainly affect the head and upper limbs. In the tongue, it is visible as the characteristic darting movements.

Medications used to treat Sydenham chorea are:

• Mild cases - Providing a calm environment.

• Severe cases:

• Dopamine antagonists - haloperidol, pimozide

• Antiepileptic medications - sodium valproate, carbamazepine.

• Severe, refractory cases - Steroids (thought to be immune-mediated)

• IV immunoglobulins, plasmapheresis - limited studies, decreased efficacy (not currently recommended)

Solution to Question 17:

Basiliximab exhibits a high affinity towards IL-2 receptor and inhibits it. It is an anti- CD-25 monoclonal antibody.

It has a short plasma half-life of 1 week. It is used to prevent transplant rejection reactions. Its adverse effects are anaphylactic reactions & opportunistic infections.

Solution to Question 18:

Pirenzepine is used for treating gastric ulcers. It is a selective M1 anticholinergic drug. It decreases gastric acid secretions at doses that have little effect on salivation or heart rate.

Solution to Question 19:

Piperacillin-tazobactam is an antimicrobial that is effective against organisms producing extended-spectrum beta-lactamases.

Extended-spectrum beta-lactamase (ESBL) enzyme produced by bacteria causes resistance to most of the penicillins, cephalosporins (third and fourth generation), and monobactams. ESBL is most prevalent among gram-negative organisms like Klebsiella and E. coli.

ESBL can be inhibited by combining beta-lactamase inhibitors like clavulanic acid or tazobactam. Carbapenems and quinolones have inherent activity against ESBL.

Note: As piperacillin has greater activity against Klebsiella and other gram-negative organisms than amoxicillin, piperacillin-tazobactam can be considered as a better option in this question.

Solution to Question 20:

Metyrapone inhibits the synthesis of glucocorticoids by inhibiting steroid 11β-hydroxylase reversibly.

It is used to test the hypothalamic-pituitary feedback pathway in the diagnosis of Cushing syndrome.

Solution to Question 21:

Isosorbide dinitrate (ISDN) is indicated in combination with hydralazine for CHF.

Hydralazine is a direct vasodilator and its mechanism of action is not known. It also reduces nitrate tolerance.

The combination of ISDN and hydralazine showed mortality benefits in a specific group of population (African-Americans).

Solution to Question 22:

Colchicine acts by binding to tubulin and inhibits granulocyte migration.

Colchicine produces its anti-inflammatory effects by binding to the intracellular protein tubulin. This prevents the polymerization of tubulin into microtubules which is necessary for leukocyte migration and phagocytosis. Inhibition of microtubule formation prevents granulocyte migration and also arrests cell division at G1 phase.

Solution to Question 23:

Prasugrel is a prodrug.

It acts as an irreversible ADP (P2 Y12) antagonist and is converted into its active metabolite by CYP3A4 and CYP2B6. Since omeprazole is an inhibitor of CYP2C19, it does not interfere with its activation.

It is 10-times more potent than clopidogrel and has a more rapid onset of action because of better absorption and more predictable metabolic activation.

Solution to Question 24:

Apixaban, a novel oral anticoagulant (NOAC), is a factor Xa inhibitor that is used in the treatment of VTE (venous thrombo embolism)/DVT and in its prophylaxis after major orthopedic surgeries.

Direct factor Xa inhibitors:

• Rivaroxaban

• Apixaban

• Edoxaban

• Betrixaban

Parenteral direct thrombin inhibitors:

• Bivalirudin

• Argatroban

• Desirudin

Oral direct thrombin inhibitor—Dabigatran is the only oral direct thrombin inhibitor available for clinical use.

The FDA recently approved the drug Apixaban for a new indication. The drug is now used to reduce the risk of stroke and systemic embolism in patients with nonvalvular atrial fibrillation.

Pharmacology NEET 2019

Question 1:

Calculate the therapeutic index of a drug whose dose-response curves are plotted as shown below:



a) 1

b) 2

c) 4

d) 8

Question 2:

Which of the following statements is true regarding the dose-response curve shown in the image below?



a) C is competitive antagonist

b) C is more potent than A

c) B is more potent than A

d) B is competitive antagonist

Question 3:

Which of the following antiglaucoma medications is unsafe in infants?

a) Timolol

b) Brimonidine

c) Latanoprost

d) Dorzolamide

Question 4:

Apparent volume of distribution of a drug is very high (6L/kg). Which of the following is true regarding the distribution of that drug?

a) Highly bound to plasma proteins

b) Confined to vascular compartment

c) Sequestered in body tissues

d) Both A and B

Question 5:

Which of the following phases of a clinical trial is also referred to as efficacy trial?

a) Phase 1

b) Phase 2

c) Phase 3

d) Phase 4

Question 6:

Which of the following is an oral factor Xa inhibitor?

a) Bivalirudin

b) Dabigatran

c) Rivaroxaban

d) Enoxaparin

Question 7:

What is the drug of choice for diphtheria carrier state?

a) Amoxicillin

b) Erythromycin

c) Rifampicin

d) Tetracycline

Question 8:

What is the mechanism of action of oseltamivir and zanamivir?

a) DNA polymerase inhibition

b) Protein synthesis inhibition

c) Nucleotide analogue

d) Neuraminidase inhibition

Question 9:

Which of the following is a synthetic PGF2 alpha derivative used as an abortifacient?

a) Dinoprost

b) Dinoprostone

c) Misoprostol

d) Carboprost

Question 10:

Which of the following drugs causes activation of K+-ATP channels?

a) Nicorandil

b) Molsidomine

c) Isosorbide dinitrate

d) Nitroglycerin

Question 11:

Most effective antiemetic for chemotherapy-induced vomiting is_____.

a) Domperidone

b) Tegaserod

c) Granisetron

d) Doxylamine

Question 12:

True statement about teduglutide is :

a) It is a GLP- 2 analogue

b) It's half life is 6 - 8 hours

c) It is administered orally

d) It is recommended for patients with colorectal cancer

Question 13:

Cisatracurium is preferred over atracurium because?

a) It has rapid onset of action

b) It has less depressant action on heart

c) It has short duration of action

d) It causes less release of histamine

Question 14:

Which one of the following drugs inhibits neutrophil recruitment?

a) Montelukast

b) Sodium cromolyn

c) Colchicine

d) Febuxostat

Question 15:

Which of the following fluoroquinolones has maximum bioavailability?

a) Ciprofloxacin

b) Levofloxacin

c) Gatifloxacin

d) Norfloxacin

Question 16:

What is the mechanism of triazoles?

a) Interferes with DNA and RNA synthesis

b) Inhibition of ergosterol synthesis

c) Forms pores in fungal membranes

d) Inhibits epoxidation of squalene in fungi

Question 17:

A dermatologist wants to prescribe acitretin to a patient with psoriasis. Which of the following statements is false regarding the drug?

a) Contraindicated in persons with hyperlipidemia

b) It can cause increased sensitivity to sunlight

c) Female patients should avoid getting pregnant for 6 months after therapy

d) Acitretin has a half life of about 50 hours

Question 18:

Cilastatin is administered to prolong the action of which of the following drugs?

a) Meropenem

b) Imipenem

c) Cefazolin

d) Piperacillin

Question 19:

What is the drug used to differentiate myasthenia gravis from cholinergic crisis?

a) Obidoxime

b) Edrophonium

c) Donepezil

d) Atropine

Question 20:

Which of the following is true about pegylated-filgrastim?

a) Duration of action is shorter than normal filgrastim

b) Added advantage is that it can be taken orally

c) It should not be administered in sickle cell patients

d) It is not effective in the treatment of severe neutropenia

Question 21:

The best approach in the treatment of chronic SIADH is _____

a) Tolvaptan

b) Reducing fluid intake to <500 ml per day

c) Demeclocycline

d) Fludrocortisone

Question 22:

Intravenous mannitol is used in ____?

a) Pulmonary edema

b) Congestive cardiac failure

c) Acute kidney injury with anuria

d) Glaucoma

Question 23:

Which of the following carbapenems has increased risk of causing seizures?

a) Meropenem

b) Imipenem

c) Etrapenem

d) Doripenem

Question 24:

Which of the following helps in maintaining the patency of ductus arteriosus?

a) PGF2α

b) PGE1

c) PGH2

d) PGI2

Answer Key

Question No.	Correct Option
1	c

2	d
3	b
4	c
5	b
6	c
7	b
8	d
9	d
10	a
11	c
12	a
13	d
14	c
15	b
16	b
17	c
18	b
19	b
20	c
21	a
22	d
23	b
24	b

Detailed Explanations

Solution to Question 1:

The therapeutic index is 4.

In the above graph,

Median lethal dose (LD50) = 400

Median effective dose (ED50) = 100

Therapeutic index (TI) = LD50/ED50.

Hence,

Therapeutic index (TI) = 400/100 = 4

Clinically, the therapeutic range bound by the dose that produces the minimal therapeutic effect and the dose that produces the maximal acceptable adverse effect is more relevant.

Solution to Question 2:

In the above image, drug B is a competitive antagonist.

In competitive antagonism, the DRC is shifted to the right whereas in non-competitive antagonism the DRC is flattened.

Thus in the above image Drug B is the Competitive antagonist while drug C is the non-competitive antagonist.

Potency:

• It is defined as the amount of drug needed to produce a given response.

• A highly potent drug evokes a given response at low concentrations and vice-versa.

• The more the curve is on the left in a dose-response curve (DRC), the greater is its potency.

• Thus, in the above image, the potency is drug A > drug C > drug B.

Efficacy:

• It is defined as the maximum effect produced by the drug.

• It expresses the effect produced by the receptors when bound by a certain drug.

• In the log dose-response curve, the perpendicular height represents the drug efficacy.

• Thus, in the above image, the efficacy is drug A = drug B > drug C.

Solution to Question 3:

The use of brimonidine is unsafe in infants. It is an alpha-2 agonist that can cross the blood-brain barrier. This may result in CNS depression and apnoea in neonates. Therefore it is contraindicated in children less than 2 years.

α2 agonists, through the activation of G-protein coupled receptor activation, inhibit the activity of adenylate cyclase. This reduces cAMP and hence reduces aqueous humor production by ciliary action.

Solution to Question 4:

In the present case, the volume of distribution of the drug is 6L/kg (420 L in a normal adult with reference weight 70kg) which is very high when compared to the total body water of 42L. Therefore, it can be implied that the drug is having high sequestration in tissues.

The apparent volume of distribution (Vd) is an imaginary volume throughout which the drug is distributed if the plasma concentration remains constant throughout.

Vd = Dose administered IV/ plasma concentration

The difference in the volume of distribution based on drug type:

• Low Vd

• Restricted to the intravascular compartment

• Seen in drugs with high molecular weight (e.g., heparin) or high protein binding (e.g., warfarin - 9.8L)

• Medium Vd

• Lipid insoluble drugs (e.g., gentamicin - 20L) do not enter the cells and are distributed in the extracellular fluid volume

• Small water-soluble molecules (e.g., ethanol - 42L) are distributed in the total body water

• High Vd

• Sequestered and accumulated in the tissues

• Small lipophilic molecules (e.g., chloroquine - 13000L)

Solution to Question 5:

Efficacy and appropriate dose range is the main aim of phase 2 of clinical trials. It is evaluated on a small group of patients with the target disease.

Phase II of clinical trials involve a small group of patients (50-500). Most of the new drugs will be administered for the first time in patients with the target disease. The study is designed to obtain proof of concept (potential of the drug to treat the disease) and dose-ranging study (provides input for choosing the most appropriate dose for phase 3 studies).

Phase 3 studies are also known as pivotal studies as they establish treatment benefits.

Solution to Question 6:

Rivaroxaban is an oral direct factor Xa inhibitor. They represent a new class of oral anticoagulant drugs that do not require monitoring.

This group of drugs inhibit factor Xa in the final common pathway of clotting [conversion of Prothrombin (Factor II) to thrombin (Factor IIa)]. They are given as fixed doses and have a rapid onset of action.

Oral direct factor Xa inhibitors are indicated in:

• Prevention of embolic stroke in patients with atrial fibrillation (without valvular heart disease)

• Venous thromboembolic disease (VTE)

• Postoperative thromboprophylaxis in patients undergoing hip and knee arthroplasty

Solution to Question 7:

The drug of choice for diphtheria carrier state is erythromycin.

Erythromycin for 7 days is very effective for eradicating the diphtheria carrier state or acute infections.

Solution to Question 8:

Oseltamivir and zanamivir are neuraminidase inhibitors.

They competitively and reversibly interact with the enzyme sites to inhibit viral neuraminidase activity. This interferes with the release of the progeny from infected host cells, halting the spread of infection.

Early administration hastens the resolution of symptoms if given within 48 h of infection. Oseltamivir is given at a dose of 75 mg, BD for 5 days for the treatment of Influenza A and B.

For prophylaxis, it is given at a dose of 75mg, OD and continued for the duration of exposure.

Solution to Question 9:

Carboprost is a 15-methyl synthetic derivative of PGF2 alpha. It is used as an abortifacient and in refractory PPH.

Options A: Dinoprost is a natural PGF2 alpha preparation.

Option B: Dinoprostone is natural PGE2 preparation.

Option C: Misoprostol is a synthetic PGE1 analog. Gemeprost is also a synthetic PGE1 analog.

Solution to Question 10:

Nicorandil causes activation of K+-ATP channels. This leads to the hyperpolarization of the smooth muscle cells leading to vasodilatation.

It also has nitrate like cGMP dependent action. Nicorandil dilates both arteries and veins leading to a decrease in afterload and preload of the heart.

Molsidomine, isosorbide dinitrate, and nitroglycerin are nitrates. They do not act on K-ATP channels.

Solution to Question 11:

Among the given options, the most effective antiemetic for chemotherapy-induced vomiting is granisetron.

Granisetron is a 5HT3 receptor antagonist. They are the most effective drugs for the treatment of chemotherapy-induced and post-operative nausea and vomiting.

Palonosetron is a second-generation 5HT3 antagonist that has the highest affinity for 5HT3 receptors and has the maximum potency. It is also the longest acting that makes it effective in delayed emesis associated with chemotherapy. Palonosetron has minimal/no propensity to cause QT interval prolongation.

Solution to Question 12:

Teduglutide is a GLP-2 analogue.

It has been recently approved for the treatment of short bowel syndrome. Teduglutide binds to enteric neurons and endocrine cells, stimulating the release of a number of hormones such as insulin-like growth factor, this stimulates mucosal epithelial growth and enhances fluid absorption.

The drug has a t1/2 of 1–2 hours and is administered subcutaneously once daily. It can potentially cause bowel cancer, therefore it is not recommended for patients with active malignancies.

Note: GLP-1 analogs are used for the treatment of Type II diabetes mellitus.

Solution to Question 13:

Cisatracurium is preferred over atracurium because it causes less release of histamine.

Other options:

Option A- Cisatracurium has a slower onset of action

Option B and C- Both atracurium and cisatracurium have similar duration of action and cardiac action.

Solution to Question 14:

Colchicine inhibits neutrophil (granulocyte) recruitment.

Colchicine acts by preventing the polymerisation of microtubules by binding to tubulin. This reduces neutrophil recruitment to inflamed tissue and also decreases the secretion of chemotactic factors by activated neutrophils, thus reducing the inflammatory response overall.

Colchicine is used in gout and Mediterranean fever.

Solution to Question 15:

Levofloxacin and ofloxacin have the maximum bioavailability among fluoroquinolones. Hence, intravenous and oral doses are the same for these drugs.

Solution to Question 16:

Triazoles act by inhibition of ergosterol synthesis.

The azole group inhibits the fungal cytochrome P450 enzyme called 14-α-sterol demethylase and results in an impairment of conversion of lanosterol to ergosterol. This results in the depletion of membrane ergosterol.

Azole antifungals are of two types:

• Imidazoles - clotrimazole, ketoconazole, luliconazole

• Triazoles - itraconazole, fluconazole, voriconazole, posaconazole and isavuconazole

Solution to Question 17:

Female patients should avoid conception for 3 years after discontinuing acitretin treatment to prevent retinoid-induced embryopathy.

Acitretin is approved for use in the cutaneous manifestations of psoriasis.

Absolute contraindications:

• Pregnant women

• Women who are planning to get pregnant

• Breastfeeding

Relative contraindications:

• Leukopenia

• Alcoholism

• Hyperlipidemia

• Hypercholesterolemia

• Hypothyroidism

• Significant hepatic or renal disease

Note: Female patients should avoid conception for 1 month after discontinuation of isotretinoin.

Solution to Question 18:

Cilastatin is administered to prolong the action of imipenem.

Imipenem is rapidly hydrolyzed by the enzyme renal dehydropeptidase. It is administered with cilastatin, a reversible inhibitor of dehydropeptidase to inhibit the breakdown and prolong the action of imipenem.

Meropenem is not hydrolyzed by renal dehydropeptidase. Cilastatin administration is not necessary with meropenem.

Solution to Question 19:

The drug used to differentiate myasthenia gravis from a cholinergic crisis is edrophonium.

Muscle weakness is a common presentation in patients with myasthenia gravis and cholinergic crisis (excessive dose of anti-ChE drugs). Intravenous injection of edrophonium improves muscular weakness in a patient with myasthenia gravis and not in a case of cholinergic crisis.

The test used for this differentiation is known as the edrophonium test.

Initially, 2 mg is administered; if improvements of symptoms do not occur within 45 seconds, an additional 8 mg can be administered. Improvement of strength is a positive test for myasthenia gravis.

Solution to Question 20:

Patients with sickle cell anemia should not receive pegylated filgrastim as it can trigger a sickle cell crisis.

Filgrastim is a recombinant human G-CSF (granulocyte colony-stimulating factor). The pegylated forms of filgrastim, pegfilgrastim, and lipefilgrastim, are longer-acting forms. It is administered subcutaneously or by slow intravenous infusion only.

Indications of recombinant G-CSF:

• Severe neutropenia after autologous hematopoietic stem cell transplantation

• Severe neutropenia after high dose cancer chemotherapy

• Severe congenital neutropenias

• Neutropenia of patients with AIDS receiving zidovudine

Solution to Question 21:

The best approach in the treatment of chronic SIADH is the oral administration of a vaptan like tolvaptan, which is a selective V2 receptor antagonist. This increases urinary water excretion by blocking the antidiuretic effect of AVP.

Other treatment options for chronic SIADH include:

• Fluid intake < 500 ml/day

• Demeclocycline

• Fludrocortisone

Increased thirst is a frequently observed characteristic of hyponatremia in SIADH. Fluid restriction is usually a burden to the patient. It is unreliable, impractical, and slow to work. As thirst is particularly difficult to suppress, water restriction in SIADH frequently leads to non-compliance in the long run. Vaptans would be especially suited for the treatment of these patients. Serum sodium concentrations are also corrected by vaptans and hence, vaptans substantially improve the quality of life in these patients.

Solution to Question 22:

Intravenous mannitol is used for the treatment of acute attacks of glaucoma.

Uses of mannitol are:

• To reduce cerebral edema and intracranial pressure

• Enhancement of urinary excretion of salicylates, barbiturates, bromides, and lithium

• Diagnosis of bronchial hyperreactivity (inhalation)

• Antihemolytic urological irrigation during transurethral procedures

• Treatment of acute attacks of glaucoma.

Solution to Question 23:

Imipenem is the carbapenem that is associated with an increased risk of causing seizures, especially when high doses are given to patients with CNS lesions.

Excretion of imipenem is mainly renal and dosage modification is required to avoid seizures in renal failure patients.

Solution to Question 24:

Prostaglandin E1 (PGE1) maintains the patency of ductus arteriosus (PDA). This is used in term neonates with underlying cyanotic congenital heart disease awaiting surgical intervention as they have duct-dependant circulation.

Indomethacin and Ibuprofen are the NSAID indicated for closing the ductus arteriosus in premature infants and neonates.

Pharmacology NEET 2020

Question 1:

Which of the following is an inhibitor of DNA synthesis?

a) 6-mercaptopurine

b) Mitomycin

c) Actinomycin

d) Asparaginase

Question 2:

A patient on lithium therapy developed hypertension. He was started on thiazides for hypertension. After few days, he developed coarse tremors and other symptoms suggestive of lithium toxicity. Explain the likely mechanism of this interaction.

a) Thiazide inhibits the metabolism of lithium

b) Thiazides act as an add on drug to lithium

c) Thiazides increase the tubular reabsorption of lithium

d) Thiazides cause water loss therby increase lithium levels

Question 3:

Which of the following should not be prescribed with theophylline?

a) Erythromycin

b) Cefotaxime

c) Cotrimoxazole

d) Amoxicillin

Question 4:

Which of the following drugs gets deposited in cornea?

a) Leflunamide

b) Chloroquine

c) Methotrexate

d) Sulfasalazine

Question 5:

A patient was prescribed an antipsychotic drug following which he developed parkinsonism. Which of the following drugs■ can be given?

a) Trihexiphenidyl

b) Entacapone

c) Rasagiline

d) Pramipexole

Question 6:

Which of the following antiarrhythmic drugs is contraindicated in a patient with interstitial lung disease?

a) Amiodarone

b) Sotalol

c) Quinidine

d) Lignocaine

Question 7:

Which of the following drugs acts by inhibiting the late sodium current in the heart?

a) Ivabradine

b) Ranolazine

c) Trimetazidine

d) Fasudil

Question 8:

A patient with diabetes and COPD developed post-operative urinary retention. Which of the following drugs can be used for short-term treatment of the condition?

a) Tamsulosin

b) Bethanechol

c) Terazosin

d) Methacoline

Question 9:

Which of the following drugs is used for the treatment of motion sickness?

a) Cetrizine

b) Promethazine

c) Misolastin

d) Loratidine

Question 10:

Which of the following is not a prokinetic?

a) 5HT4 agonist

b) Dopamine antagonist

c) Motilin

d) Diphenylmethane

Question 11:

Which of the following is used in the first-line management of postmenopausal osteoporosis?

a) Estrogen

b) Bisphosphonates

c) Raloxifene

d) Combined oral contraceptives

Question 12:

What is the mechanism of resistance by lactamases■ to β-lactam antibiotics?

a) Alteration of penicillin binding protein

b) Degradation of antibiotics

c) Drug efflux

d) Alteration in 50s ribosome structure

Question 13:

A female patient is on first-line pharmacological therapy for bipolar disorder for 6 months. She is fasting for a few days due to religious reasons and presented with seizures, coarse tremors, confusion, and weakness of limbs. Which of the following investigations should be done to help diagnose her condition?

a) Serum potassium level

b) Serum lithium level

c) ECG

d) MRI

Question 14:

Pegloticase is used in management of:

a) Chronic gout

b) Rheumatoid arthritis

c) Psoriatric arthritis

d) Calcium pyrophosphate dihydrate crystal deposition (CPPD)

Question 15:

Which of the following is an inhibitor of RNA synthesis?

a) 5-Fluorouracil

b) Rifampicin

c) Nitrofurantoin

d) Novobiocin

Question 16:

Which of the following fluoroquinolone is not given in liver disease?

a) Ofloxacin

b) Levofloxacin

c) Pefloxacin

d) Lomefloxacin

Question 17:

Theophylline causes diuresis by:

a) Adenosine antagonism

b) PDE 4 inhibition

c) Beta 2 agonism

d) PDE 3 inhibition

Question 18:

The graph below shows the relationship between the log(dose) and the response of three drugs, A, B, and C. Which of these drugs has the highest potency?



a) A

b) B

c) Both A and B

d) C

Question 19:

A patient presented with biliary colic. The intern has administered an analgesic, following which the pain got aggravated. Which of the following could be responsible for the aggravation of the patient's symptoms?

a) Morphine

b) Diclofenac

c) Nefopam

d) Etoricoxib

Question 20:

A patient with gout develops hypertension. The uric acid level was found to be elevated. Which of the following is not administered?

a) Hydrochlorothiazide

b) Atenolol

c) Enalapril

d) Prazosin

Question 21:

Voriconazole is not effective against which of the following?

a) Candida albicans

b) Mucormycosis

c) Candida tropicalis

d) Aspergillosis

Question 22:

A patient who was diagnosed with ADPKD was prescribed tolvaptan. After a few weeks, he presented with dry mouth and increased thirst. What is the likely cause?

a) Increased free water clearance

b) V2 receptor agonism

c) Increased renal cAMP levels

d) Increase in urine osmolality

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	b
5	a
6	a
7	b
8	a
9	b
10	d
11	b
12	b
13	b
14	a
15	b
16	c
17	a
18	a
19	a
20	a
21	b
22	a

Detailed Explanations

Solution to Question 1:

6-Mercaptopurine is an inhibitor of DNA synthesis.

6-Mercaptopurine is inactive in its parent form. It must be metabolized by hypoxanthine-guanine phosphoribosyltransferase to form monophosphate nucleotide

6-thioinosinic acid. This inhibits several enzymes of de novo purine nucleotide synthesis.

Option B- Mitomycin acts as an alkylating agent and crosslinks DNA. It also generates free radicals which damage DNA.

Option C- Actinomycin causes single-strand breaks in DNA. It blocks RNA transcription as well.

Option D- L-Asparaginase causes the breakdown of asparagine and thus interferes with protein synthesis.

Solution to Question 2:

Thiazides act by increasing the tubular reabsorption of lithium.

Thiazide use is accompanied by sodium loss which is compensated in a few days by increased reabsorption of Na+ in the proximal tubule.

The kidney handles both lithium and Na+ in the same way. Thus, along with Na+ reabsorption, there will be an increase in the reabsorption of lithium also, resulting in decreased excretion of lithium.

Solution to Question 3:

Erythromycin should not be prescribed with theophylline.

Theophylline is mainly metabolized by CYP1A2 in the liver. Erythromycin inhibits CYP1A2 and thereby decreases the clearance of theophylline.

Factors decreasing the clearance of theophylline are:

• Drugs - erythromycin, cimetidine, ciprofloxacin, allopurinol, fluvoxamine, zileuton, zafirlukast

• Congestive heart failure

• Liver disease

• Pneumonia

• Viral infection and vaccination

• High-carbohydrate diet

• Old age

Solution to Question 4:

Chloroquine, an anti-malarial drug may get deposited in the cornea leading to corneal opacity.

Ocular toxicity of chloroquine:

• Short-term therapy - blurring of vision and diplopia, reversible on stoppage of drug.

• Long-term use (high-dose):

• Keratopathy

• Pigmentary retinopathy

• Optic atrophy

Hydroxychloroquine has lower ocular toxicity compared to chloroquine.

Drugs causing corneal opacity:

• Amiodarone

• Phenothiazine

• Indomethacin

• Chloroquine

Long-term treatment with these drugs can cause a whorl-like opacity in the corneal epithelium called cornea verticillata. It is also seen in Fabry's disease.

Solution to Question 5:

Trihexyphenidyl is the drug of choice for drug-induced parkinsonism. It is a central anticholinergic.

Drug-induced parkinsonism is a condition caused by certain drugs having D2 antagonistic action (typical antipsychotics). The decrease in dopaminergic activity results in an unbalanced cholinergic activity. Anticholinergics act by restoring the balance.

Solution to Question 6:

Amiodarone is contraindicated in a patient with interstitial lung disease.

The most serious adverse effect during chronic amiodarone therapy is pulmonary fibrosis, which can be rapidly progressive and fatal. Underlying lung disease, doses of 400 mg/day or more, and recent pulmonary insults such as pneumonia appear to be risk factors for this condition.

Amiodarone is a class III antiarrhythmic drug.

The side effects of amiodarone include:

• Pulmonary fibrosis

• Hypotension, depressed ventricular function

• Torsades de pointes

• Corneal microdeposits

• Hepatotoxicity

• Neuropathies

• Photosensitivity

• Thyroid dysfunction

Solution to Question 7:

Ranolazine acts by inhibiting the late sodium current in the heart.

Late sodium current (INa) facilitates entry of calcium via the sodium-calcium exchanger. Inhibition of INa, therefore, reduces the intracellular calcium concentration which in turn reduces the diastolic tension, cardiac contractility and work.

At higher doses, it may also modify the fatty acid oxidation, which improves the efficiency of cardiac oxygen utilization.

It is used in the prophylaxis of angina.

Solution to Question 8:

Tamsulosin is the drug used to relieve the symptoms in cases of postoperative urinary retention in a patient with COPD.

It is also efficacious in the treatment of benign prostatic hyperplasia (BPH) with little effect on blood pressure.

Option B: Bethanechol is the drug of choice to relieve the symptoms in cases of postoperative urinary retention. However, it can cause bronchoconstriction in patients with COPD. Hence tamsulosin is the best answer here.

Option C: Terazosin is an $\alpha 1$ receptor antagonist, used in the treatment of benign prostatic hyperplasia (BPH).

Option D: Methacholine is a muscarinic agonist but acts specifically on bronchial smooth muscle causing its contraction. It is available as an inhalational agent and is used for the diagnosis of bronchial hyper-reactivity in patients who do not have clinically apparent asthma.

Solution to Question 9:

Promethazine is used for motion sickness. It is an antihistaminic agent that acts as an antagonist of H1 receptors. It also has central anticholinergic activity that is responsible for its antiemetic action.

Solution to Question 10:

Diphenylmethane is not a prokinetic agent. Its derivative (Bisacodyl) is used as a laxative.

Prokinetic agents are the drugs that increase GI motility and transit of material in the GI tract. These agents include:

• Dopamine receptor antagonists- metoclopramide and domperidone

• Serotonin receptor agonists- cisapride and prucalopride

• Motilin

• Macrolide antibiotics mimicking motilin - erythromycin

• Miscellaneous agents:

• Sincalide (C-terminal octapeptide of CCK)

• Camicinal (motilin receptor agonist)

• Relamorelin (ghrelin receptor agonist)

• Velusetrag and naronapride (5HT4 agonists)

Solution to Question 11:

Bisphosphonates are used in the first-line management of postmenopausal osteoporosis.

Bisphosphonates are analogs of pyrophosphates. They specifically inhibit bone resorption and thus are very effective for the treatment of postmenopausal osteoporosis.

Bisphosphonates concentrate at sites of active remodeling. They remain in the matrix until the bone is remodeled. Then, they are released in the acid environment of the resorption lacunae and induce apoptosis in osteoclasts.

Classification of bisphosphonates (mainly based on potency):

First-generation:

• Etidronate (least potent)

• Clodronate

• Tiludronate

Second-generation: 10-100 times more potent than the first generation

• Alendronate

• Pamidronate

• Ibadronate

Third-generation: 10,000 times more potent than the first generation

• Risedronate

• Zoledronate

Solution to Question 12:

Resistance by lactamases is through the degradation of antibiotics.

The bacterial cell wall is made of heteropolymeric peptidoglycan that confers mechanical stability. The β-lactam antibiotics inhibit the last step in peptidoglycan synthesis by acylating the transpeptidase via cleavage of the –CO–N– bond of the β-lactam ring. These transpeptidase targets are collectively called Penicillin Binding Proteins (PBPs)

Beta-lactamases are enzymes produced by certain bacteria that inactivates β-lactam antibiotics by hydrolysis of the β-lactam ring in their structure.

The resistance to penicillins and other beta-lactam antibiotics can be conferred through one of the four mechanisms:

• Inactivation of antibiotic by lactamases

• Modification of target PBPs

• Impaired penetration of drug to target PBPs

• Antibiotic efflux

Solution to Question 13:

The clinical features given in the question are most likely suggestive of lithium toxicity, and hence serum lithium levels should be measured to confirm it.

Lithium is a mood stabilizer with a very narrow therapeutic index and is the first line of management in bipolar disorder.

The kidney handles both lithium and Na+ in the same way via ENac channel. Thus during fasting/dehydration, along with Na+ reabsorption in renal tubules, there will be an increase in the reabsorption of lithium leading to lithium toxicity.

Lithium has a very narrow therapeutic index of 0.8-1.2 mEq/L. The severity of lithium toxicity is often divided into the following three grades:

A fine tremor is common in patients taking lithium at therapeutic doses. Therapeutic lithium tremor is generally symmetric, limited to the arms, and nonprogressive. The tremor from lithium toxicity is coarser and more disabling, may also affect the legs, and occurs with other symptoms of toxicity. Other adverse effects include mental confusion, hypothyroidism, nephrogenic diabetes insipidus, and edema.

As lithium is readily dialyzed, lithium overdose or toxicity is treated with hemodialysis.

*Note - Serum Na+ levels are significant with respect to lithium toxicity, where as serum potassium levels becomes significant in digoxin toxicity. Hypokalemia increases the risk of digoxin toxicity.

Grade	Sr. lithium concentration(in mEq/L)	Symptoms
Mild symptoms	1.5-2.5	nausea, vomiting,tremor, and fatigue
Moderate intoxication	2.5-3.5	confusion, agitation, delirium, tachycardia, and hypertonia

Grade	Sr. lithium concentration(in mEq/L)	Symptoms
Severe intoxication	>3.5	coma,seizures, hyperthermia, and hypotension

Solution to Question 14:

Pegloticase is used in the treatment of chronic gout.

Pegloticase is a pegylated uricase (urate oxidase). It catalyzes the enzymatic oxidation of uric acid into allantoin, which is a more soluble and inactive metabolite.

It is used in the treatment of severe,refractory chronic gout or in cases where other urate-lowering therapies are contraindicated.

Solution to Question 15:

Rifampicin is an inhibitor of RNA synthesis. It binds to the beta subunit of DNA dependent RNA-polymerase (rpoB) to form a stable drug-enzyme complex. The binding suppresses the chain formation in RNA synthesis.

Option A: 5-Fluorouracil inhibits thymidylate synthesis.

Option C: Nitrofurantoin produces highly reactive intermediates that cause DNA damage

Option D: Novobiocin is a potent inhibitor of bacterial DNA gyrase.

Solution to Question 16:

Pefloxacin is mainly metabolized in the liver and hence avoided in patients with liver disease.

Quinolones with primarily hepatic metabolism include moxifloxacin, pefloxacin, and trovafloxacin (withdrawn from the market due to hepatotoxicity).

Quinolones primarily excreted by the kidneys include ciprofloxacin, levofloxacin, norfloxacin, lomefloxacin, and ofloxacin.

Solution to Question 17:

The diuretic effect of theophylline is due to blockade of adenosine (A1) receptors.

Theophylline is a methylxanthine drug that acts by:

• Inhibition of phosphodiesterase enzyme (PDE). PDE3 inhibition causes relaxation of bronchial smooth muscle and PDE4 inhibition causes inhibition of the release of cytokines and chemokines

which further leads to a decrease in immune cell migration and activation.

• Blockade of adenosine receptors, which inhibits bronchoconstriction. However, blockade of A1 receptors in the glomerular afferent arteriole and PCT causes diuresis.

• Activating enzyme histone deacetylase-2, which is responsible for its anti-inflammatory action

Solution to Question 18:

Drug A has the highest potency.

Potency:

• It is defined as the amount of drug needed to produce a given response. A highly potent drug evokes a given response at low concentrations and vice-versa.

• The more the curve is on the left in a dose-response curve (DRC), the greater is its potency.

• Thus, in the above image, the potency is Drug A >Drug B> Drug C.

Efficacy:

• It is defined as the maximum effect produced by the drug. It reflects the capacity of the drug to activate a receptor and produce the response.

• In the log dose-response curve, the perpendicular height represents the drug efficacy.

• Thus, in the above image, the efficacy is Drug A = Drug B > Drug C.

• Drugs having a similar pharmacological effect can have different efficacies e.g furosemide and metolazone are both diuretics but furosemide has greater efficacy than metolazone.

Solution to Question 19:

Morphine is the analgesic which could be responsible for the aggravation of the patient's symptoms.

It causes constriction of the sphincter of Oddi causing an increase in fluid pressure in the common bile duct and gall bladder.

Note: Pethidine causes lesser spasm of the sphincter of Oddi and can be used in biliary colic.

Solution to Question 20:

Hydrochlorothiazides are not given to this patient.

Gout is a metabolic disease associated with hyperuricemia.Thiazides increase direct urate reabsorption in the proximal renal tubules and are associated with elevated serum uric acid levels and may precipitate gout. Hence, hydrochlorothiazides are not used to treat hypertension in gout patients.

Solution to Question 21:

Voriconazole is not effective against mucormycosis.

Mucormycosis is treated by amphotericin B. Posaconazole can also be used. However, other azoles are not effective against this fungus.

Voriconazole is the drug of choice for the treatment of aspergillosis. It can also be used for Candida species (including glabrata and krusei), Scedosporium and Fusarium.

Solution to Question 22:

Tolvaptan causes increased free water clearance, which causes dehydration. Hence, the patient may present with dry mouth and increased thirst.

Tolvaptan is a vasopressin V2 receptor antagonist. Decreased binding of vasopressin to the V2 receptor in the kidney lowers the adenyl cyclase activity leading to decreased intracellular cAMP. Decreased cAMP concentrations prevent the fusion of aquaporin 2 containing vesicles with the plasma membrane, leading to an increase in urinary water excretion (free water clearance), and a decrease in urine osmolality.

The common side effects of tolvaptan include:

• dehydration

• hypernatremia

• hypovolemia

• liver injury

Tolvaptan is used to delay disease progression in patients with ADPKD.

Pharmacology INI-CET 2021

Question 1:

Which of the following is false about the selection of essential drugs?

a) Adequate safety profile needs to be established

b) Cost to benefit has to be considered

c) Fixed drug combination is preferred over single drugs

d) Disease prevalence is considered

Question 2:

A 34-year-old man with hypertension is diagnosed with stage 4 chronic kidney disease, with eGFR of <30. His doctor wants to prescribe a thiazide diuretic. Which of the following can be given?

a) Chlorothiazide

b) Chlorthalidone

c) Metolazone

d) Hydrochlorothiazide

Question 3:

A patient who was diagnosed with epilepsy was put on retigabine TDS. Now phenytoin is being added. Which of the following changes should be made to retigabine?

a) Increase the dose

b) Decrease the dose

c) Stop the drug

d) Stop retigabine and start on carbamazepine

Question 4:

A patient with history of ischemic stroke was started on clopidogrel. However, she had another attack of stroke after 6 months. Which of the following is likely to be responsible for

the failure of clopidogrel in this patient?

a) Downregulation of CYP2C19

b) Downregulation of CYP2D6

c) Upregulation of CYP1A1

d) Downregulation of CYP2E1

Question 5:

Which of the following can be given for long term management of obesity?

a) Liraglutide

b) Metformin

c) Sibutramine

d) Fenfluramine

Question 6:

An HIV-positive woman on ART was prescribed ergotamine for a migraine attack 4 days back. She presented with complaints that she is unable to feel her legs from the mid-thigh to her toes, for the past two days. The angiography image is given below. What is the likely diagnosis?



a) Atherosclerosis

b) Ergotism

c) Polyradicular neuropathy

d) Herniated disc

Question 7:

Topiramate is used for

a) Treatment of migrane

b) Treatment of Lennox Gastaut syndrome

c) Prophylaxis of ADHD

d) Treatment of dementia

Question 8:

Which of the following is incorrectly matched?

a) Andexanet alfa: Edoxaban

b) Andexanet alfa: Rivaroxaban

c) Idarucizumab : Dabigatran

d) Ciraparantag: Fondaparinux

Question 9:

Inhaled Nitric oxide is approved for treatment of

a) Labile hypertension

b) Pulmonary hypertension

c) Treatment of resistant migraine

d) Resistant Parkinson's disease

Question 10:

Nivolumab is used as a checkpoint inhibitor in

a) Retinoblastoma

b) Medulloblastoma

c) Pleuropulmonary blastoma

d) Hodgkin's lymphoma

Question 11:

Paradoxical bronchoconstriction seen with ipratropium bromide is due to all except

a) Hypertonic saline

b) EDTA

c) Benzalkonium chloride

d) Prejunctional M2 receptor blockade

Question 12:

Antimicrobial combinations are used in all except

a) Gonorrhea

b) Tuberculosis

c) Intra abdominal infections

d) Malaria

Question 13:

Which of the following steps does Vancomycin act on?



©MARROW

a) 1

b) 2

c) 3

d) 4

Question 14:

What would happen to the half-life and plasma concentration of a drug which follows first-order kinetics, if the dose is doubled?

a) Half-life doubles and plasma concentration remains the same

b) Half-life remains the same and plasma concentration doubles

c) Half-life and plasma concentration remains the same

d) Half life and plasma concentration doubles

Question 15:

Which of the following is not an off-label use of risperidone?

a) OCD

b) Dementia

c) PTSD

d) Bipolar disorder

Question 16:

The elimination rate constant of a drug is 0.05/hr. What is its half-life?

a) 20.8 hours

b) 6.9 hours

c) 13.8 hours

d) 9.8 hours

Question 17:

A patient is on phenytoin for a seizure disorder. He was prescribed sucralfate 4 times a day for peptic ulcers. What should be the minimum duration of time between consumption of these drugs?

a) 30 minutes

b) 60 minutes

c) 90 minutes

d) 120 minutes

Question 18:

Category A, B, C, D, X in drug classification are indicative of which of the following?

a) Safety in pregnancy

b) Dose adjustment in renal failure

c) Cost ratio from cheap to expensive

d) As per safety or therapeutic index

Question 19:

Identify the false statement regarding suicidal inhibition.

a) The inhibitor can bind with any site resulting in suicidal inhibition

b) The binding of the enzyme to the substrate analogue is irreversible

c) They are enzyme specific and used in rational drug design

d) The inhibitor forms a product with the enzyme and the product inhibits it

Question 20:

The dose at which dopamine acts on beta 1 receptors is ____.

a) 2-10 mcg/kg/min

b) <2 mcg/kg/min

c) 10-20 mcg/kg/min

d) >20 mcg/kg/min

Question 21:

Which of the following is the incorrect statement regarding clomiphene citrate?

a) Causes monofollicular development

b) Inhibits negative feedback on GnRH

c) Causes endometrial thinning

d) Can be used with gonadotropin

Question 22:

What is the mechanism of action of remdesivir?

a) RNA-dependent DNA polymerase inhibition

b) RNA-dependent RNA polymerase inhibition

c) Fusion inhibition

d) Entry inhibition

Question 23:

Administration of which of the following drugs would increase the bioavailability of saquinavir?

a) Ritonavir

b) Vitamin C

c) Cimetidine

d) Ganciclovir

Question 24:

The table below gives the AUC of drug B alone and that of drug B when combined with another drug A. If the ■ value is <0.01, then which of the following statements is true?

a) Drug A decreases intestinal absorption of Drug B

b) Drug A reduces first pass metabolism of Drug B

c) Drug A increases the metabolism of Drug B

d) Drug A increases renal elimination of Drug B

Question 25:

What is the mechanism of action of cyclosporine?

a) Inhibition of calcineurin

b) AMP kinase stimulation

c) IMP dehydrogenase inhibition

d) Dihydro-orotate dehydrogenase inhibition

Question 26:

Anionic and weakly acidic drugs bind to which of the following?

a) Albumin

b) Globulin

c) Lipoprotein

d) Alpha-1 glycoprotein

Question 27:

A 14-month-old boy was diagnosed with Hemophilus influenza meningitis. The initial treatment plan was to start on ampicillin but later was shifted to ceftriaxone. What could be the probable reason?

a) Hemophilus alters penicillin binding protein sites, hence resistant to ampicillin

b) Easier to administer IV ceftriaxone than IV ampicillin

c) Cotrimoxazole is the drug of choice, but patient is allergic to sulfa drugs

d) Hemophilus produces beta lactamase, hence resistant to ampicillin

Question 28:

A patient who is a known case of cirrhosis of the liver presents with abdominal distension and swelling of the lower limbs. The preferred diuretic for this patient would be

a) Triamterene

b) Eplerenone

c) Chlorthalidone

d) Furosemide

Question 29:

Which of the following is the most effective drug in smoking cessation?

a) Varenicline

b) Buspirone SR

c) Nicotine gum

d) Rimonabant

Question 30:

A patient who was on warfarin therapy for atrial fibrillation is prescribed a cephalosporin antibiotic for the treatment of a lower respiratory tract infection. He reports back to the hospital after having developed hemarthrosis. Which of the following cephalosporins is likely to have caused this?

a) Cefixime

b) Cefoperazone

c) Ceftobiprole

d) Ceftazidime

Question 31:

Which of the following is a novel anti-depressant?

a) Lurasidone

b) Vilazodone

c) Asenapine

d) Blonanserin

Question 32:

A 57-year-old man was brought to the hospital with complaints of dizziness and breathlessness. During his physical examination, you notice the following discolouration over his face. He is presently being treated with amiodarone, metoprolol, verapamil, lisinopril, and amlodipine. Which of the following drugs is most likely responsible?

a) Amiodarone

b) Amlodipine

c) Metoprolol

d) Verapamil

Question 33:

Which of the following is a centrally acting opioid antitussive?

a) Ambroxol

b) Diphenoxylate

c) Levopropoxyphene

d) Levorphanol

Question 34:

The anti-epileptic drugs which can also be used for the treatment of neuropathic pain are

a) 1, 2 and 3

b) 1, 2 and 4

c) 1, 3 and 4

d) 2 and 4 only

Question 35:

A breast cancer patient post-chemotherapy developed a low-grade fever with anemia and leukocytopenia (<4000/cm3; DNC: 30). Which of the following antibiotics will not be used in the empirical management of this patient?

a) Cefepime

b) Piperacillin-Tazobactam

c) Linezolid

d) Meropenem

Answer Key

Question No.	Correct Option
1	c
2	c
3	a
4	a
5	a
6	b
7	b
8	d
9	b
10	d
11	a
12	a
13	d
14	b
15	d
16	c
17	d
18	a
19	a
20	a
21	a
22	b

23	a
24	b
25	a
26	a
27	d
28	b
29	a
30	b
31	b
32	a
33	c
34	b
35	c

Detailed Explanations

Solution to Question 1:

Fixed drug combinations are not preferred over single drugs while selecting essential drugs.

Criteria for selection of essential medicines as per WHO:

• There should be adequate data on the safety and efficacy of the drug.

• The drug should be available in a form where its quality including bioavailability and stability on storage is assured.

• The choice of the drug depends on the pattern of prevalent diseases, financial resources, availability of trained personnel and facilities, genetic, demographic, and environmental factors.

• The cost-benefit ratio is an important factor.

• Pharmacokinetic properties of the drug and the local manufacture and storage facilities are also considered.

• Essential drugs should preferably be single compounds.

• Fixed combinations are used only when the dosage of each component meets the requirement of the population in which it is used. It should also have the advantage in safety, therapeutic effect, adherence, and prevention of drug resistance when compared to single compounds.

• The selection of essential drugs is a continuous process and changes are to be made with developments in public health and the availability of better drugs.

• The selection should be based on rationally developed treatment guidelines.

Note:

In 1977 WHO brought the first "Model List of Essential Drugs". India made its "National Essential Drug List" in 1996 and revised the list in 2011 and renamed it as "National List of Essential Medicines". The latest edition of WHO Model List of Essential Drugs is the 21st edition published in 2019.

Solution to Question 2:

The thiazide diuretic that can be used as an antihypertensive in this patient with chronic kidney disease is metolazone.

In patients with chronic renal failure, thiazide diuretics are ineffective. But metolazone, despite being a thiazide diuretic, is effective as antihypertensives in patients with chronic renal failure.

Metolazone is useful even in severe renal failure (GFR of ~15 mL/min). It has an additive action when combined with loop diuretics like furosemide. It also has an additional proximal tubular action (thiazides mainly act on DCT). It is commonly used for edema and is occasionally used as an antihypertensive.

Diuretics in hypertension:

Thiazides are the diuretics of choice for uncomplicated hypertension.

Mechanisms of antihypertensive action:

• Decreases plasma and ECF volume by diuresis thereby decreasing cardiac output

• Even when the compensatory mechanisms restore the Na+ levels and plasma volume, a low BP is maintained by a reduction in total peripheral resistance.

• This reduction in total peripheral resistance is due to the persistence of a small sodium and volume deficit even after compensation. The decreased intracellular sodium is responsible for increased compliance of the vessel wall and a reduced response to constrictor stimulus.

Advantages of thiazide as an antihypertensive:

• It can be given as once a day dose and has a flat dose-response curve.

• Decreased incidence of postural hypotension and relative freedom from CNS side effects

• No tolerance

• No fluid retention

• Decreased risk of osteoporotic fractures due to its hypocalciuric action

• Low cost

• Effectiveness in isolated systolic hypertension

Thiazides are ineffective both as antihypertensive and diuretic in patients with chronic renal failure.

High ceiling diuretics: Furosemide is the prototype drug in this class and it is a weaker antihypertensive agent than thiazides. They are given in hypertension only in the presence of the following complications:

• Chronic renal failure - As thiazides are ineffective

• Coexisting refractory congestive heart failure

• Resistance to combination regimen with thiazides

• Marked fluid retention with the use of potent vasodilators.

Potassium-sparing diuretics - They are only used in combination with thiazides to prevent the loss of potassium.

Side effects of diuretics include hypokalemia, carbohydrate intolerance, dyslipidemia, and hyperuricemia.

Solution to Question 3:

The dose of retigabine should be increased when phenytoin is added because phenytoin can decrease the plasma concentration of retigabine when given together.

Retigabine is also known as ezogabine. Metabolism of the drug occurs by glucuronidation and acetylation. Its half-life is about 7-11 hours and hence requires thrice-daily dosing. When it is administered with phenytoin or carbamazepine there may be a reduction in the plasma concentration of the drug. So the dose of retigabine should be increased when given along with these drugs.

It is a potassium(K+) channel opener that enhancer transmembrane potassium currents mediated by the KNCQ family of ion channels. It was approved as an adjunctive agent for focal onset seizures of patients above 18 years. But because of its adverse effects like retinal abnormalities and blue pigmentation, the production of the drug was discontinued by the manufacturer.

Other drug interactions of phenytoin:

It is an inducer of the enzymes CYP2C8/9, CYP3A4/5, and other CYPs. It is also an inhibitor of CYP2C9/19.

• Phenobarbitone can inhibit the metabolism of phenytoin. But the overall interaction is unpredictable as both the drug enhance each other's degradation by enzyme induction.

• Phenytoin and carbamazepine induce each other's metabolism.

• Valproate increases the level of unbound phenytoin by displacing the protein-bound form and decreasing its metabolism.

• Phenytoin increases the metabolism of steroids, oral contraceptives, doxycycline, theophylline, etc by enzyme induction.

• It produces competitive inhibition of warfarin metabolism. (Increased risk of bleeding when given together)

• Drugs like chloramphenicol, warfarin, isoniazid, and cimetidine can inhibit the metabolism of phenytoin and cause its toxicity.

• Sucralfate decreases the absorption of phenytoin by binding to it in the gastrointestinal tract.

Solution to Question 4:

Downregulation of CYP2C19 is responsible for the failure of clopidogrel in this patient.

Clopidogrel is a prodrug and its metabolic activation occurs in the liver. It is bioactivated by the CYP2C19 enzyme. If there is downregulation of this enzyme it will cause a decrease in the efficacy of clopidogrel and can lead to thrombotic events like stroke. Drugs that inhibit CYP2C19 can also decrease the antiplatelet effect of clopidogrel. Proton-pump inhibitors may cause inhibition of CYP2C19 and when they are given in combination there would be a decrease in the efficacy of clopidogrel.

Therapeutic uses: Clopidogrel is used to reduce the rate of stroke and myocardial infarction. It also decreases the rate of deaths in patients with recent myocardial infarction or stroke. Other uses are acute coronary syndrome and peripheral arterial disease.

Adverse effects: Increased risk of bleeding especially when given with an anticoagulant or aspirin. Rarely thrombotic thrombocytopenic purpura can occur.

CYP2C19: Common drugs that are metabolized include omeprazole, phenytoin, diazepam, and propanolol. Inducers of the CYP2C subfamily are rifampicin and carbamazepine. Omeprazole is an inhibitor.

Other options:

CYP2D6: Drugs metabolized include SSRIs, tricyclic antidepressants, antiarrhythmic drugs, beta-blockers, and opiates. It is inhibited by quinidine and it will prevent the formation of morphine from codeine. Hence quinidine decreases the analgesic effect of codeine.

CYP1A1/2: It metabolizes few drugs like theophylline, paracetamol, caffeine and carbamazepine. It plays an important role in the activation of procarcinogens. Inducers are rifampicin, carbamazepine, polycyclic hydrocarbons, charbroiled meat, and cigarette smoke.

CYP2E1: It is involved in the oxidation of alcohol, halothane, and minor metabolism of other drugs. It is important to note that it is involved in the formation of hepatotoxic N-acetyl benzoquinone imine from paracetamol. Chronic alcoholism is responsible for the induction of this enzyme.

Solution to Question 5:

Liraglutide is used in the long-term management of obesity.

Mechanism of action:

• Increase glucose-mediated insulin release

• Lowers the glucagon levels

• Delays gastric emptying

• Decreases feeding by stimulating GLP-1 receptors in the arcuate nucleus of the hypothalamus.

It is a GLP-1 analog used as a parenteral hypoglycemic agent and anti-obesity drug. It is administered subcutaneously. It is used as an adjuvant in patients who are not achieving glycemic control with oral agents. It can be also added with basal insulin.

It is found to decrease the risk of cardiovascular death, nonfatal myocardial infarction, and stroke in patients with type 2 diabetes and cardiovascular disease.

Adverse effects: Nausea, vomiting, diarrhea, or constipation. It should not be given in patients with a personal or family history of MEN syndrome or medullary carcinoma of the thyroid.

Other options:

Metformin: It belongs to the biguanides class of oral hypoglycemic agents. It is used as a first-line agent in the management of type 2 diabetes and it acts by reducing hepatic glucose production. It is also used to treat infertility in PCOD patients. IT is usually not associated with weight gain and in some cases, it may cause mild weight reduction.

Sibutramine: It has both serotonergic and noradrenergic action. It is also an anorectic agent with a similar mechanism of action to fenfluramine. But due to severe adverse effects, US-FDA has banned its use.

Fenfluramine: It is a serotonergic anorectic agent. It reduces food-seeking behavior due to increased serotonergic transmission in the hypothalamus. The use of is not recommended now by the US-FDA due to its severe side effects.

Solution to Question 6:

The clinical features and the CT angiography image showing diffuse symmetrical luminal narrowing of the lower limb vessels are suggestive of ergotism.

In this case, a patient on antiretroviral therapy is given ergotamine. HIV protease inhibitors used in ART like ritonavir, lopinavir, etc are potent inhibitors of the CYP3A4 enzyme. This enzyme is responsible for the metabolism of ergot compounds. Hence inhibition of this enzyme leads to decreased metabolism of ergotamine. This causes the accumulation of toxic levels of ergotamine in the patient and leads to ergotism.

Ergotamine is a partial agonist and antagonist at alpha-adrenergic, 5HT1; and 5HT2 receptors. It produces sustained vasoconstriction, vasomotor center depression, and visceral smooth muscle contraction.

Adverse effects: Nausea, vomiting, and abdominal pain may be seen. Vasoconstriction caused by the drug is responsible for most of the adverse effects. These are usually seen with long-term therapy with higher doses.

• It may precipitate or aggravate angina pectoris or myocardial infarction due to coronary insufficiency.

• Some patients develop cold, numb, and painful extremities.

• Leg claudication may be seen

• Gangrene of the extremities

Contraindications for ergotamine:

• Peripheral vascular disease

• Coronary artery disease

• Uncontrolled hypertension

• Sepsis

• Pregnancy

• Liver and kidney diseases

Solution to Question 7:

Topiramate is used in the treatment of Lennox-Gastuat syndrome.

Uses of topiramate:

• In newly diagnosed focal and primary generalized epilepsy in adults and children

• As monotherapy for refractory focal epilepsy and generalized tonic-clonic seizures

• Treatment of tonic-clonic seizures in patients with Lennox-Gastuat syndrome

• It is also used in the prophylaxis of migraine(not for treatment)

• It can be given as monotherapy for children of 10 years or more. But if used as adjunctive therapy it can be used in children as young as 2 years.

Mechanism of action: It reduces voltage-gated sodium currents in the cerebellar granule cells. It also activates a hyperpolarizing potassium current, enhances GABAA receptor currents, and limits the activation of the AMPA-kainate subtype of glutamate receptors. It also produces weak inhibition of the carbonic anhydrase enzyme.

Pharmacokinetics: Only 10-20% is bound to plasma proteins and the drug is largely excreted unchanged in the urine. Half-life is about 1 day. Topiramate is found to decrease the plasma concentrations of oral contraceptives, hence a higher dose of OCPs should be given along with topiramate.

Adverse effects:

• Somnolence, weight loss, fatigue, and nervousness are commonly seen

• It can precipitate renal calculi. It may be due to the weak carbonic anhydrase inhibition.

• Cognitive impairment

Solution to Question 8:

Ciraparantag is a reversal agent for all direct anticoagulants and is not useful with fondaparinux which is an indirect factor Xa inhibitor.

Ciraparantag, andexanet alfa, and idarucizumab are all reversal agents of direct anticoagulants.

Ciraparantag: It is considered as a potential reversal agent against all direct anticoagulants including dabigatran, rivaroxaban, edoxaban, apixaban, LMWH, and heparin. This drug is in the early stages of development.

Andexanet alfa: It is a recombinant analog of factor Xa and is proposed for reversal of oral factor Xa inhibitors like rivaroxaban, edoxaban, and apixaban. It rapidly reverses the action of these drugs and restores thrombin generation.

Idarucizumab: It is a licensed monoclonal antibody that specifically acts as a reversal agent for dabigatran. It has a 350 times higher affinity for dabigatran than thrombin. Hence it forms a complex with dabigatran and is excreted by the kidneys.

If a patient on direct oral anticoagulants like dabigatran, rivaroxaban, edoxaban, and apixaban presents with life-threatening bleeds and the reversal agents are not available then other options are the administration of prothrombin complex concentrate, activated prothrombin complex concentrate, or recombinant factor VII-A. Hemodialysis is useful in severe bleeding due to dabigatran.

Other reversal agents for anticoagulants:

• Protamine sulphate - Heparin

• Vitamin K - Warfarin

Note:

Andexanet alfa has received US-FDA approval in May 2018 as a reversal agent for rivaroxaban and apixaban.

Solution to Question 9:

Inhaled nitric oxide (NO) is used in the management of pulmonary hypertension.

The only FDA-approved therapeutic indication of inhaled nitric oxide is in persistent pulmonary hypertension of the newborn. It has the ability to selectively dilate pulmonary vasculature. Hence it can be used in conditions with increased pulmonary vascular resistance.

It is proposed for the:

• Long-term management of pulmonary artery hypertension(PAH) and other forms of pulmonary hypertension

• Rescue therapy for severely hypoxemic patients with or without pulmonary hypertension

Diagnostic indications:

• In heart failure patients and infants with congenital heart disease during cardiac catheterization to evaluate pulmonary vasoreactivity.

• In determining the diffusion capacity across the alveolar-capillary unit. It is considered superior to carbon dioxide in this because it has a greater affinity for hemoglobin and higher water solubility at body temperature.

Mechanism of action:

It activates soluble guanylyl cyclase and produces cGMP, which then activates cGMP-dependent protein kinase. This leads to the activation of multiple regulatory mechanisms that decrease intracellular calcium and cause vasodilation in precapillary resistance arterioles.

Adverse effects:

• Methemoglobinemia at higher concentrations.

• It can worsen left ventricular performance in patients with an already impaired left ventricular function.

Solution to Question 10:

Nivolumab is used as a checkpoint inhibitor in Hodgkin's lymphoma.

FDA approved uses of nivolumab:

• Advanced melanoma

• Advanced non-small cell lung cancer(NSCLC) that are previously treated

• Renal cell carcinoma

• Advanced head and neck cancers

• Refractory/relapsed classical Hodgkin's lymphoma

Nivolumab is a checkpoint inhibitor that blocks the interaction between PD-1 and its ligands. It is a fully human monoclonal IgG4 antibody. It has a good response rate in patients with pretreated solid tumors and hematological malignancies. Similar effectiveness is also seen with pembrolizumab which is also an anti-PD-1 monoclonal antibody.

Adverse effects:

• In melanoma patients rash is a commonly seen side effect

• fatigue, dyspnea, musculoskeletal pain, cough, decreased appetite, and constipation can be seen in advanced squamous NSCLC patients

• Pneumonitis, colitis, nephritis, hepatitis, and renal dysfunction are also seen

Important note:

The FDA on May 2021 has approved the use of nivolumab in patients having completely resected esophageal or gastroesophageal junction cancers with residual pathologic disease who have received neoadjuvant chemoradiotherapy.

Solution to Question 11:

Paradoxical bronchoconstriction with ipratropium bromide is seen with a hypotonic nebulizer solution and not hypertonic saline.

Paradoxical bronchoconstriction with ipratropium bromide is seen especially when given by a nebulizer and may be due to the following factors:

• Hypotonic nebulizer solution

• Antibacterial additives like benzalkonium chloride and EDTA

• It may be also due to prejunctional M2 receptor blockade on the cholinergic nerves present in the airway. These prejunctional M2 receptors normally inhibit acetylcholine release into the synapse. So their blockade can cause increased Ach release and bronchoconstriction.

Paradoxical bronchoconstriction is not reported with tiotropium bromide or any other long-acting muscarinic antagonists. Tiotropium is selective for M3 and M1 receptors and therefore less likely

to produce prejunctional M2 receptor blockade.

Other adverse effects of inhaled muscarinic antagonists:

• A bitter taste is seen with ipratropium

• Nebulization with ipratropium bromide may precipitate glaucoma in elderly patients. This is due to the direct effect of the drug on the eyes and can be easily prevented by using a mouthpiece instead of a facemask.

• Long-acting agents can cause dryness of the mouth.

• In elderly patients, urinary retention may be seen occasionally.

Solution to Question 12:

Gonorrhea is not managed by antimicrobial combinations. According to the latest CDC guidelines, single-agent therapy with ceftriaxone is the preferred treatment for gonococcal infections.

Single-dose of 500mg intramuscular(IM) ceftriaxone is given for gonorrhea. It is important to note that the previous guidelines that recommended dual therapy with ceftriaxone and azithromycin are no longer preferred.

Monotherapy with ceftriaxone is given when concurrent chlamydia infection is ruled out by molecular testing. If this is not done then presumptive treatment for chlamydia should be given along with the treatment of gonorrhea. According to CDC guidelines, 100mg doxycycline twice daily for 7 days is given as presumptive treatment of chlamydia in nonpregnant individuals.

Other options:

Tuberculosis: Multidrug therapy is given in tuberculosis for proper cure and to prevent the selection of drug-resistant mutants. ATT regimen is given in detail in the pearl attached.

Intraabdominal infections: It is usually polymicrobial and can result in an intraabdominal abscess or secondary peritonitis. These are managed with either single-agent or combination therapy. Broad-spectrum agents with gram-negative cover such as meropenem or piperacillin-tazobactam can be used alone. Combination therapy with metronidazole and other agents like ciprofloxacin, cefepime, etc are used as empirical agents in intraabdominal infections.

Malaria: It is also treated with antimicrobial combinations. In plasmodium vivax and ovale malaria, schizontocides like chloroquine are combined with agents that attack the hypnozoites like primaquine for radical cure. Primaquine is also given in combination with Artemisinin combined therapy(ACT) for plasmodium falciparum infections as a gametocidal agent. Treatment regimens are given in the pearl attached.

Solution to Question 13:

Vancomycin acts on step 4 in the given image.

Mechanism of action of vancomycin: It is a cell wall synthesis inhibitor. It binds to the D-Ala-D-Ala terminus of the newly formed peptidoglycan pentapeptide and inhibits the transglycosylase enzyme. This would prevent further elongation of the peptidoglycan and its cross-linking.

Vancomycin is a bactericidal agent for gram-positive organisms. It is also active against many gram-positive anaerobes like Clostridium difficile. Resistance against vancomycin in enterococci is due to the alternation in its binding site where D-Ala is replaced by D-Lactate.

Clinical uses:

• Parenteral vancomycin is given for bloodstream infections and endocarditis caused by MRSA.

• Vancomycin plus gentamicin is given as an alternative regimen for enterococcal endocarditis with penicillin allergy.

• In combination with other agents for suspected meningitis or meningitis due to penicillin-resistant pneumococcus.

• Oral vancomycin is given for Clostridium difficile colitis.

Adverse effects:

• Injection site phlebitis

• Nephrotoxicity

• "Redman" syndrome due to histamine release following infusion

Other options:

Fosfomycin inhibits step 1 in the image. It acts at a very early stage in cell wall synthesis and inhibits the cytoplasmic enzyme enol pyruvate transferase. This would block the addition of phosphoenolpyruvate to UDP-N-acetylglucosamine. This prevents the formation of UDP-N-acetylmuramic acid which is the precursor N-acetylmuramic acid. N-acetylmuramic acid is a component of the bacterial cell wall and hence cell wall synthesis is inhibited.

Cycloserine inhibits step 2 in the image. It is a structural analog of D-alanine and inhibits the enzymes alanine racemase that converts L-alanine to D-alanine and d-alanyl-d-alanine ligase. This would thereby inhibit the incorporation of D-alanine into the peptidoglycan pentapeptide. It is mainly used in the management of tuberculosis where there is resistance to the first-line agents.

Bacitracin inhibits step 3 in the image. It is a cell wall synthesis inhibitor that interferes with dephosphorylation in the cycling of the lipid carrier which transfers peptidoglycan subunits to the growing cell wall. It is only given for topical application because of its severe nephrotoxicity when given systemically.

Note:

Beta-lactam antibiotics prevent cross-linking by inhibiting the transpeptidases. (Last step in the images)

Solution to Question 14:

In this case, the half-life remains the same, and the plasma concentration doubles.

Half-life is constant in drugs following first-order kinetics irrespective of the dose as a constant fraction of the drug is eliminated per unit time.

The plasma concentration of the drug increases when the dose given is increased. In this case, as the drug dose is doubled plasma concentration is also doubled.

Solution to Question 15:

Risperidone is FDA approved for the treatment of bipolar affective disorders.

Uses of risperidone:

• Long-term management of psychosis and schizophrenia.

• Maintenance therapy in mania bipolar disorders.

• Management of irritability associated with autism.

Off-label uses of risperidone:

• Management of psychotic symptoms in delirium and dementia.

• Obsessive-compulsive disorders.

• Post-traumatic stress disorders.

Risperidone is an atypical antipsychotic that shows both D2 and 5 HT2 receptor blockade. It also has an affinity for alpha 1, alpha 2, and H1 receptors. It produces a more potent D2 blockade than clozapine. Hyperprolactinemia is an important adverse effect. It is also associated with an increased risk of stroke in elderly patients. At low doses, extrapyramidal side effects are less.

Solution to Question 16:

The half-life of the given drug is 13.8 hours.

The half-life of a drug or t1/2 is the time taken for the plasma concentration to be reduced by 50%. The elimination rate constant or k value of a drug reflects the fraction of drug removed from the body compartment per unit of time.

The half-life of a drug and its elimination rate constant are inversely proportional to each other. They are represented by the equation:

k t1/2 = 0.693

Therefore, t1/2 = 0.693/ k

The given value for elimination rate constant ■ is 0.05, therefore the half-life of the given drug would be:

0.693/ 0.05 = 13.8

Therefore, the half-life of the drug would be 13.8 hours.

Solution to Question 17:

The minimum duration of time between the consumption of phenytoin and sucralfate should be 120 minutes.

Sucralfate, in an acidic environment (pH < 4) acts by producing a viscous, sticky polymer that adheres to ulcer craters and epithelial cells for up to 6 hours after a single dose. This viscous layer produced in the stomach leads to inhibition of absorption of other drugs, like phenytoin, digoxin, cimetidine, ketoconazole, and fluoroquinolone antibiotics. Therefore, sucralfate should be taken after a gap of at least 2 hours / 120 minutes after the administration of phenytoin.

Constipation is the most common side effect of sucralfate. It should be avoided in patients with renal failure who are at risk of aluminium overload. Aluminum-containing antacids should not be combined with sucralfate in these patients. As the viscous gel produced by sucralfate is sticky, it may eventually lead to bezoar formation.

It is generally used for prophylaxis of peptic ulcers, but rarely so because of its side effects and its risks outweigh its benefits.

Solution to Question 18:

Category A, B, C, D, X in drug classification are indicative of their safety in pregnancy.

The united states food and drug administration (FDA) system for teratogenic potential attempts to quantify teratogenic risk from category A (safe) to category X (definite human teratogenic risk).

Category A: The possibility of fetal harm is remote and studies in women have failed to demonstrate any evidence of risk to the fetus. Examples: doxylamine, folic acid, levothyroxine.

Category B: Either animal-reproduction studies have not demonstrated a fetal risk, but there are no controlled studies in pregnant women, or animal-reproduction studies have shown an adverse effect that was not confirmed in controlled studies in pregnant women. Example: Amoxicillin, loratadine, ondansetron.

Category C: Either studies in animals have revealed adverse effects on the fetus, that is, teratogenic or embryocidal but there are no controlled studies in women, or studies in women and animals are not available. These drugs should be given only if the potential benefit justifies the potential risk to the fetus. Example: fluconazole, metoprolol, sertraline.

Category D: Though there is evidence of human fetal risk, the benefits from use in pregnant women may be acceptable despite the risk, for example, if the drug is needed in a life-threatening situation or for a disease for which safer drugs cannot be used or are ineffective. Example: lisinopril, lithium, phenytoin.

Category X: There is evidence of fetal risk based on human or animal experience and the risk of the use of the drug in pregnant women clearly outweighs any possible benefit. The drug is contraindicated in women who are or may become pregnant. Example: methotrexate, simvastatin, warfarin.

Solution to Question 19:

The false statement regarding suicidal inhibition is the inhibitor can bind with any site resulting in suicidal inhibition.

In suicidal inhibition, the inhibitor can only bind with the active site of an enzyme and is unreactive outside the confines of the enzyme's active site.

The enzyme inhibition helps to modify the rate of reactions it catalyzes and this inhibition can either be reversible or irreversible. Reversible inhibitors bind to enzymes through non-covalent bonds. Irreversible inhibitors usually bind to an enzyme through covalent bonds and do so strongly so that it can no longer dissociate.

Suicide inhibitors are a category of irreversible inhibitors also called mechanism-based inhibitors. They are specialized substrate analogs that contain a chemical group that reacts with the catalytic machinery of the target enzyme. In these types of inhibitors, the substrate analog irreversibly ties up with the binding site of an enzyme, deceiving the enzyme by concealing its functional group resulting in the formation of a highly reactive molecule, which brings about self-inactivation of the enzyme.

Suicide inhibitors bind to the enzyme by means of covalent bonds; they are enzyme-specific and unreactive outside the confines of the enzyme’s active site.

An example of this would be the binding of clavulanic acid to the active site of beta-lactamases in bacteria. Clavulanic acid mimics the structure of a beta-lactam antibiotic and forms a covalent adduct with the β-lactamase active site. This leads to a rearrangement that creates a much more reactive derivative, which is subsequently attacked in its active site to irreversibly acylate the enzyme and inactivate it. Thus, amoxicillin and clavulanic acid are combined to overcome bacterial resistance to amoxicillin alone.

The development of clavulanic acid to target beta-lactamase-producing bacteria is also an example of the rational design of drugs. Rational design of drugs refers to the ability to predict the appropriate molecular structure of a drug on the basis of information about its biological receptor.

Solution to Question 20:

The dose at which dopamine acts on β-1 receptors is 2-10 mcg/kg/min.

The systemic effects of dopamine vary with its concentration in the body.

When administered at an infusion rate of 2-5 mcg/kg/min, it directly stimulates cardiac β receptors. This enhances myocardial contractility and is useful in the management of cardiogenic shock.

At doses of 5-15 mcg/kg/min and above, it stimulates α adrenergic receptors. This causes peripheral arterial and venous vasoconstriction.

At lower doses of <2 mcg/kg/min, it directly stimulates both D1 and D2 receptors. The D1 receptors in renal and mesenteric blood vessels are the most sensitive: i.v. infusion of low dose of DA dilates these vessels subsequently, causing increased renal blood flow and increased urine output.

The clinical indications of dopamine include cardiogenic and vasodilatory shock, congestive cardiac failure and symptomatic bradycardia unresponsive to atropine or pacing.

The major side effects are:

• Severe hypertension (especially when taken with nonselective β-blockers)

• Ventricular arrhythmias

• Cardiac ischemia

• Tissue ischemia/gangrene (high doses/extravasation into tissues)

Solution to Question 21:

The incorrect statement regarding clomiphene citrate is that it causes monofollicular development; rather, it causes multi follicular development.

Clomiphene is an unequal mixture of two isomers, enclomiphene and zuclomiphene. Zuclomiphene is a non-steroidal compound closely resembling an estrogen, blocks the negative inhibitory feedback of estrogens on the hypothalamus, causing a surge of gonadotropins (FSH, LH), which leads to ovulation. The continued flow of FSH encourages multiple follicular development which is relatively common.

Being an anti-estrogen, it causes thickening of cervical mucus and thinning of the endometrium.

Clomiphene citrate (CC) is used for the treatment of absent or irregular ovulation due to hypothalamic-pituitary dysfunction associated with normal levels of endogenous estradiol; the patient must be treated repeatedly until pregnancy is achieved since the normal cyclic ovulatory function does not usually resume. Clomiphene citrate has no effect on patients with ovarian or pituitary failure. Ovulation induction with clomiphene citrate and a gonadotropin like FSH increases the likelihood of multiple concurrent ovulations.

Side effects of CC include hot flushes and some nausea. However, it is usually very well tolerated.

Solution to Question 22:

The mechanism of action of remdesivir is inhibition of the enzyme RNA-dependent RNA polymerase.

Remdesivir is an adenosine nucleotide prodrug. It distributes into cells where it is metabolized to form remdesivir triphosphate (RDV-TP). It binds to the viral RNA-dependent RNA polymerase and inhibits viral replication by prematurely terminating RNA transcription.

Remdesivir (under Emergency Use Authorization) is used in the treatment of COVID-19. It may be considered in patients with moderate to severe disease, requiring supplemental oxygen, within 10 days of onset of the illness.

Dose: 200 mg IV on day 1 followed by 100 mg IV daily for 4 days (total 5 days).

The following are contraindications to the use of remdesivir:

• Renal or hepatic dysfunction (eGFR <30 ml/min/m2; AST/ALT >5 times upper limit of normal) (Not an absolute contraindication)

• Pregnant or lactating females

• Patients <12 years of age

• Patients who are not on oxygen support or in-home settings

Remdesivir can cause gastrointestinal symptoms, elevated transaminase levels, increased prothrombin time, and hypersensitivity reactions. Hence, liver function tests and prothrombin time should be obtained before and during treatment with remdesivir.

Other options:

Option A- RNA-dependent DNA Polymerase / reverse transcriptase inhibitors prevent the conversion of viral RNA into proviral DNA. They help to prevent infection of susceptible cells but do not eradicate the virus from cells that are integrated with proviral DNA. Examples: zidovudine, lamivudine, stavudine, tenofovir, didanosine, zalcitabine, and emtricitabine. These are drugs used to manage HIV infection.

Options C and D- Entry/fusion inhibitors include enfuvirtide and maraviroc. Enfuvirtide inhibits the fusion of viral and cell membranes mediated by gp41 and CD4 interactions. Maraviroc is a chemokine receptor antagonist and binds to host cell CCR5 receptor to block binding of viral gp120. They are also used to manage HIV infection.

Solution to Question 23:

Administration of ritonavir would increase the bioavailability of saquinavir.

Saquinavir is an HIV protease inhibitor that inhibits replication of both HIV-1 and HIV-2. It is metabolized predominantly by the hepatic enzyme CYP3A4. Inhibition of CYP3A4 results in an increase or a boost in the concentration of saquinavir. Ritonavir is one such drug that inhibits CYP3A4. Therefore combining it with a protease inhibitor like saquinavir results in:

• Enhanced bioavailability

• Inhibition of systemic clearance

• Inhibition of first-pass metabolism, and

• Longer half-life of the co-administered HIV protease inhibitor.

This allows for a reduction of drug dose and dosing frequency of saquinavir.

All protease inhibitors except for nelfinavir are predominantly metabolized by CYP3A4. Therefore, the administration of CYP3A4 inhibitors like ritonavir or cobicistat can increase the bioavailability of these protease inhibitors. This phenomenon is known as boosting protease inhibitors by CYP inhibitors. The dosage of ritonavir, when given as a booster drug, is 100 to 200mg once or twice daily.

Other options :

Option B: Vitamin C / Ascorbic acid is a water-soluble vitamin that can actually reduce the steady-state concentration of protease inhibitors. It increases the absorption of dietary iron.

Option C: Cimetidine is a histamine 2 receptor antagonist which is used to promote gastric and duodenal ulcer healing. It is used in the management of gastroesophageal reflux disease. It also inhibits CYPs, but only ritonavir and cobicistat are used for boosting protease inhibitors.

Option D: Ganciclovir is a guanine nucleoside analogue which has inhibitory activity against all herpes viruses, especially cytomegalovirus (CMV).

Solution to Question 24:

Drug A reduces the first-pass metabolism of Drug B.

The area under the plot of plasma concentration of a drug versus time after dosage is called the area under the curve or AUC. It can be used as a measure of the bioavailability of a drug and to calculate its clearance rate from the body.

Bioavailability refers to the extent to which an administered dose of the drug reaches its site of action; if the metabolic and excretory capacity of the intestine or liver for the drug is large, bioavailability is reduced substantially and this is called the first-pass metabolism whereas bioavailability is increased if intestinal absorption of the drug is increased.

The greater the AUC, the greater is the bioavailability of the drug.

In the question, is observed that when drug A was given along with drug B, the total AUC has increased (820 seconds ± 124 seconds) when compared to the AUC when drug B is administered alone (570 seconds ± 127 seconds).

The null hypothesis for the above question would be that drug A doesn't increase the AUC for drug B and for this to be true, ■ value should be >0.05.

As given ■ value is <0.01, it is significant and the null hypothesis is rejected, which means that drug A has indeed increased the AUC for drug B.

This indicates increased bioavailability of drug B when administered with drug A, which shows that drug A has either increased the intestinal absorption of drug B or has reduced its first-pass metabolism.

Other options:

Option A: If drug A decreased the intestinal absorption of drug B, this would mean that the AUC for drug B, when administered with drug A, would be reduced. This is due to decreased bioavailability of drug B when given with drug A.

Option C: If drug A increases the metabolism of drug B, this would lead to a shorter t1/2 and duration of action of drug B, thereby decreasing the AUC for when drug B and drug A are given together.

Option D: If drug A increased the elimination of drug B, the AUC for drug B when given with drug A would be reduced due to reduced duration of action of drug B.

Solution to Question 25:

The mechanism of action of cyclosporine is inhibition of calcineurin activity.

Calcineurin is a calcium/calmodulin-dependent phosphatase required for the induction of T cell lymphokine production and proliferation. Cyclosporine is an immunomodulatory drug

that inhibits calcineurin activity by forming a complex with cyclophilin, a cytoplasmic-receptor protein present in target cells, to form a cyclosporine-cyclophilin complex. This drug-receptor complex specifically and competitively binds to and inhibits calcineurin, a calcium and calmodulin-dependent phosphatase. This in turn, inhibits the translocation of a family of transcription factors (NF-AT), leading to reduced transcriptional activation of cytokine genes for interleukin (IL)-2, tumour necrosis factor (TNF)-alpha, IL-3, IL-4, CD40L, granulocyte-macrophage colony-stimulating factor, and interferon-gamma. Ultimately, the proliferation of T lymphocytes is reduced, resulting in immunomodulation.

Its uses are :

• In solid organ transplantation, to treat organ rejection in kidney, liver, and heart allogeneic transplants.

• In patients with rheumatoid arthritis, when the disease has not adequately responded to methotrexate.

• It is approved to treat amyotrophic lateral sclerosis (ALS or Lou Gehrig disease) and its variants.

• In nephrotic syndrome, to treat focal segmental glomerulosclerosis not responding to corticosteroids.

Adverse effects of cyclosporine include a decrease in glomerular filtration rate (GFR) and metabolic disturbances like hypomagnesemia, hyperkalemia, dyslipidemia, gynecomastia, hypertrichosis; neurotoxicity in the form of convulsions. There is an increased risk of developing skin and lymphoproliferative malignancies in cyclosporine-treated psoriasis patients.

Other options:

Option B: AMP kinase is stimulated by metformin, a first-line orally administered drug in the management of type-2 diabetes mellitus.

Option C: IMP dehydrogenase is inhibited reversibly by the drug mycophenolate mofetil, which is indicated in the prophylaxis of transplant rejection.

Option D: Dihydro-orotate dehydrogenase is inhibited by leflunomide, a pyrimidine synthesis inhibitor indicated for the management of rheumatoid arthritis in adults.

Solution to Question 26:

Anionic and weakly acidic drugs bind to albumin.

After being distributed in the circulating blood, drugs bind to serum proteins to varying degrees. Among serum proteins, albumin and alpha-1 acid glycoprotein play important roles in protein binding for many drugs, which affects their distribution in the body.

Albumin is a major carrier of and binds to acidic drugs like phenytoin, NSAIDs. Whereas, α-1 acid glycoprotein binds basic drugs like lignocaine,disopyramide, quinidine and propranolol.

Importance of plasma protein binding:

Conditions resulting in acute phase reaction response leading to elevated levels of alpha-1 acid glycoprotein like Crohn's disease, myocardial infarction, arthritis, cancer, etc. enhance binding of basic drugs. Conditions causing hypoalbuminemia result in decreased binding sites for acidic

drugs.

When a drug is bound to the plasma proteins, it is kept in the blood, while the unbound (free) drug can produce a therapeutic effect, be metabolized, or excreted. In the majority of the drugs, this does not have a significant effect, but in drugs with a narrow therapeutic index like warfarin, it is clinically relevant as excessive free warfarin can produce toxicity.

Many drugs displace other drugs from their protein binding sites. This may cause an increase in the pharmacological effect of the displaced drug.

A drug bound to a plasma protein cannot cross membranes and this bound fraction of a drug is also not available for metabolism or glomerular filtration which can make the drug long-acting.

Other options:

Option B- Globulins are of three types - α,β, and γ-globulins. Globulins bind glucocorticoids and progesterone in the blood.

Option C- Lipoproteins like VLDL, LDL can bind anti-cancer drugs like 5-fluorouracil and doxorubicin.

Solution to Question 27:

The probable reason why the drug to be administered was changed from ampicillin to ceftriaxone is because Hemophilus produces beta-lactamase and hence, is resistant to ampicillin.

A considerable percentage of isolates of Hemophilus influenzae produce beta-lactamase, rendering them resistant to ampicillin, a beta-lactam antibiotic. However, these strains are still sensitive to second and third-generation cephalosporins.

In H. influenzae, the resistance to ampicillin and other β-lactam antibiotics is generally due to the production of a β-lactamase. In the case of β-lactamase-negative ampicillin-resistant strains, resistance is due to the presence of altered penicillin-binding proteins.

Ceftriaxone is a third-generation cephalosporin that is effective against Hemophilus Influenzae and considerably resistant to beta-lactamase. It has a very long half-life which permits once-daily dosing of the drug. Because of the efficacy of third-generation cephalosporins in the therapy of meningitis caused by Hemophilus Influenzae type b, ceftriaxone administered at the rate of 50 mg/kg/dose given 12th hourly should be a part of the initial empirical therapy.

Vancomycin combined with a third-generation cephalosporin like ceftriaxone is the recommended empirical antibiotic regimen in a suspected case of meningitis, beyond the neonatal period.

Solution to Question 28:

The preferred diuretic for this patient with cirrhosis presenting with abdominal distension and lower limb swelling is eplerenone.

Eplerenone is a mineralocorticoid receptor/aldosterone antagonist. High doses of mineralocorticoid receptor/aldosterone antagonists decrease total body sodium in cirrhotic patients with ascites by natriuresis.

Patients with liver cirrhosis have a decreased effective arterial blood volume because of splanchnic vasodilatation. This stimulates the renin-angiotensin-aldosterone system and results in hyperaldosteronism. This causes increased sodium and water retention and leads to the development of ascites. This also contributes to resistance to loop diuretics.

Normally, aldosterone (AL) enters the cells of the late distal tubule and collecting ducts by diffusion and binds to a specific mineralocorticoid receptor (MR) in the cytoplasm. This AL-MR complex mediates Na+ reabsorption, which drives the secretion of K+ and H+into the tubules.

Eplerenone competitively blocks this MR, preventing it from binding with aldosterone. The net effect is sodium diuresis and potassium retention. It is a potassium-sparing diuretic. Other mineralocorticoid receptor antagonists include spironolactone, canrenone, potassium canrenoate.

Some clinical uses of MR antagonists are -

• Hypertension due to primary hyperaldosteronism

• Edema associated with liver cirrhosis, congestive cardiac failure, chronic kidney disease

• They may reduce mortality in patients with myocardial infarction

Hyperkalemia is the major side effect of MR antagonists, the risk being greater in patients with renal disease or those already taking ACE inhibitors. Other important side effects are gynecomastia and decreased libido due to the anti-androgenic effect of spironolactone but eplerenone does not cause gynecomastia and decreased libido.

Other options :

Option A- Triamterene directly inhibits renal epithelial sodium channels (ENaC) in the late distal tubule and collecting duct. It is a potassium-sparing diuretic and is generally used along with other diuretics in the treatment of hypertension and edema. Triamterene is metabolized in the liver, hence triamterene toxicity may be enhanced in patients with hepatic disease. It is used in the management of Liddle syndrome.

Option C- Chlorthalidone is a thiazide-like diuretic. It inhibits Na+-Cl- symporter in the early distal tubule and increases the excretion of sodium and chloride. Thiazide diuretics may be used for the treatment of hypertension, heart failure, nephrolithiasis due to idiopathic hypercalciuria, and nephrogenic diabetes insipidus.

Option D- Furosemide is a loop diuretic. It acts by inhibiting the Na+-K+-2Cl- symporter in the thick ascending limb of the loop of Henle resulting in increased excretion of sodium and chloride. It also has mild carbonic anhydrase inhibitory activity and also increases urinary excretion of bicarbonate and phosphate. Cirrhotic patients are often resistant to loop diuretics due to decreased secretion of the drug into tubular fluid and high aldosterone levels. Furosemide may be given along with eplerenone, but the best answer here is eplerenone since aldosterone antagonists are the main therapy for cirrhotic edema.

Solution to Question 29:

Varenicline is the most effective drug in smoking cessation.

Nicotine is the principal constituent of tobacco responsible for its addictive behaviour. One of the newest agents introduced to aid in smoking cessation is varenicline.

Varenicline works by blocking the effects of nicotine on the brain. It blocks α-4, β-2 nicotinic acetylcholine receptor subtypes and varenicline inhibits dopaminergic activation produced by smoking through their partial agonism, and decreases the craving and withdrawal syndrome that occurs with cessation attempts. It prevents nicotine stimulation of the mesolimbic dopamine system associated with nicotine addiction, thus helping to occlude the "rewarding" effect of smoking.

Varenicline is prescribed only to patients 18 years or older. Varenicline is taken orally and the dose taken is 0.5 mg once daily on days 1 to 3, increased to 0.5 mg twice daily on days 4 through 7 and 1 mg twice daily after day 8 for 11 weeks. Therapy should start one week before the target quit date. It can be continued for another 12 weeks (1 mg BD) to maintain abstinence.

The adverse effects include nausea, which can be mitigated by starting at lower doses and gradually increasing the dose. Other symptoms include disturbed sleep, sleepwalking, agitation, drowsiness, and constipation and may also increase the risk of pancreatitis. Varenicline has also been implicated in specific skin syndromes such as Stevens-Johnson syndrome, erythema multiforme, and photosensitivity. Varenicline can cause renal failure and kidney stones, thus renal function must be monitored.

The FDA has issued a warning about mood and behavioral changes associated with its use, and there is some evidence of increased cardiovascular risk.

Other options:

Option B: Buspirone sustained release is a partial agonist at 5HT receptors but affinity for D2 receptors may also be present. Its effects are slow onset, over a couple of weeks and is used in the management of generalised anxiety states.

Option C: Nicotine gum has nicotine which is an agonist at Nn type nicotinic cholinergic receptors and is also advised for smoking cessation along with psychotherapy. However, varenicline was more effective when compared with single■product nicotine replacement therapy (NRT) and was similarly effective compared with combination NRT.

Option D: Rimonabant is an inverse agonist at cannabinoid receptor CB1 which decreases neurotransmitter release at GABAergic and glutamatergic synapses; its usage for smoking cessation and treat obesity was stopped after serious side effects notably severe depression and increased risk of suicide. However, it continues to be used off-label for smoking cessation.

Solution to Question 30:

The cephalosporin antibiotic most likely to have caused hemarthrosis in this patient on warfarin therapy is cefoperazone.

Cephalosporins containing a methylthiotetrazole (NMTT) group can prolong the prothrombin time. This results in clinically significant bleeding in patients already on anticoagulant therapy or having vitamin K deficiency.

This is because NMTT cephalosporins inhibit the enzyme vitamin K epoxide reductase in the liver, which is the key enzyme for vitamin K-dependent carboxylation, a posttranslational modification that is essential for the biological functions of coagulation factors.

NMTT group containing cephalosporins include latamoxef, cefmenoxime, cefoperazone, cefotetan, cefamandole.

Vitamin K epoxide reductase is also the target of anticoagulants like warfarin. Oral administration of vitamin K can prevent this problem.

Solution to Question 31:

Vilazodone is the novel antidepressant.

The antidepressant effect of vilazodone is due to its enhancement of serotonergic activity in the CNS through selective inhibition of serotonin reuptake. It binds with high affinity to the serotonin reuptake site but not to the norepinephrine or dopamine reuptake site. As a result, vilazodone potently and selectively inhibits the reuptake of serotonin.

Vilazodone is also a partial agonist at serotonergic 5-HT1A receptors; the result of this activity on serotonergic transmission also plays a role in its antidepressant effect.

Thus it offers a novel combination of selective serotonin reuptake inhibition (SSRI) and serotonergic (5-HT1A) receptor partial agonist activity. Because of these characteristics, vilazodone has been termed a serotonin partial agonist–reuptake inhibitor (SPARI).

Vilazodone is never combined with a monoamine oxidase (MAO) inhibitor or given within 14 days of discontinuing or initiating MAO inhibitor therapy due to the serious risk of serotonin toxicity.

Other options:

Option A: Lurasidone is a second-generation antipsychotic that is a 5-HT2A and D2 receptor antagonist. it is used in the treatment of schizophrenia and bipolar depression (as monotherapy or as an adjunct).

Option C: Asenapine is a second-generation antipsychotic which is also a 5-HT2A and D2 receptor antagonist, used in the management of schizophrenia and acute mania.

Option D: Blonanserin is a second-generation antipsychotic, which acts as an antagonist at D2, D3, and 5-HT2A receptors. Blonanserin has a low affinity for 5-HT2C, adrenergic α1, histamine H1, and muscarinic M1 receptors, but displays a relatively high affinity for 5-HT6 receptors. It is used in the management of schizophrenia.

Solution to Question 32:

The image shows grey-blue discoloration over this patient's face and the drug responsible for this is amiodarone.

Long-term treatment with amiodarone is responsible for skin pigmentation. Elderly men are at an increased risk of developing this.

Ultraviolet light reacts with amiodarone and its metabolites to cause local vasodilatation. This increases the diffusion of amiodarone and its metabolites, resulting in their deposition in the tissues, causing pigmentation. Therefore, it is usually seen in sun-exposed areas.

Cessation of amiodarone, avoiding exposure to sunlight using sunscreen cream is recommended.

Amiodarone is an iodine-containing compound that is structurally related to thyroid hormone. It is a class III antiarrhythmic drug, which increases the duration of an action potential by blocking potassium channels. It thereby increases the time taken for repolarization. It also prolongs the refractory period and suppresses abnormal automaticity. It also has weak calcium channel blocking, sodium channel blocking, and adrenergic blocking actions.

It is used to maintain normal sinus rhythm during atrial fibrillation and prevent recurrent ventricular tachycardia.

Adverse effects of amiodarone:

• Prolonged usage of amiodarone causes interstitial lung disease.

• Heart blocks, bradycardia and QT prolongation

• Hyperthyroidism or hypothyroidism

• Corneal micro-deposits and vortex keratopathy, optic neuropathy

Drug interactions of amiodarone

• Causes digoxin toxicity by decreasing its renal clearance.

• Increases concentration of quinidine and procainamide

• Potentiates the anticoagulant effect of warfarin when given simultaneously

Solution to Question 33:

Levopropoxyphene is a centrally acting opioid antitussive.

Levopropoxyphene is an opioid derivative that is a strong and partial μ receptor agonist that helps to reduce the cough reflex and therefore, is used in the management of acute debilitating cough. The usual antitussive dose is 50–100 mg every 4 hours. It has no analgesic activity. However, it is known to cause some sedation.

Other options:

Option A: Ambroxol is derived from bromhexine. It is a mucoactive agent that helps alter the viscoelastic properties of mucus and increase mucociliary clearance, thereby providing relief in productive cough. It is commonly available as intramuscular or intravenous, lozenges, syrups, or tablets.

Option B: Diphenoxylate is an opioid agonist that is structurally similar to meperidine. It is used in the treatment of acute, chronic, and traveler's diarrhea. Upon administration, it undergoes de-esterification to difenoxin, which is its active metabolite. Its side effects include constipation, toxic megacolon, paralytic ileus and can give rise to CNS depression in children.

Option D: Levorphanol is a synthetic opioid that is also a 5HT/Norepinephrine reuptake inhibitor and an NMDA receptor antagonist which is chiefly used in the management of moderate to severe pain and has a relatively rapid onset of action. Its side effects include delirium and hallucinations.

Solution to Question 34:

The anti-epileptic drugs which can also be used for neuropathic pain are 1, 2, and 4 which are pregabalin, carbamazepine and gabapentin respectively.

Pregabalin is an anti-epileptic drug that acts by binding to α2δ, a protein that is a subunit of voltage-gated calcium channels and causes a decrease in glutamate release at excitatory synapses. It is used in the management of neuropathic pain associated with diabetic peripheral neuropathy, spinal cord injury or postherpetic neuralgia.

Gabapentin is an anti-epileptic drug that, like pregabalin, acts by binding to α2δ protein. It is used in the management of neuropathic pain due to postherpetic neuralgia in adults.

The binding of gabapentin and pregabalin, also called "gabapentinoids" due to their structural similarity to GABA, to calcium channel subunit a2δ protein, is considered to mediate the anti-convulsant effects of gabapentin and pregabalin. They are used in focal onset seizures with or without progression to bilateral tonic-clonic seizures when used in addition to other anti-epileptic drugs.

Carbamazepine is an iminostilbine derivative. It blocks voltage-activated sodium channels, slowing their rate of recovery and limiting the repetitive firing of action potentials. It is used in the treatment of:

• Generalized, focal-to-bilateral and idiopathic tonic-clonic seizures

• Focal seizures

• Neuropathic pain associated with trigeminal and glossopharyngeal neuralgias.

Lamotrigine is an anti-folate agent whose mechanism of action is similar to that of carbamazepine i.e. it delays the recovery from inactivation of recombinant sodium channels. It also inhibits the release of glutamate by acting directly on the sodium channels. Therefore, when used concurrently with carbamazepine, is associated with toxicity in some patients due to increase in 10,11- epoxide of carbamazepine. It is useful in the management of:

• Focal and secondarily generalized tonic-clonic seizures in adults

• Lennox-Gastaut syndrome - a childhood disorder characterized by multiple type seizures, mental retardation, and refractoriness to anti-seizure medication.

Solution to Question 35:

Linezolid cannot be used in this patient as bone marrow suppression is a side effect of this drug and hence it can worsen the post-chemotherapy bone marrow suppression in this patient.

Linezolid can cause myelosuppression which can present as anemia, thrombocytopenia, leukopenia, and pancytopenia. It is commonly associated with thrombocytopenia and is seen 7-10 days following administration. This is usually reversed on discontinuation of the drug.

Other adverse effects of linezolid:

• It inhibits mitochondrial protein synthesis and may lead to peripheral neuropathy, optic neuritis, and lactic acidosis in patients receiving the drug for longer durations.

• It can cause serotonin syndrome when coadministered with adrenergic or serotonergic drugs like SSRIs due to its nonspecific MAO inhibition.

It is a protein synthesis inhibitor of the class oxazolidinones. It mainly acts against gram-positive organisms. It is approved for the treatment of the following conditions:

• Skin and skin structure infections by staphylococcus(Both MRSA and MSSA) and streptococcus.

• Community-acquired pneumonia due to streptococcus pneumoniae.

• Nosocomial pneumonia due to staphylococcus aureus.

Other options:

Cefepime: It is a fourth-generation antipseudomonal cephalosporin. It is useful in enterobacter and pseudomonas infections that are resistant to other cephalosporins.

Piperacillin-Tazobactam- It is a combination of piperacillin which is a ureidopenicillin and tazobactam which is a beta-lactamase inhibitor. It is useful in various infections including nosocomial infections, intraabdominal infections, pneumonia, and UTI. It is highly effective against E Coli and Klebsiella.

Meropenem: It belongs to the carbapenem group of beta-lactam antibiotics. They are more active against gram-negative organisms. Coadministration of cilastatin is not required with this drug. (Cilastation should be coadministered with imipenem as imipenem is sensitive to renal peptidase degradation)

Pharmacology NEET 2021

Question 1:

A female patient on oral contraceptive pills (OCP) was diagnosed to have tuberculosis and started on antitubercular therapy (ATT). She was advised to use an additional barrier method of contraception. What is the reason for this advice?

a) Teratogenic

b) Failure of ATT

c) Rifampicin decreases OCP metabolism

d) Rifampicin induces OCP metabolism and cause failure

Question 2:

A 35-year-old chronic hypertensive woman visits the outpatient department for pre-conceptional counseling. Which of the following anti-hypertensives should she discontinue?

a) Labetalol

b) Calcium channel blockers

c) Angiotensin-converting enzyme inhibitors

d) Alpha methyldopa

Question 3:

A pregnant woman was prescribed indomethacin. Which of the following effects does the drug have on the fetus?

a) Premature closure of ductus arteriosus

b) Keeps ductus arteriosus patent

c) Closure of fossa ovalis

d) Premature closure of ductus venosus

Question 4:

Which of the following can be prescribed in case of an open angle glaucoma with bronchial asthma?

a) Timolol

b) Gemeprost

c) Latanoprost

d) Mannitol infusion

Question 5:

As a medical practitioner in India, how will you prescribe alprazolam to a patient?

a) 1 mg OD before bedtime for 7 days

b) 1/2 mg HS for 7 days

c) 0.5 mg BD for 7 days

d) 0.5 mg daily before bedtime for 7 days

Question 6:

A patient started on antihypertensive presented with the following picture. Which of the following drugs is the most likely cause?

a) Lisinopril

b) Nifedipine

c) Atenolol

d) Clonidine

Question 7:

A patient on aspirin presented with occult blood in the stool. This is due to inhibition of____ by aspirin

a) Phospholipase A2

b) Thromboxane

c) Phospholipase C

d) Phospholipase D

Question 8:

A 60-year-old postmenopausal female patient with a previous history of Colle's fracture came with complaints of lower backache. On evaluation, her T score was -2.5. Which of the following statements are false about the management of her condition?

a) Teriparatide should be started before supplementing with bisphosphonates

b) Bisphosphonates are not given for more than a year

c) Calcium requirement is 1200 mg per day

d) Oral vitamin D3 is given along with oral calcium

Question 9:

A patient presented with a history of multiple episodes of vomiting. He was given an intravenous drug, and later, he developed abnormal movements. Which of the following can be used to treat him?

a) Hyoscine

b) Cyproheptadine

c) Levodopa

d) Benzhexol

Question 10:

What formula is the dosing rate of a drug based on?

a) Clearance / plasma concentration

b) Clearance x plasma concentration

c) Clearance x loading dose

d) Clearance / loading dose

Question 11:

A patient on digoxin, potassium chloride, clarithromycin, atenolol, and triamterene developed atrial fibrillation with controlled ventricular rate. Serum digoxin levels were raised. Which of his medications is responsible for the same?

a) Potassium chloride

b) Clarithromycin

c) Atenolol

d) Triamterene

Question 12:

What is the mechanism of action of methotrexate in osteosarcoma?

a) Analog of UMP

b) Competitive inhibitor of DHFR

c) Blocks purine synthesis

d) Blocks pyrimidine synthesis

Question 13:

What should not be consumed by patients with bacterial vaginosis and trichomoniasis on metronidazole?

a) Grapefruit juice

b) Alcohol

c) Benzodiazepine

d) MAOI/SSRI

Question 14:

A patient with chronic kidney disease underwent a nephrectomy. Which of the following can be administered as a postoperative analgesic in this patient?

a) Naproxen

b) Indomethacin

c) Diclofenac

d) Acetaminophen

Question 15:

A patient with renal insufficiency presented with very low urine output, pedal edema, headache and BP 160/90. Which anti-hypertensive can be administered?

a) Aliskiren

b) Chlorthalidone

c) Amlodipine

d) Prazosin

Question 16:

Which of the following interleukin (IL) is inhibited by tocilizumab?

a) IL-6

b) IL-2

c) IL-4

d) IL-5

Answer Key

Question No.	Correct Option
1	d
2	c
3	a
4	c
5	d

6	a
7	b
8	b
9	d
10	b
11	b
12	b
13	b
14	d
15	b
16	a

Detailed Explanations

Solution to Question 1:

Rifampicin is a potent cytochrome P450 (CYP450) enzyme inducer. When given along with oral contraceptive pills(OCP) it would induce the metabolism of OCPs leading to contraceptive failure. This is the reason for the patient being advised to use an additional method of contraception.

Important enzyme inducers are:

• Barbiturates

• Phenytoin

• Carbamazepine

• Rifampicin

• Cigarette smoking

• Chronic alcoholism

Important drugs that inhibit CYP450 isoenzyme are:

• Macrolide antibiotics

• Azole antifungals

• Chloramphenicol

• Omeprazole

• Selective serotonin reuptake inhibitors (SSRIs)

• HIV protease inhibitors

• Cimetidine

• Ciprofloxacin

• Metronidazole

Solution to Question 2:

Angiotensin-converting enzyme (ACE) inhibitors should be discontinued in pregnancy as they can cause hypoplasia of organs, growth retardation, and fetal loss.

ACE inhibitors cause reduced perfusion and oligohydramnios leading to fetal growth retardation, calvarium maldevelopment, hypoplasia of lungs and kidneys, and limb contractures.

Antihypertensive drugs to be avoided in pregnancy:

• ACE inhibitors and Angiotensin receptor blockers(ARB) - Can cause growth retardation and fetal damage.

• Diuretics - Can decrease uteroplacental perfusion

• Nonselective beta-blockers

• Sodium nitroprusside - It is contraindicated in eclampsia.

Antihypertensive drugs that can be used in pregnancy:

• Labetalol - Combined $\alpha + \beta$ blocker

• Nifedipine - Dihydropyridine calcium channel blocker. It is useful in preeclampsia but should be stopped before labor as they weaken the uterine contractions.

• Methyldopa

• Hydralazine

Note:

Thiazide diuretics like hydrochlorothiazide are considered to be safe in pregnancy and are useful in the management of chronic hypertension in pregnant women.

Solution to Question 3:

Indomethacin can cause premature closure of ductus arteriosus if given to a pregnant woman.

Indomethacin is approved for the closure of persistent ductus arteriosus in premature infants. It is given intravenously. One of its limiting factors is the development of renal toxicity in neonates.

Indomethacin is a nonselective COX inhibitor. It has anti-inflammatory, analgesic, and antipyretic properties. It can be used in the management of conditions like rheumatoid arthritis, gouty arthritis, osteoarthritis, ankylosing spondylitis, etc. But its use is limited due to the high rate of intolerance and availability of safer agents.

Adverse effects: Diarrhea with ulcerative bowel lesions, acute pancreatitis, and hepatitis are seen. Frontal headache is the most frequent CNS effect. Dizziness, light-headedness, vertigo, and mental confusion can occur. Neutropenia, thrombocytopenia, and rarely aplastic anemia can also occur.

Note:

Prostaglandin analogs are indicated in the maintenance of patency of the ductus arteriosus.

Solution to Question 4:

Latanoprost can be given in a case of open-angle glaucoma with bronchial asthma.

Latanoprost is a PGF2α analog and it is not associated with systemic side effects like bronchospasm and hence can be given for asthmatic patients. Prostaglandin analogs are the first choice drugs for open-angle glaucoma.

Mechanism of action: It reduces intraocular pressure by altering the ciliary muscle tension and improving the outflow of the aqueous humor.

Advantages of latanoprost:

• Good efficacy

• Once-daily application

• Absence of systemic side effects

Disadvantage: High cost

Adverse effects:

• Blurring of vision

• Iris pigmentation

• Thickening and darkening of eyelashes

• Macular edema can occur especially in aphakic patients.

Other options:

Option A- Timolol is a non-selective beta-blocker used topically in glaucoma. It is associated with systemic effects by absorption through the nasolacrimal duct and can cause life-threatening bronchospasm in asthmatics and COPD patients. Hence they are not given in in this case.

Option B- Gemeprost is a PGE1 analog used for the termination of early pregnancy.

Option D- Mannitol infusion is given for acute congestive glaucoma and not for open-angle glaucoma. It causes decongestion of the eye by its osmotic action.

Solution to Question 5:

The correct method of prescribing alprazolam is 0.5mg daily before bedtime for 7 days.

The drugs are prescribed in generic names legibly and preferably in capital letters. The prescription should have details to identify the patient and medication that needs to be dispensed and the therapeutic purpose. The use of abbreviations, especially Latin is discouraged. The dosage should be written in metric measure.

Guidelines for drug prescription:

The parts of an ideal prescription are as follows.

• Date

• Name and address of the doctor

• Superscription(Rx symbol)

• Information about the patient

• Name and dose of the drug prescribed(Inscription)

• Dispencing directions to the pharmacist(Subscription)

• Instructions for the patient

• Signature of the doctor

Solution to Question 6:

The image shows edema over the lips and face and its occurrence following intake of an antihypertensive is suggestive of drug-induced angioedema. Lisinopril being an ACE inhibitor is most likely to be responsible for this condition.

Angioedema with ACE inhibitors:

In very few patients(0.1%-0.5%), ACE inhibitors cause rapid swelling in the nose, throat, mouth, glottis, larynx, lips, or tongue. Once the drug is stopped symptoms improve drastically over hours. Angioedema of the intestine can occur very rarely.

ACE inhibitor-associated angioedema is a class effect, and patients who develop this adverse event should not be prescribed any other drugs within the ACE inhibitor class.

Other adverse effects of ACE inhibitors:

Hypotension - A sudden fall in blood pressure can occur following the first dose of ACE inhibitors.

Cough - It can produce a bothersome dry cough due to the accumulation of bradykinin, substance P, or prostaglandins in the lungs. The cough usually disappears within 4 days of the stoppage of the drug.

Hyperkalemia due to K+ retention can occur.

Acute renal failure can occur with ACE inhibitors in patients with bilateral renal artery stenosis, stenosis of the artery to a single remaining kidney, heart failure, or volume depletion due to diarrhea or diuretics.

It should be discontinued in pregnant women as it can cause developmental defects.

It can also cause maculopapular skin rash in some cases.

Other options:

Option B: Nifedipine is a calcium channel blocker. Common adverse effects include headache, flushing, and dizziness. Peripheral edema may rarely occur.

Option C: Atenolol is a beta 1 selective antagonist. Bradycardia, fatigue, cold extremities are common adverse effects. CNS side effects like nightmares and depression are less frequent with atenolol than other beta-blockers.

Option D: Clonidine is an alpha-2 adrenergic agonist. The main adverse effects of clonidine are dry mouth and sedation. Bradycardia and sexual dysfunction are other effects.

Solution to Question 7:

Aspirin causes inhibition of cyclooxygenase-1(COX-1) which would inhibit the formation of thromboxane A2. Thromboxane A2 is an inducer of platelet aggregation and is a potent vasoconstrictor. Hence inhibition of its formation is responsible for the bleeding complications of aspirin.

Aspirin is a noncompetitive, irreversible inhibitor of the cyclooxygenase(COX) enzyme. It is a nonselective COX inhibitor and inhibits both COX-1 and COX-2.

Uses of aspirin:

• Analgesic and antipyretic uses.

• Acute rheumatic fever

• Rheumatoid arthritis

• Post myocardial infarction and stroke due to its antiplatelet action.

• Prevention of preeclampsia

Adverse effects:

Nausea, vomiting, epigastric discomfort, and increased occult blood in stools are seen with analgesic doses(0.5-2g/day). Antiinflammatory doses(3-5g/day) can produce salicylism characterized by dizziness, tinnitus, vertigo, hyperventilation, reversible loss of hearing and vision, etc. It can cause elevation of liver enzymes and hepatic damage. Hypersensitivity reactions are also seen.

Reye's syndrome is a rare condition that can occur when salicylates like aspirin are given in children having viral infections like influenza.

Solution to Question 8:

Bisphosphonates are given for periods of more than one year (Option B is false). In this case, as the patient is having a history of osteoporotic fracture, teriparatide can be given before bisphosphonates.

Pharmacologic management of osteoporosis:

Bisphosphonates: These are pyrophosphate analogs that act by directly inhibiting bone resorption. The antiresorptive action is mainly due to its direct inhibitory effect on osteoclasts. It causes osteoclast apoptosis as well as the prevention of hydroxyapatite dissolution. Long-term bisphosphonates therapy is given for durations of 3-5 years, depending on the fracture risk of the

patient. Since they are associated with osteonecrosis of the jaw and atypical femur fractures, a drug holiday may be advised after long-term management.

Teriparatide is a recombinant PTH analog that has an anabolic effect on bone homeostasis by increasing osteoblastic activity. It increases bone mineral density and decreases the risk of fractures in postmenopausal women. Indications of teriparatide therapy include the history of osteoporotic fracture, multiple risk factors for fracture, intolerance, or failure of previous osteoporosis therapy. It is only approved for 2 years of use. Clinical trials that used teriparatide prior to bisphosphonate have been proven effective.

Denosumab is a human monoclonal antibody with a mechanism of action similar to that of osteoprotegerin (OPG). It binds to RANK Ligand (RANKL), thereby inhibiting its binding to the RANK receptor of the surface of osteoclasts. This prevents RANK receptor activation and subsequent bone resorption.

Selective estrogen receptor modulators (SERMs) - Estrogen has a protective effect that prevents accelerated bone loss. This is carried out by reducing the bone resorption effects of parathyroid hormone (PTH). Keeping in line with these effects, osteoporosis prevention in postmenopausal women has been attributed to the use of SERMs such as raloxifene. It acts as an estrogen agonist on the bone and liver.

Calcium and Vitamin D supplementation: Calcium supplementation in the elderly suppresses bone turnover and improves bone mineral density. 1200mg of calcium is daily required in adults of more than 50 years of age. Vitamin D supplementation with 400-800IU/day improves intestinal calcium absorption, suppresses bone remodeling, and improves bone mineral density.

Solution to Question 9:

Benzhexol should be used in the management of this patient. The development of abnormal movements following an anti-emetic injection suggests that the antiemetic used in this case is probably metoclopramide.

Metoclopramide is a dopamine receptor antagonist used as a prokinetic and anti-emetic agent. It can produce extrapyramidal side effects and dystonia is seen acutely following intravenous injections. Anticholinergics like benzhexol (trihexyphenidyl) are used in the management of these side effects. Benzhexol is a centrally-acting anticholinergic agent. It acts by restoring the cholinergic balance in the striatal interneurons.

Extrapyramidal side effects of metoclopramide are commonly seen in children and young adults with higher doses. Acute dystonia occurs after intravenous administration, while parkinsonian-like symptoms develop weeks after therapy. It is managed with the discontinuation of metoclopramide and with anticholinergics or antihistaminics. Irreversible tardive dyskinesia can occur with chronic treatment.

Other adverse effects:

It can cause galactorrhea by its dopamine antagonism. Methemoglobinemia can occur if used in infants.

Other options:

Option A: Hyoscine (scopolamine): It is a muscarinic receptor antagonist used in the management of motion sickness and vomiting.

Option B: Cyproheptadine is an H1 receptor antagonist used in the management of various allergy-related conditions like allergic rhinitis, anaphylaxis, urticaria, etc.

Option C: Levodopa is the precursor of dopamine and it is given in combination with carbidopa for the management of parkinsonism.

Solution to Question 10:

The formula for drug dosing rate is clearance x plasma concentration.

The equation for dose rate given here is based on steady-state concentration. When a drug is given repeatedly at short intervals it will accumulate in the body until the elimination of the drug matches its input. So steady-state concentration is the plasma concentration at which the drug elimination and drug input balance each other.

Steady-state concentration(Cpss) = Dose rate / Clearance

From this equation dose rate for a target steady-state plasma concentration is calculated as

Dose rate = Clearance(CL) x target plasma concentration(Cpss)

Solution to Question 11:

Elevated serum digoxin levels and the development of atrial fibrillation with controlled ventricular rate are suggestive of digoxin toxicity. Clarithromycin when given along with digoxin can cause digoxin toxicity.

P-glycoprotein pumps play a major role in the elimination of digoxin. Drugs that inhibit P-glycoprotein can thereby decrease the elimination of digoxin and lead to digoxin toxicity. Macrolide antibiotics like clarithromycin and erythromycin are inhibitors of the P-glycoprotein. So clarithromycin when given along with digoxin can cause digoxin toxicity.

Other drugs that inhibit P-glycoprotein are - Amiodarone, quinidine, verapamil, diltiazem, cyclosporine, itraconazole, propafenone, and flecainide.

Digoxin is one of the cardiac glycosides which is derived from the digitalis plant and used clinically.

It inhibits Na+/K+ ATPase, a membrane-bound transporter to increase intracellular sodium. As a result, there is less expulsion of calcium from the cell by sodium-calcium exchanger (NCX). Increased calcium inside the myocyte leads to an increase in cardiac contractility.

The most common cardiac manifestations of digitalis toxicity include atrioventricular junctional rhythm, premature ventricular depolarizations, ventricular bigeminy, ventricular tachycardia, and second-degree AV blockade.

Factors that increase the risk of digoxin toxicity are:

• Hypercalcemia

• Hypomagnesemia

• Hypokalemia- It increases the binding of the digoxin to the Na+/K+ ATPase pump.

• Drugs causing hypokalemia- loop and thiazide diuretics, amphotericin B, corticosteroids

• Drugs that decrease digoxin elimination (inhibitors of P-glycoprotein)

• Renal failure

• Hypothyroidism

Triamterene, atenolol, and potassium chloride cause hyperkalemia, and therefore do not cause digoxin toxicity.

Solution to Question 12:

The anticancer activity of methotrexate is by inhibition of dihydrofolate reductase (DHFR).

Methotrexate belongs to the antimetabolite class of anticancer drugs. The antifolate action of the drug is responsible for its anti-cancer properties.

Mechanism of action:

It is a folic acid analog and binds to the active catalytic site of dihydrofolate reductase(DHFR). This would inhibit the formation of tetrahydrofolate(THF) which is needed for purine nucleotides, thymidylate, and the amino acids serine and methionine. This would interfere with the production of DNA, RNA, and important cellular proteins.

Low doses of methotrexate are used in the management of rheumatic diseases. It is due to the inhibition of amino-imidazole carboxamide ribonucleotide (AICAR) transformylase and thymidylate synthetase. This results in increased levels of adenosine that inhibits inflammation.



Adverse effects:

Common side effects are nausea and mucosal ulcers. Others include anemia, leukopenia, stomatitis, GI ulcerations, and alopecia. Dose-related hepatotoxicity is also seen. The GI effects and hepatotoxicity can be reduced by the use of leucovorin or folinic acid.

Solution to Question 13:

Alcohol should not be consumed by a patient on metronidazole. Metronidazole produces a disulfiram-like reaction. If the patient consumes alcohol during or within 3 days of therapy, they may experience abdominal distress, vomiting, headache, and flushing.

Disulfiram and disulfiram-like reactions:

Disulfiram inhibits the aldehyde dehydrogenase(ALDH) activity and thereby increases the concentration of acetaldehyde in the blood. When a person treated with disulfiram consumes alcohol he or she would develop symptoms of acetaldehyde poisoning. The symptoms are:

- Flushing
- Pulsating headache
- Nausea and vomiting
- Sweating
- Tachycardia
- Hypotension
- Orthostatic syncope
- Vertigo
- Blurred vision

Certain drugs can also produce disulfiram-like reactions if alcohol is consumed along with these drugs. Patients should be warned not to use alcohol along with these drugs. The drugs include:

- Metronidazole
- Tinidazole
- Cefoperazone
- Sulfonylureas

Solution to Question 14:

Acetaminophen can be given as an analgesic for the patient as it is the least nephrotoxic among the given options.

Analgesia in CKD patients:

Acetaminophen is considered a safer non-narcotic analgesic in chronic kidney disease (CKD) patients. For more severe pains opioid analgesics like tramadol or fentanyl are used.

Even though acetaminophen is considered safe, with chronic high dosage it can also become nephrotoxic.

NSAIDs like naproxen, indomethacin, diclofenac, etc are not used in CKD patients because of their direct nephrotoxic effects. In healthy normal individuals, NSAIDs have little effect on renal function and blood pressure. But in conditions where the regulatory systems are impaired as hypovolemia, congestive heart failure, hepatic cirrhosis, chronic kidney disease, or other conditions where the renin-angiotensin system is activated, renal function by the formation of prostaglandins are very important.

NSAIDs cause impairment of prostaglandin-induced inhibition of chlorine reabsorption and antidiuretic hormone action. This will cause salt and water retention. It also inhibits COX-2 derived prostaglandins which regulate the renal medullary blood flow and would lead to a rise in blood pressure, increased cardiovascular thrombotic events, and cardiac failure. It can also cause electrolyte abnormalities like hyponatremia and hyperkalemia.

Solution to Question 15:

The antihypertensive that can be given in this case is chlorthalidone. The patient in this case is having renal insufficiency but there is no proteinuria mentioned in the question stem. So among the given options, the thiazide diuretic chlorthalidone can be administered as antihypertensive in this patient.

Management of hypertension in chronic kidney disease(CKD):

ACE inhibitors or ARBs are considered the first-line drugs in patients with proteinuria. These agents effectively reduce blood pressure, proteinuria and prevent kidney decline. Calcium channel blockers can be added if necessary.

For patients who do not have proteinuria, there is no established first-line therapy. One or two of the following drugs can be used:

• Thiazides/thiazide-like diuretics

• ACE inhibitors or ARBs

• Calcium channel blockers

In patients presenting with fluid retention or fluid overload, thiazide diuretics such as chlorthalidone and hydrochlorothiazide are recommended for management. In CKD stages 1 to 3 (GFR >=30ml/min), thiazides are recommended, while loop diuretics such as bumetanide, furosemide are used in stages 4-5 (GFR < 30ml/min).

Other options:

Option A: Aliskiren is a direct renin inhibitor. It effectively lowers blood pressure, reduces albuminuria, and limits left ventricular hypertrophy.

Option C: Amlodipine is a calcium channel blocker. A common side effect of amlodipine is the production of peripheral edema.

Option D: Prazosin is a selective α1 antagonist. It is generally not used as an initial agent to treat hypertension, except in symptomatic prostatism or nightmares linked to posttraumatic stress disorder. Headache is one of the side effects of prazosin.

Solution to Question 16:

Tocilizumab inhibits IL-6.

Tocilizumab is a newer humanized antibody. It binds to the soluble, membrane-bound IL-6 receptors and inhibits IL-6 mediated signaling. IL-6 can cause T-cell activation, acute-phase protein synthesis, and stimulation of inflammatory processes in diseases like rheumatoid arthritis.

Uses:

• Rheumatoid arthritis - It is a biological DMARD and is indicated in adult patients having moderate to severe disease with inadequate response to one or more DMARDs.

• Juvenile idiopathic arthritis - It can be given for active systemic juvenile idiopathic arthritis(SJIA) or polyarticular juvenile idiopathic arthritis(PJIA) in patients of 2 years of age or more.

Adverse effects:

Common adverse effects include upper respiratory infections, hypertension, headache, and elevation of hepatic enzymes. It is also associated with serious infections including TB and opportunistic infections. Neutropenia and reduction in platelet count can also occur.

Other options:

Option B- IL-2 receptor inhibitors are daclizumab, basiliximab. They are used as immunosuppressive agents following renal and heart transplants.

Option C- IL-4 receptor inhibitor is dupilumab. They have been shown to reduce exacerbations in asthma.

Option D- IL-5 inhibitors are mepolizumab, reslizumab, benralizumab. They have been shown to reduce exacerbations in asthma.

Pharmacology INI-CET 2022

Question 1:

The extent of drug absorption is represented by:

a) Tmax

b) Area under the curve

c) Cmax

d) t1/2

Question 2:

Which of the following statement is correct regarding the given graph?



a) Drug 1 represents agonist and drug 2 represents inverse agonist

b) Drug 3 represents agonist and drug 4 represents inverse agonist

c) Drug 1 represents agonist and drug 4 represents inverse agonist

d) Drug 2 represents partial agonist and drug 3 represents inverse agonist

Question 3:

Which among the following drugs is the new FDA approved immune check point inhibitor for carcinoma endometrium?

a) Nivolumab

b) Pembrolizumab

c) Trastuzumab

d) Ipilimumab

Question 4:

Which of the following is not an immune checkpoint inhibitor?

a) Nivolumab

b) Pembrolizumab

c) Atezolimumab

d) Cetuximab

Question 5:

NAPQI is the metabolite of which of the following drug toxicity?

a) Aspirin

b) Paracetamol

c) Diclofenac

d) Ketorolac

Question 6:

Identify the correct match, regarding the drug and its adverse effect.

a) Hydralazine - heart failure

b) Atenolol - hemolytic anemia

c) Aliskiren - hypokalemia

d) Verapamil - constipation

Question 7:

A lady who had undergone mastectomy for breast cancer is being treated with tamoxifen. How long should it be stopped for before she can conceive?

a) 1 month

b) 3 months

c) No need to stop

d) Can be stopped and conceived soon after stopping

Question 8:

A young boy with a history of multiple blood transfusions presents with bone pain for the past 6 months. A peripheral smear showed sickle-shaped RBCs. Which of the following drugs are not used in the treatment of his condition?

a) Bebtelovimab

b) Voxelotor

c) Hydroxyurea

d) L-glutamine

Question 9:

The half-life of letrozole is:

a) 45 hours

b) 70 hours

c) 90 hours

d) 120 hours

Question 10:

Identify the false statement about eicosanoids.

a) Synthesized from arginine

b) Synthesized from arachidonic acid

c) Prostaglandins have 5 carbon ring

d) Synthesized in platelets

Question 11:

The given diagram shows the steps in bacterial cell wall formation. Identify the site of action of beta-lactam antibiotics.



a) 1

b) 2

c) 3

d) 4

Question 12:

Phospholipase C acts via which of the following G protein-coupled receptors?

a) Gs

b) Gi

c) Gq

d) Go

Question 13:

A patient was diagnosed with pulmonary tuberculosis and was started on first-line antitubercular drugs. Which of the following drug acts by inhibiting the synthesis of the marked layer shown in the image?



a) Isoniazid

b) Rifampicin

c) Pyrazinamide

d) Ethambutol

Question 14:

A man was brought to the emergency room after poisoning with an unknown substance. Muscarinic poisoning was suspected and he was treated for the same. What is the possible presenting feature which led to the diagnosis?

a) Mydriasis

b) Diuresis

c) Muscle fasciculations

d) Bradycardia

Question 15:

A 29-year-old lady came to psychiatry OPD with symptoms of hypomania. There is a past history of maniac episode. Now, she is planning to conceive. Which drug should be avoided for being highly teratogenic to the fetus?

a) Valproate

b) Lithium

c) Oxcarbazepine

d) Olanzapine

Question 16:

True statement regarding the administration of 5 alpha-reductase inhibitors is/are:

a) 1, 2, 3

b) 1, 3, 4

c) 1, 4

d) 2, 3

Question 17:

The manufacturer of a drug company labels a drug as having 500 mg paracetamol. On quantitative analysis of the drug, it actually contained only 200 mg of paracetamol. What is this type of drug called?

a) Spurious drug

b) Adulterated drug

c) Unethical drug

d) Misbranded drug

Question 18:

Which of the following anti-diabetic drugs is associated with the increased risk of fractures?

a) Rosiglitazone

b) Liraglutide

c) Voglibose

d) Canagliflozin

Question 19:

The active metabolite of cyclophosphamide is:

a) N - hydroxy cyclophosphamide

b) 4 - hydroxy cyclophosphamide

c) 4 - acetyl cyclophosphamide

d) N - acetyl cyclophosphamide

Question 20:

Urinary alkalinization will be most useful in toxicity of which of the following drugs?

a) Beta blockers

b) Calcium channel blockers

c) Acetaminophen

d) Aspirin

Question 21:

A hospital is conducting a study to check the efficacy of statins in multiple centers. There are 4-5 patients per center and half of the patients receive the drug and the other half receive the placebo. Neither the patients nor clinicians know which drug is being administered. What phase of the clinical trial describes the above scenario?

a) Phase 1

b) Phase 3

c) Phase 2

d) Phase 4

Question 22:

A patient who is a known addict is brought to the hospital with the intoxication of an agent. An ocular examination shows the following finding. Which of the following is true regarding this condition?



a) This action can be reversed by phenylephrine.

b) It cannot be reversed by levallorphan.

c) It is caused by activation of the Edinger-Westphal nucleus.

d) Tolerance develops due to long-term use.

Question 23:

Match the following drugs in Column A with their contraindications in Column B.

a) A-1, B-3, C-2, D-4

b) A-4, B-1, C-3, D-2

c) A-2, B-4, C-1, D-3

d) A-3, B-2, C-4, D-1

Question 24:

Which of the following statements is true regarding memantine?

a) 1, 2, 3, 4

b) 1, 2

c) 3, 4

d) 1, 2, 3

Question 25:

The image below shows the life cycle of a virus. Which of the following drugs acts at step 5?



a) Raltegravir

b) Enfuvirtide

c) Tenofovir

d) Zidovudine

Question 26:

Which of the following statements is correct regarding clinical trials?

a) New drug application is done after phase 4

b) Phase 4 is also called post-marketing surveillance

c) Phase 3 is always double-blinded

d) Phase 2 is done on healthy individuals

Question 27:

Match the following teratogenic drugs with their teratogenic effects.

a) 1-a, 2-b, 3-c, 4-d

b) 1-c, 2-d, 3-a, 4-b

c) 1-d, 2-a, 3-d, 4-c

d) 1-d, 2-c, 3-b, 4-a

Question 28:

A young man sustained a scorpion sting 2 hours ago. On examination, his blood pressure is 108/ 60 mm Hg; HR, 72 beats/minute, and respiratory rate, 32 breaths/minute. He also complains of breathlessness. How do you manage this patient?

a) Metoprolol

b) Ivabradine

c) Prazosin

d) Frusemide

Question 29:

An experiment is being conducted on a cat spine to study the effect of acetylcholine. The blood pressure is recorded at each intervention and is shown below. What does phase 4 in the graph indicate?



a) Muscarinic action of acetylcholine

b) Potentiation of acetylcholine

c) Antagonism of acetylcholine

d) Nicotinic action of acetylcholine

Question 30:

A patient who is a known case of HIV is started on combined active antiretroviral therapy (cART). After 5 months of treatment, he develops hyperpigmentation in the palms and soles. What is the most likely drug causing this?

a) Nelfinavir

b) Emtricitabine

c) Abacavir

d) Zidovudine

Question 31:

Dorzolamide is a _____.

a) Topically applied sulfonamide antibacterial

b) Topical ocular carbonic anhydrase inhibitor

c) Second-generation sulfonylurea

d) Opioid receptor agonist

Question 32:

A patient with erectile dysfunction is given sildenafil. The penile erectile function is found to be improved after the therapy. Which of the following mediates the action of sildenafil?

a) cAMP

b) cGMP

c) Calcium

d) Phosphatidyl glycerol and inositol phosphate

Question 33:

Match the following diuretics in column A to their site of action in column B.

a) A-2, B-3, C-1, D-4

b) A-3, B-4, C-2, D-1

c) A-1, B-2, C-4, D-3

d) A-4, B-1, C-3, D-2

Question 34:

Alkalinization of urine will be helpful in the removal of _____.

a) Weakly basic drugs

b) Strong electrolytes

c) Polar drugs

d) Weakly acidic drugs

Question 35:

All of the following side effects are more commonly seen with carbamazepine than oxcarbazepine, except_____.

a) Hypotension

b) Hyponatremia

c) Rash

d) Elevated levels of gamma glutamyl transpeptidases

Question 36:

Match the following drugs with the targets of their actions.

a) A-2, B-3 , C-1 , D-4

b) A-4, B-3 , C-1 , D-2

c) A-3, B-4 , C-2 , D-1

d) A-4, B-3 , C-2 , D-1

Question 37:

A patient is brought with altered sensorium and seizures. Attender shows the laboratory report which reveals elevated transaminases. Which of the following can be used in this condition?

a) Midazolam

b) Diazepam

c) Alprazolam

d) Oxazepam

Question 38:

The lipid profile of a patient who suffered from acute coronary syndrome is given below. What drug can be used to treat this condition?

a) Fenofibrate 160 mg

b) Rosuvastatin + Fenofibrate

c) Rosuvastatin 10 mg

d) Atorvastatin 80 mg

Question 39:

An anti-arrhythmic drug is shown to have the following effect. It is effective in both atrial and ventricular arrhythmias. Identify the drug.



a) Tocainide

b) Propranolol

c) Quinidine

d) Lignocaine

Question 40:

Which among the following is the false statement regarding statins?

a) Although HMG-CoA reductase inhibitors substantially reduce the risk of cardiovascular events, there is mild increase in lipoprotein a (Lpa) levels.

b) These drugs should not be stopped even in severe conditions like injury, surgery, etc.

c) They can be given with verapamil and other enzyme inhibitors.

d) With the long term use, there is slight increase in the incidence of type 2 diabetes mellitus.

Question 41:

Identify the drugs marked as A, B, and C.



a) A - Aspirin, B -Vorapaxar, C - Prasugrel

b) A - Aspirin, B - Eptifibatide, C - Vorapaxar

c) A - Aspirin, B - Eptifibatide, C - Prasugrel

d) A - Aspirin, B - Prasugrel, C - Eptifibatide

Question 42:

Which of the following is false regarding disulfiram?

a) Only 3

b) 1, and 3

c) 1, 3, and 4

d) 2, 3, and 4

Question 43:

G6PD can manifest with which of the following drugs?

a) Primaquine

b) Chloroquine

c) Penicillin

d) Tetracycline

Question 44:

Choose the correctly matched pairs regarding the drugs used in schizophrenia:

a) 1,3

b) 1,4

c) 1,2,4

d) 1,2,3,4

Question 45:

A patient presented with a lesion, as shown in the image below. She is an immunocompromised patient currently on HAART. What are the systemic antifungal drugs that can be added along with the HAART in this patient?

a) Inj. Amphotericin B 4-8 weeks, followed by oral itraconazole

b) Inj. Amphotericin B 8 weeks, followed by oral posaconazole

c) Inj. Fluconazole, followed by oral fluconazole

d) Inj. Amphotericin B, followed by flucytosine

Question 46:

A patient who is on treatment for hyperlipidemia develops gallstones. What is the mechanism of action of the causative drug that was given to this patient?

a) Inhibits HMG CoA reductase

b) Activates PPAR alpha

c) Binds to deoxycholic acid

d) Decreases VLDL

Question 47:

The dose-response curve for three drugs in the same tissue is shown in the image below. Which of the following statements is true?



a) Efficacy of drug A is more than that of drug B.

b) Potency of drug B is more than that of drug A.

c) Drug A & drug B are agonists.

d) Drug C is a non-competitive antagonist.

Question 48:

Which of the following drugs is a β2 agonist?

a) 1, 2, 3

b) 4, 3, 2, 1

c) 1, 2

d) 3, 4

Question 49:

What is the mechanism of action of sildenafil?

a) Inhibition of 5α-reductase

b) Inhibition of phosphodiesterase-5

c) Release of nitrous oxide radicals

d) Blockage of voltage-gated calcium channels

Question 50:

Choose the correct options regarding the route of administration and bioavailability.

a) A,B,D

b) A,C,D

c) A and C

d) A and D

Question 51:

A hypertensive patient on medication presents with the lab values as shown below. Which of the following drugs should be stopped?

a) Amlodipine

b) Metoprolol

c) Losartan

d) Furosemide

Question 52:

Which of the following is the most effective oral drug in smoking cessation?

a) Varenicline

b) Cevimeline

c) Muscarine

d) Atropine

Question 53:

Match the following antiarrhythmic drugs with their mechanism of action:

a) 1-d, 2-b, 3-a, 4-c

b) 1-a, 2-c, 3-d, 4-b

c) 1-a, 2-d, 3-b, 4-c

d) 1-d, 2-c, 3-a, 4-b

Question 54:

Which of the following drugs increases trabecular outflow?

a) Timolol

b) Netarsudil

c) Brimonidine

d) Acetazolamide

Question 55:

Finasteride is used to treat male-pattern baldness acts by inhibiting which of the following enzymes?

a) Aromatase

b) 17-hydrolase

c) 5-alpha reductase

d) Phosphodiesterase

Question 56:

Rituximab is a:

a) Polyclonal antibody which acts against CD20

b) Polyclonal antibody which acts against CD50

c) Monoclonal antibody which acts against CD20

d) Monoclonal antibody which acts against CD50

Question 57:

A new drug application is filed after which phase of clinical trials?

a) 4

b) 3

c) 2

d) 1

Answer Key

Question No.	Correct Option
1	b
2	c
3	b
4	d
5	b
6	d
7	b
8	a
9	a
10	a
11	d
12	c
13	a

14	d
15	a
16	c
17	d
18	d
19	b
20	d
21	b
22	c
23	b
24	b
25	a
26	b
27	d
28	c
29	d
30	b
31	b
32	b
33	d
34	d
35	b
36	d
37	d
38	d
39	c
40	c
41	b
42	a
43	a
44	c
45	b
46	b
47	c
48	c
49	b
50	b

51	c
52	a
53	c
54	b
55	c
56	c
57	b

Detailed Explanations

Solution to Question 1:

The extent of drug absorption is represented by the area under the curve.

The rate and extent of absorption of a drug from dosage form administered by any route refer to the bioavailability of a drug. Bioavailability is determined by the concentration-time curve in blood or by its excretion in urine. The area under the blood concentration-time curve (AUC) is proportional to the dose and the extent of bioavailability of a drug.

Bioavailability is defined as the fraction of unchanged drug that reaches systemic circulation. It can be represented by the formula:

Bioavailability(F) = Quantity of drug reaching systemic circulation/ Quantity of drug administered.

Bioavailability depends on two factors:

• Absorption

• 1st pass metabolism

The area under the plot of plasma concentration of a drug versus time after dosage is called the area under the curve or AUC. It can be used as a measure of the bioavailability of a drug and to calculate its clearance rate from the body.

The image given below shows the AUC (bioavailability) in a plasma concentration-time curve:



Bioavailability of drug injected intravenously is 100%, but it is lower after oral ingestion because of incomplete absorption and first-pass metabolism in the intestinal wall/liver.

The image below shows the difference between the bioavailability of drugs injected intravenously(iv) and orally(PO). Note the area under the respective curves is shaded in red and blue colours which represent the absorption of the drug.



Solution to Question 2:

The correct statement is Drug 1 represents agonist and drug 4 represents inverse agonist.

Graph 1 represents the agonist: An agonist is an agent which initiates the normal physiological response when combined with a receptor.

Example: Salbutamol is a beta-2 receptor agonist.

Graph 2 represents a partial agonist: An agent which produces a submaximal or partial effect when combined with a receptor.

Example: Buprenorphine and butorphanol are partial agonists at opioid receptors.

Graph 3 represents the antagonist: An agent which blocks or reduces the action of an agonist on a receptor, but does not have any effect of its own.

Example: Naltrexone and naloxone are antagonists at opioid receptors

Graph 4 represents an inverse agonist: An agent which activates a receptor to produce an effect in the opposite direction to that of the agonist.

Example: β-Carboline is an inverse agonist at benzodiazepine site of GABA-A receptor.

The image given below shows action of various types of selective receptor modulators.



Solution to Question 3:

The new FDA-approved immune checkpoint inhibitor for carcinoma endometrium is pembrolizumab.

Pembrolizumab acts by inhibiting programmed death receptor - 1 (PD-1) and was used for the treatment of advanced melanoma. Now, the FDA has approved the use of pembrolizumab for any cancer with mismatch repair (MMR) deficiency or high microsatellite instability (MSI-H).

On March 21, 2022, the food and drug administration approved pembrolizumab, as a single agent, for patients with advanced endometrial carcinoma that is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR), as determined by an FDA-approved test.

Other options:

Option A: Nivolumab is an immune checkpoint inhibitor that acts by inhibiting programmed death receptor - 1 (PD-1) and is used for the treatment of advanced melanoma and relapsed/refractory classical Hodgkin lymphoma.

Option C: Trastuzumab is a monoclonal antibody that inhibits the HER2 receptor. It is used in the treatment of HER2-positive breast cancer.

Option D: Ipilimumab is a monoclonal antibody that binds to cytotoxic T lymphocyte–associated Protein 4 (CTLA-4) and is approved for the treatment of late-stage melanoma.

Solution to Question 4:

Cetuximab is not an immune checkpoint inhibitor. Cetuximab is a monoclonal antibody against epidermal growth factor receptor(EGFR). It is used for the treatment of colon cancer and head and neck squamous cell cancer.

Other options.

Option A: Nivolumab is an immune checkpoint inhibitor. It acts by inhibiting programmed death receptor - 1 (PD-1) and is used for the treatment of advanced melanoma and classical Hodgkin lymphoma.

Option B: Pembrolizumab is an immune checkpoint inhibitor. It acts by inhibiting programmed death receptor - 1 (PD-1) and is used for the treatment of advanced melanoma. In 2021, approval was granted for the treatment of patients with endometrial carcinoma that progressed after systemic treatment, in combination with lenvatinib.

Option C: Atezolimumab is an immune checkpoint inhibitor. It acts by inhibiting programmed death-ligand - 1 (PDL-1) and is used for the treatment of non-small cell lung cancer.

Important note:

The FDA on May 2021 approved the use of nivolumab in patients having completely resected esophageal or gastroesophageal junction cancers with residual pathologic disease who have received neoadjuvant chemoradiotherapy.

Solution to Question 5:

N-acetyl-p-benzoquinone imine (NAPQI) is a metabolite of paracetamol drug toxicity.

Paracetamol is an analgesic and antipyretic with weak anti-inflammatory action. It is mainly metabolized in the liver by conjugation with glucuronic acid and sulfate and is excreted in the urine.

At toxic doses, the glucuronide and sulfate conjugation pathway becomes saturated. The metabolites of paracetamol get converted into NAPQI. NAPQI is highly reactive and causes enzymatic dysfunction in the liver. Normally, minor amounts of NAPQI produced in our body are rapidly eliminated by conjugation with reduced glutathione(GSH) and then excreted into the urine.

Therefore, paracetamol toxicity is treated with N-acetyl cysteine(NAC), which replenishes GSH. NAC thereby helps in the excretion of NAPQI from the body. N-acetyl cysteine is an anti-oxidant and functions by detoxifying NAPQI.

Paracetamol poisoning is the most common cause of liver failure necessitating transplantation.

The maximum dose of acetaminophen that can be given is 4 g/day.

Solution to Question 6:

The correct match is verapamil causes constipation.

Verapamil is a calcium channel blocker used for the treatment of paroxysmal supraventricular tachycardia, atrial fibrillation, and atrial flutter. Bradycardia and constipation are the important side effects of verapamil.

Other options:

Option A: Hydralazine is a vasodilator used in the treatment of heart failure, and hypertensive emergencies, especially preeclampsia. The drug is used as a combination pill containing isosorbide dinitrate (BiDil) in the treatment of heart failure. The important side effect of hydralazine is drug-induced lupus.

Option B: Atenolol is a cardioselective beta-1 blocker used in the treatment of angina and hypertension. Atenolol doesn't cause hemolytic anemia.

Option C: Aliskiren is a direct renin inhibitor used as an alternative antihypertensive drug for those who do not respond or do not tolerate 1st line drugs. The side effect of aliskiren is hyperkalemia and not hypokalemia.

Solution to Question 7:

In a post-mastectomy patient undergoing therapy with tamoxifen, it should be stopped for 3 months before she can conceive.

Tamoxifen is a selective estrogen receptor modulator (SERM) and is used as a chemotherapeutic agent in receptor positive breast cancer patients to prevent recurrence after surgery. It is the drug of choice in pre-menopausal women. It has an agonistic action on the endometrium, leading to an increased risk of endometrial cancers (2 fold risk). Hence, it is not used in post-menopausal women.

It is contraindicated in pregnancy due to teratogenic effects such as craniofacial abnormalities and reproductive tract abnormalities. For the same reason, it has to be stopped for 3 months (due to long half-life) prior to conceiving. It is also stopped during lactation as it is secreted in breast milk. Other complications of tamoxifen include deep vein thrombosis, pulmonary embolism, and hot flashes.

Other breast cancer risk reduction techniques:

• Lifestyle modification: Low-calorie diet, daily exercise, no smoking, no drinking, etc.

• More frequent screening with mammogram +/- MRI

• Chemoprevention: Preferred in women >35 years of age in patients who are ER/PR positive. Given for 5-10 years.

• SERMS (selective estrogen receptor modulators) - Tamoxifen and raloxifene. These have adverse effects.

• Aromatase Inhibitors (AIs): They are used only in post-menopausal women.

• Risk-reduction surgeries: preferred in patients with BRCA1/2 mutations or patients with first-degree relatives with breast cancers. Provides >90% risk-reduction. preferred in patients who have completed their family.

• Bilateral mastectomy

• Salpingo-oophorectomy: Best results when done < 40 years of age.

Solution to Question 8:

Bebtelovimab is not used in the treatment of sickle cell disease. The clinical features of multiple blood transfusions, bone pain, and sickle cells on peripheral smear are suggestive of sickle cell disease.

Bebtelovimab is the new monoclonal antibody used for the treatment of COVID-19. It works by binding to the spike protein of the COVID-19 virus. It has been found useful in reducing the risk of hospitalization or death. Bebtelovimab retains the activity against the omicron variant.

Other options:

Option B: Voxelotor is an inhibitor of hemoglobin S polymerase and stabilizer of sickle cell hemoglobin (HbS) that is used in the therapy of sickle cell disease.

Option C: Hydroxyurea improves fetal haemoglobin (HbF) and is the primary drug for controlling HbS in sickle cell disease. Thus improving the oxygen-carrying capacity of the blood. Hydroxyurea also inhibits the enzyme ribonucleotide reductase (RNR), which is a rate-limiting step in the biosynthesis of DNA. This property of hydroxyurea is used in the treatment of various cancers, and myeloproliferative disorders like chronic myelogenous leukemia, and polycythemia.

Option D: L-glutamine is a naturally occurring amino acid with antioxidant properties.
It increases the availability of reduced glutathione which may lessen oxidative damage in sickle red blood cells.

Solution to Question 9:

The half-life of letrozole is 45 hours.

Letrozole is a reversible aromatase inhibitor. It is a non-steroidal compound that inhibits the aromatization of estrogen thereby causing estrogen depletion. Letrozole is rapidly absorbed with 99.9% oral bioavailability and has a t½ of around 41 hours.

Letrozole is used as the first-line drug for estrogen-positive breast cancer following mastectomy in post-menopausal women.

Adverse effects of letrozole are due to the depletion of estrogen. The commonly seen adverse effects are hot flashes, vaginal dryness, nausea, diarrhea, dyspepsia, accelerated bone loss, and

thinning of hair.

Solution to Question 10:

A false statement about eicosanoids is they are synthesized from arginine. Eicosanoids are synthesized from arachidonic acid (Option B)

Eicosanoids are the most universally distributed autocoids in the body. Prostaglandins, thromboxanes, and leukotrienes comprise eicosanoids. Every cell is capable of synthesizing one or more eicosanoids. Platelets primarily synthesize thromboxanes (Option D).

Eicosanoids are synthesized from arachidonic acid. Arachidonic acid is derived from membrane phospholipids. The release of arachidonic acid (AA) from the membrane phospholipids is mainly mediated by phospholipase A.

AA is metabolized by cyclooxygenases to produce prostaglandins and lipoxygenases to produce leukotrienes and lipoxins.

Inhibitors of metabolism of arachidonic acid:

• Corticosteroids cause inhibition of prostaglandin synthesis by inhibiting phospholipase A2.

• Non-steroidal anti-inflammatory (NSAIDs) drugs inhibit the synthesis of prostaglandins by inhibiting the enzyme cyclooxygenase.

• Zileuton inhibits the enzyme lipoxygenase thereby decreasing the synthesis of leukotrienes.

Chemically, all prostaglandins are considered to be derivatives of prostanoic acid, having a five-carbon ring (Option C).

The image given below shows the synthesis of various eicosanoids from arachidonic acid:



Solution to Question 11:

Beta lactams act on step 4 in the given image.

Mechanism of action of beta-lactams: They are the cell wall synthesis inhibitor. The β-lactam antibiotics inhibit this last step in peptidoglycan synthesis. They inhibit the enzyme transpeptidase which is required for cross-linking of the peptidoglycan chain. Transpeptidases cleave the terminal D-alanine of the peptide chains and form cross-linkages between them. This cross-linking provides rigidity and stability to the cell wall.

Transpeptidases and other related proteins constitute penicillin-binding proteins(PBP) and, are located in the bacterial cell membrane. β-lactam antibiotics are bactericidal. This is because bacterias swell and burst due to deficient cell walls when they divide in the presence of β-lactam antibiotics. Penicillins, cephalosporins, monobactams, and carbapenems are examples of β-lactam antibiotics.

Gram-positive bacteria are more susceptible to β-lactam antibiotics than gram-negative because their cell wall is almost entirely made of peptidoglycan.

Resistance to β-lactam antibiotics develops due to the production of the beta-lactamase enzyme and the development of altered penicillin-binding proteins like in methicillin-resistant Staphylococcus aureus (MRSA). β-lactamases inactivate β-lactam antibiotics by opening the β-lactam ring. This could be overcome by using drugs that inhibit this enzyme. Examples of β-lactamase inhibitors are clavulanic acid, sulbactam, and tazobactam.

Other options:

Option A: Fosfomycin inhibits step 1 in the image. It inhibits the cytoplasmic enzyme enol pyruvate transferase that catalyses the initial step in the bacterial cell wall and hence cell wall synthesis. It is used for the treatment of urinary tract infections.

Option B: Bacitracin inhibits step 2 in the image. It is a cell wall synthesis inhibitor that interferes with dephosphorylation in the cycling of the lipid carrier which transfers peptidoglycan subunits to the growing cell wall. It is only given for topical application because of its severe nephrotoxicity when given systemically.

Option C: Vancomycin inhibits step 3 in the image. It binds to the D-Ala-D-Ala terminus of the newly formed peptidoglycan pentapeptide. This prevents further elongation of the peptidoglycan and its cross-linking. It is used against MRSA and Clostridium difficile infection. Redman syndrome due to histamine release following infusion is an important side effect.

Note: The FDA has approved the use of cefepime-enmetazobactam, a combination of a fourth-generation cephalosporin and beta-lactamase inhibitor for use in complicated UTIs (including pyelonephritis).

Solution to Question 12:

Phospholipase C acts via the Gq type of G protein-coupled receptors.

G protein-coupled receptors (GPCRs) are cell membrane receptors that span across the membrane 7 times. They also serve as receptors for various peptide hormones, neurotransmitters, amino acids, etc.

Subtypes of GPCRs are distinguished by their alpha subunits. The important subtypes of GPCRs are Gs, Gp, Gq, and Go.

GPCRs act via various second messengers. The important ones with their action on the effector are:

• Gs causes activation of adenylyl cyclase and results in Ca2+ channel opening.

• Gi causes inhibition of adenylyl cyclase and results in K+ channel opening.

• Go causes Ca2+ channel inhibition.

• Gq causes activation of phospholipase C which generates the second messengers inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG). The water-soluble IP3 diffuses to the cytosol and mobilizes Ca2+ from endoplasmic reticular depots.

The given below image shows the effects of phospholipase C after activation by Gq



Solution to Question 13:

Isoniazid is the first-line antitubercular drug that acts by inhibiting the synthesis of mycolic acid.

Isoniazid is a first-line bactericidal drug that is used for the treatment of tuberculosis. This drug enters bacilli by passive diffusion. It then gets activated to its toxic form by the enzyme catalase-peroxidase(the enzyme encoded by the Kat gene).

Normally, InhA and KasA are involved in the synthesis of mycolic acid. Upon activation, the reactive metabolite of isoniazid forms complexes with NAD that inhibit InhA and KasA. Hence, isoniazid inhibits the production of mycolic acid.

The reactive isoniazid metabolite also forms complexes with NADP. This complex inhibits mycobacterial dihydrofolate reductase which results in the interruption of DNA synthesis.

Side effects of isoniazid include:

• Peripheral neuritis due to interference with the production of active coenzyme pyridoxal phosphate from pyridoxine. Therefore, pyridoxine 10mg/day (vitamin B6) is given prophylactically along with isoniazid to prevent neurotoxicity.

• Hepatitis - isoniazid must be stopped at the first sign of hepatotoxicity.

• Rashes

• Anaemia

• Arthralgia

Toxicity of isoniazid results in metabolic acidosis, seizures, and coma.

Other options:

Option B: Rifampin acts by binding to DNA-dependent RNA polymerase to form a stable drug-enzyme complex. This complex suppresses RNA synthesis. Side effects of rifampin are hepatitis, flu-like illness, orange-red colored urine, and body secretions.

Option C: Pyrazinamide is a prodrug, it is converted into an active metabolite pyrazinoic acid which makes the intracellular environment acidic and collapses the transmembrane proton motive force, thereby killing the bacteria. Side effects include hyperuricemia and hepatotoxicity.

Option D: Ethambutol inhibits arabinose transferase III, thereby decreasing arabinogalactan biosynthesis. This results in disruption in mycobacterial cell wall synthesis. Optic neuritis is an important side effect of ethambutol. It results in decreased visual acuity and loss of red-green discrimination.

Solution to Question 14:

Bradycardia is the possible presenting feature which led to the diagnosis of muscarinic poisoning.

Muscarinic poisoning occurs due to the consumption of poisonous mushroom-like Inocybe. Muscarinic symptoms appear within an hour of eating the mushroom. These symptoms (along with nicotinic symptoms) can also be seen in organophosphate poisoning.

There are 5 subtypes of muscarinic receptors they are M1, M2, M3, M4, and M5. The effects exerted by the M receptors are given below:

• M1: Increased seizure activity.

• M2: Tremors, bronchoconstriction, bradycardia, abdominal cramping

• M3: Increased bladder contraction, miosis, increased salivary secretion

• M4: Cataplexy(sudden muscle weakness)

Atropine is used for the management of muscarinic poisoning. It competitively blocks the acetylcholine at the muscarinic receptors.

Solution to Question 15:

Valproate should be avoided for being highly teratogenic to the fetus.

Valproate is an anti-epileptic medication. It is also used as a mood stabilizer in manic episodes as well as in bipolar disorder. The repetitive firing caused by depolarization of cortical or spinal cord neurons is inhibited by valproate. It also prolongs the sodium channel inactivation.

The side effects of valproate (mnemonic: VALPROATE)

• Vomiting

• Alopecia

• Liver toxicity

• Pancreatitis

• Rash

• Obesity

• Anorexia

• Teratogenicity

• Oedema

It causes congenital anomalies like neural tube defects (spina bififda), clefts, and cardiac anomalies. The intelligence quotient score of school-going children who were exposed in utero to valproate is found reduced. It also contributes to poor cognitive development in children. Folate supplementation reduces the chances of neural tube defects.

The images below show a baby and a USG with spinal bifida:



Other options:

Option B: Lithium is used for the treatment of bipolar disorder as well as mania. It was earlier implicated in causing Ebstein anomaly, however, recent studies suggest that lithium has a weak teratogenic effect.

Option C: Oxcarbazepine was earlier associated with teratogenicity and fetal malformations, however, recent studies suggest that oxcarbazepine monotherapy during pregnancy does not carry a significant risk of fetal malformations.

Option D: Olanzapine is an antipsychotic and an antimanic and it is not associated with teratogenicity.

Solution to Question 16:

True statements regarding the administration of 5 alpha-reductase inhibitors are 1,4. They reduce dihydrotestosterone levels and decrease prostate-specific antigen (PSA) levels.

5 alpha-reductase inhibitors are the competitive inhibitors of the enzyme 5 α-reductase which converts testosterone into the more active dihydroxy testosterone (DHT). This enzyme inhibition results in:

• Decreased DHT levels

• Decreased PSA levels

• Reduces prostate volume

However, plasma testosterone levels and LH (luteinizing hormone) levels remain unchanged

Finasteride and dutasteride are the 5 α-reductase drugs currently in use. They are used in benign prostatic hyperplasia (BPH) along with surgery and adrenergic α1 blockers for symptomatic relief. They can also be used in male pattern of baldness and hirsutism.

Side effects of 5 α-reductase drugs are:

- Impotence
- Decreased libido
- Gynecomastia
- Skin rashes

Solution to Question 17:

This type of drug is called a misbranded drug. A drug stating that 500mg of paracetamol is present in each tablet, but contains only 200mg is a misbranded drug, according to the Drugs and Cosmetics Act, 1940 definition of misbranded drugs.

According to the Drugs and Cosmetics act, 1940, a misbranded drug is a drug if its label or container or anything accompanying the drug bears any statement, design, or device which makes any false claim for the drug or which is false or misleading in any particular.

Other options:

Option A: Spurious drugs include drugs manufactured under another drug or manufacturer's name, imitations of other drugs, and drugs with labels from a fictitious company or individual. It also includes drugs substituted partly or wholly by another drug or substance.

Option B: Adulterated drugs include those drugs that contain filthy, or decomposed substances, substances that are injurious to health, or if the drug harmful substances to reduce the quality or strength of the drug. It also includes drugs that were prepared in unsanitary conditions.

Option C: The term unethical drug is not standard terminology in pharmacology. However, the term ethical drug is sometimes used to describe those drugs that are available only by prescription of a medical practitioner.

Solution to Question 18:

Canagliflozin is associated with an increased risk of fractures.

Canagliflozin is a sodium-glucose transporter-2 (SGLT2) inhibitor. SGLT2 is a Na+-glucose cotransporter that is located in the proximal convoluted tubule. They reabsorb glucose and sodium from the tubular lumen back into the blood through the epithelial cells. SGLT2 inhibitors block the transport of glucose in the proximal tubule and promote glycosuria and hence, lowering blood glucose.

The currently available SGLT2 inhibitors are canagliflozin, dapagliflozin, and empagliflozin. Other than decreasing blood glucose they also decrease blood pressure and, cause weight loss of 2–4 kg. Empagliflozin and canagliflozin reduce the risk for major cardiovascular events.

The main side effect of SGLT2 inhibitors is an increase in lower urinary tract infections and genital mycotic infections. This is because of their glycosuric properties. A recent warning from the FDA says that SGLT2 inhibitors may increase the risk of fractures. They also affect circulating levels of parathyroid hormone, and 1,25-hydroxy vitamin D thereby

affecting the mineral balance in bone.

Other options:

Option A: Rosiglitazone is a thiazolidinedione that acts as a peroxisome proliferator-activated receptor γ agonist in the adipose tissue which increases insulin sensitivity and adipose differentiation. Side effects are weight gain, edema, and increased risk of fractures. Compared to rosiglitazone, canagliflozin is commonly used in clinical practice. American diabetes association recommends canagliflozin as the first-line drug along with metformin, in patients with a high risk of cardiovascular disease, renal failure, and heart failure. Even when thiazolidinediones are required, clinicians prefer pioglitazone over rosiglitazone due to the higher risk of cardiovascular-related mortality in rosiglitazone. Therefore, canagliflozin would be a better option compared to rosiglitazone.

Option B: Liraglutide is a long-acting glucagon-like peptide-1 agonist. Nausea, diarrhea, and acute pancreatitis are some frequent side effects.

Option C: Voglibose reduces the intestinal absorption of starch, dextrin, and disaccharides by inhibiting the action of α-glucosidase in the intestinal brush border. The most prominent adverse effects are malabsorption, flatulence, diarrhea, and abdominal bloating.

Solution to Question 19:

The active metabolite of cyclophosphamide is 4 - hydroxy cyclophosphamide.

Cyclophosphamide is an alkylating agent with antitumor and immunosuppressant properties. It is used for many solid tumors. The important side effects are cystitis due to metabolite acrolein and bladder carcinoma.

Cystitis caused by cyclophosphamide is treated by systemic administration of mesna and by irrigating the bladder with acetylcysteine. Both these drugs contain –SH which detoxifies the toxic metabolites. Plenty of fluids and frequent bladder voiding are also helpful.

Solution to Question 20:

Alkalinization of urine will be helpful in the removal of aspirin, which is a weakly acidic drug and is susceptible to ion trapping in the urine.

Aspirin is a weak acid, with a pKa of 3.0. Drugs that are subjected to renal clearance are excreted into the urine by glomerular filtration and active tubular secretion. Nonionized compounds are reabsorbed more rapidly than ionized polar molecules. When the pH of the urine increases, more salicylate is in its ionized form at equilibrium. Hence, salicylic acid diffuses into the tubular lumen of the kidney. Urine alkalinization is recommended as a first-line treatment only for moderately severe salicylate poisoning that does not meet the criteria for hemodialysis.

Alkalinization of the urine is achieved by administration of 100–150 mEq of sodium bicarbonate in 1 L of 5% dextrose in water (D5W). It is infused intravenously at twice the maintenance fluid requirements.

Alkalinization of urine is contraindicated in renal failure or if fluid administration may worsen pulmonary edema or congestive heart failure.

Solution to Question 21:

A hospital conducting a double-blinded study in multiple centers to check the efficacy of statins describes phase 3 of the clinical trial. A phase 2 clinical trial is generally conducted at a few centers (2-4) by physicians.

Phase 3 of a clinical trial is also known as a multisite trial. It is conducted on a large patient population (500–3000) by several physicians at multiple centers. This phase involves patients who have been diagnosed with the disease that the drug claims to treat. Its purpose is to confirm the efficacy of the newly developed drug in a larger population. Phase 3 is randomized, controlled, and may be blinded.

Solution to Question 22:

The given clinical scenario and the image showing a pin-point pupil (miosis) suggest opioid overdose. The true statement among those given is that it is caused by activation of the Edinger-Westphal nucleus.

Edinger-Westphal nucleus is a part of the light reflex pathway and is located in the midbrain. This nucleus is responsible for causing pupillary constriction when the retina is exposed to light.

Opioid overdose symptoms are a triad of coma, respiratory depression, and pinpoint pupils (miosis). Other features are hypotension, bradycardia, shallow breathing, and hypothermia.

Tolerance to miosis does not develop even in addicts who develop a high tolerance to opioids. Miosis is induced by all opioid agonists.

Naloxone is the drug of choice for opioid overdose. It is administered after adequate ventilation is established. Airway protection and ventilatory maintenance are the most important steps in the treatment of opioid overdose; this is because respiratory depression is the major cause of mortality in such cases. Single-dose activated charcoal can be administered if the opioid ingestion occurred within one hour.

Levallorphan is an opioid antagonist, with properties similar to those of naloxone. It can be used for opioid overdose. It reverses severe opioid-induced respiratory depression.

Phenylephrine is an α1-selective agonist that causes mydriasis, but it is not useful in the reversal of opioid-induced miosis.

Solution to Question 23:

The correct answer is option B: A-4, B-1, C-3, D-2

In a patient with head injury, morphine is contraindicated because it increases intracranial tension due to CO■ retention, precipitates respiratory depression, and may cause vomiting, miosis, and altered mental state which interferes with assessment of mental state.

Amiodarone should not be used in patients with prolonged QT interval because it may cause marked prolongation of the QT interval, bradycardia, and torsades de pointes.

Vigabatrin, an inhibitor of GABA-transaminase, an anti-convulsant, is contraindicated in pregnancy due to its potential teratogenic effect.

Estrogen preparations can contribute to the incidence and severity of thromboembolism, particularly if other risk factors are present. The risk of serious cardiovascular side effects is particularly marked in women more than 35 years of age who smoke heavily.

A. Morphine	4. Head Injury
B. Amiodarone	1. QT prolongation
C. Vigabatrin	3. Pregnancy
D. Estrogen preparations	2. Thromboembolism

Solution to Question 24:

1 and 2 are true statements regarding memantine.

Memantine is a non-competitive NMDA receptor antagonist. It inhibits glutamate excitotoxicity. It delays the progression of dementia in Alzheimer's disease. Commonly, it is started in patients with moderate-to-severe disease.

It can be given both as monotherapy or can be given along with cholinesterase inhibitors. It can be started at a dose of 5 mg once daily and then progressively increased up to 10 mg twice daily for a minimum of 6 months.

It can cause dizziness and headache. Since it is excreted in the kidney, careful dose adjustments should be made in patients with renal failure.

Solution to Question 25:

Step 5 in the given life cycle shows the integration of the viral DNA into the host genome. Raltegravir is an integrase inhibitor and is active against both HIV-1 and HIV-2. It can be used in adults and children >1 month old.

It is advised for patients to consume antacids and sucralfate, 6 hours before or 2 hours after raltegravir intake.

It may cause headaches, nausea, weakness and fatigue. Rarely, it can lead to drug reactions with eosinophilia and systemic symptoms (DRESS) syndrome.

The steps involved in the lifecycle of HIV as given in the image are described as follows:

• Step 1: Binding of HIV-1 to the cell membrane of a CD4+ cell

• Step 2: HIV-1 enters the cell via interaction with the membrane glycoproteins

• Step 3: Reverse transcription of the viral genetic material using the virus' reverse transcriptase enzyme

• Step 4: The HIV-1 genome travels into the nucleus

• Step 5: The viral genome integrates into the host's DNA through the integrase enzyme

• Step 6: The HIV-1 proviral genome undergoes transcription with the host DNA

• Step 7: The viral messenger RNA (mRNA) undergoes translation to produce new viral proteins

• Steps 8 and 9: Assembly and maturation of the virion



Other options:

Option B: Enfuvirtide acts at step 2 in the given image. It blocks gp41 of the cell membrane and blocks the entry of the virus into the cell. It prevents the free viral particles from invading new CD4+ cells. It is effective against HIV-1.

Option C: Tenofovir acts at step 3 in the given image. It is a nucleotide reverse transcriptase inhibitor (NRTI) that is effective against HIV-1,2 and hepatitis B infections. It is used in adults and children >2 years old. It can also be used for pre-exposure prophylaxis.

Option D: Zidovudine also acts at step 3 in the given image. It is an NRTI and is effective against HIV-1,2 and Human T-cell Lymphoma Virus (HTLV). It can be used in adults and children. It can also be used to prevent mother-to-child transmission.

Solution to Question 26:

Phase 4 clinical trials are also called post-marketing surveillance is the correct statement.

Other options:

Option A: New drug application is done after phase 3 of clinical trials.

Option C: Phase 3 trials may or may not be blinded.

Option D: Phase 2 trials are the first round of trials in patients. Phase 1 trials are done in healthy volunteers.

Solution to Question 27:

The correctly matched pairs of teratogenic drugs and their effects is 1-d, 2-c, 3-b, 4-a.

Lithium use in early pregnancy causes Ebstein anomaly in the fetus. Fetal lithium exposure is associated with neonatal goiter, CNS depression, hypotonia (floppy baby syndrome), and cardiac murmur.

Gray baby syndrome occurs in neonates, especially premature ones, exposed to high doses of chloramphenicol (100 mg/kg), which are administered prophylactically.

Fetal warfarin syndrome or warfarin embryopathy presents with CNS abnormalities, depressed nasal bridge/ saddle nose at birth, bleeding diathesis, and growth retardation. Hypoplasia of the nose, eye sockets, and long bones with stippling of epiphyses are also seen.

Fetal thalidomide exposure causes phocomelia. This disorder is associated with underdeveloped limbs and absent pelvic bones. Both upper and lower limbs are affected, and a part of the limbs may be missing. Fused fingers and missing thumbs are also seen. Thalidomide was originally used for the treatment of pregnancy-associated morning sickness but was withdrawn due to this adverse effect. It re-entered the market for the treatment of erythema nodosum leprosum. It also has antiangiogenic and immunomodulatory effects.

Solution to Question 28:

The drug used to manage this condition is prazosin as it can counteract the effects of breathlessness most probably caused by pulmonary edema.

The venom is a potent autonomic stimulator that causes the release of massive amounts of catecholamines from the adrenals. It also has a direct effect on the myocardium.

It can present as neurotoxicity with restlessness or seizure-like activity, abnormal oculomotor function, and loss of pharyngeal muscle control that leads to respiratory compromise. Tongue fasciculations and hypersalivation are also common. Cardiovascular toxicity from systemic envenomation includes tachycardia, hypertension, pulmonary edema, and cardiogenic shock.

If the scorpion has hemolytic venom, the reaction is mainly local, with edema, pain, and reddening. This usually lasts for one to two hours.

Treatment:

• Pain - Paracetamol, NSAIDs, or opioids

• Sedation for agitation, oculomotor abnormalities, uncoordinated neuromuscular activity, and muscle spasms - Benzodiazepines

• Hypertension and pulmonary edema - Prazosin counteracts the adrenergic surge that can lead to these effects

• Cardiogenic shock - Antivenom, prazosin, nitroglycerin, dobutamine

• Cholinergic effects - Atropine

• Supportive care, wound care

Solution to Question 29:

Phase 4 in the graph indicates the nicotinic action of acetylcholine at higher doses.

In phases 1 and 2, acetylcholine (Ach) acts on muscarinic receptors (at 2 micrograms and 50 micrograms) bringing down blood pressure. Atropine blocks the muscarinic receptor, so this action of acetylcholine is not seen in phase 3. However, in phase 4, higher doses of Ach (like 5 mg) act via the nicotinic receptor, shooting up the blood pressure. This phenomenon is called the reversal of Ach in blood pressure and was studied by Dale.

Nicotinic receptors are ligand-gated ion channels. It can be divided into two groups: muscle receptors (NM), which are found at the skeletal neuromuscular junction, and neuronal receptors (NN), which are found in the ganglia and CNS. When stimulated at higher doses, Ach acts on NN receptors, leading to the stimulation of sympathetic ganglia and adrenal medulla. This increases the heart rate, blood pressure, and myocardial contractility.

Muscarinic receptors are G-coupled protein receptors involved in the parasympathetic nervous system. M1, M2, M3, M4, and M5 are the receptor subtypes. They act via Gi or Gq subunits. When Ach is administered, it acts on muscarinic receptors, primarily the M3 receptors, which are located on vascular endothelial cells. It stimulates guanylyl cyclase and promotes relaxation via a cyclic GMP–dependent mechanism. At higher doses, it acts on M2 receptors, which have direct effects on the heart. It also inhibits the release of norepinephrine from sympathetic nerve terminals and decreases cyclic AMP levels.

Solution to Question 30:

The most likely drug causing hyperpigmentation of the hands and soles would be emtricitabine.

Emtricitabine is an anti-retroviral drug that acts by inhibiting nucleoside reverse transcriptase (NRTI). It is well-tolerated and has minimal toxic effects. Adverse effects include hyperpigmentation of the skin (especially sun-exposed areas, palms, and soles), headache, vomiting, and nausea.

Other options:

Option A: Nelfinavir is a protease inhibitor that causes diarrhea or loose stools. It usually resolves within a month of starting the drug.

Option C: Abacavir is an NRTI that can cause a life-threatening hypersensitivity syndrome. The syndrome presents with fever, abdominal pain, and maculopapular rash within 6 weeks of initiating abacavir. The drug must be immediately discontinued and must not be administered to the patient thereafter. Rarely, there might be respiratory and musculoskeletal side effects.

Option D: Zidovudine is an NRTI that causes malaise, fatigue, myalgia, anorexia, and insomnia. Chronic use leads to nail hyperpigmentation. It can also cause bone marrow suppression.

Solution to Question 31:

Dorzolamide is a topical ocular carbonic anhydrase inhibitor.

It acts on carbonic anhydrase in the ciliary processes of the eye and mediates the formation of HCO_3^- in aqueous humor. Inhibition of carbonic anhydrase decreases the rate of formation of aqueous humor and consequently reduces intraocular pressure.

The major indication for carbonic anhydrase inhibitors is open-angle glaucoma. Dorzolamide and brinzolamide, which are available only as ophthalmic drops, are used for this purpose.

Dorzolamide may be used for secondary glaucoma and in acute angle-closure glaucoma to lower intraocular pressure before surgery.

Solution to Question 32:

Sildenafil is a phosphodiesterase 5 inhibitor and the action of the drug is mediated through cGMP. Erectile dysfunction is treated by phosphodiesterase-5 (PDE-5) inhibitors like sildenafil, vardenafil, and tadalafil.

The corpus cavernosum produces nitric oxide during arousal that stimulates the formation of cGMP. cGMP causes relaxation of the smooth muscle of the corpus cavernosum and penile arteries, engorgement of the corpus cavernosum, and erection.

The above-mentioned drugs compete for cyclic GMP, binding at the site of cyclic GMP hydrolysis on PDE5, enhancing the accumulation of cGMP and hence used in the management of erectile dysfunction. They are useful in treating pulmonary arterial hypertension.

Adverse effects of the drug are headache, flushing, nasal congestion, nasopharyngitis, and dyspepsia. Sildenafil and Vardenafil also cause retinal PDE-6 inhibition and can result in blurring of vision and blue-green tinting of the vision.

Solution to Question 33:

The correct match is as follows:

Acetazolamide, a carbonic anhydrase inhibitor, acts on the enzyme in epithelial cells of the proximal convoluted tubule. Acetazolamide is mainly used in patients with secondary glaucoma to reduce intra-ocular pressure, provide symptomatic relief in acute mountain sickness, and as

treatment for absence seizures.

Torsemide, a loop diuretic, inhibits the Na+-K+-ATPase symporter in the thick ascending loop of Henle. It is used in acute pulmonary edema, edema associated with congestive cardiac failure, hypertension with renal insufficiency, and hyponatremia.

Chlorthalidone, a thiazide-like diuretic, inhibits the Na+-Cl- symporter in the distal convoluted tubule. It is used in hypertension and in edema associated with congestive cardiac failure, hepatic cirrhosis, nephrotic syndrome, and chronic renal failure.

Vasopressin, also known as anti-diuretic hormone, is secreted by the posterior pituitary and mainly acts on the collecting duct, and increases the reabsorption of water (anti-diuresis). Vasopressin or its agonists (such as desmopressin) are used in cases of diabetes insipidus, type I and type II von Willebrand's disease, primary nocturnal enuresis, and postoperative ileus.

Column A	Column B
A. Acetazolamide	4. Proximal convoluted tubule
B. Torsemide	1. Thick ascending loop of Henle
C. Chlorthalidone	3. Distal convoluted tubule
D. Vasopressin	2. Collecting duct

Solution to Question 34:

Alkalinization of urine will be helpful in the removal of weakly acidic drugs as they are susceptible to ion trapping in the urine.

Drugs that are subjected to renal clearance are excreted into the urine by glomerular filtration and active tubular secretion. Nonionized compounds are reabsorbed more rapidly than ionized polar molecules. When the pH of the urine increases, more salicylate is in its ionized form at equilibrium. Hence, salicylic acid diffuses into the tubular lumen of the kidney. Urine alkalinization is recommended as a first-line treatment only for moderately severe salicylate poisoning that does not meet the criteria for hemodialysis.

Alkalinization of the urine is achieved by administration of 100–150 mEq of sodium bicarbonate in 1 L of 5% dextrose in water (D5W). It is infused intravenously at twice the maintenance fluid requirements.

Alkalinization of urine is contraindicated in renal failure or if fluid administration may worsen pulmonary edema or congestive heart failure.

Solution to Question 35:

Hyponatremia occurs more frequently in the case of oxcarbazepine than carbamazepine.

Both carbamazepine and oxcarbazepine are sodium-channel blockers.

Carbamazepine is most effective in complex partial seizures. It is also used in GTCS and simple partial seizures, trigeminal and related neuralgias, and manic depressive psychosis.

Carbamazepine potentiates the action of antidiuretic hormone (ADH). It can result in water retention with decreased osmolarity and hyponatremia, especially in elderly patients with cardiac disease. Carbamazepine and oxcarbazepine can cause leukopenia, aplastic anemia, hepato-toxicity and hyponatremia in the elderly.

Oxcarbazepine shows lesser hypersensitivity reaction. Hence, rashes are more commonly seen with carbamazepine than with oxcarbazepine.

Studies have shown that carbamazepine elevates enzyme gamma-glutamyl transpeptidases (GGT) more frequently than oxcarbazepine because of enzyme P450 induction.

Mechanism of action of some common antiepileptics:

Mechanism of action	Antiepileptics
Sodium-channel blockers	PhenytoinCarbamazepine, Oxcarbazepine Eslicarbazepine LamotrigineTopiramateFelbamateValproateRufinamideLevetiracetam
Calcium-channel blockers	ValproateLamotrigineEthosuximide
GABAAreceptor modulators	BenzodiazepinesPhenobarbitalFelbamatePrimidoneTopiramateCarbamazepine, OxcarbazepineStiripentolClobazam
GABA uptake inhibitors	TiagabineVigabatrin
NMDA receptor antagonist	Felbamate
AMPA/Kainate receptor antagonist	PhenobarbitalTopiramatePerampanel
HCN channel activity enhancers	Lamotrigine
Inhibitors of brain carbonic anhydrase	AcetazolamideZonisamideTopiramate
Potassium-channel openers	RetigabineTopiramate

Solution to Question 36:

The correct pairs are as follows:

Trastuzumab is a humanized monoclonal antibody against the external domain of HER2/neu. It is indicated as an adjuvant in the management of metastatic HER-2-positive breast cancer.

Infliximab binds with high affinity to TNF-α and prevents the cytokine from binding to its receptors. Infliximab is used for the treatment of rheumatoid arthritis; it is used in combination with methotrexate in patients who do not respond to methotrexate alone. Infliximab is also used for the treatment of the symptoms of moderate-to-severe Crohn's disease.

Sirolimus is an antimetabolic and antiproliferative drug. It acts by forming a complex with an immunophilin called FKBP-12, which in turn inhibits the mammalian target of rapamycin (mTOR). mTOR is a protein kinase that blocks cell cycle progression at the G1 phase to S phase transition. It is used in the prophylaxis of organ transplant rejection.

Imatinib is a first-generation tyrosine kinase inhibitor. Imatinib attacks BCR-ABL translocation and is used in chronic myelocytic leukemia. It also inhibits c-kit and produces extended control of GI stromal tumors.

A. Trastuzumab	4. HER2/neu
B. Infliximab	3. TNF α
C. Sirolimus	2. mTOR
D. Imatinib	1. BCR-ABL tyrosine kinase

Solution to Question 37:

Oxazepam can be used in patients with elevated transaminases. It is short-acting and so, can be used in patients with liver impairment or brain damage.

Oxazepam is an orally available benzodiazepine used to treat anxiety and alcohol withdrawal states. Hepatic oxidation is decreased in persons with liver disease and the elderly. Oxazepam and lorazepam undergo only hepatic glucuronidation.

Other options:

Option A: Midazolam is the benzodiazepine of choice for induction of anesthesia. The onset of anesthesia is within 30 to 60 seconds. It can also be used to control seizures, but there is no difference in the efficacy of intravenous midazolam and intravenous diazepam.

Option B: Diazepam is a long-acting benzodiazepine. It is metabolized in the liver to active metabolites like nordazepam which are long-acting. Diazepam and its long-acting metabolites can accumulate significantly in elderly patients with liver disease and therefore should not be used in such cases.

Option C: Alprazolam is metabolized by hepatic oxidation which is significantly decreased in liver impairment and elderly patients.

Solution to Question 38:

Atorvastatin 80 mg can be used to treat this condition. The first line of therapy in controlling cholesterol in patients with atherosclerotic cardiovascular disease (ASCVD) is starting a high-intensity statin. High-intensity statins include atorvastatin (40-80 mg) and rosuvastatin (20-40 mg).

Statins are competitive inhibitors of HMG-CoA reductase. This leads to a decrease in cholesterol biosynthesis. Statins are also effective in reducing LDL and triglyceride levels.

Other options:

Option A. Fenofibrate reduces lipoprotein levels by interacting with Peroxisome proliferator-activated receptor (PPAR-α) and stimulating lipoprotein lipase. They are used as drugs of choice for treating hyperlipoproteinemia and severe hypertriglyceridemia (>1000 mg/dL).

Option B. Rosuvastatin and fenofibrate combination was previously used to reduce atherosclerotic cardiovascular disease risk. But recent studies have indicated no change in reduction and hence is not recommended anymore.

Option C: Rosuvastatin 10 mg is a low-intensity statin and is mainly used for primary prevention against atherosclerotic cardiovascular disease.

Solution to Question 39:

Quinidine is a class IA anti-arrhythmic agent. Class 1a agents are sodium channel blockers with significant potassium channel blocking properties. This slows conduction and prolongs the QRS duration and action potential duration.



Lignocaine and tocainide (Options D and A) are class 1b anti-arrhythmic drugs. These are sodium channel blockers that inhibit both inactivated and open sodium channels and can open potassium channels. It reduces phase 4 slope and reduces automaticity. It also alters the excitation threshold.

The below image shows the changes in the action potential of a cardiac muscle when a class 1b anti-arrhythmic drug is used.



Class 1c agents like flecainide and propafenone block sodium channels with insignificant potassium channel-blocking properties.

The image below illustrates the action potential changes when a class 1c anti-arrhythmic is used.



Option B: Propranolol is a β-receptor-blocker and a class II antiarrhythmic.

Solution to Question 40:

The false statement regarding statins is that they can be given with verapamil and other enzyme inhibitors.

Statins like atorvastatin and rosuvastatin are metabolized by the cytochrome 3A4 enzymes of the liver. Verapamil is a cytochrome 3A4 inhibitor and may interfere with statin degradation. This leads to an increase in serum statin levels. The most important feature of statin toxicity is myopathy. It can present as myalgia but can sometimes also lead to fatal rhabdomyolysis.

Statins are the most commonly used hypolipidemic drugs. Statins act by inhibiting the rate-limiting HMG-CoA reductase enzyme, which in turn inhibits cholesterol synthesis. They decrease levels of low-density lipoproteins (LDL) and triglycerides and increase the levels of high-density lipoproteins (HDL) in plasma.

Other options:

Option A: Lipoprotein (lp) a is an LDL molecule that has apolipoprotein b and an additional apolipoprotein a. Studies indicate that these molecules are atherogenic and may independently cause cardiovascular disease. Although HMG-CoA reductase inhibitors substantially reduce the risk of cardiovascular events, there is a mild increase in lipoprotein a levels.

Option B: Statins can be continued during surgery. These drugs have proved to decrease overall mortality due to cardiovascular events. Statins also influence inflammatory responses and can be safely used in trauma.

Option D: Chronic administration of high statin doses also leads to an increased incidence of type 2 diabetes mellitus.

Solution to Question 41:

The drugs marked A, B and C are aspirin, eptifibatide, and vorapaxar, respectively.

Aspirin causes inhibition of cyclooxygenase-1 (COX-1), which would inhibit the formation of thromboxane A2. Thromboxane A2 is an inducer of platelet aggregation and is a potent vasoconstrictor. Hence, inhibition of its formation is responsible for antiplatelet action.

Aspirin is a non-competitive, irreversible inhibitor of the cyclooxygenase (COX) enzyme. It is a nonselective COX inhibitor and inhibits both COX-1 and COX-2.

Abciximab, tirofiban, and eptifibatide act by blocking Gp IIb/IIIa receptors. They are mainly used during percutaneous coronary intervention for acute coronary syndrome. They carry a high risk of bleeding and can cause thrombocytopenia.

Vorapaxar primarily inhibits platelet aggregation caused by thrombin. PAR-1 activation by thrombin contributes to platelet aggregation. Vorapaxar is a competitive antagonist of protease-activated receptor-1 (PAR-1). Vorapaxar is indicated in combination with aspirin and/or clopidogrel to reduce thromboembolic events in patients with MI or peripheral vascular disease.

Prasugrel is an irreversible P2Y12 inhibitor and lasts for the entire lifetime of a platelet.

Prasugrel is a prodrug that is used in patients with acute coronary syndrome who have undergone percutaneous coronary intervention. It reduces the rate of thrombotic cardiovascular events in such patients.



Solution to Question 42:

The wrong statement/s regarding disulfiram is only 3.

Disulfiram is an irreversible inhibitor of both cytosolic and mitochondrial forms of aldehyde dehydrogenase. It is used in behavioral modification for alcohol de-addiction. The principle of action is aversive conditioning (not anti-craving).

Disulfiram can be given only to highly motivated individuals. It should not be administered until the person has abstained from alcohol for at least 12 hours. The disulfiram-alcohol reaction may persist as long as 14 days after the last dose of disulfiram.

Alcohol consumption during disulfiram therapy can lead to increased concentrations of acetaldehyde. This can cause aversive reactions, resulting in flushing, pulsating headache, throbbing pain in the head and neck, nausea, vomiting, rashes, respiratory difficulties, chest pain, and autonomic manifestations.

Disulfiram is usually given in the dose of 250 mg/day (range of 125–500 mg/day).

Solution to Question 43:

G6PD deficiency will manifest with primaquine.

G6PD deficiency is an X-linked recessive (XLR) disorder that causes congenital hemolytic anemia due to enzyme deficiency. Hemolysis occurs in conditions of high oxidative stress that are triggered by infections, fava beans, and certain drugs.

G6PD is an enzyme in the hexose monophosphate pathway (HMP) that maintains the cellular levels of NADPH. NADPH then provides reducing equivalents needed for the conversion of oxidized glutathione to reduced glutathione. Reduced glutathione helps neutralize free radicles, protecting against oxidant injury.

Clinically, most patients are asymptomatic. Some patient develops episodic hemolysis, causing hemolytic anemia, hemoglobinemia, and hemoglobinuria.

Both primaquine and chloroquine cause hemolysis in G6PD-deficient individuals, but hemolysis is rare with chloroquine and it can be used in patients with G6PD deficiency, whereas primaquine is contraindicated in these individuals.

Hematological findings in G6PD deficiency are shown below:



©MARROW



©MARROW

Drugs that carry a risk of clinical hemolysis in persons with G6PD deficiency:

CLASS	DRUGS
Antimalarials	Primaquine >ChloroquineDapsone/chlorproguanil
Sulphonamides/sulphones	SulfamethoxazoleSulfasalazine, Sulfisoxazole

CLASS	DRUGS
Antibacterial/antibiotics	Cotrimoxazole,Ciprofloxacin, Chloramphenicol, Nalidixic a cid, Norfloxacin,Nitrofuranto in
Others	Naphthalene, RasburicaseMe thylene blue

Solution to Question 44:

The correct option regarding the drugs used to treat schizophrenia symptoms is option C.

Drugs used to treat schizophrenia can be classified:

• Typical or 1st generation antipsychotics.

• Atypical or 2nd generation antipsychotics.

All typical antipsychotics markedly improve positive symptoms of schizophrenia, which is attributed to their antagonism at the D2 receptor. Some of the drugs included in this class are chlorpromazine and haloperidol.

Atypical antipsychotics are antagonists or inverse agonists at the 5HT2A receptor. 5HT2A antagonism is responsible for reducing negative symptoms in schizophrenia. Some of the drugs that belong to this class are clozapine, aripiprazole, and brexpiprazole.

Atypical antipsychotics have lower extrapyramidal symptoms as compared to typical antipsychotics as the former has a lesser affinity to the D2 receptor. Muscarinic antagonists can be given to decrease the extrapyramidal symptoms. For example, promethazine or hydroxyzine is given for acute muscular dystonia, which is a form of extrapyramidal symptoms.

5HT1A receptor antagonism, not agonism, has been found to reduce food intake and hence reduce body weight.

Aripiprazole and brexpiprazole are atypical antipsychotics that have partial agonist effects at the 5HT1A receptor. They have been approved for the treatment of schizophrenia and as an adjunctive treatment for depression.

Solution to Question 45:

The given clinical scenario is suggestive of mucormycosis. The patient can be given injection amphotericin B for 8 weeks, followed by oral posaconazole.

Mucormycosis is an opportunistic mycosis. It is typically seen in immunocompromised patients and invades blood vessels.

The rhino-orbital-cerebral is the most common form of mucormycosis. Patients present with rhinitis, eye or facial pain, and facial numbness, followed by the onset of conjunctival suffusion and swelling along with blurry vision.

Diagnosis is made by biopsy with histopathologic examination. The biopsy shows characteristic wide, ribbon-like aseptate hyphal elements that branch at right angles.

The image below shows biopsy findings seen in mucormycosis:



Treatment of the infection includes 4 steps:

• Initiation of therapy (amphotericin B) as early as possible. Posaconazole can also be used in the management of this condition.

• Surgical debridement

• Reversal of underlying conditions and risk factors

• Treatment of underlying malignancy if present

Amphotericin B is a broad-spectrum antifungal. It binds to ergosterol in the fungal membrane and forms pores or channels in the fungal cell membrane. This allows the leakage of cytosolic molecules and ions, leading to a loss of membrane integrity.

Other options:

Option A: Itraconazole can be used for the treatment of chronic aspergillosis if there is resistance or failure of voriconazole. Ketoconazole is used for both dermatophytosis and deep mycosis, but it has been replaced by itraconazole.

Option C: Fluconazole is used for the treatment of candidiasis and cryptococcosis.

Option D: Amphotericin B in combination with 5-flucytosine is used as induction treatment for cryptococcal meningitis.

Solution to Question 46:

The antilipidemic drugs likely to cause gallstones are the peroxisome proliferator-activated receptors (PPARs) that act by activating PPAR alpha.

PPARs are a superfamily of nuclear receptors involved in the regulation of energy homeostasis and metabolic function. It has three subtypes:

• PPARα - reduces triglyceride level

• PPARγ - causes insulin sensitization

• PPARδ - enhances fatty acids metabolism

Lipoprotein lipase (LPL) is a key enzyme in the degradation of VLDL. Fibrates bind to PPAR alpha and cause stimulation of lipoprotein lipase enzyme. Increased LPL synthesis enhances the clearance of triglyceride-rich lipoproteins. Additionally, fibrates also stimulate fatty acid oxidation and reduce the expression of apo C-III (inhibitor of lipolysis). All fibrates increase the lithogenicity of bile, thereby increasing the risk of gall stones

Other options:

Class	HMG- CoA reductase inhibitors (statins)	Niacin	Bile acid-binding resins
Examples	Lovastatin, atorvastatin	Niacin	Colestipol,colesevelam
Mechanism of action	Inhibits HMG-CoA reductase	Decreases the production of VLDLs and the catabolism of HDLs	Binds to bile acids in the intestinal lumen and increases excretion.
Major effects on lipoprotein levels	Reduces LDL levels	Reduces triglycerides, LDL, and lipoprotein a levelsSignificantincrease HDLlevels	Reduces LDL levels
Important side effects	MyopathyElevations of serum amino-transferase activityElevation of creatine kinase activity	Cutaneous vasodilationElevations in amino-transferases levels	ConstipationBloating

Solution to Question 47:

In the given image, drug A & drug B are agonists.

Drugs that bind to physiological receptors and mimic the regulatory effects of endogenous signaling compounds' are agonists. Drugs that are only partially as effective as agonists are partial agonists.

The efficacy of a drug indicates the capacity of a drug to activate a receptor and generate a cellular response. A drug with high efficacy is a full agonist, eliciting a full response at some concentration. A drug with a low intrinsic efficacy will be a partial agonist. Drug A & drug B are agonists (Option C) that have the same efficacy (Option A), although, at different concentrations.

Drug C does not produce a full response at any concentration, making it a partial agonist. In the presence of a full agonist, a partial agonist behaves like a competitive antagonist (Option D).

When two drugs produce equivalent responses, the drug whose dose-response curve lies to the left is said to be the more potent since it requires a lower dose to produce the same effect. Drug A

& drug B produce the same action and level of response in the tissue. The curve of drug A on the left indicates that it produces the same response at a lower concentration, making it more potent than drug B (Option B).

Solution to Question 48:

Dipivefrine and dopexamine are β2-adrenergic receptor agonists.

Dipivefrine causes a decrease in aqueous production and increases outflow. It was used in the treatment of open-angle glaucoma and ocular hypertension but has since been replaced.

Dopexamine is a synthetic analogue related to dopamine with activity at D1 and D2 receptors as well as at β2 receptors. It is used in patients with severe congestive heart failure, sepsis, and shock.

Other options:

B. Fenoldopam is a peripheral D1 agonist with moderate affinity for α2-adrenergic receptors. It is a rapidly acting vasodilator that is used in severe hypertension.

D. Mirabegron is a β3-adrenergic receptor agonist which is used for bladder incontinence. Activation of this receptor in the bladder leads to detrusor muscle relaxation and increased bladder capacity. This action prevents voiding and provides relief for those with an overactive bladder and urinary incontinence.

Solution to Question 49:

Sildenafil acts by inhibiting phosphodiesterase-5 (PDE-5) and is used in the treatment of erectile dysfunction.

PDE-5 inhibitors cause the accumulation of cGMP in the corpus cavernosum, which leads to the relaxation of smooth muscles of penile arteries that fill the corpus cavernosum, thereby improving erectile function in patients with erectile dysfunction.

In addition to erectile dysfunction, sildenafil can be used in the treatment of pulmonary arterial hypertension.

Adverse effects include headache, flushing, nasal congestion, blue-green tinting of vision, and erections lasting more than 4 hours.

Sildenafil is contraindicated in patients using nitrate vasodilators, which could result in dangerously low blood pressure.

Solution to Question 50:

The correct option among the following is A, C, and D.

Bioavailability is defined as the fraction of a drug reaching systemic circulation in an unchanged form after administration by any route. The area under the blood concentration-time curve (AUC) is proportional to the dose and the extent of bioavailability for a drug.

Bioavailability (F) = Quantity of drug reaching systemic circulation/Quantity of drug administered

Bioavailability depends on two factors, absorption and 1st pass metabolism. The bioavailability of the drug injected intravenously is 100%. But it is lower after oral ingestion because of incomplete absorption and first-pass metabolism in the intestinal wall or liver and lies between 5% to <100%. The bioavailability of subcutaneous and intramuscular routes lies between 75% to ≤100%.

The image given below shows the difference between the bioavailability of drugs injected intravenously (IV) and administered per orally (PO). Note the area under the respective curves shaded in red and blue colors which represent the absorption of the drug.



Solution to Question 51:

In this hypertensive patient with hyperkalemia, losartan must be stopped.

Losartan is competitive angiotensin I (AT1) receptor antagonist, which binds to the AT1 receptor with high affinity. Uses of losartan are:

• Antihypertensive

• Renoprotective drug in diabetes mellitus

• Stroke prophylaxis

• Portal hypertension in cirrhosis without compromising renal function

• Anti-aggregant as it is a competitive antagonist of the thromboxane A2 receptor.

Hyperkalemia is an important side effect of ARBs. Losartan appears to be a safe and efficacious agent to lower serum uric acid levels in patients with hyperuricemia.

Option A: Gingival hyperplasia and ankle swelling are side effects of calcium channel blockers like amlodipine. Ankle edema occurs due to increased hydrostatic pressure in the capillaries present in the dependent parts of the body. This is due to reflex constriction of postcapillary vessels. Other common side effects of amlodipine include headache, dizziness, and abdominal pain.

Option B: Metoprolol is a cardioselective beta blocker. Cardioselective beta blockers are less likely to cause hyperkalemia as an adverse effect. The risk of metoprolol-induced hyperkalemia is increased if the patient is also diabetic.

Option D: Furosemide is a loop diuretic. It acts by inhibiting the Na+-K+-2Cl- symporter in the thick ascending limb of the loop of Henle resulting in increased excretion of sodium, potassium, and chloride. Hypokalemia, hyperuricemia, hypocalcemia, and hypomagnesemia are the important side effects.

Solution to Question 52:

Varenicline is the most effective oral drug for smoking cessation.

Nicotine is the principal constituent of tobacco responsible for its addictive behaviour. Varenicline works by blocking the effects of nicotine on the brain. It blocks $\alpha 4$, and $\beta 2$ nicotinic acetylcholine receptor subtypes and inhibits dopaminergic activation produced by smoking through their partial agonism. It decreases the craving and withdrawal syndrome that occurs with cessation attempts. It prevents nicotine stimulation of the mesolimbic dopamine system associated with nicotine addiction, thus helping to occlude the rewarding effect of smoking.

Varenicline is prescribed only to patients 18 years or older. It is taken orally and the dose taken is 0.5 mg once daily on days 1 to 3, increased to 0.5 mg twice daily on days 4 through 7, and 1 mg twice daily after day 8 for 11 weeks. Therapy should start one week before the target quit date. It can be continued for another 12 weeks (1 mg BD) to maintain abstinence.

The adverse effects include nausea, disturbed sleep, sleepwalking, agitation, drowsiness, and constipation, and may also increase the risk of pancreatitis. Varenicline has also been implicated in specific skin syndromes such as Stevens-Johnson syndrome, erythema multiforme, and photosensitivity. Varenicline can cause renal failure and kidney stones, thus renal function must be monitored.

The FDA has issued a warning about mood and behavioural changes associated with its use, and there is some evidence of increased cardiovascular risk.

Cevimeline is a muscarinic agonist with an affinity for both M1 and M3 receptors on the lacrimal and salivary glands. While M1 activation results in increased secretion, M3 activation results in smooth muscle contraction, and expulsion of the secretions. Cevimeline enhances lacrimal secretions in Sjogren's syndrome.

Solution to Question 53:

The correct option is Option C: 1-a, 2-d, 3-b, 4-c

Quinidine is a class IA anti-arrhythmic agent. Class 1a agents are sodium channel blockers with significant potassium channel blocking properties. This slows conduction and prolongs the QRS duration and action potential duration.

Digoxin is a Na+ K+ ATPase inhibitor that works by increasing AV node refractoriness.

Esmolol is a β-receptor-blocker and a class II antiarrhythmic.

Ibutilide is a class III anti-arrhythmic agent. It works by blocking K+ channels.

Solution to Question 54:

The drug that increases trabecular outflow is netarsudil.

Netarsudil is a rho-kinase inhibitor used in the treatment and management of glaucoma and ocular hypertension. It lowers the intra-ocular pressure directly through the relaxation of the trabecular meshwork, leading to an increase in aqueous outflow. It also reduces aqueous humor production.

Netarsudil ophthalmic solution is instilled in one drop into the affected eye once daily in the evening. Adverse effects are conjunctival hyperemia, and corneal verticillata.

Other options:

Option A- Timolol is a non-selective β-blocker that decreases the IOP by reducing the aqueous humor production through the blockade of the β-receptors on the ciliary epithelium. Ocular adverse effects are allergy, decreased corneal sensation, and punctate keratitis.

Option C- Brimonidine is a selective alpha-2 agonist which decreases aqueous synthesis and increases uveoscleral outflow. It can cross the blood-brain barrier and, hence, causes drowsiness and CNS depression. It is contraindicated in infants and children.

Option D- Acetazolamide is a topical carbonic anhydrase inhibitor that decreases IOP by reducing aqueous formation. The main adverse effect is the bitter taste following administration. It is contraindicated in people with allergies to sulfonamide antibiotics.

Solution to Question 55:

Finasteride is used to treat male pattern baldness acts by inhibiting 5-alpha reductase.

Finasteride is a competitive inhibitor of the 5-alpha-reductase enzyme that inhibits the conversion of testosterone into its active form dihydrotestosterone (DHT).

Finasteride inhibits type 2 5-alpha reductase within the hair follicle and reduces the production of DHT. Hence, it is used in the treatment of male pattern hair loss. But, new hair growth will be lost when finasteride is discontinued, similar to minoxidil.

On long-term use (nearly 6 months), it produces an effect by reducing prostatic epithelial cell proliferation, thereby reducing the size of the prostate and improving urine flow.

Adverse effects of finasteride:

• Decreased libido

• Erectile dysfunction

• Ejaculation disorder and decreased ejaculate volume

Finasteride is highly teratogenic and pregnant women should not be exposed to the drug because of the potential for inducing genital abnormalities in male fetuses.

Solution to Question 56:

Rituximab is a monoclonal antibody that acts against CD20 B-cell antigen expressed on the surface of B-lymphocytes, plasma cells, and B-cell lymphomas.

It promotes apoptosis through transmembrane signalling as well as by antibody dependent and complement dependent cytotoxicity. It is used in the treatment of B-cell lymphoma, non-Hodgkin lymphoma, and chronic lymphocytic leukemia. It can be used both as a single agent as well as in combination with cytotoxic chemotherapy.

Adverse effects are infusion reactions like chills, fever, urticarial rashes, pruritus, dyspnea, and hypotension. Pretreatment with antihistaminics and corticosteroids prevents the reactions.

Solution to Question 57:

New drug application is filed after phase 3 of clinical trials.

Other options:

Option A: Phase 4 clinical trials are also called post-marketing surveillance.

Option C: Phase 2 trials are the first round of trials in patients.

Option D: Phase 1 trials are done in healthy volunteers.

Pharmacology NEET 2022

Question 1:

A male patient with chronic obstructive pulmonary disease (COPD) was prescribed theophylline. He noticed that his urine output had increased the following day. This action of the drug is mediated through which of the following receptors?

a) Interleukin-10

b) Histone deacetylase

c) Adenosine A1

d) Beta 2 adrenergic receptors

Question 2:

Which of the following drugs is not likely to cause a pleural pathology?

a) Metformin

b) Methysergide

c) Bleomycin

d) Nitrofurantoin

Question 3:

A patient with deep vein thrombosis was started on a new drug. After 2 days, he presented with the given finding. Which of the following drugs is implicated in causing the above condition?

a) Warfarin

b) Heparin

c) Dabigatran

d) Rivaroxaban

Question 4:

A patient undergoing chemotherapy was given an antiemetic, after which he developed symptoms like acute dystonia, bradykinesia, and tremors. Which of the following drugs would have caused these symptoms?

a) Ondansetron

b) Metoclopramide

c) Meclizine

d) Scopolamine

Question 5:

A young patient started to take a weight loss medication that acts by inhibiting fat absorption from food. After a few weeks, she developed easy bruising and increased menstrual bleeding. Deficiency of which of the following vitamins is responsible for her condition?

a) Vitamin E

b) Vitamin K

c) Vitamin B6

d) Vitamin D

Question 6:

A woman presenting with symptoms of urinary tract infection was prescribed a drug that causes tendon rupture and arthropathy. What is the mechanism of action of the drug?

a) DNA gyrase inhibition

b) Ribosomal inhibition

c) Cell wall synthesis inhibition

d) Inhibition of folic acid synthesis

Question 7:

A patient was brought to the emergency with a history of consumption of 8 tablets of digoxin. On examination, he was unstable and his heart rate was 56 bpm. ECG showed a 3rd-degree heart block. What is the next step in the management of this patient?

a) Antibody against digoxin

b) Lidocaine

c) DC cardioversion

d) Phenytoin

Question 8:

An elderly man, who is a known case of diabetic nephropathy, presented to the emergency with palpitations and chest discomfort. ECG showed tall T-waves. Laboratory investigations showed elevated potassium levels. Which of the following drugs will cause the shift of potassium back into the cell?

a) Epinephrine

b) Glucagon

c) Atropine

d) Lactic acid

Question 9:

A patient with a pituitary tumor that overproduced growth hormone underwent surgical removal of the tumor. The resection was found to be incomplete. What is the first-line treatment for this patient?

a) Leuprolide

b) Goserelin

c) Nafarelin

d) Octreotide

Question 10:

A middle-aged male patient presents with protrusion of the chin, excessive sweating, impaired glucose tolerance, and enlargement of hands and feet. Which of the following is a growth hormone receptor antagonist used to treat this condition?

a) Octreotide

b) Pegvisomant

c) Cabergoline

d) Olcegepant

Question 11:

A diabetic patient presented with rhinitis and facial swelling. An intranasal biopsy revealed the presence of broad-based aseptate hyphae with branching at right angle. Evidence of vascular invasion was present. What will be the drug of choice for this condition?

a) Fluconazole

b) Amphotericin B

c) Ketoconazole

d) Griseofulvin

Question 12:

A tuberculosis patient on anti-tubercular treatment presents with a tingling sensation and paresthesia in the lower limbs. He is not a diabetic and occasionally consumes alcohol. Which of the following vitamins must be supplemented to this patient?

a) Vitamin B6

b) Vitamin B12

c) Vitamin B2

d) Vitamin B3

Question 13:

An elderly woman received a botox injection for the treatment of wrinkles. Which of the following is the mechanism of action of botulinum toxin?

a) Inhibits the release of acetylcholine

b) Release of noradrenaline at synaptic cleft

c) Selectively and irreversibly inhibits nicotinic receptors

d) Stimulates muscarinic and nicotinic receptors

Question 14:

Which of the following is a PCSK9 inhibitor?

a) Evolocumab

b) Ezetimibe

c) Bempedoic acid

d) Clofibrate

Question 15:

An elderly man presents with rigidity and tremors. On examination, he has blank facial expressions. Which of the following drugs can be used to manage this condition?

a) Clozapine

b) Donepezil

c) Selegiline

d) Haloperidol

Answer Key

Question No.	Correct Option

1	c
2	a
3	a
4	b
5	b
6	a
7	a
8	a
9	d
10	b
11	b
12	a
13	a
14	a
15	c

Detailed Explanations

Solution to Question 1:

An increased urine output after administering theophylline is due to the antagonism at adenosine A1 receptors.

The adenosine A1 receptors are present in the afferent arteriole of the glomeruli. When activated, they cause vasoconstriction that reduces the glomerular filtration rate (GFR) and renal blood flow. This reduces urine output. Antagonistic action on these receptors will cause an increased GFR and renal blood flow and hence increased urine output.

Methylxanthines like theophylline and aminophylline are the bronchodilators used mainly in chronic obstructive pulmonary disease (COPD) and bronchial asthma. The pharmacological actions of theophylline are:

• It inhibits phosphodiesterase (PDE) non-selectively, thereby increasing cellular cAMP and cGMP, causing bronchodilation.

• Adenosine causes bronchoconstriction by releasing histamine and leukotrienes. Theophylline antagonizes adenosine A1 receptors and causes bronchodilation.

• It increases IL-10 release, which has anti-inflammatory actions.

• It causes a decrease in gene transcription of pro-inflammatory cytokines.

• It promotes apoptosis of granulocytes and thereby, reduces the inflammatory load in the airways.

Adverse effects of theophylline:

• Headache, nausea, and vomiting (due to inhibition of PDE4)

• Increased acid secretion (due to PDE inhibition)

• Diuresis (due to inhibition of adenosine A1 receptors)

• Cardiac arrhythmias (due to inhibition of cardiac PDE3 and antagonism of cardiac A1 receptors).

• Seizures (due to central A1 receptor antagonism).

Theophylline is mainly metabolized mainly by the enzyme CYP1A2. Therefore, it has various drug interactions, which include:

• Agents that lower its plasma level are smoking, phenytoin, rifampicin, and phenobarbitone.

• Drugs that increase its plasma level are erythromycin, ciprofloxacin, oral contraceptives, and allopurinol.

Solution to Question 2:

Metformin is not associated with pleural pathology.

Metformin belongs to the biguanides class of oral hypoglycemic agents. It acts by activation of AMP-dependent protein kinase (AMPK). This results in reduced hepatic glucose production.

Side effects of metformin are lactic acidosis, gastrointestinal disturbance, and vitamin B12 deficiency.

Contraindications for metformin include :

• GFR <30 ml/min

• Unstable CHF

• Radiographic contrast studies

• Hospitalized patients who cannot take anything orally

• Acidosis

Other options:

Option B: Methysergide is a serotonin(5-HT) receptor antagonist. It was used for the treatment of migraine prophylaxis and carcinoid syndrome. But its use has been discontinued due to side effects like abdominal, pulmonary and endocardial fibrosis.

Option C: Bleomycin is an anticancer antibiotic used for the treatment of squamous cell cancer and Hodgkin lymphoma. It chelates copper or iron and produces superoxide ions that damage DNA. Side effects are mucocutaneous toxicity and pulmonary fibrosis.

Option D: Nitrofurantoin is a urinary antiseptic. They are used for the treatment of lower urinary tract infections (UTIs). It acts by damaging DNA and inhibiting protein synthesis in susceptible bacteria. Important side effects are gastrointestinal disturbances, liver damage and pulmonary fibrosis.

Solution to Question 3:

The clinical scenario of the development of purple toe syndrome, after starting a new drug for the treatment of deep vein thrombosis, is suggestive of the adverse effect of warfarin.

Purple toe syndrome is a rare complication of warfarin therapy. It occurs due to cholesterol microemboli released from atheromatous plaques, which cause arteriolar occlusion. Warfarin must be immediately stopped and replaced with an oral factor Xa inhibitor (e.g. apixaban).

Activation of coagulation factors II, VII, IX, and X and proteins C and S are done by carboxylation of their glutamate residues, with the help of the reduced form of vitamin K as a co-factor. This reduced form of vitamin K needs to be constantly recycled back from its oxidized form for the reaction to continue. This recycling is done by the enzyme vitamin K epoxide reductase, which is competitively inhibited by warfarin. Hence, there will be a reduction in the activation of coagulation factors.

The image below shows warfarin's mechanism of action.



The half-life of proteins C and S are 8 and 24 hours, respectively, which is relatively shorter than the half-lives of other clotting factors, particularly factor II. Therefore, the full antithrombotic effect of warfarin is not achieved for 4 to 5 days. For this reason, warfarin must be bridged with a rapidly acting parenteral anticoagulant, such as heparin, low-molecular-weight heparin (LMWH), or fondaparinux, in patients with thrombosis or at high risk for thrombosis.

Uses of warfarin are acute deep vein thrombosis, pulmonary embolism, and preventing stroke or systemic embolization in patients with a mechanical heart valve.

Adverse effects of warfarin are bleeding, skin necrosis, purple toe syndrome, and fetal warfarin syndrome (hypoplasia of the nose, eye socket, hand bones, and growth retardation).

The therapeutic dose of warfarin is monitored by an INR range of 2-3. Warfarin toxicity can occur when the INR exceeds 4 and can lead to fatal bleeding. Treatment includes withholding warfarin, with vitamin K, and blood transfusions or fresh frozen plasma to replenish clotting factors.

Warfarin, LMWH, and UFH do not accumulate in breast milk and do not have an anticoagulant effect in the newborn. They are compatible with breastfeeding.

Warfarin is mainly metabolized by the CYP2C9 enzyme. Drugs that affect the CYP2C9 enzyme can alter the anticoagulant function of warfarin and these include the following:

Reduced efficacy:

• CYP2C9 activation by drugs: barbiturates, carbamazepine, rifampin

• Oral contraceptives: increase blood levels of clotting factors

Increased efficacy:

• Drugs causing CYP2C9 inhibition:

• Amiodarone

• Azole antifungals

• Cimetidine

• Clopidogrel

• Cotrimoxazole

• Disulfiram

• Isoniazid

• Metronidazole

• Fluoxetine

• Sulfinpyrazone

• Zafirlukast

Other options:

Option B: Heparin acts by activating antithrombin III, thereby inactivating the active form of clotting factors II, IX, X, XI, XII, and XIII. Adverse effects are bleeding and heparin-induced thrombocytopenia (HIT) due to the formation of antibodies against the heparin-platelet complex.

Option C: Dabigatran is a direct thrombin inhibitor. It is used for the prevention of embolism and stroke in patients with atrial fibrillation. Bleeding and dyspepsia are the most important adverse effects.

Option D: Rivaroxaban is a direct inhibitor of factor Xa. It is used for prophylaxis and treatment of deep vein thrombosis. Side effects reported are bleeding, nausea, hypotension, tachycardia, and edema.

Solution to Question 4:

In the above scenario, the patient has developed extrapyramidal adverse effects like acute dystonia, bradykinesia, and tremors. This is likely due to the antagonist action on dopamine receptors by metoclopramide.

Metoclopramide is a prokinetic and antiemetic. It has the following pharmacological actions:

• Dopamine receptor antagonist

• Serotonin (5HT4) receptor agonist

• Serotonin (5HT3) antagonist

• Sensitization of muscarinic receptors on smooth muscle. This results in increased lower esophageal sphincter tone and stimulates gastric antral contractions.

Metoclopramide acts on both the gastrointestinal system and central nervous system. It is used as an upper antiemetic for many types of vomiting, especially migraine, drug-induced, and chemotherapy-induced. It is also used as a prokinetic in gastroparesis to increase gastric emptying.

Adverse effects of metoclopramide are mainly extrapyramidal effects, and these include acute dystonia, Parkinson-like symptoms (tremors, bradykinesia, rigidity), tardive dyskinesia, galactorrhea due to hyperprolactinemia, and methemoglobinemia in neonates.

Extrapyramidal symptoms can be treated by discontinuation of metoclopramide and administration of anticholinergics or antihistaminics.

Other options:

Option A: Ondansetron is a 5-HT3 receptor antagonist. It blocks impulses at the peripheral origin in GIT. It is most commonly indicated for the treatment of postoperative and chemotherapy-induced nausea and vomiting. Headache and dizziness are the common adverse effects.

Option C: Meclizine is a long-acting H1 antihistaminic drug. It is used for seasickness. It is less sedative and less anticholinergic.

Option D: Scopolamine (hyoscine) is a muscarinic antagonist. It blocks neural pathways from the vestibular apparatus in the inner ear to the emetic center in the brainstem. A transdermal patch of scopolamine is used as an antiemetic in motion sickness. Constipation, drowsiness, dry mouth, and blurred vision are the common adverse effects.

Solution to Question 5:

The clinical scenario of easy bruising and increased menstrual bleeding following weight loss medication that inhibits fat absorption from food is suggestive of vitamin K deficiency.

Vitamin K is a fat-soluble vitamin. Hence, drugs that inhibit the absorption of fat (eg; ezetimibe) also inhibit the absorption of fat-soluble vitamin K. The following are the vitamers of vitamin K:

• Phylloquinone (Vitamin K1) is the normal dietary source of Vitamin K that is found in green vegetables.

• Menaquinone (Vitamin K2) is synthesized by intestinal bacteria.

• Menadione (Vitamin K3) is a synthetic compound that is metabolized to phylloquinone.

Vitamin K is required for γ-carboxylation of glutamate residues of clotting factors like prothrombin (factor II), factors VII, IX, and X. Warfarin inhibits γ-carboxylation by preventing the conversion of vitamin K to its active hydroquinone form.

Vitamin K deficiency in adults may be seen in patients with liver disease, obstructive jaundice, malabsorption, and prolonged antimicrobial therapy. Vitamin K deficiency is usually diagnosed

on the basis of an elevated prothrombin time or reduced clotting factors.

Vitamin K deficiency is treated with a parenteral dose of 10 mg followed by 5mg (duration depends on the severity of vitamin K deficiency).

Other options:

Option A: Vitamin E is a fat-soluble vitamin. The levels of vitamin E also will get decreased following taking weight loss medication that inhibits fat absorption from food. But the symptoms of vitamin E deficiency include hemolytic anemia and neuromuscular manifestations.

Option C: Vitamin B6 is a water-soluble vitamin. Drugs that cause vitamin B6 deficiency are isoniazid, hydralazine, cycloserine, penicillamine, and oral contraceptives. Patients usually present with peripheral neuritis, anemia, seborrheic dermatitis, glossitis, lowered seizure threshold, or convulsions (due to fall in brain GABA levels).

Option D: Vitamin D is a fat-soluble vitamin. Vitamin D levels will also decrease following weight loss medication that inhibits fat absorption from food. Patients with vitamin D deficiency will typically develop rickets in children and osteomalacia in adults.

Solution to Question 6:

In the above scenario, the antimicrobial prescribed for urinary tract infection, which has a propensity to cause tendon rupture and arthropathy, is likely to be a fluoroquinolone. The mechanism of action of the drug is DNA gyrase inhibition.

Normally, during replication, to prevent excessive positive supercoiling of the strands, DNA gyrase nicks double-stranded DNA and introduces negative supercoils. Fluoroquinolones, by inhibiting DNA gyrase (in gram-negative bacteria), result in the production of damaged DNA. This is finally digested by exonucleases, resulting in bactericidal action.

Fluoroquinolones inhibit topoisomerase IV in gram-positive bacteria.

Fluoroquinolones are used in the treatment of urinary tract infections, prostatitis, and gastrointestinal, respiratory infections, and as a second-line drug in anti-tubercular therapy. They are avoided in children and pregnant women.

Adverse effects of fluoroquinolones are arthralgia, tendon rupture, Clostridium difficile colitis, dysglycemia, prolonged QT interval, rashes, hallucinations, delirium, and seizures.

Other options:

Option B: Ribosomal inhibition is the mechanism of action of various antimicrobials like tetracyclines, aminoglycosides, chloramphenicol, and macrolides.

Option C: Cell wall synthesis inhibition is done by beta-lactam antibiotics.

Option D: Inhibition of folic acid synthesis is the mechanism of action of sulfonamides and co-trimoxazole.

Solution to Question 7:

Administration of digibind (antibody against digoxin) is the next step in managing this patient. An unstable patient with bradycardia and 3rd-degree heart block following digoxin toxicity should be treated with digibind.

Digibind is a nonimmunogenic antibody used to treat cardiotoxicity due to digoxin. It is nonimmunogenic because it lacks the Fc fragment.

Digoxin is the cardiac glycoside used to treat heart failure and control ventricular rate in atrial fibrillation/flutter.

It inhibits Na+/K+ ATPase, a membrane-bound transporter to increase intracellular sodium. As a result, there is less expulsion of calcium from the cell by the sodium-calcium exchanger (NCX). Increased calcium inside the myocyte leads to an increase in cardiac contractility.

The most common cardiac manifestations of digitalis toxicity are ventricular tachycardia, partial to complete A-V block, severe bradycardia, and atrial fibrillation/flutter.

Factors that increase the risk of digoxin toxicity are:

• Hypokalemia

• Drugs that decrease digoxin elimination (inhibitors of P-glycoprotein) like quinidine

• Renal/hepatic failure

• Thyrotoxicosis

Option B: Lidocaine is commonly used as a local anesthetic. It has anti-arrhythmic properties and is used to suppress ventricular tachycardia caused by digitalis toxicity.

Option C: DC cardioversion is used for tachyarrythmias when pharmacologic treatment fails. It is not used in bradyarrhythmias.

Option D: Phenytoin is an antiepileptic used for the treatment of generalized tonic-clonic seizures and partial seizures. It can also be used for the treatment of polymorphic ventricular tachycardia due to QT prolongation.

Solution to Question 8:

The clinical scenario with ECG showing tall T-waves and elevated serum potassium levels are suggestive of hyperkalemia. Epinephrine will cause the shift of potassium back into the cell.

Epinephrine is an agonist at alpha 1, beta 1, and beta 2 receptors.

The β actions are mediated through cAMP. Epinephrine binds to Gs → ATP and is broken down to cAMP by the enzyme adenylyl cyclase. This results in:

• In the liver and muscle- hyperglycemia and hypokalemia due to potassium uptake in muscle and liver

• In adipose tissue- increased plasma free fatty acids

• In the heart- an increased force of contraction reinforces the positive inotropic action

• In pancreatic islets- increases glucagon secretion

Epinephrine is used in anaphylaxis, to prolong the action of local anesthetics, cardiac arrest, and as inhalational epinephrine in the treatment of post-intubation and infectious croup.

Epinephrine is contraindicated in anesthesia with halothane due to the risk of arrhythmias and in patients receiving beta-blockers because a marked rise in BP can occur due to unopposed alpha action.

Hyperkalemia refers to an increase in the levels of serum potassium >5.5mEq/L.

The image given below shows sequential ECG changes seen in hyperkalemia.



Hyperkalemia is a medical emergency and has to be treated when ECG changes are present or severe hyperkalemia (serum K+ >6.5) without ECG changes. Management is as follows:

• Intravenous infusion of 10 ml of 10% calcium gluconate for 2-3 minutes with cardiac monitoring. It helps to antagonize the cardiac effects.

• Intravenous regular insulin 10 units followed by 50mL of 50 % dextrose. This helps in shifting the plasma K+ into the cells.

• Nebulisation with β2 agonists such as albuterol or salbutamol.

• Cation exchange resins, diuretics, and/or dialysis are useful to remove the excess potassium.

Solution to Question 9:

The above scenario with a pituitary tumor producing growth hormone is likely to cause acromegaly in the patient. Octreotide is the first-line treatment for this condition.

Octreotide is a synthetic analog of somatostatin.

Octreotide can be used in the following conditions:

• Acromegaly

• Secretory diarrheas associated with carcinoid, AIDS, cancer chemotherapy, or diabetes- by suppression of hormones that enhance intestinal mucosal secretion.

• Bleeding esophageal varices- reduces hepatic blood flow and helps stop bleeding.

• Dumping syndrome in post gastric surgery patients

Adverse effects of octreotide are:

• Abdominal pain

• Nausea

• Steatorrhea

• Gall stones (due to biliary stasis)

• Hyperglycemia

Other options:

Option A: Leuprolide is a long-acting GnRH agonist. It is used for palliation of carcinoma prostate, uterine fibroids, endometriosis, and precocious puberty. Side effects include hot flashes, decreased bone mineral density, vaginal dryness, and erectile dysfunction.

Option B: Goserelin is also a GnRH agonist used for the treatment of endometriosis, dysfunctional uterine bleeding, and for prostate and breast cancer.

Option C: Nafarelin is a long-acting GnRH agonist. It is used for the treatment of endometriosis, uterine fibroids, and central precocious puberty.

Solution to Question 10:

The clinical features of protrusion of the chin, excessive sweating, impaired glucose tolerance, and enlargement of hands and feet are suggestive of acromegaly. Pegvisomant is a growth hormone receptor antagonist used to treat this condition.

Acromegaly is commonly caused due to growth hormone-secreting somatotroph adenoma. It occurs secondary to mutations of the pituitary-specific gene PIT-1.

Patients present with frontal bossing, increased hand and foot size, mandibular enlargement with prognathism, widened space between the lower incisor teeth, soft tissue swelling resulting in increased heel pad thickness, coarse facial features, and large fleshy nose.

Serum IGF-1 levels (elevated) and GH suppression test with oral glucose are useful in the diagnosis. The levels of GH do not reduce in response to the oral glucose load.

Transsphenoidal surgical resection of the tumor, octreotide, and pegvisomant will be helpful in the management of this condition.

Other options:

Option A: Octreotide is a synthetic analog of somatostatin. It is used for the treatment of acromegaly and secretory diarrhea.

Option C: Cabergoline is a dopamine D2 receptor agonist. It is used as a first-line drug for the treatment of hyperprolactinemia.

Option D: Oglcegepant is a calcitonin gene-related peptide (CGRP) antagonist. It is used for the treatment of migraine.

Solution to Question 11:

The above scenario and biopsy findings showing broad-based aseptate hyphae with branching at right angles are suggestive of mucormycosis. The drug of choice for mucormycosis is amphotericin B.

Mucormycosis is an opportunistic mycosis. It is typically seen in patients with diabetes mellitus and/or diabetic ketoacidosis and invades blood vessels.

The rhino-orbital-cerebral is the most common form of mucormycosis. Patients present with rhinitis, eye or facial pain, and facial numbness followed by the onset of conjunctival suffusion and swelling along with blurry vision.

Diagnosis is made by biopsy with histopathologic examination. The biopsy shows characteristic wide, ribbon-like aseptate hyphal elements that branch at right angles.

The below image shows biopsy findings seen in mucormycosis.



Treatment of the infection includes 4 steps:

• Initiation of therapy (amphotericin B) as early as possible.

• Surgical debridement.

• Reversal of underlying conditions and risk factors.

• Treatment of underlying malignancy if present.

Amphotericin B binds to ergosterol in the fungal membrane and forms pores or channels in the fungal cell membrane. This allows the leakage of cytosolic molecules and ions, leading to a loss of membrane integrity.

It is a broad-spectrum antifungal that can be used against Candida albicans, Cryptococcus neoformans, Blastomyces dermatitidis, Histoplasma capsulatum, Sporothrix schenckii, Coccidioides immitis, Aspergillus.

Amphotericin B also has antiprotozoal action along with antifungal properties and is effective against Leishmania.

Adverse effects of amphotericin B:

• Infusion-related fever and chills

• Azotemia/nephrotoxicity- all lipid formulations of amphotericin B are less nephrotoxic than conventional amphotericin B

• Renal tubular acidosis

• Renal wasting of potassium and magnesium

• Hypochromic normocytic anemia- due to decreased erythropoietin production from the kidney

Note: Mucormycosis recently emerged as an important superinfection in COVID-19 patients, especially in India. The surge in cases is most likely due to diabetes mellitus and steroid use.

Other options:

Option A: Fluconazole is used for the treatment of candidiasis and cryptococcosis.

Option C: Ketoconazole is used for both dermatophytosis and deep mycosis, but it has been replaced by itraconazole.

Option D: Griseofulvin is used for dermatophytosis. It is effective against the mycotic disease of the skin, hair, and nails due to Microsporum, Trichophyton, and Epidermophyton.

Solution to Question 12:

Vitamin B6 must be supplemented for this patient. The clinical features of tingling sensation and paresthesias in a patient on anti-tubercular treatment are suggestive of the adverse effect of isoniazid.

Isoniazid is the first-line antitubercular drug that inhibits the production of mycolic acid. It is a bactericidal drug against tubercle bacilli.

Adverse effects of isoniazid:

• Peripheral neuritis (paresthesias, numbness, tingling sensation) due to interference with the production of active coenzyme pyridoxal phosphate from pyridoxine. Therefore, pyridoxine 10 mg/day (vitamin B6) is given prophylactically along with isoniazid to prevent neurotoxicity.

• Hepatitis - isoniazid must be stopped at the first sign of hepatotoxicity.

• Rashes

• Anemia

• Arthralgia

Toxicity of isoniazid results in metabolic acidosis and seizures.

Note: Vitamin deficiency in alcoholics includes the following:

• Folate causes megaloblastic anemia. It is treated by supplementation of oral doses of 5-15 mg of folic acid.

• Vitamin B6 results in peripheral neuritis, anemia, and glossitis, and lowers the seizure threshold. It is treated by supplementation of oral doses of 2-5 mg of pyridoxine.

• Thiamine results in Wernicke's encephalopathy. It is treated with 200 mg of intravenous thiamine three times daily, followed by 10 mg/day until complete recovery.

• Vitamin A results in xerophthalmia and night blindness. It is treated with supplementation of oral vitamin A 3000-5000 IU/day.

Solution to Question 13:

The botulinum toxin acts by inhibiting the release of acetylcholine into the synaptic cleft.

Botulinum toxin cleaves SNARE protein, which is involved in the exocytosis of acetylcholine. Thus, it inhibits the release of acetylcholine.

The image given below shows the site of action of botulinum toxin:



This toxin causes a long-lasting loss of cholinergic transmission. Hence, it can be employed for the treatment of several spastic and neurological conditions caused by the overactivity of cholinergic nerves, such as age-related facial wrinkles, blepharospasm, spastic cerebral palsy, spasmodic torticollis, post-stroke spasticity, and axillary hyperhidrosis.

Adverse effects occur due to incorrect injection and overdose; they include ptosis, diplopia, facial swelling, dry mouth, muscle weakness, and respiratory paralysis.

Botulinum toxin is an exotoxin produced by the bacteria Clostridium botulinum. It is responsible for a type of food poisoning called botulism.

Solution to Question 14:

Evolocumab is a PCSK9 inhibitor.

PCSK9 refers to proprotein convertase subtilisin/kexin type 9. They are mainly involved in the downregulation of low-density lipoprotein (LDL) receptors in the liver.

Evolocumab is a monoclonal antibody that inhibits PCSK-9 (proprotein convertase subtilisin/kexin). Inhibition of PCSK-9 prevents the degradation of LDL receptors and thus increases LDL-cholesterol clearance by the liver. They are used in patients with atherosclerotic cardiovascular disease (ASCVD) to decrease LDL levels.

Side effects of PCSK9 inhibitors are injection-site reactions, nasopharyngitis, urinary tract infection, and upper respiratory tract infection and it does not increase the risk of myopathy.

Other options:

Option B: Ezetimibe inhibits cholesterol absorption by enterocytes in the small intestine. It lowers the total and LDL-C levels.

Option C: Bempedoic acid inhibits adenosine triphosphate-citrate lyase (ACL) and results in decreased LDL in patients not responding to statin therapy.

Option D: Clofibrate binds to PPAR-alpha and causes stimulation of fatty acid oxidation and increased LPL synthesis. This results in reduced triglycerides, LDL and increased high-density lipoprotein (HDL) levels. It causes myopathy when combined with statins and is associated with an increased risk of gallstone formation.

Solution to Question 15:

The clinical features of rigidity, tremors, and blank facial expressions are suggestive of Parkinson's disease. Monoamine oxidase-B (MAO-B) inhibitors like selegiline are used for the treatment of Parkinson disease.

Normally, MAO-B oxidizes dopamine. Selegiline is a selective and irreversible MAO-B inhibitor. Therefore, it increases dopamine levels in the brain. It is used as an adjuvant to levodopa in Parkinson's disease. It prolongs and enhances the action of levodopa.

Adverse effects of selegiline are postural hypotension, nausea, confusion, involuntary movements, and psychosis. Selegiline is contraindicated in patients with convulsive disorders.

Other options:

Option A: Clozapine is an atypical antipsychotic that selectively blocks dopamine (D4) receptors in the brain. It is used in the treatment of refractory schizophrenia. Agranulocytosis and metabolic syndrome are the important adverse effects.

Option B: Donepezil is a reversible cholinesterase inhibitor used for the treatment of Alzheimer's disease.

Option D: Haloperidol is a typical antipsychotic used for the treatment of acute schizophrenia, Huntington’s disease, and Gilles de la Tourette syndrome.

Pharmacology NEET 2023

Question 1:

A bronchial asthma patient on inhalational steroids presented with white patchy lesions on the tongue and buccal mucosa. What is the drug that can be used to treat this condition?

a) Clotrimazole

b) Griseofulvin

c) Terbinafine

d) Flucytosine

Question 2:

A pregnant woman with a history of bronchial asthma is in the third stage of labor. Which of the following drugs should be avoided in the management of postpartum hemorrhage in this mother?

a) Carboprost

b) Oxytocin

c) Dinoprostone

d) Methyl ergometrine

Question 3:

Name the drug that acts on both the marked areas.



a) Sacubitril

b) Omapatrilat

c) Losartan

d) Nesiritide

Question 4:

A patient with a previous history of myocardial infarction and ventricular arrhythmia is on treatment for a few months. He developed fatigue, dyspnea, and weight gain. He had also developed pulmonary fibrosis. Which of the following drugs is responsible for the above-mentioned side effects?

a) Amiodarone

b) Atenolol

c) Aspirin

d) Spironolactone

Question 5:

A young female comes to the emergency department after consuming 100 tablets of aspirin. What should be the next step in management?

a) N-acetyl cysteine to replenish glutathione stores

b) Pralidoxime

c) Glucagon to control bradycardia and hypoglycemia

d) Give sodium bicarbonate to alkalize urine

Question 6:

A chronic smoker was on nicotine replacement therapy and clonidine tablets for smoking de-addiction. He stopped taking clonidine tablets and now presents with a headache. What is the reason behind this condition?

a) Postural hypotension

b) Receptor upregulation

c) Rebound hypertension

d) Receptor hypersensitivity

Question 7:

A patient with recurrent attacks of gout was started on a therapy that inhibits the synthesis of uric acid. His symptoms were reduced after therapy. Which drug was he started on?

a) Probenecid

b) Colchicine

c) Diclofenac

d) Allopurinol

Question 8:

A patient with a malignancy is undergoing chemotherapy. The platelet counts were reduced after the previous cycle of chemotherapy. Which of the following drugs can be used in the treatment of this patient?

a) Oprelvekin (IL-11)

b) Filgrastim

c) Erythropoietin

d) Amifostine

Question 9:

The true statement among the following is _____

a) The dose of telmisartan should be reduced in renal failure but not in hepatic failure.

b) The dose of irbesartan should be reduced in case of mild-moderate hepatic failure and renal failure.

c) The dose of candesartan should be reduced in mild-moderate liver failure but not in renal failure.

d) Losartan acts as thromboxane A2 antagonist and inhibits platelet aggregation.

Question 10:

A patient given digoxin started having side effects like nausea and vomiting. The serum concentration of digoxin was 4 mg/dL. The plasma therapeutic range is 1 mg/dL. If the half-life of digoxin is 40 hours, how long one should wait before resuming the treatment?

a) 40 hours

b) 80 hours

c) 120 hours

d) 140-180 hours

Question 11:

A patient comes to the casualty with organophosphate poisoning. He was started on atropine infusion and pralidoxime. After 2 hours, the patient had a sudden rise in temperature. What is the likely cause of fever?

a) Atropine toxicity

b) Side effect of pralidoxime

c) Due to organophosphate poisoning

d) Idiopathic

Question 12:

The mechanism of action of tissue plasminogen activator is _____

a) Inhibit extrinsic pathway

b) Inhibits platelet aggregation

c) Enhance fibrin degradation

d) Inhibit clot formation

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	a
5	d
6	c
7	d
8	a
9	d
10	b
11	a
12	c

Detailed Explanations

Solution to Question 1:

The given clinical scenario of white patchy lesions on the tongue and buccal mucosa in a patient using inhalational steroids is oral candidiasis. The drug that can be used for the treatment is clotrimazole.

MDI results in the deposition of steroids in the oropharyngeal space, which can cause cough, dysphonia, and oropharyngeal candidiasis.

Clotrimazole is a topical imidazole used in the treatment of tinea infections like ringworm. It is particularly used for the treatment of vaginitis as a once-daily application because of its long-lasting residual effect.

For oral candidiasis, 10 mg troche of clotrimazole is allowed to dissolve in the mouth 3-4 times a day. Alternatively, lotion or gel is applied and swirled in the mouth for as long as possible. Side effects are local irritation and a burning sensation.

Other options:

Option B: Griseofulvin acts by inhibiting microtubule assembly, disrupting the fungal cell division. It is mainly used against dermatophytes and is the drug of choice for tinea capitis in children. It is not effective against Candida.

Option C: Terbinafine acts by inhibiting epoxidation of fungal squalene that reduces ergosterol biosynthesis. It is helpful for treating nail onychomycosis and tinea capitis.

Option D: Flucytosine acts by inhibiting fungal DNA synthesis. It is used exclusively in combination with amphotericin B for cryptococcal meningitis.

Solution to Question 2:

The drug that should be avoided in the management of postpartum hemorrhage in this mother with a history of bronchial asthma is carboprost.

Carboprost is an analog of PGF2α. It is used in obstetrics to induce second-trimester abortions and to control postpartum hemorrhage. Vomiting, diarrhea, and bronchoconstriction can occur.

Contraindications of carboprost:

Absolute	Relative
AsthmaPulmonary hypertensionSuspected amniotic fluid embolism	Pre-eclampsiaRenal diseaseLiver diseaseHeart disease

Other options :

Option B: Oxytocin is the first-line drug to prevent and treat PPH according to WHO guidelines. It is administered by intravenous route and may lead to hypotension and reflex tachycardia.

Option C: Dinoprostone is a prostaglandin E2 analog used for cervical ripening, induction of labor, and in atony treatment. It should be avoided in patients with a history of asthma, glaucoma, or myocardial infarction. One major adverse effect is uterine hyperstimulation.

Option D: Ergot derivatives are used in the control of postpartum uterine bleeding. It is given intramuscularly and is effective within 1–5 minutes. It is given at the time of delivery of the placenta or immediately after if bleeding is significant. Side effects are gastrointestinal disturbances like diarrhea, nausea, and vomiting.

Solution to Question 3:

The drug that acts on both the marked areas is omapatrilat.

Omapatrilat is a dual inhibitor of neutral endopeptidase and Angiotensin-converting enzyme (ACE). It decreases angiotensin II and increases atrial and brain natriuretic peptide, promoting vasodilation, diuresis, and anti-hypertrophic effect.

Other options:

Option A - Sacubitril is a neprilysin inhibitor. It reduces the breakdown of natriuretic peptides such as ANP, BNP, and CNP, thereby increasing their levels. Sacubitril combined with valsartan reduces mortality in heart failure with reduced ejection fraction (HFrEF).

Option C - Losartan is the angiotensin receptor blocker used in the treatment of hypertension. Hyperkalemia is an important side effect of ARBs. It appears to be a safe and efficacious agent to lower serum uric acid levels in patients with hyperuricemia.

Option D - Nesiritide is a recombinant human brain natriuretic peptide (BNP). It causes vasodilatation and natriuresis. It is administered intravenously and used in acute decompensated heart failure.

Solution to Question 4:

The drug that is responsible for the above-mentioned side effects is amiodarone.

Amiodarone is a class III antiarrhythmic drug. The most serious adverse effect during chronic amiodarone therapy is pulmonary fibrosis, which is rapidly progressive and fatal. The side effects of amiodarone include:

• Pulmonary fibrosis

• Hypotension, depressed myocardial function

• Corneal microdeposits

• Hepatotoxicity

• Neuropathies

• Photosensitivity

• Hypo- or hyperthyroidism

Other options:

Option B: Atenolol is a beta-1 selective antagonist. Bradycardia, fatigue, and cold extremities are common adverse effects. It is used in combination with a diuretic, in elderly patients with isolated systolic hypertension. CNS side effects like depression and nightmares are seen.

Option C: Aspirin is a nonsteroidal anti-inflammatory drug (NSAID) that blocks the formation of thromboxane A2 by acetylation of COX1. Through this mechanism, it shows antipyretic, analgesic, and anti-inflammatory actions. Higher doses can result in bleeding.

Option D: Spironolactone is a potasium-sparing diuretic. It is a competitive inhibitor of the mineralocorticoid receptor in the late distal tubule and collecting duct of the kidneys. It promotes sodium diuresis but maintains the body's potassium levels. It is contraindicated in patients with hyperkalemia. Side effects are gynecomastia, impotence, decreased libido, and menstrual irregularities.

Solution to Question 5:

The next step in the management of this patient who consumed 100 tablets of aspirin is to give sodium bicarbonate to alkalize the urine.

Alkalinization of urine will be helpful in the removal of aspirin, which is a weakly acidic drug and is susceptible to ion trapping in the urine. It is done by administration of sodium bicarbonate in 1

L of 5% dextrose in water (D5W). It is infused intravenously at twice the maintenance fluid requirements.

Alkalinization of urine is contraindicated in renal failure or if fluid administration may worsen pulmonary edema or congestive heart failure.

Other options:

Option A: N-acetyl cysteine is used as an antidote for the treatment of paracetamol poisoning which replenishes glutathione stores in the liver.

Option B: Pralidoxime is also used in organophosphate poisoning. It regenerates active acetylcholinesterase and relieves skeletal muscle end-plate block. It is usually used as an adjuvant to atropine.

Option D: Glucagon is used in beta-blocker toxicity.

Solution to Question 6:

The mechanism that causes headache in this patient, who stopped taking clonidine is rebound hypertension.

Clonidine is an alpha-2 receptor agonist and a centrally acting antihypertensive agent. It treats high blood pressure by stimulating presynaptic α2 receptors in the vasomotor center in the brainstem. This binding has a sympatholytic effect, by suppressing the release of norepinephrine and other sympathomimetics.

Abrupt withdrawal after long-term therapy can result in rebound hypertension, restlessness, and sympathetic overactivity. Dry mouth and sedation are the major side effects of clonidine. It can also cause impotency and constipation.

Phentolamine is used for the control of hypertension due to clonidine withdrawal.

Solution to Question 7:

Allopurinol which inhibits uric acid synthesis is the likely drug that was added to the regimen.

It is a purine antimetabolite and a competitive inhibitor of the xanthine oxidase enzyme. It prevents the conversion of hypoxanthine and xanthine to uric acid reducing plasma uric acid levels. The metabolite of allopurinol called oxypurinol is a non-competitive inhibitor of xanthine oxidase.

It is used in the treatment of hyperuricemia in patients with gout and also in patients with hematological malignancies about to undergo chemotherapy.

Other options:

Option A: Probenecid is a uricosuric agent that promotes the excretion of uric acids by inhibiting the organic anion transporter, particularly URAT-1 in the renal tubule. It is used to decrease the body pool of urate in patients with chronic gout. It is used as an adjuvant to allopurinol in chronic gout and hyperuricemia. Plenty of fluids are given to avoid urate

calculi formation.

Option B: Colchicine is an antimitotic drug that is used in the treatment of acute gout. It acts by inhibiting microtubule and spindle formation. Side effects are nausea, vomiting, myelosuppression, aplastic anemia, and rhabdomyolysis.

Option C: Diclofenac is a commonly used NSAID that has analgesic, antipyretic, and anti-inflammatory activities. It acts by inhibiting the cyclooxygenase-2 enzyme. Side effects are rashes, fluid retention, edema, and renal function impairment.

Solution to Question 8:

The most probable diagnosis for this patient is chemotherapy-induced thrombocytopenia. Oprelvekin can be used in the treatment of this patient.

Oprelvekin is an IL-11 analog used in the treatment of chemotherapy-induced thrombocytopenia. It increases thrombopoiesis. It is used in the treatment of thrombocytopenia due to non-myeloid malignancies. Important adverse effects are fluid retention, tachycardia, palpitation, edema, and shortness of breath.

Other options:

Option B: Filgrastim is a granulocyte-colony-stimulating factor (G-CSF) that is used for the treatment of severe neutropenia after high-dose cancer chemotherapy.

Option C: Epoetin alfa and darbepoetin alfa are recombinant human erythropoietin. They are used in the treatment of anemias associated with surgery, AIDS, cancer chemotherapy, and certain chronic inflammatory conditions.

Option D: Amifostine is used in the reduction of nephrotoxicity associated with cisplatin.



Solution to Question 9:

The true statement is losartan acts as a thromboxane A2 antagonist and inhibits platelet aggregation.

Losartan is a competitive angiotensin receptor antagonist. It binds to the AT1 receptor with high affinity. It is also a competitive antagonist of the thromboxane A2 receptor and attenuates platelet aggregation.

Losartan is the angiotensin receptor blocker used in the treatment of hypertension. Hyperkalemia is an important side effect of ARBs. It appears to be a safe and efficacious agent to lower serum uric acid levels in patients with hyperuricemia.

Other options:

Option A: The clearance of telmisartan is mainly by biliary secretion. The clearance is affected by hepatic insufficiency. The dose of telmisartan should be reduced in hepatic failure but not in renal failure.

Option B: Irbesartan is metabolized to glucuronide conjugate, and is cleared mainly by biliary excretion. The plasma clearance is unaffected by either renal or mild-to-moderate hepatic insufficiency. Hence, the dose need not be reduced in either of the cases.

Option C: The plasma clearance of candesartan is by biliary excretion and renal elimination. The clearance is affected by renal insufficiency. Hence, the dose should be reduced in case of renal failure and not in mild-moderate liver failure.

Solution to Question 10:

The treatment can be resumed after 80 hours.

The half-life (T1/2) is the time it takes for the plasma concentration of the drug to be reduced by 50%. In the given scenario, the half-life of digoxin is given as 40 hours. The serum concentration of digoxin is given as 4 mg/dL.

After 40 hours, the serum concentration of digoxin becomes 2 mg/dL (half of the initial concentration), and after 80 hours, it will be reduced to 1 mg/dL. So, to reach a serum concentration of 1 mg/dL (therapeutic range), it will take 80 hours.

Solution to Question 11:

The most likely cause of fever in this patient is atropine toxicity.

The clinical scenario is suggestive of organophosphate consumption and atropine is used for its management. Atropine is a muscarinic receptor antagonist, and it competitively blocks the acetylcholine at muscarinic receptors.

Organophosphates are powerful acetylcholinesterase inhibitors. It is absorbed by inhalation or through the skin. Mucous membranes and the gastrointestinal tract are also the sites for absorption.

The predominant initial signs are of muscarinic excess which includes: (DUMBBELSS)

• Diarrhoea

• Urination

• Miosis

• Bronchospasm

• Bradycardia

• Emesis

• Lacrimation

• Salivation

• Sweating

Nicotinic manifestations include twitching of eyelids, tongue, and facial muscles which then leads to paralysis. CNS stimulation results in headache, tremors, ataxia, coma, and death.

Atropine in larger doses can lead to atropine toxicity. The features are depicted in the following image:



Solution to Question 12:

The mechanism of action of the tissue plasminogen activator is to enhance fibrin degradation.

Tissue plasminogen activators activate plasminogen bound to fibrin, and catalyze the conversion of plasminogen to plasmin, resulting in increased formation of plasmin. This results in fibrinolysis of formed thrombus (thrombolysis).

It is indicated in pulmonary embolism with hemodynamic instability, acute ischemic stroke, and acute myocardial infarction. Recombinant variants of t-PA like reteplase and tenecteplase have a longer half-life.

All thrombolytics can cause bleeding, which may be counteracted by the administration of antifibrinolytics, such as aminocaproic acid and tranexamic acid.

Pharmacology INI-CET 2023

Question 1:

A patient underwent a hysterectomy and was given morphine in the post-operative period and in the ward for pain. She now presents with respiratory depression and pin point pupils. What is the drug used to treat this condition?

a) Naloxone

b) Pethidine

c) Atropine

d) Buprenorphine

Question 2:

Drug used for trastuzumab-resistant her 2 positive breast cancer is:

a) Vismodegib

b) Lapatinib

c) Vemurafenib

d) Erlotinib

Question 3:

Nomogram used for vancomycin dosing is?

a) Salazar

b) Halsted

c) Rummack-mathew

d) Matzke

Question 4:

Which of the following is true statement regarding lithium?

a) It is not absorbed from the stomach

b) It can cause fine tremors in the therapeutic range

c) Approved for the treatment of absent seizures

d) It is not teratogenic

Question 5:

Match the following

a) A-2, B-3, C-4, D-1

b) A-3, B-4, C-1, D-2

c) A-4, B-2, C-3, D-1

d) A-1, B-2, C-3, D-4

Question 6:

True about therapeutic index of a drug is:

a) Denotes safety and efficacy after experiments on humans

b) Denotes safety and efficacy in the population

c) Denotes safety and efficacy in an individual

d) Ratio of median lethal dose in 50% of population to median effective dose in 50% of population

Question 7:

Osimertinib is used in NSCLC with which of the following mutation?

a) T790M mutation

b) L858R mutation

c) T890M mutation

d) M790T mutation

Question 8:

A patient was started on a drug for mood disorder and he developed priapism. The causative drug is:

a) Trazodone

b) Venlafaxine

c) Bupropion

d) Mirtazapine

Question 9:

Cyclosporine-induced cholestasis is due to which receptor?

a) MDR 1

b) BRCA

c) MRP 1

d) OAT1B1

Question 10:

A diabetic patient was started on a drug that decreased her HbA1C from 7.6 to 6.7. She started complaining of itching around vulvovaginal area . What was the most likely drug she was started on?

a) Acarbose

b) Canagliflozin

c) Linagliptin

d) Liraglutide

Question 11:

An anti-diabetic drug causes a decrease in HbA1C by causing an increase in GIP and GLP-1 levels and a decrease in glucagon levels. It is an oral drug causing a reduction in both pre and post-prandial glucose. The drug is:

a) Pramlinitide

b) Alogliptin

c) Liraglutide

d) Metformin

Question 12:

What is true about keynote 189 trial?

a) Pembrolizumab plus chemotherapy used in NSCLC

b) Pembrolizumab used in NSCLC

c) Nivolumab plus chemotherapy used in NSCLC

d) Nivolumab is used in NSCLC

Question 13:

Which of the following is not a mechanism for thalidomide teratogenicity?

a) Release of reactive oxygen species and free radicles

b) Supression of angiogenesis

c) Cereblon binding

d) Inhibition of IL2, IL4, IL5

Question 14:

Which is a rare but very severe side effect of clozapine that requires regular monitoring?

a) Suicidal tendency

b) Agranulocytosis

c) Myocarditis

d) Cerebral bleeding

Question 15:

Which of the following newly approved anti-diabetic drugs can be given for heart failure with reduced ejection fraction and impaired renal function?

a) Empaglifozin

b) Degludec

c) Linagliptin

d) Rosiglitazone

Question 16:

A patient diagnosed with ulcerative colitis presents with 2-3 episodes of bloody stools per day. What should be given to manage this patient ?

a) Budesonide

b) Azathioprine

c) 6-mercaptopurine

d) 5-aminosalicylic acid

Question 17:

A 25 year old male patient presents with several purpuric skin lesions and is diagnosed with idiopathic thrombocytopenic purpura. Which of the following tyrosine kinase inhibitors can be used in his condition?

a) Eltrombopag

b) Romiplostim

c) Bosutinib

d) Fostamatinib

Question 18:

A 40-year-old woman with multidrug-resistant tuberculosis presents with fatigue, constipation, weight gain, and neck swelling. Which of the following drugs is implicated in this case?

a) Pyrazinamide

b) Ethionamide

c) Cycloserine

d) Streptomycin

Question 19:

A patient on amitriptyline was found in an unconscious state. Which of the following does not suggest amitriptyline toxicity?

a) Arrhythmias

b) Confusion

c) Hypothermia

d) Mydriasis

Question 20:

Which of the following is beneficial in reducing the painful crisis of sickle cell anemia?

a) Hydroxyurea

b) Methotrexate + leukovorin

c) Methionine

d) Eltrombopag

Question 21:

What is the most useful drug to treat doxorubicin induced cardiotoxicity?

a) Dexrazoxane

b) Dexamethasone

c) Aprepitant

d) Amifostine

Question 22:

An elderly patient with dementia is not compliant with oral medications. Which of the following will be beneficial in this case?

a) Donepezil

b) Rivastigmine

c) Galantamine

d) Memantine

Question 23:

Sodium valproate has an inhibitory action at which of the following sites?



a) A

b) B

c) C

d) D

Question 24:

Which of the following drugs acts via the ATP-binding cassette transporter?

a) Verapamil

b) Diltiazem

c) Nifedipine

d) Amlodipine

Question 25:

A 12 year old boy presents with complaints of profuse watery stools and an elevated WBC count of 16000 cells/mm3. He has been on treatment with amoxicillin for the past 1 week. What is the treatment of choice for his condition?

a) Fidaxomycin

b) Clindamycin

c) Zanamivir

d) Cilastatin

Question 26:

Which of the following mood stabilisers causes hepatotoxicity and hyperammonemia?

a) Lithium

b) Valproate

c) Carbamazepine

d) Topiramate

Question 27:

Which of the following pathogens is correctly matched with its intrinsic drug resistance?

a) Aspergillus niger - Voriconazole

b) Candida krusei - Fluconazole

c) Candida Glabrata - Amphotericin B

d) Sporothrix - Itraconazole

Question 28:

A child was receiving a blood transfusion. At what serum iron level will desferrioxamine be administered to him?

a) 350 mcg/dL

b) 400 mcg/dL

c) 450 mcg/dL

d) 500 mcg/dL

Question 29:

A patient on fluoxetine was started on a pain control drug for neuropathic pain. He developed an episode of syncope after 2 weeks. ECG reveals QT interval of 477 ms. Which of the following is the causative drug?

a) Methadone

b) Morphine

c) Meperidine

d) Propoxyphene

Question 30:

All of the following side effects are more commonly seen with carbamazepine than oxcarbazepine except

a) Rashes

b) Hyponatremia

c) Blood dyscrasia

d) Hepatitis

Question 31:

A patient with hemophilia on antiretroviral therapy is brought in altered mental status following complaints of nausea, vomiting and headache. Which of the following medications was he taking?

a) Lopinavir

b) Tipranavir

c) Darunavir

d) Ritonavir

Question 32:

A 60-year-old patient with chronic kidney disease suddenly develops chest pain and palpitations. He had previously complained about feeling weak for the past week. Serum potassium was found to be 6.5mEq/L. Which of the following is likely to be found in this patients ECG?

a) Tall peaked T waves

b) Flat T waves

c) Tall QRS complexes

d) Prolonged PR intervals

Question 33:

Which of the following drugs can be used in diabetic neuropathy, post herpetic neuralgia and spinal cord injury?

a) Amitriptyline

b) Phenytoin

c) Carbamazepine

d) Pregabalin

Question 34:

Which of the following antibiotics acts by inhibiting the 50s ribosome?

a) Aztreonam

b) Daptomycin

c) Linezolid

d) Rifampicin

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	b
5	b
6	b
7	a
8	a
9	a
10	b
11	b
12	a
13	d
14	b
15	a

16	d
17	d
18	b
19	c
20	a
21	a
22	b
23	b
24	a
25	a
26	b
27	b
28	d
29	a
30	b
31	b
32	a
33	d
34	c

Detailed Explanations

Solution to Question 1:

The drug of choice for the reversal of respiratory depression is naloxone. Naloxone is a competitive antagonist at μ-opioid receptors and rapidly reverses the effects of opioid toxicity when administered.

Opioid overdose symptoms:

• Triad of coma, respiratory depression, and pinpoint pupils (miosis).

• Hypotension, bradycardia

• Shallow breathing

• Hypothermia

Naloxone is a pure opioid antagonist which reverses all effects, except sedation. Naloxone is also used for reversing neonatal asphyxia due to opioid usage in labor.

Other options

Option B - Meperidine (pethidine) is a mu-opioid receptor (MOR) agonist. It has a strong analgesic effect on the central nervous system. It is mainly used as an analgesic but can also be

used in post-anesthetic shivering. It has a greater ability to enter the CNS and produce analgesia at lower systemic concentrations compared to morphine. Therefore, constipation, urinary retention, and nausea are less frequent compared to morphine.

Meperidine can cross the placenta and cause significant respiratory depression in the fetus. In patients or addicts who are tolerant to the depressant effects of meperidine, large doses repeated at short intervals may produce an excitatory syndrome that includes hallucinations, tremors, muscle twitches, etc.

Option C -Atropine is an anticholinergic drug. It acts through reversible, competitive inhibition of the postsynaptic muscarinic acetylcholine receptors. It is indicated in the treatment of organophosphate/muscarinic poisoning, in bradycardia, and to reduce salivary secretions.

The adverse effects of atropine are due to the antimuscarinic effects, including xerostomia, blurring of vision, photophobia, tachycardia, flushing, warm, dry skin, constipation, urinary retention, anhidrosis, delirium, and coma. Intoxication with atropine results in all the above symptoms along with convulsions, and circulatory collapse due to respiratory failure. Death may occur secondary to respiratory paralysis.

Option D - Buprenorphine is a partial μ agonist and an antagonist at δ and κ receptors. It is used as an analgesic and in the management of opioid dependence.

Solution to Question 2:

Lapatinib is used for trastuzumab-resistant HER-2 neu-positive breast carcinoma.

Lapatinib is an orally bioavailable, small-molecule inhibitor of the EGFR and HER2 tyrosine kinases. Lapatinib is metabolized by CYP3A4.

Lapatinib in combination with capecitabine is approved for the treatment of patients with metastatic HER2-positive trastuzumab-refractory breast cancer. Lapatinib is also used in combination with the aromatase inhibitor letrozole to treat postmenopausal women with hormone receptor-positive, metastatic breast cancer that overexpresses HER2.

Frequent adverse effects include acneiform rash, diarrhea, cramping, and exacerbation of gastroesophageal reflux. Cardiac toxicity appears less pronounced than with trastuzumab, although lapatinib should be used with caution in combination with cardiotoxic drugs. A dose-dependent prolongation of the QT interval has been reported; thus, careful monitoring of patients with heart disease is recommended. Hepatotoxicity that may be severe has been observed.

Trastuzumab is a monoclonal antibody that inhibits the HER2 receptor. It is used in the treatment of HER2-positive breast cancer.

Other options:

Option A: Vismodegib is a hedgehog pathway inhibitor used to treat metastatic and locally advanced BCCs.

Option C: Vemurafenib is a BRAF inhibitors used in metastatic melanoma and thyroid cancers with BRAF V600E mutation.

Option D: Erlotinib is a protein kinase antagonists which is the first line treatment of non-small cell lung carcinoma with EGFR deletion of exon 19 or L858R mutation

Solution to Question 3:

The nomogram used for vancomycin dosing is the Matzke nomogram.

The Matzke nomogram includes dosing intervals ranging from 0.5 to 12 days on a continuum of CLcr(Creatinine clearance) values from 0 to 120 mL/min. It is designed to guide the determination of the initial empiric dose and interval of intravenous (IV) vancomycin in hospitalized adults who are not receiving hemodialysis. The nomogram does not apply to patients who exhibit highly variable vancomycin pharmacokinetics. These patients include the critically ill, those with burn injuries affecting more than 20 percent of their body surface, patients with rapidly changing renal function, pregnant women, those with liver failure/ascites, individuals at the extremes of weight, patients who are acutely post-transplant, and those with cystic fibrosis

The Rumack-Matthew nomogram uses the serum acetaminophen concentration, in relation to the time after ingestion, to assess potential hepatotoxicity after a single, acute ingestion of acetaminophen.

Solution to Question 4:

The true statement about lithium is it can cause fine tremors in the therapeutic range.

Lithium has a very narrow therapeutic index of 0.8-1.2 mEq/L. The severity of lithium toxicity is often divided into the following three grades:

A fine tremor is common in patients taking lithium at therapeutic doses. Therapeutic lithium tremor is generally symmetric, limited to the arms, and nonprogressive. The tremor from lithium toxicity is coarser and more disabling, may also affect the legs, and occurs with other symptoms of toxicity. Other adverse effects include mental confusion, hypothyroidism, nephrogenic diabetes insipidus, and edema. Lithium use in early pregnancy causes Ebstein anomaly in the fetus(Option D). Fetal lithium exposure is associated with neonatal goiter, CNS depression, hypotonia (floppy baby syndrome), and cardiac murmur.

As lithium is readily dialyzed, lithium overdose or toxicity is treated with hemodialysis.

Lithium is a mood stabilizer with a very narrow therapeutic index. Lithium is administered orally. Lithium is almost completely absorbed from the GI tract(Option A). It is the first line of management in bipolar disorder.

Lithium is also used in the management of mania and has anti-suicidal properties. (Option C) It has a narrow therapeutic index, and blood levels in excess of this may cause symptoms of toxicity. Therapeutic drug monitoring is done by measuring lithium levels 12 hours after the last dose.

Grade	Sr. lithium concentration(in mEq/L)	Symptoms
Mild symptoms	1.5-2.5	Nausea, vomiting,tremor, and fatigue
Moderate intoxication	2.5-3.5	Confusion, agitation, deliriu m, tachycardia, and hyperton ia
Severe intoxication	>3.5	Coma,seizures, hyperthermia, and hypotension

Solution to Question 5:

Mirabegron is a β3-adrenergic receptor agonist which is used for bladder incontinence. Activation of this receptor in the bladder leads to detrusor muscle relaxation and increased bladder capacity. This action prevents voiding and provides relief for those with an overactive bladder and urinary incontinence.

Indacaterol is a very long acting beta 2 agonists (V-LABA) and can be used once daily in COPD maintenance.

Betaxolol is a β1 selective third-generation β-blocker with additional calcium entry blockade activity and a slight membrane-stabilizing property. It does not have agonist activity.

Isoproterenol is a non-selective beta agonist (β1=β2) with no α action. The direct action on cardiac β1 receptors causes an increased risk of tachycardia and arrhythmias. The decrease in peripheral vascular resistance is due to β2 action causing a fall in diastolic blood pressure. Isoproterenol currently may be used as a cardiac stimulant in patients with bradycardia or heart block, particularly in anticipation of inserting an artificial cardiac pacemaker.

A. Mirabegron	3. Beta 3
B. Indacaterol	4. Beta 2
C. Betaxolol	1. Beta 1
D. Isoproterenol	2. Beta

Solution to Question 6:

The therapeutic index of a drug denotes the safety and efficacy of the drug in the population (option B), not the individual (option C).

The therapeutic index is calculated with the following equations:

LD50/ED50 or TD50/ED50

where LD50 is the median lethal dose, TD50 is the mean toxic dose, and ED50 is the median effective dose

The therapeutic index (TI) describes a relationship between the doses of a drug that causes lethal or toxic effects with the dose that causes therapeutic effects. It is also called the therapeutic ratio. The therapeutic index is used to assess the safety of the drug administered.

Median lethal dose (LD50) and median toxic dose (TD50)

• LD50 is the dose of the drug at which 50% of the experimental animals die

• In human trials, TD50 can be considered. It is the dose of the drug that causes adverse effects in 50% of the population (Option D)

• The larger the LD50 or TD50, the safer the drug

Median effective dose (ED50) is the quantity of a drug that can produce the desired therapeutic effects in 50% of the population.

Therapeutic window refers to a range of concentrations at which the likelihood of efficacy is high and the probability of adverse effects is low.

Note: Option D is incorrect as it does not make the necessary distinction that the therapeutic index is the ratio of the median lethal dose in 50% of animals (as the human population cannot be used for lethal studies) to the median effective dose in 50% of the population. LD50 is measured in animals, not in humans. Option B is a better choice.

Solution to Question 7:

The T790M mutation in the EGFR gene is known to cause resistance to first and second-generation EGFR inhibitors. Osimertinib, as a third-generation EGFR inhibitor, is effective against this resistance, hence it is often used for patients with the T790M mutation after they've progressed on first or second-generation EGFR inhibitors.

The question seems to be asking specifically for the mutation that osimertinib is particularly known for being used against, and that would be the T790M mutation.

Osimertinib is a kinase inhibitor that is approved for

• Patients with metastatic non small cell lung carcinoma(NSCLC) that has progressed after prior EGFR TKI treatment and is positive for the EGFR T790M mutation.

• First-line treatment of patients with metastatic NSCLC and adjuvant therapy after resection of NSCLC with EGFR exon 19 deletions or exon 21 L858R mutations

So, osimertinib is used both for the L858R mutation and T790M mutation in different settings. But T790M mutation is often highlighted because of its association with resistance to first and second-generation EGFR inhibitors, and because osimertinib can overcome this resistance.

Osimertinib is administered orally and is a substrate of CYP3A. The drug is primarily eliminated in the feces and to a lesser extent in the urine. Concurrent administration of inducers of CYP3A4 may necessitate an increase in the dose.

The adverse effects after osimertinib treatment are leukopenia, diarrhea, skin rash, interstitial lung disease, QTc prolongation, and left ventricular dysfunction.

Solution to Question 8:

Trazodone causes priapism.

Trazodone is an atypical antidepressant drug. It is a serotonin receptor antagonist. Other drugs which belong to this include nefazodone, mirtazapine, and mianserin. Low doses of trazodone (50–200 mg) have been used widely, both alone and concurrently with SSRIs or SNRIs, to treat insomnia. Trazodone use is associated with priapism in rare instances. Intracavernous injection of phenylephrine can be used to treat priapism.

Other options:

Option B: Venlafaxine is a second-line drug used in generalized anxiety disorder. It belongs to the class of serotonin-norepinephrine reuptake inhibitors (SNRIs). Venlafaxine can induce sustained diastolic hypertension (diastolic blood pressure >90 mmHg at consecutive weekly visits). Venlafaxine also is associated with an increased risk of perinatal complications.

Option C: Bupropion is a first-line drug used for nicotine cessation and has the added benefit of causing weight loss. Weight gain is commonly seen after stopping nicotine use. Hence this drug can be used in this patient as it treats both of these.

Option D: Mirtazapine is used in major depressive disorder. It increases appetite and causes significant weight gain.

Solution to Question 9:

Cyclosporine induced cholestasis is due to MDR1 transporter.

ABC transporters such as MRP2, MDR1, BCRP, BSEP, and MDR3 in the bile canalicular membrane of hepatocytes mediate the efflux (excretion) of drugs and their metabolites, bile salts, and phospholipids against a steep concentration gradient from liver to bile. This primary active transport is driven by ATP hydrolysis.

Cyclosporine, an MDR1 substrate can cause cholestatic liver injury through the following mechanisms:

• Competitive inhibition of ATP dependent transporters.

• Inhibition of intrahepatic vesicle transport and targeting of ATP dependent transporters to the canalicular membrane.

• Impairment of bile secretion partly by increasing canalicular membrane fluidity without affecting the expression of canalicular transporters.

Cyclosporine is an immunomodulatory drug that inhibits calcineurin activity by forming a complex with cyclophilin, a cytoplasmic-receptor protein present in target cells, to form a cyclosporine-cyclophilin complex.Calcineurin is a calcium/calmodulin-dependent phosphatase required for the induction of T cell lymphokine production and proliferation.

This drug-receptor complex specifically and competitively binds to and inhibits calcineurin, a calcium and calmodulin-dependent phosphatase. This, in turn, inhibits the translocation of a family of transcription factors (NF-AT), leading to reduced transcriptional activation of cytokine genes for interleukin (IL)-2, tumor necrosis factor (TNF)-alpha, IL-3, IL-4, CD40L, granulocyte-macrophage colony-stimulating factor, and interferon-gamma. Ultimately, the proliferation of T lymphocytes is reduced, resulting in immunomodulation.

Its uses are as follows:

• In solid organ transplantation, to prevent organ rejection in kidney, liver, and heart allogeneic transplants

• In patients with rheumatoid arthritis, when the disease has not adequately responded to methotrexate

• Cyclosporine reportedly is effective in Behc■et's disease, acute ocular syndrome, endogenous uveitis, atopic dermatitis, inflammatory bowel disease, and nephrotic syndrome.

Adverse effects of cyclosporine include a decrease in glomerular filtration rate (GFR) and metabolic disturbances like hypomagnesemia, hyperkalemia, dyslipidemia, gynecomastia, hypertrichosis; neurotoxicity in the form of convulsions. There is an increased risk of developing skin and lymphoproliferative malignancies in cyclosporine-treated psoriasis patients.

Solution to Question 10:

In the above clinical scenario, the most likely cause of itching around the vulvovaginal area is canagliflozin. The side effects associated with canagliflozin are urinary tract infections, genital mycotic infections due to the high glucose content of the urine, increased risk of fractures, and lower extremity amputations.

Canagliflozin is a sodium-glucose transporter 2 (SGLT-2) inhibitor. Other SGLT-2 inhibitors include dapagliflozin and empagliflozin. They act by inhibiting the SGLT-2 transporter which is present on the proximal segment of the renal tubule. It is responsible for the reabsorption of 80-90% of filtered glucose. It inhibits glucose reabsorption resulting in glycosuria and a change in the renal threshold.

SGLT2 inhibitors can cause mild weight loss and reduction in blood pressure. Drugs like empagliflozin and canagliflozin are found to reduce major cardiovascular events.

The common side effects include urinary tract infections and genital mycotic infections due to the high glucose content of the urine. There is a risk of the development of diabetic ketoacidosis. It increases the risk of fractures and lower limb amputations due to the creation of an imbalance with the minerals and circulating parathormone and vitamin D. Since SGLT2 inhibitors depend on the rate of glucose filtration to be effective, potency decreases if given in stage 3 kidney disease (GFR 60–30 mL/min).

Other options:

Option A: Acarbose is an alpha-glucosidase inhibitor. It reduces the intestinal absorption of glucose by inhibiting the intestinal brush border enzyme, alpha-glucosidase. These drugs also increase GLP-1 levels in circulation. They reduce post-prandial glucose levels.

Option C: Linagliptin is a dipeptidyl-peptidase-4 (DPP-4) inhibitor. It is used in type 2 diabetes as monotherapy or in combination therapy when metformin is not tolerated or contraindicated. It binds extensively to plasma proteins and is cleared primarily by the hepatobiliary system. Hence, no dosage adjustment is needed in renal failure. Dipeptidyl peptidase-4 inhibitors enhance the release of insulin postprandially, decrease glucagon, and thereby lower mealtime as well as fasting glucose. They are weight neutral.

Option D: Liraglutide is also a GLP-1 receptor agonist and is given only in the subcutaneous injectable form. It was recently approved in the treatment of obesity.

Solution to Question 11:

Alogliptin is a dipeptidyl peptidase-4 (DPP) inhibitor. Glucagon-like peptide (GLP) and glucose-dependent insulinotropic polypeptide (GIP) are both incretins that promote the exocytosis of insulin into circulation. Both these incretins are rapidly inactivated by dipeptidyl peptidase. So, inhibition of dipeptidyl peptidase enhances the action of incretins and facilitates insulin release. Dipeptidyl peptidase-4 inhibitors enhance the release of insulin postprandially, decrease glucagon, and thereby lower mealtime as well as fasting glucose.

Other DPP-4 inhibitors include saxagliptin, linagliptin, alogliptin, and vildagliptin. They are weight neutral.

DPP-4 inhibitors are associated with severe joint pain which reduces on discontinuation of the drug. Linagliptin is cleared primarily by hepatobiliary system, with little renal clearance. Only saxagliptin is metabolized by the liver and must be used in lower doses if used along with a cytochrome p450 enzyme inhibitor drug. Other drugs are renally excreted and hence must be used with caution in renal insufficiency.

Other options:

Option A - Pramlintide is an amylin agonist. It reduces postprandial glycemia and leads to weight loss.

Option C - Liraglutide is also a GLP-1 receptor agonist and is given only in the subcutaneous injectable form. It was recently approved for the treatment of obesity.

Option D - Metformin belongs to the biguanides class of oral hypoglycemic agents. It acts by activation of AMP-dependent protein kinase (AMPK). This results in reduced hepatic glucose production. Side effects of metformin are lactic acidosis, gastrointestinal disturbance, and vitamin B12 deficiency.

Solution to Question 12:

Keynote 189 trial is a efficacy and safety study of pembrolizumab and chemotherapy in metastatic non-small cell carcinoma of lung.

Keynote 189 trial is an efficacy and safety study of pembrolizumab (MK-3475) combined with pemetrexed/platinum chemotherapy versus pemetrexed/platinum chemotherapy alone in participants with advanced or metastatic nonsquamous non-small cell lung cancer (NSCLC) who

have not previously received systemic therapy for advanced disease. Participants will be randomly assigned to receive pembrolizumab combined with pemetrexed/platinum.

Methods

In this double-blind, phase 3 trial, patients were randomly assigned (in a 2:1 ratio). 616 patients with metastatic nonsquamous NSCLC without sensitizing EGFR or ALK mutations who had received no previous treatment for metastatic disease to receive pemetrexed and a platinum-based drug plus either 200 mg of pembrolizumab or placebo every 3 weeks for 4 cycles, followed by pembrolizumab or placebo for up to a total of 35 cycles plus pemetrexed maintenance therapy. Crossover to pembrolizumab monotherapy was permitted among the patients in the placebo-combination group who had verified disease progression. The primary endpoints were overall survival and progression-free survival, as assessed by a blinded, independent central radiologic review.

Conclusions

In patients with previously untreated metastatic nonsquamous NSCLC
without EGFR or ALK mutations, the addition of pembrolizumab to standard chemotherapy of pemetrexed and a platinum-based drug resulted in significantly longer overall survival and progression-free survival than chemotherapy alone.

Solution to Question 13:

Thalidomide does not cause teratogenicity by inhibition of IL2, IL4, and IL5.

The mechanism of thalidomide teratogenicity include oxidative stress, anti-angiogenesis, and thalidomide-binding protein, cereblon which has a role in the development of cerebral tissues and because its expression in the hippocampus among other areas is associated with memory and learning processes (CRBN). Fetal thalidomide exposure causes phocomelia. This disorder is associated with underdeveloped limbs and absent pelvic bones. Both upper and lower limbs are affected, and a part of the limbs may be missing. Fused fingers and missing thumbs are also seen. Thalidomide was originally used for the treatment of pregnancy-associated morning sickness but was withdrawn due to this adverse effect. It is now used for the treatment of erythema nodosum leprosum.

Thalidomide is an immunomodulator. Thalidomide absorption from the GI tract is slow and highly variable. It distributes throughout most tissues and organs, without significant binding to plasma proteins. Thalidomide and its metabolites are excreted in the urine, while the nonabsorbed portion of the drug is excreted unchanged in feces.

Common adverse effects of thalidomide include sedation and constipation. However, one of the most serious adverse effects is peripheral sensory neuropathy, characterized by asymmetrical, painful, peripheral paresthesia with sensory loss. Symptoms often present as numbness of the toes and feet, muscle cramps, weakness, signs of pyramidal tract involvement, and carpal tunnel syndrome.

Solution to Question 14:

Agranulocytosis is a rare but very severe side effect of clozapine that requires regular monitoring.

A patient who is on clozapine should undergo regular hematological monitoring due to the risk of agranulocytosis. Avoid combining clozapine with other agranulocytosis-causing drugs like carbamazepine, phenytoin, propylthiouracil, and sulfonamides.

Adverse effects of clozapine are:

• Sedation and weight gain- H1 antagonism

• Syncope, tachycardia, hypotension- α1 antagonism

• Hypersalivation/sialorrhea- M4 agonism

• Seizures- dose-dependant

• Agranulocytosis- dose-independent

Clozapine reduces the seizure threshold thus increasing the risk of seizures. Clozapine, an atypical antipsychotic, is an antagonist of 5HT2A, D1, D3, D4, and α-receptors. Clozapine is used in treatment-resistant schizophrenia, suicidal patients with schizophrenia, severe psychotic depression, and Huntington's disease.

Solution to Question 15:

Empaglifozin, which is a sodium-glucose transporter-2 (SGLT-2) inhibitor has been recently approved for use in patients with Heart Failure with Reduced Ejection Fraction, and can be initiated in patients with impaired renal function with estimated glomerular filtration rate > 20 mL/min/1.73 m2.

Empagliflozin works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) in the proximal renal tubules resulting in reduced reabsorption and increased urinary excretion of glucose. The glucose-lowering effect is independent of insulin. Empagliflozin acts as a diuretic and natriuretic by reducing sodium and volume load, causing intravascular contraction. Empagliflozin is also associated with weight loss, with reductions in blood pressure without increasing heart rate. Empagliflozin has been found to decrease the risk of mortality from cardiovascular causes in diabetic patients.

SGLT-2 inhibitors have a cardioprotective effect resulting from improved cardiac cell metabolism, improvement of ventricular loading mechanism, reduction of cardiac cell necrosis and fibrosis. It also has a renoprotective property.

Except for empaglifozin which can be used in chronic kidney disease, use of other SGLT-2 inhibitors might lead to acute kidney injury and should be avoided in patients with renal failure. Side effects of empaglifozin includes dehydration, euglycemic diabetic ketoacidosis, urinary tract infections, genital mycotic infections, hypoglycemia and necrotizing fasciitis.

Side effects of SGLT-2 inhibitors include dehydration, hyperkalemia, worsening of diabetic ketoacidosis, increased risk of fractures, leg and foot amputation with canagliflozin, and a possible increased risk of bladder cancer with dapagliflozin. Dapagliflozin has also been recently approved by the FDA for treatment of Heart failure with Reduced Ejection Fraction in both diabetic and non-diabetic patients, it has been associated with increased risk of bladder cancer and breast cancer.

Other options :

Option B : Degludec is a form of long acting insulin, and is not indicated for the treatment of heart failure.

Option C : Linagliptin is a dipeptidyl peptidase-4 (DPP-4) inhibitor, which enhances the action of incretins that promote the release of insulin into circulation. Dipeptidyl peptidase-4 inhibitors enhance the release of insulin postprandially, decrease glucagon, and thereby lower mealtime as well as fasting glucose. Linagliptin is safe to use in chronic kidney disease as it is cleared primarily by hepatobiliary system, but is not approved for heart failure.

Option D : Rosiglitazone is a thiazolidinedione, which activates PPAR-γ receptors in adipose tissue and skeletal muscles. These are insulin sensitizers and increase insulin-mediated glucose uptake in patients with type 2 diabetes. Rosiglitazone is currently banned in India as it has shown an increased risk of developing myocardial infarction, stroke, congestive heart failure, and death.

Solution to Question 16:

5-aminosalicylic acid is is effective in inducing and maintaining remission and is considered to be the first line agents for mild to moderate ulcerative colitis, seen in this patient with 2-3 bloody stools per day.

Treatment of ulcerative colitis differs based on the severity of the disease and involves medical and surgical management. Patients with mild to moderate disease are treated with topical or systemic 5-aminosalicylate drugs such as Salazopyrine/Sulfasalazine and Mesalamine (5 ASA) which are first-line therapy. Glucocorticoids (option A), immunosuppressants, and biologics are effective in patients with moderate to severe disease.

Azathioprine and cyclosporin are used as steroid-sparing agents. Anti-TNF alpha inhibitors (infliximab, adalimumab) can be used as anti-inflammatories.

Surgery is recommended for patients who do not improve after medical treatment or with toxic megacolon, and perforation. Indications for elective surgery include failure of medical therapy, growth retardation, extraintestinal disease, and malignant change.

Ulcerative colitis is an inflammatory bowel disease resulting in superficial erosions confined to the mucosa and submucosa of the colon. It starts from the rectum and extends proximally in a continuous manner. The clinical features include abdominal pain, diarrhea, rectal bleeding, tenesmus, and passage of mucus. Other symptoms include anorexia, nausea, vomiting, fever, and weight loss.

Ulcerative colitis is graded based on the frequency of bowel movements and systemic involvement.

- Mild disease - ≤4 stools per day with/without rectal bleeding;

- Moderate disease - ≥4 stools per day with few systemic signs of illness;

- Severe disease - ≥6 blood stools per day with fever, tachycardia (≥90 beats/min), anemia (hemoglobin <10.5 g/dL), and elevated levels of inflammatory markers (ESR ≥30 mm/h)

- Fulminant disease - >10 stools per day with severe anemia requiring blood transfusion and progressive colonic dilation (toxic megacolon)

Investigations show elevated C-reactive protein, platelet count, erythrocyte sedimentation rate, fecal calprotectin, and fecal lactoferrin. A colonoscopy reveals loss of typical vascular pattern, erythema, friability, regenerative mucosal nodules, pseudopolyps, and ulcerations.

Extraintestinal manifestations associated with IBD include ocular features (episcleritis, and uveitis), dermatological manifestations (erythema nodosum, pyoderma gangrenosum), rheumatological (arthropathies, sacroiliitis, ankylosing spondylitis) and primary sclerosing cholangitis. Complications include toxic megacolon, hemorrhage, strictures, and perforations. These patients are also at increased risk of colon cancer.

Other options

Option B & C : Azathioprine and 6-mercaptopurine are not first line agents for ulcerative colitis.

Solution to Question 17:

Fostamatinib is a tyrosine kinase inhibitor used in the treatment of chronic immune thrombocytopenia (idiopathic thrombocytopenic purpura).

Fostamatinib has activity against spleen tyrosine kinase (SYK). The major metabolite of fostamatinib, R406, inhibits signal transduction of Fc-activating receptors and B-cell receptors. The fostamatinib metabolite R406 reduces antibody-mediated destruction of platelets.

Idiopathic thrombocytopenic purpura (ITP) is an acquired disorder characterized by the immune-mediated destruction of platelets and possibly inhibition of platelet release from the megakaryocyte. It is an autoimmune disorder caused by antibodies directed against platelet membrane glycoproteins IIb-IIIa or Ib-IX. Secondary causes of ITP include systemic lupus erythematosus (SLE), HIV-AIDS, and Hepatitis C.

ITP usually presents with mucocutaneous bleeding, very low platelet count, and normal peripheral blood cells. Patients typically exhibit ecchymoses and petechiae or they may incidentally discover thrombocytopenia during a routine complete blood count. Rarely, life-threatening bleeding may occur. The presence of wet purpura (blood blisters in the mouth) and retinal hemorrhages can serve as warning signs of potentially life-threatening bleeding.

The treatment of ITP is as follows:

Other options :

Option A and B: Eltrombopag and Romiplostim are thrombopoietin (TPO) receptor agonists used in the treatment of immune thrombocytopenia which have failed to respond to conventional treatments. Both drugs act via the Mpl receptor to increase platelet counts and not via tyrosine kinase inhibition.

Option C: Bosutinib is a BCR-ABL tyrosine kinase inhibitor indicated in CML.

	Clinical features	Management
1.	Minor bleeding manifestations (and/or)Platelet count > 5000/ μL	Prednisone (OR)Anti-D immunoglobulin therapy - effective only in Rh-positive patients (OR)Intravenous immunoglobulin (IVIG)
2.	Severe ITP (any of the following):- Significant GI / intracranial bleeding- Platelet count < 5000/μL- Signs of impending bleeding (retinal or large oral mucosal bleeds)	Platelet transfusion (AND)High-dose glucocorticoids (AND)IVIG / Anti-D immunoglobulins + Rituximab
3.	Relapse after first-line therapy	Splenectomy
4.	Unresponsive to all other therapy or relapse after splenectomy	Thrombopoietin receptor agonists - romiplostim (subcutaneous), eltrombopag (oral)

Solution to Question 18:

The given clinical features are suggestive of hypothyroidism with goiter, which is a known adverse effect of ethionamide.

Mutidrug-resistant tuberculosis (MDR-TB) is treated with 2nd line agents that include ethionamide (ETH). ETH is not a first-line agent due to its gastrointestinal side effects. Ethionamide inhibits activity of the InhA gene product, resulting in inhibition of mycolic acid synthesis. It is structurally related to thionamides (propylthiouracil and methimazole) and therefore causes hypothyroidism by preventing organification of iodine in thyroid hormone synthesis. Another antitubercular drug that can cause hypothyroidism is Para-amino salicylic acid (PAS).

ETH/PAS-induced hypothyroidism should be treated with thyroxine replacement therapy, instead of changing the regimen. Thyroid function usually optimizes within 2 months after completing MDR therapy.

Ethionamide also causes hepatotoxicity and neurological adverse effects and may require pyridoxine supplementation.

MDR-TB is when there is resistance to atleast isoniazid and rifampicin which are the most potent first-line drugs. The treatment of MDR-TB is curated with drugs grouped into different classes by the world health organization (WHO).

Group A: Levofloxacin, bedaquiline, linezolid

Group B: Clofazimine, cycloserine

Group C: Ethambutol, pyrazinamide, delamanid, ethionamide

In patients treated previously or with additional drug resistance, a longer MDR-TB regimen of 18-20 months is recommended. Drugs belonging to group A are preferred along with one drug from group B whenever possible. A shorter regimen of 6 months including bedaquiline with other

drugs can be used in patients not treated before and without extensive disease. In pregnant women with MRD-TB, drugs such as bedaquiline and delamanid should be avoided.

Other options :

Option A : Pyrazinamide inhibits fatty acid synthase-I leading to inhibition of cell membrane synthesis. Pyrazinamide causes hyperuricemia and hepatotoxicity.

Option C : Adverse effects of cycloserine are mainly neurological. Half of the patients on cycloserine (1g/day) experience neuropsychiatric symptoms. It ranges from headaches to psychosis, seizures, and suicidal ideas.

Option D : Streptomycin is an aminoglycoside which binds to the 30S ribosomal subunit and interferes with the initiation of protein synthesis. It is a second-line agent for the treatment of active tuberculosis. Streptomycin produces ototoxicity, vestibulotoxicity, optic nerve dysfunction, including scotomas, and peripheral neuritis.

Solution to Question 19:

Hypothermia is not associated with amitriptyline use or tricyclic antidepressant toxicity.

Tricyclic antidepressants (TCAs) such as amitriptyline are used in the treatment of psychotic depression, insomnia, and several pain conditions. Their mechanism of action is inhibition of norepinephrine and serotonin reuptake. They have a quinidine-like sodium channel blockade effect on cardiac conduction. Hence, they cause QT prolongation. Other cardiotoxic effects include atrioventricular (AV) block and hypotension. Rightward-axis deviation is also an associated finding.

TCA toxicity can result in ventricular tachyarrhythmias (Option A), status epilepticus, hyperthermia, and coma. It can also result in anticholinergic syndrome due to muscarinic receptor antagonism. Symptoms include dilated pupils (Option D), high temperature, confusion (option B), dry mouth, urinary retention, tachycardia, and myoclonic movements. It is managed with intravenous sodium bicarbonate bolus (50-100 mEq). This helps in the reversal of sodium channel blockade. Gastric lavage and administration of activated charcoal can be carried out in cases of recent ingestion.

Other side effects of TCAs include postural hypotension from anti-alpha 1 adrenergic action and sedation from anti-histaminic action. ECG changes such as T wave inversion or suppression are also common.

In some cases, it can result in polymorphic ventricular tachycardias with QT prolongation known as torsades de pointes. Recurrent episodes can be treated with intravenous magnesium sulfate.

Solution to Question 20:

Hydroxyurea is beneficial in reducing the painful vasoocclusive crisis of sickle cell anemia.

Hydroxyurea is a myelosuppressive agent that is effective in reducing the frequency of painful episodes like dactylitis and acute chest syndrome, decreasing the rate of hospitalization, and requirement for blood transfusion. It is safe and well-tolerated in children and infants.

Hydroxyurea is approved for long-term maintenance of sickle cell disease, and is recommended for patients of all ages regardless of symptoms.

Hydroxyurea increases fetal hemoglobin levels by an unknown mechanism and interferes with the polymerization of abnormal HbS.

Newer drugs for management of acute painful crisis of sickle cell anemia include L-glutamine and crizanlizumab. For individuals >5 years of age with sickle cell disease who have pain despite hydroxyurea (or who cannot tolerate hydroxyurea), L-glutamine is recommended, regardless of whether they are receiving hydroxyurea. The mechanism of action of this drug, is that it reduces oxidative stress in sickle erythrocytes. Crizanlizumab is a P-selectin-blocking monoclonal antibody which is given intravenously every month to reduce acute painful episodes.

Sickle cell anemia is an autosomal recessive disorder that occurs due to the point mutation in the β-globin gene which results in the replacement of glutamate residue with valine. This causes the production of sickle-shaped red blood cells due to the polymerization of deoxygenated hemoglobin (HbS). Factors that predispose to sickling are infection, dehydration, and acidosis. Sickle cell disease causes hemolytic anemia due to increased erythrocyte destruction. The anemia is generally well compensated. Complications occur due to abnormal erythrocyte configurations that increase hemoglobin polymerization leading to a painful vaso-occlusive state.

Clinical features of sickle cell anemia :

- Moderately severe hemolytic anemia with jaundice.

- Painful bone crisis : hand-foot syndrome or dactylitis of the bones of hands and feet.

- Acute chest syndrome typically presents with fever, cough, chest pain, and pulmonary infiltrates.

- Hyposthenuria : inability to concentrate urine.

- Aplastic crisis stems from the infection of red cell progenitors by parvovirus B19, which causes a transient cessation of erythropoiesis and a sudden worsening of the anemia.

- Spleen : In children, splenic sequestration crises occur which results in splenomegaly due to red pulp congestion. In adolescence or early adulthood, autosplenectomy occurs due to splenic infarction, fibrosis, and progressive shrinkage.

- Increased susceptibility to infection with encapsulated organisms like pneumococcal, meningococcal, and Haemophilus influenza septicemia, but can be reduced by vaccination and prophylactic antibiotics.

- Other clinical features include priapism, leg ulcers, stroke, retinopathy, pigment gallstones, and growth retardation.

Other options :

Option B : Methotrexate interferes with folate metabolism by acting as a competitive inhibitor of the enzyme dihydrofolate reductase. This starves cells for tetrahydrofolate and makes them unable to synthesize purines and thymidine monophosphate. It does not treat painful crisis of sickle cell anemia.

Option C : Methionine is an essential amino acid involved in the synthesis of folate and vitamin B12. It is not used in the management of sickle cell anemia.

Option D : Eltrombopag is a thrombopoetin analog used to treat immune thrombocytopenia. It is not used in the treatment of painful sickle cell crisis.

Solution to Question 21:

Dexrazoxane is the drug of choice to treat doxorubicin induced cardiomyopathy.

Doxorubicin is an anthracycline group of anticancer antibiotics that acts by inhibiting topoisomerase II, and is known to cause cardiotoxicity. Dexrazoxane is a cardioprotective iron-chelating agent that is indicated for use along with high dose doxorubicin to avoid cardiotoxicity.

Dexrazoxane is currently approved to prevent or reduce anthracycline-induced cardiotoxicity in women with metastatic breast cancer who have received a total cumulative dose of doxorubicin of 300 mg/m2 .

Doxorubicin-induced cardiomyopathy takes two forms:

- Acute: pericarditis-myocarditis syndrome, arrhythmias, ST and T wave changes in ECG. Occurs within the first 2–3 days, is transient and in most cases asymptomatic.

- Chronic: Dose dependendent dilated cardiomyopathy associated with heart failure.

Doxorubicin, daunorubicin and epirubicin share similar cardiotoxicity profile. Idarubicin has least cardiotoxicity when compared to other anthracyclines.

Endomyocardial biopsy showing vacuolar degeneration and myofibrillar loss is seen in cardiotoxicity due to anthracyclines.

Difference between cardiotoxicity due to anthracyclins and trastuzumab:

Anthracyclines like doxorubicin can result in a radiation recall reaction, where erythema and desquamation of skin are observed at sites of previous radiation injury. The risk of cardiotoxicity also increases with previous irradiation. Therefore, careful monitoring and dosage adjustments are required for doxorubicin.

Other options :

Option B, C and D : Dexamethasone, aprepitant and amifostine are not used in the management of anthracycline induced cardiotoxicity.

Anthracyclins	Trastuzumab
Irreversible	Reversible
Dose-dependent	Not dose-dependent
Biopsy shows myocyte destruction	Biopsy does not show myocyte destruction

Solution to Question 22:

Rivastigmine transdermal patch can be prescribed to patients with dementia who are non compliant on oral medications.

Rivastigmine is a cholinesterase inhibitor that is available as a transdermal patch for use in alzheimer's disease and dementia. It can be applied for over 24 hours. Transdermal delivery reduces the fluctuation and avoids direct gastric irritation effects associated with oral formulations.

Alzheimer's disease is the most common cause of dementia in the elderly. The characteristic feature of Alzheimer's is the degeneration of cholinergic neurons leading to reduced levels of acetylcholine.

Treatment includes cholinesterase inhibitors like Donepezil, Rivastigmine and Galantamine and NMDA receptor antagonist like memantine. The cholinesterase inhibitors cause significant side effects such as nausea, vomiting, diarrhea and peripheral cholinergic effects, which can be avoided by using a transdermal formulation.

Anticholinesterase drugs

Other options :

Options A, C and D: Donepezil, galantamine and memantine are currently available in the oral form and transdermal formulations are under development and yet to be available in the market.

Reversible inhibitors	
Carbamates	Non-Carbamates
Physostigmine, Neostigmine, Pyridostigmine, Rivastigmine	Edrophonium, Tacrine, Donepezil, Galantamine
Irreversible inhibitors	
Carbamates	Organophosphates
Carbaryl, Propoxur	Dyflos, Echthiophate, Malathion, Diazinon, Tabun, Sarin, Soman

Solution to Question 23:

Sodium valproate has an inhibitory action on the metabolism and degradation of GABA, and also increases its synthesis.

Valproate is an anti-epileptic medication and also used as a mood stabilizer in bipolar disorder. Valproate acts via multiple mechanisms. It blocks both sodium and calcium channels. It prolongs sodium channel inactivation and attenuates calcium-mediated 'T' current. It inhibits metabolism and degradation of the inhibitory neurotransmitter GABA and increases its synthesis from glutamic acid (Option A). It also inhibits excitatory NMDA glutamate receptors. The repetitive firing caused by the depolarization of cortical or spinal cord neurons is inhibited by valproate.

Valproate is a broad spectrum antiseizure drug used in partial seizures, generalised tonic-clonic seziures (GTCS), absence seizures, myoclonic seziures and atonic seziures. It can be used in absence seizures, but it is preferred when the child also suffers from GTCS. It is the drug of choice for juvenile myoclonic epilepsy.

The side effects of valproate (mnemonic: VALPROATE)

Vomiting

Alopecia

Liver toxicity

Pancreatitis

Rash

Obesity

Anorexia

Teratogenicity

Edema

It causes congenital anomalies like neural tube defects (Spina Bifida), clefts, and cardiac anomalies. The intelligence quotient score of school-going children who were exposed in utero to valproate is found reduced. It also contributes to poor cognitive development in children. Folate supplementation reduces the chances of neural tube defects.

Other options :

Option C : Drugs such as vigabatrine and tiagabine inhibit GABA reuptake by inhibiting the GABA reuptake transporter. Vigabatrine is the drug of choice for infantile spasms (West syndrome) seen in tuberous sclerosis.

Option D : Flumazenil acts as an antagonist at the GABA receptor and is used as an antidote for benzodiazepine overdose. Benzodiazepines act by promoting GABA inhibition by binding to the GABA receptor itself and inhibiting nerve transmission.

Solution to Question 24:

Verapamil is a potent inhibitor of P-glycoprotein drug transporter which belongs to the family of ATP-binding cassette (ABC) transporters.

A group of primary active transporters, the ATP-binding cassette (ABC) family, hydrolyze ATP to export substrates across membranes. P-glycoprotein (P-gp) is an ATP-binding cassette (ABC) transporter, which acts as an efflux membrane transporter. It is responsible for limiting cellular uptake and the distribution of xenobiotics and toxic substances. P-glycoprotein is also called ABCB1 or MDR1, and it exports bulky neutral or cationic compounds from cells. MDR1 exports many drugs as well. It is associated with multidrug resistance (MDR) to cancer chemotherapeutics through the multidrug resistance protein 1 or MDR1.

Verapamil can cause interactions with drugs whose disposition is dependent upon P-glycoprotein. Drugs that inhibit P-glycoprotein are -

• Clarithromycin

• Erythromycin

• Amiodarone

• Quinidine

• Verapamil

• Diltiazem

• Cyclosporine

• Itraconazole

• Propafenone

• Flecainide.

P-glycoprotein pumps play a major role in the elimination of certain drugs such as digoxin. Drugs that inhibit P-glycoprotein can thereby decrease their elimination and lead to toxicity. For eg, digoxin is metabolized by P-glycoprotein, and drugs that inhibit it increase the plasma levels of digoxin and may result in toxicity.

Increased P-gp expression: Reduces the absorption of drugs that are substrates for P-gp.

Decreased P-gp expression: Leads to supratherapeutic plasma concentrations and drug toxicity.

Other options :

Option B, C, and D: Although diltiazem, nifedipine, and amlodipine have been reported in the literature to act as p-glycoprotein inhibitors, verapamil is more potent and commonly used for p-glycoprotein inhibition in trials, and would be the best answer.

Solution to Question 25:

Oral Fidaxomicin is the antibiotic indicated for this patient's condition which is suggestive of Clostridium difficile infection, characterized by profuse watery diarrhea and leukocytosis following long term antibiotic use such as ampicillin.

C difficle infection is most commonly associated with prolonged antimicrobial use which interferes with the commensal colonic microflora. The most commonly implicated antibiotics are cephalosporins, ampicillin, clindamycin, and fluoroquinolones. Spores of toxigenic C. difficile when ingested, survive gastric acidity and colonize the lower intestinal tract. Here they produce two toxins: toxin A (an enterotoxin) and toxin B (a cytotoxin). These toxins initiate processes that disrupt the epithelial-cell barrier function, diarrhea, and pseudomembrane formation. The patient presents with non bloody diarrhea, fever, abdominal pain, and leukocytosis.

Clinical features can range from asymptomatic carriage to fulminant colitis.

• Nonsevere disease - Watery diarrhea (≥3 loose stools in 24 hours) is the cardinal symptom. Other manifestations include lower abdominal pain and cramping, low-grade fever, nausea, anorexia, mucus, or occult blood in the stool. Physical examination may demonstrate lower abdominal tenderness. Unexplained leukocytosis in hospitalized patients (even in the absence of diarrhea) may reflect underlying CDI

• Severe colitis - Clinical manifestations of severe colitis include diarrhea, lower quadrant or diffuse abdominal pain, abdominal distention, fever, hypovolemia, lactic acidosis, hypoalbuminemia, elevated creatinine, and marked leukocytosis.

• Fulminant colitis - Characterized by hypotension or shock, ileus, or megacolon.

• Recurrent disease - resolution of CDI followed by recurrence of symptoms within two to eight weeks after treatment has been stopped

Nucleic acid amplification tests including polymerase chain reaction (PCR) assays, for C. difficile toxin A or B gene in stool, are widely used diagnostically. Glutamate dehydrogenase is an enzyme immunoassay to detect C. difficile in stool. Confirmation of the presence of toxin in the stool in addition to PCR or glutamate dehydrogenase positivity is recommended.

Oral vancomycin, fidaxomicin, and metronidazole are used in the management of this condition. Fidaxomicin acts by inhibiting protein synthesis through the blockade of RNA polymerase.

Management of nonsevere to severe initial disease:

Management of recurrent disease:

Management of fulminant disease -

Other options:

Option B : Clindamycin is implicated as a risk factor for development of clostridium difficile colitis and is not used to treat it.

Option C : Zanamavir is a neuraminidase inhibitor used against influenza A and B virus.

Option D : Cilastatin is a reversible dehydropeptidase inhibitor and used adjunctively with imipenem to prolong its action as it rapidly hydrolysed by renal dehydropeptidase. It is not used in the treatment of clostridium difficile colitis.

Nonsevere diseaseSupportive clinical data: White blood cell count ≤15,000 cells/mL and serum creatinine <1.5 mg/dL	Vancomycin 125 mg orally 4 times daily for 10 days,ORFidaxomicin200 mg orally twice daily for 10 daysIf above agents are unavailable: Metronidazole 500 mg orally 3 times daily for 10 days
Severe diseaseSupportive clinical data: White blood cell count >15,000 cells/mL and/or serum creatinine ≥1.5 mg/dL	Vancomycin 125 mg orally 4 times daily for 10 days,ORFidaxomicin200 mg orally twice daily for 10 days

First recurrence	If vancomycin was used for t he initial episode: Vancomyci n pulsed-tapered regimen:12 5 mg orally 4 times daily for 10 to 14 days, then125 mg or ally twice daily for 7 days, the n125 mg orally once daily for 7 days, then125 mg orally eve ry 2 or 3 days for 2 to 8 week s,ORFidaxomicin 200 mg ora lly twice daily for 10 daysIf fi daxomicin or metronidazole was used for the initial episo de: Vancomycin 125 mg orall y 4 times daily for 10 days
Second or subsequent recurr ence	Vancomycin pulsed-tapered r egimen (outlined above),ORF idaxomicin200 mg orally twi ce daily for 10 days,ORVanco mycin followed by rifaximin: Vancomycin 125 mg orally 4 t imes per day for 10 days, the nRifaximin 400 mg 3 times d aily for 20 days,ORFMT

Initial episode	Enteric vancomycin plus pare nteral metronidazole:Vancom ycin 500 mg orally or via nas ogastric tube 4 times daily,A NDMetronidazole 500 mg int ravenously every 8 hoursIf ile us is present, additional cons iderations include:FMT (adm inistered rectally)Rectal vanc omycin (administered as a re tention enema 500 mg in 10 0 mL normal saline per rectu m; retained for as long as pos sible and readministered ever y 6 hours)
Recurrent episo de	Antibiotics as for initial fulmi nant episode aboveSome favo r FMT in the context of a firs t or second recurrence that p resents as a fulminant episod e

Solution to Question 26:

Valproate is a mood stabilizer with the side effect of hepatotoxicity and hyperammonemia.

Valproate is an anti-epileptic medication and also used as a mood stabilizer in manic episodes as well as in bipolar disorder. The repetitive firing caused by the depolarization of cortical or spinal cord neurons is inhibited by valproate. It also prolongs the sodium channel inactivation. Valproate can cause an idiosyncratic hepatic toxicity and interfere with conversion of ammonia to urea leading to hyperammonemia. Valproate is contraindicated in patients with genetic defects in urea metabolism.

Valproate therapy reduces carnitine levels in tissues, potentially impacting mitochondrial function and leading to hyperammonemia and microvesicular steatosis. Genetic factors also play a role, as individuals who are heterozygous for mutations in the gamma polymerase gene, which is primarily present in mitochondria, are more susceptible to valproate-induced liver damage. Intravenous administration of carnitine can be used to manage this condition.

The side effects of valproate (mnemonic: VALPROATE)

Vomiting

Alopecia

Liver toxicity

Pancreatitis

Rash

Obesity

Anorexia

Teratogenicity

Edema

It causes congenital anomalies like neural tube defects (Spina Bifida), clefts, and cardiac anomalies. The intelligence quotient score of school-going children who were exposed in utero to valproate is found reduced. It also contributes to poor cognitive development in children. Folate supplementation reduces the chances of neural tube defects.

Other options :

Option A : Lithium is a mood stabilizer with a very narrow therapeutic index and is the first line of management in bipolar disorder. Its adverse effects can be acute such as tremors, ataxia, weakness, polyuria, polydipsia or chronic such as nephrogenic diabetes insipidus, thyroid dysfunction, and hyperparathyroidism. It is not associated with hepatotoxicity or hyperammonemia.

Option C : Carbamazepine is a sodium channel blocker used in the management of complex partial seizures, GTCS, simple partial seizures and trigeminal neuralgia. It is known to be hepatotoxic but is not associated with hyperammonemia. Other adverse effects include hypersensitivity reactions, eosinophilia, agranulocytosis, aplastic anemia, hyponatremia and stevens-johnson syndrome.

Option D : Topiramate is an anti epileptic and non-opioid pain medication which is associated with significant weight loss of up to 7% as a side effect.

Solution to Question 27:

Among the given options, Candida krusei shows an intrinsic resistance to fluconazole.

The primary or intrinsic resistance of C. krusei to azoles (fluconazole and itraconazole) is mediated through reduced susceptibility of the target enzyme (lanosterol 14α-demethylase) to inhibition by these drugs. However, improved susceptibility is seen with newer generation azoles such as voriconazole.

Life-threatening systemic infections caused by Candida and Aspergillus species are usually seen in severely ill and immunocompromised patients such as HIV, chemotherapy-induced neutropenia, and transplant recipients on immunosuppressants. C. albicans and C. dubliniensis are most susceptible to currently available azoles, C. glabrata is less susceptible, whereas C. krusei is intrinsically resistant to fluconazole. Resistance occurs due to overexpression or point mutation in the gene encoding the target enzyme lanosterol 14α-demethylase. Voriconazole is the preferred therapy for multi-drug resistant infection with C. krusei.

C. albicans and C. glabrata (Option C) showing resistance to Amphotericin B has been increasingly reported over the past several years. But this is a secondary resistance due to alteration or a decrease in the amount of ergosterol in the cell membrane. Amphotericin B is indicated to treat C. glabrata that has developed resistance to azoles.

Aspergillus fumigatus, the most common species to cause Aspergillus infection worldwide, also shows an intrinsic resistance to fluconazole. It is susceptible to echinocandins like micafungin, which acts by inhibiting beta-(1,3)-D-glucan synthase, an enzyme necessary for the synthesis of the cell wall.

Other options

Option A : Aspergillus niger is a common cause of otomycosis and is susceptible to azoles like voriconazole, amphotericin B, and echinocandins.

Option D : The drug of choice for treatment of sporotrichosis is itraconazole.

Solution to Question 28:

IV desferrioxamine (also known as deferoxamine) is indicated when serum iron levels are greater than 500 mcg/dL in cases of iron toxicity.

Desferrioxamine is selectively used as a chelating agent to manage cases of acute iron poisoning, hemochromatosis, and patients with iron overload from repetitive blood transfusions. Desferrioxamine can be administered intramuscularly, intravenously, or subcutaneously but not orally.

Deferiprone and deferasirox are both oral drugs and can be given in transfusion related siderosis but are less effective than desferrioxamine. Although, they can be preferred because of the ease of administration through oral route and multiple transfusions required for beta thalassemia patients.

Iron exhibits toxic effects by hydration of the ferric (F3+) ion which generates H+. Non transferrin bound iron catalyses free radical formation which causes mitochondrial injury, lipid peroxidation, increased capillary permeability, vasodilation, and organ toxicity.

Clinical features of iron toxicity include initial nausea, vomiting, abdominal pain and diarrhea. In severe cases there is high anion gap metabolic acidosis, cardiovascular and CNS depression, hepatitis, coagulopathy, and seizures.

In the case of toxic iron ingestion, radiopaque iron tablets may be seen on abdominal x-ray.

Management of iron toxicity includes the following :

• Whole-bowel irrigation for large ingestions, endoscopy and gastrostomy if clinical toxicity and large number of tablets are still visible on x-ray.

• IV hydration.

• Sodium bicarbonate for acidemia.

• IV desferrioxamine for systemic toxicity, iron level >90 μmol/L (500 μg/dL).

Solution to Question 29:

Methadone is the causative drug for this patient's presentation of syncope and prolonged QT interval after coadministration other QT prolonging drugs such as fluoxetine.

Unlike other opioids, QT prolongation is primarily a side effect of methadone, which is additive with other agents known to cause QT prolongation such as fluoxetine. Serious cardiac arrhythmias including torsades de pointes have been noted with methadone use. Normal QTc is about 440 ms. Values of more than 440 ms in males and 460 ms in females are considered to be prolonged. QT prolongation can present as syncope or cardiac arrest.

Methadone is a synthetic opioid that is chemically different but has similar pharmacological action to that of morphine (Option B). It is a long-acting mu-opioid receptor agonist. Although an effective analgesic, the primary use of methadone hydrochloride is detoxification and maintenance therapy treatment for opioid use disorder. Methadone is also used for the management of chronic pain.

Side effects of methadone are similar to that of morphine such as lethargy, constipation, respiratory depression, and hypotension. Like other opioids, methadone will produce tolerance and dependence. Development of physical dependence during the long-term administration of methadone can be demonstrated following abrupt drug withdrawal or administration of an opioid antagonist.

Other options :

Option B : Morphine is not associated with QT prolongation.

Option C : Meperidine, also known as pethidine, though chemically unrelated to morphine interacts with μ opioid receptors and its actions are blocked by naloxone. Apart from usual opioid side effects, pethidine produces some atropinic effects like dry mouth, blurred vision, tachycardia. Pethidine injected in patients receiving a selective serotonin reuptake inhibitor (SSRI) may produce serotonin syndrome. It is occasionally used to control shivering during recovery from anaesthesia.

Option D : Propoxyphene is a methadone analogue that was used to treat mild-tomoderate pain. Due to reports of cardiotoxicity, it was withdrawn in UK, Europe, and USA and it was banned in India in 2013.

Solution to Question 30:

Hyponatremia occurs more frequently in the case of oxcarbazepine than carbamazepine.

Both carbamazepine and oxcarbazepine are sodium-channel blockers that stabilize neuronal membranes. Oxcarbazepine is the 10-keto analog of carbamazepine and and cannot form an epoxide metabolite. Carbamazepine potentiates the action of antidiuretic hormone (ADH). It can result in water retention with decreased osmolarity and hyponatremia, especially in elderly patients with cardiac disease. Although hyponatremia may occur more commonly with oxcarbazepine than with carbamazepine, most adverse effects of oxcarbazepine are similar to those of carbamazepine.

Carbamazepine and oxcarbazepine can cause leukopenia, aplastic anemia, hepato-toxicity in the elderly.

Oxcarbazepine shows lesser hypersensitivity reaction. Hence, rashes (option A) are more commonly seen with carbamazepine than with oxcarbazepine.

Studies have shown that carbamazepine elevates enzyme gamma-glutamyl transpeptidases (GGT) more frequently than oxcarbazepine because of enzyme P450 induction. Hepatotoxicity from oxcarbazepine occurs less commonly compared to carbamazepine (Option D).

Side effects of carbamazepine therapy include: mnemonic: HEADS

Hypersensitivity reactions

Eosinophilia

Agranulocytosis: report symptoms like sore throat/fever

Aplastic anemia

Ataxia

Dilutional hyponatremia

Diplopia, blurred vision

Splenomegaly

Stevens-Johnson syndrome

Carbamazepine is most effective in complex partial seizures. It is also used in GTCS and simple partial seizures, trigeminal and related neuralgias, and manic depressive psychosis. Oxcarbazepine has been used to treat intractable trigeminal neuralgia refractory to other anticonvulsants.

Mechanism of action of some common antiepileptics:

Mechanism of action	Antiepileptics
Sodium-channel blockers	PhenytoinCarbamazepine, OxcarbamazepineEslicarbazepineLamotrigineTopiramateFelbamateValproateRufinamideLevetiracetam

Mechanism of action	Antiepileptics
Calcium-channel blockers	ValproateLamotrigineEthosuximide
GABA A receptor modulators	BenzodiazepinesPhenobarbitalFelbamatePrimidoneTopiramateCarbamazepine, OxcarbamazepineStiripentolClobazam
GABA uptake inhibitors	TiagabineVigabatrin
NMDA receptor antagonist	Felbamate
AMPA/Kainate receptor antagonist	PhenobarbitalTopiramatePerampanel
HCN channel activity enhancers	Lamotrigine
Inhibitors of brain carbonic anhydrase	AcetazolamideZonisamideTopiramate
Potassium-channel openers	RetigabineTopiramate

Solution to Question 31:

This patient was most likely taking tipranavir, as it is known to be associated with an increased risk of intracranial hemorrhage and hepatotoxicity which is the likely cause of altered mental status in this patient with hemophilia following complaints of nausea, vomiting and headache.

Tipranavir is a protease inhibitor (PI) used in patients who are resistant to other PIs. It is often combined with ritonavir, to achieve better therapeutic outcomes. It should be avoided in patients with head trauma or bleeding diathesis, such as hemophilia as it is associated with an increased risk of intracranial hemorrhage. Other common adverse effects of tipranavir include nausea, vomiting, diarrhea, abdominal pain, urticarial or maculopapular rash, and liver toxicity.

Other Options :

Option A, C and D : Lopinavir, Darunavir and Ritonavir are protease inhibitors that can cause nausea and vomiting, although they are not contraindicated in patients with bleeding diathesis and not associated with an increased risk of intracranial hemorrhage.

The illustration below shows the mechanism of action of various antiretroviral drugs.

Antiretroviral drug	Mechanism of action
Nucleotide/ nucleoside reverse transcriptase inhibitors	Inhibits reverse transcriptase
Non-nucleoside reverse transcriptase inhibitors	Inhibits reverse transcriptase by binding to an allosteric site of the enzyme

Antiretroviral drug	Mechanism of action
Protease inhibitors	Inhibits the viral protease en zyme necessary to produce m ature virions upon budding fr om the host membrane.
Entry inhibitors	Interferes with binding, fusio n, and entry of HIV -1 into th e cell by blocking a target.Ma raviroc works by targetting C CR5, a coreceptor located on human helper T-cells
Integrase inhibitors	Inhibits the viral enzyme inte grase, which is responsible fo r the integration of viral DNA into the DNA of the infected cell.



Solution to Question 32:

A chronic kidney disease (CKD) patient with chest pain, palpitations, muscle weakness, and serum potassium of 6.4mEq/L indicates hyperkalemia.

Hyperkalemia can be defined as potassium levels >5.5 mM. In normal patients, hyperkalemia is prevented by aldosterone-dependent secretion and potassium excretion via the gastrointestinal tract. But in patients with CKD, certain situations overcome these preventive mechanisms and cause hyperkalemia. These include high potassium intake, hemolysis, red blood cell transfusion, metabolic acidosis, and drugs that inhibit potassium excretion. The most common drugs that can precipitate hyperkalemia in CKD patients are renin-angiotensin system (RAS) inhibitors, spironolactone, and other potassium-sparing diuretics like triamterene, eplerenone, and amiloride.

ECG shows tall and peaked T waves, widening of QRS complexes, loss of P waves, and sine wave pattern.



Management of hyperkalemia involves an immediate reversal of cardiac effects by administration of 10mL of 10% calcium gluconate intravenously. This leads to the reversal of the depolarization blockade due to hyperkalemia. Redistributing the excess potassium by administering 10U of intravenous insulin followed by 50mL of 50% dextrose can also be given, as insulin facilitates the intracellular shift of potassium. Beta-2-agonists can also be used in the management of hyperkalemia. But patients with end-stage renal disease are resistant to beta-2-agonists and hence must be used with caution. Removal of potassium can be done using cation exchange resins like patiromer and sodium polystyrene sulfonate, diuretics, and dialysis.

Solution to Question 33:

Pregabalin can be used for diabetic neuropathy, post herpetic neuralgia and spinal cord injury. Although amitriptyline can also be used for these conditions, pregabalin is found to be recommended over amitriptyline considering safety and efficacy and hence is the best option.

Pregabalin is an anti-epileptic drug. Its analgesic effect is mediated by binding to α2δ, a protein that is a subunit of voltage-gated calcium channels and causes a decrease in glutamate release at excitatory synapses. Decreased neurotransmitter release is associated with reduced neuronal hyperexcitability. It is used in the management of neuropathic pain associated with diabetic peripheral neuropathy, spinal cord injury or postherpetic neuralgia.

Currently, only duloxetine and pregabalin are approved by the U.S. Food and Drug Administration and the European Medicines Agency to treat painful diabetic neuropathy. Diabetic Neuropathy is the most common cause of peripheral neuropathy in developed countries. Risk factors include long-standing, poorly controlled DM and the presence of retinopathy and nephropathy.

Side effects of pregabalin include sedation and peripheral edema.

Other options :

Option A : Tricyclic antidepressants (TCAs) such as amitriptyline are used in the treatment of several pain conditions such as diabetic neuropathy, post herpetic neuralgia and following spinal cord injury. Their mechanism of action is inhibition of norepinephrine and serotonin reuptake. Although both pregabalin and amitriptyline are among the first line drugs for neuropathic pain, amitriptyline would be the next best option in this question, owing to its side effect profile and good yet controversial efficacy in pain control. It is a non FDA approved drug for diabetic neuropathy (off label indication). Side effects associated with amitriptyline include anticholinergic side effects (dry mouth, tachycardia, hyperthermia), cardiac effects (arrhythmias) and sedation.

Option B : Phenytoin is an antiepileptic drug that is not used in the treatment of pain.

Option C : Carbamazepine is an iminostilbine derivative. It blocks voltage-activated sodium channels, slowing their rate of recovery and limiting the repetitive firing of action potentials. It is used in the treatment of generalized, focal-to-bilateral and idiopathic tonic-clonic seizures, focal seizures, neuropathic pain associated with trigeminal and glossopharyngeal neuralgias.

Solution to Question 34:

Linezolid is a protein synthesis inhibitor that acts by inhibiting the 50s ribosomal subunit.

Linezolid binds to the P site of the 50s ribosomal subunit and inhibits bacterial protein synthesis. It is an oxazolidinone that is active against various resistant organisms including MRSA. It has excellent oral bioavailability. Linezolid is used to treat infections caused by gram-positive bacteria, particularly vancomycin-resistant enterococcus and methicillin-resistant Staphylococcus aureus. Linezolid is also approved for the treatment of community-acquired pneumonia due to S. pneumoniae and nosocomial pneumonia due to S. aureus.

The intrinsic resistance of gram-negative bacteria to linezolid is mediated by endogenous efflux pumps.

Linezolid is associated with the side effect of myelosuppression (anemia, leukopenia, pancytopenia, and thrombocytopenia). Thrombocytopenia is the most common side effect. Linezolid has MAOI (monoamine oxidase inhibitors) activity and therefore can precipitate serotonin syndrome with concomitant use of SSRI (paroxetine) or with other serotonergic medications.

Other adverse effects include peripheral neuropathy, optic neuropathy, lactic acidosis.

Other options :

Option A : Aztreonam is a monobactam, and acts similar to penicillin by inhibiting peptiglycan crosslinking and inhibiting bacterial cell wall synthesis.

Option B : Daptomycin is a cyclic lipopeptide antibiotic. It acts by depolarising bacterial membranes resulting in loss of membrane potential and cell death.

Option D : Rifampicin is an anti tubercular antibiotic that acts by inhibiting bacterial DNA-dependent RNA polymerase.


Mechanism of action of antibiotics
PABA
DHF
THF
DNA
RNA polymerase
mRNA
DNA gyrase
Ribosomes
50S
30S
Cell membrane
Cell wall
1 Sulphonamides: Dihydropteroate synthase inhibitors.
2 Trimethoprim: Selective microbial dihydrofolate reductase inhibitor.
3 Cell wall inhibitors: Peptidoglycan synthesis inhibitors: Vancomycin & Bacitracin. Peptidoglycan cross linking inhibitors: Penicillins, Cephalosporins, Carbapenems and Monobactams.
8 Metronidazole: Disrupts DNA integrity using free radicals.
7 RNA polymerase inhibitors: Rifampin inhibits mRNA synthesis.
6 DNA gyrase inhibitors: Fluoroquinolones.
5 50s subunit inhibitors: Macrolides, Clindamycin, Linezolid and Chloramphenicol.
4 30s subunit inhibitors: Aminoglycosides and Tetracyclines.
Bacterial cell
©MARROW

Pharmacology INI-CET 2024

Question 1:

Which of the following drugs used in the treatment of diabetes is known to cause urinary tract infections?

a) Dapagliflozin

b) Sitagliptin

c) Pioglitazone

d) Glimeperide

Question 2:

Which of the following medications used in treating leprosy causes skin pigmentation?

a) Oflaxacin

b) Dapsone

c) Clofazamine

d) Rifampicin

Question 3:

A patient is set to receive intermittent treatment with teriparatide. Which of the following complications does this drug predispose him to?

a) Osteonecrosis of the jaw bone

b) Subtrochanteric fractures

c) Net loss of bone mass

d) Osteosarcoma

Question 4:

Which of the following monoclonal antibodies is not a HER - 1 inhibitor?

a) Pertuzumab

b) Cetuximab

c) Panitumumab

d) Necitumumab

Question 5:

A G2P1 delivered a preterm baby at 36 weeks. The baby was admitted in the NICU for a week and was given expressed breast milk. Unfortunately the baby did not survive. Which of the following drugs can be used to reduce the engorgement and pain of the lactating breasts?

a) Cabergoline

b) Sumatriptan

c) Chlorpromazine

d) Pramipexole

Question 6:

Coagulation profile monitoring is required for which of the following drugs?

a) Lepirudin

b) Enoxaparin

c) Fondaparinux

d) Dabigatran

Question 7:

A patient known case of diabetes, shows microalbuminuria on urine analysis, done twice at an interval of 6 months. Renal function tests are normal. Which of the following drugs is not useful in preventing the progression of renal damage?

a) Finerenone

b) Hydrochlorothiazide

c) Angiotensin receptor blockers

d) SGLT2 inhibitor

Question 8:

Which of the following causes a decrease in serum parathyroid hormone?

a) Calcitriol

b) Risedronate

c) Furosemide

d) Lithium

Question 9:

Which of the following acts by suicide inhibition?

a) 1,2

b) 3,4

c) 1,2,4

d) 1,3,4

Question 10:

Which of the following diuretics is matched correctly with its location of action?

a) 1,2,3

b) 3,4

c) 2, 4

d) 1,2

Question 11:

Which of the following drugs acts as an inhibitor of PCSK-9?

a) Bempedoic acid

b) Lomitapide

c) Ezetemibe

d) Evolocumab

Question 12:

Inhaled nitric oxide (iNO) is used for?

a) Persistent pulmonary hypertension of newborn

b) Unstable angina

c) Obstructive sleep apnea

d) Erectile dysfunction

Question 13:

A patient presents with complaints of a cough producing tenacious white sputum. Which of the following statements is true about the drug that could be prescribed for this condition?

a) Codeine as it is a potent drug and quickly relieves symptoms.

b) Bromhexine and Dextromethorphan syrup as they liquify sputum and reduce cough due to antitussive effect

c) Dextromethorphan alone as it does not have addictive potential like codeine

d) Bromhexine as it liquifies sputum

Question 14:

Which of the following antidiabetic drugs is known to cause unexplained diarrhea?

a) Sitagliptin

b) Metformin

c) Glimepiride

d) Pioglitazone

Question 15:

A young female weighing 46 kg is on digoxin with vd-6 L/kg and clearance 4.6 L/h. Calculate the half-life of digoxin.

a) 4h

b) 68h

c) 42h

d) 0.9h

Question 16:

Choose the correct option regarding phenytoin.

a) Acute oral overdose results primarily in cerebellar and vestibular symptoms and high doses have been associated with marked cerebellar atrophy

b) High dose causes cardiotoxicity in old and nephrotoxicity in young patients

c) These complications can be minimized by administering fosphenytoin at a rate of less than 250 mg of phenytoin sodium equivalents per minute

d) Given in all patients with seizures regardless of cardiac arrhythmias, hypotension and/or CNS depression

Question 17:

True statement about Iron Isomaltoside 1000 is

a) Iron requirement can be fulfilled in single dose

b) More immunogenic than iron dextran

c) Given intravenously and gets deposited in adipose tissue therefore less toxic

d) Cannot be given with erythropoietin

Question 18:

The drug that binds to CD4 and has action on both CCR5 and CXCR4 in tropic HIV infections is

a) Maraviroc

b) Ibalizumab

c) Elvitregravir

d) Fosamprenavir

Question 19:

Inhibitors of Poly ADP ribose polymerase (PARP) are approved for BRCA-positive breast cancers. Which among the following drugs is used?

a) Pablociclib

b) Rucaparib

c) Regorafenib

d) Vandetinib

Question 20:

A 60-year-old patient who is a known case of hypertension and diabetes underwent PCI stent placement for NSTEMI last year. He is currently on aspirin, metoprolol and atorvastatin. He comes for a follow-up, and his lipid profile is as follows- Total cholesterol 252mg/dl LDL: 156mg/dl, triglyceride 158mg/dl and HDL- 40 mg/dl. Which drugs can be added based on his lipid profile?

a) 1, 2 and 4

b) Only 1 and 2

c) 1,3 and 4

d) Only 1 and 3

Question 21:

Which of the following is a true statement regarding Pretomanid?

a) Given in XDR TB with Bedaquiline and Linezolid

b) Contraindicated in patients on enzyme inducer drugs

c) Acts by reactive oxygen species

d) For resistant cases of malaria

Question 22:

Which of the following drugs act on sodium channels and potentiate slow sodium inactivation?

a) Phenytoin

b) Lamotrigine

c) Lacosamide

d) Topiramate

Question 23:

Which drug acts on the marked receptor?



a) Tiagabine

b) Ganaxolone

c) Felbamate

d) Stiripentol

Question 24:

Match the treatment with its respective condition.

a) a-i, b-ii, c-iii, d-iv

b) a-ii, b-iii, c-i, d-iv

c) a-iii, b-i, c-iv, d-ii

d) a-iv, b-ii, c-iii, d-i

Answer Key

Question No.	Correct Option
1	a
2	c
3	d
4	a
5	a

6	a
7	b
8	a
9	b
10	b
11	d
12	a
13	d
14	b
15	c
16	a
17	a
18	b
19	b
20	a
21	b
22	c
23	a
24	b

Detailed Explanations

Solution to Question 1:

Dapagliflozin is known to cause urinary tract infections.

Dapagliflozin is a sodium-glucose transporter 2 (SGLT-2) inhibitor. It acts by inhibiting the SGLT-2 transporter which is present on the proximal segment of the renal tubule and is responsible for the reabsorption of 80-90% of filtered glucose. This results in glycosuria. Other SGLT-2 inhibitors include canagliflozin and empagliflozin.

Apart from their effect on lowering blood glucose levels, SGLT2 inhibitors can cause mild weight loss and reduction in blood pressure. They have also been shown to reduce the risk of hospitalization in patients with heart failure.

The common side effects include urinary tract infections and genital mycotic infections due to the high glucose content of the urine. There is a risk of the development of diabetic ketoacidosis.

Other options :

Option B - Sitagliptin is a dipeptidyl peptidase-4 inhibitor. Its use is associated with an increased risk of heart failure.

Option C - Pioglitazone is an activator of the peroxisome proliferator-activated receptor γ (PPARγ), which is involved in the regulation of genes related to glucose and lipid metabolism. Its adverse effects include increased incidence of heart failure, peripheral edema, weight gain, macular edema, and osteoporosis.

Option D - Glimepiride belongs to the class of sulfonylureas. It stimulates insulin release by inhibiting the activity of the β cell KATP channel complex. Its side effects include hypoglycemia and weight gain.

Solution to Question 2:

Clofazimine is a dye with leprostatic and antiinflammatory properties.The mechanism of antimicrobial activity of clofazimine against mycobacteria appears to be multifactorial and includes membrane disruption, inhibition of mycobacterial phospholipase A2, inhibition of microbial K+ transport, generation of hydrogen peroxide, interference with the bacterial electron transport chain, and efflux pump inhibition.

Clofazimine is orally active. It accumulates in macrophages and gets deposited as needle-shaped crystals in many tissues including subcutaneous fat.

Clofazimine is used as a component of multidrug therapy (MDT) for leprosy. Because of its anti-inflammatory properties, it is valuable in lepra reactions. Occasionally, it is used as a component of a combination regimen for extensive drug resistant TB (XDR-TB).

Adverse effects -

• Skin - The major disadvantage is reddish-black discoloration of skin, especially on exposed parts. Discoloration of hair and body secretions may also occur. Dryness of the skin, itching and scaling are often troublesome. Acneform eruptions and phototoxicity have been noted. Conjunctival pigmentation may create cosmetic problems.

• GI symptoms - Nausea, anorexia, abdominal pain, weight loss and enteritis with intermittent loose stools can occur, particularly when higher doses are used to control lepra reaction. Deposition of clofazimine crystals in the intestinal submucosa may produce gastrointestinal symptoms.

Clofazimine is to be avoided during early pregnancy and in patients with liver or kidney damage.

Other options :

Option A- Ofloxacin is a fluroquinolone. As a component of MDT, it has been found to hasten the bacteriological and clinical response, where it is cidal to M.leprae. Its side effects include nausea, vomiting, and abdominal discomfort. CNS side effects like dizziness, anxiety and insomnia are also seen. Tendinitis is another adverse effect seen with the use of fluoroquinolones.

Option B- Dapsone exerts its antimicrobial effect by the inhibition of PABA incorporation into folic acid by folate synthase. Apart from being used in leprosy, dapsone is also used for the prophylaxis of T. gondii. The anti-inflammatory effects are the basis for therapy of dermatitis herpetiformis, linear immunoglobulin A bullous disease, and relapsing chondritis. Mild hemolytic anemia is common, especially in those with G-6-PD deficiency. Isolated instances of reversible peripheral neuropathy, drug fever, hematuria, pruritus, psychosis, and a variety of skin rashes have been reported.

Option D - Rifampicin interrupts RNA synthesis by binding to a subunit of mycobacterial DNA-dependent RNA polymerase. Apart from being used to treat leprosy, it is used in the management of tuberculosis, brucellosis, and prophylaxis of Meningococcal meningitis. Hepatitis is a major adverse effect. Other side effects include pruritus with rash, redness and watering of eyes,

flu-like symptoms and abdominal cramps. Urine and secretions may become orange-red.

Solution to Question 3:

The use of teriparatide predisposes an individual to the development of osteosarcoma.

Teriparatide is a parathormone analog, which is given as subcutaneous injections. Pharmacokinetics and systemic actions of teriparatide on mineral metabolism are the same as for PTH. Its main target of action are the osteoblasts.

Teriparatide is administered by once-daily subcutaneous injection of 20 μg into the thigh or abdomen. The site of administration should be rotated, but the time at which the injection is made should be the same each day.

Teriparatide and abaloparatide (PTHrP analog) are the only anabolic agents currently available that increase new bone formation (option C). Candidates for treatment include women who have a history of osteoporotic fracture, who have multiple risk factors for fracture, or who failed or are intolerant of previous osteoporosis therapy. Men with primary or hypogonadal osteoporosis are also candidates for treatment with these agents.

Adverse effects include hypercalcemia, exacerbation of nephrolithiasis, and elevation of serum uric acid levels. The development of osteosarcoma has been a serious concern in patients treated with teriparatide. It comes with a black-box warning, and use should be limited to no more than 2 years and avoided in patients who are at increased baseline risk for osteosarcoma (including those with Paget disease of bone, unexplained elevations of alkaline phosphatase, open epiphyses, or prior radiation therapy involving the skeleton).

Other options :

Options A and B - Osteonecrosis of the jaw bone and subtrochanteric fractures are the adverse effects of bisphosphonates.

Solution to Question 4:

Pertuzumab is a humanized IgG1 monoclonal antibody directed against HER2.

Pertuzumab is approved for the treatment of patients with HER2-positive, locally advanced, inflammatory, or early-stage breast cancer in combination with trastuzumab and docetaxel. Its adverse effect includes cardiotoxicity.

The other three antibodies act against epidermal growth factor receptor (EGFR)/ HER-1.

Option B - Cetuximab is used to treat patients with metastatic colon cancers and head and neck squamous cell carcinoma (HNSCC).

Option C - Panitumumab is approved for the treatment of EGFR-expressing metastatic colorectal cancer.

Option D - Necitumumab is approved for first-line treatment of patients with metastatic, squamous non-small cell lung cancer in combination with gemcitabine and cisplatin.

Solution to Question 5:

In the given scenario, the woman requires suppression of lactation, for which cabergoline is indicated.

In the postpartum period, prolactin promotes lactation. Dopamine(DA) is the primary regulator of prolactin secretion from the pituitary gland. DA released from the hypothalamus into the hypophyseal portal blood supply acts on lactotroph D2 receptors to decrease prolactin secretion. The DA agonists like bromocriptine and cabergoline are used in conditions requiring suppression of prolactin.

Cabergoline is an ergot derivative with a longer t1/2 (~65 h), higher affinity, and greater selectivity for the DA D2 receptor compared to bromocriptine. Cabergoline is the preferred drug for the treatment of hyperprolactinemia because of greater efficacy and lower adverse effects. Therapy is initiated at a dose of 0.25 mg twice a week or 0.5 mg once a week. Cabergoline induces remission in a significant number of patients with prolactinomas. At higher doses, cabergoline is used in some patients with acromegaly alone or in conjunction with somatostatin analogs.

Adverse Effects -

Compared to bromocriptine, cabergoline has a much lower tendency to induce nausea, although it still may cause hypotension and dizziness. Cabergoline has been linked to valvular heart disease, an effect proposed to reflect agonist activity at the serotonin 5HT2B receptor. However, this effect is seen primarily at the high doses used in patients being treated for Parkinson's disease and is not seen in the conventionally used doses (≤2 mg/week) for patients with prolactinomas.

Other options:

Option B - Sumatriptan is a 5HT1B/1D receptor agonist used in the treatment of acute attacks of migraine.

Option C - Chlorpromazine is a typical antipsychotic, which has D2 antagonistic activity. It is used in the treatment of schizophrenia, acute mania, and refractory nausea and vomiting. Hyperprolactinemia is an adverse effect associated with its use.

Option D - Pramipexole is a non-ergot alkaloid with a broader DA receptor agonist activity, with the highest affinity to D3 receptors. It is used in the treatment of Parkinson's disease and restless leg syndrome. It is not indicated for the treatment of breast engorgement.

Solution to Question 6:

Coagulation profile monitoring is required while using lepirudin.

Lepirudin is a parenteral direct thrombin inhibitor, which is a recombinant form of hirudin extracted from the saliva of leech. Its action is independent of anti-thrombin, which means it can reach and inactivate fibrin-bound thrombin in thrombi. It is excreted by the kidney and should be used with great caution in patients with renal insufficiency as no antidote exists.

Like heparin, lepirudin must be administered parenterally and is monitored by aPTT. Lepirudin was approved for use in patients with thrombosis related to heparin-induced thrombocytopenia (HIT). Its production was discontinued by the manufacturer in 2012.

Other options :

Option B - Enoxaparin is a low molecular weight heparin (LMWH) that inhibits activated factor X with less effect on thrombin. Hence, it has a smaller effect on the coagulation profile and does not need monitoring.

Option C - Fondaparinux avidly binds antithrombin, resulting in efficient inactivation of factor Xa. It is effective in the prevention and treatment of venous thromboembolism and does not appear to cross-react with pathologic HIT antibodies in most individuals. Like LMWH, it does not require monitoring of the coagulation profile.

Option D - Dabigatran is an oral direct thrombin inhibitor approved for the reduction in risk of stroke and systemic embolism with nonvalvular atrial fibrillation, treatment of venous thromboembolism (VTE) and VTE prophylaxis following hip or knee replacement surgery. It does not require monitoring of coagulation profile.

Solution to Question 7:

The given clinical scenario suggests chronic kidney disease secondary to diabetes, and hydrochlorothiazide is not useful in preventing the progression of renal damage.

Chronic kidney disease (CKD) is defined as abnormalities of kidney structure or function, present for a minimum of 3 months, with implications for health. CKD is classified based on cause, glomerular filtration rate (GFR) category (G1–G5), and albuminuria category (A1–A3). Some of the leading causes of CKD are diabetes, hypertension, glomerulonephritis and polycystic kidney disease.

In patients with diabetes mellitus, often there exists a state of glomerular hyperfiltration due to decreased vascular resistance of afferent arterioles and increased vascular resistance of efferent arterioles resulting in increased glomerular hydrostatic pressure. This contributes to the progression of the CKD.

The following drugs have been approved in the management of CKD :

• Angiotensin-converting enzyme inhibitors / Angiotensin receptor blockers - These medications appear to slow the rate of decline of kidney function in a manner that extends beyond the reduction of systemic arterial pressure and that involves a reduction in the intraglomerular hyperfiltration and hypertension.

• Finerenone - It is a non steroidal mineralocorticoid receptor antagonist which decreases proteinuria. It has been approved to reduce the risk of end-stage kidney disease, cardiovascular death, and heart failure in diabetic patients with CKD.

• SGLT-2 inhibitors - These drugs inhibit sodium-glucose cotransporter type 2 in the proximal convoluted tubule. By increasing NaCl delivery to the thick ascending limb, gliflozins trigger tubuloglomerular feedback and thereby increase preglomerular vascular resistance and decrease glomerular hypertension. This contributes to the effectiveness of gliflozins to prevent diabetic kidney disease.

Solution to Question 8:

The active form of vitamin D, 1,25-dihydroxyvitamin D (calcitriol), directly suppresses PTH gene expression

Calcitriol is the active form of vitamin D. It increases calcium absorption from the intestines and helps maintain adequate serum calcium levels. When calcium levels rise, this triggers a negative feedback mechanism on the parathyroid glands, leading to decreased secretion of parathyroid hormone (PTH). PTH's primary role is to raise blood calcium levels, so when calcium is sufficient, the need for PTH decreases, resulting in lower PTH levels.

Vitamin D and its derivatives are the primary treatment for hypoparathyroidism. Dihydrotachysterol (DHT), a reduced form of vitamin D2, has a quicker onset, shorter duration, and stronger impact on bone mobilization than traditional vitamin D, making it a preferred option. While most hypoparathyroid patients respond well to any form of vitamin D, calcitriol may be preferred for temporary treatment of hypocalcemia while waiting for the effects of a slower-acting vitamin D form. Calcium is used to address calcium deficiency and as a dietary supplement. Vitamin D is the primary treatment for hypoparathyroidism.

Severe hypermagnesemia or hypomagnesemia can also inhibit PTH secretion.

Other options:

Option B: Risedronate is a bisphosphonate that inhibits bone resorption, but it doesn't directly affect PTH secretion.

Option C: Furosemide is a diuretic that can increase calcium excretion, potentially raising PTH as the body tries to conserve calcium.

Option D: Lithium can actually increase PTH secretion by affecting calcium-sensing receptors in the parathyroid glands.

Solution to Question 9:

Ritonavir and benzbromarone are both suicide inhibitors.

Suicide inhibition is when an inhibitor binds to an enzyme and then undergoes a reaction to form an irreversible complex with the enzyme, inactivating it permanently. Ritonavir is a protease inhibitor used in HIV treatment and benzbromarone is a uricosuric agent used to treat gout by increasing uric acid excretion. It also inhibits CYP2C9 via mechanism-based (suicide) inhibition.

Other options:

Ethinyl estradiol is a synthetic estrogen used in oral contraceptives and hormone replacement therapy. It acts through receptor-mediated mechanisms, not through suicide inhibition.

Spironolactone is a potassium-sparing diuretic and aldosterone antagonist. It works by competitively inhibiting aldosterone receptors in the distal tubules of the kidney, not through suicide inhibition.

Solution to Question 10:

The osmotic diuretics act at the proximal convoluted tubule and the potassium-sparing diuretics act on the collecting tubule. Hence C and D are the right statements.

A - Osmotic diuretics, like mannitol, retain water isoosmotically in the proximal convoluted tubule (PCT) and descending loop of Henle which are freely permeable to water. Luminal fluid is diluted and opposes NaCl reabsorption. They are indicated in raised intracranial tension and in dialysis disequilibrium.

B - Thiazides act on the distal convoluted tubule (DCT). They block the Na+/Cl- co-transporter in the DCT. This results in the loss of sodium, chloride, and water contributing to the diuretic effect. Some of the examples are hydrochlorothiazide, chlorthalidone and metolazone. Their indications include hypertension, edema, diabetes insipidus and hypercalciuria.

C - Carbonic anhydrase inhibitors act on the proximal convoluted tubule (PCT). By inhibiting carbonic anhydrase, HCO_3^- reabsorption is blocked and diuresis occurs. Examples are acetazolamide, methazolamide, dichlorphenamide, and dorzolamide. They are used in the treatment of glaucoma, urinary alkalinization in cystinuria, management of metabolic alkalosis, and acute mountain sickness.

D - Potassium-sparing diuretics include aldosterone antagonists (spironolactone, eplerenone) and inhibitors of renal epithelial Na+ channel (triamterene, amiloride). They act on the late distal tubule and the collecting duct. Aldosterone antagonists are used in the treatment of edema, hypertension, heart failure and Conn's syndrome. Amiloride has a role in lithium-induced diabetes insipidus.

Solution to Question 11:

Evolocumab is a monoclonal antibody that acts by inhibiting PCSK9.

PCSK9 (proprotein convertase subtilisin/kexin type) is a serine protease that causes lysosomal degradation of the low-density lipoprotein (LDL) receptors present on hepatocytes. Monoclonal antibodies against PCSK9 such as alirocumab and evolocumab counter this and cause persistence of LDL receptors on hepatocytes. This leads to more LDL uptake from the circulation, bringing down the LDL-cholesterol levels.

Evolocumab is administered as a 140-mg injection every 2 weeks or as a 420-mg injection once a month. PCKS9 inhibitors should not be used in pregnancy because transmission across the placenta is expected.

PCSK9 inhibitors are indicated only as adjunctive therapy in patients with familial hypercholesterolemia(FH) or established atherosclerotic cardiovascular disease(ASCVD) who have elevated LDL-cholesterol levels despite being on maximum statin therapy.

Other options :

Option A - Bempedoic acid acts by inhibiting de novo hepatocyte cholesterol biosynthesis, by blocking the enzyme adenosine triphosphate citrate lyase. The drug is approved as an adjunct to diet and maximally tolerated statin therapy to further lower LDL-C in patients with FH or established ASCVD.

Option B - Lomitapide acts by inhibiting microsomal triglyceride transfer protein (MTP), which is essential for the intracellular transfer of triglycerides into triglyceride-rich lipoproteins, thus inhibiting the formation of VLDLs in hepatocytes and chylomicrons in intestinal epithelial cells. It is approved as an adjunct to diet for lowering LDL-C levels, total cholesterol, apo B, and non–HDL lipoproteins in patients with FH.

Option C - Ezetimibe inhibits luminal cholesterol uptake by jejunal enterocytes, by inhibiting Niemann-Pick C1–like 1 transport protein. The role of ezetimibe as monotherapy for patients with elevated LDL-C levels is generally limited to a small group of statin-intolerant patients.

Solution to Question 12:

Inhaled nitric oxide(iNO) is used for persistent pulmonary hypertension in newborns. (Option A)

Inhaled NO selectively dilates the pulmonary vasculature, hence used in diseases associated with increased pulmonary vascular resistance. Inhaled NO is an FDA-approved drug for persistent pulmonary hypertension in the newborn.

Inhaled NO can also be used during cardiac catheterization to evaluate the pulmonary vasodilating capacity of patients with heart failure and infants with congenital heart disease. Another use of inhaled NO is to determine the diffusion capacity (DL) across the alveolar-capillary membrane. It is more effective than CO_2 because of its greater affinity for hemoglobin and its higher water solubility at body temperature.

NO is produced from the nasal passages and the lungs of normal human subjects and can be detected in exhaled gas. The measurement of fractional exhaled NO is a noninvasive marker for airway inflammation in the assessment of respiratory tract diseases, including asthma, respiratory tract infection, and chronic lung disease.

Inhaled NO is administered at low concentrations (0.1–50 ppm) since high concentrations of 50 to 100 ppm can cause significant methemoglobinemia that can cause death.

Other options:

Option B: Unstable angina is treated by anti-platelet drugs such as aspirin and clopidogrel, and anticoagulants such as heparin.

Option C: Obstructive sleep apnoea is mainly treated by using a BiPAP machine during sleep and CNS stimulants such as armodafinil for daytime sleepiness associated with the condition.

Option D: Erectile dysfunction is treated by PDE5 inhibitors such as sildenafil, tadalafil, etc.

Solution to Question 13:

Bromhexine should be prescribed for a patient with a productive cough, as it liquifies the sputum.

Bromhexine belongs to a class of drugs called mucolytics, capable of inducing thin copious bronchial secretion. It depolymerizes mucopolysaccharides directly as well as by liberating lysosomal enzymes- the network of fibers in tenacious sputum is broken. It is particularly useful if mucus plugs are present.

Other examples of mucolytics are ambroxol, carbocisteine, and N-acetyl cysteine.

Other options :

Codeine and dextromethorphan belong to the class of drugs called antitussives. These are drugs that act in the CNS to raise the threshold of the cough center or act peripherally in the respiratory tract to reduce tussal impulses, or both these actions. Because they aim to control rather than eliminate cough, antitussive drugs should be used only for dry nonproductive cough or if it is unduly tiring, disturbs sleep, or is hazardous (hernia, piles, cardiac disease, ocular surgery).

Cough is a defensive reflex and its suppression may be inappropriate in bacterial lung infection. Hence, before treatment with antitussives, it is important to identify underlying causal mechanisms that may require therapy.

Solution to Question 14:

Metformin is known to cause unexplained diarrhoea as its adverse effect.

Metformin is a member of the biguanide class of oral hypoglycemic drugs. It acts by primarily by inhibiting hepatic gluconeogenesis. It activates AMP-dependent protein kinase, leading to stimulation of hepatic fatty acid oxidation, glucose uptake, and nonoxidative glucose metabolism and reduction of lipogenesis and gluconeogenesis. It also appears to act in the gut.

Metformin is accepted as the first-line treatment of type 2 diabetes and is the most commonly used oral agent for this condition. It has been used as a treatment of infertility in women with polycystic ovarian syndrome.

The most common side effects of metformin are gastrointestinal: nausea, indigestion, abdominal cramps, bloating, and diarrhea. The use of metformin is also associated with lower blood levels of vitamin B12.

Most adverse GI effects of metformin abate over time and can be minimized by starting at a low dose and gradually titrating to a target dose over several weeks, as well as by having patients take the drug with meals. Extended-release metformin has decreased GI side effects and can be substituted for immediate-release metformin. Lactic acidosis has been linked with the use of metformin.

Other options:

Option A - Sitagliptin is a dipeptidyl peptidase-4 inhibitor. Its use is associated with an increased risk of heart failure.

Option C - Glimepiride belongs to the class of sulfonylureas. It stimulates insulin release by inhibiting the activity of the β cell KATP channel complex. Its side effects include hypoglycemia, and weight gain.

Option D - Pioglitazone is an activator of the peroxisome proliferator-activated receptor γ (PPARγ), which is involved in the regulation of genes related to glucose and lipid metabolism. Its adverse effects include increased incidence of heart failure, peripheral edema, weight gain, macular edema, and osteoporosis.

Solution to Question 15:

The half-life of Digoxin is 42 hours.

In first-order kinetics:

$$\text{Plasma half-life (t1/2)} = 0.693 \times \frac{\text{Volume of distribution } (V_d)}{\text{Clearance (CL)}}$$

Therefore,

t1/2 ~ 0.7 x 46x6/4.6 = 0.7 x 60

t1/2 ~ 42 hours

Where,

t1/2 - Plasma half-life

Vd - Volume of distribution

CL - Clearance

Solution to Question 16:

Acute oral overdose results primarily in cerebellar and vestibular symptoms and high doses have been associated with marked cerebellar atrophy.

Dose-related toxicity of phenytoin causes ataxia, vertigo, diplopia, and nystagmus.

Other options:

Option B: High dose causes cardiotoxicity in both old and young, healthy patients. However, cardiotoxicity is more pronounced and frequent in older patients and those with known cardiac diseases.

Option C: These complications can be minimised by administering fosphenytoin at a rate of less than 150 mg of phenytoin sodium equivalents per minute, not 250 mg.

Option D: Phenytoin is not effective in generalised absent seizures.

Solution to Question 17:

Iron requirement can be fulfilled in a single dose is true regarding Isomaltoside 1000.

Iron isomaltose 1000 is a new IV iron formulation that can supply a complete replacement dose in a short, single visit. It is used in adult patients with an intolerance to oral iron or for patients with CKD not requiring hemodialysis.

Other options:

Option B: Isomaltoside 1000 is not more immunogenic than iron dextran.

Option C: The strength of iron binding causes a controlled slow release of bioavailable iron, decreasing the risk of toxicity from labile-free iron. Its decreased toxicity cannot be attributed to deposition in adipose tissue.

Option D: Isomaltoside 1000 can be given with erythropoietin.

Solution to Question 18:

The drug that binds to CD4 and has an action on both CCR5 and CXCR4 in tropic HIV infections is Ibalizumab.

Ibalizumab is an anti-CD4 monoclonal antibody approved for use in the treatment of multidrug-resistant HIV-1. It has action on both CCR5 and CXCR4 in tropic HIV infections and is administered intravenously.

It acts as an entry inhibitor. It blocks HIV-1 from infecting CD4+ T cells by binding to domain 2 of CD4 cells. It also interferes with post-attachment steps required for the entry of HIV-1 virus particles into host cells and thus prevents the viral transmission that occurs via cell-cell fusion.

Other options:

Option A: Maraviroc binds to the host cell CCR5 receptor to block the binding of viral gp120. It prevents the entry of the viral particles. It does not bind to CD4.

Option C: Elvitegravir is an integrase inhibitor that inhibits HIV 1 viral integrase enzyme and prevents proviral DNA integration into host cells.

Option D: Fosamprenavir is a protease inhibitor that blocks the viral protease enzyme necessary to produce mature virions upon budding from the host membrane.

Solution to Question 19:

Rucaparib is an inhibitor of Poly ADP ribose polymerase (PARP) used for BRCA-positive breast cancers.

Rucaparib is a poly ADP-ribose polymerase/ PARP inhibitor approved for the treatment of breast and advanced ovarian cancer.

PARP is a nuclear protein that transfers ADP-ribose from NAD+ to target proteins and this plays an important role in the DNA damage response. Rucaparib and olaparib are PARP inhibitors.

Inhibitors of PARP (Poly ADP Ribosyl Polymerase) approved for breast cancer:

• Olaparib

• Rucaparib

• Talazoparib

Other options:

Option A: Palbociclib is a selective inhibitor of the cyclin-dependent kinases CDK4 and CDK6. It is used in the treatment of metastatic breast cancer.

Option C: Regorafenib is an angiogenesis inhibitor that is indicated as a second-line treatment for unresectable hepatocellular carcinoma. (The first line is Sorafenib).

Option D: Vandetanib is an oral tyrosine kinase inhibitor recently FDA-approved for unresectable locally advanced or metastatic medullary thyroid carcinoma.

Solution to Question 20:

1,2 and 4 can be added based on his lipid profile.

A patient with hypertension and diabetes who underwent PCI stent placement for NSTEMI has atherosclerotic cardiovascular disease (ASCVD). The treatment goal in patients with ASCVD is to decrease LDL levels.

Ezetemibe, Evolocumab, Bemepedoic acid decrease LDL. Gemfibrozil mainly decreases triglycerides and not LDL.

1. Evolocumab is a monoclonal antibody that inhibits PCSK-9 (proprotein convertase subtilisin/kexin). Inhibition of PCSK-9 prevents the degradation of LDL receptors and thus increases LDL-cholesterol clearance by the liver. They are used in patients with atherosclerotic cardiovascular disease (ASCVD) to decrease LDL levels.

2. Ezetimibe inhibits cholesterol absorption by enterocytes in the small intestine. It lowers the total and LDL-C levels.

3. Gemfibrozil belongs to the fibrate group of drugs that binds to peroxisome proliferator-activated receptors (PPARα), which stimulate fatty acid oxidation and thereby reduce triglyceride levels mainly.

4. Bempedoic acid inhibits adenosine triphosphate-citrate lyase (ACL) and results in decreased LDL in patients not responding to statin therapy.

Note - Ezetimibe, colestipol, bempedoic acid and PCSK-9 inhibitors like alirocumab have been proven to reduce atherosclerosis risk when added to statins, whereas the fibrate group of drugs and niacin show no significant effect in addition to statins.

Solution to Question 21:

Contraindicated in patients on enzyme inducer drugs is the true statement regarding Pretomanid.

Pretomanid is metabolized in part by CYP3A. So co-administration with strong or moderate CYP3A inducers should be avoided. Hence, it is contraindicated in patients on enzyme-inducer drugs.

Pretomanid is a bicyclic nitroimidazole that acts by inhibiting mycolic acid (a component of the mycobacterial cell wall) and protein synthesis in the mycobacteria.

Pretomanid in combination with Bedaquiline and Linezolid, was approved by the FDA in 2019 for the treatment of adult patients with Pre (XDR) TB (resistant to rifampicin and any fluoroquinolones) and for patients who are treatment-intolerant or nonresponsive to multidrug-resistant pulmonary (MDR) TB treatment.

Other options:

Option A: Pretomanid is given for Pre -XDR TB (not XDR TB) in combination with Bedaquiline and Linezolid (BPaL regimen). Pre-XDR TB is resistant to rifampicin and any fluoroquinolone, and therefore, BPaL is given to such patients.

Option C: Pretomanid acts by reactive nitrogen species like NO and not by reactive oxygen species.

Option D: Pretomanid is not used for resistant cases of malaria.

*Note: For people with MDR/RR-TB, the regimen comprises bedaquiline (B), pretomanid (Pa), linezolid (L) and moxifloxacin (M), and is referred to as BPaLM. For people who have pre-XDR-TB, the regimen can be used without moxifloxacin (BPaL).

Solution to Question 22:

Lacosamide is a drug acting on sodium channel and potentiating slow sodium inactivation.

Lacosamide acts by blocking the voltage-gated sodium channels. It is used as adjunctive therapy for focal-onset seizures in patients older than 17 years.

It is the first anti-seizure medication that limits persistent repetitive firing, the neuronal firing pattern typical of focal seizures, and prolongs the gradual inactivation of voltage-gated Na+

channels. Additionally, lacosamide binds to the phosphoprotein collapsing response mediator protein-2 (CRMP-2), which is mostly expressed in the nervous system and has a role in controlling axonal outgrowth and neuronal differentiation.

The role of CRMP-2 binding in seizure control is unknown.

Therapeutic Uses:

• It is used for both monotherapy and add-on therapy for focal-onset seizures in patients 17 years and older.

• It is mostly beneficial in hospitalized patients because it is available in an intravenous formulation, has minimal hepatic metabolism, and has no adverse respiratory effects.

Antiepileptic Drugs	Mechanism of Action
Phenytoin (Option A)CarbamazepineLamotrigine (Option B)	Inhibit Na channels
PhenobarbitalVigabatrinTiagabine	Enhance GABA inhibitionVigabatrin – block GABA aminotransferase irreversiblyTiagabine – block GABA reuptake
EthosuximideZonisamideGabapentinPregabalin	Block Ca channelsEthosuximide, Zonisamide –block T-type Ca channelsGabapentin, Pregabalin –block voltage sensitive Ca channels
Levetiracetam	Binds synaptic vesicle protein 2A
Valproate	Multiple MOA:GABA related actionNMDA receptor antagonistHistone deacetylase inhibitor
Topiramate (Option D)	Multiple MOA:GABAergic actionAMPA receptor antagonistActs on Na, Ca and K channels
Drugs acting on glutamate receptors	NMDA-type glutamate receptors:Valproate, FelbamateAMPA-type glutamate receptors: Topiramate, Perampanel
Newer drugs:RufinamideLacosamideEzogabinePerampanel	Prolongs inactivation of voltage-gated Na channels Enhances inactivation of voltage-gated Na channels Inhibits collapsin response mediator protein 2Potassium channel openerAMPA-type glutamate receptor antagonist

Solution to Question 23:

The marked receptor is GABA Transporter 1 Receptor (GAT-1), and Tiagabine acts on this receptor.

Tiagabine inhibits the neuronal uptake of GABA and prolongs its time in the synaptic cleft, where the GABA molecules can activate GABA receptors.

Other drugs which act on the GABAergic pathway are:

• Activation of GABA A receptors - Barbiturates, Phenobarbitone

• Activation of GABA B receptors - Baclofen

• Prevents the uptake of GABA - Tiagabine

• Prevents the degradation of GABA (inhibits GABA transaminase) - Vigabatrin



Other options:

Option B: Ganaxolone is a recently approved drug for the treatment of seizures associated with CDKL5 deficiency. The exact mechanism of action of Ganaxolone is unknown, but its anticonvulsant effects are thought to result from positive allosteric modulation of the GABA-A receptor in the CNS. It can be used in adults as well as children older than 2 years of age. The most common adverse reactions associated with the drug are somnolence, pyrexia, salivary hypersecretion and seasonal allergy.

Option C: Felbamate is the NMDA receptor antagonist prescribed to this patient. It also potentiates GABA action. It is indicated in refractory primary or secondary generalised seizures and Lennox-Gastaut syndrome. Felbamate is associated with aplastic anemia and liver failure.

Option D: Stiripentol is a GABA A receptor modulator with GABA-enhancing activity. It is clinically used in conjunction with valproate and clobazam for refractory generalized tonic-clonic seizures with severe myoclonic epilepsy in infants (Dravet syndrome).

Solution to Question 24:

The correct set of matching is a-ii, b-iii, c-i, d-iv.

Naloxone is an opioid antagonist used to reverse the effects of opioid overdose, such as respiratory depression.

Oxazepam is a benzodiazepine used for the management of delirium tremens, a severe alcohol withdrawal syndrome.

Varenicline is a partial agonist at the nicotinic acetylcholine receptors, used for smoking cessation by reducing withdrawal symptoms and cravings.

Acamprosate is a modulator of glutamate and GABA neurotransmission, used for the treatment of alcoholism by reducing cravings and the risk of relapse in alcohol abstinence.

a. Naloxone	ii. Opioid overdose
b. Oxazepam	iii. Delirium tremens
c. Vareniclin e	i. Smoking cessation
d. Acampros ate	iv. Treatment of alcoholism

Pharmacology NEET 2024

Question 1:

A man presents with mixed infection of Plasmodium vivax and Plasmodium falciparum in a health centre in Rajasthan. What is the dosage of the treatment given on the second day?

a) 4 tablets of Artesunate 50mg + 6 tablets of Primaquine 2.5mg

b) 4 tablets of Artesunate 50mg + 6 tablets of Primaquine 7.5mg

c) 4 tablets of Artesunate 50mg + Sulphadoxine 1500mg + 6 tablets of Primaquine 2.5mg

d) 4 tablets of Artesunate 45mg + Sulphadoxine 1500mg + 6 tablets of Primaquine 7.5mg

Question 2:

Which drug is used to treat multiple myeloma that is known to cause reactivation of Herpes Zoster?

a) Bortezomib

b) Lenalidomide

c) Daratumumab

d) Elotuzumab

Question 3:

A patient presented with breathing difficulty and nasal polyposis when he took aspirin for fever. He faced this issue previously too when he took aspirin. Which among the following is not true about this?

a) There is no allergic sensitization to aspirin and its metabolites

b) Responds to Monteleukast

c) Occur on exposure to only aspirin

d) Occurs with exposure to other NSAIDs too that block COX 1

Question 4:

A 75-year-old female presented with urinary incontinence. She also has cognition decline and dementia. Which of the following drugs is useful for treating urinary incontinence?

a) Tiotropium

b) Ipratropium

c) Tolterodine

d) Telenzipine

Question 5:

A 7-year-old boy presents with a two-week history of persistent fever, fatigue, and pallor. His parents also report easy bruising and gum bleeding. On examination, he is pale with petechiae on his legs, hepatosplenomegaly, and generalized lymphadenopathy. Blood tests show severe anemia, thrombocytopenia, and elevated white blood cell count with blast cells. What is the most appropriate initial treatment regimen for this patient?

a) IVIG

b) Doxorubicin + Idarubicin

c) Prednisolone + vincristine

d) Radiotherapy of lymph nodes

Question 6:

Lignocaine is used in a loading dose for the treatment of arrhythmias. The loading dose of this drug depends upon which of the following factors?

a) Clearance

b) Volume of distribution

c) Plasma half life

d) Bioavailability

Question 7:

A hypertensive patient with chronic kidney disease was having frequent attacks of renal colic. Which of the following diuretics is used?

a) Furosemide

b) Mannitol

c) Hydrochlorothiazide

d) Spironolactone

Question 8:

Identify the drug marked as 'X' in the image.



a) Clopidogrel

b) Amino caproic acid

c) Tenecteplase

d) Heparin

Question 9:

Which of the following beta-adrenergic blockers are beneficial for patients with reduced ejection fraction and can reduce mortality in chronic heart failure without causing exacerbations over a few years?

a) Propanolol

b) Pindolol

c) Esmolol

d) Bisoprolol

Question 10:

Which drug is used for maintenance therapy in opium addiction?

a) Buprenorphine

b) Naloxone

c) Butarphanol

d) Clonidine

Question 11:

A patient who recently underwent surgery has complaints of nausea and vomiting. He was prescribed a drug called aprepitant. What is its mechanism of action?

a) NK-3 antagonist

b) NK-2 agonist

c) NK-1 antagonist

d) NK-1 agonist

Question 12:

An HIV patient developed a fever and dry cough. He was diagnosed with TB. He is currently on Lamivudine, Zidovudine and Nevirapine. Which Anti -TB drug should be changed in this patient?

a) Isoniazid

b) Rifampicin

c) Streptomycin

d) Ethambutol

Question 13:

A 32-year-old woman diagnosed with diarrhoea-predominant irritable bowel syndrome presented to the casualty with severe abdominal pain. She was prescribed a drug that exerts a direct spasmolytic effect on the smooth muscles of the gastrointestinal tract. Which drug could it be?

a) Dicyclomine

b) Scopolamine

c) Galantamine

d) Rivastigmine

Question 14:

A patient is started on a 3 HP TPT regimen. Which of the following is true regarding the regimen?

a) There is more compliance in 3HP than 9H and 6H

b) No significant difference in 3HP, 6H, 9H

c) Chance of abortion in 3 HP is same as 6H

d) 3HP regimen is less effective than 6H and 9H in preventing active TB

Question 15:

A female suffered a head injury from a road traffic accident. On examination, there is elevated intracranial pressure. Which diuretic among the following can be used in her condition?

a) Mannitol

b) Hydrochlorothiazide

c) Spironolactone

d) Acetazolamide

Question 16:

A 2-year-old child presents to the emergency department with lethargy, difficulty feeding, and dehydration. His vitals are stable and his ECG is shown below. What is the next step in his management?



a) Synchronized cardioversion with 2J of current

b) Adenosine 0.1 mg/kg

c) Adenosine 0.2 mg/kg

d) Synchronized cardioversion with 1J of current

Question 17:

Which of the following drugs when added with erythromycin can lead to torsades de pointes?

a) Doxylamine

b) Loratidine

c) Promethazine

d) Astemizole

Question 18:

A child was started on ciclesonide inhalational therapy for bronchial asthma. Which of the following statements is incorrect regarding his treatment?

a) It has fewer side effects compared to other inhalational steroids

b) It is a prodrug metabolised by bronchial esterase

c) The occurence of oral candidiasis is more compared to other inhaled steroids

d) Oral bioavailability is negligible

Question 19:

A patient diagnosed with hypertriglyceridemia was started on a medication, following which, he developed flushing, hyperglycemia, and elevated liver enzymes. Identify the drug consumed by the patient.

a) Nicotinamide

b) Rosuvastatin

c) Ezetemibe

d) Fenofibrate

Question 20:

Which of the following reasons best explains the combination of Atropine (0.025 mg) and Diphenoxylate (2.5 mg) for treating secretory diarrhoea?

a) Atropine potentiates the action of smooth muscle relaxation

b) It prevents overdose and abuse of Diphenoxylate

c) It antagonizes the side effect of Diphenoxylate

d) Atropine increases the antimotility action of diphenoxylate

Question 21:

A hypertensive patient is started on a drug to control his BP. He had complaints of dry mouth, drowsiness, and confusion after taking this drug, but he still continued to take the drug for 1 year. He had to travel abroad for work, and he forgot to take his medications. He now presents with a blood pressure of 200/120mmHg. Identify the drug he was on?

a) Amlodipine

b) Atenolol

c) Clonidine

d) Enalapril

Question 22:

In which phase of clinical trials is drug dosing done?

a) Phase I

b) Phase II

c) Phase III

d) Phase IV

Question 23:

A patient with a gynecological malignancy is started on Methotrexate. Later she develops oral and gastric ulcers. Which nucleotide synthesis is affected in her?

a) UDP

b) CTP

c) TMP

d) GMP

Question 24:

The drug 'X' is a GHRH analogue that is FDA-approved for the treatment of HIV-associated lipodystrophy and not used for Growth hormone deficiency. Identify the drug 'X'.

a) Octreotide

b) Somatropin

c) Tesamorelin

d) Mecasermin

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	c
5	c
6	b
7	c
8	b
9	d

10	a
11	c
12	b
13	a
14	a
15	a
16	b
17	d
18	c
19	a
20	b
21	c
22	a
23	c
24	c

Detailed Explanations

Solution to Question 1:

The dosage of the treatment given on the second day for mixed infection with Plasmodium vivax and Plasmodium falciparum is 4 tablets of Artesunate 50mg + 6 tablets of Primaquine 2.5mg.

For treating mixed infections of Plasmodium vivax and Plasmodium falciparum, the following treatment regimen is recommended. This approach ensures comprehensive treatment by addressing both the blood stages of Plasmodium falciparum with ACT and the liver stages of Plasmodium vivax with Primaquine.

Dosage chart for treatment of mixed (vivax and falciparum) malaria with ACT-SP

Age	Day 1			Day 2		Day 3		Day 4-14
	AS tablet (50 mg)	SP tablet	PQ (2.5 mg)	AS tablet (50 mg)	PQ (2.5 mg)	AS tablet (50 mg)	PQ (2.5 mg)	PQ (2.5 mg)
Less than 1 year	½	½	0	½	0	½	0	0
1-4 years	1	1	1	1	1	1	1	1
5-8 years	2	1½	2	2	2	2	2	2
9-14 years	3	2	4	3	4	3	4	4
15 years or more	4	3	6	4	6	4	6	6

©MARROW

Solution to Question 2:

The drug used to treat multiple myeloma and that has the potential to cause reactivation of Herpes Zoster is bortezomib.

Bortezomib is an inhibitor of proteasome-mediated protein degradation. It is central to the treatment of multiple myeloma, both untreated and relapsed. Bortezomib is given intravenously or subcutaneously and dosage depends upon the disease, route of administration, and the patient's history. It is known to predispose patients to the reactivation of herpesvirus, including Herpes Zoster.

Adverse effects include thrombocytopenia, fatigue, peripheral neuropathy, neutropenia, anaemia, anorexia, nausea, limb pain, dizziness, etc. Other adverse events like cardiac tamponade, disseminated intravascular coagulation, toxic epidermal necrolysis, etc. have been reported.

Other options:

Option B: Lenalidomide is the drug of choice in patients with myelodysplastic syndrome (MDS) associated with 5q deletion, and is an immunomodulatory analogue of thalidomide. It suppresses bone marrow and is often associated with significant leukopenia.

Option C: Daratumumab is a monoclonal antibody against CD-38.

Option D: Elotuzumab is a monoclonal antibody against SLAMF7, a glycoprotein expressed in multiple myeloma cells and natural killer cells. It is used in combination with lenalidomide and dexamethasone for the treatment of multiple myeloma.

Solution to Question 3:

The diagnosis for this patient with breathing difficulty and nasal polyposis, triggered by aspirin, a COX-1 inhibitor is aspirin-induced asthma, also known as Samter's triad. The statement that is

not true is that "it occurs on exposure to only aspirin", as this condition can also be triggered by other NSAIDs that inhibit COX-1, and is not restricted to just aspirin alone.

Samter's triad is also known as aspirin-induced asthma or aspirin-sensitive asthma. It consists of:

• Nasal polyposis

• Bronchial asthma

• Aspirin intolerance

A patient with Samter's triad experiences severe asthma and nasal polyposis triggered by aspirin or other COX-1 inhibitors. This condition involves nasal congestion, rhinorrhea, anosmia, and recurrent chronic rhinosinusitis, with acute asthma attacks often occurring a few hours after ingestion of these medications.

These patients overproduce leukotrienes due to COX-1 inhibition and decreased prostaglandin E2 (PGE2) levels. Therefore, they should avoid COX-1 inhibitors like aspirin and most NSAIDs but can generally tolerate COX-2 inhibitors and acetaminophen.

Treatment typically includes leukotriene modifiers, and in some cases, aspirin desensitization may be used to manage symptoms and allow the use of aspirin or NSAIDs if necessary for other conditions. Recent advances in treatment include biologics such as Dupilumab and IL-5 inhibitors, which are increasingly effective in managing aspirin-induced asthma and may replace aspirin desensitization except when ongoing aspirin or NSAID therapy is required.

Other options:

Option A: The reaction in aspirin-induced asthma is not due to an allergic sensitization but rather due to the inhibition of COX-1 and subsequent overproduction of leukotrienes.

Option B: Montelukast, a leukotriene receptor antagonist, is effective in managing symptoms of aspirin-induced asthma by counteracting the elevated levels of leukotrienes that contribute to inflammation and respiratory issues.

Option D: It occurs with exposure to other NSAIDs too, that block COX-1. Patients with aspirin-induced asthma can react to other NSAIDs that inhibit COX-1, not just aspirin.

Solution to Question 4:

For managing urinary incontinence in a 75-year-old female with cognitive decline and dementia, tolterodine is useful.

Tolterodine is a muscarinic receptor antagonist specifically effective for treating urinary incontinence. It exhibits selectivity for the urinary bladder, which helps to minimize side effects compared to other antimuscarinic agents. This selectivity is beneficial for patients with cognitive decline or dementia, as it reduces the risk of adverse effects. Tolterodine is metabolized by CYP2D6, so its dosage may need adjustment based on individual CYP2D6 activity, making it a suitable option for managing urinary incontinence while minimizing potential side effects.

Other muscarinic antagonists for urinary incontinence include oxybutynin, trospium chloride, darifenacin, solifenacin, and fesoterodine. Oxybutynin, though effective, often causes antimuscarinic side effects such as xerostomia. Trospium chloride, being a quaternary amine, is better tolerated and excreted primarily via the kidneys, requiring dosage adjustments in renal

impairment. Darifenacin and solifenacin have selectivity for M3 receptors, which helps balance efficacy and side effects, while fesoterodine, a prodrug, provides a stable pharmacokinetic profile due to its conversion to tolterodine's active metabolite.

Other Options:

Option A: Tiotropium and Option C: Ipratropium are muscarinic antagonists used primarily for respiratory conditions such as chronic obstructive pulmonary disease (COPD) and asthma, rather than urinary incontinence. Tiotropium is a long-acting agent with selectivity for M1 and M3 receptors, while Ipratropium is used more frequently for its short-acting effects.

Option D: Telenzepine, an analogue of Pirenzepine, displays higher potency and selectivity for M1 receptors compared to pirenzepine and is used in treating acid-peptic disease, not urinary incontinence.

Solution to Question 5:

The given clinical scenario suggests a diagnosis of Acute Lymphoblastic Leukemia (ALL), and standard initial treatment in children includes a chemotherapy regimen with drugs like prednisolone and vincristine during the induction phase.

Patients diagnosed with ALL present with symptoms of impaired marrow function like anaemia, infections secondary to neutropenia, and bleeding due to thrombocytopenia. Other symptoms include bone pain, generalized lymphadenopathy, splenomegaly, hepatomegaly, testicular enlargement, headache, vomiting, and nerve palsies.

Treatment of ALL

Induction: L-Asparaginase, vincristine, prednisolone, and anthracycline

CNS prophylaxis with intrathecal methotrexate

Consolidation: Methotrexate, cytarabine

Maintenance: 6-Mercaptopurine, methotrexate plus intrathecal therapy

Other Options:

Option A: IVIG: Intravenous immunoglobulin (IVIG) is not used for treating ALL.

option B: Both doxorubicin and idarubicin are anthracyclines used in chemotherapy. While doxorubicin can be part of certain ALL treatment protocols, idarubicin is more commonly used in Acute Myeloid Leukemia (AML) rather than ALL. This combination is not typically used in the induction phase for ALL children.

Option D: Radiotherapy is not a standard initial treatment for ALL. It may be considered in specific cases, such as central nervous system involvement or certain high-risk cases, but the primary treatment for ALL is systemic chemotherapy, not localized radiotherapy.

Solution to Question 6:

The volume of distribution of the drug determines the loading dose of the drug.

Loading Dose = Volume of distribution x Desired steady-state plasma concentration.

When the time to reach a steady-state is long, as it is for drugs with long half-lives, it may be desirable to administer a loading dose that promptly raises the concentration of the drug in plasma to the target concentration.

Other options:

Option A: Clearance is used to determine the maintenance dose.

Maintenance dose = clearance x desired steady-state plasma concentration.

Options C and D: Plasma half-life primarily determines the frequency of dosing, while bioavailability determines the route of administration. Though bioavailability may affect the dose necessary, it will not be the primary factor considered during the calculation of the loading dose.

Solution to Question 7:

For a hypertensive patient with chronic kidney disease experiencing frequent attacks of renal colic, hydrochlorothiazide, a thiazide diuretic, is often used.

Thiazide diuretics are particularly effective in cases of calcium oxalate kidney stones because they reduce urinary calcium excretion. They work by acting on the distal convoluted tubule (DCT) of the nephron, where they inhibit the Na+/Cl- co-transporter. This inhibition decreases the intracellular sodium concentration, which, through compensatory mechanisms, leads to increased calcium reabsorption into the bloodstream.

While this mechanism helps in decreasing urinary calcium and managing calcium oxalate stones, thiazide diuretics can also cause adverse effects such as hypokalemia, hyperuricemia, and hypercalcemia. The increased reabsorption of uric acid due to the diuretic effect can exacerbate conditions like gout and should be monitored carefully.

In addition to their use in treating calcium oxalate kidney stones, thiazide diuretics are commonly prescribed for managing uncomplicated hypertension and treating edema associated with heart, liver, and kidney diseases. They are also used for conditions like osteoporosis and nephrogenic diabetes insipidus.

Other options:

Option A: Furosemide is a diuretic that can increase calcium excretion, potentially raising PTH as the body tries to conserve calcium.

Option B: Osmotic diuretics, like mannitol, retain water isoosmotically in the proximal convoluted tubule (PCT) and descending loop of Henle which are freely permeable to water. Luminal fluid is diluted and opposes NaCl reabsorption and promotes calcium excretion. They are indicated in raised intracranial tension and in dialysis disequilibrium.

Option D: Spironolactone is a potassium-sparing diuretic and aldosterone antagonist. It works by competitively inhibiting aldosterone receptors in the distal tubules of the kidney.

Solution to Question 8:

The drug marked as 'X' in the image is aminocaproic acid.

ε-Aminocaproic acid, a lysine analogue that competes with plasminogen for binding sites on fibrin. By blocking this interaction, ε-Aminocaproic acid prevents the conversion of plasminogen into plasmin, thereby inhibiting fibrinolysis. This action makes it a useful agent for reversing the effects of fibrinolytic therapy.

However, ε-Aminocaproic acid should be administered with caution as it can suppress intrinsic fibrinolytic activity, potentially leading to complications such as disseminated intravascular coagulation (DIC). To address this, fresh frozen plasma and cryoprecipitate may be used to restore fibrin and clotting factors.



Solution to Question 9:

The beta-adrenergic blocker that is useful in patients with reduced ejection fraction and can reduce mortality in chronic heart failure without causing exacerbations over a few years is bisoprolol.

Bisoprolol is preferred for use in patients having heart failure with reduced ejection fraction (HFrEF).

The exacerbation mentioned in the context of beta-blockers for heart failure refers to the potential worsening of heart failure symptoms in some patients, despite the overall benefits of these medications. They can sometimes lead to a decline in left ventricular function, particularly in individuals with limited cardiac reserve who rely heavily on adrenergic stimulation. This deterioration can manifest as increased heart failure symptoms or a decrease in cardiac function. The only beta-blockers that have been proven to improve the survival of patients with HFrEF or chronic HF are -

• Bisoprolol

• Metoprolol

• Carvedilol

HFrEF is a complex clinical syndrome where the ventricle fails to adequately fill with or eject blood, typically defined by an ejection fraction (EF) less than or equal to 40%. Key etiologies include coronary artery disease, valvular heart disease, congenital heart diseases, and autoimmune diseases like giant cell myocarditis. Patients with HFrEF commonly experience symptoms like dyspnea on exertion, orthopnea, and paroxysmal nocturnal dyspnea, and may present with pedal edema, elevated jugular venous pressure (JVP), ascites, and hepatomegaly on examination.

The pharmacological management of HFrEF involves drugs with both symptomatic and mortality benefits, including Angiotensin receptor–neprilysin inhibitors (ARNIs) such as sacubitril/valsartan, angiotensin-converting enzyme (ACE) inhibitors like ramipril and enalapril, β-blockers such as metoprolol and carvedilol, mineralocorticoid receptor antagonists like spironolactone and eplerenone, sodium-glucose cotransporter 2 (SGLT2) inhibitors like dapagliflozin, and hydralazine plus isosorbide dinitrate (noting a mortality benefit specifically in African-American patients). Drugs used primarily for symptomatic relief, such as loop diuretics (e.g., furosemide), and digoxin, are reserved for specific indications or during decompensation. Non-pharmacological therapies for HFrEF include implantable cardioverter-defibrillators (ICD) and cardiac resynchronization therapy.

Solution to Question 10:

The drug used among the following options for maintenance therapy in opium addiction is buprenorphine.

Buprenorphine is a partial μ-opioid agonist and an antagonist at δ and κ receptors. It is widely used both as an analgesic for chronic pain conditions, such as cancer pain, and as a treatment for opioid dependence, serving as an alternative to methadone.

It effectively controls cravings and blocks the effects of higher-efficacy opioids. It is often administered in combination with naloxone to prevent misuse, although it may be given alone to pregnant women or individuals with naloxone allergies. Initial dosing typically starts at 2-4 mg daily, with adjustments made to stabilize the dose at approximately 12 to 16 mg/day.

For opioid use disorder, effective management includes both withdrawal and long-term treatment strategies. Withdrawal and detoxification generally start with managing physical dependence using medications like methadone or buprenorphine. Methadone, a full agonist, is used to stabilize and reduce withdrawal symptoms, while buprenorphine helps with cravings and has a lower risk of overdose.

Clonidine and lofexidine, alpha-2 agonists, are used to ameliorate withdrawal symptoms. Lofexidine, approved by the FDA, is the first non-opioid treatment specifically for opioid withdrawal in adults. Naltrexone, a long-acting opioid antagonist, is used as an adjunct in patients who are already opioid-free; however, it will precipitate acute withdrawal symptoms if administered to those still dependent on opioids.

Solution to Question 11:

The class of drugs most preferred in patients suffering from postoperative nausea and vomiting (PONV), following surgery are the NK1 antagonists like aprepitant.

Substance P is a neuropeptide that interacts with neurokinin-1 (NK-1) receptors. NK1 antagonists inhibit substance P at central and peripheral receptors. Aprepitant, fosaprepitant are examples of NK1 antagonists. It also acts on the chemotrigger zone whose location is shown below.



PONV Physiology: Afferents from the chemoreceptor trigger zone (CTZ), vagal neurons in the gastrointestinal system, neurons from the vestibular system and midbrain afferents relay within the vomiting centre in the reticular formation of the brainstem which then project into the nucleus tractus solitarius.

Risk factors for PONV are divided into:

• Patient factors

• Female>male

• Age<50 years

• Prior history of motion sickness

• Non-smokers > smokers (CTZ is desensitized)

• Preoperative factors

• (Fasting and anxiety, while possibly contributing to PONV, have yet to be demonstrated)

• Intraoperative factors

• Surgical - Cholecystectomy, gynaecological and laparoscopic procedures have higher incidences

• Drugs - Nitrous oxide, Opioids, Cyclopropane,

• General anaesthesia has a times more risk of developing PONV than regional anaesthesia.

• Postoperative factors

• Pain and early ambulation contribute to a higher risk of PONV.

Scoring systems used to grade PONV are the Apfel score and the Koivuranta score.

Drugs used in the prophylaxis and treatment of PONV:

• 5-HT 3 antagonists - ondansetron, palonosetron

• Butyrophenones- droperidol

• Dexamethasone

• Transdermal scopolamine

• Aprepitant

Note: Double-blind studies have found no difference in the efficacy between aprepitant(NK1 antagonists) and ondansetron (5HT3 antagonist) in the first 24 hours following surgery. But after 24 hours aprepitant has proven to be more productive than ondansetron in the management of PONV.

Solution to Question 12:

Rifampicin should be avoided as it is a potent inducer of CYP enzymes including CYPs 1A2, 3A4, 2C9, and 2C19, which in turn can induce the metabolism of nevirapine and results in therapeutic failure of ART.

Rifabutin causes the least induction of cytochrome enzymes and is therefore preferred in HIV patients who are on protease inhibitors/NNRTIs.

Cytochrome enzyme-inducing potency of rifamycins is rifampicin > rifapentine > rifabutin.

Note:

Cytochrome p-450 inducers	Cytochrome p-450 inhibitors
BarbituratesCarbamazepineRifampinPhenytoinAminoglutethimideCigarette smokingChronic alcoholism	Macrolide antibioticsAzole antifungalsChloramphenicolOmeprazoleSelective serotonin reuptake inhibitors (SSRIs)HIV protease inhibitorsCimetidineCiprofloxacinMetronidazole

Solution to Question 13:

Dicyclomine is a weak muscarinic receptor antagonist with non-specific direct spasmolytic effects on the smooth muscle of the gastrointestinal tract.

Other options:

Option B: Scopolamine exerts spasmolytic activity indirectly through its antimuscarinic action. It is not preferred due to its adverse effects like drowsiness, dryness, and blurred vision.

Option C and D: Galantamine and rivastigmine are central cholinesterase inhibitors and do not possess spasmolytic activity.

Solution to Question 14:

The 3HP regimen, which involves once-weekly doses of Isoniazid and Rifapentine for 12 weeks, has been found to have higher patient compliance than the 9-month regimen of Isoniazid (9H) and the 6-month regimen (6H).

The shorter duration and reduced number of doses in the 3HP regimen make it more convenient and thus lead to better adherence. Studies have shown that patients are more likely to complete the 3HP regimen due to the shorter treatment duration and lower pill burden.

TPT stands for Tuberculosis Preventive Therapy. It is a treatment given to individuals who are at high risk of developing active tuberculosis (TB) to prevent the progression from latent TB infection (where the person is infected with the TB bacteria but does not have symptoms) to active TB disease.

There are different regimens for TPT, including:

• 3HP: A regimen consisting of once-weekly doses of Isoniazid (H) and Rifapentine (P) for 12 weeks.

• 6H: A regimen consisting of daily Isoniazid for 6 months.

• 9H: A regimen consisting of daily Isoniazid for 9 months.

TPT is particularly important for individuals with latent TB who have a higher risk of developing active TB, such as those with HIV, close contacts of TB patients, or those with certain medical conditions that weaken the immune system.

Other options:

Option B: Research has demonstrated that the 3HP regimen offers better compliance and similar effectiveness compared to the traditional 6H and 9H regimens. The shorter duration of 3HP (once weekly for 12 weeks) makes it more convenient and tolerable for patients, which leads to higher adherence rates. Thus, there is a significant difference in patient compliance between these regimens, with 3HP showing clear advantages.

Option C: Although 3HP is generally considered safe, data suggests that the risk profile for adverse effects, including potential reproductive impacts like abortion, differs between regimens. Specifically, 6H may have a different side effect profile due to the prolonged exposure to the drug over six months, compared to the shorter and intermittent dosing of 3HP.

Option D: Studies have confirmed that the 3HP regimen is as effective as 6H and 9H in preventing active tuberculosis (TB). Both regimens aim to prevent latent TB from progressing to active disease, and 3HP has been shown to achieve similar outcomes with fewer doses, enhancing patient compliance.

Solution to Question 15:

In this patient with raised intracranial pressure after a road traffic accident, mannitol is the diuretic of choice in her management to reduce intracranial pressure.

Mannitol is an osmotic diuretic that draws fluid out of the brain tissue and into the bloodstream as it forms a hypertonic solution when it reaches the bloodstream, thereby reducing intracranial pressure (ICP). It is commonly used in cases of traumatic brain injury or other conditions that cause elevated ICP. This effect helps to relieve pressure on the brain, preventing further damage.

Other options:

The other diuretics listed Hydrochlorothiazide, Spironolactone, and Acetazolamide are not typically used for managing elevated ICP. They are not hypertonic solutes and thus do not lead to a rapid reduction of the ICP as mannitol does.

Solution to Question 16:

This clinical history along the EKG showing narrow QRS complexes without P waves and pseudo R waves in aVR lead suggests paroxysmal supraventricular tachycardia (PSVT) and the next step in his management would be an initial dose of adenosine at 0.1 mg/kg.

PSVT is a regular, narrow QRS complex tachycardia, most often caused due to an atrioventricular nodal re-entrant circuit. Patients may present with sudden onset palpitations, dizziness, and diaphoresis.

Management involves slowing AV node conduction and vagal maneuvers (e.g., Valsalva maneuver, splashing cold water on the face) are the first step. If this does not work, intravenous (IV) adenosine is the drug of choice used to terminate the majority of PSVT episodes.

Adenosine is a very short-acting endogenous purine signal molecule. It is administered as a 6 mg IV bolus over 2-3 seconds followed by a rapid flush with 20 mL NS. It can be followed up with a 12 mg IV bolus dose if required. It causes membrane hyperpolarization of cardiac muscle cells to terminate the re-entrant circuit through the AV node.

Since the action of the drug lasts for less than a minute, adverse effects are short-lived. These include transient shortness of breath, chest pain, hypotension, and cutaneous flushing due to vasodilation. It can however precipitate bronchospasm in asthmatics and verapamil is preferred in such patients.



When the patient is hemodynamically unstable, i.e., hypotension with unconsciousness or respiratory distress, QRS-synchronous direct current cardioversion is done.



The given ECG shows pseudo R' waves:



Solution to Question 17:

Among the given drugs, astemizole increases the risk of Torsades de pointes when used concomitantly with erythromycin.

Drugs that block potassium channels and prolong QT increase arrhythmogenic potential. Astemizole is a second-generation antihistamine that blocks delayed potassium channels and prolongs the QT interval.

Other options:

Option A: Doxylamine is an antihistamine commonly used as a short-term treatment for insomnia. It is combined with pyridoxine (vitamin B6) to treat nausea and vomiting during pregnancy (morning sickness). It does not prolong the QT interval.

Option B: Loratadine is a second-generation antihistamine used to treat allergic rhinitis (hay fever) and urticaria (hives). It is known for being non-sedating due to its minimal penetration into the central nervous system. It does not prolong the QT interval.

Option C: Promethazine is a first-generation antihistamine with multiple uses, including the treatment of allergies, motion sickness, nausea and vomiting, and as a sedative or sleep aid. It also has antiemetic properties and is sometimes used in the management of postoperative nausea and vomiting.

Solution to Question 18:

The incorrect statement is option C as ciclesonide has a lower risk of oral candidiasis as compared to other inhaled steroids.

Ciclesonide is an inhaled corticosteroid that is unique because it is a prodrug. It is metabolised by bronchial esterases in the lungs to its active form, which allows for targeted action within the airways. This feature reduces systemic side effects and the risk of local side effects, such as oral candidiasis (thrush), compared to other inhaled steroids.

Other options:

Option A: Ciclesonide's activation in the lungs minimises systemic absorption, leading to fewer side effects.

Option B: Ciclesonide is inactive until it is converted to its active form in the lungs.

Option D: Due to its prodrug nature and metabolism in the lungs, ciclesonide has low oral bioavailability, reducing systemic exposure.

Solution to Question 19:

The drug most likely responsible for causing flushing, hyperglycemia, and elevated liver enzymes in a patient being treated for hypertriglyceridemia is Nicotinamide (Niacin).

Niacin (nicotinic acid), is commonly used to treat hypertriglyceridemia and hypercholesterolemia. However, it is associated with side effects such as flushing, which is due to prostaglandin-mediated vasodilation. Niacin can also lead to hyperglycemia by impairing insulin sensitivity and may cause elevated liver enzymes, indicating potential hepatotoxicity.

Other options:

Option B: Rosuvastatin is a statin, primarily used to lower LDL cholesterol, and while it may elevate liver enzymes, it is less commonly associated with flushing and hyperglycemia. It can cause myopathy in patients.

Option C: Ezetimibe inhibits cholesterol absorption in the intestines and is not typically associated with these side effects.

Option D: Fenofibrate is used to lower triglycerides can elevate liver enzymes and can increase the risk of developing myopathy if used concomitantly with statins, but it does not commonly cause flushing or hyperglycemia.

Solution to Question 20:

Atropine is included in combination with diphenoxylate to prevent abuse and overdose of diphenoxylate.

Diphenoxylate is structurally similar to opioids and has the potential for misuse at higher doses. Additionally, it crosses the blood-brain barrier, making it more likely to cause dependence. The anticholinergic effects of atropine (e.g., dry mouth, tachycardia, blurred vision) become prominent at higher doses, discouraging overdose.

Diphenoxylate is an opioid agonist primarily used as an antidiarrheal agent. It works by slowing down intestinal motility, allowing more time for water and electrolytes to be absorbed from the intestines, which helps reduce diarrhea. However, because of its structural similarity to opioids, diphenoxylate has the potential for abuse and dependence if taken in large doses.

Solution to Question 21:

The adverse effects seen in this patient and the presentation of rebound hypertension on sudden drug withdrawal indicate that the drug is clonidine.

Clonidine is an alpha-2 receptor agonist and a centrally-acting antihypertensive agent. It treats high blood pressure by stimulating presynaptic $\alpha 2$ receptors in the vasomotor center in the brainstem. This binding has a sympatholytic effect, by suppressing the release of norepinephrine and other sympathomimetics.

Abrupt withdrawal after long-term therapy can result in rebound hypertension, restlessness, and sympathetic overactivity. Dry mouth and sedation are the major side effects of clonidine. It can also cause impotency and constipation. Rebound hypertension is a potentially serious complication and occurs if the drug is missed for 1–2 days. This is associated with an alarming rise in blood pressure which may be more than the pretreatment level.

Phentolamine is used for the control of hypertension due to clonidine withdrawal.

Solution to Question 22:

The dosing of a new drug is tested in phase I of a clinical trial.

Phase I trials are usually performed on healthy volunteers to check the safety and pharmacokinetic properties of a drug. The participants in this phase do not gain any direct health benefits from taking part. They are generally paid a fee for their time, with the payments being

regulated and not related to the risk involved.

In clinical trials, drug dosing typically occurs in Phase I, where researchers establish the optimal dose to achieve desired effects while minimizing side effects. In these phases, they focus on safety, tolerability, and pharmacokinetics.

Dose-ranging, primarily conducted in Phase II trials, involves testing multiple doses to identify the most effective and safe dose range. This phase helps determine the minimum effective dose and maximum tolerated dose, providing critical data for subsequent studies in Phase 3, where the chosen dose is tested for efficacy and safety in a larger population.

Solution to Question 23:

TMP synthesis is affected due to methotrexate.

Methotrexate inhibits dihydrofolate reductase (DHFR), which is required to synthesize tetrahydrofolate. Tetrahydrofolate is essential for converting dUMP to dTMP (thymidine monophosphate) in the nucleotide synthesis pathway. Therefore, the nucleotide synthesis affected in this patient is TMP (thymidine monophosphate).

By inhibiting DHFR, methotrexate reduces the availability of THF, thereby disrupting the production of thymidylate (TMP), which is necessary for DNA replication and repair. This disruption leads to impaired DNA synthesis in rapidly dividing cells, such as cancer cells. However, it also affects other rapidly dividing cells, including those in the gastrointestinal tract and oral mucosa, leading to side effects like oral and gastric ulcers.



Solution to Question 24:

Tesamorelin is approved for use in HIV-associated lipodystrophy but not for GH deficiency. It is a synthetic N-terminal modified form of human GHRH. It does not get degraded by dipeptidyl peptidase 4 and therefore has a longer duration of action.

It increases the levels of growth hormone (GH) and insulin-like growth factor (IGF-1) but reduces visceral fat accumulation.

Tesamorelin is approved for use in HIV-associated lipodystrophy but not for GH deficiency.

Other options:

Option A: Octreotide is a somatostatin analogue that decreases growth hormone levels. It is used in the treatment of acromegaly, variceal bleeding, diarrhoea due to adenomas secreting VIP (vasoactive intestinal peptide), and sulfonylurea toxicity. One of its adverse effects is gallstone formation.

Option B: Somatropin is a growth hormone analogue. It is used in growth hormone deficiency and other short-stature conditions like Turner's syndrome, Noonan syndrome, and Prader-Willi syndrome. Their use is contraindicated in acute critical illness, active malignancy, and proliferative retinopathy.

Option D: Mecasermin is a recombinant human IGF-1. It is used in IGF-1 deficiency, growth hormone receptor defects, and antibodies against growth hormone receptors. Adverse effects include hypoglycemia and lipohypertrophy. Another recombinant human IGF-1 with its IGFBP-3, mecasermin rinfabate has been FDA-approved for HIV-associated adipose redistribution syndrome.

Microbiology AIIMS 2017

Question 1:

A 28 year old male admitted with history of altered sensorium and hydrophobia. A suspicion of rabies was made and you want to diagnose the condition. You have taken corneal scrapings from the patient and which of the following tests will you do to confirm your diagnosis

a) Indirect immunofluorescence

b) Seller stain for negri bodies

c) Viral culture

d) Real time PCR

Question 2:

Identify the parasite from the given life cycle.



a) Fasciola hepatica

b) Fasciolopsis buski

c) Pargaonimus westermani

d) Clonorchis sinensus

Question 3:

The life cycle given below is of which of the following:



a) Entamoeba vermicularis

b) Ancylostoma duodenal

c) Balantidium coli

d) Entamoeba histolytica

Question 4:

A 25 year old male with history of fever and enlarged lymph nodes and sore throat was suspected to have infectious mono-nucleosis and was ordered a Paul Bunnel test. Which of the following is the principle of the test?

a) Complement mediated cell damage

b) Agglutination reaction

c) Heterophile antibody test

d) Neutralisation reaction

Question 5:

A blood spill in the operation theatre must be cleaned with _____.

a) Chlorine compound

b) Phenolic compound

c) Quarternary ammonium compounds

d) Alcoholic compounds

Question 6:

Urine LAM is used for the diagnosis of _____

a) M. tuberculosis

b) M. leprae

c) Mycoplasma

d) Listeria monocyogenes

Question 7:

Identify the organism in the given slide stained with Gomori methenamine silver stain.



a) Obtuse angled mucor

b) Acute angled penicillium

c) Acute angled branching aspergillus

d) Obtuse angled rhizopus

Question 8:

A confirmed case of which of the following viral haemorrhagic fevers was found in India?

a) Ebola hemorrhagic fever

b) Yellow fever

c) Crimean congo hemorrhagic fever

d) Marburg hemorrhagic fever

Question 9:

The given life cycle is suggestive of which of the following viruses?



a) HIV

b) Hepatitis B

c) Influenza

d) Herpes simplex

Question 10:

1,3 beta D- glucan is helpful for identification of?

a) Invasive candidiasis

b) Rhizopus

c) Cryptococcus

d) Mucormycosis

Question 11:

Relapse is seen in ?

a) Plasmodium vivax and falciparum

b) Plasmodium ovale and falciparum

c) Plasmodium vivax and ovale

d) Plasmodium vivax and malariae

Question 12:

Identify the organism.



a) Balantidium coli

b) Entamoeba dispar

c) Cryptosporidium

d) Giardia

Question 13:

A female from Himachal Pradesh presented with H/O thorn prick, a year back has a verrucous lesion in the skin with the following microscopic findings. Identify the agent?



a) Blastomycosis

b) Pheohyphomycosis

c) Chromoblastomycosis

d) Sporotrichosis

Question 14:

The following slide is given as a spotter in your microbiology practical examination. Identify the following organism.



a) Mycobacteria

b) Nocardia

c) Actinomyces

d) Corynebacterium

Question 15:

Identify the organism from the given image?



a) Vibrio

b) Leptospira

c) Helicobacter

d) Salmonella

Question 16:

Which of the following organisms causes the following disease?



a) Coxsackievirus

b) Human herpes virus 7

c) Pox virus

d) Molluscum contagiosum

Question 17:

Which of the following is not a dimorphic fungus?

a) Blastomyces dermatidis

b) Histoplasma capsulatum

c) Pneumocystis jirovecii

d) Penicillium marneffi

Question 18:

There are 3 to 5% healthy hepatitis B carriers in India who are asymptomatic. They are at the risk of developing hepatocellular carcinoma in the future because _____

a) They are unable to mount inflammation against the virus

b) Virus can integrate with host DNA and form complementary DNA

c) There is a risk of elevation of transaminases

d) Liver parenchymal cells are in a state of high proliferation

Question 19:

Identify the following?



a) Entamoeba dispar

b) Giardia lamblia

c) Balantidium coli

d) Entamoeba histolytica

Question 20:

Best method of diagnosis for Clostridium difficile?

a) Pure strain isolation from culture

b) Immunoflourescence

c) Toxin gene detection by PCR

d) Toxin Detection by ELISA

Question 21:

A patient with HIV develops diarrhoea and faecal examination shows Isospora belli. He was treated with cotrimoxazole. Diarrhoea subsided but fever persisted. Bone marrow examination of this patient is shown in the given image. Which of the following statement is false?



a) It cannot be grown in sabouraud dextrose agar

b) Spores are infective form

c) It is intracellular budding yeast

d) It can cause systemic disease

Question 22:

Bio-film confer resistance to antibiotics by all except _____

a) Mechanical barrier

b) Increased excretion of antibiotics

c) Slow metabolism

d) Adherence

Question 23:

Identify the egg in the picture given below:

a) A. lumbricoides

b) Strongyloides stercoralis

c) E. vermicularis

d) Ancylostoma duodenale

Question 24:

Identify the organism from the given staining technique.



a) Mycobacterium leprae

b) Mycobacterium tuberculosis

c) Actinomyces israelli

d) Nocardia asteroids

Question 25:

Which of the following staphylococcal infections is not toxin mediated?

a) Food poisoning

b) Toxic shock syndrome

c) Septicemia

d) Staphylococcal scalded skin syndrome

Question 26:

All are features of Zika Fever except

a) Petechial rash

b) Guillan barre syndrome

c) Fever with arthritis

d) Conjunctivitis

Question 27:

(1–3)-β-D glucan assay can not be used in

a) Invasive cryptococcosis

b) Invasive aspergillosis

c) Pneumocystis jiroveci pneumonia

d) Invasive candidiasis and candidemia

Question 28:

A patient was sputum negative for tuberculosis, but chest X-ray was suggestive of tuberculosis. Which is the next step according to RNTCP ?

a) Line probe assay

b) CB - NAAT

c) Liquid culture

d) None of the above

Answer Key

Question No.	Correct Option
1	c
2	a
3	d
4	c
5	a
6	a
7	c
8	c
9	b
10	a
11	c
12	a
13	c
14	a
15	b
16	a
17	c
18	b
19	b
20	c
21	a
22	d
23	d
24	d
25	c
26	a
27	a
28	b

Detailed Explanations

Solution to Question 1:

Viral culture from corneal scrapings is used to confirm diagnosis.

Multiple tests in succession/simultaneously are the best. Usually Direct Fluorescent Antibody test followed by virus isolation/culture is useful.

Virus isolation through culture: Best specimens are brain and saliva. CSF, corneal smears and centrifuged urine can also be used.

Option A: Serological tests like indirect immunofluorescence are insensitive.

Option B: Negri bodies can only be demonstrated in brain (pyramidal cells of Ammon's horn, and the Purkinje cells of the cerebellum are the most common site).

Option D: Real time-PCR Amplification Detection of rabies virus RNA is highly sensitive and specific. This technique can detect virus in freshsaliva samples, skin, CSF, and brain tissues. However it is not confirmatory test.

Solution to Question 2:

The given life cycle belongs to Fasciola hepatica.

Fasciola hepatica (sheep liver fluke) causes disease primarily in sheep and other domestic animals. Humans are infected by eating watercress (or other aquatic plants) contaminated by larvae (metacercariae) that excyst in the duodenum, penetrate the gut wall, and reach the liver, where they mature into adults.

The adult flukes in the bile ducts produce eggs, which are excreted in the feces. The eggs hatch in freshwater and miracidia enter the snails. Miracidia develop into cercariae, which then encyst on aquatic vegetation. Sheep and humans eat the plants, thus completing the life cycle. Most infections are asymptomatic. Halzoun is a painful pharyngitis caused by the presence of adult flukes on the posterior pharyngeal wall after eating raw sheep liver. Obstructive jaundice can occur months or years later.

Diagnosis is made by identification of the eggs in the feces.

Solution to Question 3:

The given life cycle is of Entamoeba histolytica.

Its life cycle includes 2 stages - the motile trophozoite and the non-motile cyst, both of which are seen in the image shown above. The transmission is primarily by the fecal-oral route. There is no animal reservoir. The most frequent site of systemic disease is the liver.

The ingested cysts differentiate into trophozoites in the ileum which invade the colonic epithelium. The trophozoites penetrate the submucosal layer and form abscesses which break down to form flask-shaped ulcers.

The trophozoites multiply by binary fission and produce cysts, and both stages are passed in the feces. Trophozoites are rapidly destroyed outside the body. Cysts survive in the external environment and are responsible for transmission.

Note: The genus Entamoeba has four morphologically identical species - E. histolytica, E. dispar, E. bangladeshi, and E. moshkovskii. E. moshkovskii which was initially thought to be non-pathogenic has recently been shown to be a causative agent of non-invasive diarrhea.

Solution to Question 4:

Paul Bunnell test is a heterophile antibody reaction test and is used in glandular fever like Infectious Mononucleosis.

Heterophile antibodies have the ability to agglutinate RBCs of different animal species.

Paul Bunnell test uses sheep erythrocytes for agglutination and Monospot test uses horse erythrocytes for agglutination.

In Infectious Mononucleosis, IgM heterophile antibodies are detectable for the first 3 months of infection and they can agglutinate sheep erythrocytes.

Type of test	Name of the test	Test used for
Tube agglutination	Widal	Typhoid
	Standard agglutination test	Brucellosis
	Microscopic agglutination tes t	Leptospirosis
	Paul-Bunnell test	Infectious mononucleosis
	Weil-felix test	Rickettsial infection
	Streptococcal MG agglutinati on test	Primary atypical pneumonia
Antiglobulin test	Direct coomb's test	Detecting fetal RH Antibody
	Indirect Coomb's test	Detecting maternal Rh antib ody
Passive agglutination test	Hemagglutination	Rose Waaler test
	Latex agglutination test	Detection of ASO, CRP, RA f actor, HCG
	Coagglutination test	Diagnosis of legionellosis, Go norrhea

Solution to Question 5:

Sodium hypochlorite, a compound of chlorine, is the best disinfectant for surface blood spills. It is used with a concentration of 0.5-5%.

If the spill contains large amounts of blood or body fluids, clean the visible matter with disposable absorbent material and discard the contaminated materials in appropriate, labelled containment.

A 1:10–1:100 dilution of 5.25%–6.15% sodium hypochlorite (i.e., household bleach) or an EPA-registered tuberculocidal disinfectant has been recommended for decontaminating blood spills.

Solution to Question 6:

Urine lipoarabinomannan (LAM) assay is used for the diagnosis of tuberculosis.

Lipoarabinomannan (LAM) antigen is a lipopolysaccharide present in mycobacterial cell wall. It is released from metabolically active cells or degenerating cells of patients having tuberculosis. Urine LAM testing is especially useful in diagnosing TB in patients with HIV and low CD 4 counts.

Solution to Question 7:

The given slide stained with GMS shows acute-angled (45o) uniform, branching septate hyphae which is suggestive of Aspergillus species.

Aspergillus species grow rapidly, producing aerial hyphae that bear conidia in chains. Long aseptate conidiophores (specialised conidia-bearing hyphae) are seen with terminal vesicles. Phialides (vase-shaped cells producing conidia) with conidia are arranged in basipetal chains on the terminal vesicle. The following image shows the lactophenol cotton blue solution (LPCB) mount of Aspergillus fumigatus showing the long conidiophores with terminal vesicles.



Option A: Mucor produces nonseptate, broad hyphae with aerial sporangiophores which end in a sporangium.

Option B: Penicillium produces septate hyphae with branched, septate and brush-like conidiophores.

Option D: Rhizopus appears similar to Mucor but it has rhizoids with sporangiophores arising in groups directly above the rhizoids.

Solution to Question 8:

Crimean Congo hemorrhagic fever has been found in India.

Four cases were first described in India in 2011 in the state of Gujarat. It is a severe form of viral hemorrhagic fever caused by negative sense ssRNA Nairovirus species. It is zoonotic in origin with Ixodes ticks as vectors with livestock as primary reservoirs. The outbreak was caused due to person-to-person transmission in a tertiary care hospital where the index patient (first identified case) was admitted.

Solution to Question 9:

The given life cycle is of the hepatitis B virus.

Steps:

• The infectious virion attaches to cells and uncoating occurs.

• Cell enzymes in the hepatocytes convert dsDNA to covalently closed circular double-stranded DNA (cccDNA)

• HBV mRNA and pre-genome RNA is synthesized using cccDNA as a template.

• pre-genome RNA is encapsidated by packaging signal into new core particles.

• These new core particles are templates for HBV reverse transcriptase.

• Negative-strand DNA is synthesized by viral polymerase by reverse transcription.

• The cores bud from the pre-Golgi membranes, acquiring HBsAg-containing envelopes.

• Once viral particles are assembled, they either exit the cell or cause reinfection of the same cell.

Solution to Question 10:

1,3 beta D- glucan is helpful for the identification of Invasive candidiasis (invasive fungal infections).

1-3 beta D-glucan assay test detects the presence of (1-3)-β-d-glucan (BDG) which is an abundant cell wall polysaccharide found in most fungi except

• Cryptococci

• Zygomycetes

• Blastomyces dermatitidis

Solution to Question 11:

Relapse is seen in cases of Plasmodium vivax and ovale.

Malaria relapse :

In case of Plasmodium vivax and ovale two kinds of sporozoites are seen, some of which multiply inside hepatic cells to form schizonts and others persist and remain dormant as hypnozoites.

True relapse is caused due to hypnozoites (resting forms). Some get activated to become schizonts and release merozoites, which go on infecting RBCs producing clinical relapse.

In case of Plasmodium falciparum and malariae, the initial tissue phase disappears completely and no hypnozoites are found, hence relapse is not seen.

Solution to Question 12:

The given image is the trophozoite form of the Balantidium coli.

Balantidium coli occurs in two stages

• Trophozoites are oval in shape and contain two nuclei. There is the presence of cilia over the entire surface and are larger around the mouth. Its anterior end is narrow and the posterior end is broad. At the anterior end, there is a groove leading to the mouth (cytostome). Posteriorly, there is a small anal pore (cytopyge).

• Cysts are smaller, round with a micronucleus, a macronucleus and many contractile vacuoles.

Living trophozoites/cysts are yellowish or greenish in color on wet mount.

Solution to Question 13:

The given clinical history and histopathological image are suggestive of Chromoblastomycosis

Chromoblastomycosis (chromomycosis) is a subcutaneous mycotic infection that is usually caused by traumatic inoculation.

Over months to years, the primary lesion becomes verrucous and wart-like with extension along the draining lymphatics.

Histologically, the lesions are granulomatous and the dark sclerotic bodies may be seen within leukocytes or giant cells (medlar/copper penny bodies)

The following are the various dematiaceous species of fungi

• Phialophora verrucosa

• Fonsecaea pedrosoi

• Fonsecaea compacta

• Rhinocladiella aquaspersa

• Cladophialophora carrionii

Treatment: Surgical excision with wide margins is the treatment of choice for small lesions. Chemotherapy with flucytosine or itraconazole may be efficacious for larger lesions. The application of local heat is also beneficial.

Solution to Question 14:

The image shows pink non-branching acid-fast bacilli on a blue background. These findings are consistent with the staining characteristics of mycobacteria. The stain used is the Ziehl-Neelsen stain.

Other options:

Option B: The image below shows Nocardia stained using an acid-fast staining technique. Nocardia is partially acid-fast and appears as branching filaments.



Option C: The image given below is of a gram-stained specimen of Actinomyces israelii. A. israelii is non acid-fast



Option D: The images below show the staining characteristics of Corynebacterium diphtheriae. It is stained with Albert's stain which shows the needle appearance.



Solution to Question 15:

The organisms in the image are leptospira.

In spirochetes, the presence of two axial filaments (periplasmic flagella) with polar insertions are located in periplasmic space called as endoflagella.

Endoflagella are seen in :

• Treponema

• Borrelia

• Leptospira

Solution to Question 16:

The image shows vesicles on the palms, soles, and oral mucosa, consistent with hand-foot-and-mouth disease. It is caused by coxsackievirus.

Hand-foot-and-mouth disease (HFMD) is an exanthematous fever affecting mainly young children, characterized by oral and pharyngeal ulcerations and a vesicular rash of the palms and soles that may spread to the arms and legs. Vesicles heal without crusting, which clinically differentiates them from the vesicles of herpesviruses and poxviruses. It is usually a benign illness resolving in 1-2 weeks.

Coxsackie A16, A9, and B 1-3 are common causative agents.

Other diseases caused by Coxsackie A virus:

• Aseptic meningitis is associated with a maculopapular rash.

• Herpangina

• Minor respiratory infections

Note: Conditions associated with the rash on palms and soles: Secondary syphilis, Hand foot, and mouth disease, Rickettsial feature caused by Rickettsia rickettsiae

Solution to Question 17:

Pneumocystis jiroveci is not a dimorphic fungi, it is a yeast like organism.

Dimorphic fungi :

• Blastomyces dermatitidis

• Histoplasma capsulatum

• Paracoccidioides brasiliensis

• Penicillium marneffei

• Sporothrix schenckii

• Coccidioides immitis

• Coccidioides posadasii

Solution to Question 18:

In most tumors, HBV DNA is integrated into the cell genome, and in most cases, the virus DNA has undergone rearrangements, including deletions. After integration, it promotes the overexpression of cell genes that can also contribute to oncogenesis.

The only viral gene product that is consistently present in the tumor cells is the X protein.

Solution to Question 19:

The above image shows giardia lamblia.

Cysts are resistant forms and are responsible for transmission of giardiasis. Both cysts and trophozoites can be found in the feces (diagnostic stages). The cysts are hardy and can survive several months in cold water.

Infection occurs by the ingestion of cysts in contaminated water, food, or by the fecal-oral route (hands or fomites).

In the small intestine, excystation releases trophozoites (each cyst produces two trophozoites).

Trophozoites multiply by longitudinal binary fission, remaining in the lumen of the proximal small bowel where they can be free or attached to the mucosa by a ventral sucking disk.

Encystation occurs as the parasites transit toward the colon. The cyst is the stage found most commonly in nondiarrheal feces. Because the cysts are infectious when passed in the stool or shortly afterward, person-to-person transmission is possible.

Giardia lamblia exists in :

• Infective form – Quadrinucleate cyst

• Tennis racket shaped trophozoite is actively motile with central axostyle with pair of nuclei

• Oval shaped cyst denoted with cyst wall

Solution to Question 20:

Nucleic acid amplification tests, including PCR assays, for C. difficile toxin A or B gene in stool have been approved for diagnostic purposes of Clostridium difficile infection due to high sensitivity, specificity and rapid turnaround time.

The diagnosis of CDI is based on a combination of clinical criteria:

(1) diarrhea (≥3 unformed stools per 24 h for ≥2 days) with no other recognized cause plus

(2) toxin A or B detected in the stool, toxin- producing C. difficile detected in the stool by PCR or culture, or pseudomembranes seen in the colon.

Other diagnostic methods include :

• Stool Culture in selective media

• Demonstrating Toxin by Hep-2 cell cultures

• Toxin gene identification by Enzyme immunoassay

• Colonscopy or sigmoidoscopy

Despite the array of tests available for C. difficile and its toxins, no single traditional test has high sensitivity, high specificity, and rapid turnaround.

Most laboratory tests for toxins, including enzyme immunoassays, lack sensitivity.

Nucleic acid amplification tests, including PCR assays, for C. difficile toxin A or B gene in stool have now been approved for diagnostic purposes and appear to be both rapid and sensitive while retaining high specificity.

Solution to Question 21:

The given clinical findings and bone marrow examination is suggestive of Histoplasma capsulatum.

It can be grown in sabouraud dextrose agar and blood agar.

In SDA – white cottony mycelia growth can be seen.

Mode of infection – Inhalation of spores

Yeast cells occur within budding yeasts as intracellular fungi

Disseminated infection can occur in immunocompromised patients with reticuloendothelial system being affected the most.

Solution to Question 22:

Adherence is a mechanism of bacteria to attach to host cell for penetration. It is not a mechanism of resistance.

Mechanisms of biofilm-mediated resistance to antibiotics:

• Mechanical barrier: The lipopolysaccharide biofilms prevent entry of water-soluble drugs.

• Efflux pumps: These pumps on the biofilms actively excrete drug out.

• Slow metabolism: Decreased oxygen and nutrients passage through the biofilms slows down the metabolism of bacteria.

• Anaerobic pathway: Because of the above reason bacteria follow anaerobic metabolism.

• Persisters: Some bacteria undergo phenotypic modification into dormant, spore-like structure and survive antibiotic insult. These are called as persisters which are activated later.

• Enzymatic inactivation of drugs by biofilms

• Genetic adaptation to antibiotic insults

• Quorum sensing: It is cell to cell signaling among the bacteria to coordinate among themselves for contribution in biofilm synthesis.

Solution to Question 23:

The egg in the given picture is that of a hookworm. Eggs of both Ancylostoma Duodenale (old world hookworm) and Necator Americanus (new world hookworm) are identical.

The eggs of hookworm are :

• Oval or elliptical

• Colorless

• Not bile stained

• With a thin transparent hyaline shell membrane

• Thin shell

• Segmented ovum - 4 to 8 blastomeres

• There is clear space between segmented ovum and egg shell.

• Eggs float in saturated salt solution.

Humans are infected when filariform larvae in moist soil penetrate the skin.

Solution to Question 24:

The organism shown above is the Nocardia species. The above image shows acid-fast staining, where the organisms are shown appear as acid-fast (stained with fuschia) and branching rods or filaments. They are aerobes found particularly in the soil and are found to produce lung infections in the immunocompromised.

Options A, B: Mycobacteria species are acid-fast but do not have branching filaments.

Option C: Actinomyces species is gram-positive but not acid-fast.

Solution to Question 25:

Staphylococcal septicemia is not toxin mediated.

Septicemia occurs due to the invasion of bacterial organisms into the bloodstream. This causes an inflammatory response to bacterial products, leading to various manifestations seen in sepsis - fever, hypotension, alkalosis.

Toxin mediated staphylococcal infections include food poisoning caused by enterotoxin, toxic shock syndrome caused by toxic shock syndrome toxin and staphylococcal scalded skin syndrome caused by exfoliative/epidermolytic toxin.

Solution to Question 26:

Petechial rash is not a feature of Zika Virus. Though petechiae may rarely appear in Zika virus disease it is however not included in the CDC case definition of Zika virus disease. Petechial rash points towards a diagnosis of Dengue fever while a maculopapular rash suggests a Zika virus

disease.

Zika virus infection is characterised by influenza-like clinical signs including fever, headache and malaise. A maculopapular rash, conjunctivitis, myalgia and arthralgia usually accompany or follow those manifestations.

The complications of Zika virus are:

• Gullian Barre Syndrome

• Fetal microcephaly in pregnant women.

Solution to Question 27:

(1-3) - β - D glucan assay is used for diagnosis of candida infections, invasive aspergillosis and Pneumocystis jirovecii pneumonia.

It is not useful in case of invasive cryptococcosis.

Solution to Question 28:

• CB-NAAT and Line probe assay are rapid molecular diagnostic techniques used for diagnosis of tuberculosis and also to assess resistance to isoniazid and rifampicin.

• CB NAAT (Cartridge Based Nucleic Acid Amplification Test) provides rapid and accurate diagnosis of TB by detecting Mycobacterium tuberculosis and mutations responsible for rifampicin resistance.

• It can be used for sputum samples as well as samples from extrapulmonary sites.

• It is mainly used for diagnosis of DR-TB in presumptive DR-TB patients and TB in key populations such as pediatric population, PLHIV and extrapulmonary TB.

• It is done when smear is negative for TB but chest X-ray is suggestive of tuberculosis and also in cases smear is negative and chest X-ray is not suggestive of TB but there is high index of clinical suspicion.

Diagnostic algorithm for pulmonary TB:


PLHIV
Presumptive TB patient
Smear Examination
CXR
Smear Positive and CXR suggestive of TB
Smear Positive, but CXR not suggestive of TB
Smear Negative but CXR suggestive of TB
Smear Negative or Not Available & CXR not suggestive of TB or not available
Clinical Suspicion High
CBNAAT
PMDT criteria, high MDR settings
MTB detected
MTB not detected or CBNAAT result not available
Consider alternate diagnosis and refer to specialist
Clinically Diagnosed TB
Alternate diagnosis
Rif sensitive
Rif Indeterminate
Rif Resistant
Refer to management of Rif Resistance
Repeat CBNAAT on 2nd sample
Microbiologically Confirmed TB
Indeterminate on 2nd sample, collect fresh sample for Liquid Culture / LPA
*All presumptive TB cases should be offered HIV counseling and testing: however diagnostic work up for TB must not be delayed.

Microbiology AIIMS 2018

Question 1:

A kinyoun positive oocyst (25 µm) was isolated from an HIV patient with diarrhea. Identify the organism.



a) Cystoisospora

b) Cyclospora

c) Cryptosporidium

d) Microspora

Question 2:

The test shown below, which uses a bacitracin disk, is useful in identifying _____.



a) Staphylococcus aureus

b) Streptococcus pyogenes

c) Clostridium botulinum

d) Bacillus

Question 3:

What is the incubation period of lymphogranuloma venereum?

a) 1-5 days

b) 7-10 days

c) 10-30 days

d) 30-60 days

Question 4:

Which is the agent used to treat the infection caused by the cestode egg in the image?



a) Praziquantel

b) Ivermectin

c) Albendazole

d) Diethylcarbamazine

Question 5:

Human herpesvirus 8 is associated with all except _____.

a) Kaposi sarcoma

b) Castleman's disease

c) Adult T-cell leukemia

d) Primary effusion lymphoma

Question 6:

Identify the condenser in the following image.

a) Above the stage

b) Below the stage

c) Near the eyepiece

d) Above the objective lenses

Question 7:

The following test is positive for identification of _____



a) Candida albicans

b) Candida glabrata

c) Candida parapsilosis

d) Candida tropicalis

Question 8:

Choose the correct statement regarding CMV.

a) Ganciclovir resistance is due to mutation in UL97 phosphotransferase gene

b) CMV can be cultured in vitro in any mammalian cell

c) Commonest manifestation of CMV associated disease is watery diarrhea

d) UL54 mutated CMV strains do not show cross resistance to foscarnet

Question 9:

Solvent/Detergent (SD) plasma destroys lipid-enveloped viruses. So, on SD plasma transfusion, which of the following infection is likely to occur?

a) Human Immunodeficiency Virus (HIV)

b) Hepatitis A Virus (HAV)

c) Hepatitis B Virus (HBV)

d) Hepatitis C Virus (HCV)

Question 10:

Plasma sterilization makes use of _____.

a) Hydrogen peroxide

b) Ethylene oxide

c) Gamma rays

d) UV rays

Question 11:

Identify the organism below that causes effacement and attachment of intestinal cells.



a) ETEC

b) EIEC

c) EPEC

d) EAEC

Question 12:

Which of the following organisms is represented by the given form of replication?



a) Hepatitis virus

b) Influenza virus

c) HIV

d) Herpes virus

Question 13:

Which method of diagnosis is shown in the image?



a) Darkfield microscopy

b) Phase contrast microscopy

c) Fluorescent microscopy

d) Electron microscopy

Question 14:

Identify the helminth from the life cycle given below.



a) Ancylostoma duodenale

b) Strongyloides stercoralis

c) Ascaris lumbricoides

d) Enterobius vermicularis

Question 15:

Northern blotting is used for detection of which of the following?

a) DNA

b) RNA

c) Proteins

d) None

Question 16:

A cook prepares sandwiches for 10 people going for picnic. Eight out of them develop severe gastroenteritis within 4-6 hrs of consumption of the sandwiches. It is likely that on investigations the cook is found to be the carrier of:

a) Salmonella typhi

b) Vibrio cholera

c) Entamoeba histolytica

d) Staphylococcus aureus

Question 17:

The following image depicts:



a) Production of monoclonal antibodies

b) Cell lines for viral culture

c) Vaccine synthesis

d) Process of genetic engineering

Question 18:

A patient has complains of fever with chills. A throat examination reveals swollen glands and a thick, whitish membrane over his tonsils. Albert staining is to be done. What are the constituents of Albert's stain?

a) Carbol fuschin and methylene blue

b) Carbol fuschin and malachite green

c) Toluidine and malachite green

d) Toluidine and methylene blue

Question 19:

Which of the following infection presents with mucus in stools and abdominal pain?

a) Giardia

b) Entamoeba

c) Campylobacter

d) Staphylococcus aureus

Question 20:

A patient's plasma has high levels of interferon beta. He most likely has an infection due to _____

a) Virus

b) Bacteria

c) Fungi

d) Mycoplasma

Question 21:

Which of the following is the most potent stimulator of naive T-cells?

a) Mature dendritic cells

b) Follicular dendritic cells

c) B lymphocytes

d) NK cells

Question 22:

Which of the following is an immunologically privileged site?

a) Seminiferous tubule

b) Area postrema

c) Optic nerve

d) Loop of Henle

Question 23:

Bone marrow aspirate of a patient shows the following features. Which of the following is true?



a) Buffy coat is used to diagnose the parasite

b) It is an obligate extracellular parasite

c) It is transmitted from sheep

d) Stibogluconate is the first line of treatment in India.

Question 24:

Which of the following drugs is not used for typhoid fever in adults?

a) Azithromycin

b) Ciprofloxacin

c) Ceftriaxone

d) Amikacin

Answer Key

Question No.	Correct Option
1	a
2	b
3	c
4	a
5	c

6	b
7	a
8	a
9	b
10	a
11	c
12	b
13	a
14	b
15	b
16	d
17	a
18	c
19	b
20	a
21	a
22	a
23	a
24	d

Detailed Explanations

Solution to Question 1:

Diarrhea in an HIV patient is suggestive of a coccidean parasitic infestation. Kinyoun stain positive oocysts in the image are identified as Cystoisospora.

Points in favor:

• HIV patient with diarrhea

• Oocysts are long and oval-shaped, which stain pink to red with Kinyoun staining.

Comparison of coccidian parasites causing diarrhea in an immunocompromised host:

Property	Cryptosporidium	Cyclospora	Cystoisospora
Sporulated oocyst	4-6 µm, round, contains f our sporozoites	8-12 µm, round, contains 2 sporocyst each having t wo sporozoites	23-36 µm, oval, contain t wo sporocysts, each havin g four sporozoites.
Acid-fast detection	Uniformly acid-fast	Variable acid-fast	Uniformly acid-fast
Sporulation of the ooc yst	Occurs inside the host cell	Occurs in soil	Occurs in soil

Property	Cryptosporidium	Cyclospora	Cystoisospora
Diagnostic form	Sporulated cyst	Unsporulated cyst	Unsporulated oocyst



Modified acid-fast staining (like Kinyoun stain) showing oocysts of Cyclospora cayetanensis.



Modified acid-fast staining (like Kinyoun stain) showing oocysts of Cryptosporidium species.

In immunocompromised patients, the infection causes severe diarrhea similar to cholera and require hospitalization.

The diagnosis is made with stool microscopy. The cyst wall and sporoblasts are dyed red with the Ziehl–Neelsen (warm) or Kinyoun (cold) staining methods. Both methods have good specificity and therefore are useful as confirmatory tests.

The treatment is fluid replacement and cotrimoxazole.

Solution to Question 2:

The given image shows blood agar culture with bacitracin disk with a zone of inhibition around the disk, which means that the organism is sensitive to bacitracin.

Among the options given, Streptococcus pyogenes is susceptible to bacitracin.

Streptococci belonging to group A are identified (presumptively) by inhibition of growth by bacitracin.

Solution to Question 3:

The incubation period of Lymphogranuloma venereum ranges from 10-30 days.

Chlamydia trachomatis serovars L1, L2, and L3 cause lymphogranuloma venereum (LGV). The most common presentation is the inguinal syndrome, which is characterized by painful inguinal lymphadenopathy.

Solution to Question 4:

The egg in the image is of Hymenolepis nana (dwarf tapeworm) and it is treated by praziquantel (5–10 mg/kg once).

The egg is oval, colorless, non-bile stained, and has a single shell with two distinct membranes. Inner membrane has two small knobs or poles from which 4 to 8 polar filaments arise and spread out between the two membranes. It is embryonated and the oncosphere has six hooks (three pairs of hooklets).

Life cycle of Hymenolepis nana:



Hymenolepiasis is the most common of all the cestode infections and is usually asymptomatic.

Solution to Question 5:

Adult T-cell leukemia is caused by HTLV-1 (and not HHV8).

Human Herpesvirus 8 (HHV8), also known as Kaposi sarcoma-associated herpesvirus (KSHV), is lymphotropic virus and is closely related to Ebstein–Barr virus (EBV).

Diseases caused by HHV:

• Kaposi sarcomas

• Vascular tumors of mixed cellular composition

• Body cavity-based lymphomas occurring in AIDS patients (Primary effusion lymphoma)

• Multicentric Castleman disease

Solution to Question 6:

This is a modern compound microscope, and the condenser is located below the stage.

The purpose of the condenser lens is to focus the light from the illuminator onto the specimen. Microscopes with a stage condenser lens render a sharper image than those with no condenser lens.

The simple microscope was invented first by Antonie van Leeuwenhoek, the father of microscopy and microbiology. It uses 1 lens for magnification. The compound microscope, introduced by Hans and Zacharias Janssen uses 2 lenses, an objective lens and an eyepiece, to provide better magnification power than the simple microscope.

Solution to Question 7:

The image shows the germ tube test used for identifying Candida albicans.

Candida species, when cultured, grow as oval, budding yeast cells with pseudohyphae. Unlike other species of candida, Candida albicans can also produce true hyphae or germ tubes. This can be demonstrated after incubation in serum maintained at 37°C for about 90 minutes. This is known as the Reynolds-Braude phenomenon. This test is positive for Candia dubliniensis as well. Another distinguishing feature of C. albicans is that it produces large, spherical chlamydospores on nutritionally deficient media.

Solution to Question 8:

Resistance to ganciclovir most commonly occurs due to mutations in the UL97 phosphotransferase gene.

Cytomegalovirus (CMV) is a beta-herpes virus. It is the most common congenital infection. In adults, its disease is mostly subclinical (option C). An apparent CMV infection resembles infectious mononucleosis (fever, sore throat, enlarged lymph nodes and splenomegaly). It is the most common infection in the donor organ seen 1-4 months after solid organ transplantation. In immunosuppressed hosts, it causes pneumonia and colitis.

Diagnosis is by PCR assays. Human CMV replicates in vitro only in human fibroblasts (option B) and can thus be cultured in them, although the virus is often isolated from epithelial cells of the host. The cultured fibroblasts show owl-eye inclusion bodies after 2-3 weeks.

UL54 (DNA polymerase) gene mutations can confer CMV resistance to ganciclovir, cidofovir and foscarnet (option D).

Solution to Question 9:

Solvent detergent plasma is a form of blood plasma made from plasma collected from many people which is then processed with solvents to try to get rid of viruses. HAV is a non-enveloped virus, and hence it’s not affected by solvent/detergent (SD) plasma.

Only the viruses with an envelope are susceptible to SD plasma destruction.

Hepatitis B, C, and HIV are enveloped.

Solution to Question 10:

A variety of hydrogen peroxide gas–based processes that use plasma as part of the sterilization process are known as "plasma sterilization."

The mechanisms of antimicrobial action in these processes are primarily due to hydrogen peroxide gas (or condensed gas) and not plasma, although the plasma generation phase of these processes is known to assist in the breakdown of residual peroxide in the chamber load.

Solution to Question 11:

Attachment and effacement lesions on intestinal cells are caused by enteropathogenic Escherichia coli (EPEC).

E. coli is the second most common cause of diarrhea in the pediatric age group. Rotavirus is the most common cause. EPEC causes watery diarrhea which is self-limiting.

EPEC adheres to the mucosal cells of the small bowel and causes loss of microvilli (effacement). This leads to the formation of filamentous actin pedestals or cuplike structures and occasionally may cause entry of the EPEC into the mucosal cells.

Pathogenicity requires the presence of two important factors:

- Bundle-forming pilus encoded by plasmid containing EPEC adherence factor (EAF).
- Chromosomal locus of enterocyte effacement (LEE) pathogenicity island.



The image below shows an electron microscopy image of E Coli.



Solution to Question 12:

The given image shows generation of mRNA from negative sense RNA within the nucleus. This occurs during the replication of influenza virus.

Orthomyxoviruses (influenza virus) contain segmented (8 segments) negative sense ssRNA that is replicated in the nucleus. Stages of viral replication include the following:

• Attachment, penetration, uncoating: The virus attaches to cell-surface sialic acid via hemagglutinin. It is then internalized by receptor-mediated endocytosis. There is fusion of the viral envelope and cell membrane, resulting in uncoating or release of nucleocapsid into the host cell cytoplasm.

• Transcription and translation: mRNAs are produced from nucleocapsids via a virus-encoded polymerase. This process occurs in the nucleus. The mRNAs thus produced are translated into viral proteins in the cytoplasm.

• Viral RNA replication: The viral genome is replicated by the same virus-encoded polymerase involved in transcription. The negative sense RNA is converted to its positive sense form. The positive sense copies serve as templates for replication of the viral genome.

• Maturation: The virus matures by budding from the surface of the cell.

Solution to Question 13:

The given image is of darkfield microscopy as it has a dark background with a refractile organism.

The dark ground microscope has an optical system that enhances the contrast of unstained bodies.

Dark ground microscopy is the most specific and sensitive technique to diagnose syphilis when an active chancre or condyloma lata is present to visualize Treponema pallidum.

Solution to Question 14:

The image shows the life cycle of Strongyloides stercoralis.

Strongyloides stercoralis is the smallest nematode known to cause human infection. It has both a free-living and a parasitic life cycle. Its natural host is man. The infective form to man is the filariform larva. Infection occurs through penetration of the skin by larvae in the soil. Then, the larvae migrate to the small intestine. They molt twice and become adult female worms in the small intestine. The eggs laid in the mucosa of the intestine hatch immediately releasing the rhabditiform larva (L1 stage). Thus, it is ovoviviparous. These rhabditiform larvae in the bowel can develop directly into filariform larvae that penetrate the colonic wall or perianal skin and enter the circulation to repeat the migration. This establishes ongoing internal reinfection.

The rhabditiform larva can also be passed in the stool. It may directly metamorphose to the infective filariform larva in the soil, or develop into adult worms. The free-living adult worms then pass eggs, which hatch and mature to form infective filariform larvae.

In Ancylostoma duodenale infestation, eggs containing segmented ova (not rhabditiform larvae) are passed in the stool. The rhabditiform larva hatches 2 days after the eggs are passed in the soil.

Note: Based on the method of reproduction, nematodes are classified as follows:

• Oviparous - They lay eggs:

• Unsegmented eggs: Ascaris lumbricoides, Trichuris trichuria

• Segmented eggs: Ancylostoma duodenale, Necator americanus

• Eggs containing larva: Enterobius vermicularis

• Viviparous - They produce larvae. e.g., Trichinella spiralis, Wuchereria bancrofti, Brugia malayi

• Ovoviviparous – They lay eggs containing fully formed larvae that hatch immediately. e.g., Strongyloides stercoralis

Solution to Question 15:

Northern blotting is used for the detection of RNA.

Southern blotting is used for the detection of DNA and Western blotting is used for the detection of protein.



Solution to Question 16:

The cook is most likely to be carrier of Staph. aureus.

The onset of food poisioning with in 4-6 hrs of food is mainly due to:

• Staphylococcus aureus

• Bacillius cereus (has no carriers)

Solution to Question 17:

The image depicts the production of monoclonal antibodies by hybridoma technology.

The following are the steps involved in hybridoma technology:

• Mice are immunized with an antigen. After several weeks of immunization, blood samples are obtained from mice for measurement of serum antibodies.

• When the antibody titer is high enough, mice are commonly boosted by injecting antigen, without adjuvant, intraperitoneally or intravenously. Then the mice are euthanized and their spleens removed for in vitro hybridoma cell production.

• Fusing antibody-producing spleen cells, which have a limited life span, with myeloma cells (derived from an immortal tumor of lymphocytes) results in a hybridoma that is capable of unlimited growth.

• The cells are cultured in a medium containing hypoxanthine, aminopterin, and thymidine (HAT medium) which allows only the fused cells to survive in culture.

• Monoclonal antibodies are then harvested.



Solution to Question 18:

The given scenario is suggestive of diphtheria caused by Corynebacterium diphtheria. This organism is stained using Albert's stain which contains toluidine blue 'O' and malachite green. Both these dyes have an affinity for acidic cytoplasm as they are basic dyes.

Albert's, Neisser's, and Ponder's stain are used to demonstrate the metachromatic granules(Volutin / Babes Ernst granules) of Corynebacterium diphtheria.

Corynebacterium diphtheria appears as green-coloured bacilli with purple-blue-coloured metachromatic granules in cuneiform arrangement.



Solution to Question 19:

Stools with mucus and abdominal pain are seen in infection due to Entamoeba.

Clinical features of intestinal amebiasis:

• Acute intestinal amebiasis presents as dysentery accompanied by lower abdominal discomfort and flatulence. Stools usually have blood and mucus.

• Chronic amebiasis presents with low-grade symptoms such as occasional diarrhea, weight loss, and fatigue.

• A granulomatous lesion called an ameboma may form in the cecal or rectosigmoid areas of the colon which can resemble an adenocarcinoma of the colon.

Other options:

Option a - Giardia is often asymptomatic, but in some cases can lead to fat malabsorption, dull epigastric pain, and flatulence.

Option c - Campylobacter presents with fever, pain and bloody or watery diarrhea. Stool contains leucocytes and blood.

Option d - Staphylococcal food poisoning is toxin-mediated and presents with nausea, vomiting, and diarrhea.

Solution to Question 20:

Interferon-beta is secreted during infection due to viruses.

Interferons are classified as cytokines. Interferons are non-toxic, non-antigenic and species-specific molecules. They are inactivated by proteases. They are stable at pH over 2-10, except INF-γ (labile at pH 2) and are non-dialysable and non-sedimentable.

There are three types of interferons - α, β and γ. IFN-α or leukocyte interferon is produced by leucocytes. It has antiviral activity, is immunoregulaotry and antiproliferative. IFN-β or fibroblast interferon is produced by fibroblasts and epithelial cells. It has antiviral activity and is thus increased in viral infections. It enhances MHC class 1 and NK cell activity. IFN-γ or immune interferon is produced by T-lymphocytes, NK cells . It activates macrophages, increases neutrophil function, enhances both MHC class 1 and class II expression and has anti-proliferative (anti-tumour) functions.

Solution to Question 21:

Mature dendritic cells are the most potent stimulator of naive T cells.

Naive CD4+ T cells are activated when they come in contact with two ligands - MHC class II and a costimulary membrane protein (eg: CD28). Both of these are present in antigen-presenting cells, the most potent of which is the mature dendritic cell. Other professional antigen-presenting cells include macrophages and B lymphocytes (option C).

Other options:

Option B: Follicular dendritic cells are functionally different from dendritic cells. They are not antigen-presenting-cells and do not activate naive T cells.

Option D: NK cells do not stimulate naive T cells.



Solution to Question 22:

The seminiferous tubules are an immunologically privileged site.

Immunologically privileged sites can tolerate the introduction of antigens without eliciting an immune response. This allows a higher likelihood of graft survival, even without perfect HLA matching. As a result, an allograft can be placed in these sites without endangering a rejection reaction.

The unique immune tolerance in these regions is primarily due to the lack of effective lymphatic drainage, which prevents antigen presentation to the immune system, and the presence of blood-tissue barriers that restrict immune cell infiltration. Low vascularity contributes to immune privilege by minimising antigen exposure and immune cell recruitment.

Examples of immune-privileged sites are the anterior chamber of the eye, cornea, testes, and brain.

Note: Although part of the brain, the blood-brain barrier is absent at the area postrema, therefore it is not an immune-privileged site.

Solution to Question 23:

The given image shows the amastigote form of leishmaniasis located intracellularly in macrophages. Buffy coat is used to diagnose the parasite.

Visceral leishmaniasis (kala-azar) is a slowly progressing indigenous disease caused by a protozoan parasite of genus Leishmania. Leishmaniasis is a parasitic disorder transmitted by the

bite of an infected female Phlebotomus sandfly in developing countries. The disease is transmitted by sand flies, which inoculate the flagellated promastigotes into the skin of the host.

In human beings, the disease presents in four different forms with a broad range of clinical manifestation:

• Visceral leishmaniasis, or kala-azar,

• Cutaneous leishmaniasis,

• Mucocutaneous leishmaniasis, and

• Diffuse cutaneous leishmaniasis.

The life cycle of Leishmania involves two forms,

• The promastigote which develops and lives extracellularly in the sandfly vector and

• The amastigote which multiplies intracellularly in the reticuloendothelial cells of the host.

It lowers immunity, causes persistent fever, anemia, liver and spleen enlargement, and if left untreated, death may occur due to opportunistic infection.

Diagnosis:

• Rapid diagnostic test: rK39 Rapid diagnostic test and Kala azar dipstick test strip.

• Parasitological diagnosis: Demonstration of LD bodies in buffy coat smear - confirmatory.

• Aldehyde test of Napier: Widely used for surveillance.

• Serological tests for diagnosis: Rapid dipstick test, Direct agglutination test, ELISA, Indirect Fluorescent Antibody Test

• Leishmanin (Montenegro) test: Induration of 5mm or more is considered positive, usually positive 4-6 weeks after onset in cutaneous leishmaniasis, valuable for distinguishing immune from nonimmune subjects and endemicity.

Drugs used in the treatment of visceral leishmaniasis are sodium stibogluconate, miltefosine, and amphotericin. Liposomal amphotericin single dose 10mg/Kg IV is the first line of treatment in India.

Solution to Question 24:

Amikacin is not used in the treatment of typhoid fever.

Amikacin is an aminoglycoside. Despite showing efficient in vitro killing of Salmonella, first- and second-generation cephalosporins and aminoglycosides have proven to be ineffective in actual clinical cases.

Antibiotic therapy for enteric fever in adults			
Indication	Agent	Dosage (Route)	Duration, Days
Empirical Treatment			

Antibiotic therapy for enteric fever in adults			
	Ceftriaxone	2 g/d (IV)	10-14
Azithromycin	1 g/d (PO)	5	
Fully Susceptible			
Optimal treatment	Ciprofloxacin	500 mg bid (PO) or 400 mg q12 h (IV)	5-7
Azithromycin	1 g/d (PO)	5	
Alternative treatment	Amoxicillin	1 g tid (PO) or 2 g q6h (IV)	14
Chloramphenicol	25 mg/kg tid (PO or IV)	14-21	
Trimethoprim-sulfamethoxazole	160/800 mg bid (PO)	7-14	
Multidrug-resistant, fluroquinolone-susceptible			
Optimal treatment	Ceftriaxone	2 g/d (IV)	10-14
Azithromycin	1 g/d (PO)	5	
Alternative treatment	Ciprofloxacin	500 mg bid (PO) or 400 mg q12h (IV)	5-14
Quinolone-resistant			
Optimal treatment	Ceftriaxone	2 g/d (IV)	10-14
Azithromycin	1 g/d (PO)	5	
Alternative treatment	High dose ciprofloxacin	750 mg bid (PO) or 400 mg q 12h (IV)	10-14
Ceftriaxone-resistant			
Optimal treatment	Meropenem	1 g q8h (IV)	10-14
Azithromycin	1 g/d (PO)	5	

Microbiology AIIMS 2019 (May & Nov)

Question 1:

A farmer from the sub-Himalayan belt had a thorn prick injury a year back and developed a warty, verrucous growth at the site of prick. A skin biopsy sample was taken and sent for histopathological examination. The characteristic microscopic body as shown in the image was seen. Which of the following could be the most likely diagnosis?



a) Blastomycosis

b) Pheohyphomycosis

c) Chromoblastomycosis

d) Sporotrichosis

Question 2:

Choose the wrong statement about Plasmodium species:

a) Drug resistant falciparum is seen in India

b) Presence of duffy blood group antigen is protective against vivax

c) Severe malaria with shock is called algid malaria

d) Relapse is seen in P. ovale

Question 3:

A HIV patient with fever, chronic diarrhea presented to OPD. His stool was examined which showed the following oocysts. Identify the components used for staining the stool.



a) Crystal violet, iodine, acid alcohol, carbol fuchsin

b) Methanol, Kinyoun's carbol fuchsin, acid alcohol, methylene blue

c) Methanol, dilute carbol fuchsin, acid alcohol, crystal violet

d) Methylene blue, iodine, acid alcohol, carbol fuchsin

Question 4:

Which immunoglobulin is elevated in a case of chronic allergy?

a) IgA

b) IgM

c) IgE

d) IgG

Question 5:

A patient is diagnosed with Hepatitis B infection. Which of the following markers will always be present in his serum, even during recurrences, or if he develops chronic hepatitis?

a) HBc Ag

b) Anti HBs Ab

c) Anti HBc Ab

d) HBe Ag

Question 6:

Which among the following occupation is a risk factor for this presenting illness?

a) Kennel Worker

b) Lifeguard in a swimming pool

c) Veterenarian

d) Poultry worker

Question 7:

Transmission assessment survey is carried out for:

a) P. vivax

b) P. falciparum

c) Filariasis

d) Leishmania

Question 8:

Which among the following is a subcutaneous fungal agent?

a) Cryptococcus neoformans

b) Histoplasma capsulatum

c) Sporothrix schenckii

d) Talaromyces marneffei

Question 9:

A hospital has reported an outbreak of MRSA infection. On investigation, it was found out that a staff nurse had nasal carriage of MRSA. Which of the following drugs helps in the removal of colonization of bacteria?

a) Oral vancomycin

b) Inj cephalosporin

c) Topical mupirocin

d) Inhaled colistin

Question 10:

Mw vaccine is extracted from which bacteria, mentioned below:

a) M. indicus pranii

b) M. welchii

c) M. leprae

d) M. bovis

Question 11:

Which toxin acts via the mechanism depicted in the image below?



a) Botulinum

b) Tetanus

c) Diphtheria

d) Pertussis

Question 12:

The toxin produced by the organisms shown below acts via?



a) Increase in cAMP

b) Increase in cGMP

c) ADP ribosylation of EF2

d) RNA glycosidase

Question 13:

A 25-year old man presents with a history of urethral discharge for the last three days. A gram stain smear of the discharge is shown below. All of the following are true regarding the likely etiology except?



a) Catalase and oxidase positive

b) Pili is a virulence factor

c) Obligate intracellular organism

d) Shows twitching motility

Question 14:

The CLED medium is preferred over MacConkey agar for the culture of organisms in case of UTI because:

a) It inhibits proteus swarming

b) It promotes the growth of candida, streptococcus and staphylococcus

c) It differentiates between lactose-fermenting and non-lactose fermenting bacteria

d) It promotes the growth of pseudomonas

Question 15:

Which of the following organisms cause multiple alveolar-like lesions in the liver?

a) Echinococcus multilocularis

b) Echinococcus granulosus

c) Entamoeba histolytica

d) Cysticercus cellulosae

Question 16:

All of the following can be seen intracellularly in hepatocytes except:

a) Toxoplasma

b) Plasmodium

c) Babesia

d) Leishmania

Question 17:

A Giemsa stain of a thin peripheral smear is prepared. Which of the following cannot be diagnosed?

a) Coxiella burnetii

b) Toxoplasma gondii

c) Bartonella henselae

d) Ehrlichia chaffeensis

Question 18:

Which of the following is false about influenza?

a) Virus shedding is present before symptoms begins

b) 5-12 years is a high risk group

c) Aquatic birds act as reservoir

d) Secondary attack rate is 5-15%

Question 19:

β 1-3 glucan assay is used to diagnose all except?

a) Candidiasis

b) Aspergillosis

c) Pneumocystis pneumonia

d) Mucormycosis

Question 20:

Which of the following is a xenodiagnostic method?

a) Rabbit ileal loop assay for ETEC

b) Injecting Aedes thorax with blood of a patient with suspected dengue

c) Injecting a hamster with splenic biopsy for diagnosis of leishmaniasis

d) Intradermal test on guinea pigs for toxigenicity of Corynebacterium diphtheria

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	c
5	c
6	a
7	c
8	c
9	c
10	a
11	a
12	c
13	c
14	b
15	a

16	c
17	a
18	b
19	d
20	b

Detailed Explanations

Solution to Question 1:

The history of a slowly spreading warty, verrucous lesion following a thorn prick injury and the histopathological image showing medlar bodies is suggestive of chromoblastomycosis.

Chromoblastomycosis (chromomycosis) is a subcutaneous mycotic infection that is usually caused by traumatic inoculation. Over months to years, the primary lesion becomes verrucous and wart-like with extension along the draining lymphatics.

Histologically, the lesions are granulomatous and the dark sclerotic bodies may be seen within leukocytes or giant cells called medlar/copper penny bodies.

Subcutaneous mycoses	Causative agent/s
Sporotrichosis	Sporothrix schenckii
Chromoblastomycosis	Phialophora verrucosaFonsecaea pedrosoiFonsecaea compactRhinocladiella aquaspersa Cladophialophora carrionii
Phaeohyphomycosis	Exophiala jeanselmeiPhialophora richardsiaeBipolaris spiciferaWangiella dermatitidis
Mycetoma	Pseudallescheria boydii (Anamorph Scedosporium apiospermum).Madurella mycetomatisMadurella griseaE jeanselmeiAcremonium falciforme

Solution to Question 2:

The absence, not the presence, of the Duffy blood group antigen, is protective against P. vivax because the Duffy antigen normally serves as the receptor for Plasmodium vivax.

Drug Efflux pumps are the primary mode of resistance for Chloroquine. It is mediated by an ABC transporter encoded by P. falciparum multidrug resistance gene 1 (Pfmdr1). The resistance of P. falciparum to chloroquine increased over time and is present across all regions of the country.

Algid malaria is a rare complication of severe malaria characterized by hemodynamic collapse manifesting as shock with pronounced metabolic changes and hypothermia.

Relapse is seen in cases of Plasmodium vivax and Plasmodium ovale.

Malaria relapse :

True relapse is caused due to Hypnozoites.

These are resting forms.

In the case of Plasmodium vivax and ovale two kinds of sporozoites are seen, some of which :

• multiply inside hepatic cells to form - Schizonts

• others persist and remain dormant as - hypnozoites

From time to time some are activated to become schizonts and release merozoites, which go on infecting RBCs producing clinical relapse.

In the case of Plasmodium falciparum and Plasmodium malariae, the initial tissue phase disappears completely and no hypnozoites are found, hence technically relapse is not seen.

Solution to Question 3:

Diarrhea in an HIV patient is suggestive of a coccidian parasitic infestation. The image shows a blue background and red-stained oocysts, suggestive of modified acid-fast stain positive Cryptosporidium parvum.

The modified Ziehl-Neelsen (ZN) method is used for staining cryptosporidium and isospora species. Unlike the ZN-stain, the modified ZN stain does not require heating of the reagents.

The reagents used in modified ZN stain include:

• Absolute methanol for fixation

• Kinyoun's Carbol fuchsin - Primary stain

• Acid alcohol - Decolourising agent

• Methylene blue or Malachite green - Counterstain

Solution to Question 4:

IgE is elevated in cases of chronic allergy.

Solution to Question 5:

A patient who has had exposure to HBV will have anti-HBc antibody even when he has moved into the chronic or recurrent stages. IgM anti-HBc reappears during acute recurrences while IgG anti-HBc is found in chronic states.

The HBsAg antigen is present in chronic hepatitis B, while the antibody anti-HBs is not.

Solution to Question 6:

The above image shows cutaneous larva migrans. Being a kennel worker is a risk factor for this job as exposure to animal feces is high.

As Larvae are found on sandy beaches, and under dwellings, swimming in natural water bodies would be a risk and not artificial ones like swimming pools.

CLM is caused by various animal hookworm species (e.g., Ancylostoma brazilienses), and the parasite is acquired from direct skin contact with soil contaminated by dog or cat feces. The parasitic infection usually begins with an inflamed papule, often located between the toes, where the hookworm species (e.g., Ancylostoma brazilienses) larvae enter the skin.

Serpentine or linear single-track lines later mark the course of the larvae as they migrate through the epidermis of the trunk and lower extremities.

Ivermectin is the first-line drug for the treatment of cutaneous larva migrans, taken as a single 200 μg/kg oral dose.

Solution to Question 7:

Transmission assessment survey is carried out for filariasis.

Mass Drug Administration is used to reduce infection in the community to levels below a threshold at which mosquitoes are unable to continue spreading the parasites from person to person and thus new infections are prevented.

The transmission assessment survey (TAS) is used to determine when infections have been reduced below target thresholds and Mass Drug Administration can stop. Once MDA has stopped, TAS is used as a surveillance tool to determine that infection levels are sustained below target thresholds.

Solution to Question 8:

Sporothrix schenckii is a subcutaneous fungal agent. The other three mentioned are systemic mycoses.



Solution to Question 9:

Topical mupirocin is the preferred agent for the elimination of nasal carriers of MRSA.

As Mupirocin has no structural similarities with existing systemic antibiotics that might lead to the development of cross-resistance and has minimal toxicity in its topical form, it is approved for the prevention of MRSA outbreaks

Mixed therapy with systemic agents is not preferred as it would cause unnecessary antibiotic exposure potentially paving way for drug resistance.

Solution to Question 10:

Mw vaccine is extracted from M. indicus pranii

MIP immunotherapy is used as an adjunct to chemotherapy for category II tuberculosis.

Here, indicus stands for India, where the M. indius pranii was isolated; and pranii stands for the person who discovered the species (Prof. Pran Talwar) and National Institute of Immunology (nii), where trials were conducted (pran + nii).

Administration: 0.1 ml intradermal. It is divided into two equal doses and given on both arms. Booster doses are injected only on one arm.

Solution to Question 11:

The image given above depicts the mechanism of action of botulinum toxin.

It causes proteolysis of snare proteins. Snare proteins have three components- synaptobrevin, SNAP25 and syntaxin. The snare protein is required for the excocytosis of acetylcholine vesicles. Thus it leads to presynaptic inhibition of neuronal transmission.

There are 7 different strains of clostridium botulinum and each strain acts by a different mechanism. Toxins of C. Botulinum A, C and E cleave SNAP25. Type C also cleaves syntaxin. Type B, D, F and G cleave synaptobrevin.

Solution to Question 12:

The given image shows gram-positive rods with club-shaped ends, beaded appearance, and a Chinese-letter pattern of arrangement indicative of Corynebacterium diphtheriae.

The toxin produced by the bacteria - diphtheria toxin - inhibits polypeptide chain elongation and protein synthesis by inactivating the elongation factor-2 (EF-2) by ADP ribosylation. It has two fragments - fragment B helps in binding of the toxin to host cells and fragment A inactivates EF-2.



Solution to Question 13:

The given slide shows gram-negative diplococci within the polymorphs, characteristic of Neisseria gonorrhoeae infection. Neisseria is not an obligate intracellular organism, it can also occur extracellularly i.e, it is a facultative intracellular organism. The other neisseria that are found as normal inhabitants of the human respiratory tract occur extracellularly.

Neisseria gonorrhoeae is a gram-negative, non-sporulating, non-motile diplococcus that is aerobic but may grow anaerobically also. It is oxidase and catalase-positive.

Although Neisseria are non-motile organisms, they show twitching motility. This is a flagella-independent form of bacterial translocation over moist surfaces that occurs by the extension and retraction of polar type IV pili. These pili act as virulence factors by promoting attachment to host cells and inhibiting phagocytosis.

Virulence factors for gonococci:

• Outer Membrane Proteins (OMPs)

• IgA proteases

• Pili

• Lipo-oligosaccharide (LOS)

• Transferrin

• Lactoferrin-binding protein

Solution to Question 14:

CLED (Cysteine lactose electrolyte-deficient) medium is preferred over MacConkey agar for the culture of organisms involved in urinary tract infections because it is less inhibitory and promotes the growth of gram-positive bacteria (except β hemolytic streptococcus) and candida.

A combination of blood agar and MacConkey agar is used for the processing of urine specimens. CLED medium is used as an alternative for this combination. It can also prevent swarming of Proteus, and is thus different from blood agar. Like MacConkey agar, it differentiates between lactose fermenters and non-fermenters. Lactose fermenters change the colour of the medium from green to yellow. However, these two facts are not the reason it is better than MacConkey agar for evaluation of UTI.



Solution to Question 15:

Echinococcus multilocularis causes multiple alveolar-like lesions in the liver.

E. multilocularis leads to the formation of a multilocular hydatid cyst, most commonly affecting the liver. It presents as multiple small cysts in the liver.

Echinococcus granulosus causes a hollow fluid-filled unilocular cyst in the liver. The CT below shows water lily sign, which is seen in the hydatid disease of the lung. It is caused by Echinococcus granulosus. (Option B)



The water-lily sign occurs due to the detachment of the endocyst membrane from the cyst wall, giving the appearance of floating membranes within the pericyst. This mimics the appearance of a water lily. It is classically visualised on plain radiographs (mainly chest X-ray) when the collapsed membranes are calcified but may be seen on ultrasound, CT, or MRI.

Solution to Question 16:

Babesia is an intracellular parasite of RBCs, not of the hepatocytes.

Its trophozoites invade the RBCs.

The liver is usually not affected as babesia lack an exoerythrocytic phase.

Solution to Question 17:

The diagnosis of Coxiella burnetii cannot be done using a Giemsa stain of a thin peripheral smear.

It is done mainly by serology, which shows a rise in the titer of specific antibodies to C. burnetii.

Microorganisms such as Histoplasma, Leishmania, Toxoplasma, and Pneumocystis can also be detected with Giemsa, and in gastric tissues, Helicobacter pylori (H. pylori) appear thin and distinctly blue. Giemsa's stain is frequently used for diagnostic purposes in haematology to differentiate the nuclear and cytoplasmic morphology of platelets, RBCs, WBCs, and parasites.

Solution to Question 18:

Children <5-years, especially those <2-years of age are at the highest risk of serious complications related to influenza infection.

Other high-risk groups for influenza include adults ≥60 years of age, persons with chronic diseases, persons who are immunocompromised.

Option A: The virus is shed in respiratory secretions a day before the onset of illness and for 3-4 days thereafter.

Option C: Birds, especially aquatic birds act as reservoirs of influenza viruses.

Option D: Secondary attack rate is 5- 15%.

Solution to Question 19:

The β-1-3 glucan assay cannot be used to diagnose mucormycosis. This condition is usually diagnosed based on clinical features and culture.

The β-1-3 glucan assay detects the presence of (1-3)-β-d-glucan (BDG), an abundant cell wall polysaccharide found in most fungi. It is a chromogenic, quantitative enzyme immunoassay (EIA) designed to detect beta-D-glucan by using purified, lysed horseshoe crab (Limulus polyphemus) amebocytes. 1,3 Beta D Glucan level of ≥80 pg/mL is considered a positive result.

Fungi that do not contain BDG in their cell walls include Cryptococcus neoformans, zygomycetes like rhizopus and mucor and Blastomyces dermatitidis. Mucormycosis is an opportunistic mycosis caused by zygomycetes, which do not contain (1-3)-β-d-glucan, and hence, the β 1-3 glucan assay cannot be used for its diagnosis.

Solution to Question 20:

Injecting Aedes thorax with blood of a patient with suspected dengue is the xenodiagnostic method among the given options.

Xenodiagnosis (xenos- foreign) is an indirect method of identifying the infectious disease agents (i.e. the pathogen). In xenodiagnosis, the vector of a pathogenic agent is infected and the pathogen is allowed to multiply in the vector. The vector acts as the “foreign” host. Later, the vector is examined for the presence of the pathogen.

This method was first used by Brumpt for the diagnosis of Chagas disease. Laboratory bred Reduviid bugs (vector/ foreign host) are allowed to feed on the suspected patients (original host). After 4-5 weeks, the feces of the bugs are microscopically examined for amastigote forms of Trypanosoma cruzi (pathogen).

Another method of xenodiagnosis is the intrathoracic inoculation of the mosquitoes (vector) with viruses (pathogen). The virus multiplies in different organs of the mosquito and can be easily demonstrated. Aedes aegypti, Aedes albopictus, and Toxorhynchites mosquitoes are used for the identification of the dengue virus.

In Trichinella spiralis infection, muscle biopsy bits from suspected patients are fed to laboratory rats (vector for T. spiralis). The larvae of T. spiralis can be demonstrated in the muscles of the

infected lab rats.

Advantage of xenodiagnosis: It is useful in cases where a direct microscopic diagnosis is difficult due to the low count of pathogens in the blood of the host/patient.

Other options:

Option A: Rabbit ileal loop assay for ETEC is a method of detection of toxigenicity/ virulence testing.

Option C: Injecting a hamster with splenic biopsy for diagnosis of leishmaniasis is a method of animal inoculation.

Option D: Intradermal test on guinea pigs for toxigenicity of Corynebacterium diphtheria is a method of virulence testing.

Microbiology AIIMS 2020 (May and Nov INI-CET)

Question 1:

A resident had a needle stick injury while dealing with a Hepatitis B positive patient. Serology was done and HBsAg was negative and the Anti-HBs was >105 mIU/ml. What is the next step of management?

a) Booster dose of vaccine

b) Immunoglobulin with full vaccination

c) Repeat full course of vaccine

d) No treatment required as he already has Hep B antibody titres from vaccination

Question 2:

The life cycle of an organism is given below. Identify the vector involved.



a) Hard tick

b) Soft tick

c) Trombiculid mite

d) Louse

Question 3:

What is the causative organism for the condition shown in the image below?



a) Echinococcus granulosus

b) Schistoma japonicum

c) Trypanosoma cruzi

d) Schistosoma mansoni

Question 4:

The presence of antibodies against SARS CoV 2 in the blood likely indicates

a) Previous infection even if patient has been asymptomatic

b) Presence of acute viral load

c) A high risk of reinfection

d) That the patient is in window period

Question 5:

Which of the following is the enzyme conjugate in the given ELISA technique?



a) A

b) B

c) C

d) D

Question 6:

A patient who underwent a renal transplant 3 months ago comes with complaints of diarrhea for the last 9 days. The stool sample was subject to Kinyoun stain revealing 4-6µm structures shown below. Which of the following is true about the organism?



a) Pathogen does not shows autoinfection

b) Cryptosporidium parvum is the most common organism implicated worldwide

c) Oocysts are immediately infective after being excreted in stool

d) It is an obligate intracellular organism

Question 7:

GeneXpert detects mutations in which of the following genes?

a) inhA

b) rpoB

c) KatG

d) pncA

Question 8:

A stool sample is collected. The specimen is processed for examination and is labeled at different levels as A, B, C, and D. Identify the level at which parasitic ova will be seen when viewed under a microscope.



a) A

b) B

c) C

d) D

Question 9:

Type IV hypersensitivity is due to?

a) Cell mediated response

b) Cell mediated and antibody mediated response

c) Antibody mediated response

d) Innate response

Question 10:

A child was brought to the OPD with a swollen, red knee joint that is tender to touch. 2 weeks prior, he had an episode of fever with bloody diarrhea. Which of the following could be the possible organism?

a) E. coli

b) Shigella

c) Giardia

d) Rotavirus

Question 11:

Specimen handling for molecular testing of SARS CoV 2 would require

a) BSL1

b) BSL2

c) BSL3

d) BSL4

Question 12:

Which of the following method is employed to measure the viral load?

a) ELISA

b) Western blot

c) Real time PCR

d) Immunoelectrophoresis

Question 13:

All of the following is true regarding scrub typhus except:

a) Transmitted by chiggers

b) Lymphadenopathy is a feature

c) Produces NLF producing colonies in MacConkey agar

d) Man is an accidental host

Question 14:

False about the Ebola virus is:

a) It is a filovirus

b) Transmitted by mosquito

c) Bats acts as reservoir

d) Nosocomial transmission seen

Question 15:

Identify the condenser _____

a) A

b) C

c) B

d) D

Question 16:

A young male presents to the STD clinic with painful lesions as shown in the image. What is the most likely microscopic finding of the lesion?

a) Tzank cells

b) Pund cells

c) Henderson-Patterson bodies

d) Guarnieri bodies

Question 17:

A patient has complains of fever with chills. A throat examination reveals swollen glands and a thick, whitish membrane over his tonsils. Albert staining is to be done. What are the constituents of Albert's stain?

a) Toluidine blue 'O' and malachite green

b) Carbol fuschin and methylene blue

c) Carbol fuschin and malachite green

d) Toluidine and methylene blue

Question 18:

Pneumonia due to a gram-negative pleomorphic bacillus growing on chocolate agar which is catalase and oxidase-positive is:

a) E.coli

b) Pseudomonas

c) Haemophilus influenzae

d) Actinomyces

Question 19:

A 30-year-old male presented to the OPD with painful genital ulcers. Which of the following can cause a painful ulcer on the glans penis?

a) Syphilis

b) Chlamydia

c) Chancroid

d) Lymphogranuloma venereum

Question 20:

Which protein is present in the HPV vaccine?

a) E6, E7

b) L1

c) E1, E2

d) L2

Question 21:

A patient presented with meningitis with a history of fever for 4 days and erythematous rashes on the chest. There was no history of travel and tick bite. His CSF findings showed increased protein, normal glucose, mild lymphocytosis and few RBCs. Which of the following is the most likely organism to cause this clinical picture?

a) West nile virus

b) Meningococci

c) Strep pneumoniae

d) Entero virus

Question 22:

A patient presents with seizures. His CT head reveals multiple parenchymal cysts. Which of the following is correct regarding management in this case?

a) Steroids followed by antiparasitic drugs

b) Antiparasitic drugs followed by steroids

c) Steroids before antiepileptics

d) Continue antiepileptics

Question 23:

The transplacental transmission of the organism given in the image which causes microcephaly is-



a) Nipah

b) Zika virus

c) CMV

d) Rubella

Question 24:

The disruption of which of the following is responsible for watery diarrhea in cholera caused by Vibrio?

a) Intercalated discs

b) Macula densa

c) Zonula occludens

d) Hemidesmosomes

Question 25:

Affinity maturation is due to:

a) Isotype switching

b) Somatic hypermutation

c) mRNA splicing

d) VDJ rearrangement

Question 26:

A farmer presents with a swollen foot and multiple sinuses as shown in the below image. What is the most likely diagnosis?

a) Mycetoma

b) Squamous cell carcinoma

c) Nocardia

d) Sporotrichosis

Question 27:

The given life cycle belongs to which of the following organisms?



a) Leptospira

b) Hanta virus

c) Yersinia pestis

d) Salmonella typhi

Question 28:

Identify the organism shown in this image.



a) Strongyloides stercoralis

b) Enterobius vermicularis

c) Ancylostoma duodenale

d) Ascaris lumbricoides

Question 29:

A patient presents with urethral discharge. A gram stain smear is made from the discharge sample and the following microscopic image is seen. What is the most likely causative organism?



a) Trichomonas

b) Chlamydia trachomatis

c) Neisseria gonorrheae

d) Gardenella

Question 30:

Which of the following disease has the highest mortality once clinically manifested?

a) Nipah

b) Rabies

c) Ebola

d) Zika

Question 31:

A patient presents with a mobile, warm to touch nodule in the neck. On the anaerobic culture of the biopsy material from this swelling, it showed colonies as shown in the image which had gram-positive branching rods on staining. Which of the following is the drug of choice in this condition?



a) Doxycycline

b) Cotrimoxazole

c) Penicillin

d) Surgical drainage

Question 32:

In which of the following situations will you use a sanitizer?

a) A, B & C only

b) A & B only

c) A, B C & D

d) C & D only

Question 33:

The following are true about chancroid:

a) C & D only

b) A, B,C & D

c) A B & C

d) A & B only

Question 34:

Match the following E.coli with their respective pathogenesis.

a) A-4,B-3,C-2,D-1

b) A-1,B-3,C-2,D-4

c) A-1,B-2,C-3,D-4

d) A-4,B-2,C-3.D-1

Question 35:

Match the following:

a) A-2, B-1, C-3, D-4

b) A-3, B-2, C-1, D-4

c) A-4, B-1, C-2, D-3

d) A-1, B-2, C-4, D-3

Question 36:

Galactomannan in serum is seen in:

a) 1 & 2 only

b) 1 & 3 only

c) 1 only

d) 1, 2 & 3 only

Question 37:

Match the following parasites with their respective hosts:

a) A-3, B-2, C-4, D-1

b) A-2, B-1 ,C-4 ,D-3

c) A-1, B-2, C-3, D-4

d) A-4, B-3, C-1, D-2

Answer Key

Question No.	Correct Option
1	d
2	c
3	a
4	a
5	b
6	c
7	b
8	d
9	a
10	b
11	b
12	c
13	c
14	b
15	c

16	a
17	a
18	c
19	c
20	b
21	d
22	a
23	b
24	c
25	b
26	a
27	a
28	d
29	c
30	b
31	c
32	a
33	c
34	c
35	b
36	c
37	a

Detailed Explanations

Solution to Question 1:

The given clinical scenario of needle-stick injury shows that HBsAg is negative and the anti-HBs is >105 mIU/ml. This suggests that the resident has had routine vaccination for Hep B infection. Since the antibody titers are more than 10 mIU/ml, no treatment is required.

Solution to Question 2:

The given life cycle is of trombiculid mite which acts as a vector for scrub typhus. Humans are seen to be infected when they are bitten by the mite larvae (chiggers).



Solution to Question 3:

The given image shows a hydatid cyst in the liver. It is caused by Echinococcus granulosus.

Echinococcus granulosus is the most common causative agent in hydatid cysts, while Echinococcus multilocularis and Echinococcus ligartus account for a small number of cases.

Dogs are the definitive hosts of E. granulosus. Sheep are the usual intermediate hosts, but humans are accidental intermediate hosts. Humans are end-stage hosts of the parasite since there is no human to human transmission.

Approximately 75% of hydatid cysts are located in the right lobe of the liver and are solitary. The clinical presentation of a hydatid cyst is largely asymptomatic until complications occur. The most common presenting symptoms are abdominal pain, dyspepsia, and vomiting. The most frequent sign is hepatomegaly.

The life cycle of Echinococcus is depicted in the image below:



Solution to Question 4:

The presence of antibodies against SARS CoV 2 in the blood likely indicates a previous infection even if the patient has been asymptomatic.

Antibodies most commonly become detectable 1–3 weeks after symptom onset, at which time evidence suggests that infectiousness likely is greatly decreased and that some degree of immunity from future infection has developed. But it is imperative to note that the information on SARS CoV 2 is still emerging.

Solution to Question 5:

The enzyme conjugate in the given ELISA technique is 'B'.

Other options:

A- Substrate - it will be cleaved by the enzyme present on the conjugate antibody.

B - Conjugate - it is an antibody that binds to the detection antibody. It is labeled with an enzyme that cleaves the substrate to form a color product/dye.

C - Detection antibody-binds the antigen. We direct our conjugate against this.

D - Capture Antibody-forms the base, on which antigen (say rotavirus from feces) from the patient sample binds.

Solution to Question 6:

Oocysts are immediately infective after being excreted in the stool.

Cryptosporidium is a coccidian parasite and implicated in causing traveler's diarrhea in immunocompromised patients. The two cryptosporidium species responsible are C.hominis and C.parvum. As the patient has undergone a renal transplant, he/she would most likely be on immunosuppressive medications, making the infection likely. Cryptosporidium hominis is the most common organism implicated worldwide

The infective form is oocysts, excreted in the affected patient's feces, acquired through drinking or recreational water. The oocysts are immediately infectious (unlike Isospora oocysts, which require maturation), and may cause autoinfection or implicated in person to person transmission. Because of its small size (4-5micron), it is difficult to see in routine staining, hence is subjected to acid-fast/kinyoun's stain.

Once ingested, the 4 sporozoites come out of the cyst and attach to the small intestinal mucosa. It causes crampy abdominal pain and non-inflammatory watery diarrhea.

Treatment is mainly supportive, though nitazoxanide has been FDA approved for use in cryptosporidium.

Comparative features of coccidian parasites:

Property	Cryptosporidium	Cyclospora	Cystoisospora
Sporulated oocyst	4-6 µm, round, contains f our sporozoites	8-12 µm, round, contains 2 sporocysts each having t wo sporozoites	23-36 µm, oval, contain t wo sporocysts, each havin g four sporozoites.
Acid-fast detection	Uniformly acid-fast	Variable acid-fast	Uniformly acid-fast
Sporulation of the ooc yst	Occurs inside the host cell	Occurs in soil	Occurs in soil
Diagnostic form	Sporulated cyst	Unsporulated cyst	Unsporulated oocyst

Solution to Question 7:

Gene Xpert identifies mutations in the rpoB gene to detect rifampicin resistance (RR).

Other options:

inhA and KatG- involved in resistance against isoniazid

pncA- involved in resistance against pyrazinamide

Solution to Question 8:

The parasitic ova will be seen in a specimen taken from level D.

The image shows a formol-ether sedimentation technique to concentrate the ova in the bottom (D) part of the tube. It is used for samples in which the concentration of parasitic ova is scanty. It is useful for all helminth ova/eggs and parasitic cysts. The ether adsorbs the fecal debris and floats. Formalin fixes and preserves the specimen.

This method is useful for all helminth eggs and protozoan cysts.

Solution to Question 9:

Type IV hypersensitivity is due to the cell-mediated response.

Type IV or cell-mediated hypersensitivity reaction: These reactions are induced by sensitized T cells which on contact with the specific antigen releases cytokines that produce biological effects on leucocytes, macrophages, and tissue cells.

Solution to Question 10:

The given clinical scenario points towards a diagnosis of reactive arthritis after shigella infection.

Shigellosis is a spectrum of symptoms caused by shigella consisting of 4 phases:

Other options:

E.coli - it may cause diarrhea associated with fever or blood in stools, but reactive arthritis is not typically seen.

Giardia - it may cause persistent watery diarrhea with waxing and waning course. Fever, blood in stools should prompt alternate diagnosis. Also, reactive arthritis is not seen.

Rotavirus - it is a well known viral cause of fever with diarrhea in children. But not known to have reactive arthritis as a sequela.

Incubation period	1-4 days (up to 8 days)
Watery diarrhea	Fever, watery diarrhea, anore xia
Dysentry	Bloody mucopurulent stools

Incubation period	1-4 days (up to 8 days)
Post-infectious complication	Hemolytic-uremic syndrome Reactive arthritisToxic megac olon

Solution to Question 11:

Specimen handling for molecular testing of SARS CoV 2 would require BSL-2 or equivalent facilities. Attempts to culture the virus requires minimum BSL-3 facilities.

SARS CoV 2 virus can cause coronavirus disease 2019 (COVID-19)

Microbiology and pathology laboratories performing diagnostic tests on a stool or respiratory specimen should handle potential SARS-CoV specimens using standard biosafety level (BSL)-2.

Note: Biosafety levels include a specific combination of work practices, safety equipment, and facilities that are designed to minimize the exposure of workers and the environment to infectious agents. There are four biosafety levels.

BSL-1: Work can be done on an open bench or table and PPE needed. Eg: Non-pathogenic E.coli.

BSL-2: Work to be done in the biosafety cabinet with PPE. Eg: Staphylococcus aureus.

BSL-3 laboratory has appropriate PPE and respirators might be required. All work performed within BSC(biosafety cabinet). Entrance is through 2 sets of self-closing and locking doors. Eg: when dealing with Mycobacterium tuberculosis.

BSL-4: This is the highest level of biological safety, where containment of microbes is done, which are fatal without treatment and vaccines. Eg: Ebola and Marburg viruses.

Solution to Question 12:

Viral load can be measured by the real time-PCR technique.

Real time-PCR assays are commonly used for viral load quantification of HIV, hepatitis C virus, hepatitis B virus, and CMV.

Other options:

• Western blotting is used for the detection of protein.

• ELISA is an immunosorbent assay used to detect antigens of infectious agents present in clinical specimens.

• Immuno electrophoresis includes electrophoretic separation of antigens followed by an antigen-antibody reaction.

Solution to Question 13:

The organism causing scrub typhus i.e Orientia tsutsugamushi is non-cultivable on cell-free media like MacConkey agar.

Scrub Typhus is also known as chigger borne typhus. The vector is trombiculid mite. Humans are infected when they are bitten by the mite larvae known as chiggers.

The incubation period is 1- 3 weeks. The patient typically develops a characteristic eschar at the site of the mite bite, with regional lymphadenopathy and maculopapular rash.

Solution to Question 14:

Ebola virus is not transmitted by mosquito.

Ebola virus is the cause of an emerging disease called Ebola virus disease and is widespread in Africa. It belongs to the Filoviridae family.

Fruit bats act as reservoir hosts of the virus and it is transmitted to humans only accidentally. Human to human transmission is through direct contact with blood or body fluids of an infected symptomatic person.

The incubation period of the disease is 2 to 21 days. Symptoms include fever, intense weakness, vomiting, diarrhea, rashes, and, in extreme cases, internal and external bleeding. The current treatment is mainly supportive. Nosocomial transmission seen.

Solution to Question 15:

Condensor lens of microscope is marked as B in the image. The purpose of the condenser lens is to focus the light onto the specimen.

A is coarse focusing knob.

C is objective lens. There are usually 3 or 4 objective lenses on a microscope. They almost always consist of 4X, 10X, 40X and 100X powers.

D is Illuminator. It is a steady light source used in place of a mirror.



Solution to Question 16:

Clusters of vesicles are seen on an erythematous base in the image, suggesting a diagnosis of genital herpes. Tzanck cells or multinucleated giant cells are present on microscopy.

The Tzanck smear is a rapid, sensitive, and inexpensive diagnostic method for herpes simplex infections. Smears are prepared from the lesions, from the base of vesicles and stained with a 1% aqueous solution of O-toluidine blue for 15 seconds. Multinucleated giant cells with faceted nuclei and homogeneously stained ground glass chromatin (Tzanck cells) are seen.

Mucocutaneous herpetic lesions are thin-walled, umbilicated vesicles, the roof of which breaks down, leaving tiny superficial ulcers which heal without scarring. Painful and pruritic lymphadenopathy may be seen. The most common site of cutaneous lesions is the face - on the cheeks, chin, around the mouth, or on the forehead - and is caused by herpes simplex virus-1. Genital infections are more commonly caused by herpes simplex virus-2.

Tzanck smear



©MARROW

Solution to Question 17:

Albert's stain contains toluidine blue 'O' and malachite green which have an affinity for acidic cytoplasm, both being basic dyes.

Albert's stain, Neisser's and Ponder's stain are used to demonstrate the metachromatic granules (Volutin / Babes Ernst Granules) of Corynebacterium diphtheria.

Procedure for Albert's staining:

• A heavy smear of bacterial isolate was prepared on glass slide and allowed to dry without fixing.

• Albert's stain was applied for (3-4) minutes then it was washed with water and Albert's iodine was applied for one minute.

• Wash with water and let dry with blotting paper.

• Examine the slide under oil immersion, it was examined under microsope showed bacili with numerous metachromatic granules.

Results:

Metachromatic granules: Bluish- black

Bacillary body or protoplasm: Green

Uses:

• Albert staining is useful in the demonstration of metachromatic granules.

• Presence of metachromatic granules helps to distinguish C.diphtheriae from diphtheriods which lack them.

Solution to Question 18:

Pneumonia due to a gram-negative pleomorphic bacillus growing on chocolate agar which is catalase and oxidase-positive is most likely Haemophilus influenza.

Haemophilus pneumonia typically occurs in infants and is accompanied by empyema and sometimes meningitis as well. In older children and adults, the picture is of lobar pneumonia. Sputum is collected for the detection and isolation of Hemophilus. As the bacillus is very sensitive to low temperatures, specimens should never be refrigerated before inoculation. H.infiuenzae is a Gram-negative, non-motile, non-sporing bacillus, exhibiting considerable pleomorphism. In sputum, it usually occurs as clusters of coccobacillary forms.

The bacillus has fastidious growth requirements. The accessory growth factors, named X and V, present in the blood are essential for growth. Catalase and

oxidase reactions are positive. Nitrates are reduced to nitrites. Biotyping has been done on the basis of indole production.

Cultures may be preserved for about a month on chocolate agar slopes. Chocolate agar is made by heating blood agar to 80-90°C for 15-20 minutes. This causes the release of V factor from within the erythrocytes.

Treatment- Amoxycillin-clavulanate or clarithromycin is more effective in treating respiratory infections by Haemophilus influenzae.

Solution to Question 19:

Painful genital ulcers are caused by Haemophilus ducreyi, and are known as chancroid.

The regional lymph nodes are enlarged and painful. Microscopic appearance seen in H. ducreyi is called the school of fish appearance or railroad appearance. Chocolate agar with isovitaleX is a suitable culture medium for Haemophilus ducreyi.

The given image shows the microscopic appearance of H. ducreyi.

Solution to Question 20:

The HPV vaccines use virus-like particles (VLPs) that consist of the HPV L1 major capsid protein. The L1 protein self-assembles into VLPs when expressed in eukaryotic cells.

These VLPs contain the same epitopes as the HPV virion. But they do not contain genetic material and cannot transmit infection. The immunogenicity of the HPV vaccines relies on the development of conformational neutralizing antibodies to epitopes displayed on viral capsids.

The nonavalent HPV vaccine works against strains 6,11,16,18,31,33,45,52 and 58. It is usually commercialized in the name of Gardasil 9.

HPV VACCINE	HPV TYPES COVERED
Quadrivalent	6, 11,16, 18
Bivalent	16, 18
Nonavalent	6, 11, 16, 18, 31, 33, 45, 52, 58

Recent update: Gardasil 9 has received expanded indication which now includes the prevention of oropharyngeal and other head and neck cancers caused by HPV types 16,18,31,33,45,52,58.

Solution to Question 21:

Meningitis with normal glucose, increased protein, and mild lymphocytosis is suggestive of viral meningitis, most probably by an Enterovirus.

Coxsackieviruses are typical enteroviruses. Two groups of Coxsackie viruses A and B, are known. Aseptic meningitis may be caused by a majority of group A and all group B viruses. A maculopapular rash may be present. The disease can also occur as epidemics.

CSF findings in Viral meningitis:

• A mild increase in CSF protein/pressure

• Normal glucose

• Mild increase in lymphocytes in CSF.

Other options:

West Nile virus is endemic in certain regions in North America. But there is no history of travel mentioned in the clinical scenario above. Current hotspots in North America are the California Central Valley, southwestern Arizona, southern Texas, and Louisiana, which have both compatible climates and avian reservoirs for the disease.

	Normal	Bacterial Meningitis	Viral Meningitis	Fungal Meningitis	Parasitic Meningitis	Tuberculo Meningitis
WBC Count (per μL)	<5	>1000	25-500	40-600	150-2000	25-100
Differential of WBC	60-70% lymphocytes, ≤30% monocytes/ macrophages	↑↑PMNs (≥80 %)	Predominantly lymphocytes	Lymphocytes or PMNs, depending on the specific organism	↑↑ Eosinophils (≥50 %)	Predominantly lymphocytes
Gram,s stain	Negative	Positive (in > 60% of cases)	Negative	Rarely positive	Negative	Occasionally positive
Glucose (mg/dL)	40-85	<40	Normal	↓ to normal	Normal	<50 in 75% of cases

	Normal	Bacterial Meningitis	Viral Meningitis	Fungal Meningitis	Parasitic Meningitis	Tuberculo Meningitis
Protein (mg/dL)	15-45	>100	20-80	150-300	50-200	100-200
Opening pressure (mmH2O)	50-180	>300	100-350	160-340	Normal	150-280

Solution to Question 22:

The given clinical scenario is suggestive of neurocysticercosis caused by larvae of T. solium (cysticercosis cellulosae). Glucocorticoid therapy is usually begun before initiating albendazole (antiparasitic drug). This is continued for several days after the institution of therapy to reduce the incidence of side effects from inflammatory reactions to dead and dying cysticerci of Taenia solium.

Albendazole is the drug of choice for neurocysticercosis.

There are various types of cysticercosis based on the site affected such as:

• Subcutaneous nodules (asymptomatic)

• Neurocysticercosis

• Muscular cysticercosis

• Ocular cysticercosis.

Neurocysticercosis is considered the second most common cause of intracranial space-occupying lesion (ICSOL) in India following tuberculosis. Its clinical features include:

• Epilepsy

• Raised intracranial tension

• Hydrocephalus

• Psychiatric disturbances

• Meningoencephalitis

• Transient paresis

• Behavioral disorders

• Aphasia

• Visual disturbances.

The below MRI of the brain shows multiple cysts containing scolices in a patient with neurocysticercosis.



Solution to Question 23:

The image shows transplacental transmission of a virus, leading to microcephaly, with the Aedes mosquito as its vector. Among the given options, Zika virus has the above features.

Zika virus is a mosquito-borne Flavivirus mainly transmitted through Aedes aegypti. Zika virus infection presents with mild symptoms of fever, rash, conjunctivitis, malaise, headache and muscle and joint pain. These symptoms usually last for 2-7 days. Maternal infection during pregnancy leads to congenital Zika syndrome or microcephaly.

The complications of Zika virus infection include Guillain-Barré syndrome, neuropathy, and myelitis.

Note: NCDC Delhi and the National Institute of Virology Pune are the only institutes that have the capacity to provide the diagnosis in an acute febrile stage. RT-PCR is the standard test available for the diagnosis of Zika virus infection.

Solution to Question 24:

The disruption of zonula occludens is responsible for watery diarrhea in cholera caused by Vibrio cholerae.

Zonula occludens toxin (Zot) is an enterotoxin elaborated by Vibrio cholerae that increases intestinal permeability by interacting with a mammalian cell receptor with subsequent activation of intracellular signaling leading to the disassembly of the intercellular tight junctions.

Mechanism of action of cholera toxin:

• The cholera toxin molecule consists of 1 A and 5 B subunits. The B (binding) units attach to the GM1 ganglioside receptors on the surface of jejunal epithelial cells. The A (active) subunit, on being transported into the enterocyte, dissociates into A1 and A2. A2 fragment only links

biologically active A1 to the B subunit.

• A1 causes prolonged activation of cellular adenylate cyclase and accumulation of cAMP, leading to outpouring into the small intestinal lumen, of large quantities of water and electrolytes. This leads to watery diarrhea which is isotonic with plasma but contains much more potassium and bicarbonate.

• The toxin also inhibits the intestinal absorption of sodium and chloride.

• All clinical manifestations and complications in cholera result from the massive water and electrolyte depletion thus caused.



Solution to Question 25:

Affinity maturation is due to somatic hypermutation.

Affinity maturation is the process by which antibodies gain increased affinity, avidity, and anti-pathogen activity. It is a result of somatic hypermutation (SHM) of immunoglobulin genes in B cells, coupled to selection for antigen binding.

This process occurs in the germinal centers and proceeds for weeks after acute infection or vaccination, or for many cycles during chronic infection. The resulting antibodies can be highly mutated from their germline-encoded counterparts, with increases of several orders of magnitude in affinity for antigen compared to the corresponding naïve B cell receptors (BCRs).

Other options:

Option A: Isotype switching, also known as class switch recombination, is a process where the constant region of the antibody heavy chain is changed, leading to a switch in the antibody isotype (e.g., from IgM to IgG). This process is crucial for the functional properties of antibodies but is not directly involved in affinity maturation.

Option C: mRNA splicing is a cellular process that involves the removal of introns and the joining of exons to generate mature mRNA. While this process is essential for the production of functional antibodies, it is not specifically associated with affinity maturation. Affinity maturation

primarily occurs through somatic hypermutation.

Option D: VDJ rearrangement is the process by which the variable regions of immunoglobulin genes are assembled from gene segments (V, D, and J segments). This process contributes to the diversity of antibodies but is distinct from affinity maturation. Affinity maturation occurs after VDJ rearrangement and is driven by somatic hypermutation in the variable regions.

Solution to Question 26:

The most likely diagnosis here is Mycetoma which is caused by Eumycetoma.

Mycetoma:-

• Mycetoma is a chronic, granulomatous infection of the skin, subcutaneous tissues, fascia and bone, which most often affects the foot or the hand. It is, therefore, commonly known as Maduramycosis or Madura foot. Mycetoma occurs worldwide.

• Etiology: It can be divided into three types, eumycetomas, actinomycetomas and botryomycosis.

Pathogenesis:-

• Triad of symptoms: Infection follows traumatic inoculation of the organism into the subcutaneous tissue from soil or vegetable sources, usually on thorns or splinters and results in tumefactions, deformities and draining sinuses discharging fungal colonies called grains or granules (triad of symptoms).

• Subcutaneous tissues of the feet, lower extremities, hands, and exposed areas are most often involved.

• Grains: Within host tissues the organisms develop to form compacted colonies (grains) 0.5–2 mm in diameter, the color of which depends on the organism responsible.

Laboratory Diagnosis

• A. Direct Examination: The presence of grains in pus collected from draining sinuses or in biopsy material is diagnostic.

• B. Culture Samples should also be cultured, at both 25–30°C and 37°C, on brain-heart infusion agar or blood agar for actinomycetes and on Sabouraud agar (without cycloheximide) for fungi.

Treatment:-

The management of eumycetoma is difficult, involving surgical debridement or excision and chemotherapy. Actinomycetoma responds well to rifampicin in combination with sulfonamides or cotrimoxazole.

Solution to Question 27:

The life cycle shown in the above image is suggestive of leptospirosis.

It is caused by contact with the urine or blood of an infected organism. It is a zoonotic disease, with rats being the most well-known reservoirs.

Illness is classically biphasic after an incubation period of up to 1 month. The stages are:

• Acute leptospiremic phase- organism can be cultured from blood.

• Resolving immune phase- organism can be cultured from urine.

The stages may not be clinically distinguishable.

Most cases are asymptomatic but a few may present with severe manifestations. Clinical features include:

• High-grade fever

• Hepatic dysfunction

• Renal impairment

• Hemorrhages- gastrointestinal, pulmonary (most common), petechiae and purpura

The severe form is known as Weil's disease, with a triad of hemorrhage, jaundice, and acute kidney injury.

Treatment of choice is IV penicillin or IV ceftriaxone.



Solution to Question 28:

The given image shows large cylindrical worms with tapering ends, and thus represents Ascaris lumbricoides (roundworm).

Ascaris lumbricoides is the most common of human helminths and is distributed worldwide. It is the largest nematode parasite in the human intestine. The adult worms are large cylindrical worms, with tapering ends. The anterior end is more pointed than the posterior. It appears white outside the body.

Its natural host is humans. There is no intermediate host. Adult worms live in the small intestine - jejunum and ileum. Humans acquire infectionby consuming food or water contaminated with nightsoil or human feces containing eggs. The eggs contain the infective rhabditiform larva.

Solution to Question 29:

The above image shows gram■negative intracellular diplococci within phagocytes in a stained smear from the urethral discharge sample, which is seen in infection with Neisseria gonorrhoeae.

According to CDC recommendations 2021, the first line of treatment in uncomplicated gonococcal infections is a single dose of ceftriaxone 500 mg IM.

Note: BASHH (British Association for Sexual Health and HIV) in the United Kingdom also recommends higher doses of ceftriaxone of about 500 mg along with azithromycin because it would result in fewer failures.

Solution to Question 30:

Rabies has the highest mortality once clinically presented.

Rabies is a vaccine-preventable, zoonotic, viral disease of the central nervous system. Rabies virus is a bullet-shaped rhabdovirus containing ssRNA. It spreads to humans by bites or contact with the saliva of rabid animals (bats, dogs, cats). Once clinical symptoms appear, rabies is virtually 100% fatal. The disease can be divided into 3 phases:

• Prodromal phase: It lasts 2-10 days. Patients present with malaise, headache, photophobia, vomiting, and fever. They may also have parasthesias, pain or pruritus near the site of exposure.

• Acute neurologic phase: It lasts 2-7 days. In the more common encephalitic rabies, Hydrophobia (fear of water) or aerophobia (fear of breeze) can be seen. Swallowing liquids precipitates painful spasms of the throat muscles. Patients may also have nervousness, hallucinations, excessive lacrimation, salivation, perspiration and pupillary dilation. Convulsive seizures may occur

• Coma and death (due to cardiorespiratory arrest)

In about 20% of patients, paralytic rabies is seen. Flaccid muscle weakness is seen, which progresses to quadriparesis and facial palsy.

Note: In rabid animals, the virus depicts two modes of spread:

• Centripetal spread: The virus enters the peripheral neurons. It moves via retrograde fast axonal transport towards the spinal cord or brainstem a.k.a the central nervous system (CNS) at a speed of up to 250 mm/day. Once within the CNS, it spreads quickly to other regions in the CNS through fast axonal transport. It infects neurons primarily, but may rarely infect astrocytes.

• Centrifugal spread: After CNS infection is established, it spreads along the sensory and autonomic nerves to salivary glands, adrenal glands, skin

Rabies virus replicates in acinar cells of the salivary glands and is secreted in the saliva of rabid animals that serve as vectors.

Solution to Question 31:

The given image shows molar tooth colonies that are characteristic of Actinomyces species. The given clinical scenario of a mobile, warm to touch nodule in the neck describes cervicofacial actinomycosis also known as 'lumpy jaw syndrome'. Penicillin is the drug of choice for actinomyces infection.

Actinomyces are Gram-positive anaerobic bacilli that produce molar tooth colonies on incubation for 72 hours on brain-heart infusion agar. They cause a chronic suppurative and granulomatous infection called actinomycosis. It can present as one of the following clinical types:

• Cervicofacial type- Fluctuant mass in the jaw area producing draining fistulas.

• Thoracic type- Pulmonary infection with fever, cough, and purulent sputum.

• Abdominal type- Occurs following ruptured appendix or ulcer. It may spread throughout the peritoneum and involve multiple organs.

Actinomycosis can spread to bones and surrounding tissue and produce multiple discharging sinuses. Treatment requires prolonged administration of penicillin for 6–12 months. In penicillin-allergic patients, clindamycin or erythromycin is effective. Surgical excision and drainage may be required in some cases.

The given image shows cervicofacial actinomycosis.

The given image shows a patient with multiple cutaneous lesions caused by Actinomyces:

Solution to Question 32:

You should use a sanitiser before and after examining the patient and after touching the patients' surroundings. The hands should be washed with soap and water thoroughly in case they are visibly soiled.

Sanitization is the process through which pathogenic organisms are reduced to safe levels on inanimate objects to reduce the likelihood of cross-infections. It is a component of hand-hygiene, which is demonstrated by the image below.



Note: Disinfectants like chlorhexidine and alexidine have a biguanide functional group. They are mostly used as oral disinfectants and hand products. They are bactericidal but not sporicidal. Mycobacteria are also resistant to these compounds.

Solution to Question 33:

Chancroid is a painful genital ulcer caused by Haemophilus ducreyi, which is a gram-negative coccobacillus. Microscopic appearance of H. ducreyi is called the school of fish appearance or railroad appearance. Chancroid ulcers have a soft base that bleeds easily to touch. They also cause painful lymphadenopathy.

The image below shows the 'school-of-fish' appearance characteristic of H. ducreyi.



Note: Groove sign is a characteristic sign seen in lymphogranuloma venereum. It refers to the depression created by the inguinal ligament as it passes between the groups of inflamed inguinal lymph nodes.

Solution to Question 34:

The correctly matched pairs are as follows:

The various strains of Escherichia coli are:

• Enterotoxigenic E. coli (ETEC) is the most common organism that causes diarrhea in travelers. It produces the following toxins:

• LT (heat-labile toxin)- acts by ↑cAMP

• ST (heat-stable toxin)- two types of ST are known, ST-A (ST-I) and ST-B (ST-II)

• ST-A is soluble in methanol and acts by ↑cGMP in the intestine.

• ST-B is insoluble in methanol, and the mode of action is not known.

• Enteroinvasive E. coli (EIEC) invade interstitial epithelial cells in the intestine and cause a condition resembling shigellosis. This ability is determined by a plasmid, detection of which can

also be a diagnostic test. Sereny's test was employed to diagnose EIEC but it is not in use anymore.

• Enterohemorrhagic E. coli (EHEC) produces verocytotoxin, which targets vascular endothelium. It causes hemorrhagic colitis and hemolytic uremic syndrome.

• Enteropathogenic E. coli (EPEC) causes infantile diarrhea in developing countries. It adheres to the mucosal surface of small intestine using EAF and LEE and causes effacement of microvilli. This leads to attachment and effacement lesions of intestinal cells.

• Enteroaggregative E. coli (EAEC) produces the stacked brick pattern of adherence on hep-2 cells. EAEC has been associated with persistent diarrhea. Intestinal colonization of EAEC is mediated by AAF (aggregative adherence fimbriae). EAEC produces EAST-1 toxin (enteroaggregative heat-stable toxin) which is considered to be a major virulence factor.

E. coli	Mechanism of action
1. ETEC	A. Produce a heat-stable toxin and cause activation of guanylyl cyclase
2. EIEC	B. Invades intestinal mucosal cells
3. EHEC	C. Produces a toxin against vascular endothelium cells
4. EPEC	D. Adhere to the mucosal surface and cause effacement of microvilli

Solution to Question 35:

Identification features of Trichomonas vaginalis:

• Pear-shaped or ovoid

• Short undulating membrane reaching upto the middle of the body

• It has four anterior flagella and fifth running along the outer margin of the undulating membrane, which is supported at its base by a flexible rod, costa

• A prominent axostyle runs throughout the length of the body and projects posteriorly like a tail

• The cytoplasm shows prominent siderophillic granules, which are most numerous alongside the axostyle and costa

• Jerky or twitching type of mobility

Parasite	Organ of locomotion
Balamuthia mandrillaris	Pseudopod

Parasite	Organ of locomotion
Balantidium coli	Cilia
Trichomonas vaginalis	Flagella
Toxoplasma gondi	No Particular organ



The above image shows Trichomonas vaginalis : 1-anterior flagella, 2-kinetosomes, 3-parabasal body, 4-costa, 5-parabasal fibers, 6-undulating membrane, 7-posterior flagellum, 8-hydrogenosomes, 9-axostyle, 10-nucleus, 11-pelta.

Obligate parasite - Organism cannot survive outside host

• No cystic stage. Trophozoites are infective forms

• Sexual transmission is the main route of infection. Other modes are fomites, babies during passage through infected birth canal

• Affects squamous epithelium - Intracellular edema and so called chicken-like epithelium, is the most characteristic feature of trichomoniasis

• Habitat: Vagina and cervix in female and urethra in males

• Clinical features: Often asymptomatic in males. In females, it leads to pruritic vaginitis with greenish yellow discharge, strawberry mucosa and dysuria

• Diagnosis: By wet mount microscopy of vaginal or urethral discharge, Culture (gold standard - Cysteine-peptone-liver-maltose (CPLM) medium and plastic envelope medium (PEM)), PCR, and by demonstration of antigen in vaginal smear by ELISA

• Treatment: Metronidazole is the drug of choice and simultaneous treatment of both partners is recommended



The given image is the trophozoite form of the Balantidium coli.

Balantidium coli occurs in two stages

• Trophozoites are oval in shape and contain two nuclei. There is the presence of cilia over the entire surface and are larger around the mouth. Its anterior end is narrow and the posterior end is broad. At the anterior end, there is a groove leading to the mouth (cytostome). Posteriorly, there is a small anal pore (cytopyge).

• Cysts are smaller, round with a micronucleus, a macronucleus and many contractile vacuoles.

Living trophozoites/cysts are yellowish or greenish in color on wet mount.



The image given above shows Toxoplasma gondii which has two types of life cycles:

• Enteric cycle

• Exoenteric cycle

The enteric cycle occurs in definitive hosts such as cats. Exoenteric cycle occurs in intermediate hosts such as humans, mice, rats, etc.

Both sexual reproduction (gametogony) and asexual reproduction (schizogony) occur within the mucosal epithelial cells of the small intestine of the cat.

Sporozoites from the oocysts and bradyzoites from the tissue cysts enter the intestinal mucosa and multiply asexually. Tachyzoites are formed within the intermediate host.

Tachyzoites continue to multiply and spread locally by lymphatic system and blood, while the bradyzoites multiply slowly inside the tissue cyst.

The dormant bradyzoites inside the cyst may be reactivated in immune suppression causing renewed infection in the host.



Images b-1 and b-3 showing the Trophozoite and Cystic form of Balamuthia mandrillaris in Salmonella Shigella (SS) Agar respectively.

Balamuthia Mandrillaris

B. mandrillaris is a free-living ameba, which is a newly identified species reported to cause granulomatous amebic encephalitis (GAE).

Morphology

It exists in ameboid trophozoite stage. The flagellate stage is absent

• It is relatively large (12-60 μm), irregular in shape and actively motile by broad pseudopodia.

• Cyst of B. mandrillaris are usually spherical (6-20 μm), surrounded by a three-layered cyst wall: (1) outer irregular ectocyst, (2) a middle mesocyst and (3) an inner endocyst round wall. Under light microscopy, it appears to have two walls: (1) an outer irregular wall and (2) an inner smooth wall.

• Infection is transmitled through respiratory tract, skin lesions, or eyes.

Solution to Question 36:

The galactomannan assay is used for the diagnosis of invasive (deep-seated) mycoses caused by Aspergillus species.

Galactomannan is a component of the cell wall of Aspergillus. It can be detected in serum or bronchoalveolar lavage by ELISA for the diagnosis of invasive aspergillosis in adults and children. This test is especially useful in hematopoietic stem cell transplant recipients or patients with hematologic malignancies. Invasive aspergillosis affects the lungs and presents with fever, cough, dyspnea, and hemoptysis. The organism may spread to the eye, gastrointestinal tract, kidney, liver or brain, leading to abscesses and necrotic lesions.

Other diagnostic features of invasive aspergillosis are:

• Positive culture from the affected organ

• Microscopy showing narrow, septate hyphae with branching at 45o and conidiophores

• Halo sign on CT scan

Solution to Question 37:

The correct answer is A-3, B-2, C-4, D-1.

The hosts of the above parasites are given below:

A. Diphyllobothrium	3. Cyclops
B. Paragonimus	2. Crayfish
C. Brugia	4. Mansonia
D. Fasciola	1. Sheep

Parasite	Definitive host	First intermediate host	Second intermediate host
Diphyllobothrium	Humans	Cyclops	Cyclops and fish
Paragonimus	Humans	Snail	Crayfish and crab
Fasciola	Sheep and humans	Snail	Aquatic vegetations
Brugia	Humans	Mansonia mosquito	-

Microbiology NEET 2018

Question 1:

Nude mice are not resistant to xenograft due to the absence of _____

a) B cell

b) T cell

c) Both B and T cell

d) NK cell

Question 2:

Which of the following is Burkholderia cepacia resistant to?

a) Ceftazidime

b) Trimethoprim-sulfamethoxazole

c) Temocillin

d) Cefotetan

Question 3:

Which of the following organisms causes scombroid fish poisoning?

a) Salmonella

b) Staphylococcus

c) Pseudomonas aeruginosa

d) Peptostreptococcus

Question 4:

Which of the following is the predominant pathogen to infect renal transplant recipients one to four months following transplantation?

a) Cytomegalovirus

b) JC virus

c) BK virus

d) Epstein-Barr virus

Question 5:

Sabin-Feldman dye test is used for diagnosis of which of the following condition?

a) Paragonimiasis

b) Hydatid cyst

c) Toxoplasmosis

d) Cryptococcus

Question 6:

Which of the following is called as Australia antigen?

a) HBsAg

b) HBeAg

c) HBcAg

d) HBV-DNA

Question 7:

Bacterial transduction occurs by

a) Plasmids

b) Sex pili

c) Bacteriophage

d) Uptake of genetic material by other bacteria.

Question 8:

Which fungus is most commonly associated with orbital cellulitis in patients with diabetic ketoacidosis ?

a) Candida

b) Rhizopus

c) Aspergillus

d) Trichophyton

Question 9:

The adult form of echinococcus is seen in which of the following?

a) Dog

b) Cat

c) Swine

d) Sheep

Question 10:

Which of the following can grow in an acidic pH?

a) Vibrio

b) Lactobacilli

c) Salmonella

d) Shigella

Question 11:

What is the vector for the condition diagnosed by the image shown below?



a) Anopheles mosquito

b) Ixodes Tick

c) Sandfly

d) Tsetse fly

Question 12:

TB test done only on whole blood sample is _____.

a) TrueNat TB

b) Gene xpert

c) Interferon gamma release assay

d) All the above

Question 13:

Multiple abscesses with discharging sinuses, demonstrating sulphur granules in pus, are caused by

a) Actinomyces

b) Nocardia

c) Salmonella

d) Tularemia

Question 14:

Cutaneous larva migrans is most commonly caused by which organism?

a) Strongyloides stercoralis

b) Toxocara canis

c) Ancylostoma braziliense

d) Necator americanus

Question 15:

Which type of hypersensitivity reaction is involved in the hyperacute rejection of renal transplants?

a) Type IV

b) Type III

c) Type II

d) Type I

Question 16:

Which of the following organisms produce urease?

a) Helicobacter pylori

b) Salmonella

c) Escherichia coli

d) Shigella

Question 17:

Mycoplasma is inherently resistant to which of the following antibiotics?

a) Beta lactam antibiotics

b) Macrolides

c) Tetracyclines

d) Flouroquinolones

Question 18:

Pasteurization is done at _____.

a) 73 degree Celsius for 20 minutes

b) 63 degree Celsius for 30 minutes

c) 72 degree Celsius for 30 seconds

d) 63 degree Celsius for 30 seconds

Answer Key

Question No.	Correct Option
1	b
2	d
3	c
4	a
5	c
6	a
7	c
8	b
9	a
10	b
11	c
12	c
13	a
14	c
15	c
16	a
17	a
18	b

Detailed Explanations

Solution to Question 1:

Nude mice are not resistant to xenograft due to absence of T cell.

A xenograft is a tissue graft or organ transplant from a donor of a different species.

Nude mice lack body hair, hence the name. They lack a thymus and thus cannot generate mature T lymphocytes. The absence of T-lymphocytes makes them unable to mount adaptive immune responses requiring helper T cells and cytotoxic T cells. The absence of functioning T cells also prevents the rejection of allografts & xenografts.

Solution to Question 2:

Burkholderia cepacia is resistant to cefotetan.

B. cepacia complex strains are intrinsically resistant to aminoglycosides, polymixin, first and second-generation cephalosporins (e.g. cefotetan) and carboxypenicillins. This antimicrobial resistance is mediated by efflux pumps, antimicrobial degradation, modifying enzymes or altered

membrane functions.

B. cepacia is a Gram-negative bacillus which can cause severe necrotizing pneumonia in patients with chronic granulomatous disease. It can lead to fatal septicemia in patients with cystic fibrosis.

Antimicrobial agents that are effective against B. cepacia complex include meropenem, ceftazidime, trimethoprim-sulfamethoxazole, piperacillin and temocillin.

Solution to Question 3:

Scombroid fish poisoning is associated with Pseudomonas aeruginosa.

It follows the consumption of both scombroid (mackerel-like) fish and nonscombroid fish that have not been refrigerated properly. The musculature of these dark- or red-fleshed fish undergoes decomposition by Morganella morganii, Escherichia coli, Proteus species, Klebsiella species, and Pseudomonas aeruginosa. There is consequent decarboxylation of the amino acid L-histidine to form histamine.

A second compound is thought to play a causative role in this intoxication because large doses of oral histamine do not reproduce the affliction. The toxin or toxins involved are heat stable and are not destroyed by cooking or freezing.

Symptoms develop within 15–90 min of ingestion. Most cases are mild, with tingling of lips and mouth, mild abdominal discomfort, and nausea. The more severe and commonly described presentation includes flushing, conjunctival hyperemia, pruritus, urticaria, and angioedema. Without treatment, the symptoms generally resolve within 8–12 h.

Solution to Question 4:

The predominant pathogen to infect renal transplant recipients in the period around 1 to 4 months following the transplantation is cytomegalovirus (CMV).

CMV is the most common virus to affect recipients after organ transplantation (kidney, heart, lung etc). The maximal risk occurs between 1 and 4 months after transplantation (in the absence of prophylaxis). The most common CMV presentation in kidney transplant patients includes fever, leukopenia, hepatosplenomegaly, myalgia, and arthralgia. Antiviral prophylaxis with ganciclovir or valganciclovir is used in renal transplant recipients whenever indicated to prevent CMV disease.

Given below are the most common opportunistic infections in renal transplant recipients:

Peritransplant (less than 1 m onth)	Early (1-6 months)	Late (more than 6 months)
Wound infectionHerpes virus Oral candidiasisUrinary tract infection	Pneumocystis jiroveciCytome galovirusHepatitis BListeria	AspergillusNocardiaBK Virus (polyoma)Herpes Zoster

Solution to Question 5:

Sabin-Feldman dye test is used for the diagnosis of toxoplasmosis.

It is based on the principle that the presence of anti-toxoplasma antibodies in the patient's serum inhibits the staining of trophozoites by alkaline methylene blue dye. Toxoplasma trophozoites and complement are added to the patient's serum which is then incubated for an hour at 37°C. After incubation, methylene blue dye is added.

• If anti-toxoplasma antibodies are present in the serum, these are activated by complement and the parasite membrane is lysed. Thus, the trophozoites are not stained (positive result).

• If there are no antibodies, trophozoites will take up the stain and appear blue under the microscope (negative result).

The dilution of the test serum at which 50% of the tachyzoites are not stained is reported as the antibody titer of the test serum. The test is highly sensitive and specific. However, it cannot differentiate between recent or past infections as it detects only IgG antibodies.

Solution to Question 6:

HBsAg (Hepatitis B surface antigen) is also known as the Australia antigen. It is called so because it was first isolated from the serum of an Australian person.

Sera from hepatitis B patients when observed under an electron microscope show three types of particles:

• A spherical particle: 22nm, most abundant, contains only HBsAg.

• Filamentous or tubular particle: 22nm, contains only HBsAg.

• A double-walled spherical structure: 42nm diameter, is the complete Hepatitis B virus. It was described by Dane and is called Dane particle.



Solution to Question 7:

Bacterial transduction occurs by bacteriophages.

Transduction is a process of acquiring bacterial DNA from a bacteriophage (virus) that has incorporated bacterial DNA from previous host bacteria.

Other methods of exchange of genetic material are transformation and conjugation. Transformation involves the uptake of genetic material from the environment, which is released by other bacteria. Conjugation is the direct transfer of DNA from one bacterial cell to another bacterial cell through a sex pilus or bridge. The transferred DNA is a plasmid, a circle of DNA that is distinct from the main bacterial chromosome.



Solution to Question 8:

Rhizopus is the most common fungus associated with orbital cellulitis in patients with diabetic ketoacidosis.

Diabetic patients are immunocompromised and so prone to develop opportunistic infections like mucormycosis - caused by Rhizopus and Mucor spp. Rhino-orbital- cerebral disease is the most common form of mucormycosis. It mostly occurs in diabetics. Patients initially present with eye or facial pain, facial numbness, conjunctival suffusion and swelling, and blurry vision. Infection then spreads from the ethmoid sinus to the orbit, leading to proptosis and orbital cellulitis.

Rhizopus is angioinvasive, and thus the infection may spread to the frontal lobe and cavernous sinus, leading to septicemia and thromboses. The sinus tissue in such patients often develops into a black necrotic eschar.

Solution to Question 9:

The adult form of echinococcus is seen in dogs and other canids, which are its definitive hosts.

The larval stage is passed in intermediate hosts such as sheep, cattle, goat, swine, horses, camels, and man. Man acts as an accidental intermediate host (dead end). Human infection follows the ingestion of eggs due to the intimate handling of dogs or by eating raw vegetables or other food contaminated with dog feces.

Note: Humans can become accidental definitive hosts, second intermediate hosts, or paratenic hosts for spirometra. These are cestodes belonging to the Diphyllobothriidae family. They cause a condition called sparagnosis.

The image below shows the life cycle of Echinococcus granulosus.



Solution to Question 10:

Among the given options, lactobacilli is the one that can grow in an acidic pH.

It is a gram-positive, facultatively anaerobic, non-motile, non–spore-forming bacterium. Lactobacillus derives lactic acid from glucose, creating an acidic environment, and can grow in an acidic medium. It is found in the vagina as part of its normal microbiota.

Solution to Question 11:

The image shown is of the Leishman-Donovan bodies whose presence are diagnostic of Kala Azar. The vector of Kala Azar is a sandfly.

Leishman-Donovan bodies are amastigotes with a double dot appearance containing a nucleus and small rod-like kinetoplast. They are present inside the macrophage.

Solution to Question 12:

TB test done on a whole blood sample is the interferon-gamma release assay. It has been used for the diagnosis of latent tuberculosis in patients with HIV(who frequently have a negative Mantoux test).

Interferon-gamma release assays include:

• QuantiFERON-TB Gold

• T-SPOT.TB

QuantiFERON-TB Gold quantitates the amount of IFN-γ produced by CD4+ T cells in response to the antigens from Mycobacterium tuberculosis. QuantiFERON-TB Gold Plus (QFT- Plus) is an improved version of the QuantiFERON-TB Gold test. QuantiFERON-TB Gold detects only TB-specific CD4+ responses. However, QuantiFeron-TB Gold Plus detects both CD8+ and CD4+ T-cell responses to TB-specific antigens.

T-SPOT.TB determines the total number of individual effector T cells expressing IFN-γ.

The TrueNat TB test is a new molecular test that can diagnose TB in one hour as well as test for rifampicin resistance. This test uses a sputum sample taken from the patient. Only about 0.5 ml of the sample is required. Gene Xpert is also a molecular method of testing that cannot use blood as a sample. (Option A and B)

Solution to Question 13:

Multiple abscesses with discharging sinuses, demonstrating sulfur granules in pus, are characteristic of actinomycosis caused by Actinomyces.

Actinomycosis is a chronic granulomatous infection characterized by indurated swellings in the connective tissue with suppuration and discharge of sulfur granules. The most common causative agent is Actinomyces israelli.

Various clinical forms of actinomycosis include cervicofacial, thoracic, abdominal, and pelvic actinomycosis. Among them, oral-cervicofacial actinomycosis is most common, presenting as a soft tissue swelling, abscess, or mass lesion involving the angle of the jaw.

Solution to Question 14:

Cutaneous larva migrans is most commonly caused by Ancylostoma braziliense.

Cutaneous larva migrans or ground itch or creeping eruption is caused by nematode larvae that infect by skin penetration.

The most common cause is non-human species of Hookworm Ancylostoma braziliense and A. caninum.

Necator americanus and Strongyloides sp also cause cutaneous larva migrans, and Toxocara canis usually causes viscreal and ocular larva migrans.

Differences between cutaneous larva migrans and visceral larva migrans:

	Cutaneous larva migrans	Visceral larva migrans
Tissue involved	Skin	Various organs of the body li ke the liver, lungs, and eyes
Infecting Organis m	Mostly by non-human nemat odes	Mostly by non-human nemat odes (mainly by dog and cat)
Portal of Entry	Penetration of skin	Ingestion of infected eggs
Eosinophilia	Mild	Persistent high
Serodiagnosis	Not developed	Well developed
Treatment	Thiabendazole	Diethylcarbamazine and pred nisolone

Solution to Question 15:

Hyperacute graft rejection is an example of type II hypersensitivity reaction.

It is mediated by preformed antibodies. It is a rare event that occurs in kidney transplantation.

Graft rejectio n	Time taken for rejection	Immune mechanisms involve d	Type of hypersensitivity
Hyperacute	Minutes to hours	Preformed antibodies	II
Acute	Weeks to months	Cytotoxic T-cell mediated	IV
Chronic	Months to years	Chronic DTH mediated and Ab mediated	IV

Solution to Question 16:

Helicobacter pylori produces urease.

Urease-producing organisms: (Mnemonic-PUNCH)

• Proteus

• Ureaplasma urealyticum

• Nocardia

• Cryptococcus, Corynebacterium urealyticum

• H. pylori

Solution to Question 17:

Mycoplasma are inherently resistant to beta-lactam antibiotics.

Mycoplasma lack cell wall, and are therefore inherently resistant to antibiotics acting on the bacterial cell wall e.g. beta-lactams (penicillins, cephalosporins).

Solution to Question 18:

Pasteurization: Milk is heated at either

• 63°C for 30 minutes: the Holder method or

• 72°C for 15–20 seconds: the flash process

followed by cooling quickly to 13°C or lower.

All the non-sporing pathogens are killed except Coxiella burnetti, which may survive in Holder method.

Microbiology NEET 2019

Question 1:

BCYE medium is used to culture:

a) Leptospira

b) Bacillus cereus

c) Campylobacter

d) Legionella

Question 2:

A 16-year-old boy is admitted with fever, icterus, conjunctival suffusion, and hematuria for 15 days. Which serological test should be done for diagnosis?

a) Widal test

b) Microscopic agglutination test

c) Weil felix reaction

d) Paul Bunnell test

Question 3:

A patient presents with gastrointestinal problems including abdominal pain and distension, bloody and mucus-filled diarrhea, and tenesmus, with rectal prolapse. A stool examination reveals the presence of typical barrel-shaped eggs. What is the possible causative agent?

a) Giardia lamblia

b) Trichuris trichiura

c) Ascaris lumbricoides

d) Ancylostoma duodenale

Question 4:

Viral infection mimicking erythroblastosis fetalis is:

a) HSV

b) Epstein-Barr virus

c) Picornavirus

d) Parvovirus B19

Question 5:

An 11-year-old boy was brought to the pediatric OPD with multiple abscesses over his face, chest, and back. The child has a history of recurrent■ respiratory infections. On examination, he has atopic excoriating skin and multiple cold abscesses on his back. What is the most probable diagnosis?

a) Hyper IgM syndrome

b) Wiskott-Aldrich syndrome

c) Hyper IgE syndrome

d) Carcinoid syndrome

Question 6:

Cowdry type A inclusions are seen in:

a) Herpes simplex

b) Rabies

c) Molluscum contagiosum

d) Adenovirus

Question 7:

Which of the following statements is false regarding MHC molecules?

a) MHC class II presents peptide antigen to CD4 T cells

b) The action of MHC II occurs via endocytic pathway

c) MHC Class II presents endogenous antigens

d) MHC II is present on antigen presenting cells

Question 8:

Which of the following investigations does not use antibody marking?

a) ELISA

b) Radioimmunoassay

c) Immunofluorescence

d) Hemagglutination

Question 9:

A 3-year-old child presents with a history of repeated sinopulmonary infections caused by encapsulated organisms. Which of the following is most likely to be deficient in this case?

a) IgG1

b) IgG2

c) IgG3

d) IgG4

Question 10:

A veterinarian developed an ulcer with central black eschar over the forearm as shown in the image. Which of the following is the most probable causative organism?

a) Pseudomonas aeruginosa

b) Yersinia pestis

c) Bacillus anthracis

d) Francisella tularensis

Question 11:

Which of the following is not used as an immediate test in the screening of urinary tract infections (UTI)?

a) Pour plate culture

b) Griess Nitrite test

c) Leukocyte esterase test

d) Wet mount examination

Question 12:

In which of the following is the disc diffusion method for antibiotic susceptibility testing applied?

a) Agar dilution method

b) Kirby-Bauer method

c) Broth dilution method

d) PCR based assay

Question 13:

What is the investigation of choice for neurosyphilis?

a) Venereal disease research laboratory (VDRL) test

b) FTA-ABS- Fluorescent Treponemal Antibody Absorption test

c) RPR- Rapid Plasma Reagin test

d) TPI-Treponema pallidum immobilization test

Question 14:

Contact based isolation is employed for which of the following diseases?

a) Diphtheria

b) MRSA

c) Mumps

d) Asthma

Question 15:

An 18 year old female presents with severe headache and high fever of 101°F. On examination, papilloedema and nuchal rigidity was present. Gram staining of the CSF is shown below. Causative organism is



a) Haemophilus influenza type B

b) Neisseria meningitidis

c) Streptococcus pneumoniae

d) Listeria monocytogenes

Question 16:

What is the stain used for identifying cryptococcus?

a) Giemsa stain

b) Ziehl-Nielsen stain

c) Gram staining

d) Mucicarmine

Question 17:

Which of the following organisms follow a bird-arthropod-human transmission route?

a) Malaria

b) Japanese encephalitis

c) Plague

d) Paragonimiasis

Question 18:

A 23-year-old patient has a history of travel to China and the consumption of pickled crab. He now complains of rusty sputum and cough. What is the most likely causative organism?

a) Fasciola buski

b) Fasciola hepatica

c) Paragonimus westermani

d) Schistosoma japonicum

Question 19:

Which type of virus is rubella?

a) Flavivirus

b) Togavirus

c) Paramyxovirus

d) Orthomyxovirus

Question 20:

A 30 year old female developed pruritic papules with excoriated plaques in the interdigital web spaces. Later the lesion spread to the groin and under-surface of the breasts. The lady gives a history of intense pruritus which is more severe at night. Given below is the organism responsible for this condition. Identify the wrong statement about the same.



a) Life history has 4 stages

b) Larvae have 3 pairs of legs

c) The entire life cycle takes about 15 days

d) Male parasite burrows into the epidermis

Question 21:

A HIV patient presents with persistent diarrhea. Microscopic examination of the stool is given below. Identify the causative organism:



a) Cryptosporidium

b) Cyclospora

c) Microspora

d) Cystoisospora

Question 22:

A patient is brought to the hospital, with complaints of seizures and the MRI image of the brain is given below. The disease is most probably caused by?



a) Cysticercosis cellulosae

b) T. saginata

c) Cysticerocosis bovis

d) T. asiatica

Question 23:

Vaccine that requires strain change due to evolving organism is?

a) Influenza

b) Diphtheria

c) Polio

d) Typhoid

Answer Key

Question No.	Correct Option
1	d
2	b
3	b
4	d
5	c
6	a
7	c
8	d
9	b
10	c
11	a
12	b
13	a
14	b
15	b
16	d
17	b
18	c
19	b
20	d
21	a
22	a
23	a

Detailed Explanations

Solution to Question 1:

BCYE medium (buffered charcoal yeast extract agar) is used to culture Legionella pneumophila.

Legionella pneumophila has fastidious growth requirements and grows on complex media such as BCYE medium with L-cysteine and antibiotic supplements (e.g., polymyxin B, cefazolin, vancomycin). The charcoal acts as a detoxifying agent. It grows slowly, and the colonies may take 3–6 days to appear.

Other options:

Option A: Leptospira is cultured on EMJH (Ellinghausen, McCullough, Johnson and Harris) medium

Option B: Bacillus cereus is cultured on PEMBA (Polymyxin pyruvate egg-yolk mannitol–bromothymol blue) agar

Option C: Campylobacter is cultured on Skirrow's, Campy BAP or Butzler's agar.

Solution to Question 2:

A history of fever with liver and kidney involvement should arouse suspicion of leptospirosis (Weil's disease). A microscopic agglutination test is used for the diagnosis of leptospirosis.

Antibodies against leptospirosis appear in the blood 5-7 days after infection. Serotype-specific tests for leptospirosis includes:

• Microscopic agglutination test

• Macroscopic agglutination test

Other options:

Option A: Widal test is used to screen for typhoid fever

Option C: Weil-Felix reaction is used to diagnose rickettsial fevers

Option D: Paul-Bunnell test is the investigation of choice for infectious mononucleosis.

Solution to Question 3:

The given clinical picture and the barrel-shaped eggs are suggestive of infection with Trichuris trichiura (whipworm).

The characteristic features of whipworm eggs are:

• Eggs are barrel-shaped in the middle with a projecting mucus plug at each pole containing an unsegmented ovum. The plugs are colorless.

• They float in the saturated salt solution.

• It has a triple shell, the outermost layer of which is stained brown.

The below image shows the egg of Trichuris trichiura.



In moderate infections of 20 adult worms or less, the worms are confined to the caecum and ascending colon. The symptoms include epigastric and lower abdominal pain, diarrhea (rarely bloody), vomiting, flatulence and distention, weight loss, and increased blood and iron loss.

In heavy infections of 200 adults worms or more (trichuris dysentery syndrome), the worms are in the distal colon and rectum. The symptoms include bloody diarrhea with profuse mucus, abdominal pain, tenesmus, weight loss leading to cachexia, severe anemia, and rectal prolapse. On examination, clubbing may be present. Blood investigations may show moderate eosinophilia.

Chronic infection leads to growth retardation and nutritional deficiencies. Around 0.005 ml of blood is lost per day per worm. In heavy infections, as much as 4 to 5 ml can be lost per day per worm.

Treatment is mebendazole or albendazole.

Solution to Question 4:

Parvovirus B19 infection mimics erythroblastosis fetalis.

Parvovirus B19, belonging to the parvovirus group consisting of the smallest animal DNA viruses, primarily infect immature cells of the erythroid lineage. It is associated with erythema infectiosum (fifth disease), transient aplastic crisis, pure red cell aplasia and hydrops fetalis.

During pregnancy, infection with parvovirus B19 can lead to severe anemia, non-immune hydrops fetalis and fetal death. This is because parvovirus preferentially infects RBC progenitor cells. This mimics erythroblastosis fetalis, in which maternal antibodies against fetal RBCs cause varying degrees of anemia and immune-mediated hydrops fetalis in sever cases.

Picornavirus, Epstein-Barr virus and HSV infections do not mimic erythroblastosis fetalis.

Solution to Question 5:

From the above clinical history and examination, the most probable diagnosis is hyper IgE syndrome (Job's syndrome).

It is an autosomal dominant primary immunodeficiency disease caused by a mutation of STAT3 gene. Neutrophils do not generate an immune response, probably due to a defect in chemotaxis. Patients present with recurrent staphylococcal skin abscesses, recurrent lung infections (pneumatoceles), eczema, eosinophilia and high serum levels of lgE.

Solution to Question 6:

Cowdry type A inclusions are seen in herpes simplex infection.

The classic Cowdry type A (Lipshutz) inclusion appears as a light purple, homogeneous intranuclear structure surrounded by a clear halo. The inclusions stain with antibodies to HSV, permitting a rapid, specific diagnosis of HSV infection.

HSV inclusion bodies



©MARROW

Solution to Question 7:

MHC class II presents exogenous antigens while MHC class I presents endogenous antigens.



Solution to Question 8:

The hemagglutination test is not based on the principle of antibody marking.

In tests based on the principle of antibody marking, the antigens or antibodies are labelled or marked with enzymes, fluorescent dyes, radioisotopes etc. In radioimmunoassay, fixed amounts of antibody and radiolabelled antigen react in the presence of unlabelled antigen. The labelled and unlabelled antigens compete for binding sites on the antibody. Thus, the concentration of the test antigen can be calculated from the ratio of the bound and total antigen counts using a standard dose-response curve. In immunofluorescence, fluorescent dyes are conjugated to antibodies or antigens. The labelled antibodies or antigens can then be used to identify antigens and antibodies, respectively. In ELISA, an adsorbing material - cellulose, agarose, polystyrene, polyvinvyl etc - is used to bind to either antigens or antibodies.

When antibodies bind to antigens on the surface of red blood cells, the resultant clumping reaction is referred to as hemagglutination. It is routinely performed to type red blood cells in ABO blood-typing. The image below shows hemagglutination.



Note: If no hemagglutination occurs it means the patient's serum has anti-haemagglutinin antibodies that bind to hemagglutinin and prevent the clumping reaction. This hemagglutination inhibition reaction is a method of detection of viruses and antiviral-antibodies and is used to make a diagnosis of influenza.

Solution to Question 9:

Deficiency of IgG2 results in recurrent sinopulmonary infections with encapsulated organisms.

IgG is the only maternal immunoglobulin that is normally transported across the placenta and provides natural passive immunity to the newborn. IgG1 is the antibody maximally transported across the placenta, whereas IgG2 has the least transplacental movement.

Deficiencies of other IgG:

• IgG1- bacterial infections (diphtheria, tetanus)(Option A)

• IgG3- viral infections (Option C)

• IgG4- parasitic infections (Option D)

Solution to Question 10:

The given clinical scenario is suggestive of cutaneous anthrax which is caused by Bacillus anthracis.

Human anthrax presents in one of three ways:

• Cutaneous Anthrax(hide porter's disease)- Black eschar at the base, with a very prominent inflammatory ring around the eschar, is referred to as a malignant pustule.

• Pulmonary Anthrax(Wool sorter's disease)- Hemorrhagic pneumonia.

• Intestinal Anthrax- Hemorrhagic diarrhoea. It occurs mainly in primitive communities that eat the carcasses of animals dying of anthrax.

Robert Koch validated the germ theory of disease after discovering the anthrax bacillus as the cause of anthrax. He also discovered the causative agents of tuberculosis and cholera. Hence, he is known as the Father of Bacteriology.

Koch's postulates, which link specific microorganisms to specific diseases, state that the microorganism must be:

• Found in abundance in diseased organisms, but not in healthy organisms.

• Isolated from a diseased organism and grown in pure culture

• Able to cause disease when inoculated into a healthy organism

• Re-isolated from the diseased experimental host and identical to the original specific causative agent

Solution to Question 11:

Pour plate culture (quantitative test) is not used as an immediate test for screening of UTI. However it is considered the best diagnostic test for confirmation of UTI due it's high sensitivity and specificity.

The screening tests for UTI are as follows:

• Wet mount examination: It is done to demonstrate pus cells in urine.

• Leukocyte esterase test: It is a rapid and cheaper method, and detects leukocyte esterases secreted by pus cells present in urine.

• Nitrate reduction test (Griess test): Nitrate reducing bacteria like E. coli give a positive result.

• Catalase test: E. coli and other bacteria that produce the enzyme catalase give a positive result.

Quantitative Culture:

• Standardized loop technique

• Pour plate method

Solution to Question 12:

The disc diffusion method for antibiotic susceptibility testing is applied in the Kirby-Bauer method.

In Kirby-Bauer's disc diffusion method, a suitable standard dilution of a broth culture of the test bacterium is inoculated on the surface of a solid medium (Cation-adjusted Mueller-Hinton agar) as a lawn culture. After drying, various antibiotic discs are placed on its surface. Zones of inhibition around the disc are recorded and interpreted according to the zone diameters available in the tables of the Clinical and Laboratory Standards Institute (CLSI) guidelines. The results obtained are qualitative (resistant, sensitive, or partially sensitive).

The image below shows Kirby-Bauer disc diffusion method.

Laboratory testing for antibiotic sensitivity can be done by:

• Diffusion methods: The drug diffuses through a solid medium so that a gradient is established. The concentration is highest near the site of drug application and decreases with distance. The test bacterium is cultured on the medium and its sensitivity to the antibiotic is determined by the zone of growth inhibition. Examples include Kirby–Bauer method and Stokes method.

• Dilution methods: Serial dilutions of the drug are prepared and inoculated with bacteria. Examples include tube dilution method and agar dilution method

• Diffusion and dilution method: E–test (Epsilometer)

Solution to Question 13:

The investigation of choice for neurosyphilis is VDRL.

The CSF-VDRL test is highly specific and, when reactive, is considered diagnostic of neurosyphilis.

Involvement of the CNS in syphilis is detected by examination of CSF for:

• Mononuclear pleocytosis (>5 white blood cells/µL).

• Increased protein concentration (>45 mg/dL).

• CSF VDRL reactivity.

Solution to Question 14:

Contact based isolation is employed for patients who are infected with MRSA (methicillin-resistant Staphylococcus aureus).

The three categories of transmission-based precautions include:

• Contact-based isolation is used for infections, diseases, or germs that are spread by touching the patient or items in the room. Examples include MRSA, vancomycin-resistant enterococci (VRE), diarrheal illnesses, open wounds, and respiratory syncytial virus (RSV).

• Droplet-based isolation is employed for diphtheria and mumps.

• Airborne precautions are required to protect against conditions like measles, severe acute respiratory syndrome (SARS), varicella (chickenpox), and Mycobacterium tuberculosis.

Solution to Question 15:

History and physical examination findings are suggestive of meningitis.

The given image shows intracellular gram negative diplococci i.e. Neisseria meningitidis.

Option a:Haemophilus influenzae type B is a gram negative bacillus. In CSF, long bacillary filamentous forms predominate.



Option c: Streptococcus pneumoniae is a gram positive, capsulated, flame shaped or lanceolate shaped diplococci.



Option d: Listeria monocytogenes is a gram positive bacillus.



Solution to Question 16:

Mucicarmine stain is used in the identification of cryptococcus.

Mucicarmine stain helps in the detection of the cryptococcal capsule, as it stains mucins of epithelial origin (PAS, alcian blue). Given below is an image of Mucicarmine stained specimen of lung tissue containing cryptococci:



Solution to Question 17:

Japanese encephalitis follows bird-arthropod-human transmission.

JE virus infects several extra-human hosts, e.g., pigs and birds (like pond herons). The basic cycles of transmission are:

(a)Pig → Mosquito → Pig

(b) Bird → Mosquito → Bird

Humans are incidental "dead-end" hosts and are infected by the bite of infected Culex mosquitoes.

Solution to Question 18:

The given clinical scenario is suggestive of infection with Paragonimus westermani.

Paragonimus westermani is a trematode. The transmission to humans occurs by eating raw or undercooked crab or crayfish containing metacercariae. The metacercariae excyst in the human gut and young worms migrate to the lungs, where they become encapsulated in lung tissue.

The patient complains of cough, chest pain, and hemoptysis. The viscous sputum is speckled with golden-brown eggs.

Praziquantel (25 mg/ kg TDS for 1-2 days) is the drug of choice.

The image below shows the life cycle of Paragonimus westermani.



Solution to Question 19:

Rubella belongs to the Togaviridae family.

Rubella (German measles)

• Causative organism belongs to family Togaviridae

• Period of infectivity is -5 to +6 days after the onset of the rash

• Incubation period: 2-3 weeks

• Rash appears on day 1

• Forchheimer spots are seen, which are tiny, rose-colored lesions in the oropharynx

• Associated with post infectious thrombocytopenia

• Most serious complication: Encephalitis

• Vaccination: 27/3 live attenuated

Congenital rubella syndrome:

• When a woman in the 1st trimester gets infected with rubella, the resulting complications may include miscarriage, fetal death, premature delivery, or live birth with congenital defects.

• The commonest malformations caused by rubella are cardiac defects, cataract, and deafness, which constitute the classical congenital rubella syndrome.

• Thrombocytopenic purpura present in the babies gave rise to the term “blueberry muffin syndrome.”

Expanded rubella syndrome:

• Some babies present with several other features like hepatosplenomegaly, thrombocytopenic purpura, myocarditis, and bone lesions constituting the expanded rubella syndrome.

• This term also refers to a baby with metastatic neuroblastoma.

Note: Rubella has been reclassified into the family of Matonaviridae in 2020. This is an MCQ from NEET 2019, hence the best answer out of the given options would be Togaviridae.

Solution to Question 20:

The clinical background is probably suggestive Scabies. The given image depicts Itch mite also called Sarcoptes scabiei. The female parasite burrows into the stratum corneum layer of the epidermis and lays eggs in the burrow.

A single female may lay about 30 eggs at a rate of 2 to 3 per day, and ultimately dies at the end of the burrow.

The life cycle has 4 stages: Egg, Larva, Nymph, and Adult. The adult mites live for 1 to 2 months.

Itching is a hypersensitivity reaction to the eggs and dead parasite in the epidermis.

Permethrin cream - 5% is the treatment of choice.

Solution to Question 21:

Diarrhea in an HIV patient is suggestive of a coccidean parasitic infestation. Kinyoun stain - positive oocysts as seen in the given image is suggestive of Cryptosporidium.

Points in favor:

• HIV patient with diarrhea

• Oocysts are round, contains four sporozites which is uniformly acid fast on staining.

Comparison of coccidian parasites causing diarrhea in an immunocompromised host:

Treatment is fluid replacement and cotrimoxazole.

Property	Cryptosporidium	Cyclospora	Isospora
Sporulated oocyst	4-6 µm, round contains fo ur sporozoites	8-12 µm, round contains 2 sporocyst each having tw o sporozoites	23-36 µm, oval contain tw o sporocyst, each having f our sporozoites.
Acid-fast detection	Uniformly acid fast	Variable acid fast	Uniformly acid fast
Sporulation of the ooc yst	Occurs inside host cell	Occurs in soil	Occurs in soil
Diagnostic form	Sporulated cyst	Unsporulated cyst	Unsporulated oocyst



- ***Cryptosporidium parvum***

Oocyst with 4 sporozoites
Size: 4 -5 um
Staining : Ziehl-Neelsen or Auramin

- ***Cyclospora cayetanensis***

Young oocyst

Mature oocyst with 2 sporocysts containing
2 sporozoites each
Size: 8 - 10 um
Staining : Ziehl-Neelsen or Auramin

- ***Isospora belli***



Young oocyst



Mature oocyst with 2 sporocysts containing
4 sporozoites each
Size: up to 20 x 30 um

Solution to Question 22:

The given clinical scenario and MRI brain showing scolex within multiple cystic lesions are suggestive of neurocysticercosis caused by the larval form of T. solium called Cysticercosis cellulosae.

Neurocysticercosis is the most common central nervous system parasitic infection.

The clinical features of neurocysticercosis include epilepsy, features of raised intracranial tension, and evident hydrocephalus. Patients can also have psychiatric disturbances, transient paresis, and visual disturbances.

The CT and MRI findings seen in neurocysticercosis are:

• Scolex within a cyst

• Cystic lesions with or without ring enhancement

• Nodular calcifications

• Focal enhancing lesions

• Multilobulated cystic lesions in the subarachnoid space.

Note: About 70% of adult-onset epilepsy is due to neurocysticercosis.

Albendazole is the drug of choice for neurocysticercosis. Glucocorticoid therapy is usually begun before initiating albendazole (an antiparasitic drug). This is continued for several days after the institution of therapy to reduce the incidence of side effects from inflammatory reactions to dead and dying cysticerci of Taenia solium.

Solution to Question 23:

Influenza vaccine requires periodic strain change due to antigenic change in the surface glycoprotein of the virus.

The change may be gradual (antigenic drift) or drastic (antigenic shift).

Microbiology NEET 2020

Question 1:

A 35-year-old HIV positive■ patient presents with mucosal lesions in the mouth as shown in the image below. On microscopy, budding yeasts and pseudohyphae are seen. What is the likely diagnosis?

a) Oral candidiasis

b) Hairy leukoplakia

c) Lichen planus

d) Diphtheria

Question 2:

Zika virus is transmitted by:

a) Aedes aegypti

b) Culex

c) Anopheles

d) Phlebotomus papatasii

Question 3:

Which of the following organisms causes disease following traumatic inoculation into the skin?

a) Sporothrix schenckii

b) Blastomyces dermatitidis

c) Coccidioides immitis

d) Paracoccidioides brasiliensis

Question 4:

Identify the class of immunoglobulin■ shown in the image below.



a) Ig A

b) Ig G

c) Ig M

d) Ig E

Question 5:

A 14-year-old boy presents with a history of vomiting within 3 hours of consumption of food at a party. Which of the following organism is most likely to cause these symptoms?

a) Staphylococcus aureus

b) Clostridium botulinum

c) Clostridium perfringes

d) Salmonella typhi

Question 6:

Which of the following organisms produces superantigen?

a) Staphylococcus epidermidis

b) Streptococcus pyogenes

c) Clostridium perfringens

d) Vibrio cholera

Question 7:

A female patient who is from the north-eastern part of the country came to the hospital with complaints of progressive swelling of the upper limbs. She has a history of coronary artery disease for which she underwent percutaneous coronary intervention. On examination, the upper limbs are more swollen proximally than distally. What is the next best step in the management of this patient?

a) Echocardiography

b) Night blood test for worms

c) Duplex venous scan of limbs

d) ECG

Question 8:

Which of the following helminthic infections, that is commonly transmitted by soil, is characterized by■ the egg in stool as shown in the image below?



a) Trichuris trichiura

b) Ancyclostoma duodenale

c) Necator americanus

d) Ascaris lumbricoides

Question 9:

A 30-year-old female from tribal area presented with fever for the last 3 days. On examination of the peripheral smear, a diagnosis of malaria was made. Which of the following is most likely to be the causative agent?



a) P. falciparum

b) P. malariae

c) P. ovale

d) P. vivax

Question 10:

The test kit shown in the image below works on the principle of:

©MARROW

a) ELISA

b) Immunochromatography

c) Immunofluroscence

d) Chemiluminiscence

Question 11:

A 22-year-old student presented with high-grade fever, and purpuric rash as shown in the image below. CSF sample showed gram-negative diplococci. Which of the following is the most likely aetiology?

a) Neisseria meningitidis

b) Pseudomonas aeruginosa

c) Streptococcus pneumoniae

d) E.coli

Question 12:

Which of the following is the predominant pathogen to infect renal transplant recipients one to four months following transplantation?

a) Cytomegalovirus

b) JC virus

c) BK virus

d) Epstein-Barr virus

Question 13:

Which of the following is not true about Japanese encephalitis?

a) Transmitted by Culex vishnui

b) Pig is the amplifier host

c) Man acts as the reservoir

d) Vaccinating pigs is helpful in disease control

Question 14:

Which of the following helps in bacterial adhesion?

a) Cytoplasmic membrane

b) Mesosomes

c) Fimbriae

d) Lipopolysaccharides

Question 15:

Which of the following is not involved in urethritis?

a) Trichomonas

b) Chlamydia

c) Gonococcus

d) H.ducreyi

Question 16:

A 30-year-old HIV-positive patient presented with a chronic history of watery diarrhea. Histopathology of the ■duodenum is shown in the image below. Which of the following is the causative organism?



a) Giardia lamblia

b) Entamoeba histolytica

c) Microsporidia

d) Cryptosopridia

Question 17:

What is the drug of choice for invasive pulmonary aspergillosis?

a) Itraconazole

b) Voriconazole

c) Lyophylised Amphotericin B

d) Posaconazole

Question 18:

Which of the following culture media is used to differentiate other E.coli species from E.coli O157: H7?

a) Mannitol salt agar

b) Xylose lysine deoxycholate agar

c) Sorbitol MacConkey agar

d) CLED medium

Question 19:

A 5 year old boy was suffering from low grade fever, inspiratory stridor and barking cough for past 5 days. His chest X-ray is given below. What is the cause of this condition?



a) Moraxella

b) Parainfluenza

c) Hemophilus influenza

d) Staphylococcus aureus

Question 20:

Which of the following is a bioterrorism category A agent?

a) Nipah virus

b) Anthrax

c) Coxiella

d) Brucella

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	a
5	a

6	b
7	b
8	a
9	a
10	b
11	a
12	a
13	c
14	c
15	d
16	a
17	b
18	c
19	b
20	b

Detailed Explanations

Solution to Question 1:

An HIV patient with white patches over the tonsils and posterior pharynx showing budding yeasts and pseudohyphae on microscopy suggests a diagnosis of oral candidiasis.

Oral candidiasis or thrush is a patchy, pseudomembranous lesion that develops over the tongue, lips, gum or palate. It is composed of epithelial cells, yeasts, and pseudohyphae and can be scraped off. The causative agent is usually Candida albicans. It occurs in almost all individuals with AIDS. Other risk factors include hyperglycemia, cellular immunodeficiency, corticosteroids or antibiotics. It is treated with topical nystatin, oral ketoconazole or fluconazole.

Option B: Hairy leukoplakia is a wart-like growth on the tongue, caused by Epstein-Barr virus. It is commonly seen as a white lesion on the lateral side of the tongue.

Option C: Lichen planus is a pruritic disease characterized by papules commonly seen over the flexor surfaces and trunk. Oral and vulvovaginal lesions may also occur. Oral lesions appear as lacy, white, and raised lesions.

Option D: Diphtheria is characterized by the presence of a grey pseudomembrane covering the tonsils and pharynx. It is caused by Corynebacterium diphtheriae.

Solution to Question 2:

Zika virus is transmitted by the bite of infected Aedes aegypti and Aedes albopictus.

Zika virus is an RNA virus belonging to the Flaviviridae group. The symptoms of Zika virus disease include fever, maculopapular rash, conjunctivitis, muscle and joint pain, malaise and headache. These symptoms are mild and usually last for 2-7 days. It can be misdiagnosed as dengue or influenza. In severe cases, it can lead to complications like Guillain-Barré syndrome, neuropathy, and myelitis.

Zika virus infection during pregnancy may lead to microcephaly or congenital Zika syndrome.

Solution to Question 3:

Sporothrix schenckii causes infection followed by traumatic inoculation into the skin.

It is a dimorphic fungus that lives on vegetation such as grasses, trees, rose bushes, etc. Following traumatic inoculation into the skin as in thorn prick, S schenckii causes sporotrichosis, a chronic granulomatous infection. This is usually further followed by spread along the draining lymphatics and lymph nodes as shown in the image below.



Subcutaneous mycoses are usually caused by traumatic inoculation of the causative fungus. These include:

Subcutaneous mycoses	Causative agent/s
Sporotrichosis	Sporothrix schenckii
Chromoblastomycosis	Phialophora verrucosaFonsecaea pedrosoiFonsecaea compactaRhinocladiella aquasperisaCladophialophora carrionii
Phaeohyphomycosis	Exophiala jeanselmeiPhialophora richardsiaeBipolaris spiciferaWangiella dermatitidis
Mycetoma	Pseudallescheria boydii (Anamorph Scedosporium apiospermum).Madurella mycetomatisMadurella griseaE jeanselmeiAcremonium falciforme

Solution to Question 4:

The above image shows a dimeric structure, characteristic of secretory IgA.

IgA immunoglobulin is the major antibody responsible for mucosal immunity. Its serum levels are usually low. It is predominantly present in the extravascular secretions. It is present in secretions such as milk, saliva, and tears, and in secretions of the respiratory, intestinal, and genital tracts.

When present in the serum, it exists as a monomer similar to IgG. However, when it is secreted, it exists as a dimer. The dimeric structure consists of two monomers, which are stabilized by two additional polypeptides:

• J chain, which stabilizes the molecule.

• Secretory component, which is incorporated into the IgA when transported through the epithelial cell.



Solution to Question 5:

Vomiting 2 hours after the consumption of food is likely food poisoning by Staphylococcus aureus.

Staphylococcus aureus can grow in carbohydrate and protein food sources, producing enterotoxins. There are 15 types, all of which are superantigens. They are preformed toxins that are relatively heat stable. They stimulate the vagus nerve, vomiting centre of the brain and intestinal peristaltic activity. Type A enterotoxin is responsible for most cases of food poisoning.

The common sources of infection are the foods that are not cooked after handling. These include sliced meats, puddings, pastries, and sandwiches. Ingestion of such contaminated food causes food poisoning within 30 minutes to 6 hours after consumption. Patients present with nausea, vomiting, stomach cramps, and diarrhoea.

Solution to Question 6:

Streptococcus pyogenes produces streptococcal pyrogenic exotoxin (SPE), which is a superantigen.

Superantigens are protein toxins that stimulate large numbers of T cells, irrespective of antigen specificity. They bind to the class II MHC molecule outside the peptide-binding groove and Vβ region of the TCR. Some examples of superantigens are staphylococcal enterotoxins, toxic shock syndrome toxin, and group A streptococcal pyrogenic exotoxin.

The below image shows the binding of superantigen to T-cell receptor.



Solution to Question 7:

The given clinical scenario is suggestive of filariasis. The next best step would be to test the night blood for worms since the filarial species is nocturnally periodic (between 10 pm and 4 am) in India.

Solution to Question 8:

The given image shows the egg of Trichuris trichiura (whipworm).

The characteristic features of whipworm eggs are:

• Barrel-shaped in the middle

• Projecting colourless mucus plug at each pole

• Contains an unsegmented ovum.

• Floast in the saturated salt solution.

• It has a triple shell. The outermost layer is stained brown.

Humans acquire infection by swallowing embryonated eggs in contaminated food or water. The eggs hatch in the small intestine, and the larvae produced pass into the cecum. The larvae become mature adults in 2-3 months. Adult females pass immature un-embryonated eggs in the stool. Thus, diagnosis of whipworm infection is done by microscopic identification of the eggs in a stool specimen.

Most infections are asymptomatic. Moderate whipworm infection (20 adult worms or less), the worms are confined to the caecum and ascending colon. The symptoms include epigastric and lower abdominal pain, diarrhoea (rarely bloody), vomiting, flatulence and distention, weight loss, and increased blood and iron loss. In heavy infections of 200 adult worms or more - trichuriasis

dysentery syndrome -, the worms are in the distal colon and rectum. The symptoms include bloody diarrhoea with profuse mucus, abdominal pain, tenesmus, weight loss leading to cachexia, severe anemia, and rectal prolapse. On examination, clubbing may be present. Blood investigations may show moderate eosinophilia.

Chronic infection leads to growth retardation and nutritional deficiencies. Around 0.005 ml of blood is lost per day per worm. In heavy infections, as much as 4 to 5 ml can be lost per day per worm. Treatment is mebendazole or albendazole.



Solution to Question 9:

The image shows ring forms/ early trophozoites in erythrocyte, which are binucleate (double chromatin) resembling stereo headphones in appearance, characteristic of P. falciparum.

There may be several rings present within a single erythrocyte. The rings are often attached to the margins of the red cells and are described as form applique or accole.

Solution to Question 10:

The above image shows a HbsAg card test kit which works on the principle of immunochromatography.

In immunochromatography, a nitrocellulose membrane is used to absorb the antigen from a specimen. The membrane is impregnated with anti-HBsAg antibody and colloidal gold dye. The specimen (serum) travels through the membrane by capillary action. If the serum contains HBaAg, a coloured reaction (in the form of a band) appears directly on the membrane at the test site. These assays are rapid and include a built-in internal control.

Solution to Question 11:

This patient has a high-grade fever and is a student, so he presumably spends a lot of time in crowded areas. The above image shows a purpuric rash. This patient's features are seen in meningococcemia caused by Neisseria meningitidis, which are gram-negative diplococci.

The organism enters the body through the nasopharynx by attaching to the epithelial cells with the help of pili. From the nasopharynx, they enter the bloodstream producing bacteremia, and the symptoms may resemble upper respiratory tract infection.

Fulminant meningococcemia is the severe form, where the patient presents with high-grade fever and hemorrhagic rash. There is thrombosis of several small blood vessels in various organs with perivascular infiltration and petechial hemorrhages. Some patients with fulminant meningococcemia may also develop disseminated intravascular coagulation, and circulatory collapse, a condition known as Waterhouse–Friderichsen syndrome.

Solution to Question 12:

The predominant pathogen to infect renal transplant recipients in the period around 1 to 4 months following the transplantation is cytomegalovirus (CMV).

CMV is the most common virus to affect recipients after organ transplantation (kidney, heart, lung etc). The maximal risk occurs between 1 and 4 months after transplantation (in the absence of prophylaxis). The most common CMV presentation in kidney transplant patients includes fever, leukopenia, hepatosplenomegaly, myalgia, and arthralgia. Antiviral prophylaxis with ganciclovir or valganciclovir is used in renal transplant recipients whenever indicated to prevent CMV disease.

Given below are the most common opportunistic infections in renal transplant recipients:

Peritransplant (less than 1 m onth)	Early (1-6 months)	Late (more than 6 months)
Wound infectionHerpes virus Oral candidiasisUrinary tract infection	Pneumocystis jiroveciCytome galovirusHepatitis BListeria	AspergillusNocardiaBK Virus (polyoma)Herpes Zoster

Solution to Question 13:

Humans are the dead-end hosts for the Japanese encephalitis virus.

Japanese encephalitis is an endemic viral infection transmitted by mosquitoes. There is an increased risk in travelers to rural areas. JE has an inactivated vero culture-derived vaccine, given as 2 primary series to adults who are travelling to endemic areas.

Other options:

Option A: The disease is transmitted by Culex tritaeniorhynchus and Culex vishnui (common in India) species.

Option B: Herons act as reservoir hosts and pigs as amplifier hosts.

Option D: The control measures include mosquito control, locating piggeries away from the human dwellings and vaccinating pigs.

The image below shows the life cycle of the Japanese B encephalitis virus.



Solution to Question 14:

Fimbriae help in bacterial adhesion.

Fimbriae, also called pili, are hair-like surface appendage that projects from the cell surface as straight filaments. They are primarily seen in Gram-negative bacteria. They are polymers of pilin. Fimbriae are classified based on the function as ordinary pili - for adhesion - and sex pili - involved in bacterial conjugation. The functions of pili are given below:

• Bacterial conjugation through sex pili, which transfer of genetic material occurs.

• Bacterial adhesion of symbiotic and pathogenic bacteria to host cells using adhesins.

• Resistance to phagocytosis.

• Antigenic pilin proteins are responsible for the immune response.

• Twitching motility to bacteria.

Solution to Question 15:

Haemophilus ducreyi does not cause urethritis. It causes chancroid (soft chancre), which presents as a genital ulcer.

Urethritis is classified into two types:

• Gonococcal urethritis - caused by Neisseria gonorrhoea.

• Non-gonococcal urethritis - caused by:

• Chlamydia trachomatis (most common)

• Ureaplasma urealyticum

• Adenovirus

• Haemophilus vaginalis

• Mycoplasma genitalium

• Tricomonas vaginalis (rare)

• Herpes simplex virus (rare)

Solution to Question 16:

The histopathology shows trophozoites of Giardia lamblia in the duodenal lumen (marked in arrows).



• Giardia lamblia, also called Giardia duodenalis, is the causative agent of giardiasis. It is the only common pathogenic protozoan found in the duodenum and jejunum of humans.

• Infection is acquired by the consumption of water or food contaminated with cysts and may also be transmitted by the faeco-oral route.

• It is weakly pathogenic in humans. Secretory IgA and IL-6 are responsible for clearing the infection. Hence, giardiasis usually occurs in immunosuppressed, agammaglobulinemia, or malnourished individuals.

• The symptoms include acute or chronic diarrhea, weight loss, malaise, abdominal cramps, distension, and weakness. The stools may be watery, greasy, bulky, or foul-smelling at various times during the course of the infection.

• Histopathology of the duodenum reveals several heart-shaped or pear-shaped trophozoites with two equally sized nuclei. The small intestinal morphology is usually normal. However, in heavy infections, there may be villous blunting and intraepithelial infiltrates.

Solution to Question 17:

Voriconazole is the drug of choice in the treatment of invasive pulmonary aspergillosis.

Voriconazole can also be used against Candida species (including glabrata and krusei), Scedosporium and Fusarium. The other options - itraconazole, posaconazole, lyophylised Amphotericin B - can be used as alternatives for the treatment of invasive pulmonary aspergillosis if voriconazole is unavailable.

Solution to Question 18:

E.coli O157: H7 can be differentiated from other E.coli species using sorbitol MacConkey agar.

There are more than 150 serotypes of Shiga toxin-producing E.Coli (STEC). Of these, O157:H7 is the most common type. Unlike most other E.Coli serotypes, the STEC O157:H7 does not use sorbitol and produces clear colonies on the Sorbitol MacConkey agar.

O157: H7 strains also are negative for β-glucuronidase using the substrate 4-methylumbelliferyl-β-glucuronide (MUG) test. More than 90 per cent of other E.coli tests positive with this substrate.

Solution to Question 19:

Barking cough is a feature of laryngotracheobronchitis, which is caused by the parainfluenza virus. The X-ray finding is the 'Steeple sign' due to subglottic narrowing.

The age of presentation of laryngotracheobronchitis ranges from 6 months-6 years. The clinical features include barking cough, inspiratory stridor, hoarseness, low-grade fever.

Management:

• Racemic epinephrine nebulization leads to the rapid improvement of upper airway obstruction. Epinephrine constricts pre-capillary arterioles in the upper airway mucosa and decreases capillary hydrostatic pressure, leading to fluid resorption and improvement in airway edema.

• Dexamethasone-by the least invasive route (Oral>IV>IM). It is indicated in all the grades of croup.

• Oxygen should be administered to children who are hypoxemic (SpO2 <92 %)

Antibiotics are not indicated in laryngotracheobronchitis.

Solution to Question 20:

Bacillus anthracis which causes anthrax is categorized under category A bioterrorism agent.

Microbiology INI-CET 2021

Question 1:

Which of the following causes aplastic crisis in hereditary spherocytosis?

a) Poxvirus

b) Ebstein barr virus

c) Adenovirus

d) Parvovirus

Question 2:

A stool sample is collected. The specimen is processed for examination and is labeled at different levels as A, B, C, and D. Identify the level at which parasitic ova will be seen when viewed under a microscope.



a) A

b) B

c) C

d) D

Question 3:

Identify the organism shown in this image.



a) Strongyloides stercoralis

b) Enterobius vermicularis

c) Ancylostoma duodenale

d) Ascaris lumbricoides

Question 4:

A patient presented with meningitis, and the CSF sample was subjected to gram staining and microscopy and appears as shown in the microscopic image given below. Which of the following features/tests will be characteristic of the organism?



a) Oxidase and catalase negative, does not ferment inulin

b) Ferments glucose and maltose but not inulin

c) Oxidase and catalase positive and does not ferment maltose or glucose

d) Catalase negative, optochin sensitive

Question 5:

Which cell is involved in type 4/delayed type hypersensitivity?

a) Cytotoxic T cell

b) Macrophage

c) Dendritic cell

d) Neutrophil

Question 6:

A 32-year-old laborer working at a construction site presented with fever and hemoptysis. The sputum sample collected for examination showed the following. The smear will be stained by which of the following sequences?



a) Gentian violet - iodine - alcohol - saffranin

b) Methylene blue - malachite green- acetic acid-water

c) Methanol-methylene blue-acid-water

d) Carbol fuschin - acid-alcohol - methylene blue

Question 7:

A 55-year-old patient had a tooth extraction three days back. He now presents with a mass in the submandibular region which is otherwise asymptomatic. There is no ulcer on the oral mucosa. On drainage of the discharge, it was foul-smelling with a lot of yellow granules. They showed gram-positive filamentous rods on grams staining. What is the most likely implicated pathogen?

a) Chromoblastomycosis

b) Mucormycosis

c) Actinomycetes israelli

d) Histoplasmosis

Question 8:

Which of the following components of the fungal cell wall is not seen in mucormycosis?



a) Mannoproteins

b) Beta 1, 3, glucan

c) Chitin

d) Ergosterol

Question 9:

Which of the following statements is True regarding trematodes?

a) Fasciola hepatica are seen in the intrahepatic biliary radicles

b) Schistosoma are monoecious

c) Fasciola hepatica acquired through contaminated water

d) Triclabendazole is the treatment of choice for all trematode infections

Question 10:

A 45-year-old man developed generalized muscle spasm with arching of his back as shown in the image, a few days after sustaining a puncture wound to his leg. The toxin responsible for this type of abnormal posturing acts by which of the following mechanisms?

a) Presynaptic inhibition of GABA and glycine release

b) Presynaptic inhibition of Ach receptors

c) Postsynaptic inhibition of GABA

d) Cleavage of SNARE proteins, preventing Ach release

Question 11:

A man with FUO has a job that requires traveling around the world. The bone marrow picture is shown below. Which of the following investigation would be the most appropriate in this case?



a) rk 39 antigen test, Leishmaniasis

b) pLDH, Plasmodium malaria

c) Biopsy, Histoplasmosis

d) ELISA, HIV

Question 12:

Identify the organism from the lifecycle shown in the image given below.



a) Cystoisospora

b) Cryptosporidium

c) Plasmodium knowlesi

d) Toxoplasma gondii

Question 13:

All of the following oncogenic viruses can integrate with the human genome to initiate carcinogenesis except?

a) HPV

b) HBV

c) HTLV

d) EBV

Question 14:

A patient presented with fever, dyspnea, palpitations, and tachycardia. The echocardiography revealed vegetation in the septal leaflets. Tissue sections from them reveal the following findings as shown in the image. What is the most likely implicated agent?



a) Pigmented septate hyphae at 90 degrees, Mucormycosis

b) Right angle branching septate hyphae, Fusarium

c) Acute angle branching hyphae, Rhizopus

d) Acute angle branching septate hyphae, Aspergillus

Question 15:

Which pathogen causes attachment-effacement lesion in the intestinal mucosa as shown in the image?



a) Enterotoxigenic Escherichia coli

b) Enteropathogenic Escherichia coli

c) Enteroinvasive Escherichia coli

d) Diffusely adherent Escherichia coli

Question 16:

Identify the component responsible for endotoxic shock in gram-negative bacteria.



a) 2

b) 4

c) 1

d) 3

Question 17:

A 6-year old boy complains of cough and throat pain. On examination of the oral cavity, a white pseudomembrane was visualized. A swab was taken from it and stained as given in the image below. What is the procedure for the following staining?

a) 1, 2, 5, 8

b) 6, 3, 8, 4

c) 7, 8, 4, 3

d) 3, 2, 1, 8

Question 18:

Identify the virus given in the following replication cycle.



a) Human immunodeficiency virus

b) Herpes simplex virus

c) Hepatitis B virus

d) Coronavirus

Question 19:

Which of the following is a xenodiagnostic method?

a) Rabbit ileal loop for enterotoxigenic Escherichia coli

b) Injecting Aedes thorax with blood of a suspected dengue patient

c) Injecting a hamster with splenic biopsy for diagnosis of leishmaniasis

d) Intradermal test on guinea pigs for toxigenicity of Corynebacterium diphtheria

Question 20:

A 24-year-old male patient presented to the outpatient department with cervical lymphadenopathy. As a part of the evaluation, the Paul-Bunnell test was done. What is the principle of this test?

a) Complement fixation

b) Super-antigens

c) Heterophile antibodies

d) Haptens

Question 21:

Match the following in the Japanese encephalitis virus life cycle.

a) 1-d, 2-c, 3-a, 4-b

b) 1-c, 2-a, 3-d, 4-b

c) 1-c, 2-d, 3-a, 4-b

d) 1-d, 2-c, 3-b, 4-a

Question 22:

Which of the following is a dematiaceous fungus?

a) Mucor species

b) Aspergillus niger

c) Cladosporium

d) Blastomyces dermatitidis

Question 23:

The strict anaerobe among the following is _____.

a) Burkholderia cepacia

b) Yersinia enterocolitica

c) Stenotrophomonas maltophilia

d) Prevotella

Question 24:

Match the following eggs to the organism.



a) 1-D, 2-C, 3-A, 4-B

b) 1-C, 2-D, 3-A, 4-B

c) 1-D, 2-C, 3-B, 4-A

d) 1-C, 2-D, 3-B, 4-A

Question 25:

A man presents with the following lesion after traumatic inoculation. Identify the causative organism.

a) Mycobacterium scrofulaceum

b) Sporothrix schenckii

c) Madurella mycetomatis

d) Phialophora verrucosa

Answer Key

Question No.	Correct Option
1	d
2	d
3	d
4	d
5	a
6	d
7	c
8	b
9	a
10	a
11	a
12	d
13	c
14	d
15	b
16	c
17	b
18	b
19	b
20	c
21	c
22	c
23	d
24	d
25	b

Detailed Explanations

Solution to Question 1:

Aplastic crisis in a child with hereditary spherocytosis is most likely due to infection of parvovirus. This aplastic crisis is due to the transient arrest of erythropoiesis where the parvovirus infects and kills the red cell precursors. This generally leads to a sudden worsening of anemia. Transfusions may be necessary to support the patient until the immune response clears the infection.

Parvovirus:

Parvovirus-B19 is a non-enveloped single-stranded DNA virus.

Clinically, parvovirus B-19 infection may have several presentations-

• Erythema infectiosum - Also known as the fifth disease. The incubation period after infection is typically around 4-14 days. The hallmark of erythema infectiosum is the characteristic rash, which appears initially as erythematous facial flushing, often described as a “slapped cheek” appearance. Later the rash spreads rapidly or concurrently to the trunk and proximal extremities as diffuse macular erythema. The child may be febrile.

• Transient aplastic crisis - It may be seen in patients with underlying chronic hemolytic anemia like sickle-cell disease, thalassemia, hereditary spherocytosis, and RBC enzyme deficiencies like pyruvate kinase deficiency. Patients with aplastic crisis appear ill with fever, malaise, and pale with tachycardia and tachypnea due to profound anemia. The rash may or may not be present. Children with sickle-cell disease may also develop vaso-occlusive crises.

• Pure red cell aplasia - Persistent B19V infection that results in pure red cell aplasia has been reported in a wide variety of immunosuppressed patients, including patients with congenital immunodeficiency, AIDS, and lymphoproliferative disorders and transplant recipients.

• Primary maternal infection is associated with non-immune fetal hydrops and intrauterine fetal demise

It is a clinical diagnosis not requiring investigations usually. However, in the case of pregnant women or with other manifestations an IgM ELISA or PCR assay may be done. It is self-limited and resolves on its own.

The given image shows the classical slapped cheek appearance of erythema infectiosum caused by parvovirus B19. During the rash period, the patient is generally non-infectious.

Solution to Question 2:

The parasitic ova will be seen in a specimen taken from level D.

The image shows a formol-ether sedimentation technique to concentrate the ova in the bottom (D) part of the tube. It is used for samples in which the concentration of parasitic ova is scanty. It is useful for all helminth ova/eggs and parasitic cysts. The ether adsorbs the fecal debris and floats. Formalin fixes and preserves the specimen.

This method is useful for all helminth eggs and protozoan cysts.

Solution to Question 3:

The given image shows large cylindrical worms with tapering ends, and thus represents Ascaris lumbricoides (roundworm).

Ascaris lumbricoides is the most common of human helminths and is distributed worldwide. It is the largest nematode parasite in the human intestine. The adult worms are large cylindrical worms, with tapering ends. The anterior end is more pointed than the posterior. It appears white outside the body.

Its natural host is humans. There is no intermediate host. Adult worms live in the small intestine - jejunum and ileum. Humans acquire infectionby consuming food or water contaminated with nightsoil or human feces containing eggs. The eggs contain the infective rhabditiform larva.

Solution to Question 4:

The given clinical vignette of meningitis with the grams stain slide showing gram-positive diplococci some in lancet-shaped and some arranged in chains narrows down the aetiological/ microbial agent to Pneumococcus / Streptococcus pneumoniae. It is catalase-negative and optochin sensitive.

The Gram stain would reveal Gram-positive, lancet-shaped diplococci, single cocci, or chains. Pneumococci undergo autolysis with the addition of surface-active agents like ox bile (10%) or sodium deoxycholate (2%). The Viridans streptococci do not lyse and are thus easily differentiated from pneumococci. The growth of pneumococci is inhibited by optochin (Ethyl hydrocuprein hydrochloride) while that of viridans streptococci is not. It is catalase-negative, ferments inulin, and optochin sensitive.

Pneumococci are commensals of the upper respiratory tract and can cause pneumonia, sinusitis, otitis, bronchitis, bacteremia, meningitis, peritonitis, and other infectious processes. The pneumococcal infection results in inflammation and exudation of fibrinous materials resulting in edema and consolidation. The onset of pneumococcal pneumonia is usually sudden, with fever, chills, and sharp pleural pain. The sputum is similar to the alveolar exudate, being characteristically bloody or rusty colored. Disseminated infection and bacteremia can present as meningitis, endocarditis, and septic arthritis.

The other diagnostic parameters include the C-polysaccharide found in the cell wall of all S. pneumoniae that can be detected in the urine and cerebrospinal fluid (CSF) and a positive Quellung reaction. Quellung reaction/ Neufeld reaction is used to identify capsulated organisms such as pneumococci, Neisseria meningitis, Haemophilus influenza, etc. When a suspension of a capsulated bacterium is mixed with its specific anticapsular serum and examined under the microscope, the capsule becomes very prominent and appears 'swollen' due to an increase in its refractivity.

High-dose penicillin G is effective in treating pneumococcal infection.

The culture showing alpha hemolysis on blood agar, with optochin (Op) sensitivity, and microscopy showing gram-positive, lancet-shaped diplococci,
suggests Streptococcus pneumoniae (pneumococcus).



The image shows a swollen capsule, which is demonstrated by the Quellung reaction



The flowchart demonstrates the algorithm used for identification of gram-positive organisms in the lab.



Solution to Question 5:

Cytotoxic T cells are involved in type 4/delayed type hypersensitivity

Solution to Question 6:

The given clinical vignette of fever and hemoptysis in a labourer with the sputum smear showing acid-fast bacilli is suggestive of tuberculosis. Mycobacterium tuberculosis, detected on sputum smear, is appreciated by Ziehl-Neelsen staining, which would includes carbol fuschin - acid - alcohol - methylene blue.

Acid-fast bacteria (like MTB) in a stained Ziehl-Neelsen (ZN) smear appear bright red against a blue background. Steps involved in ZN staining are:

• Place the sample on slide evenly spread it with rotational movement

• Air-dry the slide and fix it by passing it over a flame

• Apply carbol fuschin stain and heat to fix it.

• Wash with water to remove excess stain

• Decolourise with alcohol

• Apply Methylene blue/Malachite green stain

• Wash with water to remove excess stain

• View under the microscope

Kinyoun modified the classical ZN staining such that heat fixing is not required. It is called the cold method. It involves increasing the concentration of phenol acid and the duration of staining. The slide is then washed with water and decolourized with acid alcohol. Counterstaining with

Loeffler's methylene blue or malachite green is done for a minute and viewed under a microscope. Acid-fast bacteria in a stained smear appear bright red against a blue background.

The image attached below summarizes the steps.



Option A: It depicts the order of Gram staining, which was discovered by Hans Christian Gram in 1884.

• Primary staining with para-rosaniline dye (crystal violet, methyl violet or gentian violet).

• Application of dilute solution of iodine.

• Decolorisation with an organic solvent (ethanol, acetone, aniline).

• Counterstaining with the dye of contrasting colour (carbol fuchsin, safranine, neutral red).

Option B: It includes the steps for performing Albert's stain which is used for detected Corynebacterium diphtheriae.

• A heavy smear of the bacterial isolate is prepared on a glass slide and allowed to dry without fixing.

• Albert's stain is applied for (3-4) minutes. Then it is washed with water and Albert's iodine is applied for one minute.

• Wash with water and let dry with blotting paper.

• Examine the slide under oil immersion.

Corynebacterium diphtheriae are slender bacilli with an arrangement resembling the letters V or L. They have metachromatic granules at the poles of bacilli which are also called polar bodies, volutin, or Babes-Ernst granules. Albert staining is useful in the demonstration of metachromatic granules. The presence of metachromatic granules helps to distinguish C.diphtheriae from diphtheroids which lack them. The images below show Corynebacterium diphtheriae.



Solution to Question 7:

This is a case of Cervicofacial actinomycosis caused by Actinomyces israeli. It presents as a painless nodule that indurates with purulent discharge that contains sulfur granules giving it a yellow color. Sulfur granules are composed of a matrix of bacteria, calcium phosphate, and host tissue.

Actinomycosis is an indolent, slowly progressive suppurative and granulomatous infection that produces pyogenic lesions with interconnecting sinus tracts that contain sulfur granules. It is an anaerobic or microaerophilic bacteria.

It presents as one of the following clinical types:

• Cervicofacial actinomycosis- (most common site) fluctuant mass in the jaw area producing draining fistulas. (lumpy jaw)

• Thoracic actinomycosis- pulmonary infection with fever, cough, and purulent sputum. It usually occurs following aspiration or trauma leading to inoculation of the infective material.

• Abdominal actinomycosis- following ruptured appendix or ulcer. It may involve multiple organs.

• Pelvic actinomycosis- due to the colonization of intrauterine contraceptive devices.

• Actinomycosis can spread to bones and surrounding tissue and produce multiple discharging sinuses.

Diagnosis is by Gram staining of sulfur granules that shows a dense network of thin gram-positive filaments, surrounded by club-shaped structures. (sunray appearance). Culture shows the "molar tooth" appearance of the colonies. On solid media, A. israeli produces spidery colonies.

Penicillin G is the drug of choice and the duration of therapy is between 6-12 months. The other drugs that can be used include amoxicillin, erythromycin, tetracycline, doxycycline, clindamycin, and minocycline.

The image below shows a case of cervicofacial actinomycosis with discharging sinuses.

The image below shows the "molar tooth" appearance of the Actinomyces colonies.



The image below shows gram-positive filaments of Actinomyces.



The given image shows a patient with multiple cutaneous lesions caused by actinomycosis:

Solution to Question 8:

Beta-(1,3)-d-glucan (BG), a common component of many pathogenic fungi is absent in Zygomycetes/ Mucormycosis.

Fungal cell wall:

The fungal cell wall is rigid and determines the shape of the fungi. These protect them from osmotic and environmental stresses. They are composed of glycoproteins and lipids. Some sugar polymers such as chitin, glucans, and mannans are also found in the cell wall. All these components remain cross-linked to form a multilayered cell wall matrix. They exhibit pathobiological properties and are responsible for provoking an inflammatory response during

infections. These are also species-specific and release immunodominant antigens that could aid in immunological testing. In addition, some yeasts and molds are described as dematiaceous because their cell walls contain melanin which forms host defenses.

Clinical significance of Beta-1, 3-d-glucan:

G-test:

G-test is done for detection of α-3 glucan. It is helpful in establishing a diagnosis of candidiasis.

Tests for beta-d-glucan:

The presence of an elevated level of β-d-glucan in the serum can be seen in candidiasis, aspergillosis, and pneumonia due to Pneumocystis carnii pneumonia. The same test cannot be effectively applied in the evaluation of Mucorales. This assay is also not evaluated in routine in cases of cryptococcosis, not because of the absence of BG but because the capsule captures BG even before it is released into the bloodstream.

Effectiveness of echinocandins:

Echinocandins inhibit the synthesis of glucan synthase, since there is no much of a role for beta-glucans in mucormycosis, this class of drugs is seldom effective in the management of mucormycosis. The drug of choice for the management of mucormycosis is amphotericin-B.

Solution to Question 9:

Of the given statements regarding trematodes, the only true statement is that Fasciola hepatica is seen in the intrahepatic biliary radicles.

Schistosomes are dioecious, Fasciola hepatica is acquired through metacercariae encysted on raw aquatic vegetation and triclabendazole is not the treatment of choice for all trematode infections as praziquantel is used in the management of Schistosomiasis and lung fluke infections.

Trematodes are leaf-shaped unsegmented, flat, and broad helminths. The males are shorter and stouter than the females. The unique feature of flukes is the presence of two muscular cup-shaped suckers (hence called distomata)- the oral sucker surrounding the mouth at the anterior end and the ventral sucker or acetabulum in the middle, ventrally. Three of the most important pathognomic species include Schistosoma hematobium, Fasciola hepatica, and Paragonimus westermani

Features	Schistosoma hematobium	Fasciola hepatica	Paragonimus westermani
General feature	Male is broader than fema le	Largest and most common liver fluke.	Endemic foci were seen in Manipur
Sexes	Dioecious(Option-B), sexe s are separate	Hermaphroditic parasite	Hermaphroditic parasite
Habitat	Mesenteric venous plexus (S. mansoni and S. japonic um)vesical, and prostatic v enous plexus (S.haematobi um).	The biliary tract of man.	Cystic spaces in the lung.

Features	Schistosoma hematobium	Fasciola hepatica	Paragonimus westermani
Body	Leaf-like unsegmented bo dy with two cup-like sucke rs with delicate spines	Large leaf-shaped with a d orsoventrally flattened bo dy.	Egg-shaped, reddish, brow n, and covered with a scal e-like spine.
Eggs	Nonoperculated eggs cont aining a ciliated embryo, miracidium	Ovoid, operculated, and bi le-stained.	Oval, operculated, and gol den brown
Definitive host	Man	Sheep	Man and domestic animal s
First intermediate ho st	Freshwater snails	Freshwater snails (Lymna ea)	Snails of genera Semisu/co spira (Melania species)
Second intermediate host	-	Aquatic vegetation	Crab or crayfish
Infective form	Fork-tailed cercariae	Metacercariaeencysted on raw aquatic vegetation	Encysted metacercaria in c rab or crayfish
Clinical features	Swimmer's itchKatayama f everhematuriaportal hyper tension	Acute phase-fever and hep atomegaly.Chronic phase-biliary obstruction, obstru ctive jaundice, cholelithias is, and anemia	Peribronchial granuloma a nd cystic dilation of bronc hi.DyspneaHemoptysisPne umonitisBronchiectasisAb scessPneumothoraxExtrap ulmonary lesions in the br ain and intestine.
Diagnosis	Detection of eggs in urine or stoolbiopsyimagingdete ction of antigen and antib ody	Detection of eggs in stool and aspirated bileUSGER CPELISA	Ova in sputumX-ray and C T scan of the chestCFTIH AELISA
Treatment	Praziquantel	Oral triclabendazole	Praziquantel

Solution to Question 10:

The toxin responsible for the abnormal posturing shown in the image is tetanospasmin produced by Clostridium tetani. It acts by causing presynaptic inhibition of GABA and glycine release.

Tetanospasmin, after secretion by the bacterium is cleaved into two peptides linked by a disulphide bond. The larger peptide diffuses to terminals of neurons and migrates by the retrograde axonal transport system to the spinal cord and brainstem. It diffuses to the glycinergic interneurons and GABA secreting neurons there. The smaller peptide degrades synaptobrevin and blocks the release of the inhibitory glycine and GABA. This results in hyperreflexia, muscle spasms, and spastic paralysis. Even small amounts of the toxin can be lethal.

The incubation period of C. tetani ranges from 4-5 days up to 3 weeks. Muscle spasms usually involve the area of injury first and then the muscles of the jaw causing trismus or lockjaw. Gradually, other voluntary muscles become involved, resulting in tonic spasms. Any external stimulus can precipitate a tetanic generalized muscle spasm. The patient is conscious but may be in pain. Death usually results from respiratory paralysis.

Other options: Proteolysis of target SNARE proteins in the neurons inhibits the release of Ach at the synapse - this is the mechanism of action of Botulinum toxin.

Solution to Question 11:

The given clinical vignette of Pyrexia of Unknown Origin (PUO) / Fever of unknown origin (FUO) with a history of frequent travel is suggestive of an infectious cause of FUO. The given slide shows macrophages infested with intracellular Leishmania donovani bodies (arrows) characterized by a kinetoplast and characteristic double dot appearance (Leishman stain). The most likely diagnosis is leishmaniasis and it can be established by the rk 39 antigen test.

Pyrexia/ Fever of unknown origin - criteria:

• Fever >38.3 degrees C (101 F) on 2 occasions

• Duration > 3 weeks

• Diagnosis is not certain after history, physical examination, and routine and obligatory investigations in spite of three OPD visits or after 3 days of hospitalization.

Causes of FUO are enlisted in the table.

Leishmaniasis is a slowly progressing indigenous disease caused by a protozoan parasite of the genus Leishmania. It is transmitted by the bite of an infected female Phlebotomus sandfly which inoculates the flagellated promastigotes into the skin of the host in developing countries.

In human beings, the disease presents in four different forms with a broad range of clinical manifestations:

• Visceral leishmaniasis, or kala-azar,

• Cutaneous leishmaniasis,

• Mucocutaneous leishmaniasis, and

• Diffuse cutaneous leishmaniasis.

The life cycle of Leishmania involves two forms,

• The promastigote which develops and lives extracellularly in the sandfly vector and

• The amastigote multiplies intracellularly in the reticuloendothelial cells of the host.

It lowers immunity, causes persistent fever, anemia, liver and spleen enlargement, and if left untreated, death may occur due to opportunistic infection.

The microscopic demonstration of amastigotes from bone marrow is the gold standard for the diagnosis of Kala-azar.

For microscopic demonstration of the parasite, the materials collected are:

-Peripheral blood

-Bone marrow

-Splenic aspirate

-Enlarged lymph node.

Other diagnostic tests are:

• Rapid diagnostic test: rK39 Rapid diagnostic test and Kala-azar dipstick test strip. This is recommended by National Vector Borne Disease Control Programme (NVBDCP) in India.

• Parasitological diagnosis: Demonstration of LD bodies in buffy coat smear - confirmatory.

• Aldehyde test of Napier: Widely used for surveillance.

• Serological tests for diagnosis: Rapid dipstick test, Direct agglutination test, ELISA, Indirect Fluorescent Antibody Test

• Leishmanin (Montenegro) test: Induration of 5mm or more is considered positive, usually positive 4-6 weeks (delayed-hypersensitivity) after onset in cutaneous leishmaniasis, valuable for distinguishing immune from nonimmune subjects and endemicity.

The given images show amastigotes in the bone marrow.

Infectious cause	Non-infectious cause
Bacterial:M.tuberculosisSalmonella sppBrucella sppChlamydia psittaciLeptospira sppRickettsia sppCoxiella burnetti Mycoplasma sppAtypical mycobacteriaViral:CytomegalovirusEpstein - Barr virusArbovirusesEnterovirusesHIVFungal:Candida albicansCryptococcus neoformansHistoplasma capsulatumAspergillus sppCoccidiodes immitisParasitic:Plasmodium sppLeishmania spp Trypanosoma sppToxoplasma gondiiWuchereria bancrofti Brugia malayi	Neoplasm:LymphomaLeukemiaMyelomaRenal cancerColon cancerLiver cancerMetabolic disorderGoutPorphyriaConnective tissue disorder:Systemic lupus erythematosus (SLE)Polyarteritis nodosaGranulomatous disease:Crohn's diseaseSarcoidosisRheumatoid arthritisPolymyositis



Drugs used in the treatment of visceral leishmaniasis are sodium stibogluconate, miltefosine, and amphotericin. Liposomal amphotericin single dose 10mg/Kg IV is the first line of treatment in India.

Solution to Question 12:

The given image shows the lifecycle of Toxoplasma gondii.

Toxoplasma gondii is an obligate intracellular parasite that causes toxoplasmosis.

• The life cycle of T. gondii

• The enteric cycle occurs in definitive hosts such as cats. Exoenteric cycle occurs in intermediate hosts such as humans, mice, rats, etc.

• Both asexual reproduction (schizogony) and sexual reproduction (gametogony) occur within the mucosal epithelial cells of the small intestine of the cat. The tachyzoites and tissue cyst represent stages in asexual multiplication, while the oocyst is formed by sexual reproduction.

• All three forms, trophozoite, tissue cyst, and oocyst occur in the definitive host. Only the asexual forms, tachyzoites, and tissue cysts are present in the intermediate host.

• Asexual stage: Oocysts containing sporozoites or tissue cysts containing bradyzoites are ingested by intermediate hosts like humans, sheep, etc. Bradyzoites and sporozoites are released in the intestinal epithelium and form tachyzoites, which divide rapidly and replicates in all mammalian cells except for red blood cells. Tissue cysts are then formed which can occur in various organs but principally persist in the central nervous system. This chronic stage completes the asexual cycle and thus the parasite completes its life cycle in the intermediate host.

• The definitive host is the cat in which the sexual stage takes place. In this stage, the oocysts are formed, which after sporulation are excreted and if ingested by an intermediate host, cause pathological changes.

• Tachyzoites continue to multiply and spread locally by the lymphatic system and blood, while the bradyzoites multiply slowly inside the tissue cyst. The dormant bradyzoites inside the cyst may be reactivated in immune suppression causing renewed infection in the host.

• Transmission of human Toxoplasma infection

• This can be through oral routes from contaminated water or food, etc; it can be transmitted through blood or organs (transfusion or transplantation) and it can be transmitted through the trans-placental route.

The below image shows the life cycle of Toxoplasma gondii.



The image below shows the life cycle of Cystosisospora belli or Isospora belli.



The image below shows the life cycle of Cryptosporidium parvum.



The image below shows the life cycle of Plasmodium species.



Solution to Question 13:

HTLV - Human T cell lymphotropic virus causes microbial carcinogenesis by indirect mechanisms. Even, Hepatitis C virus, a flavivirus, does not generate a provirus and appears to induce cancer indirectly.

Oncogenic viruses:

Integration of viral genome into the host cell thereby leading to the formation of a pro-virus is commonly a mechanism of viral microbial carcinogenesis. DNA tumor viruses encode viral oncoproteins that are important for viral replication. Some portion of this may become integrated into the host cell chromosome. For RNA tumor viruses that mostly belong to the retrovirus family

are known to cause microbial carcinogenesis by indirect mechanisms.

Note:

HPV causes carcinogenesis by both integrations and by the E6, E7 protein systems.

Solution to Question 14:

The given clinical vignette is suggestive of infective cardiac valvulitis with mycotic vegetation. The histopathological slide shows fungal hyphae of Aspergillus species with non-pigmented, narrow, septate, hyphae with branching at 45o (acute angle).

Invasive aspergillosis is an opportunistic infection seen in the immunosuppressed hosts. Aspergillus has a tendency to invade blood vessels. Thus, widespread hematogenous dissemination with involvement of the heart valves and brain is common. Aspergillus form fruiting bodies and septate filaments, 5 to 10 μm thick, branching at acute angles (40 degrees). Aspergillus hyphae without the distinct fruiting body cannot be distinguished from Pseudallescheria boydii and Fusarium spp. by morphology alone.

Rhinocerebral Aspergillus infection in immunosuppressed individuals resembles that caused by Mucormycoses (e.g., Mucor spp., Rhizopus spp.). (Option-A) Mucormycetes form broad ribbon-like non-septate or sparsely septate pigmented hyphae of variable width (6 to 50 μm) with frequent right-angle branching, distinct from Aspergillus hyphae.

Option B: Disseminated fusariosis differs from disseminated aspergillosis as the former presents with nodular or necrotic painful skin lesions are extremely common with fusariosis. No such history is given in the clinical stem. The organism is difficult to differentiate from Aspergillus species in tissues; thus, identification with culture is imperative.

Option C: Rhizopus shows aseptate hyphae with sporangium and sporangiophores. It is similar in appearance to that of mucormycosis.

Solution to Question 15:

The attachment-effacement lesion of the intestinal mucosa as shown in the image is caused by EPEC (Entero-pathogenic Escherichia coli).

E. coli is a gram-negative, flagellated rod. It is a lactose fermenter and hence, grows on MacConkey medium to produce pink colonies.

Clinical infections are of four main types:

• Urinary tract infection

• Diarrhea

• Septicemia, neonatal sepsis and neonatal meningitis

• Pyogenic infections

There are mainly six types of diarrheagenic E.coli:

• Enteropathogenic E.coli (EPEC)

• Enterotoxigenic E.coli (ETEC)

• Enteroinvasive E.coli (EIEC)

• Enterohemorrhagic E.coli (EHEC)

• Enteroaggregative E.coli (EAEC)

• Diffusely adherent E.coli (DAEC)

Enteropathogenic E.coli (EPEC): It causes diarrhea mainly in infants and children. EPEC adherence factor (EAF), a plasmid-encoded protein is responsible for adherence. Another mechanism is adherence to enterocyte membrane by a locus of enterocyte effacement (LEE), which is regulated chromosomally. This promotes tight adherence of the pathogen with loss of microvilli leading to effacement.

The electron micrograph below shows EPEC adherent to effaced microvilli of the small bowel.

©MARROW

Enterotoxigenic E. coli: Causes endemic diarrhoea in developing countries and travellers diarrhoea. ETEC serotypes responsible for diarrhea are mostly 06, 08, 015, 025, 027, 0167. It attaches to intestinal mucosa with fimbriae called colonization factor antigens. The pathology is due to the heat-labile toxin (LT) produced by some and heat-stable toxin (ST) produced by others.

Enteroinvasive E.coli (EIEC): The clinical features are similar to shigellosis. It produces disease by invading the intestinal mucosa.

Enteroaggregative E. coli (EAEC): It causes acute and chronic diarrhea (>14 days in duration) in developing countries. These strains appear aggregated in a stacked brick formation. Some strains of EAEC produce ST-like toxins, others produce a plasmid-encoded enterotoxin that produces cellular damage, and still others, a hemolysin.

Enterohemorrhagic E.coli (EHEC): These strains produce two potent toxins, verocytotoxin (VT) or Shiga-like toxin (SLT). Especially in young children and the elderly, they cause diarrhea ranging from mild to fatal hemorrhagic colitis and hemorrhagic uremic syndrome (HUS). EHEC primarily targets vascular endothelial cells leading to HUS. E.coli serotype 0157: H7 is associated with EHEC diarrhea and its complications.

Diffusely adherent E.coli (DAEC): These are less established as pathogens.

Solution to Question 16:

The component responsible for endotoxic shock in a gram-negative cell wall is 1, which represents the lipid A part of lipopolysaccharide (LPS).

The numbers labelled in the above image of the cell wall indicate:

• 1- Lipid A part of LPS

• 2- O polysaccharide part of LPS

• 3- Outer membrane

• 4- Plasma membrane

Gram-negative cell wall is divided into 3 regions:

• Outer membrane consisting of LPS, proteins (porins) and phospholipids

• Peptidoglycan

• Plasma membrane consisting of phospholipids

Lipopolysaccharide (LPS) is a part of the outer membrane of the cell walls of gram-negative bacteria. It accounts for their endotoxic activity and O antigen specificity. They were formerly known as the Boivin antigen. LPS consists of:

• O polysaccharide portion determining O antigen specificity

• Core polysaccharide

• Lipid A (glycolipid). It is responsible for endotoxic activities such as pyrogenicity, lethal effect, tissue necrosis, anticomplementary activity, B cell mitogenicity, immunoadjuvant property, and antitumor activity.

The differences between exotoxins and endotoxins are summarised below.

Exotoxins	Endotoxins
Actively secreted by cells and diffuse into the surrounding	An integral part of the cell wall of gram-negative bacteria; death and in part during growth; may not need to be released to have biological activity
Produced by both gram-positive and gram-negative bacteria	Found only in gram-negative bacteria
Proteins	Lipopolysaccharide complexes
Heat labile	Heat stable
Highly antigenic, stimulate the formation of high-titer antitoxin.Antitoxin neutralizes the toxin	Weakly antigenic

Exotoxins	Endotoxins
Converted to toxoid, used to i mmunize (eg, tetanus toxoid)	Not converted to toxoids
Highly toxic	Moderately toxic
Usually bind to specific recep tors on cells	Specific receptors not found on cells
Usually do not produce fever in the host	Usually, produce fever in the host



Other options:

Option A: 2 is the O polysaccharide portion of the LPS which determines O antigen specificity.

Option B: 4 is the plasma (cytoplasmic) membrane. It is a thin semipermeable layer that lines the inner surface of the cell wall and separates it from the cytoplasm.

Option D: 3 is the outer membrane. It is the outermost layer of the gram-negative bacterial cell wall. It contains various proteins known as outer membrane proteins.

Solution to Question 17:

The image shows Gram's stain of Corynebacterium diphtheria and the staining procedure is crystal violet, Gram's iodine, acetone, and safranin.

Corynebacterium diphtheria (also called Klebs-Loeffler bacillus) is a gram-positive bacillus with clubbing at both ends. They possess characteristic polymetaphosphate granules also called metachromatic granules.

Gram's stain procedure:

• Fix the smear using heat or methanol.

• Cover with crystal violet stain for 10–30 seconds.

• Gram's iodine stain used as mordant for 10–30 seconds.

• Decolorize with 30% acetone–alcohol for 10–30 seconds, until the stain cannot flow off the slide.

• Cover with safranin stain for 10–30 seconds.

Image shows gram staining of Corynebacterium diphtheria



Other stains used for corynebacterium diphtheria:

• Albert's stain shows green bacilli with purple-blue metachromatic granules arranged in typical Chinese letter or cuneiform arrangement. The stain requires two reagents:

• Albert's I solution contains toluidine blue, malachite green.

• Albert's II contains iodine and potassium iodide.

• Loeffler's methylene blue: The granules take up the bluish-purple color.

• Immunofluorescence: This is more specific as the smear is stained with a specific antibody.

Image shows Albert's stain of C. diphtheria

The incubation period of diphtheria is usually 2–5 days (range: 1–10 days).

Respiratory diphtheria is characterized by mild fever, sore throat, difficulty swallowing, malaise, loss of appetite, hoarseness (if the larynx is involved). The hallmark is a gray-white pseudomembrane over the mucous lining of the tonsils (most commonly), pharynx, larynx, or nares. It is tightly adherent to the underlying tissue, and any attempt to dislodge the membrane may cause bleeding.

A few patients develop massive swelling of the tonsils and present with bull-neck diphtheria, due to edema of the submandibular and paratracheal region. Cutaneous diphtheria may present as a scaling rash or as ulcers with clearly demarcated edges and membrane.

The images below shows pseudomembrane, bull-neck, and cutaneous diphtheria respectively.

Treatment of diphtheria includes administration of diphtheria anti-toxin. Recommended antibiotics include penicillin G and erythromycin. Prevention is with the diphtheria toxoid vaccine. It is coadministered with tetanus and acellular pertussis (DPT) and should be given according to the National Immunization Schedule.

Solution to Question 18:

The given image shows the life cycle of the herpes simplex virus (HSV).

The herpes virus capsid is icosahedral and encloses a core containing a linear double-stranded DNA genome. The nucleocapsid is surrounded by a lipid envelope derived from the modified host cell nuclear membrane. The envelope carries glycoprotein spikes.

Steps of HSV replication:

The virus enters the cell by fusing with the plasma membrane, and viral DNA is released from the capsid at the nuclear pore. This is followed by circularization of the genome and transcription of immediate-early genes.

α-Proteins are products of immediate-early genes. These proteins stimulate the transcription of early genes. β-proteins are products of early genes. These proteins function in DNA replication, to yield concatemeric DNA. Transcription of late genes gives rise to γ-proteins. These primarily consist of viral structural proteins and participate in virion assembly.

Viral DNA is cleaved from concatemers and then packaged into capsids. Enveloped viral particles collect in the endoplasmic reticulum and are then transported by a vesicular movement to the cell surface.

Image shows replication cycle of herpes simplex virus.



Other options:

Option A) The human immunodeficiency virus (HIV) virus is spherical, enveloped RNA virus. The nucleocapsid has an icosahedral shell and a cone-shaped core, enclosing the ribonucleoproteins. The genome icomprises two identical single-stranded, RNA copies, with the reverse transcriptase enzyme, which is a characteristic feature of retroviruses.

The images below show the structure of the HIV virus and its life cycle.



Option C: The hepatitis B virus is a DNA virus with an outer envelope and an inner nucleocapsid. The viral genome consists of two linear strands of DNA in a circular configuration. One of the strands (the plus strand) is incomplete resulting in partially double stranded and partially single-stranded DNA. The images below show the structure and life cycle of the hepatitis B virus.





Option D: Coronavirus are enveloped RNA viruses with a genome consisting of positive sense unsegmented single-stranded RNA. They have petal-shaped surface projections. Coronavirus nucleocapsids develop in the cytoplasm and mature by budding into cytoplasmic vesicles. The images below show the structure and life cycle of coronaviruses.





Solution to Question 19:

Injecting Aedes thorax with blood of a suspected dengue patient is the xenodiagnostic method among the given options.

Xenodiagnosis (Xenos- foreign) is an indirect method of identifying the infectious disease agent or pathogen. In xenodiagnosis, the vector of a pathogenic agent is infected and the pathogen is allowed to multiply in the vector. Vectors are insects such as mosquitoes, ticks, mites, flies, fleas, and lice that transmit infections and they act as the 'foreign host.' Later, the vector is examined for the presence of the pathogen.

For example, diagnosis of Chagas disease: Laboratory bred Reduviid bugs (vector/ foreign host) are allowed to feed on the suspected patients (original host). After 4-5 weeks, the feces of the bugs are microscopically examined for amastigote forms of Trypanosoma cruzi (pathogen).

Another method of xenodiagnosis is the intrathoracic inoculation of the mosquitoes (vector) with viruses (pathogen). The virus multiplies in different organs of the mosquito and can be easily demonstrated. Aedes aegypti, Aedes albopictus, and Toxorhynchites mosquitoes are used for the identification of the dengue virus.

In Trichinella spiralis infection, muscle biopsy bits from suspected patients are fed to laboratory rats (vector for T. spiralis). The larvae of T. spiralis can be demonstrated in the muscles of the infected lab rats.

Advantage of xenodiagnosis: It is useful in cases where a direct microscopic diagnosis is difficult due to the low count of pathogens in the blood of the host/patient.

Other options:

Option A: Ligated rabbit ileal loop is used as a method of detection of Enterotoxigenic E.coli enterotoxins. It is not a method of xenodiagnosis as the vector of the pathogen is not being infected.

Option C: Injecting a hamster with splenic biopsy for diagnosis of leishmaniasis is a method of animal inoculation to help in confirmation of the diagnosis.

Option D: Intradermal test on guinea pigs for toxigenicity of Corynebacterium diphtheria is a method of virulence testing.

Solution to Question 20:

The principle of the Paul-Bunnell test is based on heterophile antibodies. Heterophile antibodies have the ability to agglutinate shared antigens.

Paul-Bunnell test is the standard diagnostic procedure for infectious mononucleosis. In infectious mononucleosis, heterophile antibodies agglutinate sheep erythrocytes. However, such antibodies may also occur after injections of sera and sometimes even in normal individuals. Infectious mononucleosis antibodies may be differentiated by absorption tests. An agglutination titer of 100 or above is suggestive of infectious mononucleosis.

The Weil-Felix reaction is also a heterophile agglutination test. It is based on the sharing of a common antigen between typhus rickettsiae and some strains of proteus bacilli. It is used for the serodiagnosis of typhus fevers. Other examples based on the same principle are Streptococcus MG agglutination test for the diagnosis of primary atypical pneumonia and the cold agglutination test for diagnosis of mycoplasmal (primary atypical) pneumonia.

Other options:

Option A: Complement fixation test is based on the ability of antigen-antibody complexes to fix complement. This is a very sensitive test capable of detecting various types of antigens and antibodies.

Option B: Superantigens are certain protein molecules, that activate very large numbers of T cells irrespective of their antigenic specificity such as staphylococcal enterotoxins. Superantigens bind outside the antibody binding groove directly to the lateral aspect of the TCR beta chain. Microbial superantigens are medium-sized proteins characterized by high resistance to proteases and to denaturation by CD4+ T cells. They cause the release of cytokines (IL-2) which results in massive proliferation of T lymphocytes. This ultimately leads to further release of a variety of cytokines which can have profound effects on the immune system.

Option D: Haptens are substances that can react specifically with antibodies but are incapable of inducing antibody formation alone. Haptens become immunogenic (capable of inducing antibodies) by combining with a larger molecule carrier. They may be complex or simple. While complex haptens can precipitate with specific antibodies, simple haptens are non-precipitating.

Solution to Question 21:

The correct match is 1-c, 2-d, 3-a, 4-b

Japanese encephalitis virus is a virus from the family Flaviviridae. It is the leading cause of vaccine-preventable encephalitis in Asia and the western Pacific.

Transmission of Japanese encephalitis virus: Natural infection has been demonstrated in Ardeid birds (herons and egrets). Bird-to-bird transmission takes place through the vectors Culex tritaeniorhynchus and Culex vishnui (common in India). Ardeid birds act as reservoir hosts, while pigs and cattle act as amplifier hosts. Humans are accidental dead-end hosts.

The image below shows the life cycle of Japanese encephalitis virus.

1. Accidental hos t	c. Human
2. Amplifier	d. Pigs
3. Reservoir	a. Ardeid bird s
4. Vector	b. Culex



LIFE CYCLE OF JAPANESE ENCEPHALITIS

Most people infected with JE do not have symptoms or have only mild symptoms. Less than 1% of people infected with Japanese encephalitis (JE) virus develop clinical illness. The incubation period is typically 5-15 days in persons who develop symptoms. Initial symptoms often include fever, headache, and vomiting. Mental status changes, neurologic symptoms, weakness, and movement disorders might develop over the next few days. Seizures are common, especially among children.

No specific treatments have been found to benefit patients with JE, but hospitalization for supportive care and close observation is generally required. Treatment is symptomatic. Rest, fluids, and the use of pain relievers and medication to reduce fever may relieve some symptoms.

Prevention: In India, inactivated vero cell culture-derived SA-14-14-2 is used. Japanese encephalitis vaccine has been introduced in selected endemic districts. 2 doses of 0.5 ml are given in the left upper arm subcutaneously.

If the first dose is delayed beyond 12 months, a minimum of 3 months gap should be there between the two JE doses.

Vaccine	Due age	Maximum age
Japanese encephalitis - 1	9-12 months	15 years
Japanese encephalitis - 2	16-24 months	15 years

Solution to Question 22:

Cladosporium is the dematiaceous fungus in the given options.

Dematiaceous fungi are fungi whose cell walls contain melanin, which imparts a brown to black color. Phialophora verrucosa, Fonsecaea pedrosoi, Fonsecaea compacta, Rhinocladiella aquaspersa, and Cladophialophora (Cladosporium) carrionii are examples of dematiaceous fungi.

These fungi cause chromoblastomycosis, a chronic subcutaneous infection caused due to traumatic inoculation of fungal agents. The lesion is verrucous and wart-like with extension along the draining lymphatics. Cauliflower-like nodules with crusting abscesses eventually cover the area.

On histological examination, the lesions contain the fungus seen as round or irregular, dark brown, yeast-like bodies with septae, called sclerotic bodies or medlar bodies. Diagnosis can be established by demonstration of these sclerotic bodies in KOH mounts or in tissue sections, and by culture on Sabouraud agar.

Chemotherapy with amphotericin B, thiabendazole, 5-fluorocytosine, itraconazole, and recently voriconazole are useful for the treatment of larger lesions. Small lesions can be excised surgically.

The image below shows sclerotic bodies seen in chromoblastomycosis.



Other options:

Option A: Mucor species causes mucormycosis. It is called black fungus because it causes serious angioinvasion leading to tissue necrosis and gangrene. It is not a dematiaceous fungus. They appear as non pigmented (hyaline), broad non-septate hyphae on microscopy.

Option B: Aspergillus species usually appear as non pigmented (hyaline), septate hyphae with acute angle branching on microscopy. The conidia of A.niger are black in color.

Option D: Blastomyces dermatitidis appear as broad-based budding yeasts.

Solution to Question 23:

The strict anaerobe in the given options is Prevotella.

Prevotella species are gram-negative rods or coccobacilli. The most commonly isolated Prevotella species are P melaninogenica, Prevotella bivia, and Prevotella disiens. P melaninogenica is associated with upper respiratory tract infections. P bivia and P disiens occur in the female genital tract. Prevotella species are also found in brain and lung abscesses, empyema, pelvic inflammatory disease and tubo-ovarian abscesses.

Other options

Option A: Burkholderia cepacia belongs to the pseudomonad group and is gram-negative, aerobic rods.

Option B: Yersinia enterocolitica are gram-negative facultative anaerobes.

Option C: Stenotrophomonas maltophilia also belong to the pseudomonad group and are gram-negative, aerobic rods.

Solution to Question 24:

The correct match is 1-C, 2-D, 3-B, 4-A.

The Hymenolepis nana egg is roughly spherical or ovoid. It has a thin colorless outer membrane and inner embryophore that encloses the hexacanth oncosphere. They are non bile-stained. The eggs float in saturated salt solution.

The image below shows an egg of H. nana.



Ascaris lumbricoides (Roundworm) are round or oval in shape. They are always bile-stained which gives them golden-brown color. They have a thick smooth translucent shell with:

• an outer coarsely mammillated albuminous coat

• a thick transparent middle layer, and

• an inner lipoidal vitelline membrane.

The eggs float in a saturated salt solution. The image below shows the egg of A. lumbricoides.



The unfertilized eggs of Ascaris are elliptical, narrower, and contain a small atrophied ovum. It has a mass of disorganized refractile granules. The shell is thinner and the egg does not float in a saturated salt solution.

The egg of Ancylostoma duodenale or hookworm is oval or elliptical. They are colorless, not bile-stained. The eggs are surrounded by a thin transparent hyaline shell membrane. It contains a segmented ovum, usually with 4-8 blastomeres. There is a clear space between the ovum and the

shell. It floats in saturated salt solution. The image below shows the egg of A. duodenale.



The eggs of Taenia solium and Taenia saginata are indistinguishable. The egg is spherical. The inner embryophore is radially striated and is yellow-brown due to bile staining. A fully-developed embryo (oncosphere) with three pairs of hooklets (hexacanth embryo) is present in the center.

The eggs of T. saginata are infective only to cattle and not to humans, whereas the eggs of T. solium are infective to pigs and humans. The eggs do not float in a saturated salt solution. The image below shows the egg of Taenia.



Solution to Question 25:

The causative organism of the lesion given in the image is Sporothrix schenckii (S. schenckii)

S. schenckii causes sporotrichosis, chronic granulomatous disease by traumatic inoculation into the skin. This is usually followed by spread along the draining lymphatics and lymph nodes.

It is a thermally dimorphic fungus. It is in the yeast phase in tissues and in cultures at 37°C, and in the mycelia phase in nature and cultures at 25°C. When it is cultured, the colonies are blackish brown, fuzzy. The organism produces branching, septate hyphae, and microconidia clustered at the ends of conidiophores. It exists as cigar-shaped yeast cells. Stains used for identifying yeast forms:

• Gomori methenamine silver - cell wall appears black

• Periodic acid-Schiff stain- cell wall appears red

• Fluorescent antibody staining

The image given below shows, (A)- (i) culture at 37°C (ii) culture at 25°C. (B) fungi with clustered microconidia with conidiophore. (C) magnified image showing branching and septate hyphae.



Clinical features: The initial infection develops as a nodule, which may break down to form a necrotic or ulcerative lesion. Untreated, the disease progresses and new nodules develop along the lymphatics leading to a chain of nodules. Less commonly it may also cause systemic infection following inhalation, which causes local pulmonary disease or may lead to widely disseminated lesions in joints, meninges, and skin.

Treatment: Itraconazole is the drug of choice for lympho-cutaneous sporotrichosis, also terbinafine can be used. A saturated solution of potassium iodide (SSKI) is a second-line drug.

Other options:

Option A: Mycobacterium scrofulaceum is a scotochromogen. It causes chronic cervical lymphadenitis in children.

Option C: Eumycetoma is a mycetoma caused by fungi like Madurella mycetomatis. A mycetoma is a chronic, post-traumatic infection that begins as a small subcutaneous swelling that enlarges and burrows into the deeper tissues, forming multiple discharging sinuses. It is shown below.

Option D: Phialophora verrucosa causes chromoblastomycosis. The lesions appear as warty, cutaneous nodules usually confined to the subcutaneous tissue of the lower limbs. Histologically, the lesions contain the fungus, which is seen as round or irregular, dark brown, yeast-like bodies with septae, known as sclerotic bodies. A skin lesion of chromoblastomycosis is seen below, followed by a histopathological image showing sclerotic bodies.



Microbiology NEET 2021

Question 1:

A patient from Uttar Pradesh presented with fever, pallor, and hepatosplenomegaly. Peripheral smear examination showed pancytopenia. Buffy coat examination showed macrophages laden with organisms with a kinetoplast. What is the vector for the likely disease?

a) Sandfly

b) Tse-Tse fly

c) Triatomine bug

d) Female anopheles mosquito

Question 2:

A patient who was on steroids for the treatment of chronic urticaria presents to the emergency department with acute worsening of respiratory symptoms. He says that he has been having nocturnal cough for last several weeks. A hemogram revealed elevated eosinophils. Bronchoalveolar lavage (BAL) specimen following bronchoscopy showed the presence of certain larvae as shown in the image below. Identify the causative organism.

a) Ancylostoma caninum

b) Capillaria philippinensis

c) Strongyloides stercoralis

d) Enterobius vermicularis

Question 3:

A patient admitted with 50% burns develops an infection at the burn site. The swab was cultured, the isolate is a strict aerobe and the test that is shown in the image is positive. What is the likely etiology of the burn infection?



a) Klebsiella

b) Escherichia coli

c) Pseudomonas aeruginosa

d) Salmonella

Question 4:

One week after having unprotected sexual intercourse, a 25-year-old male develops urethral discharge as shown in the image below. What is the most likely causative organism for this condition?

a) Neisseria gonorrheae

b) Ureaplasma urealyticum

c) Herpes simplex virus

d) Trichomonas vaginalis

Question 5:

An unimmunized 2-year old child presented with coryza, conjunctivitis, and bluish-white spots in his buccal mucosa near the lower molar teeth. A day later, he developed a maculopapular rash on the face and neck. What is the nature of the causative virus?

a) Enveloped ss RNA

b) Naked ss RNA

c) Naked ds RNA

d) Enveloped ds RNA

Question 6:

A group of 20 friends kept partying till late at night. Two of them ate pastries from outside. They present with nausea and vomiting in the early morning. What is the most likely causative agent implicated?

a) Enterotoxin of Staphylococcus

b) Emetic toxin of Bacillus cereus

c) Verocytotoxin from E. coli

d) Shiga toxin from Shigella

Question 7:

A 3-year old child presents with fever, loose stools, and blood in the stool. What is the enrichment medium used to isolate the organism from the stool sample?

a) Robertson cooked meat broth

b) Alkaline peptone water

c) Selenite F broth

d) Brain heart infusion broth

Question 8:

A truck driver came with complaints of a painless indurated genital ulcer after unsafe intercourse. Motility of the most likely causative organism can be detected by which of the following microscopes?

a) Dark field microscope

b) Electron microscope

c) Light microscope

d) Fluorescent microscope

Question 9:

Name the parasite whose microfilaria have a sheath with no nuclei at the tail end.

a) Wuchereria bancrofti

b) Brugia malayi

c) Brugia timori

d) Mansonella perstans

Question 10:

In a case of tinea capitis, white powdery colonies were isolated from the hair specimen. The hair perforation test was positive on the isolate. Which of the following is the likely etiology?

a) Epidermophyton floccosum

b) Microsporum audonii

c) Trichophyton mentagrophytes

d) Microsporum gypseum

Question 11:

A patient was suspected to have acute brucellosis. A serum sample was sent for a standard agglutination test. It came out to be negative initially but after serial dilutions of the serum sample, the test turned out to be positive. Which of the following is responsible for the initial negative test?

a) Incomplete antibodies

b) Complement inactivation

c) Prozone phenomenon

d) Post-zone phenomenon

Answer Key

Question No.	Correct Option
1	a
2	c
3	c
4	a
5	a
6	a
7	c
8	a
9	a
10	c
11	c

Detailed Explanations

Solution to Question 1:

From the given clinical features, the most likely disease is visceral leishmaniasis or Kala azar caused by Leishmania donovani, and the vector is sandfly. A patient from Uttar Pradesh presenting with fever, pallor, and hepatosplenomegaly with a smear examination showing pancytopenia and macrophages containing organisms with kinetoplasts point towards the diagnosis.

Leishmaniasis is transmitted by the bite of an infected female Phlebotomus sandfly. In India, it is endemic to Bihar, West Bengal, Jharkhand, and Uttar Pradesh. Patients present with high fever, hepatosplenomegaly, weight loss, and emaciation. The skin becomes dry, rough, and darkly pigmented. There is anemia, leukopenia, and thrombocytopenia.

The parasite exists in 2 forms:

• Amstigote form (also called LD bodies)- It is seen within macrophages and contains a kinetoplast.

• Promastigote form- It is the infective form in the sandfly. The kinetoplast is seen at the anterior end.

Kinetoplast are non-nuclear DNA present in addition to nucleus. They consist of blepharoplast and parabasal body. Flagellum originates near the kinetoplast.

Diagnosis of kala-azar is mainly by demonstration of amastigotes within macrophages in tissue aspirates such as blood smears, bone marrow, splenic, or lymph node aspirate. It is the gold standard for diagnosis.

The image below shows amastigotes located intracellularly within macrophages.



Treatment is with a pentavalent antimonial compound. In Bihar, there is resistance to antimony so amphotericin-B-deoxycholate or miltefosine is preferred.

In human beings, different species of Leishmania can cause the following diseases:

• Visceral leishmaniasis, or kala-azar,

• Cutaneous leishmaniasis,

• Mucocutaneous leishmaniasis, and

• Diffuse cutaneous leishmaniasis.

The life cycle of Lesihmania is shown below.



Other options:

Option B: Tse-Tse fly (Glossina species) is involved in the transmission of sleeping sickness or African trypanosomiasis. It is caused by Trypanosoma brucei. The patient can have hepatosplenomegaly and lymphadenopathy in the early stages and later on CNS involvement is seen. Blood routine examination would reveal anemia, leucocytosis, and thrombocytopenia.

Option C: Triatomine bug (Reduviid bug/ kissing bug) transmits Trypanosoma cruzi infection leading to Chaga's disease or American trypansomiasis. The acute phase of the disease is characterized by a typical subcutaneous lesion at the site of inoculation called chagoma. In the chronic phase, multiorgan involvement occurs leading to cardiac myopathy, megaesophagus, and megacolon.

Option-D: Female anopheles mosquito leads to the transmission of malaria caused by Plasmodium species. Malaria can present with fever and hepatosplenomegaly but the morphological forms commonly include the ring forms or the gametocytes.

Solution to Question 2:

The presence of eosinophilia in an immunosuppressed patient with respiratory symptoms and BAL revealing rhabditiform larvae is suggestive of disseminated strongyloidiasis caused by Strongyloides stercoralis.

Strongyloidiasis:

Strongyloides are ovoviviparous. The transmission of infection occurs through penetration of the skin by filariform larvae in the soil. Then, they migrate to the small intestine. The larvae moult twice and become adult female worms in the small intestine. The eggs laid in the mucosa of the intestine hatch immediately releasing the rhabditiform larva (L1 stage). These rhabditiform larvae in the bowel can develop directly into filariform larvae that penetrate the colonic wall or perianal

skin and enter the circulation to repeat the migration. This establishes ongoing internal reinfection.

The life cycle of strongyloides is shown below.



It is generally a benign and asymptomatic condition. In some cases, there may be cutaneous, pulmonary, and intestinal manifestations. Pulmonary manifestations occur when the larva escapes from the pulmonary capillaries into the alveoli leading to bronchopneumonia, chronic bronchitis, and asthmatic symptoms.

The term larva currens (meaning ‘racing larvae’) has been applied to the rapidly progressing linear or serpiginous urticarial tracks caused by migrating filariform larvae of Strongyloides stercoralis. These often follow autoinfection and start perianally. Disseminated Strongyloidiasis is seen in immunodeficient individuals.

Larva currens is shown in the image below.

Visceral larva migrans is caused by the migration of larvae of nonhuman species of nematodes that infect via the oral route.

Etiological agents causing visceral larva migrans:

• Human nematode: Ascaris lumbricoides, Strongyloides stercoralis.

• Zoophilic nematode: Toxocara canis, toxocara cati, Angiostrongylus cantoniensis, Angiostrongylus costaricensis, Anisakis.

• Non human nematode: Filarial species, Brugia pahangi, Brugia pate

Laboratory diagnosis of strongyloidiasis:

• Eosinophilia and raised IgE levels.

• Microscopic examination of the direct wet mount of the stool demonstrating rhabditiform larvae forms the definitive diagnosis

• Stool culture and serological tests are also done

Treatment: Ivermectin is the preferred agent in the management of strongyloidiasis.

Other options:

Option A: Ancylostoma canium presents with creeping eruption. Cutaneous larva migrans or creeping eruption is caused by nematode larvae that infect by skin penetration.

Option B: Capillaria phillipinesis can also cause auto-infection. It commonly presents with severe diarrhea.

Option D: Enterobius vermicularis or pinworm infection presents as itching (pruritus) and restless sleep (due to nocturnal perianal or perineal pruritus)

Solution to Question 3:

Pseudomonas aeruginosa is the most common cause of infection in the case of a burn. The given image shows a positive oxidase test. Psuedomonas aeruginosa is a strict aerobe and shows a positive oxidase reaction.

Pseudomonas aeruginosa is a gram-negative, catalase-positive, oxidase-positive, non-lactose fermenting bacteria that shows swarming motility. They produce a characteristic blue-green pigment called pyocyanin and a greenish-yellow pigment called fluorescein/pyoverdin. Other pigments produced by these organisms include pyorubin (Red), pyomelanin (Brown).

Diseases caused by Pseudomonas aeruginosa include-

• Malignant otitis externa in diabetics

• Pneumonia and sepsis in cystic fibrosis

• Wound infection and sepsis in burns

• Hot tub folliculitis/Pseudomonas aeruginosa folliculitis/ Jacuzzi syndrome is a type of folliculitis, occurring after hot tub exposure

• Shanghai fever is a febrile illness that manifests as severe enteric disease, sepsis with invasive disease.

• Ecthyma gangrenosum

Pathogenesis - It is an obligate aerobe that exerts its pathogenicity by the production of diphtheria-like exotoxin A and a slime layer. Breach in primary body defenses as in burns predisposes to Pseudomonas infection.

Clinical features - It is characterized by blue-green pus. It is notorious for being resistant to antimicrobials and antiseptics and hence is a frequent contaminant in hospital samples. Repeated isolation and associated clinical features should prompt the diagnosis of Pseudomonas.

Diagnosis - It grows well on most media and pigment production makes its identification easy. Species identification is made possible by biochemical tests - arginine hydrolysis and oxidase reaction.

Pseudomonas aeruginosa growth in a nutrient agar showing green coloured colonies are shown in the image below.



Oxidase test can be used to distinguish organisms by detecting the presence or absence of a respiratory enzyme, cytochrome oxidase. Cytochrome oxidase catalyzes the oxidation of reduced cytochrome by oxygen. Enterobacteriaceae (Klebsiella, E. coli, Salmonella) are oxidase negative.

Oxidase positive organisms:

• Pseudomonas

• Vibrio

• Helicobacter

• Campylobacter

• Pasteurella

• Legionella

• Neisseria

• Brucella

• Moraxella

• Burkholderia

Solution to Question 4:

The given image shows purulent urethral discharge that developed one week after unprotected sexual intercourse is suggestive of acute urethritis. The most likely causative organism for acute urethritis in males is Neisseria gonorrheae.

Purulent urethral discharge can be seen in gonorrhea caused by Neisseria gonorrhea or lymphogranuloma venerum caused by Chlamydia trachomatis.

Gonorrhea:

Gonorrhea is a disease acquired by sexual contact. The first step in infection is the infection of gonococci to the urethra or other mucosal surfaces. Pilli is involved in this adhesion. The cocci penetrate the intracellular spaces and reach subepithelial tissue by the third day of infection. The incubation period is 2-8 days. The major symptoms include urethral discharge and dysuria, usually without urinary frequency or urgency. The specific symptoms seen in males and females are as follows:

In males :

• The disease starts as acute urethritis. This infection may extend along the urethra to involve the prostate, seminal vesicles, and epididymis.

• Chronic urethritis may lead to stricture formation.

• The infection may spread to the periurethral tissues causing abscesses and multiple discharging sinuses (watercan perineum).

In females :

• The initial infection involves the cervix uteri and urethra which may extend to Bartholin glands, endometrium, and fallopian tubes.

• The vaginal mucosa is not usually affected in adults because the stratified squamous epithelium is resistant to infection by the cocci, and also because of the acidic pH of vaginal secretions.

• Pelvic inflammatory disease and salpingitis may lead to sterility.

• Rarely peritonitis may develop with perihepatic inflammation (Fitz Hugh Curtis syndrome)

Lab diagnosis:

The discharge is collected and subjected to Gram's stain which reveals gram■negative intracellular diplococci within phagocytes. Stuart's transport medium is used. Cultures can be obtained from chocolate agar or Mueller-Hinton agar. Specifically, Thayer-Martin agar is used to isolate Neisseria gonorrhoeae.

Treatment: Ceftriaxone (250mg, IM) single-dose + azithromycin 1 g PO, single-dose, or doxycycline 100 mg PO BD for 7 days

The image below shows the Gram-stain of a urethral exudate showing typical intracellular gram-negative diplococci and pleomorphic extracellular gram-negative organisms



Non-gonococcal urethritis is caused by:

• Chlamydia trachomatis

• Ureaplasma urealyticum (Option-B)

• Mycoplasma genitalium

• Trichomonas vaginalis (Option-D)

• Herpes simplex virus (rare)

• Adenovirus

• Haemophillus vaginalis

• Mycoplasma genitalium

Other options:

Option B: Ureaplasma urealyticum is the second most common cause of non-gonococcal urethritis next to chlamydia. That is why the initial etiological agent under suspicion always remains gonorrhea.

Option C: Herpes Simplex will be associated with vesicular lesions.

Option D: Trichomonas vaginalis generally presents with a frothy discharge.

Solution to Question 5:

Fever, coryza, conjunctivitis, maculopapular rash and Koplik spots are suggestive of measles. Measles virus is an enveloped, single-stranded RNA (ssRNA) virus.

Koplik spots are small, bluish-white ulcerations on the inner aspect of the cheek at the level of premolars. These are pathognomic for measles. The measles virus, a paramyxovirus, has an incubation period of 7-18 days. After this, the prodrome phase begins with mild fever, conjunctivitis, upper respiratory tract infection, and pathognomic Koplik spots. This is followed

by the eruptive stage, characterised by the appearance of maculopapular rash on the 4th day of fever. The rash begins on the face and spreads to the torso and extremities. Lymphadenopathy may be present.

These are multinucleated giant cells formed by the fusion of infected cells. The images below show the characteristic rashes and the Koplik spots seen in measles.



©MARROW

The most common complication of measles is acute otitis media. Sinusitis and mastoiditis are also seen. The most common cause of death in measles is pneumonia, which may be giant cell pneumonia. It is caused by the virus directly or can be due to superadded bacterial infection. Black measles or hemorrhagic measles is a fatal complication characterized by hemorrhagic skin eruptions. It is rarely seen nowadays.

Subacute sclerosing panencephalitis is a rare but fatal complication characterized by cognitive decline, spasticity of limbs, and seizures. It is insidious in onset and clinical manifestations usually begin 7-10 years after primary measles infection. It is caused due to persistent infection with a genetically altered variant of the measles virus within the central nervous system. Other

complications include croup, tracheitis, bronchiolitis, diarrhea and vomiting, febrile seizures, and encephalitis.

The diagnosis of measles infection is mainly clinical but may be confirmed by serological demonstration of IgM antibodies. IgM antibodies appear 1-2 days after the rash and are detectable for 1 month. The management is mainly supportive. There is no role for antivirals. Low serum retinol levels in children with measles are associated with higher morbidity and mortality and hence vitamin A therapy is indicated for all patients with measles.

Live attenuated vaccine for measles is safe and effective with no evidence of person-to-person transfer of vaccine strains. It is a freeze-dried product requiring reconstitution. 0.5ml is administered subcutaneously in the right upper arm in 2 doses- at the 9th month and between the 16th - 24th month.

Solution to Question 6:

The most likely causative agent in the given scenario is enterotoxin of Staphylococcus aureus. It causes food poisoning within 30 minutes to 6 hours after consumption of the contaminated food.

The common sources of infection are the foods that are not cooked after handling. These include sliced meats, puddings, pastries, and sandwiches. Patients present with nausea, vomiting, stomach cramps, and diarrhea.

Food poisoning is caused by enterotoxin produced by S. aureus. It is a preformed toxin that is relatively heat stable while most of the other exotoxins are heat-labile. It stimulates the vagus nerve and vomiting center of the brain and also stimulates intestinal peristaltic activity. Eight antigenic types of enterotoxin are currently known of which Type A toxin is responsible for most cases.

Other options:

Option B: Food poisoning after consumption of fried rice or chinese street food is strongly suggestive of Bacillus cereus (Serotypes 1, 3, 5). Two mechanisms of action have been described for the enterotoxin of B.cereus, one involving stimulation of the cyclic adenosine monophosphate (cAMP) system and the other independent of it.

Food poisoning after eating food contaminated with Bacillus cereus is of two types, namely diarrheal type and emetic type, caused by an enterotoxin and an emetic toxin respectively.

Bacillus cereus	Diarrheal type	Emetic type
Incubation period	8-16 hours	1-5 hours
Toxin	Secreted in intestine	Preformed
Heat	Heat labile	Heat stable
Food items	Meat	Rice
Clinical feature	Diarrhea, abdominal cramps	Vomiting, abdominal cramps

Option C: Verocytotoxin from E.coli is also known as Shiga-like toxin and it causes bloody diarrhea. It is produced by enterohemorrhagic E.Coli. It can cause illnesses ranging from mild diarrhea to fatal hemorrhagic colitis and hemorrhagic uremic syndrome.

Option D: Shiga toxin acts by inactivating 60s ribosomal subunit thereby inhibiting protein synthesis. Shigella dysenteriae causes acute onset of symptoms(fever, abdominal cramps) with dysentery and tenesmus. It also gives a positive Sereny test and is non-lactose fermenting.

Solution to Question 7:

The given clinical vignette of dysentery (bloody stools) and abdominal pain in a child is suggestive of bacillary dysentery most likely caused due to Shigella. Selenite F broth is used as an enrichment media for dysentery-causing bacteria like Shigella/ Salmonella.

Enrichment media:

It is used to suppress commensal bacteria while allowing the pathogen to remain viable and grow. Common examples include:

• Tetrathionate broth – It inhibits coliforms and allows typhoid and paratyphoid bacilli to grow.

• Selenite F broth – used for dysentery causing bacteria.

• Alkaline peptone water (Option-B) - used for vibrio cholera causing watery diarrhea.

Other options:

Robertson's cooked meat broth (Option-A): It is used for the culture of anaerobic organisms.

Brain-heart infusion broth (Option-D): It is an enriched media. It is used for the cultivation of fastidious organisms.

Dysentery/Shigellosis is a type of food poisoning caused by infection with the Shigella species. It is spread through faeco-oral transmission. They are catalse-negative, non-lactose-fermenting, gram-negative rods, and are non-motile. It acts by inactivating the 60s ribosomal subunit thereby inhibiting protein synthesis. It causes acute onset of symptoms(fever, abdominal cramps) with dysentery and tenesmus. It also gives a positive Sereny test.

Solution to Question 8:

The clinical vignette of a painless, indurated genital ulcer in a truck driver is most likely suggestive of syphilitic chancre / hard chancre caused by Treponema pallidum. The motility of the organism can be demonstrated by dark ground/ field microscopy.

Dark ground microscopy can be used to diagnose syphilis when an active chancre or condyloma lata is present to visualize Treponema pallidum. Dark-field microscopy images have a dark background with a refractile organism. The dark ground microscope has an optical system that enhances the contrast of unstained bodies. It is generally less sensitive.

Various types of Spirochetes and their appearances under a dark ground microscope:

• Treponemes generally are short, slender with regular and sharp spirals.

• Leptospira have a large number of closely wound spirals and characteristic hooked ends.

• Borrelia is large with irregular and wide open coils.

The image given below shows Treponema pallidum demonstrated by dark ground microscopy and the various appearances of other spirochetes respectively.





Other options:

Electron microscopy (Option-B) is not commonly used for diagnostics in laboratories. It generally helps in appreciating viral structures.

Light microscopes (Option-C) are commonly used in the labs to visualize the stained smears of various bacteria, fungi in 100x magnification. They are also used during the examination of stool specimens for identifying ova, eggs, or larvae.

Fluorescent microscope (Option-D) uses fluorescent dyes like auramine-rhodamine to appreciate Mycobacterial species and malarial parasites.

Syphilis:

Syphilis is caused by Treponema pallidum, a spirochete bacillus.

It shows corkscrew motility which can be demonstrated by darkfield microscopy. Smear specimens are stained using Fontana stain and pathological tissues are stained using Levaditi stain.

Serologic tests for syphilis recommended by the Centers for Disease Control and Prevention (CDC):

• Screening or diagnosis (RPR or VDRL),

• Quantitative measurement of antibody to assess clinical syphilis activity or to monitor response to therapy (RPR or VDRL)

• Confirmation of a syphilis diagnosis in a patient with a reactive RPR or VDRL test (FTA-ABS, TPHA, EIA/CIA)

Treatment of syphilis is with 2.4 million units of benzathine penicillin IM.

Note:

VDRL- Venereal disease Research Laboratory test

RPR- Rapid Plasma Reagin test

TA-ABS- Fluorescent Treponemal Antibody Absorption test

TPI-Treponema pallidum immobilization test

TPPA-Treponema pallidum particle agglutination assay

TPHA-Treponema pallidum haemagglutination test

EIA-Enzyme-linked ImmunoAssay

Solution to Question 9:

A sheathed microfilaria with a pointed tail free of nuclei is suggestive of Wuchereria bancrofti.

Wuchereria bancrofti causes lymphatic filariasis and most cases are seen in India.

Clinical features include massive eosinophilia, hepatosplenomegaly, pulmonary manifestations like dry nocturnal cough, asthmatic wheezing, and dyspnea. It can also cause arthritis, glomerulonephritis, thrombophlebitis, etc. The classical features seen in lymphatic filariasis are usually absent.

Elephantiasis refers to the painful swelling of the legs and genital organs seen in late-stage disease.

Image showing elephantiasis-

Investigations:

Demonstration of the microfilaria in peripheral blood, chylous urine, hydrocele fluid, etc is done. The specimen of choice is peripheral blood. Night blood samples between 10 pm and 4 am are preferred. This is because the microfilaria shows nocturnal periodicity.

DEC provocation test: In this test, a small dose of DEC is given 20-50 minutes before sample collection so that the microfilaria appears in the blood even during the daytime.

The image below shows the life cycle of Wuchereria bancrofti.



Solution to Question 10:

A positive hair perforation test is caused by Trichophyton mentagrophytes.

Hair perforation test:

It is done to differentiate Trichophyton rubrum from T.mentagrophytes. Few strands of human hair are placed in a petri dish with 20 ml of distilled water and autoclaved. A few drops of sterile 10% yeast extract and a few fragments of the test fungus are added to the hair strands. The entire dish is then incubated at 250C for 2-3 weeks. The hair is removed and examined microscopically in lactophenol cotton blue-stained wet preparation. T.mentagrophytes cause surface erosion of the shaft resulting in the wedge-shaped appearance of the shaft.

The image below shows surface erosion of shaft caused by Trichophyton mentagrophytes.



Tinea capitis refers to ringworm of the scalp where there is an invasion of the hair shafts by dermatophyte fungi (Trichophyton, Microsporum, and Epidermophyton). It presents as hair loss with inflammation, with or without itching. Diagnosis is based on clinical features, Wood's lamp, and laboratory investigations such as culture with microscopic examination of the colonies.

• Trichophyton: Colonies may be powdery, velvety, or waxy with characteristic pigment based on the species

• T. rubrum: Red

• T. mentagrophytes: White-tan

• T. tonsurans: Cream-yellow

• Microsporum: Colonies are white, cotton-like, velvety, or powdery

• M. audouinii: Brownish

• M. gypseum: Buff colored

• Epidermophyton: Colonies are greenish-yellow and powdery

Treatment is with antifungals like terbinafine, itraconazole, and griseofulvin.

Solution to Question 11:

The initial negative test is explained by the prozone phenomenon. The standard agglutination test is a tube agglutination test that mainly identifies IgM antibodies. It is used in establishing a diagnosis of acute brucellosis.

Prozone phenomenon refers to the failure of a visible precipitation reaction due to inhibition of lattice formation by the presence of excessive antibodies. This gives rise to a false-negative test. This excess in antibodies can be overcome by diluting the serum sample, leading to an increase in the antigen-antibody interactions, eventually giving rise to a positive result.

Lattice hypothesis:

The lattice hypothesis explains precipitation reactions. The amount of precipitate formed is greatly influenced by the relative proportions of antigens and antibodies.

• Prozone phenomenon: Caused by excessive antibodies in the test system. (weak/absent precipitation- false-negative test)

• Zone of equivalence: Antigens and antibodies are in optimum proportions and maximum precipitation occurs.

• Post–zone phenomenon: (Option-D) Due to the presence of excessive antigen in the test system. (weaker/absent precipitation)



Standard agglutination test identifies mainly IgM antibodies, hence it is useful in acute brucellosis. Following 7-10 days of infection, both IgM and IgG antibodies appear. The IgM antibodies then gradually fall and only IgG antibodies persist in chronic brucellosis. The complement fixation test (Option-B) is more useful in chronic cases as it detects the IgG antibody also.

Brucellosis/ Undulant fever:

Brucellosis is a zoonotic disease caused by gram-negative, capnophilic coccobacilli. The most important species that are pathologic to humans are B.melitensis, B. suis, and B.abortus. It is transmitted to humans from contact with or consumption of contaminated products.

These bacteria proliferate within the reticuloendothelial system, which accounts for its resistance to simple antimicrobial therapy.

Patients with acute infection may present with undulant fever, arthralgia, muscular pain, sweating, loss of weight, and appetite. CNS symptoms such as fatigue, irritability, and depression might also occur. Chronic infection tends to be episodic with non-specific symptoms.

Laboratory diagnosis of brucellosis includes:

• Polymerase chain reaction

• Blood culture with Castaneda method

• Serological methods include standard agglutination test (SAT), complement fixation test, Coomb's antiglobulin test, and enzyme-linked immunosorbent assay (ELISA).

• Tests for animals and animal products: Milk ring tests and Rose Bengal card tests.

• Milk ring test- A sample of milk is mixed with stained brucella antigen and incubated. If the sample contains antibodies, agglutination occurs seen as a blue ring.

Treatment includes antimicrobial therapy with IM streptomycin and oral doxycycline. Prevention of brucellosis can be achieved by pasteurization of milk and milk products.

The images below show the Casteneda's medium and the milk ring test respectively.

Castaneda's bottle

©MARROW

Milk ring test in Brucellosis



©MARROW

Microbiology INI-CET 2022

Question 1:

Identify the pathognomonic structure in the slide given below.



a) Sclerotic body

b) Budding yeast

c) Negri body

d) Asteroid body

Question 2:

Antigen binding to MHC class I activates which of the following cells?

a) Cytotoxic T cells

b) B cells

c) NK cells

d) Helper cells

Question 3:

Which of the following agents can be used as a disinfectant but not a sanitizer?

a) Hydrogen peroxide

b) Penicillin

c) Methyl paraben

d) Bleaching powder

Question 4:

Vaccination against which of the following is recommended in the elderly?

a) 1,2,3,4

b) 1,3

c) 1,4

d) 2,3

Question 5:

Which of the following is wrong regarding primary and secondary immune response ?

a) 1

b) 2

c) 3

d) 4

Question 6:

Which of the following is true about anti-CMV IgG antibodies?

a) Denotes latent CMV infection

b) Denotes chronic CMV infection with immunity to other serotypes

c) Indicates acute CMV infection

d) IgG avidity assay helps in differentiating past and primary infection

Question 7:

Which of the following is an example of Mollicutes?

a) Ureaplasma urealyticum

b) Orientia tsutsugamushi

c) Chlamydophila pneumoniae

d) Chlamydia trachomatis

Question 8:

Which of the following statements regarding the given image is correct?



a) Albendazole is the drug of choice

b) Majority of infections are asymptomatic in humans

c) Infection is acquired by ingestion of aquatic plants

d) Non-segmented leaf like parasite

Question 9:

A patient presents to you with left-sided flank pain. A CT scan was done which is shown below. Which of the following statements is incorrect regarding the condition?



a) Albendazole can be started preoperatively

b) Cyst contents can produce anaphylactic reaction

c) A segmented dog tapeworm is the causative agent

d) FNAC or biopsy to be done before nephrectomy

Question 10:

Which of the following has been reported to have endemic transmission in India?

a) Yellow fever

b) Crimean-Congo hemorrhagic fever

c) Ebola

d) Hendra virus

Question 11:

Which of the following are acid-fast staining organisms?

a) 1, 2, 3, 4, 5

b) 1, 2, 4, 5

c) 1, 2, 3

d) 3, 4, 5

Question 12:

The cell wall depicted below is most likely to belong to which of the following organisms?



a) Aspergillus fumigatus

b) Mycobacterium tuberculosis

c) Mycoplasma pneumoniae

d) Rhinosporidium seeberi

Question 13:

The organism with following life cycle is:



a) Strongyloides stercoralis

b) Ancylostoma duodenale

c) Ascaris lumbricoides

d) Enterobius vermicularis

Question 14:

What is the ratio of T cells to B cells in a healthy adult?

a) 1:1

b) 3:1

c) 1:2

d) 1:3

Question 15:

A microbiologist stained a tissue sample and found thin, filamentous bacteria, which resembled Nocardia sp. What other staining technique should be done to confirm the presence of Nocardia?



a) Ziehl-Neelsen staining

b) Silver impregnation

c) Albert stain

d) Modified acid-fast stain

Question 16:

What is the microscopy technique to produce high contrast image of transparent, live bacteria?

a) Phase contrast microscopy

b) Dark field microscopy

c) Electron microscopy

d) Polarized microscopy

Question 17:

Match the following.

a) A-4, B-1, C-3, D-2

b) A-4, B-1, C-2, D-3

c) A-4, B-2, C-3, D-1

d) A-2, B-1, C-3, D-4

Question 18:

Vesicular rash is seen in _____.

a) Hand, foot, and mouth disease

b) Dengue

c) Zika fever

d) Roseola infantum

Question 19:

Which of the following viruses show genetic change similar to influenza virus?



a) 1

b) 2

c) 3

d) 4

Question 20:

A forest worker developed skin lesions over the forearm, which initially started as macules but then became nodules. Histology of the nodules shows the following findings. Which of the following is true regarding this condition?



a) The organism does not invade underlying fascia, tendon, muscle, or joints.

b) Angioinvasion is common, especially in people with hemolytic anemia.

c) It is a dimorphic organism, and yeast forms are seen in clinical sample.

d) These bodies are formed by engulfment of the dead fungi by the macrophages.

Question 21:

An HIV patient came to the clinic with a history of diarrhea. Stool microscopy showed the oocysts that were 4-8 µm in size. The image is shown below. Identify the organism.



a) Isospora

b) Cyclospora

c) Cryptosporidium

d) Microspora

Question 22:

A patient presents to the hospital with complaints of abdominal pain. A stool examination was done, and it showed the presence of the organism shown in image A. The body has segments broader than the length. Which of the eggs shown in column B corresponds to this organism?



a) 1

b) 2

c) 3

d) 4

Question 23:

The image below shows the life cycle of a virus. Which of the proteins of the virus act as oncogenes?



a) E1, E2, E5

b) E6, E7

c) L1, L2

d) E1, E2

Question 24:

Image of Rickettsial morula in a monocyte is shown below. Identify the organism.



a) Rickettsia rickettsii

b) Rickettsia conorii

c) Ehrlichia chaffeensis

d) Anaplasma

Question 25:

A young man presents with skin lesions as shown in the image below. All of the following organisms can spread through dermal and subcutaneous lymphatics, except _____.

a) Sporothrix schenckii

b) Staphylococcus aureus

c) Nocardia asteroides

d) Mycobacterium marinum

Question 26:

The following are parvovirus-related manifestations. Identify the incorrect pair.

a) 4

b) 3

c) 1

d) 2

Question 27:

Which virus can be identified by a PCR method and is endemic to India?

a) Hendra virus

b) West nile virus

c) Ebola virus

d) Yellow fever

Question 28:

SARS-CoV-2 can be detected by all except _____.

a) ELISA to detect spike proteins

b) Real-time PCR to detect viral mRNA

c) Rapid antigen test to detect spike proteins

d) Southern blotting

Question 29:

A patient comes with a history of high fever for 5 days and altered sensorium. On further probing, it was found that he had two seizures one day ago. Peripheral smear shows the following findings. How will you manage this patient?



a) Doxycycline 100 mg intravenously for 7 days

b) Clindamycin 600 mg intravenously 8 hourly for 7 days

c) Artesunate 2.4 mg/kg intravenously at 12 and 24 hours, followed by daily once

d) Chloroquine intramuscularly 8 hourly for 3 days

Question 30:

Delayed hypersensitivity is due to _____.

a) Interferon gamma

b) Interleukin-2

c) Interleukin-12

d) Interleukin-4

Question 31:

Which disease causes the lesion shown in the image below?

a) Rocky mountain spotted fever

b) Malaria

c) COVID-19

d) Scrub typhus

Question 32:

Which of the following method of disinfection and sterilization can also kill spores?

a) 1, 3 and 4

b) 1, 2 and 4

c) 2 ,3 and 4

d) 1,2,3 and 4

Question 33:

The inclusion body rickettsial morula is found in neutrophils during infection caused by:

a) Rickettsia rickettsii

b) Rickettsia conorii

c) Anaplasma phagocytophilum

d) Ehrlichia chaffeensis

Question 34:

A petechial rash is associated with all the following, except _____.

a) Zika virus

b) Roseola infantum

c) Kyasanur forest disease

d) West Nile fever

Question 35:

Mode of transmission of abdominal TB includes all except

a) Direct spread from mesenteric lymph nodes

b) Hematogenous spread

c) Feco-oral route

d) Direct contact with infected animal

Question 36:

Choose the egg belonging to the following parasite.



a) 1

b) 2

c) 3

d) 4

Question 37:

Which of the following is a gram-positive organism that shows the following appearance on Ziehl-Neelsen staining?



a) Nocardia

b) Mycocbacterium tuberculosis

c) Mycobacterium leprae

d) Actinomyces

Question 38:

The following are parvovirus-related disease-population pairs. Identify the incorrect pair.

a) 1

b) 2

c) 3

d) 4

Question 39:

A patient with a history of fever, jaundice, hepatosplenomegaly, and laboratory test result of leptospira IgM reactivity, now presented with breathlessness, fever, and jaundice that worsened after treatment with penicillin. What is the possible diagnosis?

a) Resistant leptospirosis

b) Penicillin allergy

c) Co-infection with dengue

d) Jarisch-Herxheimer reaction

Question 40:

On inoculation in the skin, all of the following organisms can lead to dermal, subcutaneous, and lymphatic involvement except_____

a) Sporothrix schenckii

b) Corynebacterium minutissimum

c) Nocardia brasiliensis

d) Mycobacterium marinum

Question 41:

On Ziehl-Neelsen (ZN) staining oocysts of size 8-10 µm cysts are visible. Identify the organism.



a) Isospora

b) Cryptosporidium

c) Cyclospora

d) Microspora

Question 42:

Vaccine can be prepared against which of the following proteins of the virus?



a) E6, E7

b) L1

c) E1, E2

d) E3

Question 43:

The Epstein-Barr virus interacts with B cells via which of the following?

a) CD21

b) CD20

c) CD19

d) CD23

Question 44:

A forest worker developed skin lesions over the forearm, which initially started as macules but then became nodules. Histology of the nodules shows the following findings. Which of the following is true regarding this condition?



a) It is a dematiaceous fungus

b) Infection can spread and involve underlying subcutaneous tissue, tendon, muscle or joints

c) Angioinvasion is common especially in people with hemolytic anemia

d) These bodies are formed by engulfment of the dead fungi by the macrophages

Question 45:

The viruses of the filoviridae family like Ebola and Marburg resemble which of the following morphologies?



a) 3

b) 2

c) 1

d) 4

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	c
5	d
6	d
7	a
8	b
9	d
10	b
11	b
12	a
13	a
14	b

15	d
16	a
17	b
18	a
19	a
20	a
21	c
22	d
23	b
24	c
25	b
26	d
27	b
28	d
29	c
30	a
31	d
32	b
33	c
34	b
35	c
36	c
37	a
38	d
39	d
40	b
41	c
42	b
43	a
44	a
45	a

Detailed Explanations

Solution to Question 1:

The given image shows the presence of brown round sclerotic bodies which is characteristically seen in chromoblastomycosis.

Chromoblastomycosis is a chronic fungal infection of the skin and subcutaneous tissue. It is caused by pigmented fungi (most commonly by Phialophora verrucosa) and is commonly seen in tropical areas among agricultural workers and woodcutters who walk barefoot.

Patients present with warty cutaneous nodules (cauliflower-like) on the feet and lower legs. Diagnosis is by demonstration of sclerotic bodies in tissue sections. Sclerotic bodies appear as round or irregular, dark brown bodies with septae. It is usually seen in clusters of 4-8 cells due to septation in different planes and delayed separation.

Chromoblastomycosis is managed by:

• First-line treatment involves itraconazole (100-200mg daily) or terbinafine (250mg daily)

• Second-line treatment includes the use of cryotherapy or the local application of heat by heat-retaining gels for a period of 2-3 months.

• Surgery is only indicated in very small lesions and should always be combined with chemotherapy

• Flucytosine can also be used in combination with amphotericin B, but should not be used alone due to the development of resistance.

Other options:

Option B - The below image shows multiple round or oval-shaped yeast cells, pseudohyphae, and buds attached to the parent cell. They are seen in budding yeast.



Option C - Negri bodies are pathognomonic of rabies. It is an acidophilic, intracytoplasmic inclusion body with characteristic basophilic inner granules as shown below.



Option D - Asteroid bodies is seen in sporotrichosis, caused by Sporothrix schenckii. It can be seen as an oval, basophilic structure with eosinophilic radiation. These are depositions of antigen-antibody complexes.



Solution to Question 2:

Antigen binding to major histocompatibility complex (MHC) class I activates cytotoxic T cells.

The major histocompatibility complex (MHC), is a genetic locus that encodes a group of antigens. The MHC molecules also bind peptide antigens and present them to the T cells. The human MHC is called the human leukocyte antigen (HLA) complex.

The MHC class I proteins present their antigens to CD8 T cells (cytotoxic T cells). The antigens bound to MHC class II proteins activate the CD4 T cells(T helper cells). There is no cross

recognition between the T cells and MHC classes. This is known as MHC restriction. For example, a CD4 T cell would recognize an antigen presented by a macrophage only if it's bound to MHC class II molecules.

Other options:

Option B - B cells interact with an antigen based on the immunoglobulin surface receptors on the B cell. The receptor along with the antigen is embodied and degraded and the peptides produced are returned to the cell surface. It is then bound to the MHC class II molecules present on the cell surface. This is then recognized by CD4 T cells. Ultimately, the B cells are stimulated to proliferate and differentiate into plasma cells which secrete antibodies.

Option C - Natural killer (NK) cells are granular lymphocytes that confer innate immunity by protecting against viruses and intracellular pathogens.

Option D - T helper cells are responsible for the production of cytokines which activate the humoral and cell-mediated branches of immunity. These cells will recognize only those antigens that are presented along with MHC class II molecules on APC such as macrophages and B lymphocytes.

Solution to Question 3:

Hydrogen peroxide is a disinfectant but cannot be used as a sanitizer.

Sanitization is a process by which levels of pathogenic organisms on inanimate surfaces are brought down to reduce the chances of cross-infection.

Disinfection is a process by which all pathogenic organisms are destroyed.

Sterilization is a process in which all microorganisms in both vegetative and spore state are eliminated from a surface or medium.

Hydrogen peroxide can be used for disinfection or sterilization in the liquid form for cystoscopes, endoscopes, tubings, and metal instruments. The gaseous form can be used for fogging, which has largely replaced fumigation. It has a shorter cycle time and is non-toxic. The mechanism of bactericidal action is the oxidation of the cell wall of the microbe. It is also used in plasma sterilization.

Other options:

Option B - Penicillin, an antibiotic interferes with the synthesis of the bacterial cell wall. It is not used as a disinfectant or sanitizer.

Option C - Methyl parabens are antimicrobials that are commonly used in food and personal care products.

Option D - Household bleaching powder (sodium hypochlorite) is a widely used disinfectant in hospitals. They are oxidizing agents that interfere with cellular protein activity. Diluted solutions are also used for sanitizing surfaces.

Solution to Question 4:

Vaccination against varicella-zoster and pneumococcal meningitis (Options 1 and 4) is recommended in the elderly.

Vaccinations against the following diseases are recommended for the elderly:

• Influenza vaccine

• Pneumococcal vaccine

• Herpes zoster

• Tetanus, diphteria, and pertussis

• Hepatitis B

Solution to Question 5:

Statement 4 is wrong regarding primary and secondary immune responses. Primary immune response has lower antibody affinity and vice versa.

An immune response is a specific reactivity that occurred in a host as a result of an antigenic stimulus. It is of two types: Humoral or antibody-mediated response and cellular or cell-mediated response.

The response to the antigenic stimulus can be primary or secondary. The features of the primary and secondary immune responses are as follows

Features	Primary immune response	Secondary immune response
1. Lag period	4-10 days	1-3 days
2. Peak response time	7-10 days	3-5 days
3. Antibody response	Depends on the antigen	More potent than the primary immune response
4. Isotype generate	IgM	IgG
5. Antigens	Both thymus-dependent and independent	Depends only on the thymus
6. Antibody affinity	Lower	Higher

Solution to Question 6:

IgG avidity assay helps in differentiating past and primary infection with cytomegalovirus (CMV).

Avidity level measured by avidity index is used to describe the percentage of IgG bound to the antigen after being treated with denaturing agents.These assays are based on the fact that low avidity IgG antibodies are present during the first few months after the infection and avidity increases over time indicating maturation of the immune response. So the presence of high avidity IgG denotes a past infection rather than a primary one. Low avidity IgG signals a primary

infection within the last 3-4 months.

IgM detection can be used as an indicator of acute infection, but it lacks specificity for primary infection. This is because it persists for years in some individuals and is also produced during reactivation.

Cytomegalovirus is a type of herpes virus, which causes severe disease in immunocompromised patients. In most other cases, it is inapparent and marked by prolonged latency with intermittent reactivation.

Solution to Question 7:

Ureaplasma urealyticum belong to the class Mollicutes which are also known as cell-wall-free bacteria. This class contains around 200 species of mycoplasma.

Mycoplasma are the smallest microorganisms that can thrive in a cell-free medium. They are pleomorphic and lack cell walls. There are 4 species that are of importance in humans.

• Ureaplasma urealyticum

• Mycoplasma pneumoniae

• Mycoplasma hominis

• Mycoplasma genitalium

It can be grown in fluid or solid media. Colonies typically have a fried egg appearance. They mainly cause pneumonia and genital infections. Ureaplasma urealyticum is unique in it's ability to hydrolyse urea. They cause non-gonococcal urethritis. It can be resistant to commonly used antibiotics such as tetracycline and doxycycline.

Other options:

Option B - Orientia tsutsugamushi belongs to the family rickettsiaceae. It causes scrub typhus, which is spread by mites.

Option C - Chlamydophila pneumoniae belongs to the genus Chlamydia. It is a common cause of respiratory diseases such as sinusitis, bronchitis, and pneumonia.

Option D - Chlamydia trachomatis also belongs to the genus Chlamydia. It causes trachoma, inclusion conjunctivitis, and genital chlamydiasis.

Solution to Question 8:

The given image shows a spherical or ovoid, non-bile stained egg which is diagnostic of Hymenolepis nana infection. It has a thin colorless outer membrane and a thin embryophore covering the hexacanth embryo. The majority of infections are asymptomatic and are more commonly seen in children.

H. Nana or dwarf tapeworm is the smallest and most common cestode that infects man. There is no intermediate host. The infection is acquired by ingestion of food and water contaminated with eggs.

Patients with heavy infections may present with nausea, abdominal pain, anorexia, diarrhea, and irritability. An allergic response to the parasite can cause pruritus as well.

A single dose of Praziquantel given at the rate of 25mg/kg is the drug of choice. Nitazoxanide (500 mg twice daily for 3 days) may be given as an alternative.

Image below shows egg of H. Nana.



Other options:

Option C - Following are the parasites with aquatic plants as the source of infection :

• Fasciola hepatica

• Fasciolopsis buski

• Watsonius watsoni.

• Gastrodiscoides hominis

Option D - The adult worm of Fasciola hepatica is a flat leaf-shaped liver fluke.

Solution to Question 9:

FNAC or biopsy to be done before nephrectomy is the incorrect statement regarding cystic echinococcosis. The given CT image shows left-sided renal hydatid cyst.

Echinococcosis (hydatidosis) is caused by Echinococcus granulosus (larval stage). It is a segmented tapeworm. Dogs are the definitive hosts and pass the eggs in their feces. Intermediate hosts include humans, sheep, and cattle. Patients present with abdominal pain, palpable mass, jaundice, cough, and dyspnea. Rupture of a hydatic cyst can result in fever, pruritus, or anaphylaxis.

Daughter cysts within the larger cyst seen on CT imaging is diagnostic. A biopsy is not recommended because of the risk of fluid leakage which can result in anaphylaxis or dissemination of the infection.

Percutaneous aspiration, infusion of scolicidal agents, and reaspiration (PAIR) is the primary treatment. It is contraindicated in superficial loculated cysts and in cysts that communicate with the biliary tree.

Isolated involvement of the kidneys is rare. Patients can present with nonspecific symptoms, renal colic, and hydatiduria. Due to the lack of an effective systemic scolicidal agent, surgical treatment is necessary when kidneys are involved. Simple excision of the cyst is preferred. If there is extensive involvement, partial or total nephrectomy is warranted. It is recommended to start albendazole one week to one month before the surgery to decrease the risk of spillage.

Solution to Question 10:

Crimean-Congo hemorrhagic fever caused by nairovirus has an endemic transmission in India. An outbreak was recognized in Gujarat in 2011. In India, antibodies against the virus have been detected in human and animal sera. It is transmitted by hyalomma ticks.

Other options:

Option A - Yellow fever is mostly confined to the south and central America, and certain parts of Africa. It is caused by the flavivirus. It is an arthropod-borne hemorrhagic fever. In the urban cycle, Aedes aegypti transmits the disease.

Option C - Ebola virus causes a hemorrhagic fever first reported in Sudan and Zaire. A recent outbreak was reported in West Africa.

Option D - Hendra virus was responsible for the death of many horses in Australia.

Solution to Question 11:

Nocardia, Mycobacterium leprae, Cryptosporidium parvum, and Cytoisospora belli (Options 1, 2, 4, and 5) are acid-fast staining organisms.

Acid-fast staining is based on the principle that tubercle bacilli resist decolorization with acids after staining with aniline dyes. Acid fastness is attributed to the variety of fatty acids, alcohols, and lipids in the tubercle bacilli. Acid-fast organisms usually contain mycolic acid, which is acid-fast in the free state. The steps to carry out acid-fast staining as modified by Ziehl and Neelson are:

• The slide is stained with carbolfuchsin along with the application of heat.

• It is then decolorized with 20% sulphuric acid. Later methylene blue is used for counterstaining.

• The acid-fast bacteria retain the fuchsin color, while other organisms take up the counterstain.

The genus Nocardia are aerobic organisms and weakly acid-fast positive rods. They stain with carbolfuchsin and retain it when 1-4% sulfuric acid is used.



Mycobacterium leprae, the causative agent of leprosy is a gram-positive bacillus. It is acid-fast when 5% sulphuric acid is used instead of 20%.



Cryptosporidium parvum, commonly causes diarrhea in immunocompromised individuals. The oocysts in the stool can be visualized by modified Ziehl-Neelsen staining. It appears as red acid-fast spheres. It can also be used for specimens such as sputum and bronchial wash.



Isospora belli is a coccidian parasite that can cause protracted diarrhea in immunocompromised patients. Modified Ziehl-Neelsen staining reveals pink-colored acid-fast oocyst in stool samples.



Actinomyces are gram-positive, anaerobic, and filamentous bacteria that are non-acid-fast in contrast to Nocardia.



Solution to Question 12:

The organism most likely to have the cell wall depicted is Aspergillus fumigatus.

Fungi usually have a rigid cell wall composed of polysaccharides, glycoproteins, and lipids. Most of them contain sugar polymers such as chitin, glucans, and mannans. The cytoplasmic membrane is usually composed of sterols. Antifungals such as amphotericin and nystatin bind to the ergosterol in the cell membranes which results in membrane damage and leakage.

Aspergillus fumigatus is a filamentous fungi, with hyphae that may have cross-walls or septa. It can cause allergic bronchopulmonary aspergillosis, aspergilloma, and superficial infections of the ear, eye, and sinuses. It causes invasive aspergillosis in immunocompromised individuals.

Other options:

Option B - Mycobacterium tuberculosis, are acid-fast staining bacteria, owing to the mycolic acid in the cell wall. They have a three-layered cell wall enclosing a trilaminar plasma membrane.

Option C - Mycoplasma pneumoniae lacks cells wall or even cell wall precursors. They are bound by a trilaminar membrane that contains sterols.

Option D - Rhinosporidium seeberi causes a granulomatous condition known as rhinosporidiosis. It exists as a trophocyte with clear cytoplasm and a covering of chitin. Mature trophocytes develop into sporangium with an outer chitinous and inner cellulose layer.

Solution to Question 13:

The given life cycle is of Strongyloides stercoralis.

The organism has a parasitic cycle and a free-living soil cycle. Humans are the natural host but morphologically indistinguishable strains may be seen in dogs and cats.

Filariform larvae are the infective form and there are two modes of infection:

• Penetration of skin by 3rd stage filariform larva.

• Autoinfection (external and internal).

The female worms present in the intestine lays eggs in the mucosa which hatches immediately, releasing the rhabditiform larva. The larva migrates into the lumen of the intestine and passes down to be released in the feces. The rhabditiform larva may change into filariform larva while passing through the bowel.

The filariform larvae in the bowel do not leave the host and become a source of autoinfection as they penetrate the colonic mucosa or perianal skin.

The rhabditiform larvae passed in the feces may undergo two types of development in the soil:

• Direct development: The rhabditiform larva moults twice on reaching the soil to become the infective filariform larva. The filariform larva can penetrate the skin and enter the circulation.

• Indirect development: The rhabditiform larvae develop into free-living males and females in the soil.

Other options:

Option B: Humans are the only natural host and the life-cycle of Ancylostoma duodenale is completed in a single host. The eggs laid by the female worms in the intestine are passed in feces. The embryo develops inside the egg and a rhabditiform larva hatches out in about 2 days.

The rhabditiform larva moults twice to form the filariform larva (infective stage). It penetrates the skin and enters the subcutaneous tissue.

Option C: Humans are the only natural host for Ascaris lumbricoides and have no intermediate host. The infection is acquired by ingestion of embryonated eggs (infective stage).

Option D: Enterobius vermicularis completes its entire life cycle in the human host and has no intermediate host. Embryonated eggs are the infective form. Infection is acquired by ingestion of the eggs. Autoinfection and retroinfection are seen.

Solution to Question 14:

The ratio of T cells to B cells in the blood of a healthy adult is approximately 3:1.

T lymphocytes constitute 60-70% of lymphocytes in the blood while mature B cells account for 10-20%.

T cells are of three types: Helper T cells, cytotoxic T cells, and regulatory T cells. T lymphocytes develop in the thymus.

B lymphocytes develop from precursors present in the bone marrow. They differentiate to form either plasma cells or memory B cells. Plasma cells produce antibodies after stimulation by the antigen.

Lymphocyte recirculation is particularly important for T cells, as B cells do not have to move to the site of infection as the antibodies produced are circulated through blood.

Solution to Question 15:

The presence of Nocardia species can be confirmed using a modified acid-fast stain.

For microscopy, sputum smears are most commonly used. Nocardia species are seen as filamentous organisms, with hyphae-like branching. These bacteria are gram-positive and catalase positive, and they produce urease.

The image below shows acid-fast staining of Nocardia.



On direct microscopy of sputum or pus samples, crooked, branching, beaded filaments are seen; these filaments are 1 μm wide and up to 50 μm long.

After incubation at 35–37°C in laboratory media, they develop heaped, irregular, waxy colonies. Strains may vary in their pigmentation from white to orange to red. Blood cultures may be used to detect Nocardia species, although it may take up to 2 weeks to detect growth.

Solution to Question 16:

The microscopy technique used to produce high contrast images of transparent, live bacteria is phase contrast microscopy.

It improves the contrast by making use of the difference in thickness or refractive index of structures within and around the cell. The rays passing through different refractive indices are varied by fraction of wavelength, producing a phase difference. This creates a difference in intensity of light, producing contrast and making the structures within the cells and its surroundings clearly visible.

Option B: In Dark field microscope only the reflected light enters the objective lens whereas the unreflected light does not. Hence the specimen gets brightly illuminated against a dark background. It is used for visualizing very slender, live organisms such as spirochetes which are not visible under ordinary illumination.

Option C: Electron microscopy uses a beam of electrons instead of the beam of light. It is used for studying dead and dried objects. Hence, it may lead to distortion in cell morphology.

Option D: Polarized microscopy is used to study the intracellular structures using differences in birefringence.

Solution to Question 17:

The correct pairs are as follows:

Tuberculosis: Mycobacterium tuberculosis can be stained with Ziehl-Neelsen (ZN) stain. Steps involved in the ZN stain:

• Place the sample on the slide and evenly spread it with rotational movement.

• Air-dry the slide, and fix it by passing it over a flame.

• Apply carbol fuchsin stain.

• Heat the underside of the slide until steam rises.

• Wash with water to remove excess stain.

• Decolorize with alcohol.

• Apply methylene blue stain.

• Wash with water to remove excess stain.

• View under a microscope.

Malaria: Giemsa stain is used as a gold standard procedure for the detection of malarial parasites. This stain helps visualize Maurer's clefts associated with Plasmodium falciparum and Schuffner's dots associated with Plasmodium vivax or Plasmodium ovale.

Leprosy: Acid fastness in general is related to the carbon chain length of the mycolic acid found in any particular species. Acid and alcohol fastness of the lepra bacilli is less than that of the tubercle bacilli. Fite-Faraco method is the staining procedure of choice in Mycobacterium leprae. In this method, alcohol is eliminated from the hydrating and dehydrating steps, and 10% sulphuric acid is used as a decolorizer in place of acid/alcohol solution.

Actinomycosis: Actinomyces is a gram-positive bacillus that is filamentous and non-sporing in nature, and the Gram stain is more sensitive in its diagnosis than culture.

Column A	Column B
A. Malaria	1. Giemsa stain
B. Tuberculosis	2. ZN stain
C. Leprosy	3. Fite-Faraco stain
D. Actinomycosis	4. Gram stain

Solution to Question 18:

Vesicular rash is seen in hand, foot, and mouth disease.

It is a mild exanthematous fever usually occurring in children but may also occur in adults. It is characterized by papulovesicular rashes that involve the oral mucosa, palms, and soles.

The virus is shed in the stool and is transmitted by the feco-oral route. It is a self-limiting disease and subsides in 1-2 weeks. Treatment is symptomatic as the rash heals spontaneously.

Coxsackieviruses are enteroviruses and belong to the Picornaviridae family. They are classified into groups A and B. Group A viruses are classified into 24 types and group B viruses are classified into 6 types.

Diseases caused by Coxsackie A viruses:

• Hand, foot, and mouth disease (HFMD)

• Herpangina- It is febrile vesicular pharyngitis. It is characterized by the presence of ulcerating small vesicles on the oral fauces, or palate or uvula, and the posterior pharyngeal wall.

• Aseptic meningitis

Diseases caused by Coxsackie B viruses:

• Pleurodynia or Bornholm disease- It is viral myalgia defined by sudden severe lancinating pain, typically on one side of the chest. It is commonly associated with fever, malaise, and headaches.

• Myocarditis and pericarditis in the newborn.

• Orchitis, juvenile diabetes

The image below shows the rash in the hand, foot, and mouth disease.



Other options:

Option B: Cutaneous and mucosal findings in dengue are erythema, morbilliform eruptions, and hemorrhagic lesions.

Option C: Zika fever is characterized by fever, maculopapular rash, arthralgia, and conjunctivitis.

Option D: Roseola infantum presents as a nonpruritic, pink papular rash that begins on the trunk.

Solution to Question 19:

Rotavirus exhibits genetic reassortment/antigenic drift similar to that of the influenza virus.

Rotaviruses appear as cartwheel-shaped structures with short spokes radiating from them. 11 segments of double-stranded RNA constitute the genome. Influenza viruses have a linear, segmented, single-stranded RNA. Reassortment of the genome occurs readily due to the presence of this segmented RNA.



Other options:

Option B - Adenovirus contains double-stranded DNA. It has icosahedral symmetry, with structures known as fibers extruding from the vertices.



Option C - Ebola viruses are filamentous viruses with nonsegmented single-stranded RNA as genetic material.



Option D - Rabies viruses belonging to the family Rhabdoviridae are bullet-shaped viruses, with nonsegmented single-stranded RNA.



Solution to Question 20:

The most likely diagnosis based on the clinical vignette and the histopathology showing sclerotic bodies is chromoblastomycosis. It is primarily a subcutaneous mycosis and does not commonly invade underlying tissue, tendon, or joints.

Chromoblastomycosis is caused by dematiaceous fungi, primarily
Fonsecaea spp and Cladophialophora spp. Not all species of dematiaceous fungi are known to be

thermally dimorphic fungi (Option C)

Chromoblastomycosis, the most common chromomycosis, is more commonly seen in tropical areas among agricultural workers and woodcutters who walk barefoot. Traumatic implantation is the mode of infection. Most often, the infection occurs in trauma-prone sites, such as exposed areas of the lower and upper limbs.

Patients present with warty cauliflower-like cutaneous nodules on the feet and lower legs. The disease generally remains localized, with slow peripheral extension. Lymphatic spread can occur.

Diagnosis is made by the demonstration of sclerotic bodies, also called copper-penny bodies, in tissue sections. Sclerotic bodies appear as round or irregular, dark brown bodies with septae.



First-line management includes itraconazole or terbinafine. Flucytosine used on its own or combined with amphotericin B may also be effective. For small lesions, surgical excision is the therapy of choice.

Solution to Question 21:

The given clinical scenario and the stool microscopy show oocysts of Cryptosporidium.

Cryptosporidium is a coccidian parasite implicated in causing traveller's diarrhea in immunocompromised patients. The two cryptosporidium species responsible are C. hominis and C. parvum.

The infective form is the oocyst excreted in the patient's feces, which can be acquired through drinking water or recreational water activities. The oocysts are immediately infectious (unlike isospora oocysts, which require maturation), and may cause autoinfection.

Because of their small size, they are difficult to see using routine staining and hence, acid-fast staining is done.

Once ingested, the 4 sporozoites are released from each cyst and attach to the small intestinal mucosa. It causes crampy abdominal pain and non-inflammatory watery diarrhea.

Treatment is mainly supportive, though nitazoxanide has been approved for use in cryptosporidium infection.

Other options:

Option A: Oocysts of isospora are 23-36 μm, oval, and contain two sporocysts, each having four sporozoites. They are uniformly acid-fast.

Option B: Sporulated oocysts of cyclospora are 8-12 μm in size, round, and contain 2 sporocysts each having two sporozoites. They are acid-fast.

Option D: Microspora oocysts are 2-4 μm in size, oval to cylindrical in shape with a polar filament. Spores are gram-positive and acid-fast.

Solution to Question 22:

The given image shows a scolex with 4 suckers and a retracted rostellum with a single row of hooklets. The proglottids are broader than long. These features are characteristic of Hymenolepis nana.

Option 4 shows the appearance of the egg of H. nana on stool microscopy. These eggs have a thin, colorless outer membrane and thread-like polar filaments. They have a hexacanth embryo. The eggs are non-bile stained and float in a saturated salt solution.

H. nana, also called dwarf tapeworm, is the smallest tapeworm. Humans are definitive hosts. Infection is acquired by the ingestion of food and water contaminated with eggs. The egg attaches to the intestine and develops into the larva and adult worm. The adult worm resides in the small intestine. No intermediate host is required. H. nana is unusual in that it undergoes multiplication in the body of the definitive host.

Patients are usually asymptomatic. But, in heavy infection, they present with nausea, anorexia, abdominal pain, diarrhea, and irritability. Pruritus may also be present due to an allergic response.

The image given below shows the life cycle of Hymenolepis nana.



Solution to Question 23:

The image shows the life cycle of the human papilloma virus, and the E6 and E7 proteins act as oncogenes.

The E6 protein binds to and helps in the degradation of p53 and stimulates the expression of telomerase reverse transcriptase (TERT), which is the catalytic subunit of telomerase, thereby helping in the longevity of the cells.

E7 protein binds to the RB protein and displaces the E2F transcription factors that are normally sequestered by RB, thus promoting progression through the cell cycle.

L1 forms the major portion of the capsid. L2 plays an important role in papillomavirus assembly and the infectious process. E1 and E2 are involved in replication and transcription.

Solution to Question 24:

The organism given in the image is Ehrlichia chaffeensis. The peripheral blood smear shows a morula, an intracellular inclusion body in a monocyte.

Ehrlichia chaffeensis is a gram-negative, obligate intracellular bacteria that are transmitted by hard ticks such as the lone star tick.

Clinically, the symptom onset is typically within one-three weeks of exposure. Patients may present with fever, headache, myalgia, nausea, vomiting, loss of appetite, and weight loss. Rashes are generally absent. Severe human monocytic ehrlichiosis is characterized by meningoencephalitis, renal failure, respiratory failure, and shock.

Diagnosis is by demonstration of the characteristic morulae in the mononuclear cells such as monocytes. Serological studies such as indirect fluorescent antibody tests and polymerase chain reaction (PCR) detection can also be used.

The treatment of choice for human monocytic ehrlichiosis is doxycycline.

Other options:

Option A: Rickettsia rickettsii causes rocky mountain spotted fever. It is transmitted by ticks. Patients present with fever, headache and rash.

Option B: Rickettsia conorii causes Mediterranean spotted fever. The parasite is transmitted by ticks and the patients present with fever, headache, rash, and eschar (tache noire).

Option D: Anaplasma causes human granulocyte anaplasmosis. It is transmitted by ticks. Clinical features are fever, headache, and myalgia. Inclusions in granulocytes are seen.

Solution to Question 25:

The given image shows multiple vesicles on hand and multiple eruptions along lymphatics. All of the organisms mentioned in the options can spread through dermal and subcutaneous lymphatics except Staphylococcus aureus

Staphylococcus aureus is a gram-positive coccus arranged in grape-like clusters. It causes various localized pyogenic infections like folliculitis, furuncle, abscess, impetigo, paronychia, osteomyelitis, and even septicemia.

Other options:

Option A: Sporothrix schenckii is a dimorphic fungus. It causes sporotrichosis, a chronic granulomatous disease, by traumatic inoculation into the skin. This is usually followed by spread along the draining lymphatics and lymph nodes. Itraconazole is the drug of choice for treatment.

Option C: Nocardia asteroides is a gram-positive aerobic bacteria. It is frequently found in the soil and infection may be exogenous. Cutaneous infections can present as local abscesses or lymphocutaneous lesions.

Option D: Mycobacterium marinum is a non-tuberculous mycobacterium. It causes swimming pool granuloma. The lesion begins as a papule and breaks down to form an indolent ulcer. It usually occurs after abrasions and is therefore seen on the prominences like elbows, knees, ankles, or toes.

The image below shows sporotrichosis.

Solution to Question 26:

Polymyalgia rheumatica is not associated with parvovirus B19, and more commonly occurs in women aged 50 and above. It is closely associated with giant cell arteritis.

The main manifestation of symptomatic parvovirus B19V infection is erythema infectiosum, also known as fifth disease or slapped-cheek disease, which most commonly occurs in children of early school age or infants. Fever and mild constitutional symptoms may accompany the rash, which has a typical “slapped cheek” appearance.

Pure red cell aplasia and chronic hemolytic anemia have been reported in patients with immunosuppression due to a wide range of clinical settings, including AIDS, lymphoproliferative disorders, and transplantation.

Parvovirus B19V infection during pregnancy can lead to hydrops fetalis and/or fetal loss. The risk of transplacental fetal infection is 30%, and the risk of fetal loss, predominantly early in the second trimester, is 9%.

Solution to Question 27:

West Nile virus can be identified by polymerase chain reaction (PCR) and is endemic to India.

West Nile virus is a flavivirus that is transmitted by Culex mosquitoes. It can be subclinical or may present as a mild influenza-like illness. Encephalitis occurs in severe forms. Reverse-transcriptase PCR has been used to identify the virus.

Other options:

Option A- Hendra virus is endemic to Australia, and mostly affects horses. Some human cases are also reported.

Option C - Ebola virus belongs to the genus Filovirus. It causes hemorrhagic fever. It is endemic to Africa.

Option D - Yellow fever is mostly confined to South and Central America, and certain parts of Africa. It is caused by flavivirus. It is an arthropod-borne hemorrhagic fever. Aedes aegypti transmits the disease.

Solution to Question 28:

SARS-CoV-2 can be detected by all except Southern blotting.

Southern blotting cannot be used to detect SARS-CoV-2 because there is usually not enough DNA in the collected sample to obtain a pattern on gel electrophoresis.

Real-time PCR uses nasopharyngeal secretions to detect viral genomic RNA or DNA in respiratory secretions. It is the most sensitive and specific test available.

Rapid-antigen test detects spike proteins in nasopharyngeal secretions.

Enzyme-linked immunosorbent assay (ELISA) against spike protein is also found to be sensitive and specific for SARS-CoV-2.

Solution to Question 29:

The given scenario and image from the peripheral smear showing rings (headphones appearance) point towards the diagnosis of severe Plasmodium falciparum malaria, and the treatment of choice for this condition is artesunate 2.4 mg/kg IV at 12 and 24 hours, followed by daily once for 4 days.

Major manifestations of severe falciparum malaria include more than two convulsions in 24 hours, unarousable coma/ cerebral malaria (failure to localize or respond appropriately to stimuli or coma persisting for >30 min after generalized convulsion), acidosis, severe normocytic normochromic anemia, renal failure, pulmonary edema, hypoglycemia, shock, and disseminated intravascular coagulation.

Other manifestations include hemoglobinuria, hyperparasitemia, jaundice, and extreme weakness.

The treatment of choice is artesunate given at a dose of 2.4 mg/kg stat IV, followed by 2.4 mg/kg at 12 and 24 hours, and then daily if necessary.

If artesunate is unavailable, artemether is given at a dose of 3.2 mg/kg stat IM, followed by 1.6 mg/kg QD, or quinine dihydrochloride given at a dose of 20 mg of salt/kg infused over 4 hours, followed by 10 mg of salt/kg infused over 2–8 hours.

Solution to Question 30:

Delayed hypersensitivity is due to interferon-gamma (IFN-gamma).

Delayed hypersensitivity is also called cell-mediated/type 4 hypersensitivity reaction. It is mediated by T cells and occurs by 2 mechanisms:

• CD4+ T helper cells mediated

• CD8+ T helper cells mediated

CD4+ T helper cells mediated pathway:

Antigen peptides recognized by CD4+ T cells secrete interluekin -2. Interleukin -2 helps in further proliferation of T cells that respond to the antigen peptide. These T cells differentiate into Th1 or Th17 cells as a result of the cytokines produced from the antigen presenting cells (APC).

APCs also secrete interleukin - 12 which enhances differentiation of the T cells to the Th1 subset. Inflammatory cytokines like interluekins -1, 6, 23 induce differentiation of T cells to the Th17 subset. These responses lead to the release of interferon gamma that alters the macrophages in several ways to ensure eliminating the antigen.

Interferon-gamma-induced changes in the macrophages:

• Augmented ability of the macrophages to phagocytose and kill micro-organisms.

• Increased expression of class 2 MHC molecules to enhance antigen presentation.

• Increased secretion in presence of tumor necrosis factor, interleukin-1, and other inflammatory chemokines.

• Amplification of the Th1 response by producing more interleukin-12.

CD8+ T helper cells mediated pathway:

CD8+ T cells destroy antigen-presenting target cells using perforins and granzymes. These mediators are preformed in the lysosomes of a cell. The perforin-granzyme complex enters the target cell by endocytosis. Granzymes cause the activation of caspases inducing apoptosis of the target cell. CD8+ T cells also produce interferon gamma following viral infections and contact allergens. Interferon-gamma then proceeds to stimulate the production of inflammatory mediators and bring about the hypersensitivity reaction.

Examples of delayed hypersensitivity reactions include the tuberculin reaction and contact dermatitis.

Interferon-gamma is an important effector involved in both immune mechanisms of delayed hypersensitivity reactions and hence is the best answer for this question.

Solution to Question 31:

The lesion shown in the image is eschar. It is seen in scrub typhus.

Scrub typhus is caused by Orientia tsutsugamushi and transmitted by trombiculid mites. Humans are infected when they are bitten by the mite larvae known as chiggers. Transovarian transmission of microbes occurs in mites.

The incubation period is 1-3 weeks. The patient typically develops a characteristic eschar which is a punched-out ulcer covered with a blackened scab at the site of the mite bite, with regional lymphadenopathy and maculopapular rash.

Other options:

Option A: Rocky Mountain spotted fever is a rickettsial infection transmitted by ticks (Dermacentor andersoni). It is the most serious type of fever. Symptoms include fever, headache, Rash, muscle pain and lack of appetite. Early treatment with doxycycline is recommended to prevent death and severe illness.

Option B: Malaria is a protozoal disease caused by Plasmodium species. It is transmitted by the female anopheles mosquito. Here, the definitive host is the anopheles mosquito and humans are the intermediate hosts. Clinical features are periodic bouts of fever with chills and rigors, followed by anemia, splenomegaly, and hepatomegaly.

Option D: COVID-19 is a highly contagious viral disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The primary mode of transmission is through respiratory droplets. The median incubation period is around 5 days. Patients commonly present with fever, cough, and shortness of breath and less commonly with a sore throat, anosmia, anorexia, nausea, malaise, myalgias, and diarrhea.

Solution to Question 32:

Ethylene oxide, glutaraldehyde, and ortho-phthalaldehyde are disinfectants with sporicidal activity.

Ethylene oxide is a gas used to sterilize heat-sensitive articles such as syringes, needles, catheters, and gloves. Bacillus atrophaeus and Bacillus subtilis are used as sterilization control.

Glutaraldehyde and ortho-phthalaldehyde are liquid-chemical sterilizing agents. They are used for the disinfection and sterilization of cystoscopes, bronchoscopes, rubber anesthetic tubes, endotracheal tubes, polythene tubing, and other metal instruments.

Pasteurization is when milk is heated at either 63°C for 30 minutes (the holder method) or at 72°C for 15–20 seconds (the flash process), followed by cooling quickly to 13°C or lower. It has no sporicidal activity and kills non-spore-forming bacteria like Mycobacteria, Brucella, and Salmonella. Coxiella burnetii is relatively heat resistant and may survive the holder method.

Solution to Question 33:

The inclusion body rickettsial morula is found in granulocytes (neutrophils) during infection caused by Anaplasma phagocytophilum.

Anaplasma causes human granulocyte anaplasmosis. It is transmitted by ticks. Clinical features are fever, headache, confusion, vomiting and myalgia. Inclusions in granulocytes are seen. The image below shows a morula of Anaplasma in a neutrophil:



Ehrlichia chaffeensis is a gram-negative, obligate intracellular bacterium that is transmitted by hard ticks such as the lone star tick. Clinical presentation of Ehrlichia chaffeensis infection is similar to that of Anaplasma infection. Severe human monocytic ehrlichiosis is characterized by meningoencephalitis, renal failure, respiratory failure, and shock. The treatment of choice for human monocytic ehrlichiosis is doxycycline.

The image below shows a morula of Ehrlichia in a monocyte:



Note: Infection by both Anaplasma and Ehrlichia can lead to the formation of rickettsial morulas in leukocytes. However, Anaplasma leads to the formation of morulas predominantly in the granulocytes, whereas Ehrlichia predominantly affects the monocytes.

Other options:

Option A: Rickettsia rickettsii causes rocky mountain spotted fever. It is transmitted by ticks. Patients present with fever, headache, and rash.

Option B: Rickettsia conorii causes Mediterranean spotted fever. The parasite is transmitted by ticks, and the patients present with fever, headache, rash, and eschar (tache noire).

Solution to Question 34:

Roseola infantum is not associated with petechial rashes.

Roseola infantum, also known as exanthem subitum or sixth disease, is caused by human herpes viruses 6 and 7 and is an acute febrile illness that presents with a rose-pink maculopapular eruption after the fever subsides. It initially develops on the neck and trunk and may spread to the arms, face, and leg. They are surrounded by pallor and rarely become vesicular.

The causative agents of Zika, Kyasanur Forest disease, and West Nile fever belong to Flaviviridae group of viruses, which are known to cause hemorrhagic fever and present with petechial rashes.

Note: Although the Zika virus classically presents with a maculopapular rash. A minority of cases present with petechial rashes as well. Therefore, the better answer to this question is roseola infantum.

Solution to Question 35:

Abdominal tuberculosis (TB) is not transmitted by feco-oral route.

The terminal ileum and the cecum are the sites most commonly involved in abdominal TB. It causes abdominal pain (at times similar to that associated with appendicitis) and obstruction, hematochezia, palpable mass in the abdomen along with fever, weight loss, anorexia, and night sweats.

Modes of spread:

• By ingestion of infected food or milk (primary intestinal TB) or infected sputum (secondary intestinal TB)

• Contagious spread from infected adjacent foci

• Hematogenous spread from distant TB focus

• Through lymphatic channels

• Direct contact with an infected animal (Mycobacterium bovis)

There is no evidence of spread of TB through the feco-oral route.

Solution to Question 36:

The correctly matched egg for the given parasite - Enterobius vermicularis is 3

The picture on the left shows a worm with a pointed end. The mouth is surrounded by wing-like cuticular expansions- cervical alae which are transversely striated suggesting it is Enterobius vermicularis. The eggs are colourless, non-bile stained, and elongated with a

characteristic planoconvex shape-flattened on one side and convex on the other. It has an outer albuminous layer making the eggs stick to each other, to clothing and to other objects.

It presents as intense irritation and pruritus of the perianal and perineal area which is more at night when the worm crawls out of the anus to lay eggs. In females, the worm may crawl into the genital tract causing irritation, mucoid discharge and symptoms of chronic salpingitis.

The eggs are infrequently found in stools as they are deposited on the perianal and perineal skin. They are diagnosed using NIH swabs (National Institutes of Health). The treatment of choice is albendazole, mebendazole, or pyrantel pamoate.

Solution to Question 37:

The image showing filamentous, branching acid-fast rods is suggestive of Nocardia.

Nocardia appears as filamentous organisms with extensive hyphae-like branching. They are gram-positive, catalase-positive, and produce urease. They are weakly acid-fast because they stain with carbolfuchsin (routine acid-fast stain) and retain when decolorized with 1-4% sulfuric acid instead of the stronger acid-alcohol decolorant.

Acid-fast staining is based on the principle that tubercle bacilli resist decolorization with acids after staining with aniline dyes. Acid fastness is attributed to the variety of fatty acids, alcohols, and lipids in the tubercle bacilli. Acid-fast organisms usually contain mycolic acid, which is acid-fast in the free state. The steps to carry out acid-fast staining as modified by Ziehl and Neelsen are:

• The slide is stained with carbolfuchsin along with the application of heat.

• It is then decolorized with 20% sulphuric acid. Later methylene blue is used for counterstaining.

• The acid-fast bacteria retain the fuchsin color, while other organisms take up the counterstain.

Other options:

Option B: The image given below is a smear with Ziehl-Neelsen staining showing non-branching and acid-fast Mycobacterium tuberculosis bacilli.



Option C- Mycobacterium leprae bacilli are less acid and alcohol fast than the tubercle bacillus; therefore, alcohol is removed from the hydrating and dehydrating steps and 5% sulphuric acid is used as a decolorizer in place of acid/alcohol solution. The image below shows Mycobacterium leprae on ZN staining.



Option D: Actinomyces are gram-positive, anaerobic, and filamentous bacteria that are non-acid-fast in contrast to Nocardia.

The following image shows the filamentous gram-positive stained actinomyces:



Solution to Question 38:

The incorrectly matched pair is Non-erosive arthritis - Patients of SLE.

Polyarthropathy occurs in ~50% of adults infected with parvovirus B19 who are healthy without underlying SLE. Although parvovirus B19 sometimes mimics rheumatoid arthritis with detectable rheumatoid factor, it has not shown any significant association with SLE.

Parvovirus B19 infection occurs usually in children 5–10 years causing erythema infectiosum. It is also known as fifth disease. It begins as a prominent erythema of the cheeks- slapped cheek disease, spreading to the trunk and limbs, followed by lymphadenopathy and arthralgia. Fever and mild constitutional symptoms may accompany the rash.

Parvovirus B19 replicates in immature erythroid cells and causes a transient aplastic crisis in cases of chronic hemolytic anemia, such as sickle cell disease, thalassemias, and acquired hemolytic anemias in adults.

Similarly, infection during pregnancy targets fetal red blood cells causing hemolysis leading to hydrops fetalis and/or fetal loss. Since the hydrops is not caused by antigen-antibody incompatibility, it is termed as non-immune hydrops.

It can also cause pure red cell aplasia in immunocompromised patients manifesting as chronic anemia and bone marrow suppression.

The following is an image of erythema infectiosum affecting a child:

Solution to Question 39:

The possible diagnosis based on the given clinical scenario is Jarisch-Herxheimer reaction.

JHR is a condition that occurs in patients infected by spirochetes who undergo antibiotic treatment. The reaction occurs within 24 hours of antibiotic treatment. Clinical features are fever, chills, nausea and vomiting, tachycardia, hypotension, flushing, myalgia, and exacerbation of skin lesions. JHR is due to IL6, TNF-alpha, and IL 8.

Syphilis is the leading spirochetal infection associated with JHR. Other infections are Lyme disease, louse-borne relapsing fever, leptospirosis, and tick-borne relapsing fever. Antibiotics that have been most associated are penicillins, tetracyclines, and erythromycin.

Mild reactions are self-limiting and often resolve spontaneously within 24 hours. Prevention of the JHR is by giving TNF-alpha antibodies and steroids.

Other options:

Option A: All of the strains of leptospira are found to be sensitive to ampicillin, cefotaxime, ciprofloxacin, doxycycline, erythromycin, and streptomycin. Resistance is shown to amphotericin B, 5-fluorouracil, trimethoprim, sulfamethoxazole, and vancomycin.

Option B: Penicillin belongs to the beta-lactam group of antibiotics. Hypersensitivity is the most important adverse reaction, resulting in nausea, vomiting, pruritus, wheezing, laryngeal edema, and, ultimately, cardiovascular collapse.

Option C: Dengue fever presents with headache, musculoskeletal pain, and occasional biphasic fever.

Solution to Question 40:

The given image shows multiple vesicles on hand and multiple eruptions along lymphatics characteristic of sporotrichoid lymphcutaneous infection. All of the organisms mentioned in the options can cause this except Corynebacterium minutissimum.

Corynebacterium minutissimum causes a localized infection of the stratum corneum called erythrasma. It presents as superficial skin lesions in the axilla and groin. This is a lipophilic corynebacterium and can be grown readily in media containing 20% fetal calf serum.

Sporotrichoid lymphocutaneous infection classically presents as superficial cutaneous lesions following cutaneous inoculation or trauma and progresses along dermal and subcutaneous lymphatics. Also known as nodular lymphangitis, the most common causes include:

• Sporothrix schenckii

• Nocardia brasiliensis

• Mycobacterium marinum

• Leishmania brasiliensis

Option A: Sporothrix schenckii is a dimorphic fungus. It causes sporotrichosis, a chronic granulomatous disease, by traumatic inoculation into the skin. This is usually followed by spread along the draining lymphatics and lymph nodes. Microscopy shows asteroid bodies. It can be seen as an oval, basophilic structure with eosinophilic radiation. Itraconazole is the drug of choice for treatment.

The image below shows asteroid bodies.



Option C: Nocardia brasiliensis is a gram-positive aerobic bacteria frequently found in the soil and infection may be exogenous. Cutaneous infections can present as local abscesses, cellulitis, or lymphocutaneous lesions. It causes mycetoma which is a localized chronic, granulomatous condition involving subcutaneous and deeper tissues, commonly affecting the foot. It presents as a tumor with multiple discharging sinuses.

Option D: Mycobacterium marinum is a non-tuberculous mycobacterium that causes swimming pool granuloma. The lesion begins as a papule which breaks down to form an indolent ulcer. It usually occurs after abrasions and is seen on the prominences like elbows, knees, ankles, or toes.

Solution to Question 41:

The organism with oocysts of size 8-10 µm is Cyclospora.

Cyclospora is a coccidian parasite.The sporulated oocysts are infectious to humans. They reside in the enterocytes in the small intestine. The infection causes prolonged diarrhea, abdominal pain, and vomiting, which is more severe in immunocompromised individuals.

On a wet mount of the feces, oocysts are seen as red spheres when stained with acid-fast stain. Sporulated oocysts of Cyclospora are 8-10 µm in size, round, and contain 2 sporocysts, each having two sporozoites. Treatment is by giving cotrimoxazole twice daily for seven days.

Other options:

Option A: Oocysts of Isospora are 23-36 µm, oval, and contain two sporocysts, each having four sporozoites. They are uniformly acid-fast.

Option B: Sporulated oocysts of Cryptosporidium are spherical, 4-5 µm in size, and acid-fast. It is implicated in causing traveler's diarrhea and chronic persistent diarrhea in immunocompromised patients. The two cryptosporidium species responsible are Cryptosporidium hominis and Cryptosporidium parvum.

Option D: Microspora oocysts are 2-4 µm in size, oval to cylindrical in shape with a polar filament. Spores are gram-positive and acid-fast. They cause a wide range of opportunistic

infections in immunocompromised patients.

Solution to Question 42:

L1 protein is used in the formulation of HPV vaccines.

Viral proteins can be divided into early and late proteins.

• Early proteins are called non-structural proteins as they do not form any structural component of the virus. They play a role within the infected cell during virus replication or act in the regulation of virus replication or assembly.

• Late proteins are also called structural proteins as they form viral components like capsids and envelopes.

Human papillomavirus (HPV) is a DNA virus that is spread by close contact. Serotypes 6 and 11 cause anogenital warts and laryngeal papilloma. The risk of these lesions can be reduced by vaccination against HPV. Three vaccines approved by FDA include bivalent (Cervarix-types 16 and 18), quadrivalent (Gardasil- 6, 11,16, 18), and 9-valent (Gardasil 9- additional 31, 33, 45, 52, and 58). All three provide protection against infection and the development of precancerous lesions.

The vaccine is prepared by using the L1 capsid protein. It is given to girls aged 11 to 12 years (can start at 9 years of age) and can also be given to boys.

E6 and E7 are oncogenic proteins that interact with p53 and Rb genes respectively. E1, E2, and E3 are other early proteins which help in viral replication and are not used to formulate vaccines.

Solution to Question 43:

The Epstein-Barr virus (EBV) binds to CD21 and acts as a B-cell mitogen.

It affects adolescents and younger adults more frequently but may occur from childhood to old age. It is transmitted primarily through close contact with oropharyngeal secretions. The incubation period of EBV infectious mononucleosis is 4-6 weeks.

Pathogenesis of Epstein Barr virus infection:

• The virus infects nasopharyngeal epithelial cells, salivary tissues, and lymphoid tissues.

• Lymphocytosis is caused by a proliferation of these EBV-infected B lymphocytes.

• There is a production of atypical reactive T cells which are CD8+ cytotoxic T cells, that are also called Downey cells.

• Heterophile antibodies are also produced due to B cell mitogenesis.

Clinically, patients may present with

• Fever, malaise, nausea, and anorexia.

• Lymphadenopathy- bilateral and symmetrical. Bilateral posterior cervical lymphadenopathy is characteristic.

• Pharyngitis.

• Tonsillar enlargement.

• Hepatomegaly and splenomegaly. Splenic rupture may occur even with minor trauma and is highly fatal.

• A drug-induced rash may develop on using beta-lactam antibiotics.

Solution to Question 44:

The most likely diagnosis based on the clinical vignette and the histopathology showing sclerotic bodies is chromoblastomycosis. The true statement regarding the condition is that it is caused by a dematiaceous fungus, which means that this fungus produces a brown pigment.

The various dimorphic pigmented fungi are Phialophora verrucosa, Fonsecaea pedrosoi, Fonsecaea compacta, and Cladophialophora carrionii.

Chromoblastomycosis, the most common chromomycosis, is more commonly seen in tropical areas among agricultural workers and woodcutters who walk barefoot. Traumatic implantation is the mode of infection.

Patients present with warty cauliflower-like cutaneous nodules on the feet and lower legs. The disease generally remains localized with slow peripheral extension. Lymphatic spread can occur.

Diagnosis is made by the demonstration of sclerotic bodies, also called copper-penny bodies in tissue sections. Sclerotic bodies appear as round or irregular, dark brown bodies with septae.



First-line management includes itraconazole or terbinafine. Flucytosine used on its own or combined with amphotericin B may also be effective. For small lesions, surgical excision is the therapy of choice.

Solution to Question 45:

Viruses belonging to the filoviridae family, such as Ebola and Marburg, resemble the morphology shown in image 3.

Image 1- Influenza virus - Influenza viruses have a linear, segmented, single-stranded RNA.



Image 2- Adenovirus contains double-stranded DNA. They have an icosahedral symmetry with structures known as fibers extruding from the vertices.





Image 3 - Ebola viruses are filamentous viruses with nonsegmented single-stranded RNA as genetic material.



Image 4 - Corona virus belonging to the family are shaped viruses with nonsegmented single-stranded RNA. Corona virus has petal/club-shaped peplomers radiating like sunrays.




Spike
Nucleocapsid
Envelope protein E
Membrane protein M

Microbiology NEET 2022

Question 1:

A photographer who recently returned from Africa presents to the emergency department with abdominal pain, hepatomegaly, and hemorrhagic manifestations. He died despite treatment, and an autopsy revealed intranuclear Torres bodies in the liver. Which of the following vaccines could have prevented it?

a) 17D

b) Nakayama vaccine

c) Weigl's vaccine

d) Jeryl Lynn strain

Question 2:

A militant presents with rashes all over his body sparing the palms and soles. On examination, he was febrile and lice were noted. Which of the following is responsible for his condition?

a) Rickettsia typhi

b) Rickettsia prowazekii

c) Rickettsia akari

d) Rickettsia conorii

Question 3:

Microfilariae with sheathed tail and two nuclei at the tail tip is suggestive of?

a) Wuchereria bancrofti

b) Brugia malayi

c) Loa loa

d) Onchocerca volvulus

Question 4:

A patient presents with cervical lymphadenopathy and is found to be retropositive. A fungal culture depicts a velvety growth with red diffusible pigment on the underside, as shown below. Which is the most likely causative organism?



a) Talaromyces marneffei

b) Blastomyces

c) Aspergillus

d) Pneumocystis jirovecii

Question 5:

A patient presents with itching in the axilla. On microscopic examination of the skin scrapings, a red pigment-producing fungus with pencil-shaped macroconidia is seen. Which of the following is the most likely organism?



a) Trichophyton violaceum

b) Trichophyton rubrum

c) Trichophyton tonsurans

d) Trichophyton schoenleinii

Question 6:

A female patient presents with complaints of thick white vaginal secretions. Which of the following can be used to identify the likely species of the causative agent?

a) Bird seed agar

b) Brain heart infusion agar

c) CHROMagar

d) Sabouraud dextrose agar

Question 7:

A 12-year-old boy presents with right upper quadrant pain, calf pain, conjunctival suffusion, icterus, and fever. Examination shows tender hepatomegaly. What is the most likely diagnosis?

a) Chikungunya

b) Leptospirosis

c) Dengue hemorrhagic fever

d) Encephalopathy caused by hepatitis A

Question 8:

A newly joined teacher develops rice water stools. The causative agent acts on which of the following receptors?

a) GM1 ganglioside receptor

b) GM2 ganglioside receptor

c) Sphingomyelin

d) Cerebroganglioside

Question 9:

Which of the following is considered to be the best for the detection of Clostridium difficile?

a) Aerobic culture of stool

b) Glutamate dehydrogenase and toxin assay

c) Glutamate dehydrogenase assay

d) None of the above

Question 10:

A diabetic truck driver presents to you with cough and breathlessness. A diagnosis of pneumonia was confirmed, and histopathological examination revealed dichotomous branching, as shown below. What is the likely causative agent?



a) Rhizopus

b) Mucor

c) Candida

d) Aspergillus

Question 11:

A woman with recurrent diarrhea is prescribed a broad-spectrum antibiotic. Which of the following is not true regarding Clostridium difficile infection?

a) Oral fidaxomicin is used for treatment

b) It is toxin mediated

c) IgM assay is used to confirm the diagnosis

d) Pseudomembrane consists of leukocytes, fibrin, and cellular debris

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	a

5	b
6	c
7	b
8	a
9	b
10	d
11	c

Detailed Explanations

Solution to Question 1:

The above clinical vignette is suggestive of yellow fever and the use of 17D vaccine could have prevented this.

17D vaccine is a live-attenuated vaccine administered through the subcutaneous route. It is contraindicated in infants less than 9 months of age, in pregnancy, and in individuals with egg allergy.

Yellow fever is caused by flavivirus and is endemic to certain parts of Africa and America. It occurs in two patterns: the urban cycle and the sylvatic cycle. Aedes aegypti is the vector in the urban cycle, while Aedes africanus is responsible for transmission in the sylvatic cycle.

Patients present with acute onset fever, headache, and vomiting. Jaundice and hemorrhagic manifestations can also be present. Ultimately, hepatic or renal failure can result in death.

Torres bodies are intranuclear eosinophilic inclusion bodies in hepatocytes. These are rarely seen in yellow fever. To prevent the establishment of the disease in India, strict vaccination and quarantine protocols for travelers from endemic areas are enforced.

Other options:

Option B: Nakayama vaccine is used for immunization against Japanese encephalitis. Culex tritaeniorhynchus is the principal vector. There is a sudden onset of fever and vomiting, which later progresses to encephalitis. In India, it was first reported in Vellore and continues to be active in the country.

Option C: Weigl's vaccine is the earliest used vaccine against epidemic typhus caused by Rickettsia prowazekii. The main foci of the disease are Europe, Africa, South America, and Asia. Kashmir is the endemic spot in India. Pediculus humanus corporis (human body louse) is the vector. Symptoms include fever and chills. A characteristic rash on the trunk and limbs, sparing the face, palms, and soles, may also be seen.

Option D: Jeryl-Lynn strain of the mumps virus is used against the disease as a live-attenuated vaccine. It is contraindicated in pregnancy and individuals allergic to eggs. Mumps present as parotid swelling associated with fever and tenderness.

Solution to Question 2:

The given clinical scenario is suggestive of epidemic or louse-borne typhus caused by Rickettsia prowazekii.

Pediculus corporis (human body louse) is the vector for the parasite, although the head louse may also transmit the disease. After the incubation period of 5 - 15 days, the disease typically starts with fever and chills. By around 4 - 5th day of fever, a characteristic rash appears, which starts on the trunk and spreads over the limbs but spares the face, palms, and soles.

The infection can be transmitted by

• Contaminated louse feces rubbed through abrasions caused by scratching

• Inhalation of aerosols of dried louse feces or directly through the conjunctiva

The disease is often severe with the fever lasting for over 2 weeks and is more fatal in patients older than 40 years of age.

Other options:

Option A: R. typhi causes murine (flea-borne) or endemic typhus. Rats are the reservoirs of infection and the parasite is transmitted by Xenopsylla cheopis (flea). The infection is transmitted when the flea feces is rubbed into the bite wound. The clinical features are similar to that of epidemic typhus but are milder and rarely fatal. Patients can present with fever, headache, myalgia, and rash.

Option C: R. akari causes rickettsial pox, which is a mild disease resembling varicella. The disease is transmitted by mites and the patients may present with fever, headache, and a vesicular rash.

Option D: R. conorii causes the Mediterranean spotted fever. The parasite is transmitted by ticks and the patients present with fever, headache, rash, and eschar (tache noire).

Solution to Question 3:

The most likely organism based on the description of the microfilariae is Brugia malayi.

The microfilariae of B. malayi are sheathed and have kinks and secondary curves. The tail tip has two nuclei - a terminal and a subterminal one. It has nocturnal periodicity.

The female filarial worms, which belong to the phylum Nematoda, give birth to larvae known as microfilariae. Some of the microfilariae retain their egg membranes and are known as sheathed microfilariae. They can be further differentiated on the basis of the arrangement of nuclei in the tail.

Classical filariasis is due to obstruction of lymphatics by the adult worms resulting in lymph stasis. This leads to leakage of lymph into the tissues causing hard pitting (brawny) edema. It can present as acute adenolymphangitis, hydrocele, or elephantiasis which is shown below.

Occult filariasis (Meyers-Kouwenaar syndrome) results from a hypersensitivity reaction to the microfilarial antigens. Here, the microfilariae are not found in the blood. Symptoms include hepatosplenomegaly, nocturnal cough, or arthritis. Tropical pulmonary eosinophilia is a manifestation of occult filariasis which presents as low-grade fever, weight loss, and dry nocturnal cough. There is marked eosinophilia and raised serum IgE levels. It is managed with diethylcarbamazine.

Other options:

Option A - Microfilariae of Wuchereria bancrofti are sheathed and the tail tip lacks nuclei. It has nocturnal periodicity.

Option C - Loa loa causes subcutaneous filariasis which presents as swellings on the extremities (Calabar/fugitive swellings). It can also cause conjunctival granulomas and proptosis. The nuclei are arranged in a row up to the tail tip of the sheathed microfilariae. It exhibits diurnal periodicity.

Option D - Onchocerca volvulus (blinding filaria) causes onchocerciasis. The microfilariae are unsheathed and the nuclei do not extend to the tail tip. Patients present with subcutaneous nodules (onchocercoma), dermatitis, photophobia, or even blindness. It is non-periodic.

Solution to Question 4:

The most likely causative organism based on the history and image given is Talaromyces marneffei (Penicillium marneffei).

T. marneffei is a dimorphic fungus that causes talaromycosis (penicilliosis). It is commonly seen in individuals with advanced AIDS. It is diagnosed by blood culture, which shows a mold with a powdery surface and characteristic red pigment that diffuses into the agar. Histopathological examination reveals yeast-like organisms with central septation.

Clinical features include fever, weight loss, lymphadenopathy, and hepatosplenomegaly. Umbilicated papules that resemble molluscum contagiosum may also be seen.

Mild and moderate infections can be treated with itraconazole. Amphotericin B is recommended for severe infections.

Other options:

Option B - Blastomyces are thermally dimorphic fungi and cause blastomycosis. Primary infection usually affects the lungs, which can later disseminate through the blood to form multiple abscesses at sites such as skin and bone. Culture at 25–30 degrees Celsius promotes hyphal growth. Microscopic examination of stained specimens shows broad-based budding yeast cells.

Option C - Aspergillus colonies have a velvety powdery surface, of which the color varies according to the species. Aspergillus fumigatus is seen as dark green colonies, while Aspergillus niger appears black colored. Yellow-green growth is seen in the case of Aspergillus flavus. On microscopy, dichotomous branching at acute angles seen after staining is characteristic. It can cause invasive aspergillosis in immunocompromised patients.

The image below shows Aspergillus niger colonies.

Option D - Pneumocystis jirovecii cannot be cultured. On being stained by methenamine silver, the cyst wall appears black under the microscope. Trophozoites can also be seen after staining. Clinical features include dyspnea, fever, and cough. It can cause lethal pneumonia in immunocompromised individuals. Diffuse bilateral perihilar interstitial infiltrates can be seen on chest radiograph.



Solution to Question 5:

The most likely organism based on the given history and image is Trichophyton rubrum. The colonies appear velvety and a deep red, non-diffusible pigment is seen when viewed from the reverse side of the colony. A few long and pencil-shaped macroconidia can be seen.

Trichophyton is a dermatophytic fungi that can infect hair, skin, or nails. Dermatophytes infect only the keratinized tissue and have keratinases that can digest keratin. There are 3 genera of

dermatophytes:

• Trichophyton - The colonies may appear powdery, waxy, or velvety with pigmentation characteristics of different species.

• Epidermophyton - It infects the skin and nails, but not the hair. The colonies appear powdery and greenish-yellow.

• Microsporum - It infects the hair and skin, but not the nails. The colonies may appear cotton-like, velvety, or powdery, with white to brown pigmentation.

T. rubrum causes tinea corporis. It results from the invasion and proliferation of the causal fungi in the stratum corneum, with lesions on the trunk and limbs. The lesions are usually well-defined and circular with a raised edge. It can also cause tinea pedis, tinea cruris, and onychomycosis.



Other options:

Option A: The colonies of T. violaceus are very slow-growing, waxy with a violet/purple pigment. They have distorted hyphae, with conidia seen rarely.



Option C: T. tonsurans has a cream or yellowish colony with central furrows. Microconidia are abundantly seen. Thick-walled, irregular macroconidia are also seen.



Option D: Colonies of T. schoenleinii are smooth, waxy and brownish. Hyphal swellings are seen along with chlamydospores and favic chandelier.

Solution to Question 6:

The clinical scenario points to a diagnosis of candidiasis. CHROMagar can be used for the rapid identification of Candida species. It is based on the fungal enzymatic action on various chromogenic substrates in the medium.

After incubation on CHROMagar for 1–4 days, colonies are seen: Candida albicans colonies appear green, as shown below; Candida glabrata colonies appear dark purple; and Candida parapsilosis colonies have a pinkish hue.

The most common species implicated in candidal infections are C. albicans, C. parapsilosis, and C. glabrata. Vaginal mucosal invasion by the organism causes vulvovaginitis, marked by pruritus and vaginal discharge. A white pseudomembranous patchy lesion called thrush can occur on the tongue, lips, or gums.



Immunocompromised individuals are at an increased risk of systemic candidiasis, which includes candidal endocarditis and urinary tract infections.

It appears as budding gram-positive cells on wet films and smears. Mycelial forms suggest colonization. Candida albicans, the most commonly implicated pathogenic species form chlamydospores on corn meal agar.

The image below shows candidal hyphae and chlamydospores.



Mucocutaneous manifestations are managed with topical nystatin or oral ketoconazole.

Other options:

Options A and B: Bird seed agar and brain heart infusion agar are non-specific media used in fungal culture.

Option D: Sabouraud dextrose agar is the traditional mycological medium, which contains glucose and modified peptones. It supports the growth of fungi, while restricting bacterial growth due to the addition of antibacterials.

Solution to Question 7:

The most likely diagnosis based on the clinical vignette is leptospirosis.

Leptospirosis is a zoonosis caused by spirochetes of the genus Leptospira. Patients with mild disease may present with flu-like symptoms of sudden onset, fever with chills, headache, abdominal pain, conjunctival suffusion, and myalgia. The muscle pain usually affects the calves, abdomen, and back. Mild cases can be treated with oral doxycycline, ampicillin, or amoxicillin.

Severe leptospirosis is rapidly progressive and is associated with high fatality. The classic presentation of the severe disease called Weil's syndrome includes the triad of hemorrhage, jaundice, and acute kidney injury. Jaundice when present can be profound and physical examination may reveal an enlarged, tender liver. Temperature-pulse dissociation /relative bradycardia can be seen. Moderate to severe cases are treated with intravenous penicillin, ampicillin, or ceftriaxone.

The image below shows icterus and conjunctival suffusion in a patient with leptospirosis.



Leptospira are actively motile organisms and are best seen using a dark-field microscope. They are thin, tightly coiled, and flexible spirochetes with one end forming a hook. Diagnosis is by microscopic agglutination test.

Transmission occurs through cuts or breaks in the skin and mucous membranes (conjunctival and oral mucosa). Several outbreaks have been reported that resulted from exposure to floodwaters contaminated by urine from infected animals. The organism enters the bloodstream and lodges in the parenchymatous organs (liver and kidneys) leading to hemorrhage and necrosis of the tissue causing dysfunction of the organs.

The lifecycle of Leptospira is shown below.



Illness is classically biphasic after an incubation period of up to 1 month. The stages are:

• Acute leptospiremic phase- the organism can be cultured from blood.

• Resolving immune phase- the organism can be cultured from urine.

The stages may not be clinically distinguishable.

Other options:

Option A - Chikungunya is a viral disease that presents with saddle-back fever, conjunctival suffusion, and photophobia. Migratory arthralgia, especially involving small joints may also be seen. A maculopapular or petechial rash is frequently seen on the trunk.

Option C - Dengue hemorrhagic fever is a form of severe dengue with bleeding manifestations. Evidence of plasma leakage and thrombocytopenia is also generally seen.

Option D - Encephalitis due to Hepatitis A is seen in fulminant hepatitis, a rare complication. Patients present with confusion, disorientation, ascites, and edema.

Solution to Question 8:

The given clinical scenario of rice-water stools is suggestive of cholera, caused by Vibrio cholerae. The toxin produced by the organism acts on GM1 ganglioside receptors.

Vibrio cholerae is a comma-shaped, gram-negative bacillus with darting motility. It has a single polar flagellum. It causes cholera, an acute diarrheal disease. It is marked by painless, profuse, watery diarrhea with flecks of mucus (rice water stools) associated with effortless vomiting. Pulmonary edema and arrhythmias are some of the complications.

Infection is acquired by ingestion of contaminated food and water. The organisms produce an enterotoxin known as cholera toxin (CT). It has 1 A and 5 B subunits. The B subunit (binding units) fixes to the GM1 ganglioside receptors on jejunal epithelial cells. The A subunit dissociates into the A2 fragment, which attaches the A1 fragment to the B subunit. This results in prolonged accumulation of cAMP, which causes an outpouring of large amounts of water and electrolytes

into the intestinal lumen. Ultimately, it results in profuse watery diarrhea.

In addition to the cholera toxin, additional toxins such as zona occludens toxin and accessory cholera toxin are also produced. The former alters intestinal permeability by its action on intestinal epithelial cells.

The characteristic darting mobility and its hindrance by antiserum can be used for rapid microscopic diagnosis.

Fluid resuscitation by rice-based ORS is recommended. Medical management includes erythromycin or azithromycin.

Solution to Question 9:

A combination of glutamate dehydrogenase and toxin assay can be used for the detection of Clostridium difficile.

Clostridium difficile is a gram-positive anaerobic bacillus that causes Clostridioides (formerly Clostridium) difficile infection after ingestion of spores of the organism. The more severe form of the condition is known as pseudomembranous colitis (PMC). Enzyme immunoassay to detect glutamate dehydrogenase is less specific compared to toxin detection for diagnosis. So a combination of the two can be used for diagnostic confirmation.

The condition is most commonly associated with prolonged antimicrobial use, which interferes with the commensal colonic microflora. The most commonly implicated antibiotics are cephalosporins, ampicillin, and clindamycin. The spores are spread by contact with the feces of affected individuals or asymptomatic carriers. It can persist on surfaces for months. Risk factors include poor hand hygiene, old age, and gastrointestinal surgery.

The spores germinate in the small intestine and produce two toxins—toxin A (enterotoxin) and toxin B (cytotoxin). These interfere with the epithelial-cell barrier function and result in diarrhea and pseudomembrane formation. The pseudomembranes appear as small whitish-yellow plaques initially which later coalesce to involve the entire colon wall. It contains necrotic leukocytes, fibrin, and cellular debris. The anamnestic IgG response determines which patients develop the disease or remain asymptomatic.

Symptoms include diarrhea, fever, and abdominal pain. The diagnosis is based on a combination of clinical criteria and detection of toxin or the organism in stool samples. This can be done by nucleic acid amplification testing (NAAT) methods such as polymerase chain reaction (PCR) or by culture.

Discontinuation of any ongoing antimicrobials and adequate hydration are general measures that are recommended. Oral fidaxomicin or vancomycin is the first-line treatment. Oral metronidazole can be used when these are not available.

Other options:

Option A: Aerobic culture of stool is the most sensitive test and is highly specific. It is not used practically as the turnaround time is slow.

Solution to Question 10:

The most likely causative agent based on the clinical features and image showing dichotomous branching at acute angles given is Aspergillus.

Aspergillus fumigatus is the most common pathogenic species. It can cause invasive aspergillosis in immunocompromised patients. This can be due to prolonged treatment with steroids or retro-positive status. It starts as pneumonia, which can later spread to involve other organs. Voriconazole is the drug of choice.

Other manifestations include allergic bronchopulmonary aspergillosis where central bronchiectasis is usually seen. Aspergilloma and superficial infections like otomycosis or mycotic keratitis are the other presentations.

Histopathological examination of the tissue section shows characteristic dichotomous branching at an acute angle. The hyphae stain poorly with hematoxylin and eosin, so periodic acid–Schiff (PAS) stain is used.

Other methods for diagnosis include detection of galactomannan (Aspergillus cell-wall component) and culture isolation. Aspergillus fumigatus grows dark green colonies, while Aspergillus niger appears black. Yellow-green colonies are seen in case of Aspergillus flavus. Microscopic examination of colonies stained with lactophenol cotton blue would show septate hyphae and conidiophores.

Other options:

Options A and B - Rhizopus and Mucor are zygomycetes that are responsible for mucormycosis. Non-septate, broad hyphae are seen on tissue samples stained with hematoxylin and eosin.

The image below shows broad aseptate and irregularly-branched non-septate hyphae of Mucor.



Option C - Candida, the causative agent for candidiasis, appear as budding gram-positive cells on wet films and smears. Mycelial forms suggest colonization. Culture colonies are creamy-white and have a yeasty odor. Candida albicans, the most commonly implicated pathogenic species, form chlamydospores on corn meal agar.

The image below shows candidal hyphae and chlamydospores.



Solution to Question 11:

IgM assay cannot be used to confirm a diagnosis of Clostridium difficile infection. The diagnosis is based on a combination of clinical criteria and detection of toxin or the organism in stool samples.

C. difficile, a gram-positive anaerobic bacillus, causes Clostridioides (formerly Clostridium) difficile infection (CDI) after ingestion of spores of the organism. The more severe form of the condition is known as pseudomembranous colitis (PMC). It is most commonly associated with prolonged antimicrobial use which interferes with the commensal colonic microflora. The most commonly implicated antibiotics are cephalosporins, ampicillin, and clindamycin.

The spores are spread by contact with the feces of affected individuals or asymptomatic carriers. It can persist on surfaces for months. Risk factors include poor hand hygiene, old age, and gastrointestinal surgery.

The spores germinate in the small intestine and produce two toxins - toxin A (enterotoxin) and toxin B (cytotoxin). These interfere with the epithelial-cell barrier function and result in diarrhea and pseudomembrane formation. The pseudomembranes appear as small whitish-yellow plaques initially which later coalesce to involve the entire colon wall. It contains necrotic leukocytes, fibrin, and cellular debris. The anamnestic IgG response determines which patients develop the disease or remain asymptomatic.

Symptoms include diarrhea, fever, and abdominal pain. The diagnosis is based on a combination of clinical criteria and detection of toxin or the organism in stool samples. This can be done by nucleic acid amplification testing (NAAT) methods such as polymerase chain reaction (PCR) or by culture. Some cases of PMC can be visualized through endoscopy.

Discontinuation of any ongoing antimicrobials and hydration are general measures that are recommended. Oral fidaxomicin or vancomycin is the first-line treatment. Oral metronidazole can be used when these are not available.

Microbiology NEET 2023

Question 1:

A patient hailing from Delhi presents with fever, arthralgia, and extensive petechial rash for 3 days. Lab investigations revealed a hemoglobin of 9 g/dL, a white blood cell count of 9000 cells/mm3, a platelet count of 20000 cells/mm3, and a prolonged bleeding time. The clotting time was normal. What is the most likely diagnosis?

a) Dengue

b) Malaria

c) Scrub typhus

d) Typhoid

Question 2:

Which of the following statements is true about Trichomonas vaginalis?

a) It cannot be cultured

b) Twitching motility is seen on wet saline mount

c) Cysts are seen on wet saline mount of vaginal secretions

d) It is not a sexually transmitted infection

Question 3:

A lady from West Rajasthan presented with an ulcer surrounded by erythema on the right leg. Microscopy of the biopsy from the edge of the ulcer showed organisms with dark staining nuclei and kinetoplast. What is the most likely causative agent?

a) Leishmania tropica

b) Babesia

c) Trypanosoma

d) Histoplasma

Question 4:

A child presents with a fever and a rash. Urine examination showed cells with owl's eye appearance. What is the most likely diagnosis?

a) Cytomegalovirus

b) Epstein-Barr virus

c) Herpes simplex virus

d) Toxoplasma gondii

Question 5:

Infection with Clonorchis sinensis is associated with an increased risk of _____.

a) Cervical cancer

b) Cholangiocarcinoma

c) Gastric carcinoma

d) Bladder carcinoma

Question 6:

A patient presented with complaints of persistent cough and weight loss. He had a history of contact with multiple sexual partners. He was diagnosed as HIV positive 6 months back and is poorly adherent to his medications. The fungal culture of sputum was negative. On Gomori methamine silver staining, the cells showed darkly stained crescent-shaped cysts. What is the most likely causative organism?

a) Paracoccidiodes brasiliensis

b) Histoplasma capsulatum

c) Coccidiodes immitis

d) Pneumocystis jerovecii

Question 7:

A sewage worker presented with abdominal pain, jaundice, conjunctival suffusion, and blood in the urine for the past 5 days. Which of the following is the investigation of choice?

a) Widal test

b) Microscopic agglutination test

c) Weil-Felix reaction

d) Paul-Bunnell test

Question 8:

Which of the following laboratory test is used to diagnose rotavirus diarrhea?

a) Antigen detection in the stool sample

b) Antigen detection in blood

c) Antibody detection in serum

d) Light microscopy of stool specimen

Question 9:

A boy presented with a fever and chills. Rapid test was positive for specific antigen HRP-2. Which of the following species of Plasmodium is the most likely causative agent?

a) Plasmodium falciparum

b) Plasmodium malariae

c) Plasmodium vivax

d) Plasmodium ovale

Question 10:

A child was brought with complaints of high fever, multiple seizures, headache, and neck rigidity. CSF analysis showed low glucose, high protein, and many polymorphs. The CSF culture showed pleomorphic gram-negative bacilli. What is true about the causative organism?

a) It is bacitracin sensitive.

b) Produces non-lactose fermenting colonies on MacConkey agar

c) Exhibits satellitism around Staphylococcus aureus colonies

d) It exhibits alpha hemolysis on sheep blood agar

Question 11:

A patient presented with a high fever, altered sensorium, headache, neck rigidity, and seizures. He was diagnosed with HSV encephalitis. Which of the following tests can be used

to confirm the diagnosis?

a) IgM in CSF

b) Tzanck smear of CSF

c) CSF PCR for viral DNA

d) CSF culture on chick embryo lines

Question 12:

In a patient presented with a fever and a positive filarial antigen test, what is the next appropriate method of management?

a) Bone marrow biopsy

b) DEC provocation test

c) Detection of microfilariae in the blood smear

d) Ultrasound of the scrotum

Question 13:

A man presents with a history of dysuria and urethral discharge after having unprotected sexual intercourse. Gram-staining of the purulent discharge is seen in the following image. What is the treatment of choice for this infection?



a) Erythromycin

b) Azithromycin

c) Ceftriaxone

d) Penicillin G

Answer Key

Question No.	Correct Option
1	a
2	b
3	a
4	a
5	b
6	d
7	b
8	a
9	a
10	c
11	c
12	c
13	c

Detailed Explanations

Solution to Question 1:

The given clinical vignette is suggestive of dengue fever.

Dengue is an acute febrile disease caused by the mosquito-borne dengue viruses (DENVs), consisting of four serotypes (DENV 1 to 4), that are members of the Flaviviridae family, genus Flavivirus.

Dengue virus is transmitted from person to person by Aedes aegypti mosquitoes.

After an incubation period of 3–15 days, classical dengue fever begins with an abrupt onset of high fever. The condition is self-limiting though debilitating with headache, retro-orbital pain, myalgia, arthralgia, petechiae rash, and leucopenia. A macular-papular recovery rash appears 3–4 days after the onset of fever, and it usually starts on the trunk before spreading peripherally. The rash in dengue appears as 'white islands in a red sea' due to erythema with petechiae.

Shown below is an image of the rash in dengue:

Solution to Question 2:

Trichomonas vaginalis has a jerky or twitching type of mobility on the wet saline mount. Option B is the correct answer.

T. vaginalis is the only protozoan with one stage in its life cycle, i.e., trophozoite form (no cyst stage is seen). It is an obligate parasite and cannot live without close association with vaginal, urethral, or prostatic tissues. Sexual transmission is the main route of infection. Other modes are fomites, and babies during passage through the infected birth canal.

The incubation period ranges from 4 days to 4 weeks.

The common presentation of trichomoniasis is foul-smelling frothy vaginal discharge, itching, dysuria, and dyspareunia. Petechial hemorrhage and mucosal capillary dilation give the appearance of a strawberry cervix. Chicken-like epithelium occurs due to intracellular edema

which is also a characteristic feature.

The gold standard for diagnosis is culture on cysteine-peptone-liver-maltose (CPLM) medium or plastic envelope medium (PEM). The treatment of choice is metronidazole 2 g orally as a single dose.

T. vaginalis infection in women may lead to infertility, preterm birth, and low-birth-weight infants. In men, trichomoniasis has been also associated with infertility through inflammatory damage to the genitourinary tract or interference with sperm function.

The below image shows the trophozoite form of Trichomonas vaginalis.



Solution to Question 3:

The above clinical vignette is suggestive of cutaneous leishmaniasis.

Leishmaniasis is a slowly progressing indigenous disease caused by a protozoan parasite of the genus Leishmania. It is transmitted by the bite of an infected female Phlebotomus sandfly which inoculates the flagellated promastigotes into the skin of the host in developing countries. In India, it is endemic to Bihar, West Bengal, Jharkhand, and Uttar Pradesh.

Causative organisms include:

• Old world leishmaniasis: Leishmania donovani, Leishmania major, and Leishmania tropica

• New world leishmaniasis: Leishmania mexicana complex and Leishmania braziliensis complex

The image given below shows a phlebotomus sandfly.

The image given below shows Chiclero ulcer/New World cutaneous leishmaniasis.

The image below shows Lutzomyia sandfly responsible for the New World cutaneous leishmanisis.

In human beings, the disease presents in four different forms with a broad range of clinical manifestations:

• Visceral leishmaniasis, or kala-azar,

• Cutaneous leishmaniasis,

• Mucocutaneous leishmaniasis, and

• Diffuse cutaneous leishmaniasis.

The life cycle of Leishmania involves two forms:

• The promastigote which develops and lives extracellularly in the sandfly vector and

• The amastigote multiplies intracellularly in the reticuloendothelial cells of the host.

When the infected sandfly bites a person, infectious promastigotes are released into the host dermis along with the sandfly saliva, which potentiates parasite infectivity. Amastigote form (as shown in the image below) is also known as Leishman-Donovan (LD) body. It is an intracellular form found inside macrophages, monocytes, and reticuloendothelial cells. It contains a kinetoplast, which consists of a parabasal body and a blepharoplast. Flagellum is absent.



Promastigote form (as shown in the below image) is present in the midgut of the female sandfly. The kinetoplast lies transversely near the anterior end. A single flagellum is seen. It is the infective form of L. donovani for humans.



Given below is the lifecycle of L. donovanii.



This condition causes fever with rigor and chills associated with splenomegaly and hepatomegaly. It lowers immunity, causes persistent fever, anemia, and liver and spleen enlargement, and if left untreated, death may occur due to opportunistic infection.

Diagnosis of kala-azar is mainly by demonstration of amastigotes within macrophages in tissue aspirates such as blood smears, bone marrow, splenic, or lymph node aspirate. It is the gold standard for diagnosis.

The image below shows amastigotes located intracellularly within macrophages.



Tests to detect leishmaniasis include:

• Rapid diagnostic test: rK39 Rapid diagnostic test and Kala-azar dipstick test strip. This is recommended by National Vector Borne Disease Control Programme (NVBDCP) in India.

• Parasitological diagnosis: Demonstration of LD bodies in buffy coat smear - confirmatory.

• Aldehyde test of Napier: Widely used for surveillance.

• Serological tests for diagnosis: Rapid dipstick test, Direct agglutination test, ELISA, indirect fluorescent antibody test

• Leishmanin (Montenegro) test: Induration of 5mm or more is considered positive, usually positive 4-6 weeks (delayed-hypersensitivity) after onset in cutaneous leishmaniasis, valuable for distinguishing immune from nonimmune subjects and endemicity.

Drugs used in the treatment of visceral leishmaniasis are

• Sodium stibogluconate

• Miltefosine

• Amphotericin B

In Bihar, there is resistance to antimony so amphotericin-B-deoxycholate or miltefosine is preferred.

Other options:

Option B: Babesia is a protozoan parasite that causes babesiosis, a tick-borne disease. Babesiosis manifests as a malaria-like illness with fever, chills, and hemolytic anemia.

Option C: Trypanosoma is a genus of protozoan parasites responsible for diseases like African trypanosomiasis (sleeping sickness) and American trypanosomiasis (Chagas disease). These diseases present with different clinical features, such as fever, swollen lymph nodes, and neurological symptoms in African trypanosomiasis, and fever, swelling at the infection site, and cardiomyopathy in Chagas disease.

Option D: Histoplasma is a dimorphic fungus that causes histoplasmosis, a systemic mycosis. It primarily affects the lungs and can disseminate to other organs in immunocompromised individuals. The clinical presentation of histoplasmosis is variable, but it does not typically present as a skin ulcer with erythema as described in the clinical scenario.

Solution to Question 4:

The most likely causative agent is cytomegalovirus (CMV) based on the clinical symptoms and the presence of the owl's eye appearance on urinary examination.

CMV is the largest virus in the herpes family of viruses. It causes the following conditions:

• Immunocompetent individuals - asymptomatic or mononucleosis-like infection such as fever and rash

• Neonates - devastating systemic infections

• Immunosuppressed individuals - serious, life-threatening, disseminated CMV infections of the lungs (pneumonitis) and gastrointestinal tract (colitis)

CMV spreads through contact with salivary secretions and sexual contact, blood transfusion, and organ transplantation.

Laboratory diagnosis is by recovery of the virus from urine, saliva, or other bodily fluids into human fibroblast cells. Antibody demonstration using ELISA is used for diagnosis of primary infection.

A cytomegalic pneumocyte containing characteristic intranuclear basophilic inclusion surrounded by a clear halo known as an owl's eye appearance.

Other options:

Option B: Epstein Barr virus: Conditions associated with EBV infection are

• Glandular fever (infectious mononucleosis): fever, pharyngitis, and posterior cervical lymphadenopathy

• Hodgkin lymphoma

• Non-Hodgkin lymphoma

• Burkitt's lymphoma

• Nasopharyngeal carcinoma

• EBV■associated lymphoproliferative disease

Option C: Herpes simplex virus causes cutaneous infections like herpes labialis caused by viral reactivation in febrile patients. It presents as blisters around the mouth, cheeks, and forehead. Gingivostomatitis and pharyngitis are the most frequent manifestations of primary herpes infections. HSV esophagitis presents as dysphagia, sub-sternal pain, and weight loss.

Option D: Acute congenital toxoplasmosis may manifest as fever, jaundice, petechial rashes, microphthalmia, cataract, glaucoma, lymphadenopathy, hepatosplenomegaly, myocarditis, cerebral calcifications, and chorioretinitis.

Most infections in adults who are immunocompetent are usually asymptomatic. Active progression of infection is more likely in immunocompromised individuals.



Solution to Question 5:

Clonorchis sinensis or Chinese liver fluke causes clonorchiasis. It is also associated with an increased risk of cholangiocarcinoma.

Clonorchis sinensis infection is endemic in south China, Japan, Korea, and Taiwan due to a long tradition of consuming raw freshwater fish or shellfish. Humans are infected by eating raw or undercooked fish containing encysted larvae (metacercariae).

The immature flukes after excystation in the duodenum enter the biliary ducts and differentiate into adults.

The hermaphroditic adults produce eggs that are excreted in the feces. Upon reaching freshwater the eggs are ingested by snails, which are the first intermediate hosts. The eggs hatch within the gut and differentiate first into larvae (rediae) and then into many free-swimming cercariae. Cercariae encyst to form metacercariae under the scales of certain freshwater fish (second intermediate host) which are then ingested by humans.

In humans, it infects the common bile duct and gallbladder. Liver imaging shows a dilated biliary tract. It is a risk factor for cholangiocarcinoma.

The life cycle of Clonorchis sinensis is given below:



Other options:

Option A: Cervical cancer - Human papilloma virus (HPV) infection, particularly with high-risk strains like HPV-16 and 18, is the main risk factor for cervical cancer. Clonorchis sinensis is not associated with cervical cancer.

Option C: Gastric carcinoma - Helicobacter pylori infection is a major risk factor for the development of gastric carcinoma. Clonorchis sinensis is not linked to gastric carcinoma.

Option D: Bladder carcinoma - The vesical blood fluke Schistosoma haematobium primarily infects the urinary bladder and increases the risk of squamous cell carcinoma of the bladder. Clonorchis sinensis is not associated with bladder carcinoma.

Solution to Question 6:

The given clinical vignette is suggestive of pneumonia caused by Pneumocystis jirovecii.

P. jirovecii pneumonia (PCP) is an opportunistic infection in immunocompromised patients. It usually occurs when the CD4 count falls below 200 cells/µL. Patients usually present with fever and nonproductive cough with progressive shortness of breath. It can also affect other organs like lymph nodes, spleen, and liver.

Laboratory and radiographic findings are also non-specific and show an increase in lactate dehydrogenase levels and symmetric, perihilar, and diffuse bilateral interstitial infiltrates, respectively. The classic radiographic appearance of symptomatic Pneumocystis pneumonia consists of diffuse bilateral symmetric interstitial infiltrates that are perihilar. CT will show diffuse ground-glass opacities. Giemsa stain, toluidine blue, methenamine silver, and calcofluor white stains demonstrate trophozoites. The cyst wall appears black with Gomori's methenamine silver stain.

Shown below is an image of the staining showing the black-stained crescent-shaped cysts resembling the crushed ping-pong balls of P. jirovecii. The stain used for staining the causative organism is Gomori's methenamine silver stain.



Trimethoprim-sulfamethoxazole combination (commonly known as co-trimoxazole) is the treatment of choice for PCP, and it is administered for 21 days, followed by a course for secondary prophylaxis. Concomitantly, glucocorticoids should also be administered to reduce pulmonary inflammation.

Patients who are susceptible to PCP are also given co-trimoxazole as prophylaxis. For patients who are unable to tolerate co-trimoxazole, IV pentamidine is the drug of choice.

Prophylaxis is indicated for other opportunistic infections like Mycobacterium avium complex (MAC) infection if CD4 counts fall below 50 cells/µL; Coccidioides, if CD count is <250/µL; and Cryptococcus, when the CD count is <100/µL.

Other options:

Option A: Paracoccidioides brasiliensis causes paracoccidioidomycosis, which usually presents with ulcerations in the oral and nasal mucosa, lymphadenopathy, and pulmonary lesions, but it does not form crescent-shaped cysts.

Option B: Histoplasma capsulatum causes histoplasmosis and may be associated with similar symptoms as Pneumocystis jirovecii, but it does not show crescent-shaped cysts on Gomori methenamine silver staining.

Option C: Coccidioides immitis causes coccidioidomycosis, which is characterized by respiratory symptoms, skin lesions, and joint pain. It also does not show crescent-shaped cysts on Gomori methenamine silver staining.

Solution to Question 7:

The given clinical scenario is suggestive of leptospirosis (Weil's disease). A microscopic agglutination test is the investigation of choice for the diagnosis of leptospirosis.

Leptospirosis is a zoonotic disease caused by Leptospira interrogans. Humans are infected when the leptospirae in water contaminated with the urine of carrier animals enter the body through cuts or abrasions on the skin or through the intact mucosa of the mouth, nose, or conjunctiva. Hence, a sewage worker is at an increased risk of it.

Clinical features include fever, jaundice, vomiting, headache, and intense injection of the eyes. Some may present as aseptic meningitis or with hepatorenal damage (Weil's disease). There are two clinical types of leptospirosis: icteric and non-icteric. It can also manifest as aseptic meningitis.

The life cycle of Leptospira is depicted below:



Solution to Question 8:

Antigen detection in the stool sample is the laboratory test used to detect rotavirus diarrhea.

Rotavirus is the most common identifiable viral cause of diarrhea in children below the age of 5 years and is most frequent between 6-24 months of age. The incubation period is 1-3 days. Most

cases are asymptomatic or with mild symptoms like watery diarrhea, vomiting, low-grade fever, and temporary lactose intolerance.

Diagnosis can be made by rapid antigen detection of rotavirus in stool specimens. Laboratory diagnosis depends on the demonstration of the virus in stool samples which is collected at an earlier course of illness. Antigen enzyme immunoassays (EIAs) and polymerase chain reaction (PCR) are used to detect the virus in the stool. The polymerase chain reaction is the most sensitive detection method for genotyping of rotavirus nucleic acids from stool specimens.

Treatment is supportive including the correction of fluid and electrolyte imbalance. The main goals of treatment are avoiding and treating dehydration. Maintaining nutrition is a secondary goal.

Solution to Question 9:

The given clinical scenario and the HRP-2 positivity are suggestive of P. falciparum malaria.

Malaria is a parasitic infection caused by the Plasmodium species transmitted by the Anopheles mosquito. It is characterized by many nonspecific symptoms, including but not limited to tachycardia, tachypnea, chills, malaise, fatigue, diaphoresis (sweating), and headache. The fever of malaria is characterized by periodicity depending on the infective Plasmodium species. The fever returns every other day for P. vivax, P. ovale, and P. falciparum, and every three days for P. malariae.

The newer tests include rapid antigen detection tests that employ immunochromatographic methods. The parasite-F test is one of them. This test is based on the detection of histidine-rich protein-2 (HRP-2) which is produced only by the asexual forms of Plasmodium falciparum that are present on the surface of RBCs. The test strip has monoclonal antibodies against the HRP-2 antigen. This test has very high sensitivity and specificity. The disadvantages of the test include that it can detect only Plasmodium falciparum HRP-2 antigen but not the other three species of Plasmodium. Also, the test remains positive until 2 weeks after the cure.

Another diagnostic test of significance is the dual antigen test. This test detects both the parasite lactate dehydrogenase of trophozoites and gametocytes of all the Plasmodium species and HRP-2 produced by P. falciparum simultaneously.

Solution to Question 10:

This clinical vignette is suggestive of meningitis caused by Haemophilus influenzae. H. influenzae is a gram-negative pleomorphic bacillus that is non-motile and non-spore-forming. The correct statement about this causative organism is that it exhibits satellitism around Staphylococcus colonies, i.e option C.

When present in the sputum, H. influenzae appears as a coccobacillus, but in cerebrospinal fluid (CSF), it appears as a filamentous bacillus.

H. influenzae requires growth factors V and X that are present in the blood.

• Factor V is a nicotinamide adenine dinucleotide (NAD), a coenzyme, and is heat labile.

• Factor X is hemin. It is required for cytochrome synthesis and the function of enzymes like catalase and peroxidase. It is heat stable.

When H. influenzae is cultured on the same plate as Staphylococcus aureus. The colonies of H. influenzae that are closer to S. aureus, have larger growth. This is termed satellitism (Option C). This occurs because factor V is released by the hemolytic action of S. aureus.

The image below shows satellitism.



©MARROW

H. influenzae has a capsular polysaccharide, which is a major antigenic determinant. Capsular typing is done to identify strains. Importantly, the type b capsular polysaccharide has a polyribosyl ribitol phosphate (PRP) antigen. PRP is targeted during immunization.

H. influenzae can also cause meningitis, laryngo-epiglottitis, pneumonia, bronchitis, and suppurative lesions like arthritis, endocarditis, and pericarditis.

Cefotaxime or ceftazidime is the drug of choice for the treatment of Haemophilus meningitis. H. influenzae produces beta-lactamase, hence resistant to beta-lactam antibiotics like ampicillin.

Solution to Question 11:

HSV-DNA demonstration in CSF by PCR is the investigation of choice for HSV encephalitis.

Herpes simplex virus (HSV) encephalitis is defined as inflammation of brain parenchyma caused by the herpes virus, and it typically presents as an acute febrile illness.

Patients typically present with acute onset of fever, altered levels of consciousness, personality changes, and evidence of either focal or diffuse neurologic signs and symptoms, especially in the temporal lobe. Focal or generalized seizures occur in many patients with HSV encephalitis. Other most commonly encountered focal findings include aphasia, ataxia, upper motor neuron or lower motor neuron type lesions, and cranial nerve deficits.

HSV-DNA demonstration in CSF by PCR is the investigation of choice for HSV encephalitis. Other investigations that can help include CSF examination, CSF culture, serological studies, MRI, CT, EEG, and brain biopsy.

The T2-weighted MRI of the brain will show hyperintensity in the temporal lobe in most of the patients. Bilateral involvement of the temporal lobe is more common in HSV encephalitis.

The image below is a coronal T2-weighted MR image showing a high signal intensity in the temporal lobes in a patient with HSV encephalitis.



Management includes supportive therapy (monitoring of intracranial pressure, fluid restriction, avoidance of hypotonic IV fluids), anticonvulsants for seizures, and intravenous acyclovir is beneficial in the treatment of HSV encephalitis.

Other options:

Option A: IgM antibodies in CSF - Serological tests to detect antibodies are used for primary herpes infections.

Option B: Tzanck smear of CSF - A positive smear shows multinucleated giant cells with faceted nuclei and ground glass appearance. It is a rapid, inexpensive, and fairly sensitive diagnostic

test. The Tzanck smear is typically used to diagnose cutaneous herpes infections, such as herpes simplex or varicella-zoster virus. It is not an appropriate diagnostic test for HSV encephalitis, as it does not provide definitive confirmation of the presence of HSV in the CSF.

Option D: CSF culture in chick embryos - It is not a sensitive test used for the diagnosis of HSV encephalitis. Negative CSF viral cultures have no value in ruling out HSV encephalitis. Although this method can be used to isolate HSV, it is not as sensitive or specific as PCR for viral DNA and it can take a longer time to obtain results.

Solution to Question 12:

A patient with fever and filarial antigen test positive is suggestive of filariasis. The next most appropriate step for this patient is the detection of microfilariae in the blood smear.

Classical filariasis occurs due to the obstruction of lymphatics by the adult worms, which results in lymph stasis. The disease initially causes acute attacks of lymphangitis and lymphadenitis. As time goes by the lymphedema worsens and is associated with the thickening of the skin and fibrosis of lymph vessels or blockage by adult worms. The lymphatic vessels of the male genitalia are most commonly affected in bancroftian filariasis, producing episodic funiculitis, epididymitis, and orchitis. Hydrocele or elephantiasis is a very common sign of chronic bancroftian filariasis.

The standard method for diagnosing active infection is the identification of microfilariae in a blood smear by microscopic examination. The microfilariae that cause lymphatic filariasis circulate in the blood at night showing a nocturnal periodicity. Microfilariae can be detected in blood smears of samples obtained at night between 10 PM to 2 AM. A thick smear should be made and stained with Giemsa or hematoxylin and eosin. For increased sensitivity, concentration techniques like membrane filter concentration can be used.

DEC provocation test (Option B) refers to the administration of Diethylcarbamazine (DEC) at a dose of 100 mg or 2 mg/kg body weight orally and subsequently collecting blood samples. This causes the mobilization of microfilariae into the peripheral blood and thus microfilariae can be detected 20-50 minutes after the DEC administration in the sample of peripheral blood.

Although the DEC provocation test has the advantage of blood sample collection even during day time, the guidelines from CDC, NVBDCP, and IAPSM suggest blood sample collection at night, followed by identification of microfilariae in a blood smear by microscopic examination as the standard method for diagnosis (Option C).

Methods to detect adult worms include:

• Lymph node biopsy to detect adult worms. Bone marrow biopsy is not a standard diagnostic procedure for filariasis, as adult worms typically reside in the lymphatic system rather than the bone marrow (Option A).

• X-ray (if worms are calcified)

• High-frequency ultrasound and doppler of the scrotum (Option D).

Serologic techniques provide an alternative to the microscopic detection of microfilariae for the diagnosis of lymphatic filariasis. Patients with active filarial infection typically have elevated levels of anti-filarial IgG4 in the blood and these can be detected using routine assays.

• Antigen detection by enzyme-linked immunosorbent assay (ELISA) and immunochromatographic test (ICT)

• Antibody detection tests have low sensitivity and specificity

• Molecular diagnosis is done by PCR (polymerase chain reaction). This test is positive only when microfilaria are present in peripheral blood. It is negative in chronic filariasis.

Diethylcarbamazine is the drug of choice. WHO has recently recommended a triple-drug combination of ivermectin, DEC, and albendazole as a faster and more effective alternative to expedite the elimination of filariasis.



Solution to Question 13:

This clinical scenario along with the microscopic findings of gram■negative intracellular diplococci indicates infection with Neisseria gonorrhoeae. The treatment of choice for the infection is ceftriaxone.

Gonorrhea is a disease acquired by sexual contact. The first step in infection is the adhesion of gonococci to the urethra or other mucosal surfaces. The incubation period is 2-8 days. The major symptoms include purulent urethral discharge and dysuria.

In males, the disease starts as acute urethritis presenting with purulent discharge. The patients may also have penile lymphangitis when the soft tissue inflammation is extensive. This infection may extend along the urethra to involve the prostate, seminal vesicles, and epididymis.

In females, the initial infection involves the cervix uteri and urethra which may extend to Bartholin glands, endometrium, and fallopian tubes. Pelvic inflammatory disease and salpingitis may lead to sterility. Rarely peritonitis may develop with perihepatic inflammation (Fitz-Hugh-Curtis syndrome).

The discharge is collected and subjected to Gram stain, revealing gram■negative intracellular diplococci within phagocytes. Stuart's transport medium is used. Cultures can be obtained from chocolate agar or Mueller-Hinton agar. Specifically, Thayer-Martin agar is used to

isolate N. gonorrhoeae.

Single-agent therapy with ceftriaxone is preferred for the treatment of gonococcal infections.

According to CDC guidelines (2021), treatment of uncomplicated gonococcal infections is as follows:

• Weight <150 kg – Ceftriaxone 500 mg intramuscular (IM) in a single dose

• Weight ≥150 kg – Ceftriaxone 1 g IM in a single dose

The doses of ceftriaxone are higher than the previous guidelines as the gonococcal minimum inhibitory concentrations (MICs) are rising worldwide. The British BASHH guidelines also recommend a high dose of 1g.

Concurrent chlamydial infection must also be treated unless it has been excluded by molecular testing.

First-line drug administered is

• Oral doxycycline 100 mg BD for 7 days

Partner treatment is also required.

Other options:

Option A: Erythromycin: This is not the preferred treatment for gonorrhea, as it is less effective and has a higher rate of resistance compared to ceftriaxone.

Option B: Azithromycin: Although it can be used in combination with ceftriaxone for the treatment of gonorrhea, it is not the treatment of choice as a single agent.

Option D: Penicillin G: This was once a treatment option for gonorrhea, but it is no longer recommended due to widespread resistance.

Microbiology INI-CET 2023

Question 1:

A microorganism is resistant to meropenem and aminoglycoside and sensitive to ticarcillin-clavulanate, and cotrimoxazole. What is the likely microorganism?

a) Stenotrophomonas maltophila

b) Burkholderia cepacia

c) Pseudomonas aeruginosa

d) Acinetobacter baumannii

Question 2:

A patient with a history of bone marrow transplant presented with a lesion in the lung. On microscopic examination, the lesion is shown below. What is the likely causative agent?



a) Histoplasmosis

b) Cryptococcosis

c) Mucormycosis

d) Aspergillosis

Question 3:

Identify the organism in the given image.

a) Aspergillus

b) Mucor

c) Talaromyces

d) Blastomyces

Question 4:

A pregnant woman comes for an antenatal check-up. She has a history of stillbirth at 28 weeks in her previous pregnancy. She also has a history of multiple sexual partners and vaguely remembers having a painless ulcer on her vulva prior to her last pregnancy. What would be the first step of investigation for this lady?

a) HIV ELISA

b) VDRL

c) Dark ground microscopy

d) Culture

Question 5:

Match the diseases with their maximum incubation period.

a) 1-a, 2-b, 3-c, 4-d

b) 1-d, 2-c, 3-b, 4-a

c) 1-d, 2-c, 3-a, 4-b

d) 1-b, 2-a, 3-d, 4-c

Question 6:

A CSF sample of an elderly male presenting with meningitis is subjected to gram staining and revealed gram-positive bacilli. Which of the following is true regarding the motility of the causative agent?

a) Motile at 22°C and 37°C

b) Motile at 37°C, non motile at 22°C

c) Motile at 25°C, non motile at 37°C

d) Nonmotile at 22°C and 37°C

Question 7:

Which of the following is a protozoan parasite, that can be seen inside Kupffer cells?

a) Toxoplasma

b) Leishmania donovani

c) Babesia

d) Plasmodium vivax

Question 8:

Which of the following organism is transmitted by the vector shown in the image below?

a) Orientia tsutsugamushi

b) Kyasanur forest disease

c) Chandipura virus

d) Anaplasma

Question 9:

A lady cleans her cat litter regularly to avoid infection by toxoplasma during her pregnancy. What is the infective stage of this organism in the cat feces?

a) Bradyzoites

b) Tissue cysts

c) Tachyzoites

d) Oocysts

Question 10:

Which of the following is a false statement according to the CDC classification of bioterrorism agents?

a) Category A - Easily transmitted and are associated with high mortality rates

b) Category D - Re-emerging diseases that are high priority agents

c) Category C - Emerging diseases that are easy to disseminate and are associated with high mortality rates

d) Category B - Moderately easy to disseminate and are associated with low mortality rates

Question 11:

The given image of the cell wall belongs to which of the following organism?



a) Staphylococcus aureus

b) Escherichia coli

c) Neisseria meningitides

d) Pseudomonas aeruginosa

Question 12:

A patient presents with PUO. He is suspected to be suffering from brucellosis. Multiple dilutions of his serum sample are put up for a standard agglutination test. The results came as follows: Negative agglutination for the serum dilutions 1:20 and 1:40; But 1:80 and 1:160 were positive for agglutination. Which of the following is responsible for the negative test in the first 2 dilutions?

a) Incorrect antigen

b) Incorrect antibody

c) Antibody excess

d) Antigen excess

Question 13:

A patient presented with a lesion on the left buttock. He has a history of heart transplant 15 years back. Aspirate from the lesion shows gram-positive filamentous bacteria. What is the likely causative agent?

a) Nocardia

b) Bacillus anthracis

c) Mycobacterium leprae

d) Mycobacterium tuberculosis

Question 14:

An elderly male patient who underwent a kidney transplant 10 years back is on immunosuppressive therapy. He gives a history of cough and dyspnea for the past 1 week, and the sputum microscopy shows gram-positive budding yeasts cells and pseudohyphae. What is the likely diagnosis?

a) Invasive mucormycosis

b) Invasive aspergillosis

c) Invasive candidiasis

d) Information is not enough for the diagnosis

Question 15:

The toxin produced by which of the following organisms acts by the mechanism shown in the image below?



a) Vibrio cholerae

b) Corynebacterium diphtheriae

c) Bacillus anthracis

d) Shigella

Question 16:

A 13-year-old girl presented with fever and retro-orbital pain for 2 days. The tourniquet test and NS1 antigen test were positive. Which of the following features is considered a criterion for the severe form of this disease?

a) AST and ALT >1000

b) Leucopenia plus thrombocytopenia

c) Persistent vomiting

d) Hepatomegaly >1cm

Question 17:

A 25-year-old patient presented with a 3-day history of fever, headache, myalgia and retro-orbital pain with no signs of bleeding. On examination, maculopapular rash is seen. Which is the next appropriate step in management?

a) NS1 antigen detection

b) Widal test

c) Rapid influenza antigen test

d) IgM ELISA

Question 18:

Which of the following diseases is not transmitted by the vector shown in the image below?

a) Ehrlichia

b) Babesiosis

c) Orientia

d) Crimean-Congo haemorrhagic fever (CCHF)

Question 19:

Which of the following is a biosafety level-3 infection?

a) Extensively drug-resistant (XDR)- TB

b) Ebola Virus

c) Salmonella typhi

d) Crimean-Congo haemorrhagic fever (CCHF) virus

Question 20:

Which of the following statements are advantages of saline wet mount over iodine wet mount?

a) 1 and 2

b) 1 and 3

c) 2 and 3

d) 1, 2 and 3

Question 21:

Which of the following helps the Human immunodeficiency virus (HIV) to penetrate into the host cell?

a) gp 120

b) gp 41

c) CCR5

d) CXCR4

Question 22:

An HIV-positive patient presented with headache, and neck stiffness. Fundoscopy revealed papilledema. India ink staining of the CSF showed an encapsulated organism. Which of the following statements is true regarding this condition?

a) Start ART after 2-3 weeks

b) Opening pressure will be normal

c) Start amphotericin monotherapy

d) 80% patients will have positive radiological findings

Question 23:

Which of the following techniques is used to detect autoantibodies?

a) ELISA

b) Indirect immunofluorescence

c) Flow cytometry

d) Fluorescent in-situ hybridization

Question 24:

Which of the following toxins acts by the mechanism shown below?



a) Clostridium difficile exotoxin

b) Diphtheria toxin

c) Botulinum toxin

d) Streptococcal pyrogenic exotoxin

Question 25:

In which of the following organisms' genetic material is adenine not equal to thymine?

a) HIV

b) E coli

c) Human mitochondrial cells

d) Herpes virus

Question 26:

An 18-month-old child presents with fever, irritability, neck stiffness and Kernig's sign positive. The mother gives a history of painful genital ulcers a year ago which had resolved spontaneously. What is the most likely causative agent?

a) Herpes simplex virus-2

b) Cytomegalovirus (CMV)

c) Human papilloma virus (HPV)

d) Varicella zoster virus

Question 27:

Which of the following organisms exhibits pathogenesis by cell-to-cell transfer, facilitated by actin filaments, eliminating the need for extracellular transport?

a) Enterohemorrhagic E. coli

b) Salmonella typhi

c) Listeria monocytogenes

d) H. pylori

Question 28:

A 19-year-old boy from a rural area presents with a 12-day history of altered mental status, decreased vision, ataxia, multiple cranial nerve palsies and hemiplegia. The electron microscopy image of the organism is given below. What is the likely organism?



a) Acanthamoeba

b) Naegleria fowleri

c) Sappinia

d) Balamuthia

Question 29:

An immunocompromised patient presents with a white corrugated plaque on the lateral border of the tongue that cannot be rubbed off. Which of the following is the causative agent?

a) Candida

b) Human herpes virus-8

c) Herpes simplex virus (HSV)-1

d) Epstein Barr virus

Question 30:

Which of the following organisms requires the use of reverse transcriptase in PCR?

a) Parvovirus B19

b) Adenovirus

c) Cytomegalovirus (CMV)

d) Respiratory Syncytial virus (RSV)

Question 31:

A farmer presented with fever and a skin lesion as shown in the image below. Blood investigations revealed creatinine levels of 1.4 mg/dl and platelet levels of 86,000 cells/mm3. What is the likely diagnosis?

a) Anthrax

b) Leptospirosis

c) Brucellosis

d) Scrub typhus

Question 32:

A COVID-19 patient who was treated with steroids, presented with blood-tinged nasal discharge, 3 weeks after recovery. The histopathological examination of the discharge showed the following findings. Which of the following organisms is likely to be involved?

a) Aspergillus

b) Mucor

c) Candida

d) Coccidioides

Question 33:

You suspect dengue in a patient who presented with a 3-day history of fever, headache, and retro-orbital pain. Blood investigations revealed Hb=14 g/dl, WBC= 4000 cells/ mm3 and platelet count = 8900 cells/mm3. Which of the following tests will not be useful in confirming the diagnosis of this condition?

a) NS1 antigen detection

b) Viral culture and isolation in aedes albopticus c6/36 cell line

c) RT-PCR

d) IgM ELISA

Question 34:

Which of the following viruses has the least teratogenic effect?

a) Chickenpox

b) Rubella

c) Cytomegalovirus (CMV)

d) Herpes simplex virus (HSV)-2

Answer Key

Question No.	Correct Option
1	a
2	d
3	b
4	b
5	c
6	c
7	b
8	a
9	d
10	b
11	c
12	c
13	a
14	c
15	b
16	a
17	a
18	c
19	a
20	b
21	a
22	a

23	b
24	d
25	a
26	a
27	c
28	a
29	d
30	d
31	d
32	b
33	d
34	d

Detailed Explanations

Solution to Question 1:

Stenotrophomonas maltophila, formerly called Pseudomonas maltophila, is an opportunistic pathogen that causes wound infection, urinary tract infection, and septicemia. It is sensitive to cotrimoxazole.

Intrinsic resistance is the innate ability of a bacterial species to resist the activity of a particular antimicrobial agent through its inherent structural or functional characteristics, which allow tolerance of a particular drug or antimicrobial class.

Burkholderia cepacia, formerly called Pseudomonas cepacia, is an opportunistic pathogen that causes fatal necrotizing pneumonia. B. cepacia complex strains are intrinsically resistant to aminoglycosides, polymyxin, and first and second-generation cephalosporins but sensitive to meropenem, co-trimoxazole, doxycycline, minocycline, chloramphenicol and ceftazidime.

Solution to Question 2:

The given clinical scenario of lung lesion and the image showing septate hyphae with branching at acute angles is suggestive of aspergillosis.

Invasive aspergillosis occurs due to infection with Aspergillus species. It is often seen in immunocompromised patients. Most cases of invasive aspergillosis involve the lungs. The clinical features include fever, cough which may be productive, chest discomfort, hemoptysis, and shortness of breath. Voriconazole is the drug of choice.

Other manifestations include allergic bronchopulmonary aspergillosis where central bronchiectasis is usually seen. Aspergilloma and superficial infections like otomycosis or mycotic keratitis are the other presentations.

Histopathological examination of the tissue section shows characteristic dichotomous branching at an acute angle. The hyphae stain poorly with hematoxylin and eosin, so periodic acid–Schiff (PAS) stain is used.

Other methods for diagnosis include detection of galactomannan (Aspergillus cell-wall component) and culture isolation. Aspergillus fumigatus grows dark green colonies, while Aspergillus niger appears black. Yellow-green colonies are seen in case of Aspergillus flavus. Microscopic examination of colonies stained with lactophenol cotton blue would show septate hyphae and conidiophores.

Solution to Question 3:

Microscopy showing broad, aseptate, ribbon-like hyphae with irregular branching is suggestive of mucor.

Mucormycosis/ Zygomycosis is an opportunistic infection caused principally by species of rhizopus (most prevalent), mucor, and absidia. The common clinical form is rhino-orbital-cerebral mucormycosis in which there is an invasion of the blood vessels in the nasal cavity, eyes, and brain causing thrombosis, infarction, and necrosis. The blackish discoloration is due to arterial and venous thrombosis leading to gangrene.

Diagnosis can be made by staining them with 10% potassium hydroxide and culture. On Sabouraud agar at 30-37°C, they appear as grey-white colonies with a thick cottony, fluffy surface. Under the microscope, they appear as irregularly branched broad hyphae with aerial sporangiophores which end in a sac containing spores called sporangium.

Other options:

Options A and C: Aspergillus and Talaromyces (previously known as Penicillium) both demonstrate septate hyphae with conidiophores.

Option D: Blastomyces is a dimorphic fungus that exists in the yeast form in body tissues and cultures maintained at 37°C and in the filamentous mold form in cultures at room temperature (25°C). In the yeast form, it shows thick-walled round yeast cells with single broad-based budding, giving a figure of 8 appearance.

Solution to Question 4:

Venereal disease research laboratory test (VDRL test) is the first investigation in this patient with a history of stillbirth and painless genital ulcer, which is likely suggestive of syphilis.

Syphilis is caused by Treponema pallidum, a spirochete that is actively motile. It exhibits characteristic corkscrew-like motility.

Primary syphilis occurs subsequent to inoculation of treponema pallidum following sexual contact. It manifests as a painless, non-tender, single papule over the genitalia, that rapidly becomes eroded and usually becomes indurated, and is called a hard chancre. It can be accompanied by lymphadenopathy. The chancre generally heals within 10-40 days even without treatment, leaving a scar.

The obstetric history in an untreated syphilitic woman typically gets better with subsequent pregnancies. Initial few pregnancies can result in abortions and stillbirths, subsequent pregnancies can result in live births of infants with stigmata of syphilis. Finally, after a few pregnancies, healthy infants are born.

It can be diagnosed using VDRL, and RPR tests. It can be confirmed by FTA – ABS testing or TPHA, TPI testing.

Dark ground microscopy is the most specific and sensitive technique to diagnose syphilis when an active chancre or condyloma lata is present. Since the ulcer is not active in this patient, it is not useful in this scenario.

Smear specimens can be stained using Fontana stain and pathological tissues are stained using Levaditi stain. Silver impregnation technique can also be used.

Solution to Question 5:

The correct match is:

The incubation period is the time interval between the invasion of the infectious agent and the appearance of the first symptom or signs in the body.

The incubation period of syphilis is 10-90 days and varies inversely with the size of the Spirochaete inoculum.

The incubation period of SARS infection ranges from 5 to 10 days.

The incubation period of Hepatitis A infection ranges from 10 to 50 days (usually 14-28 days).

Chickenpox is caused by varicella zoster virus (HSV-3). The incubation period is between 10 to 21 days.

Diseases	Incubation period
1. Hepatitis A	d. 50 days
2. Chickenpox	c. 21 days
3. SARS-CoV-2	a. 10 days
4. Syphilis	b. 90 days

Solution to Question 6:

The given scenario of an elderly male with meningitis and smear showing gram-positive bacillus is suggestive of Listeria infection. Listeria is motile at 25°C and non-motile at 37°C.

Listeria exhibits a characteristic slow tumbling end-over-end motility at 22–28°C but is non-motile at 37°C. This is because the peritrichous flagella are produced by the bacillus optimally between 20°C and 30°C. This is known as differential motility as the motility pattern changes with a change in temperature.

Listeria monocytogenes is a gram-positive bacillus. It can survive at refrigerator temperatures (4°C), under conditions of low pH and high salt. So it is able to overcome food preservation and safety barriers, making it an important foodborne pathogen. The virulence factor produced by Listeria is called listeriolysin O. This helps the bacteria to escape and causes septicemia. They are catalase-positive, will not produce H_2S, and shows beta hemolysis. This helps to differentiate Listeria from other gram-positive bacilli.

L. monocytogenes enters the body after ingestion of contaminated foods such as refrigerated milk, cheese, fruit, or vegetables. Healthy persons exposed to L. monocytogenes in food may develop mild, self-limiting febrile gastroenteritis lasting 1–3 days. This develops after an incubation period of 6–48 hours. Symptoms include fever, chills, headache, myalgias, abdominal pain, and diarrhea.



Solution to Question 7:

The protozoan parasite, that can be seen inside Kupffer cells is Leishmania donovani.

Kupffer cells are a component of the mononuclear phagocytic system found in the liver. The amastigote form of leishmania also known as the Leishman-Donovan (LD) body (shown in the image below) is an intracellular form found inside macrophages, monocytes, and other reticuloendothelial cells. It contains a kinetoplast, which consists of a parabasal body and a blepharoplast. Flagellum is absent.



Leishmaniasis is a slowly progressing indigenous disease caused by a protozoan parasite of the genus Leishmania.

The life cycle of Leishmania involves two forms:

• The promastigote develops and lives extracellularly in the sandfly vector, and

• The amastigote multiplies intracellularly in the reticuloendothelial cells of the host

It is transmitted by the bite of an infected female Phlebotomus sandfly which inoculates the flagellated promastigotes into the skin of the host. This condition causes fever with rigor and chills associated with splenomegaly and hepatomegaly. It lowers immunity, causes persistent fever, anemia, and liver and spleen enlargement, and if left untreated, death may occur due to opportunistic infection.

Diagnosis of leishmaniasis is mainly by demonstration of amastigotes within macrophages in tissue aspirates such as blood smears, bone marrow, splenic, or lymph node aspirate. It is the gold standard for diagnosis.

Solution to Question 8:

The image shows the trombiculid mite which is responsible for the transmission of Orientia tsutsugamushi.

Scrub typhus, also known as chigger-borne typhus, is caused by Orientia tsutsugamushi. The vector is trombiculid mite. Humans are infected when they are bitten by the mite larvae known as chiggers. The incubation period is 1- 3 weeks. Patients develop a characteristic eschar at the site of the mite bite, along with regional lymphadenopathy and maculopapular rash. It can be associated with fever, headache, and conjunctival injection. Meningoencephalitis, myocarditis, and pneumonia can be seen in severe cases.

Serologic assays are used for diagnosis. Weil-Felix reaction is a heterophile agglutination reaction used for diagnosis, wherein the sera are tested for agglutinins to the O antigens of certain Proteus strains OX19, OX2, and OXK. OXK is agglutinated only in scrub typhus. Doxycycline is the

treatment of choice for both adults and children.

Solution to Question 9:

The infective stage of Toxoplasma found in the cat feces is oocysts.

The oocysts develop from the sexual reproduction of Toxoplasma in the gut lumen of the cat and pass out via the feces. It takes 1 to 5 days for the oocysts to become infective after being excreted in the feces. Hence, daily removal of cat litter and safe disposal can prevent infection.

Both asexual reproduction (schizogony) and sexual reproduction (gametogony) occur within definitive hosts such as cats. The tachyzoites and tissue cysts represent stages in asexual multiplication, while the oocyst is formed by sexual reproduction, which occurs only in the definitive hosts such as cats.

Only the asexual forms i.e., tachyzoites, and tissue cysts are present in intermediate hosts such as humans, mice, rats, etc.





Humans acquire the infection upon consumption of meat containing tissue cysts, food or water containing mature oocysts, or fingers contaminated by cat feces.

Manifestations are mostly asymptomatic and if symptomatic can be congenital or acquired. Features such as fever, headache, myalgia, splenomegaly, and lymphadenopathy commonly affect the cervical lymph nodes.

Central nervous system involvement is observed in toxoplasma infection affecting immunocompromised individuals. Toxoplasma infection in pregnancy can lead to congenital toxoplasmosis that most frequently presents with chorioretinitis.

Solution to Question 10:

According to CDC, the agents of bioterrorism are classified only into 3 categories (Category A, B, and C), there is no category D.

Bioterrorism is the intentional release or threat of release of biological agents (i.e. viruses, bacteria, fungi, or their toxins) in order to cause disease or death among the human population or food crops and livestock to terrorize a civilian population or manipulate the government.

The agents of bioterrorism are classified into 3 categories:

Category A bioterrorism agents are those which are easily transmitted from one person to the other and are associated with high mortality rates and thus have the potential to cause major public health impacts.

Category B bioterrorism agents are those which are moderately easy to disseminate and are associated with moderate morbidity and low mortality rates.

Category C bioterrorism agents are those which are emerging pathogens such as Nipah and Hantavirus, that could be engineered for mass dissemination in the future due to ease of availability, high mortality and morbidity rates, and ease of production and dissemination.

Solution to Question 11:

The image showing a cell wall with a capsule and lipooligosaccharides (LOS) is suggestive of Neisseria meningitidis.

Unlike lipopolysaccharide (LPS), LOS does not have a long O-antigen side chain. It is an important virulence factor and is found in a few organisms such as N. gonorrhoeae, N. meningitidis, H. ducreyi, etc.

The genus Neisseria consists of Gram-negative, aerobic, non-sporulating, non-motile, oxidase-positive cocci arranged typically in pairs (diplococci). It includes two pathogens, Neisseria meningitidis, and Neisseria gonorrhoeae. The important difference is the presence of a capsule in N. meningitidis.

Solution to Question 12:

The negative test in the first 2 tubes is due to the prozone phenomenon. Here, antibodies are in excess compared to antigens leading to weak or absent precipitation.

The lattice hypothesis explains precipitation reactions and agglutination reactions. The amount of precipitate formed is greatly influenced by the relative proportions of antigens and antibodies. Precipitation occurs when a lattice consisting of alternating antigen and antibody molecules is formed.

• The prozone phenomenon is caused by excessive antibodies in the test system. There is weak/absent precipitation, giving a false-negative test.

• In the zone of equivalence, antigen and antibody are in optimum proportions and maximum precipitation occurs. Lattice formation is enhanced.

• Post-zone phenomenon is due to the presence of excessive antigens in the test system. There is weaker/absent precipitation.

Prozone phenomenon and post-zone phenomenon can give rise to false-negative results in precipitation reactions. This can be overcome by making serial dilutions of serum, thereby optimizing the concentrations of antigens and antibodies in the test system.



Solution to Question 13:

The given clinical scenario of local abscess and aspirate showing filamentous gram-positive bacteria are suggestive of Nocardia.

Nocardia species appear as filamentous organisms with extensive hyphae-like branching. They are aerobic, gram-positive, catalase-positive, and urease positive. They are weakly acid-fast because they stain with carbolfuchsin and retain when decolorized with 1–4% sulfuric acid instead of the stronger acid-alcohol decolorant.

Infections caused by Nocardia include cutaneous, subcutaneous, or systemic lesions. It may lead to local abscesses, cellulitis, or lymphocutaneous lesions. Systemic nocardiosis occurs more often in immunodeficient persons.

Systemic nocardiosis usually caused by N.asteroides manifests primarily as pulmonary disease, pneumonia, lung abscess, or other lesions resembling tuberculosis. Extrapulmonary manifestation may involve the brain, kidneys, and other organs.

The image given below is a smear of modified acid-fast Kinyoun staining showing filamentous red-stained partially acid-fast rods suggestive of Nocardia.



Solution to Question 14:

Cough and dyspnea in an immunosuppressed patient with microscopy showing gram-positive budding yeasts cells and pseudohyphae suggest a diagnosis of invasive candidiasis.

Candida is the most common cause of opportunistic mycoses in the world. Immunocompromised individuals are at an increased risk of systemic candidiasis. The most common species implicated in candidal infections are C. albicans, C. parapsilosis, and C. glabrata. Morphologically, they are yeast-like fungi. They exist as yeast and pseudohyphae (image below). It appears as budding gram-positive cells on wet films and smears. Mycelial forms suggest colonization. Candida albicans form chlamydospores on corn meal agar.



When colonies of Candida albicans are inoculated in serum at 37°C, the yeast cells form elongations called germ tubes within 2 hours. The image below shows germ tube formation, also

known as the Reynolds Braude phenomenon.

Clinically, infection with Candida spp. can present as cutaneous and mucosal candidiasis, systemic candidiasis, or rarely, as chronic mucocutaneous candidiasis. Mucocutaneous infections manifest as thrush affecting the tongue, lips, gums, and palate. It is seen as a patchy to confluent whitish pseudomembranous lesion and is commonly associated with AIDS. Candida may also invade the vaginal mucosa, leading to vulvovaginitis, and the nails and nail plate, leading to onychomycosis. Systemic candidiasis occurs when Candida enters the bloodstream. It can manifest as kidney infections, urinary tract infections, infective endocarditis, arthritis, meningitis, endophthalmitis etc.

Solution to Question 15:

The above image depicts the inactivation of elongation factor-2 which is the mechanism of action of diphtheria toxin.

Corynebacterium diptheriae produces the diphtheria toxin, which is an AB subunit toxin. While the B subunit helps in attachment to cell surface receptors, the A subunit causes ADP-ribosylation of elongation factor-2 (EF-2) resulting in the inhibition of protein synthesis.



The exotoxin A produced by Pseudomonas aeruginosa, also an AB subunit toxin, acts by the same mechanism to cause inhibition of protein synthesis.

This infection presents as fever and sore throat, which progresses to dyspnea due to obstruction by the pseudomembrane. Bull's neck appearance is seen due to cervical lymphadenitis. Complications like damage to the heart (myocarditis), kidney (tubular necrosis), liver, and adrenal glands can develop due to the toxin.

Other options:

Option A: Cholera toxin is an enterotoxin produced by Vibrio cholerae. It is an AB subunit toxin where the B subunit binds to GM1 ganglioside receptors on the surface of jejunal epithelial cells. The A subunit is internalized and causes prolonged activation of adenylate cyclase resulting in increased cAMP activity which results in watery diarrhea.



Option C: Bacillus anthracis produces toxins that are made of 3 proteins-

• Protective antigen (PA) binds to specific cell receptors and mediates entry of edema factor and lethal factor into the cell.

• Edema factor (EF) is an adenylate cyclase, which is activated only inside the target cells, leading to intracellular accumulation of cyclic AMP and causing tissue edema.

• Lethal factor (LF) causes tissue death.

Option D: Shiga toxin or verotoxin is an exotoxin produced by Shigella species. It consists of two subunits A and B. The B subunit binds to globotriaosylceramide receptors on the target cell, and then, the A subunit gets internalized. The A subunit leads to inhibition of 60S ribosome, leading to shut down of protein synthesis.

Solution to Question 16:

The given clinical scenario of a 13-year-old girl presenting with fever and retro-orbital pain with a positive tourniquet and NS1 antigen tests is suggestive of dengue fever. AST or ALT >=1000 is a criterion for severe dengue.

Dengue is an acute febrile disease caused by the mosquito-borne dengue viruses (DENVs), consisting of four serotypes (DENV 1 to 4), that are members of the Flaviviridae family, genus Flavivirus. It is transmitted from person to person by Aedes aegypti mosquitoes.

The World Health Organisation released guidelines in 2009 regarding the diagnosis of probable dengue, dengue with warning signs, and a category known as severe dengue. In addition to the classic DHF/DSS syndrome, rare cases of encephalitis or encephalopathy, liver damage, or myocardial damage are also included in the category of severe dengue.

Criteria for severe dengue include:

• Severe plasma leakage leading to shock (DSS) and fluid accumulation with respiratory distress

• Severe bleeding

• Severe organ involvement- Liver: AST or ALT >= 1000, CNS: Impaired consciousness, heart and other organs.

Probable dengue includes those people who live in/ travel to dengue-endemic areas. Fever and 2 of the following criteria: nausea, vomiting, rash, aches and pain, Tourniquet test positive, leukopenia and any warning sign.

Warning signs include:

• Abdominal pain or tenderness

• Persistent vomiting

• Clinical fluid accumulation

• Mucosal bleed

• Lethargy, restlessness

• Hepatomegaly >2 cm

• Laboratory: increase in HCT concurrent with thrombocytopenia.

After an incubation period of 3–15 days, classical dengue fever begins with an abrupt onset of high fever. The condition is self-limiting though debilitating with headache, retro-orbital pain, myalgia, arthralgia, petechiae rash, and leucopenia. A macular-papular recovery rash appears 3–4 days after the onset of fever, and it usually starts on the trunk before spreading peripherally. The rash in dengue appears as 'white islands in a red sea' due to erythema with petechiae.

Shown below is an image of the rash in dengue:

Solution to Question 17:

The given clinical scenario of a 25-year-old patient presenting with a 3-day history of fever, headache, myalgia, retro-orbital pain with no signs of bleeding and a maculopapular rash on examination is suggestive of dengue fever. The next appropriate step in management is getting an NS1 antigen detection test done.

Dengue is an acute febrile illness caused by the dengue virus that is transmitted by the Aedes aegypti mosquito.

It presents as sudden-onset fever, retroorbital pain, myalgia, arthralgia, and a rash. The initial rash in dengue may appear as flushing erythema of the face in the first 2 days of symptoms.
A second rash occurs 3-6 days after the onset of fever and appears maculopapular or morbilliform. Sometimes, lesions may coalesce and appear as 'white islands in a red sea' due to generalized erythema with petechiae and islands of sparing.

Before day 5 of illness, during the febrile period, dengue infections may be diagnosed by detection of viral antigens like non-structural viral protein antigen(NS 1) by ELISA or rapid tests. This is considered the mainstay of diagnosis.

Virus isolation in cell culture and viral RNA detection by nucleic acid amplification tests (NAAT) can also be done.

Serological tests, such as IgM ELISA (option D), are performed after day 5 when dengue viruses and antigens are no longer present in the blood due to the appearance of specific antibodies.

Other options:

Option B - Widal test is used to diagnose typhoid fever caused by Salmonella Typhi and Paratyphi. It is characterized by gradual onset headache, malaise, step ladder pyrexia and gastrointestinal symptoms like abdominal pain with either constipation or diarrhoea. They also present with rose spots on the skin during the 2nd to 3rd week, that fade on applying pressure.

The following image depicts rose spots in enteric fever.



Option C: The rapid influenza antigen test uses immunofluorescence to demonstrate the influenza viral antigen on the surface of nasopharyngeal cells. The typical clinical disease has a sudden onset of fever, headache, and generalised myalgia. The presence of respiratory symptoms is prominent. It is not associated with retro-orbital pain and maculopapular rash, as given in the above scenario.

Solution to Question 18:

The vector shown in the image is hard tick. Scrub typhus caused by Orientia tsutsugamushi is transmitted by trombiculid mites.

Hard tick has an oval-shaped body, no antennae with 4 pairs of legs and is covered by a chitinous shield on its dorsal surface(scutum). It transmits diseases such as Ehrlichia, babesiosis, viral hemorrhagic fevers like Crimean-Congo haemorrhagic fever (CCHF) and Kyasanur forest disease (KFD), tick typhus, viral encephalitis (e.g., Russian spring-summer encephalitis), viral fevers (e.g., Colorado tick fever), tularemia, tick paralysis.

The following image shows a soft tick.

Orientia tsutsugamushi is transmitted by trombiculid mite which is as shown below.



Solution to Question 19:

Extensively drug-resistant(XDR)-TB is a biosafety level-3 infection.

Biohazard levels, also known as biosafety levels, are classifications of safety precautions required in the clinical microbiology laboratory based on specific pathogens handled when performing laboratory procedures.

There are 4 classifications of biosafety levels (BSLs). Each level contains specific recommendations for a clinical microbiology laboratory, with a focus on laboratory practises, safety equipment, and facility construction. As each level progresses, it includes additional biosafety considerations from the previous level.

Microorganisms classified as biosafety level-3 can be lethal to humans, but there are vaccines and other treatments available for them. They include Mycobacterium tuberculosis, anthrax, many types of viral encephalitis, hantavirus, Rift Valley fever, malaria, Rocky Mountain spotted fever and yellow fever.

GROUP	DEFINITION	BACTERIA	VIRUS	FUNGI
Group1	Biological agents unlikely to cause human disease	No pathogenic organisms	-	-
Group 2	Biological agents that can cause human disease and may be hazardous to workers but are unlikely to spread to the community; effective treatment or prophylaxis is usually available.	Bacillus (except anthrax)StreptococciStaphylococciE. coliClostridiumspeciesCornybacterium diphtheriaSalmonella typhi	AdenovirusCalcivirusCornavirusHerpesvirusInfluenza virus	CryptococcusCandidaDermatophytesAspergillus
Group 3	Biological agents that can causesevere human diseaseand are a serious hazard to workers; they may spread to the community;but effective treatment or prophylaxis is usually available.	B.AnthracisBrucellaCoxiella burnettiTularemiaM.tuberculosis	Lymphocytic choriomeningitisHaanta virusSt.louis encephalitis virusJapanese encephalitis virusWest nile encephalitis virus	
Group 4	Same as group 3 expect that effective treatment or prophylaxis is usually not available		LassaEbolaMarbugHerpes simiaeCrimean-Congo haemorrhagic fever (CCHF) virus	

Solution to Question 20:

The advantages of saline wet mount over iodine wet mount include its ability to demonstrate the motility of trophozoites and to differentiate between bile and non-bile-stained eggs.

Saline wet mount is an unstained preparation made by using physiological saline, while iodine wet mount is prepared either using diluted Lugol's iodine or Dobell and O'Connor iodine solution.

Saline wet mount aids in illustrating the motility of trophozoites such as E. histolytica, Balantidium coli, and Giardia lamblia because it does not dry up as quickly as iodine wet mount.

It also helps differentiate between bile-stained and non-bile-stained eggs. Bile stained eggs appear brown, while non-bile stained eggs appear colourless. However in iodine wet mounts, it is challenging to distinguish between bile-stained and non-bile-stained eggs because they are both stained brown.

Since iodine clearly stains the glycogen vacuoles and nuclei, the idoine wet mount helps in confirming the identification of cysts. Iodine-stained protozoan cysts display pale refractile nuclei, brown glycogen material, and yellow-gold cytoplasm.

Solution to Question 21:

The HIV virus can penetrate the host's CD4+ cells with the help of the viral glycoprotein gp120, also known as the spike antigen which is present in its envelope.

HIV-1 virion is spherical and contains an electron-dense, cone-shaped core surrounded by a lipid envelope derived from the host cell membrane. Target cells of HIV include CD4+ helper T cells (worst affected), monocyte-macrophages, and dendritic cells.

The viral envelope is studded by two viral glycoproteins, gp120 and gp41, which are coded by the env gene. Gp120 also known as the principal envelope antigen forms surface spikes while gp41 functions as a transmembrane anchoring protein. Both of these are critical for HIV infection of cells.

Viral glycoprotein gp120 binds to CD4+ cells in the host. Binding to the CD4+ receptor requires the participation of a co-receptor molecule, which has been identified as CXCR4 for the T cell-tropic HIV strains (X4 strains) and CCR5 for the macrophage-tropic strains (R5 strains).

This binding leads to a conformational change in gp41 causing exposure of a hydrophobic region called the fusion peptide at the tip of gp41. This peptide inserts into the cell membrane of the target cells (e.g., T cells or macrophages), leading to the fusion of the virus with the host cell.

The virus core contains:

• Major capsid protein p24

• Nucleocapsid protein p7/p9

• Two copies of viral genomic RNA

• The three viral enzymes (protease, reverse transcriptase, and integrase)

The viral core is surrounded by a matrix protein called p17, which lies underneath the virion envelope.

The image below shows the pathogenesis of HIV infection.



Solution to Question 22:

The above clinical scenario and the stain findings are suggestive of cryptococcosis and the causative organism is Cryptococcus neoformans. In a patient with Cryptococcus infection, antiretroviral therapy should be started 2-3 weeks after initiation of treatment. This is due to a risk of immune reconstitution which can lead to severe inflammation and further worsen symptoms.

C. neoformans spreads from inhalation of the yeast cells in pigeon faeces causing infection in HIV/AIDS and other immunosuppressive conditions. In causes influenza-like respiratory infection in immunocompetent persons. In immunocompromised
it disseminates to CNS causing cryptococcal meningoencephalitis and patients may complain of headaches, neck stiffness, and disorientation as in this case. Other common sites of dissemination include the skin, adrenals, bone, eye, and prostate gland.



Diagnosis is made by demonstration of yeast cells or cryptococcal antigen in CSF. Cryptococcus presents with raised intracranial pressure so precautions must be taken before taking a CSF sample. (Option B) India ink stain is used to demonstrate capsulated budding yeast cells in CSF as they have a distinctive appearance because their capsules exclude ink particles. Rapid detection of cryptococcal antigen in CSF and blood can also be done. Colonies develop within a few days on most media at room temperature or 37°C. Media with cycloheximide should be avoided as it inhibits Cryptococcus. MRIs and CT scans can be used but they're only reliable in about 21-27% of cases(Option D).

Combination therapy of amphotericin B and flucytosine has been considered the standard treatment for cryptococcal meningitis(Option C).

Solution to Question 23:

Circulating autoantibodies in serum are best detected by indirect immunofluorescence.

The two-step indirect immunofluorescence assay is used to find out if a serum specimen contains antibodies specific to a particular organism. A known antigen is first attached to a slide. After incubating the slide with a serum sample, the sample is removed, the slide is washed, and a second fluorescein-labelled anti-immunoglobulin is added. If the patient's serum contains the organism-specific antibodies, the organism will appear green under the fluorescence microscope.

This assay is the accepted method for identifying autoantibodies in autoimmune disorders (such as antinuclear antibodies) and detecting antibodies to certain microorganisms (eg, T. pallidum)

Other options:

Option A: Enzyme-linked Immunosorbent Assay (ELISA) is a type of heterogenous enzyme immunoassay that involves the use of an immunosorbent, an absorbing material specific for one of the components of the reaction: the antigen or antibody. It mainly detects antigens of infectious agents present in clinical specimens like rotavirus detection in fecal specimens and HIV detection.

Option C: Flow cytometry is a laser-based method that analyzes the cells and cell components based on the presence of certain cell surface molecules which are named as per cluster of differentiation (CD) nomenclature. It is very effective for the study of the immune system and its response to infectious diseases and cancer.



Option D: Fluorescent In Situ Hybridisation (FISH) is a simple technique for the detection of specific DNA sequences in cells/tissues. It uses fluorescent labelled probes to bind to complementary DNA. FISH can be performed on prenatal samples, peripheral blood cells, touch preparations from cancer biopsies, and even fixed archival tissue sections.



Solution to Question 24:

The given image depicts a superantigen which binds to the variable β region of the T-cell receptor. Streptococcal pyrogenic exotoxin (SPE) is a superantigen.

Superantigens are a third variety of the biological class of antigens that activate a very large number of T cells, irrespective of antigen specificity. This means they can activate T-cells directly without being processed by antigen-presenting cells (APCs).

• The variable β region of the T-cell receptor (vβ of TCR) appears to be the receptor for superantigens.

• They directly bridge non-specifically between MHC–II of APCs and T-cells.



The overproduction of cytokines, particularly IFN-γ, is a result of the massive activation of T cells. IFN-γ in turn activates macrophages to produce IL-1, IL-6, and TNF-α, all of which may contribute to a “cytokine storm” causing severe symptoms of shock and multiple organ failure.

Examples of superantigens:

• Bacterial

• Staphylococcal toxic shock syndrome toxin-1

• Streptococcal toxin

• Mycoplasma arthritidis mitogen 1

• Yersinia enterocolitica

• Yersinia pseudotuberculosis

• Viral

• Epstein–Barr virus (EBV)

• Cytomegalovirus (CMV))

• Rabies nucleocaspid

• HIV-encoded negative regulatory factor.

• Fungal

• Malassezia furfur.

Solution to Question 25:

The genetic material of HIV consists of a single-stranded RNA and therefore adenine will not be equal to thymine.

The pyrimidine components of RNA can differ from those of DNA. Although RNA contains the ribonucleotides adenine (A), guanine (G), and cytosine (C), it contains uracil (U) instead of thymine (T).

Because RNA is a single strand that is complementary to only one of two strands of a gene, its adenine content will not be equal to its uracil content and its guanine will not be equal to its cytosine content.

Chargaff's rule states that double-stranded DNA, such as that found in E. coli, human mitochondrial cells and herpes virus, will always have equal concentrations of (A = T) and (G = C).

The pairings of purine and pyrimidine nucleotides on opposite strands are very specific and rely on hydrogen bonds between A and T and G and C. The base pairs A-T and G-C are frequently referred to as Watson-Crick base pairs.

Solution to Question 26:

The given clinical scenario of an 18-month-old child presenting with fever, irritability, neck stiffness and positive Kernig's sign (CNS manifestations indicating meningitis) with the mother giving a past history of painful genital ulcers, is suggestive of neonatal herpes due to herpes simplex virus (HSV)-2. CMV, HPV and varicella zoster virus do not present with painful genital ulcers and can be easily ruled out.

Painful genital ulcers are commonly seen in genital herpes and chancroid due to Hemophilus ducreyi ((ducreyi makes you cry).

Herpes infection can be caused by HSV-1 and HSV-2 viruses. HSV-1 causes orofacial lesions, i.e. herpes labialis, while HSV-2 mainly causes genital herpes which is characterised by vesiculoulcerative lesions of the penis of the male or of the cervix, vulva, vagina, and perineum of the female.

The typical herpes lesions are thin-walled, umbilicated vesicles, the roof of which breaks down, leaving tiny superficial ulcers which heal without scarring. The most common site of cutaneous lesions is in the face, cheeks, chin, around the mouth and forehead. A cesarean section is performed in a case of active genital herpes at the time of labour in order to prevent neonatal herpes infection.

One of the following three disease patterns can be seen in infants infected during delivery or in the postpartum period: (1) disease limited to the skin, eyes, or mouth; (2) encephalitis with or without skin, eye, and mouth disease; or (3) disseminated infection affecting several organs, such as the brain, lungs, liver, heart, and adrenal glands.

Infants suffering from encephalitis usually exhibit clinical symptoms that point to bacterial meningitis at 8- 17 days of life. These symptoms include seizures, poor feeding, lethargy, neck stiffness, irritability, and positive Kernig's sign. When left untreated, HSV encephalitis results in 50% of infant deaths and severe neurologic sequelae in the majority of survivors.

Acyclovir at a high dose of 60 mg/kg/day, administered intravenously every eight hours, should be the first line of treatment for any infant with a confirmed or suspected neonatal HSV infection.

Other options

Option B - The transplacental spread of cytomegalovirus (CMV) primarily affects the foetus, resulting in microcephaly, intracranial calcification, hepatosplenomegaly, chorioretinitis, mental retardation, and sensorineural deafness. It is not associated with meningitis.

Option C - Adolescents who are sexually active can contract genital HPV infection, which is typically caused by HPV types 6 and 11. It can develop as a result of inoculation during childbirth through an infected birth canal or as a result of unintentional dissemination from cutaneous warts. It results in moist, fleshy, verrucous growths on the skin of the labia majora, perineum, perianal region, and vulva which are known as genital warts/venereal warts/condyloma acuminata.

Option D - Congenital varicella syndrome (CVS) is characterized by hypoplasia of limbs, psychomotor retardation, IUGR, chorioretinal scarring, cataracts, microcephaly and cutaneous scarring. It is not associated with meningitis.

Solution to Question 27:

Listeria monocytogenes exhibit pathogenesis by cell-to-cell transfer with the help of the host cell's actin filaments.

L. monocytogenes is a short, gram-positive, non-spore-forming bacteria. It enters the body through the GIT after ingestion of contaminated food such as cheese or vegetables. It has two cell-surface proteins, internalins A and B, which interact with the epithelial E-cadherin and promote its own phagocytosis into the gut epithelial cells, leading to the formation of the phagolysosome.

Low pH activates the bacterium to produce listeriolysin O, which lyses the phagolysosome and allows Listeria to escape into the cytoplasm of the epithelial cell. Here, the organisms proliferate and produce ActA, a listerial protein that induces host cell actin polymerization, which propels them to the host cell membrane.

While pushing against the host cell membrane, they produce elongated protrusions called filopods, which are ingested by adjacent epithelial cells, macrophages, and hepatocytes, resulting in the release of listeriae and the cycle starts again.

In this way, Listeria moves from cell-to-cell using the host's actin framework without being exposed to antibodies, complement systems, or neutrophils. Gram-negative bacteria like Shigella flexneri and rickettsiae also make use of the host cells' actin and contractile system to spread their infections, just like Listeria.

Other options:

Option A: Enterohemorrhagic Ecoli (EHEC) is one of the six different types of diarrheagenic E.coli. They produce verocytotoxin (VT) or Shiga-like toxin (SLT), which can lead to a spectrum of illnesses, from mild diarrhea to more severe, life-threatening conditions such as hemorrhagic colitis (bleeding in the colon) and hemolytic uremic syndrome (HUS), which can cause kidney failure.

Option B: Salmonella typhi is a non-spore-forming, facultative anaerobic, gram-negative bacilli. Following ingestion of contaminated foods or beverages, salmonellae enter the small intestine, where they pass through the epithelium via specialised M-cells overlying the Peyer's patches, and then enter the intestinal lymphatics, with subsequent invasion into the bloodstream. Unlike Listeria, it does not utilise the actin filament of the host cell.

Option D: H.pylori is a gram-negative, flagellated, helical bacterium. Virulence factors of H.pylori include:

• Flagella- which allows motility in viscous gastric mucous

• Urease production- urease generates ammonia and bicarbonate from endogenous urea, creating an alkaline microenvironment that protects it from gastric acidic pH.

• Adhesins- for bacterial attachment to surface foveolar cells

• Toxins- such as cytotoxin-associated gene A (CagA) and Vacuolization cytotoxin (VacA). CagA is translocated into the gastric epithelial cells where it interferes with the epithelial cells' cytoskeletal structure and induces IL-8 production, leading to neutrophil attraction. VacA causes gastric mucosal injury by inhibiting T cells.

Solution to Question 28:

In the given clinical scenario of a 19-year-old boy presenting with a 12-day history of altered mental status, decreased vision, ataxia, multiple cranial nerve palsies and hemiplegia with electron microscopy showing an organism with characteristic spine-like pseudopodia (acanthopodia) is suggestive of infection caused by Acanthamoeba species.

Acanthamoeba exists as an active trophozoite form and a resistant cystic form, both of which are infective. The large, 20–50 μm trophozoite form is distinguished by spine-like pseudopodia called acanthopodia.

Human beings acquire infection by inhalation of cysts or trophozoites, ingestion of cysts or through traumatized skin or eyes. Following inhalation, the trophozoites reach the lungs and invade the central nervous system through the bloodstream, producing granulomatous amebic encephalitis (GAE).

It presents chiefly as two chronic conditions: (1) keratitis and (2) encephalitis.

With a slow relapsing course, keratitis resembles severe herpetic keratitis; however, in the case of an amebic infection, the eye is extremely painful. The initial symptoms are unilateral photophobia, excessive tearing, redness and foreign body sensation. Keratitis and uveitis can cause permanent vision loss or blindness. Diagnosis includes demonstrating the cyst in corneal scrapings using wet mount, histology, and culture.

Granulomatous amebic encephalitis is a serious infection of the brain and spinal cord that usually affects people who have a weakened immune system. The clinical picture is that of intracranial space-occupying lesions with seizures, pareses, and mental deterioration. It is diagnosed by demonstration of trophozoites and cysts in brain biopsy, culture and immunofluorescence microscopy using monoclonal antibodies.

No effective treatment is available for GAE. With limited success, multidrug combinations such as pentamidine, sulfadiazine, rifampicin, and fluconazole are used.

Other options:

Option B: The electron microscopy image of the infective form of Naegleria fowleri shows rounded pseudopodia called lobopodia. The infection usually follows swimming in freshwater rivers or ponds. The amoebae invade the nasal mucosa and pass through the olfactory nerve branches in the cribriform plate into the meninges.

It is mainly associated with primary amoebic meningoencephalitis. The disease advances rapidly, causing fever, headache, vomiting, stiff neck, ataxia, seizure and coma.

Option C: Sappinia, a newly discovered human pathogen of the central nervous system (CNS), can cause amoebic encephalitis in humans. In the single case of Sappinia amoebic encephalitis that has been reported, a sinus infection occurred prior to the onset of symptoms. Sappinia can be distinguished from other free-living amoeba by the presence of a distinctive double nucleus in which the 2 nuclei are closely apposed with a central flattening.

Option D: Balamuthia is a leptomixid free-living amoeba, reported to cause GAE. It exists in the ameboid trophozoite stage and is not associated with acanthopodia as seen in Acanthamoeba species.

Solution to Question 29:

The given clinical scenario of an immunocompromised patient presenting with a white corrugated plaque on the lateral border of the tongue that cannot be rubbed off is suggestive of oral hairy leukoplakia caused by Epstein Barr Virus (EBV).

Hairy leukoplakia is a distinct oral lesion on the lateral border of the tongue caused by EBV that typically occurs in immunocompromised patients. It appears as white, confluent patches of fluffy or hairy, hyperkeratotic thickenings, almost always on the lateral border of the tongue. Unlike candida-caused oral thrush, the lesion cannot be rubbed away.

Predisposing factors include Immunodeficiency, haematological malignancies, and steroid abuse.

The Epstein-Barr virus replicates in maturing epithelial cells but not in basal or parabasal keratinocytes. On microscopy, the upper spinous layer shows the characteristic hyperparakeratosis and acanthosis with balloon cells.

Investigations show high levels of EBV DNA in the blood with circulating antibodies against viral capsid antigen (VCA) and early antigen (EA)■restricted proteins.

Other options:

Option A: Oral Thrush/ oral candidiasis associated with Candida albicans presents as white plaque-like lesions with cheesy exudate on an erythematous mucosa. They are typically painful and can be easily brushed off.



Option B: AIDS-associated Kaposi sarcoma (KS) due to human herpes virus-8 (HHV-8) is an AIDS-defining illness and is the most common malignancy associated with HIV infection. Kaposi sarcoma mainly affects the hard■palate mucosa in particular. The gingiva, tongue and buccal mucosa are also involved. The initial lesion is a reddish-purple macule, which develops into a purple nodule swelling that could be widespread and ulcerated.



Option C: Herpes simplex virus (HSV) infection is common in and around the mouth. HSV-1 is the primary cause of gingivostomatitis, which is most common in children. The gums are swollen, inflamed and bleed easily. The mucous membranes of the tongue, pharynx, palate, and buccal region are covered in vesicles that appear as white plaques. The plaques turn into ulcers with a pseudomembrane that is yellow in colour. The regional lymph nodes are enlarged and tender.

Solution to Question 30:

Respiratory Syncytial virus (RSV) which is the only RNA virus in the given list of viruses, requires the use of reverse transcriptase in PCR.

Reverse Transcriptase PCR (RT-PCR) is a molecular analysis method used for the detection of RNA and thus helps in detecting RNA-containing viruses. It can be used to quantify mRNA levels by generating complementary DNA (cDNA) copies of the mRNA using the reverse transcriptase enzyme.

Respiratory Syncytial virus belongs to the Pneumovirus genera of the Paramyxoviridae family. They contain a linear, single-stranded, nonsegmented, negative-sense RNA.

The northern blot technique can also be used to detect RNAs. Here, the mixture of mRNA molecules is separated by electrophoresis and then transferred to a nitrocellulose membrane and hybridized with radiolabeled probes.

Other options:

Option A: Parvovirus B19 is a member of the Parvoviridae family, which consists of very small viruses with a particle size ranging from 18 to 26 nm. They have single-stranded, linear DNA in them.

Option B: Adenoviruses are members of the Adenoviridae family and have double-stranded, linear DNA in their genomes.

Option C: The Herpesviridae family includes the cytomegalovirus (CMV), which has a double-stranded, linear DNA genome.

All DNA viruses can be detected by polymerase chain reaction (PCR). It is a technique used to amplify a selected DNA sequence. It can be used to synthesize millions of copies of the selected DNA sequence in a few hours, enabling the detection of even trace amounts of DNA in a sample.

Solution to Question 31:

The given clinical scenario of a farmer presenting with fever and a black eschar lesion with elevated creatinine levels of 1.4 mg/dl, indicating kidney injury, is suggestive of scrub typhus.

Scrub typhus is caused by Orientia tsutsugamushi and transmitted by trombiculid mites. Humans are infected when they are bitten by the mite larvae known as chiggers. Transovarian transmission of microbes occurs in mites.

The incubation period is 1-3 weeks. The patient typically develops a characteristic eschar which is a punched-out ulcer covered with a blackened scab at the site of the mite bite, with regional lymphadenopathy and maculopapular rash.

Renal involvement is a common systemic manifestation of scrub typhus. The severity of renal involvement can range from minor urinary abnormalities such as albuminuria and active urinary sediments such as cellular casts, pyuria, and hematuria to acute kidney injury (AKI) necessitating the immediate initiation of renal replacement therapies.

Other options:

Option A: Anthrax- Cutaneous anthrax (Hide porter's disease) present as a black eschar at the site of entry. At the site of entry of organism or spores, a papule develops in about 1–7 days. This develops into a vesicle that later breaks down into an ulcer with a black, central, necrotic eschar

surrounded by edematous skin. The disease used to be common in dock workers, carrying loads of hides and skins on their bare backs. Renal involvement is unlikely to be seen in this condition.

Leptospirosis (option B) and Brucellosis (option C) can be easily ruled out as they do not present with the black eschar lesions as shown in the image.

Solution to Question 32:

The given clinical vignette of a COVID-19 patient who was treated with steroids, presenting with blood-tinged nasal discharge, 3 weeks after recovery and histopathological examination showing broad, aseptate, ribbon-like hyphae with right angle branching is suggestive of mucor.

Mucormycosis is a fungal infection of the nose and sinuses caused by Mucor. It is an opportunistic infection seen even in mild to moderate COVID-19 patients with strong predisposing factors like uncontrolled diabetes and those immunosuppressed or on steroids.

The common clinical form is rhino-orbital-cerebral mucormycosis which occurs due to invasion of the blood vessels in the nasal cavity, eyes, and brain causing thrombosis, infarction, and necrosis. The typical finding is black necrotic material filling the nasal cavity and eroding the nasal septum, hard palate and turbinates due to arterial and venous thrombosis leading to gangrene.

Diagnosis can be made by staining them with 10% potassium hydroxide and culture. On Sabouraud agar at 30-37°C, they appear as grey-white colonies with a thick cottony, fluffy surface. Under the microscope, they appear as broad, ribbon-like, non-septate hyphae with hyphal elements branching at right angles. They also contain aerial sporangiophores which end in a sac containing spores called sporangium.

The drug of choice is the antifungal Amphotericin, administered either as Amphotericin B 0.6-1 mg/kg/day or Liposomal Amphotericin B 5 mg/kg/day which has higher CNS penetration. Surgical debridement of all dead necrotic tissues should be done endoscopically.

Other options:

Options A: Aspergillus shows narrow, septate hyphae with typical dichotomous branching at acute angles (45°) on microscopy.



Option C: Candida, the causative agent for candidiasis, appears as budding gram-positive cells on wet films and smears. Culture colonies are creamy-white and have a yeasty odour. Candida albicans, the most commonly implicated pathogenic species, form chlamydospores on corn meal agar. They exist as yeast and pseudohyphae (buds do not separate from parent cells) and true hyphae form.

The image below shows candidal hyphae and chlamydospores.



Option D: Coccidioides immitis is a dimorphic fungus, known to cause Valley Fever. It exists in two distinct forms, saprophytic and parasitic. In the soil, the organism exists as a mold with septate hyphae resembling a shape of a barrel.

When entering a host, the arthroconidia (spores) break off from the hyphae and evolve into round structures called spherules. Spherules are large sac-like structures filled with endospores as shown below.



Solution to Question 33:

In a suspected case of dengue, where a patient presented with a 3-day history of symptoms like fever, headache, and retro-orbital pain, all the above investigations can be done except for IgM-ELISA, which is used to identify antibodies against the dengue virus.

Dengue viruses belong to flaviviruses and present with sudden onset of fever (saddleback type), frontal headache, retro-orbital pain, back pain, and severe myalgias. It is transmitted from person to person by the bite of the Aedes aegypti mosquitoes.

The virus can be found in serum, plasma, circulating blood cells, and other tissues for 4-5 days following the onset of the illness. Virus isolation, nucleic acid detection, or antigen detection can be used to diagnose the infection in its early stages. Serology is the preferred method of diagnosis at the end of the acute phase of infection.

Before day 5 of illness, during the febrile period, diagnosis can be made by the following tests:

• Virus isolation can be done during the period of viremia i.e. usually before day 5. The mosquito cell line C6/36 (cloned from Aedes albopictus) or AP61 are the host cells of choice for routine isolation of the dengue virus.

• RT-PCR - They have higher sensitivity than virus isolation and a much faster turnaround time.

• NS1 antigen can be detected up to 7-10 days after the onset of illness.

After day 5, dengue viruses and antigens disappear from the blood with the appearance of specific antibodies. The presence of IgM antibody or rising titres of IgG antibody on ELISA is used for detecting and distinguishing primary and secondary dengue infections.

Solution to Question 34:

In the given list of viruses, herpes simplex virus (HSV)-2 has the least teratogenic effect.

HSV-2 presents as a genital tract infection transmitted via sexual contact. Premature labour or IUGR may occur if the primary infection is acquired during the last trimester. As the transplacental route of spread is uncommon, it is considered to have a less teratogenic effect on the baby.

Caesarean delivery is recommended in cases of active primary genital HSV infection because the virus shed from the cervix or lower genital tract during vaginal delivery is a major source of infection in the fetus.

Babies with neonatal herpes exhibit three categories of disease: (1) localized lesions of the skin, eye, and mouth, (2) encephalitis with or without skin involvement and (3) disseminated disease. It is treated with IV acyclovir.

Other options:

Option A: The varicella-zoster virus, which causes chickenpox, can pass through the placenta and result in congenital varicella syndrome (CVS). Hypoplasia of the limbs, psychomotor retardation, IUGR, chorioretinal scarring, cataracts and microcephaly are the characteristics of CVS. If maternal infection develops after 20 weeks, the risk of congenital malformation is almost absent. Varicella zoster immunoglobulin (VZIG) should also be given to newborns exposed within 5 days of delivery.

Option B: The transplacental spread of the rubella virus is known to cause congenital rubella syndrome (CRS) in the baby. When this infection develops in the first, second or third months, there is a 60%, 25%, or 10% chance of significant abnormalities, respectively. Congenital rubella syndrome predominantly includes sensorineural deafness, patent ductus arteriosus (PDA), cataracts and retinopathy. The virus predominantly affects the fetus and is extremely teratogenic if contracted within the first trimester.

Option C: Cytomegalovirus (CMV) replicates in respiratory epithelial cells, salivary glands, and kidneys and persists in lymphocytes. The transplacental route primarily affects the foetus, resulting in microcephaly, intracranial calcification, hepatosplenomegaly, chorioretinitis, mental retardation, and sensorineural deafness. Unlike rubella, CMV may damage fetal organs throughout gestation.

Microbiology INI-CET 2024

Question 1:

A 24-year-old female comes to the OPD with dysuria. She has a non-irritant profuse white discharge from her vagina. On examination, her labia were inflamed and swollen. The openings of her urethra and Bartholin's glands were congested. On further collection of her endocervical discharge, intracellular gram-negative diplococci were found. Which of the following is the most likely diagnosis?

a) Neisseria Gonnorhea

b) Chlamydia Trachomatis

c) Treponema Pallidum

d) Hemophilus Ducreyi

Question 2:

Drug resistance is seen with which of the following STD?

a) Syphilis

b) Gonorrhoea

c) Chancroid

d) Donovanosis

Question 3:

Which of the following parasites reproduce directly from egg to larva without fertilisation through parthenogenesis?

a) Strongyloides stercoralis

b) Ancylostoma doudenale

c) Enterobius

d) Trichuris trichura

Question 4:

Which is the best test used for Malarial Parasite speciation?

a) Acridine orange stained thin smear

b) Giemsa-stained thin smear

c) Giemsa-stained thick smear

d) Quantitative Buffy Coat (QBC)

Question 5:

Assertion(A): Late trophozoite and schizont stage in Plasmodium falciparum are not seen in blood smears. Reason(R): These two stages are trapped in the capillary endothelium of internal organs due to cytoadherence.

a) Both Assertion (A) and Reason (R) are true and reason is Reason (R) is the correct explanation of Assertion (A)

b) Both Assertion (A) and Reason (R) are true and reason is Reason (R) is not the correct explanation of Assertion (A)

c) Assertion (A) is true and Reason (R) is false

d) Assertion (A) is false and Reason (R) is true

Question 6:

Which of the following pathological features is incorrectly matched with the disease?

a) No reaction: Immunodeficiency disease

b) Cytopathic and cytoproliferative effect : Chickenpox

c) Mononuclear and granulomatous lesions: Berylliosis

d) Suppurative and purulent lesions: Syphilis

Question 7:

Out of the given lab sterilisation techniques, which of the following can be used for a disposable pipette tip?

a) A only

b) D only

c) A,B

d) C,D

Question 8:

Which of the following is transmitted by all the components of blood?

a) Syphilis

b) Malaria

c) Filaria

d) Toxoplasma

Question 9:

The following statements are connected with 'because'

a) Statement 1 and Statement 2 are true and Statement 2 is correct explanation of Statement 1

b) Statement 1 and Statement 2 are true but Statement 2 is not the correct explanation of Statement 1

c) Statement 1 is true and statement 2 is false

d) Both the statements 1 and 2 are false

Question 10:

A construction worker presents with a lock jaw and muscle rigidity. He gives a history of exposure to a rusted iron nail through an existing wound. Which of the following is the mechanism of action of the toxin responsible for this presentation ?

a) Inhibition of Acetyl choline release

b) Inhibition of glycine and GABA release

c) Inhibition of epinephrine

d) Inhibition of Nor-epinephrine

Question 11:

Which of the following immunodeficiency diseases predisposes to infection by catalase-positive organisms?

a) Severe combined immunodeficiency (SCID)

b) Hyper IgE syndrome

c) X-linked agammaglobulinemia

d) Chronic granulomatous disease

Question 12:

Which of the following is not a clinical manifestation of Parvovirus B19 infection?

a) Hydrops fetalis

b) Persistent anemia or pure red cell aplasia in immunodeficient individuals

c) Transient aplastic crisis in children with hemolytic anemia

d) Febrile illness with petechial rash

Question 13:

A 50-year-old patient who underwent a renal transplant recently came with complaints of watery diarrhoea. The stool sample revealed oocysts of size 30x15 microns. Identify the organism



a) Cyclospora cayatenensis

b) Cystoisospora

c) Cryptosporidium

d) Microsporidium

Question 14:

An 18-day-old baby presents with altered sensorium, and neck rigidity with vesicular lesions on the anterior abdomen and the forearm. The mother revealed a history of treatment for painful genital ulcers a year ago and the baby was born through vaginal delivery. CSF examination reveals multinucleate giant cells. Which of the following is the most likely involved organism?

a) HSV - 2

b) Coxsackie virus

c) Enterovirus

d) VZV

Question 15:

Which of the following is true with respect to the organism shown below?



a) Most commonly involves the lungs

b) Associated with budding yeast cells and pseudohyphae

c) Only reservoir is man

d) Opportunistic infections are seen in patients with CD4 count < 900

Question 16:

Which of the following is most commonly associated with S.agalactiae ?

a) Neonatal sepsis

b) Pharyngitis with or without tonsillitis

c) Dental caries

d) Endocarditis

Question 17:

Match the following:

a) 1-b, 2-a, 3-c, 4-d

b) 1-a, 2-c, 3-b, 4-d

c) 1-d, 2-c, 3-a, 4-b

d) 1-c, 2-a, 3-d, 4-b

Question 18:

Assertion (A) - Mycobacterium is detected in sputum samples with N-acetyl cysteine (NAC) and NaOH. Reason(R) - The sputum samples are treated by NAC to kill commensals and NaOH to liquefy the sputum.

a) Assertion(A) is true but Reason(R) is false

b) Both assertion(A) and Reason(R) are true and Reason(R) is the correct explanation of assertion(A)

c) Assertion(A) is false but Reason(R) is true

d) Both assertion(A) and Reason(R) are true but Reason(R) is not the correct explanation of assertion(A)

Question 19:

A gardener in Himachal notices a black ulcer in the right upper arm. Axillary lymph nodes are enlarged and matted. Aspirate from the lymph nodes shows elongated yeats 3-5 micrometres. Mycelium was isolated from the aspirate and grown at 22 degrees Celsius in SDA, and the image is given below. What is this condition?



a) Coccidiomycosis

b) Histoplasmosis

c) Sporothrix schenkeii

d) Blastomycosis

Question 20:

A 2-year-old male child presents with fever, hematuria, thrombocytopenia, and diarrhea for the last 3 days. What is isolated from the stool sample?

a) Shigella flexneri

b) Salmonella enteritis

c) Aeromonas

d) E. Coli O157:H7

Question 21:

A long-beaded, filamentous, branching, weakly acid-fast organism is isolated from a lesion caused by trauma to the chest by a blunt farming tool. What is the isolated organism?



a) Bacillus anthrax

b) Listeria

c) Nocardia

d) Mycobacterium tuberculosis

Question 22:

What is not true regarding naive T-cell interaction with CD4 effector T-cells?

a) T cell interaction with APC

b) Interaction between CD 80 and CD 28

c) PD-1 on T cell interacts with PD ligand 1 on APC

d) Cytokine mediated proliferation and differentiation

Question 23:

Which of the following is a method of antibiotic sensitivity testing?

a) E test

b) Elek test

c) Latex agglutination test

d) CAMP test

Question 24:

Which of the following is transmitted by walking barefoot in soil?

a) Madurella mycetomatis

b) Curvularia

c) Coccidioides immitis

d) Aspergillus fumigatus

Question 25:

Which of the following cancers can be prevented by vaccination?

a) A,B,C,D

b) A and C only

c) A and D only

d) C and D only

Question 26:

Which of the following statements is true about Cryptococcus?

a) 1,2,3

b) 2,3,4

c) 2,4

d) 1,2

Question 27:

What is the culture medium for M. tuberculosis?

a) Lowenstein-Jenson medium

b) Blood agar

c) Novy-MacNeal-Nicolle medium

d) Sabouraud dextrose agar

Question 28:

A 2-year-old presents with excessive watery diarrhoea and severe dehydration. What is the causative organism?

a) Trichuris Trichura

b) Shigella

c) Rotavirus

d) Yersenia

Question 29:

Match the causative organism with the disease.

a) i-c,ii-b,iii-d,iv-a

b) i-b,ii-c,iii-d,iv-a

c) i-b,ii-c,iii-a,iv-d

d) i-d,ii-a,iii-b,iv-c

Question 30:

Which of the following does not confer passive immunity?

a) Toxoid

b) Antitoxin

c) IgA

d) Maternal monoclonal antibody

Question 31:

Which of the following is not a cold sterilization technique?

a) Plasma sterilization

b) Autoclave

c) UV radiation

d) Gamma radiation

Question 32:

Which of the following is not true regarding Ascaris lumbricoides?

a) Most infected humans are asymptomatic

b) Larvae in the lungs produce asthma-like symptoms

c) High larval load can cause rectal prolapse

d) Unembryonated eggs in stool are not infective and embryonated eggs in soil are infective

Question 33:

What is the drug of choice for fasciola hepatica?

a) Triclabendazole

b) Praziquantel

c) Mebendazole

d) Albendazole

Question 34:

A 20-year-old male presented with a pustule that developed into an ulcer which was diagnosed to be a Necrotising fascitis. A sample of the ruptured pustule showed gram-positive cocci in chains. Identify the organism.

a) Streptococcus pyogenes

b) Staphylococcus aureus

c) Enterococcus

d) Micrococcus

Question 35:

A 25-year-old male presented with abdominal pain. On examination, a painful growing mass is felt in the right hypochondrium. Aspiration of the cyst is shown below. Identify the organism.



a) Echinococcus granulosus

b) Hymenolopsis nana

c) T.solium

d) T.saginata

Question 36:

Which of the following drugs is not given in uncomplicated falciparum malaria?

a) Chloroquine

b) Primaquine

c) Arthemeter

d) Pyrimethamine Sulfadoxine

Question 37:

How do you sterilise bacteriological loops used for inoculation?

a) Heat till metal turns red hot

b) Pass over flame 3 times

c) Autoclave

d) Microwave

Answer Key

Question No.	Correct Option
1	a
2	b
3	a
4	b
5	a
6	d
7	a
8	b
9	c
10	b
11	d
12	d
13	b
14	a
15	a
16	a
17	c
18	a
19	c
20	d
21	c
22	c
23	a
24	a
25	c
26	c
27	a
28	c
29	b
30	a
31	b
32	c
33	a
34	a

35	a
36	a
37	a

Detailed Explanations

Solution to Question 1:

The above clinical scenario and endocervical examination of gram-negative diplococci denote a diagnosis of Neisseria gonorrhoea.

Gonorrhoea is a sexually transmitted disease where infection occurs due to the adhesion of gonococci to the urethra or other mucosal surfaces with the help of pili. By the third day of infection, the cocci have penetrated intracellular spaces and reached subepithelial tissue. The incubation period ranges from 2 to 8 days. The main symptoms are urethral discharge and dysuria.

In females :

The infection begins in the cervix uteri and urethra and may spread to the Bartholin glands, endometrium, and fallopian tubes. The vaginal mucosa is usually unaffected in adults because the stratified squamous epithelium is resistant to cocci infection and also because vaginal secretions have an acidic pH. Rarely peritonitis may develop with perihepatic inflammation (Fitz Hugh Curtis syndrome).

In males :

The disease starts as acute urethritis. This infection has the potential to spread down the urethra and into the prostate, seminal vesicles, and epididymis. Chronic urethritis may lead to stricture formation. Infections in the periurethral tissues can lead to abscesses and multiple discharging sinuses (watercan perineum).

Lab diagnosis:

The discharge is collected and subjected to Gram's stain which reveals gram■negative intracellular diplococci within phagocytes. Stuart's transport medium is used. Specifically, Thayer-Martin agar is used to isolate Neisseria gonorrhoeae.

Treatment: Ceftriaxone (250 mg, IM) single-dose + azithromycin 1 g PO, single-dose, or doxycycline 100 mg PO BD for 7 days.

The image below shows the Gram-stain of a urethral exudate showing typical intracellular gram-negative diplococci and pleomorphic extracellular gram-negative organisms



Solution to Question 2:

Neisseria gonorrhoea endures maximum drug resistance among the given organisms causing sexually transmitted diseases.

The ability of N. gonorrhoea to change its antigenic properties and adapt to different environments has rendered it resistant to the majority of antibiotics. Gonococci develop complete resistance to antibiotics, either through chromosomal mutations or by acquiring R factors, which are plasmids.

• Often β-Lactamase producing Gonococci and tetracycline-resistant strains occur together. Hence Penicillin, ampicillin, and tetracycline are no longer reliable for the treatment of gonorrhea and should not be used.

• Changes in DNA gyrase and topoisomerase IV have been identified as mechanisms contributing to fluoroquinolone resistance.

• Mechanisms leading to enhanced drug efflux pump and reduced drug influx through PorB, collectively contribute to resistance against third-generation cephalosporins.

Currently, the effective first-line treatment of choice for uncomplicated Gonococcal infection remains Ceftriaxone (500 mg IM, single dose) plus Azithromycin (1 g PO, single dose) .

Other options :

A) Syphilis : The drug of choice for treating syphilis is benzathine penicillin G. Penicillin has been the mainstay of treatment for syphilis for many years due to its effectiveness in curing the infection and its relatively low cost.

c) Chancroid: The drug of choice for treating chancroid, a sexually transmitted infection caused by the bacterium Haemophilus ducreyi, is azithromycin or ceftriaxone. These antibiotics are effective in treating chancroid and preventing its complications. The choice between azithromycin and ceftriaxone may depend on factors such as patient allergies, local resistance patterns, and individual patient factors.

d) Donovanosis : The drug of choice for treating donovanosis, also known as granuloma inguinale, is typically azithromycin or doxycycline. Donovanosis is a sexually transmitted infection caused by the bacterium Klebsiella granulomatis, and prompt treatment is essential to prevent complications and transmission to others.

Solution to Question 3:

The females of Strongyloides stercoralis are known to be parthenogenetic i.e. capable of generating offspring without the need for fertilisation by a male.

Strongyloides stercoralis has a parasitic cycle and a free-living cycle in soil. Humans are the natural host but morphologically indistinguishable strains may be seen in dogs and cats. Filariform larvae are the infective form and there are two modes of infection: penetration of the skin by 3rd stage filariform larva, and autoinfection (external and internal).

The female worms present in the intestine lay eggs in the mucosa which hatches immediately, releasing the rhabditiform larva. The larva migrates into the lumen of the intestine and passes down to be released into the faeces. The rhabditiform larva may change into a filariform larva while passing through the bowel.

The filariform larvae in the bowel do not leave the host and become a source of autoinfection as they penetrate the colonic mucosa or perianal skin. The rhabditiform larvae passed in the faeces may undergo two types of development in the soil:

• Direct development: The rhabditiform larva moults twice on reaching the soil to become the infective filariform larva. The filariform larva can penetrate the skin and enter the circulation.

• Indirect development: The rhabditiform larvae develop into free-living males and females in the soil.

Other options :

Option B: Ancyclostoma duodenale are dioecious parasites, there is no multiplication within the host, and a single infective larva matures into either a male or a female adult to undergo mating through copulation. It typically takes around 6 weeks from the point of infection for the adult worms to reach sexual maturity and initiate egg-laying.

Option C: Enterobius vermicularis is dioecious.The male is usually unseen as it does not migrate from caecum or appendix. It usually dies after mating and is passed in the faeces. The gravid female migrates down the colon to the rectum and a single worm lays from 5000 to 17,000 eggs

Option D: Trichuris trichuria or the whipworm has a distinct male and female parasite and is typically dioecious. The fertilised female lays about 5000 eggs per day.



Solution to Question 4:

Malarial parasite speciation is best done by a Giemsa-stained thin smear.

Microscopy is the best method for the diagnosis of malaria due to its high sensitivity and specificity.A diluted Giemsa stain is applied to both thick and thin films and left to dry for thirty minutes to two hours. Afterwards, the slide is rinsed and dried.

Thin smear is used to identify the particular infecting species of Plasmodium. It is prepared by air drying for 10 minutes and fixed in methanol. Thus, the plasmodia morphology is not distorted.

Thick smear is used to detect infection and estimate parasite concentration. The correct thickness is the thickness through which a newsprint is barely visible. It is prepared by slow-drying for 30 minutes, which leads to hemolysis and distortion of the plasmodia morphology. The thick film is not fixed using methanol as it can become dehaemoglobinised. Hence, species cannot be detected.

For rapid staining, especially beneficial for fieldwork, the Field's stain is utilized. In India, the frequently employed rapid staining method is the JSB stain, named after Jaswant Singh and Bhattacharji.

The following image shows the difference between thick and thin smear.

• When properly prepared, the thick film will just permit printed letters to be visible when viewed through it.

• A well-prepared thin film will consist of a continuous smear of a single layer of red cells, terminating in a tapered tongue end that falls slightly before reaching the edge of the slide.

Other options:

Option D: The QBC method is a rapid test for malaria which is highly sensitive and specific. In this method, a small amount of blood (50 to 110 µl) is centrifuged in the QBC centrifuge, concentrating parasites near the tip of the red blood cell column.

Option A: Acridine orange is a fluorescent stain used in the QBC test of malaria.RBC tubes are pre-coated with acridine orange, and parasites fluoresce, making them easily visible under an oil immersion microscope. It is hence not the ideal stain used for speciation.

Solution to Question 5:

Late trophozoite, early and mature schizonts are stages in the asexual life cycle of P.falciparum. These stages are usually not seen in the peripheral smear unless the malaria is exceptionally severe or pernicious. In regular conditions, These two stages are trapped in the capillary endothelium of internal organs due to cytoadherence.

Plasmodium falciparum is the most highly pathogenic strain of malaria, causing fatal complications. Hence, it is also called malignant tertian or pernicious malaria. The life cycle of P.falciparum can be divided into three primary stages based on whether parasites are in the liver, the bloodstream or the mosquito's stomach.

Exo-erythrocytic cycle-Human liver :

• The female anopheles mosquito injects parasites (sporozoites) into the human after a blood meal. The sporozoites are sickle-shaped and invade into hepatocytes.

• Sporozoites mature into schizonts which release merozoites

Erythrocytic cycle:

• Merozoites leave the liver and invade red blood cells, in which they reproduce asexually and develop into trophozoites.

• Early ring trophozoites are often seen attached along the margin of the red cell, the so-called form applique or accole. Binucleate rings are common resembling stereo headphones in appearance. Several rings may be seen within a single erythrocyte.

• Typically, the trophozoites vanish from the bloodstream within approximately 24 hours.

• By this time, a strain-specific high molecular weight antigen emerges on the surface of infected red cells resembling a knob-like projection on the erythrocyte membrane.

• Infected red cells with presenting antigen, exit the bloodstream and stick to the walls of small blood vessels(cytoadherence) of internal organs, including the brain, heart, kidneys, lungs, spleen, intestines, bone marrow, and placenta.

• This adherence leads to the sequestration of infected cells in these areas, contributing to severe complications like cerebral malaria.

Sporogonic cycle- Mosquito :

• Gametocytes are taken up by a mosquito when it feeds. (9-12) Gametocytes undergo sexual reproduction in the midgut of the mosquito and develop into sporozoites, which migrate to the salivary glands.

• The mosquito injects sporozoites through saliva into another human source.

Major manifestations of severe falciparum malaria include more than two convulsions in 24 hours, unarousable coma/ cerebral malaria (failure to localise or respond appropriately to stimuli or coma persisting for >30 min after generalised convulsion), acidosis, severe normocytic normochromic anaemia, renal failure, pulmonary oedema, hypoglycemia, shock, and disseminated intravascular coagulation.

Other manifestations include hemoglobinuria, hyperparasitemia, jaundice, and extreme weakness.

The treatment of choice is artesunate given at a dose of 2.4 mg/kg stat IV, followed by 2.4 mg/kg at 12 and 24 hours, and then daily if necessary.

The life cycle of plasmodium falciparum is given below :



Solution to Question 6:

Syphilis is not known to cause suppurative and purulent lesions. It causes macular, papular or mixed maculopapular lesions.

Syphilis presents with the following lesions as it progresses through different stages :

• Primary syphilis - Hard chancre, which is painless papule that becomes eroded and indurated.

• Secondary Syphilis - Skin rashes on palms and soles which may be macular, papular, papulosquamous and nodular, but never vesicular or bullous.

• Condylomata lata- mucocutaneous papules that coalesce to form large pink to grey lesions in warm moist intertriginous areas)

• Generalized lymphadenopathy is seen

• Tertiary Syphilis - Gummas, which are locally destructive granulomatous lesions. They are most commonly seen in bone and skin.

Other options :

Option A - Immunodeficient individuals lack the normal response to antigens. This is the reason for the false negative Mantoux test seen in such individuals.

Option B - Chickenpox belongs to the alpha subfamily of herpes group of viruses which has a cytopathic effect of formation of multinucleated giant cells, with rounding and ballooning of infected cells

Option C - Berryliosis is an occupational lung disease, causing a chronic granulomatous inflammatory disease similar to sarcoidosis. Biopsy reveals non-caseating granulomas with monocytic infiltration.

Solution to Question 7:

The most common and effective method to sterilise disposable pipette tips is by using an autoclave.

Autoclaving involves subjecting the pipette tips to high-pressure steam at temperatures above 121°C (250°F) for around 20 minutes. This process effectively kills microorganisms and sterilises the pipette tips for reuse. However, it's important to check the manufacturer's instructions to ensure that the pipette tips are autoclavable and to follow proper sterilisation protocols to maintain their integrity and functionality.

Solution to Question 8:

Among the options provided, Malaria is the only disease transmitted by all components of blood.

Malaria is primarily spread through the bite of infected Anopheles mosquitoes, which inject sporozoites into the bloodstream but can also be transmitted by transfusion of blood and blood products.

other options:

Option A : Syphilis is primarily transmitted through direct contact with syphilitic sores (chancres), which can occur on the genitals, lips, mouth, or anus during sexual activity.In most cases, transmission via blood transfusion is rare.

Option C: Filarial parasites are primarily transmitted through the bite of infected mosquitoes. However, they can also be found in the blood of infected individuals, which can potentially transmit the infection through blood transfusion. However, plasma and platelets may not typically harbor filarial parasites unless there's concurrent infection or contamination.

Option D: Toxoplasma is transmitted through various routes, including ingestion of contaminated food or water, contact with infected cat feces, and transplacental transmission from mother to fetus. However, it is not primarily transmitted through blood

Solution to Question 9:

The statement Faucial diphtheria can cause death due to myocarditis is true. However, Corynebacterium diphtheria causes myocarditis due to toxaemia and not bacteremia.

Corynebacterium diptheriae produces the diphtheria toxin, which is an AB subunit toxin. While the B subunit helps in attachment to cell surface receptors, the A subunit causes ADP-ribosylation of elongation factor-2 (EF-2) resulting in the inhibition of protein synthesis.



The exotoxin A produced by Pseudomonas aeruginosa, also an AB subunit toxin, acts by the same mechanism to cause inhibition of protein synthesis.

This infection presents as fever and sore throat, which progresses to dyspnea due to obstruction by the pseudomembrane. Bull's neck appearance is seen due to cervical lymphadenitis.

Complications of C.diphtheriae:

• damage to the heart (myocarditis)

• kidney (tubular necrosis)

• liver and adrenal glands can be affected due to the toxin

In respiratory diphtheria patients, cardiac complications, particularly myocarditis, arise in around 10-25% of cases. Cardiac involvement is the leading cause of death in diphtheria, with a mortality rate of 50-75%. Manifestations of cardiac involvement vary and may encompass myocardial dysfunction, rhythm disturbances such as bradyarrhythmias or tachyarrhythmias, and heart blocks necessitating pacing. Elevated cardiac enzymes and ECG abnormalities may signal asymptomatic myocardial involvement.

Solution to Question 10:

The above clinical scenario, coupled with a history of contact with a rusty iron nail, suggests a diagnosis of tetanus, which is caused by the anaerobic bacterium Clostridium tetani. The toxin responsible for muscle rigidity and lockjaw is tetanospasmin produced by Clostridium tetani. It acts by causing presynaptic inhibition of GABA and glycine release.

Tetanospasmin, after secretion by the bacterium is cleaved into two peptides linked by a disulphide bond. The larger peptide diffuses to terminals of neurons and migrates by the retrograde axonal transport system to the spinal cord and brainstem. It diffuses to the glycinergic interneurons and GABA secreting neurons there. The smaller peptide degrades synaptobrevin and blocks the release of the inhibitory glycine and GABA. This results in hyperreflexia, muscle spasms, and spastic paralysis. Even small amounts of the toxin can be lethal.

The incubation period of C. tetani ranges from 4-5 days up to 3 weeks. Muscle spasms usually involve the area of injury first and then the muscles of the jaw causing trismus or lockjaw. Gradually, other voluntary muscles become involved, resulting in tonic spasms. Any external stimulus can precipitate a tetanic generalized muscle spasm. The patient is conscious but may be in pain. Death usually results from respiratory paralysis.

Other options:

Option A : Botulinum toxin targets the presynaptic terminals of motor neurons and inhibit the release of acetylcholine. As a result, muscle paralysis occurs, leading to the temporary weakening or cessation of muscle activity in the affected area.

Option D: The venom of the black widow spider, specifically the component α-latrotoxin, has been found to induce the massive release of neurotransmitters, including acetylcholine, norepinephrine, dopamine, and others, from presynaptic nerve terminals rather than inhibition of nor-epinephrine.

Solution to Question 11:

Chronic granulomatous disease (CGD) is a rare inherited immune system disorder predisposing individuals to infection by catalase-positive organisms. It is mostly acquired as X-linked recessive seen in boys or very rarely autosomal recessive seen in girls.

The most common organism is S. aureus, but any catalase-positive microorganism may be involved. Other organisms frequently causing infections include Serratia marcescens, B. cepacia, Aspergillus, Candida albicans, Nocardia, and Salmonella. There may also be increased

susceptibility to mycobacteria, including the bacille Calmette-Gue■rin vaccine. Catalase-negative bacteria like streptococci and pneumococci are efficiently handled by the immune system.

This disease manifests itself as chronic suppurative granulomatous lesions of the skin and lymph nodes, along with hepatosplenomegaly, progressive lung infilterates and granulomatous septic osteomyelitis.

CGD is usually diagnosed in early childhood:

• Phagocytes cannot efficiently produce reactive oxygen species (ROS), which are crucial for destroying pathogens engulfed by these cells.

• Reduced hydrogen peroxide (H_2O_2) production seems to be the primary cause of the impaired bactericidal function.

• Failure of leucocyte degranulation after phagocytosis, leading to delayed granule rupture and faulty release of myeloperoxidase further contributes to ineffective bactericidal activity.

• Bacteria proliferate within the cells, shielded from antibodies and antibiotics due to their intracellular location, leading to the establishment of chronic suppurative infection.

• Humoral and cellular immune response is normal

Nitroblue-tetrazolium test: This is used as a screening method for the diagnosis of chronic granulomatous disease. Leucocytes of induviduals with CGD fail to reduce nitroblue tetrazolium in CGD, due to a defect in the NADPH oxidase complex, neutrophils are unable to perform this reduction, leading to a negative or deficient result in the NBT test.

Treatment typically involves long-term antibiotic and antifungal therapy to prevent and manage infections. In some cases, bone marrow transplantation may be considered a potential cure for CGD. Corticosteroids may be useful for the treatment of children with antral and urethral obstruction or severe granulomatous colitis.

Solution to Question 12:

The characteristic clinical manifestation of Parvovirus is febrile illness with an erythematous rash. Petechial rash is seen only in some patients.

Parvovirus B19 is the smallest single-stranded DNA virus belonging to the family Parvoviridae. It is a non-enveloped virus with an icosahedral capsid. The virus is primarily transmitted through respiratory secretions, such as saliva and respiratory droplets. It can also be transmitted vertically from mother to fetus during pregnancy, leading to congenital infection.

Clinical manifestations:

Erythema infectiosum (Fifth Disease): Characterized by a "slapped cheek" rash on the face, the most common manifestation of Parvovirus B19 infection. This is due to erythematous flushing of the cheeks. This is followed by a lacy, macular erythematous rash on the trunk and limbs. The rash typically resolves within 1-3 weeks but may recur with exposure to sunlight or heat.

Arthropathy: Transient arthritis, predominantly affecting the joints of the hands, wrists, knees, and ankles. Arthropathy typically occurs in adults, especially women, and may persist for several weeks to months.

Aplastic Crisis: In individuals with underlying hemolytic disorders, such as sickle cell disease or hereditary spherocytosis, Parvovirus B19 infection can precipitate a transient aplastic crisis. This results in a severe reduction in red blood cell production, leading to anaemia.

Hydrops fetalis: Vertical transmission of Parvovirus B19 during pregnancy can lead to severe complications in the fetus, including hydrops fetalis.

Hydrops fetalis is characterised by fetal oedema, ascites, pleural effusion, and pericardial effusion, often resulting in intrauterine death or neonatal complications.

Parvovirus B19 mainly attacks the erythroid progenitor cells and leads to transient arrest of erythropoiesis. However, immunocompromised individuals are at risk of persistent Parvovirus B19 infection which leads to chronic suppression of bone marrow, resulting in persistent anemia or pure red cell aplasia.

Diagnosing Parvovirus B19 infection is primarily based on clinical presentation and confirmed through serological testing, including detecting specific IgM antibodies. Polymerase chain reaction (PCR) assays can also detect viral DNA in blood or other tissues.

The following image shows a slapped cheek appearance seen in Erythema infectiosum (Fifth Disease):

Note: This is a best-out-of-four type of question where all 4 options are factually correct. Petechial rash is seen in several viral infections including Parvovirus B19 infection. However, the characteristic rash of this infection is erythematous. This makes option D the best possible answer.

Solution to Question 13:

The given clinical scenario of a patient with a history of renal transplant and complaints of watery diarrhoea along with a Kinyoun stain positive image of pink coloured, acid-fast oocyst with the size of 30x15 microns is suggestive of infection due to Cystoisospora.

Cystoisospora usually causes a self-limiting but protracted diarrhoea, lasting for several years, majorly in immunocompromised persons and particularly among the HIV infected. Human infection is acquired by ingestion of mature oocysts in food or drink. The parasites generally dwell in the epithelial cells of the small intestine .

Diagnosis involves identifying oocysts in fecal smears, which may sometimes be sparse and overlooked due to their transparent appearance in unstained films. They stain red using the cold acid-fast technique. Concentration methods like zinc sulfate or formol-ether techniques can be used.

The first line of management for isosporiasis is cotrimoxazole for 7-10 days. Patients who do not tolerate cotrimoxazole are treated with nitazoxanide or ciprofloxacin.

A comparison of these coccidian parasites is given in the table below:

Property	Cryptosporidium	Cyclospora	Isospora
Sporulated Oocyst	4-6 µm, round contains fo ur sporozoites.	8-12 µm, round contains 2 sporocysts each having t wo sporozoites.	23-36 µm, oval contains t wo sporocysts, each havin g four sporozoites
Acid-fast detection	Uniformly acid-fast	Variable acid-fast	Uniformly acid-fast
Sporulation of the ooc yst	Occurs inside the host cell	Occurs in soil	Occurs in soil
Diagnostic form	Sporulated cyst	Unsporulated cyst	Unsporulated cyst

Solution to Question 14:

The given clinical scenario of an 18-day-old child presenting with altered sensorium, neck rigidity, and vesicular lesions on the anterior abdomen and the forearm with the mother giving a history of painful genital ulcers, is suggestive of neonatal herpes due to herpes simplex virus (HSV)-2. CSF examination with multinucleate giant cells with CNS manifestations indicates neonatal meningitis to the DNA virus.

Painful genital ulcers are commonly seen in genital herpes and chancroid due to Hemophilus ducreyi ((ducreyi makes you cry).

Herpes infection can be caused by HSV-1 and HSV-2 viruses. HSV-1 causes orofacial lesions, i.e. herpes labialis, while HSV-2 mainly causes genital herpes which is characterised by vesiculo ulcerative lesions of the penis of the male or of the cervix, vulva, vagina, and perineum of the female.

The typical herpes lesions are thin-walled, umbilicated vesicles, the roof of which breaks down, leaving tiny superficial ulcers which heal without scarring. The most common site of cutaneous lesions is in the face, cheeks, chin, around the mouth and forehead. A cesarean section is performed in a case of active genital herpes at the time of labour in order to prevent neonatal herpes infection.

One of the following three disease patterns can be seen in infants infected during delivery or in the postpartum period: (1) disease limited to the skin, eyes, or mouth; (2) encephalitis with or without skin, eye, and mouth disease; or (3) disseminated infection affecting several organs, such

as the brain, lungs, liver, heart, and adrenal glands.

Infants suffering from encephalitis usually exhibit clinical symptoms that point to bacterial meningitis at 8- 17 days of life. These symptoms include seizures, poor feeding, lethargy, neck stiffness, irritability, and positive Kernig's sign. When left untreated, HSV encephalitis results in 50% of infant deaths and severe neurologic sequelae in the majority of survivors.

Acyclovir at a high dose of 60 mg/kg/day, administered intravenously every eight hours, should be the first line of treatment for any infant with a confirmed or suspected neonatal HSV infection.

Other options:

Coxsackie virus, Enterovirus and VZV are not attributed to painful genital ulcers and hence ruled out.

Solution to Question 15:

The above image shows Cryptococcus neoformans stained with India ink and its infection most commonly involves the lungs. Indian ink is a type of negative stain that is not picked up by the capsule, making it evident. Although Cryptococcosis usually presents clinically as chronic meningo-encephalitis, the primary infection always involves the lungs.

Cryptococcus is an opportunistic fungus that is found worldwide as a saprophyte in soil enriched with pigeon droppings.

Pathogenesis:

• Infection is initiated by inhalation of the yeast cells.

• The primary pulmonary infection may be asymptomatic or may mimic an influenza-like respiratory infection, often resolving spontaneously.

• In immunocompromised, the yeasts may multiply and disseminate to other parts of the body, but preferentially to the CNS, causing cryptococcal meningoencephalitis.

• Other common sites of dissemination include the skin, adrenals, bone, eye, and prostate gland.

Clinical findings:

• Chronic meningitis (Headache, nuchal rigidity, disorientation)

• CSF pressure, protein concentration and cell count are elevated

• Cutaneous ulcers, granulomata.

Diagnosis:

• Specimens - CSF, tissue, exudates, sputum, blood, urine, skin scrapings

• Microscopy - Gram staining shows gram-positive round budding yeast cells

• India ink preparation reveals capsulated budding yeast cells; capsules are prominent in this staining.

• Culture - Colonies on SDA at 37°C appear as smooth, mucoid and creamy white.

• Serology - Demonstration of cryptococcal antigen by latex agglutination test is a sensitive, specific and rapid method for identificcation of cryptococcal meningoencephalitis.

Treatment:

• Cryptococcosis without CNS involvement: Fluconazole is the drug of choice

• HIV patient with CNS involvement: Induction: Liposomal Amphotericin B with Flucytosine for at least 2 weeks

Maintenance: Oral Fluconazole

Other Options:

Option B: Cryptococcus neoformans is a round or ovoid budding yeast cell, 4-20 μm in diameter, with a prominent polysaccharide capsule.Although the fungus grows primarily as a yeast, hyphae are produced during the sexual phase and during a process called monokaryotic fruiting, which is also believed to involve sexual reproduction, but no pseudohyphae are produced during normal circumstances.

Option C: C.neoformans is frequently found in soils contaminated with avian excreta and can easily be recovered from shaded and humid soils contaminated with pigeon droppings. Hence, can be found in more than one reservoir other than man.

Option D: C.neoformans usually affects patients with advanced HIV infection and CD4+ T lymphocyte counts of as low as <200/μL .

Solution to Question 16:

S. agalactiae belonging to Group B streptococci are the most common cause of meningitis and neonatal sepsis followed by E. coli.

S. agalactiae is a capsulated Gram-positive coccus occurring in chains. It causes meningitis in the neonatal period. Infection is commonly acquired from the maternal vagina during birth. There are two clinical types seen in neonates:

• Early-onset type (more common) presents within a week of birth as meningitis or septicemia and is often fatal. Early-onset infection presents similarly to other types of neonatal sepsis, with common symptoms such as respiratory distress, lethargy, and hypotension. Almost all infants with this condition have bacteremia, while one-third to one-half also experience pneumonia , respiratory distress syndrome, and one-third develop meningitis.

• Late-onset type presents between the second to twelfth week of life commonly as meningitis, and in most cases is associated with a strain of capsular type III.

Group B streptococci can also lead to infections in non-pregnant immunocompromised adults. Common infections include sepsis, skin and soft tissue infections, respiratory tract infections, and urinary tract infections in women.

Identification :

A streptococcal isolate can be classified presumptively as GBS based on biochemical tests, including hydrolysis of sodium hippurate, hydrolysis of bile esculin, and bacitracin susceptibility.

The CAMP test (Christie, Atkins and Munch-Peterson) is used to identify group B β-hemolytic-streptococci based on their ability to form CAMP factor. A characteristic arrowhead zone of hemolysis is seen when Streptococcus agalactiae is inoculated perpendicular to

Staphylococcus aureus grown on blood agar. The group B streptococci hydrolyse sodium hippurate and give a positive CAMP test.

The image given below shows a positive CAMP test





Solution to Question 17:

The correct answer is option C: 1-d, 2-c, 3-a, 4-b

1. SARS COV -2	d. Single-stranded RNA
2. Rotavirus	c. Double-stranded RNA

1. SARS COV -2	d. Single-stranded R NA
3. Human Papilloma Vi rus	a. Double-stranded D NA
4. Parvovirus	b. Single-stranded D NA

Solution to Question 18:

After sputum sample collection, Mycobacterium is detected in sputum samples with N-acetyl cysteine (NAC) and NaOH where N-acetyl cysteine is used for liquefaction of sputum and NaOH kills the contaminating bacteria. Therefore in the above question assertion(A) is true but reason(R) is false.

Sputum microscopy is the most reliable single method in the diagnosis and control of tuberculosis. NALC (N-acetyl cysteine) with 2% NaOH is superior for detecting bacilli in microscopy.

Sample collection:

• Sputum Collection: Sputum yields abundant bacilli in caseation, less in lesions organised lesion not connected to airways. Best collected in the morning pre-meal; scanty samples may be pooled over 24 hours. Sampling over three days enhances detection.

• Laryngeal aspiration and bronchial washing is done in situations where sputum is not available

• In children who are unable to expectorate sputum, gastric lavage is an ideal method for sample collection.

Collected Sputum and non-sterile samples need initial pre-treatment to prevent microbial overgrowth during incubation. Processes such as liquefaction, decontamination, and concentration are essential to enhance the yield from the sputum samples. Hence, techniques preserving bacilli viability are necessary. Always follow Class II biosafety methods while performing any procedures related to sputum.

• Petroff's method: In this method, sputum is incubated with 4% sodium hydroxide at 37°C, shaken till clear (~20 mins), then centrifuged (3000 rpm, 20 mins). The sediment is neutralised with N/10 HCl for further use. Avoid excessive alkali exposure.

• NALC (N-acetyl cysteine) with 2% NaOH: This method is superior to Petroff's, NALC liquefies sputum while NaOH eliminates contaminating bacteria. The sample is then, neutralised with buffer and centrifugation follows. This method is compatible with automated culture systems."

The image given below shows a sputum smear stained with Ziehl-Neelsen stain showing Mycobacterium tuberculosis.



Solution to Question 19:

The presence of ulcerative eruptions along the lymphatics in a gardener from the sub-Himalayan region and the microscopic finding of flower-like clusters of conidia are suggestive of sporotrichosis caused by Sporothrix schenckii. It is a thermally dimorphic fungus that lives on vegetation.

The lesions occur following traumatic inoculation of the fungus. A granulomatous nodule is first formed at the site of inoculation and may progress to form a necrotic or ulcerative lesion. The infection can spread via lymphatic channels from the skin and subcutaneous tissue to the lymph nodes, resulting in the formation of secondary ulcers on the lymph nodes. This condition is known as lymphocutaneous sporotrichosis. This disease is commonly encountered among gardeners, forest workers, and manual labourers.

Other options:

Option A: Coccidiomycosis is caused by Coccidioides immitis which is a thermally dimorphic fungus. It occurs as a spherule in tissues and culture at 37°C and as the mycelial form in soil and culture at room temperature. They are endemic to the arid desert regions of the southwestern USA and California. Infection is acquired by inhaling dust containing arthrospores of the fungus.

Option B: Histoplasmosis is caused by Histoplasma capsulatum. It has tiny oval yeast cells with narrow-based budding within the macrophages with an underlying granulomatous response. This is a dimorphic fungus that exists in the yeast form in body tissues and cultures maintained at 37°C, and in the filamentous mold form in cultures at room temperature (25°C).

Option D: Blastomyces cause North American Blastomycosis/ Chicago disease. Pulmonary involvement is the most common form followed by skin. Histopathological staining shows thick-walled yeast, with single broad-based budding (figure 8 appearance).

The image given below is that of Blastomyces which is a thick-walled yeast, with single broad-based budding (figure 8 appearance).



Solution to Question 20:

The given scenario is indicative of hemolytic uremic syndrome, and E. Coli O157:H7 is the likely causative organism.

O157: H7 is a strain of E.coli called enterohemorrhagic E. coli (EHEC), which produces verocytotoxin or Shiga-like toxin. They produce two antigenic forms of the cytotoxic toxin known as Shiga-like toxin 1 and Shiga-like toxin 2. Shiga-like toxin 1 is identical to the Shiga toxin of Shigella dysenteriae type 1. E. coli O157:H7 has a small infective dose (around 100 bacilli) and spreads easily from child to child by the fecal-oral route.

It can cause mild non-bloody diarrhoea to hemorrhagic colitis, or a severe form of diarrhoea, with hemolytic uremic syndrome, a disease resulting in acute renal failure, microangiopathic hemolytic anemia, and thrombocytopenia. The most common cause of hemolytic uremic syndrome in the world is E.coli O157:H7. The most common cause of hemolytic uremic syndrome in India is Shigella dysenteriae type I.

EHEC (O157: H7) dysentery is commonly self-limiting, and management is conservative, with the mainstay being the correction of dehydration and electrolyte imbalances and other symptomatic supportive treatment such as blood transfusion in case of severe blood loss anaemia, but antibiotics are not to be used. Therefore, the treatment of HUS is primarily supportive care, which may include:

• Intravenous fluids to prevent dehydration

• Blood transfusions to correct anemia and thrombocytopenia

• Dialysis to treat kidney failure

In some cases, plasma exchange or eculizumab may be used to treat HUS. Plasma exchange is a procedure that removes harmful substances from the blood. Eculizumab is a medication that blocks the complement system, which is part of the body's immune system.

Other options:

Option A: Shigella flexneri produces Shigella enterotoxin (ShET). It enhances local vascular damage of the intestine, kidney, and brain. It is commonly implicated in reactive arthritis, also called Reiter syndrome (post-infectious arthritis).

Option B: Salmonella enteritidis is commonly implicated in osteomyelitis in patients with sickle cell anemia.

Option C: Aeromonas is implicated in inflammatory diarrhoea and does not present with fever, hematuria, and thrombocytopenia.

Solution to Question 21:

Nocardia is the isolated organism in this patient based on the given histopathological features.

Nocardia species appear as filamentous organisms with extensive hyphae-like branching. They are aerobic, gram-positive, catalase-positive, and urease-positive. They are weakly acid-fast because they stain with carbolfuchsin and retain when decolourized with 1–4% sulfuric acid instead of the stronger acid-alcohol decolorant.

Infections caused by Nocardia include cutaneous, subcutaneous, or systemic lesions. It may lead to local abscesses, cellulitis, or lymphocutaneous lesions. Systemic nocardiosis occurs more often in immunodeficient persons.

Infection follows traumatic inoculation of the organism into the subcutaneous tissue from soil or vegetable sources. They can present with a characteristic powdery substance that appears as gritty or necrotic material on the skin or within soft tissues. Accompanying this are multiple discharging sinuses, which are abnormal openings that form as the body attempts to expel infected material. These sinuses can drain purulent, often foul-smelling discharge typically associated with localised abscesses, and the discharge may be viscous, sometimes containing necrotic tissue.

Systemic nocardiosis usually caused by N.asteroides manifests primarily as pulmonary disease, pneumonia, lung abscess, or other lesions resembling tuberculosis. Extrapulmonary manifestation may involve the brain, kidneys, and other organs.

Diagnosis is by microscopy of sputum, bronchoalveolar lavage specimens, and tissue biopsy. Gram stain shows beaded gram-positive filamentous rods. In Modified ZN staining they are weakly acid-fast.

The drug of choice for nocardiosis is cotrimoxazole. In severe cases, 3rd generation cephalosporins/ carbapenems and amikacin are recommended.

The image given below is a smear of modified acid-fast Kinyoun staining showing filamentous red-stained partially acid-fast rods suggestive of Nocardia.



Other options:

Option A: Gram's staining of Bacillus anthrax shows bacilli arranged end to end in the form of long chains. This is called bamboo stick appearance.

Option B: Listeria monocytogenes is a short, non sporing, gram positive bacillus. It shows characteristic tumbling motility at 25°C, but is non motile at 37°C.



Option D: The image given below is a smear with Ziehl-Neelsen staining showing non-branching and acid-fast Mycobacterium tuberculosis bacilli.



Solution to Question 22:

PD-1 on T cell interacts with PD ligand 1 on APC is not true regarding naive T cell interaction with CD4 effector T-cells.

T-cell activation requires two signals that are delivered by APCs (Option A). They include:

• Antigens displayed by APCs in the form of peptides bound to MHC molecules, that are recognised by T-cell receptors

• Costimulatory signal (as it stimulates T cells in conjunction with antigen) provided by molecules on APCs that engage particular costimulatory receptors on T cells

CD80 (B7-1) and CD86 (B7-2) are the costimulatory molecules present on antigen-presenting cells (APCs). They bind to the CD28 protein present on T cells (Option B). This promotes the expansion of antigen-stimulated T cells and their differentiation into effector and memory cells. (Option D).



Solution to Question 23:

E-test is a method of antibiotic sensitivity testing.

E-test (epsilometer) is a diffusion and dilution method of testing antibiotic sensitivity. It is a quantitative technique for determining antimicrobial susceptibility and minimum inhibitory concentration (MIC).

An absorbent strip containing a predefined gradient (serial dilution) of antibiotic concentration immobilized along its length is placed on a lawn inoculum of a bacterium. An elliptical zone of inhibition is produced surrounding the strip. The antibiotic concentration at which the ellipse edge intersects the strip is taken as the MIC value.

The given image shows the E-test.



The antibiotic sensitivity tests are classified as follows:

1. Diffusion methods:

• Kirby–Bauer disk diffusion method

• Stokes disk diffusion metho

2. Dilution methods:

• Broth dilution

• Agar dilution

3. Diffusion and dilution method:

• E–test (Epsilometer)

4. Automated methods:

• VITEK 2

• Phoenix system

Other options:

Option B: Elek test is an in vitro immunoprecipitation (immunodiffusion) test used to determine whether or not a strain of Corynebacterium diphtheriae is toxigenic.



Option C: In latex agglutination tests, an antigen-specific antibody (either polyclonal or monoclonal) is fixed to latex beads. When the clinical specimen from the patient is added to a suspension of the latex beads, the antibodies bind to the antigens on the microorganism forming a lattice structure, and agglutination of the beads occurs. Latex agglutination tests are mainly used for the detection of carbohydrate antigens of encapsulated microorganisms.



Option D: The Christie-Atkins-Munch-Peterson (CAMP) test is used to identify group B, β-streptococci based on their ability to form CAMP factor, which causes synergistic hemolysis with β-hemolysin formed by Staphylococcus aureus. When Streptococcus agalactiae is inoculated perpendicular to Staphylococcus aureus grown on blood agar, a characteristic arrowhead zone of hemolysis is seen. The group B streptococci hydrolyze sodium hippurate and give a positive CAMP test.



Solution to Question 24:

Madurella mycetomatis is transmitted by walking barefoot in soil. It causes Eumycetoma.

Mycetoma is a chronic, granulomatous infection of the skin, subcutaneous tissues, fascia, and bone, which most often affects the foot or the hand. It can be divided into three types- eumycetomas, actinomycetomas, and botryomycosis.

Infection follows traumatic inoculation of the organism into the subcutaneous tissue from soil or vegetable sources. Within host tissues, the organisms develop to form compacted colonies (grains) 0.5–2 mm in diameter, the colour of which depends on the organism responsible. It results in tumefactions, deformities, and draining sinuses, discharging these grains or granules.

Laboratory Diagnosis:

A. Direct Examination: The presence of grains in pus collected from draining sinuses or in biopsy material is diagnostic.

B. Culture Samples should also be cultured, at both 25–30°C and 37°C, on brain-heart infusion agar or blood agar for actinomycetes and on Sabouraud agar (without cycloheximide) for fungi.

Treatment:

The management of eumycetoma is difficult, involving surgical debridement or excision and chemotherapy. Actinomycetoma, on the other hand, responds well to rifampicin in combination with sulfonamides or cotrimoxazole.

Curvularia, Coccidioides immitis, and Aspergillus fumigatus are not transmitted by walking barefoot in soil.

The image below shows a mycetoma of the foot:

Solution to Question 25:

A and D can be prevented by vaccination.

Cervical cancer is caused by Human papillomavirus (HPV). Currently, there are four vaccines against HPV approved for clinical use which are:

• Cervarix is a bivalent vaccine active against HPV types 16 and 18. Anogenital warts are commonly caused by subtypes 6 and 11. Thus, Cervarix does not confer protection against anogenital warts.

• Gardasil is quadrivalent and additionally protects against HPV 6 and 11.

• Gardasil-9 is nonavalent and protects against 9 strains.

• Cervavac is India's first indigenously prepared quadrivalent HPV (qHPV) vaccine against HPV strains 6, 11, 16, and 18.

Hepatitis B and C can cause Hepatocellular carcinoma. Hepatitis B vaccine is an example of a subunit vaccine made using HBsAg. The first dose (0.5 mL, intramuscular) is given at birth or as early as possible within 24 hours on the anterolateral side of the mid-thigh. Subsequent doses are given as a part of the pentavalent vaccine at 6 weeks, 10 weeks, and 14 weeks (can be given until one year of age).

Burkitt lymphoma and Merkel cell carcinoma cannot be prevented by vaccination.

Solution to Question 26:

Statements 2,4 are true about Cryptococcus.

Cryptococcus is an opportunistic fungus that is found worldwide as a saprophyte in soil enriched with pigeon droppings. C.gatti sources from Eucalyptus trees, not bird Guano. (Statement 3)

Pathogenesis:

• Infection is initiated by inhalation of the yeast cells.

• The primary pulmonary infection may be asymptomatic or may mimic an influenza-like respiratory infection, often resolving spontaneously.

• In immunocompromised, the yeasts may multiply and disseminate to other parts of the body, but preferentially to the CNS, causing cryptococcal meningoencephalitis.

• Other common dissemination sites include the skin, adrenals, bone, eye, and prostate gland.

• Extrapulmonary cryptococcosis is an AIDS-defining illness (Statement 2)

• It is seen when CD4 <200 cells/microliter (Statement 1)

Clinical findings:

• Chronic meningitis (Headache, nuchal rigidity, disorientation)

• CSF pressure, protein concentration, and cell count are elevated

• Cutaneous ulcers, granulomata.

Diagnosis:

• Specimens - CSF, tissue, exudates, sputum, blood, urine, skin scrapings

• Microscopy - Gram staining shows gram-positive, urease-positive (Statement 4) round budding yeast cells

• India ink preparation reveals capsulated budding yeast cells; capsules are prominent in this staining.

• Culture - Colonies on SDA at 37°C appear as smooth, mucoid and creamy white.

• Serology - Demonstration of cryptococcal antigen by latex agglutination test is a sensitive, specific and rapid method for identification of cryptococcal meningoencephalitis.

Treatment:

• Cryptococcosis without CNS involvement: Fluconazole is the drug of choice

• HIV patient with CNS involvement: Induction: Liposomal Amphotericin B with Flucytosine for at least 2 weeks

• Maintenance: Oral Fluconazole

Solution to Question 27:

Lowenstein-Jenson medium is used for M. tuberculosis.

Lowenstein-Jenson medium is a selective medium used for the culture of Mycobacterium tuberculosis which causes tuberculosis. The presence of malachite green imparts a green colour and makes Lowenstein Jenson medium a selective media by inhibiting other bacteria.

Components of LJ medium:

• Egg: Provide fatty acids and protein required for metabolism.

• Malachite green: Prevents growth of contaminants.

• Mineral salt solution: Buffer.

• Glycerol: Source of carbon, Favours growth of M. tuberculosis, but not M. Bovis.

• L-asparagine: Source of Nitrogen.

Colony characteristics on LJ Medium:

Mycobacterium tuberculosis colonies on LJ medium are rough, raised, irregular colonies that are creamy white, and buff-colored. (rough, tough, buff). They do not emulsify in a saline solution.

Culture media used to cultivate Mycobacterium tuberculosis:

• Solid culture media: Lowenstein–Jensen (A commonly used, egg-based, selective media).

• Liquid culture media: Dubos' media, Middle Brook's media. Virulent strains tend to form long serpentine cords in liquid media, while avirulent strains grow in a more dispersed manner.

• Automated Culture Media:

BACTEC 460: Uses radioisotopes

BacT/ALERT: A colorimetric method

BACTEC MGIT: A fluorometric detection method.

The culture medium shown in the image is the Lowenstein-Jenson.

Other options:

Option B: Blood agar is a differential medium as it helps in differentiating various organisms based on the degree of hemolysis observed.

• Alpha hemolysis- Partial lysis of red blood cells and haemoglobin

• Beta hemolysis- Complete lysis of the red blood cells and haemoglobin

• Gamma hemolysis- No hemolysis

Option C: Novy-MacNeal-Nicolle medium is used for isolating parasites like Leishmania and Trypanosoma cruzi.

Option D: Sabouraud dextrose agar is the traditional mycological medium, which contains glucose and modified peptones. It supports the growth of fungi while restricting bacterial growth due to the addition of antibacterials.

Solution to Question 28:

The given clinical scenario with complaints of non-bloody diarrhoea and dehydration in a child suggests rotavirus.

Rotavirus is the most common identifiable viral cause of diarrhoea in children below the age of 5 years and is most frequent between 6-24 months of age. The incubation period is 1-3 days. Most cases are asymptomatic or with mild symptoms like watery diarrhoea, vomiting, low-grade fever, and temporary lactose intolerance. Diagnosis can be made by rapid antigen detection of rotavirus in stool specimens.

Treatment is supportive, including hydration and correction of fluid and electrolyte imbalance. The main goals of treatment are avoiding and treating dehydration. Maintaining nutrition is a secondary goal.

Other options:

Option A: Trichuris trichiura causes rectal prolapse in children (characteristic feature), chronic dysentery, anemia, poor growth, and developmental defects. The disease may result either due to the mechanical effect of the worms or an allergic reaction to the parasite.

Option B: Shigella causes Shigellosis, which is a spectrum of symptoms consisting of 4 phases:

1. Incubation

2. Watery diarrhoea: Fever, watery diarrhoea, anorexia

3. Dysentry: Bloody mucopurulent stools

4. Post-infectious complication: Hemolytic-uremic syndrome, reactive arthritis, and toxic megacolon.

Option D: Yersinia enterocolitica causes diarrhoea with symptoms resembling appendicitis (pseudoappendicitis). The patients present with abdominal pain, which can be localised to the right iliac fossa, thus mimicking appendicitis.

Solution to Question 29:

i-b,ii-c,iii-d,iv-a is the correct match of the causative organism with the disease.

i. Klebsiella Granulomatosis	b.Donovanosis
ii. Bartonella bacilliformis	c. Oroya fever
iii. Treponema carateum	d. Pinta disease

i. Klebsiella Granulomatosis	b.Donovanosis
iv. Streptobacillus monoliformis	a. Rat-bite fever

Solution to Question 30:

Toxoid does not confer passive immunity but plays a role in active immunity.

Toxoid vaccines are a type of subunit vaccine prepared by detoxifying the toxins produced by organisms like tetanus bacilli. In general, they are highly efficacious and safe immunising agents.

Passive immunity is the resistance that is transmitted passively to a recipient in readymade/ preformed antibodies. Here, the recipient's immune system plays no active role, and there is no antigenic stimulus.

Types of passive immunity:

• Natural passive immunity: It is the immunity passively transferred from mother to baby. Maternal IgG antibodies (Option D) are transmitted through the placenta. The human colostrum is rich in lgA (Option C) antibodies and gives protection to the neonate.

• Artificial passive immunity: It is the resistance passively transferred to a recipient by the administration of antibodies. For example, antitoxins (Option B) like anti-tetanus serum and anti-diphtheria serum.

Solution to Question 31:

Autoclave is not a cold sterilization technique

Autoclaving is a form of moist heat sterilization that creates an increased temperature of steam under pressure. This method of sterilization is effective against spores also. Autoclaving is done for instruments that cannot withstand high temperatures of hot air oven or dry heat. Surgical instruments are sterilized at 121° C for 10-15 minutes. The given image shows an autoclave.

Cold sterilisation is a process in which sterilisation is carried out at low temperatures with the help of chemicals, radiation, membranes (filters) and all other means excluding high temperatures.

Cold sterilisation techniques:

1. Ionising Radiation:

• X-ray

• Gamma rays (Option D)

• Cosmic rays

2. Non-ionising Radiation:

• Infrared

• Ultraviolet (Option C)

Other options:

Option A: Plasma sterilisation is a new method of sterilisation for heat-sensitive articles. A radio frequency energy is applied to create an electromagnetic field. Into this, hydrogen peroxide vapours are introduced, which generate a state of plasma containing free radicals of hydrogen and oxygen. This state renders the articles sterile by denaturing all microorganisms. Arthroscopes, urethroscopes, etc., are sterilised by plasma sterilisation.

Solution to Question 32:

A high larval load can cause rectal prolapse is not true regarding Ascaris lumbricoides.

Ascaris lumbricoides (roundworm) reside in the small intestine and not the large intestine to cause rectal prolapse. Trichuris Trichura (whipworm) cause rectal prolapse.

Ascaris lumbricoides is an intestinal nematode that causes ascariasis. Infection occurs through food or water contaminated with nightsoil or human faeces having eggs of Ascaris that contain the infective rhabditiform larva. Man is the only natural host. There is no intermediate host. Most infected humans are asymptomatic. (Option A)

The life cycle is as follows:

• Adult worms live in the lumen of the small intestine. A female produces eggs, which are passed with the faeces.

• Unfertilized eggs in the stool may be ingested but are not infective. The infective form of Ascaris is the embryonated egg in the soil (Option D). Infection typically arises from consuming food or water contaminated with embryonated eggs. These eggs harbour the rhabditiform larva.

• After infective eggs are swallowed, the larvae hatch.

• The larvae invade the intestinal mucosa and are carried through the portal system, then systemic circulation to the lungs.

• The larvae mature further in the lungs (10 to 14 days), penetrate the alveolar walls, ascend the bronchial tree to the throat, and are swallowed. This may present with asthma-like symptoms (Option B).

• Upon reaching the small intestine, they develop into adult worms. Between 2 and 3 months are required from ingestion of the infective eggs to oviposition by the adult female.

• Adult worms can live for 1 to 2 years.

• Adult worms can be occasionally detected in the stool or sputum of the patient. A definitive diagnosis is made by demonstration of eggs in faeces. Fertilised eggs are always bile-stained.

Effective and safe drugs for the treatment of ascariasis are Pyrantel pamoate 11 mg/kg once, Albendazole 400 mg once, Mebendazole 100 mg twice daily for 3 days, Ivermectin 150-200 mcg/kg once.

The image below shows the life cycle of Ascaris lumbricoides:



Solution to Question 33:

Triclabendazole is the drug of choice for fasciola hepatica.

Triclabendazole belongs to the benzimidazole group. The primary mechanism of action of benzimidazoles is by inhibition of microtubule polymerization. Benzimidazoles bind to beta-tubulin and cause inhibition of microtubule polymerization.

Drugs belonging to the benzimidazole group are:

• Albendazole

• Mebendazole

• Triclabendazole

• Thiabendazole

Mechanism of action:

• Inhibition of mitochondrial fumarate reductase

• Reduced glucose transport

• Uncoupling of oxidative phosphorylation

Other options:

Option B: Praziquantel is the drug of choice for all trematodes except Fasciola hepatica and Fasciola gigantica.

Option C: Mebendazole is used in ascariasis, trichuriasis, hookworm, pinworm infections, and other helminthic infections.

Option D: Albendazole is the drug of choice for the treatment of hydatid disease and cysticercosis. It is also used in the treatment of pinworm, hookworm, ascariasis, trichuriasis, and strongyloidiasis.

Solution to Question 34:

The gram-positive cocci in chains is Streptococcus pyogenes (Group A β-hemolytic Streptococcus), which causes Necrotizing fasciitis.

Necrotising fasciitis is a rapidly progressive necrosis of the subcutaneous tissues and fascia. The source of the infection is either the skin following trauma or a surgical incision. Diabetes mellitus is the most common comorbidity associated with the condition.

Local symptoms and signs are pain, erythema, edema, warmth, crepitus, blisters, bullae, greyish drainage (‘dishwater pus’), necrosis, and gangrene. Systemic features are fever, tachycardia, tachypnea, shock, coagulopathy, and multiorgan failure.

Treatment involves appropriate intravenous antibiotics with urgent radical surgical debridement.

The image given below is of Streptococcus pyogenes, which is a gram-positive cocci in chains.



Other options:

Option B: Gram-positive cocci arranged in clusters is suggestive of Staphylococcus aureus. It causes toxic shock syndrome and staphylococcal scalded skin syndrome.

Option C: Enterococcus are catalase-negative, bile-esculin-positive, gram-positive cocci that grow in chains and are characterised by their ability to grow in the presence of 6.5% NaCl (salt tolerance). They are implicated in Infective endocarditis.

Option D: Micrococcus is a non-pathogenic skin commensal organism that appears as gram-positive tetrads on microscopy.

Solution to Question 35:

The patient presenting with abdominal pain and a growing mass in the right hypochondrium has a hydatid cyst of the liver. Aspirate of the cyst shows Echinococcus granulosus (with hooks).

Echinococcus granulosus is the most common causative agent in hydatid cysts, while Echinococcus multilocularis and Echinococcus ligartus account for a small number of cases.

Dogs are the definitive hosts of E. granulosus. Sheep are the usual intermediate hosts, but humans are accidental intermediate hosts. Humans are end-stage hosts of the parasite since there is no human-to-human transmission.

Approximately 75% of hydatid cysts are located in the right lobe of the liver and are solitary. The clinical presentation of a hydatid cyst is largely asymptomatic until complications occur. The most common presenting symptoms are abdominal pain, dyspepsia, and vomiting. The most frequent sign is hepatomegaly.

The life cycle of Echinococcus is depicted in the image below:



The hydatid cyst wall consists of 3 layers:

• Pericyst- the outer dense fibrous capsule that is formed as a result of host inflammatory response.

• Ectocyst - the intermediate layer consisting of acellular, chitinous, laminated hyaline material.

• Endocyst - the inner germinal layer that is cellular and consists of several nuclei embedded in protoplasmic mass and is extremely thin. It is the vital layer of the cyst, and it is the site of asexual reproduction, giving rise to brood capsules with scolices.

The brood capsules are cellular masses in the hydatid fluid, which consists of scoleces. In a definitive host, these scoleces develop into adult tapeworms, while in the intermediate hosts, these develop into new hydatid cysts. The brood capsules are attached to the endocyst by a pedicle or stalk.

The brood capsules may mature and detach from the mother cyst and become the daughter cysts, which are replicas of the mother cysts. The freed brood capsules and scoleces present in the hydatid fluid are called hydatid sand.



Other options:

Option B: Hymenolepis nana (dwarf tapeworm) is the smallest tapeworm. Humans are the definitive host. It has no intermediate host. It occurs more commonly in children, and the patient acquires this infection by ingesting food or water contaminated with eggs. The egg attaches to the intestine and develops into the larva and adult worm. The adult worm resides in the small intestine. Patients are usually asymptomatic. However, in heavy infections, they present with nausea, anorexia, abdominal pain, diarrhoea, and irritability. Pruritus may also be present due to an allergic response.

Option C: Taenia solium is a parasitic tapeworm transmitted through undercooked pork, causing intestinal infection (taeniasis) or more severe cyst formation in tissues, including the brain (cysticercosis). It's prevalent in regions with poor sanitation and where pork is consumed.

Option D: Taenia saginata is a parasitic tapeworm transmitted through undercooked beef containing cysticerci ("measly" beef). Humans are the definitive host, and cows are the intermediate hosts. Taenia solium and Taenia saginata eggs are morphologically indistinguishable.

Solution to Question 36:

Chloroquine is not given in uncomplicated falciparum malaria.

NVBDCP guidelines for the treatment of uncomplicated falciparum cases:

• North-eastern states: Artemether-lumefantrine-based ACT (Option C)

• Rest of the country: Artesunate-sulfadoxine+pyrimethamine (Option D)

The artemisinin-based combination therapy (ACT) recommended by NVBDCP in the northeastern states is a combination of artemether and lumefantrine. In the northeastern states (Arunachal Pradesh, Assam, Manipur, Meghalaya, Mizoram, Nagaland, and Tripura), the presently recommended ACT for P. falciparum malaria is a fixed-dose combination of artemether-lumefantrine due to the recent reports of late treatment failures with the current

combination of AS+SP.

Primaquine 0.75 mg/kg on day 2 is given as a single dose in uncomplicated falciparum malaria.(Option B)

Solution to Question 37:

Bacteriological loops used for inoculation are sterilised by heating till metal turns red hot.

Sterilisation of bacteriological inoculating loops, the tip of spatulas and forceps is done by placing them over a Bunsen flame until they get red hot. Inoculation loops carrying infective material should be dipped in a disinfectant before flaming to prevent spattering.

Other options:

Option B: Disinfection of slides and cover slips are done by passing over the flame 3 times.

Option C: Autoclaving is a method of heat sterilisation. The autoclave is a device that uses steam under pressure to sterilise linen, instruments, laboratory ware, media, pharmaceutical products, aqueous solutions, surgical instruments, cardiac catheters, and implants.

Option D: In Microwave irradiation, microorganisms are killed using high-frequency microwaves. This cannot be used for animal or human body parts, metal items or toxic or radioactive material.

Microbiology NEET 2024

Question 1:

Which of the following is used for syndromic diagnosis of infected cases like meningitis?

a) Real time PCR

b) Nested PCR

c) Reverse transcriptase PCR

d) Multiplex PCR

Question 2:

A patient receiving chemotherapy develops pneumonia with skin lesions. Microscopy of specimen from skin lesions showed broad based budding with a thick capsule. What is the likely diagnosis?

a) Blastomycosis

b) Histoplasmosis

c) Coccidioidomycosis

d) Candidiasis

Question 3:

A 30-year-old man came to the OPD with a febrile illness of 2 days, associated with abdominal tenderness, malaise and diarrhoea. He had a fever with bradycardia and BP was noted as 110/70 mmHg. He lives in a crowded neighborhood with poor sanitation where many others have reported similar complaints recently. Which of the following investigations would be the most appropriate choice for this patient?

a) Stool culture

b) Widal test

c) Urine culture

d) Blood culture

Question 4:

A child is brought with complaints of abdominal pain, diarrhoea and perianal itching for the last 3 days. The mother said she noticed a white thread-like structure in the child's night clothes. She collected the worm and brought it to the clinician. A perianal swab was collected. Microscopy of the swab and the worm are shown in the images. What is the likely causative organism?

a) Enterobius vermicularis

b) Ascaris lumbricoides

c) Ancylostoma duodenale

d) Strongyloides stercoralis

Question 5:

A male patient comes to you with a history of UTI, the cause of which is suspected to be bacteria. Which of the following is used by the bacteria for adherence?

a) Cytoplasmic membrane

b) Cilia

c) Fimbriae

d) Flagella

Question 6:

A thalassemia patient has received multiple blood transfusions. Aplastic crisis can occur in this patient due to infection from?

a) Parvovirus

b) Epstein barr virus

c) Cytomegalovirus

d) Herpes simplex virus

Question 7:

A 45-year-old farmer comes with complaints of high-grade fever with chills, retro-orbital pain and conjunctival suffusion. On examination, he has severe calf tenderness. Lab examination shows hypokalemia and neutropenia. What is the likely diagnosis?

a) Chikungunya

b) Leptospirosis

c) Dengue

d) Malaria

Question 8:

A patient with a history of consumption of oysters comes with complaints of nausea, vomiting and crampy abdominal pain. The culture of the organism showed the Kanagawa phenomenon. What is the causative organism?

a) Vibrio cholera

b) Salmonella typhi

c) Vibrio parahemolyticus

d) E coli

Question 9:

A man is having petechial rashes with a previous history of meningitis. He was vaccinated at the age of 5 years. Now, Brudzinski and Kernig's signs were positive. Microscopy showed gram negative diplococci. What is the likely causative organism?

a) Hemophilus influenzae

b) Neisseria Meningitidis

c) Cytomegalovirus

d) Klebsiella

Question 10:

A patient who has a central line on long-term IV antibiotics develops a fever. Blood cultures show Staphylococcus epidermidis along with biofilm formation. Which of the following describes its features of antibiotic resistance?

a) Biofilm increased susceptibility to antibiotics

b) Biofilm offers resistance so catheter is to be removed

c) Biofilm does not affect susceptibility to antibiotics

d) Biofilm can be washed off from the catheter

Question 11:

Laboratory investigations in a patient with desert rheumatism showed dimorphic and thermal fungi with arthrospores. What is the causative organism of desert rheumatism?

a) Histoplasma capsulatum

b) Paracoccidioides brasiliensis

c) Blastomyces dermatitidis

d) Coccidioides immitis

Question 12:

A farmer from South East India presents with fever, respiratory distress and neurological deficits. The vector of the causative organism is shown below. what is the associated organism?

a) Ebola virus

b) Zika virus

c) Malaria

d) Nipah virus

Question 13:

A 21-year-old female complains of fever, headache and vomiting from the past 3 days. She has been living in the army recruitment camp for the last three weeks. On examination, she has neck stiffness and a maculopapular rash was noted over the trunk and extremities. Which of the following is false about the disease she's likely suffering from?

a) Carriers of this disease can be treated with Rifampicin

b) It can lead to adrenal hemorrhage

c) Terminal complement deficiency increases the risk of this disease

d) Metronidazole is the drug of choice

Question 14:

A young sexually active female presented with a symptomatic UTI. Urinalysis and cultures showed novobiocin-resistant coagulase negative staphylococcus with a count of 10^4 cells/HPF. What is true about this condition?

a) Asymptomatic bacteriuria

b) Test must be repeated with a fresh specimen

c) Diagnose and treat as urinary tract infection

d) Normal skin commensal grown on culture that needed to proved

Question 15:

A group of hostel boys consumed hamburgers from a local street food shop following which they developed abdominal pain, diarrhoea, vomiting, and fever. They were diagnosed with hemolytic uremic syndrome (HUS). Escherichia coli O157:H7 was detected in the stool samples and the ham. What does H represent in H7?

a) Fimbriae

b) Capsule

c) Flagella

d) Lipopolysaccharide

Question 16:

An elderly female with a central line undergoes surgery. On postoperative day 3, she develops fever and tachycardia. A culture from the tip of the central line revealed brown buff colonies. Microscopy showed gram-positive organisms that reproduce by budding. What is the likely causative organism?

a) Staphylococcus epidermidis

b) Streptococcus

c) Candida

d) Aspergillus

Question 17:

A 22-year-old farmer develops an ulcer on his foot with a white powdery substance and multiple discharging sinuses. Microscopy reveals a gram-positive organism with radiating filaments. What is the most likely organism isolated from the discharging sinus?

a) Nocardia

b) Staphylococcus aureus

c) Mucormycosis

d) Sporothrix schenckii

Question 18:

A middle-aged man with a history of unprotected sexual intercourse presents with complaints of painful urination and itching. On examination, Gram-negative intracellular cocci are observed. What is the most likely causative organism?



a) Staphylococcus aureus

b) Neisseria gonorrhoeae

c) Proteus vulgaris

d) Chlamydia trachomatis

Question 19:

A patient diagnosed with HIV presents with complaints of recurrent diarrhoea. The stool sample reveals only larvae and no ova. Which of the following is the most probable causative organism?

a) Ancylostoma

b) Ascaris

c) Strongyloides

d) Enterobius

Question 20:

A young man who has just returned from the USA develops fever, cough, and breathlessness. A chest radiograph shows bilateral hilar lymphadenopathy and mediastinal lymphadenopathy. Culture on Sabouraud Dextrose Agar (SDA) reveals white to light brown puffy colonies. Lactophenol blue staining shows the findings as depicted in the image. What is the most likely causative organism?



a) Cryptococcosis

b) Histoplasmosis

c) Blastomycosis

d) Mucormycosis

Question 21:

A middle-aged man presents with complaints of itching around the perianal region and unilateral lower limb lymphadenopathy. Larvae of approximately 200 microns in size were identified, and these larvae lacked nuclei at their tail tips. What is the most likely causative organism?

a) Wuchereria bancrofti

b) Brugia malayi

c) Brugia timori

d) Mansonella perstans

Question 22:

A middle-aged man presented to the OPD with dysuria and purulent urethral discharge for the last 3 days. Gram stain of the discharge is shown below. He admits to having unprotected sexual intercourse one week ago. What is the most appropriate treatment?



a) Tab. Acyclovir 500 mg TDS for 1 week

b) Tab. Doxycycline 100 mg BD for 14 days

c) Tab. Cefixime 400 mg Stat

d) Inj. Benzathaine Penicillin 2.4 MU IM Stat

Question 23:

A young man presents with burning and painful micturition. Microscopy showed gram-negative intracellular kidney bean-shaped organisms. Which of the following culture media is used to isolate the organism?

a) Blood agar

b) Thayer-Martin media

c) MacConkey agar

d) TCBS agar

Question 24:

A 35-year-old man presented with high fever, bleeding gums, myalgia, and hepatosplenomegaly. His TLC is 3100, Hb- 11 g/dl, hematocrit- 55%, and platelets - 25,000 cells. What is the most important prognostic factor in this condition?

a) Hematocrit

b) Total Leukocyte Count

c) Hemoglobin

d) Platelets

Answer Key

Question No.	Correct Option
1	d
2	a
3	d
4	a
5	c
6	a
7	b
8	c
9	b
10	b
11	d
12	d
13	d
14	c
15	c
16	c
17	a
18	b
19	c
20	b
21	a
22	c
23	b
24	a

Detailed Explanations

Solution to Question 1:

Multiplex PCR can be used for syndromic diagnosis of meningitis.

A syndromic diagnosis is a method of diagnosing infectious diseases that uses a single test to screen for multiple pathogens that could be causing a specific set of symptoms.

Multiplex PCR (Polymerase Chain Reaction) is a variation of the standard PCR technique that allows simultaneous amplification of multiple target sequences in a single reaction. It uses more than one set of primers (specific to different target sequences) to amplify different DNA fragments at the same time.

Other options:

Option A: Real-time PCR helps quantify the PCR and gives an estimate of the pathogen DNA load in the specimen. It uses fluorescent dyes or specific DNA probes. It is also known as qPCR (Quantitative PCR).

Option B: Nested PCR involves the sequential use of two primer sets. The first primer set is used for amplification and the amplicon obtained is then used as a target sequence for second amplification. This ensures specificity. Used in GeneXpert to detect Mycobacterium TB and resistance to rifampicin.

Option C: Reverse transcriptase PCR: The enzyme reverse transcriptase (RT) is used to synthesize a complimentary copy of DNA (cDNA) from RNA and this cDNA is then used as a template for PCR. Unlike conventional PCR which can only amplify DNA, this one can even amplify RNA. It is used to detect H1N1 from Nasopharyngeal swab.

Solution to Question 2:

In this case of an immunocompromised patient with pneumonia and skin lesions, the microscopy of specimens from skin lesions showing broad-based budding with a thick capsule suggests the diagnosis of Blastomycosis caused by Blastomyces dermitidis.

Blastomycosis is a chronic infection, characterized by the formation of suppurative and granulomatous lesions, especially, in the lungs and skin.

Blastomycosis predominantly affects individuals in regions where the fungus Blastomyces is endemic, such as the Midwest and southeastern United States. It is most common in North America and so is also called North American Blastomycosis.

The disease has a broad demographic impact, occurring in both immunocompetent and immunocompromised individuals. However, those with impaired immune systems—such as patients with AIDS, those on glucocorticoid therapy, or those receiving TNF-α inhibitors—are at higher risk for severe or disseminated disease involving the skin, bone, or central nervous system. Healthy individuals may experience a range of symptoms from mild pulmonary illness to more severe forms of pneumonia. It is more commonly observed in males aged 30-50 years, possibly due to occupational or recreational exposures.

Blastomycosis can present in three forms:

• Pulmonary disease:

• Presents with cough, sputum production, chest pain, and fever

• Hilar lymphadenopathy may be present

• Focal or diffuse consolidation, miliary lesions, or abscess formation may occur.

• Resembles TB and Histoplasmosis

• Cutaneous disease:

• Involvement of face and exposed parts of the skin

• Skin lesion: ulcerated verrucous granulomas with an advancing border and central scarring. The border is filled with microabscesses and has a sharp, sloping edge.

• Disseminated disease:

• Multiple skin lesions are the most common presentation

• Highly fatal

• The genitourinary tract (especially the prostate) is frequently involved

• Bone involvement resembles chronic osteomyelitis.

It can be diagnosed based on culture findings. Characteristic findings include the presence of thick-walled round yeast cells with a single broad-based bud when cultured at 37°C. (figure of 8 appearance)

Treatment of localised mild/moderate disease includes Itraconazole (6 months) and for disseminated/ severe disease- amphotericin B.

The image given below shows the skin lesions and microscopy of broad-based budding by Blastomyces dermitidis.



Other options

Option B: Histoplasmosis is caused by Histoplasma capsulatum. It has tiny oval yeast cells with narrow-based budding within the macrophages with an underlying granulomatous response. This

is a dimorphic fungus that exists in the yeast form in body tissues and cultures maintained at 37°C, and in the filamentous mould form in cultures at room temperature (25°C).

Option C: Coccidiomycosis is caused by Coccidioides immitis which is a thermally dimorphic fungus. It occurs as a spherule in tissues and culture at 37°C and as the mycelial form in soil and culture at room temperature. They are endemic to the arid desert regions of the southwestern USA and California. Infection is acquired by inhaling dust containing arthrospores of the fungus.

Option D: Candidiasis is caused by candida species which are yeast-like fungi. They exist as yeast and pseudohyphae. It appears as budding gram-positive cells on wet films and smears. Mycelial forms suggest colonization. Candida albicans form chlamydospores on corn meal agar.

Solution to Question 3:

This patient's history of fever with relative bradycardia, abdominal tenderness, and similar cases in the neighbourhood suggests enteric fever. It is caused by Salmonella typhi or paratyphi. Out of the given options, blood culture is the most sensitive test in this patient.

Diagnosis of enteric fever:

Blood culture: It is more sensitive during 1st week than in the subsequent weeks. It rapidly becomes negative on treatment with antibiotics.

Antibody testing: WIDAL test to detect rising titres of H and O agglutinins against salmonella typhi and paratyphi. Agglutinins appear by the end of the first week and rise steadily up to the 3rd or 4th week, after which they disappear.

Stool culture: It yields positive results from the 3rd week onwards.

Urine culture: Results may be positive after the 2nd week.

Bone marrow culture remains positive for 5 days even after the start of antibiotics and is the most sensitive test in the first week. The culture of duodenal aspirate may yield positive results when bone marrow culture is inconclusive.

Relative bradycardia is seen when patients have a lower heart rate than expected for a given rise in body temperature. (Faget's sign)

Infectious causes of relative bradycardia include:

• Salmonella typhi

• Francisella tularensis

• Coxiella burnetii (Q fever)

• Leptospira interrogans

• Legionella pneumophila

• Rickettsia spp.

Noninfectious causes include:

• Drug fever

• Beta-blocker use

- Central nervous system lesions
- Malignant lymphoma
- Factitious fever

Solution to Question 4:

The likely causative organism for the child's abdominal pain, diarrhoea, perianal itching, and the white thread-like worm found in the night clothes is Enterobius vermicularis, as evidenced by the worm's pointed end and wing-like cuticular expansions (cervical alae) shown in the left image, and the colourless, non-bile stained, planoconvex eggs with an outer albuminous layer that stick to each other and other objects.

Enterobius vermicularis completes its entire life cycle in the human host. They are oviparous. Man acquires infection by ingesting embryonated eggs containing larva through contaminated fingers and by autoinfection.

At night, the gravid female worm comes out through the anus and lays its sticky eggs on the perianal skin. This leads to pruritus and the patient scratches the affected perianal area. The worm can crawl into the vulva and vagina causing irritation and a mucoid discharge. It can also cause chronic salpingitis, cervicitis, peritonitis, recurrent urinary tract infections, and appendicitis.

Since pinworms are rarely released in faeces, the faecal examination is not useful in the diagnosis.

The NIH swab (National Institutes of Health, USA) consists of a glass rod with a piece of transparent cellophane attached at one end with a rubber band. The cellophane part is used for swabbing by rolling over the perianal area. The cellophane piece is spread over a glass side and examined microscopically.

The image given below shows the NIH swab.

Another method of demonstrating eggs is the scotch tape method. Here, the tape is firmly pressed against the anal margin, covering all sides, and transferred to a glass slide. A drop of toluene is added and examined under the microscope.

The below image shows the scotch tape method.

The adult worms may sometimes be noticed on the surface of stools, can be found crawling out of the anus, or may be detected in stools collected after an enema.

Frequent hand washing, fingernail cleaning, regular bathing, and frequent washing of night clothes, and bed linen should be done. Infection can be treated with pyrantel pamoate, albendazole, or mebendazole as single-dose therapy.

Other options:

Option B: Ascaris lumbricoides causes infection when the egg containing the infective rhabditiform larva is swallowed. The eggs hatch in the duodenum to release the rhabditiform larvae, which then migrate to the liver via the portal vein, heart, and then to the lungs, pass up the respiratory tract, and are swallowed. They finally develop into adults in the small intestine. Here, the gravid females lay eggs, which are shed via the feces, and the cycle repeats. Definitive diagnosis is by demonstration of eggs in feces. The life cycle is shown below.



Option C: Ancylostoma duodenale or hookworm infects a person by direct penetration of the skin by filariform larvae. They develop into adult worms in the small intestine. Segmented eggs are passed in feces and these develop into filariform larvae in the soil. They can cause severe local itching, cutaneous larva migrans, and respiratory symptoms. Adult hookworm disease is characterized by anaemia, epigastric pain, dyspepsia, vomiting, and reddish black stools. Diagnosis is by demonstration of eggs or larvae in feces. The life cycle is shown below.



Option D: Strongyloides stercoralis infection is by direct penetration of the skin by the filariform larva or by autoinfection. The larvae enter lay eggs in the small intestine mucosa. The eggs hatch and release rhabditiform larvae which pass through feces into the soil and start another cycle. External autoinfection occurs when the larvae penetrate the perianal skin. Internal autoinfection occurs in immunocompromised patients. Here, the larvae mature to the infective form in the bowel itself. Diagnosis is by demonstration of rhabditiform larvae in feces.



Solution to Question 5:

Bacteria utilise fimbriae for adherence.

Fimbriae, also called pili are hair-like surface appendage that projects from the cell surface as straight filaments. Fimbriae are classified based on the function of ordinary pili and sex pili.

Functions of pili:

• Bacterial adhesion of symbiotic and pathogenic bacteria to host cells using adhesins.

• Resistance to phagocytosis.

• Antigenic pilin proteins are responsible for the immune response.

• Twitching motility to bacteria.

• Bacterial conjugation through sex pili, in which transfer of genetic material occurs.

Other options:

Option A: The cytoplasmic membrane, or plasma membrane, is a fundamental structure of all cells, including bacteria. It acts as a selective barrier, regulating the entry and exit of substances.

Option B: In organisms like protozoa, cilia are used for locomotion and feeding. These organisms use their cilia to move through water and to sweep food particles towards their oral groove.

Option D: Flagella are thread-like appendages composed of protein. They are the organs of locomotion but motility via pili is completely different from flagellar motion because of the helical arrangement of pilin molecules.

Solution to Question 6:

Aplastic crisis in this patient is triggered by parvovirus infection.

Parvovirus infects and kills the red cell precursors. This creates a transient arrest of erythropoiesis i.e. aplastic crisis as the life span of RBCs is shortened. The immune response clears the virus in 1-2 weeks. It may be seen in patients with underlying chronic hemolytic anaemia like sickle-cell disease, thalassemia, hereditary spherocytosis, and RBC enzyme deficiencies like pyruvate kinase deficiency. Patients with aplastic crisis appear ill with fever, malaise, and paleness with tachycardia and tachypnea due to profound anaemia. The rash may or may not be present.

Parvovirus B19 infection occurs usually in children 5–10 years causing erythema infectiosum. It is also known as the fifth disease. It begins as a prominent erythema of the cheeks- slapped cheek disease, spreading to the trunk and limbs, followed by lymphadenopathy and arthralgia. Fever and mild constitutional symptoms may accompany the rash.

Other options:

Option B: Epstein Barr virus affects adolescents and younger adults more frequently but may occur from childhood to old age. It is transmitted primarily through close contact with oropharyngeal secretions. The incubation period of EBV infectious mononucleosis is 4-6 weeks. Clinically, patients may present with, fever, malaise, nausea, anorexia, lymphadenopathy- bilateral and symmetrical, pharyngitis, tonsillar enlargement, hepatomegaly and splenomegaly. A drug-induced rash may develop on using beta-lactam. antibiotics.

Option C: Cytomegalovirus causes mononucleosis in healthy individuals and disseminated diseases like leukopenia, hepatitis, colitis, pneumonitis, esophagitis, gastritis, and retinitis in immunocompromised patients.

Option D: Herpes simplex virus type 1 which belongs to the subfamily alpha viruses causes Herpes labialis (also known as cold sores). HSV type 1 (human herpes virus type 1 or HHY type 1) is isolated from lesions in and around the mouth and is transmitted by direct contact or droplet spread from cases or carriers. The most common site is the face- around the mouth, on the cheeks, chin, or the forehead.

Solution to Question 7:

The most likely diagnosis based on the clinical vignette of high-grade fever with chills, retro-orbital pain, conjunctival suffusion, severe calf tenderness, and laboratory findings of hypokalemia and neutropenia is suggestive of leptospirosis.

Leptospirosis is a zoonosis caused by spirochetes of the genus Leptospira. Patients with mild disease may present with flu-like symptoms of sudden onset, fever with chills, headache, abdominal pain, conjunctival suffusion, and myalgia. The muscle pain usually affects the calves, abdomen, and back. Severe cases of leptospirosis, which can progress rapidly and have a high fatality rate, are associated with Weil's syndrome, characterized by triad of:

• Hemorrhage

• Jaundice

• Acute kidney injury

Jaundice, when present, can be profound, and physical examination may reveal an enlarged, tender liver. Temperature-pulse dissociation or relative bradycardia may also be observed.

The image below shows icterus and conjunctival suffusion in a patient with leptospirosis:

Leptospira are actively motile organisms and are best seen using a dark-field microscope. They are thin, tightly coiled, and flexible spirochetes with one end forming a hook. Diagnosis is by microscopic agglutination test.

Transmission occurs through cuts or breaks in the skin and mucous membranes (conjunctival and oral mucosa). Several outbreaks have been reported that resulted from exposure to floodwaters contaminated by urine from infected animals. The organism enters the bloodstream and lodges in the parenchymatous organs (liver and kidneys) leading to haemorrhage and necrosis of the tissue causing dysfunction of the organs. The progression and lifecycle of Leptospira are shown below.





Treatment for leptospirosis is based on the severity of the disease. In mild cases, oral doxycycline, ampicillin, or amoxicillin is preferred and in severe cases, intravenous penicillin, ampicillin, or ceftriaxone is commonly used.

Other options:

Option A: Chikungunya is a viral disease that presents with saddle-back fever, conjunctival suffusion, and photophobia. Migratory arthralgia, especially involving small joints may also be seen. A maculopapular or petechial rash is frequently seen on the trunk.

Option C: Dengue hemorrhagic fever is a form of severe dengue with bleeding manifestations. Evidence of plasma leakage and thrombocytopenia is also generally seen.

Option D: Malaria is a parasitic infection caused by the Plasmodium species transmitted by the Anopheles mosquito. It is characterized by many nonspecific symptoms, including but not limited to tachycardia, tachypnea, chills, malaise, fatigue, diaphoresis (sweating), and headache. The fever of malaria is characterized by periodicity depending on the infective Plasmodium species. The

fever returns every other day for Plasmodium vivax, Plasmodium ovale, and Plasmodium falciparum, and every three days for P. malariae.

Solution to Question 8:

Kanagawa phenomenon is shown by Vibrio parahemolyticus.

On Wagatsuma agar (blood agar with 2% salt), pathogenic strains of V. parahemolyticus are hemolytic, while non-pathogenic strains are non-hemolytic. This is due to a heat-stable hemolysin. They produce green-coloured colonies on the TCBS agar as they do not ferment sucrose.



On staining with Giemsa stain or methylene blue (Wayson's stain), the bacteria has bipolar staining with two ends densely stained and the central area clear. This is the characteristic 'safety-pin appearance' seen in Vibrio parahemolyticus.

Other options:

Option A: Vibrio cholerae is a facultative anaerobe, which produces yellow-colored colonies on thiosulfate citrate bile salts sucrose agar (TCBS). It is a selective and differential medium used for the culture of Vibrio species.

Option B: Wilson and Blair's medium is selective for Salmonella species. It is a type of selective and indicator media. Salmonella typhi reduces sulphite to sulphide in the presence of glucose and produces jet-black colonies with a metallic sheen due to the production of H_2S on Wilson and Blair bismuth sulphite medium.

Option D: E. coli is a gram-negative, flagellated rod. It is the most frequent cause of acute complicated urinary tract infections. It is a lactose fermenter and hence, grows on MacConkey medium to produce pink colonies.

Solution to Question 9:

The clinical vignette is suggestive of fulminant meningococcemia. The organism causing this condition is Neisseria meningitidis.

Neisseria meningitidis is an aerobic, gram-negative, nonmotile diplococcus that is catalase and oxidase-positive. It oxidizes glucose and maltose but not lactose or sucrose. The bacteria typically spread from the nasopharynx to the meninges via the perineural sheath of the olfactory nerve following an initial flu-like infection.

Meningococcemia often starts with acute fever, chills, malaise, and a petechial rash. As the disease advances, symptoms of meningitis, such as severe headache, vomiting, photophobia, confusion, and a stiff neck, may develop, potentially progressing to coma within a few hours. In its fulminant form, meningococcemia presents with high fever and a hemorrhagic rash and can lead to severe complications like disseminated intravascular coagulation, circulatory collapse, and Waterhouse-Friderichsen syndrome, characterized by bilateral adrenal gland necrosis and subsequent adrenal failure.

Meningitis can rapidly progress to fulminant meningococcemia, evolving from acute illness to severe disease in hours. Classic meningitis symptoms include fever, headache, neck stiffness, and decreased consciousness. Nuchal rigidity, along with Kernig's and Brudzinski's signs, may indicate meningeal irritation, though their sensitivity can vary. Seizures, occurring in 15-40% of patients, may result from complications like raised intracranial pressure, leading to severe outcomes such as cerebral herniation. The rash associated with meningococcemia evolves quickly from a diffuse maculopapular appearance to petechiae on the trunk, extremities, mucous membranes, and sometimes palms and soles.

Other options:

Option A: Haemophilus influenzae is a gram-negative, non-motile, and non-sporing coccobacillus. H.influenzae causes meningitis, a potentially fatal disease that is most common in children of two months to 3 years old. It also causes laryngo-epiglottitis, pneumonia, suppurative lesions, and bronchitis.

Option C: Cytomegalovirus infection is the most common congenital infection. The clinical manifestations are hepatosplenomegaly, petechial rashes, jaundice, chorioretinitis, and microcephaly. One of the long-term sequelae is sensorineural hearing loss

Option D: Klebsiella pneumonia causes fatal pneumonia, which occurs in middle-aged or older persons with risk factors such as alcoholism, chronic bronchopulmonary disease or diabetes mellitus. Gram stain and culture are used for diagnosis.

Solution to Question 10:

In a patient with a central line on long-term IV antibiotics along with culture showing biofilm formation, the correct measure is to remove the catheter as biofilm offers resistance.

Biofilms are structured microcolonies of organisms that are enclosed in a self-produced glycocalyx matrix. Bacteria within biofilms are usually metabolically less active.

Biofilms are aggregates of bacteria that are enclosed in an exopolysaccharide matrix that is produced by there bacteria itself. The biofilms are attached to solid surfaces and form a slimy layer over them. Organisms that produce biofilm include:

• Staphylococcus epidermidis on medical devices and implants

• Streptococcus viridians on heart valves

• Pseudomonas aeruginosa lung infections

• Staphylococcus aureus on catheters

• Legionella pneumophila colonization of hospital water systems

The exopolysaccharide helps in protection against the host's immunity. It acts diffusion barrier for antimicrobials and makes the organism relatively insusceptible to drug action(Option A and C). Bacteria embedded in biofilms frequently become metabolically less active or dormant. It is thus harder to eradicate these organisms with antibiotics.

These bacteria within biofilms also produce small molecules, such as homoserine lactones. These molecules are taken up by adjacent bacteria and function as a communication system called Quorum Sensing and help turn on certain genes at a particular time.

Domboro's solution, containing 0.25% acetic acid, helps prevent biofilm formation but is not effective at removing established biofilm. Biofilm, once established, adheres strongly to surfaces and is resistant to simple washing or flushing(option D). More aggressive methods, such as mechanical debridement or specialized treatments, are required to remove it.

Solution to Question 11:

Coccidioidomycosis, commonly known as Valley fever or desert rheumatism, is a mammalian fungal disease caused by Coccidioides immitis or Coccidioides posadasii.

The primary infection is often respiratory, with symptoms resembling bronchitis or pneumonia that resolve over a few weeks. The classic triad of coccidioidomycosis, known as 'desert rheumatism', includes the combination of fever, joint pains, and erythema nodosum.

Fluconazole and intravenous amphotericin B are used in progressive or disseminated disease, or immunocompromised individuals.Itraconazole or ketoconazole may be used for milder disease. Fluconazole is the preferred medication for coccidioidal meningitis, due to its penetration into CSF. Intrathecal or intraventricular amphotericin B therapy is used if infection persists after fluconazole treatment. The image showing mature spherules of coccidioides is given below.



Other options

Option A: Histoplasma causes histoplasmosis/ Darling's disease/ reticuloendotheliosis. It has tiny oval yeast cells with narrow-based budding within the macrophages with an underlying granulomatous response. This is a dimorphic fungus that exists in the yeast form in body tissues and cultures maintained at 37°C, and in the filamentous mould form in cultures at room temperature (25°C).

Option B: Paracoccidioidomycosis, also known as Brazilian blastomycosis, South American blastomycosis,Lutz-Splendore-de Almeida disease and paracoccidioidal granuloma, is a fungal infection caused by the fungus Paracoccidioides brasiliensis.

Option C: Blastomycosis is a chronic infection caused by a dimorphic fungus called Blastomyces dermatitidis. It is characterized by the formation of suppurative and granulomatous lesions in the lungs and skin. It is most common in North America and is also called North American Blastomycosis.

Solution to Question 12:

The given clinical scenario along with the vector bat is suggestive of infection from the nipah virus.

Nipah virus is an RNA virus that belongs to the family Paramyxoviridae, genus Henipavirus. It is a zoonotic virus that shows both animal-to-human (bats, pigs) and human-to-human transmission. The natural hosts are fruit bats of the Pteropodidae family.

The clinical manifestations include asymptomatic subclinical infection, acute respiratory infection, and fatal encephalitis. There is no treatment or vaccine available for either people or animals. The primary treatment for humans is supportive care. Nipah virus is on the WHO list of Blueprint priority diseases.

The life cycle of nipah virus is shown below:

Other options

Option A: Ebola virus belongs to the family Filoviridae and is transmitted by African fruit bats. Human-to-human transmission through blood or body fluids can occur. It causes fever, abdominal pain, diarrhoea, and haemorrhage

Option B: Zika virus is transmitted to human beings by the bite of an infected Aedes mosquito. It can also be transmitted trans-placentally, causing microcephaly and other severe fetal brain defects.

Option C: Malaria is a parasitic infection caused by the plasmodium species and is transmitted by the female anopheles mosquito. Symptoms include fever, chills, and myalgia with periodicity differing for each species. The fever returns every two days for P. ovale, P. vivax, and P. falciparum and every three days for P. malariae.

Solution to Question 13:

The clinical features are suggestive of meningococcal meningitis and metronidazole is not the drug of choice. The drug of choice is a third-generation cephalosporin such as ceftriaxone. Chloramphenicol is used in patients with penicillin allergy.

It is caused by Neisseria meningitidis, a gram-negative, oxidase-positive, catalase-positive diplococci. Outbreaks usually occur in mass gatherings such as in army camps or hostel dorms.

The risk of infection is increased in patients with terminal complement deficiency, hypogammaglobulinemia and hyposplenism. This is because opsonization by complement is a major mechanism by which the bacteria is cleared from the blood.

Characteristic clinical findings include:

• Maculopapular or purpuric rash

• Shock due to capillary leak

• Bilateral adrenal hemorrhage (Waterhouse–Friderichsen syndrome)

Rifampicin, ciprofloxacin and cephalosporins can be used to treat carriers.

Solution to Question 14:

A young sexually active female presenting with a symptomatic UTI and urinalysis and cultures showing novobiocin-resistant coagulase-negative staphylococcus with a count of 10^4 cells/HPF should be diagnosed as a urinary tract infection and should be treated.

Staphylococcus saprophyticus is a gram-positive, coagulase-negative staphylococcus. It is non-pigmented, novobiocin resistant, and non-hemolytic on blood agar. It is a leading cause of cystitis in women. It is a part of the normal flora of human skin and periurethral area and thus causes cystitis in sexually active women.

Other options

Option A: A colony count of >10^5/mL of a clean catch sample is considered significant bacteriuria with no clinical symptoms of UTI. In men, a count of 100,000 CFU/ml is sufficient to diagnose a UTI, while for women a UTI is diagnosed when the count is 100,000 CFU/ml in two consecutive midstream urine samples.

Option B: As the patient already has symptoms of UTI and the culture is showing a significant count redoing the test is not required.

Option D: Though staphylococcus saprophyticus is a normal commensal of skin since the patient already has symptomatic UTI with a proven culture,it need not be proved again.

Solution to Question 15:

H in E coli O157:H7 represents flagellar antigen.

O represents somatic antigen.

E. Coli O157:H7 is an enterohemorrhagic bacterial strain that causes hemorrhagic diarrhoea and cramping in the abdomen. It is transmitted through the fecal-oral route after consumption of contaminated, undercooked liquids and foods.

E. Coli O157:H7 uses the fimbriae to adhere to the microvilli of intestinal epithelial cells. When bacteria attach to an epithelial layer, they rearrange the cytoskeletal actin, which flattens the microvilli and interferes with normal intestinal cell function. This is known as an effacing lesion.

Three days after exposure to the contaminated food specimen, diarrhoea—often the hemorrhagic kind- occurs due to exposure to the Shiga toxin. The patients will experience bloody diarrhoea after three days of symptoms. After seven days, the symptoms of hemorrhagic diarrhoea usually go away on their own for 85% of patients; the remaining 15%, who are usually children, will experience systemic manifestations, most commonly HUS.

EHEC are gram-negative bacilli which are lactose fermenting. Unlike most other E. coli serotypes, O157: H7 does not use sorbitol and produces clear colonies on the Sorbitol MacConkey agar, whereas other serotypes produce pink colonies. O157: H7 is a strain of E.coli called enterohemorrhagic E. coli (EHEC) which produces verocytotoxin or Shiga-like-toxin. Shiga toxin

acts by inactivating the 60s ribosomal subunit thereby inhibiting protein synthesis. It is absorbed from the inflamed gastrointestinal mucosa into the circulation and causes endothelial dysfunction, and causes platelet activation and aggregation. Children and older adults are at the highest risk. It causes mild non-bloody diarrhea to hemorrhagic colitis, or a severe form of dysentery, with hemolytic uremic syndrome, a disease resulting in acute renal failure, microangiopathic hemolytic anemia, and thrombocytopenia.

Solution to Question 16:

The given clinical scenario along with microscopy showing gram-positive budding yeasts cells and brown buff colonies is suggestive of invasive candidiasis.

Candida species are yeast-like fungi that can cause bloodstream infections, particularly in patients with central lines or other invasive devices. It is a common cause of central line-associated bloodstream infections (CLABSI) and frequently presents with fever and signs of systemic infection..

The most common species implicated in candidal infections are C. albicans, C. parapsilosis, and C. glabrata. Clinically, infection with candida can present as cutaneous & mucosal candidiasis, systemic candidiasis, or rarely, as chronic mucocutaneous candidiasis.

Cutaneous and mucosal infections manifest as thrush affecting the tongue, lips, gums, and palate. It is seen as a patchy to confluent whitish pseudomembranous lesion and is commonly associated with AIDS. Candida may also invade the vaginal mucosa leading to vulvovaginitis, and the nails, and around the nail plate leading to onychomycosis.

Morphologically, they are yeast-like fungi. They exist as yeast and pseudohyphae (Shown in the image below). It appears as budding gram-positive cells on wet films and smears. Mycelial forms suggest colonization. Candida albicans form chlamydospores on corn meal agar.



When colonies of Candida albicans are inoculated in serum, the yeast cells start to form elongations called germ tubes within 2 hours, when kept at 37 °C. The image below

shows germ tube formation, also known as the Reynolds Braude phenomenon, seen in Candida albicans.

Other options:

Option A: Staphylococcus epidermidis is a gram-positive cocci that commonly colonizes the skin and can cause infections related to indwelling devices, such as central lines. However, it does not reproduce by budding, and the description of brown buff colonies does not fit.

Option B: Streptococcus species are gram-positive cocci that are typically found in chains. They do not reproduce by budding, and the brown buff colonies seen in the culture do not align with the typical appearance of Streptococcus in culture.

Option D: Aspergillus is a mold that forms septate hyphae, not yeast-like cells, and does not reproduce by budding. Additionally, Aspergillus forms colonies with a different characteristic appearance, usually described as greenish or fuzzy, rather than brown buff.

Solution to Question 17:

The given clinical scenario of multiple discharging sinuses and aspirate showing filamentous gram-positive bacteria are suggestive of Nocardia.

Nocardia species appear as filamentous organisms with extensive hyphae-like branching. They are aerobic, gram-positive, catalase-positive, and urease-positive. They are weakly acid-fast because they stain with carbolfuchsin and retain when decolourized with 1–4% sulfuric acid instead of the stronger acid-alcohol decolorant.

Infections caused by Nocardia include cutaneous, subcutaneous, or systemic lesions. It may lead to local abscesses, cellulitis, or lymphocutaneous lesions. Systemic nocardiosis occurs more often in immunodeficient persons.

They can present with a characteristic powdery substance that appears as gritty or necrotic material on the skin or within soft tissues. Accompanying this are multiple discharging sinuses, which are abnormal openings that form as the body attempts to expel infected material; these

sinuses can drain purulent, often foul-smelling discharge typically associated with localized abscesses, and the discharge may be viscous, sometimes containing necrotic tissue.

Systemic nocardiosis usually caused by N.asteroides manifests primarily as pulmonary disease, pneumonia, lung abscess, or other lesions resembling tuberculosis. Extrapulmonary manifestation may involve the brain, kidneys, and other organs.

The image given below is a smear of modified acid-fast Kinyoun staining showing filamentous red-stained partially acid-fast rods suggestive of Nocardia.



Other options:

Option B: Staphylococcus aureus is a common cause of skin and soft tissue infections, but it appears as gram-positive cocci in clusters, not as branching filaments. It would not typically be associated with discharging sinuses and white powdery discharge.

Option C: Mucormycosis is a fungal infection that typically affects immunocompromised individuals, particularly those with diabetes or undergoing chemotherapy. It presents with necrotizing infections but does not exhibit branching filaments in microscopy and is not associated with white powdery discharge from sinuses.

Option D: Sporothrix schenckii is a dimorphic fungus that causes sporotrichosis, typically following exposure to soil or plants. It can cause skin ulcers, but it appears as cigar-shaped yeast under microscopy, not as gram-positive filamentous bacteria.

Solution to Question 18:

The above clinical scenario along with the Gram stain showing gram■negative intracellular diplococci within phagocytes is suggestive of infection by Neisseria gonorrhea.

Gonorrhea is a sexually transmitted disease caused by Neisseria gonorrhea. It uses pilli in adhesion to the urethra or other mucosal surfaces. The cocci penetrate the intracellular spaces to reach the subepithelial tissue by the third day of infection. The incubation period is 2-8 days.

The major symptoms include vaginal discharge, dyspareunia, and dysuria, usually without urinary frequency or urgency. The specific symptoms seen in males and females are as follows:

In female patients:

About half the female patients remain asymptomatic. But, patients typically present with urethritis, cervical discharge, pruritus, dysuria, and intermenstrual bleeding.

The ectocervix and vagina are not usually affected in adults as the stratified squamous epithelium is resistant to infection by the cocci. The acidic pH of vaginal secretions is also protective. The principal sites of invasion for gonococcal infection are the columnar and transitional epithelium.

Pelvic inflammatory disease and salpingitis may lead to sterility. Other complications are tubo-ovarian abscesses and ectopic pregnancies. Rarely peritonitis may develop with perihepatic inflammation (Fitz Hugh Curtis syndrome). The patient presents with right upper quadrant pain and tenderness with abnormal liver function tests.

In male patients :

• Acute urethritis

• Genital discharge: It is mucopurulent that can be described as that resembling the flow of seed (semen).

• Strictures

• Water-can perineum: The infection may spread to the periurethral tissues causing abscesses and multiple discharging sinuses.

Lab diagnosis:

The discharge is collected and subjected to Gram's stain which reveals gram■negative intracellular diplococci within phagocytes. It is non-sporulating, non-motile and aerobic but may grow anaerobically also. It acidifies glucose but not maltose in serum sugars. Stuart's transport medium is used. Cultures can be obtained from chocolate agar or Mueller-Hinton agar. Specifically, Thayer-Martin agar is used to isolate Neisseria gonorrheae.

Treatment:

According to CDC guidelines (2021), treatment of uncomplicated gonococcal infections is as follows:

• Weight <150 kg – Ceftriaxone 500 mg intramuscular (IM) in a single dose

• Weight ≥150 kg – Ceftriaxone 1 g intramuscular (IM) in a single dose

Concurrent chlamydial infection must also be treated unless it has been excluded by molecular testing. First-line drugs used are as follows:

• Oral azithromycin 1g single dose

• Oral doxycycline 100 mg BD for 7 days (non-pregnant patients only)

The sexual partners of the patient must be treated to reduce the risk of recurrence. Expedited partner therapy refers to the index patient delivering therapy to one's own partners. However, this type of empirical therapy ignores the importance of direct counseling.

Other options:

Option A: Staphylococcus aureus is a gram-positive bacterium that can cause various infections, but it is not typically associated with sexually transmitted infections (STIs) or intracellular gram-negative cocci.

Option C: Proteus vulgaris is a gram-negative bacillus and is not typically associated with sexually transmitted infections. It is more often linked to urinary tract infections but would not appear as gram-negative intracellular cocci.

Option D: Chlamydia trachomatis is another common cause of sexually transmitted infections. It is an obligate intracellular bacterium that is difficult to visualize using standard Gram staining.

Solution to Question 19:

The presence of recurrent diarrhea in HIV patient along with the presence of only larvae in stool is suggestive of infection by Strongyloides stercoralis.

Strongyloides stercoralis is a parasitic nematode known for causing chronic diarrhea, particularly in immunocompromised individuals such as those with HIV. In this case, only larvae (and not ova) are seen in the stool, which is characteristic of Strongyloides infection due to its autoinfective cycle, where the larvae hatch inside the intestine and are passed in the stool.

Strongyloidiasis:

Strongyloides are ovoviviparous. The transmission of infection occurs through penetration of the skin by filariform larvae in the soil. Then, they migrate to the small intestine. The larvae moult twice and become adult female worms in the small intestine. The eggs laid in the mucosa of the intestine hatch immediately releasing the rhabditiform larva (L1 stage). These rhabditiform larvae in the bowel can develop directly into filariform larvae that penetrate the colonic wall or perianal skin and enter the circulation to repeat the migration. This establishes ongoing internal reinfection.

The life cycle of strongyloides is shown below.



It is generally an asymptomatic condition. There may be cutaneous, pulmonary, and intestinal manifestations in some cases. Pulmonary manifestations occur when the larva escapes from the pulmonary capillaries into the alveoli leading to bronchopneumonia, chronic bronchitis, and asthmatic symptoms.

The term larva currens (meaning 'racing larvae') has been applied to the rapidly progressing linear or serpiginous urticarial tracks caused by migrating filariform larvae of Strongyloides stercoralis. These often follow autoinfection and start perianally. Disseminated Strongyloidiasis is seen in immunodeficient individuals.

Larva currens is shown in the image below.

Visceral larva migrans is caused by the migration of larvae of nonhuman species of nematodes that infect via the oral route.

Etiological agents causing visceral larva migrans:

• Human nematode: Ascaris lumbricoides, Strongyloides stercoralis.

• Zoophilic nematode: Toxocara canis, toxocara cati, Angiostrongylus cantoniensis, Angiostrongylus costaricensis, Anisakis.

• Non human nematode: Filarial species, Brugia pahangi, Brugia pate

Laboratory diagnosis of strongyloidiasis:

• Eosinophilia and raised IgE levels.

• Microscopic examination of the direct wet mount of the stool demonstrating rhabditiform larvae forms the definitive diagnosis

• Stool culture and serological tests are also done

Treatment: Ivermectin is the preferred agent in the management of strongyloidiasis.

Other options:

Option A: Ancylostoma presents with creeping eruption. Cutaneous larva migrans or creeping eruption is caused by nematode larvae that are infected by skin penetration.

Option B: Ascaris is a large intestinal worm, and its ova are typically seen in the stool, not larvae. It is not commonly associated with chronic diarrhoea, especially in immunocompromised patients.

Option D: Enterobius or pinworm infection presents as itching (pruritus) and restless sleep (due to nocturnal perianal or perineal pruritus)

Solution to Question 20:

The image shows the tuberculate macroconidia, with finger-like projections which, along with the history, is a characteristic feature of Histoplasma.

Histoplasmosis (Darling's disease) is a granulomatous disease caused by a dimorphic fungus, Histoplasma capsulatum. It is endemic in the USA, in the states bordering the Ohio River valley and the lower Mississippi River. In India, it is reported from the region of West Bengal, along the Ganga River. The fungus inhabits humid and acidic soil that contains a large amount of bird or bat droppings.

H. capsulatum is transmitted by inhalation of spores (i.e. microconidia) which usually circulate in the air after the contaminated soil is disturbed. After the spores enter into the lungs, they are engulfed inside the alveolar macrophages and then transform into yeast forms.

Clinically, the classical histoplasmosis ranges from asymptomatic infection (in immunocompetent people) to life-threatening illness seen in people with low CMI. The various clinical types include:

• Pulmonary histoplasmosis - Acute form presents with mild flu-like illness, pulmonary infiltrates in the chest X-ray with hilar or mediastinal lymphadenopathy. Chronic cavitary histoplasmosis may be seen in smokers with underlying structural lung disease.

• Mucocutaneous histoplasmosis - Skin and oral mucosal lesions may develop secondary to pulmonary infection. Oral lesions are particularly seen in Indian patients.

• Disseminated histoplasmosis - It develops if CMI is very low (e.g. untreated HIV-infected people or following organ transplantation). The common sites affected are bone marrow, spleen, liver, eyes, and adrenal glands.

Diagnosis :

Direct microscopy - Histopathological staining of the specimens reveals tiny oval yeast cells (2–4 μm size) with narrow-based budding within the macrophages with an underlying granulomatous response.



Culture - It is the gold standard method of diagnosis. Specimens should be inoculated onto media such as SDA, blood agar, and BHI agar in duplicate and incubated simultaneously at 25°C and 37°C.

At 25°C it forms a mycelial phase and produces white to buff brown colonies that consist of two types of conidia or spores - tuberculate macroconidia (with typical thick walls and finger-like projections which is a characteristic feature) and microconidia(smaller, thin, and smooth-walled).

At 37°C it gets converted into yeast form (creamy white colonies), which is best developed in special media like Kelley's media.

Treatment :

Liposomal amphotericin B is the antifungal agent of choice in acute pulmonary and disseminated histoplasmosis. Itraconazole is recommended for chronic cavitary pulmonary histoplasmosis.

Other options :

Option A - Cryptococcosis: Cryptococcus neoformans typically presents as a lung infection or meningitis in immunocompromised individuals. On SDA, Cryptococcus forms mucoid colonies, and Lactophenol blue would show encapsulated yeast, which differs from the presentation in this case.

Option C - Blastomyces causes North American Blastomycosis/ Chicago disease. Pulmonary involvement is the most common form followed by skin. Histopathological staining shows thick-walled yeast, with single broad-based budding (figure 8 appearance).



Option D: Mucormycosis typically affects immunocompromised patients and presents with invasive disease, often causing necrotizing infections in the sinuses or lungs. The fungal hyphae seen on Lactophenol blue are broad, non-septate, and ribbon-like, which does not match the culture description here.

Solution to Question 21:

The given clinical presentation of perianal itching and unilateral lower limb lymphadenopathy is suggestive of Wuchereria bancrofti.

It is a major cause of lymphatic filariasis, often presenting with lymphadenopathy and lower limb swelling. Its microfilariae are approximately 200-300 microns long and are characterized by the absence of nuclei at the tail tip. Clinical features include massive eosinophilia, hepatosplenomegaly, pulmonary manifestations like dry nocturnal cough, asthmatic wheezing, and dyspnea. It can also cause arthritis, glomerulonephritis, thrombophlebitis, etc. The classical features seen in lymphatic filariasis are usually absent.

Elephantiasis refers to the painful swelling of the legs and genital organs seen in late-stage disease.

Image showing elephantiasis-

Investigations:

Demonstration of the microfilaria in peripheral blood, chylous urine, hydrocele fluid, etc is done. The specimen of choice is peripheral blood. Night blood samples between 10 pm and 4 am are preferred. This is because the microfilaria shows nocturnal periodicity.

DEC provocation test: In this test, a small dose of DEC is given 20-50 minutes before sample collection so that the microfilaria appears in the blood even during the daytime.

The image below shows the life cycle of Wuchereria bancrofti.



Solution to Question 22:

This clinical scenario along with the microscopic findings of gram■negative intracellular diplococci indicates infection with Neisseria gonorrhoeae. The treatment of choice for the infection among the given options would be Tab. Cefixime 400 mg Stat.

Gonorrhea is a disease acquired by sexual contact. The first step in infection is the adhesion of gonococci to the urethra or other mucosal surfaces. The incubation period is 2-8 days. The major symptoms include purulent urethral discharge and dysuria.

In males, the disease starts as acute urethritis presenting with purulent discharge. The patients may also have penile lymphangitis when the soft tissue inflammation is extensive. This infection may extend along the urethra to involve the prostate, seminal vesicles, and epididymis.

In females, the initial infection involves the cervix uteri and urethra which may extend to Bartholin glands, endometrium, and fallopian tubes. Pelvic inflammatory disease and salpingitis may lead to sterility. Rarely peritonitis may develop with perihepatic inflammation (Fitz-Hugh-Curtis syndrome).

The discharge is collected and subjected to Gram stain, revealing gram■negative intracellular diplococci within phagocytes. Stuart's transport medium is used. Cultures can be obtained from chocolate agar or Mueller-Hinton agar. Specifically, Thayer-Martin agar is used to isolate N. gonorrhoeae.

Single-agent therapy with ceftriaxone is preferred for the treatment of gonococcal infections.

According to CDC guidelines (2021), treatment of uncomplicated gonococcal infections is as follows:

• Weight <150 kg – Ceftriaxone 500 mg intramuscular (IM) in a single dose

• Weight ≥150 kg – Ceftriaxone 1 g IM in a single dose

The doses of ceftriaxone are higher than the previous guidelines as the gonococcal minimum inhibitory concentrations (MICs) are rising worldwide. The British BASHH guidelines also

recommend a high dose of 1g. Other cephalosporins like Cefixime can be given orally.

Concurrent chlamydial infection must also be treated unless it has been excluded by molecular testing.

First-line drug administered is

• Oral doxycycline 100 mg BD for 7 days

As per Suraksha clinic, syndromic management of Urethral discharge is done by Gray kit which contains Tab. Azithromycin 1gm stat and Tab. Cefixime 400mg stat. Partner treatment is also required.

Other options:

Option A: Acyclovir is given for viral infections. It has no role in gonococcal urethritis

Option B: Doxycycline is given alongside Ceftriaxone or cefixime to concurrently treat chlamydial infection along with gonorrhoea.

Option D: Benzathine Penicillin was once a treatment option for gonorrhoea, but it is no longer recommended due to widespread resistance.

Solution to Question 23:

The clinical scenario is suggestive of gonococcal urethritis caused by Neisseria gonorrhoeae. Thayer-Martin and modified Thayer-Martin media are used for the culture of this organism.

Gonorrhoea may manifest as a mucopurulent genital discharge which can be described as resembling the 'flow of seed' (semen). The discharge is collected and subjected to Gram's stain which reveals gram■negative intracellular diplococci within phagocytes. Stuart's media is used as transport media and Thayer Martin media is used for culture. Thayer-Martin medium contains vancomycin, colistin and nystatin.

Gonorrhoea is a sexually transmitted disease where infection occurs due to the adhesion of gonococci to the urethra or other mucosal surfaces with the help of pili. By the third day of infection, the cocci have penetrated intracellular spaces and reached subepithelial tissue. The incubation period ranges from 2 to 8 days. The main symptoms are urethral discharge and dysuria.

In females: The infection begins in the cervix uteri and urethra and may spread to the Bartholin glands, endometrium, and fallopian tubes. The vaginal mucosa is usually unaffected in adults because the stratified squamous epithelium is resistant to cocci infection and also because vaginal secretions have an acidic pH. Rarely peritonitis may develop with perihepatic inflammation (Fitz Hugh Curtis syndrome).

In males: The disease starts as acute urethritis. This infection has the potential to spread down the urethra and into the prostate, seminal vesicles, and epididymis. Chronic urethritis may lead to stricture formation. Infections in the periurethral tissues can lead to abscesses and multiple discharging sinuses (watercan perineum)

Treatment: Ceftriaxone (250 mg, IM) single-dose + azithromycin 1 g PO, single-dose, or doxycycline 100 mg PO BD for 7 days.

The image below shows the Gram-stain of a urethral exudate showing typical intracellular gram-negative diplococci and pleomorphic extracellular gram-negative organism



Solution to Question 24:

The given clinical history of a patient with a fever, myalgia, gum bleeding, hepatosplenomegaly, and laboratory investigations showing anaemia, leukopenia, raised hematocrit and low platelets is suggestive of dengue with warning signs and hematocrit is the most important prognostic factor in this patient.

Hematocrit is the percentage of red blood cells in a given blood volume, and a rising hematocrit is a sign of hemoconcentration due to plasma leakage. As plasma leakage signifies the severity of the disease, hematocrit is monitored to assess prognosis.

Dengue is an arbovirus disease transmitted by Aedes aegypti and causes a clinical spectrum:

• Probable dengue: fever with myalgia, nausea, vomiting, leukopenia or a positive tourniquet test.

• Dengue with warning signs: abdominal pain, tenderness, persistent vomiting, fluid accumulation (pleural effusion, ascites), mucosal bleed, hepatomegaly(>2cm), rapid rise in hematocrit and fall in platelet count

• Severe dengue: severe plasma leakage causing dengue shock syndrome, respiratory distress, severe hemorrhagic manifestations, severe organ involvement (AST or ALT >1000U/L, CNS manifestations, AKI, pancreatitis, DIC)

Other options:

Options B and C- Although dengue causes anaemia and leukopenia, they do not determine the prognosis of the disease.

Option D- Reduced platelet count is a warning sign but does not correlate with plasma leakage. Therefore hematocrit monitoring would be a better option.

Pathology AIIMS 2017

Question 1:

What is the function of Acetone free Methanol in Leishman stain?

a) Causes metabolic and enzymatic activity to stop

b) Prevents the cells from sticking to the slide surface

c) Colours the red cells

d) Fixes cells to slide

Question 2:

A 30-year-old woman presented with painful blisters on the skin and oral mucosa. Direct immunofluorescence is given below. Which of the following is false regarding the same?



a) Antibodies against desmoglein 3

b) Antibodies against desmoglein 1

c) Suprabasal acantholytic blister is a characteristic feature

d) Antibodies against hemidesmosomes

Question 3:

The stain done on a section of liver shown below is:



a) Warthin starry stain

b) Steiner silver stain

c) Sweet's reticulin stain

d) Grimelius silver stain

Question 4:

ALK positivity is seen in which of the following neoplasms:

a) Fibromatosis

b) Synovial sarcoma

c) Ewing sarcoma

d) Inflammatory myofibroblastic tumour

Question 5:

Which of the following is true regarding the internal iron homeostasis in iron-deficiency anaemia?

a) Ferritin mRNA concentration iron-responsive element increases and ferritin synthesis decreases

b) Transferrin receptor 1 iron-responsive elements increases transferrin receptor mRNA concentration and synthesis

c) Transferrin receptor 1 iron responsive elements decreases transferrin receptor mRNA concentration and synthesis

d) Ferritin mRNA concentration iron responsive element decreases and ferritin synthesis increases

Question 6:

A 14-year-old patient presents with bilateral proptosis. The biopsy from the region shows this pattern. What is the likely diagnosis?



a) Leiomyoma

b) Schwannoma

c) Rhabdomyosarcoma

d) Fibromatosis

Question 7:

Endoscopic and barium swallow images of a lesion in the oesophagus of a patient are as shown below. What is the expected histopathological finding?



a) Image 2

b) Image 1

c) Image 4

d) Image 3

Question 8:

A patient has a HbA2 of 3% and HbF of 97%. A peripheral blood smear is taken. Which of the following blood pictures will be seen in this patient?



a) 1, 2 and 3

b) 1 and 2

c) 2 and 3

d) 1 and 3

Question 9:

In breast cancer, FISH for gene amplification will be done based on which of the following IHC staining for Her-2/neu?

a) 1+

b) 2+

c) 3+

d) Gene Amplification is done irrespective of Her 2 Neu status

Question 10:

A 30-year-old male patient presents with severe dyspnoea. Mitral stenosis with left atrial hypertrophy is seen. Vegetations present on the mitral valve were excised. What is the most likely diagnosis based on the given HPE image?



a) Rheumatic valvular disease

b) Infective endocarditis

c) Myxoma

d) Sarcoidosis

Question 11:

A child presents with bilateral proptosis. Which investigation will help you in arriving at a diagnosis of chloroma?

a) Peripheral smear

b) Platelets

c) Hb concentration

d) Leucocytic count

Question 12:

All are reduced in Iron Deficiency Anaemia except

a) Ferritin

b) TIBC

c) Iron

d) Transferrin saturation

Question 13:

A 35-year-old depressed man came with complaints of sudden onset breathlessness, with desaturation. Chest x-ray showed bilateral diffuse infiltrates with predominant involvement of right middle and lower lobes. The patient could not be revived despite treatment. On autopsy, the lungs appeared grossly normal. The histopathology image of the lung is as shown below. Which among the following is the appropriate cause?



a) Necrotizing TB granuloma

b) Vegetative matter aspiration pneumonia

c) Sarcoid granuloma

d) Severe fungal pneumonia

Question 14:

Which of the following can be used to store blood for up to 35 days?

a) Acid Citrate dextrose

b) Citrate phosphate dextrose

c) Citrate phosphate 2 dextrose

d) Citrate phosphate dextrose adenine - 1

Question 15:

Identify the correct sequence:

a) Bedside blood collection, label the samples, fill the laboratory requisition forms, Identify the patient

b) Fill the laboratory requisition forms, Identify the Patient, Bedside blood collection, label the samples

c) Label the sample tubes, Bedside blood collection, fill the laboratory requisition forms, Identify the patient

d) Identify the Patient, Fill the laboratory requisition forms, Bedside blood collection, label the samples

Question 16:

Which anticoagulant is used for electrolyte estimation?

a) EDTA

b) Citrate

c) Sodium Fluoride

d) Lithium Heparin

Question 17:

When you add calcium and thromboplastin to platelet poor plasma which pathway is assessed?

a) Extrinsic

b) Intrinsic

c) Fibrinolysis

d) Common

Question 18:

Origin of B cell in CLL/SLL is:

a) Centrocytes

b) Extranodal mature Peripheral B cells

c) Naive B cells in Interfollicular zones

d) Bone Marrow Progenitors

Question 19:

A young boy from high altitude came with fever and weakness and had the following reports - Hemoglobin 17 g%, TLC is 21,000 with neutrophils 25, lymphocytes 30, eosinophils 5, myelocytes and metamyelocytes 40 in peripheral smear. Next step in the investigation should be:

a) Philadelphia chromosome

b) JAK mutation

c) Erythropoietin levels

d) Bone Marrow biopsy with reticulin stain

Question 20:

A 40-year-old man presents with complaints of facial puffiness and frothy urine. Given below is the histological image of the condition. What do you think is the diagnosis?



a) Membranous nephropathy

b) Focal segmental glomerulosclerosis

c) Membranoproliferative nephritis

d) Dense deposit disease

Question 21:

A 50-year-old man presented with the blurring of vision. Urine examination showed proteinuria. Fundus examination showed dot and blot hemorrhages, microaneurysms, and cotton wool spots. Histopathology of the kidney was as shown below. What is your diagnosis?



a) Renal amyloidosis

b) Focal segmental glomerulosclerosis

c) Kimmelstiel-Wilson disease

d) Lupus nephritis

Question 22:

If the RB gene is hypophosphorylated, which of the following will happen?

a) Cell cycle will stop at G2

b) Cell cycle will stop at G1

c) There will be no effect on cell cycle as RB gene phosphorylation is not needed

d) The cell cycle progresses and cell divides

Question 23:

A 35-year-old patient presented with heartburn and increased salivation. Upper gastrointestinal endoscopy was done, and the biopsy result is shown below. What is the diagnosis?



a) Esophagitis

b) Squamous cell carcinoma

c) Adenocarcinoma

d) Barrett's esophagus

Question 24:

A 60-year-old male presented with a history of intractable diarrhea. Based on the histological image shown below which of the following is the most appropriate diagnosis:



a) Lymphoma

b) Leishmaniasis

c) Urate nephropathy

d) Multiple myeloma

Question 25:

A 10-year-old boy with fever, weight loss came to the hospital. On examination, he was pale and had significant lymphadenopathy. The bone marrow histology of the person is as given below. Your diagnosis is?



a) Acute monocytic leukemia

b) Acute myelomonocytic leukemia

c) Acute lymphoblastic leukemia

d) Juvenile myelomonocytic leukemia

Question 26:

A 32 year old female with history of back ache and polyarthralgia was investigated and found to have nucleolar pattern in IF. Which of the following is the best fit in her case?

a) Raynaud phenomenon and sclerodactyly

b) Genital and oral painful ulcers

c) Glomerulonephritis and heart failure

d) Joint pains with nodules on skin

Question 27:

A young male presented with cervical lymphadenopathy. The biopsy of the lymph node showed RS cells with vague nodule formation and a background of reactive T-lymphocytes. The cells were positive for CD20, LCA, EMA and negative for CD 15 and CD30. What is the diagnosis?

a) Nodular Sclerosis - Hodgkin Lymphoma

b) CLL

c) T cell rich B cell lymphoma

d) Nodular Lymphocyte predominant Hodgkin Lymphoma (NLPHL)

Question 28:

Which of the following is a tool used in gene editing:

a) CRISPR

b) Big data

c) Gene Xpert

d) GWAS

Question 29:

A 37-year-old presented with fever. On examination,■ he had enlarged and ulcerated tonsils. His peripheral blood smear showed lymphocytosis. Monospot test was negative. Tonsillectomy was done and it showed large cells mixed with lymphocytes. The cells were positive for CD20, EBV LMP1, MUM1, CD 79a. Background cells were positive for CD3. The cells are negative for CD15. Your most probable diagnosis is



a) Hodgkin lymphoma

b) EBV positive – DLBCL, NOS

c) Infectious mononucleosis

d) EBV positive mucocutaneous ulcer

Question 30:

The scraping of a painless genital ulcer, when stained with Gram stain and viewed under the microscope, shows a gram-negative coccobacillus surrounded by neutrophils. What is your diagnosis?

a) Chlamydia trachomatis

b) Hemophilus ducreyi

c) Klebsiella granulomatis

d) Neisseria gonorrhoea

Question 31:

When papain acts on a molecule of immunoglobulin, which of the following are the end products?

a) 2 FC and 1 FAB fragments

b) 1 Variable chain and 1 constant chain

c) 1 Fc and 2 Fab fragments

d) 2 Fc and 2Fab fragments

Question 32:

Which of the following Immunohistochemistry marker is used in Cyclin D1 negative Mantle cell lymphoma?

a) SOX11

b) ITRA 1

c) MYD88

d) Annexin V

Question 33:

During development, thymic cells recognizing self-antigens are removed to prevent autoimmunity. Which gene among the following is involved in the process?

a) NOTCH1

b) RB gene

c) CPK gene

d) AIRE

Question 34:

Among the following, which cells have MHC class II?

a) NK cells

b) All nucleated cells

c) B cells, dendritic cells, mononuclear phagocytes

d) Platelets

Answer Key

Question No.	Correct Option
1	a

2	d
3	c
4	d
5	b
6	b
7	c
8	d
9	b
10	a
11	a
12	b
13	b
14	d
15	b
16	d
17	a
18	c
19	c
20	a
21	c
22	b
23	d
24	d
25	c
26	a
27	d
28	a
29	b
30	c
31	c
32	a
33	d
34	c

Detailed Explanations

Solution to Question 1:

Acetone free methanol is used in fixation (preservation of biological tissues from decay which acts by preventing metabolic and enzyme activity and not by fixing the cells to the slide).

Acetone (lipid solvent) is not used as it causes shrinkage and destruction of cell membrane making identification of cells difficult:

Fixation can be achieved by

A. Physical method:

• Heating

• Microwaving

• Cryopreservation

B. Chemical method:

• Immersion of specimen in fixative

• Perfusing vascular system with a fixative

The most commonly used fixative is 10% neutral buffered formalin.

Technique/Tissue	Fixative
Electron microscopy	Glutaral dehyde
Trephine biopsy-bone marr ow	Zenker's fluid (AZF fixative)
GI biopsies	Bouin's fixative
Brain	Formalin ammonium chlorid e
Adrenal biopsy	Orth's fluid
PAP smear	Ethanol

Solution to Question 2:

Antibodies against hemidesmosomes is the false statement as the given image shows a characteristic fish-net appearance on direct immunofluorescence, which is diagnostic of pemphigus vulgaris. The antibodies are against the desmosomal adhesion proteins such as desmogleins 1 and 3.

This pattern is due to intraepidermal intercellular deposition of IgG and C3.

The first-line investigation in pemphigus vulgaris is the Tzanck smear. It allows us to visualize the histology by collecting material from the floor of a fresh blister that is stained with Giemsa or any Romanowsky stain. Tzanck cells are seen on microscopy.



Direct immunofluorescence microscopy (DIF) is the most accurate test.

Antibodies against hemidesmosomes that link basal keratinocytes to the underlying basement membrane are seen in bullous pemphigoid.

Solution to Question 3:

The above stain demonstrates reticular fibers of the liver and hence has to be reticulin stain.

Reticulin stain

• Stained by 2 procedures, Gordon and Sweet's reticulin stain (most common) and Gomori reticulin stain.

• They demonstrate reticulin fibers and basement membrane.

• Reticulin fibers: thin, type 3 collagen, widespread in connective tissue

• Basement membrane: type 4 collagen fibers and laminin.

Reticulin stain is used to shows the normal architecture of the liver and thereby help detect disrupted architecture in hepatocellular carcinoma and hepatocellular proliferation.

Solution to Question 4:

ALK (anaplastic lymphoma kinase) positivity (translocation) is seen in Inflammatory myofibroblastic tumor of lung.

ALK positivity means the rearrangements in the ALK gene on chromosome 2p23.

These rearrangements break the ALK locus leads to the formation of chimeric genes encoding signalling pathways.

ALK rearrangements carry a very good prognosis as they show good response to treatment with ALK kinase inhibitors. The tumours lacking ALK rearrangements have a substantially worse prognosis.

ALK positivity is seen in:

• Adenocarcinoma of Lung

• Neuroblastoma

• Inflammatory myofibroblastic tumor of Lung

Solution to Question 5:

In iron deficiency anaemia, there is an increase in transferrin receptor mRNA concentration and synthesis.

Iron absorbed through diet or which is released from circulation is bound to transferrin (iron transfer protein). When erythropoiesis is stimulated, iron-transferrin complexes act as transferrin receptors on the surface of bone marrow erythroid cells. Through receptor-mediated endocytosis, this complex is internalised.

This iron is then made available for heme synthesis while the transferrin-receptor complex is recycled to the surface of the cell, from where the transferrin is released back into circulation and the transferrin receptor re-anchors into the cell membrane. The excess iron after hemoglobin synthesis binds to apoferritin (storage protein), forming ferritin.

Senescent RBCs undergo phagocytosis in the cells of the reticuloendothelial system. The hemoglobin is broken down, the globin and other proteins are returned to the amino acid pool, and the iron is shuttled back to the surface of the reticuloendothelial cell, where it is taken up by transferrin via ferroportin (iron export channel).

There is no regulated excretory pathway for iron, and the only mechanisms by which iron is lost are blood loss (via gastrointestinal bleeding, menstruation, or other forms of bleeding) and the loss of epithelial cells from the skin, gut, and genitourinary tract. The normal daily iron loss from the body is 1-2 mg.

Variables	Iron-deficient state	Iron-excess state
Iron uptake	↑	↓
Transferrin receptor synthesis	↑	↓
Transferrin receptor mRNA concentration	↑	↓
Iron storage	↓	↑
Ferritin synthesis	↓	↑
Ferritin mRNA concentration	No change	No change

Solution to Question 6:

The given clinical history and biopsy picture is characteristic of schwannoma.

The biopsy specimen in schwannoma shows hypocellular (Antoni B) and hypercellular areas (Antoni A). Antoni A area has the nuclei of tumor cells aligned in palisading rows. The nuclear-free zones that lie between nuclear palisading are known as Verocay bodies.

The histological section of a schwannoma showing Antoni A, Antoni B and Verocay bodies is given below:



Solution to Question 7:

The given images showing an esophageal exophytic growth on endoscopy and apple core appearance in the middle 1/3 of the esophagus on barium swallow are suggestive of squamous cell carcinoma. The characteristic histopathological finding in this condition are nests of stratified malignant cells and keratin pearls, which are noted in image 4.

Squamous cell carcinoma



Normal Esophagus

Barrett's Esophagus





Solution to Question 8:

The presence of HbA2 3% and HbF 97% indicates that it is a case of thalassemia major. In thalassemia major, hematological findings in image 1 showing poikilocytes and image 3 showing target cells are seen. Image 2 showing Howell Jolly bodies are seen in post-splenectomy thalassemia patients and not in this case.

Hematological findings in thalassemia major include the following:

- Anisopoikilocytosis
- Target cells (codocytes)
- Nucleated RBC's
- Dacrocytes
- Basophilic stippling
- Polychromatophilic cells







3.

Solution to Question 9:

• Fluorescence in situ hybridization (FISH) should be performed in all cases of breast tumors with immunohistochemistry (IHC) score of 2+.

• Her-2/neu IHC interpretation:

• 1+ = Negative

• 2+ = Equivocal-> FISH is done to differentiate whether the tumor cells are positive or negative for Her-2 Neu expression.

• 3+ = +ve Her 2 Neu. (Circumferential complete membrane staining in >10% of tumor cells)

Solution to Question 10:

The given image depicts Aschoff's body, which is suggestive of rheumatic valvular disease.

Aschoff's bodies have the following components:

• Anitschkow cells or plump macrophages

• Abundant cytoplasm

• Caterpillar cells - refer to centrally condensed chromatin into slender, wavy ribbons

• Foci of fibrinoid necrosis

• Aschoff's giant cells - multinucleated larger macrophages

• T-cells

• Plasma cells





Solution to Question 11:

• Chloroma is an extramedullary manifestation of acute myeloid leukemia (AML). It is a solid collection of leukemic cells occurring outside of the bone marrow.

• The best investigation among the given options would be a peripheral blood smear.

• But the Investigation of Choice for AML is Flow Cytometry

Solution to Question 12:

In Iron deficiency Anaemia, Total Iron Binding Capacity(TIBC) is increased in order to compensate to the deficient state.

• Diagnostic criteria include of Iron deficiency anaemia include :

• Hypochromic and microcytic red cell indices

• Low serum ferritin(<15mg/dl) and iron levels(<30mg/dl)

• Low transferrin saturation(<15%)

• Increased total iron-binding capacity

• The platelet count often is elevated

• Erythropoietin levels are elevated, but the marrow response is blunted by the iron deficiency; thus, marrow cellularity usually is only slightly increased.

Solution to Question 13:

The given histopathological image clearly shows a granuloma with refractile granules indicative of a foreign body. It is seen in aspiration pneumonia due to vegetative matter (such as food).

In this condition, the chest x-ray shows bilateral diffuse infiltrates especially in the right middle and lower lobes. Patients with altered sensorium are at high risk (stroke, medications).



Solution to Question 14:

Citrate dextrose phosphate adenine-1 (CDPA-1) can be used to store blood for up to 35 days. It helps in improving ATP synthesis during storage. Hence, it increases the shelf life.

The disadvantage of using CDPA-1 is that it increases uric acid stone formation.

The shelf life of storage with other components are:

• Acid Citrate dextrose – 21 days

• Citrate-phosphate dextrose – 21 days

Solution to Question 15:

After making a provisional diagnosis by the physician, some laboratory tests are requested. Requisition forms are filled accordingly. Later the patient is identified and samples are collected by laboratory personnel/ health care staff and labeled appropriately.

The WHO guidelines for taking a blood sample is as illustrated below:



Solution to Question 16:

Lithium heparin is used for Electrolyte estimation.

Heparin

• It is the anticoagulant causing the least interference with the test.

• It is the best anticoagulant to use when minimal hemolysis is desired e.g. for sodium and potassium.

• Heparin accelerates the action of antithrombin III which neutralizes thrombin and thus prevents the formation of fibrin from fibrinogen.

• Heparin is added (0.2 mg /mL of blood) or a drop of heparin is drawn into the syringe.

• Usually found as green top tubes

• Drawbacks

• It inhibits the acid phosphatase activity.

• It gives a blue background for Wright's stain smears so not good for peripheral blood smear interpretation.

• It also affects the binding of triiodothyronine and thyroxine to their carrier protein and produces a higher free concentration of these hormones.

• It interferes with binding of calcium to EDTA.

Solution to Question 17:

• Platelet poor plasma is used in assessing Coagulation pathway.

• Thromboplastin (Factor III) binds with Factor VII. When calcium is added, it gears up the coagulation process via Extrinsic pathway.

• Coagulation Parameters used to assess :

• Extrinsic Pathway - Prothrombin time (PT)

• Intrinsic Pathway - Activated Partial Thromboplastin time (aPTT)

• Common Pathway - Both PT and aPTT



Solution to Question 18:

In CLL/SLL, B cells originate from naive B cell or Pre/Post Germinal Centre memory B cell. CLL arises in the lymph node.

DNA sequencing has revealed that the Ig genes of some CLL/SLL are somatically hypermutated, whereas others are not, suggesting that the cell of origin may be either a postgerminal center memory B cell or a naive B cell. For unclear reasons, tumors with unmutated Ig segments (those putatively of naive B-cell origin) pursue a more aggressive course.

Centrocytes are present in Germinal Centre. They are not the origin of CLL.

Solution to Question 19:

The given clinical scenario is suggestive of polycythemia (Hb>16 g/dL). Hence, erythropoietin levels (EPO) are estimated next, to differentiate between primary or secondary polycythemia. Based on EPO levels:

• Primary polycythemia - normal/ decreased

• Secondary polycythemia - increased

Options A and B: JAK 2 mutation and Philadelphia chromosome analysis is only done as a last resort.

Option D: Bone marrow estimation with reticulin is done for myelofibrosis which can be seen in the spent/ late stage of the disease. However, peripheral smear findings are not suggestive of primary myelofibrosis. Leukoerythroblastic reaction with dacrocytes is seen in myelofibrosis.

Solution to Question 20:

The image is characteristic of membranous nephropathy. The given hematoxylin and eosin (H&E)-stained image shows uniform, diffuse thickening of the glomerular capillary wall. (intense pink color)

Membranous nephropathy is one of the most common causes of nephrotic syndrome in older adults, with a peak incidence in 30–50 years and a male:female ratio of 2:1.

It is a form of chronic immune-complex-mediated disease.

It can be primary (75% of cases) or secondary (associated with systemic diseases/identifiable etiologic agents).

Microscopy:

• Light microscopy: diffuse thickening of the glomerular capillary wall without cellular proliferation. Irregular spikes projecting from the basement membrane are best seen with silver stains.

• Immunofluorescence microscopy: granular deposits containing Ig and complement along the glomerular basement membrane

• Electron micrograph: electron-dense subepithelial deposits (epimembranous spikes) and effacement of foot processes of podocytes

Image: The silver stain of the glomerulus highlights the proteinaceous basement membranes in black. There are characteristic "spikes" of basement membrane between the immune deposits of membranous nephropathy. The black basement membrane material shown here appears as projections in the capillary loops.



Solution to Question 21:

Histopathological image shows acellular periodic acid Schiff (PAS)-positive nodules suggestive of nodular glomerulosclerosis/intercapillary glomerulosclerosis/Kimmelstiel–Wilson disease, which occurs in patients with diabetic nephropathy.

These lesions appear as ovoid or spherical, often laminated, PAS-positive nodules within the mesangial core of the glomerular lobules.

These nodular lesions are frequently accompanied by prominent accumulations of hyaline material in capillary loops (“fibrin caps”) or adherent to Bowman capsules (“capsular drops”).

Renal failure is the second common cause of death in diabetes mellitus, the first being myocardial infarction.

Three renal lesions are encountered in diabetic nephropathy:

1. Glomerular lesions:

• Capillary basement membrane thickening

• Diffuse mesangial sclerosis

• Nodular glomerulosclerosis/Kimmelstiel–Wilson disease

2. Renal vascular lesions, principally hyaline arteriolosclerosis

3. Pyelonephritis, including necrotizing papillitis

Renal histopathology in

• Amyloidosis: Glomerular deposits appearing as thickenings of the mesangial matrix, forming interlacing broad ribbons of amyloid

• Lupus nephritis: Immune complex deposits in the mesangium or along the entire basement membrane (wire loop lesions), with mesangial cell proliferation

• Focal segmental glomerulosclerosis: Focal and segmental hyalinosis and sclerosis of the capillary wall; lipid droplets in the sclerosed area.

Solution to Question 22:

If the retinoblastoma (RB) gene is hypophosphorylated, it is active and the cell cycle will stop at G1.

RB is a cell-cycle regulator because it binds to and inactivates a transcription factor (E2F). This is necessary for the transcription of certain genes necessary for the progression from the G1 to S phase. The phosphorylation of RB results in the release of E2F. This activates gene transcription and cell-cycle progression occurs.

The image given below shows the cell cycle regulation by RB gene.



Solution to Question 23:

This image indicates the diagnostic features of Barrett's esophagus.

The histological image above shows the transition between normal squamous epithelium and the mucin-filled glands, indicating intestinal-type metaplasia with plenty of goblet cells.

Barrett's esophagus is a complication of chronic gastroesophageal reflux disease (GERD) characterized by intestinal metaplasia within the esophageal squamous mucosa. It is a premalignant condition that confers an increased risk of esophageal adenocarcinoma.

Hence, patients diagnosed with Barrett's Esophagus must be subjected to a regular surveillance endoscopy annually or with 2-year intervals, provided no dysplasias have been detected. The presence of dysplasia is associated with prolonged symptoms, longer segment length, increased patient age, and Caucasian race.

Solution to Question 24:

The given clinical scenario points to the diagnosis of multiple myeloma.

The image shows Russell bodies which are eosinophilic, large, homogeneous immunoglobulin-containing inclusions usually found in a plasma cell undergoing excessive synthesis of immunoglobulins. Russell bodies are distended endoplasmic reticulum.

Globular inclusions are Russell bodies (if cytoplasmic) and Dutcher bodies (if nuclear).

The below image shows Russell bodies:



Solution to Question 25:

The given scenario with bone marrow histology showing lymphoblasts (condensed nuclear chromatin, small nucleoli, scanty agranular cytoplasm) is suggestive of acute lymphoblastic leukemia.

ALL is more common in childhood. Clinical features of ALL include:

• Bone marrow failure - anemia, fever, infections, bleeding

• Mass effects due to neoplastic infiltration - bone pain, generalized lymphadenopathy, splenomegaly, hepatomegaly, and testicular enlargement

• CNS manifestations - headache, vomiting, and nerve palsies

• B-ALL - Pancytopenia due to marrow involvement

• T-ALL - Mediastinal mass, compression of large vessels and airways in the mediastinum, lymphadenopathy, splenomegaly

Solution to Question 26:

The nucleolar pattern on IF (immunofluorescence) is suggestive of RNA antibodies which point towards the diagnosis of systemic sclerosis. Hence, Raynaud phenomenon and sclerodactyly are the best fit in the given case.

The nucleolar pattern in immunofluorescence:



Some sclerodactyly patients develop a combination called the CREST syndrome:

• Calcinosis

• Raynaud phenomenon

• Esophageal dysmotility

• Sclerodactyly

• Telangiectasia

Patients with this syndrome have relatively limited involvement of skin, often confined to fingers, forearms, and face, and calcification of the subcutaneous tissues. Involvement of the viscera may not occur at all or occur late. In general, these patients live longer than those with systemic sclerosis with diffuse visceral involvement at the outset.

Solution to Question 27:

The given clinical scenario is suggestive of nodular lymphocyte-predominant Hodgkin's lymphoma (NLPHL).

Axial group (cervical) lymphadenopathy and along with the L&H variant of Reed-Sternberg cells is characteristic.

RS cells are neoplastic giant cells. In NLPHL, the cells have vague nodular formation. Immunophenotyping is positive for CD20 (B cell-associated), LCA (CD45), EMA (epithelial

marker), and negative for CD15 (monocyte or macrophage-associated) and CD30 (activation marker).

The image given below shows the L&H variant of RS cells seen in NLPHL.



Solution to Question 28:

Guide RNAs called CRISPRs help to selectively alter or correct DNA sequences.

CRISPR is Clustered Regularly Interspaced Short Palindromic Repeats

• CRISPR-Cas9 is a genome editing tool and this system enables select bacteria to respond to and eliminate invading genetic material.

• CRISPRs are transcribed into an RNA sequence that directs the Cas9 to a sequence leading to its cleavage.

• Gene editing uses an artificial guide RNA that binds Cas9 and is complementary to a DNA sequence of interest and Cas9 induces the break of double-stranded DNA.

One of the first successful diseases used on experimental basis is Sickle Cell Anemia.

Other gene editing tools are:

• Zinc-finger nucleases (ZNFs)

• Transcription-Activator Like Effector Nucleases (TALEN)

• Meganucleases

Solution to Question 29:

The given clinical scenario suggests a likely diagnosis of EBV positive DLBCL, NOS (Epstein Barr virus-positive Diffuse Large B cell Lymphoma, Not otherwise specified).

The given slide shows sheets of large lymphoid cells with irregular vesicular nuclei and prominent nucleoli typical of DLBCL.

Approach to the question:

• In the given case, Monospot test negative rules out Infectious Mononucleosis and the identification of Reed-Sternberg cell and their variant is essential for the diagnosis of Hodgkin Lymphoma, which can also be ruled out.

• EBV-positive mucocutaneous ulcer is a localized lymphoproliferative disorder in immunocompromised patients. It is a self-limiting ulcer which reverts on treatment in Immunocompromised people. Histologically characterized by atypical large B cells with RS cell-like morphology. Hence, this is also ruled out.

• EBV positive DLBL(Diffuse Large B cell Lymphoma) occurs in patients more than 50 years of age and without a previous history of lymphoma. DLBCL typically presents as a rapidly enlarging mass at a nodal or extranodal site.

• Immunohistochemical studies indicate that EBV+ DLBCL, NOS is molecularly heterogeneous with a prominent background of T lymphocytes. The cells are positive for CD 20, CD 79a, IRF4/MUM1. EBV LMP-1(Latent membrane protein-1) is positive in about 2/3rdof cases, while EBNA-2 (Epstein-Barr nuclear antigen) is positive in the remaining 1/3rd of cases. It is negative for CD10 and BCL6.

Solution to Question 30:

The above description of a gram stain showing Gram-negative Coccobacillus surrounded by neutrophils is suggestive of Klebsiella granulomatis.

Granuloma inguinale, or donovanosis, is a sexually transmitted chronic inflammatory disease caused by Klebsiella granulomatis, a minute, encapsulated, Coccobacillus. It can exhibit a characteristic safety pin appearance on microscopy.

Klebsiella granulomatis is difficult to culture and diagnosis needs visualization of dark-staining Donovan bodies on tissue crush preparation or biopsy. The organisms are demonstrable in Giemsa-stained smears of the exudate as a minute, encapsulated Coccobacilli (Donovan bodies) in macrophages.

Options 1, 4: Gram-negative Diplococci

Option 2: School of fish appearance

Solution to Question 31:

When papain acts on a molecule of immunoglobulin, the end products are 1 Fc and 2 Fab fragments.

If an antibody molecule is digested with papain, cleavage occurs above the disulfide bonds that hold the heavy chains together. This generates 3 separate fragments, two of which are called Fab (fragment antigen binding), and one of which is called Fc (fragment crystallizable).

The image given below shows the cleavage of immunoglobulin by papain.



Cleavage of the antibody molecule with pepsin generates one large fragment called F(ab´)2 and a digested Fc fragment.

The image given below shows the cleavage of immunoglobulin by pepsin.



Solution to Question 32:

SOX11 is used in Cyclin D1 negative Mantle cell lymphoma. Mutation of SOX11 is associated with worse prognosis in mantle cell lymphoma.

Other options are not associated with mantle cell lymphoma.

Solution to Question 33:

The above process is called central tolerance and the AIRE (AutoImmune REgulator) protein is necessary for this process.

Immature self-reactive T and B lymphocyte clones recognize self-antigens during their maturation in the central lymphoid organs (thymus for T cells, bone marrow for B cells). They are killed or rendered harmless.

The removal of self reacting T lymphocytes in the thymus is called negative selection/ deletion. For this process, AIRE protein is needed. Mutations in the AIRE gene results in autoimmune disorders (autoimmune polyendocrinopathy).

B cells undergo a process called receptor editing. When developing B cells encounter an antigen, it formulates a change in the receptor that makes the antigen unrecognizable. Self-reacting B cells can also undergo apoptosis.

Solution to Question 34:

Class II major histocompatibility complex (MHC) molecules are mainly expressed on macrophages, B lymphocytes, and dendritic cells. These cells present ingested antigens to T-cells.

The function of MHC molecules is to display peptide fragments of protein antigens for recognition by antigen-specific T cells. In humans, the MHC molecules are called human leukocyte antigens (HLA) because they were initially detected on leukocytes by the binding of antibodies. The combination of HLA alleles in each individual is called the HLA haplotype.

The genes encoding HLA molecules are clustered on the short arm of chromosome 6.

Uses of MHC/ HLA typing include:

• Anthropology

• Transfusion

• Paternity testing

• Forensic science

• Transplantation

• Disease correlation

Pathology AIIMS 2018

Question 1:

Choose the correct option regarding the storage temperatures of blood components.

a) RBC 20-22°C, Platelet 2-6°C, FFP -30°C

b) RBC 2-6°C, Platelet 20-22°C, FFP -30°C

c) RBC -30°C, FFP 2-6 °C, Platelet 20-22°C

d) RBC 20-22°C, FFP below -30°C, Platelet 2-6 °C

Question 2:

Acid mucin is stained by _____.

a) PAS

b) Alcian blue

c) Masson trichrome

d) PTAH

Question 3:

Toluidine blue staining is used in the Identification of which of the following cells?

a) Lymphocyte

b) Fibroblast

c) Macrophage

d) Mast Cell

Question 4:

t(11,22) is best diagnosed by _____

a) Conventional karyotyping

b) FISH

c) Bone marrow aspiration

d) Next generation sequencing

Question 5:

Gamma delta T cell(γδ TCRs) receptors recognizes

a) Lipids without MHC proteins

b) Peptides with MHC proteins

c) CD20

d) CD3

Question 6:

Cancer that least commonly spreads by lymphatic route is _____.

a) Squamous cell carcinoma

b) Merkel Cell Carcinoma

c) Basal Cell Carcinoma

d) Malignant melanoma

Question 7:

What is correct regarding transfusion of the blood received from blood bank?

a) It should be started within 4 hours of receiving it from the blood bank.

b) It should be completed within 4 hours of receiving from the blood bank.

c) No such time restraint and can be done anytime.

d) It should be completed within 6 hours of receiving from the blood bank.

Question 8:

A patient working in a factory presented with lung fibrosis and thickening of pleura. Biopsy was done. Diagnosis from the histopathology picture is:



a) Asbestosis

b) Silicosis

c) Tuberculosis

d) Mesothelioma

Question 9:

All leukemias come under provisional entity in WHO 2016 classification of hematolymphoid tumours except_____.

a) Natual Killer cell lymphoblastic leukemia/lymphoma

b) B-lymphoblastic leukemia/lymphoma with hypodiploidy

c) Acute myeloid leukemia with RUNX1 mutation

d) Splenic B-cell lymphoma/leukemia

Question 10:

A 65-year-old HIV-positive man presented with fever and breathlessness for 2 weeks. Imaging shows dense infiltration in the lower lobe and the lung biopsy showed the following finding. What is the diagnosis?



a) Small-cell lung cancer

b) CMV-associated organizing pneumonia

c) Cryptococcal pneumonia

d) Tuberculosis

Question 11:

Which of the following is a true statement about apoptosis?

a) There is enzymatic digestion of cellular contents

b) It is associated with adjacent inflammation

c) Karyolysis is present

d) Plasma membrane remains intact

Question 12:

Which of the following is the color of vacutainer containing sodium fluoride as a content?

a) Red

b) Grey

c) Blue

d) Yellow

Question 13:

What does the following image show?



a) Autophagy

b) Emperipolesis

c) Phagocytosis

d) Amoebic autophagy

Question 14:

A patient presents with abdominal distension and intermittent fever. Peripheral smear findings are normal. A tumor is resected and the histopathology image of the resected lymph node is shown below. What is the most probable diagnosis?



a) Angiolymphoma

b) Hodgkin's lymphoma

c) IgG4-related disease

d) Castleman disease

Question 15:

Which of the following is not usually involved in the immunohistochemistry of acute lymphoblastic leukemia?

a) Tdt

b) CD1a

c) CD3

d) Cytokeratin

Question 16:

Antibody-dependent cellular cytotoxicity (ADCC) is seen with:

a) Natural killer (NK) cells only

b) Macrophages only

c) NK cells, neutrophils and eosinophils

d) Neutrophils and eosinophils only

Question 17:

A middle-aged male patient has proteinuria. Looking at the immunofluorescence image below, what is the diagnosis?



a) Focal segmental glomerulosclerosis (FSGS)

b) Post-streptococcal glomerulonephritis (PSGN)

c) Good pasture syndrome

d) Lupus nephritis

Question 18:

Which of the following is involved in recognition and clearance of necrotic cells?

a) Inflammasome

b) Toll-like receptors (TLRs)

c) P selectin

d) Manoreceptor

Question 19:

A patient has been brought with severe bleeding to your ER. His blood group is unknown. Fresh frozen plasma from which of the following blood groups can be given?

a) O

b) AB

c) B

d) A

Question 20:

Which of the following is not a proto-oncogene implicated in carcinoma of thyroid?

a) RAS

b) RET

c) BRAF

d) p53

Question 21:

NARP syndrome is a?

a) Lysosomal storage disorder

b) Mitochondrial function disorder

c) Glycogen storage disorder

d) Golgi body transport disorder

Question 22:

Which amongst the following is true about Turner's syndrome?

a) Webbing of hands and toes are associated with risk of visceral anomalies

b) Noonan syndrome is more associated with cardiac defects

c) Absence of neck webbing is associated with higher risk of co-arctation of aorta

d) Webbing of neck is associated with higher risk of cardiac anomalies

Answer Key

Question No.	Correct Option
1	b

2	b
3	d
4	b
5	a
6	c
7	b
8	a
9	b
10	b
11	d
12	b
13	b
14	d
15	d
16	c
17	d
18	a
19	b
20	d
21	b
22	d

Detailed Explanations

Solution to Question 1:

The recommended storage temperatures of different blood components is as follows:

Blood component	Storage temperature
Whole blood and red blood cells	2–6°C
Fresh frozen plasma	-30°C or lower
Cryoprecipitate	Below -30°C
AlbuminFreeze-dried factor VIII	2–8°C
Platelet	20–24°C

Solution to Question 2:

Acid mucin is stained by Alcian blue.

Solution to Question 3:

Toluidine blue is the special stain for mast cells.

It stains the mast cells violet in a blue background.

Mast cells may be difficult to differentiate from lymphocytes in routine, hematoxylin, and eosin–stained sections, and special metachromatic stains (toluidine blue or Giemsa) are used to visualize their granules.

Other Options:

Melanocytes can be identified using melan A/ MART-1, MITF, S100, HMB45.

Macrophages can be identified using MITF, SOX 10, CD31, CD68.

Fibroblast can be identified using Fibroblast Surface Antigen, Vimentin, HSP47, Fibroblast-specific protein 1.

Solution to Question 4:

t(11,22) can be best detected using fluorescence in situ hybridization (FISH).

t(11,22) is a characteristic translocation giving rise to Ewing's Sarcoma. In 85 to 90 percent of cases of Ewing sarcoma family of tumors (EFT), a recurrent chromosomal translocation, t(11;22)(q24;q12), fuses the 5' portion of the EWSR1 gene on chromosome 22 to the 3' portion of the FLI1 gene on chromosome 11.

Solution to Question 5:

γδ TCRs (T-cell receptors) have been shown to recognize lipid antigens presented by CD1d in an MHC-independent manner

T lymphocytes develop in the thymus from precursors that arise from hematopoietic stem cells. Mature T cells are found in the blood, where they constitute 60% to 70% of lymphocytes, and in T-cell zones of peripheral lymphoid organs.

• Each T cell recognizes a specific cell-bound antigen by means of an antigen-specific T-cell receptor (TCR).

• In approximately 95% of T cells, the TCR consists of a disulfide-linked heterodimer made up of α and β polypeptide chain, each having a variable (antigen-binding) region and a constant region.

• The αβ TCR recognizes peptide antigens that are presented by major histocompatibility complex (MHC) molecules on the surfaces of antigen-presenting cells. By limiting the specificity of T cells for peptides displayed by cell surface MHC molecules, called MHC restriction, the immune system ensures that T cells see only cell-associated antigens.

•

• A small population of mature T cells expresses another type of TCR composed of γ and δ polypeptide chains. The γδ TCR recognizes lipids, and small molecules, without a requirement for display by MHC proteins.

Solution to Question 6:

Basal cell carcinoma is a locally aggressive tumor that rarely spreads by the lymphatic route.

Mutations in sonic hedgehog pathway are associated with it.

It presents as small, slowly growing pearly nodule, often with tortuous telangiectatic vessels on its surface, rolled borders, and a central crust (nodular BCC).

The occasional presence of melanin in this variant of nodular BCC (pigmented BCC) may lead to confusion with melanoma.

The most common site is the inner canthus of the eye.

Basal cell carcinoma does not show lymphatic or hematogenous metastasis. It is primarily locally invasive.

Solution to Question 7:

Blood received from blood bank should be completely transfused within 4 hours of receiving.

Principles of blood administration:

• All blood products should be given through an appropriate blood administration set containing a filter.

• A careful aseptic technique should be practised at all times.

• Vital signs should be documented before and after transfusion and as clinically required. For the first 15 minutes after the infusion has begun, the patient should be kept under close observation to detect any signs of a serious transfusion reaction. If none is observed, the infusion rate may be increased.

• One unit of red cells is often given in 1 to 4 hours, depending on the amount to be transfused and on the patient's cardiovascular status. Infusion of a unit for longer than 4 hours is not recommended, as there is a risk of bacterial proliferation because the opened unit is at room temperature.

• No other solution and no medication should be added to or infused through the same tubing as a blood product.

• Ringer's lactate or other solutions containing calcium must never be added to a blood product because the calcium present leads to coagulation.

• After the transfusion has been completed, the transfusion tag or record should become part of the patient's chart.

Solution to Question 8:

This clinical scenario and image showing typical asbestos bodies are suggestive of asbestosis.

Asbestosis is a type of pneumoconiosis associated with asbestos exposure. The disease-causing capabilities of the different forms of asbestos depend on concentration, size, shape, and solubility. This condition is marked by diffuse pulmonary interstitial fibrosis.

Asbestos bodies are a typical finding. They are golden brown, fusiform, or beaded rods with a translucent center. They consist of asbestos fibers coated with an iron-containing proteinaceous material and arise when these fibers are phagocytosed by macrophages.





Other inorganic particulates may become coated with similar iron-protein complexes and are called ferruginous bodies.

Solution to Question 9:

Recent update: As per WHO 2022 Classification of Hematopoietic and Lymphoid Tumors, provisional entities were not created as they lack sufficient evidence. However, we have retained the question and the answer as it is a previous year question.

B-lymphoblastic leukemia/lymphoma is not a provisional entity under the WHO 2016 Classification of Hematopoietic and Lymphoid Tumors.

Provisional entities are tumors that appear to be clinicopathologically distinct, but require additionally published studies for full acceptance.

The following are included as provisional entities in the classification.

Tumor class	Provisional Entity
Myeloid/Lymphoid Neoplasms with eosinophilia and rearrangement	Myeloid/Lymphoid neoplasm with PCM1-JAK2
Myelodysplastic Syndromes	Refractory cytopenia of childhood
Acute myeloid leukemia with related neoplasms	AML with mutated RUNX1
Precursor Lymphoid neoplasm	Natural killer cell lymphoblastic leukemia/lymphoma

Tumor class	Provisional Entity
Mature B cell neoplasm	Splenic B cell lymphoma/leukemiaPediatric nodal marginal zone lymphomaLarge B-cell lymphoma with IRF4 rearrangementEBV-positive mucocutaneous ulcerBurkitt-like lymphoma with 11q aberration
Mature T and NK cell neoplasm	Chronic lymphoproliferative disorder of NK cellsIndolent T-cell lymphoproliferative disorder of the gastrointestinal tractPrimary cutaneous CD8-positive aggressive epidermotropic cytotoxic T-cell lymphomaPrimary cutaneous acral CD8-positive T-cell lymphoma Primary cutaneous CD4-positive small/medium T-cell lymphoproliferative disorderBreast implant-associated anaplastic large cell lymphoma

Solution to Question 10:

The given image shows a cytomegalic pneumocyte containing characteristic intranuclear basophilic inclusion surrounded by a clear halo known as owl's eye appearance. Along with the given clinical scenario, it is suggestive of cytomegalovirus (CMV) pneumonia.





CMV is a β-group herpesvirus. It causes the following conditions:

• Immunocompetent individuals - asymptomatic or mononucleosis-like infection

• Neonates - devastating systemic infections

• Immunosuppressed individuals - serious, life-threatening, disseminated CMV infections of the lungs (pneumonitis) and gastrointestinal tract (colitis)

Option A: Small-cell carcinoma has small cells with scant cytoplasm, salt-and-pepper chromatin with inconspicuous nucleoli.

Option C: In cryptococcal infection, encapsulated yeasts are seen on direct microscopy.

Option D: Tuberculosis is associated with well-defined epithelioid granuloma.

Solution to Question 11:

The plasma membrane remains intact in apoptosis.

Apoptosis literally means falling-off in Greek. It is a pathway of programmed cell death.

Apoptotic cells break up into fragments called apoptotic bodies. These apoptotic bodies have intact plasma membrane but altered structures. Hence, they are subjected to phagocytosis without eliciting inflammation.

Necrosis is an unprogrammed form of cell death that includes:

• Enzymatic digestion

• Adjacent inflammation

• Karyolysis

Solution to Question 12:

Sodium fluoride is present in the grey coloured tube.

Sodium fluoride inhibits glycolysis. Hence, is used for the estimation of blood glucose and alcohol.

Solution to Question 13:

The image shows the phenomenon of emperipolesis. Here, intact hematopoietic cells are seen within the cytoplasm of the host cell without any damage.

The cells frequently involved are neutrophils, lymphocytes, and plasma cells. Both the cells are viable and there are no physiological and morphological consequences to either of them.





Solution to Question 14:

The given clinical scenario and histopathological image showing follicular hyperplasia of lymph nodes are suggestive of Castleman disease.

Castleman disease (angiofollicular lymph node hyperplasia) is a lymphoproliferative disorder that can be associated with HHV-8 (multicentric Castleman disease or MCD) or idiopathic (unicentric Castleman disease or UCD)

There are 3 major histological subtypes:

• Hypervascular (UCD) -

• Lollipop appearance - atrophic germinal centers with radially penetrating blood vessels

• Onion skin appearance of mantle zone around germinal centers

• Plasma cell subtype (UCD or MCD) - hypertrophic germinal centers, increased vascularity

• Mixed variant (MCD) - mix of hypervascular and plasma cell subtype features in the same lymph node

The clinical features include:

• Multiple enlarged lymph nodes

• Night sweats

• Weight loss

• Weakness and fatigue

• Hepatosplenomegaly

As this condition is associated with elevated levels of IL-6, the treatment involves IL-6 inhibitors like siltuximab and tocilizumab.

Note: Hodgkin's lymphoma has characteristic Reed-Sternberg cells in the histopathological section of lymph nodes.

Solution to Question 15:

Cytokeratin is not a marker of acute lymphoblastic leukemia (ALL). It is an epithelial cell marker for thymoma.

Tdt, CD1a, and CD3 are markers of ALL.

Solution to Question 16:

Antibody-dependent cellular cytotoxicity (ADCC) is mediated by NK cells, neutrophils and eosinophils (killing certain parasitic worms like helminths via IgE antibodies).

In ADCC, cells/ antigens are coated with IgG antibodies. These are killed by effector cells like NK cells and macrophages. These effector cells bind to the target by their receptors for the Fc

fragment of IgG and cell lysis proceeds without phagocytosis.

Solution to Question 17:

The given immunofluorescence staining of the kidney biopsy specimen shows positive staining for fibrinogen, IgG, IgA, IgM, C1q, C3, λ chain, and k light chain. This is known as full-house effect. It is seen in lupus nephritis.



Solution to Question 18:

Inflammasomes are involved in the recognition and clearance of necrotic cells.

Inflammasomes are involved in pyroptosis, a process of programmed cell death distinct from apoptosis and necrosis. This process is dependent on caspases 1, 4, and 5 and is responsible for killing of infected cells. It occurs in the following sequence:

• Activation of inflammasome complex by infected cells

• This causes activation of caspase-1

• This results in generation of IL-1, which is responsible for fever, inflammation, leukocyte recruitment

Option B: Toll-like receptors (TLR)- recognizes molecules shared by pathogens

Option C: P-selectins - an important role in the initial recruitment of WBCs at the site of inflammation & in platelet recruitment and aggregation at the site of injury.

Solution to Question 19:

Fresh Frozen Plasma (FFP) from AB blood group can be given to this patient as it is the universal plasma donor. AB plasma does not contain antibodies against antigens on RBCs of AB blood group.

O Rh-negative blood is the universal blood donor. It is used during emergency packed cell blood transfusion with an unknown blood group. It is preferred because it lacks the A and B and Rh antigens. Hence, it cannot be hemolyzed by anti-A, anti-B or anti-D antibodies in the recipient's blood.

Solution to Question 20:

p53 is is a tumor-suppressor gene and not an oncogene.

The following are the proto-oncogenes implicated in carcinoma of the thyroid.

Proto-oncogene	Tumor
RAS	Papillary thyroid carcinomaFollicular thyroid carcinomaHurthle cell carcinoma
RET	Papillary thyroid carcinoma Medullary thyroid carcinoma
MET	Papillary thyroid carcinomaFollicular thyroid carcinoma
BRAF	Papillary thyroid carcinomaTall-cell poorly differentiated cancerAnaplastic carcinoma

Solution to Question 21:

NARP syndrome - Neuropathy Ataxia Retinitis Pigmentosa syndrome is described as a mitochondrial function disorder.

NARP syndrome is due to a point mutation of the mitochondrial ATPase-6 gene. It has 2 phenotypes:

• NARP

• Maternally inherited Leigh syndrome (MILS)

It is characterized by moderate, diffuse cerebral and cerebellar atrophy and symmetric lesions of the basal ganglia on MRI. The severity of the syndrome corresponds to the mutant mitochondrial DNA load in the tissues.

Solution to Question 22:

In Turner's syndrome, webbing of the neck is associated with a higher risk of cardiac anomalies.

The image below summarises the clinical features of Turner's syndrome.



The image given below shows the webbing of neck.

Noonan syndrome is characterised by mutations in the RAS-MAPK pathway. The most common of these is the PTPN11 which encodes a protein tyrosine phosphatase on chromosome 12.

Clinical features include:

• Short stature

• Short webbed neck

• Cystic hygroma

• Tall forehead

• Low posterior hairline

• Dental malocclusion

Although it shares many clinical features with Turner's syndrome, Noonan syndrome affects both sexes. It follows an autosomal dominant pattern and it also has a different pattern of congenital heart disease, typically involving right-sided lesions.

Pathology AIIMS 2019 (May & Nov)

Question 1:

A young boy from high altitude came with fever and weakness and had the following reports - Hemoglobin 17 g%, TLC is 21,000 with neutrophils 25, lymphocytes 30, eosinophils 5, myelocytes and metamyelocytes 40 in peripheral smear. Next step in the investigation should be:

a) Philadelphia chromosome

b) JAK mutation

c) Erythropoietin levels

d) Bone Marrow biopsy with reticulin stain

Question 2:

A 36-year-old male with skin-colored umbilicated papules. Microscopy is as shown below. The most likely diagnosis is_____.



a) Herpes virus

b) Verruca vulgaris

c) Molluscum contagiosum

d) Lichen planus

Question 3:

Which of the following is not an intermediate filament?

a) Vimentin

b) Tubulin

c) Desmin

d) Neurofilament

Question 4:

A 20-year-old male patient is diagnosed with acute myeloid leukemia. The peripheral blood smear picture is given in the image. The cell indicated by the arrow is best described as:



a) Auer rod

b) Flaming promyelocyte

c) Faggot cell

d) Dusky promyelocyte

Question 5:

Which test evaluates the extrinsic pathway?

a) APTT

b) PT

c) TT

d) Bleeding time

Question 6:

A 58-year-old man died 24 hours after the onset of substernal chest pain radiating to the left arm. On autopsy, an atheromatous plaque was seen obstructing the left anterior descending artery. Which type of calcification is most likely to be seen in this condition?

a) Dystrophic with increased serum calcium

b) Metastatic with increased serum calcium

c) Dystrophic with normal serum calcium

d) Metastatic with normal serum calcium

Question 7:

Hepcidin inhibits iron transfer from the enterocyte to the plasma by binding to _____

a) Ferroportin

b) Transferrin

c) Divalent metal transporter 1 (DMT1)

d) Ferritin

Question 8:

A CKD patient had to undergo dialysis and his Hb was 5.2gm/dL. So two blood transfusions were to be given. The first bag was completed in 2 hours. The second was started and in the midway, he developed shortness of breath and hypertension. Vitals were BP - 180/120 mmHg and pulse rate - 110/min. What is the cause?

a) Allergic

b) Transfusion related circulatory overload

c) Transfusion-related acute lung injury

d) Febrile non-hemolytic transfusion reaction

Question 9:

A patient who receives recurrent transfusions for thalassemia develops fever and chills during this time. There is no evidence of hemolysis. Which of the following measures can decrease the rate of transfusion reactions?

a) Leukoreduced RBCs

b) Antibiotics

c) Irradiation

d) Washed RBCs

Question 10:

If DNA is damaged in the cell cycle, which gene causes cell cycle arrest?

a) Rb

b) MYC

c) p53

d) K-RAS

Question 11:

Which of the following is not a hemostatic agent?

a) Zeolite

b) Chitosan

c) Boric acid

d) Kaolin

Question 12:

Acute inflammatory response is not seen in:

a) Pyroptosis

b) Necroptosis

c) Apoptosis

d) Necrosis

Question 13:

An affected male does not have affected children but an affected female always has affected children. Type of inheritance?

a) X linked recessive

b) Autosomal recessive

c) X linked dominant

d) Mitochondrial

Question 14:

Which of the following is the most common cause of liver failure?

a) Alcohol

b) Paracetamol

c) CCl_4

d) Pesticides

Question 15:

Which is the most important light microscopic feature of irreversible cell injury?

a) Amorphous densities in mitochondria

b) Cell membrane blebs and loss of microvilli

c) Ribosomal detachment from endoplasmic reticulum

d) Nuclear changes like pyknosis and karyolysis

Question 16:

Which of the following are features of iron deficiency anaemia?

a) TIBC increased, Ferritin decreased, Transferrin saturation increased

b) TIBC increased, Soluble transferrin receptor decreased, Ferritin increased

c) TIBC increased, Transferrin saturation decreased, Ferritin decreased

d) TIBC increased, Transferrin saturation decreased, Ferritin increased

Question 17:

Which of the following is the major source of plasma von-Willebrand factor?

a) Platelets

b) Lymphocytes

c) Macrophages

d) Endothelial cells

Question 18:

Which of the following causes vasodilation?

a) Histamine

b) Thromboxane A2

c) Serotonin

d) Leukotriene C4

Question 19:

Both intrinsic and extrinsic pathways are seen in:

a) Apoptosis

b) Necrosis

c) Necroptosis

d) Pyroptosis

Question 20:

What is the most likely diagnosis based on the given serum electrophoresis image?



a) Multiple myeloma

b) Tuberculosis

c) Hepatic cirrhosis

d) Nephrotic syndrome

Question 21:

All are small round blue cell tumors except:

a) Hemangioblastoma

b) Ewing's sarcoma

c) Retinoblastoma

d) Neuroblastoma

Question 22:

A 3-year-old boy presents with recurrent bacterial infections and Pneumocystis jiroveci pneumonia. The flow cytometry analysis of the boy (Boxes 3& 4) is shown below (Boxes 1 & 2 are normal). What is the most likely diagnosis?



a) Bare lymphocytes

b) Chronic granulocyte deficiency

c) Hyper IgM syndrome

d) DiGeorge syndrome

Question 23:

Which gene is not involved in SCID?

a) BTK

b) JAK-3

c) RAG1

d) IL-7Rα

Question 24:

True about anaphylactoid reaction are all except:

a) Involves degranulation of mast cells

b) IgE mediated

c) Can occur on first exposure

d) Most commonly caused by NSAIDs

Question 25:

The mutations/changes that convert proto-oncogene to oncogene include all except:

a) Promoter insertion

b) Enhancer deletion

c) Gene amplification

d) Point mutation

Question 26:

Hepcidin inhibits iron transfer from the enterocyte to the plasma by binding to _____

a) Transferrin

b) Ferroportin

c) Ferritin

d) Divalent metal transporter 1

Question 27:

Which of the following about γ■ T-cells is incorrect:

a) Kills extracellular bacteria by granulysin and perforin

b) Need antigen processing and presentation for it recognition

c) Direct killing of infected macrophages

d) Provide protection against mycobacteria

Question 28:

Which of the following is not true regarding the procedure done using the instrument given in the image below?

a) Contraindicated if platelet count <40,000

b) Breath holding is required

c) Procedure can be done in both lateral & prone position

d) Diagnosing an infiltrative or granulomatous disease

Question 29:

Ciliocytophthoria is seen in:

a) Acute respiratory infection

b) Kartagener syndrome

c) Situs inversus

d) Cystic fibrosis

Question 30:

Maltese cross crystals in the urine are seen in which of the following diseases other than nephrotic syndrome?

a) Friedrich's ataxia

b) Felty syndrome

c) Fanconi anemia

d) Fabry's disease

Question 31:

Basophils are activated by:

a) IL-5

b) INF-gamma

c) TNF-alpha

d) IL-2

Question 32:

Which transporter is required for the transport of iron into the enterocyte?

a) Ferroportin

b) Divalent metal transporter 1

c) Divalent metal transporter 2

d) Haptocorrin

Question 33:

Berry aneurysm occurs due to:

a) Endothelial injury of vessel due to HTN

b) Arterial internal elastic lamina defect

c) Endothelial layer defect

d) Adventitia defect

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	c
5	b

6	c
7	a
8	b
9	a
10	c
11	c
12	c
13	d
14	b
15	d
16	c
17	d
18	a
19	a
20	d
21	a
22	c
23	a
24	b
25	b
26	b
27	b
28	a
29	a
30	d
31	a
32	b
33	b

Detailed Explanations

Solution to Question 1:

The given clinical scenario is suggestive of polycythemia (Hb>16 g/dL). Hence, erythropoietin levels (EPO) are estimated next, to differentiate between primary or secondary polycythemia. Based on EPO levels:

• Primary polycythemia - normal/ decreased

• Secondary polycythemia - increased

Options A and B: JAK 2 mutation and Philadelphia chromosome analysis is only done as a last resort.

Option D: Bone marrow estimation with reticulin is done for myelofibrosis which can be seen in the spent/ late stage of the disease. However, peripheral smear findings are not suggestive of primary myelofibrosis. Leukoerythroblastic reaction with dacrocytes is seen in myelofibrosis.

Solution to Question 2:

The given microscopic image shows large, ellipsoid, homogeneous, cytoplasmic inclusion in cells of the stratum granulosum and the stratum corneum classically seen in molluscum contagiosum.

Molluscum contagiosum is a common, self-limited viral disease of the skin caused by a poxvirus. Poxvirus is brick-shaped and is about 300 nm in size, the largest pathogenic poxvirus in humans

Tzanck smear shows intracytoplasmic molluscum bodies (Henderson–Patterson bodies), the largest known inclusion bodies (30–35 μm).

Individual lesions are firm, pruritic, pink to skin-colored umbilicated papules (shown below). A curd-like material can be expressed from the central umbilication



Solution to Question 3:

Tubulin is a part of microtubules and is not an intermediate filament.

There are three major classes of cytoskeletal proteins: microfilaments, intermediate filaments, and microtubules.

1. Actin microfilaments:

These fibrils formed from the globular protein actin (G-actin) noncovalently polymerize into long filaments (F-actin) that intertwine to form double-stranded helices.

2. Intermediate filaments:

These rope-like intermediate filament fibers are found predominantly in a polymerized form within cells and do not possess the quality of active reorganization like actin and microtubules. They are responsible for cellular tensile strength, which allows the cells to bear mechanical stress.

3. Microtubules:

These are formed by non-covalent polymerization of a dimer of two globular proteins, alpha-tubulin, and beta-tubulin. The microtubules in most cells extend outward from a microtubule-organizing center (MTOC), in which the minus ends of microtubules are anchored.

Type of intermediate filament	Description
Lamin A, B, and C	These proteins are localized in thenuclear laminaof all cells. Mutations in lamin proteins are associated with certain disorders such as familial partial lipodystrophy type 2, Hutchinson-Gilford progeria syndrome, and familial dilated cardiomyopathy.
Vimentin	These are mainly found inmesenchymal cells(fibroblasts, endothelium).
Desmin	Present insarcomeresin muscle cells.
Neurofilament	These are found in neuronal axons, imparting strength and rigidity.
Glial fibrillary acidic protein	Present in glial cells around neurons.
Cytokeratins	These proteins are the most diverse among IFs and constitutetype I (acidic)andtype II (basic)IF proteins.There are at least 30 isoforms of this protein divided into two groups:Epithelial keratins(found in epithelial cells) andtrichocytic keratins(found in hair-keratin).

Solution to Question 4:

The given cell shows multiple Auer rods clustered together within a single cell resembling a bundle of sticks. Such cells have been termed as faggot cells (faggot in French means bundle of sticks).

Since the question specifically asks about the cell, the best answer is faggot cell (not Auer rods).

Option A: Auer rods are distinctive needle-like azurophilic granules present in many cases but particularly numerous in AML with the t(15;17) aberration (acute promyelocytic leukemia).

Option B: The term flaming promyelocyte refers to cells that appear to be breaking apart, taking on a vibrant reddish-purple hue with the apparent liberation of granules into the surrounding cellular matrix.

Option C: Promyelocytes containing fine granules appear dusky or hazy and are referred to as dusky promyelocytes.

Solution to Question 5:

Prothrombin time (PT) evaluates the extrinsic pathway of coagulation.

PT measures the clotting time of plasma in the presence of an optimal concentration of tissue extract (thromboplastin). This indicates the overall efficiency of the extrinsic clotting system.

Causes of prolonged PT:

• Administration of oral anticoagulant drugs - vitamin K antagonists

• Liver disease - obstructive jaundice

• Vitamin K deficiency

• Disseminated intravascular coagulation

• Previously undiagnosed factor VII, X, V or prothrombin deficiency or defect (rare)

Activated partial thromboplastin time (APTT) measures the clotting time of plasma after the activation of contact factors and the addition of phospholipid and $CaCl_2$, but without added tissue thromboplastin. It indicates the overall efficiency of the intrinsic pathway.

Causes of prolonged APTT:

• Disseminated intravascular coagulation

• Liver disease

• Massive transfusion with plasma-depleted red blood cells

• Administration of or contamination with heparin or other anticoagulants

• A circulating anticoagulant (inhibitor)

• Deficiency of a coagulation factor other than factor VII

• Intake of oral anticoagulant drugs

• Vitamin K deficiency

Solution to Question 6:

Atheromatous plaques are associated with dystrophic calcification. This is seen in the presence of normal calcium levels in the serum.

The deposition of calcium in dead or dying tissues is called dystrophic calcification. Although it occurs in the presence of normal serum calcium, it can be worsened by hypercalcemia.

Dystrophic calcification is seen in areas of necrosis of any type. Some examples include atheromas of advanced atherosclerosis and areas of caseous necrosis in tuberculosis.

Solution to Question 7:

Hepcidin inhibits iron transfer from the enterocyte to plasma by binding to ferroportin.

Hepcidin is released by the liver in response to increases in intrahepatic iron levels. It inhibits iron transfer from the enterocyte to plasma by binding to ferroportin and causing it to be endocytosed and degraded. As a result, iron becomes trapped within duodenal cells in the form of mucosal ferritin and is lost as these cells are sloughed.



Note: In hereditary hemochromatosis, there will be impaired hepcidin synthesis resulting in increased iron absorption.

Solution to Question 8:

The given clinical scenario is suggestive of transfusion-associated circulatory overload (TACO).

TACO is caused by rapid or excess transfusion of red cell preparations or plasma products. This condition is characterized by:

• Dyspnea with O2% <90 at room air

• Bilateral infiltrates on chest X-ray

• Systolic hypertension

• Elevation of brain natriuretic peptide, 1.5 times that of pre-transfusion levels

It can be prevented by administering the transfusion slowly, at the rate of 1 to 2 ml of blood/kg of body weight per hour. Diuretics are also used if necessary.

Solution to Question 9:

The given scenario is suggestive of febrile non-hemolytic transfusion reaction (FNHTR). Such reactions can be prevented by using leukoreduced RBCs.

Removal of leukocytes from the blood is called leukoreduction. It is done before storage to prevent cytokine formation.

FNHTR may occur due to the following:

• Sensitization to the WBC antigens by previous blood transfusions or pregnancy

• Cytokines produced by the donor leukocytes during storage

• Bacterial contamination

Note: Luspatercept is an erythroid maturation agent, a receptor fusion protein consisting of a modified extracellular domain of human activin receptor type-2B linked to the human Ig-G1 Fc domain. The drug was recently approved by the FDA for the treatment of transfusion-dependent thalassemia.

Solution to Question 10:

If the DNA is damaged in the cell cycle, the p53 gene causes arrest of the cell cycle in G1 phase to provide time for DNA repair. In case the DNA damage could not be repaired, p53 triggers apoptosis.

In conditions where there is mutation in p53 gene, the cells with damaged DNA continue to replicate and thus may undergo neoplastic transformation.

Solution to Question 11:

Boric acid is not a hemostatic agent.

Topical hemostatic agents used in practice include:

• Hemostatic field dressings

• Surgical hemostatic agents

1. The hemostatic field dressings are the granular and gauze-based agents which are used to facilitate coagulation and hemostasis. The active ingredients used in various hemostatic field dressings are kaolin, mineral zeolite, and chitosan.

2. The surgical hemostatic agents are further divided into the following groups based on the nature of the agent:

• Caustic agents: They act by causing local tissue destruction, which leads to protein coagulation and precipitation. This further leads to thrombus formation and sealing of the small blood vessels. The agents include Monsel solution (a20% ferric nitrate solution), silver nitrate, and aluminum chloride.

• Biological physical agents: They act by promoting platelet activation and formation of three-dimensional matrix for the formation of clots. These agents are used in several operations including neurosurgical, orthopedic, thoracic, abdominal, and vascular procedures. The agents include gelatin, oxidized cellulose, and microfibrillar collagen.

• Physiologic agents: These agents activate a specific step of the coagulation cascade and promotes the clot formation. These agents include thrombin and fibrin glue/sealant.

• Combination or polymer agents: These agents include several different combinations of the above agents and various synthetic hemostatic agents. Examples include gelatin-thrombin agent, polyethylene glycol hydrogel, and glutaraldehyde cross-linked albumin.

	Mechanism of action	Advantage	Disadvantage
Kaolin	Cation based activationof the intrinsic pathwayof the coag ulation cascade	It isnon-granularcompound. Hence, it does not leave any residue upon removal of the dressing	Clots may be disruptedupon removal. It is less effective in functional or dilutional coag ulopathies.
Mineral zeolit e	Adsorbs the water, thereby i ncreasing the concentration of the clotting factors, platele ts, RBCs.	It is available in two forms:G ranular zeoliteis effective for pouring into low-flow wound s.Themesh-embeddedproduc t is useful in high flow woun ds.	The chemical reaction which causes adsorption of water is anexothermic reaction.This causes local tissue damage. T he granules may be difficult t o be removed.
Chetosan	Poly-N-acetylglucosamine is commonly known as chitin. The deacetylated form of chit in is called chitosan. It has m ucoadhesive properties whic h causes the dressing to adhe re to the tissues. It induces v asospasm and attracts RBCs and platelets to the site of bl eeding.	It has antimicrobial properti es. It can be easily applied an d has a long shelf-life.	As the dressing material adh eres to the tissues, it may cau sedisruption of the clotsupon removal.

Solution to Question 12:

An inflammatory response is not seen in apoptosis. In this process, the cell membrane remains intact. Hence, there is no leakage of cell contents and no surrounding inflammation.

On the other hand, other forms of cell death including necrosis, necroptosis, and pyroptosis involve the disintegration of the cell membrane. This causes the cell contents to leak out of the cell leading to an acute inflammatory response.

Solution to Question 13:

In mitochondrial inheritance, if the mother is affected and the father is normal, then all their children (both male and female) will be affected. However, children born to an affected father and a normal mother will not be affected.

The mitochondrial inheritance pattern is as shown in the image below:



The unique feature of mitochondrial inheritance is maternal inheritance, as the disease is transmitted through the affected mitochondrial DNA (mtDNA), which is derived from the maternal mitochondria.

The ovum contains several mitochondria, while the spermatozoa has very few mitochondria if present. Hence, the mitochondrial DNA of the zygote is entirely derived from the ovum.

Therefore, affected mother transmits the disease to all the offsprings (both male and female), and only the daughters of the affected mother, will further transmit the disease.

Characteristic features of mitochondrial inheritance:

• The mitochondrial DNA encodes enzymes required for oxidative phosphorylation. Therefore, organs that are most dependent on the oxidative phosphorylation are the ones most affected in diseases due to mutations in mtDNA. These organs include central nervous system, skeletal muscle, cardiac muscle, liver, and kidneys.

• There are several thousands of copies of mtDNA in each mitochondria, of which only a few copies are affected in case of mutation. Therefore, a diseased individual possesses copies of both wild type mtDNA and mutant mtDNA in each mitochondria This phenomenon is called heteroplasmy.

• The disease manifests if a minimum number of copies of mutant mtDNA are present in the mitochondria that can lead to oxidative dysfunction, called the threshold effect. In most cases, this threshold effect is reached in the organs as mentioned above.

• Also, during a cell division, the wild and mutant copies of the mtDNA are unequally distributed among the daughter cells. As a result, the expression of the disease varies even among different cells of an organ.

Solution to Question 14:

The most common cause of liver failure is paracetamol poisoning.

The most common hepatotoxin causing chronic liver disease is alcohol.

Solution to Question 15:

Nuclear changes like pyknosis, karyorrhexis, and karyolysis are features of irreversible cell injury visible on light microscopy.

The ultrastructural findings visible on electron microscopy include:

• Large amorphous floccular densities in mitochondria

• Cell membrane blebs and loss of microvilli

• Ribosomal detachment from endoplasmic reticulum

Solution to Question 16:

In iron deficiency anemia, TIBC is increased, transferrin saturation is decreased, and ferritin is decreased.

Solution to Question 17:

The major source of plasma von-Willebrand factor is endothelial cells.

vWF is also found in:

• Platelet granules - lesser compared to endothelial cells

• Endothelial cells within cytoplasmic vesicles called Weibel-Palade bodies

• Subendothelium - binds to collagen

Solution to Question 18:

Histamine causes marked dilatation of smaller blood vessels that include arterioles, capillaries, and venules.

Solution to Question 19:

Intrinsic and extrinsic pathways are both seen in apoptosis.

Apoptosis has two pathways:

APAF-1 - Apoptosis activating factor-1

TNF - Tumor necrosis factor

Mitochondrialorintrinsicpathway	Death receptororextrinsicpathway
Activation inside the cell	Activation outside the cell
Occurs in response to:DNA damageRadiationInfection	Involved in:Elimination of self-reactive lymphocytesKilling of target cells by cytotoxic T-cells
Involves release of pro-apoptotic proteins (BAX, BAK)They form aBAX-BAKchannel in the mitochondrial membrane	Involves death receptors on cell surfaces (have a death domain)These includeTNF-1and Fas(CD95) of the TNF family
Leakage ofcytochrome-cresults in the formation of Cyt.c-APAF1 complex, followed byapoptosomeformation	FasLon T-cells binds to Fas bind adapter proteins via death domain
Activation ofcaspase 9	Activation ofcaspases 8and10
Both pathways converge into a common pathway that involves the activation of caspases 3, 6, 7 (executioner caspases)	



Solution to Question 20:

The given electrophoresis shows hypoalbuminemia and increased alpha-2 globulin, which is classically seen in nephrotic syndrome.

Solution to Question 21:

Hemangioblastoma is not a small round blue cell tumor.

Hemangioblastomas are tumors of central nervous system that originate from vascular tissues. They demonstrate the proliferation of capillaries with variable sized, closely packed, and thin walled vessels as seen in the image below.



Small round blue cell tumors consist of small round cells that stain blue on routine H&E staining sections. They include:

• Neuroblastoma

• Retinoblastoma

• Ewing's sarcoma

• Wilms tumor

• Rhabdomyosarcomas

• Acute leukemia

The following is the image of neuroblastoma showing Homer-Wright pseudorosettes.



Solution to Question 22:

The deficiency of the CD40 and its absence in B cells interferes with class switching and the production of memory cells. The B-cell response to antigens, is unaffected, accounting for the presence of increased IgM antibodies. Hence, the condition is known as Hyper IgM syndrome.

The given clinical scenario and the flow cytometry is suggestive of Hyper IgM syndrome. The flow cytometry shows the following findings:

• In comparison of image 3 to 1, where side scatter is plotted against CD 19: B-cell gating is present.

Gating of B cells means selecting a cell during a flow cytometry experiment for further analysis or data presentation.CD 19 is widely expressed on B cells. Hence the graph shows B cell gating. Side scatter measurement provides information about the granularity of a cell.

• In comparison of image 4 to 2, where CD 40 is plotted against CD 19: B cells are present only in quadrant 4, which means that there is CD 40 deficiency.

The deficiency of the CD40 and its absence in B cells interferes with class switching and the production of memory cells. The B-cell response to antigens is unaffected, accounting for the presence of increased IgM antibodies. Hence, the condition is known as Hyper IgM syndrome.

Given below is the explanation of how to interpret flow cytometry results:



Interpretation of results:

Quadrant 1- Cells marked in green: -VE for A antigen, +VE for B antigen

Quadrant 2- Cells marked in purple: -VE for both the antigens.

Quadrant 3- Cells marked in yellow: +VE for both the antigens

Quadrant 4- Cells marked in red: +VE for A antigen, -VE for B antigen

Solution to Question 23:

BTK mutation causes X-linked agammaglobulinemia and not SCID (severe combined immunodeficiency).

Severe combined immunodeficiency is a defect in both humoral and cell-mediated immune responses. In this condition, the following genes are involved:

• JAK3

• RAG1

• IL-7R

Persons with SCID are extremely susceptible to recurrent, severe infections by a wide range of pathogens, including:

• Candida albicans

• Pneumocystis jiroveci

• Pseudomonas

• Cytomegalovirus

• Varicella

Solution to Question 24:

Anaphylactoid reactions are those reactions that are similar to the anaphylactic reactions but are not IgE mediated. These reactions are caused by direct degranulation of the mast cells or basophils, which causes the release of mediators.

Unlike the classic anaphylaxis which occurs on re-exposure to the allergen, the anaphylactoid reaction can occur even on the first exposure. The most common agents that can cause an anaphylactoid reaction include NSAIDs and radiocontrast media.

In 2014, World Allergy Organisation has introduced two new terms:

• Immunologic anaphylaxis - It includes anaphylactic reactions mediated by IgE, IgG (in animals), and immune complex/ complement-mediated reactions.

• Non-immunologic anaphylaxis (previously called anaphylactoid reaction) - It includes reactions caused by agents which cause massive degranulation of mast cells or basophils in the absence of immunoglobulin.

The image given below shows the sequence of events in type 1 hypersensitivity reaction.



Solution to Question 25:

Enhancer deletion does not convert a proto-oncogene to an oncogene.

Mechanisms of activating oncogenes:

Mechanism	Activation of oncogenes
Mutation	Mutation activates the gene. A classic example is the point mutation of the RAS oncoge ne.

Mechanism	Activation of oncogenes
Promoter insertion	Insertion of a viral promoter region near a gene activates it.
Enhancer insertion	Insertion of a viral enhancer region near a gene activates it.
Chromosomal translocation	One chromosome is split off and joined to another. An example is the Philadelphia chromosome.
Gene amplification	Abnormal multiplication of a gene resulting in many copies. This can occur with oncogenes and genes involved in tumor drug resistance.

Solution to Question 26:

Hepcidin inhibits iron transfer from the enterocytes into the plasma by binding to ferroportin, which is an iron efflux channel in the enterocyte.

Iron metabolism is regulated by hepcidin, a protein and acute phase reactant produced by the liver. It binds to ferroportin and degrades it. This inhibits the transfer of iron from the enterocyte to the plasma via ferroportin. As a result, iron becomes trapped within the duodenal cells (enterocytes) in the form of mucosal ferritin and is lost in feces when these cells are shed off.

Transcription of hepcidin is increased by the following factors:

• Increased levels of iron

• Inflammatory cytokines

Transcription of hepcidin is decreased by the following factors:

• Iron deficiency

• Hypoxia

• Ineffective erythropoiesis

Hepcidin is encoded by the HAMP gene and its mutations are associated with hereditary hemochromatosis.



Solution to Question 27:

γ■ T-cells do not need antigen processing and presentation for its recognition.

There are two types of T cells: γ■ T-cells and αβ T-cells.

The functions of γ■ T-cells:

• Provide protection against mycobacteria by direct killing of infected macrophages

• Kills extracellular bacteria by granulysin and perforin

• Plays a role in maintaining self-tolerance

• Capable of recognising and lysing diverse cancers in an MHC-unrestricted manner

αβ T-cells have MHC restricted action.

Solution to Question 28:

The image shows a bone marrow biopsy needle. Thrombocytopenia is not in itself a contraindication for bone marrow biopsy. However, a platelet count >20,000 is ensured for safety purposes.

Indications of bone marrow biopsy:

• Evaluation of unexplained anemia, leukopenia, thrombocytopenia, pancytopenia

• Evaluation of polycythemia, thrombocytosis, leukocytosis

• Diagnosis and staging of lymphoma or solid tumors

• Diagnosis and evaluation of plasma cell disorders and leukemias

• Evaluation of suspected infiltrative and storage diseases

• Evaluation of fever of unknown origin, suspected mycobacterial, fungal, or parasitic infections, or granulomatous diseases

Contraindications of bone marrow biopsy:

• Severe hemophilia

• Severe disseminated intravascular coagulopathy

• Severe bleeding disorders

Procedure for bone marrow biopsy:

Prone or lateral position is chosen for the procedure. Posterior superior iliac spine is identified. After disinfection and local anaesthesia, the patient is asked to inhale slowly through the nose and exhale slowly through the mouth to relieve anxiety .The needle is inserted and rotated on its axis.

On reaching the desired depth, the stylet is removed and needle is advanced. The needle is rotated in the opposite direction to loosen the specimen. It is then withdrawn using the same rotational movement.



Solution to Question 29:

Ciliocytophthoria refers to the degeneration of ciliated cells. It occurs in viral infections, such as acute respiratory infections.

Options B and C: Kartagener syndrome is also known as primary ciliary dyskinesia/ immotile cilia syndrome. It occurs due to a defect in the dynein arm of cilia. It is characterized by the triad of:

• Sinusitis

• Situs inversus

• Bronchiectasis

Option D: In cystic fibrosis ciliary ultrastructure remains normal.

Solution to Question 30:

Maltese cross crystals in the urine are seen in Fabry's disease.

Urine analysis in Fabry's disease may reveal oval fat bodies and birefringent glycolipid globules under polarized light showing a Maltese cross appearance. It can also be seen in nephrotic syndrome.

The following image shows Maltese cross casts:



Note: Maltese cross appearance is also seen in short-range shotgun wounds and babesiosis.

Solution to Question 31:

Basophils are activated by IL-5 and IL-3.

Note: IL-5 primarily activates eosinophils.

Solution to Question 32:

DMT1 (divalent metal transporter 1) is required for the transport of iron into the enterocyte.

Dietary iron is mostly in Fe^{3+} (ferric state) and must be reduced to Fe^{2+} (ferrous state) by ferrireductases. Fe^{2+} iron destined for circulation is transported from the cytoplasm across the basolateral enterocyte membrane by ferroportin 1.

Hephaestin is an iron oxidase that converts Fe^{2+} to Fe^{3+}



Solution to Question 33:

Berry aneurysm occurs due an arterial internal elastic lamina defect.

As an aneurysm develops, it typically forms a neck with a dome and the arterial internal elastic lamina disappears at the base of the neck. The media thins and connective tissue replaces the smooth-muscle cells. The site of rupture is most often the dome. Here the wall thins, and the tear that allows bleeding is usually ≤0.5 mm long.

Risk of aneurysm rupture is higher if:

• Aneurysm is >7 mm in diameter

• Aneurysm at the top of the basilar artery

• Aneurysm at the origin of the posterior communicating artery

Pathology AIIMS 2020 (May and Nov INI-CET)

Question 1:

Which of the following is true about celiac disease?

a) Most common cause of malnutrition in developing countries

b) Can be diagnosed with CECT

c) Improves with food rich in carbohydrates

d) Increased risk of GI lymphoma

Question 2:

Cells lining the isthmus of the gastric pits

a) Chief cells

b) Stem cells

c) Parietal cells

d) Mucous cells

Question 3:

Which of the following is not true about stem cells?

a) They can differentiate

b) They can replicate

c) They can divide without any regulatory stimulus

d) They can regenerate damaged tissues

Question 4:

Which of the following is true regarding liquid-based cytology?

a) Urine specimen is used

b) Sanger sequencing is used

c) Study of fluid in tumour microenvironment

d) DNA is collected in microvessels

Question 5:

Which of the following is true regarding glucose-6-phosphate dehydrogenase deficiency?

a) Spectrin defect

b) Cells are spherical

c) Cells don't lyse in hypotonic solution

d) Cells lyse in hypertonic solution

Question 6:

A woman who had undergone total knee replacement surgery comes with complaints of knee pain with the loosening of implants. The following histopathological slide was obtained which showed foreign body giant cells along with the appearance highlighted in the image below. What is your diagnosis?



a) Ochronisis

b) Metallosis

c) Malignant melanoma

d) Rosai dorffman disease

Question 7:

A 24-year-old man met with an RTA and was brought to the casualty after 6 hours. His B.P. was 90/60 mm Hg, heart rate was 110/min, and SpO2 was 92%. A blood transfusion was done. After a few hours, the patient is febrile, tachypneic, and starts bleeding from the IV and NG tube sites. His SpO2 levels started falling. What could be the cause?

a) Transfusion reaction

b) Acute adrenal crisis

c) Fat embolism

d) Hemorrhagic shock

Question 8:

Which of the following is not a component of Papanicolaou stain?

a) Light green SF

b) Orange G

c) Eosin B

d) Hematoxylin

Question 9:

The vacutainer tube shown in the given image contains:

a) Na-EDTA

b) Trisodium citrate

c) NAF-heparin

d) Clot activator

Question 10:

All of the following are crosslinking fixatives except?

a) Glutaraldehyde

b) Formaldehyde

c) Methanol

d) Osmium tetraoxide

Question 11:

All of the following conditions are related to the same core pathology except

a) Parkinson's disease

b) Alzheimers disease

c) Lewy body dementia

d) Multisystem atrophy

Question 12:

Isotype switching is seen in _____

a) Helper T cells

b) Activated B cells

c) Cytotoxic t cells

d) B cells

Question 13:

Which of the following caspase binds to Cyt c and APAF-1 complex?

a) Caspase 1

b) Caspase 8

c) Caspase 9

d) Caspase 4

Question 14:

During rounds, you come across the urinalysis report of the 30-year-old male diagnosed with Fabry's disease. What type of urinary casts would you expect to see in the report?

a) Waxy cast

b) RBC cast

c) WBC cast

d) Maltese cross cast

Question 15:

JAK2 mutation is most commonly associated with _____

a) Polycythemia vera

b) Essential thrombocytosis

c) CML

d) PMF

Question 16:

A 20-year-old female with easy fatiguability and pallor presented to the OPD. The image of her hand is as given below. What is the most likely diagnosis?

a) Aplastic anemia

b) Vitamin B12 deficiency

c) Iron deficiency anemia

d) Hypoalbuminemia

Question 17:

Rees and Ecker diluting fluid is used in _____

a) Wintrobe tube

b) WBC Pipette

c) RBC Pipette

d) Hemoglobin pipette

Question 18:

Which of the following conditions will have the least chances of a dry tap on bone marrow aspiration?

a) Hairy cell leukemia

b) Follicular lymphoma

c) AML M7

d) Myelodysplastic syndrome

Question 19:

The most common lung finding in COVID 19 infection is?

a) Pulmonary infarction

b) Diffuse alveolar damage

c) Endothelial injury

d) Fibrin clots

Question 20:

A patient presented to the OPD with long hands and feet, and a protruding lower jaw. Which of the following genes is most likely implicated in this scenario?

a) GATA1

b) PIT1

c) TPIT

d) SF1

Question 21:

What is your diagnosis with respect to the findings shown in the peripheral smear given below?



a) Iron deficiency anemia

b) B12 deficiency

c) Hereditary spherocytosis

d) Acute myeloid leukemia

Question 22:

Which of the following is not true about nodular lymphocyte-predominant Hodgkin lymphoma?

a) EBV Negative

b) CD 30 Negative

c) CD 20 positive

d) Poor prognosis compared to classical HL

Question 23:

Which fixation technique is used for fat in the histopathological examination?

a) IHC

b) Frozen section

c) Liquid paraffin

d) Formalin fixed

Question 24:

Identify the cells marked in the given image.



a) Lymphocyte- 1, Monocyte-4, Eosinophil-2, Basophil-3

b) Lymphocyte- 2, Monocyte-4, Eosinophil-1, Basophil-3

c) Lymphocyte- 1, Monocyte-2, Eosinophil-3, Basophil-4

d) Lymphocyte- 2, Monocyte-1, Eosinophil-3, Basophil-4

Question 25:

Which of the following clotting factor deficiency will not affect clotting in vivo?

a) Factor VII

b) Factor V

c) Factor XII

d) Factor IX

Question 26:

A child was brought to the emergency department with a big lesion on his forearm. As a surgery resident you decide to excise the lesion in the emergency operation theatre. Which of the following investigations will test the entire coagulation pathway?

a) Activated partial thromboplastin time

b) Prothrombin time

c) Thromboelastography

d) Bleeding time

Question 27:

The absence of the structure marked as A will present with all except:



a) Pernicious anemia

b) Carcinoid syndrome

c) Vit B12 Deficiency

d) Zollinger ellison syndrome

Question 28:

The most appropriate biopsy specimen in case of a thyroid biopsy is?

a) 15 follicular cell clusters contain 10-12 cells

b) 10 follicular cell clusters contain 6-8 cells

c) 3 follicular cell clusters contain 10-15 cells

d) 6 follicular cell clusters contain 10-15 cells

Question 29:

Identify the correct statement regarding immune surveillance.

a) Anti CTLA 4 Ab is a new modality in the treatment of advanced cancers

b) Mutation of tumor supressor genes decreases antigen toxicity

c) PD1/ PDL1 activation leading to tumor supression

d) Binding of CD28 leads to immune tolerance

Question 30:

Membranoproliferative glomerulonephritis is seen in:

a) 1, 2, 3

b) 1, 2, 3, 4

c) 2, 3, 4

d) 1, 2

Question 31:

Match the following:

a) 1-b, 2-d, 3-a, 4-c

b) 1-d, 2-b, 3-c, 4-a

c) 1-a, 2-b, 3-c, 4-d

d) 1-a, 2-b, 3-d, 4-c

Question 32:

Blood examination shows pancytopenia, which can be correct?

a) B, D

b) A, B, C

c) B, C, D

d) A, B, D

Question 33:

Which of the following statements is correct regarding Peutz Jeghers syndrome?

a) 2, 4, and 5

b) 1, 2, and 3

c) 1, 3, and 4

d) 2, 3, and 5

Question 34:

A patient with cervical lymphadenopathy has t(11;14) positive. Which of the following markers should be done to confirm the diagnosis? A. CD10 B. CD 200 C. Sox11 D. Cyclin D1

a) A,B

b) B,D

c) A,C

d) C,D

Answer Key

Question No.	Correct Option
1	d
2	b
3	c
4	a
5	b

6	b
7	a
8	c
9	d
10	c
11	b
12	b
13	c
14	d
15	a
16	b
17	c
18	b
19	b
20	b
21	b
22	d
23	b
24	b
25	c
26	c
27	d
28	d
29	a
30	b
31	b
32	b
33	b
34	d

Detailed Explanations

Solution to Question 1:

Individuals with celiac disease have an increased risk of GI lymphoma.

Celiac disease is a common cause of diarrhea in developed countries. It is due to permanent sensitivity to gluten in susceptible individuals. It results in partial or total villous atrophy that can

be seen on histopathological examination of biopsy samples. CECT is not a modality of diagnosis. The treatment is lifelong gluten free diet.

Solution to Question 2:

Stem cells are the cells lining the isthmus of the gastric pits.

Isthmus is the narrow, upper-third segment of the gastric gland. This region has stem cells and mucous neck cells. Most stem cells constantly regenerate and primarily mature to form mucous neck cells that migrate upwards and form mucosa. Some stem cells differentiate into mature cells (unknown) and move to the base of the gastric gland.



Solution to Question 3:

Stem cells do not divide without any regulatory stimulus.

Stem cells give rise to all the various differentiated tissues. In adults, stem cells replace damaged cells and maintain tissue populations.

The properties of stem cells are:

• Self-renewal

• Asymmetric division - In which one daughter cell enters a differentiation pathway and gives rise to mature cells, while the other remains undifferentiated and retains its self-renewal capacity.

The types of stem cells are:

• Embryonic stem cells which are totipotent

• Adult stem cells present in stem cell niches

The pluripotency of stem cells is due to factors like:

• Oct3/4

• Klf 4

• Sox 2

• cMyc

Solution to Question 4:

Liquid-based cytology can also be done using a urine specimen. It is commonly used in cervical cancer screening in place of a pap smear.

The advantages of liquid-based cytology are:

• The process reduces obscuring inflammatory cells and mucous, blood, and debris

• The cells are deposited on the slide in a monolayer.

• Test sensitivity is improved compared with conventional pap tests.

Liquid-based cytology has found broad acceptance in non-gynecologic cytopreparation such as:

• Thyroid cyst fluid examination.

• Oral pathology: diagnostic for various types of oral lesions.

• This technique is also used to body fluids (e.g. urine, pleural effusions), brushing samples (e.g. gastrointestinal tract, lung), and fine needle aspiration.

Solution to Question 5:

G6PD deficiency has spherical red cells (Spherocytes).

Glucose-6-phosphate dehydrogenase (G6PD) is an enzyme that helps in the synthesis of NADPH. NADPH helps in keeping glutathione in a reduced state (GSH). This reduced glutathione acts as a defense against oxidative damage. In G6PD deficiency, RBCs are exposed to oxidants. These cross-linking of the reactive sulphydryl groups in the globin chain of hemoglobin. These are then deposited as inclusions in RBC membranes as Heinz bodies. When these RBCs pass-through spleen, macrophages pluck out these membrane inclusions. The remaining cells are seen as bite cells. Less severely damaged cells lose a part of their membrane, forming spherocytes.

Spectrin is a part of the RBC membrane skeleton. It is commonly defective in hereditary spherocytosis.

Normal biconcave RBCs have a bit of an 'extra space'. They can accommodate a little fluid when exposed to a hypotonic solution. But spherocytes have already lost a part of its membrane, so are relatively full. Hence if they are placed in a hypotonic solution, they absorb more fluid and result in lysis of RBCs. This forms the basis of testing spherocytes in Hereditary spherocytosis or G6PD and is known as osmotic fragility testing.

On the other hand, placing RBCs in a hypertonic solution may cause them to lose water and shrink in size. But hardly any hemolysis takes place.

Solution to Question 6:

The given clinical scenario and image with the said foreign body giant cell point towards a diagnosis of metallosis.

Metallosis is an uncommon complication following total joint replacements of high wear joints like hip and knees. The metal from the prosthesis infiltrates the bone and soft tissue surrounding it. It presents with pain or squeaking on weight bearing or loosening of implant. Sometimes it may be asymptomatic with imaging findings (cloud sign or bubble sign). Aspiration reveals dark fluid consisting of metallic debris.

Solution to Question 7:

The symptoms of this patient (bleeding from the IV catheter and nasogastric tube sites) are suggestive of DIC (disseminated intravascular coagulation) within a few hours of blood transfusion. Hence, the most likely cause of this presentation is an acute hemolytic transfusion reaction.

These reactions are mediated by pre-existing IgM antibodies that act against the polysaccharide blood group antigens A or B. They bind to the red cells and cause complement-mediated lysis which further leads to intravascular hemolysis and hemoglobinuria. Severe cases may present with DIC, which is a consumptive coagulopathy with bleeding manifestations. Other complications include shock, acute kidney injury, and sometimes, death.

The various blood transfusion reactions according to their time of occurrence are as follows:

• Anaphylactic reaction - Within a few seconds to minutes of the transfusion

• Allergic/Urticarial reactions - Within 2-3 hours of transfusion

• Acute hemolytic transfusion reaction (AHTR) - Within 24 hours of transfusion

• Febrile non-hemolytic transfusion reaction (FNHTR) - Within 6 hours of transfusion

• Delayed hemolytic transfusion reaction (DHTR) - After 24 hours of transfusion

Other options:

Option b- Acute adrenal crisis usually presents as an unexplained shock that is refractory to fluids and vasopressors.

Option c- Fat embolism is associated with fractures of the long bones. It develops within 2-3 days of fracture/trauma. Moreover, a history of petechial rash is characteristic of fat embolism.

Option d- Hemorrhagic shock presents as tachypnea, tachycardia, narrow pulse pressure, and oliguria. Cold extremities and delayed capillary refill time are other features. In this case, hemorrhagic shock might occur as a consequence of bleeding due to DIC, but the cause of this presentation is transfusion reaction.

Solution to Question 8:

Eosin B is not a component of Papanicolaou stain.

Papanicolaou stain is the most popular stain used in the practice of histopathology. It is a polychrome stain containing multiple dyes to stain the various components of cells differentially. Papanicolaou stain includes both acidic and basic dyes:

• Hematoxylin is a basic dye and it stains the nucleus.

• Orange G is an acidic counterstain and it stains matured and keratinized cells.

• Eosin A is the second counterstain and it is a polychrome mixture of Eosin Y and Light green SF.

Solution to Question 9:

The plastic tube shown in the given image has a red stopper, hence it contains clot activator.

Red stopper plastic tubes contain silica as a clot activator. They are used for serum chemistry tests, serology tests, and in blood banks, where both serum and red blood cells may be used.

The tubes are inverted five times to initiate the clotting process.

Note: Red stopper glass tubes are often referred to as clot or plain tubes because they contain no anticoagulants or additives.

Some important plastic tubes and their additives:

Cap colour	Additive present
Light blue top	Sodium citrate
Dark green to p	Heparin
Purple top	EDTA
Pale yellow to p	Acid citrate dextrose
Light grey top	Sodium fluoride and potassiu m oxalate

Solution to Question 10:

Methanol is not a crosslinking fixative. It is a dehydrant coagulant fixative.

Fixation of tissues can be done by physical or chemical methods.

Physical methods include heating, microwaving, and freeze-drying.

Chemical methods: These methods are commonly used in the processing of tissue for various histopathological diagnoses. It includes the use of:

• Cross-linking fixatives: These fixatives add covalent reactive groups which may induce cross-links between proteins, individual protein moieties within nucleic acids, and between nucleic acids and proteins. Eg: Formaldehyde and Glutaraldehyde.

• Dehydrant coagulant fixatives: These fixatives remove free water from tissues and precipitate or coagulate the proteins. Eg: ethanol, methanol, and acetone.

• Compound fixatives: Fixatives, which are the mixtures of reagents are referred to as compound fixatives. Eg: alcoholic formalin fixes tissues in two ways i.e firstly, by adding covalent hydroxymethyl groups and crosslinking proteins and secondly, by coagulation and dehydration.

Note: Osmium tetraoxide causes additive cross-links and is used for ultrastructural visualization of membranes as an additive for post-fixation after glutaraldehyde.

Solution to Question 11:

Alzheimer's disease, unlike the rest, is not an alpha-synucleinopathy. It is a β-amyloidopathy characterized by the accumulation of Aβ amyloid protein.

Alpha-synucleinopathies: These are a group of neurodegenerative disorders, that are characterized by aggregation of α-Synuclein in neurons, nerve fibers, or glial cells to form inclusion bodies.

Solution to Question 12:

Isotype switching is seen in activated B cells.

Isotype switching: Conversion of one antibody class (isotype) to another resulting from the genetic rearrangement of heavy-chain constant-region genes in activated B cells.

Isotype switching is dependent on the presence of both antigens and cytokines secreted by T-cells.

Note: Defect in isotype switching can cause immunodeficiencies known as Hyper IgM syndrome.

Solution to Question 13:

Caspase 9 binds to Cyt c and APAF-1 complex.

Apoptotic stimuli cause the release of cytochrome c into the cytosol. Cytochrome-C in turn binds to a protein called APAF-1 (apoptosis-activating factor-1) to form the apoptosome.

This complex is able to bind caspase-9, the critical initiator caspase of the mitochondrial pathway, and this step is responsible for the activation of other caspases.

Solution to Question 14:

Maltese cross urinary casts are seen in a patient diagnosed with Fabry's disease.

Urine analysis in Fabry's disease may reveal oval fat bodies and birefringent glycolipid globules under polarized light showing a maltese cross appearance. It can also be seen in nephrotic syndrome.

The image below shows Maltese cross appearance under polarized light:



Maltese cross appearance elsewhere:

• Short-range shotgun wound

• Peripheral smear of babesiosis

Solution to Question 15:

Polycythemia vera is strongly associated with activating point mutations in the tyrosine kinase JAK2.

A mutation in JAK2, a key member of the cytokine intracellular signaling pathway, can be found in 90–95% of patients with polycythemia vera.

It is characterized by :

• Increased marrow production of red cells, granulocytes, and platelets (panmyelosis)

• Low erythropoietin levels and elevated hematocrit

• Blood viscosity and sludging lead to both thrombosis and bleeding.

• Headache, dizziness, hypertension, dizziness, venous thrombosis (Budd Chiari syndrome due to hepatic vein thrombosis), bowel infarction, and stroke.

• Intense pruritus especially aquagenic pruritus

• Peptic ulceration, hyperuricemia

• 15% to 20% of patients after an average period of 10 years ->Myelofibrosis and extensive extramedullary hematopoiesis,

• PCV transforms into AML in 2% of patients.

Primary myelofibrosis

• Marrow fibrosis associated with extramedullary hemopoiesis in the spleen.

• Activating mutation in JAK2 (more commonly) or MPL.

• Presence of neoplastic megakaryocytes, which are responsible for the release of fibrogenic factors like PDGF (platelet-derived growth factor) and TGF-β.

• Bone marrow aspiration yields dry tap due to fibrosis of the bone marrow.

• So, bone marrow biopsy is the investigation of choice, which shows the presence of hypocellularity with increased deposition of reticulin inside the marrow, abnormal megakaryocytes and the characteristic finding of dilated marrow sinusoids.

• Blood shows mild leukocytosis, decreased Hb and hematocrit levels, and elevated LAP scores.

• Acute myeloid leukemia (AML) occurs in 5% to 20% of cases.

Essential thrombocytosis (ET) may also be associated with JAK2, MPL, or CALR mutations.

• Essential thrombocytosis (ET) is often associated with activating point mutations in JAK2 (50% of cases) or MPL (5% to 10% of cases), a receptor tyrosine kinase that is normally activated by thrombopoietin.

• In addition, recent DNA sequencing studies have revealed that most of the remaining cases have mutations in calreticulin (CALR), a protein with several functions in the endoplasmic reticulum and the cytoplasm. Since JAK2 and calreticulin mutations are mutually exclusive, it is hypothesized that the calreticulin mutations also increase JAK-STAT signaling through mechanisms that are currently unknown.

CML is the consequence of the balanced translocation between chromosomes 9 and 22 (t[9;22]).

PV, PMF, and ET are characterized by driver mutations that directly or indirectly constitutively activate JAK2, a tyrosine kinase essential for the function of the erythropoietin andthrombopoietin receptors and also utilized by granulocyte colony stimulating factor receptor.

Solution to Question 16:

Skin hyperpigmentation especially on the dorsum of hands involving the knuckles is seen in cyanocobalamin or vitamin B12 deficiency which is a causative factor of megaloblastic anaemia.

In megaloblastic anaemia, there is impairment of DNA synthesis that leads to ineffective hematopoiesis. It also causes distinctive morphologic changes, including abnormally large erythroid precursors and red cells.

Clinical features of megaloblastic anaemia (cyanocobalamin and folate deficiency) include:

Peripheral smear findings:

• Macro-ovalocytes

• Most macrocytes lack the central pallor of normal red cells and even appear hyperchromic, but the MCHC is not elevated.

• Reticulocyte count is low

• Nucleated red cell progenitors occasionally appear in the circulating blood when anaemia is severe

• Neutrophils are also larger than normal (macro polymorphonuclear) and show nuclear hypersegmentation, having five or more nuclear lobules

The image given below shows hypersegmented neutrophils seen in megaloblastic anaemia.

System	Clinical features
Oral	GlossitisAngular cheilosis
Gastrointestinal	AnorexiaWeight lossDiarrhea Constipation
Skin	Hyperpigmentation(reversible)
Hematological	ThrombocytopeniaLeucopeniaIneffective erythropoiesis and unconjugatedjaundice
Cardiovascular	Atherosclerosis leading to PVD, CAD (due to elevated homocysteine)
Neurological (only B12)	Peripheral neuropathySubacute combined degeneration (degeneration of the posterior and pyramidal tracts of the spinal cord)
Pregnancy (only folate)	Recurrent fetal lossNeural tube defects
Others	Mildfeverin severe casesPsychiatric disturbances



Bone marrow findings:

• Markedly hypercellular as a result of increased hematopoietic precursors.

• Sieve-like pattern of chromatin of immature precursors (Megaloblasts)

• Granulocytic precursors also display dysmaturation in the form of giant metamyelocytes and band forms

• Megakaryocytes can be abnormally large and have bizarre, multilobate nuclei

• The derangement in DNA synthesis causes most precursors to undergo apoptosis in the marrow (ineffective hematopoiesis) and leads to pancytopenia

Solution to Question 17:

Rees and Ecker's solution is used as a diluting fluid in the RBC pipette for performing platelet counts. Its composition is:

The steps involved in platelet count are:

I - Filling the pipette

• Diluting fluid is drawn to mark 1 of the RBC pipette and expelled

• Blood is aspirated to 0.5 mark of the RBCs diluting pipette

• Diluting fluid is aspirated to 101 mark

• Blood and diluting fluid are mixed mechanically for 5 min

II - Filling the counting chamber

• Allow the counting chamber to stand for 20-30 minutes to allow the platelets to settle.

• To prevent drying out of the preparation, it should be kept in a moist atmosphere, this is most easily arranged by placing the filled counting chamber in the bottom of a clean Petri dish, alongside which arrange a small piece of cotton wool which has been soaked in water and squeezed out. Petri dish lid is additionally put over the dish.

• After 20-30minutes, the counting chamber can be removed and carefully transferred to the microscope stage.

III - Counting the number of platelets

• Focus on the surface of the hemocytometer chamber with a high power dry objective. Dim the light to make the thrombocytes visible, and count all of the thrombocytes in the entire center 1 mm^2 (erythrocyte counting area) of both sides of the hemocytometer. This gives the number of thrombocytes in 0.2 l of 1: 200 dilution of blood.

• The number of thrombocytes counted X 1000 = number of thrombocytes/l of undiluted blood.

Components	Value
Sodium citrate	3.8 g

Components	Value
Formaldehyde (40 %)	0.2 ml
Brilliant cresyl blue	0.1 g
Distilled water	100 ml

Solution to Question 18:

Follicular lymphoma is the least likely cause of a dry tap on bone marrow aspiration.

Causes of dry tap are:

• Metastatic carcinoma infiltration

• Chronic myeloid leukemia

• Myelofibrosis

• Hairy cell leukemia

• Acute leukemia

• Hodgkin's lymphoma

• AML- M7

Causes of pancytopenia with cellular marrow are:

Primary disease of marrow-

• Myelodysplastic syndrome

• Paroxysmal nocturnal hemoglobinuria (PNH)

• Myelofibrosis

• Aleukemic leukemia

• Myelophthisis

• Bone marrow lymphoma

• Hairy cell leukemia

Secondary to systemic disease-

• SLE, sarcoidosis

• B12, folate deficiency

• Copper deficiency

• Hypersplenism

• Alcohol

• HIV infection, TB

• Brucellosis, leishmaniasis

Note: Although marrow is hypercellular in CML, MDS etc., extensive marrow fibrosis (rare) has been proposed to account for a dry tap on bone marrow biopsy.

Solution to Question 19:

Diffuse alveolar damage may be seen in patients suffering from COVID-19.

COVID-19 related histopathological changes in the lungs include:

• Diffuse alveolar damage

• Hyaline membrane formation

• Desquamation of pneumocytes

• Fibrin deposits in lungs

• Inflammation with mononuclear cells

Note: Diffuse alveolar damage is not specific to COVID-19. It may be seen in all cases of acute respiratory distress syndrome (ARDS) including pneumonia, burns, near drowning, etc.

Solution to Question 20:

Mutations of the pituitary-specific gene PIT-1 can lead to the development of a somatroph adenoma which produces elevated levels of GH Growth hormone (and IGF-1) which can lead to acromegaly as given in the clinical scenario above.

Somatotrophs, mammosomatotrophs, and lactotrophs are derived from stem cells that express the pituitary transcription factor, PIT-1.

If the increased levels of GH are present after closure of the epiphyses, acromegaly develops. In this condition, growth is most conspicuously seen in

• Skin and soft tissues, viscera (thyroid, heart, liver, and adrenals).

• The bones of the face, hands, and feet are thickened and enlarged.

• Bone density may increase (hyperostosis) in both the spine and the hips.

• Enlargement and protusion of the jaw is seen (prognathism), and broadening of the lower face.

• The feet and hands are enlarged, and the fingers become thickened and sausage-like.

GH excess can also be associated with a variety of other manifestatiions, like gonadal dysfunction, diabetes mellitus, generalized muscle weakness, hypertension, arthritis, congestive heart failure, and an increased risk of gastrointestinal cancers.

Other options:

Steroidogenic factor -1 (SF-1) and GATA-2 are factors that are required for gonadotroph differentiation.

Solution to Question 21:

The peripheral smear showing macro-ovalocytes and a hypersegmented neutrophil are suggestive of megaloblastic anaemia. Vitamin B12 deficiency is a cause of megaloblastic anaemia.

Peripheral smear findings in megaloblastic anaemia:

• Macro-ovalocytes

• Most macrocytes lack the central pallor of normal red cells and even appear hyperchromic, but the MCHC is not elevated.

• Reticulocyte count is low

• Nucleated red cell progenitors occasionally appear in the circulating blood when anaemia is severe

• Neutrophils are also larger than normal (macro polymorphonuclear) and show nuclear hypersegmentation, having five or more nuclear lobules

Solution to Question 22:

Nodular lymphocyte-predominant Hodgkin's lymphoma (NLPHL) has an excellent prognosis.

Nodular lymphocyte-predominant Hodgkin's lymphoma (NLPHL) is seen in young males <35 years. They present with cervical and axillary lymphadenopathy. The involved nodes have a nodular infiltrate of mainly small lymphocytes and macrophages, eosinophils, and plasma cells are scanty. It is not associated with EBV.

L&H variant of Reed-Sternberg (RS) cell that has multilobed nucleus resembling a popcorn kernel (popcorn cell) is seen. Immunophenotyping shows CD20 +, BCL6 +, CD15 -, CD30 -. It has an excellent prognosis.

The image shows the L&H variant of RS cell (popcorn cell) which is seen in NLPHL type of Hodgkin’s lymphoma.



Solution to Question 23:

A frozen section is done for the fixation of fats. This ensures that the tissue need not go through routine processing, which can remove lipids.

When the tissue sample goes through rapid freezing, it becomes firmer. The ice acts as an embedding media allowing the tissue to be sectioned.

The temperature range in the cryostat machine usually ranges from 0■C to -35■C. The majority of the tissue can be appropriately sectioned between -15■C to -25■C. Tissue samples that contain water can be sectioned at a higher temperature. The tissues that contain more fat will need to be cut at a lower temperature.

The stain used for fat is the oil red O stain. It can identify neutral lipids and fatty acids in smears and tissues. It is used in the following conditions:

• Fat emboli in lung tissue

• Renal cell carcinoma

• Sebaceous gland tumors of the skin

• Lipid-rich carcinomas

Fixatives containing alcohols, or routine processing with clearing, will remove lipids. Fixation can be achieved by the following methods:

• Physical methods - heating, microwaving, cryopreservation

• Chemical methods - immersion of specimen in fixative, perfusing vascular system with a fixative

Given below are a few fixatives with their uses:

The most commonly used fixative is 10% neutral buffered formalin.

Technique/Tissue	Fixative
Electron microscopy	Glutaraldehyde
Trephine biopsy-bone marrow	Zenker's fluid (AZF fixative)
GI biopsies	Bouin's fixative
Brain	Formalin ammonium chloride
Adrenal biopsy	Orth's fluid
Pap smear	Ethanol

Solution to Question 24:

A peripheral blood smear shows RBCs, WBCs (granulocytes and agranulocytes), and platelets.

Granulocytes:

• Neutrophil - 3 to 5 lobed nucleus with small, bluish granules in the cytoplasm

• Eosinophil - Binucleate/spectacle shaped nuclei with brick-red granules in the cytoplasm. (1)

• Basophil- Large bluish granules in the cytoplasm which obscure the nucleus. (3)

Agranulocytes:

• Lymphocytes- size just larger than an RBC. No granules, no nucleoli present. (2)

• Monocytes - largest cells with a kidney-shaped nucleus. (4)

Solution to Question 25:

Factor XII deficiency will not affect the clotting in vivo.

Unlike the clotting cascade in the lab (in vitro), the cascade in the body (in vivo) differs slightly - as tissue factor (TF) is the major initiator of the cascade. The TF/VIIa complex is the most important activator of factor IX and factor IXa/factor VIIIa complex is the most important activator of factor X. Factor V is involved in both the pathways as shown in the image below.

The coagulation cascade in the laboratory:

• Extrinsic pathway sequence (in a test tube): Tissue factor → Factor VII → Factor X.

• Intrinsic pathway sequence (in a test tube): Factor XII → Factor XI → Factor IX → Factor X

Laboratory methods to initiate clotting include:

• Intrinsic pathway - adding phospholipids, calcium, and either a negatively charged substance such as glass beads

• Extrinsic pathway - a source of tissue factor



Solution to Question 26:

Thromboelastography (TEG) tests the entire coagulation pathway. It measures coagulation cascade, platelet function, and fibrinolysis.

It is a test that quantitatively measures the ability of whole blood to form a clot. The advantages are:

•

Global assessment of blood coagulability, including coagulation cascade, platelet function, and fibrinolysis

•

Rapid real-time bedside test with a simple methodology (point-of-care testing)

•

Diagnosis of coagulopathic bleeding

•

Guide transfusion therapy and decrease the use of blood products

•

Detect dynamic changes in blood coagulation during resuscitation

•

Predict the clinical efficacy of therapeutic agents affecting blood coagulability

Solution to Question 27:

The eosinophilic appearing cells marked as A are the parietal cells, the absence of which is not associated with Zollinger-Ellison Syndrome. The cells marked as B are Chief cells.

Zollinger-Ellison Syndrome is caused by gastrin-secreting neuroendocrine tumors, which leads to parital cell hyperplasia. Gastrinomas are almost always found in the small intestine or pancreas. The most remarkable feature found in the stomach is an increase in oxyntic mucosal thickness due to massive parietal cell hyperplasia.



Other options:

Option A and C: Pernicious anemia is an autoimmune disease characterized by the presence of autoantibodies that impairs the production or activity of the intrinsic factor (IF) that is released from the parietal cells, which is required for the uptake of vitamin B12. There are three types of autoantibodies namely, type I, II, and III. The action of each of these include:

• Autoantibody type I prevents the binding of vitamin B12 to the intrinsic factor (IF).

• Autoantibody type II prevents the binding of vitamin B12-IF complex to the ileal receptors.

• Autoantibody type III acts against the α and β subunits of the gastric proton pump, which impairs the production of intrinsic factor.

These patients also develop chronic atrophic gastritis, characterized by loss of parietal cells, infiltration of lymphocytes and plasma cells, and megaloblastic changes in mucosal cells.

Option B: Gastric carcinoid tumors originate from gastric enterochromaffin-like (ECL) cells. They are referred to as well-differentiated neuroendocrine tumors. The peak incidence of carcinoid tumors is in the 6th decade. These tumors occur throughout the GI tract, most commonly in the appendix, ileum, and rectum, in decreasing order of frequency.

Gastric carcinoid tumors may be associated with endocrine cell hyperplasia, autoimmune chronic atrophic gastritis, MEN-I, and Zollinger–Ellison syndrome. Ileal carcinoid may cause carcinoid syndrome which is characterized by cutaneous flushing, sweating, bronchospasm, colicky abdominal pain, diarrhea, and right-sided cardiac valvular fibrosis.

Solution to Question 28:

The following cytologic findings have been used to define specimen adequacy:

• a minimum of five or six groups of well-preserved cells, with each group containing approximately 10–15 cells

• six clusters of benign cells on at least two slides prepared from separate FNA biopsy samples

• 10 clusters of follicular cells, with each cluster containing at least 20 cells .

Solution to Question 29:

Anti-CTLA-4 antibody ipilimumab was the first drug to achieve a significant improvement in survival of advanced-stage melanoma. Cytotoxic T lymphocyte antigen 4 (CTLA-4) plays an important inhibitory role in the control of T-cell activation.

Ipilimumab is a fully-humanized IgG1 monoclonal antibody that inhibits cytotoxic T lymphocyte antigen-4 (CTLA-4).

It has been postulated that anti-CTLA-4 antibodies act by blocking inhibitory signals from effector T-cells, allowing costimulatory signaling and generation of antitumor T-cell responses. In vitro studies demonstrated also that the antitumor activity of CTLA-4 blockade is mediated by Fc gamma receptor-expressing macrophage and hypothesized a possible role of regulatory T-cell depletion in the tumor microenvironment.

Option B - Mutation of tumor supressor genes is associated with increased risk of cancer. Antigen toxicity is when the immune response to an antigen is excessive or misdirected, causing damage to tissues or organs.

Option C - PD1/PDL1 is an immune checkpoint whose activation inhibits T-cell function and helps with immune tolerance. Blocking PD-1/PD-L1 interaction using immune checkpoint inhibitors reactivates T cells and enhances immune responses against tumors.

Option D - CD28 is primarily involved in providing the second signal required for T cell activation. The first signal comes from the interaction between the T cell receptor (TCR) and the antigen presented by the major histocompatibility complex (MHC) on antigen-presenting cells (APCs). Without CD28-mediated costimulation, T cells that only receive the TCR signal (without the second signal) can become anergic.

Solution to Question 30:

Membranoproliferative glomerulonephritis (MPGN) is seen in HIV, SLE, CLL, and Hepatitis B.

MPGN is caused due to immune-mediated injury. It often presents with a nephritic-nephrotic picture.

The causes of MPGN are as follows:

• Primary MPGN (Idiopathic)

• Secondary MPGN

• System lupus erythematosus (SLE)

• α1-antitrypsin deficiency

• Infections - hepatitis B, hepatitis C, HIV, schistosomiasis, endocarditis, infected ventriculoatrial shunts, chronic visceral abscesses

• Malignancies - chronic lymphocytic leukemia (CLL)

MPGN is characterized by the proliferation of cells predominantly in the mesangium and the capillary loops. Hence it is also called mesangiocapillary glomerulonephritis. There are 2 types of MPGN:

• Type I - It is characterized by the deposition of immune complexes and mesangiocapillary proliferation.

• Type II - It is now known as dense-deposit disease. It belongs to a group of disorders called C3 glomerulopathies.

Light microscopy findings of MPGN are as follows:

• Lobular glomeruli with proliferating mesangial cells, capillary endothelium, and infiltrating leukocytes (endocapillary proliferation)

• Duplication of the basement membrane gives a double contour or tram-track appearance. It is especially evident in silver or PAS stains.

Membranoproliferative Glomerulonephritis - Light microscopy





Solution to Question 31:

The correctly matched associations of the following conditions are:

• Silicosis: Crazy paving pattern

• Mesothelioma: Malignant pleural effusion without mediastinal shift

• Caplan syndrome: Initially demonstrated in coal workers

• Asbestosis: Basal lobes are involved

Caplan's syndrome also called rheumatoid pneumoconiosis, is a combination of rheumatoid arthritis and pneumoconiosis. It manifests as pneumoconiotic nodules along with features of seropositive rheumatoid arthritis. Caplan syndrome was first described in coal miners and later in patients with silicosis too.

	Silicosis	Asbestosis	Malignant mesothelioma
Exposure	Crystallinesilica	Asbestos	Most specific neoplasm as sociated withasbestosexpo sure
Radiologic/ Gross findi ngs	Simple silicosis - small ro undedopacitiesin upper lo besComplicated silicosis - Larger opacities > 1cm an deggshell calcificationof h ilar lymph nodesAcute sili cosis - ground-glass opaci ties on X-ray andcrazy pa ving patternon HRCT (int erlobular septal thickenin g with background of gro und-glass opacity)	X-ray shows peripheralret icular opacitiespredomina ntly in thebasal lobesHRC T shows subpleural fibroti c lines, irregular opacities, tractionbronchiectasis, an dhoneycombing	Diffuse gelatinous,greyish -pink lesionarising either from the visceral or pariet al pleura.

	Silicosis	Asbestosis	Malignant mesothelioma
Clinical manifestations	Increased risk of developi ng TB and bronchogenic c arcinoma	It mainly affects the lower lobesPFT shows arestricti ve patternwith a decrease in lung volumes and diffu sing capacity	Strong male predilection 3:1Can causemalignant pl eural effusion

Solution to Question 32:

Megaloblastic anemia (hypocellular phase), hairy cell leukemia, and hypocellular MDS can show pancytopenia. Acute promyelocytic leukemia is not associated with pancytopenia.

Causes of pancytopenia with hypocellular bone marrow:

• Acquired aplastic anemia

• Hypocellular myelodysplastic syndrome

• Rare aleukemic leukemia

• Some acute lymphoid leukemia

• Rare lymphomas of bone marrow

• Copper deficiency

Causes of pancytopenia with cellular bone marrow:

• Myelodysplastic syndromes

• Systemic lupus erythematosus

• Paroxysmal nocturnal hemoglobinuria (PNH)

• Hypersplenism

• B12, folate deficiency, Copper deficiency

• Myelofibrosis

• Aleukemic leukemia

• HIV, Brucellosis, Leishmaniasis, Tuberculosis Sepsis

• Myelophthisis

• Bone marrow lymphoma

• Sarcoidosis

• Hairy cell leukemia

Solution to Question 33:

Peutz-Jeghers syndrome is an autosomal dominant, hereditary disorder characterized by multiple intestinal hamartomatous polyps having an arborizing pattern of smooth muscle. Germline

heterozygous loss-of-function mutations in the STK11 gene are present in approximately half of individuals with familial Peutz-Jeghers syndrome.

It is also characterized by mucocutaneous dark blue to brown melanin pigmentation, which is particularly evident on the face, buccal mucosa, hands, feet, perianal and genital regions. This rare syndrome presents at a median age of 11 years.

The hamartomatous polyps seen in Peutz-Jeghers syndrome are characterized by a branching network of smooth muscle fibers continuous with the muscularis mucosa that supports the glandular epithelium of the polyp.

Complications:

• Peutz-Jeghers polyps can cause intussusception, which is occasionally fatal.

• It is associated with a markedly increased risk of many malignant conditions. Lifetime risk is approximately 40% and regular surveillance is recommended from a very young age for sex cord tumors of the testes.

• During later life, there is a risk for gastric and small intestinal cancers, and subsequently for colon, pancreatic, breast, lung, ovarian, and uterine cancers.

Histologic image showing the characteristic arborizing network of connective tissue, smooth muscle, lamina propria, and glands lined by the normal-appearing intestinal epithelium.



Note: Congenital hypertrophy of retinal pigment epithelium is seen in Familial adenomatous polyposis (FAP), not Peutz-Jeghers syndrome.

Solution to Question 34:

Mantle cell lymphoma has its origin from naïve B cell which is characterized by a lack of somatic hypermutation of the IgH gene (Due to the tumor arising from Naive B Cells).

Mantle cell lymphoma :

• Tumor cells resemble the normal mantle zone B- cells which surround the germinal center

• Seen in fifth and sixth decades of life

• Has male predominance

• Has (11;14) translocation

• IgH locus on – chromosome 14

• Cyclin D1 locus on – chromosome 11

• Overexpression of Cyclin D1 is seen

• a more indolent, SOX11 variant that often presents with circulating disease with splenomegaly but without significant lymphadenopathy and with a low Ki67 (<10%).

Features:

• Painless lymphadenopathy

• Splenomegaly

• Multifocal mucosal involvement of small bowel and colon – lymphomatoid polyposis

Histology :

• Nodal tumor cells may surround reactive germinal centers to produce a nodular appearance at low power, or diffusely efface the node.

• Typically, the proliferation consists of a homogeneous population of small lymphocytes with irregular to occasionally deeply clefted (cleaved) nuclear contours

• Large cells resembling centroblasts and proliferation centers are absent, distinguishing mantle cell lymphoma from follicular lymphoma and CLL/SLL, respectively

Immunophenotyping :

• CD 5+

• CD 23 negative

• CD19 and CD20 positive

• Moderately high levels of surface Ig (usually IgM and IgD with κ or λ light chain)

• The IgH genes lack somatic hypermutation, supporting an origin from a naive B cell

Type of neoplasm	Cytogenetics	Immunophenotyping
Follicular lymphoma	t(14;18)Increased expression ofbcl-2	Bcl-2 protein, surface Ig, CD19+, CD20+, CD 10+(CALLA), CD 5-
Mantle cell lymphoma	t(11;14)Increased expression ofcyclin D1	Cyclin d1. Surface Ig, CD 5+, CD 23-
Burkitt's lymphoma	t(8;14)increased expression o fc-myc	BCL-6 protein, surface Ig, CD 19+, CD20+, CD 10+(CALLA)
Marginal zone lymphoma	t(11;18)	CD20+, CD 10-, CD 5-

Type of neoplasm	Cytogenetics	Immunophenotyping
Hairy cell leukemia	?TNF alpha	Surface Ig, CD19+, CD20+, C D 25+, CD 103+, CD 11c+; TR AP positive

Pathology NEET 2018

Question 1:

The following cells seen in the lymph node are indicative of



a) Infectious Mononucleosis

b) Measles

c) CMV infection

d) Influenza

Question 2:

What is the most common site for an extragonadal germ cell tumor?

a) Pineal gland

b) Retroperitoneum

c) Sacro coccegeal region

d) Mediastinum

Question 3:

Identify the condition represented in the image below.



a) Emphysema

b) Miliary tuberculosis

c) Sarcoidosis

d) Asbestosis

Question 4:

Which of the following is an epithelial tumor of the stomach?

a) Carcinoid tumor

b) Lymphoma

c) Gastrointestinal stromal tumor

d) Gastric adenocarcinoma

Question 5:

What is the maximum duration for TRALI to occur after blood transfusion?

a) 12 hours

b) 24 hours

c) 6 hours

d) 48 hours

Question 6:

Prolonged bleeding time is seen in ___.

a) Hemophilia A

b) Hemophilia B

c) von Willebrand disease

d) Both (b) and (c)

Question 7:

The single most important long-term limitation of cardiac transplantation is _____

a) Allograft rejection

b) Allograft arteriopathy

c) Infection

d) Malignancy

Question 8:

Which chromosome is involved in myotonic dystrophy?

a) Chromosome 19

b) Chromosome 20

c) Chromosome 21

d) Chromosome 22

Question 9:

Cowden syndrome is associated with mutation in _____

a) PTEN

b) WT1

c) p53

d) Ras

Question 10:

What is the most common cause of agranulocytosis?

a) Marrow infiltration

b) Autoimmune

c) Drug toxicity

d) Radiation exposure

Question 11:

All of these are characteristic features of pancreatic carcinoma except:

a) Cigarette smoking is a risk factor

b) Carcinoma of body of pancreas often present with jaundice

c) They elicit intense desmoplastic response

d) Perineural invasion is common

Question 12:

Stellate granuloma is seen in :

a) Sarcoidosis

b) Cat-scratch disease

c) Cryptococcosis

d) Histoplasmosis

Question 13:

L & H type Reed-Sternberg (RS) cell is seen in:

a) Mixed celluar type

b) Nodular sclerosis

c) Lymphocyte predominant

d) Classical Hodgkin Lymphoma

Question 14:

The most consistent and persistent feature of bladder carcinoma is:

a) Painless hematuria

b) Painful hematuria

c) Dysuria

d) Suprapubic mass

Question 15:

Fever occurs during inflammation due to all except:

a) IL 1

b) Prostaglandins

c) TNF

d) Histamine

Question 16:

Anaplasia refers to:

a) Lack of differentiation

b) Variation in cell size and shape

c) Replacement of one type of cell by other

d) Disordered arrangement of tumor cells

Question 17:

Which of the following is an opsonin?

a) C3a

b) C3b

c) C5a

d) C6

Question 18:

CD-59 is involved in _____.

a) Paroxysmal nocturnal hemoglobinuria

b) Chediak-Higashi syndrome

c) Essential thrombocythemia

d) Primary myelofibrosis

Question 19:

Leiden mutation is of factor:

a) V

b) VIII

c) I

d) X

Question 20:

Which of the following is the most important determinant of malignant potential in the papillary type of renal tumor?

a) Immnunohistology

b) Size

c) Cytogenetics

d) Part of kidney involved

Answer Key

Question No.	Correct Option
1	b
2	d
3	b
4	d
5	c
6	c
7	b
8	a
9	a

10	c
11	b
12	b
13	c
14	a
15	d
16	a
17	b
18	a
19	a
20	b

Detailed Explanations

Solution to Question 1:

The given slide is pathognomonic for measles. This is a slide of a lymph node biopsy showing a Warthin–Finkeldey giant cell with large grapelike clusters of inclusion bodies.

The lymphoid organs typically have marked follicular hyperplasia, large germinal centers, and randomly distributed multinucleate giant cells (Warthin-Finkeldey cells) which have eosinophilic nuclear and cytoplasmic inclusion bodies.

These are pathognomonic of measles and are also found in the lung and sputum.

Solution to Question 2:

Extragonadal germ cell tumor is most commonly seen in mediastinum.

Extragonadal germinal cell tumors (EGGCTs) are rare tumors that predominantly affect young males. Seminomas account for 30-40% of these tumors, and nonseminomatous germ cell tumors (NS-GCTs) account for 60-70%.

Common sites:

• Mediastinum (50-70%)

• Retroperitoneum (30-40%)

• Pineal gland (5%)

• Sacrococcygeal area (less than 5%)

Solution to Question 3:

The above image represents "miliary tuberculosis". Individual lesions are small (2-mm) foci of yellow-white consolidation scattered through the lung parenchyma

Miliary tuberculosis (TB) is the widespread dissemination of Mycobacterium tuberculosis via hematogenous spread. It occurs when organisms draining through the lymphatics enter the venous blood and circulate back to the lung. Miliary TB may infect any number of organs, including the lungs, liver and spleen.

Solution to Question 4:

Gastric adenocarcinoma is an epithelial tumor of the stomach.

Adenocarcinoma is the most common malignancy (90% cases) of the stomach.

Option A: Carcinoid tumor is a neuroendocrine tumor.

Option B: Lymphoma is a lymphoid tumor.

Option C: GIST is the most common mesenchymal tumor of the abdomen.

Solution to Question 5:

The maximum duration for TRALI (transfusion-related acute lung injury) to occur after blood transfusion is upto 6 hours. TRALI is sudden-onset respiratory failure that occurs during or soon after blood transfusion.

The pathogenesis involves a two-hit hypothesis:

• Priming event - increased sequestration and sensitization of neutrophils in the microvasculature of the lung

• Second hit - the factor (most commonly MHC class I antigens) present in the transfused blood product activates the primed neutrophils

Solution to Question 6:

Prolonged bleeding time is seen in vonWillebrand disease.

Solution to Question 7:

The single most important long-term limitation of cardiac transplantation is allograft arteriopathy.

Complications following cardiac transplantation:

Allograft rejection: It is the primary concern that requires close surveillance. Both classic cellular rejection and antibody-mediated rejection is recognized. Cellular rejection is characterized by interstitial lymphocytic inflammation, with associated myocyte damage.

Allograft arteriopathy: It is the single most important long-term limitation for cardiac transplantation. It is a late, progressive, diffusely stenosing intimal proliferation in the coronary arteries, leading to ischemic injury. It can lead to silent myocardial infarction since transplant patients have denervated hearts and do not experience angina. Progressive congestive heart failure and sudden cardiac death can also occur.

Other postoperative problems include infection and malignancies, especially Epstein–Barr virus-associated B-cell lymphomas.

The image given below shows allograft arteriopathy. There is severe diffuse concentric intimal thickening producing critical stenosis. The internal elastic lamina and media are intact (Movat pentachrome stain, elastin black)



Solution to Question 8:

Chromosome 19 is involved in myotonic dystrophy.

Solution to Question 9:

Cowden syndrome is associated with PTEN mutation.

Important mutations and associated diseases:

Mutation	Disease
PDGF	Astrocytoma

Mutation	Disease
PDGFR	Glioma
EGFR-2	Breast carcinoma
RET	MEN syndrome
KIT	Gastrointestinal stromal tum or (GIST)
K-Ras	Colon carcinoma
H-Ras	Bladder carcinoma
N-Ras	Hematological malignancy
ABL	Chronic myeloid leukemia
BRAF	Hairy-cell leukaemia, colorec tal carcinoma, Melanoma, Pa pillary carcinoma thyroid
Beta- cateni n	Hepatocellular carcinoma(liv er)
C-myc	Burkitt lymphoma
N-myc	Neuroblastoma
L-myc	Lung Ca
NF-1	NF-1
NF-2	NF-2
PTEN	Prostate and Endometrial ca ncer
APC	Colon cancer (Polyposis)
SMAD	Pancreatic Carcinoma
p53	Li Fraumeni syndrome
Rb	Retinoblastoma, osteosarcom a, Pineoblastoma
WT-1	Wilms tumor
BRCA-1	Breast, ovarian Ca
BRCA-2	Breast(Male>female), prostat e, Gall bladder Ca

Solution to Question 10:

The most common cause of agranulocytosis is drug toxicity.

Causes of agranulocytosis include:

• Inadequate or ineffective granulopoiesis -

• Suppression of hematopoietic stem cells - aplastic anemia, infiltrative marrow disorders

• Suppression of committed granulocytic precursors

• Ineffective hematopoiesis - megaloblastic anemia, myelodysplastic syndromes

• Congenital conditions - Kostmann syndrome

• Increased destruction or sequestration of neutrophils -

• Immunological injury to neutrophils - idiopathic, systemic lupus erythematosus, drug exposure

• Increased peripheral utilization - bacterial, fungal, or rickettsial infections

• Splenomegaly

Solution to Question 11:

Patients with carcinoma of the head of the pancreas develop jaundice due to obstruction of the biliary tract.

Cigarette smoking doubles the risk of pancreatic cancer and is the leading preventable cause. The clinical features depend on the location of the tumor.

• Head - painless, palpable gallbladder with mild jaundice is suggestive of malignancy (according to Courvoisier's law)

• Body and tail - may remain asymptomatic until later stages as they do not impinge on the biliary tract and do not cause jaundice

Other common pathological features of this condition are:

• Desmoplastic reaction - growth of fibrous or connective tissue

• Perineural invasion, as well as lymphatic and large vessel invasion

Solution to Question 12:

Stellate granuloma is seen in cat-scratch disease. It is also known as subacute regional lymphadenitis.

It is a bacterial infection caused by Bartonella henselae. It is acquired following contact with infected animals (commonly cats).

Regional lymph nodes demonstrate:

• Follicular hyperplasia with circumscribed central stellate necrosis with neutrophils

• Surrounded by palisading histiocytes (suppurative granulomas)

• Sinuses packed with monocytoid B cells

• Usually, there are no perifollicular and intrafollicular epithelioid cells

Cat Scratch Disease



Cat Scratch Disease



Solution to Question 13:

L & H type Reed–Sternberg cell is seen in lymphocyte-predominant Hodgkin lymphoma.

It is characterized by the presence of lymphohistiocytic (L&H) variant RS cells that have a delicate multilobed, puffy nucleus resembling popped corn (popcorn cell). Other reactive cells, such as eosinophils, are scanty or absent, and typical RS cells are rare.

Unlike the RS variants in classical Hodgkin lymphoma, L&H variants express B cell markers (e.g., CD20) and usually fail to express CD15 and CD30.

Subtypes of Hodgkin's lymphoma:

Type	Reed–Sternberg cell variety	EB Virus	Prognosis
Mixed cellularity(Most common in India)	Mononuclear and Diagnostic RS cells	70%	Good
Nodular sclerosis(Most common in world)	Lacunar	5%	Very good
Lymphocyte Rich	Mononuclear and Diagnostic RS cells	40%	Very good
Lymphocyte depleted	Reticular	90%(Max)	Poorest
Lymphocyte predominant	L&H variant (Popcorn)	0%(Min)	Best

Solution to Question 14:

The most consistent and persistent feature of carcinoma bladder is painless hematuria.

Bladder tumors produce a wide range of clinical features including:

• Painless hematuria - classical feature and may be the only clinical manifestation

• Irritative urinary symptoms (frequency, urgency, dysuria) - diffuse tumors or tumors at the bladder neck

• Pyelonephritis or hydronephrosis - when the ureteral orifice is involved

• Obstructive symptoms, palpable suprapubic mass, or lower extremity edema - late presentation with a large tumor mass

• Weight loss and/or localized bone pain - metastatic disease

Solution to Question 15:

Histamine is involved in the mediation of vascular events of inflammation, not fever.

Fever is characterized by an elevation of body temperature, usually by 1° to 4°C. It is one of the most prominent manifestations of the acute-phase response, especially if associated with infection.

Substances that induce fever are called pyrogens. They may be exogenous or endogenous.

• Endogenous pyrogens - IL-1, TNF, prostaglandins

• Exogenous pyrogens -

• Lipopolysaccharides of bacterial walls

• They stimulate leukocytes to release endogenous pyrogens (IL-1 and TNF)

Cytokines result in an increase in cyclooxygenases that convert arachidonic acid into prostaglandins. These prostaglandins are produced in the vascular and perivascular cells of the

hypothalamus. In the hypothalamus, they stimulate the production of neurotransmitters that reset the temperature setpoint to a higher level. This is especially by PGE2.

Solution to Question 16:

Anaplasia refers to a lack of differentiation.

The term anaplasia means to form backward implying a reversal of differentiation to a more primitive level. The extent to which neoplastic cells resemble normal cells, both morphologically and functionally, is called differentiation.

Anaplasia is an important morphological feature of malignant transformation.

The morphological features of malignancy include:

• Anaplasia

• Pleomorphism - variation in the size and shape of cells

• Increased N: C ratio (nucleus: cytoplasmic ratio) - around 1:4 in normal cells, increases to 1:1 in malignant cells

• Hyperchromasia - coarsely clumped and darkly stained chromatin within irregular nuclei

• Increased mitosis - atypical, bizarre mitotic figures

• Loss of polarity - sheets or large masses of tumor cells grow in an anarchic, disorganized fashion

Solution to Question 17:

C3b functions as an opsonin.

Opsonization is the process of coating a foreign particle and preparing it for phagocytosis. The substances involved in this process are called opsonins and include:

• Mannose-binding lectin

• C3b, C4b, C5b

• IgM and Fc receptor of IgG

• CRP

• Fibrinogen

Elie Metchnikoff was the first person to describe the process of phagocytosis, and is hence known as the Father of Innate Immunity.

Solution to Question 18:

CD-59 is involved in paroxysmal nocturnal hemoglobinuria (PNH).

Factors regulating complementary pathways:

• DAF or CD-55 inhibits C3 activation

• MIRL or CD-59 inhibits the membrane-activating complex

In PNH, DAF and MIRL are absent. Hence, there will be unregulated complement activation.

Clinical features of PNH include:

• Intravascular hemoglobinuria - cola-colored urine

• Pancytopenia + hypercellular bone marrow

• Thrombosis (if hepatic vein involved → Budd–Chiari syndrome)

Diagnosis of PNH is by the following methods:

• HAM'S test - acidified serum lysis

• Sucrose lysis test

• Flow cytometry - absence of CD 55 and CD 59 on RBC

Eculizumab is used to alleviate the hemolysis associated with PNH and its sequelae. Eculizumab does not alter the underlying defect of the disease.

Note: FDA has recently approved Pegcetacoplan for adults with PNH which acts by binding to complement protein C3 and increasing hemoglobin concentration.

Solution to Question 19:

Leiden mutation is a single nucleotide mutation that occurs in factor V. This is named after the city in The Netherlands where it was discovered.

Of the inherited causes of hypercoagulability, point mutations in the factor V gene and prothrombin gene are the most common.

Solution to Question 20:

The most important determinant of malignant potential in the papillary type of renal tumor is the size of the tumor.

The papillary type of renal tumors includes:

• Renal papillary adenoma (≤ 1.5 cm)

• Papillary type of renal cell carcinoma (RCC) - > 1.5 cm

Small renal papillary growths are fairly common and incidentally found. Ruling out of malignancy is one of the processes carried out during the evaluation of donor transplant kidneys.

Renal Papillary Adenoma



Papillary architecture (Nipple-like projections)

Papillary renal cell carcinoma (RCC) is the second most common type of renal cell carcinoma and it arises from proximal tubules. It can be multifocal or bilateral. It is typically hemorrhagic and cystic, especially when large.

It is often a circumscribed tumor composed of papillae formed by delicate fibrovascular cores that often contain foamy macrophages and Psammoma bodies.

It has two histological types:

• Type 1 papillary RCC - single layer of cells on the basement membrane and low-grade nuclear features

• Type 2 papillary RCC - pseudostratification of cells and usually has high-grade nuclear features

Pathology NEET 2019

Question 1:

All of the following are features of fragile X syndrome except:

a) Trinucleotide mutation in FMR-1 gene

b) Mental retardation

c) Micro-orchidism

d) Large everted ears

Question 2:

Predominantly osteoblastic secondaries are seen in _____

a) Prostate carcinoma

b) Breast carcinoma

c) Stomach carcinoma

d) Bone carcinoma

Question 3:

A 55 yr old man is found to have hepatomegaly on physical examination. He also complains of dark tan despite avoiding sun exposure. His liver microscopy is given on the slide below. This patient most likely suffers from _____



a) Wilson's disease

b) Autoimmune hepatitis

c) Alpha-1 antitrypsin deficiency

d) Hemochromatosis

Question 4:

Punched-out ulcers in the esophagus are caused by:

a) Herpes-simplex virus

b) Cytomegalovirus

c) Candida

d) Acid

Question 5:

α1-antitrypsin is the major inhibitor of the enzyme _____.

a) Catalase

b) Collagenase

c) Elastase

d) Phospholipase

Question 6:

All are true about Barrett oesophagus except:

a) Complication of chronic GERD

b) Seen in 10% of individuals with symptomatic GERD

c) Increases risk of squamous cell carcinoma

d) Intestinal metaplasia

Question 7:

Which of the following diseases is most commonly associated with Down's syndrome?

a) Parkinson Disease

b) Alzheimer's disease

c) Schizophrenia

d) Huntington's Disease

Question 8:

Which of the following statements is true regarding the modified Duke's criteria?

a) Single positive blood culture for HACEK group is a major criterion

b) Single positive blood culture for Coxiella Burnetti is a major criterion

c) Complete dehiscence of prosthetic valve

d) Documentation of four minor criterion allows a clinical diagnosis

Question 9:

IL-1 is activated by :

a) Caspase - 1

b) Caspase - 3

c) Caspase - 9

d) Caspase - 8

Question 10:

Flask shaped ulcers in the colon are caused by:

a) Entamoeba Histolytica

b) Giardia lamblia

c) Helicobacter pylori

d) Entamoeba vermicularis

Question 11:

Which of the following is true about fever?

a) Aspirin increases fever by inhibiting prostaglandin synthesis

b) PGD2 is responsible for resetting the temperature to a higher level

c) IL-1 is an endogenous pyrogen

d) Fever occurs during inflammation due to release of histamine

Question 12:

Identify the anti-apoptotic factor among the following:

a) P-53

b) K-ras

c) Bcl-2

d) BAK

Question 13:

What is the type of necrosis seen in the vessel wall of a patient with polyarteritis nodosa?

a) Fibrinoid necrosis

b) Gangrenous necrosis

c) Fat necrosis

d) Caseous necrosis

Question 14:

Which of the following is a risk factor for Alzheimer's disease?

a) Apo E1

b) Apo E2

c) Apo E3

d) Apo E4

Question 15:

Alcohol abuse is strongly associated with the development of:

a) Pericarditis

b) Hypertrophic cardiomyopathy

c) Dilated cardiomyopathy

d) Myocarditis

Question 16:

Which lymphoma is strongly associated with overexpression of BCL2?

a) Follicular lymphoma

b) Burkitt lymphoma

c) Diffuse large-cell lymphoma

d) Chronic lymphocytic leukemia

Question 17:

The X-ray of a 65-year-old man is given below. Clinical examination, and hematological and urinary tests were normal. Which of the following is the probable diagnosis?



a) Multiple myeloma

b) Langerhans cell histiocytosis

c) Metastasis

d) Hyperparathyroidism

Question 18:

A 47-year-old female patient with a long-standing history of rheumatic heart disease was admitted to hospital due to severe breathlessness. She died on the 3rd day following admission. The mitral valve shows the following findings on autopsy. What is the cause of these findings?



a) Rupture of valve

b) Calcification and fibrosis

c) Hypertrophy of ventricular wall

d) Hypertrophy of the atrial wall

Question 19:

Which of the following myocardial infarcts cause aneurysm as a post-MI complication?

a) Subendocardial

b) Anterior transmural

c) Posterior transmural

d) Inferior wall

Answer Key

Question No.	Correct Option
1	c
2	a
3	d
4	a
5	c
6	c
7	b
8	b
9	a
10	a
11	c
12	c
13	a
14	d
15	c
16	a
17	b
18	b

19	b

Detailed Explanations

Solution to Question 1:

Macro-orchidism (large testicles) are seen in fragile X syndrome, not micro-orchidism.

Fragile X syndrome is the second most common genetic cause of mental retardation (Down syndrome being the most common). It is caused by a trinucleotide mutation, characterised by multiple tandem repeats of the CGG sequence in the familial mental retardation-1 (FMR1) gene. It is associated with the phenomenon of anticipation, where the disease phenotype worsens over generations within a family.

The affected males are mentally retarded, with an IQ in the range of 20 to 60.

The characteristic phenotypic features include the following:

• Long face with a large mandible

• Large everted ears

• Macro-orchidism

• Hyperextensible joints

• High arched palate

• Mitral valve prolapse

The given image shows a child suffering from fragile X syndrome.

Solution to Question 2:

Predominantly osteoblastic secondaries are seen in prostate carcinoma.

Metastatic tumors of bone:

Metastatic tumors are the most common form of skeletal malignancy, greatly outnumbering primary bone cancers.

Skeletal metastases are typically multifocal. However, carcinomas of the kidney and thyroid may present with solitary lesions. Most metastases involve the axial skeleton (vertebral column, pelvis, ribs, skull, and sternum). The red marrow in these areas, with its rich capillary network and slow blood flow, facilitates implantation and growth of the tumor cells.

The radiographic appearance of metastases may be purely lytic (bone destroying), purely blastic (bone forming), or mixed lytic and blastic. Furthermore, some cancers are associated with predominantly one pattern or the other.

For example, prostatic adenocarcinoma is predominantly blastic whereas carcinomas of the kidney, lung, and gastrointestinal tract and malignant melanoma produce lytic lesions.

Solution to Question 3:

The patient most likely suffers from hemochromatosis. In the given scenario, the liver microscopy image shows hepatocellular iron, which is stained blue with Perl's Prussian blue stain.

Hemochromatosis is an inherited or acquired metabolic disorder caused by excessive iron absorption. Iron overload leads to the accumulation of iron in parenchymal organs such as the liver and pancreas followed by the heart, joints, and endocrine organs.

The image below shows the triad of haemochromatosis.



The image below is the liver microscopy image showing hepatocellular iron, stained with Perl's Prussian blue.

Hemochromatosis - Prussian blue staining



The total body iron pool ranges from 2 to 6 gm in normal adults. In hemochromatosis, total iron accumulation may exceed 50 gm, more than one-third of which accumulates in the liver.

Solution to Question 4:

Infection with Herpes simplex virus (HSV) typically results in punched-out ulcers in the esophagus.

HSV esophagitis commonly occurs in immunocompromised patients. For diagnosis, the biopsy sample should be taken from the edge of the ulcer. The following microscopic findings are noted:

- Ballooning degeneration
- Ground-glass changes in nuclei
- Cowdry type A intranuclear inclusion bodies

Herpes simplex virus infection



Note: CMV esophagitis causes shallow ulcerations. The biopsy is taken from the base of the ulcer and shows characteristic nuclear and cytoplasmic inclusions bodies.

Solution to Question 5:

α1-antitrypsin is the major inhibitor of the enzyme neutrophil elastase.

• Lysosomal enzymes, if unchecked, can potentiate further inflammation by damaging tissues.

• These harmful proteases, however, are normally controlled by a system of antiproteases in the serum and tissue fluids.

• α1-antitrypsin is the prototype member of the serine protease inhibitor (serpin) superfamily of proteins.

Clinical correlate:

• Alpha 1 anti-trypsin deficiency (α1-ATD) is caused by mutations in the SERPINA1 gene located in the long arm of chromosome 14.

• This genetic defect alters the configuration of the α1-antitrypsin molecule and prevents its release from hepatocytes.

• As a result, serum levels of α1-antitrypsin are decreased, leading to low alveolar concentrations, where the α1-antitrypsin molecule normally would provide protection against proteases such neutrophil elastase.

• The resulting protease excess in alveoli destroys alveolar walls and causes emphysema.

Solution to Question 6:

Barrett's esophagus increases the risk of esophageal adenocarcinoma, not squamous cell carcinoma.

Barrett's esophagus is a complication of GERD and it is estimated to occur in as many as 10% of individuals with symptomatic GERD. Microscopically, it is seen as the replacement of the squamous esophageal epithelium with goblet cells (intestinal metaplasia).



The following images show endoscopic and microscopic images of Barrett's esophagus:





Solution to Question 7:

Among the choices provided, Alzheimer's disease is most commonly associated with Down's syndrome.

Conditions most commonly associated with Down's syndrome:

• Congenital heart disease (in about 35% of cases)

• Gastrointestinal anomalies (in about 10%)

• Chronic serous otitis media (in >50%)

• Hypothyroidism

• Alzheimer's disease (in the 30's and 40's)

• Epilepsy (in about 10%)

• Ocular disorders

• Reduced fertility

• Reduced life span (Often due to complications like infections)

Solution to Question 8:

Single positive blood culture for Coxiella burnetti is considered as one of the major criteria to establish the diagnosis of infective endocarditis.

Solution to Question 9:

Caspase- 1 activates the biologically active form of IL-1.

IL-1 is involved in pyroptosis, a form of programmed cell death that is different from apoptosis and necrosis. It is involved in the clearance of necrotic and microbe-infected cells. It occurs in the following sequence:

• Activation of the inflammasome complex

• Activation of caspase-1

• Generation of IL-1 - results in fever, inflammation, leukocyte recruitment

Solution to Question 10:

Flask shaped ulcers are classically seen in amoebiasis caused by Entamoeba histolytica.

In amoebiasis, amoeba attach themselves to the colonic epithelium, induces apoptosis, invade crypts, and burrow laterally into the lamina propria. This recruits neutrophils and causes tissue damage, resulting in flask-shaped ulcers with a narrow neck and broad base.



Flask shaped ulcer

Clinical manifestations:

• Amebic Colitis

• Colicky abdominal pains and frequent bowel movements.

• Diarrhea associated with tenesmus and heme-positive stool.

• Trophozoites invade the intestinal mucosa, produce flask-shaped ulcers.

• Amebic Liver Abscess

• Fever is the hallmark of amebic liver abscess

• It is associated with abdominal pain, abdominal distention, and enlargement and tenderness of the liver.

• Changes at the base of the right lung, such as elevation of the diaphragm and atelectasis or effusion, may also occur.

• The most common finding is a single abscess in the right hepatic lobe, detected on ultrasonography,computed tomography, or magnetic resonance imaging.

Solution to Question 11:

IL-1 is an endogenous pyrogen

Substances that induce fever are called pyrogens. They may be exogenous or endogenous.

• Endogenous pyrogens - IL-1, TNF, prostaglandins

• Exogenous pyrogens -

• Lipopolysaccharides of bacterial walls

• They stimulate leukocytes to release endogenous pyrogens (IL-1 and TNF)

In the hypothalamus, prostaglandins such as PGE2 stimulate the production of neurotransmitters that reset the temperature setpoint at a higher level. Aspirin decreases fever by inhibiting prostaglandin synthesis.

Solution to Question 12:

BCl-2 is an anti-apoptotic factor.

The intrinsic pathway of apoptosis is initiated by the permeabilization of the mitochondrial membrane. This is facilitated by pro-apoptotic proteins (BAX and BAK) and inhibited by anti-apoptotic proteins.

Solution to Question 13:

Immune reactions involving blood vessels, as seen in the case of polyarteritis nodosa, usually show fibrinoid necrosis. Fibrinoid means fibrin-like.

In some immune disorders, the antigen-antibody complexes get deposited in the blood vessels. These complexes result in complement activation (type III hypersensitivity) leading to cell injury. This leads to leakage of fibrin out of the blood vessel, giving a pinkish appearance around the blood vessels.

Fibrinoid necrosis is seen in the following conditions:

• SLE

• Libman-sacks endocarditis

• Arthus reaction

• Serum sickness

• Polyarteritis nodosa
• Post-streptococcal glomerulonephritis
• Malignant hypertension



Solution to Question 14:

Apolipoprotein E4 is a risk factor for Alzheimer's disease.

Solution to Question 15:

Alcohol abuse is strongly associated with the development of dilated cardiomyopathy.

Alcohol and its metabolites, especially acetaldehyde have a direct toxic effect on the myocardium which leads to the development of dilated cardiomyopathy.

Causes of dilated cardiomyopathy:

• Acquired causes :

• Infectious and non-infectious myocarditis: most common cause globally

• Alcohol is the most common cause in India

• Pregnancy:

• Pregnancy-associated cardiomyopathy (PACM)

• Peripartum cardiomyopathy (PPCM)

• Stress-induced: Takotsubo cardiomyopathy

• Drug-induced:

• Doxorubicin (anthracycline)

• Trastuzumab

• Cyclophosphamide

• Nutritional deficiency:

• Beriberi (thiamine),

• Keshan disease (selenium),

• Carnitine deficiency

• Endocrine: Hypo and hyperthyroidism, acromegaly, diabetes mellitus

• Others:

• Cobalt toxicity,

• Heavy metals (lead, mercury, arsenic, lithium)

• Familial causes (50%):

• Mutation in the cardiac myocyte titin gene

• Haemochromatosis

• Syndromes: Duchenne muscular dystrophy, Becker's muscular dystrophy

Solution to Question 16:

Follicular lymphoma is associated with overexpression of BCL2.

Follicular lymphoma

• Arises from germinal center B cells

• Usually, presents in the middle age.

• (14;18) translocation

• BCL-2 is the inhibitor of apoptosis and causes promotion of the follicular lymphoma cell.

Clinical features

• It presents as painless generalized lymphadenopathy with less commonly the involvement of CNS, GIT, or testes.

• Median survival is for 7-9 years

• In almost 50% of patients, this cancer gets transformed to diffuse large B-cell lymphoma

Lymph node biopsy

• Closely packed follicles

• Centrocytes (small cell with the cleaved nucleus and scant cytoplasm) and centroblasts (large cell with open nuclear chromatin and multiple nucleoli).

Bone marrow

• Para-trabecular lymphoid aggregates

Solution to Question 17:

The given image showing a solitary punched-out lytic lesion without a sclerotic rim with a beveled edge appearance, along with the age of the patient with history of normal hematological and urinary tests is suggestive of Langerhans cell histiocytosis.

Langerhans cell histiocytosis occurs due to the proliferation of a special type of immature dendritic cells, called Langerhans cells. Their normal function is to present antigens.

It is characterized by single or multiple osteolytic bone lesions. It can also infiltrate the skin (of the scalp), lungs, liver, spleen, bone marrow, or the central nervous system.

Solution to Question 18:

The given image is of mitral stenosis showing fish mouth or buttonhole appearance in rheumatic heart disease. It is due to calcification and fibrosis.

Rheumatic heart disease is also associated with subendocardial irregular thickenings known as McCallum plaques, as shown below.

Solution to Question 19:

Anterior transmural infarcts are involved in causing aneurysm as a post-MI complication.

Posterior transmural infarcts are more likely to be complicated by conduction blocks, right ventricular involvement, or both.

Pathology NEET 2020

Question 1:

Which of the following is the correct statement regarding Huntington's chorea?

a) It is a trinucleotide repeat disorder

b) There is loss of function type of mutation

c) There are abnormal repeats of CUG

d) Abnormality is seen due to mutation in chromosome 6

Question 2:

Which of the following statements is true about the pathology of achalasia cardia?

a) Relaxation of lower esophageal sphincter

b) Defective peristalsis

c) Decreased release of NO and VIP in the distal part of esophagus

d) Decreased tone of the lower esophageal sphincter

Question 3:

A patient presents with midline swelling in the neck. Biopsy revealing Orphan Annie eye nuclei is shown in the image below. What is the diagnosis?



a) Follicular carcinoma

b) Papillary carcinoma

c) Toxic nodular goitre

d) Medullary carcinoma

Question 4:

A 51-year-old male patient came with complaints of hematuria to the clinic. On examination, he was normotensive and had pedal edema. Investigations revealed the patient had no glycosuria and had a creatinine of 9. His renal biopsy is shown below. Which of the following investigations should be done to identify the disease?



a) ANA

b) Anti GBM antibodies

c) HIV RNA

d) Urine immunoelectrophoresis

Question 5:

A 50-year-old man presented with a retroperitoneal mass. Biopsy from the lesion is shown in the image below. Molecular analysis revealed t(12;16). What is the most likely condition?



a) Myxoid Liposarcoma

b) Lipoma

c) Spindle lipoma

d) Polymorphic lipoma

Question 6:

Large, friable, irregular vegetations on heart valves are seen in which condition?

a) Infective endocarditis

b) Rheumatic heart disease

c) Libman-Sacks endocarditis

d) Non-bacterial thrombotic carditis

Question 7:

A 23-year-old male patient presented with a history of fatigue and tiredness. On investigation, he was found to have Hb = 9 gm% and MCV =101 FL. Peripheral smear examination showed macrocytic RBC and hypersegmented neutrophils. Which of the following is the most likely etiology?

a) Lead poisoning

b) Iron deficiency anemia

c) Hemolytic anemia

d) Chronic alcoholism

Question 8:

A 5-year-old child was admitted to the hospital with a history of prolapsing rectal mass and painless rectal bleeding. The histopathological image of the mass is as shown below. What is the most likely diagnosis?



a) Peutz Jegher syndrome

b) Villous adenoma

c) Juvenile polyp

d) Tubular adenoma

Question 9:

A 20-year-old patient presented to the clinic with a history of malabsorption, diarrhea, and poor appetite. Duodenal biopsy of the patient showed crypt hyperplasia, villi atrophy, and CD8+ cells in lamina propria. What is the most probable diagnosis?

a) Environmental enteropathy

b) Celiac disease

c) Whipple's disease

d) Chronic pancreatitis

Question 10:

A 3-year-old child was brought to the hospital with complaints of fever, rash and swelling of her hands and feet since past 5 days. There is desquamative skin rash, oedema of hands and feet, erythema of palms and soles, oral erythema, and cervical lymphadenopathy. What is the likely condition?

a) Kawasaki disease

b) Measles

c) Scarlet fever

d) Henoch Schonlein Purpura

Question 11:

Effacement of podocyte foot processes is seen in which of the following diseases?

a) Steroid resistant glomerulonephritis

b) Minimal change disease

c) Rapidly progressive glomerulonephritis

d) Chronic glomerulonephritis

Question 12:

Thrombosis is initiated by:

a) Platelet activation

b) Endothelial injury

c) Coagulation cascade

d) Vasoconstriction of vessels

Question 13:

A middle-aged woman presents with thyroid swelling. Blood tests show elevated TSH levels and the presence of thyroid peroxidase antibodies. Histopathological examination revealed the presence of Hurthle ■cells. What is the most likely diagnosis?

a) Hashimoto thyroiditis

b) Graves' disease

c) Follicular carcinoma

d) Medullary carcinoma of thyroid

Question 14:

A 52-year-old male patient comes with complaints of abdominal pain, nausea, and vomiting. A peripheral blood smear reveals macrocytes and hypersegmented neutrophils and an endoscopy reveals atrophic gastritis. Which of the following conditions is likely to be seen this patient?

a) Vitamin B12 deficiency

b) Folate deficiency

c) Pyridoxine deficiency

d) Niacin deficiency

Question 15:

A 35-year-old woman presents with a lump in the upper outer quadrant of her breast. Histopathological examination of the mass showed cells in mucin pools and faint nuclei. What is the most likely diagnosis?

a) Colloid carcinoma

b) Papillary carcinoma

c) Medullary carcinoma

d) Lobular carcinoma

Question 16:

Which of the following is true about the pathogenesis of Hirschsprung disease?

a) Failure of migration of neural crest cells

b) Loss of ganglion cells

c) Visceral myopathy

d) Obstruction by inflammatory stricture

Question 17:

Premature aging in Werner syndrome is due to:

a) Increased telomerase activity

b) Defect in DNA helicase

c) Increase in length of telomere due to decrease in telomerase activity

d) None of the above

Question 18:

A 24-year-old presented with swelling in the wrist joint. Histopathological examination of the swelling revealed spindle cells with palisading of the nuclei and Verocay bodies. Which of the following is the most likely condition?

a) Schwannoma

b) Neurofibroma

c) Neuroma

d) Glioma

Question 19:

B-cells are induced to produce IgE by which of the following:

a) IL-2

b) IL-4

c) IL-1

d) IL-6

Question 20:

The lining epithelium of the respiratory tract of a middle-aged man who is a chronic smoker shows a change from columnar to stratified squamous epithelium, as depicted in the image below. The change is called:



a) Dysplasia

b) Hypertrophy

c) Atrophy

d) Metaplasia

Question 21:

Graft from identical twin is called:

a) Allograft

b) Autograft

c) Isograft

d) Xenograft

Question 22:

Histology of reperfused myocardium:

a) Wavy fibres

b) Neutrophils in cardiac muscle

c) Swelling of cells

d) Eosinophilc contraction band

Question 23:

All-trans retinoic acid is used in the treatment of tumor that is associated with which of the following genetic aberrations?

a) PML-RARA

b) BCR ABL

c) cMYC

d) RUNX1 ETO

Question 24:

A 12-year-old child presented with 15 days of cough. On examination, cervical lymphadenopathy was noted and the lymph node biopsy showed the following features. What is the most probable diagnosis?



a) Tuberculosis

b) Leprosy

c) Sarcoidosis

d) Syphilis

Question 25:

A 30-year-old woman with a long history of dyspnea, chronic cough with sputum, and wheezing died of respiratory failure following a bout of lobar pneumonia. She was not a

smoker or an alcoholic. The lung at autopsy is shown in the image below. Which of the following is most likely to cause the pathological changes shown?



a) Alpha-1-antitrypsin deficiency

b) Cystic fibrosis

c) Mutation in dynein arms

d) Antibodies against type-IV collagen

Question 26:

A 6-foot-5-inch man undergoing a routine physical examination was found to have an early diastolic murmur. His family pedigree chart is given below. Identify the mode of inheritance of this condition.



a) Autosomal dominant

b) Autosomal recessive

c) X-linked

d) Mitochondrial

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	b
5	a
6	a
7	d
8	c
9	b
10	a
11	b
12	b
13	a
14	a

15	a
16	a
17	b
18	a
19	b
20	d
21	c
22	d
23	a
24	a
25	a
26	a

Detailed Explanations

Solution to Question 1:

Huntington's chorea is a trinucleotide repeat disorder.

It is an autosomal dominant neurodegenerative disorder. It is caused by CAG trinucleotide repeats (>40). It occurs due to a gain-of-function (expansion) mutation in the huntingtin gene on chromosome 4p.

The disease shows genetic anticipation and has an early-onset (2-3 decades) if inherited from the father.

Cognitive impairment is an early feature. Pathologically, atrophy of caudate and putamen with ex vacuo ventriculomegaly is seen. Decreased acetylcholine leads to dementia and decreased GABA and increased dopamine lead to chorea.

Treatment consists of haloperidol and tetrabenazine.

Trinucleotide repeat disorders:

Disease	Repeats	Clinical features
Fragile X- syndrome	CGG	Mental retardation, large ears, and jaws, macroorchidism in males
Myotonic Dystrophy(chromosome 19)	CTG	Muscle loss, cardiac arrhythmia,cataracts, frontal balding
Huntington’s disease	CAG	Affective disorder
Spinocerebellar ataxia type 1	CAG	Progressive ataxia, dysarthria, dysmetria.

Disease	Repeats	Clinical features
Friedreich's ataxia	GAA	Progressive ataxia,dysarthria, hypertrophiccardiomyopathy.

Solution to Question 2:

Achalasia cardia is characterized by decreased release of NO(nitric oxide) and VIP(vasoactive intestinal polypeptide) in the distal part of esophagus.

Normally, NO and VIP are secreted by the inhibitory neurons of the Auerbach's plexus in the distal esophagus. This leads to relaxation of the lower esophageal sphincter (LES). In achalasia cardia, these inhibitory neurons degenerate leading to decreased release of NO and VIP. This causes increased LES tone.

The triad of achalasia cardia is as follows:

• Incomplete relaxation of LES

• Increased tone of LES

• Absent peristalsis (aperistalisis)

Solution to Question 3:

The presence of empty appearing nuclei, also called Orphan Annie eye nuclei as seen in the above image is diagnostic of papillary carcinoma of the thyroid.

Microscopic features of papillary thyroid carcinoma include:

• Branching papillae with fibrovascular stalk

• Covering epithelium is simple or stratified cuboidal

• Orphan Annie eye nuclei -

• Empty-appearing nuclei due to dispersed chromatin

• They are diagnostic, regardless of papillary architecture

• Psammoma bodies - concentric calcified structures

• Foci of lymphatic invasion

The image below shows papillary thyroid carcinoma with characteristic Orphan Annie eye nuclei:



Papillary Carcinoma of Thyroid



Psammoma bodies



Note: Orphan Annie refers to a comic book character with empty appearing eyes.

Solution to Question 4:

Based on the given clinical history and microscopy showing crescents, rapidly progressive glomerulonephritis (RPGN) is suspected. Among the given options, anti- GBM antibody testing should be done, since it is a part of the work-up of RPGN.





Type 1 RPGN includes renal limited disease and Goodpasture's disease. In this condition, antibodies are produced against the non-collagenous domain of the α3 chain of collagen IV of the GBM.

Microscopy findings of Goodpasture's disease are:

• Immunofluorescence - linear deposits of Ig and complement along the glomerular basement membrane

• Electron microscopy - wrinkling of the glomerular basement membrane with focal disruptions without any immune complex deposition.

A variant of Goodpasture's disease with pulmonary involvement is known as Goodpasture's syndrome.

Option A: ANA testing is done if SLE is suspected. Electron microscopy will show immune complex deposits.

Option C: HIV RNA testing is done in immunodeficiency syndrome associated with weight loss. Electron microscopy will show diffuse podocyte injury with complete foot process effacement. Tubuloreticular inclusions will be seen in endothelial cells.

Option D: Urine electrophoresis is done if nephrotic syndrome is suspected.

Solution to Question 5:

The given clinical scenario is suggestive of myxoid liposarcoma.

The image shows abundant ground substance, with branching capillaries and a few immature adipocytes.

Liposarcoma is the most common sarcoma in adults and is usually seen in the 5th to 6th decade of life. They commonly occur in the deep soft tissues of the proximal extremities and in the retroperitoneum.

Histologically the liposarcomas are divided into three subtypes:

• Well-differentiated liposarcoma: It consists of mature adipocytes and scattered spindle cells with hyperchromatic nuclei. It is associated with amplification of 12q13-q15.

• Myxoid liposarcoma: It contains abundant basophilic extracellular matrix with branching capillaries. Few immature adipocytes are found scattered within the matrix. It is associated with t(12;16).

• Pleomorphic liposarcoma: It consists of sheets of anaplastic cells with bizarre nuclei and few immature adipocytes.

Among the three subtypes, the pleomorphic liposarcoma is the most malignant and often metastasizes, while the well-differentiated is least malignant. Myxoid liposarcoma has intermediate malignancy behaviour.

Solution to Question 6:

Large, friable, irregular vegetations on heart valves are seen in infective endocarditis.

Infective endocarditis is characterized by the presence of large, friable, irregular vegetations on heart valves. Aortic and mitral valves are the most commonly affected, however, in some cases, the right heart valves may be involved.

Vegetations may be single or multiple and may involve one or more valves. Embolization often occurs and may lead to septic infarcts or mycotic aneurysms at the site of lodgement due to the presence of virulent organisms in the emboli.

Solution to Question 7:

The given findings are suggestive of macrocytic anaemia, which can be seen in chronic alcoholics.

Anemic patients with a peripheral smear showing macrocytic RBCs and hypersegmented neutrophils are important characteristic features of megaloblastic anemia.

Chronic alcoholics also have deficiency of folic acid due to:

• Trapping of folate within the liver

• Excessive urinary loss

• Disordered folate metabolism

• Inadequate dietary intake is frequently encountered in chronic alcoholics

Solution to Question 8:

The clinical features and the histopathological image showing characteristic cystically dilated crypts and glands are suggestive of a juvenile polyp.





Juvenile polyps are malformations of epithelium and lamina propria commonly located in the rectum. It is seen in children <5 years of age. The typical presentation is painless rectal bleeding. Sometimes there may be a proliferating rectal mass.

The disease may be sporadic or syndromic.

• Sporadic juvenile polyp occurs as a single mass and is called retention polyp.

• Juvenile polyposis syndrome is characterized by multiple hamartomatous polyps, which may be premalignant.

Morphologically, both of these types are indistinguishable.

• Gross appearance: They appear as pedunculated, smooth reddish lesions, and with cystic spaces.

• Microscopic appearance: The cysts appear as dilated glands filled with mucin and inflammatory debris.

Note: Complex glandular architecture and the presence of smooth muscle cutting through the lamina propria are seen in Peutz-Jeghers polyps.

Solution to Question 9:

The clinical presentation and the histopathological findings are suggestive of celiac disease.

Celiac disease is an autoimmune gluten-induced enteropathy where HLA DQ2 and DQ8 are expressed. In the small intestine, peptides of gluten are presented to helper T cells which mediate mucosal inflammation. Antiendomysial and anti-tTG antibodies are found in serological studies.

Clinically, malabsorption syndrome with diarrhea, abdominal pain, and marked weight loss is seen along with chronic refractory anemia. Dermatitis herpetiformis is seen in 10% of patients with celiac disease. Microscopic findings show villi atrophy, villi flattening, crypt hyperplasia, and increased intraepithelial T lymphocytes (IEL).

Option A: Environmental enteropathy / tropical sprue is seen in tropical areas with poor sanitation. It is believed to be caused by one or more infectious agents that respond to antibiotics. Mucosa shows variable villous atrophy.

Option C: Whipple's disease is a multisystemic disease caused by T.whipplei. Small intestinal biopsy shows PAS-positive macrophages.

Solution to Question 10:

The given clinical scenario is suggestive of Kawasaki disease or mucocutaneous lymph node syndrome.

Kawasaki disease is an acute febrile illness associated with arteritis of medium to large-sized vessels. It is commonly seen in infants and children <4 years of age.

Clinical features include:

• Fever

• Conjunctival and oral erythema

• Blistering, edema of hands and feet

• Erythema of palms and soles

• Desquamative rash

• Cervical lymphadenopathy

The disease is usually self-limiting. In some cases, the disease may involve coronary artery leading to thrombosis and acute myocardial infarction. Intravenous immunoglobulin and aspirin may be given to reduce the risk of coronary artery disease.

Solution to Question 11:

Effacement of the podocyte foot processes is characteristically seen in minimal change disease (MCD), which is also called lipoid nephrosis.

It is also seen in other conditions such as membranous nephropathy and focal segmental glomerulosclerosis. Since the podocytes are affected in these diseases, they are also known as podocytopathies.

The effacement of foot process can only be observed on electron microscopy.

Solution to Question 12:

Thrombosis is initiated by endothelial injury.

Thrombosis is is the formation of a blood clot inside a blood vessel, obstructing the flow of blood through the circulatory system.

Three primary abnormalities, often referred to as Virchow triad, are known to induce thrombosis. They include:

• Endothelial injury - most important factor

• Stasis or turbulent blood flow

• Hypercoagulability of the blood

Solution to Question 13:

This clinical scenario with the presence of Hurthle cells is suggestive of Hashimoto's thyroiditis.

It is an autoimmune disease, characterized by the gradual destruction of the thyroid gland leading to thyroid failure. It is commonly seen in women between 45 to 65 years of age. It is caused by the presence of circulating autoantibodies against thyroglobulin and thyroid peroxidase.

Morphologically, there is a diffuse enlargement of the thyroid gland, with an intact capsule. It is well demarcated from the adjacent structures.

Microscopic features include:

• Intense lymphocytic infiltrate containing small lymphocytes, plasma cells, and well developed germinal centers

• Atrophied thyroid follicles

• The epithelial cells lining the follicles contain eosinophilic granular cytoplasm

• This is called oncocytic metaplasia

• These cells are called Hurthle cells





Solution to Question 14:

The given findings are suggestive of pernicious anemia, which is associated with atrophic gastritis and vitamin B12 deficiency.

Anemic patient with a peripheral smear showing macrocytic RBCs and hypersegmented neutrophils are important characteristic features of megaloblastic anemia. Vitamin B12 and folic acid deficiency are the main causes of megaloblastic anemia.

Pernicious anemia is an autoimmune disease characterized by the presence of autoantibodies which impairs the production or activity of the intrinsic factor (IF). This is required for the uptake of vitamin B12. There are three types of autoantibodies namely, type I, II, and III. The action of each of these include:

• Autoantibody type I prevents the binding of vitamin B12 to the intinsic factor (IF).

• Autoantibody type II prevents the binding of vitamin B12-IF complex to the ileal receptors.

• Autoantibody type III acts against the α and β subunits of the gastric proton pump, which impairs the production of intrinsic factor.

These patients also develop chronic atrophic gastritis, characterized by loss of parietal cells, infiltration of lymphocytes and plasma cells and megaloblastic changes in mucosal cells.

Solution to Question 15:

The given scenario is suggestive of colloid or mucinous carcinoma of the breast.

Colloid carcinoma is an invasive breast carcinoma, which shows intracellular and extracellular mucin deposition.

On histopathological examination, extracellular type shows tumor cells with faint nuclei, arranged in clusters, floating in lakes or pools of mucin. In intracellular type, nuclei are pushed to the periphery to form a signet ring appearance.





Option B: Microscopic image of papillary carcinoma of the breast showing true papillae with a leaf-like arrangement of fibrovascular tissue lined by the tumor cells is shown below.

Papillary Carcinoma of Breast



Papillary Carcinoma of Breast



Option C: Microscopic image of medullary carcinoma of the breast consisting of sheets of highly pleomorphic cells with mitotic figures (anaplastic cells), lymphoplasmacytic infiltrate is shown below.



Option D: Microscopic image of lobular carcinoma of the breast shows infiltrating small monomorphic discohesive tumor cells arranged in an Indian-file or single-file pattern is shown below.



Solution to Question 16:

Hirschsprung disease occurs due to the failure of migration of the neural crest cells to the bowel wall.

During normal embryogenesis, the neural crest cells migrate from the caecum to the rectum. The enteric neuronal plexuses develop from these migrated neural crest cells.

In Hirschsprung disease, there is an arrest in the migration of these cells, which leads to the development of a distal intestinal segment that is devoid of Meissner (submucosal) and the

Auerbach (myenteric) plexus. Hence this segment is referred to as the aganglionic segment.

As a result, there is a lack of coordination in the peristaltic contractions and functional obstruction, which further leads to the dilation of the intestine proximal to the affected segment, leading to congenital megacolon. Diagnosis is confirmed by demonstrating the absence of ganglion cells in biopsy sections.



Some conditions may cause acquired megacolon. These include:

- Chagas disease, due to the loss of ganglion cells
- Visceral myopathy
- Obstruction by neoplasm or inflammatory stricture
- Toxic megacolon complicating ulcerative colitis
- Functional psychosomatic disorders

Solution to Question 17:

Premature aging in Werner syndrome is due to a defect in DNA helicase.

DNA helicase is involved in DNA replication and repair and other functions requiring DNA unwinding. A defect in this enzyme causes rapid accumulation of damaged chromosomes that may mimic the injury that normally accumulates during cellular aging.

Solution to Question 18:

The histopathological examination is suggestive of schwannoma. Features showing spindle cells with palisading nuclei and Verocay bodies are characteristic of schwannoma.

Schwannomas are well-circumscribed, encapsulated masses that abut the associated nerve without invading it. They are commonly associated with inactivating mutations in the NF2 gene on chromosome 22q12.

Microscopically, it consists of dense areas called Antoni A, and loose areas called Antoni B.

• The Antoni A areas contain spindle cells that are arranged into intersecting fascicles. Palisading of nuclei is seen, with nuclear-free zones between them, forming structures called Verocay bodies.

• The Antoni B areas contain spindle-shaped cells that are loosely arranged separated by prominent myxoid extracellular matrix.

The image below shows cellular Antoni A and paucicellullar Antoni B areas in schwannoma.



The image below shows verocay bodies in the Antoni-A area of vestibular schwannoma.



Solution to Question 19:

IL-4 stimulates B-cells to differentiate into IgE secreting plasma cells.

Solution to Question 20:

The change described in the question is called metaplasia. It is defined as a reversible change in which one type of differentiated cell is replaced by another type.

The table below shows various types of metaplasia:

GERD: Gastroesophageal reflux disease

Tissue	Metaplasia	Cause
Airways	Squamous metaplasia: Normal pseudostratified epithelium is replaced by the squamous epithelium.	Chronic smoking,Deficiency of retinoic acid
Urinary bladder	Squamous metaplasia: Normal transitional epithelium is replaced by squamous epithelium.	Bladder stones
Esophagus	Columnar metaplasia: Normal squamous epithelium is replaced by the glandular columnar epithelium (Barret's esophagus)	Chronic GERD
Cervix	Squamous metaplasia: Normal glandular epithelium may be replaced by the squamous epithelium.	Low pH of the cervix
Muscle	Connective tissue metaplasia: Formation of bone in the muscle tissue; (myositis ossificans).	After intramuscular hemorrhage

Solution to Question 21:

A graft from an identical twin is called an isograft.

Isograft is defined as an organ or tissue that is taken from a donor and is transplanted to a recipient who has the same genetic constitution as the donor. For example, graft between identical twins.

Solution to Question 22:

Reperfused myocardium shows eosinophilic contraction bands on light microscopy. They occur in irreversibly injured myocytes with damaged membranes, when blood supply is restored 12-24 hours after MI.

During ischemia, the intracellular levels of calcium are increased due to impaired calcium cycling and sarcolemmal damage. This leads to augmented and uncontrolled contraction of myofibrils after reperfusion.

On microscopic examination, the irreversibly injured myocytes exhibit contraction bands that are intensely eosinophilic intracellular stripes. These contraction bands are composed of closely packed sarcomeres.

Reperfused infarcts are usually hemorrhagic because the vasculature is injured during ischemia. This leads to bleeding after the flow is restored.

Solution to Question 23:

All-trans retinoic acid (ATRA) is used in the treatment of acute promyelocytic leukemia (APML), which is associated with PML-RARA fusion gene.

Acute promyelocytic leukemia (APML) is associated with a reciprocal translocation between chromosomes 15 and 17 producing a PML-RARA fusion gene. This fusion gene encodes for a chimeric protein, which consists of a part of the PML protein and a part of the retinoic acid receptor-α (RARα).

The normal RARα binds to the DNA and activates the transcription in the presence of retinoids. However, the chimeric protein produced by PML-RARA has reduced affinity for retinoids. This causes inhibition of transcription, thereby preventing gene expression for cellular differentiation. This leads to the accumulation of myeloid progenitor cells.

The condition is treated with all-trans retinoic acid. It binds to the PML-RARA chimeric protein. This causes conformational change such that the protein regains its normal function and activates DNA transcription.

Solution to Question 24:

The histopathological picture showing the caseating granuloma, eosinophilic necrotic debris known as caseation, bordered by collections of epithelioid cells, Langhan giant cells (horse shoe shape) and lymphocytes and clinical history is suggestive of tuberculosis.

Clinically, tuberculosis presents with chronic cough, fever, weight loss, swollen lymph nodes.

Other options:

Option B- Leprosy presents with skin lesions and nerve involvement with loss of sensation.

Option C- Sarcoidosis shows giant cells with cytoplasmic inclusions, non-caseating granuloma with fibroblastic proliferation at the periphery.

Option D- Syphilis shows thickened capsule/ hyperplasia of capsule accompanied by fibrosis, perivascular inflammation, and infiltration of chronic inflammatory cells.

The below image shows histopathological findings seen in tuberculosis



Solution to Question 25:

The above gross specimen shows emphysematous damage involving enlargement of the entire acinus suggestive of panacinar emphysema. It is caused due to alpha-1-antitrypsin deficiency.

In panacinar (panlobular) emphysema, the acini are uniformly enlarged from the level of the respiratory bronchiole to the terminal blind alveoli. It occurs commonly in the lower zones and in the anterior margins of the lung and is most severe at the bases.

The image below shows panacinar emphysema.

Solution to Question 26:

The patient's height and the presence of early diastolic murmur in the given clinical scenario are suggestive of Marfan syndrome. It is inherited in an autosomal dominant disorder. The family pedigree confirms the mode of inheritance.

The following flow chart helps in making out the pattern of inheritance from a given pedigree chart.



Patients with Marfan syndrome have pleiotropic manifestations:

• Skeletal systems - Tall and decreased upper to lower segment ratio with long arms, hands, and fingers

• Deformities of the chest wall, vertebral column, and foot arches

• Cardiovascular abnormalities - mitral valve prolapse leading to mitral regurgitation and dilatation of aortic root leading to aortic regurgitation, aortic dissection, and rupture.

• Ocular - superio-temporal lens dislocation

Pathology INI-CET 2021

Question 1:

What is the cell of origin of Chronic Lymphocytic Leukaemia/Small Lymphocytic Lymphoma?

a) Mature B cells

b) Mature T cells

c) Progenitor T cells

d) Naive B cells

Question 2:

Human Papilloma Virus inactivates RB gene by which of the following proteins?

a) E6

b) E7

c) L1

d) L2

Question 3:

In cystic fibrosis, which of the following structure is affected in pancreas?



a) A

b) B

c) C

d) D

Question 4:

Mediators involved in the early phase of type 1 hypersensitivity are?

a) 1, 5

b) 1, 2, 3, 4,

c) 1, 2, 4, 5

d) 2, 3, 4

Question 5:

Given below is an image of activation of the coagulation cascade, following closed endothelial injury. Identify the factor 'L' which combines with tissue factor to activate factor IX.



a) Factor XII

b) Factor VII

c) Factor VIII

d) vWF

Question 6:

The vacutainer tube shown in the given image contains:

a) Na-EDTA

b) Trisodium citrate

c) NAF-heparin

d) Clot activator

Question 7:

Arrange the following cellular events of inflammation in the correct sequence:

a) 1, 2, 3, 4

b) 2, 1, 4, 3

c) 4, 1, 2, 3

d) 3, 4, 1, 2

Question 8:

Which of the following is used in the treatment of sideroblastic anemia?

a) Vitamin B12

b) Vitamin B6

c) Vitamin B1

d) Iron

Question 9:

A 5-year-old child who presented with proptosis of one of the eyes was found to have a desmin positive tumour. What is the probable diagnosis?

a) Ewing's sarcoma

b) Embryonal rhabdomyosarcoma

c) Leukemia

d) Retinoblastoma

Question 10:

A 17-year-old boy presented with intussusception and was taken up for surgery. He was found to have multiple polyps. The resected section was sent for histopathology and the whole mount image of the specimen is given below. Identify the lesion.



a) Tubulovillous adenoma

b) Hamartomatous polyp

c) Adenocarcinoma

d) Hyperplastic polyp

Question 11:

Which of the following is true?

a) 1, 2

b) 1, 3

c) 2, 3

d) All of the above

Question 12:

Identify the given condition.



a) Pemphigus vulgaris

b) Leprosy

c) Mycosis fungoides

d) Visceral leishmaniasis

Question 13:

A karyotyping image is given below. What are the clinical features most likely to be expected in such patients?



a) Gynecomastia and long limbs

b) Flat occiput

c) Webbing of neck and lymphedema

d) Short stature and polydactyly

Question 14:

A 20-year-old athlete suddenly collapsed while playing football and was taken to the hospital where he was declared dead. His autopsy image is given below. Which of the following is the most likely cause of death?



a) Valvular heart disease due to RHD

b) Sudden cardiac death due to arrhythmia

c) Sudden cardiac death due to Hypertrophic cardiomyopathy

d) Sudden cardiac death due to atherosclerosis

Question 15:

Which of the following can be prevented by transfusing irradiated RBCs?

a) Graft versus host disease

b) HLA Alloimmunization

c) Immunomodulation

d) Transfusion Related Acute Lung Injury (TRALI)

Question 16:

Which of the following statements are correct regarding the effect of Glucose-6-Phosphate Dehydrogenase deficiency?

a) Only 1 is correct.

b) 1 & 2 are correct.

c) 2 & 3 are correct.

d) 1, 2 & 4 are correct.

Question 17:

Which of the following is not likely to be seen in a patient with Paroxysmal Nocturnal Hemoglobinuria (PNH)?

a) Aplastic anemia

b) Leukemia

c) Thrombosis

d) Hemolysis

Question 18:

Given below is an electron microscopic image of the lung tissue. A patient infected with SARS-COV2 presents with ARDS. The infection would affect and expand which of the labeled

layers?



a) B

b) A

c) C

d) D

Question 19:

What is the gold standard investigation for the diagnosis of myocarditis?

a) Endo-myocardial biopsy

b) Magnetic Resonance Imaging

c) Brain Natriuretic Peptide

d) LDH1 to LDH2 ratio

Question 20:

A child presented with bilateral proptosis. From the peripheral smear, he was suspected to have chloroma. What is the next investigation of choice?

a) Platelet count

b) WBC count

c) Flow cytometry

d) Bone marrow biopsy

Question 21:

A person working in an industry for a long time developed progressive breathlessness. The histopathological image of his lung biopsy is given below. What industry was he most likely working in?



a) Silica

b) Asbestos

c) Coal dust

d) Cotton

Question 22:

A patient presented with complaints of dyspnea on exertion for 1 month. Echocardiography showed the presence of mitral stenosis and left ventricular hypertrophy. Identify the histopathological finding given below and the probable diagnosis.



a) Aschoff body; Rheumatic heart disease

b) Calcification of valve; Mitral stenosis

c) Granuloma; Sarcoidosis

d) Non-caseating granuloma; Tuberculosis

Question 23:

A patient with lytic lesions on the skull is suspected of a diagnosis of Langerhan cell histiocytosis. Which of the following is a characteristic finding on electron microscopy?

a) Histiocytes

b) Birbeck granules

c) Eosinophils

d) Giant cells

Question 24:

A 60-year-old patient presented with scrotal swelling and ataxia. Which among the following could likely be a cause?

a) Glioblastoma multiforme

b) Hodgkin lymphoma

c) Non-Hodgkin lymphoma

d) Non seminomatous germ cell tumor

Question 25:

A patient with cirrhosis and ascites presented to you. The gross appearance of his liver is as shown below. Which among the following is true about it?



a) Dark areas represent necrosis, white areas represent viable liver tissue

b) Dark areas represent viable liver tissue, white areas represent necrosis

c) White areas represent central hepatocytes

d) Dark areas represent peri-portal hepatocytes

Question 26:

A 6-year-old boy came with bilateral cervical lymphadenopathy. What is the probable diagnosis, cell, and virus associated with the image given below?



a) Non-Hodgkin's lymphoma, RS cell, EBV

b) Kaposi sarcoma, Giant cell, HIV

c) Hodgkin's lymphoma, RS cell, EBV

d) Infectious mononucleosis, RS cell, CMV

Question 27:

A person was brought to the emergency department with facial swelling, itching, and hypotension following a bee sting. The mentioned clinical features are due to the increase in which immunoglobulin?

a) IgA

b) IgE

c) IgM

d) IgG

Question 28:

All of the following are associated with massive splenomegaly except :

a) Chronic myeloid leukemia

b) Aplastic anemia

c) Hairy cell leukemia

d) Lymphoma

Question 29:

A 42-year-old female patient presents with cough, low-grade fever, and hemoptysis. Investigations reveal a cavitary lesion on her right lung apex, which on biopsy reveals caseous necrosis. The underlying pathophysiology is :

a) Enzyme degradation

b) Type 4 hypersensitivity reaction

c) Fibrinoid deposition

d) Sudden cut-off of blood supply

Question 30:

A 45-year-old male patient was diagnosed with iron deficiency anemia. Which of the following is true about intracellular regulation of iron in the above condition?

a) Ferritin translation is increased by binding IRP to IRE

b) Ferritin translation is decreased by binding IRP to IRE

c) Transferrin receptor 1 translation is unaffected by binding IRP to IRE

d) Transferrin receptor 1 translation is decreased by binding IRP to IRE

Question 31:

A 65-year-old male patient with chronic fatigue and lymphadenopathy presents to the outpatient department. His peripheral smear shows the presence of smudge cells. Which of the following additional investigations will you do to arrive at a diagnosis?

a) Polymerase chain reaction

b) Fluorescent in situ hybridization

c) Flow cytometry

d) Cytogenetics

Question 32:

Oil red O stain is used in which of the following?

a) Formalin fixation

b) Frozen section

c) Alcohol fixation

d) Gluteraldehyde fixation

Question 33:

A boy presents with fever, night sweats, and neck swelling. The biopsy of swelling showed a starry sky appearance. What is the most likely genetic abnormality seen in this case?

a) MYC

b) RAS

c) BCR-ABL

d) p53

Question 34:

An electron microscopic picture of a cut section of a nerve fiber is given. Identify the area which will be affected in the most common variant of Guillain-Barré syndrome.



a) 1

b) 2

c) 3

d) 4

Answer Key

Question No.	Correct Option
1	d
2	b
3	b
4	c
5	b
6	d
7	a
8	b
9	b
10	b
11	c
12	a
13	a
14	b
15	a
16	b
17	b
18	b
19	a
20	d
21	b
22	a
23	b
24	c
25	a
26	c
27	b
28	b
29	b
30	b
31	c
32	b

33	a
34	b

Detailed Explanations

Solution to Question 1:

The cell of origin of Chronic Lymphocytic Leukaemia (CLL)/ Small Lymphocytic Lymphoma (SLL) is either naive B cells or post-germinal centre memory B cell.

The CLL and SLL differ only in the degree of peripheral blood lymphocytes. CLL is the most common leukemia of adults in the western world. The median age of diagnosis is 60 years. The most common genetic anomalies seen are deletions of 13q14.3, 11q, and 17p and trisomy 12q. Loss of tumor suppressor genes in chromosome 13 results in overexpression of the anti-apoptotic protein BCL2.

The growth of CLL/SLL cells is confined to the proliferation centers (pathognomonic for CLL/SLL), where the tumor cells receive critical factors from the surrounding environments, such as NF-κB for cell survival, MYC for cell growth, Bruton Tyrosine Kinase (BTK) signaling pathway, etc.,

The tumor cells express CD19, CD20, CD23, and CD5. Smudge cells are also characteristic of CLL. Clinical features include easy fatiguability, weight loss, anorexia, generalized lymphadenopathy, hepatomegaly, hemolytic anemia, etc. CLL/SLL has a tendency to transform to a more aggressive diffuse large B-cell lymphoma (DLBCL) - this is called Richter syndrome.

Factors with a bad prognosis include:

• Presence of deletions of 11q and 17p.

• Lack of somatic hypermutation.

• Expression of ZAP-70, a protein that augments signals produced by the Ig receptor.

• Presence of NOTCH1 mutations

BTK inhibitors and BCL2 inhibitors produce sustained therapeutic responses in a higher fractions of people with CLL/SLL.

Solution to Question 2:

E7 protein of Human Papilloma Virus (HPV) binds to the hypophosphorylated RB protein and displaces the E2F transcription factors, thereby promoting G1/S transition.

RB gene is a key negative regulator of the cell cycle at the G1/S transition. RB gene exists in an active hypophosphorylated state in resting cells and in an inactive hyperphosphorylated state in cells undergoing G1/S transition.

E7 protein from the high-risk types of HPV (such as HPV 16) binds to the hypophosphorylated RB through the same pocket that RB uses to bind and sequester the E2F transcription factors, thereby

displacing the E2F which leads to the progression of the cell cycle and thus the proliferation of tumor cells. E7 proteins also inactivate the CDK inhibitors p21 and p27. E7 proteins from high-risk HPVs (types 16,18 and 31) also bind and presumably activates cyclins A and E.

Oncogenic activities of protein E6:

The E6 protein binds to and helps in the degradation of p53 and stimulates the expression of telomerase reverse transcriptase (TERT), which is the catalytic subunit of telomerase, thereby helping in the longevity of the cells. E6 from high-risk HPV types has a higher affinity for p53 than E6 from low-risk types.



Solution to Question 3:

The part labeled 'B' in the image is the exocrine pancreas which is affected in cystic fibrosis.

Features of cystic fibrosis in the pancreas:

• Accumulation of mucus in the ducts leads to dilation of exocrine glands.

• Complete blockage of ducts with mucus leads to atrophy of the exocrine pancreas and progressive fibrosis

• Due to atrophy, there will be loss of exocrine secretions, which in turn leads to impaired absorption of fat and fat-soluble vitamins

• Associated loss of vitamin A leads to squamous metaplasia of the pancreatic duct lining epithelium



Other options:

A- Endocrine pancreas

C- Pancreatic ducts

D- Blood vessels

Solution to Question 4:

The mediators involved in the early phase of type 1 hypersensitivity are,

• Mast cells,

• Histamine,

• Prostaglandins (D2) and

• Leukotrienes (C4 and D4).

Early phase of type 1 hypersensitivity:

The mediators produced by the mast cells are responsible for most of the immediate hypersensitivity reactions. Histamine and leukotrienes are released rapidly from the sensitized mast cells and trigger intense immediate reactions characterized by edema, mucus secretion and smooth muscle spasm.

Other mediators involved in the early phase are,

• Enzymes: neutral proteases (chymase, tryptase) and acid hydrolases

• Proteoglycans: heparin and chondroitin sulfate

• Lipid mediators: Platelet-activating factor (PAF)

Late-phase of type 1 hypersensitivity:

The late phase reaction is characterized by the recruitment of leukocytes, that amplify and sustain the inflammatory response without exposure to the triggering antigen.

Mediators involved in the late phase of type 1 hypersensitivity reaction are:

• Cytokines: TNF, IL-1, IL-4, IL-5, and chemokines (eotaxin)

• Leukocytes (eosinophils)

• Major Basic Protein (MBP)

• Eosinophil cationic protein



Solution to Question 5:

The factor marked as 'L' is factor VII, which combines with tissue factor that is produced following a close endothelial injury to activate factor IX.

As the image represents the in vivo coagulation cascade, the tissue factor is the major initiator of coagulation. It is believed that in vivo, factor VIIa/tissue factor complex is the most important activator of factor IX and that factor IXa/factor VIIIa complex is the most important activator of factor X.

Tissue factor is expressed on the surfaces of the subendothelial cellular components of the vessel wall, such as smooth muscle cells and fibroblasts. This tissue factor is exposed to blood when there is a vascular damage. This activates the in vivo extrinsic pathway of coagulation cascade, which is as follows:

• Tissue factor binds with serine protease factor VIIa

• This complex then activates factor IX to factor IXa

• Factor IXa then activates factor X to factor Xa

• Factor Xa can be formed through the actions of either the tissue factor-factor VIIa complex or factor IXa (with factor VIIIa as cofactor)

• Factor Xa then converts prothrombin to thrombin, which is the pivotal protease of the coagulation cascade.

• The cofactor essential for this conversion is factor Va.

• The thrombin then formed converts the soluble plasma fibrinogen to an insoluble fibrin matrix.

• Thrombin also activates factor XIII, which is the fibrin stabilizing factor, to factor XIIIa. This factor XIIIa forms covalent cross-links and thereby stabilizes the fibrin clot.

The below image shows the in vivo coagulation cascade.



Solution to Question 6:

The plastic tube shown in the given image has a red stopper, hence it contains clot activator.

Red stopper plastic tubes contain silica as a clot activator. They are used for serum chemistry tests, serology tests, and in blood banks, where both serum and red blood cells may be used.

The tubes are inverted five times to initiate the clotting process.

Note: Red stopper glass tubes are often referred to as clot or plain tubes because they contain no anticoagulants or additives.

Some important plastic tubes and their additives:

Cap colour	Additive present
Light blue top	Sodium citrate
Dark green to p	Heparin
Purple top	EDTA
Pale yellow to p	Acid citrate dextrose

Cap colour	Additive present
Light grey top	Sodium fluoride and potassium oxalate

Solution to Question 7:

The correct sequence of cellular events that occurs during inflammation is as follows: Rolling, Cytokine-mediated integrin activation, Adhesion, Migration.

The recruitment of leucocytes to the site of inflammation is a multi-step process that is mediated by adhesion molecules and cytokines called chemokines. This process can be divided into sequential phases:

• In the lumen: margination, rolling and adhesion to the endothelium occurs.

• Migration of the leucocytes across the endothelium and vessel wall

• Migration in the tissues towards a chemotactic stimulus.

The process of leucocyte recruitment is as follows:

• Margination: Because of the dilation of the inflamed postcapillary venues, there is stasis of blood flow, and hence, more leucocytes assume a peripheral position along the endothelial surface; this process of leucocyte redistribution is called margination.

• Rolling: The slowed down leucocytes then sense the signals from the endothelium and start rolling on the vessel wall.

• Cytokine-mediated integrin activation: The attachment of leucocytes to the endothelial cells requires adhesion molecules that are activated by cytokines. The major families of proteins involved in adhesion and migration are selectins and integrins and their ligands.

• Adhesions: The weak rolling interactions slow down the leucocytes and allow them to bind more firmly to the endothelium. Chemokines that were produced at the site of injury bind to the endothelial cell proteoglycans. These chemokines then bind to the rolling leucocytes and activate them. This, in turn, converts the ligands for β1 and β2 integrins, VLA-4, and LFA-1 present on the leucocytes to a high-affinity state.

• This combination of cytokine-induced expression of integrin ligands on the endothelium and increased integrin affinity on the leucocytes leads to firm adhesion of leucocytes to the endothelium at the site of inflammation.

• Leucocyte migration: This is the movement of leucocytes through the intact endothelium, called transmigration or diapedesis. This occurs in postcapillary venules. This migration occurs through the inter-endothelial gap towards the chemical concentration gradient created by the chemokines at the site of injury. This occurs with the help of adhesion molecules present in the intercellular junctions such as platelet endothelial cell adhesion molecule (PECAM-1) or CD31.



Solution to Question 8:

Vitamin B6 (Pyridoxine) is used in the treatment of sideroblastic anaemia.

Pyridoxine 200mg per day for 2-3 months is administered routinely to all cases of sideroblastic anaemia. This cofactor is essential for ALA-synthase activity, and mutations in the pyridoxine binding site of the enzyme have been found in several responsive patients. Blood transfusion and other supportive therapy are also indicated. In cases of secondary sideroblastic anaemia, the primary focus is the removal of the offending agent. But no definitive treatment is available for hereditary and idiopathic types of sideroblastic anaemia.

Sideroblasts are nucleated red cells (normoblasts) containing siderotic granules which stain positively with Prussian blue stain. Based on aetiology, sideroblastic anaemia are classified into hereditary and acquired types.

Hereditary sideroblastic anaemia is an X-linked disorder caused due to a defect in the activity of aminolevulinic acid (ALA) synthetase required for heme synthesis. Affected females act as carriers while males have moderate to marked anaemia and manifest in childhood or during early adulthood

Acquired sideroblastic anaemia:

I) Primary acquired:

• Also called idiopathic or refractory acquired sideroblastic anaemia

• Occurs in middle-aged and older individuals of both sexes

• Due to the reduced activity of the enzyme ALA synthetase, there is disturbed growth and maturation of the erythroid precursors at the level of hematopoietic stem cell

• Moderate to severe anaemia, which appears insidiously

• Neutropenia and thrombocytopenia may also be seen

• Regarded as a myelodysplastic disorder in the FAB classification and thus considered preleukaemic

II) Secondary acquired:

May appear secondary to

• Drugs, chemicals and toxins: Isoniazid, Cycloserine, Chloramphenicol, Cyclophosphamide, Alcohol, Lead, etc.,

• Haematological disorders: myelofibrosis, polycythemia vera, acute leukaemia, myeloma, lymphoma and hemolytic anaemia.

• Miscellaneous: Carcinoma, myxoedema, rheumatoid arthritis and SLE.

Laboratory findings:

• Peripheral smear: Hypochromic, microcytic (sometimes normocytic) anaemia

• Absolute values: MCV, MCH and MCHC are reduced except in acquired type, where MCV is raised.

• Bone marrow: Erythroid hyperplasia with macronormoblastic erythropoiesis. Marrow iron stores are raised. Pathognomonic ring sideroblasts are present.

• Serum ferritin: Raised

• Serum iron: Raised

• TIBC: Normal

• Haemoglobin electrophoresis: Normal

Solution to Question 9:

The clinical picture of a 5-year-old child with proptosis and a desmin positive tumor suggests a diagnosis of embryonal rhabdomyosarcoma.

Rhabdomyosarcoma is a malignant mesenchymal tumor with skeletal muscle differentiation. There are four subtypes: alveolar, embryonal, pleomorphic, and spindle cell/sclerosing. Rhabdomyosarcoma is the most common soft tissue sarcoma of childhood and adolescence and is usually of the alveolar and the embryonal variety. Pleomorphic rhabdomyosarcoma is seen predominantly in adults. Spindle cell/sclerosing type affects all ages. The pediatric forms often arise in the sinuses, head and neck, and genitourinary tract.

Embryonal rhabdomyosarcoma has sheets of both primitive round and spindled cells in a myxoid stroma. Rhabdomyoblasts with visible cross-striations may be present. Sarcoma botryoids is a variant of embryonal rhabdomyosarcoma in the walls of hollow, mucosal-lined structures, such as the nasopharynx, common bile duct, bladder, and vagina with cambium layer. Alveolar rhabdomyosarcoma has a network of fibrous septae that divide the cells into clusters or aggregates, creating a crude resemblance to pulmonary alveoli.

Clinically, the tumor is usually aggressive, with unilateral proptosis when the orbit is involved. Swelling and redness of the overlying skin, diplopia may be present. Immunohistochemistry is positive for desmin, myoglobin, myogenin, MyoD1, and vimentin

Treatment involves surgical resection and chemotherapy, with or without radiotherapy. The best prognosis is for the botryoid variant while the pleomorphic has variant has the worst prognosis.

Other options:

A) Ewing's sarcoma: CD99 positive

C) Leukaemia: CD45 positive

Solution to Question 10:

The given clinical feature of intussusception and multiple polyps and the histologic picture showing an arborizing pattern of connective tissue, smooth muscle, lamina propria, and glands are suggestive of Peutz–Jeghers Syndrome, which is a type of hamartomatous polyp of the small intestine.

It is an autosomal dominant condition characterized by hamartomatous polyps of the intestinal tract and melanosis of the mouth and the lips.

Clinical features:

• Hamartomatous polyps of the intestinal tract

• Hyperpigmentation of the buccal mucosa, lips, perianal skin, and digits

• Increased (2% to 10%) risk for cancer of the intestinal tract, with cancers reported throughout the intestinal tract, from the stomach to the rectum.

• Increased risk for extraintestinal malignant neoplasms, including cancer of the breast, ovary, cervix, fallopian tubes, thyroid, lung, gallbladder, bile ducts, pancreas, and testicles.

Management:

• Surgery is indicated in cases where there are bleeding, obstruction, intussusception, or adenomatous changes of the polyp.

• If surgery is done for these indications attempts should be made to remove as much as polyps as possible.

• Regular screening with endoscopic studies should be done due to the increased risk of malignancies.

• Screening of other malignancies like breast and cervix are also important.

Other options:

A) Tubulovillous adenoma:

These are a mixture of tubular and villous adenoma and consist of small, pedunculated polyps composed of rounded or tubular glands and larger, sessile polyps which are covered by slender villi. These adenomas are characterized microscopically by the presence of epithelial dysplasia. The hallmarks of epithelial dysplasia are nuclear hyperchromasia, elongation and stratification.



C) Adenocarcinoma:

Presents with occult bleeding, changes in bowel habits, or cramping and left lower quadrant discomfort. Morphologically, they present as polypoid, exophytic masses or annular lesions that produce ‘napkin-ring’ construction and luminal narrowing. Microscopically, these tumours consist of tall columnar cells with elongated, hyperchromatic nuclei and necrotic debris in the gland lumen.



D) Hyperplastic polyp:

Hyperplastic polyps, most commonly found in the left colon are smooth, nodular protrusions of the mucosa, often on the crests of mucosal folds. Histologically, they are composed of mature goblet and absorptive cells with a serrated surface architecture.



Solution to Question 11:

Of the given statements, 2 and 3 are true. BRCA1 is a tumor suppressor gene and MYC is a proto-oncogene.

BRCA1 is a tumor suppressor gene located on chromosome 17q21 and its function is to repair the double-stranded breaks in DNA. It is associated with familial breast and ovarian carcinoma. BRCA1-associated breast cancers are commonly poorly differentiated.

All breast cancers are separated into 3 major groups based on the expression of two proteins, ER and HER2 (also known as ERBB2). HER2 cancers are defined as cancers overexpressing HER2 and can be either ER-positive or ER-negative. Thus HER2neu is expressed only in a fraction of people with breast cancer.

EGF-receptor family has two proto-oncogenes: ERBB1 that codes for EGFR and ERBB2 that codes for HER. Mutation in EGFR results in adenocarcinoma of the lung, which is one of the main subtypes of non-small cell lung carcinoma.

MYC is a proto-oncogene, that is expressed in all eukaryotic cells and belongs to the immediate-early response genes, which are rapidly induced by RAS/MAPK signaling following growth factor stimulation of quiescent cells. Normally, MYC protein concentration are tightly controlled at the level of transcription, translation and protein stability. The mechanisms by which MYC promotes neoplastic cell growth may be follows:

• MYC activates the expression of many genes like D cyclins, that are involved in cell cycle progression

• MYC regulates the expression of telomerase

• MYC is one of the transcriptional factors that can act together to reprogram somatic cells into pluripotent stem cells

MYC gene is deregulated in a number of cancers such as Burkitt lymphoma, breast, colon, lung and other cancers. The functionally identical NMYC and LMYC genes are also amplified in

neuroblastomas and small cell cancers of the lung, respectively.

Solution to Question 12:

The given histologic picture shows a suprabasal blister suggesting the diagnosis of pemphigus vulgaris.

It is caused due to the production of antibodies mainly against desmoglein 3. Rarely antibodies against desmoglein 1 can also be found. The antibodies are commonly IgG antibodies.

Desmoglein 3 is mainly present in the basal and suprabasal layers of the skin and mucosa. The patients, therefore, present with flaccid bullae over the skin and mucosal ulcerations. Oral mucosa is commonly affected. Other areas where ulceration can be seen are conjunctiva, nasopharynx, larynx, etc. Cutaneous lesions are more common over the scalp, face, neck, upper chest, and back.

Direct immunofluorescence will show the deposition of antibodies around the keratinocytes in the epidermis. It is treated with steroids, azathioprine, mycophenolate mofetil, rituximab, etc.



Other options:

B) Leprosy: Tuberculoid leprosy shows granulomatous lesions, without much bacilli. Hence called paucibacillary leprosy. Lepromatous lesions contain large aggregates of lipid-laden macrophages (lepra cells), often filled with masses (globi) of acid-fast bacilli. Because of the presence of abundant bacteria, it is also called as multibacillary.

Given below are the images that depict lepromatous lesion showing foamy macrophages with globi of pink/red acid-fast bacilli and tuberculoid leprosy showing granulomatous lesions, respectively.





C) Mycosis fungoides: Mycosis fungoides is a cutaneous T-cell lymphoma, characterized by polymorphic patches and plaques with fine scaling usually present on the limbs, breast, and buttocks. The lesions may be pruritic or even asymptomatic. Histological hallmarks include: 1. Sézary-Lutzner cells - cells with cerebriform nuclei (infolded/ hyper-convoluted nuclear membrane), 2. Pautrier's microabscesses - tumor cells invade the epidermis as single cells and small clusters, forming a collection of atypical lymphocytes in the epidermis and 3. Epidermotropism - pronounced invasion of the tumor cells into the epidermis.

Given below is the image of Sezary-Lutzner cells.



Given below is the arrow pointing at Paurtier's microabcess.



D) Visceral leishmaniasis: In visceral leishmaniasis, parasites invade and activate the macrophages throughout the mononuclear phagocyte system. Hence, these phagocytic cells are enlarged and filled with leishmania species; many plasma cells are also present.

The given slide shows macrophages infested with intracellular Leishmania donovani bodies (arrows) characterized by a kinetoplast and characteristic double dot appearance (Leishman stain).



Solution to Question 13:

The given karyotyping image shows '47, XXY' abnormality, which is seen in Klinefelter's syndrome. Gynecomastia and long limbs are seen in Klinefelter's syndrome.

Other clinical features seen in Klinefelter's syndrome are:

• Disproportionately long arms and legs,

• Gynecomastia

• Eunuchoid body habitus

• Mitral valve prolapse

• Mental retardation; IQ: 60-80,

• Minor developmental and learning disabilities,

• Lack of secondary sexual characteristics (due to a decrease in androgen production)

• Higher risk of breast carcinoma

• Testicular dysgenesis with small penis

• Infertility/azoospermia

• Type 2 diabetes mellitus

• Metabolic syndrome

• Higher risk of extragonadal germ cell tumors

Klinefelter syndrome:

• Most common sex aneuploidy in males

• Classically karyotype is 47, XXY. Other variants can have 48 XXXY; rarely, 49 XXXXY; or mosaics with some cells containing normal 46, XY and others 47, XXY

• Results from meiotic nondisjunction of sex chromosomes (mostly during 1st meiotic division)

Note:

A variant of Klinefelter's syndrome—49, XXXXY—presents with short stature





Other options:

B) Flat occiput:

It is usually seen in Down's syndrome or trisomy 21.

Clinical features: (Mnemonic: I C A PROBLEM Somewhere)

• Intellectual disability/Incurved or short fifth finger with clinodactyly

• Congenital heart disease

• Acute megakaryoblastic leukemia/Alzheimer's disease/Atlantoaxial instability/Atresia of the duodenum

• Protruding tongue

• Round face

• Occiput flat/Open wide fontanelle/Open mouth

• Brachycephalic/Brushfield's spots

• Low (depressed) nasal bridge

• Epicanthal fold/Ears low set and dysplastic

• Mongoloid slant (oblique palpebral fissure)

• Simian palmar crease and a wider gap between first and second toes

C) Webbing of neck and lymphedema:

It is usually seen in Turner's syndrome.

Important features associated with Turner's syndrome: (mnemonic: See A Baby CLOWN Here)

• Short stature, Short fourth metacarpal, Sensorineural hearing loss, Shield chest

• Amenorrhea (primary), infertility

• Barr body absent

• Cardiac defects, Cubitus valgus, Congenital lymphedema

• L: Learning disabilities

• Ovaries underdeveloped (streak ovaries), Gonadoblastoma (increased risk if Y chromosome material is present)

• Webbed neck

• Nipples widely spaced

• Horseshoe kidneys

D) Short stature and polydactyly:

Short stature with polydactyly is most commonly seen in Patau syndrome or trisomy 13.

It's clinical features include:

• Cleft lip often midline

• Flexed fingers with postaxial polydactyly

• Ocular hypotelorism

• Bulbous nose

• Low-set, malformed ears

• Microcephaly

• Cerebral malformation, especially holoprosencephaly

• Microphthalmia

• Cardiac malformations

• Scalp defects

• Hypoplastic or absent ribs

• Visceral and genital anomalies



Solution to Question 14:

The given image shows asymmetric septal hypertrophy and along with the scenario, it is suggestive of sudden cardiac death due to arrhythmia. Patients with hypertrophic cardiomyopathy have an increased risk of sudden cardiac death from ventricular tachyarrhythmias. In fact, arrhythmias are the most important/most common cause of death in hypertrophic cardiomyopathy.

Hypertrophic cardiomyopathy is one of the most common causes of sudden, otherwise unexplained death in young athletes. HCM is characterized by reduced stroke volume due to impaired diastolic filling, which is a consequence of the massively hypertrophic left ventricle. There is a dynamic obstruction to the left ventricular outflow as the anterior mitral leaflet moves towards the ventricular septum during systole. This compromised cardiac output in combination with a secondary increase in pulmonary venous pressure is the reason for exertional dyspnea at times of increased oxygen demand.

Major clinical problems in HCM are atrial fibrillation, stroke, intractable cardiac failure, ventricular arrhythmias and also sudden death. There is a harsh systolic ejection murmur caused by the ventricular outflow obstruction. Mutations causing hypertrophic cardiomyopathy (HCM) are most commonly seen in the genes encoding β-myosin heavy chain (MHC/MYH7) or myosin-binding protein C (MYBP-C).

HCM is mostly an autosomal dominant disorder. The most important histologic features of HCM myocardium are as follows:

• Massive myocyte hypertrophy particularly of transverse myocyte diameters

• Myofiber disarray including that of bundles of myocytes, individual myocytes, and contractile elements

• Interstitial and replacement fibrosis

Most of the patients can be treated with β blockers to decrease heart rate and contractility. Implantable cardioverter-defibrillators are used in patients with increased risk of ventricular arrhythmias. Some patients can also be treated by surgical excision of the septal muscle or by carefully controlled septal infarction through a catheter-based infusion of alcohol.

The below-given images show normal cardiac myocytes and myocytes in HCM respectively.







Solution to Question 15:

Transfusion of irradiated RBCs prevent the incidence of graft versus host disease (GVHD) associated with the transfusion.

GVHD occurs when immunologically competent cells or their precursors are transplanted into immunologically compromised recipients and the transferred cells recognize the alloantigens in the host and attack host tissues. It is seen most commonly during hematopoietic stem cell transplantation but may occur following transplantation of solid organs too.

GVHD maybe of two types: acute and chronic.

Acute GVHD occurs within days to weeks after allogeneic hematopoietic stem cell transplantation. It mainly involves the immune system and the epithelia of the skin (in the form of generalized rash), liver(in the form of jaundice), and intestines (in the form of bloody diarrhea). But the affected tissues are not heavily infiltrated by lymphocytes.

Chronic GVHD may follow acute GVHD or may occur insidiously. These patients have extensive skin damage, with destruction of skin appendages and fibrosis of the dermis, resembling systemic sclerosis. Cholestatic jaundice and esophageal strictures are also common. There will be involution of the thymus and depletion of lymphocytes in the lymph nodes. There will also be recurrent bouts of life-threatening infections. Some may even develop autoimmunity, owing to the grafted CD4+ helper T cells reacting with host B cells and stimulating them, which may result in autoantibodies production.

Depletion of donor T cells by irradiation, before transfusion can eliminate the occurrence of GVHD. But irradiation also has its side effects such as recurrence of tumor in leukemic patients, increased incidence of graft failures and EBV-related B-cell lymphoma.

Solution to Question 16:

Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency causes decrease in reduced glutathione and decreased NADPH.

G6PD is an important enzyme involved in the hexose monophosphate shunt. Defective enzyme function reduced the ability of red cells to protect themselves against oxidative injuries, thereby increasing lipid peroxidation and lead to hemolysis.

• G6PD reduces NADP to NADPH while oxidizing glucose-6-phosphate.

• This NADPH then provides reducing equivalents needed for conversion of oxidized glutathione to reduced glutathione.

• This reduced glutathione acts as a cofactor in reactions that neutralize compounds such as H_2O_2, thereby protecting against oxidant injury.

• Exposure of G6PD deficient red cells to oxidative stress leads to crosslinking of reactive sulfhydryl groups on globin chains, which then becomes denatured and form membrane-bound precipitates called 'Heinz bodies', thereby producing membrane damage.

• When these inclusion bearing RBCs pass through the splenic cords, macrophages pluck out these Heinz bodies.

• The resultant cells are called 'Bite cells'.

Clinical features:

• Episodic hemolysis, both intravascular and extravascular

• Hemolytic anaemia

• Hemoglobinemia

• Hemoglobinuria

The episode of hemolysis is self-limiting, since only the older RBCs are at risk for lysis, leaving the G6PD-replete younger RBCs in circulation. The recovery phase is heralded by reticulocytosis.

Solution to Question 17:

Leukemia is least likely to be seen in patients with Paroxysmal Nocturnal Hemoglobinuria (PNH).

Clinical manifestations of PNH include:

• Hemolytic anemia

• Pancytopenia

• Intermittent hemoglobinuria- acute hemolytic episodes occur at night owing to a slight fall in pH during sleep

• Haemosiderinuria

• Venous thrombosis (a leading cause of death in PNH)

• Patients diagnosed with PNH can later develop aplastic anemia and vice versa

PNH is an acquired disorder resulting from the mutation in the phosphatidylinositol glycan complementation group A gene (PIGA), an enzyme that is essential for the synthesis of

membrane-associated complement regulatory proteins. PNH blood cells are abnormally susceptible to lysis or injury by complement since they are deficient in three GPI- linked proteins that regulate complement activity:

• Decay-accelerating factor or CD55

• Membrane inhibitor of reactive lysis or CD59 (Inhibitor of C3 convertase) and

• C8-binding protein

Diagnosis:

• Ham's test: The abnormal sensitivity of red blood cells, leukocytes, and platelets to complement can be demonstrated in vitro using red cell lysis at acidic pH.

• Flow cytometry detects the red cells that are deficient in GPI-linked proteins such as CD59.

Treatment:

Eculizumab, a monoclonal antibody, that prevents the conversion of C5 to C5a has been proven effective in PNH. The definitive treatment is hematopoietic stem cell transplantation.

Solution to Question 18:

The image given in the question demonstrates the electron microscopic image of the blood-air barrier in the lungs. Acute respiratory distress syndrome (ARDS) has been set in this patient with COVID-19. Damage to the endothelial (Labelled as A) cells, with inflammation, is the key initiating event and the basis of lung damage. Neutrophils play a significant role in tissue injury in this condition.

Shock lung also known as ARDS/HMD/ALI (acute lung injury) is characterized histologically by diffuse alveolar damage (DAD). The characteristic histological picture is that of a hyaline membrane lining the alveolar walls. Edema, scattered neutrophils and macrophages, and necrotic epithelial cells are also present.

Histologic findings in COVID-19 patients :

• Diffuse alveolar damage

• Inflammation with mononuclear cells

The given HP slide shows diffuse alveolar damage which is the most common finding in the histopathological analysis of COVID-19 infected lungs with ARDS.



The image given in the question demonstrates the electron microscopic image of the blood-air barrier in the lungs. The structures that form the same are labeled in the image given below. The side of location of endothelium can be deduced by the location of the RBC.



P1 - Type I pneumocyte
E - Endothelial cell
BM - Fused basal lamina
Er - Erythrocyte

©MARROW

Solution to Question 19:

An endo-myocardial biopsy is the gold standard investigation for the diagnosis of myocarditis.

Myocarditis can be due to

• Infections: Coxsackie A and B, HIV, C.diphtheriae, Ricketssiae, Candida, Protozoa and helminths.

• Immune-mediated reactions: Post-viral, post-streptococcal, drug hypersensitivity (e.g: methyl dopa, sulfinamides), transplant rejection, etc.,

• Unknown: Sarcoidosis, Giant cell myocarditis

Active myocarditis is characterized by an interstitial inflammatory infiltrate associated with degenerating or apoptotic myocytes.

Solution to Question 20:

The next investigation of choice, in a child suspected to have a chloroma, is a bone marrow biopsy.

Chloroma, also called myeloid sarcoma, granulocytic sarcoma, or myeloblastoma is an extramedullary tumor that occurs in 2% to 14% of patients with acute myeloid leukemia (AML). Frequent sites of involvement are the skin, lymph nodes, gastrointestinal tract, bone, soft tissue, testis, and peritoneum.

Isolated orbital involvement of myeloid sarcoma is rare. Proptosis is the most common presentation. Other symptoms include lacrimal swelling, conjunctival mass lesion, retinal hemorrhages, ptosis, dystopia. Decreased visual acuity and ocular movement restriction have also been reported.

Most cases of chloroma can be diagnosed on peripheral smear examination, which is easily available and is relatively inexpensive. Sometimes it may not be associated with any peripheral smear abnormality, and hence bone marrow aspirate and biopsy should be done to confirm AML. The diagnosis can also be made if Auer rods are detected. Myeloid sarcomas have immunophenotypic features that are identical to AML.

Although myeloid sarcomas are radiosensitive, they are managed with systemic chemotherapy in most cases, even in the absence of bone marrow involvement. Surgery and/or radiation therapy may be indicated for symptomatic lesions or tumors causing local organ dysfunction or obstruction.

The image shows Auer rods contained within myeloblasts.



Solution to Question 21:

From the clinical history of progressive breathlessness and the histopathological image showing asbestos bodies, ferruginous bodies, and fibrosis, the patient is most likely working in the asbestos industry.

Asbestosis occurs due to the inhalation of crystalline hydrated silicates (asbestos). It is characterized by diffuse parenchymal interstitial fibrosis. It usually begins in the lower lobes of the lungs and may present with dyspnoea and cough, usually after 20-30 years of exposure. Other asbestos-related diseases include pleural plaques, recurrent pleural effusions, lung carcinoma, and mesothelioma.

Asbestos bodies are a hallmark finding. They appear as golden brown, fusiform, or beaded rods. These are formed when the asbestos fibers are phagocytosed by the macrophages and are coated by iron-containing proteinaceous material derived from phagocyte ferritin. Similar iron-protein complexes coat other inorganic particulates other than asbestos which are termed as ferruginous bodies.

The image below shows asbestos bodies.



Other options:

Option A: Silicosis is due to the inhalation of crystalline silicon dioxide (silica) as seen in individuals employed in sandblasting or rock mining. Macrophages phagocytose inhaled silica particles, thereby activating and releasing inflammatory mediators (IL-1, IL-8) that cause fibroblast recruitment. Subsequently, collagen is deposited in the interstitium.

Microscopy reveals coalescent nodules of whorled collagen fibers surrounded by a peripheral zone of silica-laden macrophages.

Shown below is an image of a silicotic nodule with collagen whorls as seen in this condition.



Option C: Anthracosis/coal workers' pneumoconiosis is a pulmonary condition seen in coal miners. The inhaled carbon pigment is engulfed by alveolar or interstitial macrophages, which accumulate in the connective tissue. Initially it causes simple coal workers pneumoconiosis with little or no pulmonary dysfunction. With continued exposure, lung tissue undergoes progressive massive fibrosis which is called as complicated coal workers pneumoconiosis. The upper lobes of the lungs are commonly involved.

Microscopic features consist of coal-laden macrophages, black pigment, and dense collagenous interstitial fibrosis.

The image below shows anthracosis- black pigment engulfed within macrophages as well as in the form of deposits.



Option D: Cotton dust causes byssinosis and can induce occupational asthma.

Solution to Question 22:

The lesion shown in the histopathological slide is Aschoff's body and the most probable diagnosis is rheumatic heart disease (RHD).

Aschoff's bodies are the lesions found in the heart during acute rheumatic fever. They are composed of foci of T lymphocytes, activated macrophages called Anitschkow cells, and occasional plasma cells. Anitschkow cells have abundant cytoplasm and central round-to-ovoid nuclei with chromatin that is centrally condensed into a slender, wavy ribbon. Hence, they are also called caterpillar cells.

Rheumatic fever is the leading cause of mitral stenosis. During acute rheumatic fever, Aschoff bodies may be found in any of the three layers of the heart that results in pericarditis, myocarditis, or endocarditis.

Chronic inflammation leads to diffuse thickening and fibrosis of the valve leaflets, often with calcific deposits. Fibrinoid necrosis occurs due to inflammation of the endocardium and the left-sided valves. Over these areas of necrosis, small vegetations called verrucae develop along the lines of closure. Irregular thickenings present in the subendocardium are called MacCallum plaques.

In chronic RHD, the mitral valve leaflet is thickened, with commissural fusion and shortening. There is thickening and fusion of the tendinous cords. This can lead to narrowing the mitral orifice, leading to a fish mouth appearance (button hole).

Some major manifestations of rheumatic Fever are:

• Migratory polyarthritis of the large joints

• Pancarditis (myocarditis, pericarditis, or endocarditis)

• Subcutaneous nodules (typically on extensor surfaces of extremities)

• Erythema marginatum, an irregular circinate skin rash

• Sydenham chorea, a neurologic disorder with involuntary rapid movements

Image showing button hole appearance seen in mitral stenosis in case of rheumatic heart disease.

Image show Aschoff bodies seen in RHD.



Other options:

Option B: Degenerative calcific deposits in the mitral valve typically develop in the fibrous annulus. They are seen grossly as irregular, stony hard, occasionally ulcerated nodules at the base of the leaflets. Usually, it does not affect valvular function. However, occasionally it can lead to regurgitation, stenosis, or arrhythmias.

The image below shows severe calcification of the mitral valve annulus with focally thickened shortened and fused chordae.

Option C: In sarcoidosis, the involved tissues contain well-formed non-necrotizing granulomas. They are composed of aggregates of tightly clustered epithelioid macrophages, often with giant cells, usually without central necrosis. As the lesions become chronic, fibrous rims or hyaline fibrous scars enclose the granulomas.

The image below shows non-necrotizing granulomas in the wall of a bronchiole in sarcoidosis.



Option D: In tuberculosis, sites of active involvement of tuberculosis are marked histologically by a characteristic granulomatous inflammatory reaction composed of both caseating and non-caseating tubercles. Usually, granuloma shows central granular caseation surrounded by multinucleate giant cells. Central caseation may be absent in non-caseating granulomas. Staining should be done to detect acid-fast bacilli when granulomas are present.

The image below shows a caseating granulomatous lesion with areas of caseation (on the right half) bordered by collections of epithelioid cells, Langhans giant cells, and lymphocytes.



Image showing non-necrotizing granuloma seen in pulmonary tuberculosis.



Solution to Question 23:

The presence of Birbeck granules in the cytoplasm is a characteristic finding of Langerhans cell histiocytosis (LCH) on electron microscopy.

Birbeck granules have a dilated terminal end producing a tennis racket–like appearance.
It contains the protein langerin (CD207). It is seen on electron microscopy.

The image below shows Birbeck granules on electron microscopy seen in LCH.



Histopathologically, Langerhans cells have a single large nucleus with a characteristic folded or grooved nucleus (coffee-bean appearance) with small inconspicuous nucleoli. Birbeck granules are seen in the cytoplasm. The tumor cells specifically express HLA-DR, S-100, and CD1a and langerin (CD207) markers.

The image below shows histopathological image of Langerhan cells with folded nuclei.



Langerhans cell histiocytosis occurs due to the proliferation of a special type of immature dendritic cells, called Langerhans cells. Their normal function is to present antigens. The most common mutation is BRAFV600E gain-of-function mutation.

It is characterized by single or multiple osteolytic bone lesions and can also infiltrate nearly every organ, like the skin (of the scalp), lungs, liver, spleen, bone marrow, or the central nervous system.

LCH can present as the following clinical entities:

• Multifocal multisystem Langerhans cell histiocytosis (Letterer-Siwe disease)

• Unifocal and multifocal unisystem Langerhans cell histiocytosis (eosinophilic granuloma)- The combination of bone defects, diabetes insipidus, and exophthalmos is called the Hand-Schüller-Christian triad.

• Pulmonary Langerhans cell histiocytosis

The below image shows lytic lesion of skull seen in LCH.



Solution to Question 24:

The most likely cause in the following case is Non-Hodgkin lymphoma(NHL). Diffuse large B cell lymphoma (DLBCL), a type of NHL is the most common lymphoma. NHL affects the elderly age group and extranodal presentation is common. DLBCL can involve CNS, and CNS involvement is particularly common if there is involvement of the testes, paranasal sinuses, or bone marrow as well.

Primary testicular lymphoma is common in NHL when compared to Hodgkin lymphoma. Primary testicular lymphoma(PTL) is an aggressive and uncommon form of non-Hodgkin lymphoma(NHL) usually affecting the elderly age group(>60y). It typically presents as a painless, firm testicular mass. Constitutional symptoms are uncommon, and if present, warrant a systemic involvement. Metastasis involves the central nervous system (CNS), skin, contralateral testis, and pleura resulting in local complications.

Even though ataxia is seen as a paraneoplastic symptom due to cerebellar degeneration in Hodgkin lymphoma, testicular involvement is very rare and so far only 5 cases have been reported. Hodgkin lymphoma arises in a single node or chain of nodes and spreads first to anatomically contiguous lymphoid tissues. The extranodal presentation is rare. The extranodal disease can occur at any site like lung, liver, bone, and bone marrow is most common if it occurs. The central nervous system and the testis are exceptionally rare sites for Hodgkin's lymphoma. Thus Hodgkin's lymphoma can be ruled out.

Other options:

A) Glioblastoma multiforme is a highly malignant tumor, common in older adults in the fifth and sixth decades. The most common locations are the frontal and temporal lobes. The tumor may spread and invade the other hemisphere, resulting in a characteristic butterfly appearance. Extracranial metastases of malignant gliomas are extremely rare due to:

• Absence of a lymphatic system in the brain and spinal cord for systemic dissemination

• Dense dura around intracranial veins, preventing the penetration of the tumor

• Lack of nurturing stroma in other organs prevents the proliferation of glioblastoma cells

Therefore the tumor metastasizing to the testis is very much unlikely and glioblastoma can be ruled out.

Image showing MRI(magnetic resonance imaging) of glioblastoma



Option D) Testicular cancer is the most common cancer in men aged from 15 to 35 years. Germ cell tumors(GCT) account for 90%–95% of cases. Non-seminomatous form (NSGCT) metastasizes to the lungs, liver, central nervous system, and bone in order of frequency. The frequency of brain metastases is very low, not exceeding 1.2%.

Note: This is a controversial question. A more detailed clinical history and examination than what is provided in the stem are required for us to arrive at a definitive diagnosis.

Solution to Question 25:

The gross image of the liver shows a nutmeg appearance, seen in heart failure. The dark areas represent necrosis, white areas represent viable liver tissue.

Nutmeg liver refers to the mottled appearance of the liver which resembles the cut section of nutmeg. This occurs as a result of hepatic venous congestion following right heart failure. The dark centrilobular zones reflect sinusoidal congestion. Congestion is seen around the central veins usually from right-sided heart failure. The pale periportal zones reflect normal or fatty liver tissue.

The liver derives its blood supply from the portal venous system and the hepatic artery. The portal triad constitutes:

• Branches of the hepatic artery

• Portal vein

• Bile duct

The blood from the portal vein and the hepatic artery flow through the intervening sinusoids and drains into the central vein. The central veins from adjacent lobules form interlobular veins, which then unite as hepatic veins and finally drain into the inferior vena cava.

The hepatic lobule is divided into three zones:

• Zone 1 is the periportal area

• Zone 2 is the intermediate zone

• Zone 3 is the area closest to the central vein

Right-sided cardiac failure leads to passive congestion of the liver. Left-sided cardiac failure leads to hypoperfusion and hypoxia of hepatocytes around central veins. The combination of hypoperfusion and retrograde congestion causes centrilobular hemorrhagic necrosis, leading to this mottled appearance of the liver, known as nutmeg liver.

The microscopic image below of a nutmeg liver shows perisinusoidal fibrosis, hepatic venule dilation, and dilation of the sinusoids in zone III.



Solution to Question 26:

The probable diagnosis, cell, and virus from the given clinical scenario and the histopathological image are Hodgkin's lymphoma, Reed-Sternberg cells (RS cells), and Epstein-Barr virus (EBV) respectively.

Hodgkin's lymphoma arises in a single lymph node or chain of nodes and spreads first to anatomically contiguous lymphoid tissues. It usually presents as painless lymphadenopathy. The distinctive feature of Hodgkin's lymphoma is the presence of Reed-Sternberg cells.

RS cells are giant neoplastic cells derived from germinal centre or post–germinal centre B cells. They appear as large cells with multiple nuclei or a single nucleus with multiple nuclear lobes. Each cell contains large inclusion-like prominent nucleoli, and abundant, usually slightly eosinophilic, cytoplasm, depicting an owl-eye appearance, as shown in the image below.



WHO classifies Hodgkin's lymphoma into 5 subtypes:

• Nodular sclerosis: Frequent lacunar cells

• Mixed cellularity

• Lymphocyte-rich

• Lymphocyte depletion

• Nodular lymphocyte predominance

The first 4 subtypes are considered classical Hodgkin's lymphoma. The RS cells are CD 15+ and CD30+. The lymphocyte predominant type is considered non-classical Hodgkin's lymphoma with CD15-, CD30-, CD20+ and EB -.

The image below shows a popcorn cell seen in lymphocytic predominant Hodgkin's lymphoma.

Type	RS-cell	EB Virus	Prognosis
1. Nodular sclerosis (most common in the world)	Frequent lacunar cells and occasional diagnostic RS cells	Rare	Very good
2. Mixed cellularity (most common in India)	Frequent mononuclear and diagnostic RS cells	70%	Good
3.Lymphocyte Rich	Frequent mononuclear and diagnostic RS cells	40%	Very good
4.Lymphocyte depleted	Reticular variant: frequent diagnostic RS cells and variants and a paucity of background reactive cell	90%(Max)	Poorest
5.Lymphocyte predominant	L&H variant (Popcorn cell) variants in the background of follicular dendritic cells	0%(Min)	Excellent



Other options:

Option A: RS cells are diagnostic of Hodgkin's lymphoma, not non-Hodgkin's lymphoma. The histopathologic findings in non-Hodgkin lymphoma (NHL) are varied and depend on the subtype.

Option B: Kaposi sarcoma is a vascular neoplasm caused by human herpesvirus 8 (HHV8), also known as Kaposi sarcoma herpesvirus. Cutaneous lesions of Kaposi sarcoma are composed of spindle cells, mostly in the dermis or subcutaneous tissues, with small vessels and slit-like spaces with red cells.

The image shows lesions seen in Kaposi's sarcoma.



Option D: Infectious mononucleosis is caused by EBV. The peripheral blood smear shows absolute lymphocytosis with mononucleosis cells which are atypical activated T lymphocytes. They have abundant cytoplasm with multiple vacuolations and azurophilic granules, and an oval, indented, or folded nucleus and are shown below.



About 5-7% of the cases of infectious mononucleosis can be caused by the cytomegalovirus (CMV), which can also cause pneumonia. CMV-infected pneumocytes with large, intranuclear, basophilic inclusions with haloes at their periphery as well as small, granular cytoplasmic inclusions are seen below with an owl-eye appearance.



Solution to Question 27:

The given clinical features suggest type 1 hypersensitivity reaction and are mediated by immunoglobulin E (IgE).

Immediate (Type I) hypersensitivity reactions refer to allergic reactions or allergies.

It is a rapid immunologic reaction that is triggered by the binding of an antigen to IgE antibody on the surface of mast cells. It occurs in a previously sensitized individual. These reactions are

induced by environmental antigens (allergens) that stimulate strong TH2 responses and IgE production in genetically susceptible individuals.

Activation of TH2 cells is mediated by IL-4 and in turn, TH2 cells produce IL-4, IL-5, and IL-13. These cytokines act on B cells to stimulate IgE and activation of eosinophils. IgE coats mast cells by binding to the FcεRI receptor. Re-exposure to the allergen leads to cross-linking of the IgE and FcεRI which in turn causes activation of mast cells releasing powerful mediators. The principal mediators are:

• Histamine

• Proteases

• Prostaglandins

• Leukotrienes

• Cytokines like TNF, IL-1, and chemokines.

The clinical manifestations may be local or systemic and range from mildly annoying rhinitis to fatal anaphylaxis.

It occurs in two phases:

• The immediate phase is characterized by vasodilation, vascular leakage, smooth muscle spasm (in asthma), or glandular secretions. These changes occur within minutes after exposure to an allergen and subside in a few hours.

• Late-phase reaction is characterized by infiltration of tissues with eosinophils, neutrophils, basophils, monocytes, and CD4+ T cells. Late-phase reaction sets in 2 to 24 hours later and may last for several days.

Image: Sequence of events in immediate (type 1) hypersensitivity.



Solution to Question 28:

All the above are associated with massive splenomegaly except aplastic anemia.

Aplastic anemia is a syndrome of chronic primary hematopoietic failure. This results in pancytopenia (anemia, neutropenia, and thrombocytopenia). Important causes include

• Alkylating agents, antimetabolites

• Benzene

• Whole-body irradiation

• Viral infections: Hepatitis (unknown virus), Cytomegalovirus, Epstein-Barr virus, Herpes zoster

• Fanconi anemia

The patient will have

• Decreased red blood cells leading to anemia

• Decreased white blood cells leading to recurrent infections

• Decreased platelets leading to petechiae and ecchymoses.

Many other leukemias and lymphomas may have a similar presentation. However, splenomegaly is characteristically absent in these patients.

The bone marrow is markedly hypocellular. It is largely devoid of hematopoietic cells. Marrow aspirates yield a dry tap having only fat cells, fibrous stroma, lymphocytes, and plasma cells. Aplasia is best appreciated in marrow biopsies. The following image shows a hypocellular bone marrow with little hematopoiesis and several fat cells.



Stem cell transplantation is the treatment of choice in those with a suitable donor. Immunosuppressive therapy is responsive in old patients or in patients without suitable donors.

Other options:

Option A- Chronic myeloid leukemia (CML) is a myeloproliferative neoplasm with the presence of BCR-ABL fusion gene translocation mutation. The marrow is markedly hypercellular because of increased numbers of maturing granulocytic precursor cells. This results in decreased hematopoiesis within the bone marrow. The spleen is often greatly enlarged as a result of

extensive extramedullary hematopoiesis. Extramedullary hematopoiesis can also produce mild hepatomegaly and lymphadenopathy.

Option C- Hairy cell leukemia is a memory B-cell neoplasm predominantly seen in middle-aged male patients. It occurs due to activating BRAF mutations. Infiltration of the bone marrow, liver, and spleen with tumor cells result in splenomegaly. Sometimes, this can be the only finding in the patient. Pancytopenia results from marrow involvement.

Hairy cells have round, oblong, or reniform nuclei and moderate amounts of pale blue cytoplasm with thread-like or bleb-like extensions. Immunophenotyping shows hairy cell leukemias typically express the pan-B-cell markers CD19 and CD20 and certain relatively distinctive markers, such as CD11c, CD25, CD103, and annexin A1.

Option D- Lymphomas cause splenomegaly due to the infiltration of neoplastic cells in the spleen.

Solution to Question 29:

The clinical scenario with complaints of cough, low-grade fever, hemoptysis is suggestive of pulmonary tuberculosis. The diagnosis is further supported by a cavitary lesion in the apex of the lung and a biopsy showing caseous necrosis. The pathophysiology is a type 4 hypersensitivity reaction mediated by activated macrophages.

Tuberculosis is a chronic pulmonary and systemic disease caused by Mycobacterium tuberculosis. Granulomatous inflammation and tissue damage are by Th-1 mediated macrophage activation and killing of bacteria. Interferon-gamma (IFN-γ) released by Th-1 cells is the critical mediator that activates the macrophages. This activation differentiates the macrophages into epithelioid cells that aggregate to form granuloma. This is a T-cell mediated (Type 4) hypersensitivity reaction.

Pulmonary TB due to exogenous (new infection) or endogenous (reactivation of a dormant primary lesion) infection in a sensitized individual is called post-primary or reactivation or secondary or adult-type TB.

The apical and posterior segments of the upper lobe and superior segments of the lower lobe are most commonly involved in post-primary TB due to increased ventilation in these segments than the lower zones. Patients often present with fever, productive cough, blood-streaked sputum, night sweats, malaise, loss of appetite, and weight loss.

Radiological findings include ill-defined opacification in the upper lobes, consolidation, collapse, and cavitation development.

• Typically, the disease is parenchymal without nodal enlargement.

• Cavitation is a distinguishing feature of postprimary TB.

• Typical cavities are thick-walled and irregular.

• Air-fluid levels are uncommon in TB. The presence of air-fluid levels indicates superadded infections.

• Upper-lobe involvement with cavitation and the absence of lymphadenopathy distinguishes postprimary TB from primary TB.

The image below shows chest X-ray findings of a post-primary TB with a cavity (black arrows) in the right upper lobe and bilateral pulmonary infiltrates (white triangles) in the apical regions.



Image shows a gross picture of caseous necrosis. Notice the characteristic "cheesy appearance" due to its white appearance of the area of necrosis.



Image below shows tuberculoid granuloma associated with caseous necrosis (upper right) and surrounding epithelioid cells, Langhans' multinucleated giant cells.



Other options:

Option A: Enzyme degradation is seen in liquefactive necrosis which is characterized by digestion of the dead cells, resulting in transformation of the tissue into a viscous liquid. It is seen in focal bacterial or fungal infections. Eg: Brain abscess.

Option C: Fibrinoid deposition is seen in fibrinoid necrosis. It is seen in immune reactions involving blood vessels with deposition of fibrinoid material which is combination of immune complexes and plasma proteins. Eg: Vasculitis syndromes

Option D: Sudden cut-off of blood supply causes coagulative necrosis. It is a form of necrosis in which the original architecture of dead tissue is preserved with firm texture. Eg: Ischemic injury in all organs, except the brain.

Solution to Question 30:

In iron deficiency anemia, ferritin translation is decreased by binding iron regulatory protein to iron response elements.

Dietary iron is carried in the blood by transferrin and enters cells via endocytosis, mediated by the transferrin receptor. Excess iron is stored by binding to ferritin. The levels of transferrin, transferrin receptor (TfR), and ferritin are therefore crucial to cellular iron homeostasis.

The synthesis of these three proteins is regulated in response to iron availability and aconitase (enzyme in citric acid cycle).

When a cell is depleted of iron, the aconitase enzyme is converted to apoaconitase. Apoaconitase is identical to iron regulatory protein 1 (IRP-1) and is closely related to IRP-2. It regulates iron metabolism.

The iron regulatory proteins (IRPs) are two cytoplasmic RNA-binding proteins that control iron metabolism. They bind to regions in the mRNA containing the ferritin and transferrin receptor. These mRNA sequences are part of hairpin structures known as iron response elements (IREs).

• When the IRPs bind to the 5' untranslated IRE sequence in the ferritin mRNA, IRPs block ferritin translation, thus decreasing ferritin.

• When IRPs are bound to the 3' end of the untranslated IRE sequence, they stabilize the mRNA, prevent its degradation and increase the copies of transferrin receptor per mRNA molecule.

Hence, in iron-deficient cells, iron storage (bound to ferritin) is reduced and iron uptake (via transferrin receptor) becomes more efficient.

Solution to Question 31:

An elderly patient with chronic fatigue and lymphadenopathy, and a peripheral smear showing the presence of smudge cells suggest a diagnosis of chronic lymphocytic leukemia. Immunophenotyping using flow cytometry is the investigation done to arrive at the diagnosis. The classical immunophenotypic feature of CLL is CD 19+, CD20+, CD 23+, CD5+.

Chronic Lymphocytic Leukemia (CLL) is a peripheral B-cell neoplasm usually affecting adults, more commonly men. Patients are often asymptomatic at diagnosis. When symptomatic, it presents non specifically with easy fatigability, weight loss, and anorexia. Generalized lymphadenopathy and hepatosplenomegaly are present in most symptomatic patients.

The chromosomal defects in CLL include:

• Deletion of 13q (commonest and has a good prognosis)

• Deletion of 11q

• Trisomy 12q

• Deletion of 17p (poor prognosis)

The peripheral blood smear in cases of chronic lymphocytic leukemia (CLL) contains small round lymphocytes having condensed chromatin and scant cytoplasm. Some of these cells may get disrupted while preparing the smears, producing smudge cells. An absolute lymphocyte count of >5000/mm3 is required for diagnosis. Hypogammaglobulinemia is commonly seen, making the patient susceptible to infections. A few patients may develop hemolytic anemia or thrombocytopenia due to autoantibodies made by non-neoplastic B cells.

On lymph node biopsy: Lymph nodes are composed of small lymphocytes admixed with larger activated lymphocytes. The large lymphocytes contain mitotically active cells and gather in loose aggregates referred to as proliferation centers. These proliferation centers are pathognomic to CLL when present.

Expectant management is the treatment of choice for asymptomatic patients with chronic lymphocytic leukemia. The gold standard for therapy is a combination of fludarabine, cyclophosphamide, and rituximab.

Richter's syndrome is a complication of CLL in which there is a transformation to an aggressive lymphoma, most commonly diffuse large B-cell lymphoma (DLBCL).

The image shows small round lymphocytes and smudge cells seen in CLL.



Solution to Question 32:

Oil red O (ORO) stain is used in frozen section.

Oil red O stains lipid/ fats and gives a characteristic orange/red appearance. It is used in the unfixed frozen sections as the fixation of fats with other routine processes may lose the lipid content. 10% Neutral buffer formaldehyde (NBF) is the preferred fixative in histopathological sections staining with Oil red O. Zenker's fixative and Helly's fixative are best avoided during staining with ORO.

Oil red O stains the lipid in the vacuoles in the pulmonary vasculature. It can be useful in identifying fat emboli in lung tissue, diagnosing renal cell carcinoma, and sebaceous gland tumors of the skin.

Other stains used for lipids or fats:

• Nile blue sulfate. The preferred fixative is formal calcium.

• Osmic acid is used in the frozen section. 10% NBF is the preferred fixative

• Sudan black B is used in the frozen section. The preferred fixative is 10% NBF.

Other Options:

Option A: Formalin fixation- Formalin as 10% neutral buffered formalin (NBF) is the most common fixative used in diagnostic pathology.

Option C: Ethyl alcohol- Ideal fixative for cytological smears because they act quickly and give good nuclear detail.

Option D: Glutaraldehyde- Preferred fixative in electron microscopy. The standard solution is a 2.5% buffered glutaraldehyde.

Image below shows microscopy of a oil red O stained bronchoalveolar lavage sample from a patient with acute lung injury.



The image shows the lipid vesicles in the adipocytes stained using oil-red-O staining.



Solution to Question 33:

The given clinical scenario and the classical histopathological finding of starry sky appearance suggest the diagnosis of Burkitt's lymphoma and the most likely genetic abnormality is the translocation of the MYC gene on chromosome 8.

Burkitt's lymphoma arises from mature germinal center B-cells that express surface IgM, CD19, CD20, CD10, and BCL6.

It is divided into endemic and sporadic types, both of which occur mainly in children or young adults.

• African (endemic) Burkitt lymphoma often presents as a mass involving the mandible and may involve the abdominal viscera, particularly the kidneys, ovaries, and adrenal glands.

• Sporadic (non-endemic) Burkitt lymphoma most often appears as a mass involving the ileocecum and peritoneum.

• A subset of aggressive lymphomas occurs in individuals infected with HIV.

All forms of Burkitt lymphoma are associated with translocations of the MYC gene on chromosome 8 that lead to increased MYC protein levels. Essentially all endemic Burkitt lymphomas are latently infected with Epstein-Barr virus (EBV).

The tumor shows a high mitotic index with numerous apoptotic cells. The nuclear remnants of these apoptotic cells are phagocytosed by interspersed benign macrophages. The phagocytes have abundant clear cytoplasm, creating a characteristic starry sky pattern.

Burkitt lymphoma is very aggressive but responds well to intensive chemotherapy.

The image below shows the starry sky appearance seen in Burkitt's lymphoma.



Solution to Question 34:

The most common variant of Guillain-Barré syndrome is acute inflammatory demyelinating polyneuropathy (AIDP). The area affected in this condition is the myelin sheath (marked as 2 in the image).



Guillain-Barré syndrome (GBS) is an acute autoimmune disease that causes fulminant polyradiculoneuropathy. It is a T-cell mediated immune response.

GBS is usually associated with a history of upper respiratory infection (URI) or acute gastroenteritis (AGE) 1-3 weeks prior. The causative organisms may be Campylobacter jejuni, cytomegalovirus, Epstein-Barr virus, and Mycoplasma pneumoniae.

It presents with the following features:

• Ascending motor paralysis (lower limbs > upper limbs)

• Sensory deficits may or may not be seen

• Deep tendon reflexes - absent

• Transient bladder dysfunction - severe cases

• Autonomic manifestations - loss of vasomotor control, postural hypotension, cardiac dysrhythmias

Histological examination shows inflammation of peripheral nerves, manifested as perivenular and endoneurial infiltration by lymphocytes, macrophages, and a few plasma cells. On electron microscopy, an early effect on myelin sheaths is seen. There is segmental demyelination induced by the actions of activated macrophages.

There are several types of Guillain–Barre syndrome:

• Demyelinating

• Acute inflammatory demyelinating polyneuropathy (AIDP) - most common

• Axonal

• Acute motor axonal neuropathy (AMAN)

• Acute motor-sensory axonal neuropathy (AMSAN)

• Regional GBS syndromes: Miller–Fisher syndrome (MFS)

Pathology NEET 2021

Question 1:

A 70-year-old patient presented with cough, fatiguability, and weight loss. He was diagnosed with squamous cell carcinoma (SCC) on bronchoscopy. The resected specimen had a hilar lymph node that was 1cm in size and showed a black pigment. What is the black pigment likely to be?

a) Anthracotic pigment

b) Melanin

c) Lipochrome

d) Hemosiderin

Question 2:

A 34-year-old female patient presented to the outpatient department with migratory arthritis and examination revealed a pan-systolic murmur. An echocardiography was performed which was suggestive of mitral regurgitation. A biopsy specimen was obtained and the image is provided below. What is the most likely lesion shown in the histopathology slide?



a) Aschoff's bodies

b) Epithelioid granuloma

c) Granulomatous vasculitis

d) Granuloma inguinale

Question 3:

A 30-year-old male complained of fatigue for the past 1 year. Examination revealed massive splenomegaly. His blood workup shows the following findings. Which of the following is the most sensitive investigation in this case?

a) Immunophenotyping

b) LAP score

c) FISH and PCR

d) Flow cytometry

Question 4:

A girl presents with short stature, webbed neck, thick calves, shield-shaped chest, swelling of the ankle, and increased carrying angle. What is the likely diagnosis?

a) Turner syndrome

b) Down syndrome

c) Patau syndrome

d) Edward syndrome

Question 5:

A patient with occupational exposure to benzene presents with repeated infections, easy fatigability, weight loss, and evening rise of temperature. This occupational exposure will predispose him to which of the following cancers?

a) Leukemia

b) Lung cancer

c) Bladder cancer

d) Skin cancer

Question 6:

In iron deficiency anemia, all of the following is increased except

a) Transferrin saturation

b) RBC protoporphyrin

c) Total iron binding capacity

d) Ferritin soluble receptors

Question 7:

Which of the following is a poor prognostic marker of multiple myeloma?

a) β-2 microglobulin

b) Calcium levels

c) Serum creatinine

d) Protein levels

Question 8:

A pap smear taken from a patient revealed hyperchromatic nuclei, irregular nuclear borders, and low maturation index involving the whole thickness but did not breach the basement membrane. What is the diagnosis?

a) Dysplasia

b) Carcinoma

c) Metaplasia

d) Hyperplasia

Question 9:

The mother of a 12-year-old girl was worried that her daughter is shorter when compared with her classmates. On examination, the child had a webbed neck, low hairline, and shield chest. A provisional diagnosis of Turner's syndrome was made. Which of the following is a likely finding to be seen in this patient?

a) Aortic dissection

b) Subluxation of lens

c) Mitral valve prolapse

d) Bilateral small streak ovaries

Question 10:

An 18-year-old male presented to the outpatient department with gum bleeding and fever for the past 2 months. General examination showed pallor and the systemic examination was unremarkable. Laboratory examination revealed Hb level- 3 gm/dl, TLC- 1500/microL, platelet count- 15000/microL. Peripheral smear shows macrocytes and the reticulocyte count is 0.5. Bone marrow examination revealed fatty streaks and absent megakaryocytes but no immature cells. What is the likely diagnosis?

a) Idiopathic acquired aplastic anemia

b) Paroxysmal nocturnal hemoglobinuria

c) Myelodysplastic syndrome

d) Tuberculosis

Question 11:

A child is presenting with palpable blanching purpura, abdominal pain, and pain in the knee joint resulting from arthritis. Urine examination showed hematuria and proteinuria. What is the most likely diagnosis?

a) Immune thrombocytopenic purpura

b) Dengue

c) Churg- Strauss syndrome

d) Henoch-Schonlein purpura

Question 12:

A 54-year-old patient presents to the outpatient department with fatigue, tiredness and cervical lymphadenopathy. The TLC and DLC were normal. No immature cells. A lymph node biopsy was performed which shows diffuse effacement of the lymph node architecture, cells with an indented nucleus, prominent nucleolus, and are atypical. CD10 and BCL2 are positive. What is the likely diagnosis?

a) Follicular lymphoma

b) Burkitt lymphoma

c) Mycosis fungoides

d) Chronic myeloid leukemia

Question 13:

A software engineer presents to the outpatient department with easy fatiguability. He gives a history of sitting in front of the computer for 12 - 14 hours. He consumes a lot of junk food with very less fruits and vegetables. Hb is 9gm/dl and MCV 120 fl. What is the most likely cause of this patient's anemia?

a) Cyanocobalamin deficiency

b) Folic acid deficiency

c) Iron deficiency

d) Hereditary spherocytosis

Question 14:

A chronic alcoholic patient presented with increasing abdominal girth. On examination, shifting dullness was present and the liver span was less than 7cm. Histopathological examination revealed the presence of intracytoplasmic eosinophilic inclusions in hepatocytes. What is the content of these inclusions?

a) Intermediate filaments

b) Actin

c) Microtubules

d) Fibronectin

Question 15:

A 65-year-old female patient is diagnosed with pure red cell aplasia and a mediastinal mass. Which of the following can be the likely cause?

a) Thymoma

b) Non Hodgkin lymphoma

c) Bronchogenic Carcinoma

d) Germ cell tumor

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	a
5	a
6	a
7	a
8	a
9	d
10	a
11	d
12	a
13	b
14	a
15	a

Detailed Explanations

Solution to Question 1:

The black-colored pigment in the hilar lymph node of the resected bronchoscopic specimen is likely to be an anthracotic pigment.

The black color is due to the deposition of carbon particles. The inhaled carbon particles are engulfed by the lung macrophages and then transported to the hilar lymph nodes.
This accumulates and causes the blackening of the lung tissues or anthracosis.

It is commonly seen in people who work in environments where there is high air pollution with carbon particles, such as coal mines. It is also the most common exogenous pigment found in the human body as a result of air pollution in urban areas. The aggregates of the carbon dust may induce fibroblastic reaction or emphysema that may lead to a condition called coal worker's pneumoconiosis.

The following image shows anthracosis with deposition of black carbon particles, within the cytoplasm of macrophages.





Option B: Melanin is brown-black in color which occurs as a result of oxidation of tyrosine in melanocytes, as shown in the following histopathology of a lymph node.



Option C: Lipochrome or lipofuscin (brown lipid) is seen as a yellow-brown pigment in histopathological sections of the liver and heart of normal aging patients, patients with severe malnutrition, and cancer cachexia. It is derived through lipid peroxidation and indicates free radical injury and lipid peroxidation. The following image shows a cardiac myocyte with fine yellow-brown lipofuscin granules in the perinuclear region.





Option D: Hemosiderin is a hemoglobin-derived golden-yellow colored pigment seen in conditions associated with iron overload such as hemochromatosis. The following image shows extensive intra-alveolar hemosiderin deposition following pulmonary hemorrhage.



Solution to Question 2:

The given clinical findings of migratory arthritis, pan-systolic murmur, and mitral regurgitation are suggestive of acute rheumatic fever. The most likely lesion shown in the histopathological slide is Aschoff's bodies.

Aschoff's bodies are the lesions found in the heart during acute rheumatic fever. They are composed of foci of T lymphocytes, occasional plasma cells, and plump-activated macrophages called Anitschkow cells. Anitschkow cells have abundant cytoplasm and central round-to-ovoid nuclei with chromatin that is centrally condensed into a slender, wavy ribbon. Hence they are also

called caterpillar cells.

During acute rheumatic fever, Aschoff bodies may be found in any of the three layers of the heart that results in pericarditis, myocarditis, or endocarditis.

Fibrinoid necrosis can occur due to inflammation of the endocardium and the left-sided valves. Over these areas of necrosis, small vegetations can be seen along the lines of closure, called verrucae. Irregular thickenings called MacCallum plaques can be present In the subendocardium.

In chronic RHD, the mitral valve shows leaflet thickening, commissural fusion and shortening, and thickening and fusion of the tendinous cords. Calcification and fibrous bridging of the valve can lead to narrowing the mitral orifice and giving it a fish mouth appearance.



Solution to Question 3:

The clinical presentation with fatigue, splenomegaly, and the peripheral blood smear findings are suggestive of chronic myelogenous leukemia(CML). The BCR-ABL gene which is specific for CML is detected by fluorescence in situ hybridization(FISH) or polymerase chain reaction(PCR) based tests.

Chronic myelogenous leukemia (CML) is a myeloproliferative disorder with the peak incidence in the fifth to sixth decade of life. The chimeric BCR-ABL gene is created by a reciprocal (9;22)(q34;q11) translocation(Philadelphia chromosome). This BCR-ABL gene distinguishes CML from other myeloproliferative disorders and is detected by fluorescent in situ hybridization (FISH) or polymerase chain reaction (PCR).

Clinical features:

Mild to moderate anemia and hypermetabolism, fatigability, weakness, weight loss, and anorexia. Dragging sensation in the abdomen caused by splenomegaly, or the acute onset of left upper quadrant pain due to splenic infarction. The spleen is often greatly enlarged as a result of extensive extramedullary hematopoiesis. Extramedullary hematopoiesis can also produce mild hepatomegaly and lymphadenopathy.

Peripheral blood:

• Leukocytosis with leukocyte count, often exceeding 100,000 cells/mm3, which consists predominantly of neutrophils, band forms, metamyelocytes, myelocytes, eosinophils, and basophils.

• Blasts are usually less than 10% of the circulating cells.

• Platelet count is usually increased

Bone Marrow:

• Markedly hypercellular marrow because of massively increased numbers of maturing granulocytic precursors. There is usually an elevated proportion of eosinophils and basophils.

• There is also an increase in megakaryocytes and usually include small, dysplastic forms and erythroid progenitors are present in normal or mildly decreased numbers. Typically there is an increased reticulin deposition, but overt marrow fibrosis is rare early in the course.

• Sea-blue histiocytes - These are scattered macrophages with abundant wrinkled,green-blue cytoplasm.

Other options:

Immunophenotyping and flow cytometry: Flow cytometry refers to the measurement of cellular antigens as they flow in a fluid stream. Immunophenotyping is employed using labeled antibodies against specific epitopes of cellular antigens. The technique that is usually used in immunophenotyping is flow cytometry. It has a limited role in the diagnosis of CML. It is mainly used in identifying CML blast transformation.

LAP score: Leucocyte alkaline phosphatase(LAP) score is decreased in cases of CML. It is only a first-line test in the diagnosis of CML and the diagnosis needs to be confirmed by other tests including bone marrow biopsy, molecular diagnostics, and cytogenetic analysis.

Solution to Question 4:

The clinical presentation of short stature, webbed neck, thick calves, shield-shaped chest, swelling of the ankle, and increased carrying angle is suggestive of Turner syndrome.

Turner syndrome occurs due to complete or partial monosomy of X-chromosomes. It produces hypogonadism in phenotypic females. It is the most common sex chromosome abnormality in females.

Important features associated with Turner's syndrome:

• Short stature

• Webbing of neck

• Broad chest and widely spaced nipples

• Cubitus valgus

• Streak ovaries, infertility, amenorrhea

• Peripheral lymphedema at birth

• Low posterior hairline

• Poor breast development

• Pigmented nevi

• Cardiac manifestations – Coarctation of aorta, bicuspid aortic valve

• Horse shoe kidney

Image showing the features of Turner syndrome



Other options:

Option B: Important features associated with Down's syndrome (Trisomy 21) include

• Intellectual disability

• Congenital heart defects

• Epicanthic folds and flat facial profile

• Simian crease

• Umbilical hernias

• Predisposition to leukemia

Option C: Important features associated with Patau syndrome (Trisomy 13):

• Microphthalmia

• Microcephaly and intellectual disability

• Cleft lip and palate

• Rocker-bottom feet

• Cardiac and renal defects

Option D: Important features associated with Edward syndrome (Trisomy 18):

• Prominent occiput, micrognathia, low set ears

• Intellectual disability

• Renal and cardiac malformations

• Limited hip abduction, rocker-bottom feet

Solution to Question 5:

Occupational exposure to benzene predisposes to leukemia. The given clinical history of repeated infections, easy fatigability, weight loss, and evening rise of temperature with benzene exposure points towards a diagnosis of leukemia.

Benzene is oxidized to an epoxide through hepatic CYP2E1. The epoxide and other metabolites disrupt progenitor cell differentiation in the bone marrow and lead to marrow aplasia and acute myeloid leukemia.

Acute myeloid leukemia is a tumor of hematopoietic progenitors caused by acquired oncogenic mutations that impede differentiation. This leads to the accumulation of immature myeloid blasts in the marrow. The replacement of the marrow with blasts produces marrow failure and complications related to anemia, thrombocytopenia, and neutropenia.

Other options:

Option B: Exposure to arsenic and its compounds, asbestos, beryllium and its compounds, chromium compounds, nickel compounds, radon, and its decay products can cause lung cancer.

Option C: Exposure to aryl amines, particularly 2-naphthylamine and related compounds, cyclophosphamide, cigarette smoking, irradiation can cause bladder cancer.

Option D: Exposure to UV rays, arsenic, and its compounds can cause skin cancer.

Solution to Question 6:

Transferrin saturation is decreased in iron deficiency anemia.

Clinical presentation may include fatigue, pallor, reduced exercise capacity. Cheilosis (fissures at the corners of the mouth) and koilonychia (spooning of the fingernails) are signs of advanced tissue iron deficiency.

Laboratory findings are:

• Reduced hemoglobin and hematocrit values.

• hypochromia, microcytosis, and poikilocytosis.

• serum iron and ferritin levels are low.

• The total plasma iron-binding capacity that reflects the elevated transferrin levels is high.

• Low serum iron with increased iron-binding capacity results in a reduction of transferrin saturation to below 15%.

• RBC protoporphyrin levels are increased. Protoporphyrin is an intermediate in the heme synthesis pathway. Under conditions in which heme synthesis is impaired, protoporphyrin

accumulates within the red cell.

• Bone marrow reveals a mild to moderate increase in erythroid progenitors. The absence of stainable iron in macrophages is a diagnostically significant finding.

Peripheral smear of microcytic hypochromic anemia:





Solution to Question 7:

β-2 microglobulin is a poor prognostic marker of multiple myeloma.

Multiple myeloma is a plasma cell neoplasm commonly associated with lytic bone lesions, hypercalcemia, renal failure, and acquired immune abnormalities.

Poor prognostic factors for multiple myeloma:

• Serum β2-microglobulin. It is the single most powerful predictor

• Deletion of 17pITP53 sequences and the translocations t(14;16) and t(14;20)

• Polyclonal serum Immunoglobulins

• Elevated lactate dehydrogenase

• High C-reactive protein

• Increased plasma cell proliferative activity

• High degree of bone marrow replacement

• Presence of plasmablastic morphology

• Elevated serum soluble receptor for IL6

Multiple myeloma usually presents as destructive plasma cell tumors involving the axial skeleton. The bones most commonly affected in descending order of frequency are the vertebral column, ribs, skull, pelvis, femur, clavicle, and scapula.

Radiographically, bone lesions appear as punched-out defects, usually 1 to 4 cm in diameter, and consist of soft, gelatinous, red tumor masses.

Radiograph of the skull showing punched out lesions:

Morphology:

• Malignant plasma cells have a perinuclear clearing due to a prominent golgi apparatus and an eccentrically placed nucleus.

• Relatively normal-appearing plasma cells, plasmablasts with vesicular nuclear chromatin and a prominent single nucleolus, or bizarre, multinucleated cells may predominate.

• Other cytologic variants include flame cells, Mott cells.

• Flame cells have fiery red cytoplasm, Mott cells contain multiple grapelike cytoplasmic droplets, a variety of other inclusions, including fibrils, crystalline rods, and globules. The globular inclusions are referred to as Russell bodies (if cytoplasmic) or Dutcher bodies (if nuclear).

• Rouleaux formation is characteristic but not specific. High levels of M protein in the blood causes red cells to stick to one another in linear arrays in smears leading to rouleaux formation.

Image showing Mott cell:



Image showing Russell and Dutcher bodies:



Peripheral smear of multiple myeloma:



Bence Jones proteins are excreted in the kidney and causes myeloma kidney. Most myelomas are associated with more than 3 g/dL of serum Ig and/or more than 6 mg/dL of urine Bence Jones protein. The most common monoclonal Immunoglobulin (M protein) is IgG followed by IgA.

The clinicopathologic diagnosis of multiple myeloma can be done based on the identification of clonal plasma cells in the marrow and the presence of CRAB criteria (hypercalcemia, renal dysfunction, anemia, and bone lesions).

Solution to Question 8:

The hyperchromatic nuclei, irregular nuclear borders, and intact basement membrane are all suggestive of dysplasia.

Dysplasia is characterized by the loss in the uniformity of the individual cells and their architectural orientation. The features of dysplasia include:

• Pleomorphism

• Hyperchromatic nuclei

• High nuclear-to-cytoplasmic ratio

• Increased mitotic figures

When the dysplastic changes involve the entire thickness of the epithelium without breaching the basement membrane it is considered preinvasive and is termed carcinoma-in-situ. If the tumor invades the basement membrane it is said to be invasive.

Dysplasia can be considered as a precursor to malignant but it does not always progress to cancer. Dysplasia usually occurs in the metaplastic epithelium but all metaplastic epithelium is not dysplastic.

Other options:

Metaplasia: It refers to the replacement of one cell type with another type. It is almost always associated with tissue damage, repair, and regeneration. The most common epithelial metaplasia is columnar to squamous.

Carcinoma: In carcinoma, there is the invasion of the basement membrane.

Hyperplasia: It refers to the increase in the number of cells in a tissue or organ. It can be physiologic or pathological. Enlargement of the female breast during pregnancy is physiologic hyperplasia. Endometrial hyperplasia is an example of pathological hyperplasia.

Solution to Question 9:

The likely finding seen in a patient with Turner syndrome is bilateral small streak ovaries.

Turner syndrome results from complete or partial monosomy of the X chromosome. It is characterized by hypogonadism in phenotypic females. It is the most common sex chromosome abnormality in females. Three types of karyotypic abnormalities are seen

• Missing an entire X chromosome - this results in a 45X karyotype.

• Structural abnormalities of the X chromosomes - isochromosome of the long arm and loss of the short arm, 46X,i(X)(q10), deletion of portions of both long and short arms that lead to the formation of a ring chromosome, 46X,r(X), deletion of portions of the short or long arm, 46X,del(Xq) or 46X,del(Xp).

• Mosaics - they will have 45X along with one or more karyotypically normal or abnormal cell types. Examples are 45X/46XX, 45X/46XY, 45X/47XXX, 45X/46X,i(X)(q10).

Important features associated with Turner's syndrome:

• Short stature

• Webbing of neck

• Broad chest and widely spaced nipples

• Cubitus valgus

• Streak ovaries

• Infertility, amenorrhea

• Peripheral lymphedema at birth

• Low posterior hairline

• Broad chest and widely spaced nipples

• Pigmented nevi

• Coarctation of aorta

There is failure to develop normal secondary sex characteristics at puberty. The genitalia remain infantile, breast development is inadequate, and there is little pubic hair. Short stature and amenorrhea are of particular importance in establishing the diagnosis in an adult.

Other options:

Aortic dissection, subluxation of lens, and mitral valve prolapse are seen in Marfan syndrome.



Solution to Question 10:

The given clinical scenario showing a young male with gum bleeding, fever, pancytopenia, and a low reticulocyte count with bone marrow showing fatty streaks and no immature cells is suggestive of idiopathic acquired aplastic anemia.

Aplastic anemia is a syndrome characterized by chronic hematopoietic failure and pancytopenia. The common causes include:

• Acquired

• Idiopathic (most common cause, about 65% of the cases)

• Acquired stem-cell defects

• Immune-mediated

• Chemical agents

• Dose-related (alkylating agents, antimetabolites, benzene, chloramphenicol, inorganic arsenals)

• Idiosyncratic (chloramphenicol, phenylbutazone, organic arsenicals, carbamazepine, penicillamine, gold salts)

• Physical agents

• Whole-body irradiation

• Viral infections (hepatitis, Cytomegalovirus, Epstein–Barr virus, Varicella-Zoster virus)

• Inherited

• Fanconi anemia

• Telomerase defects

Pathogenesis involves an extrinsic immune-mediated suppression of bone marrow and an intrinsic abnormality of the stem cells.

Clinical features and diagnosis:

There is no age or sex predisposition for aplastic anemia. It presents insidiously with features of pancytopenia such as pallor, petechiae, and ecchymosis. Recurrent infections and fever occur as a result of neutropenia. Reticulocytopenia is seen and splenomegaly is characteristically absent. Bone marrow biopsy is used for diagnosis.

Histology:

Bone marrow aspirate usually results in a dry tap. Examination shows a hypocellular marrow devoid of hematopoietic cells with remnant fat cells, fibrous stroma, and scattered lymphocytes. Red cells are usually slightly macrocytic and normochromic. The following image shows a hypocellular bone marrow with little hematopoiesis and several fat cells.



Other options:

Option B - Paroxysmal nocturnal hemoglobinuria (PNH) is an acquired disorder resulting from the mutation in the phosphatidylinositol glycan complementation group A gene (PIGA), an enzyme that is essential for the synthesis of membrane-associated complement regulatory proteins. Red cells deficient in these proteins are susceptible to intravascular hemolysis. Patients with PNH usually present with pancytopenia and reticulocytosis with bone marrow biopsy showing a cellular marrow with marked to massive erythroid hyperplasia.

Option C - Myelodysplastic syndromes (MDS) are a group of clonal stem cell disorders. It is a consequence of maturation defects that result in ineffective hematopoiesis with a high predisposition to conversion to AML. They are characterized by a hypercellular marrow with dysplastic cells on bone marrow aspirate. Marrow changes seen in MDS are given in the pearl below.

Option D - Pulmonary tuberculosis (TB) is caused by Mycobacterium tuberculosis and patients often present with fever, productive cough, blood-streaked sputum, night sweats, malaise, loss of appetite, and weight loss.

Solution to Question 11:

The clinical presentation and the investigations are suggestive of Henoch-Schonlein purpura.

Henoch-Schönlein purpura is a systemic immune disorder characterized by a purpuric rash, abdominal pain, polyarthralgia, and renal abnormalities. It is most commonly seen in the age group of 3-8 years. The patient may present with the following clinical features:

• Skin lesions- characteristically involve the extensor surfaces of arms and legs, and also the buttocks

• Abdominal manifestations- include pain, vomiting, and intestinal bleeding.

• Renal manifestations- include gross or microscopic hematuria, nephritic syndrome, or nephrotic syndrome. Renal manifestations are more severe in adults. A rapidly progressive form of glomerulonephritis is seen in some cases.

This condition usually occurs following an upper respiratory tract infection and there is a history of atopy in one-third of the cases. IgA is deposited in the glomerular mesangium in a distribution similar to that of IgA nephropathy.

Microscopic findings:

• IgA deposition in the mesangial region. Sometimes there is the deposition of IgG and C3 along with IgA.

• Renal lesions show focal or diffuse mesangial proliferation, and/or endocapillary proliferation to crescentic glomerulonephritis.

• Skin lesions show subepidermal hemorrhages and necrotizing vasculitis of small vessels.

Other options:

Option A: Immune thrombocytopenic purpura(ITP): It occurs due to antibody-mediated destruction of platelets. It can be acute or chronic. Acute ITP is usually self-limiting in nature. Cutaneous bleeding can be in the form of petechiae and ecchymoses. The patient can also have hematuria, melena, or excessive menstrual flow.

Option B: Dengue: It is a viral illness caused by the dengue virus. The clinical manifestations include fever, headache, musculoskeletal pain, and rash. The rash which is maculopapular in nature begins on the trunk and then spreads centrifugally to the face and extremities.

Option C: Churg- Strauss syndrome (Eosinophilic granulomatosis with polyangiitis): It is characterized by eosinophilic vasculitis of multiple organs. The features of this condition include asthma, peripheral eosinophilia, pulmonary infiltrates, neuropathy, paranasal sinus abnormality, and the presence of eosinophilic vasculitis.

Solution to Question 12:

The given clinical scenario showing fatigue, tiredness and cervical lymphadenopathy with CD10 positive BCL2 positive cells is highly suggestive of follicular lymphoma.

Follicular lymphoma presents at middle age with no sex predilection. The most common presenting symptom is painless generalized lymphadenopathy. A lymph node biopsy is adequate for the diagnosis of follicular lymphoma. Immunophenotyping of the cells showing CD19, CD20, CD10, BCL2 positive, and CD5 negative is suggestive that the cells originate from the germinal centre.

The majority of follicular lymphomas have characteristic (14;18) translocation. This juxtaposes the IGH locus on chromosome 14 and the BCL2 locus on chromosome 18. This chromosomal rearrangement results in the overexpression of BCL2 protein, which is an inhibitor of apoptosis that contributes to lymphoma cell survival.

Histopathology:

Lymph nodes exhibit a nodular or nodular and diffuse growth pattern. Two cell types are predominantly present in varying proportions:

• Centrocytes (small cleaved cells) - Small cells with a cleaved nucleus and scant cytoplasm

• Centroblasts - Large cells with open nuclear chromatin, several nucleoli, and modest amounts of cytoplasm

In most cases, there is a majority of centrocytes.

The peripheral blood smear shows lymphocytosis and bone marrow tap reveals paratrabecular lymphoid aggregates.

The following image shows a lymph node biopsy in follicular lymphoma:



Other options -

Option B - Burkitt lymphoma is an aggressive type of non-Hodgkin lymphoma (NHL). It is also known as small non-cleaved-cell lymphoma and is characterized by the translocation and deregulation of the c-MYC gene on chromosome 8 to chromosome 14 or 22 - t(8:14) or t(8:22). Histopathology of the tumor reveals a starry sky appearance created by apoptotic cells and macrophages containing clear cytoplasm.

Option C - Mycosis fungoides is the most common type of cutaneous T-cell lymphoma. The histological hallmark of cutaneous T-cell lymphoma (mycosis fungoides type) is the presence

of Pautrier microabscess. The presence of B-cell markers in the question rules out the differential of mycosis fungoides.

Option D - Chronic Myeloid Leukemia (CML) is a myeloproliferative neoplasm characterized by the presence of chimeric BCR-ABL gene which is a result of a reciprocal (9;22)(q34;q11) translocation (Philadelphia chromosome). The presence of lymphadenopathy in the given scenario rules out the possibility of CML as it is a myeloid disorder.

Solution to Question 13:

The given laboratory values showing anemia and increased MCV are suggestive of megaloblastic anemia. This could be due to cyanocobalamin deficiency or folic acid deficiency. The history of inadequate intake of fruits and vegetables points towards folic acid deficiency.

Dietary sources of Vit-B12 are animal products such as meat, fish, milk, and eggs. Dietary sources of folic acid are green vegetables such as lettuce, spinach, asparagus, and broccoli.

Megaloblastic anemia can occur due to Vit-B12 or folic acid deficiency. Inadequate intake, intrinsic factor deficiency, pernicious anemia, gastrectomy, malabsorption, ileal resection, alcoholism, anticonvulsants, oral contraceptives, pregnancy, infancy, disseminated cancer, markedly increased hematopoiesis, folic acid antagonists are some of the causes.

Patient can present with anorexia, weight loss, diarrhea, constipation, glossitis, angular cheilosis, reversible melanin skin hyperpigmentation, paresthesias, muscle weakness, dementia, or psychotic disturbances.

In megaloblastic anemia, DNA synthesis is impaired that leading to ineffective hematopoiesis and distinctive morphologic changes.

Peripheral blood examination shows oval macrocytes, usually with anisocytosis and poikilocytosis. MCV is usually >100 fL. The normal value of MCV is between 80-100 fL. Most macrocytes lack the central pallor of normal red cells and even appear hyperchromic, but the MCHC is not elevated. The reticulocyte count is low. Neutrophils are larger than normal and show nuclear hypersegmentation with five or more nuclear lobules.

Marrow is markedly hypercellular. Megaloblastic changes are seen at all stages of erythroid development. There is nuclear-to-cytoplasmic asynchrony, where nuclear maturation is slower than cytoplasmic maturation. Granulocytic precursors are seen in the form of giant metamyelocytes and band forms. Megakaryocytes may be abnormally large with bizarre, multilobate nuclei.

Peripheral blood smear showing hypersegmented neutrophil:



Other options:

Option A - In cyanocobalamin deficiency, MCV will be increased. But the dietary sources of cyanocobalamin are mainly of animal products. In the given clinical scenario there is inadequate intake of fruits and vegetables. Hence, this is not the answer.

Option C - In iron deficiency anemia, MCV will be decreased.

Option D - In hereditary spherocytosis, MCV will be normal.

Solution to Question 14:

The clinical scenario given here describes alcoholic hepatitis. In this condition, there are intracytoplasmic eosinophilic inclusions called Mallory Denk bodies and they contain intermediate filaments.

The liver undergoes inflammation and fibrosis in alcoholic hepatitis. This leads to injured and swollen hepatocytes with cleared-out cytoplasm and cytoskeletal damage. These hepatocytes are called ballooned hepatocytes. When the cytoskeletal damage is extensive, there is the formation of Mallory Denk/ Mallory hyaline bodies. These are remnants of partially degraded intermediate filaments and are composed of keratin filaments.

Cytoskeletal proteins include intermediate filaments, microfilaments (made up of actin - option B), and microtubules (option - C).

Intermediate filaments provide a flexible scaffold to organize cytoplasm and prevent external forces from acting on the cell. They are of five types:

- Keratin filaments (epithelial cells)
- Neuro filaments (neurons)
- Desmin filaments (muscles)
- Vimentin filaments (connective tissues)

• Glial filaments (glial cells)

Fibronectin (option D) is an adhesive glycoprotein. Its presence in cervicovaginal secretions before rupture of membranes is a possible marker for impending preterm labor.

The given image shows intracytoplasmic eosinophilic Mallory bodies.



Solution to Question 15:

The most probable diagnosis for the clinical presentation of mediastinal mass with pure red cell aplasia is thymoma.

Thymoma refers to the tumors of the thymic epithelial cells. Most of the thymic tumors occur in the anterior mediastinum. Clinical features can be chest pain, dyspnea, cough, or superior vena cava syndrome.

It is associated with paraneoplastic syndromes like myasthenia gravis, pure red cell aplasia, or hypogammaglobulinemia. It is rarely associated with limbic encephalitis, hemolytic anemia, aplastic anemia, and other autoimmune conditions like Sjogren's syndrome.

Pure red cell aplasia: It is a marrow disorder where the erythroid progenitors are only suppressed. The complete absence of red cell progenitors is seen in severe cases.

Causes of pure red cell aplasia: It can be hereditary or acquired. Diamond-Blackfan anemia or congenital pure red cell aplasia is the hereditary cause. Causes of acquired pure red cell aplasia are given below:

• Cancer

• Thymoma

• Lymphoid malignancies (eg. Non-Hodgkin's lymphoma, large granular lymphocytosis)

• Paraneoplastic to solid tumors

• Connective tissue disorders with immunologic abnormalities

• Systemic lupus erythematosus, rheumatoid arthritis

• Viruses

• Persistent B19 parvovirus, hepatitis, adult T cell leukemia virus, Epstein-Barr virus

• Drugs

• Phenytoin, azathioprine, chloramphenicol, procainamide, isoniazid

• Multiple endocrine gland insufficiency

• Antibodies to erythropoietin

• Idiopathic

Other options:

Option B: Non-Hodgkin's lymphoma can cause pure red cell aplasia but it commonly presents as peripheral adenopathy rather than mediastinal mass.

Options C and D: Germ cell tumors and bronchogenic carcinoma are not commonly known to be associated with red cell aplasia.

Pathology INI-CET 2022

Question 1:

All of the following are seen in anemia of chronic disease except:

a) High serum hepcidin

b) Low serum ferritin

c) Chronic infection

d) Adequate iron stores

Question 2:

Neurosecretory bodies are seen in all except

a) Medullary thyroid carcinoma

b) Pituitary adenoma

c) Paraganglioma

d) Adrenal cortical tumor

Question 3:

Identify the true statement/s regarding clear cell carcinoma of the kidney

a) 1, 2, 4

b) 2, 3

c) 1, 3

d) 1, 4

Question 4:

The phenomenon of a differential effect of the same gene from different parents is called:

a) Gonadal mosaicism

b) Genomic imprinting

c) Anticipation

d) Incomplete penetrance

Question 5:

Blood transfusion requires screening of donor blood for all of the following infections except:

a) Hepatitis C virus

b) Dengue virus

c) Human immunodeficiency virus

d) Hepatitis B virus

Question 6:

Match the microscopic appearance in column A with the type of tumor in column B

a) 1-b, 2-d, 3-a, 4-c

b) 1-a, 2-d, 3-c, 4-b

c) 1-b, 2-d, 3-c, 4-a

d) 1-c, 2-d, 3-a, 4-b

Question 7:

A patient suffering from cholelithiasis underwent cholecystectomy. The image of the specimen is shown below. Which of the following is the correct diagnosis?

a) Cholesterolosis

b) Gallbladder polyp

c) Carcinoma gallbladder

d) Strawberry gallbladder

Question 8:

A 15-year-old boy presents with swelling in the upper diaphyseal part of the tibia. X-ray of the swelling showed a periosteal reaction. A histopathological image of the biopsy is given below. Immunohistochemistry was positive for synaptophysin and mic 2. What is the likely diagnosis?



a) Osteosarcoma

b) Ewing sarcoma

c) Metastatic neuroblastoma

d) Osteoclastoma

Question 9:

Which of the following is a marker of natural killer cell?

a) CD 68

b) CD 3

c) CD 25

d) CD 94

Question 10:

Which of the following conditions is not caused by high-risk strains of human papillomavirus?

a) Cervical cancer

b) Anal cancer

c) Oropharyngeal cancer

d) Laryngeal papillomatosis

Question 11:

A boy presents with recurrent episodes of melena. On examination, there was eczema of the skin with multiple petechiae. His platelet count was low but RBC and WBC counts were within normal limits. Which of the following is the diagnostic investigation in this patient?

a) WASP gene mutation

b) Anti platelet antibodies

c) Bone marrow biopsy

d) Electron microscopy

Question 12:

Which of the following is not a part of innate immunity?

a) Alpha defensins

b) B lymphocytes

c) Gamma Delta T lymphocytes

d) Lectin pathway

Question 13:

Which of the following disorders is inherited in an X-linked dominant pattern?

a) Rett syndrome

b) Duchenne muscular dystrophy

c) Huntington disease

d) Tuberous sclerosis

Question 14:

Cell in cell appearance is seen in:

a) Necrosis

b) Apoptosis

c) Necroptosis

d) Entosis

Question 15:

Identify the cell marked in the image below



a) Mast cell

b) Macrophage

c) Plasma cell

d) Fibroblast

Question 16:

Match the malignancies in column A with the genes in column B

a) 1-c, 2-b, 3-d, 4-a

b) 1-b, 2-c, 3-d, 4-a

c) 1-d, 2-c, 3-b, 4-a

d) 1-b, 2-d, 3-c, 4-a

Question 17:

How does phagolysosome cause cell death?

a) Is both intra and extra cellular

b) Can fuse with other organelles directly

c) Fuses with lysosome to form autophagolysosome

d) Is a single membrane structure

Question 18:

Which of the following features is not observed in pleomorphic adenoma?

a) Myoepithelial cells

b) Keratin pearls

c) Chondroid differentiation

d) Perineural invasion

Question 19:

A 60-year-old woman presents with complaints of fatigue and dyspnea for 2 weeks. On examination, pallor was seen and hemoglobin was found to be 6 gm/dL. What finding will be seen on peripheral smear?

a) Poikilocytes

b) Microcytic hypochromic cells

c) Macrocytic cells

d) Target cell

Question 20:

The image below is indicative of which of the following tumors?



a) Yolk sac tumor

b) Dysgerminoma

c) Krukenberg tumor

d) Granulosa cell tumour

Question 21:

In a patient with bare lymphocytic disease, graft vs host reaction does not occur due to:

a) Complement deficiency

b) Deficiency of CD4 T lymphocytes

c) MHC class II deficiency on lymphocytes

d) MHC class I deficiency on lymphocytes

Question 22:

A child presents with multiple skin lesions over the inner thigh and genital areas. It initially started as macule and later developed to become nodules. Some nodules have central umbilication. The histology of the lesion is shown below. What would be found on biopsy?



a) Multinucleate giant cells (Tzanck cells)

b) Donovan bodies

c) Henderson-Patterson bodies

d) Koilocytes

Question 23:

Investigations of a patient being evaluated for leukemia revealed a WBC count of 4,50,000/µL, with eosinophilia and basophilia. What is the likely diagnosis in this case?

a) Acute lymphoblastic leukemia

b) Chronic myelogenous leukemia

c) Chronic lymphocytic leukemia

d) Acute myeloid leukemia

Question 24:

Which of the following molecules helps the leukocytes roll over the endothelial cells in the vessels?



a) Cadherins

b) Platelet endothelial cell adhesion molecule-1 (PECAM-1)

c) E-selectin

d) Cd11a

Question 25:

Identify the given bone marrow biopsy instrument.

a) Klima needle

b) Salah's needle

c) Jamshidi needle

d) Islam needle

Question 26:

Match the following stains and the tissue.

a) 1- d, 2- c, 3- b , 4- a

b) 1- c, 2- a, 3- d , 4- b

c) 1- a, 2- b, 3- d, 4- c

d) 1- b, 2- a, 3- c , 4- d

Question 27:

Which of the following statements are true about hereditary spherocytosis?

a) Gall stones develop by 4-5 years of age.

b) Splenectomy does not protect against developing gall stones.

c) At least 25 % cases have a family history of HS.

d) High MCH is pathognomonic of HS.

Question 28:

B cells express IgM and IgD antibodies at the same time due to which phenomenon?

a) Somatic hypermutation

b) Alternative RNA splicing

c) Affinity maturation

d) Allelic exclusion

Question 29:

Which of the following are true regarding sirtuins?

a) 1, 2, 3

b) 1, 2, 3, 4

c) 2, 3

d) 2, 4

Question 30:

Match the following cells with their locations.

a) 1-a, 2-b, 3-c, 4-d

b) 1-c, 2-d, 3-a, 4-b

c) 1-d, 2-a, 3-d, 4-c

d) 1-b, 2-a, 3-c, 4-d

Question 31:

Which of the following cytokines is not involved in Th1 response?

a) IFN gamma

b) IL- 2

c) IL- 12

d) IL- 4

Question 32:

A 19-year-old girl presents with a midline neck swelling that moves on deglutition. USG revealed a 2 cm x 2 cm nodule on the left lobe of the thyroid, extending into the strap muscles. The histopathology image is given below. What is the most likely diagnosis?



a) Medullary CA thyroid

b) Follicular CA thyroid

c) Thyroglossal cyst

d) Papillary CA thyroid

Question 33:

A chronic smoker with a smoking history of 24 packs presents with breathlessness and chest pain. Further investigations revealed the presence of a lung tumor. On staining, cells are positive for chromogranin. What is the diagnosis?

a) Adenocarcinoma

b) Squamous cell carcinoma

c) Large cell carcinoma

d) Small cell carcinoma

Question 34:

A male patient presented with hemoptysis. On examination, hilar mass was present. The histopathological image is shown below. Immunohistochemistry revealed that the cells were positive for p40. What is the diagnosis?



a) Squamous cell carcinoma

b) Adenocarcinoma lung

c) Small cell cancer lung

d) Large cell carcinoma

Question 35:

Which of the following is true about proto-oncogenes?

a) 2 only

b) 3 only

c) 1 and 4

d) 2 and 3

Question 36:

The given peripheral smear is associated with which of the following leukemias?



a) Acute lymphoblastic leukemia

b) Chronic lymphocytic leukemia

c) Acute myeloid leukemia

d) Chronic myelogenous leukemia

Question 37:

Which of the following statements are true about familial adenomatous polyposis?

a) 2 and 4

b) 1 and 4

c) 2 and 3

d) 3 and 4

Question 38:

Which of the following shows maximum amplitude on thromboelastography?

a) Platelet

b) Intrinsic

c) Extrinsic

d) vWD

Question 39:

A man presented with fever, night sweats, and weight loss for 2 months. On examination, non-tender, rubbery lymph nodes were palpable in the supraclavicular and inguinal regions on both sides. Peripheral smear showed the presence of bi-nucleated cells. Which markers will be seen in this condition?

a) 1, 2, 4

b) 2, 3, 4

c) 2 and 3

d) 1 and 2

Question 40:

A 10-year-old boy presented with fatigue. Investigations revealed: Hemoglobin, 9 g/dL; MCV, 60 fL; MCH, 20 pg; and serum ferritin, 185 µg/L. The TLC was elevated and showed predominant lymphocytes and neutrophils. What is the likely diagnosis in this patient?

a) Thalassemia

b) Anemia of chronic disorder

c) Iron-deficiency anemia

d) Sickle cell anemia

Question 41:

Senile atrophy occurs due to _____.

a) Denervation

b) Decreased nutrition

c) Decreased workload

d) Reduced blood supply

Question 42:

Cyclooxygenase plays a role in which pathway?

a) Kreb's cycle

b) Prostaglandin

c) Leukotriene

d) Vitamin K

Question 43:

Metastatic calcification can be explained by all, except _____.

a) Renal failure

b) Accumulation of calcium phosphate

c) Vitamin D deficiency

d) Parathyroid activation/ secretion

Question 44:

Match the cytokines with the effector cells .

a) 1-D;2-A;3-C;4-B

b) 1-C;2-A;3-B;4-D

c) 1-C;2-B;3-D;4-A

d) 1-D;2-A;3-B;4-C

Question 45:

The following is a histopathological image of thyroid pathology. What is the diagnosis?



a) Medullary carcinoma of thyroid

b) Papillary carcinoma of thyroid

c) Follicular carcinoma of thyroid

d) Anaplastic carcinoma of thyroid

Question 46:

A pediatric patient is diagnosed with B cell leukemia. Which of the following are poor prognostic factors?

a) A, C, D

b) C, D

c) A, B, C, D

d) B, C, D

Question 47:

The Hb is 5g/dL and the absolute reticulocyte count percentage is 9%. What is the corrected reticulocyte count?

a) 5

b) 4.5

c) 3

d) 1.8

Question 48:

A 7-year-old child presents with fever, erythematous rash, and bone pain. X-ray of the skull shows multiple lytic lesions. Skin biopsy shows prominent nuclear grooves with eosinophils as in the image below. Immunohistochemistry for S100, CD1a is also shown in the images below. What would be the diagnosis?



a) Rosai-Dorfman disease

b) Kimura disease

c) Langerhans cell histiocytosis

d) Juvenile Xanthogranuloma

Question 49:

Match the following cell types with their associated malignancy.

a) 1-b, 2-a, 3-d, 4-c

b) 1-a, 2-d, 3-c, 4-b

c) 1-b, 2-d, 3-a, 4-c

d) 1-a, 2-b, 3-d, 4-c

Question 50:

Hereditary angioedema is due to a defect in which of the proteins mentioned below?

a) C1 inhibitor

b) Decay accelerating factor

c) CD59

d) C5 convertase

Question 51:

A 35-year-old woman presents with fatigue. Investigations revealed the following: Hb, 5 gm %; MCH, 24; low MCV; leukocytes, 11,000/uL, and platelets, 5 lakhs. The peripheral smear is shown below. What is the diagnosis?



a) Megaloblastic anemia

b) Iron-deficiency anemia

c) Thalassemia major

d) Essential thrombocytosis

Question 52:

Which of the following statements is true about interleukin-12?

a) 1 and 2

b) 1 and 4

c) 2 and 3

d) 3 and 4

Question 53:

A chronic smoker presents to the clinic with shortness of breath. A mass is noted at the lower lobe periphery of the lung. A diagnosis of lung adenocarcinoma is made. The biopsy will be positive for:

a) 1, 3

b) 2, 4

c) 3, 4

d) 2, 3

Question 54:

The following image shows an instrument used to estimate hemoglobin levels. What does it measure?

a) Alkaline hematin

b) Acid hematin

c) Oxyhemoglobin

d) Deoxyhemoglobin

Question 55:

Iron in tissues is stained by:

a) Masson's trichrome

b) Prussian blue

c) PAS

d) Fite-Faraco

Question 56:

A woman presents with lesions on the inner thighs and peri-anal region. They are nodular, 4-6 mm in size, and appear pale. The histopathological image shows multiple intracytoplasmic inclusion bodies consistent with Henderson-Patterson bodies. The diagnosis is:



a) Trichodysplasia spinulosa

b) Condyloma acuminata

c) Molluscum contagiosum

d) Donovanosis

Question 57:

Jarisch-Herxheimer reaction is caused by all of the following cytokines except:

a) TNF alpha

b) IL-6

c) IL-8

d) INF- γ

Answer Key

Question No.	Correct Option
1	b
2	d
3	d
4	b
5	b
6	d
7	b
8	b
9	d
10	d
11	a
12	b
13	a
14	d
15	a
16	c
17	c
18	d
19	b
20	a
21	c
22	c
23	b
24	c
25	a
26	c
27	c
28	b
29	b

30	d
31	d
32	d
33	d
34	a
35	b
36	b
37	a
38	a
39	d
40	b
41	d
42	b
43	c
44	b
45	a
46	d
47	c
48	c
49	a
50	a
51	b
52	b
53	a
54	b
55	b
56	c
57	d

Detailed Explanations

Solution to Question 1:

Low serum ferritin levels are not seen in anemia of chronic disease.

Anemia of chronic disease is anemia resulting due to chronic diseases which produce systemic inflammation. During chronic inflammation, inflammatory mediators like interleukin - 6(IL-6) stimulate the production of hepcidin from the liver. Hepcidin is an acute phase reactant that

inhibits ferroportin on macrophages and also decreases iron from stores to developing RBC precursors. This results in decreased proliferation of erythroid progenitors due to impaired iron utilization.

By decreasing available serum iron, the body defends itself from certain bacteria(Eg, Hemophilus influenza) that require iron for their pathogenicity. The lab findings seen in anemia of chronic disease are:

• Low serum iron

• Decreased total iron-binding capacity

• Abundant stored iron

• Increased hepcidin levels

• Increased serum ferritin

The chronic illnesses associated with anemia of chronic disease can be grouped into three categories:

• Chronic microbial infections, such as osteomyelitis, bacterial endocarditis, and lung abscess

• Chronic immune disorders, such as rheumatoid arthritis and regional enteritis

• Neoplasms, such as carcinomas of the lung and breast and Hodgkin lymphoma

Note: Hepcidin is the main regulator of iron absorption which is encoded by the HAMP gene and is defective in hemochromatosis.

Given below is an image of iron absorption that shows ferroportin being inhibited by hepcidin at intestinal mucosal cells



Solution to Question 2:

Neurosecretory bodies are not seen in adrenal cortical tumours.

Most of the adrenocortical neoplasms are aldosterone-producing adenomas. Conn syndrome is primary hyperaldosteronism caused by a solitary aldosterone-secreting adenoma.

The clinical features of hyperaldosteronism are secondary hypertension and various neuromuscular manifestations resulting from hypokalemia.

Microscopically, aldosterone-producing adenomas are found to have eosinophilic, laminated cytoplasmic inclusions, known as spironolactone bodies, found after treatment with the antihypertensive drug spironolactone.

The image given below shows a microscopic image of spironolactone bodies found in adrenal adenomas



Other options:

Option A: Medullary thyroid carcinoma arises from parafollicular C cells and they produce calcitonin. They are associated with a point mutation in the RET proto-oncogene and multiple endocrine neoplasias (MEN 2A and 2B). Microscopically, polygonal cells in an amyloid stroma are seen. Electron microscopy reveals neurosecretory granules within the cytoplasm.

Option B: Pituitary adenoma may present with visual field abnormalities or pituitary apoplexy (acute haemorrhage into an adenoma). Prolactinoma is the most common type of hyperfunctioning pituitary adenoma. Histologically, prolactinomas are composed of chromophobe cells with cytoplasmic neurosecretory granules.

Option C: Adrenal medullary pheochromocytomas are the most common paraganglioma. Extra-adrenal paragangliomas occur in the head and neck at carotid bodies. Microscopically, carotid body tumours contain nests (zellballen) of chief cells. Electron microscopy shows well-demarcated neurosecretory granules.

Solution to Question 3:

True statements regarding clear cell carcinoma of the kidney are 1, 4. Clear cell carcinoma is the most common sporadic renal cell cancer and cells show intracellular accumulation of glycogen

and fat.

Clear cell cancer arises from the proximal tubular epithelium and usually occurs as solitary unilateral lesions.

Most of the cases are sporadic, but can also occur either in familial forms or in association with von Hippel-Lindau (VHL) disease. They are associated with the loss of sequence of the short arm of chromosome 3. Those with VHL disease inherit a germline mutation of the VHL gene on chromosomal band 3p25 and lose the second allele by somatic mutation.

The VHL protein causes the degradation of hypoxia-induced factors (HIFs), which are transcription factors that contribute to carcinogenesis by stimulating the expression of vascular endothelial growth factor (VEGF) that helps drive tumour cell growth.

Histologically, the tumour cells have a rounded or polygonal shape and abundant clear or granular cytoplasm, which contains glycogen and lipids.

Given below is the histopathological image of clear cell carcinoma.



Cancerous cells produce various abnormal hormones resulting in paraneoplastic syndrome. These include polycythemia, hypercalcemia, cushing syndrome, amyloidosis and hypertension.

The most common site of metastasis is the lungs and bones. Radical nephrectomy is the treatment of choice. But T1a(size less than 4cm) tumours are treated by nephron-sparing surgery or partial nephrectomy to preserve renal function. Drugs that inhibit VEGF and tyrosine kinase inhibitors are used as an adjunct therapy for metastatic disease.

Solution to Question 4:

The phenomenon of a differential effect of the same gene from different parents is called genomic imprinting.

Genomic imprinting refers to gene silencing/ inactivation. Here either the maternal or the paternal allele is inactivated resulting in only one functional allele. It occurs by the process of

epigenetics, where gene expression is lowered by:

• DNA methylation

• Histone deacetylation

Genomic imprinting is best illustrated by two uncommon genetic disorders: Prader-Willi syndrome and Angelman syndrome.

The image given below is a child with Prader-Willi syndrome

	Prader-Willi syndrome	AngelMan syndrome
Deletion(most common)	Paternal chromosome 15	Maternal chromosome 15
Uniparental disomy(less common)	Maternal disomy(duplication)	Paternal disomy(duplication)
Clinical features	Elevated ghrelin levels - hyperphagiaTruncal obesitySevere hypotoniaShort statureSmall hands and feet with hypotoniaHypogonadotropic hypogonadismCognitive impairment	Short statureSeveremental retardationwith lack of speechStiff arm movementsSpastic, uncoordinated walk (ataxia)SeizuresInappropriate outbursts of laughter (happy puppets)

The image given below is a child with Angelman syndrome:

Diagnosis of these syndromes is based on the assessment of the methylation status of marker genes and fluorescence in-situ hybridisation(FISH).

Other options:

Option A: Gonadal mosaicism is a mutation that occurs only in egg or sperm cells and the individual will be phenotypically normal. These individuals with gonadal mosaicism can transmit the disease-causing mutation to the offspring through their mutated

gametes.

Option C: Anticipation refers to earlier onset of disease or increased severity in succeeding generations. This is seen in trinucleotide repeat disorders like Huntington's disease.

Option D: Incomplete penetrance refers to variations in penetrance and expressivity of clinical features. Not all individuals who inherit the mutant gene show a mutant phenotype.

Mnemonic:

Prader–Willi Syndrome: Paternal gene deletion - P for P

Angelman syndrome: Maternal gene deletion (Mothers are Angels, Maternal gene deletion leads to Angelman syndrome)

Solution to Question 5:

Blood transfusion does not requires screening of donor blood for dengue virus infection.

Some of the transfusion transmitted infections routienly tested are:

• Hepatitis C virus

• Human immunodefeciency virus(HIV)

• Hepatitis B virus

Transfusion-transmitted disease not currently, routinely tested are:

- Hepatitis A virus
- Hepatitis E virus
- Parvovirus B19
- Dengue fever virus
- Malaria
- Zika virus

Solution to Question 6:

The correctly matched option of the microscopic appearance with the type of tumor is 1-c, 2-d, 3-a, 4-b.

- Schiller Duval bodies - Yolk sac tumor
- Presence of teeth, hair, etc - Teratoma
- Signet ring cells - Kruckenberg tumor
- Call-Exner bodies - Granulosa cell tumor

Schiller Duval bodies are seen in yolk sac tumors. A yolk sac tumor/endodermal sinus tumor is an aggressive malignant germ cell tumor of the ovaries. Most of the yolk sac tumors have Schiller-Duval bodies and they are pathognomonic. These are glomerulus-like structures consisting of a central vessel surrounded by tumor cells within a space that is also lined by tumor cells.

The image given below shows Schiller-Duval bodies seen in yolk sac tumors.



Presence of hair, teeth, etc is seen in teratomas. These are germ cell tumors. They contain embryonic tissues leading to the differentiation of cells along these pathways. Apart from teeth, hair may also contain skin, sebaceous glands, fat, bone, cartilage, etc.

There are three classes of teratomas:

• Mature teratomas (eg - dermoid cyst)

• Immature teratomas (eg - solid teratoma)

• Monodermal highly specialized tumors (eg - Struma ovarii)

Given below is the histopathological image of immature teratoma of the ovary, with immature tissue of the bone (asterisk), cartilage (arrow), and islands of adipose tissue (arrowhead)



Signet-ring cells are seen in the Krukenberg tumor. Krukenberg tumor is the carcinoma of the stomach that metastasize to the ovary. Signet ring cells are mucin-producing cells. The appearance of a signet ring often originates from the gastric mucosa.

The image below shows signet-ring cells in the Kruckenberg tumor.



Option D: Granulosa cell tumours arise from granulosa cells which form part of the developing ovarian follicle. They are unilateral in nature and the mass is cystic and encapsulated. Call-Exner

bodies are seen in granulosa cell tumors. These are gland like small structures with acidophilic material and immature follicles.

The image given below shows Call-Exner bodies seen in granulosa cell tumours.



Solution to Question 7:

The correct diagnosis in the patient is gallbladder polyp.

Gallbladder polyp is a pre-malignant condition that usually presents with biliary colic and cholecystitis. Cholesterol polyps are more common while adenomatous polyps have malignant potential.

Risk factors of malignancy in adenomatous polyp:

• Age >60 years

• Coexistence of gall stones

• The documented increase in size

• Size of >10 mm.

Polyps less than 1 cm can be followed with serial ultrasonography. The USG given below depicts gallbladder polyps that are fixed and do not cast any acoustic shadow. The polyps larger than 1 cm have an increased risk of malignancy and should be treated by performing an open cholecystectomy.

The image below shows an intraoperative photograph of gallbladder polyps.



The image given below shows the management of gall bladder polyp



Other options:

Options A and D: Cholesterosis or strawberry gallbladder is a condition characterized by a collection of submucous aggregations of cholesterol crystals and cholesterol esters in the gallbladder. The interior of the gallbladder looks like a strawberry with yellow specks.

The image given below shows the gross appearance of cholesterosis of the gallbladder



Option C: Gallbladder adenocarcinoma is the most common malignancy of the extrahepatic biliary tract. Gallstones are the most important risk factor. Her-2/neu is overexpressed and CA 19-9 is used for monitoring. The most important prognostic factor is the depth of invasion (T-stage). On gross appearance, they resemble chronic cholecystitis.

Solution to Question 8:

The given clinical scenario along with the image showing small round blue cells and immunohistochemistry (IHC) positive for synaptophysin and mic-2 point to a likely diagnosis of Ewing sarcoma.

Ewing sarcoma is the 2nd most common bone sarcoma in children. The majority of the patients affected are usually <20 years of age. It has a slight male predominance.

It commonly involves the diaphysis of the long bones like the femur, tibia, fibula, and humerus and flat bones of the pelvis. The majority of Ewing sarcomas have a balanced translocation between the EWSR1 gene on chromosome 22 and the FLI1 gene on chromosome 11.

Patients present with painful enlarging mass, fever, and anemia. The affected area is usually tender, swollen, and warm. Laboratory findings may show an elevated WBC count and erythrocyte sedimentation rate (ESR).

A histopathological examination of Ewing sarcoma shows sheets of uniform, round cells arranged in Homer-Wright rosettes. These cells are of neuroectodermal origin and have minimal cytoplasm and abundant glycogen. When they are arranged around a central fibrillary core in the form of clusters, the appearance is termed Homer-Wright rosettes. Immunohistochemistry is usually positive for synaptophysin and mic-2.

Given below is the histopathological appearance of Ewing sarcoma.



As tumor cells are rich in glycogen, they stain positive for periodic acid Schiff (PAS) stain however, they stain negatively for reticulin.

The image given below shows PAS staining done in Ewing's sarcoma:



Radiological examination of Ewing sarcoma shows a destructive lesion in the diaphysis of long bones. Deposition of periosteal reaction in layers gives an onion-peel appearance (shown below).

Other options:

Option A: Osteosarcoma is the most common primary malignancy of bone. The histological features of osteosarcoma include pleomorphic cells with hyperchromatic nuclei and abundant mitoses. The diagnosis is definitive with the presence of malignant tumor cells producing unmineralized osteoid or mineralized bone arranged in a typically fine, lacelike pattern.

Given below is the histopathological image of osteosarcoma:



X-ray shows periosteal reaction and it may take the form of a Codman's triangle or it may have a sunray appearance as shown below.

Option C: Neuroblastoma is a tumor of the sympathetic ganglia and adrenal medulla. It is derived from primordial neural crest cells. It is the most common extracranial solid tumor in childhood. On histopathology small round blue cells are seen. These cells are concentrically arranged around a faintly eosinophilic fibrillary material (neuropil). This appearance is called Homer-Wright pseudorosettes (shown below).



Option D: Osteoclastoma is also known as a giant cell tumour. The tumor cells are primitive osteoblast precursors and express high levels of RANKL. Histologically, the tumors consist of osteoclast-type giant cells with 100 or more nuclei. These cells are usually present against a backdrop of mononuclear stromal cells.

The image given below shows the histopathological appearance of giant cell tumour/osteoclastoma:



The image below shows an X-ray of a giant cell tumour showing the typical soap bubble appearance.

Solution to Question 9:

The marker of the natural killer cells is CD94. It is expressed in natural killer cells and a small subset of T lymphocytes.

Natural killer cells (NK cells) are essential cells of innate immunity. They act against malignant and virus-infected cells. NK cells are identified by CD16 and CD56 cell surface receptors.

NK cells' activity is augmented by interferons and upon activation, they bind to glycoprotein receptors on the surface of target cells and release perforins. Perforins create pores in the membrane of the target cells. Through these pores, cytotoxic factors like tumor necrosis factor-beta (TNF-β) enter the cells and destroy them by apoptosis.

NK cells express a phenomenon known as antibody-dependent cellular cytotoxicity (ADCC). Here, NK cells have the capability of killing target cells that are coated by IgG antibodies.

The image given below shows the actions of NK cells on cancer cells.



Other options:

Option A: CD68 is expressed on macrophages and monocytes.

Option B: CD 3 is a T-cell receptor marker. They are involved in the transduction of signals to the T cells when an antigen binds to the T cell receptor.

Option C: CD 25 is a component of the IL2 receptor and is important in T cell proliferation and activation.

Solution to Question 10:

Laryngeal papillomatosis is not caused by high-risk strains of human papillomavirus (HPV). It is caused by low-risk HPV strains 6 and 11.

Laryngeal papillomatosis or juvenile papillomatosis is the most common benign neoplasm of the larynx in children. It is caused by HPV 6 and 11. It involves supraglottic and glottic regions of the larynx. Adult-onset laryngeal papillomatosis involves the anterior part of the vocal cords.

It usually occurs between the ages 3–5 years, but less commonly can also occur in men aged between 30-50 years. Clinical features include hoarseness or aphonia and respiratory difficulty. The recurrence rate is high but they rarely transform to SCC.

Diagnosis is confirmed by direct laryngoscopy and biopsy. Treatment consists of micro laryngoscopy, CO_2 laser excision, and topical cidofovir.

The image given below shows laryngeal papilloma.

Other options:

Cervical cancer, anal cancer and oropharyngeal cancers are caused by high risk HPV types 16 and 18.

Solution to Question 11:

The given scenario of recurrent episodes of melena, eczema, and multiple petechiae along with low platelet counts is suggestive of Wiskott-Aldrich syndrome. The diagnostic investigation in this patient is WASP gene mutation testing.

Wiskott-Aldrich syndrome(WAS) is an X-linked recessive immunodeficiency disorder. It is caused by mutations in the Wiskott-Aldrich syndrome protein (WASp) that primarily affects the T lymphocytes.

Normally, WASp regulates the actin cytoskeleton and thus helps in cell adhesion, migration, and the formation of synapses between antigen-presenting and target cells. In WAS, there is a progressive loss of T lymphocytes in the peripheral blood resulting in defects in cellular immunity.

The clinical features of WAS are recurrent bacterial infections, eczema, and bleeding due to thrombocytopenia. Patients with WAS are prone to develop B-cell lymphomas and autoimmune disorders.

The image given below shows eczema seen in WAS

Diagnosis is based on intracellular immunofluorescence analysis of WAS protein (WASp) expression in blood cells. IgM levels in the serum are low, but levels of IgG are usually normal. Reduced-sized platelet is a typical feature of WAS.

The only curative treatment is an allogeneic hematopoietic stem cell transplant. Prophylactic antibiotics, immunoglobulin G (IgG) supplementation, and topical treatment for eczema can be tried. Gene therapy trials are also under research.

Solution to Question 12:

B lymphocytes are not a part of innate immunity. B lymphocytes are part of adaptive immunity.

Innate immunity is an intrinsic mechanism that provides the first line of defence against infections. It is non-specific, provides immediate defence against microbes, and helps eliminate damaged cells.

Components of Innate Immunity are:

• Epithelial cells - these produce antimicrobial molecules such as alpha-defensins

• Macrophages and neutrophils

• Dendritic cells - these have receptors that sense microbes and cell damage. They stimulate secretion of cytokines, which play important role in inflammation and antiviral defence

• Innate lymphoid cells (ILCs) - these are local tissue lymphocytes that lack T-cell antigen receptors. They produce inflammatory cytokines during the early phases of immune reactions. Gamma delta T lymphocytes and natural killer (NK) cells belong to this class

• Mast cells

• Complement pathway - it include complement proteins of alternative and lectin pathways.

Solution to Question 13:

Rett syndrome is inherited in an X-linked dominant pattern.

Rett syndrome is a neurodevelopmental disorder. It is caused due to mutation in the MECP-2 gene. The disorder is mostly seen in girls because affected boys usually die in infancy due to a defective single X chromosome.

Symptoms of Rett syndrome are microcephaly, developmental delay, autistic-like behavior, cognitive dysfunction, seizures, and repetitive motor actions.

Other options:

Option B: Duchenne muscular dystrophy (DMD) is inherited in X linked recessive pattern. It occurs due to a mutation in the dystrophin gene. Typical features include calf muscle pseudohypertrophy and Gower's sign (using arms to lift themselves from a seated position). Laboratory investigations show elevated serum creatine kinase levels. Casimersen has been recently approved by the FDA for the treatment of DMD.

Option C: Huntington's disease is inherited in an autosomal dominant pattern. It is a trinucleotide (CAG) repeat disorder and shows genetic anticipation (earlier onset of disease or increased severity in succeeding generations). Patients present with involuntary movements known as chorea. Tetrabenazine is used for treatment.

Option D: Tuberous sclerosis also known as Bournville disease is inherited in an autosomal dominant pattern. It occurs due to mutation in TSC1 and TSC2 genes. It is characterized by subependymal nodules, angiomyolipomas, cafe au lait spots, shagreen patches, and cardiac rhabdomyomas.

Solution to Question 14:

Cell in cell appearance is seen in entosis.

Entosis is a phenomenon where a cell engulfs a cell of the same type. While cells that are internalised by this mechanism are initially viable. But, the majority of them undergo a non-apoptotic form of cell death. It plays a physiological role in the elimination of cells detached from a surface.

Other options:

Option A: Necrosis is a pathologic process of irreversible cell death following severe injury. It is associated with local inflammation due to damaged cell membranes. Some microscopic features of necrosis are as follows:

• Dilation of mitochondria

• Nuclear changes: karyolysis, pyknosis, and karyorrhexis.

• Myelin figures replacing dead cells.

Option B: Apoptosis is a pathway of programmed cell death. In apoptosis, Mitochondrial and death receptor pathways activate caspases resulting in the formation of apoptotic bodies which are then phagocytosed.

Option C: Necroptosis is a type of cell death which is a hybrid of both necrosis and apoptosis. It is also called programmed necrosis/caspase-independent programmed cell death(since it does not result in caspase activation).

Solution to Question 15:

The cell marked in the image is a mast cell stained with toluidine blue. Mast cell granules contain acid proteoglycan that binds basic blue such as toluidine blue.

Mast cells may be difficult to differentiate from lymphocytes in routine, hematoxylin, and eosin-stained sections, and special metachromatic stains (toluidine blue or Giemsa) are used to visualize their granules.

Mast cells are a part of innate immunity and are widely distributed in connective tissues. They play an important role in immediate hypersensitivity and both acute and chronic inflammation.

Mast cells express a receptor on their surface that binds to the Fc portion of the IgE antibody. When an antigen binds to IgE antibodies on the surface of the mast cell, they get activated and degranulate. This results in the release of:

• Granule contents: Histamine, chymase, tryptase, heparin.

• Lipid mediators: Leucotriene C4, D4, prostaglandin D2, platelet-activating factor.

• Cytokines: Tumor necrosis factor, interleukin-1

These mediators released from mast cells play an important role in the pathogenesis of immediate/type 1 hypersensitivity.

The image given below shows the sequence of events in immediate (type 1) hypersensitivity.



Some of the important disorders associated with mast cell activity are bronchial asthma, urticaria, mastocytosis, interstitial cystitis and neurofibroma.

Solution to Question 16:

The correctly matched option is option C 1-d, 2-c, 3-b, and 4-a.

Pleomorphic adenoma is associated with chromosomal rearrangements that induce overexpression of PLAG1. It is the most common salivary gland and lacrimal gland neoplasm. Microscopically, it contains a mixture of ductal (epithelial), myoepithelial, and mesenchymal cells.

Burkitt's lymphoma is a non-Hodgkin lymphoma (NHL). It is associated with the translocation cMYC gene on chromosome 8 to chromosome 14 or 22 - t(8:14) or t(8:22). Histopathology of the tumour reveals a starry sky appearance.

Neuroblastoma is a tumour derived from primordial neural crest cells. It is the most common extracranial solid tumour in childhood. They are associated with the translocation nMYC gene. On histopathology small round blue cells and Homer-Wright pseudorosettes are seen.

Retinoblastoma is the most common primary intraocular malignancy in children. It is associated with chromosomal rearrangements of the Rb gene. Microscopillay, Flexner-Wintersteiner rosettes are seen.

1. Pleomorphic adenoma	d. Plag 1
2. Burkitt lymphoma	c. cMyc
3. Neuroblastoma	b. nMyc
4. Retinoblastoma	a. Rb

Solution to Question 17:

Phagolysosomes cause cell death by the fusion of lysosomes to form autophagolysosomes.

Autophagy is a process in which a cell eats its own contents. Cytoplasmic materials are delivered and degraded by the lysosome. It is a survival technique of the cell in nutrient deprivation states.

The steps are:

• An isolation membrane called phagophore is formed and nucleated.

• Phagophore is derived from the ER with the help of plasma membrane and mitochondria

• A double membrane-bound vesicle is formed from the isolation membrane called the autophagosome

• Maturation of the autophagosome by fusion with lysosomes, called autophagolysosome.

Apoptosis is a regulated mechanism of eliminating unwanted and damaged cells. It is initiated by enzymes called caspases and is executed by two different pathways. The mitochondrial pathway which is also known as the intrinsic pathway carries out the majority of the apoptosis. The death receptor pathway or the extrinsic pathway is initiated by plasma membrane death receptors. Both these pathways combine together and activate the caspase cascade which results

in the final phase of apoptosis.

Pyroptosis is a form of apoptosis mediated by fever-inducing cytokine IL-1. A multiprotein complex called inflammasome is activated when microorganisms enter the cell. It will in turn activate caspase 1 which cleaves the IL-1 precursor and releases its biologically active form.

Solution to Question 18:

Perineural invasion is not observed in pleomorphic adenoma. Perineural invasion is seen in malignant salivary gland tumours like adenoid cystic carcinoma.

Pleomorphic adenoma is the most common salivary gland and lacrimal gland neoplasm. They are commonly seen in the parotid. They are painless, slow-growing, mobile, discrete masses within the parotid.

Histologically, pleomorphic adenoma consists of epithelial and mesenchymal components.

• Epithelial components consist of

• Ductal cells, myoepithelial cells organized as ducts, acini, irregular tubules, or sheets of cells

• Squamous metaplasia with the formation of keratin pearls may be present

• Mesenchymal components consist of

• Loose myxoid and hyaline tissue containing cartilage (due to chondroid differentiation) and bone

The image below shows pleomorphic adenoma of the salivary gland consisting of mixed epithelial (left) and mesenchymal cell components (right).



Clinical features

• Painless, slow-growing swelling

• Lobulated, firm, elastic-to-hard in consistency

• Lift the ear lobe

• Free from deeper structures

FNAC is the investigation of choice in diagnosing pleomorphic adenoma. Open biopsy contraindicated due to chances of injury to the facial nerve. Surgery is the mainstay of treatment by performing superficial parotidectomy.

Solution to Question 19:

Microcytic hypochromic cells will be seen on the peripheral smear. An elderly female presenting with fatigue and dyspnea with severe anaemia is suggestive of iron deficiency anaemia(IDA). The peripheral smear of IDA shows microcytic hypochromic red blood cells.

Iron deficiency is the most common nutritional disorder in the world. IDA results from dietary deficiency, impaired absorption, increased requirement and, chronic blood loss. The cause of iron deficiency anaemia in an older man or post-menopausal woman is always gastrointestinal cancer until proven otherwise.

Clinical presentation may include fatigue, pallor, dyspnea, cheilosis (fissures at the corners of the mouth) and koilonychia (spooning of the fingernails).

Peripheral smear in IDA shows:

• Microcytic hypochromic red blood cells with an enlarged zone of central one-third pallor.

• Poikilocytes: pencil-shaped RBCs and, target cells.

The image given below shows a microcytic hypochromic blood picture in a peripheral smear



IDA is treated by giving oral, parenteral and packed cell transfusions depending on the severity of anaemia.

For oral iron replacement therapy,

• Up to 200mg of elemental iron is given.

• According to the Anemia Mukt Bharat initiative, adult women need to take 60 mg elemental iron + 500 mcg folic acid tablets once a week.

For parenteral iron therapy, the amount of iron needed by an individual is calculated by the formula:

Body weight(kg) x 2.3 x (15 - patient's hemoglobin, g/dL) + 500 or 1000mg (for stores)

Solution to Question 20:

The image shows the characteristic Schiller Duval body consistent with a yolk sac tumor.

A yolk sac tumor/endodermal sinus tumor is an aggressive malignant germ cell tumor of the ovaries. It originates from embryonal structures of the yolk sac by selective differentiation.

It is most oftenly seen in children and young women. It usually presents as abdominal distention and a palpable pelvic mass.

Grossly, the yolk-sac tumor appears yellow, friable (hemorrhagic), and forms solid mass.

Most of the yolk sac tumors have Schiller-Duval bodies and they are pathognomonic. These are glomerulus-like structures consisting of a central vessel surrounded by tumor cells within a space that is also lined by tumor cells.

The image given below shows Schiller-Duval bodies seen in yolk sac tumors.



Yolk-sac tumors are rich in alpha-fetoprotein. Therefore, serum levels of alpha-fetoprotein can serve as a reliable tumor marker in post treatment surveillance.

Other options:

Option B: Dysgerminoma is the ovarian counterpart of seminoma. They are unilateral tumors composed of vesicular cells and clear cytoplasm. The tumor cells grow in cords or sheets that are separated by scanty fibrous stroma. They are usually infiltrated by lymphocytes.

Option C: Krukenberg tumor is the carcinoma of the stomach that metastasize to the ovary. It is characterized by mucin-producing signet ring cells. The appearance of a signet ring often originates from the gastric mucosa.

The image below shows signet-ring cells in Kruckenberg tumor.



Option D: Granulosa cell tumours arise from granulosa cells which form part of the developing ovarian follicle. They are unilateral in nature and the mass is cystic and encapsulated. Call-Exner bodies are seen in granulosa cell tumours. These are gland like small structures with acidophilic material and immature follicles.

The image given below shows Call-Exner bodies seen in granulosa cell tumours.



Solution to Question 21:

In a patient with bare lymphocytic disease, graft vs host reaction does not occur due to MHC class II deficiency on the lymphocytes.

Bare lymphocyte disease occurs due to mutations in transcription factors required for class II MHC gene expression. This results in the lack of class II MHC molecules, leading to the impaired development of CD4+ T cells. As these CD 4+ T cells are required for both for cellular immunity and for assisting B cells, class II MHC deficiency results in a combined immuno- deficiency.

Solution to Question 22:

The given clinical scenario and histological image are diagnostic of molluscum contagiosum lesions, which show intracytoplasmic inclusion bodies, called Henderson–Patterson bodies or molluscum bodies.

Infection is usually spread by direct contact, particularly among children and young adults. Multiple lesions occur on the skin and mucous membranes, with a predilection for the trunk and anogenital areas. Individual lesions are firm, pruritic, pink to skin-colored umbilicated papules. A curd-like material can be expressed from the central umbilication.

Smearing this material onto a glass slide and Giemsa staining shows diagnostic molluscum bodies, which are ellipsoid, homogeneous, cytoplasmic inclusions that are found in the cells of the stratum granulosum and the stratum corneum and are up to 35 μm in size.



Other options:

Option A: Multinucleate giant cells (called Tzanck cells) are seen in herpes simplex virus infections. They are formed when individual epidermal cells containing viral inclusions fuse to produce giant cells. These can be demonstrated by the diagnostic Tzanck test, which is based on the microscopic examination of the vesicle fluid.

Option B: Donovan bodies are seen in donovanosis, caused by Klebsiella granulomatis. Donovan bodies are minute, encapsulated coccobacilli in macrophages, which can be demonstrated in Giemsa-stained smears of the exudate.

Option D: Koilocytes are characteristically seen in HPV infections, such as condylomata acuminata. Koilocytes are lining cells that have perinuclear cytoplasmic vacuolization.

Solution to Question 23:

A WBC count of 4,50,000/μL with eosinophilia and basophilia indicates a probable diagnosis of chronic myelogenous leukemia.

CML is a myeloproliferative neoplasm. It is caused by the translocation of ABL gene on chromosome 9 onto BCR gene on chromosome 22, resulting in the formation of the BCR-ABL fusion gene. This chromosomal abnomality, called Philadelphia chromosome, is characteristic of CML.

The resulting BCR-ABL translocation results in the activation of tyrosine kinase that signals pro-growth and pro-survival pathways like RAS and JAK/STAT pathways resulting in the proliferation of hematopoietic stem cells.

Patients typically present with a dragging sensation abdomen due to splenomegaly or splenic infarct, fatigue, malaise, and weight loss.

On peripheral smear there is leukocytosis with leukocyte count, often exceeding 100,000 cells/mm3 (neutrophils, band forms, metamyelocytes, myelocytes, eosinophils, and basophils), blasts are usually less than 10% of the circulating cells, the platelet count is usually increased.

Basophils and eosinophils are often increased.



Other options:

Option D: Acute myeloid leukemia (AML) is a hematopoietic progenitor malignancy that causes immature myeloblast accumulation in the bone marrow. Incidence increases with age and peaks after 60 years of age.

The WBC count is about 15,000/µL. Few cases have either <5000 or >1,00,000/µL WBC count. Clinically, the patients present with symptoms resulting due to pancytopenia (anemia, neutropenia, thrombocytopenia). For acute myeloblastic leukemia (AML) diagnosis, the blast cell count should be ≥20% in the peripheral blood or bone marrow

Other distinctive features of AML include

• Bone marrow/ peripheral smear shows a large, uniform population of cells (myeloblasts) with prominent nucleoli (yellow arrow). It shows the presence of primary cytoplasmic granules

• Auer rods: needle-shaped azurophilic intracytoplasmic inclusion bodies formed by the fusion of lysosomes. (black arrow)

• Multiple Auer rods clustered together within a single cell resembling a bundle of sticks known as faggot cells (seen in peripheral smear).



Option A: Acute lymphoblastic leukemia (ALL) arises from immature B (pre-B) or T (pre-T) cells called lymphoblasts. Most cases present with a reduced or normal lymphocyte count. Blood counts show anemia, thrombocytopenia, and neutropenia.

Immunohistochemistry for terminal deoxynucleotidyl transferase (TdT) is positive in pre-B and pre-T lymphoblasts. Peripheral smear shows lymphoblasts that have condensed chromatin, less conspicuous nucleoli, and smaller amounts of cytoplasm which lacks granules as shown below.



Option C: Chronic lymphocytic leukemia presents with a mean WBC count of 30*109/L. It is associated with the deletions of 13q, 11q, 17p, and trisomy 12q.

Lymph nodes are diffusely effaced by small lymphocytes with proliferation centers. The image shows small round lymphocytes and smudge cells seen in CLL.



Solution to Question 24:

E-selectin, which is present over the endothelium, helps the leukocyte roll over the endothelium.

Selectins are of 3 types: L-selectin present on leukocytes, E-selectin present on the endothelium, and P-selectin present in platelets and on the endothelium.

The weak rolling action slows down the leukocytes and allows them to bind more firmly to the endothelium with the help of integrins.

The recruitment of leukocytes to the site of injury involves the following steps:

• Lumen of the vessel to the endothelium:

• Margination: Leukocytes assume a peripheral position along the endothelial surface because of the stasis of blood flow.

• Rolling: Leukocytes start rolling on the vessel wall because of signaling molecules from the endothelium.

• Adhesion: Leukocytes are attached to the endothelium because of adhesion molecules that are activated by cytokines.

• Transmigration of the leukocytes across the vascular endothelium.

• Migration towards a chemotactic stimulus in the extravascular tissues.

Other options:

Option A: Cadherins are a type of glycoprotein cell adhesion molecule that allows cells to adhere together.

Option B: Platelet endothelial cell adhesion molecule-1 (PECAM-1), aids in diapedesis, which is the movement of leukocytes through intact endothelium of the capillaries to reach the site of injury.

Option D: Cd11a, a molecule from the integrin family, helps firmly adhere the leukocyte to the endothelial cell.



Solution to Question 25:

The given instrument is a Klima needle.

Klima and Salah's needles are reusable bone marrow aspiration & biopsy needles, both are made of stainless steel. Klima needle has a guard that screws along the length of the needle. It has no side screw. Whereas, Salah's needle has a guard situated 2 cm from the tip and has a side screw.

The image below shows Klima needle.

The image below shows Salah's needle.

Islam and Jamshidi needles have a t-bar at the end for easier maneuvering. Jamshidi needle has a tapering end to reduce the crush artefacts in the sample. Islam needle has a core securing device, i.e., its cutting edge is shaped to hold the core of the bone during the procedure.

The image below shows the Jamshidi needle.

The image below shows the Islam needle.



Solution to Question 26:

The correct match is 1- a, 2- b, 3- d, 4- c.

Perl's Prussian blue stain is used in histopathological studies to detect the presence of iron in the specimen. Iron is stained blue with this stain.

It is used in hereditary hemochromatosis to stain liver tissue, which shows hepatocellular iron. It also stains the iron-laden mitochondria of ring sideroblasts seen in lead toxicity and myelodysplastic syndrome.

The following image shows the blue staining of iron (in the form of hemosiderin) deposited in hepatocytes:

1. Prussian blue stain	a. Iron
2. PAS stain	b. Glycogen
3. Congo red stain	d. Amyloid
4. Fite- Faraco stain	c. Leprosy



PAS stain (Periodic acid Schiff stain) is used for the detection of glycogen in tissues such as the liver, skeletal muscle, etc., and also in tumor cells.

The image below shows intense PAS +ve staining of hepatocytes in glycogen storage disorder:



The stain used for amyloid deposits is congo red staining, which imparts pink to red color to the amyloid under ordinary light and apple-green birefringence under polarized microscopy.

The image shows a congo-red stained biopsy specimen with the following:

(A) Focal positive staining (orange-red) of amorphous extracellular material

(B) Apple-green birefringence under polarized light



Fite-Faraco stain has been used in cases of leprosy for staining Mycobacterium leprae. It can also be used to stain Nocardia.

The image below shows acid-fast bacilli inside foamy macrophages:



Solution to Question 27:

Among the given statements about hereditary spherocytosis, the true statement is option c– at least 25% of cases have a familial inheritance (as 75% of cases show an autosomal dominant inheritance pattern)

Hereditary spherocytosis is a type of hereditary hemolytic anemia that is mostly inherited in an autosomal dominant pattern. Some of the cases can be autosomal recessive or sporadic. Mutations of the red blood cell surface proteins spectrin and ankyrin can result in

unstable cell membranes, leading to hemolysis.

The following mutations have been identified in hereditary spherocytosis:

Autosomal dominant (more common):

• Ankyrin - most common

• Band 3

• β-Spectrin

Autosomal recessive:

• α-Spectrin

• Band 4.2

This disorder can present at a young age with severe jaundice and fatigue. The unconjugated hyperbilirubinemia predisposes the patient to develop gallstones in early adulthood. Pigment stones are found in 40-50% of the affected adults.

Laboratory findings include normocytic anemia with unconjugated hyperbilirubinemia. The peripheral smear will show large spherical red blood cells without central pallor.

MCHC (not, MCH) is increased in hereditary spherocytosis; however, it is not pathognomonic as it could be found in other conditions as well. MCV is mildly decreased because of cell shrinkage. There is mild to moderate anemia, therefore, MCH could be slightly lower.

The diagnosis can be confirmed by the acid glycerol lysis test and the eosin-5′-maleimide (EMA)–binding test.

There is no specific treatment available besides splenectomy. Spleen is the major site where spherocytes undergo hemolysis. Hence, splenectomy corrects the anemia and its associated complications, thereby playing a role in preventing gallstones. It is done at 4–6 years of age in severe cases and after puberty in moderate cases. It should be followed up with the appropriate antibiotic and vaccine prophylaxis.

Solution to Question 28:

B cells express IgM and IgD antibodies at the same time due to alternative RNA splicing.

Alternative RNA splicing is the process by which different proteins are produced from the same RNA transcript by splicing it in different ways.

Mature circulating B cells initially express IgM and IgD antibodies on their surfaces before they are stimulated to produce specific antibodies. The heavy chains in IgM and IgD are mu (μ) and delta (δ), respectively. The mu and delta genes for these heavy chains are located proximal to the VDJ encoding regions (i.e., variable, diversity, and joining regions) of the heavy chain locus. A primary RNA transcript of this entire region is produced, which then undergoes alternative RNA processing, enabling the production of both transmembrane IgM and IgD on B cells.

Other options:

Options A and C: Somatic hypermutation is a cellular response by which the immune system adapts to the new foreign elements that confront it. They do so by adding nucleotides in V gene

regions of heavy and light chains, resulting in a better fit of antigen with the antibodies. This hence increases the affinity of antibodies to antigens. Due to this, the subset of plasma cells with these improved hypervariable regions are more strongly selected by the antigen and therefore constitute a larger part of the population of antibody-producing cells. This process is called affinity maturation.

Option D: Allelic exclusion is the process by which a cell expresses one of the two alleles of the gene and excludes the other allele.

B cells are diploid, therefore, each B cell has six gene segment pools related to antibody chains:

• Two heavy-chain pools (one from each parent) and

• Four light-chain pools (one κ and one λ from each parent).

However, each B cell uses only two of the six gene segment pools: one out of the two heavy-chain pools (maternal and paternal) and one out of the four light-chain pools. Thus, each B cell must choose not only between its κ and λ light-chain pools but also between its maternal and paternal light-chain and heavy-chain pools. This process of choosing is by allelic exclusion.

Solution to Question 29:

The true statements regarding sirtuins are 1, 2, 3, and 4 (option B).

Sirtuins are a class of proteins that possess either mono-ADP-ribosyltransferase or deacetylase activity. They have been shown to slow down the aging process.

Sirtuins promote the expression of several genes whose products increase longevity. These include proteins that:

• Inhibit metabolic activity

• Reduce apoptosis

• Stimulate protein folding

• Inhibit the harmful effects of oxygen-free radicals

• Increase insulin sensitivity and glucose metabolism (maybe targeted for the treatment of diabetes)

• Ensure that damaged DNA is not propagated, preventing the accumulation of mutations

• Promotes replicative senescence under chronic stress

Sirtuins level increases by caloric restriction and are present in red wine.

Sirtuins are responsible for an increase in insulin sensitivity. Hence, they can play an important role in the treatment of diabetes.

Solution to Question 30:

The correctly matched set of the cells and their locations is 1-b, 2-a, 3-c, 4-d.

Kupffer cells are a component of the mononuclear phagocytic system and are located in the liver.

Parietal cells, also called oxyntic cells, are found in the upper half of the gastric glands. They have numerous mitochondria and are actively involved in the secretion of hydrochloric acid and intrinsic factor - a glycoprotein that aids in the absorption of vitamin B12 from the ileum.

Dust cells are macrophages in the walls of the alveoli. They phagocytose dust particles in the alveoli. In CHF these macrophages phagocytose erythrocytes that escape from the capillaries. The cells acquire a brick-red color and are called heart failure cells.

Clara cells are non-ciliated cuboidal epithelial cells that secrete important defense markers and serve as stem cells after injury. These cells make up a larger portion of the epithelial lining in the later portions of the conducting airway.

Solution to Question 31:

The cytokine not involved in TH1 response is IL-4.

Dendritic cells and macrophages present antigens to T helper cells (TH) after which they differentiate into TH1 and TH2 cells. They are responsible for cell-mediated and humoral immune responses, respectively. IL-12 is secreted by macrophages and promotes the proliferation of TH1 cells.

TH1 cells secrete:

• IL-2 which activates natural killer cells (NK cells) and cytotoxic T cells

• IFN gamma which stimulates macrophages and is also responsible for delayed hypersensitivity along with TGF beta.

TH2 cells secrete:

• IL- 4 stimulates IgE production from B cells

• IL- 5 stimulates proliferation of eosinophils and together play a role in helminthic and allergic reactions

• IL- 6 promotes proliferation of B cells and production of antibodies.

Solution to Question 32:

The clinical scenario and the image showing papillae lined by cells with overlapping nuclei and Orphan Annie eye nuclei indicate a diagnosis of papillary carcinoma of thyroid.

Papillary carcinoma of thyroid presents as a painless nodule or mass in the neck or cervical node in women of reproductive age. The risk factors include exposure to ionizing radiation and Hashimoto thyroiditis.

It usually presents as a cold nodule on a scan. Histopathological examination shows branching papillae and follicles with fibrovascular stalk. Nuclear changes include ground glass or Orphan Annie eye nuclei, pseudo-inclusions, and nuclear grooves. Psammoma bodies are often present.

Diagnosis can be made based on these nuclear features, even in the absence of papillary architecture. The prognosis is excellent, with a 10-year survival rate in excess of 95%.



The image below shows Orphan Annie eye nuclei with central clearing and margination of chromatin:



Other options:

Option A: Medullary carcinoma of the thyroid is a neuroendocrine tumor that is derived from parafollicular C cells, that secrete calcitonin.

Most of these tumors occur sporadically. Some are associated with familial multiple endocrine neoplasia (MEN) syndrome: MEN 2A and MEN 2B. They frequently present as a neck mass that may be associated with cervical lymphadenopathy. With the enlargement of the nodule, patients may have dysphagia and dyspnea.

Histopathology shows polygonal to spindle-shaped cells which form nests, trabeculae, or follicles. Amyloid deposits that are derived from the altered calcitonin molecules are present in the stroma.

Calcitonin in the tumor cells and stroma is easily demonstratable by immunohistochemistry.



Option B: Follicular carcinoma thyroid on histopathology may show uniform cells forming small follicles containing colloid (resembling normal thyroid gland) or sheets of cells without colloid.



Anaplastic carcinomas are poorly differentiated thyroid carcinomas with varied morphology including giant cells or spindle-shaped cells or both. There is minimal follicular differentiation with intratumoral necrosis.



Option C: A thyroglossal duct cyst is a congenital midline neck swelling that typically presents during childhood often noticed after an upper respiratory infection.

It represents the persistence of the track of the thyroid gland and can be found adjacent to the midline. It presents as smooth, fluctuant (cystic) swelling that is often transilluminant. Classically, the cyst moves upwards on swallowing and on tongue protrusion.

Sistrunk's operation is performed to excise the cyst and prevent a recurrence.



Solution to Question 33:

The clinical scenario of a lung tumor associated with chronic smoking and tumor cells staining positive for chromogranin indicates that the most likely diagnosis in this patient is small cell carcinoma of the lung.

Small cell carcinoma of the lung belongs to the neuroendocrine tumors and originates from neuroendocrine progenitor cells, Kulchitsky cells that line the bronchial epithelium. It is a highly malignant tumor with a high association with cigarette smoking. It usually arises from the major bronchi.

Histopathology of small cell lung carcinoma shows clusters of small cells which are round, oval, or spindle-shaped, and a prominent nuclear molding. These cells have scant cytoplasm, ill-defined cell borders, finely granular nuclear chromatin (salt and pepper pattern), and absent or inconspicuous nucleoli. The Azzopardi effect is described as basophilic staining of vascular walls due to coating by DNA from necrotic tumor cells.

Neuroendocrine markers seen in small cell lung carcinoma include:

• Chromogranin

• Synaptophysin

• CD56

Paraneoplastic syndromes associated with small cell carcinoma of the lung are Cushing syndrome due to ACTH production and hyponatremia due to inappropriate ADH secretion. Small cell lung cancer-associated SIADH is due to the ectopic release of antidiuretic hormone. It is associated with increased thirst and hyponatremia.

Other options:

Option A- Adenocarcinoma is a malignant epithelial tumor with differentiation in glands or mucin production by the tumor cells. These tumors are commonly found in women and in non-smokers. These tumors are more peripherally located and are smaller. On histopathology, the tumor shows a lepidic pattern of spread with tumor cells crawling along the normal-appearing alveolar septa. The tumor cells are positive for thyroid transcription factor 1 (TTF-1) and Napsin A.

Option B- Squamous cell carcinoma is found to be more common in men and has a strong association with smoking. They have precursor lesions like squamous dysplasia, squamous metaplasia, or in situ carcinomas. On histopathology, it is characterized by keratinization and/or intercellular bridges. Keratinization occurs in the form of squamous pearls. It is positive for p40 and p63 markers. Hypercalcemia is the common paraneoplastic syndrome associated with this type.

Option C- Large cell carcinoma is an undifferentiated malignant epithelial carcinoma. On histopathology, the cells are large, with large nuclei, prominent nucleoli with a moderate amount of cytoplasm.

Solution to Question 34:

The given clinical history and the histopathological findings showing keratin pearls are suggestive of squamous cell carcinoma of the lung.

Squamous cell carcinoma is most commonly found in men and is strongly associated with smoking.

Squamous cell carcinoma predominantly arises in the central segmental or subsegmental bronchi. The tumor may grow into the bronchial lumen and produce an intraluminal mass. In

certain cases, the bronchial obstruction may lead to atelectasis or infection. The tumor may penetrate the wall of the bronchus and infiltrate the peribronchial tissue into the carina or the mediastinum.

This tumor may be associated with paraneoplastic syndromes like hypercalcemia. The symptoms could include anorexia, nausea, vomiting, constipation, lethargy, polyuria, polydipsia, and dehydration.

Squamous metaplasia and dysplasia often precede frank squamous cell carcinoma. Carcinoma in-situ persists for many years and this may often be recognized on cytologic smears or in BAL fluid or brushings.

Gross appearance: The tissue is grey-white, firm, with areas of necrosis and hemorrhages. Red or yellow-white mottling and softening may be seen. The necrotic foci may cavitate.

Histopathology: They have precursor lesions like squamous dysplasia, squamous metaplasia, or in situ carcinomas. On histopathology, it is characterized by keratinization and/or intercellular bridges. Keratinization occurs in the form of squamous pearls or individual cells with dense eosinophilic cytoplasm. It is positive for p40 and p63 markers.



Solution to Question 35:

The true statement about proto-oncogenes is statement 3. ERBB2 amplification (gene that encodes HER2) is associated with breast cancer, while ERBB 1 mutation (gene that encodes EGFR) is associated with adenocarcinoma of the lung.

Other statements:

Statement 1: The most common genetic change in GISTs is the gain-of-function mutation of the tyrosine kinase KIT gene. These are present in 75% to 85% of all GISTs. Therefore KIT has increased expression in GIST.

The second most common mutation results in the activation of platelet-derived growth factor receptor A (PDGFRA).

Statement 2: KRAS mutation is associated with colon, lung, and pancreatic tumors. Among lung cancers, about 30% of cases of adenocarcinoma have gain of function mutations of the KRAS gene.

Small cell carcinomas are associated with the inactivation of TP53 and RB genes.

Statement 4: PDGFR-β amplification is associated with gliomas and leukemias. PDGF-β overexpression is associated with astrocytoma.

Mutations involved in colon cancer pathogenesis:

• Familial adenomatous polyposis usually has mutations involving the APC gene.

• HNPCC- majority of the cases have mutations in MSH2 or MLH1.

• Adenocarcinomas- the two pathways implicated are the APC pathway and MSI pathway.

Solution to Question 36:

The peripheral smear findings point toward a diagnosis of chronic lymphocytic leukemia.

The peripheral smear shows smudge cells, increased lymphocytes with scant cytoplasm, and round to slightly irregular nuclei containing clumped chromatin. This large number of monotonous, uniform-looking, mature lymphocytes is often described as a "convent-girl" appearance.

Chronic lymphocytic leukemia (CLL)/ Small lymphocytic lymphoma (SLL) is the most common leukemia among adults in the western world. The differentiating point between CLL and SLL is that the absolute lymphocyte count is >5000/mm³ in CLL and <5000/mm³ in SLL.

The chromosomal defects:

• Deletion of 13q (commonest and has a good prognosis)

• Deletion of 11q

• Trisomy 12q

• Deletion of 17p (poor prognosis)

It is usually asymptomatic at diagnosis but can present with easy fatigability, weight loss, and anorexia. Generalized lymphadenopathy and hepatosplenomegaly are usually seen.

Total blood count:

• Total leukocyte count is highly variable; leukopenia is seen when the marrow is involved.

• Hypogammaglobulinemia is common and contributes to increased susceptibility to infections.

• 10% to 15% of patients develop hemolytic anemia or thrombocytopenia due to autoantibodies made by non-neoplastic B cells.

Findings of peripheral smear:

• Predominantly shows small lymphocytes with scant cytoplasm.

• Contains a large number of monotonous, uniform-looking, mature lymphocytes, which is often described as a convent-girl appearance.

• Smudge cells or basket cells or parachute cells - artifacts of the lymphocytes which are disrupted during the process of making smears.

• Spherocytes (due to coexisting autoimmune hemolytic anemia) and nucleated RBCs (due to premature release of progenitors in face of severe anemia, marrow infiltration by the tumor, or both)



Lymph node biopsy: Diffuse effacement of lymph node architecture is seen. Small lymphocytes admixed with variable numbers of larger activated lymphocytes that often gather in loose aggregates. Proliferation centers containing mitotically active cells are pathognomonic for CLL / SLL.



The histopathological image shows small, darker staining lymphocytes with prominent nucleoli, which are typically seen in chronic lymphocytic leukemia (CLL).



AML is a hematopoietic progenitor malignancy that causes immature myeloblast accumulation in the bone marrow. Incidence increases with age and peaks after 60 years of age.

Clinically, the patients present with symptoms resulting due to pancytopenia (anemia, neutropenia, thrombocytopenia). There is generalized fatigue, bone pain, bleeding tendencies, and an increased risk of recurrent infections. For acute myeloblastic leukemia (AML) diagnosis, the blast cell count should be ≥20% in the peripheral blood or bone marrow.

Other distinctive features of AML:

• Bone marrow/ smear shows a large, uniform population of cells (myeloblasts) with prominent nucleoli (yellow arrow). It shows the presence of primary cytoplasmic granules.

• Auer rods: needle-shaped azurophilic intracytoplasmic inclusion bodies formed by the fusion of lysosomes. (black arrow)

• Multiple Auer rods clustered together within a single cell resembling a bundle of sticks known as faggot cells (seen in peripheral smear).



Acute lymphoblastic leukemia (ALL) arises from immature B (pre-B) or T (pre-T) cells called lymphoblasts. Immunohistochemistry for terminal deoxynucleotidyl transferase (TdT) is positive in pre-B and pre-T lymphoblasts.

Peripheral smear shows lymphoblasts that have condensed chromatin, less conspicuous nucleoli, and smaller amounts of cytoplasm which lacks granules as shown below.



Chronic myeloid leukemia is characterized by the presence of a Philadelphia chromosome resulting from reciprocal translocation t(9;22) forming the BCR-ABL fusion gene. They have hypercellular marrow due to a massively increased number of maturing granulocytic precursors.

The image below shows peripheral smear findings of CML with neutrophils, myelocytes, metamyelocytes, and band cells.



Solution to Question 37:

The true statements about familial adenomatous polyposis (FAP) are statements 2 and 4 (option-A).

Familial adenomatous polyposis is an inherited autosomal dominant disorder presenting numerous colorectal adenomatous polyps. It is an autosomal dominant disease caused by mutations in the APC gene, which is a regulator in the Wnt signaling pathway. The diagnosis of FAP requires the presence of at least 100 polyps.

Patients are usually teenagers and multiple polyps are visible in the colon during sigmoidoscopy. It is associated with a variety of extraintestinal manifestations including congenital hypertrophy of the retinal pigmental epithelium. Colorectal adenocarcinoma occurs in almost all cases before the age of 50.

Prophylactic colectomy is recommended in patients with APC gene mutations. The surgeries used are colectomy with ileorectal anastomosis (IRA), restorative proctocolectomy with an ileal pouch-anal anastomosis (RPC), or total proctectomy and end ileostomy. The first 2 surgeries are preferred in young patients as it avoids a permanent stoma.

Note - FAP along with brain tumors is called Turcot syndrome. FAP along with multiple osteomas, skin, and soft tissue tumors is called Gardner syndrome.

Solution to Question 38:

The maximum amplitude in thromboelastography represents platelets. A decrease in maximum amplitude signifies a quantitative or qualitative deficiency of platelets.

Thromboelastography (TEG) is a non-invasive test that quantifies the ability of whole blood to form a clot. It is useful in intraoperative coagulation status assessment. The dynamic changes in the viscoelastic properties of a blood sample during the clotting can be detected and quantified by thromboelastography. Underlying coagulation deficiencies show different shapes in the TEG. They are shown in the image below:





Advantages of thromboelastography:

- Blood coagulability, including coagulation cascade, platelet function, and fibrinolysis can be assessed.
- It is a simple bedside test
- Can diagnose coagulopathic bleeding
- Can be used as a guide in transfusion therapy and decrease the use of blood products
- Can detect dynamic changes in blood coagulation during resuscitation
- Predict the clinical efficacy of therapeutic agents affecting blood coagulability

Solution to Question 39:

The probable diagnosis from the given clinical scenario and the histopathological description is Hodgkin's lymphoma. The bi-nucleated cells are called the Reed-Sternberg cells which are CD 15+ and CD30+ (option D- 1 and 2).

Hodgkin's lymphoma arises in a single lymph node or chain of nodes and spreads first to anatomically contiguous lymphoid tissues. It usually presents as painless lymphadenopathy. The distinctive feature of Hodgkin's lymphoma is the presence of Reed-Sternberg cells.

WHO classifies Hodgkin's lymphoma into 5 subtypes:

• Nodular sclerosis: Frequent lacunar cells

• Mixed cellularity

• Lymphocyte-rich

• Lymphocyte depletion

• Nodular lymphocyte predominance

The first 4 subtypes are considered classical Hodgkin's lymphoma. The RS cells are CD 15+ and CD30+. The lymphocyte predominant type is considered non-classical Hodgkin's lymphoma with CD15-, CD30-, CD20+, and EB -.

The image below shows various cells like eosinophils, RS cells, plasma cells, and histiocytes in an FNA specimen of Hodgkin lymphoma:

Type	RS-cell	EBVirus	Prognosis
1. Nodular sclerosis (most common in the world)	Frequent lacunar cells and occasional diagnostic RS cells	Rare	Very good
2. Mixed cellularity (most common in India)	Frequent mononuclear and diagnostic RS cells	70%	Good
3. Lymphocyte Rich	Frequent mononuclear and diagnostic RS cells	40%	Very good
4. Lymphocyte depleted	Reticular variant: frequent diagnostic RS cells and variants and a paucity of background reactive cell	90%(Max)	Poorest
5. Lymphocyte predominant	L&H variant (Popcorn cell) variants in the background of follicular dendritic cells	0%(Min)	Excellent



Reed-Sternberg cell (RS cell) is the characteristic tumor giant cell of Hodgkin Lymphoma. It is derived from the germinal or post-germinal center B cell. It has a classic owl-eyed appearance due to the presence of a symmetric or mirror-image bilobed nucleus, with prominent nucleoli surrounded by a clear space. It has the following variants:

- Lacunar
- Classic
- L&H popcorn cells
- Pleomorphic
- Mononuclear

RS cells should be present in a background of non-neoplastic inflammatory cells like lymphocytes, plasma cells, and eosinophils, to make a diagnosis of Hodgkin's lymphoma.

The image given below shows the diagnostic Reed-Sternberg cell, with two nuclear lobes, large inclusion-like nucleoli, and abundant cytoplasm, surrounded by lymphocytes, macrophages, and an eosinophil.



The image below shows the Reed-Sternberg cell, mononuclear variant.

Mononuclear variant of RS cell



The image below shows Reed-Sternberg cell, lacunar variant. This variant has a folded or multilobated nucleus and lies within an open space, which is an artifact created by disruption of the cytoplasm during tissue sectioning.

Lacunar variant of RS cell



The image below shows the Reed-Sternberg cell, lymphohistiocytic variant. Several such variants with multiple infolded nuclear membranes, small nucleoli, fine chromatin, and abundant pale cytoplasm are present.



Solution to Question 40:

The given clinical scenario and the investigation findings are suggestive of microcytic hypochromic anemia. The most likely diagnosis in this patient is anemia of chronic disorder.

Anemia of chronic disorders more commonly presents as normochromic normocytic anemia that later develops into microcytic hypochromic anemia rarely. It is caused due to production of erythrocytes.

Anemia of chronic disease is anemia resulting due to chronic diseases which produce systemic inflammation. During chronic inflammation, inflammatory mediators like interleukin - 6 (IL-6) stimulate the production of hepcidin from the liver. Hepcidin is an acute phase reactant that inhibits ferroportin on macrophages and also decreases transport of iron from stores to developing RBC precursors. This results in decreased proliferation of erythroid progenitors due to impaired iron utilization.

Other options:

Option B: Thalassemia also presents as iron deficiency anemia but it does not have elevated luekocytosis.

Option C: Iron deficiency anemia also presents as microcytic hypochromic anemia. But since there is deficient iron, the serum ferritin level will also be low. This patient has normal serum ferritin and hence does not have iron deficiency anemia.

Note:

MCV/RBC count (Mentzer index) can also be used to distinguish thalassemia from iron-deficiency anemia (IDA). MCV (mean corpuscular volume) and MCH (mean corpuscular hemoglobin) are low in both IDA and thalassemia. RBC count may be normal in thalassemia, whereas it is reduced in IDA. A lower RBC count in IDA means the denominator is smaller, yielding a greater MCV/RBC ratio.

Option D: Sickle cell anemia is caused by a mutation in β-globin that creates sickle hemoglobin (HbS). It does not present as microcytic hypochromic anemia.

Solution to Question 41:

Senile atrophy is seen when there is a reduced blood supply to a tissue. This reduction in the blood supply occurs in a gradual manner because of slowly progressing atherosclerotic lesions.

Atrophy is a process of reduction in the size of a tissue or an organ resulting from a decrease in the cell size and number. It is a type of cellular adaptation. In this process, there is a decrease in the synthesis of cellular proteins and an increase in their degradation, mainly mediated by the ubiquitin-proteasome pathway. Atrophy can either be physiological or pathological:

Physiologic atrophy:

• Loss of hormonal stimulation: Atrophy of the uterus after parturition; atrophy of the endometrium, vaginal epithelium, and breast after menopause.

• During embryonic development: Certain structures, like the notochord and thyroglossal duct, undergo atrophy during fetal development.

Pathologic atrophy:

• Disuse Atrophy (decreased workload): e.g., atrophy in a limb that has been immobilized following a fracture

• Denervation atrophy (loss of nerve supply): e.g., skeletal muscle atrophy following nerve injury

• Ischemic atrophy: e.g., senile atrophy of the brain and heart due to atherosclerosis

• Inadequate nutrition: e.g., severe muscle wasting in marasmus

• Pressure atrophy: e.g., atrophy of surrounding uninvolved tissues caused by an enlarging benign tumor

Solution to Question 42:

Prostaglandins are produced from arachidonic acid present in the membrane phospholipids. This process is mediated by 2 cyclooxygenases: COX-1 and COX-2. Prostaglandins are produced by mast cells, macrophages, and endothelial cells.

COX-1 is commonly found in most tissues. It plays a role in maintaining normal homeostasis. It is also induced by inflammatory stimuli.

COX-2 is found in cells that are involved in inflammatory reactions.

The most important prostaglandins in inflammation are PGE_2, PGD_2, PGF_{2a}, PGI_2 (prostacyclin), and thromboxane A_2.

Other options:

Options A: Kreb's cycle is also called the citric acid cycle and tricarboxylic acid cycle. It is the terminal pathway for the oxidative catabolism of carbohydrates, amino acids, and fatty acids.

Option C: Leukotrienes are also produced from arachidonic acid with the help of lipoxygenases.

Solution to Question 43:

Metastatic calcification cannot be explained by vitamin D deficiency. Metastatic calcification is associated with hypercalcemia, which may be caused by vitamin D intoxication rather than deficiency.

Pathologic calcification is the abnormal tissue deposition of calcium salts, together with smaller amounts of iron, magnesium, and other mineral salts. It is of two types namely metastatic and dystrophic calcification.

Metastatic calcification is the deposition of calcium salts in otherwise normal tissues.

• Hypercalcemia secondary to some disturbance in calcium metabolism.

• Principal causes of hypercalcemia:

• Increased secretion of parathyroid hormone (PTH)

• Resorption of bone tissue

• Vitamin-D-related disorders- excess or abnormal sensitivity

• Renal failure

• Aluminum intoxication

• Milk-alkali syndrome- due to excess ingestion of calcium

Metastatic calcification can be seen in interstitial tissues of the gastric mucosa, kidneys, lungs, systemic arteries, and pulmonary veins.

Solution to Question 44:

The correct answer is option B:

Cytokines are proteins produced by many cell types (principally activated lymphocytes, macrophages, and dendritic cells, but also endothelial, epithelial, and connective tissue cells) that mediate and regulate immune and inflammatory reactions.

Three subsets of CD4+ T cells secrete different types of cytokines and elicit different types of inflammation.

TH1 cells - activate macrophages by the classical pathway by secreting IFNγ.

TH2 cells - recruit and activate eosinophils and are responsible for the alternative pathway of macrophage activation by secreting IL4, IL5, and IL13.

• IL4 acts on B cells to stimulate class switching to IgE and promotes the development of additional TH2 cells.

• IL5 is involved in the development and activation of eosinophils, which are essential effectors of type I hypersensitivity

(discussed later).

• IL13 enhances IgE production and acts on epithelial cells to stimulate mucus secretion. In addition, TH2 cells (as well as mast cells and epithelial cells) produce chemokines that attract more TH2 cells, as well as other leukocytes, to the reaction site. It also shares some functional similarities to IL-4 and can thus induce differentiation in dendritic cells.

TH17 cells secrete IL17 and other cytokines responsible for neutrophils and monocyte recruitment.

1. IL 4	C. B cells
2. IL 5	A. Eosinophils
3.IL9	B. Mast cells
4. IL 13	D. Dendritic cells

Solution to Question 45:

The histopathological image shows the presence of amyloid-rich stroma (pinkish material) and nests of polygonal cells, which are findings suggestive of medullary carcinoma of thyroid.



Medullary carcinoma of the thyroid (MTC) arises from the parafollicular C cells. Most of the MTC are sporadic, but they may also be associated with inheritable syndromes like familial MTC, MEN2A, and MEN2B. All the inherited variants are secondary to RET gene mutations.

Most patients present at 50–60 years and have a female preponderance. MTC secretes calcitonin, carcinoembryonic antigen (CEA), prostaglandins E2 and F2α, histaminase, serotonin, and adrenocorticotrophic hormone (ACTH).

Features of MTC

• Painful neck mass with cervical lymphadenopathy

• Dysphagia, dysphonia, or dyspnea if there is local invasion

• Diarrhea due to increased intestinal motility

• Cushingoid habitus due to ectopic ACTH production

• Blood-borne metastasis

• Microscopically, they consist of sheets of neoplastic cells separated by collagen and amyloid stroma

Diagnosis is by careful family history, physical examination, raised calcitonin and CEA levels, and fine-needle aspiration biopsy of the mass.

Other options:

Option B: The image below shows Orphan Annie-eye nuclei with central clearing and margination of chromatin in papillary carcinoma of the thyroid.



Option C: Follicular carcinoma thyroid on histopathology may show uniform cells forming small follicles containing colloid (resembling normal thyroid gland) or sheets of cells without colloid.



Option D: Anaplastic carcinomas are poorly differentiated thyroid carcinomas with varied morphology, including giant cells or spindle-shaped cells or both. There is minimal follicular differentiation with intratumoral necrosis.



Solution to Question 46:

Age less than 1 year, t(9,22), and t(4,11) are poor prognostic factors for acute lymphoblastic leukemia.

The given clinical scenario of B cell leukemia in a pediatric patient is acute lymphoblastic leukemia, since it is the most common cancer in children. The prognostic factors for a patient of ALL are as follows:

Poor prognostic factors	Good Prognostic factors
t(4;11)	t(12;21)
t(9;22)	t(1;19)
Hypodiploidy	Hyperdiploidy
11q23	t(10;14)
	Trisomies 4, 10, and 17
Age less than 2 years	Age between 2 and 10 yea rs
Presentation in adolescence o r adulthood	
Peripheral blood blast count >1 lakh	Low WBC count

Solution to Question 47:

The corrected reticulocyte count (CRC) of this patient is 3.

CRC = Reticulocyte % x (Actual Hb / Normal Hb)

According to the values given in this question:

CRC = 9 x 5/15 = 3

Alternatively, CRC can be calculated using hematocrit (PCV) as follows:

CRC = Reticulocyte % x (Actual PCV / Normal PCV)

Note: With these calculations, take normal Hb as 15g/dl and normal PCV as 45%.

Solution to Question 48:

The above clinical scenario along with the histopathological image showing Langerhans cells with folded nuclei (image A), CD1a (image B), and positive S100 (image C), is suggestive of Langerhans cell histiocytosis (LCH).

LCH occurs due to the proliferation of a special type of immature dendritic cells, called Langerhans cells. Their normal function is to present antigens. The most common mutation is the BRAFV600E gain-of-function mutation.

It is characterized by single or multiple osteolytic bone lesions and can also infiltrate nearly every organ, like the skin (of the scalp), lungs, liver, spleen, bone marrow, or the central nervous system.

LCH can present as the following clinical entities:

• Multifocal multisystem Langerhans cell histiocytosis (Letterer-Siwe disease) - mostly seen in infants

• Unifocal and multifocal unisystem Langerhans cell histiocytosis (eosinophilic granuloma) - The combination of bone defects, diabetes insipidus, and exophthalmos is called the Hand-Schüller-Christian triad - mostly seen in younger children

• Pulmonary LCH - mostly seen in adults

Clinical features include pain and swelling in the affected area. When calvarium is affected, it results in otitis media and mastoiditis. Spinal involvement results in compression fractures and neurological involvement. Vertebral involvement results in a rapid flattening of the vertebral body (vertebra plana).

Histopathologically, Langerhans cells have a single large nucleus with a characteristic folded or grooved nucleus (coffee-bean appearance) with small inconspicuous nucleoli. Birbeck granules are seen in the cytoplasm. The tumor cells specifically express HLA-DR, S-100, and CD1a and langerin (CD207) markers.

The image below shows a histopathological image of Langerhans cells with folded nuclei.



The presence of Birbeck granules in the cytoplasm is a characteristic finding of Langerhans cell histiocytosis (LCH) on electron microscopy.

Birbeck granules have a dilated terminal end producing a tennis racket–like appearance. It contains the protein langerin (CD207). It is seen on electron microscopy. The image below shows Birbeck granules on electron microscopy seen in LCH.



The below image shows lytic lesion of skull seen in LCH.



Option A: Rosai-Dorfman syndrome is otherwise called sinus histiocytosis with massive lymphadenopathy. Rosai-Dorfman disease is not a histiocytic neoplasm. It is an idiopathic disease characterized by Painless lymphadenopathy, usually of cervical nodes. The disorder predominantly affects children, adolescents, and young adults. It may present with nonspecific symptoms and management mostly involves observation rather than specific treatment.

Option B: Kimura disease is s a chronic inflammatory disorder of unknown etiology that most commonly presents as painless lymphadenopathy.

Option D: Juvenile xanthogranuloma is a non-Langerhans cell histiocytosis occurring in early infancy and childhood. It presents as reddish-brown papular lesions, which evolve to become orange. The lesions are most frequently seen in the face, scalp, and upper torso. They may be single or multiple lesions. Single lesions are the most common. The characteristic multinucleated giant cell, Touton cell is observed in 85% of juvenile xanthogranulomas.

Solution to Question 49:

The correct matches are as follows:

Popcorn cell is a lymphohistiocytic (L&H) variant of Reed Steenberg (RS) cells that are characteristic of lymphocyte-predominant Hodgkin lymphoma. It has a delicate multilobed, puffy nucleus resembling popped corn. The image below shows a popcorn cell seen in lymphocytic predominant Hodgkin's lymphoma.

1. Popcorn cell	b. Lymphocyte-predominant Hodgkin's lymphoma
2. Faggot cell	a. Acute promyelocytic leuke mia
3. Cerebriform nucle i	d. Sezary syndrome
4. Starry sky	c. Burkitt lymphoma



Multiple Auer rods clustered together within a single cell resembling a bundle of sticks are known as faggot cells and are highly suggestive of acute promyelocytic leukemia. Histopathology from bone marrow aspirate reveals numerous Auer rods (needle-like azurophilic granules) in the cytoplasm of the promyelocytes. Faggot cells are shown below.



Sézary-Lutzner cells - cells with cerebriform nuclei (infolded/hyper-convoluted nuclear membrane).

The histological hallmark of cutaneous T-cell lymphoma (mycosis fungoides type) is the presence of Sezary-Lutzner cells, which form band-like aggregates within the superficial dermis and invade the epidermis as single cells and small clusters called Pautrier microabscess (marked by arrow). These cells have markedly infolded nuclear membranes giving them a cerebriform contour as seen below.



Microscopy in Burkitt lymphoma shows a characteristic starry sky pattern. The neoplastic cells are homogeneous, medium-sized B cells with frequent mitotic figures and reactive macrophages scattered through the tumor, with their pale cytoplasm in a background of blue-staining tumor cells giving the tumor a so-called "starry sky" appearance.

The image given below shows a starry sky pattern seen in Burkitt lymphoma.



Solution to Question 50:

Hereditary angioedema is an autosomal dominant condition characterized by a deficiency of C1 esterase inhibitors. C1 esterase inhibitors usually target proteases like C1r and C1s, factor XII and the kallikrein system. Therefore, their deficiency leads to unregulated activation of kallikrein, leading to increased bradykinin.

Clinical features include swelling of the skin and mucosal surfaces, nausea, vomiting, and diarrhea, which may be aggravated after minor trauma or emotional stress. The condition can be treated by giving C1 inhibitor concentrates from human plasma.

The main inflammatory mediator responsible for these symptoms is bradykinin. Bradykinin is an inflammatory mediator formed by the cleavage of kininogen, in the presence of the enzyme kallikrein. Bradykinin has the following effects:

• Increases the vascular permeability

• Causes contraction of smooth muscle

• Dilates the blood vessel

• Mediates pain

Other options:

Options B and C: Decay-accelerating factor (DAF) and CD59 are two proteins that are linked by a glycophosphatidylinositol (GPI) anchor to the plasma membrane. DAF prevents the formation of C3 convertases and CD59 inhibits the formation of the MAC. An acquired deficiency of the enzyme that creates GPI anchors leads to a deficiency of these regulators and excessive complement activation and lysis of red cells resulting in Paroxysmal nocturnal hemoglobinuria(PNH).

During inflammation, C5a (Option D), C3a, and C4a, which are all cleavage products of their corresponding complement components, stimulate mast cells to release histamine. They have effects similar to those of mast cell mediators involved in anaphylactic reactions and therefore, are

called anaphylatoxins. C5a is also chemotactic for neutrophils, monocytes, eosinophils, and basophils. Additionally, it activates the lipooxygenase pathway of arachidonic acid metabolism in neutrophils and monocytes, resulting in the further release of inflammatory mediators.

Solution to Question 51:

The given scenario of a 35 year old woman, her iron profile and peripheral smear showing microcytic hypochromic anemia are suggestive of iron-deficiency anemia (IDA).

IDA is an example of microcytic hypochromic anemia. The causes of IDA can be inadequate diet, impaired reabsorption, increased requirement and increased blood loss.

The manifestations of IDA depend upon the underlying cause. In general, IDA presents with koilonychia, alopecia, atrophic changes in the tongue, gastric mucosa, malabsorbtion and pica (craving for consumption of non-edible materials)

Peripheral smears show small and pale red cells typically described as microcytic (reduced MCV) and hypochromic (enlarged zone of central pallor). Poikilocytosis can be seen in the form of elongated red cells, known as pencil cells. Target cells can be found in cases of severe IDA.

Diagnosis is mainly by laboratory studies. The typical findings are:

• Low hemoglobin and hematocrit

• Low serum iron and ferretin

• Increased Total Iron Binding Capacity (TIBC)

• Hepcidin is low (decreased stores of iron inhibit hepcidin synthesis)

IDA is treated by giving oral and parenteral preparations if iron or packed cell transfusion depending upon the severity of the anemia.

Option A: Megaloblastic anemia is defined by a high MCV

Option C: Thalassemia presents much earlier in infancy, with reduced MCV and a normal/increased RBC count

Option D: Essential thrombocytosis is characterized by isolated rise in platelet count (>450,000) along with normal hemoglobin, hematocrit and iron stores.

Solution to Question 52:

The true statements are that interleukin-12 (IL-12) is produced by macrophages and increases the production of interferon-γ (IFN-γ).

IL-12 is produced by dendritic cells and macrophages. Its principle actions are to promote the production of IFN-γ, which drives TH1-type responses and further stimulates macrophage activation. It also promotes the differentiation of CD4+ T-cells to Th1 cells.

Solution to Question 53:

The biopsy of adenocarcinoma of the lung will be positive for thyroid transcription factor-1 (TTF-1) and napsin A.

Adenocarcinoma is an invasive malignant epithelial tumor with glandular differentiation or mucin production by the tumor cells. It is associated with tobacco smoking, but less so than other histologic subtypes. It is the most common subtype in non-smokers and is associated with the loss of function of EGFR, ALK, ROS1, MET or RET genes.

The majority express TTF-1, which is a protein required for normal lung development. Napsin A is a new marker and is more sensitive and specific than TTF-1 in the diagnosis of adenocarcinoma.

Adenocarcinomas grow in various patterns: acinar, lepidic, papillary, micropapillary, and solid. The lesions are small and are usually located in the periphery. At the periphery of the tumor there is often a lepidic pattern of spread, in which the tumor cells “crawl” along normal-appearing alveolar septa.

Clinical features usually have an insidious onset. The incidence is usually at age 50 or more. The major presenting complaints are cough, weight loss, chest pain, and dyspnea.

Other options:

Option B: p40, along with p63 is a marker for squamous cell carcinoma of the lung.

Option D: Chromogranin is a marker for small cell carcinoma of the lung.

Solution to Question 54:

The instrument in the image is Sahli's hemoglobinometer and is used to measure acid hematin levels of a given blood sample.

In Sahli's method, oxyhemoglobin is converted to acid hematin with the action of dilute hydrochloric acid (N/10 HCl). The brown color of the acid hematin is matched to its intensity with a standard brown glass comparator in a visual colorimeter called Sahli’s colorimeter.

Sahli's method is a simple bedside test and is cost-effective. However, it has some disadvantages which include visual errors, inability to measure other forms of hemoglobin, fading of the comparator over time, and the color of acid hematin fading quickly.

Solution to Question 55:

Perl's Prussian blue stain is used in histopathological studies to detect the presence of iron in the specimen. Iron is stained blue with this stain.

It is used in hereditary hemochromatosis to stain liver tissue, which shows hepatocellular iron. It also stains the iron-laden mitochondria of ring sideroblasts seen in lead toxicity and myelodysplastic syndrome

The following image shows the blue staining of iron (in the form of hemosiderin) deposited in hepatocytes:



Other options:

Option A: Masson's trichrome stain is used to stain collagen.

Option C: PAS stain (Periodic acid Schiff stain) is used for the detection of glycogen in tissues such as the liver, skeletal muscle, etc., and also in tumor cells.

Option D: Fite-Faraco stain has been used in cases of leprosy for staining Mycobacterium leprae. It can also be used to stain Nocardia.

The image below shows acid-fast bacilli inside foamy macrophages:



Solution to Question 56:

The diagnosis based on the given clinical scenario and the histological image is molluscum contagiosum, which shows intracytoplasmic inclusion bodies, called Henderson–Patterson

bodies or molluscum bodies.

Infection is usually spread by direct contact, particularly among children and young adults. Multiple lesions occur on the skin and mucous membranes, with a predilection for the trunk and anogenital areas. Individual lesions are firm, pruritic, pink to skin-colored umbilicated papules. A curd-like material can be expressed from the central umbilication.

Smearing this material onto a glass slide followed by Giemsa staining shows diagnostic molluscum bodies, which are ellipsoid, homogeneous, cytoplasmic inclusion bodies that are found in the cells of the stratum granulosum and the stratum corneum and are up to 35 μm in size.

The image below shows umbilicated papules of molluscum contagiosum.



Other options:

Option A: Trichodysplasia spinulosa is a condition characterized by painless erythematous papules on the face with folliculocentric keratin spines. It is caused by trichodysplasia spinulosa-associated polyomavirus. Histologic features are keratin plugging, abundant dystrophic

and cornfield inner root sheath epithelium, and numerous, large eosinophilic trichohyalin granules.

Option B: Koilocytes are characteristically seen in HPV infections, such as condylomata acuminata. Koilocytes are lining cells which have perinuclear cytoplasmic vacuolization.

Option D: Donovan bodies are seen in donovanosis, caused by Klebsiella granulomatis. Donovan bodies are minute, encapsulated coccobacilli in macrophages, which can be demonstrated in Giemsa-stained smears of the exudate.

Solution to Question 57:

INF-γ is not involved in Jarisch-Herxheimer reaction (JHR).

JHR is a condition that occurs in patients infected by spirochetes who undergo antibiotic treatment. The reaction occurs within 24 hours of antibiotic treatment. Clinical features are fever, chills, nausea and vomiting, tachycardia, hypotension, flushing, myalgia, and exacerbation of skin lesions.

Syphilis is the leading spirochetal infection associated with JHR. Other infections are Lyme disease, louse-borne relapsing fever, leptospirosis, and tick-borne relapsing fever. Antibiotics that have been most associated are penicillins, tetracyclines, and erythromycin.

Mild reactions are self-limiting and often resolve spontaneously within 24 hours. Prevention of the JHR is by giving TNF-alpha antibodies and in some cases steroids.

Pathology NEET 2022

Question 1:

Choose the correct statement regarding the telomerase theory of aging.

a) Telomere stability is associated with aging

b) Abnormal telomerase activation is associated with aging

c) Decreased telomere length is associated with aging

d) Increased telomere length is associated with aging

Question 2:

A child presents with recurrent chest infections and abdominal pain. There is a history of 1 blood transfusion in the past. On examination, he had icterus and mild splenomegaly. Electrophoresis shows increased HbA2, HbF, and S spike. What is the likely diagnosis?

a) Beta thalassemia

b) HbC disease

c) Sickle cell disease

d) Acute coronary disease

Question 3:

Why do neoplastic cells utilize Warburg metabolism?

a) It decreases glucose utilisation by neoplastic cells

b) It forms metabolic intermediates which are needed for cell growth and multiplication

c) It provides more energy in the form of increased ATP production

d) It prevents apoptosis and makes the cancer cells immortal

Question 4:

An elderly patient presents with anemia and hemoglobinuria. Investigations reveal increased lactate dehydrogenase (LDH). The peripheral smear image is given below. Which of the following physical examination findings can support the likely diagnosis?



a) Splenomegaly

b) Frontal bossing

c) Mechanical second heart sound

d) Goitre

Question 5:

All of the following statements are true regarding neutrophil extracellular trapping (NET) except that

a) It is detected in blood during sepsis.

b) It is produced in response to bacterial infection.

c) Mitochondrial DNA is seen.

d) It is chromatin with antibacterial enzymes.

Question 6:

A patient underwent cystoscopy, which showed multiple yellow-white plaques in the trigone of the bladder. The histopathology image is given below. What is the diagnosis?



a) Interstitial cystitis

b) Malakoplakia

c) Polypoid cystitis

d) Acute cystitis

Question 7:

Which of the following measures are associated with an increased life span?

a) Moderate or regular exercise for 30 min

b) Decrease stress

c) Decreasing calorie intake by 30 percent

d) Pharmacological intervention with proton pump inhibitors

Question 8:

A 56-year-old man was diagnosed with COVID-19 and was put on mechanical ventilation. He passed away after a week. What is the likely post-mortem change seen in the lungs?

a) Thick layer of fibrin lining the alveoli

b) Acute and chronic alveolar hemorrhage

c) Perivascular cuffing

d) Pulmonary artery hypertrophy with increased resistance

Question 9:

A female patient presented with fatigue. Routine complete blood count analysis showed hemoglobin of 9 g/dL, MCV 60fL, and RBC count of 5.2 million. A peripheral smear showed microcytic hypochromic anemia. Which of the following is the next best investigation for this patient?

a) Serum ferritin levels

b) HbA2 levels

c) Serum folate levels

d) Serum homocysteine levels

Question 10:

A 56-year-old man presents with dragging pain in the abdomen. On examination, there is massive splenomegaly. Peripheral smear shows leukocytosis with increased myelocytes, metamyelocytes and basophils. Which of the following translocation is seen in this condition?

a) t(9;22)

b) t(8;22)

c) t(15;17)

d) t(8;14)

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	c
5	c
6	b
7	c
8	a
9	b
10	a

Detailed Explanations

Solution to Question 1:

A decreased telomere length is associated with aging.

Telomeres are short TG-rich repeated sequences of DNA present at the ends of linear eukaryotic chromosomes. Telomeres do not code for any proteins. They are important for ensuring the complete replication of chromosome ends and for protecting the ends from fusion and degradation.

Normally, with every cell division, there is some shortening of telomeres at the ends of chromosomes. Once the telomeres are shortened beyond a certain point, it may cause cell-cycle arrest or apoptosis. This is called replicative senescence.

Telomerase is the enzyme that prevents the shortening of telomeres. It is a specialized RNA-protein complex that uses its own RNA as a template for adding nucleotides to the ends of chromosomes.

This enzyme is usually present in germ cells but is absent in most somatic cells. Therefore most of the somatic cells age, their telomeres become shorter and they exit the cell cycle, resulting in an inability to generate new cells to replace damaged cells. Abnormal telomerase reactivation in somatic cells leads to evasion of senescence and apoptosis. It enables the cells to divide several times than usual.

In stem cells and cancer cells, telomerase is usually reactivated and telomere length is stabilized, allowing the cells to proliferate indefinitely.

Solution to Question 2:

The given scenario with electrophoresis showing increased HbA2, HbF, and S spike is suggestive of sickle cell disease.

Sickle cell anemia is an autosomal recessive disorder that occurs due to the point mutation in the β-globin gene which results in the replacement of glutamate residue with valine. This causes the production of sickle-shaped red blood cells due to the polymerization of deoxygenated hemoglobin (HbS).

Factors that predispose to sickling are infection, dehydration, and acidosis.

The clinical features of sickle cell disease are due to the tendency of HbS molecules to stack into polymers when deoxygenated. The features are:

• Moderately severe hemolytic anemia with jaundice.

• Painful bone crisis- hand-foot syndrome or dactylitis of the bones of hands and feet.

• Acute chest syndrome typically presents with fever, cough, chest pain, and pulmonary infiltrates.

• Hyposthenuria- inability to concentrate urine.

• Aplastic crisis stems from the infection of red cell progenitors by parvovirus B19, which causes a transient cessation of erythropoiesis and a sudden worsening of the anemia.

• Spleen

• In children- splenic sequestration crises occur which results in splenomegaly due to red pulp congestion.

• Adolescence or early adulthood- autosplenectomy occurs due to splenic infarction, fibrosis, and progressive shrinkage.

• Increased susceptibility to infection with encapsulated organisms like pneumococcal, meningococcal, and Haemophilus influenza septicemia, but can be reduced by vaccination and prophylactic antibiotics.

• Other clinical features- priapism, leg ulcers, stroke, retinopathy, pigment gallstones, and growth retardation.

The diagnosis is suggested by the clinical findings, the presence of irreversibly sickled red cells on peripheral smear, and hemoglobin electrophoresis showing increased levels of HbA2, HbF, and S spike. The changes in the skull resemble a crewcut in radiographic studies as a result of compensatory erythroid hyperplasia.

The image given below shows the sickling of RBCs in sickle cell disease.



The image given below shows Howell-Jolly bodies seen in asplenia.



Note: The sickle cell trait protects against falciparum malaria in certain African populations.

Other options:

Option A: Beta-thalassemia is an autosomal codominant disorder, caused by mutations in alpha or beta globin that reduce hemoglobin synthesis. Peripheral smear shows microcytic, hypochromic cells with anisocytosis, poikilocytosis, target cells, and basophilic stippling. Hemoglobin electrophoresis shows increased HbF and HbA2.

Option B: HbC disease is caused by mutation in the beta-globin chain in which glutamate is replaced by lysine in the sixth position of the beta-globin chain. Patients usually present with very mild anemia and microcytosis. Peripheral smear shows hexagonal HbC crystals. Hemoglobin electrophoresis shows increased HbC levels.

Option D: Acute coronary syndrome is very rare in children and may be associated with Kawasaki disease. Children with Kawasaki disease present with fever, conjunctival ingestion, strawberry tongue, polymorphous rash, cervical lymphadenopathy, and hand and foot changes. They are treated with intravenous immunoglobulin and aspirin.

Solution to Question 3:

Neoplastic cells utilize Warburg metabolism for the generation of metabolic intermediates, which are needed for cell growth and multiplication.

The Warburg Effect is a phenomenon in which there is an increase in the rate of glucose uptake and conversion into lactate(fermentation), even in the presence of oxygen. Cancer cells derive energy mainly from glycolysis.

Cancer cells utilize glycolysis (2 ATP) over mitochondrial oxidative phosphorylation (36 ATP) for glucose metabolism, even though the latter produces more ATP. It is because pure oxidative phosphorylation produces abundant ATP, but also burns glucose to CO_2 and H_2O, leaving no carbon moieties suitable for use as building blocks for lipids or proteins.

For this reason, rapidly growing cells, by undergoing glycolysis, produce carbon moieties that are needed for growth and cellular maintenance.

The image given below shows the comparison of glucose metabolism in normal cells versus cancer cells.



As a result of decreased ATP production, the main metabolic function of mitochondria in growing cells is not to generate ATP, but to carry

out reactions that generate cellular building blocks. For example, lipid biosynthesis.

Warburg effect is produced by growth factor receptors downstream signalling pathways. These include receptor tyrosine kinase/PI3K/AKT and MYC.

Clinically, cancer cells exhibiting the Warburg effect show a property of glucose hunger. This is used to visualize them via positron emission tomography (PET) scanning.

Solution to Question 4:

The above scenario with anemia, increased LDH, and hemoglobinuria along with peripheral smear showing schistocytes are suggestive of macroangiopathic hemolytic anemia (MAHA). The presence of a mechanical second heart sound can support the diagnosis.

Macroangiopathic hemolytic anemia is a type of hemolytic anemia that results from trauma to red cells by artificial mechanical cardiac valves that produce turbulent blood flow and pressure gradients that cause hemolysis from shear forces. This traumatic damage leads to intravascular hemolysis and the appearance of red cell fragments (schistocytes), burr cells, helmet cells, and triangle cells in blood smears.

The given below peripheral smear shows schistocytes and helmet cells.



Laboratory findings in hemolytic anemia resulting from trauma to red cells:

• Hemoglobin- decreased

• Reticulocytes- increased

• Platelets- normal

• Unconjugated bilirubin- increased

• Lactate dehydrogenase- increased

• Haptoglobin- decreased

• Hemoglobinuria due to intravascular hemolysis

Treatment is conservative. But, if more than mild anemia develops, reintervention to correct regurgitation may be required.

Option A: Splenomegaly is seen in various disorders of white and red cells. But splenomegaly in intravascular hemolysis is less likely.

Option B: Frontal bossing is seen in thalassemia due to expansion of hematopoietically active marrow. An X-ray of the skull shows a 'crewcut' appearance. Peripheral smear in thalassemia shows microcytic hypochromic cells with anisocytosis, poikilocytosis, and target cells.

Option D: Goiter is usually associated with iron deficiency anemia. A peripheral smear shows microcytic hypochromic cells.

Solution to Question 5:

Mitochondrial DNA is not seen in neutrophil extracellular traps (NET).

Neutrophilic extracellular traps are networks of extracellular fibers that provide a high concentration of antimicrobial substances at the site of infection and prevent the spread of the microbes by trapping them in the fibrils.

Neutrophilic extracellular traps (NET) are the meshwork of nuclear chromatin of neutrophils. The loss of nuclei leads to the death of the neutrophils. They concentrate antimicrobial peptides and enzymes to fight against bacteria and fungus (not viruses). They are detected in the blood during sepsis. They are produced by neutrophils in response to:

• Infectious pathogens: mainly bacteria and fungi

• Inflammatory mediators: chemokines, cytokines (mainly interferons), complement proteins, and reactive oxygen species (ROS)

Note: The nuclear chromatin in the NETs, which includes histones and associated DNA, has been postulated to be a source of nuclear antigens in systemic autoimmune diseases, particularly systemic lupus erythematosus (SLE).

Solution to Question 6:

The cystoscopy findings of multiple yellow-white plaques in the trigone of the bladder and histopathology image showing Michaelis–Gutmann bodies are consistent with the diagnosis of malakoplakia.

Malakoplakia is a rare inflammatory reaction seen in chronic cystitis. It is caused due to acquired defects in phagocyte function, chronic bacterial infection by Escherichia coli or Proteus species, and immunosuppression (renal transplant patients).

The term malakoplakia means soft plaques. Grossly, malakoplakia appears as soft, yellow, slightly raised mucosal plaques.

Microscopically, they are composed of large, foamy macrophages mixed with occasional multinucleate giant cells and lymphocytes. The macrophages in malakoplakia have abundant granular cytoplasm due to phagosomes stuffed with undigested bacterial products and intracellular deposition of calcium and iron within enlarged lysosomes form laminated basophilic concretions. These are known as Michaelis–Gutmann bodies, pathognomonic of malakoplakia (shown below).



The image given below shows the Von Kossa stain demonstrating calcium and iron-containing Michaelis-Gutman bodies (MG bodies) in malakoplakia.



Note: Xanthogranulomatous cystitis is morphologically and probably pathogenetically similar to malakoplakia, but it lacks Michaelis–Gutmann bodies.

Other options:

Option A: Interstitial cystitis, also known as chronic pelvic pain syndrome, is a disorder of the urinary bladder that presents with urinary tract symptoms of more than six weeks duration of unknown etiology. Grossly, it is associated with chronic mucosal ulcers called Hunner's ulcers. Microscopically, there is a non-specific increase in mast cells in the submucosa. Biopsy should be done in all cases to rule out carcinoma in situ.

The cystoscopy image shows Hunner's ulcer, seen in interstitial cystitis.

Option C: Polypoid cystitis is an inflammatory lesion resulting from irritation of the bladder mucosa, most commonly due to indwelling catheters. The urothelium is thrown into broad

bulbous polypoid projections due to marked submucosal edema. They must be differentiated from papillary urothelial carcinoma.

The given image below shows polypoid cystitis showing broad polypoid growth, prominent blood vessels, and metaplastic urothelium in H&E stain.



Option D: Acute cystitis is characterized by an inflammation of the urinary bladder following infection, most commonly by Escherichia coli, Proteus, Klebsiella, and Enterobacter. It is more commonly seen in women due to the short urethra. Microscopically, there is hyperemia and neutrophilic infiltration of the mucosa.

Solution to Question 7:

Decreasing calorie intake by 30 percent will increase lifespan.

Caloric restriction reduces both the degree of oxidative damage and the associated decline in function. It decreases the steady-state concentrations of products of oxidative damage to proteins, DNA, and lipids. This is due to decreased production of mitochondrial superoxide anion and hydrogen peroxide and increased antioxidant production. Overall, caloric restriction depresses the rate of energy metabolism.

Caloric restriction increases longevity both by reducing the signaling intensity of the insulin-like growth factor (IGF-1) pathway and by increasing sirtuins (NAD-dependent protein deacetylases).

The attenuation of IGF-1 signaling leads to lower rates of cell growth and metabolism that reduces cellular damage. A similar effect is produced by rapamycin. An increase in sirtuins (particularly sirtuin-6) promotes the expression of several genes whose products increase longevity by activating DNA repair enzymes through deacylation.

Caloric restriction delays the onset of age-associated diseases such as cancer (including lymphomas and breast and prostate cancers), nephropathy, and cataracts. The onset of diabetes, hypertension, hyperlipidemia, and the development of autoimmune diseases is also decreased by caloric restriction.

Solution to Question 8:

The clinical scenario of a patient with COVID-19 pneumonia on mechanical ventilation is suggestive of acute respiratory distress syndrome (ARDS). A thick layer of fibrin or waxy hyaline membranes lining the alveolar walls are characteristic of ARDS.

COVID-19 is a respiratory disease caused by SARS-CoV-2. These are enveloped positive-sense RNA viruses. The spread of SARS-CoV-2 is believed to be primarily via respiratory droplets, direct contact, airborne, and fomites.

The major complication in patients with severe COVID-19 is ARDS requiring oxygen therapy and mechanical ventilation. ARDS is a clinical syndrome characterized by acute onset, severe dyspnea, tachypnea, hypoxemia, and pulmonary infiltrates leading to respiratory failure.

Etiological factors can be divided into direct and indirect causes of lung injury as follows:

Damage to the endothelial and alveolar epithelial cells, with inflammation, is the key initiating event and the basis of lung damage. Neutrophils play a significant role in tissue injury in this condition.

There are three stages in the development of ARDS:

• Exudative phase- occurs for the first 7 days of the illness. This is associated with damage to alveolar-capillary endothelial cells and type I pneumocytes. This results in a breach in the tight alveolar barrier. The release of proinflammatory cytokines leads to the recruitment of neutrophils and the development of a hyaline membrane in the alveoli.

• Proliferative phase- occurs from day 7-21. Recovery from mechanical ventilation is seen in most patients.

• Fibrotic phase- is associated with increased mortality. Extensive alveolar duct and interstitial fibrosis are seen.

The image given below shows hyaline membrane formation in the alveolar walls seen in ARDS.

Direct lung injury	Indirect lung injury
Pneumonia	Sepsis
Aspiration of gastric contents	Transfusions
Near drowning	Drug overdose
Toxic inhalational injury	Multiple transfusions
Pulmonary contusion	Pancreatitis
COVID-19	Severe trauma likeburns, flail chest, head trauma, multiple bone fractures



The chest X-ray features in ARDS usually develop 12-24 hours after initial lung insult as a result of proteinaceous interstitial edema. Within one week, a hyaline membrane occurs due to type 1 pneumocyte damage. There are diffuse bilateral coalescent opacities consistent with pulmonary edema.

The image given below shows chest x-ray findings seen in ARDS.



Berlin's definition of ARDS predicts increased mortality. It considers the following parameters:

• Chest X-ray showing bilateral opacities

• Respiratory failure within 1 week of alveolar insult

• Decreased PaO2/FiO2 ratio to less than 300

• Symptoms of respiratory failure are not due to heart failure/fluid overload

Diagnosis of COVID-19 is done by nucleic acid amplification testing of respiratory tract secretions by nasopharyngeal swab. It is associated with elevated inflammatory markers, such as interleukin

6 (IL-6), tumor necrosis factor, ferritin, C-reactive protein, lactate dehydrogenase, and D-dimer.

Chest CT may be more sensitive than radiographs, but CT should be used principally for medical management of respiratory disease, not as a primary diagnostic tool for COVID-19.

The image given below shows the management of COVID-19 infection.



Solution to Question 9:

The next best investigation for this patient would be determining the HbA2 levels, as the history of reduced hemoglobin, reduced MCV (microcytic anemia) and normal to high RBC count is suggestive of thalassemia.

The MCV count divided by the RBC count is called the Mentzer index, which is lower than 13 in thalassemia due to the normal to high RBC count in the denominator. In iron deficiency anemia, it is higher than 13 as the MCV and the RBC count both are reduced.

Alpha- and beta-thalassemia are inherited disorders in which there is reduced production of alpha-globin or beta-globin chains, respectively. This leads to precipitation of the excess (unpaired) chains, interfering with red blood cell (RBC) production and survival.

Clinical features in thalassemia include anemia, jaundice, skeletal deformities including facial deformities, changes in body habitus, growth impairment, and hepatosplenomegaly. Due to frequent blood transfusions, iron overload is also seen.

Peripheral smear is likely to show microcytic hypochromic anemia with target cells, teardrop cells, and some coarse basophilic stippling in certain cases.

The image given below shows peripheral smear findings in thalassemia.



An X-ray of the skull shows a crewcut appearance due to expansion of hematopoetically active marrow.

Thalassemia is usually managed by blood transfusions every 2-4 weeks and chelation therapy with deferoxamine, deferasirox, and deferiprone for iron chelation. Splenectomy is indicated when the transfusion requirement to maintain ideal hemoglobin increases. Hematopoietic stem cell transplantation (HSCT) is the only known curative treatment for thalassemia.

Other options:

Option A: Serum ferritin level is used to diagnose iron-deficiency anemia (IDA). They are usually low in IDA.

Option C: Serum folate level is used to diagnose macrocytic anemia due to a deficiency of folic acid.

Option D: Serum homocysteine levels are increased in vitamin B12–deficient macrocytic anemia.

Solution to Question 10:

The clinical scenario of massive splenomegaly and peripheral smear shows leukocytosis with increased myelocytes, metamyelocytes and basophils are suggestive of chronic myeloid leukemia (CML). A translocation between t(9;22) is seen in this condition.

CML is a myeloproliferative neoplasm. It is caused due to BCR (on chromosome 22) - ABL (on chromosome 9) fusion gene translocation mutation. It results in the formation of the Philadelphia chromosome which is of CML.

The image given below shows Philadelphia chromosome showing reciprocal (9;22)(q34;q11) translocation seen in CML.



The resulting BCR-ABL translocation results in the activation of tyrosine kinase that signals pro-growth and pro-survival pathways like RAS and JAK/STAT pathways resulting in the proliferation of hematopoietic stem cells.

Patients typically present with a dragging sensation abdomen due to splenomegaly or splenic infarct, fatigue, malaise, and weight loss.

On peripheral smear there is leukocytosis with leukocyte count, often exceeding 100,000 cells/mm3 (neutrophils, band forms, metamyelocytes, myelocytes, eosinophils, and basophils), blasts are usually less than 10% of the circulating cells, the platelet count is usually increased.

The image below shows peripheral smear findings of chronic myeloid leukemia (CML) with neutrophils, myelocytes, metamyelocytes, and band cells.



Bone marrow is hypercellular due to increased numbers of maturing granulocytic precursors with sea-blue histiocytes (macrophages with wrinkled, green-blue cytoplasm).

Fluorescence in situ hybridization (FISH) or polymerase chain reaction(PCR) to detect the BCR-ABL gene. Leucocyte alkaline phosphatase (LAP) score is decreased in cases of CML.

Sokal scoring is used in assessing the prognosis of this patient. The prognosis of CML can be assessed based on:

• Age

• Spleen size

• Platelet count

• Percentage of blast cells in the blood

FDA-approved tyrosine kinase inhibitors for the treatment of CML are imatinib, nilotinib, dasatinib, bosutinib, and ponatinib.

Complications, if left untreated:

• Accelerated phase- increased anemia, thrombocytopenia, and basophilia

• A blast crisis- may transform into acute leukemia (AML/ALL)

Other options:

Options B and D: t(8;14) and t(8;22) are seen in Burkitt lymphoma. It is associated with increased MYC levels. All endemic cases of Burkitt lymphoma are caused by the Ebstein-Barr virus. Microscopy shows a characteristic starry sky pattern.

The image given below shows a starry sky pattern seen in Burkitt lymphoma.



Option C: t(15;17) is seen in acute promyelocytic leukemia. It is characterized by promyelocytes with Auer rods on bone marrow aspirate. All-trans-retinoic acid is used for the treatment.

The image given below shows bone marrow findings in acute promyelocytic leukemia.



Pathology NEET 2023

Question 1:

Which of the following is true regarding Mendelian mode of inheritance?

a) Every 10th patient has a Y-linked disorder

b) 1/3rd of Mendelian disorders is autosomal recessive

c) Every 5th patient has an X-linked recessive disorder

d) 2/3rds of Mendelian disorders are autosomal dominant

Question 2:

An elderly male who is a known smoker presented with chronic cough, significant weight loss, and fatigue. Serum calcium level is raised. A lung biopsy was done, and it showed large atypical cells with hyperchromasia. What is the probable diagnosis?

a) Large cell neuroendocrine tumor

b) Small cell carcinoma

c) Adenocarcinoma

d) Squamous cell carcinoma

Question 3:

A young patient presents with a large retroperitoneal hemorrhage. He also gives a frequent history of intermittent swelling of knees on strenuous exercise. There is no history of any mucosal bleed. Which among the following clotting factors is responsible for the above-mentioned condition?

a) Factors VIII and IX

b) Factors XI and XII

c) Von Willebrand factor

d) Lupus anti-coagulant

Question 4:

An elderly male went for a medical examination. Laboratory investigations showed a marked elevation of lymphocytes with marked leukocytosis. The peripheral smear of the patient is given below. What is the appropriate next step for this patient?



a) Stem cell transplantation

b) Bone marrow biopsy

c) Flow cytometry

d) Start high-dose chemotherapy

Question 5:

A 25-year-old male presented with low hemoglobin, platelet count of 25,000/mL, gum bleeding, and raised PT & aPTT. The image of his peripheral smear is given below. What is the likely diagnosis?



a) AML with t(8;21)

b) AML with inv(16)

c) AML with t(15;17)

d) AML with t(11;22)

Question 6:

A chronic smoker who worked in a cement factory for the past 20 years developed mesothelioma. The association is likely to be due to _____.

a) Silicosis

b) Asbestosis

c) Coal workers' pneumoconiosis

d) Bagassosis

Question 7:

A child presents with intermittent jaundice and splenomegaly. There is a history of similar complaints in the elder brother. Peripheral smear shows the following finding. How will you investigate this condition?



a) Osmotic fragility test - Hereditary spherocytosis

b) Coombs test - AIHA

c) G6PD deficiency - Genetic testing

d) Paroxysmal nocturnal hemoglobinuria - Flow cytometry

Question 8:

An African child is found to have a mass in the jaw. A biopsy of the mass was done, and it appears as shown in the image below. The child also had a translocation of t(8;14). What is the probable diagnosis?



a) Acute myeloid leukemia

b) Follicular lymphoma

c) Multiple myeloma

d) Burkitt lymphoma

Question 9:

A child presented to the hospital with cola-colored urine, hypertension, and puffiness of eyes. Laboratory investigations were done and creatinine was 2.5 mg/dL. Treatment was started, and despite treatment, the patient did not improve for the next 3 weeks. The creatinine value increased to 4.5 mg/dL. Which among the following electron microscopic findings will be seen in this patient?

a) Subendothelial deposits

b) Subepithelial deposits

c) Crescent formation

d) Mesangial deposits

Question 10:

A patient was transfused 4 units of blood 2 hours ago, and now she complains of chest discomfort. X-ray shows bilateral fissural thickening. Which of the following is the most probable cause?

a) ABO incompatibility reactions

b) Fluid overload

c) Hypocalcemia

d) Graft-versus-host disease

Question 11:

A 48-year-old man presents with complaints of facial puffiness, frothy urine, and hypertension. He gives a history of infection with hepatitis B. Urine examination reveals microscopic hematuria. The histopathological image of the kidney biopsy shows a spike and dome pattern. What is the diagnosis of this condition?



a) Minimal change disease

b) Membranous nephropathy

c) FSGS

d) PSGN-associated disease

Question 12:

A 5-year-old child is brought with fever, fatigue, and left ventricular dysfunction. An endomyocardial biopsy is shown below. What is the probable diagnosis?



a) Acute rheumatic fever

b) Chagas disease

c) Pyogenic myocarditis

d) Lymphocytic myocarditis

Answer Key

Question No.	Correct Option
1	d
2	d
3	a
4	c
5	c
6	b
7	a
8	d
9	b
10	b
11	b
12	d

Detailed Explanations

Solution to Question 1:

2/3rds of Mendelian disorders, which corresponds to approximately 66% disorders, are autosomal dominant is the true statement about the Mendelian mode of inheritance. Autosomal dominant patterns of inheritance comprise 65% of all Mendelian disorders (monogenic human diseases).

Mendelian disorders (classic inheritance)

• Autosomal dominant

• Autosomal recessive

• X-linked dominant

• X-linked recessive

Of all the Mendelian disorders, also known as monogenic human disease, about 65% are autosomal dominant, 25% are autosomal recessive, and 5% are X-linked.

Other options:

Option A: Every 10th patient has a Y-linked disorder - This statement is incorrect. Y-linked disorders are extremely rare, and their prevalence is much lower than 1/10 (10%).

Option B: 1/3rd of Mendelian disorders is autosomal recessive - This statement is incorrect. Approximately 25% of Mendelian disorders are autosomal recessive, not 1/3rd (33.33%).

Option C: Every 5th patient has an X-linked recessive disorder - This statement is incorrect. Approximately 5% of Mendelian disorders are X-linked, not 1/5 (20%).

Solution to Question 2:

The most probable diagnosis based on the given clinical scenario is squamous cell carcinoma of the lung.

Squamous cell carcinoma is most commonly found in men and is strongly associated with smoking. It predominantly arises in the central or hilar region. Histologically, it is characterized by the presence of keratin pearls and intercellular bridges.

Hypercalcemia is one of the most common paraneoplastic syndromes associated with squamous cell cancer (SCC) of the lung. Increased levels of parathyroid hormone-related protein (PTH-rP) are the most likely mechanism of his hypercalcemia. The symptoms could include anorexia, nausea, vomiting, constipation, lethargy, polyuria, polydipsia, and dehydration.

The image given below shows keratin pearls seen in squamous cell carcinomas.



Other options:

Option A: Large cell neuroendocrine tumor - These tumors are a subtype of large cell carcinoma and show neuroendocrine features on immunohistochemistry. They are aggressive and can be associated with paraneoplastic syndromes, but they do not commonly cause hypercalcemia.

Option B: Small cell carcinoma - This is a highly aggressive neuroendocrine tumor strongly associated with smoking. It frequently presents with paraneoplastic syndromes, including SIADH

and ectopic ACTH production, but hypercalcemia is less common. Histologically, it is characterized by small cells with scant cytoplasm, ill-defined cell borders, and finely granular nuclear chromatin.

Option C: Adenocarcinoma - This is the most common type of lung cancer and is more common in non-smokers and women. It usually arises in the peripheral lung tissue and is associated with glandular differentiation or mucin production. Histologically, adenocarcinoma is characterized by glandular or acinar structures and/or intracytoplasmic mucin.

Solution to Question 3:

The diagnosis based on the given clinical scenario hemophilia A. The clotting factors responsible for the above-mentioned condition are factors VIII and IX.

Hemophilia A is an X-linked recessive disorder. It occurs due to a deficiency of clotting factor VIII, which causes inadequate thrombin generation and results in the inability to form a tight fibrin clot to support the platelet plug.

Symptoms include easy bruising, intramuscular hematomas, hemarthroses, hematuria, and bleeding from minor traumatic lacerations of the mouth can persist for hours or days. Retroperitoneal hemorrhages can accumulate large volumes of blood and may cause a hypovolemic shock.

Laboratory findings in hemophilia A:

• Factor VIII - reduced

• Partial thromboplastin time (PTT) - prolonged

• Platelet count - normal

• Bleeding time - normal

• Prothrombin time - normal

Mild cases of hemophilia are treated by the administration of desmopressin acetate. Moderate to severe cases need the replacement of deficient clotting factors (factor VIII in hemophilia A). Monoclonal antibody emicizumab can restore activated factor VIII activity in patients with hemophilia.

Other options:

Option B: Factors XI and XII - Deficiencies of these factors are not associated with the clinical presentation described in the question. Factor XI deficiency (hemophilia C) can cause mild bleeding disorders, but it is generally not associated with severe hemorrhages or joint swelling. Factor XII deficiency is typically asymptomatic and not associated with bleeding disorders.

Option C: Von Willebrand factor - A deficiency in von Willebrand factor results in von Willebrand disease, which can cause excessive bleeding similar to hemophilia. However, von Willebrand disease usually presents with mucosal bleeding (e.g., epistaxis, menorrhagia, and gum bleeding), which is not described in this case.

Option D: Lupus anticoagulant - This is an acquired condition seen in antiphospholipid antibody syndrome (APS), which can cause thrombotic events rather than bleeding. It is not associated with

retroperitoneal hemorrhage or joint swelling upon strenuous exercise.

Solution to Question 4:

The likely diagnosis based on the given clinical scenario and the peripheral smear showing smudge cells is chronic lymphocytic leukemia. The next step in his management should be flow cytometry.

Chronic lymphocytic leukemia (CLL) is a peripheral B-cell neoplasm usually affecting adults, more commonly men. Patients are often asymptomatic at diagnosis. When symptomatic, it presents non-specifically with easy fatigability, weight loss, and anorexia. Generalized lymphadenopathy and hepatosplenomegaly are present in most symptomatic patients.

The chromosomal defects in CLL include

• Deletion of 13q (commonest and has a good prognosis)

• Deletion of 11q

• Trisomy 12q

• Deletion of 17p (poor prognosis)

The peripheral blood smear in cases of chronic lymphocytic leukemia (CLL) contains small round lymphocytes having condensed chromatin and scant cytoplasm. Some of these cells may get disrupted while preparing the smears, producing smudge cells. An absolute lymphocyte count of >5000/mm3 is required for diagnosis. Hypogammaglobulinemia is commonly seen, making the patient susceptible to infections. A few patients may develop hemolytic anemia or thrombocytopenia due to autoantibodies made by non-neoplastic B cells.

Immunophenotyping using flow cytometry is the investigation done to arrive at the diagnosis. The classical immunophenotypic feature of CLL is CD19+, CD20+, CD 23+, and CD5+.

On lymph node biopsy: Lymph nodes are composed of small lymphocytes admixed with larger activated lymphocytes. The large lymphocytes contain mitotically active cells and gather in loose aggregates, which are referred to as proliferation centers. These proliferation centers are pathognomic to CLL when present.

Expectant management is the treatment of choice for asymptomatic patients with chronic lymphocytic leukemia. The gold standard for therapy is a combination of fludarabine, cyclophosphamide, and rituximab.

Richter's syndrome is a complication of CLL in which there is a transformation to an aggressive lymphoma, most commonly diffuse large B-cell lymphoma (DLBCL).

Other options:

Option A: Stem cell transplantation - This is not the first step in the management of CLL. It may be considered in certain high-risk patients or those with refractory disease, but only after confirming the diagnosis and assessing the disease stage.

Option B: Bone marrow biopsy - While it can provide valuable information about the extent of infiltration, it is not the initial test for diagnosing CLL. Flow cytometry is the preferred diagnostic method.

Option D: Start high-dose chemotherapy - This is not the first step in managing CLL. Treatment decisions should be based on clinical stage, prognostic factors, and the patient's overall health. Moreover, asymptomatic patients with CLL are often managed with a watchful waiting approach.

Solution to Question 5:

The peripheral blood smear shows numerous Auer rods in the cytoplasm of progranulocytes, which are characteristic of acute myeloid leukemia (AML). The presentation of the patient with gum bleeding points to the diagnosis of AML with t(15;17).

Auer rods (needle-like azurophilic granules) are predominantly seen in acute promyelocytic leukemia, which is a subtype of acute myeloid leukemia. It is associated with t(15;17), which results in the formation of the fusion gene called PML-RARA. The condition is associated with a high incidence of disseminated intravascular coagulation (DIC). Treatment includes the administration of all-trans retinoic acid.

Solution to Question 6:

The association in this case is between occupational exposure to cement for 20 years and the occurrence of mesothelioma, which is mediated by the development of asbestosis as a result of prolonged inhalation of asbestos.

Asbestosis occurs due to the inhalation of crystalline hydrated silicates (asbestos). It occurs in two distinct geometric forms, serpentine and amphibole. It is characterized by diffuse parenchymal interstitial fibrosis. It usually begins in the lower lobes of the lungs and may present with dyspnea and cough, usually after 20-30 years of exposure. Other asbestos-related diseases include pleural plaques, recurrent pleural effusions, lung carcinoma, and mesothelioma.

Asbestos bodies are a hallmark finding. They appear as golden brown, fusiform, or beaded rods. These are formed when the asbestos fibers are phagocytosed by the macrophages and are coated by iron-containing proteinaceous material derived from phagocyte ferritin.

The image below shows asbestos bodies.



Other options:

Option A: Silicosis is due to the inhalation of crystalline silicon dioxide (silica) as seen in individuals employed in sandblasting or rock mining. Macrophages phagocytose inhaled silica particles, thereby activating and releasing inflammatory mediators, like IL-1 and IL-8.

Histopathology reveals coalescent nodules of whorled collagen fibers surrounded by a peripheral zone of dust-laden macrophages. Radiologically, egg-shell calcification of the lymph nodes is seen.

Shown below is an image of a silicotic nodule with collagen whorls as seen in this condition.



Option C: Anthracosis/coal workers' pneumoconiosis is a pulmonary condition seen in coal miners. The inhaled carbon pigment is engulfed by alveolar or interstitial macrophages, which accumulate in the connective tissue. On prolonged exposure, lung tissue undergoes progressive massive fibrosis, which is called complicated coal workers' pneumoconiosis. The upper lobes of the lungs are commonly involved.

Microscopic features consist of coal-laden macrophages, black pigment, and dense collagenous interstitial fibrosis.

The image below shows anthracosis—black pigment engulfed within macrophages as well as in the form of deposits.



Option D: Bagassosis is an occupational lung disease caused by the inhalation of bagasse or sugarcane dust. A thermophilic actinomycete, Thermoactinomyces sacchari, is the causative organism. The symptoms are breathlessness, cough, hemoptysis, and fever. Initially, it can result in acute diffuse bronchiolitis. If left untreated, it can progress to diffuse fibrosis, emphysema, and bronchiectasis.

Solution to Question 7:

The most likely diagnosis based on the given clinical condition and the peripheral smear showing spherocytes is hereditary spherocytosis. The investigation that can be done in this condition is the osmotic fragility test.

Hereditary spherocytosis is a type of hereditary hemolytic anemia that is mostly inherited in an autosomal dominant pattern. The mutations commonly affect ankyrin, band 3, spectrin, or band 4.2.

This disorder can present at a young age with severe jaundice, anemia, and splenomegaly. The unconjugated hyperbilirubinemia predisposes the patient to develop gallstones in early adulthood. Pigment stones are found in 40-50% of the affected adults.

Laboratory findings include normocytic anemia with unconjugated hyperbilirubinemia. The peripheral smear will show large spherical red blood cells without central pallor.

Laboratory investigations that are done to confirm the diagnosis:

• Osmotic fragility test

• Acid glycerol lysis test

• Eosin-5'-maleimide (EMA)–binding test

• SDS-gel electrophoresis test

There is no specific treatment available besides splenectomy.

Other Options:

Option B: Coombs test - AIHA (Autoimmune Hemolytic Anemia) - This test is performed to determine if the patient's red blood cells are coated with immunoglobulins or complement, leading to their destruction. AIHA typically does not present with a family history.

Option C: G6PD deficiency - This condition is characterized by hemolytic anemia triggered by oxidative stress, such as infections or exposure to certain drugs. G6PD deficiency is diagnosed by measuring G6PD enzyme activity or genetic testing.

Option D: Paroxysmal nocturnal hemoglobinuria - This rare acquired hemolytic anemia is characterized by episodes of hemolysis and hemoglobinuria, particularly during sleep. The diagnosis is made by testing for the presence of GPI-anchored proteins using flow cytometry.

Solution to Question 8:

The most probable diagnosis based on the given clinical scenario and the histopathology showing a "starry sky" pattern is Burkitt lymphoma.

Burkitt lymphoma is a highly aggressive B-cell non-Hodgkin lymphoma. It is characterized by the translocation of the c-Myc gene on chromosome 8. This leads to increased MYC protein levels. The translocation most commonly seen in Burkitt's lymphoma is t(8;14).

It can occur in 3 forms that have distinct clinical characteristics:

• African (endemic) Burkitt lymphoma presents as a mass involving the mandible. It may also involve the abdominal viscera, particularly the kidneys, ovaries, and adrenal glands.

• Sporadic (non-endemic) is often seen as a mass involving the ileocecum and peritoneum.

• Aggressive lymphoma or acute leukemia is seen in immunocompromised individuals

Histopathology shows tumor cells with round or oval nuclei, coarse chromatin, small nucleoli, minimal cytoplasm, and a high mitotic index. The phagocytes have abundant clear cytoplasm, creating a characteristic "starry sky" pattern. Bone marrow aspiration shows tumor cells with royal blue cytoplasm containing clear cytoplasmic vacuoles.

The tumor cells express surface IgM, CD19, CD20, CD10, and BCL6, a phenotype consistent with a germinal center B-cell origin.



Other options:

Option A: Acute myeloid leukemia is most commonly seen in elderly people. Histopathology from bone marrow aspirate reveals Auer rods (needle-like azurophilic granules) in the cytoplasm of the progranulocytes. Myeloblasts have two to four nucleoli and multiple coarse azurophilic granules.

The criterion for making a diagnosis of AML according to WHO requires one of the following:

• Blast cell count of ≥20% either in the peripheral blood or bone marrow

• Blast cell count of <20% in the presence of any one of the following three cytogenetic abnormalities t(15;17), t(8;21), and inversion 16 or t(16;16)

The peripheral blood smear shows numerous Auer rods in the cytoplasm of progranulocytes, characteristic of acute myeloid leukemia.



Option B: Follicular lymphoma presents in middle age with no sex predilection. The most common presenting symptom is painless generalized lymphadenopathy. Most follicular

lymphomas have characteristic (14:18) translocation, leading to overexpression of BCL2. Lymph nodes exhibit a nodular or nodular and diffuse growth pattern.

Two cell types are predominantly present in varying proportions:

• Centrocytes - Small cells with a cleaved nucleus and scant cytoplasm

• Centroblasts - Large cells with open nuclear chromatin, several nucleoli, and modest amounts of cytoplasm

The following image shows a lymph node biopsy in follicular lymphoma:



Option C: Multiple myeloma is a plasma cell neoplasm. It is associated with lytic bone lesions, hypercalcemia, renal failure, and acquired immune abnormalities.

Histopathological findings include

• Flame cells with fiery red cytoplasm

• Mott cells with multiple grapelike cytoplasmic droplets

• Inclusions bodies: Russell bodies (if cytoplasmic) or Dutcher bodies (if nuclear)

Solution to Question 9:

The diagnosis based on the above clinical scenario is post-streptococcal glomerulonephritis (PSGN). Subepithelial deposits will be seen on electron microscopy.

The rapid decline in renal function and raise in creatinine value to 4.5 mg/dL within 3 weeks, is suggestive of PSGN progressing to RPGN (Rapidly progressive glomerulonephritis). If the histopathology finding of the patient was asked then the answer would be crescent formation, which is characteristic of RPGN. Since the electron microscopy finding in the given patient is asked, the answer is subepithelial deposits, which is the characteristic finding of underlying pathology i.e. PSGN.

Post-streptococcal glomerulonephritis (PSGN) is caused by Group A β-hemolytic streptococcal infections. It is uncommon before the age of 3 years and most prevalent in kids between the ages of 5 and 12 years.

The patient typically develops an acute nephritic syndrome

• 1 to 2 weeks after streptococcal pharyngitis or

• 3 to 6 weeks after streptococcal pyoderma

From asymptomatic microscopic hematuria with normal renal function to gross hematuria with acute renal failure, the severity of kidney involvement varies. Patients may experience different levels of edema, hypertension, and oliguria depending on the extent of the renal involvement.

Laboratory findings: More than 90% of patients see a considerable reduction in serum C3 levels during the acute phase, which lasts 6 to 8 weeks before returning to normal.

In order to confirm the diagnosis, evidence of preceding streptococcal infection is required. The antistreptolysin O titer frequently rises following pharyngeal infections. The anti-deoxyribonuclease B level is the most effective single antibody titer to demonstrate cutaneous streptococcal infection.

Microscopic Findings in PSGN:

• Light microscopy: Enlarged, hypercellular glomeruli with diffuse mesangial and endothelial proliferation and leukocyte infiltration.

• Immunofluorescence: Granular deposits of IgG and complement C3 along the glomerular basement membrane (GBM) and in the mesangium

• Electron microscopy: Primarily subepithelial humps are seen. Subendothelial deposits can be seen in early disease stages.

Below is the image showing light microscopy findings of post-streptococcal glomerulonephritis.



Other options:

Option A: Subendothelial deposits - These are characteristic of membranoproliferative glomerulonephritis (MPGN) type I. MPGN typically presents with nephrotic or nephritic features

but does not necessarily follow a streptococcal infection.

Option C: Crescentic glomerulonephritis, also known as rapidly progressive glomerulonephritis (RPGN)is a type of nephritic syndrome associated with severe glomerular injury with necrosis and formation of crescents. The disease progresses rapidly and leads to renal failure in weeks to months when not treated.

Option D: Mesangial deposits - These are characteristic of IgA nephropathy, which typically presents with recurrent episodes of gross hematuria following upper respiratory infections, rather than acute nephritic syndrome following streptococcal infection.

Solution to Question 10:

The given clinical condition is most likely due to fluid overload.

Fluid overload also called transfusion-associated circulatory overload (TACO) is the main cause of death from transfusion. Risk factors include older age, renal failure, preexisting fluid overload, cardiac dysfunction, administration of a large volume of RBCs, and an excessive rate of transfusion in relation to the patient's hemodynamic tolerance.

TACO results in dyspnea, hypoxia, bilateral and predominantly alveolar infiltrates on chest x-ray, frequent systolic hypertension, and elevated brain natriuretic peptide. Fever may also present.

Prevention involves identifying at-risk patients, close monitoring, a slow transfusion rate, and the use of diuretics in hemodynamically stable patients with a history of TACO. Treatment requires stopping the transfusion and administrating oxygen and diuretics.

Other options:

Option A: ABO incompatibility reactions may present with hypotension, tachypnea, tachycardia, fever, chills, chest, and back pain, hemoglobinuria, and hemoglobinemia. In the most severe cases, DIC, acute renal failure, shock, and death may occur.

Option C: Hypocalcemia can be caused by citrate, which is used as an anticoagulant. It manifests as circumoral paresthesia, and changes in cardiac function may result from multiple rapid transfusions.

Option D: Graft-versus-host disease typically presents 5–10 days after transfusion, and it manifests as cytopenia, fever, a characteristic skin rash, diarrhea, and abnormal liver function.

Solution to Question 11:

The diagnosis of the patient based on the given clinical scenario and the histopathology showing spike and dome pattern is membranous nephropathy.

Membranous nephropathy is one of the most common causes of nephrotic syndrome in adults. It occurs commonly in men aged between 30-50 years. The majority of cases are idiopathic/primary that occur due to the autoantibodies against the renal antigen phospholipase A2 receptor. These autoantibodies are absent in secondary membranous nephropathy.

The various conditions leading to secondary membranous nephropathy are

• Drugs: Captopril, non-steroidal anti-inflammatory drugs NSAIDs, penicillamine, etc.

• Infections: Hepatitis B and C, syphilis, malaria

• Malignancy of lung, colon, and melanoma

• Autoimmune diseases like SLE, Sjorgen syndrome, sarcoidosis, primary biliary cirrhosis, etc.

As in this case, there is a history of hepatitis B infection, the likely etiology is secondary membranous nephropathy. Hence, PLA2R autoantibodies are absent.

Patients present with symptoms of nephrotic syndrome, nonselective proteinuria, and microscopic hematuria.

Diagnosis:

• Light microscopy shows diffuse thickening of the glomerular capillary wall without cellular proliferation. Irregular spikes and dome pattern on silver staining.

• Immunofluorescence microscopy shows granular deposits containing IgG and complement proteins along the glomerular basement membrane. IgG4 is the principal immunoglobulin deposited in most cases of primary membranous nephropathy.

• Electron micrograph: electron-dense subepithelial deposits (epimembranous spikes) and effacement of foot processes of podocytes

Management is primarily by the administration of immunosuppressive drugs such as steroids, cyclophosphamide, chlorambucil, and mycophenolate mofetil.

Image: The silver stain of the glomerulus highlights the proteinaceous basement membranes in black. There are characteristic "spikes" of the basement membrane between the immune deposits of membranous nephropathy. As the disease progresses, these spikes thicken to form dome-like protrusions. The black basement membrane material shown here appears as projections in the capillary loops.

Other options:

Option A: Minimal change disease is the most common cause of nephrotic syndrome in children (peaks between 2–6 years).

Biopsy findings in this condition are as follows:

• Light microscopy - glomeruli appear normal (hence called 'minimal change')

• Immunofluorescence studies - no Ig or complement deposits

• Electron microscopy - diffuse effacement of foot processes of visceral epithelial cells (podocytopathy)

Option C: FSGS is the most common cause of nephrotic syndrome in adults. It is a podocytopathy, with characteristic degeneration and focal disruption of visceral epithelial cells as well as effacement of foot processes. The electrophoretic pattern is more likely to yield a less prominent band in A, a more prominent band in C as there is a fall in the level of albumin with rising in the level of $\alpha 2$.

The histology of FSGS reveals focal and segmental hyalinosis and sclerosis of the capillary wall. Lipid droplets and foam cells are present in the sclerosed area.

Option D: Post-streptococcal glomerulonephritis (PSGN) is characterized by the sudden onset of gross hematuria, edema, hypertension, and renal insufficiency. It usually occurs following

a throat infection by nephritogenic strains of group A β-hemolytic streptococci.

Microscopic findings in PSGN:

• Light microscopy: Enlarged, hypercellular glomeruli with diffuse mesangial and endothelial proliferation and leukocyte infiltration.

• Immunofluorescence: Granular deposits of IgG and complement C3 along the glomerular basement membrane(GBM)and in the mesangium.

• Electron microscopy: Electron-dense deposits, or humps, are observed on the epithelial side of the GBM.

Solution to Question 12:

The clinical scenario of a child with complaints of fever, fatigue, and left ventricular dysfunction, and a biopsy showing lymphocytic infiltration in myocardial tissue, is suggestive of lymphocytic myocarditis.

The inflammatory process of the myocardium is termed myocarditis. Patients mostly present with fatigue, chest pain, cardiogenic shock, and even cardiac arrest. Viruses are the most common among the infectious causes of myocarditis. Endomyocardial biopsy is the gold standard for the diagnosis, and it shows diffuse, mononuclear, lymphocytic infiltration, as shown below.



Myocarditis begins with the direct invasion of an infectious agent. it subsequently replicates within and around the myocardium, causing myonecrosis. This infiltration and replication of the infectious agent lead to the destruction of cardiac tissue.

They are treated with supportive management.

Other Options:

Option A: Aschoff's bodies are the lesions found in the heart during acute rheumatic fever. They are composed of foci of T lymphocytes, occasional plasma cells, and plump-activated macrophages called Anitschkow cells. Anitschkow cells have abundant cytoplasm and central

round-to-ovoid nuclei with chromatin that is centrally condensed into a slender, wavy ribbon. Hence, they are also called caterpillar cells.

This condition classically occurs a few weeks after an episode of group A streptococcal pharyngitis or, less commonly, skin infections, and Aschoff's bodies are seen in the image below.



Option B: The myocarditis of Chagas disease has distinctive features of the parasitization of scattered myofibers by trypanosomes accompanied by a mixed inflammatory infiltrate of neutrophils, lymphocytes, macrophages, and occasional eosinophils. The clusters of amastigotes cause swelling of individual myocardial fibers and create intracellular pseudocysts. There is focal myocardial cell necrosis accompanied by extensive, dense, acute interstitial inflammatory infiltration throughout the heart.

The image below shows the amastigotes within a myocyte (arrow) and the surrounding inflammatory response.



Option C- In most bacterial infections, the heart can be involved through direct invasion and abscess formation. Causes include diphtheria, brucellosis, legionella, Meningococcus, and Mycoplasma, and require treatment of the systemic infection.

Pathology INI-CET 2023

Question 1:

Which of the following disorders is caused due to abnormal folding of proteins?

a) Cirrhosis

b) Creutzfeldt-Jakob disease

c) Nephritic syndrome

d) Sickle cell anemia

Question 2:

A 65-year-old male patient with chronic fatigue and lymphadenopathy presents to the outpatient department. His peripheral smear shows the presence of smudge cells. Which of the following additional investigations will you do to arrive at a diagnosis?

a) Polymerase chain reaction

b) Fluorescent in situ hybridization

c) Flow cytometry

d) Cytogenetics

Question 3:

Which of the following statements is true about blood products?

a) Platelets is stored between 20°C to 24°C with a shelf life of 5 days.

b) Cryoprecipitate is stored at -20°C with a shelf life of 6 months

c) Packed red cells stored with adenine, glucose, mannitol and saline is stored between -5°C to -15°C with a shelf life of 5 weeks

d) Fresh frozen plasma is stored between -5°C to -15°C with a shelf life of 2 years

Question 4:

A patient with lytic lesions on the skull is suspected of a diagnosis of Langerhan cell histiocytosis. Which of the following is a characteristic finding on electron microscopy?

a) Histiocytes

b) Birbeck granules

c) Eosinophils

d) Giant cells

Question 5:

Identify the mode of inheritance from the pedigree chart given below.



a) Autosomal dominant

b) Autosomal recessive

c) Pseudodominant inheritance

d) Incomplete penetrance

Question 6:

Match the following disorders with their genetic abnormalities.

a) 1-a, 2-d, 3-b, 4-c

b) 1-c, 2-a, 3-b, 4-d

c) 1-b, 2-d, 3-a, 4-c

d) 1-c, 2-b, 3-a, 4-d

Question 7:

Most specific indicator of alcohol-induced liver dysfunction is _____

a) Serum bilirubin

b) GGT

c) SGOT

d) SGPT

Question 8:

Most common cause of adrenal incidentaloma is_____

a) Subclinical Cushing's syndrome

b) Aldosterone producing adenoma

c) Pheochromocytoma

d) Non-functioning adenoma

Question 9:

A patient on long term hemodialysis due to renal failure is likely to develop which type of amyloidosis?

a) Amyloid light chain (AL)

b) Amyloid A (AA)

c) Amyloid β2 microglobulin (Aβ2M)

d) Amyloid transthyretin (ATTR)

Question 10:

Given below is an electron microscopy image of a cell. Identify the cell.



a) Target cell

b) Hairy cell

c) Gaucher cell

d) Sickle cell

Question 11:

Which of the following statements is true regarding the immunological abnormality that is seen with the lack of CD40 in B cells is?

a) Total lack of natural killer cells

b) Lack of CD8 mediated cytotoxicity

c) Decrease in IgG and increase in IgM

d) Inability of neutrophils to fight against infections

Question 12:

Match the following:

a) 1-b, 2-d, 3-a, 4-c

b) 1-d, 2-b, 3-c, 4-a

c) 1-a, 2-b, 3-c, 4-d

d) 1-a, 2-b, 3-d, 4-c

Question 13:

A 28-year-old female presenting with 36 weeks of gestation, underwent a C-Section for placenta accreta spectrum. During the procedure, there was profuse bleeding and she was transfused blood according to the massive blood transfusion protocol. Which of the following is not seen after a massive blood transfusion?

a) Hypomagnesemia

b) Hypokalemia

c) Hypocalcemia

d) Hypothermia

Question 14:

Which of the following are the true statements about amyloidosis?

a) 2,3,4

b) 1,3,4

c) 1,2,3

d) 1,2,4

Question 15:

Which of the following is an anti-apoptotic factor?

a) PUMA

b) BAD

c) BAK

d) MCL1

Question 16:

Thymus is designed to recognize the MHC molecules presented to it and distinguish the cells as self and foreign. Which gene is responsible for this?

a) NOTCH1

b) E2A

c) Rag 1/2

d) AIRE

Question 17:

Which of the following are the features of necrosis?

a) 1,2,4

b) 1,2,3

c) 2,3,4

d) 1,3,4

Question 18:

Which of the following is not essential in the workup of a mosaic Turner syndrome?

a) Anti-nuclear antibodies

b) Glucose tolerance test

c) Audiometry

d) Echocardiography

Question 19:

Most common genetic cause of hypercoagulability is _____

a) Factor V Leiden mutation

b) Protein C deficiency

c) Antithrombin III deficiency

d) G20210A prothrombin gene mutation

Question 20:

Identify the histopathological image.



a) Elastic fibres

b) Routine HPE

c) Amyloidosis

d) Collagen

Question 21:

BRCA 1 gene is associated with_____

a) 1,2,3

b) 1,3,4

c) 2,3,4

d) 1,2,4

Question 22:

A phlebotomist has been asked to take samples from a patient for multiple tests. Which of the following order of blood collection for testing will not contaminate the sample based on the additives present in the collecting tubes?

a) Light blue -> Red-> Lavender-> Grey

b) Light blue -> Lavender-> Red-> Grey

c) Grey-> Red->Light blue -> Lavender

d) Red-> Lavender-> Grey->Light blue

Question 23:

A transbronchial biopsy of a patient with fever, cough and weight loss shows the findings of the biopsy below. Which of the following chemical mediators are involved in its pathogenesis?



a) IFN gamma

b) Bradykinin

c) C5a

d) Prostaglandin

Question 24:

A patient with rapidly progressive renal failure (RPRF) presents with crescentic glomerulonephritis which is found to be pauci-immune. Which of the following antibodies are involved in this condition?

a) ANA

b) ANCA

c) Anti-GBM

d) Anti-PLA2R

Question 25:

Sickle cells have _____:

a) High mechanical fragility and high osmotic fragility

b) High mechanical fragility and low osmotic fragility

c) Low mechanical fragility and high osmotic fragility

d) Low mechanical fragility and low osmotic fragility

Question 26:

B cells express IgM and IgD antibodies at the same time due to which phenomenon?

a) Somatic hypermutation

b) Alternative RNA splicing

c) Affinity maturation

d) Allelic exclusion

Question 27:

A 75-year-old woman with diabetes and hypertension has a biopsy of her heart that is shown below. What is the diagnosis?



a) Dilated cardiomyopathy

b) Restrictive cardiomyopathy

c) Hypertrophic obstructive cardiomyopathy

d) Atrial fibrillation

Question 28:

A 62-year-old woman presents with a hard, irregular, fixed, and non-mobile lump in her breast. Her nipple is retracted, her skin is discolored, and her lymph nodes are enlarged. What does it mean if the tumor is luminal type A?

a) ER+ / HER2 neu-, mild to moderate proliferation

b) ER+ / HER2 neu+

c) ER- / HER2 neu-

d) ER+ / HER2 neu-, high proliferation

Question 29:

A 60-year-old vegetarian presents with fatigue and weakness that have been present for a few months. On physical examination, pallor is noted. Hemoglobin is 9 g/dL and MCV is increased at 110 fL. What is the most likely diagnosis?

a) Iron deficiency anemia

b) Megaloblastic anemia

c) Aplastic anemia

d) Thalassemia

Question 30:

Which of the following proteins shows a marked increase in size in nephrotic syndrome on serum electrophoresis?

a) Albumin

b) Alpha 2 globulin

c) Beta globulin

d) Gamma globulin

Question 31:

Seminomas are positive for which of the following immunohistochemistry markers?

a) 1, 2, 3

b) 1, 2, 3, 4

c) 2, 3, 4

d) 1, 3, 4

Question 32:

Match the following genes with their associated cancer types:

a) 1B, 2C, 3A, 4D

b) 1C, 2D, 3A, 4B

c) 1C, 2B, 3A, 4D

d) 1A, 2B, 3D, 4C

Question 33:

Which of the following stains are used in the cervical smear below?

a) Papanicolau stain

b) Toluidene blue

c) H & E stain

d) Giemsa stain

Question 34:

A woman presents with a fixed nodule in her breast with nipple retraction. Biopsy shows HER2/neu positivity. What chromogen is responsible for the brown color in the immunohistochemical stain?

a) FITC

b) Diaminobenzidine

c) Cyanate

d) Phycoerythrin

Question 35:

Transferrin saturation is decreased in which of the following conditions?

a) Hemosiderosis

b) Iron deficiency anemia

c) Hemochromatosis

d) Pernicious anemia

Question 36:

Post-orchidectomy, a patient's histopathology slide showed rods of Reinke's crystalloids. Which of the following conditions is associated with this?



a) Gonadoblastoma

b) Sertoli cell tumor

c) Leydig cell tumor

d) Yolk sac tumor

Question 37:

Match the following genetic disorders with their inheritance patterns:

a) A2, B4, C3, D1

b) A3, B4, C2, D1

c) A1, B4, C3, D2

d) A2, B4, C1, D3

Question 38:

Which of the following conditions will not show gastric acid secretion after the injection of pentagastrin?

a) Normal healthy individual

b) Gastrinoma

c) Pernicious anemia

d) Duodenal ulcer

Question 39:

You have a 55-year-old male patient with a history of infective endocarditis presents with high-grade fever, new-onset murmurs, and splinter hemorrhages. What is the correct way to collect blood culture samples?

a) 3 sets of culture samples, 2 from the peripheral lines and 1 from the affected valve with a gap of 30 minutes between the collection of each

b) 2 samples, 1 from the periphery and 1 from atrium, with a gap of 30 minutes between the collection of each

c) 1 from the periphery

d) 1 from the blood and 1 from the urinary catheter

Question 40:

A 26-year-old patient presents with a history of headache, fever, and neck stiffness for several days. Cerebrospinal fluid (CSF) analysis reveals the following parameters: protein 62, glucose 20 mg/dl (plasma glucose 140 mg/dl), and white blood cells composed of lymphocytes. What is the most likely diagnosis?

a) Coxsackievirus meningitis

b) HSV1 meningitis

c) Bacterial meningitis

d) Tubercular meningitis

Question 41:

The protease of V.cholera disrupts the extracellular part of which of the following proteins?

a) Occludin

b) Integrin

c) Cadherin

d) Claudin

Answer Key

Question No.	Correct Option
1	b
2	c
3	a
4	b
5	a
6	c
7	b
8	d
9	c
10	b
11	c
12	b
13	b
14	c

15	d
16	d
17	b
18	a
19	a
20	c
21	a
22	a
23	a
24	b
25	b
26	b
27	b
28	a
29	b
30	b
31	a
32	b
33	a
34	b
35	b
36	c
37	a
38	c
39	a
40	d
41	a

Detailed Explanations

Solution to Question 1:

Creutzfeldt-Jakob disease is caused due to abnormal folding of proteins.

Creutzfeldt-Jakob disease (CJD) is a prion disease that causes higher cortical function deficits. This infection can be acquired by corneal transplantation, contaminated electroencephalogram (EEG) electrode implantation, dura mater grafts, and contaminated human growth hormone (hGH) preparations.

Symptoms are almost always rapidly progressive, and the clinical presentation includes the following:

• Myoclonus

• Higher cortical deficits

• Psychiatric manifestations - depression, psychosis, and visual hallucinations

• Profound dementia - memory loss, impaired judgment, intellectual decline

• Visual impairment or cerebellar gait and coordination deficits in a few patients

• Nonspecific prodromal symptoms - fatigue, sleep disturbance, weight, loss, headache, anxiety, vertigo, malaise, and ill-defined pain

• Extrapyramidal dysfunction - rigidity, masklike facies, choreoathetoid movements;

• Seizures

• Hypoesthesia

• Supranuclear gaze palsy, optic atrophy

Patients commonly present with myoclonus and dementia. EEG shows characteristic periodic electrical bursts. MRI shows increased intensity in the basal ganglia and cortical ribboning (ribbon-like signal hyperintensity of cerebral cortical gyri). On light microscopy, the pathologic hallmarks of CJD are spongiform degeneration and astrocytic gliosis.



Other options:

Option C: Nephritic syndrome classically presents with tea- or cola-colored urine (hematuria), edema, hypertension, and oliguria.

Conditions which present as glomerulonephritis:

• Postinfectious glomerulonephritis

• IgA nephropathy (Option D)

• Membranoproliferative glomerulonephritis

• Henoch-Schönlein purpura (HSP) nephritis

• Systemic lupus erythematosus (SLE) nephritis

• Granulomatosis with polyangiitis

• Goodpasture syndrome (Option C)

• Hemolytic-uremic syndrome (HUS)

Option D: Sickle cell anemia is an autosomal recessive disorder that occurs due to the point mutation in the β-globin gene, which results in the replacement of glutamate residue with valine. This causes the production of sickle-shaped red blood cells due to the polymerization of deoxygenated hemoglobin (HbS). Factors that predispose to sickling are infection, dehydration, and acidosis. Clinical features are moderately severe hemolytic anemia with jaundice and painful bone crisis. The aplastic crisis stems from the infection of red cell progenitors by parvovirus B19, which causes a transient cessation of erythropoiesis and a sudden worsening of the anemia. In children, splenic sequestration crises occur which results in splenomegaly due to red pulp congestion.

Solution to Question 2:

An elderly patient with chronic fatigue and lymphadenopathy, and a peripheral smear showing the presence of smudge cells suggest a diagnosis of chronic lymphocytic leukemia. Immunophenotyping using flow cytometry is the investigation done to arrive at the diagnosis. The classical immunophenotypic feature of CLL is CD 19+, CD20+, CD 23+, CD5+.

Chronic Lymphocytic Leukemia (CLL) is a peripheral B-cell neoplasm usually affecting adults, more commonly men. Patients are often asymptomatic at diagnosis. When symptomatic, it presents non specifically with easy fatigability, weight loss, and anorexia. Generalized lymphadenopathy and hepatosplenomegaly are present in most symptomatic patients.

The chromosomal defects in CLL include:

• Deletion of 13q (commonest and has a good prognosis)

• Deletion of 11q

• Trisomy 12q

• Deletion of 17p (poor prognosis)

The peripheral blood smear in cases of chronic lymphocytic leukemia (CLL) contains small round lymphocytes having condensed chromatin and scant cytoplasm. Some of these cells may get disrupted while preparing the smears, producing smudge cells. An absolute lymphocyte count of >5000/mm3 is required for diagnosis. Hypogammaglobulinemia is commonly seen, making the patient susceptible to infections. A few patients may develop hemolytic anemia or thrombocytopenia due to autoantibodies made by non-neoplastic B cells.

On lymph node biopsy: Lymph nodes are composed of small lymphocytes admixed with larger activated lymphocytes. The large lymphocytes contain mitotically active cells and gather in loose aggregates referred to as proliferation centers. These proliferation centers are pathognomic to CLL when present.

Expectant management is the treatment of choice for asymptomatic patients with chronic lymphocytic leukemia. The gold standard for therapy is a combination of fludarabine, cyclophosphamide, and rituximab.

Richter's syndrome is a complication of CLL in which there is a transformation to an aggressive lymphoma, most commonly diffuse large B-cell lymphoma (DLBCL).

The image shows small round lymphocytes and smudge cells seen in CLL.



Solution to Question 3:

The statement that is true about blood products is platelets can be stored between 20°C to 24°C with a shelf life of 5 days.

Platelet concentrates can be stored for up to 5 days at +20°C to +24°C, with agitation. It is used in thrombocytopenia, platelet function defects, disseminated intravascular coagulation, during massive blood transfusion to prevent dilutional thrombocytopenia, and in bone marrow failure. Agitation during storage is necessary to maintain their viability and function. This is because platelets are stored at room temperature, where they continue to metabolize glucose and produce lactate. Agitation helps to maintain a uniform distribution of platelets, prevent their clumping together, and facilitate gas exchange.

Fresh frozen plasma and cryoprecipitate are generally stored at a temperature of –30 °C. They are thawed at temperatures between +30 °C and +37 °C in the blood bank before issue and transported to the ward at ambient temperature. They must be used immediately and should never be refrozen.

Whole blood and packed red cells are always stored at +2 °C to +6 °C and transported between +2 °C and +10 °C. Packed red cells stored with SAGM (sodium, adenine, glucose, and mannitol as components) provide a shelf-life of up to 42 days.

Solution to Question 4:

The presence of Birbeck granules in the cytoplasm is a characteristic finding of Langerhans cell histiocytosis (LCH) on electron microscopy.

Birbeck granules have a dilated terminal end producing a tennis racket–like appearance.
It contains the protein langerin (CD207). It is seen on electron microscopy.

The image below shows Birbeck granules on electron microscopy seen in LCH.



Histopathologically, Langerhans cells have a single large nucleus with a characteristic folded or grooved nucleus (coffee-bean appearance) with small inconspicuous nucleoli. Birbeck granules are seen in the cytoplasm. The tumor cells specifically express HLA-DR, S-100, and CD1a and langerin (CD207) markers.

The image below shows histopathological image of Langerhan cells with folded nuclei.



Langerhans cell histiocytosis occurs due to the proliferation of a special type of immature dendritic cells, called Langerhans cells. Their normal function is to present antigens. The most common mutation is BRAFV600E gain-of-function mutation.

It is characterized by single or multiple osteolytic bone lesions and can also infiltrate nearly every organ, like the skin (of the scalp), lungs, liver, spleen, bone marrow, or the central nervous system.

LCH can present as the following clinical entities:

• Multifocal multisystem Langerhans cell histiocytosis (Letterer-Siwe disease)

• Unifocal and multifocal unisystem Langerhans cell histiocytosis (eosinophilic granuloma)- The combination of bone defects, diabetes insipidus, and exophthalmos is called the Hand-Schüller-Christian triad.

• Pulmonary Langerhans cell histiocytosis

The below image shows lytic lesion of skull seen in LCH.



Solution to Question 5:

The pedigree chart shows an autosomal dominant mode of inheritance.

Autosomal dominant conditions are manifested in the heterozygous state, hence at least one parent of an index case is usually affected. In this mode of inheritance, both males and females are affected and both can transmit the condition. When an affected person marries an unaffected one, every child has one in two chance of having the disease.



Solution to Question 6:

The correct match is:

Huntington's disease (HD) is an autosomal dominant progressive brain disorder that causes uncontrolled movements, emotional problems, and loss of thinking ability. It occurs due to atrophy, with loss of neurons and gliosis in the caudate lobe, putamen, and globus pallidus. There is an abnormal expansion of the trinucleotide CAG in the Huntington gene on chromosome 4.

Trisomy 18 results in Edward's syndrome in which a complete extra copy of chromosome 18 is inherited as a result of meiotic nondisjunction. The clinical features include micrognathia, low-set ears, prominent occiput, and rocker bottom feet. Patients have significant hypertonia including clenched hands with overlapping fingers. Cardiac anomalies (ventricular septal defects and patent ductus arteriosus) and gastrointestinal anomalies (Meckel's diverticulum, malrotations) are common. Prenatal ultrasound reveals fetal growth restriction.

Genetic abnormalities	Disorders
1. Trinucleotide repeat	b. Huntington chore a
2. Trisomy 18	d. Edward syndrome
3. Trisomy 13	a. Patau syndrome
4. Hb beta chain Glu→ Val	c. Sickle cell anemia



Trisomy 13 results in Patau syndrome which presents with midline facial clefts(Cleft lip/palate) and musculoskeletal abnormalities,eg-polydactyly. Phenotypic features include microcephaly, low-set ears, hypertelorism, and a broad nasal bridge.

Sickle cell anemia is an autosomal recessive disorder that occurs due to the point mutation in the β-globin gene, which results in the replacement of glutamate residue with valine. This causes the production of sickle-shaped red blood cells due to the polymerization of deoxygenated hemoglobin (HbS). Factors that predispose to sickling are infection, dehydration, and acidosis. Clinical features are moderately severe hemolytic anemia with jaundice and painful bone crisis. The aplastic crisis stems from the infection of red cell progenitors by parvovirus B19, which causes a transient cessation of erythropoiesis and a sudden worsening of the anemia.

Solution to Question 7:

Among the given options, the most specific indicator of alcohol-induced liver dysfunction is gamma-glutamyl transferase (GGT).

GGT is commonly used in patients with alcoholic liver disease as a marker of excessive alcohol intake. GGT is present in many tissues, including the kidneys, pancreas, and liver. It is mainly derived from the liver and this is often used as a marker of hepatobiliary diseases. Chronic alcohol consumption leads to the induction of hepatic microsomal GGT which leads to a rise in GGT in alcoholics.

Laboratory investigations in patients with alcohol-induced liver dysfunction:

• AST (SGOT) and ALT (SGPT) elevations that do not exceed 400 IU/L.

• AST/ALT ratio of >1.5.

• Serum bilirubin >3 mg/dL.

Note that while GGT is a sensitive marker for alcohol-induced liver disease, it is not completely specific to this condition. Other conditions such as biliary obstruction or use of certain medications can also elevate GGT levels.

Solution to Question 8:

The most common cause of adrenal incidentaloma is non-functioning adenoma

Incidentalomas are evaluated to classify them as:

• Non-functional: most common

• Functional: may present with Cushing's syndrome, pheochromocytoma, aldosteronoma, cortisol-producing adenoma, etc.

• Secondary metastasis: from the breast, lung, and kidneys are seen in 2% of the patients.

For patients without a confirmed diagnosis, it is recommended to conduct specific tests to identify underlying hormonally active tumors. These tests include:

(a) Performing a low-dose (1 mg) overnight dexamethasone suppression test to exclude subclinical Cushing's syndrome.

(b) Collecting a 24-hour urine sample to measure levels of catecholamines, metanephrines, or VMA, or measuring plasma metanephrine levels to rule out the presence of a pheochromocytoma.

(c) In hypertensive patients, conducting tests for serum electrolytes, plasma aldosterone, and plasma renin to exclude the presence of an aldosteronoma.

Management of incidentalomas:

• Non-functional: Tumors with a size <4cm are managed conservatively. MRI surveillance is done every 3-6 months. For size >4cm, laparoscopic adrenalectomy is done.

•

Functional tumors: Hormonal assay studies are done in suspected functional tumors or patients presenting with symptoms. If positive, it should be managed accordingly. If not managed, functional tumors can become malignant and the incidence increases 4-fold if the size of the lesion crosses 4 cm.

•

Secondary metastasis: Fine-needle aspiration and biopsy is performed. Tests for pheochromocytoma should always be done before this procedure to avoid precipitating a hypertensive crisis.

Other options:

Option A: Patients with incidentaloma may present with subclinical Cushing’s syndrome. This condition is identified by subtle indications of excessive cortisol, including weight gain, thinning of the skin, facial rounding, diabetes, and high blood pressure. These symptoms are accompanied by the absence of the typical daily variation in cortisol secretion, autonomous cortisol production, and insensitivity to suppression by dexamethasone. Although the total amount of cortisol produced and the levels of cortisol in 24-hour urine samples may appear normal, other clinical signs and hormone irregularities indicate cortisol excess.

Option B: Aldosterone-producing adenomas originate in the zona glomerulosa of the adrenal glands. These tumors are typically 10 to 20 mm in diameter when initially detected, and they have

a distinct appearance—well-defined and exhibiting a golden yellow color upon slicing, often referred to as the "canary tumor."

The most commonly observed mutations occur in the KCNJ5 gene, which encodes the potassium channel. These mutations are believed to cause an influx of calcium into glomerulosa cells, leading to increased secretion of aldosterone and cellular proliferation. These tumors are more frequently found in females and are associated with larger tumor sizes and higher levels of aldosterone. However, they are not linked to adrenal hyperplasia.

Option C: Pheochromocytomas and paragangliomas are catecholamine-producing tumors. The dominant clinical feature in pheochromocytoma is hypertension. Most patients have paroxysmal episodes, which are abrupt elevations in blood pressure. They are accompanied by tachycardia, palpitations, headache, and sweating. Lab diagnosis of pheochromocytoma is based on the demonstration of increased urinary excretion of free catecholamines and their metabolites such as VMA.

Solution to Question 9:

Patients on long-term hemodialysis for renal failure can develop amyloidosis as a result of the deposition of β2-microglobulin.

β2-microglobulin is a component of the MHC class I molecule and a normal serum protein. In the past, it was retained in the circulation of patients with chronic kidney disease because it could not be filtered through dialysis membranes. In these patients, it would get deposited in the joints, muscles, tendons, and ligaments. One of the complications which can occur as a result is carpal tunnel syndrome. However, with the advent of new dialysis filters, this complication has become uncommon now.

Solution to Question 10:

The given electron microscopic image shows a hairy cell.

Hairy cell leukemia is so-called because of the appearance of the leukemic cells, which have fine hair-like projections that are best recognized under the phase-contrast microscope.

Hairy cell leukemia is a distinctive B cell neoplasm, that predominantly affects middle-aged males. The markers expressed in hairy cell leukemia include the usual B-cell markers- CD-19 and 20, and specific markers such as CD 11c, CD 25, CD103, and annexin A1.

On peripheral blood smear, hairy cells have round or reniform nuclei and moderate amounts of pale blue cytoplasm with thread or bleb-like extension. In the bone marrow, these cells are usually trapped in a mesh of reticulin fibers and so it is difficult to aspirate and hence termed a dry tap.

Patients commonly present with massive splenomegaly due to infiltration of the leukemic cells. Pancytopenia due to marrow involvement can also be seen. There can be atypical monocytopenia due to which these patients experience an increased incidence of atypical mycobacterial infections. These tumors respond well to chemotherapy and BRAF (B-type Raf proto-oncogene) inhibitors and they have an excellent prognosis.

The classic hairy cells can be seen in the slide is shown below:



Other options:

Option A: Target cells have a centrally located disk of hemoglobin surrounded by an area of pallor. There is an outer rim of hemoglobin adjacent to the cell membrane giving the cell the appearance of a target. Conditions in which target cells appear are thalassemia, hepatic disease with jaundice, hemoglobin C disorders, and postsplenectomy state.

Option C: Gaucher disease is a lysosomal storage disorder with a deficiency of acid-β-glucosidase. It has an autosomal recessive mode of inheritance. It leads to the accumulation of glucosylceramide in the reticuloendothelial system. The features of this disease include growth retardation, hepatosplenomegaly, thrombocytopenia, anemia, and Erlenmeyer flask deformity of the distal femur. Histology shows the presence of Gaucher cells, a characteristic wrinkled-paper appearance resulting from the presence of intracytoplasmic substrate inclusions. The intelligence of the child is normal.



Option D: Sickle cell anemia is an autosomal recessive disorder that occurs due to the point mutation in the β-globin gene, which results in the replacement of glutamate residue with valine. This causes the production of sickle-shaped red blood cells due to the polymerization of deoxygenated hemoglobin (HbS).



Solution to Question 11:

The deficiency of the CD40 and its absence in B cells leads to Hyper IgM syndrome. A decrease in IgG and an increase in IgM is seen in this syndrome.

The patients most often have loss-of-function mutations involving either CD40 or activation-induced cytidine deaminase (AID), a DNA-editing enzyme expressed in B cells that are required for Ig class switching and affinity maturation.

The deficiency of the CD40 and its absence in B cells interferes with class switching and the production of memory cells. The B-cell response to antigens, is unaffected, accounting for the presence of increased IgM antibodies. Hence, the condition is known as Hyper IgM syndrome.

The serum of the patients has normal or elevated levels of IgM but no IgA or IgE and extremely low levels of IgG. Clinical features are recurrent pyogenic infections because of low levels of opsonizing IgG antibodies. They are highly susceptible to pneumonia caused by Pneumocystis jirovecii.

Solution to Question 12:

The correctly matched associations of the following conditions are:

• Silicosis: Crazy paving pattern

• Mesothelioma: Malignant pleural effusion without mediastinal shift

• Caplan syndrome: Initially demonstrated in coal workers

• Asbestosis: Basal lobes are involved

Caplan's syndrome also called rheumatoid pneumoconiosis, is a combination of rheumatoid arthritis and pneumoconiosis. It manifests as pneumoconiotic nodules along with features of seropositive rheumatoid arthritis. Caplan syndrome was first described in coal miners and later in patients with silicosis too.

	Silicosis	Asbestosis	Malignant mesothelioma
Exposure	Crystallinesilica	Asbestos	Most specific neoplasm as sociated withasbestosexpo sure
Radiologic/ Gross findi ngs	Simple silicosis - small ro undedopacitiesin upper lo besComplicated silicosis - Larger opacities > 1cm an deggshell calcificationof h ilar lymph nodesAcute sili cosis - ground-glass opaci ties on X-ray andcrazy pa ving patternon HRCT (int erlobular septal thickenin g with background of gro und-glass opacity)	X-ray shows peripheralret icular opacitiespredomina ntly in thebasal lobesHRC T shows subpleural fibroti c lines, irregular opacities, tractionbronchiectasis, an dhoneycombing	Diffuse gelatinous,greyish -pink lesionarising either from the visceral or pariet al pleura.
Clinical manifestations	Increased risk of developi ng TB and bronchogenic c arcinoma	It mainly affects the lower lobesPFT shows arestricti ve patternwith a decrease in lung volumes and diffu sing capacity	Strong male predilection 3:1Can causemalignant pl eural effusion

Solution to Question 13:

Hyperkalemia is seen after a massive blood transfusion.

Massive transfusion is defined as the transfusion of 50% of the patient's total blood volume over 3 hours or >5–10 units of RBCs.

Complications are:

• Citrate toxicity

• Hypothermia (Rapid transfusion of blood components at 4°C)

• Hypocalcemia that presents as circumoral paresthesia.

• Hyperkalemia(RBC leakage during longer storage, and irradiation increases the concentration of potassium in the unit)

• Dilutional coagulopathy

Prevention includes the usage of washed or plasma-reduced RBC components or a storage age of <7–10 days and the avoidance of RBC components stored for >24 h after irradiation. Treatment includes administration of insulin, glucose, calcium gluconate, and furosemide.

Note: As the question was a part of the INI-CET May 2023 exam, the best possible answer has been marked.

Solution to Question 14:

The true statements about amyloidosis are:

• It is an extracellular protein.

• It is called amyloid because of starch deposition.

• Beta-pleated sheets are seen on X-ray crystallography.

Amyloidosis refers to extracellular deposits of fibrillar proteins.

Abnormal fibrils are produced by the aggregation of misfolded proteins. The fibrillar deposits bind a wide variety of proteoglycans and glycosaminoglycans, including heparan sulfate and dermatan sulfate, and plasma proteins, notably the serum amyloid P component (SAP). Amyloid appears as an amorphous, eosinophilic, hyaline, extracellular substance.

The adsorbed proteins contain abundant charged sugar groups that give the deposits staining characteristics that resemble starch. Hence, these deposits were called amyloid.

It could be classified as systemic, localized, or hereditary amyloidosis. Microscopically, all types of amyloid deposits are made up of continuous, nonbranching fibrils. A characteristic cross-β-pleated sheet conformation is visualized on X-ray crystallography and infrared spectroscopy.

The stain used for amyloid deposits is congo red staining, which imparts pink to red color to the amyloid under ordinary light and apple-green birefringence under polarised microscopy.

AL type(Immunoglobulin light chain amyloidosis) of protein belongs to primary amyloidosis. It is seen in multiple myeloma and other monoclonal plasma cell proliferation.

The image shows a congo-red stained biopsy specimen with the following:

(A) Focal positive staining (orange-red) of amorphous extracellular material

(B) Apple-green birefringence under polarized light.

The next best step is to determine the type of precursor protein by immunohistochemistry, immunoelectron microscopy, or extraction and biochemical analysis using mass spectrometry.



Solution to Question 15:

Among the given options MCL1 is an anti-apoptotic factor.

Anti-apoptotic factors are BCL2, BCL-XL, and MCL1. They reside in outer mitochondrial membranes as well as the cytosol and endoplasmic reticulum membranes. By keeping the mitochondrial outer membrane impermeable they prevent leakage of Cytochrome c and other death-inducing proteins into the cytosol.

Pro-apoptotic factors are BAX and BAK. Upon activation, BAX and BAK oligomerize within the outer mitochondrial protein and promote mitochondrial outer membrane permeability which leads to leakage of death-inducing proteins into the cytosol.

Regulated apoptosis initiators are BAD, BIM, BID, Puma, and Noxa. During cellular stress and damage, they get activated and initiate apoptosis.

Solution to Question 16:

The above process is called central tolerance and the AIRE (AutoImmune REgulator) gene is necessary for this process.

Central tolerance is a process in which immature self-reactive T and B lymphocyte clones that recognize self-antigens during their maturation in the central lymphoid organs (Thymus for T cells, Bone marrow for B cells) are killed or rendered harmless.

The removal of self-reacting T lymphocytes in the thymus is called negative selection/ deletion. For this process, AIRE protein is needed. Mutations in the AIRE gene result in autoimmune polyendocrinopathy.

B-cells undergo a process called receptor editing. When developing B cells encounter a self-antigen, it formulates a change that makes the antigen unrecognizable. Self-reacting B cells

can also undergo apoptosis.

Other options:

Option A: The NOTCH1 gene is associated with signal transduction. Point mutations and translocations of this gene are associated with leukemias, lymphomas, and breast carcinoma. NOTCH1 is the most commonly affected gene in T-cell acute lymphoblastic leukemia (T-ALL).

Option B: E2A also known as transcription factor(TCF3) is an important regulator of gene expression in germinal center B cells. It is believed that E2A drives the expression of a set of genes, including cyclin D, that collaborate with MYC to enable the rapid growth that characterizes Burkitt lymphoma.

Option C: RAG1 and RAG2 are developed by B and T cells that permit the assembly of functional antigen receptor genes. Adaptive immunity of B and T cells relies on their ability to diversify their antigen receptor genes.

Solution to Question 17:

The features of necrosis are:

• Disrupted cell membrane

• Induces inflammation

• Swelling of the cell

The characteristics of necrosis are denaturation of cellular proteins, leakage of cellular contents through damaged membranes, local inflammation, and enzymatic digestion.

The causes are loss of oxygen supply, exposure to microbial toxins, burns, and other forms of chemical and physical injury.

When the cell membrane is damaged, lysosomal enzymes enter the cytoplasm and digest the cell. The contents from the cell leak through the damaged plasma membrane into the extracellular space, where they elicit inflammation.

Morphological changes that occur in necrosis are:

• The cell size is increased.

• Cytoplasmic changes -

• Increased eosinophilic staining on H&E specimens

• Moth-eaten appearance of the cytoplasm - due to vacuolated cytoplasmic organelles formed by enzymatic degradation

• Glassy and homogenous cytoplasm - due to loss of glycogen in the cytoplasm

•

• Nuclear changes -

• Pyknosis - nucleus first shrinks and becomes more condensed

• Karyorrhexis - fragmentation of the pyknotic nucleus

• Karyolysis - nucleus fades away leaving behind only the cell membrane or empty cells

Solution to Question 18:

Among the given options, anti-nuclear antibodies are not essential in the workup of a mosaic Turner syndrome.

Turner syndrome occurs when females have a deletion or non-functional X chromosome. Approximately 50% of individuals with Turner syndrome have a condition called monosomy X (45, XO), where they only have one X chromosome. The remaining 50% have a mosaic chromosomal component (45, X with mosaicism).

Mosaic Turner syndrome does not stem from inheritance. Instead, it is the outcome of a random occurrence during cell division in the early development of the fetus. This results in a combination of cells in the affected individual's body, where some cells possess the usual two sex chromosomes, while others have only one copy of the X chromosome.

The clinical features of Turner's syndrome are mentioned in the image below:



Prenatal diagnosis of Turner syndrome can be conducted through procedures like chorionic villus sampling or amniocentesis. If a prenatal ultrasound reveals fetal hydrops, cystic hygroma, or cardiac defects, it may raise suspicions of Turner syndrome. However, the definitive confirmation of the diagnosis is typically done after birth through karyotype testing.

After receiving a diagnosis of Turner syndrome, it is essential to manage the condition by assessing for any associated abnormalities such as cardiac anomalies, renal anomalies, and learning disabilities. Screening should be conducted as part of the initial evaluation, followed by regular periodic screenings.

During the initial diagnosis, patients should undergo renal ultrasonography and cardiovascular evaluation. Echocardiography is typically performed in infants and children, while MRI is recommended for older girls and women. Regular monitoring of hearing, including serial audiology evaluations, is recommended throughout life, with an audiology evaluation every 3 yrs

in children and every five years in adults.

Several laboratory tests are recommended for screening purposes at specific ages:

For patients aged four years and above, screening involves checking serum TSH to detect autoimmune thyroiditis and conducting tissue transglutaminase testing with total IgA to screen for celiac disease.

For patients aged ten years and above, screening includes fasting blood glucose, glycated hemoglobin, ALT, AST, serum creatinine, and urinalysis. These tests are performed to screen for conditions such as diabetes mellitus, fatty liver, and kidney disease.

For evaluation of vitamin D deficiency, serum 25-hydroxyvitamin D should be measured between 9 to 11 years of age and every 2 to 3 years thereafter.

Solution to Question 19:

The most common genetic cause of hypercoagulability is the Factor V Leiden mutation.

Primary (inherited) hypercoagulability is most often caused by mutations in the factor V and prothrombin genes. Factor V Leiden mutation is the most common cause followed by prothrombin G20210A mutation.

The Leiden mutation is inherited in an autosomal dominant pattern. The mutation makes factor V resistant to cleavage and inactivation by protein C. This leads to the inactivation of an important antithrombotic counterregulatory pathway.

Protein C, Protein S, and antithrombin III deficiency are rare causes of primary hypercoagulability.

Other options:

Option B: Protein C deficiency causes thrombosis in teenagers. It may also be induced by liver dysfunction, vitamin K antagonists, renal failure, DIC, and active thrombosis.

Option C: Antithrombin III forms a complex with thrombin by binding to heparin on endothelial cells. This complex inhibits coagulation. Its deficiency presents as early-age thrombosis (less than 50 years old) and it carries the highest risk for thrombotic events among the inherited thrombophilias.

Option D: The prothrombin G20210A mutation results in increased levels of prothrombin which demonstrates a higher risk for arterial and venous thrombotic events.

Solution to Question 20:

The given histopathological slide shows congo red staining that is used in staining amyloid.

Amyloidosis refers to extracellular deposits of fibrillar proteins. Abnormal fibrils are produced by the aggregation of misfolded proteins. The fibrillar deposits bind a wide variety of proteoglycans and glycosaminoglycans, including heparan sulfate and dermatan sulfate, and plasma proteins, notably serum amyloid P component (SAP). Amyloid appears as an amorphous,

eosinophilic, hyaline, extracellular substance.

It could be classified as systemic, localized, and hereditary amyloidosis. Microscopically, all types of amyloid deposits are made up of continuous, nonbranching fibrils. A characteristic cross-β-pleated sheet conformation is visualized on X-ray crystallography and infrared spectroscopy.

The stain used for amyloid deposits is congo red staining, which imparts pink to red color to the amyloid under ordinary light and apple-green birefringence under polarised microscopy.

The image shows a congo-red stained biopsy specimen with the following:

(A) focal positive staining (orange-red) of amorphous extracellular material

(B) apple-green birefringence under polarized light

The next best step is to determine the type of precursor protein by immunohistochemistry, immunoelectron microscopy, or extraction and biochemical analysis using mass spectrometry.



Other options:

Connective tissue fibers	Stain	Appearance
Collagen fibers	Unaided dye	White
Hematoxylin and eosin	Light pink	
Masson's trichrome	Blue	
Verhoeff-Van Gieson	Red	
Silver impregnation	Brown	
Reticular fibers	Silver impregnation	Black
Elastic fibers	Verhoeff-Van Gieson	Black
Aldehyde fuchsin	Purple	

Connective tissue fibers	Stain	Appearance
Orcein	Purplish brown	



Solution to Question 21:

The statements associated with the BRCA1 gene are:

• Repair of ds DNA break

• Risk of development of ovarian cancer

• Tumor suppressor gene

BRCA1 and BRCA2 are tumor suppressor genes. Mutations of these genes are associated with an increased risk of breast cancer in females. They play an important role in the repair of double-stranded breaks in DNA.

In addition to breast cancer, women with BRCA1 mutations have a higher risk of developing epithelial ovarian cancers, and men have a slightly higher risk of prostate cancer. BRCA2 is associated with cancers of the ovary, prostate, pancreas, bile ducts, stomach, melanocytes, and B lymphocytes.

Few germline BRCA2 mutations can cause Fanconi anemia. Both BRCA proteins and Fanconi proteins function together in a DNA damage response network linked to homologous recombination repair. Defects in this pathway can lead to the activation of a nonhomologous end-joining pathway that can result in telomere shortening.

Solution to Question 22:

The correct color sequence of the blood collection tube is light blue-> red-> lavender-> grey.

The Clinical & Laboratory Standards Institute (CLSI) has recommended a color sequence for multiple collections of venous and/or arterial blood samples in a single patient.

Yellow, pink, and blue→ light Blue→ red→ green→ lavender→ grey.



Solution to Question 23:

Interferon gamma (IFN-γ) is a pro-inflammatory cytokine that is involved in the formation of granulomas. It activates macrophages to differentiate into epithelioid cells and multinucleated giant cells, which are the main components of granulomas.

IFN-γ is the most important chemical mediator involved in the pathogenesis of granulomas. It is produced by T helper 1 (Th1) cells and other immune cells in response to a variety of stimuli, including infections, autoimmune diseases, and foreign bodies.

Interferon-gamma-induced changes in the macrophages:

• Augmented ability of the macrophages to phagocytose and kill micro-organisms.

• Increased expression of class 2 MHC molecules to enhance antigen presentation.

• Increased secretion in presence of tumor necrosis factor, interleukin-1, and other inflammatory chemokines.

• Amplification of the Th1 response by producing more interleukin-12.

Granulomas are characteristic of a number of diseases, including tuberculosis, sarcoidosis, and Crohn's disease. In these diseases, IFN-γ plays a central role in the formation of granulomas, which help to contain and eliminate the offending agent. However, granulomas can also be formed in response to non-harmful stimuli, such as foreign bodies or allergens. In these cases, granulomas can be harmful and can lead to tissue damage and organ dysfunction.

B. Bradykinin: Bradykinin is a vasodilator that is involved in the inflammatory response. It is not directly involved in the formation of granulomas.

C. Complement C5a: Complement C5a is a chemoattractant that attracts neutrophils and other inflammatory cells to the site of infection or injury. It is not directly involved in the formation of granulomas.

D. Prostaglandin: Prostaglandins are a group of signaling molecules that are involved in a variety of physiological processes, including inflammation. Prostaglandins can promote or inhibit the formation of granulomas, depending on the context.

Solution to Question 24:

ANCA - antineutrophil cytoplasmic antibody is the most common antibody associated with pauci-immune crescentic glomerulonephritis.

There are two main types of ANCA: myeloperoxidase (MPO)-ANCA and proteinase-3 (PR3)-ANCA. MPO-ANCA is more strongly associated with renal-limited vasculitis, while PR3-ANCA is more strongly associated with systemic vasculitis.

Rapidly progressive glomerulonephritis (RPGN) is a clinical entity with features of nephritic syndrome and rapid loss of renal function. It is a syndrome associated with severe glomerular injury with necrosis and the formation of crescents, but it does not denote a specific etiologic form of glomerulonephritis.

In most cases, RPGN is immune-mediated. Circulating anti-neutrophil cytoplasmic antibodies (ANCAs) are detected in the pauci-immune type of RPGN. Serum analyses for anti-GBM antibodies, antinuclear antibodies, and ANCAs are helpful in the diagnosis of specific subtypes

Based on immunologic findings, RPGN is classified as:

Anti-GBM antibody-mediated disease (Type I) is characterized by linear GBM deposits of IgG and C3 on immunofluorescence and includes renal limited disease and Goodpasture's disease. Here,

the antibodies are produced against a peptide present in the noncollagenous domain of the α3 chain of collagen IV of the GBM. This serves as the antigen which is common to the alveoli and the glomerulus.

Immune complex deposition (Type II) is characterized by granular deposits of antibodies and complement on immunofluorescence and can be a complication of any of the immune complex nephritides which include:

• Postinfectious glomerulonephritis

• Lupus nephritis

• IgA nephropathy

• Henoch-Schönlein purpura

Pauci-immune (Type III) exhibits no characteristic immunoglobulin or complement deposits on immunofluorescence. It includes

• ANCA-associated

• Granulomatosis with polyangiitis

• Microscopic polyangiitis

Other options:

Option A: ANA - antinuclear antibody: ANA is a non-specific antibody seen in most autoimmune disorders; including lupus, rheumatoid arthritis, and scleroderma. ANA is not typically associated with pauci-immune crescentic glomerulonephritis.

Option C: Anti-GBM - anti-glomerular basement membrane antibody: directed against the glomerular basement membrane. Anti-GBM is associated with anti-GBM disease/Goodpasture syndrome, a type of rapidly progressive glomerulonephritis that is characterized by linear deposits of IgG along the BM.

Option D: Anti-PLA2R - anti-phospholipase A2 receptor antibody: Anti-PLA2R is an antibody that is directed against the phospholipase A2 receptor; associated with membranous nephropathy, where thickening of the glomerular basement membrane is characteristic.

Solution to Question 25:

Sickle cells have high mechanical fragility and low osmotic fragility.

When subjected to mechanical stress, such as shearing forces. trauma, turbulence. Sickle cells have high mechanical fragility because they are more rigid and less deformable than normal red blood cells. This makes them more likely to break apart when subjected to mechanical stress.

The higher concentration of hemoglobin in sickle cells makes them less likely to break apart when subjected to osmotic stress; such as changes in the concentration of salt - de/overhydration, CKD, etc. This is because the sickle cell hemoglobin (HbS) protein is more resistant to swelling in hypotonic solutions.

Conditions that can cause increased mechanical fragility of red blood cells:

• Sickle cell anemia

• Thalassemia: Thalassemia can cause RBCs to become smaller and less flexible. This can make them more mechanically fragile.

• Red blood cell membrane disorders

• G6PD deficiency

• Autoimmune hemolytic anemia

• Malaria

Conditions that can cause increased osmotic fragility of red blood cells:

• Hereditary spherocytosis

• Elliptocytosis

• Ovalocytosis

• Autoimmune hemolytic anemia

Disease	Mechanical fragility	Osmotic fragility
Sickle cell anemia	High	Low
Thalassemia	High	Low
Malaria	High	High

Solution to Question 26:

B cells express IgM and IgD antibodies at the same time due to alternative RNA splicing.

Alternative RNA splicing is the process by which different proteins are produced from the same RNA transcript by splicing it in different ways.

Mature circulating B cells initially express IgM and IgD antibodies on their surfaces before they are stimulated to produce specific antibodies. The heavy chains in IgM and IgD are mu (μ) and delta (δ), respectively. The mu and delta genes for these heavy chains are located proximal to the VDJ encoding regions (i.e., variable, diversity, and joining regions) of the heavy chain locus. A primary RNA transcript of this entire region is produced, which then undergoes alternative RNA processing, enabling the production of both transmembrane IgM and IgD on B cells.

Other options:

Options A and C: Somatic hypermutation is a cellular response by which the immune system adapts to the new foreign elements that confront it. They do so by adding nucleotides in V gene regions of heavy and light chains, resulting in a better fit of antigen with the antibodies. This hence increases the affinity of antibodies to antigens. Due to this, the subset of plasma cells with these improved hypervariable regions are more strongly selected by the antigen and therefore constitute a larger part of the population of antibody-producing cells. This process is called affinity maturation.

Option D: Allelic exclusion is the process by which a cell expresses one of the two alleles of the gene and excludes the other allele.

B cells are diploid, therefore, each B cell has six gene segment pools related to antibody chains:

• Two heavy-chain pools (one from each parent) and

• Four light-chain pools (one κ and one λ from each parent).

However, each B cell uses only two of the six gene segment pools: one out of the two heavy-chain pools (maternal and paternal) and one out of the four light-chain pools. Thus, each B cell must choose not only between its κ and λ light-chain pools but also between its maternal and paternal light-chain and heavy-chain pools. This process of choosing is by allelic exclusion.

Solution to Question 27:

The hyaline, eosinophilic deposits seen on the biopsy point to a diagnosis of cardiac amyloidosis, which most commonly leads to restrictive cardiomyopathy.

Transthyretin is the major component of senile cardiac amyloid deposits. It
is a normal serum protein synthesized in the liver that transports thyroxine and retinol-binding protein. The deposits may accumulate subendocardially, and slowly expand - compressing the myocardial fibres - leading to pressure atrophy of myocaridial fibres. The stain used for amyloid deposits is congo red staining, which imparts pink to red color to the amyloid under ordinary light and apple-green birefringence under polarised microscopy.

Presentation of cardiac amyloidosis:

• Restrictive cardiomyopathy- most common

• Asymptomatic

• Dilation

• Arrhythmias

• Ischemic disease

The echocardiographic findings in cardiac amyloidosis may include a distinctive granular, sparkling appearance of the thickened cardiac walls, representative of the infiltrative process. Despite the thickened ventricular walls, the ECG in cardiac amyloidosis typically demonstrates low limb lead voltage, often accompanied by ventricular conduction delay and pseudoinfarction patterns. Suspect cardiac amyloidosis when there is thickened ventricular walls with an ECG that shows low voltage. Amyloid infiltration can be detected with gadolinium enhancement on MRI.

A. Dilated cardiomyopathy/DCM is a condition in which the heart muscle becomes enlarged and weakened. This makes it difficult for the heart to pump blood effectively. DCM can be caused by a variety of factors, including coronary artery disease, high blood pressure, and diabetes.

B. Hypertrophic cardiomyopathy/HOCM is a condition in which the IV septum becomes thickened and is typically genetic. On biopsy, myocardial disarray would be noted.

D. Atrial fibrillation by rapid, irregularly irregular contractions of the atria that can lead to tachycardia induced cardiomyopathy. AF can be caused by a variety of factors, including heart disease, high blood pressure, and diabetes.

Solution to Question 28:

Luminal type A breast cancer is defined as estrogen receptor (ER) and/or progesterone receptor (PR) positive, HER2-negative, and low to moderate Ki-67 proliferation index.

This means that the tumor cells have ER and/or PR proteins on their surface and that they are not overexpressing the HER2 gene. The Ki-67 proliferation index is a marker of the rate of tumor proliferation. A low to moderate Ki-67 index indicates that the tumor cells are dividing slowly; ie: less aggressive.

Luminal type A breast cancer is the most common subtype of breast cancer and has the best prognosis. It is typically a low-grade, slow-growing tumor that is responsive to hormone therapy.

Luminal A type is more commonly associated with BRCA2 mutations. Hormone receptor-positive tumors are responsive to hormonal chemotherapeutic agents like tamoxifen and aromatase inhibitors. In Her-2 negative cases, trastuzumab is not effective as it is a recombinant humanized monoclonal antibody directed against HER-2.

Option B: ER+ / HER2neu+: This is a separate subtype that is more aggressive, and responds to targeted chemotherapy.

Option C: ER- / HER2neu-: ER-negative breast cancer is a subtype that is not responsive to hormone therapy.

Option D: ER+ / HER2neu-, high proliferation: This is a characteristic of luminal type B breast cancer, which is a more aggressive subtype than luminal type A.

Molecular subtypes	ER	PR	HER 2	Ki-67	Cytokeratin 5 and 6
Luminal A(Most common type and has the best prognosis)	+ve	+ve	-ve	Low	-ve
Luminal B	+ve	+ve	-ve	High	-ve
	+ve	+ve	+ve	Any	-ve
HER2 enriched	-ve	-ve	+ve	Any	-ve
Basal-like or Triple-negative breast cancer(Aggressive, has worst prognosis but shows good response to chemotherapy, high PCR rates)	-ve	-ve	-ve	Any	+ve
Unclassified, now also known asClaudin-low	-ve	-ve	-ve	Any	-ve

Solution to Question 29:

The most likely diagnosis in this case is megaloblastic anemia, given the patient's vegetarian diet, macrocytosis, and low hemoglobin level.

Megaloblastic anemia can be caused by a deficiency of vitamin B12 or folate. Vegetarians are at increased risk of vitamin B12 deficiency because vitamin B12 is primarily found in animal products. Other conditions can also cause megaloblastic anemia, such as celiac disease, Crohn's disease, and ulcerative colitis. Other causes of high MCV, such as liver disease, alcoholism, and hypothyroidism.

Iron deficiency anemia is characterized by a low hemoglobin level and a microcytic (small cell) anemia. Iron is essential for the production of hemoglobin, and a deficiency of iron can lead to a decrease in hemoglobin levels. However, IDA is typically associated with a low MCV, not a high MCV.

In Aplastic anemia the bone marrow does not produce enough progenitor cells. It can be caused by autoimmune diseases, medications, and radiation. Aplastic anemia can lead to a decrease in all types of blood cells and is associated with a low MCV and a low hemoglobin level, not a high MCV.

Thalassemia is a group of genetic blood disorders that affect the production of hemoglobin. It is characterized by a microcytic anemia. However, thalassemia is not typically associated with a high MCV.

Solution to Question 30:

Alpha 2 globulin becomes relatively more prominent in nephrotic syndrome.

Alpha 2 globulin, being a relatively large compared to others is not lost through urine as it is unable to pass through the glomeruli in nephrotic syndrome. Hence the α2 band which includes α2 macroglobulin is very prominent in serum protein electrophoresis of a nephrotic syndrome patient.

ALL proteins are lost in nephrotic syndrome and their size is relatively reducied, except alpha 2 globulin:

Albumin is the most abundant protein in the blood and is responsible for maintaining osmotic pressure. In nephrotic syndrome, albumin leaks out of the glomeruli into the urine. This leads to a decrease in serum albumin levels, but the remaining albumin molecules may become larger due to increased glycosylation.

Beta globulins are a group of proteins that play a role in transporting hormones and other molecules in the blood.

Gamma globulin: Gamma globulins are a group of proteins that play a role in immunity.

The following image shows the bands with their corresponding proteins in normal serum electrophoresis.

Certain common diseases and their electrophoretic pattern:

Condition	Electrophoretic pattern
Chronic infections	Smooth increase in gamma gl obulins
Multiple myeloma	Gamma globulin part is incre ased (M BAND)

Condition	Electrophoretic pattern
Liver cirrhosis	Reduced albumin and compensatory increase in globulin
Primary immune deficiency	Gamma globulin reduced
α1 antitrypsin deficiency	α1 band is thin and even lost sometimes

Solution to Question 31:

A seminoma would exhibit positivity for PLAP, OCT4, and KIT (Also NANOG)

• PLAP (placental alkaline phosphatase): PLAP is a germ cell tumor marker that is positive in seminomas, dysgerminomas, and embryonal carcinomas.

• OCT4 (octamer-binding transcription factor 4): OCT4 is a pluripotency marker that is positive in seminomas, dysgerminomas, and embryonal carcinomas.

• KIT (KIT proto-oncogene receptor tyrosine kinase): KIT is a tyrosine kinase receptor that is positive in seminomas and dysgerminomas.

• hCG (human chorionic gonadotropin): HCG is a pregnancy hormone that is produced by the syncytiotrophoblast cells of the placenta. HCG is sometimes positive in other types of germ cell tumors, such as choriocarcinomas and yolk sac tumors, but not typically in seminomas. About 15% of seminomas that contain syncytiotrophoblasts may be associated with a raised beta hCG but even in these patients the levels are not as high as those seen in choriocarcinoma.

• NANOG (homeobox protein NANOG): NANOG is a pluripotency marker that is positive in seminomas, dysgerminomas, and embryonal carcinomas.

Seminoma is the most common type of germ cell tumor (~50% of GCTs). The peak incidence is seen in the fourth decade. It is the testicular equivalent of an ovarian dysgerminoma, or the pineal gland tumor termed germinoma. The gross specimen shows a homogeneous, grey-white, lobulated cut surface, devoid of hemorrhage or necrosis.

Histopathological examination shows sheets of uniform cells with clear cytoplasm and central nucleus, with a prominent nucleolus (fried-egg appearance). The sheets of cells are poorly demarcated into lobules by delicate fibrous septa containing a lymphocytic infiltrate. Immunohistochemistry is positive for KIT, OCT3/4, placental alkaline phosphatase (PLAP), and podoplanin and negative for cytokeratin. Lactate dehydrogenase (LDH) levels are also measured to assess tumor burden. It usually presents as painless testicular enlargement. A heaviness may be felt if the size is 2-3 times normal. Rarely, it may present with symptoms of metastatic disease. Sonography is mandatorily done in all suspected cases.

Tumor markers which may be elevated in up to 50% of cases, should be investigated prior to orchiectomy. This serves as useful in follow-up after surgery. An orchidectomy through an inguinal incision is performed as a definitive treatment. Surveillance using CT imaging and tumor marker to look for metastasis or relapse is done. Histopathologic diagnosis also helps to plan radiotherapy or chemotherapy.

Note: Since ~85% of seminomas are not positive for HCG, option 1,2,3 is a better one to choose.

Solution to Question 32:

Gene	Cancer type	Explanation
KIT	Gastrointestinal stromal tumors (GISTs)	KIT mutations are found in about 85% of GISTs. GISTs are the most common mesenchymal tumors of the gastrointestinal tract.
KRAS	Pancreatic cancer	KRAS mutations are found in about 90% of pancreatic cancers. Pancreatic cancer is a highly aggressive cancer with a poor prognosis.
JAK 2	Myeloproliferative disorders	JAK 2 mutations are found in a variety of myeloproliferative disorders, including polycythemia vera, essential thrombocythemia, and myelofibrosis. Myeloproliferative disorders are characterized by an overproduction of blood cells.
ALK	Leukemia/myeloproliferative disorders	ALK mutations are found in a variety of leukemias and myeloproliferative disorders, including anaplastic large cell lymphoma and chronic myeloid leukemia. It is also seen in some non small cell lung cancers.

Solution to Question 33:

The cervical smear shown above is stained with Papanicolau stain. Pap stain is the most commonly used stain for cervical smears. It is a polychrome stain that stains different cell types different colors, making it easy to identify abnormal cells.

The above image shows superficial cells (arrow mark) along with other basophilic cells.

Toluidine blue stain: Toluidine blue stain is a metachromatic stain that stains acidic and basic substances different colors. It is commonly used to stain mast cells and other cells that contain acidic and basic granules.

Hematoxylin and eosin stain (H&E stain): H&E stain is a polychrome stain that stains nuclei blue and cytoplasm pink. It is commonly used to stain tissue sections for histological examination.

Giemsa stain: Giemsa stain is a polychrome stain that stains nuclei blue and cytoplasm pink. It is commonly used to stain blood smears and bone marrow aspirates.

Type of cell	Identification feature
Basal & Para Basal	Cells are small, rounded,basophilicwith large nuclei
Middle layer (Intermediate)	Cells are squamous cells, transparent andbasophilic (Blue/Green)with vesicular nuclei
Superficial layer	Cells are acidophilic (Pink), with pyknotic nuclei
Malignant cells	Hyperchromatic, nuclei vary in size, small cytoplasm in undifferentiated malignant cells

Stain	Uses
Papanicolaou stain (Pap stain)	Screening for cervical cancer and precancerous lesions, diagnosis of other gynecological conditions such as endometrial cancer and vulvar cancer, diagnosis of urinary tract infections and other conditions of the bladder and urethra
Toluidine blue stain	Diagnosis of mast cell disorders, detection of acidic and basic granules in cells, diagnosis of certain fungal infections
Hematoxylin and eosin stain (H&E stain)	Histological examination of tissue sections for a wide variety of diagnostic purposes
Giemsa stain	Staining blood smears and bone marrow aspirates for the diagnosis of blood disorders, diagnosis of parasitic infections, diagnosis of certain fungal infections

Solution to Question 34:

DAB (3,3■-diaminobenzidine) is a chromogen that is used in immunohistochemistry. It produces a brown precipitate when exposed to hydrogen peroxide and horseradish peroxidase. DAB is the most commonly used chromogen in immunohistochemistry because it produces a strong and stable color. It is also relatively inexpensive and easy to use.

In IHC, a primary antibody is applied to the tissue sample. The primary antibody binds to the protein of interest. A secondary antibody is then applied to the tissue sample. The secondary antibody binds to the primary antibody. The secondary antibody is conjugated to an enzyme, such as horseradish peroxidase. When the tissue sample is exposed to hydrogen peroxide, the enzyme catalyzes the production of a free radical. The free radical then reacts with the chromogen to produce a colored precipitate.The color of the precipitate depends on the chromogen that is used.

Other options:

FITC (fluorescein isothiocyanate): FITC is a fluorescent chromogen that is used in immunofluorescence staining and flow cytometry. It produces a green fluorescence when excited by light at 488 nm.

Cyanate is a colorless compound that can be used to conjugate antibodies to other molecules. Alkaline phosphatase is an enzyme that is used in immunohistochemistry to amplify the signal from the primary antibody. It does not produce a color by itself.

Phycoerythrin is a fluorescent chromogen that is used in flow cytometry. It produces a red fluorescence when excited by light at 561 nm.

Solution to Question 35:

The correct answer is iron deficiency anemia. Transferrin saturation is a measure of the percentage of transferrin molecules that are saturated with iron. Transferrin is a protein that transports iron in the blood - normal saturation is between 20% and 50%.In iron deficiency anemia, the body does not have enough iron to saturate all of the transferrin molecules -> decreased transferrin saturation. Note: transferrin saturation can be affected by inflammation, infection, and liver disease.

Hemosiderosis: Hemosiderosis is a condition in which there is excess iron deposition in the tissues. Transferrin saturation is typically normal/increased in hemosiderosis.

Hemochromatosis: Hemochromatosis is a condition in which the body absorbs too much iron. Transferrin saturation is typically elevated in hemochromatosis.

Pernicious anemia: Pernicious anemia is a type of anemia caused by a deficiency of vitamin B12. Transferrin saturation is typically normal in pernicious anemia.

Test	Normal ran ge	Increased in:	Decreased in:
Transferrin	200-400 m g/dL	Iron deficiency anemia	Inflammation, infection, liver disease, nephrotic syndrome
Total iron binding capacity (TIBC)	250-450 µg /dL	Iron deficiency anemia	Chronic inflammatory diseas e, anemia of chronic disease, kidney disease
Ferritin	20-250 ng/ mL	Hemochromatosis, infection, inflammation, malignancy	Iron deficiency anemia, chro nic inflammatory disease, an emia of chronic disease

Solution to Question 36:

Reinke's crystalloids are rod-shaped or corn-shaped structures that are found in Leydig cells, and seen in Leydig cell tumors. They are made of calretinin. Leydig cell tumors are usually benign, but they can sometimes be malignant.

This is the most common sex-cord stromal type of testicular neoplasm. It is mostly benign and invariably unilateral. Most cases occur between 20 and 60 years of age. However, they can present at any age. The most common presenting feature is testicular swelling or gynecomastia (~15% of cases). In children, hormonal effects manifested primarily as sexual precocity, are the dominant features. Tumors of Leydig cells may secrete androgens and estrogens, and even corticosteroids.

Grossly it shows golden brown, homogeneous circumscribed nodules <5 cm in diameter. Histologically:

• large, round, or polygonal cells

• round central nucleus and prominent nucleoli

• abundant granular eosinophilic cytoplasm containing lipid droplets, vacuoles, or lipofuscin pigment

• crystalloids of Reinke, rod-shaped eosinophilic cytoplasmic bodies are seen in about 25% of the tumors

Other options:

Option A: Gonadoblastoma is a rare mixed germ cell-sex cord stromal tumor that occurs in the gonads of patients with dysgenetic gonads.

Option B: Sertoli cell tumor is a rare tumor that arises from Sertoli cells, which are supporting cells in the testes. Sertoli cells, with a thickening of the tubular basement membrane and absence of germ cells which is characteristic of Sertoli cell-only syndrome.

Option D: Yolk sac tumor is a type of germ cell tumor that is most common in children. Schiller Duval bodies are seen in yolk sac tumors and resemble endodermal sinuses. They resemble glomerulus-like structures in the visceral and parietal layer while the central core consists of capillaries.

Solution to Question 37:

Cystic fibrosis: Cystic fibrosis is an autosomal recessive genetic disorder; both parents must carry a copy of the mutated gene in order for the child to be affected.

Leber optic neuropathy: Leber optic neuropathy is a mitochondrial genetic disorder. This means that the mutated gene is located on the mitochondrial DNA, which is DNA that is outside of the nucleus of the cell. Mitochondrial DNA is inherited solely from the mother.

Duchenne muscular dystrophy: Duchenne muscular dystrophy is an X-linked recessive genetic disorder. This means that the mutated gene is located on the X chromosome. Males have only one X chromosome, so if they inherit a mutated copy of the gene, they will be affected. Females have

two X chromosomes, so if they inherit one mutated copy of the gene, they will be carriers.

Myotonic dystrophy: Myotonic dystrophy is an autosomal dominant genetic disorder. This means that only one parent needs to carry a copy of the mutated gene in order for their child to be affected.

Genetic disorder	Inheritance patter n
A. Cystic fibrosis	Autosomal recessi ve
B. Leber optic neuropathy	Mitochondrial
C. Duchenne muscular dystro phy	X-linked recessive
D. Myotonic dystrophy	Autosomal domin ant

Solution to Question 38:

The correct answer is pernicious anemia that can have atrophic gastritis.

Pentagastrin is a synthetic hormone that stimulates the release of gastrin from the G cells of the stomach. Gastrin stimulates gastric acid secretion. In pernicious anemia there are:

• Antibodies to parietal cells and intrinsic factor

• Reduced serum pepsinogen I concentration

• Hypergastrinemia

• Endocrine cell hyperplasia → Carcinoid tumors

• Vitamin B12 deficiency (pernicious anemia)

• Defective gastric acid secretion (achlorhydria)

After the injection of pentagastrin, patients with atrophic gastritis/pernicious anemia will typically show a weak or absent gastric acid response due to destruction of the parietal cells.

Other options:

Normal healthy individual: A normal healthy individual will show gastric H+ secretion after the injection of pentagastrin. This is because gastrin is a potent stimulator of gastric acid secretion.

Gastrinoma: A gastrinoma is a tumor of the G cells of the stomach. Gastrinomas produce excess gastrin, which leads to hypergastrinemia and increased gastric acid secretion. Therefore, patients with gastrinomas will show a vigorous gastric acid response after the injection of pentagastrin.

Duodenal ulcer: Duodenal ulcers are caused by an imbalance between gastric acid secretion and mucosal defense mechanisms. Duodenal ulcer patients typically have normal or increased gastric acid secretion. Therefore, duodenal ulcer patients will typically show a vigorous gastric acid response after the injection of pentagastrin.

Solution to Question 39:

Correct way to collect sample is 3 sets of culture samples, 2 from the peripheral lines and 1 from the affected valve with a gap of 30 minutes between the collection of each

The American Heart Association recommends collecting three sets of blood culture samples for patients with suspected infective endocarditis. One set should be collected from the affected valve, if possible. The other two sets should be collected from peripheral veins. There should be a gap of at least 30 minutes between the collection of each set.

Solution to Question 40:

The lymphocytic predominance and low CSF glucose points to a diagnosis of tubercular meningitis.

Note: CSF glucose/Serum glucose < 0.4 -> Hypoglycorrhachia; Secondary to glycolysis by bacteria or leukocytes, and/or by altered glucose transport.

Tuberculous meningitis caused by Mycobacterium tuberculosis is caused by the formation of tubercles in the brain parenchyma. This happens during the hematogenous spread of the bacilli during primary infection. It can also be due to the rupture of a subependymal tubercle into the subarachnoid space.

Tuberculous meningitis presents as a subacute infection and has a longer course than bacterial meningitis. Ocular nerve palsy is a common association as there is marked meningeal involvement at the base of the brain. The patient can also develop hydrocephalus.

The CSF sample shows a pellicle or cobweb-like clots. Acid-fast bacilli can be best isolated from these. There is usually lymphocytic leukocytosis, low glucose concentration, and elevated proteins. CSF culture remains the gold standard for diagnosis.

The clinical triad of fever, headache, and nuchal rigidity is common to all types of meningitis. Nuchal rigidity is the resistance of the neck to passive flexion and is pathognomonic of meningeal irritation. Kernig's and Brudzinski's signs are other features of meningeal irritation. Raised intracranial pressure (ICP) is a complication that presents as papilledema, disorientation, or dilated poorly reactive pupils.

Important features of TB meningitis:

• Common between 6 months and 4 years of age

• Usually spread by hematogenous route

• Gelatinous exudate infiltrates corticomeningeal blood vessels leading to hydrocephalus.

• Three stages

• Prodromal stage

• stage of meningitis

• stage of coma

• CSF study shows reduced sugar level, increased protein level, and lymphocytosis.

• Treatment includes antitubercular drugs for 12 months and corticosteroids (to reduce neurologic sequelae).

Other options:

Coxsackie meningitis is a viral meningitis. (aseptic meningitis) The CSF findings in Coxsackie meningitis are typically normal-elevated protein and lymphocytic pleocytosis. The glucose level in the CSF is usually normal.

The CSF findings in HSV encephalitis are typically normal-elevated protein and lymphocytic pleocytosis, and RBCs due to hemorrhagic necrosis of the temporal lobe. CSF Glucose levels would be normal.

The CSF findings in bacterial meningitis are typically elevated protein, neutrophilic pleocytosis and low glucose level.

Solution to Question 41:

The protease of Vibrio cholera disrupts the extracellular part of occludin.

Occludin is a transmembrane protein that is found in tight junctions; prevents the leakage of fluids and molecules across the epithelium. The protease of V. cholera cleaves occludin, which disrupts the tight junctions and allows for the leakage of fluids and electrolytes into the intestinal lumen. This leads to the profuse diarrhea that is characteristic of cholera.

Vibrio cholerae in addition to the cholera toxin (CT) produces some additional toxins such as,

• Zonula occludens toxin, which alters the intestinal permeability by acting on intestinal epithelial cell tight junctions (which are also called zonula occludens)

• Accessory cholera enterotoxin.

Zonula occludens:

• Also known as tight junctions

• Characteristically surround the apical margins of the cells in epithelia such as the intestinal mucosa, the walls of the renal tubules, and the choroid plexus.

• Also important to the endothelial barrier function and endothelium-dependent vasodilation.

• Made up of ridges- half from one cell and a half from another cell- which adhere so strongly that there is no space between the cells.

• Three families of transmembrane proteins are involved:

Other options:

Integrin: transmembrane proteins that mediate cell adhesion to the extracellular matrix. They are not involved in tight junctions.

Cadherin: transmembrane proteins that mediate cell adhesion to other cells. They are not involved in tight junctions.

Claudins are transmembrane proteins that are also found in tight junctions. However, the protease of V. cholera does not cleave claudins.

Protein	Function	Disease association
Integrin	Cell adhesion and migration	Leukocyte adhesion deficiency, Glanzmann thrombasthenia, muscular dystrophies
Cadherin	Cell-cell adhesion	E-cadherin mutations: gastric cancer
Occludin	Tight junction protein	Inflammatory bowel disease, celiac disease, Vibrio cholerae infection
Claudin	Tight junction protein	Claudin-1 mutations: familial hypomagnesemia with hypercalciuria, nephrocalcinosis syndrome

Pathology INI-CET 2024

Question 1:

Which of the following is not a marker for gastrointestinal stromal tumors?

a) KIT

b) SDH

c) PDGFRA

d) NRG-1

Question 2:

Identify the given histopathological image of a carcinoma of the lung.



a) Adenocarcinoma

b) Small cell carcinoma

c) Squamous cell carcinoma

d) Sarcoma

Question 3:

Aneurysm due to excessive degradation of connective tissue is seen in which of these disorders?

a) Marfan's syndrome

b) Ehler Danlos syndrome

c) Scurvy

d) Atherosclerosis

Question 4:

A known case of hemophilia-B comes with pain and swelling of the knee joint. Which of the following will be used in his management?

a) i, ii

b) ii, iv

c) i, iii

d) ii, iii, iv

Question 5:

A patient presents with truncal obesity, abominal striae, poor wound healing. What is the likely cause of his symptoms?

a) Hypercortisolism

b) Addison's disease

c) Hyperestrogenism

d) Increased growth hormone

Question 6:

Match the following blood products with their corresponding shelf life.

a) 1-B, 2-A, 3-D, 4-C

b) 1-D, 2-C, 3-B, 4-A

c) 1-A, 2-C, 3-B, 4-D

d) 1-C, 2-D, 3-B, 4-A

Question 7:

Which of the following statements about Hodgkin's Lymphoma is not true?

a) Waldeyer's ring is the most common site to be involved

b) RS cells are immunopositive for CD30 and PAX5

c) RS Cells are of B cell origin

d) Nodular sclerosis is the most common subtype and presents as a mediastinal mass

Question 8:

A patient presents to the ER with anaphylaxis. Which of the following is a part of the clinical criteria to diagnose anaphylaxis?

a) 2,3,4,5

b) 1,2,3,4

c) 1,2,3,5

d) 3,4,5

Question 9:

A patient presents with a neck swelling. An excision biopsy is performed and is shown below. What is the likely diagnosis?



a) Tuberculosis

b) Sarcoidosis

c) Langerhans cell histiocytosis

d) Hodgkin's lymphoma

Question 10:

A patient was brought to the emergency following a heavy bleed and was transfused with multiple units of blood. Which of the following is not seen as a result of delayed hemolytic transfusion reaction?

a) Cola coloured urine in the first 2 hours

b) Failure of rise in Hemoglobin level

c) Increased Indirect Bilirubin

d) Increased LDH

Question 11:

Given below is the image of a survival analysis curve in patients with organ transplants. Which of the following statements are true about the graph?



a) Both CD4 and CD8 cells take part in allograft rejection

b) Both CD4 and CD8 participate in the protection against allograft rejection

c) CD4 cells are not involved in allograft rejection

d) CD8 cells have a predominant role in graft rejection

Question 12:

A patient was brought to the emergency care following severe hemorrhage and was unable to get the matching blood type. Which blood group can be used for immediate transfusion?

a) 1,2,3,4

b) 2,4

c) 3,4

d) 1,2

Question 13:

Which of the following does not differentiate between necrosis and apoptosis?

a) Intact cell contents

b) Altered cell shape

c) Intact plasma membrane

d) Lysosomal enzymatic digestion

Question 14:

In comparison to Ulcerative colitis, Crohn's disease is associated with

a) Continuous lesions rather than segmented lesions

b) Less likely to present with daily hematochezia

c) Increased risk of developing cancer

d) Lower risk of perianal fistulas

Question 15:

Match the following organisms with their association :

a) 1 - a ; 2 - b ; 3 - c ; 4 - d

b) 1 - c ; 2 -d ; 3 - a ; 4 - b

c) 1 - c ; 2 -a ; 3 - d ; 4 - b

d) 1 - d ; 2 -c ; 3 - a ; 4 - b

Question 16:

Which of the following is a nuclear neuroendocrine immunohistochemical marker?

a) Synaptophysin

b) NCAM 1

c) INSM1

d) Chromogranin

Question 17:

Which of the following is incorrect regarding myelodysplastic syndrome?

a) More commonly seen in females

b) Treatment induced MDS is not dependent on age

c) Median age of onset 70 years

d) It can be radiation induced

Question 18:

Which among the following is not a growth-promoting oncogene?

a) FGF

b) PDGF

c) TGF beta

d) TGF alpha

Question 19:

Which vacutainer is used in electrolyte estimation?

a) Green

b) Lavender

c) Yellow

d) Gray

Question 20:

Match the following with appropriate disorder.

a) i-b, ii-c, iii-a, iv-d

b) i-d, ii-c, iii-b, iv-a

c) i-b, ii-a, iii-c, iv-d

d) i-a, ii-b, iii-d, iv-c

Question 21:

Match the following

a) a-i, b-ii, c-iii, d-iv

b) a-iii, b-ii, c-iv, d-i

c) a-ii, b-iii, c-iv, d-i

d) a-iv, b-iii, c-ii, d-i

Question 22:

Epithelial-mesenchymal transition induced by which pair of transcription factors?

a) TF IIA & TF IIB

b) Cathepsin & MMP8

c) SNAIL & TWIST

d) STAT-5 & HIF- 1alpha

Question 23:

A young child presents with bleeding gums and bruising for two weeks. Bone marrow biopsy and flow cytometry were done. What is the diagnosis?



a) AML

b) CLL

c) ALL: B Phenotype

d) ALL: T phenotype

Question 24:

A patient presents with a lesion on the hand. Histopathology of the lesion is given. Which of the following is incorrect?



a) The lesion arises from cranial nerve

b) Verocay bodies are seen in light microscopy

c) Associated with neurofibromatosis

d) The lesion is malignant and metastasises frequently

Question 25:

A 30-year-old female presented with weight loss and a cough. On investigation, it was found that she had pleural effusion, lymphadenopathy, and tree-in-bud appearance on the CT scan. On aspiration of the pleural fluid sample, adenosine deaminase was low, lymphocytes were high. What would be the finding in the aspirate of the mediastinal lymph node?

a) Necrotized epithelioid giant cells with necrosis

b) Reactive lymphoid hyperplasia

c) Fungal hyphae surrounded by lymphocytes

d) Anthracotic pigment

Question 26:

Match the following:

a) a-iv, b-ii, c-iii, d-i

b) a-iii, b-i, c-iv, d-ii

c) a-i,b-iv,c-ii,d-iii

d) a-ii, b-iii, c-i, d-iv

Question 27:

All are hereditary causes of thrombosis except?

a) PNH

b) Protein S deficiency

c) Antithrombin deficiency

d) Homocystinuria

Question 28:

Which anticoagulant is used in the test tube for estimating Hba1c?

a) Warfarin

b) Sodium Fluoride

c) EDTA

d) Heparin

Question 29:

Most common cause of inherited coagulopathy

a) Factor V leiden mutation

b) Antithrombin III deficiency

c) Protein C deficiency

d) Prothrombin G20210A

Question 30:

Match the Following diseases with their respective gene mutation

a) a-i, b-ii, c-iii, d-iv

b) a-ii, b-iii, c-i, d-iv

c) a-iii, b-i, c-iv, d-ii

d) a-iv, b-ii, c-iii, d-i

Question 31:

Which of the following is not a component of MEN2B syndrome?

a) Medullary thyroid carcinoma

b) Marfanoid habitus

c) Mucosal ganglioneuromatosis

d) Parathyroid hyperplasia

Answer Key

Question No.	Correct Option
1	d
2	a
3	a
4	b
5	a
6	b
7	a
8	a
9	a
10	a
11	a
12	c
13	b
14	b
15	b
16	c
17	a
18	c
19	a
20	b
21	b
22	c
23	c
24	d
25	a
26	b
27	a
28	c
29	a
30	c
31	d

Detailed Explanations

Solution to Question 1:

NRG-1 (Neuregulin-1) is not a marker for gastrointestinal stromal tumors (GISTs).

GIST are the most common mesenchymal tumor of the abdomen, the stomach being the most common site. They arise from the interstitial cells of Cajal, the pacemaker cells of the GIT. GIST is seen in the 5th to 6th decade of life and has an equal preponderance among males and females. Most patients present with haematemesis, melena, or vague abdominal pain. The most common indications for surgery include bleeding and obstruction.

The GIST may occur in a sporadic form or as part of familial syndromes. In sporadic GIST, the most common mutations seen are gain-of-function mutation in KIT (75-80%) and PDGFRA mutation (8%). Microscopically, GIST has elongated spindle cells or plumper epithelioid cells, or a mixed pattern.

The below image shows the microscopy of GIST composed of bundles, or fascicles, of spindle-shaped tumor cells.



Succinate dehydrogenase (SDH)-deficient GIST is a subset of wild-type GIST seen with Carney triad or Carney-Stratakis syndrome.

Imatinib can be used to treat GIST with mutations in KIT or PDGFRA. Succinate dehydrogenase (SDH) - deficient GIST is imatinib-resistant, so sunitinib, sorafenib, or regorafenib can be used instead.

During surgical resection of GIST, care must be taken to prevent capsule rupture, which causes seeding of nearby tissues leading to relapse in 100% of the cases. If capsule rupture occurs, adjuvant therapy with imatinib is indicated regardless of tumor extent.

Other options:

Option A: KIT (CD117): This is the most well-known marker for GISTs. About 85% of GISTs show mutations in the KIT gene, leading to the expression of the KIT protein on the cell surface. KIT is a receptor tyrosine kinase, and its activation is a key driver in the pathogenesis of GISTs.

Option B: SDH (Succinate Dehydrogenase): SDH mutations are associated with a subset of GISTs, particularly those that are KIT and PDGFRA wild-type. Loss of SDH function is characteristic of this subgroup, and SDH deficiency can be demonstrated through immunohistochemistry.

Option C: PDGFRA (Platelet-Derived Growth Factor Receptor Alpha): Mutations in the PDGFRA gene account for approximately 5-10% of GISTs. Similar to KIT, PDGFRA is a receptor tyrosine kinase, and mutations in this gene can lead to the development of GISTs.

Solution to Question 2:

The microscopic image of the tissue shows an increased mucinous type of cells and its characteristic growth along the pre-existing alveolar septa indicating adenocarcinoma of the lung.

Smoking is a major risk factor for developing lung cancer. Other risk factors that can contribute are industrial chemical exposure, air pollution, etc.

They are broadly divided into

• Small cell lung cancer

• Non-small cell lung cancerAdenocarcinoma

• Squamous cell carcinoma

• Large cell carcinoma

Though adenocarcinoma is the most common tumor of the lung in non-smokers, it can be seen in smokers as well. It is also more common in women. Adenocarcinoma of the lung occurs in the periphery of the lung, frequently in areas of chronic scarring caused by chronic inflammatory changes.

Clinical features include cough, chest pain, hemoptysis, and symptoms due to metastasis. Some lung cancers are responsible for paraneoplastic syndromes.

Other options:

Option B: Small cell lung carcinoma is usually identified by small, round cells that lack glandular formation. It is associated with smoking and is known for its aggressive behavior and rapid growth.



Option C: Squamous cell carcinoma originates in the squamous cells that line the airways and typically shows keratinization and the formation of intercellular bridges, rather than glandular structures.



Option D: A sarcoma arises from connective tissues and not from epithelial cells, so it would not show glandular structures on histological examination.

Solution to Question 3:

Aneurysm due to excessive degradation of connective tissue is seen in Marfan's syndrome.

Marfan's syndrome arises from a mutation in the FBN1 gene, responsible for encoding fibrillin, a crucial protein essential for maintaining the structural integrity of connective tissues and regulating TGF-β signaling. The condition primarily affects the skeleton, eyes, and cardiovascular system. Clinical manifestations commonly include tall stature, elongated fingers, bilateral lens subluxation, mitral valve prolapse, aortic aneurysm, and potentially life-threatening aortic dissection.

Cardiovascular complications pose the greatest threat to this disorder. The two primary concerns are mitral valve prolapse and dilation of the ascending aorta due to cystic medial necrosis. Progressive dilation of the aortic root and valve ring results from the loss of medial support, leading to severe aortic incompetence. Additionally, medial weakening increases the risk of intimal tears, which can initiate intramural hematomas. These hematomas may cleave through layers of the aortic media, potentially causing aortic dissection. In some cases, the hemorrhage extends extensively, from the aortic root to the iliac arteries and may ultimately rupture through the aortic wall.

Solution to Question 4:

For Hemophilia B, fresh frozen plasma and cryo-poor plasma can be used.

Hemophilia B, also known as Christmas Disease, occurs due to a deficiency of Factor IX, also called Christmas Factor, named after the first patient identified with the disease.

Fresh Frozen Plasma (FFP)contains all coagulation factors, including factor IX, and can be used to treat Hemophilia B, and cryo poor plasma is the plasma remaining after the cryoprecipitate has been removed, and it still contains factor IX, making it useful for treating Hemophilia B.

Hemophilia is a coagulation disorder occurring due to a deficiency of clotting factor VIII (Hemophilia A) or clotting factor IX (Hemophilia B), both of which present in the same manner. The disorder is X-linked recessive, meaning males are affected, and females are asymptomatic carriers.

Patients present with ecchymoses, epistaxis, haematuria, menorrhagia, prolonged bleeding after dental or minor surgical procedures, haemarthrosis. However, Factor IX assays are more specific to diagnose Hemophilia B.

Definitive treatment is by infusions of recombinant Factor IX.

Other options:

Factor VIII: This is used in Hemophilia A, not Hemophilia B, because Hemophilia A is caused by a deficiency of factor VIII.

Cryoprecipitate: This is used primarily for Hemophilia A and von Willebrand disease as it is rich in factor VIII and fibrinogen, but it does not contain significant amounts of factor IX, making it unsuitable for Hemophilia B.

Solution to Question 5:

This patient with abdominal striae, truncal obesity, and poor wound healing most likely has Cushing's syndrome. This is characterized by hypercortisolism.

Cushing's syndrome is a group of clinical features that occur due to chronic exposure to excess glucocorticoids of any etiology. Cushing's disease is specifically Cushing's syndrome occurs due to pituitary corticotrope adenoma.

Management of Cushing's syndrome includes: Surgical removal of the tumor followed by hydrocortisone replacement with gradual tapering is preferred.

Medical management includes metyrapone and ketoconazole.

System	Clinical features
Body fat	Centralor truncalobesity(most common)Buffalo hump(fat deposition behind the neck) Moonfacies
Skin	Purplestriae (over abdomen & flanks)Acneand Hirsutism Easy bruisingFacial plethora Hyperpigmentation
Musculoskeletal	Proximal myopathyOsteoporosis

System	Clinical features
Metabolic	Poorly controlled hyperglycemiaDyslipidemia
Reproductive	Low libidoAmenorrhea
Renal	Hypokalemic alkalosis
CVS	AtherosclerosisDiastolic hypertensionEdema
Hematologic	Hypercoagulable stateNeutrophiliaEosinopenia
CNS	Depression, anxietyPsychosis Irritability

Solution to Question 6:

The correct answer is option B 1-D, 2-C, 3-B, 4-A

1. Packed RBC in additive solution	D. 42 days
2. Granulocytes	C. 24 hours
3. Single donor platelet solution	B. 5 days
4. Cryoprecipitate	A. 1 year

Solution to Question 7:

The statement that Waldeyer's ring is the most common site involved in Hodgkin's Lymphoma is incorrect; the most commonly affected sites are the cervical and supraclavicular lymph nodes.

The slide below shows the classic appearance of the RS cells showing the owl's eye appearance.



WHO classifies Hodgkin's lymphoma into 5 subtypes:

• Nodular sclerosis: Frequent lacunar cells

• Mixed cellularity

• Lymphocyte-rich

• Lymphocyte depletion

• Nodular lymphocyte predominance

The first 4 subtypes are considered classical Hodgkin's lymphoma. The RS cells are CD15+, PAX5 and CD30+. The lymphocyte predominant type is considered non-classical Hodgkin's lymphoma with CD15-, CD30-, CD20+, and EB -.

Other options:

Option B: RS cells are immunopositive for CD30 and PAX5: Reed-Sternberg cells typically express CD30 and PAX5, among other markers. These immunohistochemical markers help in confirming the diagnosis of Hodgkin's Lymphoma.

Option C: RS Cells are of B cell origin: Despite their atypical appearance, RS cells are derived from B lymphocytes, specifically from germinal center B cells.

Option D: Nodular sclerosis is the most commonly present as a mediastinal mass: Nodular sclerosis is the most common subtype of Hodgkin's lymphoma and often presents with a mediastinal mass, particularly in younger patients.

Solution to Question 8:

Anaphylaxis is characterised by all of the above features except for hypertension. It causes hypotension.

Anaphylaxis is a severe, potentially life-threatening allergic reaction that occurs rapidly and affects the whole body. It is characterized by a sudden release of histamines and other chemicals

from the immune system in response to an allergen, such as certain foods, medications, insect stings, or latex. The onset of symptoms can occur within minutes to a few hours after exposure to the allergen.

Clinically, anaphylaxis can present with a combination of symptoms that typically include:

• Respiratory Symptoms: Patients often experience wheezing and respiratory distress due to bronchoconstriction. Other signs may include shortness of breath, coughing, and a feeling of tightness in the chest.

• Skin Symptoms: Urticaria (hives) and angioedema (swelling of the skin and mucous membranes) are common. These skin reactions usually develop rapidly, often within minutes to two hours of exposure to the allergen.

• Cardiovascular Symptoms: Hypotension (low blood pressure) is a key indicator of anaphylaxis. Patients may also experience tachycardia (increased heart rate) and, in severe cases, shock.

• Gastrointestinal Symptoms: Abdominal pain, vomiting, diarrhea, and cramps can occur, particularly if the anaphylaxis is triggered by food allergens.

• Neurological Symptoms: Dizziness, lightheadedness, or loss of consciousness can occur due to the sudden drop in blood pressure and reduced oxygen supply to the brain.

Specific diagnostic criteria for anaphylaxis include the presence of acute onset symptoms involving the skin, mucosal tissue, or both (e.g., generalized hives, pruritus, or flushing, swollen lips-tongue-uvula), plus respiratory compromise (e.g., dyspnea, wheezing-bronchospasm, stridor, reduced peak expiratory flow), or hypotension or associated symptoms of end-organ dysfunction (e.g., hypotonia, collapse, syncope, incontinence). Alternatively, it can be diagnosed if two or more of the following occur rapidly after exposure to a likely allergen: involvement of the skin-mucosal tissue, respiratory compromise, hypotension, or persistent gastrointestinal symptoms.

Prompt treatment is critical in managing anaphylaxis. The first-line treatment is the immediate administration of intramuscular epinephrine, which counteracts the effects of histamines and other chemicals released during the reaction. Additional treatments may include antihistamines, corticosteroids, and bronchodilators to manage symptoms and prevent further complications.

Patients with a history of anaphylaxis are typically advised to carry an epinephrine auto-injector at all times and to seek emergency medical care immediately if they experience symptoms of anaphylaxis. Long-term management may involve identifying and avoiding known allergens, as well as potentially undergoing allergen immunotherapy in certain cases.

Solution to Question 9:

The given clinical scenario and the slide showing granulomatous inflammation with central caseous necrosis are suggestive of tuberculosis of cervical lymph nodes.

Once Mycobacterium tuberculosis enters the macrophage, it inhibits the formation of the phagolysosome. This allows it to replicate unchecked. M. tuberculosis recruits a host protein coronin to block the fusion of phagosome and lysosome. Th1 response to M. tuberculosis is mounted about three weeks following the infection. The antigen-presenting cells (APCs) present mycobacterial antigens to the T-cells in the lymph nodes and release IL-12 and IL-18 to cause the differentiation of T-cells into Th1 cells. Th1 cells produce IFN-γ. IFN-γ is the critical step in the

body's defense against M. tuberculosis. It acts by activating macrophages, causing phagolysosome maturation, and inducing nitric oxide synthase enzyme to form more reactive nitrogen species. It also mobilizes anti-bacterial compounds like defensins and stimulates autophagy.

Th1 response leads to the formation of a granuloma and caseous necrosis. Macrophages that are activated by the action of IFN-γ differentiate into epithelioid histiocytes. These are the cells that actually aggregate to form a granuloma. Some epithelioid cells may also fuse to form a giant cell.

Other options:

Option B: Sarcoidosis: This granulomatous disease can involve lymph nodes but typically does not produce caseous necrosis. Non-caseating granulomas are more characteristic of sarcoidosis.

Option C: Langerhans Cell Histiocytosis: This rare disorder involves the proliferation of Langerhans cells and can affect lymph nodes, but it does not typically show caseous necrosis.

Option D: Hodgkin's Lymphoma: This type of lymphoma often presents with lymph node enlargement and may show Reed-Sternberg cells on biopsy. Necrosis can be seen, but it is not typically caseous in nature.

Solution to Question 10:

Cola-colored urine in the first 2 hours is not a feature of delayed transfusion hemolytic reaction. This is characteristic of an acute hemolytic transfusion reaction. Acute reactions present immediately or within a few hours of transfusion with symptoms like dark urine, fever, chills, and back pain.

Delayed hemolytic transfusion reactions (DHTR) occur days to weeks after a blood transfusion, unlike acute reactions that manifest immediately or within a few hours. In DHTR, the recipient's immune system recognizes the transfused red blood cells as foreign and mounts an immune response against them, leading to their destruction. This delayed immune response is often due to the formation of alloantibodies against minor antigens on the donor red blood cells. Clinically, DHTR presents with symptoms such as a failure of rise in hemoglobin levels despite transfusion, increased levels of indirect bilirubin, and elevated lactate dehydrogenase (LDH) levels, indicative of ongoing hemolysis. While DHTR is generally less severe than acute reactions, it can still cause significant morbidity and necessitates careful monitoring and management.

Solution to Question 11:

From the given graph is it evident that with anti CD8 the increase in graft survival is negligible compared to the control, however with anti CD4 the graft survival doubles. When anti-CD4 and anti-CD8 are coadministered, the survival becomes four times that of control. From this, it can be inferred that both CD4 and CD8 cells take part in allograft rejection.

Rejection is a process in which T lymphocytes and antibodies produced against graft antigens react against and destroy tissue grafts. The major antigenic differences between a donor and recipient that result in the rejection of transplants are differences in HLA alleles. Following transplantation, the recipient’s T cells recognize donor HLA antigens from the graft by two pathways. The graft antigens are either presented directly to recipient T cells by graft

antigen-presenting cells (APCs), or the graft antigens are picked up by host APCs, processed, and presented to host T cells. These are called the direct and indirect pathways of recognition of alloantigens. Both lead to the activation of CD8+ T cells, which develop into cytotoxic T lymphocytes and CD4+ T cells, which become cytokine-producing effector cells, mainly Th1 cells.

Graft rejection is classified into hyperacute, acute, and chronic, based on clinical and pathologic features :

• Hyperacute rejection is mediated by preformed antibodies specific for antigens on graft endothelial cells.

• Acute rejection is mediated by T cells and antibodies that are activated by alloantigens in the graft.

• Chronic rejection is an indolent form of graft damage that occurs over months or years, leading to progressive loss of graft function.

Methods of increasing graft survival :

In kidney transplants, there is substantial benefit if all the polymorphic HLA alleles are matched. However, HLA matching is usually not done for transplants of the liver, heart, and lungs.

Except for identical twins, immunosuppressive therapy is essential in all donor-recipient combinations. Immunosuppressive drugs in current use include steroids (which reduce inflammation), mycophenolate mofetil (which inhibits lymphocyte proliferation), and tacrolimus (inhibitor of calcineurin).

Solution to Question 12:

In this patient who was unable to get matched donor blood, O+ve and O-ve blood types can be administered to him.

In cases of severe hemorrhage where there is an immediate need for a blood transfusion but the matching blood type is not available, both O-negative and O-positive blood can be used for immediate transfusion, with specific considerations. O-negative blood is the universal donor type for all patients because it lacks A, B, and Rh antigens on the surface of its red blood cells, minimizing the risk of an adverse reaction in the recipient. This makes O-negative blood particularly valuable in emergency situations where the patient's blood type is unknown.

O-positive blood, on the other hand, can also be used in emergencies but is generally preferred for transfusing Rh-positive patients to conserve the limited supply of O-negative blood for those who are Rh-negative or when the blood type is entirely unknown. While O-positive blood is not universally compatible with O-negative, it is still a safe option for patients who are known to be Rh-positive, thus preventing an Rh incompatibility reaction.

Blood types A+ve and B+ve are not suitable for emergency universal transfusion because they contain A and B antigens, respectively, which can cause severe transfusion reactions if the recipient has antibodies against these antigens. For example, if a patient with type B blood (who has anti-A antibodies) receives type A+ve blood, it can lead to a dangerous immune response. Similarly, a patient with type A blood (who has anti-B antibodies) cannot safely receive type B+ve blood.

Solution to Question 13:

Necrosis and apoptosis cannot be differentiated based on the altered cell shape. They can however be differentiated based on cell size, where necrosis is associated with an enlarged cell and apoptosis with cell shrinkage.

Necrosis is a pathologic process that is the consequence of severe injury. Necrosis is characterized by denaturation of cellular proteins, leakage of cellular contents through damaged membranes, local inflammation, and enzymatic digestion. It is characterized by denaturation of cellular proteins, leakage of cellular contents through damaged membranes, local inflammation, and enzymatic digestion.

Apoptosis is a type of cell death that is induced by a tightly regulated suicide program in which cells destined to die activate intrinsic enzymes called caspases that degrade the cells' genomic DNA and nuclear and cytoplasmic proteins. Apoptotic cells break up into plasma membrane-bound fragments, called apoptotic bodies, which contain portions of the cytoplasm and nucleus. While the plasma membrane remains intact, its surface components are altered. As a result, the dead cell and its fragments are rapidly devoured, before the contents leak out, and therefore apoptosis does not elicit an inflammatory reaction.

Other options :

Option 1 - Apoptosis is associated with intact cellular contents, while necrosis is associated with enzymatic digestion, and contents leak out.

Option 3 - Intact plasma membrane is a feature of apoptosis; however, in necrosis, there is a disrupted plasma membrane.

Option 4 - Lysosomal enzymatic digestion is a feature of necrosis and not apoptosis.

Feature	Necrosis	Apoptosis
Cell size	Cells swell and enlarge.	Cells shrink in size.
Nucleus	Undergoes condensation, fragmentation, and dissolution.	Breaks into smaller fragments.
Membrane	Membrane integrity is lost.	Remains intact but lipid orientation may change
Contents	Cell contents are enzymatically digested and may leak out.	Contents remain intact and may be released in apoptotic bodies.
Inflammation	Often accompanied by adjacent inflammation.	No adjacent inflammation typically occurs.
Role	Typically pathological, marking irreversible cell injury.	Oftenphysiological,used to eliminate unwanted cells.

Solution to Question 14:

Patients with Crohn's disease are less likely to present with daily hematochezia compared to patients with ulcerative colitis.

CD is characterized by acute or chronic inflammation of the bowel. The most commonly involved site is the terminal ileum because of which patients present with diarrhea and right lower quadrant pain. Extensive inflammation can result in malabsorption and steatorrhea.

Blood investigations may reveal anemia, elevated ESR and CRP. Fecal calprotectin and lactoferrin levels are used to rule out irritable bowel syndrome and as a marker of ileo-colonic inflammation. A colonoscopy is done as part of the evaluation. It shows characteristic skip lesions and a cobblestone appearance. Biopsy shows transmural inflammation with non-caseating granulomas.

Serological markers that are usually positive in patients with CD include the anti–Saccharomyces cerevisiae antibody (ASCA). Other positive markers (however, with low sensitivity) include antibodies to bacterial proteins (OmpC and I2), flagellin, and bacterial carbohydrates.

Solution to Question 15:

The correct answer is option B: 1 - c ; 2 -d ; 3 - a ; 4 - b.

MALT (mucosa-associated lymphoid tumor) lymphoma is an extranodal marginal zone lymphoma, most commonly seen in the stomach, but other common sites include the skin, salivary glands, small bowel, ocular adnexa, and synovium. It is associated with states of chronic inflammation due to either autoimmune diseases (Sjo■gren's syndrome or Hashimoto's thyroiditis) or chronic infections with organisms like H. pylori (gastric), Borrelia burgdorferi (skin), C. psittaci (conjunctiva), C. jejuni (intestines), and hepatitis C virus.

The essential pathologic feature of MALT lymphoma is the presence of lymphoepithelial lesions, which result from the invasion of mucosal glands and crypts by the neoplastic lymphocytes. These cells are positive for CD19, CD20, and CD79a and negative for CD5 and CD10. Recurrent cytogenetic abnormalities include t(11;18), t(14;18), t(1;14), t(3;14), and trisomy 8. The t(11;18) is most common, occurring in up to 50% of MALT lymphomas.

Unlike other indolent B-cell lymphomas, MALT lymphomas present most commonly with stage I or II disease. In these cases, radiation therapy may be curative. Alternatively, patients may respond to antibiotics for the associated underlying infection.

Alpha heavy chain disease / immunoproliferative small intestinal disease (IPSID) is recognized as an infectious pathogen-associated human lymphoma associated with Campylobacter jejuni. It involves mainly the proximal small intestine, resulting in malabsorption, diarrhea, and abdominal pain. IPSID is associated with excessive plasma cell differentiation and produces truncated alpha heavy chain proteins lacking the light chains. Early-stage IPSID responds to antibiotics (30–70% complete remission). Most untreated IPSID patients progress to lymphoplasmacytic and immunoblastic lymphoma.

1) H. pylori	c)Gastric MALT
2) C.psittaci	d)Ocular MALT
3) Borrelia burgdorferi	a)Cutaneous MALT

1) H. pylori	c)Gastric MALT
4) Campylobacter jejuni	b)Alpha heavy chain disease

Solution to Question 16:

INSM1 is a nuclear marker for neuroendocrine tumours.

Insulinoma-associated protein 1 is a nuclear protein that serves as a transcription factor and has gained prominence as a neuroendocrine marker in pathology. Originally identified in insulinomas, INSM1 has been to be a reliable marker for various neuroendocrine tumors (NETs).

Other options:

Option A: Synaptophysin is a glycoprotein located in the membrane of presynaptic vesicles. It is a marker for neuroendocrine cells and tumors, but it is found in the cytoplasm, not the nucleus.

Option B: NCAM 1 (Neural Cell Adhesion Molecule 1), also known as CD56, this marker is a cell surface glycoprotein involved in cell adhesion. It is often used in the diagnosis of neuroendocrine tumors, but it is not a nuclear marker.

Option D: Chromogranin A is a protein found in the secretory vesicles of neuroendocrine cells. It is another commonly used marker for neuroendocrine tumors and is located in the cytoplasm.

Solution to Question 17:

Myelodysplastic syndrome (MDS) is more commonly seen in males than females.

The MDS is a heterogeneous group of hematologic disorders characterized by both cytopenias due to bone marrow failure and a high risk of development of acute myeloid leukemia. Anemia due to ineffective erythropoiesis, often with thrombocytopenia and neutropenia, is seen.

MDS is a disease of the elderly; the mean age at onset is older than 70 years (option C). There is a slight male predominance (option A). Secondary or therapy-related MDS is not age-related (option B). It occurs as a late toxicity of cancer treatment; radiation (option D) and the radiomimetic alkylating agents such as busulfan, nitrosourea, or procarbazine (with a latent period of 5–7 years); or the DNA topoisomerase inhibitors (2-year latency). Acquired aplastic anemia, Fanconi anemia, and other constitutional marrow failure diseases can evolve into MDS.

Anemia dominates the early course. Most symptomatic patients complain of the gradual onset of fatigue and weakness, dyspnea, and pallor. Platelets also are large and lack granules. In functional studies, they may show marked abnormalities, and patients may have bleeding symptoms despite seemingly adequate numbers. Neutrophils are hypo-granulated; have ringed, or abnormally segmented nuclei; contain Do■hle bodies; and may be functionally deficient. Circulating myeloblasts usually correlate with marrow blast numbers, and their quantity is important for classification and prognosis.

The bone marrow is usually normal or hypercellular, but in about 20% of cases, it is sufficiently hypocellular to lead to confusion with aplastic anemia. The following are commonly observed:

dyserythropoietic changes (especially nuclear abnormalities) and ringed sideroblasts in the erythroid lineage; hypogranulation and hyposegmentation in granulocytic precursors, with an increase in myeloblasts; and megakaryocytes showing reduced numbers of or disorganized nuclei.

Only hematopoietic stem cell transplantation offers a cure for MDS. The hypomethylating agents azacitidine and decitabine are frequently used in bone marrow failure clinics. Lenalidomide, a thalidomide derivative with a more favorable toxicity profile, is particularly effective in reversing anemia in MDS patients with 5q– syndrome.

Solution to Question 18:

TGF beta is not a growth-promoting oncogene among the listed options.

TGF beta (Transforming Growth Factor beta) primarily acts as a regulatory molecule In normal cells, TGF beta functions mainly as a tumor suppressor by inhibiting cell proliferation, inducing apoptosis (programmed cell death), and promoting differentiation. It helps maintain tissue homeostasis and prevents the uncontrolled cell growth that can lead to cancer. In the early stages of tumorigenesis, TGF beta exerts anti-proliferative effects.

FGF (Fibroblast Growth Factor) is a family of growth factors involved in various processes, including angiogenesis, wound healing, and embryonic development. Some members of the FGF family can act as oncogenes, meaning they promote cell proliferation and survival, which can contribute to the development and progression of cancers. (Option A)

PDGF (Platelet-Derived Growth Factor) is another crucial growth factor that regulates cell growth and division. In the context of cancer, PDGF is often implicated due to its role in promoting cellular proliferation and survival. Dysregulation of PDGF signaling pathways is commonly observed in various tumors, making it a significant contributor to oncogenesis. (Option B)

TGF alpha (Transforming Growth Factor alpha) is a growth factor that binds to the EGF (Epidermal Growth Factor) receptor, promoting cell proliferation and survival. TGF alpha is frequently overexpressed in various types of cancers, where it stimulates the growth of tumor cells. Its ability to drive cellular proliferation through the activation of EGF receptors classifies it as a growth-promoting oncogene, contributing to the malignancy of cancers where it is overexpressed.(Option D)

Solution to Question 19:

The green vacutainer contains heparin as an anticoagulant and is commonly used for electrolyte estimation.

Green vacutainer contains heparin as an anticoagulant. It does not interfere with the levels of electrolytes in the blood sample.



Note: If serum electrolytes asked in the question - yellow bulb . If electrolyte only asked in the question- green bulb.

Yellow bulb is ideal for testing serum electrolytes like sodium, potassium, and chloride because they allow easy separation of serum after centrifugation.

Vacutainer	Additive	Uses
Blue	Sodium citrate	Coagulation tests (APTT and prothrombin time)
Lavender	EDTA	CBC, HbA1C(Option A)
Yellow	Acid-citrate-dextrose	HLA tissue typing, paternity testing, DNA studies(Option C)
Gray	Sodium fluoride and potassium oxalate	Blood glucose estimation(Option D)
Green	Lithium heparin	Chemistries(Option B)

Solution to Question 20:

The correct marking is i-d, ii-c, iii-b, iv-a.

Cystic fibrosis: An autosomal recessive disorder caused by mutations in the CFTR gene. Affects the lungs, pancreas, and other organs, causing thick mucus buildup that leads to breathing difficulties and digestive issues.

Leber's hereditary optic neuropathy (Leber's disease): A mitochondrial disorder, as it is inherited maternally due to defects in mitochondrial DNA. Sudden vision loss due to damage to the optic nerve, often affecting young adults.

Duchenne's muscular dystrophy: An X-linked recessive disorder caused by mutations in the dystrophin gene on the X chromosome. A severe form of muscular dystrophy that primarily affects boys, leading to progressive muscle weakness and loss of mobility.

Spinomuscular atrophy: Associated with trinucleotide repeat expansions (e.g., CAG repeats in some forms) leading to progressive muscle wasting. Affects motor neurons in the spinal cord, leading to muscle weakness and atrophy, often caused by mutations in the SMN1 gene.

i. Autosomal recessive	d. Cystic fibrosis
ii. Mitochondrial	c. Leber's disease
iii. X-linked recessive	b. Duchenne's muscular dyst rophy
iv. Trinucleotide repeats	a. Spinomuscular atrophy

Solution to Question 21:

a-iii, b-ii, c-iv, d-i is the correct match

Medullary thyroid carcinoma is a tumor of parafollicular (C cells) derived from the neural crest. Histological analysis shows characteristic amyloid stroma. High levels of calcitonin and carcinoembryonic antigen are produced by many medullary tumors.

Hepatoblastoma presents as an asymptomatic mass in the right upper quadrant of the abdomen. It mostly arises from the right lobe of the liver. Patients have non-specific symptoms such as fever, fatigue, weight loss, and anorexia. A biopsy of the liver mass is necessary to make the diagnosis. Alpha-fetoprotein (AFP) is a useful tumor marker that is used in diagnosis and monitoring.

Choriocarcinoma is a type of mixed germ cell tumor. It is highly vascular. Histologically, the tumor shows syncytiotrophoblasts and cytotrophoblasts. It secretes large quantities of human chorionic gonadotropin (hCG) hormone, which is used as a tumor marker in the diagnosis and management of the tumor.

Pheochromocytoma is a catecholamine-secreting tumor derived from the chromaffin cells of the adrenal medulla. It most commonly presents as a classic triad of intermittent headache, palpitation, and profuse sweating, along with hypertension.

a. Medullary thyroid cancer	iii. Calcitonin
b. Hepatoblastoma	ii. AFP
c. Choriocarcinoma	iv. Beta HCG
d. Pheochromocytoma	i. Catecholamin es

Solution to Question 22:

Epithelial-mesenchymal transition is induced by SNAIL & TWIST.

Epithelial-mesenchymal transition (EMT) is a process where epithelial cells lose their cell polarity and adhesion properties, transitioning into a more motile mesenchymal cell phenotype. This is critical in embryogenesis, wound healing, and cancer metastasis.

SNAIL and TWIST encode transcription factors whose primary function is to promote epithelial-to-mesenchymal transition (EMT). SNAIL and TWIST downregulate E-cadherin expression.

• In EMT, carcinoma cells downregulate certain epithelial markers (e.g., E-cadherin) and upregulate certain mesenchymal markers (e.g. vimentin and smooth muscle actin).

• This favors the development of a promigratory phenotype that is essential for metastasis.

• Loss of E-cadherin expression seems to be a key event in EMT.

• EMT has been documented mainly in breast carcinoma

Other Options:

Option A: TF IIA & TF IIB: These are general transcription factors involved in the assembly of the transcription initiation complex, not specific to EMT.

Option B: Cathepsin & MMP8: These are proteolytic enzymes involved in extracellular matrix remodeling, which may assist invasion during EMT but do not directly induce it.

Option D: STAT-5 & HIF-1α: While these factors are involved in cellular responses to cytokines and hypoxia, respectively, they are not directly linked to the initiation of EMT.

Solution to Question 23:

Bone marrow biopsy and flow cytometry are suggestive of ALL: B Phenotype.

In B ALL, the cells show markers such as intracellular tdt, CD 22, CD 19, and CD 10 on flow cytometry. Tdt represents that the cells are blasts, and CD 10, 19, and 22 represent that cells are B cells.

Peripheral smear shows lymphoblasts that have condensed chromatin, less conspicuous nucleoli, and scant basophilic cytoplasm, which lacks granules.

ALL is the most common cancer in childhood. B-cell ALLs usually present as early childhood leukemias, while T-cell ALLs are usually seen in adolescents as thymic lymphomas. Clinical features include fatigue, infections, bleeding diathesis (due to impaired marrow function), hepatosplenomegaly, fever, and dyspnea due to mediastinal or thymic infiltration.

Initial evaluation involves a routine complete blood count and peripheral smear. It typically reveals anemia, thrombocytopenia, and the presence of blasts (immature WBCs) on the smear. Morphologically, myeloblasts and lymphoblasts may be difficult to distinguish. Immunophenotyping thus plays an important role in confirming the diagnosis.

Other options:

Option A: Acute myeloid leukemia leads to the accumulation of immature myeloid blasts in the marrow, resulting in marrow failure and complications related to anemia, thrombocytopenia, and

neutropenia. Bone marrow/peripheral smear shows a large, uniform population of cells (myeloblasts) with prominent nucleoli. It shows the presence of primary cytoplasmic granules. Auer rods are Needle-shaped azurophilic intracytoplasmic inclusion bodies particularly seen in acute promyelocytic leukemia (AML M3) with t(15;17).



Option B: CLL is associated with the deletions of 13q, 11q, 17q, and trisomy 12q. Lymph nodes are diffusely effaced by small lymphocytes with proliferation centers. The peripheral smear shows small round lymphocytes with scant cytoplasm and smudge cells as seen below.



Option D: In T ALL, the cells show markers such as intracellular tdt, CD 1, CD 2, CD 5, and CD 7 on flow cytometry. Tdt represents that the cells are blasts, CD 1, CD 2, CD 5 and CD 7 represent that cells are T cells.

Solution to Question 24:

The given clinical and histopathological features are suggestive of Schwanomma, and they are benign and not malignant and do not metastasise frequently.

Schwannomas are benign tumors that often arise from cranial (Option A) and peripheral nerves and exhibit Schwann cell differentiation. They can be sporadic or associated with neurofibromatosis-1 (NF1) or occur as vestibular schwannoma (acoustic neuroma) in cases of neurofibromatosis-2 (NF2) (Option C).

They present with features of nerve compression involving both the adjacent structures and the nerve involved. Most tumors occur in the cranium as acoustic neuromas that affect the vestibular branch of the VIII cranial nerve near the cerebellopontine angle. They present with tinnitus, hearing loss, and symptoms resulting from compression of the brainstem and cerebellum.

Microscopically, it consists of dense areas called Antoni A area and loose areas called Antoni B area.

• The dense eosinophilic Antoni A areas contain spindle cells that are arranged into cellular intersecting fascicles. Palisading of nuclei is seen, with the nuclear-free zones between them, called Verocay bodies (Option B)

• The Antoni B areas contain spindle-shaped cells that are loosely arranged and separated by a prominent myxoid extracellular matrix.



Solution to Question 25:

The given clinical presentations of weight loss, cough, pleural effusion, and lymphadenopathy are suggestive of tuberculosis. Tree-in-bud appearance on the CT scan and elevated lymphocytes in the pleural fluid also point to TB. Necrotized epithelioid giant cells with necrosis would be the finding in the aspirate of the mediastinal lymph node, which is characteristic of tuberculosis.

The given symptoms are classic symptoms of pulmonary tuberculosis. Elevated ADA levels are more commonly associated with TB-related pleural effusions. Low ADA can occur in some cases of TB-related pleural effusions. Lymphocytic pleural effusions are common in TB pleuritis.

In tuberculosis, the characteristic histological feature in involved lymph nodes would be caseating granulomas, which consist of necrotized epithelioid giant cells with central necrosis. These granulomas are a hallmark of mycobacterial infection and are usually seen in the mediastinal lymph nodes of patients with active pulmonary tuberculosis.

Other options

Option B: Reactive lymphoid hyperplasia is typically seen in viral infections or other causes of non-specific lymphadenopathy, but it is not typical of tuberculosis.

Option C: Fungal hyphae surrounded by lymphocytes would suggest a fungal infection (e.g., histoplasmosis or coccidioidomycosis), but the overall presentation in the given scenario does not point toward a fungal infection.

Option D: Anthracotic pigment is seen in the lungs and lymph nodes of individuals with chronic exposure to coal dust or pollution but is not associated with tuberculosis.

Solution to Question 26:

a-iii, b-i, c-iv, d-ii is the correct match.

Inclusion bodies are structures with distinct sizes and shapes that are seen in infected cells under light microscopy.

Negri bodies seen in Rabies:

Disease	Inclusion bodies
a. Rabies	iii. Negri bodies
b. Parkinson's disease	i. Lewy bodies
c. Multiple myeloma	iv. Dutcher bodies
d. Malakoplakia	ii.Michaelis gutmann bodies



Lewy bodies seen in Parkinson's disease:



Dutcher bodies seen in multiple myeloma:



Michaelis Gutmann bodies seen in Malakoplakia:



Solution to Question 27:

Paroxysmal nocturnal hemoglobinuria (PNH) is not a hereditary cause of thrombosis.

PNH is an acquired disorder where RBCs are deficient in membrane-bound glycosylphosphatidylinositol (GPI)- CD55 (decay accelerating factor)and CD59 (membrane inhibitor of reactive lysis), which causes complement-mediated intravascular lysis of RBCs. It is characterised by a triad of pancytopenia with hypercellular marrow, nocturnal hemoglobinuria (decrease in blood pH during sleep), and acquired thrombosis.

Hypercoagulable states or thrombophilias (thrombosis) can be inherited or acquired:

Inherited causes include:

• Factor V Leiden mutation

• Prothrombin mutation

• Protein C deficiency

• Protein S deficiency (Option B)

• Antithrombin III deficiency (Option C)

• Homocystinuria (Option D)

Acquired causes include:

• Surgery

• Malignancy

• Antiphospholipid syndrome

• Trauma

• Pregnancy

• Long-haul travel

- Obesity
- Oral contraceptives/hormone replacement
- Myeloproliferative disorders
- Polycythemia vera

Solution to Question 28:

EDTA is used in test tubes as an anticoagulant for estimating HbA1C.

EDTA (ethylenediaminetetraacetic acid) is used for estimating HbA1c because it prevents clotting and preserves the integrity of the red blood cells for accurate measurement of glycated hemoglobin. It is a chelating agent, that works by binding to calcium ions in the blood.

Other Options:

Option A: Warfarin is an oral anticoagulant. It is not used in blood collection tubes. It is a therapeutic anticoagulant, not suitable for preserving blood samples for laboratory testing, such as HbA1c.

Option B: Sodium Fluoride is used in grey-top tubes for blood glucose testing. It is specific for glucose preservation, not for HbA1c estimation, as it does not effectively prevent clotting or preserve red blood cells.

Option D: Heparin is used in green-top tubes to collect plasma samples. It can interfere with some analytical methods for measuring glycated hemoglobin and does not preserve red blood cells effectively for this purpose.

Solution to Question 29:

The most common genetic cause of inherited coagulopathy is the Factor V Leiden mutation.

Primary (inherited) hypercoagulability is most often caused by mutations in the Factor V and prothrombin genes. Factor V Leiden mutation is the most common cause, followed by prothrombin G20210A mutation.

The Leiden mutation is inherited in an autosomal dominant pattern. The mutation makes Factor V resistant to cleavage and inactivation by protein C. This leads to the inactivation of an important antithrombotic counterregulatory pathway.

Protein C, Protein S, and Antithrombin III deficiency are rare causes of primary hypercoagulability.

Other options:

Option B: Protein C deficiency causes thrombosis in teenagers. It may also be induced by liver dysfunction, vitamin K antagonists, renal failure, DIC, and active thrombosis.

Option C: Antithrombin III forms a complex with thrombin by binding to heparin on endothelial cells. This complex inhibits coagulation. Its deficiency presents as early-age thrombosis (less than 50 years old) and carries the highest risk for thrombotic events among the inherited

thrombophilias.

Option D: The prothrombin G20210A mutation results in increased levels of prothrombin, which demonstrates a higher risk for arterial and venous thrombotic events.

Solution to Question 30:

a-iii, b-i, c-iv, d-ii is the correct match of the diseases with their respective gene mutations

Tumor suppressor genes are the genes that control the proliferation of the cells, but abnormalities in these genes lead to failure of growth inhibition, which is one of the molecular hallmarks of carcinogenesis.

Mutated tumor suppressor genes lead to many familial syndromes, such as :

Familial Syndrome	Mutated Tumor suppressor gene	Components of syndromes
Familial adenomatous polyposis	APC	Carcinoma of the stomach, colon, pancreas
Neurofibromatosis I	NF1	Neurofibroma, Malignant peripheral nerve sheath tumors
Neurofibromatosis II	NF2	Acoustic schwannoma and meningioma
Tuberous sclerosis	TSC1	subependymal nodules, giant cell astrocytoma, hypomelanotic macules (ash-leaf spots), facial angiofibroma, cardiac rhabdomyoma, renal angiomyolipoma
Familial Retinoblastoma	RB	Retinoblastoma, osteosarcoma
Cowden syndrome	PTEN	Variety of benign skin, GI, and CNS growths
Von-hippel lindau syndrome	VHL	Renal cell carcinoma
Peutz jeughers syndrome	STK11	GI polyps, GI cancers, and pancreatic cancers
Juvenile polyposis	SMAD 4	-
Familial Gastric cancer	CDH 1	Gastric carcinoma
Li Fraumeni syndrome	TP53	Diverse cancers in the body
Familial and male breast cancer, Ovarian carcinoma	BRCA1, BRCA 2	-
Wilms tumor	WT 1	-
Multiple Endocrine Neoplasia (MEN - 1)	MEN 1	Pituitary, parathyroid and Pancreatic endocrine tumours

Solution to Question 31:

Parathyroid hyperplasia is not a component of MEN2B syndrome. It is a feature of MEN2A.

MEN2B syndrome is a subtype of Multiple Endocrine Neoplasia type 2 (MEN2) caused by mutations in the RET proto-oncogene. Its characteristic features include:

• Medullary Thyroid Carcinoma (MTC): (Option A): Aggressive and occurs early in life.

• Pheochromocytoma: Tumor of the adrenal medulla

• Mucosal Ganglioneuromatosis: (Option C): A classic finding that is characterized by benign neuromas on mucosal surfaces such as the tongue, lips, and gastrointestinal tract.

• Marfanoid Habitus: (Option B): Long limbs, tall stature, and joint hypermobility.

Pathology NEET 2024

Question 1:

Which condition is inherited according to the given pedigree?



a) Wilson disease

b) Wiskott–Aldrich syndrome

c) Prader Willi syndrome

d) Achondroplasia

Question 2:

Which of the following is not true about vWD?

a) VWD type 3 has severely low levels of VWF

b) VWD type 1 is the most severe disease in children

c) VWD type 3 is the most severe disease in children

d) VWD type 2 is associated with moderate bleeding tendency

Question 3:

Which of the following is most common mutation associated with the microscopy image of the thyroid shown?



a) BRAF V600E

b) RET

c) MET

d) KRAS

Question 4:

A patient presented with abdominal pain, hepatosplenomegaly and WBC count >50,000/mm3. Which of the following genes is involved in this case?

a) t (15:17)

b) Del 1q

c) t (9:22)

d) Del 3q

Question 5:

A 55-year-old chronic smoker presents with weight loss, hemoptysis and TTF-1 positivity. What is the possible diagnosis?

a) Squamous cell carcinoma

b) Small cell carcinoma

c) Adenocarcinoma

d) Large cell carcinoma

Question 6:

A patient presents with flank pain and hematuria. The histology slide of the patient is provided. Based on the clinical presentation and histological findings, which of the following conditions is most likely?



a) Chromophobe renal cell carcinoma

b) Clear cell carcinoma

c) Metastasis

d) Papillary cell carcinoma

Question 7:

A patient came to the OPD with complaints of vomiting and diarrhoea along with skin eruptions. He gives the history of similar episodes in the past. After switching to gluten free diet his symptoms resolved. Which marker is useful in diagnosing this condition?

a) ASCA

b) ANCA

c) Anti tissue transglutaminase antibody

d) Anti-parietal Antibody

Question 8:

A 2 day old child is brought to the OPD. Mother complains that child is not feeding well and has bilious vomiting. The child is diagnosed with congenital megacolon and aganglionic segment of the colon is shown. Which of the following layers are affected in this condition?



a) Layer 1 and 2

b) Layer 2 and 3

c) Layer 3 and 4

d) Layer 1 and 4

Question 9:

A COPD patient with a history of smoking presents with respiratory symptoms. High resolution CT showed panacinar emphysema. Which enzyme deficiency is associated with this condition?

a) Alpha-1 antitrypsin

b) Elastase

c) Chymotrypsin

d) Type 2 surfactant

Question 10:

A 34-year-old female presented with low Hb, platelet count of 25000/mm3, raised PT and aPTT. A peripheral smear is given. Which of the following fusion genes is affected?



a) BCR ABL

b) IGH NSD2

c) RUNX1 RUNX1T1

d) PML RARA

Question 11:

A 12-year-old child comes with cervical lymphadenopathy. Lymph node biopsy shows CD 10 + , CD 19 + , CD 20 +, CD 5 -, CD 23 -, and Tdt -. What is the diagnosis?



a) Diffuse large B cell lymphoma

b) Acute Lymphoid Leukemia

c) Mantle cell lymphoma

d) Burkitt lymphoma

Question 12:

Identify the disease with the chromosomal abnormality?



a) Angelman syndrome

b) Prader Willi syndrome

c) Cri-du Chat syndrome

d) Rett syndrome

Question 13:

A 49-year-old patient comes with complaints of left flank pain, hematuria, weight loss, headache, nausea, and convulsions. A family history of brain tumours and kidney tumours is present. MRI showed a cerebellar mass. What is the diagnosis?

a) Li fraumeni syndrome

b) Tuberous sclerosis complex

c) Von Hippel-Lindau syndrome

d) Neurofibromatosis

Question 14:

A patient with COPD presented with ARDS symptoms. What is the role of IL-8 in inflammation?

a) Endothelial activation

b) Recuritement of neutrophils

c) Macrophage activation

d) Promote surfactant production

Question 15:

Which of the following is incorrect about iron metabolism?

a) Free serum iron is toxic to cells

b) Iron is stored in the body in the form of ferritin

c) Serum ferritin denotes the iron levels in the body

d) Iron is transported by binding with apoferritin

Question 16:

A patient passed away following a myocardial infarction. An autopsy was conducted and the biopsy findings of his cardiac tissue are shown below. What is the number of days since the injury, based on the given findings?



a) 7-10 days

b) 1 day

c) Around 3-4 weeks

d) 3-4 days

Question 17:

A 38-year-old woman came with complaints of a slowly growing breast lump for the last 6 months. She had a history of a similar lump for which excision was done and medical and mammographic reports are not available. On examination, the lump was smooth, well-defined, mobile and with regular margins. The cytology and biopsy of the same were taken and are shown below. What is the most likely diagnosis?



a) Leiomyosarcoma

b) Fibroadenoma

c) Sclerosing adenosis

d) Phyllodes tumor

Question 18:

A 72-year-old man came with generalised anascarca and icterus. He had hepatosplenomegaly. Kidney function tests were found to be deranged and his urine gammaglobulins were more than 3gm/dl. Electrophoresis of the urine protein is shown below. Which is the most likely diagnosis?



a) Multiple Myeloma - IgM

b) Multiple Myeloma - IgA

c) Multiple Myeloma - IgG

d) Monoclonal gammopathy of undifferentiated Proteins

Question 19:

A 53-year-old woman came with complaints of fatigue, nausea, anasarca and proteinuria. She had anaemia and neutropenia. Her Bone marrow examination showed 22% plasma cells and acellular eosinophilic deposits. The bone marrow biopsy is shown below. What is the associated pathology?



a) Bone marrow - fibrinoid necrosis

b) Bone marrow - coagulative necrosis

c) Amyloidosis

d) Myxomatous degeneration of the bone marrow

Question 20:

A 45 year old man came with complaints of tearing chest pain, tachycardia, and hypotension with negative Trop I and T. The wall of the aorta was biopsied and the image is shown below. What is the most likely pathology behind it?



a) Connective tissue damage of the aortic wall

b) Fibrinoid necrosis and smooth muscle hyperplasia

c) Fat deposition and smooth muscle degeneration

d) Immune complex vasculitis

Question 21:

A 10-year-old boy comes with a fever and pain in his right upper limb. He has an expansile lesion in his right radius in the diaphysis. The histopathology of the lesion is shown below. What is the most accurate test among the following?



a) t15;17 (q22 11-12)

b) RB mutation

c) EWSR1 (FUS- ETS fusion gene)

d) t11:22 (q24,q12)

Question 22:

A couple comes for prenatal genetic counselling. They have a family history of Hemophilia. They have a male child with Hemophilia and a normal female child. The mother's family has Hemophilia and the father's family is normal. What are the chances of their offspring being affected?

a) All males will be affected and all females will be normal

b) Half of the males can be affected and half of the females can be carriers

c) All males will be affected and half of the females will be carriers

d) All of males and females will be affected

Question 23:

A 10-year-old boy comes with visual disturbances and a headache. On imaging, he was found to have a suprasellar lesion. On biopsy and examination, the tumour exuded "black motor oil" like fluid. The MRI image is shown below. What is your likely diagnosis?



a) Glioblastoma

b) Craniopharyngioma

c) Meningioma

d) Pituitary adenoma

Question 24:

Which protein accumulation is seen in Alzheimer's disease?

a) Alpha Synuclein

b) Aβ amyloid

c) GFAP

d) Albumin

Question 25:

Which method is more effective than measuring clotting factors for assessing the intrinsic pathway of coagulation?

a) PT

b) APTT

c) Fibrinogen levels

d) CBC

Question 26:

An adult male presented with protruding abdomen, diarrhoea, visual symptoms and neurological manifestations. Based on the peripheral smear finding what is the likely diagnosis?



a) Burns

b) Abetalipoproteinemia

c) Uremia

d) EDTA changes

Question 27:

Which among the following statements is true regarding PNH?

a) Anemia with heart failure is the most common cause of mortality

b) It is an inherited condition

c) Sucrose lysis test is diagnostic

d) Flow cytometry is the gold standard for diagnosis

Question 28:

A patient presented with an arm span of more than height, a subluxated lens, and a pigeon chest appearance. What is the likely protein involved in this condition?

a) Fibrillin

b) Collagen

c) Elastin

d) Laminin

Question 29:

A 6-month-old child presented with leucocoria (cat's eye reflex) in the right eye. A cranial pathology was detected. The lesion is positive for synaptophysin, NSE, and chromgranin, and shows a palisade in a spoke-and-wheel arrangement, surrounding a central core. Identify the condition.

a) Retinoblastoma

b) Medulloblastoma

c) Pineoblastoma

d) Epidermoid carcinoma

Question 30:

A middle aged man came with complaints of dyspnea, chest tightness and dry cough. On imaging, he was found to have diffuse interstitial fibrosis. The biopsy of his lung was taken and is shown below. Which of the following industries he has likely been working in for the last 20 years?



a) Asbestos

b) Silica

c) Coal

d) Beryllium

Question 31:

A 45 year old woman came with complaints of fever, night sweats, generalised itching and unintentional weight loss of 10% in the last 3 months. On examination, she has enlarged cervical and axillary lymph nodes. The lymph node examination and biopsy showed the following picture. What would the likely diagnosis and treatment option be?



a) Non Hodgkins Lymphoma - Rituximab monotherapy

b) Non Hogkins Lymphoma - Poor prognosis due to advanced stage - treated with radiotherapy

c) Hodgkins Lymphoma - ABVD regimen

d) Sarcoidiosis - Started on Prednisolone 40 mg/day

Question 32:

A 28 year old woman P1L1, came with complaints of weakness and generalised fatigue. The peripheral blood picture showed microcytic hypochromic anemia. Her serum iron was low and she had increased RDW. Her MCV and MCH were also low and TIBC was high. What would her serum transferrin saturation be?

a) Reduced

b) Cannot be determined

c) Equivocal

d) Increased

Question 33:

A 38-year-old woman came with complaints of weakness, lethargy, and heavy menstrual bleeding. Her peripheral blood picture showed small RBCs with pallor. Lab investigations are given below. What is the diagnosis?

a) Megaloblastic anemia

b) Iron-deficiency anemia

c) Hemolytic anemia

d) Hypoplastic anemia

Answer Key

Question No.	Correct Option
1	b
2	b
3	a
4	c
5	c
6	a
7	c
8	b
9	a
10	d
11	d
12	b
13	c
14	b
15	d
16	b
17	d
18	a

19	c
20	a
21	c
22	b
23	b
24	b
25	b
26	b
27	d
28	a
29	a
30	a
31	c
32	a
33	b

Detailed Explanations

Solution to Question 1:

The given pedigree chart points towards an X-linked recessive mode of inheritance and among the options provided Wiskott–Aldrich syndrome is inherited via this mode.

A Triad of micro-thrombocytopenia, eczema, and recurrent infections points towards Wiskott–Aldrich syndrome.

Other options:

Option A: Wilson's disease is caused by a defect in the gene ATP7B on chromosome 13, which results in decreased biliary copper excretion and its accumulation in the liver. It is known to cause hepatolenticular degeneration. Serum ceruloplasmin levels are decreased.

Option C: Prader-Willi syndrome belongs to a group of disorders called genomic imprinting i.e. differential expression of a gene based on chromosomal inheritance from maternal or paternal origin. It is due to the deletion of paternally derived Chr 15.

Option D: Achondroplasia is the most common skeletal dysplasia that follows the autosomal dominant inheritance pattern (. It characteristically presents with macrocephaly, disproportionately short limbs, facial dysmorphism (e.g., mid-face hypoplasia, frontal bossing), joint laxity, and hypotonia.

Solution to Question 2:

Von Willebrand disease type 1 is not the most severe disease in children. On the other hand, it has been found that it is VWD type 3 is the most severe.

Features of VWD:

• Most common type is Type 1

• The most severe type is Type 3

• Most common inheritance: Autosomal dominant

• Type 1- partial quantitative deficiency of VWF

• Type 2- qualitative deficiency of VWF

• Type 3- complete quantitative deficiency of VWF

Characteristics of the subtypes of VWD are summarized below :

AD: autosomal dominant

AR: autosomal recessive

Type	Inheritance	Mechanism	Important characteristics
1	AD	Partial, quantitative deficiency of VWF	Most common type
3	AR	Complete quantitative deficiency	Most severe VWD
2A	AD	Loss of VWF multimers (increased cleavage by ADAMS13)	Platelet adhesion defect
2B	AD	Increased binding of VWF to platelets causing VWF-platelet complex destruction	Thrombo-cytopenia
2M	AD	Decreased affinity of VWF to platelets	Platelet adhesion defect, very rare
2N	AR	Decreased affinity of VWF to Factor VIII	Autosomal hemophilia

Solution to Question 3:

The most common mutation associated with the microscopy image which shows the Orphan Annie eye nuclei which is characteristic of papillary carcinoma of the thyroid is the BRAF V600E mutation.

Papillary thyroid carcinoma is the most common thyroid cancer. It is also the most predominant thyroid cancer in children and iodine-sufficient areas. Radiation exposure is an important risk factor.

Genetics :

BRAF gain-of-function mutation (V600E amino acid substitution) is commonly associated.

RET/PTC rearrangements

NTRK gene fusions

The clinical features of Papillary thyroid carcinoma are as follows :

• It is more common in females

• The mean age of presentation is 30-40 years

• Most patients are euthyroid, with painless swelling. Initial presentation may be with cervical lymphadenopathy due to lymph node metastases.

• These are usually multifocal (85% cases)

• They appear as cold masses on thyroid scan

• Metastasis occurs via the lymphatic route

• Skip metastasis from papillary thyroid carcinoma to the ipsilateral cervical lymph nodes, known as lateral aberrant thyroid, is managed with modified radical neck dissection

Microscopic features of papillary thyroid carcinoma include :

• Branching papillae with fibrovascular stalk

• Covering epithelium is simple or stratified cuboidal

• Orphan Annie eye nuclei - Empty-appearing nuclei due to dispersed chromatin, they are diagnostic, regardless of papillary architecture

• Psammoma bodies - concentric calcified structures, foci of lymphatic invasion



BRAF V600E mutation is also seen in :

• Langerhans cell histiocytosis

• Hairy cell leukemia

• Colon adenocarcinoma

• Melanoma

Solution to Question 4:

The clinical scenario of abdominal pain, hepatosplenomegaly, a TLC of > 50,000/mm3 points toward the diagnosis of chronic myeloid leukemia (CML) which results from reciprocal translocation t(9:22) forming the BCR-ABL fusion gene.

CML is a hematopoietic stem cell disorder that is common in males 55-65 years of age. It occurs due to the translocation between the long arms of chromosomes 9 and 22, forming the Philadelphia chromosome. This results in the activation of tyrosine kinase, which causes abnormal cellular proliferation. Common presenting features are fatigue, malaise, weight loss, and splenomegaly.

Blood investigations show anemia, leucocytosis, and thrombocytosis. It has decreased levels of leucocyte alkaline phosphatase due to a relative increase in abnormal myeloid cells. The emergence of tyrosine kinase inhibitors as first-line therapy has dramatically improved survival for these patients. Imatinib is the most commonly used tyrosine kinase inhibitor used for CML.

The image below shows peripheral smear findings of CML with neutrophils, myelocytes, metamyelocytes, and band cells.



Option A: translocation (15:17) - This is seen in the acute promyelocytic variant of acute myeloid leukaemia. Clinically, the patients present with symptoms resulting due to pancytopenia (anemia, neutropenia, thrombocytopenia). For acute myeloblastic leukaemia (AML) diagnosis, the blast cell count should be ≥20% in the peripheral blood or bone marrow. Peripheral smears (as depicted below) would show the classic intracytoplasmic Auer rods (red arrows) :



Option C: The 1q21.1 deletion syndrome itself does not lead to a clinically recognizable syndrome. Some patients can present with dysmorphic facial features, mild intellectual disability, microcephaly, and eye abnormalities.

Option D: The 3q29 deletion syndrome clinically presents with psychiatric manifestations including mild-to-moderate intellectual disability, autism spectrum disorder, anxiety disorders, attention-deficit/hyperactivity disorder, executive function deficits, and psychosis.

Solution to Question 5:

The clinical scenario cough, hemoptysis, weight loss with image showing immunohistochemical marker thyroid transcription factor-1 (TTF-1) positivity is suggestive of adenocarcinoma of the lung.

Adenocarcinoma is the most common type of lung cancer seen in women and nonsmokers. Adenocarcinomas are usually located in the periphery as compared to other lung tumors and are relatively smaller in size. Atypical adenomatous hyperplasia is a precursor lesion of adenocarcinoma.

Molecular genetics show frequent mutations in EGFR, ALK, ROS, MET, and RET (Receptor tyrosine kinase genes) or KRAS mutations. The majority express TTF-1, which is a protein required for normal lung development. Napsin A is a new marker and is more sensitive and specific than TTF-1 in the diagnosis of adenocarcinoma.

Microscopic examination shows a lepidic pattern of growth at the periphery of the tumor. The tumor cells appear to crawl along the normal-appearing alveolar septa, as shown in the image below.



Thyroid transcription factor 1 (known as NKX2-1 or thyroid-specific enhancer binding protein) is a nuclear immunohistochemistry marker with preferential expression in thyroid, lung and brain structures of diencephalic origin. It is positive in adenocarcinoma of the lungs.

The image below shows the immunohistochemistry marker TTF-1 in lung adenocarcinoma



Other options:

Option A: p40 and p63 are markers of squamous cell carcinoma lung. It is the most common cancer found in smokers. Histologically it is characterized by the presence of keratinization and markedly eosinophilic cytoplasm as shown below.



Option B: Chromogranin, synaptophysin, and CD56 are the markers of small cell carcinoma lung. Due to the encrustation by DNA from the necrotic cancer cells, vascular walls in the necrotic area take up basophilic staining as shown below. This is called the Azzopardi effect.



Option D: Large cell carcinoma is an undifferentiated malignant epithelial tumour that lacks the cytologic features of other forms of lung cancer. They do not express markers of adenocarcinoma or squamous cell carcinoma.

The image below shows large cell undifferentiated carcinoma



Solution to Question 6:

The given histological image is of chromophobe renal cell carcinoma. It can be identified by the presence of pale eosinophilic cells arranged in the form of solid sheets with some cells having a perinuclear halo.

Chromophobe carcinoma is thought to arise from the intercalated cells of the collecting ducts. It is composed of cells with prominent cell membranes and pale eosinophilic cytoplasm, usually with a halo around the nucleus. It shows large cells having a plant-cell-like appearance. They are arranged in solid sheets with a concentration of the largest cells around blood vessels.

Cytogenetic examination of the tumour shows multiple chromosome losses and extreme hypodiploidy. It has an excellent prognosis compared to clear cell and papillary cancers.

Close differential diagnoses are as follows:

• Oncocytoma: They also consist of large eosinophilic cells having small, round, benign-appearing nuclei that have large nucleoli. However, they do not show a perinuclear halo.

• Clear cell RCC: Pale eosinophilic cytoplasm in chromophobe RCC distinguishes it from the clear appearance of the cytoplasm due to the accumulation of glycogen and lipids in clear-cell RCC.

The below image shows clear cell renal carcinoma:



Solution to Question 7:

The given scenario of vomiting, diarrhoea, skin eruptions and resolving symptoms with gluten free diet along with crypt hperplasia, villous atrophy and intra epithelial lymphocyte as seen in histopathology are suggestive of celiac disease. Anti tissue transglutaminase antibody will help in diagnosing this condition.



Small intestinal adenocarcinoma is also associated with celiac disease.

Celiac disease or celiac sprue or gluten-sensitive enteropathy is triggered by the ingestion of gluten, which is the major storage protein of wheat and similar grains. It usually affects the 2nd part of the duodenum or proximal jejunum.

Antibodies associated with celiac disease are:

• Anti-gliadin antibodies

• Anti tissue transglutaminase antibodies

• Anti endomysial antibodies

MARSH's score is used to assess celiac disease histologically.

Dermatitis herpetiformis occurs in about 10% of patients with celiac disease.

Other options :

Option A : Serological markers that are usually positive in patients with Crohn's disease include the anti–Saccharomyces cerevisiae antibody (ASCA).

Option B : Anti-neutrophil cytoplasmic antibodies (ANCA)-associated vasculitis is categorized into 3 distinct forms: Granulomatosis with polyangiitis (GPA; formerly Wegener granulomatosis), Microscopic polyangiitis (MPA), Eosinophilic granulomatosis with polyangiitis, formerly called Churg–Strauss syndrome (CSS).

Option D : Anti-parietal cell antibody is positive in patients with Pernicious anemia.

Solution to Question 8:

The clinical presentation of bilious vomiting, congenital megacolon and aganglionic segment of the colonis suggestive of Hirschsprung's disease. The layers 2 and 3 representing the submucosa and muscularis, respectively, are affected by this condition.

Hirschsprung disease or congenital aganglionic colon is characterized by the absence of ganglion cells in the submucosal (Meissner) plexus - Layer 2 and myenteric (Auerbach) plexus - Layer 3 of the distal intestinal segment. This is because of the premature arrest of normal migration of neural crest cells from the cecum to the rectum.

It causes a muscular spasm of the distal intestinal segment, resulting in functional obstruction of the affected bowel and proximal dilation (megacolon). The area between the dilated and contracted segments is referred to as the transition zone.

The image below shows layers of the colon



Hirschsprung disease is classified into -

• Short-segment - limited to the rectosigmoid segment

• Long-segment - proximal extension beyond the rectosigmoid segment

Note: while the gold standard investigation for Hirschsprung's disease is rectal suction biopsy, an imaging study will first be required to figure out the location from where the sample needs to be taken.

The image below shows dilated ganglionic and contracted aganglionic segments in Hirschsprung's disease,



It is more common in males. Most infants present with abdominal distension and bilious vomiting with failure to pass meconium within the first 24 hours of life. Enterocolitis is a potentially fatal complication.

It may be familial or associated with Down syndrome. The following gene mutations can be seen,

• Mutations in chromosome 10 (loss-of-function mutations in the receptor tyrosine kinase RET) or chromosome 13

• glial cell line-derived neurotrophic factor (GDNF) mutations

GDNF promotes the survival, proliferation, and migration of neural crest cells in culture. It is expressed in the gut in advance of migrating neural crest cells and is chemoattractive for neural crest cells in culture.

The investigations for Hirschsprung's disease include:

Water-soluble contrast enema is the initial investigation. It shows a narrow aganglionic segment, a conical transition zone, and a dilated proximal bowel. It helps in excluding other causes of distal intestinal obstruction, such as colonic atresia, meconium ileus, and meconium plug syndrome. It is both diagnostic and therapeutic for the latter.

A contrast enema image in Hirschsprung's disease is shown below,



Rectal biopsy is the gold standard investigation. It shows the following:

• Absent ganglia

• Hypertrophied nerve trunks

• Positive immunostaining for acetylcholinesterase

• Calretinin immunostaining (superior to acetylcholinesterase staining)

The ideal site to obtain a rectal biopsy in a patient with Hirschsprung's disease is from a point that is at least 2 cm above the dentate line to avoid sampling the normal aganglionated region of the internal sphincter.

Anorectal manometry shows the failure of the relaxation of the internal anal sphincter. It can be performed in older children, but it is relatively inaccurate and rarely used.

The management of Hirschsprung's disease is as follows:

Surgery is the definitive treatment. A multi-stage approach may be adopted. A colostomy is done in the newborn period, followed by a definitive pull-through operation when the child is over 10

kgs.

Definitive surgeries for Hirschsprung disease are as follows:

• Swenson

• Duhamel

• Soave

Solution to Question 9:

Alpha-1 antitrypsin deficiency is associated with panacinar emphysema.

Panacinar or panlobular emphysema is associated with α1 antitrypsin (α1AT) deficiency. It is exacerbated by smoking. Panacinar occurs more commonly in lower zones and the anterior margins of the lung. In this type, the acini from respiratory bronchiole to terminal blind alveoli are uniformly enlarged.

α1AT deficiency is an autosomal-recessive disorder of protein misfolding. Deficiency of α1AT protein results in excessive degradation of alveoli by neutrophil elastase causing panacinar emphysema. It also causes liver disease as a consequence of the abnormal accumulation of the misfolded protein within hepatocytes. α1AT deficiency is characterized by the presence of round-to-oval cytoplasmic globular inclusions in hepatocytes. These hepatocytes are strongly Periodic acid–Schiff positive and diastase resistant.

Trypsin, chymotrypsin, and elastase are endopeptidases. They cleave the internal peptide bonds. Trypsin is specific for cleavage at the sites of basic amino acids, while chymotrypsin and elastase are specific for cleavage at neutral amino acids.

Other options:

Option B: Elastase is inhibited by alpha-1 antitrypsin. In the case of alpha 1 antitrypsin deficiency, there is the destruction of lung tissue (elastin) by the elastase released by macrophages.

Option C: Chymotrypsin degrades trypsin and protects the pancreas from trypsin-related injury.

Option D: Type 2 surfactant deficiency is associated with respiratory distress syndrome as type 2 pneumocytes produce the surfactant which decreases the surface tension.

Solution to Question 10:

The image showing numerous Auer rods and the lab findings pointing towards DIC are suggestive of acute myeloid leukaemia - M3 or acute promyelocytic leukaemia. The pathogenesis behind this is the PML-RARA fusion gene or translocation t(15,17).

AML-M3, or acute promyelocytic leukemia, is a subtype of acute myeloid leukemia associated with PML-RAR■ fusion gene or t(15;17). The patient presents with fever, fatigue, and weight loss. On examination, findings consistent with anemia (pallor) and thrombocytopenia (easy bruising) are seen. This is due to the accelerated destruction of RBC and platelet precursors in the bone marrow.

Histopathology of the bone marrow aspirate reveals numerous Auer rods (needle-like azurophilic granules) in the cytoplasm of the promyelocytes. Sometimes, multiple Auer rods cluster together within a single cell resembling a bundle of sticks. Such cells have been termed faggot cells (faggot in French means bundle of sticks).

The condition is associated with a high incidence of disseminated intravascular coagulation (DIC).

Treatment includes the administration of all-trans retinoic acid.

Other options:

Option A: CML is a myeloproliferative neoplasm. It is caused by BCR (on chromosome 22) - ABL (on chromosome 9) fusion gene translocation mutation. It results in the formation of the Philadelphia chromosome.

Option B: Multiple myeloma patients are affected by the translocation t(4;14) resulting in the IGH::NSD2 fusion.

Option C: AML with t(8:21) shows the RUNX1/RUNXT1 fusion gene.

Solution to Question 11:

The clinical scenario of a child with cervical lymph node, which is positive for CD 10, 19, and 20 and negative for CD 5, 23, and tdt and the histopathology showing tumour cells showing a diffuse monotonous pattern with clumped and dispersed chromatin in the nuclei along with deeply basophilic cytoplasm containing lipid vacuoles points towards a diagnosis of Burkitt's lymphoma.

Burkitt lymphoma is a highly aggressive B-cell non-Hodgkin lymphoma. It is characterized by the translocation of the c-Myc gene on chromosome 8. This leads to increased MYC protein levels. The translocation most commonly seen in Burkitt's lymphoma is t(8;14).

It can occur in 3 forms that have distinct clinical characteristics:

• African (endemic) Burkitt lymphoma presents as a mass involving the mandible. It may also involve the abdominal viscera, particularly the kidneys, ovaries, and adrenal glands.

• Sporadic (non-endemic) is often seen as a mass involving the ileocecum and peritoneum.

• Aggressive lymphoma or acute leukaemia is seen in immunocompromised individuals

Histopathology shows tumour cells with round or oval nuclei, coarse chromatin, small nucleoli, minimal cytoplasm, and a high mitotic index. The phagocytes have abundant clear cytoplasm, creating a characteristic “starry sky” pattern. Bone marrow aspiration shows tumour cells with royal blue cytoplasm containing clear cytoplasmic vacuoles.

The tumour cells express surface IgM, CD19, CD20, CD10, and BCL6, a phenotype consistent with a germinal centre B-cell origin.



Other options:

Option A: Diffuse large B cell lymphoma/ DLBCL is the most common type of non-Hodgkin lymphoma and is characteristic of large cell type and shows a diffuse pattern of growth.are positive for CD10, Cd23 and Bcl-6.



Option B: Acute lymphoblastic leukaemia (ALL) arises from immature B (pre-B) or T (pre-T) cells called lymphoblasts. Immunohistochemistry for terminal deoxynucleotidyl transferase (TdT) is positive in pre-B and pre-T lymphoblasts.

Peripheral smear shows lymphoblasts that have condensed chromatin, less conspicuous nucleoli, and smaller amounts of cytoplasm which lacks granules as shown below.



Option C: Mantle cell lymphoma is a lymphoid neoplasm in which tumour cells closely resemble the normal mantle zone B cells surrounding the germinal centre. Mantle cell lymphomas express high levels of cyclin D1. These cells are positive for CD19, CD20, CD5, and surface immunoglobulins (Ig). They are negative for CD23. In cases, where cyclin D1 is negative, SOX11 marker can be used to identify this lymphoma.

Lymph node biopsy shows medium-sized lymphoma cells with irregular nuclear outlines, scanty cytoplasm, and condensed chromatin in mantle cell lymphoma.



Solution to Question 12:

Paternal deletion of 15q12 is seen in Prader-Willi syndrome as shown in the image.

Prader-Willi syndrome is a type of non-Mendelian mode of inheritance and genomic imprinting disorder. It occurs due to the deletion of band 15q12 on the paternal chromosome and the silencing of the same set of genes derived from maternal chromosomes. It is characterized by mental retardation, obesity, hyperphagia, hypogonadism, short stature, small hands and feet, and abnormal facial features like a small upturned nose, narrow bifrontal distance and thin upper lip.

Other options:

Option A: Angelman syndrome is characterized by the deletion of the ubiquitin ligase gene (UBE3A) on maternal chromosome 15q12 and the silencing of the paternal chromosomal counterpart. It is characterized by intellectual disability, microcephaly, ataxic gait, seizures, inappropriate laughter and so is also known as the 'happy puppet syndrome'.

Option C: Cri-du Chat syndrome due to partial deletion or monosomy of short arm chromosome 5.

Option D: Rett syndrome is a neurodevelopmental disorder. It is caused by a mutation in the MECP-2 gene. The disorder is mostly seen in girls because affected boys usually die in infancy due to a defective single X chromosome.

Prader Willi syndrome	Angelman syndrome
Chromosome 15	Chromosome 15
Maternal imprinting	Paternal imprinting
Paternal deletion	Maternal deletion
Maternal disomy	Paternal disomy
SNORP gene	UBE3A gene
Hyperphagia, obesity, mental retardation	Happy puppet syndrome

Solution to Question 13:

The given clinical history of left flank pain, hematuria, weight loss, headache, nausea, and convulsions along with the family history of brain tumours and kidney tumours is suggestive of Von Hippel-Lindau syndrome. (VHL)

VHL syndrome is inherited as an autosomal dominant trait and is characterized by a predisposition to a variety of neoplasms, particularly hemangioblastomas of the cerebellum and retina. Hundreds of bilateral renal cysts, often multiple, clear cell carcinomas develop in 40% to 60% of affected persons. Those with VHL syndrome inherit a germline mutation of the VHL gene on chromosome 3p. A 'second hit' can lead to manifestations as described below.

Other options:

Option A: Li-Fraumeni syndrome is an autosomal dominant disorder due to a defect in the tumour suppressor gene P53 located on chromosome 17. It is characterised by Sarcomas, Leukaemia, Osteosarcomas, Brain tumours and Adrenocortical carcinomas.

Option B: Tuberous sclerosis is caused by to mutation of genes coding for hamartin (chromosome 9) and tuberin (chromosome 16), which leads to the formation of hamartomas, subependymal nodules, and astrocytomas. Skin manifestation includes facial angiofibroma (adenoma sebaceum), shagreen patch and confetti skin lesions. It can also be associated with renal angiomyolipoma and cardiac rhabdomyoma.

Option D: Neurofibromatosis-1 is caused by the autosomal dominant loss-of-function mutations in the NF1 gene on chromosome 17. It is a tumour suppressor gene that encodes neurofibromin, a negative RAS regulator. It can present with café-au-lait macules (CALMs), axillary or inguinal freckling, Lisch nodules, neurofibromas, pseudoarthrosis, and optic gliomas.

ORGAN SY STEM	LESION
Eye	Retinal angiomas (benign)
CNS	Hemangioblastomas (benign)
Kidney	Clear-cell renal cell carcinom aRenal cyst
Adrenal gla nd	Pheochromocytoma
Pancreas	Neuroendocrine tumours (be nign and malignant)Pancreat ic cysts
Epididymis	Cystadenoma
Ear	Endolymphatic sac tumour

Solution to Question 14:

IL-8 (CXCL8) is secreted by activated macrophages and endothelial cells which causes activation and recruitment of neutrophils.

Once leukocytes have exited the circulation, they move towards the inciting stimulus by chemotaxis. This process is mediated by:

Exogenous chemoattractants, such as bacterial products as well as endogenous chemoattractants: cytokines of the chemokine family (e.g., IL-8), components of the complement system (particularly C5a), and arachidonic acid (AA) metabolites (mainly leukotriene B4).

Solution to Question 15:

Iron is transported by ferroportin and not apoferritin. Apoferritin is a storage form of iron.

Apoferritin is present in the intestinal mucosa membrane and it binds and stores iron, by combining with a ferric hydroxide–phosphate compound to form ferritin.(Option B)

Iron absorption occurs in the duodenum. Iron in our gut exists in 2 forms, they are non-heme and heme iron. Luminal non-heme iron is mostly in the Fe3+ (ferric) state and is reduced to Fe2+ (ferrous) iron by ferrireductases such as duodenal cytochrome B. Fe2+ iron is then transported across the apical membrane by divalent metal transporter 1 (DMT1). Heme iron is moved across the apical membrane into the cytoplasm through heme transporters such as heme carrier protein 1 and heme oxygenase 2.

In the duodenal cells, Fe2+ iron can be transported to the blood or stored as mucosal iron. Fe2+ form of iron is transported from the cytoplasm into circulation by the protein ferroportin. A protein called hephaestin is associated with ferroportin 1. It is not a transporter itself, but it facilitates basolateral transport.

Once in circulation, Fe2+ is again oxidized back to Fe3+ iron by hephaestin and ceruloplasmin. The newly absorbed Fe3+ iron binds rapidly to transferrin and is delivered to red cell progenitors in the marrow. Hepcidin is a protein secreted from the liver that plays a major role in the regulation of iron absorption. When the body's iron stores are adequate, it downregulates the expression of ferroportin and thereby decreases iron absorption.



Other options:

Option A: Free serum iron is toxic to cells. After entering cells, free iron gathers in the mitochondria. This produces free radicals, catalyses lipid peroxidation, interferes with oxidative phosphorylation, and eventually results in cell death.

Option C: Serum ferritin denotes the iron levels in the body.

Solution to Question 16:

The biopsy of the cardiac tissue showing neutrophilic infiltration is suggestive that the injury is 1 day old.

In a patient with myocardial infarction around 1-3 days heart will show mottling accompanied by a yellow-tan infarct and histopathology will show coagulative necrosis with striations and loss of

nuclei, and a rapid neutrophil infiltration of the interstitial space.

Solution to Question 17:

The given clinical scenario and the image showing leaf-like architecture, stromal cellularity and mitotic figures are suggestive of phyllodes tumor.



Phyllodes tumour is a fibroepithelial tumour arising from the intralobular stroma. It presents as a smooth, rapidly growing, multinodular, painless breast lump that is usually palpable. It most commonly presents in the 6th decade of life and is associated with:

• Gain in chromosome 1q

• HOXB13 overexpression

Phyllodes tumours show stromal cellularity next to the epithelium (subepithelial or peri epithelial regions). This feature helps to differentiate it from intracanalicular fibroadenoma.

The prognosis of phyllodes tumours depends on their histological grade. Most phyllodes tumours are low-grade, with limited stromal cellularity. They recur only occasionally and do not metastasize.

The features of malignant (intermediate and high grade) phyllodes tumours are:

• Markedly increased stromal cellularity

• Stromal overgrowth - the absence of epithelial elements

• Nuclear atypia

• Increased mitosis (>10/ 10 hpf)

• Highly recurrent

Regardless of the grade, lymphatic spread is rare in phyllodes tumours. Hence, axillary lymph node dissection is contraindicated.

Note: Mitotic figures indicate active cell division, reflecting the tumour's proliferative nature. Their presence, particularly when abundant or atypical, can suggest a higher tumour grade and potential aggressiveness.

Other options:

Option A: Leiomyosarcoma is a malignant smooth muscle cell tumour that arises from the myometrium or endometrial stromal precursor cells. Grossly polypoid masses that protrude into the uterine lumen, or large, fleshy masses that invade the uterine wall. Cytologic atypia is seen in histology, ranging from highly anaplastic to extremely well differentiated. Histopathology of leiomyosarcoma shows variable atypia, often with cytoplasmic vacuoles at both ends of nuclei, which is given below:



Option B: Fibroadenoma is a form of aberrations of normal development and involution of the breast (ANDI), and it is the most common cause of a breast lump in women aged 15–25 years. It arises from hyperplasia of a single lobule and presents as a well-circumscribed, smooth or lobular, firm, non-tender, freely mobile mass (mouse in the breast). The histopathology shows the proliferation of intralobular stroma, and the epithelium may be surrounded by stroma (pericanalicular pattern) or compressed and distorted (intracanalicular pattern). The histopathology image is given below:



Option C: Sclerosing adenosis is a benign condition where the breast lobules contain fibrous tissue that resembles scars. Histopathology shows a complex entity with a prominent myoepithelium and stromal fibrosis, comprised of enlarged and distorted lobules with duplicated and crowded acini. The image is given below:



Solution to Question 18:

The given clinical scenario along with deranged RFT, urine gammaglobulins >3gm/dl, and urine electrophoresis showing IgM kappa paraprotein is suggestive of multiple myeloma.

The electrophoresis pattern of multiple myeloma would typically show:

• Monoclonal Spike (M-spike): A distinct peak in the gamma region, representing the high concentration of the IgM kappa paraprotein.

• Reduced or Absent Polyclonal Immunoglobulins(IgG, IgA), indicating a suppression of normal antibody production.

• Increased kappa light chains, with a corresponding decrease in lambda light chains, confirm the kappa light chain restriction.

Multiple myeloma is a plasma cell neoplasm commonly associated with lytic bone lesions, hypercalcemia, renal failure, and acquired immune abnormalities.

Poor prognostic factors for multiple myeloma:

• Serum β2-microglobulin. It is the single most powerful predictor

• Deletion of 17pITP53 sequences and the translocations t(14;16) and t(14;20)

• Polyclonal serum Immunoglobulins

• Elevated lactate dehydrogenase

• High C-reactive protein

• Increased plasma cell proliferative activity

• High degree of bone marrow replacement

• Presence of plasmablastic morphology

• Elevated serum soluble receptor for IL6

Multiple myeloma usually presents as destructive plasma cell tumors involving the axial skeleton. The bones most commonly affected in descending order of frequency are the vertebral column, ribs, skull, pelvis, femur, clavicle, and scapula.

Radiographically, bone lesions appear as punched-out defects, usually 1 to 4 cm in diameter, and consist of soft, gelatinous, red tumor masses.

Radiograph of the skull showing punched out lesions:

Morphology:

• Malignant plasma cells have a perinuclear clearing due to a prominent golgi apparatus and an eccentrically placed nucleus.

• Relatively normal-appearing plasma cells, plasmablasts with vesicular nuclear chromatin and a prominent single nucleolus, or bizarre, multinucleated cells may predominate.

• Other cytologic variants include flame cells, Mott cells.

• Flame cells have fiery red cytoplasm, Mott cells contain multiple grapelike cytoplasmic droplets, a variety of other inclusions, including fibrils, crystalline rods, and globules. The globular inclusions are referred to as Russell bodies (if cytoplasmic) or Dutcher bodies (if nuclear).

• Rouleaux formation is characteristic but not specific. High levels of M protein in the blood causes red cells to stick to one another in linear arrays in smears leading to rouleaux formation.

Image showing Mott cell:



Image showing Russell and Dutcher bodies:



Peripheral smear of multiple myeloma:



Bence Jones proteins are excreted in the kidney and causes myeloma kidney. Most myelomas are associated with more than 3 g/dL of serum Ig and/or more than 6 mg/dL of urine Bence Jones protein. The most common monoclonal Immunoglobulin (M protein) is IgG followed by IgA.

The clinicopathologic diagnosis of multiple myeloma can be done based on the identification of clonal plasma cells in the marrow and the presence of CRAB criteria (hypercalcemia, renal dysfunction, anemia, and bone lesions).

Solution to Question 19:

The clinical scenario along with the bone marrow biopsy showing increased plasma cells(poor prognosis) showing PAS positive eosinophilic substance is suggestive of amyloidosis.

Amyloidosis refers to extracellular deposits of fibrillar proteins. Abnormal fibrils are produced by the aggregation of misfolded proteins. The fibrillar deposits bind a wide variety of proteoglycans and glycosaminoglycans, including heparan sulfate and dermatan sulfate, and plasma proteins, notably serum amyloid P component (SAP).

Amyloid appears as an amorphous, eosinophilic, hyaline, extracellular substance. The most common organ involved in AL amyloidosis is renal tissue. Kidneys are the most frequently involved organ and are affected in 70–80% of patients. The heart is the second most commonly affected organ (50–60% of patients), and cardiac involvement is the leading cause of death from AL amyloidosis. Renal amyloidosis usually manifests as proteinuria, often in the nephrotic range, and is associated with hypoalbuminemia, secondary hypercholesterolemia, hypertriglyceridemia, and oedema or anasarca.

Note: The plasma cells in amyloidosis usually will be between 7% to 10%, higher plasma cells indicate poor prognosis.

Other options:

Option A: Fibrinoid necrosis is usually the result of the deposition of immune complexes and plasma proteins in the arterial wall. On biopsy, fibrinoid necrosis appears as bright pink and amorphous deposits in the vessel wall.

Option B: Coagulative necrosis is the most common form of necrosis which mostly is found to occur in solid organs (except the brain). The affected tissue has a preserved architecture and firm texture for several days. The injury leads to the denaturation of structural protein leading to eosinophilic cells with reddish nuclei and a glossy appearance due to loss of glycogen. The cells may also present with a moth-eaten vacuolated appearance due to digestion of organelles by lysosomal enzymes. The dead cell debris is ultimately removed by phagocytosis.

Option D: Bone marrow necrosis (BMN) and myxomatous stromal changes are rare histologic findings. These changes have been observed in several illnesses, including disseminated carcinomatosis, cases following chemotherapy, persistent infections, and marrow infiltrative disorders.

Solution to Question 20:

The given clinical scenario with features such as tearing chest pain, tachycardia, and hypotension with negative Trop I and T(indicating no myocardial infarction) with biopsy showing separation of the tunica intima and tunica media is suggestive of aortic dissection secondary to connective tissue damage of the aortic wall.

Dissection occurs when blood enters into the media to form a blood-filled channel between the laminar planes of the media. It usually occurs due to a tear in the intima which allows blood in the lumen to flow into the media. It may also occur due to the rupture of vasa vasorum.

It occurs due to:

• Hypertension

• Connective tissue defects like Marfan syndrome

• Iatrogenic trauma during coronary catheterization or cardiopulmonary bypass

Note: Fibrinoid necrosis, commonly seen in Malignant hypertension, Rheumatic carditis and Vasculitis - Polyarteritis nodosa, Kawasaki disease and not seen in the cases of aortic dissection, is usually the result of the deposition of immune complexes and plasma proteins in the arterial wall. On biopsy, fibrinoid necrosis appears as bright pink and amorphous deposits in the vessel wall.

Solution to Question 21:

This 10-year-old boy presents with fever and pain in his right upper limb, accompanied by an expansile lesion in the diaphysis of the radius, with the characteristic histopathology showing uniform, round blue cells arranged in Homer-Wright rosettes is suggestive of Ewing's sarcoma. The most accurate test to confirm this diagnosis is the detection of the EWSR1 (FUS-ETS fusion gene).

Ewing sarcoma is the 2nd most common bone sarcoma in children. The majority of the patients affected are usually <20 years of age. It has a slight male predominance. The majority of Ewing

sarcomas have a balanced translocation between the EWSR1 gene on chromosome 22 and the FLI1 gene on chromosome 11.

It commonly involves the diaphysis of the long bones like the femur, tibia, fibula, humerus, and flat bones of the pelvis. Patients present with painful enlarging mass, fever, and anaemia. The affected area is usually tender, swollen, and warm.

Laboratory findings may show an elevated WBC count, LDH levels (lactate dehydrogenase) and erythrocyte sedimentation rate (ESR). The common diagnostic test used for Ewing's sarcoma is the detection of translocation-EWSR1 (FUS-ETS fusion gene), t(11;22) (q24;q12).

Radiological examination of Ewing's sarcoma shows a destructive lesion in the diaphysis of long bones. Deposition of periosteal reaction in layers gives an onion-peel appearance.

A histopathological examination of Ewing sarcoma shows sheets of uniform, round blue cells arranged in Homer-Wright rosettes. Immunohistochemistry is usually positive for synaptophysin and mic-2. The tumor cells contain glycogen and are hence PAS positive and they are reticulin negative.



Fluorescent in situ hybridisation (FISH) demonstrates a positive EWSR1 gene rearrangement, characteristic of Ewing sarcoma. In the image given below, the fused region denoted by yellow corresponds to a normal intact gene, while a split signal pattern showing separate red and green signals is indicative of EWSR1 gene rearrangement.



Preoperative chemotherapy with vincristine, ifosfamide, doxorubicin and etoposide results in extensive tumour necrosis and shrinkage. Wide excision may be combined with adjuvant radiotherapy if surgical margins are poor. Definitive radiotherapy rather than surgical excision may be advocated if it is non-resectable or metastasis has occurred. It is a radio-sensitive tumour.

Indicators of poor prognosis in patients with Ewing's sarcoma:

• Presence of distant metastasis - worst prognostic factor

• Size and location of the primary lesion

• Presence of fever, anaemia

• Leukocytosis and raised ESR.

Solution to Question 22:

The chances of their offspring being affected by hemophilia, an X-linked recessive disorder, are that half of the males can be affected and half of the females can be carriers.

The couple's family history reveals that the mother is likely a carrier, given her family background of hemophilia, while the father's family is unaffected. Their male child has hemophilia, indicating he inherited the affected X chromosome from his mother. The normal female child may or may not be a carrier, depending on whether she inherited the affected X from her mother.

X-linked recessive disorders have the following features:

• Males are affected more commonly and more severely than females.

• Female carriers have a 25% risk of having an affected son, a 25% risk for a carrier daughter, and a 50% chance of having a child that does not inherit the mutated X-linked gene.

• The affected father will have all daughters (100%) to be carriers and unaffected sons because they pass their X chromosome to all of their daughters and their Y chromosome to all of their sons.



Hemophilia B, also known as Christmas disease, results from mutations in coagulation factor IX. In addition to hemophilia B, it's important to note hemophilia A, which is caused by a deficiency in factor VIII. Hemophilia A is also inherited in an X-linked recessive manner, presenting similar clinical features to hemophilia B, such as prolonged partial thromboplastin time (PTT) and normal prothrombin time (PT). Both conditions require different factor assays for accurate diagnosis, but both lead to increased bleeding risks. Treatment options, including recombinant factor infusions for both hemophilia A and B, are essential for managing the conditions and minimizing complications.

Solution to Question 23:

The given clinical scenario along with MRI showing intracranial mass with calcification at sella turcica and histopathology showing black motor oil point to a diagnosis of craniopharyngioma.

Craniopharyngiomas are tumors that arise from the remnants of Rathke's pouch. It is an intracranial space-occupying lesion. It has a bimodal age of presentation. Children usually present with frequent headaches, short stature, bitemporal hemianopsia (due to optic chiasm compression), and polyuria.

These tumors are usually large and heterogeneous, displaying both solid and cystic components, and occur within the suprasellar area. The cysts contain a cholesterol-rich, thick brownish-yellow fluid and are compared to machine oil. They also show radiologically demonstrable calcification.

Treatment of central diabetes insipidus caused by craniopharyngioma includes supplementation with vasopressin analogues desmopressin (DDAVP) and resection of the tumour.

Other options:

Option A: Glioblastoma affects people usually in the sixth or seventh decade. These tumours may arise de novo or may result from a malignant transformation of lower-grade astrocytomas. Mutations of isocitrate dehydrogenase (IDH), PTEN, and epidermal growth factor receptor gene (EGFR) are implicated in the pathogenesis. The majority of the tumours (90%) have IDH

wild-type mutation, and the remaining have IDH mutant type. Clinically, the patient presents with headaches, seizures due to raised intracranial pressure, or focal neurologic deficits, depending upon the site involved.

Radiologically, glioblastomas appear as a large ring-enhancing heterogeneous mass with central necrosis and surrounding oedema that crosses the corpus callosum into both frontal lobes. The image given below shows MRI findings seen in glioblastomas. Note the butterfly morphology of the tumour:



Option C: Meningiomas are predominantly benign tumors, usually attached to the dura. They are predominantly seen in females and in patients with neurofibromatosis type 2, which is an autosomal disorder associated with multiple meningiomas, schwannomas, and ependymomas. The image below shows a well-defined, hyper-intense, and homogeneously enhanced mass close to the midline suggestive of meningioma.



Option D: Pituitary adenomas are usually found in adults. The most common genetic alterations in this condition are mutations in G proteins (particularly GNAS). Depending on their size, they are classified as microadenomas(diameter < 1 cm) and macroadenomas(diameter > 1 cm). They may be functioning or non-functioning. The most common type of functioning pituitary adenomas are prolactinomas. The brain MRI showing a pituitary adenoma is given below:



Solution to Question 24:

Aβ amyloid deposition is seen in Alzheimer's disease.

Aβ constitutes the core of cerebral plaques found in Alzheimer's disease. It also gets deposited in the walls of cerebral blood vessels. The Aβ protein is derived by proteolysis from a much larger transmembrane glycoprotein, called amyloid precursor protein (APP).

Other options

Option A: Alpha-synuclein is an unfolded protein that aggregates in Lewy Body Dementia, particularly in the subcortical regions and in the hippocampus and parahippocampal gyrus.

Option C: GFAP is the marker for glial origin tumors like glioblastoma, and oligodendroglioma.

Option D: Albumin levels may rise during fasting and diarrhoea.

Solution to Question 25:

Activated partial thromboplastin time (APTT) is more effective than measuring clotting factors for assessing the intrinsic pathway, i.e. factors XII, XI, IX, VIII, X, V, II, and fibrinogen.

Causes of prolonged APTT:

• Disseminated intravascular coagulation

• Liver disease

• Massive transfusion with plasma-depleted red blood cells

• Administration of or contamination with heparin or other anticoagulants

• A circulating anticoagulant (inhibitor)

• Deficiency of a coagulation factor other than factor VII (since it is part of the extrinsic pathway)

• APTT is also moderately prolonged in patients taking oral anticoagulant drugs and in the presence of vitamin K deficiency

APTT is also moderately prolonged in patients taking oral anticoagulant drugs and in the presence of vitamin K deficiency (in which factors II, VII, IX, and X will fail to activate). A patient with previously undiagnosed haemophilia or another congenital coagulation disorder can also present with an isolated prolonged APTT.

Other options:

Option A: Prothrombin time evaluates the extrinsic pathway, i.e. factors VII, X, V, II (prothrombin), and fibrinogen. It measures the clotting time of plasma in the presence of an optimal concentration of tissue extract (thromboplastin).

Causes of prolonged PT:

• Administration of oral anticoagulant drugs (vitamin K antagonists)

• Liver disease, particularly obstructive jaundice

• Vitamin K deficiency

• Disseminated intravascular coagulation

• Rarely, a previously undiagnosed factor VII, X, V or prothrombin deficiency or defect

Isolated prolonged prothrombin time occurs due to the deficiency of clotting factor VII.

Option C: Fibrinogen levels are used for measuring clotting.

Option D: Complete blood count is used to measure total RBC count, WBC count, and Platelet count.

Solution to Question 26:

The given clinical scenario along with the peripheral smear showing acanthocytosis is suggestive of abetalipoproteinemia.

Abetalipoproteinemia is a rare autosomal recessive disease. It is caused due to the mutation in microsomal triglyceride transfer protein (MTTP). MTTP is necessary to transfer lipids to nascent chylomicrons in the small intestine and to VLDL in the liver. In abetalipoproteinemia, there is an absence of chylomicrons, VLDL, LDL (apoB containing lipoproteins), and deficient levels of plasma cholesterol and triglycerides.



In abetalipoproteinemia, the usual presentation is fat malabsorption, diarrhea, and failure to thrive in early childhood. Vitamin E deficiency secondary to abetalipoproteinemia can lead to spinocerebellar degeneration. It manifests as loss of deep tendon reflexes, progressive ataxia, and lower limb spasticity by adulthood.

Intestinal biopsy shows lipid-laden vacuolated epithelium due to triglyceride accumulation. Accumulated lipids disappear after infants stop taking feed. Early treatment with supplemental vitamins, especially vitamin E, may significantly slow the development of neurological sequelae.

Acanthocytes (spur cells) are spiculated red cells with a few projections of varying size and uneven surface distribution. The image below shows acanthocytes in the peripheral smear.



Other options:

Option A: Burns patient's peripheral smear will show spherocytes due to direct physical or chemical injury to the cells. The image below shows spherocytes.



Option C: Uremia patients present with nausea, vomiting, fatigue, anorexia, weight loss, muscle cramps, pruritus, mental status changes, visual disturbances, increased thirst, and peripheral smear will show burr cells/echinocytes due to anemia. Echinocyte or Burr cells are characterized by projections of the red cell membrane that are smaller, regularly spaced, smoothly rounded crenulations, and much more uniform in shape and distribution in contrast to the acanthocytes.



Option D: Ethylenediamine tetraacetic acid (EDTA) changes include structural, biochemical and functional damage to platelets. It also causes irreversible dissociation of the fibrinogen receptor, glycoprotein alpha IIb beta 3 (GPIIb/IIIa).

Solution to Question 27:

Flow cytometry is the gold standard for diagnosis and is the correct statement regarding PNH. It provides a sensitive means for detecting red cells that are deficient in GPI-linked proteins such as CD59.

Investigations in PNH include:

• Flow cytometry (gold standard) - deficiency of CD59 and CD55

• Ham test (acidified serum lysis) - lysis of RBCs in a mildly acidic pH due to complement fixation

• Bone marrow biopsy -

• Hypercellularity with erythroid hyperplasia is seen in classical presentations

• Patients who develop pancytopenia will have hypocellular bone marrow

• Imaging - to evaluate venous thrombosis

Paroxysmal nocturnal hemoglobinuria (PNH) is due to acquired mutations in the phosphatidylinositol glycan A gene (PIGA), leading to a deficiency of glycosylphosphatidylinositol (GPI)-linked proteins. Blood cells are deficient in three GPI-linked proteins that regulate complement activity making them susceptible to lysis or injury by complement. These are a decay-accelerating factor or CD55, membrane inhibitor of reactive lysis or CD59 (most important), and C8-binding protein.

Clinical features in paroxysmal intravascular hemolysis occur nocturnally in 25% of cases. Clinically, the patient would have hemolysis, pancytopenia, and a tendency for venous thrombosis. Intravascular hemolysis may be nocturnal and paroxysmal or chronic hemolysis without the dramatic hemoglobinuria. Some patients may complain of recurrent attacks of abdominal pain that may be related to thrombosis in abdominal veins.

Other options:

Option A: Thrombosis is the most common cause of death in persons with PNH and not anaemia with heart failure.

Option B: It is an acquired condition and not an inherited condition.

Option C: Sucrose lysis test is used for screening of PNH.

Solution to Question 28:

The given clinical scenario is suggestive of Marfan's syndrome and the likely protein involved in this condition is fibrillin.

Marfan's syndrome is an inherited connective tissue disorder that is caused by a mutation in a gene that codes for an extracellular matrix protein-fibrillin-1 (FBN1). The FBN1 locus is on the long arm of chromosome 15. It is a clinical diagnosis based on skeletal, cardiovascular, and ocular manifestations.

Skeletal manifestations:

• Overgrowth of long bones (dolichostenomelia) with an arm span to height ratio of more than 1.05

• Pectus carinatum or pectus excavatum

• Long and slender fingers (arachnodactyly), contracture of fingers (camptodactyly)

• Malar hypoplasia

• Joint laxity

Cardiovascular manifestations:

• Thickened atrioventricular valves, valvular prolapse (mitral regurgitation)

• Aortic valve dysfunction due to aortic root aneurysm

• Aortic dissection and rupture due to cystic medial necrosis of the aorta

Ocular manifestations:

• ectopia lentis

• myopia

• flat cornea

• retinal detachments

• early cataracts and glaucoma

Other systemic manifestations:

• thoracic scoliosis

• spontaneous pneumothorax

• striae atrophicae in the skin

Image (a) shows Walker-Murdoch (wrist) sign - the ability to join the thumb and fifth finger around the wrist. Image (b) shows Steinberg (thumb) sign - thumb tip extends from the palm of the hand in a closed fist

(a) (b)

Diagnosis is based on criteria depending on the family history, aortic root X score, ectopia lentis, mitral valve prolapse, and presence of FBN1 mutation.

Homocystinuria can be one of the differential diagnoses for this condition. It is an inborn error of methionine metabolism caused by a deficiency of the enzyme cystathionine β-synthase. It is characterized by tall stature, long limbs, dislocation of the lens of the eye, and predisposition to thromboembolism, but it does not lead to black urine.

Other options:

Option B: Collagen is a connective tissue. Defects in collagen lead to conditions like osteogenesis imperfecta, stickler syndrome, Ehler-Danlos syndrome, and Alport syndrome.

Option C: Elastin is a crucial protein found in connective tissue predominantly located in ligaments and the walls of blood vessels, especially large arteries like the aorta. The maturation of elastin involves the formation of desmosine cross-links from four lysine residues, contributing to its stability and low turnover rate. Williams-Beuren syndrome is an associated developmental disorder (deletions of the elastin gene) affecting connective tissue and the central nervous system. Fragmentation or a decrease of elastin is found in conditions such as pulmonary emphysema, cutis laxa, and ageing of the skin. Copper deficiency can also affect the formation of crucial cross-linking in elastin.

Option D: Laminin is a component of the basement membrane, providing structural support and anchoring the epidermis to the dermis. Mutations in its genes cause epidermolysis bullosa junctionale.

Solution to Question 29:

The clinical history along with histology showing palisade in a spoke-and-wheel arrangement, surrounding a central core(Flexner Wintersteiner rosettes) and the lesion being positive for synaptophysin, NSE, and chromogranin is suggestive of retinoblastoma.

Retinoblastoma is a malignant proliferation of immature retinal neural cells and is the most common intraocular malignancy in children. It is associated with mutations in the RB1 gene at chromosome 13q14. It is a tumor of a group called small round blue cell tumors because the immature retinal neural cells are seen as small round cells with large nuclei.

White pupillary reflex or leukocoria is the most common presenting feature, followed by a squint which is usually convergent. Rarely, the child may present with nystagmus and defective vision.

Image showing leukocoria of left eye.



Histopathology of the tumor shows small round cells with large nuclei. Microscopic features of a well-differentiated tumor include Flexner-Wintersteiner rosettes (highly specific of retinoblastoma), Homer-Wright rosettes, pseudorosettes, and fleurettes formation. Areas of necrosis and calcification can also be seen.

The histopathological image of retinoblastoma is shown below.



Treatment typically involves chemotherapy plus focal therapy. The commonly used three-drug regimen is the CVE regimen consisting of vincristine, carboplatin, and etoposide. Focal therapy includes cryotherapy, laser photocoagulation, thermotherapy, plaque radiotherapy, and external beam radiotherapy. Enucleation is done when:

- the tumor involves more than half of the retina
- the optic nerve is involved
- glaucoma is present and the anterior chamber is involved

In the stage of extraocular extension, the globe bursts through the sclera, usually near the limbus or the optic disc. This is followed by rapid fungation and involvement of extraocular tissues resulting in marked proptosis. Distant metastasis occurs through the lymphatic spread and the direct extension by continuity to the optic nerve and brain.

Other options:

Option B: Medulloblastoma is a malignant embryonal tumor of childhood that occurs exclusively in the cerebellum. Patients usually present with headaches, ataxia, and signs of brainstem involvement.

Histologically, the tumor is very densely cellular. Individual tumor cells have scant cytoplasm and elongated or crescent-shaped hyperchromatic nuclei. Tumor cells form Homer Wright rosettes. They express neuronal markers such as synaptophysin.



Option C: Pineoblastomas are tumours in the pineal region that are best described as small, round, blue-cell tumours. They closely resemble medulloblastomas and retinoblastomas. They are regarded as WHO grade 4 tumours and are the most aggressive and high-grade tumour among pineal parenchymal tumours. The below image shows tightly packed small round blue cells (high nuclear to cytoplasmic ratio).



Option D: Epidermoid carcinoma arises from ectodermal cells misplaced during the closure of the neural tube in the process of embryogenesis. There is an accumulation of keratin and cholesterol which gives the tumours their pearly or milky white appearance. A substantial region of keratin is encased in a thin layer of epithelial cells on the outside of an epidermoid tumour. Fatty acids and keratin are found inside the tumour, while stratified squamous epithelium, or normal skin cells, make up the outside of the growth.

Solution to Question 30:

This clinical scenario and image showing typical asbestos bodies are suggestive of that he has been likely to be working in the asbestos industry for 20 years.

Asbestosis is a type of pneumoconiosis associated with asbestos exposure. The disease-causing capabilities of the different forms of asbestos depend on concentration, size, shape, and solubility. This condition is marked by diffuse pulmonary interstitial fibrosis. Asbestos bodies are a typical finding. They are golden brown, fusiform, or beaded rods with a translucent center. They consist of asbestos fibers coated with an iron-containing proteinaceous material and arise when these fibers are phagocytosed by macrophages.

The image below shows asbestos bodies.



Asbestos occurs in two distinct forms - serpentine chrysotile (more common) and amphibole (more pathogenic). Exposure may result in the following conditions:

• Localized fibrous plaques or, rarely, diffuse pleural fibrosis

• Pleural effusions

• Parenchymal interstitial fibrosis (asbestosis)

• Lung carcinoma

• Mesotheliomas

• Extrapulmonary neoplasms including laryngeal, ovarian, and colon carcinomas

Other options

Option B: Silicosis is an occupational disease caused by exposure to silica or crystalline quartz. It is seen in people who work in mining, stone cutting, sandblasting, and quarrying. Silicosis presents as small rounded opacities in the upper lobe. There may be calcification of hilar lymph nodes which leads to the appearance of eggshell calcification. HRCT will show a characteristic crazy paving pattern due to interlobular septal thickening on the background of ground-glass opacity.

Option C: Anthracosis/coal workers' pneumoconiosis is a pulmonary condition seen in coal miners. The inhaled carbon pigment is engulfed by alveolar or interstitial macrophages, which accumulate in the connective tissue. On prolonged exposure, lung tissue undergoes progressive massive fibrosis, which is called complicated coal workers' pneumoconiosis. The upper lobes of the lungs are commonly involved.

Option D: Berylliosis is an occupational disease due to chronic exposure to beryllium. It is seen in people who work on processing alloys for high-tech industries like aerospace industries. Chest radiography shows nodules along the septal lines with hilar and mediastinal lymph node enlargement. HRCT will show pulmonary nodules, diffuse interstitial fibrosis, and interlobular septal thickening.

Solution to Question 31:

The clinical vignette with the lymph node biopsy showing a classical Reed-Sternberg cell is suggestive of a diagnosis of Hodgkin's lymphoma and an ABVD regimen (Doxorubicin (Adriamycin), Bleomycin, Vinblastine, and Dacarbazine) is used for the treatment.

The Reed Sternberg cell is a giant cell. It has a classic owl-eyed appearance due to the presence of a symmetric or mirror-image bilobed nucleus with prominent nucleoli surrounded by a clear space.



Hodgkin's lymphoma -

• It is of B-cell origin

• It has a bimodal distribution of age at diagnosis, with one incidence in young adults in their twenties and another incidence in older patients in their eighties.

• Typical presentation involves lymphadenopathy/mass with B symptoms which include fever, night sweats, and weight loss

• HIV infection is a risk factor. Another virus that has been implicated in the pathogenesis of Hodgkin's lymphoma is the Epstein-Barr virus (EBV). Further EBV can be detected in almost all cases of HIV-associated Hodgkin's lymphoma.

• Majority of the patients can be treated with chemotherapy and radiotherapy or chemotherapy alone. The most widely used chemotherapy regimen in Hodgkin's lymphoma is the ABVD regimen (Doxorubicin (Adriamycin), Bleomycin, Vinblastine, and Dacarbazine).

Other options

Option A and C: R-CHOP (Rituximab + Cyclophosphamide, Doxorubicin, Vincristine (Oncovin), and Prednisone) regimen is used in the treatment of Non-Hodgkin's lymphoma.

Option D: Sarcoidosis is a granulomatous inflammatory condition with multisystem involvement. The most common organs affected are the lungs, liver, eyes, and skin. The most widely accepted etiological hypothesis is an inflammatory response to an infectious or non-infectious trigger in a susceptible host.

Solution to Question 32:

The clinical scenario along with lab investigations showing microcytic hypochromic anemia is suggestive of iron deficiency anemia. Transferrin saturation is decreased in iron deficiency anemia.

Clinical presentation may include fatigue, pallor, reduced exercise capacity. Cheilosis (fissures at the corners of the mouth) and koilonychia (spooning of the fingernails) are signs of advanced tissue iron deficiency.

Laboratory findings are:

• Reduced hemoglobin and hematocrit values.

• hypochromia, microcytosis, and poikilocytosis.

• serum iron and ferritin levels are low.

• The total plasma iron-binding capacity that reflects the elevated transferrin levels is high.

• Low serum iron with increased iron-binding capacity results in a reduction of transferrin saturation to below 15%.

• RBC protoporphyrin levels are increased. Protoporphyrin is an intermediate in the heme synthesis pathway. Under conditions in which heme synthesis is impaired, protoporphyrin accumulates within the red cell.

• Bone marrow reveals a mild to moderate increase in erythroid progenitors. The absence of stainable iron in macrophages is a diagnostically significant finding.

Peripheral smear of microcytic hypochromic anemia:



Below is the schematic representation of iron absorption and transportation:



Solution to Question 33:

The given clinical profile of a woman with low hemoglobin, MCV, MCH, transferrin saturation, and high TIBC suggests a diagnosis of iron-deficiency anemia (IDA).

IDA is an example of microcytic hypochromic anemia. The causes of IDA can be inadequate diet, impaired reabsorption, increased requirement and increased blood loss.

The manifestations of IDA depend upon the underlying cause. In general, IDA presents with koilonychia, alopecia, atrophic changes in the tongue, gastric mucosa, malabsorption, and pica (craving for consumption of non-edible materials)

Peripheral smears show small, pale red cells typically described as microcytic (reduced MCV) and hypochromic (enlarged zone of central pallor). Poikilocytosis can be seen as elongated red cells, known as pencil cells. Target cells can be found in cases of severe IDA.

Diagnosis is mainly by laboratory studies. The typical findings are:

• Low hemoglobin and hematocrit

• Low serum iron and ferritin

• Increased Total Iron Binding Capacity (TIBC)

• Hepcidin is low (decreased stores of iron inhibit hepcidin synthesis)

IDA is treated by giving oral and parenteral preparations if iron or packed cell transfusion, depending on the anemia's severity.

Other options:

Option A: Megaloblastic anemia is defined by a high MCV

Option C: Hemolytic anaemia causes the body to produce more erythropoietin, which in turn causes erythroid differentiation and the marrow to produce more erythroid precursors or normoblasts. A noticeable reticulocytosis in the peripheral blood is the consequence of

compensatory increases in erythropoiesis. Hemosiderin, an iron-containing pigment, accumulates as a result of red cell phagocytosis, especially in the spleen, liver, and bone marrow.

Option D: Hypoplastic anaemia will have markedly hypocellular marrow. Aplastic anaemia is one of the most important examples. Marrow aspirate will show a dry tap.

Community Medicine AIIMS 2017

Question 1:

Which of the following wastes are disposed in the bag shown below?

a) Urine bags

b) Anatomical waste

c) Soiled waste

d) Sharp waste

Question 2:

A researcher discovered a new drug which is effective in chronic hypertensives with a p value of <0.10. which of the following is true regarding the same

a) Not more than 10% of the people benefitted by the drug could be due to chance

b) 90% of patients will be benefitted by giving the drug

c) The test is 90% reproducible

d) 90% of test results could have occurred by chance

Question 3:

Below is the graph of CD4 count in HIV patients. From the graph below identify the mean CD 4 count



a) 300-400

b) 200-300

c) Below 250

d) Above 350

Question 4:

Which of the following is true about triage?

a) Yellow – least priority

b) Red – mortality

c) Green – ambulatory

d) Blue – ambulatory

Question 5:

A randomized trial comparing the efficacy of two drugs showed a difference between the two with a p value of <0.005. In reality, both drugs do not differ at all. This is an example of:

a) Alpha error

b) Beta error

c) 1 – α

d) $1 - \beta$

Question 6:

Which of the following vaccine vials can be used based on the vaccine vial monitor (VVM) color changes shown?

a) Only vial 1 can be used

b) Vials 1 and 2 can be used

c) Vials 1, 2, and 3 can be used

d) All vials can be used

Question 7:

Under the Mid-Day Meal (MDM) scheme, supplementary nutrition provided should cover at least _____

a) 1/3 of the total protein requirement + ½ of total energy requirement

b) 2/3 of the total protein requirement + ½ of total energy requirement

c) ½ of the total protein requirement + 1/3 of total energy requirement

d) ½ of the total protein requirement + 2/3 of total energy requirement

Question 8:

In a subcentre area with a crude birth rate (CBR) of 20, what would be the expected number of antenatal care (ANC) registrations by the village health nurse (VHN) in a year?

a) 60

b) 80

c) 100

d) 110

Question 9:

As per the syndromic approach to sexually transmitted infections (STIs) and reproductive tract infections (RTIs), what is the treatment of choice of gonococcal and non-gonococcal urethritis?

a) Azithromycin + Metronidazole

b) Cefixime + Azithromycin

c) Azithromycin + Doxycycline

d) Doxycycline + Cefixime

Question 10:

Annual new case detection rate (ANCDR) of leprosy in India, as of 31st March 2016 was _____.

a) 0.66/10,000 population

b) 0.66/1 lakh population

c) 9.7/10,000 population

d) 9.7/1 lakh population

Question 11:

The recommended calorie intake for a pregnant woman in the second trimester, who is a sedentary worker would increase by _____

a) 600 kcal/day

b) 520 kcal/day

c) 450 kcal/day

d) 350 kcal/day

Question 12:

Match the following declarations to their significance.

Declaration	Significance
I. Geneva	A. Guidelines for physicians concerning torture and other cruel, inhuman or degrading treatment or punishment in relation to detainment or imprisonment
II. Oslo	B. Set of ethical principles directed to the medical community for human experimentation.
III. Tokyo	C. Physician's dedication to the humanitarian goals of medicine
IV. Helsinki	D. Statement on therapeutic abortion

a) I – A, II – B, III – C, IV – D

b) I - C, II - D, III - A, IV - B

c) I – D, II – C, III – A, IV - B

d) I – B, II – A, III – D, IV - C

Question 13:

Chronic liver disease is classified using the Child-Pugh score as follows: Class A: 5-6 , Class B: 7-9, Class C: 10-15. What type of data is this scoring system?

a) Nominal

b) Ordinal

c) Qualitative

d) Metric

Question 14:

An 18 month old child is brought with no history of vaccination. From the following, which vaccines would be appropriate to give this child according to the National Immunization

Schedule (NIS), India?

a) DPT -1, OPV-1 only

b) BCG, DPT-1, OPV-1

c) BCG, DPT-1, OPV-1, Measles – 1, Vit. A -1

d) DPT-1, OPV-1, Measles – 1, Vit. A – 1

Question 15:

P-value is the probability of which of the following?

a) Not rejecting a null hypothesis when it is true

b) Rejecting a null hypothesis when it is true

c) Not rejecting a null hypothesis when it is false

d) Rejecting a null hypothesis when it is false

Question 16:

While testing the deleterious health hazards on a group of people after giving a new drug and a placebo, the following results were obtained as tabulated. Based on these results, which of the following is true?

a) Absolute risk reduction (ARR) is 1% and relative risk reduction (RRR) is 0.1

b) Absolute risk reduction (ARR) is 10% and relative risk reduction (RRR) is 0.9

c) Absolute risk reduction (ARR) is 1% and relative risk reduction (RRR) is 10

d) Absolute risk reduction (ARR) is 10% and relative risk reduction (RRR) is 9

Question 17:

A new drug combination of chemotherapy and immunotherapy for metastatic melanoma was prescribed by a radiotherapist. It prolongs the survival. Which of the following is true in this situation?

a) Incidence increases and prevalence reduces

b) Incidence reduces and prevalence remains the same

c) Incidence remains the same and prevalence increases

d) Incidence reduces and prevalence increases

Question 18:

Observe the following curves, which of the following is a true statement about sensitivity and specificity when the curve changes from green to pink?



a) Both sensitivity and specificity increase

b) Both sensitivity and specificity decrease

c) Sensitivity increases while specificity decreases

d) Sensitivity decreases while specificity increases

Question 19:

A study was conducted to compare the incidence of postpartum depression in mothers as compared to the sex of the child. For assessment, Edinburg Depression scaled score (EDSS) was used. The study was divided into mothers with male babies as one group to be compared with mothers with female babies. Which of the following statistical test will be used to determine the association?

a) Students t test

b) Paired t test

c) ANOVA

d) Chi square test

Question 20:

A one-and-half-year-old child was brought to the hospital with a history of breathlessness, fever, cough, and cold. On examination, respiratory rate was 46/min, and there were no chest retractions. Which is the most appropriate management?

a) Give IV antibiotics and observe

b) Give IV antibiotics and refer to higher centre

c) Oral antibiotics, warn of danger signs, and send home

d) No need of treatment only home remedies

Question 21:

The following are the disease frequencies in a population. Based on the number of cases which of the following is the correct match?

a) Pandemic, endemic, sporadic

b) Sporadic, endemic, epidemic

c) Endemic, epidemic, sporadic

d) Endemic, sporadic, epidemic

Question 22:

Below is a diagram depicting the natural history of the disease. Which point marks the onset of symptoms?



a) Point A

b) Point B

c) Point C

d) Point D

Question 23:

A researcher finds out a sensitive test to be positive for a person. For diagnosis, which of the following will hold true?

a) It confirms that the person does not have the disease

b) It suggests that if the disease is rare, the person surely has the disease

c) It confirms that the person has the disease if the disease is common

d) It is an expected finding in case the disease is prevalent

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	c
5	a
6	b
7	c
8	a
9	b
10	d
11	d
12	b
13	b
14	d
15	b
16	a
17	c
18	a

19	d
20	c
21	c
22	c
23	c

Detailed Explanations

Solution to Question 1:

The above picture shows a red coloured bag which is used to dispose contaminated wastes (recyclable) such as tubing, bottles, intravenous tubes and sets, catheters, urine bags, syringes (without needles) and gloves.

Option B: Anatomical wastes go into yellow non-chlorinated plastic bags.

Option C: Soiled wastes go into yellow coloured non-chlorinated plastic bags.

Option D: Sharp wastes go into white puncture proof, leak proof, tamper proof containers.

Solution to Question 2:

P value <0.10 indicates that not more than 10% of people benefitted by the drug could be due to chance.

It can also be interpreted as

• If the test is repeated 100 times, the difference will be statistically non significant only 10 times and difference will be statistically significant 90 times.

• Probability of false positive conclusion that drug is beneficial which in reality is not, is 10%.

It arises in case of statistical studies to accept or reject a null hypothesis.

P value is compared with a predetermined value α; P value < α means the test is statistically significant and null hypothesis is rejected.

Solution to Question 3:

The given data should be converted into numerical table. Small approximations can be made while converting into numerical form.

Since intervals are given, midpoint of each interval has to be calculated.

It should then be multiplied with frequency of that particular interval.

Mean is given by the formula: Mean = Sum of all observations / number of observations.

Mean = 34500/100 = 345.

Between 300 -400 is the most suitable answer.

CD4 Count	Number of people
<100	5
100 - 200	10
200 - 300	25
300 - 400	35
400 - 500	20
>500	5

CD4 Count	Mid point	Number of people (n)	Mid point × n
0 - 100	50	5	250
100 - 200	150	10	1500
200 - 300	250	25	6250
300 - 400	350	35	12250
400 - 500	450	20	9000
500 - 1600	1050	5	5250
TOTAL		100	34500

Solution to Question 4:

In triage, green signifies low priority or ambulatory patients and those with minor illnesses.

An internationally accepted four-color coded system, with red having the highest priority is used commonly here.

Solution to Question 5:

• In this scenario, let's take the null hypothesis as the two drugs being the same, which the questions states, is true in reality.

• The alternative hypothesis will then be that both drugs are different.

• The questions states that p value was found to be < 0.005, which is statistically significant and so the null hypothesis is rejected.

• But we already know that the null hypothesis is true and we are forced to accept the alternative hypothesis even though it is false.

• This is a type I error or alpha error.

• When an alternative hypothesis is accepted even when it is false and the null hypothesis is true, it is a type I error or false positive error.

Solution to Question 6:

From the above image, only vials 1 and 2 can be used as the inner squares of the VVMs are lighter than the outer circles.

Vial 3 has crossed the discard point as the inner square and outer circle are the same color and thus cannot be used. Vial 4 cannot be used as the VVM shows a darker inner square compared to the outer circle.

Solution to Question 7:

Under the Mid-Day Meal (MDM) scheme, supplementary nutrition provided must cover at least ½ of the total protein requirement and 1/3 of total energy requirement to children in schools.

The MDM scheme is also called School Lunch Programme and has been operational since 1961 in India. Currently, it functions under the broad coverage of the Minimum Needs Programme.

Some other principles of the MDM are:

• The meal provided under the MDM should only be a supplement and not a substitute to home diets.

• The cost of the meals should be reasonably low.

• The meal should be easy to cook in schools.

• Locally available foods must be used, for as far as possible.

• The menu should be regularly changed to avoid monotony.

Solution to Question 8:

• According to the Indian Public Health Standards (IPHS) laid down by the Government of India, a subcentre is to serve a population of 5000 in plains.

• In this scenario, the CBR is 20.

• Probable number of pregnancies (Y) = Birth rate x population of the area / 1000 = 20 x 5000/1000 = 100.

• But, we add a correction factor of 10% to adjust for pregnancy wastage in the form of abortions or stillbirths.

• Thus, total number of expected pregnancies= Y + 10% of Y= 110

• As all the pregnancies are not registered, and at any given time an ANM should have atleast half the expected pregnancies registered with her, the number of ANC registrations is taken as 50% of the expected pregnancies.

• Therefore, Total ANC registrations = total number of expected pregnancies/2 = 55

• Hence, the closest answer is 60.

Solution to Question 9:

As per the syndromic approach to STIs/RTIs, the treatment of choice of for gonococcal and non-gonoccocal urethritis is azithromycin + cefixime.

Solution to Question 10:

For the year 2015 – 2016, the annual new case detection rate (ANCDR) of leprosy in India was 9.7/100,000 population.

Solution to Question 11:

The recommended calorie intake for a pregnant woman who is a sedentary worker would increase by ~ 350 kcal/day.

Solution to Question 12:

Declaration	Significance
Geneva	Physician's dedication to the humanitarian goals of medicine.
Oslo	Statement on therapeutic abortion.

Declaration	Significance
Tokyo	Guidelines for physicians concerning torture and other cruel, inhuman or degrading treatment or punishment in relation to detainment or imprisonment.
Helsinki	Set of ethical principles directed to the medical community for human experimentation.

Solution to Question 13:

The Child-Pugh score is an ordinal scale of data as:

There is a rank order (Class A having a better prognosis than classes B and C) but no regular interval between each class. The final division of prognosis is based on class and not the numerical values arrived at by the summation of a set of variables.

Other options:

Option A: The Child-Pugh score cannot be a nominal scale of data as there is a rank order with class A having better survival statistics than B or C. Nominal data do not have a rank order.

Option C: This is also not merely a qualitative scale as there are scores given to each variable to arrive at the final class.

Option D: The data used in the Child-Pugh score cannot be merely metric as the final division of prognosis is based on class and not the numerical values. Also, there is no definite interval between the classes.

Solution to Question 14:

• An 18 month old, completely un-immunized child should receive first doses of DPT, OPV, measles vaccine and vitamin A, according to the National Immunization Schedule (NIS), India.

• BCG can be given only up to one year of age.

Solution to Question 15:

p-value is the probability of rejecting a null hypothesis when it is true.

The p-value is the criterion or significance level (α), below which an outcome can be said to be statistically significant. From this, we can also deduce that the p-value is the probability that a null hypothesis will be rejected even when it is true, thereby making a type I error.

Solution to Question 16:

In the given scenario, absolute risk reduction (ARR) is 1% and relative risk reduction (RRR) is 0.1.

Let us reconstruct the design table in the following way:

Relative risk (RR) is given as a/(a+b) ÷ c/(c+d).

RR = 1603/15225 ÷ 1804/15225 = 0.106/0.118 = 0.89 ≈ 0.90.

Relative risk reduction (RRR) = 1 – Relative risk. In this scenario, RRR = 1 – 0.9 = 0.1 or 10%.

Incidence of a disease is the absolute risk of contracting it.

Absolute risk rate of the group using new drug = a/(a+b) = 1603/15225 = 0.106 or 10.6% ≈ 11%.

Absolute risk rate of the group using placebo = c/(c+d) = 1804/15225 = 0.118 or 11.8% ≈ 12%.

Absolute risk reduction (ARR) is the difference in the absolute risk rates between the placebo and the drug. In this scenario, ARR = 12% - 11% = 1%.

	Health hazard (+)	Health hazard (-)	Total
New drug	1603 (a)	13622 (b)	15225 (a +b)
Place bo	1804 (c)	13421 (d)	15225 (c+ d)

Solution to Question 17:

By the new drug combination, the survival of the patient has increased, hence incidence remains the same while prevalence increases.

Incidence: Number of new cases in a defined population in a specified period of time.

Prevalence: Number of old and new cases at a given point of time over a period of time.

Options A, B, D: The treatment only prolongs the disease, it has nothing to do with the occurrence of cases thereby nothing to do with incidence.

Solution to Question 18:

Both sensitivity and specificity are higher in the test represented in pink (Test B) compared to the test represented in green (Test A).

Sensitivity is the proportion of all people with the disease who test positive. It is the probability of the test being positive when the disease is present.

Sensitivity = TP / (TP + FN) or Sensitivity = 1 – false-negative rate.

Specificity is the proportion of all people without the disease who test negative. It is the probability of the test being negative when the disease is absent.

Specificity= TN / (TN + FP) or specificity= 1 – false-positive rate.

Test B has an increased number of true positives and true negatives and decreased number of false positives and negatives, compared to Test A.

Hence leading to increased sensitivity and specificity.

Solution to Question 19:

The Chi-square test is used to compare qualitative data and compares 2 groups with independent measures. The variables used here are yes/no signifying the presence/absence of postpartum depression. Hence it is the most appropriate test here.

Option A: Student's t-test is for quantitative data. Here the data is qualitative, hence it can't be used.

Option B: Paired t-test compares paired data or pre-intervention and post-intervention data of the same group. This doesn't apply here as the groups are not paired.

Option C: ANOVA is used to compare more than 2 groups.

Solution to Question 20:

From the given clinical scenario, it can be easily inferred that the child has pneumonia. The treatment for pneumonia for a child between 2 months and 5 years of age is oral amoxicillin and home care advice.

The following image gives the IMNCI classification for cough/difficulty in breathing in children.



Solution to Question 21:

Disease 1- the cases have been almost the same so it has to be endemic.

Disease 2- there is an alarming increase in the number of cases. It has to be an epidemic.

Disease 3- the number of cases has decreased without any explanation so it has to be sporadic.

Epidemic refers to the occurrence of a disease in excess of expected frequency.

A pandemic is an epidemic affecting a large population over a large geographical area crossing international boundaries.

An endemic disease is a regularly occurring disease in a population in expected frequency without any importation from outside.

Sporadic diseases present as scattered cases of infection without relation temporally/spatially.

Solution to Question 22:

Point C represents a point where signs and symptoms appear.

Various points marked in the image represent the following:

Point A: Entry of infectious agent into the host.

Point B: Pathological and physiological changes occur in the stage of the subclinical phase. These changes are induced by the causative agent but are not sufficient enough to cause clinical signs and symptoms.

Point C: Marks onset of clinical signs and symptoms.

Point D: Usual time of diagnosis at which clinical signs and symptoms are well evident.



Solution to Question 23:

A sensitive test, if it is positive it confirms that the person has the disease if the disease is common.

The sensitivity of a screening test is the ability of the test to identify correctly those who have the disease. In other words, Sensitivity is the ability of a test to detect true positives (TP) out of total disease (TP+FN).

Sensitivity = TP/ TP+FN

Community Medicine AIIMS 2018

Question 1:

A patient from a North-Eastern state was diagnosed to have an infection with P. falciparum malaria. What is the most appropriate treatment for this patient?

a) Artemether plus lumefantrine

b) Sulfadoxine plus pyrimethamine

c) Choroquine

d) Mefloquine

Question 2:

Choose the correct order of steps of disaster management.

a) Disaster impact – Mitigation – Rehabilitation – Response

b) Disaster impact – Response– Rehabilitation – Mitigation

c) Rehabilitation – Response – Disaster Impact – Mitigation

d) Response – Disaster impact – Mitigation – Rehabilitation

Question 3:

Chi square test is used to test the significance of association between:

a) Age and serum cholesterol level

b) Heart rate and body mass index

c) Sex and type of cancer

d) Weight of a patient before and after a treatment

Question 4:

Sample sizes of samples 1, 2 and 3 are 1000, 800, 600 respectively. The order of margin of error will be _____.



a) 3 >2>1

b) 3>1>2

c) 1>2>3

d) 1=2=3

Question 5:

Which of the following is a heat-sensitive indicator of vaccine viability?

a) e-VIN

b) Freeze indicator

c) Vaccine vial monitor

d) Fridge indicator

Question 6:

Which of the following is set up for health planning at the village level?

a) Panchayat Health Committee

b) Village Health Planning and Management Committee

c) Rogi Kalyan Samiti

d) Village Health Sanitation and Nutrition Committee

Question 7:

You have gone to a subcenter as part of an audit. How many infants should be registered in a year with a health worker working there?

a) 50

b) 100

c) 150

d) 200

Question 8:

As per the national immunization schedule, which vaccines are indicated at the first visit for immunization of a 13-month-old baby who has not been previously immunized?

a) BCG, OPV 1st dose, Pentavalent 1st dose, Rota virus 1st dose

b) OPV 1st dose, Pentavalent 1st dose, Measles 1st dose

c) BCG, OPV 1st dose, Measles 1st dose

d) OPV 1st dose, DPT 1st dose, Measles 1st dose

Question 9:

Regarding the given box-whisker plot, the true statement is:



a) Mean > Median > Mode

b) Mode > Median > Mean

c) Mode = Mean = Median

d) Peaked symmetrical distribution

Question 10:

A randomised control trial is being conducted in patients with direct inguinal hernia, to compare the outcomes of a new surgery 'A' and the gold standard surgery 'B'. The p value of this trial is found to be 0.04. What can we conclude from this?

a) Type II error is small and we can accept the findings of the study

b) The probability of false negative conclusion that operation A is better than B is 4%

c) The power of the study to detect the difference between operation A and B is 96%

d) The probability of a false positive conclusion that operation A is better than B is 4%

Question 11:

A large number of tests were done in series. Which of the following will be true?

a) Sensitivity and specificity will increase

b) Sensitivity will decrease, specificity will increase

c) Sensitivity and specificity will decrease

d) Sensitivity will increase, specificity will decrease

Question 12:

Blood bag is to be disposed in:

a) Red bin

b) Yellow bin

c) Blue bin

d) Black bin

Question 13:

Phase 1 of a clinical trial is done to test:

a) Safety

b) Efficacy

c) Dose range

d) Pharmacoeconomics

Question 14:

A child was brought to you with complaints of fever, malaise and a rash. You diagnose the child with measles. What is the recommended period of isolation for this disease?

a) From onset of rash to 3 days after rash

b) 2 weeks before rash to 1 week after rash

c) From onset of rash to 5 days after rash

d) From catarrhal stage to 5 days after rash

Question 15:

BCG vaccine provides protection against which of the following?

a) Pulmonary TB

b) CNS and disseminated TB

c) Pulmonary and CNS TB

d) Pulmonary and skeletal TB

Question 16:

In a community, there are usually 40-50 reported cases of dengue every week. This week, there were 48 reported cases. This would classify as:

a) Epidemic

b) Endemic

c) Sporadic

d) Outbreak

Question 17:

Which of the following vaccines has not been introduced to the immunization programme in India?

a) Adult Japanese encephalitis

b) Influenza

c) Measles-Rubella

d) Rotavirus

Question 18:

The life cycle of which of the following is depicted in the image below?

a) Japanese encephalitis

b) Zika virus

c) Rabies

d) Nipah virus

Question 19:

While collecting blood for samples, you have spilled some on the floor. What would be the immediate next step?

a) Call the infection control team of the hospital

b) Mop it

c) Cover it with 1% sodium hypochlorite

d) Clean with absorbable material

Question 20:

The following drug packet is used in:

a) TB

b) Leprosy

c) Vaginal discharge

d) Gonorrhea

Question 21:

The symbol shown in the image depicts:

a) Cytotoxic waste

b) Biomedical waste

c) Radiation hazard

d) Biohazard

Question 22:

The Mid Day Meal Programme aims to supplement:

a) 1/2 protein and 1/3 calorie requirements

b) 1/3 protein and 1/3 calorie requirements

c) 1/2 protein and 1/2 calorie requirements

d) 1/3 protein and 1/2 calorie requirements

Question 23:

What test is used to check for freeze damage to vaccines?

a) Shake test

b) Schick test

c) Habel test

d) Tine test

Question 24:

Study A(red) and Study B(blue) done in a given population yields two different distribution curves as shown below. Comparing both the curves, which of the following will be true?



a) Mean = mode = median

b) Mean = median, not equal to mode

c) Mean, median, mode not equal

d) Mode = median, not equal to mean

Question 25:

A clinical trial was done between two groups to measure their blood pressure after treatment. Which of the following tests would be appropriate to use?

a) Paired t test

b) Unpaired t test

c) ANOVA

d) Chi-square test

Question 26:

Maternal mortality ratio (MMR) for a country is best given as _____

a) Maternal deaths per 100 live births

b) Maternal deaths per 10,000 live births

c) Maternal deaths per 100,000 live births

d) Maternal deaths per 1,000 live births

Question 27:

SPIKES protocol is for:

a) Triage

b) Consent for study participation

c) Hospital infection control

d) Breaking bad news

Question 28:

According to the WHO Mental Health Gap Action Programme, which of the following is not included in the Mental Health Care Act, 2017?

a) Screening of family members

b) Mobilization of resources

c) Social support

d) Interpersonal therapy

Question 29:

Which of the following is the appropriate schedule for the treatment of multi-drug resistant tuberculosis?

a) 6 drugs for 4-6 months followed by 4 drugs for 12 months

b) 7 drugs for 4-6 months followed by 4 drugs for 5 months

c) 4 drugs for 4 months followed by 6 drugs for 10-12 months

d) 5 drugs for 6 months followed by 4 drugs for 12 months

Question 30:

The POCSO Act is related to which of the following?

a) Women empowerment

b) Human trafficking

c) Juvenile delinquency

d) Child sexual abuse

Question 31:

Droplet precaution and isolation is to be done for all of the following conditions except:

a) Pertussis

b) Diptheria

c) Localized varicella zoster

d) Mycoplasma pneumonia

Question 32:

A patient presents with fever for 1 week, with abdominal distension, loss of appetite, and lymphadenopathy. There is a poor response to antibiotics and antimalarials. The Widal test is negative but rk39 is positive. Which drug can be used in the treatment of this patient?

a) Liposomal Amphotericin B

b) Fluconazole

c) Linezolid

d) Bedaquiline

Answer Key

Question No.	Correct Option
1	a
2	b
3	c
4	a
5	c
6	d
7	c

8	d
9	a
10	d
11	b
12	b
13	a
14	d
15	b
16	b
17	b
18	d
19	d
20	b
21	c
22	a
23	a
24	b
25	b
26	c
27	d
28	a
29	b
30	d
31	c
32	a

Detailed Explanations

Solution to Question 1:

The most appropriate treatment for Plasmodium falciparum infection in North-Eastern states is artemether (20 mg) plus lumefantrine (120 mg).

As per National Vector Borne Disease Control Program (NVBDCP):

• Plasmodium malariae: treat as Plasmodium falciparum

• Plasmodium ovale: treat as Plasmodium vivax

• Severe and complicated malaria

-Initial parenteral treatment x 24-48 hr: quinine/artemether/artesunate/arteether

-Oral treatment after 48 hrs-after parenteral quinine: quinine + doxycyclinex 7 days or quinine + clindamycin x 7 days (pregnancy and chidren<8yrs)

-After parenteral artemisinin derivates: in NE states: ACT -AL X 3 days+primaquine (day2), in other states: ACT-SPX 3 days+primaquine (day 2)

• Chemoprophylaxis

Short-term, less than or equal to 6 weeks: Doxycycline OD (start one day before travel and continue for 4 weeks after return)

Long-term, more than 6 weeks: mefloquine weekly (start 2 weeks before travel and 4 weeks after return).

Solution to Question 2:

The correct sequence of steps is disaster impact – response – rehabilitation – mitigation.





Solution to Question 3:

Chi square test is a nonparametric test that assesses the significance of association between two qualitative variables.

Among the given options, sex and type of cancer is the only pair of qualitative variables. Hence the answer.

Solution to Question 4:

Curve 1 has the least margin of error, followed by curve 2, and then curve 3. So the order is 3>2>1.

The image shows three normal distribution curves of different sample sizes, having the same mean but different standard deviations.

As the sample size is decreasing, the peak decreases and the curve becomes more flat, implying a wider confidence of interval for the given mean.

The width of the base of the curve is directly proportional to the margin of error.

Solution to Question 5:

Vaccine vial monitors (VVM) are the only heat-sensitive indicator of vaccine viability that are routinely used in vaccines throughout the supply chain.

VVM records cumulative heat exposure through a gradual change in color. If the color of the inner square is the same or is darker than the outer circle, the vaccine has been exposed to too much temperature and should be discarded.

Solution to Question 6:

Village Health Sanitation and Nutrition Committee (VHSNC) has been formed to take collective actions on issues related to health and its social determinants at the village level (local level community action).

The committee is formed at the revenue village level and it should act as a sub-committee of the gram panchayat. It should have a minimum of 15 members and ASHA residing in the village shall be the member secretary and convener of the committee.

Solution to Question 7:

150 infants should be registered in a subcentre per year.

Annual infant population = (total population) x (% infants in population or expected birth rate)

The percentage of infants in the population, or the expected birth rate, should be obtained from local data. If a specific local percentage is not available, it is estimated to be 3% of the total.

Total population in a subcenter is 5000.

Here, annual infant population = 5000 × 3/100 = 150

Solution to Question 8:

DPT-1, OPV-1 and Measles 1st dose will be given to this child.

BCG can only be given upto 1 years of age, according to the National Immunisation Schedule. BCG will not be given to this child, if the National Immunisation Schedule is followed.

Please note that BCG can be given upto 5 years of age, as part of the catch-up immunization schedule, which is put forth by the Indian Academy of Paediatrics and that is not followed in government setups.

OPV can be given till 5 years of age.

Measles can be given till 5 years of age.

DPT can be given till 7 years of age (with booster doses).

Solution to Question 9:

The given box-whisker plot represents a positively (or right) skewed distribution. In a positively skewed distribution, the mean > median > mode.

To further understand and correlate the fact, have a look at the following images:





Solution to Question 10:

Since the p-value of the study is found to be 0.04, we can infer that the probability of a false positive conclusion that the operation A is better than B is 4%.

In this scenario, the p value is 0.04 which makes it, by convention, statistically significant. We can reject the null hypothesis and accept the alternative hypothesis that operation A is better than operation B.

Solution to Question 11:

When a large number of tests are done in series, net sensitivity is decreased and net specificity is increased.

Screening tests are done in series when the result of the first test determines whether to run the second test or not. e.g., HIV screening is done using consecutive ELISA.

Screening tests are done in parallel when the second test is independently performed regardless of what the first test result is. Both the tests are done at the same time. For e.g., In LFT, multiple tests are done together like SGOT, SGPT, total bilirubin, alkaline phosphatase, etc.

Following are the trends that are observed while evaluating screening tests:

	Tests in Series	Tests in parallel
Combined sensitivity	Decreases	Increases
Combined specificity	Increases	Decreases
Combined PPV	Increases	Decreases
Combined NPV	Decreases	Increases

Solution to Question 12:

Blood bag should be disposed off in the yellow bin.

Solution to Question 13:

The phase I of clinical trial is done to test Safety.

Solution to Question 14:

The isolation period for measles is from the catarrhal (prodromal) stage to 5 days after the onset of the rash.

The period of communicability is approximately 4 days before and 5 days after the appearance of the rash. Isolation of the patient for a week from the onset of rash adequately covers the period of communicability.

Solution to Question 15:

BCG vaccine provides protection against CNS and disseminated TB.

It does not prevent primary infection. More importantly, it does not prevent reactivation of latent pulmonary infection, the principal source of bacillary spread in the community. The impact of BCG vaccination on transmission of mycobacterium tuberculosis is therefore limited.

Solution to Question 16:

The given scenario is an example of an endemic disease.

Endemic diseases are those that have a constant presence and /or usual prevalence of a disease or infectious agent in a population within a geographic area.

Epidemic refers to an increase, often sudden, in the number of cases of a disease above what is normally expected in that population in that area.

Outbreak carries the same definition of an epidemic, but is often used for a more limited geographic area.

Sporadic diseases occur in a haphazard and irregular manner from time to time.

Solution to Question 17:

Influenza vaccine (Flu shot) has not been introduced to the National Immunization Programme in India.

The Influenza vaccine provides protection against viral flu caused by the influenza virus.

Note: Bacterial Hemophilus influenzae type-b (Hib) vaccine is different from the viral influenza (Flu) vaccine. HiB is a part of the pentavalent vaccine.

Solution to Question 18:

The given image depicts the life cycle of the Nipah virus.

Nipah virus is an RNA virus that belongs to the family Paramyxoviridae, genus Henipavirus. It is a zoonotic virus that shows both animal-to-human (bats, pigs) and human-to-human transmission. The natural hosts are fruit bats of the Pteropodidae family.

The clinical manifestations include asymptomatic subclinical infection, acute respiratory infection, and fatal encephalitis. There is no treatment or vaccine available for either people or animals. The primary treatment for humans is supportive care.

Nipah virus is on the WHO list of Blueprint priority diseases.

Solution to Question 19:

The immediate next step in this scenario is to clean the spilled blood with absorbable material.

This absorbable material is then discarded in the yellow bin. Following this, sodium hypochlorite solution is used to decontaminate the surface. The concentrations of sodium hypochlorite used are:

• < 10 mL blood spill - 1:100 dilution to decontaminate non-porous surfaces

• > 10 mL blood spill or culture spill - 1:10 dilution for the first application, followed by 1:100 dilution

Solution to Question 20:

The given image shows the adult blister packet used in the treatment of leprosy.

Solution to Question 21:

The black propellers on a yellow background is the logo for radiation hazard.

This symbol is used to let everyone be aware of the hazard in an area where there is a source of radiation. This indicates the need for precautionary measures to avoid contamination.

The other logos representing cytotoxic waste and biohazard are given in the image below.

Solution to Question 22:

Under the Mid-Day Meal (MDM) scheme, supplementary nutrition provided must cover at least 1/2 of total daily protein requirement and 1/3rd of total daily energy requirement of children in schools.

The MDM scheme is also called School Lunch Programme and has been operational since 1961 in India. Currently, it functions under the broad coverage of the Minimum Needs Programme.

Some other principles of the MDM are:

• The meal provided under the MDM should only be a supplement and not a substitute for home diets.

• The cost of meals should be reasonably low.

• The meal should be easy to cook in schools.

• Locally available foods must be used, for as far as possible.

• The menu should be regularly changed to avoid monotony.

Note: In September 2021, the Mid-Day Meal Scheme was renamed PM POSHAN or Pradhan Mantri Poshan Shakti Nirman.

Solution to Question 23:

The freeze damage to vaccines is detected by Shake test.

• A frozen vaccine is used as a control vial and the vial suspected to be damaged by freezing is taken as the test vial.

• Both are shaken after thawing and the rate of sedimentation is compared.

• If the rate of sedimentation in test vial is slower than the control vial, then it is usable and has not been damaged.

• Some of the most widely used vaccines are freeze-sensitive. Examples: Diphtheria, Tetanus, Pertussis, Liquid Haemophilus influenza type b (Hib), Hepatitis B.

Solution to Question 24:

On comparing the 2 graphs, we can say mean=median for both study A and B but the mode is different.

Study A is a unimodal symmetrical distribution, so mean=median=mode.

Study B is a bimodal distribution. Here, mean=median, but the curve has two highest points, therefore it has 2 modes.

Solution to Question 25:

In the given scenario, two groups are being compared independent of each other.
Hence, students t test/ unpaired t test should be used.

T tests are used if 2 groups are being compared. If the same group is compared before and after, paired T test is used. Otherwise, unpaired T test is used.

Solution to Question 26:

Maternal mortality ratio is the number of maternal deaths in a given period per 100,000 live births during the same period.

Solution to Question 27:

SPIKES protocol is for breaking bad news.

Each letter represents a phase in the six-step sequence.

• S - setting

• P - perception

• I - invitation or information

• K - knowledge

• E - empathy

• S - summarizing or strategizing

Key components of the SPIKES strategy include:

• Demonstrating empathy

• Acknowledging and validating the patient's feelings

• Exploring the patient's understanding and acceptance of the bad news

• Providing information about possible interventions

Solution to Question 28:

According to the WHO Mental Health Gap Action Programme (mhGAP), screening of family members is not included in the Mental Health Care Act, 2017.

The Mental Healthcare Act of 2017 has provisions that include:

• Rights of persons with mental illness to access health care, live without discrimination and enjoy confidentiality.

• Advance directives to allow mentally ill people to make decisions about their treatment

• Procedures for admission, treatment (behavioral and interpersonal therapy, medications), and discharge

• Decriminalizing suicide and prohibiting electroconvulsive therapy unless anesthesia is used

These are achieved through the mobilization of resources to required areas, and by providing social support.

Solution to Question 29:

The appropriate schedule for the treatment of multi-drug resistant tuberculosis (MDR-TB) is 7 drugs for 4-6 months followed by 4 drugs for 5 months.

Note: Pearl 1720 has been updated according to NTEP 2021 guidelines.

Solution to Question 30:

The POCSO (Protection of Children from Sexual Offences) Act is related to child sexual abuse.

POCSO Act defines a child as any person below eighteen years of age. It defines different forms of sexual abuse, including penetrative and non-penetrative assault, as well as sexual harassment and pornography.

The Act deems a sexual assault to be "aggravated" under certain circumstances, such as when the abused child is mentally ill or when the abuse is committed by a person in a position of trust or authority vis-à-vis the child, like a family member, police officer, teacher, or doctor.

People who traffic children for sexual purposes are also punishable under the provisions relating to abetment in the Act. The Act prescribes stringent punishment graded as per the gravity of the offence, with a maximum term of rigorous imprisonment for life and a fine.

Solution to Question 31:

Droplet precaution and isolation is not done for localized varicella zoster, as it is only transmitted through direct contact.

Varicella-zoster localized in the patient with an intact immune system with a lesion that can be contained/covered do not require droplet precaution. Disseminated varicella in any patient requires airborne and contact precautions.

Diphtheria, pertussis, and mycoplasma are transmitted by droplets and hence require droplet precaution and isolation.

Although patients are not contagious starting 48 hours after antibiotic therapy, droplet precautions are typically maintained until the completion of the antibiotic course and are culture-negative on two cultures taken 24 hours apart. Droplet precautions include the use of masks, gowns, gloves and protective eye gear when providing direct patient care.

Solution to Question 32:

This clinical scenario is suggestive of visceral leishmaniasis. This patient should be managed with liposomal amphotericin B.

This condition causes fever with rigor and chills associated with splenomegaly and hepatomegaly. Though lymphadenopathy is common in most endemic regions, it is rare in India.

It is diagnosed by ELISA, indirect immunofluorescent antibody test (IFAT), and rk39 which detects antibodies to Leishmania donovani.

The treatment is as follows:

• In most endemic regions:

• Drug of choice - pentavalent antimonial (sodium stibogluconate or meglumine antimoniate)

• If it fails or is unavailable - amphotericin B

• In India and Mediterranian countries, the drug of choice is liposomal AmB.

• Other drugs - paromomycin and miltefosine

Option B: Fluconazole is an antifungal and is not useful in visceral leishmaniasis

Option C: Linezolid is used in the treatment of MRSA and related infections.

Option D: Bedaquiline is a newer anti-TB drug used in MDR-TB.

Community Medicine AIIMS 2019 (May & Nov)

Question 1:

Scope of modern concept of family planning services include all of the following except:

a) Screening for cervical cancer

b) Providing services for unmarried mothers

c) Screening for HIV infection

d) Providing adoption services

Question 2:

Sandfly does not transmit:

a) Kala azar

b) Oriental sore

c) Oraya fever

d) Trench fever

Question 3:

The seasonal variations of a disease were studied in children, in a group from June to July and in another group of similar characteristics from August to September. The test of significance that should be used to compare the data is:

a) Chi square

b) Paired t test

c) Wilcoxan test

d) ANOVA

Question 4:

The bone marrow density along with the gestational age is plotted from two different studies. Which of the following is true for the relation between bone marrow density and gestational age?



a) Strength of correlation is same in A and B

b) Strength of the correlation is more in A than in B

c) Strength of the variation is more in B than A

d) Variation in variables is more with A than B

Question 5:

A study was conducted to assess the effect of an intervention for certain disease. The p values for two similar studies A and B with 95% confidence limits, was found to be 0.02 and 0.001 respectively using interventions "X" and "Y" with a control group. Based on preliminary analysis, the result may be interpreted as:

a) If A, B and C are correct

b) If A and C are correct

c) If B and D are correct

d) If all (A, B, C and D) are correct

Question 6:

In a study, group A received placebo while group B received a new drug. In group A, 36 deaths were reported out of a sample of 120. In group B, 26 deaths were reported out of a sample size of 130. How many patients should be treated to avert 1 death?

a) 100

b) 10

c) 250

d) 160

Question 7:

Regarding ASHA, all are true except:

a) Malaria slide preparation

b) Ensure immunization as per schedule

c) Accompany pregnant women to hospital

d) Spread awareness about contraception

Question 8:

A study was conducted on two different samples for comparing variation between DBP and Vitamin D. The mean diastolic blood pressure is 110 with standard deviation 11 and the mean vitamin D level is 18 with standard deviation 3. Which of the following statement is true regarding the coefficient of variation (CV)?

a) CV of DBP is more than Vitamin D

b) CV of Vitamin D is more than DBP

c) Both DBP and Vitamin D have equal variability

d) It cannot be ascertained from information provided

Question 9:

Which is the study design of choice for finding out circadian variation of fat content in expressed breast milk of mother of preterm babies?

a) Prospective Cohort

b) Ambispective Cohort

c) Case control

d) Cross sectional

Question 10:

Mass vaccination is most effective in which of■ the following outbreaks?

a) Cholera

b) Typhoid

c) Polio

d) Measles

Question 11:

Program for improvement of quality of labor room comes under:

a) LaQshya

b) Ayushman Bharat

c) Janani Shishu Suraksha Karyakram

d) Pradhan Mantri Surakshit Matritva Abhiyan

Question 12:

Look at the image below and select the linear regression equation corresponding to the graph.



©MARROW

a) y=10-x

b) x=10+y

c) y=11+2x

d) x=11+2y

Question 13:

All of the following can be done to remove confounders except:

a) Randomization

b) Blinding

c) Restriction

d) Matching

Question 14:

A study was done to assess the risk factors for breast cancer in 100 females from urban cities. A statistical group and analysis are shown below. The values indicate the odds ratio and the 95% confidence interval. How many risk factors may be ascertained from the statistical analysis of the research?



a) 3

b) 4

c) 5

d) 6

Question 15:

In a town of 20,000 populations, total 456 births were there in a year out of which 56 were dead born. The total death were 247 out of which 56 deaths were within 28 days of life and

another 34 had dies after 28 days and before completing the first birthday. Calculate the infant mortality rate of the area?

a) 197

b) 392

c) 225

d) 344

Question 16:

A nurse sustained a needle stick injury while sampling the blood of a patient who is HIV-positive. Which of the following is incorrect?

a) Washing hands with soap and water

b) Administration of tenofovir monotherapy

c) Measure baseline viral markers of health care worker immediately

d) Measure viral markers after 6 weeks

Question 17:

The renal feed of a patient suffering from end-stage renal disease (stage 5) should ideally be:

a) Low volume, high calories

b) High volume, low calories

c) High volume, high calories

d) Low volume, low calories

Question 18:

Which of the following is best method to survey the injection practices of health workers?

a) Direct observation with a structured checklist

b) Closed circuit camera surveillance

c) In-depth interview with the patient

d) In-depth interview with the healthcare provider

Question 19:

A polysaccharide vaccine against Streptococcus pneumoniae has 23 serotypes. In which of the following patients is this vaccine most beneficial?

a) A child < 2 years

b) A patient with sickle cell disease

c) A patient with cystic fibrosis

d) A patient with recurrent otitis media

Question 20:

A 5-year-old boy had received a full course rabies vaccine taken in December 2018. His pet dog which has been fully vaccinated now (November 2019) bit him on the toe. Bleeding was present at the site of the bite. What would be the line of treatment?

a) No treatment

b) 2 doses of vaccine on Day 0 & 3

c) 5 doses of vaccine with Immunoglobulin

d) 5 doses of vaccine

Question 21:

Mass drug administration is not done for which of the following conditions?

a) Worm infestation

b) Vitamin A deficiency

c) Scabies

d) Filariasis

Question 22:

Primary level of prevention for specific protection includes all of the following except?

a) Meningitis prophylaxis in exposed contacts/relatives

b) Hand washing

c) Vaccination

d) Adding iron to wheat flour

Question 23:

The treatment of paucibacillary leprosy consists of MDT (which comprises rifampicin, dapsone, and clofazimine). The latest recommended regimen is:

a) 2 drug therapy for 6 months

b) 3 drug therapy for 12 months

c) 2 drug therapy for 12 months

d) 3 drug therapy for 6 months

Question 24:

The pathogen transmitted by the vector is ?

a) Kyasanur forest disease virus

b) Orientia tsutsugamushi

c) Chandipura virus

d) Anaplasma phagocytophilia

Question 25:

The numerator of perinatal mortality rate includes _____

a) Post-neonatal period death

b) Early neonatal death of fetus > 1000 g

c) Stillbirth of fetus < 500 g

d) Abortion after 16 weeks

Question 26:

What is the cornerstone of treatment for a patient of dengue with warning signs?

a) IV Fluids

b) Platelets

c) Cryoglobulins

d) FFP

Question 27:

When can a patient with dengue hemorrhagic shock be discharged?

a) No episode of fever for more than 24 hours with paracetamol

b) After the return of appetite

c) Urine output more than 200ml

d) After 24hrs from recovery of shock

Answer Key

Question No.	Correct Option
1	c
2	d
3	a
4	b
5	b
6	b
7	a
8	b
9	a
10	c
11	a

12	c
13	b
14	a
15	c
16	b
17	a
18	a
19	b
20	b
21	b
22	b
23	d
24	b
25	b
26	a
27	b

Detailed Explanations

Solution to Question 1:

Screening for HIV infection does not come under the purview of family planning.

The modern concepts of family planning services include:

• Proper spacing and limitation of birth

• Advice on sterility

• Education for parenthood

• Sex education

• Screening for pathological conditions related to the reproductive system (e.g. cervical cancer)

• Genetic counseling

• Premarital consultation and examination

• Pregnancy tests

• Marriage counseling

• Preparation of couples for the first child

• Providing services to unmarried mothers

• Teaching home economics and nutrition

• Providing adoption services

Solution to Question 2:

Trench fever is not transmitted by sandflies. It is transmitted by lice and caused by Bartonella quintana.

Solution to Question 3:

Chi-square test is used to test the significance of the difference between two proportions.

• Analysis of seasonal or cyclic variation of a disease provides the data, which is expressed as proportion of children affected by the disease in various months. Though the data seems to be expressed in numbers, it is actually a qualitative or nominal variable as we assess the proportion of children who are affected by the disease and those who are not affected by the disease.

• The chi-square test can be used to compare two or more groups.

Solution to Question 4:

The strength of the correlation is more in graph A than in graph B.

Graphs A and B show a positive correlation between bone marrow density and gestational age.

In graph A: The data points in the scattergram are closer to the line of best fit, indicating a strong correlation.

In graph B: The data points are scattered farther away from the line of best fit, indicating a weak correlation.

Therefore, the strength of the correlation is more in graph A than B.

Solution to Question 5:

Statements A and C are correct.

The p-value of study A with intervention X at 95% confidence interval is 0.02.

The p-value of study B with intervention Y at 95% confidence interval is 0.001

The probability (p) values of both the studies are less than 0.05 (level of significance - α(alpha)). Therefore, both studies A, and B are significant, thus, rejecting the null hypothesis.

However, study B has a lower p-value compared to study A, which suggests that the study B with intervention Y is more significant and precise than study A with intervention X.

Solution to Question 6:

10 patients need to be treated with the new drug to avert 1 death.

The number needed to treat (NNT) is defined as the number of individuals to be treated with the new drug to prevent 1 death (or to save 1 life).

NNT = 100/ARR

Where, ARR is the absolute risk reduction, which is the difference in the absolute risk rates between the placebo group and drug group. The absolute risk is usually expressed in percentage.

ARR = (No. of deaths/Sample population) x 100

Calculation:

Step 1: Calculation of absolute risk rate

The absolute risk rate in group A (placebo)

= (36/120)x100

= 30%

The absolute risk rate in group B (new drug)

= (26/130)x100

= 20%

Step 2: Calculation of absolute risk reduction (ARR)

ARR = Absolute risk rate in group A - Absolute risk rate in group B

= 30 - 20

= 10%

Step 3: Calculation of number needed to treat

NNT = 100/ARR

= 100/10

= 10

Therefore, 10 individuals need to be treated with the new drug to save 1 life.

Solution to Question 7:

Preparation of malaria slides is the responsibility of health worker male (HWM), not ASHA.

Roles and responsibilities of ASHA:

A. Health education: The ASHA provides information and awareness to the community regarding nutrition, basic sanitation and hygienic practices, healthy living and working conditions, and timely utilization of health and family welfare services.

B. Maternal and child health:

• Facilitate utilization of health services such as immunization, antenatal and postnatal checkup and supplementary nutrition, sanitation and other services provided by the government.

• Provide counseling to women on birth preparedness, safe delivery, breastfeeding and complementary feeding, immunization, contraception, and prevention of common infections.

• Escort/accompany pregnant women and children requiring treatment or admission to the nearest health center.

• Upon graded training, provide newborn care and management of common childhood illnesses.

C. Primary care:

• Provide primary care for minor ailments like diarrhea and fever and provide first aid for injuries.

• Provide DOTS under RNTCP.

• Depot holder of essential provisions such as oral rehydration therapy, iron and folic acid tablets, chloroquine, disposable delivery kits, oral pills, condoms, etc.

D. Sanitation:

• Work with village health and sanitation committee to create a comprehensive village health plan.

• Promote the construction of household toilets under total sanitation campaign

E. Notification: ASHA will inform births, deaths, and any disease outbreaks to the sub-center or PHC.

Solution to Question 8:

The coefficient of variation of vitamin D is more than that of diastolic blood pressure.

The coefficient of variation (CV) is given by the formula:

$CV = 100(s/\bar{x})$

where,

CV = Coefficient of variation

s = standard deviation

$\bar{x}$ = mean

Substituting the data given in the question, in the above formula,

The CV of diastolic blood pressure is:

$CV(DBP) = 100(11/110)$

$= 100(1/10)$

$= 10\%$

The CV of vitamin D is

$CV(\text{Vitamin D}) = 100(3/18)$

$= 100(1/6)$

= 16.67%

Therefore, the CV of vitamin D is more than the CV of DBP.

Solution to Question 9:

The study of choice is a prospective study.

In the above scenario, the study group would consist of the mothers of preterm babies, while the comparison group would consist of the mothers of term babies. To study the circadian variation of the fat content in the expressed breast milk, the samples of breast milk would be taken at specific time intervals every day and followed up for several days and would be analyzed. This is a prospective study.

Other options:

Option b: An ambispective cohort is the combination of both retrospective and prospective cohort studies.

Option c: In a case-control study, the study begins from the outcome and goes retrospectively to find out the exposure among the cases and controls.

Option d: Cross-sectional study is done to identify the prevalence of a disease.

Solution to Question 10:

Mass immunization is effective in case of an outbreak of poliomyelitis.

The WHO has laid down standard operating procedures to be followed for responding to a poliovirus event or outbreak.

In the case of detection of an event or outbreak, the WHO recommends the following step-wise approach:

• Notification of the disease to international health regulations (IHR) focal point at the WHO regional centre.

• Detailed investigation and risk assessment.

• Enhanced surveillance to detect ongoing person-person transmission.

• Vaccination response to reach every child.

The vaccination response includes:

• A rapid response (RR) vaccination campaign: This campaign is initiated with 14 days of the laboratory sequencing result and targets the immediate area of the virus isolation to rapidly stop further transmission. The target population is about 2 lakh to 5 lakh children.

• Supplementary immunization activities (SIA): Two high-quality large-scale SIA campaigns (SIA1and SIA2) are carried out, to ensure that >90% of children are vaccinated. These campaigns should be completed within 8 weeks of laboratory sequencing result. The target population in the large scale rounds is about 2 million children.

• Mop-up round or additional SIAs: If the monitoring suggests that children have been missed in certain areas, then a mop-up round of immunization is done.

The target age group for immunization under SIAs is children less than five years of age. If there is evidence of virus circulation among older age groups, then the target age group can be expanded to 10 or 15 years or even to the whole population depending on the local context.

Other options:

Options a: In case of a cholera outbreak, the mainstay of response includes health education on the role of hygiene, rehydration therapy, and sanitation measures.

Option b: The mainstay of typhoid control in case of an outbreak includes health education, water quality and sanitation improvement, and training of health professionals in diagnosis and treatment. Although WHO recommends immunization as a complementary method to the above measures, there is insufficient data to ascertain the effectiveness of vaccination in case of a typhoid outbreak.

Option d: In case of measles outbreak, the control measures include isolation for seven days after the onset of rash and immunization of the contacts within 2 days of exposure.

Solution to Question 11:

The Laqshya initiative launched recently by the government of India refers to initiative for labor room quality improvement.

There is enough evidence to show that improving the quality of care in labor rooms, especially on the day of birth, is central to maternal and neonatal survival.

• Laqshya is expected to improve the quality of care that is being provided to the pregnant mother in the labor room and maternity operation theatres, thereby preventing the undesirable adverse outcomes associated with childbirth.

• This initiative will be implemented in government medical colleges (MCs) besides district hospitals (DHs), and high delivery load sub-district hospitals (SDHs) and community health centers (CHCs).

• The initiative plans to conduct quality certification of labor rooms and also incentivize facilities achieving the targets outlined.

• The goal of this initiative is to reduce preventable maternal and new-born mortality, morbidity, and stillbirths associated with the care around delivery in the labor room and maternity OT and ensure respectful maternity care.

Solution to Question 12:

Among the given options, the equation $y=11+2x$ corresponds to the linear regression equation of the form $y=\alpha + \beta x$.

When 2 variables are correlated, such that the dependent variable is represented on the y-axis and the independent variable on the x-axis, then the value of y can be obtained by using the linear

regression equation $y = \alpha + \beta x$, where:

y - dependent variable

α - intercept

β - slope of the line

x - independent variable

In the equation $y=11+2x$, the intercept(α) = 11, while the slope of the line (β) = 2.

Option A: $y=10-x$, represents an intercept of 10, and a slope of -1 indicating a negative correlation. However, the given image shows a positive correlation.

Options B and D: They do not fit the linear regression equation of the form $y = \alpha + \beta x$.

Solution to Question 13:

Blinding does not eliminate confounding factors. Blinding is done to eliminate the bias (such as subject bias, observer bias, or investigator bias).

A confounding factor is defined as the one associated with both exposure and disease and is distributed unequally in study and control groups.

• For example, a prospective study is being conducted to know the effect of alcohol on the development of lung cancer.

• However, individuals who drink alcohol are most likely to smoke. So, smoking is associated with the exposure. Also, smoking is an independent causative factor for lung cancer. Thus, smoking is a confounding factor.

• In the above example, if the study is conducted without eliminating the confounding factor, then the result may falsely suggest that alcohol is a causative factor for lung cancer.

Techniques to eliminate confounding factors include

• Randomization

• Matching

• Stratified randomization

• Restriction

Randomization is the random allocation of the participants
in experimental studies (Ex.randomized control trial) into the study group and control group.

Matching, which is usually done in analytical studies (for example, case–control and cohort studies), is the process where the participants in two groups are matched based on factors such as age, gender, race, smoking status, etc. to eliminate the potential confounders.

Stratified randomization, which is also known as prestratification or blocking, is the process that combines both randomization and matching techniques. In this method, the participants are first grouped into subgroups which internally match with potential confounding factors such as age, race, gender, etc. An equal number of participants form each subgroup are then allocated to the study and control groups.

Restriction: This method restricts the study to only those participants with particular characteristics. For example, a study can be restricted to participants of only one gender, race, age group, geographical area, disease severity, etc., thus eliminating the potential confounders.

Solution to Question 14:

Among the factors shown in the above statistical group, the odds ratio of early menarche, BRCA mutation, and obesity show positive association and are statistically significant. Hence these 3 factors can be ascertained as risk factors.

High socioeconomic status is not a risk factor. Even though there is a positive association, it is not statistically significant. Any association that has a 95% confidence interval range that includes the value of 1.0 is considered to be non-significant. The value of 1.0 in the range means that the two factors may co-exist together but one is not causing the other to happen.

Odds ratio (OR) is the key parameter in case-control studies to assess the strength of association between the exposure and outcome. The interpretation of the odds ratio can be done as below:

• OR > 1 indicates a positive association. If both upper and lower limits of the 95% confidence limits are also >1, it suggests that the odds ratio is significant and the factor can be ascertained as a risk factor.

• OR = 1indicates no association. Irrespective of the odds ratio, if the lower limit of 95% CI is <1 and the upper limit of 95% CI >1, the OR is non-significant and cannot be ascertained either as a protective or a risk factor.

• OR < 1 indicates a negative association. If both upper and lower limits of the 95% confidence limits are also <1, it suggests that the odds ratio is significant and the factor can be ascertained as a protective factor.

Interpretation of the individual factors given in the question:

Factor	Odds ratio(95%confidence limits)	Interpretation
Early menarche	1.3(1.2-1.9)	Positive association, significant,risk factor
Age <50 years	0.04 (0.01-0.32)	Negative association, significant, protective factor
Smoking	0.5 (0.02-0.9)	Negative association, significant, protective factor
BRCA mutation	1.9 (1.1-2.8)	Positive association, significant,risk factor
Alcohol	1.0 (0.8-1.8)	No association, non-significant
High socioeconomic status	1.4 (0.5-3.5)	Positive association, non-significant
Obesity	1.24 (1.1-1.7)	Positive association, significant,risk factor

Factor	Odds ratio(95%confidence limits)	Interpretation
No family history of Ca. breast	0.3 (0.01-0.9)	Negative association, significant, protective factor

Solution to Question 15:

The infant mortality rate is 225 per 1000 live births.

Infant mortality rate (IMR) is defined as the ratio of infant deaths registered in a given year to the total number of live births in the same year. IMR is usually expressed as a rate per 1000 live births. Infants include those under the age of 1 year.

Calculation:

• IMR = (Number of deaths of children under 1 year of age in a year x 1000)/(Number of live births in the same year).

• In the given question, Out of total 456 births, 56 were dead born. Therefore, the number of live births= 456-56 = 400.

• In the same year, it is given that the number of deaths before 28 days is 56, and the number of deaths after 28 days but before 1 year. Therefore, the total number of deaths of children under the age of 1 year is 56 + 34 = 90

• IMR = (90x1000)/400 = 900/4 = 225

• Hence, the IMR is 225 per 1000 live births.

Solution to Question 16:

Triple therapy (tenofovir 300mg + lamivudine 300mg + dolutegravir 50mg) is recommended for HIV post-exposure prophylaxis (PEP), not monotherapy.

Steps to be undertaken after a needlestick injury are:

• Wash the wound and skin with water and soap. Do not scrub and avoid antiseptics.

• Assess exposed individual:

• Check baseline viral markers

• Monitor for acute seroconversion illness within 3-6 weeks after exposure

• If testing is positive, refer to ART center

• Assess exposure source and determine the risk of transmission

• Counseling and support

• Initiate PEP at the earliest - preferably < 2 hours and ideally < 72 hours

• 28-day prescription with drug information

• HIV test at 3 months after exposure

Solution to Question 17:

The renal feed of a patient suffering from end-stage renal disease should ideally be low volume, high calories.

Dietary factors may have an effect on the progression of kidney disease and its complications. A low-volume feed is preferred in order to avoid:

• Sodium and volume overload

• Hyperkalemia

• Hyperphosphatemia

• Accumulation of toxic metabolites of protein degradation

A high-calorie feed is preferred in order to combat the patient's risk of malnutrition and wasting seen in ESRD.

Solution to Question 18:

Direct observation with a structured checklist is the best method to survey the injection practices of health workers.

Checklist of practices for assessment or evaluation include:-

• In order not to harm the patient, each procedure should be administered with a new sterile single-use device, using the right medication, vaccine or fluid for infusion.

• In order not to expose the provider to any avoidable risk, in general, any needles used during a procedure should be placed in a puncture-proof closed container immediately after use without recapping. In addition, providers of these procedures should be fully vaccinated against Hepatitis-B.

• In order that any waste produced during the performance of a procedure does not become a hazard for other people, used sharps waste and infectious non-sharps waste should be safely managed.

Solution to Question 19:

The polysaccharide vaccine against Streptococcus pneumoniae (PPSV-23) is most beneficial for patients with sickle cell disease.

S. pneumoniae is an encapsulated organism that is detected by the spleen and is removed. However, in sickle cell disease, the sickled cells block the vessels in the spleen which results in splenic infarction. This hyposplenism results in an increased risk of pneumococcal infection. Hence, pneumococcal vaccine (PPSV-23) is given to these patients as a prophylactic measure to

reduce this risk.

Solution to Question 20:

This boy should receive 2 doses of vaccine on days 0 & 3 as he has received a prior course of post-exposure prophylaxis (PEP) > 3 months ago.

Management of a new dog bite in a previously immunized individual with either pre-exposure prophylaxis (PrEP) or post-exposure prophylaxis (PEP) is as follows:

• Category I: Wound treatment only

• Category II/ III:

• Wound treatment with soap and running water for 15 min and apply antiseptic

• Receive Anti-rabies vaccine (ARV) 2 doses ID (0.1 ml, 1 site)/ 2 doses IM (1 VIAL, 1 site) on days 0 and 3.

*Note: In developing countries like India, the vaccination status of animals alone, doesn't play a significant role in determining the initiation of prophylaxis in humans.

Solution to Question 21:

Mass drug administration is not done in case of vitamin A deficiency.

Mass Drug Administration (MDA) is the administration of a drug to an entire concerned population irrespective of age and gender. For example, antimalarials are given as MDA in malaria-endemic island communities and in low-endemic non-island settings approaching elimination, where there is minimal risk of re-introduction of infection, good access to treatment, and implementation of vector control and surveillance.

Vitamin A prophylaxis is given under National Immunization Schedule (NIS) only to children up to 5 years. It is given as mass prophylaxis and not as a drug for treatment. Vitamin A as a treatment is given only to a child with features of vitamin A deficiency and such doses are not instituted to everyone in the community. Therefore, it is not considered under MDA.

On the other hand, MDA is done in scabies (ivermectin), worm infestation (albendazole), and filariasis (DEC + albendazole).

Solution to Question 22:

Handwashing is not a specific protection measure under primary prevention. All other options are protection measures against specific diseases.

Solution to Question 23:

The latest recommendation is 3 drug therapy for 6 months to treat paucibacillary (PB) leprosy by WHO. The regimen consists of rifampicin, dapsone, and clofazimine.

Meanwhile, multibacillary (MB) leprosy is treated using the same 3 drugs for a duration of 12 months. While the NLEP guidelines earlier mentioned a 2-drug MDT for PB leprosy, the National Strategic Plan and Roadmap for Leprosy (2023-2027) now recommends a uniform MDT of 3 drugs for both PB and MB leprosy.

Note: The central government has approved a new treatment regimen for paucibacillary (PB) leprosy. The Ministry of Health and Family Welfare has introduced a 3-drug regimen for PB leprosy instead of a 2-drug regimen. This revised drug regimen will be applicable in India from April 1, 2025.

Solution to Question 24:

The given image is of a Trombiculid mite and it transmits Orientia tsutsugamushi.

Orientia tsutsugamushi is an obligate, intracellular bacterium. It causes scrub typhus disease in humans.

Hard ticks are covered by a chitinous shield on their dorsal surface which is called scutum. They have a head or capitulum present at the anterior end, which is visible from above. Hard ticks transmit certain diseases like tick typhus, viral encephalitis, hemorrhagic fever, etc.

Solution to Question 25:

Early neonatal death of fetus > 1000 g is included in the numerator of perinatal mortality rate .

Perinatal mortality rate is described as late foetal and early neonatal deaths weighing over 1000 grams at birth expressed as a ratio per 1000 live births weighing over 1000 gram at birth.

$$\text{Perinatal mortality rate} = \frac{\text{Late foetal and early neonatal deaths weighing over 1000g at birth}}{\text{Total live births weighing over 1000g at birth}} \times 1000$$

Solution to Question 26:

The primary modality of treatment in patients with dengue with warning signs is IV fluids.

The main pathology in dengue is plasma leakage resulting in intravascular volume depletion and thereby shock. Warning signs in dengue include:

• Abdominal pain or tenderness

• Persistent vomiting

• Clinical fluid accumulation (ascites, pleural effusion)

• Mucosal bleeding

• Lethargy or restlessness

• Hepatomegaly >2 cm

• Increase in hematocrit concurrent with a rapid decrease in platelet count

Inpatient management is advised to all patients of dengue with warning signs. The management of dengue includes:

• IV crystalloids - mainstay of treatment

• Platelet transfusion - severe thrombocytopenia (<10,000/mm3) and active bleeding

FFP and cryoglobulins are not the preferred for treatment in dengue.

Solution to Question 27:

A patient with dengue hemorrhagic shock can be discharged after the return of appetite.

Criteria for the discharge of patients recovering from dengue:

• Absence of fever for at least 24 hours without the use of antipyretic drugs

• Platelet count >50,000/cu.mm

• Return of appetite

• Adequate urine output

• Visible clinical improvement

• Minimum of 2-3 days after recovery from shock

• No respiratory distress from pleural effusion or ascites

Community Medicine AIIMS 2020 (May and Nov INI-CET)

Question 1:

A resident doctor gets pricked with a needle while attending an HIV positive patient. The regimen to be given is?

a) Zidovudine + Lamivudine

b) Zidovudine + Lamivudine + Nevirapine

c) Zidovudine + Lamivudine + Indinavir

d) Zidovudine + Nevirapine

Question 2:

Airborne transmission precaution is necessary for all of the following except:

a) Tuberculosis

b) Herpes Zoster

c) Measles

d) Chicken Pox

Question 3:

The maternal mortality ratio is defined as?

a) Maternal Deaths per 10,000 live births

b) Maternal Deaths per 100,000 live births

c) Maternal Deaths per 10,000 women in reproductive age group

d) Maternal Deaths per 100,000 women in reproductive age group

Question 4:

What is the dosage of iron and folic acid for adult women under the Anemia Mukth Bharath initiative?

a) 60 mg elemental iron + 500 mcg folic acid daily

b) 60 mg elemental iron + 500 mcg folic acid weekly

c) 100 mg elemental iron + 500 mcg folic acid weekly

d) 100 mg elemental iron + 500 mcg folic acid daily

Question 5:

Which of the following is true regarding nikshay poshan yojana?

a) Rs 500 is given per week

b) Rs 500 is given per month

c) Mid day meals are given

d) Dietary advice is given

Question 6:

Diphtheria vaccine is a:

a) Polysaccharide vaccine

b) Toxoid

c) Killed vaccine

d) Live attenuated vaccine

Question 7:

Under RBSK screening which one of the following diseases is not covered?

a) Congenital cataract

b) Congenital glaucoma

c) ROP

d) Vitamin A deficiency

Question 8:

The serum level of tetanus antitoxin that gives clinical protection is?

a) >0.1 IU/ml

b) >0.01 IU/ml

c) >0.001 IU/ml

d) >0.0001 IU/ml

Question 9:

Which of the following is true about the reference woman used in the calculation of the recommended daily allowance (RDA)?

a) Weight 55 kgs

b) Height 1.73 m

c) Works 10 hours/day

d) Age of 18-25 years

Question 10:

Which of the following methods is not effective against preventing the spread of COVID 19?

a) Face mask

b) 1% glutaraldehyde

c) 70% alcohol

d) Hand washing

Question 11:

According to biomedical waste management, used pacemakers should be disposed of in which of the following colored bag?

a) Yellow

b) White

c) Black

d) Blue

Question 12:

In which of the following bags are latex gloves disposed of after use?

a) Yellow

b) Red

c) Blue

d) Transparent

Question 13:

Which of the following is a non-parametric test?

a) Student t test

b) Friedman test

c) ANOVA

d) Pearson

Question 14:

Glasgow coma scale is a type of?

a) Nominal scale data

b) Ordinal scale data

c) Alphanumerical data

d) Interval scale data

Question 15:

A new drug combination of chemotherapy and immunotherapy for metastatic melanoma was prescribed by a radiotherapist. It prolongs the survival. Which of the following is true in this situation?

a) Incidence increases and prevalence reduces

b) Incidence reduces and prevalence remains the same

c) Incidence remains the same and prevalence increases

d) Incidence reduces and prevalence increases

Question 16:

The HIV sentinel surveillance includes all of the following populations except

_____

a) Single male migrants

b) Truck drivers

c) STD clinic attendees

d) Antenatal woman

Question 17:

You are a medical officer at a PHC. Which of the following would you advise for mass prophylaxis/ treatment for anemia?

a) $FeSO_4$ + Vit C + Folate + Vit B12

b) Iron (III) hydroxide polymaltose

c) Ferrous fumarate

d) Inj. iron sucrose

Question 18:

Which of the following quantitative methods is the best to study previous research studies?

a) Cohort study

b) Systematic review

c) Meta-analysis

d) RCT

Question 19:

Discrete probability distribution that expresses probability of a given number of events in a particular time frame is known as:

a) Binomial distribution

b) Gaussian distribution

c) Poissons distribution

d) Normal distribution

Question 20:

The data which best explains the difference between the highest and the lowest values is:

a) Variance

b) Coefficient of variation

c) Range

d) Interquartile range

Question 21:

Which of the following is a marker of the heat stability of vaccines?

a) VMV

b) VVM

c) MVV

d) VMM

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	b
5	b
6	b
7	b
8	b
9	a
10	b
11	d
12	b
13	b
14	b
15	c
16	c
17	a
18	c

19	c
20	c
21	b

Detailed Explanations

Solution to Question 1:

Amongst the given options, a three-drug combination of Zidovudine+Lamivudine+Indinavir is preferred.

Three drug combination is preferred as per the US Public health service guidelines, WHO guidelines, and MOHFW 2020 guidelines.

A PEP regimen for HIV should have Lamivudine + Tenofovir + Lopinavir/ritonavir to be taken for 4 weeks after exposure. It should ideally be started within 2 hours and definitely within 72 hours. Here as the Lopinavir/ritonavir is not part of the options, one can choose, the indinavir-based combination as the best of the available options.

Nevirapine is not preferred for use in PEP. Previous NACO 2018 guidelines mentioned the use of 2 drug combinations but it is no longer preferred.

Exposures that may warrant occupational PEP include:

• Parenteral or mucous membrane exposure (splashes to the eye, nose, or oral cavity)

• The following bodily fluids may pose a risk of HIV infection: blood, blood-stained saliva, breast-milk, genital secretions, and cerebrospinal, amniotic, rectal, peritoneal, synovial, pericardial, or pleural fluids.

Exposures that do not require PEP include:

• When the exposed individual is already HIV-positive

• When the source is established to be HIV-negative

• Exposure to bodily fluids that do not pose a significant risk: tears, non-blood-stained saliva, urine, and sweat.

To ensure no transmission has occurred, follow-up with HIV antibody testing should be done for at least 6 months post-exposure (e.g. at baseline, 6 weeks, 3 months, and 6 months).

Note: This is an AIIMS 2020 recall question. It may not reflect the latest guidelines. Kindly refer to the attached pearl for the same.

Solution to Question 2:

Measles, Tuberculosis, and chickenpox require airborne transmission precaution as does the disseminated herpes zoster.

The patient should be made to wear a mask for source control. A fit-tested NIOSH-approved N95 or higher level respirator should be used by the healthcare personnel whilst care delivery.

CDC recommends the display of signs similar to the one below in room doors of patients with the above conditions:

Solution to Question 3:

Maternal mortality ratio is defined as the number of maternal deaths per 100,000 live births during a given time period.

Maternal mortality rate is the number of maternal deaths per 100,000 women of reproductive age in a given time period.

Solution to Question 4:

According to the Anemia Mukt Bharat initiative, adult women need to take 60 mg elemental iron + 500 mcg folic acid tablets once a week. These tablets are sugar-coated, and red in color.

Solution to Question 5:

Under nikshay poshan yojana, a financial incentive of Rs.500 per month is provided for nutritional support to each notified TB patient for the duration of anti-TB treatment.

For pediatric TB patients who do not have bank accounts, the money shall be deposited in parent's/guardian's accounts whose details are to be sought and entered in nikshay against the patient's records.

The processing of payment is on the 7th of every month.

Solution to Question 6:

Diptheria vaccine is a toxoid.

Toxoids: Certain organisms like diphtheria and tetanus bacilli produce exotoxins. The toxins produced by these organisms are detoxicated and used in the preparation of vaccines.
The antibodies produced to neutralize the toxic moiety produced during infection, rather than acting upon the organisms. In general, toxoid preparations are highly efficacious and safe immunizing agents.

Solution to Question 7:

All of the above conditions are included in child health screening & early intervention services under RBSK (Rashtriya Bal Swasthya Karyakram) except congenital glaucoma.

The RBSK aims at the universal screening of the 4 'D's i.e Defects at birth, Deficiencies, Diseases, Developmental delays including disabilities that would lead to early identification, timely management, and ultimately a reduction in morbidity, mortality, and lifelong disability in children.

Health conditions selected for the child (0-18 years) health screening and early intervention services are:

Defects:

• Neural tube defect

• Down's syndrome

• Cleft lip and palate

• Talipes

• Developmental dysplasia of the hip

• Congenital cataract, congenital deafness, congenital heart disease

• Retinopathy of prematurity

Deficiencies:

• Anemia especially severe anemia

• Vitamin A deficiency

• Vitamin D deficiency

• Severe acute malnutrition

• Goitre

Diseases:

• Scabies, fungal infections, and eczema

• Otitis media

• Rheumatic heart disease

• Reactive airway disease

• Dental caries

• Convulsive disorders

Developmental delays and disabilities:

• Vision, hearing, and neuro-motor impairment

• Motor, cognitive and language delay

• Autism

• Learning disorder

• Attention deficit hyperactivity disorder

Solution to Question 8:

The serum level of tetanus antitoxin that gives clinical protection is 0.01 IU/ml.

Tetanus is best prevented by tetanus toxoid. It provides active immunization by stimulating the production of protective antitoxin. Vaccination should be aimed at maintaining the serum protective level of antitoxin (approximately 0.01 IU/ml) throughout life and covering the entire community.

Solution to Question 9:

The reference woman used in the calculation of recommended daily allowance (RDA) weighs about 55 Kgs and is aged between 19-39 years, nonpregnant, non-lactating with a height of 1.61 meters, and a BMI of 21.2, is free from disease and physically fit for active work.

On each working day, she must be engaged in 8 hours of occupation/work, which usually involves moderate activity. When not at work she spends 8 hours in bed, 4-6 hours in sitting and moving about, 2 hours in walking, and in active recreation or household duties.

Solution to Question 10:

1% glutaraldehyde is not effective against preventing the spread of COVID 19.

Effective everyday practices that prevent the spread include actions like keeping a social distance (at least 6 feet) from others, washing your hands with soap and water for 20 seconds or using a hand sanitizer with at least 70% alcohol, and wearing a face mask when you have to go out in public.

Solution to Question 11:

According to biomedical waste management, used pacemakers should be disposed of in a blue colored bag. All the metallic body implants and glasswares are discarded in a blue colored bag.

Solution to Question 12:

Latex gloves are disposed of in red-colored bag after use.

Solution to Question 13:

Friedman test is a non-parametric test. It is used to test for differences between groups when the dependent variable being measured is ordinal.

Other options:

Student t-test, analysis of variation (ANOVA) test, and pearson correlation are all based on quantitative data and hence grouped under parametric tests.

Solution to Question 14:

Glasgow coma scale is a type of ordinal data as the GCS score can be graded as mild, moderate, and severe based on the order of severity.

Solution to Question 15:

By the new drug combination, the survival of the patient has increased, hence incidence remains the same while prevalence increases.

Incidence: Number of new cases in a defined population in a specified period of time.

Prevalence: Number of old and new cases at a given point of time over a period of time.

Options A, B, D: The treatment only prolongs the disease, it has nothing to do with the occurrence of cases thereby nothing to do with incidence.

Solution to Question 16:

Sentinel surveillance of HIV is usually not done for STD clinic attendees.

Sentinel surveillance is used to augment the notification systems used in the surveillance of diseases.

Till 2012-13, STD sites were also included in the HIV sentinel surveillance but from 2014-15, these sites were entirely discontinued, since the STD data had significant biases.

The study population for HIV sentinel surveillance includes:

• Female sex worker: Women who are engaged in consensual sex for money or payment in kind, as a means of livelihood in the last six months

• Men who had anal or oral sex with a male partner in the last month.

• Injection drug users: Men and women who use addictive substances or drugs for recreational or non-medical reasons, through injections, at least once in the last three months.

• Hijra/Transgender: Person whose identity does not conform unambiguously to conventional notions of male or female gender roles, but combines or moves between these.

• Single male migrants: Single male, living at a place other than the place of usual residence without his spouse or family, for the purposes of work and visiting his home town at least once a year.

• Long-distance truckers: Truckers who travel more than 800 km one way between source and destination.

• A pregnant woman of age 15-49 years and attending the antenatal clinic for the first time.

Solution to Question 17:

Among the given options, $FeSO_4$ + vitamin C + Folate + vitamin B12 may be considered for mass prophylaxis against anemia.

Vit C enhances the absorption of iron from the stomach by acting as a reducing substance.

Subclinical folate deficiency is seen in up to 30% of North Indian pregnant women. Moreover, it may be masked by widespread iron deficiency. It is, for this reason, iron and folate supplementation is carried out in national programmes.

Clinical vitamin B12 deficiency is not commonly encountered, but subclinical cases might be seen in up to 30% of adults and children due to widespread restriction to consumption of vegetarian products only. Hence supplementing vitamin B12 is a good idea provided the cost factor is not an issue.

Option B: Iron (III) hydroxide polymaltose was marketed as being superior to iron sulfate in terms of efficacy, but it lacked any data to back up the evidence.

Option C: Ferrous fumarate may be considered for the treatment of iron deficiency anemia with similar efficacy as that of ferrous sulfate.

Option D: Injection iron sucrose or any parenteral form is not used as a first-line option. It is only indicated if the oral form is not tolerated.

Solution to Question 18:

The quantitative method which is the best to study previous research studies of a selected topic/research question is meta-analysis.

A systematic review is a process of collecting, reviewing, and presenting information from earlier research to answer a question. It is a qualitative technique. It may or may not involve a meta-analysis.

Solution to Question 19:

Poissons distribution is the probability distribution that describes the chances of an event in a particular time frame.

For example, if you plot the number of patients visiting the OPD of your hospital for a week on a chart, you can conclude that on a particular day(say Monday) there are more number of patients. And if the pattern of patients is similar you can estimate the number of patients in OPD next week or so.

It is usually done for rare events and is depends on $\mu = \lambda t$ (where μ defines the number of events in happening in time t)

Other options

Gaussian distribution is another name for Normal distribution

Binomial distribution examples consist of those events which can only have two possible results given as yes or no / success or failure. It is usually done for finite number of events unlike Poissons distribution.

Solution to Question 20:

Range gives the difference between the highest and lowest values in a given set of data.

In a dataset of marks of students in a subject, if the highest score is 100 and the lowest is 51, the range would be 49.

Option A: Variance (var or SD^2) is the mean of the squares of all the deviation scores of the distribution. It is used to measure how far data is spread out.

Option B: Coefficient of variation (CV) is used to compare the variability. For example, whether the pulse varies more in old or young and growth varies more in boys or girls. It is calculated by expressing SD as a percentage of the mean, i.e., CV = SD/mean x 100. Since it is expressed as a percentage it doesn't have any units.

Option D: Quartiles divide the distribution into 4 equal groups. The difference between mid values of each quartile gives the interquartile range.

Solution to Question 21:

Vaccine Vial Monitors (VVM) are heat-sensitive indicators that are routinely used to monitor vaccine viability as the vaccine vial moves across the supply chain.

VVM records cumulative heat exposure through a gradual change in color. If the color of the inner square is the same or is darker than the outer circle, the vaccine has been exposed to too much temperature and should be discarded.

Community Medicine NEET 2018

Question 1:

The Chadah committee recommended all of the following except:

a) PHC at block level

b) One PHC for 50,000 population

c) One basic health worker per 10,000 population

d) Responsibility of PHCs in malaria eradication

Question 2:

Which of the following is the most peripheral centre under the Revised National Tuberculosis Control Programme organization structure?

a) District TB centre

b) Intermediate Reference Laboratory

c) Tuberculosis Unit

d) Designated Microscopy Centre

Question 3:

Which of the following is true about Factories Act?

a) Children under 14 years of age must not be employed in factories

b) A child of 16 years can be employed only between 6 AM and 8 PM

c) The maximum working hours for an adult is 72 hours per week

d) A child of 16 years can work till 5 hours per day

Question 4:

The dose of iodine for lactating women in India according to the recommended dietary allowance is:

a) 200 μg/day

b) 220 μg/day

c) 250 µg/day

d) 280 µg/day

Question 5:

Which of the following is true for non-parametric tests?

a) ANOVA is an example of a non parametric test

b) It is used for skewed distributions

c) It involves the assumption that the data has a normal distribution

d) It cannot be used for small sample sizes

Question 6:

Which article of the Indian Constitution confers Right to Life to the citizens of India?

a) Article 11

b) Article 21

c) Article 23

d) Article 25

Question 7:

According to the WHO, what percentage of immunization coverage is needed for eradication of measles?

a) 68%

b) 72%

c) 84%

d) 96%

Question 8:

In a well-designed cancer study, similar remission rates were obtained for a new treatment (33.2%) and the usual treatment (32.2%), with the p-value being 0.04. Which of the following statements is true regarding the study?

a) Both the treatments are equally effective

b) Neither of the treatments is effective

c) The new treatment is more effective than the usual treatment

d) The information given is not adequate to compare the efficacy of the treatments

Question 9:

A 9-month-old baby with diarrhea is feeding well and is thirsty. Skin pinch when done returns in 2 seconds. What is the treatment?

a) 200-400 ml of ORS to be given in the first four hours.

b) 400-700 ml of ORS to be given in the first four hours.

c) 700-900 ml of ORS to be given in the first 4 hours.

d) In addition to the usual fluid intake, 100-200 ml should be given after each loose stool.

Question 10:

What value of BMI is considered as 'lethal' in men?

a) < 12

b) < 13

c) < 14

d) < 15

Question 11:

Which of the following is not used in the assessment of body fat?

a) Quetelet's index

b) Total body potassium

c) Total body water

d) Breslow index

Question 12:

What is the ratio of the incidence of the disease in the exposed to that of the non-exposed?

a) Attributable risk

b) Odds ratio

c) Relative risk

d) Population attributable risk

Question 13:

In order to reach the demographic goal of NRR=1, the couple protection rate should exceed?

a) 50%

b) 40%

c) 70%

d) 60%

Question 14:

Which is the new online software launched under the national tuberculosis elimination program (NTEP) by the government to monitor TB patients?

a) NIKSHAY

b) NISCHAY

c) E-DOTS

d) NIKUSTH

Question 15:

Cytotoxic and expired drugs are disposed by:

a) Dumping

b) Autoclave

c) Incineration

d) Chemical disinfection

Question 16:

In a screening test for DM out of 1000 population, 90 were positive. However, a gold standard test for DM was done of which 100 were positive. Calculate the sensitivity of the screening test?

a) 90/100

b) 100/110

c) (90-10)/1000

d) 90/1000

Question 17:

The incidence of a disease is 4 per 1000 of the population with an average duration of 2 years. What is the prevalence?

a) 8/100

b) 2/1000

c) 4/1000

d) 8/1000

Question 18:

The probability that infection will occur among susceptible persons following contact with an infectious person within the incubation period is known as the_____.

a) Secondary attack rate

b) Case fatality rate

c) Primary attack rate

d) Tertiary attack rate

Question 19:

Which of the following does not cause hardness of water?

a) Calcium carbonate

b) Calcium sulphate

c) Calcium bicarbonate

d) Magnesium bicarbonate

Question 20:

Which of the following is not a method of direct transmission of communicable diseases?

a) Vertical transmission

b) Contact with soil

c) Droplet nuclei

d) Droplet transmission

Question 21:

What should be the minimum interval between the administration of two live vaccines?

a) 2 weeks

b) 4 weeks

c) 8 weeks

d) 12 weeks

Question 22:

Kala-Azar is endemic in all these areas except:

a) West Bengal

b) UP

c) Bihar

d) Assam

Question 23:

What is the unit of observation in ecological studies?

a) Population

b) Patient

c) Healthy inividuals

d) Case

Answer Key

Question No.	Correct Option
1	b
2	d
3	a
4	d
5	b
6	b
7	d
8	c
9	b
10	b
11	d
12	c
13	d
14	a
15	c
16	a
17	d
18	a
19	a
20	c
21	b
22	d
23	a

Detailed Explanations

Solution to Question 1:

The recommendation of one PHC for a 50,000 population was given by Kartar Singh Committee.

The recommendations given by the Chadah Committee of 1963:

• PHC at the block level

• One basic health worker per 10,000 population

• Responsibility of PHCs in malaria eradication

Solution to Question 2:

Among the options given, Designated Microscopy Centre is the most peripheral centre.



Solution to Question 3:

Factories act prohibits children of <14 years from being employed.

Children between 15–18 years can be employed only in certain industries and work only between 6 AM–7 PM. They should not be made to work for more than four and a half hours.

The maximum working hours for an adult industrial worker should be not more than 48 hours per week, not exceeding 9 hours a day; if overtime is paid, then it should be not more than 60 hrs per week.

Solution to Question 4:

According to the National Institute of Nutrition (NIN) 2020 guidelines, the recommended dietary allowance (RDA) for dietary iodine in lactating women in India is 280 µg/day.

NIN guidelines on recommended dietary allowance for iodine per day:

Note: According to the WHO guidelines, the RDA during both pregnancy and lactation is 250 mcg.

Age	Male	Female
Birth to 6 mont hs	100 mcg	100 mcg
7–12 months	130 mcg	130 mcg

Age	Male	Female
1–9 years	90-120 mcg	90-120 mcg
10–12 years	150 mcg	150 mcg
14–18 years	150 mcg	150 mcg
> 19 years	150 mcg	150 mcg
Pregnancy	—	250 mcg
Lactation	—	280 mcg

Solution to Question 5:

Non-parametric tests are tests that do not assume that the data is normally distributed. Thus, they are used for skewed (asymmetric) distributions {skewed either to the lesser side (left-sided skewed) or towards the greater side (right-sided skewed)}.

The chi-square test is an example of a non-parametric test that is used to test hypotheses in relation to nominal scale data or categorical data.

ANOVA is an example of a parametric test.

Solution to Question 6:

The Right to Life is given in Article 21 of the Indian Constitution.

Option A: Article 11 defines the right of the parliament to regulate the right of citizenship by law.

Option C: Article 23 defines the prohibition of traffic in human beings and forced labor.

Option D: Article 25 defines the freedom of conscience and free profession, practice, and propagation of religion.

Solution to Question 7:

According to the WHO, eradication of measles requires achieving an immunization coverage of at least 96% of children under 1 year of age.

Solution to Question 8:

As the p-value is 0.04, we can conclude that the new treatment is more effective than the usual treatment.

Even though both treatments have similar rates of remission, a valid statistical test is required to determine the difference in their effectiveness. Statistically, p-value < 0.05 implies that the alternate hypothesis is true.

In this scenario, we can consider:

• Null hypothesis: Both the treatments are equally effective (in other terms, there is no difference between the effect of the two treatments).

• Alternative hypothesis: New treatment is more effective than the usual treatment.

As the p-value is 0.04, which is statistically significant (<0.05), the null hypothesis is rejected and the alternative hypothesis is accepted. Hence the new treatment is more effective than the usual treatment.

Solution to Question 9:

According to the IMNCI classification, the baby has some dehydration and comes under Plan B. The amount of ORS given is 75ml/kg body weight.

Here, the weight of the child is not specified. Hence, 400-700 ml of ORS at home is to be given in the first four hours according to the age of the child. The below table shows the amount of ORS to be given according to the age of the child.

In addition to this, 50-100 ml extra fluid is given after each loose stool. After four hours, the baby is reassessed and the appropriate plan is selected again for the continuation of the treatment.

The four rules of home treatment are:

• Give extra fluid

• Give zinc supplements (1 tablet per day for 14 days, dissolved in a small amount of ORS or breast milk or clean water)

• Continue breastfeeding

• Counsel the mother to return if there is blood in stools and if the child is feeding poorly

Age	under 4 months	4-11 months	1-2 years	2-4 years	5-14 years	15 years and over
Weight(kg)	under 5	5-7.9	8-10.9	11-15.9	16-29.9	30 or over
ORS solution (ml)	200-400	400-600	600-800	800-1200	1200-2200	2200-4000

SIGNS	CLASSIFY AS	IDENTIFY TREATMENT (Urgent pre-referral treatments are in bold print.)
Two of the following signs: • Lethargic or unconscious • Sunken eyes • Not able to drink or drinking poorly • Skin pinch goes back very slowly	SEVERE DEHYDRATION	➤ If child has no other severe classification: — Give fluid for severe dehydration (Plan C). OR **If child also has another severe classification:** **— Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way.** **Advise the mother to continue breastfeeding** ➤ **If child is 2 years or older and there is cholera in your area, give antibiotic for cholera.**
Two of the following signs: • Restless, irritable • Sunken eyes • Drinks eagerly, thirsty • Skin pinch goes back slowly	SOME DEHYDRATION	➤ Give fluid and food for some dehydration (Plan B). ➤ **If child also has a severe classification:** **— Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way.** **Advise the mother to continue breastfeeding** ➤ Advise mother when to return immediately. ➤ Follow-up in 5 days if not improving.
Not enough signs to classify as some or severe dehydration.	NO DEHYDRATION	➤ Give fluid and food to treat diarrhoea at home (Plan A). ➤ Advise mother when to return immediately. ➤ Follow-up in 5 days if not improving.

Solution to Question 10:

A body mass index (BMI) of 13 or less is considered lethal in men.

However, females can survive even at a lower BMI up to 11.

Solution to Question 11:

Breslow's staging is used for the staging of malignant melanoma, not for the assessment of body fat.

Solution to Question 12:

The ratio of incidence of disease or death among exposed to that among the non-exposed is called relative risk (RR).

Solution to Question 13:

In order to reach the demographic goal of NRR=1, the couple protection rate (CPR) should exceed 60%. This coincides with every couple having 2 children.

Solution to Question 14:

Nikshay is the new online software launched under the national tuberculosis elimination program for monitoring TB patients.

Nikusth is the new online software for monitoring leprosy patients.

Nischay is a pregnancy test kit.

Solution to Question 15:

Cytotoxic and expired drugs are collected in a yellow bag and then disposed of by incineration.

Expired cytotoxic drugs are to be returned to the manufacturer and subjected to incineration, encapsulation, or plasma pyrolysis. Other discarded medicines can be disposed by incineration.

Solution to Question 16:

The screening test only identified 90 people as positive for diabetes, while the gold standard test identified 100 people. This means that out of the 100 confirmed, 90 have been identified as truly positive by the screening test and 10 are falsely negative. Therefore,

• True positive = 90

• False negative = 100-90= 10

• Sensitivity = TP/(TP+FN) =90/(90+10) = 90/100 = 90%

Solution to Question 17:

The prevelance in the given situation is 8/1000 population.

Prevalence is given by the formula :

Prevalance = Incidence x Mean Duration

Prevalence = 4/1000 x 2 = 8/1000 population

Solution to Question 18:

The probability that infection will occur among susceptible persons following contact with an infectious person within the incubation period is called secondary attack rate.

It is calculated by the following formula:

Secondary attack rate (SAR) = (Number of exposed or susceptible persons developing the disease within the incubation period x 100)/(Total number of close or exposed contacts that are susceptible)

Primary cases are usually excluded from the numerator and denominator.

Secondary attack rate measures the communicability of the disease.

Solution to Question 19:

Calcium carbonate does not cause hardness of water. Hardness of water is caused by bicarbonates and sulphates of calcium and magnesium.

There are two types of hardness of water:

• Temporary (carbonate) hardness is due to bicarbonates of calcium and magnesium.

• Permanent (non-carbonate) hardness is due to sulphates, chlorides and nitrates.

Solution to Question 20:

Airborne transmission by droplet nuclei is not a method of direct transmission of diseases.

It is a method of indirect transmission of diseases.

Droplet nuclei are tiny particles of size 1-5 microns range that may be formed by either due to evaporation of droplets coughed or sneezed into the air or are formed from aerosols.

These droplet nuclei remain airborne for long periods of time and can transmit diseases like tuberculosis, influenza, chickenpox, measles, Q fever and other respiratory infections.

Droplet transmission, vertical transmission, and contact with soil are methods of direct transmission of diseases.

Differences between droplet and droplet nuclei are summarised below:

DROPLET NUCLEI	DROPLET
Indirect transmission	Direct transmission
It is the residue from evapora ted droplets	It is produced while coughing, sneezing, talking
<5microns in diameter	≥5 microns in diameter
No need for close contact for transmission	Requires closeness and direct contact for transmission
Diseases transmitted: chicke npox, measles etc.	Diseases transmitted: diphth eria, meningococcal meningit is etc.

Solution to Question 21:

The minimum interval between the administration of two live vaccines is 4 weeks.

This is because decreasing the interval between two doses may interfere with the antibody response and protection. Thus, live vaccines can either be given on the same day according to the schedule or 4 weeks apart.

Note: In Park 27th ed. the minimum interval between two live vaccines is given as 3 weeks. However, several other standard sources including CDC and UNDP suggested an interval of 4 weeks. When both are given in options, it is recommended to mark 4 weeks as the correct answer.

Solution to Question 22:

Kala-Azar is not endemic in Assam. It is endemic in the following states in 54 districts:

- Uttar Pradesh
- West Bengal
- Bihar
- Jharkhand

Solution to Question 23:

In ecological studies, the unit of observation is the population or community.

Type of study	Unit
Ecological/ Correlational	Population
Cross-sectional/ Prevalence	Individual
Case-control/ Case reference	Individual
Cohort/ Follow up	Individual
Randomised Controlled Trials	Patients
Field Trial	Healthy people
Community trial	Community

Community Medicine NEET 2019

Question 1:

A 4-year-old male child developed sore throat and difficulty in swallowing and mild fever. On examination, the clinical picture shown below is seen. Which of the following sentences is false regarding the vaccine that could have prevented this condition?

a) Its administered on the anterolateral aspect of thigh

b) A prior adverse reaction with temperature above 37 degrees is a contraindication

c) Cellular pertussis component in the vaccine is not recommended after 7-years of age

d) The diphtheria toxoid dose in adults is much lesser than the pediatric dose

Question 2:

Which of the following parameters would you use to check the efficiency of the surveillance system for malaria under the National Vector Borne Disease Control Programme?

a) Annual Parasite Index

b) Annual Blood Examination Rate

c) Slide positivity rate

d) Slide falciparum rate

Question 3:

Which of the following is false regarding confounding factor?

a) It can be reduced by matching

b) It is associated individually with both cause and effect

c) It is distributed equally in both study and control groups

d) It is associated with the exposure of the study

Question 4:

Which of the following is the best level of prevention of breast cancer?

a) Specific protection

b) Early diagnosis and treatment

c) Disability limitation

d) Rehabilitation

Question 5:

What is the dosage of vitamin A given for a 2-year-old baby with keratomalacia?

a) 2,00,000 IU immediately, followed by same dose after 24 hours

b) 1,00,000 IU immediately, followed by the same dose after 1 week

c) 1,00,000 IU immediately, followed by the same dose after 24 hours and after a week

d) 2,00,000 IU immediately, followed by the same dose 24 hours later and after 2 weeks

Question 6:

In a normal curve, what is the percentage of distribution of one standard deviation?

a) 68%

b) 34%

c) 99%

d) 95%

Question 7:

According to IMNCI, a 6-month-old child is said to have pneumonia if the child has fast breathing with a respiratory rate more than:

a) 40

b) 60

c) 30

d) 50

Question 8:

Which of the following is part of the concurrent list of the Indian Constitution?

a) Adulteration of food

b) Fishing and fisheries beyond territorial waters

c) Regulating labor in mines

d) Public health and sanitation

Question 9:

The principal unit of administration in India is _____

a) Village

b) Centre

c) District

d) State

Question 10:

Cross product ratio can be arrived from:

a) Ecological study

b) Cohort study

c) Cross sectional study

d) Case control study

Question 11:

In controlled tipping, the amount of land required for the trench method to be practiced for a population of 10,000 is:

a) 4 acres

b) 5 acres

c) 2 acres

d) 1 acre

Question 12:

The test of choice to measure the difference in means between two groups in a study, with one group being a control is:

a) Chi-square test

b) Z test

c) Unpaired T test

d) Paired T test

Question 13:

The test of significance done for two or more proportions:

a) Chi- square test

b) Student's test

c) Z test

d) ANOVA test

Question 14:

If a patient tests positive for a disease, which of the following represents the probability of her actually having the disease?

a) Sensitivity

b) Specificity

c) Positive predictive value

d) Negative predictive value

Question 15:

In the following natural history of disease flowchart, point A to C is called as?



a) Screening time

b) Lead time

c) Lag time

d) Generation time

Question 16:

The following diagram charts the population change over years from 1983 to 1989 in a given place. What is the statistical diagram used here?



a) Histogram

b) Line diagram

c) Component bar graph

d) Multiple bar graphs

Question 17:

Triage is done during disaster management to ensure treatment of casualties is done properly. What is true about triage done at the disaster site?

a) First come first treated basis

b) Green labels are for those who need to be transported on priority

c) Moribund patients receive the lowest priority

d) Most commonly used system is three color code system

Question 18:

The risk of genetic diseases in consanguineous marriage between first cousins is?

a) 1-2%

b) 4-8%

c) 8-10%

d) 12-14%

Question 19:

A paradoxical carrier is defined as:

a) A person who sheds pathogens during incubation period

b) A carrier who acquired pathogen from another carrier

c) A person who acquired pathogen from patient

d) A patient who became a carrier

Question 20:

Which of the following vaccine is contraindicated in pregnancy?

a) Hepatitis A

b) Hepatitis B

c) Rabies

d) Varicella

Question 21:

Which of the following is not a personal protective equipment?

a) Goggles

b) Gloves

c) Face shield

d) Lab coat

Question 22:

What is the best way to plot the changes in incidence of a disease in a given area over time?

a) Line graph

b) Histogram

c) Ogive

d) Tree diagram

Question 23:

Which of the following is false about Rubella infection?

a) It is a type of droplet infection

b) Vertical transmission is possible

c) Infection in early pregnancy causes a milder form of disease in the fetus

d) Fetus affected in late pregnancy may have only deafness

Question 24:

A resident doctor sustained a needlestick injury while sampling blood of a patient who is HIV positive. A decision is taken to offer him post-exposure prophylaxis. Which one of the following would be the best recommendation?

a) Zidovudine + Lamivudine for 4 weeks

b) Lamivudine + Tenofovir + Efavirenz for 4 weeks

c) Lamivudine + Tenofovir + Dolutegravir for 4 weeks

d) Zidovudine + Lamivudine + Nevirapine for 4 weeks

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	b
5	d
6	a
7	d
8	a
9	c
10	d
11	d
12	c
13	a
14	c

15	b
16	c
17	c
18	b
19	b
20	d
21	d
22	a
23	c
24	c

Detailed Explanations

Solution to Question 1:

The image shows a dirty white pseudomembrane classic of diphtheria. A prior adverse reaction with a temperature above 37 is not a contraindication.

Precautionary measures are taken when the temperature is above 40 degrees, but it is not considered a contraindication for subsequent doses of DPwT.

Anaphylaxis is a contraindication for subsequent vaccination with diphtheria, pertussis, or tetanus vaccine. Encephalopathy following vaccination is a contraindication for pertussis vaccine. DT is given for successive doses.

Events such as inconsolable cry, fever more than 40.5°C, seizures with or without fever are considered as precautions, and not contraindications for DTwP.

The cellular component of pertussis is not recommended after 7 years.

• Unvaccinated child 5-7 years: Give DPT 1, 2 and 3 at 1-month intervals. Administer booster dose of DPT at a minimum of 6 months

after administering DPT-3 up to the age of 7 years.

• Unvaccinated child above 7 years: Administer Tdap (Tetanus and diphtheria toxoids and acellular pertussis) at 0, and Td (Tetanus, Diphtheria vaccine) at 1 and 6 months, respectively.

The adult dose of diphtheria toxoid is almost 12 times lesser in comparison with the pediatric dosage.

Note: IAP guidelines state that "temperature more than >40.5°C is a precaution and not contraindication". Park's 27th Ed states that "temperature above 40°C is a contraindication to DPT". This topic is controversial, however, IAP guidelines are commonly followed by practitioners.

Solution to Question 2:

The Annual Blood Examination Rate (ABER) is used to check the efficiency of the surveillance system for malaria. It takes into account the number of slides examined for the population under surveillance.

Solution to Question 3:

A confounding factor is a factor that is associated with both exposure and diseases and is distributed unequally in study and control groups. It can be reduced by matching.

For example, if the study is that a sedentary lifestyle can lead to coronary artery diseases, then age can act as a confounding factor as it is individually linked to both sedentary lifestyle and coronary artery diseases.

Solution to Question 4:

Early diagnosis and treatment is the best level of prevention of breast cancer. It is a secondary level of prevention.

The earlier a disease is diagnosed and treated, the better is the prognosis. The occurrence of secondary cases or any long term disability can be prevented. This is of particular importance in chronic diseases.

Early diagnosis and treatment is not as effective and economical as primary prevention. However, it may be critically important in reducing the high morbidity and mortality in diseases such as essential hypertension, cancer cervix and breast cancer.

Solution to Question 5:

The recommended vitamin A dosage for a 2-year-old child with keratomalacia is 2,00,000 IU given immediately, followed by the same age-specific dose 24 hours later and after 2 weeks.

Solution to Question 6:

Approximately 68% of a normal distribution falls within ±1 standard deviations of the mean. This is as per the empirical rule of a normal distribution.

Approximately 95% of the distribution falls within ±2 standard deviations of the mean.

Approximately 99.7% of the distribution falls within ±3 standard deviations of the mean.

Solution to Question 7:

Three key clinical signs are used to assess a sick child with cough or difficult breathing:

• Respiratory rate (RR): distinguishes children who have pneumonia from those who do not. Children are classified as pneumonia if:

• > 50/min RR for infants 2 months to 12 months

• > 40/min RR for children 12 months to 5 years of age

• Lower chest wall indrawing indicates severe pneumonia

• Stridor indicates those with severe pneumonia requiring hospital admission.

The following image gives the IMNCI classification for cough/difficulty in breathing in children.



† Not able to drink, persistent vomiting, convulsions, lethargic or unconscious, stridor in a calm child or severe malnutrition.

Signs of severe pneumonia are :

• Not able to drink

• Persistent vomiting

• Convulsions

• Lethargic or unconscious

• Stridor in a calm child

• Severe malnutrition

Solution to Question 8:

Adulteration of food is included under the concurrent list.

The legislative section of the seventh schedule is divided into three lists:

• Union list - a list of 100 items on which the parliament has exclusive power to legislate

• State list - a list of 61 items on which the state government has exclusive power to legislate

• Concurrent list - a list of 52 items on which both central and state governments have the power to legislate

Solution to Question 9:

• The principal unit of administration in India is the district under a Collector.

• Under districts are the sub-divisions, tahsils or talukas, municipalities or corporations, villages, panchayats, etc.

• As of the 2019 census, there are 725 districts in India.

Solution to Question 10:

Cross product ratio (Odds ratio) can be obtained from the case-control study.

Solution to Question 11:

In controlled tipping, one acre of land is required by the trench method to be practiced for a population of 10,000.

It is also known as a sanitary landfill. It is the most satisfactory method of refuse disposal where suitable land is available. In this method, waste material is placed in a trench or other prepared area. Then, it is adequately compacted and covered with earth at the end of the day.

Methods used in controlled tipping are:

• Trench method: Done in places where level ground is available

• Ramp method: Well suited where the terrain is moderately sloping

• Area method: Used for filling land depressions, disused quarries, and clay pits

Solution to Question 12:

The test of choice to measure the difference in means between two groups in a study, with one group being a control is unpaired T test.

Solution to Question 13:

Chi-Square($\chi 2$) test is a test used for proportions and associations.

It is a non-parametric (non-numeric) test and it is used to test hypotheses in relation to nominal scale data or categorical data.

Solution to Question 14:

Positive predictive value implies the probability of a person with a positive test, actually having the disease.

It is the ability of a test to detect true positive(TP) out of total positive(TP+FP) and it is directly proportional to the prevalence of a disease.

PPV= TP/TP+FP

Solution to Question 15:

Lead time is the period between the first possible time of detection (point A) and the usual time of diagnosis (point C).

Screening time is the time difference between the first point of diagnosis and the critical point of diagnosis.

Diagnosis of a disease before the critical point has a good outcome and reduces the severity by reversing the disease process. Diagnosis beyond the critical point treatment would be unsuccessful and/or permanent damage would have occurred.

Screening must be done in conditions where there is a considerable time lag between the onset of disease and the usual time of diagnosis. That is conditions with a long lead time.

Generation time is defined as the interval of time between the onset of infection by a host and the maximal infectivity of that host.

Solution to Question 16:

The given image shows a component bar graph. It is used to represent data in which the total magnitude is divided into different subsets or components.

A bar chart or bar graph is used mainly for nominal or ordinal scale data. They can be simple, multiple or component bar graphs.

The image given below shows a multiple bar graph showing the human losses of world war II by country. In a multiple bar chart, 2 or more sets of data are shown simultaneously.



A bar chart can be distinguished from a histogram because the rectangles are separated by the spaces. This indicates that the data forms discrete categories.

In a histogram, the rectangles are not separated by spaces as the data represented forms a continuous set instead of discrete groups.

Solution to Question 17:

Moribund patients receive the lowest priority.

This is because despite paying quite a lot of attention, the benefit is still questionable.

Solution to Question 18:

The risk of genetic diseases in consanguineous marriage between first cousins is 4-8%.

The risk of genetic diseases in non-consanguineous marriage is 2-4%.

Solution to Question 19:

A paradoxical carrier is a carrier that acquired a pathogen from another carrier.

Incubatory carriers are those who shed the infectious agents during the incubation period of the disease.

Convalescent carriers are those who continue to shed the disease agent during the period of convalescence.

Healthy carriers are those who had a subclinical infection and developed a carrier state without suffering from overt disease.

Solution to Question 20:

Varicella vaccine is contraindicated in pregnant women as it is a live vaccine.

The live attenuated varicella virus vaccine is currently recommended for children between 12 to 18 months of age who have not had chickenpox.

Solution to Question 21:

A lab coat is not a personal protective equipment.

Personal Protective Equipment (PPE) are:

• Gloves – protect hands.

• Gowns/ aprons – protect skin and clothing.

• Masks protect the mouth/nose and respirators protect the respiratory tract from airborne infectious agents.

• Goggles – protect eyes.

• Face shields – protect face, mouth, nose, and eyes.

Solution to Question 22:

Line graphs or line diagrams are the best way to plot the changes in the incidence of a disease in a given area over time.

Solution to Question 23:

Earlier the rubella infection occurs in pregnancy, greater is the damage to the foetus.

The major determinant of the extent of fetal infection in congenital rubella syndrome is the gestational age at which fetal transmission occurs.

Rubella infection in 1st trimester can cause:

• Abortions

• Still birth

• Skin lesions - blueberry muffin rash

• Congenital rubella syndrome

Solution to Question 24:

Triple therapy with tenofovir 300mg, lamivudine 300mg, and dolutegravir 50mg for 4 weeks is recommended for HIV post-exposure prophylaxis (PEP).

Steps to be undertaken after a needlestick injury are:

• Wash the wound and skin with water and soap. Do not scrub and avoid antiseptics.

• Assess exposed individual:

• Check baseline viral markers

• Monitor for acute seroconversion illness within 3-6 weeks after exposure

• If testing is positive, refer to ART center

• Assess exposure source and determine the risk of transmission

• Counseling and support

• Initiate PEP at the earliest - preferably < 2 hours and ideally < 72 hours

• 28-day prescription with drug information

• HIV test at 3 months after exposure

Community Medicine NEET 2020

Question 1:

Sharp waste is disposed in:

a) Yellow bag

b) Red bag

c) Blue container

d) White container

Question 2:

According to the Factories Act 1948, what is the maximum work hours per week including overtime work hours?

a) 48 hours

b) 72 hours

c) 60 hours

d) 66 hours

Question 3:

Which of the following was launched to ensure complete immunization of children?

a) Mission Dhanush

b) Mission Indradhanush

c) Mission Indira Yojana

d) Mission Suraksha Yojana

Question 4:

Which of the following is an example of prospective screening?

a) Cervical Pap smear in a 40 year old patient

b) Neonatal screening of a new-born baby for hypothyroidism

c) Screening of immigrants to a country

d) Urine for sugar screening in a 40 year old man

Question 5:

Monetary benefit is measured in which of the following analyses?

a) Cost effective analysis

b) Cost benefit analysis

c) Network analysis

d) Input Output analysis

Question 6:

A 24-month-old child weighing 11 kg was brought to the pediatric OPD. He had a respiratory rate of 38/min. On examination, chest indrawing was observed. What is the next step of management?

a) Oral amoxicillin for five days

b) Urgent referral to the tertiary care centre immediately

c) Administer IV antibiotics and ask the patient to be brought after 24 hours

d) Start antipyretics urgently

Question 7:

An employee diagnosed with TB gets extended sickness benefit for:

a) 1 year

b) 2 years

c) 3 years

d) 4 years

Question 8:

Chemical waste is disposed in _____ bags.

a) Yellow

b) White

c) Red

d) Green

Question 9:

Which of the following techniques is based on behavioral sciences?

a) Decision making

b) Systems analysis

c) Network analysis

d) Management by objective

Question 10:

A study was done to assess nutritional status among young children in rural and urban areas. 30 children in rural and 20 children in urban areas were malnourished. Which of the following tests of significance is the most appropriate in this case?

a) Chi Square test

b) Paired T test

c) Standard error of the mean

d) ANOVA

Question 11:

As per NCEP- ATP III, which of the following is not a criterion for metabolic syndrome?

a) Hypertriglyceridemia

b) High LDL

c) Central obesity

d) Hypertension

Question 12:

In a population study, the value of the mean is 200. If the standard deviation is 20, 68% of the population will lie between?

a) 180- 220

b) 160- 240

c) 170- 230

d) 190- 210

Question 13:

Which of the following is disposed in the yellow bag?

a) Blood bag

b) Gloves

c) Sharps

d) Urine bag

Question 14:

Calculating one variable using another variable is done by

a) Coefficient of correlation

b) Coefficient of Regression

c) Coefficient of variation

d) Coefficient of determination

Question 15:

The best indicators for routine monitoring of air pollution are:

a) Sulfur dioxide, smoke, and suspended particles

b) Sulfur dioxide, lead, and particulate matter

c) Sulfur dioxide and carbon monoxide

d) Carbon monoxide and hydrogen sulfide

Question 16:

A person wants to study about the relationship between smoking and lung cancer, so he collects data about people diseased with lung cancer from government hospitals and the number of cigarette packets sold at the same time period. Which type of study is this?

a) Cross sectional

b) Ecological

c) Experimental

d) Quasi-experimental

Question 17:

Project MONICA is related to:

a) Cardiovascular diseases

b) Cervical Cancer

c) Breast Cancer

d) Road Traffic Accidents

Question 18:

Which of the following compounds is predominantly responsible for the disinfecting property of bleaching powder?

a) Hypochlorite ion

b) Hydrochloric acid

c) Hypochlorous acid

d) Chloride ion

Question 19:

The difference between the incidence of disease among exposed and incidence among non exposed is given by which of the following variables?

a) Relative risk

b) Attributable risk

c) Population Attributable risk

d) Odds Ratio

Question 20:

In Sustainable Development Goal 3, the target 3.1 aims to reduce the Maternal Mortality Ratio by 2030 to ____ (per 1 lakh live births):

a) <70

b) <100

c) <50

d) <130

Question 21:

A patient comes with a ■history of clean-cut injury which is not lacerated. He says that he received his last dose of tetanus vaccination about 10 years ago. What should be given next?

a) Full course of tetanus vaccination

b) Single dose tetanus toxoid

c) Tetanus toxoid + immunoglobulin

d) No vaccination needed

Question 22:

The variation in a data set is compared with that of another data set using:

a) Variance

b) Coefficient of variation

c) Standard error of mean

d) Standard deviation

Question 23:

Pandemic that occurred in 2009 was due to which influenza virus?

a) H1N1

b) H5N1

c) H5N7

d) H3N2

Answer Key

Question No.	Correct Option
1	d
2	c
3	b
4	c
5	b
6	a
7	b
8	a
9	d
10	a
11	b
12	a
13	a
14	b
15	a
16	b
17	a
18	c
19	b
20	a
21	b
22	b
23	a

Detailed Explanations

Solution to Question 1:

Sharp waste is disposed of in the puncture and leak-proof white containers.

Solution to Question 2:

Maximum work hours per week including overtime work hours is 60 hours.

Solution to Question 3:

Mission Indradhanush was launched by the Ministry of Health and Family Welfare in 2014 to fully immunize children (>90%) who are either partially immunized or completely unimmunized by 2020. It was later modified in 2017 to the Intensified Mission Indradhanush, to achieve the same goal by December 2018.

Intensified Mission Indradhanush includes 12 vaccine-preventable diseases (updated from the 7 diseases that Mission Indradhanush focussed on). 4 rounds of immunization are proposed to be held for 7 days each month (from the 7th- 13th of each month).

The diseases included in Mission Indradhanush are:

• TB

• Polio

• Hep B

• Diphtheria

• Pertussis

• Tetanus

• Measles

The diseases included recently in Intensified Mission Indradhanush are:

• Rotavirus diarrhoea

• Rubella (as MR vaccine)

• H. influenzae B diseases (as part of the pentavalent vaccine)

• Pneumococcal meningitis

• Japanese Encephalitis

The Intensified Mission Indradhanush 2.0:

The government of India introduced IMI 2.0 to accelerate the vaccine drive and extend the coverage to children and pregnant women in identified districts and blocks from December 2019-March 2020.

Note: Currently, the government has completed Intensified Mission Indradhanush 5.0:

Intensified Mission Indradhanush (IMI 5.0), the flagship routine immunization campaign of the Union Ministry of Health and Family Welfare concluded all 3 rounds in October 2023. This is the first campaign to be conducted across all the districts in the country and includes children up to 5 years of age (Previous campaigns included children up to 2 years of age). It also had a special focus on improvement of Measles and Rubella vaccination coverage.

Solution to Question 4:

Screening of immigrants to a country is an example of prospective screening.

Screening of disease is done to detect unrecognized disease or defect, in apparently healthy individuals, by means of rapidly applied tests, examinations, or other procedures.

The difference between prospective and prescriptive screening is given below:

Note: Most common neonatal disease to be screened for is neonatal hypothyroidism.

	Prospective screening	Prescriptive screening
Definition	People screened for others benefit	People screened solely for their own benefit
Essential purpose	Disease control	Case detection
Request for screening	Specific request from the authority	Request from physician
Example	Immigrantsfor infectious diseases like tuberculosis syphilis, yellow fever, etc;HIV screeningduring antenatal visits, sex workers.Screening ofblood donors.	Neonatalscreening for hypothyroidism.Pap smear for detection of cervicalcancer.Testing of urine sugar for diabetes mellitus.

Solution to Question 5:

The monetary benefit is measured in a cost-benefit analysis.

In cost-benefit analysis, the economic benefits of any program are compared with the cost of the program. The benefits are expressed in monetary terms to determine whether a given programme is economically sound and to select the best out of several alternate programmes.

The main drawback with this technique is that the benefits in the health field, as a result of a particular programme, cannot always be expressed in monetary terms. We generally express the benefit in terms of births or deaths prevented, or illness avoided or overcome.

In cost-effective analysis, the benefit is expressed in terms of results achieved. (Eg: number of lives saved or number of days free from disease).

Input-output analysis: how much input is needed to produce a unit amount of output. In the health field, input refers to all health service activities which consume resources (manpower, money, materials, and time); and output refers to useful outcomes such as cases treated, lives saved, or inoculations performed.

Network analysis: a network is a graphic plan of all events and activities to be completed in order to reach an end objective.

Management Methods in Health Programmes:

Methods based on behavioral sciences	Quantitative methods
Organisational designPersonnel managementCommunicationInformation systemsManagement by objectives (MBO)	Cost–Benefit analysisCost-Effective analysisCost–AccountingInput–Output analysisModelSystems analysisNetwork analysis- Includes Programme Evaluation and Review Technique (PERT) and Critical Path Method (CPM)Planning–Programming–Budgeting System (PPBS)Work samplingDecision making

Solution to Question 6:

From the given clinical picture, it appears that the child has pneumonia as diagnosed from the presence of chest indrawing. The next best step of management is to give oral amoxicillin for five days and send the child home.

This is based on the revised WHO classification and treatment of childhood pneumonia at health facilities. Pneumonia with general danger signs like inability to drink, persistent vomiting, convulsions, lethargic or unconscious, stridor in a calm child or severe malnutrition needs antibiotics with urgent referral to higher centre.

The following image distinguishes the previous and revised classification and treatment of the childhood pneumonia.



Solution to Question 7:

Extended sickness benefit is given for a period of 2 years.

Solution to Question 8:

Chemical waste is disposed of in yellow-colored containers or non-chlorinated plastic bags.

Solution to Question 9:

Management by objective is based on behavioral sciences.

Management methods used in health programmes are either quantitative or based on behavioral sciences.

Methods based on behavioral sciences include:

- Organizational design
- Personnel management
- Communication
- Information systems
- Management by objectives (MBO)

Quantitative methods include:

- Cost-benefit analysis
- Cost-effective analysis

• Cost–accounting

• Input-output analysis

• Model

• Systems analysis

• Network analysis - includes programme evaluation and review technique (PERT) and critical path method (CPM)

• Planning–programming–budgeting system (PPBS)

• Work sampling

• Decision making

Solution to Question 10:

The most appropriate test of significance, in this case, is the chi-square test. It is used to compare qualitative data between 2 groups with independent measures.

Option B: Paired t-test compares paired data or pre-intervention and post-intervention data of the same group. This doesn't apply here as the groups are not paired.

Option C: The standard error of the mean is calculated for quantitative data, with a measure of central tendency as mean.

Option D: ANOVA is used to compare more than 2 groups.

Solution to Question 11:

As per NCEP- ATP III, high LDL is not included in the definition criteria of metabolic syndrome.

Metabolic syndrome is present if three or more of the following five criteria are met, which is given by NCEP-ATP III (National Cholesterol Education Program and Adult Treatment Panel III). It includes:

• Central obesity: waist circumference >102 cm (M), >88 cm (F)

• Hypertriglyceridemia: triglyceride level ≥ 150 mg/dL

• Low HDL cholesterol: <40 mg/dL and <50 mg/dL for men and women, respectively.

• Hypertension: blood pressure ≥130 mmHg systolic or ≥ 85 mmHg diastolic.

• Fasting plasma glucose level: ≥100 mg/dL or previously diagnosed type 2 diabetes.

Solution to Question 12:

68% of the population will lie between 180- 220.

In a normal distribution curve, 68% of the values lie within the mean +/- 1 standard deviation and the mean=median=mode. Since the mean = 200 and the standard deviation = 20:

68% of population will lie between Mean +/- 1SD, that is, 200 +/- 20 = 180-220

95% of population lies between Mean+/-2SD, 200+/- 2*20 = 160- 240

99% of population lies between Mean+/-3SD, 200+/- 3*20 = 140- 260

Solution to Question 13:

Blood bag is disposed in the yellow bag.

Option B: Gloves are disposed in the red bag.

Option C: Sharps are disposed in the white container.

Option D: Urine bags are disposed in the red bag.

Solution to Question 14:

Calculating one variable using another variable is done using coefficient of regression.

Solution to Question 15:

The best indicators for routine monitoring of air pollution are sulfur dioxide, smoke, and suspended particles.

Solution to Question 16:

The type of study described is an ecological study as the unit of study is the population. The number of cigarette packets sold at the same time is data about the population.

Option A: The unit of study in a cross-sectional study is an individual.

Option C: There is no experiment done here. Hence, it is not experimental study.

Option D: Quasi-experiments are nonrandomized, pre-post intervention studies done when it is not feasible or ethical to conduct a randomized controlled trial. The unit of study is therefore patients or individuals and not population.

Solution to Question 17:

Project MONICA is related to cardiovascular diseases.

It stands for multinational monitoring of trends and determinants in cardiovascular diseases. It was done to study the reasons for the changes in the trends of cardiovascular diseases in different countries.

Solution to Question 18:

The disinfectant property of bleaching powder is predominantly due to the action of hypochlorous acid. It is 70-80 times more effective than hypochlorite ions.

Bleaching powder, also known as chlorinated lime ($CaOCl_2$), has 33 per cent of chlorine in it. When chlorine is added to water, there is the formation of hydrochloric and hypochlorous acids. The hydrochloric acid is neutralized by the alkalinity of the water. The hypochlorous acid ionizes to form hydrogen ions and hypochlorite ions, as follows:

$H_2O + Cl_2 \rightarrow HCl + HOCl$

$HOCl \rightarrow H + OCl$

Solution to Question 19:

The difference between the incidence of disease among exposed and incidence among non-exposed is given by attributable risk.

Attributable risk is often expressed as a percentage.

$$\text{Attributable risk (AR)} = \frac{\text{Incidence of disease rate among } \mathbf{exposed} - \text{Incidence of disease rate among } \mathbf{non\text{-}exposed}}{\text{Incidence of disease rate among } \mathbf{exposed}} \times 100$$

Solution to Question 20:

In Sustainable Development Goal 3, the target 3.1 aims to reduce the Maternal Mortality Ratio by 2030 to <70 per 1 lakh live births.

There are 17 global goals and 169 targets set for 2030 as part of the official Sustainable Development Goals of the Agenda 2030.

Solution to Question 21:

A single dose of tetanus toxoid is given as the wound is clean and the last dose was received more than 10 years ago.

Solution to Question 22:

The variation in one data set is compared with that of another data set using the coefficient of variation (CV). It measures the relative variability by using the mean and standard deviation.

Solution to Question 23:

Recent influenza pandemic occurred in 2009 due to the emergence of a novel H1N1 influenza A virus.

Past influenza pandemics:

- 1968 pandemic due to H3N2 virus
- 1957-58 pandemic due to H2N2 virus
- 1918 pandemic due to H1N1 virus

Community Medicine INI-CET 2021

Question 1:

Select the correct option regarding Mobile Medical Units (MMUs).

a) Only 1

b) 1 and 2

c) 1, 2, and 3

d) 1, 2, 3, and 4

Question 2:

Which of the following statements is true?

a) In a population with low prevalence of the disease, positive result with a highly sensitive test means true positive

b) Sensitivity depends on prevalence of the disease

c) Positive predictive value depends on prevalence of disease

d) Specificity depends on prevalence of the disease

Question 3:

You are starting services for hypertension in your PHC. 50 patients who required antihypertensive treatment were transferred from another center. 40 of them were on amlodipine (5mg PO) and 10 were on lisinopril (10 mg PO) as they had contraindications to the use of amlodipine. The drugs are supplied at the PHC on a monthly basis and you have to place an order for their medications. What is the number of tablets that you will order and the reorder factor?

a) 1000, rf= 3

b) 1200, rf=2

c) 1400, rf= 3

d) 1600, rf= 2

Question 4:

Match the following STI kit colors and their use:

a) A-4, B-3, C-2, D-1

b) A-4, B-1, C-3, D-2

c) A-1, B-4, C-3, D-2

d) A-3, B-2, C-4, D-1

Question 5:

Compared to a pregnant female, a lactating female would require a higher level of nutrient supplementation for which of the following?

a) Vitamin A

b) Calcium

c) Iron

d) Folic acid

Question 6:

What does the point labeled C denote?



©MARROW

a) Onset of symptoms

b) Onset of pathological changes

c) Usual time of diagnosis

d) Exposure to etiologic agent

Question 7:

Certain new vaccines are being studied in the market and their efficacies are compared and presented in the image as shown below. Which of the newer vaccines can be used with the same efficacy as compared to the previous vaccine?



a) D only

b) D and E

c) C, D and E

d) A, B, C and D

Question 8:

As per the verbal autopsy report 2010-2013, arrange the most common cause of death among 1-4-year-old children in descending order:

a) 2 > 1 > 3 > 4

b) 4 > 1 > 2 > 3

c) 1 > 2 > 4 > 3

d) 2 > 1 > 4 > 3

Question 9:

The most common mode of absorption of inorganic lead in industries leading to lead poisoning is:

a) Absorption through skin

b) Inhaled lead dust

c) Ingestion of contaminated food and water

d) Contaminated hands

Question 10:

Pentavalent vaccine includes which of the following?

a) Diphtheria, Pertussis, Tetanus , HBV, Injectable Polio

b) Diphtheria, Pertussis, Typhoid, HBV, Hib

c) Diphtheria, Pertussis, Tetanus, Haemophilus influenza B, Pneumococcal vaccine

d) Diphtheria, Tetanus, HBV, Haemophilus influenza B, Pertussis

Question 11:

A nurse keeps the bins as shown in the image in the hospital ward. Which of the following items would go into the black bin?

a) Soiled linen bedsheet

b) Glove paper cover

c) Gloves

d) Contaminated gloves

Question 12:

A homogenous sample of 4 groups was taken as shown below. The height and weight of each group had a correlation coefficient of 0.6. What will be the total coefficient of correlation of the whole group?



a) Equal to 0.6

b) Less than 0.6

c) More than 0.6

d) Skewed coefficient of 0.6

Question 13:

The graphical representation for flow cytometry analysis is done by which of the following?

a) Bar diagram, dot plot

b) Line diagram, dot plot

c) Histogram, dot plot

d) Pie chart, dot plot

Question 14:

Administration of which of the following vaccines is associated with the risk of intussussception?

a) Rotavirus vaccine

b) Oral Polio vaccine

c) Adenovirus vaccine

d) Measles vaccine

Question 15:

Interpret the following graph.



a) Normal, negatively skewed, positively skewed, skewed with outliers

b) Normal, positively skewed, negatively skewed, normal with outliers

c) Normal, negatively skewed, positively skewed, normal with outliers

d) Skewed with outliers, positively skewed, negatively skewed, normal

Question 16:

The standardized age mortality ratio is an indicator for which of the following?

a) Death rate of a population adjusted to a standard age distribution

b) Death rate of a population adjusted to a standard sex distribution

c) Removal of bias due to different age group population in different countries

d) To eliminate age structure of different countries

Question 17:

According to WHO, which of the following are components of the Human Development Index (HDI)?

a) Life expectancy at birth, literacy and standard of living

b) Life expectancy at 1 year, literacy and standard of living

c) Infant mortality rate, literacy and standard of living

d) Life expectancy at 1 year, literacy and infant mortality rate

Question 18:

The external validity of a screening test is related to?

a) Stability

b) Objectivity

c) Generalizability

d) Repeatability

Question 19:

All the following tests are related to pasteurization of milk except

a) Phosphatase test

b) Iodine test

c) Standard plate count

d) Methylene blue reduction test

Question 20:

Over the past few years, there is an increase in the incidence of non-communicable diseases. What is this type of trend called?

a) Seasonal

b) Cyclical

c) Periodic

d) Secular

Question 21:

Which of the following is not a part of case-control studies?

a) Follow up

b) Matching

c) Strength of association

d) Selection of study subjects

Question 22:

Which of the following is the best study design to demonstrate the relationship between cause and effect?

a) Case control study

b) Cohort study

c) Cross-sectional study

d) Clinical trial

Question 23:

Hepatitis B subunit vaccine contains which of the following antigen?

a) HbeAg

b) HbsAg

c) HbcAg

d) HBV DNA

Question 24:

Arrange the following causes of neonatal mortality from maximum to the least (according to the verbal autopsy report 2010-2013 data).

a) 1>2>4>3

b) 1>4>2>3

c) 4>1>2>3

d) 1>2>3>4

Question 25:

Which of the following is not involved in meta-analysis?

a) Selection

b) Analysis

c) Randomization

d) Abstraction

Question 26:

A new drug has been introduced into the market which was found to decrease mortality but it does not cure the disease. Which of the following is a true statement regarding prevalence and incidence?

a) Decrease in incidence

b) Decrease in prevalence

c) Increase in incidence

d) Increase in prevalance

Answer Key

Question No.	Correct Option
1	c
2	c
3	d
4	b
5	b
6	a
7	c
8	d
9	b

10	d
11	b
12	a
13	c
14	a
15	b
16	a
17	a
18	c
19	b
20	d
21	a
22	b
23	b
24	a
25	c
26	d

Detailed Explanations

Solution to Question 1:

Statements 1, 2, and 3 are correct. MMUs can be run by either Government or an external agency, but the drugs and medical supplies are to be provided always by Government only.

Mobile Medical Units (MMU) is a major initiative under NRHM to provide a range of health care services for people living in remote, inaccessible, un-served, and underserved areas in both rural as well as urban areas, with the aim of taking healthcare services to the doorsteps of such people.

There are 3 models of operationalizing MMUs:

• Government operated MMU

• Operation of MMU on outsourcing basis by an external agency but capital expenditure, drugs, and supplies provided by Government.

• Outsourcing by an external agency which provides both capital expenditure and operational expenditure. However, drugs and supplies are to be provided by Government.

MMUs provide primary care services for common diseases (communicable and non-communicable), RCH services, screening tests, and referral to higher facilities and specialists as required. Clinical services are provided by a Medical Officer and team, and mobilization and outreach services are provided by ANMs and ASHA.

Solution to Question 2:

The positive predictive value (PPV) of a screening test depends on the prevalence of the disease. The positive predictive value (PPV) of a test indicates the probability that a patient with a positive test result has actually contracted the disease i.e., the likelihood that a person with a positive test result truly has the disease.

If the prevalence of a disease in a population is high, the more accurate will be the positive predictive value. Therefore, in a population with a high prevalence of the disease, a positive result with a highly sensitive test means true positive. Example: Hypothyroidism is more prevalent in hilly areas than in coastal areas. If someone from a hilly area tests positive he is more likely to have the disease than someone who tests positive in a coastal area.

Likewise, if the prevalence decreases, the PPV will also become low. So in a population with a low prevalence of the disease, positive result with a highly sensitive test need not always mean true positive as it can produce an unacceptably high proportion of false positive results.

Predictive value is affected the most by prevalence. However, both sensitivity and specificity can change the predictive value.

Solution to Question 3:

The Quantity to Order (QO) or the reorder quantity of medicines to be placed at each order interval is calculated using the reorder formula as given below:

QO = CA x (LT + PP) = CA x Rf

where QO is the quantity to order, CA is average consumption in a month, LT is lead time, PP is procurement period and Rf is the reorder factor. Reorder factor (Rf) is the sum of lead time (LT) and procurement period (PP). Lead time (LT) is the time between the initiation of a purchase order and receipt at the warehouse from the supplier. The procurement period (PP) covers the time until the next regular order will be placed.

In the given question, lead time (LT) and procurement period (PP) are 1 month each.

Average consumption (CA) in a month = (40 tablets of amlodipine + 10 tablets of lisinopril) x 30 days = 1500 tablets

Reorder factor (Rf) = LT + PP = 1 + 1 = 2

Since these patients got transferred from another center there is no safety stock for them at present at the facility. So in order to prevent stockouts for them, an extra number of tablets must be ordered, which makes 1600 tablets with a reorder factor (Rf) of 2 the correct option.

Factors determining the reorder formula are:

• Average consumption or demand of the drug (CA) in a month

• Supplier Lead time (LT): is the time between initiation of a purchase order and receipt at the warehouse from the supplier.

• Safety stock (SS): is the stock that should always be on hand to prevent stockouts. The safety stock level and reorder level will be the same if consumption and lead times are predictable and stable.

• Procurement period (PP): it covers the time until the next regular order will be placed. In a scheduled system this is in multiples of one month and in a perpetual system, it is counted in days or weeks. Note: Quantity ordered (QO) + Safety Stock (SS) must cover the time until the next order is received, which is given by the Reorder factor (Rf) = Procurement Period (PP) + Lead Time (LT).

• Reorder level: is the quantity of remaining stock that should trigger a reorder of the item. In a basic purchasing formula, reorder level is calculated by multiplying the average lead time by the average consumption of the drug during the lead time.

• Maximum stock level: is the same as the target stock level, which is required to satisfy demand until the next order is received.

• Stock position: is the sum of stock on hand (working and safety stock) and stock on order, minus any stock back-ordered to clients.

Solution to Question 4:

The correct match is A-4, B-1, C-3, D-2

The National Aids Control Organization (NACO) has provided color-coded kits for the syndromic management of sexually transmitted diseases:

Kit	Syndrome/s
Kit 1(Grey)	Urethral dischargeCervical di schargeAno-rectal dischargeP ainful scrotal swelling
Kit 2 (Green)	Vaginal discharge/ Vaginitis
Kit 3 (White)	Non-herpetic genital ulcers
Kit 4(Blue)	Non-herpetic genital ulcers if allergic to penicillin
Kit 5(Red)	Herpetic genital ulcers
Kit 6 (Yellow)	Lower abdominal pain
Kit 7(Black)	Inguinal bubo
Kit 8(Brown)	Anorectal discharge syndrom e

Solution to Question 5:

According to the Recommended daily allowance (RDA) guidelines set by the National Institute of Nutrition (NIN), the requirement of calcium intake is increased in a lactating woman (1200mg/d) when compared to that of a pregnant woman (1000mg/d). Calcium intake of 1200mg/d is advised in postmenopausal women as well.

The RDA of vitamin A is increased by 50mcg in lactation as compared to pregnancy. But the increased vitamin A demand is met from the diet and it is not given as a supplement nutrient. Vitamin A supplements are given only when there is vitamin A deficiency.

The RDA of iron and folic acid is decreased during lactation compared to pregnancy.

The salient features of updated RDA guidelines, 2020 by NIN are:

Variable	Non pregnant female	Pregnancy	Lactation			
1st TM	2nd TM	3rd TM	0-6 m	6-12 m		
Body weight	55 kg	55 kg + 10 kg in entire pregnancy				
Energy (Kcal/d)SedentaryModerateHeavy	166021302720	+350	+600	+520		
Protein (g/d)	45.7	45.7	+9.5g	+22g	+16.9 g	+13.2 g
Carbs (g/d)	130	175g	200g			
Iron (mg/d)	29 mg/d	27 mg/d	23 mg/d			
Iodine (mcg/d)	150	250	280			
Calcium (mg/d)	1000	1000	1200			
Vit A (mcg/d)	840	900	950			
Folate (mcg/d)	220	570	330			

Solution to Question 6:

The given diagram shows the natural history of disease in an individual over time, in which the point labeled C denotes the onset of symptoms. The natural history of disease refers to the progression of a disease in an individual over time, in the absence of treatment.

Exposure (point A): the disease process begins with appropriate exposure to or accumulation of factors sufficient for the disease process to begin in a susceptible host. For example in infectious disease, the exposure is a microorganism. For lung cancer, it is any factor that initiates the process, such as asbestos fibers or components in tobacco smoke.

Pathological changes (point B): after the disease process has been triggered, pathological changes occur without the individual being aware of it. This stage of subclinical disease, extending from the time of exposure to onset of symptoms, is usually called the incubation period for infectious diseases, and the latency period for chronic diseases. During this stage, the disease remains asymptomatic.

The onset of clinical symptoms (point C): it is the beginning of the clinical stage of the disease.

The usual time of diagnosis (point D): the onset of symptoms is followed by a short interval (varies from disease to disease) after which the patient presents to a healthcare setup where a diagnosis of the disease is made. Based on treatment and prognosis, the patient can progress to either

recovery, disability, or death.



Solution to Question 7:

Vaccines C, D, and E can be used with the same efficacy as compared to the previous vaccine, as their confidence interval lies within the zone of equivalence.

In a classical randomized controlled trial (RCT), the aim is to evaluate a difference between two drugs by proving the superiority of the new drug over the standard drug. However, an equivalence study is carried out with the aim of showing there are no significant differences in the efficacy between the new drug and the standard drug.

To define 'no significant differences' the limit of equivalence (δ) must be defined first. δ must be set as superiority margin (+δ) and non-inferiority margin (-δ), so that they are clinically appropriate and the differences can be tolerated as clinically irrelevant. To conclude equivalence the confidence interval (CI), i.e., the whisker (tail) of the box-whisker plot must lie within the zone of equivalence.

Therefore in the given question, vaccines C, D, and E have CI in the zone of equivalence and thus have the same efficacy as the previous vaccine. Vaccines F and G lie outside the zone of equivalence but well within +δ, making them superior to the previous vaccine. Vaccines A and B lie outside the zone of equivalence but well within -δ, making them non-equivalent and at the same time non-inferior to the previous vaccine. If the CI of any vaccine includes or crosses the -δ limit, they will be considered inferior to the standard vaccine.



Note: A superiority trial is done to show that the new drug is superior to the standard drug. A non-inferiority trial is done to show that the new drug is not worse than the standard drug. This is done for newer interventions or drugs that may be less costly, less invasive, and have fewer side effects.

Solution to Question 8:

According to the verbal autopsy report of 2010-2013, the most common cause of death among 1-4 year old children in descending order is Pneumonia > Diarrhea > Injuries > Malaria.

According to the Verbal autopsy report 2010-2013, the top 5 causes of death among children aged 1-4 years are:

• Pneumonia

• Diarrheal diseases

• Injuries

• Other non-communicable diseases

• Malaria

1-4 year mortality rate also known as the child death rate is the number of deaths of children aged 1-4 years per 1000 children in the same age group in a given year. It thus excludes infant mortality. It is given by the formula,

1-4 year mortality rate (child death rate) = (no. of deaths of children aged 1-4 years during a year x 1000) / Total no. of children aged 1-4 years at the middle of the year.

It is a more refined indicator of the social situation than the infant mortality rate as it reflects adverse environmental health hazards like malnutrition, poor hygiene, infections, and accidents. It is not affected by perinatal hazards. In the 1-4 year group, second year is the period with the highest risk of death (mainly due to childhood infections like measles, whooping cough, diphtheria, diarrhea, and ARI), after which the rates decline progressively.

Solution to Question 9:

The most common mode of industrial lead poisoning (plumbism) is due to inhalation of fumes and dust of lead. Lesser common modes include ingestion of small quantities of lead trapped in the upper respiratory tract, ingestion due to contaminated hands, and absorption through the skin (only organic lead). Lead poisoning is a notifiable and compensable disease in India.

Lead exerts its toxic action by combining with the SH-group of enzymes involved in porphyrin synthesis and carbohydrate metabolism, disrupting normal membrane permeability, and causing potassium leakage in RBCs. Symptoms of lead toxicity include:

Inorganic lead toxicity:

• Abdominal colic, also known as Painter's colic

• loss of appetite, obstipation, constipation

• Anemia (stippling of red cells)

• Wrist drop or foot drop

• Blue-line on the gums known as Burtonian line

Organic lead toxicity (mainly affects CNS):

• Headache

• Delirium, confusion

• Insomnia

Earliest and consistent sign: Facial pallor.

Clinical symptoms of lead poisoning occur when blood levels are >70µg/dl. Measurement of Coproporphyrin (CPU) in urine is a useful screening test. Periodic examination of workers must be done and atmosphere lead concentration must be kept <2mg/cubic meter at the workspace. Handwashing before eating and prohibition of taking food in workplaces is essential. D-penicillamine is used as a chelating agent for treatment.

Solution to Question 10:

Pentavalent vaccine is a combination of five vaccines that provide protection against 5 life-threatening diseases such as diphtheria, pertussis, tetanus, hepatitis B, and Haemophilus influenzae type b (Hib). Administration of pentavalent vaccine reduces the number of pricks to a child.

It is given at 6, 10, and 14 weeks in the immunization program, and can replace the need for separate administration of Hepatitis B and DPT vaccines. However, the Hepatitis B birth dose and DPT booster doses should be continued. It is given 0.5ml intramuscularly in the anterolateral aspect of the mid-thigh.

Pentavalent vaccine is a freeze-sensitive vaccine and therefore should be stored and transported at 2- 8° C temperature. It is a very safe vaccine with a few mild side effects like pain, redness and

swelling at the injection site, fever, vomiting, abnormal crying, and loss of appetite.

Solution to Question 11:

Glove paper cover is a general solid waste that is to be collected in the black bin.

All non-infectious general wastes like food waste, paper waste, and mineral water bottles should be discarded in black bins.

Soiled linen bedsheets contaminated by blood or body fluids should be disposed in the yellow bin. All plastic recyclable wastes are discarded in red bins (gloves and contaminated gloves).

Solution to Question 12:

In a homogenous sample or population, the total coefficient of correlation (r) will be the average of the individual correlation coefficients. Therefore in the given condition, it will be the same as 0.6

The given scattergram shows the correlation coefficient of height and weight of 4 homogenous samples. Averaging methods are appropriate when several different samples are independently drawn from the same or a homogenous population and the interest is in aggregating those values.

Therefore the total correlation coefficient of the whole group will be:

r = (sum of correlation coeffiecient of each sample) ÷ (total no. of samples)

= (0.6 + 0.6 + 0.6 + 0.6) ÷ 4 = 2.4 ÷ 4

= 0.6

Note: A correlation between 2 variables does not demonstrate any causal relationship between those variables, no matter how strong it is. Correlation is merely a measure of a statistical association between the variables. Therefore inferring a causal relationship between two variables on the basis of a correlation is a common and fundamental error.

Solution to Question 13:

Flow cytometry results are typically presented by two-dimensional dot plots or one-dimensional histograms.

Histograms compare only a single parameter. A large number of events detected at one particular intensity will be displayed as a spike on the histogram. They are fast to read and easy to understand.



Dot plots can compare 2 or more parameters simultaneously on a 2D scatterplot. Each distinct event is represented as a single dot and events with similar intensities will cluster together in the same region on the scatter plot. They are a more complex, illustrative representation of data.



Pie charts and bar diagrams are used for representing qualitative data. A line diagram is used to show the trend of events or changes over passage of time.

Solution to Question 14:

Rotavirus vaccine is associated with a potential risk of intussusception, especially when the first dose of these vaccines is given to infants aged >12 weeks. Because of this risk, the rotavirus vaccine is not a part of catch-up immunization programs, as the age of the child is difficult to ascertain in such programs.

Rotavirus vaccines are oral, live attenuated vaccines and are of 3 types:

• Rotarix (human monovalent) vaccine is administered in 2 oral doses:

• First dose: At 6 weeks, no later than 12 weeks (due to the risk of intussusception).

• Second dose: With a gap of 4 weeks from the first dose, to be given by 16 weeks and no later than 24 weeks.

• RotaTeq (pentavalent bovine-human reassortant) vaccine is administered in 3 oral doses at 2, 4, and 6 months. The first dose is given between 6-12 weeks and subsequent doses within a 4-10 weeks gap. All the 3 doses should be given before 32 weeks.

• Rotavac (monovalent liquid frozen) vaccine is administered in 3 oral doses, 4 weeks apart at 6,10 and 14 weeks. It should not be administered to children more than 8 months of age.

Solution to Question 15:

The graph shown is of a boxplot, and option B represents the correct answer. 1 is normal, 2 is positively skewed, 3 is negatively skewed, and 4 is normal with an outlier.

A boxplot is a method for graphically depicting groups of numerical data through their quartiles. Box plots may also have lines extending vertically from the boxes (whiskers) indicating variability outside the upper and lower quartiles, hence the terms box-and-whisker plot.

It shows the five-number summary of a set of data including,

• Minimum score

• First (lower) quartile

• Median

• Third (upper) quartile and

• Maximum score



In the box plot, a simple box extends from the 25th percentile (lower limit of the box) to the 75th percentile (upper limit of the box) and the line through the box indicates the median value (50th percentile).

The top and bottom ends of the whiskers are the highest and lowest value of that group excluding outliers

Interpreting the box plot:

• When the median is in the middle of the box, and the whiskers are about the same on both sides of the box i.e when each quartile is of the same length, then the distribution is symmetric or normal (as shown in box plot-1).

• When the median is closer to the bottom of the box, and if the whisker is shorter on the lower end of the box, then the distribution is positively skewed (skewed right as shown in boxplot-2).

• When the median is closer to the top of the box, and if the whisker is shorter on the upper end of the box, then the distribution is negatively skewed (skewed left as shown in boxplot-3).

• Small circles or unfilled dots are drawn on the chart to indicate where suspected outliers lie. Filled circles are used for known outliers (as shown in boxplot-4).



Solution to Question 16:

The age standardized mortality ratio (SMR) is an indicator of the death rate of a population adjusted to a standard age distribution. It is obtained by indirect standardization and cannot be used to compare multiple groups with different age structures.

Standardized death rate (SDR) obtained by direct standardization enables comparisons of mortality rates between populations with different age compositions by eliminating the confounding effect caused by variations in age structures (Options C and D). This adjustment process can also account for factors such as sex, race, and parity.

Standardization can be achieved through two methods: direct or indirect. In both cases, a standard population is selected as a reference, characterized by known numbers in each age and

sex group.

The direct method of standardization requires age-specific death rates (ASDR) along with the corresponding subgroup sizes. By applying these rates to the standard population, the expected deaths across all age groups can be calculated assuming that both populations share the same age-specific death rates. Dividing the total expected deaths by the standard population yields the SDR.

The calculated standardized death rate in direct standardization can be compared with different population groups with different age group compositions.

The table below shows the calculation of age-specific death rates per 1000 population for a sub-population:

Crude death rate = 64/7500 = 8.533 per 1000 population.

Applying ASDR to the standard population, we can get the SDR

SDR per 1000 population = (212/23500) x 1000 = 9.021

Indirect standardization allows adjustment when age-specific rates are either unavailable or unstable due to small numbers. Examples of indirect standardization methods include standardized mortality ratio (SMR), life table analysis, survival analysis, regression analysis, and multivariate analysis.

SMR estimates expected deaths, assuming that the study group experiences the same death rates as a standard population.

The table below shows the calculation of the standardized mortality ratio for a study group population.

SMR = Observed deaths / Expected deaths x 100 = (65/59.85) x 100= 108.6

If the SMR exceeds 100, it indicates that the study group population carries a higher mortality risk compared to the overall population. However, it's important to note that the SMR cannot be directly compared to other groups with different age compositions.

Age(years)	Mid-year population	Deaths in the year	ASDR	Standard population	Expected deaths
0-15	2000	10	5	6000	30
15-45	4000	24	6	12000	72
>45	1500	30	20	5500	110
Total	7500	64		23500	212

Age(years)	National population death rates per 1000	Study group population	Observed deaths	Expected deaths
0-15	3.0	1200	*	3.6
15-45	6.5	2500	*	16.25
>45	10	4000	*	40
Total			65	59.85

Solution to Question 17:

The components of the Human Development Index (HDI) include life expectancy at birth, literacy, and standard of living.

Human Development Index (HDI) is a measure of a country's average achievements in three dimensions of human development:

• A long and healthy life, as measured by life expectancy at birth

• Knowledge (literacy), as measured by mean years of schooling and expected years of schooling, and

• A decent standard of living, as measured by GNI (gross national income) per capita.

HDI values range from 0 to 1, with 1 as the maximum value.

Note- Life expectancy at 1 year, literacy and infant mortality rate are components of the Physical quality of life index (PQLI).

Solution to Question 18:

External validity of a screening test is related to generalizability.

Validity (accuracy) refers to what extent the test accurately measures which it sets out to measure. It refers to the closeness with which measured values agree with "true" values.

While assessing the validity of a research study, two kinds of validity are involved:

• External validity: If the study sample is truly representative of the population from which it was drawn, then the study is said to have external validity. Its findings can be applied to the entire population as well and therefore it is said to be generalizable.

• Internal validity: Here the results of the study are valid for the sample of patients who were actually studied.

Note: Precision (reliability or reproducibility) is a measure of the repeatability of a test. That is, the test must give consistent results when repeated more than once on the same individual or material, under the same conditions.

Solution to Question 19:

All the above tests are related to the pasteurization of milk except the iodine test. It is done to test if milk is adulterated with starch or cereal flour.

The iodine test is done by adding 2 drops of iodine solution containing potassium iodide and iodine to 1ml of milk. The appearance of a blue color indicates that the milk is adulterated with starch. A control sample of milk that is not adulterated with starch will develop a slightly yellowish color.

Pasteurization can be done by the following methods,

• Holder/Vat method: 63-66 °C for 30 minutes and then quickly cooled to 5°C.

• HTST method (High-temperature short time): 72°C for at least 15 seconds, and then cooled rapidly to 4°C.

• UHT method (Ultra-high temp): Rapidly heated to 125 °C for a few seconds, then cooled and bottled as quickly as possible.

Tests in pasteurization:

• Phosphatase test: It is based on the fact that raw milk contains phosphatase enzyme which gets destroyed at temperatures corresponding to the standard time and temperature required for pasteurization.

• Standard plate count: It is used to determine the bacteriological quality of pasteurized milk.

• Coliform count: The presence of coliform organisms in pasteurized milk indicates either improper pasteurization or post-pasteurization contamination.

• Methylene blue reduction test is an indirect method for the detection of microorganisms in milk. This test is carried out on the milk accepted for pasteurization. Definite quantities of methylene blue are added to 10 ml of milk at a uniform temperature of 37°C. The milk is considered satisfactory if it fails to reduce the methylene blue dye in 30 minutes under standard conditions.

Solution to Question 20:

An increase in the incidence of non-communicable diseases over a few years is described as a secular trend.

The secular trend describes the occurrence of disease over a prolonged period, usually years. Even though it may have short-term fluctuations, there is a tendency of the disease occurrence to change consistently in a particular direction or a definite movement in one direction.

Examples of the secular trend include coronary heart disease, lung cancer, and diabetes, which have shown a consistent upward trend in incidence as well as prevalence, in developed countries. Similarly, there has been a consistent decline in the diseases such as tuberculosis, typhoid fever, diphtheria, and polio.

Other options:

Option A: Seasonal trends reflect seasonal changes in disease occurrence following changes in environmental conditions that enhance the ability of the agent to replicate or be transmitted. For example, measles (usually occurs in early spring), varicella, etc. Non-infectious conditions like hay fever and snake bites may also show seasonal variations.

Option B: Cyclic trend is shown by some diseases that occur in cycles spread over short periods of time which may be, days, weeks, months, or years. Examples include the influenza pandemic, which is known to occur at intervals of every 7–10 years due to antigenic variations.

Option C: Periodic fluctuations include both seasonal and cyclic trends.

Solution to Question 21:

Among the given options, follow-up is not a part of case-control studies.

Case-Control Studies are also called retrospective studies. This study proceeds from effect to cause and is done after the exposure and outcome have occurred. They focus on diseases or health problems that have already developed. They are less time-consuming and have no risk of attrition because they do not require follow-up.

There are four basic steps in conducting a case-control study:

• Selection of cases and controls

• Matching- This is done to ensure comparability between cases and controls. Controls are selected such that they are similar to cases with respect to certain pertinent selected variables that influence the disease outcome.

• Measurement of exposure - This is obtained by interviews, questionnaires, or study of the past records of cases.

• Analysis in the form of exposure rates, risk ratio, and odds ratio.

Odds Ratio (OR) calculated from a case-control study, is a measure of the strength of the association between risk factor and outcome.

Note: Follow-up is a part of cohort studies. The elements in cohort studies include selecting study subjects, obtaining data on exposure, selecting comparison groups, follow up and analysis.

Solution to Question 22:

Cohort study is the best study design to demonstrate the relationship between cause and effect.

A cohort study is a type of analytical study, which progresses from cause to effect, as exposure has occurred but not disease. It is also known as forward-looking study, incidence study, prospective study, and longitudinal study.

A well-designed cohort study is considered the most reliable means of showing an association between a cause or suspected risk factor and subsequent disease (or effect) because it eliminates many of the problems of the case-control study.

Relative risk (RR) calculated using cohort study demonstrates the relationship between cause and effect. It is a direct measure of the strength of association between suspected cause and effect. The larger the RR, the greater the strength of the association between the suspected factor and disease.

The advantages of cohort study are the following:

• The incidence can be calculated.

• Several possible outcomes related to exposure can be studied simultaneously.

• Provides a direct estimate of relative risk.

• Dose-response ratios can be calculated.

• Bias like misclassification can be minimized.

Other options:

Option A: In a case-control study, the study begins from the outcome and goes retrospectively to find out the exposure among the cases and controls.

Option C: A cross-sectional study is a prevalence study which is more useful for the study of chronic diseases. It is more appropriate for the study of the distribution of a disease in a population than studying its etiology.

Option D: Clinical trials are research studies performed on people that are aimed at evaluating new tests and treatments and their effects on human health outcomes.

Solution to Question 23:

Hepatitis B subunit vaccines are composed of the HBsAg (Hepatitis B virus surface antigen).

Subunit vaccines are produced from macromolecules derived from immunogenic components of pathogens by recombinant DNA technology. They differ from inactivated whole-cell vaccines, by containing only the antigenic parts of the pathogen. These parts are necessary to elicit a protective immune response.

Hepatitis B vaccine is made by inserting the HbsAg gene into the yeast cell. This modified yeast cell produces large amounts of hepatitis B surface antigen, which is purified and harvested and used to produce the vaccine. The recombinant hepatitis B vaccine is identical to the natural hepatitis B surface antigen, but does not contain virus DNA and is unable to produce infection.

Solution to Question 24:

According to verbal autopsy report 2010-2013 data, the correct sequence of causes of neonatal mortality from maximum to least is prematurity and low birth weight> birth injuries and birth asphyxia> sepsis> congenital anomalies. (1>2>4>3)

Neonatal deaths are deaths that occur during the neonatal period, that is between birth and 28 completed days after birth. It is calculated by using the following formula:

Neonatal mortality rate = Number of deaths of children under 28 days of age in a year x 1000/Total live births in the same year

Important causes of neonatal mortality include (from maximum to least)

• Prematurity & low birth weight

• Birth asphyxia & birth trauma

• Neonatal pneumonia

• Other non-communicable diseases

• Sepsis

• Congenital anomalies

• Diarrhoeal diseases

• Injuries

• Tetanus

Note: Recent 2020 SRS (Sample Registration System) report shows the following:

• Neonatal mortality rate to be 20.

• Infant mortality rate to be 28.

• Under 5 mortality rate to be 32.

Solution to Question 25:

All the above steps are involved in meta-analysis except for randomization.

Meta-analysis is a set of statistical techniques for combining quantitative results from multiple studies to produce a summary of empirical knowledge on a given topic. It refers to a method of a systematic review.

Four basic steps involved in a meta-analysis are:

• Identification: This is the first step in a meta-analysis and it includes finding all of the articles or studies related to the topic.

• Selection: After assembling all the related studies, it is important to select the right or necessary studies by using inclusion (also called eligibility) criteria. Examples are minimum sample size (as very small studies may be unrepresentative), study design (controlled trials only vs randomized controlled trials only), etc.

• Abstraction: Once an appropriate group of studies has been identified, the author has to abstract the relevant data from each study.

• Analysis: This is the last step of meta-analysis.

Objectives of meta-analysis include,

• Summarize results from individual studies.

• Analyze differences in results among studies.

• Overcome small sample sizes of individual studies to detect the effects of interest.

• Determine if new studies are needed to investigate an issue.

• Generate new hypotheses for future studies.

Note- Randomization involves randomly assigning patients to different groups, like experimental and control groups. Clinical trials usually involve randomization and are hence called randomized clinical trials (RCTs). It is done to prevent confounding.

Solution to Question 26:

In the given scenario, the new drug was found to decrease mortality but not cure the disease so the duration of the disease is increased thereby increasing the prevalence. There is no change in

the incidence of the disease.

Incidence refers to the number of new cases in a defined population in a specified period of time.

Prevalence refers to the number of old and new cases at a given point of time or over a period of time in a given population.

Prevalence depends on two factors, the incidence, and duration of illness. If the population is stable and incidence and duration are unchanging, the relationship between incidence and prevalence can be expressed as:

P = I X D

(I - Incidence and D - Mean duration)

This equation shows longer the duration of the disease, the greater the prevalence.

Note:

• In chronic diseases like TB, prevalence will be higher than incidence.

• If a new treatment for a disease is effective, incidence does not change but prevalence decreases as the duration of disease decreases.

• If the treatment only prevents death but doesn't cure the disease, it may cause a paradoxical increase in prevalence.

Community Medicine NEET 2021

Question 1:

Which of the following thermometers is used to measure the low air velocity rather than the cooling power of the air?

a) Kata thermometer

b) Globe thermometer

c) Wet globe thermometer

d) Dial thermometer

Question 2:

There is an outbreak of buboes in the community. What is the vector responsible for this condition?

a) Xenopsylla cheopis

b) Phlebotomus argentipes

c) Ixodes tick

d) Female Anopheles mosquito

Question 3:

The Pathanamthitta district of Kerala was affected by floods and the government distributed doxycycline tablets for prophylaxis. Which other chemical will be distributed along with it?

a) Zinc phosphide

b) Malathion

c) Lindane

d) Paris green

Question 4:

Following admission of a road traffic accident case, there is spillage of blood on the hospital floor. Which disinfectant will you use to clean the floor?

a) Ethyl alcohol

b) Chlorhexidine

c) Sodium hypochlorite

d) Formaldehyde

Question 5:

The air pollution index chart of 4 consecutive days in Delhi station is given below. The air quality index of November 23 is classified as?

a) Moderately polluted

b) Poor

c) Very poor

d) Severe

Question 6:

There is an outbreak of acute encephalitis in the community and a vaccination drive is launched. Which of the following is true about the vaccine given in this condition?

a) Live and subcutaneous

b) Killed and intramuscular

c) Live and intramuscular

d) Killed and subcutaneous

Question 7:

The evaluation based on the treatment given to the patient and their clinical management in a health care facility measures which of the following?

a) Outcome

b) Process

c) Structure

d) Input

Question 8:

A girl child has had recurrent yeast infections and respiratory virus infections since she was 3 months old. Which of the following vaccine is contraindicated considering her immune status?

a) Killed IPV

b) Measles/MMR

c) DPT

d) TT/Td

Question 9:

Which of the following is a conditioning influence for malnutrition?

a) Child rearing

b) Infectious diseases

c) Food habits

d) Socioeconomic factors

Question 10:

Which of the following is not a component of the global hunger index (GHI)?

a) Infant mortality rate

b) Under 5 mortality

c) Percentage of undernourished population

d) Percentage of undernourished children under 5 years

Question 11:

A researcher wants to know whether there is an association of CRP values with the risk of myocardial infarction/stroke. The study group was divided into quintiles and the following table is formed based on the study results. Which of the following statements is true regarding the relationship between CRP and the risk of myocardial infarction/stroke?

a) CRP has no association with risk of MI/stroke

b) Increase in CRP increases the risk of MI/stroke

c) Increase in CRP decreases the risk of MI/stroke

d) In quintile 1, there is no risk of MI/stroke

Question 12:

A child presented with pharyngitis. A throat swab was obtained and sent for culture. The used swab should be discarded in which color bin?

a) Red

b) Yellow

c) White

d) Blue

Question 13:

Which of the following is the SI unit for the measurement of brightness of light from a point source?

a) Candela

b) Lux

c) Lambert

d) Lumen

Question 14:

A pregnant woman whose niece lives with her in the same house contracted Varicella. She tested negative for serum antibodies against varicella. What would this mean?

a) She is immune to zoster

b) She is susceptible to zoster

c) She is susceptible to chicken pox

d) She is immune to chicken pox

Question 15:

The school health programs are managed by which of the following?

a) Primary health center

b) District hospital

c) Sub-centre

d) Sub-divisional hospital

Question 16:

A man from Chattisgarh presented with progressive muscle weakness and leg spasms. Pure motor paresis was seen on examination. What is the most appropriate history to be elicited in this patient?

a) History of similar illness in the past

b) History of fever

c) History of vaccination

d) History of diet

Question 17:

In a 10-year-old school child, which of the following vaccines is given as a part of the immunization programme?

a) BCG

b) MMR

c) TT/Td

d) DPT

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	c
5	d
6	b
7	b
8	b
9	b

10	a
11	b
12	b
13	a
14	c
15	a
16	d
17	c

Detailed Explanations

Solution to Question 1:

Kata thermometer is used to measure the low air velocity rather than the cooling power of the air.

It was originally used to measure the cooling power of the air (air temperature, humidity, and air movement) but currently, it is used only as an anemometer to measure low air velocities.

A wet kata reading of 20 and above and a dry kata reading of 6 and above were taken as indices of thermal comfort.

Kata thermometer is shown in the image below.

Other options:

Option B: Globe thermometer is used for the direct measurement of mean radiant temperature of the surroundings.

Options C and D: Wet globe thermometer consists of a dial thermometer. It provides a measure of the cooling capacity of the working environment.

Solution to Question 2:

The vector for bubonic plague is Xenopsylla cheopis. It is also called a rat flea. It also transmits murine typhus, chiggerosis, and hymenolepsis diminuta. It can be identified by the presence of a conical-shaped head, the presence of thorax, abdomen, and three pairs of legs.

Plague is a zoonotic disease caused by Yersinia pestis. It is a gram-negative, non-motile, coccobacillus that exhibits bipolar staining with special stains like Wayson's stain. The reservoir of infections is wild rodents like field mice, gerbils, skunks, etc. In India, Tatera indica is the main reservoir. The important vector of plague is rat flea, also called Xenopsylla cheopis.

Humans acquire the disease by the bite of an infected flea, by direct contact with the tissues of the infected animal, or by droplet infection from cases of pneumonic plague.

Human plague mainly presents as:

Bubonic plague- This is the most common type of disease. The patient develops a sudden fever, chills, headache, and painful lymphadenitis. Within a few days, the lymph nodes in the groin, axilla, or neck enlarge and are called buboes. This is followed by the suppuration of the buboes. Bubonic plague cannot spread from person to person as the bacilli are locked up in the buboes and do not find a way or easy exit.

Pneumonic plague- It develops as a complication of bubonic-septicemic plague.

Septicemic plague- It occurs when the infection spreads to blood.

Other options

Option B- Species Phlebotomus argentipes, belongs to the group of sand fly. It transmits kala azar. Diseases caused by other species of sand flies are sandfly fever and oriental sore.

Below image shows sand fly

Option C- Ixodes ticks (hard ticks) transmit rocky mountain spotted fever, Russian spring-summer encephalitis, Colorado tick fever, tularemia, tick paralysis and human babesiosis.

Below image shows Ixodes tick

Option D- Female Anopheles mosquitoes are vectors for malaria and filariasis (not in India). They can be identified by the presence of spotted wings. When resting on a surface, they make an angle with the surface.

The below image shows anopheles mosquito.



Solution to Question 3:

Leptospirosis, followed by acute diarrhoeal diseases and dengue are most commonly prevalent following floods in Kerala. Malathion will be the choice of insecticide used in this scenario.

Leptospirosis is a bacterial zoonotic disease transmitted via water contaminated with rodent urine. Infected rodents excrete large amounts of leptospira in their urine and floods facilitate the

spread of the organism.

Prevention of leptospirosis:

• Chemoprophylaxis with Doxycycline 200 mg once a week for 6 weeks is recommended for high-risk populations in endemic areas. In case of fever, Doxycycline 100 mg twice a day for seven days is the recommended dose.

• Rubber shoes and gloves should be used.

• Anti-rodent measures must be taken.

• Mapping of water bodies must be done to establish a proper drainage system.

Other diseases usually transmitted after floods are typhoid fever, malaria, cholera, hepatitis A & E.

General measures to control disease spread during floods are:

• Chlorine tablets are distributed to disinfect the household water supply. One tablet can chlorinate 20 liters of water. Drinking water disinfectants like bleaching powder are also used.

• Malathion is an organophosphorus synthetic insecticide used for vector control (adult mosquitoes). It is a contact poison that is sprayed using ultra-light volume fogging machines.

• Indoor residual sprays with insecticides and larvicides like abate for drinking water are recommended.

• Tetanus booster dose for previously vaccinated people with open injuries may be indicated.

• General hygiene and sanitary measures must be followed.

Other health risks posed by floods are hypothermia, respiratory tract infections, snake bites, and injuries/drowning.

Other options:

Option A: Zinc phosphide is a rodenticide used to kill animals like mice and rats. It is a highly toxic compound and can produce toxicity by oral ingestion, inhalation, and through damaged skin.

Option C: Lindane is used as an insecticide for hardwood logs, lumber, and crops. It also has topical pediculicidal and scabicidal activities and is used in pet treatment.

Option D: Paris green is an insecticide that is not much used due to the availability of lesser phytotoxic agents.

Solution to Question 4:

The disinfectant used in blood and body fluid spillage management is sodium hypochlorite.

Procedure to be followed in the event of blood/body fluid spillage :

• Wear a pair of non-sterile gloves

• Broken glass (if any) of the container is cleaned first using tongs or a pan and brush and discarded in the sharps container. Caution is taken to not use hands.

• Disposable paper towels are used to absorb as much as possible.

• Water and detergent are used to clean the surface until visibly clean.

• The area is saturated with 0.5% Sodium Hypochlorite. This is prepared by diluting 5.25% sodium hypochlorite bleach at a 1:10 ratio.

• The tongs, pan, and brush are washed under running water and left to dry

• The gloves are removed and discarded.

• Hands are washed with soap and water using proper technique and dried with a disposable towel.

• The spillage is recorded in an incident book.

Solution to Question 5:

According to the data presented in the question, the air quality index for the 23rd of November is 407 which is classified as severe.

Air Quality Index (AQI) -

A tool used for effective communication of air quality status on a scale that is easily comprehensible. Complex air quality data of various pollutants are compiled to give a single score (index value), nomenclature and colour that determines the air pollution of that particular location.

Eight pollutants are taken into consideration to calculate the AQI -

• PM10 (Airborne particulate matter with a diameter of 10 microns or less)

• PM2.5 (Airborne particulate matter with a diameter of 2.5 microns or less)

• NO_2 (Nitrogen dioxide)

• SO_2 (Sulfur dioxide)

• CO (Carbon monoxide)

• O_3 (Ozone/Trioxygen)

• NH_3 (Ammonia)

• Pb (Lead)

AQI categories and the range are as follows :

AQI Category	AQI Range
Good	0 - 50
Satisfactory	51 - 100
Moderately polluted	101 - 200

AQI Category	AQI Range
Poor	201 - 300
Very poor	301 - 400
Severe	401 - 500

Solution to Question 6:

The given scenario is suggestive of Japanese encephalitis for which a killed vaccine is administered intramuscularly.

The Japanese encephalitis (JE) virus is transmitted by the bite of Culex mosquitoes, specifically Culex tritaeniorhynchus and Culex vishnui (common in India) species. They present with fever and headache in mild disease form. Still, severe disease presents with rapid onset of high fever, headache, neck stiffness, disorientation, coma, seizures, spastic paralysis, and ultimately death.

Killed JE vaccines (Inactivated SA 14-14-2 cell culture-derived and inactivated Vero cell culture-derived) are gradually replacing live attenuated vaccines. This change is because the killed vaccines are reported to have better immunogenicity than the live attenuated vaccine.

The SA 14-14-2 (live attenuated) vaccine is based on a stable neuro-attenuated strain of the JE virus with a Yellow Fever vaccine acting as a viral vector. Adverse effects of the vaccine include fever and rash.

Dose: 0.5 ml

Site: Live vaccine: Subcutaneous, left upper arm

Killed vaccine: Intramuscular, anterolateral side of left mid-thigh

They can be administered up to 15 years of age with a month gap between the two doses.

The JE virus vaccine has been included in the National Immunization Schedule (NIS) to be administered in endemic areas (Uttar Pradesh, Assam, West Bengal, Karnataka) in two doses. The first dose is administered at 9-12 completed months. The second dose is administered at 16-24 months. The schedule however mentions the use of both live and killed vaccines.

Note: According to the national immunisation schedule, both live attenuated, killed-inactivated vaccines are now used for vaccination against the JE virus.

Solution to Question 7:

The evaluation based on the treatment given to the patient and their clinical management in a health care facility measures the "process" of medical care.

The basic steps of evaluation are as follows :

• Determine what is to be evaluated

• Establish standards and criteria

• Plan the methodology to be applied

• Gather information

• Analyze the results

• Take action

• Re-evaluate

The three types of evaluation are:

• Evaluation of "structure": This is an evaluation of whether facilities, equipment, manpower, and organization meet a standard approved by experts.

• Evaluation of "process": The process of medical care includes the problems of recognition, diagnostic procedures, treatment and clinical management, care, and prevention. They are evaluated by comparing with a predetermined standard. Eg; Evaluation of the physician (or nurse) performance is known as "Medical (or nursing) Audit".

• Evaluation of "outcome": This is concerned with the benefits gained by people using the health services. The traditional outcome components are the "5 Ds" of ill-health, i.e; disease, discomfort, dissatisfaction, disability, and death.

Elements of evaluation are relevance, adequacy, accessibility, acceptability, effectiveness, efficiency and impact.

Solution to Question 8:

The clinical scenario showing an immunocompromised child should not receive the Measles/MMR vaccine from the given options as it is a live attenuated vaccine which is contraindicated in immunocompromised people.

Contraindications for live vaccines:

• Immune deficiency diseases

• Suppressed immune responses as a result of

• Leukaemia

• Lymphoma

• Malignancies

• Corticosteroid therapy

• Treatment with alkylating agents and antimetabolic agents

• Therapeutic radiation

• Pregnancy

Measles vaccine:

Live attenuated vaccine for measles is safe and effective with no evidence of person to person transfer of vaccine strains. It is freeze-dried product requiring reconstitution. 0.5ml is administered subcutaneously in the right upper arm as two doses at the 9th month and between the 16th-24th month. They also contain a small amount of neomycin.

Other options :

Option A - Killed IPV is a form of killed/inactivated vaccine where the absolute contraindication is a severe local or general reaction to a previous dose.

Options C and D - DPT and TT/Td fall under toxoids/protein vaccines that are usually highly efficacious and safe immunizing agents.

Solution to Question 9:

From the given options, infectious diseases are the conditioning influence for malnutrition.

Ecology of Malnutrition :

The ecological factors related to malnutrition as follows

• Conditioning influences: A conditioning factor that plays a huge role in malnutrition is infectious diseases which causes a vicious cycle where infections cause malnutrition which in turn causes minor childhood ailments. Diarrhoeal diseases, intestinal parasites, measles, whooping cough, malaria, tuberculosis contributes significantly to malnutrition. The occurrence of infections are largely based on the environmental conditions that the child is exposed to.

• Cultural influences: Poor diets as a result of cultural and regional influences also play a role in malnutrition. They are as follows with an example for each:

• Food habits, traditions, customs, beliefs and attitudes: Rice and wheat are largely staple in India. Papaya is avoided as it is thought to cause abortion.

• Food fads: Personal likes and dislikes to a certain group of foods.

• Cooking practices: Prolonged boiling of cereals in open pans which leads to a loss of nutritive value

• Child-rearing practices: Premature curtailment of breastfeeding

• Socio-economic factors: Poverty plays a huge role in causing malnutrition. This is usually a result of ignorance, insufficient education, lack of knowledge regarding the nutritive value of foods, inadequate sanitary environment, large family size, etc. which affects the quality of life.

• Food production: The amount of food produced is directly proportional to the amount of food consumed. Developed countries have 5.8 hectares of arable land compared to the 0.6 hectares of land per Indian. Even distribution of food among all sectors of society is also a factor.

• Health and other services: Remedial actions that can be taken by the health sector to reduce malnutrition are as follows -

• Nutritional surveillance - Continuous monitoring of factors that affect nutrition; groups affected are identified.

• Nutritional rehabilitation - Urgent care for individuals suffering from severe malnutrition.

• Nutrition supplementation - Supplementary feeds for mothers and children

• Health education - Health education programmes to increase awareness about malnutrition and nutrition value of food.

Solution to Question 10:

The infant mortality rate is not a component of the global hunger index.

The Global Hunger Index (GHI) has three dimensions. These dimensions are measured by indicators.

GHI score = (1/3 X standardized PUN) + (1/6 X standardised CWA) + (1/6 X standardised CST) + (1/3 X Standardised CM)

Where,

• PUN is the proportion of the population that is undernourished

• CWA is the prevalence of wasting in children under 5 years of age

• CST is the prevalence of stunting in children under 5 years of age

• CM is the proportion of children dying under the age of 5 years of age.

The result lies on a scale from 0 to 100, with 0 meaning no hunger and 100 being the worst score. The GHI is given by the International Food Policy Research Institute. In 2021, India's GHI was 27.5. It ranks 101 among 116 countries. In the 2022,India's GHI stands at 29.1, and India ranks 107th out of the 121 countries.

Dimensions	Indicators
Inadequate food supp ly	Proportion of the population undernourished
Child mortality	Under 5 mortality rate
Child undernutrition	Wasting in children under 5 yearsStunting in children un der 5 years

Solution to Question 11:

From the study results, option B is the true statement i.e; an increase in the CRP increases the risk of myocardial infarction/stroke.

Relative risk (RR)/risk ratio is the ratio of the incidence of the disease (or death) among exposed and the incidence among non-exposed.

$$RR = \frac{\text{Incidence of disease (or death) among exposed}}{\text{Incidence of disease (or death) among non-exposed}}$$

It is a direct measure of the strength of the association between suspected cause and effect. The larger the RR, the greater the strength of the association between the suspected factor and disease. As there is an increase in the RR with increasing CRP levels, the risk of an MI/stroke also increases with elevated CRP levels.

RR = 1 indicates no association.

RR >1 indicates a positive association.

RR of 2 or higher indicates a >/=100% increase in risk.

Relative risk calculation (RR) = [a/(a+b)]/ c/(c+d)] where,

In the question, the table shows an increase in relative risk with an increase in CRP values. This thereby indicates that an increase in the CRP values increases the risk of myocardial infarction/stroke.

Note: Relative risk provides information on the increase or decrease in the likelihood of a condition based on exposure. It does not provide the absolute risk of the event occurring. The last option states that "In quintile 1 there is no risk of MI/stroke". Even though the relative risk is 1, it only indicates no association but does not exclude the occurrence of MI/ Stroke.

	Disease		
Exposure		yes	no
yes	a	b	
no	c	d	

Solution to Question 12:

The used throat swab should be discarded in the yellow color bin as it is a clinical laboratory waste.

Microbiology, biotechnology, and clinical laboratory waste are collected in autoclave-safe bags. They must be pre-treated before disposal by incineration, plasma pyrolysis, or deep burial.

Solution to Question 13:

The SI unit for the measurement of brightness of light from a point source is Candela.

The "power" of light from a given point source which radiates in all directions is also known as it's luminous intensity of the point of source. This is measured in candela or candle light.

The following table shows different units used for the measurement of light as recommended by the International Organization for Standardization:

Quantity Measured	Description	SI Unit	Other Units Used
Luminous intensity	Brightness of a point source	Candela	Candle power
Luminous flux	Flow of light	Lumen	-
Illumination(Illuminance)	Amount of light reaching a s urface	Lux (per unit area)	Foot candle, phot, lm/cm ^{2}
Luminance	Amount of light reflected fro m a surface	Lambert	Foot lambert, candle/cm 2

Solution to Question 14:

The given clinical scenario showing the absence of serum antibodies is indicative of susceptibility to chickenpox.

Varicella zoster virus causes two host responses which present as chickenpox and herpes zoster. They are characterized by a vesicular rash that is accompanied by fever and malaise (chickenpox). The immune system eliminates the infection except for some virions that establish latent infections inside nerve cells which are reactivated when immunity is compromised and present as herpes zoster.

One attack with the virus gives durable immunity while a second attack of the virus is rare. In the absence of immunity, all ages are equally susceptible to the disease. The acquisition of maternal antibodies protects the infant during the first few months of life. The IgG antibodies produced during the first attack with the virus persist for life and their presence is correlated with protection against varicella. The absence of antibodies in the given scenario signifies that the patient is susceptible to chicken pox. Hence, the serologic screening of serum for lgG antibodies is used to assess immunity or susceptibility of a person to varicella, especially in unvaccinated people, e.g. in healthcare workers.

The presence of cell-mediated immunity appears to be important in the recovery from VZ infections and in protection against the reactivation of latent VZ virus which causes Herpes zoster.

Varicella in pregnant women :

The contraction of the virus by a pregnant woman increases the risk of the fetus to congenital varicella syndrome. This is seen in mothers who are affected during the first 20 weeks of gestation. Affected babies present with the following :

• Foetal wastage

• Cutaneous scars

• Atrophied limbs

• Microcephaly

• Low birth weight

• Cataract

• Microphthalmia

• Chorioretinitis

• Deafness

• Cerebra-cortical atrophy

Prevention:

• Varicella-Zoster lmmunoglobulin (VZIG) is given as post-exposure prophylaxis within 72 hours of exposure and has been recommended for prevention of chickenpox in exposed susceptible individuals particularly in immunosuppressed persons such as:

• Persons receiving immunosuppressive therapy

• Persons with congenital cellular immunodeficiency

• Persons with acquired immunodeficiency including HIV/AIDS

• Exposed pregnant women

• Newborns

• Premature infants of low birth weight

• Live attenuated vaccine is administered to children who have not had chicken pox between the age of 12 to 18 months.

Solution to Question 15:

The school health programs are managed by the primary health centre (PHC).

The School Health Committee (1961) in India recommended that the school health services should be a part of the general health services. Since the general health services are administered through PHCs, school health programs are also managed by the PHCs under their jurisdiction.

The School Health Committee advocated the formation of school health committees at the village level, block level, district level, state level, and national level. The National School Health Council will be an advisory and coordinating body.

PHCs provide the following school health services:

• Screening of general health assessment of anemia/nutritional status, visual acuity, hearing problem, dental check-up, physical disabilities, learning disorders, and behavior problems, etc.

• Basic medicines for common ailments

• Immunization

• Micronutrient (Vitamin A, iron, and folic acid) management

• Deworming

• Mid-day meal

Note: National Programme for Mid-Day Meal in Schools has recently been renamed Pradhan Mantri Poshan Shakti Nirman (PM POSHAN).

Solution to Question 16:

The given clinical features are suggestive of neurolathyrism and excessive consumption of the pulse Lathyrus sativus is the main cause. Hence, a diet history should be elicited in this patient to confirm the diagnosis.

Lathyrism is a paralyzing disease of humans and animals. It is due to the consumption of Khesari dal/Lathyrus sativus. When the husk is removed, it looks similar to red gram dal or Bengal gram dal. It is more prevalent in parts of Madhya Pradesh, Uttar Pradesh, Odisha, and Bihar.

In humans, it is referred to as neurolathyrism because it affects the nervous system. It is characterized by gradually developing spastic paralysis of the lower limbs.

In animals, it is referred to as osteolathyrism/odoratism because the pathological changes in the bones lead to skeletal deformities.

It usually affects young males aged 15-45 years and it is manifested in stages:

• Latent stage

• No-stick stage

• One-stick stage

• Two-stick stage

• Crawler stage

BOAA (Beta oxalyl amino alanine) is the toxic compound causing Lathyrism. The water-soluble property of the toxin is used in removing it from the pulses by soaking it in hot water and rejecting the soaked water.

Parboiling can be used for detoxifying the crop on large-scale operations. Overnight soaking in lime water followed by boiling is also effective in destroying the toxin.

Vitamin C in doses of 500-1000mg can be used for the prevention and treatment of lathyrism.

The Prevention of Food Adulteration Act in India has banned Lathyrus sativus in all forms (whole/split/flour).

Solution to Question 17:

In the given scenario, a ten-year-old schoolchild will be administered the TT/Td vaccine according to the National Immunization Schedule.

The Tetanus and adult diphtheria (Td) vaccine is a combination of tetanus and diphtheria with a lower concentration of diphtheria antigen (d) as recommended for older children and adults. The MoHFW recommends the replacement of TT vaccine with Td vaccine for all age groups, including pregnant women. A dose of 0.5ml is administered in the intramuscular route in the upper arm.

Scheduled to be administered at 10 and 16 years of age.

For pregnant women :

• Td-1: early in pregnancy

• Td-2: 4 weeks after Td-1

• Td (Booster): if pregnancy occurs within 3 years of last pregnancy and 2 Td doses were received.

Other options:

Option A - According to the National Immunization Schedule, BCG should be given at birth or as early as possible up to 1 year of age. It provides protection against CNS and disseminated TB. A dose of 0.1 ml is administered intradermally in the left upper arm.

Option B - Measles/MR is given at 9 completed months (1st dose) and a second dose is given at 16-24 months. Measles can be given till 5 years of age. A dose of 0.5ml is administered subcutaneously in the right upper arm.

Option D - DPT vaccine is administered as a part of the pentavalent vaccine at the 6th, 10th and 14th weeks of life. DPT booster doses are administered at 16-24 months (first booster) and 5-6 years (second booster). It can be administered until 7 years of age (with booster doses). A dose of 0.5ml is administered in the anterolateral aspect of the mid-thigh. The 2nd booster dose can be administered in the upper arm.

Community Medicine INI-CET 2022

Question 1:

As per the National Tuberculosis Elimination Program (NTEP), HIV TB prevalence of _____ would deem a district as a high-priority district?

a) >10%

b) >12%

c) >15%

d) >20%

Question 2:

India is a country with different cultures and diverse languages. Which of the following steps should a physician take to address the patient for better outcomes?

a) 1, 2

b) 2, 3

c) 3, 4

d) 1, 4

Question 3:

5-year-old Ram and 3-year-old Sham are Rama Devi's sons. Sham developed a rash, but Rama Devi ignored it, assuming it to be an allergy. After 3 days, Ram also developed a rash along with a fever. Out of fear, Rama Devi took Ram to the primary health center (PHC). The healthcare workers at the PHC suspected it to be measles. Choose all correct options from among the following:

a) 1 and 4 are correct

b) 1 only is correct

c) 1, 2, 3 and 4 are correct

d) 1 and 3 are correct

Question 4:

You are measuring a patient's blood pressure at a primary health center when the sphygmomanometer fell and broke and mercury was spilled onto the ground. How will you clean the mercury?

a) Collect it by using a card onto a sheet of paper and put the mercury into a jar

b) Collect it by sweeping it with a broom and throw it outside

c) Collect it by sweeping it and flush it in the toilet

d) Collect it by hand and then wash hands with water

Question 5:

The true statement about screening tests for genetic diseases is:

a) It is always invasive

b) Screening test has better accuracy than diagnostic test

c) It defines risk of transmission of disease to the child

d) Screening requires genetic mapping

Question 6:

Which of the following is the true statement regarding measures to prevent typhoid transmission in the community?

a) Typhoid vaccine administration is the best method of preventing transmission

b) Hygiene practices and clean sanitation control is more important than typhoid vaccine

c) In 10% cases, person to person transmission occurs

d) Drug resistance in typhoid is not as big a problem as in TB

Question 7:

Which of the following is true about the management of a baby born to an HIV-positive mother?

a) If replacement feeding, the neonate should be tested for HIV, if negative, ART is not required.

b) In exclusive breastfeeding, give nevirapine for 6 months only when baby is positive

c) If replacement feeding, the neonate need not receive nevirapine if HIV negative

d) If exclusive replacement feeding, give nevirapine for 6 weeks, even if the baby is negative

Question 8:

What is the branch of statistics dealing with describing the internal collection of data?

a) Theoretical statistics

b) Descriptive statistics

c) Analytical statistics

d) Inferential statistics

Question 9:

Isolation in a separate room is mandatory for which of the following air-borne infections?

a) Measles

b) Rabies

c) Nipah

d) Aseptic meningitis

Question 10:

All the following cause widening of confidence interval except?

a) Increase in sample size

b) Decrease in sample size

c) Increase in variability

d) Increase in confidence level

Question 11:

False statement regarding rabies immunoglobin is:

a) It is exclusively given in the deltoid

b) The dose is 20 IU/kg

c) Not given after 7 days of the 1st dose of the vaccine

d) If human RIG is unavailable, equine is preferred

Question 12:

A 1-year-old child is brought to the healthcare unit that you are working on. His mother mentions that he had received the first dose of DPT vaccine at 6 weeks of age but has missed vaccination since then. What would you advise the patient?

a) Do not give DPT

b) Restart DPT

c) Give second dose of DPT

d) Give dT alone

Question 13:

The following image shows a health belief model. What does box C stand for?



a) Perceived control

b) Behavioral modification

c) Health motivation

d) Self-efficacy

Question 14:

A point-of-care diagnostic test with a sensitivity of 98% shows a negative test result in a patient. What does this imply?

a) The patient would need another point-of-care diagnostic test for the same clinical condition.

b) Prevalance of the disease is 2%.

c) Patient is less likely to have the disease.

d) Patient has been wrongly ruled out due to false negativity.

Question 15:

The table given below shows a multivariate analysis of the relation between the risk of developing blindness and age, literacy rate, residence, and gender. Which of the following is true?

a) 2 and 3

b) 1 and 3

c) 1 and 2

d) 3 and 4

Question 16:

A patient with lymphatic filarisis is being managed as shown in the image below. What is the level of prevention done here?

a) Specific protection

b) Disability limitation

c) Primary prevention

d) Health promotion

Question 17:

Which of the following indicators can help in determining whether the health system is giving importance to identifying leprosy in the community?

a) Treatment completion rate

b) Annual new case detection rate per lac

c) Treatment initiation rate

d) Proportion of newly diagnosed patients with grade-2 disability

Question 18:

Which of the following is true regarding the vaccine vial monitor (VVM)?

a) 3,4

b) 1,2

c) 1, 2, 3, 4, 5

d) 5,6

Question 19:

Which of the following models of disease causation is depicted in the image below?



a) Sufficient component cause model

b) Epidemiological triad

c) PERT model

d) Web of causation model

Question 20:

Attributable risk calculated to understand the relation between smoking and lung cancer is 90%. Which of the following is true?

a) 10% cases of lung cancer can be eliminated in smokers if smoking is eliminated

b) 90% cases of lung cancer can be eliminated in population if smoking is eliminated

c) 90% cases of lung cancer can be eliminated in smokers if smoking is eliminated

d) 10% cases of lung cancer can be eliminated in population if smoking is eliminated

Question 21:

Following an earthquake, the building codes were changed to earthquake-resistant buildings. Which of the following step of disaster management does this fall under?

a) Disaster response

b) Disaster reconstruction

c) Disaster rehabilitation

d) Disaster mitigation

Question 22:

Which of the following statements are true regarding drugs used in anti-tubercular treatment?

a) A, B, C, D

b) B, C

c) A, C

d) A, D

Question 23:

Which of the following criteria should be satisfied to start MDT?

a) 1,4

b) 1,3

c) 1,3,4

d) 1,2,3,4

Question 24:

Match the following stages with their relevant descriptions.

a) 1-c, 2-d, 3-a, 4-d

b) 1-d, 2-b, 3-a, 4-c

c) 1-b, 2-d, 3-c, 4-d

d) 1-a, 2- c, 3-d, 4-b

Question 25:

Contraceptive failure is assessed using which of the following methods?

a) Pharmacological index

b) Pearl index

c) Performance index

d) Efficacy index

Question 26:

Match the following scientists with their discoveries:

a) 1-a, 2-b, 3-a

b) 1-b, 2-a, 3-c

c) 1-c, 2-b, 3-a

d) 1-b, 2-c, 3-a

Question 27:

Identify the formula given below:



a) Standard deviation

b) Standard error

c) Median

d) Variance

Question 28:

A child presents with a dog bite from a street dog. On examination, an abrasion of 1.5 cm is present, with slight oozing. Which of the following statements are correct?

a) 2, 3, 5, 6

b) 1, 2, 5, 6

c) 1, 2, 3, 5

d) 1, 2, 3, 6

Question 29:

Water from a broken septic tank is collected in a container and breeding mosquitoes are seen. The image of the larva is given below. Identify the species of the mosquito.



a) Anopheles

b) Aedes

c) Mansonia

d) Culex

Question 30:

The drug shown in the image below is administered for:

a) Malaria

b) Dengue

c) Sexually transmitted infections

d) AIDS

Question 31:

Under the Anemia Mukt Bharat initiative, mild to moderate anemia in pregnant women < 34 weeks of gestation is treated using:

a) 3 iron and folic acid (IFA) tablets, taken OD

b) IV iron sucrose for non-compliance with oral tablets

c) 2 iron and folic acid tablets OD + IV iron sucrose

d) IM ferric carboxy maltose (FCM)

Answer Key

Question No.	Correct Option
1	a
2	d
3	c
4	a
5	c
6	b
7	d
8	b
9	a
10	a
11	a
12	c
13	c
14	c
15	d
16	b
17	d
18	b
19	a

20	c
21	d
22	d
23	b
24	b
25	b
26	d
27	a
28	d
29	d
30	a
31	b

Detailed Explanations

Solution to Question 1:

As per the National Tuberculosis Elimination Programme (NTEP), an HIV-TB prevalence of >10% would deem a district as a high-priority district.

RNTCP (Revised National Tuberculosis Control Programme) has been revised by the government of India as NTEP. The vision and goals of the NTEP programme respectively are

• Tuberculosis (TB) free India with zero deaths, diseases and poverty due to tuberculosis

• To achieve a rapid decline in the burden of TB, morbidity and mortality while working towards the elimination of TB in India by 2025.

NTEP has categorized and prioritized the districts in the country based on the following criteria to intensify tuberculosis control.

Categorization	Criteria	Category
High TB	Total TB case notification rate > 180 per lakh	A
High TB-HIV(Tuberculosis and HIV co-infected patients)	>10%proportion of known HIV positives amongst TB patients tested for HIV	B
High DR-TB (Drug-resistant tuberculosis)	> 25% proportion of relapse out of incident smear-positive TB cases	C
Very low case finding effort	Annual TB suspect examination rate of < 400 per lakh population	D

Categorization	Criteria	Category
Average	None of above	E
High case finding but Low TCNR	Annual suspect examination rate >1200 per lakh population and total case notification rate < 80 per lakh population	F

Solution to Question 2:

The physician should have good communication and consider the patient's religious, and cultural perceptions for better outcomes (statements 1 and 4 are correct).

Statement 1 is true since the proper usage of vocabulary that is easy for the patient to understand is essential for the patient. The doctor should clarify the patient's interpretation of the words used. Thus insisting on good communication is essential for better health outcomes.

Statement 2 is not true. An interpreter is needed if the patient is not a native speaker of the language spoken in that region. The interpreter can be a family member above the age of 16 or an independent interpreter. Communication gets affected with open-ended questioning getting lost in the double translation. The answer returned might not be adequate for the doctor and the consultation as such becomes longer than usual. The doctor will have to change to a more direct style of questioning to prevent any misunderstanding.

Statement 3 is not true while statement 4 is true. Personality, culture, and beliefs are important in creating trust between the doctor and patient and are an important component in the medical interview for a better outcome. The doctor must listen to the patients' concerns, and preferences.

Health communication may fail if the message is not adequate. A good message should have the following:

• It should be in line with the objective

• It needs to be meaningful

• Based on the felt needs

• Clear and understandable

• Specific and accurate

• Culturally and socially appropriate.

Solution to Question 3:

All 4 statements are correct from the given clinical scenario.

Statement 1 is correct: Ram is the index case as he was the first case to come to the attention of the investigating authority (PHC)

Statement 2 is correct: From the given clinical scenario, considering the family of Ramadevi as a unit, Sham is the primary case (the first person who gets the communicable disease and introduces it to the community).

Statement 3 is correct:

• Incubation period from the time of exposure to the development of rash is 7 to 18 days with an average of 14 days.

• Ram developed the rash 3 days after Sham. Considering the period of communicability of rash as 4 days prior to or 4 days later than the onset of rash, it is evident that Sham was in an infectious state and Ram has contracted the infection from Sham

Statement 4 is correct: Post-exposure prophylaxis (PEP) can be given in the form of a vaccine or immunoglobulin. The vaccine can be given within 72 hours of exposure, while immunoglobulin (IVIG) can be given up to 6 days following exposure. Hence, Ram could would have been eligible to receive PEP if he were brought to the PHC along with Sham.

Solution to Question 4:

Mercury is cleaned by collecting it by using a card onto a sheet of paper and putting the mercury into a jar.

After a mercury spill, the following steps should be followed:

• Isolation of the spill and ventilation of the area

• Collection of items to clean the spill: zipper-top plastic bags, trash bags, gloves, paper towels, cardboard or squeegee, eye dropper, tape, flashlight, and powdered sulfur.

• Clean up the spill.

• After putting on the gloves, the glass pieces are picked and placed on a towel, rolled and disposed of into a zipper-top plastic bag, and finally sealed.

• The beads of mercury are cleaned by using cardboard or a squeegee. They are rolled onto a sheet of paper, disposed into the zipper bag, and sealed.

• Powdered sulfur may be used to absorb small beads that are not visible

• Looking for any missed spill using a flashlight

• Remove other mercury contaminated items and wipe off any mercury beads with a wet towel into a plastic trash bag

• Proper disposal of contaminated clean-up materials.

• Keep the area well ventilated for 24 hours, keep pets and children away from the area

The use of a vacuum cleaner or broom to clean the spill should be avoided. This will break the mercury into smaller droplets and spread it further. Mercury should not be poured down any drain. Do not use a washing machine to clean the clothes contaminated with mercury. This can contaminate the water and sewage. Mercury-contaminated shoes or clothing should not be worn.

Solution to Question 5:

The true statement about screening tests for genetic diseases is it defines the risk of transmission of disease to the child.

A screening test is an initial examination done in apparently healthy individuals while diagnostic tests are done on individuals with positive test results for further workup and treatment.

The aim of performing a screening test is to favorably change the natural history of the disease by an early diagnosis and treatment in individuals who are positive for the particular screening test. They are less invasive and less accurate than diagnostic tests.

Genetic screening is a type of public health program done in high-risk asymptomatic individuals in which tests are used to detect an inherited predisposition to disease, risk of transmitting the disease, prevention of the disease, early treatment, or family planning. For example, prenatal screening during pregnancy.

Genetic mapping is a tool used to identify new genes and their functions. It also may give information on which chromosome contains the defective gene and where the gene is present on that chromosome. Gene maps have been used to pinpoint the gene responsible for rare single-gene inherited disorders. Screening does not require genetic mapping.

Solution to Question 6:

The true statement regarding typhoid transmission preventive measures is that hygiene practices and clean sanitation control are more important than the typhoid vaccine. It helps in breaking the chain of transmission.

Typhoid fever is caused by Salmonella typhi and is mainly transmitted to humans by the fecal-oral route.

Clinical features include high fever with chills associated with headache, malaise, cough, sore throat, abdominal pain, and constipation or diarrhea are seen in the early phase. The fever occurs in a step-ladder fashion reaching a plateau in 7-10 days. A maculopapular rash may be seen on the trunk and chest known as rose spots. In the late stages, hepatosplenomegaly, abdominal distension, relative bradycardia, and a dicrotic pulse can occur.

Diagnosis is by isolation of the organism from blood, bone marrow, and stools. Blood culture is mainly used for the diagnosis. Other tests such as the Widal tests are useful.

Management is mainly by sanitation. It is the ultimate method to control typhoid. It breaks the transmission of the disease. It is combined with proper health education.

Antibiotic therapy is started early and the choice of antibiotics depends on the susceptibility of the organism to the drug. In the case of drug-susceptible typhoid, ciprofloxacin is useful.

Drug resistance to typhoid has been emerging. Similar to drug resistance in TB, there are multidrug-resistant (MDR) and extensively drug-resistant (XDR) S. Typhi strains. MDR strains are resistant against at least 1 out of 3 or >3 categorically differentiated antimicrobials, such as ampicillin, cotrimoxazole, and chloramphenicol. XDR are those which are resistant to all except 1-2 antimicrobials.

Ceftriaxone, cefotaxime, and (oral) cefixime are effective for the treatment of MDR enteric fever.

Immunization is a complementary approach in management since the protective efficacy can be overcome by a high inoculum in the case of food-borne diseases.

The typhoid vaccines available are the Vi polysaccharide vaccine and the Ty21a vaccine. Ty21a is a live attenuated vaccine recommended after 6 years and is given orally on days 1,3,5 and 7. Vi polysaccharide vaccine can be given after 2 years as a single dose with a booster dose for every 2 years.

Solution to Question 7:

If exclusive replacement feeding, giving nevirapine for 6 weeks, even if the baby is negative is true regarding the management of a baby born to an HIV-positive mother.

The national AIDS control organization (NACO) has recommended the following for an infant born to an HIV-positive mother.

According to the National AIDS Control Organization (NACO), antiretroviral therapy (ART) should be given to all HIV-positive pregnant women including during labor hours and breastfeeding. They are started on a triple-drug ART. ART should be given irrespective of the CD4 count, and clinical stage and should be continued lifelong.

Immediate care of HIV-exposed infants includes wiping of baby's mouth and nostrils, handling of infants with gloves, immediate cord clamping, and initiation of feeding. An infant born to an HIV-positive mother should be breastfed exclusively or by replacement feeds exclusively for 6 months of age.

Infants should receive ARV prophylaxis. A single drug or dual drug regimen is followed depending on the risk of transmission which can be low or high risk.

• Low-risk infants are those born to mothers with a low viral load which is detected after 32 weeks of pregnancy till delivery.

• High-risk infants are those born to HIV-positive mothers, not on ART, with viral load not estimated beyond 32 weeks, viral load not suppressed after 32 weeks, or diagnosed with HIV within 6 weeks of delivery.

HIV Risk	ARV prophylaxis	Duration
Low-risk infants	Syrup nevirapine or zidovudine	Birth to6 weeksof ageirrespective of feeding
High-risk infants	Syrup nevirapine + syrup zidovudine	Exclusive replacement feeding: Birth to 6 weeksExclusive breastfeeding: Birth till 12 weeks

Solution to Question 8:

Descriptive statistics is the branch that deals with describing the internal collection of data.

Statistical methods are mainly of two types

• Descriptive statistics is a type of statistical method which helps to describe, organise or summarise the data. It refers only to the actual data. They are described by the measures of central tendency and the degrees of dispersion.

• Inferential statistics involves going beyond the actual data and making inferences to test the hypothesis.

Other types of statistics:

• Analytical statistics is mainly used to test the hypothesis.

• Theoretical statistics is a mathematical approach which involves describing, predicting the events or analyzing the relationship between things.

Solution to Question 9:

Isolation in a separate room is mandatory for measles among the given options. It is transmitted by droplet infection as well as via the airborne route. The period of isolation is from the onset of symptoms (catarrhal stage) through the fourth day of the rash.

Isolation is defined as the separation of an infected person or animal for the period of communicability in places to prevent direct or indirect transmission to susceptible individuals.

Measles is an infectious disease mainly seen in the childhood age group. It is caused by an RNA paramyxovirus and is transmitted to humans primarily by droplet infection and also through airborne transmission.

Features include

• The prodromal phase-characterized by fever, cough, coryza, and Koplik's spots which are rashes seen in the buccal mucosa opposite the first and second molar.

• The eruptive phase- macular or maculopapular rash begins behind the ears and spreads to the face, neck, and extremities.

• Post measles phase is characterized by weight loss.

Some of the complications of measles include otitis media, laryngotracheobronchitis (croup), diarrhea, post-infectious measles, or subacute sclerosing pan encephalitis (SSPE). Subacute sclerosing panencephalitis is a rare but fatal complication.

Treatment is mainly aimed at supportive care such as relieving the symptoms, treating dehydration, and isolating patients.

The below images show a characteristic rash in measles and Koplik's spots on the buccal mucosa.





Other options:

Option B: Rabies is transmitted via a bite or scratch from an infected animal. Contraction of rabies by inhalation of virus-containing aerosols is extremely rare. Human-to-human transmission is possible but has not been confirmed.

Option C: Nipah is transmitted mainly from direct contact with sick pigs or their contaminated tissues. To prevent human-to-human transmission, close unprotected contact with infected people is avoided. Regular hand hygiene should be followed when caring for or visiting infected people.

Option D: If the diagnosis is confirmed as aseptic meningitis, the patient can often be discharged home. Isolation is required if the meningitis is of infectious etiology.

Solution to Question 10:

All the given options cause a widening of confidence interval except for an increase in sample size. An increase in sample size leads to a narrower confidence interval.

Research is usually done with samples drawn from a population, rather than by using the entire population. In order to get a fair idea of the population parameters, sample statistics and estimates of error in the sample are used to get a range of values. This range is the confidence interval (CI). It is estimated based on a desirable confidence level.

The confidence interval is calculated as the difference between upper and lower confidence limits.

The width of the CI is an indication of the utility of our population parameter estimation. Factors affecting the width of the confidence interval include:

• Desired confidence level- width of CI is directly proportionate to the confidence level

• Variability in the sample- width of CI is directly proportionate to the variability in the data

• Sample size- width of the CI is inversely related to the sample size

Therefore, increases in variability and confidence levels and decreases in sample size lead to wider confidence intervals.

Solution to Question 11:

The false statement regarding rabies immunoglobulin is it is exclusively given in the deltoid. Rabies immunoglobulin is given as much as possible into the wound site or around it.

Rabies immunoglobulin (RIG) is passive immunization. It is administered only once, preferably within 24 hours of the exposure, ie, on day 0 of the anti-rabies vaccine (ARV). If RIG was not given with the ARV 1st dose, it can be given up to the 7th day after the first dose of ARV. It is not given after 7 days of receiving the 1st dose of ARV.

The dose is 20 IU/kg body weight for human rabies immunoglobulin (HRIG) and for equine rabies immunoglobulin (ERIG), the dose is 40 IU/kg body weight. When HRIG is not available, ERIG can be used. They are potent, safe, and less expensive.

The entire RIG dose or as much as anatomically possible should be infiltrated into the wound or as close as possible. If it is likely that there are additional small wounds, the remaining RIG should be given IM as close as possible to the site of wound or exposure.

All patients with category III exposures should be given RIG. This includes:

• single/multiple transdermal bites/scratches

• licks on broken skin

• contamination of mucous membranes with saliva

• contact with bats

In immunocompromised patients, RIG should be given to both category II and III exposures.

Solution to Question 12:

The advice for the patient is to give the second dose of DPT (Diphtheria-pertussis-tetanus).

DPT is a combined vaccine given for immunization against the three diseases diphtheria, pertussis, and tetanus. It is given to infants as early as 6 weeks following birth. The pertussis component of the vaccine is not recommended beyond 6 years due to the decrease in the severity of pertussis infection with age. 7 years of age onwards, only Td combinations are used.

It is given in 3 doses with an interval of 4 weeks followed by booster doses at 18-24 months and at 5 years of age. The dosage is 0.5 mL and is given in the anterolateral aspect of the thigh intramuscularly. The immunization can be resumed without repeating the previous dose in case of any delayed or interrupted immunization.

Adverse effects include fever, and neurological complications such as encephalitis, convulsions, infantile spasms and Reye's syndrome.

Anaphylaxis is a contraindication for subsequent vaccination with diphtheria, pertussis, or tetanus vaccine. Encephalopathy following vaccination is a contraindication for pertussis vaccine. DT is given for successive doses.

Events such as inconsolable cry, fever more than 40.5°C, seizures with or without fever are considered as precautions, and not contraindications for DTwP.

Note: IAP guidelines state that "temperature more than >40.5°C is a precaution and not contraindication". Park's 27th Ed states that "temperature above 40°C is a contraindication to DPT". This topic is controversial, however, IAP guidelines are commonly followed by practitioners.

Solution to Question 13:



Box C stands for health motivation.

The health belief model was proposed from the idea that a person's health behavior changes in accordance with certain beliefs.

It can be categorized as follows:

• Individual perceptions.

• Variable factors that modify beliefs.

• Action.

Individual perceptions:

Perceived susceptibility refers to a person's own beliefs about the probability of developing a disease. If the perceived susceptibility is high, an individual is more likely to engage in health-promoting behavior to reduce the chances of contracting the illness.

Perceived severity is an individual's assessment of the severity of a particular health problem. If the disease is perceived to have a high severity, then the person is more likely to adopt a healthier lifestyle.

Perceived threat comprises perceived susceptibility and severity. People with a higher perceived threat tend to take on health-promoting activities.

Action:

Perceived benefits refer to a person's subjective judgment on how a particular health habit is modifying his chances of developing a disease. If the person believes that a habit is likely to reduce his/her chances of susceptibility to a disease or lower the severity of a disease, he/she is more likely to follow that healthy habit.

Perceived barriers to action also influence if a person wants to follow healthy behavior. The benefits of a particular health habit must overweigh the barriers that a person must overcome to continue following that behavior.

Cues to action may be internal or external. Internal cues include pain and clinical symptoms. External cues include awareness through media, physicians, and friends.

Behavioral modification (Option B) is part of the action after assessing all of the above-mentioned concepts.

Self-efficacy (Option D) can be defined as a person's capacity in achieving his health goals. A person's confidence in his/her ability to overcome limitations to attain his/her health goals was found to be a major factor in determining the outcome of his behavioral changes.

The health belief system is thought to be useful in analyzing health behavior patterns and formulating disease prevention plans based on these beliefs. All internal and external factors, including demographic and psychological characteristics, aim at health motivation to attain the action phase of a health belief system.

Other options:

Option A: Perceived control refers to a person's internal belief in the control they would have over things that are part of the past or the future or over a process, behavior, and outcome. Perceived control can influence all of the categories of the health belief system.

Solution to Question 14:

A negative result in a diagnostic test with a sensitivity of 98% implies that the patient is less likely to have the disease.

Sensitivity is the proportion of all people with the disease who test positive. It is the probability of the test being positive when the disease is present. The sensitivity of a test is given by those who have the disease and test positive (true positives) out of the total diseased population.

Other statements:

Option A: The patient would not need another point-of-care diagnostic test for the same clinical condition as the sensitivity of the diagnostic test is 98%.

Option B: Prevalence refers to the number of old and new cases at a given point in time or over a period of time in a given population. It cannot be derived from the given data.

Option D: The higher the sensitivity of a diagnostic test, the lower will be the number of false negatives. Since the test has a 98% sensitivity, the patient most likely does not have the disease.

Solution to Question 15:

The correct statements drawn from the values given in the table are 3 and 4.

An odds ratio is a measure of the association between an exposure and an outcome. It represents the odds that an outcome will occur given a particular exposure in comparison to the odds of the outcome occurring in the absence of that exposure.

An odds ratio of 1 indicates that the risk factor is not related to the disease.

An odds ratio of less than 1 indicates that a person with the disease is less likely to have been exposed to the factor, implying that the factor may be a protective factor against the disease.

An odds ratio greater than 1 indicates that the odds ratio is significant, and the factor can be ascertained as a risk factor.

The p-value is the probability of obtaining a result at least as extreme as the one that was actually observed in the biological or clinical experiment. A p-value of less than 0.05 indicates that the test result is statistically significant.

Other options:

Statement 1: Female population is at greater risk of developing blindness is a false statement; since the odds ratio is <1, the female population will be at lesser risk of developing blindness than the male population.

Statement 2: People dwelling in rural areas are at increased risk of developing blindness is also false; a p-value of >0.05 means that this association is not statistically significant.

Solution to Question 16:

The image shows an established case of chronic lymphatic filariasis, where hygiene is being maintained to prevent further disability by secondary bacterial infection. Disability limitation is a mode of intervention in tertiary prevention.

Other options:

Option A: Specific protection is a mode of intervention in primary prevention which aims at limiting the number of cases of a particular disease. This includes immunization, use of specific nutrients, chemoprophylaxis, accident protection, and avoidance of allergens.

Option C: Primary prevention is a level of prevention which is defined as action taken prior to the onset of a disease, which removes the possibility that a disease will ever occur. It is accomplished by measures designed to promote general health and well-being.

Option D: Health promotion is a mode of intervention in primary prevention, and is the process of enabling people to increase control over, and improve health. Health promotion may be achieved through health education, environmental modification, nutritional interventions or lifestyle changes.

Solution to Question 17:

The proportion of newly diagnosed patients with grade 2 disability is useful in determining whether the health system is giving importance to identifying leprosy in the community.

The National Leprosy Eradication Programme (NLEP) was introduced in 1983. The strategy of NLEP is to control the disease through a reduction in the infection and the infective source, thus breaking the chain of disease transmission. It functions under National Health Mission (NHM).

The vision of the NLEP is 'Leprosy-free India'. Disability prevention and medical rehabilitation (DPMR) is the priority of the programme. The mission of NLEP is to provide quality services free of cost to all people through the integrated healthcare system.

Main indicators for evaluating case detection:

• Proportion of grade-2 disabilities among new cases

• Proportion of females among new cases

• Proportion of MB among new cases

• Proportion of child (0-14 years) among new cases

• Child rate per 1,00,000 population

• Scheduled caste new case detection rate

• Scheduled tribe new case detection rate

Solution to Question 18:

The correct statements are 1 and 2.

Vaccine vial monitors (VVM) are heat-sensitive indicators routinely used to monitor heat exposure of the vaccine and determine vaccine viability as the vaccine vial moves across the supply chain. It records cumulative heat exposure through a gradual change in color.

If the color of the inner square is the same or is darker than the outer circle, it indicates that the vaccine has been exposed to high temperatures and should be discarded.

A VVM can be used to assess vaccine potency and not vaccine efficacy. Vaccine potency is the ability of the vaccine to provide adequate immunity to the patient. Vaccine potency is irreversibly lost when the vaccine is exposed to unfavorable temperatures; hence, VVM indicating heat exposure can be used to assess whether the vaccine potency is likely to be intact or not. Vaccine efficacy, on the other hand, is a measure of the immunity conferred by the vaccine or the ability of the vaccine to prevent the disease in a vaccinated person under ideal conditions, such as clinical trials. The term vaccine efficacy is used in relation to findings of clinical trials, whereas the term vaccine potency is used for regular usage of the vaccine. Vaccine efficacy is determined only during clinical trials, whereas vaccine potency is tested for every batch of the vaccine that is manufactured.

There are currently 4 types of VVMs; they are chosen to match the heat sensitivity of the vaccines based on the VVM number:

• VVM 30: High stability

• VVM 14: Medium stability

• VVM 7: Moderate stability

• VVM 2: Least stable

Purpose of VVMs:

• Ensures that vaccines damaged by heat are not administered.

• Decision on which vaccines to be kept after a cold chain break occurs, minimizing unnecessary vaccine wastage.

• Decision on which vaccine is to be used first. Vaccines showing significant heat exposure should be used before those that show lower heat exposure, even if their expiry date is longer.

The expiry date of a vaccine cannot be relaxed even if VVM is in an acceptable range. Expired vaccines must be discarded. The expiry date of a vaccine cannot be determined using a VVM as it does not monitor the vaccine components. It only ensures that the vaccines have been exposed to the required temperatures and are viable for use.

There are two different locations for VVMs:

• On the label of the vaccine: The vaccine vial, once opened, can be kept for subsequent immunization sessions for up to 28 days, regardless of the formulation of the product.

• In a location other than on the label (e.g., cap or neck of ampoule): The vaccine vial, once opened should be discarded at the end of the immunization session or within six hours of opening, whichever comes first, regardless of the formulation of the product.

Solution to Question 19:

The model of disease causation depicted below is a sufficient component cause model.

It was developed by Rothman based on the idea that an outcome can only happen when a sufficient cause is present. A disease can have multiple sufficient causes. But, each sufficient cause

can consist of a pie chart of component causes, a causal pie. One component cause can be a part of more than one sufficient cause. A component cause can be considered necessary if it is a part of a disease's sufficient causes.

In the given model, each pie chart represents a sufficient cause made of component causes. Since B is present in all the charts, it can be considered a necessary cause. All the others are contributing causes.

Other options:

Option B: An epidemiological triad depicts the interrelationship between the causative agent, the host, and the environment. All of them contribute to disease development. The aim of an epidemiological triad is to identify how to break this relationship and prevent the outbreak of the disease.

Option C: PERT is used for network analysis. It is a type of quantitative management technique that helps in detailed planning and comprehensive supervision.

Option D: The web of causation considers all the predisposing factors and their complex interrelationship with each other. This model is ideally suited to study chronic disease. The web of causation does not imply that the disease is controlled only when all the multiple chains of causation are removed. Sometimes removal of one significant link is sufficient to control the disease.

Solution to Question 20:

Attributable risk (AR) describes the extent of outcome (lung cancer) that can be attributed to exposure (smoking). In the given scenario, an AR of 90% means that 90% of lung cancer cases among smokers are attributable to smoking. Therefore, 90% cases of lung cancer can be eliminated in smokers if smoking is eliminated.

AR is the difference in incidence rates of disease between an exposed group and an unexposed group. AR is indicative of the extent of the disease attributed to the exposure. It also refers to the amount of disease that could be eliminated in the exposed persons if the exposure is controlled or eliminated. It also helps to determine the impact a successful preventive or public health program might have in case the exposure to cause is reduced or eliminated.

Population-attributable risk (PAR) is the difference in the incidence of disease in the total population and the incidence of disease among those not exposed to the supposed cause. PAR is useful in providing an estimate of the amount by which the disease could be reduced in that whole population if the suspected factor was eliminated or modified.

For example, a PAR of 90% in the context of smoking and lung cancer means that 90% cases of lung cancer in the population can be attributed to smoking.

Relative risk is defined as the ratio of incidence of the disease (outcome) among exposed and unexposed groups. It helps in establishing the association between a cause and a disease. However, it does not help in assessing the extent of the cause bringing about the disease (outcome).

Solution to Question 21:

The measures that are taken to prevent disasters from causing emergencies or to lessen the likely effects of emergencies are part of disaster mitigation.

Disaster mitigation measures aim to reduce the vulnerability of the system. It complements disaster preparedness and response activities and includes measures taken to lessen the likely effects of emergencies. Some of these measures are:

• Flood mitigation

• Land-use planning

• Improved building codes

• Reduction and protection of vulnerable populations or structures.

Other options:

Option A - Disaster response includes activities such as rescue and first aid, field care, triage and tagging that form a part of the emergency care needed in the first few hours after a disaster.

Option B - Disaster reconstruction is a part of the recovery phase after a disaster.

Option C - Disaster rehabilitation refers to measures that lead to the restoration of pre-disaster conditions. These measures include water supply restoration, food safety, basic sanitation, personal hygiene, and vector control.

The disaster management cycle is given below:



Solution to Question 22:

Isoniazid inhibits mycolic acid synthesis and Bedaquiline inhibits ATP synthase.

Other options:

Pyrazinamide is not an enzyme inducer. Pyrazinamide is converted into its active form pyrazinoic acid. Pyrazinoic acid disrupts mycobacterial cell membrane metabolism and transport functions.

Amongst the anti-tubercular drugs, rifampicin is an enzyme inducer.

Rifampicin can be used in treatment of meningococcal carriers, prophylaxis in contacts with Haemophilus influezae type b, and serious staphylococcal infections.

Solution to Question 23:

To start multidrug therapy (MDT), the criteria that should be satisfied are a hypopigmented patch with sensory loss and peripheral nerve thickening with sensory loss.

Leprosy (Hansen's disease) is a chronic infectious disease caused by Mycobacterium leprae. It mainly affects the skin and peripheral nerves. The disease has two polar forms, lepromatous leprosy, and tuberculoid leprosy, lying at the two ends of a long spectrum of the disease.

Leprosy is diagnosed by the detection of at least one of the following cardinal signs:

• Definite loss of sensation in a pale hypopigmented patch

• Thickened or enlarged peripheral nerve, with loss of sensation and/or weakness of the muscles supplied by that nerve (statement 2)

• Presence of acid-fast bacilli in a slit-skin smear.

However, a skin smear is not a prerequisite for implementing MDT. Since MDT has been introduced, many procedures have been simplified, so that leprosy patients can be detected by health workers in the field and MDT can be immediately started.

Solution to Question 24:

The correct match is 1-d, 2-b, 3-a, and 4-c.

Natural history of disease refers to the progression of a disease in an individual over time, in the absence of treatment. It has two phases, a period of pre-pathogenesis and a period of pathogenesis.



The period of pre-pathogenesis is before the onset of disease in humans. An interaction between the agent, host and the environment occurs in this phase before the pathogenesis. It starts with appropriate exposure to or accumulation of factors sufficient for the disease process to begin in a susceptible host. The exposure is a microorganism in an infectious disease. For lung cancer, it is any factor that initiates the process, such as asbestos fibres or components in tobacco smoke. Primary prevention (health promotion and specific protection) is applicable in this period.

The period of pathogenesis starts after the disease process has been triggered and pathological changes occur without the individual being aware of it. It begins with the entry of the organism till recovery, death, or disability. The subclinical or asymptomatic period between exposure and onset of symptoms is called the incubation period for infectious diseases, and the latency period for chronic diseases. The onset of clinical symptoms is the beginning of the clinical stage of the disease. Secondary (halting of the disease process) and tertiary prevention (measures to limit disability) are applicable in this period.

Rehabilitation has been defined as the use of medical, social, educational, and vocational measures for training the individual to the highest possible level of functional ability. It is a part of tertiary prevention like providing callipers for walking patients and arranging schooling after being affected by polio.

Solution to Question 25:

Pearl index is the common method used to measure contraceptive failure rate.

It is defined as the number of failures per 100 woman-years of exposure (HWY). This rate is given by the formula:

Pearl index = Total accidental pregnancies x 1200/Total months of exposure

Solution to Question 26:

The correctly matched pairs are as follows:

• Walter Reed showed that yellow fever is transmitted by the Aedes mosquito.

• Edward Jenner - smallpox vaccination

• James Lind - prevention of scurvy by intake of citrus fruits

Scientist/ Doctor	Achievements
Susruta	Father of Indian SurgeryFather of Plastic and Cosmetic SurgeryAuthor of “Sushruta Samhita”
Hippocrates	First True EpidemiologistFather of MedicineAuthor of “Airs, Water and Place”
Charaka	Father of Indian MedicineAuthor of “Charaka Samhita”
Avicenna	Author of “Canon of Medicine” and “The Book on Healing”
Andreas Vesalius	Father of Modern AnatomyAuthor of “Fabrica”
Hammurabi	Author of “Babylonian Code of Hammurabi”
John Snow	Father of Epidemiology
Louis Pasteur Robert Koch / Ferdinand Cohn	Father of Bacteriology
Samuel Hahnemann	Father of Homeopathy

Solution to Question 27:

The formula given above is used to calculate the standard deviation.

Standard deviation (or σ) is a measure of how dispersed the data is in relation to the mean. Data clustered around the mean have a low standard deviation, while data that are more spread out have a high standard deviation. A standard deviation close to zero indicates that data points are close to the mean, whereas a high or low standard deviation indicates data points are respectively above or below the mean.

Standard error = SD/√n, where SD is the standard deviation and n is the sample size

Standard error will reduce when the sample size is increased.

The median divides the frequency distribution in half when all the scores are listed in order. When a distribution has an odd number of elements, the median is the middle value. When the distribution has an even number of elements, the median is the average of the two middle scores.

Variance is the mean of the squares of all the deviation scores of the distribution.

Solution to Question 28:

A child presenting with abrasion with slight oozing after a dog bite comes under category III. The correct statements are 1, 2, 3, and 6.

The following measures should be taken immediately to stop the contraction of rabies:

• Cleansing: The most important step in the prevention of rabies is washing the affected area with plenty of soap and water under a running tap for 15 minutes. This is a standard procedure that minimizes the contraction of rabies.

• Chemical treatment: After washing and drying the wound, disinfectants like povidone-iodine and spirit can be applied.

• Suturing: Bite wounds are not sutured generally. If suturing is unavoidable, rabies immunoglobulin should be infiltrated in the depth and around the wound, and suturing should be delayed by a few hours.

• In unavoidable situations, minimal loose sutures may be applied to arrest bleeding.

• Other measures: Local antibiotics and tetanus toxoids should be given.

Different wound categories for post-exposure prophylaxis:

Post-exposure prophylaxis regimen for rabies:

Intramuscular regimen: In this regimen, 1 ml or 0.5 ml of vaccine is injected into the deltoid in patients with category II and III wounds.

• Essen regimen: A 5-dose regimen in which 1 dose of vaccine is given on days 0, 3, 7, 14, and 28.

• Zagreb regimen: A 4-dose multisite regimen in which 2 doses of vaccine are given on day 0, followed by 1 dose on days 7 and 21.

Intradermal regimen: In this regimen, 0.1 ml of injection is given at 2 sites on days 0, 3, 7, and 28.

Equine Rabies Immunoglobulin at a dose of 40 IU/kg (a maximum of 3000 IU) body weight is given once. It is given to all Category III bites and also Category II bites in the case of immune-compromised persons.

Categories of wound	Post-exposure prophylaxis
Category I- touching or feedi ng animals, licks on intact ski n	None
Category II- nibbling of unco vered skin, minor scratches, or abrasionswithout bleeding	Immediate vaccination andlo cal treatment of the wound
Category III- single or multip le transdermal bites or scratc hes, licks on broken skin cont acts with bats	Immediate vaccination anda dministration of rabiesimmu noglobulin; localtreatment of the wound

Solution to Question 29:

The species of mosquito that breeds in dirty water is Culex.

Culex fatigans are found everywhere in India and they transmit Bancroftian filariasis. Culex mosquitoes are also vectors for Japanese encephalitis, West Nile fever, and viral arthritis.

It can be identified by the presence of unspotted wings and hunched back. The larva has a head that is suspended downwards at an angle to the water surface and a siphon tube is present.

It breeds in dirty water collections like stagnant drains, septic tanks, and burrow pits. The peak biting time is during midnight. The preferred biting site is the legs, mainly below the knee.

Solution to Question 30:

The image given below shows artemisinin-based combination therapy (ACT), which is used in the treatment of falciparum malaria.

ACT consists of artesunate given for 3 days and sulphadoxine-pyrimethamine given for 1 day. It is supplied by the government of India under the national vector-borne disease control program (NVDCP).

It is given to all confirmed cases of Plasmodium falciparum infection, found positive by microscopy or rapid diagnostic tests. Primaquine with dose of 0.75 mg/kg of body weight is given on day 2 along with it. Artemether-lumefantrine is used for the treatment of falciparum malaria in the northeastern states of India.

This drug is not given in pregnancy and in children less than 5 kg of body weight.

Solution to Question 31:

According to the Anemia Mukt Bharat initiative, pregnant women with mild and moderate anemia is treated with

• Two iron and folic acid tablets (60 mg elemental iron and 500 mcg folic acid) daily, orally given by the health provider during the ANC contact

• Parenteral iron (IV iron sucrose or ferric carboxy maltose [FCM]) may be considered as the first line of management in pregnant women who are detected to be anemic late in pregnancy or in whom compliance is likely to be low (high chance of loss to follow-up)

Anemia Mukt Bharat 2018 scheme:

This is an initiative that aims to strengthen the existing mechanisms and foster newer strategies for tackling anemia. It focuses on six target beneficiary groups, through six interventions and six institutional mechanisms to achieve their target by 2022.

Interventions:

• Prophylactic iron and folic acid supplementation.

• Deworming.

• Intensified year-round Behaviour Change Communication Campaign (Solid Body, Smart Mind) including ensuring delayed cord clamping in newborns.

• Testing of anemia using digital methods and point-of-care treatment.

• Mandatory provision of Iron and Folic Acid fortified foods in government-funded health programs.

• Addressing non-nutritional causes of anemia in endemic pockets, with a special focus on malaria, hemoglobinopathies, and fluorosis.

Beneficiaries:

• Children, 6-59 months of age

• Children, 5-9 years of age

• Adolescent girls and boys, 10-19 years of age

• Women of reproductive age, 20-49 years of age

• Pregnant women

• Lactating mothers, 0-6 months

Institutions through which activities are carried out:

• Intra-Ministerial Coordination

• National Anemia Mukt Bharat Unit

• National Centre of Excellence and Advanced Research on Anemia Control (NCEAR-A)

• Convergence with other Ministries

• Strengthening Supply Chain and Logistics

• Anemia Mukt Bharat Dashboard and Digital Portal

Community medicine NEET 2022

Question 1:

The average life expectancy for a woman in Japan is 87 years. Due to recent advances in testing for cervical cancer, there is an increase in life expectancy by 15 years. The healthcare utility value is 0.8. Which of the following can be calculated from the parameters given?

a) HALE

b) DALY

c) DFLE

d) QALY

Question 2:

You are working in a primary health center (PHC) situated in a high seismic zone. Which of the following will you do as part of preparedness for an emergency?

a) 1, 2, 3, 4

b) 2, 3, 4

c) 1, 2, 3

d) 1, 2, 4

Question 3:

Although many animals are implicated in the spread of rabies, dogs are the most common ones. Also, it usually affects children in developing countries. Knowing this, what is the most cost-effective and logical way to reduce the incidence of rabies?

a) Testing all the dogs for rabies

b) Reduce stray dog population and vaccinate all dogs

c) Increase the laboratory facilities

d) Increase capacity of healthcare workers for surveillance

Question 4:

Which of the following steps is not included in STEP approach of WHO?

a) Therapeutic assessment

b) Physical assessment

c) Psychological assessment

d) Behavioral assessment

Question 5:

A male patient diagnosed with tuberculosis took complete treatment. Sputum examination was done after the completion of the intensive and the continuation phases. It was found to be negative. What is the status of the patient?

a) Cured

b) Treatment completed

c) Lost to follow up

d) Treatment failed

Question 6:

A cohort study was conducted with drinkers and non-drinkers of green tea to study its effect on diabetes mellitus. The risk ratio was found to be 0.84. Which of the following statements is correct?

a) Green tea reduces the risk of diabetes

b) Green tea increases the risk of diabetes

c) Data insufficient to establish causal association

d) The value (0.85) tends to be close to 1, hence there is no effect

Question 7:

Which statement refers best to the criteria for starting an urban community health center?

a) Caters to a population of 1-1.5 lakh

b) Referral centre for 2- 3 primary health centers

c) No sub-district and district hospitals present in the area

d) Should have a 100-bed facility in metro cities

Question 8:

How is a broken vaccine vial disposed, according to biomedical waste management?

a) Puncture proof blue bin

b) White container

c) Yellow container

d) Red container

Question 9:

In an urban area in the state of Madhya Pradesh, a primigravida goes for institutional delivery after being motivated by an ASHA worker. What are the benefits they will receive (in terms of money in rupees) under the Janani Suraksha Yojana?

a) 1000 for mother and 400 for ASHA

b) 1400 for mother and 600 for ASHA

c) 600 for mother and 400 for ASHA

d) 400 for mother and 600 for ASHA

Question 10:

Which of the following agency provides seed and manure in applied nutrition programs in schools?

a) CARE

b) UNDP

c) UNICEF

d) WHO

Question 11:

Many children from a particular community coming to a hospital were detected to have acute lymphoblastic leukemia (ALL). It was assumed that it is due to the presence of cytotoxic waste in the water of that community. If a case-control study has to be done to find whether the chemical and ALL are associated, what will be taken as the control?

a) Children from the area exposed, but unaffected with the disease

b) Children from the area not exposed and affected with the disease

c) Children coming to your OPD, who do not have the disease

d) All children with ALL irrespective of exposure status

Question 12:

An auxiliary nurse midwife has to conduct a vaccination camp in a village. She received 2 open vials, one of which is a pentavalent vaccine and the other is an MR vaccine. What can she do regarding the utilization of these vials?

a) Use MR vaccine and discard pentavalent vaccine

b) Use pentavalent vaccine and discard MR vaccine

c) Use both

d) Discard both

Answer Key

Question No.	Correct Option
1	d
2	c
3	b
4	a
5	a
6	a
7	d
8	a
9	a
10	c
11	c
12	b

Detailed Explanations

Solution to Question 1:

QALY (quality-adjusted life year) can be calculated by the utility value.

QALY is a disability rate indicator. It is based on the number of years that would be added to life due to medical intervention. A utility value of 1.0 is given for each year of perfect health and 0 for death. It is the measure of the disease burden which includes both quality and quantity of life.

Disability rates that are related to illness and injury are used as supplementation to mortality and morbidity indicators. It is based on the concept that a full range of daily activities implies good health. It is divided into:

• Event-type indicators: It counts the number of days of restricted activity, bed disability days, and work loss days.

• Person-type indicators: It counts limitation of mobility and limitation of activity.

Other options:

Option A: HALE (health adjusted life expectancy) is an indicator that estimates the number of years a newborn is expected to live in full health based on the current rates of ill health and mortality. HALE is calculated using the life expectancy at birth with an adjustment for time spent in poor health. It was known as DALE (disability-adjusted life expectancy) previously.

Option B: DALY (disability adjusted life years) is an indicator that estimates the number of years lost due to ill-health, disability, or early death. It is calculated by:

DALY= Years of life lost (YLL) + Years lived with disability (YLD)

Option C: DFLE (disability-free life expectancy) is an indicator that estimates the average number of years an individual is expected to live free of disability if the current pattern of mortality and disability continues to apply. It is also known as active life expectancy and is calculated by:

DFLE = Life expectancy - Duration of disability

Solution to Question 2:

Steps 1, 2, and 3 are done as part of preparedness for an emergency by someone working in a PHC situated in a high seismic zone.

Emergency preparedness refers to long-term development activities that strengthen the capacity of a country to effectively manage all kinds of emergencies. It involves the following steps:

• Disaster preparedness by making sure all financial and other resources are available

• Increase public awareness through campaigns and loudspeakers

• Conduct a simulation for the disaster and assess the response

• Evaluate the risk of disaster

• Information sessions coordination with news media

• Adopt standards and regulations

Disaster risk management is a systematic process using administrative decisions, policies, and strategies to lessen the impacts of natural hazards and disasters. The steps are:

• Response - This includes activities such as:

• Search, rescue, and first aid: This step is important since the organized relief services will be able to address only a small fraction of demand after a major disaster.

• Field care: This includes making health care available regardless of facilities and operating status.

• Triage: Classifying the injured based on the severity of the injuries and chances of survival with medical intervention.

• Tagging: Tags with name, age, triage category, diagnosis and treatment are put for identification purposes.

• Identification of the dead: The dead bodies should be shifted from the disaster site to the mortuary.

• Rehabilitation - This includes measures that lead to the restoration of pre-disaster conditions. These measures include water supply restoration, food safety, basic sanitation, personal hygiene, and vector control.

• Mitigation - This includes measures to prevent disasters from causing emergencies or to lessen the likely effects of emergencies. Some of them are flood mitigation, land use planning, improved building codes, and the reduction and protection of vulnerable populations or structures.

• Preparedness - This is a program that aims at strengthening the capacity of a country to manage all types of emergencies efficiently. Its objective is to ensure that appropriate measurements are taken for the relief and rehabilitation of disaster victims.

Some measures that should be followed by all in case of an emergency include:

• Avoiding telephone use, unless it is to call for help

• Listen to and carry out the radio or loudspeaker broadcast messages

• Carry out the official instructions given over the radio or by loudspeaker

• Keep a family emergency kit ready

Note: Vaccination of the general population post-disaster against typhoid, cholera, and tetanus is not recommended.



Solution to Question 3:

Reducing the stray dog population and vaccinating all dogs is the most cost-effective and logical way to reduce the incidence of rabies. The Animal Birth Control initiative from the Government of India is directed towards this goal.

Rabies is a highly fatal viral disease caused by Lyssavirus type 1, affecting the central nervous system. Dogs are considered the major source of rabies infection. Thus immunization of dogs at the age of 3-4 months and booster doses at regular intervals is recommended to control rabies. The type of vaccines used are:

• BPL inactivated nervous tissue vaccine - it is a single-dose vaccine. Booster is advised after 6 months and subsequently every year.

• Modified live virus vaccine - it is also a single-dose vaccine and a booster is given every 3 years.

Other methods to control rabies include:

• Registration and licensing of all domestic dogs

• The immediate killing of dogs and cats bitten by rabid animals

• 6 months quarantine for imported dogs

• Educating people regarding dog care and prevention of rabies.

Note: Strengthening laboratory facilities and surveillance are also the methods used to reduce the incidence according to the National Rabies Control Program (NRCP)

Humans get infected after a deep bite or scratch by an infected animal. The transmission also occurs when the saliva of the infected dog comes into direct contact with human mucosa or fresh skin wounds.

Rabies in man is also known as hydrophobia. The clinical features usually begin as prodromal symptoms like headache, and fever followed by neurological symptoms. The patients usually present with confusion, hallucination, irritability, hydrophobia, aerophobia, and muscle spasms. They also present with symptoms of autonomic dysfunction such as hypersalivation, lacrimation, cardiac arrhythmias, etc.

Laboratory tests for rabies include:

• Fluorescent antibody test is a highly reliable and best test for rapid detection of rabies antigen in the infected specimens

• Microscopic examination of the brain tissue to demonstrate Negri bodies.

Solution to Question 4:

Therapeutic assessment is not included in the STEP approach of WHO.

STEP approach of WHO is a survey methodology developed to assess risk factors of non-communicable diseases(NCD). It helps the countries to establish an NCD surveillance system. The steps include the following:

• Questionnaires contain questions about behavioral risk factors like tobacco use, alcohol use, diet, and physical activity.

• Physical measurements contain blood pressure, height, weight, and waist size

• Biochemical measurements contain blood glucose, blood lipids, urinary sodium, and creatinine.

Note: Mental health and suicide are included in optional behavioral risk factors (step 1). Hence, psychological assessment is also included in the STEP approach.

Surveillance is defined as systematic and continuous data analysis and taking appropriate preventive actions according to the results obtained. Identification of cases that are missed and not notified is known as sentinel surveillance. It is used in communicable diseases such as HIV, malaria, etc. The study population for HIV sentinel surveillance are sex workers, IV drug abusers, transgender people, migrant workers, truck drivers, etc.

Solution to Question 5:

The negative sputum examination at the end of the treatment implies the status of the patient is cured.

Tuberculosis (TB) is a communicable disease caused by Mycobacterium tuberculosis. The disease mainly affects the lungs and causes pulmonary tuberculosis.

A bacteriologically confirmed TB case is one in which the biological specimen tests positive on smear microscopy and culture. A clinically diagnosed TB case is one that is diagnosed as an active TB case by a clinician, who has decided to give the patient a full course of TB treatment.

There are two phases in the treatment of TB. The intensive phase lasts for 2 months and four drugs are used. They are isoniazid, rifampicin, pyrazinamide, and ethambutol. The continuation phase lasts for 4 months, where isoniazid is given along with rifampicin and Ethambutol.

After the completion of anti-tubercular treatment, the sputum/culture is tested and one of the outcomes given below is assigned to the patient:

• Cured - A TB patient whose smear or culture turned negative in the last month of treatment and on at least one previous occasion

• Treatment completed - A TB patient who completed treatment without any failure, but has no record to show that sputum smear or culture results in the last month of treatment and on at least one previous occasion were negative; it may be either because tests were not done or because results are unavailable

• Treatment failed - A TB patient whose sputum smear or culture is still positive after treatment for 5 months or more

• Lost to follow-up - A TB patient who did not start treatment or whose treatment was interrupted for 2 consecutive months or more

• Died - A TB patient who dies before starting or during the course of treatment for any reason

• Not evaluated - A TB patient for whom no treatment outcome is assigned; this includes cases "transferred out" to another treatment unit and cases for whom the treatment outcome is unknown to the reporting unit whose treatment I interrupted

Solution to Question 6:

Green tea reduces the risk of diabetes mellitus is the correct statement since the risk ratio is 0.84.

Relative risk (RR) or risk ratio is the ratio of the incidence of the disease (or death) among exposed and the incidence among non-exposed. It can be determined from a cohort study.

It is a direct measure of the strength of the association between suspected cause and effect. The larger the RR, the greater the strength of the association between the suspected factor and disease.

• RR < 1 indicates inverse association(protective effect).

• RR = 1 indicates no association

• RR > 1 indicates a positive association.

• RR of 2 indicates a 100% increase in risk.

In the question, the risk ratio is less than 1. Hence it implies the incidence of diabetes mellitus is reduced when exposed to green tea.

Solution to Question 7:

An urban community health center (U-CHC) has a 100-bed facility in metro cities.

U-CHC acts as a referral center for 4–5 urban primary health centers (U-PHC). It is usually a 30–50 bedded facility that caters to a population of 2,50,000.

When it comes to metro cities, it caters to a population of 5,00,000 with a 100-bed facility. These hospitals will have surgeons, gynecologists, physicians, and pediatricians, along with x-ray and laboratory facilities.

The specialists at the community health center can refer a patient to the state-level hospital or the nearest medical college hospital directly. The subdistrict and district hospitals operate in addition to CHC to cater to the population.

Services of CHC:

• Care of routine and emergency cases in medicine and surgery

• Maternal, newborn, and child health

• Family planning

• All the national health program functions

• Physical medicine and rehabilitation services

Solution to Question 8:

A broken vaccine vial is disposed of in a puncture-proof blue bin, according to biomedical waste management (BMW).

According to BMW, broken or discarded contaminated glass materials such as vials, ampules, etc are disposed of in boxes marked with blue color. It is then disinfected either by washing with detergent, autoclaving, or microwaving and sent for recycling.

The methods used for treatment and disposal of healthcare waste are:

• Incineration: This is a dry oxidation process done using high temperatures.

• Chemical disinfection: Chemicals are added to waste to kill or inactivate the microorganisms.

• Wet and dry thermal treatment: Shredded infectious waste is exposed to high temperature, high-pressure steam and it resembles an autoclave sterilization process.

• Microwave irradiation: Microorganisms are killed using high-frequency microwave.

• Land disposal.

• Inertization: Waste is mixed with cement and other substances before disposal to reduce toxicity.

Solution to Question 9:

According to Janani Suraksha Yojana (JSY) scheme, the mother will receive 1000 rupees and the ASHA worker will receive 400 rupees.

Janani Suraksha Yojana (JSY) is a program under the National Rural Health Mission (NRHM), whose aim is to reduce maternal mortality by encouraging delivery in government health facilities. Under the scheme, a financial incentive is also provided for giving birth in a government health facility.

The Accredited Social Health Activist (ASHA) is the link between pregnant women and health institutions. She is responsible for making institutional antenatal and postnatal care available.

States are divided into low-performing and high-performing on the basis of institutional delivery rate. The 10 low-performing states (LPS) with a low institutional delivery rate are Uttar Pradesh, Uttarakhand, Madhya Pradesh, Jharkhand, Bihar, Rajasthan, Chattisgarh, Odisha, Assam, and Jammu & Kashmir. All the other states are high-performing states (HPS).

The incentives under JSY are as follows:

Under this scheme, the cost of cesarean section and management of obstetric complications is also subsidized, up to Rs 1500 per delivery to the government institutions, where government specialists are not present.

Category	Rural area	Urban area
LPS	Mother package: 1400/-ASHA's package:600/-Total:2000/-	Mother package: 1000/-ASHA's package:400/-Total:1400/-
HPS	Mother package: 700/-ASHA's package:600/-Total:1300/-	Mother package: 600/-ASHA's package:400/-Total:1000/-

Solution to Question 10:

UNICEF provides seed and manure in applied nutrition programs in schools.

Applied Nutrition Program is implemented with the aim of producing various types of productive food by the community for the community itself.

UNICEF assists the program by providing implements, seeds, manure, and water supply equipment. If free land is available in schools, the facilities provided by UNICEF should be used in developing school gardens.

The produce obtained from the gardens can be utilized for school feeding programs and for nutrition education.

Another strategy called GOBI was also implemented by UNICEF with the aim of a child health revolution.

Other options:

Option A- CARE- Cooperative for Assistance and Relief Everywhere aims at providing food for children in the age group of 6-11 years. It is also helping in projects like integrated nutrition and health project, better health and nutrition projects, anemia control projects, etc.

Option B- UNDP- The United Nations Development Program provides funds for technical assistance. Its main objective is to help poorer nations in developing their human and natural resources.

Option D- WHO- World Health Organization aims at the attainment of the highest level of health by all people, which helps them to lead a socially and economically productive life.

Solution to Question 11:

Children coming to the OPD, who do not have the disease, will be taken as the control for the case-control study.

The prerequisite for selecting a control group is that they must be free from the disease under study. They must be as similar as possible to the cases, except for the fact that they should not have the disease. Usually, the control group consists of persons who have not been exposed to the disease or factor whose influence is under study. Therefore, among the options, the best answer would be the children coming to the OPD, who do not have the disease as controls for the study.

Case-control studies are also called retrospective studies. This study proceeds from effect to cause and is done after the exposure and outcome have occurred.

They focus on diseases or health problems that have already developed. They are less time-consuming and have no risk of attrition because they do not require follow-up.

There are four basic steps in conducting a case-control study:

• Selection of cases and controls.

• Matching- This is done to ensure comparability between cases and controls. Controls are selected such that they are similar to cases with respect to certain factors, like age, that influence the disease outcome.

• Measurement of exposure - This is obtained by interviews, questionnaires, or study of the past records of cases.

• Analysis in the form of exposure rates, risk ratio, and odds ratio.

The Odds Ratio (OR) calculated from a case-control study, is a measure of the strength of the association between risk factor and outcome.

Solution to Question 12:

She can use the pentavalent vaccine and discard the MR vaccine. According to the open vial policy, the pentavalent vaccine can be reused, whereas MR can not be reused.

Open vial policy was implemented to allow the reuse of partially used multidose vials under the Universal Immunization Program (UIP) for up to 28 days if certain conditions are met. This was implemented to reduce vaccine wastage.

It applies only to DPT, TT, hepatitis B, oral polio vaccine (OPV), liquid pentavalent, PCV, injectable IPV, inactivated Japanese encephalitis (JE) vaccine, and rotavirus vaccine. It does not apply to Measles/MR and BCG vaccines.

Note: Open vial policy does not apply to rotavirus vaccine with VVM on cap, it applies only to new rotavirus vaccine with VVM on label.

Community Medicine NEET 2023

Question 1:

In a village, it is observed that several farmers have crossed gait and use a stick for support to stand up and walk. Due to poor yield from farms, they consume meals containing rice and pulses only. Supplementing their diet with which of the following vitamins could have prevented this?

a) Vitamin A

b) Vitamin D

c) Vitamin C

d) Vitamin B

Question 2:

A 35-year-old homeless man presented with a 1-month history of fever, cough, and weight loss. Both sputum smears turned out to be negative, but the chest x-ray ordered was suggestive of tuberculosis. According to the recent NTEP guidelines, which is the next best line of management?

a) Repeat sputum smears

b) Ask for CBNAAT

c) Ask for line probe assay

d) Wait until TB culture results to start ATT

Question 3:

Research is being conducted to find the association between aniline dye exposure and bladder cancer in workers who have worked in the industry for >20 years. Two groups were formed: one directly involved with dye handling and the other group consisting of office clerks not directly exposed to the dye. Years of occupation were noted from records. What type of study is being performed?

a) Retrospective cohort study

b) Prospective cohort study

c) Case-control study

d) Intervention and response

Question 4:

The blood pressure of a population was tracked from childhood to adulthood. It was observed that those who had lower BP in childhood had low BP in adulthood, while those who had higher BP in childhood had high BP in adulthood. This can be best described as____

a) Rule of halves

b) Tracking of blood pressure

c) STEPwise approach

d) Primordial approach

Question 5:

A 30-week primigravida c/o reduced vision at night. She has been avoiding papaya, mango, and other fruits throughout her pregnancy as she thinks they could be abortifacients. It is the primary duty of which of the following workers to provide counseling and information to the patient?

a) ANM

b) AWW

c) Trained birth attendant

d) ASHA

Question 6:

A 22-year-old female comes to the STI clinic with minimal vaginal discharge. On speculum examination, erosions are seen on the cervix. Which of the following kit should be given to this patient?

a) Green

b) Red

c) Gray

d) Yellow

Question 7:

The years of potential life lost could be attributed to_____.

a) Years lost to morbidity

b) Years lost due to premature death

c) Years lost to disability

d) Years lost to poor quality of life

Question 8:

In a 10-year-old school child, which of the following vaccines is given as a part of the school immunization program?

a) Measles vaccine

b) Rotavirus vaccine

c) TT/Td vaccine

d) Hepatitis B vaccine

Question 9:

Which of the following statements is true about cancer treatment according to the Colombo plan?

a) Help with PET scan units for diagnosis of cancer

b) Human resource strengthening

c) Setting up chemotherapy units

d) Setting up cobalt therapy units

Question 10:

A poor farmer with a history of successive crop failure develops progressive spastic paraparesis, signs of upper motor neuron paralysis, and gait instability. Name the toxin responsible for this condition.

a) Aflatoxin

b) Beta-oxalyl-amino-alanine

c) Ergot alkaloids

d) Fusarium toxin

Question 11:

A young male came to the hospital with a clean-cut wound without any bleeding. The patient received a full course of tetanus vaccination 10 years ago. What is the best management for this patient?

a) Human tetanus immunoglobulin and full course of vaccine

b) Human tetanus immunoglobulin only

c) Single-dose tetanus toxoid

d) Full course tetanus toxoid

Question 12:

How is water collected for bacteriological examination during a disease outbreak?

a) Collect water from already leaking taps

b) Before collecting, let water flow for at least 1 minute

c) Water sample container is kept close to the tap avoid spillage

d) Collect from a gentle stream of water to avoid splashing

Question 13:

The average daily dietary nutrient intake level sufficient to meet the nutrient requirements of nearly all (97-98%) healthy individuals in a particular life stage and gender group is known as _____.

a) Adequate intake

b) Dietary goal

c) Estimated average requirement

d) Recommended dietary allowance

Question 14:

A boys' hostel has an outbreak of fever cases with headache, followed by the development of pleomorphic rashes sparing palms and soles. What is the next best step in the management of suspected cases?

a) Isolate for 6 days after giving acyclovir, followed by VZIG within 72 hours of exposure

b) Isolate for 12 days after giving acyclovir, followed by VZIG within 48 hours of exposure

c) Isolate for 6 days

d) Only give VZIG

Question 15:

Which of the following is the sensitive indicator to assess the availability, utilization, and effectiveness of healthcare in a community?

a) Infant mortality rate

b) Maternal mortality rate

c) Immunization coverage

d) Disability-adjusted life years

Answer Key

Question No.	Correct Option
1	c
2	b
3	a
4	b
5	d
6	c
7	b
8	c
9	d
10	b
11	c
12	d
13	d
14	c
15	a

Detailed Explanations

Solution to Question 1:

The likely diagnosis based on the given clinical scenario is neurolathyrism. Administration of vitamin C at doses of 500-1000 mg is beneficial for this.

Lathyrism is a paralyzing disease of humans and animals. It is due to the consumption of khesari dal or Lathyrus sativus. When the husk is removed, it looks similar to red gram or Bengal gram. It is mostly consumed by poor agricultural laborers because it is relatively cheap. Lathyrism is more prevalent in parts of Madhya Pradesh, Uttar Pradesh, Odisha, and Bihar. The Prevention of Food Adulteration Act in India has banned Lathyrus sativus in all forms (whole/ split/ flour).

In humans, it is referred to as neurolathyrism because it affects the nervous system. It is characterized by gradually developing spastic paralysis of the lower limbs. Beta-oxalyl amino alanine (BOAA) is the toxic compound causing this disease. It is water-soluble and can be removed by the following methods:

• Soaking pulses in hot water and discarding the water

• Overnight soaking in lime water, followed by boiling

• Parboiling - for large-scale detoxification

Solution to Question 2:

The given case with the classical symptoms of evening rise of temperature and productive cough of more than 2 weeks in duration prompts a diagnosis of pulmonary tuberculosis. According to the NTEP guidelines, the next best line of management is CBNAAT.

The initial diagnosis of pulmonary TB is done with sputum smear examination and chest radiography. Chest radiography features suggestive of TB include upper lobe infiltrates and cavities. However, these are not very specific to TB. The presence of these features should prompt microbiological evaluation for Mycobacterium tuberculosis.

A positive smear examination, irrespective of CXR findings, is considered as microbiologically positive with sensitivity status unknown. Drug sensitivity status via NAAT is to be done following a sputum smear before initiation of treatment. This helps in the early detection and treatment of drug-resistant TB. This has been integrated into the regular TB services to detect rifampicin resistance at the time of diagnosis. Depending on the sensitivity of the tubercle bacilli, appropriate ATT is started.

In cases of high clinical suspicion, positive CXR findings, or in people with HIV, a CB-NAAT is done. Two sputum samples are sent for cartridge-based nucleic acid amplification (CBNAAT), which detects the presence of Mycobacterium DNA along with resistance against rifampicin within 2 hours. It analyses sputum or any other biological specimen (except blood and blood-contaminated specimens) using real-time PCR technology.

The interpretation of CBNAAT is as follows:

• If the CBNAAT turns out to be positive/MTB detected, the 2nd sample is sent for culture and drug sensitivity testing (C&DST) immediately.

• If the CBNAAT turns out to be negative for rifampicin resistance, a first-line line probe assay is ordered, to look for isoniazid resistance.

• If FL-LPA shows no resistance to isoniazid, the patient is treated as drug-sensitive, and a conventional daily regime of 2HRZE and 4HRE is given.

• If FL-LPA shows resistance to isoniazid, reflex testing is done using second-line line probe assay (SL-LPA).

• If SL-LPA turns out to be negative, treat as isoniazid or H-mono-/poly-resistance

• If SL-LPA turns out to be resistant to some drugs, modify the treatment regime accordingly.

• If CBNAAT is positive for rifampicin resistance, the second sample is sent for first-line line probe assay (FL-LPA), second-line line probe assay (SL-LPA), and BACTEC MGIT 960 to look for isoniazid (H) mono-/poly-resistance

• If there is no resistance/H-mono-/poly- resistance, treat as shorter oral bedaquiline-containing MDR regime.

• If there is resistance, then treat as longer oral bedaquiline-containing MDR regime.

• If the 2nd sample also turns out to be indeterminate, a fresh sample is taken for LPA or liquid culture, to ascertain drug resistance. Meanwhile, treatment initiation is done and modified subsequently, as per results.

Other options:

Option A: Repeat sputum smears are not the next best line of management in this case, as the patient already had two negative sputum smears and a chest x-ray suggestive of tuberculosis.

Option C: Line probe assay would not be the immediate next step in management. First, CBNAAT should be performed to detect Mycobacterium tuberculosis and identify rifampicin resistance.

Option D: Waiting for TB culture results to start ATT is not the best management strategy, as it can take several weeks for the results to come back, and the patient may deteriorate in the meantime.

Solution to Question 3:

The study given here is a retrospective cohort study.

A retrospective cohort study is when both the exposure and outcome have occurred before the investigations and study is undertaken.

In this scenario, the workers in the industry were divided into two groups, viz those who are directly exposed and the clerical staff who are not directly exposed to the suspected carcinogen. The incidence of carcinogenesis among these two groups is studied going backward tracking their records.

An increased risk of bladder cancer was observed among long-term workers directly exposed to the chemical in aniline dye industries when compared with the other study group. It is also called a historical cohort study, a prospective study in retrospect, or a non-concurrent prospective study. The cohort is identified from past records and is assessed to date for the outcome. The same cohort is followed up prospectively in the future for further assessment of the outcome.

Other options:

Option B - Prospective cohort study: This type of study involves following a group of participants over time and monitoring their exposure to a risk factor and the development of an outcome. In this case, the outcome (bladder cancer) and exposure (aniline dye) have already occurred, making this a retrospective cohort study.

Option C - Case-control study: A case-control study involves comparing individuals with a specific outcome (cases) to those without the outcome (controls) and assessing their exposure history. This study design is not being used in the given scenario, as the workers are divided into two groups based on their exposure status and followed for the development of bladder cancer.

Option D - Intervention and response: This option is not a type of epidemiological study design. It could refer to an intervention study, in which an intervention is applied to a group of participants, and the response is measured. However, this is not applicable to the current scenario.

Solution to Question 4:

The above phenomenon is best described as the "tracking" of blood pressure.

In this phenomenon, the blood pressure of individuals is followed up from early childhood into adulthood. The BP in individuals with low BP levels tend to remain low, and in those with high BP levels, they tend to remain high. This phenomenon of persistence of rank order of blood pressure has been described as "tracking".

Other options:

Option A: The rule of halves depicts the disease burden of hypertension in the community. It states that only about half of the hypertensive subjects in the general population are aware of the condition. Only half of those aware are being treated and only half of those being treated are considered to be adequately treated.

A pictorial representation of the rule of halves in hypertension is shown below:



Option B: The WHO STEPwise approach to non-communicable disease (NCD) risk factor surveillance (STEPS) is a simple, standardized method for collecting, analyzing, and disseminating

data on key NCD risk factors such as tobacco use, alcohol use, physical inactivity, unhealthy diet, overweight and obesity, raised blood pressure, raised blood glucose, and abnormal blood lipids.

Solution to Question 5:

Amongst the given options, it is the duty of the ASHA to provide counseling and information to this pregnant patient

An ASHA is a trained female community health activist and a resident woman of the village with a formal education of at least up to 10th grade. She should preferably be in the age group of 25 to 45 years (married/divorced/widow) and have qualities such as good communication skills and leadership skills. The general norm of selection recommends having one ASHA for 1000 population, but in tribal, hilly, and desert areas, the norm could be relaxed to one ASHA per habitation.

Roles of ASHA:

• To create awareness and provide information to the community on determinants of health such as nutrition, basic sanitation, and hygienic practices

• To counsel women on birth preparedness, the importance of safe delivery, breast-feeding and complementary feeding, immunization, and contraception

• To counsel women on the prevention of common infections including reproductive tract infection/sexually transmitted infection

• To mobilize the community and help them in accessing health and health-related services.

• To arrange and escort pregnant women and children requiring treatment/admission to the nearest pre-identified health facility

• To inform sub-center/primary health center about the births and deaths in her village and any unusual health problems/disease outbreaks in the community

• To promote the construction of household toilets under the total sanitation campaign

Other Options:

Option A (ANM): An Auxiliary Nurse Midwife (ANM) is a trained health worker who provides basic healthcare services, including antenatal care, delivery, and postnatal care. Although an ANM can provide counseling and information to pregnant women, the primary responsibility for this role lies with the ASHA worker.

Option B (AWW): An Anganwadi Worker (AWW) is responsible for providing basic health care, nutrition, and preschool education to children under the age of six, pregnant women, and lactating mothers. While AWWs play a vital role in maternal and child health, counseling pregnant women on health-related issues is not their primary responsibility.

Option C (Trained birth attendant): A trained birth attendant is responsible for assisting with childbirth and providing postpartum care. Although they may provide some counseling and information, this role is primarily focused on the delivery process rather than ongoing counseling throughout pregnancy.

Solution to Question 6:

This patient with vaginal discharge and erosions on the cervix has cervicitis from a sexually transmitted infection. She should be given the gray kit which contains cefixime and azithromycin.

Under the STD control program, pre-packed color-coded kits have been provided for free supply to treat sexually transmitted (STI) or reproductive tract infections (RTI). Management of STDs under the syndromic approach is based on specific signs and symptoms and is not dependent on laboratory investigations.

A patient with just vaginal discharge is presumed to have vaginitis and is given a green kit that contains secnidazole and fluconazole. But this patient also has cervical erosion, which is a sign of cervicitis. She should be treated with the grey kit. This kit contains a single tablet of 1 gram azithromycin for chlamydia and a single tablet of 400 mg cefixime for gonorrhea.

Solution to Question 7:

The years of potential life lost could be attributed to the years lost due to premature death.

It is a mortality indicator that shows the number of years that a dying person could have expected to survive (usually taken as 75 years). Mortality indicators are used as the starting point in the evaluation of health status. They are used as indirect indicators of health.

The concept of years of potential life lost (YPLL) is used to estimate the societal impact of premature death. It helps prioritize public health interventions and allocate resources to address significant health problems. YPLL, along with other metrics like DALYs and QALYs, aids in assessing the burden of disease and the effectiveness of healthcare interventions. These measurements can help inform policymakers and healthcare providers in making informed decisions about healthcare delivery and resource allocation.

Strategies to reduce YPLL include prevention and early intervention, improving access to healthcare, and addressing social determinants of health that contribute to health disparities and premature death.

Other options:

Option A: Years lost to morbidity refers to the time spent living with a disease or disability, which is not directly related to the concept of years of potential life lost.

Option C: Years lost to disability is a component of disability-adjusted life years (DALY). DALY indicates the number of years lost due to ill health, disability, or early death.

DALY = YLL +YLD

• Years lost to disability (YLD): Here, the number of cases due to injury and illness is multiplied by the average duration of the disease and a weighting factor. (Severity of the disease on a scale of 0 (perfect health) to 1 (dead))

• Years of lost life (YLL) - Here, the number of deaths at each age is multiplied by the expected remaining years of life. (according to a global standard of life expectancy)

Option D: Quality-adjusted life year (QALY) indicates how the health status of an individual impacts his quality and quantity of life. It shows the years of life lost due to injury or illness. It is

commonly used in evaluating the outcomes of medical interventions.

Solution to Question 8:

A ten-year-old school child will be administered the TT/Td vaccine, as part of the school immunization program, as per the National Immunization Schedule.

Tetanus and adult diphtheria (Td) vaccine is a combination of tetanus and diphtheria antigens, with a lower concentration of diphtheria antigen (d), as recommended for older children and adults. The MoHFW recommends the replacement of the TT vaccine with Td vaccine for all age groups, including pregnant women. A dose of 0.5 mL is administered in the intramuscular route in the upper arm.

Scheduled to be administered at 10 and 16 years of age.

For pregnant women:

• Td-1: early in pregnancy

• Td-2: 4 weeks after Td-1

• Td (Booster): if pregnancy occurs within 3 years of the last pregnancy and 2 Td doses were received.

Other options:

Option A - According to the National Immunization Schedule, Measles/MR vaccine is given at 9 completed months (1st dose), and a second dose is given at 16-24 months. Measles vaccine can be given till 5 years of age. A dose of 0.5 mL is administered subcutaneously in the right upper arm.

Option B - Rotavirus vaccines are oral, live attenuated vaccines and are of 3 types: Rotavac, RotaTeq, and Rotarix. Rotavac (monovalent liquid frozen) vaccine is administered in 3 oral doses, 4 weeks apart at 6, 10, and 14 weeks. Though CDC guidelines recommend that it should not be administered to children more than 8 months of age, the National Immunisation Guidelines in India mention that the first dose of rotavirus vaccine can be administered up to the age of 1 year.

Option D - Hepatitis B vaccine is an example of a subunit vaccine made using HBsAg. The first dose (0.5 mL, intramuscular) is given at birth or as early as possible within 24 hours on the anterolateral side of mid-thigh. Subsequent doses are given as a part of the pentavalent vaccine at 6 weeks, 10 weeks, and 14 weeks (can be given until one year of age).

Solution to Question 9:

Aid to set up cobalt therapy units for cancer treatment at medical institutions in India was a part of the Colombo plan.

The Colombo Plan is a regional organization that upholds the concept of a collective intergovernmental effort to strengthen the economic and social development of member countries in the Asia-Pacific region.

It was formed at a meeting of the commonwealth foreign ministers with the headquarters at Colombo in January 1950. It comprises 20 developing countries within the region and 6 non-regional members - Australia, Canada, Japan, New Zealand, the UK, and USA. It involves foreign aid and technical assistance for the economic and social development of the region.

The contribution of Canada in supplying cobalt therapy units to medical institutions in India was an important item of aid under the Colombo plan. AIIMS New Delhi was established with financial assistance from New Zealand under the Colombo plan. Cobalt therapy is a form of radiotherapy that uses cobalt-60, a radioactive isotope, to generate high-energy gamma rays for treating cancer. Cobalt therapy units were an important tool for cancer treatment in the mid-20th century, and their provision under the Colombo plan significantly improved cancer treatment capabilities in India.

The Colombo Plan has evolved over the years and now focuses on a broader range of development goals, including education, public administration, agriculture, and industry. This reflects the changing needs of the Asia-Pacific region and the importance of a comprehensive approach to development.

Other Options:

Option A (Help with PET scan units for diagnosis of cancer): While PET scan units are important for cancer diagnosis, they were not part of the Colombo plan's cancer treatment initiatives in India.

Option B (Human resource strengthening): Although the Colombo plan aims to improve the economic and social development of member countries, human resource strengthening in the context of cancer treatment is not explicitly mentioned as part of the plan.

Option C (Setting up chemotherapy units): The Colombo plan's focus in cancer treatment was on providing aid for cobalt therapy units, not chemotherapy units.

Solution to Question 10:

The diagnosis based on the given clinical scenario is neurolathyrism, and the toxin responsible for the condition is beta-oxalyl-amino-alanine.

Lathyrism is a paralyzing disease of humans and animals. It is due to the consumption of Khesari dal/Lathyrus sativus. It is mostly consumed by agricultural laborers of poor economic status because it is relatively cheap. It is more prevalent in parts of Madhya Pradesh, Uttar Pradesh, Odisha, and Bihar.

In humans, it is referred to as neurolathyrism because it affects the nervous system. It is characterized by gradually developing spastic paralysis of the lower limbs. In animals, it is referred to as osteolathyrism as it leads to skeletal deformities. It usually affects young males aged 15-45 years and it is manifested in stages:

- Latent stage
- No-stick stage
- One-stick stage

• Two-stick stage

• Crawler stage

Beta-oxalyl-amino-alanine (BOAA) is the toxic compound causing lathyrism. The water-soluble property of the toxin is used in removing it from the pulses by soaking it in hot water and rejecting the soaked water. Vitamin C in doses of 500-1000 mg can be used for the prevention and treatment of lathyrism.

Other options:

Option A: Aflatoxins are produced by certain fungi, such as Aspergillus flavus and Aspergillus parasiticus, are the most potent hepatotoxins. These fungi can contaminate food grains such as groundnut, maize, parboiled rice, sorghum, wheat, and tapioca. It is mainly due to improper storage. It can be prevented by proper storage after drying. Moisture content should be kept below 10%. They do not cause the neurological symptoms described in the question.

Option B: Ergot alkaloids are produced by ergot or Claviceps purpurea, a field fungus. Food grains like bajra, rye, sorghum, and wheat have a tendency to get contaminated with this ergot fungus. These get harvested along with the food grains and consumption of this ergot-infested grain leads to ergotism.

Acute cases present with nausea, vomiting, giddiness, and drowsiness for a period of up to 24 to 48 hours after the ingestion of ergot grain. In cases of chronic consumption, they present with painful cramps in limbs, and peripheral gangrene due to vasoconstriction of capillaries. However, they do not cause the progressive spastic paraparesis and upper motor neuron paralysis characteristic of neurolathyrism. Ergot-infested grains are removed by floating them in 20% saltwater. Handpicking can also be done.

Option D: Fusarium toxin is produced by the fungus Fusarium incarnatum. They are known to contaminate sorghum and rice.

Solution to Question 11:

The patient with a clean-cut wound without any bleeding, who received a full course of tetanus vaccination 10 years ago, should be given a single dose of tetanus toxoid vaccine and appropriate wound care.

Wound management involves cleaning the wound thoroughly to remove soil, foreign bodies, and necrotic tissue.

Note: In cases where combined immunization is necessary, the patient is given 250 units of human tetanus immunoglobulin in one arm and 0.5 mL of adsorbed tetanus toxoid (PTAP or APT) in the other arm or gluteal region. The antitoxin is given for immediate temporary protection, and the tetanus toxoid is given for long-lasting protection.

Solution to Question 12:

During a disease outbreak, for collection of a water sample for bacteriological examination from a tap in regular use, water is collected from a gentle stream of water to avoid splashing.

If water samples are to be collected from a tap that is not regularly used, they should be first sterilized by heating. Then, the tap is cooled and opened fully and the initial water stream is let out to flow before the sample is collected.

The sample bottle must be held near the base with a hand and the stopper and paper cover over it are removed together and held in the fingers.

Other options:

Option A (Collect water from already leaking taps): Collecting water from leaking taps is not recommended as it increases the risk of contamination and may not provide an accurate representation of water quality.

Option B (Before collecting, let water flow for at least 1 minute): This option is close to the correct answer but falls short, as it is recommended to let the water flow for at least two minutes to flush out stagnant water in the pipe.

Option C (Water sample container is kept close to the tap to avoid spillage): While keeping the sample container close to the tap is a good practice to avoid spillage, it is not the main concern during the collection of water samples for bacteriological examination.

Solution to Question 13:

The average daily dietary nutrient intake level sufficient to meet the nutrient requirements of nearly all healthy individuals in a particular life stage and gender group is known as recommended dietary allowance (RDA).

Recommended dietary allowances (RDAs) are developed by the Food and Nutrition Board of the Institute of Medicine, National Academy of Sciences, and are part of the Dietary Reference Intakes (DRIs). DRIs include not only RDAs but also adequate intakes (AIs), tolerable upper intake levels (ULs), and estimated average requirements (EARs).

RDAs are essential for planning and assessing diets for healthy individuals and serve as the basis for food labeling, nutrition policy, and dietary guidelines. Meeting the RDA for a specific nutrient helps ensure that an individual's nutrient intake is sufficient to prevent deficiency and maintain optimal health. However, individual nutrient needs may vary depending on factors such as age, gender, health status, and activity level.

Other options

Option A: Adequate intake is the recommended average daily intake level which is based on experimentally found estimates of nutrient intake by a group of apparently healthy population (assumed to be adequate). It is used when RDA cannot be determined. It is referred to as acceptable intake in the indian context.

Option B: Dietary goals are country-specific quantitative targets for selected nutrients aimed at preventing long-term chronic diseases. They are not directly focused on meeting the nutrient requirements of individuals but rather focus on population-level nutritional health.

Option C: The estimated average requirement (EAR) is the daily nutrient intake level estimated to meet the requirements of half the healthy individuals in a specific life stage and gender group. It is used to calculate the RDA but is not the same as the RDA.

The RDA is calculated as the estimated average requirement (EAR) plus twice the standard deviation (SD).

RDA = EAR + (2 × SD)

Solution to Question 14:

The scenario describes an outbreak of fever cases with headache followed by the development of pleomorphic rashes sparing palms and soles, which points to the diagnosis of chickenpox. In a boys' hostel, it is reasonable to assume that the majority of the affected individuals are not in the high-risk groups for severe varicella infection. For the management of suspected cases in this situation, isolation for 6 days from the onset of the rash is the appropriate measure to prevent further spread of the infection. Acyclovir and VZIG are not necessary for healthy individuals.

Varicella zoster immunoglobulin (VZIG) is not indicated for everyone. VZIG is a passive immunization used as post-exposure prophylaxis for individuals at high risk of developing severe varicella (chickenpox) infection after being exposed to the varicella-zoster virus. It should be administered within 72 hours of exposure to provide the best protection, although it may still be effective if given up to 10 days after exposure.

High-risk individuals who may benefit from VZIG include

• Immunocompromised individuals, such as those receiving immunosuppressive therapy, those with congenital cellular immunodeficiency, and those with acquired immunodeficiency, including HIV/AIDS.

• Pregnant women who have been exposed to the virus and have no history of chickenpox or vaccination, as varicella infection during pregnancy can lead to congenital varicella syndrome or severe neonatal varicella infection.

• Newborns whose mothers developed chickenpox within five days before delivery or within 48 hours after delivery, as they are at risk of severe neonatal varicella infection.

For healthy individuals who have been exposed to the varicella-zoster virus, the varicella vaccine is usually recommended as post-exposure prophylaxis, rather than VZIG. The vaccine should be administered within 3 to 5 days of exposure to help prevent or reduce the severity of the disease.

Live attenuated vaccine is administered to children who have not had chickenpox between the age of 12 to 18 months.

The below image shows multiple vesicular lesions on the trunk is suggestive of chickenpox.

Solution to Question 15:

The infant mortality rate (IMR) is a sensitive indicator of the availability, utilization, and effectiveness of health care in a community.

Infant mortality rate (IMR) = Number of deaths in the first year of life/Total number of live births x 1000.

It is one of the universally accepted healthcare indicators for the whole population and the socioeconomic conditions under which they live.

Other options:

Option B - Maternal mortality rate: While maternal mortality rate is an important indicator of maternal healthcare services, it is not as sensitive as the infant mortality rate in reflecting the availability, utilization, and effectiveness of healthcare in a community.

Option C - Immunization coverage: Immunization coverage is an essential factor in preventing diseases and promoting public health. However, it is a narrower indicator than infant mortality rate, focusing primarily on preventive healthcare services for children. Full immunization coverage is defined as the proportion of infants 9-11 months old who have received the 3 doses of DPT vaccine, BCG vaccine, 3 doses of OPV, and one dose of measles vaccine against the estimated number of infants during a specific year.

Option D - Disability-adjusted life years: DALYs are a useful measure of the overall burden of disease in a population, but they are less sensitive in assessing the availability, utilization, and effectiveness of healthcare services in a community compared to infant mortality rate.

Community Medicine INI-CET 2023

Question 1:

All of the following statements are true for descriptors of central tendency except:

a) Mode is affected by extreme values

b) Mean is affected by extreme values

c) Median is calculated by overall observation

d) Mode can be used as a qualitative data descriptor

Question 2:

The formula given below corresponds to:



a) Standard deviation

b) Variance

c) Mean

d) Median

Question 3:

All of the following may be used to find the normalcy of data except:

a) Plot histogram

b) Mantel-Haenszel test

c) Box whisker

d) Shapiro Wilk test

Question 4:

In Kaplan Meier survival curve, the bands used for the confidence interval is?

a) Unequal precision band

b) Hirshberg band

c) Holmes band

d) Hall-Wellner Band

Question 5:

30,000 women were followed up for 10 years for the development of breast cancer. 1200 women developed cancer and were given a questionnaire to assess possible risk factors. Additionally, 2000 women from the same study were used as controls and were also given the questionnaire. What is this type of study called?

a) Case cohort

b) Retrospective cohort

c) Nested case control

d) Cross control cohort

Question 6:

A person had a dog bite. Vaccination is up to date for both the dog and the man. The person's history is significant for having undergone splenectomy five years ago. What is the best management in this case?

a) Observation

b) Amoxiclav

c) Metronidazole

d) Ciprofloxacin

Question 7:

A child came for DPT vaccination at 10 weeks. He has a history of fever of more than 40°C and an inconsolable cry at 6 weeks of age following vaccination. What should be done next?

a) Avoid DPT vaccination

b) Give DT vaccine

c) Give DPT vaccine

d) Defer by 4 weeks

Question 8:

High fat intake according to FDA is?

a) Diet should contain 400 kcal from carbohydrates

b) Diet should contain total 1000 kcal

c) Diet should contain 800 kcal from fat

d) Diet should contain 80% fats

Question 9:

The slogan coined by WHO 'Do it Right' in the first aid of a snake bite includes all of the following except?

a) Reassurance

b) Tell sequence of events in detail after the bite

c) Go to hospital

d) Incisions

Question 10:

Calculate the kappa statistics for the following data set comparing IGRA and Mantoux test.

a) 0.4

b) 0.6

c) 0.8

d) 0.5

Question 11:

Which of the following is not correctly matched?

a) Observational study — MOOSE

b) Systematic review — PRISMA

c) Diagnostic Study - CONSORT

d) Case report — CARE

Question 12:

Which of the following best represents the definition of culture?

a) It is static and not related to health and disease

b) A set of beliefs which are static

c) Socially acquired learned behaviour

d) Rules set according to religion which are dynamic

Question 13:

According to National Programme for Control of Blindness and Visual Impairment which of the following is incorrect about the reach in approach?

a) People with cataract are treated in a makeshift operation centre

b) IOL placement by small incision cataract surgery is preferred

c) Reach in approach has lesser post operative complications

d) People with cataract are identified and sent to higher centre

Question 14:

A 68-year-old lady was brought to the hospital by her daughter, with episodes of falls, confusion, chronic diarrhea, and cough. The intern decided to conduct a geriatric assessment. Which of the following factors indicated such an examination?

a) Age, sex, falls

b) Age, cough, chronic diarrhoea

c) Age, fall, confusion

d) Age, sex, cough, chronic diarrhea

Question 15:

Which of the following best helps assess the quality of healthcare provided?

a) Multiple logistic regression model

b) Bar graph

c) Scattergram

d) Ishikawa diagram/ Fish bone diagram

Question 16:

A mother brought her 6 month old infant baby to an Anganwadi worker Lata for advice regarding the vaccination of her child. The last vaccines he had received were OPV3, PENTA 3, IPV2, and ROTA3 at 14 weeks of age. BCG and Hep B were given at birth. When is the next vaccination due?

a) MR vaccine at left arm intramuscularly and Vitamin A drops at the end of 9 months

b) MR vaccine at right arm subcutaneously and Vitamin A drops at the end of 8 months

c) MR vaccine at left arm intramuscularly and Vitamin A drops at the end of 8 months

d) MR vaccine at right arm subcutaneously and Vitamin A drops at the end of 9 months

Question 17:

Which of the following statements is true about palliative care?

a) A,B,C,D

b) C,E,F

c) B,D

d) A,B,E,F

Question 18:

Which of the following is the highest level of community participation?

a) Considering their ideas

b) Felt needs of community considered

c) Contribution of resources

d) Involving in planning of health related campaigns

Question 19:

A patient hailing from an area infested with mosquitoes presents with fever, chills, and myalgias. WBC count was 4000 cells/cu.mm, RBC was normal, and platelets were 90000 cells/cu.mm. He also had a maculopapular rash. What is the likely diagnosis?

a) Dengue

b) Typhoid

c) Ebola haemorrhagic fever

d) Rift valley fever

Question 20:

Which of the following statements are correctly matched?

a) Statement 1,2

b) Statement 3,4

c) Statement 2,3

d) Statement 1,2,3

Question 21:

Two doctors examined 100 chest X - Rays of patients. The kappa statistic was found to be 0.71. What does it mean?

a) Good agreement

b) Perfect agreement

c) Poor agreement

d) No agreement

Question 22:

Which of the following is an example of a longitudinal observational and analytical study?

a) Case control Study

b) Cohort Study

c) Randomized Controlled Trial

d) Ecological Study

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	d
5	c
6	b
7	c
8	b
9	d
10	a
11	c
12	c
13	a
14	c
15	d
16	d
17	c
18	d
19	a
20	c
21	a
22	b

Detailed Explanations

Solution to Question 1:

Mode is not affected by extreme values in a given set of data.

The mean of a given set of data is responsive to each and every value in the distribution. Thus, the mean is the measure of central tendency which is maximally affected by extreme values (option B). It is usually an inappropriate measure for describing skewed distributions. However, repeated samples from the same population will always tend to have similar means. Mean is able to resist fluctuations between different samples drawn from the same population better than mode or median.

To obtain the median of a given set of data, the values are first arranged in ascending or descending order of magnitude, and then the value of the middle observation is noted. Since the data has to be first arranged in order before the median can be determined, an overall observation is necessary for its calculation (option C).

Mode is the observed value in the frequency distribution that occurs with the greatest frequency. It is completely uninfluenced by extreme values in the distribution. The mode can be used to describe qualitative data (option D).

Solution to Question 2:

The formula given above is used to calculate the standard deviation.

Standard deviation (or σ) is a measure of how dispersed the data is in relation to the mean. Data clustered around the mean have a low standard deviation, while data that are more spread out have a high standard deviation. A standard deviation close to zero indicates that data points are close to the mean, whereas a high or low standard deviation indicates data points are respectively above or below the mean.

Variance (option B) is the mean of the squares of all the deviation scores of the distribution. It is numerically equal to the square of the standard deviation.

Mean (option C) is obtained by adding together the individual observations and then dividing by the total number of observations.

Median (option D) divides the frequency distribution in half when all the scores are listed in order. When a distribution has an odd number of elements, the median is the middle value. When the distribution has an even number of elements, the median is the average of the two middle scores.

Solution to Question 3:

Mantel-Haenszel method is used to calculate adjusted odds ratio during stratified analysis. It is not a test to detect normalcy of data.

A plot histogram and a box whisker plot are visual methods to assess if a given set of data is distributed normally. A normal distribution curve is an example of a plot histogram.

Sometimes just the visual representation of the data might not be enough to tell whether the data is normally distributed. This is where Shapiro-Wilk test comes in. It examines how closely a given set of data fits a normal distribution. The sample data is first ordered and standardized. An assessment is then made how the sample quantiles fit the standard normal quantiles.

Shapiro-Wilk test is commonly used on small samples.

Stratified analysis of the data is done to reduce confounding. In such a scenario, multiple odds ratios are obtained — one for each stratum of the data. These have to be combined to give an adjusted odds ratio. This represents the effect of the exposure adjusted for the confounding factor. Mantel-Haenszel method is used for this purpose.

Solution to Question 4:

The bands used for confidence interval in the Kaplan-Meier survival curve are Hall-Wellner bands.

When using standard statistical software, at each point in time, the upper and lower limits of the confidence interval are also obtained. Hall-Wellner bands, Gill bands, and Nair's Equal Precision confidence bands are used for this purpose.

Kaplan-Meier plot is a graphical representation of the number of people surviving during a given length of time. Often there are two or more groups represented in the plot, such as those treated versus those not treated. The graph often has a number of steps as the number of people surviving is a discrete variable. The image given below shows a Kaplan-Meier plot. The dotted lines represent the upper and lower limits of the 95% confidence interval.



Solution to Question 5:

The type of study described in the question is a nested case-control study.

A nested case-control study is essentially a case-control study that is present inside a cohort study. The cases are all the cohort members who developed the disease being studied at a given point in time during the follow-up period. However, each case is matched to one or more controls — individuals who did not develop the disease but are otherwise comparable to cases in terms of age and gender. Cases and controls are then compared in terms of their respective exposures to the

risk factor. Nested case-control studies thus represent an interim analysis of the exposure and outcome data of a cohort study.

A case-cohort study (option A) involves selecting a subset of the entire cohort at the beginning of the study. This sub-cohort acts as the control group for making comparisons with multiple case groups. For example, if a cohort study is being done to assess smoking as a risk factor for ischemic heart disease and lung cancer, the same sub-cohort can be used as the control group for both outcomes. If it was a nested case-control study, it would require two separate control groups. Case-cohort studies are more efficient than nested case-control studies when multiple outcomes are being studied.

In a retrospective cohort study (option B), the outcomes have already occurred before the study begins. The investigator has to analyze the history of prior exposures and the duration of such exposures. Even though the exposure and the outcome have already occurred, the investigator goes from the exposure → outcome in a retrospective cohort study. This is in contrast to case-control studies, where the investigator goes from outcome → exposure.

Cross control cohort (option D) is not a type of study design.

Solution to Question 6:

Amoxicillin-clavulanate (Amoxiclav) is the drug of choice in this patient.

Antibiotic prophylaxis is indicated in:

• High-risk animal bites: cat bites in any location and hand bites by any animals, including humans. Cat and human bites have a higher risk of infection versus dog bites. Moreover, hand bites have a risk of developing an infection in the closed spaces of the hand.

• High-risk patients: comorbidities such as diabetes, liver disease, and absence of functional spleen. Patients without a functional spleen are at an increased risk for overwhelming sepsis primarily due to Capnocytophaga species.

Pasteurella species is the most common isolate from cat and dog bites. Streptococci, S. aureus and Eikenella corrodens are found in human bites. Amoxicillin-clavulanate is the first-line drug of choice for prophylaxis. In patients with penicillin allergy, clindamycin plus doxycycline/co-trimoxazole/fluoroquinolone can be used.

Solution to Question 7:

The next step for this patient should be to administer DPT vaccine.

Events such as persistent inconsolable crying (more than 3 hours in duration), hyperpyrexia (fever > 40°C), hypotonic-hyporesponsive episodes within 48 hours of DTwP administration, and seizures, with or without fever, within 72 hours DTwP administration are considered precautions but not contraindications to future doses of DTwP. These events don't recur with the next dose and have been shown to have no permanent sequelae.

Absolute contraindications to any pertussis vaccination are a history of anaphylaxis or development of encephalopathy within 7 days following previous DTwP vaccination. In case of

anaphylaxis, further immunization with diphtheria, tetanus, and pertussis vaccines are contraindicated as it cannot be said with certainty which component caused anaphylaxis. For patients with a history of encephalopathy following vaccination, any pertussis vaccine is contraindicated, and only diphtheria and tetanus vaccines may be used.

Note: This answer might be controversial. The above answer is as per the IAP Guidebook on Immunization.

Solution to Question 8:

High fat intake according to FDA is a diet that should contain a total of 1000 kcal.

High fat diets have low carbohydrates, <50 grams/day. This equates to <250 kcal/day in the form of carbohydrates, not 400 kcal (option A). The percentage of calories from fat is >50% (option D), which would equate to about 500-600 kcal from fats (option C). The diet should contain a total of 800 - 1000 kcal.

Ketogenic diets mainly incorporate low carbohydrates (<50 grams/day), low protein, and high fats. Gluconeogenesis from the glucogenic amino acids in the diet and fat oxidation are utilized to meet the metabolic demands of the body. Ketogenic diets are especially useful for the treatment of intractable epilepsy.

Solution to Question 9:

Incisions are not a part of the management of a snake bite.

The first aid following a snake bite consists of the mnemonic: carry no right.

Carry: The victim is not allowed to walk, even for a short distance. Walking generates muscle-pumping activity which disperses the venom into the systemic circulation.

No: No tourniquet, no electrotherapy, no cutting, no pressure immobilization, no suction to the bite site

R: Reassure the patient since 70% of the bites are from non-venomous species.

I: Immobilize in the same way as a fractured limb. A splint can be applied to the bitten extremity to limit the movement and assist with positioning. The extremity should be maintained in a neutral position at approximately heart level.

GH: Get to a hospital immediately. The most important step is to rapidly transport the patient to a medical facility equipped to provide supportive management.

T: Tell the doctor of any systemic symptoms which may have appeared on the way to the hospital.

Note that elapid envenomations that are primarily neurotoxic and have no significant effects on local tissue may be managed by pressure immobilization. In this technique, the entire limb is immediately wrapped with a bandage and then immobilized. This technique restricts lymphatic drainage and has been shown to delay the absorption of venom into the systemic circulation. However, the offending snake should have been reliably identified as neurotoxic and the rescuer should be trained in applying this pressure bandage.

Solution to Question 10:

Kappa statistic is used to evaluate the agreement between IGRA and TST test results.

Kappa is calculated as:

$$\kappa = \frac{p_O - p_E}{1 - p_E}$$

where, pO is observed agreement and pE is expected agreement.

The table in the question has been redrawn with alphabets in the cells to better illustrate the formulae:

	IGRA (+)	IGRA (–)
Mantoux (+)	80(a)	40(b)
Mantoux (–)	20(c)	60(d)

In this question, the pO will be all the situations in which both IGRA and Mantoux tests are positive or negative — that is, both of them are in agreement with one another.

$$p_0 = \frac{a+d}{a+b+c+d} = \frac{140}{200} = 0.7$$

pE represents the probability of random agreement between the tests. It is calculated as a sum of both the tests being positive together at random and negative together at random.

$$p_E = p_{pos} + p_{neg}$$

$$p_{pos} = \frac{a+b}{a+b+c+d} \times \frac{a+c}{a+b+c+d}$$

$$p_{pos} = \frac{120}{200} \times \frac{100}{200} = 0.3$$

$$p_{neg} = \frac{c+d}{a+b+c+d} \times \frac{b+d}{a+b+c+d}$$

$$p_{neg} = \frac{80}{200} \times \frac{100}{200} = 0.2$$

$$p_E = 0.3 + 0.2 = 0.5$$

Substituting the calculated values in the formula for kappa:

$$K = \frac{0.7 - 0.5}{1 - 0.5} = \frac{0.2}{0.5} = 0.4$$

Strength of agreement varies with kappa coefficient as follows:

• 0.81 to 1: perfect agreement between the two tests/methods

• 0.61 to 0.80: substantial agreement

• 0.41 to 0.60: moderate agreement

• 0.21 to 0.40: fair agreement

• 0.10 to 0.20: slight agreement

Solution to Question 11:

The guidelines used for reporting the results of RCT are CONSORT (consolidated standards of reporting trials).

To improve the quality of reporting of health research, certain standards have been developed. These standards vary according to the type of study in question. Many journals now require the authors to complete the relevant checklist according to these standards prior to submitting their paper for publication. These standards include:

Many of these guidelines are available via the EQUATOR network.

Guideline	Expansion	Area
CONSORT	Consolidated standards of reporting trials	Randomized controlled trials
TREND	Transparent reporting of non-randomized designs	Behavioural and public health interventions with non-randomized designs

Guideline	Expansion	Area
STROBE	Strengthening the reporting of observational studies in epidemiology	Cohort, case-control, and cross-sectional studies
STRAEGA	Strengthening the reporting of genetic association studies	Modification of STROBE for genetic studies
STARD	Standards for the reporting of diagnostic accuracy studies	Diagnostic tests
CARE	Case reporting	Clinical case reports and reports of patient encounters
SAMPL	Statistical analyses and methods in published literature	Basic statistical reporting for articles published in biomedical journals
PRISMA	Preferred reporting items for systematic reviews and meta-analyses	Meta-analyses of randomized controlled trials
MOOSE	Meta-analysis of observational studies in epidemiology	Meta-analyses of observational studies

Solution to Question 12:

Culture is the socially acquired learned behaviour. It is passed onto generations through formal and informal learning. It is the representation of customs, beliefs, laws, arts, and moral values of a society. Cultural factors have a large part to play in health.

Other closely related terms include:

Customs are the traditionally accepted ways of behaving that are specific to a society. They can be folkways, which are referred to as the right ways of doing things in less vital areas of human conduct. Mores are stringent customs followed by a society.

Socialization is the process by which a person in a society imbibes its beliefs, customs, etc. through daily interaction with its members.

A community is a social group determined by geographical boundaries and/or common values and interests.

Acculturation is the process by which there is the diffusion of cultures between two culturally different groups of people. Acculturation does not take place by folklore.

Folklore is the passage of customs and beliefs of a community through generations, merely by word of mouth.

Solution to Question 13:

Patients with cataract are not operated in makeshift operation centres but are rather shifted to base hospitals (reach-in approach) as the visual acuity results indicate that the Base Hospital had significantly higher final corrected visual acuity compared to the Peripheral Eye Camps.

The Base Hospital approach, also known as the reach-in approach, and the Peripheral Eye Camp approach, also known as the reach-out approach, are two community-oriented strategies implemented within the framework of the National Programme for the Control of Blindness. These techniques aim to address the existing backlog of cataract-related blindness. Both methods have demonstrated efficacy and possess distinct advantages and disadvantages. Both community participation and intersectoral coordination, together with the utilization of suitable technology at an affordable price, are essential components for both endeavours.

Based on recent developments, there is a shift in emphasis from the 'Reach-out' approach to the 'Reach-in' approach. The Reach-in Approach involves the organization of screening camps in rural and remote locations that lack access to eye-care institutions. Comprehensive eye care treatments, including refraction, are offered to patients. Subsequently, those requiring cataract surgery are conveyed to the nearest hospitals that possess adequate resources and facilities, following the base hospital strategy. To conduct ocular surgery, it is imperative to comply with the guidelines established by the Government of India.

• It is imperative to have at least one anesthesiologist who is well prepared to address typical crises to ensure patient safety and optimal care.

• It is recommended that a minimum of one, ideally two, operating surgeons be present, with each surgeon limiting their daily caseload to no more than 30 procedures.

• Currently, the preferred approach for cataract extraction is extracapsular surgery, with a preference for manual small incision cataract surgery (SICS) or phacoemulsification, together with the implantation of a posterior chamber intraocular lens.

• It is ideal to limit the number of surgical procedures conducted in a day to no more than 50.

• Simultaneous surgery on both eyes should be avoided.

• Patients who have a high surgical risk due to conditions such as severe diabetes, severe hypertension, and cardiac issues should not undergo surgery in camp settings. Cases related to such issues should be appropriately referred to the base hospitals.

National Programme for the Control of Blindness and Visual Impairment (NPCBVI). The levels of eye care, according to NPCBVI are:

Primary level: Sub-district level hospitals, community health centres, mobile ophthalmic units, upgraded PHCs, link workers, and Panchayats.

Secondary level: District hospitals, NGO eye hospitals

Tertiary levels: Regional institutes of Ophthalmology, medical colleges.

The goal is to reduce the prevalence of blindness to 0.3% by 2020.

Solution to Question 14:

This patient's age, history of falls and episodes of confusion necessitate a comprehensive geriatric assessment.

Various comprehensive geriatric assessment programs use the following before performing a full-fledged assessment:

• Age

• The presence of concurrent medical conditions, such as heart failure or cancer.

• The presence of psychosocial illnesses like depression or isolation.

• Diseases unique to the elderly, include cognitive decline, mobility issues, and falls

• Prior or anticipated high health-care utilization

• Changes in living situation (for example, from independent living to assisted living, nursing facility, or in-home caretakers)

The inclusion of geriatric evaluation is imperative within the broader framework of the patient's health trajectory, taking into account their future care objectives and personal preferences. Functional status refers to the comprehensive assessment of an individual's health effect within the framework of their surroundings and social support network. It encompasses the individual's capacity to do the physical and social tasks required for their typical activities and duties. The constituents of functional status encompass activities of daily living (ADLs), instrumental activities of daily living (IADLs), and advanced activities of daily living (AADLs). Two often employed measures in the field are the Katz index, which assesses activities of daily living (ADLs), and the Lawton scale, which evaluates instrumental activities of daily living (IADLs). The most effective approach for ascertaining AADLS involves the utilization of open-ended inquiries, such as "In what manner do you allocate your time throughout the day?" or "What activities do you find enjoyable?"

There is a growing trend in geriatric assessment to integrate perspectives from non-clinical office workers, patients, and their relatives or loved ones. This collaborative approach may encompass the use of standing orders, forms, and questionnaires. An advisable strategy involves doing an annual evaluation of functional status, commencing at the age of 65 years, using the Medicare Annual Wellness Visit. For individuals below the age of 75 years with multiple comorbidities or those aged 75 years and above, it is recommended to augment this assessment with supplementary components. Furthermore, it is recommended that some components of geriatric assessment, such as evaluating activities of daily living (ADLs) and instrumental activities of daily living (IADLs), measuring gait, balance, and the risk of falls, evaluating mood and affect, as well as assessing cognition, be conducted following significant illnesses, particularly those necessitating hospitalization.

Solution to Question 15:

Ishikawa diagram or else better known as the fishbone diagram helps sketch out problems that affect healthcare in a community and thus uses those as pointers to better address them in the community.

The Ishikawa diagram, often known as the fishbone diagram, is employed by teams aiming to comprehend the numerous potential factors that contribute to a quality of care issue. This tool enables teams to concentrate their development endeavours on these identified causes. The utilization of this tool is applicable across several phases of the quality improvement framework, with its primary application commonly observed within the context of root cause analysis.

The fishbone graphic bears a resemblance to the skeletal structure of a fish. To depict the diagram about a quality of care issue, the problem (effect) is encapsulated within a box positioned on the extreme right of the diagram. A vertical line, referred to as the centre line or spine, is then positioned to the left of the box where the problem is documented. Subsequently, diagonal lines resembling fish bones are sketched emanating from the central line, also known as the spine. The diagonal lines depicted in the diagram signify distinct categorizations of the underlying factors contributing to the issue. Supplementary branches may also be delineated from the primary skeletal structure to symbolize underlying factors leading to the causes.

The groupings are typically structured into categories, commonly known as the five Ps (patients/clients, providers, policies, processes and procedures, and place/equipment), the six Ms (machine, method, materials, measurement, man, and Mother Nature), and the four Ss (surroundings, suppliers, systems, and skills). Nevertheless, the groupings need to be tailored to the organization and problem being investigated. The responsibility for ensuring this specificity lies with the quality improvement team. Once the categories have been delineated, the quality improvement team proceeds to partake in a brainstorming session to identify causes that are specific to each respective grouping. The aforementioned procedure persists until the team reaches a point where they are unable to ascertain any further underlying factors.



Other options:

Option A: Logistic regression may be used in some situations where we are not interested in a numerical outcome (like risk, etc) but we want a nominal or categorical one (like death/survival, developing a disease/not developing a disease). Here, the relationship between the variables will not be a linear one as the outcome from survival to death in relationship with the risk factor can change suddenly. Thus these outcomes are represented by 0 and 1.

Option B: A bar graph is a pictorial way of presenting a set of numbers, the magnitudes of which are proportional to the length of the bars in the graph. It can be simple, multiple, or component bar graphs and enable values to be compared visually. They are used to compare a single variable value between several groups.

Option C: A scatter plot is a statistical map used to study the correlation between two given variables. In scatter plots or scatter diagrams, the pattern made by plotted points along the two axes indicates the possible relationship. If the dots cluster along a straight line, then the relationship is linear. If there is no cluster, then there might be no relationship. Below is an example of a scatter diagram without a linear relation.



Solution to Question 16:

According to the National Immunization Schedule, the Measles/MR vaccine is given at 9 completed months (1st dose), and a second dose is given at 16-24 months. Measles vaccine can be given till 5 years of age. A dose of 0.5 mL is administered subcutaneously in the right upper arm.

Vitamin A schedule in the National Immunisation Schedule(NIS)

• 1 lakh IU at 9 completed months along with the Measles-Rubella vaccine

• 2 lakh IU is administered bi-annually for children between 1-5 years of age

Total number of doses administered- 9 doses,17 lakhs IU

Solution to Question 17:

True statements are:

• The patient is sovereign

• The aim is to improve the quality of life

Palliative care helps patients and their families cope with physical, psychological, social, and spiritual obstacles of life-threatening disease. It is interesting to note that even the quality of life of the caregiver also improves. Palliative care is needed by 56.8 million individuals annually, including 25.7 million in their final year according to WHO, while only 14% of palliative care patients worldwide receive it.

Palliative care improves the quality of life of life-threatening illness patients and their families. Early detection, assessment, and treatment of physical, psychosocial, and spiritual suffering and additional concerns diminish discomfort. Addressing suffering goes beyond physical symptoms. Palliative care supports patients and caregivers collaboratively. This includes practical assistance

and bereavement counselling. It supports patients to live as fully as possible till death.

The human right to health includes palliative care. Patient-centred and integrated health services should focus on individual needs and preferences.

Many diseases require palliative care. Most palliative care patients have chronic conditions like cardiovascular disease (38.5%), cancer (34%), respiratory disease (10.3%), AIDS (5.7%), and diabetes (4.6%). Kidney failure, chronic liver illness, multiple sclerosis, Parkinson's, rheumatoid arthritis, neurological disease, dementia, congenital defects, and drug-resistant TB may require palliative care.

Most common and serious symptoms that patients undergoing palliative care experience are pain and respiratory difficulties. Almost 80% of AIDS, cancer, and 67% of cardiovascular and COPD patients would experience moderate to severe agony at death.

Opioids also relieve dyspnea and diminish the increased pain experienced by patients. Early symptom control is ethical to reduce suffering and respect the dignity of the patient.

Solution to Question 18:

The highest level of community participation is for the community to be involved in the planning of health-related campaigns.

Engaging in the strategic planning of health-related campaigns signifies a substantial degree of community involvement, surpassing the mere contemplation of ideas or the provision of resources. The process of planning campaigns necessitates proactive participation and cooperation, signifying a heightened level of engagement in decision-making procedures that have a direct influence on the welfare of the community. This level of participation also includes engaging with the community's perceived needs through actively contributing to the strategic design and implementation of initiatives that aim to enhance health outcomes.

Earlier socialisation was envisaged as an idea for the utilisation of services but recently WHO and UNICEF deemed community participation would be of even greater help. They have defined it as "the process by which individuals and families assume responsibility for their health and welfare for those of the community and develop the capacity to contribute to their and the community's development." It also signifies the involvement of the community in the process of planning, organizing, and managing their healthcare services. This concept is commonly referred to as "Health by the People."

There are three ways in which a community would be able to participate:

1) The community can offer various resources such as facilities, human resources, logistical assistance, and maybe financial help.

2) Active community engagement is crucial in the processes of planning, management, and assessment.

3) Another significant contribution that individuals may make is by actively participating in and utilizing healthcare services.

Other options:

Options A, B and C are also methods by which a community will be able to participate but planning of health-related campaigns is considered the highest form of community participation.

Solution to Question 19:

The given clinical scenario with a patient hailing from a mosquito-infested area with fever and decreased platelet counts is diagnostic of dengue.

Symptoms of severe dengue are headache, myalgias, retroorbital pain, end-organ damage, pleural & ascitic effusions and thrombocytopenia (<100,000/ uL). Complications include mild hepatic injury, seizures, arrythmias and acute kidney injury.

Diagnosis is made by NS1 antigen detection and IgM antibody assays. Treatment includes crystalloid infusion in mild cases and colloid infusion in severe cases, oxygen administration and close monitoring.

Dengue is classified into dengue with or without warning signs and severe dengue. Dengue is to be suspected when a high fever (40°C/104°F) is accompanied by any two of the following symptoms during the febrile phase (2-7 days) which include severe headache, retro-orbital pain, arthralgia or myalgia, nausea, vomiting and maculopapular rash.

Warning signs in dengue are:

• severe abdominal pain

• persistent vomiting

• tachypnea

• bleeding gums or nose

• fatigue

• restlessness

• hepatomegaly

• malena or hematemesis

Solution to Question 20:

True statements are:

• Contemplation phase - A smoker is considering giving up his habit of smoking. (Recognizes his problem and thinks about to bring about a change)

• Maintenance phase - A person who quit smoking one year back and has abstained from smoking ever since. (He has sustained his abstinence from smoking)

Motivation enhancement therapy (MET) is a systematic intervention approach for bringing about change in the addictive behaviour seen in people with substance use disorders. It is based on the principle of motivational psychology and aims to enable the patient to deal with his addiction by producing internally motivated change.

Addictive behaviors do not change abruptly, and go through gradual stages. The stages of motivation enhancement therapy (MET) include:

Precontemplation - individuals in this stage are often unaware that their behaviour (eg. alcohol abuse) is problematic or is producing negative consequences.

Contemplation - individuals in this stage recognize their problems and begin to think about changing their behaviour. They weigh the pros and cons of change. They may still feel ambivalent about changing their behaviour.

Preparation or determination - individuals in this stage make decisions to change their habits and are ready to take action. They start to take small steps toward behaviour change.

Action - individuals in this stage change their behaviour and intend to keep moving forward with this change in behaviour

Maintenance - individuals in this stage sustain their change in behaviour for a while and intend to maintain this change going forward. People work to prevent relapse to earlier stages.

Other statements:

Statement 1: This statement is false as people in the pre-contemplation stage are unaware that their behaviour is problematic.

Statement 4: This statement is also false as people in the action stage make strides to change their behaviour.

Solution to Question 21:

A kappa statistic of 0.71 indicates a good agreement between the two doctors who collected the data.

The strength of agreement varies with the kappa coefficient as follows:

0.81 to 1: perfect agreement between the two tests/methods

0.61 to 0.80: substantial agreement/ good agreement

0.41 to 0.60: moderate agreement

0.21 to 0.40: fair agreement

0.10 to 0.20: slight agreement

Most of the studies test the association between an exposure variable and the disease. In certain cases, it is anticipated that there will be a correlation between the variables, and the challenge is to determine the extent of this correlation or association. This assertion holds special significance in the realm of reliability studies, wherein the researcher seeks to quantitatively assess the extent to which a given variable (such as a particular finding on a chest X-ray) can be reproduced through several measurements which is the utility of the kappa statistic. It is also important to note that a chi-square test can also be used but would only qualitatively assess the data and not quantitatively which of course is accomplished by the use of kappa statistic.

Kappa statistic can be calculated as:

$$\kappa = \frac{p_O - p_E}{1 - p_E}$$

where pO is observed agreement and pE is expected agreement.

Solution to Question 22:

Cohort study is an example of a longitudinal observational and analytical study.

A cohort study is a type of analytical study, which progresses from cause to effect, as exposure has occurred but not disease. It is also known as a forward-looking study, incidence study, prospective study, and longitudinal study. A well-designed cohort study is considered the most reliable means of showing an association between a cause or suspected risk factor and subsequent disease (or effect).

Types of epidemiological studies:

• Observational studies - allow nature to take its course, the investigator measures but does not intervene.

• Descriptive studies - limited to the description of the occurrence of a disease in a population- (hypothesis formulation).

• Analytical studies - go further by analyzing the relationship between health status and other variables- (hypothesis testing).

• Ecological studies

• Cross-sectional studies

• Case-control studies

• Cohort studies

• Experimental studies or intervention studies involve an active attempt to change a disease determinant or the progress of a disease and are similar in design to experiments in other sciences- (hypothesis confirmation).

• Randomized controlled trials

• Field trials

• Community trials

Other options:

Option A: Although case-control studies are inherently analytical, they are not longitudinal. Longitudinal studies involve the collection of variables about a subject over an extended duration.

Option C: A randomized control trial is an experimental study in which an intervention is given to a group of patients participating in the study and compared to another group of patients with similar characteristics who are not given that intervention to be studied. It uses patients as the unit of study.

Option D: An ecological study is a type of analytical study that analyzes the health status and its relation with other variables. It is also called a correlational study, and it uses the population as a unit of study. Longitudinal studies involve the collection of variables about a subject over an extended duration.

Community Medicine INI-CET 2024

Question 1:

Which of the following statements is incorrect about rotavirus vaccine?

a) Available as monovalent and pentavalent vaccines

b) It is a subunit vaccine

c) Not recommended after 6 months of age

d) Oral route of administration

Question 2:

Which of the following is not a polysaccharide vaccine?

a) Pneumococcal

b) Meningococcal

c) Hepatitis B

d) Haemophilus influenza type B

Question 3:

Which of the following vaccines is not recommended after 65 years?

a) Influenza

b) Herpes Zoster

c) Pneumococcal

d) Human papilloma virus

Question 4:

Arrange the steps of the experimental study :

a) C - B - A - D

b) C - A - B - D

c) A - B - C - D

d) D - C - B - A

Question 5:

Which of the following groups is not included in the high-risk core population for HIV transmission?

a) IV drug users

b) Female sex workers

c) Men who have sex with men

d) Long-distance truck drivers

Question 6:

A student went on a trip to Jaisalmer, where the temp is 40 degrees, and the relative humidity is 20%. What will the mechanism of heat loss be?

a) Evaporation

b) Radiation

c) Conduction with air

d) Conduction with objects

Question 7:

Malaria detection on the field is done by which of the following?

a) Thick smear

b) Thin smear

c) HRP2 antigen

d) Quantitative buffy coat

Question 8:

Which of the following is the best study design as per the hierarchy of evidence?

a) Meta analysis

b) Case control study

c) Cohort study

d) Large RCT

Question 9:

Dr. Sarah made a presentation on Infectious diseases. She spoke about the prevention and treatment of Zika virus. Which of the following is true with respect to the prevention and treatment of Zika virus?

a) Sexual abstinence, using condoms, and delaying pregnancy helps control the disease

b) Purified, inactivated FDA-approved vaccine is found to be helpful in prevention

c) Symptomatic treatment with NSAIDs

d) Immunoglobulin and plasmapheresis are used in the treatment of Zika

Question 10:

ICER is a unit for which of the following economic analysis?

a) Cost effectiveness

b) Cost benefit

c) Cost utility

d) Cost margin

Question 11:

You have gone from your clinic to a village for screening where people have diabetes and hypertension. What type of screening is done for family members of these patients who show up in large numbers at your clinic?

a) Multiphasic screening

b) Mass screening

c) High-risk screening

d) Anonymous screening

Question 12:

All are live vaccines except

a) Rota virus

b) Hepatitis B

c) Measles

d) Varicella

Question 13:

A pus swab has been sent for microbiological examination to your lab. After testing the sample, how will you discard it?

a) Red bin

b) Yellow bin

c) Blue bin

d) Black bin

Question 14:

Universal health coverage (UHC) means that all people and communities receive the quality health services they need, without financial hardship. Which of the following are the challenges to implementing UHC?

a) Inequalities in the distribution, adequate support, and access to quality-assured products only in low-income settings

b) The purpose of UHC is to find a balance between countries' income, health provision, and financial burden

c) To provide healthcare to everyone regardless of the country's income

d) UHC only deals with the financial aspect of health and cannot cover health provisions for all

Question 15:

Which is not a part of the Anemia Mukt Bharat Programme?

a) 6 beneficiaries

b) 6 interventions

c) 6 institutional mechanisms

d) 6 health functionaries

Answer Key

Question No.	Correct Option
1	b
2	c
3	d
4	a
5	d
6	a
7	c
8	a
9	a
10	a
11	c
12	b
13	b
14	c
15	d

Detailed Explanations

Solution to Question 1:

Rotavirus vaccine is not a subunit vaccine.

Rotavirus vaccines are oral, live attenuated vaccines and are of 3 types:

• Rotarix (human monovalent) vaccine is administered in 2 oral doses:

• First dose: At 6 weeks, no later than 12 weeks (due to the risk of intussusception).

• Second dose: With a gap of 4 weeks from the first dose, to be given by 16 weeks and no later than 24 weeks.

• RotaTeq (pentavalent bovine-human reassortant) vaccine is administered in 3 oral doses at 2, 4, and 6 months. The first dose is given between 6-12 weeks and subsequent doses within a 4-10 weeks gap. All the 3 doses should be given before 32 weeks.

• Rotavac (monovalent liquid frozen) vaccine is administered in 3 oral doses, 4 weeks apart at 6,10 and 14 weeks. It should not be administered to children more than 8 months of age.

Option C: According to the MoHFW, the rotavirus virus vaccine can be given till the age of 1 year and the potential benefit from vaccination outweighs the risk of intussusception. However, per the CDC guidelines, the maximum age for all rotavirus vaccines is 8 months, 0 days. Since this is a recall question, the options appeared as such in the exam. Additionally, rotavirus is a live vaccine,

making option 2 (which states that it is a subunit vaccine) the distinctly wrong statement and the better answer.

Solution to Question 2:

Hepatitis B vaccine is a recombinant protein vaccine.

Polysaccharide vaccines are a type of subunit vaccine containing the capsular polysaccharide antigen. They generate a T-cell independent response.

Examples include:

• Pneumococcus

• Meningococcus

• Haemophilus influenzae type b (Hib)

Recombinant protein vaccines: With the help of recombinant DNA technology, the purified recombinant proteins from the cultured microorganisms result in cleaner vaccine preparations with a better safety profile.

Examples include:

• HBV

• HPV

• Lyme disease

• Cholera toxin-B

• COVID-19

Solution to Question 3:

Human papillomavirus vaccine is not recommended after 65 years.

In otherwise healthy elderly patients, the recommended vaccines are:

• Inactivated or recombinant influenza vaccine, 1 dose annually

• Td or Tdap booster, every 10 years

• Zoster recombinant vaccine

• Pneumococcal vaccine

• COVID-19 vaccine

The zoster recombinant vaccine is not indicated for the prevention of varicella. This vaccine is given to prevent episodes of herpes zoster and post-herpetic neuralgia. It is indicated regardless of a previous episode of Zoster.

All adults ≥65 years of age should also receive pneumococcal vaccination. This is in the form of pneumococcal conjugate vaccine (PCV) first, followed by pneumococcal polysaccharide vaccine

(PPSV23) after an interval.

Solution to Question 4:

C - B - A - D is the correct order of steps in an experimental study.

Defining inclusion and exclusion criteria → Randomization → Blinding → Statistical analysis

Phases of experimental study:

1. Planning phase

• Literature review

• Trial design

• Protocol

• Budget, funding

• Approvals - ethics, collaborating institutes, funding agencies, sponsors

2. Preparatory phase

• Organization, staffing

• Delegation, responsibility, and accountability

• Standardization of protocols, training, data management, privacy

• Pilot testing, quality control

3. Operational phase

• Selection of target population

• Choosing a sample study population

• Eligibility screen: Exclusion and inclusion criteria

• Informed consent

• Subject registration, baseline evaluations

• Randomization

• Blinding

• Conduct a study on the groups allocated

• Follow-ups, evaluation protocol

• Exit plan, termination point

• Trial monitoring

4. Analysis phase

• Preliminary clarifications

• Initial Data Analysis, definitive analysis

• Statistical inference

5. Conclusion

• Validity, interpretation

• Writing reports, manuscripts, dissemination

Randomization is the random allocation of the participants into the study and control groups in the experimental studies.

Blinding is done after randomization to reduce the possibility of bias, such as subject variation bias (due to participants), observer bias, and evaluation bias (by the investigator).

Solution to Question 5:

Long-distance truck drivers are not included in the high-risk core population for HIV transmission. They are part of the bridge population.

Target interventions for the control of AIDS transmission are done for high-risk core populations like:

• sex workers

• Men who have sex with men

• Transgender

• Injecting drug users

Bridge populations like migrants and long-distance truckers also receive targeted interventions for the control of AIDS transmission.

The services offered through targeted interventions include detection and treatment for sexually transmitted diseases and condom distribution.

Solution to Question 6:

Evaporation is the mechanism of heat loss in this student.

Evaporation happens when perspiration evaporates from the skin. This is a significant source of heat loss during activity but not at rest in normal temperature settings.

When the temperature of the surroundings is more than that of the skin, the body gains heat by both radiation and conduction. Under these conditions, the heat can be lost from the body only by evaporation or sweating.

Different methods of heat loss:

• Radiation is heat loss in the form of infrared rays. (Option B)

• Conduction is heat loss that occurs by direct contact with an object. (Option C and D)

• Convection is heat loss resulting from air movement

Solution to Question 7:

In field settings, rapid diagnostic tests, such as those detecting Histidine-Rich Protein 2 (HRP2) antigen, are commonly used for malaria detection.

Rapid diagnostic tests are simple, quick, and require minimal resources, making them ideal for field use where laboratory facilities may be unavailable. HRP-2 antigen detection used in rapid diagnostic tests for malaria is specific for P. falciparum.

Rapid diagnostic tests are based on the detection of antigens using immunochromatographic methods. They are less sensitive than microscopy and cannot differentiate between species except P. falciparum. The 2 antigens detected are:

• pLDH (parasite lactate dehydrogenase)- pan malarial antigen

• HRP-2 (histidine-rich protein 2)- specific for P.falciparum.

Other options

Option A: Thick smears are more sensitive for detecting malaria parasites, they require laboratory facilities and skilled personnel, making them less practical for field settings.

Option B: Thin smears are used primarily for species identification and quantification of parasites, but like thick smears, they are unsuitable for field settings due to resource requirements.

Option D: Quantitative buffy coat (QBC): This method is highly sensitive but requires special equipment (a fluorescent microscope and centrifuge), making it unsuitable for field use.

Solution to Question 8:

Meta-analysis is the best study design as per the hierarchy of evidence.

The correct order of ranking of the hierarchy of evidence is:

Meta-analysis and systemic review > Randomised control trial > Cohort study > Case-control study > Case reports.

Solution to Question 9:

Sexual abstinence, using condoms, and delaying pregnancy help control Zika virus infections.

Zika virus can be transmitted by the bite of infected Aedes aegypti and Aedes albopictus. Zika virus belongs to the Flaviviridae family. It is an arbovirus that is transmitted by Aedes mosquitoes, primarily Aedes aegypti. The reservoir of infection is unknown and has not yet been reported in India. Zika virus can also be transmitted via sexual contact, blood transfusion, and also from mother to fetus during pregnancy.

Infection with Zika virus is usually asymptomatic. The symptoms are usually mild and include fever, skin rashes, conjunctivitis, and muscle and joint pain. These symptoms usually resolve in 2-7 days. Complications include Guillain-Barre■ syndrome, neuropathy, and myelitis, and are particularly seen in adults and older children.

Zika virus infection during pregnancy is associated with complications such as fetal loss, preterm birth, and stillbirth. The transmission of the infection to the baby can lead to congenital Zika syndrome which is characterized by microcephaly. Other malformations include limb contractures, high muscle tone, eye abnormalities, and hearing loss. Congenital malformations can follow both symptomatic and asymptomatic infections.

RT-PCR is the standard test for the diagnosis of Zika virus infection. In India, NCDC Delhi and the National Institute of Virology Pune, are the only institutes that can provide the diagnosis in an acute febrile stage.

There is no treatment available for Zika virus infection(options C & D). Plenty of rest and fluids along with antipyretics and/or analgesics are indicated for symptom relief. Medical attention should be sought if symptoms worsen. There is no vaccine available for the prevention of Zika virus infection(option B). The key measure for prevention is protection against mosquito bites. WHO also recommends contraceptive counselling for sexually active men and women, especially in regions with active Zika virus transmission.

The image below depicts the transplacental transmission of Zika virus infection.



Solution to Question 10:

ICER is a unit for Cost-effectiveness.

Incremental cost-effectiveness analysis (ICER) is a difference in cost/difference in effectiveness. Effectiveness is usually measured in Quality Adjusted Life Years (QALYs), which reflect both the quality and quantity of life.

The Cost-effectiveness analysis measures the costs and outcomes of different interventions and calculates the cost-effectiveness ratio. The benefits of the health care system are expressed in terms of results achieved (Eg, the number of lives saved or the number of days free from disease). This is a more promising tool in the health field than cost-benefit analysis. However, in many cases, it is not possible to use cost-effective analysis.

Other options:

Option B: Cost-benefit analysis: Here, the economic benefits of any program are compared with the cost of the program. The benefits are expressed in monetary terms (whereas in cost-effectiveness analysis, it is expressed in terms of results achieved)to identify whether it is economically sound and to select the best one from several alternate programs. The main drawback of this technique is that the benefits of a particular program in the health field cannot always be expressed in monetary terms. Usually, the benefits are expressed in terms of births or deaths prevented or illnesses avoided or overcome.

Option C: Cost-utility: compares the cost of a program to the health improvement it provides.

Option D: Cost margin: the difference between a product's revenues and how much it costs to make.

Solution to Question 11:

The given scenario of screening the family members of diabetic and hypertensive patients is an example of high-risk screening.

High-risk or selective screening is done on high-risk groups like in this case. Another example: MRI screening of the breast is done annually in 25-year-old women with BRCA 1, and BRCA 2 mutations in the first-degree relative.

Other options

Option A: Multiphasic screening is the application of two or more screening tests in combination at one time to a large number of people instead of screening separately for individual diseases. This procedure includes a health questionnaire, clinical examination, and investigations that can be performed rapidly with the appropriate staffing organization and equipment.

Option B: Mass screening is a screening of a whole or large population like using chest X-rays for TB screening in large populations. It is done for diseases with high prevalence.

Option D: Anonymous screening is a screening that allows people to take tests without using their name, commonly used for HIV testing. They will receive a number to access the test results instead of providing their name.

Solution to Question 12:

Hepatitis B is a recombinant vaccine, not a live vaccine.

Live attenuated vaccines are prepared from organisms that have lost their capacity to induce full-blown disease but retain their immunogenicity and are contraindicated in pregnancy unless the risk of infection exceeds the risk of harm to the fetus. Varicella, Mumps, BCG, Measles, and Rubella vaccines are examples of live attenuated vaccines.

Recombinant protein vaccines are prepared by recombinant DNA technology by expressing protective protein antigens in yeast cells like the hepatitis B vaccine. They are generally better purified from cultured microorganisms resulting in cleaner vaccine preparations with a better

safety profile.

Inactivated or killed vaccines are prepared by growing the organism in culture media and inactivating them with heat and chemicals (usually formalin). They are usually safe in pregnancy. Examples include cholera, rabies, and polio (IPV) vaccines.

Toxoid vaccines are a type of subunit vaccine prepared by detoxifying the toxins produced by organisms like tetanus bacilli. In general, they are highly efficacious and safe immunising agents.

Solution to Question 13:

The used pus swab should be discarded in the yellow colour bin as it is clinical laboratory waste.

Microbiology, biotechnology, and clinical laboratory waste are collected in autoclave-safe bags. They must be pre-treated before disposal by incineration, plasma pyrolysis, or deep burial.

Solution to Question 14:

Providing healthcare to everyone, regardless of the country's income, is one of the challenges to implementing Universal Health Coverage. A challenge related to this is mobilising sufficient resources in low-income countries to provide equitable care.

UHC aims at ensuring that all people have access to quality health services without suffering financial hardship.

Components:

• Equity in access: Everyone receives the healthcare they need.

• Quality of services: Delivered in a manner that improves health outcomes.

• Financial protection: Avoiding financial hardship for those accessing services.

Challenges:

• Health inequities.

• Resource constraints in low- and middle-income countries.

• Balancing financial protection and service delivery.

Strategies to Achieve UHC:

• Strengthening health systems.

• Universal health insurance.

• Public-private partnerships.

Other options

Option A: Inequalities in the distribution, adequate support, and access to quality-assured products are a challenge, but they occur across income settings, not only in low-income ones.

Option B: The purpose of UHC to find a balance between countries' income, health provision, and financial burden describes a goal of UHC and not a challenge.

Option D: UHC encompasses both financial protection and access to quality health services.

Solution to Question 15:

Among the listed options, 6 health functionaries are not a part of the 6x6x6 strategy employed by the Anemia Mukt Bharat Program.

The Anemia Mukt Bharat scheme is an initiative that aims to strengthen the existing mechanisms and foster newer strategies for tackling anaemia. The 6 x 6 x 6 strategy represents:

• 6 beneficiaries (option A)

• 6 interventions (option B)

• 6 institutional mechanisms (option C)

The 6 beneficiaries include:

• Preschool children (6-59 months of age)

• Children (5-9 years of age)

• Adolescents (10-19 years of age)

• Pregnant women

• Lactating mothers

• Young women in the reproductive age group (20-24 years of age)

The 6 interventions include:

• Prophylactic iron and folic acid supplementation

• Deworming

• Intensified year-round "behaviour change communication campaign"

• Testing of anaemia using digital methods and point-of-care treatment

• Mandatory provision of iron and folic acid-fortified foods in government-funded health programs

• Addressing non-nutritional causes of anaemia in endemic pockets, with a special focus on malaria, hemoglobinopathies, and fluorosis

The 6 institutional mechanisms include:

• Intra-ministerial coordination

• National Anemia Mukt Bharat unit

• National Center of Excellence and Advanced Research on Anemia Control

• Convergence with other ministries

• Strengthening supply chain and logistics

• Digital Anemia Mukt Bharat dashboard and online portal

Community Medicine NEET 2024

Question 1:

Identify the image representing the National Leprosy eradication program:

a) 1

b) 2

c) 3

d) 4

Question 2:

What is calculated with the following information?



©MARROW

a) Maternal Mortality Rate

b) Maternal Mortality Ratio

c) Death rate in reproductive women

d) Total fertility rate

Question 3:

Every pregnant woman receives a free antenatal check-up on the 9th of each month, along with free medicines and delivery services, including cesarean sections, under the JSSK program. What does JSSK stand for?

a) Janani Shishu Swasthya Karyakram

b) Janani Shishu Suraksha Kendra

c) Janani Shishu Suraksha Karyakram

d) Janani Shishu Swasthya kendra

Question 4:

What is the communication barrier if interpretation of a words are understood in a different manner?

a) Linguistic barrier

b) Semantic barrier

c) Psychological barrier

d) Cultural barrier

Question 5:

30 young volunteers with previous lifestyle modifications were contributing to drug trial for cholesterol levels based on statin therapy for 9 months. Which of the following tests is performed?

a) Chi-square test

b) Paired t-test

c) McNemar's test

d) Unpaired T-test

Question 6:

A case-control study is conducted on the association between intake of Vitamin C and the occurrence of respiratory illness. What is the order in which the following steps are conducted?

a) 3-2-1-4

b) 1-2-3-4

c) 4-3-2-1

d) 3-1-2-4

Question 7:

Which among the following diseases will be the best representation of the submerged portion?



a) Tetanus

b) Influenza

c) Chicken Pox

d) Rabies

Question 8:

What does line B represent in the graph given?



a) Death rate

b) Birth rate

c) Total population

d) Population growth rate

Question 9:

In a sudden onset disaster, the 'recovery' phase consists of which of the following?

a) i, ii, iii and iv

b) ii, iii and iv only

c) ii and iii only

d) i, ii and iv only

Question 10:

A pharmacist randomly selects from three groups of medicines: plant-based, animal-derived, and synthetic. What type of sampling is this?

a) Simple random sampling

b) Stratified sampling

c) Systemic sampling

d) Cluster sampling

Question 11:

The children in a school underwent a health check-up whose ages were 2,3,4,4,5,6,7,8. What is their median age?

a) 4

b) 4.5

c) 4.8

d) 5

Question 12:

Identify the use of this image

a) Cooling power of air

b) Air temperature

c) Air humidity

d) Radiant heat

Question 13:

A student was making a presentation with charts but lost titles while editing. What could be the heading for the following chart?

a) Physical Quality of Life Index

b) Multidimensional Poverty Index

c) Disability-adjusted life years

d) Human Development Index

Question 14:

A mother delivers a baby in a rural area with the assistance of a skilled birth attendant. The attendant cuts the umbilical cord cleanly, ties the ends with new threads, bathes the baby with warm water, wipes the baby's eyes with a sterile wet cloth from the medial to lateral corner, and then hands the baby to the mother for skin-to-skin contact. Which of these steps is incorrect?

a) Using new threads to tie the ends of the umbilical cord

b) Wiping the baby's eyes from medial to lateral corner

c) Skin-to-skin contact with the mother

d) Bathing the child with warm water

Question 15:

A homeless man comes with diarrhea, dementia, and dermatitis. He consumes alcohol occasionally. He admits to having 'X' as his staple diet. Which of the following is X most likely to be?

a) Maize

b) Wheat

c) Milk

d) Peanuts

Question 16:

In the Motivation model of health education, there are three stages for the adoption of new ideas. What is the third one other than awareness & motivation?

a) Contemplation

b) Reflection

c) Action

d) Dedication

Question 17:

An unvaccinated 5-year-old child presents with a sore throat and difficulty in swallowing. On examination, she has a greyish membrane over her tonsils. She has a fully vaccinated 2-year-old brother who was asymptomatic. What is the appropriate management of this contact?

a) No intervention

b) Booster dose of toxoid

c) Single dose of immunoglobulin

d) Prophylactic dose of penicillin

Question 18:

A disease has an incubation period of 3-14 days. It has a median incubation of 10 days. A person had to travel abroad and he was not vaccinated. As the disease is of international significance, how long will he have to be quarantined?

a) 20 days

b) 14 days

c) 28 days

d) 10 days

Question 19:

The image shown corresponds to which health program and disease?

a) AROHI for HIV

b) AMITHA for polio

c) SAPANA for leprosy

d) GAURI for cataract

Question 20:

A study is investigating the relationship between diet and the incidence of a particular disease. Among 100 vegetarians, 30 individuals are found to have the disease, while among 100 non-vegetarians, 50 individuals are affected. Which biostatistical method is most appropriate to determine if there is no significant association between diet and disease incidence?

a) Mann-Whitney test

b) Correlation coefficient

c) T-squared test

d) Chi-square test

Question 21:

Which of the following options best defines the correct sequence of the phases of the demographic cycle?

a) High expanding→ Early stationary → Late expanding → Low stationary → Declining

b) High stationary → Early expanding → Late expanding → Low stationary → Declining

c) Low stationary → Early expanding → Declining

d) Late expanding → High stationary → Declining

Question 22:

A man presents with sudden onset bilateral leg swelling, diarrhea, and difficulty breathing. He reports consuming mustard oil purchased from a local vendor prior to the onset of symptoms. A health inspector is investigating the cause of his condition. Which test is the most appropriate to detect the adulterant in the mustard oil?

a) Paper Chromatography Test

b) Methylene Blue Test

c) Phosphatase Test

d) Nitric Acid Test

Question 23:

In a population of 30,000, there were 500 live births & 14 still births in the year 2023. There were 18 children that died before the age of 1 year, 12 before 28 days of life & among that, 9 died in the first 7 days of life. Calculate the early neonatal mortality rate?

a) 24

b) 18

c) 32

d) 36

Question 24:

A 37-year-old woman from Bihar presents with fever and unexplained weight loss over the past 6 months. Examination reveals massive splenomegaly and blood investigations show anemia, thrombocytopenia, and hypergammaglobulinemia. What is the vector of the organism causing this condition?

a) Phlebotomus species

b) Aedes mosquito

c) Mite

d) Female anopheles mosquito

Question 25:

A hospital collected nasopharyngeal swabs from 1000 patients every day from the onset of the disease. The organism count was found to be maximum on the 5th day, which also showed the maximum infectivity (infecting 825 patients). What is this period of 5 days from the onset of the disease known as?

a) Generation time

b) Median Incubation period

c) Serial interval

d) Lead time

Question 26:

Which of the following are not included in the School Health Programme under Ayushman Bharat?

a) Monthly IFA tablets for all children

b) Health screening

c) Deworming for school going children

d) Sanitary napkins for adolescent girls

Question 27:

Which of the following is true about longevity dividend?

a) The intervention can delay the onset of the disease

b) Small amounts of damage is accumulated over time

c) The older you are the more prone you are to disease

d) Economic and health benefit for the society

Question 28:

For standards of ventilation, the widely quoted standard is that of De Chaumont who advocated a fresh air supply of 3000 cubic feet per person per hour. It is calculated based on which parameter?

a) Humidity of air

b) O2 content of air

c) Temperature of air

d) CO2 content of air

Question 29:

Which vaccine will be placed at zone 3 in the device shown below?



a) Pentavalent

b) Hep B

c) OPV

d) DPT

Question 30:

Pick the correct statement regarding the Sanchol cholera vaccine

a) Bivalent vaccine given in 2 liquid doses orally to children >=1 year of age

b) Bivalent given in single dose orally to children >1 years of age

c) Monovalent given in 2 liquid doses orally with a bicarbonate buffer

d) Monovalent given in a single dose orally with a bicarbonate buffer

Question 31:

In a town, there were 1000 live births in a period of 6 months. During the same period, 12 women died due to the following causes. What is the maternal mortality ratio?

a) 800 per 100000 live births

b) 20 per 1000 live births

c) 125 per 1000 live births

d) 400 per 100000 live births

Question 32:

Which of the following steps of disaster management is in correct order?

a) Disaster impact – Mitigation – Rehabilitation – Response

b) Rehabilitation – Response – Disaster Impact – Mitigation

c) Disaster impact – Response – Rehabilitation – Mitigation

d) Response – Disaster impact – Rehabilitation – Mitigation

Question 33:

Which of the following are components of the physical quality of life index (PQLI)?

a) 1,3 and 5

b) 2 and 5

c) 1,2 and 4

d) 1,3 and 4

Answer Key

Question No.	Correct Option
1	b
2	a
3	c
4	b
5	b
6	a
7	b
8	a
9	c
10	b
11	b
12	a
13	d
14	d
15	a
16	c
17	d
18	b
19	c
20	d
21	b
22	a
23	b
24	a
25	a
26	a
27	a
28	d
29	c
30	a
31	a
32	c
33	d

Detailed Explanations

Solution to Question 1:

The given image 2 represents the National Leprosy Eradication Programme (NLEP).

Characteristics of the NLEP emblem:

• The lotus in the NLEP emblem symbolizes beauty and purity.

• A partially affected thumb, a normal forefinger, and the shape of a house symbolize that leprosy can be cured and that a patient with leprosy can be a useful member of society.

• The rising sun symbolizes hope and optimism.

• The emblem captures the spirit of hope and positivity in the eradication of leprosy.

NLEP is a centrally sponsored health scheme of the Ministry of Health and Family Welfare, Government of India.

Other options:

Option A: Poshan Abhiyaan: It aims to enhance key nutritional indicators for both children and women.

Option C: National Health Mission: It is a public health initiative focused on ensuring universal access to affordable, equitable, and high-quality healthcare services.

Option D: Ayushman Bharat: It provides nationwide medical coverage and cashless hospitalization benefits for enrolled families in both public and private hospitals.

The given image shows the correct logos for these programs



Solution to Question 2:

The given expression is used for calculating the maternal mortality rate. The maternal mortality rate is expressed as the "number of maternal deaths in a given period per 100,000 women in reproductive age during the same time period".

Other options:

Option B: The maternal mortality ratio is the "number of maternal deaths during a given period per 100,000 live births during the same time".

Option C: The death rate in reproductive women is expressed as the "number of deaths among women aged 15-49 years during a given period per 1,000 women of the same age group during the same time period.

Option D: The Total Fertility Rate (TFR) is expressed as the "average number of children a woman would have over her lifetime if she were to experience the current age-specific fertility rates throughout her reproductive years (typically ages 15-49).

Solution to Question 3:

JSSK stands for Janani Shishu Suraksha Karyakram. It is an initiative by the Indian government aimed at reducing maternal and infant mortality.

The provisions of JSSK include:

• Free transport

• Free treatment

• Free drugs and consumables

• Free diagnostics and diet

• Free and cashless delivery

• Free C-Section

• Free provision of blood

Note: Previously under JSSK, the free entitlements were provided only for sick newborns till 30 days after birth. Now it is expanded to cover infants also.

Solution to Question 4:

The communication barrier that occurs when the interpretation of words is understood differently is called a semantic Barrier.

Semantic barriers arise from misunderstandings due to differences in the meaning of words, phrases, or symbols. These barriers can cause confusion and misinterpretation, leading to ineffective communication.

example: In a public health campaign, "herd immunity" may be misunderstood by some as meaning they don't need vaccination or as related to animals. This confusion can reduce vaccination rates and is a perfect example of a semantic barrier.

other options:

Option A: Linguistic Barrier: This barrier occurs when the language or dialect used in communication is not understood by one or more parties. It can include differences in language, vocabulary, or grammar, making it difficult to convey the intended message.

Option C: Psychological Barrier: Psychological barriers involve emotions, attitudes, or mental states that affect communication. These could include stress, fear, anger, or low self-esteem, which can hinder the ability to send, receive, or interpret messages accurately.

Option D: Cultural Barrier: Cultural barriers arise when differences in cultural backgrounds, beliefs, values, or practices lead to misunderstandings. These barriers can affect how messages are interpreted and can lead to miscommunication if cultural contexts are not considered.

Solution to Question 5:

This study involves a quantitative variable, i.e., blood cholesterol, and compares the before and after results of the same group. Hence, a paired T-test is used.

Tests of significance are used to check for significant association between two or more variables. Examples are the Student t-test, Paired t-test, Chi-square test, ANOVA, and Fischer-exact test.



McNemar's test is a non-parametric test for analysing paired nominal data. This test examines the marginal homogeneity of two dichotomous variables using a 2 x 2 contingency table. One independent variable with two dependent groups and one nominal variable with two categories (dichotomous) are needed for the test.

Solution to Question 6:

The correct order of steps in this case-control study is: Selecting case and control groups (3) -> Matching (2) -> Measurement of exposure (1) -> Analysis (4)

Case-Control Studies are also called retrospective studies. This study proceeds from effect to cause and is done after the exposure and outcome have occurred. They focus on diseases or health problems that have already developed. They are less time-consuming and have no risk of attrition because they do not require follow-up.

There are four basic steps in conducting a case-control study:

• Selection of cases and controls

• Matching- This is done to ensure comparability between cases and controls. Controls are selected such that they are similar to cases with respect to certain pertinent selected variables that influence the disease outcome.

• Measurement of exposure - This is obtained by interviews, questionnaires, or study of the past records of cases.

• Analysis in the form of exposure rates, risk ratio, and odds ratio.

Solution to Question 7:

The best representation of the submerged portion of the iceberg phenomenon would be influenza, as it often has a large number of asymptomatic or mild cases that go undetected, especially in milder seasons.

Tetanus (option A), chickenpox(option C), rabies (option D) and measles are typically acute and severe, with most cases being symptomatic and well-recognized when they occur. They do not show the iceberg phenomenon.

Iceberg phenomenon is shown by:

• Polio

• Influenza

• Rubella

• Mumps

• Japanese Encephalitis

• Diphtheria

• Hepatitis A and B

• Tuberculosis

• Diabetes

• Malnutrition

• Hypertension

Note: If a disease has more number of subclinical infections, it shows the iceberg phenomenon of disease depicted in the image below.



Solution to Question 8:

The line B here represents the death rate.

Line A represents the Birth rate

Line C represents the Total population

Solution to Question 9:

In disaster management, the recovery phase consists of response and rehabilitation.

Disaster preparedness refers to long-term development activities that strengthen the capacity of a country to manage all kinds of emergencies effectively.

• Response - This includes activities such as:

• Triage: Classifying the injured based on the severity of the injuries and chances of survival with medical intervention.

• Tagging: Tags with name, age, triage category, diagnosis and treatment are put for identification purposes.

• Identification of the dead: The dead bodies should be shifted from the disaster site to the mortuary.

• Rehabilitation - This includes measures that lead to the restoration of pre-disaster conditions. These measures include water supply restoration, food safety, basic sanitation, personal hygiene, and vector control.

• Mitigation - This includes measures to prevent disasters from causing emergencies or to lessen the likely effects of emergencies. Some of them are flood mitigation, land use planning, improved building codes, and the reduction and protection of vulnerable populations or structures.

• Preparedness - This is a program that aims at strengthening the capacity of a country to manage all types of emergencies efficiently. Its objective is to ensure that appropriate measurements are taken for the relief and rehabilitation of disaster victims.

Some measures that should be followed by all in case of an emergency include:

• Avoiding telephone use, unless it is to call for help

• Listen to and carry out the radio or loudspeaker broadcast messages

• Carry out the official instructions given over the radio or by loudspeaker

• Keep a family emergency kit ready

Note: Vaccination of the general population post-disaster against typhoid, cholera, and tetanus is not recommended.



Solution to Question 10:

The type of sampling described is stratified random sampling.

In stratified random sampling, the population is divided into distinct subgroups or "strata" (in this case, plant-based, animal-derived, and synthetic medicines). A random sample is then taken from each subgroup.

Other options:

Option A: In simple random sampling, every member of the population has an equal chance of being selected,

Option C: In systemic sampling, elements from a larger population are selected at regular intervals, known as sampling intervals.

Option D: In cluster sampling, the population is divided into distinct groups, or "clusters", and then a random sample of these clusters is selected

Solution to Question 11:

The median age here is 4.5 years.

In a distribution that has an odd number of elements, the median is the middle value. In a distribution that has an even number of elements, the median is the average of the middle values, i.e., the arithmetic mean of the two middle values.

Here: elements: 2,3,4,4,5,6,7,8, therefore median = (4+5)/2= 4.5

Solution to Question 12:

The image shows a Kata thermometer, it is a specialized device used to measure the cooling power of air.

Solution to Question 13:

The index that is calculated using the given indicators such as life expectancy at birth, mean and expected years of schooling, and the per capita income is the Human Developmental Index (HDI).

Human Development Index (HDI) is defined as a composite index focusing on three basic dimensions of human development. It assesses abilities

• To lead a healthy long life - measured by life expectancy at birth

• To acquire knowledge- measured by mean years of schooling

• To achieve a decent standard of living- measured by gross national income per capita

Hence the 3 dimension indices used to calculate the human development index are as follows life expectancy index, education index, and gross national index.

Other options:

Option A- Physical Quality of Life Index (PQLI) uses three indicators which include infant mortality, life expectancy at age one, and literacy rate.

Option B- Multidimensional Poverty Index (MPI) - It is an international measure of acute multidimensional poverty by capturing the acute deprivations in health, education, and living standards that a person faces simultaneously. It is assessed using 3 dimensions and 10 indicators which are as follows:

• Health - Nutrition and child mortality

• Education- Years of schooling and school attendance

• Standard of living -

• Cooking fuel

• Sanitation

• Drinking water

• Electricity

• Housing

• Assets

Option C- Disability-adjusted life years (DALY): It is the amount of time lost due to ill health. It is expressed as the number of years lost due to ill-health, disability, or early death. The components of DALY are:

• Years of life lost due to premature death

• Years lived with disability (YLD)

• Years lived in a state less than full health

Solution to Question 14:

The incorrect step is bathing the child with warm water immediately after birth. It is generally recommended to delay bathing in the immediate newborn period to help maintain the baby's body temperature and prevent hypothermia.

Other options:

Option A: Using new threads to tie the ends of the umbilical cord is correct and helps prevent infection and ensure proper cord care.

Option B: Wiping the baby's eyes from the medial to lateral corner is appropriate to prevent the spread of any potential infection from the eye's inner to outer corners.

Option C: Skin-to-skin contact with the mother is beneficial as it helps stabilize the baby's temperature, promotes bonding, and supports early breastfeeding.

Solution to Question 15:

This patient with symptoms of dementia, diarrhea and dermatitis most likely has niacin/vitamin B3 deficiency. X is likely to be maize in his diet.

Niacin deficiency causes pellagra, found mostly among people eating corn/maize-based diets, which is low in niacin and tryptophan. Excess leucine in jowar and maize disrupts the conversion of tryptophan to niacin.

Niacin can be synthesized in the body from tryptophan. Two compounds- nicotinic acid, and nicotinamide have the biological activity of niacin. The nicotinamide ring of the coenzymes NAD (nicotinamide adenine dinucleotide) and NADP (nicotinamide adenine dinucleotide phosphate) participate in oxidation/reduction reactions.

The initial symptoms of pellagra include loss of appetite, generalized weakness, irritability, abdominal pain, and vomiting. This is followed by bright red glossitis and a characteristic pigmented, scaly skin rash in sun-exposed areas. This rash is known as Casal's necklace when it forms a ring around the neck.

The 4Ds of pellagra:

• Diarrhea

• Dermatitis

• Dementia, leading to

• Death

It is diagnosed by measuring urinary levels of 2-methyl nicotinamide and 2-pyridone.

It is treated by oral supplementation of 100–200 mg of nicotinamide or nicotinic acid three times daily for 5 days.

Casal's necklace finding in niacin deficiency

Solution to Question 16:

The third stage in the Motivation model of health education, following awareness and motivation, is action.

In the process of adopting new ideas or behaviours, individuals progress through distinct stages. The first stage, awareness, involves gaining general information about health needs and issues, which serves as the foundation for further exploration. Without awareness, individuals may not recognize the importance of making changes to their health behaviours.

The second stage, motivation, encompasses interest, evaluation, and decision-making. Here, individuals express curiosity about the information presented, seek out additional details, and weigh the pros and cons of adopting the new behaviour. This stage often requires interpersonal communication and support from peers and family, as individuals contemplate the implications of their choices.

Finally, in the action stage, individuals actively adopt and implement the new behaviour or idea into their lives. This involves not just the decision to change but also taking concrete steps to integrate the new practice into their daily routine. It is during this stage that the new behaviour

becomes part of an individual's values and lifestyle, leading to sustained change. Effective communication strategies are essential to support individuals as they navigate through these stages, ensuring a successful transition to healthier behaviours.

Note: The three models of health education:

• Medical model: Transfer of proven medical information into education

• Motivational model: Awareness → Motivation → Action or Internalization

• Social intervention model: Changing of a social environment

Solution to Question 17:

The given clinical features of sore throat and difficulty in swallowing with a greyish-white membrane over the tonsils in an unvaccinated child is suggestive of pharyngeal diphtheria. The appropriate management for the asymptomatic 2-year-old brother of the child with diphtheria is to initiate prophylactic antibiotic treatment with penicillin, regardless of his immunization status.

Diphtheria is caused by the bacteria Corynebacterium diphtheriae which spreads via droplets through the respiratory tract (most important), cutaneous lesions, and fomites. Management includes antitoxin therapy as a single empirical dose of 20,000-100,000 units and antibiotics. Penicillin G/ Benzylpenicillin is the drug of choice for diphtheria. However, in case of an allergy to the same, Erythromycin is used. Other antibiotics used are clindamycin, rifampin, and tetracycline.

All household contacts of diphtheria should be closely monitored for 7 days. Culture should be taken from both the throat and nose. After culture, all contacts should receive the following:

• Antibiotic prophylaxis (benzathine penicillin G or erythromycin), regardless of immunization status.

• If fewer than three doses of diphtheria toxoid have been given, or vaccination history is unknown, an immediate dose of diphtheria toxoid should be given and the primary series completed according to the current schedule.

• If more than 5 years have elapsed since the administration of the diphtheria toxoid-containing vaccine, a booster dose should be given.

• If the most recent dose was within 5 years, no booster is required.

At least two successive negative cultures obtained 24 hr apart after completion of therapy are needed to confirm the elimination of the organism.

Solution to Question 18:

A person who is not vaccinated and has been exposed to a disease of international significance (like COVID) with an incubation period of 3-14 days should be quarantined for 14 days(longest incubation period).

Quarantine is defined as the limitation of freedom of movement of well-persons or animals, exposed to a communicable disease for a period of time not longer than the longest usual incubation period, to prevent contact with those not exposed. This is done to prevent the potential spread of the disease during its incubation period, which is the time between exposure and the onset of symptoms.

The quarantined person:

• Should stay in a well-ventilated single room and stay away from high-risk individuals

• Should not attend any religious or social gatherings

• Must wash hands often

• Their surroundings must be disinfected with 1% sodium hypochlorite solution

In case the person being quarantined becomes symptomatic, all his contacts should be home quarantined for 14 days and followed up for another 14 days or till the report turns out negative on lab testing.

Note: The median incubation period refers to the time between exposure to a pathogen and the onset of symptoms in 50% of cases. And isolation is the separation and restriction of movements or activities of persons who are sick, to prevent the transmission of disease.

Solution to Question 19:

The given image is of SAPANA for leprosy. Sapana refers to a local school-going girl who will help to spread awareness in the community.

The other developments under NLEP are:

• Mycobacterium Indicus Prani (MiP) vaccine for multibacillary leprosy patients and their contacts (it is in a project mode).

• Nikusth: A web-based system for the data management of leprosy cases.

Note: Meena is a nine-year-old girl, who is the symbol of UNICEF's Girl Empowerment programme to impart important messages on gender, child rights, education, health and development in the South Asian region.

The posters below depict the role play by Sapna:

Solution to Question 20:

To determine whether there is an association between diet (vegetarians vs. non-vegetarians) and the incidence of disease (present or absent), the best biostatistical method to use is the Chi-square test.

Other options:

Option A: Mann-Whitney test is used for comparing two independent groups on a continuous or ordinal outcome, but in this case, both variables (diet and disease) are categorical.

Option B: The correlation coefficient measures the strength of a linear relationship between two continuous variables, but it is not appropriate for categorical data.

Option C: T-squared test is typically used for multivariate hypothesis testing and is not suited for testing relationships between categorical variables like diet and disease incidence.

Solution to Question 21:

The correct sequence of the phases of the demographic cycle is High stationary → Early expanding → Late expanding → Low stationary → Declining.

India is currently in the "late expanding" stage which is the third stage in the demographic cycle where declining death rates and declining birth rates are seen.

Another important concept of the demographic cycle is the demographic gap which is the difference between birth rate and death rate. The demographic gap is maximum in the early-expanding phase, characterized by a high birth rate and low death rate. As the demography of the country transitions to the late-expanding phase, both birth rate and death rate falls causing a decreasing demographic gap. Although the birth rate is falling in the late-expanding phase, it is still higher than the death rate, which causes the population to increase.

Solution to Question 22:

The clinical scenario is suggestive of epidemic dropsy. Paper chromatography test is used to identify the presence of toxic alkaloids from Argemone mexicana seeds that adulterate mustard oil.

Epidemic dropsy is caused by the toxic compound sanguinarine, which is present in argemone oil extracted from argemone seeds. It is commonly used as an adulterant in mustard oil.

Epidemic dropsy results in sudden onset of swelling of lower limbs, diarrhea, and breathlessness. Some patients may develop hypersecretory glaucoma. It can lead to death due to cardiac failure.

Adulteration with argemone oil can be detected by the following tests:

- Paper chromatography - most sensitive test
- Nitric acid test

Solution to Question 23:

The early neonatal mortality rate here would be 18 per 1,000 live births.

It can be calculated as shown:



©MARROW

The neonatal mortality rate (NMR) refers to the rate of deaths within the first 28 days of life. It is calculated using the following parameters:

- Early neonatal deaths- death in the first 7 days of life
- Late neonatal deaths - after 7th day but before 28th completed day of life

Solution to Question 24:

From the given clinical features, the most likely disease is visceral leishmaniasis or Kala azar caused by Leishmania donovani, and the vector is a female Phlebotomus sandfly. A patient from Bihar presenting with fever, weight loss, and hepatosplenomegaly with a smear examination showing anemia, thrombocytopenia, and hypergammaglobulinemia point towards the diagnosis.

Leishmaniasis is transmitted by the bite of an infected female Phlebotomus sandfly. In India, it is endemic to Bihar, West Bengal, Jharkhand, and Uttar Pradesh. Patients present with high fever, hepatosplenomegaly, weight loss, and emaciation. The skin becomes dry, rough, and darkly pigmented. There is anemia, leukopenia, and thrombocytopenia.

The parasite exists in 2 forms:

• Amstigote form (also called LD bodies)- It is seen within macrophages and contains a kinetoplast.

• Promastigote form- It is the infective form in the sandfly. The kinetoplast is seen at the anterior end.

Kinetoplast are non-nuclear DNA present in addition to nucleus. They consist of blepharoplast and parabasal body. Flagellum originates near the kinetoplast.

Diagnosis of kala-azar is mainly by demonstration of amastigotes within macrophages in tissue aspirates such as blood smears, bone marrow, splenic, or lymph node aspirate. It is the gold standard for diagnosis.

The image below shows amastigotes located intracellularly within macrophages.



Treatment is with a pentavalent antimonial compound. In Bihar, there is resistance to antimony so amphotericin-B-deoxycholate or miltefosine is preferred.

In human beings, different species of Leishmania can cause the following diseases:

• Visceral leishmaniasis, or kala-azar,

• Cutaneous leishmaniasis,

• Mucocutaneous leishmaniasis, and

• Diffuse cutaneous leishmaniasis.

The life cycle of Lesihmania is shown below.



Other options:

Option B: The Aedes mosquito is the vector for diseases such as dengue, chikungunya, and Zika virus, none of which fit the clinical profile of visceral leishmaniasis.

Option C: Mites are vectors for diseases such as scrub typhus, which typically presents with fever and an eschar, but it does not explain the massive splenomegaly or hematologic findings seen in this case.

Option D: Ixodid ticks are vectors for tick-borne diseases such as Lyme disease and babesiosis. These conditions do not present with the clinical features observed in this patient.

Solution to Question 25:

The time interval between the receipt of infection by a host and the maximal infectivity of that host is known as the generation time.

The time of maximum communicability may precede or follow the incubation period.

Other Options:

Option B: The median incubation period refers to the time between exposure to a pathogen and the onset of symptoms in 50% of cases.

Option C: Serial interval is the gap in time between the onset of the primary case and the second case. The term serial interval is used in an outbreak of disease.

Option D: Lead time is the period between the first possible time of detection and the usual time of diagnosis.

Solution to Question 26:

Weekly Iron Folic Acid Supplementation is included in School Heath Programme not monthly IFA tablets for all children.

The Weekly Iron Folic Acid Supplementation (WIFS) program is a school health program that provides iron and folic acid (IFA) tablets which aims to reduce the prevalence of anemia in school going children.

The school health programs are managed by the primary health center (PHC).

The School Health Committee (1961) in India recommended that the school health services should be a part of the general health services administered through PHCs. Formation of school health committees at the village, block, district, state, and national levels was recommended. The National School Health Council was created as an advisory and coordinating body.

PHCs provide the following school health services:

• Screening and general assessments for

• Anemia/nutritional status

• Visual acuity and hearing problems

• Dental check-up

• Physical disabilities

• Learning disorders and behavior problems

• Basic medicines for common ailments

• Immunization

• Micronutrient management (vitamin A, iron, and folic acid)

• Deworming

• Mid-day meal

• Provison of sanitary napkins

Note: National Programme for Mid-Day Meal in Schools has recently been renamed Pradhan Mantri Poshan Shakti Nirman (PM POSHAN).

Solution to Question 27:

The intervention can delay the onset of the disease (Statement 1) is true about longevity dividend.

The longevity dividend is a horizontal approach to public health which is based on a broader strategy which means to foster health for all generations. This can be done by developing a new model for health promotion and disease prevention. The longevity-dividend model aims to prevent or delay the root causes of disease and disability by decimating the one major risk factor which is biological aging.

Solution to Question 28:

For standards of ventilation, CO_2 content of air is the paramater used to calculate fresh air supply of 3000 cubic feet per person per hour which was advocated by De Chaumont.

Ventilation involves the quality control of incoming air but not quality control of outgoing air. It also involves the replacement of vitiated air(air with decreased oxygen content) by a supply of fresh outdoor air. It aims to provide a comfortable thermal environment and reduce the risk of infections.

Fresh air supply of f 3000 cubic feet per person per hour is a good standard for air requirement. It is observed that the amount of CO_2 due to respiration is not more than 2 parts in 1000 parts of air, then the air in the room is considered fresh.

Solution to Question 29:

The image shown in the question is of an ice-lined refrigerator (ILR). The OPV vaccines are stored in the zone 3 that corresponds to the lowest

ILRs are the main method of vaccine storage in primary health centers. They are an important part of the cold chain and they maintain a temperature of 2 to 8 °C for the storage of vaccines.

They have different temperature zones for different vaccines. The upper zone is for freeze-sensitive vaccines like cholera, hep B, rotavirus(liquid), etc while the lower zone is for heat-sensitive vaccines like OPV, measles, etc.

All vaccines are kept in the basket provided with ILR. Vaccines like OPV, BCG, measles, and JE can be kept at the bottom of the basket while, DPT, TT, Hep B, IPV, and pentavalent vaccines and diluents are kept in the upper part of the basket.



Solution to Question 30:

The Sanchol cholera vaccine is a bivalent vaccine given in 2 liquid doses orally to children >=1 year of age.

The Sanchol cholera vaccine is a bivalent oral cholera vaccine (OCV) that targets both Vibrio cholerae O1 and Vibrio cholerae O139 serogroups, which are the primary causes of cholera outbreaks. It is designed to provide immunity by stimulating the production of antibodies in the intestines, the site of infection.

Key Features of the Sanchol Vaccine:

Bivalent Nature: The vaccine protects against two strains of the cholera bacterium: O1, which is responsible for most global cholera cases, and O139 which is seen in South Asia.

Dosing Schedule: The Sanchol vaccine is administered in two oral doses. The doses are given 14 days apart for children aged 1 year and above as well as adults. This two-dose regimen is essential for achieving effective immunity. The vaccine is provided in liquid form, making it easy to administer. It does not require any special preparation, such as mixing with a bicarbonate buffer, which is needed for some older cholera vaccines.

Age Recommendation: The vaccine is recommended for individuals aged 1 year and older. It is particularly useful in endemic regions or during outbreaks to prevent the spread of the disease.

Efficacy and Duration: Sanchol has been shown to provide significant protection against cholera for up to 3-5 years after vaccination, although booster doses may be required in areas with ongoing transmission.

The vaccine is widely used in mass vaccination campaigns, especially in areas prone to cholera outbreaks, such as refugee camps or regions with poor sanitation. It has been a critical tool in reducing the incidence of cholera in many endemic areas.

Solution to Question 31:

This town's maternal mortality ratio (MMR) is 800 per 100,000 live births.

Maternal mortality refers to deaths during pregnancy or within 42 days of termination of pregnancy, from causes related to pregnancy. Excluding the deaths from traffic accidents and snake bite, the total number of maternal deaths is 8.

MMR = (Number of maternal deaths in a given year / Number of live births in the same area and year) x 1,00,000 (or 1000).

Hence, MMR = 8 x 100000/1000 = 800 per 100,000 live births.

Solution to Question 32:

The correct order is: Disaster impact – Response – Rehabilitation – Mitigation

• Disaster impact and response includes:

• Search, rescue and first aid must be done first.

• Field care

• Triage

• Lagging

• Identification of dead

• Rehabilitation or recovery includes:

• Water supply

• Sanitation and personal hygiene

• Food safety

• Vector control

• Mitigation:

• Measures designed to either prevent hazards or reduce the effects of the disaster leading to disaster preparedness.



Solution to Question 33:

Infant mortality rate, literacy rate, and life expectancy at 1 year of age are the components of the physical quality of life index (PQLI).

Forensic Medicine AIIMS 2017

Question 1:

A 35-year-old male came to the OPD with complaints of an increase in his heart rate and palpitations. On examination, there was episodic tachycardia and occasional extrasystole. Ocular examination showed visual field defects. Which of the following is the likely cause?

a) Cocaine

b) Nicotine

c) Cannabis

d) Atropine

Question 2:

Which of the following is true regarding corporobasal index of the sacrum?

a) Breadth of fifth lumbar vertebra X 100 / breadth of all sacral vertebra

b) Breadth of first sacral vertebra X 100 / breadth of fifth lumbar vertebra

c) Breadth of fifth lumbar vertebra X 100 / breadth of base of sacrum

d) Breadth of first sacral vertebra X 100 / breadth of base of sacrum

Question 3:

A person was brought dead and an autopsy was performed to find the cause of death. The following findings were noted on examination. What is the probable cause of death?

a) Throttling

b) Strangulation

c) Hanging

d) Mugging

Question 4:

First carpal bone to ossify:

a) Lunate

b) Capitate

c) Hamate

d) Scaphoid

Question 5:

A 14-year-old rape victim presents with 26 weeks of pregnancy. What should not be done?

a) A female doctor should examine the patient

b) Vaginal swab examination is not necessary

c) No need to confirm pregnancy by UPT

d) Termination of pregnancy with her consent by a Gynaecologist

Question 6:

Match the following:

a) 1=a, 2=b, 3=c, 4=d

b) 1=b, 2=c, 3=d, 4=a

c) 1=d, 2=a, 3=b, 4=c

d) 1=c, 2=d, 3=a, 4=b

Question 7:

In a case of a gun shot injury to skull with entry wound at left temporal region surrounded by abrasion collar & tattooing and a exit wound at right temporal region along with metal fragments of bullet inside the bullet track. Which of the following should be preserved to determine the range of firing ?

a) Hair

b) Bullet pieces

c) Blood

d) Clothes

Question 8:

Order of appearance of rigor mortis:

a) Orbicularis oculi, facial muscles, thorax, upper limb

b) Orbicularis oculi, neck, facial muscles, upper limb, thorax

c) Neck, muscles of eyelid, upper limb, thorax

d) Neck, muscles of eyelid, facial muscles, thorax, upper limb

Question 9:

Black foot is seen in which poisoning

a) Arsenic

b) Lead

c) Mercury

d) Phosphorus

Question 10:

Criminal responsibility of Insane is defined in:

a) McNaughten's Rule

b) Currens Rule

c) Durham's Rule

d) American Law Institute Test

Question 11:

An autopsy was performed to find the cause of death of a person brought dead to the casualty and it showed a fracture of the thyroid cartilage and bruising around the neck. The cause of death is?

a) Hanging

b) Throttling

c) Strangulation

d) Mugging

Question 12:

A 30-year-old drowsy male was brought to the casualty. His pulse was 130/min, respiratory rate -30/min. His lab reports are given below. He showed improvement on nasogastric aspiration and administration of sodium bicarbonate, calcium gluconate, and 4-methyl pyrazole. What substance did he most likely consume?

a) Formaldehyde

b) Ethylene glycol

c) Paraldehyde

d) Methyl alcohol

Question 13:

Which of the following is false regarding teeth features and ethnicity?

a) In Africanoids the cusps of molars are wide and deep and shovel shaped cusps in incisors

b) Caucasians have carabelli cusps

c) Upper third molar is most commonly absent in Mongolians

d) Prominent lingual ridge and labial ridge in mongols

Question 14:

Jack knife position causes death due to:

a) Wedging

b) Burking

c) Positional asphyxia

d) Traumatic asphyxia

Answer Key

Question No.	Correct Option
1	b
2	d
3	a
4	b
5	d
6	b
7	a
8	a
9	a
10	a
11	b
12	b
13	a
14	c

Detailed Explanations

Solution to Question 1:

The given clinical scenario is suggestive of nicotine poisoning.

Chronic poisoning of nicotine also known as nicotine addiction leads to a condition called tobacco heart. It is characterized by irregularities of the heart with occasional extrasystoles and angina. It also causes amblyopia and narrowing of the field of vision.

Effects on CVS and eyes by different poisons are given below:

Poison	Effect on CVS	Effect on ocular system
Cocaine	TachycardiaSystemic arterial hypertensionMI (due to coronary vasoconstriction)TachyarrhythmiasDilated cardiomyopathyAortic dissection & rupture	Madarosis (Loss of eyebrow/eyelashes)Corneal abrasions/ulcerationsCRAOBilateral blindness
Nicotine	Irregularities of heart with occasional extrasystoleEpisodes of chest pain suggestive of angina	AmblyopiaNarrowing of field of visionBlindness
Cannabis	TachycardiaPalpitationHypertension	Bloodshot eyes (conjunctival congestions),Occasional mydriasisNystagmus
Atropine	TachycardiaArrhythmias	Dilated pupil (Mydriasis)Diplopia

Solution to Question 2:

Corporobasal index is given by the formula:

Corporobasal index = Breadth of first sacral vertebra x 100 / breadth of the base of the sacrum.

Corporobasal index of the sacrum is one of the indices used to determine the sex of the individual.

In males, it is >42 and in females, it is <42.

Other indices to determine the sex of the individual are:

Index	Formula	Male	Female
Washburnischiopubic index	Pubic length in mmx100Ischial length in mm	73-94	91-115
Sciatic notch index	Width of the sciatic notchx100Depth of sciatic notch	4-5	5-6
Sacral index	Transverse diameter of base of sacrumX100Anterior length of sacrum	<114	>114

Index	Formula	Male	Female
Sternal index	Length of manubriumX100Length of body	46.2	54.3
Medullary index oflong bones	Diameter of medullaX 100Diameter of whole bone		

Solution to Question 3:

The given autposy images are suggestive of throttling.

Thumbnail impression over the neck

Petechial haemorrhages in the eye

Extravasated blood clots surrounding larynx

In throttling, the following mechanisms may lead to death:

• Occlusion of blood supply to the blood to the brain - carotid arteries

• Occlusion of the airway - minor role

• Carotid sinus pressure may result in vagus nerve stimulation resulting in cardiac arrest (vasovagal shock)

Other signs in deaths due to asphyxia:

• Petechial hemorrhages in the face, eyebrows, eyelids, forehead, ears, conjunctiva, sclera

• When they occur in the visceral pleura, they are called Tardieu's spots

• Congested larynx with slight oozing

• Marked venous congestion of internal organs

• Emphysematous, congested lungs with serosanguinous fluid exudates in alveoli

• Dark-coloured blood due to accumulation of CO_2

Option B: Strangulation is a very generalised term used to describe deaths due to violent asphyxia. Eg- ligature strangulation, throttling etc

Option C: In the case of hanging, there will be a ligature mark, injury to the cervical spine etc.

Option D: In case of mugging the neck is compressed or squeezed by holding it between the crook of the elbow or knee. The attack is usually made from behind. Though a victim killed by this method shows signs of asphyxia and other struggle marks, the examination neck will not show any evidence of violence, as the compression was affected by the soft muscles of the arm or leg.

Solution to Question 4:

Capitate is the first carpal bone and pisiform is the last carpal bone to ossify.

Age of ossification centres of carpal bones:

Carpal bone	Age of ossification
Capitate	1-3 months
Hamate	2-4 months
Triquetrum	2-3 years
Lunate	2-4 years
Scaphoid	4-6 years
Trapezium	4-6 years
Trapezoid	4-6 years
Pisiform	8-12 years

Solution to Question 5:

According to the MTP Act 2021 amendments, pregnancy beyond 24 weeks of gestation cannot be terminated. Also, the guardian's written consent is required in the case of a minor girl.

• Up to 20 weeks of pregnancy, any qualified practitioner can perform an abortion. However, he must have undergone compulsory training.

• Between 20-24 weeks of pregnancy, the opinion of a second practitioner must be obtained to justify performing the abortion.

• The woman to be wanting an abortion should be above 18 years of age to consent. If not, consent can be obtained from the parents or guardian when she is a minor or mentally unsound.

In the cases of sexual assault of a female child, a female doctor is appointed for the examination (Option A). In case, such a doctor is unavailable, then the head of the institution can nominate a woman for that purpose. These examinations are carried out in the presence of a parent or guardian.

Vaginal swab samples are typically collected within 96 hours (ideally up to 72 hours), as the chances of detecting evidence decrease significantly after this period. (Option B)

Solution to Question 6:

• Common popular street names for cocaine are—Snuff, Rock, Crack, Coke, Snow, Cadillac, White lady, etc

• Common popular street names for LSD are –acid, blotter acid, blue caps, blue dots, brown caps, crackers, Deeda, green caps, orange wedges, Paisley caps, pink dots, pink wedges, purple Owsleys, purple wedges, The Beast, The Chief, The Ghost, The Hawk, white lightning, window panes, yellow caps, yellow dots, 25.

• Common names of Abrus precatorius are Jequirity bean, rosary pea, Buddhist rosary bead, rosary bead, Indian bead, Indian liquorice, Seminole bead, prayer head, crab's eye, weather plant, lucky bean, ojo de pajaro, gulagunchi, rati, gunja

• Capsaicin, active principle of red chilly or capsicum cause Dermatitis and burning of hands and fingers known as 'Hunan hand' which is common among the pickle industry workers, who use their hands for handling chilly paste or powder for prolonged period.

Solution to Question 7:

The main objective of the study of a firearm wound (Ballistics) is to establish the range and direction of the firing. The ultimate purpose is to determine the weapon used for firing.

Firearm wounds are of two fundamental types: wounds of entry and wound of exit.

The wound of entry is the wound through which the bullet or pellets enter the body, and the wound of exit is the wound through which the bullet emerges out of the body.

To determine the Range of firing, the following parameters are used:

• Entry wound size and shape

• Blast effect – Muzzle end imprint and edge eversion

• Blackening, Tattooing, Singeing of hair, Abrasion collar, and Grease collar.

In the above options, Hair should be preserved in order to determine the range of firing which can be examined for carbonization/desiccation in close range, also from lead deposition around the wound to assess the range of firing.

Characteristics	Contact shot	Close Shot	Near Shot	Distant Shot
Range	In-touch with skin	2.5-7.5 cm	30-60 cm	>60 cm
Size	Largest size	Bullet size	Smaller	Smallest
Shape	Stellate	Circular	Circular	Circular
Blast effect				
Muzzle end imprint	+ve	-ve	-ve	-ve
Edge eversion	+ve	-ve	-ve	-ve
Other findings				
Blackening	-ve	+ve	-ve	-ve
Tattooing	-ve	+ve	+ve	-ve
Singeing	-ve	+ve	-ve	-ve
Abrasion collar	-ve	+ve	+ve	+ve
Grease collar	-ve	+ve	+ve	+ve

Solution to Question 8:

The correct order is orbicularis oculi, facial muscles, thorax, and upper limb.

Nysten's rule describes the sequential onset of rigor mortis in various muscle groups.

Rigor mortis begins from the head and progresses down the body:

• The order of appearance of rigor mortis is heart (left chamber) → eyelids → muscles of the neck → lower jaw → muscles of the face → muscles of the chest(thorax) → upper limb → abdomen → lower limb → fingers and toes (proximo-distal progression)

It disappears in the same order of appearance.

Note that in option B, the upper limb is mentioned before thorax and hence is not the answer.

Solution to Question 9:

Blackfoot's disease is a form of arsenic induced peripheral vascular disease resulting in gangrene.

• Arsenic is a heavy metallic inorganic irritant poison. Metallic arsenic is not poisonous as it is insoluble in water and cannot be absorbed from the gastrointestinal tract. However arsenious oxide or arsenic trioxide (sankhyal or somalkar) is poisonous.

• They are found in shell fish, cod, and haddock.

• Arsenic compounds act by inactivating the sulfhydril enzymes, which in turn interfere with the cellular metabolism, in the liver, lungs, intestinal wall, and spleen. Arsenic can replace phosphorus in the bones where it may remain for years. It also gets deposited in the hairs.

• Chronic poisoning with arsenic presents with a sequence of five different set of manifestations.

• Gastrointestinal : The victim presents with gradual weight loss, malnutrition, fatigue, loss of appetite, cirrhosis of liver, nausea, vomiting, etc.

• Catarrhal changes : The victim presents with running nose, headache, conjunctivitis, bronchial catarrh, etc.

• Raindrop pigmentation : It is known to produce milk and roses complexion initially, followed by patchy brown pigmentation of the skin (especially face), which resembles the raindrops. It might also show hyperkeratosis of the skin of the palm and soles (Black Foot), which is prone to change into basal cell carcinoma at a later stage. The scalp may also show alopecia (baldness).

• Meese's lines : The victim's nail manifests with whitish lines 1 to 2 mm breadth across the nail of the finger and toes representing the deposition of the poison as a result of high sulphhydril content of the keratin.

• Arsenical neuritis : The victim presents with polyneuritis, optic neuritis, anesthesias, paresthesias, atrophy of extensors resulting in wrist and foot drop, etc.

Solution to Question 10:

• McNaughten's Rule was formed by the Judges of McNaughten's case, to clarify the common law of insanity in England during 1843. It is similar to IPC 84/ BNS 22 which states "Nothing is an offence which is done by a person who at the time of doing it by reason of unsoundness of mind is incapable of knowing the nature of the act, or that he is doing what is either wrong or contrary to law".

• This forms the basis of the 'test' of insanity that the jury should be instructed to apply in each case of alleged insanity.

• The rules are:

• Every man is presumed to be sane and to possess a sufficient degree of reason to be responsible for his crimes, until the contrary be proved to their satisfaction.

• To establish an insanity defence, it must be clearly proved that, at the time of the committing of the act, the accused was laboring under 'such a defect of reason, from a disease of the mind, as not to know the nature and quality of the act he was doing; or if he did know it, that he did not know he was doing what was wrong'.

• Curren's rule of 1964 states that an accused person is not criminally responsible, if at the time of commiting the act, he did not have the capacity to regulate his conduct to the requirements of the law, as a result of mental disease or defect.

• Durham 's rule of 1954 States that an accused person is not criminally responsible if his unlawful act is the product of mental disease or mental defect.

Solution to Question 11:

Both bruising around the neck and the fracture of the thyroid cartilage are findings seen in throttling. Throttling/manual strangulation is when bare hands are used to compress the neck and strangulate.

Mechanism of death:

• Occlusion of blood vessels supplying blood to the brain (carotid arteries).

• Occlusion to airway probably plays a minor role if any, in causing death

• Carotid sinus pressure may result in vagus nerve stimulation, resulting in cardiac arrest (vagal inhibition)

Autopsy findings:

• Bruising of neck: due to the grasping of the neck by the assailant.

• Abrasions on the neck: fingernail marks from assailant or victim

• Fracture of thyroid cartilage

• Inward fracture of greater cornu of the hyoid bone.

Solution to Question 12:

The given scenario point towards ethylene glycol poisoning.

Ethylene glycol is converted to glyoxyaldehyde by alcohol dehydrogenase. Glyoxyaldehyde causes kidney damage.

Methanol is converted to formaldehyde by alcohol dehydrogenase. Formaldehyde causes retinal damage like blurring of vision/loss of vision.

The principles of treatment for both these conditions are the same. It predominantly involves using fomipezole or ethanol to inhibit the metabolic conversion of these relatively non-toxic alcohols to their toxic metabolites.

Solution to Question 13:

The cusps of molars are wide and deep and cusps of incisors are shovel-shaped in Mongoloid races.

Africanoid races tend to have large teeth and often have more cusps on their molars, even up to eight, with two lingual cusps on the mandibular first premolars as an additional common finding.

In Caucasian races, the lateral incisors in the upper jaw are usually smaller than the central. They also have long pointed canine roots. Small nodules on the lingual surface of maxillary molars, called Carabelli's cusp are most common in Caucasian races.

In mongoloid teeth, enamel pearls and small nodules of enamel on the tooth surface, are much more frequent. The condition of bull-tooth or taurodontism is most common in Mongoloid people. The pulp cavity of molars is wide and deep, and the roots are fused and bent. A congenital lack of the third upper molar is most commonly seen. The lingual and labial ridges are more prominent.

Solution to Question 14:

Jack-knife position leads to positional asphyxia.

It is a position where the victim's thighs and knees are driven against the chest causing indirect compression of the chest and restriction of breathing movements.

Forensic Medicine AIIMS 2018

Question 1:

Which of the following is incorrect regarding the image shown?

a) Only the roots of this plant are poisonous

b) Causes AV block

c) Drug of choice for poisoning is Atropine

d) Mitha zaher is one of the variety of this plant

Question 2:

A girl was allegedly kidnapped by a man. In court, she says that she is a major and is married to the man. X-ray images of her elbow, wrist and pelvis are given. What is the approximate age of the girl ?

a) 14 to 15 years

b) 16 to 17 years

c) 18 to 19 years

d) 21 to 22 years

Question 3:

First-hand knowledge rule is applicable to:

a) Common witness

b) Hand writing expert

c) Doctor

d) Hostile witness

Question 4:

Crown Rump Length is 21 cm, length of lower limb is 10 cm, gestational age of the fetus will be:

a) 6 to 7 months

b) 4 to 5 months

c) 7 to 8 months

d) Term

Question 5:

Identify the correct pair i- Mummification, ii- Adipocere, iii- Marbling, iv- Lividity of death

a) i-4, ii-3, iii-2, iv-1

b) i-3,ii-4, iii-2, iv-1

c) i-3, ii-1, iii-4, iv-2

d) i-3, ii-4, iii-1, iv-2

Question 6:

Match the following: i- Heroin ii- Eve iii-Joint iv-LSD 1-Ganja 2-Acid 3-Brown sugar 4-Amphetamine

a) i-2, ii-1, iii-4, iv-3

b) i-3, ii-2, iii-1, iv-4

c) i-4, ii-2 iii-1, iv-3

d) i-3, ii-4, iii- 1, iv-2

Question 7:

Which of the following trait is a feature of the female skull?

a) Larger teeth

b) Frontal and maxillary sinuses are larger

c) Frontal and parietal eminences are larger

d) Supratympanic crest is present

Question 8:

If a person starts to serve a second employer, the new employer is responsible for his actions by:

a) Vicarious liability

b) Doctrine of negligent choice

c) Contributory negligence

d) Borrowed servant doctrine

Question 9:

Identify the following post-mortem change.

a) Tattooing

b) Suggilation

c) Putrefaction

d) Decomposition

Question 10:

All of the following statements are true about the given condition except:



a) Swabs should be taken immediately after moistening with sterile water

b) It is most commonly caused by incisors and canines

c) Photograph is taken with two scales at right angle to one another in the vertical plane

d) 3D imaging is possible in the given condition

Question 11:

An 18-year-old girl was brought to the OPD. Genital examination revealed the findings as given below. These findings are indicative of:

a) True virgin

b) False virgin

c) Premenstrual stage

d) Molestation

Question 12:

In case of professional misconduct, the patient's records on demand should be provided within:

a) 72 hours

b) 42 hours

c) 24 hours

d) 7 days

Question 13:

A registered medical practitioner is requested by a survivor of sexual violence to examine her. Within what time must the practitioner take a vaginal swab to look for the presence of spermatozoa?

a) Within 72 hours

b) Within 36 hours

c) Within 5 days

d) Within 48 hours

Question 14:

According to POCSO, aggravated sexual assault includes:

a) Sexual assault done by police officer

b) Threatening

c) Gang penetrative sexual assault

d) All of the above

Question 15:

Name the type of skull fracture.



a) Depressed fracture

b) Hinge fracture

c) Pond fracture

d) Comminuted fracture

Question 16:

Bullet fingerprinting involves:

a) Primary markings

b) Secondary markings

c) Both

d) None

Question 17:

Identify the injury in the image below.

a) Abrasion

b) Laceration

c) Bruise

d) Imprint

Question 18:

What is the cause of death in smoke inhalation?

a) Histotoxic hypoxia

b) Ischemic hypoxia

c) Anemic hypoxia

d) All of the above

Question 19:

Paradoxical undressing is seen in:

a) Heat stress

b) Voyeurism

c) Hypothermia

d) LSD overdose

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	a
5	d
6	d
7	c
8	d
9	b
10	c
11	b
12	a
13	a
14	a
15	b

16	c
17	a
18	c
19	c

Detailed Explanations

Solution to Question 1:

The image is of aconite. All parts of this plant (especially the roots and seeds) are poisonous.

Varieties of this plant are monkhood, mitha zaher, bish, and bikh.

It is a cardiac poison and causes AV block, bradycardia. Atropine is used in the treatment of its toxicity.

Solution to Question 2:

The most approximate age of this girl is 21-22 years.

Since the ischial tuberosity is fused in this scenario the girl must be aged >21 years.

Bone/ joint	Maximum age at fus ion	Interpretation of given x-ray
Elbow joint centres (lower en d of humerus, upper end of r adius, olecranon to ulna)	16 years	Fused, >16 years
Wrist joint centres (lower en d of ulna and radius)	19 years	Fused, > 19 years
Hip bone centres (head of fe mur, greater and lesser troch anter, ischial tuberosity)	22 years	Fused, > 21 years

Solution to Question 3:

The first-hand knowledge rule is applicable to a common witness who gives evidence about the facts observed or perceived by him.

An expert witness is someone who has been trained or is skilled or has knowledge and experience and is able to draw conclusions from the facts observed by him. Doctors, handwriting/ fingerprints/ firearms experts come under this category.

A hostile witness is a person who is supposed to have some interest or motive for concealing a part of the truth or for giving completely false evidence.

Solution to Question 4:

Gestational age of this fetus is 6-7 months.

Crown rump length + length of lower limb = crown heel length

Rule of Haase enables the estimation of the age of the fetus in lunar months from its crown-heel length (in cm).

According to the rule,

• Age in months = Square root of crown-heel length (when the crown-heel length is 25 cm or less/ first 5 months).

• Age in months = crown-heel length divided by 5 (when the crown-heel length is more than 25 cm/ last 5 months).

Here, crown heel length = Crown rump + lower limb length = 31 cm.

Hence, gestational age = 31 / 5 = 6-7 months.

Solution to Question 5:

Image 1. Marbling: It is the discoloration of superficial veins due to the formation of sulphmethemoglobin, which starts in 24 hours and becomes prominent in 36-48 hours.

Image 2. Lividity of death: Postmortem lividity (suggillation, postmortem hypostasis/postmortem staining/livor mortis/suggillation/vibices) develops in the dependent portions of the body due to engorged vessels showing through the skin.

Image 3. Mummification: It occurs in the absence of moisture in the air and continuous action of dry or warmed air.

Image 4. Adipocere: It forms under warm and humid weather due to hydrolysis of lipids.

Solution to Question 6:

Heroin	Brown sugar, Smack, Dope
Ganja	Joint , Reefer, Weed, Pot, Gr ass, Purple haze
Ethylene dioxymethampheta mine	Eve
LSD	Acid

Solution to Question 7:

Frontal and parietal eminences are larger in a female skull.

Differentiating features between male and female skull:

Note: The features given above develop after puberty and are modified by senility, so are applicable only between the 20th and the 55th year.

Trait	Male	Female
General size	Large	Small
Architecture	Rugged	Smooth
Supraorbital ridges	Prominent	Less prominent
Mastoid process	Prominent	Less prominent
Occipital region	Marked muscle markings	No marked muscle markings
Frontal eminences	Small	Large
Parietal eminences	Small	Large
Orbits	Square with round margins	Round with sharp margins
Mandible	Large, broad ramus	Small, small ramus
Chin shape	U-shaped	V-shaped

Solution to Question 8:

When an employee is being borrowed, the lending employer temporarily surrenders control over this worker and the borrowing employer temporarily assumes control. This is known as borrowed servant doctrine.

Vicarious liability is by which the employer is responsible for his employees' negligence if such activity occurs in the course of the employment and within its scope.

Solution to Question 9:

The image shows suggilation.

Postmortem lividity (suggilation/ postmortem hypostasis/ postmortem staining/ livor mortis/ vibices) develops in the dependent portions of the body due to engorged vessels showing through the skin.

Solution to Question 10:

The above image shows a human bite mark. The photograph is taken with two scales at the right angle to one another in the horizontal plane.

Human bites are usually semi-circular or crescentic caused by the front teeth (incisors, canine).

Swabs of bite marks should be taken immediately using a swab moistened with sterile water. A sample of the control area adjacent to the mark and of the victim's saliva should also be taken using swabs moistened with sterile water.

3D imaging is done in which the image is measurable with high precision.

A plastic substance such as rubber is laid over the bite mark which produces a permanent negative cast.

Solution to Question 11:

The given findings are suggestive of a false virgin.

The examination findings of a separated labia majora, a flabby labia minora, and a fourchette tear are indicative of forceful penetration.

Hymen usually gets ruptured with the first act of sexual intercourse. However, in some cases, hymen remains intact in spite of sexual intercourse. It is indicative of a false virgin.

Solution to Question 12:

In case of professional misconduct, the patient's records on demand should be provided within 72 hours.

If a doctor does not maintain the medical records of the patients for a period of three years as per the regulation, and refuses to provide the same within 72 hours, when the patient or his/her authorized representative makes a request for it, then it is professional misconduct.

Solution to Question 13:

A vaginal swab of a rape victim demonstrates spermatozoa if the vaginal swab is taken within 72 hours after an assault.

However, a vaginal swab can be taken within 96 hours to demonstrate semen.

Evidence on the outside of the body and on materials such as clothing can be collected even after 96 hours.

Solution to Question 14:

Sexual assault done by a police officer is an aggravated sexual assault according to POCSO (Protection of Children from Sexual Offences).

The POCSO act defines a child as any person below eighteen years of age. It defines various forms of sexual offenses which include:

• Sexual harassment - Threatening

• Sexual assault - Any act with sexual intent that involves physical contact without penetration.

• Aggravated sexual assault - Similar to sexual assault along with the criteria explained under the term aggravated

• Penetrative sexual assault - An act with sexual intent that involves physical penetration of the penis or any object into the vagina, mouth, urethra, or anus.

• Aggravated penetrative sexual assault - Similar to penetrative sexual assault along with the criteria explained under the term aggravated

• Pornography

The term aggravated is used for those assaults done:

• By a person in a position of trust or authority such as a police officer, security force, relative, doctor

• In the course of communal violence

• On mentally & physically disabled

• Gang sexual assault

• Using deadly weapons

• Causing grievous hurt

Depending on if there was penetration or not, it can be aggravated sexual assault (if there is no penetration) or aggravated penetrative sexual assault (if there was penetration). Both are different.

The Act prescribes stringent punishment graded as per the gravity of the offence, with a maximum term of rigorous imprisonment for life and fine.

Note- Gang penetrative sexual assault comes under aggravated penetrative sexual assault and not under aggravated sexual assault. Therefore, option C cannot be correct in this case.

Solution to Question 15:

The image shows hinge fracture/ motorcyclists fracture. Here, the base of the skull is divided into 2 halves, each moving independently of the other like a hinge.

Solution to Question 16:

Bullet fingerprinting involves both primary and secondary markings.

Primary markings are formed when a bullet passes through a barrel and the sides are marked by the rifling of the barrel (which is specific to the make and model of the gun).

Irregularities on the inner surface of barrel itself which are specific for that particular weapon are also seen on the bullet and are called secondary markings.

Solution to Question 17:

The given image shows an abrasion.

An abrasion is the most superficial of injuries. It does not penetrate the full thickness of the epidermis.

A pure abrasion does not bleed, as blood vessels are confined to the dermis. However, the dermal papillae are corrugated in nature. Thus, many abrasions enter the dermis and bleeding commonly occurs.

Solution to Question 18:

Victims of smoke inhalation are affected by carbon monoxide poisoning. In carbon monoxide poisoning, the cause of death is anemic hypoxia.

Carbon monoxide has a higher affinity to hemoglobin compared to oxygen and hence the amount of hemoglobin available for oxygen transport is markedly reduced. It displaces oxygen from its combination with hemoglobin and forms a relatively stable compound known as carboxyhemoglobin. It thus reduces the oxygen content of the blood, and hence, that of the tissues. It acts as a chemical asphyxiant and produces death due to anemic anoxia.

It is important to note that in the case of anemic hypoxia, O_2 content is low but dissolved O_2 is within normal limits so it does not stimulate peripheral chemoreceptors.

Note: Carbon monoxide poisoning causes both anemic (due to CO-Hb) and histotoxic (due to inhibition of cytochrome C oxidase) hypoxia. But the main cause of death is due to anemic hypoxia.

Option A - Histotoxic hypoxia- The most common example is cyanide poisoning. Cyanide inhibits important enzymes in the oxidative phosphorylation process, which is essential to the survival of our cells. Additional oxygen supplementation will not be able to overcome this inhibition, so it will have limited benefit. Hyperbaric oxygen may be helpful to some extent.

Option B - Stagnant hypoxia refers to ischemic hypoxia that results from inadequate circulation due to conditions such as heart failure and hypoxia in the setting of shock. In such conditions, administering intravenous fluids can improve the patient's condition because it is the circulation that is affected. If the shock is prolonged it can result in acute respiratory distress syndrome (ARDS), which can be treated with positive pressure ventilation and not simply by increasing oxygen administration.

Solution to Question 19:

The phenomenon of paradoxical undressing is seen in hypothermia victims.

Its occurrence is attributed to a paradoxical feeling of heat and may lead to an erroneous assumption of a sexual offense.

Fatal hypothermia-induced stupor, loss of powers of judgment, disorientation, and increasing incapacitation are associated with this phenomenon.

Forensic Medicine AIIMS 2019 (May & Nov)

Question 1:

Identify the following post-mortem change.

a) Tattooing

b) Suggilation

c) Putrefaction

d) Decomposition

Question 2:

Paradoxical undressing is seen in:

a) Heat stress

b) Voyeurism

c) Hypothermia

d) LSD overdose

Question 3:

Identify the plant which produces a deliriant toxin.

a) Erythroxylum cocca

b) Datura

c) Hyoscine

d) Digitalis purpura/cannabis

Question 4:

An 11-year-old girl was found to be quiet and shy in her classroom with intermittent crying, for the past few days. It was discovered that her uncle was touching her genitalia inappropriately. The duty of the principal is to report this to?

a) Magistrate

b) Police

c) Child Welfare Committe

d) Parents

Question 5:

Which of the following postmortem changes is shown in the given image?

a) Marbling

b) Algor mortis

c) Rigor mortis

d) Postmortem lividity

Question 6:

What is bullet wipe?

a) Residual from barrel of gun

b) Blackening

c) Gutter fracture of skull

d) Tattooing

Question 7:

A 28-year-old man presented with dry mouth, urinary retention, and constipation after consuming an unknown substance. On examination, his PR - 110bpm, and his pupils were dilated. What did he most likely consume?

a) Heroin

b) Morphine

c) Malathion

d) Belladona

Answer Key

Question No.	Correct Option
1	b
2	c
3	a
4	b
5	a
6	a
7	d

Detailed Explanations

Solution to Question 1:

The image shows suggilation.

Postmortem lividity (suggilation/ postmortem hypostasis/ postmortem staining/ livor mortis/ vibices) develops in the dependent portions of the body due to engorged vessels showing through the skin.

Solution to Question 2:

The phenomenon of paradoxical undressing is seen in hypothermia victims.

Its occurrence is attributed to a paradoxical feeling of heat and may lead to an erroneous assumption of a sexual offense.

Fatal hypothermia-induced stupor, loss of powers of judgment, disorientation, and increasing incapacitation are associated with this phenomenon.

Solution to Question 3:

The given plant is Erythroxylum cocca.

Cocaine is the principal alkaloid found in the cultivated varieties of coca.

Solution to Question 4:

The principal has to report this case to the police.

The POCSO Act, 2012 is a comprehensive law for the protection of children from the offences of sexual assault, sexual harassment, and pornography. This is done while safeguarding the interests of the child at every stage of the judicial process by incorporating child-friendly mechanisms for reporting, recording of evidence, investigation, and speedy trial of the offence through designated special courts.

Section 21(1) of the POCSO Act, 2012 requires mandatory reporting of cases of child sexual abuse to the law enforcement authorities and applies to everyone including parents, doctors, and school personnel. Failure to report a suspicion of child abuse is an offence under the Act.

Solution to Question 5:

The postmortem change shown above is marbling.

The blood in the vessels is hemolyzed, which stains the vessel walls and the adjacent tissues, giving rise to a marbled appearance. The marbling of the skin becomes prominent about 36–48 hours after death in summer and is distinctly appreciable in the superficial veins of the abdomen, shoulders, chest, and inguinal region.

Solution to Question 6:

Bullet wipe is due to the grease, soot or debris from the barrel of the gun.

While examining the clothing of a victim of bullet injury, there is a deposition of various residues surrounding the entrance defect at various ranges known as Bullet wipe. It helps in determining the wounds of inlet and outlet upon the body.

Solution to Question 7:

The given clinical scenario with dilatation of the pupil indicates Atropa belladonna consumption.

Atropa Belladonna acts by inhibiting the muscarinic effects of acetylcholine. All the parts of the plants are toxic. The active principle is 1-hyoscyamine.

Heroin, morphine, and malathion cause constriction of the pupil.

Forensic Medicine AIIMS 2020 (May and Nov INI-CET)

Question 1:

Identify the type of skull fracture in the given image?



a) Ring

b) Depressed

c) Fissured

d) Comminuted

Question 2:

A middle-aged man was brought to the ER after an assault with a sharp axe. Identify the type of wound?

a) Incised wound

b) Chop wound

c) Lacerated wound

d) Abrasion

Question 3:

A patient comes to the casualty with life-threatening injuries. There was no time for taking consent. The doctor started doing life-saving procedures with all precautions, and despite best efforts, the death of the patient occurred. The doctor is not responsible under which doctrine?

a) Res ipsa loquitor

b) Doctrine of anticipation

c) Doctrine of extended consent

d) Doctrine of conjugated consent

Question 4:

Which of the following is false regarding the management of snakebite in India?

a) Atropine and neostigmine are very useful in the management of krait bite

b) Neostigmine and ventilatory support are used in treatment along with ASV

c) ASV is the mainstay of treatment

d) Current ASV in India is not effective against humped pit viper

Question 5:

A 14-year-old girl comes to the OPD with her mother. History of penovaginal penetration by the neighbour is reported to the doctor. The girl admits to giving the neighbor the consent for the same and says she does not want to be examined. As a medical officer, which of the following should you do?

a) 1,2,3 are true

b) 2,3,4 are true

c) 1,3,5 are true

d) 3,4,5 are true

Question 6:

A surgeon performs a hysterectomy for uterine fibroids after obtaining relevant informed consent from the patient. Despite adequate precaution, the ureter was injured intraoperatively. Under which of the following principles will the doctor not be held responsible for this act?

a) Novus actus interveniens

b) Medical maloccurrence

c) Res ipsa loquitur

d) Physician error

Question 7:

The punishment for the death of a patient by spurious drug:

a) 5 years

b) 7 years

c) Life imprisonment

d) 1 year

Question 8:

Identify the seeds shown in the image below.

a) Tropicana canna

b) Nigella sativa

c) Datura seeds

d) Opium seeds

Question 9:

A 12-year-old boy presents with the following injury that he sustained on being tackled by an opponent during the course of a football match. As the Casualty medical officer, you would enter the wound description in the case sheet as _____

a) Incised wound

b) Abrasion

c) Laceration

d) Bruise

Question 10:

You are posted as an intern in casualty when an MLC case is brought by the police with the following appearance. She was declared brought dead. What would you mark as the likely cause of death before sending the patient for postmortem?

a) Drowning

b) Hanging

c) Strangulation

d) Traumatic asphyxia

Question 11:

What is the approximate age of a person in whom the medial end of the clavicle is fused and also the sternum is completely fused

a) <18 years

b) 18-22 years

c) 22-25 years

d) >25 years

Question 12:

Identify the condition shown in the image.

a) Postmortem lividity

b) Marbling

c) Filigree burns

d) Subcutaneous emphysema

Question 13:

Identify the image shown below



a) Shot gun wound

b) Tattooing

c) Arsenic poisoning

d) Graze wound

Question 14:

A 22-year-old woman was found dead at home. She was brought to the mortuary by the police and the following findings were noted on examination. Which of the following is the likely cause of her death?



a) Strangulation by ligature

b) Hanging

c) Throttling

d) Drowning

Question 15:

Which of the following is seen in freshwater drowning?

a) A,B,C,D

b) A,B,C

c) B,C,D

d) A,C,D

Question 16:

Match the following weapons with the possible injuries caused by them:

a) A-4, B-3, C-1, D-2

b) A-1, B-3, C-4, D-2

c) A-1, B-2, C-4, D-3

d) A-4, B-1, C-3, D-2

Question 17:

You are working as a medical officer at a PHC. A 17-year-old girl presents to you after an alleged sexual assault, the girl refuses physical examination. Which of the following steps would you take to proceed in this case?

a) 1, 2 and 3

b) 2, 3 and 5

c) 1, 2 and 5

d) 2 and 3

Question 18:

Match the following poisons with their probable mechanism of action: A. Zinc chloride B. Chloral hydrate C. Quinine D. Potassium carbonate 1.Abortifacient 2. Irritant 3. Corrosive 4.

Stupefying

a) A-2, B-4, C-1, D-3

b) A-4,B-3, C-2,D-1

c) A-4, B-2, C-1,D-3

d) A-2, B-3, C-4,D-1

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	a
5	c
6	b
7	c
8	b
9	c
10	a
11	d
12	b
13	a
14	c
15	d
16	a
17	a
18	a

Detailed Explanations

Solution to Question 1:

The given image shows a fissured fracture of skull.

Solution to Question 2:

The given wound caused by an axe, is likely to be a chop wound.

Chop wounds are so-called because of the action in which they are produced. Depending on the type of instrument, they may resemble an incised or a lacerated wound. The shape of the wound may give clues to the type of weapon. It would be a deep, well-margined wound, gaping with limited damage to the surrounding skin.

For example, in the given image, the wound resembles an injury produced by a sharp axe. Such wounds may be inflicted with homicidal intent on the skull/scalp.

Solution to Question 3:

Under IPC 92, the doctor can perform life-saving procedures in a patient with life-threatening injuries without obtaining consent. The guiding principle is whether the delay in treatment in order to obtain consent would result in harm to the patient. This is the doctrine of anticipation.

Under some circumstances like emergencies, informed consent may be neither possible nor necessary.

IPC 92 states that nothing is an offense by reason of any harm which it may cause to a person for whose benefit it is done in good faith, even without that person's consent, if the circumstances are such that it is impossible for that person to signify consent, or if that person is incapable of giving consent, and has no guardian or other person in lawful charge of him from whom it is possible to obtain consent in time for the thing to be done with benefit.

Other options:

'Res Ipsa Loquitor' means 'the thing or fact speaks for itself'. It is obvious evidence for the negligent act committed. For example, swabs left in the abdominal cavity post laparotomy.

The doctrine of extended consent: It is used in unanticipated conditions during surgery. This comes into play when a surgeon encounters an unanticipated abnormality within the field of surgery. It is called the "extension doctrine," and it assumes that the surgeon is using reasonable judgment. For example, a surgeon incurred no liability for draining some ovarian cysts during the course of an appendectomy. The condition must be one that was unforeseen, and the patient must not have expressly refused such an intervention.

Doctrine of conjugated consent - Dummy option (No such consent has been defined).

Solution to Question 4:

Atropine is not useful in the management of krait bite. The toxin in common krait's venom acts presynaptically, hence atropine is not used in the treatment of envenomation by krait.

As the cobra venom acts postsynaptically, atropine can be used in the management.

ASV(Anti Snake Venom) is the mainstay of snakebite treatment.

ASV used in India is effective against:

- Cobra
- Common krait
- Russel viper
- Saw scaled viper

Neostigmine and ventilatory support are used in the treatment of snakebite.

Solution to Question 5:

In the given scenario 1, 3, 5 are true.

The scenario describes a 14-year-old girl with a history of sexual intercourse with her neighbor albeit with consent. Consent for intercourse is valid only after 18 years in an unmarried girl. As she is less than 18 years, it is presumed to be a medicolegal case of sexual assault.

Examination of a victim of sexual assault should be done after the consent of the victim (if she is less than 12 years of age or unsound mind, consent from the guardian needs to be taken). If she denies examination, counseling has to be done regarding the need for examination and how it could be held up against her in the court of law. If she still refuses for part or whole of examination or sample collection, an informed refusal needs to be documented.

Information to the police is given in all medicolegal cases.

Solution to Question 6:

The doctor will not be responsible for the above act under medical maloccurrence.

Medical Maloccurrence/Inevitable Accident: Every human being responds differently to different drugs as well as to different treatment modalities. On certain occasions, despite all proper care given by the doctor during treatment, the patient fails to respond properly or may suffer some adverse reactions to the drug which may lead to severe injuries or permanent deformities. This is known as a medical maloccurrence/inevitable accident.

If the doctor can prove this before the court, it will be an absolute defence against malpractice.

'Res ipsa loquitur' means 'the thing or fact speaks for itself'. It is obvious evidence of the negligent act committed. For example, swabs left in the abdominal cavity post laparotomy.

'Novus actus interveniens': This essentially means that the negligence at the core of the lawsuit stems from the actions of a third party and that this negligence was unforeseeable. The viability of this defence heavily relies on the specific details of each case. For instance, if a doctor writes a prescription that is allegedly illegible, and the pharmacist misreads it, fails to verify it with the doctor, and then dispenses the wrong medication to the patient, resulting in an injury, the doctor might use this defence in such a scenario.

Solution to Question 7:

The punishment for the death of a patient by a spurious drug is imprisonment for life.

In case of an adulterated, spurious, or any drug that when used by any person for or in the diagnosis, treatment, mitigation, or prevention of any disease or disorder is likely to cause ill health or is likely to cause such harm on his body as would amount to grievous hurt within the meaning of Section 320 of IPC, shall be punishable with imprisonment for a term that shall not be less than 3 years but that may extend to a term of life.

Solution to Question 8:

The seeds shown in the given image are of the plant Nigella sativa. These are also known as black cumin or kalonji.

The seeds are the most commonly used parts of the plant in traditional medicine. Seeds have a pungent and bitter taste with a considerable amount of oil inside. The seeds are used as a spice in cooking. They are also used as a food preservative and the most often used therapeutic agent is its oil.

Solution to Question 9:

The given image shows a laceration.

A laceration is a tear or a split in the skin, mucous membrane, muscle, or any internal organ, with a depth spanning more than the covering epithelium of the skin or an organ, and is produced by blunt force.

Features of laceration:

• Margins are irregular with pieces of tissue attached in between called tissue tags or bridges

• Bruising is seen around the margin

• Deeper tissues are unevenly divided with tissue tags

• Hair bulbs are crushed

• Depth of the wound contains abundant foreign material such as dust particles, paint material of the vehicle involved, etc

• In skin lacerations, usually one of the margins overhangs the other, which could help in assessing the direction of the force

• A laceration, on healing, produces a permanent scar.

Classification of a laceration:

• Split laceration (incised - like laceration)

• This occurs when the skin is crushed between two hard objects

• For example, during a fall, the scalp gets crushed between the skull from inside and hard ground outside. This results in a wound, which usually mimics an incised one unless examined closely.

• Stretch laceration

• As the name implies, it occurs due to the over-stretching of the skin, beyond its elasticity.

• This typically occurs in the setting of a fractured femur, where the fracture fragment may stretch and pierce the skin to project out

• Avulsed laceration

• An avulsion is a lesion where the skin only separates by tearing from the underlying tissues or peels off and overhangs the wound

• This is common in road traffic accidents.

Solution to Question 10:

The given image shows frothing from nostrils and no clear signs of facial or conjunctival congestion. It is likely a case of drowning.

Frothing may also be seen in throttling or strangulation, but it might be associated with signs of conjunctival congestion, facial congestion, patterned bruise on neck etc. In the absence of these features, it is unlikely to be caused by strangulation.

Traumatic asphyxia is unlikely as the face of this patient is all whitened without signs of congestion or any obvious signs of trauma

Solution to Question 11:

Approximate age for the given person with fused medial end of clavicle and completely fused sternum is most likely >25 years.

The sternum is formed from the fusion of Manubrium, 4 sternebrae(body), and the xiphoid process. The fusion of sternebrae occurs from below upwards, followed by its fusion to the xiphoid process and then fusion of manubrium to the fused sternum.

Age	Age determination
<5 years	The appearance of ossificatio n centers
5-16 years	Analysis of hand and wrist
>16 years(male)	Complete fusion of distal radi us or ulna
>16 years(femal e)	Complete fusion of distal uln a
>18 years(male)	Complete fusion of distal fem ur or proximal tibia
>18 years(femal e)	Complete fusion of iliac crest
>21 years	Complete fusion of medial en d of clavicle

Manubrio-sternal joint	60-70 ye ars
1st and 2nd Sternebrae	25 years
2nd and 3rd sternebrae	20 years
3rd and 4th sternebrae	15 years
Xiphoid process and bo dy	40-45 ye ars

Solution to Question 12:

The image shows postmortem marbling. It is a visible postmortem change that is characterized by linear branching patterns on skin that correspond to the course of superficial veins.

Intestinal bacteria infest the venous system as blood acts as a nutrient medium for their proliferation. There is subsequent hemolysis which stains the vessel walls and the adjacent tissues, giving rise to marbling. This becomes prominent in about 36–48 hours after death in the summer and is distinctly appreciable in the superficial veins of the abdomen, shoulders, chest, and inguinal region.

Solution to Question 13:

The given image shows entry wound of a shot gun from a mid-range or long distance.

• There is a central round hole, with multiple pellet holes scattered around the central hole.

• This wound is most likely to have been caused by a shotgun from a range of 1-5 m.

Following are the appearances of shot gun entry wound from different distance:

Appearance of shotgun entry wounds at different ranges:

Contact shot

• Soft tissues - single circular wound about the diameter of the muzzle with muzzle imprint and no soot soiling, tattooing, singeing.

• Bone - same as with soft tissues, but wound may be ragged and split because of gas rebound.

<30 cm

• Circular wound with surrounding soot soiling.

• Burnt hairs (singeing).

• Punctate skin lesions from powder particles (tattooing).

30cm - 1m

• Circular rat-hole wound.

• Soot soiling.

• Powder tattooing.

1 - 5 meters

• Central rat-hole wound.

• No soot, tattooing +/-.

• Satellite pellet holes around the periphery.

>5 meters

• Diffuse pellet pattern

• No central hole, no soot soiling, tattooing, singeing.

Solution to Question 14:

In the given case scenario and autopsy images showing carotid intimal tear and muscle hematoma with nail marks on neck and absence of ligature mark point towards a diagnosis of throttling.

Other options

Option A- Strangulation by ligature - ligature mark with signs of a struggle would've made this diagnosis more likely.

Option B- Hanging - ligature mark with a suicide note would've made it more likely.

Option C- Drowning- Body recovered from a pond or other water body with other suggestive features like frothing etc would've made this more likely.

Solution to Question 15:

In cases of freshwater drowning, hyperkalemia is seen due to hemolysis of swollen RBCs and not hypokalemia.

Freshwater is hypotonic to plasma and is rapidly absorbed from the alveoli and into the RBCs. The RBCs finally swell up and lyse resulting in the release of intracellular potassium causing hyperkalemia. Increased intravascular blood volume combined with hyperkalemia causes arrhythmia.

Seawater is hypertonic to plasma and causing a shift of fluid from the blood to alveoli causing pulmonary edema. There is a net decrease in blood volume causing a fall in blood pressure.

The table below summarises the cause of death in different types of drowning.

Type of drowning	Cause of death
Freshwater drowningSeawater drowning	Disruption of surfactant
Dry drowning	Laryngeal spasm/Vagal inhibition
Post immersion Syndrome / Secondary drowning	Cerebral anoxiaPneumoniaPulmonary edemaChemical pneumonitisMetabolic acidosis

Solution to Question 16:

The correctly matched sets of the above-mentioned weapons and the possible injuries are as follows:

Mechanism of injury with the above mentioned weapons:

Axe - it is a heavy weapon that may or may not be sharpened. Injury is produced by chopping action of the weapon and it may be in a spectrum of an incised wound to laceration depending upon the sharpness of the blade.

Road traffic accident - rubbing of epidermis tangential to a rough or hard surface gives rise to graze abrasion also called brush burn. It can occur in a road traffic accident after a pedestrian is being hit by a vehicle and the victim is slid/grazed for some distance before he comes to a halt.

Blade - it gives rise to a clean-cut wound which is usually deepest at its starting. Tailing of the wound indicates the direction of injury. Length is the biggest parameter of this type of wound.

Lathi- It is a blunt force injury giving rise to a patterned abrasion with 2 parallel lines and a normal unbruised area on site of impact or stick/lathi. The lathi forces the blood from the site of impact to either side which gives rise to tramline appearance.

The image below shows a chop wound inflicted on the scalp:

A. Axe	4. Chop wounds
B. Road traffic accident	3. Graze abrasions
C. Blade	1. Incised wound

A. Axe	4. Chop wounds
D. Lathi	2. Tram track bruise

The image below shows a schematic illustrating heaping up of epidermis in a graze abrasion or brush burn:



The image below shows production of a tramline bruise associated with a lathi strike:



Solution to Question 17:

In the given scenario, you need to start treatment for STI, inform the police, and record refusal of consent for examination.

Examination of a sexual assault victim shouldn't be forced upon, but an informed refusal is to be documented after explaining to them the consequences of choice.

Medical treatment and testing for pregnancy have to be done for all victims of sexual assault free of cost and police should be informed as per Section 397 of BNSS (357-C of CrPc).

As the girl is under 18 years of age, as per POCSO (The Protection of Children from Sexual Offences) act 2012, the medical practitioner should initiate treatment for exposure to sexually transmitted diseases (STDs) including prophylaxis for identified STDs (including HIV if suspected) and possible pregnancy and emergency contraceptives should be discussed with the victim and her parent/guardian.

Solution to Question 18:

For the above-given options, the correct match is given as

Rationale for the classification

Zinc chloride also known as butter of zinc is highly irritant to the eyes on topical application. It can cause erosive esophagitis on ingestion. It has a fatal dose of 350mg. It could've also been classified as a corrosive, but since potassium carbonate is an alkali with stronger corrosive action, it is better classified as an irritant for this question

Chloral hydrate is a CNS depressant. It is also known as knockout drops because of its rapid action and used in robbery or rape to quickly render the victim helpless.

Quinine is a cardiotoxic drug with the additional action of being an abortifacient. It's overdose may lead to an echolic action (contract the pregnant uterus)

Potassium carbonate is a caustic alkali with a predominant action of corrosive due to its high pH.

Poison	Mechanism
A.Zinc Chloride	2. Irritant
B.Chloral hydrate	4. Stupefying
C. Quinine	1. Abortifacient
D.Potassiumcarbonate	3. Corrosive

Forensic Medicine NEET 2018

Question 1:

Which is the first organ to putrefy?

a) Brain

b) Heart

c) Kidney

d) Prostate

Question 2:

Which of the following constitutional article is not related to children?

a) 21A

b) 24

c) 45

d) 42

Question 3:

McNaughten's rule falls under which Section?

a) BNSS Section 85 (CrPC 83)

b) BNS Section 22 (IPC 84)

c) BNS Section 219 (IPC 184)

d) BNSS Section 204 (CrPC 184)

Question 4:

Luminescent stools are associated with which of the following?

a) Mercury

b) Cadmium

c) Phosphorus

d) Oxalic acid

Question 5:

Which of the following poisons cause bluish discoloration of the stomach mucosa on post mortem examination?

a) Oxalic acid

b) Sodium amytal

c) Soneryl

d) Arsenic

Question 6:

Saturnine gout is seen in:

a) Lead poisoning

b) Cadmium poisoning

c) Beryllium poisoning

d) Mercury poisoning

Question 7:

At what age, basiocciput fuses with basisphenoid?

a) 18-22 years

b) 22-25 years

c) 14-16 years

d) 12-14 years

Question 8:

Edmond Locard is most famous for?

a) Study of fingerprints

b) Theory of exchange

c) Stature estimation

d) Forensic ballistics

Question 9:

What is the odor of the mummified body?

a) Odorless

b) Pungent

c) Putrid

d) Offensive

Question 10:

Which metallic poisoning causes proximal convoluted tubule damage with proteinuria and muscle pain ?

a) Arsenic

b) Chromium

c) Mercury

d) Lead

Question 11:

Posthumous child is a

a) Child delivered after the death of parents

b) Child delivered after the death of biological father

c) Child born out of wedlock

d) Fictitious child

Answer Key

Question No.	Correct Option
1	a
2	d

3	b
4	c
5	b
6	a
7	a
8	b
9	a
10	a
11	b

Detailed Explanations

Solution to Question 1:

Among the given options, the first organ to putrefy is the brain.

The order of organ putrefaction (from first to last):

• Larynx and trachea

• Stomach, intestine, spleen

• Liver and lungs

• Brain

• Heart

• Kidney and the bladder

• Prostate and uterus

• Skin, muscle, tendons

• Bones

Solution to Question 2:

Article 42 is not related to children. Article 42 says the state shall make provision for securing just and humane conditions of work and maternity relief.

Constitutional Guarantees that are meant specifically for children include:

• Article 21 A - Right to free and compulsory elementary education in the 6 -14 year age group.

• Article 24 - Right to be protected from any hazardous employment till the age of 14 years.

• Article 39 E - Right to be protected from being abused.

• Article 39 F - Right to equal opportunities and facilities.

• Article 45 - Right to early childhood care and education for all children until they complete the age of 6 years.

Solution to Question 3:

McNaughten's rule falls under Section 22 of the Bharatiya Nyaya Sanhita (IPC 84).

It states, "A person of unsound mind is not liable for an act."

Defect: In Mc Naughthen's rule, only intellectual factors are considered but emotions, delusion, and control of impulses are ignored.

Solution to Question 4:

Luminescent stools are seen in acute phosphorus poisoning. The vomitus and stools will be luminous in the dark.

The stools may give rise to faint fumes, constituting the smoky stool syndrome.

Solution to Question 5:

In sodium amytal poisoning, the capsules stain the stomach mucosa and its contents, turquoise blue. Bluish discolouration of the stomach mucosa is also seen in copper sulphate poisoning.

The appearance of the gastric mucosa in specific poisons:

Poison	Appearance of Gastric Mucosa
Arsenic	Red velvety mucosa
Sulphuric acid	Brown or black with erosion perforation and Carbonization
Carbolic acid	Grey leathery and thickened mucosa
HNO_3	Yellow
HCN	Pink
KCN	Brick red to brown
Phosphorous	Yellow or grey white
Sodium amytal, Copper sulphate	Blue

Solution to Question 6:

Saturnine gout is seen in chronic lead intoxication or poisoning.

The features of saturnine gout are the same as primary gout but it affects the knee more than the first metatarsophalangeal joint and tophi rarely develop. Lead exposure should be suspected if a patient presents with the triad of:

• Saturnine gout

• Hypertension

• Renal insufficiency (renal failure and interstitial fibrosis)

Solution to Question 7:

Basiocciput fuses with basisphenoid around 18-22 years.

Solution to Question 8:

Edmond Locard is famous for his theory of exchange or Locard's exchange principle.

This theory states that every contact leaves a trace. When two objects come into contact with one another, each will take something from the other object or leave something behind.

Locard is also renowned for his contribution to the improvement of dactylography, an area of study that deals with fingerprints. He developed the science of poroscopy, the study of fingerprint pores, and the impressions produced by these pores.

Solution to Question 9:

The mummified body is odorless.

Solution to Question 10:

Arsenic poisoning causes PCT damage with proteinuria and muscle pain.

Nephropathy is caused by most of the heavy metals. However, muscle pain is seen with arsenic poisoning.

Salient features of Arsenic poisoning:

Acute poisoning: Resembles cholera; garlicky odor

Chronic poisoning: Following features are seen:

• Raindrop pigmentation seen on temples, eyelids and neck

• Aldrich-Mee's lines (transverse white striae on the fingernails)

• Hyperkeratosis of palms and soles

• Painful paraesthesia of hands and feet

• Red velvety appearance of stomach

• Blackfoot disease - a gangrenous condition affecting the extremities

Tests to detect Arsenic poisoning:

• Marsh test and Reinsch test

Treatment:

Freshly prepared hydrated ferric oxide.

BAL(British Anti Lewisite)

Penicillamine(If BAL not available)

Note: Mercury does not cause muscle pain though it commonly causes PCT damage.

Mee's lines- Transverse white striae of the fingernails

Hyperkeratosis of palms and soles

Rain drop pigmentation

Solution to Question 11:

The child delivered after the death of the biological father, or a child delivered after the death of the biological mother, usually when delivered by a cesarean section is a posthumous child.

Illegitimate child is a child born out of wedlock i.e. when the parents are not married.

Supposititious child or fictitious child is when a woman pretends to go through pregnancy and delivery. Later, she produces a living child as her own for obtaining money or for the purpose of claiming the property.

Forensic Medicine NEET 2019

Question 1:

Which Section of the IPC/BNS deals with the punishment of perjury?

a) 191/ BNS 227

b) 192/ BNS 228

c) 193/ BNS 229

d) 197/ BNS 234

Question 2:

Gastric lavage can be performed in case of poisoning by:

a) Sulphuric acid

b) Strychnine

c) Carbolic acid

d) Kerosene oil

Question 3:

Frotteurism is _____

a) Obtaining sexual pleasure by wearing clothes of opposite sex

b) Desire to seek surgery to become member of opposite sex

c) Sexual gratification by rubbing his private parts against another person

d) Exposure of one's genitals to an unsuspecting stranger

Question 4:

Identical twins have similarities in all of the following except:

a) Blood group

b) DNA fingerprinting

c) Fingerprint pattern

d) Iris colour

Question 5:

Which of the following statements is true about battered baby syndrome?

a) Skin lesions and bruises of various ages are seen on the body

b) It occurs due to the tendency of a child to fall repeatedly

c) Multiple stab wounds are seen

d) The syndrome is a result of firearm injuries

Question 6:

Which of the following tests is based on the principle that prior knowledge of the events/activity will influence the suspect's reaction?

a) Truth Serum

b) Polygraph

c) Brain Mapping

d) Hypnosis

Question 7:

In alleged history of gunshot injury, there is burning, blackening, tattooing around the wound. The injury is:

a) Close shot entry wound

b) Close shot exit wound

c) Distant shot entry wound

d) Distant shot exit wound

Question 8:

Rigor mortis occurs first in:

a) Eyelids

b) Intestines

c) Myocardium

d) Neck

Question 9:

A patient died from bronchopneumonia following an intracerebral hemorrhage caused by cerebral metastases from a squamous cell carcinoma of the left main bronchus. What should be recorded in the line I (c) of the death certificate?

a) Bronchopneumonia

b) Cerebral metastasis

c) Intracerebral hemorrhage

d) Squamous cell carcinoma of the left main bronchus

Question 10:

Which of the following acts as a physiological antidote to scorpion venom?

a) Physostigmine

b) Norepinephrine

c) Prazosin

d) Atropine

Question 11:

Which of the following is commonly used for preservation of viscera in toxicology autopsy?

a) Glycerin

b) Rectified sprit

c) Formalin

d) Saturated salt solution

Answer Key

Question No.	Correct Option

1	c
2	c
3	c
4	c
5	a
6	c
7	a
8	c
9	b
10	c
11	d

Detailed Explanations

Solution to Question 1:

The punishment of perjury is dealt with under IPC 193, has now been updated to BNS 229.

Perjury is defined as the offence of wilfully telling an untruth or making a misrepresentation under oath.

Solution to Question 2:

Gastric lavage can be performed in case of carbolic poisoning.

In carbolic acid poisoning, if the patient survives for 48 hrs, there will be carboluria followed by anuria. The victim will pass dark, smoky urine which soon turns olive green on standing due to oxidation of hydroquinone and pyrocatechol.

It is a corrosive acid, however, it hardens the GI mucosa producing a leather bottle stomach and anesthetizes the nerve endings. This allows for gastric lavage.

Contraindications of gastric lavage:

• Corrosives poisoning (except carbolic acid)

• Convulsants/CNS stimulants

• Comatose patients

• Volatile poison

• Upper GI pathology (eg. esophageal varices)

• Hypothermic patient

• Severe heart disease

• Advanced pregnancy

Sulphuric acid and kerosene are corrosives and strychnine is a convulsant. Hence gastric lavage is not done in these poisonings.

Solution to Question 3:

• Frotteurism is obtaining sexual gratification by contact with another person (rubbing his private parts against female buttocks, breasts, thighs etc. in crowds etc). It is punishable under Section 292 BNS (290 IPC, with a fine of up to 200 Rs). Under the BNS, it is permissible to be fined up to Rs. 1000.

• Eonism (also known as Transvestism) is a sexual perversion where a person's whole personality is dominated by a desire to be identified as a member of the opposite sex. Usually seen in males who obtain sexual pleasure by wearing female clothes. There is no hormonal/genital abnormality.

• Transsexualism: It is the desire to become members of the opposite sex and seek surgery for anatomical alterations.

• Exhibitionism: It involves behaviours over a period of 6 months or more, involving the exposure of one's genitals to an unsuspecting stranger. This is a punishable offence. Section 296 BNS (294 IPC with imprisonment up to 3 months or fine.)

Under the BNS, this is punishable with imprisonment up to 3 months, or a fine of Rs. 1000, or with both.

Solution to Question 4:

Fingerprint pattern differs in identical twins.

Fingerprints are formed during intrauterine life due to friction ridges. These ridges are created by pressure on the fingers from touching the surroundings. Hence, it is unique to every individual.

Solution to Question 5:

Skin lesions and bruises of various ages are a classical feature of a child with battered baby syndrome.

A battered child is one who has received repetitive physical injuries as a result of non-accidental violence, produced by a parent or guardian, there may also be deprivation of nutrition, care, and affection.

The syndrome must be considered in any child:

• In whom the degree and type of injury varies with the history given

• When injuries are of different ages and in different stages of healing

• When there is a purposeful delay in seeking medical attention despite serious injury

• Who exhibits evidence of fracture of any bone, subdural hematoma, failure to thrive, soft tissue swelling or skin bruising

• Who dies suddenly

Skeletal survey, which includes the following, should be done in a case of suspected child abuse:

• AP views of the entire skeleton

• Dedicated views of hands and feet

• Lateral views of appendicular skeleton, skull and spine

Solution to Question 6:

Brain Mapping is based on the principle that if the suspect has prior knowledge of the event/activity being described it will reflect in the suspect's brain wave responses.

It is a technique that measures recognition of familiar stimuli by measuring electrical brain wave responses(P300) to words, phrases, or pictures that are presented on a computer screen.

Truth serum : This is based on the principle, that at a point very close to unconsciousness, the subject will be mentally incapable of resisting questioning and will be incapable of lying.

Polygraph : It is based on the theory, that when a person lies in the answer to a question, if he/she experiences fear, the emotion of fear will result in certain physiological changes.

The physiological changes are commonly observed are :

• Relative rise in blood pressure

• Pulse rate increases

• Slowing down of breathing/erratic breathing

• Suppression of involuntary muscular movements

• Increased activity of sweat glands

Solution to Question 7:

Based on the ballistics, the gunshot injury is close shot entry wound.

• The main objective of study of a firearm wound (Ballistics) is to establish the range and direction of firing, the ultimate purpose is to determine the weapon used for firing.

• Firearm wounds are fundamentally of two mains types: wounds of entry and wound of exit. The wound of entry is the wound through which the bullet or pellets enter the body, and the wound of exit is the wound through which the bullet emerges out of the body.

• To determine the Range of firing, following parameters are used:

• Entry wound size and shape

• Blast effect – Muzzle end imprint and edge eversion

• Blackening, Tattooing, Singeing of hair, Abrasion collar and Grease collar

• In the above options, Hair should be preserved in order to determine the range of firing which can be examined for carbonisation/dessication in close range, also from lead deposition around the wound to assess range of firing.

Characteristics	Contact shot	Close Shot	Near Shot	Distant Shot
Range	In-touch with skin	2.5-7.5 cm	30-60 cm	>60 cm
Size	Largest size	Bullet size	Smaller	Smallest
Shape	Stellate	Circular	Circular	Circular
Blast effect				
Muzzle end imprint	+ve	-ve	-ve	-ve
Edge eversion	+ve	-ve	-ve	-ve
Other findings				
Blackening	-ve	+ve	-ve	-ve
Tattooing	-ve	+ve	+ve	-ve
Singeing	-ve	+ve	-ve	-ve
Abrasion collar	-ve	+ve	+ve	+ve
Grease collar	-ve	+ve	+ve	+ve

Solution to Question 8:

Rigor mortis occurs first in the myocardium.

The order of appearance of rigor mortis: Left chamber of heart is affected first. Other muscles are affected in the given order:

• Involuntary muscles: Heart

• Eyelids

• Neck and lower jaw

• Face

• Chest

• Upper extremities

• Abdomen

• Legs

• Fingers and toes

Note- Rigor mortis first occurs in the myocardium but is first seen in eyelids. Here the question is first occurs in and the answer is myocardium.

Solution to Question 9:

Cerebral metastasis will be filled in the line I (c) of the death certificate, as one of the antecedent causes in the order of the events that took place.

The death certificate will be filled in the order of the events that took place, as follows:

Cause of Death		Approximate interval between onset and death
I Disease or condition directly leading to death*	(a) Bronchopnemonia	
	due to (or as a consequence of)	
Antecedent causes Morbid conditions, if any, giving rise to the above cause, stating the underlying condition last	(b) Intracerebral hemorrhage	
	due to (or as a consequence of)	
	(C) Cerebral metastasis	
	due to (or as a consequence of)	
	(d) Squamous cell carcinoma of the left main bronchus	
II Other significant conditions contributing to the death, but not related to the disease or condition causing it.		

**This does not mean the mode of dying e.g. heart failure, respiratory failure. It means the disease, injury, or complication that caused death.*

Source : Reference No. 6

Fig. 1 : International form of medical certificate of cause of death.

Solution to Question 10:

Prazosin acts as a physiological antidote for scorpion venom.

The venom is a clear, colorless, proteinous toxalbumen, having a hemolytic and neurotoxic effect. Its toxicity is more than that of snakes, but only a small quantity is injected. The venom is a potent autonomic stimulator, resulting in the release of massive amounts of catecholamines from the adrenals.

If the scorpion has hemolytic venom, the reaction is mainly local and simulates the viper snake bite. The scorpion sting will have only one hole in the center of the reddened area. The extremity will have edema, pain, and reddening. This usually lasts for one to two hours.

The symptoms produced by a neurotoxic venom are similar to a cobra bite. There is usually no marked reaction in the local area. Nausea, vomiting, extreme restlessness, fever, various types of paralysis, cardiac arrhythmias, convulsions, coma, respiratory depression, and death may occur within hours from pulmonary edema or cardiac failure.

Prazosin can counteract the alpha-adrenergic effects of autonomic stimulation, such as hypertension.

Solution to Question 11:

A saturated salt solution is used commonly for the preservation of viscera in toxicology autopsy.

Rectified spirit is indicated for most of the poisons, exceptions being alcohol, kerosene, phosphorus, paraldehyde, carbolic acid, acetic acid, etc.

Forensic Medicine NEET 2020

Question 1:

A deceased male was brought to the morgue for a ■post-mortem examination. His relatives reported that he had a tattoo over his skin, which was not found on examination. Which of the following structures would you examine?

a) Lymph nodes

b) Liver

c) Arteries

d) Skin

Question 2:

Which Section of the IPC/BNS deals with the punishment of perjury?

a) 191/ BNS 227

b) 192/ BNS 228

c) 193/ BNS 229

d) 197/ BNS 234

Question 3:

The most common drug which causes physical dependence is?

a) Ketamine

b) Heroin

c) LSD

d) Phencyclidine

Question 4:

A 5-year-old child with a ■history of perianal pain is brought to the OPD. On subjecting the specimen from the perianal region to a test, it produces yellow rhombic crystals of spermine picrate. What is the test done?

a) Barberio test

b) Florence test

c) Takayama test

d) Teichmann test

Question 5:

In the case of rape of a young child, the hymen is usually:

a) Ruptured since it is superficially situated

b) Ruptured since it is very thin

c) Unruptured since it is deeply situated

d) Unruptured since it is highly elastic

Question 6:

In the civil negligence case against a doctor who bears the onus of proof?

a) Patient

b) Police not under rank of sub inspector

c) Magistrate

d) Doctor

Question 7:

Which of the following poison can be found even in skeletonised remains of the body?

a) Arsenic

b) Carbon monoxide

c) Hydrogen sulphide

d) Phosphorus

Question 8:

A child who is a victim under POCSO act is brought to the department of forensic medicine for age estimation. The X-ray image of the hand is shown below. What is the likely age of the child?



a) 4 years

b) 7 years

c) 10 years

d) 13 years

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	a
5	c
6	a
7	a
8	d

Detailed Explanations

Solution to Question 1:

The lymph nodes should be examined in this case.

Some amount of material or dye used for creating tattoo marks is transported through lymph to the regional lymph nodes. Hence, in cases where a tattoo mark has disappeared naturally or has been removed, the lymph nodes must be examined.

Solution to Question 2:

The punishment of perjury is dealt with under IPC 193, has now been updated to BNS 229.

Perjury is defined as the offence of wilfully telling an untruth or making a misrepresentation under oath.

Solution to Question 3:

Heroin is the most common drug which can cause physical dependence.

Physical dependence is defined is a condition where withdrawal of a drug after its prolonged use causes physiological and emotional illness, often referred to as the withdrawal syndrome. Neither physical dependence nor withdrawal symptoms occur with hallucinogens. Although it is necessary to note that a user can develop a psychological dependence on the insight-inducing experiences of episodes of hallucinogen use. Hence, all 3 hallucinogens - ketamine, LSD, and Phencyclidine (PCP) does not cause physical dependence.

According to the world drug report 2019, released by the United Nations office on drugs and crime (UNODC), the most common drugs used across the world (in decreasing order) include:

• Cannabis

• Opioids

• Amphetamine and prescription stimulants

• Ecstacy

• Cocaine

Among the opioids, the commonly used group of drugs are opiates which include drugs like opium, morphine, and heroin.

Solution to Question 4:

The test done in the scenario is the Barberio test.

It is used to detect seminal fluid. When saturated aqueous or alcoholic solution of picric acid is added to the spermatic fluid, yellow needle-shaped rhombic crystals of spermine picrate are produced.

Florence test:

• Used to detect: semen

• Stain used: 10% hydrochloric acid and Florence solution (potassium iodide, iodine, and water)

• Finding: If semen is present, dark brown rhombic or needle-shaped crystals of choline iodide appear immediately

Takayama test (Haemochromogen crystal test):

• Used to detect: bloodstains

• Stain used: Takayama reagent

• Finding: If bloodstain is present, pink feathery crystals of haemochromogen or reduced alkaline haematin appear in a few minutes

Teichmann test (Haemin Crystal Test):

• Used to detect: bloodstains

• Stain used: glacial acetic acid and a small crystal of sodium chloride

• Finding: If bloodstain is present, faint yellowish-red to brownish-black rhombic crystals of haemin or haematin chloride appear. When hydrogen peroxide is added, the haematin crystals give bubbles of gas

Solution to Question 5:

In the case of a young child, the hymen is deeply situated and usually intact after intercourse. As the vagina is very small, it is impossible for the penetration of the adult organ to take place.

In children, there are few or no signs of general violence, as the child is unaware of the assault and incapable of resisting. A few indicators for inquiry are:

• Pain on urination and/or defecation

• Abdominal pain

• Inability to sleep

• Sudden withdrawal from peers/adults

• Feelings of anxiety

• Nervousness

• Helplessness

• A feeling of ending one's life

Solution to Question 6:

In the case of civil negligence, the burden of proof is on the patient.

The question of civil negligence arises when a suit is filed in a civil court:

• By a patient or his/her relatives against the doctor for getting compensation from the doctor if he/she has suffered an injury due to the doctor's negligence.

• By a doctor against the patient or his/her relatives to collect the fees, the payment of which has been refused due to alleged professional negligence by the doctor.

In either case, the patient has to prove that the doctor has committed professional negligence.

To prove negligence, all the following four elements must be present:

• Duty - A duty of care by the doctor must exist.

• Dereliction - The physician must maintain the standard of a 'prudent physician' under similar circumstances. If the physician fails to maintain the standard of care, it is called dereliction.

• Direct causation - The negligent act or omission must directly cause the injury without any intervening cause.

• Damage - The negligent act must cause damage of a type that would have been foreseen by a reasonable physician.

Solution to Question 7:

Arsenic can be found even in skeletonized remains of the body.

It becomes fixed in the cancellous tissues of the long bones by the conversion of phosphates to arsenates. The elimination of arsenic in this form is much slower and can be detected in the bones for several years after death.

Heavy metals such as lead, copper, mercury, including metalloids like arsenic and antimony, may be found in bones for a prolonged period in cases of chronic poisoning. They are also deposited in hair, nails, etc. For these reasons, when heavy metal poisoning is suspected, these specimens are collected in addition to the routine viscera and body fluids.

Solution to Question 8:

The child's age is likely to be 13 years.

In the given X-ray image of the wrist and hand, all the carpal bones are seen. The lower epiphysis of the first metacarpal has not united with the shaft. This suggests that the age of the child is between 8 to 15 years.

Additionally, the presence of the hook of the hamate in the radiograph suggests that the age is between 13-14 years.



The age of ossification of the carpal bones:

The age of fusion of joints:

Carpal bone	Age of ossification
Capitate	1-3 months
Hamate	2-4 months
Triquetrum	2-3 years
Lunate	2-4 years
Scaphoid	4-6 years
Trapezium	4-6 years
Trapezoid	4-6 years
Pisiform	8-12 years
Hook of the Ham ate	13-14 years

Bone/ joint	Maximum age at fus ion
Elbow joint centersMetacarp als	16 years
Wrist joint centers	19 years
Hip bone centers	22 years

Forensic Medicine INI-CET 2021

Question 1:

Arrange the following in the correct order of symptomatology seen in iron toxicity:

a) 2 -> 3 -> 1 -> 4

b) 2 -> 3 -> 4 -> 1

c) 3 -> 2 -> 4 -> 1

d) 3 -> 4 -> 1 -> 2

Question 2:

Match the following weapons with their possible injury type

a) A-4,B-3,C-1,D-2

b) A-1,B-3,C-4,D-2

c) A-1,B-2,C-4,D-3

d) A-4,B-1,C-3,D-2

Question 3:

Match the following postmortem findings and their time of appearance:

a) A-3, B-2, C-1, D-4

b) A-3, B-1, C-2, D-4

c) A-4, B-2, C-1, D-3

d) A-1, B-2, C-3, D-4

Question 4:

Which of the following is true about the the cause of the given finding?

a) Lightning strike

b) Electrocution

c) Marbling

d) Trickling down of acid

Question 5:

A 24-year-old man who was accused of murder was taken into police custody. As a part of the interrogation, the prisoner was kept in a prone position with both his wrists and ankles bound behind his back and secured with a rope. What is this method known as?

a) Choke/carotid hold

b) Carotid sleeper hold

c) Hog tie

d) Bar arm

Question 6:

An 18-year-old female was sexually assaulted by a 20-year-old male. There was presence of white stains on the clothes of the victim. Barberio's test was performed on the stains. Which of the following is detected by this test?

a) Spermine

b) Acid phosphatase

c) Choline

d) Fructose

Question 7:

Match the following with the respective ages of fusion:

a) A-1, B-2, C-3, D-4

b) A-4, B-1, C-2, D-3

c) A-4, B-3, C-1, D-2

d) A-1, B-4, C-3, D-2

Question 8:

A skull was recovered from a forest. According to inquest papers, a girl had gone missing 15 days back. The skull was sent to the forensic research lab. Which of the following would identify it as a female skull?

a) 1 and 4 only

b) 2, 3 and 4

c) 1, 3 and 5

d) 3, 4 and 5

Question 9:

The dead body of a 20-year-old man found in the sea was brought in for postmortem examination. Which of the following findings would you see in seawater drowning?

a) 1 and 3

b) 1 and 4

c) 2, 3 and 5

d) 2, 4 and 5

Question 10:

A pelvis is brought for forensic examination. Which of the following characteristics will identify the given pelvis as a male?

a) 2,3,4,5

b) 4,5,6

c) Only 5

d) 1 and 3

Question 11:

Match the following in relation to fingerprint changes.

a) 1-e, 2-c, 3-b, 4-a, 5-d

b) 1-d, 2-b, 3-c, 4-e, 5-a

c) 1-c, 2-a, 3-d, 4-b, 5-e

d) 1-d, 2-c, 3-a, 4-e, 5-b

Question 12:

A patient came to the casualty after a road traffic accident. A lacerated wound of 4x1 cm, which is bone-deep, is noted over the scalp of the forehead. Which of the following is the correct mechanism of this lacerated wound?

a) Shearing force

b) Impact of a semi sharp-edged object

c) Overstretching of skin and subcutaneous tissue

d) Tissue crushed between weapon and underlying bone

Question 13:

The friend and victim had alcohol and got into an argument over a common girlfriend, following which the friend kills the victim by sitting on his chest and covering his nose and mouth. This is an example of?

a) Burking

b) Smothering

c) Traumatic asphyxia

d) Overlaying

Question 14:

A person comes to the OPD with a burn injury to the right thigh. On examination, a soddened injury is seen along with erythema, blisters, and a line of demarcation that extends down the limb. What is the most likely cause of the injury?

a) Electric burn

b) Chemical burn

c) Dry heat

d) Moist heat

Question 15:

Which of the following rules are used in case as tests of insanity?

a) 1, 2, 3 and 4

b) 2, 3, 4 only

c) 1, 2 only

d) 3, 4 only

Question 16:

A person working in a dye factory presented with nausea, vomiting, dark bloody stools, conjunctivitis, and a burning sensation in the throat and stomach. Which poisoning do you suspect in this case?

a) Potassium permanganate

b) Arsenic

c) Thallium

d) Lead

Question 17:

Following a blast, a man was brought from the scene to the emergency with multiple small puncture wounds to the right side of the chest, below the nipple, and multiple small abrasions and bruising over the left lower abdomen. Which of the following mechanisms is responsible for these injuries?

a) Flying missiles

b) Blast of air

c) Explosion burns

d) Building collapse

Question 18:

On conducting the autopsy on a victim of hanging, the ligature mark is seen at the lower 1/3rd of the neck. The victim is seen to have a protruded tongue. He was found with his head hanging to his left side with saliva dribbling from the left angle of his mouth. Both the pupils are dilated and the right eye is open. Compression of which of the following structures is the probable reason for an open right eye in this case?

a) Right internal jugular vein

b) Cervical sympathetic chain

c) Left vagus nerve

d) Right internal carotid artery

Answer Key

Question No.	Correct Option
1	c
2	a
3	a
4	a
5	c
6	a
7	b
8	c
9	b
10	c
11	a
12	d
13	a
14	d
15	c
16	b

17	a
18	b

Detailed Explanations

Solution to Question 1:

The correct order of symptomatology in iron toxicity is 3 -> 2 -> 4 ->1.

Symptoms of iron toxicity are divided into 4 stages:

• Stage 1 (few hours after ingestion): vomiting, abdominal pain, gastrointestinal bleeding, shock, acidosis, and coma

• Stage 2 (6-24 hours): Asymptomatic phase

• Stage 3 (24-48 hours): Metabolic acidosis, jaundice, hypoglycemia, shock, coma with hepatic and renal failure.

• Stage 4 (1-2 weeks): Late complications such as gastric stricture and pyloric stenosis.

Iron exerts its toxic effects by increasing capillary permeability, release of hydrogen ions, inhibition of mitochondrial function, and direct corrosive action on the gastric mucosa. Ingestion of 20 to 30 grams is considered to be fatal and the fatal period is around 24-30 hours. Desferrioxamine is the mainstay of treatment with supportive gastric lavage, administration of egg and milk to form iron-protein complexes, oral MgOH solution, and hemodialysis in severe cases.

Postmortem changes in iron toxicity include:

• Hemorrhagic necrosis of gastric mucosa with perforation of gastric or jejunal wall

• Acute hepatic necrosis

• Degenerative renal tubular changes

Solution to Question 2:

For the above-given weapons, possible injury type is matched as under:

Mechanism of injury with the above weapons

Axe - it is a heavy weapon that may or may not be sharpened. Injury is produced by chopping action of the weapon and it may be in a spectrum of an incised wound to laceration depending upon the sharpness of the blade.

RTA - rubbing of epidermis tangential to a rough or hard surface gives rise to graze abrasion also called brush burn. It can occur in an RTA after a pedestrian is being hit by a vehicle and the victim is slid/grazed for some distance before he comes to a halt.

Blade - it gives rise to a clean-cut wound which is usually deepest at its starting. Tailing of the wound indicates the direction of injury. Length is the biggest parameter of this type of wound.

Lathi- It is a blunt force injury giving rise to a patterned abrasion with 2 parallel lines and a normal unbruised area on site of impact or stick/lathi. The lathi forces the blood from the site of impact to either side which gives rise to tramline appearance.

A. Axe	4. Chop wounds
B. RTA	3. Graze abrasions
C. Blade	1. Incised wound
D. Lathi	2. Tram track bruise

The above image shows a chop wound inflicted on scalp.



The above image shows a schematic illustrating heaping up of epidermis in a graze abrasion or brush burn



The above image shows production of a tramline bruise associated with a lathi strike

Solution to Question 3:

The correct match is A-3, B-2, C-1, D-4.

Adipocere formation starts in 4-5 days.

Relaxation of urethral sphincter by 2-3 days.

Marbling of veins starts in 24 hours.

Liquefaction of tissues by 5-10 days.

In postmortem marbling, blood in the vessels is haemolysed to sulfmeth-hemoglobin, which stains the vessel walls and the adjacent tissues, giving rise to a marbled appearance. The marbling of the skin begins by about 24 hours and becomes prominent in 36–48 hours after death. The image below shows postmortem marbling which is distinctly appreciable in the superficial veins of the abdomen, shoulders, chest, and inguinal region.

Urethral sphincter relaxation is of two types;- Primary and Secondary. Primary relaxation occurs before rigor mortis sets in. Secondary relaxation occurs at the time of decomposition i.e. around 2-3 days.

Adipocere formation, also called corpse wax, grave wax, or mortuary wax is a late sign of death. A wax-like organic substance is formed by the anaerobic hydrolysis or hydrogenation of fat in tissues by bacteria like Clostridium welchii. It has a greyish-white appearance with a distinctive offensive or sweetish smell. In a country like India, adipocere formation begins within 4-5 days and takes several weeks to complete.

Liquefaction or colliquative changes begin 5 days after death and become prominent in 10 days. Soft tissues change into semisolid masses which are black and get separated from bones and fall off.

Note: The first visible sign of putrefaction is greenish (due to sulfmeth-hemoglobin) discolouration of the anterior abdominal wall, usually in the right iliac fossa.

Solution to Question 4:

The given finding is known as Lichtenberg's figures or filigree burns which are pathognomonic for lightning injury. It is a superficial, thin, tortuous, fern-like pattern of erythema in the skin. It indicates the path taken by the electrical discharge and tends to follow skin creases and long axis of the body.

They are also known as arborescent burns or feathering as they have a resemblance to the branches of a tree. Lichtenberg's figures occur due to the rupture of small vessels. They are usually seen over the shoulder or flank within an hour of injury and disappear in 1-2 days.

Medico-legal aspect: Its recognition may be lifesaving in the unaccompanied comatose patient and is important because even delayed resuscitation of lightning victims can be very successful.

Lesions seen in electrical burns or electrocution are:

• Joule burns: When the skin has been in firm contact with an electrical conductor, this may split the layers of the epidermis or the epidermal-dermal junction and produce a raised blister, called Joule burns.

• Spark lesion: When the contact is less firm, an air gap (albeit narrow) exists between skin and conductor, the current jumps the gap as a spark. This causes the outer skin keratin to melt over a small area. On cooling, the keratin fuses into a hard brownish nodule, usually raised above the surrounding surface, the so-called spark lesion. The heat generated by the current may melt the calcium phosphate, which is radiologically seen as typical round dense foci known as bone pearls or wax drippings.

• Crocodile skin effect is caused by high-tension current burns. In high-voltage burns, such as those sustained from high-tension grid transmission cables, where the voltage is in the multi-kilovolt range, sparking may occur over many centimeters. This can cause multiple spark lesions giving rise to a crocodile-skin effect.

The image given below shows Joule burns

The image given below shows high voltage burns due to electrocution causing multiple spark lesions. Inset shows crocodile skin effect

Image below shows electrical marks



Marbling is a postmortem finding seen when blood in the vessels is haemolysed to sulfmeth-hemoglobin, which stains the vessel walls and the adjacent tissues, giving rise to a marbled appearance.

Solution to Question 5:

This type of physical restraint in which a person is placed in prone position with his wrists and ankles bound behind his back and secured by a rope is known as hog-tying or total appendage restraint procedure. This can cause breathing difficulty and the person can become unresponsive. Rarely, death may occur due to asphyxia, soon following the physical restraint.



Carotid hold also known as choke hold or bar arm involves placing the forearm straight across the front of the neck. The free hand grips the wrist and pulls it back causing obstruction of the airway and carotids resulting in immediate unconsciousness. Fracture of the hyoid or larynx can occur with greater force. Pressure on the carotid bodies can cause sudden cardiac arrest.



In carotid sleeper hold, also known as lateral vascular neck restraint, the arm is placed about the neck with the antecubital fossa centered at the midline of the neck. The free hand grips the wrist and the arm is pulled backward, compressing carotid arteries and jugular veins but not vertebral arteries or trachea. The person loses consciousness in 10 to 15 seconds and regains consciousness within 10 to 20 seconds upon releasing the pressure. Here, the neck structures are not damaged. Death may rarely occur due to catecholamine release or hypoxia which can trigger myocardial infarction or dysrhythmias in patients with ischemic heart disease.



Solution to Question 6:

Barberio's test detects the presence of spermine which is present in prostatic secretions and thus helps in the identification of seminal fluid.

In Barberio's test, a saturated aqueous or alcoholic solution of picric acid is added to the spermatic fluid. This produces yellow needle-shaped rhombic crystals of spermine picrate.

Option B: Acid phosphatase test is a quantitative test for the presence of acid phosphatase enzyme. High levels of acid phosphatase from prostatic secretions are present in seminal stains (The normal vaginal secretions have low levels of acid phosphatase). Positive reactions are found till 36 hours after which the enzyme levels fall and disappear within 72 hours. In the absence of demonstrable sperms or aspermia, this test can be considered conclusive.

Option C: Florence test detects choline which is secreted by seminal vesicles. The suspected seminal stain is extracted with 10% hydrochloric acid and a drop of it is placed over a slide. A coverslip is placed over it and a drop of Florence solution (potassium iodide, iodine, and water) is made to run under the coverslip. If semen is present, dark brown rhombic or needle-shaped crystals of choline iodide appear immediately. The test is not confirmatory for the presence of semen, however, a negative reaction rules out the presence of semen.

Option D: Fructose acts as an energy source for sperm. Its presence in seminal fluid indicates a patent efferent ductal system, as fructose will be absent in its obstruction.

The absolute proof of semen is finding at least one unbroken spermatozoon or electrophoretic LDH isoenzyme detection of sperms.

Solution to Question 7:

The correct match is A-4, B-1, C-2, D-3.

The respective ages of fusion are:

• The medial end of clavicle at 21-25 years

• Sacrum as a single bone at 22-25 years

• Crista scapulae at 50 years

• Lambdoid suture at 45 years

The secondary ossification center at the medial end of the clavicle appears at 18-19 years of age and fuses by 20-22 years of age. This occurs 2 years earlier in females. Thus the complete fusion of the medial end of clavicle can be used for establishment of age of 21 years in medicolegal cases.

The 5 sacral vertebrae are separated by cartilage till puberty. Fusion of epiphysis begins after puberty and ossification start from below upwards. Sacrum becomes a single bone by the age of 25 years.

Ossification of scapulae:

• 30-35 years: lipping starts on ventral margin of glenoid cavity

• 35-40 years: irregular lipping around the clavicular facet and inferior surface of acromion process.

• 45 years: localized bony atrophy

• 50 years: Cristae scapulae

Lambdoid suture starts closing endocranially at 25-35 years and completes fusion by 45-50 years. Endocranial fusion occurs before ectocranial fusion.

The sagittal suture is used for age determination even though age estimation by skull sutures is not considered to be very reliable. The last skull suture to fuse is the squamous temporal suture, which fuses after 80 years of age.

Age	Age determination
<5 years	Appearance of ossification ce nters
5-16 years	Analysis of hand and wrist
>16 years(male)	Complete fusion of distal radi us or ulna
>16 years(femal e)	Complete fusion of distal uln a
>18 years(male)	Complete fusion of distal fem ur or proximal tibia
>18 years(femal e)	Complete fusion of iliac crest
>21 years	Complete fusion of medial en d of clavicle

Solution to Question 8:

Larger frontal and parietal eminences, smooth glabella, and narrow mastoid are features of a female skull.

Differentiating features between male and female skull:

Note: The features given above develop after puberty and are modified by senility, so are applicable only between the 20th and the 55th year.

Trait	Male	Female
General size	Larger, longer (dolichocrania)	Smaller, rounder (brachycran ia)
Architecture	Rugged	Smooth
Glabella	Rough	Smooth
Supraorbital ridge s	Prominent	Less prominent
Cheek bones	Heavier, laterally arched	Lighter, compressed
Zygomatic arch	More prominent	Less prominent
Mastoid process	Prominent, wide and long	Less prominent, narrow and short

Trait	Male	Female
Occipital region	Marked muscle markings	No marked muscle markings
Frontal eminences	Small	Large
Parietal eminences	Small	Large
Orbits	Squarewith round margins	Roundwith sharp margins
Mandible	Large, broad ramus	Small, small ramus
Chin shape	U-shaped	V-shaped

Solution to Question 9:

In cases of seawater drowning, hypernatremia and myocardial anoxia are seen. Seawater being hypertonic to plasma causes a shift of fluid from the blood to alveoli causing pulmonary edema. There is a net decrease in blood volume causing hemoconcentration and a fall in blood pressure.

This hemoconcentration in seawater drowning causes increased viscosity of blood and myocardial anoxia leading to heart failure. Hyperkalemia is not seen in seawater drowning because RBCs get shrunken in contrast to freshwater drowning.

Freshwater is hypotonic to plasma and is rapidly absorbed from the alveoli into the RBCs resulting in hemodilution and hyponatremia. The RBCs finally swell up and lyse resulting in the release of intracellular potassium causing hyperkalemia. Increased intravascular blood volume combined with hyperkalemia causes arrhythmia.

The table below summarises the cause of death in different types of drowning.

Type of drowning	Cause of death
Freshwater drowningSeawater drowning	Disruption of surfactant
Dry drowning	Laryngeal spasm/Vagal inhibition
Post immersion Syndrome / Secondary drowning	Cerebral anoxiaPneumoniaPulmonary edemaChemical pneumonitisMetabolic acidosis

Solution to Question 10:

Among the given options, a large acetabulum (only 5) is the characteristic of a male pelvis.

Triangular obturator foramen, U-shaped pubic angle, everted ischial tuberosity, large greater sciatic notch, deep preauricular sulcus are characteristics of a female pelvis.

The differences between male and female pelvis are summarized in the table below.

Characteristic	Male pelvis	Female pelvis
General Morphology	Massive, rougher, marked muscle attachments	Less massive, slender, smoother
Overall shape	Deep funnel	Flat bowl
Preauricular sulcus	Narrow,shallow	Broad,deep
Acetabulum	Large, wider, deeper	Small, narrower
Body of pubis	Narrow, triangular, thick	Broad, square
Obturator foramen	Large,oval with base upward	Small,triangularwith apex forwards
Subpubic angle	V-shaped, narrow, (700)	U-shaped, rounded, wide, (900)
Greater sciatic notch	Smaller, narrower, deeper	Larger, wider, shallower
Ischial Tuberosity	Inverted	Everted, more widely separated
Pelvic Brim	Heart-Shaped	Circular or elliptical
Pelvic outlet	Smaller	Larger
Sacroiliac articulation	Large extends to 21/2to 3 vertebrae	Small, oblique, extends to 2 to 21/2vertebrae
Sacroiliac joint surface	Largeand less sharply angulated	L-shapedand elevated anteriorly
Sacrum	Longer, narrower, with more evenly distributed curvature, promontory well marked. Body of first sacral vertebra larger.	Shorter, wider; upperhalf almost straight, curve forward in the lower half, promontory less marked. Body of first sacral vertebra small

Solution to Question 11:

The correctly matched option is 1-e, 2-c, 3-b, 4-a, 5-d.

The study of ridge patterns in the skin is Dactylography/ Galton-Henry system/ Dermatoglyphics. Fingerprints are the impressions of patterns produced by the papillary or epidermal ridges present on the fingertips.

Some changes in fingerprints include:

• Moderate epidermal ridge atrophy and loss of pattern in celiac disease

• Incomplete atrophy of the ridges in dermatitis

• Ridge alteration occurs in eczema, acanthosis nigricans, scleroderma, and dry or atrophic skin

• Permanent impairment of the fingerprint pattern occurs in leprosy, electric injury, and after exposure to radiation

• Distance between the ridges is changed but the pattern is not altered in infantile paralysis, rickets, and acromegaly

Classification of fingerprint ridges:

• Loops: radial or ulnar

• Whorls: concentric, spiral, double spiral, or almond-shaped

• Arches: plain, tented, exceptional

• Composite: They are a combination of 2 or more types

The fingerprint patterns are not inherited and cannot be used to prove paternity. The fingerprint pattern differs even between identical twins.

Edgeoscopy is the study of edges of friction ridges which normally show seven characteristic features. Ridgeology is the study of friction ridges which are composed of the edges of ridge units that vary in size, shape, and alignment.

1. Incomplete atrophy of ridges	e. Dermatitis
2. Loss of pattern with ridge atrophy	c. Celiac disease
3. Altered ridges	b. Scleroderma
4. Permanent loss of fingerprints	a. Radiation
5. Distance between ridges changed but pattern retained	d. Acromegaly



Solution to Question 12:

The description of the wound given in the question suggests a split laceration and the mechanism is the tissue being crushed between the underlying bone (i.e. skull) and the weapon.

A laceration is a tear or a split in the skin, mucous membrane, muscle, or any internal organ produced by blunt force. The margins are irregular and may show minute tears and bruising. Tears that are seen at the ends, diverging away from the main laceration are called swallowtails. In skin lacerations, usually one of the margins overhangs the other. This helps in assessing the direction of the force.

Classification of a laceration:

• Split laceration (incised-like laceration): It occurs when the skin is crushed between two hard objects. For example, during a fall, the scalp gets crushed between the skull from inside and hard ground outside. This results in a wound, which usually mimics an incised one unless examined closely.

• Incised-looking lacerated wound is a wound with the linear splitting of the tissue caused by blunt force. It is seen where the skin overlies an area of the bone where there is scanty underlying subcutaneous tissue such as the scalp, forehead, shin, etc.

• Avulsed laceration: A lesion where the skin separates or peels off by tearing from the underlying tissues. An avulsion is a laceration produced by a shearing force delivered at an acute angle to tear off a portion of a traumatized surface or viscus from its attachments. It may overhang the wound. It is also called flaying. This is common in road traffic accidents. (Option A)

• Tears: Tearing of skin and subcutaneous tissue can occur from localized impact by or against irregular or semi-sharp objects. It may be due to blows by broken glass, or a fall on a rough projected object. (Option B)

• Stretch laceration: It occurs due to the over-stretching of the skin, beyond its elasticity. This is seen in wounds produced from running over by a motor vehicle, and when a fracture produces a sudden deformity of the bone. (Option C)

• Cut laceration: Injury by a heavy sharp-edged instrument. For example, cut from ax, chopper, etc

The image shows a split laceration which is incised looking laceration.



Solution to Question 13:

The given scenario is an example of burking.

Burking is a method of homicidal smothering and traumatic asphyxia where the assaulter kneels or sits on the chest (traumatic asphyxia) and close the nose and mouth with his hands (smothering). This was the method of killing performed by William Burke and William Hare.

Image shows burking

Other options:

Option B: Smothering is a form of asphyxia that is caused by closing the external respiratory orifices either by the hand or by other means or blocking up the cavities of the nose and mouth by a foreign substance, such as mud, paper, cloth, etc.

Below image shows smothering



Option C: Traumatic asphyxia results from respiratory arrest due to mechanical fixation of the chest, by a heavy substance or object. The normal movements of the chest wall are prevented. There is a gross compression of the chest and usually abdomen by a powerful force, due to which chest expansion and diaphragmatic lowering are prevented. For example stampede deaths, crushing by falls of the earth in coal mines, or earthquakes.

Option D: Overlaying or compression suffocation results due to compression of the chest, so as to prevent breathing. It occurs when an intoxicated parent or other person shares a bed with an infant. The thoracic movements are limited and respiratory exchange is either reduced or completely prevented which leads to death. It is always accidental.

Below image below shows overlaying.

Solution to Question 14:

The soddened injury with erythema, blisters, and a line of demarcation that extends down the limb is highly suggestive of a scald injury caused by moist heat.

Scald injury is an injury that results from the application of liquid above 60°C or from steam i.e moist heat. Redness appears immediately followed by blistering in a few minutes. Removal of the blistered skin leaves a raw, pink surface. Blisters are surrounded by a hyperaemic zone. Scalds appear soddened and bleached but the hair does not get singed. The skin does not get blackened or charred.

Scald injuries are limited to the area of contact and are more severe at the initial point of contact. It can occur through clothing, which worsens damage due to a longer period of contact with the hot liquid and a reduced cooling effect of evaporation.

It can be

• Immersion burns: Accidental or deliberate immersion in hot liquid, usually water.

• Splash or spill burns: Usually accidental.

• Steam burns: Exposure to superheated steam.

Clinical degree of scalding:

• Erythema (reddening)

• Vesication (blister formation)

• Necrosis of dermis when deeper layers of skin are involved.

The image below shows a scald burn due to moist heat.

Other options:

Option A: Electric burns are injuries caused due to electric current, characterized by puckering of the skin at the edges of the burns. It is surrounded by an area of pallor. There is no red line at the point of entry and exit. The characteristic marks seen at the entry point are called Joule burns, or electrical burns/marks. They are chalky white, shallow, centrally collapsed blisters with a raised

border.

The image below shows electric marks.



Option B: Chemical burns by acid or alkalis appear as ulcerated patches without blisters. The hair is not singed and the red line of demarcation is absent. Sometimes, the burn shows distinct coloration and is usually uniform in character. Strong acids produce dark leathery burns on the skin. Strong alkalis cause the skin to slough and leave moist, slimy, greyish areas.

Option C: Dry heat injuries are caused by a flame, via direct conduction or radiation to the skin. These are characterized by singeing of the hair and blackening of the skin. Vesication may or may not be present.

Solution to Question 15:

Durham's rule and Curren's rule are used in cases as tests for insanity.

Durhams's rule states that an accused person is not criminally responsible if, his unlawful act is the product of mental disease or mental defect.

Curren's rule states that an accused person is not criminally responsible if, at the time of committing the act, he did not have the capacity to regulate his conduct to the requirements of the law as a result of mental disease or defect.

In India, the law of Criminal responsibility is under Section 84 I.P.C. It states that "Nothing is an offense which is done by a person who, at the time of doing it, by reason of unsoundness of mind, is incapable of knowing the nature of the act, or what he is doing is either wrong or contrary to the law". This is slightly different from McNaughten's rule followed in Britain.

McNaughten's rule states that every man is to be presumed to be sane, and possesses a sufficient degree of reason to be responsible for his crimes until the contrary is proved.

Other rules to assess the criminal responsibility of insane persons:

• The Irresistible Impulse

• Federal rule or American Law Institute test (ALT)

• The Brawner rule.

Note: Ashley's rule is used to differentiate males and females based on the combined length of the manubrium and sternum. It is considered to be male sternum when the combined length is >149mm, while if the combined length is <149mm, then it is considered female.

Solution to Question 16:

The given clinical scenario of an industrial worker presenting with nausea, vomiting, dark bloody stools, conjunctivitis, burning in the throat and stomach is suggestive of acute arsenic poisoning.

There are 3 types of arsenic poisoning,

• Gastroenteric type is a common form and resembles bacterial food poisoning. There is a sweetish metallic taste. Features include:

• GIT: Constriction in the throat and dysphagia, burning, and colicky pain in the esophagus, stomach, and bowel occur. Intense thirst and severe vomiting are constant symptoms. Involuntary and frequent dark-colored, stinking, and bloody stools that later become colorless, odorless, and watery resembling rice-water stools of cholera. A garlicky odor of breath and feces may be noted

• Renal: Oliguria, uremia, albuminuria, red cell casts.

• Cardiovascular system: Acute circulatory collapse with vasodilation, increased vascular permeability, ventricular tachycardia, and fibrillation.

• Central nervous system: Headache, vertigo, hyperthermia, convulsions, coma. Peripheral neuropathy is more sensory than motor.

• Death is usually due to circulatory failure.

• Fulminant type: Massive doses of arsenic when rapidly absorbed cause death in one to three hours. Death occurs due to shock and peripheral vascular failure. Arsenic also has a direct action

on the heart muscle. In this type gastrointestinal symptoms are absent.

• Narcotic type: The gastrointestinal symptoms are minimal. There is giddiness, formication, delirium. Arseniuretted hydrogen when inhaled as a gas, destroys hemoglobin and causes hemolysis, hemoglobinuria, and renal failure. Death is instantaneous.

Treatment:

• Repeated stomach washes with warm water and milk.

• Emetics should not be used

• Oral hydrated ferric oxide is given in small doses. It converts toxic arsenic to non-toxic ferric arsenite

• Dimercaptosuccinic acid (DMSA) is preferred, if not available British anti-Lewisite (BAL) is used.

Other options:

Option A: Potassium permanganate causes burning pain from the mouth to the stomach, intense thirst, dysphagia, stridor, and a persistent cough. The oral cavity is discolored, inflamed, and superficially corroded. Soon after poisoning the color is purple-brown, but later it appears coal-black.

Option C: Thallium poisoning is characterized by a triad of symptoms of acute gastrointestinal upset, painful combined polyneuropathy, and hair loss after poisoning. Other symptoms include muscle and joint pain, convulsions, optic neuritis, delirium. Loss of the outer third of eyebrows is significant. Maculo-papular butterfly-shaped skin eruption on the face is characteristic. There may be Mees lines on the nails.

Option D: Acute Lead poisoning causes dry throat, burning abdominal pain, peripheral circulatory collapse, insomnia, paraesthesias, depression, coma, and death. Cerebellar ataxia is common in children with acute lead poisoning. A screening test for lead poisoning is the presence of porphyrinuria, mainly due to coproporphyrin III. Peripheral smear shows basophilic stippling. A Burtonian line may be visible on the gums.

Solution to Question 17:

The given history of multiple small puncture wounds to the right side of the chest, below the nipple, and multiple small abrasions and bruising over the left lower abdomen are highly suggestive of blast injuries sustained due to flying missiles.

Blast injury is a complex type of physical trauma resulting from direct or indirect exposure to an explosion causing a sudden increase or decrease in atmospheric pressure.

T.K.Marshall's (Marshall's triad) described the characteristic injuries associated with bomb explosions. It includes:

• Abrasions

• Bruises/contusions

• Puncture lacerations

Blast injuries are classified as:

• Primary injuries: Effect of transmitted blast waves on gas-containing structures. The external injuries are absent and are characterized by internal injuries. The middle, ear, lungs, and gastrointestinal tract are the most susceptible to this.

• Secondary injuries: Injuries caused by shrapnel and other airborne debris propelled by the explosion. These injuries may affect any part of the body and sometimes result in penetrating trauma. Most casualties are caused by secondary injuries.

• Tertiary injuries: These are caused by blast wind that can throw victims against solid objects.

• Quaternary injuries: All other injuries caused by explosions are not included in the above three categories. eg, flash burns, crush injuries, respiratory effects due to toxic dust, gas, etc.

Option B: Blast of air or blast wave cause primary injuries which involve internal organs without external injury.

Option C: Explosion burns: When the bomb explodes, the temperature of the explosive gases can exceed 2000°C, and the heat radiated momentarily can cause flash burns. It is a quaternary type of blast injury.

Option D: Building collapse can cause crush injuries which is a quaternary blast injury.

Solution to Question 18:

Compression of the cervical sympathetic chain is the probable reason for an open right eye in this case.

Dilated pupils are seen when the ligature knot presses on the cervical sympathetic chain, the eye on the same side may remain open and its pupil dilated termed as le facie sympathique. It indicates antemortem hanging.

Dribbling of saliva is the surest sign of antemortem hanging. This occurs due to the increased salivation before death due to the stimulation of the salivary glands by the ligature or by congestive hypoxia. Stimulation of pterygopalatine ganglion also causes increased salivation.

Dribbling of saliva occurs from the angle of the mouth opposite to the side of the knot. When the knot is on the nape of the neck, it occurs across the middle of the lower lip.

The ligature mark is usually above the hyoid bone. It runs obliquely, backward, upwards, is non - continuous, and towards the point of suspension. The mark is noncontinuous because of a gap at the nape of the neck. When the knot is in an inverted V shape when it is in contact with the skin, because the ligature material extends downward on both sides. The skin is usually depressed/grooved, yellowish-brown, dry, hard, with small abrasions called rope burns at its edges.

Protrusion of tongue results from the pressure on floor of the mouth by ligature. It is usually swollen and blue. Injuries include bite marks with or without underlying small hemorrhages (‘marginal’ hemorrhages).

Tardieu’s spots may be present on the forehead, over the eyelids, under the conjunctiva, and near the temple.

Other options:

Option A- In partial hanging, internal jugular veins are compressed, but vertebral arteries continue to supply blood to the head, resulting in congestion of the face and petechiae. When suspension is complete, face is pale and petechial haemorrhages are relatively uncommon due to the complete obstruction of the arterial system.

Option C- Vagus nerve compression leads to a sudden halt of the progression of the pure asphyxial process due to the superimposition of vagal inhibition. In these cases, the face remains pale.

Option D- Right internal carotid artery compression can cause transverse tears of carotid artery intima, known as the Amussat's sign. It is seen with antemortem hanging. These are usually seen in obese victims and in posteriorly placed knots.



Forensic Medicine NEET 2021

Question 1:

A guy was playing in a garden. Suddenly, he collapsed and was rushed to the hospital. His friend who played with him in the garden informed the doctors that he ate one of the fruits in the garden post which he seemed to have developed these symptoms. He also had irritability, restlessness, dry hot skin, and was unable to pass urine and stools. Identify the poison and its appropriate antidote.

a) Datura, Pralidoxime

b) Datura, Physostigmine

c) Yellow oleander, Digoxin

d) Yellow oleander, Physostigmine

Question 2:

Identify the lesion shown in the image below.

a) Laceration

b) Incised wound

c) Laceration looking incised wound

d) Incised looking laceration

Question 3:

A man was working in a field and he suddenly collapsed on a hot summer afternoon. On examination, there were no physical signs of dehydration. His serum electrolytes level was normal. The doctor found that the body temperature was 106 degrees Fahrenheit. Which of the following symptoms is least likely to be seen in this patient?

a) Hot skin

b) Hypotension

c) Sweating

d) Disorientation

Question 4:

A young teenage boy with a history of drug abuse had been instilled with a shot of intravenous cocaine at a party. He is rushed to the emergency room by his friends. Some of the symptoms were diaphoresis, headache, and acute coronary spasm. Which of the following symptoms noted would go against the likely diagnosis of cocaine poisoning?

a) Myocardial ischemia

b) Bradycardia

c) Hyperthermia

d) Agitation

Question 5:

The below phenomenon is due to

a) Case of torture falanga

b) Case of suspension/hanging more than 72 hours

c) Case of immersion in water for 36 hours

d) Case of colliquative liquefaction due to immersion in hot water

Question 6:

What is the active principle of the poison shown in the image?

a) Ricin

b) Bhilawanol

c) Abrin

d) Calotropin

Question 7:

Which of the following is true regarding the post mortem change shown in the image given below?

a) Seen within 24 hours of death

b) Due to sulphemoglobin accumulation and formation

c) Due to aseptic non bacterial cause

d) It is associated with a case of lightning

Question 8:

A guy was sitting on the balcony which was overlooking a park. He always used to sit nude on his balcony and derived sexual gratification by observing the people that look at him. This is best described as?

a) Voyeurism

b) Fetishism

c) Exhibitionism

d) Masochism

Question 9:

A woman gave birth to twins. The father who believed that they did not belong to him requested DNA testing. After testing, it was found that one child did not belong to him. This case would be best described as?

a) Superfetation

b) Superfecundation

c) Supposititious child

d) Atavism

Question 10:

While recording evidence in the court of law, the lawyer asked the witness, "Did A kill B in your presence?" The witness said, "Yes." This type of question would be permitted in which procedural route in the court of law?

a) Examination-in-chief

b) Direct examination

c) Cross examination

d) Re-direct examination

Answer Key

Question No.	Correct Option
1	b
2	d
3	c
4	b
5	c
6	b
7	b
8	c
9	b
10	c

Detailed Explanations

Solution to Question 1:

The clinical vignette in the question is suggestive of Datura poisoning for which the appropriate antidote is physostigmine, a reversible acetylcholine esterase antagonist. It is an effective treatment for severe anticholinergic poisoning.

Datura poisoning:

The common names include thorn apple, Jimson weed, and Devil's trumpet. The active ingredients include the anticholinergic alkaloids atropine, scopolamine, and hyoscyamine. The seeds or other parts of the plant can be ingested or smoked.

Toxic manifestations of Datura:

• Mnemonic: "hot as a hare, blind as a bat, dry as a bone, red as a beet, and mad as a hatter"

• Also, remember the D's of Datura

• Dry hot skin (Temp rises up to 42 C, a scarlet rash may also be appreciated).

• Dilatation of pupils along with loss of accommodation

• Diplopia

• Dryness of the mouth and throat, unquenchable thirst

• Dysphagia

• Dilatation of cutaneous blood vessels

• Dysarthria (Difficulty in talking)

• Drunken gait

• Delirium

• Drowsiness

• Death due to respiratory or cardiac failure

Treatment

• Gastric lavage or decontamination with activated charcoal or tannic acid. Gastric emptying is often delayed, and the small, plentiful seeds can become trapped among the gastrointestinal folds after ingestion; thus gastric decontamination can be an important therapy.

• Physostigmine in severe toxicity given 1-2 mg IM/IV after every half hour as required. It is a physiological antidote. Pilocarpine may also be given.

Medico-legal importance:

In India, datura is used as a stupefying poison prior to robbery, kidnapping, and rape. It is also known as a rail-road poison, as it is commonly encountered during a journey. Robbers disguised as saints may offer ‘prasad’ mixed with datura seeds and rob the passengers.

Yellow oleander:

All parts of the plant are poisonous. The active principles are thevatin, and thevatoxin, which are cardiac glycosides. The nectar of the flower yields poisonous honey. Symptoms include

burning sensation in the mouth, tingling of the tongue, dryness of throat, vomiting, diarrhea, headache, dilated pupils, irregular action of heart followed by drowsiness, coma, collapse, and death.

The given image shows the flowers of Cerebra thevatia.

Solution to Question 2:

The image depicts a wound with linear splitting of the tissue, sharp margins, and crushed hair follicles, which is suggestive of an incised-looking lacerated wound.

Incised-looking lacerated wounds are seen at sites where the skin overlies an area of the bone where there is scanty underlying subcutaneous tissue. A blunt force causes linear splitting of the tissue which results in the laceration resembling an incised wound. There will be extensive tissue crushing but the margins will be clear and delineable.

Common areas are Scalp, Forehead, Eyebrows, Zygoma, Lower jaw, Perineum, Shin, and Iliac crest.

Laceration (Option-A)

A laceration is a tear or a split in the skin, mucous membrane, muscle, or any internal organ produced by blunt force. The full thickness of the skin is penetrated.

Features of laceration:

• Margins are irregular with pieces of tissue attached in between called tissue tags or bridges. Tears that are seen at the ends, diverging away from the main laceration are called swallow tails.

• Bruising is seen around the margins.

• Deeper tissues are unevenly divided with tissue tags.

• Hair bulbs are crushed.

• Depth of the wound contains abundant foreign material such as dust particles, paint material of the vehicle involved, etc

• In skin lacerations, usually one of the margins overhangs the other. This helps in assessing the direction of the force.

• A laceration, on healing, produces a permanent scar.

Classification of a laceration:

• Split laceration (incised - like laceration)

• This occurs when the skin is crushed between two hard objects

• For example, during a fall, the scalp gets crushed between the skull from inside and hard ground outside. This results in a wound, which usually mimics an incised one unless examined closely.

• Stretch laceration

• As the name implies, it occurs due to the over-stretching of the skin, beyond its elasticity.

• This typically occurs in the setting of a fracture, where the fracture fragment may stretch and pierce the skin to project out

• Avulsed laceration

• An avulsion is a lesion where the skin separates or peels off by tearing from the underlying tissues. It may overhang the wound.

• This is common in road traffic accidents.

• Tears - Tearing of skin and subcutaneous tissue can occur from localized impact

• Cuts - Injury leads to a deep and wide injury leading to a cut



Incised wound (Option-B)

An incised wound is a clean-cut wound made with a sharp-edged instrument, through the skin and subcutaneous tissues, including blood vessels. It is more in length than depth. The edges are well-defined and usually everted, free from contusions and abrasions. It is generally spindle-shaped. The depth is more at the commencement and progressively becomes shallower,

known as tailing. Tailing determines the direction of the force applied when creating the wound.

The image below shows two spindle-shaped incisions.

Laceration-looking incised wounds (Option C)

In areas of the body where a loose layer of muscle fibres is attached to the skin, the edges of the incision may become inverted, adherent, and wrinkled leading to it simulating a lacerated wound. Such wounds are called lacerated-looking incised wounds. These are commonly seen in Scrotum and Axilla.

Solution to Question 3:

The given clinical vignette in a patient with no signs of dehydration and body temperature of 106oF is highly suggestive of heatstroke. Sweating is absent in heatstroke.

Heatstroke/ thermic fever/ heat hyperpyrexia is due to the failure of the heat regulation center of the hypothalamus. It is caused by failure of cutaneous circulation and sweating. The predisposing factors include high temperature, increased humidity, minor infections, old age, lack of acclimatization, alcoholism.

Heatstroke can be:

• Exertion-induced heat stroke is seen in people with prolonged exposure to or physical exertion in high temperatures, usually without adequate training.

• Sedentary heatstroke is a disease of elderly or infirm people whose cardiovascular systems are unable to adapt to the stress of a hot environment and release sufficient heat such that body temperature rises.

All the signs of shock are present. Patients usually present with a sudden collapse accompanied by loss of consciousness. Typical vital sign abnormalities include tachypnea, tachycardias, hypotension, and a widened pulse pressure. Other specific signs include rapid Kussmaul type of breathing, the absence of sweating with dry, hot skin, delirium, coma, convulsions,

rhabdomyolysis, multiorgan failure.

Although there is no single specific diagnostic test, clinical features of exposure to heat stress, CNS dysfunction, and core temperature >104.9-degree Fahrenheit (> 40.5 degrees Celsius) help establish the preliminary diagnosis. The body temperature must be measured rectally.

Treatment includes rapidly cooling the body with ice water till the rectal temperature falls below 1020F.

Autopsy findings include the presence of edema and reduction of Purkinje's cells in the cerebellum.

Solution to Question 4:

Cocaine intoxication presents with tachycardia and not bradycardia. Thus, the presence of bradycardia would go against the diagnosis of cocaine intoxication.

Cocaine is an alkaloid deliriant extracted from Erythroxylum coca.

Acute intoxication with cocaine leads to symptoms spanning 2 stages-

• Stage of excitement/euphoria - associated with hypervigilance, anxiety, hyperactivity along with hyper-adrenergic signs of mydriasis, tachycardia, elevated blood pressure, hyperthermia, and cardiac arrhythmias.

• Stage of depression - associated with loss of reflexes, muscle paralysis, circulatory and respiratory failure, and coma.

Chronic intoxication is characterized by toxic delirium and tactile hallucination of bugs (cocaine bugs) or vermin crawling under the skin (formication).

Common complications include cerebrovascular accidents and myocardial ischemia.

Treatment includes intravenous diazepam for controlling seizures. For ventricular arrhythmias, propranolol may be given intravenously. If the patient has psychosis, haloperidol may be given. Treatment of chronic cocaine abuse requires psychiatric support.

Cocaine withdrawal is a triphasic syndrome that presents after the sudden stoppage of chronic cocaine use. The phases are as follows -

• Crash phase (9 hours - 4 days): Anorexia, depression, excessive craving, hypersomnia, fatigue.

• Normal mood, anxiety, and anhedonia (4 - 7 days)

• Extinction phase (7 - 10 days): Absence of withdrawal symptoms but increased susceptibility to relapse.

Treatment for withdrawal includes bromocriptine and amantadine. They are useful in reducing cocaine craving.

Solution to Question 5:

The given image shows maceration of the skin of the soles suggestive of washerwoman's feet/ washerwoman's skin. It is seen in a case of immersion in water for 36 hours.

Washerwoman's skin occurs if the body of the person has been immersed in water for a few hours to 3 days. The first signs tend to be on areas with an appreciable keratin layer (such as the fingertips, palms, backs of the hands, and later the soles) where the surface becomes wrinkled, pale, and sodden. This is called washerwoman's skin. The first site affected is the fingerpads. It affects the hands earlier than the feet.

When a body falls into water in average temperate climates, the following is an approximate guide to timing:

The other signs of immersion are:

• Cutis anserina - goose-flesh or piloerection: This is usually found when the body is immersed in cold water.

• Post-mortem hypostasis - If the water is still, the postmortem hypostasis affects the heads and the limbs. This is because the chest and abdomen float higher because of the lungs and the abdominal gases.

• Foreign bodies in respiratory passages and stomach - Sand may be found deep in the respiratory passages and stomach, especially if the body has been rolled by the waves on a beach. This is not evidence of live aspiration.

Note: Signs of immersion just indicate that the body was immersed in water, they are not evidence of death from drowning.

Finding	Time since immersion
No wrinkling of finger pads	Less than a few hours
Wrinkled fingers, palms, and feet	Few hours to 3 days
Early decomposition (Head and abdomen)	4-10 days
The marbling of veins, peeling of the epidermis	2-4 weeks
Muscle loss with bone exposure	1-2 months

Falanga (Option-A) also known as Falaka or Bastinado, is a method of torture that involves repeatedly striking the soles of the feet with an object such as a truncheon, iron rod, cable, bamboo cane, wooden pole, or rifle butt.



Glove-stocking-like staining (Option-B) is used to describe the purple-colored postmortem staining visualized in the dependent parts of the upper and lower limbs of a hanging victim. It is absent in postmortem hanging, where the pattern of hypostasis does not correspond with the position of suspension of the body

Liquefaction or colliquative changes begins by 5 days after death and becomes prominent in 10 days. Soft tissues change into semisolid masses which are black in color and get separated from bones and fall off.

Solution to Question 6:

The given image shows the seeds of Semecarpus anacardium, and its active principles are bhilawanol and semicarpol. Common names are marking nut, bhilwan, bibva, bhela, and oriental cashew.

The image shows black, heart-shaped seeds with hard rinds that identify them as seeds of S. anacardium. Within the fruit is a thick fleshy pericarp that yields the active principle, which is a brown oily resinous fluid called bhilawanol.

The juice when applied locally to the skin produces irritation, painful blisters, itching, and eczematous eruption. The seeds are used to produce an artificial bruise. Symptoms such as pain, fever, and painful passage of brownish urine may be seen.

A large dose taken orally can produce blisters in the mouth and throat, with gastroenteritis. It also produces dyspnea, cyanosis, tachycardia, coma, and death. The fatal period is 12 to 24 hours.

Medicolegal importance is the use of the juice by malingerers to fabricate wounds. Instilled in the eyes it can cause conjunctivitis or ophthalmia due to its local irritant effect.

Other options:

Option A: Ricin is the active principle of Ricinus communis. It is a toxalbumin or phytotoxin that inhibits protein synthesis by affecting the ribosomal RNA polymerase. The seeds are flat to oval-shaped and have dark brown markings (shown below). It is a chemical warfare agent.



Option C: Abrin is the active principle of Abrus precatorius. The seeds are egg-shaped, scarlet in color, and have a black spot at one end (shown below). On ingestion, it causes gastroenteritis with bloody diarrhea, rectal bleeding, tachycardia, and circulatory collapse. Injecting the active substance can mimic viperine snakebite. Rarely, it may lead to tetanic convulsions.

The seeds of A. precatorius are decocted and mixed with datura, opium, onion, spirit, and water to form a paste. From this, small sharp needles called suis are made. They are used for hunting animals and even for homicide purposes.

Option D: Calotropin is one of the active principles of Calotropis. The toxic parts of the plant are the stem, branches, leaves, and milky white latex (Madar juice). When applied locally, it can mimic bruises, and form pustules and vesicles. In the eyes, it can lead to conjunctivitis. On ingestion, it causes abdominal pain, vomiting, mydriasis, and tetanic convulsions and can eventually lead to collapse and death.

It is shown in the image below.

Solution to Question 7:

The postmortem change shown in the given image is marbling. It is seen when blood in the vessels is haemolysed to sulfmeth-hemoglobin, which stains the vessel walls and the adjacent tissues, giving rise to a marbled appearance.

Marbling:

The marbling of the skin begins at about 24 hours and becomes prominent in 36–48 hours after death. It is a visible postmortem change that is characterized by linear branching patterns on skin that correspond to the course of superficial veins. Intestinal bacteria infest the venous system and there is subsequent gas production and hemolysis which stains the vessel walls. This becomes prominent in the summer and is distinctly appreciable in the superficial veins of the abdomen, shoulders, chest, and inguinal region.

Lichtenberg's figures or filigree burns are pathognomonic for lightning injury. It is a superficial, thin, tortuous, fern-like pattern of erythema in the skin. It indicates the path taken by the electrical discharge and tends to follow skin creases and the long axis of the body. They are also known as arborescent burns or feathering as they have a resemblance to the branches of a tree. Lichtenberg's figures occur due to the rupture of small vessels. They are usually seen over the shoulder or flank within an hour of injury and disappear in 1-2 days.

Medico-legal aspect: Its recognition may be lifesaving in the unaccompanied comatose patient and is important because even delayed resuscitation of lightning victims can be very successful.

The image below shows the filigree burns seen in a case of lightning.

Solution to Question 8:

The given scenario is suggestive of exhibitionism.

It involves behaviors over a period of 6 months or more, involving the exposure of one's genitals to an unsuspecting stranger. The individual obtains sexual gratification from this. This is a punishable offense (Section 294 IPC/ Section 296 BNS) with imprisonment up to 3 months or a fine.

The important and common sexual perversions are summed in the pearl attached below. The various punishments in the legal provisions include:

IPC section	BNS Section	Section it deals with	Punishment
290	292	Punishment for public nuisance (Frotteurism)	Fine Up to Rs.1000
294	296	Obscene acts and songs (Exhibitionism)	Up to 3 months of imprisonment or fine (upto Rs. 1000) or both
297	301	Trespassing on burial places/ wounding the feeling of any person /indignity to any human corpse(Necrophilia, Necrophagia)	Up to 1 year of imprisonment or fine or both

Solution to Question 9:

The given clinical scenario where one twin is found to not belong to the father of the other twin is suggestive of heteroparental superfecundation.

Superfecundation is the fertilization of 2 different ova in the same menstrual cycle (as suggested by the same gestational age) by separate acts of coitus. It can occur even after artificial reproductive therapy. Hence, after embryo transfer, intercourse must be avoided. Heteropaternal superfecundation is used to refer to instances of two different males fathering fraternal twins. This leads to the possibility of twins also being half-siblings. This may give rise to the doubt of adultery and infidelity.

Option A: Superfetation is the fertilization of 2 different ova released in different menstrual cycles (i.e ovulation and fertilization during pregnancy). This can occur within 12 weeks of pregnancy after which the decidual space is obliterated and a detailed evaluation will reveal disparities in gestational ages.

Option C: Supposititious child is also known as a fictitious child. In such cases, a woman pretends to be pregnant and generally substitutes a male child for a female child born of her, or after an abortion. This is done for the purpose of claiming property.

Option D: Atavism is the reappearance of a characteristic in an individual after several generations of absence. For example, the child may not resemble his parents but resemble his ancestors. This is attributed to the recombination of genes.

Solution to Question 10:

In the given scenario, the kind of question asked by the lawyer to the witness is an example of a yes/no question or leading question. Leading questions are allowed during cross-examination according to Section 143 IEA (Indian Evidence Act).

Leading questions (Sec. 141 IEA) are defined as questions that suggest the answer that the person asking the question wishes or expects to receive. It is favourable for the case of the summoning counsel or in alignment with their claim. They are not allowed during the direct examination(Examination-in-Chief) conducted by the summoning counsel (Option A, B). The only exception to this is in the event that the Judge is satisfied that the witness is hostile, leading questions may be allowed during direct examination.

In cross-examination, the witness is examined by the defence lawyer or adverse party with the objective to elicit favourable facts to their case, to test the accuracy of the witness statements, unearthing new or old facts, and also, to discredit the witness. While leading questions are allowed in cross-examination, the Judge can disallow questions that are indecent or scandalous (Sec. 151 IEA) or questions asked with an intention to insult, annoy or offend the witness in any form (Sec. 152 IEA). There is no time limit to the duration of cross-examination.

The sequence of examination in the recording of evidence is as follows -

Sequence of Examination	Objective
Examination-in-Chief/Direct examination by the party tha t summoned the witness	To elicit the facts of the case from the witness. Leading qu estions are not allowed.

Sequence of Examination	Objective
Cross-examination by the defence lawyer	To elicit facts favourable to their case and test the reliability of the witness. Leading questions are allowed.
Re-examination by the party that summoned the witness	To resolve doubts that formed during cross-examination and re-emphasize the party's perspective

Forensic Medicine INI-CET 2022

Question 1:

Sagar was being tried in a court for the murder of Mohan in the park. Shikhar appeared in court and said he saw Sagar carrying a knife in his hand while in the park. What is this evidence known as?

a) Direct

b) Indirect

c) Hostile

d) Hearsay

Question 2:

Fingerprint first develops in how many weeks of intrauterine life?

a) 12-16

b) 20-24

c) 28-32

d) 32-36

Question 3:

Identify the given plant.

a) Datura alba

b) Argemone mexicana

c) Nerium odorum

d) Papaver somniferum

Question 4:

A male patient was presented with a history of consumption of an unknown substance. He had increased sweating, lacrimation, diarrhea, muscle weakness, and bradycardia. What's the management?

a) Atropine

b) Calcium gluconate

c) Pralidoxime

d) Obidoxime

Question 5:

Which of the following is not a feature of cocaine intoxication?

a) Tingling and numbness of hands and feet

b) Constricted pupils and bradycardia

c) Hypertension

d) Hyperthermia

Question 6:

Which of the following are seen in patients with chronic arsenic poisoning?

a) 1,2,3 are correct

b) 2,3 are correct

c) 1,4 are correct

d) Only 1 is correct

Question 7:

Post-mortem examination of a patient reveals the following. What is this finding known as?

a) Hesitation cuts

b) Grazed abrasion

c) Laceration

d) Patterned bruise

Question 8:

Poisoning with which of the following agents would cause hypertension?

a) Organophosphorus

b) Paracetamol

c) Arsenic

d) Cocaine

Question 9:

Treatment of choice for acute arsenic poisoning is :

a) Ipecac

b) Penicillamine

c) Activated charcoal

d) Dimercaprol

Question 10:

To detect vaginal cells on a person who is accused of rape, which test will you use?

a) Florence

b) Barberio

c) Lugol's

d) Precipitin

Question 11:

Which of these is not a cardiac poison?

a) Aconite

b) Cerbera thevetia

c) Nicotiana tabacum

d) Atropa belladonna

Question 12:

The image given below is a type of _____.

a) Contusion

b) Abrasion

c) Laceration

d) Stab wound

Question 13:

Match the following:

a) A-1, B-3, C-2, D-4

b) A-3, B-4, C-1, D-2

c) A-3, B-2, C-1, D-4

d) A-2, B-3, C-4, D-1

Question 14:

Match the following plant poisons with their active principles.

a) 1-b,2-c,3-d,4-a.

b) 1-a,2-c,3-b,4-d.

c) 1-a,2-b,3-c,4-d.

d) 1-c,2-b,3-a,4-d.

Question 15:

The dead body of a child was recovered from a river with froth around his mouth, which reappeared even after wiping it off. Which postmortem finding is not seen in this case?



a) Voluminous and ballooned out lungs

b) High concentration of arsenic from shaft of hair

c) Paltauf's hemorrhage in sub pleura

d) Cadaveric spasm with mud in closed fist

Question 16:

Spanish windlass is a type of which strangulation method?

a) Bansdola

b) Garrotting

c) Mugging

d) Throttling

Question 17:

A person of eonism derives pleasure from _____.

a) Wearing clothes of opposite sex

b) Kissing and licking of anus by a sexual partner

c) Underclothing of female

d) Seeing a female undressing

Question 18:

A lady underwent cholecystectomy and now she presents with abdominal pain, one month later. An erect abdomen x-ray was taken, and the image is given below. The following offense comes under _____.

a) Novus actus interveniens

b) Res ipsa loquitur

c) Res judicata

d) Respondeat superior

Question 19:

Tattooing around the entry wound of a bullet is due to?

a) Unburnt particles

b) Smoke

c) Flame

d) Muzzle impression

Question 20:

Pick the mechanism of cold water drowning.

a) Asphyxia

b) Laryngeal spasm

c) Vagal inhibition

d) Cardiac arrest

Question 21:

Identify the pattern of abrasion shown in the image below.

a) Ligature mark

b) Graze abrasion

c) Pressure abrasion

d) Imprint abrasion

Question 22:

Gas shadow is present in the great vessels of a fetus in the case of intrauterine demise. What is this sign called?

a) Roberts sign

b) Spalding sign

c) Ball sign

d) Halo sign

Question 23:

A man was brought to the emergency department with a history of assault by a mob using heavy blunt objects and sharp objects. On examination, crushing of the hair bulb was seen. Identify the type of wound.

a) Clean cut incised wound

b) Lacerated wound

c) Chop wound

d) Avulsion injury

Question 24:

Identify the postmortem change given in the image.

a) Marbling

b) Tattooing

c) Lividity

d) Filigree burns

Question 25:

A male patient working in the matchbox industry for the past 20 years reports to the dentist with multiple sinuses and fistulae in and around his jaw with foul-smelling discharge. Which of the following chemicals are responsible for this?

a) Bromine

b) Antimony

c) Fluorine

d) Phosphorus

Question 26:

Which of the following rules of law are associated with insanity?

a) 1, 2, 3 and 4

b) 2, 3, 4 only

c) 1, 2 only

d) 3, 4 only

Question 27:

The steps in the management of snake bite include:

a) 1, 2, 3, 4, and 5

b) 1, 2, and 4

c) 4 only

d) 4 and 5

Question 28:

Hooch tragedy is related to:

a) Phosphorus

b) Mercury

c) Ethanol

d) Methanol

Answer Key

Question No.	Correct Option
1	b
2	a
3	b
4	a
5	b
6	a
7	a
8	d
9	d
10	c
11	d
12	a
13	c

14	b
15	b
16	b
17	a
18	b
19	a
20	c
21	c
22	a
23	b
24	a
25	d
26	c
27	d
28	d

Detailed Explanations

Solution to Question 1:

This evidence is known as indirect evidence. Indirect evidence is also known as circumstantial evidence. It is not the direct testimony of an eyewitness. It has an effect on other facts and subsidiary facts which are depended upon as consistent.

Other options:

Option A: Direct evidence is the evidence of a fact that is actually in issue. Eg: The witness saw A shooting B, prescription, consent form

Option C: Hostile evidence is given by someone who is supposed to have some interest or motive for hiding part of the truth. He might give completely false evidence too.

Option D: Hearsay evidence is given by someone about something that he did not personally witness. Sometimes it might be obtained from a third party. It is presented in the court to claim that the facts contained in the statement are true.

Solution to Question 2:

Fingerprint first develops in 12-16 weeks of intrauterine life. It is complete by 24 weeks. In identical twins also the pattern is different.

The image given below shows the different patterns of fingerprints:



Changes might occur in fingerprints in certain conditions, some of them are given below:

Fingerprint changes	Condition
Incomplete atrophy of ridges	Dermatitis
Loss of pattern with ridge atr ophy	Celiac diseas e
Altered ridges	Scleroderma
Permanent loss of fingerprint s	Radiation
Distance between ridges cha nged but pattern is retained	Acromegaly

Solution to Question 3:

The given plant is Argemone mexicana.

The flowers are yellow in color. All parts of the plant are poisonous. Oil extracted from argemone seeds is commonly used as an adulterant in mustard
oil. Sanguinarine and dihydrosanguinarine are the toxic alkaloids present in the oil and it causes epidemic dropsy.

Epidemic dropsy is characterized by vomiting, diarrhea, edema of the feet, and generalized anasarca. Some patients may develop hypersecretory glaucoma. It can lead to death due to cardiac failure.

Other options:

Option C: Nerium odorum - Flowers are usually white, pink, dark red, or pale yellow in colour. All parts of the plant including nectar, are poisonous. It contains several cardiac glycosides such as neriodorin, neriodorein, and karabin. The action of neriodorin mimics digitalis causing death

due to cardiac failure. It can resist heat and hence can be found even in burnt dead bodies.

Option D: Papaver somniferum - The flowers are white in color. Opium is the dry juice of this plant. Morphine is isolated from opium. The clinical features include euphoric feeling in the stage of excitement followed by headache, giddiness, and a tendency to sleep in the stage of stupor. The stage of narcosis is characterized by a patient in a deep coma, who cannot be aroused.

Solution to Question 4:

The clinical scenario is suggestive of organophosphate consumption and atropine is used for its management. Atropine is a muscarinic receptor antagonist and it competitively blocks the acetylcholine at muscarinic receptors.

Organophosphates are powerful acetylcholinesterase inhibitors. It is absorbed by inhalation or through the skin. Mucous membranes and the gastrointestinal tract are also the sites for absorption.

The predominant initial signs are of muscarinic excess which includes: (DUMBBELSS)

• Diarrhoea

• Urination

• Miosis

• Bronchospasm

• Bradycardia

• Emesis

• Lacrimation

• Salivation

• Sweating

Nicotinic manifestations include twitching of eyelids, tongue and facial muscles along with tachycardia which then leads to paralysis. CNS stimulation results in headache, tremors, ataxia, coma and death.

Atropine in larger doses can lead to atropine toxicity. The features are depicted in the following image:



Other options:

Option B: Calcium gluconate is used as the antidote for oxalic acid poisoning.

Option C: Pralidoxime is also used in organophosphate poisoning which is used to counter the nicotinic effects of organophosphates.

Option D: Obidoxime is used in organophosphate poisoning if pralidoxime is not available.

Note: In OP poisoning, the frequency of occurrence is Muscarinic --> CNS --> Nicotinic signs and symptoms. The definitive treatment, however, is atropine.

Solution to Question 5:

Constricted pupils and bradycardia are not a feature of cocaine intoxication. Dilated pupils and rapid pulse are seen in cocaine intoxication.

Cocaine is obtained from the leaves of Erythroxylum coca. Cocaine can be administered by snorting, intravenous route, or smoking.

The manifestations of cocaine intoxication occur in two stages- the stage of excitement and the stage of depression.

In the stage of excitement, the features seen are dry mouth, dysphagia, tachycardia, tachypnea, hypertension, dilated pupils, and increased temperature. There may be a feeling of well-being, and the patient may be excited, restless, and talkative. This progresses into a calm, dull state. Neurological manifestations include exaggerated reflexes, tremors or convulsions, hallucinations, tingling, and numbness in the hands and feet.

The stage of depression is characterized by feeble respiration, excessive sweating, collapse, convulsions, and death.

The fatal dose of cocaine is 1g when taken orally. The fatal period ranges from a few minutes to hours.

Chronic cocaine users may have the feeling of insects crawling on their skin or the presence of grains of sand under the skin. This is known as cocaine bugs or Magnan's symptom. Ulceration of the nasal septum may be seen in chronic abusers who sniff cocaine.

Treatment includes gastric lavage with warm water containing potassium permanganate, charcoal, or tannic acid if taken orally. When it is through the nose or throat, a wash with water is done. In case of intravenous intoxication, a ligature is applied above the site of the injection.

The antidote for cocaine intoxication is inhalational amyl nitrite. Other measures include control of convulsions, airway and circulatory stabilization, administration of iv thiamine, and naloxone.

Solution to Question 6:

Among the given options, raindrop pigmentation, peripheral neuritis, and mees line are seen in patients with chronic arsenic poisoning. Luminous vomit is a feature of yellow phosphorous poisoning.

Arsenic is an odorless, tasteless, metallic poison. Gastrointestinal features of acute arsenic poisoning include nausea and vomiting, colicky abdominal pain, and profuse diarrhea with rice water stools.

In chronic arsenic poisoning, skin symptoms include erythematous lesions, flushing, and can also lead to hyperkeratotic skin and desquamation. Pigmentation is patchy and resembles raindrops on a dusty road. In the distal parts of the body, hyperkeratosis is present. Mees lines are

transverse white striae of the fingernails, called striate leukonychia. Renal manifestations include chronic nephritis and urine contains albumin, casts, etc.

Arsenic can be detected in the bones for prolonged periods. Hence it can be found even in skeletonized remains of the body. Dimercaprol is the treatment of choice for acute arsenic poisoning.

Solution to Question 7:

The finding is known as hesitation cuts. It is a feature of suicide. They are small multiple cuts which involve only the superficial skin. They are seen at the beginning of the incised wound. The throat, wrist and front of the chest are the most common sites. Unintentional cuts are seen on the fingers in which the razor blade are held.

Other options:

Option B: Grazed abrasion occurs when there is movement between skin and some rough surface. The epidermis is destroyed and shows uneven longitudinal parallel lines in the direction of the force. Cleaning and dressing is the treatment.

Option C: In laceration tearing of the skin, mucous membrane, muscles or internal organs occurs due to blunt force application. The tissues are crushed and stretched beyond their elasticity limit.



Option D: In Patterned bruise, it mirrors the pattern of the object used. The tissue is crushed if heavier objects are used and will produce a solid contusion pattern.

Solution to Question 8:

Poisoning with cocaine cause hypertension. This is due to the inhibition of norepinephrine at the adrenergic terminals leading to increased sympathetic stimulation. This effect occurs at all concentrations of cocaine and causes hypertension and ventricular ectopics.

Cocaine is obtained from the plant Erythroxylum coca. The symptoms of cocaine poisoning are divided into two stages:

• Stage of excitement: This stage is characterized by features like dysphagia, dryness in the mouth, tachycardia, and dilated pupils. The reflexes are exaggerated and are associated with tingling and

numbness in the hands and feet.

• Stage of depression: This stage presents with feeble respiration, profuse sweating, convulsions, and death. In most cases, the cause of death is cardiopulmonary failure or vascular collapse.

Chronic users develop a feeling of insects crawling through the skin or the presence of grains of sand under the skin. This is pathognomic for cocaine poisoning and is known as cocaine bugs or Magnan's symptoms. Ulceration and sometimes perforation of the nasal septum are seen in chronic abusers due to the sniffing habit.

The image of the nasal septum of a chronic abuser is shown below:



The image of the plant is shown below:

Other options:

Option A: Organophosphorous poisoning produces autonomic instability. Both tachycardia and bradycardia can be seen, however, usually it is the parasympathetic effects which predominate.

Option B: Paracetamol poisoning causes hypotension, tachycardia, diaphoresis, etc.

Option C: Arsenic poisoning causes vasodilation, ventricular tachycardia, ventricular fibrillation, etc.

Solution to Question 9:

Dimercaprol is the treatment of choice for acute arsenic poisoning.

Arsenic is an odorless, tasteless, metallic poison. Gastrointestinal features of acute arsenic poisoning include nausea and vomiting, colicky abdominal pain, and profuse diarrhea with rice water stools.

In chronic arsenic poisoning, skin symptoms include erythematous lesions, and flushing and can also lead to hyperkeratotic skin and desquamation. Pigmentation is patchy and resembles raindrops on a dusty road. In the distal parts of the body, hyperkeratosis is present. Mees lines are transverse white striae of the fingernails, called striate leukonychia. Renal manifestations include chronic nephritis and urine contains albumin, casts, etc.

Arsenic can be detected in the bones for prolonged periods. Hence it can be found even in skeletonized remains of the body.

Other options:

Option A: Ipecac is an emetic used in ergot and hydrocarbon poisoning.

Option B: Penicillamine is used in lead poisoning as a chelating agent.

Option C: Activated charcoal acts as an adsorbing agent in many poisonings like aspirin, phenobarbital poisoning, etc.

Solution to Question 10:

Lugol's iodine test will be used to detect the vaginal cells present on the penis of a rape accused. It shows brown color on filter paper due to the presence of glycogen in epithelial cells of the vagina.

Other options:

Option A: Florence test is used to detect choline in that originates from seminal vesicles. It produces dark brown, rhombic, or needle-shaped crystals of choline iodide if semen is present.

Option B: Barberio test is used to detect spermatic fluid. It produces yellow rhombic crystals of spermine picrate when picric acid is added to spermatic fluid.

Option D: Precipitin test is used for species identification of semen stain. A positive test will confirm human origin.

Solution to Question 11:

Atropa belladonna is a deliriant poison that acts by inhibiting the muscarinic effects of acetylcholine. All the parts of the plants are toxic. The active principle is 1-hyoscyamine.

Clinical manifestations of A. belladonna poisoning:

• Fever, flushing, rash

• Neurological symptoms - delirium, slurred speech, headache, choreoathetosis, myoclonus, visual and auditory hallucinations, convulsions, and coma

• Eye symptoms - photophobia, blurred vision, dilated pupils poorly reactive to light

• Cardiovascular symptoms - arrhythmia, tachycardia, hypertension

• Gastrointestinal symptoms - severely dry mouth and throat, decreased bowel sounds, constipation, ileus

• Renal symptoms - urinary retention

Anticholinergic symptoms include tachycardia, hypertension, flushing, dry mouth, mydriasis, and urinary retention.

Other options:

Option A: Aconite is a cardiac poison derived mainly from the root and root tubers of the plant Aconitum napellus (shown below). Symptoms of intoxication are due to action on the voltage-gated sodium channels on the cell membrane of nerves and skeletal and smooth muscles. Symptoms include intestinal (nausea, vomiting, perioral tingling and numbness, abdominal pain, and diarrhea), cardiovascular (chest pain, palpitations, hypotension, bradycardia), and neurological (vertigo and restlessness) manifestations.

Option B: Thevetin is a cardiac poison derived from the plant Cerbera thevetia (shown below). Symptoms of intoxication are characterized by vomiting, diarrhea, headache, dizziness, mydriasis, and loss of muscular power. Cardiovascular signs include a weak and irregular pulse rate with hypotension. Possible complications are heart block and cardiorespiratory failure.

Option C: Nicotine is a plant alkaloid derived from the Nicotiana tabacum plant, shown in the image below. It is a cardiac poison.

All components of this plant are poisonous except its ripe seeds. It is a cardiac poison that initially stimulates the autonomic ganglia to produce symptoms of acute intoxication (tachycardia, hypotension, tachypnea, nausea, vomiting, abdominal pain, and headache). These symptoms later subside to give way to respiratory depression and collapse.

Solution to Question 12:

The type of injury shown in the image is a contusion.

A contusion is characterized by the infiltration of extravasated blood into the subcutaneous and/or subepithelial tissues. It results from the rupture of small blood vessels due to the application of blunt force.

• A pure bruise lies beneath an intact epidermis.

• Bruises are caused by damage to veins, venules, and small arteries.

• The usual bruise from a blunt impact is situated in the subcutaneous tissues, often in the fat layer. (Because of the greater volume of soft subcutaneous tissue in fat people, they tend to bruise more easily than thin ones).

• Delayed bruising - It is common knowledge among lay persons, as well as doctors, that a bruise may ‘come out’ - that is, become more prominent with the passage of hours or days. This is partly caused by continued bleeding from the ruptured vessels, but mainly by the percolation of free blood from its origin deeper in the tissues upwards towards the epidermis.

Other options:

Option B: Abrasion, also known as gravel rash, is the destruction of superficial layers of the epidermis. It is caused by friction against a rough surface or by compression. It is of four types:

• Scratches

• Grazes

• Pressure abrasions

• Impact abrasions

Option C: Laceration refers to a tear or split of the skin, mucous membranes, muscle, or internal organs. It is due to the application of blunt force to a broad area of the body. They are also called tears or ruptures. The laceration may indicate the cause of the injury and the shape of the blunt weapon. The age of the injury can also be determined.

Option D: A stab wound is formed when force is delivered along the long axis of a narrow or pointed object. The object can be a knife, sword, chisel, scissors, or needle. It is deeper than its length and width on the surface of the skin. The edges are clean-cut and inverted.

Solution to Question 13:

The correct option is Option C: A-3, B-2, C-1, D-4.

The time of emergence of secondary dentition or permanent teeth:

The time of emergence of primary dentition or temporary teeth:

The deciduous teeth all emerge into the mouth by 2.5-3 years.

Note: 6-13 years is the age of mixed dentition. The first molar emerges at 6-7 years. The second molar emerges at 12 years, and the third molar emerges beyond 18 years of age.

Tooth	Emergence
First molar	6-7 years
Central incisors	6-8 years
Lateral incisors	7-9 years
First premolar	9-11 years
Second premolar	10-12 years
Canine	11-12 years
Second molar	12-14 years
Third molar	17-25 years

Tooth	Emergence
Mandibular central incisor	6-8 months
Maxillary central incisor	7-9 months
Maxillary lateral incisor	7-9 months
Mandibular lateral incisor	10-12 months
First molar	12-14 months
Canine	17-18 months
Second molar	20-30 months

Solution to Question 14:

The correct match is 1-a,2-c,3-b,4-d.

The active ingredients of Datura include the anticholinergic alkaloids atropine, scopolamine, and hyoscyamine. The seeds or other parts of the plant can be ingested or smoked.

Manifestations of Datura toxicity:

• Dry hot skin (Temperature rises up to 42 C, a scarlet rash may also be appreciated).

• Dilatation of pupils along with loss of accommodation

• Diplopia

• Dryness of the mouth and throat, unquenchable thirst

• Dysphagia

• Dilatation of cutaneous blood vessels

• Dysarthria (Difficulty in talking)

• Drunken gait

• Delirium

• Drowsiness

• Death due to respiratory or cardiac failure

Morphine, codeine, and thebaine are natural opioids derived from poppy Papaver somniferum.

Abrin is a toxalbumin of the plant Abrus precatorius, and it acts by inhibiting protein synthesis.

Toxic principles of Abrus precatorius:

• Abrin (a toxalbumen)

• Abrine (an amino acid)

• Abralin (a glucoside)

Abrin acts by inhibiting protein synthesis, thereby, resulting in cell death. The clinical features after oral ingestion of these seeds include abdominal pain, nausea, vomiting, bloody diarrhea, and miosis. Faintness, dyspnea, and convulsions can also occur.

The poison obtained from Strychnos nux-vomica seeds is strychnine. Strychnine acts at the ventral horn by blocking the postganglionic glycine receptor sites in the spinal cord and brainstem. Glycine is an inhibitory neurotransmitter that acts at the postsynaptic membranes of ventral horn motor neurons. Hence, the blockade of these receptors results in unopposed muscle contraction, resulting in the classical clinical features:

• Tonic-clonic seizures

• Opisthotonus (body arched in hyperextension)

• Risus sardonicus

Solution to Question 15:

The above case scenario with frothing is suggestive of antemortem drowning. High concentration of arsenic in the shaft of hair is seen in patients with chronic arsenic poisoning.

Given below are the features of antemortem drowning.

• Froth - The presence of copious amounts of fine, shaving-lather-like froth at the mouth and the nostrils is the most characteristic finding of antemortem drowning. It is caused by the vigorous agitation of the tracheal and bronchial mucous secretions, edema fluid, and alveolar surfactant during respiratory efforts.

• Cutis anserina - The contraction of the erector pilorum muscles causes puckering and a grainy appearance of the skin. This resembles goose skin; hence, it is called cutis anserina.

• Washerwoman's hand - Maceration of the skin on the palms and soles gives a wrinkled, sodden, and bleached appearance due to submersion.

• Cadaveric spasm - Gravel, weed, soil, grass, etc. is firmly held in the clenched hands due to cadaveric spasm. This is a vital sign of antemortem drowning. The material also indicates the place of submersion.

• Rigor mortis - This appears early in cases of a violent antemortem struggle.

• Face - It is bloated and discolored, with cyanosis.

• Eyes - Conjunctiva are suffused and pupils dilated.

• Tongue - It may be swollen and protruded.

• Antemortem injuries - These may be sustained by striking objects, such as rocks, during the fall into the water body. Examination for such injuries must be performed only after the body is dry.

Internal findings:

• Lungs - They appear voluminous and distended show ballooning. The cut section will exude copious amounts of frothy, bloodstained fluid.

• Larynx, trachea, and bronchioles - The presence of sand, silt, and aquatic vegetation in the trachea and lower respiratory tract are characteristic findings of antemortem drowning.

• Heart and blood vessels- Gettler test can be done to estimate the chloride content in the right and left sides of the heart to differentiate between freshwater and saltwater drowning.

• Diatom test- Diatoms are microscopic algae. Their presence in the organs, particularly in the bone marrow, is suggestive of antemortem drowning.

• Stomach and small intestines- These organs may contain water in some cases. Water present in the small intestine is positive evidence of death by drowning.

• Middle ear - The presence of water in the middle ear is highly suggestive of antemortem drowning.

• Brain - It is congested, and edematous, with loss of gray-white junction.

Solution to Question 16:

Spanish windlass is an improvised version of garrotting, and it involves tightening an iron collar (Spanish windlass) around the neck and forcing a spike against the neck into the spinal cord.

This was a form of judicial execution once employed in Spain. The rod-like object used to tighten the ligature is called a Spanish windlass; this may be a ruler, stick, or screwdriver.

Garrotting involves attacking the victim from behind using a ligature. The ligature is thrown over the victim's neck and then tightened by twisting with a lever.

The image given below shows garrotting.

Option A: Squeezing to death using sticks is called bansdola. The neck or the chest is squeezed between two sticks, one placed in front, and one behind. The neck may also be squeezed with a

single stick in front if the victim is lying down.

Option C: Squeezing the neck with the arm and forearm is called mugging. The image below shows mugging.

Option D: Throttling or manual strangulation occurs due to compression of the neck by humans leading to asphyxia. Thumb impression and finger marks may be seen over the victim's neck.

Solution to Question 17:

A person of eonism (transvestism) derives pleasure from wearing clothes of the opposite sex. It is usually found in males. His dress, manner, occupational interests, and associations are all designed to increase his feeling of being a woman.

Other options:

Sexual pleasure from

• Kissing and licking of anus by a sexual partner - Anilingus

• Sexual interest in nonliving objects that are used to generate sexual excitement. -Fetichism

• Seeing a female undressing - Voyeurism or scoptophilia

Solution to Question 18:

The x-ray shows a pair of surgical scissors left behind in the abdomen after surgery. The offense comes under res ipsa loquitur. It is obvious evidence of the negligent act committed.

"Res ipsa loquitur" means "the thing or fact speaks for itself." The negligent act is so obvious that the patient need not prove the negligence. He has to merely state what according to him was the negligent act.

This doctrine applies to both civil and criminal negligence. All of the following conditions should be fulfilled for this doctrine:

• In the absence of negligence, there would not have been any injury usually

• The doctor had complete control over the injury-producing instrument or treatment.

• The patient was not a part of contributory negligence.

Other options:

Option A: Novus actus interveniens: A person is responsible not only for his actions but also for the logical consequences of those actions. This principle applies to cases of assault and accidental injury.

Option C: Res judicata (Section 337, BNSS) -If a question of negligence against a doctor has already been decided by a court in a dispute between the doctor and his patient, the patient will not be allowed to contest the same in another legal proceeding, based on the same set of facts. Multiple appeals cannot be made.

Option D: The principle of respondeat superior is used to explain vicarious liability, which explains that the employer is responsible for his employees' negligence if such an act occurs in the course of the employment and within its scope.

Solution to Question 19:

Tattooing around the entry wound of a bullet is due to the deposition of unburnt powder particles. These are difficult to remove. They are also called stippling or peppering. This is seen up to 50 cm of firing

When the distance between the firing and the victim is between 30-50 cm (in the case of pistols and revolvers), it is a near-shot wound. The victim is within the range of powder deposition, so there is some amount of tattooing present. But it is outside the range of flame, so there will not be any burning or singeing. There will not be any blackening, as it is outside the range of smoke as well.

The injuries related to firearms are:

• Burn injuries - occur due to flame released during the firing. It is seen in contact shot wounds or distances up to 5-8 cm. Singeing of hair is caused due to burns

• Blackening - occurs due to the deposit of smoke. It can be removed with a wet cloth. This can be seen up to around 15 cm of firing

• Tattooing - occurs due to the deposition of unburnt powder particles. These are difficult to remove. They are also called stippling or peppering. This is seen up to 50 cm of firing

• Greasing and collaring - occur due to the deposition of the lubricant of the bullet in the tissues, during its penetration. It can be found in contact, near-shot, or distant shot wounds.

In a firm contact shot, the imprint of the muzzle of the gun is seen on the skin around the wound. The victim is within the range of flame, hence singeing of the hair will be present. Tattooing begins only when the muzzle-to-target distance is >1cm, like in a loose contact shot.

When the distance between the victim and the firing is 5-8cm, it is called a close shot. The victim is within the range of the flame, hence, features related to flame, smoke, and powder deposition will be seen. Singeing, burning, blackening and tattooing will be present.

When the distance between the victim and the firing is >50 cm, it is a distant shot. The victim lies outside the range of flame and powder deposition. Hence, features related to flame, smoke, and powder deposition are absent. Burning, blackening, and tattooing are absent.

Solution to Question 20:

In cold water drowning or immersion syndrome or hydrocution, death occurs due to vagal inhibition, leading to cardiac arrest.

Hydrocution/ immersion syndrome refers to sudden death, following immersion in cold water. Death results from vagal inhibition of the heart, i.e. cardiac arrest occurring due to increased stimulation of the vagus nerve.

Vagus stimulation occurs due to the following mechanisms:

• Sudden entry of water into the nasopharynx or larynx

• Falling or diving into the water in a manner, so that it suddenly strikes the abdomen, especially the epigastric region

• Sudden in-rush of cold water into the ears

Types of drowning:

• Wet drowning/ typical drowning

• Atypical drowning/dry drowning

• Immersion syndrome/ hydrocution

• Post-immersion syndrome/ secondary drowning/ near-drowning

• Submersion of the unconscious/ shallow-water drowning

Solution to Question 21:

The given image shows an abrasion due to ligature mark, which is a pressure abrasion.

Abrasions are superficial wounds produced due to injury to the epidermis. They are usually caused by friction against a rough surface or by compression.

Types of abrasions:

• Linear abrasions (scratches): Caused by relative movement of a pointed object on the skin

• Examples: nail, thorn, the point of sword or knife

• Grazed abrasions (sliding /scrapping/grinding abrasion): Caused by relative movement of a rough surface on a wide area of skin

• brush burns- tangential rubbing over the surface, usually the ground

• friction burns are extensive excoriated areas without bleeding

• Examples: fall, a victim being dragged on the ground, road traffic accidents

• Imprint abrasions (patterned abrasions): Caused by the impact of a rigid hard rough surface over a localized area of the skin without any movement; usually show the pattern of the object causing them

• Examples: tire, grills, bicycle chain, bite marks

• Pressure abrasions (pressure abrasion): Caused by crushing of the superficial layers of the epidermis

• Examples: ligature marks around the neck in hanging

Solution to Question 22:

Gas shadow in the great vessels of a fetus in intrauterine death is called the Roberts sign. It is the earliest sign and is seen within 12 hours of death. The gas shadow may be present within the chambers of the heart or in large vessels of the fetus.

Signs of intrauterine death:

• Earliest sign - reddening of the skin with peeling in 12 hours

• Robert's sign - radiographic finding showing gas in the great vessels like the aorta in 12 hours

• Large blebs containing a red serous or serosanguineous fluid appear at around 24 hours

• Serous cavities may contain a turbid reddish fluid after 48 hours

• Spalding's sign (Option B) radiographic finding is seen as
a loss of alignment and over-riding of bones of the cranial vault. This occurs due to shrinkage of the brain after death. It is seen 7 days after fetal death.

• A sweetish and disagreeable odor

• Sodden and coppery red/purplish skin (never greenish as with putrefaction) with blisters, and cuticle peeling. The skin detaches easily and leaves moist and greasy areas

• Distended abdomen and thick, red, and soft umbilical cord

• Abnormally mobile or flexible joints

• Soft, edematous, and loss of morphology of the organs or viscera (lungs and uterus remain unaffected for long period)

• The body is soft and pliable and flattens out when placed on an even surface

Option C: Ball sign is hyperflexion of the spine with crowding of ribs seen in intrauterine death.

Option D: Halo sign is a radiological sign of intrauterine fetal death showing an increase in soft tissue width between subcutaneous fat and the cranial vault.

Solution to Question 23:

The type of wound in which the crushing of the hair bulb is seen is a lacerated wound.

Cut lacerations may be produced by blunt weapons or heavy relatively sharp-edged weapons such as axe, hatchets, and choppers. Scalp lacerations can be found in association with bruising and abrasions. Fascial strands, hair bulbs, nerves, and vessels, that run in the depth of the wound, are irregularly torn and crushed.

Margins are irregular, ragged, and uneven. The ends are pointed or blunt. The edges of lacerations, especially over the skull are undermined. The depth of the wound varies according to the thickness of the soft parts at the site of the injury and the degree of force applied.

Medicolegal importance of laceration:

• The type of laceration indicates the cause of the injury and the shape of the weapon.

• Foreign bodies found in the wound may indicate the circumstances in which the crime was committed.

• The age of the injury can be determined.

The image below shows a lacerated wound.



Other options

Option A: An incised wound is clean-cut, regular, and well-defined. Bruising does not occur when caused by light-cutting weapons but bruising and some irregularity of the margins may be present when heavy and not-so-sharp-edged weapons like a shovel, Kassi, axe, tangi, etc. have been used.

Option C: Chop wounds are caused by a blow with the sharp cutting edge of a fairly heavy weapon like an axe, sword, or hatchet. The margins of the wound depending on the condition of the weapon, a sharp margin with slight bruising is seen with a sharp-edged weapon, and a ragged and bruised margin is seen with a blunt edge. Undermining occurs in the direction towards which the chop is made.

Option D: Avulsion or shearing laceration is a lesion where the skin separates or peels off by tearing from the underlying tissues. It is due to shearing force delivered at an acute angle. Commonly seen in road traffic accidents.

Solution to Question 24:

The postmortem change shown above is marbling, which is an external sign of putrefaction.

In marbling, the superficial veins of roots of the limb, thighs, sides of the abdomen, and shoulders are stained greenish-brown or purplish-red, in a linear branching pattern. It is usually prominent 36–48 hours after death occurs due to the formation of sulfhemoglobin subsequent to hemolysis of red blood cells.

In putrefaction, bacteria from the gut enter into and multiply in the surrounding tissues, eventually spreading through the blood vessels to other parts of the body. The bacterium Clostridium welchii is largely responsible for the putrefactive changes spreading through the body, due to its production of the enzyme lecithinase. The first external sign of putrefaction in a body lying outside in the air is a greenish discoloration of the skin over the region of the cecum.

Other options:

Option B: Tattooing around the entry wound of a bullet is due to the deposition of unburnt powder particles. These are difficult to remove. They are also called stippling or peppering. This is seen up to 50 cm of firing

Option C: Postmortem hypostasis/Postmortem staining/Postmortem lividity/Livor mortis/Suggilation refers to bluish-purple or reddish-purple discoloration of the superficial layers of the dermis. It is an early postmortem change that occurs due to the stagnation of blood in the vessels of the dependent parts due to the effect of gravity.

Option D: Filigree burns also known as Lichtenberg's flowers are superficial, thin, irregular, and tortuous markings on the skin indicative of a lightning strike. They resemble the branches of a tree and are usually found over the skin of the shoulders or the flanks.

Solution to Question 25:

The clinical scenario is suggestive of a phossy jaw, which occurs due to chronic phosphorus poisoning. Phossy jaw refers to osteomyelitis and necrosis of the jaw, with foul-smelling pus discharging multiple sinuses.

Frequent inhalation of phosphorus fumes causes necrosis of the lower jaw in the region of a decayed tooth. Patients initially present with toothache, followed by swelling of the jaw, loosening of the teeth, necrosis of gums, and sequestration of the bone in the mandible producing considerable facial disfigurement.

Chronic phosphorus poisoning is common among the workers of phosphorus industries (matchbox factories), who have had chronic exposure to phosphorus fumes for months to years. It is managed surgically and by prevention of further exposure to the poison.

Phosphorus is a non-metallic, inorganic irritant poison that affects cellular oxidation. It exists in two forms, white phosphorus, a highly toxic form, and red phosphorus, the non-toxic form.

Fulminant poisoning is seen when more than one gram is consumed, and death occurs within 12 hours due to peripheral vascular collapse. In acute poisoning, they present with pain in the throat and abdomen, with intense thirst, nausea, diarrhea, and severe abdominal pain. System toxicity occurs after 2–3 days and results in hematemesis, enlarged tender liver, jaundice, pruritus, and hemorrhages. The gastric and intestinal content have garlic odor and are luminous.

Gastric lavage with potassium permanganate oxidizes phosphorus into phosphoric acid and phosphates. Activated charcoal can be used to adsorb the poison.

Other options:

Option A: Bromine is a corrosive substance that causes necrosis of skin and tissues.

Option B: The symptoms of antimony poisoning, both acute and chronic are similar to that of arsenic. Inhalation of fumes causes a flu-like illness called metal fume fever. In cases of chronic poisoning with heavy metals such as lead, arsenic, mercury, antimony, copper, etc., the shaft of a long bone, strands of scalp hair, fingernails or toenails, and a wedge of the quadriceps muscle should be taken for sampling.

Option C: Prolonged ingestion of fluorides results in the mottling of teeth, an increase in the density of bones, lipping of bony margins of vertebrae, which sometimes produce pressure on the spinal cord leading to paraplegia.

Solution to Question 26:

Durham's rule and Curren's rule are laws associated with insanity.

Durhams's rule states that an accused person is not criminally responsible if his unlawful act is the product of mental disease or mental defect.

Curren’s rule states that an accused person is not criminally responsible if, at the time of committing the act, he did not have the capacity to regulate his conduct to the requirements of the law as a result of mental disease or defect.

In India, the law of Criminal responsibility is under BNS Section 22 (Section 84 IPC) It states that "Nothing is an offense which is done by a person who, at the time of doing it, by reason of unsoundness of mind, is incapable of knowing the nature of the act, or what he is doing is either wrong or contrary to the law”. This is slightly different from McNaghten's rule followed in Britain.

M'Naghten's rule states that every man is to be presumed to be sane, and possesses a sufficient degree of reason to be responsible for his crimes until the contrary is proved.

Other rules to assess the criminal responsibility of insane persons:

• The Irresistible Impulse

• Federal rule or American Law Institute test (ALT)

• The Brawner rule

Note: Ashley's rule is used to differentiate males and females based on the combined length of the manubrium and sternum. It is considered to be a male sternum when the combined length is >149 mm, while if the combined length is <149 mm, then it is considered female.

Solution to Question 27:

The management of snakebites includes immobilization and administration of anti-snake venom (ASV) after hospitalization.

Immobilization is done to minimize the spread of the venom. Hospitalization, followed by ASV administration is the definitive management of snake bites.

Measures that can be taken in the case of snakebite:

• Apply firm pressure over the bitten area, which delays the absorption of venom

• Pressure immobilization is recommended for elapid and sea-snake bites, but not for viper bites as it may cause local necrosis

• Immediately apply a broad firm bandage (Sutherland wrap) on the bitten area and around the limb. As much of the limb should be bandaged as possible. It should be tight enough to occlude the superficial venous and lymphatic return, but not the arterial or deep venous flow. A pressure of 50 to 70 mm Hg is maintained.

• In bites on the trunk, head, or neck, apply firm pressure over the bitten area

• Immobilize the limb, as movement can accelerate the spread of venom

• Make the patient lie on one side in the recovery position so that the airway is clear, in case of vomiting or fainting.

Measures contraindicated in the case of snakebite:

• Suction and incision of the wound

• Electric shock treatment of wound site

• Application of tourniquets because they obstruct arterial flow and cause ischemia

• Ice water immersion of the bitten limb

• Do not attempt to kill or catch the snake as it may be dangerous.

• Administration of ASV locally at the wound site.

• Herbal/local remedies

Solution to Question 28:

Hooch tragedy is due to the consumption of cheap illicit liquor containing methanol, particularly by the lower socio-economic classes.

The central organ for metabolism is the liver. Methanol or methyl alcohol is first oxidized to formaldehyde with the help of alcohol dehydrogenase. It is further metabolized to formic acid via aldehyde dehydrogenase. These metabolites are responsible for the adverse effects of intoxication.

Features seen are ophthalmological complications such as blindness, and optic neuropathy. Brain edema, acute renal failure, and severe metabolic derangement are also seen.

Forensic Medicine NEET 2022

Question 1:

Identify the gun that caused the wound and the range of the shot.



a) Shot gun, intermediate range

b) Shot gun, close range

c) Pistol, near shot

d) Pistol, close shot

Question 2:

Identify the type of wound from the image.

a) Defense wounds

b) Antemortem wound

c) Postmortem wound

d) Hesitation cuts

Question 3:

A 56-year-old man presents with fatigue, pallor, and abdominal pain. He reports memory loss and reveals that he works at a battery recycling plant. On examination, a noticeable foot drop in both lower extremities is seen. Laboratory testing shows elevated blood lead levels. Which enzyme is affected in the given scenario?

a) Aminolevulinic acid (ALA) dehydratase

b) Uroporphyrinogen III

c) Uroporphyrinogen I

d) Pyruvate dehydrogenase

Question 4:

A 45-year-old female patient is told about the benefits and complications of hysterectomy and she agrees to the procedure. What kind of consent is this?

a) Informed consent

b) Implied consent

c) Opt out

d) Passive consent

Question 5:

A patient came with complaints of hair loss. His wife mentions that she has noticed some behavioral changes. The doctor notices that there is a loss of eyebrows on the lateral side. He then comes to a conclusion by examining the nails. What is the type of poisoning in this case?

a) Thallium

b) Arsenic

c) Mercury

d) Lead

Question 6:

A dead body is brought for evaluation. On post-mortem examination, a ligature that was completely encircling the neck, horizontal, and below the level of the thyroid was seen. There was no dribbling of saliva. What is the cause of death?

a) Throttling

b) Ligature strangulation

c) Gagging

d) Hanging

Question 7:

A surgeon returns home from a party after many pegs of alcohol and was called to perform an emergency operation. During the operation, the assisting staff noticed the surgeon's hand shaking and instruments falling. He eventually nicks an artery and the patient collapses. Under which of the following terms will this incident be tried?

a) Criminal negligence

b) Civil negligence not amounting to criminal negligence

c) Therapeutic misadventure

d) Dichotomy

Question 8:

During the court proceedings, the defense lawyer asks a leading question, which the prosecutor appeals against. The judge grants the appeal. Leading questions are not allowed in all of the following except?

a) Re-examination

b) Cross examination

c) Examination in chief

d) Dying declaration

Answer Key

Question No.	Correct Option
1	c
2	a
3	a
4	a
5	a
6	b
7	a
8	b

Detailed Explanations

Solution to Question 1:

The given image showing a gunshot entry wound with powder burn around the wound is likely due to a near-shot injury by a pistol.

A near-shot injury is one when the range of flame is within 60 cm.

The features of a near-shot wound are a circular entry wound with the size same as that of the bullet. The edges are lacerated, inverted, and surrounded by a narrow zone of abrasion. There are no signs of burning or singeing. Powder tattooing around the entry wounds is another important feature.

Other options:

Option A: In an intermediate range which is up to 7 feet, the edge of the wound is abraded and the appearance is referred to as a rat hole. There is no burning and the tattooing is appreciable till 3-4 feet and then starts disappearing.

Option B: In shotgun injuries due to a close-range i.e between contact and 3 feet, the entry wound is circular or elliptical with features of the scorching, singing, and less dense tattooing.

Option D: In a close shot pistol injury, the range of flame is within 5-8 cm with a circular entry wound. Other features such as burning, singeing, blackening and tattooing of skin are seen in and around the entry wound.

Solution to Question 2:

The given image shows irregular and ragged cut wounds over the palmar surface of the hands and fingers, which is suggestive of defense wounds.

Defense wounds are the wounds that occur as a result of the victim's attempt to save himself from an injury.

It can be active or passive. Active defense wounds occur when the victim tries to grasp the weapon, while passive wounds result when the victim raises the hands, arms, or legs in an attempt to protect the eyes, head, or neck.

The morphology of the wound depends upon the type of weapon and the attack. In a sharp injury due to a single-edged weapon, a single cut may be seen on the palm or the bends of fingers. In the case of a double-edged sharp weapon, cuts are seen on both palm and fingers. The cuts are irregular in-depth and are ragged.

The defense wounds signify homicide, and in females, they often indicate sexual assault.

Other options:

Options B and C: Antemortem and postmortem wounds are of medicolegal importance while evaluating trauma. They can be differentiated by the following features:

Option D: Hesitation cuts are a feature of self-inflicted wounds or suicide. It typically appears as multiple, small, and superficial wounds on the skin, at the beginning of an incised wound. The common sites are the throat, wrist, and front of the chest.

Feature	Antemortem wounds	Postmortem wounds
Edges	Swollen edges with wound ga pes	Closely approximated with n o wound gapes
Hemorrhage	Arterial type of hemorrhage	Venous hemorrhage
Spurting	Spurting of blood seen with fi rmly coagulated blood	No spurts and no clotting
Vital reactio n	Inflammation and repair see n	No signs of inflammation an d repair

Solution to Question 3:

The given scenario is suggestive of chronic lead poisoning, and the enzyme affected is aminolevulinic acid dehydratase or ALA dehydratase.

Lead is a heavy metal, which mainly acts by depressing the enzymes in heme synthesis, blocking the enzyme aminolevulinic acid dehydratase and inhibiting ferrochelatase.

Chronic lead poisoning also referred to as plumbism or saturnism occurs following inhalation of lead dust or vapors that arise as a result of burning of paints, batteries, smouldering, glass blowing, and polishing.

Features of lead poisoning:

• Facial pallor, which is the earliest and most consistent sign

• Anemia

• Burtonian line, or the lead line, is seen on the gums as a blue line. It occurs as a result of lead sulphide formation, which is deposited as granules in the upper jaw at the junction of the teeth.

• Abdominal colic, constipation

• Lead palsy is a late manifestation, commonly seen in adults and is characterized by numbness, hyperesthesia, and muscle cramps, followed by weakness. Other features like foot drop, wrist drop, and fatigue may be seen.

• Lead encephalopathy is characterized by vomiting, convulsions, headache, insomnia, visual disturbances, and irritability.

Diagnosis is mainly by history and clinical findings. Increased urinary lead and porphyrin excretion are suggestive of chronic lead poisoning. Radiograph, erythrocyte porphyrin assays, and basophilic stippling in the peripheral smear are also useful.

Prophylaxis in the form of proper ventilation, personal hygiene, periodic medical examination, supplementation of calcium-rich diet and sulphuric acid in the water, and weekly saline purgative in factory works is useful.

Management includes British Anti-Lewisite, or BAL, for severe acute poisoning, with and without encephalopathy; ethylenediamine tetraacetic acid (EDTA) for moderate poisoning;
and D-penicillamine in mild cases.

The below image shows basophilic stippling.



Burtonian line - Lead poisoning

Solution to Question 4:

This kind of consent is informed consent since the patient has been made aware of the benefits and complications and has agreed to the procedure.

Informed consent is one given by the patient after understanding the nature of the condition, proposed treatment, outcomes of the recommended treatment, alternative options to treatment, and the risks and benefits involved. The patient can accept or reject the procedure. This is also called the rule of full disclosure.

Consent refers to a voluntary agreement, compliance, or permission. It can be expressed or implied.

Implied consent (option B) is that which is not written or expressly stated. It is implied that the patient is giving consent to examination in a general sense. For example, when a patient visits the

hospital with an illness, it implies that he has given consent to a physical examination to find out the nature of his illness. Another example of implied consent is when a patient exposes his arm for injection. It is legally effective.

Expressed consent can be oral or written and refers to consent in which the terms are clearly stated in distinct and explicit language.

Opt out (Option C) refers to a patient's refusal of the procedure, test, or treatment. It is the physician's duty to disclose the risks or consequences of refusal adequately and appropriately to the patient.

Passive consent (Option D) is not commonly described. However, in regard to research, it refers to parents signing and returning a form if they do not want to allow their child to participate in research.

Solution to Question 5:

The given scenario of hair loss, loss of eyebrows on the lateral side along with behavioral and nail changes point to thallium poisoning.

Thallium poisoning occurs due to an overdose or idiosyncrasy. Thallium sulfate and thallium acetate are the two salts of thallium used in dyes, in the glass industry, and as rodenticides and pesticides.

Mild cases of poisoning have features of joint pain, loss of appetite, and drowsiness. Severe cases present with colic, vomiting, muscle pain, convulsions, psychosis, tremors, delirium, and coma are seen in severe cases. Hair loss is seen after 2 weeks and a characteristic feature is loss of the outer thirds of the eyebrows. Mees lines are seen on the nails. Maculopapular rashes in a butterfly pattern on the face are characteristic.

In chronic poisoning, alopecia, skin rashes, peripheral neuropathy, and mental confusion are present.

Treatment is mainly by stomach wash with potassium iodide or Prussian blue, activated charcoal. Hemodialysis is useful and forced diuresis can be given.

The image below shows Mees lines.

Solution to Question 6:

The most likely cause of death in a case where the ligature is completely encircling the neck, horizontal, and below the level of the thyroid, with no saliva dribbling, is ligature strangulation.

Strangulation is a type of asphyxia where there is neck constriction without suspension of the body. The pressure on the neck is applied either by bare hands, a ligature, or by any other instrument.

When strangulation is done by using a ligature like a constricting band or rope, it is called ligature strangulation. The constricting band is tightened by a force other than the victim's body weight. It is characterized by the ligature mark, which is transverse and completely encircles the neck below the thyroid cartilage. There is no dribbling of saliva. Bleeding may be present in the ears, nose, and mouth.

Ligature and manual strangulations are mostly homicidal. Marks of violence are usually seen on other parts of the body.

Other options:

Option A: Throttling or manual strangulation occurs due to compression of the neck by humans leading to asphyxia. Thumb impression and finger marks may be seen over the victim's neck.

Option C: Gagging is a type of asphyxia due to forcible closure of the mouth by a cloth or closing the mouth and nose and is tied around the head.

Option D: Hanging is characterized by an incomplete and oblique ligature mark on the neck. It is usually seen higher, between the chin and larynx. Saliva often dribbles from the mouth.

Solution to Question 7:

The incident mentioned in the given scenario will be tried under criminal negligence.

Criminal negligence is when a doctor lacks a gross skill or care while treating, which results in serious injury/death of the patient, or when an illegal act is performed by a doctor or when an assaulted person dies. Most cases of criminal negligence occur due to drunkenness or usage of drugs, which leads to impaired efficiency.

Some of the examples are amputation of the wrong finger; leaving an instrument, tube, or sponge in the abdomen; incompetent administration of general anesthetic; and administrating the wrong substance in the eye, leading to visual loss.

The doctor is prosecuted by the police and charged in criminal court, when it is proved that no medical doctor in his ordinary senses would have done it. Contributory negligence cannot be used as a defense in criminal negligence.

Other options:

Option B: Civil negligence occurs as a result of lacking care and skill in the profession. Contributory negligence can be cited as a defense in civil negligence.

Option C: Therapeutic misadventure is a case that causes injury or death to the individual due to an unintentional act by a doctor or hospital. The physician is not liable for the injuries unless some negligence on his part contributed to the occurrence of the injury.

Option D: Dichotomy is a type of unethical act, wherein a physician offers or receives any gift or bonus for referring or procuring any patient for medical or surgical treatment.

Solution to Question 8:

Leading questions are not allowed in all of the given options except cross-examination, i.e leading questions are allowed during cross-examination.

As per section 141 of the Indian Evidence Act, a leading question is one which is suggestive of the answer that is desired or one which includes a fact and leads to the admitting of a conclusive answer, such as yes or no.

Cross-examination is one wherein the witness is questioned by the opposite party's lawyer, i.e the lawyer for the accused or the defense lawyer. It is undertaken with the objective to demonstrate favorable facts to their case, to test the accuracy of the witness statements, to unearth new or old facts, and to discredit the witness.

The sequence of examination while recording evidence is as follows:

• Examination-in-Chief/direct examination by the party that summoned the witness to elicit all the facts from the witness. Leading questions are not allowed (option C). The only exception is if the judge is satisfied that the witness is hostile.

• Cross-examination is done by the defence lawyer.

• Re-examination is done by the party that summoned the witness to correct or clarify or detail the statements made by the witness during cross-examination. Leading questions are not allowed (option A).

Other options:

Option D: A dying declaration is an oral or written statement made by a person who is dying. The doctor should certify that the person is conscious and with normal mental faculties before the declaration is recorded. It is recorded by the executive magistrate, or a doctor in the presence of two witnesses, if there is no time for the magistrate to come. Leading questions should not be asked.

Forensic Medicine NEET 2023

Question 1:

A child before playing consumed fruit from the garden. After some time he developed a high fever, confusion, photophobia, and unable to urinate. What are the likely causative agent and the appropriate antidote used in this case?

a) Datura, Pralidoxime

b) Datura, Physostigmine

c) Yellow oleander, Pralidoxime

d) Yellow oleander, Physostigmine

Question 2:

The method of autopsy carried out en masse to remove from tongue to prostate is_____.

a) Virchow technique

b) Rokitansky technique

c) Ghon technique

d) Letulle technique

Question 3:

A 42-year-old HIV-positive patient died, and his body was brought for post-mortem. The method of autopsy to be done is_____.

a) Ghon technique

b) Letulle technique

c) Virchow technique

d) Rokitansky technique

Question 4:

A 10-year-old male child came to the casualty with difficulty in walking and pain in the perianal region. On subjecting the specimen from the perianal region to a test, it produces

yellow needle-shaped rhombic crystals with picric acid. What is the test done?

a) Barberio test

b) Florence test

c) Teichmann test

d) Acid phosphatase test

Question 5:

A patient was brought to casualty with seizures. He had multiple episodes and was taking herbal medicines for the same. He was conscious in between the episodes. During the episodes, he had arching of his back. What is the likely cause?

a) Strychnine

b) Ricinus

c) Nerium odorum

d) Datura

Question 6:

BNS Section 90 (IPC 314) deals with

a) Causing miscarriage with the consent of the mother

b) Causing miscarriage without the consent of the mother

c) Death of the mother by act done with intent to cause miscarriage

d) Causing the death of the quick unborn child by an act amounting to culpable homicide

Question 7:

The temperature of a body of a deceased person is found to be 39 degrees Celsius. Which of the following is the most probable reason?

a) Cyanide poisoning

b) Septicemia

c) Corrosive poisoning

d) Intra-abdominal hemorrhage

Question 8:

A farmer was sleeping in the field, and he felt a sting on his leg. He saw something moving away quickly. He then got drowsy and was taken to the hospital. He developed pain around the site and continued to bleed profusely from the wound site. The wound became red with blisters. Which of the following is the most likely cause?

a) Viper

b) Cobra

c) Wasp bite

d) Scorpion

Answer Key

Question No.	Correct Option
1	b
2	d
3	d
4	a
5	a
6	c
7	b
8	a

Detailed Explanations

Solution to Question 1:

The clinical vignette in the question is suggestive of Datura poisoning, for which the appropriate antidote is physostigmine, a reversible acetylcholine esterase antagonist. It is an effective treatment for severe anticholinergic poisoning.

Datura poisoning:

The common names include thorn apple, Jimson weed, and Devil's trumpet. The active ingredients include the anticholinergic alkaloids atropine, scopolamine, and hyoscyamine. The seeds or other parts of the plant can be ingested or smoked.

Toxic manifestations of Datura:

• Mnemonic: "hot as a hare, blind as a bat, dry as a bone, red as a beet, and mad as a hatter"

• Also, remember the D's of Datura

• Dry hot skin (Temp rises up to 42 C, a scarlet rash may also be appreciated).

• Dilatation of pupils along with loss of accommodation

• Diplopia

• Dryness of the mouth and throat, unquenchable thirst

• Dysphagia

• Dilatation of cutaneous blood vessels

• Dysarthria (Difficulty in talking)

• Drunken gait

• Delirium

• Drowsiness

• Death due to respiratory or cardiac failure

Treatment

• Gastric lavage or decontamination with activated charcoal or tannic acid: Gastric emptying is often delayed, and the small, plentiful seeds can become trapped among the gastrointestinal folds after ingestion; thus, gastric decontamination can be an important therapy.

• Physostigmine in severe toxicity given 1-2 mg IM/IV after every half hour as required: It is a physiological antidote. Pilocarpine may also be given.

Medico-legal importance:

In India, Datura is used as a stupefying poison prior to robbery, kidnapping, and rape. It is also known as a rail-road poison, as it is commonly encountered during a journey. Robbers disguised as saints may offer ‘prasad’ mixed with Datura seeds and rob the passengers.

Yellow oleander:

All parts of the plant are poisonous. The active principles are thevatin and thevatoxin, which are cardiac glycosides. The nectar of the flower yields poisonous honey. Symptoms include a burning sensation in the mouth, tingling of the tongue, dryness of the throat, vomiting, diarrhea, headache, dilated pupils, irregular action of the heart followed by drowsiness, coma, collapse, and death.

Treatment includes

• Gastric lavage

• Single-dose activated charcoal

• To combat acidosis IV sodium lactate and 5% glucose can be given

• Atropine, adrenaline, or noradrenaline (if blood pressure is low)

• Digoxin-specific antibodies

The given image shows the flowers of Cerebra thevatia.

Solution to Question 2:

The en masse removal of organs is done in the Letulle technique.

It is the removal of the cervical, thoracic, abdominal, and pelvic organs as a single organ mass. They are subsequently dissected into organ blocks.

Solution to Question 3:

Rokitansky's technique of autopsy will be done on patients with highly transmissible diseases such as HIV, Hepatitis B, etc.

It involves in situ dissection in part, combined with en block removal of organs.

Solution to Question 4:

The test done in the scenario is the Barberio test.

Barberio is used to detect seminal fluid. When a saturated aqueous or alcoholic solution of picric acid is added to the spermatic fluid, yellow needle-shaped rhombic crystals of spermine picrate are produced.

Other options:

Option B: Florence test is used to detect semen. If semen is present, dark brown rhombic or needle-shaped crystals of choline iodide appear immediately

Option C: The Teichmann test (hemin crystal test) is used to detect bloodstains. If bloodstain is present, faint yellowish-red to brownish-black rhombic crystals of hemin or hematin chloride

appear. When hydrogen peroxide is added, the hematin crystals give bubbles of gas.

Option D: The acid phosphatase test is a quantitative test for the presence of acid phosphatase enzyme in the semen. High levels of acid phosphatase from prostatic secretions are present in seminal stains (Normal vaginal secretions have low levels of acid phosphatase). Positive reactions are found till 36 hours after which the enzyme levels fall and disappear within 72 hours. In the absence of demonstrable sperm or aspermia, this test can be considered conclusive.

Solution to Question 5:

The given clinical vignette suggests the diagnosis of strychnine poisoning.

Strychnine poisoning occurs if a person ingests crushed seeds of Strychnos nux vomica. If swallowed uncrushed, the seeds of nux vomica have no poisonous action. The fatal dose is 15-30 mg of strychnine or one crushed seed.

Strychnine is a competitive antagonist of glycine at the postganglionic receptors in the motor neurons of the anterior horn of the spinal cord. This leads to increased activity in the motor neurons.

Patients typically present with

• Convulsions are clonic initially but eventually become tonic.

• Risus sardonicus results from the contraction of the jaws and facial muscles in which the corners of the mouth are drawn back.

• The convulsions are most marked in anti-gravity muscles so that the body typically arches in hyperextension, the posture known as opisthotonus.

• If the spasm involves the abdominal muscles, the body bends forward (emprosthotonos), or to the side (pleurosthotonus).

• If death is preceded by a severe convulsion, as in strychnine poisoning, it causes an increase in the body temperature known as postmortem caloricity.

The first step in the management of strychnine poisoning is to control convulsions with benzodiazepines or barbiturates to control seizures, and the patients should be placed in a dark room to avoid triggering further convulsions. Activated charcoal can be used to adsorb unabsorbed poison after a thorough stomach wash. Acidification of urine can be done to increase the excretion of strychnine.

Other options:

Option B: Ricinus poisoning leads to symptoms such as the burning of the oral mucosa, salivation, nausea, vomiting, severe abdominal pain, and bloody diarrhea. It may also cause drowsiness, convulsions, delirium, and death.

Option C: Nerium poisoning presents with headache, dizziness, respiratory difficulty, and nausea. Ingestion causes difficulty in swallowing and articulation, abdominal pain, vomiting, profuse frothy salivation, and diarrhea. Blood pressure falls, fibrillation, AV block, coma, respiratory paralysis, and death can occur due to cardiac failure.

Option D: Datura poisoning presents with

• Dry hot skin (Temperature rises up to 42°C, a scarlet rash may also be appreciated)

• Dilatation of pupils along with loss of accommodation

• Diplopia

• Dryness of the mouth and throat, unquenchable thirst

• Dysphagia

• Dilatation of cutaneous blood vessels

• Dysarthria (Difficulty in talking)

• Drunken gait

• Delirium

• Drowsiness

• Death due to respiratory or cardiac failure

Solution to Question 6:

BNS Section 90 (IPC 314) deals with the death of the mother due to miscarriage.

Important sections of IPC relating to miscarriages:

Note: A quick child is a fetus that has developed to such a stage that it moves within the womb of the mother. It shall mean an unborn child whose heart is beating, who is experiencing electronically measurable brain waves, who is discernibly moving, and who is so far developed and matured as to be capable of surviving the trauma of birth with the aid of usual medical care and facilities available in this state.

BNS Section	IPC section	Deals with
88	312	Causing miscarriage with consent
89	313	Causing miscarriage without consent
90	314	Death of the mother by act done with intent to cause miscarriage
91	315	Preventing the child from being born alive or causing it to die after birth
92	316	Causing the death of the quick unborn child by an act amounting to culpable homicide

Solution to Question 7:

The condition in which the temperature of the body remains raised for about the first two hours after death is termed postmortem caloricity and the most probable reason for this is septicemia.

The condition occurs

• When the regulation of body temperature has been severely disturbed before death as in sunstroke or pontine hemorrhage.

• When heat production in muscles is increased due to convulsions as in tetanus and strychnine poisoning.

• When there is increased bacterial activity as in septicemia and cholera.

Solution to Question 8:

Based on the given clinical findings, the most likely cause is a bite of viper snake, resulting in a haemotoxic envenomation.

The Russell's viper has a flat and triangular head with a V-shaped mark, white in color, with the angle of the V pointing forwards. Along the back, it has three rows of diamond-shaped spots. Other points of identification include broad plates on its belly, the small scales on the head, and a shield beneath the tail divided into two rows. It is commonly found throughout India.

Patients with Russell's viper bite usually present with severe pain, and redness around the bite site. Swelling usually develops followed by blisters in and around the wound site. Persistent bleeding from the bite site is a constant feature. Patients may develop hemorrhagic syndrome due to bleeding from various sites like gums, nose, rectum, etc. In severe cases, shock may develop. Bleeding and clotting time are usually prolonged. Death usually occurs due to shock or hemorrhage.

Other options:

Option B: Cobra venom usually produces neurotoxic effects with minimal local effects. Neurological manifestations of cobra venom include ptosis, paralysis of the face, palate, jaws, tongue, neck muscles, muscles of deglutition, and respiratory muscles.

Option C: Wasp stings cause immediate pain, a wheal and flare reaction, and local edema and swelling that subside in a few hours. Multiple stings can lead to vomiting, diarrhea, generalized edema, dyspnea, hypotension, and collapse.

Option D: Scorpion sting usually presents with localized redness and burning pain which is unable to localize due to its radiation along the dermatomes involved, localized swelling, and punctate hemorrhagic spots. Other features include headache, profuse perspiration, chest discomfort, paraesthesias, hypersalivation, cold extremities, and sometimes, priapism. Hypertension usually develops early, while pulmonary edema and myocarditis may develop later.

Forensic Medicine INI-CET 2023

Question 1:

Identify the following image.

a) Blister due to barbiturate poisoning

b) Blister due to antemortem burns

c) Edema due to chemical burn

d) Pemphigus vulgaris

Question 2:

A 5-year-old boy went to the temple in a village with his grandmother. He was crying inconsolably on the way back. He was taken to the emergency department within 3 hours. He had altered sensorium, cold clammy extremities, increased salivation, and excessive sweating. On examination, he was pale and had priapism with the following vitals - BP - 140/78 mmHg, HR - 152/min, RR - 36/min sPO2 of 96%. Which of the following would you use in his management?

a) Anti-snake venom

b) Methylprednisolone

c) Prazosin

d) Adrenaline

Question 3:

In an examination for drunkenness, a person smells of alcohol and his conjunctiva appears red. During the examination of motor coordination, it is noted that he can walk in a straight line and can talk normally. Which of the following statements is true?

a) He has consumed alcohol

b) He has consumed alcohol and is under the influence of alcohol

c) He has consumed alcohol but is not under its influence

d) He has not consumed alcohol

Question 4:

The Blackfoot disease is an endemic peripheral vascular disease seen in toxicity of which of the following metal?

a) Arsenic

b) Antimony

c) Barium

d) Cadmium

Question 5:

A 12-year-old girl is brought by a lady police sub-inspector with an alleged history of sexual violence 2 days back. She was accompanied by her mother. The registered medical practitioner should do which of the following as the next step?

a) Take a detailed history

b) Preserve clothing

c) Take consent for genital examination

d) Documentation of injuries and examination findings

Question 6:

Which of the following is not true about postmortem changes?

a) In drowning, postmortem hypostasis is seen only in the lower limbs

b) Greenish discolouration in right iliac fossa is the earliest sign of decomposition

c) Adipocere is seen in bodies recovered from water source

d) Cadaveric spasm is seen only in specific group of muscles

Question 7:

A female was brought for autopsy with an alleged history of being found at a lake in the village. All of the following are suggestive of antemortem drowning except:

a) Water in stomach

b) Washerwoman's hands and feet

c) White leathery foam around mouth and nostrils

d) Positive diatom test

Question 8:

In death due to firearm injury, the entry wound shows only an abrasion collar. The range of the firearm is:

a) Contact range

b) Close range

c) Near range

d) Distant range

Question 9:

In the 18th century, the British East India Company smuggled which of the following drugs to China?

a) Cocaine

b) Magic mushroom

c) Opium

d) Marijuana

Question 10:

An industrial worker sustained burn injuries to the anterior region of his chest and abdomen, as well as burns to his right upper limb and a circumferential burn to his right thigh, as a result of a cylinder explosion. What is the percentage of total burns surface area?

a) 27

b) 36

c) 45

d) 53

Question 11:

In a Firearm injury, blackening surrounding the gunshot entry wound is due to?

a) Flame effect

b) Unburnt powder

c) Smoke deposition

d) Muzzle of the firearm

Question 12:

Match the following poisons to their site of action:

a) A-3, B-1, C-4, D-2

b) A-2, B-3, C-1, D-4

c) A-4, B-2,C-1, D-3

d) A-4, B-2,C-3, D-1

Question 13:

A 13-year-old female was brought by her mother to the emergency with abdominal pain for a few days. On clinical examination by the RMP, they found that she is 8 weeks pregnant. What should be your immediate next step?

a) File an MLC and Inform Police

b) Obtain consent form her mother for genital examination.

c) Preserve the clothing

d) Plan/perform MTP

Question 14:

A young man is brought to the casualty with pinpoint pupils. Which of the following is the most likely substance he would have consumed?

a) Datura

b) Opium

c) Alcohol

d) Cocaine

Question 15:

A 25 year old man was brought to the casualty in a semiconscious state following a road traffic accident. Which type of injury is unlikely to be seen in this patient?

a) Graze abrasions

b) Laceration Wound

c) Multiple Contusions

d) Chop Wound

Question 16:

An autopsy was performed on a person with a history of drowning; however, the report revealed that there was no water in his lungs, hence the case was labeled as dry drowning. What is the most likely cause of his death?

a) Vagal inhibition

b) Laryngospasm

c) Pulmonary edema

d) Electrolyte imbalace

Question 17:

Cadaveric spasm is not seen in?

a) Drowning

b) Traumatic asphyxia

c) Firearm suicidal wounds

d) Electrocution

Question 18:

Which of the following is not considered a grievous hurt?

a) Unilateral permanent hearing loss

b) Emasculation

c) Fracture of limb

d) 7 days admission in yellow zone of hospital

Question 19:

All of the following can be used for the identification of a dead body, except?

a) Blood

b) Nail

c) Muscle

d) Teeth

Question 20:

Which chelating agent is not used in heavy metal poisoning?

a) British anti-Lewisite

b) Ethylenediaminetetraacetic acid

c) Pralidoxime

d) Penicillamine

Answer Key

Question No.	Correct Option
1	b
2	c
3	c

4	a
5	a
6	a
7	b
8	d
9	c
10	b
11	c
12	a
13	a
14	b
15	d
16	b
17	b
18	d
19	b
20	c

Detailed Explanations

Solution to Question 1:

The given image of burns with blisters with surrounding hyperemia points to a diagnosis of antemortem burns.

A zone of hyperemia (line of redness), which varies in width is present at the edge of the burnt area in antemortem burns. It is due to edema of tissues and capillary dilatation which merges with the edge of the burn. It involves the whole thickness of the actual skin. It is permanent and persists even after death. If the entire body is burnt, a line of redness will be absent.

The antemortem blister appears as a raised dome. It contains serous fluid with proteins and chlorides. The base shows reddening with swollen papillae. In antemortem burns, the skin adjacent to the burnt area shows an increase in enzyme reaction. The reaction is increased for sulfhydryl groups in all layers. There is also an increase in ATP, esterases, aminopeptidases, and acid and alkaline phosphatase. Marked cellular exudation and reactive tissue changes are called vital reactions. This is characteristic of antemortem burns.

Other options-

Option A: Blister due to barbiturate poisoning, is commonly seen in frictional areas such as interdigital cleft, genital regions, etc. It is due to the dermo necrotic effect of barbiturates.

Option C: Chemical burns by acid or alkalis appear as ulcerated patches without blisters. The hair is not singed and the red line of demarcation is absent. Sometimes, the burn shows distinct

coloration and is usually uniform in character. Strong acids produce dark leathery burns on the skin. Strong alkalis cause the skin to slough and leave moist, slimy, greyish areas.

Option D: Pemphigus vulgaris is an intraepidermal immunobullous disease. It occurs commonly between the 4th to 6th decades. They present with flaccid blisters. Mucosal involvement will be seen in the form of buccal and palatal erosions. Firm pressure over the lesion separates the epidermis from the dermis. Nikolsky's sign is positive.

Solution to Question 2:

The clinical stem points to a likely diagnosis of a scorpion bite, for which Prazosin can be used in the management. It acts as a physiological antidote counteracting the symptoms of the adrenergic surge.

The venom is a potent autonomic stimulator that causes the release of massive amounts of catecholamines from the adrenals. It also has a direct effect on the myocardium.

Scorpion sting usually presents with localized redness and burning pain which is unable to localize due to its radiation along the dermatomes involved, localized swelling, and punctate hemorrhagic spots. Other features include headache, profuse perspiration, chest discomfort, paraesthesias, hypersalivation, cold extremities, and sometimes, priapism. Hypertension usually develops early, while pulmonary edema and myocarditis may develop later.

Treatment:

• Paracetamol, NSAIDs, or opioids for pain.

• Benzodiazepines for sedation for agitation, oculomotor abnormalities, uncoordinated neuromuscular activity, and muscle spasms.

• Hypertension and pulmonary edema- Prazosin counteracts the adrenergic surge that can lead to these effects.

• Cardiogenic shock- Antivenom, prazosin, nitroglycerin, dobutamine.

• Cholinergic effects- Atropine.

• Supportive care, wound care.

Solution to Question 3:

The given scenario of a person who smells of alcohol but whose motor coordination is normal with no slurring of speech is suggestive that he has consumed alcohol but is not under its influence.

The effects of alcohol intoxication at various blood alcohol concentrations are listed below.

Smell of alcohol	Motor coordination	Inference
Present	Normal	Consumed alcohol, not under the influence
Present	Abnormal	Consumed alcohol, under the influence
Absent	Normal	Not consumed alcohol
Absent	Abnormal	Cerebellar disorders

Blood alcohol concentration (mg/dL)	Effects
30-50	Impairment of driving and similar skills
50-100	Reduced inhibitions, slurred speech, laughter
100-150	Incoordination, ataxia
150-200	Obvious drunkenness, nausea, staggering gait
200-300	Vomiting, stupor
300-350	Stupor, coma, risk of aspirating vomitus
350-450 and above	Danger of death from respiratory paralysis

Solution to Question 4:

The Blackfoot disease is due to chronic arsenic poisoning.

In China (Province of Taiwan) exposure to arsenic via drinking water has been shown to cause a severe disease of the blood vessels, which leads to gangrene, known as 'black foot disease'. The Blackfoot disease is due to platelet activation & hypercoagulability of blood in peripheral

arteries and is observed in endemic areas of arsenicism.

Clinicopathological findings in acute and chronic arsenic poisoning:

Option B: The symptoms of antimony poisoning, both acute and chronic are similar to that of arsenic. Inhalation of fumes causes a flu-like illness called metal fume fever. In cases of chronic poisoning with heavy metals such as lead, arsenic, mercury, antimony, copper, etc., the shaft of a long bone, strands of scalp hair, fingernails or toenails, and a wedge of the quadriceps muscle should be taken for sampling.

Option C: Barium poisoning causes nausea, vomiting, and diarrhea. It causes cramps and stiffness of the muscles, dilation of the pupil, paralysis of the tongue and larynx, and vertigo. There is increased blood pressure with slow and irregular pulse.

Option D: Cadmium poisoning causes proteinuria and painful bone lesions known as Ouch-Ouch disease.

System	Acute	Chronic
Skin	Delayed hair loss (alopecia)	Melanosis, Bowen's disease f acial edema, hyperkeratosis, cutaneous cancer, hyperpigm entation.
Nails	nil	Mees's lines
Neurologic	Hyperpyrexia, convulsions, tr emors, coma	Encephalopathy, polyneurop athy, tremors, axonal degene ration
GI tract	Abdominal pain, dysphagia, vomiting, bloody or rice wate r diarrhea, mucosal erosions.	Nausea, vomiting, diarrhea, a norexia, weight loss
Liver	Fatty infiltration	Hepatomegaly, jaundice, cirr hosis
Kidney	Tubular and glomerular chan ges – oliguria and uremia	Nephritic finding
Haematologic	thrombocytopenia, impaired folate metabolism	Bone marrow hypoplasia, an emia, leukopenia, basophilic stippling, and karyorrhexis.
Cardiovascular	The heart will show S-T wave abnormalities, prolonged QT interval, ventricular fibrillati on, atypical ventricular tachy cardia with humic acid in the Southwest coast of Taiwan	Blackfoot disease due to plat elet activation and Hypercoa gulability of blood in periphe ral arteries is observed in end emic areas of arsenicism.
Reproductive syst em	Maternal ingestion of inorga nic arsenic crosses the placen tal barrier and infant death s hortly after birth.	-

Solution to Question 5:

The registered medical practitioner should take a detailed history as the next step.

The registered medical practitioner must take a detailed history and then take informed consent for examination. In case of refusal, an informed refusal is to be documented after explaining to them the consequences of choice. As part of the examination, they should collect specimens, preserve clothing, and document injuries and examination findings. Medical treatment and testing for pregnancy have to be done for all victims of sexual assault free of cost and police should be informed as per section 357-C of CrPc.

The Protection of Children from Sexual Offences (POCSO) Act, 2012 considers all individuals (both male and female) below the age of 18 years as children and protects them from sexual abuse. It defines various forms of sexual offenses which include:

• Sexual harassment: A person is said to commit sexual harassment upon a child when such person with sexual intent,

• Carries out any act with sexual connotations

• makes a child exhibit his/her body

• shows any object to a child in any form/media for pornographic purposes.

• Sexual assault

• Any act with sexual intent that involves physical contact without penetration.

• Aggravated sexual assault

• Similar to sexual assault along with the criteria explained under the term "aggravated".

• Penetrative sexual assault

• An act with sexual intent that involves physical penetration.

• Aggravated penetrative sexual assault

• Similar to penetrative sexual assault along with the criteria explained under the term "aggravated".

• Pornography

The term aggravated is used for those assaults done:

• by a person in a position of trust or authority such as police officer, security forces, relative, doctor.

• in the course of communal violence.

• on mentally & physically disabled.

• gang sexual assault.

• using deadly weapons.

• causing grievous hurt

The Act prescribes stringent punishment graded as per the gravity of the offense, with a maximum term of rigorous imprisonment for life and fine. Section 21(1) of the POCSO Act, 2012 requires mandatory reporting of cases of child sexual abuse to law enforcement authorities and applies to

everyone including parents, doctors, and school personnel. Failure to report a suspicion of child abuse is an offense under the Act.

Solution to Question 6:

In the case of drowning, if the water is still, the postmortem hypostasis affects the head, chest and the limbs. This is because the chest and abdomen float higher because of the lungs and the abdominal gases. Drowning in fast-flowing water results in absent postmortem staining as the position of the body keeps on changing.

Postmortem hypostasis/Postmortem staining/Postmortem lividity/Livor mortis/Suggilation refers to bluish-purple or reddish-purple discoloration of the superficial layers of the dermis. It is an early postmortem change that occurs due to the stagnation of blood in the vessels of the dependent parts due to the effect of gravity.

Sites of postmortem hypostasis depending on the position of the body:

Postmortem changes are classified on the basis of time:

• Immediate signs of death (at the time of somatic or clinical death):

• Insensibility and loss of voluntary power

• Cessation of respiration

• Cessation of circulation

• Early signs of death. It is seen in cellular death:

• Pallor and loss of elasticity of the skin

• Eye changes

• Primary flaccidity of muscles

• Algor mortis/Postmortem cooling

• Postmortem lividity/Suggillation

• Rigor mortis

• Late signs of death:

• Putrefaction

• Adipocere

• Mummification

Other options-

Option B: The first external sign of putrefaction in a body lying outside in the air is a greenish discoloration of the skin over the region of the right iliac fossa. Sulphmethemoglobin is the agent imparting greenish color.

Option C: Adipocere is a modification of putrefaction. The transformation is brought about by the gradual hydrolysis and hydrogenation of pre-existing fats, including olein, into higher fatty acids. It is characterized by the formation of a waxy, soft, moist, and whitish substance with a

sweetish offensive or ammoniacal (early stages) odor. It is more commonly seen in bodies recovered from damp, warm environments or immersed in water.

Option D: Cadaveric spasm - This is a vital sign of antemortem drowning. Cadaveric spasm is limited to select muscles i.e., the muscles that were in a state of contraction at the time of death. The mechanism of cadaveric spasm is thought to be due to neurogenic phenomena. Gravel, weed, soil, grass, etc. is firmly held in the clenched hands due to cadaveric spasm. The material also indicates the place of submersion.

Position of the body	Sites of hypostasis
Supine	Back except for shoulder blades, buttocks, and calves (contact pallor)
Vertical as in hanging	Legs, feet, forearm, and hands (glove and stocking hypostasis)
Drowning	Face, the upper part of chest, hands, lower arms
Inverted like in drunken persons	Head and neck

Solution to Question 7:

Washerwoman's hands and feet are a non-specific sign of submersion. They occur due to the imbibition of water into the outer skin layers. They can be seen in antemortem as well as postmortem drowning victims.

The presence of water in the stomach, white leathery foam around the mouth and nostrils, and a positive diatom test are suggestive of antemortem drowning.

Given below are the features of antemortem drowning.

• Froth - The presence of copious amounts of fine, shaving-lather-like froth at the mouth and the nostrils is the most characteristic finding of antemortem drowning. It is caused by the vigorous agitation of the tracheal and bronchial mucous secretions, edema fluid, and alveolar surfactant during respiratory efforts.

• Cutis anserina - The contraction of the erector pilorum muscles causes puckering and grainy appearance of the skin. This resembles goose skin; hence, it is called cutis anserina.

• Washerwoman’s hand - Maceration of the skin on the palms and soles gives a wrinkled, sodden, and bleached appearance due to submersion.

• Cadaveric spasm - Gravel, weed, soil, grass, etc. is firmly held in the clenched hands due to cadaveric spasm. This is a vital sign of antemortem drowning. The material also indicates the place of submersion.

• Rigor mortis - This appears early in cases of a violent antemortem struggle.

• Face - It is bloated and discolored, with cyanosis.

• Eyes - Conjunctiva are suffused and pupils dilated.

• Tongue - It may be swollen and protruded.

• Antemortem injuries - These may be sustained by striking objects, such as rocks, during the fall into the water body. Examination for such injuries must be performed only after the body is dry.

Internal findings:

• Lungs - They appear voluminous and distended and show ballooning. The cut section will exude copious amounts of frothy, bloodstained fluid.

• Larynx, trachea, and bronchioles - The presence of sand, silt, and aquatic vegetation in the trachea and lower respiratory tract are characteristic findings of antemortem drowning.

• Heart and blood vessels- Gettler test can be done to estimate the chloride content in the right and left sides of the heart to differentiate between freshwater and saltwater drowning.

• Diatom test- Diatoms are microscopic algae. Their presence in the organs, particularly in the bone marrow, is suggestive of antemortem drowning.

• Stomach and small intestines- These organs may contain water in some cases. Water present in the small intestine is positive evidence of death by drowning.

• Middle ear - The presence of water in the middle ear is highly suggestive of antemortem drowning.

• Brain - It is congested, and edematous, with loss of gray-white junction.

Solution to Question 8:

The entry wound shows only an abrasion collar when the range of the firearm is distant.

To determine the range of firing, the following parameters are used:

• The entry wound size and shape

• Blast effect – Muzzle end imprint and edge eversion

• Blackening, Tattooing, Singeing of hair, Abrasion collar, and Grease collar

• Hair should be preserved to determine the range of firing which can be examined for carbonization/desiccation in close range, also from lead deposition around the wound to assess the range of firing.

Abrasion collar: Gyroscopic movement of the bullet, the skin's elasticity, and the grease and dirt particles on the bullet's surface are the major causative factors of abrasion and grease collar. They are seen on the skin around the wound of entry at all ranges. Depending on the angle of entry made by the bullet, i.e.

• A perpendicular entry of the bullet develops the abrasion collar of rounded shape.

• An acute angular entry of the bullet develops the abrasion collar of an oblique or oval shape.

• Some contusion present in the abraded collar is known as contusion collar

Characteristics	Contact shot	Close Shot	Near Shot	Distant Shot
Range	In-touch with skin	2.5-7.5 cm	30-60 cm	>60 cm
Size	Largest size	Bullet size	Smaller	Smallest
Shape	Stellate	Circular	Circular	Circular
Blast effect				
Muzzle end imprint	+ve	-ve	-ve	-ve
Edge eversion	+ve	-ve	-ve	-ve
Other findings				
Blackening	-ve	+ve	-ve	-ve
Tattooing	-ve	+ve	+ve	-ve
Singeing	-ve	+ve	-ve	-ve
Abrasion collar	-ve	+ve	+ve	+ve
Grease collar	-ve	+ve	+ve	+ve

Grease collar: Mostly seen on the clothing and is seen as a black deposit, sharply outlined as if printed (better-seen on light color clothes and on fair-skinned individuals).

Medicolegal importance: Helpful in deriving information such as wound of entry and direction of firing



Note: A grease collar need not always be present. Hard metal jacketed bullets are associated less commonly with soiling. Often, the ring is not appreciable during gross inspection and may only be demonstrable on microscopic examination or other laboratory tests.

Solution to Question 9:

The British East India Company smuggled opium into China in exchange for tea, silk and porcelain pots.

Opium derived from the immature seedpods of the opium poppy (Papaver somniferum), was originally indigenous to the present-day territory of Turkey. The introduction of opium to China occurred during the late 6th or early 7th century CE, facilitated by Turkish and Arab traders.

During this time, tobacco smoking moved from North America to China, resulting in widespread opium smoking. Opium addiction and imports increased in the first century of the Qing era (1644–1911/12). To address the issue, the Yongzheng emperor (1722–35) banned opium trafficking and consumption in 1729. Despite hurdles, the trade continued, and in 1796, the Jiaqing emperor banned opium imports and cultivation. Opium sales survived despite these laws.

In the early 18th century, the Portuguese made a lot of money importing opium from India and selling it in China. British traders became China's main suppliers in 1773. The British East India Company monopolized Bengali opium farming. The company developed an inexpensive and productive opium poppies cultivation method in this location. The US and other Western nations traded Turkish and Indian opium.

The opium trade was pursued by Britain and other European nations as a means to address their persistent trade deficit with China. In Europe, there existed a substantial and notable demand for Chinese commodities such as tea, silks, and porcelain pots. Conversely, in China, there was a very limited market for Europe's manufactured goods and other trade items. As a result, Europeans were required to utilize gold or silver as payment for Chinese goods. The opium trade, which resulted in a consistent demand for opium among Chinese individuals addicted to the substance, effectively addressed the persistent trade imbalance by facilitating the importation of opium from Western countries.

Opium was not carried by the East India Company. Due to the Chinese restriction, the corporation outsourced it to “country traders”—private traders entitled to transport commodities from India to China. Smugglers on the Chinese shore bought opium from country traders. The traders gave the East India Company their sales of gold and silver. The corporation spent its gold and silver in China on profitable things to sell in England.

Opium imports into China rose from 200 chests in 1729 to 1,000 in 1767 and 10,000 between 1820 and 1830. Although chest weight varied by origin, it averaged 140 pounds (63.5 kg). By 1838, China imported 40,000 chests annually. China lost the balance of payments to Britain for the first time. Opium distribution networks had evolved across China, frequently with corrupt officials' help. High opium addiction levels affected imperial troops and officials.

Solution to Question 10:

The affected percentage of body surface area (BSA) is calculated using Wallace's rule of 9:

• Right upper limb - 9%

• Circumferential right thigh - 9% (Anterior thigh 4.5% and Posterior thigh 4.5% in Adults)

• Anterior chest - 9%

• Anterior abdomen - 9%

Thus, totaling to 36%.

Note: According to Bailey's Short Practice of Surgery, 28th edition, the anterior thigh accounts for 4%, while according to Current Practice of Surgery, 14th edition, the anterior thigh accounts for 4.75 %. Either percentage, when totalled gives us the best possible answer to this question to be 36%.

The following image shows Wallace's rule of 9:



Lund and Browder's chart is an accurate method to measure burns if it is used correctly. In this, different proportions of body surface areas in children are taken into consideration. This chart compensates for the changes in the shape of the body with age and can accurately assess the burn area in children.

The below image shows the Lund and Browder chart.



	0	1	5	10	15	Adult
A: half of head	9½	8½	6½	5½	4½	3½
B: half of thigh	2¾	3¼	4	4¼	4½	4¾
C: half of leg	2½	2½	2¾	3	3¼	3½

Solution to Question 11:

Blackening occurs due to the deposits of smoke over the epidermis which can be removed with a wet cloth. This can be seen up to around 15 cm of firing.

Other injuries related to firearms are:

• Burn injuries - occur due to flame released during the firing. It is seen in contact shot wounds or distances up to 5-8 cm. Singeing of hair is caused due to burns.

• Tattooing - occurs due to the deposition of unburnt powder particles. These are difficult to remove. They are also called stippling or peppering. This is seen up to 50 cm of firing

• Greasing and collaring - occur due to the deposition of the lubricant of the bullet in the tissues, during its penetration. It can be found in contact, near-shot, or distant shot wounds.

In a firm contact shot, the imprint of the muzzle of the gun is seen on the skin around the wound. The victim is within the range of flame, hence singeing of the hair will be present. Tattooing begins only when the muzzle-to-target distance is >1cm, like in a loose contact shot.

When the distance between the victim and the firing is 5-8cm, it is called a close shot. The victim is within the range of the flame, hence, features related to flame, smoke, and powder deposition will be seen. Singeing, burning, blackening and tattooing will be present.

When the distance between the victim and the firing is >50 cm, it is a distant shot. The victim lies outside the range of flame and powder deposition. Hence, features related to flame, smoke, and powder deposition are absent. Burning, blackening, and tattooing are absent.

Solution to Question 12:

Datura poisoning

The common names include thorn apple, Jimson weed, and Devil's trumpet. The active ingredients include the anticholinergic alkaloids atropine, scopolamine, and hyoscyamine. The seeds or other parts of the plant can be ingested or smoked.

Toxic manifestations of Datura:

Mnemonic: "Hot as a hare, blind as a bat, dry as a bone, red as a beet, and mad as a hatter"

Also, remember the D's of DaturaDry hot skin (Temp rises up to 42 C, a scarlet rash may also be appreciated).

• Dilatation of pupils along with loss of accommodation

• Diplopia

• Dryness of the mouth and throat, unquenchable thirst

• Dysphagia

• Dilatation of cutaneous blood vessels

• Dysarthria (Difficulty in talking)

• Drunken gait

• Delirium

• Drowsiness

• Death due to respiratory or cardiac failure

Medico-legal importance:

In India, Datura is used as a stupefying poison before robbery, kidnapping, and rape. It is also known as a rail-road poison, as it is commonly encountered during a journey. Robbers disguised as saints may offer ‘prasad’ mixed with Datura seeds and rob the passengers.

The following are images of the Datura plant and seeds:

Poison	Site of action
A. Datura	CNS
B. Nicotine	Cardiac muscle
C. Nux Vomica	Spinal Cord
D. Nitric acid	Skin

Nicotine is the principal constituent of tobacco responsible for its addictive behaviour. One of the newest agents introduced to aid in smoking cessation is varenicline.

Adverse effects of nicotine intoxication:

• Doses of 60 mg are toxic to the person and can cause respiratory paralysis.

• Nausea, vomiting, abdominal pain, diarrhoea

• Salivation

• Increased blood pressure, tachycardia, tremors and cardiac arrhythmias.

Other cardiac poisons include Oleander and Digitalis.

Strychnine poisoning occurs if a person ingests crushed seeds of Strychnos nux vomica. If swallowed uncrushed, the seeds of nux vomica have no poisonous action. The fatal dose is 15-30 mg of strychnine or one crushed seed.

Strychnine is a competitive antagonist of glycine at the postganglionic receptors in the motor neurons of the anterior horn of the spinal cord. This leads to increased activity in the motor neurons. Patients typically present with risus sardonicus, convulsions that are clonic initially but eventually become tonic and are most marked in anti-gravity muscles so that the body typically arches in hyperextension, the posture known as opisthotonus. If the spasm involves the abdominal muscles, the body bends forward (emprosthotonos), or to the side (pleurosthotonus)

Nitric acid is a corrosive inorganic acid. Due to the xanthoproteic reaction, the nitric acid combines with aromatic amino acids like tyrosine to form picric acid which is responsible for the yellowish discolouration of the skin.

Solution to Question 13:

The registered medical practitioner should take register a medico-legal case and immediately inform the police in this case. All three other options are steps to be taken eventually in this case but registering an MLC takes priority.

The Protection of Children from Sexual Offences (POCSO) Act, 2012 considers all individuals (both male and female) below the age of 18 years as children and protects them from sexual abuse. It defines various forms of sexual offenses which include:

• Sexual harassment: A person is said to commit sexual harassment upon a child when such person with sexual intent,

• Carries out any act with sexual connotations

• makes a child exhibit his/her body

• shows any object to a child in any form/media for pornographic purposes.

• Sexual assault

• Any act with sexual intent that involves physical contact without penetration.

• Aggravated sexual assault

• Similar to sexual assault along with the criteria explained under the term "aggravated".

• Penetrative sexual assault

• An act with sexual intent that involves physical penetration.

• Aggravated penetrative sexual assault

• Similar to penetrative sexual assault along with the criteria explained under the term "aggravated".

• Pornography

The term aggravated is used for those assaults done:

• by a person in a position of trust or authority such as police officer, security forces, relative, doctor.

• in the course of communal violence.

• on mentally & physically disabled.

• gang sexual assault.

• using deadly weapons.

• causing grievous hurt

The Act prescribes stringent punishment graded as per the gravity of the offense, with a maximum term of rigorous imprisonment for life and fine. Section 21(1) of the POCSO Act, 2012 requires mandatory reporting of cases of child sexual abuse to law enforcement authorities and applies to everyone including parents, doctors, and school personnel. Failure to report a suspicion of child abuse is an offense under the Act.

Solution to Question 14:

Pinpoint pupils can be observed in the overdose of opium.

Opioid overdose symptoms:

• Triad of coma, respiratory depression, and pinpoint pupils (miosis)

• Hypotension, bradycardia

• Shallow breathing

• Hypothermia

Naloxone is also used for reversing neonatal asphyxia due to opioid usage in labour. Naloxone is a pure opioid antagonist which reverses all effects except sedation.

Note: Naloxone is contraindicated in infants born to mothers with opiate addiction as it may precipitate acute neonatal withdrawal with seizures.

Causes ofMiosis	Causes ofMydriasis
Carbolic acid poisoning	Daturapoisoning
Morphinepoisoning	Cyanides
Organophosphorous poisoning	Cocaine
Pontine haemorrhage	Ether
Barbiturate poisoning	Pethidine
	Nux vomica
	Viper venom
	Amphetamine

Solution to Question 15:

A chop wound would be unlikely to be found in this patient who has succumbed to a road traffic accident.

Injuries that are common following a blunt impact (here road and traffic accident) include:

• Bruises and Contusions

• Laceration

• Abrasion (patterned due to tyre marks)

Chop wounds or slash wounds are the wounds produced by chopping motion, made with heavy and sharp weapons like axes, cleave, etc. The dimensions of the wound correspond to the crosssection of the penetrating part of the weapon. The margins are sharp and show slight abrasion and bruising with the destruction of underlying organs. The presence of an incised wound on the skin with an underlying fracture or deep grove in the bone indicates a chop wound. The majority of these injuries are homicidal in nature and are typically inflicted upon the vulnerable regions of the body, such as the head, face, neck, shoulders, and extremities. Accidental injuries can be attributed to various sources such as power fans, band saws, or ship and aeroplane propellers. These incidents have the potential to cause significant lacerations to soft tissues or result in the amputation of body parts. Instances of self-inflicted lacerations with suicidal intent have seldom been observed.

Individuals who sustain injuries as a result of road accidents can be categorized into three categories: pedestrians, cyclists (both pedal-powered and motorized), and occupants of cars, including both drivers and passengers. Among these three categories, pedestrians are frequently the ones who sustain injuries, however, the percentage of pedestrian casualties exhibits significant variation depending on the traffic patterns observed in different nations.

Injuries to pedestrians:

• Primary impact injuries are caused by the first impact of the vehicle on the victim.

• Secondary impact injuries are further injuries caused by the vehicle.

• Tertiary injuries are caused when the victim strikes other objects or falls on the ground after being hit by the vehicle.

Note: In case a pedestrian is hit by the vehicle only once (i.e only primary impact and no secondary impact by the vehicle) and then sustains injuries by falling on the road, it would be an example of secondary injury.



Injuries to occupants:

Approximately eighty per cent of vehicle impacts are against another vehicle or a stationary object and involve the front or front corners. Vehicle deceleration is rapid in this impact. Rarely, the car is hit from behind, causing ‘acceleration’. The least common impacts are side and 'roll-overs'.

In a typical unrestrained deceleration condition, front-seat occupants surge forward and hit car structures in front of them.

An unrestrained impact may cause the following:

• The face and head following a collision with the windscreen glass, frame, or side pillars, cause skull and cervical spine fractures along with brain injury.

• Contact with the fascia or steering wheel can cause rib, sternal, heart, and liver damage.

• Hyperflexion may cause aorta tears at the descending part of the arch, where the vessel attaches to the vertebral column.

• The passenger's knees may hit the parcel shelf, causing fractures.

• The driver's legs, supported by stop and clutch pedals, may transfer impact force to the pelvis via the tibia and femur.

• The rebounding force from the contact may cause the head to violently swing backwards, damaging the cervical or thoracic spine.

Unrestrained rear occupants are projected forwards and impact the back of the front seats, including headrests if equipped, during severe deceleration. They may be hurled over the seats, striking and injuring the front-seat occupants, or they may be expelled through the windscreen, which has been damaged by them or the persons in front.

Injuries inflicted on motorcyclists following collision with other vehicles include patterned abrasions and various head, chest and spine injuries. A distinct injury can arise from the practice of "tail-gating" in which a motorcycle manoeuvres beneath the rear of a vehicle, resulting in significant cranial trauma or, in extreme cases, decapitation. The implementation of protective bars positioned at the sides and rear of trucks has effectively mitigated the occurrence of injuries resulting from bikes and vehicles passing beneath the vehicle.

Solution to Question 16:

In dry drowning death is due to sustained laryngeal spasm and subsequent airway closure. This occurs when there is sudden entry of water into the nasopharynx or larynx.

Dry drowning:

In this type of drowning, water does not enter the lungs, but death results from immediate sustained laryngeal spasm due to an inrush of water into the nasopharyngeal or larynx. Thick mucus, foam and froth may develop, producing a plug.

Immersion syndrome (Hydrocution/Submersion inhibition):

Death results from cardiac arrest due to vagal inhibition as a result of

• Cold water stimulates the nerve endings of the surface of the body.

• Water striking the epigastrium.

• Cold water enters ear drums, nasal passages, pharynx and the larynx which cause stimulation of the nerve endings of the mucosa.

Secondary drowning/Post immersion-syndrome/Near drowning:

It refers to a submersion victim who is resuscitated and survives for 24 hours. The patient may or may not be conscious. These patients may develop hypoxemia resulting in brain damage, electrolyte disturbances, pulmonary edema, sepsis, myocardial anoxia, etc., Death may occur from half to several hours after resuscitation in about 10-20% of people. Those who survive suffer from serious neurological damage.

Type of drowning	Cause of death
Freshwater drowningSeawater drowning	Disruption of surfactant
Dry drowning	Laryngeal spasm
Immersion syndrome	Vagalinhibition
Post immersionSyndrome/ Secondary drowning	Cerebral anoxiaPneumoniaPulmonary edemaChemical pneumonitisMetabolic acidosis Electrolyteimbalance

Solution to Question 17:

Cadaveric spasm cannot be observed in traumatic asphyxia as the occurrence of traumatic asphyxia is sudden and hence does not give the patient or victim time to contract his muscles during the time of death.

Cadaveric spasm has been observed in the following situations:

• Drowning death

• Brain Injury

• Fear or Excitement

• Electrocution

Perte's syndrome, also known as traumatic asphyxia, is a clinical syndrome characterized by craniocervical cyanosis, subconjunctival haemorrhage, numerous petechiae, and accompanying neurological symptoms. This syndrome manifests due to the abrupt or intense compression of the thorax or upper abdomen, or both. Frequently observed injuries encompass pulmonary contusion, hemothorax, and pneumothorax. Effective management of the related injuries is crucial for optimal treatment outcomes. The expeditious initiation of treatment with a focus on restoring adequate oxygenation and perfusion has the potential to yield a favourable prognosis. The recoveries of the patients were shown to be correlated with the severity of their injuries as well as the presence of concomitant injuries.

Cadaveric spasm is an antemortem change that indicates the cause and manner of death. This phenomenon is alternatively referred to as instantaneous rigor or cataleptic rigidity

The victims do not undergo postmortem flaccidity (unlike rigor mortis), thereby directly entering a state of instantaneous rigor. This results in muscle stiffening which indicates the victim's attitude at the time of death. Cadaveric spasm is limited to select muscles i.e., the muscles that were in a state of contraction at the time of death. The mechanism of cadaveric spasm is thought to be due to neurogenic phenomena.

Predisposing factors for this postmortem change include excitement, fear, fatigue, electrocution, nervous system injury, and convulsant poisons such as strychnine. It is a unique phenomenon that cannot be replicated by any other situation, and it is not possible to induce it through any means following the cessation of life. A significant amount of force is necessary to overcome stiffness. The phenomenon seamlessly transitions into typical rigor mortis and ceases to exist after rigor mortis subsides.The process of protein coagulation can be observed under microscopic analysis in burns, but not in cases of cadaveric spasm.

Solution to Question 18:

Among the options provided, a seven-day admission in the yellow zone of the hospital is not an example of grievous hurt, but any hurt which causes the sufferer to be in severe body pain for 15 days is considered to be an example of grievous hurt.

Grievous hurt (Sec 116 BNS) is inflicted voluntarily on another person and comprises any of the eight kinds (clauses) enumerated below:

Clause 1 — Emasculation.

Clause 2 — Permanent privation of the sight of either eye.

Clause 3 — Permanent privation of the hearing of either ear.

Clause 4 — Privation of any member or joint.

Clause 5 — Destruction or permanent impairment of powers of any member or joint.

Clause 6 — Permanent disfigurement of head or face.

Clause 7 — Fracture or dislocation of bone or tooth.

Clause 8 — Any hurt that endangers life or causes the sufferer to be in severe bodily pain or unable to follow his or her ordinary pursuits for 15 days.

Punishment for grievous hurt :

BNS Section 117(2) - previously, IPC 325: Punishment for voluntarily causing grievous hurt –imprisonment up to seven years and fine.

BNS Section 118(2) - previously, IPC 326: Punishment for voluntarily causing grievous hurt by dangerous weapons or means—life imprisonment or imprisonment of a minimum of one year, extendable up to ten years, and fine.

Related IPC's:

BNS Section 115(1) - previously, IPC 321: Voluntarily causing hurt.

BNS Section 117(1) - previously, IPC 322: Voluntarily causing grievous hurt.

BNS Section 115(2) - previously, IPC 323: Punishment for voluntarily causing hurt: Imprisonment up to one year, or with a fine up to one thousand rupees or both.

BNS Section 118(1) - previously, IPC 324: Voluntarily causing hurt by dangerous weapons or means (3 years imprisonment) or with fine or both.

Solution to Question 19:

Among the options, nail is the least preferred in the identification of a dead body.

Nails have been used in the identification of dead bodies, however, their use has been deemed uncommon. Skeletal muscles are resistant to degradation and are therefore more likely to persist during decomposition, thus making them useful for the identification of corpses. It is also important to note that few studies show the amount of identical loci decreases with time.

Suitable samples for DNA analysis encompass several types of nucleated biological material, including but not limited to white blood cells, hair root bulb cells, and spermatozoa. In the past, a substantial quantity of samples was necessary for doing this analysis. However, the advent of the polymerase chain reaction (PCR), a technique capable of amplifying minute quantities of DNA, has enabled the identification of DNA from even the most limited samples.

When obtaining samples, whether from a deceased individual, a living victim, or the accused, the size of the sample that can be collected is typically not a significant concern. The collection of samples from various locations of staining inside the scene, the deceased, the surviving victim, or the accused can provide distinct challenges. These areas may consist of minute traces of blood, semen staining, and similar substances, necessitating meticulous collection procedures to mitigate the risk of contamination.

Blood and hair with intact follicles are utilized in both live and deceased individuals, while samples of the spleen can be obtained during post-mortem examinations. Additionally, buccal smears can be collected from living individuals. In situations where sexual assault is a potential occurrence, or in instances of suspicious deaths, it is advisable to collect vaginal, anal, and penile swabs. In the event of a potential delay in transporting samples to the laboratory, it is recommended that the samples be subjected to freezing conditions at a temperature of -20°C or as specifically directed by the laboratory.

Solution to Question 20:

Pralidoxime is used in organophosphate poisoning and not as a chelator in heavy metal poisoning.

In organophosphate poisoning, oximes like pralidoxime (2-PAM) regenerates active acetylcholinesterase restoring neuromuscular transmission and relieves skeletal muscle end-plate block. It is usually used as an adjuvant to atropine which act as competitive inhibitors to counter the nicotinic and muscarinic effects of organophosphates.

Pralidoxime should not be administered without concurrent atropine, in order to prevent worsening of symptoms due to transient oxime-induced acetylcholinesterase inhibition.

Pralidoxime attaches to the anionic site of the cholinesterase enzyme, which remains unoccupied in the presence of organophosphates. It binds and forms oxime-phosphonate which diffuses away, leaving the reactivated cholinesterase. Pralidoxime is not an effective antidote to carbamates like carbaryl and may potentiate its toxicity, as the anionic site required for attachment of oximes is occupied.

Other options :

Option A : British anti-Lewisite (BAL) also known as Dimercaprol is used as an antidote for gold, arsenic (except arsine), or acute mercury poisoning (except non-alkyl mercury). It is also an adjunct to edetate calcium disodium in acute lead poisoning.

Option B : Ethylenediaminetetraacetic acid (EDTA) is used for acute lead poisoning. It is not suitable for the management of chronic lead poisoning. The calcium salt of EDTA can be used in the management of heavy metal poisonings where the metal in question has a higher affinity for EDTA than calcium in vivo. Because of their lower affinity to EDTA, mercury and arsenic cannot be effectively chelated.

Option D : Penicillamine is used as a chelator in the treatment of mild cases of lead poisoning. D-penicillamine (Cuprimine) can also be used to treat scleroderma, Wilson's disease, and rheumatoid arthritis. Side effects include nephrotoxicity, hematologic derangements, and a characteristic rash that involves the neck and axillae—elastosis perforans serpiginosa. It is associated with membranous glomerulonephritis, drug-induced lupus, and myasthenia gravis.

Forensic Medicine INI-CET 2024

Question 1:

What is the Declaration of Oslo related to?

a) Human Torture

b) Therapeutic abortion

c) Human experimentation

d) Hunger strikes

Question 2:

A middle-aged housewife consumed rodenticide powder and presented with K+ levels of 2.5meq. On examination, irregular pulse with muscle paralysis is present. ECG shows ventricular ectopics. What is the suspected poison?

a) Aluminium phosphide

b) Zinc phosphide

c) Barium carbonate

d) Warafarin tablets

Question 3:

Which of the following antidotes are used to treat cyanide toxicity?

a) 1 and 2

b) 1, 2 and 3

c) 1, 3 and 4

d) 3 and 4

Question 4:

What is the correct order for opening cavities in case of a foetal autopsy?

a) 1, 3, 2

b) 3, 2, 1

c) 2, 1, 3

d) 1, 2, 3

Question 5:

Match the following:

a) 1-b, 2-c, 3-d, 4-e

b) 1-a, 2-b 3-d, 4-c

c) 1-d, 2-c, 3-a, 4-e

d) 1-c, 2-b, 3-d, 4-e

Question 6:

The following finding was found in an autopsy. Identity the weapon.

a) Bullet

b) Hammer

c) Iron rod

d) Scissors

Question 7:

Marshall's triad (punctate lacerations, bruises, and abrasions) is seen in which type of blast injury?

a) Primary blast injury

b) Secondary blast injury

c) Tertiary blast injury

d) Quaternary blast injury

Question 8:

What is not seen in heat exhaustion?

a) Loss of thermoregulatory mechanism

b) Altered sensorium

c) Orthostatic Hypotension

d) Core body temp below 105°F

Question 9:

Correctly matched antidote with toxicity

a) Organophosphorus- Physostigmine

b) Atropine - L-carnitine

c) Benzodiazepene-Flumazenil

d) Arsenic-Desferrioxamine

Question 10:

The compound given in the image below can be used in the treatment of which of the following poisonings?

a) Aluminium Phosphide

b) Mercury

c) Phophorus

d) Arsenic

Question 11:

A student consumed an unknown substance and came with shock and pulmonary edema. The silver nitrate breath test is positive. What is the antidote?

a) BAL

b) Pralidoxime

c) $MgSO_4$

d) Atropine

Question 12:

Cytochrome C oxidase present in ETC can be blocked by all except?

a) Carbon monoxide

b) Cyanide

c) Nitric Oxide

d) Hydrogen sulphide

Answer Key

Question No.	Correct Option
1	b
2	c
3	b
4	a
5	a
6	a
7	b
8	a
9	c
10	c
11	c
12	c

Detailed Explanations

Solution to Question 1:

In 1970, the World Medical Association adopted a resolution on therapeutic abortion known as the Declaration of Oslo.

Other options:

Option A: Declaration of Tokyo: related to human torture and other degradation

Option C: Declaration of Helsinki: concerned with human experimentation.

Options D: Declaration of Malta: related to hunger strikes.

Solution to Question 2:

The above clinical scenario with a presentation of hypokalemia, cardiac arrhythmia and paralysis after consumption of rodenticide poison is caused by Barium carbonate.

Barium carbonate is used in rat poised, a local corrosive irritant that, after absorption, mainly acts on muscles including the heart muscles.

The initial symptoms manifest as intense gastrointestinal distress, including nausea, vomiting, and diarrhoea. Subsequently, muscle cramps and stiffness, pupil dilation, tongue and larynx paralysis, and vertigo occur. Early impacts on the heart lead to elevated blood pressure, a slow and irregular pulse, and potential cardiac arrest in systole. Among the most notable changes are a loss of reflexes and paralysis.

Treatment involves the following steps:

• Gastric lavage to remove contents and administration of either sodium sulfate or magnesium sulfate solution to precipitate barium as insoluble sulfate.

• Intravenous administration of 10 ml of 10% sodium sulfate every 15 minutes to convert the soluble salt into insoluble sulfate.

• Purging with magnesium sulfate and repeated bowel lavage.

• Symptomatic management.

Other options:

Option A: Aluminium phosphide (ALP) is a solid fumigant pesticide, insecticide and rodenticide. ALP has a garlicky odour. Phosphine is a toxic agent. It acts by inhibiting the electron transport resulting from preferential inhibition of cytochrome oxidase.Severe toxicity produces ARDS, cardiac arrhythmias, congestive heart failure, pulmonary oedema, convulsions and coma. Cardiogenic shock is the most common cause of death, associated with higher mortality(35%-100%) but does not cause hypokalaemia.

Option B: Zinc phosphide poisoning can cause symptoms such as abdominal pain, nausea, vomiting, diarrhoea, and weakness. However, these symptoms are generally less severe compared to aluminium phosphide poisoning due to the highly toxic nature of phosphine gas released by aluminium phosphide.

Option D: Warfarin is an anticoagulant used in rat poison but is not directly associated with hypokalemia.

Solution to Question 3:

Amyl nitrite, sodium nitrite and sodium thiosulfate are antidotes used to treat cyanide toxicity.

Cyanide acts by inhibiting the activity of enzyme systems like cytochrome oxidase and carbonic anhydrase. It inhibits the final step of oxidative phosphorylation, preventing the synthesis of ATP and its usage as an energy source. Cyanide reduces the blood's oxygen-carrying capacity and combines with the ferric iron atom of intracellular cytochrome oxidase, inhibiting oxygen intake for cellular respiration.

Treatment primarily aims at reversing the cyanide-cytochrome combination by providing nitrites which convert haemoglobin to methaemoglobin.

Cyanides combine with methaemoglobin to produce non-toxic cyanmethaemoglobin, which, in the presence of rhodanase and sulphate donors such as thiosulphate, transforms cyanide to thiocyanate, which is eliminated in the urine.

The Cyanide kit contains:

• 21 amyl nitrite pearls

• Two 10 ml vials of 3% sodium nitrite

• Two 50 ml vials of sodium thiosulphate.

Using a handkerchief, a 0.2 ml ampoule of amyl nitrite is broken and held over the patient's nose for 15 to 30 seconds of every minute, until sodium nitrite infusion is started.

IV administration of 0.3 g of sodium nitrite in 10 ml of sterile water is given slowly over a 5-minute interval. Sodium nitrite forms methaemoglobin (Hb-Fe2+), then competes with cytochrome oxidase for the cyanide ion, thus protecting cytochrome oxidase. Folllowing this, 25 g of sodium thiosulphate in 50% solution is given i.v. over a period of 10 minutes. It converts cyanide to non-toxic thiocyanate, which is excreted in urine.

Other antidotes such as Hydroxocobalamin (Vit B12) and Dicobalt EDTA can also be used.

Solution to Question 4:

During the postmortem examination of a foetus, cranium is examined first intact along with the dura followed by the abdomen and thorax. The abdomen should be opened before the thorax to note the highest point of the diaphragm. Hence the answer is HEAD> ABDOMEN>THORAX.

During the removal of body cavities, preference is given depending upon the findings. Cavities that give trace evidence are opened first as compared to cavities that give maximum evidence to reduce the various artefacts.

Body cavities to be opened first:

• New Born: Abdomen> thorax

• Poisoning: Cranium

• Asphyxial deaths: Cranium > Thorax > Abdomen > Neck

The protocol for fetal autopsies differs slightly from that of adult autopsies. It includes examining the umbilicus, umbilical cord, placenta, scalp hairs, lanugo hairs, nails and their length, skin color and texture, scrotal sac wrinkling, as well as specific fetal measurements of the head, chest, and abdominal circumference.

Solution to Question 5:

The correct answer is option A: 1-b, 2-c, 3-d, 4-e

Paraphilias or perversions are sexual stimuli or acts that are deviations from normal sexual behaviors but are necessary for some people to experience arousal and orgasm.

1. Eonism (also known as transvestism) is a sexual perversion, where a person's whole personality is dominated by a desire to be identified as a member of the opposite sex. This is usually seen in males who obtain sexual pleasure by wearing female clothes. There is no hormonal/genital abnormality.

2. Algolagnia is sexual pleasure derived from inflicting pain (active algolagnia) or enduring pain (passive algolagnia).

Active algolagnia also called Sadism, is the sexual gratification obtained from acts of physical cruelty or infliction of pain upon one's partner. It is seen more commonly in men. To obtain sexual gratification the sadist may bite, beat, cut, burn, or otherwise mistreat or abuse his sexual partner.

3. Frotteurism refers to the sexual gratification achieved by touching or rubbing against a non-consenting person. This behavior often occurs in busy and crowded places. It is predominantly described in men.

4. Necrophilia is an obsession with obtaining sexual gratification from dead bodies.

Bestiality is an act of using a lower animal for sexual gratification. It is an unnatural sexual offence and is punishable in India.

1. Eonism	b. Sexual pleasure by wearing clothes of the opposite sex
2. Algolagnia	c. Sexual gratification in inflicting pain
3. Frotteurism	d. Sexual gratification by rubbing private parts against an other person
4. Necrophilia	e. Sexual gratification by intercourse with dead bodies

Solution to Question 6:

The given image shows a well-defined punched-out hole in the inner table of the skull. Hence, it could most probably be an entry wound produced by a bullet.

Entry wound: The initial contact of the missile punches a clean hole through the outer table of the skull. Where the bullet emerges internally, the inner table is then unsupported and a cone-shaped plug is detached, forming a crater that is appreciably larger than the external hole.

Bevelling: inner table

Punched out hole: outer table

Exit wound: If the bullet traverses the cranium and penetrates the opposite side, then the same pattern occurs, but this time with the small hole on the inner table and the crater on the outside.

Bevelling: outer table

Punched out hole: inner table



The sequence of bullet injuries is given by Puppe's rule.

Puppe's rule: It is possible to determine the sequence of injuries in skull fractures caused by blunt injuries to the head because the second fracture ends at the point where a break in cohesion is already present. The fracture lines created by every other shot extend only up to the fractured lines produced by the previous shot without continuing over them. Even though this rule is true for all blunt injuries to the skull, it is of particular importance in gunshot wounds.

Other options :

Option B: Hammer: When a hammer is used, the fracture is circular or in the shape of an arc of a circle. The diameter of the fracture is the same as the striking surface. There are generally no radiating linear fractures. A depressed fracture is caused by blows from a heavy weapon with a small striking surface. It is also called a signature fracture as the pattern of fracture resembles the weapon or object used.

The following is an image of a depressed fracture:



Option C: Iron Rod: An irregular missile such as an iron rod would produce several irregular lacerated wounds, and the exit wounds are larger than the entrance wounds.

The following is an image of the famous 3D model of Phineas Gage's skull, who survived a workplace injury of a tamping iron rod through his skull.

Solution to Question 7:

Marshall's triad in a bomb explosion is seen due to secondary blast injuries.

Marshall described the characteristic injuries associated with bomb explosions. It includes:

• Abrasions

• Bruises/ contusions

• Puncture lacerations

Blast injuries are classified as:

• Primary injuries- effect of transmitted blast waves on gas-containing structures. External injuries are absent and are characterized by internal injuries. The middle ear, lungs, and gastrointestinal tract are the most susceptible to this.

• Secondary injuries- injuries caused by shrapnel and other airborne debris propelled by the explosion. These injuries may affect any part of the body and sometimes result in penetrating trauma. Most casualties are caused by secondary injuries.

• Tertiary injuries- are caused by blast wind that can throw victims against solid objects.

• Quaternary injuries- all other injuries caused by explosions are not included in the above three categories. eg, flash burns, crush injuries, respiratory effects due to toxic dust, gas, etc.

Solution to Question 8:

Loss of the thermoregulatory mechanism is not seen in heat exhaustion. It is seen in heat stroke.

Heat exhaustion is caused by excessive exposure to heat, leading to nausea, vomiting, dizziness, altered sensorium(option B), thirst, fatigue, headache, sweating, and decreased urine output. Increases in sweating and dehydration can cause orthostatic hypotension(option C). Body temperature may rise but does not exceed 40.5°C (105°F). In heat exhaustion, the body maintains thermoregulation and no CNS dysfunction occurs. Prompt treatment is essential to prevent the progression of life-threatening heat stroke.

Solution to Question 9:

Benzodiazepine-Flumazenil is the correct match of toxicity with its antidote.

Flumazenil is a competitive antagonist at the benzodiazepine binding site of the GABA A receptor. It antagonizes both the agonist (benzodiazepines) and inverse-agonist (beta-carbolines) of this receptor.

The duration of clinical effects usually is only 30–60 min. It is absorbed rapidly after oral administration, less than a quarter of the drug reaches the systemic circulation. This is due to the extensive first-pass hepatic metabolism. Flumazenil is eliminated almost entirely by hepatic metabolism to inactive products with a t1/2 of about 1 hour.

It is primarily indicated for the management of benzodiazepine overdose and the reversal of benzodiazepine anaesthesia. Flumazenil is only available for intravenous administration. It reliably reverses the sedation, respiratory depression, and amnesia caused by benzodiazepines.

Other options:

Option A: In organophosphate poisoning (insecticide) with both muscarinic and nicotinic manifestations, antidotes administered are atropine and pralidoxime. Atropine is effective against muscarinic receptors. Pralidoxime (2-PAM) is a cholinesterase-reactivating agent. Hence, both drugs must be co-administered to treat both muscarinic and nicotinic symptoms.

Option B: Treatment of atropine toxicity includes administration of a short-acting barbiturate or diazepam to control convulsions. Physostigmine is used as an antidote to atropine poisoning to treat delirium and coma. Other measures for symptomatic management are respiratory support and control of hyperthermia.

Option D: ■■In Arsenic toxicity, the antidote preferred is freshly prepared hydrated ferric oxide, which converts toxic arsenic to non-toxic ferric arsenite. DMSA (Succimer), British anti-Lewisite (BAL), and penicillamine can also be used.

Solution to Question 10:

The compound given in the image is copper sulphate. It can be used in the treatment of phosphorus poisoning.

The initial stage of acute phosphorus poisoning through ingestion involves severe abdominal discomfort, vomiting, and diarrhoea. Excreta and breath have a garlicky odour. Luminescent faeces and vomit are characteristically seen (smoky stool syndrome). Direct contact with the skin

can result in second and third-degree burns. Shock, hematemesis due to liver failure, and renal insufficiency are potential causes of death.

Regular inhalation of phosphorus fumes, leading to chronic phosphorus poisoning, results in necrosis of the lower jaw around a decayed tooth. This condition is known as phossy jaw and is associated with the discharge of foul-smelling pus from multiple sinuses.

Gastric lavage with potassium permanganate oxidizes phosphorus into phosphoric acid and phosphates. Activated charcoal can be used to adsorb the poison.

Stomach wash with 0.2% copper sulphate solution or 0.2 g of copper sulphate every 5 minutes until vomiting occurs. It coats the particles of phosphorus with a film of copper phosphide, which is relatively harmless. As copper has caustic properties and can cause acute poisoning, adequate care should be taken. Copper sulphate was also used for topical application in phosphorus poisoning. It is not recommended now.

Other options:

Option A: Aluminium Phosphide has no antidote. Magnesium sulphate is given in ALP poisoning to correct hypomagnesemia and arrhythmias and reduce organ toxicity.

Option B and D: British anti-Lewisite (BAL), also known as Dimercaprol, is used as an antidote for Mercury poisoning (except non-alkyl mercury) and Arsenic poisoning.

Solution to Question 11:

The patient presents with clinical features suggestive of Aluminium phosphide (ALP) poisoning. Magnesium sulphate is given in ALP poisoning to correct hypomagnesemia, arrhythmias and reduce organ toxicity.

Aluminium phosphide (ALP) is a solid fumigant pesticide, insecticide and rodenticide. ALP has a garlicky odour. Phosphine is a toxic agent. It acts by inhibiting the electron transport resulting from preferential inhibition of cytochrome oxidase. Aluminium phosphite and Aluminium hypophosphite are non-toxic agents. Phosphine is a systemic poison and affects all organs of the body. The presence of HCl in the stomach accelerates this reaction.

Signs and Symptoms include irritation of mucous membranes and acute respiratory distress in mild toxicity. Moderate toxicity produces ataxia, numbness, tremors, muscular incoordination and paralysis. Severe toxicity produces ARDS, cardiac arrhythmias, congestive heart failure, pulmonary oedema, convulsions and coma. Cardiogenic shock is the most common cause of death and is associated with higher mortality(35%- 100%)

There is no specific antidote. Gastric lavage should not be done with warm water as on coming in contact with water, aluminium phosphide releases phosphine and increases the concentration of toxic agent phosphine, so it is done with potassium permanganate and activated charcoal. Magnesium sulphate is used to correct hypomagnesemia, arrhythmias and reduces organ toxicity. Supportive measures to treat acidosis, shock and hemodialysis can be done.

On post-mortem examination, a garlic-like odour is present at the mouth, nostrils and in the gastric contents. Silver nitrate paper test is done to detect phosphine which turns the paper black.

Silver nitrate paper test: This test is used to detect aluminium phosphide poisoning.

• Aluminium phosphide (ALP)is used as a pesticide, insecticide, and rodenticide.

• When ALP comes in contact with moisture, it releases an active component called phosphine.

• When the silver nitrate-impregnated filter paper, comes in contact with phosphine the paper turns black in colour.

• The test is done in two methods:

a. Gastric aspirate: 5 mL of gastric aspirate and 15 mL of water are put in a flask and the mouth of the flask is covered with silver nitrate-impregnated filter paper. The flask is heated at 50°C for 15 to 20 minutes. If the filter paper turns black, it indicates the presence of phosphine.

b. Breath: The patient is asked to breathe in and out through a face mask which is impregnated with silver nitrate. If the face mask turns black, it indicates the presence of phosphine. The breath test is positive only if the patient ingests more than 6 grams of aluminium phosphide.

Solution to Question 12:

Nitric Oxide cannot block cytochrome C oxidase present in ETC.

Nitric Oxide: While NO can modulate ETC activity and inhibit respiration temporarily, it does not universally block Complex IV like the other inhibitors. (Option C)

Cytochrome c oxidase (Complex IV) in the electron transport chain (ETC) is the terminal enzyme that facilitates the transfer of electrons to molecular oxygen, reducing it to water. Various inhibitors can block this process, leading to impaired oxidative phosphorylation and ATP synthesis:

Carbon monoxide: CO binds strongly to the heme iron of cytochrome c oxidase in its reduced (ferrous, Fe^{2}■) state. This prevents oxygen from binding and accepting electrons, halting the production of water and the proton gradient. Poisoning with CO can lead to severe hypoxia at the cellular level, even in the presence of normal oxygen levels. (Option A)

Cyanide: Cyanide binds irreversibly to the ferric (Fe^{3}■) iron in the heme group of cytochrome c oxidase. It completely blocks electron transfer to oxygen, making it a potent inhibitor. Cyanide poisoning results in rapid cellular asphyxiation and is life-threatening. (Option B)

Hydrogen Sulfide: Inhibits Complex IV at high concentrations by binding to its heme groups, similar to cyanide. Its inhibition is dose-dependent, but it remains a strong inhibitor at high levels. (Option D)

Forensic Medicine NEET 2024

Question 1:

During the post mortem examination following finding is seen. What is this indicative of _____

a) Choking

b) Throttling

c) Hanging

d) Smothering

Question 2:

An 18-year-old boy was brought by police, and accused of sexual assault, but the boy claims to be a minor (16 years old). Which radiographs will you order to solve this discrepancy?

a) Elbow and hip joint

b) Elbow and ankle joint

c) Shoulder and skull joint

d) Wrist and knee joint

Question 3:

A man sexually aroused when he wears his female partner's dress and heels. He denies getting attracted to male. What is this condition called?

a) Gay sexual preference

b) Transvestic fetishism

c) Gender dysphoria

d) Voyeurism

Question 4:

A woman gave birth to a child 8 months after the death of her husband. The in-laws claim that the child is not their son, however, the DNA testing confirms the paternity. The child is known as _____

a) Illegitimate child

b) Posthumous child

c) Surrogate child

d) Supposititious child

Question 5:

The railway police has bought a person with suspected poisoning. On examination, he was confused with slurred speech, hot skin, dry mouth, dilated pupils. What is the probable poisoning ?

a) Cocaine

b) Datura

c) Cannabis

d) Alcohol

Question 6:

A police officer brought a child to the hospital. There is redness and tenderness around perianal region. A rectal swab is taken and the image given below is the finding. What test is it indicative of?

©MARROW

a) Barberio test

b) Florence test

c) Teichmann test

d) Takayama test

Question 7:

Rigor mortis is well established in 12-24 hours after death. What is the order of appearance of rigor mortis?

a) Toe to head

b) Head to toe

c) Centre to periphery

d) All the body parts simultaneously

Question 8:

A 5-year-old child has consumed overdose of iron syrup, around 60mg/kg. Which among the following is contraindicated?

a) BAL

b) Deferoxamine

c) Calcium EDETATE

d) DTPA

Question 9:

A sexual assault survivor had to testify in court and an in-camera trial was conducted. What is an "in-camera" trial?

a) Open court proceedings

b) Closed court proceedings

c) Proceedings recorded on camera

d) Proceeding held in a different location

Question 10:

A 36-year-old man died following a RTA. The doctor informed the police officer and shifted the body to the mortuary. Autopsy in this case will be conducted at the request of _____

a) Forensic expert

b) Investigating police officer

c) Defense lawyer

d) Prosecution lawyer

Question 11:

A person comes to emergency after consumption of a substance, the form and dose of which is not known to the relatives. The patient has tachypnea and hypotension. On metabolic assessment, high anion gap metabolic acidosis is noted with hypocalcemia. What is the diagnosis?

a) Methyl alcohol

b) Ethylene glycol

c) Datura

d) Digoxin

Question 12:

A patient presented to the OPD with findings as shown below. Identify the type of poisoning in this patient.



1 2

a) Chronic malnutrition

b) Copper

c) Mercury

d) Arsenic

Question 13:

The following findings were noted during the autopsy of a patient with a gunshot injury. What is the likely range of the gunshot?



a) Cannot be opined

b) Distant range

c) Contact range

d) Close range

Question 14:

A large number of people who consumed cheap alcohol presented to the emergency department with abdominal pain, confusion, and decreased vision. Methyl alcohol level in blood was detected to be around 20 mg %. On assessment which of the following metabolites will be found in blood?

a) Glycolic acid and Glyoxylic acid

b) Formic acid and lactic acid

c) Glycolic acid and lactic acid

d) Oxalic acid and formic acid

Question 15:

In the case of drowning, following autopsy, an organism was isolated from the water present in the internal organs. Which of the following is true regarding the isolated organism?



a) Present only in the bone marrow

b) Dried outer wall contains silica and chlorophyll

c) It is a microscopic unicellular bacteria

d) It is not heat and acid resistant

Question 16:

A pregnant female comes to a gynaecologist, who sends her for USG. On USG, twin pregnancy was noted with about a one-month difference in the age of fetuses. Which of the following is true with regard to this patient?

a) Superfecundation

b) Superfetation

c) Supposititious child

d) Posthumous child

Question 17:

A 32-year-old woman G3P2L2 came to you in active labor. The labour process was arrested at around cervical dilatation of 8 cm. She was counselled regarding the need for an emergency cesarean section. The baby was delivered by cesarean section and was doing well but the mother developed intractable postpartum hemorrhage. Conservative measures were tried initially but failed. Emergency hysterectomy was done as a life-saving procedure. What are the medicolegal implications of this situation?

a) Patient could sue the doctor for removing her uterus

b) Hysterectomy was done as a life saving procedure, so the doctor is legally protected

c) Consent can be taken from the patient on OT table

d) Counsellling of patient and attendants is not needed

Question 18:

A 30-year-old man who was found dead was brought in for autopsy. The doctor notices a greyish-white waxy layer over the body with preserved facial features. Which of the following statements is true concerning adipocere formation?

a) It occurs in hot and dry temperatures

b) It occurs in cold temperatures

c) It is a type of body preservation with saponification of fats

d) Occurs within few hours of death

Answer Key

Question No.	Correct Option
1	d
2	d
3	b
4	b
5	b
6	a
7	b
8	a
9	b
10	b
11	b
12	d
13	d
14	b
15	b
16	b
17	b
18	c

Detailed Explanations

Solution to Question 1:

Abrasions around the mouth, cheeks, lips or lesions within the lips may reliably indicate smothering.

The non-specific signs of death due to asphyxia:

• Petechial haemorrhages in the face, eyebrows, eyelids, forehead, ears, conjunctiva, sclera - When they occur in the visceral pleura, they are called Tardieu's spots

• Emphysematous, congested lungs with serosanguinous fluid exudates in alveoli.

• Cyanosis or dark-coloured blood due to accumulation of CO_2.

• Congestion of the larynx with slight oozing

Pressure marks on the face can occur as a part of normal postmortem changes. Circumoral and circumnasal pallor can be caused merely by passive pressure of the dependent head after death.

Note: The coexistence of these findings, in themselves, does not prove that death resulted from mechanical asphyxia. All these phenomena are non-specific and are in no way peculiar to this mode of death. They are of no value in proving that death resulted from mechanical asphyxia.

Solution to Question 2:

Wrist and knee joint radiographs will be ordered to solve this discrepancy as lower end of radius and upper end of tibia fuse at 18-19 years. If these joints are fused, It is certain that the boy is more than 18 years and his claim is false.

Bone/ joint	Max age of fusion
Wrist joint centers (lower end of ulna and radius)	19 years
Knee joint centres(lower end of femur, upper end of tibia and upper end of fibula)	20 years
Shoulder joint (head of humerus, greater tuberosity and lesser tuberosity)	19 years
Hip joint (head of femur, greater trochanter)	17 years
Elbow joint centers (lower end of humerus, upper end of radius, olecranon to ulna)	17years
Ankle joint	17 years

Solution to Question 3:

The given scenario is suggestive of transvestic fetishism.

It is a paraphilia characterized by recurrent sexually arousing fantasies, sexual urges, or behavior involving the wearing of clothes of the opposite sex and appearing as a member of the opposite sex. It is also called transvestic fetishism. It is usually seen in males who obtain sexual pleasure by wearing female clothes.

Fetishistic transvestism is a paraphilia whereas transvestism is a gender identity disorder. There is no hormonal/genital abnormality in fetishistic transvestism.

Option A : Gay sexual preference is due to attraction to people of one's own gender.

Option C: Gender dysphoria is a term that describes a sense of unease that a person has due to a mismatch between their biological sex and their gender identity. This sense of unease or dissatisfaction can lead to depression and anxiety and can have a harmful impact on daily life.

Option D : Voyeurism/Peeping Tom/Scotophilia is sexual gratification by secretly watching others getting undressed, taking a bath, performing sexual intercourse, etc.

Solution to Question 4:

The child delivered after the death of the biological father, or a child delivered after the death of the biological mother, usually when delivered by a cesarean section is a posthumous child.

Legal issues involved with a posthumous child are legitimacy, inheritance of property, and the case of compensation for slander against the mother.

Other options :

Option A: Illegitimate child is a child born out of wedlock i.e. when the parents are unmarried.

Option C: A surrogate child is born out of a pregnancy in which a woman gives birth to a baby for a person who is not able to have children.

Option D: Supposititious child or fictitious child is when a woman pretends to go through pregnancy and delivery. Later, she produces a living child on her own to obtain money or claim the property.

Solution to Question 5:

The given clinical scenario is suggestive of Datura poisoning.

The two species are jimson weed (Datura stramonium) and angel's trumpet (Datura candida).

Seeds or other parts of the plant can be ingested or smoked. Poisoning leads to delirium, hallucinations, and seizures. Additionally, other classic anticholinergic effects, such as mydriasis, tachycardia, dry mouth and skin, blurred vision, urinary retention, and hyperthermia may present.

Gastric emptying is often delayed, and the small, plentiful seeds can become trapped among the gastrointestinal folds after ingestion. Thus gastric decontamination can be an important therapy.

Physostigmine, a reversible acetylcholine esterase antagonist, is an effective treatment for severe anticholinergic poisoning.

Solution to Question 6:

The test done in this scenario is the Barberio test.

Barberio test is used to detect seminal fluid. When a saturated aqueous or alcoholic picric acid solution is added to the spermatic fluid, yellow needle-shaped rhombic crystals of spermine picrate are produced.

Other options:

Option B: The Florence test detects semen. If semen is present, dark brown rhombic or needle-shaped crystals of choline iodide appear immediately.

Option C: The Teichmann test (hemin crystal test) is used to detect bloodstains. If bloodstain is present, faint yellowish-red to brownish-black rhombic crystals of hemin or hematin chloride appear. When hydrogen peroxide is added, the hematin crystals give bubbles of gas.

Option D: Hemochromogen crystals are seen in the Takayama test. It is a microchemical test used to detect bloodstains. The stain used is Takayama reagent (Sodium hydroxide, pyridine, glucose). If bloodstain is present, pink feathery crystals of hemochromogen or reduced alkaline hematin appear in a few minutes. The test is negative if the crystals are not formed in 30 minutes. This test is more reliable and gives positive results with old stains also but not a definitive test to detect blood stains.



Solution to Question 7:

The order of appearance of rigor mortis is from head to toe.

Rigor mortis (cadaveric rigidity or death stiffening) refers to the stiffening and shortening of muscles following death. The basic sequence of the rigor mortis begins from the head and continues downwards the body. It disappears in the same order of appearance. Nysten's rule describes the sequential onset of rigor mortis in the various muscle groups.

The muscles are affected in the given order:

• Involuntary muscles - Heart (left chamber of the heart is affected first)

• Eyelids

• Neck and lower jaw

• Face

• Chest

• Upper extremities

- Abdomen
- Legs
- Fingers and toes

Solution to Question 8:

The clinical scenario points to iron toxicity, which is managed with deferoxamine (also known as desferrioxamine), calcium EDETATE and DTPA whereas Dimercaprol, also known as BAL (British anti-Lewisite), is used to treat arsenic, mercury poisoning, and Wilson's disease.

Large amounts of ferrous salts can be fatal in children. Initial symptoms include abdominal pain, diarrhea, vomiting, and gastrointestinal bleeding. In severe cases, cyanosis and hyperventilation due to acidosis can develop. This can sometimes be followed by an asymptomatic phase. Later, hypoglycemia and hepatic and renal failure can develop. Pyloric stenosis and gastric scarring can develop due to its corrosive nature. Ultimately, if left untreated, the child can die.

The gastric contents can be used for an iron color test to estimate the iron concentration in plasma. As iron tablets are radiopaque, an X-ray can be used to assess the number of pills taken. Deferoxamine (Option B) is selectively used as a chelating agent to manage cases of acute iron poisoning, hemochromatosis, and patients with iron overload from repetitive blood transfusions. Desferrioxamine can be administered intramuscularly, intravenously, or subcutaneously but not orally. Side effects include pruritus, wheals, rash, and anaphylaxis.

Deferiprone and deferasirox are other oral iron chelators used to treat iron overload in thalassemia patients.

Other options:

Option C : Ethylenediaminetetraacetic acid (EDTA) is used for acute lead poisoning. It is not suitable for the management of chronic lead poisoning. The calcium salt of EDTA can be used in the management of heavy metal poisonings where the metal in question has a higher affinity for EDTA than calcium in vivo. Because of their lower affinity to EDTA, mercury and arsenic cannot be effectively chelated.

Option D : Diethylenetriamene pentaacetate (DTPA) is a chelating agents which works by binding and holding on to radioactive materials (plutonium, americium, and curium) and poisons like iron. Calcium DTPA is well tolerated and effective as desferrioxamine in mobilising iron than Zinc DTPA.

Solution to Question 9:

'In-camera trial' is a closed court proceeding in which the case is heard in private i.e. in court where general public is excluded.

BNSS sections related to court proceeding are tabulated below:

BNSS	CrPC	Deals with
63-71	61-69	Summons
181	162	Oral statements made to the police and recorded by the police should not be signed
195	175	Power to summon persons by a police officer
196	176	Magistrate inquest
350	312	Expenses of complaints and witnesses
366366(2)	327	Criminal court is generally deemed to be an open court.In-camera trial for rape
389	350	Summary procedure for punishment for non-attendance by a witness in obedience to the summons
395	357	Order to pay compensation

Solution to Question 10:

In the case of the 36-year-old man who died following an RTA, the autopsy will be conducted at the request of the investigating police officer.

According to BNSS Section 194 (Section 174 CrPC), a police inquest is the standard procedure for investigating unnatural deaths in India. This process is typically carried out by a police officer, not below the rank of Senior Head Constable, who is responsible for conducting a preliminary inquiry into the circumstances surrounding the death.

During the police inquest, the investigating officer visits the scene of the incident, examines the body, and documents any visible injuries or signs of trauma. The officer's report will detail these findings and assess whether the death appears to be suicidal, homicidal, accidental, or suspicious. If the situation warrants, the body will be sent for a postmortem examination to be conducted by a qualified medical professional to determine the exact cause of death.

In certain cases, a magistrate inquest [BNSS 196 (Section 176 CrPC)] may be conducted instead of, or in addition to, a police inquest. This is particularly true in cases involving deaths under special circumstances, such as lock-up deaths, deaths while in police custody, custodial rape, death in a psychiatric hospital, dowry death or deaths resulting from police actions. The magistrate inquest is carried out by a District Magistrate, Subdivisional Magistrate, or another authorized magistrate, and involves a more formal judicial inquiry into the cause of death.

Solution to Question 11:

The patient's presentation of tachypnea, hypotension, high anion gap metabolic acidosis, and hypocalcemia, is suggestive of ethylene glycol poisoning.

Ethylene glycol is a clear, colourless, and odourless liquid commonly used as an antifreeze. It is known for its bitter-sweet taste and is absorbed mainly through ingestion rather than skin contact. Upon ingestion, ethylene glycol is metabolized in the liver to several toxic metabolites, including glycolic acid and oxalic acid, which contribute to its harmful effects.

Initially, ethylene glycol poisoning can cause symptoms such as vomiting, lethargy, and ataxia, progressing to severe effects including tachycardia, tachypnea, and circulatory collapse within 12-24 hours. As the condition progresses, typically within one to three days, patients may experience hypocalcemia, oliguria, tubular necrosis, and acute kidney injury (AKI). A notable feature of ethylene glycol poisoning is the formation of calcium oxalate crystals in the urine, which can lead to renal failure.

The postmortem findings often include cerebral edema, chemical meningoencephalitis, and significant damage to the liver and kidneys. Oxalate crystals may be observed in the brain, spinal cord, and kidneys, further confirming the diagnosis.

For treatment, immediate measures include gastric lavage, activated charcoal, and administration of ethanol or 4-methyl pyrazole (fomepizole), which inhibit the enzyme alcohol dehydrogenase to prevent further metabolism of ethylene glycol into toxic compounds. Additional treatments such as intravenous sodium bicarbonate and calcium gluconate are used to address metabolic acidosis and hypocalcemia, respectively, while hemodialysis is indicated in severe cases with significant renal impairment or high serum ethylene glycol concentrations.

Other options:

Option A: Methyl alcohol (Methanol) and Ethylene glycol poisoning can present very similarly. However, the key distinguishing feature of methanol poisoning is the presence of eye symptoms (as opposed to kidney damage in ethylene glycol) - like blurring of vision/loss of vision due to retinal damage by formate metabolites.

Methanol--Alcohol Dehydrogenase (ADH) ⇒ Formaldehyde ⇒ (Retinal damage)

Ethylene glycol--Alcohol Dehydrogenase (ADH) ⇒ Glyoxyaldehyde ⇒ (Kidney damage)

Option C: Datura is used as a stupefying poison before robbery, kidnapping, and rape. The active ingredients include the anticholinergic alkaloids atropine, scopolamine, and hyoscyamine. The seeds or other parts of the plant can be ingested or smoked. Toxic manifestations include dilatation of pupils along with loss of accommodation, diplopia, dryness of the mouth and throat, unquenchable thirst, dysphagia, dilatation of cutaneous blood vessels, dysarthria, drunken gait, delirium, drowsiness and death due to respiratory or cardiac failure.

Option D: Digoxin toxicity can occur in the elderly and patients with renal impairment and presents with nausea, vomiting, diarrhoea, confusion, blurred vision with a characteristic yellowing of vision (compared to the “starry night” painting), and irregular heart rhythms. Overdose and resultant arrhythmias are managed with anti-digoxin Fab fragments (Digibind or Digifab).

Solution to Question 12:

The image showing the raindrop pattern of skin pigmentation and the Aldrich-Mees lines on the fingernails of the patient are suggestive of chronic arsenic poisoning.

Arsenic in its metallic form is not poisonous, but when it unites with oxygen, it forms poisonous vapours of arsenic trioxide. These interfere with cellular respiration, damaging mitochondrial enzymes. It is present in fruit sprays, copper and gold mineral deposits, weed killers, insecticides, rat poisons, wallpapers, and artificial flowers.

There are several types of arsenic poisoning:

• Fulminant - When massive doses are rapidly absorbed, death occurs in 1–3 hours due to shock and peripheral vascular failure.

• Gastrointestinal - This is the most common form of acute poisoning and resembles bacterial food poisoning. There is a constriction in the throat and difficulty in swallowing. The stools are dark-colored, foul smelling, and bloody initially and later take a colorless odorless watery form similar to the rice-water stools of cholera.

• Chronic poisoning - It occurs due to repeated daily exposure. The main features include finely mottled, raindrop pigmentation that appears on the skin of flexures, temples, eyelids, and neck. In prolonged contact, hyperkeratosis of the palms and soles occurs along with the development of a transverse white line in the fingernails called the Aldrich-Mees lines.

Tests to detect arsenic poisoning:

• 24-hour urinary excretion of arsenic

• Detection of arsenic by atomic absorption spectroscopy

• Marsh and Reinsch's tests are now considered obsolete.

The preferred treatment option is freshly prepared hydrated ferric oxide, which converts toxic arsenic to non-toxic ferric arsenite. DMSA (Succimer), British anti-Lewisite (BAL), and penicillamine can also be used.

Other options:

Option A: Chronic malnutrition leads to distinct skin and nail changes depending on the specific condition. In kwashiorkor, there is flaky paint dermatosis with reddish-brown scaling, thin and sparse hair featuring a "flag sign," and soft nails. In marasmus, patients exhibit dry, wrinkled skin, slow-growing brittle hair, and nails, often accompanied by growth retardation.

Option B: Acute copper poisoning is due to inhalation of a large amount of copper dust or fumes, which leads to upper respiratory tract irritation, conjunctivitis, and sinus irritation. Contact with the skin may lead to irritant contact dermatitis and a greenish-blue discolouration of the skin.

In chronic poisoning, symptoms mostly resemble those of Wilson's disease, mainly including gradual anaemia, peripheral neuritis, and muscle atrophy.

Option C: Acute mercury poisoning presents with corrosive bronchitis with fever, chills, and dyspnea, which may progress to pulmonary edema and fibrosis. The symptom profile in children is similar to that of Kawasaki disease (mucocutaneous lymph node syndrome). Chronic exposure to elemental mercury leads to a classic triad of gingivitis, tremors, and neuropsychiatric changes.

Solution to Question 13:

The likely range of the gunshot injury shown above is close range, characterized by a circular wound with blackening and tattooing.

To determine the range of firing, the following parameters are used:

• The entry wound size and shape

• Blast effect – Muzzle end imprint and edge eversion

• Blackening, Tattooing, Singeing of hair, Abrasion collar, and Grease collar

• Hair should be preserved to determine the range of firing which can be examined for carbonization/desiccation in close range, also from lead deposition around the wound to assess the range of firing.

Blackening occurs due to the deposit of smoke and can be removed with a wet cloth. This can be seen up to around 15 cm of firing. Whereas, tattooing occurs due to the deposition of unburnt powder particles and are difficult to remove. They are also called stippling or peppering. This is seen up to 50 cm of firing.

Abrasion collar: Gyroscopic movement of the bullet, the skin's elasticity, and the grease and dirt particles on the bullet's surface are the major causative factors of abrasion and grease collar. They are seen on the skin around the wound of entry at all ranges. Depending on the angle of entry made by the bullet, i.e.

• A perpendicular entry of the bullet develops the abrasion collar of a rounded shape.

• An acute angular entry of the bullet develops the abrasion collar of an oblique or oval shape.

• Some contusion present in the abraded collar is known as contusion collar

Grease collar: Mostly seen on the clothing and is seen as a black deposit, sharply outlined as if printed (better seen on light colour clothes and fair-skinned individuals). Its medicolegal importance is that it helps derive information such as entry wound and direction of firing.

Solution to Question 14:

The given case scenario is suggestive of methanol poisoning. In these cases, formic acid and lactic acid are typically the metabolites found in the blood.

Methyl alcohol, or methanol, is commonly present in cheap or illicit alcoholic beverages due to its lower cost compared to ethyl alcohol. These inexpensive alcohols may be improperly distilled or adulterated, leading to methanol contamination. When ingested, methanol is absorbed into the bloodstream and metabolized by the liver, where it is first converted into formaldehyde, a highly toxic intermediate. Formaldehyde is then further oxidized into formic acid, which is responsible for many of the severe symptoms associated with methanol poisoning.

The accumulation of formic acid leads to significant metabolic acidosis, which manifests as confusion, abdominal pain, and visual disturbances. The visual symptoms can be particularly severe, including blurred vision or even blindness, due to the toxic effects of formic acid on the optic nerve. In addition to formic acid, methanol's disruption of normal oxidative enzyme activity results in the buildup of lactic acid, contributing further to acidosis. Brain edema, acute renal failure, and severe metabolic derangement are also seen.

The principles of treatment for this condition predominantly involve using fomepizole or ethanol to inhibit the metabolic conversion of these relatively non-toxic alcohols to their toxic metabolites.

Solution to Question 15:

Diatoms enter pulmonary circulation on antemortem drowning. Diatoms are a class of microscopic unicellular algae (option C) found suspended in most natural waters. Their cell walls are impregnated with silica, chlorophyll, diatomic, and a brown pigment.

The drowning fluid and the particles in it, e.g., diatoms and plankton, pass from the ruptured alveolar wall into lymph channels and pulmonary veins and thus enter the left heart. Only a live body with circulation could transport diatoms from the lungs to the brain, bone marrow, liver and other viscera, and skeletal muscle. They are also found in bile and urine. Bone marrow is highly suitable and reliable.

Isolation of diatoms involves acid digestion of tissue - bone marrow, lung, blood, and kidneys with subsequent centrifugation and the final residue is aspirated and examined for diatoms, which are birefringent, using dark ground illumination. A water sample is collected at the time of body retrieval in a clean container and the similarity of different species of diatom is compared. Diatoms are both heat and acid-resistant (option D).

• The presence of diatoms in the lung, brain, liver, kidneys, and bone marrow (option A) of the femur (best site for analysis) or humerus is suggestive proof of antemortem drowning.

• Since diatoms resist putrefaction, the diatom test may have some value in the examination of decomposed bodies.

• The test is negative in postmortem submersion and in dry drowning.

Diatoms are ubiquitous in the environment and may enter circulation via the gastrointestinal tract (as contaminants of foods)or via the respiratory tract (which are present in the air). Still, the diatom test is considered the gold standard for the diagnosis of typical drowning.

Solution to Question 16:

The above phenomenon is known as superfetation. It occurs when there is fertilization of 2 different ova released in different menstrual cycles. The consequence is two children are born at the same time and one of them might be mature while another might be immature or two fully developed fetuses are born at intervals ranging from one to three months apart.

Ovulation usually stops with the onset of pregnancy, but in a few individuals, ovulation may be possible till the first trimester, after which the decidual cavity is obliterated. The fertilization of the newly released ovum in a woman who is already pregnant may take place following coitus.

Other options:

Option A: Superfecundation is the fertilization of 2 different ova in the same menstrual cycle by separate acts of coitus. Heteropaternal superfecundation is used to refer to instances of two different males fathering fraternal twins. This leads to the possibility of twins also being half-siblings. This may raise doubts regarding adultery and infidelity.

Option C: A supposititious child is also known as a fictitious child. In such cases, a woman pretends to be pregnant and substitutes a living child as her own, or may substitute a male child for her female child, or abortion. This is done to claim property. This can be identified by verifying the signs of pregnancy and delivery in the woman and by DNA fingerprinting.

Option D: A posthumous child is a child born alive after the death of his or her father who was conceived before the death of the father.

Solution to Question 17:

According to BNS Section 30 (IPC 92), a doctor is permitted to perform life-saving procedures on a patient with life-threatening injuries. For example, in this case where conservative measures have failed to control intractable postpartum haemorrhage, an emergency hysterectomy can be performed without obtaining consent in order to save the patient's life.

The guiding principle is whether the delay in treatment in order to obtain consent would result in harm to the patient. This is the doctrine of anticipation. Under some circumstances like emergencies, informed consent may be neither possible nor necessary.

BNS Section 30 (IPC 92) states that nothing is an offence by reason of any harm which it may cause to a person for whose benefit it is done in good faith, even without that person's consent, if the circumstances are such that it is impossible for that person to signify consent, or if that person is incapable of giving consent, and has no guardian or other person in lawful charge of him from whom it is possible to obtain consent in time for the thing to be done with benefit.

Solution to Question 18:

The given scenario indicates adipocere formation, which is a late postmortem change where body fats undergo saponification through anaerobic hydrolysis and hydrogenation, resulting in a waxy substance. Therefore, the correct answer is that it is a type of body preservation with saponification of fats.

Adipocere formation or corpse wax, is a late sign of death, that occurs days to months after death. A wax-like organic substance is formed by the anaerobic hydrolysis or hydrogenation of fat in tissues. It has a greyish-white appearance with a distinctive offensive or sweetish smell.

Adipocere formation varies widely, taking as little as three weeks in temperate regions during summer, but typically requires several months for significant changes. In rare cases, the entire body may be affected, with complete conversion of an adult limb taking three to six months.

Medicolegal importance is as follows:

• It is the surest sign of death

• It gives a rough estimate of the time since death

• The cause of death can be identified.

• In the face, the features are well preserved and therefore used in identification.

Postmortem changes are classified based on time:

• Immediate signs of death (at the time of somatic or clinical death):

• Insensibility and loss of voluntary power

• Cessation of respiration

• Cessation of circulation

• Early signs of death (cellular death that is observed in the first 12 to 24 hours of death):

• Pallor and loss of elasticity of the skin

• Eye changes

• Primary flaccidity of muscles

• Algor mortis/Postmortem cooling

• Postmortem lividity/Suggillation

• Rigor mortis

• Late signs of death:

• Putrefaction

• Adipocere

• Mummification

Ophthalmology AIIMS 2017

Question 1:

A 62-year-old presented with diminished vision 1 year after cataract surgery. The findings are shown below. What is the most likely diagnosis?



a) After cataract

b) Irvine-Gass syndrome

c) End ophthalmitis

d) Uveitis-glaucoma-hyphema syndrome

Question 2:

Defect at the level of the optic chiasma will result in which of the following visual field defects?

a) Central scotoma

b) Bilateral hemianopia

c) Binasal hemianopia

d) Bitemporal hemianopia

Question 3:

A 60-year-old diabetic male presents with a history of decreased vision. Fluorescein angiography showed the following image. What is the most likely diagnosis?



a) Mild non-proliferative diabetic retinopathy

b) Severe non-proliferative diabetic retinopathy

c) Bird shot choroidopathy

d) Proliferative diabetic retinopathy

Question 4:

What is the most probable diagnosis based on the given image?

a) 3rd cranial palsy

b) 6th nerve palsy

c) Bell's palsy

d) Horner's syndrome

Question 5:

A patient presents with endophthalmitis 5 days after cataract surgery. All of the following treatments can be used, except _____.

a) Intravitreal antibiotic injection

b) Intravenous steroids

c) Pars plana vitrectomy

d) Topical antibiotics

Question 6:

Identify the refractive error.



a) Myopia

b) Presbyopia

c) Hypermetropia

d) Astigmatism

Question 7:

What is the diagnosis of the condition seen in the image?



a) Pinguecula

b) Pterygium

c) Chemical burn

d) Thermal burn

Question 8:

A patient with glaucoma presents with a bulging cornea. What is the most likely diagnosis?

a) Keratoconus

b) Keratomalacia

c) Staphyloma

d) Granular dystrophy

Question 9:

A patient presents with unilateral ptosis with hypotropia in the same eye. He was given an IV injection, following which the symptoms go away. Which of the following is the most likely diagnosis?

a) Myasthenia gravis

b) 6th nerve palsy

c) 3rd nerve palsy

d) Horner syndrome

Question 10:

Which is the most common lacrimal gland tumor?

a) Adenoid cystic carcinoma

b) Pleomorphic adenoma

c) Mucoepidermoid carcinoma

d) Malignant mixed tumor

Question 11:

A patient recently underwent LASIK surgery for his myopia. He is now complaining of headaches and eye pain. Identify the test being performed on him.

a) Pachymetry

b) Tonometry

c) Auto Refractometery

d) A Scan

Question 12:

A patient presents to the emergency department and is found to have a corneal abrasion, corneal ulcer, corneal perforation, and foreign body. Which of the following will not cause hypotonic maculopathy in this patient?

a) Suprachoroidal haemorrhage

b) Cyclodialysis of uvea

c) Corneal perforation

d) Filtration site leak

Question 13:

Name the location of the nucleus supplying the muscle marked by the arrow in the image given below.



a) Superior olivary

b) Inferior colliculus

c) Superior colliculus

d) Facial colliculus

Question 14:

Silent choroid on FFA is a feature of which of the following disorders?

a) Stargardt's disease

b) Best's disease

c) Age related macular degeneration

d) Cystoid macular edema

Question 15:

The physician noticed the following finding in a patient that came to the OPD for an eye exam. What is the most likely diagnosis?



a) Entropion of right eye

b) Lagophthalmos of right eye

c) Ptosis of right eye

d) Exophthalmos of left eye

Answer Key

Question No.	Correct Option
1	a
2	d
3	d
4	b

5	b
6	a
7	b
8	c
9	a
10	b
11	b
12	a
13	c
14	a
15	c

Detailed Explanations

Solution to Question 1:

Diminished vision 1 year after cataract surgery, supported with the image of vacoulated clusters seen behind IOL is suggestive of Elschnig's pearls. It is a type of after cataract.

After cataract, or secondary cataract or posterior capsular opacification, is a late postoperative complication that follows extracapsular extraction of the lens. It is the most common late postoperative complication. It may be caused by residual opaque lens matter or proliferation of left out anterior epithelial cells. Different types of after cataract include:

• Elschnig's pearls: The subcapsular epithelial cells proliferate and develop into vacuolated soap bubble like clusters along the posterior capsule.

• Soemmering's ring: A peripheral dense ring is formed behind the iris and is enclosed between layers of the capsule.

• Dense membranous after-cataract: It is formed in the pupillary area impairing the vision.

• Thin membranous after-cataract: It is thickened posterior capsule.

Treatment of after cataract is by Nd-YAG laser capsulotomy of the posterior capsule, which works on the principle of photodisruption, or surgical posterior capsulotomy.

Option B: Irvine–Gass syndrome is cystoid macular edema (CME) that develops in post-cataract surgery patients. Though it is mostly clinically insignificant, some cases can present with blurring of vision, one to three months after surgery. In CME, on fundoscopy a honeycomb appearance and on fundus fluorescein angiography(FFA) flower petal pattern can be seen. The below image depicts the flower petal pattern seen on FFA.



Option C: Post operative endophthalmitis presents as ocular pain, diminished vision and hypopyon within months to years after the surgery. Fungal endophthalmitis can rarely occur with puff ball exudates in the vitreous cavity.

Option D: UGH syndrome is where uveitis, glaucoma and hyphema occurs simultaneously. It is seen commonly with rigid anterior chamber intraocular lens (IOL) implantation.

Solution to Question 2:

Lesions at the level of the optic chiasma cause bitemporal hemianopia.

The Nasal fibers of each retina decussate at the optic chiasma. Hence, a lesion at this level affects the temporal quadrants of both eyes.

The optic chiasma lies directly above the tuberculum and diaphragma sellae & is affected in

• Pituitary adenoma

• Craniopharyngioma

• Meningioma

• Gliomas

• Multiple sclerosis, sarcoidosis

Option A: Central scotoma is seen in lesions affecting the optic disc (intraocular part of optic nerve) such as ethambutol toxic neuropathy, macular degeneration and lesions of visual cortex with bilateral homonymous macular defects.

Option C: Binasal hemianopia is seen in lesions affecting bilateral lateral chiasma where the temporal fibers of both the retina continue forward without crossing. This affects vision in the nasal quadrants of both eyes.

Solution to Question 3:

The given clinical vignette and Fluorescein angiography findings of neovascularization are suggestive of proliferative diabetic retinopathy (PDR).

Neovascularization is a hallmark of PDR. So this rules out Non-proliferative diabetic retinopathy-NPDR (Option A,B).

Option C: Birdshot retinochoroidopathy is an idiopathic bilateral chorioretinitis characterized by creamy yellow spots of chorioretinal hypopigmentation resembling a birdshot scatter. It most commonly affects middle aged females who are HLA-A29 positive.

PDR is most common in patients with Type ■ diabetes after a period of 15-20 years and is characterized by proliferation of new vessels at the optic disc (NVD) or elsewhere in the fundus (NVE).

(NVD- neovascularization at the optic disc ; NVE-neovascularization elsewhere in the fundus)

Complications of PDR include recurrent vitreous hemorrhage, tractional retinal detachment and pre retinal hemorrhages.

It is treated by administration of Anti-VEGF drugs followed by pan retinal photocoagulation which destroys the hypoxic retina that has been producing the vasoformative factors responsible for neovascularization.

Duration of diabetes mellitus is the most important factor for the development of diabetic retinopathy.

Ophthalmoscopic features of NPDR include:

• Microaneurysms: due to loss of pericytes around the capillaries

• Retinal hemorrhages: both superficial (flame shaped) and deep (dot & blot) hemorrhages.

• Hard exudates: composed of lipoproteins and lipid laden macrophages. They are yellow waxy looking and arranged in clumps or circinate patterns.

• Cotton wool spots: which are small whitish fluffy lesions which represent areas of nerve infarct.

• Intra retinal microvascular abnormalities(IRMA): are arteriovenous shunts in retina

• Venous abnormalities such as beading,looping and dilatation.



The below given image is that of Hard Exudates.

Solution to Question 4:

The given image depicts restricted lateral movement of the left eye when the person is in left lateral gaze. This is suggestive of left lateral rectus muscle paralysis due to isolated 6th cranial nerve palsy.

Isolated abducens(6th cranial) nerve palsy presents with esotropia / convergent squint in primary position with limitation of abduction on the side of the lesion.

This is due to unopposed action of the medial rectus muscle in the affected eye. There is also a compensatory head turn towards the side of the lesion to reduce the symptom of diplopia. The adduction is normal in 6th nerve palsy.

The various causes of 6th nerve palsy includes:

• Raised intracranial pressure

• Nasopharyngeal tumors

• Gradenigo syndrome as seen in mastoiditis or acute petrositis

• Basal skull fracture



Option A: 3rd Cranial nerve palsy presents with

• ptosis due to weakness of levator palpebrae superioris muscle(LPS)

• Abduction and depression in the primary position(down and out) due to unopposed action of the lateral rectus(6th nerve supply) and superior oblique muscles(4th nerve supply)

• Dilated pupil and defective accommodation due to parasympathetic palsy(involvement of Edinger Westphal parasympathetic supply)

• Limited adduction, elevation due to ipsilateral paralysis of medial rectus, inferior oblique and inferior rectus .

Option D: Bell's palsy is an acute onset, idiopathic lower motor neuron lesion of the facial nerve which involves the ipsilateral upper and lower halves of the face.

It presents with an inability to close the eye and any attempts to close the eye results in the eye turning upwards and outwards which is known as the Bell's phenomenon. Other features of Bell's palsy include drooling of saliva, deviation of angle of the mouth and facial asymmetry. Additionally there may be noise intolerance and a loss of taste sensation.

The following image shows a patient with Bell's palsy:



Option D: Horner syndrome, also called oculosympathetic paresis, is produced by a lesion anywhere along the sympathetic pathway supplying the head, neck and eyes. It is characterized by the classical triad of ptosis, miosis and anhidrosis(reduced ipsilateral sweating). Other features include mild enophthalmos, hypochromic heterochromia and anisocoria that is accentuated in dim light.

Tests for confirmation of Horner's syndrome -

• Cocaine drops (2-4%)

Horner pupil does not dilate

Normal pupil dilates

• Apraclonidine drops (0.5-1%)

Horner pupil dilates - due to upregulation of α1 receptors in the dilator pupillae.

Normal pupil does not dilate

Solution to Question 5:

Intravenous steroids are not used for the treatment of endophthalmitis.

Endophthalmitis is defined as an intraocular inflammation that predominantly affects the inner spaces of the eye and their contents, i.e. the vitreous and/or the anterior chamber. The cardinal features of endophthalmitis are pain, swelling of the lid and diminished vision. Once the endophthalmitis sets in it is important to treat all the cases empirically with a combination of antibiotics and steroids from the beginning.

Steroids are given in endophthalmitis to reduce the inflammation of the tissues. The preferred route of administration is either by intravitreal injections, topical concentrations or oral route. Intravenous route of administration of steroids is not used in endophthalmitis.

Acute postoperative bacterial endophthalmitis occurs within 1-7 days after cataract surgery and the most important factor to prevent it is preoperative administration of 5% povidone iodine.

The various drugs,routes and surgical management used to treat endophthalmitis are as follows:

• The mainstay of treatment is intravitreal injections of antibiotics.

First choice: Vancomycin 1 mg in 0.1 mL plus ceftazidime 2.25mg in 0.1ml.

Second choice: Vancomycin 1 mg in 0.1 mL plus amikacin 0.4mg in 0.1ml.

If there is no improvement, a repeat intravitreal injection should be given after 48 hours

• Topical Solutions of antibiotics are started immediately and used frequently

Vancomycin (50mg/ml) or cefazolin (50mg/ml)

Amikacin (20mg/ml) or tobramycin

• Systemic antibiotics have a limited role in the treatment but are commonly used in practice.

• Steroid therapy includes

Intravitreal injection of Dexamethasone(0.4mg in 0.1ml) which is started along with antibiotics.

Topical Dexamethasone (0.1%)

Oral steroids such as prednisolone are started after 24 hours of intensive antibiotic therapy.

• Supportive therapy are

Cycloplegics such as Atropine 1%

Anti-Glaucoma drugs such as oral acetazolamide or timolol in patients with raised intraocular pressure

• Vitrectomy- Pars Plana Vitrectomy is done if the patient does not improve with the above therapy in 48-72 hours or when the visual acuity is reduced to hand movements near the eye.

Solution to Question 6:

In the above image, parallel rays are focused in front of the retina and hence the refractive error shown is Myopia.

Myopia, also known as short sightedness, is that dioptric condition of the eye in which, with the accommodation at rest, incident parallel rays come to a focus anterior to the light-sensitive layer of the retina. In myopia the defect in vision is more for objects present at a distance.

American Optometric Association (AOA) Grading of Myopia:

• Low myopia, when the error is ≤ –3D.

• Moderate myopia, when the error is between -3D to -6D.

• High myopia, when the error is ≥ –6D.



Option B: Presbyopia is used to define physiological insufficiency of accommodation that causes progressive loss of near vision. It is commonly seen in elderly due to age-related decline in ciliary muscle power. Bifocals spectacles are prescribed in presbyopia.

Option C: In hypermetropia, parallel rays of light are focussed behind the retina causing defective vision which is more for near objects.

Option D: In astigmatism, the rays of light entering the eye fail to converge at a point focus but they form focal lines. Strum's connoid of astigmatism represents the configuration of a bundle of light rays formed by a regularly astigmatic(toric) surface and helps in understanding the optics of regular astigmatism.



Solution to Question 7:

The above image depicting a wing shaped encroachment of conjunctiva onto the cornea is suggestive of Pterygium.

Pterygium is a triangular fibrovascular ingrowth that is seen following an elastotic degeneration of the subconjunctival tissue. The apex of the pterygium lies towards the cornea. The pterygium is usually seen on the nasal side in the palpable aperture and can give rise to symptoms such as foreign body sensation in the eye, diplopia and defective vision due to corneal astigmatism. In some cases, iron deposition can be seen in the corneal epithelium anterior to the advancing head of the pterygium known as the stocker's line.

Pterygium can be left untreated if there is no visual impairment. In cases with impaired vision, procedures such as surgical excision with free conjunctival limbal graft & surgical excision with amniotic membrane graft and mitomycin-C can be done.

Option A: Pinguecula is an asymptomatic yellow-white mound of degenerated collagen fibers of the conjunctival stroma. It more commonly starts on the nasal limbus and can later be seen on the temporal side. The apex of the Pinguecula is directed away from the cornea. It is mostly a stationary condition, so no treatment is usually required. If inflamed, topical steroids can be applied.



Option C: Chemical burns are caused by alkalis and acids. In mild-moderate burns, the end result is corneal ulceration followed by opacification. Severe burns result in sloughing of the whole cornea followed by staphyloma formation. Complications of chemical burns include symblepharon where the palpebral & bulbar conjunctiva are fused.

Option D: Thermal burns from fire & hot fluids mainly affect the lids. Cornea & conjunctiva will be affected in severe cases.

Solution to Question 8:

High intraocular pressure leads to the thinning of the outer tunics of the eyeball such as cornea and sclera, thereby resulting in their ectasia (bulging) which is known as

staphyloma. Ciliary and equatorial staphylomas are most commonly associated with glaucoma.

Staphyloma is an abnormal protrusion of the uveal tissue through a weak and thin portion of the cornea or sclera. So, staphyloma is lined internally by uveal tissue and externally by weak cornea or sclera.

Types of Staphyloma:

Types	Pathology	Causes
Anterior staphyloma	Due to bulging of the pseudo cornea, internally lined by th e iris & externally by the newl y formed epithelium.	Sloughing corneal ulcer
Intercalary staphylom a	Due to bulging of weakened l imbal tissue at the limbus, in ternally lined by the root of ir is & the anterior most part of the ciliary body	Developmental glaucomaSec ondary angle closure glauco maHealing of perforating inj uryPeripheral corneal ulcer
Ciliary staphyloma	Due to bulging of the weaken ed sclera, internally lined by the ciliary body, can extend u p to 8mm behind the limbus	ScleritisEnd stage glaucomaP erforating injuries
Equatorial staphylom a	Due to bulging of sclera with choroidal lining, seen around 14 mm behind the limbus, in the equatorial region	ScleritisChronic uncontrolled glaucomaScleral degeneratio n in pathological myopia
Posterior staphyloma	Due to bulging of sclera lined by the choroid, seen behind the equator	Posterior scleritisPathologica l myopiaPerforating injuries



Keratoconus (Option A) is a progressive bilateral corneal ectatic disorder due to defective synthesis of mucopolysaccharide and collagen tissue. It usually starts at puberty and progresses slowly, not associated with risk factors like glaucoma. Therefore, we can rule out keratoconus in

this patient.

Solution to Question 9:

Only Myasthenia gravis among the conditions mentioned above, would show dramatic improvement after administration of an IV injection like Anti-Cholinesterase, edrophonium.

Myasthenia gravis is an autoimmune disease that is characterized by generalized muscle weakness and rapidly progressing fatigue that improves after short periods of rest. Ocular myasthenia gravis causes progressive external ophthalmoplegia which affects only the extraocular muscles. Early ocular manifestations include failure of convergence & ptosis which is mostly bilateral and increases on prolonged fixation or attempts to look upwards. In the initial phase, there can be unilateral myogenic ptosis, which improves after short rest rapidly.

Diagnosis of myasthenia gravis can be made by performing the tensilon test where the symptoms improve for a short period on administering a short acting Anticholinesterase drug such as edrophonium.

Note:

Single most sensitive test for Myasthenia Gravis - Single fiber Electromyography

The single most specific test for Myasthenia Gravis - AChR antibodies

Option B: 6th nerve palsy- isolated 6th nerve palsy presents with convergent squint with limitation of abduction in the affected eye, due to the lateral rectus muscle palsy.



Option A: 3rd Cranial nerve palsy presents with

• ptosis due to weakness of levator palpebrae superioris muscle(LPS)

• Abduction and depression in the primary position(down and out) due to unopposed action of the lateral rectus(6th nerve supply) and superior oblique muscles(4th nerve supply)

• Dilated pupil and defective accommodation due to parasympathetic palsy(involvement of Edinger Westphal parasympathetic supply)

• Limited adduction, elevation due to ipsilateral paralysis of medial rectus, inferior oblique and inferior rectus .

Option D: Horner syndrome, also called oculosympathetic paresis, is produced by a lesion anywhere along the sympathetic pathway supplying the head, neck and eyes. It is characterized by the classical triad of ptosis, miosis and anhidrosis(reduced ipsilateral sweating). Other features include mild enophthalmos, hypochromic heterochromia and anisocoria that is accentuated in dim light.

Tests for confirmation of Horner's syndrome -

• Cocaine drops (2-4%)

Horner pupil does not dilate

Normal pupil dilates

• Apraclonidine drops (0.5-1%)

Horner pupil dilates - due to upregulation of α1 receptors in the dilator pupillae.

Normal pupil does not dilate

Solution to Question 10:

Pleomorphic adenoma is the most common lacrimal gland tumor. It is also the most common epithelial tumor of the lacrimal gland.

Pleomorphic adenoma, or benign mixed tumor, presents commonly in middle age as a slowly progressive painless swelling in the upper outer quadrant of the eyelid. The bone is usually not invaded in these tumors. They are treated by surgical removal along with their intact pseudo-capsule.

Malignant tumors of the lacrimal gland such as adenoid cystic carcinoma (option A), mucoepidermoid carcinoma (option C) and malignant mixed tumor (option D) generally occur in the elderly with a short history of appearance, rapid progression and pain. These tumors also present in the upper eyelid and are known to invade the adjacent bones. Radical surgical procedures and high dose radiotherapy are necessary for the removal of the tumor. Adenoid cystic carcinoma is the most common malignant epithelial tumor of the lacrimal gland.

Both malignant and benign tumors are known to push the eyeball downwards and inwards while restricting the upwards and outwards movement of the eyeball.

The most common epithelial tumors, in order of frequency, are as follows:

• Benign mixed tumor (BMT), also known as a pleomorphic adenoma (PA)

• Adenoid cystic carcinoma (ACC)

• Pleomorphic adenocarcinoma (ACA)

• Mucoepidermoid carcinoma

Solution to Question 11:

The above image is of the dynamic contour tonometer such as the Pascal tonometer. The dynamic contour tonometer(Pascal) offers a non-invasive and direct method of intraocular pressure (IOP) measurement with an additional measurement of the ocular pulse amplitude. This type of tonometer is unaffected by the corneal thickness and it is the tonometer of choice in post-Lasik patients.

Option A: Pachymetry is used to measure the corneal thickness.

The below image is of pachymetry.



Option C: Auto Refractotemetry is a computerized refractometer that is used to find out the error of refraction. This gives rapid information about the kind of refractive error in terms of sphere, cylinder with axis and inter pupillary distance.

Option D: A-Scan ultrasonography is a useful diagnostic tool for intraocular evaluation especially in the setting of opaque media. The report of the A scan is displayed as a unidimensional display of spikes along a baseline where the distance between the spikes gives the indirect measurement of the tissues. A-scan is used to measure

• The axial length of the eyeball for Intraocular lens (IOL) power calculation

• Anterior chamber depth

• Thickness of intraocular mass.

A-scan ultrasonography is shown in the image below.

Solution to Question 12:

Suprachoroidal hemorrhage will not cause hypotonic maculopathy in this patient.

Maculopathy is common in eyes developing hypotony, which is defined as IOP <6 mmHg. The most common cause is excessive drainage following glaucoma filtration surgery (Option D). Other causes include trauma such as cyclodialysis of the uvea (Option B), penetrating injury (Option C), chronic uveitis, retinal detachment, and other systemic causes such as dehydration, uncontrolled diabetes, uremia, and treatment with hyperosmotic agents. They usually cause bilateral hypotonic maculopathy.

Suprachoroidal hemorrhage is a bleed into the suprachoroidal space that results from the rupture of the posterior ciliary artery. When the bleed is severe it causes extrusion of intraocular contents that is termed as expulsive hemorrhage. It is a rare but dreaded complication of the phacoemulsification procedure. Signs seen in suprachoroidal hemorrhage are:

• Progressive increase in intraocular pressure

• Prolapse of iris

• Loss or partial obscuration of the red reflex

• In severe cases there can be extrusion of the posterior segment contents into the anterior chamber and through the incision such as vitreous extrusion

Immediate treatment involves rapid closure of the incision followed by administration of IOP-lowering medications such as oral acetazolamide and topical or systemic steroids to reduce inflammation. Standard post-operative antibiotics can be added in these patients.

Solution to Question 13:

The muscle marked is the medial rectus. The medial rectus is supplied by the oculomotor nerve, the nucleus of which is situated at the level of the superior colliculus in the posterior midbrain.

The oculomotor (3rd cranial) nerve nucleus consists of

• Single central nuclei for the levator palpebrae muscle of both the eyes

• Bilateral subnuclei with crossed projections supplying the contralateral superior rectus

• Bilateral subnuclei with uncrossed projections supplying the ipsilateral inferior rectus, inferior oblique, medial rectus in the dorso-ventral order.

• Bilateral subnuclei known as the Edinger-Westphal nucleus which provides parasympathetic input to the ciliary body and the iris sphincter.

Option A: Superior olivary nucleus is an important relay station in the auditory pathway, it is responsible for detection of direction and localization of sound.

Option B: Inferior colliculus- The trochlear(4th cranial) nerve nucleus lies in the midbrain at the level of the inferior colliculus. The trochlear nerves leave the midbrain from its posterior surface and decussate before supplying the contralateral superior oblique muscle of the eyeball.

Option D: Facial colliculus- the facial colliculus is located in the cranial part of the pons. It is produced by the facial nerve(7th cranial nerve) winding around the nucleus of the abducens nerve (6th cranial nerve).

Solution to Question 14:

Silent/Dark choroid on FFA is a feature of Stargardt's disease. The classic feature of 'silent choroid' is due to the masking of background choroidal fluorescence by lipofuscin deposits in the subretinal space.

Stargardt disease constitutes the most common type of macular dystrophy. It has mutations in a photoreceptor cell-specific ATP-binding cassette gene (ABC4) that is usually inherited in an autosomal recessive manner. The disease usually starts in the juvenile group with progressive central vision loss and color vision loss that occurs due to the atrophy of the macular retinal pigment epithelium with an accumulation of discrete pisciform flecks of lipofuscin.

Signs of Stargardt’s disease:

• Beaten bronze appearance of the macula that is caused by the atrophic changes in the RPE.

• Silent/Dark choroid on FFA.

• Visual field testing shows central vision loss.

• The electroretinogram (ERG) is normal early in the disease but may be reduced in more advanced cases.

Fundus flavimaculatus is genetically considered to be a part of the same spectrum as Stargardt's disease with late onset i.e. usually after 40 years of age.

Other options:

Option B: Best's disease also known as vitelliform macular dystrophy is an autosomal dominant macular dystrophy in which lipofuscin accumulates in the central macula, causing progressive central vision loss. This disease is linked to mutations in the bestrophin (VMD2) gene and they do not exhibit silent choroid on FFA. Findings seen in Best's disease are:

• Vitelliform(yolk like) appearance of the fundus

• Relatively normal ERG results associated with abnormal EOG

The given below is the fundus picture of Best's disease showing an egg yolk lesion in the macula.



Option C: Age related Macular degeneration(ARMD) also known as senile macular degeneration is a chronic and progressive bilateral degenerative disorder of the macula that affects older individuals >50 yrs. ARMD is classified as dry and wet ARMD where the hallmarks are the presence of drusen and choroidal neovascularization respectively. Drusen are extracellular eosinophilic deposits between RPE and Bruch's membrane.

Clinical features include loss of central vision, distorted vision (lines seen as wavy), and sudden painless loss of vision as seen in wet ARMD. The image below shows Drusens in the macular area in dry macular degeneration.

Option D: Cystoid Macular edema (CME) is a collection of fluid between the outer plexiform layer and the inner nuclear layer of the retina following a break in the blood-retinal barrier. In CME on fundoscopy, a honeycomb appearance and on FFA flower petal pattern can be seen. The below image shows the flower petal pattern on FFA as seen in CME.



Solution to Question 15:

The given image of the patient shows ptosis of the right eye. Ptosis is abnormal drooping of the upper eyelid to a level that covers more than 2mm of the superior cornea, it commonly occurs due to paralysis or defective development of the levator palpebrae superioris (LPS).

Types:

• Congenital ptosis

• Acquired ptosis

• Neurogenic ptosis is caused by innervational defects such as third nerve palsy, Horner's syndrome.

• Myogenic ptosis occurs due to acquired disorders of the LPS muscle or the myoneural junction- Myasthenia gravis.

• Aponeurotic ptosis develops due to defects of the levator aponeurosis.

• Mechanical ptosis may result from the excessive weight on the upper lid as seen in patients with lid tumors, multiple chalazia, and lid edema.

Treatment of ptosis:

• The Fasanella–Servat operation is indicated for cases of minimal ptosis of 1.5 –2 mm with good function of the levator and a good lid fold.

• In moderate-to-severe ptosis with a moderate levator action, levator resection can be carried out.

Note: Pseudoptosis is apparent but not true drooping of the eyelid. Causes include lack of support of the lids by the globe(orbital volume deficit like an artificial eye, microphthalmos, or enophthalmos), contralateral lid retraction, ipsilateral hypotropia or brow ptosis.

Option A: Entropion is defined as inward rolling of the lid margin. The constant rubbing of the eyelashes onto the cornea can cause irritation, punctate corneal erosions and in severe cases ulceration and pannus formation. Weiss operation is done in senile entropion.

Option B: Lagophthalmos- in lagophthalmos there is incomplete closure of the palpable aperture when the patient attempts to close the eyes due to congenital deformity, paralysis of the orbicularis muscle, ectropion or contraction of the lid. Treatment includes the management of exposure keratopathy with frequent instillation of lubricants.



Option D: Exophthalmos- Although exophthalmos and proptosis are used interchangeably, the term exophthalmos is used to describe the protrusion/prominence of the eyes due to thyroid diseases while proptosis can be due to any other causes. Most common causes of proptosis:

Unilateral proptosis	Bilateral proptosis
In children: orbital cellulitis	Neuroblastoma & leukemia
In adults: Thyroid ophthalmopathy	Thyroid opthalmopathy

Ophthalmology AIIMS 2018

Question 1:

All of the following are true about oculomotor nerve palsy except _____.

a) Ptosis

b) Mid-dilated pupil

c) Loss of light reflex

d) Lateral and upward gaze

Question 2:

The center for vertical gaze is _____

a) Pontine Paramedian Reticular Formation

b) Raphe Nuclues

c) Rostral Interstitial Nucleus of Cajal

d) Nucleus of Perlia

Question 3:

You can use a stenopaeic slit for all of the following except _____.

a) Fincham's test

b) Corneal tattooing

c) Optical Iridectomy

d) To find out the axis in astigmatism

Question 4:

A 55-year-old male patient came to OPD with the complaints of glare during night drives. He had undergone cataract surgery 1 year back. The best-corrected visual acuity is RE = 6/12 and LE = 6/9. No improvement with pinhole. What is the probable diagnosis?

a) Cystoid macular edema

b) Pseudophakic bullous keratopathy

c) ARMD

d) Posterior capsular opacification

Question 5:

Identify the lesion in the histopathological image of retina.



a) Drusen

b) Hard exudates

c) Cotton wool spots

d) Macular Edema

Question 6:

Identify the picture

a) Macular Hole

b) Cystoid Macular Edema

c) Smoke stack pattern

d) Age related Macular degeneration

Question 7:

What is the adjuvant therapy in fungal corneal ulcer?

a) Pilocarpine e/d

b) Lignocaine e/d

c) Dexamethasone e/d

d) Atropine Sulfate eye ointment

Question 8:

All are ocular manifestations of dengue except _____.

a) Cataract

b) Subconjunctival hemorrhage

c) Uveitis

d) Retinal hemorrhage

Question 9:

A 5-year-old boy was brought with the complaints of protrusion of the right eye with no h/o fever. CT Scan showed a well-defined mass in the orbit with an irregular border and adjacent bony destruction. Biopsy showed small, round cells which were positive for Desmin in Immunohistochemistry. What is the probable diagnosis?

a) Retinoblastoma

b) Cavernous Hemangioma

c) Orbital Cellulitis

d) Rhabdomyosarcoma

Question 10:

A 65-year-old female patient with a history of chronic arthritis on treatment came with complaints of defective vision in both eyes. On examination, vision in RE = 6/12, LE = 6/18, IOP = 18 mmHg in both eyes, the anterior segment was normal, fundus examination and fundus fluorescein angiography (FFA) findings given in the figure. What is the probable diagnosis?

a) Choroiditis

b) Chloroquine toxicity

c) Steroid-induced glaucoma

d) Macular edema

Question 11:

A 30-year-old man presents with a history of injury to the eye caused by a leaf 5 days ago, resulting in pain, photophobia, and redness of the eye for 2 days. What would be the most likely pathology?

a) Anterior uveitis

b) Conjunctivitis

c) Fungal corneal ulcer

d) Corneal laceration

Question 12:

What is the angle subtended by the largest alphabet on the Snellen's chart at 60m distance?

a) 5

b) 30

c) 50

d) 10

Question 13:

Identify the marked layer of the retina.



a) Inner plexiform layer

b) External limiting layer

c) Internal limiting layer

d) Outer plexiform layer

Question 14:

The spectacles shown in the image are not used for the management of which of the following?

a) Presbyopia

b) Aphakia

c) Pediatric pseudophakia

d) Hypermetropia

Question 15:

A 6-year-old girl is admitted for evaluation for recurrent seizures. She has a large port-wine stain on the right side of her face as shown below. The MRI of the head shows a leptomeningeal angioma. What is the most common ocular manifestation of the underlying condition?

a) Uveitis

b) Retinitis pigmentosa

c) Keratitis

d) Glaucoma

Question 16:

The child in the picture has large corneas, photophobia, and lacrimation. What is the likely diagnosis?

a) Congenital glaucoma

b) Retinoblastoma

c) Congenital hereditary endothelial dystrophy

d) Megalocornea

Question 17:

The following image indicates:

a) Trochlear nerve palsy

b) Lateral rectus palsy

c) Internuclear ophthalmoplegia

d) Occulomotor palsy

Question 18:

A 26-year-old female patient presents with 6/60 visual acuity in one eye and 6/18 in the other eye. She has recurrent episodes of diminution of vision which recovers on treatment with steroids. She also has a history of transverse myelitis. What is the most probable diagnosis?

a) Hereditary spastic quadriplegia

b) Subacute combined degeneration

c) Neuromyelitis optica

d) Multiple sclerosis

Question 19:

Which of the following is the site of lesion in internuclear ophthalmoplegia?

a) Medial Longitudinal Fasciculus

b) 6th Nerve nucleus

c) Pontine paramedian reticular formation

d) A & C

Answer Key

Question No.	Correct Option
1	d
2	c
3	b
4	d
5	a
6	b
7	d
8	a
9	d
10	b
11	c
12	a
13	a
14	d
15	d
16	a
17	d
18	c
19	a

Detailed Explanations

Solution to Question 1:

Lateral and upward gaze is a sign of trochlear (4th) nerve palsy and not of oculomotor palsy. Oculomotor (3rd) nerve palsy leads to lateral and downward gaze.

3rd Cranial nerve palsy presents with

• Ptosis(option A) due to weakness of levator palpebrae superioris muscle (LPS)

• Abduction and depression in the primary position ("down and out") due to unopposed action of the lateral rectus (6th nerve supply) and superior oblique muscles (4th nerve supply)

• Dilated pupil(option B), loss of light reflex(option C) and defective accommodation due to parasympathetic palsy (involvement of Edinger Westphal parasympathetic supply)

• Limited adduction, elevation and depression due to ipsilateral paralysis of medial rectus, inferior oblique and inferior rectus .

4th cranial nerve palsy presents with superior oblique muscle paralysis.

Clinical findings of trochlear nerve palsy include:

• Hypertropia (elevation) in the primary position

• Extorsion that is greatest in abduction

• Compensatory head posture with contralateral face turn and head tilt with chin slightly depressed. This position is assumed to avoid Diplopia.

• Limitation of depression

• Normal abduction and elevation

The below given image depicts 4th cranial nerve palsy.

Solution to Question 2:

The Rostral Interstitial Nucleus of Cajal is the center for vertical gaze.

Vertical eye movement initiation in the frontal lobes is mediated by the impulses generated via the vertical gaze center called Rostral interstitial Nucleus of Cajal. This nucleus is a prominent group of cells present within the Medial Longitudinal Fasciculus (MLF) in the midbrain.

Pontine Paramedian Reticular Formation(Option A) is the center for horizontal gaze. Classically, initiation occurs in the premotor cortex frontal eye fields from where impulses for horizontal movement pass to the contralateral paramedian pontine reticular formation (PPRF).

Both horizontal and vertical gaze are saccadic types of conjugate eye movements. Conjugate eye movements, also known as versions, are binocular movements in which the eyes move synchronously in the same direction.



Solution to Question 3:

Stenopaeic slit is not used for corneal tattooing.

Stenopaeic slit is an ophthalmological instrument that is used in:

• Fincham's test (Option A) : This test is used to differentiate between the colored halos due to immature cataract and corneal abnormalities such as corneal edema due to glaucoma.

• To find out the axis in astigmatism (option D) for prescribing the correction glasses.

• To find out the optimal site for optical iridectomy (option C) for treating cases of central macular or leucomatous corneal opacities.

Corneal tattooing was done mainly in the past for cosmetic purposes. It is commonly done for firm scars of corneal opacities in a eye without any useful vision.

The below given image is of stenopaeic slit.



Solution to Question 4:

The given clinical picture of a post cataract surgery patient coming with symptoms of glare and reduced visual acuity which doesn't improve on pinhole testing is suggestive of posterior capsular opacification (PCO).

Pinhole testing is done to differentiate between the various causes of reduced visual acuity.On pinhole testing if the visual acuity remains the same or worsens it is suggestive of either a macular disease or posterior capsular opacification.

Posterior capsular opacification also known as secondary cataract or After cataract is a late postoperative complication that follows extracapsular extraction of the lens.

Different types of after cataract include:

• Elschnig's pearls: where the subcapsular epithelial cells proliferate and develop into vacuolated soap bubble like clusters along the posterior capsule

• Soemmering's ring: A peripheral dense ring is formed behind the iris and is enclosed between layers of the capsule.

• Dense membranous after cataract: formed in the pupillary area impairing the vision.

• Thin membranous after cataract (thickened posterior capsule)

Treatment of after cataract is by:

• Nd-YAG laser capsulotomy of the posterior capsule whose principle is photodisruption.

• Surgical posterior capsulotomy.



The below given image is of Elschnig's pearls type of after catarct.



Option A: Cystoid Macular edema (CME) is a collection of fluid between the outer plexiform layer and the inner nuclear layer of the retina following a break in the blood-retina barrier. In CME, on fundoscopy a honeycomb appearance and on fundus fluorescein angiography (FFA) flower petal pattern can be seen.

The below given image shows the flower petal pattern on FFA as seen in CME

Option B: Pseudophakic bullous keratopathy is a postoperative complication of cataract surgery where the patient presents with corneal edema. This occurs due to a surgical or chemical insult to the corneal endothelium during the cataract surgery. It is treated by penetrating keratoplasty.

Option C: Age related Macular degeneration (ARMD) also known as senile macular degeneration is a chronic and progressive bilateral degenerative disorder of the macula that affects older individuals >50 yrs. ARMD is classified as dry and wet ARMD where the hallmarks are presence of drusen and choroidal neovascularization respectively.

Drusen are extracellular eosinophilic deposits between RPE and Bruch's membrane. Clinical features include loss of central vision, distorted vision (lines seen as wavy) and sudden painless loss of vision as seen in wet ARMD. The below given image shows Drusens in the macular area in Dry macular degeneration.

Solution to Question 5:

The below given image shows extracellular eosinophilic deposits between retinal pigment epithelium(RPE) and bruch's membrane which is suggestive of drusens.

Drusens are seen in dry age related macular degeneration(ARMD) and are composed of lipids, complement factors and other cellular components.

Age related Macular degeneration(ARMD) also known as senile macular degeneration is a chronic and progressive bilateral degenerative disorder of the macula that affects older individuals >50 yrs.

It is the leading cause of blindness in elderly in developed countries and its risk factors include smoking, exposure to sunlight, hypertension, female gender and nutritional deficiencies.

ARMD generally starts as dry ARMD which is characterized by the presence of drusen and gradually progresses to the advanced form known as wet ARMD whose hallmark is choroidal neovascularization with additional findings such as geographic atrophy of retinal pigmentatory epithelium (RPE) and photoreceptors at the macula.

Clinical features include loss of central vision, distorted vision(lines seen as wavy) and sudden painless loss of vision as seen in wet ARMD.

On fundus fluorescein angiography (FFA) the Druscens shows hyperfluorescent spots arranged in ‘stars in the sky’ appearance.

Disease management is by:

• Preventative measures to delay progression such as cessation of smoking, intake of good nutrition and supplementation of vitamins C,E and zinc.

• Treatment modalities with intravitreal Anti-VEGF injection- Bevacizumab (avastin) and Ranibizumab (Lucentis), photodynamic therapy (PDT) and submacular surgery.

The below given images show drusens in macular area and FFA depicting the 'stars in sky appearance' of drusens respectively.



Option B: Hard exudates are composed of lipoproteins and lipid laden macrophages. They are yellow waxy looking and arranged in clumps or circinate patterns. They are seen in diabetic retinopathy, hypertensive retinopathy and coat's disease.

The below given image is that of Hard Exudates.

Option C: Cotton wool spots also known as soft exudates are small whitish fluffy lesions which represent areas of nerve infarcts in the nerve fiber layer of the retina. They are seen in central retinal vein occlusion (CRVO), hypertensive retinopathy, diabetic retinopathy, toxaemic retinopathy of pregnancy and papilledema.



Option D:Cystoid Macular edema(CME) is a collection of fluid between the outer plexiform layer(Henle's layer) and the inner nuclear layer of the retina.

In CME, on fundoscopy a honeycomb appearance and on FFA, flower petal pattern can be seen. Optical coherence tomography(OCT) of CME shows loss of foveal depression and intraretinal cystoid spaces.

The below given OCT image shows intraretinal cystoid spaces seen in CME.



Solution to Question 6:

The given Fundus fluorescein angiography(FFA) image depicts the flower petal pattern that is seen in Cystoid macular edema.

Cystoid Macular edema(CME) is a collection of fluid between the outer plexiform(Henle's) layer and the inner nuclear layer of the retina following a break in the blood-retina barrier.

CME can manifest in the following conditions

• Post cataract surgery- Pseudophakics (called Irvine-Gass syndrome)

• Topical prostaglandins side effect(especially in diabetics)

• Niacin(Vitamin B3) toxicity

• Epinephrine(especially in Aphakia)

• Uveitis

• Retinitis pigmentosa

• Central and branched retinal venous occlusion

• Diabetic retinopathy

• Retinopathy of prematurity

Clinical features include sudden painless diminution of vision, blurring of vision and metamorphosis such as micropsia.

In CME, on fundoscopy, a honeycomb appearance and on FFA, flower petal pattern can be seen. On Optical coherence tomography(OCT), loss of foveal depression and intraretinal cystoid spaces can be seen.

Treatment of CME is by treatment of causative factor, administration of anti-prostaglandins such as NSAIDS(ketorolac) preoperatively and topical or systemic steroids.

The below given OCT image shows intraretinal cystoid spaces seen in CME.



Option A:Macular hole is a partial or full-thickness hole seen in the neurosensory retina of the foveal region. It could be idiopathic, senile or caused due to trauma or CME. There is a decrease in vision and metamorphopsia seen in this condition. Amsler's grid testing shows a central scotoma. FFA shows absent foveal reflex with a macular hole.

An OCT image of a full-thickness macular hole is shown below:



Option B: smoke stack pattern is seen on FFA in central serous retinopathy(CSR). CSR is the spontaneous serous detachment of the sensory neuroretina in the foveal region with or without retinal pigment epithelium (RPE) detachment occuring due to choroidal vascular hyperpermeability.

On FFA it is initially seen as an ink-blot pattern or smoke stack pattern of hyperfluorescence which then spreads to take the configuration of mushroom or umbrella pattern.

The below given images depict the smoke stack pattern and mushroom pattern respectively.





Option D:Age related Macular degeneration (ARMD) also known as senile macular degeneration is a chronic and progressive bilateral degenerative disorder of the macula that affects older individuals >50 yrs. ARMD generally starts as dry ARMD which is characterized by the presence of drusen and gradually progresses to the advanced form known as wet ARMD whose hallmark is choroidal neovascularization with additional findings such as geographic atrophy.

Drusen are extracellular eosinophilic deposits between retinal Pigment Epithelium and Bruch's membrane. The below given FFA image shows stars in sky appearance of Drusen.



Solution to Question 7:

Atropine sulfate (1%) eye ointment or eye drops is used as the nonspecific supportive treatment in both bacterial as well as a fungal corneal ulcer.

Atropine sulphate (1%) is a cycloplegic drug that is used as an adjunct in the treatment of corneal ulcers to

• reduce pain from ciliary spasm

• Prevent posterior synechiae formation that can develop from secondary iridocyclitis

• Increase blood supply to the anterior chamber that gets more antibodies into aqueous humor

• Reduce exudation by decreasing hyperpermeability of the vessels

Another cycloplegic that can be used as an adjunct is homatropine 2%.

Topical steroids such as Dexamethasone(option C) will increase fungal replication and corneal invasion and hence are contraindicated in the treatment of fungal corneal ulcers.

Management of fungal corneal ulcer includes

Definitive treatment:

• Topical antifungal eye drops: used for 6-8 weeks

• Filamentous fungi such as Aspergillus and Fusarium- Drug of choice (DOC) is natamycin (5%). Others include amphotericin B (0.1-0.3%), fluconazole (0.2%) or voriconazole 10%.

• Yeast such as Candida- Drug of choice (DOC) is Amphotericin B. Others include Nystatin 3.5%.

• Intracameral or intracorneal administration of Voriconazole in cases with intraocular extension or anterior chamber involvement.

• Systemic antifungals such as tablet fluconazole are given in severe cases of deeper fungal keratitis.

• Therapeutic penetrating keratoplasty for unresponsive cases.

Adjunct therapy:

• Cycloplegics : Atropine sulphate 1% or homatropine 2%

• Systemic analgesics and anti-inflammatory drugs such as paracetamol, ibuprofen to decrease pain and edema.

• Vitamins A,B-complex,C- to aid healing of the ulcer.

• Hot fomentation, dark goggles, rest, good diet.

Solution to Question 8:

Cataract is not an ocular manifestation of dengue.

Dengue has varied manifestations, in different organ systems, mainly of inflammation and bleeding. The mechanisms for ocular manifestations in dengue include a direct viral effect on the eye, an immune-mediated vasculitis, and a variety of secondary effects related to dengue-induced vascular leakage and hemostatic abnormalities.

Ocular manifestations of dengue are:

• visual impairment which can be either painless or painful

• Retro orbital pain

• Conjunctival injection and subconjunctival hemorrhage(option B)

• Uveitis(option C)

• Complications such as retinal hemorrhage(option D), retinal edema, macular ischemia, macular edema, vitreous hemorrhage, vitritis, and optic neuritis or papillitis.

Depending on the severity of the disease, gradual improvement in vision occurs, although in rare cases permanent visual impairment may be seen.

Solution to Question 9:

The clinical scenario of a patient with complaints of protrusion of eye with well-defined mass in the orbit and adjacent bony destruction on CT scan and biopsy showing small, round cells positive for Desmin favors rhabdomyosarcoma.

Cavernous hemangioma(option B) and orbital cellulitis(option C) usually are associated with fever, hence ruled out. Retinoblastoma(option A) is associated with Flexner–Wintersteiner rosettes on microscopy, therefore its not correct.

Rhabdomyosarcoma is the most common soft tissue sarcoma of childhood usually in the head and neck region, most common primary orbital malignancy in children. They are malignant tumors arising from voluntary muscle and produce a rapidly increasing proptosis. It extends in all directions. Diagnosis is done using biopsy in which cross-striations in the tumor cells are pathognomonic. Myoglobulin is a specific marker but it can also be positive for desmin.

Subtypes include embryonal(majority), alveolar, botryoid and pleomorphic.

Treatment includes combination of chemotherapy and radiotherapy.

• Two injections of vincristine, cyclophosphamide, and actinomycin D are given at weekly intervals before radiotherapy.

• Radiotherapy is given during which vincristine and cyclophosphamide are administered weekly.

• After radiotherapy, a combination of vincristine, cyclophosphamide, and doxorubicin is given for a year or longer in those patients in whom metastases were detected.

Other options:

Option A: Retinoblastoma is the most common intraocular malignancy in children. It is associated with mutations in the RB1 gene at chromosome 13q14. White pupillary reflex or leukocoria is the most common presenting feature, followed by a squint which is usually convergent. Histopathology of the tumor shows small round cells with large nuclei. Microscopic features includes Flexner-Wintersteiner rosettes(highly specific of retinoblastoma), Homer-Wright rosettes, pseudorosettes, and fleurettes formation. Areas of necrosis and calcification can also be seen.

Option B: Cavernous hemangioma is the most common primary benign orbital tumor in adults. They are composed of large, dilated vascular channels; are more infiltrative; frequently involve deep structures; and do not spontaneously regress. On histologic examination, the mass is sharply delineated but unencapsulated with large, cavernous blood-filled vascular spaces, separated by connective tissue stroma.

Option C: Orbital cellulitis is the most common cause of unilateral proptosis in children. It refers to the acute infection of the contents of the orbit behind the orbital septum. Features include pain, lid edema, erythema, chemosis, and axial proptosis. Increased pain with movement of eyes and ophthalmoplegia(decreased ocular motility) is a classic symptom. With posterior exenteration cavernous sinus thrombosis, subdural empyema, and intracranial abscess develop.

Solution to Question 10:

The given clinical viginette of female patient on treatment for chronic arthritis which is most likely the drug chloroquine given for rheumatoid arthritis, now presenting with decreased visual acuity, fundoscopy with a typical bull's eye sign and Fundus fluroscein angiography showing increased fluroscence in the macular area is suggestive of chloroquine toxicity.

Chloroquine and it's derivative hydroxychloroquine (HCQ) are antimalarial drugs which is also used to treat connective tissue disorders such as Rheumatoid arthritis, Systemic Lupus Erythematosus (SLE) and Discoid Lupus.

The maximum dose allowed for chloroquine is 6 mg/kg in 24 hours while for hydroxychloroquine it is 4.5 mg/kg. Hydroxychloroquine has a lower risk of ocular toxicity and hence is percived to be a safer drug but with weaker anti-inflammatory effect.

Manifestations of Chloroquine toxicity:

• Reversible on stopping the drug: difficulty in night vision, blurring of vision, diplopia, keratopathy

• Central and paracentral scotomas

• Earliest fundus findings: blunting of foveal reflex, irregularity in the macular pigmentation.

• Advanced stages: 'Bull's eye maculopathy' where the central irregular pigmentation may become surrounded by a concentric zone of hypopigmentation

• End-stage appearance includes bullseye maculopathy, pigmentary degeneration of retina and bone spicule formation, vascular attenuation, and optic atropy.

Screening should include a baseline examination for all patients before starting these drugs followed by annual screening starting after 5 years of use.

Pre-existing maculopathy is considered a contraindication to treatment with chloroquine and HCQ because underlying abnormalities may mask early toxicity.

Screening examinations include visual acuity testing, dilated fundus examination, spectral-domain optical coherence tomography (SD-OCT) testing, and functional assessment with Humphrey test.

Various causes of Bull's eye maculopathy are:

• Chloroquine and HCQ toxicity

• Cone-Rod dystrophy

• Advanced stargardt's disease

• Clofazimine retinopathy

Solution to Question 11:

The given clinical vignette of the patient presenting with pain, photophobia, and redness of the eye following trauma by a vegetative matter is suggestive of a fungal corneal ulcer.

Fungal corneal ulcer also known as mycotic keratitis is usually caused by trauma from vegetative matter such as branches of trees, leaves and animal tails. Injudicious use of topical steroids and antibiotics increases the predisposition to develop secondary fungal keratitis.

Mycotic keratitis can be caused by

• Filamentous fungi such as Aspergillus and Fusarium

• Yeasts such as Candida and cryptococcus

• Dimorphic fungi such as Histoplasma and Blastomyces

Typically in fungal corneal ulcers, the signs are more prominent than the symptoms. The symptoms of fungal keratitis are similar to that of bacterial ulcers which include pain, lacrimation, photophobia, blurred vision, and redness of the eyes.

A typical fungal ulcer has the following signs:

• Greyish white looking dry corneal ulcer with rolled out margin.

• Feathery finger-like extensions into the corneal stroma under intact epithelium

• Wessley ring which is a sterile immune ring that occurs due to the deposition of immune complexes

• Multiple small satellite lesions around the ulcer

• Hypopyon: The hypopyon in fungal ulcer is immobile and unsterile unlike bacterial ulcer (sterile hypopyon)

• Perforation is rare

• Corneal vascularization is usually absent.

A diagnosis of fungal ulcer is made by a typical history of trauma with vegetative matter followed by clinical examination and KOH mount or SDA agar cultures from corneal scrappings.

The image below depicts a fungal corneal ulcer with the arrowhead indicating a satellite lesion.



Other options:

Option A: Anterior uveitis is the inflammation of the uveal tissue from the iris up to the pars plicata of the ciliary body. Infectious uveitis can be caused by the exogenous entry of the organisms following penetrating injuries, perforation of corneal ulcer, and post-intaocular operations.

Acute anterior uveitis presents with symptoms of pain, photophobia, lacrimation, blurring of vision and redness of the eye. Signs of anterior uveitis include:

• Aqueous flare - It is the earliest sign of acute anterior uveitis, it occurs following the leakage of protein particles into the aqueous humor.

• Keratic precipitates(KP) - These are pathognomonic for anterior uveitis. They are proteinaceous cellular deposits at the back of the cornea arranged in a triangular fashion. These include mutton fat KP's as seen in granulomatous uveitis and granular KP's of non granulomatous uveitis.

• Iris nodules: typically occur in granulomatous uveitis. Koppe's nodules are at the pupillary border and Busacca's nodules are situated near the collarette.

Option C: Conjunctivitis is the inflammation of the conjunctiva of which the bacterial conjunctivitis is the most common. Acute bacterial conjunctivitis is characterized by marked

conjunctival hyperemia, swelling of the conjunctiva(chemosis), edematous eyelids, sticking together of lid margins and mucopurulent discharge from the eye.

The image shows a patient with bacterial conjunctivitis.

Option D: Corneal laceration: Partial corneal laceration or lamellar corneal laceration occurs from a blunt trauma and is treated by patching and antibiotics.

Solution to Question 12:

The angle subtended by the largest alphabet on the Snellens chart at 60m distance is 5 minutes.

Visual acuity is a measure of the spatial resolution of the eye or an estimation of its ability to discriminate between two points. Snellens chart is used for Visual acuity assesment.

In snellens chart the alphabets decrease in size as we move from above-downwards and these lines, from above-downwards should be read at 60, 36, 24, 18, 12, 9, 6, 5 and 4 m respectively. At these distances, the respective letters subtend a visual angle of 5 minutes. Hence, the largest alphabet that should be read at 60 m subtends a visual angle of 5 minutes in the eye.



Solution to Question 13:

The inner plexiform layer is the marked layer in the given histological image of retina, identified as the fourth layer from the uppermost (innermost) layer.

The retina consists of 10 layers. The retinal pigment epithelium (RPE) is the outermost layer - below the Bruch's membrane of the choroid - and the internal limiting membrane is the innermost layer - separating the retina from the vitreous.

The retinal pigment epithelium consists of a single layer of cells containing pigment. It is separated from the neurosensory retinal layers by the potential space between them which is known as interphotoreceptor matrix.

The inner plexiform layer consists of synapsing nerve fibers. The axons of bipolar cells synapse with dendrites of ganglion cells; and both these processes synapse with processes

of amacrine cells. It also contains some horizontally placed internal plexiform cells; and also a few ganglion cells.



Solution to Question 14:

The given image shows spectacles made of lenses with two different optical power. Thus, it is a pair of bifocal spectacles which is not used in hypermetropia or myopia.

Bifocal glasses have different corrective powers in the upper segment for distant vision and lower segment for near vision. Thus, they are useful in conditions where both distance and near vision are affected, like the following:

1. Presbyopia (option A): It is a definitive indication. It is used to correct age-related near and far vision defects.

2. Aphakia (option B): It provides both near and far vision correction when intra-ocular lens insertion is contraindicated.

3. Paediatric pseudophakia (option C): It is used to correct refractive errors or accommodative fluctuations post intra-ocular lens (IOL) insertion.

4. Non refractive accomodative esotropia

In hypermetropia there is defect in near vision only and hence a single vision glasses having uniform corrective power throught the entire surface is used instead of bifocal glasses.



Solution to Question 15:

The given clinical picture of a young child presenting with unilateral port wine stain in the region of trigeminal nerve supply and leptomeningeal angioma as seen on MRI is indicative of Sturge-Webber syndrome (SWS). Glaucoma is the most common ocular manifestation of SWS.

Sturge weber syndrome is a neurocutaneous disorder(phakomatoses) that is characterized by unilateral capillary malformation at the region of trigeminal supply known as port wine stain or nevus flammeus, ipsilateral diffuse cavernous hemangioma of the choroid and ipsilateral leptomeningeal hemangiomatosis, which causes seizures and mental retardation.

Ocular manifestation of SWS include:

• Glaucoma - most common ocular manifestation. It develops anywhere between infancy to adulthood. When developed in infancy it presents as buphthalmos.

• Diffuse choroidal hemangioma

• Serous retinal detachment

• Visual field defects- homonymous hemianopia due to the leptomeningeal hemangiomas affecting the optic tract or occipital lobe.

• Heterochromia of the iris

• Episcleral and conjunctival lesions

Option A: Uveitis is the inflammation of the uveal tissue of the eyes. It can be associated with a number of systemic conditions such as sarcoidosis, behcet's disease, ankylosing spondylitis, reiter's syndrome,Toxoplasmosis and Posner–Schlossman syndrome (PSS).

Option B: Retinitis pigmentosa is an inherited or acquired pigmentatory retinal dystrophy affecting the rods more. The earliest manifestation is worsening of night blindness which gradually progresses to tunnel vision and ultimately loss of central vision. The classic triad of findings in RP comprises bone-spicule retinal pigmentation, arteriolar attenuation and waxy disc

pallor.

Systemic Associations of RP include:

• Laurence-Moon-Biedl syndrome

• Cockayne's syndrome

• Refsum's syndrome

• Usher's syndrome

• Bassen-Kornzweig syndrome

The below given image shows the classical triad findings of Retinitis pigmentosa.

Option C:Keratitis is the inflammation of cornea which is characterised by corneal edema, cellular infiltration and ciliary congestion. It can be either infectious or non-infectious in nature which includes allergic keratitis, neurotrophic keratitis, exposure keratitis and peripheral ulcerative keratopathies such as Rosacea keratitis and Mooren's ulcer.

Solution to Question 16:

The given clinical scenario of an infant with large corneas, lacrimation and photophobia supported with the image is suggestive of congenital glaucoma.

Primary congenital glaucoma (PCG) also known as developmental glaucoma refers to abnormally high intraocular pressure(IOP) which results from the developmental anomaly of the angle of the anterior chamber.

It can be classified as:

• True congenital or newborn glaucoma in which IOP is elevated during intrauterine life and the baby is born with ocular enlargement.

• Infantile glaucoma: which manifests prior to age 3 years.

• Juvenile glaucoma: IOP rises between 3 and 16 years of age.

Clinical findings include:

• Blepharospasm

• Buphthalmos: Large eyeball due to stretching of the globe from elevated IOP prior to the age of 3 years.

• Corneal enlargement:

• Corneal haze is due to corneal edema.

• Blue sclera - due to the shining of the underlying uveal tissue through the thinned-out sclera

• Haab striae are healed breaks in Descemet's membrane

Treatment:

• Goniotomy - when the angle of the anterior chamber is visible through the clear cornea.

• Trabeculectomy when the corneal clouding prevents visualization of the angle.

• Filtration procedures such as Trabeculectomy or combined trabeculectomy with trabeculectomy if the above procedures fail.

• Glaucoma drainage devices in unresponsive cases.

Other options:

Option A: Retinoblastoma is the most common primary intraocular malignancy of childhood that is seen between 1-2 years.

Findings of Retinoblastoma include:

• Leukocoria (white pupillary reflex)- the most common manifestation

• Strabismus

• Painful red eye with secondary glaucoma

• An endophytic tumor grows into the vitreous as a white mass that may ‘seed’ into the gel. If calcification is present it gives a "cottage cheese appearance". An exophytic tumor grows outwards and causes overlying exudative retinal detachment.

The image below shows the leucocoria of the left eye.



Option C: Congenital hereditary endothelial dystrophy (CHED) is an inherited corneal dystrophy that manifests since birth with endothelial degeneration. It appears as diffuse bilateral corneal edema in the absence of elevated IOP. The clinical features of the disease include corneal clouding, visual impairment, and nystagmus.

Option D: Megalocornea is a bilateral non-progressive X-linked condition that most commonly affects males. In this condition, the horizontal corneal diameter is ≥12mm(<2 years) and is ≥13 mm(>2 years). It is characterized by a clear cornea, normal corneal thickness, and no breaks in the descemet's membrane. Systemic associations of Megalocornea include Marfan's syndrome, Apert syndrome, Ehlers-Danlos, and Down syndrome.

Solution to Question 17:

The given image is suggestive of oculomotor nerve palsy.

The central image shows the primary position of gaze where the patient's left eye is deviated down and out with ptosis of the left eyelid. From the eye movements in other positions, it can be seen that adduction and elevation are absent and depression is reduced in the left eye. All of these findings are indicative of oculomotor nerve palsy.

3rd Cranial nerve(oculomotor nerve) palsy presents with:

• Ptosis due to weakness of levator palpebrae superioris muscle (LPS)

• Abduction and depression in the primary position (down and out) due to unopposed action of the lateral rectus(6th nerve supply) and superior oblique muscles(4th nerve supply)

• Dilated pupil and defective accommodation due to parasympathetic palsy (involvement of Edinger Westphal parasympathetic supply)

• Limited adduction, elevation and depression due to ipsilateral paralysis of medial rectus, inferior oblique and inferior rectus .

Option A: Trochlear(4th cranial) nerve palsy presents with superior oblique muscle paralysis.

Clinical findings of trochlear nerve palsy include:

- Hypertropia (elevation) in the primary position
- Extorsion that is greatest in abduction
- Compensatory head posture with contralateral face turn and head tilt with chin slightly depressed. This position is assumed to avoid Diplopia.
- Limitation of depression
- Normal abduction and elevation.

The below given image depicts 4th cranial nerve palsy.



Option B: Lateral rectus palsy is seen in cases of isolated Abducens nerve palsy. These patients present with esotropia/convergent squint in primary position with limitation of abduction on the

side of the lesion. This is due to unopposed action of the medial rectus muscle in the affected eye.

There is also a compensatory head turn towards the side of the lesion to reduce the symptom of Diplopia. The adduction is normal in lateral rectus palsy.



Option C: Internuclear opthalmoplegia occurs following an injury to the medial longitudinal fasciculus (MLF). A unilateral lesion to the MLF leads to the failure of adduction in the ipsilateral eye with a normal abduction of the contralateral eye.

The below given image depicts the pathogenesis of Inter nuclear opthalmoplegia in detail.





Solution to Question 18:

The given clinical vignette of a patient presenting with decreased visual acuity and a history of transverse myelitis which is treated with steroids is indicative of Neuromyelitis optica (NMO).

Neuromyelitis optica also known as Devic's disease is an autoimmune demyelinating disease in which antibodies are formed against Aquaporin 4 present in the cell membranes of Astrocytes. It is characterized by the findings of unilateral or bilateral optic neuritis developing within a few weeks of transverse myelitis(demyelination of the spinal cord). The disease may be monophasic or recurrent.

Diagnostic Criteria for Neuromyelitis Optica Spectrum Disorder (NMOSD):

Diagnostic Criteria for NMOSD associated with AQP4-IgG

• At least 1 core clinical characteristic

• A positive test for AQP4-IgG using the best available detection method (cell-based assay)

• Exclusion of alternative diagnoses

Core Clinical Characteristics:

• Optic neuritis is usually bilateral and causes severe visual loss.

• Acute myelitis- may be severe, transverse and typically longitudinally extensive involving three or more contiguous vertebral segments.

• Area postrema syndrome: episodes of otherwise unexplained hiccups or nausea or vomiting

• Acute brainstem syndrome

• Symptomatic narcolepsy associated with NMOSD-typical diencephalic MRI lesions

For acute cases of NMO, high-dose glucocorticoids or plasma exchange is used. Mycophenolate mofetil, rituximab, and glucocorticoids with azathioprine are used for prophylaxis against relapses.

The FDA has approved satralizumab, a humanized IgG2 monoclonal antibody, for the treatment of NMO spectrum disorder with AQP4 antibody positivity.

Eculizumab, a complement inhibitor, and inebilizumab, a B-cell depletion agent, are two more new drugs that have recently been approved for use in NMO attack prevention.

A history of transverse myelitis with optic neuritis(decreased visual acuity) rules out Multiple sclerosis(option D). It is an autoimmune inflammatory and demyelinating disease of the CNS that affects both upper motor neurons (UMN) and lower motor neurons (LMN) with involvement of sensory system, ocular system and bowel and bladder function. Ocular manifestations of MS include Optic neuritis(usually retrobulbar)- decreased visual acuity, internuclear ophthalmoplegia and nystagmus.

Solution to Question 19:

In internuclear ophthalmoplegia (INO) the lesion is at the medial longitudinal fasciculus (MLF).

In INO, there is impaired adduction on the ipsilateral side of the lesion, and nystagmus on abduction on the contralateral side. When the gaze is initiated in the motor cortex, the horizontal gaze center or the paramedian pontine reticular formation (PPRF) generates the eye movements.

The PPRF has two outputs. The first is to the ipsilateral 6th cranial nerve nucleus which causes abduction of the ipsilateral eye. The second output is to the contralateral MLF which sends an impulse to the contralateral 3rd cranial nerve nucleus causing contralateral adduction. These two outputs from PPRF lead to conjugate movement of the eyes to the same side.

Therefore, a lesion in the right MLF leads to failure of the right eye adduction. The left eye, however, abducts normally. This is called internuclear ophthalmoplegia.

The following image shows the pathway for horizontal eye movements as described above.



Anatomical pathway for horizontal eye movements



PPRF – Paramedian pontine reticular formation
FEF – Frontal eye field
MLF – Medial longitudinal fasciculus
MR – Medial rectus
LR – Lateral rectus

©MARROW

Ophthalmology AIIMS 2019 (May & Nov)

Question 1:

A patient presented with painless decrease in vision and metamorphopsia. Fundus examination revealed the lesion shown in the image. What is the expected finding on angiography?



a) Smoke-stack appearance

b) Inkblot appearance

c) Collarstud appearance

d) Double circulation

Question 2:

The test given below in the image is used for:

a) Ocular deviation

b) Stereopsis test

c) Refractive errors

d) Colour Blindness

Question 3:

A patient presents to the emergency and his ocular examination reveals the following. Which of the following is not done initially?



a) Removal of the foreign body

b) Check for visual acuity

c) Give antibiotics

d) Do primary survey

Question 4:

An Intraocular lens was placed uneventfully in a young patient during cataract surgery. What should be done then?

a) IOL replaced every 10 years

b) Never replaced

c) Replaced when secondary cataract develops

d) Replaced when presbyopia develops

Question 5:

What is the first step in the management of an 18-month-old child with both eyes deviated medially:

a) Forced ductions test

b) Examination under anaesthesia

c) Refractive error testing

d) Examination of fundus

Question 6:

Which of the following is correct regarding the image given below:



a) It is called Argyl Robertson pupil

b) Both eyes are abnormal

c) The vision is normal in right eye

d) Seen in optic neuritis, the pupil can paradoxically dilate in the both eyes

Question 7:

A person is diagnosed to have diabetes on his 45th birthday. When will you recommend a dilated fundus examination for him?

a) Immediately

b) Before his 50th birthday

c) After his 50th birthday

d) When he complains of decrease in vision

Question 8:

ETDRS study was related to?

a) Central retinal artery occlusion

b) Diabetic macular edema

c) Primary open angle glaucoma

d) Non-infectious uveitis

Question 9:

A patient with a history of diabetes and hypertension underwent fundus fluorescent angiography (FFA) to evaluate his decreasing vision. The FFA yields an image as shown below, which is characteristic of:



a) Macular ischemia

b) Macular edema

c) Submacular haemorrhage

d) Premacular haemorrhage

Question 10:

Causes for expanding blind spot are all except _____

a) POAG

b) Hypoplasia of optic disc

c) Papilledema

d) Medullated nerve fibres

Question 11:

Identify the structure marked with red arrow



a) Outer limiting membrane

b) Inner plexiform layer

c) Outer plexiform layer

d) Outer nuclear layer

Answer Key

Question No.	Correct Option
1	d
2	b
3	a
4	b
5	c
6	d
7	a
8	b
9	a
10	b
11	c

Detailed Explanations

Solution to Question 1:

The clinical vigenette is of a patient presenting with painless dimunition of vision and metamorphopsia with the right fundus showing choroidal melanoma superotemporal to the optic disc and involving the macula.

In this patient a “Double circulation” pattern consisting of an internal circulation within the lesion and the normal vascularity of the overlying retina will be seen on angiography.

Although collar stud(Option C) appearance is pathagnomic for choroidal melanoma, it can be seen on B-scan and not on angiography.

Choroidal melanoma is a primary intraocular malignancy in adults that peaks around 60 years of age.

Symptoms include a range of visual disturbances such as painless decrease in vision and metamorphosia when macula is involved.

Other features of Choroidal melanoma includes:

• A solitary elevated subretinal grey-brown dome shaped mass

• Clumps of orange pigment are common over the mass

• Collar stud appearance on B-scan (option C): It is pathagnomic of choridal melanoma. It is seen if the tumour breaks through the Bruch's membrane into the subretinal space .

• Associated haemorrhage

Smoke stack pattern (option A) and ink blot (option B) pattern is seen on FFA in central serous retinopathy (CSR).

CSR is the spontaneous serous detachment of the sensory neuroretina in the foveal region with or without retinal pigment epithelium (RPE) detachment.

This occurs due Choroidal vascular hyperpermeability.

On FFA it is initially seen as an ink-blot pattern or smoke stack pattern
of hyperfluorescence which then spreads to take the configuration
of mushroom or umbrella pattern.

The below given images depict the smoke stack pattern and mushroom pattern respectively





Solution to Question 2:

The given image is of Titmus stereotest, which is done for the assesment of stereopsis (perception of depth).

• The pictures consist of a housefly for children and circular patterns for adults.

• The picture is held in front of the eyes of the patient at a distance of about 16 inches and the patient is asked to wear polarized glasses.

• If the patient's perception of the depth is normal, the housefly appears to rise above the plane, like a 3D image. If the patient is asked to pinch the wings of the housefly, he/she will try to pinch in the air, above the plane of the image.

• If the patient's perception of depth in either eye is defective, then the image appears just like normal 2D image.

Option A: Tests for ocular deviation include

• Cover tests

• Prism cover test- This will tell the amount of deviation in prism dioptres.

• Maddox rod test

• Hirschberg corneal reflex test

The below given image shows Maddox rod test.



Option C: Test for refractive error includes

• Retinoscopy

• Auto refractometry

The below given image shows the findings of Retinoscopy in different refractive errors.



Option D: Tests for colour blindness include

• Ishihara test: It is frequently used to screen for a red–green colour vision deficit. It consists of a test plate followed by 16 plates, each with a series of dots arranged to show a shape or number that the subject is asked to identify.

• Hardy–Rand–Rittler test: It is similar to ishihara test but is used to detect all three color vision defect.

• Farnsworth–Munsell 100-hue test: Is the most sensitive test for both acquired and congenital colour blindness. It consists of 85 caps of different hue that needs to be rearranged.

The below given image depicts the ishihara test.



Solution to Question 3:

The above image shows penetrating injury by a metallic fish-hook.

In cases of penetrating injuries, the foreign body is removed only after ascerting it's location through a CT scan.

Hence, Removal of the foreign body is not done initially in these injuries. It is surgically explored & repaired under OT settings.

Management of penetrating injuries of the eye include:

• Primary survey (Option D) of the patient and the wound

• Visual acuity assesment (Option B) in both normal and injured eye

• A CT scan is performed to localize the foreign body and detect any intaorbital or intracranial extensions.

• MRI is contraindicated for metallic foreign bodies.

• Prophylactic intravitreal antibiotics (option C) are prescribed

Solution to Question 4:

Uneventful intraocular lens placement, is not an indication for replacing the IOL.

An IOL replacement (IOL exchange) may be done under the following conditions:

• Incorrect IOL placed

• Posterior or anterior dislocation of IOL

• Hyphema

• Asymmetrical Haptic Placement

• Sunset Syndrome

• Lens-Bag Decentration

Option C: In the event that secondary catarct develops, it is treated by surgical or laser (Nd:YaG) posterior capsulotomy.

Option D: Presbyopia is a disorder of accomdation seen in old age. It is usually corrected by prescription unifocal or Bifocal or multifocal glasses or contact lenses.

Solution to Question 5:

In children, who present with a convergent squint, the first step is to check the visual acuity and estimate the error of refraction.

Convergent squint also known as Esotrpia is commonly seen in children with hyperopia (also known as hypermetropia) of which the accomdative type of esotropia is more common in the age group below 3 years.

The first line of management in the treatment of esotropia includes refractive error correction with glasses.

If a child with accomdative issues, is unable to use spectacles (such as a child ≤ 2 years) or is non compliant with wearing glasses we can eliminate accommodation by temporarily treating him with miotic drugs such as echothiophate.

Further treatment may include orthoptic training with specific exercises or surgical correction when the angle of squint is 10° or more with correcting lenses and when orthoptic training doesnt help in reasonable correction of the squint.

Mnemonic:

The algorithm for the management of accommodative esotropia is as follows:

• R: Refractive correction: Glasses/Miotics

• O: Orthoptics/Occlusion: for amblyopia

• O: Operative correction: for non-accommodative component

• P: Prism correction: for remaining error

Classification of Accomdative esotropia:

The below image shows accomdative esotropia before and after correction.

Type	Criteria
Refractive	Associated withhigh hyperme tropia(+4 to +7 D)Typically e sotropia is for both near and distance.Fullycorrectablebysp ectacles

Type	Criteria
Non-refractive	Occurs independent of refractive errorsAbnormal accommodative convergence/accommodation (AC/A)ratioTypically esotropia isgreater for near than for distanceFully corrected bybifocal glasses

Option A: Forced Duction Test (FDT) is performed to differentiate between the paralytic and restrictive type of inconcomitant squint.

FDT is positive in restrictive squint where a resistance is encountered during passive rotation.

Solution to Question 6:

The given image of "swinging flashlight test" shows both pupils constricting when the left eye is stimulated (normal response) followed by both pupils abnormally dilating when the right eye is stimulated (abnormal response).

This finding is suggestive of Marcus Gunn pupil which is also known as Relative afferent pupillary defect (RAPD).

In this patient the right eye is defective (Option A) and the left eye is normal (rules out option B).

Marcus Gunn pupil refers to the paradoxical response of the pupil to light and it is the earliest sign of optic nerve disease such as optic neuritis or optic nerve damage in glaucoma.It is caused due to a defect in the afferent pathway (optic nerve) of the pupillary light reflex in one eye.

Normal response to stimulation of the eye is constriction of the pupil of the stimulated eye (direct light reflex) and also of the contralateral eye (indirect or consensual light reflex).

Causes of RAPD:

• Optic nerve lesions (e.g.All types of optic neuritis, glaucoma)

• Optic tract lesions (e.g., infarction)

• Severe retinal disease (e.g., central retinal artery occlusion, extensive retinal detachment)

• Maculopathy (macular degeneration)

The below given image depicts the findings seen in Marcus Gunn pupil (RAPD).



Option A:Argyl Robertson pupil is seen in neurosyphilis. In this condition,on light stimulation both the pupils will not constrict, but on accommodation (near target) both the pupils will constrict.

Argyll Robertson pupil is attributed to a dorsal midbrain lesion of neurosyphilis that affects the pupillary light pathway but spares the more ventrally placed pupillary near reflex pathway(Accomdation pathway), thereby causing a light-near dissociation.

Features of Argyll Robertson pupil (ARP) include:

• Bilateral small, irregular pupils in dim light

• Accommodation Reflex Present (ARP)- The eye accomdates by constricting the pupil for a near target.

• Light Reflex absent: pupils do not constrict when exposed to bright light.

• Cocaine eye drops(mydriatics) induced mydriasis.

Solution to Question 7:

A 45 year old diagnosed with diabetes mellitus is likely to have type 2 diabetes and thus, he should undergo screening for diabetic retinopathy immediately.

Duration of diabetes is the most important risk factor for developing diabetic retinopathy, especially proliferative diabetic retinopathy. The recommendations for first fundus examination in diabetic retinopathy is:

• Within 3-5 years of diagnosis in type 1 diabetes

• Immediately in type 2 diabetes.

Later, the screening is continued once every year in both Type 1 and Type 2 diabetes. In the event proliferative diabetic retinopathy develops, screening is done more frequently - once every 2 months.

Solution to Question 8:

Early Treatment Diabetic Retinopathy Study (ETDRS) was a randomized clinical trial that studied the effect of photocoagulation on diabetic macular edema.

The results of the study demonstrated the beneficial effects of focal photocoagulation on diabetic macular edema. The study resulted in the recommendation that all eyes with clinically significant diabetic macular edema should be considered for focal photocoagulation.

Solution to Question 9:

The fundus fluorescent angiography (FFA) image shows enlarged foveal avascular zone, seen as a hypofluorescent patch in the macula (white arrow), with surrounding capillary non perfused areas (blue arrow). Both of these are suggestive of macular ischemia.



The above mentioned findings when combined with the history of diabetes in this patient points towards the diagnosis of diabetic macular ischemia.

FFA images in a case of diabetic retinopathy are given below:

• Non proliferative diabetic retinopathy (NPDR)

• Punctate or pinpoint areas of hyperfluorescence - microaneurysms

• Abnormal branching of arteriovenous shunt - intraretinal microvascular abnormalities (IRMA)

• Patchy filling defects of ischemia due to capillary atrophy or non perfusion

• Increase in the size of hypoperfusion in foveal avascular zone in macular ischemia

2. Proliferative diabetic retinopathy(PDR)

• Fluorescein leakage - in retinal neovascularization.

• Leakage in the disc region suggests neovascularization of the disc (NVD) and leakage in other regions suggests neovascularization elsewhere (NVE).

The image below shows pinpoint areas of hyperfluorescence and areas of fluorescein leakage in the fundus.



Option B: Macular edema also known as cystoid macular edema (CME) is a collection of fluid between the outer plexiform layer and the inner nuclear layer of the retina following a break in the blood-retina barrier. On fundoscopy, a honeycomb appearance is seen, and on FFA, flower petal pattern can be seen. The below image shows the flower petal pattern on FFA as seen in CME.



Option C: Submacular hemorrhage also known as subretinal hemorrhage will obscure the choroidal dye visualization and appaers as ill defined areas of hypoperfusion in the corresponding area. The retinal vessels overlying the area would, however will be visualized. There are no assosciated features of pinpoint hyperfluorescence and vascular leakage that are usually seen with diabetic macular ischemia. The below given images show fluorescein angiography which reveal blocked fluorescence in the maculae of both eyes due to subretinal hemorrhages.



Option D: Premacular hemorrhage is a term used to define vitreous hemorrhage that occurs in the subhyaloid space. It is seen as a region of hypofluorescence with no overlying retinal vasculature visualization on FFA.



Solution to Question 10:

Optic disc hypoplasia does not cause enlargement of the blind spot.

The physiological blind spot is an area of absolute scotoma in the visual field, seen in the region of optic disc. It is due to the absence of photoreceptors (rods and cones) in that region. Enlarged blind spot is seen in primary open-angle glaucoma (POAG) (Option A), papilledema (Option C), medullated nerve fibers (Option D), optic disc drusen, coloboma of optic disc and

myopic disc with a crescent.

Optic nerve hypoplasia is the most common congenital optic disc anomaly. There is underdevelopment of the optic disc in the disease and it typically presents with a small disc that is surrounded by a yellowish pigmented ring, called the 'double ring sign'.

Option A: In primary open-angle glaucoma(POAG), the optic disc enlargement is seen secondary to increased intra-ocular pressure (IOP) in the eye.

Option C: Papilledema is a passive disc swelling seen with raised intracranial pressure. The blind spot is enlarged due to the edema of the optic disc.

Option D: Medullated or opaque nerve fibers are the abnormal myelination of nerve fibers of the retina. Normally, optic nerve myelination stops at the level of lamina cribrosa. If the myelination continues beyond the optic disc it leads to the expansion of the blind spot. On ophthalmoscopic examination, these are seen as a whitish patch with feathery margins.

Solution to Question 11:

The marked structure in the slide is the outer plexiform layer of the retina.

The outer plexiform layer (outer synaptic layer) is present between the outer nuclear layer and the inner nuclear layer. This layer stains lightly. This layer consists of the axons of rods and cones which synapse with dendrites of bipolar neurons and horizontal cells.



Ophthalmology AIIMS 2020 (May and Nov INI-CET)

Question 1:

A patient presents with left-sided head tilt which on straightening leads to right hypotropia. This increases on dextroversion and right head tilt. Paralysis of which of the following muscles is involved?

a) Right superior oblique

b) Right superior rectus

c) Right inferior oblique

d) Left inferior oblique

Question 2:

Which of the following procedure is shown in the image given below?



a) Nasolacrimal duct syringing

b) Anterior ethmoidal nerve block

c) Infraorbital nerve block

d) Trephination of frontal sinus

Question 3:

What is the correct order of using eye drops and eye ointment?

a) Eye drops first, ointment 15 min later

b) Ointment first, eye drops 15 min later

c) Eyedrops first, then ointment immediately

d) Ointment first, then eyedrops immediately

Question 4:

Which of the following statements is false about corneal epithelium?

a) Regeneration is limited to the limbus

b) Bowman's membrane regenerates

c) Lined by stratified squamous epithelium

d) Apical cells have microvilli

Question 5:

A patient underwent LASIK for myopia and now is undergoing cataract surgery. What is the formula used for lens power calculation in this patient?

a) SRK 1

b) SRK 2

c) Hoffer Q

d) Haigis L

Question 6:

Which of the following pupillary abnormalities is most likely to be observed in a patient with primary open-angle glaucoma?

a) Marcus Gunn pupil

b) Adie's tonic pupil

c) Argyll-Robertson pupil

d) Hutchinson's pupil

Question 7:

Which of the following helps to differentiate a fixed dilated pupil? A. Phenylephrine B. 1% Pilocarpine C. 0.1% Pilocarpine D. Cocaine 0.5% E. Epinephrine

a) A, B, C, D and E

b) B and C only

c) A , D and E only

d) A, C and D

Question 8:

True regarding POAG:

a) A,B, D and E

b) A, B and C

c) B and E only

d) B, C and D

Answer Key

Question No.	Correct Option
1	b
2	b
3	a
4	b
5	d
6	a
7	b
8	c

Detailed Explanations

Solution to Question 1:

The given clinical scenario is suggestive of right superior rectus palsy.

The Park's-Bielschowsky three-step test is employed to diagnose the muscle paralysis responsible for vertical eye deviation. In this clinical scenario:

•

Identify the hypotropic eye in primary gaze

•

Right eye is hypotrophic.

•

This could be due to either right eye elevator weakness or left eye depressor weakness.

•

Possible muscles: Right superior rectus, right inferior oblique, left inferior rectus, or left superior oblique

•

Observe if hypotropia worsens in specific gazes

•

Hypotropia worsens in right gaze (dextroversion).

•

Paralyzed muscle is one that elevates the right eye in right gaze or depresses the left eye in right gaze.

•

Possible muscles: Left superior oblique and right superior rectus.

•

Determine if hypotropia worsens in head tilts

•

Hypotropia worsens during right head tilt.

•

The right superior rectus is the muscle involved.

Mnemonic for superior oblique paralysis:

• Step 2: Vertical deviation (hypertropia/hypotropia) is Worse On Opposite Gaze – WOOG

• Step 3: Vertical deviation (hypertropia/hypotropia) deviation is Better On Opposite Tilt – BOOT

Muscle	Primary action	Secondary action	Tertiary action
Medial rectus (MR)	Adduction	-	-
Lateral rectus (LR)	Abduction	-	-
Superior rectus (SR)	Elevation(bestwhen the eye is inabduction)	Intorsion (mnemonic: SIN)	Adduction (mnemonic: R Add)

Muscle	Primary action	Secondary action	Tertiary action
Inferior rectus (IR)	Depression(bestwhen the eye is inabduction)	Extorsion	Adduction (mnemonic: R Add)
Superior oblique (SO)	Intorsion (mnemonic: SIN)	Depression(bestwhen eye is inadduction)	Abduction
Inferior oblique (IO)	Extorsion	Elevation(bestwhen eye is in adduction)	Abduction

Solution to Question 2:

The procedure shown in the given image is an anterior ethmoidal nerve block.

The anterior ethmoidal nerve supplies the roof of the nasal cavity. It is a branch of the nasociliary nerve which in turn is a branch of the ophthalmic nerve (V1). It is sometimes used in endonasal surgery.

Other options:

Option A: In nasolacrimal duct syringing, the puncta of the nasolacrimal duct near the medial canthus of the eye is probed using a lacrimal cannula. The below image shows lacrimal syringing.

Option C: Infraorbital nerve block causes anesthesia in the distribution of infraorbital nerve which includes lower lid, cheek, side of nose, upper lip, and upper teeth. It is most commonly done for dental procedures.

Option D: Trephination of the frontal sinus is a surgical procedure done for the drainage of pus in acute frontal sinusitis. In this procedure, the incision is made on the floor of the frontal sinus.

Solution to Question 3:

Eye drops first followed by ointment 15 minutes later is the correct way of usage.

An ointment is thicker in consistency and needs to stay on the eyes for a longer time for absorption of the medication. Hence, it is generally put in the eyes at bedtime.

Eyedrops are watery thin and can be absorbed faster. Hence eyedrops are put first, and after 10-15 minutes, once they are completely absorbed, eye ointment can be applied.

Eye ointments are usually avoided during the daytime as they cause marked blurring of vision.

Solution to Question 4:

Bowman's membrane is acellular. Hence, it will not regenerate but heals by scarring when damaged.

Corneal stem cells are located at the corneoscleral limbus (Option A), so the regeneration is limited to the limbus.

The cornea comprises distinct layers:

• Corneal epithelium is lined by non-keratinized stratified squamous epithelium (Option C), with microvilli on the apical layer (Option D) to enhance surface area and tear film attachment. It consists of basal, columnar, wing, and squamous cells.

• Bowman's membrane, a superficial stromal layer, is acellular, collagen-based, and incapable of regeneration.

• The stroma, the thickest layer constituting 90% of corneal thickness, houses collagen fibrils with ground substances such as chondroitin sulfate and keratan sulfate, maintaining transparency. It cannot regenerate and heals with scarring.

• Dua's layer or Pre-Descemet's membrane is the toughest.

• Descemet's membrane extends anteriorly to Schwalbe's line and possesses regenerative potential.

• Endothelium, a metabolically active monolayer, lacks regenerative ability. It maintains deturgescence through active pumps, ensuring transparency with a usual cell density of approximately 3000 cells/mm² measured by specular microscopy.





Solution to Question 5:

Among the given options, the Haigis-L regression formula is used for calculating the lens power in a patient who is undergoing cataract surgery with a previous history of refractive surgery.

Haigis formula uses Axial Length (L), Corneal curvature (K), and Anterior Chamber Depth (ACD) for the calculation of IOL power (biometry).

• Keratometry measures the curvature of the anterior corneal surface.

• A-scan ultrasonography is used to determine the axial length.

Standard IOL calculations are unsuitable in any type of corneal refractive surgery as they are likely to result in residual refractive errors even after IOL implantation in these patients. In these patients, several other methods of calculating corneal refractive power are used which are then

added to the standard and specific formulas such as the Masket, modified masket, Haigis L formula, and double-K method.

Other options:

Options A and B: SRK 1 and SRK 2 are the basic formulas used for IOL power calculations.

SRK formula: P = A-2.5L-0.9K where P is the power of IOL, A is a constant for the lens, L is the axial length and K is corneal curvature.

Option C: Hoffer Q formula is used for IOL power calculation in short eyes (hypermetropia). It uses only two variables; Axial length (L) and corneal curvature (K).

Solution to Question 6:

Primary open-angle glaucoma commonly causes bilateral and asymmetric optic nerve damage. Marcus Gunn pupil also known as relative afferent pupillary defect (RAPD) is the earliest sign of optic nerve disease as seen in optic nerve neuritis or primary open-angle glaucoma(POAG).

RAPD is caused due to a defect in the afferent pathway (optic nerve) of the pupillary light reflex in one eye.

It is diagnosed by the swinging flashlight test where the flashlight is swung to and fro, and the pupillary response is observed. Normally, both pupils constrict when any one eye is stimulated (due to direct and consensual light reflex). In the presence of RAPD, both pupils will abnormally dilate when the affected eye is stimulated. This paradoxical response of the pupil to light is called Marcus Gunn pupil.

Causes of RAPD:

• Optic nerve lesions (e.g.All types of optic neuritis, glaucoma)

• Optic tract lesions (e.g., infarction)

• Severe retinal disease (e.g., central retinal artery occlusion, extensive retinal detachment)

• Maculopathy (macular degeneration)

Other options:

Option B: Adie tonic pupil is a unilateral dilated pupil with light-near dissociation due to postganglionic parasympathetic pupillomotor nerve supply damage. In these patients the light reflex is absent and the accomdation (near) reflex is sluggish. It is commonly seen in young females and is associated with absent knee jerk.

Option C: Argyll Robertson pupils are bilateral small pupils with light-near dissociation. It is commonly seen in patients with neurosyphilis. In these patients the pupillary reflex is absent but the accommodation reflex is present (ARP).

Option D: Hutchinson's pupil is a clinical sign where the pupil on the side of an intracranial mass lesion is dilated and unreactive to light, due to compression of the oculomotor nerve on that side.

Solution to Question 7:

Among the given options, 1% pilocarpine and 0.1% pilocarpine can be used to differentiate fixed dilated pupil.

Fixed dilated pupil is seen in conditions such as third nerve palsy, Hutchinson's pupil (increased cranial pressure), Adie's pupil, and due to pharmacological mydriasis.

Pilocarpine acts as an agonist at cholinergic muscuranic receptors and ganglion stimulator to cause miosis of the pupil.

Normal strength pilocarpine (1-2%) constricts the dilated pupil of 3rd nerve palsy, whereas there will be no response if the dilation is due to pharmacological mydriasis (anticholinergic causes) such as accidental application of atropine.

0.1% pilocarpine is used to diagnose Adie's pupil. Adie's pupil constricts with weak concentrations of pilocarpine (0.1%), while the normal pupil does not. This is because the denervated sphincter of the iris is very sensitive to topical parasympathomimetics such as pilocarpine. 0.1% pilocarpine also helps differentiate Adie's pupil from Argyll Robertson pupil (ARP).

Note:

Cocaine test (2 to 4%) is done in Horner's syndrome. In these patients, the small pupil fails to dilate on instillation of mydriatics such as Cocaine which is in contrast to the normal pupil which dilates on instilling cocaine. Alternatively, the small pupils of Horner's syndrome dilate with 1% apraclonidine.



Solution to Question 8:

In primary open-angle glaucoma (POAG), family members are likely to develop steroid responsiveness (statement B) and visual field defect can exist with a normal C:D ratio (statement E).

Statement A: Gonioscopy cannot visualize the specific abnormalities that occur in the trabecular meshwork in POAG. Gonioscopy diagnoses POAG because of the visibility of all the structures of the angle of the anterior chamber in contrast to the non-visibility of different structures in angle

closure glaucoma.

Statement C: Primary angle-closure glaucoma is associated with exacerbation of IOP with pupillary dilatation and not POAG.

Statement D: There is a 10% increased risk of POAG in siblings and a 4% increased risk in children of patients with POAG.

POAG also known as chronic simple glaucoma of adult-onset is characterized by the presence of any 2 of the following 3 categories of changes:

• Increased intraocular pressure:

• IOP >21mmHg on more than one occasion, and/or

• Circadian variation in IOP of >8 mmHg, or

• Asymmetry in IOP > 5 mmHg between both eyes

• Optic nerve head changes suggestive of glaucomatous damage. These may include one or more of the following changes:

• Vertically oval cup

• Cup-disc (C:D) ratio > 0.6 (average cup: disc ratio 0.3-0.4)

• Asymmetry in C:D ratio > 0.2 between both eyes.

• Narrowing, notching, or pallor of the neuroretinal rim.

• Disc haemorrhages

• Nasal shifting of retinal vessels-Bayonetting sign

• Lamellar dot sign

• Visual field changes - isopter contraction, barring of blind spot, paracentral scotoma, Seidel's scotoma, Bjerrum's scotoma, double arcuate scotoma, and Roenne's central nasal step.

A patient presenting with just raised IOP and visual field defects with normal optic nerve head findings such as a normal C:D ratio can still be diagnosed as POAG (statement E).

Ophthalmology NEET 2018

Question 1:

Which of the following would be prescribed for simple myopic astigmatism?

a) +1.00 DS

b) -1.00 DC X 180 degree

c) -1.00 DS

d) -1.00 DS -1.00 DC X 180 degree

Question 2:

What is the causative organism of angular conjunctivitis?

a) Moraxella lacunata

b) Haemophilus influenzae

c) Adeno virus type 32

d) Coxackie B virus

Question 3:

What is the most common cause of ophthalmia neonatorum in developed countries?

a) Chlamydia

b) Gonorrhoea

c) Chemical conjunctivitis

d) Staphylococcus aureus

Question 4:

What is the cause of acute hemorrhagic conjunctivitis?

a) Enterovirus 68

b) Enterovirus 69

c) Enterovirus 70

d) Enterovirus 71

Question 5:

100-day glaucoma is seen in which of the following conditions?

a) Central retinal vein occlusion

b) Eale's disease

c) Central retinal artery occlusion

d) Steroid-induced glaucoma

Question 6:

Yoke muscle of the right lateral rectus in dextroversion is the _____.

a) Left superior oblique

b) Left medial rectus

c) Left inferior rectus

d) Left superior rectus

Question 7:

A patient, 36 hours after cataract surgery, presents with complaints of pain and visual disturbance. His eye is as shown below. What is the management of this condition?

a) Intravenous steroids

b) Intravenous antibiotics

c) Bandage and padding of the eye

d) Intravitreal antibiotics

Question 8:

Which of the following is the most common feature of third nerve palsy, in a case of long-standing, uncontrolled diabetes?

a) Proptosis

b) Pseudoptosis

c) Ptosis

d) Pupillary dysfunction

Question 9:

What is the most common site of an orbital blowout fracture?

a) Medial wall

b) Lateral wall

c) Roof

d) Floor

Question 10:

A male patient came with a history of unprotected sexual intercourse around 20 years ago. Now, he complains of neurological symptoms and visual disturbances. Which of the following will be seen on exposure of his pupils to light?

a) Pupils react but do not accommodate.

b) Pupils accommodate but do not react.

c) Pupils neither react not accommodate.

d) Pupils react and accommodate.

Question 11:

Roth spots are seen in _____.

a) Acute leukemia

b) Uveal melanoma

c) Uveal lymphoma

d) All of the above

Question 12:

A 50-year-old man is presbyopic. The power of his prescription glasses should be _____.

a) +1.75D

b) +1.5D

c) +2D

d) +2.25D

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	c
5	a
6	b
7	d
8	c
9	d
10	b
11	a
12	b

Detailed Explanations

Solution to Question 1:

Among the above options, -1.00 DC × 180 degrees is the correct prescription for simple myopic astigmatism (SMA).

There are 3 components for prescription of glasses.

1. The first number is the refractive power of the spherical component. It is represented as convex (+) or concave (-) dioptre sphere (DS).

2. The second number denotes the power of the cylinder, represented as convex (+) or concave (-) dioptre cylinder (DC).

3. The above two components are followed by an x representing the axis (orientation) of the cylinder.



©MARROW



©MARROW

In the most common type of SMA - with-the-rule astigmatism - meridian A (represented by 90 degrees) is more curved than meridian B (represented by 180 degrees). The light rays refracting

from B focus on the retina, whereas those refracting from A focus in front of the retina.

To correct this error, a negative cylindrical lens (DC) is used to focus the myopic rays of meridian A on the retina. This correction needs to occur at 90 degrees (meridian A). The axis of a cylindrical lens is perpendicular to the required correction and thus, the spectacles will have an axis of 180 degrees.

There is no spherical refractive error since the light rays from meridian B lies on the retina. Thus DS is 0.

Solution to Question 2:

Moraxella species, a type of gram-negative diplobacilli, are involved in causing angular conjunctivitis which is a type of chronic conjunctivitis.

Angular conjunctivitis is a chronic type of conjunctivitis where there is inflammation of the conjunctiva at the lid margins and the inner and outer canthi of the eyes with red, scaly, excoriated, macerated and fissured skin at the canthi. This is because of the proteolytic enzymes secreted by the Moraxella species.

Patients with angular conjunctivitis usually experience eye irritation, a burning sensation and collection of dirty-white, foamy discharge at the angles. These patients can also have associated blepharitis and papillary and follicular conjunctivitis.

Causative organisms are Moraxella lacunata, Moraxella axenfeld and Staphylococcus aureus (Rare cause). The source of infection is typically the nasal tract or the nasal discharge of patients with angular conjunctivitis. The infection spreads from the nasal cavity to the eyes through contaminated hands or handkerchiefs. Treatment includes oxytetracycline 1% ointment for 10 to 14 days, and zinc oxide as it inhibits proteolytic enzymes.

The image given below shows the eye of a patient infected with angular conjunctivitis.

Other options:

Option B: Haemophilus influenzae causes acute conjunctivitis.

Option C: Adenovirus is the most common cause of viral conjunctivitis and it most commonly causes follicular conjunctivitis, pharyngoconjunctival fever, and epidemic keratoconjunctivitis.

Option D: Coxsackie B virus is known to cause aseptic meningitis, encephalitis, pericarditis, and myocarditis. There are no known mentions of ocular disease caused by the virus.

Note: Pathogens that cause bacterial conjunctivitis

The below-given table elaborates on the type of conjunctivitis caused by different serotypes of adenovirus.

Hyperacute(symptoms develop within 24 hours; aggressive course)	Acute(resolving in less than 4 weeks)	Chronic(of greater than 4 weeks duration)
Neisseria gonorrhoeae	Staphylococcus aureus	Staphylococcus aureus
Neisseria meningitidis	Streptococcus pneumoniae	Moraxella lacunata
	Haemophilus influenzae	Enteric bacteria

Conjunctivitis type	Adenovirus serotype
Follicular Conjunctivitis	Adenovirus serotypes 1-11 and 19
Pharyngoconjunctival Fever	Adenovirus serotypes 3, 4, and 7
Epidemic Keratoconjunctivitis	Adenovirus serotypes 8, 19, and 37

Solution to Question 3:

Chlamydia is at present the most common cause of ophthalmia neonatarum in developed countries.

Ophthalmia neonatarum also called neonatal conjunctivitis is inflammation of the conjunctiva occurring in an infant <30 days after birth.It should be ruled out in any infant having discharge or watering within the first week of birth.

Causes include:

• Chlamydia(Option A): neonatal inclusion conjunctivitis caused by serotypes D-K is at present the most common cause in developed countries.

• Gonorrhoea(Option B): effective prophylaxis against gonococcal infection and its subsequent decline has almost eliminated it's incidence in developed countries.

• Staphylococcus aureus(Option D), Haemophilus species, Streptococcus haemolyticus, and Streptococcus pneumoniae are other less commonly seen causes of ophthalmia neonatarum

• Herpes simplex ophthalmia neonatrum rarely occurs and is caused by Herpes simplex type 2 virus from the mother.

• Chemical conjunctivitis(Option C) is caused by the application of prophylactic antibiotics or silver nitrate at the time of birth.

Time of presentation	Causative agent
Within 48 h	Chemical, Gonorrhoea
48–72 h	Staphylococcus aureus,Streptococcus pneumoniae
5–7 days	Herpes simplex II
>1 week	Chlamydia

Prophylactic treatment: 1% tetracycline or erythromycin eye ointment is applied to the eyes, if the mother is suspected to have gonococcal or chlamydia infection.

Curative treatment:

• Chlamydia trachomatis (D– K)- Erythromycin eye ointment

• Neisseria gonorrhoeae- Ceftriaxone injection, gentamicin drops, bacitracin eye ointment

The below image shows an infant presenting with ophthalmia neonatarum.

Solution to Question 4:

Acute hemorrhagic conjunctivitis is caused by picornoviruses namely enterovirus 70.

Acute hemorrhagic conjunctivitis also known as "Apollo conjunctivitis" is caused by picornoviruses such as enterovirus 70 and coxsackievirus A24. Rarely it is also caused by adenovirus type 11.

It manifests with severe conjunctival congestion, chemosis, subconjunctival hemorrhages, painful follicular conjunctivitis, tearing, lid edema and pre-auricular lymphadenopathy. The cornea may demonstrate a fine punctate keratopathy. The disease is usually self-limiting, running a course of 3-7 days.

The image given below shows the eyes of a patient with acute hemorrhagic conjunctivitis.



Enterovirus 71 (Option D) is associated with fatal meningitis, encephalitis, and paralysis. It is also implicated in Hand-Foot-Mouth disease(HFMD) and Herpangia which is papulo-vesiculo-ulcerative rash in oral cavity.

Solution to Question 5:

100-day glaucoma is a neovascular glaucoma that occurs in ischemic type of central retinal vein occlusion (CRVO).

It is known as 100-day glaucoma because it occurs 3 months after the episode of CRVO. Although neovascular glaucoma occurs as a complication in Eale's disease (Option B) and central retinal artery occlusion (Option C) and long-term use of topical steroid therapy causes steroid-induced glaucoma (Option D) none of them are termed as 100-day glaucoma.

CRVO is further classified into two types:

• Non-ischemic CRVO: occurs secondary to venous stasis.It is the most common type of CRVO of the two.

• Ischemic CRVO: occurs due to acute occlusion of the central retinal vein. It is also termed as hemorrhagic retinopathy.

Ischemic CRVO presents with sudden loss of vision and Relative afferent pupil defect(RAPD).

Fundus examination shows massive enlargement of retinal veins and massive retinal hemorrhages giving a "splashed tomato appearance". Other findings include cotton wool spots, optic disc edema and neovascularization at the disc and in the periphery.

Pathagnomic features of ischemic CRVO that differentiates it from non-ischemic CRVO are:

• RAPD

• Reduced amplitude of B-wave on electroretinogram

• Visual field defects

Complications of the disease:

• Neovascular Glaucoma known as 100 day glaucoma

• Neovascularisation of iris known as Rubeosis iridis

• Vitreous hemorrhage

• Proliferative retinopathy

The below-given fundus image is of the "splashed tomato" appearance of CRVO.



Option B: Eales disease is an idiopathic inflammatory disease of the peripheral retinal veins.It is considered to be a hypersensitivity reaction to the tubercular proteins and it is commonly seen in young males. It is characterized by recurrent vitreous hemorrhage.

Complications include:

• Neovascular glaucoma

• Proliferative vitreoretinopathy

• Tractional retinal detachment,

• Rubeosis iridis

Option C: Central retinal artery occlusion(CRAO) is most often caused due to an embolus from an atherosclerotic plaque. It presents as sudden painless loss of vision and amaurosis fugax (transient monocular blindness that is due to ischemia). Fundus examination shows an opaque, milky-white retina with a cherry-red spot.

Neovascular glaucoma may occur as a delayed complication of CRAO but it is not termed as 100 day glaucoma.

The below image depicts the fundus image seen in CRAO with the cherry red spot.



Option D:Steroid induced glaucoma is a type of secondary open angle glaucoma that usually follows weeks-months of topical steroid therapy.

Solution to Question 6:

The yoke muscle of the Right lateral rectus in right dextroversion is the left medial rectus.

Yoke muscles also known as contralateral synergists include a pair of muscles, one from each eye, which contract simultaneously during version movements.

Versions are simultaneous movements of both eyes in the same direction, such as dextroversion in which both the eyes move to the right.

Hence, in dextroversion, the right eye is in abduction due to the action of right lateral rectus and the left eye is in adduction by the action of the left medial rectus.

An easy method to determine yoke muscle pairs:

Just take the opposite of each word-

Left $\rightarrow$ Right

Superior $\rightarrow$ Inferior

Oblique $\rightarrow$ Rectus

“Left SO and Right IR” are a yoke muscle pair.

Other examples of yoke muscles are:

• Right MR and Left LR for levoversion

• Right IO and Left SR - Levoelevation

• Right SR and Left IO - Dextroelevation

• Right IR and Left SO - Dextrodepression

[IO-inferior oblique;SO-superior oblique;SR-superior rectus;IR-inferior rectus;MR-medial rectus;LR-lateral rectus]



Solution to Question 7:

A post cataract surgery patient presenting within 36 hours of surgery with inflammation, congestion and hypopyon as seen in the image, is suggestive of acute bacterial endophthalmitis. The treatment of acute bacterial endophthalmitis is intravitreal antibiotics.

Endophthalmitis is an intraocular inflammation affecting the inner structures of the eyeball. Acute postoperative bacterial endophthalmitis occurs within 1-7 days after cataract surgery and the most important factor to prevent it is preoperative administration of 5% povidone iodine. The cardinal features of endophthalmitis are pain, swelling of the lid, and a decrease in vision. The patient classically shows inflammation of the eye, corneal edema and a hypopyon due to the pouring of exudates in the vitreous cavity, anterior chamber, and posterior chamber.

Once the endophthalmitis sets in it is important to treat all the cases empirically with a combination of antibiotics and steroids from the beginning. The various drugs, routes and surgical management used to treat endophthalmitis are as follows:

• The mainstay of treatment is intravitreal injections of antibiotics.

• First choice: Vancomycin 1 mg in 0.1 mL plus ceftazidime 2.25mg in 0.1ml.

• Second choice:Vancomycin 1 mg in 0.1 mL plus amikacin 0.4mg in 0.1ml.

• If there is no improvement, a repeat intravitreal injection should be given after 48 hours

• Topical antibiotics are started immediately and used frequently.

• Vancomycin (50mg/ml) or cefazolin (50mg/ml)

• Amikacin (20mg/ml) or tobramycin

• Systemic antibiotics have a limited role in the treatment but are commonly used in practice.

• Steroid therapy includes:

• Intravitreal injection of dexamethasone (0.4mg in 0.1ml) which is started along with antibiotics.

• Topical dexamethasone (0.1%)

• Oral steroids such as prednisolone are started after 24 hours of intensive antibiotic therapy.

• Supportive therapy includes:

• Cycloplegics such as atropine 1%

• Anti-glaucoma drugs such as oral acetazolamide or timolol in patients with raised intraocular pressure

• Pars plana vitrectomy is done if the patient does not improve with the above therapy in 48-72 hours or when the visual acuity is reduced to hand movements near the eye

Other options:

Option A: Steroids are given in endophthalmitis to reduce the inflammation of the tissues. The preferred route of administration is either by intravitreal injections, topical concentrations or oral route. Intravenous route (option A) of administration of steroids is not used in endophthalmitis.

Option B: Systemic administration (intravenous) of antibiotics have a limited role in the treatment but are commonly used in practice as an add on.

Solution to Question 8:

The most common feature seen in diabetic oculomotor palsy among the given options is ptosis.

In the anatomical arrangement of the nerve fibers of the 3rd cranial nerve, the fibers controlling the pupillary reflex are present superficially and are thus spared from diabetic ischemia-induced injury which starts from the central nerve fibres. Thus, pupil sparing is commonly seen in diabetic patients.

The etiology of diabetic neuropathy is hyperglycemia-induced damage to nerve cells and neuronal ischemic change due to microvascular angiopathy.

Involvement of the third cranial nerve is most common and it often presents with early features of ptosis and diplopia with pupil sparing.

The involvement of levator palpebrae superioris muscle of the eyelid manifests as ptosis and the involvement of extraocular muscles such as medial rectus,inferior rectus and inferior oblique presents as opthalmoplegia with diplopia. All the muscles mentioned above are supplied by the Occulomotor nerve.

Note: Aneurysm of the posterior communicating artery is an important cause of isolated third nerve palsy with involvement of the pupil. This is because the aneurysm impinges on the superficial pupil nerve fibres of the 3rd cranial nerve.

Solution to Question 9:

The most common site of an orbital blowout fracture is the floor of the orbit.

Blow-out fractures are isolated comminunited fractures caused from direct trauma to the orbit that results in a backward displacement of the eyeball and an increase in the intraorbital pressure. This causes a fracture at the weakest point of the orbital wall which is usually the posteromedial portion of the maxillary bone in the orbital floor.

The second most common site for orbital blowout fracture is the medial wall of the orbit (Option A).

The fractures of the roof(Option C) are relatively rare and most often do not require any treatment.The lateral wall(Option B) of the orbit is the most solid orbital wall and a fracture here is usually seen in injuries with extensive facial damage.

The patient with blow out fracture usually presents with immediate periorbital edema and blood extravasation in and around the orbit. Proptosis of the eyeball secondary to orbital edema or hemorrhage with paresthesia along the infraorbital nerve are also very common.

Imaging by a plain x-ray in water's view or by CT demonstrates depression of bony fragments and orbital contents herniating through the orbital floor into the maxillary antrum. This is called as the "hanging drop sign" or "tear drop sign".

Definitive treatment is by surgical repair 10-14 days after injury with initial supportive management with analgesics and antibiotics.

The below given CT image shows hanging drop sign.



The below given X-ray image in water's view shows "hanging drop sign".

Solution to Question 10:

The patient most likely has acquired syphilis due to unprotected sex and has developed neurological and visual disturbances secondarily as a complication of neurosyphilis. Argyll-Robertson pupils(ARP) are seen associated with neurosyphilis where the pupils accommodate but do not react to light.

Argyll Robertson pupil is attributed to a dorsal midbrain lesion of neurosyphilis that affects the pupillary light pathway but spares the more ventrally placed pupillary near reflex pathway(Accomdation pathway), thereby causing a light-near dissociation.

Features of Argyll Robertson pupil (ARP) include:

• Bilateral small, irregular pupils in dim light

• Accommodation Reflex Present (ARP)- The eye accomdates by constricting the pupil for a near target.

• Light Reflex absent: pupils do not constrict when exposed to bright light.

• Cocaine eye drops(mydriatics) induced mydriasis.

Other clinical features in neurosyphilis:

• Fleeting and repetitive lancinating pains, primarily in the legs

• acute abdominal pain with vomiting(visceral crisis)

• Tabes dorsalis with Ataxia and gait disturbances due to loss of position sense

• Loss of reflexes in the legs

• Paresthesias, bladder disturbances

Other causes of light-near dissociation are:

In unilateral eye

• Adie pupil

• Herpes zoster ophthalmicus

• Aberrant regeneration of the third cranial nerve

In bilateral eyes

• Neurosyphilis

• Type 1 diabetes mellitus

• Myotonic dystrophy

Note: Adie's pupil more commonly manifests in young women with unilateral large regular pupil(Anisocoria) with absent light reflex and sluggish and tonic accomdation reflex.

It is caused by post-ganglionic parasympathetic pupillary supply damage and is usually associated with absent knee jerk.

Adie's pupil constricts with weak pilocarpine (0.1%) drops, while normal pupil and Argyll robertson pupil does not.

Solution to Question 11:

Roth's spots are seen in acute leukemia.

Roth's spots are retinal hemorrhages with white or pale center. The white centre due to coagulated fibrin-platelet plug is surrounded by red rim that is caused by hemorrhage. Roth's spots may be seen in:

• Subacute bacterial endocarditis

• Leukemia

• Diabetes

• Sickle cell retinopathy

• Hypertensive retinopathy

• HIV retinopathy

• Systemic lupus erythematosus

The fundus image below shows Roth's spots.



Option B: Uveal melanoma devolps due to malignant transformation of the normal melanocytes present in the uveal tissue. Primary posterior uveal melanoma (choroidal melanoma) which forms the majority of uveal melanoma manifests as:

• A solitary elevated subretinal grey-brown dome shaped mass

• "Collar stud" appearance on B-scan - when the tumor breaks through the Bruchs membrane and protrudes into subretinal space

• "Double circulation" sign

Primary anterior uveal melanoma frequently involves the iris and presents as nodular melanocytic mass on the iris. The below given fundus image shows the "double circulation sign" of choroidal melanoma.



Option C: Uveal lymphoma is characterized by localized or diffuse uveal infiltrates of malignant lymphoma cells. Fundus examination in these patients show diffuse creamy thickening of posterior uvea with nodular accentuation.

Solution to Question 12:

The power of prescription glasses for a 50-year-old woman should be +1.5D.

Presbyopia is the condition of decreasing near vision due to an age-related decrease in the magnitude of accommodation.

Basic principles for correction in presbyopic patients are given below:

• Correct any refractive error for distance present in the patient first.

• Add the presbyopic correction(correction for near vision) of each eye to any distant correction that is present.

• For presbyopic correction, the weakest convex lens(plus lens) with which the patient can see clearly at the near point is prescribed, since over-correction will also result in eye fatigue symptoms.

Presbyopic spectacles may be unifocal, bifocal or Multifocal.

A rough prescription of presbyopic glasses can be made by the age of the patient in an emmetrope as follows:

Age	Power
45 years	+1 to +1.25 D
50 years	+1.5 to 1.75 D
55 years	+2 to +2.25 D
60 years	+2.5 to +3D

Ophthalmology NEET 2019

Question 1:

Which of the following would be prescribed for simple myopic astigmatism?

a) +1.00 DS

b) -1.00 DC X 180 degree

c) -1.00 DS

d) -1.00 DS -1.00 DC X 180 degree

Question 2:

Treatment of ptosis caused by Horner's syndrome is:

a) Levator resection

b) Frontalis sling

c) Fasanella servat operation

d) None of the above

Question 3:

Which of the following is not true about retinitis pigmentosa?

a) Retinal pigmentation

b) Pale waxy disc

c) Narrowing of vessels

d) ERG – normal

Question 4:

A 45-year-old patient comes with complaints of progressive weakness as the day progresses. He is found to have antibodies against nicotinic acetylcholine receptors. Which is the most common ocular finding in this disease?

a) Diplopia

b) Lagophthalmos

c) Exophthalmos

d) Ptosis

Question 5:

A patient with a history of trauma, presented with diplopia, epistaxis, and restricted eye movements. Looking at the CT image given below, what will be your diagnosis?



a) Le- fort fracture

b) Fracture maxilla

c) Fracture zygomatic

d) Blow out fracture

Question 6:

What is the likely diagnosis from the image given below?



a) Pinguecula

b) Concretions

c) Pterygium

d) Bitots spots

Question 7:

Which of the following statements is false regarding accommodative esotropia?

a) It is the most common type of squint.

b) Miotics are used as the treatment strategy in the high AC/A ratio

c) It is associated with high myopia.

d) Refractive type can be fully corrected by use of spectacles.

Question 8:

Which of the following regimen is commonly used■ in the treatment of retinoblastoma?

a) Vinblastine, etoposide and bleomycin

b) Vinblastine, vincristine and etoposide

c) Vinblastine, vincristine and cisplatin

d) Vincristine, carboplatin and etoposide

Question 9:

A 50-year-old male presented with progressive blurred vision, photophobia, and eye pain. On examination, there was conjunctival redness, a mid-dilated pupil, and a shallow anterior chamber in the right eye. He has been treated both medically and surgically for the condition but showed no improvement. The following procedure incorporates a device to aid drainage. Identify the device.



a) Ex-Press™ implant

b) Ahmed glaucoma valve

c) Molteno implant

d) Baerveldt implant

Question 10:

Identify the instrument.

a) Maddox wing

b) Maddox glass

c) Maddox rod

d) Phoropter

Question 11:

Identify the test being shown below.



a) Obscure test

b) Prism test

c) Cover uncover test

d) Maddox rod test

Answer Key

Question No.	Correct Option
1	b
2	c
3	d
4	d
5	d
6	c
7	c
8	d
9	a
10	c
11	c

Detailed Explanations

Solution to Question 1:

Among the above options, -1.00 DC × 180 degrees is the correct prescription for simple myopic astigmatism (SMA).

There are 3 components for prescription of glasses.

1. The first number is the refractive power of the spherical component. It is represented as convex (+) or concave (-) dioptre sphere (DS).

2. The second number denotes the power of the cylinder, represented as convex (+) or concave (-) dioptre cylinder (DC).

3. The above two components are followed by an x representing the axis (orientation) of the cylinder.



©MARROW



©MARROW

In the most common type of SMA - with-the-rule astigmatism - meridian A (represented by 90 degrees) is more curved than meridian B (represented by 180 degrees). The light rays refracting from B focus on the retina, whereas those refracting from A focus in front of the retina.

To correct this error, a negative cylindrical lens (DC) is used to focus the myopic rays of meridian A on the retina. This correction needs to occur at 90 degrees (meridian A). The axis of a cylindrical lens is perpendicular to the required correction and thus, the spectacles will have an axis of 180 degrees.

There is no spherical refractive error since the light rays from meridian B lies on the retina. Thus DS is 0.

Solution to Question 2:

Fasanella servat operation is done for cases of mild ptosis (1.5-2 mm) with retained levator palpabrae superioris muscle function as seen in Horner's syndrome.

Ptosis in Horner's syndrome is due to the loss of sympathetic supply to the muller's muscle of the lid and not due to levator muscle palsy. Fasanella servat operation is not done in patients with severe ptosis and poor levtaor function.

Option A:Levator resection is performed for moderate and severe grades of ptosis but with retained levator function. This operation is contraindicated in patients with poor levator function.

Option B: Frontalis sling also known as Brow suspension is done in patients with severe ptosis with poor or absent levator muscle function. Here, the lid is anchored to the frontalis muscle via a sling.

Surgical treatment of various types of ptosis

Surgery	Amount of ptosis that can be treated	Indications
Fasanella Servat	< 2 mm -mild ptosis	Horner Syndrome
Levator Resection– Anterior approach	Any amount (but with retaine d levator function)	Larger resection in ptosis
Levator resection – Conjuncti val approach	Any amount (but with retaine d levator function)	Moderate resection in ptosis
Levator resection with apone urotic reinsertion	Any amount (but with retaine d levator function)	Acquired ptosis
Frontalis suspension– Fascia lata	>4 mm -severe ptosis	Congenital ptosis– Marcus G unn ptosis

Solution to Question 3:

Electroretinogram (ERG) is abnormal in retinitis pigmentosa (RP).

ERG is a sensitive diagnostic test in retinitis pigmentosa that shows reduced scotopic, subnormal responses in early disease and becomes completely extinguished in advanced disease. Electroocoulogram (EOG) is also abnormal in retinitis pigmentosa.

Retinitis pigmentosa (RP) is an inherited or acquired pigmentatory retinal dystrophy affecting the rods more than the cones. The earliest manifestation is worsening of night blindness and delayed adaptation which gradually progresses to tunnel vision and ultimately loss of central vision.

The classic triad of findings in RP comprises:

• Bone-spicule retinal pigmentation

• Arteriolar attenuation

• 'Waxy' disc pallor

The image below shows the classical triad findings of retinitis pigmentosa.



Solution to Question 4:

The given clinical scenario is suggestive of myasthenia gravis and ptosis is the most common ocular feature of the disease.

Myasthenia gravis (MG) is an autoimmune disorder of the neuromuscular junction most commonly caused by anti-acetylcholine receptor antibodies. The disorder is characterized by weakness and fatigability of skeletal muscles where the weakness increases during repeated use and may improve following rest.

Ptosis and diplopia are common initial complaints due to early involvement of the lids and extraocular muscles (EOMs). Of them, Ptosis is a more common manifestation than diplopia. Ptosis in myasthenia gravis is insidious, bilateral, and frequently asymmetrical.

Investigations of myasthenia gravis include:

• Ice pack test -This tests for a transient improvement of the ptotic eyelid on placing an ice pack for 2 minutes, as the cold inhibits the breakdown of acetylcholine by acetylcholinesterase.

• Tensilon (Edrophonium) test - This tests for a transient improvement in the weakness on administration of a short-acting anticholinesterase such as edrophonium.

• Antibody testing - for Acetylcholine receptor (AChR) antibodies, MuSK protein antibodies helps support a diagnosis of myasthenia gravis.

Solution to Question 5:

The given clinical scenario plus the supportive CT image showing the "hanging drop sign" is suggestive of a blowout fracture of the orbital wall.

Blow-out fractures are isolated comminunited fractures caused from direct trauma to the orbit that results in a backward displacement of the eyeball and an increase in the intraorbital pressure.

This causes a fracture at the weakest point of the orbital wall which is usually the posteromedial portion of the maxillary bone in the orbital floor. The second most common site for orbital blowout fracture is the medial wall of the orbit.

The patient with blow out fracture usually presents with immediate periorbital edema and blood extravasation in and around the orbit. Proptosis of the eyeball secondary to orbital edema or hemorrhage with paresthesia along the infraorbital nerve are also very common.

Imaging by a plain x-ray in water's view or by CT demonstrates depression of bony fragments and orbital contents herniating through the orbital floor into the maxillary antrum. This is called as the "hanging drop sign" or "tear drop sign".

Definitive treatment is by surgical repair 10-14 days after injury with initial supportive management with analgesics and antibiotics.

Le fort fracture- Is the term used to describe different types of maxillary fractures.

Solution to Question 6:

The likely diagnosis, based on the above image, is pterygium.

Pterygium is a subconjunctival fibrovascular tissue encroaching the cornea. It is a degenerative condition of the sub-conjunctival tissue and is never inflammatory. It occurs due to exposure to UV-B radiation and the most common site is the nasal.

It is made up of three parts, "cap" (an avascular halo-like zone at the advancing edge), a head, and a body.

Clinical features:

• Astigmatism with the rule - most common finding

• Stocker's line - deposition of iron on the advancing edge

• Fuchs islets - small, discrete, whitish flecks consisting of clusters of pterygial epithelial cells, often present at the advancing edge.

The differential diagnosis is pseudo pterygium.

The treatment for pterygium is usually excision. To prevent a recurrence, mitomycin C or 5 fluorouracil is used. Other methods:

• Auto-grafting- Conjunctival limbal autograft (CLAG) from the superior limbal conjunctiva is the best method.

• Bare sclera technique- high recurrence rate of 30 to 80%.

• PERFECT surgery- zero rates of recurrence. (PERFECT stands for Pterygium Extended Removal Followed by Extended Conjunctival Transplant)

Pterygium	Pseudo-pterygium
Degenerativeprocess	Inflammatoryprocess
Occurs inelderly	Can occur at any age

Pterygium	Pseudo-pterygium
Situated at thepalpebral aper ture	Can occur at any site
Either progressive, regressive, or stationary	Always stationary
Aprobe cannotbepassedunde rneath	A probe can be passed under the neck

Solution to Question 7:

Accommodative esotropia is associated with a high degree of hypermetropia (+4 to +7 D).

Accommodative esotropia is the most common type of squint. Typically presents around 2 to 3 years of age.

Types:

• Refractive

• Associated with high hypermetropia

• Fully correctable by spectacles

• Non-refractive type

• Abnormal accommodative convergence/accommodation (AC/A) ratio

• Fully corrected by bifocal glasses

• Miotics useful in cases where glasses are unsuitable

• Echothiophate causes iris cyst

• Prevention of iris cysts- Phenylephrine drops

• Mixed type

The accommodative convergence/accommodation (AC/A) ratio is defined as the amount of convergence measured in prism diopters per unit (diopter) change in accommodation.

The image below shows accommodative esotropia with and without correction.

Solution to Question 8:

Chemotherapy commonly used is a three-drug regimen containing vincristine, carboplatin, and etoposide.

Solution to Question 9:

The clinical vignette points to a diagnosis of glaucoma not responsive to medical treatment and trabeculectomy and the device used is an ex-Press™ glaucoma implant (mini shunt).

It is a form of minimally invasive glaucoma surgery (MIGS) to increase aqueous drainage.

Ex-PRESS glaucoma implant is a small stainless steel device. It is inserted under a scleral flap during a modified trabeculectomy. Complications such as hypotony are lower than with standard trabeculectomy.

Glaucoma drainage devices are devices that allow aqueous outflow by establishing communication between the anterior chamber and sub-Tennon's space/sub-conjunctival space. While these devices were traditionally reserved for high-risk patients, or after trabeculectomy had failed, they are becoming increasingly popular.

The commonly used devices are as follows:

1. Shunts using episcleral explants

• Ahmed glaucoma valve: Silicone

• Molteno implant - Polypropylene plate and silicone plate

• Baerveldt implant - Silicone

2. Mini-shunts

• Ex-Press™ Mini-Shunt: Stainless steel

• iStent® - Titanium

Solution to Question 10:

The above instrument is a Maddox rod.

Maddox Rod Test

A Maddox rod is a specially constructed device, consisting of a series of parallel cylinders, that converts a point source of light into a line image. It can be used to test for horizontal and vertical deviations, and when used in conjunction with another Maddox rod, for cyclo deviations.

A patient is asked to fix on a point light in the center of Maddox tangent scale placed 6 meters away. A Maddox rod is placed in front of one eye with the axis of the rod parallel to the axis of deviation. The patient will see a point light with one eye and a red line with the other. Due to dissimilar images of the two eyes, fusion is broken and heterophoria becomes manifest. The number on the Maddox tangent scale where the red line falls will be the amount of heterophoria in degrees.

Maddox tangent scale



©MARROW

Maddox rod



©MARROW

Maddox rod test



©MARROW



Solution to Question 11:

The test shown is the cover - uncover test.

Ophthalmology NEET 2020

Question 1:

Which of the following layers is responsible for maintaining the hydration and transparency of the cornea?

a) Descemet's membrane

b) Endothelial cells

c) Stroma

d) Corneal epithelium

Question 2:

What is the characteristic feature of a fungal ulcer?

a) Reverse hypopyon

b) Dendritic ulcer on a fluorescien dye

c) Ring abscess

d) Satellite lesion

Question 3:

What is the term used to describe an extra row of eyelashes behind the grey line?

a) Trichiasis

b) Distichiasis

c) Tylosis

d) Madarosis

Question 4:

A middle-aged woman presented with bilateral proptosis, restriction of eye movements, and chemosis. Her thyroid profile showed she was euthyroid. What is the most probable cause?

a) Orbital pseudotumor

b) Thyroid ophthalmopathy

c) Orbital lymphoma

d) Orbital cellulitis

Question 5:

Which of the following drugs increases the uveoscleral outflow?

a) Timolol

b) Latanoprost

c) Pilocarpine

d) Carbonic anhydrase inhibitor

Question 6:

Photostress test is used to differentiate between:

a) Macula and optic nerve disease

b) Cataract and glaucoma

c) Lens and cornea

d) Retina and vitreal pathologies

Question 7:

Shifting fluid sign is seen in?

a) Exudative RD

b) Traction RD

c) Rhegmatogenous RD

d) Retinal dialysis

Question 8:

A 40-year-old diabetic patient presented with the ■squint. On examination, the secondary deviation was found to be more than the primary deviation. Forced duction test is negative. Which of the following types of squint is most likely to be present in this patient?

a) Comitant squint

b) Restrictive squint

c) Paralytic squint

d) Pseudosquint

Question 9:

A physician observed the following finding in a small child who came for an eye exam. This finding is associated with which of the following conditions?

a) Myopia

b) Astigmatism

c) Hyperopia

d) Emmetropia

Question 10:

Sympathetic ophthalmia is seen after _____.

a) Penetrating trauma

b) Blunt trauma

c) Chemical injury

d) Urinary tract infection

Question 11:

A patient presented with unilateral proptosis and bilateral abducent palsy. What is the most probable cause ?

a) Retinoblastoma

b) Orbital pseudotumor

c) Orbital cellulitis

d) Cavernous sinus thrombosis

Answer Key

Question No.	Correct Option
1	b
2	d
3	b
4	b
5	b
6	a
7	a
8	c
9	c
10	a
11	d

Detailed Explanations

Solution to Question 1:

The hydration level and transparency of the cornea is maintained mainly by the endothelial cells of the cornea.

The endothelium possesses Na+/K+ ATPase, which keeps the corneal stroma in a relatively dehydrated state. This helps maintain corneal transparency. Endothelial cells decrease in number with age. If the cornea becomes overhydrated, it begins to scatter light and gradually begins to lose its transparency

Other factors contributing to the transparency of the cornea are as follows:

• Regular arrangement of corneal lamellae (Maurice theory)

• The peculiar refractive index of corneal lamellae (Goldmann theory)

• Avascularity of the cornea

• A balance between the intraocular pressure (IOP) and swelling pressure

The different layers of the cornea are as follows:

• Epithelium

• Bowman's membrane

• Stroma (substantia propria)

• Dua's layer (pre-Descemet's layer)

• Descemet's membrane

• Endothelium

Solution to Question 2:

Satellite lesions are characteristic of fungal ulcer.

The signs of a fungal ulcer include:

• Irregular margins of the ulcer.

• Presence of satellite lesions surrounding the ulcer.

• An immune ring (Wesseley) may be seen around the ulcer due to deposition of the immune complexes.

• Slough is usually dry, with feathery borders surrounded by a yellow line of demarcation.

• Hypopyon may be present or absent. If present, it is thick, immobile due to the direct invasion of the fungal hyphae into the anterior chamber.

Solution to Question 3:

The extra row of eyelashes behind the grey line is called distichiasis.

Distichiasis can occur due to congenital or acquired causes. These lashes may irritate the cornea. Management is by lid splitting with cryotherapy to the base of the hair follicles.

Other options:

Option A: Trichiasis is inward misdirection of cilia, with the normal position of the lid margin.

Option C: Tylosis is the thickening of the tarsal plate.

Option D: Madarosis is the loss of eyelashes or eyebrows.

Solution to Question 4:

The given clinical scenario is suggestive of thyroid ophthalmopathy.

Thyroid eye disease (TED) is the most common cause of both bilateral and unilateral proptosis in an adult in the euthyroid state. The main features of TED include lid retraction, proptosis, periorbital edema, chemosis, and restrictive myopathy.

Other options-

Option A- Orbital pseudotumor presents with painful eye movements periorbital edema, erythema, proptosis, ptosis, diplopia.

Option C- Orbital lymphoma usually presents in the 7th decade, most commonly observed as a painless palpable mass in the superolateral quadrant.

Option D- Orbital cellulitis is usually a unilateral condition with pain, proptosis, and chemosis.

NOTE: Thyroid eye disease can also, though less commonly, occur in euthyroid patients.

Solution to Question 5:

Latanoprost is a prostaglandin drug, which acts by increasing the uveoscleral outflow and thereby lowers the intraocular pressure.

Mechanism of action: Latanoprost is a prodrug and gets activated during its passage through the cornea. It acts by stimulating the prostanoid receptors on the ciliary muscles. This leads to the release of matrix metalloproteinases, which increases the uveoscleral outflow.

Solution to Question 6:

Photostress test is used to differentiate between the disease involving the macula and optic nerve.

This test is of particular use in early macular diseases such as central serous retinopathy.

In this test, one eye is covered and the patient is asked to read the smallest line on the near chart. A bright light is then shone into the eye for 15 seconds, and the patient is asked to read the line again, and the time taken to recover is noted. The test is repeated with the other eye.

In normal people and in patients with optic nerve disease, the recovery time is not affected. Whereas, in patients with macular disease the recovery time is prolonged and is usually more than one-third of the normal recovery time. This is because, in macular disease, the photoreceptors are damaged, and there is a delay in the visual pigment regeneration process.

Solution to Question 7:

The presence of shifting fluid is a hallmark feature of serous or exudative retinal detachment.

In serous retinal detachment, there is a subretinal accumulation of the serous fluid mostly due to the breakdown in the blood-retinal barrier. When the position of the head is altered, the fluid shifts to the dependent portion. Serous retinal detachment has a smooth convex configuration. A retinal tear or preretinal traction is absent.

Other options:

Option B: Traction RD has a concave configuration and breaks are absent. Retinal mobility is severely reduced and shifting fluid is absent.

Option C: Rhegmatogenous RD presents with tobacco dust in anterior vitreous with substantial vitreous blood or inflammatory cells. Retinal breaks are also seen.

Solution to Question 8:

The above clinical scenario is suggestive of paralytic squint. It is characterized by the presence of secondary deviation more than the primary deviation and a negative forced duction test.

The squint may be either comitant or incomitant.

• Comitant squint: The angle of squint measured is same in all gazes.

• Incomitant squint: The angle of squint measured is different in different gazes.

The differences between comitant and incomitant squint are shown in the table below:

The incomitant squint is further divided into two main types: Paralytic and restrictive.

Paralytic squint occurs due to the paralysis of extraocular muscle while restrictive squint is due to the mechanical restriction caused by tight or fibrosed muscle.

Forced duction test(FDT) differentiates between these two types. In this test, the eye is moved mechanically with the help of a forceps.

• The test is negative if there is no resistance to the movement. It is seen in a paralytic squint.

• The test is positive if there is resistance to the movement. It is seen in a restrictive squint.

Features	Comitantsquint	Incomitantsquint
Magnitude of squint	Same in all gazes	Varies with the gaze
Diplopia	Absent	Present
Ocular movements	Full	Restricted
False projection	Absent	Present
Abnormal head post ure	Absent	Present
Secondary deviation	Equal to primary deviatio n	More than primary deviati on

Solution to Question 9:

The given image shows convergent squint or esotropia, which is commonly seen in children with hyperopia, also known as hypermetropia.

Esotropia results due to increased accommodative effort in cases where there is under correction of hypermetropia. The condition may be prevented by full hypermetropic correction. If the child is unable to wear the glasses due to flat nasal bridge or if the child is less than 2 years old, the accommodation may be treated with drugs. Miotic drugs with anticholinesterase activity such as di-isopropyl fluorophosphate (DFP) or echothiophate can be used in these children. Further treatment may include orthotropic training or surgical correction of the squint.

Solution to Question 10:

Sympathetic ophthalmia is seen after a penetrating trauma.

Sympathetic ophthalmia is a bilateral granulomatous panuveitis seen after a penetrating trauma or less frequently, following intraocular surgery, usually multiple vitreoretinal procedures. Patients present between 2 weeks and 3 months after initial injury.

The earliest symptom is blurred near vision in the sympathizing (uninjured) eye. The earliest sign is a retrolental flare in the sympathizing eye. Anterior uveitis develops in both eyes. The fundus shows multifocal choroidal infiltrates in the mid-periphery. Sub-retinal pigment epithelium (RPE) infiltrates corresponding to Dalen–Fuchs nodules (image)(a granuloma situated between Bruch membrane and the RPE) are also present.

Steroids are the basis of treatment. High-dose oral prednisolone is given for several months. Immunosuppressants such as azathioprine, cyclosporine, and methotrexate can be used in resistant cases or as steroid-sparing agents. Enucleation of a severely injured eye with no visual potential within 2 weeks of injury prevents the occurrence of sympathetic ophthalmia.



Solution to Question 11:

The given clinical scenario is suggestive of cavernous sinus thrombosis.

Features of cavernous sinus thrombosis include early involvement lateral rectus palsy (abducent nerve involvement) since it runs through the cavernous sinus. Oculomotor and trochlear nerve palsy occur later as they are related to the lateral wall of the cavernous sinus. Other signs are proptosis, severe supraorbital pain, edema in the mastoid region, absence of corneal sensation, and dilated pupil.

Options A- Retinoblastoma occurs in young children and presents with unilateral proptosis with leukocoria, buphthalmos, and squint.

Option B- Orbital pseudotumor presents with unilateral proptosis with diplopia, conjunctival chemosis. The inflammation is responsive to systemic corticosteroids.

Option C- Orbital cellulitis presents with unilateral proptosis with pain and decreased ocular motility.



Ophthalmology INI-CET 2021

Question 1:

A 30-year-old female presents with redness and pain in the eye. Examination revealed an IOP of 38 mm of Hg, aqueous flare, and corneal precipitates. Which of the following drugs must be avoided in her?

a) Beta blockers

b) Mannitol

c) Carbonic anhydrase inhibitors

d) Prostaglandin analogs

Question 2:

Identify the condition:

a) Medial rectus palsy

b) Trochlear nerve palsy

c) Abducens nerve palsy

d) Oculomotor nerve palsy

Question 3:

Identify the site of the lesion.



a) Left occipital lobe

b) Right occipital lobe

c) Left geniculate body

d) Optic chiasma

Question 4:

What is the most common route of the spread of retinoblastoma?

a) Optic nerve invasion

b) Hematogenous

c) Direct invasion

d) Lymphatic spread

Question 5:

The most common tumor of the lacrimal gland is?

a) Adenoid cystic carcinoma

b) Non Hodgkin Lymphoma

c) Pleomorphic adenoma

d) Mucoepidermoid carcinoma

Question 6:

SAFE strategy for trachoma includes all except?

a) Surgery for trichiasis

b) Antibiotics

c) Facial cleanliness

d) Evaluation of control program

Question 7:

Identify the surgical step shown in the image given below.



a) Capsulorrhexis

b) Hydrodissection

c) Intraocular lens implantation

d) Lens aspiration

Question 8:

The most common ocular manifestation of COVID-19 is:

a) Retinal vein thrombosis

b) Viral keratouveitis

c) Follicular conjunctivitis

d) Pseudomembranous conjunctivitis

Question 9:

A patient who is a known case of angle-closure glaucoma has been diagnosed with major depressive disorder. Which of the following drugs have to be avoided in this patient?

a) Amitriptyline

b) Sertraline

c) Mirtazapine

d) Fluvoxamine

Question 10:

A patient is not able to see the lateral fields. He was having a mild headache for a few weeks and now presented to the clinic. Visual field examination shows the following finding. What is the probable site of the lesion based on the image displayed?



a) Optic chiasma

b) Occipital lobe

c) Optic tract

d) Optic radiation

Question 11:

Identify the finding marked by the arrow in the below image.



©MARROW

a) Vossius ring

b) Weiss ring

c) Fleischer's ring

d) Kayser Fleischer ring

Question 12:

Identify the chart given in the picture below.



a) Hess chart

b) Snellen chart

c) Pelli-Robson chart

d) ETDRS chart

Question 13:

A 14-day-old neonate brought to the outpatient department is found to have bilateral conjunctivitis. Chest X-ray done showed lung infiltrates. Which of the following organisms is most likely responsible for the infection?

a) Chlamydia trachomatis

b) Hemophilus influenzae

c) Streptococcus

d) Mycoplasma

Answer Key

Question No.	Correct Option
1	d
2	d
3	a
4	c
5	c
6	d
7	a
8	c
9	a
10	a
11	a
12	a
13	a

Detailed Explanations

Solution to Question 1:

The given clinical vignette is suggestive of anterior uveitis with a raised intraocular pressure. Among the given options, latanoprost is the least suited for use in this patient as it can result in worsening of the anterior uveitis.

Latanoprost is a PGF 2 alpha analog that acts by increasing the uveoscleral outflow and thereby lowers the intraocular pressure. Latanoprost is a prodrug and gets activated during its passage through the cornea. It acts by stimulating the prostanoid receptors on the ciliary muscles. This leads to the release of matrix metalloproteinases, which act to increases the uveoscleral outflow. It should be used with caution in those with uveitis, iritis, cystoid macular edema, and herpes simplex keratitis.

Topical prostaglandins like latanoprost, travoprost, bimatoprost, and tafluprost are now used as first-line agents to lower IOP. The most common ocular side effects are conjunctival hyperemia, eyelash thickening, iris hyperpigmentation, and loss of periorbital fat (bimatoprost). The systemic side effects are uncommon and include occasional headache, malaise, myalgia, skin rash, and mild upper respiratory infections.

Solution to Question 2:

The image is suggestive of oculomotor nerve palsy.

Isolated 3rd nerve palsy

Causes

• Microvascular disease such as hypertension and diabetes is the most common cause of third nerve palsy.

• Aneurysm of the posterior communicating artery

• Episodic cases of isolated palsy may be seen due to migraine, compression, ischemia, and alterations in intracranial pressure

• Trauma

• Inflammatory diseases such as syphilis, Lyme disease, sarcoidosis; giant cell arteritis

• Vasculitis

The signs of left oculomotor nerve palsy are summarized in the image attached below.

Solution to Question 3:

The visual field defect shows right homonymous hemianopia with macular-sparing, which is caused by lesions of the left occipital lobe (contralateral/crossing of fibers).

Such lesions are generally due to the occlusion of the posterior cerebral artery. These are commonly seen in head injury and may be associated with visual hallucinations, denial of blindness (Anton syndrome), and the Riddoch phenomenon (only moving visual targets are perceived). These lesions generally spare the central vision/macular vision. The anatomical reason for this sparing is attributed to the uniqueness in the blood supply. While the part of the occipital lobe representing the peripheral field vision is supplied by the posterior cerebral artery, the area at the tip of the calcarine cortex, representing the central macular vision is supplied by a branch of the middle cerebral artery.

Solution to Question 4:

Direct invasion is the most common mode of spread of retinoblastoma.

Retinoblastoma arises from the neurosensory retina. It is a proliferation of neural cells that have failed to evolve normally. Retinoblastoma is the most common primary intraocular malignancy in children. Tumours are composed of small basophilic cells (retinoblasts), with large hyperchromatic nuclei and scanty cytoplasm.

It is of 4 types:

• Endophytic (into the vitreous)with seeding of tumor cells throughout the eye

• Exophytic (into the subretinal space) leading to retinal detachment

• Mixed type

• Retinocytoma: retina may be diffusely infiltrated

Solution to Question 5:

Pleomorphic adenoma is the most common tumor of the lacrimal gland while adenoid cystic carcinoma is the most common carcinoma of the lacrimal gland.

Solution to Question 6:

E in the SAFE strategy stands for Environmental improvement with better hygiene and sanitation.

Solution to Question 7:

Capsulorrhexis is the step shown in the given image.

Capsulorrhexis is a method of anterior capsulotomy, where an opening is made in the anterior capsule of the lens. It is one of the steps involved in conventional extracapsular cataract extraction (ECCE).

Anterior capsulotomy can be done by one of three methods - can opener's technique, linear capsulotomy or envelope technique, or continuous curvilinear capsulorhexis.

Continuous curvilinear capsulorrhexis (CCC), is considered to be superior to all the preexisting methods because when done correctly, it does not leave any edges or tears. Trypan blue dye can be used to stain the lens capsule for better visibility.

In ECCE, a portion of the anterior capsule of the lens is removed, allowing extraction of the lens nucleus and cortex. This leaves the remainder of the anterior capsule, the posterior capsule, and the zonular support intact. The intact lens nucleus is then removed from the eye through a limbal incision.

An anterior capsulotomy is followed by hydro dissection using a balanced salt solution. This helps to separate the corticonuclear mass from the capsule. Removal of lens nucleus and aspiration of the cortex is done using a two-way irrigation and aspiration cannula. An intraocular lens (IOL) is then implanted and the incision is then closed.

Some of the IOLs used are shown in the image below.



Solution to Question 8:

The most common ocular manifestation of COVID-19 is follicular conjunctivitis.

It has been observed that SARS-CoV-2 needs the ACE-2 receptors for cell invasion. The ACE-2 receptors are found not only in human type 2 alveolar epithelial cells but also in the cornea and conjunctiva. This suggests that ocular surface tissue may be a potential target tissue for SARS-CoV-2.

The ocular manifestations of COVID-19 vary greatly. The most prevalent ocular symptoms were dry eye or foreign body sensation, redness, tearing, itching, eye pain, and discharge.

The most common ocular complication of COVID-19 is mild follicular conjunctivitis. Hemorrhagic conjunctivitis, dry eye disease, episcleritis, chemosis, and photophobia can also occur, but less frequently. Intraocular complications include retinal artery occlusions, hemorrhages, and cotton wool spots caused by complement-mediated thrombotic angiopathy.

Sometimes, conjunctivitis may appear as the only sign and symptom of COVID-19. Patients should therefore undergo a nasopharyngeal RT-PCR test.

All physicians and ophthalmologists should be cautious when addressing a patient with conjunctivitis and should wear eye-protective gear in addition to face masks and other protective devices during clinical examination of these patients.

Solution to Question 9:

Amitriptyline is the drug to be avoided in patients having angle-closure glaucoma with major depressive disorder.

In angle-closure glaucoma, there is apposition of peripheral iris against the trabecular meshwork resulting in obstruction of aqueous outflow by the closure of an already narrow-angle of the

anterior chamber, with an increase in the intraocular pressure being the aftermath.

Amitriptyline is a tricyclic antidepressant that also has a marked anticholinergic activity due to a high affinity for muscarinic acetylcholine receptors and the subsequent blockade of the receptors. These muscarinic cholinoceptors, specifically M3 type, upon stimulation, are responsible for contraction of the iris sphincter, ciliary muscle and trabecular meshwork as well as the increased outflow of aqueous humour. Therefore, blocking of these receptors would result in the decreased outflow of aqueous humour, which ultimately leads to an increase in intraocular pressure.

Therefore, amitriptyline when administered in a patient with pre-existing angle-closure type glaucoma can exacerbate the disorder by further impeding the drainage of aqueous humour.

Other options:

Options B and D: Sertraline and Fluvoxamine are selective serotonin reuptake inhibitors with no activity at the muscarinic cholinergic receptors and therefore have no effect on the outflow of aqueous humour and thereby, on the intraocular pressure.

Option C: Mirtazapine is an atypical antidepressant that is a serotonin-norephedrine reuptake inhibitor with the ability to precipitate angle-closure glaucoma by itself; however, due to lack of any activity at the muscarinic cholinergic receptors, it does not increase the intraocular pressure to the extent that Amitryptaline does.

Solution to Question 10:

The visual field defect in the image is indicative of bitemporal hemianopia, so the probable site of the lesion is the optic chiasma.

Hemianopia or hemianopsia refers to loss of sight in half the visual field. Bitemporal hemianopia refers to loss of sight in the temporal halves of the visual field, seen in lesions affecting the optic chiasma. The most common cause of this is pituitary gland tumours. Other causes include suprasellar aneurysms, chronic arachnoiditis, suprasellar tumours like craniopharyngioma and suprasellar meningioma.

Damage to the middle part of the optic chiasm affects fibres crossing at the nasal retina of each eye, leaving the uncrossed fibres from the temporal retinas intact. The resulting loss of vision is confined to the temporal visual field of each eye and this is known as bitemporal hemianopia. It is also called heteronymous hemianopia to emphasize the fact that the parts of the visual field that are lost in each eye do not overlap.

Individuals with this condition are able to see in both left and right central visual fields, with both eyes open. However, all information from the most peripheral parts of visual fields (which are seen only by the nasal retinas) is lost.

Other options:

Lesions at the occipital lobe, optic tract and optic radiation would all cause homonymous hemianopia. Here, either the right or left half of the binocular field of vision is lost, due to the loss of the temporal half of one field and the nasal half of the other.

Solution to Question 11:

The finding given in the image is that of a Vossius ring.

Vossius ring is a circular ring of faint or stippled opacity, seen on the anterior surface of the lens due to brown amorphous granules of pigment lying on the capsule of the lens. It is usually seen after blunt trauma to the eye and orbit.

It usually has about the same diameter as the contracted pupil. The force of the blow drives the cornea and iris backward and thus producing an impression of the iris on the lens. The resorption of the pigment results in discrete subcapsular opacities.

Other options:

Option B: Weiss ring is a floater which may be seen by the patient as a circle or other large solitary lesion. It is the detached former attachment to the margin of the optic disc. It is seen in posterior vitreous detachment and is not confirmatory. It is seen in the image below.



Option C: A Fleischer's ring is usually observed in keratoconus, representing epithelial iron deposits. It is seen best with a cobalt blue filter.

Option D: Kayser-Fleischer ring is observed in Wilson's disease, due to intraocular deposition of copper. A brownish-yellow zone is seen in the peripheral Descemet membrane of the cornea. Anterior capsular cataract, also knowns as a sunflower cataract may be seen in some patients.



Solution to Question 12:

The chart given in the picture is Hess's chart.

Hess chart is used to diagnose and monitor patients with incomitant strabismus by assessing the field of a binocular single vision. In its interpretation, a smaller chart indicates the eye with the paretic muscle and a larger chart indicates the eye with an overacting muscle.

Incomitant strabismus is one where squint generally results from the limitation of eye movement associated with paralytic or mechanical etiologies like extraocular muscle palsy, thyroid ophthalmopathy, blow-out fracture, myasthenia gravis, etc. The field of binocular single vision is the area where bifoveal fusion of an object of interest occurs.

Other options:

Option B: The Snellen chart is used to test visual acuity. Visual acuity is the eye's ability to discriminate between two points.

A picture of the Snellen chart is given below.



Option C: The Pelli-Robson contrast sensitivity letter chart consists of letters of constant size but decreasing contrast. It is viewed by the patient from a distance of 1 meter until the lowest resolvable group of three letters on the chart is reached. It is used to measure contrast sensitivity, a measure of the ability to distinguish an object against its background.

An image of the Pelli-Robson chart used in the middle east is given below.



Option D: The early treatment diabetic retinopathy study or ETDRS chart presents a series of 5 letters of equal difficulty on each row, with standardized spacing between letters and rows. It consists of a total of 14 lines having 70 letters. It is also used to test visual acuity. An image of the ETDRS chart is given below.



Solution to Question 13:

Chlamydia trachomatis is most likely responsible for the infection.

Chlamydia trachomatis inclusion conjunctivitis usually manifests 1 week after birth. It is usually a venereal infection derived from the cervix or urethra of the mother during delivery.

On examination, the conjunctiva is swollen and edematous. The discharge is initially watery, later becomes purulent or bloody.

Newborns with suspected chlamydia infection should have samples drawn from both the conjunctivae and oropharynx. Since it is an obligate intracellular organism, swabs should also include conjunctival epithelial cells if possible. Subsequent Giemsa staining shows characteristic intracellular inclusion bodies formed by Chlamydia oculogenitalis.

Treatment involves administration of suspension of erythromycin ethylsuccinate 50 mg/kg daily in four divided doses before feeds for 2–3 weeks.

If left untreated, associated systemic diseases such as chlamydial otitis and pneumonia may develop later. In all cases, both parents must receive appropriate treatment for genital infection.

Neonatal conjunctivitis, also called ophthalmia neonatorum, typically presents during the first month of life. The age of the neonate is an important clue towards the etiology of the infection.

Neonates are at higher risk of conjunctivitis due to predisposing factors such as:

• Decreased tear production

• Absence of IgA in tears

• Decreased immune function

• Absence of lymphoid tissue in the conjunctiva

• Decreased lysozyme activity.

Other risk factors for developing neonatal conjunctivitis are premature rupture of membranes, prolonged delivery, prematurity, poor prenatal care, mechanical ventilation, poor hygiene conditions, and history of midwife interference.

Other options :

Option B: Hemophilus influenza is one of the most common causes of acute bacterial conjunctivitis in children < 6 years of age and is responsible for causing conjunctivitis-otitis syndrome where conjunctivitis is followed by acute otitis media.

Option C: Streptococcus usually manifests 48-72 hours after birth. It is a common cause of acute bacterial conjunctivitis in children.

Option D: Mycoplasma, specifically M.genitalium can cause conjunctivitis due to contamination of the eye with genital infections in adults.

Age	Cause of conjunctivitis
First 24 hours	Chemical causes- erythromycin/gentamycin drops,
24 to 48 hours	Neisseria gonorrhea (most common), Staphylococcus aureus, Streptococcus, coliforms
5 to 14 days	Chlamydia trachomatis, Herpes keratoconjunctivitis
5 to 18 days	Pseudomonas aeruginosa

Ophthalmology NEET 2021

Question 1:

A 27-year-old IT professional has presented to you with intense watering, redness, and foreign body sensation in the eye for the past 2 months. He took out his contact lens and applied eye lubricants to help alleviate his ailment, but it didn't provide any relief. What is the most probable diagnosis?

a) Acute trachoma

b) Follicular conjunctivitis

c) Giant papillary conjunctivitis

d) Spring catarrh

Question 2:

A 60-year-old patient with progressive painless loss of vision presents to the ophthalmology OPD. The fundus image of the patient is given below. What could be the probable finding and the cause of it?



a) Soft exudate, hypertension

b) Hard exudate, diabetes mellitus

c) Flame-shaped hemorrhages, hypertension

d) Soft exudate, central retinal vein occlusion

Question 3:

A 33-year-old woman presents with complaints of progressive loss of vision in the right halves of both eyes. Where is the lesion located in the optic pathway?

a) Left optic tract

b) Right visual cortex

c) Optic chiasma

d) Right optic nerve

Question 4:

A 2-month-old infant was brought by his mother with complaints of inability to open the eyes in light and excessive watering. The clinical picture is given below. What is the most probable diagnosis?

a) Ophthalmia neonatorum

b) Congenital glaucoma

c) Mucopolysaccharidosis

d) Cataract

Question 5:

A 15-year-old girl is not compliant with spectacles for her myopic astigmatism. What would be the appropriate management in her case?

a) LASIK

b) Spherical equivalent spectacles

c) Femto LASIK

d) Implantable collamer lens

Question 6:

What is the most common complication associated with the following abnormality?

a) Exposure keratopathy

b) Restricted eye movements

c) Diplopia

d) Amblyopia

Question 7:

A 7-year-old boy presented with a white pupillary reflex for which enucleation was done. The histopathological examination of the specimen showed Flexner Winterstein rosettes. What is the most likely diagnosis?

a) Retinoblastoma

b) Medulloblastoma

c) Rhabdomyosarcoma

d) Astrocytoma

Question 8:

An industrial worker was admitted to the hospital after an eye injury sustained while cutting stone with a hammer and iron chisel. A foreign body was suspected to be impacted in his eye. Which investigation can be detrimental to this patient?

a) MRI

b) CT

c) X-ray

d) B mode scan

Answer Key

Question No.	Correct Option
1	c
2	b
3	a
4	b
5	b
6	a
7	a
8	a

Detailed Explanations

Solution to Question 1:

The clinical history of watering, redness, and foreign body sensation in a contact lens user, along with the image showing giant papillae in the conjunctiva, suggests a probable diagnosis of giant papillary conjunctivitis.

Giant papillary conjunctivitis is a severe form of mechanically induced papillary conjunctivitis. It is most commonly associated with contact lens wear. It can also be associated with ocular prostheses, exposed sutures, and scleral buckles.

Patients usually present with redness, foreign body sensation, increased mucus production, itching, blurring, and loss of contact lens tolerance. On examination, giant papillae are seen on the upper tarsal conjunctiva with hyperemia. Focal apical ulceration and whitish scarring may develop on larger papillae.

Papillae are reddish, flat-topped, raised areas that give a red and velvety appearance to the tarsal conjunctiva. These are caused by the hyperplasia of the normal vascular system, with glomerulus-like bunches of capillaries growing into the epithelium.

The image below shows papillae in the tarsal conjunctiva.

Treatment includes the removal of the offending cause. Mast cell stabilizers, such as sodium cromoglycate, and drugs like azelastine and olopatadine, which have combined antihistamine and mast cell–stabilizing effects, are effective. Steroids can be used for resistant cases.

Other options:

Option A: In acute trachoma, sago-like follicles are characteristically seen. It is caused by Chlamydia trachomatis. The SAFE strategy is used in the management of trachoma.

The image below shows sago-like follicles in trachoma.

Option B: In follicular conjunctivitis, follicles are seen. It is most commonly caused by herpes and adenoviruses. Follicles are greyish-white raised areas on fornices and palpebral conjunctiva. These are formed due to the localized aggregations of lymphocytes in the sub-epithelial adenoid layer.

The image shows follicles.



Option D: In spring catarrh (vernal catarrh), cobblestone appearance is a papillary reaction in the palpebral conjunctiva, ropy discharge, and Maxwell-Lyon sign are present. Horner-Trantas spots that are discrete, whitish, raised dots along the limbus can also be seen. It is more common during summer.

The image below shows Horner-Trantas spots.



Solution to Question 2:

The given image shows small, discrete yellowish, waxy areas with crenated margins that are suggestive of hard exudates, seen in diabetes mellitus.

In diabetic retinopathy, hard exudates occur due to chronic localized edema and are composed of leaked lipoproteins and lipid-filled macrophages.

Other common causes of hard exudates include hypertensive retinopathy, Coats' disease, and circinate retinopathy.



Cotton wool spots (soft exudates) appear as whitish fluffy spots with indistinct margins that represent areas of infarcts in nerve fiber layer. Common causes are hypertensive retinopathy, toxemic retinopathy of pregnancy, diabetic retinopathy, anemias, and collagen disorders like discoid lupus erythematosus, polyarteritis nodosa, and scleroderma.

The image shows cotton wool spots



Flame-shaped hemorrhages are superficial retinal hemorrhages that occur due to disruption of the capillaries in the retinal nerve fiber layer. Common causes are diabetic retinopathy, non-ischemic central retinal vein occlusion, traumatic retinopathy, Eale's disease, hypertensive retinopathy, and anemic retinopathy.



Fundus examination of central retinal vein occlusion (CRVO) reveals massive engorgement, congestion, and tortuosity of retinal veins. Ischemic CRVO has massive retinal hemorrhages giving a splashed tomato appearance involving the whole fundus.



Solution to Question 3:

The patient presenting with progressive visual loss in the right halves of both eyes is suggestive of right homonymous hemianopia. The site of the lesion in this case is the left optic tract.

Here, vision is lost in the right half of both eyes, which means:

• in the left eye, the nasal visual field is lost, and

• in the right eye, the temporal visual field is lost.

It means left temporal and right nasal fibers are destroyed which leads to right homonymous hemianopia. Anatomically, this can be explained by the fact that each optic tract contains fibers from the ipsilateral temporal half of the retina and the contralateral nasal half of the retina.

Optic tract lesions can occur due to pituitary adenomas (most common), craniopharyngioma, tumors of optic thalamus, large aneurysm of superior cerebellar or posterior cerebral arteries, demyelinating diseases, and infarction.

Other options:

Option B - Lesion at right visual cortex causes left homonymous hemianopia with macular sparing.

Option C - Lesion at optic chiasma causes bitemporal hemianopia.

Option D - Lesion at the right optic nerve causes blindness in the right eye.



Solution to Question 4:

The clinical history of inability to open eyes in light (photophobia), excessive watering, and the image showing hazy cornea are suggestive of congenital glaucoma.

Primary congenital glaucoma refers to abnormally high intraocular pressure (IOP) which occurs as a result of a developmental anomaly of the angle of the anterior chamber. It is not associated with any other ocular or systemic anomaly. It can be classified as follows:

• True congenital glaucoma: The IOP is elevated during intrauterine life.

• Infantile glaucoma: Manifests prior to age 3.

• Juvenile glaucoma: The IOP rises between 3 and 16 years of age. It is the least common type.

The CYP1B1 gene is implicated in the pathogenesis of congenital glaucoma.

Clinical presentation: Lacrimation, photophobia, and blepharospasm form the classic triad of symptoms of congenital glaucoma. A corneal haze is seen due to diffuse edema secondary to

raised IOP. Buphthalmos occur as a result of stretching due to elevated IOP prior to the age of 3 years. Haab striae that are curvilinear healed breaks in Descemet's membrane are seen. Corneal scarring can also occur.

Evaluation under general anesthesia is generally required. IOP measurement, anterior chamber examination, optic disc examination, corneal diameter measurement, gonioscopy, and refraction should be evaluated.

Management is essentially surgical with goniotomy or trabeculotomy. If angle surgery fails, trabeculectomy, tube shunt implantation, and ciliary body ablative procedures are performed.

Other options:

Option A - In ophthalmia neonatorum, there will be inflammation of the conjunctiva. Clinical presentation includes conjunctival discharge, hyperemia chemosis, swollen eyelids, tender eyeball. Corneal involvement is rare which may occur in the form of superficial punctate keratitis.

Option C - In mucopolysaccharidosis, there will be bilateral cloudy cornea.

Option D - In cataract, there will be opacification of lens presenting as whitish reflex/ leukocoria.

Solution to Question 5:

The appropriate management for myopic astigmatism in a 15-year-old girl (<18 years) would be spherical equivalent spectacles.

Prescription in astigmatism has 3 components:

• Power of the spherical component: Convex (+) or concave (-) in dioptre sphere (DS)

• Power of the cylinder: Convex (+) or concave (-) dioptre cylinder (DC)

• Axis or orientation of the cylinder

The cylinder power and axis are correcting the astigmatic part of the refractive error. If an astigmatic eye were to be corrected with a spherical lens alone, the result would be a blur circle on the retina, called the circle of least confusion.

The spherical lens power places the circle of least confusion on the retina and is called the spherical equivalent. It is defined as the spherical power whose focal point coincides with the circle of least confusion, where one would see minimal blurring in their vision. To calculate the spherical equivalent, add half the cylinder power to the sphere power:

Spherical equivalent= spherical power + 1/2 (cylindrical power)

For example, for spherical power +2 and cylindrical power +3,

+2.00 +(+3.00 /2) = +3.50.

Spherical equivalent spectacles are given if one has to give less cylinder power than was measured.

Other options:

Options A and C - LASIK and Femto LASIK should be done in patients above 18 years of age.

Option D - Implantable Collamer lens is indicated for patients 21-45 years of age.

The image demonstrates the steps of LASIK



Solution to Question 6:

The given image shows a coloboma of the upper eyelid. The most common complication associated with this condition is exposure keratopathy since the defect leaves the cornea exposed. If left untreated, it can lead to corneal ulceration.

Coloboma of the eyelid is a partial or full-thickness eyelid defect. It usually occurs near the nasal side. It more frequently involves the upper lid than the lower lid. It occurs when eyelid development is incomplete, due to failure of migration of lid ectoderm to fuse with the lid folds.

Small defects can be managed by primary closure, while large defects require skin grafts and rotation flaps.

Upper lid colobomas occur at the junction of the middle and inner thirds. It has a relatively strong association with cryptophthalmos, facial abnormalities, and Goldenhar syndrome. Cryptophthalmos is a congenital anomaly in which the eyelids are absent, replaced by a continuous layer of skin.

Lower lid colobomas occur at the junction of the middle and outer thirds and are frequently associated with systemic conditions.

Treacher Collins syndrome (mandibulofacial dysostosis) is a genetically heterogeneous condition. It is characterized by mandibular and ear anomalies. lower eyelid coloboma, slanted palpebral apertures, cataract, microphthalmos, and lacrimal atresia.



Solution to Question 7:

The most likely diagnosis with a histopathological finding of Flexner Winterstein rosettes in a 7-year-old child with a whitish pupillary reflex is retinoblastoma. Flexner Winterstein rosettes refer to an arrangement of columnar cells around a central lumen, which is highly characteristic of retinoblastoma.

Retinoblastoma is a malignant proliferation of immature retinal neural cells and is the most common intraocular malignancy in children. It is associated with mutations in the RB1 gene at chromosome 13q14. It is a tumour of a group called small round blue cell tumours because the immature retinal neural cells are seen as small round cells with large nuclei.

White pupillary reflex or leukocoria is the most common presenting feature, followed by a squint which is usually convergent. Rarely, the child may present with nystagmus and defective vision.

Image showing leukocoria of left eye



Histopathology of the tumor shows small round cells with large nuclei. Microscopic features of a well-differentiated tumor include Flexner-Wintersteiner rosettes (highly specific of retinoblastoma), Homer-Wright rosettes, pseudorosettes, and fleurettes formation. Areas of necrosis and calcification can also be seen.

The histopathological image of retinoblastoma is shown below.



Treatment typically involves chemotherapy plus focal therapy. The commonly used three-drug regimen is the CVE regimen consisting of Vincristine, Carboplatin, Etoposide. Focal therapy includes cryotherapy, laser photocoagulation, thermotherapy, plaque radiotherapy, and external beam radiotherapy.

Enucleation is done when

• the tumor involves more than half of the retina

• the optic nerve is involved

• glaucoma is present and the anterior chamber is involved

In the stage of extraocular extension, the globe bursts through the sclera, usually near the limbus or near the optic disc. This is followed by rapid fungation and involvement of extraocular tissues results in marked proptosis. Distant metastasis occurs through the lymphatic spread and the direct extension by continuity to the optic nerve and brain.

Other options:

Option B - Medulloblastoma is a malignant embryonal tumor of childhood that occurs exclusively in the cerebellum. Patients usually present with headaches, ataxia, and signs of brainstem involvement.

Histologically, the tumor is very densely cellular. Individual tumor cells have scant cytoplasm and elongated or crescent-shaped hyperchromatic nuclei. Tumor cells form Homer Wright rosettes. They express neuronal markers such as synaptophysin.

Medulloblastoma



©MARROW

Option C - Rhabdomyosarcoma is the most common soft tissue sarcoma of childhood. The child usually presents with rapidly progressive unilateral proptosis. There are 4 subtypes - embryonal, alveolar, botyroid, and pleomorphic. Cells show light microscopic features of striated muscle differentiation. Immunohistochemistry is positive for desmin.



Option D - Astrocytoma of the retina and optic nerve head is rare. It is usually asymptomatic and is most frequently seen in tuberous sclerosis. Fundus examination reveals a yellowish or white plaque, nodule, or mulberry-like lesion. These lesions are semitransparent peripherally and calcified centrally.

Image showing calcified astrocytoma:



Solution to Question 8:

MRI can be detrimental for a patient with a history of an intraocular foreign body (IOFB) after cutting stone with a hammer and iron chisel. It can cause further damage by producing movements of the foreign body if it is magnetic. The use of MRI is contraindicated for metallic (iron) foreign bodies and is not recommended as a screening procedure.

The foreign bodies most likely to penetrate and be retained in the eye are minute chips of iron or steel. While chipping stone with an iron chisel, the chip of the iron chisel is more likely to enter the eye than the stone. A foreign body entering the eye may cause damage by:

• mechanical effects

• introduction of infection

• reacting with the intraocular tissues

Anteroposterior and lateral view X-rays are recommended for locating an IOFB as most foreign bodies are radio-opaque. Ultrasonographic localization can tell the position of even non-radiopaque foreign bodies. CT scan is the best method of IOFB localization. MRI is contraindicated for metallic IOFB.

Management: A foreign body should always be removed unless:

• it is inert and probably sterile

• little damage has been done to vision

• the process of removal will destroy sight

Siderosis bulbi refers to the ocular degenerative changes produced by an iron foreign body. It usually occurs 2 months to 2 years after the injury. This condition is due to the electrolytic dissociation of the metal by an intrinsic resting current in the eye which disseminates the metal, combining it with cellular proteins and cause atrophy.

Clinical features:

• Iris: first stained greenish and later turns reddish-brown, causing heterochromia iridis.

• Pupil: iron deposition in sphincter leads to mydriasis.

• Lens: iron deposition in a radial fashion on the anterior capsule of the lens, leading to anterior capsular cataract.

• Retina: pigmentary retinopathy; followed by atrophy of the retina and retinal pigmentary epithelium.

• Secondary open-angle Glaucoma: due to damage to the trabecular meshwork.

Ophthalmology INICET 2022

Question 1:

What is the action of the superior oblique muscle?

a) Intorsion, abduction, depression

b) Intorsion, adduction, elevation

c) Extorsion, abduction, depression

d) Extorsion, adduction, elevation

Question 2:

Which of the following triad is seen after an acute attack of angle-closure glaucoma?

a) Vogt's triad

b) Virchow triad

c) Hutchinson's triad

d) Gaucher triad

Question 3:

In which of the following conditions do you see punched-out lesions of the retina?

a) Posterior staphyloma

b) Fundal coloboma

c) Congenital rubella

d) Toxoplasmosis

Question 4:

Which of the following has familial inheritance?

a) 1, 2, 3, 4

b) 1, 2, 3

c) 1, 3

d) 3, 4

Question 5:

A 45-year-old man presented with lagophthalmos, orbicularis oculi palsy, and loss of corneal sensation. Which of the following is the most likely causative organism?

a) Acanthamoeba castellanii

b) Mycobacterium leprae

c) Onchocerca volvulus

d) Chlamydia A to C

Question 6:

A 27-year-old female patient presents with sudden diminishing vision associated with relative afferent pupillary defect in the right eye. On examination, the left eye is normal. Which of the following investigations would you do next?

a) 1, 2, and 3

b) 1 and 2

c) 2 and 3

d) 3 and 4

Question 7:

Which of the following is not a feature of the optic disc in glaucoma?

a) Roenne's nasal step

b) Bayonetting

c) Increased cup-disc ratio

d) Baring of nasal arteries

Question 8:

A farmer presents with pain and discharge in his left eye. He says that he sustained an injury caused by a leaf a few days back. What is the finding seen in the image?

a) Hyphema

b) Aqueous flare

c) Arlt line

d) Hypopyon

Question 9:

What is the most common cause of moderate-severe visual impairment according to the National Blindness and Visual Impairment Survey 2015-19?

a) Refractive error

b) Untreated cataract

c) Glaucoma

d) Diabetic retinopathy

Question 10:

Which of the following statements is true about optic neuritis?

a) Relative afferent pupillary defect is seen.

b) There is no pain on ocular movement.

c) Multiple myeloma is a common cause.

d) Oral corticosteroids are the mainstay of treatment.

Question 11:

The most common ocular manifestation of COVID-19 is _____.

a) Uveitis

b) Mucormycosis

c) Follicular conjunctivitis

d) Papillary conjunctivitis

Question 12:

Identify the condition given in the image.

a) Symblepharon

b) Ankyloblepharon

c) Congenital blepharophimosis

d) Congenital ptosis

Question 13:

Match the following:

a) 1-d, 2-b, 3-a, 4-c

b) 1-a, 2-c, 3-d, 4-b

c) 1-b, 2-a, 3-d, 4-c

d) 1-d, 2-c, 3-a, 4-b

Question 14:

In which of the following is atropine contraindicated?

a) Acute congestive glaucoma

b) Acute anterior uveitis

c) Malignant glaucoma

d) Corneal ulcer

Question 15:

What is the mechanism of action when the retina is exposed to light?

a) Hyperpolarization, increase in neurotransmitter release

b) Hyperpolarization, decrease in neurotransmitter release

c) Depolarization, decrease in neurotransmitter release

d) Depolarization, increase in neurotransmitter release

Question 16:

Which of the following is not a feature of keratoconus?

a) Haab's striae

b) Munson's sign

c) Irregular astigmatism

d) Scissoring reflex

Question 17:

A patient has presented for a routine eye evaluation. You have checked his visual acuity using Snellen's chart and found it to be 6/6. What is the minimum angle of resolution?

a) 15 minutes of arc

b) 5 minutes of arc

c) 10 minutes of arc

d) 20 minute of arc

Question 18:

Which of the following is not a feature of pterygium?

a) Rate of recurrence is 95%

b) It is a fibrovascular growth

c) The parts of a pterygium are head, neck and body

d) Has progressive and regressive types

Question 19:

Which of the following is not a feature of complicated cataracts?

a) Krukenberg spindle

b) Polychromatic lusture

c) Occurs after uveitis

d) Breadcrumb appearance

Question 20:

Oculocephalic reflex assesses the integrity of which of the following cranial nerves?

a) A and C

b) A and E

c) A and D

d) A, C, and D

Question 21:

A pregnant woman presents with fever, retroorbital pain, headache, pulsatile proptosis of the right eye, and tinnitus. BP and fundus were normal. Which of the following structures are involved?

a) A,B,C

b) B,C,D

c) A,C,D

d) A,B,C,D

Answer Key

Question No.	Correct Option
1	a
2	a
3	d
4	a
5	b
6	a
7	a
8	d
9	b
10	a
11	c
12	b
13	c
14	a
15	b
16	a
17	b
18	a
19	a
20	d
21	d

Detailed Explanations

Solution to Question 1:

The actions carried out by the superior oblique muscle are intorsion, depression, and abduction.

There are six extraocular muscles for each eye. Four rectus muscles namely - superior, inferior, medial, and lateral recti. The oblique muscles are two - superior oblique and inferior oblique.

These muscles are responsible for the movements of the eye which can be either unilateral or binocular movements.

The superior oblique muscle originates from the apex of the orbit, turns around the trochlea, and inserts into the sclera behind the equator. The oblique direction of the tendon is responsible for its rotatory effect (intorsion), in addition to depression and abduction.



Lateral view of the right eye

Unilateral movements (ductions) are :

1.Adduction: It is the movement towards the nasal side around the vertical axis.

2.Abduction: It is the movement towards the temporal side around the vertical axis.

3.Supraduction (elevation): It is upward movement around the horizontal axis.

4.Infraduction (depression): It is downward movement around the horizontal axis.

5.Intorsion (medial rotation): Rotatory movement along the anteroposterior axis where the superior pole of cornea moves medially.

6.Extorsion (lateral rotation): Rotatory movement along the anteroposterior axis where the superior pole of cornea moves laterally.

Binocular movements can be of two types - versions and vergences.

The oculomotor nerve supplies all the extraocular muscles except the superior oblique and lateral rectus. The superior oblique muscle is supplied by the trochlear nerve and lateral rectus muscle by the abducent nerve.



Solution to Question 2:

Vogt's triad is seen after an acute attack of primary angle-closure. It includes glaukomflecken, patches of iris atrophy, and a slightly dilated nonreactive pupil. Glaukomflecken is used to describe an anterior subcapsular lenticular opacity. The slightly dilated pupil is attributed to sphincter atrophy.

Glaucoma is used to describe a group of conditions where there is progressive optic neuropathy with irreversible visual field defects associated with raised intraocular pressure. It can be classified into congenital/developmental glaucomas, primary adult glaucomas, and secondary adult glaucomas. Primary adult glaucomas are of three types:

• Primary open-angle glaucoma (POAG)

• Primary angle-closure glaucoma (PACG)

• Primary mixed mechanism glaucoma.

In primary angle-closure disease, there is apposition of the peripheral iris with the trabecular meshwork. This interferes with the drainage of aqueous humor. It is seen more commonly in elderly females. Anatomical risk factors include hypermetropic eyes, anteriorly placed iris-lens diaphragm, and a narrow-angle of the anterior chamber. The international society of geographical and epidemiological opthalmology (ISGEO) has classified primary angle-closure disease into three types.

• Primary angle-closure suspect (PACS)

• Primary angle-closure (PAC)

• Primary angle-closure glaucoma (PACG).

PACS and PAC are seen before the development of PACG. Vogt triad is seen after an acute attack of PAC.

PAC includes irido-trabecular contact on gonioscopy and elevated intraocular pressure (IOP) and/or peripheral anterior synechiae. The optic disc and visual fields are usually normal. Due to any reason, there can be an acute rise of IOP in patients with PAC. It is a vision-threatening emergency and the sequelae of which can be ciliary body shut down, Vogt's triad, and rarely spontaneous angle reopening.

Other options:

Option B - Virchow triad is used to describe the abnormalities that result in thrombosis. These include endothelial injury, abnormal blood flow, and hypercoagulability of the blood.

Option C - Hutchinson triad is used to describe manifestations of congenital syphilis. It includes interstitial keratitis, Hutchinson teeth, and eighth nerve deafness. Hutchinson teeth are screw-driver or peg shaped small incisors.

Option D - Gaucher's triad is not a named triad. Hypertonia, strabismus, and stridor usually point to a diagnosis of type 2 Gaucher's disease.

Solution to Question 3:

Toxoplasmosis characteristically causes punched-out lesions in the retina.

Toxoplasmosis is a parasitic infestation due to Toxoplasma gondii. It primarily affects the central nervous system and retina. In humans, it occurs as congenital, acquired, or recurrent.

Congenital toxoplasmosis is acquired by the transplacental route. Infants are usually born with inactive disease. Bilateral healed punched-out chorioretinal lesions in the macular area are usually seen. In acquired toxoplasmosis, punched-out lesions similar to congenital toxoplasmosis are seen. In addition, punctate outer retinal toxoplasmosis (PORT) may also be seen.



Other options:

Option A - Posterior staphyloma is used to describe a bulge in the sclera with the choroid behind the equator. On ophthalmoscopy, there is an excavated area with retinal vessels dipping in it.

Option B - Fundal coloboma occurs due to a defect in the closure of the embryonic cleft. It appears as a comet-shaped defect with the apex towards the disc which may or may not be involved. A few choroidal or retinal vessels may also be seen on the surface.

Option C - Congenital rubella typically produces a salt and pepper appearance of the fundus.

Solution to Question 4:

All the above conditions (1, 2, 3, 4) have a familial inheritance.

Retinoblastoma is a malignancy arising from the retina more commonly seen in children. 10% of the cases are familial. These cases have an autosomal dominant mode of inheritance. The remaining 90% occurs sporadically. The clinical presentation includes leucocoria, strabismus, and painful red eye.

Retinitis pigmentosa is a pigmental retinal dystrophy where the photoreceptors, especially rods are affected. Around 50 - 60% of the cases have a familial inheritance. It has multiple modes of inheritance, autosomal dominant, autosomal recessive and X-linked recessive. Night blindness is a characteristic feature. Tubular vision is seen in advanced cases.

Myopia (shortsightedness) is a refractive error in which the light rays from infinity are focused in front of the retina. It is proven to have familial inheritance. Features include defective vision, impaired night vision, and muscae volitantes.

Primary open-angle glaucoma has a normal anterior chamber angle. Family history is a risk factor implicated in the condition. Symptoms include headache, scotoma, and delayed dark adaptation.

Solution to Question 5:

Mycobacterium leprae is the most likely causative organism based on the clinical features given which points to ocular manifestations of leprosy. These include corneal anesthesia, lagophthalmos, corneal ulceration, episcleritis, and iris atrophy. Patients can also present with cataract, glaucoma, and madarosis.

Lagophthalmos is the inability to close the eyelids. This can be due to the involvement of the zygomatic branch of the facial nerve (as in leprosy). Other causes of lagophthalmos include extreme proptosis and symblepharon. The decreased corneal sensation seen is due to the involvement of the ophthalmic branch of the trigeminal nerve which innervates the cornea.

Other options:

Option A - Acanthamoeba castellanii can lead to acanthamoeba keratitis which presents with pain, lacrimation, photophobia, and foreign body sensation. It results from direct corneal contact with infected material. Contact lens users are at an increased risk. Radial keratoneuritis is pathognomonic.

Option C - Onchocerca volvulus causes onchocerciasis commonly seen in African countries. Symptoms include sclerosing keratitis, uveitis, chorioretinitis, and optic neuritis. Ultimately it can lead to optic atrophy.

Option D - Chlamydia trachomatis serovars A, B, Ba, and C are responsible hyperendemic (blinding) trachoma. The source of infection is infected conjunctival discharge. If there is no secondary bacterial infection, symptoms include lacrimation, scanty mucoid discharge and foreign body sensation. Conjunctival congestion and follicles, papillary hyperplasia, Herbert follicles (seen in the limbal area), and progressive pannus are the characteristic features. Serovars D to K causes neonatal and adult inclusion conjunctivitis. Neonates present with conjunctival discharge, chemosis and swollen lids. Adult inclusion conjunctivitis more commonly seen in sexually active adults present with foreign body sensation, mild photophobia, and mucopurulent discharge.

Solution to Question 6:

The clinical vignette is suggestive of optic neuritis. Sudden, progressive, and profound visual loss associated with relative afferent pupillary defect (RAPD) is characteristic of optic neuritis. MRI brain & orbit, screening for circulating myelin oligodendrocyte glycoprotein (MOG) & neuromyelitis optica (NMO) antibodies, and blood sugar levels are the next best investigations for optic neuritis.

A brain MRI is recommended in every patient after the first attack of optic neuritis. The 15-year cumulative probability of developing multiple sclerosis after an attack of optic neuritis is about 50%. It may remain an isolated event in some patients.

Neuromyelitis optica (NMO) is a severe type of optic neuritis which may occur bilaterally and is associated with myelitis. Detection of antibodies like MOG & NMO (also known as aquaporin-4 antibody) is diagnostic of the condition.

Optic neuritis is primarily treated with steroids. The blood sugar levels must be checked as steroids can worsen the glycemic control in individuals with DM which may precipitate hyperglycemia.

Optic neuritis comprises inflammatory and demyelinating disorders of the optic nerve. It can be idiopathic, hereditary (Leber's disease), or due to demyelinating disorders such as multiple sclerosis or neuromyelitis optica (Devic's disease). Anatomically, it can be papillitis, neuroretinitis, or retrobulbar neuritis. It is more common in women.

Other symptoms include lowered dark adaptation and visual obscuration in bright light. Transient obscuration of vision on exposure to light (Uhthoff's symptom) and impaired depth perception (Pulfrich's phenomenon) may also be seen.

Marcus Gunn pupil which points to RAPD is a characteristic sign. It is elicited by the swinging flashlight test. One eye is illuminated with the flashlight and the response is noted. The same is done for the other eye. This is repeated multiple times (swinging). The normal response is constriction in both eyes. If there is RAPD in one eye, it will dilate when light is shone on it.



Solution to Question 7:

Roenne's nasal step is not a feature of the optic disc in glaucoma.

Roenne's nasal step is a visual field defect. It is the result of two arcuate scotomas running in different arcs and meeting at the horizontal meridian. This forms a right-angled defect in the nasal side of the visual field.

Early glaucomatous changes of the optic disc include an increase in cup-disc ratio (C/D ratio) to more than 0.4, splinter hemorrhages near the optic disc margin, and atrophy of the retinal nerve fiber layer.

Advanced glaucomatous changes of the optic disc include a marked increase in C/D ratio (0.7 or more), thinning of the neuroretinal rim, nasal shifting of the retinal vessels or bayonetting, and pulsations of retinal arterioles at the disc margin.

If the damage progresses, all neural tissue is destroyed and the optic nerve appears white and deeply excavated.

Solution to Question 8:

The finding seen in the image is hypopyon.

Hypopyon is a purulent whitish exudate consisting of inflammatory cells in the inferior part of the anterior chamber. It can be associated with bacterial or fungal corneal ulcers and HLA-B27-associated anterior uveitis.

In bacterial corneal ulcers, the hypopyon is sterile. In these lesions, the outpouring of polymorphonuclear cells is due to the toxins and not due to actual invasion by bacteria. The hypopyon is absorbed when the ulcerative process is controlled.

In fungal corneal ulcers, the hypopyon is not sterile. The fungi can penetrate into the anterior chamber without perforation.

The hypopyon associated with anterior uveitis is immobile due to high fibrin content. In Behcet disease, there is minimal fibrin content, so the hypopyon shifts according to the patient's head position.

Treatment of hypopyon involves correcting the underlying conditions.

Other options:

Option A: Hyphema is a hemorrhage in the anterior chamber. It commonly occurs after a blunt ocular injury. The bleeding is from the iris root or ciliary body face. The blood settles inferiorly forming a fluid level.

Option B: Aqueous flare refers to the haziness of the normally clear fluid in the anterior chamber. It reflects the presence of protein due to the breakdown of the blood-aqueous barrier. The presence of flare indicates active inflammation.

Option C: Arlt line is the scarring on the conjunctiva in the area of sulcus subtarsalis. It can be a single line, irregularly shaped, or star-shaped. Common causes are trachoma, healed traumatic wounds, or surgical scars.

Solution to Question 9:

According to the National Blindness and Visual Impairment Survey 2015-19, the most common cause of moderate-severe visual impairment (MSVI) is untreated cataract.

MSVI is defined as presenting visual acuity of <6/18 to 3/60 in the better eye with available correction.

Most commonly, cases of visual impairment and blindness are more prevalent in patients aged 50 years and above.

Other options:

Option A: Refractive errors are the leading cause of MSVI in patients aged 49 years and below.

Options C and D: Glaucoma and diabetic retinopathy account for a very low number of visual impairment cases in patients aged 50 years and above.

Solution to Question 10:

In optic neuritis, relative afferent pupillary defect (RAPD) or Marcus Gunn pupil is seen. It is elicited by the swinging flashlight test. Normally, both pupils remain constricted. In the presence of RAPD, both pupils dilate on moving the flashlight from the normal to the abnormal eye. This paradoxical response of the pupil to light is called Marcus Gunn pupil.

Other options:

Option B: Pain is aggravated by ocular movement, especially in the upward or downward direction due to attachment of the superior rectus fibers to the dura mater.

Option C: Multiple sclerosis, and not multiple myeloma, is a known demyelinating cause of optic neuritis.

Option D: Oral corticosteroid therapy alone is contraindicated because it does not improve the outcome and is associated with an increased risk of new attacks of optic neuritis. Intravenous methylprednisolone can be used if the patient shows radiological evidence of multiple sclerosis.

Solution to Question 11:

The most common ocular manifestation of COVID-19 is follicular conjunctivitis.

Hemorrhagic conjunctivitis, dry eye disease, episcleritis, chemosis, and photophobia may also occur, but less frequently.

The most prevalent ocular symptoms are dry eye or foreign body sensation, redness, tearing, itching, eye pain, and discharge.

Intraocular complications include retinal artery occlusions and hemorrhages, in addition to cotton-wool spots caused by complement-mediated thrombotic angiopathy.

The ACE-2 receptors present in the cornea and conjunctiva make the ocular surface a potential target tissue for SARS-CoV-2.

Sometimes, conjunctivitis may appear as the only sign and symptom of COVID-19. Patients should, therefore, undergo a nasopharyngeal RT-PCR test.

Solution to Question 12:

The condition given in the image is ankyloblepharon. It is the adhesion between the margins of the upper and lower eyelids.

It may occur as a congenital anomaly or as acquired adhesions after healing of chemical burns, ulcers, or traumatic wounds of the lid margins.

Treatment is the excision of the adhesions between lid margins. Epithelial grafts should be used if adhesions extend to angles.

Other options:

Option A: Symblepharon is the adhesion between the palpebral and bulbar conjunctiva in which the eyelids become adherent to the eyeball.

The image given below shows symblepharon.



Option C: Congenital blepharophimosis is a condition in which the extent of the palpebral fissure is decreased. It can be congenital or acquired and is due to a vertical skin fold at the lateral canthus. No treatment is required. In some cases, canthoplasty is done.

Option D: Congenital ptosis is associated with developmental defect and weakness of the levator palpebrae superioris (LPS) muscle. Drooping of one or both of the upper eyelids is seen since birth. Lid lag on downward gaze is present. Treatment involves surgical correction.

Solution to Question 13:

The correct option is Option C: 1-b, 2-a, 3-d, 4-c.

Color vision is tested with Ishihara plates, which is a pseudoisochromatic chart. There are patterns of colored and grey dots that reveal one pattern to persons with normal color vision and another to those with the color deficiency.

The below image shows Ishihara chart.



An applanation tonometer is used to measure the intraocular pressure. It is based on the Imbert-Fick law, which states that the pressure inside a sphere(P) is equal to the force (W) required to flatten its surface divided by the area of flattening (A).

$P = W/A$

The following is an image of the applanation tonometer:



Perimetry is the procedure of estimating the extent of the visual fields. There are 2 types of perimetry: kinetic and static perimetry.

In kinetic perimetry, a target of given luminance is moved from a non-seeing area in the periphery towards the center, where it is perceived. In the static method, the location of a stimulus remains fixed, with the intensity increased until it is seen by the subject (threshold is reached) or decreased until it is no longer detected.

The tangent screen perimetry evaluates the central 30 degrees field at 1 m, thereby testing the central vision.

Solution to Question 14:

Atropine is contraindicated in acute congestive glaucoma.

Acute congestive or angle closure glaucoma is caused by the opposition of the peripheral iris against the trabecular meshwork. This obstructs the aqueous outflow by closing an already narrow-angle of the anterior chamber leading to an acute increase in intraocular pressure(IOP). It can be precipitated by dim light, mydriatics like atropine.

It presents as sudden onset of severe unilateral eye pain or a headache associated with blurred vision, rainbow-colored halos around bright lights, nausea, and vomiting. The physical exam will reveal a fixed midpoint pupil and a hazy or cloudy cornea with marked conjunctival injection.

Management of acute condition includes rapid lowering of IOP by mannitol or glycerol, acetazolamide, and topical antiglaucoma drugs. Definitive management is done by laser peripheral iridotomy or filtration surgery.

Atropine is indicated as cycloplegic to reduce pain and prevent synechiae in corneal ulcers and acute anterior uveitis. It is the initial management in malignant glaucoma.

Solution to Question 15:

The exposure of the retina to light causes hyperpolarization and a decreased release of neurotransmitters in rods and cones.

When photoreceptors are exposed to light, a series of biochemical reactions result in the generation of electrical charge. This phenomenon of conversion of light energy into electrical impulse is known as phototransduction.

In dark, cyclic GMP (cGMP) gated channels are open leading to a steady release of neurotransmitters (glutamate). When light falls on the retina,11-cis-retinal is converted to all-trans-retinal. This activates transducin which is a g protein-coupled receptor with phosphodiesterase activity breaking down cGMP. With a decrease in cGMP levels sodium channels close causing hyperpolarization. Hyperpolarization reduces glutamate release, which leads to the formation of the action potential.





Solution to Question 16:

The incorrect about keratoconus is Haab's striae. Haab's striae are seen in cases of buphthalmos where the intraocular pressure is raised and the eyeball is enlarged as a whole. The enlarged cornea is associated with tears in Descemet's membrane called Haab's striae.

Keratoconus or the conical cornea is a bilateral non-inflammatory progressive condition of the axial part of the cornea. Central or paracentral thinning of the cornea and apical protrusion of the anterior cornea are pathognomonic. This is attributed to mucopolysaccharide and collagen tissue synthesis defects.

Clinical features of the condition include defective vision due to progressive myopia and irregular astigmatism. Light sensitivity and ocular irritation may also be present. Complications include acute hydrops following the rupture of Descemet's membrane.

Slit-lamp examination reveals Fleischer's ring at the base of the cone formed. Deep, vertical stromal striae (Vogt lines) that disappear with external pressure are characteristic. Retinoscopy shows a scissoring reflex. The bulging of the lower lid in the downgaze, called Munson's sign, is shown in the image below.

The severity of the condition is graded based on keratometry. Corneal topography is a diagnostic modality with the highest sensitivity for early cases.

Intacs or intracorneal ring segments are useful in moderate cases due to their flattening effect. Rigid gas-permeable contact lenses, corneal collagen cross-linking with riboflavin, and keratoplasty are the other treatment modalities.

Solution to Question 17:

The angle subtended for a patient with 6/6 vision is 5 minutes of arc.

Visual acuity is a measure of the spatial resolution of the eye or an estimation of its ability to discriminate between two points.

Snellen's chart consists of a series of letters arranged in lines, each diminishing in size. Each whole letter will subtend an

the angle of 5 minutes at the nodal point of the eye at the given distance.

In the Snellen chart, the largest letter will subtend an angle of 5 minutes at the nodal point if it is 60 m from the eye. Those in the subsequent lines will subtend an angle of 5 minutes if they are 36, 24, 18, 12, 9, and 6 m from the eye.

The patient is placed at 6 m and asked to read the visual acuity chart from the largest letters downwards to the smallest that can be visualized with each eye separately. A person with normal 6/6 vision sitting at a distance of 6 m from the types should be able to read every letter from the top to the end of the 6 m line. So the angle subtended by the 6 m line at the eye a distance of 6 meters will be 5 minutes of arc.

If a patient can only read up to the 12m line, his distant vision is defective. His visual acuity will be recorded as 6/12. The numerator denotes the distance at which the patient is able to read the particular line and the denominator denotes the distance at which a person with normal visual acuity will be able to read the given line.



Solution to Question 18:

The recurrence rate of 95% is not a feature of pterygium. The recurrence rate of pterygium is around 80% after simple surgical excision.

Pterygium is a wing-shaped, triangular fibrovascular ingrowth (Option B) of degenerative conjunctival tissue that can encroach onto the cornea. It is a response to prolonged exposure to

ultraviolet rays, dry heat, etc. It has 3 parts mainly, a head (apical part on the cornea), a body (the region between the limbus and canthus), and a cap(semilunar white infiltrate in front of the head)(Option C).. Some authors mention the neck as the constricted part near the limbal area.

Patients may complain of itching and grittiness due to Dellen (localized drying effect). If the pterygium encroaches onto the pupillary area, it may interfere with vision and also induce astigmatism. Stocker's line may be associated with a pterygium. It is a linear deposition of iron seen in the corneal epithelium, anterior to the head of a pterygium. Pterygium is of 2 types (Option D):

• Progressive which is thick, fleshy, and vascular.

• Regressive which is thin, atrophic, and attenuated.

This condition can be treated symptomatically with tear substitutes for dry eye symptoms and patients are advised to avoid ultraviolet rays. Surgical excision with free conjunctival limbal autograft is the preferred treatment to avoid recurrence.

The recurrence rate of pterygium is around 80% after simple surgical excision. Excision together with adjunctive therapies such as mitomycin C or 5-fluorouracil can reduce the risk of recurrence to approximately 10%

The image below shows a pterygium.



Solution to Question 19:

Krukenberg spindle is not a feature of complicated cataracts. The slit lamp examination in pigmentary glaucoma shows a pattern of pigment deposition on the back of the cornea-endothelium called Krukenberg spindles.

Complicated or secondary cataracts develop secondary to inflammatory/degenerative conditions of the lens or other parts of the eye like iridocyclitis and ciliary body tumors which lead to disturbances in the nutrition of the lens. These cataracts present with a breadcrumb appearance and polychromatic luster.

Even in the early stages, vision is impaired. Inflammation of the posterior segment leads to the characteristic posterior subcapsular cataract while the anterior segment inflammation causes opacification throughout the cortex.

Solution to Question 20:

The oculocephalic reflex or doll's eye reflex is used to assess the integrity of the cranial nerves III, VI, and VIII.

It is a brainstem reflex examined by observing the the motion of eyes during the sudden rotation of the head of the patient by the examiner. It is done in both directions laterally, followed by flexion and extension. In a normal positive reflex, the eyes move conjugately in the opposite direction of the movement of the head to maintain their position in space.

The reflex has afferents in the vestibular nerve (VIII) that relay to the brainstem nuclei and bring about conjugate eye movements through the oculomotor nerve (III) and abducens nerve (VI).

A negative reflex occurs in any of the above cranial nerve palsies or brainstem injuries, where the eyes show no movement or movement in the same direction as head movement. This is used to assess brainstem reflexes in a comatose patient as a criterion of brain death.

Solution to Question 21:

The structures affected in this patient are A, B, C and D.

Cavernous sinus thrombosis (CST) is a septic thrombus formed in the cavernous sinus. It spreads from infection of the sinuses, teeth, ears, nose, and skin of the face.

It presents as fever, severe headache, supraorbital pain, seizures, and edema in the mastoid region. Eye signs include proptosis, chemosis, and diplopia due to ophthalmoplegia from the involvement of oculomotor (CN-III), trochlear (CN-IV), and abducens nerve (CN-VI). Absence of corneal sensation due to the involvement of the ophthalmic division of trigeminal nerve (CN-Va). Ipsilateral ptosis, dilated pupil, and absence of direct and consensual pupillary right reflex due to oculomotor nerve palsy. Pulsatile proptosis occurs with the involvement of the internal carotid artery.

Investigation using brain magnetic resonance imaging or magnetic resonance venography should be done urgently. Following early diagnosis, management of cavernous sinus thrombosis typically includes intravenous broad-spectrum antibiotics, analgesics, anti-inflammatory drugs, and anticoagulants.

Cavernous sinus



©MARROW

Ophthalmology NEET 2022

Question 1:

What is the diagnosis?



a) Intraocular foreign body

b) Pseudoexfoliation syndrome

c) Ocular trauma

d) Vossius ring

Question 2:

A patient presents with a history of penetrating injury to the eye. A diagnosis of sympathetic ophthalmitis was confirmed. Which of the following will be seen?

a) Acute anterior uveitis

b) Pars planitis

c) Panuveitis

d) Chronic anterior uveitis

Question 3:

A diabetic patient presents to you with visual acuity of 6/9 in one eye. Further investigations revealed pre-retinal hemorrhages with neovascularization at the optic disc. What is the next step in management?

a) Focal laser photocoagulation

b) Pan-retinal photocoagulation

c) Grid laser photocoagulation

d) Scleral buckling

Question 4:

A 35-year-old woman is diagnosed with rheumatoid arthritis. What associated complication is shown in the image?



a) Scleromalacia perforans

b) Ciliary staphyloma

c) Coloboma

d) Malignant melanoma

Question 5:

What is the indication of this procedure?



a) Keratoconus

b) Vogt's limbal girdle

c) Keratoglobus

d) Corneal dystrophy

Question 6:

A 3-year-old boy presents with mental retardation and an inability to walk. The funduscopy image is given below. What is the most likely diagnosis?



a) Tay-Sachs disease

b) Hunter disease

c) Hurler syndrome

d) Gaucher disease

Answer Key

Question No.	Correct Option
1	b
2	c
3	b
4	a
5	a
6	a

Detailed Explanations

Solution to Question 1:

The diagnosis based on the image given is pseudoexfoliation syndrome.

Pseudoexfoliation syndrome is an age-related degenerative disorder, more commonly seen in women. It is marked by the deposition of exfoliative material on the iris, ciliary region, and lens capsule. They appear as flakes on the anterior lens capsule and edge of the iris.

The material can get collected in the anterior chamber angle and impede the aqueous humor drainage. Patients affected by this condition are at an increased risk of developing open-angle glaucoma. Rarely, angle-closure glaucoma can also be seen. Other ocular symptoms include iris sphincter atrophy, anisocoria, and heterochromia. Systemic manifestations due to abnormal fibrillary deposition may also be seen.

Other options:

Option A - Common intraocular foreign bodies include glass particles or metal chips. It can lodge in various sites like the anterior chamber, lens, posterior chamber, and vitreous. This can lead to cataract formation, retinal hemorrhages, and tears. Organic foreign bodies can cause infection. Iron foreign bodies can cause siderosis while copper ones can result in chalcosis. An entry wound can be identified with topical fluorescein. In cases of metallic intraocular foreign bodies, MRI is contraindicated. CT scan can be done instead.



Option C - Ocular trauma can be a closed-globe or open-globe injury. Closed-globe injury can vary from simple corneal abrasions to widespread intraocular damage. Open-globe injuries include globe rupture or laceration.

A corneal abrasion highlighted with fluorescein staining is shown below.



Option D - Vossius ring is used to describe a ring-shaped pigment dispersal observed on the anterior surface of the lens due to the deposition of brown amorphous pigment granules. It is close to the miosed pupil in diameter. The force of the blow drives the cornea and iris backward, creating an impression of the iris on the lens.



Solution to Question 2:

Panuveitis is the usual clinical manifestation seen in sympathetic ophthalmitis. Panuveitis is inflammation of the whole uvea.

Sympathetic ophthalmitis is a granulomatous pan-uveitis that usually occurs after penetrative trauma. It can also develop after intraocular surgery. Both the injured eye (exciting eye) and fellow eye (sympathizing eye) develop panuveitis. Trauma to the ciliary region (dangerous zone) poses an increased risk.

It is a type 4 hypersensitivity reaction. According to the allergic theory, the uveal pigment in the injured eye acts as an allergen that causes uveitis in the other eye. Dalen-Fuchs nodules (proliferation of pigment epithelium), exudative retinal detachment, and optic disc swelling can be seen on fundoscopy. A sunset glow appearance can persist due to residual chorioretinal scarring.

Dalen-Fuchs nodules are depicted in the image below.



The exciting eye shows features of low-grade uveitis such as ciliary congestion, lacrimation, and tenderness. The sympathizing eye is usually involved 4-8 weeks later. It has a prodromal stage with photophobia, ciliary congestion, and keratic precipitates. In the fully developed stage, symptoms of acute iridocyclitis supervene.

Prophylactic measures to be taken before the condition sets in include topical or systemic steroids and antibiotics. Early excision of the injured eye is carried out if it cannot be salvaged. Once sympathetic ophthalmitis has already set in, enucleation of the injured eye must be done. Atropine, immunosuppressants, and corticosteroids should also be administered.

Other options:

Options A and D: Anterior uveitis is inflammation of the iris up to pars plicata. Keratic precipitates are characteristic of the condition. Acute anterior uveitis, which can last for 6 weeks to 3 months, presents with pain, photophobia, and lacrimation. Chronic anterior uveitis lasts for longer than 6 months. Anterior uveitis is associated with juvenile idiopathic arthritis, sarcoidosis, and ankylosing spondylitis.

The image below shows keratic precipitates.



Option B: Pars planitis or intermediate uveitis is an inflammatory condition that affects the pars plana of the ciliary body, the peripheral retina, the choroid, and the vitreous base. Snowbanking is the hallmark feature. It is a grey-white fibrovascular plaque formed due to the coalescence of snowball opacities over the pars plana. Pars planitis can be seen in multiple sclerosis, sarcoidosis, and syphilis.

Solution to Question 3:

The clinical vignette is suggestive of proliferative diabetic retinopathy with high-risk characteristics and the next step in management is pan-retinal photocoagulation.

Diabetic retinopathy is used to describe retinal changes seen in diabetic patients. It is classified into the following types:

- Non-proliferative diabetic retinopathy (NPDR)
- Proliferative diabetic retinopathy (PDR)
- Diabetic maculopathy
- Advanced diabetic eye disease

Fundoscopy in NPDR reveals hard exudates, cotton-wool spots, and microaneurysms. Hard exudates are small, discrete yellowish, waxy areas with crenated margins.



Proliferative diabetic retinopathy is marked by NVD or NVE which is the hallmark of the condition. Condensation of connective tissue surrounding the newly formed vessels results in the formation of a fibrovascular epiretinal membrane. Vitreous detachment and hemorrhage may also occur. Preretinal hemorrhages are considered a high-risk characteristic. In addition, changes seen in NPDR may also be present.

The condition usually develops 25 years after the onset of the disease. Hence, it is more commonly seen in patients with type I diabetes.

The image below shows NVD.

PDR with high-risk characteristics is managed with pan-retinal photocoagulation. Laser treatment helps by inhibiting the release of vascular endothelial growth factors (VEGF). Anti-VEGFs are recommended before pan-retinal photocoagulation in patients with PDR.

Diabetic maculopathy can be classified as:

• Focal exudative maculopathy: Microaneursysms, hard exudates, and hemorrhages can be seen.

• Diffuse exudative maculopathy: Diffuse retinal edema with few hard exudates.

• Ischemic maculopathy: Microaneursysms, hemorrhages, few hard exudates. Fluorescein angiography reveals areas of non-perfusion.

• Mixed maculopathy: Features of both exudative and ischemic maculopathy.

An FFA image showing features of macular ischemia is given below.



Other options:

Option A: Focal laser photocoagulation is the treatment of choice for focal diabetic macular edema (DME) that does not involve the fovea.

Option C: Grid laser photocoagulation can be used in diffuse DME not responding to anti-VEGFs and intravitreal steroids.

Option D: Scleral buckling can be used in cases of advanced PDR with tractional retinal detachment.

Solution to Question 4:

Scleromalacia perforans is the associated complication of rheumatoid arthritis that is shown in the image.

Scleromalacia perforans is a specific type of necrotizing scleritis without inflammation, characterized by progressive scleral thinning. The uveal tissue appears to shine through it. Spontaneous perforation is a rare complication of the condition. It is typically seen in elderly women with long-standing rheumatoid arthritis. A yellowish patch of melting sclera can be seen. This is due to the obliteration of blood supply.

Clinical features include pain, photophobia, and lacrimation. The affected eye appears red. Medical management includes steroids and immunosuppressants. In severe cases, surgical treatment by scleral patch graft may be done.

Other options:

Option B - Ciliary staphyloma is used to describe the bulge of an area of weak sclera lined by the ciliary body. Common causes include scleritis, developmental glaucoma, and end-stage primary or secondary glaucoma. Thinning of the sclera, secondary to perforating injury, can also result in the condition.



Option C - Coloboma is a congenital anomaly, that develops due to a defect in the closure of the embryonic cleft. It is characterized by poorly developed uveal tissue. In some cases, it might be deficient. The lens, choroid, and retina may also be involved.

The image below shows coloboma of the right eyelid.

Option D - Malignant melanoma of the conjunctiva appears as a mass near the limbus. It can be pigmented or non-pigmented. It is usually seen in elderly patients. The choroid, iris, ciliary body, and eyelids can also develop the tumor.

The image below shows malignant melanoma of the choroid.

Solution to Question 5:

The given image shows the placement of Intacs, which is done for keratoconus.

Keratoconus or conical cornea is a non-inflammatory condition of the axial cornea. Central or paracentral thinning of the cornea and apical protrusion of the anterior cornea are pathognomonic. This is attributed to mucopolysaccharide and collagen tissue synthesis defects.

Clinical features of the condition include defective vision, which develops due to progressive myopia and irregular astigmatism. Light sensitivity and ocular irritation may also be present. Complications include acute hydrops due to rupture of Descemet's membrane.

Slit-lamp examination reveals Fleischer's ring at the base of the cone formed. Deep, vertical stromal striae (Vogt lines) that disappear on external pressure are characteristic. Retinoscopy

shows a yawning reflex.

The image below shows Munson's sign i.e. bulging of the lower lid on looking down.

The severity of the condition is graded based on keratometry. Corneal topography is a diagnostic modality with the highest sensitivity for early cases.

Intacs or intracorneal ring segments are useful in moderate cases, due to their flattening effect. Rigid gas permeable contact lenses, corneal collagen cross-linking with riboflavin, and keratoplasty are the other treatment modalities.

Other options:

Option B - Vogt's limbal girdle is a degenerative corneal disease that is more commonly seen in women aged ≥40 years. A white crescentic limbal band can be seen at 9 and/or 3 'o'clock position, more frequently on the nasal side. In type I, a swiss-cheese hole pattern with a clear zone separating it from the scleral margin can be observed. Type II is more common and lacks a clear zone and holes.

Option C - Keratoglobus is a familial disorder, marked by thinning and hemispherical protrusion of the entire cornea. It can be congenital or acquired. Globular protrusion is seen here, in contrast to the conical ectasia seen in keratoconus. Intrastromal ring segments and cross-linking can be used for management.

Option D - Corneal dystrophy is used to describe a group of non-inflammatory conditions, with bilateral opacification of the cornea with no vascularisation. These are divided into epithelial, Bowman layer, stromal, Descemet membrane, and endothelial dystrophies. Most conditions have genetic abnormalities.

Solution to Question 6:

The given clinical scenario, along with the fundoscopy image showing a macular cherry-red spot, indicates the most likely diagnosis to be Tay-Sachs disease.

Tay-Sachs disease is a lysosomal storage disease caused by the deficiency of β-hexosaminidase. It can present during infancy with macrocephaly, increased startle reaction, gradual loss of motor skills, and a macular cherry-red spot. Affected children usually die by the age of 10–15 years.

It is of two types—juvenile-onset and adult-onset. The symptoms of the juvenile-onset form include ataxia and dementia. Dysarthria, lower motor neuron signs, and psychiatric disorders are common in the adult-onset type.

Other options:

Options B - Hunter disease (mucopolysaccharidosis II) is an X-linked disorder that arises due to the deficiency of iduronate 2-sulfatase. Short stature, joint stiffness, and intellectual disability are the features.

Option C - Hurler syndrome (mucopolysaccharidosis I) is an autosomal recessive condition that affects multiple organs. Presentation is usually at 6–24 months with corneal clouding, coarse facial features, joint stiffness, and skeletal dysplasia. Glaucoma, retinal degeneration, and nasal discharge are the other features. The condition is generally fatal.

Option D - Gaucher disease is a glycolipid storage disorder that can present with hepatosplenomegaly, mental retardation, and erosion of long bones. It is caused by the deficiency of β-glucosidase. The Gaucher cell has a wrinkled-paper appearance due to intracytoplasmic inclusions, which is a characteristic feature.

Ophthalmology NEET 2023

Question 1:

A female presents with loss of vision in the right halves of both eyes. Where is the lesion located in the optic pathway?

a) Left optic tract

b) Optic radiation

c) Optic chiasma

d) Right optic tract

Question 2:

A 65-year-old diabetic woman presents with painless loss of vision. The ocular findings are shown in the image below. What is the likely diagnosis and management for this patient?



a) Zonular cataract and lensectomy

b) Rosette cataract and sugar control

c) Sunflower cataract and lesion extraction

d) Immature senile cataract and phacoemulsification with IOL

Question 3:

Which of the following refractive errors is associated with the image given below?



a) Presbyopia

b) Astigmatism

c) Hypermetropia

d) Myopia

Question 4:

A 5-year-old boy presents with a small nodule on the sclera as shown in the image below. What is the likely diagnosis?

a) Pterygium

b) Pinguecula

c) Dermolipoma

d) Dermoid

Question 5:

The eye examination of a 20-year-old male shows a golden ring on the iris. What is the next investigation to diagnose the condition?

a) Alpha-1-antitrypsin

b) Alpha-fetoprotein

c) Serum iron

d) Serum ceruloplasmin

Question 6:

What is the dye and filter used to diagnose corneal ulcers?

a) Lissamine dye - green filter

b) Lissamine dye - cobalt blue filter

c) Fluorescein dye - visualized under green filter

d) Fluorescein dye - visualized under cobalt blue filter

Question 7:

An infant is brought with complaints of excessive watering of the eyes and photophobia. The image is given below. What is the likely diagnosis?

a) Congenital glaucoma

b) Congenital cataract

c) Mucopolysaccharidosis

d) Ophthalmia neonatorum

Question 8:

A 60-year-old patient complains of decreased distant vision but now he does not use spectacles for near vision. The image of his ocular examination is given below. What is this type of refractive error called?



a) Index myopia

b) Axial myopia

c) Curvatural myopia

d) Positional myopia

Answer Key

Question No.	Correct Option
1	a
2	d
3	b
4	d
5	d
6	d
7	a
8	a

Detailed Explanations

Solution to Question 1:

Visual loss in the right halves of both eyes is suggestive of right homonymous hemianopia. The site of the lesion in this case is the left optic tract.

Here, vision is lost in the right half of both eyes, which means:

• in the left eye, the nasal visual field is lost, and

• in the right eye, the temporal visual field is lost.

It means left temporal and right nasal fibers are destroyed, which leads to right homonymous hemianopia. Anatomically, this can be explained by the fact that each optic tract contains fibers from the ipsilateral temporal half of the retina and the contralateral nasal half of the retina.

Optic tract lesions can occur due to pituitary adenomas (most common), craniopharyngioma, tumors of the optic thalamus, large aneurysm of superior cerebellar or posterior cerebral arteries, demyelinating diseases, and infarction.

Other options:

Option B: Optic radiation lesions result in contralateral homonymous hemianopia, which means that the visual loss would occur in the left halves of both eyes rather than the right halves, as described in the question.

Option C: Lesions at the optic chiasma typically cause bitemporal hemianopia, which involves visual loss in the outer (temporal) halves of the visual field in both eyes. This is not consistent with the patient's described symptoms.

Option D: A lesion in the right optic tract would result in left homonymous hemianopia, causing visual loss in the left halves of both eyes. This does not match the patient's presentation of vision loss in the right halves of both eyes.



Solution to Question 2:

The given clinical scenario and the above image are suggestive of an immature senile cataract and the appropriate treatment is phacoemulsification with posterior intraocular lens implantation.

An immature senile cataract is characterized by partial opacification of the lens. The cuneiform and cupuliform patterns of opacification are seen and the lens appears greyish-white. Iris shadow is visible. Slit-lamp examination shows areas of the normal and cataractous cortex.

The treatment of choice for immature senile cataracts is treated by extracapsular cataract extraction by phacoemulsification and foldable posterior chamber intraocular lens implantation.

Other options:

Option A: Zonular cataract is a developmental type of cataract in which opacification occupies a discrete zone in the lens. It is the most common type of congenital cataract presenting with visual impairment. Lensectomy is the treatment of choice for congenital cataracts.

Option B: Rosette cataract presents after close globe injuries (blunt trauma) resulting in damage to the lens fibers and capsule resulting in hydration of lens fibers by aqueous humor. This results in opacification of the posterior subcapsular cortex along the posterior sutures. It presents as a flower-shaped opacification. Iridodialysis occurs due to dehiscence of the iris from the ciliary body at its root. The pupil becomes D-shaped. Surgery is recommended in case of visual impairment.

Option C: Sunflower cataract is seen in Wilson disease due to the deposition of cuprous oxide in the anterior lens capsule and subcapsular cortex in a stellate pattern. D-penicillamine is an effective treatment modality.

Solution to Question 3:

The given image is a Sturm's conoid. It represents the configuration of a bundle of light rays formed by a regularly astigmatic (toric) surface and helps in understanding the optics of regular astigmatism.

Astigmatism is a type of refractive error that occurs when light rays enter the eye at different planes and converge at different points forming focal lines.

When light passes through a regularly astigmatic surface (surface with toric curvature), vertical and horizontal rays converge at different points and form two focal lines. The shape of a bundle of light rays at different points is as follows:

For the given Sturm's conoid,

• At point A, a horizontal oval is formed because vertical rays converge more rapidly than horizontal rays.

• At point B (first focus), vertical rays come to a focus, while horizontal rays are still converging; so, it will be a horizontal straight line.

• At points C, D, and E, the vertical rays are diverging, and horizontal rays are still converging.

• At point D, the vertical rays have diverged as much as the horizontal rays have converged; so, we get a perfect circle called the circle of least diffusion.

• At point E, a vertical oval is formed.

• At point F (second focus), a vertical straight line is formed

• At point G and beyond, both rays are diverging, and we always get a vertical oval/ prolate ellipse.

• The focal interval of Sturm refers to the distance between two foci (B and F).

The image is clearest or least confusing as point D, the circle of least confusion/ circle of diffusion. The circle of diffusion lies at the mid-point of this interval.



• Option A: Presbyopia is a refractive error caused by the loss of elasticity of the crystalline lens, leading to difficulty in focusing on near objects. Presbyopia is not associated with Sturm's conoid.

• Option C: Hypermetropia, or farsightedness, is a refractive error in which the eye does not focus incoming light properly, causing near objects to appear blurry. It is not associated with Sturm's conoid.

• Option D: Myopia, or nearsightedness, is a refractive error in which distant objects appear blurry because the eye focuses light in front of the retina rather than on it. Myopia is not associated with Sturm's conoid.

Solution to Question 4:

A dermoid is a likely diagnosis based on the ocular finding and the given image.

An orbital dermoid is a choristoma (a mass of histologically normal tissue in an abnormal location) formed due to subcutaneous displacement of the ectoderm.

It is lined by keratinized stratified squamous epithelium and contains dermal appendages like sweat glands, sebaceous glands, and hair follicles. It can be located anteriorly (superficial) or posteriorly (deep) to the orbital septum.

Clinical features: Superficial dermoid presents in infancy as a painless nodule in the superotemporal part of the orbit whereas deep dermoid presents in adulthood as a gradually protruding eye or acutely as an inflamed orbit. CT imaging would show a well-circumscribed cystic lesion.

Treatment is by excision in toto.

Other options:

Option A: Pterygium is a triangular fibrovascular ingrowth of degenerative bulbar conjunctival tissue on the limbus as a result of chronic exposure to UV rays.

Option B: Pinguecula is an asymptomatic yellow-white mound of degenerated conjunctival stroma seen adjacent to the limbus. It is more common on the nasal limbus.

Option C: Dermolipoma presents in adulthood as a subconjunctival yellowish mass near the outer canthus of the eye. It contains dermal appendages and fatty tissue.

Solution to Question 5:

The given image shows a Kayser-Fleischer ring, which is seen in Wilson's disease. Low serum ceruloplasmin level is diagnostic of Wilson's disease.

Wilson's disease (hepatolenticular degeneration) is an inherited metabolic disorder resulting due to abnormal copper deposition in tissues (mainly the liver and brain). It is caused by a deficiency of ceruloplasmin (copper-carrying blood protein).

Kayer-Fleischer ring is a brownish-yellow zone of copper deposition that is present in the peripheral Descemet membrane.

The ring is best seen on gonioscopy. Penicillamine therapy may cause the ring to disappear.

Another ocular finding in Wilson's disease is anterior capsular sunflower cataract.

Other options:

Option A: Alpha-1 antitrypsin deficiency leads to lung and liver disease.

Option B: Alpha-fetoprotein is a diagnostic marker in hepatocellular carcinoma.

Option C: Serum iron level is assessed in hemochromatosis which occurs due to increased iron deposition in various tissues. The ocular manifestation includes dry eye and rusty-brown perilimbal and conjunctival discoloration.

Solution to Question 6:

The image given is of a dendritic ulcer seen in recurrent herpetic keratitis. Corneal ulcers can be detected with the help of fluorescein sodium dye and a cobalt-blue filter. The epithelial defect of the cornea appears opaque green with a cobalt blue filter.

Fluorescein staining of the cornea is carried out with a disposable autoclaved filter paper strip impregnated with the dye. The denuded epithelial area seen in corneal abrasions or ulcers appears brilliant green when stained with fluorescein. Under a cobalt blue light, the stained area appears opaque green.

Other options:

Lissamine green dye is used to stain devitalized cells of the cornea and conjunctiva. It is used to diagnose ocular surface disorders, like dry eye.

Solution to Question 7:

The clinical scenario of excessive watering and photophobia in an infant and the given image with corneal enlargement and haze suggest a likely diagnosis of congenital glaucoma.

Primary congenital glaucoma (PCG) is caused by impaired aqueous outflow due to a maldeveloped anterior chamber angle.

It can be classified as

• True congenital glaucoma - IOP is elevated during intrauterine life.

• Infantile glaucoma - manifests before 3 years of age.

• Juvenile glaucoma (least common) - IOP rises between 3 and 16 years of age.

Clinically, the child presents with lacrimation, photophobia, and blepharospasm.

PCG may also present with:

• Corneal haze (due to diffuse edema secondary to raised IOP)

• Buphthalmos (a large eye that results from stretching due to raised IOP before 3 years of age). The thin sclera appears blue due to underlying uveal tissue.

• Haab striae (curvilinear healed breaks in Descemet's membrane)

• Corneal scarring

• Optic disc cupping

Evaluation is done under general anesthesia. Initially, IOP measurement is performed (10–12 mmHg is normal). A corneal diameter measuring greater than 12 mm is highly suggestive of an infant under the age of 1 year.

Management is essentially surgical and goniotomy and/or trabeculectomy may be done.

Other options:

Option B: Congenital cataract occurs due to disturbance in lens growth leading to the development of opaque lens fibers. It can be idiopathic, hereditary, or secondary to metabolic disorders and infections. It presents with leukocoria (white reflex in the pupillary area), wandering eye movement, squint, and nystagmus.

Option C: Mucopolysaccharidoses are inherited metabolic disorders presenting with corneal opacification, pigmentary retinopathy, glaucoma, and optic atrophy.

Option D: Ophthalmia neonatorum presents in neonates with conjunctival discharge and chemosis, and tenderness of the eyeball secondary to infections.

Solution to Question 8:

The most likely diagnosis based on the given clinical scenario and the image is index myopia seen in nuclear senile cataracts.

Index myopia occurs due to an increase in the refractive index of the crystalline lens due to nuclear sclerosis. Nuclear sclerosis is an ageing process in which the lens becomes hard and inelastic resulting in an increase in the refractive index, causing progressive myopia (decreased distant vision). A patient with presbyopia may be able to read again, without the aid of spectacles. This improvement in vision is termed second sight.

Other options:

Option B: Axial myopia is the most common type of myopia resulting due to an increase in the anteroposterior length of the eyeball.

Option C: Curvatural myopia results due to increased curvature of the cornea, lens, or both.

Option D: Positional myopia results due to anterior placement of the crystalline lens in the eye.

Ophthalmology INI-CET 2023

Question 1:

The vestibulo-ocular reflex (VOR) is seen to stabilize the image on the retina while the head moves. What is the gain of VOR (eye velocity/head velocity)?

a) 4

b) 2

c) 1

d) 3

Question 2:

In ocular surface examination and dry eye staining, which of the following order of dyes is used?

a) Fluorescein ->Rose Bengal -> Lisssamine green

b) Rose Bengal -> Lisssamine green -> Fluorescein

c) Lisssamine green -> Rose Bengal -> Fluorescein

d) Fluorescein-> Lisssamine green -> Rose Bengal

Question 3:

Which of the following are direct gonioscopes?

a) 1, 3

b) 2, 4

c) 4

d) 1, 2, 3

Question 4:

On administration of phenylephrine, the following change was noted as shown in the images below. What is the probable diagnosis?



a) Nodular scleritis

b) Nodular episcleritis

c) Tenon's cyst

d) Nodular phlycten

Question 5:

Identify the marked structure in the eye.



a) Eggers line

b) Lens capsule

c) Canal of Hannover

d) Canal of Petit

Question 6:

Which of the following is true about perimetry?

a) Kinetic perimetry is used in quantification and tracking of visual changes in established glaucoma

b) In Static perimetry, stimulus intensity is held constant and moved centrally with detection

c) Visual field defect to be classified as glaucoma, it should have the corresponding optic nerve head abnormalities

d) Early signs include generalised constriction of isopters and baring of blind spot

Question 7:

A man with known history of diabetes has come to your clinic. His vision is 6/6. However, on retinal examination, he is classified to have severe NPDR. Which of the following is not a feature of this condition?

a) Retinal microvascular changes

b) Retinal hemorrhages

c) Disc neovascularisation

d) Venous beading

Question 8:

Identify the filter, the dye and the condition shown below



a) Cobalt green - Fluorescein dye - Corneal degeneration

b) Cobalt blue - Fluorescein dye - Corneal ulcer

c) Cobalt blue - Lissamine green dye - Corneal opacity

d) Cobalt green - Rose Bengal dye - Interstitial keratitis

Question 9:

A professional boxer wants vision correction surgery. Which of the following is not recommended?

a) RK

b) PRK

c) LASIK

d) SMILE

Question 10:

A child went to an ophthalmologist and is found to have a power of -1 D spherical and -2 D cylindrical at 180 degrees. What kind of astigmatism does the child have?

a) Simple astigmatism

b) With the rule astigmatism

c) Against the rule astigmatism

d) Mixed astigmatism

Question 11:

The following instrument is used to measure which of the following?

a) Proptosis

b) Squint

c) Diplopia

d) Stereopsis

Answer Key

Question No.	Correct Option
1	c
2	a
3	b
4	b
5	d
6	d
7	c
8	b
9	a

10	b
11	a

Detailed Explanations

Solution to Question 1:

The gain of the vestibulo-ocular reflex (VOR) is 1, which means there is a compensatory eye velocity equal to the head velocity and in the opposite direction.

VOR prevents the slipping of the retinal images when the head moves; it moves the eyes at the same speed but in the opposite direction as the head. If an object is being pursued with eye and head movement, vision cancels the vestibulo-ocular reflex to prevent the eyes from moving in the opposite direction as the head.

VOR is a physiological vestibular nystagmus that forms the basis of the caloric test. If cold water is poured into the right ear the patient develops left jerk nystagmus (rapid phase towards the left), while the reverse happens with warm water, i.e. patient develops right jerk nystagmus. It can be remembered by the mnemonic 'COWS' (Cold- Opposite, Warm-Same).

Solution to Question 2:

Amongst the given options, the better order of dyes to pick for use in ocular surface examination and dry eye staining is Fluorescein -> Rose Bengal -> Lisssamine green.

Fluorescein staining of the cornea is carried out with a disposable autoclaved filter paper strip impregnated with the dye. The denuded epithelial area seen in corneal abrasions or ulcers appears brilliant green when stained with fluorescein. Under a cobalt blue light, the stained area appears opaque green.

A corneal abrasion highlighted with fluorescein staining is shown below.



Rose Bengal stains dead and devitalized cells including mucous strands on ocular surface. The dye causes a stinging and burning sensation soon after instillation.



Lissamine green stains dead and degenerate cells, but does not stain healthy epithelial cells. It is used to diagnose ocular surface disease. It stains the conjunctiva. It does not cause stinging like Rose Bengal. The Green staining is best observed two minutes after instillation.

The image below shows the lissamine staining of the eye.



Note: The answer to this question is controversial and there is no clearcut evidence based information on the sequence. The above mentioned answer is the best amongst the given options based on consensus.

Solution to Question 3:

Richardson and Koeppe are direct gonioscopes.

A gonioscope is used for the measurement of the angle of the anterior chamber. The types of gonioscopes include:

• Indirect gonioprism: Provides a mirror image of the opposite angle. Indirect goniolenses include Goldmann, Zeiss, and Posner.

• Direct gonioprism: Provides a direct view of the angle. Direct goniolenses include the Koeppe, Richardson, Medical Workshop, Barkan, and Swan–Jacob.

Applications:

• To view the angle of the anterior chamber. The structures visualized in order from front to back are:

• Schwalbe's line

• Trabecular meshwork

• Scleral spur

• Ciliary body band

• Root of iris

• The localization of foreign bodies, abnormal blood vessels or tumors in the angle.

• Demonstration of the extent of peripheral anterior synechiae.

• During goniotomy, a surgical procedure for the treatment of congenital glaucoma.

The below instrument is an indirect gonioscope of four-mirror type, which is used to visualize the angle of the anterior chamber.

Solution to Question 4:

The image shows a nodule surrounded by redness. On administration of phenylephrine, blanching of superficial episcleral vessels is seen. Hence, the probable diagnosis is nodular episcleritis.

2.5% phenylephrine can be used to differentiate episcleritis and scleritis. Since the vessels of the episclera are superficial, they blanch. Whereas, in scleritis, the vessels do not blanch.

Episcleritis is benign, recurrent, inflammation of the episclera, involving the overlying Tenon's capsule but not the underlying sclera. It is more common in women than in men. It is an allergic reaction to an endogenous toxin. It is commonly associated with rheumatoid arthritis. Episcleritis is characterized by redness, mild ocular discomfort described as a gritty, burning, or foreign body sensation

Clinically, two types of presentations may occur:

• Simple episcleritis is characterized by sectorial or diffuse inflammation of the episclera.

• Nodular episcleritis is characterized by a pink or purple flat nodule surrounded by injection, usually situated 2-3 mm away from the limbus. The nodule is firm, and tender and can be moved separately from the sclera, and the overlying conjunctiva also moves freely.

Treatment includes topical NSAIDs, corticosteroid eye drops, cold compress, and artificial tears. In recurrent cases, systemic NSAIDs are used.

Scleritis refers to an inflammation of the sclera proper. It usually occurs in elderly patients (40-70 years) involving females more than males. It is frequently associated with an underlying systemic inflammatory condition. The patient presents with moderate to severe pain, redness, lacrimation, photophobia, and diminution of vision.

Scleritis can be classified as immune-mediated non-infectious scleritis and infectious scleritis. Immune-mediated non-infectious scleritis is further classified as:

I. Anterior sclerltis (98%)

a. Non-necrotizing scleritis (85%)

• Diffuse

• Nodular

b. Necrotizing scleritis(13%)

1. With inflammation:

• Vasa-occlusive

• Granulomatous

• Surgically induced

2. Without inflammation(scleromalacia perforans).

II. Posterior scleritis (2%)

Treatment options include:

• Non-necrotizing scleritis- Topical steroids and systemic indomethacin.

• Necrotizing scleritis- Topical and oral steroids, immunosuppressive agents and surgical treatment like scleral patch graft.

• Infectious scleritis- Antimicrobial therapy, and surgical debridement.

Other options-

Option C: Tenon's cyst or conjunctival inclusion cyst is a benign condition filled with serous fluid and slimy mucous secretions of congenital or acquired origin. Acquired cysts can occur following strabismus surgery, pars plana vitrectomy, scleral buckling, and Ahmed glaucoma valve insertion. The patient presents with swelling near the limbus with no pain or redness. The consistency is cystic to firm, non-tender with no mobility suggesting that the cyst is adherent to the underlying scleral tissue.

Option D: Phlycten is a white-pinkish nodule surrounded by hyperemia on the bulbar conjunctiva near the limbus. Phlyctenular conjunctivitis is a type 4 hypersensitivity reaction due to endogenous allergens like Staphylococcus aureus (most common) and Mycobacterium tuberculosis. Treatment includes topical steroids; atropine if the cornea is involved.

Solution to Question 5:

The marked structure represents the Canal of Petit.

In front of the anterior hyaloid membrane and behind the ciliary body and zonule is the capsulohyaloid sinus (or space), or Petit's canal. This is one of the sites of retrolental hemorrhages. This is also the route along which blood sometimes enters the lens, through some rupture in the posterior lens capsule.

Other options-

Option A: Posterior capsule of lens has circular adhesions with anterior hyaloid surface called as the hyaloideocapsular ligaments of Weiger or retrolental ligament. The circular attachment is

known as Egger's line. The potential space between the anterior hyaloid face and the posterior lense capsule with the circular attachment is referred to as the Berger's space.

Option B: Lens capsule is a thin, transparent, hyaline membrane surrounding the lens which is thicker over the anterior than the posterior surface

Option C: The canal of Hannover lies between orbiculo-anterocapsular and orbiculo-posterocapsular parts of zonular fibers.



Solution to Question 6:

Early signs of glaucoma in perimetry include generalized constriction of isopters and baring of blind spot.

Progression of visual field changes seen in glaucoma:

Perimetry is the procedure of estimating the extent of the visual fields. There are 2 types of perimetry - kinetic and static. In kinetic perimetry, a target of given luminance is moved from a non-seeing area in the periphery towards the center where it is perceived. In the static method, the location of a stimulus remains fixed, with the intensity increased until it is seen by the subject (threshold is reached) or decreased until it is no longer detected.

Charting of the visual fields is useful in the diagnosis of many disease conditions such as glaucoma, neurological disorders, and retinal diseases (e.g. retinitis pigmentosa) as well as following laser treatment of diabetic retinopathy.

Other options-

Option A: Static perimetry (not kinetic perimetry) is used in the quantification and tracking of visual changes in established glaucoma.

Option B: In Kinetic perimetry (not static perimetry), stimulus intensity is held constant and moved centrally with detection.

Option C: Optic disc changes are much greater than visual field changes. Therefore, visual field defects need not correspond to optic nerve head abnormalities to be classified as glaucoma.

Early non-specific changes	Isopter contractionBaring of blind spot:Exclusion of blind spot from the central field
Early significant changes	Peripheral nasal step(may be early or late)Paracentral scotomain Bjerrum's area.Siedel's scotoma:Paracentral scotoma joins the blind spot
Late significant changes	ArcuateorBjerrum scotomaRingorDoublearcuate scotoma: When 2 arcuate scotomas join togetherRoenne's central nasal step
Advanced glaucomatous field defects	Tubular visionNo light perception: Complete visual field defect.

Solution to Question 7:

Disc neovascularization is not a feature of nonproliferative diabetic retinopathy (NPDR), and is seen in proliferative diabetic retinopathy.

The important findings in severe NPDR are one or more of severe retinal hemorrhages in 4 quadrants, significant venous beading in 2 or more quadrants, and moderate intraretinal microvascular anomalies (microaneurysms) in 1 or more quadrants. This is known as the 4-2-1 rule.

Diabetic retinopathy is used to describe retinal changes seen in diabetic patients. It is classified into the following types:

• Non-proliferative diabetic retinopathy (NPDR).

• Proliferative diabetic retinopathy (PDR).

• Diabetic maculopathy.

• Advanced diabetic eye disease.

Management of diabetic retinopathy includes intravitreal anti-VEGF drugs, intravitreal steroids, laser therapy, and or surgical management.

Surgical management is done in the following cases:

• Tractional diabetic macular edema with NPDR: pars plana vitrectomy (PPV) with the removal of posterior hyaloid.

• Advanced PDR with dense vitreous hemorrhage/ fibrovascular epiretinal membrane: PPV along with the removal of opaque vitreous gel/ membrane and endophotocoagulation.

• Advanced PDR with tractional retinal detachment: PPV with endophotocoagulation and reattachment of the detached retina by scleral buckling.

Early Treatment of Diabetic Retinopathy Study classification:

Nonproliferative diabetic retinopathy (NPDR) -

• Very mild NPDR :

- Microaneurysms only.

- Management - Review in 12 months.

• Mild NPDR :

- Any or all of : microaneurysms, retinal hemorrhages, exudates, cotton wool spots, up to the level of moderate NPDR. No intraretinal microvascular anomalies (IRMA) or significant beading.

- Management - Medical control of diabetes with follow-up in 9-12 months.

• Moderate NPDR :

- Severe retinal hemorrhages (about 20 medium-large per quadrant) in 1–3 quadrants or mild IRMA.

- Significant venous beading can be present in no more than 1 quadrant.

- Cotton wool spots commonly present.

- Management - Review in approximately 6 months.

• Severe NPDR : The 4–2–1 rule; one or more of:

• Severe hemorrhages in all 4 quadrants

• Significant venous beading in 2 or more quadrants

• Moderate IRMA in 1 or more quadrants

• Management - Review in 4 months.

• Very severe NPDR : The 4–2–1 rule, two or more of the criteria for severe NPDR

• Review in 2–3 months.

Proliferative diabetic retinopathy (PDR)-

• Mild to moderate PDR :

• New vessels on the disc (NVD) or new vessels elsewhere (NVE), but extent insufficient to meet the high-risk criteria.

• Management- Prompt Pan-retinal photocoagulation is done if type 2 DM in a patient >40 years of age, otherwise defer PRP.

• High-risk PDR :

• New vessels on the disc (NVD) (about 1/3 disc area).

• Any NVD with vitreous hemorrhage.

• NVE greater than 1/2 disc area with vitreous hemorrhage.

• Management - Prompt pan-retinal photocoagulation.



Solution to Question 8:

The image depicts cobalt blue filter, fluorescein dye and corneal ulcer.

Corneal ulcers can be detected with the help of fluorescein sodium dye and a cobalt blue filter. Fluorescein staining of the cornea is carried out with a disposable autoclaved filter paper strip impregnated with the dye. The denuded epithelial area seen in corneal abrasions or ulcers appears brilliant green when stained with fluorescein. The epithelial defect of the cornea appears opaque green with a cobalt blue filter.

Corneal epithelial lesions are visualised using vital dye staining techniques. Dyes used are:

1. Fluorescein - to delineate areas with epithelial damage. Fluorescein Staining is shown below.



2. Rose Bengal - Stain diseased and devitalised cells. Rose bengal staining is shown below.



3. Alcian Blue - Stains excess mucus selectively.

Other options :

Option C : Lissamine green dye is used to stain devitalized cells of the cornea and conjunctiva. It is used to diagnose ocular surface disorders, like dry eye.

Option D : Interstitial keratitis is a form of non ulcerative keratitis. It is a non suppurative inflammation of the corneal stroma without primary involvement of the epithelium or endothelium. It is characterized by cellular infiltration of the corneal stroma. Some causes of interstitial keratitis include herpes simplex virus, syphilis, cogan syndrome and rheumatoid arthritis.

Slit-lamp examination of the cornea and conjunctiva after rose bengal staining reveals punctuate corneal ulcerations and attached filaments of corneal epithelium in keratoconjunctivitis sicca. The image attached below sums up the various patterns of staining and their diagnostic utility.



Solution to Question 9:

RK (Radial Keratotomy) would not be recommended to this boxer who wants vision correction surgery.

Radial keratotomy (RK) involves making deep (90% of corneal thickness) radial incisions in the peripheral part of cornea leaving the central 4 mm optical zone. These incisions on healing flatten the central cornea reducing its refractive power. This procedure was done to correct low to moderate myopia (2 to 6 D). However, because of its disadvantages and advent of safe techniques (LASIK and PRK), RK is now obsolete.

Disadvantages of RK:

- Cornea is weakened, increasing chances of globe rupture following trauma (RK more than PRK). Hence this procedure should not be done for patients who are at high risk of blunt trauma, such as sports persons and military personnel.

- Irregular astigmatism from uneven healing.

- Glare at night.

Laser-assisted in situ keratomileusis (LASIK) (Option C) is the most commonly done surgery for myopia. An excimer laser beam reshapes the cornea by ablating the superficial stroma to a predetermined extent after lifting a flap of the cornea with a sharp microkeratome. It is safer than RK.

Patient selection criteria:

- In individuals aged 21 years and above.

- In individuals who have a stable refractive error for at least 1 year.

- In individuals who have had unsatisfactory results with non-surgical treatment.

- In individuals in whom corneal-thinning disorders such as keratoconus have been ruled out.

Complications:

- Tear instability (dry eye)

- Wrinkling, distortion, or dislocation of the flap

- Subepithelial haze with resultant glare, especially at night.

- Persistent epithelial defects.

- Epithelial ingrowth under the flap.

- Diffuse lamellar keratitis

- Corneal ectasia

- Keratoconus



Other options :

Option B : PRK (Photorefractive keratectomy) corrects myopia by using an excimer laser to photoablate a central optical zone of anterior corneal stroma. This causes flattening of the central cornea. Like RK, the PRK also gives very good correction for up to -6 D of myopia but is safer.

Option D : SMILE (Small incision lenticule extraction) is a technique in which a lenticule of corneal stroma is extracted using a femtosecond laser. This can correct myopia, with and without astigmatism up to 10 D.

Solution to Question 10:

The child with a power of -1 D spherical and -2 D cylindrical at 180 degrees has with the rule astigmatism, as the vertical meridian is greater and more curved than the horizontal meridian.

Step 1 : - 1 D spherical would mean a power of -1 on vertical and horizontal axes.

| (-1)

|

--------|------- (-1)

|

|

Step 2 : - 2 D cylindrical at 180 degrees, would mean a power of -2 D at 90 degrees (perpendicular to 180).

| (-1 - 2 = -3)

|

--------|------- (-1)

|

|

Hence, as the vertical meridian (-3D) is greater than horizontal meridian (-1D), this would be with the rule astigmatism.

Astigmatism refers to the condition where there is different refractory power in both the principal meridians of the eye. Astigmatism is classified into two broad varieties - regular and irregular astigmatism.

• Regular astigmatism: there are two principal meridians.

• With the rule - principal meridians are vertical and horizontal; both meridians at 90°; and the vertical meridian is more curved than horizontal.

• Against the rule (Option C) - principal meridians are vertical and horizontal; both meridians at 90°; and the horizontal meridian is more curved than vertical.

• Oblique - the principal meridians are not vertical and horizontal, but they are at 90° to each other; e.g., at 45° and 135°.

• Bi-oblique - the two principal meridians are not vertical and horizontal, and they are not at 90° each other; e.g., at 40° and 100°.

• Irregular astigmatism: there are multiple meridians.

Clinical presentation of astigmatism include asthenopia (tiredness of eyes relieved by closing the eyes), blurred vision and defective vision, elongation of objects, keeping reading material closer, and half closure of the lid.

Tilted or oval optic disc is seen on ophthalmoscopy. Retinoscopy shows different power on the two meridians. Other investigations include keratometry, astigmatic fan test, and Jackson's cross-cylinder test.

Clinical types of astigmatism



Prescription in astigmatism: There are 3 components.

1) Power of the spherical component: Convex (+) or concave (-) dioptre sphere (DS)

2) Power of the cylinder: Convex (+) or concave (-) dioptre cylinder (DC)

3) Axis (orientation) of the cylinder

Other options :

Option A : In simple astigmatism, one of the meridia focuses on the retina. The other meridian focuses either in front of the retina (simple myopic) or behind the retina (simple hyperopic). Example, -1.00 DC × 180 degree.

Option D : Mixed astigmatism: Light rays in one meridian are focused in front of the retina and those in the other meridian are focused behind the retina. Example, -1DS with +2DC at 180 degree.



Solution to Question 11:

The image shows Hertel's exophthalmometer used to measure proptosis.

An abnormal protrusion of the globe because of a space-occupying lesion in the orbit is called proptosis. Exophthalmometry measures protrusion of the apex of the cornea from the outer orbital margin (with the eyes looking straight ahead). An absolute protrusion of >20 mm or a difference of >2 mm between the two eyes is labelled as proptosis.

Instruments used to measure proptosis:

• Hertel's exophthalmometer: the most common instrument used. Its advantage is that it measures both eyes simultaneously.

• Luedde exophthalmometer : The simplest instrument to measure proptosis



• Naugle Exophthalmometer : This is a superior and inferior orbital rim-based instrument.

Pseudoproptosis occurs when the eye has a proptotic anatomic appearance that is not caused by a mass effect from a lesion displacing the globe. Pseudoproptosis is seen in high myopia (due to increase in size of the eyeball), ipsilateral lid retraction (resultant sclera creates the impression of the globe protruding forward) and contralateral enophthalmos (the normal orbit, which is situated more anteriorly, may simulate proptosis when there is none).

Ophthalmology INI-CET 2024

Question 1:

A 3 months old baby was brought with photophobia and lacrimation. Which of the following cannot be the cause?

a) Congenital entropion

b) Congenital glaucoma

c) Congenital distichiasis

d) Congenital nasolacrimal duct obstruction

Question 2:

A construction site worker comes in with red eyes and lacrimation. He got chuna particles in his eyes an hour ago. What is the initial step in the treatment?

a) Irrigation with normal saline

b) Topical antibiotics and cycloplegics

c) Patch the eye after topical lubricants

d) Temporary tarsoraphy

Question 3:

What is anisometropia ?

a) Difference in visual acuity

b) Difference in refractive power

c) Difference in axial length

d) Difference in size of eye ball

Question 4:

The following are features of third cranial nerve palsy except

a) Elevation

b) Adduction

c) Limited abduction

d) Ptosis

Question 5:

What is the most common cause of proptosis in middle aged individuals?

a) Cavernous hemangioma

b) Orbital varix

c) Thyroid eye disease

d) Retinoblastoma

Question 6:

A 40 year old with high myopia wearing thick glasses from childhood came with complaints of sudden onset of flashes of light in left eye. What could be the likely cause ?

a) Retinal detachment

b) CRAO

c) CRVO

d) Subarachnoid haemorrhage

Question 7:

Which of the following disease results in weakness of the levator palpebrae superioris, corneal anaesthesia, lagophthalmos, and ultimately blindness?

a) Lyme's disease

b) Leprosy

c) Ophthamia neonatorum

d) Trachoma

Question 8:

All of the following are actions of superior oblique except?

a) Depression

b) Adduction

c) Abduction

d) Intorsion

Question 9:

A patient presents with diminished night vision and the following finding was seen on fundoscopy. Which of the following statements is not true regarding the condition?

a) Dystrophy of rods more common than that of cones

b) ERG shows decreased amplitude of A and B waves

c) Sometimes patients present in later stages with loss of peripheral vision.

d) Night blindness is a late feature

Question 10:

A patient of rheumatic heart disease presents with superior visual field defect in left eye since 1 week, what is the probable cause?

a) Central retinal artery occlusion

b) Central retinal vein occlusion

c) Branch retinal vein occlusion

d) Branch retinal artery occlusion

Question 11:

An 80 yrs old patient presents with a painful, red right eye. On examination, the IOP of his right eye is 50mmHg and left eye is 16mmHg. Slit lamp examination reveals deep anterior chamber, presence of flare cells in the anterior chamber and cataractous lens in the right eye. What is the most likely diagnosis?

a) Phacolytic glaucoma

b) Malignant glaucoma

c) Fuch heterochromic iridocyclitis

d) Phacomorphoic glaucoma

Question 12:

Which of the following is a complication of diabetic retinopathy?

a) Posterior staphyloma

b) Retinoschisis

c) Tractional retinal detachment

d) Progressive myopia

Question 13:

Which of the following is true about macular opacity?

a) Iris and pupil are visible

b) Only the pupillary margin is visible

c) Iris details are visible

d) Iris and pupillary margins are not visible

Question 14:

Which of the following is true about viscoelastic substances?

a) Causes mechanical injury to corneal endothelium

b) Prevents the Anterior Chamber from collapsing during crucial steps like capsulorhexis and IOL placement

c) It is inert and can be left behind after surgery

d) Silicone oil, perfluoropropane, and SF6 are the most commonly used substances

Question 15:

In pterygium, there is

a) Eye pain

b) Macular edema

c) Purulent discharge

d) Astigmatism

Question 16:

A 6-month-old baby, born preterm at 33 weeks of gestation, is brought by parents with complaints of white pupillary reflex since 2 weeks. CT revealed calcification of the globe in both eyes. What is the probable diagnosis?

a) ROP

b) Retinoblastoma

c) Congenital Glaucoma

d) Congenital Cataract

Question 17:

Which of the following is a cause of bilateral sudden painless loss of vision?

a) Acute congestive glaucoma

b) Diabetic retinopathy

c) Keratoconus

d) Iridocyclitis

Question 18:

In FFA, the microaneurysms appear as:

a) Hypofluorescent dots

b) Hyperfluorescent dots

c) Hypofluorescent lines

d) Hyperfluorescent branching-like structures in the vessel

Question 19:

Normal diameter of artery to vein at the optic disc margin

a) 1:3

b) 2:3

c) 2:5

d) 1:5

Question 20:

In the case of Best disease, which of the following is the diagnostic investigation?

a) EOG

b) ERG

c) Multifocal ERG

d) Visual evoked potential

Question 21:

A factory worker presented with a chemical injury in both his eyes a few hours back. What is the first immediate management?

a) Copious irrigation of eyes

b) Take history

c) Check vision in both eyes

d) Check IOP and plan for surgery

Question 22:

Which of the following layers of the retina are correctly labeled?



a) 1- Retinal pigment epithelium; 2- Outer nuclear layer; 3- Inner plexiform layer; 4-Ganglion layer

b) 1- Retinal pigment epithelium; 2-Inner nuclear layer; 3- Outer plexiform layer; 4- Ganglion layer

c) 1- Ganglion layer; 2- Inner nuclear layer; 3- Outer plexiform layer; 4- Retinal pigment epithelium

d) 1- Ganglion layer; 2- Outer plexiform layer; 3- Inner nuclear layer; 4- Retinal pigment epithelium

Answer Key

Question No.	Correct Option
1	d
2	a
3	b
4	c
5	c
6	a
7	b
8	b
9	d
10	d
11	a

12	c
13	b
14	b
15	d
16	b
17	b
18	b
19	b
20	a
21	a
22	a

Detailed Explanations

Solution to Question 1:

Congenital nasolacrimal duct obstruction (NLDO) is least likely to cause watering and photophobia.

Nasolacrimal duct obstruction is the most common disease of the lacrimal system. It usually presents in the first few weeks or months of life as excessive tearing and ocular discharge. It mostly occurs due to incomplete canalization of the duct.

Congenital NLDO mainly results in persistent tearing and sometimes a discharge from the eyes but not photophobia, as the light sensitivity is not directly related to this condition.

Typically, diagnosis is established through a combination of patient history and physical examination. In cases where symptoms are intermittent, the presence of an obstruction can be confirmed using the fluorescein dye disappearance test.

The initial treatment involves performing nasolacrimal massage, typically 2-3 times daily, along with cleansing the eyelids using warm water and applying topical antibiotics. If it does not resolve after massaging by the age of 6 to 10 months, lacrimal duct probing is done.

If probing fails, other treatment options include intubation with silastic tubing with or without balloon dilatation of the nasolacrimal duct and dacryocystorhinostomy.

Other options:

Option A: Congenital entropion: This is a condition where the eyelid is turned inward, causing the eyelashes to rub against the cornea. This can lead to irritation, photophobia, and excessive tearing (lacrimation).

Option B: Primary congenital glaucoma (PCG) is caused by impaired aqueous outflow due to a maldeveloped anterior chamber angle. Clinically, the child presents with lacrimation, photophobia, and blepharospasm.

Option C: Congenital distichiasis: This is a condition where an extra row of eyelashes grows from the meibomian glands on the eyelid margin. These extra lashes can irritate the cornea, leading to photophobia and tearing.

Solution to Question 2:

The initial step in the management of a patient presenting with chemical burns is immediate and thorough irrigation with the available clean water or normal saline delivered through an IV tubing. This is done to prevent further damage.

Along with irrigation, contaminant elimination through mechanical means shall be done. In the event of residue, especially with lime (chuna), it is essential to meticulously eliminate them using a swab stick.

After the initial steps, maintenance of favorable conditions for rapid and uncomplicated healing by following measures:

• Topical antibiotic drops, for example, moxifloxacin 4-6 times a day to prevent infection.

• Steroid eye drops to decrease inflammation, neutrophil infiltration, and treat anterior uveitis. Caution is advised after 2 weeks, as steroids could disrupt collagen synthesis.

• Cycloplegics, such as atropine, might enhance comfort.

Grading of severity (Roper–Hall system) of chemical injuries is based on corneal clarity and severity of limbal ischemia:

• Grade 1 is characterized by a clear cornea (epithelial damage only) and no limbal ischemia (excellent prognosis).

• Grade 2 shows a hazy cornea but with visible iris detail and less than one-third of the limbus being ischemic (good prognosis).

• Grade 3 manifests as a total loss of the corneal epithelium, stromal haze obscuring iris detail, and between one-third and half of the limbus showing ischemia (guarded prognosis).

• Grade 4 manifests with an opaque cornea and more than 50% of the limbus showing ischemia (poor prognosis).

Solution to Question 3:

Anisometropia is the condition where there is difference in refractive powers of two eyes.

Anisometropia occurs when the total refraction of both eyes are unequal. A difference of 1D in the two eyes causes a 2% difference in the size of retinal images formed. A difference up to 2.5D is well-tolerated, but beyond 4D cannot be tolerated. Anisometropia can occur as a combination of emmetropia, myopia, hypermetropia, and astigmatism.

It is diagnosed after retinoscopic examination and autorefractometry.

Treatment includes the usage of spectacles (it can be tolerated up to a maximum difference of 4D), contact lenses, intraocular lenses (for uniocular aphakia), refractive corneal surgery, or refractive

lens exchange methods (used in high degree anisometropia).

Solution to Question 4:

Limited abduction is not a feature of third cranial nerve palsy.

Abduction is carried out by the lateral rectus, which is supplied by the abducens (sixth) nerve.

Third cranial nerve (oculomotor nerve) palsy presents with:

• Profound ptosis due to the weakness of the levator muscle

• Limited adduction due to medial rectus weakness

• Limited elevation due to the weakness of the superior rectus and inferior oblique

• Abduction and depression in the primary position ("down and out")

Solution to Question 5:

The most common cause of proptosis in middle age is thyroid eye disease.

Graves ophthalmopathy is one of the most serious ocular manifestations of Graves disease. Between the ages of 25 and 50, it's the leading cause of unilateral and bilateral proptosis in adults. Proptosis, eyelid retraction, restricted myopathy, and perhaps compressive optic neuropathy are clinical manifestations that could potentially be used for diagnosis.

The mechanism proposed for this is the acute precipitation of a hypothyroid state which leads to an increase in TSH secretion. This leads to stimulation of the retro-orbital fibroblasts and adipocytes which have the receptor for TSH.

An abnormal protrusion of the globe is called proptosis. Exophthalmometry measures protrusion of the apex of the cornea from the outer orbital margin (with the eyes looking straight ahead). An

absolute protrusion of >20 mm or a difference of >2 mm between the two eyes is labelled as proptosis.

Option A: Cavernous hemangioma is the most common primary benign orbital tumor in adults. They are composed of large, dilated vascular channels; are more infiltrative; frequently involve deep structures; and do not spontaneously regress. On histologic examination, the mass is sharply delineated but unencapsulated with large, cavernous blood-filled vascular spaces, separated by connective tissue stroma.

Option B: The type of proptosis seen in orbital varix is non-pulsatile, intermittent, and reducible proptosis. On activities such as forward bending or lowering of the head, turning the head forcefully, hyperextension of the neck, and coughing, the varices fill up, causing proptosis, and then, they regress on shifting to a non-dependent position.

Option D: Retinoblastoma is the most common intraocular malignancy in children. It is associated with mutations in the RB1 gene at chromosome 13q14. White pupillary reflex or leukocoria is the most common presenting feature, followed by a squint which is usually convergent. Histopathology of the tumor shows small round cells with large nuclei. Microscopic features includes Flexner-Wintersteiner rosettes(highly specific of retinoblastoma), Homer-Wright rosettes, pseudorosettes, and fleurettes formation. Areas of necrosis and calcification can also be seen.

Solution to Question 6:

The given scenario of patient with severe myopia presenting sudden onset of flashes of light is likely retinal detachment.

Retinal detachment is the separation of the neurosensory retina and the retinal pigment epithelium (RPE).

Risk factors include myopia, previous intraocular surgery such as aphakia or pseudophakia, a family history of retinal detachment, and trauma.

Clinical features include sudden painless loss of vision, flashes of light or photopsia, floaters in the eye, and curtains in the field of vision.

Examination findings will reveal:

• Normal or slightly low intraocular pressure

• Marcus Gunn pupil (relative afferent pupillary defect)

• Altered red reflex on the distant direct ophthalmoscope

• Indirect ophthalmoscopy may show grey reflex of the freshly detached retina or breaks in the retina

• Vitreous pigments on the anterior vitreous (Shaffer sign)

• Shifting fluid sign- Change in the position of the detached area with a change in gravity is the hallmark of exudative retinal detachment.

Management of rhegmatogenous retinal detachment includes sealing of retinal breaks, reducing the vitreous traction on the retina, and flattening of the retina by draining the subretinal fluid.

Scleral buckling, pneumatic retinopexy, and pars plana vitrectomy can help in maintaining the chorioretinal apposition.

Treating the cause helps in exudative retinal detachment. Enucleation can be done if intranuclear tumors are present.

Solution to Question 7:

Leprosy could result in paresis of the levator palpebrae superioris, corneal anaesthesia, lagophthalmos, and ultimately blindness.

Lagophthalmos is the inability to close the eyelids. This can be due to the involvement of the zygomatic branch of the facial nerve (as in leprosy). Other causes of lagophthalmos include extreme proptosis and symblepharon. The decreased corneal sensation seen is due to the involvement of the ophthalmic branch of the trigeminal nerve which innervates the cornea.

Leprosy or Hansen disease is a chronic granulomatous disease caused by infection with Mycobacterium leprae. Ocular features in leprosy occur due to bacterial invasion and include corneal anesthesia, lagophthalmos, corneal ulceration, episcleritis, and iris atrophy. Other features seen are:

• Keratitis with thickened, beaded corneal nerves, punctate subepithelial lesions, pannus, and vascularization

• Chronic low-grade anterior uveitis

• Miosis

• Iris atrophy

• Iris pearls

• Retinal pearls

Option B: The most common ocular manifestation in Lyme's disease is conjunctivitis. Other manifestations like keratitis, iridocyclitis, retinal vasculitis, choroiditis, optic neuropathy, and uveitis occurs very rarely.

Option C: Ophthalmia neonatarum also called neonatal conjunctivitis is inflammation of the conjunctiva occurring in an infant <30 days after birth. The clinical features include conjunctival redness, eyelid edema, and ocular discharge (purulent or mucopurulent).

Option D: In trachoma, the symptoms include lacrimation, scanty mucoid discharge and foreign body sensation. Conjunctival congestion and follicles, papillary hyperplasia, Herbert follicles (seen in the limbal area), and progressive pannus are the characteristic features.

Solution to Question 8:

Adduction is not an action of the superior oblique muscle.

The superior oblique muscle's primary, secondary and tertiary actions are intorsion, depression and abduction respectively.

It originates from the apex of the orbit, turns around the trochlea, and inserts into the sclera behind the equator. The oblique direction of the tendon is responsible for its rotatory effect (intorsion), in addition to depression and abduction. It is supplied by the trochlear nerve.



Solution to Question 9:

Diminished night vision along with fundoscopy features point to the diagnosis of retinitis pigmentosa. Option D is not true regarding retinitis pigmentosa. Night blindness is an early feature.

Retinitis pigmentosa is retinal dystrophy that results in degeneration of photoreceptors(rods more than cones). It can be inherited (autosomal dominant, autosomal recessive, or X-linked recessive) or acquired due to sporadic mutation in the rhodopsin gene. Fundoscopy shows bilateral, mid-peripheral intraretinal perivascular 'bone-spicule' changes and atrophy of the retinal pigment epithelium

Patients present with worsening night blindness initially. However, patients do not often notice night blindness and may present after much disease progression. They also develop tunnel vision, which manifests as difficulty performing activities of daily living such as reading, walking, and driving.

Ocular examination reveals normal visual activity and central vision (though central vision will be involved later in disease progression). Fundoscopy shows bilateral, mid-peripheral intraretinal perivascular 'bone-spicule' changes and atrophy of the retinal pigment epithelium. There is severe peripheral pigmentation, arteriolar narrowing, and pallor of the disc.

Retinitis pigmentosa can lead to posterior subcapsular cataracts, open-angle glaucoma, keratoconus, and posterior vitreous detachment as possible complications.

There is no specific treatment for retinitis pigmentosa

Option B:Electroretinography (ERG) is the record of changes in the resting potential of the eye induced by a flash of light.

Waves of normal ERG

• a-wave: negative wave possibly arising from the rods and cones

• b-wave: large positive wave which is generated by Muller cells, but represents the activity of the bipolar cells

• c-wave: positive wave representing the metabolic activity of pigment epithelium

In retintis pigmentosa, it is abnormal and shows reduced scotopic, subnormal responses in early disease and becomes completely extinguished in advanced disease.

Solution to Question 10:

A patient with rheumatic heart disease (RHD) presenting with superior visual defects for the past one week is likely due to branch retinal artery occlusion (BRAO).

Retinal artery occlusion usually occurs due to an embolus. Emboli can originate between the vasculature from the heart to the ophthalmic artery. Risk factors include Buerger's disease, hypertension, elevated lipid levels, cigarette smoking, RHD, and diabetes mellitus.

Retinal emboli can be of 3 types: cholesterol (Hollenhorst plaque), platelet-fibrin, and calcific. The emboli associated with cardiac thrombosis as seen in RHD are platelet-fibrin.

Features of BRAO:

• Sudden painless loss of vision in one eye corresponding to the territory of the obstructed artery. Amaurosis fugax can be seen in patients with preexisting carotid disease.

• Retinal whitening is the most notable finding seen due to ischaemia of the inner layers of the retina. Cotton wool spots can also appear.

• Fluorescein angiography (FA) shows diminution in the dye distal to the site of obstruction. There may be some leaks visible at the site of the embolus.

• OCT (optical coherence tomography) initially shows hyperreflective and thickened inner layers of the retina in the obstructed territory, which start thinning out with time.

• Visual field testing is used to assess the extent of loss of vision.

No effective treatment has been found for BRAO. Ocular massage, lasers, or paracentesis can be used to disrupt the embolus. Vasodilation with inhaled amyl nitrite can be done.

In hypercoagulative states, prophylactic anticoagulation can be started to prevent future episodes.

Other options:

Option A: Central retinal artery occlusion occurs due to obstruction at the level of lamina cribrosa. Patient complains of sudden painless loss of vision occurring over seconds.

Option B: Patients with CRVO usually complain of the acute onset of painless blurred vision in one eye. CRVO is seldom asymptomatic.

Option C: Patients with BRVO typically present with peripheral visual field defects in the defective eye. However, in settings of RHD, arterial occlusion is more likely.

Solution to Question 11:

The given clinical scenario of an 80 year old presenting with painful red eyes, increased IOP and presence of a deep anterior chamber points towards the diagnosis of secondary open-angle glaucoma - phacolytic glaucoma.

Phacolytic glaucoma is secondary open-angle glaucoma occurring in association with hyper mature morgagnian cataracts.

Trabecular obstruction is caused by high molecular-weight lens proteins that leak through the intact capsule in cases of the hyper mature cataract into the aqueous humour; macrophages containing lens proteins may also contribute to this.

Presentation:

• Eye pain

• Diminished vision

• Photophobia

• Lacrimation

• Headache

• Increased intraocular pressure

• Deep anterior chamber with fine white protein particles which may settle as pseudohypopyon

• Open-angle seen on gonioscopy

• Lens: Milky white cortex with sunken brownish nucleus (Morgagnian cataract)

Treatment: Medical therapy to lower the intraocular pressure, Surgery to extract the lens

Other options:

Option B: Malignant glaucoma is seen after intraocular surgery such as trabeculectomy. The forward flow of aqueous is blocked as the tips of ciliary processes rotate forward and press against the lens/IOL/anterior hyaloid of the vitreous. Hence, aqueous humor collects in the vitreous cavity. There is persistent shallow or flat anterior chamber

Option C: In Fuchs heterochromic iridocyclitis, the IOP is not usually so high.

Option D: Phacomorphic glaucoma is an acute secondary angle-closure glaucoma caused by an intumescent cataractous lens due to rapid maturation of cataract or following traumatic capsular rupture. The swollen lens obliterates the angle by forcing the root of the iris against the cornea. The patient presents with features of acute glaucoma with a shallow anterior chamber and a swollen cataractous lens.

Solution to Question 12:

Tractional retinal detachment is a complication of diabetic retinopathy.

Diabetic retinopathy is a complication of diabetes that affects the retina due to prolonged high blood sugar levels. Over time, it can lead to several complications, including:

• Diabetic Macular Edema (DME): Diabetic retinopathy weakens the walls of the blood vessels in your retina, causing the leak of fluid and blood into the macula.

• Vitreous Hemorrhage: New blood vessels can bleed into the vitreous. This bleeding prevents light from reaching the retina. If the bleeding is minor, may present with floaters. However, in more severe cases, the blood can fill the vitreous cavity, completely obstructing your vision.

• Neovascular glaucoma: Abnormal blood vessels can grow out of the iris and can block fluid from draining properly, leading to a type of glaucoma.

• Tractional retinal detachment occurs due to the formation of fibrovascular tissue from proliferative diabetic retinopathy. The fibrous tissue exerts traction on the retina, pulling it away from the underlying layers and causing a retinal detachment. The most common cause of tractional retinal detachment is proliferative diabetic retinopathy.

Other options

Option A: Posterior staphyloma is associated with progressive myopia.

Option B: Retinoschisis refers to the splitting of the retinal layers and is seen in degenerative retinal diseases.

Option D: Progressive myopia is unrelated to diabetic retinopathy and occurs due to elongation of the eyeball, genetic factors, environmental factors and too much near work such as reading and using digital devices for long hours.

Solution to Question 13:

Among the given options, ‘only the pupillary margin is visible’ is the true statement regarding macular opacity. Macular opacity is a type of corneal opacity.

Corneal macular opacity refers to a moderate to dense opacity in the cornea. In such cases, light transmission through the cornea is partially obstructed. This limits the visibility of deeper structures like the iris and its fine details(options A & C). However, because the pupillary margin contrasts with its surrounding structures and lies directly behind the corneal opacity and the obstruction selectively allows the pupillary margin to be visible(options B & D).

Solution to Question 14:

Viscoelastic substances prevent the anterior chamber from collapsing during crucial steps like capsulorhexis and IOL (Intra-ocular lens) placement.

Viscoelastic substances are commonly used in ocular surgery, particularly in cataract surgery, to maintain the anterior chamber (AC), protect the corneal endothelium, and facilitate surgical procedures. They prevent the collapse of the AC during crucial steps such as capsulorrhexis and IOL (intraocular lens) placement by providing a cushioning effect.

Other options

Option A: Viscoelastic substances are typically used to protect the corneal endothelium, not to cause any mechanical injury.

Option C: While some viscoelastics are inert, not all can be safely left behind. Some types of viscoelastic substances (especially those that are not fully biodegradable) should be removed from the eye after surgery to prevent complications.

Option D: Sodium hyaluronate and hydroxypropyl methylcellulose are the most commonly used viscoelastic substances, compared to silicone oil, perfluoropropane, and SF6. Silicone oil, perfluoropropane, and SF6 are the substances used for vitreoretinal surgery (such as tamponade agents for retinal detachment), not for routine cataract surgery. Viscoelastic used in cataract surgery are typically substances like sodium hyaluronate and hydroxypropyl methylcellulose.

Solution to Question 15:

In pterygium, there is astigmatism. It causes astigmatism by flattening the cornea along the axis of the pterygium.

Pterygium is a subconjunctival fibrovascular tissue encroaching on the cornea. It is a degenerative condition of the sub-conjunctival tissue and is never inflammatory. It occurs due to exposure to UV-B radiation and the most common site is the nasal.

It is made up of three parts - a "cap" (an avascular halo-like zone at the advancing edge), a head, and a body.

Clinical features:

• Astigmatism with the rule - most common finding

• Stocker's line - deposition of iron on the advancing edge

• Fuchs islets - small, discrete, whitish flecks consisting of clusters of pterygial epithelial cells, often present at the advancing edge

The image below shows a pterygium.



Solution to Question 16:

The given clinical presentations of white pupillary reflex (leukocoria) and CT revealing calcification of the globe in both eyes is suggestive of retinoblastoma.

Retinoblastoma is a proliferation of neural cells that have failed to evolve normally. It arises from the neurosensory retina. It is a malignant tumour of the retina that typically presents in children, often before the age of 5. Leukocoria and squint are the most common presentation of retinoblastoma. It may also present as painful red eyes, cataracts, buphthalmos, proptosis(not microphthalmos) and pseudohypopyon.

In this case, the presence of calcification within the globe on a CT scan supports the diagnosis of retinoblastoma, as it is known to frequently show calcifications on imaging.



Other options

Option A: ROP (Retinopathy of Prematurity) is a concern in preterm infants (especially those born before 32 weeks), and it can also present with white pupillary reflex, it does not show calcification of the globe on a CT scan. ROP can lead to retinal detachment or fibrosis, but calcification is not a typical feature.

Option C: Congenital glaucoma may present with symptoms such as enlarged corneas, increased intraocular pressure, and a shallow anterior chamber. White pupillary reflex or calcification on CT are not typical presentations of congenital glaucoma.

Option D: Congenital cataracts can present with leukocoria (white pupillary reflex) but would not show calcification on a CT scan. They are more likely to be detected by a clinical examination and would typically not result in calcifications.

Solution to Question 17:

Diabetic retinopathy is a cause of bilateral sudden painless loss of vision due to vascular complications.

Bilateral sudden painless loss of vision is a hallmark of advanced diabetic retinopathy, particularly when associated with complications such as:

• Vitreous haemorrhage: Leakage of blood into the vitreous cavity due to rupture of fragile neovascular vessels.

• Tractional retinal detachment: Occurs when fibrovascular tissue pulls the retina away from its underlying structures.

• Macular oedema: Severe swelling in the macula can lead to central vision loss.

These conditions can cause sudden and significant visual impairment, often affecting both eyes in patients with long-standing, poorly controlled diabetes.

Other options

Option A: Acute congestive glaucoma: Typically causes sudden vision loss, but it is painful and unilateral in most cases.

Option C: Keratoconus is a progressive condition that leads to gradual vision loss due to corneal thinning and irregular astigmatism, not sudden or bilateral loss.

Option D: Iridocyclitis is associated with pain, redness, and photophobia, and it is usually unilateral.

Solution to Question 18:

In Fundus Fluorescein Angiography (FFA), microaneurysms (a hallmark of diabetic retinopathy) appear as hyperfluorescent dots due to the accumulation of fluorescein dye in the localized dilations of retinal capillaries.



Other options:

Option A: Hypofluorescent dots indicate blocked fluorescence, often due to hemorrhage or ischemia.

Option C: Hypofluorescent line might occur in cases of vascular occlusion.

Option D: Hyperfluorescent branching-like structure in the vessel suggests leakage from neovascularization or vascular abnormalities.

Solution to Question 19:

The normal diameter ratio of the central retinal artery to the central retinal vein at the optic disc margin is 2:3.

The ratio 2:3 means that the artery's diameter is approximately two-thirds of the vein's diameter. This is considered the physiological normal. The ratio is assessed during the fundoscopic examination, and significant deviations can indicate pathological conditions:

• Decreased ratio (arteriolar narrowing) may suggest hypertension or arteriosclerosis.

• Increased ratio (relative venous narrowing) is less common but may occur in rare conditions.

Solution to Question 20:

EOG (electrooculogram) is the diagnostic investigation in the case of Best disease. EOG measures the function of the retinal pigment epithelium (RPE), which is typically abnormal in Best disease, even in asymptomatic carriers. The Arden ratio is significantly reduced (<1.8) in affected individuals.

Best's disease/Vitelliform macular dystrophy is an inherited macular dystrophy in which lipofuscin accumulates in the central macula causing progressive central vision loss. It is autosomal-dominant and linked to the mutation in the bestrophin (VMD2) gene.

It is characterized by bilateral yellow, yolk-like (vitelliform) macular lesions.

The stages of evolution of the macular lesions are described as progressing from:

• Pre-vitelliform stage

• Vitelliform stage- scrambled egg stage with or without hypopyon

• Vitelliruptive- the lesion breaks and visual acuity drops

• The atrophic stage

Patients with Best's disease usually have good visual acuity when the "yolk" remains intact.

Diagnosis: This is the only disease with relatively normal ERG (electroretinogram) results associated with an abnormal EOG. Optical coherence tomography (OCT) reveals that the vitelliform material appears as a dome-shaped, hyperreflective, and homogenous lesion, located below the hyperreflective photoreceptor layer.

The image below shows Best's disease - the typical vitelliform lesion



Other options

Option B: ERG (electroretinogram) assesses retinal activity and is mainly used in diagnosing pigmentary retinal dystrophy, complete occlusion of the retinal artery, early glaucomatous damage, macular degeneration in cataract, complete old retinal detachment or advanced siderosis.

Option C: Multifocal ERG assesses retinal activity and mainly provides a detailed evaluation of the central retina as in Stargardt disease or chloroquine toxicity.

Option D: Visual evoked potential (VEP) is used to evaluate the visual pathway's function

Solution to Question 21:

Copious irrigation of the eyes is the first immediate management in this case.

The initial step in the management of a patient presenting with chemical burns is immediate and thorough irrigation with the available clean water or normal saline delivered through an IV tubing. This is done to prevent further damage.

Along with irrigation, contaminant elimination through mechanical means shall be done. In the event of residue, especially with lime (chuna), it is essential to meticulously eliminate them using a swab stick.

After the initial steps, maintenance of favourable conditions for rapid and uncomplicated healing by following measures:

• Topical antibiotic drops, for example, moxifloxacin 4-6 times a day to prevent infection.

• Steroid eye drops to decrease inflammation, and neutrophil infiltration, and treat anterior uveitis. Caution is advised after 2 weeks, as steroids could disrupt collagen synthesis.

• Cycloplegics, such as atropine, might enhance comfort.

Grading of severity (Roper–Hall system) of chemical injuries is based on corneal clarity and severity of limbal ischemia:

• Grade 1 is characterized by a clear cornea (epithelial damage only) and no limbal ischemia (excellent prognosis).

• Grade 2 shows a hazy cornea but with visible iris detail and less than one-third of the limbus being ischemic (good prognosis).

• Grade 3 manifests as a total loss of the corneal epithelium, stromal haze obscuring iris detail, and between one-third and half of the limbus showing ischemia (guarded prognosis).

• Grade 4 manifests with an opaque cornea and more than 50% of the limbus showing ischemia (poor prognosis).

Solution to Question 22:

The layers of the retina marked above are:

• 1- Retinal pigment epithelium

• 2- Outer nuclear layer

• 3- Inner plexiform layer

• 4-Ganglion layer

Ophthalmology NEET 2024

Question 1:

Which of the following muscle is responsible for ptosis in horner's syndrome?

a) Levator palpebrae superioris muscle

b) Oribicularis oculi

c) Horner's muscle

d) Muller's muscle

Question 2:

An 11-year-old presented with the following growth in the eye as shown below. What is the diagnosis?

a) Papilloma

b) Myxoma

c) Pyogenic granuloma

d) Limbal dermoid

Question 3:

Patient undergoes lens implantation folllowing which there's worsening of vision after 1 year. Slit lamp examination finding is shown in the image below. Identify the complication.



a) Posterior subcapsular cataract

b) Bullous keratopathy

c) Subluxation of lens

d) Corneal opacity

Question 4:

Identify the perimetry finding.



a) Arcuate scotoma

b) Blind spot enlargement

c) Altitudinal defect

d) Roenne nasal step

Question 5:

A patient undergoes worth's four-dot test . The results are as shown in the image below. Identify the condition.



a) Crossed diplopia

b) Left suppression

c) Right suppression

d) Uncrossed diplopia

Question 6:

A 20-year-old presented with gradual loss of vision at night and a fundoscopy image is shown. What is the diagnosis?

a) Retinitis pigmentosa

b) Retinitis punctata albicuns

c) Retinal hemorrhage

d) Fundus flavimaculatus

Question 7:

A 2-month-old girl child was brought to the hospital by her parents with the following presentation. What is the appropriate management?



a) Immediate surgery

b) Wait until 4 months of age

c) Wait until 2 years of age

d) Trial of medical therapy

Question 8:

The following MRI image shows a malignancy. What is the visual field defect observed in this case?



a) Complete blindness

b) Homonymous hemianopia

c) Binasal hemianopia

d) Bitemporal hemianopia

Question 9:

A 26-year-old teacher presents with a red nodule over her sclera, which resolves with medications but is recurrent. Identify the condition and what else will you look for in this patient?



a) Pingecula; Dry eye

b) Episcleritis; Rheumatoid arthritis

c) Pterygium; Astigmatism

d) Scleritis; Rheumatoid arthritis

Question 10:

Identify the visual field defect in the given image:



a) Altitudinal hemianopia

b) Complete binasal hemianopia

c) Bitemporal hemianopia

d) Homonymous hemianopia

Question 11:

A patient complained of acute painful diminution of vision in the left eye for the past 2 days. On ophthalmic examination, circumcorneal congestion and irregularly shaped pupil were noted as shown below, with the left eye visual acuity at 6/60 and the right eye at 6/12. What is the probable diagnosis?



a) Angle closure glaucoma

b) Acute anterior uveitis

c) Acute conjunctivitis

d) Episcleritis

Question 12:

A patient who was watching a movie suddenly developed acute pain in the eye and diminution in vision. He was brought to the emergency department and on examination, a shallow anterior chamber was noted. What would be the appropriate management for this condition?



a) IV Mannitol + atropine e/d

b) IV Mannitol + pilocarpine e/d

c) IV Mannitol+ methyl cellulose e/d

d) IV Mannitol+ moxifloxacin e/d

Question 13:

Identify the defect shown in the image:

a) Herbert pits

b) Horner tranta spots

c) Gelatinous papillae

d) Pannus

Question 14:

A patient wearing contact lenses develops pain in the orbital region. He has had a very casual attitude towards cleaning his contact lenses. He presented with corneal ulcer and keratitis. Scrapings showed an organism with a spine-like pseudopodia. Which organism is the likely cause of this patient's symptoms?

a) Balamuthia

b) Acanthamoeba

c) Naegleria fowleri

d) Staphylococcus aureus

Answer Key

Question No.	Correct Option
1	d
2	d
3	a
4	b
5	c
6	a
7	a
8	d
9	b
10	d
11	b
12	b
13	a
14	b

Detailed Explanations

Solution to Question 1:

Ptosis in Horner's syndrome is due to the paralysis of Muller's muscle. Muller's muscle(superior tarsal muscle) maintains the elevation of the upper eyelid. It is innervated by the sympathetic nervous system.

Fasanella-Servat operation is indicated in cases having mild ptosis(< 2 mm) and good levator function. In this surgery, the upper tarsal border along with its attached Muller's muscle and conjunctiva are resected.

Solution to Question 2:

The image shows the presence of a limbal dermoid.

Limbal Dermoid is a choristoma (a mass of histologically normal tissue in an abnormal location). It consists of collagenous tissue containing dermal elements and covered by stratified squamous epithelium. Limbal dermoid present in early childhood as a subconjunctival mass in the infratemporal limbus. Treatment is indicated for chronic irritation, amblyopia, vision impairment and corneal dellen formation. Small dermoids can be excised, larger lesions may need lamellar keratosclerectomy.

Solution to Question 3:

The given clinical vignette supported with the image of vacoulated clusters seen behind IOL is suggestive of Elschnig's pearls which is a type of posterior subcapsular cataract (after cataract).

Posterior subcapsular cataracts occur mainly due to ageing, other causes include the use of topical steroids, trauma, excessive consumption of alcohol, and exposure to ionizing radiation. Visual impairment is more during the day or exposure to bright light as the pupil constricts which blocks light rays.

After cataract also known as secondary cataract or posterior capsular opacification is a late postoperative complication that follows extracapsular extraction of the lens.

Different types of after cataract include:

• Elschnig's pearls: where the subcapsular epithelial cells proliferate and develop into vacuolated soap bubble like clusters along the posterior capsule.

• Soemmering's ring: A peripheral dense ring is formed behind the iris and is enclosed between layers of the capsule.

• Dense membranous after cataract: formed in the pupillary area impairing the vision.

• Thin membranous after cataract : is thickened posterior capsule.

Treatment of after cataract is by

• Nd-YAG laser capsulotomy of the posterior capsule which works on the principle of photodisruption

• surgical posterior capsulotomy.

Solution to Question 4:

The perimetry image in the question above shows blind spot enlargement.

Blind spot enlargement is a rare finding in the visual field examination. The most common cause of blind spot enlargement is Idiopathic intracranial hypertension. Blind spot enlargement occurs due to displacement of the peripapillary retina and refractive changes due to significant optic nerve head swelling. Other common causes of blind spot enlargement are Syphilis, West Nile virus, or Rubella vaccination

Progression of visual field changes seen in glaucoma :

Early non-specific changes	Isopter contractionBaring of a blind spot:Exclusion of blin d spot from the central field
Early significant changes	Peripheral nasal step(may be early or late)Paracentral scot omain Bjerrum's area.Siedel' s scotoma:Paracentral scoto ma joins the blind spot
Late significant changes	ArcuateorBjerrum scotomaRi ngorDoublearcuate scotoma: When 2 arcuate scotomas joi n togetherRoenne's central n asal step
Advanced glaucomatous field defects	Tubular visionNo light perce ption: Complete visual field d efect.

Solution to Question 5:

The given image with 3 green dots likely suggests that the patient is suffering from right suppression.

Worth's four-dot test

• Sensory function test

• Patient wears goggles with

• Red lens: Right eye

• Green lens: Left eye

• A box is viewed having 4 lights: one red, two green, and one white

• Inference:

• A: Normal—All 4 lights seen

• B: Abnormal retinal correspondence—All 4 lights seen in the presence of manifest squint

• C: Left Suppression—2 red lights seen

• D: Right suppression—3 green lights seen

• E: Diplopia—5 lights seen (2 red, 3 green)



Solution to Question 6:

The fundus shows a pale waxy disc, attenuated arteries, and bony spicules along with gradual loss of vision which implies that it is case of retinitis pigmentosa (RP).

Retinitis pigmentosa (RP) is a dystrophy of rods and cones, with rods being primarily affected. It is a genetic disease. The non-hereditary/sporadic type is most common. Among the hereditary type, it can occur in an autosomal dominant (AD), autosomal recessive (AR), or X-linked recessive (XLR) pattern. AD disease generally has the best prognosis.

The presenting feature is night blindness.

Investigations:

• Perimetry- the initial stage will have ring scotoma, which further develops into tunnel vision.

• Electroretinography- reveals a decreased amplitude of both "a" and "b" waves.

Atypical types of RP are as follows:

• Sectoral RP

• Pericentric/Inverse RP

• RP sine pigmento

• RP punctata albescens

There is no effective treatment for RP.

Solution to Question 7:

For the 2-month-old girl presenting with leukocoria in the left eye, likely indicative of a congenital infantile cataract, immediate surgery is the most appropriate management.

Congenital infantile cataract refers to lens opacities present at birth, which can obstruct vision and potentially affect the child's overall development. If left untreated, congenital cataracts, especially unilateral cataracts, can lead to amblyopia, a condition where one eye fails to develop normal vision due to insufficient visual input. Immediate surgical intervention is crucial not only to remove the cataract and restore visual clarity but also to minimize the risk of amblyopia, ensuring the best possible outcomes for the child's visual and developmental health. Prompt diagnosis and treatment can significantly enhance the chances of preserving and improving the child's vision.

Leukocoria, or the white pupillary reflex or cat's eye reflex, is a critical sign that often indicates serious underlying eye disorders requiring prompt evaluation and management. Common causes of leukocoria include:

• Retinoblastoma

• Congenital cataract

• Coloboma of the choroid

• Persistent hyperplastic primary vitreous

• Toxocara infestation

• Coat's exudative retinopathy.

Note: Retinoblastoma, a malignant tumour of the retina, often affects children under 6 years of age and can present with symptoms such as leukocoria, strabismus (crossed eyes), or vision problems. Treatment options vary widely depending on factors like the tumour's size, location, and whether it is unilateral or bilateral.

Solution to Question 8:

The given MRI shows a tumour protruding out of sella turcica. This is likely to be a pituitary tumour(pituitary macroadenoma) compressing the optic chiasma, resulting in bitemporal hemianopia.

Pituitary macroadenoma can compress the optic chiasm, causing bitemporal hemianopia, which leads to the loss of vision in the temporal halves of the visual fields in both eyes. This occurs when damage to the middle part of the optic chiasm affects the crossing fibres from the nasal retina of each eye while the uncrossed fibres from the temporal retinas remain intact. As a result, individuals retain central vision but lose peripheral vision on the sides. The most common cause of bitemporal hemianopia is pituitary adenoma. Other common causes include suprasellar aneurysms, chronic arachnoiditis, and suprasellar tumours like craniopharyngioma and

suprasellar meningioma.

Individuals with this condition can see in both left and right central visual fields, with both eyes open. However, all information from the most peripheral parts of visual fields (which are seen only by the nasal retinas) is lost.

A pituitary adenoma is the most common tumour of the sellar region. Depending on their size they are broadly classified into:

• Pituitary microadenoma: < 1 cm in size

• Pituitary macroadenoma: > 1 cm in size and usually extending superiorly.

Clinical features include bi-temporal hemianopia, headache, and vomiting. Skull X-ray finding is ballooning of the sellae. MRI is the investigation of choice. Indentation at the diaphragma sellae gives rise to a snowman/figure of eight appearance, which is seen in the pituitary adenoma as shown below.

Snowman appearance – Pituitary adenoma



Other options:

Option A: Complete blindness refers to a total lack of vision, which can result from various causes, including severe damage to the eyes themselves (like retinal detachment, severe glaucoma, or optic nerve damage) or the visual pathways leading to the brain. Individuals who are completely blind have no visual perception at all.

Option B: Homonymous hemianopia (HH) involves loss of vision on the same side of the visual field in both eyes. HH is characteristically seen in lesions involving the visual pathway posterior to the optic chiasma.

Option C: Binasal hemianopia is seen in lesions affecting bilateral lateral chiasma where the temporal fibres of both the retina continue forward without crossing. This affects vision in the nasal quadrants of both eyes.

Solution to Question 9:

A red nodule on the sclera, as shown in the image, with a history of recurrence, suggests a diagnosis of nodular episcleritis. It is often associated with systemic conditions like rheumatoid arthritis.

Episcleritis is benign, recurrent, inflammation of the episclera, involving the overlying Tenon's capsule but not the underlying sclera. It is more common in women than in men. It is an allergic reaction to an endogenous toxin. Clinically, two types of presentations may occur:

• Simple episcleritis is characterized by sectorial or diffuse inflammation of the episclera.

• Nodular episcleritis is characterized by a pink or purple flat nodule surrounded by injection, usually situated 2-3 mm away from the limbus. The nodule is firm, and tender and can be moved separately from the sclera, and the overlying conjunctiva also moves freely.

Management of episcleritis typically includes topical NSAIDs, corticosteroid eye drops, cold compresses, and artificial tears. For recurrent cases, systemic NSAIDs may be required. 2.5% phenylephrine can be used to differentiate episcleritis and scleritis. Since the vessels of the episclera are superficial, they blanch, whereas, in scleritis, the vessels do not blanch.

Other options:

Option A: Pinguecula is a degenerative condition of the conjunctiva, characterized by the formation of a yellowish-white patch on the bulbar conjunctiva near the limbus. It usually presents as a triangular patch on the nasal side, with the triangle's apex commonly pointed away from the cornea.



Option C: Pterygium is a wing-shaped, triangular fibrovascular ingrowth of degenerative conjunctival tissue that can encroach onto the cornea. It is a response to prolonged exposure to ultraviolet rays, dry heat, etc. This condition can be treated symptomatically with tear substitutes for dry eye symptoms and patients are advised to avoid ultraviolet rays.



Option D: Scleritis refers to an inflammation of the sclera proper. It usually occurs in elderly patients (40-70 years) involving females more than males. It is frequently associated with an underlying systemic inflammatory condition. The patient presents with moderate to severe pain, redness, lacrimation, photophobia, and diminution of vision.

Solution to Question 10:

The visual field defect in the given image is the right homonymous hemianopia.

Right homonymous hemianopia, marked by loss of vision in the right half of the visual field of both eyes, indicates a lesion in the left optic tract. This occurs because the optic tract transmits visual information from the left nasal retina and right temporal retina, and disruption leads to right-sided visual field loss.

This condition highlights a post-chiasmal visual pathway issue, primarily affecting the optic tract, which processes contralateral visual fields. Common causes of lesions include pituitary adenomas, craniopharyngiomas, optic thalamus tumours, and aneurysms of the superior cerebellar or posterior cerebral arteries, as well as demyelinating diseases and infarctions.

Complete homonymous hemianopia affects the complete visual hemifields of both eyes due to large lesions posterior to the optic chiasma. In contrast, smaller lesions can cause incomplete homonymous hemianopia, categorized as congruous or incongruous based on the similarity of defects in each half field. Lesions in the immediate post-chiasmal region often produce incongruent defects, while more posterior lesions in the occipital cortex result in congruous homonymous hemianopia.

Altitudinal hemianopia is defined as hemianopia affecting the upper or lower half of the visual field. It is most commonly seen in anterior ischemic optic neuropathy (AION). Also seen in suprasellar tumors and occipital lobe lesions.

Solution to Question 11:

Given the patient's symptoms—acute pain, decreased vision in the left eye, circumcorneal congestion, irregularly shaped pupil and keratic precipitates, as shown in the image—the most likely diagnosis is acute anterior uveitis. This condition, also known as iritis, involves inflammation of the iris and anterior chamber.

Key features of acute anterior uveitis include unilateral eye pain, circumcorneal redness, photophobia, and reduced visual acuity. The pupil may appear irregular or constricted, and keratic precipitates (KP) on the corneal endothelium are pathognomonic for this condition. Inflammation leads to pseudo-rubeosis iridis and aqueous flare, a hazy appearance in the anterior chamber due to inflammatory exudate.

Anterior uveitis involves inflammation of the uvea, which can cause sluggish pupillary reactions and a "muddy" appearance of the iris. Aqueous flare is often the earliest sign seen during slit-lamp examination, alongside Koeppe and Busacca nodules and potential posterior synechiae formation.

Management focuses on reducing inflammation and managing symptoms, typically using topical corticosteroids and cycloplegics like homatropine or atropine to relieve pain and prevent synechiae. Monitoring intraocular pressure is also essential, as cycloplegics are contraindicated in angle-closure glaucoma due to the risk of exacerbating the condition.

Other options:

Option A: Angle-closure glaucoma involves a sudden increase in intraocular pressure due to blockage of the anterior chamber angle. It is characterized by severe eye pain, blurred vision, and a mid-dilated pupil, with diffuse redness rather than the circumcorneal congestion seen in uveitis.

Option C: Acute conjunctivitis typically presents with diffuse redness, discharge, and mild irritation, without significant pupil changes or severe vision loss.

Option D: Episcleritis features sectoral or diffuse redness with mild tenderness but does not cause severe pain, irregular pupil, or substantial visual changes seen in anterior uveitis.

Solution to Question 12:

The given clinical scenario, supported by the image of the mid-dilated pupil, with a shallow anterior chamber (AC), is suggestive of acute angle-closure glaucoma and the appropriate management for this patient includes intravenous mannitol and topical medications like pilocarpine, which is given to reduce the IOP, and constrict the pupil and help relieve the acute angle closure attack.

The primary goal in managing Primary Angle-Closure Glaucoma (PACG) is to prevent optic nerve damage by reducing IOP. In acute attacks, treatment involves aggressively lowering IOP using intravenous mannitol and oral acetazolamide to quickly control symptoms. Topical medications such as pilocarpine, beta-blockers, cholinergic agents, and carbonic anhydrase inhibitors are also used to address the acute episode.

Once the immediate crisis is managed, YAG laser iridotomy (YAG PI) is performed to create an alternate drainage pathway for the aqueous humor, both in the affected eye and the fellow eye. For ongoing management, patients with PACG may require glaucoma medications to maintain target IOP. In cases where maximal medical therapy fails, surgical options such as trabeculectomy

or glaucoma drainage devices may be necessary, particularly if there is progressive visual field loss or non-compliance with treatment. Cyclodestructive procedures may be considered for patients with a painful, blind eye or those with recurrent surgical failures.

Acute primary angle-closure glaucoma refers to the occlusion of the trabecular meshwork by the peripheral iris (iridotrabecular contact [ITC]), obstructing aqueous outflow.

Angle closure can be primary when it occurs in an anatomically predisposed eye or secondary to another ocular or systemic factor.

Classification:

Symptoms: Pain, redness, photophobia, impaired vision, watering

Signs:

• Corneal edema- coloured halos

• Conjunctiva- congestion, chemosis

• IOP- elevated 40–70 mm Hg

• AC very shallow

• Pupil- vertically oval, mid-dilated

• Angle- completely closed, Grade 0

Primary angle-closure suspect (PACS)	Gonioscopy shows posterior meshwork ITC in three or more quadrants but no peripheral anterior synechiae (PAS).Intraocular pressure (IOP), optic disc, and visual field are normal.No PAS are seenThe risk of PACG at 5 years may be around 30%.
Primary angle closure (PAC)	Gonioscopy shows three or more quadrants of ITC with raised IOP and/or PAS or excessive pigment smudging on the TM.Normal optic disc and field.
Primary angle-closure glaucoma (PACG)	ITC in three or more quadrants, with glaucomatous optic neuropathy.Optic nerve damage from an episode of severe IOP elevation, such as acute angle closure, may not appear as typical glaucomatous cupping.

Solution to Question 13:

The image shows tiny lesions at the limbus, indicating Hebert's pits. This finding is seen in trachoma, which is caused by Chlamydia trachomatis, more specifically the trachoma inclusion conjunctivitis (TRIC) serovars A, B, Ba, C.

Trachoma is a form of chronic infective conjunctivitis and is one of the leading causes of blindness. It has an ‘active’ inflammatory stage and a ‘cicatricial’ chronic stage.

The active inflammatory stage is more prevalent in children and presents with nonspecific symptoms of ocular irritation, foreign body sensation, and discharge (often mild symptoms). Examination findings at this stage show:

• Conjunctival follicles contain Leber cells (multinucleated lymphocytes)

• corneal Herbert's follicles (similar to conjunctival follicles)

Cicatricial chronic stage is seen in middle age. And when the chronic infection with sequelae sets in, the patient complains of pain, lacrimation, photophobia, and later blurring, and finally loss of vision. Conjunctiva shows concretions (whitish deposits of inspissated mucus in depressions called glands of Henle). Corneal signs include:

• Herbert pits (pitted scars in the limbal which are remnants of healed Herbert follicles)

• corneal opacity is seen in later stages.

There can be associated trichiasis (when at least one eyelash rubs the eyeball) and entropion.

At present WHO grading with 'FISTO' is in use.

Trachoma is treated according to the WHO classification using the SAFE strategy (Surgery, Antibiotics, Facial hygiene, and Environmental improvement). Azithromycin is the antibiotic of choice, and surgery involves the correction of trichiasis and entropion. In case of significant corneal scarring, penetrating keratoplasty is done.

Other options:

Option B: Horner–Trantas dots (image below) are distinct white, chalky, gelatinous nodules made up of eosinophils and epithelial debris that appear at the limbus, and they are characteristic of limbal vernal keratoconjunctivitis (VKC).

Option C: In giant papillary conjunctivitis (GPC), gelatinous papillae initially appear as small, elevated nodules on the conjunctiva. These papillae eventually grow larger, forming macropapillae and giant papillae, and may be accompanied by Trantas dots at the limbus.

The image below shows giant papillae in the conjunctiva:

Option D: Progressive pannus in trachoma involves the infiltration of the cornea with new blood vessels developing in the upper part, lying between the epithelium and Bowman's membrane. As the condition advances, these superficial vessels contribute to the destruction of Bowman's membrane, with corneal infiltration occurring before significant vascularization.

Solution to Question 14:

The given clinical history of orbital pain in a contact lens wearer and the demonstration of corneal ulcer and keratitis with scrapings showing an organism with spine-like pseudopodia suggest Acanthamoeba keratitis and the organism implicated in causing this patient's condition is Acanthamoeba.

Acanthamoeba exists as an active trophozoite form and a resistant cystic form, both of which are infective. The large, 20–50 μm trophozoite form is distinguished by spine-like pseudopodia called acanthopodia. Human beings acquire infection by inhalation of cysts or trophozoites, ingestion of cysts or through traumatized skin or eyes. Following inhalation, the trophozoites reach the lungs and invade the central nervous system through the bloodstream, producing granulomatous amebic encephalitis (GAE). It presents chiefly as two chronic conditions: (1) keratitis and (2) encephalitis.

Acanthamoeba keratitis is an infection of the eyes that typically develops from the entry of the amebic cyst through abrasions on the cornea. Most of the cases are associated with the use of contact lenses and individuals who swim or bathe in contaminated water.

Patients usually present with photophobia, excessive tearing, redness, and foreign body sensation, which can be bilateral in contact lens users. Pain is disproportionate to the degree of keratitis due to perineural invasion. It presents as a ring lesion or ring stromal infiltrate as shown below.



Diagnosis of amoebic keratitis is made by the demonstration of the cyst in corneal scrapings by wet mount, histology, and culture. The cysts are smaller (12-25 micrometres) and polygonal or star-shaped.

Topical administration of biguanides or chlorhexidine with or without diamidine agents can be used for the management of acanthamoeba keratitis. In severe cases penetrating keratoplasty can be done.

Polygonal cysts of acanthamoeba in interface contrast microscopy:



Spine like pseudopodia of acanthamoeba:



Other options:

Option A: Balamuthia is a leptomixid free-living amoeba, reported to cause GAE. It exists in the ameboid trophozoite stage and is not associated with acanthopodia as seen in Acanthamoeba species.

Option C: The electron microscopy image of the infective form of Naegleria fowleri shows rounded pseudopodia called lobopodia. The infection usually follows swimming in freshwater rivers or ponds. The amoebae invade the nasal mucosa and pass through the olfactory nerve branches in the cribriform plate into the meninges. It is mainly associated with primary amoebic meningoencephalitis. The disease advances rapidly, causing fever, headache, vomiting, stiff neck, ataxia, seizure and coma.



Option D: Staphylococcus aureus is a common bacterium(not associated with spine-like pseudopodia), that can cause phlyctenular keratoconjunctivitis, characterized by eye redness, pain, and the development of a white-pinkish nodule on the conjunctiva. This condition results from a delayed hypersensitivity reaction to the bacterial proteins.

ENT AIIMS 2017

Question 1:

A 25-year-old woman presented with hearing loss that developed shortly after her first pregnancy. Her audiometry report is given below. What is the diagnosis?



a) Ototoxicity

b) Meniere disease

c) Otosclerosis

d) Noise induced hearing loss

Question 2:

Kraissl's lines are

a) Point of maximum tension in a fracture

b) Relaxed Tension lines in skin

c) Collagen and elastin lines in stab injury

d) Point of tension in hanging

Question 3:

Identify the following line shown in the below picture:



a) Frankfort line

b) Donaldson line

c) Line of Sebileau

d) Ohngren line

Question 4:

The given image is of a patient in whom mastoidectomy is done. Identify the lateral semi-circular canal:



a) A

b) B

c) C

d) D

Question 5:

Which of the following is not a branch of the external carotid artery in Kiesselbach's plexus?

a) Anterior and posterior ethmoidal

b) Sphenopalatine artery

c) Greater palatine artery

d) Septal branch of superior labial

Question 6:

A metal factory worker suffers from noise-induced hearing loss. Which of the following structures will be affected?

a) Crista ampullaris

b) Macula

c) Cupola

d) Outer hair cells

Question 7:

Rhinosporidium seeberi is now classified as a/an _____

a) Aquatic protistan protozoa

b) Fungus

c) Virus

d) Bacteria

Question 8:

Which of the following is not ligated for the control of epistaxis

a) Internal carotid artery

b) Anterior ethmoidal artery

c) Maxillary artery

d) External carotid artery

Question 9:

A 5 year old with gradually progressive hoarseness of voice over three months and respiratory distress for 2 weeks. Diagnosis is

a) Croup

b) Respiratory papillomatosis

c) Vocal nodule

d) Acute epiglottitis

Answer Key

Question No.	Correct Option
1	c
2	b
3	d
4	b
5	a
6	d
7	a
8	a
9	b

Detailed Explanations

Solution to Question 1:

The clinical scenario describes a patient with post-pregnancy hearing loss. The audiogram shows conductive deafness along with Carhart's notch (dip in bone conduction at 2000 Hz). These findings are highly suggestive of otosclerosis.

Ototoxicity, Meniere's disease, and noise-induced hearing loss cause sensory neural hearing loss (SNHL) and not conductive hearing loss.

Otosclerosis is caused by bony overgrowth involving the enchondral layer of the bony otic capsule. It presents with bilateral progressive conductive hearing loss and tinnitus. It has an autosomal dominant inheritance and can worsen with stress and hormonal changes like during pregnancy and menopause.

Stapedial otosclerosis is the most common type of otosclerosis. Here the lesion starts just in front of the oval window in an area called fistula ante fenestrae.

The presence of a reddish hue on the promontory through the tympanic membrane, known as the Schwartze sign, suggests an active focus. PTA shows loss of air conduction with air-bone gap, more for lower frequencies and Carhart's notch may be seen. Tympanometry shows an As type of tympanogram in which there is reduced compliance at or near the ambient air pressure.

Medical management with sodium fluoride is used in the active stage. Definitive treatment is stapedectomy/stapedotomy with the placement of a prosthesis. Carhart notch usually disappears after a successful stapedectomy.

Other options:

Option A: Ototoxicity causes SNHL and shows a high-frequency hearing loss in the SNHL curve.

Option B: In Meniere's disease, pure tone audiometry shows an SNHL pattern with a rising curve in the early stages (as lower frequencies are involved first) and as the disease progresses higher frequencies are involved, the curve becomes flat or a falling type.

Option D: Noise-induced hearing loss is SNHL and PTA shows a characteristic acoustic dip at 4000 Hz both for air and bone conduction.

Solution to Question 2:

Kraissl's lines are relaxed tension lines in the skin. They lie perpendicular to the long axis of the underlying muscle. They were described in living individuals and reflect underlying muscle action.

Image of Kraissl's lines:



Langer's lines correspond to the natural orientation of collagen fibers in the dermis and are generally parallel to the orientation of the underlying muscle fibers. They were described in cadavers in reference to rigor mortis.

Image of Langer's lines:



Borge's lines are also relaxed skin tension lines observed by pinching the skin in living individuals.

Borge's lines and Kraissl's lines are a better guide for elective incisions in the face and body than Langer's lines.

Linear incisions in the skin tend to have variable degrees of gaping. When a circular skin incision is performed, the skin defect acquires an elliptical configuration. By aligning circular incisions perpendicular to the muscle action, surgeons can minimize tension on the wound edges, reduce the risk of scar contracture, and achieve better cosmetic results.

Difference between Kraissl's line and Langer's line

Kraissl's line	Langer's line
Described in living individuals	Described in cadavers
Lie perpendicular to the long axis of the underlying muscle	Correspond to the natural orientation of collagen fibers
Reflects the action of the underlying muscle	They are skin tension vectors observed in the stretched skin of cadavers exhibiting rigor mortis

Solution to Question 3:

The given image shows Ohngren's line. It is an imaginary line passing through the medial canthus of the eye and the angle of the mandible. It is used for the classification of maxillary carcinoma.

Growth situated above the plane (supra structure) tends to be more aggressive and poorly differentiated and has a poor prognosis. Growth situated below the plane (infrastructure) is more amenable to treatment and has a better prognosis.

Other options:

Option A: Proper alignment of the face is determined by the Frankfort plane, a horizontal line bisecting the inferior border of the infraorbital rim and the superior border of the external auditory canal.

Option B: Donaldson's line passes through the horizontal semicircular canal and bisects the posterior canal. It is the landmark for the endolymphatic sac which lies anterior and inferior to it.



ELS: Endolymphatic sac, HSCC: Horizontal semicircular canal, PSCC: Posterior semicircular canal, PF dura: Posterior fossa dura

Option C: Ledermann's classification uses two horizontal lines of Sebileau, one passes through the floor of the orbit and another passes through the floor of the maxillary antra dividing the face into supra structure (ethmoid, sphenoid, frontal sinuses, and the olfactory area of nose), mesostructure (maxillary sinus and respiratory part of nose) and infrastructure (alveolar process). It is used to classify maxillary carcinoma.



Solution to Question 4:

The structure marked as 'B' in the given image represents the lateral (horizontal) semi-circular canal. The tympanic (horizontal) segment of the facial nerve lies below the lateral horizontal semicircular canal.

Other markings represent:

A: Superior semicircular canal (SSC)

C: Posterior semicircular canal (PSC)

D: Incus

The semicircular canals are positioned in the superior part of the mastoid region. These canals are situated adjacent to the middle ear, with the lateral semicircular canal being the nearest to the middle ear. The superior and posterior canals share a common crescent-shaped space in between. The one that lies in close proximity to the lateral semicircular canal is the posterior semicircular canal.



The semicircular canals are part of the bony labyrinth. They house the semicircular ducts which are the end organs for angular momentum. They are 3 semicircular canals, lateral (horizontal), superior (anterior), and posterior. They lie in planes at right angles to one another.

Stimulation of semicircular canals produces nystagmus. The nystagmus is rotatory from the superior canal, horizontal from the horizontal canal, and vertical from the posterior canal.



Solution to Question 5:

Anterior and posterior ethmoidal arteries are branches of the ophthalmic artery which in turn is a branch of the internal carotid artery.

Kiesselbach's plexus is an arterial anastomosis at the anterior inferior part of the nasal septum (Little's area). It includes the anterior ethmoidal artery, septal branch of the superior labial artery,

septal branch of the sphenopalatine artery, and septal branch of the greater palatine artery. It is the usual site of epistaxis in children and young adults.

The blood supply of the nasal septum can be divided into two main systems, the internal carotid system, and the external carotid system.

• Internal carotid system: Branches of the ophthalmic artery, anterior ethmoidal artery, and posterior ethmoidal artery.

• External carotid system: Sphenopalatine artery (branch of the maxillary artery) gives nasopalatine and posterior medial nasal branches, septal branch of the greater palatine artery (branch of the maxillary artery), and septal branch of the superior labial artery (branch of the facial artery).



Solution to Question 6:

Noise-induced hearing loss (NIHL) damages the outer hair cells, starting in the basal turn of the cochlea. Outer hair cells are affected before the inner hair cells.

NIHL occurs after chronic exposure to less intense sounds. The damage caused by noise trauma depends on the frequency of noise (frequency of 2000 to 3000 Hz causes more damage) intensity, duration of the noise, and preexisting ear disease.

The audiogram in noise-induced hearing loss shows a typical notch at 4000Hz both for air and bone conduction known as acoustic dip/boiler's notch. Patients at this stage complain of high-pitched tinnitus and difficulty hearing in noisy surroundings. Pre-employment and annual audiograms for persons working at places where noise is above 85 dB (A) help in the early detection of NIHL.

The human ear can hear frequencies from about 20 to 20,000 Hz. Below 20 Hz, the vibrations are infra-audible, and above 20,000 Hz, they are ultra-sonic. Pure tone audiometry can detect only up to 8000 Hz.

Extended high-frequency audiometry is recommended for early detection of hearing loss, as it can identify hearing impairments before the frequencies necessary for speech comprehension are affected. Distortion product otoacoustic emissions is a more sensitive tool than conventional audiometry, particularly in children under 5 years old for early detection of hearing loss.

A noise of 90 dB(A) SPL (acceptable noise levels, sound pressure level), 8 hours a day for 5 days per week is the maximum safe limit as recommended by the Ministry of Labour. As per WHO, 85 dB(A) is considered the highest safe exposure level up to a maximum of 8 hours.

The silence zone extends to a distance of 100 meters around the premises of hospitals, nursing homes, educational institutions, and courts.

Permissible limits of noise as per the noise pollution (Regulation and Control) rules:

Noise levels are measured in dB(A), which prioritizes middle-range frequencies for assessing noise-induced hearing loss while giving less weightage to low and high frequencies.

The image below shows an acoustic dip at 4000Hz:

Zone/Area	Day(6 am to 10 pm) limits in dB(A)	Night(10 pm to 6 am) limits in dB(A)
Industrial	75	70
Commercial	65	55
Residential	55	45
Silence	50	40



Other options:

Option A: Crista ampullaris is a thickened ridge of neuroepithelium in the ampullated end of semicircular ducts.

Option B: Macula is the sensory epithelium of the utricle

Option C: Cupula is a gelatinous mass extending from the surface of the crista (peripheral receptor located in the ampullated end of semicircular ducts) to the ceiling of the ampulla, into which project the cilia of sensory hair cells.

Solution to Question 7:

Rhinosporidium seeberi is not a classic fungus, but rather the first known human pathogen from the DRIPs (Dermocystidium, Rosette agent, Icthyophonus, and Posorospermum) clade, a novel clade of aquatic protistan parasites (Ichthyosporea).

It causes a granulomatous disease called rhinosporidiosis. It is endemic in south India. Patients typically have a history of bathing in water bodies shared by animals and present with recurrent epistaxis, nasal obstruction by a red vascular mass (mulberry mass), and hypertrophic turbinates.

For diagnosis, a nasal smear can be taken to demonstrate sporangia.

Treatment is surgical excision of the mass with a diathermy knife and cauterization of base. Dapsone can also be used postoperatively to reduce the incidence of recurrence.

The image below shows a polypoidal mass of rhinosporidiosis:

Solution to Question 8:

The internal carotid artery is not ligated in epistaxis control due to a high percentage of cerebral complications and high mortality.

The first step in treating epistaxis is to pinch the nose with the thumb and index finger for about 5 minutes. The patient should sit leaning slightly forward over a basin to spit out blood and breathe through the mouth (Hippocratic or Trotter's method).

If the bleeding point is localized, cauterization can be done with electrocautery. In cases of uncontrolled bleeding or when the bleeding is posterior into the throat, anterior or posterior nasal packing may be required, respectively. If bleeding continues despite these measures, endoscopic procedures or vessel ligation can be considered.

The sphenopalatine artery is ligated first through a trans-nasal endoscope (Transnasal endoscopic sphenopalatine artery ligation). However, if the bleeding persists, maxillary artery ligation is done. If bleeding is still not controlled, external carotid artery ligation is done. It is important to note that if the bleeding site is located superior to the level of the middle turbinate, the anterior and posterior ethmoidal arteries should be ligated.

Solution to Question 9:

The given clinical scenario of a child presenting with chronic hoarseness (3 months) of voice and respiratory distress for 2 weeks suggests the diagnosis of respiratory papillomatosis (juvenile papillomatosis).

Croup (Option A) and acute epiglottitis (Option D), which cause acute upper respiratory tract obstruction, are ruled out due to the chronic nature of the symptoms. In croup, the upper airway obstruction becomes apparent after 1 to 3 days of low-grade fever, rhinorrhea, and pharyngitis. On the other hand, in epiglottitis, a child experiences respiratory distress within hours of developing a fever.

Vocal nodules (Option C) are not typically associated with respiratory distress.

Respiratory papillomatosis is the most common respiratory tract neoplasm in children. They are simple warts (benign tumors) caused by human papillomavirus types 6 and 11 commonly seen in children aged 3 to 5 years.

The course involves remissions and exacerbations of recurrent papillomas, mostly affecting the glottic area (vocal cords), and supraglottis. The patient presents with progressively worsening hoarseness, sleep-disordered breathing, dyspnea, stridor, and severe airway obstruction if left untreated.

Diagnosis confirmed by direct laryngoscopy, which shows grape-like projections (shown below), and biopsy. Treatment consists of micro laryngoscopy and CO_2 laser excision. Antivirals like interferon alpha2a, ribavirin, cidofovir, and acyclovir are used as adjunct therapies.



Other options:

Option A: Croup (laryngotracheobronchitis) is caused by the parainfluenza virus. The patient initially presents with an upper respiratory tract infection (rhinorrhea, pharyngitis, cough, low-grade fever) followed by a characteristic barking cough, hoarseness, and inspiratory stridor.

Option C: Vocal nodule (singer's or screamer's nodule) occurs as a result of chronic vocal abuse. They appear symmetrically at the junction of the anterior one-third and posterior two third of the vocal cord. The patient presents with chronic hoarseness and voice fatigue.

The image below shows symmetric vocal nodules:

Option D: Acute epiglottitis is caused by streptococcus pyogenes, pneumococci, Hemophilus influenzae, and staphylococcus aureus. It is characterized by an acute rapidly progressive course of high-grade fever, dyspnea, and stridor.

ENT AIIMS 2018

Question 1:

While performing a caloric test, what type of nystagmus would you expect on cold water irrigation?

a) Horizontal nystagmus towards the same side

b) Vertical nystagmus towards the same side

c) Horizontal nystagmus towards the opposite side

d) Vertical nystagmus towards the opposite side

Question 2:

In which of the following conditions is Gelle's test negative?

a) CSOM with cholesteatoma

b) Senile deafness

c) Meniere's disease

d) Otosclerosis

Question 3:

What is the definitive treatment of intractable vertigo due to Meniere's disease?

a) Acetazolamide

b) Labyrinthectomy

c) Vestibular neurectomy

d) Intratympanic gentamicin

Question 4:

A patient comes with an ulcer on the tongue with a duration of 3 months. On examination, there is a 3.5 cm x 2.5 cm lesion with induration of 8 mm. There was no lymphadenopathy. What is the stage of this lesion according to 8th AJCC guidelines?

a) Stage I

b) Stage II

c) Stage III

d) Stage IV

Question 5:

Which of the following features will not be manifested in an infection of the post-styloid compartment of the parapharyngeal space?

a) Trismus

b) Vocal cord palsy

c) Parotid swelling

d) Horner's syndrome

Question 6:

A patient presents with oral cavity findings are seen in the image below. He is a smoker. What would you choose as your next step in the management of this patient?

a) Give corticosteroids

b) Cessation of smoking and do biopsy

c) Lifestyle modifications

d) Injection vitamin B12

Question 7:

A teenager has a mass in his nasopharynx that has minimally spread into the sphenoid sinus with no lateral extension. Which stage of nasopharyngeal angiofibroma could this be?

a) Stage IA

b) Stage IB

c) Stage IIA

d) Stage IIB

Question 8:

A patient complains that he feels like the room is spinning when he gets up from lying down or turning his head. He has no history of loss of consciousness. Which of the following could be the probable diagnosis?

a) Benign paroxysmal positional vertigo

b) Meniere's disease

c) Labyrinthitis

d) Syncope

Answer Key

Question No.	Correct Option
1	c
2	d
3	b
4	b
5	a
6	b
7	b
8	a

Detailed Explanations

Solution to Question 1:

In caloric testing, cold water irrigation produces horizontal nystagmus towards the opposite side of irrigation. The direction of nystagmus can be remembered by the mnemonic, COWS (cold–opposite, warm–same).

The mechanism for the nystagmus is the creation of a temperature gradient by the cold water, which pulls the cupula away from the utricle, causing nystagmus towards the oppsite side. As the horizontal semicircular canal is closer to the outer ear canal, most of the response comes from it and the nystagmus is horizontal.

The Fitzgerald–Hallpike test (bithermal caloric test) is done to assess this phenomenon. The patient lies in a supine position with head tilted forward at a 30° angle, ensuring the horizontal canal becomes vertical. Alternatingly, water at temperatures of 30°C and 44°C (7° below and above normal body temperature, respectively) is irrigated into the ears for a duration of 40 seconds each. The examiner closely observes the patient's eyes for the emergence of nystagmus until its endpoint is reached. The time taken from the start of irrigation to the endpoint of nystagmus is recorded and plotted on a calorigram.

If no nystagmus is evoked from either ear, the test is repeated with water at a temperature of 20°C for a duration of 4 minutes before labeling the labyrinth dead. A 5-minute interval should be allowed between testing each ear to ensure accurate results.

Solution to Question 2:

Gelle's test is negative when the ossicular chain is fixed as in cases of otosclerosis.

Gelle's test is a test of bone conduction and examines the effect of increased air pressure in the ear canal on hearing. In normal individuals, due to the increased intralabrynthine pressure applied by the Seigel's speculum, there is immobility of the basilar membrane, as the tympanic membrane and the ossicles are pushed inwards (medially). This results in decreased hearing. But, in diseased individuals, where the ossicular chain and the tympanic membrane are fixed or disconnected, no change in hearing is observed.

Gelle's test is positive in normal persons and in those with sensorineural hearing loss.

Note: This test was commonly used for the diagnosis of otosclerosis, but has since been replaced by tympanometry.

Solution to Question 3:

The definitive treatment of intractable vertigo due to Ménière's disease is labyrinthectomy.

Labyrinthectomy involves uniform destruction of hearing and vestibular function. Ideal candidates have no functional hearing and have failed more conservative treatments, such as gentamicin injection. Despite this morbidity, labyrinthectomy has the highest rate of vertigo control (>95%), more than vestibular neurectomy (78-90%), and is considered the single definitive treatment.

The step-by-step management approach for Ménière's disease consists of dietary modifications and vestibular suppressants as the initial measures, followed by intratympanic gentamicin ablation, and finally, surgical intervention.

Primarily, the management of Menier's disease involves a sodium restricted diet and diuretics like acetazolamide (Option A). They are more useful for short-term control.

Gentamicin (Option D) is a selective vestibulotoxic aminoglycoside antibiotic that is preferentially taken up by type I hair cells. It is opted for only if dietary modifications and vestibular suppressants have failed to control the symptoms. If used judiciously, hearing is preserved.

The options for surgical management include - Endolymphatic sac surgery, vestibular neurectomy, and labyrinthectomy. Endolymphatic sac surgery is currently declining in popularity due to lack of evidence of vertigo control.

Vestibular neurectomy (Option C) refers to selective ablation of the vestibular nerve. Even though it is considered a hearing sparing surgery and carries comparable rates of vertigo control as labyrinthectomy, it is an intradural procedure and carries a greater risk of complications. Hearing loss will still progress as it would have in the natural course of illness.

Solution to Question 4:

The maximum diameter of the ulcer is less than 2-4 cm and the depth of invasion is less than 10mm which implies T2; There is no lymph node involvement (N0) and no metastasis (M0). A T2N0M0 tumor is classified as stage II according to the 8th AJCC guidelines.

8th AJCC guidelines for the staging of oral tumors:

The changes in the 8th AJCC update include:

• Depth of invasion (DOI) and extranodal extension have been included as independent prognostic factors.

• A new category of tumor added - HPV positive, p16 positive

• In the past, the lip was considered a primary site for oral cancers; now - it is divided into mucosal and cutaneous lip, mucosal lip is included in the oral cavity.

The gold standard investigation for oral cancer includes an incisional biopsy. An MRI is the imaging modality of choice for assessing primary extent. Surgical management with chemotherapy and adjuvant radiotherapy is the mainstay of management.

Staging	Tumor Size	Lymph Node	Metastasis
I	T1 (<2 cm, depth <5 mm)	N0 (No regional Lymph Nodes)	M0
II	T2 (2 - 4 cm, depth <10 mm)	N0	M0
III	T3 (>4 cm, depth>10 mm)	N0	M0
III	T1, 2, 3	N1	M0

Staging	Tumor Size	Lymph Node	Metastasis
IVA	T4a (cortical bone of mandible or maxillary sinus or skin of face)	N0,1	M0
IVA	T1, 2, 3, 4a	N2	M0
IVB	Any T	N3	M0
IVB	T4b (masticator space, pterygoid plates, skull base or encases carotid vessels)	Any N	M0
IVC	Any T	Any N	M1

Solution to Question 5:

Trismus is not a feature of infection of the post-styloid compartment of the parapharyngeal space. Trismus is seen in pre-styloid abscess due to the involvement of medial pterygoid muscle.

Parapharyngeal space is divided into two compartments by the styloid process and its attachments. Clinical features depend on the compartment involved.

In a post-styloid abscess, there is a bulge of the pharynx behind the posterior pillar, accompanied by paralysis of cranial nerves IX, X (resulting in vocal cord paralysis), XI, XII, and the sympathetic chain, which can manifest as Horner's syndrome. Swelling in the parotid region may also be present. Notably, trismus, the inability to fully open the mouth, is usually absent in this type of abscess.

A pre-styloid abscess exhibits distinct features including the prolapse of the tonsil and tonsillar fossa, along with trismus caused by the spasm of the medial pterygoid muscle. Marked odynophagia is common. Additionally, an external swelling can be observed behind the angle of the jaw.

Fever, sore throat, torticollis (caused by spasm of the prevertebral muscles), and signs of systemic toxicity are common features.

Diagnosis can be established using a contrast-enhanced CT scan (CECT). If there is suspicion of thrombosis of the internal jugular vein or an aneurysm of the internal carotid artery, magnetic resonance arteriography (MRA) can be useful.

Patients require intravenous antibiotic therapy with amoxicillin-clavulanic acid, carbapenem antibiotics plus drugs that target anaerobic organisms, such as clindamycin or metronidazole. A parapharyngeal abscess is drained by an external horizontal incision made on the skin a few centimeters below the angle of the mandible.

The image attached below shows the components of both spaces:



Solution to Question 6:

The patient, who is a chronic smoker, presenting with a wrinkled white patch on the ventral surface of his tongue is likely to have leukoplakia. Confirmation of this diagnosis will require a biopsy of the lesion. Leukoplakia is a premalignant lesion, with a malignant potential of 1 - 17.5%, hence it is imperative that the patient is advised on cessation of smoking.

Leukoplakia is a premalignant condition of the oral cavity characterized by whitish patches or plaques that cannot be rubbed off. They appear as small, well-circumscribed lesions and can extensively involve a large area of the oral mucosa.

Variants of leukoplakia are:

• Homogenous variant: Smooth white patch. Less association with malignancy.

• Non-homogenous variants:

• Speckled leukoplakia is whitish lesions on an erythematous base. They have the highest risk of malignant transformation.

• Proliferative verrucous leukoplakia is rare and has a high risk for malignant transformation.

• Erosive (erythroleukoplakia) is when it is interspersed with erythroplakia and has erosions and fissures. This also has malignant potential.

Management includes smoking cessation and reduction in alcohol consumption. A biopsy should be done. A CO_2 laser is often used for ablation.

A photographic record of the lesion is useful for long-term follow-up in cases of conservative treatment.

Given below is a list of premalignant lesions of the oral cavity with their risk of malignancy:

High-risk lesions	Medium risk lesions	Low-risk lesions
Erythroplakia(homogenous/speckled)Seen asred velvetylesionsSpeckledtype is the most aggressiveThe treatment of choice issurgicalablation	Oral submucous fibrosisHypersensitivereactionto betel nut product.Other risk factors are - tobacco, chili, vitamin deficiencyClinical features - trismus, ankyloglossia, soreness, and burning sensation in the mouth.Management includes antioxidants, intralesional triamcinolone, and surgical excision.	Oral lichen planusReticulate fine lace-like pattern.Management istopicalcorticosteroids.
Proliferative verrucous leucoplakiaAwhite patchylesion thatcannot be rubbed off.Speckled leucoplakia has an erythematous baseThe treatment of choice is excision	Syphilitic glossitisAssociated with tertiary syphilis	Discoid lupus erythematosus
Chronic hyperplastic candidiasisWhite patchesthatcan be rubbed offwith ahyperaemicborder.Management includes prolonged treatment withtopicalantifungalsorsystemicantifungaltherapy. If no improvement or if there are any signs of dysplasia,surgical excision is done.		Dyskeratosis Congenita

Solution to Question 7:

The mass has spread into the sphenoid sinus with no lateral extension, so it's Stage 1B according to the Radkowski classification.

Multiple classification systems exist for the staging of juvenile nasal angiofibroma, of which the most commonly used ones are the Fisch and Radkowski classification systems:

Staging of juvenile nasopharyngeal angiofibroma (Radkowski, et al):

Fisch staging system of juvenile nasopharyngeal angiofibroma:

Juvenile nasopharyngeal angiofibroma (JNA) is a highly vascular and locally aggressive tumor seen almost exclusively in adolescent males, associated with significant morbidity and occasional mortality.

Note: The Fisch system is considered to be the more robust and practical of the two.

Stage	Spread
IA	Limited to the nose and/or nasopharyngeal vault
IB	Extension into ≥ 1 sinus
IIA	Minimal extension into the pterygopalatine fossa
IIB	Full occupation of pterygopalatine fossa with or without erosion of orbital bones
IIC	Infratemporal fossa with or without cheek or posterior to pterygoid plates
IIIA	Erosion of skull base-minimal intracranial
IIIB	Erosion of skull base-extensive intracranial with or without cavernous sinus

Type	Details
1	Tumourlimited to the nasopharyngeal cavity; bonedestruction negligibleor limited to the sphenopalatine foramen
2	Tumourinvading the pterygopalatine fossaor the maxillary, ethmoid, or sphenoidsinus with bone destruction
3	A:Tumour invading theinfratemporal fossaororbital region withoutintracranial involvement
B:Tumour invading theinfratemporal fossaororbital region withintracranialextradural(parasellar) involvement	
4	A:Intracranialintraduraltumorwithoutinfiltration of the cavernous sinus, pituitary fossa, or optic chiasm
B:Intracranialintraduraltumorwithinfiltration of the cavernous sinus, pituitary fossa, or optic chiasm	

Solution to Question 8:

The given clinical scenario, with complaints of dizziness associated with postural change and no history of loss of consciousness is suggestive of benign paroxysmal positional vertigo (BPPV).

BPPV is one of the common causes of recurrent vertigo, and the episodes are brief (<1min) and are always provoked by changes in head position. The attacks are caused by free-floating calcium carbonate crystals usually dislodged from the posterior canal.

A positive Dix-Hallpike maneuver confirms the diagnosis of BPPV. In this maneuver, the patient begins sitting up with their head turned 45 degrees toward the ear to be tested. Then the patient is made to lie down quickly with their head past the end of the bed. Then, the patient's neck is extended 20 degrees below the horizontal. The clinician then watches the patient's eyes for torsional and up-beating nystagmus, which should persist for no more than one minute. This would indicate a positive test.

Epley's maneuver is performed to reposition the crystals into the macula from the semicircular canals. This is only a temporary cure and there is a high rate of recurrence.

ENT AIIMS 2019 (May & Nov)

Question 1:

Which of the following marked structures is a branch of the internal carotid artery?



a) 1

b) 2

c) 3

d) 4

Question 2:

Which of the following is not an etiological factor for head and neck malignancies?

a) EBV infection

b) HPV infection

c) Betel nut

d) Vitamin A deficiency

Question 3:

Which of the following structures are not removed in radical neck dissection?

a) Tail of parotid

b) Sublingual gland

c) Submandibular gland

d) Level 2b lymph nodes

Question 4:

Which of the following features is not associated with superior semicircular canal dehiscence syndrome?

a) Gaze dependent nystagmus

b) Positive Fistula test

c) Sensorineural hearing loss

d) Positive Tulio phenomenon

Question 5:

A patient undergoes surgery of the lateral part of the base of the skull. Postoperatively, the patient presents with aspiration but does not have a change in voice. Which of the following is the reason?

a) Superior laryngeal nerve injury

b) Glossopharyngeal nerve injury

c) Recurrent laryngeal nerve injury

d) High vagal injury

Answer Key

Question No.	Correct Option
1	a
2	d
3	b
4	c
5	a

Detailed Explanations

Solution to Question 1:

The structure marked 1 is the anterior ethmoidal artery, which is a branch of the ophthalmic division of the internal carotid artery.

Other structures marked in the diagram are as follows:

2 – Sphenopalatine artery

3 – Greater palatine artery

4 – Superior labial artery

2, 3 and 4 are derived from branches of the external carotid artery.

The nasal septum is supplied by branches of both internal and external carotid arteries.



Solution to Question 2:

Vitamin A deficiency is not a risk factor for head and neck malignancies. Vitamin A deficiency is a risk factor for esophageal cancer.

The head and neck cancers include cancers of the oral cavity, pharynx, larynx, paranasal sinuses, nasal cavity, and salivary glands. Squamous cell cancer is the most common head and neck malignancy, often linked to tobacco and alcohol use. Non-squamous malignancies include thyroid cancer, salivary gland cancer, and sarcomas.

The risk factors for head and neck malignancies are listed below:

Oral cavity cancer:

• Tobacco, smokeless tobacco like Bidi, Chutta, Paan (oral tobacco mixed with betel leaf, areca nut, and salted lime), Khaini, and marijuana

• Alcohol, poor oral hygiene, and intake of red meat and salted meat

• Infections with human papillomavirus, and HIV

• Genetic conditions like Li-Fraumeni, ataxia telangiectasia, Bloom syndrome, and Fanconi's anemia.

Oropharyngeal cancer:

• Smoking and alcohol

• HPV 16 and 18

Cancer of the larynx and hypopharynx:

• Tobacco and alcohol

• Nickel, mustard gas, and asbestos exposure

• Long-term exposure to sulphuric acid and hydrochloric acid in battery plant workers.

• Postcricoid carcinoma is associated with previous radiation and sideropenic dysphagia

• HPV infection and immunosuppression

Nasopharyngeal cancer:

• Chinese population

• Epstein Barr virus infection

• Nitrosamines in salted fish and meat

• Vitamin C deficiency

Cancers of the nasal cavity and paranasal sinus:

• Tobacco

• Occupational exposure to dust from wood, textiles, and leather. Nickel and chromium dust and mustard gas exposure.

• HPV 16 and 18

Thyroid cancer:

• Females

• Previous radiation

• Inherited conditions like Gardner syndrome, familial polyposis, and Cowden syndrome. Medullary thyroid cancer associated with MEN-2 syndrome

Salivary gland carcinoma:

• Previous radiation

• Exposure to nickel alloy dust, and silica dust

• Smoking and heavy alcohol consumption were linked to a higher risk of salivary gland cancer in men but not in women

• Early menarche and nulliparity

Solution to Question 3:

Sublingual gland is not removed in radical neck dissection (RND). Submandibular gland is removed in RND.

RND involves the removal of:

• All lymph nodes from Levels I through to V (submental, submandibular, upper, middle, and lower jugular, and posterior triangle regions) along with its fibrofatty tissue.

• The spinal accessory nerve

• Internal jugular vein

• Sternocleidomastoid muscle

• Tail of parotid

• Submandibular gland

• Omohyoid muscle

Solution to Question 4:

Sensorineural hearing loss is not seen in superior semicircular canal dehiscence syndrome (SSCDS). They may experience mild to moderate hearing conductive loss.

Conductive hearing loss is due to the abnormal transmission and dissipation of acoustic energy through the defect (third window), leading to reduced conduction to the cochlea for lower frequencies. This can manifest as a low-frequency air-bone gap at multiple frequencies. There is no defect in the cochlea or the auditory pathway.

Superior semicircular canal dehiscence (SSCD) is a condition in which bone between the apex of the superior semicircular canal and the middle cranial fossa is absent. This causes endolymph in the labyrinthine system to move with sound or pressure, causing an activation of the vestibular system.

Patients may have vestibular symptoms like vertigo triggered by loud sounds (Tullio phenomenon)or pressure such as coughing and can also experience hyperacusis for bone-conducted sounds. They may perceive the sounds of their own eye movements or footsteps in the affected ear.

The patients may exhibit nystagmus when exposed to loud sounds or during the Valsalva maneuver or pressure in the ear canal (positive fistula test). The direction of the nystagmus aligns with the affected superior semicircular canal, and some patients may tilt their heads in the plane of the affected superior semicircular canal in response to loud sounds.

Noncontrast HRCT of the temporal bone can demonstrate the defect. The click-evoked VEMP (vestibular evoked myogenic potential) response and vestibuloocular reflex are pronounced in SSCD patients, in which the evoked eye movements align with the affected superior canal.

For mild symptoms, avoidance of triggers is recommended. In case of severe disabling symptoms, surgical resurfacing of the superior semicircular canal or plugging of the superior semicircular canal may be attempted.

Solution to Question 5:

In the given clinical scenario, the patient experiences aspiration but does not exhibit any change in voice following base of skull surgery. This suggests an injury to the superior laryngeal nerve (SLN).

In SLN injury, only the cricothyroid muscle is involved which is used only in high-pitch vocalization, normal speech will not be affected. SLN also provides sensory supply to the mucosa above vocal cords, hence its lesion would lead to increased susceptibility to aspiration.

Other options:

Option B: The glossopharyngeal nerve does not provide sensory or motor supply to the larynx.

Option C and D: The patient also has normal vocalization which rules out recurrent laryngeal nerve (RLN) and high vagal injury. RLN supplies all the intrinsic muscles of the larynx except cricothyroid, therefore an RLN lesion would lead to disordered vocalization. A high vagal injury would mean that both the SLN and RLN functions are affected, therefore that can also be ruled out.

ENT AIIMS 2020 (May and Nov INI-CET)

Question 1:

A patient underwent total laryngectomy for laryngeal cancer. Identify the modality of voice production in this patient shown in the image given below.



a) Electro-larynx

b) Chicago implant

c) Tracheoesophageal prosthesis

d) Esophageal speech

Question 2:

Anterior most air cells of anterior ethmoidal sinus are known as?

a) Bulla ethmoidalis

b) Onodi cell

c) Agger nasi

d) Haller cell

Question 3:

A patient complaining of vertigo without hearing loss has consulted an ENT surgeon. The surgeon performs a diagnostic maneuver and cautiously performs a therapeutic maneuver. What is the diagnostic maneuver?

a) Hampton's maneuver

b) Dix-Hallpike maneuver

c) Epley's maneuver

d) Simon's maneuver

Question 4:

Identify the retractor shown in the below image?



a) Joll's retractor

b) Perkins retractor

c) Mastoid retractor

d) Langenbeck retractor

Question 5:

A senior resident in the hospital is performing the following clinical examination as shown in the image below. Which of the following structures will not be visible to him?



a) Adenoids

b) Arytenoids

c) Torus tubarius

d) Upper surface of soft palate

Question 6:

Which of the following is not a branch of the external carotid artery in Kiesselbach's plexus?

a) Anterior and posterior ethmoidal

b) Sphenopalatine artery

c) Greater palatine artery

d) Septal branch of superior labial

Question 7:

The radiograph shown below is done for better assessment of the frontal sinus. What is the common name of this view?

a) Water's view

b) Caldwell view

c) Pierre's view

d) Towne's view

Question 8:

Which of the following is not seen in a case of otitis media?

a) Bezold abscess

b) Griesinger's sign

c) Battle sign

d) Delta sign

Question 9:

Given below is the endoscopic view of the right nasal cavity. Identify the marked structure?



a) Inferior Turbinate

b) Middle turbinate

c) Superior Turbinate

d) Uncinate process

Question 10:

Which of the following are the objective tests for the hearing?

a) 1, 2, and 4

b) 1,2, and 3

c) 2,3, and 4

d) 1,2,3, and 4

Question 11:

Identify the correct sequence of the auditory pathways: A.Superior Olivary nucleus B. Cochlear Nucleus C. Inferior Colliculi D. Lateral lemniscus E. Medial geniculate ganglion

a) ABCDE

b) BADCE

c) BACDE

d) BAEDC

Question 12:

A 55-year-old female came to the hospital with a history of trauma to the ear. On examination, there is swelling, tenderness, and discoloration of the ear. Which of the following is false regarding the given image?



a) All cases should receive prophylactic antibiotic

b) May lead to Pugilistic ear

c) Caused by accumulation of blood in perichondrial space

d) Resolves spontaneously

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	c
5	b
6	a
7	b
8	c
9	a

10	a
11	b
12	d

Detailed Explanations

Solution to Question 1:

The modality of voice production shown in the given image is a tracheoesophageal voice prosthesis.

In a tracheoesophageal prosthesis, the exhaled air from the trachea is diverted to the esophagus through a device that connects the trachea and esophagus. It has a one-way valve, which prevents food from entering into the trachea. A variety of tracheoesophageal prostheses are available including the provox, Blom-Singer, and Panje devices.

Other options:

Option A: Electrolarynx is a vibrating device that produces a low-pitched sound in the hypopharynx. It is then articulated by the palate, tongue, and teeth.



Option B: Chicago implant is a hearing aid.

Option D: In oesophageal speech, the patient is trained to swallow air and slowly release it into the pharynx. 6-10 words can be spoken in a rough manner before re-swallowing the air.



Solution to Question 2:

Agger nasi is the anterior-most air cells of anterior ethmoid air cells. When enlarged it may cause mechanical obstruction to frontal sinus drainage.

Other options:

Option A: Bulla ethmoidalis is the largest air cell in the ethmoidal air sinus. It is situated behind the uncinate process and forms the posterior boundary of hiatus semilunaris through which the maxillary, frontal, and anterior ethmoidal sinuses drain.

The image below shows the ethmoidal bulla:



Option B: Onodi cell is formed from posterior ethmoidal air cell. The optic nerve lies in its lateral wall and can be injured while operating in the cells.

The image below shows Onodi air cell:



Option D: Haller cells are formed from either anterior or posterior ethmoid air cells. They present in relation to the floor of the orbit and roof of the maxillary sinus, enlargement may prevent drainage of the maxillary sinus.

The image below shows Haller cell:



Solution to Question 3:

The given clinical scenario of a patient with vertigo without hearing loss is suggestive of benign paroxysmal positional vertigo (BPPV). The diagnostic maneuver used in BPPV is the Dix-Hallpike maneuver, the gold standard for diagnosing BPPV. The therapeutic maneuver is Epley's maneuver.

BPPV is characterized by vertigo, recurrent brief (<1 minute) episodes triggered by specific head movements, and is not associated with hearing loss or other neurological symptoms.

It is thought to be caused by the dislodgement of crystals of calcium carbonate from the degenerating macula of the utricle. When this crystal settles on the cupula of the posterior canal, it causes displacement of the cupula and vertigo.

A patient presenting with vertigo should be evaluated for ruling out the central lesions (like a tumor in the fourth ventricle, cerebellum, temporal lobe, or multiple sclerosis) before confirming the diagnosis of BPPV. The Dix-Hallpike maneuver helps to differentiate a peripheral from a central lesion.

Dix-Hallpike maneuver: The maneuver starts with the patient in a sitting position. Their head is turned 45 degrees towards the ear to be tested. Then the examiner quickly brings the patient to a supine position with their head 30 degrees below the horizontal. The 45-degree head rotation is maintained during the procedure. Now the examiner observes the patient for nystagmus.



The features of nystagmus differentiate between BPPV and central lesions:

The therapeutic maneuver is the Epley maneuver which displaces the otolith out of the posterior canal. This ends the sensation of vertigo with positional changes relieving the symptoms.

Nystagmus	BPPV	Central lesions
Latency	2-20 seconds	No latency
Duration	<1 minute	>1 minute
Direction	fixed, towards the lower-placed ear	changing
The result on repetitions of the test	nystagmus is not seen (fatigable)	nystagmus persists (nonfatiguable)
Symptoms during the test	severe vertigo	Absent/mild

Solution to Question 4:

The instrument shown in the image is Mollison's mastoid retractor. It is used in mastoidectomy to retract soft tissues after incision and elevation of flaps. It is self-retaining and hemostatic.

Jansen's self-retaining mastoid retractor (shown below) is also used in mastoidectomy and is similar to Mollison's retractor.

Option A: Joll retractor (shown below) is a self-retaining retractor used in thyroid surgery.

Option D: Langenbeck retractor is useful for holding open wounds, as in an open appendicectomy and they are often used in pairs (one in each of the assistant's hands).

Solution to Question 5:

The senior resident would not be visualizing the arytenoids as he is performing a posterior rhinoscopy indicated by the post-nasal mirror that he is holding.

The posterior rhinoscopy mirror (St. Clair Thompson mirror) has two bends in the shaft and is bayonet shaped.

The structures seen in the posterior rhinoscopy are:

• Adenoids

• Posterior ends of nasal turbinates

• Opening of the Eustachian tube

• The posterior border of the nasal septum

• Fossa of Rosenmuller

• Torus tubarius

• The upper surface of the soft palate

The images below show the posterior rhinoscopy mirror and structures seen in the posterior rhinoscopy:



Indirect laryngoscopy is done using a laryngeal mirror. It is straight without any bends with a mirror at the end.

Structures seen during indirect laryngoscopy (in order):

• Oropharynx: Base of the tongue, lingual tonsils, valleculae, medial and lateral glossoepiglottic folds.

• Laryngopharynx: Both piriform fossae, post cricoid region, posterior wall of laryngopharynx.

• Larynx: Epiglottis, aryepiglottic folds, arytenoids, cuneiform and corniculate cartilages, glossoepiglottic folds, pharyngoepiglottic folds, aryepiglottic fold, true and false vocal cords, anterior commissure, posterior commissure, subglottis and rings of the trachea.

The images below show the laryngeal mirror and structures seen during indirect laryngoscopy:



Solution to Question 6:

Anterior and posterior ethmoidal arteries are branches of the ophthalmic artery which in turn is a branch of the internal carotid artery.

Kiesselbach's plexus is an arterial anastomosis at the anterior inferior part of the nasal septum (Little's area). It includes the anterior ethmoidal artery, septal branch of the superior labial artery,

septal branch of the sphenopalatine artery, and septal branch of the greater palatine artery. It is the usual site of epistaxis in children and young adults.

The blood supply of the nasal septum can be divided into two main systems, the internal carotid system, and the external carotid system.

• Internal carotid system: Branches of the ophthalmic artery, anterior ethmoidal artery, and posterior ethmoidal artery.

• External carotid system: Sphenopalatine artery (branch of the maxillary artery) gives nasopalatine and posterior medial nasal branches, septal branch of the greater palatine artery (branch of the maxillary artery), and septal branch of the superior labial artery (branch of the facial artery).



Solution to Question 7:

The frontal sinus is best seen in Caldwell's view (occipitofrontal view or nose-forehead position).

The structures seen in Caldwell's view include:

• Frontal sinuses (seen best)

• Ethmoid sinuses

• Maxillary sinuses

• Frontal process of the zygoma and the zygomatic process of the frontal bone

• Superior margin of the orbit and lamina papyracea

• Superior orbital fissure

• Foramen rotundum (inferolateral to superior orbital fissure)

The nose and forehead are touching the X-ray film and the X-ray beam is projected 15 degrees caudally in Caldwell's view.



Other options:

Option A: Water's view (occipitomental view or nose-chin position) is a radiographic view where an X-ray beam is angled at 45 degrees to the orbitomeatal line. Structures seen are the maxillary sinus (best seen), sphenoid sinus (if the film is taken with an open mouth), frontal sinus, zygoma, zygomatic arch, nasal bone, and infratemporal fossa.

Option C: Pierre's view is the occipitomental with an open mouth. It helps in visualizing the sphenoid sinus as well, in addition to the sinuses that can be seen in water's view.

Option D: Towne's view is an angled anteroposterior radiograph of the skull. Structures seen are the petrous part of the pyramids, arcuate eminence, superior semicircular canal, mastoid antrum, internal auditory canal, tympanic cavity, cochlea, and external auditory canal. This view is taken for acoustic neuroma and apical petrositis.

Solution to Question 8:

Battle sign refers to retro auricular or mastoid ecchymosis that typically results from a fracture of the middle cranial fossa following head trauma. It is prominent when there is a fracture of the temporal bone. Battle sign is not seen in the case of otitis media.

Option A: Bezold abscess is a complication of acute otomastoiditis where the infection/pus breaks through the thin medial side of the tip of the mastoid and presents as a swelling in the upper part of the neck.

Option B: Griesinger sign refers to edema of the postauricular soft tissues overlying the mastoid process as a result of thrombosis of the mastoid emissary vein seen in sigmoid sinus thrombosis. It is a complication of acute otomastoiditis.

Option D: Delta sign is a contrast-enhanced CT finding seen in sagittal sinus thrombosis, which can occur as a complication of acute mastoiditis or chronic suppurative otitis media. It is a triangular area with rim enhancement and the central low-density area is seen in the posterior cranial fossa on axial cuts. MR venography is useful to assess the progression or resolution of a thrombus.

The image below depicts a filling defect (thrombus) outlined by contrast medium in the sagittal sinus.



Solution to Question 9:

The marked structure is the lowermost projection in the lateral wall of the nose, the inferior turbinate. The middle turbinate (Option B) is seen as superior to it.

The lateral wall of the nose contains three or sometimes four projections of variable size namely superior, middle, and inferior turbinates along with a corresponding meatus under each turbinate.

The inferior turbinate is the largest turbinate and occupies the lower third of the lateral nasal wall. It is a separate bone. Middle and superior turbinates (Option C) are ethmoturbinal, part of the ethmoid bone.

Other important structures which could be seen in the above image are:

• IT: Inferior turbinate

• S: Septum,

• MT: middle turbinate





Uncinate process (Option D) is a hook-like structure running parallel to the anterior border of bulla ethmoidalis. The posteroinferior end of the uncinate process is attached to the inferior turbinate.

The image below shows a CT image of the uncinate process.



Solution to Question 10:

BERA, OAE, and tympanometry are the objective tests for hearing. PTA (pure tone audiometry) is a subjective test for hearing as it relies on patient responses to pure tone stimuli. Other examples of subjective tests include tuning fork tests and speech audiometry.

Tympanometry is an objective test of the middle ear and eustachian tube functions (conductive compartment of hearing). Here, the compliance of the TM, i.e., how much it moves to the pressure

differentials applied from the external ear is measured.

The image below shows the types of tympanogram:



Solution to Question 11:

Cochlear nucleus (B), superior olivary nucleus (A), lateral lemniscus (D), inferior colliculi (C), and medial geniculate ganglion (E) represent the correct sequence of the auditory pathway.

The correct sequence of structures involved in the auditory pathway from the periphery to the center is

- Vestibulocochlear nerve
- Cochlear Nucleus
- Superior olivary nucleus
- Lateral lemniscus
- Inferior colliculus
- Medial geniculate body
- Auditory cortex





Solution to Question 12:

The given clinical history of trauma to the ear along with swelling, tenderness, and discoloration of the pinna is highly suggestive of hematoma of the auricle (perichondrial hematoma). It does not resolve spontaneously.

It is a collection of blood between the auricular cartilage and its perichondrium known as perichondrial space (Option C). It is commonly seen in boxers, wrestlers, and rugby players due to blunt trauma and can result in a deformity called cauliflower ear, wrestler's ear, or pugilistic ear (Option B). Severe perichondritis may occur if the hematoma gets infected.

All cases should receive prophylactic antibiotics (Option A). Aspiration of the hematoma under strict aseptic precautions and a pressure dressing is done to prevent reaccumulation. Aspiration may need to be repeated. When aspiration fails, incision and drainage should be done.

ENT NEET 2018

Question 1:

Which of the following is an indication of high tracheostomy?

a) Vocal cord palsy

b) Tracheomalacia

c) Foreign body obstruction

d) Suspicion of Laryngeal carcinoma planned for surgery later

Question 2:

Which of the following nerves is responsible for referred otalgia from tonsillitis?

a) Glossopharyngeal nerve

b) Facial nerve

c) Trigeminal nerve

d) Vagus nerve

Question 3:

Water's view is used to best visualize which of the following sinuses?

a) Maxillary sinus

b) Ethmoidal sinus

c) Frontal sinus

d) Sphenoid sinus

Question 4:

The following instrument is used for:

a) Septoplasty

b) Myringoplasty

c) Myringotomy

d) Adenoidectomy

Question 5:

The vertical fracture of the nasal septum is known as _____.

a) Chevallet fracture

b) Citelli fracture

c) Tripod fracture

d) Jarjaway fracture

Question 6:

Where is the electrode placed in cochlear implant procedures?

a) Round window

b) Oval window

c) Scala vestibuli

d) Scala tympani

Answer Key

Question No.	Correct Option
1	d
2	a
3	a
4	c
5	a
6	d

Detailed Explanations

Solution to Question 1:

Laryngeal carcinoma is an indication of high tracheostomy. It is done at the level of the 1st tracheal ring and an opening is made above the isthmus of the thyroid gland.

The disadvantage of a high tracheostomy is subglottic stenosis at the level of the cricoid and perichondritis of the cricoid. However, in laryngeal carcinoma this would be a temporary measure as the definitive management is total laryngectomy.

Tracheostomy involves making an opening and placing a stoma in the anterior wall of the trachea.

Functions:

• Provides an alternative pathway for breathing by bypassing the obstruction.

• Improves alveolar ventilation by reducing dead space by 30-50% and reducing airway resistance

• Uses a cuff to protect the airways from secretions and blood

• Allows suction of secretions in the setting of the impaired cough reflex.

• Allows positive pressure respiration

• Allows administration of anesthesia

Indications:

• Respiratory obstruction due to

• Infections (laryngo-tracheo-bronchitis, epiglottitis)

• Trauma due to external injury, mandibular or maxillofacial fractures

• Neoplasms

• Foreign bodies or edema

• Bilateral abductor paralysis

• Congenital anomalies

- Retained secretions
- Inability to cough due to coma, spasm, or paralysis of respiratory muscles
- Aspiration of pharyngeal secretions
- Painful cough
- Respiratory insufficiency due to emphysema, chronic bronchitis

Solution to Question 2:

The tympanic branch of the glossopharyngeal nerve (cranial nerve IX) is responsible for referred otalgia from tonsillitis.

The tonsils and tonsillar fossa are supplied by the glossopharyngeal nerve. The glossopharyngeal nerve and the styloid process lie in relation to the lower part of the tonsillar fossa. Therefore, any irritation or pain can be referred to the ear along the tympanic branch of the glossopharyngeal (Jacobson's) nerve.

Causes of referred otalgia via the glossopharyngeal nerve include acute pharyngitis, tonsillitis, peritonsillar abscess, post-tonsillectomy, foreign bodies embedded in the base of the tongue or tonsillar area and an elongated styloid process (Eagle's syndrome).

The ear receives extensive sensory innervation via four cranial nerves (V, VII, IX, and X) and two spinal segments (C2 and C3).

The image below shows the anatomic relations of the tonsil and tonsillar bed:



Solution to Question 3:

Water's view is used to best visualize the maxillary sinus. Frontal and ethmoidal sinuses can also be visualized. Sphenoid sinus can be seen in Water's view with open mouth.

It is also known as the occipitomental view. The X-ray beam is angled at 37°-45° to the orbitomeatal line.

The recommended imaging views for different sinuses are as follows:

• Ethmoidal sinus: Rheese's view

• Frontal sinus: Caldwell's view

• Sphenoid sinus: Hirtz view





Solution to Question 4:

The given instrument with an arrowed blade and angled shank is a myringotome. It is used for myringotomy.

Solution to Question 5:

A Chevallet fracture usually results from a blow to the nose from below. This fracture line runs vertically. It extends from the anterior nasal spine upwards to the junction of the bony and cartilaginous dorsum of the nose.

Injuries to the nasal septum can result from trauma to the nose from the front, side, or below. The septum may get buckled on itself, fracture vertically and horizontally, or get crushed

A Jarjaway fracture (option D) results from a blow to the nose from the front. The fracture line runs horizontally backward. It begins just above the anterior nasal spine and extends just above the junction of septal cartilage with the vomer.



Early recognition and treatment of septal injuries are crucial, including drainage of hematomas.

Dislocated or fractured septal fragments should be repositioned and supported between mucoperichondrial flaps using mattress sutures and nasal packing. Nasal pyramid fractures often coincide with septal fractures and should be treated simultaneously.

Neglecting injuries to the septum can lead to deviations in the cartilaginous nose, as well as asymmetry in the nasal tip, columella, or nostrils.

Other options:

Option B: Citelli fracture is a term that does not exist. Citelli's abscess is an extratemporal complication of otitis media, occurring behind the mastoid, more towards the occipital bone. Citelli's angle refers to the sino-dural angle between the sigmoid sinus and middle cranial fossa dural plate. It is an important landmark in mastoidectomy.

Option C: A tripod fracture refers to a fracture of the zygomatic complex. There are three commonly occurring disruptions - frontozygomatic, infraorbital rim and the zygomaticomaxillary buttress.

Solution to Question 6:

The electrode of a cochlear implant is placed in the scala tympani. It allows for the electrodes to be in close proximity to the ganglion cells and their dendrites which are located in the modiolus and bony spiral lamina of the cochlea.

A cochlear implant is an electronic device for severe to profound SNHL and who cannot benefit from a hearing aid. It converts sound to an electrical signal (transduction) and delivers this to the auditory nerve. Prerequisites include an intact cochlear nerve and higher auditory pathway.

In the auditory brainstem implant, the site of the implant is the lateral recess of the fourth ventricle. It is used when the cochlear nerve is not intact.



ENT NEET 2019

Question 1:

Which of the following is not a feature of tubercular otitis media?

a) Ear ache

b) Multiple perforations

c) Pale granulation

d) Foul smelling ear discharge

Question 2:

Which among the following statements is true about keratosis obturans?

a) Failure of migration of desquamated epithelium along posterior meatal wall

b) Widening of meatus and facial nerve palsy might be seen

c) Associated bronchiectasis and sinusitis

d) All of the above

Question 3:

What is the maneuver depicted in the image given below?



a) Epley's maneuvre

b) Trotter's method

c) McGovern's technique

d) Valsalva maneuvre

Question 4:

Pott's puffy tumor is

a) Subperiosteal abscess of frontal bone

b) Subperiosteal abscess of ethmoid bone

c) Mucocele of frontal bone

d) Mucocele of ethmoid bone

Question 5:

Identify the lesion of vocal cord in the image given below :

a) Reinke's edema

b) Malignancy

c) Tracheomalacia

d) Laryngeal papilloma

Question 6:

Inspiratory stridor is found in what kind of lesions:

a) Supraglottic

b) Subglottic

c) Tracheal

d) Bronchus

Answer Key

Question No.	Correct Option
1	a
2	d
3	b
4	a
5	d
6	a

Detailed Explanations

Solution to Question 1:

Earache is not a feature of tubercular otitis media. It is a painless condition.

Tuberculosis of the ear is usually secondary to pulmonary tuberculosis. It occasionally spreads hematogenously, from a tubercular focus in the lungs, tonsils, and cervical or mesenteric lymph nodes

It presents as a painless condition with foul-smelling discharge from the ear, that does not respond to standard antibiotic treatment. Patients have severe hearing loss that is out of proportion to symptoms. Although predominantly conductive, sensorineural hearing loss may also occur if the labyrinth is involved.

A thickened tympanic membrane with multiple perforations in pars tensa of the tympanic membrane and pale granulations in the middle ear cleft is seen.

The image below shows the tympanic membrane in tuberculous otitis media:



Histopathology shows submucosal tubercles with Langhans giant cells and caseous necrosis. Smears and culture of ear discharge may be positive for acid-fast bacilli (AFB). Evidence of primary pulmonary tuberculosis may also be seen on chest radiography.

Facial nerve palsy is a common complication. Mastoiditis, postauricular fistula, and osteomyelitis are also seen.

Treatment includes aural toilet, treatment of secondary bacterial infections, and a full course of antitubercular therapy followed by reconstructive surgery.

Solution to Question 2:

All of the above statements are true regarding keratosis obturans.

Keratosis obturans occurs due to failure of migration of desquamated epithelial cells along the posterior meatal wall (Option A).

It is usually seen between the ages of 5-20 years and is commonly unilateral. Clinical features include severe otalgia and difficulty in hearing. It is associated with systemic conditions like bronchiectasis and chronic sinusitis (Option C).

On otoscopic examination, there is a thickening of the tympanic membrane and a keratin plug appearing like a pearly white mass occluding the canal. The keratin that is shed from the entire circumference of the deep ear canal appears to have an onion skin arrangement. There is also hyperemia of canal skin, sometimes with ulcerations and granulations.

Sometimes, the extent of bone erosion is such that it leaves the facial nerve exposed to injury and subsequent paralysis (Option B). Extensive erosion of the bone can also result in automastoidectomy.

Treatment involves the removal of the plug by syringing or instrumentation. Any recurrence can be prevented with the use of keratolytic such as 2% salicylic acid.

Solution to Question 3:

The given image shows Trotter's method in which the patient is made to sit, leaning a little forward over a basin to spit any blood and breathe quietly from the mouth. It is the first step to control epistaxis.

Other options:

Option A: Epley's maneuver is used in the treatment of BPPV (Benign Paroxysmal Positional Vertigo).



Option C: McGovern's technique is used in choanal atresia. A feeding nipple with a large hole provides a good oral airway.

Option D: Valsalva maneuver is performed by forceful attempted exhalation against a closed nose and mouth. It is used as a test of cardiac function and autonomic nervous control of the heart, or to clear the ears and sinuses.

Solution to Question 4:

Pott's puffy tumor is the subperiosteal abscess of frontal bone. It is a complication of frontal sinusitis.

An acute infection of the frontal sinus can cause osteomyelitis of the frontal bone. Suppurative material then breaches the frontal bone leading to subperiosteal abscess formation in the tissues of the forehead. Pus may form externally under the periosteum as a soft boggy swelling with a doughy consistency called Pott's puffy tumor. It may spread internally causing subdural, epidural, and periorbital abscesses as well as secondary septic thrombosis of the dural sinuses. It is more often seen in adults as frontal sinus is not developed in infants and children.

CT of the brain and paranasal sinuses is the most appropriate investigation. It can show the involvement of sinuses as well as intracranial or soft tissue spread of infection.

Treatment consists of large doses of antibiotics, abscess drainage, and trephining of the frontal sinus through its floor. Sometimes, the removal of sequestra and necrotic bone may also be required.

The images below show the clinical presentation and radiographic findings of the condition: red arrow - Pott's puffy tumor; blue arrow - intracranial spread

Solution to Question 5:

The image showing grape-like projections, involving the anterior part of the vocal cords depicts a laryngeal papilloma.

Laryngeal papilloma is a benign tumor of the larynx caused by the human papillomavirus (HPV type 6 and 11). They can occur in children, most commonly around 3 -5 years of age (juvenile papillomatosis), as well as in adults. They involve the supraglottic and glottic regions of the

larynx. Seeding of the lesions can occur during invasive laryngeal procedures, such as intubation. They have a high recurrence rate and very low potential for malignancy.

Adult-onset laryngeal papillomatosis is less common and usually seen in males aged 30-50 years. It is less aggressive and involves the anterior part of the vocal cords. The treatment is the same as juvenile papillomatosis.

Patients present with hoarseness, aphonia, dyspnea, or stridor. Laryngoscopy shows pinkish to white grape-like projections that can be sessile or pedunculate with a visible central vascular core. The diagnosis is confirmed by direct laryngoscopy and biopsy. Treatment consists of micro-laryngoscopy and CO_2 laser excision.

The image below depicts laryngeal papillomatosis on laryngoscopy:



Reinke's edema (Option A) is a collection of fluid in the subepithelial space of Reinke. It generally occurs in patients with vocal abuse or smoking. The vocal cords will show diffuse, symmetrical swelling.

The laryngoscopic image below shows Reinke's edema:



Solution to Question 6:

Inspiratory stridor is seen in supraglottic lesions. It is often produced in obstructive lesions of supraglottis or pharynx, e.g. laryngomalacia or retropharyngeal abscess.

Expiratory stridor is produced in lesions of thoracic trachea, primary and secondary bronchi such as bronchial foreign body, and tracheal stenosis.

Biphasic stridor is seen in lesions of glottis, subglottis, and cervical trachea like laryngeal papillomas, vocal cord paralysis, and subglottic stenosis.

Stridor is an audible respiratory noise that gets produced by turbulent airflow through an obstructed upper airway. It is an inspiratory sound that may have an expiratory component. It is distinct from stertor which is a snoring-like noise produced from an obstructed nasopharynx or oropharynx.

The intensity and pitch of sounds originating from the respiratory system vary based on their site of origin within the respiratory tract and may be heard with or without a stethoscope. Usually, the pitch of the sound increases, and the intensity decreases as one goes lower into the respiratory tract.

Types of stridor and their site of origin



©MARROW

ENT NEET 2020

Question 1:

From which of the following structure does the saccule develop?

a) Saculus anterior

b) Saculus posterior

c) Pars superior

d) Pars inferior

Question 2:

What is the surgery done to widen the cartilaginous part of the external auditory canal called?

a) Meatoplasty

b) Tympanoplasty

c) Myringoplasty

d) Otoplasty

Question 3:

Cough on scratching the external acoustic canal is due to?

a) Auriculotemporal nerve

b) Auricular branch of vagus

c) Great auricular nerve

d) Facial nerve

Question 4:

Which of the following is not a feature of tubercular otitis media?

a) Ear ache

b) Multiple perforations

c) Pale granulation

d) Foul smelling ear discharge

Question 5:

A person who met with an accident, and suffered from a skull fracture presents with the following finding. Identify this clinical finding.

a) Battle sign

b) Bezold abscess

c) Mastoiditis

d) Griesinger sign

Question 6:

Surgery where nostrils are partially or completely occluded is done for which condition?

a) Vasomotor rhinitis

b) Atrophic rhinitis

c) Invasive aspergillosis

d) Allergic rhinitis

Question 7:

What is occipitomental view with open mouth also known as?

a) Water's view

b) Towne's view

c) Law's view

d) Stenver's view

Question 8:

A 9-year-old boy comes with complaints of right ear pain, difficulty in opening the mouth, painful swallowing, and fever. The oral cavity examination reveals the following. The external facial examination is unremarkable. Which of the following is likely?



a) Pharyngitis

b) Parotid abscess

c) Bezold's Abscess

d) Quinsy

Answer Key

Question No.	Correct Option
1	d
2	a
3	b

4	a
5	a
6	b
7	a
8	d

Detailed Explanations

Solution to Question 1:

The saccule develops from pars inferior.

The development of the inner ear begins in the 3rd week of fetal life and is completed by the 16th week. The surface ectoderm on each side of the rhombencephalon thickens to form otic/auditory placodes, which then invaginate to form the otic vesicle (otocysts).

The cells of the otic vesicle differentiate to form ganglion cells, which later develop into the vestibulocochlear ganglia. Later each otic vesicle divides into a dorsal component called pars superior and a ventral component called pars inferior.

Pars superior gives rise to the utricle, semicircular canals, and endolymphatic ducts. Pars inferior gives rise to saccule and cochlear duct. The saccule is connected to the cochlea through a narrow duct called ductus reuniens.

Solution to Question 2:

The surgery done to widen the cartilaginous part of the external auditory canal is meatoplasty.

Meatoplasty is a surgical procedure that involves the removal of a crescent-shaped portion of conchal cartilage to widen the external auditory meatus. It is performed in combination with all canal wall-down procedures like modified radical and radical mastoidectomies, for easy access to the mastoid cavity for periodic inspection and cleaning.

It is also done as an isolated procedure in elderly people with sagging auricles. A sagging auricle causes obstruction to the meatus leading to the retention of wax and difficulty in hearing.

Other options:

Option B: Tympanoplasty is the surgical reconstruction to correct damage in the middle ear and restore the integrity of the tympanic membrane for improved hearing. It is ossicular reconstruction only or ossicular reconstruction with myringoplasty.

Option C: Myringoplasty refers to the reconstruction of the tympanic membrane only.

Option D: Otoplasty is the correction of external ear/deformities of the pinna.

Solution to Question 3:

Cough on scratching the external acoustic canal is due to the auricular branch of the vagus nerve (Arnold's nerve).

Arnold's nerve supplies the concha and corresponding eminence on the medial surface. It also supplies the posterior wall and floor of the external auditory canal. Scratching or mechanical stimulation of the ear leads to activation of Arnold's nerve, which evokes reflex cough (ear-cough reflex/otorespiratory reflex).

The image below shows the nerve supply of the pinna:

Causesofreferred otalgia	Nerveresponsible	Site
Temporomandibular joint and dental conditions, parotid infections/tumor, and anterior 2/3rd of tongue	Auriculotemporal nerve	Tragal area, crus of the helix, and anterior wall and roof of the external auditory canal.
Acute tonsillitis, peritonsillar abscess, carcinoma base of the tongue or tonsils (oropharynx)	The glossopharyngeal nerve (Jacobson's)	Medial surface of the tympanic membrane, and middle ear
Carcinoma of the larynx (cough), hypopharynx, epiglottis, vallecula, esophagus, and thyroid	Auricular branch of the vagus nerve (Arnold's/Alderman's nerve)	Concha, posterior wall, and floor of the external auditory canal
Cervical spondylitis	Great auricular nerve (C2,3)	Lateral and medial surfaces of the pinna
Acoustic neuroma	Facial nerve (nervus intermedius/nerve of Wrisberg)	Concha, retro auricular groove, and posterosuperior part of the external auditory meatus. Numbness in the posterosuperior meatal wall known as Hitzelberger's sign, is seen in acoustic neuroma.



Solution to Question 4:

Earache is not a feature of tubercular otitis media. It is a painless condition.

Tuberculosis of the ear is usually secondary to pulmonary tuberculosis. It occasionally spreads hematogenously, from a tubercular focus in the lungs, tonsils, and cervical or mesenteric lymph nodes

It presents as a painless condition with foul-smelling discharge from the ear, that does not respond to standard antibiotic treatment. Patients have severe hearing loss that is out of proportion to symptoms. Although predominantly conductive, sensorineural hearing loss may also occur if the labyrinth is involved.

A thickened tympanic membrane with multiple perforations in pars tensa of the tympanic membrane and pale granulations in the middle ear cleft is seen.

The image below shows the tympanic membrane in tuberculous otitis media:



Histopathology shows submucosal tubercles with Langhans giant cells and caseous necrosis. Smears and culture of ear discharge may be positive for acid-fast bacilli (AFB). Evidence of primary pulmonary tuberculosis may also be seen on chest radiography.

Facial nerve palsy is a common complication. Mastoiditis, postauricular fistula, and osteomyelitis are also seen.

Treatment includes aural toilet, treatment of secondary bacterial infections, and a full course of antitubercular therapy followed by reconstructive surgery.

Solution to Question 5:

The given image shows ecchymosis over the retro auricular and mastoid area suggestive of Battle's sign. It is typically seen in fractures of the middle cranial fossa following head trauma. Battle sign is prominent when there is a fracture of the petrous temporal bone.

Other options:

Option B: Bezold abscess is a complication of acute otomastoiditis where the infection/pus breaks through the thin medial side of the tip of the mastoid and presents as a swelling in the upper part of the neck.



Option C: Acute mastoiditis is used to refer to an acute infection that spreads from the mucosa lining the mastoid air cells to involve the bony walls of the mastoid air cell system. Mastoiditis presents as retro-auricular redness and swelling which is painful.

The image below shows a smooth ironed-out mastoid with erythema seen in acute mastoiditis.

Option D: The Griesinger sign refers to edema of the postauricular soft tissues overlying the mastoid process as a result of thrombosis of the mastoid emissary vein seen in sagittal sinus thrombosis. It is a complication of acute otomastoiditis.

Solution to Question 6:

Complete closure of the nostrils (Young's operation) or partial closure of the nostrils (Modified Young's operation) is done for the treatment of atrophic rhinitis.

In Young's operation, both nostrils are closed completely by raising flaps within the nasal vestibule. They are opened after 6 months. During this time, the mucosa becomes normal, and the crusting is reduced.

In Modified Young's operation, the procedure has been modified to avoid discomfort from complete closure of the nostrils. Here the nostrils are only partially closed. The results are the same when compared to Young's operation.

Atrophic rhinitis (ozaena) is a chronic inflammation of the nose with atrophy of nasal mucosa and turbinate bones. Roomy nasal cavities with foul-smelling crusts are seen. The patient presents with a foul smell from the nose, marked anosmia (merciful anosmia), and paradoxical nasal obstruction (nasal obstruction despite wide nasal passages). Paradoxical nasal obstruction occurs due to degenerative changes in the sensory nerves of the nasal mucosa, leading to an inability to sense airflow.

Medical management includes nasal irrigation and removal of crusts. 25% glucose in glycerine, antibiotics (quinolones or metronidazole), estradiol nasal spray, and mucolytics can be used.

Solution to Question 7:

Occipitomental view is called Water's view. Water's view with an open mouth is known as Pierre's view. Among the given options, Water's view is the best answer.

In Water's view, the patient's head is positioned such that the nose and the chin touch the film while the X-rays are projected from backward. The mouth is usually kept open, which allows visualizing the sphenoid sinus.

Structures seen in Water's view are:

• Maxillary sinuses (seen best)

• Frontal sinuses

• Sphenoid sinus (if the film is taken with an open mouth)

• Zygoma

• Zygomatic arch

• Nasal bone

• Frontal process of the maxilla

• Superior orbital fissure

• Infratemporal fossa

Other options:

Option B: Towne's view is an angled anteroposterior radiograph of the skull. Structures seen are the petrous part of the pyramids, arcuate eminence, superior semicircular canal, mastoid antrum, internal auditory canal, tympanic cavity, cochlea, and external auditory canal. This view is taken for acoustic neuroma and apical petrositis.

Option C: Law's view is a lateral oblique X-ray view of the mastoid. Structures seen are the external auditory canal, mastoid air cells, tegmen, lateral sinus plate, and temporomandibular

joint.

Option D: Stenver's view is an X-ray view of the skull where the long axis of the petrous bone is parallel to the film. Structures seen are the entire petrous pyramid, arcuate eminence, internal auditory meatus, labyrinth with its vestibule, cochlea, and mastoid antrum.

Solution to Question 8:

The given clinical scenario describes a child with fever, otalgia, trismus, and odynophagia with a normal external facial examination. The oral cavity shows congested and edematous tonsil and tonsillar pillars with edematous uvula pushed to the opposite side. These findings are highly suggestive of quinsy (peritonsillar abscess).

It is the collection of pus in the peritonsillar space between the fibrous capsule of the tonsil and the superior constrictor muscles of the pharynx. Both aerobic and anaerobic organisms are seen.

Patient presents with fever, unilateral throat pain, odynophagia, hot potato voice, torticollis, foul breath, referred otalgia through glossopharyngeal nerve and trismus due to spasm of pterygoid muscles.

Oral cavity examination may show congested and edematous tonsils, pillars, and soft palate on the involved side. Edematous uvula pushed to the opposite side. Pus may be seen on the tonsillar region. Jugulodigastric lymph nodes may be enlarged.

The management of quinsy is by antibiotics and incision and drainage to drain the abscess followed by interval tonsillectomy 4 to 6 weeks later.

Rare complications seen are parapharyngeal abscess, edema of the larynx, septicemia, lung abscess, or jugular vein thrombosis.

Other options:

Option A: Pharyngitis presents with fever, throat pain, enlarged tonsils, and lymphoid follicles on the posterior pharyngeal wall.

Option B: Parotid abscess occurs due to the suppuration of parotid space. It presents with swelling, redness, firmness, and tenderness in the parotid area and at the angle of the mandible. It is usually unilateral, but bilateral abscesses can occur. Due to the thick capsule, fluctuation is often difficult to elicit.

Option C: Bezold abscess is a complication of acute otomastoiditis where the infection/pus breaks through the thin medial side of the tip of the mastoid and presents as a swelling in the upper part of the neck.

ENT INICET 2021

Question 1:

Identify the condition shown in the image.

a) Adenoid facies

b) Goldenhar syndrome

c) Horse facies

d) Frog facies

Question 2:

The given image is of a person with a history of minor trauma with a piece of wood, 20 days back. His face is affected on the left side with facial swelling and orbital edema. CT scan revealed clear sinuses and the presence of subcutaneous nodules. Microscopy of the tissue sections was PAS-positive and stained positive with Grocott's methenamine silver stain. What is the most likely diagnosis?

a) Phycomycosis

b) Midline lethal granuloma

c) Foreign body granuloma

d) IgG4 granuloma

Question 3:

Which of the following are the objective tests for the hearing?

a) 1, 2, and 4

b) 1,2, and 3

c) 2,3, and 4

d) 1,2,3, and 4

Question 4:

A young man presented with hearing loss and tinnitus. Histology image is shown below. What is the diagnosis?



a) Neurofibroma

b) Schwannoma

c) Leiomyoma

d) Rhabdomyoma

Question 5:

All of the following are major diagnostic criteria for allergic fungal sinusitis except?

a) Presence of nasal polyps

b) Eosinophilic mucin without invasion

c) Characteristic CT findings

d) Positive fungal culture

Question 6:

Which among the following is not used in post laryngectomy rehabilitation?

a) Tracheo-esophageal prosthesis

b) Esophageal speech

c) Polite yawning

d) Super supraglottic swallow

Question 7:

Which of the following devices can be used in a patient with bilateral external auditory canal atresia?



a) 1

b) 2

c) 3

d) 4

Question 8:

A 65-year-old patient presents with right-sided nasal obstruction and occasional blood-stained nasal discharge. His CT and endoscopy findings are shown in the below image. What could be the underlying cause?



a) Inverted papilloma

b) Carcinoma maxilla

c) Esthesioneuroblastoma

d) Nasopharyngeal angiofibroma

Question 9:

A patient presents with vertigo, tinnitus, and head tilt. He underwent myringoplasty for the safe type of chronic suppurative otitis media(CSOM) 6 months back. What is your diagnosis?

a) Peri-lymphatic fistula

b) Labyrinthitis

c) Paget disease

d) Vestibular schwannoma

Question 10:

What is the probable diagnosis based on the finding given in the CT image?

a) Concha bullosa

b) Pneumatised superior turbinate

c) Onodi cell

d) Haller cell

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	b
5	d
6	d
7	a
8	a
9	a
10	b

Detailed Explanations

Solution to Question 1:

The given image shows adenoid facies. It is characterized by an elongated face with a dull expression, an open mouth, a hitched-up upper lip, prominent upper teeth causing crowding (malocclusion of teeth), a high arched hard palate (due to loss of tongue's molding action on the palate), and a pinched-in nose due to disuse atrophy of the alae nasi.



Chronic nasal obstruction resulting in mouth breathing leads to this characteristic facial appearance in children.

Adenoids (nasopharyngeal tonsils) are prone to physiological enlargement during childhood. Affected children present with symptoms of nasal obstruction, nasal discharge, sinusitis, and rhinolalia clausa (loss of nasal tone of voice). They are prone to eustachian tube obstruction, recurrent acute otitis media, chronic otitis media, and otitis media with effusion. They may also experience aprosexia (inability to concentrate) leading to declining scholastic performance. Pulmonary hypertension and cor pulmonale are potential long term complications

Other Options:

Option B: Goldenhar Syndrome, also known as fascio-auriculo-vertebral dysplasia or oculo-auriculo-vertebral (OAV) syndrome, is characterized by facial asymmetry, low set ears, atresia of the ear canal, cardiac abnormalities, pre-auricular tags/pits, hemivertebrae in the cervical region, epibulbar dermoid, coloboma of the upper lid, and mixed or conductive hearing loss.



©MARROW



©MARROW

Option D: Frog facies is a characteristic presentation observed in cases of juvenile nasopharyngeal angiofibroma. It is identified by the swelling of the cheeks, proptosis (protrusion of the eyeballs), and broadening of the nasal bridge.

Solution to Question 2:

The history of facial swelling and orbital edema following trauma with wood and positive fungal stain (Grocott-Gomori's methenamine silver) points toward phycomycosis (mucormycosis), a potentially invasive fungal infection.

Mucormycosis is an opportunistic mycosis caused by various molds in the Mucorales order which are ubiquitous thermotolerant saprobes. These molds, including Rhizopus, Mucor, and Absidia, are common in the environment. The spores can be found in air and dust.

Mucormycosis can manifest as a localized cutaneous or subcutaneous infection in immunologically normal individuals after traumatic exposure to soil or vegetation. It is a fatal condition due to its angio-invasive nature. Lung involvement is common, and the infection primarily affects immunocompromised patients.

Rhino-orbital-cerebral mucormycosis is frequently seen in individuals with diabetic ketoacidosis. It typically starts with eye and facial pain and may progress to orbital cellulitis, proptosis, and vision loss. A characteristic finding is a black necrotic mass filling the nasal cavity.

Diagnosis involves testing specimens and cultures. In exudate samples, wet preparations with 10% potassium hydroxide (KOH) reveal broad, ribbon-like hyphae. Tissue biopsy confirms the diagnosis by observing non-septate hyphae branching at right angles, characteristic of Mucorales like Rhizopus delemar.



Culture isolation is performed on Sabouraud agar without cycloheximide, resulting in grey-white colonies with a thick cottony, and fluffy surface.

Treatment is primarily with liposomal amphotericin B. Surgical debridement of the affected tissues and control of underlying predisposing cause is equally important.

Other options:

Option D: IgG4-related disease (IgG4-RD) is characterized by tissue infiltrates dominated by IgG4 antibodies, which is the hallmark of the disease. IgG4-RD leads to fibrosis and obliterative phlebitis. Patients may present with Riedel's thyroiditis, autoimmune pancreatitis, or idiopathic orbital inflammation, but not with ptosis. Diagnosis is confirmed through positive fibrosis and the presence of plasma cells on biopsy. Treatment with Rituximab is done.

Option C: Foreign body granulomas are due to inflammation induced by inert foreign bodies. These granulomas develop without T-cell-mediated immune responses. Sutures are a common site for their formation. The foreign material is at the center of the granuloma, surrounded by epithelioid cells and specific giant cells opposing the surface.

Option B: Midline lethal granuloma is characterized by necrotizing destruction and damage to the nose and other respiratory structures. It is rare and usually a diagnosis of exclusion. CT scans can detect the lesions associated with this condition. Negative c-ANCA (cytoplasmic antineutrophil

cytoplasmic antibodies) and fungal stain results contribute to the diagnosis.

Solution to Question 3:

BERA, OAE, and tympanometry are the objective tests for hearing. PTA (pure tone audiometry) is a subjective test for hearing as it relies on patient responses to pure tone stimuli. Other examples of subjective tests include tuning fork tests and speech audiometry.

Tympanometry is an objective test of the middle ear and eustachian tube functions (conductive compartment of hearing). Here, the compliance of the TM, i.e., how much it moves to the pressure differentials applied from the external ear is measured.

The image below shows the types of tympanogram:



Solution to Question 4:

The history of hearing loss and tinnitus and the histology showing Verocay bodies are suggestive of Schwannoma.

Vestibular Schwannoma (acoustic neuroma), is a benign tumor that predominantly originates from the Schwann cells in the vestibular division of the eighth cranial nerve, but it rarely arises from the cochlear division. It commonly affects individuals aged 40-60, with no gender preference. Bilateral schwannomas occur in neurofibromatosis type 2 (NF2).

The most common presenting manifestations are progressive, unilateral sensorineural hearing loss, often accompanied by tinnitus.

It can involve all cranial nerves except the first and second. Apart from the 8th cranial nerve, the earliest cranial nerve to be involved is the 5th cranial nerve leading to reduced corneal sensitivity and paraesthesia of the face. This indicates that the tumor is roughly 2.5 cm in diameter and occupies the cerebellopontine angle.

Facial nerve involvement causes hypoaesthesia of the posterior meatal wall (Hitzelberger sign), taste loss, and delayed blink reflex. The earliest sign is a delayed blink reflex.

Glossopharyngeal (IX) and vagus (X) nerve involvement result in dysphagia and hoarseness. Brainstem involvement manifests as ataxia, limb weakness, numbness, and increased tendon reflexes. Cerebellar involvement leads to an unsteady gait and dysdiadochokinesia.

Histopathologically, the tumor comprises dense Antoni A areas with spindle cells in a palisading pattern and Antoni B areas with loosely arranged spindle cells in a myxoid matrix. Verocay bodies are formed in the Antoni A region, characterized by nuclear-free zones surrounded by palisaded nuclei.

Pure tone audiometry reveals sensorineural hearing loss, particularly in high frequencies. Speech audiometry shows the roll-over phenomenon. The absence of the recruitment phenomenon is noted. Short Increment Sensitivity Index (SISI) ranges from 0-20%.

Brainstem Evoked Response Audiometry (BERA) demonstrates a significant delay of >0.2 ms in wave V between the two ears. Magnetic Resonance Imaging (MRI) with gadolinium contrast is considered the gold standard for diagnosis.

The primary management is the surgical removal of the tumor via the middle cranial fossa approach, the trans-labyrinthine approach, the suboccipital (retro sigmoid) approach, or a combined trans-labyrinthine-suboccipital approach.

Conventional radiotherapy using external beam radiation is not effective. In cases where surgery is contraindicated, refused by the patient, or when a residual tumor remains, alternative treatments such as the x-knife or gamma knife can be considered.

Other options:

Option A: Neurofibromas are also benign nerve sheath tumours and the histological picture shows the presence of bland Schwann cells admixed with stromal cells such as mast cells, perineural cells, and fibroblasts. The stroma of the tumour contains loose collagen arranged in wavy bundles resembling carrot shavings.



Option C: Leiomyomas are benign smooth muscle neoplasms that may occur single or multiple. Microscopically, leiomyomas are comprised of well-differentiated, regular, spindle-shaped smooth muscle cells associated with hyalinization. Very less mitotic figures are seen.



Option D: Rhabdomyomas are the most frequent primary tumours of the heart in children. Microscopically, they are composed of bizarre, markedly enlarged myocytes with abundant glycogen. But due to routine histological processing, the glycogen content is reduced and looks like thin strands that stretch from the nuclei to the surface membrane, which appears like spider cells.



Solution to Question 5:

A positive fungal culture is a minor and not a major diagnostic criterion for allergic fungal rhinosinusitis (AFRS).

The Bent and Kuhn diagnostic criteria for allergic fungal rhinosinusitis include:

Major criteria:

• Type I hypersensitivity

• Nasal polyposis

• Characteristic CT findings

• Eosinophilic mucin without invasion

• Positive fungal stain

Minor criteria:

• Asthma

• Unilateral disease

• Bone erosion

• Fungal cultures

• Charcot-Leyden crystals

• Serum eosinophilia

All the major criteria must be met by the patient while the minor criteria serve to support the diagnosis and describe individual patients.

Characteristic CT findings in AFRS include heterogeneous signal intensities within the paranasal sinuses filled with allergic mucin (double density sign), expansion of the paranasal sinuses/nasal cavity, unilateral or asymmetric disease load, and bony erosion.

The treatment of allergic fungal sinusitis includes avoidance of allergen exposure, antihistamines, corticosteroid nasal sprays, aggressive surgical intervention for mucin removal which improves sinus drainage, post-surgery oral corticosteroids, and immunotherapy against fungal and non-fungal allergens.

Solution to Question 6:

Super supraglottic swallow is not used in post-laryngectomy rehabilitation.

Super-supraglottic swallow is a maneuver that prevents the aspiration of food or liquid by closing the airway before swallowing. It is used in the management of dysphagia. This method involves holding one's breath to close the airway prior to swallowing, followed by coughing voluntarily immediately after swallowing to clear any residual food or liquid from the airway entrance. It is an effortful breath-holding technique, which not only results in Valsalva and subsequent leakage of bolus but can also result in arrhythmias.

Other options:

Option A: A tracheoesophageal prosthesis(TEP) involves a one-way valve that is inserted through a previously-created tracheoesophageal puncture. When the patient occludes the stoma, the exhaled air is shunted through the TEP into the oesophagus, where it induces vibration of the upper oesophagal sphincter, which is responsible for phonation. It generates the most intelligible,

fluent, natural-sounding voice in contrast to other laryngeal speech methods.



Option B: Oesophagal speech also involves vibration of the upper oesophagal sphincter and subsequent phonation.



Option C: Polite yawning also called the nasal airflow-inducing maneuver is an important olfaction rehabilitation tool after a laryngectomy.

Electrolarynx produces vibrations that are transmitted through the skin toward the throat. The sound generated is converted into low-pitch speech by the vocal tract—palate, tongue, and teeth. The advantages are that is easy and quick to learn, while the disadvantages are that the voice is monotonous and the constant occupation of one hand.



Solution to Question 7:

Device 1 shown in the above image represents bone anchored hearing aid (BAHA), which can be used in a patient with bilateral external auditory canal atresia. Cochlear implants (Device 2) are primarily used to restore sensorineural hearing loss, whereas BAHA is used in cases with defects in the outer or middle ear.

External auditory canal atresia refers to the absence of a patent ear canal and this can either be congenital or acquired.

Congenital atresia: The external ear develops from the first and second pharyngeal arch and the first pharyngeal cleft. Atresia of the external canal can occur in association with microtia (under-developed pinna) or with abnormalities of the middle ear, internal ear, and other structures. It is usually bilateral.

Acquired atresia: A rare condition, following external ear trauma, such as road traffic accidents, gunshot wounds, or recent otologic surgery. In rare instances, canal stenosis and atresia have been observed in association with neoplastic changes and idiopathic inflammatory processes.

Bilateral aural atresia will present with failed audiological testing and require early bone conduction hearing aids, one of which is BAHA, that combines a sound processor with a small titanium fixture implanted behind the ear and allows the bone to transfer sound to a functioning cochlea rather than via the middle ear – a process known as direct bone conduction.

A picture of BAHA is shown below. It consists of three parts, an innermost titanium fixture in the bone, a middle abutment, and an outer hearing aid fixed to the abutment.

Bone-anchored Hearing Aids (BAHA) are used:

• When air-conduction (AC) hearing aid cannot be used:

• People who have chronic inflammation or infection of the ear canal associated with discharge and cannot wear standard “in the ear” air-conduction hearing aids.

• Children with malformed or absent outer ear and ear canals as in microtia or canal atresia.

• In conductive or mixed hearing loss, e.g. otosclerosis and tympanosclerosis where surgery is contraindicated

• In unilateral deafness

Other options:

Option B: Device 2 is a picture of a cochlear implant. It is a medical device that uses electricity to stimulate the spiral ganglion cells of the auditory nerve to restore sensorineural hearing loss. It converts sound to an electrical signal (transduction) and delivers this to the hearing nerve, which bypasses the damaged hearing apparatus. It cannot be used in patients with cochlear aplasia or without the VIIIth cranial nerve.

Option C: Device 3 is a picture of a stapedectomy piston, used in patients with otosclerosis.

Option D: Device 4 is a picture of a partial ossicular replacement prosthesis that is designed for situations when the incus and/or malleus are eroded but the stapes is structurally intact.



Solution to Question 8:

In an elderly patient, unilateral nasal obstruction, epistaxis, and pale polypoidal mass arising from the lateral wall of the nose are suggestive of inverted papilloma (Ringertz tumor).

Inverted papilloma (transitional cell papilloma) is a tumor originating from the nonolfactory mucosa of the nose (Schneiderian membrane) and paranasal sinuses. The most common site of origin is the lateral wall of the nose in the middle meatus.The unilateral growth of hyperplastic papillomatous tissue into the stroma characterizes this condition. Human papillomavirus (HPV) is believed to be the cause.

It primarily affects men aged 40-70 and presents unilaterally with symptoms such as nasal obstruction, nasal discharge, and epistaxis. Invasive growth into the sinuses or orbit can lead to proptosis, diplopia, and lacrimation.

On nasal endoscopy, it appears as a pale polypoidal mass resembling a nasal polyp. Computed tomography (CT) and magnetic resonance imaging (MRI) are used to determine the location and extent of the lesion. MRI can also help differentiate between associated sinus secretions and the tumor mass. Biopsy is essential for diagnosis, as it is important to differentiate it from simple nasal polyps, which may be associated with or mistaken for the tumor.

The treatment of choice is medial maxillectomy, which can be performed through lateral rhinotomy, sublabial degloving approach, or preferably, an endoscopic approach. In 10%-15% of cases, the tumor is associated with malignancy. Recurrence is possible, and radiotherapy is generally not recommended due to the potential risk of inducing malignancy.

Other options:

Option B: Carcinoma maxilla arises from the lining of the maxillary sinus and its early features include additional symptoms like facial paraesthesias, pain, or epiphora and depending on the direction of spread of the growth, would cause additional symptoms.

Option C: Esthesioneuroblastoma (olfactory neuroblastoma) arises from the olfactory epithelium in the upper third of the nose. It has bimodal peaks of incidence at 10–20 years and 50–60 years. Unilateral nasal obstruction and epistaxis are the most common symptoms. The tumour also invades the orbit and the surrounding structures, which results in other symptoms like proptosis, headache, epiphora, diplopia, and blurred vision.

Option D: Juvenile nasopharyngeal angiofibroma is seen almost exclusively in males in the age group of 10–20 years and is very rarely seen in older people and females. Profuse, recurrent, and spontaneous epistaxis is the most common presentation.

Solution to Question 9:

The presentation of vertigo, tinnitus, and head tilt following myringoplasty suggests a diagnosis of perilymphatic fistula.

A perilymphatic fistula is an abnormal connection between the inner ear and middle ear, resulting in leakage of perilymph from the cochlea or vestibule. This most commonly occurs through the round or oval window.

Causes of perilymphatic fistula include post-stapedectomy, ear surgery with dislocation of the stapes, head trauma, inner ear injury due to forceful Valsalva or barotrauma, raised intracranial pressure, and occasionally spontaneous or idiopathic.

Clinically, intermittent vertigo, fluctuating sensorineural hearing loss, tinnitus, and a sense of fullness in the affected ear are present. The fistula test is positive, and high-resolution computed tomography (HRCT) of the temporal bone is highly sensitive for detecting the fistula. A combination of CT and magnetic resonance imaging (MRI) can diagnose almost all cases. Audiometry and electrocochleography may be performed additionally.

Management of perilymphatic fistula can be conservative or surgical. Conservative measures involve avoiding activities that increase inner ear or intracranial pressure and using intra-tympanic steroids for acute decompensation.

Other options:

Option B: Labyrinthitis presents with unilateral vestibular symptoms like sudden onset of persistent vertigo with associated symptoms such as nausea, vomiting, sweating, and pallor. Symptoms may progress resulting in nystagmus and hearing loss. Tinnitus is not a common finding in labyrinthitis.

Option C: Paget's disease of bone usually presents with hearing loss as a prominent feature, when the skull is involved. It is not associated with a history of surgery, trauma, or vestibulocochlear symptoms.

Option D: Vestibular Schwannoma (acoustic neuroma), originates from Schwann cells in the vestibular division of the vestibulocochlear (VIIIth cranial) nerve within the internal auditory canal. The main symptoms are progressive unilateral hearing loss and tinnitus. Other cranial nerves (V, IX, X, XI) may also be involved.

Solution to Question 10:

The given CT image reveals a pneumatised superior turbinate.

The superior turbinate, located posteriorly and superiorly to the middle turbinate can become pneumatized. It forms an important landmark to identify the ostium of the sphenoid sinus which lies medial to it. This pneumatization is common in the lateral nasal wall, although it is less frequently seen in the superior turbinate compared to the middle turbinate.

In most cases, pneumatized superior turbinate is an incidental finding during radiological examinations. It can lead to nasal obstruction, headache, epistaxis, anosmia, and postnasal drainage.

Asymptomatic cases of superior turbinate pneumatization generally do not require treatment. However, when symptoms are present, various surgical techniques can be considered, including outfracture, crushing, radio-frequency ablation, turbinoplasty, and microdebrider surgery.

Turbinoplasty is contraindicated if inferior turbinate pneumatization is associated with the maxillary sinus because it may result in the formation of a second ostium. In addition, total turbinectomy is not performed because it may cause atrophic rhinitis.

Other options:

Option A: Pneumatization of the middle turbinate leads to an enlarged middle turbinate that is ballooned out, this is called concha bullosa. The CT image shown below depicts concha bullosa.

Option C: Onodi cell is a posterior ethmoidal cell that may grow posteriorly, either by the side of the sphenoid sinus or superior to it. Onodi cell is surgically important as the optic nerve may be related to its lateral wall. A CT image of an onodi cell is shown below.



Option D: Haller cells are air cells situated in the roof of the maxillary sinus. They are pneumatized from anterior or posterior ethmoid cells. Enlargement of Haller cells may lead to maxillary sinusitis as it may encroach the ethmoid infundibulum, impeding draining of the maxillary sinus. A CT image of Haller cell is shown below.



ENT NEET 2021

Question 1:

The image of a drug is given below. It is used in which of the following conditions?

a) Subglottic stenosis

b) Rhinocerebral mucormycosis

c) Adenoidectomy

d) Tympanoplasty

Question 2:

Following total thyroidectomy, a patient started having difficulty in breathing, and repeated attempts to extubate were unsuccessful. The most probable cause is ___

a) Superior laryngeal nerve injury

b) Unilateral recurrent laryngeal nerve injury

c) Bilateral recurrent laryngeal nerve injury

d) Hematoma

Question 3:

A female patient presents with nasal obstruction, nasal discharge, and loss of smell. On examination, foul-smelling discharge and yellowish-green crusts are present in the nasal cavity. She is found to have merciful anosmia. Which of the following finding can also be seen during the examination of her nose?

a) Roomy nasal cavity

b) Nasal polyps

c) Inferior turbinate hypertrophy

d) Foreign body

Question 4:

A patient presents to the emergency with epistaxis. There was no relief on pinching the nostrils. Nasal packing was done but the patient still continues to bleed. What would be the next appropriate step in the management of this patient?

a) Ligation of external carotid artery

b) Ligation of internal carotid artery

c) Ligation of sphenopalatine artery

d) Ligation of maxillary artery

Question 5:

A patient post-tonsillectomy in the recovery room starts bleeding from the operative site. On examination, blood clots are seen. What will be your immediate management?

a) Shift to OT, remove the clots, and cauterize/ligate the vessel

b) Shift to OT, start IV antibiotics, and pack the tonsillar fossa

c) Give anticoagulants, repeated gargling, and wait for 24 hours

d) Do blood transfusion and wait and watch

Question 6:

A patient with a history of chronic ear infection presents with fever, headache, vomiting, irritability, and confusion. His CT brain is shown in the image below. What is the possible diagnosis?



a) Temporal lobe abscess

b) Cerebellar abscess

c) Subdural abscess

d) Meningitis

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	c
5	a
6	a

Detailed Explanations

Solution to Question 1:

The given drug is Mitomycin-C. It is used topically in the management of subglottic stenosis.

Mitomycin-C is derived from the Streptomyces caespitosus bacteria. It can modify wound healing at a molecular level by inhibiting DNA synthesis by cross-linking DNA, acting like an alkylating

agent (cell cycle non-specific). It has been used to inhibit post-surgical scar formation and as a chemotherapeutic agent for solid tumors.

Mitomycin-C is employed in the treatment of superficial bladder tumors, postoperative glottic and subglottic stenosis, and tracheal scarring after tracheal reconstruction surgery.

Subglottic stenosis is the narrowing of the airway in the subglottis, clinically presenting with biphasic stridor. Diagnosis is made if the subglottic diameter is <4mm in full-term infants and if the subglottic diameter is <3mm, in the case of premature infants.

Myer-Cotton grading of subglottic stenosis is as follows:

Grade	Obstruction
Grade I	≤50%
Grade II	51-70%
Grade III	71-99%
Grade IV	No detectable lumen

Solution to Question 2:

The clinical picture of breathing difficulty and unsuccessful attempts at extubation after total thyroidectomy is suggestive of bilateral recurrent laryngeal nerve injury.

Thyroid surgery may cause injury to both external (ELN) and recurrent laryngeal nerves (RLN). The upper part of the RLN is closely related to the inferior thyroid artery. Unilateral RLN injury may be asymptomatic but in a few may cause hoarseness of voice.

The relation of RLN with the inferior thyroid artery is shown in the image below:



Bilateral recurrent laryngeal nerve palsy can also occur postoperatively and the presentation is acute. Other causes of it include carcinoma of the cervical esophagus, carcinoma thyroid, and cervical lymphadenopathy. All the intrinsic muscles are paralyzed in this condition.

The vocal cords lie in the median or paramedian position due to the unopposed action of the cricothyroid muscles. This leads to dyspnoea and stridor, necessitating emergency tracheostomy.

For long-standing cases of bilateral recurrent laryngeal nerve palsy, there are two main treatment options. One approach involves performing a tracheostomy with a speaking valve. Alternatively, cord lateralizing procedures can be considered, such as transverse cordotomy, partial arytenoidectomy, injection of botulinum toxin, thyroplasty type II, or re-innervation procedures.

Transverse cordotomy (Kashima operation) involves excising laterally the soft tissue at the junction of the membranous cord and the vocal process of the arytenoid with laser.

Partial arytenoidectomy involves excising the medial part of the arytenoid with a laser.

Thyroplasty type II creates a lateral expansion of the larynx which is similar to vocal cord lateralization.

Reinnervation procedures innervate paralyzed posterior cricoarytenoid muscle by implanting a nerve-muscle pedicle of sternohyoid or omohyoid muscle with its nerve supply from ansa hypoglossi.

Solution to Question 3:

The clinical vignette of anosmia, foul-smelling discharge, and yellowish-green crusts in the nasal cavity points to a diagnosis of atrophic rhinitis. The other examination finding that is observed in atrophic rhinitis is a roomy nasal cavity.

Atrophic rhinitis is a chronic inflammation of the nose characterized by atrophy of nasal mucosa and turbinate bones. The exact etiology of this condition is not known.

The patients present with nasal obstruction, foul odor, merciful anosmia, greenish-grey crusting of the nasal passage, pale nasal mucosa, atrophied nasal turbinates, and epistaxis following the removal of the crusts. Roomy nasal cavity is seen due to the resorption of bones of the turbinates.

Treatment options for atrophic rhinitis include nasal irrigation using warm saline or an alkaline solution composed of sodium bicarbonate, sodium biborate, and sodium chloride (1:1:2) mixed with water.

Local antibiotics such as Kemicetine antiozaena solution, which contains chloromycetin, oestradiol, and vitamin D2, can be used. Topical applications of 25% glucose in glycerine, estradiol spray, and systemic administration of streptomycin, potassium iodide, or placental extract may also be employed.

Surgical management options include Young's operation, where flaps are raised within the nasal vestibule and both nostrils are completely closed, then opened after 6 months. During this time, the nasal mucosa may revert to a more normal state, and crusting is reduced. Another option is the modified Young's operation, where the nostrils are partially closed to avoid discomfort. The results of this modified approach are comparable to the original Young's operation.

Additionally, the nasal cavity can be narrowed through submucosal injection of teflon paste or by inserting fat, cartilage, or teflon strips beneath the mucoperiosteum.

The image below shows the endoscopic appearance of atrophic rhinitis:



Other options:

Option B: Nasal polyps are non-neoplastic masses of the edematous nasal or sinus mucosa. Nasal polyps can be antrochoanal polyps or ethmoidal polyps.

Option C: Inferior turbinate hypertrophy is associated with various mucosal diseases which include allergic rhinitis, infectious rhinitis, and vasomotor rhinitis. It can cause chronic nasal obstruction.

Option D: Foreign body nose is commonly seen in children and the child presents with unilateral, foul-smelling nasal discharge.

Solution to Question 4:

The next appropriate step for the management of epistaxis which was not controlled by pinching of nostrils and nasal packing would be ligation of the sphenopalatine artery, as it is the main artery contributing to Kiesselbach’s plexus.

Little's area is the most common site of epistaxis. It is present in the anteroinferior part of the septum. Four arteries anastomose here to form Kiesselbach’s plexus.

The image below shows the blood supply of the nasal septum and areas of epistaxis.



The first step in treating epistaxis is to pinch the nose with the thumb and index finger for about 5 minutes. The patient should sit leaning slightly forward over a basin to spit out blood and breathe through the mouth (Hippocratic or Trotter's method).



If the bleeding point is localized, cauterization can be done with electrocautery. In cases of uncontrolled bleeding or when the bleeding is posterior into the throat, anterior or posterior nasal packing may be required, respectively. If bleeding continues despite these measures, endoscopic procedures or vessel ligation close to the bleeding point can be considered.





The recommended hierarchy of ligation of vessels in a patient of epistaxis is the sphenopalatine artery first, the internal maxillary artery followed by the external carotid artery, and finally the anterior and posterior ethmoidal arteries.

The sphenopalatine artery is ligated first through a trans-nasal endoscope (transnasal endoscopic sphenopalatine artery ligation) at the sphenopalatine foramen which is situated at the posterior end of the middle turbinate.

However, if the bleeding persists, internal maxillary artery ligation is done at the pterygopalatine fossa via the Caldwell-Luc approach. If bleeding is still not controlled, external carotid artery ligation is done just above the origin of the superior thyroid artery.

If the bleeding site is located superior to the level of the middle turbinate, the anterior and posterior ethmoidal arteries should be ligated via Lynch incision in the medial wall of the orbit.

Solution to Question 5:

The given clinical picture suggests reactionary hemorrhage post-tonsillectomy. The immediate management is to shift the patient to OT, remove the clots, and cauterize/ligate the bleeding vessel.

Hemorrhage is a common complication of tonsillectomy. It can be primary, reactionary, or secondary.

Primary hemorrhage occurs at the time of operation. It can be controlled by pressure, ligation, or electrocoagulation of the bleeding vessels.

Reactionary hemorrhage occurs within 24 hours. It may be due to improper clipping action of the superior constrictor muscle.

The tonsillar artery perforates the superior constrictor and ramifies in the tonsil. Therefore, the presence of a clot can hinder the clipping action of the superior constrictor muscle, leading to hemorrhage. Slippage of ligature, or blood pressure returning to normal are other causes.

It is managed by removing the clot and application of pressure or vasoconstrictor. Ligation or electrocoagulation of the bleeding vessels is done under anesthesia if the above measures fail.

The image below shows the relation between the superior constrictor muscle and the tonsillar artery at the tonsillar bed.



Secondary hemorrhage is usually seen between the 5th to 10th postoperative day, due to sepsis and premature separation of the membrane. Clots are removed, IV antibiotics are given and the bleeding vessel is electrocoagulated or ligated if bleeding continues.

Solution to Question 6:

The clinical vignette depicting features of CNS infection (fever, headache, vomiting, and confusion) in a patient with a chronic ear infection and the CT image showing a ring-enhancing lesion point to a diagnosis of a temporal lobe abscess.

Temporal lobe abscess is an intracranial complication that can arise from chronic suppurative otitis media (CSOM).

Brain abscess typically progresses through four stages. The first stage is the stage of invasion (initial encephalitis), which often goes unnoticed. Symptoms such as headache, malaise, low-grade fever, and drowsiness may be present.

The second stage is the stage of localization (latent abscess), during which a capsule forms around the abscess, effectively localizing the pus.

The third stage is the stage of enlargement (manifest abscess). The abscess enlarges, leading to increased intracranial pressure (ICP) and the development of edema surrounding the abscess. As a result, symptoms worsen.

The final stage is the stage of termination (rupture of abscess). At this point, the abscess expands and eventually ruptures into the ventricle or subarachnoid space, which can result in potentially fatal meningitis.

Signs of raised ICP may include headache, nausea, vomiting, lethargy, drowsiness, confusion, papilledema, and coma.

Characteristic findings associated with temporal lobe abscess include nominal aphasia, homonymous hemianopia, seizures, contralateral motor paralysis, pupillary changes, and oculomotor palsy.

The preferred diagnostic imaging modality for temporal lobe abscess is a CT scan of the brain. This imaging study reveals a ring-enhancing lesion indicative of a brain abscess.

Medical treatment involves initiating intravenous antibiotics, typically metronidazole. Raised ICP can be addressed using dexamethasone or mannitol.

Surgical intervention is the definitive management and involves drainage of the abscess followed by radical mastoidectomy to address the underlying CSOM.

Other options:

Option B: Cerebellar abscess is a form of brain abscess and is usually associated with an otogenic source of infection. It can occur as a direct extension through Trautmann's triangle or via retrograde thrombophlebitis.

An MRI image showing cerebellar abscess is given below:



Option C: Subdural abscess refers to the focal collection of purulent material between the dura mater and the arachnoid mater.

An MRI image showing subdural abscess is given below:



Option D: Meningitis is the inflammation of the leptomeninges which includes the arachnoid mater and pia mater. It can be due to an infectious or noninfectious etiology. It presents with fever, vomiting, neck stiffness, and signs of meningeal irritation. Ring enhancing lesion on neuroimaging is not a feature of meningitis.

ENT INICET 2022

Question 1:

CT scan of a patient with laryngeal carcinoma is shown below. What is the staging?

a) T1N0M0

b) T4N1M0

c) T2N1M0

d) T3N0M0

Question 2:

Arrange the following in the sequence of auditory pathway:

a) 1-2-3-4-5

b) 5-4-3-2-1

c) 2-1-3-4-5

d) 3-4-5-1-2

Question 3:

What is the grade of tympanic membrane retraction shown in the image?

a) Grade 1

b) Grade 2

c) Grade 3

d) Grade 4

Question 4:

A patient presents with vertigo associated with horizontal nystagmus. The slow component is towards the left. What is the most likely diagnosis?

a) Posterior canal BPPV

b) Superior canal BPPV

c) Right hypoactive labyrinth

d) Left hypoactive labyrinth

Question 5:

What is the site of drainage of the marked structure?



a) Superior meatus

b) Middle meatus

c) Inferior meatus

d) Sphenoethmoidal recess

Question 6:

A 6-year-old boy presented with recurrent URTI, poor growth, high-arched palate, and impaired hearing. Tympanogram is given as follows. What would be the most appropriate management?



a) Grommet insertion

b) Adenoidectomy with grommet insertion

c) Myringotomy with grommet insertion

d) Myringotomy

Question 7:

A gentleman presents to the OPD with a saddle nose deformity. He gives a past history of cough, on/off fever, and hemoptysis. On examination, septal perforation is present with pale granuloma. His chest x-ray showed multiple cavitary lesions, and the biopsy showed granuloma, multinucleate giant cells, and caseous necrosis. What would be the probable cause?

a) Wegener's granulomatosis

b) Tuberculosis

c) Syphilis

d) Sarcoidosis

Question 8:

Foreign body entry into the laryngeal inlet is prevented by the cough reflex. This reflex is sluggish in alcohol ingestion and some other conditions. Which nerve is likely injured?

a) Internal laryngeal nerve

b) Glossopharyngeal nerve

c) External laryngeal nerve

d) Recurrent laryngeal nerve

Question 9:

A woman presents with impaired hearing. Her tympanogram findings are given below. Pick the right combination of tuning fork test results that would be seen in this patient.

a) Left Rinne's test negative, Weber's test lateralized to right ear

b) Left Rinne's test negative, Weber's test lateralized to left ear

c) Left Rinne's test positive, Weber's test lateralized to right ear

d) Left Rinne's test positive, Weber's test lateralized to left ear

Question 10:

A 35-year-old female patient presents with complaints of nasal obstruction and post-nasal drip. There is a past history of FESS for failed conservative management 5 years ago. Uncinectomy and maxillary ostium dilation was done during the previous FESS. A DNE done now shows patent ostia and mucosal edema of the maxillary sinus lining. What is the next best step in management?

a) Repeat surgery

b) Macrolides for 4 months

c) Steroid irrigation and antihistamines

d) Biological therapy

Question 11:

Which of the following tests cannot be used to assess nasal mucociliary clearance?

a) Saccharine test

b) Electron microscopy

c) Charcoal powder method

d) Scintigraphy

Question 12:

Match the following.

a) a-1, b-2, c-3, d-4

b) a-2, b-3, c-4, d-1

c) a-2, b-4, c-3, d-1

d) a-1, b-3, c-4, d-2

Question 13:

What test is not used for testing olfaction?

a) Smell diskettes

b) Arnold stick test

c) UPSIT

d) CC-SIT

Question 14:

Which of the following tests can be used if otoacoustic emissions is absent for screening?

a) 2 and 4

b) 1, 3, 4

c) 2, 3, 4

d) 2 and 3

Answer Key

Question No.	Correct Option
1	b
2	c
3	b

4	d
5	d
6	b
7	b
8	a
9	c
10	c
11	b
12	c
13	b
14	a

Detailed Explanations

Solution to Question 1:

The given CT image depicts a mass invading the thyroid cartilage as pointed by the red arrows. This indicates a tumor (T) stage of T4. Among the given options, T4N1M0 is the only option with a T4 staging and hence the best answer.

Laryngeal cancer is more commonly seen in men. Alcohol, tobacco, and previous radiation to the neck region are some of the risk factors for the development of laryngeal cancer. The most common type is squamous cell carcinoma.

Larynx is divided into three regions - supraglottis, glottis, and subglottis. In the TNM system: T indicates the extent of the tumor, N indicates lymph node involvement, and M indicates distant metastasis.

TNM classification of laryngeal carcinoma:

Tumor (T)

• T1: Only one site/subsite is involved (one named structure) with both vocal cords mobile.

• T1a: Only one vocal cord involved

• T1b: Involvement of both vocal cords

• T2: Spread to adjacent sites/subsites

• T2a: Both vocal cords are mobile

• T2b: Slight impairment in the mobility of vocal cord

• T3

• Fixed vocal cord

• Involvement of pre-epiglottic space, para-glottic space, post-cricoid space, or inner cortex of thyroid

• T4a: Local invasion.

• Anteriorly: Thyroid cartilage, thyroid gland, strap muscles

• Superiorly: Tongue muscles

• Inferiorly: Trachea

• Posteriorly: Esophagus

• T4b: Unresectable

• Involvement of prevertebral space, carotid sheath, or mediastinum

Regional lymph nodes (N)

• Nx: Cannot be assessed

• No: No regional lymph node metastasis

• N1: Metastasis in a single ipsilateral lymph node, ≤3cm

• N2: Metastasis in a single ipsilateral lymph node, >3cm but <6cm or multiple ipsilateral lymph nodes or bilateral lymph nodes none more than 6cm

• N3: Metastasis in a lymph node more than 6cm

Distant metastasis (M)

• Mx: Metastasis cannot be assessed

• Mo: No distant metastasis

• M1: Distant metastasis

Solution to Question 2:

The correct sequence of the auditory pathway is spiral ganglion (2) - cochlear nucleus (1) - superior olivary nucleus (3) - inferior colliculus (4) - medial geniculate body (5).

The spiral ganglion contains bipolar cells whose dendrites form the afferent supply for inner and outer hair cells of the organ of Corti and whose axons form the cochlear division of CN VIII that end in the cochlear nuclei.

The auditory pathway can be remembered with the mnemonic E.C.O.L.I.M.A

Eighth nerve

Cochlear nuclei

Olivary complex

Lateral lemniscus

Inferior colliculus

Medial geniculate body

Auditory cortex



Solution to Question 3:

The given image shows retraction in both pars tensa and pars flaccida. It is grade 2 retraction as the pars flaccida retraction is in contact with the neck of malleus while that in parts tensa is in contact with the incus.

Retraction pockets on tympanic membrane



Tympanic membrane can be divided into two parts. Pars tensa makes up most of the tympanic membrane. It is peripherally thickened to form annulus tympanicus (a fibrocartilaginous ring). It contains the cone of light. Pars flaccida (Shrapnell's membrane), is slightly pinkish and is located above the lateral process of malleus.

Tympanic membrane retraction can be either in the pars tensa or pars flaccida.

Tos classification is used to grade the retraction in pars flaccida or attic retraction

• Stage 1: The par flaccida is dimpled but not in touch with the malleus.

• Stage 2: The tympanic membrane is in contact with the neck of malleus and the entire extent of retraction can be visualized.

• Stage 3: The retraction is not visible fully, and there may be partial erosion of bony attic wall.

• Stage 4: The retraction is definite erosion of the bony attic wall and the full extent of retraction is not visible.

Due to difficulty in distinguishing the latter two, they are grouped as stage 3/4.

Sade classification is used to grade the retraction in pars tensa

• Stage 1: The tympanic membrane is retracted but not in touch with the incus

• Stage 2: The tympanic membrane is in contact with the long process of the incus

• Stage 3: Middle ear atelectasis - the tympanic membrane is in contact with the promontory, but not adherent. It can be raised from the promontory with suction

• Stage 4: Adhesive otitis media - the tympanic membrane is adherent to the promontory

Solution to Question 4:

The most likely diagnosis for a patient presenting with horizontal nystagmus and fast component towards the right (slow component towards the left) is left hypoactive labyrinth.

Nystagmus is an involuntary oscillatory movement of the eyes, which can occur in horizontal, vertical, or torsional directions. The direction of nystagmus is determined by the fast component.

In horizontal nystagmus, the direction of nystagmus (fast phase) is always away from the hypoactive lesions and towards the active or hyperactive lesions of the labyrinth. Benign paroxysmal positional vertigo (BPPV) is a hyperactive disease while vestibular neuronitis and vestibular Schwanomma are hypoactive diseases.

The involvement of the vertical semicircular canals (superior and posterior) leads to vertical or mixed vertical-torsional nystagmus, ruling out posterior canal BPPV (Option A) and superior canal BPPV (Option B).

Alternatively, if the right hypoactive labyrinth was one of the options, it could be considered as a differential for horizontal nystagmus with the slow component towards the right.

BPPV is a condition characterized by recurrent episodes of transient vertigo (<1 min) triggered by changes in head position relative to gravity. It is caused by dislodged calcium carbonate crystals (otoconia) from the utricular macula into one of the semicircular canals, usually the posterior semicircular canal. A diagnosis of BPPV can be made by Dix-Hallpike test and the condition is treated by repositioning maneuvers for the otoconia such as the Epley maneuver.

Solution to Question 5:

The marked structure in the given image is the sphenoid sinus and it drains into the sphenoethmoidal recess.

Paranasal sinuses are air-containing cavities that are present in the skull bones. These paranasal sinuses have been divided into two groups:

Anterior group: Their ostia drain into the middle meatus (Option B) through a common drainage pathway known as osteomeatal unit. This group includes:

• Maxillary sinus (antrum of Highmore)

• Frontal sinus

• Anterior ethmoidal sinus

Posterior group:

• Posterior ethmoidal sinuses: Opens in the superior meatus (Option A)

• Sphenoid sinus: Opens in sphenoethmoidal recess

The inferior meatus (Option C) has the Nasolacrimal duct draining into it with its terminal end being guarded by the Hasner's valve.



The images below show the location of paranasal sinuses:





Solution to Question 6:

The given clinical scenario along with type B tympanometry curve is suggestive of serous otitis media secondary to adenoid hypertrophy. Adenoidectomy with grommet insertion would be the most appropriate management.

Enlarged adenoids in children can present as nasal obstruction, rhinolalia clausa, recurrent acute otitis media, and serous otitis media. Adenoid facies, characterized by an elongated face, open mouth, teeth crowding, high arched hard palate, and pinched-in nose, may also be present.

Serous otitis media also known as secretory otitis media, mucoid otitis media, or glue ear is characterized by the accumulation of nonpurulent, serous effusion in the middle ear cleft.

Glue ear can occur either due to eustachian tube dysfunction (adenoid hyperplasia, chronic sinusitis or tonsillitis or rhinitis, cleft palate, tumors of nasopharynx) or due to increased secretory activity of the middle ear mucosa (allergy, unresolved otitis media, viral infections).

Glue ear is commonly seen in children aged 5-8 years and is associated with insidious onset of hearing loss (not exceeding 40 dB), mild ear aches, and a history of upper respiratory tract infections. Delayed or defective speech may also result from hearing loss.

Otoscopy shows:

• Dull and opaque tympanic membrane with loss of light reflex

• Thin leash of vessels along the handle of the malleus

• Fluid levels and air bubbles can be seen when the tympanic membrane is transparent.

The otoscopy image of serous otitis media is shown below:



The tuning fork test reveals a conductive type of hearing loss and audiometry reveals hearing loss in the range of 20–40 dB. The fluid accumulation in glue ear causes decreased mobility (compliance) of the tympanic membrane leading to a flat or dome-shaped curve or type B curve on a tympanogram.

Treatment is mainly aimed at the removal of fluid and preventing recurrence.

• Medical management includes decongestants, antihistamines, and antibiotics.

• Surgical management is done if the condition cannot be controlled with medications.
It includes myringotomy and aspiration of fluid with grommet insertion. Adenoidectomy and tonsillectomy may be required and are usually done at the time of myringotomy.

Solution to Question 7:

In a patient with saddle nose deformity, septal perforation with pale granuloma and caseous necrosis on biopsy is consistent with tuberculosis (TB). The history of fever, cough, and hemoptysis along with cavitary lesions on chest x-ray support the diagnosis of TB.

While other conditions like Wegener's granulomatosis (Option A), syphilis (Option C), and sarcoidosis (Option D) can also cause saddle nose deformity, caseous necrosis on biopsy makes TB a more likely diagnosis.

Saddle-nose deformity refers to a depressed nasal dorsum that can involve either the bony, cartilaginous, or both components.

The most common cause is nasal trauma resulting in depressed fractures. Other causes include excessive submucosal resection, destruction of septal cartilage due to hematoma or abscess, and infections such as leprosy, tuberculosis, or syphilis, as well as inflammatory conditions like granulomatosis with polyangiitis (Wegener granulomatosis) and sarcoidosis.

Chronic granulomatous conditions such as lupus, tuberculosis, and leprosy can affect the cartilaginous part, syphilis primarily affects the bony part, and Wegener's granulomatosis and T-cell lymphoma can lead to total (bony and cartilaginous) septal destruction involving both the

bony and cartilaginous portions.

TB is an infectious disease caused by Mycobacterium tuberculosis and primarily affects the lungs but can also involve the nose and sinuses.

Common symptoms of TB include fever, productive cough, weight loss, night sweats, and hemoptysis. Chest x-ray often shows multiple cavitation lesions affecting the upper lobe and hilar lymphadenopathy. Biopsy findings in tuberculosis typically reveal granulomas, caseous necrosis, and multinucleate giant cells (Langhan's giant cells).

The following images show the histopathological image of tuberculosis:



Other options:

Option A: Wegener's granulomatosis is characterized by vasculitis affecting small to medium-sized vessels. It affects the upper respiratory tract (ear, nose, sinuses, and throat), lower respiratory tract (lung), and kidney and is characterized by necrotizing granuloma which is non-caseating.

Option C: Although nose deformity is seen in syphilis, gummatous lesions (caseating granulomas) are present in tertiary syphilis which is not seen in this patient. Fever, cough, and hemoptysis are not features of syphilis.

Option D: Sarcoidosis most commonly affects the lungs forming non-caseating granuloma with no central necrosis. Laminated calcium and protein-containing bodies called Schaumann bodies and star-shaped inclusions called asteroid bodies are also found in the giant cells of most of these granulomas. Chest x-ray would show interstitial fibrosis and hilar lymphadenopathy.

The image below shows the characteristic non-caseating sarcoid granuloma:



Solution to Question 8:

The internal laryngeal nerve is the nerve implicated in the inhibition of the cough reflex in an alcoholic. Alcohol can cause chemical irritation of the nerve and it is also a potent inhibitor of the central nervous system.

The cough reflex is a protective mechanism of the larynx to expel irritants and keep the airway clear. Chemical or mechanical irritation of the respiratory mucosa stimulates the afferent internal laryngeal nerve, which is the branch of the superior laryngeal nerve of the vagus nerve.

Internal laryngeal nerve supplies the mucosa of the larynx above the level of the vocal cords (supraglottis). Hence, any injury to the internal laryngeal nerve results in aspiration and depressed cough reflex with no change in voice.

Recurrent laryngeal nerve (Option D) gives sensory supply to the mucosa of the larynx at the level of vocal cords (glottis) and below the level of vocal cords (subglottis) and also motor supply to all the intrinsic muscles of the larynx except cricothyroid.

Unilateral injury to the recurrent laryngeal nerve (RLN) may cause minimal or no change in voice, but it does not lead to aspiration or airway obstruction.

Bilateral injury to RLN can result in dyspnea and stridor due to the median or paramedian position of the vocal cords, which narrows and obstructs the airway. However, the voice remains unaffected.

The glossopharyngeal nerve (Option B) provides sensory supply to the oropharynx, posterior third of the tongue, and middle ear cavity. Damage to the glossopharyngeal nerve leads to the absence of the gag reflex.

The external laryngeal nerve (Option C) innervates the cricothyroid muscle, which is responsible for adjusting the pitch of sound during speech. Paralysis of the external laryngeal nerve results in the inability to increase the pitch of sound.



Solution to Question 9:

The given tympanogram findings indicate sensorineural hearing loss in the left ear. Tuning fork tests in this patient would show left Rinnie's positive and Weber lateralized to the right ear.

In left sensorineural hearing loss, Rinne's test will be positive on both left and right ears, and Weber's test will be lateralized to the better ear, which is the right.

Tuning fork tests are performed commonly with tuning forks of frequencies such as 128, 256, 512, 1024, etc. The tuning fork of 512 Hz is considered ideal as forks of lower frequencies produce a sense of bone vibration, while those of higher frequencies have a shorter decay time.

Rinne’s test is done by placing a vibrating tuning fork on the patient’s mastoid and when he/she stops hearing, it is brought beside the external auditory meatus (EAC). If he/she still hears the sound, air conduction (AC) is better than bone conduction (BC).

Weber test is done by placing a vibrating tuning in the middle of the forehead or the vertex and the patient is asked in which ear the sound is better heard.



Hearing loss	Rinne test (Conduction)	Weber test (Lateralized to)
None	Air > Bone	Both ears
Sensorineural	Air > Bone	Normal ear
Conductive	Bone > Air	Affected ear

Solution to Question 10:

The history of nasal obstruction and post-nasal drip and endoscopy showing mucosal edema is suggestive of acute rhinosinusitis. The next best step in management is symptomatic treatment with nasal steroid irrigation, decongestants, and antihistamines, regardless of the patient's past history of FESS operation.

Acute viral rhinosinusitis causes mucosal edema which causes obstruction of the sinus ostia. There is also reduced mucociliary transport in the sinuses resulting in stasis of secretions within the sinuses causing headache, facial pain, congestion, nasal obstruction or discharge, post-nasal drip, and hyposmia or anosmia.

If the sinuses remain obstructed or the mucociliary transport system does not return to normal, a bacterial infection can ensue, causing fever, purulent discharge, and persistent or worsening symptoms beyond 10 days.

In the case of bacterial sinusitis, antibiotics like amoxicillin and macrolides (erythromycin) are used for a short course (Option B).

Procedures like antral lavage and surgery (Option A) are done in cases not responding to medical therapy.

Biological therapy (Option D) is not used in the treatment of sinusitis.

Solution to Question 11:

Among the given options, electron microscopy will be the least useful in assessing nasal mucociliary clearance. Other tests mentioned can be used in humans to detect mucociliary transit time and total nasal clearance in humans.

Electron microscopy can be used to best assess the normalcy of cilia structure and its function, but it might not be able to detect any other abnormality that might interfere with nasal mucociliary clearance like any obstruction in the nasal cavity.

Test for mucociliary clearance include:

• Saccharine with or without color test in vivo (Option A)

• Radioisotope transport testing in vivo

• Measuring ciliary activity in vitro

• Nasal nitric oxide

• Scintigraphy (Option D) is the nuclear medicine equivalent of the dye disappearance test to assess functional and anatomic obstruction

The charcoal method (Option C) is a less commonly used method to detect nasal mucociliary clearance. In this method, a very small amount of charcoal is placed behind the head of the inferior turbinate. The pharynx is then observed to detect the presence of charcoal.

Mucociliary clearance tests are used to diagnose inborn disorders of mucociliary transport such as primary ciliary dysfunction and cystic fibrosis.

Solution to Question 12:

The correct match for Wullstein's classification is a-2, b-4, c-3, d-1 (Option C).

Solution to Question 13:

Arnold stick test is not a test used for olfaction.

Tests used for olfaction are as follows:

• Smell diskettes are used as a screening test for olfaction. Reusable diskettes are used with the application of 8 different odorants. A questionnaire with illustrations is used to score the patient.

• The University of Pennsylvania smell identification test (UPSIT) is a commercially available, self-administered test used to assess the function of an individual's olfactory system via smell identification. It uses microencapsulated odorants, which are released on scratching standardized odor-impregnated test booklets.

• The 12-item version of UPSIT, also known as the cross-cultural smell identification test (CC-SIT), is useful for quantifying smell loss when less time is available. This test serves as a means to reliably assess olfactory function in less than 5 minutes.

• The Sniffin' sticks test is a new test of nasal chemosensory performance based on pen-like odor-dispensing devices.

Solution to Question 14:

Among the given options, tympanometry (Test 2) and speech audiometry (Test 4) are used in cases where otoacoustic emission (OAE) is absent for screening.

Otoacoustic emissions are sounds produced by the outer hair cells of cochlea. These sounds then travel in the reverse direction from the outer hair — basilar membrane and perilymph of cochlea— oval window — ossicles and tympanic membrane (middle ear) — ear canal (external ear).

Hence in the absence of OAE, pathologies of the middle and external ear should be ruled out before making a definitive diagnosis of cochlear pathology.

Tympanometry is an objective test that can be performed in patients with absent OAE to rule out any middle ear pathologies. Speech audiometry is a subjective test that can be performed in these patients to confirm if the hearing loss is due to cochlear or retro cochlear pathology (neural hearing loss).

Impedance Audiometry also known as Tympanometry is an objective test that is based on the principle that when the sound strikes the tympanic membrane, certain amount of sound is absorbed, while the rest is reflected So a stiffer tympano-ossicular system would reflect greater sound than a compliant one. By marking the compliance of tympano-ossicular system against various pressure changes, different types of tympanograms are obtained which are diagnostic of various middle ear pathologies.

In speech audiometry, the patient's ability to hear and understand speech is measured. The speech reception threshold (SRT) is the lowest intensity of sound at which 50% of the words are repeated correctly by the patient. Speech discrimination scores test the capacity of the patient to understand speech. Hence, speech audiometry is a subjective test.

The rollover phenomenon is diagnostic of retrochochlear pathologies. Here, the phonetically balanced word score falls with an increase in speech intensity above a particular level due to nerve fatigue. The graph in cochlear disorders will rather show a plateau after a certain level unlike the fall seen in the rollover phenomenon.



A better objective test that could be used to diagnose retro-cochlear pathologies is brainstem auditory evoked response audiometry (BERA). It is a noninvasive test that can find any

pathologies of central auditory pathways through the VIIIth nerve to pons and midbrain.

Other options:

1. Pure tone audiometry (Test 1) is a test that measures the threshold of hearing by air and bone conduction and thus the degree and type of hearing loss. Pure tone audiometry is a subjective test for hearing and is not employed in cases where OAE is absent when better objective tests such as tympanometry are available.

3. Free field audiometry (Test 3), a part of behavior observational audiometry is used in the age group where children start localizing sounds when an auditory signal is presented. This test is not useful in this case as the possibility of hearing loss is already established with absent OAE.

ENT NEET 2022

Question 1:

A post-COVID patient, who is a known diabetic, develops unilateral facial pain and loosening of teeth. Which investigation would you do to confirm the diagnosis on this patient?

a) MRI

b) Biopsy with histopathologic examination

c) Serum ferritin

d) HbA1c

Question 2:

A patient presents with the complaint of inability to close the eye, drooling of saliva, and deviation of the angle of the mouth. Which of the following nerves is most likely to be affected?

a) Facial nerve

b) Trigeminal nerve

c) Occulomotor nerve

d) Glossopharyngeal nerve

Question 3:

An adult man in a restaurant suddenly begins choking on his food. He is conscious. The following procedure was performed. Identify the procedure.



a) Heimlich's maneuver

b) Back slap

c) Chest thrust

d) Blind insertion of finger

Question 4:

A patient presents with a firm, tender, slow-growing mass below the ear as shown in the image below. What could be the diagnosis?

a) Bezold abscess

b) Parotid abscess

c) Upper cervical lymphadenopathy

d) Osteoma of the mandible

Question 5:

A 5-year-old child presents with reduced hearing for the past 2-3 months. The otoscopy finding is given below. What is the most likely diagnosis?

a) Myringitis bullosa

b) Serous otitis media

c) Acute otitis media

d) Pneumotympanum

Question 6:

A patient comes with a history of asthma and sinusitis. On looking into his medical records, you notice this has been attributed to Samter's triad. Which drug should be avoided in this patient?

a) Cotrimoxazole

b) Co-amoxiclav

c) Aspirin

d) Chloramphenicol

Question 7:

A 70-year-old male patient presents with decreased hearing in higher frequencies. It was noted that the basilar membrane was affected. Which of the following structures lie near the affected structure?

a) Modiolus

b) Stria vascularis

c) Oval window

d) Helicotrema

Question 8:

Identify the structure given in the image.

a) Malleus

b) Incus

c) Stapes

d) Vomer

Question 9:

A 20-year-old male patient presents with unilateral nasal obstruction and recurrent bleeding for the past 1 year. Transnasal endoscopic results are shown below. A contrast-enhanced CT revealed a mass extending from the posterior choana to the nasopharynx. What is the most likely diagnosis?



a) Nasopharyngeal angiofibroma

b) Antrochoanal polyp

c) Rhinoscleroma

d) Concha bullosa

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	b
6	c
7	c
8	b
9	a

Detailed Explanations

Solution to Question 1:

The given scenario of a post-COVID diabetic patient with facial pain and loosening of teeth is suggestive of rhino-orbital-cerebral mucormycosis. Among the given options, Biopsy with histopathologic examination would be the investigation for definitive diagnosis on this patient.

Rhino-orbital-cerebral mucormycosis is the most common form of mucormycosis. It commonly occurs in patients with diabetes mellitus.

The presenting features are facial pain, numbness, loosening of teeth, conjunctival swelling, congestion, and blurry vision. When the disease spreads to the orbit, proptosis can be present. It can later spread to the frontal lobe of the brain and cavernous sinus.

A definitive diagnosis is done with a biopsy and histopathologic examination. It is the most sensitive and specific method. It reveals aseptate fungi with thick-walled, ribbon-like, hyphae branching at right angles.

Among the imaging techniques used for diagnosis, MRI is more sensitive to detecting orbital and nervous involvement than CT. Contrast-enhanced MRI is the imaging technique of choice. 1,3 beta D- glucan assay, which is helpful for the identification of most fungi, cannot be used for the diagnosis of mucormycosis.

Management is by amphotericin B and surgical debridement of the affected tissues. The underlying cause and risk factors should also be addressed.

Solution to Question 2:

The given clinical features of inability to close the eye, drooling of saliva, and deviation of the angle of the mouth are suggestive of paralysis of the facial nerve.

Facial nerve paralysis can be central or peripheral. Peripheral is more common and is usually idiopathic.

Bell's palsy is an acute onset idiopathic, peripheral facial paralysis. It is a lower motor neuron lesion of the facial nerve and involves the ipsilateral side of the upper and lower halves of the face.

It presents with an inability to close the eye, and if the patient attempts to close the eye, it turns up and out. This is known as the Bell's phenomenon. Other features include drooling of saliva and facial asymmetry. There may be noise intolerance and a loss of taste sensation.

Bell's palsy is a diagnosis of exclusion. All other possible causes of facial paralysis should be ruled out by investigations.

Treatment of Bell's palsy involves prednisone 60-80 mg daily during the first 5 days and then tapered over the next 5 days. Tape to depress the upper eyelid during sleep and artificial tears can be used to prevent corneal drying. Most patients recover within a few weeks or months.

The following image shows a patient with Bell's palsy.



Other options:

Option B: Lesions of the trigeminal nerve produce numbness and paresthesias, and loss of sensation in the area supplied by the trigeminal nerve.

Option C: In oculomotor nerve palsy, there is ptosis, a dilated pupil, and the eye appears to be down and out due to the unopposed lateral rectus and superior oblique.

Option D: Glossopharyngeal nerve lesions produce difficulty swallowing, and impairment of taste over the posterior one-third of the tongue and palate. There is also impaired sensation over the posterior one-third of the tongue, palate, and pharynx, and an absent gag reflex.

Solution to Question 3:

The image shows a person standing behind with arms placed around the choking person's lower chest, giving abdominal thrusts, which is suggestive of Heimlich's maneuver.

The Heimlich maneuver (also known as subdiaphragmatic abdominal thrusts or abdominal thrusts) is done for an adult or a child (1-8 years) with foreign body airway obstruction (FBAO). It is not recommended for FBAO relief in infants.

The rescuer should stand or kneel behind the patient and wrap his arms around the patient's waist. A fist is made with one hand and using the other hand, the fist is driven into the patient's abdomen with a rapid, forceful inward and upward thrust to dislodge the foreign body. This is continued until the foreign body is expelled or if the patient becomes unresponsive.

If the patient becomes unresponsive, immediately start cardiopulmonary resuscitation (CPR).

Other options:

Option B: Back slaps or back blows can also be used to relieve FBAO. They are given between the shoulder blades using the heel of the rescuer's hand, and up to 5 are given. If there is a failure of the back slaps to relieve the obstruction, up to 5 abdominal thrusts can be given.

Infants (<1 year) should be treated with alternating 5 back blows and then 5 chest thrusts. The infant's torso is placed on the rescuer's nondominant arm. It is shown below.





©MARROW

Option C: Chest thrusts can be used as an alternative to the Heimlich maneuver for pregnant women or markedly obese patients. The patient is supine, and the rescuer should kneel close to the victim's side. Hands are placed in the same manner as is done to perform CPR chest compressions. Each thrust is given in such a way so as to relieve the obstruction.

Option D: The blind finger sweep should be avoided in infants to avoid pushing the object further inside the airway. However, it can be done for unresponsive adults with complete FBAO. With the patient's face up, the mouth is opened by grasping both the tongue and lower jaw and lifting the mandible. The index finger is inserted along the inside of the cheek and using a hooking action, the foreign body is dislodged and removed.

Solution to Question 4:

The given clinical scenario and the image are suggestive of a parotid abscess.

A parotid abscess refers to an infection and pus formation in the parotid space. It is caused most commonly by Staphylococcus aureus. Post-surgical cases and debilitated patients with stasis of salivary flow are at risk of developing parotid abscesses.

Patients typically present with unilateral swelling, redness, induration, and tenderness over the parotid gland and angle of the mandible. Dehydration and fever are also present. Diagnosis is by ultrasonography or a CT scan. Multiple small loculi of pus are seen in the parenchyma. Aspiration of abscess can be done to determine the culture and sensitivity of the infective agent.

Correction of dehydration, improving oral hygiene, and promoting salivary flow. Intravenous antibiotics are also used in treatment. Surgical drainage of the abscess is done via a preauricular incision.

The parotid gland is enclosed by deep cervical fascia which splits into superficial and deep layers. Parotid space lies deep in the superficial layer. Contents of parotid space include :

• parotid gland

• lymph nodes associated with parotid gland

• facial nerve

• external carotid artery

• retromandibular vein

Other options:

Option A: Bezold abscess is a complication of acute coalescent mastoiditis and is formed when pus breaks inferiorly, through the tip of the mastoid and collects in the upper part of the neck. The onset is sudden with pain, fever, tender neck swelling, and torticollis. There will be a history of purulent otorrhea.

Option C: Osteoma of the mandible is usually slow growing and asymptomatic. It usually occurs in the mandibular condyle area. They are usually detected incidentally during a radiographic examination.

Option D: Upper cervical lymphadenopathy occurs in the neck region following infection in the area drained by these structures.

Solution to Question 5:

The given clinical scenario of a child presenting with hearing loss for the past 2-3 months and the otoscopic finding is suggestive of serous otitis media. The given otoscopic image shows a dull and opaque tympanic membrane with clear effusion.

Serous otitis media is also known as secretory otitis media or mucoid otitis media or glue ear. It is characterized by the accumulation of nonpurulent effusion in the middle ear cleft.

Glue ear is thought to happen due to either malfunctioning (adenoid hyperplasia, chronic sinusitis or tonsillitis or rhinitis, palatal defects, tumors of nasopharynx) or increased secretory activity of the eustachian tube (allergy, unresolved otitis media, viral infections).

It is often seen in children of 5-8 years of age. They present with hearing loss not exceeding 40 dB, which is insidious in onset. Speech may be delayed and defective due to hearing loss. Mild earaches with a history of upper respiratory tract infection are usually seen.

On otoscopy,

• Dull and opaque tympanic membrane with loss of light reflex

• Thin leash of vessels along the handle of the malleus

• If the tympanic membrane is transparent fluid levels and air bubbles can be seen



The tuning fork test reveals a conductive type of hearing loss and audiometry reveals hearing loss in the range of 20-40 dB. On impedance audiometry, there is reduced compliance and a flat curve shifted to the negative side due to the presence of fluid. X-ray mastoid reveals clouding of air cells.

Treatment is mainly aimed at the removal of fluid and preventing recurrence.

• Medical management includes decongestants, antihistamines, antibiotics, and middle ear aeration techniques.

• Surgical management includes myringotomy and aspiration of fluid, grommet insertion, tympanotomy, or cortical mastoidectomy.

Chronic glue ear sequelae include atrophic tympanic membrane and middle ear atelectasis, ossicular necrosis, tympanosclerosis, cholesterol granuloma, and retraction pockets.

Other options:

Option A: Myringitis bullosa is characterized by painful hemorrhagic blebs on the tympanic membrane and deep meatus. It occurs due to infection by a virus or mycoplasma.

Option C: Acute otitis media is an acute infection of the middle ear by pyogenic organisms. It presents with marked earache and hearing loss is seen in later stages. Otoscopy shows a red and bulging tympanic membrane.



Solution to Question 6:

In a patient with Samter's triad, aspirin should be avoided as he will have aspirin intolerance.

Samter's triad is also known as aspirin-induced asthma or aspirin-sensitive asthma. It consists of:

• Nasal polyposis

• Bronchial asthma

• Aspirin intolerance

Patients with Samter's triad will have persistent nasal congestion, rhinorrhea, anosmia, nasal polyps, and recurrent chronic rhinosinusitis. Asthma and aspirin intolerance develop later.

Acute asthma attacks usually occur 3 hours after intake of aspirin or NSAIDs. Severe rhinorrhea, conjunctival injection, facial flushing, and periorbital edema are also seen. Aspirin-induced hypersensitivity reactions also may occur.

Patients should avoid aspirin and NSAIDs that inhibit COX-1. Treatment may involve aspirin desensitization and endoscopic sinus surgery.

Solution to Question 7:

The affected structure in high-frequency hearing loss is the basal turn of the cochlea. Among the given options, the structure that lies near it is the oval window.

The traveling wave theory of von Bekesy states that a sound wave, depending on its frequency, reaches maximum amplitude at a segment on the basilar membrane and stimulates that segment. The high frequencies are represented in the basal turn of the cochlea and the progressively lower ones towards the apex.

The cochlea has a bony core called the modiolus, and a spiral canal that runs around it. At the base of the canal, the two main openings are the oval window and the round window. The footplate of stapes rests against the oval window. The base of the modiolus transmits nerves and vessels to the cochlea. (option A)

The inner ear consists of a bony labyrinth and a membranous labyrinth. The bony labyrinth consists of the vestibule, semicircular canals, and the cochlea. The membranous labyrinth consists of the cochlear duct, the utricle and saccule, semicircular ducts, and the endolymphatic duct and sac.

The cochlear duct has three walls- the basilar membrane, Reissner's membrane, and the stria vascularis. The basilar membrane supports the organ of Corti, which is the sensory hearing epithelium. The length of the basilar membrane increases from the base to the apex. As a result of this, higher frequency sound is heard at the basal coil while lower frequencies are heard at the apex.

The stria vascularis is involved in the secretion of endolymph and contains vascular epithelium. (Option B)



The helicotrema (option D) is an opening at the apex of the cochlea, at which the scala vestibuli and scala tympani communicate. The scala vestibuli, tympani, and media are three compartments of the bony cochlea.

Solution to Question 8:

The structure given in the image is the incus.

It is one of the three ossicles that are in the middle ear cavity i.e, malleus, incus, and stapes. Ossicles conduct sound energy from the tympanic membrane to the oval window and then to the inner ear fluid.

Incus is shaped like an anvil. It has a body, a short process, and a long process. It has a saddle-shaped articular facet on its anterior surface that articulates with the malleus. The long

process descends vertically and ends as a rounded lenticular process.

Incus has a single nutrient vessel within the long process and no collateral circulation. Thus, it is more susceptible to aseptic necrosis following a middle ear infection.

The malleus (option A) is the largest ossicle and is shaped like a mallet. The parts of the malleus are the head, neck, handle (manubrium), and anterior and lateral processes. The head and neck lie in the attic. The tympanic membrane is connected to the handle of the malleus.

The stapes (option C) has a head, neck, footplate, and anterior and posterior crura. The annular ligament holds the footplate of stapes in the oval window.

The joints between the ear ossicles are synovial joints. A saddle-shaped joint is present between the incus and malleus. Incus and stapes articulate via a ball and socket joint.





Option D- The vomer is a small bone in the posteroinferior part of the nasal septum. Along with perpendicular plate of the ethmoid, and the septal cartilage, it forms the nasal septum.

Solution to Question 9:

The above clinical scenario with recurrent spontaneous bleeding with unilateral nasal obstruction and contrast-enhanced CT showing a growth protruding into the nasopharynx, along with the transnasal endoscopic image, is suggestive of nasopharyngeal angiofibroma.

Nasopharyngeal angiofibroma is a benign but locally invasive tumor. It is predominantly seen in adolescent males and is thought to be testosterone-dependent. It arises from the posterior part of the nasal cavity near the sphenopalatine foramen.

The clinical features include epistaxis, progressive nasal obstruction, and denasal speech. Other symptoms include conductive hearing loss, broadening of the nasal bridge, proptosis leading to frog-face deformity, cheek swelling, and involvement of II, III, IV, and VI cranial nerves.

Contrast-enhanced CT scan of the head is the investigation of choice that reveals anterior bowing of the posterior wall of the maxillary sinus (Antral sign/Holman Miller sign), which is pathognomonic of angiofibroma as shown below.

Carotid angiography shows the extent of the tumor with its vascularity. The feeding vessels mostly come from the external carotid artery. In large tumors with intracranial extension, vessels come from the internal carotid system also.

Biopsy of the mass is avoided due to the risk of bleeding.

Surgical approach is the treatment of choice. Embolization of the vessels can be done prior to surgery to reduce the risk of bleeding during the surgery.

Radiotherapy, chemotherapy, and hormonal therapy can also be considered.

Other options:

Option B: Antrochoanal polyp (Killian's polyp) arises from the mucosa of the maxillary antrum. They are usually unilateral and are seen in children and young adults. They present with unilateral nasal obstruction with minimal to no bleeding. X-rays of the paranasal sinuses

show opacity of the maxillary sinus.

Option C: Rhinoscleroma is caused by a gram-negative coccobacillus Klebsiella rhinoscleromatis. It causes this progressive granulomatous infection starting in the nose and extending into the upper respiratory tract. It presents initially with foul-smelling nasal discharge, and then with granulomatous nodules in the nose giving it a woody feeling. The last stage is the cicatricial stage, with stenosis of nares, and adhesions in the nose.

Option D: Concha bullosa is pneumatization of the middle turbinate that leads to an enlarged middle turbinate, which is ballooned out.

The CT image shown below depicts concha bullosa

ENT NEET 2023

Question 1:

Identify the structure marked in the image.



a) Fossa of Rosenmuller

b) Tubal tonsil

c) Opening of eustachian tube

d) Adenoid

Question 2:

A 30-year-old male presents with nonaxial proptosis of the left eye. The patient gives a history of a road traffic accident 15 years back. The CT image is given below. What is the most likely diagnosis?



a) Frontal mucocele

b) Frontal meningioma

c) Juvenile nasopharyngeal angiofibroma

d) Pseudotumor of orbit

Question 3:

The instrument shown in the image is not used for which of the following?

a) Airway toileting

b) Upper airway examination

c) Acute nasopharyngeal obstruction

d) Prolonged mechanical ventilation

Question 4:

A 10-year-old child presents with throat pain, fever, and ear pain. He is diagnosed with recurrent tonsillitis. Which nerve is responsible for the ear pain in this patient?

a) Tympanic branch of glossopharyngeal nerve

b) Greater auricular nerve

c) Auriculotemporal nerve

d) Auricular branch of vagus nerve

Question 5:

A 55-year-old patient comes with hoarseness of voice and difficulty swallowing. The patient was diagnosed with laryngeal carcinoma and surgical management was done. The post-operative image of the patient is given below. Which of the following surgery was done on this patient?



a) Partial laryngectomy

b) Percutaneous tracheostomy

c) Standard tracheostomy

d) Total laryngectomy

Question 6:

A female patient with hearing loss is examined and is found to be Rinne negative at 256 Hz and 512 Hz, while Rinne positive at 1024 Hz. What is the expected air conduction and bone conduction gap?

a) 30-45 dB

b) 15-30 dB

c) 45-60 dB

d) >60 dB

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	a
5	d
6	a

Detailed Explanations

Solution to Question 1:

The image shown is the endoscopic view of lateral wall of the nasopharynx, and the marked structure is the fossa of Rosenmuller.

The nasopharynx is present above the soft palate and is made up of the roof, floor, posterior wall, and two lateral walls.

Important structures seen in the nasopharynx are

• Pharyngeal opening of the eustachian tube

• Torus tubarius- tubal elevation which is present behind the opening of the eustachian tube.

• Fossa of Rosenmuller - A lateral pharyngeal recess that is present above and behind the torus tubarius. It is the most common site for nasopharyngeal carcinoma

• Adenoids or nasopharyngeal tonsil, which is present in the roof and posterior wall

• Passavant's ridge

The tubal tonsil is a collection of subepithelial lymphoid tissue situated at the tubal elevation. It is a part of the Waldeyer's ring and infection can cause eustachian tube occlusion.





Solution to Question 2:

The given clinical scenario and the image findings are suggestive of a frontal mucocele.

Mucoceles are slow-growing, mucous-filled benign masses that can develop years after head injury. Post-traumatic mucocele formation occurs due to chronic obstruction of the sinus ostium. The frontal sinus is most commonly involved in mucocele formation.

The sinuses commonly affected by mucocele in the order of frequency are the frontal > ethmoidal > maxillary > sphenoid.

Frontal mucocele presents with headache, diplopia, and proptosis. A cystic non-tender swelling is noticed with eggshell crackling.

The image given below shows a coronal MRI demonstrating a frontal mucocele and an orbital cystic mass.

Other options:

Option B: Frontal meningiomas are benign brain tumors presenting with headaches, seizures, vomiting, sensory and motor weakness, and visual impairment.

Option C: Juvenile nasopharyngeal angiofibroma is a benign neoplasm of the nasopharynx presenting with spontaneous recurrent epistaxis, nasal obstruction, and nasal discharge.

Option D: Pseudotumor of the orbit is an idiopathic inflammation of the orbit presenting with painful proptosis, diplopia, and orbital congestion. Imaging shows swollen eye muscles (orbital myositis) and enlarged tendons.

Solution to Question 3:

The image shows a tracheostomy tube. It is not used for upper airway examination.

Tracheostomy involves making an opening and placing a stoma in the anterior wall of the trachea.

Functions:

• Provides an alternative pathway for breathing by bypassing the obstruction.

• Improves alveolar ventilation by reducing dead space by 30-50% and reducing airway resistance

• Uses a cuff to protect the airways from secretions and blood

• Allows suction of secretions in the setting of the impaired cough reflex.

• Allows positive pressure respiration

• Allows administration of anesthesia

Indications:

• Respiratory obstruction due to

• Infections (laryngo-tracheo-bronchitis, epiglottitis, diphtheria)

• Trauma due to external injury, mandibular or maxillofacial fractures

• Neoplasms

• Foreign bodies or edema

• Bilateral abductor paralysis

• Congenital anomalies

• Retained secretions

• Inability to cough due to coma, spasm, or paralysis of respiratory muscles

• Aspiration of pharyngeal secretions

• Painful cough

• Respiratory insufficiency due to emphysema, chronic bronchitis

Solution to Question 4:

The tympanic branch of the glossopharyngeal nerve (cranial nerve IX) is responsible for referred otalgia from tonsillitis.

The tonsils and tonsillar fossa are supplied by the glossopharyngeal nerve. The glossopharyngeal nerve and the styloid process lie in relation to the lower part of the tonsillar fossa. Therefore, any irritation or pain can be referred to the ear along the tympanic branch of the glossopharyngeal (Jacobson's) nerve.

The image below shows the anatomic relations of the tonsil and tonsillar bed.



The ear receives extensive sensory innervation via four cranial nerves (V, VII, IX, and X) and two spinal segments (C2 and C3).

Referred otalgia transmitted by the glossopharyngeal nerve may be secondary to lesions and/or inflammatory processes of the nasopharynx, palatine tonsil, soft palate, or posterior one-third of the tongue.

Causes of referred otalgia via glossopharyngeal nerve:

• Acute pharyngitis

• Tonsillitis

• Peritonsillitis

• Peritonsillar abscess

• Post-tonsillectomy

• Foreign bodies embedded in the base of the tongue or tonsillar area

• Elongated styloid process

Other options:

Option B: The great auricular nerve provides sensory sensation for the lower two-thirds of the pinna.

Option C: Causes of referred pain via the auriculotemporal nerve which is the branch of the trigeminal nerve include caries tooth, ulcerative lesions of the oral cavity, temporomandibular joint disorders, and sphenopalatine neuralgia.

Option D: Causes of referred pain via the auricular branch of the vagus nerve include malignancy or ulcerative lesion of the vallecula, epiglottis, larynx, and esophagus.

Solution to Question 5:

The given image shows a post-operative tracheal stump created for breathing in patients undergoing total laryngectomy for laryngeal carcinoma.

In total laryngectomy, the entire larynx, including the hyoid bone, pre-epiglottic space, strap muscles, and one or more rings of trachea, are removed. The pharyngeal wall is repaired, and the tracheal stump is sutured to the skin to allow respiration.

The indications of total laryngectomy are as follows:

• T3 lesions

• All T4 lesions

• Invasion of thyroid or cricoid cartilage

• Bilateral arytenoid cartilage involvement

• Lesions of the posterior commissure

• Failure after radiotherapy or conservation surgery

• Transglottic cancers with cord fixation

It is contraindicated in patients with distant metastasis.

Other options:

Option A: Partial laryngectomy is a conservation procedure that involves the excision of the vocal cord and anterior commissure region in partial front lateral laryngectomy and the excision of supraglottis in partial horizontal laryngectomy. This surgery is performed to conserve the voice and avoid a permanent tracheal opening.

Option B: Percutaneous tracheostomy is done in ICU patients under sedation when the patient is already intubated and being monitored. The image below shows the procedure of percutaneous tracheostomy.



Option C: Standard tracheostomy can be therapeutic (used for respiratory obstruction, remove tracheobronchial secretions or assisted ventilation) and prophylactic (for anticipated respiratory obstruction or aspiration of blood or pharyngeal secretions).

Solution to Question 6:

The given clinical scenario of hearing loss with a negative Rinne at 256 Hz and 512 Hz and positive Rinne at 1024 Hz indicates air conduction and bone conduction gap of 30-45 dB.

Tuning fork tests are performed commonly with tuning forks of frequencies such as 128, 256, 512, and 1024. The tuning fork of 512 Hz is considered ideal as forks of lower frequencies produce a sense of bone vibration, while those of higher frequencies have a shorter decay time.

Rinne's test is done by placing a vibrating tuning fork on the patient's mastoid and when he stops hearing, it is brought beside the external auditory meatus (EAC). If he still hears the sound, air conduction (AC) is better than bone conduction (BC).

The inferences are as follows:

• Rinne's test is considered positive (normal) when the air conduction AC>BC is seen in normal people and people with sensorineural deafness.

• It is considered negative when the BC>AC, and is seen in patients with conductive deafness. It suggests a minimum hearing loss of 15-20 dB.

A prediction of the air-bone gap can be made using tuning forks of 256, 512, and 1024 Hz:

• A Rinne test negative for 256 Hz but positive for 512 Hz indicates an air-bone gap of 20-30 dB.

• A Rinne test negative for 256 and 512 Hz but positive for 1024 Hz indicates an air-bone gap of 30-45 dB.

• A Rinne negative for 256, 512, and 1024 Hz indicates an air-bone gap of 45-60 dB.

ENT INICET 2023

Question 1:

A 22 year old male presented with progressive left sided facial swelling for 2 weeks. There is history of traumatic injury with wood two months back. CECT nose and paranasal sinuses shows mass in subcutaneous tissue of left cheek. Biopsy of specimen shows foreign body granuloma with positive PAS and GMS stain. What is the diagnosis?

a) Phycomycosis

b) Midline lethal granuloma

c) NK/T cell lymphoma

d) IgG4 granuloma

Question 2:

Arrange the following structures in the hearing pathway from the receptors to the cortex -

a) 3-2-4-6-7-5-1

b) 2-3-4-6-7-5-1

c) 2-3-6-4-5-7-1

d) 3-2-6-4-7-5-1

Question 3:

Identify the nerve block shown in the given image

a) Nasociliary nerve

b) Anterior ethmoidal nerve

c) Sphenopalatine nerve

d) Greater palatine nerve

Question 4:

Arrange the following in the sequence of auditory pathway:

a) 1-2-3-4-5

b) 5-4-3-2-1

c) 2-1-3-4-5

d) 3-4-5-1-2

Question 5:

A 14-year-old is brought to the opd with complaints that she is able to hear the sound of speech but has difficulty understanding words. Her PTA and BERA were inconsistent with each other, with PTA being more or less normal. Her middle latency and cortical responses were absent. What is the likely diagnosis?

a) Michel aplasia

b) Auditory neuropathy

c) Malingering

d) Cochlear otosclerosis

Question 6:

A South Indian male cattle breeder comes with complaints of frank intermittent epistaxis. The examination of the nose is given below. A biopsy was taken and HPE appears as shown below. What is the likely diagnosis?



a) Nasal polyp

b) Rhinosporidiosis

c) Hemangioma

d) Rhinopyma

Question 7:

Which of the following statements are correct about hair cells?

a) i, ii, iv

b) i, ii, iii

c) i, iii, iv

d) ii, iii, iv

Question 8:

A 7-year-old male presents with progressive bulging of the right eye with a diminision of vision since 1 day. He has a history of URTI and on examination, it is seen that he also has nasal congestion with active discharge. The CT scan of the PNS is seen. What is the diagnosis?

a) Sinusitis with preseptal cellulitis

b) Sinusitis with orbital cellulitis

c) Sinusitis with subperiosteal abcess

d) Sinusitis with orbital abscess

Question 9:

Which type of synovial joint is seen in the picture?



a) Ball and socket

b) Saddle

c) Pivot

d) Plane

Question 10:

The audiogram given in the image below is seen in _____.



a) Ototoxicity

b) Presbycusis

c) Noise induced hearing loss

d) Endolymphatic hydrops

Question 11:

A farmer presents with unilateral obstruction of the nasal cavity and nasal bleeding. A mass is found and removed. A biopsy is shown below. What is the likely diagnosis?



a) Rhinosporidiosis

b) Rhinoscleroma

c) Inverted papilloma

d) Angiofibroma

Question 12:

Out of the four given statements, which is true about coughing?

a) i, ii, iii

b) i, iv

c) ii, iv

d) ii, iii

Question 13:

A 32-year-old female presents with progressive bilateral hearing loss, in the right ear more than left. After the initial evaluation, a stapedotomy of the right ear was planned. What might've been the tuning fork test results of the patient?

a) Bilateral Rinne's negative, Weber's lateralized to the right

b) Bilateral Rinne's negative, Weber's lateralized to the left

c) Bilateral Rinne's positive, Weber's lateralized to the right

d) Bilateral Rinne's positive, Weber's lateralized to the left

Question 14:

A 60-year-old man can only hear when words are shouted. What is the degree of his hearing impairment?

a) Moderate hearing loss < 50%

b) Moderately severe hearing loss < 70%

c) Profound hearing loss > 80%

d) Severe hearing loss > 70%

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	c
5	b
6	b
7	b
8	c
9	b
10	d
11	a
12	b
13	a
14	d

Detailed Explanations

Solution to Question 1:

The given clinical scenario is strongly suggestive of Phycomycosis/mucormycosis. A positive fungal stain(Grocott-Gomori's methenamine silver stain) narrows down the diagnosis. The cutaneous disease can be highly invasive, penetrating into muscle, fascia, and even bone.

Mucormycosis is an opportunistic mycosis caused by a number of molds classified in the order 'Mucorales'. These fungi are ubiquitous thermotolerant saprobes. The leading pathogens among this group are species of the genera Rhizopus, Mucor, and Absidia. Their spores are present in air and dust.

Mucormycosis can occur as isolated cutaneous or subcutaneous infection in immunologically normal individuals after traumatic implantation of soil or vegetation. Angio-invasiveness is one of the characteristics of zygomycosis which makes it fatal. Lungs are usually the primary site of infection. Almost all patients are immunocompromised. Rhino-orbital-cerebral mucormycosis occurs commonly in patients with diabetic ketoacidosis. It is the most common form; starts as eye and facial pain may progress to cause orbital cellulitis, proptosis, and vision loss. The typical finding is the presence of a black necrotic mass filling the nasal cavity.

Diagnosis:

• Exudate: wet preparation is made in 10% potassium hydroxide. Broad, ribbon-like hyphae are seen.

• Tissue: Hematoxylin & eosin stain shows non-septate, broad hyphae, often invading into blood vessels. Special stains help to identify the fungus in tissue sections.

• Biopsy confirms this diagnosis by revealing non-septate hyphae branching at right-angles, which are characteristic of Mucorales such as Rhizopus delemar. The hyphae are broad (10–15 μm) with uneven thickness, irregular branching, and sparse septations.

• Isolation is by culture on Sabouraud agar with an antibacterial agent but without cycloheximide. The colonies are grey-white with a thick cotton, fluffy surface.

Treatment is primarily with liposomal amphotericin B. Surgical debridement of the affected tissues and control of underlying predisposing cause is equally important.

Other options:

Option D: IgG4-related disease (IgG4-RD) is characterized by tissue infiltrates dominated by IgG4 antibodies (the hallmark of the disease). The pathogenesis of this condition is not understood. This condition leads to fibrosis and obliterative phlebitis. The patients may present as Riedel's thyroiditis, autoimmune pancreatitis, or idiopathic orbital inflammation. But not with ptosis as shown in the image. Diagnosis is established by positive fibrosis and the presence of plasma cells on biopsy. Rituximab is known to be beneficial. Sometimes, as a rare presentation, this can mimic Wegener's granulomatosis.

Option B: Midline lethal granuloma is a rare disease. It is characterized by necrotizing destruction and mutilation of the nose and other structures of respiratory passages. Generally, it is a diagnosis

of exclusion. CT may be helpful in picking up the lesions. Negative c-ANCA and fungal stains also aid in the diagnosis.

Option C: NK/T cell lymphoma is a destructive lesion usually starting on one side of nose involving the upper lip, oral cavity, maxilla and sometimes even extending to orbit.Histologically polymorphic lymphoid tissue with angiocentric and angioinvasive features is seen. Immunochemical staining of biopsy material should be done for CD56+ marker for diagnosis. Nearly all cases of T/NK cel lymphomas are associated with EB virus. Localized T/NK-cell lymphoma is treated by radiation while a disseminated disease requires chemotherapy.

Solution to Question 2:

The correct sequence of the auditory pathway is spiral ganglion - cochlear nucleus - Olivary complex -Lateral leminiscus- Inferior colliculus-Medial geniculate body-Auditory Cotex

The spiral ganglion situated in Rosenthal's canal contains bipolar cells. The dendrites of these bipolar cells join to form the cochlear branch of the vestibulocochlear nerve which ends in the cochlear nuclei.



Solution to Question 3:

The given image shows a nasociliary nerve block.

The procedure is done by inserting the needle 1 cm superior and 1.5 cm posterolateral to the medial canthal ligament of the eye.

The nasociliary nerve is a branch of the ophthalmic nerve which in turn is a branch of the trigeminal nerve.

Branches:

• Long ciliary nerve- cornea, ciliary body, iris

• Ramus communicans to the ciliary ganglion

• Posterior ethmoidal nerve- sensory innervation to frontal sinus and posterior ethmoidal air cells

• Anterior ethmoidal nerve- sensory innervation to middle and anterior ethmoidal cells as well as lateral part of the nasal septum

• Infra-trochlear nerve (continuation of the nasociliary nerve)- provides innervation to the conjunctiva, lacrimal sac, caruncle, the medial part of the eyelids

Uses of nasociliary nerve block:

Used in

• Surgery of nasal fractures,

• Septoplasty,

• Rhinoplasty and

• Repair of skin laceration.





The above image represents an anterior ethmoidal nerve block.

Solution to Question 4:

The correct sequence of the auditory pathway is spiral ganglion (2) - cochlear nucleus (1) - superior olivary nucleus (3) - inferior colliculus (4) - medial geniculate body (5).

The spiral ganglion contains bipolar cells whose dendrites form the afferent supply for inner and outer hair cells of the organ of Corti and whose axons form the cochlear division of CN VIII that end in the cochlear nuclei.

The auditory pathway can be remembered with the mnemonic E.C.O.L.I.M.A

Eighth nerve

Cochlear nuclei

Olivary complex

Lateral lemniscus

Inferior colliculus

Medial geniculate body

Auditory cortex



Solution to Question 5:

In the clinical scenario, the history of a child presenting with complaints of inability to understand words with abnormal BERA and normal PTA(Pure Tone Audiometry) is suggestive of auditory neuropathy.

Auditory neuropathy/auditory dyssynchrony is a condition that affects the neural processing of auditory stimuli.

Pathophysiology

• Injury to the synaptic junction between inner hair cells of the cochlea and dendrites of spiral ganglion neurons

• Direct damage to the dendrites of the spiral ganglion neurons

• Direct injury to the spiral ganglion neurons

• Direct axonal damage to the auditory nerve

Auditory neuropathy should be suspected in any child with severe speech and language delay out of proportion with the presumed hearing loss.

All of the following must be present in newborns to diagnose auditory neuropathy:

• Absent or severely abnormal ABR test results at maximal stimulus (100 dBnHL)

• Normal outer hair cell function as determined by OAEs(Otoacoustic emission)

• Absent or elevated stapedial reflex thresholds

In addition, adults with mildly to moderately abnormal pure tone thresholds and speech discrimination scores out of proportion with suspected hearing loss should undergo additional evaluation. Some patients may have normal or slightly elevated audiogram pure tone thresholds

Management includes conventional hearing aids. If a child does not progress with hearing aid devices and shows limited speech discrimination abilities, cochlear implantation is the next viable option.

Other options

Option A:

Michel aplasia: There is a complete absence of bone and a membranous labyrinth. Even the petrous apex is absent but external and middle ears may be completely unaffected. The child presents with congenital sensorineural hearing loss. No hearing aids or cochlear implantation can be used.

Option B:

Malingering(Nonorganic hearing loss)

In this type of hearing loss, there is no organic lesion. Usually, there is a motive to claim some compensation for being exposed to industrial noises, head injury, or ototoxic medication. The patient may present with any of the three clinical situations: Total hearing loss in both ears, total loss in only one ear, or exaggerated loss in one or both ears.

Investigations

Inconsistent results on Repeat Pure Tone and Speech Audiometry Tests: Normally, the results of repeat tests are within +5 dB. A variation greater than 15 dB is diagnostic of nonorganic hearing loss (NOHL).

Absence of Shadow Curve: Normally, a shadow curve can be obtained while testing bone conduction, if the healthy ear is not masked. This is due to the transcranial transmission of sound to the healthy ear. The absence of this curve in a patient complaining of unilateral deafness is

diagnostic of NOHL.

Inconsistency in PTA and SRT: Normally, the pure tone average (PTA) of three speech frequencies (500, 1000, and 2000 Hz) is within 10 dB of the speech reception threshold (SRT). An SRT is better than PTA by more than 10 dB points to NOHL.

Stenger Test: It helps in identifying malingering

Acoustic Reflex: Threshold Normally, the stapedial reflex is elicited at 70-100 dB sensation level. If the patient claims total deafness but the reflex can be elicited, it indicates nonorganic hearing loss (NOHL).

Option C:

Otosclerosis is the disease of the bony labyrinth in which a normal dense endochondral layer of the bony otic capsule is replaced by irregularly laid spongy bone. It has a positive family history with an autosomal dominant trait. It is more commonly seen in females between 20-30 years of age and the hearing loss is aggravated during pregnancy and menopause. It may be associated with osteogenesis imperfecta and a viral infection like measles.

Stapedial otosclerosis is the most common type of otosclerosis. Here the lesion starts just in front of the oval window in an area called fissula ante fenestram.

Patients present with a bilateral conductive type of hearing loss, tinnitus (commonly in cochlear type and active stage), paracusis Willisii (better hearing in noisy surroundings than in a quiet place), vertigo, and a monotonous, well-modulated soft speech.

The tympanic membrane is quite normal and mobile. Eustachian tube function is also normal. Pure tone audiometry shows a dip at 2000 Hz in the bone conduction curve known as the Carhart's notch. Tympanometry shows an As type of tympanogram in which there is reduced compliance at or near the ambient air pressure.

On examination, Schwartz sign/flamingo flush showing a red blush of the tympanic membrane due to vascularity can be seen.

Treatment includes medical, surgical management, and hearing aids. Medical management with sodium fluoride is used in the active stage to speed up the maturity of the active focus and halt further cochlear loss. The treatment of choice is surgical management which includes stapedotomy/stapedectomy with the placement of a prosthesis. Hearing aids are used in patients who refuse or are unfit for the surgery.

Solution to Question 6:

The clinical image showing a polyp and the histopathology showing several sporangia is suggestive of rhinosporidiosis.

Rhinosporidiosis is a chronic granulomatous disease caused by Rhinosporidium seeberi. It is most prevalent in rural districts, among persons bathing in public ponds or working in stagnant water, such as rice fields. Most common in persons in the age range of 15–40; males are more commonly affected than females.

The rhinoscopic examination will reveal papular or nodular, smooth-surfaced lesions that become pedunculated and acquire a papillomatous or proliferative appearance. The lesions are pink, red,

or purple in color. Histopathology shows several sporangia, oval, or round in shape and filled with spores.

The treatment of choice is surgical excision of lesions, with or without cauterization. Not many drugs are effective against this disease. Dapsone has been tried with some success.

Solution to Question 7:

Statements i, ii, and iii are correct.

Statement i: There is 1 row of inner hair cells and 3 rows of outer hair cells is a correct statement.

Statement ii: Inner hair cells are responsible for transmitting auditory impulses is a correct statement. Inner hair cells are richly supplied by afferent cochlear fibres and are probably more important in the transmission of auditory impulses.

Statement iii: The majority of the afferent fibres of spiral ganglion supply the inner hair cells is a correct statement. Ninety-five percent of afferent fibres of spiral ganglion supply the inner hair cells while only five percent supply the outer hair cells. Efferent fibres to the hair cells come from the olivocochlear bundle.

Statement iv: Outer hair cells are important in the modulation of basilar membrane action is an incorrect statement. Outer hair cells mainly receive efferent innervation from the olivary complex and are concerned with modulating the function of inner hair cells.

Solution to Question 8:

In the given clinical scenario, a male patient with symptoms of an upper respiratory tract infection (URTI) exhibits opacity in the right ethmoidal sinus, as shown by the CT image, suggesting a probable diagnosis of sinusitis. CT scans show a typical lenticular rim-enhancing collection adjacent to the lamina papyracea with a fat plane between it and the displaced medial rectus muscle, suggesting a diagnosis of Sinusitis with subperiosteal abscess.

Orbital complications occur most commonly following ethmoid sinusitis as the ethmoid sinus is separated from the orbit by a thin papery bone known as lamina papyracea. Also, the venous supply of orbit and ethmoid sinuses is the same and additionally, these veins are valveless (superior and inferior ophthalmic veins). It can vary in degree and severity and is typically graded with the Chandler classification.

Contrast-enhanced computed tomography (CT) is advised as first-line imaging because of its superiority in demonstrating bony anatomy and pathology of the orbit and sinuses with its speed and ease of examination. MRI provides outstanding soft tissue detail without radiation exposure.

Chandler stages				
Stage 1	Stage 2	Stage 3	Stage 4	Stage 5
Preseptal cellulitis	Postseptal cellulitis or orbital cellulitis without abscess	Subperiosteal abscess	Orbital abscess	Cavernous sinus thrombosis/abscess

Chandler stages				
Inflammation does not extend beyond the orbital septum (where the medial orbital periosteal reflection attaches to the medial eyelid at the tarsal plate).	Inflammation extends into the tissues of the orbit.	There is abscess formation deep to the periosteum of the orbital bones, typically at the lamina papyracea from ethmoid sinusitis.	There is abscess formation within the orbit which has breached the periosteum.	The inflammatory process has extended into the cavernous sinus which thromboses and may progress to abscess formation.

Solution to Question 9:

The malleo-incudal joint is a saddle type.

The type of joint between the middle ear ossicles is a synovial joint. The incudostapedial joint is a ball and socket type.

Note: The image in the question has been updated to include the arrow which was initially missing.

Solution to Question 10:

The given image shows an upsloping/ ascending audiogram in which lower frequencies are affected first like in the early stages of Meniere's disease also known as endolymphatic hydrops.

Meniere's disease (endolymphatic hydrops) is an inner ear disorder where the endolymph distends the endolymphatic system.

Meniere's disease is usually unilateral. It is characterized by episodic vertigo, fluctuating sensorineural hearing loss, tinnitus, and a sense of ear fullness.

Pure tone audiometry reveals a sensorineural type of hearing loss with lower frequencies affected first. It shows a rising type of curve. As the disease progresses, middle and higher frequencies get involved, and the audiogram becomes flat or downsloping type.

The illustration given below shows an audiogram seen in different stages of Meniere's disease.



Other options:

Option A: Ototoxicity causes bilateral high-frequency sensorineural hearing loss. Audiometry shows a downsloping audiogram, typically involving frequencies above 8,000 Hz and progresses to lower speech frequencies. Common drugs causing the same include aminoglycoside antibiotics; loop diuretics (not thiazides); antimalarials; cisplatin; and NSAIDs like ibuprofen, salicylates, and indomethacin.

Option B: Presbycusis is a physiological sensorineural hearing loss occurring in individuals above 65 years of age. Audiometry shows a downsloping graph.

Option C: In noise-induced hearing loss, audiometry shows a typical notch at 4000 Hz. As the duration of noise exposure increases, the notch becomes deeper and wider, involving lower and higher frequencies.



Solution to Question 11:

In the above clinical scenario, a patient presenting with unilateral obstruction of the nasal cavity due to the presence of mass and nasal bleeding along with the histopathology showing several sporangia is most likely suggestive of the diagnosis of rhinosporidiosis.

Rhinosporidiosis is a chronic granulomatous disease caused by Rhinosporidium seeberi. It is most prevalent in rural districts, among persons bathing in public ponds or working in stagnant water, such as rice fields. Most common in persons in the age range of 15–40; males are more commonly affected than females.

The rhinoscopic examination will reveal papular or nodular, smooth-surfaced lesions that become pedunculated and acquire a papillomatous or proliferative appearance. The lesions are pink, red, or purple in color. Histopathology shows several sporangia, oval, or round in shape and filled with spores.

The treatment of choice is surgical excision of lesions, with or without cauterization. Not many drugs are effective against this disease. Dapsone has been tried with some success.

A patient with rhinosporidiosis is shown below.

Other Options:

Option B: Rhinoscleroma is caused by Klebsiella rhinoscleromatis (Frisch bacillus). Histopathology shows large foam cells with a central nucleus and vacuolated cytoplasm containing the causative bacilli, which are Mikulicz cells. There are also homogenous eosinophilic inclusion bodies consisting of the immunoglobulins in the plasma cells called the Russell bodies.



Option C: Inverted papilloma (transitional cell papilloma) is a tumour originating from the nonolfactory mucosa of the nose (Schneiderian membrane) and paranasal sinuses. The unilateral growth of hyperplastic papillomatous tissue into the stroma characterizes this condition. It primarily affects men aged 40-70 and presents unilaterally with symptoms such as nasal obstruction, nasal discharge, and epistaxis. Invasive growth into the sinuses or orbit can lead to proptosis, diplopia, and lacrimation. A biopsy is essential for diagnosis. Histology shows a thin respiratory epithelium overlying the surface and an underlying stroma containing multiple nodules of thick neoplastic epithelium growing inward.

Option D: Juvenile nasopharyngeal angiofibroma is seen almost exclusively in males in the age group of 10–20 years and is very rarely seen in older people and females. Profuse, recurrent, and spontaneous epistaxis is the most common presentation. Diagnosis is mostly clinical. Contrast-enhanced CT scan of the head is the investigation of choice that reveals anterior bowing of the posterior wall of the maxillary sinus (Antral sign/Holman Miller sign) which is pathognomonic of angiofibroma.

Solution to Question 12:

Coughing is a type of forceful expiration. During forceful expiration, when pressure builds up, the glottis opens and then the air comes out. The cough reflex is a protective mechanism of the larynx to expel irritants and keep the airway clear.

Coughing begins with a deep inspiration followed by a forced expiration against a closed glottis. This increases the intrapleural pressure to 100 mm Hg or more. The glottis is then suddenly opened, producing an explosive outflow of air at velocities up to 965 km (600 ml) per hour.

Both chemical and mechanical stimuli can initiate the cough reflex. Chemical or mechanical irritation of the respiratory mucosa stimulates the afferent internal laryngeal nerve, which is the branch of the superior laryngeal nerve of the vagus nerve. These sensory fibres synapse in the nucleus tractus solitarius in the medulla oblongata. Complex neural networks here generate the conscious 'urge to cough' while also sending efferent signals to the various muscles involved in

generating an effective cough which include the adductors of the vocal cord, the diaphragm, the external intercostals and the muscles of the anterior abdominal wall.

Other Options

Statement ii and iii: It is the forceful expiration, not just a normal expiration or deep inspiration which opens the glottis.

Solution to Question 13:

In the given clinical scenario, a female with progressive bilateral hearing loss, in the right ear more than the left, and a stapedotomy of the right ear was planned. Stapedotomy surgery is done for otosclerosis. Otosclerosis results in conductive hearing loss so, the tuning fork test results of the patient will be Bilateral Rinne's negative, Weber's lateralized to the right.

As there is conductive hearing loss in both ears, like that associated with otosclerosis (which may require a stapedotomy), the bilateral Rinne test will be negative. In Weber's test, the sound lateralizes to the affected ear, which, in this case, is the right ear with more significant hearing loss.

Tuning fork tests are performed commonly with tuning forks of frequencies such as 128, 256, 512, 1024, etc. The tuning fork of 512 Hz is considered ideal as forks of lower frequencies produce a sense of bone vibration, while those of higher frequencies have a shorter decay time.

Rinne’s test is done by placing a vibrating tuning fork on the patient’s mastoid and when he stops hearing, it is brought beside the external auditory meatus (EAC). If he still hears the sound, air conduction (AC) is better than bone conduction (BC) i.e. AC>BC. The inferences are as follows:

• Rinne's test is considered positive (normal) when the air conduction AC>BC is seen in normal people and people with sensorineural deafness.

• It is considered negative when the BC>AC, and is seen in patients with conductive deafness.

Weber test is done by placing a vibrating tuning in the middle of the forehead or the vertex and the patient is asked in which ear the sound is heard. The inferences are as follows:

• Normal result is achieved when the sound heard equally in both ears

• In conductive deafness, it is lateralized to the worse ear

• In sensorineural deafness, it is lateralized to the better ear.



Solution to Question 14:

In the given clinical scenario, the patient can only hear when words are shouted, which is a severe hearing loss. The degree of his hearing impairment will be Severe hearing loss > 70%.

Our normal conversational speech is only up to 60 dB. If we have to shout, it means the patient is having a hearing loss greater than 60 dB.

WHO grades of hearing loss:

World Hearing Day 2023 observed on March 3, will highlight the importance of integrating ear and hearing care within primary care, as an essential component of universal health coverage. The theme for this year's campaign is "Ear and hearing care for all, let's make it a reality"

Grade of hearing loss range	Range of hearing loss	Performance	Recommendation
0: No impairment	25 dB or better	No or very slight hearing probl ems. Able to hear whispers	None
1: Slight impairment	26-40 dB	Able to hear and repeat words spoken in normal voice at 1 me ter	Counselling. Hearing aids may be needed
2: Moderate impairment	41-60 dB	Able to hear and repeat words using raised voice at 1 meter	Hearing aids are usually recom mended
3: Severe impairment	61-80 dB	Able to hear some words when shouted into better ear	Hearing aids needed. If no hea ring aids available, lip reading should be taught
4: Profound	81 dB or greater	Unable to hear and understand even a shouted voice whether in quiet or noise.	Hearing aids may help in unde rstanding words. Additional re habilitation is needed. Lip-read ing and sometimes signing ess ential

ENT INI-CET 2024

Question 1:

All of the following are features of otitis media with effusion EXCEPT?

a) Most cases have Ad type of curve on tympanometry and a conductive hearing loss

b) Resolves on its own after the symptoms subside

c) Usually occurs after an episode of common cold, but can occur de-novo

d) It is usually bilateral

Question 2:

Which of the following is typically not seen in allergic rhinitis?

a) Allergic shiners

b) Allergic salute

c) Otto Veraguth folds

d) Dennie-Morgan line

Question 3:

A 13 year old male patient presents with unilateral nasal obstruction and recurrent episodes of epistaxis for past 6 months. What is the most likely diagnosis?

a) Antrochoanal polyp

b) Juvenile nasopharyngeal angiofibroma

c) Coagulation disorder

d) Allergic rhinitis

Question 4:

Which of the following is not recommended in an 8 yr old child with bilateral sensorineural hearing loss?

a) Adenoidectomy and grommet insertion

b) Cochlear implant

c) Hearing aid

d) Sitting in front row of the class

Question 5:

A young male patient presents to ER with complaints of daytime somnolence and repeated awakening at night from sleep and his partner gives history of snoring in bed. All of the following are true except?

a) Apnea results in awakening

b) Apnea is due to fall in oxygen saturation

c) Low oxygen stimulates respiratory effort

d) Pharngeal muscle tone increases

Question 6:

Identify the procedure.



a) Stapedotomy

b) Myringoplasty

c) Tympanoplasty

d) Ossiculoplasty

Question 7:

What is the most commonly used test in screening for neonatal hearing?

a) Conduction audiometry

b) Otoacoustic emissions (OAE)

c) Brainstem evoked response audiometry

d) Clap and smile

Question 8:

Identify the type of hearing loss.



a) Mixed hearing loss

b) Conductive hearing loss

c) Sensory hearing loss

d) Normal

Question 9:

The posterior end of the marked structure is present at the level of which of the following?



a) Eustachian tube

b) Sphenopalatine foramen

c) Odontoid process

d) Sphenoidal sinus

Question 10:

Carhart's notch is seen in _____

a) Otosclerosis

b) Noise-induced hearing loss

c) Meniere's disease

d) Serous otitis media

Question 11:

Which of the following is the most dangerous lesion?

a) U/L Partial RLN lesion

b) B/L Partial RLN lesion

c) U/L Complete RLN lesion

d) B/L Complete RLN lesion

Question 12:

Identify the nasal fracture.



a) Chevallet fracture

b) Jarjaway fracture

c) Simple fracture

d) Le fort fracture

Question 13:

What causes maximum hearing loss?

a) Malleus-incus discontinuity with intact tympanic membrane

b) Malleus-incus discontinuity with perforation of tympanic membrane

c) Partial fixation of the footplate of stapes

d) Otitis media with effusion

Answer Key

Question No.	Correct Option
1	a
2	c

3	b
4	a
5	d
6	a
7	b
8	a
9	a
10	a
11	d
12	a
13	a

Detailed Explanations

Solution to Question 1:

Type B tympanometry curve is obtained in otitis media with effusion.

Serous otitis media also known as secretory otitis media, mucoid otitis media, or glue ear is characterized by the accumulation of nonpurulent, serous effusion in the middle ear cleft.

Glue ear can occur either due to eustachian tube dysfunction (adenoid hyperplasia, chronic sinusitis or tonsillitis or rhinitis, cleft palate, tumors of nasopharynx) or due to increased secretory activity of the middle ear mucosa (allergy, unresolved otitis media, viral infections).

Glue ear is commonly seen in children aged 5-8 years and is associated with insidious onset of hearing loss (not exceeding 40 dB), mild ear aches, and a history of upper respiratory tract infections. Delayed or defective speech may also result from hearing loss.

Otoscopy shows:

• Dull and opaque tympanic membrane with loss of light reflex

• Thin leash of vessels along the handle of the malleus

• Fluid levels and air bubbles can be seen when the tympanic membrane is transparent.

The otoscopy image of serous otitis media is shown below:

Treatment is mainly aimed at the removal of fluid and preventing recurrence.

• Medical management includes decongestants, antihistamines, and antibiotics.

• Surgical management is done if the condition cannot be controlled with medications.
It includes myringotomy and aspiration of fluid with grommet insertion. Adenoidectomy and tonsillectomy may be required and are usually done at the time of myringotomy.

Hearing tests suggestive of conductive hearing loss as seen in serous otitis media:

• Rinne negative (BC>AC).

• Weber - lateralized towards the worse ear

• Audiogram - AC abnormal, BC normal, air-bone gap present

• Impedance audiometry - flat curve (reduced compliance, shift to the negative side)

B-type tympanogram is expected in a case of serous otitis media.

Here, compliance decreases. The fluid accumulation causes decreased mobility of the tympanic membrane leading to a flat or dome-shaped curve, this results in a B-type tympanogram, as shown below:



Note: Tympanograph is Flat when there is Fluid in the middle ear.

Solution to Question 2:

Otto Veraguth folds are not seen in allergic rhinitis.

Otto Veraguth fold is a physical sign of depression wherein there is a triangular fold in the nasal corner of upper eyelid.

Allergic rhinitis is an IgE-mediated immunological response of the nasal mucosa to allergens in the environment.

Patients typically present with recurrent bouts of sneezing, nasal obstruction, watery nasal discharge, and itching.

Clinical signs include:

• Allergic shiners: dark circles around the eyes.

• Allergic salute: black transverse nasal crease on the dorsum of the nose due to repeated upward rubbing.

• Dennie-Morgan lines - transverse skin creases in the infraorbital area.

Allergic rhinitis can be classified into intermittent and persistent, based on the duration of the disease:

• Intermittent- Symptoms are present for less than 4 days a week, or less than 4 weeks

• Persistent- Symptoms are present for more than 4 days a week, or more than 4 weeks

It can be classified as mild, moderate, or severe based on the severity of the disease:

• Mild- None of the following symptoms are present- sleep impairment, impairment of daily activities, school, or work, troublesome symptoms.

• Moderate- severe: If at least one of the symptoms mentioned above are present.

A step-care approach is recommended for allergic rhinitis and its impact on asthma (ARIA) for the treatment of allergic rhinitis includes the following:

• Oral antihistamines or intranasal cromolyn sodium are recommended for mild, intermittent disease. (option A).

• Intranasal corticosteroids are used for moderate disease as monotherapy. (option B).

• Oral antihistamines and intranasal steroids can be used in combination to treat severe diseases. (option C).

• A short course of oral corticosteroids and immunotherapy can be used for severe, persistent disease in case the above intervention fails

In the given image, Otto Veraguth's sign or Veraguth's fold is shown, a diagonal palpebral fold running from the lateral corners of the eyes, medially upward to the medial end of the eyebrows. It is a physical sign of depression.

Another physical sign of depression is the Omega sign. In this, there is vertical wrinkling between the eyebrows joined at the top by a horizontal crease.



Solution to Question 3:

The most likely diagnosis is juvenile nasopharyngeal angiofibroma (JNA), as there is unilateral nasal obstruction and recurrent epistaxis.

Juvenile nasopharyngeal angiofibroma (JNA) is a highly vascular and locally aggressive tumor seen almost exclusively in adolescent males, associated with significant morbidity and occasional mortality.

Patients usually present with:

• Epistaxis (most common presentation): Profuse, recurrent, and spontaneous

• Mass: A mass may be present in either the postnasal space, resulting in progressive nasal obstruction, or nasopharynx, where it may obstruct both choanae. The mass is firm in

consistency, but digital palpation of the same should not be done. If the mass obstructs the eustachian tube, conductive hearing loss and otitis media with effusion may be observed.

• Proptosis, swelling of the cheeks, broadening of nasal bridge, and involvement of 2nd, 3rd, 4th and 6th cranial nerves may occur

JNA can be diagnosed using CT, MRI, and carotid angiography.

Anterior bowing of the posterior wall of the maxillary sinus due to the presence of a mass in the pterygomaxillary space is a characteristic CT finding of JNA. This is known as the Holman-Miller sign or antral sign.

Recurrence has been reported up to 35%. Surgical excision is the treatment of choice. Recurrent angiofibroma is treated with revision surgery or radiotherapy when the tumor is inaccessible.

Solution to Question 4:

Adenoidectomy and grommet insertion is not recommended in bilateral sensorineural hearing loss (SNHL).

Adenoidectomy and grommet insertion is done in the management of serous otitis media.

Serous otitis media is also known as secretory otitis media or mucoid otitis media or glue ear. It is characterized by the accumulation of nonpurulent effusion in the middle ear cleft.

Treatment is mainly aimed at the removal of fluid and preventing recurrence.

• Medical management includes decongestants, antihistamines, antibiotics, and middle ear aeration techniques.

• Surgical management includes myringotomy and aspiration of fluid, grommet insertion, tympanotomy, or cortical mastoidectomy. Adenoidectomy and tonsillectomy may be required and are usually done at the time of myringotomy.

Option B: Cochlear implant is an electronic device which helps in hearing for those with profound sensorineural hearing loss, who cannot benefit from hearing aids. It has two components :

• External component : speech processor and transmitter

• Internal component : receiver / stimulator and electrode array



Sound signals received by the processor is then conducted to the stimulator via the transmitter. It is then decoded and sent to the electrode array which is placed over the scala tympani. Scala tympani then transmits the signal to the spiral ganglion, which in turn stimulates the auditory nerve cells. Thus, sound is conducted to the brain.

Option C: Hearing aids are used in the management of SNHL. Hearing aids are of the following types

• Conventional hearing aids

• Bone-anchored hearing aids (BAHAs)

• Implantable hearing aids (vibrant soundbridge)

Option D: Sitting in the front row of class is recommended in SNHL to help the child hear the class better.

Solution to Question 5:

Pharyngeal muscle tone decreases during snoring.

Pharyngeal muscles are relaxed during sleep and cause partial obstruction. Breathing against obstruction causes vibrations of soft palate, tonsillar pillars and base of tongue producing sound.

Types of snoring:

• Primary snoring: without association with obstructive sleep apnea (OSA). Primary snoring is not associated with excessive daytime sleepiness and has apnea—hypnoea index of less than five.

• Complicated snoring: associated with OSA.

Obstructive sleep apnea-hypopnea syndrome (OSAHS) is diagnosed if the patient has either symptoms of nocturnal breathing disturbances or daytime sleepiness, and ≥5 episodes of obstructive apnea or hypopnea per hour of sleep. Each such episode should last for at least 10 seconds, resulting in ≥3% drop in oxygen saturation or a brain cortical arousal. OSA is diagnosed

by overnight polysomnogram (PSG).

OSAHS is characterized by recurrent episodes of upper airway obstruction during sleep, leading to intermittent hypoxia and sleep fragmentation

Usually, there is a decline in neuromuscular output to pharyngeal dilator muscles during sleep. In predisposed individuals, this results in complete (apnea) or partial (hypopnea) pharyngeal collapse.

Solution to Question 6:

The above image shows a stapedotomy procedure performed in the middle ear.

It can be identified as a stapedotomy procedure by the following features in the image given:

• The footplate of the stapes is not removed

• The prosthesis is placed by creating a hole in the footplate of the stapes

Stapedotomy is the excision of the superstructure of stapes and drilling a hole in the footplate & placing a piston in the hole. It is done to bypass the immobile stapes using a piston which will take over the function of conducting sound.



Contraindications for Stapedotomy:

• Only hearing ear

• Positive Schwartze sign

• Otitis external

• Meniere's disease: can damage the inner ear while drilling a hole in the footplate

• Young children

• Professional athletes, pilots, construction workers, and divers: Sudden changes in pressure can displace the piston and cause vertigo.

• Those who work in noisy surroundings: Surgery makes them more vulnerable to noise-induced hearing loss.

Otosclerosis is a primary disease of the bony labyrinth where foci of irregularly placed spongy bone replace part of the normally dense enchondral layer of the bony otic capsule. Otosclerosis commonly involves the stapes, causing stapes fixation and conductive hearing loss. Clinically, patients present with painless progressive bilateral conductive hearing loss and paracusis willisii (patients hear better in noisy surroundings as other people raise their voices to be heard). Pure tone audiometry would reveal conductive hearing loss with a dip in bone conduction at 2000 Hz (Carhart's notch). The definitive treatment for otosclerosis involves stapedectomy, in which the otosclerotic stapes is removed, and a prosthesis is placed in its place.

The image shown below shows Carhart's notch.



Other options

Option A: Myringoplasty is the term given for the closure of perforation of pars tensa of the tympanic membrane. It is done to check repeated infections from the external ear.

Option B: Tympanoplasty is the surgical reconstruction to correct damage in the middle ear and restore the integrity of the tympanic membrane. It involves ossiculoplasty (ossicular chain reconstruction) with or without myringoplasty.

Option C: Ossiculoplasty eradicates the disease from the middle ear and reconstruction of hearing.

Solution to Question 7:

Otoacoustic emissions (OAE) are used in the universal neonatal hearing screening protocol.

In India, a two-stage screening protocol is followed according to which infants are screened first with otoacoustic emissions (OAE). Infants who fail the OAE are screened with auditory brainstem response (ABR) or Brainstem Evoked Response Audiometry (BERA), which is also the definitive test. All babies kept in NICU must be screened by BERA test as per the protocol.

The first screening is usually conducted at birth or before discharge. Conducting the first screening before discharge is preferred to ensure better coverage and reduce loss to follow-up. A second screening is scheduled anytime between 1–3 weeks or during the next scheduled visit at 6 weeks.

All babies with abnormal BERA should undergo detailed ENT evaluation and hearing-aid fitting (cochlear implants) and auditory rehabilitation must be done before 6 months of age.

Other options

Option A: Conduction audiometry is used in older children and adults to measure hearing thresholds, requiring active cooperation, which is not feasible in neonates.

Option C: Brainstem evoked response audiometry (BERA): While BERA is more precise and can detect auditory pathway issues up to the brainstem, it is more time-consuming and costly. It is typically used for confirmation if OAE results are abnormal.

Option D: Clap and smile is an outdated, subjective test and is not reliable for neonatal hearing screening.

Solution to Question 8:

The given audiogram is suggestive of left-sided mixed hearing loss.

Interpretation of the audiogram:

• Blue colour indicates the left ear.

• > indicates bone conduction (BC), and the threshold is >25 dB. Hence, BC is abnormal, which implies that the abnormality is in the sensorineural pathway.

• × indicates air conduction (AC), and the threshold is >15 dB. Hence, AC is abnormal.

• Both AC and BC are abnormal. It indicates sensorineural hearing loss.

• In addition, the air-bone gap is present, which implies conductive hearing loss.

• Hence, the interpretation is left-mixed hearing loss.

A mixed type of hearing loss presents with defects in both conductive and sensorineural pathways. This type of hearing loss is seen in chronic suppurative otitis media and otosclerosis.

Solution to Question 9:

The marked structure shown is the inferior turbinate. The posterior end of the inferior turbinate is present at the level of the opening of the eustachian tube.

The eustachian tube connects the nasopharynx and the tympanic cavity in the middle ear. It is usually collapsed and opens during activities such as yawning or swallowing. Its function is to maintain the pressure in the middle ear cavity and drain any secretions from the middle ear. It opens approximately 1.25 cm behind the posterior end of the inferior turbinate, in the lateral wall of the nasopharynx.

The inferior turbinate is the largest turbinate located in the lower part of the lateral nasal wall. The Woodruff's plexus is located inferior to the posterior end of the inferior turbinate and is responsible for posterior epistaxis in adults.



Other options:

Option B: Sphenopalatine foramen is situated at the posterior end of the middle turbinate.



Option C: Odontoid process is a protuberance of the Axis (C2 vertebra). Part of the axis is located much lower in the neck and not associated with the nasal cavity.

Option D: Sphenoidal sinus opens into the nasal cavity at the superior meatus.

Solution to Question 10:

Carhart's notch is seen in otosclerosis. Carhart's notch is a dip at 2000 Hz in Bone Conduction, seen in pure tone audiometry.



Pure Tone Audiometry in Otosclerosis:

• Air conduction is poor & worse for lower frequencies.

• Large A-B gap

• Bone conduction is normal.

• In some cases, there is a dip in the bone conduction curve.

• It is different at different frequencies but is maximum at 2000 Hz and is called Carhart's notch.

Note: Boiler's notch is a similar dip in the BC and AC curve but at 4000 Hz, as seen in noise-induced hearing loss.



Solution to Question 11:

Bilateral complete RLN (recurrent laryngeal nerve) lesion is the most dangerous lesion among the given options.

In bilateral RLN paralysis

• Both the cords lie in the median or paramedian position, the airway is inadequate causing dyspnea and stridor. This may lead to death if untreated.

• The voice is good.

• Dyspnea and stridor become worse on exertion or during an attack of acute laryngitis.

All laryngeal muscles are supplied by the ipsilateral recurrent laryngeal nerve (RLN), except cricothyroid which is supplied by the superior laryngeal nerve (SLN). Both SLN and RLN are branches of the vagus nerve.

The recurrent laryngeal nerves (RLN) on the right and the left side follow different courses-

• Right RLN takes origin in the root of the neck, winds around the first part of the subclavian artery, and then ascends in the tracheoesophageal groove.

• Left RLN takes origin in the thorax, winds around the arch of the aorta, and then ascends in the tracheoesophageal groove. Since it has a longer course, it is more likely to be injured.

Solution to Question 12:

A Chevallet fracture usually results from a blow to the nose from below. This fracture line runs vertically. It extends from the anterior nasal spine upwards to the junction of the bony and cartilaginous dorsum of the nose.

Injuries to the nasal septum can result from trauma to the nose from the front, side, or below. The septum may get buckled on itself, fracture vertically and horizontally, or get crushed

A Jarjaway fracture (option B) results from a blow to the nose from the front. The fracture line runs horizontally backwards. It begins just above the anterior nasal spine and extends just above the junction of the septal cartilage with the vomer.



Early recognition and treatment of septal injuries are crucial, including drainage of hematomas.

Dislocated or fractured septal fragments should be repositioned and supported between mucoperichondrial flaps using mattress sutures and nasal packing. Nasal pyramid fractures often coincide with septal fractures and should be treated simultaneously.

Neglecting injuries to the septum can lead to deviations in the cartilaginous nose, as well as asymmetry in the nasal tip, columella, or nostrils.

Other options:

Option C: Simple fractures refer to the fractures of the nasal bone without displacement. Simple fracture describes the nature of the fracture.

Option D: Le Fort fractures are a group of fractures that affect the midface of the skull and collectively involve a partial or complete separation of the midface from the skull.

Solution to Question 13:

Malleus-incus discontinuity with an intact tympanic membrane causes maximum hearing loss.

This condition causes maximum conductive hearing loss, up to 60 dB, as there is no alternative pathway for sound transmission to the cochlea. Even though the tympanic membrane is intact, the disconnection of the ossicular chain severely disrupts sound transmission to the inner ear.

Hearing loss severity depends on the extent to which the normal transmission of sound waves through the middle ear is disrupted.

Average hearing losses seen in different lesions of conductive apparatus

• Complete obstruction of the ear canal: 30 dB

• Perforation of tympanic membrane: 10-40 dB

• Ossicular interruption with intact drum: 54 dB

• Ossicular interruption with perforation: 38 dB

• Malleus fixation: 10-25 dB

• Closure of oval window: 60 dB

Other options

Option B: Malleus-incus discontinuity with perforation of the tympanic membrane: Although this results in conductive hearing loss (38-40 dB), the perforated tympanic membrane allows some direct transmission of sound to the inner ear, slightly reducing the severity compared to an intact tympanic membrane with ossicular discontinuity.

Option C: Partial fixation of the footplate of stapes causes a milder form of conductive hearing loss compared to ossicular discontinuity, typically around 30–40 dB because some sound transmission still occurs through the fixed stapes.

Option D: Otitis media with effusion leads to mild to moderate conductive hearing loss (20–30 dB), as fluid in the middle ear dampens sound vibrations but does not fully block sound transmission.

ENT NEET 2024

Question 1:

Which of the following stain can be used for laryngeal mass seen during laryngoscopic examination?

a) Congo red

b) Silver nitrate

c) H & E

d) Supra vital

Question 2:

An 18-year-old male presents to the emergency department with epistaxis and a nasal mass. What is the investigation of choice to evaluate his condition?

a) Plain CT

b) CECT PNS and nose

c) MRI

d) X-ray

Question 3:

A female patient presents with nausea, vomiting, and a sensation of objects moving around her. She is diagnosed with left vestibular neuritis. Her head impulse test is positive. What is the correct finding?

a) Right saccade with right side movement

b) Left saccade with right side movement

c) Right saccade with left side movement

d) Left saccade with left side movement

Question 4:

Match the following with the nerve supply of external ear:



a) A - Auriculotemporal nerve; B - Greater auricular nerve; C - Facial and Vagus nerve

b) A - Greater auricular nerve; B - Auriculotemporal nerve; C - Facial and Vagus nerve

c) A - Greater auricular nerve; B - Facial and Vagus nerve; C - Auriculotemporal nerve

d) A - Facial and Vagus nerve; B - Greater auricular nerve; C - Auriculotemporal nerve

Question 5:

A female singer presents with hoarseness of voice and difficulty singing in a high pitch. On laryngoscopy, it was noted that one vocal cord is bowed, positioned lower than the other, and the associated muscle is shortened. Which muscle is likely affected in this scenario?

a) Posterior Cricoarytenoid

b) Lateral Cricoarytenoid

c) Interarytenoid

d) Cricothyroid

Question 6:

What is the marked structure of the cochlear implant shown below?



a) Ground electrode

b) Internal magnet

c) Transmitter coil

d) Receiver stimulator antenna

Question 7:

A patient presented with nasal congestion and upon examination, nasal polyps were observed. A coronal CT scan of the patient was taken and is as shown below. Which of the following sinus drainage pathways is likely to be affected?



a) Ethmoid

b) Frontal

c) Maxillary

d) Sphenoid

Question 8:

Which of the following statements is true regarding an esophageal foreign body?



a) Impaction can not cause mediastinitis

b) More commonly seen in adults

c) Most common site of impaction is at level of cricopharyngeus

d) One of the constrictions of the esopahgus is at the level of the right bronchus

Question 9:

In which procedure is the incision showed in the image used?

a) Caldwell Luc's procedure

b) Modified Denker's

c) Open rhinoplasty

d) Midface degloving procedure

Question 10:

The patient complains of repeated episodes of crusting in the nose with occasional bleeding. She had undergone surgery for nasal obstruction four months back. On examination, the following finding is seen. Which surgery has the highest likelihood to result in this complication?



a) FESS

b) Submucosal resection

c) Denker's method

d) Inferior meatotomy

Question 11:

A 40-year-old woman presents with mild, progressive, bilateral hearing loss. A PTA showed the following findings. What is the most likely clinical finding in this patient?



a) Hennebert sign

b) Hitzelberger sign

c) Schwartze sign

d) Bryce's sign

Question 12:

A 70-year-old man presents with unilateral hearing loss. On examination, a small perforation is noted on his left tympanic membrane. What will be the left-sided Rinne's test finding in this patient?

a) Positive

b) False postive

c) Negative

d) Intermediate

Question 13:

A 50-year-old woman presents with hearing loss in the left ear and episodic pulsatile tinnitus. On examination, there was a mass in the middle ear behind the tympanic membrane. A CT performed is as shown below. What is the probable diagnosis?



a) Glomus tympanicum

b) Reparative granuloma

c) Glomus jugulare

d) Acoustic neuroma

Answer Key

Question No.	Correct Option
1	d
2	b
3	c
4	b
5	d
6	a
7	c
8	c
9	c
10	b
11	c
12	c
13	c

Detailed Explanations

Solution to Question 1:

Laryngoscopic image shows growth over the vocal cords; supravital stain is used to detect laryngeal mass seen during laryngoscopic examination.

In the larynx, it is difficult to identify carcinoma in situ, early infiltrating carcinoma, and the exact margins of an established carcinoma. However, it has been found that the use of toluidine blue as a supravital stain during direct laryngoscopy is a reliable adjunct to the early and precise diagnosis of carcinoma of the larynx.

Also, toluidine blue staining is useful in improving the rate of negative resection margins of early glottic cancer (T1a–T2) treated by TLM (transoral oral microsurgery). This technique is called chromoendoscopy, where the interpretation is based on the colour: a dark blue (royal or navy) stain is considered positive, light blue staining is doubtful, and when no color is observed, it is interpreted as a negative stain, as shown below

Solution to Question 2:

History of nasal mass and associated epistaxis in an 18-year-old raises the suspicion of nasopharyngeal angiofibroma. CECT of the PNS and nose can be done to evaluate the patient.

CECT of the PNS and nose is particularly useful for revealing characteristic findings such as the anterior bowing of the posterior wall of the maxillary sinus, known as the Antral sign or Holman-Miller sign (shown in the image below), which is indicative of nasopharyngeal angiofibroma.

Nasopharyngeal angiofibroma is a benign but locally invasive tumor that predominantly affects adolescent males aged 10-20 years. It typically originates from the posterior part of the nasal cavity near the superior margin of the sphenopalatine foramen and is thought to be testosterone-dependent. Clinically, this condition presents with profuse, recurrent spontaneous

epistaxis, progressive nasal obstruction, conductive hearing loss, and otitis media with effusion. It may also cause additional features such as proptosis, cheek swelling, and broadening of the nasal bridge.

Diagnosis is primarily based on clinical presentation, with CECT serving as the preferred imaging technique to confirm the diagnosis and assess the extent of the tumor. Carotid angiography may be performed to evaluate the vascularity of the tumor and its feeding vessels, typically arising from the external carotid artery, and potentially from the internal carotid system in cases of larger tumors with intracranial extension. Biopsy of the mass is generally avoided due to the risk of severe bleeding.

Staging of juvenile nasopharyngeal angiofibroma

Surgery is the treatment of choice. Embolization of the vessels can be done prior to surgery to reduce the risk of bleeding during the surgery. Radiotherapy, chemotherapy, and hormonal therapy can also be considered.

Stage	Spread
IA	Limited to the nose and/or nasopharynx
IB	Extension into ≥ one paranasal sinus
IIA	Minimal extension into the sphenopalatine foramen with minimal part of pterygopalatine fossa
IIB	Full occupation of pterygopalatine fossa with Holman-Miller sign, may with erosion of orbital bones
IIC	Extension into the cheek, temporal fossa, or posterior to pterygoids through the pterygomaxillary fossa
IIIA	Erosion of skull base with minimal intracranial
IIIB	Erosion of skull base with extensive intracranial with or without cavernous sinus

Solution to Question 3:

In the case of left vestibular neuritis, the correct finding on the head impulse test is a right saccade with left-side movement.

The head impulse test is to assess the vestibulo-ocular reflex (VOR). The abnormal head impulse test result is due to the VOR being impaired on the side of the vestibular lesion leading to

saccades. Saccades are rapid, ballistic movements of the eyes that abruptly try to return to the point of fixation/ gaze.

The head impulse test is done by asking the patient to focus on a target. Then, the patient's head is quickly rotated to the left or right. If there is a deficiency in the VOR, the head movement will be followed by a catch-up saccade in the opposite direction. Thus, in left vestibular neuritis, a rightward saccade is seen following a left-side head rotation.

Vestibular neuritis is distinguished from central causes of vertigo by its characteristic findings: a positive head impulse test, absence of skew deviation, and spontaneous unidirectional nystagmus. Central vertigo, on the other hand, presents with a normal head impulse test, vertical skew deviation, bidirectional nystagmus, and additional neurological signs like diplopia, weakness, or dysarthria. In acute vestibular syndrome, especially in older patients with vascular risk factors, evaluation for a possible stroke is warranted.

Vestibular neuritis resolves spontaneously, but glucocorticoids can improve outcomes if administered within 3 days of symptom onset. Antiviral medications are only suggested for Herpes zoster oticus (Ramsay Hunt syndrome). Vestibular suppressant medications can reduce acute symptoms. Patients should be encouraged to resume normal activity as soon as possible. Vestibular rehabilitation therapy can also be tried.



Solution to Question 4:

The correctly matched option regarding the nerve supply of the pinna is option B: A - Greater auricular nerve; B - Auriculotemporal nerve; C - Facial and Vagus nerve.

The nerve supply of the pinna is provided by the following:

• Greater auricular nerve (A) supplies most of the medial surface of the pinna and the posterior part of the lateral surface.

• Lesser occipital nerve supplies the upper part of the medial surface.

• Auriculotemporal nerve (B) supplies the tragal area and crus helix.

• Arnold's nerve (C) i.e. the auricular branch of the vagus nerve (CN X) supplies the concha and the corresponding area on the medial surface.

• Facial nerve (CN VII) (C) which is distributed along with Arnold's nerve also supplies the concha and the retro auricular groove.

The nerve supply has been demonstrated in the image below.





Solution to Question 5:

The affected muscle is likely the cricothyroid muscle, one of the paired intrinsic muscles of the larynx, which is responsible for elongating and tensing the vocal cords to produce higher pitches.

The cricothyroid muscle and the vocalis muscle(the internal part of the thyroarytenoid) are crucial for adjusting the tension of the vocal cords to produce higher pitches. The cricothyroid muscle, in

particular, plays a unique role by elongating and tensing the vocal cords, which is essential for achieving high-pitched phonation. This muscle's function is directly influenced by the external branch of the superior laryngeal nerve, which supplies it and facilitates the adjustment of pitch.

The external branch of the superior laryngeal nerve descends along the larynx, situated beneath the sternothyroid muscle, to innervate the cricothyroid muscle. By activating this muscle, the external branch enables the tension of the vocal cords to be increased, thereby contributing to the production of higher-pitched sounds. This intricate nerve-muscle interaction highlights the specialized role of the cricothyroid muscle in pitch modulation and vocal quality.



Solution to Question 6:

The marked structure in the radiology image of the cochlear implant is the ground electrode.

A cochlear implant is a medical device that uses electricity to stimulate the spiral ganglion cells of the auditory nerve to restore sensorineural hearing loss. It can be used in a patient with bilateral severe sensorineural hearing loss.

It converts sound to an electrical signal (transduction) and delivers this to the hearing nerve, which bypasses the damaged hearing apparatus. It cannot be used in patients with cochlear aplasia or without the VIIIth cranial nerve.

Cochlear implant:

• An electronic device for severe to profound SNHL and who cannot benefit from a hearing aid.

• Principle- surgery to place the electrode array within the scala tympani of the cochlea.

• Indication- bilateral severe to profound deafness

• Prerequisites- intact 8th nerve and higher auditory pathway

• Surgical approach- facial recess (posterior tympanotomy) approach

• The electrode of a cochlear implant is placed in the scala tympani. (Note: In the auditory brainstem implant, the site of the implant is the lateral recess of the 4th ventricle.)

The receiver-stimulator implanted during surgery (under the skin) and the electrode array (implanted in the scala tympani of the cochlea) are part of the internal component of the cochlear implant. The microphone, speech processor and transmitter are part of the external component of the cochlear implant, which remains outside the body.

Cochlear implants replace the non-functional transducer system of hair cells of the cochlea and convert the sound signal to electrical impulses, which directly stimulate the fibres of cranial nerve VIII.

X-ray of bilateral cochlear implant



Cochlear implant



Solution to Question 7:

In this patient with rhinosinusitis and nasal polyps, the CT scan indicates that the maxillary sinus is likely affected, as evidenced by the presence of a polyp in this sinus.

Antrochoanal polyp is a type of non-neoplastic nasal polypoidal masses of edematous nasal or sinus mucosa. It arises from the mucosa of the maxillary antrum near its accessory ostium, comes out of it, and grows in the choana and nasal cavity. It is usually single and unilateral which are more commonly seen in children and young adults.

It has 3 parts namely

• Thin antral stalk,

• Round and globular choanal part, and

• Flat nasal part.

The clinical features include unilateral nasal obstruction, mucoid nasal discharge, and a hyponasal voice.

Anterior rhinoscopy may miss the antrochoanal polyp when it starts to grow posteriorly. A large polyp may protrude from the nasal cavity and may appear pink and congested. It is soft and can be moved with a probe. Posterior rhinoscopy may reveal a globular mass filling the choana and hanging down behind the soft palate.

X-ray (lateral view) of soft tissue nasopharynx, reveals a globular swelling in the postnasal space, it is differentiated from angiofibroma by the presence of a column of air behind the polyp known as Dodd sign or crescent sign.

Diagnosis is made by radiograph of the paranasal sinuses reveals opacity of the involved antrum. Non-contrast CT scan reveals the extent of the polyp. Endoscopic examination reveals the antrochoanal polyp hidden posteriorly in the nasal cavity.

Endoscopic sinus surgery is the treatment of choice. Recurrence is uncommon after the complete removal of the polyp.





Solution to Question 8:

The correct statement with regard to the oesophageal foreign body in the image is that the most common site of impaction is at the level of the cricopharyngeus.

Foreign bodies commonly ingested by children are coins, disk batteries, and small toy items. In adults, it is usually a food bolus (mostly meat) that gets lodged.

The flat surface of a coin is seen on the anteroposterior view (AP view) and the edge is seen on the lateral view. This indicates that it is oriented in the coronal plane, which is seen when it is lodged in the esophagus (and not in the trachea).

Initial symptoms are choking, gagging, and coughing followed by excessive salivation, dysphagia, food refusal, vomiting, or pain in the neck and throat. Respiratory symptoms such as stridor, wheezing, cyanosis, or dyspnea are seen if the esophageal foreign body impinges on the larynx or tracheal wall. Imaging with chest and abdominal X-rays should be performed in AP and lateral positions. Chest CT should always be performed if oesophageal perforation is suspected.

Endoscopic removal of the esophageal foreign body is the procedure of choice. It should be performed

• Immediately in esophageal obstruction, disk batteries, and sharp pointed objects.

• Within 12-24 hours in non-pointed objects, pointed objects in the stomach, and coins in the esophagus.

• Non-urgent removal in objects in stomach >2.5cm in diameter, and disk battery in the stomach (up to 48h)

Other options:

Option A: If a foreign body becomes lodged in the esophagus and causes perforation, it can lead to mediastinitis, a potentially life-threatening infection of the mediastinum.

Option C: Esophageal foreign bodies are more commonly seen in children, although they do occur in adults, particularly in those with underlying esophageal abnormalities (e.g., strictures or motility disorders).

Option D: The esophagus has several natural constrictions, including at the cricopharyngeus (C6 level), at the aortic arch, at the level of the left main bronchus and at the diaphragmatic hiatus.

Solution to Question 9:

The incision shown is used in open rhinoplasty.

This procedure involves a transcolumellar incision, an incision made across the columella to lift the nasal skin, allowing full access to the underlying nasal structures. This approach is used for cosmetic or reconstructive surgery. The open technique allows better visibility of the nasal framework compared to closed rhinoplasty.

Other options:

Option A: The Caldwell-Luc operation, also known as anterior antrostomy, involves opening the maxillary sinus via the canine fossa through a sublabial approach to address various pathologies within the sinus. It's indicated for conditions like chronic maxillary sinusitis, dental cysts, oroantral fistula, neoplasms, and certain fractures. Surgery is contraindicated for patients below 17 years of age.

Option B: The modified Denker's procedure is similar to the Caldwell-Luc procedure but involves more extensive exposure of the maxillary sinus and nasal cavity. It’s used for the removal of larger sinus tumors.

Option D: Midface degloving approach is a facial incision-less approach used for accessing the midface, including the nasal cavity and sinuses.

It’s commonly used in removing large nasal and sinus tumors, and it involves incisions within the oral cavity and nasal vestibule, avoiding external scars. No incision on the nose itself is involved.

Solution to Question 10:

The given clinical vignette of a patient with complaints of repeated episodes of crusting in the nose with occasional bleeding and a history of nasal surgery for nasal obstruction is likely to have undergone correction of her DNS(Deviated Nasal Septum). The image is that of septal perforation; SMR (Submucosal resection) has a higher likelihood of causing the same.

Septal perforation can occur after nasal surgery. Features depend on the size of the perforation. Smaller perforations can present with whistling while the larger perforations will present with epistaxis and crust formation. Therefore, septoplasty has become a more favored treatment of DNS than SMR due to its conservative approach.

Septoplasty is a conservative technique in the surgical correction of septal deformity. The key advantage is that it can be carried out as a daycare procedure, most of the bone and cartilage are retained, and complications are less. The deformed septum may corrected by various techniques like scoring the concave side, shaving, and wedge excision. Sometimes realignment by joining

straight pieces of the septum outside the nose and replacing them can be done which is called extra-corporeal septoplasty.

The indications for an SMR are:

• DNS causing symptoms of nasal obstruction and recurrent headaches.

• DNS causing obstruction of paranasal sinuses and middle ear, resulting in recurrent sinusitis and otitis media.

• Recurrent epistaxis from a septal spur.

• As a part of septorhinoplasty, for cosmetic correction of external nasal deformities.

• As a preliminary step in hypophysectomy or vidian neurectomy.

It is contraindicated in children below 17 years of age, during acute respiratory infection, bleeding diathesis, untreated diabetes, or hypertension.

In this procedure, mucoperichondrial and mucoperiosteal flaps are raised on both sides of the septum. Bony and cartilaginous parts of the septum are excised. This is followed by extensive dissection of the septum removing all deformed bony and cartilaginous parts of the septum and preserving only a caudal and a dorsal strut of cartilage.

There is a greater chance of complications associated with SMR that include bleeding, septal hematoma, septal abscess, perforation, depression of the nasal bridge, retraction of the columella, and persistence of the deviation. The flapping of the nasal septum and toxic shock syndrome following SMR can also occur.

Other options:

Option A: FESS (Functional Endoscopic Sinus Surgery) is a minimally invasive procedure to enlarge nasal drainage pathways and improve sinus ventilation; rarely causes septal perforation.

Option C: Denker Approach is a surgical technique to access the maxillary sinus, pterygopalatine, and infratemporal fossae, used for large sinonasal tumors and extensive sinus pathology.

Option D: Endoscopic Surgical Treatment of Odontogenic Maxillary Sinusitis involves middle or inferior meatotomy; inferior meatotomy is preferred for foreign bodies or sinus floor cysts.

Submucous resection	Septoplasty
Non-conservative procedure with extensive dissection of c artilage	Conservative procedure
Not done in < 18 yrs	Can be done for adults and p ersons < 18 yrs if bilateral an d severe
Killian's incision is used	Freer's incision is used
Mucoperichondrial/periosteal flaps raised on both sides of the septum	Flaps raised on only one side
More Complications	Less complications
Reoperation is difficult	Reoperation is easier

Solution to Question 11:

The clinical scenario describes a 40-year-old female patient with mild, progressive, bilateral hearing loss, and a PTA showing Carhart's notch (dip in bone conduction at 2000 Hz) are all suggestive of Otosclerosis and the classical sign seen here is Schwartze sign.

The presence of a reddish hue on the promontory through the tympanic membrane, known as the Schwartze sign, suggests an active focus.

Otosclerosis is caused by bony overgrowth involving the enchondral layer of the bony otic capsule. It presents with bilateral progressive conductive hearing loss and tinnitus. It has an autosomal dominant inheritance and can worsen with stress and hormonal changes like during pregnancy and menopause.

Stapedial otosclerosis is the most common type of otosclerosis. Here the lesion starts just in front of the oval window in an area called fistula ante fenestrae.

PTA shows loss of air conduction with air-bone gap, more for lower frequencies and Carhart's notch may be seen. Tympanometry shows an As type of tympanogram in which there is reduced compliance at or near the ambient air pressure.

Medical management with sodium fluoride is used in the active stage. Definitive treatment is stapedectomy/stapedotomy with the placement of a prosthesis. Carhart notch usually disappears after a successful stapedectomy.

Other options:

Option A: Hennebert sign is pressure-induced dizziness seen in patients with Superior semicircular canal dehiscence syndrome, congenital syphilis, and Méniére's disease.

Option B: Compression of the facial nerve due to acoustic neuroma will result in hypoesthesia of the posterior wall of the external auditory canal. This is called the Hitzelberger sign.

Option D: The gurgling sound produced on pressing the swelling in the laryngocele is known as Bryce's sign.

Solution to Question 12:

This patient with unilateral hearing loss and a small perforation of his tympanic membrane is likely to have conductive hearing loss in the left ear, which will show a negative Rinne's test in the affected ear.

Tuning fork tests are performed commonly with tuning forks of frequencies such as 128, 256, 512, 1024, etc. The tuning fork of 512 Hz is considered ideal as forks of lower frequencies produce a sense of bone vibration while those of higher frequencies have a shorter decay time. Some examples of commonly done tuning fork tests are as follows:



Hearing loss	Rinne test (Conduction)	Weber test (Lateralized to)
None	Air > Bone	Both ears
Sensorineural	Air > Bone	Normal ear
Conductive	Bone > Air	Affected ear

©MARROW

Rinne's test is done by placing a vibrating tuning fork on the patient's mastoid and when he stops hearing, it is brought beside the external auditory meatus (EAC). If he still hears the sound, air conduction (AC) is better than bone conduction (BC). The inferences are as follows:

• Rinne's test is considered positive (normal) when the air conduction AC>BC is seen in normal people and people with sensorineural deafness.

• It is considered negative when the BC>AC, and is seen in patients with conductive deafness. It suggests a minimum hearing loss of 15-20 dB.

Weber test is done by placing a vibrating tuning in the middle of the forehead or the vertex and the patient is asked in which ear the sound is heard. The inferences are as follows:

• A normal result is achieved when the sound heard equally in both ears

• In conductive deafness, it is lateralized to the worse ear

• In sensorineural deafness, it is lateralized to the better ear

Other options:

Option A: Rinne's test is positive in normal subjects.

Option B: Rinne's test is falsely negative in patients with sudden severe unilateral sensorineural hearing loss.

Option B: There is no such interpretation as intermediate in Rinne's test.

Solution to Question 13:

The given clinical vignette of a woman with hearing loss and pulsatile tinnitus and examination suggesting a mass behind the tympanic membrane and CT image showing a mass in the region of jugular foramen makes Glomus Jugulare the likely diagnosis.

Glomus tumors are the most common benign tumors of the middle ear. The tumors comprise paraganglionic cells and arise from glomus bodies which resemble carotid bodies. Glomus bodies

are present in the dome of the jugular bulb and on the promontory along the course of the tympanic branch of the IX cranial nerve.

They are most commonly seen in females between the ages of 40-50 years. They are very slow-growing, thus it may take years before changes in symptoms are appreciated. It follows the rule of 10s - 10% are familial, 10% are multicentric, and 10% are functional (secrete catecholamines).

Depending on the area from which it arises, glomus tumors are classified into two types:

• Glomus jugulare, arising from the dome of the jugular bulb

• Glomus tympanicum, arising from the promontory of the middle ear

The presentation of glomus tumors is as follows:

• Intratympanic presentation

• Progressive hearing loss

• Pulsatile tinnitus is synchronous with pulse and can be stopped by carotid pressure

• Otoscopy: red reflex through the intact tympanic membrane

• Rising sign when a tumor arises from the floor of the middle ear

• Pulsation or Brown sign: when the ear canal pressure is raised with Siegel's speculum the tumor pulsates and blanches. This reverses when pressure is released.

• Polyp presentation

• Profuse bleeding

• Otorrhea if there is a secondary infection

• Red vascular polyp filling the meatus that bleeds easily

• Cranial nerve palsies

• Dysphagia and hoarseness with unilateral paralysis of the soft palate, and pharynx due to IX and X

• Vocal cord paralysis due to X

• Weakness of the trapezius and sternomastoid muscles XI

• Atrophy of half of the tongue XII

• Audible bruit over the mastoid

They can be diagnosed by CT scan, MRI, four-vessel angiography, brain perfusion and flow studies, embolization, and biopsy. It is treated by surgery, radiation, embolization, or a combination of them. Phlep sign is seen in a CT scan of the glomus jugulare tumor. It is erosion of the normal crest of bone between the carotid canal and jugular foramen.

The following image shows the rising sun sign in the case of glomus jugulare.



Other options:

Option A: Glomus tympanicum is a close differential to the jugular tumor, however, it commonly arises from the promontory of the middle ear rather than the jugular bulb and does not show the Phlep sign and it is also found in relation to the temporal bone on CT.

Option B: Giant cell reparative granuloma accounts for 1–7% of all benign lesions of the jaw. It often arises in the maxilla followed by the mandible and affects children and young adults. It is usually a slow-growing lesion. The fast-growing lesions are rare and despite the innocent histological appearance, have an aggressive behavior mimicking a malignant lesion. The following image shows the CT of the reparative granuloma of the maxilla showing the extension of tumor mass over the pterygoid plates and infratemporal fossa.



Option D: Acoustic neuromas are noncancerous, usually slow-growing tumors that form along the vestibulocochlear nerve. They arise from Schwann cells. Schwannomas can occur on any cranial or peripheral nerve in the body, but in the brain, acoustic neuromas are the most common

schwannomas. The following image depicts the classical ice cream cone appearance of acoustic neuroma on MRI.



Anesthesia AIIMS 2017

Question 1:

A 40 year old female with burns over her abdomen, both limbs and back and presented after 8 hours. Which of the following is the formula for calculation of fluid infusion?

a) 2 ml/kg X %TBSA for the next 16 hours

b) 5 ml/kg X %TSBA for the next 16 hours

c) 4 ml/kg X %TSBA for first 8 hours then 2 ml/kg X %TBSA for next 16 hours

d) 4 ml/kg X%TSBA for the next 16 hours

Question 2:

A 35-year-old woman was brought to the casualty ward with a history of road traffic accident (RTA) and head trauma. Her eye opens to pain, moans, can localize limb towards the pain stimulus on the left hand and away from the pain stimulus on the right hand and both the legs in extended posture. Calculate her GCS.

a) 11

b) 9

c) 7

d) 3

Question 3:

A patient who was on ventilator and being ventilated for past few days suddenly pulls out of the endotracheal tube. What is the next step of management?

a) Re intubate

b) Oxygenate and reintubate

c) Start CPR

d) Watch for spontaneous breaths and reintubate if needed

Question 4:

The time required for preoxygenation before tracheal intubation?

a) 1 min

b) 3 min

c) 5-7 min

d) 5 min

Question 5:

Nerve fibre least susceptible to local anesthetic is:

a) A-delta

b) A- gamma

c) A- beta

d) C

Question 6:

Which of the following can be used as an intravenous induction agent?

a) Lorazepam

b) Bupivacaine

c) Dexmedetomidine

d) Neostigmine

Question 7:

Which of the following is the correct sequence?

a) Assess the patient's response, Activate ERS, check carotid pulse, Start CPR

b) Check carotid pulse, Start CPR, Call for help and activate ERS, Defibrillate

c) Defibrillate, Assess response, Check carotid pulse, Maintain airway

d) Start CPR, Activate ERS, Defibrillate, Check pulse

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	b
5	d
6	a
7	a

Detailed Explanations

Solution to Question 1:

Fluid resuscitation in burns:

According to ATLS 10th edition, 2018, the resuscitation formula used in thermal burns is 2 mL/kg × % TBSA.

First 8 hours: 1 mL/kg × % TBSA (lactated Ringer solution)

Next 16 hours: 1 mL/kg × % TBSA (lactated Ringer solution)

For a patient presenting 8 hours after sustaining thermal burns, the total volume using resuscitation formula should be given in the next 16 hours i.e; 2 mL/kg × % TBSA in the next 16 hours while maintaining target urine output of 0.5 mL/kg/hr for adults.

In adults, urine output should be maintained between 30 and 50 cc/hr to minimize potential over-resuscitation.



Note:

The goal of resuscitation is to maintain the fine balance of adequate perfusion as indicated by urine output. It is important to understand that formulas provide a starting target rate; subsequently, the amount of fluids provided should be adjusted based on a urine output target of 0.5 mL/kg/hr for adults and 1 mL/kg/hr for children weighing less than 30 kg.

Total percentage of the burnt area is calculated clinically using the rule of nines.

Solution to Question 2:

The GCS (Glasgow Coma Scale) of this patient is E2V2M5, which comes to a total score of 9.

E2-Eye opens to pain

V2-Moans

M5-Localizes to pain on the left hand (the score represents the best performance elicited, which would be localizing towards the pain stimulus)

Revised Glasgow coma scale for head injury:

Minimum GCS is 3, maximum GCS is 15.

Solution to Question 3:

About half of self-extubated patients may not require re-intubation. Pre-extubation FiO2, PaO2/FiO2, and ventilator mode, as well as gender, may be useful in determining the likelihood that a self-extubated patient will require re-intubation and also watch for spontaneous breathing.

Solution to Question 4:

Preoxygenation with tidal volume breathing of 3 mins is required before tracheal intubation.

Preoxygenation or denitrogenation is the process of replacing nitrogen in the lungs with oxygen.

This delays the onset of critical hypoxia during the period of apnea by 5-8 minutes.

This lengthened apnea time provides an improved margin of safety while the anesthesiologist secures the airway and resumes ventilation.

Solution to Question 5:

The nerve fibre least susceptible to local anaesthetic is type C. The order of susceptibility of nerve fibres to local anaesthetics is B >A-δ > A-γ > A-β > A-α > C i.e

• B fibres are the most susceptible/least resistant

• C fibres are the least susceptible/most resistant

Factors that increase the sensitivity of nerve fibres are :

• Smaller diameter

• Unmyelinated nerves

• Anatomical and physiological factors

*Note: Although theoretically, C fibres are supposed to be the most susceptible to local anaesthetics, practically it was found to be the least susceptible.

Solution to Question 6:

Lorazepam is a benzodiazepine that can be used for induction of anesthesia at a dose of 0.1mg/kg, although midazolam is preferred among benzodiazepines.

Benzodiazepines are mainly used for sedation and less commonly for induction of anesthesia due to the availability of better induction agents (propofol, etomidate, ketamine, and thiopentone).

Dexmedetomidine is a selective alpha-2 agonist and is used primarily as an adjuvant. So, a better option would be lorazepam.

Solution to Question 7:

BLS sequence:

First step – Assess safety of the surroundings.

Second step - Assess for responsiveness

Third step – Proceed if the patient is unresponsive. Shout for help.

If the patient is unresponsive assess for breathing and carotid pulse simultaneously for no more than 10 seconds.

Anesthesia AIIMS 2018

Question 1:

A patient is undergoing modified radical neck dissection (MRND) for laryngeal malignancy. While ligating the internal jugular vein, his etCO2 dropped from 40 mmHg to 14 mmHg, and the patient developed hypotension along with cardiac arrhythmia. Which of the following is the most likely cause?

a) Sympathetic overactivity

b) Vagal stimulation

c) Venous air embolism

d) Carotid body stimulation

Question 2:

At what level does the spinal cord end in adults?

a) L3-L4

b) L1-L2

c) T12-L1

d) L2-L3

Question 3:

Which of the following is not a contraindication to the use of morphine?

a) Bronchial asthma

b) Renal failure

c) Emphysema

d) Acute myocardial infarction

Question 4:

Cardiac output can be determined using the following except?

a) Thermal dilution technique

b) Capnography

c) Esophageal doppler

d) Echocardiography

Answer Key

Question No.	Correct Option
1	c
2	b
3	d
4	b

Detailed Explanations

Solution to Question 1:

A drastic fall in etCO2 with hypotension during neck surgery is most likely due to venous air embolism (VAE).

In modified radical neck dissection (MRND), a head-up position is given to provide a bloodless surgical field. VAE into severed vessels is likely to occur when the surgical site is above the right atrium and venous pressure at the incision site is below the atmospheric pressure. Pulmonary artery pressure will rise and etCO2 will fall after a venous air embolism.

In sympathetic overactivity and carotid body stimulation, blood pressure will rise.

In vagal stimulation, a drastic drop in etCO2 is unlikely.

Solution to Question 2:

The distal termination of the spinal cord is the lower border of L1 in adults.

Because the spinal cord ends at L1, lower nerve roots course some distance before exiting the intervertebral foramina. These lower spinal nerves form the cauda equina.

Note:

In children, the spinal cord ends at L3 and moves up with age.

Solution to Question 3:

Acute myocardial infarction is not a contraindication for morphine use.

Morphine is the opioid of choice for myocardial infarction as it relieves anxiety, chest pain, and myocardial oxygen demand.

Contraindications to the use of Morphine:

• Bronchial asthma: Histamine release can cause bronchoconstriction and precipitate attacks of asthma.

• Renal failure: Morphine-6-glucuronide (metabolite) can accumulate and lead to prolonged respiratory depression.

• Emphysema: Respiratory depression and decreased response to hypoxia can cause hypercapnia in patients with COPD

• Head injury: Hypercapnia due to respiratory depression can increase ICP. Miosis, vomiting, and effect on wakefulness make interpreting the signs of head injury difficult.

• Biliary colic: Constriction of sphincter of Oddi can worsen biliary spasm

• Hypotension and shock

Solution to Question 4:

Capnography is a measurement of end-tidal CO_2, which is used to confirm tracheal intubation and assess ventilation in the perioperative period.

Cardiac output measurements technique:

• Thermodilution

• Dye dilution, using indocyanine green

• Lithium dilution

• Pulse contour devices

• Esophageal doppler

• Thoracic bioimpedance

• Echocardiography

Note: Thermal dilution method uses a non-toxic liquid of temperature different from the blood. This fluid is injected into a vein and is detected on the arterial side using a catheter with a temperature sensor. The rate of flow of this liquid is used to calculate the cardiac output.

Anesthesia AIIMS 2019 (May & Nov)

Question 1:

Which is not an obvious advantage of high pressure oxygen through nasal cannula:

a) Decreases need of intubation

b) Reduces nasopharyngeal dead space

c) Heating and humidification

d) PEEP

Question 2:

Duration of action of Proparacaine:

a) 2 min

b) 5 min

c) 40 min

d) 20 min

Question 3:

Glycopyrrolate is used as a pre-anesthetic agent for:

a) Reducing the secretions

b) Sedative effect

c) Skeletal muscle relaxant

d) Anxiolytic

Question 4:

Identify the device shown in the image below.

a) Automated External Desynchonizer

b) Automated External Defibrillator

c) Automated Electronic Defibrillator

d) Automated External decompressor

Question 5:

You are an intern posted in the ICU. While taking a blood sample for ABG, which of the following steps would you not do?

a) Normal ph can't rule out acid base disorder

b) Add 0.5 ml Heparin to the syringe before taking blood

c) If modified Allens test is negative, alternate site is chosen

d) Radial artery is the preferred site

Question 6:

Which of the following is not a risk factor for postoperative pulmonary complications?

a) BMI > 30

b) Well controlled asthma

c) Upper abdominal surgery

d) Age > 70 years

Question 7:

Which of the following is associated with respiratory alkalosis?

a) Assisted control mode ventilation

b) Controlled mandatory

c) Pressure controlled

d) SIMV

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	b
5	b
6	b
7	a

Detailed Explanations

Solution to Question 1:

High flow nasal cannula (HFNC), which delivers oxygen at high flow, and pressure through the nasal cannula.

Advantages of high-flow nasal cannula:

• Soft and pliable nasal prongs provide comfort to the patient.

• Warming and humidification of secretions: The heating of air to core temperature and humidification of the inspired air is better at high-flow. The humidification causes increased water content in the secretions, which facilitates the removal of secretions. It also decreases the work of breathing and avoids airway dessication and epithelial injury.

• Reduces dead space: The high flow of the oxygen delivered clears the upper airway of the expired air, which reduces dead space by a decrease of rebreathing and allows a higher fraction of minute ventilation to participate in alveolar gas exchange.

• Continuous positive airway pressure (CPAP): HFNC increases the positive end-expiratory pressure (PEEP), and reduces the work of breathing.

• High flow rate causes decreased entrainment of room air, allowing more accurate delivery of oxygen.

Solution to Question 2:

The duration of action of proparacaine is approximately 10-20 minutes.

Proparacaine hydrochloride ophthalmic solution is used as a topical anesthetic agent.

Indications: For procedures which require corneal anesthesia of short duration such as:

• Tonometry

• Gonioscopy

• Removal of corneal foreign bodies

• For short corneal and conjunctival procedures

Contraindication: Patients with known hypersensitivity

Solution to Question 3:

Glycopyrrolate is used as a pre-anesthetic agent for reducing the secretions.

Anticholinergic drugs e.g. atropine, glycopyrrolate are routinely administered with anticholinesterases in order to attenuate the undesirable muscarinic effects. Glycopyrrolate is usually preferred because it has less vagolytic effects than atropine at doses that inhibit secretions and does not cross the blood-brain barrier. It should be administered as early as possible to maximize its effectiveness.

Glycopyrrolate 0.2 mg IV, is administered to dry airway secretions before flexible scope intubation of the trachea, during the bronchoscopic examination.

Solution to Question 4:

The device is an Automated External Defibrillator (AED).

This is a special medical device that is commonly used for recognizing the absence/presence of rapid ventricular tachycardia or ventricular fibrillation. With no need for intervention by an operator, it determines whether an electrical shock right to the heart must be performed (defibrillation).

An automated external defibrillator can be used on those victims, who are unconscious and aren't breathing in a normal way. As a rule, it is used together with CPR in cases of sudden cardiac arrest. It is forbidden to use it on patients who are less than eight years old or those weighing less than 25 kgs.

Solution to Question 5:

0.5 ml of Heparin is not added to the syringe before taking blood.

Very little heparin is needed in the sample to prevent clotting; 0.05 to 0.10 ml of a dilute solution will help prevent coagulation of 1 ml of blood without affecting its pH. This is achieved by drawing 0.5 ml into the syringe and flushing it out. The residue that remains adherent to the syringe is sufficient for the procedure.

Heparin must be added to the syringe as an anticoagulant. Because the pH of heparin is near 7.0, and the P02 and Pco2 of the heparin solution are near room air values, excess heparin can alter ABG measurements. After flushing the syringe with heparin, a sufficient amount usually remains in the dead space of the syringe and needle for anticoagulation without distortion of the ABG determination.

Note:

• Modified Allen's test positive indicates that the collateral blood flow is normal

• Modified Allen's test negative indicates that the collateral blood flow is abnormal and an alternate site must be chosen.



Solution to Question 6:

Well-controlled asthma is not a risk factor for postoperative pulmonary complications.

Risk factors for postoperative pulmonary complications:

• Age >70 years

• COPD (uncontrolled)

• Neck, upper abdominal, Aortic, neurologic surgical procedures

- Prolonged procedure >2 hours
- General anesthesia with intubation
- BMI > 30
- Albumin concentration < 35 g/L
- Inability to walk two blocks on level land or one flight of stairs

Solution to Question 7:

Respiratory alkalosis can occur in Assist Control Mode Ventilation

Ventilatory Mode	ACMV(Assist Control Mode Ventilation)	IMV(Intermittent Mandatory Ventilation)	PSV(Pressure Support Ventilation)	NIV(Noninvasive Ventilation)
Variables set by User (Independent)	Tidal volumeVentilator rateFio2PEEP levelPressure limit	Tidal volumeMandatory ventilator rateFio2PEEP levelPressure limitSpontaneous breaths between assisted breaths	Inspiratory pressure levelFio2PEEPPressure limit	Inspiratory and expiratory pressure levelFio2
Variables Monitored by the user (Dependent)	Peak, mean, and plateau airway pressuresVEABGI/E ratio	Peak, mean, and plateau airway pressuresVEABGI/E ratio	Tidal volumeRespiratory rateVEABG	Tidal volumeRespiratory rateVEABG
Trigger cycle limit	Patient effortTimer Pressure limit	Patient effortTimer Pressure limit	Pressure limitInspiratory flow	Pressure limitInspiratory flow
Advantages	Patient controlGuaranteed ventilation	Patient controlComfort from spontaneous breathsGuaranteed ventilation	Patient controlComfortAssures synchrony	Patient control
Disadvantages	Potential hyperventilation(Leading to respiratory alkalosis) Barotrauma and volutraumaEvery effective breath generates a ventilator volume	Potential dysynchronyPotential hypoventilation	No timer backupPotential hypoventilation	Mask interface may cause discomfort and facial bruisingLeaks are commonHypoventilation

Anesthesia AIIMS 2020 (May and Nov INI-CET)

Question 1:

A patient of blunt chest trauma has been started on mechanical ventilation. Which of the following is not correct about the initial setting?

a) Respiratory rate - 12 Breaths/ min

b) PEEP - 5cm H20

c) FiO2- 100%

d) Tidal volume - 12ml/kg of actual BW

Question 2:

Glycopyrrolate is used as a pre-anesthetic agent for:

a) Reducing the secretions

b) Sedative effect

c) Skeletal muscle relaxant

d) Anxiolytic

Question 3:

Cardiac output can be determined using the following except?

a) Thermal dilution technique

b) Capnography

c) Esophageal doppler

d) Echocardiography

Question 4:

Identify the equipment shown in the image below.

a) Mechanical ventilator

b) Infusion pump

c) CPAP machine

d) High flow nasal cannula

Question 5:

The blue pipeline in the operation theatre is used for delivering which of the following gases?

a) Oxygen

b) Nitrous oxide

c) Air

d) Vacuum

Question 6:

The given set of instruments are used in:

a) Central line insertion

b) Arterial line insertion

c) Airway management

d) Umbilical catheter insertion

Question 7:

On your morning walk in the park, you notice an unconscious man lying on the ground. Which of the following options would be the next best step?

a) Check carotid pulse, start CPR, and call for help; Defibrillate if no pulse.

b) Call for help, check pulse, do CPR, defibrillate if no pulse

c) Observe

d) Ignore and continue to walk

Question 8:

Which of the following can be considered as effective CPR?

a) 1, 2, 3, and 4

b) 1, 2, and 4

c) 2, 3, and 4

d) 1 and 2 only

Answer Key

Question No.	Correct Option
1	d
2	a
3	b
4	d
5	b
6	c
7	b
8	c

Detailed Explanations

Solution to Question 1:

The tidal volume of 12ml/kg of actual body weight is an incorrect initial setting for mechanical ventilation.

Initial ventilatory settings are as given below:

• Tidal volume - 4-6 ml/kg of IBW (Ideal Body Weight)

• Respiratory rate - 12 Breaths/ min

• FiO2 - 100% to begin with and titrated to lesser values to prevent oxygen toxicity

• PEEP - Begin with 5 cm H2O, titrate to 10 cm H2O (Carefully increased as high PEEP may impair ventricular filling in a patient with blunt chest trauma)

Solution to Question 2:

Glycopyrrolate is used as a pre-anesthetic agent for reducing the secretions.

Anticholinergic drugs e.g. atropine, glycopyrrolate are routinely administered with anticholinesterases in order to attenuate the undesirable muscarinic effects. Glycopyrrolate is usually preferred because it has less vagolytic effects than atropine at doses that inhibit secretions and does not cross the blood-brain barrier. It should be administered as early as possible to maximize its effectiveness.

Glycopyrrolate 0.2 mg IV, is administered to dry airway secretions before flexible scope intubation of the trachea, during the bronchoscopic examination.

Solution to Question 3:

Capnography is a measurement of end-tidal CO_2, which is used to confirm tracheal intubation and assess ventilation in the perioperative period.

Cardiac output measurements technique:

• Thermodilution

• Dye dilution, using indocyanine green

• Lithium dilution

• Pulse contour devices

• Esophageal doppler

• Thoracic bioimpedance

• Echocardiography

Note: Thermal dilution method uses a non-toxic liquid of temperature different from the blood. This fluid is injected into a vein and is detected on the arterial side using a catheter with a temperature sensor. The rate of flow of this liquid is used to calculate the cardiac output.

Solution to Question 4:

The equipment shown in the image is high flow nasal cannula.

High flow nasal cannula is a mode of respiratory support used in the management of acute respiratory failure. It delivers humidified gas via the nasal cannula at high flow rates.

• HFNC can deliver oxygen at 40L/min, 37°C, and 99.9% relative humidity.

• The high-flow oxygen is delivered directly to the nasopharynx. This produces FiO_2 equal to that delivered by traditional mask devices.

• The benefits of HFNC include reduced work of breathing, improved gas exchange by nasopharyngeal dead-space washout, and decreased energy required to humidify and heat respiratory gases.

The HFNC equipment shown in the image below uses a heated humidifier (named PMH200), a nasal cannula (no. 2), and a ventilator (brand: vivo30).



Other options:

Option a: Mechanical ventilator

Option b: Infusion Pump

Option c: CPAP

Solution to Question 5:

The blue pipeline in operation theatre is for nitrous oxide.

Other options:

Oxygen: White

Air: Black

Vacuum/Central suction: Yellow

Solution to Question 6:

The image shows a tray in an airway cart with the instruments necessary for airway management.

An airway cart may include:

• Laryngoscope blades of various sizes, including a rigid fibreoptic.

• Tracheal tubes of various sizes, and tracheal tube guides such as semirigid stylets, ventilating tube changer, light wands, and forceps that are required to manipulate the distal portion of the tracheal tube.

• Masks of various sizes.

• Laryngeal mask airways of various sizes.

• Flexible fiberoptic intubation equipment.

• Retrograde intubation equipment.

• At least one device to perform emergency noninvasive airway ventilation such as esophageal tracheal tube, a hollow jet ventilation stylet.

• Equipment required for emergency invasive airway access.

• An exhaled CO_2 detector.

Solution to Question 7:

The next best step is to Call for help, check pulse, do CPR, defibrillate If no pulse.

Steps in basic life support:

First step – Assess safety of the surroundings.

Second step - Assess for responsiveness

Third step – Proceed if the patient is unresponsive. Shout for help.

If the patient is unresponsive, assess for breathing and carotid pulse simultaneously for no more than 10 seconds.

In the given scenario, it is already mentioned that you are in a park (scene safety has been assessed), and that the person is unconscious (responsiveness has been assessed). So, the immediate next step is to call for help, and start CPR.

Solution to Question 8:

The ideal rescue breath rate for adults is 10 breaths/min, while in infants and children up to puberty it is 20-30 breaths/min. As the person's age group is not specified in the question, assuming an adult, statement 1 is incorrect, while 2, 3, and 4 are correct.

Effective chest compression in adults:

• Provide 100-120 compressions per minute (30 compressions for every 15-18 sec).

• Place your arms, one over the other in the midline, on the lower 1/3rd of the patient's sternum, and lock your arms.

• Use both arms to press the chest to a depth of 5-6 cm (2 - 2.4 inches) or more.

• Press hard and fast

• Allow for full chest recoil after every compression.

Effective chest compressions in a child older than 1 year and up to puberty:

• The compression rate is 100 - 120/min (30 compressions for every 15-18 sec).

• One or both palms are placed in the midline.

• The depth of compression is 2 inches.

• Press hard and fast

• Allow for full chest recoil after every compression.

Effective chest compressions in infants (<1 year):

• 100 to 120 compressions per minute (30 compressions for every 15-18 sec).

• One provider: Place two fingers, one between the nipple line and the other 1 cm below.

• Two providers: Encircle the infant's torso with both hands with both thumbs pointing cephalic positioned 1cm below the nipples over the sternum.

• Depth of compressions should be at least 1.5 inches or 1/3 the depth of the infant’s chest.

• Press hard and fast

• Allow for full chest recoil after every compression.

Anesthesia NEET 2018

Question 1:

What is the mechanism of action of the curare group of muscle relaxants?

a) Persistently depolarizing at neuromuscular junction

b) Competitively blocking the binding of ACh to its receptors

c) Repetitive stimulation of ACh receptors on muscle endplate

d) Inhibiting the calcium channels on presynaptic membrane

Question 2:

Which of the following nerves is popularly used in clinical anaesthesia to monitor neuromuscular blockade?

a) Ulnar Nerve

b) Median Nerve

c) Radial nerve

d) Mandibular nerve

Question 3:

Identify the Mallampatti class of this patient.



a) Class II

b) Class I

c) Class IV

d) Class III

Question 4:

In which of the following devices is Murphy's eye seen?

a) Nasopharyngeal airway

b) Laryngoscope

c) Bronchoscope

d) Endotracheal tube

Question 5:

What is the most effective circuit for anaesthesia under spontaneous breathing?

a) Mapleson A

b) Mapleson B

c) Mapleson C

d) Mapleson D

Question 6:

The best type of anesthesia for pre eclampsia is:

a) Spinal anesthesia

b) Epidural anesthesia

c) General anesthesia

d) Caudal block

Question 7:

What alternative drug would you use for cardiac arrest if epinephrine is not effective?

a) Vasopressin

b) Atropine

c) Amiodarone

d) Adenosine

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	d
5	a
6	b
7	c

Detailed Explanations

Solution to Question 1:

The curare group of drugs includes non-depolarizing neuromuscular blocking agents that act by competitively blocking the binding of acetylcholine to its receptors.

This leads to a decrease in the effect of acetylcholine and thus, neuronal transmission to the muscle does not occur.

Solution to Question 2:

Ulnar nerve stimulation at wrist is popularly used to assess the adequacy of neuromuscular blockade.

Neuromuscular blockade is measured using a peripheral nerve stimulator.

Nerve	Response
Ulnar nerve	Contraction of adductor polli cis

Nerve	Response
Facial nerve	Contraction of orbicularis oc uli
Posterior tibial nerve	Contraction of flexor hallucis brevis

Solution to Question 3:

The image shows Mallampatti class II airway.

The modified Mallampatti classification is used in the assessment of airway. It is a predictor of the difficulty of intubation or difficult airway.

The assessment is performed with the patient sitting up straight, mouth open and tongue maximally protruded, without saying "ahh."

Modified Mallampati classification:

• Class 0: Ability to see any part of the epiglottis.

• Class I: Soft palate, fauces, uvula visible.

• Class II: Soft palate and fauces seen, tip of uvula not visible.

• Class III: Soft palate, base of uvula visible.

• Class IV: Soft palate not visible at all. Only hard palate visible.



Classes 3 and 4 are difficult intubation situations.

Initially, the Mallampatti scoring system (1985) had only 3 classes. This was modified by Samsoon and Young to include class 4. Later class 0 was also added.

Solution to Question 4:

Murphy's eye is seen in the endotracheal tube (ETT).

Murphy's eye refers to the distal opening in the side wall of the ETT.

If the distal end of the lumen of the tube is blocked by secretions, Murphy's eye serves to provide an alternate route for ventilation.



Solution to Question 5:

The most effective circuit for anesthesia under spontaneous breathing is Mapleson A or Magill's circuit.

When the Mapleson A circuit is used for spontaneous ventilation, the fresh gas flow rate of the circuit is closest to the physiological minute ventilation of the patient. Hence it is preferred over other circuits.

Solution to Question 6:

Continuous epidural anesthesia is the first choice for most patients with preeclampsia during labor, vaginal delivery, and cesarean section.

Pre-eclampsia patients have a risk of severe airway edema, which makes intubation difficult. Thus continuous epidural anesthesia is advantageous as it avoids the need for intubation. Besides, continuous epidural anesthesia can improve uteroplacental perfusion and also decrease catecholamine secretion.

But one major prerequisite for regional anesthesia in preeclamptic patients is normal platelet count and normal coagulation. Coagulopathy is a contraindication for regional anesthesia.

Minimal platelet count for regional anesthesia is 100,000/µL, but some cases can be taken up with a platelet count as low as 70,000/µL.

General anesthesia is indicated in patients with coagulopathy and in severe pre-eclampsia with complications.

Spinal anesthesia can be given in mild pre-eclampsia but hypotension associated with spinal anesthesia limits its use in moderate to severe pre-eclampsia.

Since the severity of pre-eclampsia or the type of delivery (vaginal or cesarean) is not mentioned, epidural anesthesia has an advantage over other techniques for above-mentioned reasons. Hence the best option here would be epidural anesthesia.

Solution to Question 7:

Epinephrine and amiodarone are the two main drugs approved by the AHA for the resuscitation of a cardiac arrest patient.

The first dose of epinephrine 1 mg IV is administered after the 2nd defibrillator shock is delivered and it is continued every 3-5 minutes.

If this is not effective, 300 mg IV bolus dose of amiodarone can be given and another dose of 150 mg given if necessary

Anesthesia NEET 2019

Question 1:

What is the mechanism of action of the curare group of muscle relaxants?

a) Persistently depolarizing at neuromuscular junction

b) Competitively blocking the binding of ACh to its receptors

c) Repetitive stimulation of ACh receptors on muscle endplate

d) Inhibiting the calcium channels on presynaptic membrane

Question 2:

Which among the following IV anaesthetics does not cause myocardial depression?

a) Etomidate

b) Propofol

c) Thiopentone

d) Ketamine

Question 3:

Which of the following is not used for induction of anaesthesia in pediatric patients?

a) Halothane

b) Sevoflurane

c) Desflurane

d) Nitrous oxide

Question 4:

Which of the following agents does not elicit pain on injection?

a) Ketamine

b) Propofol

c) Etomidate

d) Thiopentone

Question 5:

Which among the following is most commonly used to monitor the depth of anaesthesia?

a) EEG

b) Provoked lower oesophageal contractility

c) Entropy

d) Bispectral index

Question 6:

Which of the following is measured by the device shown in the given image?

a) Oxygen saturation

b) Oxygen content of the blood

c) Partial pressure of oxygen

d) Amount of inspired oxygen

Answer Key

Question No.	Correct Option
1	b
2	a
3	c
4	a
5	d
6	a

Detailed Explanations

Solution to Question 1:

The curare group of drugs includes non-depolarizing neuromuscular blocking agents that act by competitively blocking the binding of acetylcholine to its receptors.

This leads to a decrease in the effect of acetylcholine and thus, neuronal transmission to the muscle does not occur.

Solution to Question 2:

Etomidate, a short-acting I.V. anesthetic agent doesn't cause myocardial depression.

Etomidate is the most cardiac stable intravenous agent due to its lack of effect on the sympathetic nervous system and on the function of the baroreceptor. Hence, etomidate is the I.V. agent of choice in patients with coronary artery disease, cardiomyopathy, and cerebral vascular disease.

Side effects of Etomidate:

• The highest incidence of nausea and vomiting among inducing agents

• Superficial thrombophlebitis and pain on injection

• Adrenocortical suppression

Note: Although Ketamine has an indirect sympathetic stimulatory effect, it possesses a direct negative inotropic effect on the myocardium. Hence, ketamine has to be used cautiously in critically ill patients with shock, as they are prone to have a depletion of endogenous catecholamines. Ketamine infusion in such patients can unmask its direct negative inotropic effects and further decrease the blood pressure and cardiac output.

Hence, for this MCQ, Etomidate is the best answer.

Solution to Question 3:

Desflurane is a highly pungent gas and can produce laryngospasm if used for inhalational induction. Hence it is not used as an induction agent.

The incidence of laryngospasm following desflurane induction is almost 50%. It is usually used for maintenance of anaesthesia only and is of particular value in obese patients, neurosurgical and spinal fusion procedures where rapid awakening is required for neurological function assessment.

Option A: Halothane is hepatotoxic but can be used (Sevoflurane is preferred)

Option B: Sevoflurane is the preferred inhalational agent for induction in the pediatric age group. Cardiovascular depression, bradycardia, and arrhythmias are less frequent with sevoflurane than with halothane and hence the former is preferred.

Option D: Nitrous oxide has the highest minimum alveolar concentration and so it is the least potent. It is used in combination with other potent inhaled anaesthetics to reduce their dose. It can be used to hasten the induction of anaesthesia along with other induction agents.

Solution to Question 4:

Ketamine does not elicit pain on injection, instead produces profound analgesia.

Propofol is formulated as a lipid emulsion, which causes pain on injection. A new aqueous prodrug, fospropofol is associated with less pain on injection.

Etomidate, although it causes pain on injection, the lipid formulation of etomidate is associated with a much less frequent incidence of pain.

Thiopentone at a concentration greater than 2.5% can produce pain on injection and venous thrombosis.

Solution to Question 5:

The bispectral index is the most commonly used method to monitor the depth of anesthesia.

Note: Middle latency auditory evoked potential is the most sensitive indicator to evaluate the depth of anesthesia.

Solution to Question 6:

The given image depicts a pulse oximeter. Pulse oximetry is used for monitoring pulse rate and oxygen saturation of a patient.

A small sensor is placed on the finger, toe, earlobe, or another convenient place. The relative absorption of light by oxyhemoglobin (HbO_2) and deoxyhemoglobin is assessed by measuring the amount of red and infrared light emerging from tissues traversed by light rays and processed by the device, producing an oxygen saturation level.

Pulse oximetry does not measure the absolute content or the partial pressures of oxygen or carbon dioxide.

A pulse oximeter is not always reliable as it reads:

• Falsely high - carboxyhemoglobinemia

• Falsely low - severe anemia, low cardiac output states, hypovolemia, increased systemic vascular resistance

• 80% to 85%, irrespective of the arterial oxygen saturation - methemoglobinemia

Anesthesia NEET 2020

Question 1:

Which of the following investigations provides the most accurate prognostic information with respect to predicting risks of perioperative cardiac complications?

a) Exercise ECG testing

b) Dobutamine stress echocardiography

c) Myocardial perfusion scintigraphy

d) Coronary angiography

Question 2:

A capnograph obtained from an intubated patient undergoing controlled ventilation is given below. Which of the following is a likely cause of the finding seen?



a) Esophageal intubation

b) Inspiration with cardiac oscillations

c) Bronchospasm

d) Spontaneous respiratory effort

Question 3:

A 50-year-old patient with a history of hypertension and diabetes mellitus developed the following rhythm. His blood pressure was 90/40 mm Hg and he had a feeble pulse. What is the next step of management?



a) IV adenosine

b) IV diltiazem

c) DC cardioversion

d) Ibutilide

Question 4:

A patient is given a nicotinic receptor antagonist as a muscle relaxant. Which drug is given postoperatively to recover from muscle weakness?

a) Physostigmine

b) Neostigmine

c) Carbachol

d) Succinylcholine

Answer Key

Question No.	Correct Option
1	b

2	d
3	c
4	b

Detailed Explanations

Solution to Question 1:

Among the given investigations, dobutamine stress echocardiography provides the most accurate prognostic information with respect to predicting risks of perioperative cardiac complications.

Preoperative cardiac stress testing can help identify the presence and severity of preexisting CAD. The prognostic performance of these tests is better assessed based on positive and negative likelihood ratios. Among all the given options, dobutamine stress echocardiography has the maximum positive likelihood ratio and hence provides the most accurate prognostic information.

Even though coronary angiography is considered the gold standard, it may not absolutely risk-stratify patients because many perioperative cardiac events may be related to noncritical coronary lesions.

Solution to Question 2:

The curare notch in the plateau phase of the capnograph indicates that the patient is trying to breathe spontaneously. The patient's inspiratory effort causes some fresh gas to be sucked in the capnometer, resulting in a lower carbon dioxide concentration and a notch.

Image below shows a capnograph with clefts during phase III which indicates spontaneous breathing efforts by the patient during controlled mechanical ventilation. It is also known as a curare notch or cleft.



Solution to Question 3:

The clinical scenario suggests supraventricular tachycardia with hemodynamic instability and the next step of management is DC cardioversion (50-100J).

If the patient was hemodyamically stable, initial treatment would be vagal maneuvers (Valsalva maneuver) in awake patients. This is because vagal maneuvers alone can terminate approximately 20% to 25% of reentry supraventricular tachycardias (SVTs).

If this is unsuccessful, then iv adenosine 6 mg is given. If necessary another dose of adenosine 12 mg may be administered after 1-2 minutes.

If the drug is administered via a central venous catheter, then these doses are reduced to 3 mg and 6 mg respectively.

Solution to Question 4:

Neostigmine is given postoperatively to recover from muscle weakness.

Non-depolarizing muscle relaxants (NDMRs) act mainly by the competitive antagonism of nicotinic (Nm) receptors. Neostigmine inhibits acetylcholinesterase and inhibits the degradation of acetylcholine. The increased accumulation of undegraded acetylcholine can effectively compete with NDMRs, and thereby displace them from the site and antagonize their effects.

Anesthesia INI-CET 2021

Question 1:

Ventilatory support is required if SpO2 becomes less than 90% because

a) Tissues become vulnerable for hypoxia below 90%.

b) Peripheral chemoreceptors becomes insensitive to hypoxia below 90%.

c) Central chemoreceptors becomes insensitive to hypoxia below 90%.

d) Patient capacity to generate adequate tidal volume decreases below 90%.

Question 2:

In conventional oxygen therapy, which device will deliver highest FiO2?

a) Venturi mask

b) Hudson mask

c) Non rebreathing mask

d) Nasal cannula

Question 3:

What is the nerve blocked in the following local anesthesia technique?

a) Nasociliary nerve

b) Anterior ethmoidal nerve

c) Sphenopalatine nerve

d) Greater palatine nerve

Question 4:

How long should elective surgery be deferred for a patient who had cough and fever, after he has recovered from COVID-19, and had not required any hospitalization?

a) 6 weeks

b) 8-10 weeks

c) 12 weeks

d) 4 weeks

Question 5:

A 42-year-old man who was under home isolation after testing positive for COVID-19, develops breathing difficulty and is brought to the ER. The patient is alert and responsive, but is breathing rapidly. SpO2 is measured to be 86%. He is administered oxygen via a face mask with a reservoir bag with a flow rate of 10 L/min. After a trial of one hour, his SpO2 is 88%. What is the next step in management?

a) Increase flow rate to 15 L/min

b) High-flow nasal cannula oxygen therapy

c) Endotracheal intubation

d) No specific intervention necessary

Question 6:

What is the mechanism of action of local anesthetics?

a) Block sodium channels

b) Block potassium channels

c) Block calcium channels

d) Block chloride channels

Question 7:

You are in the operating room and notice the tracing in yellow color on this device. What does it indicate?



a) Capnography

b) Airway pressure

c) O2 pressure in inhaled air

d) O2 pressure in exhaled air

Question 8:

What is the most common cause for hospital admission after day care surgery?

a) Respiratory distress

b) Post surgical pain

c) Postoperative nausea and vomiting

d) Bleeding

Question 9:

What is the frequency of tetanic stimulation used in neuromuscular monitoring?

a) 0.2 Hz

b) 2 Hz

c) 20 Hz

d) 50 Hz

Question 10:

A 60-year-old male patient was undergoing laparoscopic cholecystectomy. During the surgery, a sudden drop in EtCO2 to 8 mmHg was noted. His SpO2 became 90%, his blood pressure dropped to 80/50 mmHg, and peak airway pressure was 18 cm of H2O. What is the diagnosis?

a) Both 1 and 2

b) Both 3 and 4

c) 1 only

d) 2 only

Question 11:

The maximum oxygen flow rate which can be kept with a conventional oxygen mask is

a) 5 L/min

b) 15 L/min

c) 10 L/min

d) 20 L/min

Question 12:

Which of the following can be used for induction of anesthesia, maintenance and can be safely used in endotracheal intubation?

a) Halothane

b) Desflurane

c) Sevoflurane

d) Isoflurane

Question 13:

During resuscitation, a patient received a 20% Intralipid injection. What could have been the most probable diagnosis?

a) Local anaesthesia toxicity

b) Asphxyia due to drowning

c) Electrocution

d) Pulseless electrical activity

Question 14:

A 50-year-old male patient has grade 3 ASA. Which of the following postbronchodilator PFT findings requires further evaluation of the lung function of the patient?

a) FEV1 < 0.85 and FEV1/FVC <0.8

b) FEV1 < 0.85 and FEV1/FVC <0.75

c) FEV1 < 0.8 and FEV1/FVC <0.7

d) FEV1 < 0.8 and FEV1/FVC <0.75

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	a
5	b
6	a
7	a
8	c
9	d
10	c
11	c
12	c
13	a
14	c

Detailed Explanations

Solution to Question 1:

Tissues become vulnerable to hypoxia when SpO2 falls below 90% and thus supplementary oxygen therapy is indicated below this oxygen saturation level.

In adults, children, and infants (older than 1 month), When SpO2 falls below 90%, the PaO2 becomes less than 60 mmHg and hence to prevent tissues from hypoxia, supplemental oxygen therapy should be started.

In neonates, when SpO2 falls below 88%, the PaO2 becomes less than 50 mmmHg and hence supplemental oxygen therapy should be initiated in these patients.

Solution to Question 2:

Non rebreathing masks provide highest FiO2 during conventional oxygen therapy.

The non-rebreather mask is a low flow device that can be used to deliver oxygen at higher concentrations (FiO2 80-95%). It uses a reservoir bag to let the oxygen get concentrated while a one-way valve doesn't allow to breathe in expired air.

This type of masks are used in patients with significant hypoxia, with relatively normal ventilation, such as victims of trauma or carbon monoxide exposure.

Other options:

A) Venturi mask:

• Venturi mask is a high-flow, fixed-performance oxygen delivery system and the highest oxygen concentration (FiO2) delivered is 60% at a flow rate of 15 L/min.

• It is based on the Bernoulli Principle, which states that in order to maintain a constant flow, a fluid must increase its velocity as it flows through a constriction.

• It is available in different colors, with the green valve venturi mask providing 60% FiO2.

• This is useful with patients who require greater FiO2 such as COPD patients, where it is important to provide controlled oxygenation.

B) Hudson mask:

• Also known as simple face mask

• Low-flow, variable performance device

• They provide an oxygen flow rate of up to 5-10 L/min and provide a variable FiO2.

• The body of the mask functions as reservoir for both oxygen and expired carbon dioxide.

• Used in patients who require oxygen therapy for shorter periods of time.

D) Nasal cannula:

• They are low-flow, variable performance devices, with flow rate of 1-4 L/min.

• Used in patients who require low supplemental oxygen.

• Nasopharynx acts as the reservoir in such cannulas.

Indications of a nasal cannula and other low-flow oxygen delivery systems include

• Minute ventilation < 8–10 L/min

• Breathing frequencies < 20 breaths/min

• Normal inspiratory flow (10–30 L/min).



Solution to Question 3:

The given image shows a nasociliary nerve block.

The procedure is done by inserting the needle 1 cm superior and 1.5 cm posterolateral to the medial canthal ligament of the eye.

The nasociliary nerve is a branch of the ophthalmic nerve which in turn is a branch of the trigeminal nerve.

Branches:

• Long ciliary nerve- cornea, ciliary body, iris

• Ramus communicans to the ciliary ganglion

• Posterior ethmoidal nerve- sensory innervation to frontal sinus and posterior ethmoidal air cells

• Anterior ethmoidal nerve- sensory innervation to middle and anterior ethmoidal cells as well as lateral part of the nasal septum

• Infra-trochlear nerve (continuation of the nasociliary nerve)- provides innervation to the conjunctiva, lacrimal sac, caruncle, the medial part of the eyelids

Uses of nasociliary nerve block:

Used in

• Surgery of nasal fractures,

• Septoplasty,

• Rhinoplasty and

• Repair of skin laceration.



The above image represents an anterior ethmoidal nerve block.

Solution to Question 4:

A patient, who has recovered from COVID-19 and hadn't required any hospitalization should be deferred from any elective surgery for a period of 6 weeks.

The American Society of Anesthesiologists has suggested the following duration of intervals for patients recovered from COVID-19. The wait time is from the date of diagnosis of COVID-19 to the date of surgery as follows:

• 4 weeks: for an asymptomatic patient or recovery from only mild, non-respiratory symptoms.

• 6 weeks: for a symptomatic patient (e.g: cough, dyspnea) who did not require hospitalization

• 8-10 weeks: for a symptomatic patient who is hospitalized or who has comorbidities such as diabetes or who is immunocompromised.

• 12 weeks: for a patient who was admitted in intensive care unit due to COVID-19 infection.

Solution to Question 5:

This patient has persistent hypoxia after an adequate trial of standard oxygen therapy and should be started on high-flow nasal cannula oxygen therapy.

When respiratory distress and/or hypoxemia of the patient cannot be alleviated after receiving standard oxygen therapy, high–flow nasal cannula oxygen therapy (HFNO), or non–invasive ventilation can be considered. Compared to standard oxygen therapy, HFNO reduces the need for intubation.

Patients on HFNO should be closely monitored, and endotracheal intubation should be done if the patient acutely deteriorates or does not improve after a short trial (about 1 hr).

Patients with hypercapnia (exacerbation of COPD), hemodynamic instability, multi-organ failure, or abnormal mental status should not receive HFNO.

Note: If a high flow nasal cannula (HFNC) or simple nasal cannula is used to administer O2, an N95 mask should be applied over it.

Some of the benefits of HFNC are

• It can deliver oxygen at 40L/min, 37°C, and 99.9% relative humidity.

• The high-flow oxygen is delivered directly to the nasopharynx. This produces FiO2 equal to that delivered by traditional mask devices.

• It also reduces the work of breathing, improves gas exchange by nasopharyngeal dead-space washout, and decreases energy required to humidify and heat respiratory gases.

The HFNC equipment shown in the image below uses a heated humidifier (named PMH200), a nasal cannula (no. 2), and a ventilator (brand: vivo30).



Solution to Question 6:

Local anesthetics primarily act by blocking the sodium channels bound to the membrane.

Sodium channels have one larger α subunit and one or two smaller β subunits. These voltage-gated sodium channels exist in 3 states :

• Resting (non conducting)

• Open (conducting) and

• Inactivated (non conducting)

The local anesthetics bind to the α subunit of sodium channels, thereby preventing their activation and inhibiting the influx of sodium.

Local anesthetics have a higher affinity for the sodium channels in open and inactivated state than in the resting state.

Local anesthetic inhibition is both voltage and frequency-dependent and is higher when nerve fibers are firing rapidly than with slow, infrequent depolarizations.

Note:

Local anesthetics may also bind and inhibit calcium, potassium, transient receptor potential vanilloid 1 (TRPV1) and many other channels and receptors. But blocking of sodium channels is the best answer, of all the given options.

Solution to Question 7:

The reading marked as X in the monitor is the capnography.

Waveform capnography is a continuous non-invasive measurement and graphical display of the EtCO2.

It is a direct measure of lung ventilation and an indirect measure of circulation and metabolism.

The normal capnograph displays the exhaled CO2 in both inspiratory and expiratory phases. The expiratory phase is divided into three distinct components:

• Phase I: exhalation of dead space gas (i.e., partial pressure of CO2, pCO2 ~0).

• Phase II: A sharp rise in pCO2 to a plateauindicates the sampling of transitional gas between the airways and alveoli.

• Phase III: The plateau region of the capnograph, corresponds to the pCO2 in the alveolar compartment.

• Following phase III, a sharp inspiratory downstroke occurs, which is referred to as the beginning of phase 0 or phase IV by some.



Solution to Question 8:

Postoperative nausea and vomiting (PONV) is the most common cause of unanticipated hospital admission following daycare surgery.

The two most common complications of daycare surgery are PONV and postoperative pain. PONV refers to the presentation of symptoms in the postoperative care unit or within 24 hours of the procedure. Pain is usually managed preoperatively and postoperatively with the administration of analgesics. While it is a common complication, postoperative pain is not the most common cause of unanticipated hospital admission. Opioid medications which are commonly used to treat postoperative pain present with side effects such as nausea and vomiting which may aggravate PONV.

The occurrence of PONV can be based on the following factors, patient, choice of anesthetic agent, and the procedure. The risk factors are:

• Patient: Women (higher incidence in pregnancy), history of PONV or motion sickness, nonsmokers, younger age (incidence decreases as the age increases).

• Anesthetic agent: General anesthesia, inhalational agents (isoflurane, sevoflurane), nitrous oxide, longer duration of anesthesia, preoperative opioid administration.

• Procedure: Laparoscopic procedures, lithotripsy, gynecologic procedures, adenotonsillectomy, strabismus surgery.

The drugs used in the prophylaxis and treatment of PONV include ondansetron, dexamethasone, and transdermal scopolamine.

Solution to Question 9:

The commonly used frequency of tetanic stimulation in neuromuscular monitoring is 50 Hz.

Tetanic stimulation is a method to evaluate residual neuromuscular blockade.

Tetanic stimulation consists of a series of extremely rapid electrical stimuli typically applied over 5 seconds. The most commonly used frequency is 50 Hz stimulation given for 5 seconds.

• During a depolarizing block, the muscle response to 50 Hz tetanic stimulation for 5 seconds is sustained.

• In contrast, during a phase II block after the injection of succinylcholine, the response following tetanic stimulation is not sustained i.e., a fade occurs.

Disadvantages of tetanic stimulation are:

• This method has only 70% sensitivity to detect residual paralysis and its specificity is only about 50%. Hence, this test is of limited value for assessing neuromuscular recovery.

• This method is very painful, which makes it difficult to use in unanesthetized patients.

• During the late phase of neuromuscular recovery, tetanic stimulation can produce lasting antagonism of neuromuscular block in the stimulated muscle. This may change the response of the tested site such that it may no longer be representative of other muscle groups.

Note:

Other patterns of nerve stimulations, which are used to assess the patients for residual blockade are

• Train of four stimulation: Here four supramaximal stimuli (2 Hz) are delivered every 0.5 seconds.

• Double-burst stimulation: Here two short bursts of 50 Hz tetanic stimuli is delivered, which are separated by 750 ms.

Solution to Question 10:

A sudden fall in EtCO2 with hypotension (fall in BP to 80/50 mmHg) along with normal airway pressure (< 35 to 40 cm of H2O) during surgery is suggestive of CO2 embolism.

Intraoperative pulmonary embolism usually presents as sudden cardiovascular collapse and hypoxemia with a sudden fall in EtCO2. Transesophageal echocardiography (TEE) is the most sensitive device for detecting intracardiac air embolus. If air is identified in the right atrium, or if it is suspected, immediate central vein cannulation and aspiration of the air may be lifesaving.

The placement of a vena cava filter should be considered postoperatively.

The best method to confirm the position of the endotracheal tube is the end-tidal CO2 concentration assessed by capnography.

Other options:

Intraoperative bronchospasm is usually manifested as wheezing, increasing peak airway pressure, and a slowly rising waveform on the capnograph.



A pneumothorax may manifest as hypercapnia along with hypoxemia, hypotension, increased peak airway pressure, and decreasing tidal volumes. It may result in abrupt cardiovascular collapse unresponsive to fluid and vasopressor administration.

Deep sulcus sign on a supine chest radiograph points towards pneumothorax. The costophrenic angle is abnormally deepened when the pleural air collects laterally, producing the deep sulcus

sign.

The below chest X-ray shows left pneumothorax. Notice the collapsed lung and absent lung markings on the left side. Also mediastinal shift towards the right side.

Endobronchial intubation shows increased peak expiratory pressure, hypoxemia, increased airway pressure, asymmetric chest expansion and the absence of breath sounds over one lung, generally the left.

Note: Peak airway pressure is the amount of airway pressure delivered by the ventilator to overcome the resistance of the endotracheal tube, airway, and alveoli.

Solution to Question 11:

The maximum oxygen flow rate which can be kept with a conventional oxygen mask is 10 L/min.

A conventional oxygen mask is a simple face mask that is used routinely in hospitals. It covers the nose and mouth of the patient. There are 2 exhalation ports through which the patient exhales carbon dioxide. This mask is held in place with elastic around the back of the head and a metal piece that fits over the nose. It provides 40% to 60% of O2 concentration and delivers O2 at the rate of 6 to 10 L/min.

Conventional oxygen therapy includes all the devices that are used in routine practices. It includes nasal prongs, low-flow nasal cannula, or masks. It is the main supportive treatment administered to patients after planned extubation.

Oxygen delivery devices can be broadly divided into low flow oxygen devices (eg: nasal cannula, simple face mask, non-rebreathing mask) and high flow oxygen devices (eg: venturi mask).

Important note- If the question is about the maximum flow rate with the conventional oxygen mask, it would be 10L/ min. Whereas if the question is asking about the maximum oxygen flow rate with conventional oxygen therapy, then it would include all the conventionally used devices. Therefore, the answer would be 15L/min.

The below image shows a nasal cannula, which delivers oxygen at 1-6 L/min.

Oxygen delivery devices	Flow rate (liter/min)	Approximate FiO2
Nasal cannula	1-6	0.24-0.44
Simple face mask	6-10	0.40-0.60
Non-rebreathing mask	10-15	0.90-1.00
Venturi mask	2-15	0.24-0.60



The below image shows a non-rebreathing mask that provides an oxygen flow rate of up to 10-15 L/min.

The below image shows a venturi mask, which delivers oxygen at 2-15 L/min.

Solution to Question 12:

Among the given options, sevoflurane can be used for induction and maintenance of anesthesia. It can also be safely used in endotracheal intubation

Inhalation agents such as sevoflurane and halothane are useful in the induction of anesthesia, But sevoflurane due to its nonpungency and rapid onset makes it an excellent choice for smooth and rapid inhalation inductions in pediatric and adult patients. Sevoflurane has low blood solubility and partition coefficients resulting in more rapid emergence.

Sevoflurane also produces adequate muscle relaxation for intubation of children following an inhalation induction.

Other options:

Option A: Halothane can also be used for gaseous induction of anesthesia. Its disadvantage is long induction time owing to the high blood:gas partition coefficient. However, the incidence of halothane hepatitis and myocardial depression has reduced its use.

Option B: Desflurane produces laryngospasm when used for inhalational anesthesia and hence it is not used for induction. Laryngospasm is because of the pungent nature of the gas.

Option D: Isoflurane causes respiratory depression as well as upper respiratory irritation.

Solution to Question 13:

Patients who are unresponsive due to local anesthesia toxicity require 20% Intralipid injection during resuscitation.

Accidental intravascular injection of local anesthetic like bupivacaine results in bupivacaine-induced cardiotoxicity.

Initial management includes immediate initiation of cardiopulmonary resuscitation (CPR) and defibrillation as per the advanced cardiac life support (ACLS) algorithm.

• Following CPR, Intralipid therapy is initiated. A rapid bolus of Intralipid 20% IV (1.5 mL/kg) is administered without delay, followed by an infusion of 0.25 mL/kg/min. The infusion should be given continuously for at least 10 minutes after hemodynamic stability is achieved.

• Intralipid decreases cardiac bupivacaine concentration by 11% within 3 minutes of administration, and cerebral bupivacaine concentration by 18% within 15 minutes.

Bupivacaine is a long-acting local anesthetic that is the most cardiotoxic among the local anesthetics. It depresses the rapid phase of depolarization in Purkinje fibers and ventricular muscle due to a decreased availability of fast sodium channels in cardiac membranes. This reduced rate of depolarization depresses myocardial contractility. Hence, bupivacaine is a potent cardio-depressant.

Note: Local anesthesia systemic toxicity (LAST) can also affect the central nervous system (CNS), which is more susceptible to local anesthetic action as compared to the cardiovascular system. Dizziness, tinnitus, and disorientation are the initial symptoms. Shivering, muscular twitching, and tremors follow. Tonic-clonic seizures occur. Finally, respiratory depression can lead to respiratory arrest.

Solution to Question 14:

FEV1 <0.8 and FEV1/FVC <0.7 are classified as moderate COPD under GOLD classification and they require further evaluation of the lung function.

Preoperative pulmonary function tests (PFTs) assess the severity of known pulmonary disease, diagnose causes of respiratory symptoms, and may help to identify patients who are at risk for postoperative pulmonary complications.

The severity of COPD is assessed by using the Global Initiative for Lung Disease (GOLD) criteria. In pulmonary function testing, a postbronchodilator FEV1/FVC ratio of <0.70 is considered diagnostic for COPD.

The Global Initiative for Chronic Obstructive Lung Disease (GOLD) system categorizes airflow limitation into stages. In patients with FEV1/FVC <0.70:

• GOLD 1: mild: FEV1 ≥80% predicted

• GOLD 2: moderate: 50% ≤ FEV1 <80% predicted

• GOLD 3: severe: 30% ≤ FEV1 <50% predicted

• GOLD 4: very severe: FEV1 <30% predicted.

This is important in the perioperative period as the diagnosis of COPD is an independent and significant risk factor in the development of postoperative pulmonary complications. Pre-operative FEV1 and DLCO of >80% imply a lower risk for a permanent postoperative pulmonary disability. Below these values, further risk-benefit assessment of the surgery needs to be examined and more information should obtained.

Anesthesia NEET 2021

Question 1:

A patient undergoing surgery is maintained on anesthesia with halothane. During the procedure, the patient suddenly developed hyperthermia and muscle rigidity. Which of the following drugs is most likely implicated in this condition?

a) D-curare

b) Suxamethonium

c) Cis-atracurium

d) Rocuronium

Question 2:

A 30-year-old male patient was intubated for surgery by a final-year resident. Which of the following is the best method to confirm the position of the endotracheal tube?

a) X-ray chest

b) Auscultation

c) End-tidal CO_2 concentration

d) Chest rise

Question 3:

What is the maneuver being performed in the given image?



a) Head tilt, chin lift

b) Jaw thrust

c) Head extension

d) In line manual stabilisation

Question 4:

A patient presented with headache, vomiting, and fever. Meningitis was suspected. While performing a lumbar puncture, which is the last structure you will pierce just before entering the lumbar cistern?

a) Ligamentum flavum

b) Dura mater

c) Arachnoid membrane

d) Pia mater

Question 5:

A young male patient was given regional block with 0.25% bupivacaine for a minor surgical procedure. Shortly after the injection, the patient suddenly became unresponsive and the pulse was not recordable. Which of the following would be the best course of management for this patient?

a) CPR with 20% intralipid

b) CPR with sodium bicarbonate

c) CPR with dobutamine

d) CPR with calcium

Answer Key

Question No.	Correct Option
1	b
2	c
3	a
4	c
5	a

Detailed Explanations

Solution to Question 1:

The given clinical scenario of a patient being operated under halothane, developing hyperthermia and muscle rigidity is suggestive of an episode of malignant hyperthermia. The drug most likely implicating/causing this condition is suxamethonium (succinylcholine).

Suxamethonium is a peripherally acting depolarizing muscle relaxant, commonly administered during the induction of anesthesia to relax muscles of the jaw, neck, and airway and thereby facilitate laryngoscopy and endotracheal intubation.

Malignant hyperthermia is a rare genetic hypermetabolic muscle disease. It occurs due to mutation of the gene for the ryanodine receptor (Ryr 1), located on chromosome 19. Ryr 1 is an ion channel responsible for calcium release from the sarcoplasmic reticulum.

The symptoms develop when the patient is exposed to inhaled anesthetics (halothane in this case) or suxamethonium/succinylcholine.

The mutated Ryr 1 leads to a sudden release of calcium from the sarcoplasmic reticulum of the skeletal muscle. This uncontrolled increase in intracellular calcium removes the inhibition of troponin, resulting in sustained muscle contraction, which in turn, leads to an increase in adenosine triphosphatase (ATPase) activity. This results in an uncontrolled hypermetabolic state with increased oxygen consumption and CO_2 production, producing severe lactic acidosis and hyperthermia.

The key clinical features are:

• Succinylcholine-induced masseter muscle rigidity (causing trismus) or generalized muscle rigidity

• Tachycardia

• Hypercarbia (increased CO_2 production) and reduced oxygen saturation

• Hyperthermia (rise of 1° Celsius can occur every 5 minutes)

• Arrhythmia

• Hypertension

• Myoglobinuria and myoglobinemia indicating muscle damage

• Metabolic derangements like hyperkalemia and mixed metabolic and respiratory acidosis

Treatment is to discontinue the triggering agents i.e; inhaled anesthetics and succinylcholine followed by administration of 100% oxygen. Metabolic acidosis and hyperkalemia need to be corrected. Hyperthermia must be controlled by cooling the patient.

Dantrolene sodium is the drug of choice. It is given at a dose of 2.5 mg/ kg intravenously and can be repeated every 5-10 minutes till the attack subsides.

Other options:

D-curare or D-tubocurarine (option A), cis-atracurium (option C), and rocuronium (option D) are non-depolarizing muscle relaxants and are not implicated in malignant hyperthermia.

Solution to Question 2:

The best method to confirm the position of the endotracheal tube is the end-tidal CO_2 concentration assessed by capnography.

Capnography is a non-invasive method of measurement and graphical display of partial pressure of carbon dioxide (CO_2) in exhaled breath or the end-tidal CO_2. Capnograph is the device used for measurement and capnogram is the CO_2 waveform that is displayed. It is the most reliable indicator of endotracheal tube placement.

The proper placement of the endotracheal tube is indicated by a normal CO_2 waveform for at least three breaths.



Other signs to confirm proper endotracheal intubation are:

Symmetrical chest rise, presence of equal bilateral breath sounds, absent breath sounds over the epigastrium, visible condensation in the ETT, large exhaled tidal volume, and appropriate compliance of the reservoir bag during manual ventilation.

Absent CO_2 waveform tracing despite correct ETT intubation is seen in the following conditions:

• Severe bronchospasm,

• Equipment malfunction,

• Cardiac arrest, or hemodynamic collapse

Clinical relevance:

• Capnography does not differentiate endotracheal from endobronchial intubation. Signs of endobronchial intubation include increased peak expiratory pressure, hypoxemia, increased airway pressures, asymmetric chest expansion and the absence of breath sounds over one lung, generally the left.

• In esophageal intubation, the capnograph can initially detect some CO_2 in the stomach from swallowing expired air. But after a few breaths, this CO_2 will be washed out and $ETCO_2$ will become zero.

• The most sensitive and earliest indicator of malignant hyperthermia is a persistent increase in $ETCO_2$.

Solution to Question 3:

The maneuver being performed in the given image is that of head-tilt chin-lift. It is an important technique in emergency airway management to open up the airway.

Basic life support involves early recognition and management of emergencies such as heart attack, stroke, and foreign body airway obstruction. The sequence of BLS is circulation (chest compression), airway, and then breathing.

The airway is usually obstructed by posterior displacement of the tongue or epiglottis. One hand (palm) is placed on the patient's forehead applying pressure to tilt the head back. Simultaneously, the chin is lifted with the forefinger and index finger of the opposite hand. This causes cervical flexion with an atlantooccipital extension (sniffing position). This lifts the tongue from the back of the throat.

Head-tilt chin-lift should not be performed in a patient with suspected cervical injury.

Other options:

Option B: Jaw thrust- Both hands of the clinician are placed on either side of the patient's head and the angles of the jaw are grasped and lifted. It can be performed in a patient with cervical injury but caution must be followed to not hyperextend the neck.

Jaw Thrust

Option D: Manual inline stabilization is a maneuver that limits external head and neck movement in patients with cervical spine injuries. Fingers and palms of both hands are used to stabilize the patient's occiput and mastoid processes to ensure successful airway intervention.



Solution to Question 4:

The arachnoid membrane is the last structure you will pierce while performing a lumbar puncture, just before entering the lumbar cistern.

Lumbar puncture is a diagnostic procedure done to obtain a sample of cerebrospinal fluid from the subarachnoid space. During the procedure, the following structures are pierced to reach the subarachnoid space, in the given order:

• Skin

• Subcutaneous tissue

• Supraspinous ligament

• Interspinous ligament

• Ligamentum flavum

• Dura mater

• Arachnoid membrane

Indications:

• Diagnosis of suspected meningitis and subarachnoid hemorrhage

• Spinal anesthesia and intrathecal administration of antibiotics/chemotherapy

• Therapeutic relief of pseudotumor cerebri, less frequently

Contraindications for lumbar puncture include:

• Thrombocytopenia or other bleeding diatheses (including anticoagulant therapy)

• Raised intracranial pressure

• Spinal epidural abscess

• Infection over the site

Site of the procedure:

In adults, the spinal cord ends at the level of the lower border of the L1 vertebra. Lumbar puncture is usually performed at L3-L4 space (below L1 vertebra) to avoid potential needle trauma to the spinal cord.

Technique:

The patient is usually placed in the lateral recumbent position. An alternative position is the sitting position, useful in obese patients to determine the midline. L3-L4 space is located by palpating the right and left posterior superior iliac crests and moving the fingers medially toward the spine at the same level.

A local anesthetic like 1% lidocaine is administered. A 22-gauge needle is advanced through the skin. In the lateral recumbent position, the bevel should face up, and in the sitting position, it should face to one side or the other, to avoid injury to the dural fibers. There is a loss of resistance as the needle enters the epidural space. Dura and then arachnoid are next pierced, to release CSF.

Complications following lumbar puncture:

Post-dural puncture headache (PDPH) is a common post-procedure symptom. It occurs due to loss of CSF through the dura which causes traction on the pain-sensitive intracranial structures. The headache in PDPH is typically frontal or occipital. The smaller the needle used, the lesser the risk of PDPH.

Other complications include infections, bleeding, cerebral herniation, and radicular pain.

The image below shows a patient undergoing lumbar puncture in the lateral recumbent position:



Solution to Question 5:

The clinical picture of an unresponsive patient with an absent pulse following bupivacaine administration is most likely due to bupivacaine-induced cardiotoxicity resulting from accidental intravascular injection of local anesthetic. The best course of management in this scenario would be administering CPR (cardiopulmonary resuscitation), along with a 20% intralipid injection.

Bupivacaine is a long-acting local anesthetic that is the most cardiotoxic among the local anesthetics. It depresses the rapid phase of depolarization in Purkinje fibers and ventricular muscle due to a decreased availability of fast sodium channels in cardiac membranes. This reduced rate of depolarization depresses myocardial contractility. Hence, bupivacaine is a potent cardio depressant.

Cardio-vascular effects of bupivacaine-induced toxicity are as follows:

• Left ventricular depression

• Atrioventricular heart block

• Ventricular arrhythmias and ventricular fibrillation

• Sinus bradycardia can lead to cardiac arrest

Initial management includes immediate initiation of cardiopulmonary resuscitation (CPR) and defibrillation as per the advanced cardiac life support (ACLS) algorithm.

Following CPR, intralipid therapy is initiated. Its mechanism is explained by the lipid sink theory which states lipophilic drugs (bupivacaine) are absorbed into the lipid emulsion of the plasma, hence, removed from tissues affected by toxicity.

A rapid bolus of intralipid 20% IV (1.5 mL/kg) is administered without delay, followed by an infusion of 0.25 mL/kg/min. The infusion should be given continuously for at least 10 minutes after hemodynamic stability is achieved.

Local anesthesia systemic toxicity (LAST) can also affect the central nervous system (CNS), which is more susceptible to local anesthetic action as compared to the cardiovascular system. Dizziness, tinnitus, and disorientation are the initial symptoms. Shivering, muscular twitching, and tremors follow. Tonic-clonic seizures occur. Finally, respiratory depression can lead to respiratory arrest.

Anesthesia INI-CET 2022

Question 1:

Arrange the following anesthetic agents on the basis of their potency.

a) Methoxyflurane> Halothane> Isoflurane> Nitrous oxide

b) Methoxyflurane> Isoflurane> Halothane> Nitrous oxide

c) Methoxyflurane> Nitrous oxide> Halothane> Isoflurane

d) Halothane> Isoflurane> Nitrous oxide> Methoxyflurane

Question 2:

Which of the following local anaesthetic causes hypertension?

a) Cocaine

b) Lignocaine

c) Bupivacaine

d) Prilocaine

Question 3:

Which of the following is required for the cannulation of a newborn?

a) 1,3,4,5

b) 2,3,4,5

c) 2,4

d) 1,4

Question 4:

Identify the incorrect statement in rapid sequence intubation?

a) Preoxygenation with 100% oxygen for 3 min is done.

b) High-volume low-pressure cuff is used.

c) It is the method of choice in cardiac arrest patient.

d) Induction agent and muscle relaxant are given simultaneously.

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	c

Detailed Explanations

Solution to Question 1:

The potency of the inhaled anesthetics given above can be arranged as methoxyflurane> halothane> isoflurane> nitrous oxide. It is due to the change in the agents' minimum alveolar concentration (MAC).

The minimum alveolar concentration (MAC) of an inhaled anesthetic is the lowest concentration required to prevent movement in response to a defined noxious stimulus in 50% of subjects. It is a measure of the potency of inhalational anesthetics. So, the lower the MAC value, the lower the concentration of anesthetic drug required, implying the drug is more potent.

MAC values (expressed as a '%' of 1 atmosphere) of some anesthetic agents:

• Nitrous oxide with a MAC value of 104 is the least potent inhaled anesthetic

• Methoxyflurane – MAC- 0.2 (most potent)

• Desflurane: 6.0

• Sevoflurane– 2.05

• Isoflurane – 1.28

• Halothane – 0.75

Solution to Question 2:

The local anesthetic that causes hypertension is cocaine. This is due to the inhibition of norepinephrine at the adrenergic terminals leading to increased sympathetic stimulation. This effect occurs at all concentrations of cocaine and causes hypertension and ventricular ectopies.

Treatment is with the help of benzodiazepines, α-adrenergic antagonists and amiodarone.

All local anesthetics are mainly cardiodepressants causing reduced contractility and vasodilation. However, at low concentrations, all local anesthetics cause vasoconstriction due to the inhibition of nitric oxide. Only cocaine produces vasoconstriction and hypertension at all concentrations.

Accidental intravascular injection of local anesthetic like bupivacaine results in bupivacaine-induced cardiotoxicity. Management is by resuscitation followed by intralipid therapy. A rapid bolus of intralipid 20% IV is administered without delay.

Other adverse effects of local anesthetics include neurologic toxicity, respiratory depression, and hypersensitivity reactions. Nervous system manifestations are circumoral numbness, tongue paresthesia, dizziness, tinnitus, and disorientation. Finally, respiratory depression can lead to respiratory arrest.

Solution to Question 3:

For cannulation in newborns, we would require a 24G needle, 5 mL syringe, adhesive tape, and normal saline.

Also, for a peripherally inserted central catheter (PICC), we use a 24G needle for >1 kg neonate, and for <1 kg neonate, we use a 28G needle. We need to avoid using 1 mL syringes for flushing as it may lead to catheter rupture; preferably, use 5 or 10 mL syringes. Flushing of the cannulation is done using 0.9% sodium chloride or so-called normal saline (NS).

Dressing change: We change PICC line dressing only when visibly soiled, damp, or loosened. For making it easier and sterile, two persons should perform a dressing change.

Solution to Question 4:

The incorrect statement regarding rapid sequence intubation (RSI) is that it is the method of choice in cardiac arrest patients.

RSI is a technique of anesthesia induction in patients who are at an increased risk of gastric regurgitation and pulmonary aspiration of gastric contents. Common cases include pregnancy, full stomach, intestinal obstruction, poorly controlled gastroesophageal reflux disease, morbid obesity, and diabetic gastroparesis. The goal is to intubate rapidly to minimize the time between the loss of consciousness and securing of the airway with a cuffed endotracheal tube.

It is achieved by pre-oxygenating the patient with 100% oxygen for 2-3 minutes. An inducing agent is administered, which is immediately followed by a muscle relaxant succinylcholine or rocuronium. The trachea is intubated without any prior bag and mask ventilation using a high-volume and low-pressure cuff which adequately cuffs the airway and prevents aspiration.

Cardiac arrest is an emergency condition with importance given to start chest compressions, defibrillation, or drugs to restore perfusion. The airway is commonly blocked by posterior displacement of the tongue which can be managed by head tilt and chin lift or giving jaw thrust. In case of airway blockade or to prevent aspiration, the airway is maintained by emergency tracheal tube intubation and does not involve the procedure that is followed in RSI. Endotracheal tube placement is commonly done during post-cardiac arrest care.

RAPID SEQUENCE INDUCTION



Anesthesia NEET 2022

Question 1:

The image given below shows neuromuscular monitoring of the patient after anesthesia. What is the most commonly used nerve for monitoring?



a) Ulnar nerve

b) Median nerve

c) Radial nerve

d) Metacarpal nerve

Question 2:

Identify the mask in the image used for patients with COVID-19 infection.

a) Venturi mask

b) Hudson mask

c) Nebuliser

d) Non-rebreathing mask

Answer Key

Question No.	Correct Option
1	a
2	d

Detailed Explanations

Solution to Question 1:

The following image shows neuromuscular monitoring and the most commonly used nerve for monitoring is the ulnar nerve.

The mechanosensor with electrodes is placed over the region of the ulnar nerve. This assesses the contraction of the adductor pollicis muscle to monitor neuromuscular activity.

Monitoring of neuromuscular blockade is required to monitor the blockade intraoperatively, ensure its complete reversal postoperatively, and prevent residual blockade. To monitor the neuromuscular function, a peripheral motor nerve is supramaximally stimulated and

the response of the muscle is assessed.

The most commonly assessed nerve-muscle unit is the ulnar nerve-adductor pollicis muscle. This is because it is easily accessible, has the lowest risk of direct muscle stimulation, and the response can be evaluated visually, by tactile methods, or by objective methods. To stimulate the ulnar nerve, the electrodes are best applied to the volar aspect of the wrist.

Other nerve-muscle units that can be assessed include:

• Posterior tibial nerve- flexor hallucis brevis muscle

• Facial nerve- orbicularis oculi muscle

• Facial nerve- corrugator supercilii muscle

Many methods can be used for objective monitoring of neuromuscular function:

• Mechanomyography (MMG)

• Electromyography (EMG)

• Acceleration of the muscle response (AMG)

• Kinemyography (KMG)

• Cuff pressure modality (CPM)

• Compressomyography (CMG)

• Phonomyography (PMG)

Solution to Question 2:

The mask in the image is a non-rebreathing mask (NRBM). It can be used for patients with COVID-19 infection.

A NRBM has a reservoir bag attached to the mask in order to provide additional reservoir volume. In addition, there is a valve attached, which provides a unidirectional flow of gases. Flutter valves on the side ports prevent room air entrainment, while a valve incorporated between the mask and the reservoir bag prevents the entry of the exhaled gases.

NRBM can provide the highest FiO2 of 0.9 to 1.0. without intubation at the fresh gas flow of 12-15 L/min. They are used for preoxygenation before intubation and post-extubation in covid patients.

Other options:

Option A: The venturi mask is a high-flow, fixed-performance device that delivers a specific oxygen concentration to the patient. It is a color-coded device that connects to the oxygen tubing distally and has a nozzle connecting to a mask proximally. Oxygen flows through a central constriction inside the device. Room air enters through surrounding apertures which dilutes the oxygen to the required concentration. The venturi mask is based on Bernoulli's principle.

Option B: Hudson mask is a simple face mask that delivers oxygen without forming a tight seal against the patient's face. There are holes on either side of Hudson's mask. This provides air entrainment, meeting the peak inspiratory flow and venting out the expired gases.

Option C: A nebulizer is used to deliver drugs to the breathing system. It generates an aerosol, producing droplets of varying sizes. It is shown below.

Anesthesia NEET 2023

Question 1:

Identify the maneuver being performed in the given image.

a) Head tilt, chin lift

b) Jaw thrust

c) Head extension

d) In line manual stabilization

Question 2:

Which of the following is the drug of choice for preoperative antibiotic prophylaxis in a patient undergoing cardiac surgery?

a) Penicillin G

b) Erythromycin

c) Azithromycin

d) Cefazolin

Question 3:

During resuscitation, when is the given position indicated?

a) Unconsciousness with pulse and breathing absent

b) Unconsciousness with pulse present and breathing absent

c) Unconsciousness with pulse and breathing present

d) Unconsciousness with pulse absent and breathing present

Answer Key

Question No.	Correct Option
1	a
2	d
3	c

Detailed Explanations

Solution to Question 1:

The maneuver performed in the given image is that of a head-tilt chin-lift. It is an important technique in emergency airway management to open up the airway.

Basic life support (BLS) involves early recognition and management of emergencies such as heart attack, stroke, and foreign body airway obstruction. The sequence of BLS is circulation (chest compression), airway, and breathing.

The airway is usually obstructed by posterior displacement of the tongue or epiglottis. One hand (palm) is placed on the patient's forehead, applying pressure to tilt the head back. Simultaneously, the chin is lifted with the forefinger and index finger of the opposite hand. This causes cervical flexion with an atlantooccipital extension (sniffing position). This lifts the tongue from the back of the throat.

Head-tilt chin-lift should not be performed in a patient with suspected cervical injury.



Other options:

Option B: Jaw thrust - Both hands of the clinician are placed on either side of the patient's head, and the angles of the jaw are grasped and lifted. It can be performed in a patient with cervical injury, but caution must be exercised to not hyperextend the neck.



Option D: Manual inline stabilization is a maneuver that limits external head and neck movement in patients with cervical spine injuries. Fingers and palms of both hands are used to stabilize the patient's occiput and mastoid processes to ensure successful airway intervention.

Solution to Question 2:

First- and second-generation cephalosporins (cefazolin) are the drugs of choice for preoperative prophylaxis for cardiac surgery. Clindamycin is reserved for cases of allergy to beta-lactams and vancomycin is recommended for methicillin-resistant Staphylococcus aureus (MRSA).

Preoperative antibiotic prophylaxis is essential in reducing the risk of surgical site infections, particularly in cardiac surgery, where the consequences of infection can be severe. The choice of antibiotic depends on the patient's medical history, allergies, and the most common pathogens involved in surgical site infections for the specific surgery. In addition to cefazolin, other cephalosporins, such as cefuroxime, can also be used for prophylaxis in cardiac surgery. The timing of antibiotic administration is crucial, with the first dose typically given within 60 minutes before the incision.

Other Options:

Option A: Penicillin G is not the drug of choice for preoperative antibiotic prophylaxis in cardiac surgery. It is mainly used to treat infections caused by susceptible gram-positive bacteria, such as Streptococcus pyogenes, Streptococcus pneumoniae, and some strains of Staphylococcus aureus.

Option B: Erythromycin is a macrolide antibiotic and is not the drug of choice for preoperative prophylaxis in cardiac surgery. It is typically used to treat respiratory tract infections, skin infections, and other infections caused by susceptible gram-positive bacteria.

Option C: Azithromycin, another macrolide antibiotic, is also not the drug of choice for preoperative prophylaxis in cardiac surgery. It is commonly used to treat respiratory tract infections, sexually transmitted infections, and other infections caused by susceptible gram-positive and gram-negative bacteria.

Solution to Question 3:

The position illustrated in the picture is known as the recovery position, and it is used in the management of patients who are unconscious but with pulse present and breathing present.

The recovery position can help prevent complications such as aspiration and airway obstruction in unconscious individuals. Additionally, it is crucial to monitor the patient's airway, breathing, and circulation continuously while they are in the recovery position and seek immediate medical assistance. An unresponsive patient who is breathing spontaneously in the supine position is at risk of airway obstruction from their tongue, mucus, vomit, etc. A modified lateral position is used and the following 6 principles are considered while choosing which variation of recovery position to use,

•

The victim should be in as near a true lateral position as possible, with the head dependent to allow free drainage of fluid.

•

The position should be stable.

•

Avoid any pressure on the chest that impairs breathing.

•

It should be possible to turn the victim on his or her side and to return to the back easily and safely, with concern for a possible cervical spine injury.

•

Good observation of and access to the airway should be possible.

•

The position itself should not cause injury to the victim.

Other options

Option A: Unconsciousness with pulse and breathing absent - In this situation, the patient is in cardiac arrest and requires immediate cardiopulmonary resuscitation (CPR), not the recovery position. CPR involves chest compressions and rescue breaths to support circulation and respiration.

Option B: Unconsciousness with pulse present and breathing absent - In this case, the patient is experiencing respiratory arrest. Rescue breathing or bag-valve-mask ventilation should be initiated to support the patient's breathing. The recovery position is not indicated in this scenario.

Option D: Unconsciousness with pulse absent and breathing present - This option is not physiologically plausible, as the absence of a pulse implies a lack of circulation, and breathing cannot be present without circulation.

Anesthesia INI-CET 2023

Question 1:

A patient who is under general anaesthesia shows the following finding on the ET CO2 monitor. Which condition can lead to this characteristic waveform on capnography?



a) Malignant hyperthermia

b) Bronchospasm

c) Rebreathing

d) Accidental extubation

Question 2:

In a patient with thrombocytopenia, platelet transfusion is done when the platelet level drops below a threshold. To proceed with surgery, platelet transfusion is done to achieve a platelet level of?

a) 30,000

b) 40,000

c) 50,000

d) 60,000

Question 3:

The presence of which of the following indicates that CPR is of high quality?

a) 2, 3 only

b) 1, 3, 4 only

c) 1, 2, 3, 4

d) 1 only

Question 4:

A 70 kg patient who presented to the emergency with acute pancreatitis was shifted to the ICU and intubated. His ventilatory settings are: FiO2 90%, PEEP 12 mmHg, respiratory rate 16/min, and tidal volume 420 ml. The results of the ABG analysis are given below. Which of the following would be the next step in the management of this patient?

a) Increase PEEP to 16

b) Increase tidal volume

c) Reduce respiratory rate

d) Reduce FiO2

Question 5:

Which of the following best represents the American Society of Anesthesiologists (ASA) physical status III?

a) Mild systemic disease

b) Moderate systemic disease

c) Severe systemic disease

d) Severe systemic disease with threat to life

Question 6:

A 5-year-old is on 3L/min O2 via a nasal cannula. What is the appropriate FiO2 level for this child?

a) 25

b) 30

c) 45

d) 60

Answer Key

Question No.	Correct Option
1	b
2	c
3	b
4	d
5	c
6	b

Detailed Explanations

Solution to Question 1:

The above capnographic waveform depicts bronchospasm.

Shark fin appearance - Prolonged phase II, increased α angle, and steeper phase III suggest bronchospasm or airway obstruction.

Solution to Question 2:

In an otherwise healthy patient, surgery can safely be performed in patients with platelet counts higher than 50,000/mm3. The risk of bleeding increases significantly if the platelet count is below this level. When platelet transfusions are used to treat thrombocytopenia, the platelet count rises by 10,000/mm3 for every unit transfused.

Thrombocytopenia is defined as a platelet count less than 1,50,000/mm3. It may be the result of immune thrombocytopenic purpura (ITP), drug-induced thrombocytopenia, rheumatological diseases (SLE, rheumatoid arthritis), pregnancy and chronic liver disease. The initial step if the patient has low platelet count is to repeat the test and examine the peripheral smear.

Anemia is also a common preoperative hematological disorder. The optimal preoperative hemoglobin concentration varies between 7.5 to 10 g/dL across the individuals, with inter-individual differences due to co-morbid conditions in the patients. Patients with anemia or suspected anemia must undergo a battery of investigations to determine the type of anemia and how best to correct it. Preexisting anemia is a risk factor for postoperative death and

complications such as AKI, stroke and infections.

Solution to Question 3:

Chest compressions at a rate of 100-120/minute, depth of compression 5-6 cm, and allowing complete chest recoil indicate high-quality CPR. The compression-to-ventilation ratio in the absence of an advanced airway should be 30:2 (30 compressions to be followed by 2 breaths). If an advanced airway (endotracheal tube or supraglottic airway device) is present, 10 breaths/minute should be given.

Solution to Question 4:

In a patient with a history of acute pancreatitis who needed to be intubated, the diagnosis of acute respiratory distress syndrome (ARDS) is the most likely. In this patient, the PaO_2 is 110 mmHg and the SpO_2 is 100%, which are significantly higher than the target recommended in patients with ARDS. The next step in the management of this patient would be to reduce the FiO_2 to prevent any further lung injury and worsening of ARDS due to oxygen toxicity. In the initial phase of the management, the FiO_2 of the patient should be kept ≤ 60% and SpO_2 should be maintained between 88-95%.

The main treatment strategy which has been shown to reduce mortality in patients is low tidal volume ventilation (6 mL/kg of predicted body weight). Increasing the tidal volume (option B) in this patient will lead to barotrauma to the alveoli because air is pushed through the alveoli which are already collapsed.

Although a high PEEP keeps the alveoli opened up, there is no rationale for increasing the PEEP further (option A) in this patient because the oxygenation is adequate. Furthermore, PEEP must be adjusted to minimize FiO_2 and provide adequate PaO_2 at the same time.

Hypercapnia is expected in a patient on low tidal volume ventilation. It can be managed by increasing the respiratory rate. However, respiratory acidosis is tolerated by the patient up to a blood pH of 7.2. This is known as permissive hypercapnia. Reducing the respiratory rate (option C) in this patient will increase the $PaCO_2$ further and cause a worsening of respiratory acidosis.

Apart from these, prone position, early neuromuscular blockade (to prevent the patient from fighting the ventilator), and use of extracorporeal membrane oxygenation (ECMO) have also been shown to improve mortality in patients with ARDS. Management strategies that have not been shown to improve the outcome in ARDS patients include glucocorticoids, surfactant replacement, anti-inflammatory therapies like NSAIDs, and inhaled vasodilators.

Solution to Question 5:

Severe systemic disease best fits into grade 3 of the ASA classification.

American Society of Anesthesiologists classification is used to define relative risk prior to conscious sedation and surgical anesthesia.

It is simple, reproducible and strongly correlates to perioperative risk.

American Society of Anesthesiologists physical status classification of patients-

Class 1: Normal healthy patient

Class 2: Patient with mild systemic disease (no functional limitations)

Class 3: Patient with severe systemic disease (some functional limitations)

Class 4: Patient with severe systemic disease that is a constant threat to life (functionality incapacitated)

Class 5: A moribund patient who is not expected to survive without the operation

Class 6: A brain-dead patient whose organs are being removed for donor purposes

Solution to Question 6:

The appropriate FiO2 for this child would be 30.

The formula for calculating FiO2 for a child on oxygen supplementation through nasal cannula is FiO2= 21%+ (nasal cannula flow in L/min) × 4

A nasal cannula is a low flow device, and FiO2 should be adjusted according to flow rates.

Examples of low-flow devices are:

• Nasal cannula

• Face mask

• Face tent

• Non-rebreather mask.

High-flow devices include:

• Venturi mask

• High flow nasal cannula

• Mechanical ventilation

Oxygen delivery devices	Flow rate (liter/min)	Approximate FiO2
Nasal cannula	1-6	0.24-0.44
Simple face mask	6-10	0.40-0.60
Non-rebreathing mask	10-15	0.90-1.00
Venturi mask	2-15	0.24-0.60

Anesthesia INI-CET 2024

Question 1:

Which of the following is a 2nd generation LMA device?

a) Intubating LMA

b) Proseal LMA

c) Flexible LMA

d) Ambu LMA

Question 2:

A patient was planned for surgery under general anaesthesia. During induction of anaesthesia, rocuronium 85 mg was given and the patient was tried to be intubated. It resulted in a failed intubation. Which of the following drugs can be given for immediate reversal of the drug?

a) Neostigmine

b) Glycopyrrolate

c) Sugammadex

d) Edrophonium

Question 3:

Which of the following are components of high-quality cardio-pulmonary resuscitation(CPR) in adults?

a) 1,2 and 3

b) 1,3 and 4

c) 2,3 and 4

d) 3 and 4 only

Question 4:

What is the inhalational anaesthetic agent used for induction in children?

a) Nitrous oxide

b) Isoflurane

c) Sevoflurane

d) Desflurane

Question 5:

Ambu Aura 40 is an example of?

a) Supraglottic laryngeal airway generation 1

b) Supraglottic laryngeal airway generation 2

c) Supraglottic laryngeal airway generation 3

d) Supraglottic laryngeal airway generation 4

Question 6:

Which of the following drugs causes adrenal suppression?

a) Thiopental

b) Ketamine

c) Etomidate

d) Propofol

Question 7:

Which of the following is incorrect regarding thiopentone?

a) It is lipid soluble

b) It achieves maximum concentration in the brain after an IV infusion

c) Redistribution to adipose tissue and muscle is seen

d) Rapid decline in its effect is due to renal excretion

Question 8:

A female following an RTA injured her chest. On X-ray, a fracture of 3 to 5th ribs was noted. Pain was not responsive to analgesics. Which block will be given to reduce the pain?

a) Cervical block

b) Thoracic epidural block

c) Supraclavicular block

d) Lumbar epidural block

Question 9:

Which of the following will lead to a decrease in diastolic BP?

a) Systemic infusion of alpha-adrenergic blocker

b) Systemic infusion of smooth muscle relaxant

c) Loss of lumbar sympathetic outflow

d) Systemic infusion of beta-1 adrenergic agonist

Question 10:

Which of the following is true regarding paediatric cardiopulmonary resuscitation?

a) Chest compression at the rate of 30:2 with 2 rescuers

b) Adenosine via interosseous is preferred over other methods

c) In chest compression, depth must be at least 1/3rd of the child's chest AP diameter

d) Adrenaline given in dose 0.01 mg/kg iv/io in 1:1000 concentration

Question 11:

Which of the following must be avoided in post-cardiac arrest resuscitation care?

a) Maintain MAP over 65

b) Maintain SpO2 above 94%

c) Revascularisation procedure if the patient develops ST elevation post-resuscitation

d) Bring the body temperature down to below 32 degrees, therapeutic hypothermia if unresponsive

Question 12:

Jackson Rees circuit is used in

a) Paediatric anaesthesia

b) Cardiac anaesthesia

c) Neuro anaesthesia

d) Obstetric anaesthesia

Answer Key

Question No.	Correct Option
1	b
2	c
3	b
4	c
5	a
6	c
7	d
8	b
9	c
10	c
11	d
12	a

Detailed Explanations

Solution to Question 1:

Proseal LMA is a 2nd generation LMA device as it has a separate gastric tube to decompress the stomach and prevent gastric aspiration associated with the classical LMA or 1st generation LMA. Other 2nd generation LMA's include LMA Supreme, I-Gel and LMA Protector.



All other options are examples of 1st generation LMA's since they do not have a separate gastric tube to aspirate for gastric secretions.

Flexible LMA allows the tube complete flexibility and resistance to compression so that the head and neck can be turned without dislodging the mask.

The intubating LMA (ILMA) and Ambu LMA(ALMA) are variants of the LMA Classic, a 1st generation LMA used to aid in intubation of the patient in difficult airways.

Note : Although Ambu LMAs are produced as 1st generation and 2nd generation LMA, Proseal is exclusively a 2nd generation LMA and hence the best answer to this question.

Solution to Question 2:

Sugammadex is used for immediate reversal of rocuronium induced neuromuscular blockade/relaxant in the given scenario.(Option C)

Sugammadex is a modified γ-cyclodextrin and the first selective relaxant–binding agent based on an encapsulating principle for inactivation of nondepolarising neuromuscular blocking drugs mainly rocuronium and vecuronium. This principle of complex formation of sugammadex and rocuronium or vecuronium in a 1:1 ratio occurs at all levels of neuromuscular blockade resulting in a more fast-acting pharmacologic reversal when compared with anticholinesterase drugs like neostigmine.

The stability of the rocuronium-sugammadex complex is a result of Van der Waals forces, hydrogen bonds and hydrophobic interactions. There is some binding affinity with pancuronium, but this interaction is too low to have a significant clinical effect.

Other options:

Option A: Neostigmine is used to reverse depolarising neuromuscular blockade such as succinylcholine, especially in the phase II block(fade of the train-of-four response) and sufficient time has passed for the circulating concentration of succinylcholine to be negligible. Therefore, neostigmine is slow-acting compared to sugammadex but the agent preferred in the case of succinylcholine.

Neostigmine administration during phase I block, increases neuromuscular Ach concentration and inhibits pseudocholinesterase (plasma cholinesterase) activity that can prolong the action of succinylcholine, hence it is given after some time during the phase II block.

Option B: To prevent cholinergic side-effects of neostigmine, glycopyrrolate is co-administered in a 1:5 ratio to neostigmine(0.2mg: 1 mg). Atropine is an alternative but glycopyrrolate is preferred

due to the incidence of tachycardia associated with atropine.

Option D: Edrophonium is a short-acting acetylcholinesterase inhibitor preferred in the case of mivacurium reversal since it does not show plasma cholinesterase inhibition present in neostigmine that can prolong mivacurium action.

Solution to Question 3:

The depth of compression should be at least 5cm to ensure high-quality CPR in an adult cardiac arrest.

The foundation of CPR is high-quality chest compressions. Compressing the chest pumps blood from the heart to the brain and then to the rest of the body. Each time the compressions are stopped, the blood flow from the heart decreases significantly. Once resumed, it takes several compressions to bring the blood flow back up to the levels present before the interruption.

The components of high-quality chest compressions in adults are as follows :

• Position victim - Position the victim face up on a firm, flat surface. This allows compression of the heart to create adequate blood flow.

• Compression-to-ventilation ratio - Single rescuers should use the compression-to-ventilation ratio of 30 compressions to 2 breaths when giving CPR to victims of any age.

• Compression rate - Compress at a rate of 100 to 120/min. This rate is the same for compression in all cardiac arrest victims.

• Compression depth - Compress the chest at least 5cm. Compressing the chest more than 6 cm in adults may decrease the effectiveness of the compression and cause injuries. Using a CPR-quality feedback device can help reach the optimal depth.

• Chest recoil - Allow the chest to recoil completely after each compression. This allows blood to flow into the heart. To help ensure complete recoil, avoid leaning on the chest between compressions. Chest compression and chest reoil times should be about equal.

• Interruptions in chest compressions - Minimize interruptions in chest compression. the proportion of time that rescuers perform chest compressions during CPR is called chest compression fraction (CCF). A CCF of 80% or higher should be the goal.

Solution to Question 4:

Sevoflurane is the inhalational anaesthetic agent used for induction in children. (Option C)

It is the preferred agent due to its non-pungent sweet-smelling nature and rapid increase in alveolar concentration on administration. Its low blood solubility helps during emergence from the effects of anaesthesia, reducing the post-operative delay in the patient's recovery from anaesthesia.

Desflurane and isoflurane are avoided for inhalation induction because they are pungent and associated with coughing, breath-holding and laryngospasm. They are usually used for the maintenance of anaesthesia. Desflurane is preferred in patients with renal and liver disease and

isoflurane is preferred in neurosurgery and cardiac surgery.

Due to its second-gas effect, nitrous oxide is mainly used as a carrier gas for other inhalational agents and not given alone.

Solution to Question 5:

The Ambu Aura 40 is an example of a first-generation supraglottic airway device.

Supraglottic airway (SGA) devices are classified into different generations based on their features and advancements:

First-generation SGA devices: Basic devices designed to provide a seal around the glottis for ventilation. They lack additional safety features, such as protection against aspiration. Examples include the original Laryngeal Mask Airway (LMA Classic) and Ambu Aura 40 (can be reused up to 40 times),

Second-generation SGA devices: These are improved devices with additional features to enhance safety and functionality. They typically have:

A gastric access channel (to allow gastric decompression or reduce the risk of regurgitation and aspiration).

Improved sealing mechanisms. Examples: Ambu Aura Gain (with drain tube), LMA ProSeal, and i-gel.

Third-generation and Fourth-generation devices: These are not standard classifications in clinical practice yet. However, devices beyond the second generation may feature advanced integration with monitoring systems, adjustable seals, or further innovations.

Solution to Question 6:

Etomidate causes adrenal suppression which can last for 72 hours.

Etomidate inhibits 11β-hydroxylase, resulting in decreased biosynthesis of cortisol and mineralocorticoids and increases the formation of intermediaries like 11-deoxycorticosterone.

It is a more potent inhibitor of steroid synthesis than as a sedative-hypnotic as the concentration which produces adrenocortical suppression (10 ng/mL)is much lower than that needed for hypnosis (>200 ng/mL).

This suppression is particularly concerning in critically ill patients, where stress-induced cortisol production is vital for survival.

Other Options:

Option A: Thiopental: Primarily causes sedation and hypnosis.

Option B: Ketamine: Does not cause adrenal suppression. Instead, it may slightly stimulate adrenal catecholamine release due to its action on the sympathetic nervous system.

Option D: Propofol: Can cause CNS depression without direct impact on adrenal hormone synthesis.

Solution to Question 7:

The rapid decline in the effect of thiopentone is due to its rapid redistribution, not due to renal excretion.

Thiopentone has a rapid onset of action because of its lipophilic nature, so it rapidly crosses the blood-brain barrier and enters highly perfused organs like the brain, heart, liver, and kidneys.

As thiopentone redistributes from the brain to tissues: it perfuses to muscle, fat, and skin. Over time, due to its high lipid solubility, thiopentone accumulates maximally in fat. Fat acts as a storage reservoir since it has low blood flow, and drug release from fat occurs slowly.

This redistribution reduces the concentration of thiopentone, terminating its anesthetic effect.

Other Options:

Option A: It is lipid-soluble: Thiopentone is highly lipid soluble, which allows it to cross the blood-brain barrier quickly and achieve rapid onset of action in the CNS.

Option B: It achieves maximum concentration in the brain after an IV infusion: Due to its lipophilicity, thiopentone rapidly accumulates in the brain after IV administration.

Option C: Redistribution to adipose tissue and muscle is seen: Thiopentone's effect is terminated by redistribution from the brain to adipose tissue and muscle. This accounts for the short duration of its anesthetic effect.

Solution to Question 8:

A thoracic epidural block will be given to reduce the rib fracture pain.

Epidural anaesthesia is used to provide adequate analgesia which is the mainstay of treatment in patients with multiple rib fractures.

The complications can be minimized by adequate pain relief with epidural anaesthesia. Rib fracture patients will have the following complications due to splinting and pain:

• Hypoventilation

• Hypoxia

Other techniques to provide analgesia following rib fractures are intercostal nerve block and intrapleural block which necessitates multiple injections, which increases the risk for complications.

The complications include vascular puncture, pneumothorax, patient discomfort due to the need for palpation of the fractured rib during local anaesthetic (LA) administration. Also, LA toxicity can occur because of the high systemic absorption.

Rib fractures can cause severe pain that does not respond well to conventional analgesics. A thoracic epidural block is effective for pain relief in this area because it provides segmental anesthesia to the thoracic dermatomes and surrounding structures, helping reduce pain and improve respiratory function.

Indications of thoracic epidural block:

• Postoperative pain relief: After thoracic or upper abdominal surgeries.

• Trauma management: Rib fractures to reduce pain and improve ventilation.

• Chronic pain syndromes: Postherpetic neuralgia or cancer-related pain.

Other Options:

Option A: Cervical block: Used primarily for pain management or anesthesia involving the neck and upper shoulders.

Option C: Supraclavicular block: Used for anesthesia of the upper limb by blocking the brachial plexus.

Option D: Lumbar epidural block: Used for lower body anesthesia, including the pelvis and lower extremities.

Solution to Question 9:

Loss of lumbar sympathetic outflow will lead to a decrease in diastolic blood pressure (DBP).

Loss or blockade of the lumbar sympathetic chain (spinal anaesthesia) causes dilation of capacitance (venous) vessels. This leads to a decrease in diastolic blood pressure and subsequent orthostatic lightheadedness or syncope.

Vasodilation is more marked on the venous than on the arterial side of the circulation, resulting in blood pooling in the venous capacitance vessels. This reduction in circulating blood volume is well tolerated at low levels of spinal anaesthesia in healthy patients. With an increasing level of block, this effect becomes more marked, and venous return becomes gravity-dependent. If venous return decreases too much, cardiac output and organ perfusion decline sharply.

Other options:

Option A: Systemic infusion of alpha-adrenergic blocker will minimally decrease diastolic BP, not significantly like in loss of lumbar sympathetic outflow.

Option C: Systemic infusion of smooth muscle relaxants like Sildenafil, which is a PDE 5 inhibitor (increases cGMP), does not cause a decrease in DBP.

Option D: Systemic infusion of beta-1 adrenergic agonist increases systolic BP and heart rate.

Solution to Question 10:

The depth of compressions in infants should be at least 1.5 inches or 1/3 the depth of the infant's chest.

Effective chest compressions in infants and pediatric population:

• 100 to 120 compressions per minute (30 compressions for every 15-18 sec).

• One provider: Place two fingers, one between the nipple line and the other 1 cm below the first finger

• Two providers: Encircle the infant's torso with both hands, thumbs pointing cephalad and positioned 1cm below the nipples over the sternum.

• Depth of compressions should be at least 1.5 inches or 1/3 the depth of the infant's chest.

• Press hard and fast

• Allow for full chest recoil after every compression



Other Options:

Option A: Chest compression at the rate of 30:2 with 2 rescuers: For children, the compression-to-ventilation ratio is 15:2 when there are two rescuers. A 30:2 ratio is used only if there is a single rescuer, similar to adult CPR.

Option B: Adenosine via interosseous is preferred over other methods: Adenosine is not used in pediatric cardiopulmonary resuscitation. It is used in PSVT treatment in intravenous routes.

Option D: Adrenaline given in dose 0.01 mg/kg iv/io in 1:1000 concentration: Epinephrine is the drug of choice for treating bradycardia in infants and children. The recommended dose is 0.01 mg/ kg of a 1:10,000 solution.

Solution to Question 11:

The body temperature in post cardiac arrest care is to be maintained between 32-36 degrees for at least 24 hours. This is known as targeted temperature managament.

Targeted temperature management (TTM) is the only intervention demonstrated to improve neurological recovery after cardiac arrest and involves maintaining core body temperature between 32-36°C for at least 24 hours.

This is done in post-cardiac resuscitation patients who have achieved a return of spontaneous circulation (ROSC) but are still comatose.

The entire body is to be cooled using a combination of methods like fluid-filled cooling blankets, ice packs, forced air blankets, cold IV fluids, and/or invasive devices or catheters.

It is the only intervention that can improve neurological recovery after cardiac arrest. It is most beneficial in ventricular fibrillation and pulseless ventricular tachycardia patients.

Cardiac arrest (especially with shockable rhythms) and neonatal hypoxic-ischemic encephalopathy are the primary established indications for therapeutic hypothermia. Stroke (in some cases), where thrombectomy is not an option, is another indication.



Other Options:

Option A: Maintain MAP over 65: A MAP of ≥ 65 mmHg is the minimum threshold recommended during post-cardiac arrest care to prevent ischemic injury and support organ function.

Option B: Maintain SpO2 above 94%: The recommended target SpO■ range in post-cardiac arrest care is 92-98%.

Option D: Revascularisation procedure if the patient develops ST elevation post resuscitation: Post-cardiac arrest patients with STEMI benefit from revascularization, as it reduces mortality and improves cardiac function.

Solution to Question 12:

The Jackson Rees circuit is used in paediatric anaesthesia.

Jackson Rees modification of Ayre's T piece is classified as a Mapleson's F system (Semi-Closed Circuits).

It is used in children <6 years or <20 kgs.



Other options:

Option B, C, and D: Cardiac anaesthesia, Neuro anaesthesia, and Obstetric anaesthesia require closed circuit. They have a provision to absorb CO_2, using soda lime. Closed circuits are used for long periods of procedures.

Mapleson Circuit	Usage
Type A (Magill's circuit)	Circuit of choice for spontaneous ventilation
Type B	Obsolete
Type C (Water's to and fro)	Obsolete
Type D (Bain's circuit)	Circuit of choice for controlled ventilation. Most commonly used circuit.
Type E (Ayre's T piece)	Circuit of 2nd choice in children
Type F (Jackson Rees Modification)	Circuit of choice in children (<20 kg)

Anesthesia NEET 2024

Question 1:

A patient undergoing surgery in the prone position is maintained on mechanical ventilation with PEEP. What is the significance of using PEEP in this scenario?

a) Prevents alveoli collapse and atelectasis of lung

b) Decreases inspiratory pressure

c) Increases venous return and maintains cardiac output

d) Reduces CO_2 in alveoli

Question 2:

Which of the following statements are true about eutectic mixture of local anesthetic (EMLA)?

a) 1gm of 0.5% of cream contains 50mg lignocaine

b) Methohemoglobinemia is a side effect

c) It is an equal mixture

d) It is made of Lignocaine and ropivacaine

Question 3:

An anesthetist used the airway device shown in the image given below on a patient. Which among the following statements is true regarding the adjunct?

a) It can cause damage to the teeth

b) It can be used in conscious patients

c) It does not cause laryngospasm

d) It can be safely used in facial trauma cases

Question 4:

In the following condition as shown in the radiograph, which of the following procedures would be difficult to perform? "



a) Nasogastric tube insertion

b) Laryngeal mask airway

c) Tracheostomy

d) Indirect laryngoscopy

Question 5:

Which of the following agents is appropriate for reversing vecuronium when renal function tests are normal?

a) Physostigmine

b) Rivastigmine

c) Sugammadex

d) Succinyl choline

Question 6:

A COVID-19 patient admitted to the ICU on mechanical ventilation suddenly develops hypoxia. The patient has decreased breath sounds on the left side and increased intrathoracic pressure. What is the most likely diagnosis?



a) Pneumothorax

b) Left-Sided Pleural Effusion

c) Acute Respiratory Distress Syndrome

d) Left-Sided Pneumonia

Question 7:

A 3-year-old child presents with drowsiness and refusal to feed. On examination, the child is in a stuporous state. The ECG is shown below. What is the next best step in the management of this patient?

a) Immediate synchronized cardioversion @ 1 joule/kg

b) Defibrillate, check rhythm and then start CPR

c) Defibrillate followed by CPR for 2min, and check rhythm

d) Stop CPR and do cardioversion at 2 joules/ kg

Question 8:

At what pressure does the marked structure open?



a) 30-40 cm H2O

b) 30-40 mm Hg

c) 10-20 cm H2O

d) 10-20 mm Hg

Question 9:

This recovery position is appropriate for:



a) Unresponsive patient, pulse and breathing absent

b) Responsive patient, pulse and breathing present

c) Unresponsive patient, pulse and breating present

d) Unresponsive with pulse, without breathing

Question 10:

Following is the image of a normal capnograph. What is true about the segment 3-4?



a) Inhalation of fresh gas

b) Exhalation of dead space air

c) Exhalation of mixed gases

d) Alveolar gas and plateau pressure

Question 11:

What is the procedure being done in the image?



a) Bier Block

b) Axillary nerve block

c) Median nerve block

d) Wrist block

Question 12:

What is true about the following position?



a) High rate of error

b) Ideal position for bougie insertion in emergencies

c) Ideal postion for neurosurgery

d) Sniffing position for intubation

Question 13:

A patient presents with stridor after prolonged intubation. The endoscopic view is shown in the image below. What is the staging system for this condition?

a) Cotton Meyer's

b) AJCC

c) Radkowski

d) TNM

Question 14:

A patient is undergoing implant surgery for his upper limb. On giving general anaesthesia he develops tachycardia, sudden raise in body temperature and rigidity of muscles. What drug is most likely to have caused this?

a) Propofol

b) Etomidate

c) Nitrous oxide

d) Halothane

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	c
5	c
6	a
7	c
8	a
9	c
10	d
11	a
12	d
13	a
14	d

Detailed Explanations

Solution to Question 1:

The significance of using positive end-expiratory pressure (PEEP) in the given clinical scenario is to help keep the alveoli open, preventing collapse (atelectasis), which is especially important in the prone position where certain lung regions might be more susceptible to collapse.

This is crucial in conditions where the lungs are prone to collapse, such as Acute Respiratory Distress Syndrome (ARDS) or when the patient is in a prone position during surgery. By maintaining open alveoli, PEEP improves gas exchange, which increases oxygen levels in the blood.

Other options:

Option B: Decreases inspiratory pressure: PEEP can increase the overall inspiratory pressure by maintaining pressure in the lungs at the end of expiration.

Option C: Increases venous return and maintains cardiac output: PEEP can decrease venous return due to increased intrathoracic pressure, which may reduce cardiac output.

Option D: Reduces CO_2 in alveoli: PEEP does not directly reduce CO_2 in the alveoli. The effect on CO_2 depends on ventilation and perfusion matching, not just PEEP alone.

Solution to Question 2:

Methemoglobinemia can develop when high doses of prilocaine are administered, typically over 600 mg.

EMLA (Eutectic Mixture of Local Anaesthetic) is a topical mixture containing prilocaine. The risk of methemoglobinemia is increased in neonates with metabolic disorders or in those receiving drugs that can impair the reduction of methemoglobin.

It is an eutectic(in fixed proportions) mixture, not necessarily equal (Option C). It consists of a 1:1 mixture of 2.5% lidocaine (Option A) and 2.5% prilocaine, not ropivacaine (Option D)

EMLA can provide adequate dermal analgesia for procedures like:

• Split skin-graft harvesting

• Venipuncture

• Intravenous cannulation in children

• Laser skin treatments

• Lithotripsy

• Circumcision

EMLA cream should not be used on mucous membranes, broken skin, infants younger than 1 month of age, or patients with a predisposition to methemoglobinemia. EMLA is applied 45-60 minutes before the procedure with an occlusive dressing. It provides analgesia up to a depth of 5 mm, which lasts for 1-2 hours.

Other side effects include- blanching of the skin, erythema, and oedema.

Solution to Question 3:

The image shows a nasopharyngeal airway and its insertion. The true statement is that it can be used in conscious patients.

A nasopharyngeal airway is used to provide artificial airway passage to the hypopharynx. To place a nasopharyngeal airway, the device should be lubricated and inserted into the nose with the bevel facing the nasal septum, at a 90-degree angle to the longitudinal axis of the body. It bypasses the gag reflex as they do not go through the mouth. Thus, it can be used in conscious patients or those who have been lightly anesthetized.

Care should be taken to avoid excessive force as damage to the nasal mucosa can result in epistaxis. The nasal approach should be avoided in conditions that put patients at higher risk for epistaxis. These conditions include but are not limited to coagulopathy, pregnancy, hereditary hemorrhagic telangiectasia, and uncontrolled hypertension. Absolute contraindications include basilar skull fractures, facial trauma, midface disruption, and injury to the roof of the mouth.

An oropharyngeal airway is used to maintain the airway in anesthetized patients. This adjunct prevents the tongue and epiglottis from falling back against the posterior pharynx and blocking

the airway. They can cause coughing, retching, or laryngospasm in conscious patients.

To place an oropharyngeal airway, the device is inserted with the curved portion facing downwards. It is then rotated 180 degrees to be put in place. The below image shows the oropharyngeal airway:



Solution to Question 4:

Tracheostomy would be difficult to perform in this case of pneumothorax with tracheal deviation.

The trachea is displaced from its usual midline position, making it challenging to accurately locate and access the trachea for the incision. The tracheostomy procedure requires precise placement to ensure proper airway management and avoid complications.

Steps of Tracheostomy:

• Incision: Make a vertical incision in the midline from the cricoid cartilage to above the sternal notch. For elective procedures, a 5 cm transverse incision two fingers' breadth above the sternal notch can be used for a better cosmetic outcome.

• Dissection: Dissect tissues in the midline. Displace or ligate any dilated veins encountered to control bleeding.

• Muscle Separation: Separate and retract the strap muscles of the neck laterally to expose underlying structures.

• Thyroid Isthmus Management: Displace the thyroid isthmus upwards or divide and ligate it with sutures.

• Local Anesthesia: Inject a few drops of 4% lignocaine into the trachea to suppress coughing.

• Tracheal Incision: Fix the trachea with a hook and make a vertical incision between the third and fourth tracheal rings, converting it into a circular opening. Avoid dividing the first tracheal ring to prevent perichondritis and stenosis.

• Inserting the Tube: Insert a tracheostomy tube of appropriate size into the tracheal opening and secure it with tapes.

• Avoiding Tight Suturing: Avoid tightly suturing or packing the skin incision to prevent subcutaneous emphysema.

• Dressing: Place a gauze dressing between the skin and the flange of the tracheostomy tube around the stoma to absorb secretions, provide a barrier against infection, and allow air to escape, promoting proper healing

Other Options:

Option A: Nasogastric tube insertion: Targets the esophagus, so tracheal deviation has minimal impact.

Option B: Laryngeal mask airway (LMA): Positioned over the larynx, not directly affected by tracheal deviation.

Option D: Indirect laryngoscopy: Focuses on the larynx and vocal cords, and is less influenced by the trachea's position.

Solution to Question 5:

Sugammadex is used for immediate reversal of vecuronium-induced neuromuscular blockade/relaxant in the given scenario.

Sugammadex is a modified γ-cyclodextrin and the first selective relaxant–binding agent based on an encapsulating principle for the inactivation of nondepolarising neuromuscular blocking drugs mainly rocuronium and vecuronium. This principle of complex formation of sugammadex and rocuronium or vecuronium in a 1:1 ratio occurs at all levels of neuromuscular blockade resulting in a more fast-acting pharmacologic reversal when compared with anticholinesterase drugs like neostigmine.

Other options:

Options A and B: Physostigmine and Rivastigmine are acetylcholinesterase inhibitors, which are not effective for reversing vecuronium.

Option D: Succinylcholine is a neuromuscular blocker itself and would not reverse vecuronium; it is used to induce paralysis rather than reverse it.

Solution to Question 6:

This patient in the above clinical scenario has developed a sudden onset of hypoxia, decreased breath sounds on one side, and increased intrathoracic pressure, along with an X-ray showing hyper lucency in the left side with tracheal and mediastinal shift to the right side is suggestive of a diagnosis of Pneumothorax.

A pneumothorax occurs when air enters the pleural space, causing lung collapse and preventing normal lung expansion, which leads to absent or diminished breath sounds over the affected area. Pneumothorax can cause a rapid increase in intrathoracic pressure, which impairs lung expansion

and leads to significant hypoxia. In patients on mechanical ventilation, it delivers positive pressure to the lungs, which can exacerbate or reveal underlying pleural issues. Increased positive pressure can contribute to the development or worsening of a tension pneumothorax.

Other options:

Option B: Left-Sided Pleural Effusion: While this condition can also cause decreased breath sounds and hypoxia, it typically does not cause acute increases in intrathoracic pressure.

Option C: Acute Respiratory Distress Syndrome (ARDS): ARDS is characterized by diffuse lung inflammation and can lead to hypoxia. However, it generally does not cause localized decreased breath sounds.

Option D: Left-Sided Pneumonia: Pneumonia can cause hypoxia and decreased breath sounds but is less likely to be associated with a sudden increase in intrathoracic pressure.

Solution to Question 7:

The given ECG showing polymorphic ventricular tachycardia, which is a shockable rhythm, in a stuporous paediatric patient warrants immediate defibrillation followed by CPR for 2 minutes.

Polymorphic ventricular tachycardia, especially in a paediatric patient, is often associated with severe underlying conditions and is a life-threatening rhythm disturbance. Immediate defibrillation is critical for restoring a normal rhythm.

After defibrillation, it is advisable to do CPR for 2 minutes before reassessing the rhythm for the next round of shock to ensure adequate blood circulation and oxygenation. .



Note: Cardioversion is not suitable for polymorphic VT; defibrillation is the preferred approach in such cases.

Solution to Question 8:

In the image, the marked structure is the adjustable pressure-limiting (APL) valve of Mapelson-C anaesthetic breathing system and it will open at the pressure of 30-40 cm H2O. This valve is designed to release gas when a certain pressure is exceeded to prevent excessive pressure from building up in the patient's airway.

The APL valve typically opens at pressures of 30-40 cm H■O to allow for safe ventilation without causing barotrauma to the lungs. This range is appropriate for manual ventilation to prevent high airway pressures that could lead to lung injury.



Solution to Question 9:

The position illustrated in the picture is known as the recovery position, and it is used in the management of patients who are unresponsive but with pulse present and breathing present.

The recovery position can help prevent complications such as aspiration and airway obstruction in unconscious individuals. Additionally, it is crucial to continuously monitor the patient's airway, breathing, and circulation while in recovery and seek immediate medical assistance. An unresponsive patient who is breathing spontaneously in the supine position is at risk of airway obstruction from their tongue, mucus, vomit, etc. A modified lateral position is used and the following 6 principles are considered while choosing which variation of recovery position to use,

•

The victim should be in as near a true lateral position as possible, with the head dependent to allow free drainage of fluid.

•

The position should be stable.

•

Avoid any pressure on the chest that impairs breathing.

•

It should be possible to turn the victim on his or her side and to return to the back easily and safely, with concern for a possible cervical spine injury.

•

Good observation of and access to the airway should be possible.

•

The position itself should not cause injury to the victim.

Other options:

Option A: Unresponsive with pulse and breathing absent - In this situation, the patient is in cardiac arrest and requires immediate cardiopulmonary resuscitation (CPR), not the recovery position. CPR involves chest compressions and rescue breaths to support circulation and respiration.

Option B: Responsive patient with pulse and breathing present- The patient should be monitored until the emergency response team arrives.

Option D: Unresponsive with pulse present and breathing absent - In this case, the patient is experiencing respiratory arrest. Rescue breathing or bag-valve-mask ventilation should be initiated to support the patient's breathing. The recovery position is not indicated in this scenario.

Solution to Question 10:

In a capnographic waveform, the plateau phase corresponds to alveolar air and plateau pressure.

The normal waveform has four different phases:

Phase I: Inspiratory baseline, represents the CO_2-free gas from the airways (anatomical and apparatus dead space).

Phase II: Expiratory upstroke, mixing of dead space gas with alveolar gas.

Phase III: Alveolar plateau, represents CO_2-rich gas from the alveoli. The last of the alveolar gas is sampled to measure $ETCO_2$.

Phase 0: Inspiratory downstroke and the beginning of inspiration.

Alpha angle: The transition from phase II to III.

The alpha angle can be used to assess the ventilation/perfusion of the lung. V/Q mismatches will have an alpha angle greater than 90 degrees.

Beta angle: The transition from phase III to 0.

The beta angle can be used to assess rebreathing. If rebreathing occurs, the angle is greater than 90 degrees.



Solution to Question 11:

The position in the image of a hand with a tourniquet occluded limb shows a Bier Block.

IVRA or Bier's block involves the intravenous administration of a local anesthetic into a tourniquet-occluded limb. It is used to provide anesthesia for short surgical procedures (<90 minutes) on the limbs.

Other options:

Option B: The axillary nerve block provides anesthesia for short surgical procedures on the hand, wrist, or forearm. It is performed by using a ultrasound-guided needle in the proximal axilla between the tendon of the lattisimus dorsi and the teres major.

Option C: Median nerve blocks are used for procedures involving the palmar aspect of the arm or the carpal tunnel. For a median nerve block at the elbow, the probe is placed in the elbow crease over the proximal radius and the anesthetic is injected aiming for the fascial plane between flexor digitorum superficialis and flexor digitorum profundus.

Option D: For a wrist block, the arm is abducted with the hand in supination, following which the anesthetic is infiltrated in the dorsal aspect of the wrist. This is used for short procedures of the wrist and the fingers.

Note: Bupivacaine is not used for Biers block. It is the most cardiotoxic local anesthetic. Accidental early deflation of the tourniquet can lead to massive systemic doses of bupivacaine resulting in cardiac arrhythmias and cardiovascular collapse.

Solution to Question 12:

The above image shows the sniffing position used for intubation, which is the true statement.

The sniffing position is a common head and neck alignment technique used during intubation to optimize the view of the glottis for easier insertion of the endotracheal tube. It is achieved by flexing the neck and extending the head, resembling the posture one assumes when sniffing. The position aligns the oral, pharyngeal, and laryngeal axes to provide a straight line of sight, making the glottis more visible during direct laryngoscopy and intubation.

Steps of intubation using the sniffing position:

• Direct laryngoscopy is performed in the sniffing position, with the head elevated 5-10 cm above the surgical table, the cervical spine flexed, and the atlantooccipital joint extended.

• The epiglottis is lifted up by the tip during laryngoscopy with Miller's blade.

• While performing laryngoscopy, the scope is lifted upwards and never levered over the upper incisors.

Other options:

Option A: The sniffing position is the most commonly used position in anesthesia as it ensures patient safety by ensuring a straight line of sight, allowing for better outcomes of the procedure.

Option B: The sniffing position is not the ideal position for bougie insertion during emergencies as there is concerns for cervical spine trauma in patients in the emergency department.

Option C: The sniffing position is not the typical position for neurosurgery. Depending on the location of the surgery, different head and neck positions are used.

Solution to Question 13:

This patient presenting with stridor after prolonged intubation most likely has subglottic stenosis post intubation. This condition is staged by the Cotton Meyer's grading.

Subglottic stenosis following prolonged intubation is a narrowing of the airway just below the vocal cords (subglottic region), which can lead to difficulty breathing. This condition is often a complication of endotracheal intubation due to the pressure exerted by the tube or cuff on the airway mucosa, leading to injury and scarring.

These patients present with stridor, dyspnea, hoarseness or changes in voice, and difficulty in breathing.

Diagnosis is by laryngoscopy or bronchoscopy which allows direct visualization of the subglottic area to assess the degree of stenosis.

CT or MRI may be used to evaluate the extent of narrowing and identify any underlying structural damage.

Treatment can be via endoscopic balloon or mechanical dilation, laser therapy or surgical resection may be needed to remove the stenosis in severe cases. In severe cases, a tracheostomy may be required to bypass the obstructed segment and allow for breathing.



Solution to Question 14:

In the clinical scenario given above, the development of tachycardia and muscle rigidity soon after administration of anesthetic agents can be due to malignant hyperthermia. Halothane is the anesthetic agent associated with this condition.

Malignant hyperthermia is a rare genetic hypermetabolic disease of the muscle. It occurs due to a mutation of the gene for the ryanodine receptor (Ryr 1), located on chromosome 19. It is an ion channel responsible for calcium release from the sarcoplasmic reticulum.

The symptoms develop when the patient is exposed to suxamethonium/succinylcholine or other inhaled anesthetics like halothane, isoflurane, sevoflurane, desflurane, etc. The mutated ryanodine receptor leads to a sudden release of calcium from the sarcoplasmic reticulum of the skeletal muscle. This uncontrolled increase in intracellular calcium removes the inhibition of troponin, resulting in sustained muscle contraction, and increasing adenosine triphosphatase (ATPase) activity. This results in an uncontrolled hypermetabolic state with increased oxygen consumption and CO_2 production, producing severe lactic acidosis and hyperthermia.

The key clinical features are:

• Succinylcholine-induced masseter muscle rigidity (causing trismus) or generalized muscle rigidity

• Tachycardia

• Hypercarbia (increased CO_2 production) and reduced oxygen saturation

• Hyperthermia (rise of 1° Celsius can occur every 5 minutes)

• Arrhythmia

• Hypertension

• Myoglobinuria and myoglobinemia indicating muscle damage

• Metabolic derangements like hyperkalemia and mixed metabolic and respiratory acidosis

Treatment is to discontinue the triggering agents i.e. succinylcholine followed by administration of 100% oxygen. Metabolic acidosis and hyperkalemia need to be corrected. Hyperthermia must be controlled by cooling the patient. Dantrolene sodium is the drug of choice. It is given at a dose of 2.5 mg/ kg intravenously and can be repeated every 5-10 minutes till the attack subsides.

Dermatology AIIMS 2017

Question 1:

The histopathological slide given below is suggestive of:



a) Psoriasis vulgaris

b) Leishmaniasis

c) Mycosis fungoides

d) Pemphigus vulgaris

Question 2:

A 23-year-old patient came with complaints of non-scarring alopecia in the vertex region of his head. Histological examination shows loss of hair with haemorrhage into the follicular areas and mild perinuclear lymphocytic infiltrate. What is the diagnosis?

a) Alopecia areata

b) Telogen effluvium

c) Tinea infection

d) Trichotillomania

Question 3:

Which of the following diseases is unlikely to present with skin lesions on the shin?

a) Sarcoidosis

b) Diabetes

c) Hyperthyroidism

d) Hypothyroidism

Question 4:

Identify the layer of skin (in the image given below) in which desmosomes are abundant.



a) B

b) A

c) D

d) C

Question 5:

The following image depicts:



a) Scabies

b) Ichthyosis

c) Contact dermatitis

d) Urticaria

Question 6:

A female came with complaints of bone and joint pain. NSAIDs were prescribed. After a few days, she presents again complaining vesicobullous lesions healing to give a brownish discoloration over the face (upper lip area) as shown in the image. This most probably suggests a diagnosis of

a) Malaria

b) Dengue

c) Chikungunya

d) Fixed drug eruption

Question 7:

Dense irregular collagen is found in:

a) Ligaments

b) Dermis

c) Tendons

d) Bones

Question 8:

A patient presented with hair loss and itching of the scalp along with mild discharge. Which of the following is useful for diagnosis?

a) Tzanck smear

b) Gram stain

c) Slit skin smear

d) KOH mount

Question 9:

Identify the condition shown below:



a) Epidermal verrucous nevus

b) Congenital melanocytic nevus

c) Melanoacanthoma

d) Malignant melanoma

Answer Key

Question No.	Correct Option
1	d
2	d
3	d
4	c
5	a
6	d
7	b
8	d
9	b

Detailed Explanations

Solution to Question 1:

The given slide shows a row of tombstones appearance, which is classically seen in pemphigus vulgaris.

In this condition, there is a suprabasal splitting of the epidermis leading to blister formation. The basal layer remains adherent to the basement membrane and gives the resemblance to a row of tombstones as shown in the histology image below.



Pemphigus vulgaris is a chronic immunobullous disorder characterized by the presence of antibodies against desmosomal proteins, desmogleins 1 and 3 expressed in the skin and mucosa.

Clinically, the patients present with ill-defined, irregular, slow-to-heal oral lesions and erosions. Flaccid blisters filled with clear fluid may arise on normal skin or an erythematous base. The rupture of such blisters produces painful erosions and heals without scarring.

Nikolsky's sign is positive which is elicited by applying firm tangential sliding pressure with a finger on lesional or perilesional skin. This leads to the separation of the normal■looking epidermis from the dermis, producing erosion.

Direct immunofluorescence is considered the most accurate confirmatory test of this condition. It shows intercellular IgG and C3 deposition on the surface of keratinocytes throughout the epidermis in normal, perilesional skin, in a characteristic fishnet/chickenwire appearance as seen below.



Solution to Question 2:

The clinical scenario depicting a young adult presenting with non-scarring alopecia in the vertex region and histology showing perifollicular hemorrhage is suggestive of trichotillomania (trichotillosis).

Trichotillomania has to be differentiated from alopecia areata (Option A). Trichotillomania presents with incomplete hair loss within a patch with varying lengths of hair whereas alopecia areata shows smooth hairless patch with sparing of white hairs. Regular pitting of nails and exclamation mark hairs are seen in alopecia areata. When the diagnosis is uncertain, a scalp biopsy can be done. Perifollicular hemorrhage is characteristic of trichotillomania while peribulbar lymphocytic infiltrate is diagnostic of alopecia areata.

Trichotillomania is classified under obsessive-compulsive and related disorders. It is characterized by recurrent body-focused repetitive behavior (hair pulling) and repeated attempts to decrease or stop the behavior. It is commonly seen in females. The hair loss is commonly seen over the vertex but can involve eyelashes, eyebrows, and beard.

The most effective treatment for trichotillosis is habit-reversal training (HRT).

Other options:

Option B: Telogen effluvium is diffuse hair loss resulting in non-scarring alopecia due to loss of telogen hairs due to fever, pregnancy, hypothyroidism or anemia.

Option C: Tinea infection shows patchy hair loss, resulting in non-scarring/scarring alopecia. Easy pluckability, broken hairs and wood's lamp positivity are features.

Solution to Question 3:

Hypothyroidism is unlikely to present with skin lesions on shin. Myxedema (severe hypothyroidism) and pretibial myxedema are different entities. Pretibial myxedema is the infiltrative dermopathy (mucin deposition) seen over shin in Grave's disease, a type of hyperthyroidism.

Other options :

Option B: Diabetes: Necrobiosis lipoidica diabeticorum is classically seen on the anterior leg.

Option A: Sarcoidosis: Erythema nodosum may be seen on the anterior leg.



Solution to Question 4:

Desmosomes are abundant in the stratum spinosum (Option C, label D in the image) layer of the epidermis of the skin.

Desmosomes are adhesion complexes in epidermis which link adjacent keratinocytes. They appear like spines on microscopy and hence this layer is also called the prickle layer. Defects of desmosomal proteins, inherited or acquired, result in blistering disorders.



Solution to Question 5:

The given image shows raised, brownish thread-like lesions in the interdigital distribution (burrows), typically seen in scabies.

Scabies is an ectoparasitic infection caused in humans by Sarcoptes scabiei. It is transmitted by close personal contact. Pruritis with nocturnal exacerbation and contact cases within a family should raise the suspicion of scabies. Typical locations of lesions are the finger webs, the flexor surfaces of the wrists, the elbows, the axillae, the buttocks and genitalia and the breasts of women (Circle of Hebra). In adults, the face is often spared. Pruritic papules and nodules on male genitalia and breasts serve as valuable diagnostic clues in the absence of burrows.

Dermoscopy of the burrows shows jet■with■contrail appearance. Microscopy of scraping from burrows can be done to demonstrate mites.

A single overnight application of 5% permethrin cream from neck to toe in adults and from head to toe in children is the drug of choice for the treatment of scabies in all ages including infants (at least 2 months old) and pregnant women. All family members, even if asymptomatic, need to be treated simultaneously for this condition to prevent reinfection. Oral ivermectin can also be used.

Solution to Question 6:

The patient has a history of consumption of NSAIDs, following which there is an appearance of vesiculobullous lesions on the lips, evolving to give a brownish discolouration. The most probable diagnosis is fixed drug eruption (FDE).

A strong differential diagnosis is post-chikungunya pigmentation (PCP) of the face seen on the nose and cheeks (chik sign). However, this is not the answer because there is no antecedent history of fever, myalgia or other symptoms suggestive of chikungunya. PCP often appears as the fever subsides and may persist for about 3-6 months. PCP mimics melasma and may be misdiagnosed if proper history is not taken.

The image below shows chik sign:



Fixed drug eruption (FDE) is a type of delayed hypersensitivity reaction. It is characterized by well-defined brown, hyperpigmented plaques that may follow vesiculobullous lesions. They occur on lips, hands, legs, face, genitalia, and oral mucosa, and can cause a burning sensation. They usually present 30 min to 8 h after drug exposure and recur at the same location on re-exposure. NSAIDs, Paracetamol and Co■trimoxazole are the most common drugs causing FDE.

Solution to Question 7:

Dense irregular collagen is found in dermis.

Irregular connective tissue is of three types: Loose, dense and adipose. Dense connective tissue is located in regions which are under consistent mechanical stress and where protection is required for the ensheathed organs. The abundant collagen fibres in the dermis run in a haphazard manner. This protects the skin.

It is usually present in :

• Reticular layer of the dermis

• Superficial connective tissue sheaths of the muscle and nerves

• Adventitia of large vessels

• Capsules of various organs and glands (e.g, testis, sclera, periosteum, perichondrium)

On the other hand, dense regular collagen fibres are found in tendons.

Solution to Question 8:

The patient has a history of itching and mild discharge and examination shows a well-demarcated patch of alopecia, with scaling and broken hair shafts. This is characteristic of tinea capitis. Though the diagnosis is clinical, a KOH mount is used for confirmation.

Scrapings from the edge of a lesion are transferred to a slide, to which KOH is added. It is then examined under the microscope for the presence of hyphae, as shown below.



Tinea capitis is a dermatophyte infection primarily occurring in children. Clinical features of tinea capitis are patchy hair loss, scaling, tender lymphadenopathy, and in severe cases, there might be pustules and crusting of the affected area along with sinus formation.

The clinical variants of tinea capitis are:

• Endothrix - a non-inflammatory type in which multiple black dots are present within the areas of alopecia. It is most commonly caused by T. tonsurans and T. violaceum.

• Ectothrix - a mild inflammatory type in which single or multiple scaly patches with hair loss are seen. Microsporum audouinii, M.canis, M.equinnum, and M.ferrugineum cause ectothrix infection.

• Kerion - a severe inflammatory type that presents with an inflammatory mass with thick crusting and matting of hair with pus and sinus formation. It is commonly caused by T. verrucosum and T. mentagrophytes.

• Favus - is characterized by the presence of yellowish, cup■shaped crusts known as scutula that develop around hair follicles. It is caused by T. schoenleinii

Treatment:

• Terbinafine: <10 Kg- 62.5 mg; 10-20 kg - 125 mg; >20 kg- 250 mg, all given daily for 4 weeks.

• Itraconazole: 2-4 mg/kg/day for 4-6 weeks.

Solution to Question 9:

The image shows a child with a well-circumscribed, black, heterogenous lesion covered with hair. This is characteristic of a congenital melanocytic nevus (CMN). They may be present at birth or develop shortly thereafter. They are benign and can be brown, black, purplish, or red, and are usually a palpable lesion.

They can be classified based on size: small (<1.5 cm), medium (1.5-19.9 cm), and large or giant (>20 cm). Larger CMN have a higher risk of developing malignant melanoma.

Management of CMN depends on the size and location of the lesion. Smaller nevi may be observed, while larger nevi or those at a higher risk for complications may require surgical excision, particularly if there is a concern for malignant transformation.

Complications of CMN include neurological abnormalities (intraparenchymal melanosis is the commonest finding), malignant melanoma, and other tumors like rhabdomyosarcoma.

The images below show different melanotic skin lesions:

Melanoacanthoma

Malignant melanoma

Epidermal verrucous naevus

Dermatology AIIMS 2018

Question 1:

Which of the following nerves is biopsied for the diagnosis of pure neuritic leprosy?

a) Median nerve

b) Superficial branch of radial nerve

c) Ulnar nerve

d) Lateral popliteal nerve

Question 2:

Which component of cement causes allergic dermatitis among construction workers in India?

a) Cobalt

b) Nickel

c) Iron

d) Chromate

Question 3:

Which of the following is a drug used in the treatment of tinea cruris for which widespread resistance is found to be developing?

a) Itraconazole

b) Griseofulvin

c) Terbinafine

d) Voriconazole

Question 4:

A patient came with a history of fever, body ache, and joint pain. He was given NSAIDs. A few days later, he developed brownish pigmentation over the nose. What is the diagnosis?

a) Melasma

b) Fixed drug eruption (FDE)

c) Dengue

d) Chikungunya

Question 5:

A mother has a pearly white papule on her forehead (as shown in the image below) which her infant also develops. What is the cause?

a) HPV

b) Pox virus

c) Adenovirus

d) Herpes virus

Question 6:

A disease associated with the occurrence of blistering and peeling of over >30% of body surface area is usually caused by:

a) Viral infection

b) Drugs

c) Fungal infection

d) Auto-immunity

Question 7:

A 23 year old patient presents with lesions in the groin as shown in the image below. Which of the following cannot be a cause of these lesions?

a) Trichophyton

b) Microsporum

c) Aspergillus

d) Epidermophyton

Question 8:

A patient presents with multiple nodulocystic lesions on the face. Which among the following is the drug of choice in this condition?

a) Steroids

b) Isotretinoin

c) Erythromycin

d) Tetracycline

Question 9:

What is the type of connective tissue present in the area marked by the arrow in the given image?



a) Dense regular

b) Dense irregular

c) Loose irregular

d) Specialised

Question 10:

The histopathological slide given below is suggestive of:



a) Psoriasis vulgaris

b) Leishmaniasis

c) Mycosis fungoides

d) Pemphigus vulgaris

Answer Key

Question No.	Correct Option
1	b
2	d
3	c
4	d
5	b
6	b
7	c
8	b
9	b
10	d

Detailed Explanations

Solution to Question 1:

Superficial sensory branch of the radial nerve is most commonly used for nerve biopsy in neuritic leprosy.

Although any nerve involved would show the pathology, it is prudent to biopsy a branch of a sensory nerve, rather than a mixed nerve or major nerve trunk. So, the ulnar nerve will not be the best answer despite it being the most common peripheral nerve involved in neuritic leprosy.

The common nerves and sites where nerve biopsy is performed in leprosy are as follows:

• Superficial sensory radial nerve branch at the wrist

• Ulnar cutaneous nerve near the ulnar border of the base of the hand

• Sural nerve near the ankle

Note: Nerve biopsy is the gold standard for confirmation of pure neuritic leprosy.

Solution to Question 2:

Chromate is the most frequent allergen in cement causing occupational contact dermatitis. The most common allergen causing allergic contact dermatitis in India is nickel. It is commonly found in artificial jewelry. However, among construction workers, chromate is more common than nickel. The reported prevalence of allergic contact dermatitis (ACD) to chromate among construction workers is usually more than 10%.

Allergic contact dermatitis (ACD) is an inflammatory condition caused due to a delayed hypersensitivity reaction (Type 4 hypersensitivity) to a specific allergen that comes into direct contact with the skin. Some common allergens include nickel (found in artificial jewelry), PPD/Para phenylene diamine (in hair dye), chromate (in cement), rubber or latex, cosmetics, etc. Patch test is used for the diagnosis of ACD.

Patients usually present with pruritus, erythema, swelling, papules, and papulovesicles. In severe cases, large vesicles or blisters may form due to the disruption of the intercellular bridges. If exposure persists the affected area becomes dry, scaly, and thicker with lichenification and fissuring develops later. There is involvement of the covered areas as well as the exposed areas.

Solution to Question 3:

The use of terbinafine as the agent of choice for tinea cruris and corporis has been limited lately due to the development of resistance.

Tinea cruris (also known as jock itch) is a dermatophyte infection involving the crural fold. The most common cause is Trichophyton rubrum.

T. rubrum isolates have been found to exhibit resistance against terbinafine.

Note: The question is based on research papers about Trichophyton rubrum strains exhibiting primary resistance to terbinafine.

Solution to Question 4:

The given history of fever, joint pain, and centrofacial hyperpigmentation points toward a diagnosis of chikungunya. Post chikungunya pigmentation (PCP) usually appears after the fever subsides and persists for about 3-6 months. Hyperpigmentation associated with chikungunya fever is macular and most commonly affects the nose and cheeks.

Another close differential is fixed drug eruptions (FDE). FDE has a shorter latency, usually presents 30 min to 8 h after drug exposure, and recurs at the same location on re-exposure. It presents with hyperpigmented plaques that may be preceded by vesiculobullous lesions.

Hyperpigmentation of the centrofacial region (nose and cheeks) mimics melasma and may be misdiagnosed if proper history is not taken.

Nose pigmentation is striking in cases of chikungunya fever and is termed chik sign. The image below shows chik sign on the nose:

FDE is a type of delayed hypersensitivity reaction. They occur on lips, hands, legs, face, genitalia, and oral mucosa, and can cause a burning sensation. NSAIDs, paracetamol and co■trimoxazole are the most common drugs causing FDE. NSAID■induced FDE mainly affects the genitals and lips.

The image below shows FDE on upper lip area:

Solution to Question 5:

The image shows pearly white umbilicated wart-like lesions with a characteristic dimple at the center, suggestive of molluscum contagiosum, which is caused by pox virus.

The age of peak incidence is between 2 and 5 years. In children, it is mainly caused by the type I molluscum contagiosum virus (MCV-1). There is a relatively higher incidence of MCV■2 type in

adults and HIV infection. Sexually transmitted molluscum contagiosum may be present over the genital area.

It has an incubation period of 2 weeks to 6 months. Widespread and refractory lesions are seen most commonly in HIV or patients on immunosuppressive therapy.

It is characterized by the presence of intracytoplasmic eosinophilic inclusion bodies called Henderson-Patterson bodies or molluscum bodies. Pseudo-Koebner phenomenon is seen, that is, new lesions develop along lines of trauma.

Treatment is usually not required, and lesions are likely to resolve spontaneously. Stimulation of the immune response can be done after destructive or inflammatory therapies with salicylic acid, trichloroacetic acid, and imiquimod. Cryotherapy when done needs to be repeated at 3-4 week intervals. Extensive lesions are treated with an antiviral agent like cidofovir.

Solution to Question 6:

Blistering and peeling involving greater than 30% of the body surface area points to the diagnosis of toxic epidermal necrolysis (TEN). TEN is primarily drug■induced, with a culprit drug being demonstrated in approximately 85% of cases.

Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) are both severe mucocutaneous reactions, characterized by blistering and epithelial sloughing. The two terms describe phenotypes within a severity spectrum, where SJS is the less extensive form and TEN is the more extensive

Common drugs that can cause SJS/TEN are:

• Allopurinol

• NSAIDs- oxicam group

• Anti-epileptics - phenobarbital, phenytoin, carbamazepine, lamotrigine

• Antibiotics - penicillins, cephalosporins, sulfamethoxazole, sulfasalazine, nevirapine

Disease	Epidermal detachment	Purpuric macules and atypical target lesions
SJS	<10%	Present
Overlap SJS-TEN	10–30%	Present
TEN with spots	>30%	Present
TEN without spots	>30% (large sheets of epidermal loss)	Absent

Solution to Question 7:

The image shows a well-circumscribed, annular plaque in the groin region which is characteristic of tinea cruris. Tinea cruris is a dermatophytosis. Trichophyton, Microsporum and Epidermophyton are dermatophytes whereas Aspergillus is not.

Dermatophytes include 3 genera:

• Trichophyton species infect skin, hair, and nails and include T. rubrum, T. mentagrophytes, T. tonsurans, T. schoenleinii, and T. violaceum.

• Microsporum contains M. gypseum, M. canis, and M. nanum. These infect only hair and skin (not nails).

• Floccosum is the only species in the genus Epidermophyton, It infects only skin and nails (not hair).

Aspergillus spp. produce large necrotic lesions like ecthyma gangrenosum, and sometimes small papules and cold abscesses.

Solution to Question 8:

The presence of multiple nodulocystic lesions implies that this is a case of severe acne. Isotretinoin is the first-line drug for the treatment of severe acne (nodular acne and acne conglobata).

Systemic retinoids are highly teratogenic and are absolutely contraindicated in pregnant women, those planning to become pregnant, or breastfeeding mothers. Two negative pregnancy tests (on separate occasions) are required for females of childbearing age before starting acitretin or isotretinoin.

Baseline evaluation of serum lipids, transaminases, and complete blood counts should be done before starting any systemic retinoids. These values should be checked monthly for the first 3–6 months and once every 3 months thereafter. Monthly pregnancy tests should be done while on the drug.

Note: Female patients should avoid conception for 3 years after discontinuing acitretin treatment for cutaneous psoriasis.

Solution to Question 9:

The area marked is the reticular dermis. It is made up of dense irregular connective tissue.

It is found in the regions that are under considerable mechanical stress and where protection is given to ensheathed organs. It is usually present in:

• Reticular dermis

• Superficial connective tissue sheaths of the muscle and nerves

• Adventitia of large vessels

• Capsules of various organs and glands (e.g., testis, sclera, periosteum, perichondrium)

On the other hand, dense regular collagen fibers are found in tendons.

Solution to Question 10:

The given slide shows a row of tombstones appearance, which is classically seen in pemphigus vulgaris.

In this condition, there is a suprabasal splitting of the epidermis leading to blister formation. The basal layer remains adherent to the basement membrane and gives the resemblance to a row of tombstones as shown in the histology image below.



Pemphigus vulgaris is a chronic immunobullous disorder characterized by the presence of antibodies against desmosomal proteins, desmogleins 1 and 3 expressed in the skin and mucosa.

Clinically, the patients present with ill-defined, irregular, slow-to-heal oral lesions and erosions. Flaccid blisters filled with clear fluid may arise on normal skin or an erythematous base. The rupture of such blisters produces painful erosions and heals without scarring.

Nikolsky's sign is positive which is elicited by applying firm tangential sliding pressure with a finger on lesional or perilesional skin. This leads to the separation of the normal■looking epidermis from the dermis, producing erosion.

Direct immunofluorescence is considered the most accurate confirmatory test of this condition. It shows intercellular IgG and C3 deposition on the surface of keratinocytes throughout the epidermis in normal, perilesional skin, in a characteristic fishnet/chickenwire appearance as seen below.



Intercellular deposition of immune complexes giving a "chicken wire/fish net" look

Dermatology AIIMS 2019 (May & Nov)

Question 1:

A 5-year-old boy had painful sores in the mouth with a rash over the hands along with fever. Which of these is the causative agent?

a) Pox virus

b) Coxsackievirus

c) HHV-7

d) Parvovirus

Question 2:

The following lesion is seen in a 25-year-old man who came to a primary health centre 5 days after he had unprotected sexual intercourse. On palpation the ulcers were painful. Suppurative inguinal lymphadenopathy was present on the right side. The patient has no previous history of drug allergy. Which kit will be used for the management of this particular sexually transmitted disease?

a) Kit 6/Yellow

b) Kit 2/green

c) Kit 3/white

d) Kit 1/grey

Question 3:

A patient with fever and joint pain developed pigmentation on the nose after a few days of taking NSAIDs as shown below. What is the diagnosis?



a) Melasma

b) Fixed drug eruption

c) Lichen planus

d) Chikungunya

Question 4:

Which cell is not present in the layer marked D?



a) Keratinocyte

b) Merkel cell

c) Melanocyte

d) Langerhans cells

Question 5:

Topical steroids are most effective in:

a) Dermal atrophy

b) Eczematous dermatitis

c) Bullous lesions due to HSV

d) Rosacea

Question 6:

You have ordered a patch test for an 8-year-old boy with a history of allergic contact dermatitis. After the application of the patch, when should you ask him to return for removal?

a) 24 hours

b) 72 hours

c) 48 hours

d) 96 hours

Question 7:

A patient was treated with steroids for psoriasis. On stopping the treatment, he develops fever, malaise, and lesion as seen in the image below. What is the most probable diagnosis?

a) Pustular psoriasis

b) Staphylococcal infection

c) Acute generalised exanthematous pustulosis

d) Subcorneal pustulosis

Question 8:

Which of the following does not present with maculopapular rash?

a) Zoster (shingles)

b) Rubella

c) Dengue (during febrile phase)

d) Measles

Question 9:

A patient presents with flaccid bullous lesions involving the oral cavity and the skin. He has lesions as shown below. Acantholytic cells are seen on Tzanck smear. What is the most probable diagnosis?

a) Pemphigus foliaceus

b) Pemphigus vulgaris

c) Dermatitis herpetiformis

d) Bullous pemphigoid

Question 10:

A 10-year-old child presents with hair loss. The hair loss is patchy as shown in the image below. Black dots are noted in areas of hair loss. Which of the following types of tinea is this child suffering from?

a) Ectothrix

b) Endothrix

c) Kerion

d) Favus

Answer Key

Question No.	Correct Option
1	b
2	c
3	d
4	d
5	b
6	c
7	a
8	a
9	b
10	b

Detailed Explanations

Solution to Question 1:

The presence of fever, along with painful sores in the mouth and small, tense vescicles with surrounding erythema on the palms and soles in a child point towards the diagnosis of hand-foot-mouth disease. The most common cause of which worldwide is coxackievirus A16.

Spread is by droplets or faecal contamination and the incubation period is 7 days. The fever is generally mild, and is followed by the appearance of lesions. The characteristic lesions are slightly oval, with a grey blister roof and a narrow rim of erythema. The oral lesions are aphthoid and are less painful than aphthous ulcers.

Only symptomatic treatment is given. The lesions generally disappear in 7 - 10 days.

Solution to Question 2:

The appearance of painful genital ulcers, along with unilateral suppurative inguinal lymphadenopathy, 5 days after unprotected sexual intercourse best fits the symptom cluster of a non - herpetic genital ulcer. The treatment of which is kit 3 / white kit in case of no penicillin allergy.

The above-shown clinical image and the findings are consistent with a diagnosis of chancroid.

Soft Chancre/Chancroid:

• The causative organism is Haemophilus ducreyi.

• The incubation period is 3 to 10 days.

• It presents with multiple painful ulcers on genitals.

• It may be accompanied by unilateral painful, suppurative inguinal lymphadenopathy.

• Treatment: Inj. Benzathine penicillin 2.4 million units + tab. Azithromycin 1 gm single dose

Note: In the syndromic approach to sexually transmitted infections, kit assignment based on the identification of a cluster of symptoms rather than making an actual diagnosis is given emphasis.

Solution to Question 3:

The given clinical history of fever and joint pain and the image showing hyperpigmented macules on nose (chik sign) is suggestive of Chikungunya.

Post-chikungunya pigmentation (PCP) most commonly affects the nose and cheeks. It often appears as the fever subsides and may persist for about 3-6 months.

PCP mimics melasma and may be misdiagnosed if proper history is not taken.

Another close differential is fixed drug eruptions (FDE). FDE is a type of delayed hypersensitivity reaction. It is characterized by well-defined brown, hyperpigmented plaques that may follow vesiculobullous lesions. They occur on lips, hands, legs, face, genitalia, and oral mucosa, and can cause a burning sensation.They usually present 30 min to 8 h after drug exposure and recur at the same location on re-exposure. NSAIDs, Paracetamol and Co■trimoxazole are the most common

drugs causing FDE.

The image below shows FDE on upper lip area:

Solution to Question 4:

The layer marked as D is stratum basale. Langerhan's cells are not present in stratum basale. Langerhan's cells are predominantly present in stratum spinosum.



Keratinocytes, melanocytes, and Merkel discs are all seen in stratum basale:

(Option A) Keratinocytes are predominant cells of the epidermis and constitute the majority of the stratum basale.

(Option B) Merkel discs are type 1 slow-adapting touch receptors.

(Option C) Melanocytes are dendritic cells which are derived from the neural crest cells. The melanosomes synthesized by the melanocytes migrate to the adjacent keratocytes through the dendritic processes

Solution to Question 5:

Topical steroids are the drug of choice in the treatment of eczematous dermatitis.

Other options:

• Steroids cause atrophy and hence is contraindicated in patients with dermal atrophy.

• Herpes infection is associated with immunosuppression so steroids are not useful.

• Routine steroid use not recommended in rosea as it can worsen the condition.

Solution to Question 6:

In patch testing, the patch is removed after 48 hours to examine the site of reaction.

Patch testing is a test to detect sensitivity to a specific allergen. A battery of suspected allergens is applied to the patient's back under occlusive dressings or patches. It is allowed to remain in contact with the skin for 48 hours.

• 48 hours: Patch removal for initial reading to detect delayed hypersensitivity reaction.

• 72-96 hours: Final reading done for confirmation.

Note: Since the question is regarding the timing of patch removal and not regarding the final diagnosis, 48 hours is the answer.





Solution to Question 7:

The given clinical scenario points towards the diagnosis of pustular psoriasis.

Pustular psoriasis is a type of psoriasis characterized by tiny pus-filled lesions (pustules) having a characteristic lakes or sheets of pus appearance (seen in the image).

The patient presents with a history of fever and malaise along with the presence of pustules on a circumscribed, fiery red, edematous or scaling plaque. The pus is usually sterile. Sudden withdrawal of systemic steroids is usually the trigger for this condition.

Pustular psoriasis can be divided into:

• Localized pustular psoriasis

• Generalized pustular psoriasis:

• Acute generalized pustular psoriasis (Von Zumbusch)

• Subacute annular and circinate pustular psoriasis

• Acute generalized pustular psoriasis of pregnancy (Impetigo herpetiformis)

Treatment: Acitretin (oral retinoid) is the drug of choice for pustular psoriasis. However, it is contraindicated in pregnancy. Systemic steroids are used instead. Cyclosporine is the second-line drug.

Note: Systemic steroids are never used for the treatment of psoriasis except in the case of Impetigo herpetiformis where they are first-line agents.

Solution to Question 8:

A maculopapular rash is not seen in zoster (shingles).

Shingles cause vesicular lesions that are grouped together and specific to a dermatome.

Vesicular lesions grouped together in a dermatome

Maculopapular rashes are seen in:

• Rubella

• Dengue (febrile phase)

• Measles

• Chikungunya (acute infectious phase)

• Erythema infectiosum

• Zika virus

• West Nile virus

• Infectious mononucleosis

Type of rash	Description
Macule	Flatlesion with a well-circumscribed change in skin color < 1 cm
Papule	Elevated solidskin lesion < 1 cm
Vesicle	Smallfluidcontaining blister

Solution to Question 9:

Flaccid lesions, mucosal involvement, and the presence of acantholytic cells are suggestive of pemphigus vulgaris.

Option A: There is no mucosal involvement in pemphigus foliaceus.

Option C: Dermatitis herpetiformis presents with multiple small vesicles and no acantholysis.

Option D: Bullous pemphigoid presents with tense bullae and no acantholysis.

Solution to Question 10:

The given clinical scenario depicting a young child with patchy hair loss and the image showing black dots is suggestive of endothrix type of tinea capitis.

Tinea capitis is a dermatophyte infection primarily occurring in children. Clinical features of tinea capitis are patchy hair loss, scaling, tender lymphadenopathy, and in severe cases, there might be pustules and crusting of the affected area along with sinus formation.

The clinical variants of tinea capitis are:

• Endothrix - a non-inflammatory type in which multiple black dots are present within the areas of alopecia. It is most commonly caused by T. tonsurans and T. violaceum.

• Ectothrix -a mild inflammatory type in which single or multiple scaly patches with hair loss are seen. Microsporum audouinii, M.canis, M.equinnum, and M.ferrugineum cause ectothrix infection.

• Kerion -a severe inflammatory type that presents with an inflammatory mass with thick crusting and matting of hair with pus and sinus formation. It is commonly caused by T. verrucosum and T. mentagrophytes.

• Favus - is characterized by the presence of yellowish, cup■shaped crusts known as scutula that develop around hair follicles. It is caused by T. schoenleinii

Treatment:

• Terbinafine: <10 Kg- 62.5 mg; 10-20 kg - 125 mg; >20 kg- 250 mg, all given daily for 4 weeks.

• Itraconazole: 2-4 mg/kg/day for 4-6 weeks.

Dermatology AIIMS 2020 (May and Nov INI-CET)

Question 1:

A female patient presents with active genital warts with a history of recurrence, but her husband doesn't have it. What advice will you give to prevent transmission?

a) Give continuous prophylaxis to the patient

b) Give prophylactic anti-virals to the husband

c) Give anti-virals whenever active lesions are present

d) Don't involve in intercourse when active lesions are present

Question 2:

A male baby born to a non-consanguineous couple develops blisters on areas of friction. Their previous baby also had similar complaints and died at the age of 2 weeks. What is the diagnosis?

a) Epidermolysis bullosa

b) Chronic bullous disorder of childhood

c) Congenital syphilis

d) Neonatal pemphigus

Question 3:

A rose gardener with a history of thorn prick presents with the following lesions. What is the possible causative agent?

a) Streptococcus pyogenes

b) Sporothrix schenkii

c) Strongyloides stercoralis

d) Staphylococcus aureus

Question 4:

A school-going child who had applied henna comes to dermatology OPD with the following lesions. What is the diagnosis?

a) Contact dermatitis due to henna

b) Dermatitis artefacta

c) Polymorphous light eruption

d) Vitiligo

Question 5:

A 40-year-old woman who presented with complaints of fever and joint pain develops the following lesion on her nose after a few days of taking NSAIDs. What is your diagnosis?



a) Dengue

b) Chikungunya

c) Melasma

d) Fixed drug eruption

Question 6:

Methylisothiazolinone (MI) in cosmetics is used as?

a) Surfactant

b) Humectant

c) Emulsifier

d) Preservative

Question 7:

The given condition is caused by which drug?

a) Actinomycin

b) Bleomycin

c) Mitomycin C

d) Doxorubicin

Question 8:

Which of the following does not present with maculopapular rash?

a) Zoster (shingles)

b) Rubella

c) Dengue (during febrile phase)

d) Measles

Question 9:

A child presents to OPD with tense bullae over the torso. Biopsy of the lesion showed a subepidermal level of blistering and neutrophil infiltration. What is the drug of choice?

a) Rituximab

b) Dapsone

c) Cyclosporine

d) Azathioprine

Question 10:

What is the probable diagnosis of this condition affecting hair?



a) Trichorhexis nodosa

b) Trichorexis invaginata

c) Trichothiodystrophy

d) Monilethrix

Question 11:

A patient with a known case of chronic plaque psoriasis with more than 50% skin involvement presents to you. Which of the following will you not recommend?

a) Oral methotrexate

b) NB UVB

c) Oral steroids

d) Oral cyclosporine

Question 12:

Identify the type of skin lesion.

a) Nummular

b) Annular

c) Discoid

d) Target

Answer Key

Question No.	Correct Option
1	d
2	a
3	b
4	b
5	b
6	d
7	b
8	a
9	b
10	c
11	c
12	b

Detailed Explanations

Solution to Question 1:

The advice to prevent transmission is to not involve in intercourse when active genital warts are present.

Genital warts are caused by HPV and transmitted through the sexual route. Antivirals are not very effective against anogenital warts. Biological treatment options like podophyllin, imiquimod, 5-fluorouracil are given. The best way to prevent transmission of infection is by avoiding sexual intercourse when active lesions are present.

Solution to Question 2:

The clinical history given in the question points to a diagnosis of epidermolysis bullosa congenita.

Epidermolysis bullosa is a mechanobullous disease characterized by blistering in areas of mechanical trauma. Blisters are commonly seen in trauma-prone areas like hands, feet, elbows, and knees. There are three types of epidermolysis bullosa:

• Epidermolysis bullosa simplex - defective K5/K14 protein

• Junctional epidermolysis bullosa - defective Laminin

• Dystrophic epidermolysis bullosa - defective Collagen-7

The image below shows epidermolysis bullosa lesions on the foot:

Other options:

Option B: Chronic bullous disorder of childhood - It typically presents with bullous lesions having a string of pearls/cluster of jewel appearance (image below). It is an IgA-mediated immunobullous disorder. It manifests in early childhood and is not a congenital condition.

Option C: Congenital Syphilis - Syphilitic pemphigus presents as a vesiculobullous disorder. The investigation of choice is VDRL.

Option D: Neonatal Pemphigus - It is an autoimmune blistering disease caused by the transfer of maternal IgG autoantibodies to desmoglein 3 to the neonate through the placenta when the mother is affected with pemphigus. It is clinically characterized by transient flaccid blisters and erosions on the skin and, rarely, the mucous membranes.

Solution to Question 3:

The given image shows the distribution of lesions along lymphatics. This distribution along with the history of thorn pick is suggestive of the causative agent Sporothrix schenkii.

Sporothrix schenkii causes rose gardener's disease. It is a saprophyte that lives on grass and plants. Hence a history of thorn prick in farmers or gardeners is an important clue.

Other causes of skin lesion distributed along lymphatics is fish tankgranuloma caused by Mycobacterium marinum.

Solution to Question 4:

The likely diagnosis in this case is dermatitis artefacta (factitious dermatitis).

Dermatitis artefacta is a psychodermatological excoriation disorder with a female preponderance, commonly affecting adolescents and young adults, where elaborate techniques (using sharp objects, cigarettes, lighters, chemicals etc.) are used to self induce skin lesions.

The morphology of the lesions are varied, with oddly linear, clear-cut and angular edges, surrounded by normal looking skin. These lesions appear with no prodromal stage or associated symptoms as they are self inflicted. They are distributed over easy to reach areas of the body and have a varied appearance based on the mode of injury.

• Option A: Contact dermatitis is due to contact with allergic agents such as paraphenylenediamine (PPD). Presentation can vary from erythema, swelling, papules, vesicles and in severe cases, blisters on the areas of contact with the allergen (henna, in this case). There exists a sequence of inflammatory changes here, unlike the lesions in dermatitis artefacta.

• Option B: Polymorphic light eruptions as the name suggests are seen over sunlight-exposed areas of the body. The lesions are polymorphic and are symmetrically distributed.

• Option C: Vitiligo is characterized by patchy involvement of skin with hypopigmentation.

Solution to Question 5:

The given history of fever and joint pain along with centrofacial hyperpigmentation shown in the image point towards a diagnosis of chikungunya. Hyperpigmentation associated with chikungunya fever is macular and most commonly affects the nose and cheeks.

Nose pigmentation is striking in cases of Chikungunya fever and is termed as Chik sign. It appears as the fever subsides and persists for about 3-6 months.

Hyperpigmentation of the centrofacial face (nose and cheeks) mimics melasma and may be misdiagnosed if proper history is not taken.

Another close differential is fixed drug eruptions (FDE). FDE is a type of delayed hypersensitivity reaction. It is characterized by well-defined brown, hyperpigmented plaques that may follow vesiculobullous lesions. They occur on lips, hands, legs, face, genitalia, and oral mucosa, and can cause a burning sensation. They usually present 30 min to 8 h after drug exposure and recur at

the same location on re-exposure. NSAIDs, Paracetamol and Co■trimoxazole are the most common drugs causing FDE.

The image below shows FDE on upper lip area:

Solution to Question 6:

Methylisothiazolinone (MI) in cosmetics is used as a preservative to inhibit bacterial growth. It is permitted to be used up to the level of 100 ppm.

Several cases of allergic contact dermatitis due to MI were reported due to its presence in wet wipes, hair cosmetics, facial cosmetics, deodorants, and sunscreens.

Furthermore, cases of airborne exposure to MI causing severe airborne and generalized dermatitis from recently painted walls and even from toilet cleaners have been described.

Solution to Question 7:

The given image shows flagellate dermatitis. It is caused by bleomycin.

Flagellate dermatitis starts as bands of pruritic, flagellate, erythema over the trunk, and extremities within few hours to 6 months after administration. The rash heals with persistent flagellate hyperpigmentation.

No treatment is necessary since the lesions usually disappear when bleomycin is discontinued. If the patient is itchy, managing pruritus with an antihistamine may limit trauma to the skin, and thus avoid further hyperpigmentation.

Another side effect that one needs to watch out for with the use of bleomycin is the development of pulmonary fibrosis.

Solution to Question 8:

A maculopapular rash is not seen in zoster (shingles).

Shingles cause vesicular lesions that are grouped together and specific to a dermatome.

Maculopapular rashes are seen in:

• Rubella

• Dengue (febrile phase)

• Measles

• Chikungunya (acute infectious phase)

• Erythema infectiosum

• Zika virus

• West Nile virus

• Infectious mononucleosis

Type of ra sh	Description
Macule	Flatlesion with a well-circum scribed change in skin color < 1 cm
Papule	Elevated solidskin lesion < 1 cm
Vesicle	Smallfluidcontaining blister

Solution to Question 9:

This clinical scenario with tense blisters showing subepidermal separation and neutrophilic infiltration at the basement membrane points towards a diagnosis of IgA pemphigoid or linear IgA disease or chronic bullous disease of childhood. The drug of choice is dapsone. It is bacteriostatic against M. leprae.

It also has an anti-inflammatory action by decreasing the action of neutrophil enzymes. Hence, it is useful in the treatment of IgA pemphigoid, linear IgA disease and chronic bullous disease of childhood.

Solution to Question 10:

The given clinical image of hair seen under polarising microscopy reveals tiger tail banding (alternating light and dark bands) characteristic of trichothiodystrophy.

Trichothiodystrophy is an autosomal recessive condition causing defective DNA repair. It causes brittle sulfur-deficient hair, photosensitivity, ichthyosis and systemic involvement. Unlike xeroderma pigmentosum (a similar condition with defective DNA repair), there is no risk of skin cancer.

Option A: Trichorrhexis nodosa is associated with node formation and fractures. The classic appearance, shown in the image below, is known as thrust paintbrush appearance as if two paintbrushes are rubbing against each other. It is associated with trauma to the hair shaft and arginosuccinic aciduria.



B- Trichorrhexis invaginata or bamboo hair is characterized by invagination of the distal hair shaft into the proximal portion. It is seen in Netherton's syndrome. It is also called golf-tee or ball and socket appearance as seen below.



Option D: Monilethrix, seen in the image below, is characterized by beaded hair and is fragile at the constricted sites. It occurs as an isolated condition or can occur in Menkes disease.



Solution to Question 11:

Oral/systemic steroids are not used in the management of chronic plaque psoriasis. Topical steroids may be used.

Systemic steroids are avoided because the withdrawal of systemic steroids causes the development of pustular psoriasis.

The only indication of using systemic steroids in psoriasis is generalized pustular psoriasis in pregnancy (impetigo herpetiformis).

Other options maybe used in treatment of extensive (>30%) chronic plaque psoriasis.

Solution to Question 12:

The given image shows ring-shaped lesions with central clearing, which are characteristic of annular lesions.

In annular lesions, only the edges are active and appear raised, or in a different color. They are seen in tinea corporis, granuloma annulare, borderline leprosy, and psoriasis.

Dermatology NEET 2018

Question 1:

A 15-year-old girl presents with itchy lesions on her arm as shown. Her family history is positive for asthma. What could be the most probable diagnosis?

a) Seborrhoeic dermatitis

b) Atopic dermatitis

c) Allergic contact dermatitis

d) Erysipelas

Question 2:

Which of the following is a primary skin lesion?

a) Crust

b) Atrophy

c) Purpura

d) Induration

Question 3:

Which of the following statements is false regarding scleredema?

a) The dermis is 3–4 times thicker than normal

b) Erythema and peau d'orange appearance of the skin can be seen

c) Can occur in association with diabetes

d) Associated with sclerodactyly or Raynauds phenomenon

Question 4:

Identify the pigmentation abnormality in the image shown below.

a) Mongolian spot

b) Naevus of Ito

c) Congenital melanocytic naevus

d) Becker's naevus

Question 5:

What is the most common causative organism for verruca vulgaris?

a) Human papillomavirus-1

b) Human papillomavirus-3

c) Human papillomavirus-10

d) Human papillomavirus-2

Question 6:

Cutis marmorata occurs due to _____

a) Exposure to cold temperature

b) Humidity

c) Exposure to hot temperature

d) An adverse reaction to drugs

Question 7:

Varicella-zoster virus remains dormant in the:

a) Medulla oblongata

b) Dorsal root ganglion

c) Skin

d) Ventral root

Answer Key

Question No.	Correct Option
1	b
2	c
3	d
4	d
5	d
6	a
7	b

Detailed Explanations

Solution to Question 1:

The clinical scenario depicting pruritic lesions at a flexural site along with a family history of atopy (asthma) is suggestive of atopic dermatitis (AD) or atopic eczema (AE). The scratch marks and excoriations shown in the image are due to pruritis, which is the hallmark feature of

AD.

AD is a chronic relapsing inflammatory condition. It presents with erythematous macules, papules and vesicles that often start in early childhood, usually less than 2 years old. The lesions later become excoriated and lichenified. It is frequently associated with dryness of the skin and secondary infections. In infancy, the lesions initially appear on face (sparing the napkin area) and later on extensor aspect of knees and elbows as the child starts crawling. In later childhood and in adults, the principal site of involvement is flexures.

The filaggrin monomers in keratinocytes in the epidermis bind and aggregate keratin bundles and intermediate filaments to form the epidermal barrier. Any damage/dysfunction of this barrier is associated with an increased risk of developing AD. Damage to this barrier can be due to environmental factors (detergents, hard water, UV light) or local inflammatory reactions. Dysfunction of the barrier can be due to genetic mutation of the FLG gene leading to reduced filaggrin production or local immunomodulatory effects leading to reduced filaggrin expression.

Other options:

Option A: Seborrheic dermatitis is a chronic inflammatory skin condition characterized by erythema, pruritus, scaling, flaking, or crusting of the skin. It is often associated with Malasezzia fungal infection. A milder form of seborrheic dermatitis of the scalp is commonly known as dandruff.

Option C: Allergic contact dermatitis (ACD) is an inflammatory condition caused due to a delayed hypersensitivity reaction (Type 4 hypersensitivity) to a specific allergen that comes into direct contact with the skin.Patients usually present with pruritus, erythema, swelling, papules, and papulovesicles. Some common allergens include nickel (found in artificial jewelry), PPD/Para phenylene diamine (in hair dye), chromate (in cement), rubber or latex, cosmetics, etc. Patch test (image below) is used for the diagnosis of ACD.



Option D: Erysipelas is a bacterial skin infection caused by Streptococcus pyogenes (Group A Streptococcus) characterized by raised, erythematous sharply demarcated rapidly spreading skin lesions, along with systemic symptoms like fever and chills. The skin may also have a peau d'orange or orange peel-like appearance.

Solution to Question 2:

Purpura is a primary skin lesion. Crust, atrophy, and induration are secondary lesions.

It is a condition with red or purple discolored spots on the skin that do not blanch on applying pressure. The spots are caused by bleeding underneath the skin. It is seen:

• Vasculitis

• Thrombocytopenia

• Vitamin C deficiency

Purpura



Option A: Crusts or scabs consist of dried-up discharge from the lesions.

Crusts (Scabs)

Option B: Atrophy refers to loss of tissue. There may be fine wrinkling and increased translucency if the process is superficial.

Option D: Induration refers to dermal thickening that clinically presents as skin that feels thicker and firmer than normal on palpation.

Solution to Question 3:

Scleredema is not associated with sclerodactyly or Raynauds Phenomenon. Scleredema and scleroderma are different entities. In contrast to systemic sclerosis (scleroderma), scleredema is not associated with sclerodactyly, Raynaud phenomenon, nail fold capillary changes, or serum

autoantibodies.

In scleredema, the dermis is 3–4 times thicker than normal due to an increase in type 1 collagen synthesis by dysfunctional fibroblast. However, fibroblast proliferation is not seen.

Erythema and a peau d'orange appearance of the skin are commonly observed. The hands and feet are characteristically spared.

Diabetic scleredema is the most common type. In diabetic scleredema, the accumulation of collagen may be due to irreversible non■enzymatic glycosylation of collagen and resistance to degradation by collagenase. It can also occur post-streptococcal infections.

Solution to Question 4:

The given image showing hyperpigmented lesion on shoulder associated with hypertrichosis is suggestive of Becker's naevus.

It is an epidermal melanocytic naevus with aggregation of melanocyte clusters in the epidermis. It is often hairy. The onset is around adolescence when there is an increased sensitivity to androgens especially in the distribution of outer arms, scapular area and chest.

A triad of features can be seen:

• Hyperpigmentation

• Hypertrichosis

• Acne

Option A: Mongolian spots are slate blue areas in the lumbosacral region that spontaneously regress before puberty.

Option B: Naevus of Ito is unilateral hyperpigmentation in the acromioclavicular distribution. It is not known to have hypertrichosis or acne.

Option C: Congenital melanocytic nevi are common moles that become raised and have a cerebriform appearance and can be hairy. They can occur anywhere. The onset is at birth or soon after birth

Solution to Question 5:

Verruca vulgaris or common wart is most commonly caused by human papillomavirus-2 (HPV-2) but is also related to types 1, 4, 27, and 57.

Common warts are usually asymptomatic. They show acanthosis, hyperkeratosis, and papillomatosis and present as firm papules with a rough horny surface. The most commonly affected sites are on the backs of the hands and fingers, and, in children on the knees. Pseudo-Koebner phenomenon may be observed.

Treatment is usually with topical agents such as salicylic acid, caustics, glutaraldehyde, formalin, or with laser, cryotherapy, surgical removal.

verruca vulgaris.

Solution to Question 6:

Cutis marmorata refers to the pinkish-blue mottled or marbled appearance when subjected to cold temperatures.

It is seen throughout infancy and in 50% of children. It is caused by simultaneous dilation and contraction of the superficial capillaries. Rewarming restores the skin to normal. This is a physiological finding and requires no treatment. However, persistent cutis marmorata may be seen in trisomies 18 and 21.

Note: Cutis marmorata telangiectatica congenita is a distinct vascular developmental disorder and is easily distinguished as it is fixed (it does not disappear on rewarming), and it may be associated with a variety of other anomalies such as limb growth or renal anomalies.

Solution to Question 7:

Varicella-zoster virus (VZV) is a human herpesvirus that causes varicella (chickenpox) as a primary infection and then remains in a latent form in the dorsal root ganglia for a variable period after which it can reactivate to cause herpes zoster (shingles).

Dermatology NEET 2019

Question 1:

A 15-year-old girl presents with itchy lesions on her arm as shown. Her family history is positive for asthma. What could be the most probable diagnosis?

a) Seborrhoeic dermatitis

b) Atopic dermatitis

c) Allergic contact dermatitis

d) Erysipelas

Question 2:

Bindi Leukoderma is caused by which chemical?

a) Mono-benzyl ether of Hydroquinone (MBH)

b) Crocein Scarlet MOO and Solvent Yellow 3

c) p-phenylenediamine (PPD)

d) Para Tertiary butylphenol (PTBP)

Question 3:

The most common triggering factor of the given condition is _____

a) Vaccination

b) Malignancy

c) Drugs

d) Infection

Question 4:

A young female presented with vaginal itching and green frothy genital discharge. Strawberry cervix is seen on examination. What will be the drug of choice?

a) Doxycycline

b) Oral fluconazole

c) Metronidazole

d) Azithromycin

Question 5:

What is correct regarding the diagnosis based on the given image?

a) The lesions are not infectious

b) Trigeminal dermatome is most commonly affected

c) Anterior nerve roots are more commonly involved

d) Mucous membranes within the affected dermatomes are involved

Question 6:

What is the diagnosis seen in the image below?

a) Candidal paronychia

b) Staphylococcal scalded skin syndrome

c) Candidal intertrigo

d) Diabetic foot ulcer

Question 7:

Which of the following is commonly associated with the condition shown below?

a) Oil drop sign

b) Leukonychia striata

c) Pterygium of nails

d) Pitting of nails

Question 8:

Identify the condition given in the image :

a) Dermatomyositis

b) Acanthosis nigricans

c) Pityriasis rotunda

d) Melasma

Answer Key

Question No.	Correct Option
1	b
2	d
3	d
4	c
5	d
6	c

7	d
8	b

Detailed Explanations

Solution to Question 1:

The clinical scenario depicting pruritic lesions at a flexural site along with a family history of atopy (asthma) is suggestive of atopic dermatitis (AD) or atopic eczema (AE). The scratch marks and excoriations shown in the image are due to pruritis, which is the hallmark feature of AD.

AD is a chronic relapsing inflammatory condition. It presents with erythematous macules, papules and vesicles that often start in early childhood, usually less than 2 years old. The lesions later become excoriated and lichenified. It is frequently associated with dryness of the skin and secondary infections. In infancy, the lesions initially appear on face (sparing the napkin area) and later on extensor aspect of knees and elbows as the child starts crawling. In later childhood and in adults, the principal site of involvement is flexures.

The filaggrin monomers in keratinocytes in the epidermis bind and aggregate keratin bundles and intermediate filaments to form the epidermal barrier. Any damage/dysfunction of this barrier is associated with an increased risk of developing AD. Damage to this barrier can be due to environmental factors (detergents, hard water, UV light) or local inflammatory reactions. Dysfunction of the barrier can be due to genetic mutation of the FLG gene leading to reduced filaggrin production or local immunomodulatory effects leading to reduced filaggrin expression.

Other options:

Option A: Seborrheic dermatitis is a chronic inflammatory skin condition characterized by erythema, pruritus, scaling, flaking, or crusting of the skin. It is often associated with Malasezzia fungal infection. A milder form of seborrheic dermatitis of the scalp is commonly known as dandruff.

Option C: Allergic contact dermatitis (ACD) is an inflammatory condition caused due to a delayed hypersensitivity reaction (Type 4 hypersensitivity) to a specific allergen that comes into direct contact with the skin.Patients usually present with pruritus, erythema, swelling, papules, and papulovesicles. Some common allergens include nickel (found in artificial jewelry), PPD/Para phenylene diamine (in hair dye), chromate (in cement), rubber or latex, cosmetics, etc. Patch test (image below) is used for the diagnosis of ACD.



Option D: Erysipelas is a bacterial skin infection caused by Streptococcus pyogenes (Group A Streptococcus) characterized by raised, erythematous sharply demarcated rapidly spreading skin lesions, along with systemic symptoms like fever and chills. The skin may also have a peau d'orange or orange peel-like appearance.

Solution to Question 2:

Bindi leukoderma is caused by para tertiary butylphenol (PTBP) present in adhesive sticker bindi.

• This is called as Contact Leukoderma or Chemical Leukoderma in which the chemical increases sensitivity and induces apoptosis of the melanocytes in the area of contact.

• There is also evidence that azo dyes, by virtue of structural similarity to tyrosine, can directly inhibit melanogenesis by competing with that amino acid for binding with tyrosinase enzyme.

Common chemicals in the Indian set up which cause Chemical leukoderma are:

Bindi Leukoderma	Para Tertiary butyl phenol (P TBP)
Synthetic leather, rubber slip pers, Latex condoms, rubber gloves	Mono-benzyl ether of Hydroq uinone (MBH)-Rubber Antio xidant
Hair dye	p-phenylenediamine (PPD)
Alta (red colour like henna)	Crocein Scarlet MOO and Sol vent Yellow 3(azo dyes)

Solution to Question 3:

The given image shows characteristic target lesions of erythema multiforme. This most commonly occurs following a herpes simplex virus infection.

The common triggers are:

• Herpes simplex infection

• Mycoplasma infection

• Bacterial infections

• Fungal infections

• Vaccinations

• Malignancy- carcinoma, lymphoma, leukemia

• Drugs- sulfonamides, rifampicin, penicillins, hydantoin derivatives

• Miscellaneous- lupus erythematosus (Rowell Syndrome), sarcoidosis, polyarteritis nodosa

Erythema multiforme is best regarded as a self■limiting cytotoxic dermatitis resulting from cell■mediated hypersensitivity.

It presents clinically with a spectrum of macular, papular, or urticarial lesions, as well as the classic acral iris or 'target lesions'.

Target lesions are less than 3 cm in diameter, rounded, and have three zones:

• Central area of dusky erythema or purpura

• Middle paler zone of edema

• Outer ring of erythema with a well■defined edge

Lesions may involve the palms and the trunk. Oral and genital mucosal membranes lesions are associated with erosions

The image below shows target lesions.



Solution to Question 4:

Vaginal itching and green frothy discharge in a young female with strawberry vagina points towards trichomoniasis. Metronidazole is the drug of choice for trichomoniasis.

It is is the most common parasitic infection in humans in the industrialized world.

It is caused by Trichomonas vaginalis, an anaerobic, flagellated protozoa, exists only as a trophozoite (no cyst stage).

Diagnosis is made by detection of motile flagella during microscopic examination of wet mounts. Direct immunofluorescent antibody staining is more sensitive (70-90%) than wet mount.

Culture is done on Feinberg-Whittington media.

Treatment: DOC- Metronidazole 2 gm single oral dose. All sexual partners must be treated to prevent reinfection, especially from asymptomatic males. Tinidazole can also be used for the treatment of trichomoniasis.

The image below shows strawberry vagina seen in trichomoniasis:

Solution to Question 5:

The above image shows unilateral, painful, grouped vesicles in a single dermatome suggestive of herpes zoster or shingles. The mucous membranes within the affected dermatomes are involved in shingles.

Herpes zoster/shingles is an acute cutaneous segmental eruption due to the reactivation of latent varicella-zoster virus from dorsal root ganglia. An earlier infection with varicella is essential before zoster can occur.

The first symptom of zoster is usually pain, 1-3 days after which the lesions appear. Occasionally, the pain may not be followed by the eruption ('zoster sine eruptione'). The patients are infectious, both from the virus in the lesions and nasopharynx.

The most commonly involved dermatomes in decreasing order of frequency are thoracic > cervical > trigeminal > lumbosacral. The incidence of ophthalmic zoster increase with age. The posterior nerve roots and ganglia show inflammatory changes.

Herpes zoster ophthalmicus is a variant of zoster involving the ophthalmic division of the trigeminal nerve (V1). Herpes zoster oticus/Ramsay Hunt syndrome is another variant involving the geniculate ganglion and presents with ipsilateral facial nerve palsy, otalgia and vesicles over external ear.

Zoster is a self-limiting infection but it is painful and carries a risk of secondary infection and post-herpetic neuralgia. Analgesia and treatment of secondary infections are sufficient for mild infections.

Indications for antivirals in VZV (Acyclovir 800 mg):

• Painful zoster infections in adults

• Facial zoster at any age

• Immunocompromised

Solution to Question 6:

The condition shown in the image is candidal intertrigo.

It is a candidal infection of opposing skin surfaces, which presents as erythema with a little moist exudation starting deep in a fold. When the web spaces of the toes or fingers are affected, marked maceration with a thick, white, horny layer is usually prominent.

As the condition progresses, it spreads beyond the area of contact, usually developing the typical features of candidosis with a fringed, irregular edge and subcorneal pustules rupturing to give tiny erosions, and then further peeling of the stratum corneum.

Treatment:

Topical antifungal agents such as nystatin, ketoconazole, and clotrimazole are routinely used.

Solution to Question 7:

The condition shown in the image above is alopecia areata. Classically, it causes fine stippled pitting of the nails.

It is a chronic inflammatory disease which causes non-scarring patches of hair loss, especially in genetically predisposed.

The most characteristic lesion is a smooth circumscribed patch of alopecia, which has short extractable broken hairs called as exclamation mark hair. However, there is always sparing of white hairs in the region affected.

Other options :

Subungal oil drop sign is a highly specific pathognomonic sign seen in Nail psoriasis. It is a yellow-red discolouration in the nail bed resembling a drop of oil.

Leukonychia striata are lines that extend across the nail and lie parallel to lunula. They are seen in Hypoalbuminemic states such as nephrotic syndrome .

Nail Pterygium is a wing-shaped deformity of nail due to a central fibrotic band which divides the nail into two parts in the proximal.

• Dorsal pterygium is seen in lichen planus.

• Ventral pterygium is seen in trauma, systemic sclerosis, Raynaud phenomenon, lupus erythematosus, familial subungual pterygium, and infections.

Solution to Question 8:

The condition in the given image is acanthosis nigricans. It is characterized by thickened, gray-brown velvety plaques seen in flexural areas such as the back of the neck, axillae,

inframammary creases, waist, and groin.

The causal mechanism is due to the secretion of epidermal growth factors in the region involved. It can be benign or malignancy-associated.

• Benign - commonly associated with obesity or insulin resistance, usually mild.

• Malignancy - much less common, associated with gastric, lung, and uterine cancers.

Acanthosis nigricans can also be seen in the following:

• Drugs such as nicotinic acid used to treat hyperlipidemia

• Sudden onset of multiple seborrhoeic keratoses

Dermatology NEET 2020

Question 1:

Identify the drug used to treat the condition shown below:

a) Anti-fungal therapy

b) Anti-tubercular therapy

c) Topical steroids therapy

d) Anti-leprosy treatment

Question 2:

Identify the finding associated with the condition shown in the image below.

a) Isomorphic phenomenon

b) Meyerson phenomenon

c) Gottron's papule

d) Nikolsky's sign

Question 3:

Identify the condition shown in the image below:

a) Herpes labialis

b) Herpangina

c) Molluscum contagiosum

d) Impetigo

Question 4:

Which of the following drugs used as nail lacquer belongs to morpholines?

a) Amorolfine

b) Oxiconazole

c) Ciclopirox olamine

d) Tioconazole

Question 5:

A child was born with membranes around the body and had ectropion and eclabium. He is brought to the OPD with lesions covering his face, trunk, and extremities. Which of the following is an unlikely diagnosis?

a) Ichthyosis vulgaris

b) Lamellar ichthyosis

c) Bathing suit ichthyosis

d) Harlequin ichthyosis

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	a
5	a

Detailed Explanations

Solution to Question 1:

The given image shows an annular plaque with central scarring points to lupus vulgaris, which is treated using anti-tubercular drugs.

It is the most common cutaneous tuberculosis in India. The characteristic lesions are annular, infiltrated plaques with serpiginous edges, central atrophy and scarring and are commonly seen on face, buttocks and extremities.

On diascopy, apple jelly nodules are seen.

Histologically, tubercles are seen in the superficial dermis showing scanty or absent central caseation with surrounding epithelioid histiocytes and giant cells.

Other differential diagnoses for annular plaque include:

• Tinea corporis shows central clearing, peripherally raised scaly margins and presents with an itch.

• Cutaneous leishmaniasis lesions have central crusting and are seen on the face in patients from areas of high leishmaniasis prevalence.

Solution to Question 2:

The image shows plaque psoriasis of the dorsum of the hand, and isomorphic or Koebner phenomenon is associated with the condition.

Option B: Meyerson phenomenon refers to the formation of an eczematous ring around a melanocytic nevus.

Option C: Gottron's papules are violaceous, erythematous papules overlying the dorsal interphalangeal or metacarpophalangeal, elbow, or knee joints.

Option D: Nikolsky sign refers to easy peeling of skin on applying tangential pressure over a bony prominence and is classically seen in pemphigus and staphylococcal scalded skin syndrome.

Solution to Question 3:

The above image shows multiple vesicles & crusts clustered on an inflamed base around the lips which is characteristic of herpes labialis or cold sore.

It is commonly caused by herpes simplex virus type 1(HSV-1) and rarely by herpes simplex virus type 2 (HSV-2). It begins with a prodrome of pain, burning, and tingling followed by the appearance of erythematous papules. The papules then become vesicular and may ulcerate.

It is treated with oral acyclovir 200 mg and 5% topical acyclovir.

Option B: Herpangina or vesicular pharyngitis is characterized by the presence of ulcerating small vesicles on the oral fauces and the posterior pharyngeal wall. It is commonly caused by the coxsackie virus.



Option C: Molluscum contagiosum is characterized by pearly white, wart-like umbilicated papules. It is caused by the pox virus.

Option D: Impetigo is characterized by golden yellow or honey-coloured crusts that form after the rupture of pustulating vesicles. Non-bullous impetigo is caused by both S. aureus and streptococcus. Bullous impetigo is caused by S. aureus.

Solution to Question 4:

Amorolfine is a morpholine that is available as a 5% nail lacquer. It is used in the treatment of onychomycosis.

Options B and D: Oxiconazole and tioconazole are classified as imidazoles

Option C: Ciclopirox olamine is a hydroxypyridone compound.

Solution to Question 5:

The clinical picture shows a neonate encased in a shiny, parchment-like membrane (collodion baby) suggestive of a disorder belonging to the autosomal recessive congenital ichthyosis spectrum.

Ichthyosis vulgaris is not a part of this disease spectrum. It does not present at birth and is therefore an unlikely diagnosis.

Ichthyosis vulgaris is an autosomal dominant condition caused due to a mutation in the filaggrin gene (FLG). It develops in the first few months of life. Patients exhibit accentuated palmar and plantar creases and have dry scaly skin. The scales appear light grey and cover the extensor surfaces of the limbs and the trunk. It usually spares the groin and larger flexures. Immunohistochemistry reveals an absent or markedly reduced filaggrin signal.

Autosomal recessive congenital ichthyosis spectrum includes the following:

- Harlequin ichthyosis (HI)
- Bathing suit ichthyosis (BSI)
- Lamellar ichthyosis (LI)
- Congenital ichthyosiform erythroderma (CIE)
- Self- improving congenital ichthyosis (SICI)
- Transient manifestations, such as collodion baby.

Dermatology INI-CET 2021

Question 1:

A forest officer develops the lesion as shown in the image. Which of the following is not a differential to consider?

a) Scrub typhus

b) KFD

c) Cutaneous anthrax

d) Healing brown recluse spider bite

Question 2:

A patient presents with following lesions on the genitals. Scrapings from the lesion would show?

a) Tzanck cells

b) Cells with clear halo around the nucleus.

c) Molluscum bodies.

d) HP bodies.

Question 3:

A young girl presents with leukotrichia and lesions as shown in the image. What is the most likely diagnosis?



a) Focal vitiligo

b) Segmental vitiligo

c) Piebaldism

d) Nevus depigmentosus

Question 4:

The most specific and typical sign of juvenile dermatomyositis in a 17-year-old patient is

a) Photosensitive skin rash.

b) Nail bed capillary changes.

c) Gottron's papules.

d) Malar rash.

Question 5:

A child presents to you with itchy lesions on the scalp with hair loss as shown in the image below. What would be the next best step?

a) KOH mount

b) Slit skin mount

c) Gram staining

d) Biopsy

Question 6:

The below-given image is characteristically seen in which of the following type of leprosy?



a) Borderline lepromatous leprosy

b) Lepromatous leprosy

c) Borderline leprosy

d) Histoid leprosy

Question 7:

A child presented with itchy plaques over the neck, the bilateral popliteal and cubital fossa. What could be the diagnosis?

a) Atopic dermatitis

b) Dermatitis herpetiformis

c) Pemphigus vegetans

d) Psoriasis

Question 8:

A 65-year-old man presents with itchy tense bullae over the limbs. The roof of the bullae collapsed after a few days. A clinical image is shown below, what is the likely diagnosis?

a) Bullous pemphigoid

b) Pemphigus vulgaris

c) Dermatitis herpetiformis

d) Linear IgA dermatosis

Question 9:

A young female presented with the following findings on the face. Which among the following will not be useful for its treatment?

a) Topical benzoyl peroxide

b) Topical fluocinolone

c) Topical tazarotene

d) Topical antibiotics

Question 10:

A 45-year-old male patient from Bihar presented with normoanesthetic hypopigmented lesion as shown below. He had a history of fever for a long duration in his childhood. No thickened nerves were present. What is the probable diagnosis?

a) Tuberculoid leprosy

b) Lepromatous leprosy

c) Mycosis fungoides

d) Post kala-azar dermal leishmaniasis

Question 11:

A child presented with asymptomatic lesions on the forearm and on the shaft of the penis. The lesions on the forearm are shown below. What is the most likely diagnosis?

a) Lichen planus

b) Lichen nitidus

c) Scrofuloderma

d) Scabies

Question 12:

Which among the following is correct regarding the healing of leprosy if left untreated?

a) No spontaneous remission is seen

b) Half of the cases resolve spontaneously after bacilli exposure

c) Spontaneous remission is more common in children than in adults

d) Indeterminate leprosy have more remission than tuberculoid

Question 13:

A young man presented with painful lesions as shown in the image below. What is the likely diagnosis?

a) Herpes Zoster

b) Molluscum contagiosum

c) Herpes Simplex

d) Lymphangioma circumscriptum

Answer Key

Question No.	Correct Option
1	b
2	a
3	b
4	c
5	a
6	c
7	a
8	a
9	b
10	d
11	b
12	d
13	a

Detailed Explanations

Solution to Question 1:

The given image shows the presence of eschar and Kyasanur Forest Disease (KFD) is not a differential diagnosis for the presence of eschar.

Kyasanur Forest Disease is a hemorrhagic fever found in the state of Karnataka, India. KFD has an abrupt onset of fever, headache, conjunctivitis, myalgia, and severe prostration. Characteristic skin lesions that are associated with KFD are petechial lesions. Some cases may develop hemorrhages in the skin, mucosa, and viscera, but there is no eschar formation.

Infection is transmitted by the bite of ticks, the principal vector being Haemaphysalis spinigera

Other options:

A) Scrub typhus:

• Scrub Typhus is also known as chigger-borne typhus.

• Scrub typhus is caused by Orientia tsutsugamushi.

• The vector is trombiculid mite.

• Humans are infected when they are bitten by the mite larvae known as chiggers.

The incubation period is 1- 3 weeks. The patient typically develops a characteristic eschar at the site of the mite bite as shown in the image below, with regional lymphadenopathy and maculopapular rash.

C) Cutaneous anthrax:

• In cutaneous anthrax, the lesion starts as a papule 1-3 days after infection and becomes vesicular, containing fluid that may be clear or blood-stained.

• The whole area is congested and edematous and several satellite lesions filled with serum or yellow fluid are arranged around a central necrotic lesion which is covered by a black eschar.

• This lesion is called a malignant pustule.

• The disease used to be common in dock workers, carrying loads of hides and skins on their bare backs and hence known as hide porter's disease.

D) Healing brown recluse spider bite:

• The bite of the Loxosceles reclusa, the brown recluse spider is usually relatively painless.

• After minutes or hours, severe pain develops at the site, accompanied by erythema, edema, and a central bulla.

• In severe envenomation, a ‘target lesion’ is seen- central blue/ purple discoloration surrounded by an ischemic halo and an outer ring of erythema (the ‘red, white and blue’ sign).

• After 3-4 days, the central area becomes necrotic and an eschar develops.

• The eschar sheds eventually, leaving behind an ulcer, which takes time to heal.

Solution to Question 2:

The given image shows clusters of vesicles seen on an erythematous base suggesting a diagnosis of genital herpes. Tzanck cells or multinucleated giant cells with intranuclear bodies are present on microscopy. Intranuclear inclusion bodies are seen when the smears are stained with Giemsa stain.

The typical herpes lesions are thin-walled, umbilicated vesicles, the roof of which breaks down, leaving tiny superficial ulcers. They heal without scarring. The most common site of cutaneous lesions in the face - on the cheeks, chin, around the mouth, or on the forehead.

The Tzanck smear technique is a rapid and inexpensive diagnostic method. Scrapings (preferably from the base of vesicles) are obtained from the lesions and smears are prepared. When stained with 1% aqueous solution of O-toluidine blue for 15 seconds, multinucleated giant cells with faceted nuclei and homogenously stained ground glass chromatin known as ‘ Tzanck cells’ are seen.

Genital herpes:

Herpes Simplex Virus (HSV) 2 is more responsible for genital herpes than HSV 1. The vesiculo-ulcerative lesions are very painful. In men, it occurs mainly on the penis or urethra. In women, the cervix, vagina, vulva and perineum can all be involved. The primary infection causes serious symptoms accompanied by fever and malaise, followed by recurrent episodes.

Solution to Question 3:

The image shows areas of localized, unilateral depigmented macules. This feature, along with a history of leukotrichia suggest that the patient is suffering from segmental vitiligo. It occurs due to a lack of melanocytes in the basal layer of the epidermis.

It is a localised depigmentation that occurs due to a progressive loss in the melanocytes. It is classified as

• Segmental vitiligo- which appears as unilateral maculae in a segmented or band-shaped pattern

• Non-segmental vitiligo- which appears as bilateral maculae distributed in an acrofacial pattern or scattered symmetrically over the entire body

The lesions become apparent between 20-30 years of age.

Hypopigmented macules in vitiligo are commonly found in the hips, dorsum of the hands/fingers, feet, elbows, knees, and ankle areas. These are the areas of repeated friction. Rarely there may be complete vitiligo (vitiligo universalis), or trichrome vitiligo where there are areas of depigmentation, hyperpigmentation, along with the normal skin are present. The Koebner phenomenon is the development of lesions at the site of trauma on the uninvolved skin.

Several autoimmune conditions can be associated with vitiligo. They are:

• Thyroid disease (hyperthyroidism and hypothyroidism)

• Pernicious anemia

• Addison disease

• Diabetes

• Hypoparathyroidism

Under the Wood's lamp examination the depigmented areas emit a bright blue-white fluorescence and appear sharply demarcated.

Treatment involves cosmetic camouflage and the use of sunscreen. The topical application of corticosteroids is the first line of treatment. Topical calcineurin inhibitors like pimecrolimus, tacrolimus have also shown to be successful. Psoralen photochemotherapy, khellin, selective ultraviolet B therapy are used as second-line treatment. In patients who do not respond to this treatment, surgical grafting techniques can be used, where the melanocytes from a normal pigmented area are transferred to a non pigmented area. In patients with extensive vitiligo where only a few pigmented areas are left, depigmentation can be done.



Other options:

A) Focal vitiligo:

These are small isolated depigmented lesions that are neither segmentally distributed nor evolved into non-segmental lesions after 1-2 years. The focal vitiligo along with isolated mucosal lesions on one site are considered as undetermined /unclassified vitiligo.

C) Piebaldism:

It commonly presents with a white V-shaped forelock. it is caused due to the impaired migration of melanocytes to the skin, resulting in hypopigmented patches.

D) Nevus depigmentosus:

It is a congenital pigmentary disorder that presents with a non-progressive hypopigmented macule. It usually is present from birth or early childhood and remains stable throughout life.

Solution to Question 4:

Gottron papules are typical and specific sign of juvenile dermatomyositis.

Gottron papules and a heliotropic rash of the eyelids are the classical criteria that are required for the diagnosis of juvenile dermatomyositis along with three of the following:

Weakness	SymmetricProximal
Muscle enzyme elevation (≥1)	Creatine kinaseAspartate aminotransferaseLactate dehydrogenaseAldolase
Electromyographic changes	Short, small polyphasic motor unit potentialsFibrillations Positive sharp wavesInsertional irritabilityBizarre, high-frequency repetitive discharges
Muscle biopsy	NecrosisInflammation

Gottron's papules are violaceous (of violet color), lichenoid (red-purple), inflammatory, flat-topped papules on proximal interphalangeal joints, distal interphalangeal joints, and metacarpophalangeal joints.

This should not be confused with Gottron's sign.

Gottron's sign is erythematous or violaceous, macules or patches in the same symmetric distribution pattern but sparing the interphalangeal spaces and sometimes involving elbows and

knees.

The heliotrope rash is a violet discoloration of the eyelids that may be associated with periorbital edema. Rarely, an erythematous, scaly rash develops on the palms of the children called 'mechanic hands'. Nail bed capillary changes are visible as individual thickened and telengiectatic capillary loops, which can be well seen with the help of capillaroscopy. Facial erythema mimicking malar rash of systemic lupus erythematosus, but with the involvement of nasolabial folds are also common.

Solution to Question 5:

The above image shows tinea capitis, which presents with a well-demarcated scaly patch in which hair shafts are broken off just above the skin resulting in alopecia. KOH mount is useful for the depicted condition.

The scrapings should be taken from the edge of a lesion and transferred to a slide, to which KOH is added and examined under the microscope for the presence of hyphae. Tinea capitis is a dermatophyte infection primarily occurring in children. Clinical features of tinea capitis are patchy hair loss, scaling, tender lymphadenopathy, and in severe cases, there might be pustules and crusting of the affected area.

The clinical variants of tinea capitis are:

• Endothrix - a non-inflammatory type in which multiple black dots are present within the areas of alopecia. It is most commonly caused by T. tonsurans and T. violaceum.

• Ectothrix - a mild inflammatory type in which single or multiple scaly patches with hair loss are seen. Microsporum audouinii, M.canis, M.equinnum, and M.ferrugineum cause ectothrix infection.

• Kerion - a severe inflammatory type that presents with an inflammatory mass with thick crusting and matting of hair with pus and sinus formation. It is commonly caused by T. verrucosum and T. mentagrophytes.

• Favus - is characterized by the presence of yellowish, cup■shaped crusts known as scutula that develop around hair follicles. It is caused by T. schoenleinii.

Treatment:

• Terbinafine: <10 Kg- 62.5 mg; 10-20 kg - 125 mg; >20 kg- 250 mg, all given daily for 4 weeks.

• Itraconazole: 2-4 mg/kg/day for 4-6 weeks.

Solution to Question 6:

The given image shows a case of borderline tuberculous leprosy, or dimorphous leprosy, as evidenced by the cutaneous, rash-like discoloration of the patient's skin. This generally overlies the back, and crosses the midline. So, the best choice would be Borderline leprosy.

Microscopically, there is diffuse epithelioid cell granuloma with very scanty lymphocytes and no giant cells; the papillary zone is clear and the nerves are swollen slightly by cellular infiltrate. Acid-fast bacilli are present in moderate numbers.

The borderline leprosy patients are immunologically unstable and are at an increased risk of developing immune-mediated reactions. Type 1 lepra reactions that occur most commonly in mid-borderline leprosy patients, are delayed hypersensitivity reactions caused by increased recognition of mycobacterial leprae antigens in skin and nerve sites.

Solution to Question 7:

The given clinical history of itchy plaques over the neck, cubital fossa and popliteal fossa suggests the diagnosis of 'atopic dermatitis'.

Atopic dermatitis is the cutaneous expression of the atopic state, characterised by a family history of asthma, allergic rhinitis or eczema.

The etiology is only partially defined, but there is a clear genetic predisposition. A characteristic defect is impaired epidermal barrier. In many patients a mutation in gene encoding filaggrin (a structural protein in the stratum corneum) is responsible. The lesions are distributed in the flexural aspects.(as shown in the image).

Patients may display many immunoregulatory abnormalities, including increased IgE synthesis, increased serum IgE levels, delayed type hypersensitivity reactions.

The cardinal symptom is itching, along with rash.

Atopic Eczema can be divided into 3 phases: Infantile, Childhood, and Adult phases.

Infantile phase: The lesions of Atopic Eczema most frequently start on the face but may occur anywhere on the skin surface. Often, the napkin area is relatively spared.

• When the child begins to crawl, the exposed surfaces, especially the extensor aspect of the knees and elbows are most involved.

• The lesions consist of erythema and discrete or confluent oedematous papules. The papules are intensely itchy and may become exudative and crusted as a result of rubbing.

• Secondary infection and lymphadenopathy are common.

Childhood Phase: From 18 to 24 months onwards, the sites most characteristically involved are the elbow and knee flexures, a mixture of erythema, crusting, excoriation, hyper■ and hypopigmentation, and warty lichenification may be seen depending on the skin type on the sides of the neck, wrists, and ankles.

• The sides of the neck may show a striking reticulate pigmentation, sometimes referred to as atopic dirty neck.

• The erythematous and oedematous papules tend to be replaced by lichenification.

Adult Phase: The picture is essentially similar to that in later childhood, with lichenification, especially of the flexures and hands.

Other options:

B) Dermatitis herpetiformis:

• Dermatitis herpetiformis is a chronic, intensely pruritic, skin condition associated with gluten■sensitive enteropathy.

• It typically appears in the 4th decade, and the patients present with itching and rash, most typically over the extensor surfaces of the elbows, knees, buttocks, and scalp.

• They may also have symptoms associated with gluten-sensitive enteropathies, like bloating, diarrhea and other gastrointestinal complaints.

• On examination, multiple erythematous papules and vesicles can be seen over the extensor aspects. They appear symmetrical and heal without scarring. The oral cavity may show mucosal abnormalities and enamel pits.

C) Pemphigus vegetans:

It is a rare form of the bullous disorder that presents with large, moist, verrucous, vegetating plaques studded with pustules on the groin, axillae and flexural surfaces.

Predominant cell	Name of microabscess	Disease
Neutrophilic microabscess	Papillary tip microabscessMunroe's microabscess (superficial)Spongiform pustules of Kogoj (deep)	Dermatitis HerpetiformisPsoriasisPsoriasis
Lymphocytic microabscess	Pautrier's microabscess	Mycosis fungoides
Eosinophillic microabscess		Pemphigus vegetans

D) Psoriasis:

• Plaque psoriasis is the most common type of psoriasis and is seen in 80-90% of patients.

• The peak on occurrence is between 16 to 22 years and between 57 to 62 years. They usually present with itching and a papulosquamous rash.

• The lesions are red, scaly, and have well demarcation from the normal skin.

• They can also be encircled by a clear peripheral zone which is called the Woronoff ring.

• They commonly appear in the scalp, trunk, and extensor surfaces of the limbs.



Solution to Question 8:

The given clinical scenario is highly suggestive of bullous pemphigoid.

Bullous pemphigoid is characterized by tense itchy bulla in the flexor areas and trunk which do not rupture easily. Elderly aged people are usually affected. It is a subepidermal immunobullous disorder, which occurs due to the production of autoantibodies (IgG) against BP180 (BPAG2) and BP 230 (BPAG1) antigens on hemidesmosomes.

Mucosal membrane involvement is rare in this disease. Most of the lesions heal without a scar. Since these bullae are subepidermal, pressure over the lesions does not separate the epidermis from the dermis. (Nikolsky sign is negative).

Immunofluorescent microscopy will show the presence of subepidermal split and deposition of IgG antibodies at the dermo-epidermal junction. Histopathology will show the presence of subepidermal bullae with eosinophils.

It is mainly treated with corticosteroids. Other drugs used are azathioprine, dapsone, doxycycline, etc.



The above image (immunofluorescence microscopy) shows linear IgG deposits at the dermo-epidermal junction seen in the bullous pemphigoid.

Other options:

Option B: Pemphigus Vulgaris is an intraepidermal immunobullous disease. It occurs commonly between the 4th to 6th decade. They present with flaccid blisters, unlike bullous pemphigoid. Firm pressure over the lesion separates the epidermis from the dermis. Nikolsky's sign is positive. Mucosal involvement is seen in the form of buccal and palatal erosions.



Option C: Dermatitis herpetiformis typically appears in the 4th decade, and the patients present with itching and rash, most typically over the extensor surfaces. The majority of the patient have symptoms related to gluten-sensitive enteropathies, like bloating, diarrhea, and other gastrointestinal complaints. Immunofluorescence microscopy will show the deposition of IgA antibodies in the papillary dermis in a granular or fibrillar pattern(as shown in the below image).



Option D: Linear Ig A dermatosis (Chronic bullous dermatosis of childhood) is an autoimmune blistering disease seen in infants and children. The disease can also be seen in adults and is known as Linear IgA disease of adults or IgA bullous pemphigoid. The blisters are arranged in an annular pattern commonly known as the crown of jewels appearance (image below).

Solution to Question 9:

The age of the patient and the presence of papules and pustules with an erythematous base on the face are suggestive of acne. All the above agents are useful in the treatment of acne except topical fluocinolone (topical steroid). Corticosteroids may provoke an acneform reaction irrespective of their route of administration.

Acne is a chronic inflammatory disease of the pilosebaceous unit. It is commonly seen in teenagers. It can vary from very mild comedonal acne to severe aggressive pustular acne. These

eventually may lead to scarring due to sinus formation.

Acne results from the combination of:

• Increased sebaceous gland activity under the influence of androgens - oily skin

• Abnormal follicular differentiation with increased keratinization- hypercornification of the duct due to increased IL-1

• Microbial hypercolonization of the follicular canal leading to its inflammation.

Management-

• Mild comedonal acne is treated with a topical retinoid. Mild papular and pustular acne is treated with topical retinoid and topical antibiotics.

• Moderate papular/pustular acne and moderate nodular acne are treated with oral antibiotic + topical retinoid with or without benzoyl peroxide.

• Severe nodular and conglobate acne are treated with oral isotretinoin.

Tazarotene is a third■generation retinoid, which is available for the treatment of acne.

Solution to Question 10:

The case presenting with hypopigmented lesions with normal sensations and a history of fever in childhood suggests a diagnosis of post-kala-azar dermal leishmaniasis.

Post-kala-azar dermal leishmaniasis (PKDL) is a condition when Leishmania donovani is found in the skin and manifests as dermal lesions. It is usually seen in the kala-azar endemic zones i.e. Bihar, Jharkhand, U.P, West Bengal (Northeastern states of India)

These patients present with erythematous rash and hypopigmented macules similar in appearance and distribution to those of lepromatous leprosy but normally less than 1cm. After a variable period of years or months, diffuse nodulation begins to develop in these macules.

The rash is progressive over many years and seldom heals spontaneously.

The image below shows post-kala-azar cutaneous leishmaniasis.

Other options:

Options A and B: The presence of hypopigmented macules over the trunk and face, suggests a differential of leprosy, but the nerves and sensations are normal, hence ruling it out.

Option C: Mycosis fungoides is characterized by polymorphic erythematous patches and plaques which are usually associated with severe pruritus. They could be asymptomatic too. The image below shows characteristic lesions in mycosis fungoides.

Solution to Question 11:

The image shows an aggregate of pinhead-sized papules over the forearm of a child which is suggestive of lichen nitidus.

Lichen nitidus presents with tiny pinhead-sized flesh-colored papules and has a flat or dome■shaped, shiny surface. They are found on any part of the body but the sites of predilection are the forearms, penis, abdomen, chest, and buttocks.

Lichen nitidus is considered to be a variant of Lichen planus as both are histologically similar in early stages. It has a "claw clutching a ball' appearance on histopathological examination. It refers to a sharply circumscribed inflammatory infiltrate in the dermis, that spans 4-5 dermal papillae.

The diagnosis can be made clinically, but skin biopsy provides a definitive diagnosis. Lesions are usually asymptomatic and rarely require any treatment. Symptomatic or generalized lesions can be treated with fluorinated topical steroids, PUVA, narrow■band UVB phototherapy, astemizole, or acitretin.

Pinhead-sized flesh-colored papules seen.

Other options:

Option A: Lichen planus is an idiopathic inflammatory disease affecting the skin and mucous membrane. It usually presents with pruritic, purple, polygonal, papules. The lesions have thin white lines on their surface known as Wickham's striae.

Option C: Scabies has typical distribution in the interdigital spaces and burrows are the characteristic lesions. The below image shows a magnified view of a burrowing trail of the scabies mite.



Option D: Scrofuloderma results from the direct invasion of the tubercle bacillus into the skin from underlying tuberculous foci.

Solution to Question 12:

Among the given options, Indeterminate leprosy have more remission than tuberculoid is the most appropriate answer.

The manifestations of leprosy depend on the immune response of an individual to the bacillus Mycobacterium leprae. About 80-90% of the people do not develop any disease after bacilli exposure. The remaining patients develop indeterminate leprosy. The lesions of indeterminate leprosy tend to be hypopigmented and ill-defined. Among those patients, 75% undergo spontaneous remission. Very few patients present clinically between tuberculoid or lepromatous leprosy. Therefore, more patients with indeterminate leprosy undergo spontaneous remission compared with those having tuberculoid leprosy. (Option D).

Leprosy (Hansen's disease) is a chronic infective granulomatous disease caused by the bacillus Mycobacterium leprae.

Leprosy is clinically characterized by one or more of the following features :

- Hypopigmented patches
- Partial or total loss of cutaneous sensation in the affected areas
- Presence of thickened nerves, and
- Presence of acid-fast bacilli in the skin or nasal smears.

Option A is incorrect because spontaneous remission is seen with leprosy. Even in the absence of specific treatment, a majority of patients tend to get cured spontaneously.

Option B is incorrect because around 80-90% of the cases (not half) resolve spontaneously after bacilli exposure.

Option C is incorrect because spontaneous remission in children is variable. Even though few studies mentions that children have more remission, however, the observation period was not long enough to establish its permanence.

Solution to Question 13:

The above image shows unilateral, red, closely grouped vesicles in a single dermatome associated with severe pain is suggestive of herpes zoster or shingles.

It is a segmental eruption due to the reactivation of latent varicella-zoster virus from dorsal root ganglia. An earlier infection with varicella is essential before zoster can occur.

The first symptom of zoster is usually pain and it may be accompanied by fever, headache, malaise, and tenderness localized to areas of one or more dorsal roots. The mucous membranes within the affected dermatomes are involved. The lymph nodes draining the affected area are enlarged and tender.

The patients are infectious, both from the virus in the lesions and nasopharynx. The most commonly involved dermatomes in decreasing order of frequency are thoracic > cervical > trigeminal > lumbosacral. The posterior nerve roots and ganglia show inflammatory changes.

Zoster is a self■limiting infection but it is painful and carries a risk of secondary infection and post■herpetic neuralgia. Analgesia and treatment of secondary infections are sufficient for mild infections.

Indications for antivirals in VZV (Acyclovir 800 mg):

• Painful zoster infections in adults

• Facial zoster at any age

• Immunocompromised

Other options:

Option B: The lesions of the molluscum contagiosum are shiny, white, and hemispherical. It is umbilicated with a central pore. It is caused by poxvirus.

The below image shows the lesions seen in Molluscum contagiosum

Option C: Herpes simplex virus are classified into two major antigenic types,

• Type 1 (HSV-1) which is classically associated with facial infections;

• Type 2 (HSV-2) which is typically genital.

Below image shows small, closely grouped vesicles on an inflamed base and is characteristic of Herpes Labialis or Cold sore.

Option D: Lymphangiomas are dilated dermal lymphatics that blister onto the skin surface. The fluid is usually clear, but occasionally maybe blood-stained. Lymphangiomas may thrombose and get fibrosed, to form hard nodules in the long term. This may raise concerns about malignancy.

If lymphangiomas are <5 cm across, they are called lymphangioma circumscriptum (shown below), and they are called lymphangioma diffusum if they are more widespread.

J.Menzies

Dermatology NEET 2021

Question 1:

A 10-year-old male child complains of mild painful swelling over the scalp for 3 months as shown in the image below. He also has a pet dog at his home. What is the diagnosis of the given condition?

a) Furuncle

b) Epidermoid cyst

c) Kerion

d) Folliculitis

Question 2:

A child with a history of night-blindness presented to you with the following findings. What is the diagnosis?

a) Phrynoderma

b) Keratosis pilaris

c) Darier disease

d) Follicular eczema

Question 3:

A 30-year-old presented with flaccid bullae on her skin which are easy to rupture. The biopsy revealed a suprabasal split. What is the most likely diagnosis?

a) Pemphigus vulgaris

b) Pemphigus foliaceus

c) Pemphigus vegetans

d) Erythema multiforme

Question 4:

What is the treatment of choice for a patient with paronychia of the index finger on the right hand, as shown in the image below?

a) Amoxiclav

b) Metronidazole

c) Amikacin

d) Norfloxacin

Question 5:

A patient taking multi-drug therapy (MDT) presents with worsening of existing lesions and nerve involvement. What will be your next best step of action?

a) Stop MDT, start systemic corticosteroids

b) Continue MDT, start systemic steroids

c) Stop MDT give thalidomide

d) Continue MDT, start thalidomide

Question 6:

A 30-year-old male patient presents to you with inflammatory alopecia. Which of the following variants of tinea capitis infection commonly result in this type of alopecia?

a) Endothrix and Favus

b) Ectothrix and Endothrix

c) Favus and Kerion

d) Kerion and Endothrix

Answer Key

Question No.	Correct Option
1	c
2	a
3	a
4	a
5	b
6	c

Detailed Explanations

Solution to Question 1:

The clinical vignette of a young boy in contact with a zoophilic organism, presenting with a boggy inflammatory mass, with thick crusting and matting of hair is suggestive of a kerion. Kerion is a severe inflammatory type of tinea capitis.

Tinea capitis is a dermatophyte infection primarily occurring in children. Clinical features of tinea capitis are patchy hair loss, scaling, tender lymphadenopathy, and in severe cases, there might be pustules and crusting of the affected area along with sinus formation.

The clinical variants of tinea capitis are:

• Endothrix - a non-inflammatory type in which multiple black dots are present within the areas of alopecia. It is most commonly caused by T. tonsurans and T. violaceum.

• Ectothrix - a mild inflammatory type in which single or multiple scaly patches with hair loss are seen. Microsporum audouinii, M.canis, M.equinnum, and M.ferrugineum cause ectothrix infection.

• Kerion - a severe inflammatory type that presents with an inflammatory mass with thick crusting and matting of hair with pus and sinus formation. It is commonly caused by T. verrucosum and T. mentagrophytes.

• Favus - is characterized by the presence of yellowish, cup■shaped crusts known as scutula that develop around hair follicles. It is caused by T. schoenleinii.

Treatment:

• Terbinafine: <10 Kg- 62.5 mg; 10-20 kg - 125 mg; >20 kg- 250 mg, all given daily for 4 weeks.

• Itraconazole: 2-4 mg/kg/day for 4-6 weeks.

The image below shows a kerion:

The image below shows the endothrix variant of tinea capitis:

Option A: Furuncle/ boil is caused by Staphylococcus. It is painful and usually situated on the neck area, and the shedding of loosened hairs is generally much less evident than in kerion. It is pustular and later becomes necrotic and forms a scar.

The image below shows a furuncle- pustule over the hair follicle with a necrotic base:



Option B: Epidermoid cyst/ sebaceous cyst results from squamous metaplasia in a damaged sebaceous gland. Many are the result of inflammation around a pilosebaceous follicle. The most common site of a sebaceous cyst is the postauricular sulcus or below and behind the ear lobule. They are fixed to the skin and usually have a central punctum. Treatment depends on the clinical state of the cyst. When inflamed or infected, they should be incised and drained initially, total surgical excision should be done once the inflammation has subsided.

The image below shows an epidermoid cyst of the post-auricular sulcus- cyst fixed to the skin, shows a central punctum:

Option D: Folliculitis would involve specific inflammation in the pilosebaceous units. It is caused by Staphylococcus aureus. They heal without a scar.

The image below shows folliculitis- inflammation of the pilosebaceous units leading to pustule formation:

Solution to Question 2:

A child having night-blindness, and multiple keratotic papules on elbows as shown in the image suggests a diagnosis of phrynoderma or toad-skin. This can occur due to the deficiency of vitamin A.

Phrynoderma is not specific to vitamin A deficiency. It may also be associated with deficiencies of B complex, riboflavin, vitamin C, vitamin E, and essential fatty acids. It manifests as groups of

papules with a central keratotic plug. The commonest sites are elbows and knees. They are present around the follicles and give the skin a rough texture. Treatment involves the correction of poor diet and administration of a multivitamin preparation containing vitamin A.

Option B: Keratosis pilaris is an inherited abnormality of keratinization. It affects the follicular orifices leading to keratotic follicular plugging, perifollicular erythema, and follicular atrophy. It usually presents during puberty. They present as small, keratotic papules on the extensor aspects of the limbs, particularly the arms which can lead to ingrowing of the hair. Treatment is purely reserved for cosmetic purposes.

Option C: Darier disease is an autosomal dominant genodermatosis characterized by keratotic papules that show acantholysis and dyskeratosis. ATP2A2 gene mutations are present in Darier disease. Histologically, acantholysis (separation of keratinocytes) and blistering are seen in the suprabasal epidermis.

It presents with discrete or confluent rough, greasy, skin-colored papules in seborrheic areas (trunk, face scalp margins) and flexures (perineum, axillae, groin). There may also be isolated nipple hyperkeratosis. Nail findings include fragility, painful longitudinal splits, or distinctive red and white longitudinal bands terminating in V-shaped nicks.

Darrier's disease

Discrete or confluent rough, greasy, skin-colored papules in seborrheic areas

Darrier's disease

Greasy papules over the flexural areas like thigh and groin

Nail finding in Darrier's disease

Distinctive red and white longitudinal bands terminating in V-shaped nicks.

Option D: Follicular eczema is a form of atopic dermatitis that presents with a discoid patch formed due to the coalescence of the follicular lesions with peripheral excoriation marks. The biopsy shows spongiotic dermatitis.

Solution to Question 3:

The given clinical vignette of a bullous disorder with the histologic picture pointing towards a suprabasal blister suggests the diagnosis of pemphigus vulgaris.

It is caused due to the production of antibodies mainly against desmoglein 3. Rarely antibodies against desmoglein 1 can also be found. The antibodies are commonly IgG antibodies. Desmoglein 3 is mainly present in the basal and suprabasal layers of the skin and mucosa.

The patients, therefore, present with flaccid bullae over the skin and mucosal ulcerations. Oral mucosa is commonly affected. Other areas where ulceration can be seen are conjunctiva, nasopharynx, larynx, etc. Cutaneous lesions are more common over the scalp, face, neck, upper chest, and back.

Direct immunofluorescence will show the deposition of antibodies around the keratinocytes in the epidermis. It presents with a characteristic fish-net appearance in the immunofluorescence assay It is treated with steroids, azathioprine, mycophenolate mofetil, rituximab, etc.

The images attached below depict the lesions, the suprabasal split, and the fish-net pattern respectively seen in a case of pemphigus vulgaris.



Option B: Pemphigus foliaceus is a benign form of pemphigus, an inflammatory blistering disorder. This is a form of a subcorneal blister, where the stratum corneum forms the roof of the bulla. It is caused by autoantibodies that result in the dissolution of intercellular attachments within the epidermis. Antibodies against desmoglein 1 are mainly seen in pemphigus foliaceous. There is no involvement of the mucosal epithelium in pemphigus foliaceus.

Lesions are most common on the scalp, face, chest, and back. Bullae are superficial and mainly present as areas of erythema and crusting. These represent superficial erosions at sites of blister rupture.

Option C: Pemphigus vegetans is a rare form of the bullous disorder that presents with large, moist, verrucous, vegetating plaques studded with pustules on the groin, axillae, and flexural surfaces. It is associated with eosinophilic microabscess. It has two types, a severe Neumann type, and a milder Hallopeau type.

Option D: Erythema multiforme is a self■limiting cytotoxic dermatitis resulting from cell■mediated hypersensitivity most commonly to infection or drugs. The common triggers are

- Herpes simplex infection
- Mycoplasma infection
- Bacterial infections

• Fungal infections

• Vaccinations

• Malignancy: carcinoma, lymphoma, leukemia

• Drug reactions: sulfonamides, cotrimoxazole, penicillins

• Miscellaneous: Lupus erythematosus (Rowell Syndrome), Sarcoidosis, Polyarteritis Nodosa

It presents clinically with macular, papular, or urticarial lesions, along with the classic acral iris or target lesions. The lesions are distributed on the distal extremities, palms or trunk, and oral and genital mucous membranes.

Target lesions are less than 3 cm in diameter, rounded, and have three zones:

• The central area of dusky erythema or purpura

• Middle paler zone of edema

• The outer ring of erythema with a well■defined edge

Atypical lesions will have only 2 zones. They fade in 1-2 weeks leaving dusky discoloration.



Classical acral iris/ Target sign

Solution to Question 4:

The above image depicts paronychia associated with a reddish erythematous streak-like lesion on the hand suggestive of paronychia associated with lymphangitis. The treatment of choice for this patient is amoxiclav.

Acute paronychia is an acute infection of the nail beds and folds. It is most commonly caused by Staphylococcus aureus. It presents as a tender, erythematous swelling around the nail.

Treatment of pyogenic paronychia:

• Protection against trauma

• Keep affected fingernails dry.

• Abscesses - Incision and drainage. (abscess is often opened by pushing the nail fold away from the nail plate)

• Antibiotics- A penicillin in combination with a β-lactamase inhibitor, e.g. amoxicillin with clavulanic acid (Amoxiclav)

Chronic paronychia is usually due to infection with Candida albicans and responds to topical antifungal therapy.

Lymphangitis:

Lymphangitis represents acute inflammation elicited by the spread of bacterial infections into lymphatics, with group A β-hemolytic streptococci being the most common causative agent. It is manifested by red, painful subcutaneous streaks (the inflamed lymphatics) and painful enlargement of the draining lymph nodes (lymphadenitis). It is managed by elevating the affected extremity and broad-spectrum antibiotics.

Solution to Question 5:

The given scenario of worsening of existing lesions in the context of leprosy treatment with multi-drug therapy (MDT) is suggestive of a type 1 lepra reaction. The next best step in the management would be to continue MDT and start systemic steroids.

Type 1 lepra reactions present with inflammation of existing skin lesions. The lesions become red, painful, and edematous. They may rarely ulcerate. New lesions may occur but are very rare. It is associated with acute neuritis, with the tenderness and loss of sensory and motor function.

MDT is never stopped in patients with lepra reactions. For pain, NSAIDs can be supplemented along with oral steroids.

The use of thalidomide is restricted to severe forms of type-2 lepra reaction or erythema nodosum leprosum. These are characterized by systemic effects like iritis, orchitis, and eruption of new skin lesions.

The summary of lepra reactions is attached in the pearl below.

Solution to Question 6:

Favus and kerion are the inflammatory variants of tinea capitis.

Tinea capitis is a dermatophyte infection primarily occurring in children. Clinical features of tinea capitis are patchy hair loss, scaling, tender lymphadenopathy, and in severe cases, there might be pustules and crusting of the affected area along with sinus formation.

The clinical variants of tinea capitis are:

• Kerion is a severe inflammatory type of tinea capitis which presents with an inflammatory mass with thick crusting and matting of hair.

• Favus is characterized by the presence of yellowish, cup■shaped crusts known as scutula that develop around hair follicles.

- Endothrix is a non inflammatory type in which black dots form around swollen hair shafts.
- Ectothrix is a mild inflammatory type in which dull grey coated broken hairs is seen.

Dermatology INI-CET 2022

Question 1:

A 10-year-old boy presents with fever, joint pain, and a lesion over his hand as seen in the image below. Which of the following is the clinical finding given and what is the likely diagnosis?

a) Gottron's papules - Juvenile dermatomyositis

b) Rheumatoid nodules - Juvenile rheumatic arthritis

c) Sclerodactyly - Systemic sclerosis

d) Mechanic's hand - Systemic lupus erythematosus

Question 2:

A 12-week pregnant woman who is already on multidrug therapy for leprosy presents with type 2 lepra reaction. What is the treatment of choice for this patient?

a) Stop MDT and start oral steroids

b) Antibiotics

c) Thalidomide

d) Continue MDT and add oral steroids

Question 3:

A patient presents to the outpatient department with a satellite skin lesion. On examination, he has one hypopigmented lesion and one thickened nerve. Images of two blister packs are given below. The treatment best suited for this patient would be:



a) Packet 1 for 12 months

b) Packet 2 for 12 months

c) Packet 1 for 6 months

d) Packet 2 for 6 months

Question 4:

Chemoprophylaxis for leprosy should be given to which of the following persons?

a) 1, 2, 4

b) 1, 3, 4

c) 2, 3, 4

d) 1, 2, 3

Question 5:

Match the following types of scales with their associated conditions.

a) 1-b, 2-c, 3-d, 4-a

b) 1-a, 2-b, 3-d, 4-c

c) 1-d, 2-c, 3-a, 4-b

d) 1-c, 2-b, 3-d, 4-a

Question 6:

A 60-year-old patient presents with the lesion given in the image below. He also describes having the sensation of spiders crawling in the area of the lesion when he woke up. Which of the following is the most likely diagnosis?

a) Herpes zoster infection

b) Irritant contact dermatitis

c) Allergic contact dermatitis

d) Herpes simplex infection

Question 7:

A 60-year-old diabetic female patient presents with progressive skin lesions, as shown in the image. What is the diagnosis?

a) Squamous cell carcinoma

b) Basal cell carcinoma

c) Discoid lupus eryhtematosus

d) Melasma

Question 8:

A diabetic patient presents with the following lesions in the armpit. Which of the following would be seen on Wood's lamp examination?

a) Yellow fluorescence

b) Coral red fluorescence

c) Blue fluorescence

d) Green fluorescence

Question 9:

A 10-year-old boy is brought with weakness in his lower limbs and rash over the eyelids. The biopsy shows inflammation of the muscle. What is the finding shown in the image below?

a) Gottron's papules

b) Shawl sign

c) Heliotrope rash

d) Scleroderma

Question 10:

Consider the causes of alopecia. Which among the following cause(s) non-scarring alopecia?

a) 3 and 4

b) 1, 2, and 3

c) 2, 3, and 4

d) Only 4

Question 11:

Match the following therapeutic methods used in the management of psoriasis with their possible side effects.

a) 1-c, 2-b, 3-a, 4-d

b) 1-c, 2-a, 3-b, 4-d

c) 1-b, 2-d, 3-c, 4-a

d) 1-b, 2-c, 3-a, 4-d

Question 12:

Skin manifestations caused by the drug administered for amebic dysentery are due to _____.

a) Fixed drug eruption

b) Urticaria

c) Maculopapular rash

d) Atopic dermatitis

Question 13:

A 40-year-old man presents with erythematous painful lesions on the trunk and face, joint pain, and glove and stocking pattern of anesthesia. Which of the following investigations is used for the diagnosis of this condition?

a) 1, 2

b) 1, 2, 3

c) 1, 3, 4

d) 1, 2, 3, 4

Question 14:

Identify the condition shown in the image below.

a) Paget disease

b) Lichen sclerosus

c) Lichen planus

d) Angioma of vulva

Question 15:

A patient comes to the OPD with the following skin manifestation of a disease associated with Raynaud phenomenon. What is the manifestation?

a) Poikiloderma

b) Salt and pepper pigmentation

c) Leukotrichia

d) Vitiligo

Question 16:

A female patient came to the hospital with a history of relapsing, remitting, mildly itchy erythematous plaques over the scalp, sacral area, and knees. She has a history of joint pain and her nail changes are shown in the image below. What is your diagnosis?

a) Adult onset Still's disease

b) Pachyonychia congenita

c) Psoriasis

d) Lichen planus

Answer Key

Question No.	Correct Option
1	a
2	d
3	d
4	a

5	a
6	a
7	b
8	b
9	a
10	b
11	a
12	a
13	b
14	b
15	b
16	c

Detailed Explanations

Solution to Question 1:

The given image showing erythematous papules over dorsal metacarpophalangeal and interphalangeal joints (Gottron's papules) is a characteristic skin lesion seen in dermatomyositis (DM). Fever and joint pain (arthralgia) are non-specific complaints.

DM is an inflammatory myopathy affecting skin and skeletal muscle. DM can be seen in children (juvenile DM) or in adults. It presents with symmetric, proximal more than distal muscle weakness associated with characteristic skin manifestations:

• Heliotrope rash: Lilac erythema of eyelids with periorbital edema

• Gottron’s papules: Violaceous, flat■topped papules on the skin overlying the interphalangeal joints and metacarpophalangeal joints

• Gottron sign: Erythematous rash over the extensor surfaces of joints such as the knuckles, elbows, knees, and ankles

• V-sign: Erythematous macules over anterior chest

• Shawl sign: Erythematous macules over posterior neck and upper back

• Mechanic's hand: Roughening and cracking of the skin of the tips and sides of the fingers, resulting in irregular, dirty-appearing lines that resemble those of a mechanic or manual laborer

• Nail bed telangiectasias

• Subcutaneous calcium deposits

DM is associated with interstitial lung diseases (ILD) and internal malignancies, most commonly adenocarcinomas. Adult onset DM has a stronger association with malignancies.

Laboratory evaluation of patients with dermatomyositis will reveal elevated serum creatinine kinase and antinuclear antibodies. Electromyography will reveal features of myopathy. Histology will show perifascicular atrophy. Immunohistochemical staining will reveal myxovirus resistance protein A, a sensitive and specific marker.

Treatment of dermatomyositis is include:

• Immunotherapy with high-dose glucocorticoids - first-line drug

• Methotrexate - second line drug. Indications include -

• When patient presents with severe weakness, or other organ system involvement

• Patients with risk of steroid complications like patients with diabetes, osteoporosis, or postmenopausal women

• Patients who do not show improvement after 2-4 months of treatment with prednisone or those who cannot be maintained on a low dose

• Triple therapy with intravenous immunoglobulin or rituximab may be required

The prognosis is favourable in the absence of associated ILD or malignancies.

Solution to Question 2:

The treatment of choice for a pregnant lady with type 2 lepra reaction is to continue multidrug therapy (MDT) and start oral steroids. MDT is not stopped in lepra reactions. MDT is proven to be safe in pregnancy. Thalidomide is contraindicated (especially in the 1st trimester) due to the risk of teratogenic changes in the fetus (phocomelia).

Type 2 lepra reaction or erythema nodosum leprosum (ENL) is an immune complex-mediated syndrome (type 3 hypersensitivity) affecting the skin, nerves, and other organs. It is mainly seen in the multibacillary form of leprosy and is commonly recurrent. It usually occurs during treatment, but can also be seen before or after treatment.

ENL presents with acute onset, painful, erythematous nodules on the face and extensor surfaces of limbs that blanch on pressure. Systemic manifestations include fever, malaise, anorexia, neuritis, uveitis, dactylitis, arthritis, myositis, etc. can also be seen.

Treatment includes:

• Thalidomide is the drug of choice in young men with severe or recurrent ENL. It is contraindicated during pregnancy due to its high teratogenicity causing phocomelia. Even in women of reproductive age, it is given with caution.

• Glucocorticoids (high doses) are routinely used in all patients and help in suppressing the acute signs and symptoms of inflammation.

• Clofazimine is useful as long-term maintenance therapy for preventing recurrent reactions. Clofazimine is not given to this patient because it is already a part of the MDT regimen containing dapsone, rifampicin, and clofazimine.

Solution to Question 3:

The treatment best suited for this patient would be packet 2-maroon for 6 months. The clinical scenario of one hypopigmented satellite skin lesion and one thickened nerve suggests paucibacillary leprosy. The treatment of paucibacillary leprosy (PB) is multidrug therapy (MDT) with rifampicin, dapsone, and clofazimine which is the maroon blister packet for 6 months.

The current recommendation by WHO is to treat PB leprosy with a combination of 3 drugs - rifampicin, dapsone, and clofazimine (maroon blister pack) for 6 months. Meanwhile, multibacillary (MB) leprosy is treated using the same 3 drugs for a duration of 12 months. While the NLEP guidelines earlier mentioned a 2-drug MDT for PB leprosy, the National Strategic Plan and Roadmap for Leprosy (2023-2027) now recommends a uniform MDT of 3 drugs for both PB and MB leprosy.

Management of leprosy is based on the bacterial load. WHO defines PB disease as the presence of no bacilli on smears or biopsy and less than or equal to 5 skin lesions with no or only one peripheral nerve involved. MB leprosy is characterized by bacilli on smears or biopsies of more than 5 skin lesions and more than 1 peripheral nerve involvement.

The image given below shows a hypopigmented satellite skin lesion seen in leprosy.

Chemoprophylaxis for leprosy should be given in those

• Age more than 2 years

• Close contact for more than 20 hours per week for more than 3 months

A single dose of rifampicin is recommended for the close contact of a patient with leprosy provided they do not develop any signs or symptoms pertaining to leprosy.

Solution to Question 4:

Chemoprophylaxis for leprosy should be given to contacts 1,2 and 4.

Chemoprophylaxis for leprosy should be given to the following contacts:

• A person who has been in close contact with the index case for 20 hours or more per week for more than 3 months.

• Close contact if the age is more than 2 years

The contact mentioned in option 4, is with the patient for more than 3 months. Therefore, even this contact will require chemoprophylaxis.

Chemoprophylaxis cannot be given if there is/are:

• Possible signs of leprosy

• Signs or confirmed tuberculosis

• Liver or kidney disease

• Pregnancy

• Received rifampicin in the last 2 years

A single dose of rifampicin is used in chemoprophylaxis. Chemoprophylaxis must be initiated only 4 weeks after treatment for leprosy has begun in the patient so as to ensure the patient is non-infectious.

Dose of rifampicin in chemoprophylaxis of leprosy	
Age	Dose
Children 2-5 years	10-15 mg/kg
Children 6-9 years (less than 20kg)	150 mg
Children 6-9 years (more than 20kg)	300 mg
10-14 years	450 mg
15 years and above	600 mg

Solution to Question 5:

The correctly matched option is option A: 1-b, 2-c, 3-d, 4-a.

Solution to Question 6:

The given image showing unilateral, red, grouped vesicles in a single dermatome associated with crawling sensation is highly suggestive of Herpes zoster infection.

Herpes zoster infection or shingles is a segmental eruption due to the reactivation of latent varicella-zoster virus from the dorsal root ganglia. An earlier infection with varicella is essential

before zoster can occur.

The first symptom of zoster is usually pain. It may also be accompanied by fever, headache, malaise, and tenderness localized to areas of one or more dorsal roots. The mucous membranes within the affected dermatomes are involved.

The patients are infectious, both from the virus in the lesions and the nasopharynx. The most commonly involved dermatomes in decreasing order of frequency are thoracic > cervical > trigeminal > lumbosacral. The posterior nerve roots and ganglia show inflammatory changes.

Zoster is a self■limiting infection but it is painful and carries a risk of secondary infection and post■herpetic neuralgia. Analgesia and treatment of secondary infections are sufficient for mild infections.

Other options:

Option B: Irritant contact dermatitis is the cutaneous response to the physical/toxic effects of a wide range of environmental exposures. The main clinical features are
usually dryness or chapping. The lesions could commonly develop into patchy or diffuse erythema with scaling and fissuring.

Option C: Allergic contact dermatitis is an eczematous reaction following exposure to a substance to which the immune system has previously been sensitized.

Option D: Herpes simplex lesions are thin-walled, umbilicated vesicles, the roof of which breaks down, leaving tiny superficial ulcers. They heal without scarring. The most common site of cutaneous lesions in the face - is on the cheeks, chin, around the mouth, or on the forehead.

Solution to Question 7:

The image above shows a central ulcer with raised borders and rolled-out edges (rodent ulcer) suggesting a diagnosis of basal cell carcinoma.

Basal cell carcinoma is a slow-growing malignant tumor with locally invasive characteristics. It affects the pilosebaceous skin.

Ultraviolet radiation exposure is the strongest predisposing factor. Other risk factors include exposure to arsenic, coal tar, aromatic hydrocarbons, and ionizing radiation.

Basal cell carcinoma mainly affects elderly or middle-aged men with excessive skin exposure, commonly in the age group of 40–80 y and in white-skinned people.

Macroscopically, it is categorized as localized and generalized. Nodular, nodulocystic (nodular and nodulocystic together account for 90% of basal cell carcinoma), pigmented, and nevoid are the localized cancers, while superficial multifocal and superficial spreading, infiltrative, and cicatrizing are the generalized types. The nodular type is described as pearly waxy nodules at the margins. A central depression (umbilication) or ulceration and rolled-out edges are characteristic of basal cell carcinoma.

The below gross image shows pearly, rolled-out borders suggestive of basal cell carcinoma.

Sites near the eye, nose, and ear and of size >2 cm are considered to be high risk. Direct invasion at these sites can reach the cranium.

Management is by Mohs microscopic surgery. Non-surgical measures, such as the administration of topical 5-fluorouracil and imiquimod, are useful. Metastasis is extremely rare.

Other options:

Option A: Squamous cell carcinoma (SCC) is the second most common skin cancer. UV radiation exposure is the primary risk factor for SCC. Classically, it appears as a scaly or ulcerated papule or plaque that can bleed with minimal trauma, but the pain is rare. It can exhibit in situ (confined to the epidermis) or invasive subtypes. Wide surgical excision including subcutaneous fat is the treatment of choice for SCC.

Option C: Discoid lupus erythematosus (DLE): DLE is typically located on the nose, cheeks, ear lobe, and concha. It may involve lips, oral mucosa, nose, or eyelids. Initial lesions of DLE are dry red patches, which then progress to indurated red or hyperpigmented plaques with adherent scale. Follicular keratosis (plugs of keratin within hair follicles) is observed when the surface scales are removed (carpet-tack sign). Older lesions are hyperpigmented, especially on the plaque edges. There is scarring, which results in central loss of pigment, white patches, and skin atrophy.

Option D: Melasma is the most common cause of facial melanosis that manifests as hyperpigmented macules on the face. It is exacerbated by sun exposure and oral contraceptive pills. The condition is predominantly seen in females. Melasma is very common in the 3rd trimester of pregnancy and is most marked in brunettes. It occurs due to increased levels of estrogen and progesterone, stimulating the activity of melanocytes. It resolves on its own after pregnancy. Hypermelanosis affects mainly the upper lip, the malar regions, forehead, and the chin.

Solution to Question 8:

The image showing brown patches in axilla in a diabetic patient is suggestive of erythrasma. Coral■red fluorescence under Wood's lamp is seen in erythrasma.

Erythrasma is a skin infection caused by Corynebacterium minutissimum. It is common in diabetics and immunodeficient patients and is associated with obesity, sweating, and poor hygiene. It affects the axillae, groins, toe webs, inter-gluteal and submammary regions. Lesions may be symptomless or show mild discomfort and itching.

The lesions may be of irregular shape with sharp margins. They may start as smooth, red patches and then become brown. The lesions are more likely to be pruritic in tropical regions and may be asymptomatic in a temperate climate.

Coral■red fluorescence with Wood's light is due to coproporphyrin III and suggests erythrasma. The fluorescence may persist after the eradication of the disease.

Treatment of erythrasma is by using topical clotrimazole and miconazole for two weeks. For more extensive lesions, erythromycin can be administered. Relapse can occur. In such cases, long-term topical povidone-iodine application with drying agents can be used. Photodynamic therapy can also be attempted.

Green and blue-green fluorescence is seen in tinea capitis infections associated with microspora species and favus.

Pityriasis versicolor is caused by Malassezia species and the typical lesions are asymptomatic hypopigmented or hyperpigmented macules and patches and produce fine scales. The color of the scales may vary from pale ochre to medium brown. Pale yellow fluorescence seen in Wood's lamp examination is characteristic of tinea versicolor or pityriasis versicolor.



Solution to Question 9:

The finding shown in the image is Gottron's papules. The given scenario of a patient with weakness, rash, muscle inflammation, and presence of Gottron's papules on the skin is highly suggestive of dermatomyositis.

Dermatomyositis (DM) is an autoimmune disorder predominantly affecting the skin and skeletal muscle. The following are various manifestations of dermatomyositis:

• Proximal myopathy: Patients have trouble performing everyday activities like climbing stairs, getting up, and lifting their arms above their heads

• Gottron's papules: Violaceous, erythematous papules overlying the dorsal interphalangeal or metacarpophalangeal, elbow, or knee joints, as shown in the image below

• Gottron's sign: Symmetric, non-scaling, violaceous, erythematous macules or plaques, often atrophic, in the same distribution as Gottron's papules

• Shawl sign/V-sign: Fixed macular erythema, across the shoulders and outer arms (Shawl distribution)

• Periungual telangiectasias: Dilated capillaries of the nail folds

• Mechanic's hand: Roughening and cracking of the skin of the tips and sides of the fingers, resulting in irregular, dirty-appearing lines that resemble those of a mechanic or manual laborer

• Less common manifestations include: Facial swelling, malignancy, erythroderma, lichen planus

Laboratory evaluation of patients with dermatomyositis will reveal elevated levels of serum creatine kinase and antinuclear antibodies. Electromyography will reveal features of myopathy. Histology will show perifascicular atrophy. Immunohistochemical staining will reveal myxovirus resistance protein A, a sensitive and specific marker.

Treatment of dermatomyositis:

• Immunotherapy with high-dose glucocorticoids - first-line drug

• Methotrexate - second-line drug; indications include the following

• When the patient presents with severe weakness, or other organ system involvement

• Patients with a risk of steroid complications like patients with diabetes, osteoporosis, or postmenopausal women

• Patients who do not show improvement after 2–4 months of treatment with prednisone or those who cannot be maintained on a low dose

• Triple therapy with intravenous immunoglobulin or rituximab may be required

Solution to Question 10:

Non-scarring alopecia is caused by androgenic alopecia, alopecia areata, and telogen effluvium (1, 2 and 3). Frontal fibrosing alopecia is a type of primary cicatricial (scarring) alopecia.

Alopecia areata is a chronic inflammatory disease that causes non-scarring hair loss. It is a T-cell-mediated inflammatory disease. It is associated with several autoimmune diseases, including thyroiditis, lupus erythematosus, vitiligo, and psoriasis. It mainly affects the scalp, but can also affect the beard, eyebrows, and eyelashes. In some cases, this progresses to a total loss of scalp hair (alopecia totalis) or a loss of all hair on the body (alopecia universalis).

Telogen effluvium is a type of nonscarring alopecia that is marked by diffuse shedding of normal hair. This occurs due to premature termination of the anagen phase of the hair cycle. It can be either acute or chronic.

• Acute telogen effluvium occurs 2–3 months after a stressful event such as high fever, surgery, starvation, and hemorrhage. Hormonal fluctuations during the postpartum period can trigger it (telogen gravidarum). Telogen effluvium can result in a diffuse reduction in hair density but never total baldness. It generally resolves spontaneously.

• Chronic telogen effluvium is used to describe the diffuse shedding of telogen hair for more than 6 months. Common causes include thyroid disorders, iron-deficiency anemia, and malnutrition.

Androgenetic alopecia is a disorder characterized by a reduction of hair fiber production by follicles and miniaturization. Dihydrotestosterone, an androgenic metabolite, is implicated in the etiology. It is of two types: male-pattern and female-pattern hair loss. Male-pattern hair loss causes frontal hairline recession and thinning of the vertex. Female-pattern hair loss involves the mid-frontal scalp diffusely. There is no or minimal involvement of the temporal region.

Solution to Question 11:

The correctly matched pairs are as follows:

PUVA is highly effective in clearing psoriasis with clearance rates of over 90%. It is associated with both acute (erythema, blistering) and chronic side effects. Long-term PUVA use is associated with the development of non-melanoma skin cancer (squamous cell cancer and basal cell cancer).

Cyclosporine is associated primarily with renal impairment (decreased GFR), metabolic disturbances, and an increased risk of malignancy. Due to its nephrotoxic effects, it is now uncommonly used for the maintenance of psoriasis.

Both acitretin and methotrexate are teratogenic.

Acitretin is associated with dose-dependent mucocutaneous side effects (mucositis) and hyperlipidemia. To prevent retinoid-induced embryopathy, female patients should avoid conception for 3 years after discontinuing acitretin treatment.

Acitretin is approved for use in the cutaneous manifestations of psoriasis.

Absolute contraindications:

• Pregnant women

• Women who are planning to conceive

• Breastfeeding

Methotrexate is also associated with mucositis and teratogenicity, it has been used as an abortifacient and is used in the management of molar pregnancies. Methotrexate is also associated with hepatotoxicity.

Column A	Column B
1. PUVA	c. Long-term use increases th e risk of skin cancer
2. Cyclospori ne	b. Effective but is highly nep hrotoxic
3. Methotrexa te	a. Abortifacient
4. Acitretin	d. Mucositis, teratogenic

Solution to Question 12:

The skin manifestations are most commonly due to metronidazole, which is associated with fixed drug eruption (FDE).

Although maculopapular rashes can also be seen with metronidazole, the incidence of a fixed drug eruption is higher with metronidazole.

Fixed drug eruptions/ reactions are characterized by sharply demarcated, dull red to brown lesions. They occur on lips, hands, legs, face, genitalia, and oral mucosa, and can cause a burning sensation. They are usually solitary or few in number and recur at the exact location. But, new lesions can occur on different body parts on re-exposure to the offending drug.

FDE is a type of delayed hypersensitivity reaction. Paracetamol/NSAIDs are some of the most common drugs causing FDE.

Drug■specific clinical patterns of FDE include:

Diagnosis is based on history and lesion morphology. Discontinuation of the offending drug is the first step in management.

The image below shows a well-defined hyperpigmented patch of a fixed drug eruption.

Drugs	Drug-specific clinical pattern of FDE
NSAIDs	Lips and genitals
Tetracycline/ Trimethoprim-sulfamethoxazole	Genitals
Metamizole	Trunk and extremities
Carbocysteine	Face

Solution to Question 13:

The clinical scenario points to a case of leprosy. The diagnosis of this condition requires nerve examination, skin biopsy, and a slit skin smear (option B- 1, 2, 3).

Leprosy is a chronic granulomatous condition caused by Mycobacterium leprae. It affects the skin, cutaneous and peripheral nerves, and mucous membranes. The most commonly affected peripheral nerves are the posterior tibial nerve, followed by the ulnar, median, and lateral popliteal nerves. The most commonly involved cranial nerve is the facial nerve.

Leprosy has a long incubation period, usually ranging from 2 to 12 years. It commonly presents as an area of numbness on the skin or a visible skin lesion which could be a hypopigmented or erythematous macule. Lesions usually involve the face, extensor aspect of the limbs, trunks, and buttocks. The lesions have impaired nerve function and poor hair growth. The patients may also present with burns or ulcers of the anesthetic limb.

Classification systems of leprosy:

• Ridley-Jopling classification:

• Tuberculoid (TT)

• Borderline tubercular (BT)

• Borderline (BB)

• Borderline lepromatous (BL)

• Lepromatous (LL)

• WHO classification:

• Paucibacillary

• Multibacillary

Cardinal signs for the diagnosis of leprosy:

• Characteristic skin lesions- hypopigmented or erythematous lesions with definite impairment of sensation.

• Peripheral nerve involvement- thickening of the nerve with sensory impairment. Nerve tenderness can also be seen.

• Detection of acid-fast bacilli in slit-skin smear, skin biopsy, or positive biopsy PCR.

The presence of two of the three signs can establish the diagnosis of leprosy. Other tests done for diagnosis of leprosy include nerve biopsy in pure neural leprosy and PGL-1 antibody in cases with low bacterial load.

Solution to Question 14:

The given image shows lichen sclerosus.

Lichen sclerosus usually presents in postmenopausal women. Multiple etiologies have been suggested (infectious, hormonal, genetic, and autoimmune). Symptoms include itching, excoriations, vulval skin thinning and crinkling, dysuria, and dyspareunia. Perianal involvement is frequently seen in the form figure of 8 appearance. The typical appearance is white, atrophic papules may coalesce into plaques and distort normal anatomy. Labia minora regression, clitoral concealment, urethral obstruction, and introital stenosis can develop. A biopsy is needed to exclude malignancy.

Topical corticosteroids remain the first line of treatment.

Premalignant lesions of vulva:

• Vulvar intraepithelial neoplasia (VIN)

• Paget's disease

• Lichen sclerosus

• Squamous cell hyperplasia

• Condyloma accuminata

• Bowen's disease (erythroplasia of Queyrat)

Lichen planus is an inflammatory disorder affecting the skin and mucosal membranes. Nail involvement includes exaggeration of the longitudinal lines and linear depression. It manifests as 6Ps pruritic, purple, polygonal, planar, papules, and plaques. It also exhibits Wickham striae (thin white lacy lines on the lesion), and the Koebner phenomenon (new lesions at sites of skin trauma). Mucosal involvement is common including mouth and tongue, rarely it can also be found on the anus and genitalia

The histological changes in lichen planus include increased Langerhans cells in the epidermis, spongiosis, and sawtooth appearance of Rete ridges. A fully developed lichen planus papule shows acanthosis of the epidermis, thickening of the granular layer, and hyperkeratosis. The image below shows the histopathological changes in lichen planus.



Vulvar Paget disease is an intraepithelial neoplasia that presents as an eczematoid, red, weeping area on the vulva. It is associated with invasive Paget disease or adenocarcinoma of the vulva or carcinoma at another non-vulvar location. It can be managed by local excision or vulvectomy. This condition needs long-term surveillance as recurrence is very common.

Angioma of vulva - Hemangiomas and angiokeratomas are benign lesions of the vulva. Angiokeratomas are benign tumors characterized by hyperplasia of the epidermis along with multiple dilated vessels in the superficial dermis. They present as well-defined, erythematous, hyperkeratotic papules. Senile (cherry) hemangiomas usually present as red macules that rapidly progress to well-circumscribed, raised, red, and soft lesions. Excision biopsy of bleeding hemangiomas, cryosurgery, or carbon dioxide laser can be used for management.

The image below shows a cherry angioma:

Solution to Question 15:

The clinical scenario is suggestive of systemic sclerosis, and the skin manifestation shown is a salt-and-pepper appearance.

Systemic sclerosis is an autoimmune disease of unknown etiology, complex pathogenesis, and variable clinical presentations. It is associated with significant disability and mortality. Virtually every organ is affected. Thickened and indurated skin, called scleroderma, is the distinguishing hallmark of systemic sclerosis.

Cutaneous features of systemic sclerosis:

• Raynaud phenomenon, with the typical white discoloration of the fingers caused by vasospasm and bluish discoloration of the fingertips

• Loss of pigmentation except in the perifollicular areas is known as the salt and pepper appearance of the skin, which is mostly seen in the upper back, chest, and scalp

• Puffy fingers in early disease

• Sclerodactyly and sclerosis extending proximal to the metacarpophalangeal joints with or without contractures

• Vasculopathic ulcers over the bony prominences

• Loss of fingertip pulp

• Severe flexion contractures

• Facial features include expressionless facies, mat■like telangiectases, microstomia, perioral furrowing, and a beak■like nose

Other options:

Option A: Poikiloderma is a skin disorder characterized by hypopigmentation and hyperpigmentation, wrinkling following atrophy, and telangiectasias. It is seen in ionizing radiation-damaged skin and dermatomyositis.

Option C: The hair in the amelanotic lesions of vitiligo can sometimes lose color and become white. This is called leukotrichia.

Option D: Vitiligo is characterized by depigmented macules that have sharp demarcation. It is due to the loss of melanocytes.

Solution to Question 16:

The diagnosis in this patient with the above clinical features is psoriasis.

The image shows nail pitting, oil drop sign, and salmon patches in fingernails

Psoriasis is a chronic inflammatory condition of the skin with systemic manifestations. The skin lesions can have exacerbations and relapse. They include red, silvery-white well-demarcated scales. A peripheral clearing known as the ring of Woronoff may be present. It is usually seen on the scalp, trunk, and extensor surfaces of the limbs. The appearance of a smooth, glossy red membrane with small bleeding points on the removal of the supra-papillary epidermis by gentle scraping is known as Auspitz's sign.

Psoriatic arthritis presents as early morning stiffness and joint swelling, heel pain due to enthesitis of the Achilles tendon, or plantar fasciitis. The clinical examination may reveal evidence of dactylitis (sausage fingers) or swollen or tender joints. The distal interphalangeal joint (DIP) is classically involved in causing pencil-in cup deformity.

Punctate pitting is the most common nail finding and others include the oil-drop sign, subungual hyperkeratosis, splinter haemorrhages, and distal onycholysis.

The topical application of steroids, retinoids, vitamin D, and emollients is the mainstay of management. Phototherapy and oral methotrexate, acitretin, and cyclosporine are used for more extensive and systemic diseases.

Dermatology NEET 2022

Question 1:

A patient presents to you with multiple anogenital warts. The biopsy of these lesions showed squamous atypia. Which of the following human papillomavirus types are considered high-risk?

a) HPV 2

b) HPV 18

c) HPV 6

d) HPV 11

Question 2:

A farmer presents to you with a swelling on the foot and multiple discharging sinuses in the lower limb. Granules from the discharge were examined under the microscope, which is shown below. Which of the following is true regarding this condition?



a) Both bacteria and fungi can be causative

b) Undergoes lymphatic spread

c) There is lymphocyte accumulation

d) Involves only superficial tissues

Question 3:

A farmer presents to you with a cauliflower-shaped mass on the foot, which developed after a minor injury. Microscopy shows copper penny bodies. What is the most likely diagnosis?

a) Chromoblastomycosis

b) Blastomycosis

c) Sporotrichosis

d) Phaeohyphomycosis

Question 4:

A patient presents with an anesthetic patch in the areas on the face as shown in the image below. Which of the following nerves is the most commonly involved in this condition?

a) Abducens nuclei

b) Facial nerve

c) Optic nerve

d) Trigeminal nerve

Question 5:

A female patient with a body mass index of 30 kg/m2 presents to you with a lesion on the neck as shown below. Which of the following condition is she most likely to be suffering from?

a) Hypothyroidism

b) Metabolic syndrome

c) Addison's disease

d) Hyperparathyroidism

Question 6:

Irregular pitting of nails with subungual hyperkeratosis is seen in _____.

a) Lichen planus

b) Psoriasis

c) Atopic dermatitis

d) Alopecia areata

Question 7:

A 35-year-old woman presents to you with hair loss for the past three months. She tested positive for COVID-19 eight months ago. What is the most likely diagnosis?

a) Tinea capitis

b) Telogen effluvium

c) Trichotillomania

d) Female-pattern androgenic alopecia

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	b
6	b
7	b

Detailed Explanations

Solution to Question 1:

Human papillomavirus (HPV) types 16 and 18 are considered high-risk types that are commonly implicated in cervical, vaginal and vulval cancer. Other high-risk types include HPV types 31, 33,

and 35.

The E6 protein of high-risk types binds to the p53 gene while the E7 protein inactivates the pRb gene. p53 and pRb are tumor suppressor genes that when inactivated results in uncontrolled cell proliferation. The most common type of cancer seen is squamous cell carcinoma. The image below depicts vulval carcinoma which is also associated with HPV infection.

The image given above shows anogenital warts pointing to condyloma acuminata (venereal warts). It is commonly caused by HPV types 6 and 11. However, rarely HPV 16 and 18 can cause it too. It is sexually transmitted. Patients are advised not to involve in intercourse when active genital warts are present. It can appear discrete or may coalesce to resemble cauliflower-like growths. 20% of cases may develop into cervical, vaginal, or vulval cancers. Risk factors include oral contraceptives and pregnancy. The condition is diagnosed by colposcopy.

The presence of koilocytes on histopathological examination is pathognomonic of the virus. These are cells with irregular and hyperchromatic nuclei, perinuclear halo, and peripheral cytoplasmic condensation. These cells disappear when there are dysplastic changes.

The image below depicts koilocytes.



Other options:

Option A - HPV type 2 is responsible for common warts in non-genital regions.

Options C and D - HPV types 6 and 11 are responsible for anogenital warts and laryngeal papillomatosis.

Solution to Question 2:

The clinical vignette along with the image showing sulfur granules is consistent with the diagnosis of mycetoma. Both bacteria and fungi can be causative agents of the disease.

Mycetoma is a chronic, slow-progressing subcutaneous infection that commonly affects the foot. Other sites which can be affected include the hands, upper back, and posterior part of the neck.

The mode of infection is usually traumatic. Agricultural workers are more prone to infection.

There are three types - eumycetoma, actinomycetoma, and botryomycosis. Eumycetoma is caused by fungal species such as Madurella mycetomatis and Madurella grisea. Adjacent fascia and bones may be involved as the disease progresses.

Actinomyces israelii and Nocardia asteroides result in the bacterial form - actinomycetoma. It is generally more invasive and can spread via the fascial planes to the underlying muscle or bone. Dissemination is rare but can occur in lesions of the head and neck.

Staphylococcus aureus and other pyogenic bacteria can cause botryomycosis.

Initially, it starts off as a small subcutaneous swelling of the foot, which then enlarges to form multiple discharging sinuses. The discharge contains granules which are microcolonies of the causative organism. Different causative organisms produce granules that are characteristic in color, consistency, and size, which helps in delineating the agent in each case.

The granules in eumycetoma are generally hard, while that in actinomycetoma is soft in consistency.

Medical management of eumycetoma includes ketoconazole and miconazole, while that of actinomycetoma includes dapsone or rifampicin. It should be continued for 18-24 months even after symptomatic recovery or the absence of laboratory evidence for infection.

Solution to Question 3:

The most likely diagnosis based on the clinical vignette is chromoblastomycosis.

Chromomycosis is a term used to describe various conditions caused by pigmented (dematiaceous) fungi. Chromoblastomycosis, the most common chromomycosis, is more commonly seen in tropical areas among agricultural workers and woodcutters who walk barefoot. Traumatic implantation is the mode of infection.

Patients present with warty cauliflower-like cutaneous nodules on the feet and lower legs. Species of the fungal genus Fonsecaea are implicated as the causative agents. Diagnosis is made by the demonstration of sclerotic bodies (copper-penny bodies) in tissue sections. Sclerotic bodies appear as round or irregular, dark brown bodies with septae.



It can be managed with itraconazole (100–200 mg daily) or terbinafine (250 mg daily). For small lesions, surgical excision is the therapy of choice.

Other options:

Option B - Blastomycosis is an infection caused by Blastomyces dermatitidis. It causes a chronic infection with granulomatous and suppurative lesions, which starts in the lungs but may disseminate to other organs. Symptoms of an acute lower respiratory tract infection with pulmonary infiltrates are the most common clinical presentations. Microscopic examination of stained specimens shows broad-based budding yeast cells.



Option C - Sporotrichosis caused by the dimorphic fungus Sporothrix schenckii, is commonly seen in gardeners followed by minor trauma. Patients present with skin nodules or lymph node ulceration. Microscopy shows asteroid bodies. It can be seen as an oval, basophilic structure with eosinophilic radiation.



Option D - Phaeohyphomycosis caused by dematiaceous fungi is frequently encountered in immunodeficient individuals. Clinical manifestations include brain abscesses, intramuscular abscesses, and cysts. On microscopy, the fungi appear as distorted hyphal strands. In contrast to chromoblastomycosis, sclerotic bodies are not found.



Solution to Question 4:

The clinical scenario points to a case of leprosy and the most commonly involved cranial nerve is the facial nerve.

Leprosy is a chronic granulomatous condition caused by Mycobacterium leprae. It affects the skin, cutaneous and peripheral nerves, and mucous membranes. The most commonly affected peripheral nerves are the posterior tibial nerve followed by ulnar, median, and lateral popliteal nerves. The most commonly involved cranial nerve is the facial nerve.

Leprosy has a long incubation period, usually ranging from 2 to 12 years. It commonly presents as an area of numbness on the skin or a visible skin lesion which could be a hypopigmented or erythematous macule. Lesions usually involve the face, extensor aspect of the limbs, trunks, and buttocks. The lesions have impaired nerve function and poor hair growth. The patients may also present with burns or ulcers of the anesthetic limb.

Classification systems of leprosy include:

• Ridley Jopling classification:

• Tuberculoid (TT)

• Borderline tubercular (BT)

• Borderline (BB)

• Borderline lepromatous (BL)

• Lepromatous (LL)

• WHO classification:

• Paucibacillary

• Multibacillary

The cardinal signs for the diagnosis of leprosy include:

• Characteristic skin lesions- hypopigmented or erythematous lesions with definite impairment of sensation.

• Peripheral nerve involvement- thickening of the nerve with sensory impairment. Nerve tenderness can also be seen.

• Detection of acid-fast bacilli in slit-skin smear, skin biopsy, or positive biopsy PCR.

The presence of two of the three signs can establish the diagnosis of leprosy. Other tests done for diagnosis of leprosy include nerve biopsy in pure neural leprosy and PGL-1 antibody in cases in low bacterial load.

Solution to Question 5:

The patient is most likely to be suffering from metabolic syndrome based on the clinical scenario given and the image depicting acanthosis nigricans.

Metabolic syndrome (syndrome X) is a group of conditions that increases the risk for cardiovascular disease. These include hypertension, hyperglycemia, hypertriglyceridemia, and central obesity. Increasing age, genetics, and sedentary lifestyle are considered risk factors.

Insulin resistance is the most accepted hypothesis used to explain the pathophysiology of the condition. An elevated level of circulating fatty acids is considered a major factor in the development of insulin resistance. It is associated with acanthosis nigricans which are velvety hyperpigmented lesions in skin flexures such as the axilla, groin, and back of the neck. Acanthosis nigricans can also be seen in malignancies, especially those of the gastrointestinal tract, acromegaly, and Cushing's syndrome.

Other conditions associated with metabolic syndrome include polycystic ovary syndrome, obstructive sleep apnea, and non-alcoholic fatty liver disease. The condition is managed by dietary and lifestyle modifications. Individual components of the syndrome also need to be addressed by the use of statins, antihypertensives, or hypoglycemic agents.

Other options:

Option A - Hypothyroidism is more common in women and can be either congenital or acquired. Autoimmune and iatrogenic hypothyroidism are the common causes of the latter. Symptoms include dry coarse skin, weight gain, poor appetite, fatigue, and alopecia.

Option C - Addison's disease (primary adrenal insufficiency), most commonly caused by autoimmune adrenalitis presents with hyperpigmentation of the skin, especially in areas exposed to sunlight or prone to friction. This is due to excessive stimulation of melanocytes by the raised adrenocorticotropic hormone (ACTH). This is seen in chronic cases.

Option D - Hyperparathyroidism is a disorder that arises due to excessive secretion of parathyroid hormone. Solitary adenomas are the most common cause. It primarily affects the renal and skeletal systems. Sestamibi scan is the imaging of choice to delineate parathyroid abnormalities.

Solution to Question 6:

Irregular pitting of nails with subungual hyperkeratosis is seen in psoriasis.

Psoriasis is a chronic inflammatory condition of the skin with systemic manifestations. Plaque psoriasis is the most common type. The nails are frequently involved, and punctate pitting is the most common finding seen. Nail beds can also be affected, giving rise to the oil-drop sign—a highly specific sign seen in this condition. Other manifestations include subungual hyperkeratosis, splinter hemorrhages, and distal onycholysis.

The skin is involved in the form of red/ salmon pink, scaly, and well-demarcated lesions. A peripheral clearing (ring of Woronoff) may be present. It is usually seen on the scalp, trunk, and extensor surfaces of the limbs. The lesions are surrounded by silvery-white scales. The appearance of a smooth, glossy red membrane with small bleeding points on the removal of the supra-papillary epidermis by gentle scraping (Auspitz's sign) is characteristic.

Psoriasis is associated with Koebner or isomorphic phenomenon. It describes the appearance of new lesions along the site of trauma (linear lesions). It is also observed in lichen planus.

Topical application of steroids, retinoids, vitamin D, and emollients is the mainstay of management. Phototherapy and oral methotrexate, acitretin, and cyclosporine can also be used. Both methotrexate and acitretin are teratogenic drugs. It is recommended that women do not conceive for three years after cessation of the acitretin.

Other options:

Option A - Lichen planus is an inflammatory disorder affecting the skin and mucosal membranes. Nail involvement includes exaggeration of the longitudinal lines and linear depression. This is due to nail plate thinning. If the nail bed is involved, it can result in hyperpigmentation, subungual hyperkeratosis, and onycholysis. In some cases, pterygium unguis can be seen due to adhesion between the dorsal nail fold and the nail bed.

Option C - Atopic dermatitis is a relapsing chronic inflammatory skin disease. Patients present with itchy erythematous papules/papulo-vesicles in flexural regions. Nail involvement is in the form of coarse pitting and ridging.

Option D - Alopecia areata is a condition that results in non-scarring hair loss. It can cause fine stippled pitting of the nails, roughening of the nail plate (trachyonychia), or atrophic dystrophy.

Solution to Question 7:

The most likely diagnosis based on the clinical scenario is telogen effluvium.

The hair follicle goes through three phases known as the hair cycle:

• Anagen: Phase of active hair growth

• Catagen: Involutionary phase

• Telogen: Time period between catagen and next anagen phase

Telogen effluvium is a type of nonscarring alopecia that is marked by diffuse shedding of normal hair. This occurs due to premature termination of the anagen phase of the hair cycle. It can be either acute or chronic.

Acute telogen effluvium occurs 2-3 months after a stressful event such as high fever, surgery, starvation, and hemorrhage. Hormonal fluctuations during the postpartum period can trigger it (telogen gravidarum). Telogen effluvium can result in a diffuse reduction in hair density but never

total baldness. It generally resolves spontaneously.

Chronic telogen effluvium is used to describe the diffuse shedding of telogen hair for more than 6 months. Common causes include thyroid disorders, iron-deficiency anemia, and malnutrition.

Other options:

Option A - Tinea capitis is used to describe an infection of the scalp by the dermatophyte group of fungi. The most common pathogenic species are Trichophyton tonsurans and Trichophyton violaceum. The presentation can vary from hair breakage associated with scaling to widespread inflammatory mass on the scalp.

Clinical variants of tinea capitis:

• Endothrix - A non-inflammatory type in which multiple black dots are present within the areas of alopecia

• Ectothrix - A mild inflammatory type in which single or multiple scaly patches with hair loss are seen

• Kerion - A severe inflammatory type that presents with an inflammatory mass with thick crusting and matting of hair with pus and sinus formation

• Favus - Characterized by the presence of yellowish, cup■shaped crusts known as scutula that develop around hair follicles

Observation under KOH mount is used for diagnosis.

Option C - Trichotillomania (trichotillosis) is considered an obsessive-compulsive disorder (OCD) spectrum disorder, marked by gratification from pulling one's own hair. It could be associated with underlying anxieties or depression. The vertex region of the scalp is most commonly involved. Scalp histology will show perifollicular hemorrhages and intrafollicular pigment casts.

Option D - Androgenic hair loss is used to describe a progressive decrease in hair fiber production. Dihydrotestosterone, an androgenic metabolite, is implicated in the etiology. Female-pattern hair loss involves the mid-frontal scalp diffusely. There is no or minimal involvement of the temporal region.

Stage 1
Stage 2
Stage 3
Stage 4
Stage 5

Dermatology NEET 2023

Question 1:

A lady developed a skin reaction as shown in the image after using a hair dye. Which of the following chemicals is responsible for this condition?

a) Pollen

b) Chromates

c) Balsam of Peru

d) p-Phenylenediamine

Question 2:

A female patient presented with acne that is not resolving on oral isotretinoin and antibiotics therapy. Which of the following is the next best investigation?

a) Look for dietary triggers

b) Evaluate for hyperandrogenism

c) Check for antibiotic resistance

d) Look for drug triggers

Question 3:

A young woman presents with complaints of a painless ulcer in the genital area. It is associated with non-tender inguinal lymphadenopathy. What is the most likely diagnosis?

a) Chancroid

b) Syphilis

c) Herpes genitalis

d) Granuloma inguinale

Question 4:

Which of the following is associated with the clinical condition shown in the image?

a) Cataract

b) Glaucoma

c) Malignant melanoma

d) Basal cell carcinoma

Answer Key

Question No.	Correct Option
1	d
2	b

3	b
4	c

Detailed Explanations

Solution to Question 1:

This is most likely a case of hypersensitivity reaction caused by p-phenylenediamine (PPD), which is present in hair dyes.

Hair dyes such as PPD have the potential to cause contact dermatitis. 10% of PPD users tend to develop contact dermatitis. A patch test is recommended 24-48 hours before its usage either on the forearm or behind the ear. The appearance of any redness, blistering, or swelling implies allergy; therefore, the application of dye is not recommended. Cross-sensitization with other aromatic benzenes is also noted, which needs to be avoided.

Hair dyes can trigger acute edema and intense pruritus, followed by desquamation. Hair dye can elicit an allergic reaction on the forehead. PPD in the hair dye can also cause an erythema multiforme–like reaction.

In PPD-associated allergies, there is edema and weeping of the scalp margins, eyes, and ears. There may be relative sparing of the scalp. Immediate hypersensitivity reactions present as urticaria, and contact anaphylaxis have also been noted to occur in patients using PPD. To minimize the risk of an allergic reaction to hair dye, individuals should follow the manufacturer's instructions, perform a patch test, and avoid over-processing the hair. Treatment for PPD-induced contact dermatitis may include topical corticosteroids, oral antihistamines, and in severe cases, oral corticosteroids.

Other options:

Option A: Pollen is not typically found in hair dyes and is not responsible for this condition. Pollen allergies usually present as hay fever or seasonal allergic rhinitis.

Option B: Chromates, which can be found in cement, leather, and paints, are not typically present in hair dyes and are not responsible for this skin reaction.

Option C: Balsam of Peru, a fragrant resin, is found in various cosmetic and food products but is not a common component of hair dyes. It is less likely to be responsible for this condition compared to PPD.

Solution to Question 2:

The next best step for a female patient with acne that is resistant to oral isotretinoin is to check for hyperandrogenism.

Hyperandrogenism usually manifests as hirsutism, acne, and/or androgenic alopecia. Acne that is persistent, moderate to severe, or late-onset raises concern for PCOS or other causes of hyperandrogenism.

The pathogenesis of acne vulgaris involves

• Blockage of the follicular opening by hyperkeratosis

• Sebum overproduction

• The proliferation of commensal Propionibacterium acnes

• Inflammation

Management is directed at minimizing inflammation, decreasing keratin production, antibiotics, and antiandrogens to diminish sebum production.

Other options:

Option A: Looking for dietary triggers could be considered, but there is limited evidence to support a direct relationship between specific food items and acne.

Option C: Since the patient has not shown improvement even with antibiotic therapy, it is reasonable to consider checking for antibiotic resistance. However, evaluating for hyperandrogenism should be prioritized due to its potential role in the patient's condition.

Option D: Looking for drug triggers may be relevant in some cases, but it is less likely to be the primary cause of acne that is not responding to isotretinoin and antibiotics. Drug-induced acne typically resolves once the offending drug is discontinued.

Solution to Question 3:

The given clinical scenario of a painless ulcer and non-tender lymphadenopathy is suggestive of primary syphilis.

The incubation period of syphilis is 9-90 days and varies inversely with the size of the Spirochaete inoculum.

Stages of syphilis:

• Primary chancre: Painless, indurated, button-like ulcer with painless lymphadenopathy.

• Secondary syphilis is the stage when generalized manifestations of syphilis appear on the skin and mucous membranes. A macular rash is commonly seen and this stage may be preceded by constitutional symptoms such as fever, sore throat, and malaise.

• Latent syphilis is the stage when there are no clinical features of the active disease but serological testing for the disease is positive.

• Late syphilis (tertiary syphilis) commonly presents as superficial or nodular punched-out ulcers or gummas (granulomas appearing as cutaneous plaques or nodules with central ulceration) that appears 3-10 years after infection. It may also present as red nodules (nodular syphilides), neurosyphilis, and cardiovascular syphilis.

Serological testing is done using standard non-treponemal antibody tests like VDRL and rapid plasma reagin (RPR). Specific treponemal antibody tests like Treponema pallidum hemagglutination assay (TPHA) and fluorescent treponemal antibody absorption test (FTA-ABS) are further required. Microscopically, T. palladium is identified with characteristic corkscrew morphology and motility on darkfield microscopy.

Parenteral penicillin G is the drug of choice for all stages of syphilis. Treatment of recent sexual contacts is recommended.

Other options:

Option A: Chancroid presents with a painful genital ulcer and tender inguinal lymphadenopathy. It is caused by the bacterium Haemophilus ducreyi.

Option C: Herpes genitalis is characterized by painful vesicular lesions that progress to ulcers in the genital region, caused by the herpes simplex virus (HSV). It is usually associated with localized tender lymphadenopathy.

Option D: Granuloma inguinale (also known as donovanosis) is caused by the bacterium Klebsiella granulomatis. It presents as a slowly progressive, painless, beefy-red ulcerative lesion in the genital region, often without lymphadenopathy.

Solution to Question 4:

The given image showing a black, heterogeneous, hairy palpable lesion in a child is suggestive of congenital melanocytic nevus. Malignant melanoma is more commonly associated with a congenital melanocytic nevus.

Congenital melanocytic nevi (CMN) are benign, pigmented, melanocytic nevi present at birth. They are most commonly brown, black, purplish, or red, usually a palpable lesion with heterogeneous color and texture. CMN often have thick hair at birth, which grows at a significantly greater rate than the surrounding scalp hair.

Congenital melanocytic nevi can be classified based on size: small (<1.5 cm), medium (1.5-19.9 cm), and large or giant (>20 cm). Larger CMN have a higher risk of developing malignant melanoma. Management of CMN depends on the size and location of the lesion. Smaller nevi may be observed, while larger nevi or those at a higher risk for complications may require surgical excision, particularly if there is a concern for malignant transformation. Complications of CMN include neurological abnormalities (intraparenchymal melanosis is the commonest finding), malignant melanoma, and other tumors like rhabdomyosarcoma.

The lifetime risk of malignant transformation in large or giant CMN ranges from 5% to 10%. Regular follow-up and monitoring for changes in size, color, or texture are essential for early detection of potential complications.

Note: More than 50% of patients with Sturge-Weber syndrome have glaucoma as an ocular complication. The image given below shows port-wine staining in a child with Sturge-Weber syndrome.

DAILYROUNDS

Dermatology INI-CET 2023

Question 1:

A patient on Anti PD1 immunotherapy(Nivolumab) comes with pruritus, eczema, and urticarial rashes. Histopathological examination reveals a subepidermal blister. The direct Immunofluorescence of the lesions showed Linear C3 and IgG on the basement membrane. What is the diagnosis?

a) Pemphigus vulgaris

b) Bullous pemphigoid

c) Dermatitis herpetiformis

d) Pemphigus foliaceous

Question 2:

A 24 year old young male presented with asymptomatic scaly lesions over back, symmetrical in distribution. Images of the lesions are given, Identify the disease and the pattern of lesions

a) Collaratte scale-Christmas tree pattern-Pityriasis rosea

b) Wickhams striae-Lichen planus

c) Christmas tree pattern-Lichen planus

d) Wornoff ring-Seborrhic dermatitis

Question 3:

Statement 1: Histopathological examination of a patient with Flaccid bulla and painful erosions showing suprabasal blister and inflammatory infiltrate in the blister cleft. Statement 2: The row of tombstones appearance is diagnostic of Pemphigus vulgaris.

a) Statement 1 is false and Statement 2 is true

b) Statement 1 is true and Statement 2 is false

c) Both the statements are true but 2 is not the conclusion of 1

d) Both the statements are true but 2 is the conclusion of 1

Question 4:

Identify the following lines in dermatology



a) Hinderer lines

b) Marionettes line

c) Relaxed skin tension lines

d) Blaschko lines

Question 5:

Match the following wood lamp findings with the disease

a) 1-a,2-c,3-d,4-b

b) 1-e,2-c,3-d,4-b

c) 1-e,2-d,3-c,4-b

d) 1-a,2-d,3-c,4-b

Question 6:

Consider the causes of alopecia. Which among the following cause(s) non-scarring alopecia?

a) 3 and 4

b) 1, 2, and 3

c) 2, 3, and 4

d) Only 4

Question 7:

A farmer from Tamil Nadu presents with fever, body ache, and cough for 3 days. Lab studies showed thrombocytopenia. On examination, a black lesion is present, as in the picture below. IgM scrub typhus-positive. What drug will be used in the management?

a) Doxycyline

b) Ceftriaxone

c) Ceftriaxone+Doxycycline

d) Doxycycline + Azithromycin

Question 8:

A young male complains of an itchy rash over his back as seen below. What is the most likely sign and diagnosis?



a) Collarette scale-pityriasis rosea

b) Wickham's striae - lichen planus

c) Woronoff's sign - psoriasis

d) Christmas tree appearance - psoriasis

Question 9:

An overweight 12-year old girl has the following blackish discolouration on her neck, as shown below. What is this condition and its' association?

a) Acanthosis nigricans - Insulin resistance

b) Stria Albicans - Cushing syndrome

c) Xeroderma pigmentosum - Malignancy

d) Cafe au lait spots - McCune Albright syndrome

Question 10:

Which of the following are cause(s) of non-scarring alopecia?

a) 3 and 4

b) 1, 2 and 3

c) 2, 3 and 4

d) Only 4

Question 11:

A middle-aged Indian man came to your OPD with the lesion as shown in the image below. The most likely diagnosis is

a) Squamous cell carcinoma

b) Basal cell carcinoma

c) Lupus vulgaris

d) Nevus

Question 12:

A patient with several skin lesions undergoes an investigation and an image of it is shown below. What is this test?



a) Indirect immunofluorescence

b) Direct immunofluorescence

c) Immunohistochemistry

d) Fluorescent in-situ hybridization

Question 13:

A patient presents with vesicles and oral lesions. Histopathological examination reveals suprabasal split. Which of the following is the most likely diagnosis?



a) Varicella

b) IgA disease

c) Pemphigus vulgaris

d) Pemphigus foliaceous

Answer Key

Question No.	Correct Option
1	b
2	a
3	c
4	d
5	b

6	b
7	d
8	a
9	a
10	b
11	b
12	b
13	c

Detailed Explanations

Solution to Question 1:

In the given clinical scenario history of use of Anti-programmed cell death 1 (PD1) targeted immune checkpoint inhibitors with HPE showing subepidermal split and direct Immunofluorescence of the lesions showed Linear C3 and IgG on basement membrane is suggestive of bullous pemphigoid.

Anti-programmed cell death 1 (PD1) targeted immune checkpoint inhibitors such as nivolumab and pembrolizumab are associated with an increased risk of development of bullous pemphigoid. Histopathology will show the presence of subepidermal bullae with eosinophils. Immunofluorescent microscopy will show the presence of subepidermal split and deposition of IgG antibodies and C3 at the dermo-epidermal junction.

Bullous pemphigoid is one of the most common subepidermal immunobullous disorders. It is mostly a disease of the elderly. It is characterised by a subepidermal split, with tissue-bound and serum IgG antibodies against BP180, more than BP 230 antigens on hemidesmosomes, necessary to maintain the structure of.

Bullous pemphigoid is associated with psoriasis and treatments of psoriasis, like phototherapy and psoralen. There was an association seen with debilitating neurological conditions like stroke, Parkinsonism, cognitive impairment and multiple sclerosis, and haematological malignancies like lymphomas and myeloid leukaemia. It is also precipitated by drugs like diuretics, penicillamine and psycholeptics.

Clinical features include:

• Prodromal phase with excoriated papules, eczematous or urticarial skin lesions and haemorrhagic crusts may be seen.

• Bullous phase is characterised by:

• Intense pruritus is almost constantly present.

• Symmetrically distributed tense bullae and vesicles over the flexural aspects and abdomen of the body.

• Partly haemorrhagic crusts, erythematous plaques with an annular pattern are seen.

• Lesions heal without scarring, provided there are no superinfections.

• The bullae contain clear, sometimes haemorrhagic exudates.

• Nikolsky's sign is negative.

• Oral or mucosal lesions and milia formation are rare.

The image given below shows the lesions seen in bullous pemphigoid.

Histopathology will show the presence of subepidermal bullae with eosinophils. Immunofluorescent microscopy will show the presence of subepidermal split and deposition of IgG antibodies and C3 at the dermo-epidermal junction.

The below image shows linear IgG deposits at dermo-epidermal junction seen in bullous pemphigoid.



Untreated bullous pemphigoid has a chronic, self-limiting course, with remissions and relapses. Complete remissions are possible with treatment.

Bullous pemphigoid is mainly treated with topical and systemic corticosteroids. Other drugs used like azathioprine, dapsone, doxycycline and immunomodulatory are also used.

Other options:

Option A-Pemphigus vulagaris is an intraepidermal immunobullous disease. It occurs commonly between the 4th to 6th decade. They present with flaccid blisters. Firm pressure over the lesion separates the epidermis from the dermis. Nikolsky's sign is positive. Mucosal involvement will be seen in the form of buccal and palatal erosions

Option C-Dermatitis herpetiformis typically appears in the 4th decade, and the patients present with itching and rash, most typically over the extensor surfaces of the elbows, knees, buttocks, and scalp. The majority of the patient have symptoms related to gluten-sensitive enteropathies, like bloating, diarrhea, and other gastrointestinal complaints. Immunofluorescence microscopy will show the deposition of IgA antibodies in the papillary dermis in a granular or fibrillar pattern. Complement 3 may also be found along with the IgA deposits.

Option D-Pemphigus foliaceous shows antidesmoglein-3 antibodies, anti-desmocollin antibodies and a predominance of anti-desmoglein-1 antibodies. Lack of mucosal involvement in pemphigus foliaceous, where anti-desmoglein-1 antibodies are prominent, is due to the low expression of desmoglein 1 in the mucosa. In pemphigus vulgaris, where anti-desmoglein-3 antibodies are prominent, mucosal involvement is extensive in early stages as desmoglein 3 is abundantly present in the mucosa.

Solution to Question 2:

The above description with central detachment and peripheral attachment of scales with the images suggestive of collarette scales and lesions in christmas tree pattern is suggestive of pityriasis rosea.

Pityriasis rosea/pityriasis circinata/roseola annulata is an acute self-limiting papulosquamous disorder. It is commonly seen in children and young adults.

Clinical features include:

• Prodrome of a sore throat, gastrointestinal disturbance, fever or arthralgia.

• Herald patch - Large and conspicuous eruptions on the thigh, upper arm, trunk or neck covered by fine scales

The image given below shows a herald patch.

• After 5-15 days, it turns into discrete oval lesions, dull pink in color and covered by fine dry silvery-grey scales.

• These secondary lesions occur as macules and papules that are elliptical or ovular, symmetric, and commonly involve the thorax, back, and abdomen.

• It is followed by a marginal collarette of scale with a central wrinkling and atrophy. The collarettes of the scales are attached peripherally. It is also called a hanging curtain sign.

The image given below shows collarette scales.

• The long axes of the lesions characteristically follow the lines of skin tension parallel to the ribs in a Christmas tree pattern on the upper chest and back.

The images given below show Christmas tree pattern along the lines of skin tension on the back and the chest.



The rashes of pityriasis rosea usually resolve by 8 weeks.

Other options:

Lichen planus is an idiopathic inflammatory skin disease affecting the skin and mucous membranes, often with a chronic course with relapses and periods of remission.Patients present with pruritic, purple, polygonal, papules. The lesions have thin white lines on their surface known as Wickham's striae. They are commonly seen on the volar aspect of the wrists, the lumbar region, and around the ankles.The mucosal involvement is seen. The commonest mucosal sites are buccal mucosa and tongue, rarely it can also be found on the anus and genitalia.



A hypopigmented ring surrounding individual psoriatic lesions is known as the Woronoff ring.It is usually associated with treatment, most commonly UV radiation or topical corticosteroids. The pathogenesis of the Woronoff ring is not well understood, but may result from the inhibition of prostaglandin synthesis.

Solution to Question 3:

Both statements 1 and 2 are correct but statement 2 is not the conclusion of statement 1. Statement 2 is not the conclusion of statement 1 because DIF is used for a definitive diagnosis.

Pemphigus vulgaris is a chronic immunobullous disorder characterized by the presence of antibodies against desmosomal proteins expressed in the skin and mucosa. In the skin, desmoglein 1 is mostly seen in all layers, while desmoglein 3 is more in the lower layers. In the oral mucosa, desmoglein 3 is seen in all layers and desmoglein 1 is present in lower layers.

Clinically, the patients present with ill■defined, irregular, slow-to-heal oral lesions and erosions. Flaccid blisters filled with clear fluid may arise on normal skin or an erythematous base. The rupture of such blisters produces painful erosions and heals without scarring. Lesions may also involve the mucosa of the conjunctiva, nasopharynx, larynx, esophagus, urethra, vulva, or cervix.

Nikolsky's sign is positive which is elicited by applying firm tangential sliding pressure with a finger on lesional or perilesional skin. This leads to the separation of the normal■looking epidermis from the dermis, producing erosion.

Direct immunofluorescence is considered the most accurate confirmatory test of this condition. It shows intercellular IgG and C3 deposition on the surface of keratinocytes throughout the epidermis in normal, perilesional skin, in a characteristic fishnet/chickenwire appearance as seen below.



Histology shows a suprabasal split of the epidermis with a characteristic row of tombstone appearance as seen below.



Other investigations include a Tzanck smear showing acantholytic cells. Anti-Dsg ELISA assays can be used to monitor the disease activity.

Systemic corticosteroids are the mainstay of treatment for controlling the disease, improving the quality of life and reducing disease activity.

Solution to Question 4:

The above image shows Blaschko lines as identified by the spiraling over the scalp. Blaschko's lines are lines of normal developmental growth patterns in the skin. They do not correspond to any known nervous, vascular, or lymphatic structures. Conditions that are distributed along these lines include epidermal nevi, and incontinentia pigmenti.



Other options:

Option A: Hinderer's lines are 2 intersecting lines—one drawn from the ala to the tragus and the other from the oral commissure to the lateral canthus.

Option B: Melomental folds, or marionette lines, are one of the consequences of facial aging. The curvilinear wrinkles form because of facial movements and the aging process extend downward from the oral commissures.

Option C: Relaxed skin tension lines run perpendicular to the direction of contraction of the underlying muscles and parallel to the dermal collagen bundles.To produce favourable aesthetic results, incisions should generally be designed to follow rhytids or the relaxed skin tension lines as the resulting scars will be stronger and less likely to stretch.

Solution to Question 5:

The correct match is

- Erythrasma e)Coral red fluorescence
- Pityriasis versicolor c)Yellow fluorescence
- Tinea Capitis d)Blue fluorescence
- Vitiligo b)Milky white florescence

Solution to Question 6:

Non-scarring alopecia is caused by androgenic alopecia, alopecia areata, and telogen effluvium (1, 2 and 3). Frontal fibrosing alopecia is a type of primary cicatricial (scarring) alopecia.

Alopecia areata is a chronic inflammatory disease that causes non-scarring hair loss. It is a T-cell-mediated inflammatory disease. It is associated with several autoimmune diseases, including thyroiditis, lupus erythematosus, vitiligo, and psoriasis. It mainly affects the scalp, but can also affect the beard, eyebrows, and eyelashes. In some cases, this progresses to a total loss of scalp hair (alopecia totalis) or a loss of all hair on the body (alopecia universalis).

Telogen effluvium is a type of nonscarring alopecia that is marked by diffuse shedding of normal hair. This occurs due to premature termination of the anagen phase of the hair cycle. It can be either acute or chronic.

• Acute telogen effluvium occurs 2–3 months after a stressful event such as high fever, surgery, starvation, and hemorrhage. Hormonal fluctuations during the postpartum period can trigger it (telogen gravidarum). Telogen effluvium can result in a diffuse reduction in hair density but never total baldness. It generally resolves spontaneously.

• Chronic telogen effluvium is used to describe the diffuse shedding of telogen hair for more than 6 months. Common causes include thyroid disorders, iron-deficiency anemia, and malnutrition.

Androgenetic alopecia is a disorder characterized by a reduction of hair fiber production by follicles and miniaturization. Dihydrotestosterone, an androgenic metabolite, is implicated in the etiology. It is of two types: male-pattern and female-pattern hair loss. Male-pattern hair loss causes frontal hairline recession and thinning of the vertex. Female-pattern hair loss involves the mid-frontal scalp diffusely. There is no or minimal involvement of the temporal region.

Solution to Question 7:

The above clinical scenario and the image show an eschar at the bite site along with positive IgM of scrub typhus caused by Orientia tsutsugamushi and transmitted by trombiculid mite. As per a study published in 2023, Doxycycline and Azithromycin combined therapy was found to be superior to Doxycycline monotherapy.

Scrub Typhus is also known as chigger-borne typhus, which is caused by Orientia tsutsugamushi. The vector is trombiculid mite. Humans are infected when they are bitten by the mite larvae known as chiggers. The incubation period is 1- 3 weeks. The patient typically develops a

characteristic eschar at the site of the mite bite, with regional lymphadenopathy and maculopapular rash.

Diagnosis is by serologic assays.

Treatment is with a combination therapy of azithromycin and doxycycline.

Solution to Question 8:

Pityriasis rosea is the most likely diagnosis in this patient presenting with christmas tree lesions, and will likely have collarette scales.

The image given below shows collarette scales.

Pityriasis rosea/pityriasis circinata/roseola annulata is an acute self-limiting papulosquamous disorder. It is commonly seen in children and young adults.

Clinical features include:

• Prodrome of a sore throat, gastrointestinal disturbance, fever or arthralgia.

• Herald patch - Large and conspicuous eruptions on the thigh, upper arm, trunk or neck covered by fine scales

The image given below shows a herald patch.

• After 5-15 days, it turns into discrete oval lesions, dull pink in color and covered by fine dry silvery-grey scales.

• These secondary lesions occur as macules and papules that are elliptical or ovular, symmetric, and commonly involve the thorax, back, and abdomen.

• It is followed by a marginal collarette of scale with a central wrinkling and atrophy. The collarettes of the scales are attached peripherally. It is also called a hanging curtain sign.

• The long axes of the lesions characteristically follow the lines of skin tension parallel to the ribs in a Christmas tree pattern on the upper chest and back.

The images given below show Christmas tree pattern along the lines of skin tension on the back and the chest.

The rashes of pityriasis rosea usually resolve by 8 weeks.

Other options:

Lichen planus is an idiopathic inflammatory skin disease affecting the skin and mucous membranes, often with a chronic course with relapses and periods of remission.Patients present with pruritic, purple, polygonal, papules. The lesions have thin white lines on their surface known as Wickham's striae. They are commonly seen on the volar aspect of the wrists, the lumbar region, and around the ankles.The mucosal involvement is seen. The commonest mucosal sites are buccal mucosa and tongue, rarely it can also be found on the anus and genitalia.

A hypopigmented ring surrounding individual psoriatic lesions is known as the Woronoff ring.It is usually associated with treatment, most commonly UV radiation or topical corticosteroids. The pathogenesis of the Woronoff ring is not well understood, but may result from the inhibition of prostaglandin synthesis.

Solution to Question 9:

The condition shown in the above image is likely acanthosis nigricans - that is strongly associated with insulin resistance/hyperinsulinemia

Acanthosis nigricans is a darkening of the skin of the intertriginous areas. The hyperpigmentation is generally velvety with poorly defined borders and thickening of the skin.

It can be grossly classified into benign and malignant.

Benign acanthosis nigricans is associated with obesity or insulin resistance. Insulin resistance is the most accepted hypothesis used to explain the pathophysiology of the condition. Elevated circulating fatty acids are considered a major factor in developing insulin resistance.

Malignant acanthosis nigricans has a rapid onset and is characterized by symmetrical, hyperpigmented, rugose, velvety plaques. It affects the axillae and flexures, nape of the neck, and mucosal surfaces. If it occurs de novo and is associated with weight loss, malignancy should be suspected. Generalized pruritus and skin changes of tripe palms (acanthosis palmaris) are also associated with malignancy. Associated malignancies are generally GI or lung carcinomas. In malignancy, tumor cells are believed to produce transforming growth factor-alpha or cytokines that activate insulin-like growth factors.

Acanthosis nigricans can only be treated by treating the underlying disease. In addition, treatment specific to acanthosis nigricans is done for aesthetic reasons. Medications like keratolytics, topical vitamin D, and melatonin can be used. Procedures like laser, dermabrasion, and chemical peels are also available.

Other options:

Option B: Stria Albicans is a stretch mark that is white or pearly in color. It is most common in pregnant women and people who have gained or lost weight quickly. Stria Albicans is not associated with Cushing syndrome, which is a hormonal disorder caused by excess cortisol production and produces a strechy, purplish rash most commonly over the abdomen/flanks.

Option C: Xeroderma pigmentosum - malignancy is a rare genetic disorder that makes the skin extremely sensitive to ultraviolet radiation from the sun. People with xeroderma pigmentosum have a greatly increased risk of developing skin cancer.

Option D: Cafe au lait spots - McCune Albright syndrome is a genetic disorder that causes multiple light brown, oval-shaped birthmarks on the skin. It is also associated with other symptoms, such as precocious puberty and bone abnormalities - polyostotic fibrous dysplasias.

Solution to Question 10:

Non-scarring alopecia is caused by androgenic alopecia, alopecia areata, and telogen effluvium.

Alopecia areata

It is a chronic inflammatory disease that causes non-scarring hair loss. It is a T-cell-mediated inflammatory disease. It is associated with several autoimmune diseases, including thyroiditis,

lupus erythematosus, vitiligo, and psoriasis.

It mainly affects the scalp, but can also affect the beard, eyebrows, and eyelashes. In some cases, this progresses to a total loss of scalp hair (alopecia totalis) or a loss of all hair on the body (alopecia universalis).

Telogen effluvium

It is a type of nonscarring alopecia that is marked by diffuse shedding of normal hair. This occurs due to premature termination of the anagen phase of the hair cycle. It can be either acute or chronic.

Acute telogen effluvium occurs 2–3 months after a stressful event such as high fever, surgery, starvation, and hemorrhage. Hormonal fluctuations during the postpartum period can trigger it (telogen gravidarum). Telogen effluvium can result in a diffuse reduction in hair density but never total baldness. It generally resolves spontaneously.

Chronic telogen effluvium is used to describe the diffuse shedding of telogen hair for more than 6 months. Common causes include thyroid disorders, iron-deficiency anemia, and malnutrition.

Androgenetic alopecia

It is a disorder characterized by a reduction of hair fiber production by follicles and miniaturization. Dihydrotestosterone, an androgenic metabolite, is implicated in the etiology. It is of two types: male-pattern and female-pattern hair loss. Male-pattern hair loss causes frontal hairline recession and thinning of the vertex. Female-pattern hair loss involves the mid-frontal scalp diffusely. There is no or minimal involvement of the temporal region.

CICATRICIAL/SCARRING ALOPECIA: Frontal fibrosing alopecia is a primary cicatrical condition that affects post-menopausal women in particular and frequently involves the eyebrows.

Solution to Question 11:

The image above shows a central ulcer with raised borders and rolled-out edges (rodent ulcer) suggesting a diagnosis of basal cell carcinoma.

Basal cell carcinoma is a slow-growing malignant tumor with locally invasive characteristics. It affects the pilosebaceous skin.

Ultraviolet radiation exposure is the strongest predisposing factor. Other risk factors include exposure to arsenic, coal tar, aromatic hydrocarbons, and ionizing radiation.

Basal cell carcinoma mainly affects elderly or middle-aged men with excessive skin exposure, commonly in the age group of 40–80 y and in white-skinned people.

Macroscopically, it is categorized as localized and generalized. Nodular, nodulocystic (nodular and nodulocystic together account for 90% of basal cell carcinoma), pigmented, and nevoid are the localized cancers, while superficial multifocal and superficial spreading, infiltrative, and cicatrizing are the generalized types. The nodular type is described as pearly waxy nodules at the margins. A central depression (umbilication) or ulceration and rolled-out edges are characteristic of basal cell carcinoma.

Sites near the eye, nose, and ear and of size >2 cm are considered to be high risk. Direct invasion at these sites can reach the cranium.

Management is by Mohs microscopic surgery. Non-surgical measures, such as the administration of topical 5-fluorouracil and imiquimod, are useful. Metastasis is extremely rare.

Other Options:

Option A: Squamous cell carcinoma is the second most common malignant tumor, seen more commonly in men. Microscopically, the proliferation of irregular masses of squamous epithelium is seen. The image is shown below.



Option C- Lupus vulgaris originates from an underlying focus of tuberculosis, typically in a bone, joint, or lymph node, or maybe from the hematogenous and lymphatic spread. It presents as a slowly enlarging plaque with an elevated border and central atrophy. The edges of the lesion gradually extend in some areas and heal with scarring in others. In India, the buttocks, and trunk are more frequently involved. Apple jelly nodules are demonstrated on diascopy.

Option D: Nevus are aggregates of melanocytes at the dermis or dermo-epidermal junction. It can get transformed into malignant melanoma and the features suggestive of malignant transformation are a change in size, shape, color, thickness, satellite lesions, and tingling/ itching/serosanguinous discharge.

Solution to Question 12:

Direct immunofluorescence test, that is shown above, is the most accurate test to diagnose Pemphigus Vulgaris.

Direct Immunofluorescence uses anti-IgG antibodies which detects the anti-desmoglein IgG antibodies deposited on the Desmosomes. These desmosomes are present at cell junctions; this leads to the fishnet pattern.

Direct immunofluorescence (DIF) is a type of immunofluorescence that is used to diagnose autoimmune blistering diseases, such as pemphigus. In DIF, antibodies are applied directly to a skin biopsy specimen. If the patient has pemphigus, the antibodies will bind to the desmosomes, which are the proteins that hold skin cells together. This binding will produce a fish net pattern of

fluorescence under a microscope.

Other options:

Indirect immunofluorescence (IIF) is a type of immunofluorescence that is used to diagnose autoimmune diseases. In IIF, antibodies are first applied to a patient's serum, and then the serum is applied to a skin biopsy specimen. The presence of antibodies in the patient's serum will cause the skin biopsy specimen to fluoresce under a microscope. Seen below is a centromeric pattern on IF in Sjogren's disease -1 and Diffuse/homogenous staining in SLE - 2.

Immunohistochemistry is a type of microscopy technique used to visualize the presence of specific proteins/receptors in cells or tissues. It uses antibodies that are labeled with enzymes or fluorescent dyes to bind to specific antigens. An example is receptor identification in breast cancer specimens.

Fluorescent In Situ Hybridisation (FISH) is a simple technique for the detection of specific DNA sequences in cells/tissues. It uses fluorescent labeled probes to bind to complementary DNA FISH can be performed on prenatal samples, peripheral blood cells, touch preparations from cancer biopsies, and even fixed archival tissue sections. It is used to detect

• numeric abnormalities of chromosomes (aneuploidy)

• subtle microdeletions and rearrangements.

• complex translocations that are not demonstrable by routine karyotyping

• gene amplification (e.g., HER2 in breast cancer or NMYC amplification in neuroblastomas)

Solution to Question 13:

The image given shows the row of tombstone appearance seen in Pemphigus vulgaris.

Pemphigus vulgaris is an autoimmune blistering disease caused by the deposition of IgG antibodies against desmoglein 3, a protein that holds skin cells together. It is characterized by widespread flaccid bullae, which are large, fluid-filled blisters that rupture easily. The lesions typically start on the trunk and spread to the extremities. Oral lesions are common in pemphigus vulgaris. Histologically, pemphigus vulgaris is characterized by a suprabasal split with acantholysis (loss of cohesion between skin cells).

Acantholysis and cell apoptosis are the key pathological processes in pemphigus. Histological changes include:

• Intercellular oedema with loss of intercellular attachments in the basal layer is the earliest change

• Suprabasal epidermal cells separate from the basal cells to form clefts and blisters.

• Basal cells remain attached to the basement membrane, in a ‘row of tombstones’ on the floor of the blister.

• Blister cavities contain rounded■up acantholytic cells, also seen in Tzank smears.

IgA disease (Linear IgA bullous dermatosis): Autoimmune blistering disease caused by the deposition of IgA antibodies at the dermal-epidermal junction. It is characterized by vesicles and bullae, which are large, fluid-filled blisters. The lesions typically start on the trunk and spread to

the extremities. Oral lesions are not common in IgA disease. Histologically, IgA disease is characterized by a suprabasal split with linear IgA deposition at the dermal-epidermal junction. Anti-transglutaminase antibodies are positive in dermatitis herpetiformis. Patients present with erythematous papules and vesicles on the extensor aspect of elbows, knees, buttocks, and scalp. Histopathology demonstrates subepidermal blister formation with neutrophils located at the tips of the dermal papillae. Direct immunofluorescence will reveal the deposition of IgA in the papillary dermis in a granular or fibrillar pattern.

Pemphigus foliaceus

Pemphigus foliaceus is an autoimmune blistering disease caused by the deposition of IgG antibodies against desmoglein 1, a protein that holds skin cells together. It is characterized by superficial, flaccid bullae that are prone to rupture and erosion. The lesions typically start on the face and scalp and then spread to the trunk and extremities. Oral lesions are not common in pemphigus foliaceus. Histologically, pemphigus foliaceus is characterized by a subcorneal split with acantholysis and minimal inflammation.

The image given below shows the lesions seen in pemphigus foliaceous.

Type of pemphigus	Cutaneous distribution	Mucosal involvement	Lesions
Pemphigusvulgaris	- Scalp, face, chest, upper t runk- May be generalised	- Oro■pharynx, conjunctiv a, genital	- Flaccid blisters- Erosions - Flexural vegetations
Pemphigusvegetans	- Flexural aspects	- Oro■pharynx, conjunctiv a, genital	- Vegetating plaques- Vesic les- Pustules- Erosions
Pemphigusfoliaceous	- Scalp, face, chest, upper back, seborrheic areas- Ma y be generalised	- None	- Scaly papules- Crusted er osions- Erythroderma
Paraneoplasticpemp higus	- Upper body- Palmoplant ar	- Severe mucositis	- Polymorphous bullae- Er osions- Target lesions

Dermatology INI-CET 2024

Question 1:

Which of the following is not associated with annular lesions?

a) Tinea corporis

b) Pityriasis rosea

c) Urticaria

d) Porokeratosis

Question 2:

Match the following

a) a-3, b-4, c-2, d-1

b) a-1, b-4, c-2, d-3

c) a-3, b-2, c-4, d-1

d) a-2, b-4, c-3, d-1

Question 3:

A hypopigmented patch which grows initially and stops progressing after sometime is seen in which type of vitiligo?

a) Focal

b) Segmental

c) Mixed

d) Acrofacial

Question 4:

What is the mechanism responsible for the release of secretions from the gland seen in the image below?



a) Apocrine

b) Merocrine

c) Eccrine

d) Holocrine

Question 5:

All are a part of NAME syndrome, a subset of Carney's complex, except?

a) Atrial myxoma

b) Nevi

c) Ebstein anamoly

d) Myxoid neurofibroma

Question 6:

A 60-year-old woman presented with a great toe infection. She had significant discoloration of the toenail with a subungual hyperkeratosis that caused color alteration and a rough surface area. She was suspected to have onychomycosis. Which topical drug is given for toenail onychomycosis?

a) Caspofungin

b) Itraconazole

c) Griseofulvin

d) Tavaborole

Question 7:

A male patient presents with urethritis. It can be caused by all except?

a) Neisseria Gonorrhea

b) Gardenella vaginalis

c) Trichomonas vaginalis

d) Mycoplasma

Answer Key

Question No.	Correct Option
1	c
2	a
3	b
4	d
5	c
6	d
7	b

Detailed Explanations

Solution to Question 1:

Urticaria is not associated with annular lesions. It is characterized by wheals.

Urticaria is a heterogeneous group of disorders characterized by itchy wheals, which develop due to evanescent edema of dermis.

A weal is a descriptive term for transient, well■demarcated, superficial pink or pale swellings of the dermis due to reversible exudation of plasma in the skin that fade, usually within hours, without leaving a mark . Weals are usually very itchy and associated with a surrounding red flare when they arise. Lesions begin as erythematous macules, which rapidly evolve into pale pink edematous wheals with a surrounding flare.

Option A: Tinea corporis/ ringworm infection is caused by the
dermatophyte Trichophyton rubrum. It results from invasion and proliferation of the causal fungi

in the stratum corneum with lesions on the trunk and limbs. Characteristic lesions are circular, usually sharply marginated with a raised edge with central resolution.

Option B: Pityriasis Rosea (PR) is an unknown etiology, self-limiting condition that typically lasts between 2 to 10 weeks. It is characterized by a herald patch, followed by a secondary eruption with typical lesions that are oval, annular, erythematous, and plaque-like with a collarette of scales. These lesions commonly appear on the trunk in a "fir tree" pattern along lines of cleavage. Differential diagnoses include guttate psoriasis, secondary syphilis, and drug rash. There is no specific treatment for PR; symptomatic treatment is usually sufficient.

Option D: Porokeratosis is a clonal expansion of keratinocytes which differentiate abnormally. But they are not hyperproliferative. It may present as single or multiple lesions and may be localized or disseminated. Porokeratoses present with single or multiple papules or plaques which develop into annular lesions with a thin raised border.

Solution to Question 2:

The correctly matched option is a-3, b-4, c-2, d-1.

Lupus vulgaris is one of the prevalent forms of cutaneous tuberculosis that occurs in previously sensitized individuals with high immunity. It is characterized by reddish-brown papules and slow-growing plaques with elevated borders and central atrophy. Diagnosis is made by biopsy. Caseating granulomas are pathognomonic of tuberculosis.

Tinea corporis/ ringworm infection is caused by the dermatophyte Trichophyton rubrum. It results from invasion and proliferation of the causal fungi in the stratum corneum with lesions on the trunk and limbs. Characteristic lesions are circular, usually sharply marginated with a raised edge with central resolution.

KOH mount is used for the diagnosis. Branching hyphae are seen in Tinea corporis.

Leprosy is a chronic granulomatous condition caused by Mycobacterium leprae. It affects the skin, cutaneous and peripheral nerves, and mucous membranes.

Cardinal signs for the diagnosis of leprosy:

• Characteristic skin lesions- hypopigmented or erythematous lesions with definite impairment of sensation.

• Peripheral nerve involvement- thickening of the nerve with sensory impairment. Nerve tenderness can also be seen.

• Detection of acid-fast bacilli in slit-skin smear, skin biopsy, or positive biopsy PCR.

Allergic contact dermatitis is an eczematous reaction that occurs as an immunological response that occurs due to exposure to a substance towards which the immune system has previously been sensitized.

The patch test is used to detect delayed (type IV) hypersensitivity to antigens that come in contact with the skin, by applying the suspected antigen under occlusive patches to the skin. Patches are removed after 48 hours (or sooner if there is itching and burning).

• 48 hours: Patch removal for initial reading to detect delayed hypersensitivity reaction.

• 72-96 hours: Final reading done for confirmation.

An image showing the patch test:

a. Patch test	3. Contact dermatitis
b. KOH mount	4. Tinea corporis
c. Biopsy	2. Lupus vulgaris
d. Slit skin smear	1. Leprosy



Solution to Question 3:

In segmental vitiligo, lesions increase initially (6–12 months) and then remain static.

Vitiligo is a localized depigmentation that occurs due to a progressive loss in the melanocytes in the basal layer of skin. It is classified as

• Segmental vitiligo- which appears as unilateral maculae in a segmented or band-shaped pattern

• Non-segmental vitiligo- which appears as bilateral maculae distributed in an acrofacial pattern or scattered symmetrically over the entire body. Non-segmental can either be focal vitiligo or undetermined/unclassified vitiligo.

Acrofacial vitiligo is a type of generalized vitiligo, predominantly seen periorificially (around eyes and on lips on the face) and acral parts (periungual area, palms, and soles). This type of vitiligo is more resistant to therapy due to absence of hair in the affected parts.

Mixed vitiligo: the coexistence of non■segmental and segmental vitiligo in one patient is called mixed vitiligo

According to a recent Vitiligo Global Issue Consensus Conference, the term ‘vitiligo’ can be used as an umbrella term for all non■segmental forms of vitiligo (including several variants: acrofacial,

mucosal, generalized, universal, mixed, and rare variants of vitiligo). The classification is shown below.

Clinical features:

The lesions become apparent between 20-30 years of age.

Hypopigmented macules in vitiligo are commonly found in the hips, dorsum of the hands/fingers, feet, elbows, knees, and ankle areas. These are the areas of repeated friction. Koebner phenomenon is seen, which is the development of lesions at the site of trauma on the uninvolved skin.

Vitiligo universalis, which is complete vitiligo, or trichrome vitiligo where there are areas of depigmentation, hyperpigmentation, as well as areas of normal skin may be seen rarely.

Several autoimmune conditions can be associated with vitiligo. They are:

• Thyroid disease (hyperthyroidism and hypothyroidism)

• Pernicious anemia

• Addison disease

• Diabetes

• Hypoparathyroidism

Examination:

Under the Wood's lamp examination the depigmented areas emit a bright blue-white fluorescence and appear sharply demarcated.

Treatment:

Treatment involves cosmetic camouflage and the use of sunscreen. The topical application of corticosteroids is the first line of treatment. Topical calcineurin inhibitors like pimecrolimus, tacrolimus have also shown to be successful. Psoralen photochemotherapy, khellin, selective ultraviolet B therapy are used as second-line treatment. In patients who do not respond to this treatment, surgical grafting techniques can be used, where the melanocytes from a normal pigmented area are transferred to a non pigmented area. In patients with extensive vitiligo where only a few pigmented areas are left, depigmentation can be done.

Vitiligo (Non segmental)	AcrofacialMucosal (>1 one mucosal site)GeneralizedUniversalMixed (associated with SV)Rare variants
Segmental vitiligo	Uni-, bi-, or pleuri-segmental
Undetermined/unclassified vitiligo	FocalMucosal (one site in isolation)

Solution to Question 4:

The gland depicted in the image is a sebaceous gland, which releases its secretions through the holocrine mechanism.

Identification features of sebaceous gland:

• Lobular structure with clusters of cells.

• Foamy cytoplasm due to lipid content.

• Holocrine secretion with disintegrated cells.

• Found near hair follicles in the dermis.

Sebaceous gland cells synthesize and accumulate lipid-rich sebum within their cytoplasm.

Holocrine secretions are produced in the cytoplasm of the cell and released by the rupture of the plasma membrane, which destroys the cell and results in the secretion of the product into the lumen.

Examples of holocrine glands are

• Sebaceous glands of the skin and

• Meibomian glands of the eyelid

Other options:

Option A: Apocrine: involves partial loss of the apical portion of the cell (e.g., mammary glands).

Option B: Merocrine: involves secretion via exocytosis (e.g. sweat glands).

Option C: Eccrine glands are a type of sweat gland responsible for thermoregulation through the secretion of sweat. They use the merocrine mechanism for secretion.

Solution to Question 5:

Ebstein anomaly is not a part of NAME syndrome.

Carney Complex is a rare, autosomal dominant genetic disorder characterized by multiple neoplasms and lentigines. It includes features like cardiac myxomas, endocrine tumors, spotty skin pigmentation, and other benign or malignant tumors, caused by mutations in the PRKAR1A gene. It includes:

• Myxomas - cardiac, skin, and/or breast

• Lentigines and/or pigmented nevi

• Endocrine overactivity - primary nodular adrenal cortical disease, testicular tumors, and/or pituitary adenomas with gigantism or acromegaly

NAME syndrome and LAMB syndrome are synonymous with the Carney complex:

• LAMB - Lentigines, Atrial Myxoma, and Blue nevi.

• NAME syndrome is a subset of Carney's complex, which is a rare autosomal dominant disorder characterized by multiple neoplasms and pigmentation abnormalities. The acronym NAME stands for:

• Nevi

• Atrial myxomas

• Myxoid neurofibromas

• Ephelides (freckle-like pigmented lesions)



Solution to Question 6:

Tavaborole is the topical drug given in toenail onychomycosis.

Tavaborole is a topical antifungal. It is an oxaborole antifungal that inhibits fungal protein synthesis by targeting leucyl-tRNA synthetase.

Onychomycosis is defined as the infection of the nail plate by fungus.

Most common cause is Trichophyton rubrum, but many fungi may be causative.

The clinical scenario of Tinea unguium presents with yellowish discolouration, which spreads proximally as a streak in the nail. Later, subungual hyperkeratosis becomes prominent and spreads until the entire nail is affected. Gradually, the entire nail becomes brittle and separated from its bed.

Diagnosis is made by KOH mount examination of dystrophic subungual debris.

Management of onychomycosis involves Oral terbinafine as the treatment of choice for Tinea unguium or onychomycosis. It acts as a non-competitive inhibitor of squalene epoxidase, which is essential in the synthesis of fungal ergosterol. Oral terbinafine at 250 mg/day is given for 6 weeks for fingernail infection and 3 months (12 weeks) for toenail infection. An oral Griseofulvin dose of 1 g/day for 4–8 months (longer for toenails) is used as a second-line treatment.

Other Options:

Option A: Caspofungin: Caspofungin is an echinocandin antifungal that works by inhibiting fungal cell wall synthesis. It is used in intravenous form for systemic fungal infections like candidiasis and aspergillosis.

Option B: Itraconazole: Itraconazole is an oral antifungal from the triazole class, used for onychomycosis due to its penetration into keratinised tissues.

Option C: Griseofulvin: Griseofulvin is an oral antifungal used for dermatophyte infections, including onychomycosis.

Solution to Question 7:

Gardenella vaginalis does not cause urethritis in male patients.

Urethritis is classified into two types:

• Gonococcal urethritis - caused by Neisseria gonorrhoeae(Option A)

• Non-gonococcal urethritis - caused by:

• Chlamydia trachomatis (most common)

• Ureaplasma urealyticum

• Trichomonas vaginalis (rare) (Option C)

• Herpes simplex virus (rare)

• Mycoplasma genitalium (Option D)

The most likely causative organism for acute urethritis in males is Neisseria gonorrhoeae.

Dermatology NEET 2024

Question 1:

A child presents with orangish hue skin with rough and scaly skin as shown in the image. On histopathological examination, orthokeratosis and parakeratosis are seen. What is the most likely diagnosis?

a) Follicular psoriasis

b) Lichen planus

c) Pityriasis rubra pilaris.

d) Keratosis follicularis

Question 2:

A patient presented with a slight loss of sensation along the ulnar nerve and a solitary nodule in the forearm. Intradermal antigen was injected. The histopathology image is shown below. Which of the following is true regarding this condition?



a) Lepromatous leprosy with dermal test positive

b) Tuberculoid leprosy with dermal test positive

c) Lepromatous leprosy with dermal test negative

d) Erythema nodosum leprosum with negative skin test

Question 3:

Which of the following is associated with the" box sign"?



a) Lichen planus

b) Lichen nitidus

c) Morphea

d) Amyloidosis

Question 4:

A 32-year-old man presents with painful vesicular lesions over the genital area. He has a history of multiple sexual partners and inconsistent condom use. What is the most likely diagnosis?

a) Chancroid

b) Genital Herpes

c) Primary syphillis

d) Granuloma inguinale

Question 5:

An adult male presented with asymptomatic macules over the chest, moist grey, perianal lesions, and findings on the scalp as shown. What is the most likely causative organism of this condition?

a) Malassezia

b) Trichophyton

c) Treponema pallidum

d) Microsporum

Question 6:

Match the structures shown in the image:



a) A- Apocrine sweat gland; B- Eccrine sweat gland; C- Sebaceous gland; D- Erector pili muscle

b) A- Eccrine sweat gland; B- Apocrine sweat gland; C- Sebaceous gland; D- Erector pili muscle

c) A- Sebaceous gland; B- Apocrine sweat gland; C- Eccrine sweat gland; D- Erector pili muscle

d) A- Eccrine sweat gland; B- Sebaceous gland; C- Apocrine sweat gland; D- Erector pili muscle

Question 7:

Which of the following causes the condition shown in the image below?



a) Candida albicans

b) Trichophyton species

c) Epidermophyton floccosum

d) Microsporum species

Question 8:

A 65-year-old treated leprosy patient came with complaints of a well-demarcated, non-tender ulcer on his foot with hyperkeratotic borders as given below for the last 3 months. What should be the appropriate next step?

a) Restart MDT

b) Rest the limb, non weight bearing splints and foot care

c) Admit the patient and aggressive debridement and antibiotics

d) Amputation

Question 9:

A man presents with multiple lesions on his face and upper back, diagnosed as acne vulgaris. Which of the following is the most effective treatment for his symptoms?

a) Systemic Minocycline

b) Systemic Isoretinoin

c) Systemic Doxycycline

d) Topical Dapsone

Question 10:

A 42-year-old man comes with vesicular eruptions in the buccal mucosa. Direct immunofluorescence is done and the image is shown below. What is the most likely diagnosis?



a) Lupus Vulgaris

b) Pemphigus vulgaris

c) Bullous Pemphigoid

d) Borderline Leprosy

Question 11:

A 3-year-old boy had patches of partial hair loss with mild itching and scales. The picture of his scalp is shown below. His sister also has a similar condition. Which of the following is the most likely diagnosis?

a) Alopecia areata

b) Tinea capitis

c) Pediculis Capitis

d) Pyoderma Scalp

Question 12:

A middle aged man comes with complaints of painless progressively enlarging lesions on his scalp and perianal region for the last 3 months. He has had liver and kidney transplant 2 years back and is on Mycophenolate Mofetil, Tacrolimus and Prednisolone for his immunosupression. Which of the following is the most likely diagnosis?



a) Drug induced eruption

b) Malakoplakia

c) Cutaneous TB

d) Pyoderma Gangrenosum

Answer Key

Question No.	Correct Option
1	c
2	b
3	c
4	b
5	c
6	b
7	b
8	b
9	b
10	b
11	b
12	b

Detailed Explanations

Solution to Question 1:

The image shows prominent erythema and scale on the palms of the hands and wrists with marked orange■yellow palmoplantar keratoderma which is a feature of pityriasis rubra pilaris.

Features of pityriasis rubra pilaris:

• Salmon red or orange red-coloured dry scaly plaques are seen characteristically, which coalesce together, leaving islands of normal skin, typically known as 'islands of sparing' in between the lesions

• Most common site - trunk

• On the elbow and wrist, nutmeg grater papules are seen due to follicular hyperkeratosis

• Palmoplantar keratoderma - keratoderma on the sole is very thick that it appears as a sandal. (keratotic sandal)

• Nails - thickened, discoloured distally, showing splinter haemorrhages

Histologically, pityriasis rubra pilaris is characterized by hyperkeratosis with follicular plugging, alternating ortho- and parakeratosis, dilated but non-tortuous dermal capillaries, sparse lymphocytic infiltrate, and possible acantholysis in the adnexal epithelium.

The image below shows islands of sparing in pityriasis rubra pilaris.

The image below shows keratotic sandal.

Other options:

Option A: Follicular psoriasis affects hair follicles on the trunk and limbs and can occur alone or alongside plaque psoriasis. The lesions are smaller than those seen in guttate psoriasis and may appear clustered or spread out. In children, it can be mistaken for pityriasis rubra pilaris.

Option B: Lichen planus is a papulosquamous lesion presenting as pruritic, purple, polygonal, planar, papules and plaques. These papules are often highlighted by white dots or lines called

Wickham striae, which are created by areas of hypergranulosis.

Option D: Keratosis follicularis or Darier's disease or dyskeratosis follicularis, is an autosomal dominant condition arising from mutations in the ATP2A2 gene which affects the SERCA2 calcium pump. It presents with greasy hyperkeratotic papules and plaques in seborrheic regions. Neuropsychiatric features such as depression, psychosis and mental retardation may also occur.

Solution to Question 2:

The patient's slight loss of sensation along the ulnar nerve, solitary forearm nodule, and histopathology showing well-defined granulomas indicate tuberculoid leprosy with a dermal test positive.

In tuberculoid leprosy, dermal granulomas, consisting of groups of epithelioid cells with giant cells, are seen. The granulomas are elongated and generally run parallel to the surface, following neurovascular bundles. The granulomas extend up to the epidermis, thereby obscuring the Grenz zone. It also shows high positivity of the lepromin skin (dermal) test.

Leprosy is a chronic granulomatous condition caused by Mycobacterium leprae. It affects the skin, cutaneous and peripheral nerves, and mucous membranes. The most commonly affected peripheral nerves are the posterior tibial nerve, followed by the ulnar, median, and lateral popliteal nerves. The most frequently involved cranial nerve is the facial nerve. It has a long incubation period, usually 2 to 12 years. It commonly presents as an area of numbness on the skin or a visible skin lesion, which could be a hypopigmented or erythematous macule.

The Grenz zone is most prominent in lepromatous leprosy, as there are scarce granulomas to fill up the space. It is also seen in the borderline spectrum.

Cardinal signs for the diagnosis of leprosy:

• Characteristic skin lesions- hypopigmented or erythematous lesions with definite impairment of sensation.

• Peripheral nerve involvement- thickening of the nerve with sensory impairment. Nerve tenderness can also be seen.

• Detection of acid-fast bacilli in slit-skin smear, skin biopsy, or positive biopsy PCR.

The presence of two of the three signs can establish the diagnosis of leprosy. Other tests done for diagnosis of leprosy include nerve biopsy in pure neural leprosy and PGL-1 antibody in cases with low bacterial load.

Solution to Question 3:

The "box sign" is associated with morphea, also known as localized scleroderma.

Morphea is a condition characterized by localized skin sclerosis and fibrosis due to increased collagen deposition. This can lead to well-defined, box-like patches of thickened, hardened skin. The "box sign" refers to the sharp, boxy edges of these lesions, which can vary in depth and extent, sometimes involving deeper tissues such as muscle and bone.

Biopsies taken during the indurative stage often display squared-off edges, resulting in a "box-shaped" or "boxed dermis" appearance under the microscope, which can vary in depth and extent, sometimes involving deeper tissues such as muscle and bone. This characteristic sign occurs because of extensive collagen sclerosis throughout the reticular dermis and extending into the subcutaneous fat septa. The pronounced sclerosis of the collagen fibers causes this distinct boxy appearance. Similar squared-off edges may also be observed in morphea-lichen sclerosus overlap and scleredema, due to comparable changes in the dermal and subcutaneous tissue structure.

The epidermis in morphea can exhibit varying changes, from normal to flattened with loss of rete ridges, or slightly acanthotic. In the inflammatory phase, the reticular dermis shows edema and a dense perivascular infiltrate, which gradually gives way to thickened, horizontally oriented collagen bundles. The loss of adnexal structures, such as hair follicles and eccrine glands, is common, and the subcutaneous fat may be replaced by collagen, contributing to the appearance of a "box sign."

Management of morphea focuses on controlling inflammation and preventing disease progression. Early intervention is crucial and may include topical treatments like corticosteroids or tacrolimus for localized lesions. For more extensive or resistant cases, phototherapy and systemic treatments, such as methotrexate or oral corticosteroids, may be considered. The choice of therapy often depends on the extent and activity of the disease, with the goal of minimizing skin damage and improving functional outcomes.

Other options:

Option A: Lichen planus is a papulosquamous lesion presenting as pruritic, purple, polygonal, planar, papules and plaques. These papules are often highlighted by white dots or lines called Wickham striae, which are created by areas of hypergranulosis.

The characteristic lesions in lichen planus is shown below:

Option B: Lichen nitidus is a rare variant of lichen planus, where both are histologically similar in early lesions. It is mostly seen in children and young adults. Lesions of lichen nitidus are typically described as a minute pinpoint to pinhead-size, flat or dome-shaped papules with a shiny surface. They are most commonly found over the forearm, dorsum of the penis, abdomen, chest, and buttocks.

Option D: Amyloidosis refers to extracellular deposits of fibrillar proteins. The kidney is the most common organ involved in amyloidosis. Abnormal fibrils are produced by the aggregation of misfolded proteins. The fibrillar deposits bind a wide variety of proteoglycans and glycosaminoglycans, including heparan sulfate and dermatan sulfate, and plasma proteins, notably serum amyloid P component (SAP). Amyloid appears as an amorphous, eosinophilic, hyaline, extracellular substance.

Solution to Question 4:

The most likely diagnosis for this patient with painful vesicular lesions over the genital area is genital herpes.

The diagnosis of genital herpes is suggested by the presence of painful, multiple vesicular ulcers clustered on the genitalia, along with inguinal lymphadenopathy. These symptoms are indicative of an infection caused by the Herpes Simplex Virus-2 (HSV-2), which is the primary cause of genital herpes.

Genital herpes is characterized by vesiculoulcerative lesions, which appear as grouped vesicles filled with clear fluid on the penis in males or on the cervix, vulva, vagina, and perineum in females. The painful nature of these lesions, combined with the history of multiple sexual partners and inconsistent condom use, supports the diagnosis. Recurrences of genital herpes are common and generally tend to be milder compared to the initial outbreak.

Other options:

Option A: Chancroid caused by Haemophilus ducreyi may also present with soft wide indurated painful genital ulcers and painful genital lymphadenopathy.

Option C: In primary syphilis, the typical primary chancre usually begins as a single painless papule that rapidly becomes eroded and usually becomes indurated, with a characteristic cartilaginous consistency on palpation of the edge and base of the ulcer.

Option D: Granuloma inguinale (also known as donovanosis) is caused by Klebsiella granulomatis. It presents with a painless papule or nodule that erodes into a beefy red granulomatous ulcer with rolled edges and often no lymphadenopathy.

Solution to Question 5:

The most likely causative organism for this condition is Treponema pallidum, which caused the asymptomatic macules on the chest, moist grey, perianal lesions, and moth-eaten pattern alopecia seen in secondary syphilis.

The causative organism for this condition is Treponema pallidum, a spirochete bacterium. In secondary syphilis, the macular or roseolar rash is often the earliest sign, which may fade, leaving hypopigmented areas around the neck, known as leucoderma syphiliticum or "necklace of Venus." This stage also features a papular rash that can appear on the palms and soles.

Additionally, secondary syphilis can present with nummular lesions that resemble psoriasis, often with easily removable scales due to serous discharge. On the oral mucosa, papules may form grey oval patches, leading to "snail-track" ulcers (as shown below).

Condyloma lata, which are moist, well-demarcated papules and plaques, are commonly seen in the genital area and are highly infectious. Syphilitic alopecia, characterized by non-scarring patchy hair loss, and generalized lymphadenopathy are also notable features.

The systemic manifestations of secondary syphilis include panuveitis, periostitis, joint effusions, and involvement of various organs such as the kidneys, liver, and heart. The lesions in secondary syphilis typically resolve spontaneously, transitioning into the latent stage of the disease.

Solution to Question 6:

The correctly matched option (B)- A- Eccrine sweat gland; B- Apocrine sweat gland; C- Sebaceous gland; D- Erector pili muscle.



Human skin contains various structures, including sweat and sebaceous glands. Eccrine sweat glands, which are crucial for temperature regulation, are distributed across most of the body and are identified early in fetal development. These glands have a coiled secretory portion in the lower dermis and a duct that opens directly onto the skin surface. They are most concentrated on the palms, soles, and forehead.

Apocrine sweat glands, found in areas like the axillae, genital, and mammary regions, differ from eccrine glands in that they release secretion via decapitation, where parts of the cell pinch off. They are connected to hair follicles and have a larger lumen compared to eccrine glands. Their primary function remains unclear, and they have a lower secretory output.

Sebaceous glands secrete sebum, providing skin lubrication and waterproofing. These glands are associated with hair follicles and are absent in non-hairy skin.

The erector pili muscle, a smooth muscle bundle, extends from the dermis to the follicle wall and is involved in hair erection.

Solution to Question 7:

The above image shows numerous fungal spores within the hair shaft suggestive of endothrix type of tinea capitis, which is caused by Trichophyton tonsurans.

Tinea capitis is of four types:

Other options:

Option A: Candida is the most common cause of opportunistic mycoses in the world. Immunocompromised individuals are at an increased risk of systemic candidiasis. Clinically, infection with candida can present as cutaneous & mucosal candidiasis, systemic candidiasis, or rarely, as chronic mucocutaneous candidiasis.

Morphologically, they are yeast-like fungi. It appears as budding gram-positive yeast cells with pseudohyphae

Option C: Epidermophyton floccosum: Distinguished by absent microconidia and club-shaped macroconidia, which are smooth, thick-walled, and occur in clusters. It is a dermatophyte responsible for superficial fungal infections of the skin, particularly tinea pedis (athlete's foot), tinea cruris (jock itch), and tinea unguium (nail infection).

Option D: Tinea cruris, a superficial fungal skin infection that affects the groin area, is primarily caused by Epidermophyton floccusum. Typical symptoms include redness and inflammation in the groin region, accompanied by sensations of itching, burning, or stinging. The characteristic lesions exhibit raised borders surrounding a central clearing, thus imparting a ring-like appearance.

Type of Infection	Causative fungus
Endothrix (infection inside s haft)	Trichophyton tonsuransT.violaceumT. schoenleiniiT.yaoundeiT. soudanense
Ectothrix(infection outside sh aft)	MicrosporumaudouiniiM.canisM.equinumM.ferrugineum

Type of Infection	Causative fungus
Favus	T. schoenleinii
Kerion	T. verrucosumT. mentagrophytes

Solution to Question 8:

The given scenario and the image indicate a neuropathic ulcer. This can be managed by limb rest, utilising non-weight-bearing splints, and administering foot care.

Leprosy, a condition affecting the skin and peripheral nerves, results in peripheral neuropathy and subsequent loss of sensation. This can lead to the development of chronic trophic or neuropathic ulcers characterised by punched-out appearance, regular margins, and calluses due to repeated trauma in the presence of anaesthesia of feet.

Management primarily focuses on relieving pressure through the application of non-weight bearing splints for even pressure distribution across the entire foot, limb elevation to mitigate swelling, and regular foot inspection to promptly identify trauma, in addition to the use of microcellular rubber footwear.

Other options:

Option A: Trophic ulcers are not an indication for restarting MDT in an already treated leprosy patient.

Option C: Due to the absence of debris or signs of infection in the ulcer, aggressive debridement and antibiotics are not indicated.

Option D: As there are no signs of vascular compromise or gangrene, amputation of the limb is not required.

Solution to Question 9:

The presence of multiple nodular and cystic lesions over the face and upper back implies that this is a case of acne vulgaris. Systemic Isotretinoin is the most effective treatment for acne.

Systemic retinoids are a very efficient treatment for acne vulgaris. Their mechanism of action si by inhibiting the formation of keratin, a protein that can clog pores and cause acne. Oral retinoids are usually reserved for severe cases of acne that do not respond to topical creams and gels.

Acne is a chronic inflammatory disease of the pilosebaceous unit. Comedonal lesions are the earliest lesions of acne. They represent the keratin plugs of the pilosebaceous duct.

When the keratin plug is inside the follicle, it is called a closed comedone (whitehead). When the keratin plug is exposed to the environment, it gets oxidized by the air and turns black which is known as open comedone (blackhead).

Solution to Question 10:

The given image shows a characteristic fish-net appearance on direct immunofluorescence, which is diagnostic of pemphigus vulgaris.

This pattern is due to intraepidermal intercellular deposition of IgG and C3.

Direct or indirect immunofluorescence cannot be used to differentiate between pemphigus vulgaris and pemphigus foliaceus, as they present similarly.

Other options:

Option A: Lupus vulgaris(Cutaneous TB) is the most common form of cutaneous tuberculosis in adults. The characteristic lesion is a plaque, composed of soft, reddish brown papules, the appearance on the diascopy being said to resemble apple jelly. The image below shows the characteristic lesion:

Option C: Bullous pemphigoid present linear or shoreline pattern is seen due to IgG and C3 deposits along the dermo-epidermal junctions, as seen in the following image.



Option D: In borderline leprosy, inverted-saucer-shaped lesions are seen and are characterized by a dome-shaped appearance with central infiltration and a gradually sloping peripheral edge.



Solution to Question 11:

Patchy hair loss and multiple black dots within the areas of alopecia along with fine scaling and itching of the scalp point towards the diagnosis of tinea capitis.

Tinea capitis is a dermatophyte infection primarily occurring in children. Clinical features of tinea capitis are patchy hair loss, scaling, tender lymphadenopathy, and in severe cases, there might be pustules and crusting of the affected area along with sinus formation.

The clinical variants of tinea capitis are:

Endothrix - a non-inflammatory type in which multiple black dots are present within the areas of alopecia. It is most commonly caused by T. tonsurans and T. violaceum.

Ectothrix - a mild inflammatory type in which single or multiple scaly patches with hair loss are seen. Microsporum audouinii, M.canis, M.equinnum, and M.ferrugineum cause ectothrix infection.

Kerion - a severe inflammatory type that presents with an inflammatory mass with thick crusting and matting of hair with pus and sinus formation. It is commonly caused by T. verrucosum and T. mentagrophytes.

Favus - is characterized by the presence of yellowish, cup■shaped crusts known as scutula that develop around hair follicles. It is caused by T. schoenleinii.

Other options:

Option A: Alopecia areata is a chronic inflammatory disease causing non-scarring hair loss. It is characterized by a circumscribed, smooth patch of hair loss. Exclamation mark hairs are often seen at the margins of the bald patches during active phases of the disease.

The image below shows alopecia areata.

Option C : Pediculis Capitis (head louse infestation) is very common in children. The majority of the transmission is by head-to-head contact, which is frequently observed with overcrowding (as in schools), poverty, and poor hygiene.

Symptoms include severe itching of the scalp. On examinations, nits are seen, which are firmly attached to hair shafts on which they can be freely glided.

Option D : Pyoderma of the scalp is an idiopathic cutaneous disorder. It is characterized by progressive painful ulceration which can cause destruction of underlying structures. It is a benign lesion, which typically develops after truama to that area.

Solution to Question 12:

Clinical scenario of painless progressive nodular, ulcerated lesions in an immuno-compromised patient with the clinical image showing yellow-erythematous-purple plaque and histopathology image showing Michaelis–Gutmann bodies are consistent with the diagnosis of cutaneous malakoplakia.

Malakoplakia is a rare inflammatory condition that typically occurs in the urinary tract. The cutaneous form of malakoplakia is less prevalent and it most commonly occurs in the perianal or genital regions. . It is caused due to acquired defects in phagocyte function, chronic bacterial infection by Escherichia coli or Proteus species, and immunosuppression (renal transplant patients).

The term malakoplakia means soft plaques. Grossly, malakoplakia appears as soft, yellow, slightly raised mucosal plaques.

Microscopically, they are composed of large, foamy macrophages mixed with occasional multinucleate giant cells and lymphocytes. The macrophages in malakoplakia have abundant granular cytoplasm due to phagosomes stuffed with undigested bacterial products and intracellular deposition of calcium and iron within enlarged lysosomes form laminated basophilic concretions. These are known as Michaelis–Gutmann bodies, pathognomonic of malakoplakia (shown below).



The image given below shows the Von Kossa stain demonstrating calcium and iron-containing Michaelis-Gutman bodies (MG bodies) in malakoplakia.



Other options :

Option A : Drug induced eruption can mimic a wide range of dermatoses. They can develop due to traditional pharmaceutical agents, biologic agents and targeted immunotherapeutic agents, food additives, and dyes. The morphologies of dryg induced eruptions can be morbilliform, urticarial, papulosquamous, pustular, and bullous.

Option C : Cutaneous TB (Lupus vulgaris) is the most common form of cutaneous tuberculosis in adults. The characteristic lesion is a plaque, composed of soft, reddish brown papules, the appearance on diascopy being said to resemble apple jelly.

Option D : Pyoderma gangrenosum is a non■infectious neutrophilic dermatosis commonly associated with inflammatory bowel disease. Central necrosis and ulceration of the epidermis and dermis surrounded by a strong inflammatory cell infiltrate are typical findings.

Psychiatry AIIMS 2017

Question 1:

A 42-year-old man had a fall followed by one episode of vomiting. When brought to the casualty, he was awake and conscious but had unrelieved anterograde amnesia. Which of the following is the most likely cause?

a) Drunkenness

b) Diffuse axonal injury

c) Concussion

d) Cerebral venous thrombosis

Question 2:

The marked area given in the sleep polysomnography is of _____



a) REM sleep

b) NREM stage 1

c) NREM stage 2

d) NREM stage 3

Question 3:

Which of the following is the test for immediate memory?

a) Digit span forward up to 7 digits with 2 skips allowed

b) Serial (100-7) subtraction test up to 5 steps

c) Digit span backwards up to 5 digits with 2 skips allowed

d) Serial (20-1) subtraction test up to 5 steps

Question 4:

A patient believes he is the most important person in the world and so his neighbours and family members are trying to harm him as they are jealous of him. His wife says otherwise and says he behaves like this only recently and before that he was working peacefully as a school teacher. He is most likely suffering from?

a) Delusion of grandiosity

b) Delusion of persecution

c) Delusion of grandiosity and persecution

d) Delusion of grandiosity, persecution and reference

Question 5:

A young patient is admitted to hospital with acute psychosis. He wakes up and asks for his wife even while she is there in the same room right next to him. Upon pointing her out, he states that this is not his wife and someone else is masquerading as his wife. The most likely diagnosis is?

a) Capgras syndrome

b) Fregoli syndrome

c) Delusion of subjective doubles

d) Othello syndrome

Question 6:

A young male patient is on 5 mg haloperidol for many days. However, recently for the last 4 days, he has an inner restlessness and urges to move. The most likely diagnosis is:

a) Akathisia

b) Tardive dyskinesia

c) Rabbit syndrome

d) Acute dystonia

Question 7:

An IT employee who has recently taken up the job is feeling guilty, hopeless and not able to concentrate on work. His symptoms started 3 years back when he started college. The most likely diagnosis is:

a) Depressive disorder

b) Dysthymia

c) Adjustment disorder

d) Cyclothymia

Question 8:

Mutism and akinesis in a person who is aware of his surroundings and somewhat alert is best described as a feature of:

a) Oneroid state

b) Twilight state

c) Stupor

d) Delirium

Question 9:

A 25 year old female has thoughts of cutting her fingers, she plans and imagines doing it but never actually does it. She says she is not having any guilt of having such thought. And also says the thoughts are distressing her and she is unable to control them. The thoughts vanish either by ending with a seizure or automatically subsides on its own. Which of the following is the cause?

a) Thought insertion

b) Forced thinking

c) Crowding of thoughts

d) Obsession

Answer Key

Question No.	Correct Option
1	c
2	a
3	a
4	c
5	a
6	a
7	b
8	c
9	b

Detailed Explanations

Solution to Question 1:

The given history of fall and anterograde amnesia points towards the diagnosis of concussion.

A concussion is the most common type of traumatic brain injury. It can result from vehicle accidents, falls, sports injuries, etc. Anterograde amnesia/post-traumatic amnesia is the hallmark of a concussion. In this type of amnesia, the events following the injury cannot be recalled.

Other clinical features of concussion include headache, dizziness, nausea, and vomiting.

Option A: Drunkenness - the patient will be drowsy or sleepy and not awake.

Option B: Diffuse axonal injury - seen in high-speed deceleration which causes disruption and shearing of axons leading to microhemorrhages. Usually results in a persistent vegetative stage.

Option D: Cerebral venous thrombosis - acute thrombosis in dural venous sinuses; manifests as headache, visual abnormalities, or seizures.

Solution to Question 2:

The marked area in the given polysomnography represents REM sleep.

The three zones in the given image represent EOG, EEG and EMG, respectively.



The following table helps identify the state of wakefulness from the graph :

In the marked area,

• EOG shows Positive activity

• EEG shows Mixed waves

• EMG shows Minimal activity

This is characteristic of REM sleep.

	EEG	EOG	EMG
Active state	Fast	+	+
REM sleep	Mixed	+	-
NREM sleep	Slow	-	+



Solution to Question 3:

Forward digit span test is the best test among the given options.

Memory is of 3 types:

• Immediate

• Recent

• Remote

Immediate memory is attention dependent. Tests of attention and memory include

• Digit span test: Most important and best test for immediate memory

• Serial (100-7) subtraction test is the best for concentration

• Days of the week backwards

• Spell WORLD backwards

• (20–3) test

Digit span test

The examiner recites numbers randomly without any sequence and asks the patient to repeat the same. The examiner starts with a 2-number sequence and the digit span increases from 2 numbers to 3 numbers and so on. The patient is asked to repeat until he/she fails to repeat all numbers correctly.

Digit forward

• Normally a person can repeat up to 7 digits (+/- 2)

• E.g., Examiner says 2 - 4 –8 –1 –5 –7 - 9. The person has to repeat in the same order 2 – 4 – 8 –1 –5 –7 –9.

Digit backward

• Reproduction of longer digits in backward direction.

• Normally a person can repeat up to 5 digits (+/- 1) in backward direction (making option C wrong)

• Eg : Examiner says 2 - 4 –8 –1 –5 –7 - 9. The person has to repeat it backwards. 9 -7 -5 -1 -8- 4- 2.

Serial (100-7) subtraction test

• 5 scores / steps are taken in to account

• 1st score – When the patient tells 93 (100 - 7 = 93)

• 2nd score - When the patient tells 86 (93 – 7 = 86)

• And so on . . 100 , 93 , 86 , 79 , 72 , 65

Hence among the given options :

Digit span forward up to 7 digits with 2 skips allowed would be the answer of choice as it is the best test for immediate memory

Solution to Question 4:

• Belief that he is the most important person in the world – Delusion of grandeur

• Belief that his neighbours and family members are trying to harm him – Delusion of persecution

• Belief that others feel jealous – Part of delusion of persecution

A scenario of this kind is most likely seen in manic patients with

• Primary delusion of Grandeur (and)

• Secondary delusion of Persecution

Delusion of reference:

• Delusion in which events, objects, or other persons in one's immediate environment have a particular and unusual significance in reference to the person experiencing the delusion.

• Examples :

• A feeling that people on television or radio are talking about or talking directly to them

• Believing that headlines or stories in newspapers are written especially for them

Solution to Question 5:

According to the given scenario,

A stranger has taken the guise of Wife (Family person)

This is characteristic of Capgras syndrome.

Capgras syndrome

The delusion in Capgras syndrome is the belief that a familiar person has been replaced by an imposter

Fregoli's phenomenon

• Fregoli syndrome is a disorder in which a person holds a delusional belief that different people are in fact a single person who changes his or her appearance or is in disguise.

Intermetamorphosis

• A very rare delusion that familiar persons can change themselves into other persons at will.

Doppelganger

• A person feels he has his own impostor (duplicate) – Delusion of subjective doubles

Othello syndrome

• It is a rare psychiatric condition marked by morbid, pathological, or delusional jealousy.

• Eg : Suspecting a faithful partner of infidelity, with accompanying jealousy.

Solution to Question 6:

The probable diagnosis is Akathisia.

Haloperidol is an antipsychotic drug and a side effect is Extrapyramidal symptoms.

Inner restlessness and urge to move are EPS referred to as Akathisia.

Akathisia

Akathisia consists of motor restlessness accompanied by subjective feelings of inner tension and discomfort referable chiefly to the limbs.

Restless leg syndrome

Closely resembles akathisia but occurs in absence of neuroleptic medication. No neurological deficit is seen.

Tardive Dyskinesia

Tardive dyskinesia is involuntary movements of the tongue, lips, face, trunk, and extremities that occur in patients treated with long-term dopaminergic antagonist medications.

Rabbit syndrome

Rabbit syndrome is an antipsychotic-induced rhythmic motion of the mouth/lips, resembling the chewing movements of a rabbit.

The movement consists of a vertical-only motion, at about 5Hz, with no involvement of the tongue.

Solution to Question 7:

The above scenario is consistent with the diagnosis of dysthymia.

Dysthymia / Persistent Depressive Disorder:

•

DSM V Criteria for diagnosis of dysthymia: Depressed mood for most of the day as indicated by either subjective account or observation by others, for at least 2 years. In children and adolescents - at least 1 year.

Presence of two (or more) of the following while being depressed.

• Poor appetite or overeating

• Insomnia or hypersomnia

• Low energy or fatigue

• Low self-esteem

• Poor concentration and/or difficulty making decisions

• Feelings of hopelessness

Early-onset, beginning in childhood or adolescence and in most cases by the age of 20 years. More common in unmarried and young persons with low income.

Cyclothymia:

• A milder form of bipolar disorder which is chronic in nature.

• Characterized by episodic mood swings of hypomania and mild depression.

Adjustment disorder:

• It is an abnormal and excessive reaction to an identifiable life stressor.

• It is a temporary condition

• Eg: Feeling sad or hopeless in an exaggerated manner after a stressful life event. Inability to cope with stressful situations.

Solution to Question 8:

Akinetic mutism is a feature of Stupor.

• In this condition , the patient is unable to initiate action and speech but appears awake and alert.

• There is severe reduction of relational function – speech and action.

Akinetic mutism is also seen in Schizophrenia with affective psychosis and dissociative states.

Stupor :

• Lesion at :

• Diencephalon

• Upper brain stem

• Frontal lobes

• Basal ganglia

• Locked In syndrome

Option A, Oneroid state

• Impaired consciousness.

• Marked emotional changes in response to visual hallucination.

• Patient lives as if in a dream state – occupational delirium.

Option B, Twilight state

• Abrupt onset of abnormal violent behavior

• Occurs in

• Epilepsy

• Alcoholism

• Brain trauma

Option D, Delirium

• Global impairment of consciousness and cognition.

Solution to Question 9:

In the above scenario, the patient has forced thinking.

Forced thinking is usually seen in patients with frontal lobe epilepsy.The thoughts are formed are not in the control of the patient and are distressing thoughts.

Option A: thought insertion is seen in schizophrenia. The delusion of thoughts being placed into patient's mind by someone else.

Option C: crowding of thoughts is also seen in schizophrenia. The patient has thoughts that are too fast and excessive and unexplainable. The patient can even locate these thoughts crowding at the back of the head.

Option D: The obsessive-compulsive symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication) or another medical condition.The patient feels the need to carry out certain behaviours to get rid of these thoughts.

Psychiatry AIIMS 2018

Question 1:

A patient was prescribed escitalopram. Which of the following adverse effect is the least likely?

a) Nausea

b) Vivid dreams

c) Anorgasmia

d) Sialorrhea

Question 2:

Abrupt stoppage of which of the following drugs causes agitation, anxiety and insomnia?

a) Valporate

b) Olanzapine

c) Imipramine

d) Venlafaxine

Question 3:

A young male who is shy and prefers social isolation appears to be emotionally cold and resides alone. There is no history of hallucinations or delusions. The most likely diagnosis is _____

a) Paranoid personality disorder

b) Schizoid personality disorder

c) Antisocial personality disorder

d) Emotionally unstable personality disorder

Question 4:

A 14 year old girl presents to the emergency room with fracture forearm and says she has tripped and fell. On examination you notice multiple cigarette burns on her forearm. What

will you do next?

a) Conduct a thorough examination

b) Seek a social worker's help

c) Inform higher authorities immediately

d) Inform your colleagues

Question 5:

Among the following, what is the most common symptom of adult obsessive-compulsive disorder?

a) Aggressive symptoms

b) Need for symmetry

c) Pathological doubt

d) Sexual symptoms

Question 6:

Which of the following is not a feature of atypical depression?

a) Increased sleep

b) Reactive mood response to positive stimulus

c) Heaviness of the body

d) Responds better to TCAs than MAO-I and SSRI

Question 7:

Under the Mental Health Care Act 2017, a patient has the right to choose his or her caretaker and future course of action in treatment. What is this provision known as?

a) Treatment directive

b) Mental will

c) Future directive

d) Advance directive

Question 8:

What does the given image represent?



a) Hypnogram

b) Electroencephalogram

c) Polysomnogram

d) Epworth sleep assessment

Question 9:

A 40-year-old woman has a history of urinating in public without having any remorse for it. She also has difficulty in getting motivated and has mild memory deficits. Which of the following lobes may be affected?

a) Parietal

b) Occipital

c) Frontal

d) Temporal

Answer Key

Question No.	Correct Option
1	d

2	d
3	b
4	a
5	c
6	d
7	d
8	a
9	c

Detailed Explanations

Solution to Question 1:

Sialorrhea is not a side effect of escitalopram.

Escitalopram is a selective serotonin reuptake inhibitor (SSRI). SSRIs are associated with dry mouth, and not sialorrhea.

System	Side effects of SSRIs
Sexual	Anorgasmia, Decreased libido
Gastrointestinal (5-HT3 receptor)	Nausea, Diarrhea, Anorexia, Vomiting, Flatulence, and Dyspepsia
Nervous system	Extremelyvivid dreamsor nightmares, Anxiety, Insomnia/sedation, Seizures, Emotional blunting
Anticholinergic	Dry mouth, Constipation, Sedation
Cardiovascular	Drug induced long QT syndrome
Hematological	Easy bruising
Endocrine	Mammoplasia, Galactorrhea

Solution to Question 2:

Abrupt stoppage of venlafaxine causes agitation, anxiety, and insomnia.

Solution to Question 3:

The diagnosis, in this case, is schizoid personality disorder.

Schizoid personality disorder is a cluster A personality disorder. The patient neither desires nor enjoys close relationships (shy and prefers social isolation, resides alone as in the question) and shows emotional coldness and detachment.

In contrast, subjects with emotionally unstable (borderline) personality disorder cannot tolerate being alone, and they search for companionship.

Solution to Question 4:

A child who has been physically abused should undergo a thorough physical examination at first.

This should be followed by a psychiatric interview.

When the child is brought to the emergency room, a detailed and spontaneous account of the injury should be obtained promptly from parents or other caregivers before secondary details and rationalizations cloud the information provided.

Solution to Question 5:

Among the above options, pathological doubt is the common symptom of adult OCD. Pathological doubt is the second most common pattern of adult obsessive-compulsive disorder (OCD), only after obsession of contamination.

Pathological doubt is an obsession with doubt, followed by a compulsion to check. The obsession often implies some danger of violence e.g., forgetting to turn off the stove or not locking a door. The checking may involve multiple trips back into the house to check the stove. These patients have obsessional self-doubt and always feel guilty about having forgotten or committed something.

The frequency of presenting patterns of OCD :

Obsession of contamination > Pathological doubt > Intrusive thoughts> Symmetry

Solution to Question 6:

Patients with atypical features do respond effectively to treatment with monoamine oxidase inhibitors (MAOIs) and selective serotonin reuptake inhibitors (SSRIs).

Mood reactivity, increased sleep, weight gain, and leaden paralysis (heaviness of body) are features of atypical depression.

Mood reactivity is the capacity to be cheered up when presented with positive events.

Leaden paralysis is the heavy, leaden feelings in arms or legs.

Solution to Question 7:

The provision by which a patient has the right to choose his caretaker and future course of action in treatment because of illness or incapacity is known as an advance directive. It is also known as a living will, personal directive, or medical directive.

Solution to Question 8:

The above image represents a hypnogram. It is a graph that depicts the stages of sleep as a function of time. It provides a visual representation of sleep structure and is useful in the diagnosis of various sleep disorders.

An electroencephalogram is the depiction of the electrical activity of the surface of the brain. Different waveforms (alpha, beta, theta, delta) are produced depending on the functional status of the brain.



Polysomnography is a multi parametric test that is considered the gold standard in the diagnosis of sleep disorders. It is a continuous recording of biophysiological changes that occur during sleep. It includes,

• Continuous EEG recording, from occipital and parietal leads

• Electro-oculography- to record eye movements

• Electromyography- for muscle potential and its activities

• ECG

• Multiple sleep latency tests- repeated measures of time taken to sleep

• Nasal-oral airflow, respiratory effort, oxyhemoglobin saturation, and leg movements during sleep are measured.



Epworth sleep assessment scale is a questionnaire used to quantify daytime sleepiness. It determines how frequently a patient is likely to doze off in 8 commonly encountered situations. It is a subjective assessment and hence less reliable.

Solution to Question 9:

The frontal lobe is affected in this patient. The woman has difficulty in getting motivated, memory deficits, and urinary incontinence.

Frontal lobe injury is commonly due to trauma, infarcts, tumors, lobotomy, multiple sclerosis, or Pick's disease. The following features may be seen in a patient with frontal lobe injury:

• Impaired executive functions such as motivation, attention, and sequencing of actions

• Aphasia, apraxia, constructional apraxia, neglect, and memory deficits

• Urinary incontinence and fecal incontinence is commonly seen

• Changes in personality

• Apathetic indifference and impulsive behavior.

Note: Frontal lobe pathology may become apparent only under unstructured, stressful, real-life situations and are difficult to detect using Intelligence scales, as IQ mostly requires parietal lobe activation.

Psychiatry AIIMS 2019 (May & Nov)

Question 1:

Under the Mental Health Care Act 2017, a patient has the right to choose his or her caretaker and future course of action in treatment. What is this provision known as?

a) Treatment directive

b) Mental will

c) Future directive

d) Advance directive

Question 2:

Which of the following is not a feature of melancholia?

a) Severe anhedonia

b) Weight loss

c) Sleep-onset insomnia

d) Profound feelings of guilt over trivial events

Question 3:

Which of the following best describes paradoxical suicide?

a) Suicide occuring at the time when the patient starts to recover

b) Suicide after taking low dose of drug

c) Accidental completion of suicide

d) None of the above

Question 4:

Which of the following is true regarding depersonalization disorder?

a) More common after life threatening trauma

b) Seen in seizure and migraine patients

c) More common in females

d) Mean age at onset is 25 years

Question 5:

A patient says that he feels that there is something wrong with him mentally and it is because of the summer season. How much would he score on the insight scale?

a) 2

b) 3

c) 4

d) 5

Question 6:

A patient was prescribed escitalopram. Which of the following adverse effect is the least likely?

a) Nausea

b) Vivid dreams

c) Anorgasmia

d) Sialorrhea

Question 7:

A 48-year-old male presents to the ER with confusion and shaking of the entire body. He has a history of excessive alcohol consumption. What is the cause of delirium tremens in an alcoholic?

a) Acute infection

b) Fatty liver

c) Gradual withdrawal of alcohol

d) Small doses of regular consumption of alcohol

Question 8:

A patient of anorexia nervosa was eating all her meals according to the prescribed diet chart but there is no change in her weight after 1 week. Which of the following needs to be done?

a) Observe the patient for 2 hours post meal

b) Ask the psychiatrist to prescribe anti-anxiety medication

c) Increase the calories from 1500 to 2000 in her diet

d) Increase water intake

Answer Key

Question No.	Correct Option
1	d
2	c
3	a
4	a
5	b
6	d
7	a
8	a

Detailed Explanations

Solution to Question 1:

The provision by which a patient has the right to choose his caretaker and future course of action in treatment because of illness or incapacity is known as an advance directive. It is also known as a living will, personal directive, or medical directive.

Solution to Question 2:

Sleep-onset insomnia is a characteristic feature of anxiety disorder, not melancholia.

Melancholia is one of the oldest terms used in psychiatry, to describe the dark mood of depression. It is still used to refer to a state of depression characterized by:

• Severe anhedonia

• Early morning awakening (terminal insomnia)

• Weight loss

• Profound feelings of guilt (often over trivial events)

Melancholia is sometimes referred to as "endogenous depression" or depression that arises in the absence of external life stressors or precipitants.

Solution to Question 3:

Paradoxical suicide is defined as suicide which occurs when the patient begins to recover from the depression

• This is because, when the patient is in depression, though he/she has the suicidal thoughts cannot act on it due to lack of energy.

• As the patient begins to recover from depression, patients gain energy to put their suicidal thoughts into action.

It is called paradoxical, as one might expect that the risk of suicide reduces as the patient starts to recover, but on contrary to that, the risk of suicide is increased during the recovery phase.

Solution to Question 4:

Depersonalization is common after a life-threatening trauma.

The DSM-V defines depersonalisation as experiences of unreality, detachment, or being an outside observer with respect to one's thoughts, feelings, sensations, body, or actions (e.g., perceptual alterations, distorted sense of time, unreal or absent self, emotional and/ or physical numbing).

Another term called derealization refers to the experiences of unreality or detachment with respect to the surroundings.

The DSM-V diagnostic criteria for depersonalization/derealization disorder is as follows:

• The presence of persistent or recurrent experiences of depersonalization, derealization, or both.

• Intact reality testing, during the episodes.

• The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

• The disturbance is not attributable to the physiological effects of a substance(e.g., a drug of abuse, medication) or another medical condition(e.g., seizures). (option b)

• The disturbance is not better explained by another mental disorder, such as schizophrenia, panic disorder, major depressive disorder, acute stress disorder, posttraumatic stress disorder, or another dissociative disorder.

Characteristic features of depersonalization/derealization disorder:

• The condition commonly occurs after a life-threatening trauma.

• There is no gender preponderance. Male:Female ratio is 1:1 (option c).

• Mean age at onset is 16 years (option d).

• The condition is commonly associated with symptoms such as an altered sense of time, difficulty in remembering past memories, anxiety and depression.

Solution to Question 5:

The patient would score 3, as he is aware of being sick but has attributed it to external factors.

Insight is rated on a 5 point scale:

• Complete denial of illness.

• Slight awareness of being sick and needing help but denying it at the same time.

• Awareness of being sick, but is attributed to physical or external factors.

• Intellectual insight - awareness of being ill, and that the symptoms are due to own particular irrational feelings and thoughts.

• Emotional insight - the highest level of insight. Intellectual insight with making significant changes in future behavior.

Loss of insight is seen in schizophrenia, mania, delusional disorders, etc.

Solution to Question 6:

Sialorrhea is not a side effect of escitalopram.

Escitalopram is a selective serotonin reuptake inhibitor (SSRI). SSRIs are associated with dry mouth, and not sialorrhea.

System	Side effects of SSRIs
Sexual	Anorgasmia, Decreased libid o
Gastrointestinal (5-HT3 rece ptor)	Nausea, Diarrhea, Anorexia, Vomiting, Flatulence, and Dy spepsia
Nervous system	Extremelyvivid dreamsor nig htmares, Anxiety, Insomnia/s edation, Seizures, Emotional blunting
Anticholinergic	Dry mouth, Constipation, Se dation
Cardiovascular	Drug induced long QT syndr ome
Hematological	Easy bruising
Endocrine	Mammoplasia, Galactorrhea

Solution to Question 7:

Among the given options, acute infection can be a cause of delirium tremens in an alcoholic patient.

Physical illness e.g., hepatitis or pancreatitis predisposes to the syndrome; a person in good physical health rarely has delirium tremens during alcohol withdrawal.

The syndrome usually develops on the third day of admission. A patient admitted for an unrelated condition may unexpectedly have an episode of delirium, which is the first sign of a previously undiagnosed alcohol-related disorder.

Episodes of delirium tremens usually begin at the age of 30-40 years, after 5 to 15 years of heavy drinking, and are typical after a binge. Gradual withdrawal of alcohol does not cause delirium tremens. However, sudden withdrawal can cause it.

Treatment of alcohol withdrawal:

• For patients showing withdrawal symptoms, chlordiazepoxide 25-50 mg every 2-4 hours can be administered orally.

• For patients with delirium tremens, 50-100 mg of chlordiazepoxide should be administered every 4 hours.

• If oral administration of drugs is not possible, intravenous lorazepam can be tried.

Solution to Question 8:

Observe the patient for 2 hours post-meal because the patient might vomit the food.

Treatment of anorexia nervosa:

• Hospitalization:

• Patients with a weight 20% less than expected weight for height should be considered for hospitalization.

• During a hospitalized treatment program, patients should be weighed daily, early in the morning after emptying the bladder.

• The daily fluid intake and urine output should be recorded.

• If vomiting is occurring, hospital staff members must monitor serum electrolyte levels regularly and watch for the development of hypokalemia.

• Because food is often regurgitated after meals, the staff may be able to control vomiting by making the bathroom inaccessible for at least 2 hours after meals or by having an attendant in the bathroom to prevent the opportunity for vomiting.

• Psychotherapy:

• Cognitive-behavioral therapy (CBT)

• Dynamic psychotherapy

• Family therapy

• Pharmacotherapy:

• Pharmacological agents are not the first-line treatment but cyproheptadine, amitriptyline, clomipramine, pimozide, and chlorpromazine show some improvement in these patients.

• Trials of fluoxetine have resulted in some reports of weight gain.

Psychiatry AIIMS 2020 (May and Nov INI-CET)

Question 1:

You are attending physiology viva. The examiner asks you who recorded the first EEG. You say "I don't know ". The examiner feels bad for you and gives you a clue that "the name sounds like something you get commonly in McDonalds". You remember the name and answer correctly. What type of memory has been implicated here?"

a) Semantic memory

b) Implicit memory

c) Associative learning

d) Priming

Question 2:

A known alcoholic patient is brought to the emergency in a disoriented state. He complains of seeing snakes and insects crawling on his skin. Which of the following should be given to this patient?

a) Haloperidol + thiamine

b) Lorazepam only

c) Lorazepam + thiamine

d) Haloperidol only

Question 3:

A drug that selectively reduces the reuptake of serotonin would likely cause which of the following side effects?

a) Constipation

b) Sexual dysfunction

c) Sialorrhea

d) Visual disturbance

Question 4:

A patient undergoing ECT for his severe depression was given thiopentone and succinylcholine. The purpose of these medications was to?

a) To bring calming effect

b) To produce mood elevation

c) To induce anesthesia and relax muscles

d) To prevent memory loss after the ECT

Question 5:

A young male who is shy and prefers social isolation appears to be emotionally cold and resides alone. There is no history of hallucinations or delusions. The most likely diagnosis is _____

a) Paranoid personality disorder

b) Schizoid personality disorder

c) Antisocial personality disorder

d) Emotionally unstable personality disorder

Question 6:

A 40-year-old male patient comes to the psychiatry OPD with complaints of repetitive thoughts that, he always feels his hands are dirty, though they are not. This gives him discomfort and hence he has to wash them again and again. This is a disorder of thought _____.

a) Flow

b) Form

c) Possession

d) Content

Question 7:

Which of the following is not a type of subcortical dementia?

a) Pick's disease

b) Parkinson's disease

c) Huntington's disease

d) Wilson's disease

Question 8:

Which of the following is not a good prognostic factor in schizophrenia?

a) Positive symptoms

b) Late age of onset

c) Insidious onset of symptoms

d) Associated with depression

Question 9:

A 7-year-old child presented with complaints of bed-wetting almost twice a week for the past 1 year. A detailed investigation was done to rule out the organic cause. What should be the treatment plan for this child?

a) Pharmacotherapy with imipramine

b) Psychodynamic psychotherapy

c) Bladder training, with reward for delaying micturition during daytime

d) Bell and pad based classical conditioning

Question 10:

A male patient who lost his job recently became irritable and had a sad mood for most of the time. Thinking about the future worried him and he felt irritation towards his family members. He occasionally went out to watch movies with his friends and was able to enjoy but on returning home he felt similar symptoms. What is the probable diagnosis?

a) Generalised anxiety disorder

b) Adjustment disorder

c) Mixed anxiety with depression

d) Moderate depression

Question 11:

A 10-year-old child is suffering from selective mutism. He is most probably suffering from?

a) Childhood depression

b) Childhood psychosis

c) Childhood anxiety disorder

d) Hyperkinetic disorder

Question 12:

A psychotic patient developed voluntary purposeless movements and was once observed to be standing still for hours with waxy flexibility and negativism. What is the appropriate medical management?

a) Haloperidol

b) Lorazepam

c) Clonidine

d) Propanolol

Question 13:

Which of the following features differentiate delirium from dementia in Alzheimer's disease?

a) Acuity of onset and agitation

b) Acuity of onset and loss of consciousness

c) Visual hallucinations and impaired memory

d) Agitation and irritation

Answer Key

Question No.	Correct Option
1	d
2	c
3	b
4	c
5	b
6	c
7	a

8	c
9	d
10	b
11	c
12	b
13	b

Detailed Explanations

Solution to Question 1:

In the given scenario, priming is the most appropriate answer.

The examiner is asking for a fact, and then prompting the answer, by giving a clue which sounds (phonological) like the answer. This is an example of priming.

Prime is any cue, which precedes the recall of the target word. It maybe similar to hear (sun-bun-gun) or may hold similarity to the target word (Sun- moon - star).

Solution to Question 2:

The given clinical scenario points towards delirium tremens likely due to acute alcohol withdrawal. Lorazepam + thiamine should be given to this patient.

Delirium tremens occurs after 2-4 days of abstinence in a heavy or chronic drinker. It is characterized by autonomic hyperactivity, tremors, delusions, and hallucinations. It usually resolves spontaneously in 7 days but may cause the death of the patient due to cardiovascular collapse or infections or self-injury.

Delirium tremens is treated using benzodiazepines starting with a higher dosage which is gradually tapered as withdrawal symptoms disappear. In cases of chronic alcohol use, vitamin deficiency (especially thiamine) may be co-existing and hence bolus doses of thiamine are given to prevent/treat Wernicke encephalopathy.

Haloperidol is generally not the first-line drug in case of alcoholic withdrawal. It might be used as an add-on in agitated patients. It decreases the seizure threshold and its use would be appropriate in a case of co-existing psychotic disorder like schizophrenia.

Drug Therapy For Alcohol Intoxication and Withdrawal			
Clinical Problem	Drug	Route	Dosage
Tremulousness and mild to moderate agitation	ChlordiazepoxideDiazepam	OralOral	25-100 mg every 4-6 hrs5-20 mg every 4-6 hrs
Hallucinosis	Lorazepam	Oral	2-10 mg every 4-6 hrs

Drug Therapy For Alcohol In toxication and Withdrawal			
Extreme agitation	Chlordiazepoxide	Intraven ous	0.5 mg/kg at 12 mg/min
Withdrawal seizures	Diazepam	Intraven ous	0.15 mg/kg at 2.5 mg/min
Delirium tremens	Lorazepam	Intraven ous	0.1 mg/kg at 2.0 mg/min

Solution to Question 3:

Selective serotonin reuptake inhibitors (SSRIs) are associated with sexual dysfunction.

Note - Sexual dysfunction is the most prominent side effect of SSRIs and weak anticholinergic effects are only shown by paroxetine, and the GI effects of SSRIs predominantly include diarrhoea.

System	Side effects of SSRIs
Sexual	Anorgasmia, Decreased libid o
Gastrointestinal (5-HT3 rece ptor)	Nausea,diarrhea, anorexia, v omiting, flatulence, and dysp epsia
Nervous system	Extremelyvivid dreamsor nig htmares, Anxiety, Insomnia/s edation, seizures, emotional blunting
Anticholinergic	Dry mouth, constipation, sed ation
Cardiovascular	Drug-induced long QT syndr ome
Hematological	Easy bruising
Endocrine	Mammoplasia, galactorrhea

Solution to Question 4:

Thiopentone and succinylcholine are given for anesthesia induction and muscle relaxation respectively.

Thiopentone sodium is a barbiturate and acts by opening chloride channels, whilst succinylcholine is a depolarising muscle relaxant.

ECT can be of 2 types:

• Direct - without anesthesia and muscle relaxant

• Modified - with anesthesia and muscle relaxant

Currently modified ECT is preferred and is given with adequate spacing i.e., no more than 3 ECT sessions per week.

After adequate preparation, a current of 200-1600mA (110V for 0.1-1 seconds) is given, resulting in an adequate seizure. Electroconvulsive therapy treatments are associated with retrograde amnesia for a period of several minutes before the treatment and anterograde amnesia after the treatment.

Solution to Question 5:

The diagnosis, in this case, is schizoid personality disorder.

Schizoid personality disorder is a cluster A personality disorder. The patient neither desires nor enjoys close relationships (shy and prefers social isolation, resides alone as in the question) and shows emotional coldness and detachment.

In contrast, subjects with emotionally unstable (borderline) personality disorder cannot tolerate being alone, and they search for companionship.

Solution to Question 6:

The clinical history is suggestive of obsessive-compulsive disorder, which is a disorder of thought possession.

The diseases under thought possession include:

• Obsession and Compulsion

• Thought Alienation

In obsessions and compulsions, although the sense of thought possession in the patient is not impaired, he/she experiences the obsessive thoughts to appear against their will. Hence, despite the patient perceiving a loss of control over the voluntary act (compulsion), they still retain personal possession (ownership) of the act.

Obsessive-compulsive disorder is defined as the presence of obsessions and/or compulsions.

• Obsessions are recurrent and persistent ideas, thoughts, urges, or images that are experienced as intrusive and unwanted and cause anxiety or distress to the patients.

• Compulsions are the repetitive and intentional behaviors or acts performed in response to obsessions or according to certain rules that must be applied rigidly (e.g., repetitive hand washing, ritualistic checking).

In the given case, the obsessional thoughts are the repetitive, unwanted thoughts that his hands are dirty, while the compulsion is to wash his hands to neutralize the obsession.



Solution to Question 7:

Pick's disease (a type of frontotemporal dementia) is a type of cortical dementia.

	Cortical dementia	Subcortical dementia
Site of lesion	Cortex	Thalamus, basal ganglia, cere bellum, brain stem
Higher cortical sympto ms	AphasiaApraxiaAgnosiaProso pagnosiaAcalculia	No
Motor symptoms	No	DysarthriaTremorsAtaxiaRigi dity
Memory	Recall and recognition are im paired	Recall is impaired
Examples	Alzheimer's diseaseFronto-te mporal dementiaCreutzfeldt Jakob disease	Parkinson's diseaseProgressi ve supranuclear palsyHuntin gton's diseaseWilson's diseas eIdiopathic basal ganglia calc ificationNormal-pressure hyd rocephalusVascular dementia

Solution to Question 8:

Insidious onset of symptoms is a poor prognostic factor of schizophrenia.

Prognostic factors of Schizophrenia:

T.J.Crow classified schizophrenia into two separate types :

Good Prognostic factors	Poor prognostic factors
Acute onset	Insidious onset
Late-onset	Early-onset
Presence of a stressor	Absence of a stressor
Short duration < 6 months	Duration >2 years
Presence of depression	No depression
Positive symptoms	Negative symptoms
Family history of mood disor der	Family history of schizophre nia
Married, female sex	Single, male sex
Catatonic schizophrenia	Hebephrenic schizophrenia
Type 1 schizophrenia	Type 2 schizophrenia
	No remissions in 3 years
	History of perinatal trauma
	History of assaultiveness

Type 1	Type 2
Positive symptoms present	Negative symptoms present
More responsive to treatment	Less responsive to treatment
No demonstrable neuropatho logy	Demonstrable neuropatholog y (CT/MRI)
Good prognosis	Poor prognosis

Solution to Question 9:

The clinical features are suggestive of nocturnal enuresis. Bell and pad based classical conditioning is the most effective treatment.

Nocturnal enuresis refers to the occurrence of involuntary urination at night after 5 years of age. It may be a normal phenomenon in children up to 5 years of age.

Treatment for nocturnal enuresis may include:

Behavioural therapy:

• Classic conditioning with the bell (or buzzer) and pad (alarm) apparatus is the most effective treatment for enuresis.

• Dryness is achieved in 50% of the cases.

• Bladder training with encouragement or reward for delaying micturition for increasing times during waking hours is useful but considered inferior to bell and pad.

Pharmacotherapy:

• Medications should be considered if behavioural therapy has failed, and if enuresis is causing impairment in social, family, and school function.

• Desmopressin (an antidiuretic) and Reboxetine (norepinephrine reuptake inhibitor) are commonly used drugs. Reboxetine is considered as a safer alternative to imipramine.

Psychotherapy:

• It is indicated in cases with chronic enuresis, to deal with any coexisting psychiatric problems and emotional and family difficulties.

Solution to Question 10:

The clinical features are suggestive of an adjustment disorder.

Adjustment disorder is an emotional response to a stressful event. Stressors may be single like loss of job or divorce; multiple like the death of a parent together with loss of a job; or recurrent like seasonal business difficulties.

Diagnosing criteria for adjustment disorder:

• Development of emotional or behavioural response within 3 months of the onset of the stressor.

• The symptoms or behaviour are clinically significant, where, any one of the following are met:

• The response is out of proportion to the severity or intensity of the stressor.

• Significant impairment in social, occupational, or other areas of functioning.

• The stress-related disturbance does not meet the criteria for another mental disorder or exacerbation of pre-existing mental disorder.

• Mostly depressive but can also have anxiety symptoms.

• The symptoms disappear within 6 months if the stressor is removed.

In the given case, it has been mentioned that the patient lost his job recently. Although a time period is not mentioned in the question, the term recently can be understood as well within the 3 months duration. The patient's behaviour such as sad and irritated affected his social functioning. As he was finding joy in watching movies, depression can be ruled out.

Solution to Question 11:

The child is most likely suffering from a childhood anxiety disorder.

Selective mutism refers to the lack of speaking (being mute) in one or more specific social situations, commonly in school, but speaking well in other situations, commonly at home. The child may be either completely silent at school or maybe just whispering. Rarely, a child may be silent at home but communicating normally at school.

The disorder may develop in children between 4 to 8 years of age and is associated with a social anxiety disorder. They may have symptoms of separation anxiety disorder, school refusal, and

delayed language acquisition.

The first-line treatment includes cognitive behavioural therapy.

Option A and B: selective mutism is not commonly associated with childhood depression or psychosis

Option D: Hyperkinetic disorder is associated with overfamiliarity.

Solution to Question 12:

The clinical features are suggestive of a catatonic disorder. Benzodiazepines such as lorazepam are effective in the treatment of catatonia.

Catatonia is a syndrome characterized by striking behavioral abnormalities such as motoric immobility or excitement, negativism, or echolalia (mimicry of speech), or echopraxia (mimicry of movement).

The clinical picture may include three or more of the following features:

• Stupor: Absence of psychomotor activity and not actively relating to the environment.

• Catalepsy: Passive induction of a posture held against gravity.

• Waxy flexibility: Slight, even resistance to positioning by examiner.

• Negativism: Opposition or no response to instructions or external stimuli.

• Posturing: Spontaneous and active maintenance of a posture against gravity.

• Mannerism: Odd, circumstantial caricature of normal actions.

• Stereotypy: Repetitive, abnormally frequent, non-goal-directed movements

• Automatic obedience: Commands are followed automatically, irrespective of their nature.

• Grimacing

• Ambitendency: Due to ambivalence, opposing actions are carried out.

• Echopraxia: Mimicking another's movement.

• Echolalia: Mimicking another's speech.

• Mutism: Very little or no verbal response to instructions or external stimuli.

• Verbigeration: Incomprehensible speech.

Catatonic disorder may occur due to:

• Advanced mood or psychotic illnesses

• Neurological disorders (e.g., nonconvulsive status epilepticus, and head trauma)

• Infections (e.g., encephalitis)

• Metabolic disturbances (e.g., hepatic encephalopathy, hyponatremia, and hypercalcemia)

• Drugs induced- like corticosteroids, immunosuppressants, and antipsychotics (i.e., neuroleptic) agents.

Management: The primary treatment includes:

• Identifying and treating underlying medical or pharmacological causes. Any offending substances should be removed.

• Benzodiazepines, especially, lorazepam, are effective in improving the symptoms.

• Electroconvulsive therapy is the definitive treatment in cases of life-threatening catatonia (if the patient is unable to eat) or if the patient has developed malignant catatonia.

Solution to Question 13:

Delirium is characterized by acute onset of symptoms and is often accompanied by diminished or loss of consciousness and cognition. This feature is useful to clinically differentiate between delirium and dementia.

Dementia is a group of diseases marked by chronic, progressive cognitive impairment with clear consciousness.

Psychiatry NEET 2018

Question 1:

A 40-year-old lady met with a car accident 2 months ago, following which she complains of screaming and waking up in the night after having nightmares of the same incident over and over again. Which of the following conditions is she suffering from?

a) Acute stress reaction

b) Adjustment disorder

c) Mania

d) Post traumatic stress disorder

Question 2:

Which of the following features is more in favour of delirium?

a) Occurs gradually over a period of time

b) Fluctuating course

c) Preserved consciousness

d) Commonly associated with auditory hallucinations

Question 3:

What is defined as the emotional release and discharge after consciously reliving a painful experience that has been repressed?

a) Catharsis

b) Abreaction

c) Venting out

d) Guided relaxation

Question 4:

What is 'mental retardation' now called according to the American Psychiatric Association in its revision of DSM-5?

a) Feeble mindedness

b) Mental subnormality

c) Intellectual disability

d) Intellectually challenged

Question 5:

The confusion assessment test is used in:

a) Dementia

b) Depression

c) Delirium

d) Schizophrenia

Question 6:

Which of these disorders was earlier known as minimal brain dysfunction?

a) Dyslexia

b) Attention deficit/hyperactivity disorder

c) Mental subnormality

d) Oligophrenia

Question 7:

Seman's squeeze technique is used for which of the following conditions?

a) Retrograde ejaculation

b) Erectile dysfunction

c) Premature ejaculation

d) Painful ejaculation

Question 8:

Addison's disease can present with symptoms similar to which of the following disorders?

a) Depression

b) Generalised anxiety disorder

c) Bipolar disorder

d) Panic disorder

Question 9:

Which of the following is not an OCD related disorder?

a) Hair pulling disorder

b) Temper tantrums

c) Hoarding disorder

d) Skin picking

Question 10:

Which of the following is not true about Alzheimer's disease?

a) Intracellular neurofibrillary tangles are seen

b) Neuritic plaques made of beta amyloid are found

c) Nucleus of Meynert is not affected

d) Short term memory is affected

Answer Key

Question No.	Correct Option
1	d
2	b
3	b
4	c
5	c
6	b
7	c
8	a
9	b
10	c

Detailed Explanations

Solution to Question 1:

The given clinical scenario is suggestive of post-traumatic stress disorder. PTSD is a psychiatric disorder occurring after a life-threatening or potentially fatal event, (e.g., war, house fire, serious accident, rape, robbery) affecting the patient or the patient's close friend or relative.

Symptoms can be divided into four types:

• Re-experiencing (e.g., intrusive memories of the event [flashbacks] and nightmares)

• Hyperarousal (e.g., anxiety, increased startle response, impaired sleep, hypervigilance)

• Negative cognition (e.g., negative beliefs, emotional numbing)

• Avoidance (e.g., avoiding stimuli associated with the traumatic event)

Psychotherapies found to be effective for PTSD include exposure therapy, trauma-focused cognitive-behavioral therapy, eye movement desensitization, and reprocessing.

Solution to Question 2:

Fluctuating course is suggestive of delirium.

Characteristic presentation of delirium:

• Acute onset

• Brief duration - usually days to weeks

• Decline in the level of consciousness.

• Fluctuating course - unpredictable fluctuations in severity and other clinical manifestations

• Sundowning - worsening of symptoms at night

• Rapid improvement - when the underlying cause is treated

• Other characteristics include asterixis, carphologia (picking movements on cover sheets and clothes)

• More commonly associated with visual hallucinations

Solution to Question 3:

Abreaction is a psychoanalytical term for reliving an experience to release it emotionally. It is a specific type of catharsis done by becoming conscious of repressed traumatic events.

Catharsis is a general term for the process of releasing strong or repressed emotions. It can be done through various mediums, like expressing emotions through art. When the unconscious repressed memories are consciously recalled and relived, this technique specifically is called abreaction.

Solution to Question 4:

Mental retardation is now known as Intellectual Disability, according to the revised DSM-5, by the American Psychiatrist Association.

Solution to Question 5:

The confusion assessment test is used in the detection of delirium.

Delirium is reported to occur in a wide range of mechanically ventilated ICU patients and can be detected by the Confusion Assessment Method (CAM)-ICU or the Intensive Care Delirium Screening Checklist.

These tools are used to ask patients to answer simple questions and perform simple tasks and can be used readily at the bedside.

Solution to Question 6:

Attention deficit/hyperactivity disorder (ADHD) was known as minimal brain dysfunction earlier.

Solution to Question 7:

Seman's squeeze technique is used for premature ejaculation, to prolong the time period until ejaculation.

Seman's technique (stop-start technique): It is a variation of the squeeze technique where a man approaches the point close to ejaculation and then stops before ejaculation and relaxes before starting stimulation again. This method prolongs the time until ejaculation by relying on behavior modification.

Other treatments for premature ejaculation:

• Squeeze technique (Master and Johnson's technique)- as a man approaches a climax, he squeezes the tip of the penis just below the head of the penis. The pressure is held there until the sensation of impending orgasm diminishes. The process can then be started over again so that over time a man prolongs the time period until he reaches ejaculation.

• Sensate focusing- refocussing on the non-coital aspects of sexual activity.

• Selective serotonin reuptake inhibitors (SSRIs)- delays ejaculation. e.g.- dapoxetine, paroxetine.

Solution to Question 8:

Addison's disease presents most similarly to depression. Apathy, social withdrawal, impaired sleep, decreased concentration, and prominent fatigue is frequently seen which is usually mistaken for depression.

Following infection, trauma, or surgery, a patient with previously unrecognized Addison's disease may suffer from an Addisonian crisis. Such patients present with either shock or hypoglycemia. This is a medical emergency requiring urgent medical referral.

Endocrinopathy	Differential psychiatric diagnosis
Addison's disease	Depression
Hyperthyroidism	Mood disorder, panic disorder, generalized anxiety disorder
Hypothyroidism	Depression,hypomania, delusions, schizophrenia (myxedema madness).
Hypoparathyroidism	Deliriumis the most common.
Hyperparathyroidism	Depression, anxiety,paranoid delusions. Tetrad of "painful bones, renal stones, abdominal groans, and psychiatric overtones".
Hypoglycemia	Anxiety, depersonalization, derealisation, aggressive behavior, amnesia.

Solution to Question 9:

Temper tantrums are not OCD-related.

Obsessive-compulsive and related disorders are:

• Obsessive-Compulsive Disorder

• Body Dysmorphic Disorder

• Hoarding Disorder

• Hair-Pulling Disorder (Trichotillomania)

• Excoriation (Skin-Picking) Disorder

Solution to Question 10:

Nucleus of Meynert is affected in Alzheimer's disease.

Biochemically, there is a reduction of acetylcholine and degeneration of cholinergic neurons in the nucleus basalis of Meynert that project throughout the cortex. There is also noradrenergic and serotonergic depletion due to degeneration of brain stem nuclei such as the locus ceruleus and the dorsal raphe.

Microscopically, there are neuritic plaques containing amyloid-β, neurofibrillary tangles, and amyloid-β accumulation in the blood vessel walls of the cortex and leptomeninges.

The FDA recently approved Flortaucipir F18 for intravenous injection, the first radioactive diagnostic agent for adult patients with cognitive impairment who are being evaluated for Alzheimer's disease. It is indicated for PET imaging of the brain to estimate the density and distribution of aggregated tau neurofibrillary tangles (NFTs), a primary marker of Alzheimer's disease.

Psychiatry NEET 2019

Question 1:

Which of the following drugs is used as an anti-smoking agent?

a) Busulfan

b) Acamprosate

c) Varenicline

d) Gabapentin

Question 2:

What is the false statement regarding somnambulism?

a) Disorder of sleep arousal

b) Person will have full consciousness

c) Usually, terminates in awakening followed by confusion

d) There may be neurological condition associated

Question 3:

Which of the following is not a risk factor for delusional disorder?

a) Recent immigration

b) Social isolation

c) Family history

d) Young age

Question 4:

Premature ejaculation is a disorder of which phase of the normal sexual cycle?

a) Desire

b) Arousal

c) Orgasm

d) Pain

Question 5:

Confusion assessment test is used in:

a) Delirium

b) Dementia

c) Schizophrenia

d) Depression

Question 6:

Which of the following has a poor prognosis with exposure and response prevention in OCD?

a) Pathological doubt

b) Magical thinking

c) Hoarding

d) Contamination obsession

Question 7:

What is the fear of heights known as?

a) Acarophobia

b) Algophobia

c) Agoraphobia

d) Acrophobia

Answer Key

Question No.	Correct Option
1	c
2	b
3	d

4	c
5	a
6	c
7	d

Detailed Explanations

Solution to Question 1:

Varenicline is an α4β2 nicotinic acetylcholine receptor partial agonist used in the treatment of nicotine addiction.

The FDA approved medication to aid in smoking cessation include:

• NRT-Nicotine Replacement Therapy

• Bupropion

• Varenicline

Drugs are only adjunctive to behavioral therapy, which is the first line of management.

Behavior therapy is the most widely accepted and well-proven psychological therapy for treating Tobacco use disorder. It consists of several techniques:

• Skill training and relapse prevention.

• Stimulus control.

• Mindfulness approaches.

Bupropion:

• It enhances both noradrenergic and dopaminergic neurotransmission by inhibiting reuptake.

• Used for aiding smoking cessation and for the treatment of depression in combination with SSRIs.

• Side effects: seizures, sedation

Varenicline:

• α4β2 Nicotinic ACh receptor partial agonist.

• Used for smoking cessation.

• Side effects: sleep disturbance, may depress mood and increase suicidal tendencies.

Mirtazapine:

• Serotonin receptor antagonist.

• Used for the treatment of depression in combination with SSRIs and for insomnia in depressive patients.

• Side effect: sedation

Solution to Question 2:

Somnambulism is characterized by sequence of behaviors without full consciousness.

Note: About 80% of sleepwalkers have a family history of sleepwalking, and an identical twin is much more likely to sleepwalk if their other twin sleepwalks.

Solution to Question 3:

Advanced age is a risk factor for delusional disorders, not young age.

Risk factors for delusional disorders:

• Advanced age

• Sensory impairment or isolation

• Personality features (e.g.unusual interpersonal sensitivity)

• Family history

• Social isolation

• Recent immigration

Solution to Question 4:

Premature ejaculation is a disorder of the orgasm phase of the normal sexual cycle.

Premature ejaculation is when ejaculation occurs before the completion of satisfactory sexual activity. In severe cases, it is characterized by ejaculation either before penile entry into the vagina or soon after penetration. The causes can be biological or psychological, like performance anxiety.

Solution to Question 5:

Confusion assessment test is used in detection of delirium.

Delirium is reported to occur in a wide range of mechanically ventilated ICU patients and can be detected by the Confusion Assessment Method (CAM)-ICU or the Intensive Care Delirium Screening Checklist. These tools are used to ask patients to answer simple questions and perform simple tasks and can be used readily at the bedside.

Solution to Question 6:

Hoarding behavior shows poor prognosis with exposure and response prevention.

Challenges posed by hoarding patients to typical cognitive behavior therapy (CBT) include poor insight into the behavior, low motivation, and resistance to treatment.

The most effective treatment for hoarding is a cognitive-behavioral model that includes:

• Training in decision making and categorizing

• Exposure and habituation to discarding

• Cognitive restructuring.

Solution to Question 7:

Acrophobia is the extreme or irrational fear of heights.

Phobia	Fear of
Acarophobia	Small bugs
Acrophobia	High places
Agoraphobia	Open or crowded places, leav ing familiar setting of home
Aichmophobia	Pointed objects
Ailurophobia	Cats
Algophobia	Pain
Androphobia	Men
Astraphobia or astrapophobi a	Thunder and lightning
Autophobia	Being alone, or of self
Bathophobia	Depths
Claustrophobia	Closed or confined places
Coprophobia	Excreta
Cynophobia	Dogs
Demophobia	Crowds
Erythrophobia	Blushing
Gynophobia	Women
Haphephobia	Being touched
Hematophobia	Sight of blood
Hydrophobia	Water
Hypnophobia	Sleep
Lalophobia or glossophobia	Speaking
Mysophobia	Dirt, contamination

Phobia	Fear of
Necrophobia	Dead bodies
Nyctophobia	Darkness, night
Panphobia	Everything
Pathophobia or nosophobia	Disease, suffering
Peccatophobia	Sinning
Pedophobia	Children
Phonophobia	Speaking aloud
Photophobia	Strong light
Sitophobia	Eating
Taphophobia	Being buried alive
Thanatophobia	Death
Toxophobia	Being poisoned
Xenophobia	Strangers
Zoophobia	Animals

Psychiatry NEET 2020

Question 1:

A 25-year-old male presented to the clinic with a complaint of the feeling of insects crawling under the skin. Which of the following drug abuse can cause the symptoms of this patient?

a) Cannabis

b) Cocaine

c) Amphetamine

d) Alcohol

Question 2:

A 60-year-old male who lost his wife 3 months back complains that his intestines have become rotten. He feels that he is responsible for his wife's death and should be sent to prison. He complains of feeling low all the time and has lost interest in daily activities since his wife's death. What is the most likely diagnosis?

a) Normal grief reaction

b) Psychotic depression

c) Delusional disorder

d) Schizophrenia

Question 3:

What is the treatment of choice for postpartum blues?

a) Fluoxetine

b) Cognitive behavioural therapy

c) Lithium carbonate

d) Support and education

Question 4:

A 25-year-old woman complains of a creepy crawling sensation in her legs. It is more at night and prevents her from sleeping. She is relieved of these symptoms by either walking or moving her legs. Which of the following drugs is used in treating the condition?

a) Gabapentin

b) Pramipexole

c) Vitamin B12

d) Iron tablets

Question 5:

A first-year MBBS student came to OPD with intermittent episodes of a feeling of impending doom accompanied by intense perspiration. Usually, the episodes occur prior to her exams. What is the most likely diagnosis?

a) Panic disorder

b) Conversion disorder

c) Social anxiety disorder

d) Generalized anxiety disorder

Question 6:

What is the maximum period of voluntary self-admission under the Mental Healthcare Act 2017?

a) 48 hours

b) 7 days

c) 30 days

d) 90 days

Question 7:

Identify the stage of sleep based on EEG finding highlighted in red.



a) Stage 1

b) Stage 2

c) Stage 3

d) REM sleep

Question 8:

A middle-aged female patient presents with a history of loss of interest in day to day activities, weight loss, insomnia, and sadness for six months. She also recalls hearing a voice once that ordered her to kill herself. What is the DSM-V diagnosis?

a) Major depressive disorder with psychotic features

b) Schizophrenia

c) Schizoaffective disorder

d) Delusional disorder

Answer Key

Question No.	Correct Option
1	b
2	b
3	d

4	a
5	a
6	c
7	d
8	a

Detailed Explanations

Solution to Question 1:

The feeling of insects crawling under the skin, also known as cocaine bugs or formication, is a feature of cocaine abuse.

Other features of acute cocaine intoxication are delusions of persecution, paranoia, and suspiciousness. It is essential to differentiate acute cocaine intoxication from acute schizophrenia which it closely resembles.

Solution to Question 2:

The given clinical scenario of persistent low mood, anhedonia, guilt, and self-critical thought content along with nihilistic delusion (rotten intestine), points towards a diagnosis of major depression with psychotic features.

Differentiating features of grief and depression have been enlisted in the following table:

Note: Tricyclic antidepressants are considered to be the first-line drugs for psychotic depression.

Grief	Depression
Feelings of emptyness and lo ss	Persistent low mood and ina bility to anticipate pleasure
Dysphoria comes in waves an d tends to decrease in intents ity over time	Low mood tends to be persist ent and consistent
Associated with thoughts and reminders of deceased/loss	Not tied to specific thoughts and pre-occupations
May be present along with po sitive emotions and humor	Pervassive unhappiness and misery
Thought content associated with thoughts and memories of deceased/loss	Thought content associated with self-critical and pessimis tic ruminations
Self-esteem is generally prese rved	Worthlessness and self-loathi ng is common

Grief	Depression
Ideas of death focused on joi ning the deceased	Ideas of death focused on en ding life due to hopelessness

Solution to Question 3:

Postpartum blues is common in mothers in the puerperal period and requires nothing more than education and support. It is characterised by a labile mood, sadness and subjective confusion. It is thought to be due to changes in hormone levels, the stress of childbirth and anxiety about the responsibilities of motherhood.

If symptoms last longer than 2 weeks, an evaluation for postpartum depression is needed.

Solution to Question 4:

The history is suggestive of restless leg syndrome and gabapentin is the first-line choice of treatment for this condition.

Gabapentinoids are considered superior to dopamine agonists due to the high frequency of intermediate and long-term augmentation with dopamine agonists, as well as their risk of impulse control disorders.

Restless legs syndrome (RLS) is an irresistible urge to move the legs when at rest or while trying to fall asleep or an uncomfortable sensation of creeping-crawling feeling. It is worse at night and moving legs or walking reduces discomfort. Uremia, neuropathies, as well as iron-deficiency and folic acid–deficiency anemias can produce secondary RLS.

In patients with restless legs syndrome, iron stores are initially repleted, and the aggravating factors are eliminated. If RLS still persists, initial therapy with a gabapentinoid (pregabalin, gabapentin, gabapentin enacarbil) is preferred, rather than a dopamine agonist (pramipexole, ropinirole, rotigotine).

FDA-approved drugs for the treatment of restless leg syndrome:

• Gabapentin

• Ropinirole

• Pramipexole

• Rotigotine

Note: A dopamine agonist is used as an alternative in patients with increased risk for side effects from gabapentinoids, such as those with obesity and its complications, a past or present history of moderate to severe depression, a disorder causing gait instability or respiratory failure, and a history of substance use disorder.

Solution to Question 5:

The most likely diagnosis for this patient is panic disorder.

A panic attack is an abrupt surge of intense fear or intense discomfort. Recurrent panic attacks for more than one month is classified as panic disorder.

Panic attacks can occur in mental disorders other than panic disorder, particularly in specific phobia, social phobia, and PTSD.

Solution to Question 6:

The maximum period of voluntary self-admission is 30 days.

When a person has a mental illness that requires admission, and if he/she is able to make his or her own decision and voluntarily requested admission, then it is called voluntary self-admission.

According to the Mental Healthcare Act, the maximum duration for which such admission can last is 30 days.

If the admission requires more than 30 days, or if it requires readmission within 7 days after discharge, then such patients should be examined by two psychiatrists and can be admitted for a maximum period of 90 days.

Note: The aforementioned question is as per the 2020 recall. This concept has been updated, and the admissions are classified as independent and supported. Please refer to the MCQ MG6497 for recent information.

Solution to Question 7:

The marked area in the given EEG shows sawtooth waves that are characteristic of REM sleep.

Sleep is comprised of two distinct physiological states, non-rapid eye movement (NREM) sleep and REM sleep. Stages I to IV make up the NREM sleep with marked lower physiological functions than during wakefulness. Dreams which occur during this state are usually lucid and purposeful.

REM sleep is characterized by a high level of brain activity. The physiological activity is similar to that during wakefulness. In REM sleep,

• Pulse, respiration, and blood pressure are increased.

• Brain oxygen use increases during REM sleep.

• Thermoregulation is altered - Poikilothermia prevails during REM sleep.

• There is loss of muscle tone - body movements are absent during REM sleep.

• There is partial or full penile erection - This is used to evaluate the cause for impotence.

• Dreams are abstract and surreal.

In normal adults, a REM episode has a latency of about 90 minutes during which the sleep is in the NREM stage. This latency is shortened in narcolepsy and depressive disorders.



Solution to Question 8:

Since the patient fulfills the DSM-V criteria for major depressive disorder and has had an episode of auditory hallucination, a diagnosis of major depressive disorder with psychotic features is made.

Diagnostic criteria for depression:

• Depressed mood most of the day

• Anhedonia- diminished interest in day to day activities

• Significant weight loss or weight gain. Change in appetite.

• Insomnia or hypersomnia

• Psychomotor agitation or retardation

• Fatigue or loss of energy

• Feelings of worthlessness or excessive, inappropriate guilt

• Diminished ability to think or concentrate, indecisiveness

• Recurrent thoughts of death, suicidal ideation

At least five out of nine symptoms must be present over a two week period and one symptom must be depressed mood or anhedonia.

In major depressive disorder with psychotic features, delusions and/or hallucinations are present along with the depression symptoms.

Psychiatry INI-CET 2021

Question 1:

A person often feels that his hands are contaminated and he is forced to wash his hands. Recently, he feels this repetitive, distressing thought of repetitive hand washing has begun affecting his performance. Which of the following is the best treatment option for this patient?

a) DBT

b) Exposure and response prevention

c) Systematic desensitization

d) CBT

Question 2:

Match the following

a) A-3,B-4,C-1,D-2

b) A-4,B-3,C-1,D-2

c) A-3,B-4,C-2,D-1

d) A-2,B-1,C-3,D-4

Question 3:

Which of the following is true regarding nicotine substitution therapy?

a) Preferably given by gastrointestinal route

b) Varenicline comes with a blackbox warning of cardiovascular death

c) There should be a 15 minute gap between nicotine gum and coffee/soda/acidic food as they decreases its absorption

d) Nicotine chewing gum is better for constant use as it gives 25% higher blood level than lozenges

Question 4:

Match the following

a) A-4, B-2, C-3, D-1

b) A-3, B-4, C-1, D-2

c) A-1, B-4, C-2, D-3

d) A-2, B-4, C-1, D-3

Question 5:

Which of the following is not classified as OCD as per ICD 11?

a) Trichotillomania

b) Body dysmorphic disorder

c) Hypochondriac disorder

d) PTSD

Question 6:

Which of the following is a true statement about obsessive compulsive disorder (OCD)?

a) Atypical antipsychotics is the first line treatment

b) Prevalence is 7 -10 percent in general population

c) Depression is a common comorbidity

d) Contamination is uncommon obsession

Question 7:

Which of the following is true statement about Korsakoff syndrome?

a) Triad of confusion, ophthalmoplegia, and ataxia

b) Loss of long term memory but intact personality

c) Anterograde amnesia with loss of recent memory

d) Loss of implicit memory and immediate recall

Answer Key

Question No.	Correct Option
1	b
2	c
3	c
4	d
5	d
6	c
7	c

Detailed Explanations

Solution to Question 1:

A patient presenting with an obsession with contamination and repetitive handwashing is suggestive of classical obsessive-compulsive disorder (OCD) and the best treatment option for this patient is exposure and response prevention (ERP).

First line management of OCD in adults can include pharmacotherapy or psychotherapy. Psychotherapy is primarily based on exposure and response prevention.

Exposure and response prevention is a form of exposure therapy that is included under behavioral therapy. It is a more specific treatment modality for OCD than CBT. It refers to exposing the patient to OCD-inducing situations and preventing them from exhibiting the usual compulsive responses. The level of exposure and control of response to it, is gradual and is decided by the therapist and the patient. The dose and duration of exposure and response prevention therapy should be sufficient for it to be effective.

Cognitive therapies aim to address distortions in thought like increased responsibility to clean hands by replacing them with adaptive thinking patterns.

Pharmacotherapy consists of SSRIs and clomipramine. Pharmacotherapy is preferred in patients with severe symptoms or with co-morbid depression.

Combined therapy with Pharmacotherapy+ Psychotherapy are effective in patients not responding to either of them alone.

Somatic therapies like targeted neurosurgical lesions to cortico-striatal-thalamic cortical tracts (CSPC) and deep brain stimulation have proven efficacious in treatment-refractory patients.

Cognitive behavioral therapy (Option D) with ERP for OCD is preferred in children and adolescents. Cognitive-Behavioral therapy comprises of a number of treatment patterns that include both cognitive and behavioral interventions. Cognitive therapy aims to correct maladaptive beliefs and behavioral therapy facilitates better functioning by reducing symptoms. Broadly, it is used to treat depression, generalized anxiety disorder, post traumatic stress disorder, panic disorder, eating disorders, substance use disorder, obsessive compulsive disorder, insomnia, smoking, low back pain.

Other options:

Option A: Dialectical behavioral therapy (DBT) is a long-term behavioral intervention that is used in borderline personality disorder and to reduce suicidal behavior in adults.

Option C: Systemic desensitization is counterconditioning the patient by approaching the feared situation gradually, in a relaxed state. It is useful in phobias, obsessions, compulsions, and some sexual disorders. It includes relaxation training, hierarchy construction and desensitization of the stimulus.

Solution to Question 2:

The correct option is A-3,B-4,C-2,D-1.

• Depression - Difficult to diagnose due to the shell of physician's pride

• Suicide - Women physicians, substance abuse, relationship troubles

• Burnout - State of emotional and physical exhaustion caused by prolonged stress, relieved after relaxation

• Overworking - The feeling goes away as soon as you leave the hospital

Depression in doctors:

• Rates of depression are high in doctors.

• It is difficult to diagnose depression in doctors as they won't seek medical help because of the fear of being judged personally and professionally.

• Depression if untreated is a significant risk factor for suicide.

Suicide in doctors:

The risk of suicide is more among female physicians. Risk factors for suicide can be:

• Personal factors: Underlying psychiatric disorders, relationship troubles, stress, trauma, etc.

• Structural factors: It refers to the condition in which the doctors work. These include increased managerialism, paperwork, rushed patient interaction, inability to time off or rest, work pressure, etc.

• Endemic factors: Exposure to trauma and deaths.

Burnout:

It is classified in ICD 11 as an occupational phenomenon. It is defined as a syndrome resulting from chronic workspace stress that has not been managed successfully. It has three dimensions:

• having a feel of energy depletion or exhaustion

• Feeling of negativism or cynicism related to the job, or having increased mental distance from the job

• decreased professional efficacy.

It is a phenomenon related to occupational context and is not classified as a medical condition.

Solution to Question 3:

When a patient is using nicotine gum for nicotine substitution therapy, there should be a gap between nicotine gum and coffee/soda/acidic food as it decreases its absorption.

Treatment of nicotine addiction

1. Behavioral therapy- Elimination of the factors that favor smoking in the patient's environment, skill training, etc.

2. Pharmacotherapy used in the treatment of nicotine addiction are:

• Nicotine replacement therapy- It is used to reduce the symptoms of nicotine withdrawal without the use of tobacco. It is given for the initial maintenance period followed by a reduction period both lasting for about 6-12 weeks. It is available in various forms like

• Nicotine gum- It releases nicotine by chewing and buccal absorption. Smokers can use 1-2 gums per hour and a maximum of 24 gums per day. Acidic beverages like coffee, tea, soda, and juice should not be used during gum use because they decrease absorption. Poor compliance is a drawback.

• Nicotine lozenges- This is especially useful for people who smoke immediately after awakening. Users must suck the lozenges until it is completely dissolved and must not swallow it. It provides the highest level of nicotine compared to other nicotine replacement therapies.

• Nicotine patches- It provides continuous delivery of nicotine and is available as 16 or 24-hour tapering preparations. It has good compliance.

• Nicotine nasal spray- Provides nicotine concentration in the blood similar to smoking. Major adverse effects are rhinitis, watering eyes, and coughing

• Nicotine inhaler - Even though it was designed to deliver the nicotine to the lungs, the actual absorption occurs in the upper throat.

• Non-nicotine medications

• Bupropion is an antidepressant medication. It has dopaminergic and adrenergic actions.

• Varenicline is a partial agonist of alpha-4 beta-2 nicotinic receptor. It blocks nicotine from binding to the receptor and decreases the rewarding aspects of cigarette smoking. At the same time, it also stimulates the receptor and reduces withdrawal and cravings. There were concerns about its neuropsychiatric and cardiovascular side effects, but according to recent studies varenicline is considered a safe drug.

• Nortriptyline - Second-line drug

• Clonidine- It decreases the sympathetic activity from locus ceruleus and is useful in controlling the withdrawal symptoms.

Nicotine addiction:

It acts on the nicotinic subtype of acetylcholine receptors. Its addictive properties are thought to be due to the activation of the dopaminergic pathway in the ventral tegmental area. It causes improved attention, problem-solving ability, learning, and lifts the mood of the users.

Withdrawal symptoms include intense craving for tobacco, irritability, difficulty concentrating, drowsiness, altered sleep, decreased heart rate, and blood pressure, decreased motor performance, and increased muscle tension.

Solution to Question 4:

The correct option is A-2,B-4,C-1,D-3.

• Twilight state - Impaired level of consciousness

• Oneiroid state - Oneirophrenia

• Oligophrenia - Subnormal intelligence

• Paranoid ideation - Persecution

Twilight state - It refers to a state of disturbed consciousness along with hallucinations.

Oneiroid state or oneirophrenia - It refers to dream-like delusional derangement of consciousness in which the patient is not fully oriented in time and place. The characteristic feature of this state is the presence of kaleidoscopic quality of the dream-like experiences where reality, illusions, and hallucinations are merged into one. It is associated with catatonic disturbances. The term oneiroid schizophrenia refers to patients engaged in their hallucinations to an extent where there is the exclusion of involvement in the real world.

Oligophrenia - It refers to a group of mental states characterized by subnormal levels of intellectual activity. It is either congenital or acquired during early infancy. It is associated with other cognitive disturbances, affective disorders, and volitional disorders.

Paranoid ideation: It refers to thinking that is dominated by persecutory, suspicious, or grandiose contents. The thought content is less than that for delusions. Paranoid delusions refer to persecutory delusions and delusions of grandeur, control, and reference.

Solution to Question 5:

Post-traumatic stress disorder(PTSD) is not classified as OCD under ICD 11.

Obsessive-compulsive or related disorders as per ICD 11

It refers to a group of disorders that are characterized by repetitive thoughts and behaviors that are similar in etiology and important diagnostic validators. Important features of these disorders are obsessions, intrusive thoughts, and preoccupations. It includes:

• Obsessive-compulsive disorder

• Body dysmorphic disorder

• Hypochondriasis

• Olfactory reference disorder

• Hoarding disorder

• Body focused repetitive behavior disorders (Skin pricking, hair pulling - Trichotillomania)

Important points regarding ICD-11

The ICD 11 was released by the WHO in June 2018 and will come into effect on 1 January 2022.

Its designations for antibiotic resistance align with the Global Antimicrobial Resistance System (GLASS).

New chapters on

• Traditional medicine

• Sexual health: Gender incongruence has been moved from mental health to sexual health.

• Immune system disorders

• Diseases of the Blood and Blood forming Organs

• Sleep-Wake Disorders

• Extension codes

Gaming disorder has been added under Addictive disorders.

Solution to Question 6:

Obsessive-compulsive disorder (OCD) is commonly associated with depression. The lifetime prevalence of the major depressive disorder in patients with OCD is 67 percent. Other common psychiatric comorbidities in patients with OCD are social phobia, alcohol use disorders, panic, and personality disorders.

Obsessive-compulsive disorder (OCD) comprises symptoms that include intrusive thoughts, preoccupations, and compulsions. These recurrent obsessions or compulsions cause severe distress to the patient. The patient might have an obsession, a compulsion, or both.

An obsession is a thought, feeling, or sensation that is repetitive and intrusive. A compulsion is a conscious, recurrent behavior, eg: counting, checking, and are done to reduce the anxiety associated with the obsession. However, this act may not reduce the anxiety and might increase the anxiety even more.

Symptom patterns:

• Contamination - This is the most common symptom pattern of OCD. The patient feels that the objects are contaminated. They repeatedly wash their hands or avoid touching objects which are thought to be contaminated. In extreme cases, patients may rub the skin off their hands due to excessive hand washing or may refuse to leave the house in fear of germs.

• Pathological doubt - This is the second most common symptom pattern. The patient presents with an obsession with doubt and a compulsion to check. This obsession arises due to possible danger or violence (e.g., forgetting to turn off the stove or not locking a door) which is followed by multiple trips to check, for example, if the stove is turned off, if the door is locked, etc.

• Intrusive thoughts -This is the third most common symptom pattern. These patients have intrusive obsessional thoughts without compulsion. These thoughts could be sexual or an aggressive act that the patient doesn't agree. The patients might report themselves to the police or confess to a spiritual leader.

• Symmetry - This is the fourth most common pattern which can lead to a compulsion of slowness. Patients can take hours to eat a meal or shave their faces.

Diagnosis:

• Presence of obsessions, compulsions or both

• Symptoms are time-consuming

• Symptoms not related to substance abuse

• Not better explained by other mental disorders

Treatment of OCD includes:

• Pharmacotherapy: Selective serotonin reuptake inhibitors (SSRI) or clomipramine

• Behavioral therapy: Approaches include:

• exposure and response prevention

• desensitization

• thought stopping

• flooding

• implosion therapy

• aversive conditioning

A combination of behavioral and pharmacotherapy is most effective.

Other options

Option A - Atypical antipsychotics are not the first-line in the treatment of OCD. They can be given only when SSRIs and clomipramine (tricyclic antidepressant) are not effective.

Option B - The prevalence of OCD among the general population is 2-3 percent and not 7-10 percent.

Option D - Contamination is the most common obsession.

Solution to Question 7:

In Korsakoff's psychosis, there is anterograde amnesia with loss of recent memory.

Korsakoff's syndrome is a chronic amnestic syndrome that can follow Wernicke's syndrome. It is caused by thiamine deficiency that is associated with chronic alcohol abuse. The typical feature of this disease is anterograde amnesia with loss of recent memory in an alert and responsive patient. Patients with Korsakoff's syndrome also typically have a change in personality, with a lack of initiative, reduced spontaneity, and a lack of interest.

The treatment is with 100 mg of thiamine orally for 3-12 months. Despite adequate treatment and nutritional support, very few of the patients fully recover. With treatment, patients may remain amnestic for up to 3 months, and then improve gradually during the next year.

Prolonged use of alcohol causes alcohol-induced persisting amnestic disorder with loss of short-term memory. When the symptoms are acute, it is called Wernicke's encephalopathy and when chronic, it leads to Korsakoff's syndrome. These conditions are caused due to thiamine deficiency and when they occur together, they are called Wernicke-Korsakoff syndrome.

Other options

Option A -Triad of confusion, ataxia, and ophthalmoplegia is seen in Wernicke's encephalopathy. There may also be vestibular dysfunction. Wernicke's encephalopathy is completely reversible. The treatment is parenteral administration of 100 mg thiamine for 1-2 weeks.

Option B - Personality is affected in Korsakoff's syndrome. Recent memory is affected more compared to long-term memory.

Option D - Implicit memory refers to memory that does not require conscious recollection. Implicit aspects of memory are preserved, however, some studies conclude that implicit memory may be intact or impaired depending on the specific task parameters.

Psychiatry NEET 2021

Question 1:

During psychotherapy, the therapist had mixed conscious and unconscious feelings towards his patient. This is known as_____.

a) Countertransference

b) Transference

c) Dissociation

d) Preoccupation

Question 2:

A patient stopped alcohol consumption for 3 days and presented with irritability, disorientation, paranoid delusions, agitation, visual hallucinations, and altered sensorium. What is the likely diagnosis in this case?

a) Delirium tremens

b) Wernicke's encephalopathy

c) Korsakoff psychosis

d) Alcohol induced psychosis

Question 3:

A teacher demonstrated the steps of handwashing to her students. Students also performed the same activity later at school and at home. What is the type of learning in this case?

a) Cognitive

b) Psychomotor

c) Affective

d) Affective and cognitive

Question 4:

Which of the following is a formal thought disorder?

a) Obsession

b) Derailment

c) Delusion

d) Thought insertion

Question 5:

A 16-year-old girl presented with overfamiliarity, flight of ideas, elevated mood, increased sexual desire, and pseudohallucination. What is the likely diagnosis?

a) Mania

b) Hypomania

c) Cyclothymia

d) Schizomania

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	b
5	a

Detailed Explanations

Solution to Question 1:

In the given clinical scenario, the therapist has mixed conscious and unconscious feelings towards his patient. This is known as counter-transference.

Counter-transference is the positive and negative feelings/emotions expressed by healthcare professionals towards their patients. It is the process of the physician unconsciously displacing onto the patient patterns of behaviors or emotional reactions as if he/she were a significant figure from earlier in the physician's life. Psychiatrists should be alert to signs of countertransference issues like missed appointments by the psychiatrist, boredom, or sleepiness during a session. Supervision and personal therapy can help the psychiatrist recognize and deal with these issues.

Transference is the positive and negative feelings expressed by the patients towards the healthcare professional. It is the process of the patient unconsciously displacing onto individuals patterns of behavior and emotional reactions that originated with significant figures from the past (generally in childhood). This displacement is onto the psychiatrist in a clinical scenario, who is a parent surrogate. The patient may be angry, hostile, demanding, or obedient to an extreme degree because of former relationships and patterns of behaviors. Failure to recognize this can make the psychiatrist react to the patient's behavior like a personal attack.

These are not defense mechanisms but are commonly encountered during clinical practice or while giving psychotherapy. In an interview, transference and counter-transference are very significant expressions of unconscious processes.

Option C: Dissociation is an immature defense mechanism. In this, involuntary splitting/suppression of mental functions occurs from the rest of the personality in a way that allows the expression of forbidden unconscious impulses without having a sense of responsibility for actions.

Solution to Question 2:

The given clinical scenario of a patient who has not consumed alcohol for the past 3 days with disorientation, paranoid delusions, visual hallucinations, and altered sensorium is suggestive of delirium tremens.

Delirium tremens is a medical emergency that occurs during alcohol withdrawal. It is an acute condition and usually develops within 72 hours after the last drink. It has symptoms of both delirium and alcohol withdrawal. The clinical features of delirium tremens are:

• Altered sensorium with disorientation

• Perceptual distortions, most frequently tactile or visual hallucinations

• Autonomic hyperactivity with tachycardia, fever, anxiety, insomnia, hypertension. And sweating

• Fluctuating levels of psychomotor activity, ranging from agitation to lethargy

• Altered sleep-wake cycle

Other options:

Option B - Wernicke's encephalopathy (alcoholic encephalopathy) is associated with thiamine deficiency. Alcohol abuse may lead to thiamine deficiency because of poor nutritional habits, or malabsorption. It is an acute neurological disorder characterized by:

• Global confusion

• Ocular abnormalities like horizontal nystagmus, gaze palsy, anisocoria, and sluggish pupillary reflex

• Ataxia - mainly affecting the gait

• Vestibular dysfunction

It may resolve with thiamine therapy or progress to Korsakoff's syndrome.

Option C - Korsakoff's psychosis can follow Wernicke's encephalopathy due to thiamine deficiency. It is a chronic amnestic syndrome that is characterized by the following clinical symptoms in an intact and responsive patient:

• Impaired mental syndrome, especially recent memory

• Anterograde amnesia

• Confabulation

Option D - Alcohol-induced psychosis is characterized by the development of auditory hallucinations or paranoid delusions in the context of a heavy drinking or alcohol withdrawal. These patients have a clear sensorium, unlike the patients having delirium tremens.

Note: Tremulousness is the first alcohol withdrawal symptom.

Solution to Question 3:

The type of learning seen in this case is psychomotor learning.

The psychomotor domain includes learning by observation, physical movements, and the usage of motor skill areas of the brain. The development of these skills requires practice, depending upon the complexity of the task. It is measured based on speed, precision, distance, and procedures in execution.

The three domains in learning are the cognitive, psychomotor, and affective domains. They are described as follows.

Cognitive learning includes the domain of knowledge. Bloom's taxonomy described six levels of cognitive skills. This was revised to include types of knowledge and cognitive processes. The cognitive processes include remembering, understanding, applying, analyzing, evaluating, and creating. Eg. Solving multiple-choice questions in the NEET PG entrance exam involves the usage of the cognitive domain.

Psychomotor learning includes the domain of skills. It includes imitation, manipulation, precision, articulation, and naturalization. Eg. Learning how to insert an IV cannula by observing the residents and nurses during the internship involves the usage of the psychomotor domain.

Affective learning includes the domain of attitude and values. It includes receiving, responding, valuing, organizing, and characterization by value. Eg. Communicating with patients with empathy, responding to their concerns, and interacting with the relatives of patients involves the usage of the affective domain.

Integration of all these three methods in teaching and learning methodology leads to a better understanding of the subject.

Solution to Question 4:

Derailment is a formal thought disorder.

It is a breakdown in both the logical connection between ideas and the overall sense of being goal-oriented. It is generally identified in schizophrenic patients as a generalized loosening of associations.

For example, a person with derailment thought disorder might jump from talking about rabbits to the hair on their head to your sweater. They talk in chains of semi-related ideas. Their communication style has a generalized loosening of associations. Their ideas often fall further and further from the topic.

Formal thought disorders are:

• Circumstantiality is the overinclusion of trivial or irrelevant details that impedes the sense of getting to the point.

• Clang association is an association of thoughts with the sound of words rather than by their meaning (e.g., through rhyming).

• Derailment is the breakdown in both the logical connection between ideas and the overall sense of goal-directedness. The words make sentences, but the sentences do not make sense.

• Flight of ideas is a succession of multiple associations so that thoughts seem to move abruptly from idea to idea, through rapid, pressured speech.

• Neologism is the invention of new words or phrases or the use of conventional words in idiosyncratic ways.

• Perseveration is the repetition of out-of-context words, phrases, or ideas.

• Tangentiality is responding to a question, that is appropriate to the general topic without actually answering the question.

• Thought blocking is a sudden disruption of thought or a break in the flow of ideas.

Other options:

Option A - An obsession is a recurrent and intrusive thought, feeling, idea, or sensation.

Option C - A delusion is defined as a false belief based on incorrect inference about external reality that firmly sustains despite evidence to the contrary.

Option D - Thought insertion is a type of delusion where the patient believes that thoughts are not his/her own and have been inserted into his/her mind.

Solution to Question 5:

In the given clinical scenario, history of overfamiliarity, elevated mood, flight of ideas, and increased sexual desire are suggestive of mania.

The diagnostic criteria for mania are:

A. Distinct period of abnormally and persistently elevated/irritable mood, and abnormally and persistently increased activity/energy, lasting at least 1 week.

B. During the period of mood disturbance and increased energy/activity, three (or more) of the following symptoms are present to a significant degree:

1. Increased self-esteem or grandiosity

2. Decreased need for sleep

3. More talkative than usual or a pressure to keep talking

4. Flight of ideas

5. Distractibility

6. Increase in goal-directed activity or psychomotor agitation

7. Excessive involvement in activities that have a high potential for painful consequences (unrestrained buying sprees, sexual indiscretions)

C. The mood disturbance is severe enough to cause marked social or occupational impairment or to necessitate hospitalization.

D. The episode is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication, other treatment) or to another medical condition.

Other options:

Option B - Hypomania is a distinct period of abnormally and persistently elevated/irritable mood, abnormally and persistently increased activity/energy, lasting for at least 4 days. It is similar to a manic episode but it is not severe enough to cause impairment in social or occupational functioning. No psychotic features are present.

Option C - Cyclothymia is a mild, chronic mood disorder with numerous depressive and hypomanic episodes over the course of at least 2 years.

Option D - Schizomania is an old term in DSM III, used to describe schizophrenic and manic symptomatology occurring in one episode, classified under schizoaffective disorder. Schizoaffective disorder is characterized by active phase symptoms of schizophrenia, like delusions, hallucinations, disorganized speech, grossly disorganized behavior, or negative symptoms. Additionally, there is a presence of manic/depressive syndrome that is not brief relative to the duration of the psychosis. The delusions or hallucinations are present for at least 2 weeks without coexisting mood symptoms.

Psychiatry INI-CET 2022

Question 1:

Which of the following is false about bipolar disorder?

a) It most commonly affects ages 35-45 years

b) Type 1 is common in both sexes

c) Prevalence in general population is 1-2%

d) Attempt to commit suicide is seen around 15% of people

Question 2:

On performing polysomnography in a patient, the waves of EOG, EEG, and EMG respectively from above downwards are seen below. Which stage of sleep does the marked area represent?



a) REM sleep

b) NREM I sleep

c) NREM II sleep

d) NREM III sleep

Question 3:

A patient diagnosed with schizophrenia was started on risperidone. After taking the medication last night, the patient rushes to OPD today with complaints of upwards rolling of his eyes. What is the management of the condition?

a) Lorazepam

b) Promethazine

c) Olanzapine

d) Stop antipsychotic

Question 4:

Which of the following disease is exclusively seen in females?

a) Rett syndrome

b) Asperger syndrome

c) Selective mutism

d) Atypical autism

Question 5:

Match the following:

a) 1-D, 2- B, 3-A, 4-C

b) 1-C, 2-B, 3-D, 4-A

c) 1-C, 2-D, 3-B, 4-A

d) 1-D, 2-C, 3-B, 4-A

Question 6:

A 14-year-old girl presented with sudden-onset blindness for the past 4 hours. However, on history taking, it is noted that she is not concerned about it. However, she is concerned that her mother passed away recently and that she should have spent more time with her. Physical examination findings are normal. Which of the following is true about the condition?

a) In children, equally among males and females.

b) In adults, equally among males and females.

c) In adults, occurs more in males than in females.

d) In children, occurs more in females than in males.

Question 7:

What is the dose range of lithium in mEq/L for acute attacks of mania and long-term maintenance?

a) 0.5 to 1 mEq/L

b) 0.1 to 0.5 mEq/L

c) 0.6 to 1.5 mEq/L

d) 2 to 3 mEq/L

Question 8:

A patient who is a chronic alcoholic presents with seizures and altered sensorium. He has been drinking for the past 10 years and takes 24 units per day. His last drink was 2 days back. He is found to have auditory hallucinations and secondary delusions. What will you use for the management of this patient?

a) 1,4

b) 1,2,4

c) 1,3,4

d) 1,2

Question 9:

Match the following:

a) 1-B;2-A;3-D;4-C

b) 1-C;2-A;3-D;4-B

c) 1-C;2-D;3-A;4-B

d) 1-C;2-B;3-D;4-A

Question 10:

Which of the following is the most appropriate definition of stress?

a) External pressure

b) Mental tension

c) It is the body's response to anything that requires attention or action causing physical, emotional, or psychological strain.

d) Internal strain

Question 11:

A pregnant mother was on SSRIs. The risk of which of the following is increased in the newborn?

a) A, C

b) A, B, C, D

c) B, C, D

d) A, B, C

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	a
5	c
6	d
7	c
8	d
9	b
10	c
11	b

Detailed Explanations

Solution to Question 1:

The false statement among the following is option A. The age of onset for bipolar type I disorder ranges from, children of age 5 or 6 years to adults of age 50 years or older, with a mean age of 18

years. The mean age of onset for bipolar type II is 20 years.

Patients with bipolar disorder can present with both manic and depressive episodes or with manic episodes alone. The bipolar disorder presented with mania and depression for at least a week is known as bipolar type 1 disorder. Whereas, if it is presented with hypomania and depression it is known as bipolar type 2 disorder.

The lifetime prevalence of bipolar type 1 disorder is 0 to 2.4%, whereas bipolar type 2 disorder is 0.3-4.8%. Bipolar type 1 disorder has an equal prevalence among men and women. Suicide attempt is seen in around 15% of people with bipolar disorder.

A manic episode is defined as a persistent elevated, expansive, or irritable mood for at least 1 week which requires hospitalization. A hypomanic episode is also similar to a manic episode except it doesn't cause social and occupational functioning impairment. It lasts for at least 4 days and doesn't show psychotic symptoms.

A person is said to be depressed if he has at least four of the following symptoms for at least 2 weeks:

• Change in appetite and weight

• Change in sleep and activity

• Lack of energy

• Guilty feeling

• recurring thoughts about death or suicide.

Lithium is used for the pharmacological management of bipolar disorder.

Solution to Question 2:

REM sleep is represented by the marked area in the given polysomnography.

The three zones in the given image represent EOG, EEG, and EMG, respectively. In the marked area,

• EOG shows Positive activity

• EEG shows Mixed waves

• EMG shows Minimal activity

This is characteristic of REM sleep.



The following table helps in identifying the state of wakefulness and sleep from the polysomnograph :

	EEG	EOG	EMG
Active state	Fast	+	+
REM sleep	Mixed	+	-
NREM sleep	Slow	-	+



Solution to Question 3:

The given clinical scenario is suggestive of oculogyric crisis and promethazine is used for the management.

Oculogyric crisis refers to the spasm of the extraocular muscles, usually causing upward and outward deviation of the eyes. It is a severe form of acute dystonia, an extrapyramidal side effect of antipsychotics like risperidone. It can be treated by drugs with antimuscarinic activities like promethazine.

Dystonia is acute in onset and occurs within 1-5 days of starting treatment. It is more common in the younger population. The proposed mechanism of action is D2 blockade in the nigrostriatal pathway. There is typical involvement of head and neck muscles and presents in a variety of forms, such as torticollis, trismus, mouth opening dystonia, grimacing, dysarthria, oculogyric crisis, blepharospasm, and swallowing difficulties.

Risperidone is an atypical antipsychotic with D2 blocking but strong 5HT2 antagonistic activity. Its main side effects include weight gain, hyperglycemia, hyperprolactinemia, dyslipidemia, fatigue, etc. At low doses, extrapyramidal side effects are less, but can be significant in higher doses. Extrapyramidal side effects include;

• Parkinsonism: Features are bradykinesia, rigidity, variable tremor, masked facies, and shuffling gait.

• Acute dystonia: Which causes spasms of muscles of the tongue, face, eyes, neck, back, etc.

• Neuroleptic malignant syndrome: Extreme rigidity, fever, unstable blood pressure,myoglobinemia, etc.

• Rabbit syndrome: Perioral tremors occur as a delayed variant of parkinsonism.

Other options:

Option A: Lorazepam is a benzodiazepine. Its adverse effects include sedation, light-headedness, confusional state, weight gain, etc.

Option C: Olanzapine is also an atypical antipsychotic. Its adverse effects are dry mouth, constipation, weight gain, impaired glucose tolerance etc.

Solution to Question 4:

Rett syndrome is exclusively seen in females. It is caused by X-linked dominant inheritance and hence affects only females, as most males die in utero.

Rett syndrome is a form of autism spectrum disorder. The child is normal at birth, but by 1 year of age deterioration in communication skills, motor behaviour, and social functioning are noticed. Symptoms are ataxia, loss of speech, teeth grinding, and facial grimacing. When the child is awake, intermittent hyperventilation and disorganized breathing pattern are observed. Stereotypical hand movements like hand-wringing, are characteristically seen. Progressive gait disturbance and seizures are seen. Decreased pigmentation of the substantia nigra, along with cerebral atrophy occurs, implying an abnormal dopaminergic nigrostriatal system.

Other options:

Option B: Asperger syndrome is a version of autism spectrum disorder. It is characterized by limited social interaction and restricted interest and behaviour. There is no delay in language

development.

Option C: Selective mutism is a social anxiety disorder. The child hesitates to speak in one or more social settings, mostly the school.

Option D: Atypical autism is autism, atypical either in the age of onset or in symptomology or both.

Solution to Question 5:

The correct option is 1-C, 2-D, 3-B, 4-A

Impulse control disorders are characterized by the repeated failure to resist an impulse, drive, or urge to perform an act that is rewarding to the person, at least in the short-term, despite consequences such as long-term harm either to the individual or to others; marked distress about the behavior pattern; or significant impairment in personal, family, social, educational, occupational, or other important areas of functioning.

It includes

• Pyromania

• Kleptomania

• Compulsive sexual behavior disorder

• Intermittent explosive disorder

Kleptomania is the repeated failure to resist impulses to steal objects that are of no value to the person. The individual can usually afford the object or does not need it, but it is the act of stealing itself that is the goal. The stolen objects are usually given away, returned, or kept hidden. The act is not planned and is carried out alone.

Pyromania is the deliberate setting of fires on more than one occasion.

Pyromania is significantly associated with the following:

• Substance abuse disorder (especially alcoholism).

• Affective disorders (depression or bipolar)

• Other impulse control disorders, such as kleptomania in female firesetters.

• Personality disturbances, such as borderline personality disorders.

• Attention-deficit/hyperactivity disorder and learning disabilities in childhood pyromania.

Dipsomania is an irresistible impulse to drink alcohol at a periodic intervals.

Mutilomania is an irresistible impulse to mutilate animals.

Solution to Question 6:

The above scenario is suggestive of conversion disorder. Option D is a true statement regarding the above condition as it is more commonly seen in females.

Conversion disorder is also called functional neurological symptom disorder, and there is no bodily cause involved in this condition. A conversion disorder is an acute and temporary loss or alteration in motor or sensory function triggered by psychosocial stress. Most of the patients have a history of traumatic events in childhood. The symptoms or deficits of conversion disorder are not intentionally produced but are a symbolic representation of an unconscious conflict (primary gain). It is more commonly seen in females.

Primary gain refers to keeping internal conflicts outside of awareness and the relief from stress. Secondary gain refers to the attention gained by the patient (from bystanders) as a result of being sick. If the secondary gain is intentional, then it would lead to the diagnosis of malingering. A related phenomenon is la belle indifference, in which the patient is not concerned about the serious symptoms.

Symptoms of conversion disorder:

• Mutism, blindness, and paralysis are the most common symptoms.

• Sensory system: Anesthesia, paresthesia, deafness, blindness, and tunnel vision

• Motor system: Abnormal movements, gait disturbance (astasia abasia), weakness, paralysis, tics, jerks, etc.

• CNS: Pseudo-seizures

Treatment: Hypnosis, anxiolytics, and behavioral relaxation exercises are effective. Psychoanalysis and insight-oriented psychotherapy help in the resolution of the symptoms.

Solution to Question 7:

Dose range of lithium in mEq/L for acute attacks of mania and long-term maintenance is 0.6 to 1.5 mEq/L

Lithium has a narrow therapeutic index (0.8 to 1.2 mEq/L) and, therefore, requires monitoring to suspect toxicity. The toxic effects of lithium usually start manifesting when the serum level crosses 1.5 mEq/L.

The range of 1.0-1.5 mEq/L is only favoured for acute attacks of mania with lower doses of 0.6-1.0 mEq/L for long term maintainance.

Acute intoxication is characterized by

• Vomiting

• Profuse diarrhea

• Coarse tremors

• Ataxia

• Coma

• Convulsions.

A fine action tremor is common in patients taking lithium at therapeutic doses and can develop at any time during treatment. Therapeutic lithium tremor is generally symmetric, limited to the arms, and nonprogressive. The tremor from lithium toxicity is coarser and more disabling, may

also affect the legs, and occurs with other symptoms of toxicity.

Serious effects of lithium toxicity will involve confusion, hyperreflexia, and seizures. Motor and cognitive adverse effects can become irreversible, with persistent cerebellar tremor being the most common.

Solution to Question 8:

The given clinical scenario is suggestive of delirium tremens. Management of this patient involves the administration of diazepam and thiamine.

Delirium tremens is the most severe form of alcohol withdrawal syndrome. It occurs usually within 3–4 days of alcohol withdrawal.

Characteristic features of delirium tremens:

• Clouding of consciousness, with disorientation to time and place

• Poor attention and distractibility

• Visual hallucinations and illusions, which are often vivid and very frightening

• Autonomic disturbances like tachycardia, fever, hypertension, diaphoresis, and pupillary dilatation

• Psychomotor agitation and ataxia

Benzodiazepines such as diazepam (intravenously) or chlordiazepoxide (orally every 4 hours) are given. It is also recommended to administer thiamine in patients with alcohol-related disorders to prevent Wernicke-Korsakoff syndrome.

Acute alcohol intoxication can be treated with the evacuation of stomach and bowel and gastric lavage with an alkaline solution. Administration of 1 L of normal saline with 10% glucose, 100 mg thiamine, and 15 U insulin may be useful. If the coma progresses deeper, stimulants like strychnine and caffeine can be used. Patients should be started on artificial respiration if there is respiratory depression or difficulty breathing.

Chronic alcohol intoxication can be treated with disulfiram (daily dose of 250 mg OD), conditioned reflex treatment, hypnosis, and psychotherapy.

According to the Motor Vehicle Amendment bill (2019), whoever while driving, or attempting to drive, a motor vehicle, has, in his/her blood, alcohol exceeding 30 mg/100 mL of blood, detected in a test by a breath analyzer, is punishable by law.

Naltrexone is an opioid receptor antagonist which is used along with cognitive behavioral therapy in cases of opioid withdrawal and alcohol withdrawal. It reduces craving and consumption, also reduces the severity of withdrawal symptoms.

Flupenthixol is an antipsychotic antagonist of both D1 and D2 dopamine receptors. The depot injections are used for the management of schizophrenia and depression.

Solution to Question 9:

The correct answer is option B: 1-C;2-A;3-D;4-B

Paraphilias or perversions are sexual stimuli or acts that are deviations from normal sexual behaviors but are necessary for some people to experience arousal and orgasm.

Eonism (also known as transvestism) is a sexual perversion, where a person's whole personality is dominated by a desire to be identified as a member of the opposite sex. This is usually seen in males who obtain sexual pleasure by wearing female clothes. There is no hormonal/genital abnormality.

Exhibitionism is the urge to expose genitals to a stranger or unsuspecting person.

Frotteurism refers to the sexual gratification achieved by touching or rubbing against a non-consenting person. This behavior often occurs in busy and crowded places. It is predominantly described in men.

Necrophilia is an obsession with obtaining sexual gratification from dead bodies.

1. Eonism	C. Sexual pleasure by wearing clothes of the opposite sex
2. Exhibitionism	A. Exposure of one's genitals to an unsuspecting stranger
3. Frotteurism	D. Sexual gratification by rubbing private parts against another person
4. Necrophilia	B. Sexual gratification by intercourse with dead bodies

Solution to Question 10:

According to the WHO, stress can be defined as any type of change that causes physical, emotional, or psychological strain. Stress is your body's response to anything that requires attention or action. Everyone experiences stress to some degree.

Solution to Question 11:

The newborn of a pregnant mother on SSRI has an increased risk for A, B, C and D. SSRIs like sertraline are the most commonly used antidepressants to treat depression in pregnant women.

SSRIs and maternal depression have both been found to cause a low APGAR score.

The development of ADHD in the newborn has been found to be commonly associated with untreated maternal depression. But studies regarding SSRIs having a causal association with the development of ADHD have been inconclusive. This can be contributed to maternal depression being a confounder.

When used in the second half of pregnancy, SSRIs have been associated with an increased risk of precipitating persistent pulmonary hypertension in the newborn.

There has been evidence of a transient delay in developmental milestones in patients who were exposed to SSRIs during pregnancy. Most studies have observed that the development returns to normal after some time. But if the mother is severely depressed, delayed developmental milestones have also been attributed to the mother's disinterest in the infant.

The other side effects of SSRIs include:

System	Side effects of SSRIs
Sexual	Anorgasmia, Decreased libido
Gastrointestinal (5-HT3 receptor)	Nausea, diarrhea, anorexia, vomiting, flatulence, and dyspepsia
Nervous system	Extremelyvivid dreamsor nightmares, Anxiety, Insomnia/sedation, seizures, emotional blunting
Anticholinergic	Dry mouth, constipation, sedation
Cardiovascular	Drug-induced long QT syndrome
Hematological	Easy bruising
Endocrine	Mammoplasia, galactorrhea

Psychiatry NEET 2022

Question 1:

A 16-year-old girl has intense cravings for food. She eats large amounts of food, which is followed by self-induced vomiting. What is the probable diagnosis?

a) Anorexia nervosa

b) Bulimia nervosa

c) Atypical depression

d) Binge eating disorder

Question 2:

A woman, who is 4 days postpartum, presented with tearfulness, mood swings, and occasional insomnia. What is the likely diagnosis?

a) Postpartum depression

b) Postpartum blues

c) Postpartum psychosis

d) Postpartum anxiety

Answer Key

Question No.	Correct Option
1	b
2	b

Detailed Explanations

Solution to Question 1:

The probable diagnosis in the given scenario of a 16-year-old girl with intense food cravings followed by self-induced vomiting is bulimia nervosa.

Bulimia nervosa is an eating disorder, often seen in young women. It is characterized by binge-eating episodes, followed by inappropriate, compensatory measures to stop weight gain, like self-induced vomiting, laxative abuse, diuretics, enema, and emetic abuse. These binge-eating episodes occur at least once a week and are present for at least 3 months. The patient fears gaining weight and has a relentless drive to be thin.

The weight is normal in most patients with bulimia nervosa. However, few patients may either be overweight or underweight. There is a risk of dehydration and electrolyte imbalance.

Treatment modalities are cognitive behavioral therapy, psychotherapy, and antidepressants such as fluoxetine.

Other options:

Option A: Anorexia nervosa is an eating disorder that is often seen between the ages of 10 and 30. Typical features are self-induced starvation, voluntary and unhealthy reduction in weight, intense drive for thinness, and fear of becoming obese. These symptoms are usually present for at least a period of 3 months. Associated features can include depression, anxiety, amenorrhea in females, hypothermia, and bradycardia.

Option C: Atypical depression is a type of depression characterized by mood reactivity, which is brightening of the mood in response to positive events; significant weight gain or increase in appetite; hypersomnia; leaden paralysis; and/or interpersonal rejection sensitivity, resulting in significant□ social or occupational impairment.

Option D: Binge-eating disorder is the most common eating disorder. It is characterized by rapid intake of abnormally large quantities of food over a short period of time, even when not hungry. There are feelings of guilt about the eating episodes, but these patients do not compensate in any way following the episodes of binge.

Solution to Question 2:

Postpartum blues is the most likely diagnosis in a postpartum woman with tearfulness, mood swings, and insomnia from day 4.

Postpartum blues, also known as baby blues, is a transient mood disorder characterized by sadness, tearfulness, confusion, and dysphoria. There are no suicidal thoughts or thoughts of harming the baby. These symptoms typically develop 3-5 days after delivery. They are thought to be due to hormonal changes, the stress of giving birth, and the responsibilities of motherhood. They resolve spontaneously within 10-14 days.

Management involves reassurance, education, and support. The patient must be followed up and assessed if symptoms persist for >2 weeks to evaluate postpartum depression.

Other options:

Option A: When the symptoms of depressed mood, anxiety, insomnia, and weight changes start within 4 weeks as per DSM 5 or 6 weeks as per ICD 11, and the symptoms persist for ≥ 2 weeks, the diagnosis will be postpartum depression. The patient may have occasional thoughts of harming the baby and suicidal thoughts. It can persist for months to years when untreated.

Option C: Postpartum psychosis is a psychiatric emergency that usually occurs within 4 weeks after delivery (DSM 5) or 6 weeks after delivery (ICD 11). The predominant symptoms are

hallucinations, delusions, suicidal thoughts, and thoughts of harming the baby. Psychotic symptoms such as insomnia, restlessness, labile mood, and agitation are seen.
Lithium, antipsychotics, and an antidepressant is the treatment of choice.

Option D: Postpartum anxiety is not defined by DSM-5. However, some platforms might describe postpartum anxiety as excessive or severe worry after childbirth.

Note: Zuranolone is the first-ever oral drug to be approved specifically for the treatment of postpartum depression. It is a neuroactive steroid that enhances the action of GABA in the brain and regulates mood.

Psychiatry NEET 2023

Question 1:

A schizophrenic patient was prescribed drug A after he did not respond to haloperidol and thioridazine. He now presents with excessive salivation, an increase in blood glucose, and hyperlipidemia. What is drug A?

a) Ziprasidone

b) Risperidone

c) Clozapine

d) Aripiprazole

Question 2:

A woman, who is 4 days postpartum, presented with tearfulness, mood swings, and occasional insomnia. What is the likely diagnosis?

a) Postpartum depression

b) Postpartum blues

c) Postpartum psychosis

d) Postpartum anxiety

Question 3:

A 25-year-old female reported symptoms of anxiety, palpitations, sweating, breathlessness, chest pain, and a feeling of impending doom. She recalls having experienced 5-6 similar episodes in the past 6 months, each episode lasting for about 20-30 minutes. What is the likely diagnosis?

a) Depression

b) Panic disorder

c) Generalized anxiety disorder

d) Phobia

Question 4:

A chronic alcoholic is brought to the emergency department with confusion, ataxia, and painful eye movements. The 6th cranial nerve is also involved. What is the likely diagnosis?

a) Wernicke's encephalopathy

b) Korsakoff psychosis

c) Delirium tremens

d) De Clerambault syndrome

Question 5:

A middle-aged man complains of very early ejaculation during intercourse. What is the non-pharmacological management that can be advised to the patient?

a) Cognitive behavioral therapy

b) Exposure and response prevention therapy

c) Squeeze technique

d) Sensate focus therapy

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	a
5	c

Detailed Explanations

Solution to Question 1:

Drug A is clozapine, an atypical antipsychotic used in the treatment of refractory schizophrenia. Excessive salivation, an increase in blood glucose, and hyperlipidemia are side effects of clozapine.

The side effects of clozapine are

• Agranulocytosis

• Weight gain

• Seizures

• Drooling

• Hyperthermia

• Hyperglycemia

• Hypertriglyceridemia

Clozapine causes the highest weight gain among all antipsychotics. Agranulocytosis is a dose-independent side effect and seizures are a dose-dependent side effect.

Other options:

Option A: Ziprasidone - Although it is an atypical antipsychotic, it does not typically cause excessive salivation, significant increases in blood glucose, or hyperlipidemia. The side effects of ziprasidone include orthostatic hypotension, QT prolongation, and minimal weight gain.

Option B: Risperidone - Risperidone is another atypical antipsychotic, but its side effect profile differs from the symptoms presented in the question. Risperidone is more likely to cause hyperprolactinemia, weight gain, sedation, dystonia, and parkinsonism features than the symptoms mentioned.

Option D: Aripiprazole - It is an atypical antipsychotic used to treat schizophrenia and bipolar disorder, but its side effects are typically less severe than those of clozapine. Aripiprazole is more likely to cause nausea, anxiety, and insomnia than the symptoms described.

Solution to Question 2:

Postpartum blues is the most likely diagnosis in a postpartum woman with tearfulness, mood swings, and insomnia from day 4.

Postpartum blues, also known as baby blues, is a transient mood disorder characterized by sadness, tearfulness, confusion, and dysphoria. There are no suicidal thoughts or thoughts of harming the baby. These symptoms typically develop 3-5 days after delivery. They are thought to be due to hormonal changes, the stress of giving birth, and the responsibilities of motherhood. They resolve spontaneously within 10-14 days.

Management involves reassurance, education, and support. The patient must be followed up and assessed if symptoms persist for >2 weeks to evaluate postpartum depression.

Other options:

Option A: When the symptoms of depressed mood, anxiety, insomnia, and weight changes start within 4 weeks as per DSM 5 or 6 weeks as per ICD 11, and the symptoms persist for ≥ 2 weeks, the diagnosis will be postpartum depression. The patient may have occasional thoughts of harming the baby and suicidal thoughts. It can persist for months to years when untreated.

Option C: Postpartum psychosis is a psychiatric emergency that usually occurs within 4 weeks after delivery (DSM 5) or 6 weeks after delivery (ICD 11). The predominant symptoms are

hallucinations, delusions, suicidal thoughts, and thoughts of harming the baby. Psychotic symptoms such as insomnia, restlessness, labile mood, and agitation are seen.
Lithium, antipsychotics, and an antidepressant is the treatment of choice.

Option D: Postpartum anxiety is not defined by DSM-5. However, some platforms might describe postpartum anxiety as excessive or severe worry after childbirth.

Note: Zuranolone is the first-ever oral drug to be approved specifically for the treatment of postpartum depression. It is a neuroactive steroid that enhances the action of GABA in the brain and regulates mood.

Solution to Question 3:

The given clinical scenario is suggestive of a panic disorder. Panic disorder refers to an acute intense attack of anxiety accompanied by feelings of impending doom. Such attacks may not only occur in response to a fearful situation, or phobic objects, or in response to obsession or separation as in other anxiety disorders.

Panic disorder is characterized by the presence of an abrupt surge of intense fear or intense discomfort that reaches a peak within minutes. During the episode, the person may experience symptoms like palpitations, sweating, trembling or shaking, shortness of breath, nausea, dizziness, chest pain or discomfort or fear of dying. The patient also might feel like losing control, having a heart attack, or going crazy.

There is a dysfunction in major neurotransmitter systems like norepinephrine, serotonin, and GABA. Panic-inducing substances or panicogens may induce such attacks and include substances like 5-35% carbon dioxide, sodium lactate, bicarbonate, yohimbine (α blocker), flumazenil (GABA antagonist), cholecystokinin and caffeine.

It is not attributable to any drug abuse, medication, medical condition or mental disorder.

Treatment of this condition can be done by cognitive behavioral therapy and drugs like benzodiazepines (alprazolam, lorazepam), selective serotonin reuptake inhibitors (paroxetine), tricyclic antidepressants and monoamine oxidase inhibitors.

Solution to Question 4:

The above clinical history is suggestive of Wernicke's encephalopathy caused by thiamine (vitamin B1) deficiency.

Wernicke's encephalopathy is an acute, neuropsychiatric syndrome characterized by nystagmus and ophthalmoplegia, mental status changes (confusion), and unsteadiness of stance and gait (ataxia). It is a medical emergency, and thiamine should be initiated immediately, either intravenously or intramuscularly. The current standard of treatment for patients with symptoms of Wernicke's encephalopathy is to give thiamine 100 mg intravenously before administering glucose-containing IV fluids.

Other options:

Option B: Korsakoff psychosis - While Korsakoff psychosis is also associated with thiamine deficiency and can occur along with Wernicke's encephalopathy, it is primarily characterized by memory loss and confabulation. The presence of confusion, ataxia, and painful eye movements with 6th cranial nerve involvement is more indicative of Wernicke's encephalopathy.

Option C: Delirium tremens - It is a severe form of alcohol withdrawal syndrome and usually occurs within 3-4 days of alcohol withdrawal. The features of delirium tremens are clouding of consciousness, disorientation to time and place, poor attention and distractibility, visual hallucinations and illusions, tachycardia, fever, hypertension, diaphoresis, and pupillary dilatation, and psychomotor agitation and ataxia.

Option D: De Clerembault syndrome (erotomania) - It is characterized by the delusional belief that someone is deeply in love with them. It is more common in women than in men, commonly a young woman who believes that a man whom she considers to be of higher social and/or professional class, is in love with her. This condition is not related to alcohol abuse and does not cause the neurological symptoms described in the question.

Solution to Question 5:

The clinical scenario of ejaculation very early on vaginal penetration is suggestive of premature ejaculation (PE), and the squeeze technique is a non-pharmacological management (behavioral method) used to prevent it.

PE is a disorder of the orgasm phase of the normal sexual cycle.

In premature ejaculation, there is a persistent or recurrent achievement of orgasm and ejaculation before the patient actually wishes to. According to DSM-5, it is classified as follows:

• Mild if ejaculation occurs within approximately 30 seconds to 1 minute of vaginal penetration

• Moderate if ejaculation occurs within approximately 15 to 30 seconds of vaginal penetration

• Severe if ejaculation occurs at the start of sexual activity or within approximately 15 seconds of vaginal penetration.

Pharmacologic therapy when combined with behavioral treatment appears to be more effective in treating premature ejaculation.

The squeeze technique is a behavioral method used on the penis to prevent premature ejaculation. Here, the penis is stimulated until just before orgasm after which it is gently squeezed.

Other options:

Option A: Although Cognitive behavioral therapy (CBT) can be beneficial for some patients with sexual dysfunction. It is not a specific technique for managing PE. Instead, CBT is more useful in addressing underlying psychological issues or relationship problems that may contribute to the condition. CBT is used in the treatment of depression, phobia, anxiety and panic disorders, OCD, personality disorders, and somatoform disorders.

Option B: Exposure and response prevention therapy is a strategy within CBT that is effective in the management of OCD. It is also useful in other conditions like tics but is not a suitable method for managing premature ejaculation.

Option D: Sensate focus therapy is a couple-oriented sex therapy that involves the exchange of sensual touch between partners to alleviate anxiety and negativity associated with sex and to build better sexual intimacy and communication. It does not specifically address the issue of premature ejaculation like the squeeze technique does and is hence not useful.

Psychiatry INI-CET 2023

Question 1:

A boy is experiencing low mood, decreased energy, and decreased interest in his surroundings. He is not taking food for the past one week. The thought sample says "I lost my head. I am dead. How can I eat?". What is the delusion seen in this case?

a) Cotard's syndrome

b) Hypochondriacal delusion

c) Nihilistic delusion

d) Bizarre delusion

Question 2:

La belle indifference is seen in:

a) Mania

b) Depression

c) Schizophrenia

d) Conversion disorder

Question 3:

SCOFF protocol is used for assessing:

a) Mental health disorder

b) Behavioural disorder

c) Sexual health disorder

d) Eating disorder

Question 4:

The tactile hallucinations seen in cases of chronic cocaine intoxication is?

a) Oneroid state

b) Formication

c) Dipsomania

d) Carphologia

Question 5:

Which of the following is used to define dependence?

a) 1, 2

b) 1 only

c) 1, 2, 3, 4

d) 1, 2, 3

Question 6:

A pregnant female with bipolar disorder came to the clinic in the third trimester. She has had 4 episodes in the past and is stable on lithium 750 mg/day since 1 year during which time she has not had any episodes. What is the next best step in management?

a) Stop lithium and let her be drug free

b) Increase lithium dosage

c) Decrease lithium dosage

d) Switch to valproate

Question 7:

HEADSSS assessment will be used by an intern for which of the following patient groups?

a) Children under 5 years

b) Neonates

c) Infants

d) Adolescents

Answer Key

Question No.	Correct Option

1	c
2	d
3	d
4	b
5	c
6	c
7	d

Detailed Explanations

Solution to Question 1:

The type of delusion seen in this patient is a nihilistic delusion.

Nihilism refers to the delusion of the non-existence of self or parts of self. It also includes an attitude of rejection of established values or extreme skepticism regarding moral and value judgments. The patient may believe that they are going to die soon or their organs are rotting. Delusion of nihilism is often seen as a mood-congruent delusion in depression with psychotic symptoms. This patient has three major symptoms of depression — low energy, depressed mood, and lack of interest in the surroundings. However, these symptoms have to be present for a period of at least 14 days to make the diagnosis of major depressive disorder.

Other options:

Option A: Cotard's syndrome was first described by Jules Cotard. He described a psychotic disorder characterized by severe depression and nihilistic delusions regarding one's own body or existence. In the DSM or ICD, Cotard's syndrome is not classified separately. Nihilistic delusions have been mentioned as examples of mood-congruent delusions in a depressive episode with psychotic features.

Option B: Patients with hypochondriacal delusions are fixated on one or more somatic symptoms as evidence of some severe, yet undetected disease. They undergo repeated and unnecessary medical examinations and tests. There are frequent deteriorations of relationships with their doctors when nothing suggesting a disease is ever found, and this leads to doctor shopping.

Option D: Delusions are false and fixed ideas that are not shared by other people. Bizarre delusions are the ones whose content is out of the realm of possibility, for example, a patient believing that aliens abduct him and take him to Mars every night. Typically, patients with schizophrenia tend to have bizarre delusions.

Solution to Question 2:

La belle indifference is seen in conversion disorder.

La belle indifference refers to a lack of concern towards the symptoms, despite the apparent severity of the disability. Although not pathognomonic of conversion disorder, it is often seen in

association with it.

Conversion disorder is also known as functional neurological symptom disorder or dissociative disorder. It includes acute and temporary loss of motor or sensory function without any underlying neurological disease. Because the onset of such symptoms often coincides with psychological issues such as internal conflict, it was speculated that these issues converted to neurological symptoms. Conversion disorder can present with symptoms such as paresis, non-epileptic seizures (NES), blindness, and anesthesia of the extremities. One of the gait disturbances which can be seen is astasia-abasia — a wildly ataxic, staggering gait with gross and irregular truncal movements, and thrashing and waving arm movements. The patient often has a carefree attitude towards his illness as mentioned above.

Patients of conversion disorder achieve primary gain by keeping internal conflicts outside of their awareness. These conflicts manifest as physical symptoms. As a result of being sick, the patients are excused from obligations and receive support that might not otherwise have been available. This is referred to as secondary gain.

Solution to Question 3:

The SCOFF (Sick, Control, One, Fat, Food) questionnaire is a screening tool for eating disorders (anorexia nervosa and bullimia nervosa).

It consists of 5 questions addressing the core features of these disorders. The SCOFF questions are:

• Do you make yourself Sick because you feel uncomfortably full?

• Do you worry that you have lost Control over how much you eat?

• Have you recently lost more than One stone (~6.35 kg) in a 3-month period?

• Do you believe yourself to be Fat when others say you are too thin?

• Would you say that Food dominates your life?

A 'Yes' answer to 2 or more questions out of the 5 indicates likely anorexia or bulimia.

Solution to Question 4:

Formication, which refers to the sensation of insects crawling beneath the skin, is a feature of cocaine abuse. It is also referred to as Magnan's symptoms or cocaine bugs.

The symptoms of acute stimulant intoxication include agitation, irritability, aggression, and an increase in psychomotor activity. On examination, tachycardia, hypertension, and mydriasis may be seen. When psychotic symptoms such as delusions and hallucinations are present due to stimulants such as cocaine, it is referred to as stimulant-induced psychotic disorder. Paranoid delusions and hallucinations are hallmarks of stimulant-induced psychosis. Tactile hallucinations in the form of formication are seen in cocaine use. The presence of these symptoms depends on the dose and duration of the substance abuse and thus may not be seen in all patients. Cocaine-induced psychotic disorders are more frequently seen in intravenous users and those who

use crack (an extremely potent and addictive form of cocaine that can be smoked).

Other options:

Option A: Oneroid state is a dream-like state which can happen because of substance use or some organic disorder.

Option C: Dipsomania refers to a compulsive urge to drink alcoholic beverages. Patients suffering from alcohol-use disorders feel a need for daily use of large amounts of alcohol for adequate functioning throughout the day.

Option D: Carphologia (also known as floccillation) refers to the action of trying to pick imaginary objects. The patient will usually be trying to pick lint from his clothes or the bedsheet. This is often seen in delirium. Datura intake can lead to a delirious state and may show carphologia.

Solution to Question 5:

According to ICD-11, dependence on any substance is characterized by:

• Strong internal drive to use the substance, which the patient is unable to control.

• Increased priority is given to the use of the substance over other activities.

• Persistence of substance use despite harmful consequences.

• Presence of the subjective sensation of urge or craving to use the substance.

• Tolerance to the effect of the substance; the patient may require higher amounts of the substance to achieve the same effect.

• Appearance of withdrawal symptoms on cessation or reduction in the dose of the substance.

The diagnosis of dependence can be made if these symptoms are present for at least a period of 3 months.

Solution to Question 6:

In this scenario of a pregnant woman in her third trimester with bipolar disorder with multiple past episodes and currently stable on lithium, the best next step in management is to decrease lithium dosage within therapeutic levels.

Lithium is a mood stabilizer with a very narrow therapeutic index and is the first line of management in bipolar disorder. Lithium therapy should be continued throughout the pregnancy in women who have severe, frequent episodes, and these patients should be counseled about the reproductive risks associated with therapy. Ideally, the next best step in this scenario would be to monitor lithium blood levels monthly until 34 weeks, and weekly monitoring thereafter until delivery. However, dose reduction can be considered closer to delivery as there may be an increased risk of lithium toxicity in the fetus. According to NICE guidelines, it is recommended to stop lithium during active labour, measure lithium levels 12 hours after the last dose and reestablish prophylactic levels after delivery.

First and second trimester are characterised by a significant decrease of lithium blood levels with a risk of subtherapeutic levels. In third trimester and the postpartum, lithium levels gradually return to their preconception level which implicates that in this period clinicians need to be aware of the risk of lithium intoxication (Option B).

Bipolar disorder has been associated with adverse perinatal outcomes such as preterm birth. Medication discontinuation in pregnancy (Option A) can pose an increased risk for relapse. Moreover, symptomatic women during pregnancy more likely have postpartum relapses that can be severe.

Typical therapy for bipolar disorder includes mood stabilizers such as lithium, valproic acid, and carbamazepine and antipsychotic medications. Bipolar disorder during pregnancy is a complicated condition that is best treated in tandem with psychiatric care. Risks and advantages of mood stabilizers, some of which are teratogenic, must be considered while making decisions.

For example, lithium use during the first trimester has been linked to Ebstein anomaly, although very rarely. Fetal echocardiography is recommended for lithium-exposed fetuses. Fetal lithium exposure is also associated with neonatal goiter, CNS depression, hypotonia (floppy baby syndrome), and cardiac murmur.

Maternal adverse effects of lithium can be acute such as tremors, ataxia, weakness, polyuria, polydipsia or chronic such as nephrogenic diabetes insipidus, thyroid dysfunction, and hyperparathyroidism.

Due to all these adverse effects, patients on lithium therapy should have routine monitoring of serum electrolytes, serum calcium, thyroid profile, renal function tests, ECG, and serum lithium levels.

Other options :

Option D : It is not necessary to switch to valproate in the third trimester as the patient is already stable on lithium. Use of valproate in the first trimester is associated with significant teratogenicity.

Solution to Question 7:

HEADSSS assessment is done in adolescents to determine early symptoms of mental disorders and functional impairment in major life domains such as home and family, school, peers, and community.

These domains are included in the HEADSSS (Home, Education, Activities, Drugs, Sexuality, Suicide/Depression, Safety) Interview Guide. An adolescent’s risk and protective factors are identified during each visit using this assessment. Risk factors increase the likelihood that an adolescent will engage in risky behaviors, whereas protective factors build an adolescent’s resiliency and ability to cope with stress.

Examples of questions in HEADSSS assessment :

Home	Who lives at home, and where?Do you have your own room?What are relationships like at home?What do parents and relatives do for a living?
Education	School/grade performance-any recent changes? Any dramatic past changes?Any years repeated/classes failedSuspension, termination, dropping out?Any current or past employment?
Activities	On own, with peers (what do you do for fun? where? when?)Hobbies-other activities?History of arrests-acting out-crime?
Drugs	Use by peers? Use by young person? (include tobacco, alcohol)Use by family members? Amounts, source, frequency, patterns of use/abuse
Sexuality	Degree and types of sexual experience and acts? Number of partners?Sexually transmitted diseases-knowledge and prevention? Contraception? Frequency of use?History of pregnancy/abortion?
Suicide / Depression / Self Image	Sleep disorders, appetite/eating behavior changes, emotional outbursts and highly impulsive behavior, history of withdrawal/isolation, hopeless/helpless feelings, suicidal ideationHistory of past suicide attempts, depression, psychological counseling
Safety	Do you ever feel unsafe?

Psychiatry INI-CET 2024

Question 1:

Which of the following is not a priority condition under the WHO Mental health gap action plan?

a) Schizophrenia and other psychotic disorders

b) Obsessive compulsive disorder

c) Depression

d) Mental health disorders in children

Question 2:

Which of the following is not true about 'rapid cyclers'?

a) Occurs commonly in men

b) Commonly seen with concomitant hypothyroidism

c) At least 4 distinct episodes per year

d) Can be triggered by antidepressants

Question 3:

Which among the following is not part of the Beck's cognitive theory on depression?

a) Cognitive decline

b) Automatic thoughts

c) Dysfunctional beliefs

d) Introjection

Question 4:

Which of these is not an essential feature for the diagnosis of autism spectrum disorder?

a) Persistent deficits in social communication

b) Abnormalities in socializing

c) Restrictive patterns of behaviour and interests

d) Cognitive impairment

Question 5:

All the following can be treated with supportive psychotherapy except

a) Person with cognitive decline and physical symptoms

b) Person who has good motivation and control over frustration

c) Patient with acute psychiatric illness along with crisis

d) Patient with severe chronic disease and ego dysfunctioning

Answer Key

Question No.	Correct Option
1	b
2	a
3	d
4	d
5	b

Detailed Explanations

Solution to Question 1:

Obseesive compulsive disorder is not a priority condition under WHO Mental health gap action plan.

WHO aims to offer health planners, policymakers, and donors a comprehensive and cohesive set of initiatives and projects to enhance the provision of care for mental, neurological, and substance use disorders through the Mental Health Gap Action Programme (mhGAP).

The priority conditions identified by the above criteria for mhGAP are depression, schizophrenia and other psychotic disorders, suicide, epilepsy, dementia, disorders due to use of alcohol, disorders due to use of illicit drugs, and mental disorders in children.

The intervention package includes evidence-based strategies for prevention and management, considering cost-effectiveness, severity of disorders, poverty reduction, and human rights protection.

Solution to Question 2:

Option A is the false statement. Rapid cycling disorder is more common in females.

Rapid cycling disorder is diagnosed when the patient has a combination of 4 or more bipolar episodes within the last 12 months. Two separate episodes can be demarcated from one another by either switch in polarity or full/partial remission for at least 2 months. Manic and hypomanic episodes are considered to be on the same pole, hence a switch in polarity includes an episode of mania/hypomania with an episode of depression.

Possible risk factors for rapid cycling disorder include female patients, diabetes, childhood abuse or family trauma, and alcoholism or substance abuse. Rapid-cycling bipolar patients are significantly more likely than non-rapid-cycling patients to have a lifetime history of comorbidities like anxiety disorders, ADHD, OCD, bulimia nervosa, and substance use disorders.

The drug of choice for prophylaxis of rapid cycling disorder is valproate

Other options

Option B: Conditions like hypothyroidism and diabetes are possibly involved in some cases of rapid cycling .

Option D: Usage of antidepressants could worsen rapid cycling in bipolar depression.

Solution to Question 3:

Introjection is not a part of the Beck's cognitive theory of depression.

Aaron Beck's cognitive triad consists of:

• Beliefs about oneself - a pessimistic self-image

• Perceptions of the surroundings - a tendency to see the world as unfriendly and challenging,

• Thoughts about the future - anticipating pain and defeat.

The psychodynamic understanding of depression, initially defined by Sigmund Freud and later expanded by Karl Abraham, is considered the classic view of depression. This theory includes four main points: (1) disturbances in the infant-mother relationship during the oral phase (the first 10 to 18 months of life) create a predisposition to depression later in life; (2) depression can arise from the real or imagined loss of an important object; (3) introjection of the lost object serves as a defense mechanism to cope with the distress of the loss; and (4) because the lost object is associated with both love and hate, feelings of anger are turned inward toward the self.

Solution to Question 4:

Cognitive impairment does not form part of the diagnostic criteria for autism spectrum disorder.

The DSM-5 diagnostic criteria for autism spectrum disorder are as follows:

• Persistent deficits in social communication and social interaction across multiple contexts, currently or by history

• Restricted, repetitive patterns of behavior, interests, or activities, currently or by history

• Symptoms must be present in the early developmental period (but may not become fully manifest until social demands exceed limited capacities, or may be masked by learned strategies in later life).

• Symptoms cause clinically significant impairment in social, occupational, or other important areas of current functioning.

• These disturbances are not better explained by intellectual disability or global developmental delay. Intellectual disability and autism spectrum disorder frequently co-occur; to make comorbid diagnoses of autism spectrum disorder and intellectual disability, social communication should be below that expected for the general developmental level.

Solution to Question 5:

Supportive psychotherapy is not indicated in a person who has good motivation and control over frustration. Such individuals are better treated by expressive psychotherapy.

Supportive psychotherapy places major etiological emphasis on external rather than intrapsychic events, particularly on stressful environmental and interpersonal influences on a severely damaged self. The general aim of supportive treatment is the amelioration or relief of symptoms through behavioral or environmental restructuring within the existing psychic framework. This often means helping the patient to adapt better to problems and to live more comfortably with his or her psychopathology.

Supportive therapy uses several methods, either singly or in combination, including warm, friendly, strong leadership; help in developing pleasurable activities (e.g., hobbies); removal of excessive strain, when possible; medication to alleviate symptoms; and guidance and advice in dealing with current issues. This therapy uses techniques to help patients feel secure, accepted, encouraged, and not anxious.

Supportive psychotherapy is generally indicated for those patients for whom classic psychoanalysis or insight-oriented psychoanalytic psychotherapy is typically contraindicated-those who have poor ego strength and whose potential for decompensation is high.

Amenable patients fall into the following major areas:

• Individuals in acute crisis or a temporary state of disorganization and inability to cope, whose intolerable life circumstances have produced extreme anxiety or sudden turmoil (e.g., individuals going through grief reactions, illness, divorce, job loss, or who were victims of crime, abuse, natural disaster, or accident);

• Patients with chronic severe pathology with fragile or deficient ego functioning (e.g., those with latent psychosis, impulse disorder, or severe character disturbance);

• Patients whose cognitive deficits and physical symptoms make them particularly vulnerable and, thus, unsuitable for an insight-oriented approach (e.g., certain psychosomatic or medically ill persons);

• Individuals who are psychologically unmotivated (e.g., patients who come to treatment in response to family or agency pressure and are interested only in immediate relief or those who need assistance in very specific problem areas of social adjustment as a possible prelude to more exploratory work).

Because support forms a tacit part of every therapeutic modality, it is rarely contraindicated as such. The typical attitude regards better-functioning patients as unsuitable not because they will be harmed by a supportive approach, but because they will not be sufficiently benefited by it.

Psychiatry NEET 2024

Question 1:

A patient comes with complaints of difficulty falling asleep. He consumes black coffee one hour before bed. Which of the following is the role of caffeine in the wakefulness?

a) Caffeine has no effect, and he can continue taking black coffee

b) Caffeine causes release of chemical from locus coeruleus that causes awakening

c) Caffeine has histaminic action in brain that leads to awakening

d) Caffeine inhibits adenosine receptors, thus leading to awakening

Question 2:

Which of the following SNRI is used in the treatment of ADHD?

a) Guanfacine

b) Methylphenidate

c) Reboxetine

d) Modafinil

Question 3:

A 45-year-old woman presents with a feeling of nervousness and has abdominal pain and diarrhoea. She has had a history of heartburn, and indigestion for the past 25 years. She gives similar complaints in the family of being "tense and nervous". What other symptoms might you expect to see in this condition?

a) Tingling of extremities

b) Neologism

c) Idea of reference

d) Hallucination

Question 4:

A girl was referred to the psychiatry department by the plastic surgeon. She had visited multiple plastic surgery departments requesting full facial reconstruction. She complains of dark blemishes and spots on her face, which are not present. She always says that she is ugly and looks constantly in the mirror and the back of the spoon and windows. Her mother assured her of no such issues, but still, she persists in getting the surgery. What should the psychiatrist do?

a) Start on atypical antipsychotics

b) Give fitness for surgery

c) Start her on selective serotonin reuptake inhibitors

d) Initiate insight-oriented psychotherapy

Question 5:

A girl presented with multiple episodes of loss of consciousness. The episodes usually happen in the presence of the family members in the morning. She regains consciousness after 15 to 30 minutes and usually remembers the events during these episodes. There is no history of tongue bite or urinary incontinence. What should be done in this scenario?

a) Start behavioural therapy

b) Initiate aversion therapy

c) Give valproate

d) Start on ketogenic diet

Question 6:

If a child with ADHD goes untreated into adolescence, what is he most likely to develop?

a) Selective mutism

b) Separation anxiety disorder

c) Conduct disorder

d) Binge eating disorder

Question 7:

A chronic alcoholic met with a vehicular accident. His blood alcohol level is >300 mg/dl. He needs hospitalization and will be withheld from taking alcohol. The drug that will be used to manage his withdrawal symptoms is:

a) Methyl alcohol

b) Fomepizole

c) Lorazepam

d) Buspirone

Question 8:

A 19-year-old girl child came agitated to the emergency department after consuming some drug with a previous history of suicidal tendencies. The parents do not know the mode and route of the drug administered. She presented with tachycardia, tachypnea, diaphoresis, and dilated pupils. What is the likely drug administered?

a) Chlorpheniramine

b) Morphine

c) Heroin

d) Cocaine

Question 9:

A 19-year-old presented to the psychiatry OPD. He has a charming persona and is well-composed and behaved in front of interviewers. His mother complains of him getting into fights and bunking classes, drinking alcohol and getting involved in police cases. What kind of personality disorder does he have?

a) Paranoid

b) Antisocial

c) Narcissistic

d) Schizotypal

Answer Key

Question No.	Correct Option
1	d
2	c
3	a

4	c
5	a
6	c
7	c
8	d
9	b

Detailed Explanations

Solution to Question 1:

The role of caffeine in the wakefulness in this patient is that caffeine inhibits adenosine receptors, thus leading to awakening.

Caffeine, a commonly used stimulant, plays a significant role in promoting wakefulness and alertness. It functions primarily as an antagonist at adenosine A2A receptors in the brain. Adenosine, a natural neuromodulator which accumulates in the basal forebrain area promotes sleepiness and relaxation by activating these receptors. By blocking adenosine A2A receptors, caffeine counteracts these sedative effects, thus helping to increase wakefulness and reduce fatigue.

In addition to its receptor antagonism, caffeine inhibits cyclic nucleotide phosphodiesterases, which leads to increased levels of cyclic AMP (cAMP) in the brain. This action enhances the release of neurotransmitters such as norepinephrine (NE) and dopamine (DA), further contributing to increased neural activity and alertness. Caffeine is rapidly absorbed from the digestive tract and crosses the blood-brain barrier easily, affecting brain function soon after consumption.

Caffeine's stimulating effects can also be influenced by factors such as metabolism, which varies widely among individuals. For instance, smoking can reduce caffeine's half-life, making it clear from the body faster, while oral contraceptives can prolong its effects. Therefore, consuming caffeine close to bedtime, as in the case of drinking black coffee an hour before bed, can interfere with sleep by sustaining wakefulness and reducing the ability to fall asleep.

Solution to Question 2:

The SNRI that can be used for the treatment of ADHD is Reboxetine.

Reboxetine, a SNRI has shown potential efficacy in treating ADHD, particularly in patients who do not respond to methylphenidate. Common side effects of reboxetine include drowsiness, sedation, and gastrointestinal issues.

In the treatment of ADHD , pharmacotherapy is the primary approach, with central nervous system (CNS) stimulants being the most effective. Stimulants such as methylphenidate and amphetamine salts are commonly used due to their high efficacy in managing symptoms,

although they come with risks of dependence and addiction. For patients who cannot tolerate stimulants or have a family history of addiction, non-stimulant options are preferred. These include atomoxetine, a norepinephrine reuptake inhibitor (NRI), and guanfacine, an alpha-2 adrenergic agonist.

Modafinil is used primarily for narcolepsy and is not a standard treatment for ADHD.

Solution to Question 3:

In this scenario, given the patient's feelings of nervousness, abdominal pain, diarrhoea, and a history of long-term digestive issues along with familial history, you might also expect to see tingling of extremities as a symptom of Generalized Anxiety Disorder (GAD).

According to DSM-5 (and ICD-11) criteria, GAD is characterized by excessive and uncontrollable anxiety and worry about various aspects of life, persisting for at least six months. This anxiety is typically accompanied by at least three additional symptoms from a specific list, which includes:

• The presence of excessive anxiety and worry about a variety of topics, events, or activities.

• Worry occurs more often than not for at least six months and is clearly excessive.

• The worry is experienced as very challenging to control.

• The worry in both adults and children may easily shift from one topic to another.

• The anxiety and worry are accompanied by at least three of the following physical or cognitive symptoms (In children, only one of these symptoms is necessary for a diagnosis of GAD):

• Edginess or restlessness

• Tiring easily; more fatigued than usual impaired concentration or feeling as though the mind goes blank

• Irritability (which may or may not be observable to others)

• Increased muscle aches or soreness

• Difficulty sleeping (due to trouble falling asleep or staying asleep, restlessness at night, or unsatisfying sleep)

Neologism, idea of reference, and hallucinations are not typical symptoms of GAD but are more associated with other psychiatric conditions such as psychotic disorders like schizophrenia.

Solution to Question 4:

The given history suggests the presence of body dysmorphic disorder (BDD) in the patient, as she is preoccupied with her appearance and self-perceived flaws, despite being told otherwise by her family. The psychiatrist should start her on selective serotonin reuptake inhibitors(SSRI).

BDD is characterized by persistent preoccupations about one or more perceived defects or flaws in one's appearance, but the defects or flaws are negligible or not even observable to others. In response to their appearance concerns, individuals with BDD demonstrate a range of mental acts:

• Comparing themselves with others

• Checking in the mirror

• Camouflaging their perceived flaws

• Avoidance- from minor social avoidance to being housebound.

The specific focus of BDD on appearance serves to differentiate it from other conditions characterized by repetitive behaviours.

The treatment for BDD typically involves a combination of SSRIs and psychotherapy. SSRIs, such as fluoxetine and clomipramine, have been shown to reduce symptoms in many patients, helping alleviate the obsessive thoughts and anxiety associated with BDD.

Psychotherapy, particularly cognitive-behavioural therapy (CBT), is crucial as it addresses the distorted self-image and underlying emotional issues. CBT helps patients reframe their thoughts about appearance and develop healthier coping strategies.

It's important to note that while surgical interventions may be sought by patients, they are often ineffective and can lead to further dissatisfaction and distress. Therefore, focusing on mental health treatment is essential. Regular follow-ups and a supportive therapeutic relationship can significantly enhance the effectiveness of treatment.

Other options:

Option A: Start on atypical antipsychotics: Antipsychotics are generally not the first-line treatment for BDD. They may be considered in cases with comorbid psychotic symptoms, but this patient does not exhibit hallucinations or delusions.

Option B: Give fitness for surgery: Giving fitness for surgery would be inappropriate and potentially harmful. Surgery is not the solution for BDD, as patients typically remain dissatisfied with the results and continue seeking more surgeries.

Option D: Initiate insight-oriented psychotherapy: While psychotherapy, particularly cognitive-behavioral therapy (CBT), is effective for BDD, insight-oriented psychotherapy (focused on understanding unconscious conflicts) is less effective than targeted approaches like CBT in managing this condition.

Solution to Question 5:

In this scenario, the girl is experiencing episodes of loss of consciousness that may suggest conversion disorder. The most appropriate management is to start behavioural therapy to help address her symptoms and underlying stressors.

Conversion disorder, also referred to as functional neurological symptom disorder in the DSM-5, involves symptoms or deficits that affect voluntary motor or sensory functions, suggesting another medical condition. However, these symptoms are believed to arise from psychological factors, often triggered by conflicts or stressors. The symptoms are not intentionally produced, are not the result of substance use, and are not limited to pain or sexual issues. The primary gain associated with conversion disorder is psychological rather than social, monetary, or legal.

Common presenting symptoms include:

• Non-epileptic seizures

• Weakness and/or paralysis

• Movement disorders

• Speech or visual impairment

• Swallowing difficulty

• Sensory disturbances

• Cognitive symptoms

Resolution of conversion disorder symptoms often occurs spontaneously but can be enhanced through supportive or behavioural therapy. For patients who are resistant to psychotherapy, it can be beneficial to focus on stress and coping strategies rather than directly addressing their symptoms as imaginary, as this may exacerbate their condition.

Additional techniques, such as hypnosis, anxiolytics, and behavioural relaxation exercises, may also be effective in managing symptoms. In certain cases, medications like parenteral amobarbital or lorazepam can help gather further information, especially if the patient has experienced a recent traumatic event.

Psychodynamic approaches, including psychoanalysis and insight-oriented psychotherapy, allow patients to explore their internal conflicts and the symbolic meanings of their symptoms.

Solution to Question 6:

If a child with ADHD goes untreated into adolescence, among the following options he is most likely to develop conduct disorder.

The course of ADHD is variable. The majority of the cases persist into adult life. The hyperactivity may sometimes disappear, but the decreased attention span and impulse-control problems usually persist. Patients with ADHD are vulnerable to conduct disorder, antisocial behaviour, substance use disorders, and mood disorders.

ADHD is a neuro psychiatric condition characterized by a pattern of diminished attention span and increased impulsivity or hyperactivity. The prefrontal cortex of the brain has been implicated in its etiology and dopamine is the main neurotransmitter.

ADHD can begin during infancy but is usually recognized between 1-3 years of age. ADHD was known as minimal brain dysfunction earlier. The characteristics of children with ADHD, in order of frequency, are

• Hyperactivity

• Attention deficit (short attention span, distractibility)

• Impulsivity (action before thought, abrupt shifts in activity, lack of organization)

• Memory and thinking deficits

• Specific learning disabilities

• Speech and hearing deficits.

According to DSM-5 criteria, a diagnosis of ADHD requires symptoms to be present before age 12, persist for at least six months, negatively impact social, academic, or occupational activities, and occur in two or more settings. Pharmacotherapy is the first-line treatment, with CNS stimulants like methylphenidate and amphetamine salts being the most effective due to their mild side effects. Non-stimulant options include atomoxetine, clonidine, and guanfacine. Additionally, psychosocial interventions—such as academic organization skills, parent training, behaviour modification, cognitive-behavioural therapy (CBT), and social skills training—are used as adjuncts to medication.

Note: Conduct disorder is characterized by a persistent pattern of behaviour that violates the basic rights of others and disregards societal rules. The severity of the behaviours can lead to emergencies, especially if they are alarming to parents. Individuals with severe and chronic symptoms may later be diagnosed with antisocial personality disorder after age 18. Treatment typically involves behaviour modification strategies and, if necessary, placement in a corrective institution.

Other options:

Option A: Selective mutism refers to the lack of speaking (being mute) in one or more specific social situations, commonly in school, but speaking well in other situations, commonly at home. The child may be either completely silent at school or maybe just whispering. Rarely, a child may be silent at home but communicate normally at school.

Option B: Separation anxiety disorder is characterized by developmentally inappropriate and excessive anxiety when the patient is separated from their attachment figure. It is a normal developmental phenomenon in infants. It peaks between 9 months and 18 months and usually diminishes by two and a half years of age.

Option D: Binge-eating disorder is the most common eating disorder. It is characterized by rapid intake of abnormally large quantities of food over a short period of time, even when not hungry. There are feelings of guilt about the eating episodes, but these patients do not compensate in any way following the episodes of binge.

Solution to Question 7:

The first-line drug used in the management of this patient's withdrawal symptoms is intravenous(IV) lorazepam.

Delirium tremens is the most severe form of alcohol withdrawal syndrome. It occurs usually within 3–4 days of alcohol withdrawal. Acute infection, hepatitis and pancreatitis predispose delirium tremens.

Clinically, patients present with confusion, agitation, and fluctuating levels of consciousness. Visual or tactile hallucinations along with features of autonomic hyperactivity like tachycardia, diaphoresis, fever, insomnia, and hypertension are also seen.

Delirium tremens is treated using benzodiazepines, starting with a higher dosage, which is gradually tapered as withdrawal symptoms disappear. Benzodiazepines such as IV lorazepam or oral chlordiazepoxide (every 4 hours) are given.

In the case of benzodiazepine-resistant cases, IV phenobarbital is the first-line drug, and IV propofol is the second-line drug.

It is also recommended to administer thiamine in patients with alcohol-related disorders to prevent Wernicke-Korsakoff syndrome

Solution to Question 8:

In the given clinical scenario, the patient presenting with tachycardia, tachypnea, diaphoresis, and dilated pupils after consuming some drug with a previous history of suicidal tendencies points towards a diagnosis of cocaine toxicity leading to sympathetic toxidrome.

Cocaine is a CNS stimulant obtained from coca plants belonging to the family Erythroxylaceae. The route of intake is by snorting, IV route and smoking. Tattoos may be made to cover the injection marks.

It inhibits the uptake of dopamine, norepinephrine, and serotonin.

At lower doses, it leads to alertness but at higher doses, it produces stereotyped behavior, involuntary actions, and paranoia. There is a transient euphoria. Tactile hallucinations (Magnan's symptom or cocaine bugs) are characteristic.

There is a dose-dependent increase in heart rate and blood pressure. It causes dilation of pupils and is associated with chills and perspiration. In severe cases, intracranial bleeding due to increased blood pressure, and myocardial infarction due to coronary vasospasm can occur. Ultimately, the patient goes into a stage of depression with loss of reflexes, paralysis of muscles, cardiorespiratory failure, and coma.

Management - Administration of activated charcoal and gastric lavage with warm water and potassium permanganate in case of ingestion. Benzodiazepines to control heart rate/seizures. Amyl nitrite is the antidote given via inhalation. Thiamine 100 mg IV and Naloxone 2 mg IV can also be given.

Other options:

Option A: Chlorpheniramine is an effective H1 antagonist that tends to be mildly sedative, making it suitable for daytime use. A notable number of patients experience side effects involving central nervous system stimulation.

Option B and C: Morphine and Heroin are isolated from opium. The clinical features include a euphoric feeling in the stage of excitement followed by headache, giddiness, and a tendency to sleep in the stage of stupor. Intoxication is characterized by bradycardia, hypotension, respiratory depression, hypothermia, and pinpoint pupils. Respiratory depression can even result in death.

The image below shows the Erythroxylum plant:

Solution to Question 9:

The patient most likely has antisocial personality disorder. Although the patient seems charming and well-composed, he is aggressive and clearly exhibits disregard for social norms, and shows no remorse for his actions. These findings are characteristic of antisocial personality disorder.

Antisocial Personality Disorder is characterized by a long-term pattern of manipulation, exploitation, or violation of others' rights. Individuals with antisocial personality disorder tend to show no regard for right or wrong and often disregard the feelings, rights, and safety of others.

Other options:

Option A: Paranoid personality disorder is characterized by unwarranted suspicion, hypersensitivity, jealousy, envy, rigidity, excessive self-importance, and a tendency to blame and ascribe evil motives to others.

Option C: Narcissistic personality disorder is marked by a heightened sense of self-importance. Individuals with this personality disorder display a lack of empathy and grandiose feelings. This is to mask their fragile self-esteem and vulnerability.

Option D: Individuals with schizotypal personality disorder appear strange to surrounding people. They have ideas of reference, odd or magical thinking, inappropriate or constricted affect, and suspiciousness. It is most commonly associated with depression in nearly half of the patients.

Radiology AIIMS 2017

Question 1:

A 30-year-old female came to OPD with complaints of backache. On examination, Schober test was positive and her eyes and nose were hyperpigmented. Probable diagnosis is

a) Hypoparathyroidism

b) Fluorosis

c) Ochronosis

d) Ankylosing spondylitis

Question 2:

A 30-year-old woman presented with complaints of bone pain and abdominal cramps. Her family says she has a history of abnormal behavior. The consultant doctor arrived at a provisional diagnosis based on the clinical features. Which of the following would be the best investigation to arrive at a definitive diagnosis?

a) MRI

b) Ultrasonogram

c) Sestamibi scan

d) CT scan

Question 3:

Based on the given image, what is the most likely diagnosis?



a) Carotico-cavernous fistula

b) Pseudoaneurysm of internal carotid artery

c) Tumour blush of angiofibroma

d) Extensive head and neck hemangioma

Question 4:

Which of the following will have reduced uptake on FDG PET scan?

a) Typical carcinoid

b) Small cell carcinoma

c) Atypical carcinoid

d) Large cell neuroendocrine carcinoma

Question 5:

A thin, 34-year-old male, presents with sharp pain and sudden onset dyspnea. Chest X-ray was performed and is shown below. What is the most probable diagnosis?

a) Pulmonary embolism

b) Pneumothorax

c) Hydropneumothorax

d) Pleural effusion

Question 6:

A 45-year-old woman comes to the emergency department with severe back pain, vomiting, and hypotension. CT findings are given below. What is the diagnosis?



a) Renal cyst

b) Renal cell carcinoma

c) Para pelvic cyst

d) Renal angiomyolipoma

Question 7:

A 25-year-old male came with complaints of acute abdominal pain. A plain X-ray abdomen erect was taken which is shown below. What is your diagnosis?

a) Subdiaphragmatic abscess

b) Pneumoperitoneum

c) Pneumopericardium

d) Intestinal obstruction

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	a
5	b

6	d
7	b

Detailed Explanations

Solution to Question 1:

Given X-ray shows intervertebral disc calcification and hyperpigmentation of nose and eyes are present. Both features are indicative of ochronosis as diagnosis.

• Ochronosis also known as alkaptonuria is an autosomal recessive disorder.

• It is due to deficiency of homogentisic acid oxidase leading to accumulation of homogentisic acid.

• Clinical features: Triad of

• Homogentisic aciduria (urine turns dark color on standing due to oxidation or alkalinisation)

• Eumelanin like pigmentation of skin, sclera, cartilages etc.

• Degenerative chronic arthropathies especially of spine, hip, knees.

Option a: Hypoparathyroidism doesn't explain hyperpigmentation and other clinical features of hypoparathyroidism aren't mentioned.

Option b: Fluorosis - dental symptoms are not mentioned and osteoporosis will be marked in case of fluorosis. Also, hyperpigmentation is not a typical feature of fluorosis.

Option d: Ankylosing spondylitis – females are unlikely to be affected and it doesn't explain hyperpigmentation.

Solution to Question 2:

The given clinical features are suggestive of primary hyperparathyroidism. Sestamibi scan is the best investigation to arrive at a definitive diagnosis. Parathyroid adenoma is the most common cause of primary hyperparathyroidism.

Clinical features of hyperparathyroidism:

• Bones- bone pain, pathological fractures

• Stones- nephrocalcinosis

• Groans- abdominal pain

• Moans- psychotic symptoms

• Fatigue overtones- fatigue, malaise

Solution to Question 3:

The given image is a digital subtraction angiogram(DSA) and it shows a pseudoaneurysm of the internal carotid artery.

The right ICA injection anteroposterior view shows the neck of the pseudoaneurysm originating from the transverse segment of the cavernous portion of the ICA.

Option A: Carotico-cavernous fistula is characterized by the premature filling of the veins as shown in the image below:



Option C: Angiofibroma is characterised by tumour blush. It is the radiological enhancement of a tumour that reveals opacification of the well-vascularized tumours.

Option D: Hemangioma can be differentiated from other vascular malformations by the presence of well-circumscribed mass demonstrating intense tissue staining, usually in a lobular pattern.

Solution to Question 4:

Typical carcinoid is more likely to have reduced uptake on FDG PET scan than an atypical carcinoid.

The low metabolic rate of tumors makes the tumor negative (show reduced uptake) on an FDG PET scan.

Typical examples are:

• Bronchoalveolar carcinoma (Adenocarcinoma in situ)

• Carcinoid tumour (Typical > Atypical)

Solution to Question 5:

The given chest radiograph clearly depicts the presence of pneumothorax, which represents air in the pleural space.

• The right hemithorax shows blackening (depicts area of low density) due to the presence of air.

• The visceral pleural line has separated from the chest wall by a transradiant zone devoid of vessels.

• The collapsed lung can be seen with a clear border.

• There is a mediastinal shift to the left side (contralateral), which is typically seen with cases of tension pneumothorax.

All the above features are suggestive of pneumothorax.

Note: Pleural effusions signify the presence of fluid accumulation in the pleural space. The fluid may have varying degrees of mobility depending on the nature of effusion (free-flowing or loculated).

While a small amount of free fluid is usually hard to detect on an erect PA chest x-ray, a significant pleural effusion (200-500 mL) is represented by obscured lateral and posterior costophrenic angles and the hemidiaphragm affected. Subsequently, homogenous opacification of the lower lung field develops, associated with concave superior margin relative to the lung. This margin is higher laterally than medially.

A massive pleural effusion cause dense opacification usually associated with a contralteral mediastinal shift.

Here is an image of a left-sided pleural effusion representing the homogenous opacification of the left lower and middle lung fields. There is associated blunting of the left sided costophrenic angle. The superior border of the effusion is concave and is higher laterally than medially.

Solution to Question 6:

The given CT image shows well-circumscribed lesions with low attenuation (black in colour) that indicate fatty tissue. This is suggestive of renal angiomyolipomas.

Angiomyolipomas (AMLs) are benign tumours that occur sporadically or in association with tuberous sclerosis. It is commonly seen in adults between 30 and 50 years of age, with a female predilection. AMLs predominantly affect the kidneys, though they can occur in other sites such as the liver (a common extra-renal location) and less frequently in the ovary, fallopian tube, spermatic cord, palate, and colon.

Histologically, angiomyolipomas consist of perivascular epitheloid cells with varying proportions of vascular, immature smooth muscle cells, and fat cells. Patients with AMLs present with flank pain, a palpable mass, hematuria, recurrent urinary tract infections, hypertension, or renal failure. Larger tumours increase the risk of complications, such as spontaneous retroperitoneal hemorrhage, which can manifest as sudden-onset abdominal or flank pain, along with symptoms like nausea, vomiting, and hypotension.

Everolimus is used for the treatment of angiomyolipomas. Treatment should be considered for asymptomatic, growing AMLs measuring >3 cm in diameter. Angiomyolipomas do not normally require surgery unless life-threatening bleeding is present.

Other options:

Renal cyst is a broader term that includes the cystic lesions of the kidney. They can be simple cysts that are benign. Complex or complicated cystic lesions need evaluation and can be malignant. Bosniak classification is used for renal cysts. Cysts appear as hypodense lesions (greyish) with a thin wall, with or without septations.

CT image showing a simple renal cyst:



Renal cell carcinoma (RCC) is the most common malignant tumour of the kidney. It usually occurs in the age group 50-70 years and macroscopic hematuria is seen in about 60% of the cases. It is a differential for angiomyolipoma but it is important to note that macroscopic fat is almost always associated with ossification or calcification.



Solution to Question 7:

The above X-ray clearly shows the presence of subdiaphragmatic free gas in the right upper quadrant. This indicates pneumoperitoneum.

Causes of pneumoperitoneum include:

• Bowel perforation

• Peptic ulcer

• Ischemic bowel

• Necrotizing enterocolitis.

• Small bowel perforation

• Trauma

• Penetrating injuries

• During surgery / laparoscopy

Signs of pneumoperitoneum:

• Bowel-related signs

• Double wall sign (also known as Rigler's sign or gas-relief sign)

• Telltale triangle sign (also known as triangle sign)

• Peritoneal ligament-related signs

• Football sign

• Falciform ligament sign

• Lateral umbilical ligament sign (also known as inverted "V" sign)

• Urachus sign

- Right upper quadrant signs
- Cupola sign
- Fissure for ligamentum teres sign
- Hepatic edge sign
- Lucent liver sign
- Morison's pouch sign (doge cap sign)
- Periportal free gas sign



Radiology AIIMS 2018

Question 1:

A child presents with history of having ingested a button battery. What is your next course of action?

a) Give oral fluids

b) Remove endoscopically

c) Reassure and to check whether passed in stools

d) Induce vomiting

Question 2:

Which of the following statements are false regarding X Rays?

a) Loss of soft tissue planes is the earliest X-ray change of infection

b) Glass pieces can be seen in X-ray images.

c) Joint space is a misnomer as there is no real space in the joint.

d) Live bone is more radio-opaque than dead bone on X- Ray images.

Question 3:

A child presented with a history of road traffic accident. The axial CT image of the child, shown below, is suggestive of _____

a) Extradural Hemorrhage

b) Subdural Hemorrhage

c) Subarachnoid Hemorrhage

d) Intraventricular Hemorrhage

Question 4:

A 39-year-old male patient presented with complaints of fever and cough. His chest radiograph shows air bronchogram sign. The probable diagnosis is :

a) Pleural effusion

b) Consolidation

c) Hilar lymphadenopathy

d) Bronchiectasis

Question 5:

The Doppler waveform of the venous blood flow in lower limb DVT is _____.

a) Monophasic

b) Biphasic

c) Triphasic

d) Multiphasic

Question 6:

A middle-aged female patient with a history of cholelithiasis, presented with vomiting and abdominal pain. The abdominal radiograph showed Rigler's triad. Which of these is not a component of Rigler's triad?

a) Pneumobilia

b) Small bowel obstruction

c) Ectopic gall stone

d) Large bowel obstruction

Question 7:

All among the following are ionizing radiations except:

a) Alpha rays

b) Beta rays

c) X-rays

d) MRI

Question 8:

Empty thecal sac sign on MRI is seen in:

a) Arachnoiditis

b) Discitis

c) Vertebral osteomyelitis

d) Tethered cord syndrome

Question 9:

From the X-ray given, pick out the probable diagnosis:



a) Subdiaphragmmatic abscess

b) Pneumothorax

c) Pneumoperitoneum

d) Intestinal obstruction

Question 10:

What is the diagnosis shown in the image below?



a) Costochondritis

b) Spondylolisthesis

c) Cervical rib

d) Rib fracture

Question 11:

Identify the procedure.



a) Myelography

b) Angiography

c) DEXA

d) Projectional radiography

Answer Key

Question No.	Correct Option
1	b
2	d
3	a
4	b
5	a
6	d
7	d

8	a
9	c
10	c
11	a

Detailed Explanations

Solution to Question 1:

This case should be managed by removing the button battery endoscopically after confirming its position on X-Ray.

Button battery ingestion causes alkaline caustic injury, more in the areas surrounding the negative pole.

Management :

• The child is to be kept NPO until the oesophageal position of the battery is ruled out by X-ray scan. Vomiting should never be induced since the battery, which has passed through the esophagus, might lodge back again into it. Reassurance can be done only if the position of the battery is confirmed and if there are no symptoms.

• Since the question does not mention that the child is asymptomatic, the management of choice would be removing the battery endoscopically.

• Management also depends on battery life. Fresh batteries need to be removed as soon as possible as they have a higher concentration of injurious substances.

Solution to Question 2:

Live bone is not more radio-opaque than dead bone on X-ray images.

In fact, dead bone looks more white (more radio-opaque) than live bone on X-Ray as there's no bone resorption activity.

Loss of soft tissue planes is the earliest X-ray change in infections and is seen 24-48 hours after the onset of disease.

Joint space in X-rays is the radiolucency of cartilage being apparently perceived as space.

Glass pieces can be seen on X-ray images as they contain lead.

Solution to Question 3:

The given CT image shows the typical bi-convex hyperdense lesion not crossing cranial sutures, characteristic of an extradural hemorrhage.

As the dura mater tends to adhere to the skull, the hematoma collecting between the skull and this dura is seen on CT sections as a biconvex dense area immediately beneath the skull vault, convex towards both the brain and the vault. They tend not to cross the suture lines.

The temporoparietal convexity is the most common site. The hematoma usually lies beneath a fracture of the squamous part of the temporal bone. The underlying brain is displaced but often appears intrinsically normal.

Monitoring an extradural hemorrhage and immediate intervention if necessary is an important step in management since the brain can herniate with increased bleeding that is causing raised ICP.

Solution to Question 4:

The radiograph shows air bronchogram sign in a lung with consolidation, seen in pneumonia, which is supported by the clinical history of this patient.

An air bronchogram is a tubular outline of an airway made visible when there is filling of the surrounding alveoli by fluid or inflammatory exudates.

The causes of air bronchogram are as follows:

• Lung consolidation

• Pulmonary edema

• Nonobstructive pulmonary atelectasis

• Severe interstitial disease

• Neoplasm

• Normal expiration

Pleural effusion is seen as a homogenous opacity on chest radiograph.

The given radiograph shows peribronchial thickening and air bronchogram sign:





Solution to Question 5:

The Doppler waveform of the venous blood flow in lower limb DVT is monophasic proximal to the level of obstruction.

The normal venous waveform shows a variation (phasicity) secondary to respiration referred to as respirophasic waveform. Loss of this phasic variation results in a monophasic waveform suggestive of venous obstruction proximal to the level of interrogation.

A normal artery has a triphasic signal. The waveform becomes biphasic in mild-moderate stenosis and monophasic in severe stenosis.

The image given below shows the normal triphasic signal from an artery.



Monophasic waveform:



Solution to Question 6:

Rigler's triad is seen in gallstone ileus and consists of

• Pneumobilia

• Small bowel obstruction

• Ectopic gall stone (usually in the right iliac fossa)

This should not be confused with Rigler's sign and Rigler's notch sign.

Rigler's Sign:

Seen in pneumoperitoneum. Here, the air is seen on both sides of the bowel wall unlike normally, where the air is seen only in the intraluminal side of the bowel.

Rigler's notch sign:

Refers to indentation seen in the border of a solid lung mass, which is thought to represents a feeding vessel, thus suggesting the presence of bronchial carcinoma. But, this is not specific and can also be seen in granulomatous infections.

Solution to Question 7:

MRI is a technique that produces topographic images by means of magnetic fields and radiofrequency waves.

The ionizing radiations

• Alpha

• Beta

• Gamma

• X-Rays

The non-Ionizing radiations are:

• Radiofrequency waves

• UV rays

• Microwaves

Alpha and beta rays are particulate radiation (particles) whereas gamma and x-rays are electromagnetic rays. All electromagnetic waves have a constant velocity of 3×108 m/s and no mass. The electromagnetic spectrum, from low frequency to high frequency consists of:

Radiowaves < Microwaves < Infrared < Visible light < Ultraviolet < X-rays < Gamma rays

Note: W. C. Roentgen is the father of X-ray and the first medical X-ray by Roentgen was of his wife Anna Bertha's left hand. Physicists Peter Mansfield and Paul Lauterbur developed MRI-related imaging techniques in medicine and they were awarded the Nobel prize in 2003.

Solution to Question 8:

An empty thecal sac sign is seen in arachnoiditis.

Adhesive arachnoiditis can be diagnosed if the empty thecal sac indication is observed. This occurs due to inflammation, which leads to clumping of nerve roots. This leads to an empty sac appearance. It is best observed on T2-weighted MRI scans of the lumbar spine when the thecal sac appears empty.



Solution to Question 9:

The above X-ray clearly shows the presence of subdiaphragmatic free gas in the right upper quadrant. This indicates pneumoperitoneum.

Pneumoperitoneum is the presence of free gas within the peritoneal cavity but outside the viscera. The most common cause of pneumoperitoneum is bowel perforation.

• Gastric and posterior duodenal perforations are seen in long-standing cases of peptic ulcers.

• Intestinal perforations are seen with ischemia, inflammation, diverticula, or obstruction.

• Other causes also include iatrogenic, penetrating injuries, prolonged mechanical ventilation, peritoneal dialysis, etc.

Signs of pneumoperitoneum:

• Bowel-related signs

• Double wall sign/Rigler's sign/gas-relief sign- Air is seen on both sides of the bowel wall.

• Telltale triangle sign/triangle sign - Triangle of gas formed between three loops of the bowel or between the abdominal wall and two loops of the bowel.

• Peritoneal ligament-related signs

• Football sign- Air outlines the confines of the peritoneal cavity

• Falciform ligament sign- Air outlines the falciform ligament

• Lateral umbilical ligament sign/inverted "V" sign- Air outlines the lateral umbilical ligament

• Urachus sign - Air outlines the urachus

• Right upper quadrant signs

• Cupola sign- Air outlines the median subphrenic space

• Fissure for ligamentum teres sign- Air outlines the ligamentum teres

• Hepatic edge sign- Cigar-shaped free-air pocket in the subhepatic region. It tracks superomedial following the contours of the liver

• Lucent liver sign - Reduction of hepatic radiodensity due to accumulation of free gas anterior to the liver

• Morison's pouch sign / Doge cap sign - Triangular/ crescentic gas lucency bound by the 11th rib

• Periportal free gas sign.

Solution to Question 10:

The given X-ray shows cervical rib.

Patients with a cervical rib usually show a lump and tenderness in the supraclavicular fossa. The pallor and paresthesias are due to vascular and neural compressions respectively. A radiograph of the neck is useful in detecting a cervical rib.

However, the absence of a rib on the x-ray does not rule out the possibility of a cervical rib. CT and MRI are more sensitive.



Solution to Question 11:

The procedure depicted in the image represents myelography.

Myelography is a procedure used to evaluate spinal conditions. X-ray myelography can be used to evaluate the spinal canal in patients with spondylolisthesis. While MRI has largely replaced X-ray myelography it is done when there is a contraindication to MRI (Eg; non-compatible pacemaker, severe claustrophobia).

Radiology AIIMS 2019 (May & Nov)

Question 1:

All lung carcinomas for diagnosis requires CECT thorax but MRI is useful in which of the following:

a) Small cell lung carcinoma

b) Adenocarcinoma

c) Carcinoid

d) Pancoast tumor

Question 2:

The following image of a FAST scan in a trauma patient suggests:



a) Positive fast

b) Negative fast

c) Equivocal fast

d) Pneumoperitoneum

Question 3:

The given USG of lung shows:



a) Stratosphere sign

b) Sea shore sign

c) Troposphere sign

d) Mirror image artifact

Question 4:

An 8-year old child presents with the history of cough for 7 months. He also gives a history of haemoptysis for 1 month and fever for 4 days. The X-ray is given below. What is the diagnosis?

a) Lung abscess

b) Consolidation

c) Hydatid cyst

d) Thoracic duct malformation

Answer Key

Question No.	Correct Option
1	d
2	a
3	b
4	b

Detailed Explanations

Solution to Question 1:

MRI is useful in Pancoast tumor because of the associated soft tissue infiltration, extent of which can be best visualised using an MRI

Pancoast syndrome/superior sulcus tumor results from a local extension of a tumor growing in the apex of the lung, with the involvement of C8, T1, and T2 nerve roots.

It presents with pain that radiates to the ulnar distribution of the arm, often with the destruction of the 1st and 2nd ribs. Patients with the central or endobronchial growth of the primary tumor may present with a cough, hemoptysis, wheeze, stridor, dyspnea, or post-obstructive pneumonitis.

Regional spread can cause

• Tracheal obstruction,

• Esophageal compression with dysphagia

• Recurrent laryngeal paralysis with hoarseness

• Phrenic nerve palsy with the elevation of the hemidiaphragm and dyspnea

• Sympathetic nerve paralysis with Horner's syndrome (enophthalmos, ptosis, miosis, and anhydrosis).

Solution to Question 2:

The image of FAST in a trauma patient suggests Positive FAST as there is a fluid collection in the abdominal cavity. The thin anechoic area located between the liver(on the left) and the Kidney(on the right) is a collection of fluid visualized in this RUQ Sonogram image.

FAST stands for Focused assessment with sonography for trauma. It is a rapid diagnostic examination to assess patients with potential thoracoabdominal injuries. It is performed within 2--4□ minutes.

Technique:

• Patient in the supine position

• 3.5-5.0 MHz convex transducer

• Following regions are scanned :

• Subxiphoid: assess for pericardial effusion and left lobe liver injuries

• Right upper quadrant: assess for right liver injuries, right kidney injury, and Morison pouch

• Left upper quadrant: assess for splenic injury and left kidney injury

• Suprapubic region: assess the bladder and pouch of Douglas

An "extended FAST or eFAST" scan may be performed: left and right thoracic views to assess for pneumothorax and haemothorax.

Sequentially order of FAST is:- pericardial sac, the Right Upper Quadrant (RUQ), Left Upper Quadrant (LUQ), and pelvis.

4 ‘Ps’ are evaluated in sequence: Pericardial sac → Perihepatic region → Perisplenic region → Pelvis.

Note:-

In Pneumoperitoneum, air is collected in the peritoneal cavity which is usually hyperechoic not anechoic and would appear white in the ultrasound image.

Solution to Question 3:

The given image shows the seashore sign seen in M-mode imaging of the normal lung.

M-mode is defined as the time-motion display of the ultrasound wave along a chosen ultrasound line. With M-Mode, even rapid motions can be displayed, measured, and recorded.

In the M-mode of USG, the motionless portion of the chest above the pleural line creates horizontal ‘waves', and the sliding below the pleural line creates a granular pattern, the ‘sand’. The resultant picture is one that resembles waves crashing in onto the sand and is therefore called the ‘seashore sign’ and is present in a normal lung.

Option A: Stratosphere sign or barcode sign is a finding in the M-mode seen as parallel horizontal lines above and below the pleural line. This is seen in pneumothorax.

Option C: There is no named sign called troposphere sign.

Option D: Mirror image artifact is seen when there is a highly reflective surface (eg: Diaphragm) in the path of the primary beam.

Solution to Question 4:

The given X-ray shows opacities in the right middle lobe and history of fever, cough and haemoptysis suggesting consolidation due to lobar pneumonia.

Radiology AIIMS 2020 (May and Nov INI-CET)

Question 1:

Which of the following statements is false regarding acute radiation syndrome?

a) GI manifestations occurs at a lower radiation dose than bone marrow depression

b) Bone marrow suppression appear at doses greater than 1 Gy

c) Bone marrow suppression occurs earlier than neurovascular effects

d) GI manifestations occur at a lower radiation dose than neurovascular effects

Question 2:

A 46-year-old male presented with chronic back pain and neck stiffness. X-ray cervical spine is given below. What is the diagnosis?

a) Ankylosing spondylitis

b) Diffuse idiopathic skeletal hyperostosis

c) Klippel feil syndrome

d) Sprengel deformity

Question 3:

An 18-year-old girl presented with seizures and her radiological image is as given below. Knowing that the MR spectroscopy of the lesion showed elevated choline, alanine, and lactate, what is the likely diagnosis?



a) Cerebral toxoplasmosis

b) Neurocysticercosis

c) Cryptococcosis

d) Multiple tuberculomas

Question 4:

Which of the following is a finding of large bowel obstruction?

a) Dilated loops of intestine with plica circularis

b) Dilated loops with haustrations

c) String of pearls sign

d) Central dilated bowel loops

Question 5:

A 25-year-old male with Marfanoid features presented with sudden onset breathlessness. The chest X-ray is given below. What is the likely diagnosis?



a) Pneumothorax

b) Pleural effusion

c) Consolidation

d) Fibrosis

Question 6:

A 22-year-old male was brought to the emergency room following a road traffic accident. On examination, his GCS was E2M2V3. He had no fractures and no hydrocephalus or raised intracranial pressure. Non-contrast CT was performed and was found to be normal. The likely diagnosis is

a) Diffuse axonal injury

b) Post concussion syndrome

c) Extra dural hemorrhage

d) Herniation of brain

Answer Key

Question No.	Correct Option
1	a
2	b

3	b
4	b
5	a
6	a

Detailed Explanations

Solution to Question 1:

GI manifestations do not occur at a lower radiation dose than bone marrow depression.

Stages of Acute Radiation Syndrome (ARS)

The prodromal phase manifests with non-specific symptoms like nausea, vomiting, anorexia. Early and persistent symptoms predict severity.

Latent phase: The patient looks and feels generally healthy for a few hours or even up to a few weeks. In severe cases, this phase may be absent.

Manifest illness phase: Symptoms are seen as per the radiation dose. Tissues with greater rates of cellular division are most radiosensitive, and hence the hematopoietic and gastrointestinal systems are the most affected.

Recovery phase or Death: The recovery phase will be seen after the manifest stage when the patient recovers from the acute exposure. The time taken for recovery will depend on the severity of the injury and the organ system affected.

Radiation dose	Sub-syndrome	Features
~1-2 Gy	Hematopoietic	Bone marrow suppression
~6-10 Gy	Gastrointestinal	Nausea, vomiting, gastrointestinal bleeding, malabsorption,
~20Gy	Neurovascular	irritability, altered mental status, seizures, coma

Solution to Question 2:

The given clinical scenario and the radiograph showing ossification of the anterior longitudinal ligament, in front of the C2 to C7 cervical vertebral bodies, point towards a diagnosis of Diffuse Idiopathic Skeletal Hyperostosis (DISH). It is also referred to as Forestier disease.

DISH is a common disorder characterized by bone and calcium formation at the sites of ligamentous insertions. It occurs in the middle-aged or elderly. The radiograph shows calcification of anterior and sometimes posterior longitudinal ligaments appearing as flowing or dripping-like

candle wax. It is usually asymptomatic but the calcification might cause compression of the spinal cord and can be better visualized on CT and MRI. In contrast to ankylosing spondylitis, the sacroiliac joints are usually spared.

Solution to Question 3:

The given radiological image and history of seizures suggests a diagnosis of neurocysticercosis.

Neurocysticercosis presents with seizures, hydrocephalus, raised ICP, and may even cause signs of chronic meningitis. The given image shows multilobulated cystic lesions (with scolex) in the subarachnoid space, typical parenchymal brain calcifications are suggestive of neurocysticercosis caused by T. solium (Cysticercosis cellulose).

• Cyst with a dot sign: seen in both MRI and CT and the dot represents the eccentrically placed scolex of the parasite.

MR Spectroscopy features of neurocysticercosis include elevated lactate, alanine, succinate, and choline levels and reduced levels of N -acetyl aspartate and creatine. Escobar's pathological stages describe 4 stages - vesicular, colloidal vesicular, granular nodular, and nodular calcified.

Other options

• Solid or ring enhancement is typical in the post-contrast images of tuberculoma. A central focus of calcification with a ring of peripheral enhancement (the "target sign") is also described in tuberculosis but is not specific to it.

• Cerebral toxoplasmosis appears as multiple hypodense regions predominantly in the basal ganglia and at the corticomedullary junction. It can be associated with ring enhancing lesions as well.

• Cryptococcosis is usually associated with normal scans but may show cerebral atrophy, hydrocephalus or mass lesions as well.

Solution to Question 4:

Dilated bowel loops with haustrations is a feature of large bowel obstruction.

In large bowel obstruction, the colon is distended proximally with the distal portion collapsed. The colon is considered dilated if it is more than 6 cm in diameter while the cut off for caecum is 9cm.

Other options:

Dilated loops of the intestine with plica circularis (valvulae conniventes) and predominantly central loops are features of the small bowel obstruction. The small bowel is considered dilated if it is more than 3 cm in diameter.

The string of pearls sign is also a feature of small bowel obstruction seen in plain radiographs or CT. It represents a row of gas bubbles trapped between the valvulae conniventes.

Differences in small and large bowel obstruction on radiography are listed below:

	Small bowel	Large bowel
Valvulae conniventes	Present in jejunum	Absent
No of Loops	Many	Few
Distribution of loops	Central	Peripheral
Haustra	Absent	Present
Diameter	3-5 cm	>5 cm

Solution to Question 5:

The chest X-ray demonstrating tracheal deviation and displacement of the heart away from the affected side along with the partial collapse of the left lung in the above condition points to a diagnosis of left-sided pneumothorax.

Marfan's syndrome is one of the causes for seconday spontaneous pneumothorax in adults.



Pneumothorax refers to the collection of air in the pleural space.

Chest radiograph in spontaneous pneumothorax demonstrates:

• Thin, sharp white line, demarcating the visceral pleural edge

• Absence of lung markings seen peripheral to this line

• Peripheral space radiolucent compared to adjacent lung

• Complete lung collapse, as well as subcutaneous emphysema and pneumomediastinum, may be seen

• Deep sulcus sign (on supine chest radiograph)

Deep sulcus sign: On a supine position(common in the ICU), air collects anteriorly and basally, within the non-dependent portions of the pleural space, abnormally deepening the costophrenic angle. This sign is indicative of pneumothorax in such patients and may be the only visible clue, especially when patient can not be imaged upright.

The image below is plain chest x-ray showing pneumothorax with deep sulcus sign (shown by arrow):



When there is a progressive accumulation of air intrapleurally in the thoracic cavity (caused by a valve effect during inspiration and expiration), it is called Tension pneumothorax. It can result in compression of mediastinal structures and compromise cardiopulmonary function. It is life-threatening and requires emergency intervention.

Solution to Question 6:

The given clinical scenario of an RTA with no imaging abnormality along with a disproportionate clinical manifestation suggests diffuse axonal injury.

It is caused due to stretching of axons in subcortical white matter due to sudden deceleration injuries. It is usually associated with mild nonspecific changes in NCCT. It is visualized on T2 MRI characterized by multiple lesions showing microbleeds or linear hypointensities. It may also show disruption of white matter tracts on tractography.

Other options

Post-concussion syndrome usually follows mild traumatic brain injury. It does not have a severe clinical status as seen in this patient.

Extradural hemorrhage is produced usually results secondary to trauma to head. The bleeding is usually arterial and blood collects between outer dura and inner table of the skull. It can very well be visualized on CT.

A cerebral herniation is defined as a shift of cerebral tissue in an adjacent space. Intracranial brain herniation can be classified into 3 types - subfalcine, transtentorial and tonsillar. A distinct

clinical picture can be produced as a result of compression of neurovascular structures by the herniating part. CT is the imaging modality of choice because it can be done rapidly.

Radiology NEET 2018

Question 1:

In which of the following conditions is the given X-ray finding seen?



a) Esophageal atresia

b) Esophageal stenosis

c) Reflux esophagitis

d) Barrett's esophagus

Question 2:

Identify the structure marked in the CT scan shown below:



a) Descending aorta

b) Ascending aorta

c) Superior vena cava

d) Right pulmonary artery

Question 3:

What condition causes the skull appearance as in the given radiograph?

a) Hyperparathyroidism

b) Multiple myeloma

c) Hyperthyroidism

d) Paget's disease

Question 4:

What dye is used for the diagnosis of esophageal perforation?

a) Gastrograffin

b) Iohexol

c) Gadolinium

d) Barium sulphate

Question 5:

A 50-year-old patient has come with sudden onset of chest pain and his CT scan shows the following picture. What is the diagnosis?



a) Pulmonary embolism

b) Aortic dissection

c) Aortic aneurysm

d) Myocardial infarction

Question 6:

The following scan is of a young woman on oral contraceptives. The most likely diagnosis is:



a) Hepatocellular carcinoma

b) Metastatic infiltrates

c) Hepatic adenoma

d) Hemangioma

Question 7:

A child presents with a left-to-right shunt and high-output cardiac failure. The following is his MRA. What is the most likely diagnosis?



a) Vein of Galen malformation

b) Dandy Walker syndrome

c) Pneumocephalus

d) Venous sinus thrombosis

Question 8:

What is the diagnosis based on the given image?

a) Uterus didelphys

b) Bicornuate uterus

c) Unicornuate uterus

d) Septate uterus

Question 9:

The Bragg peak effect is pronounced in:

a) X ray

b) Proton

c) Neutron

d) Electron

Question 10:

The image seen below shows_____.

a) Magnetic Resonance Imaging

b) Positron Emission Tomography

c) Hysterosalpingography

d) Laparoscopic chromopertubation

Question 11:

Identify the artery marked by blue arrow.



a) Superior mesenteric artery

b) Inferior mesenteric

c) Celiac artery

d) Intestinal artery

Question 12:

What is the probable diagnosis based on the image of the scan given below?



a) Pituitary apoplexy

b) Pituitary macroadenoma

c) Rathke cleft cyst

d) Craniopharyngioma

Answer Key

Question No.	Correct Option
1	c
2	c
3	a
4	b

5	b
6	c
7	a
8	c
9	b
10	c
11	a
12	d

Detailed Explanations

Solution to Question 1:

The given image is a barium swallow X-ray showing a feline esophagus suggestive of reflux esophagitis.

The feline esophagus is also known as esophageal shiver. These folds are 1-2 mm thick and run horizontally around the entire circumference of the esophageal lumen. It is thought to be due to the contraction of the muscularis mucosae of the esophagus. This results in the shortening of the esophagus and the bunching up of the mucosa in the lumen.

The appearance is almost always associated with active gastroesophageal reflux. The findings are transient, seen following reflux and not during swallowing. The appearance is confined to the distal two-thirds of the thoracic esophagus.

It can also be seen in hiatus hernia and eosinophilic esophagitis. Here, the folds are permanent and associated with strictures.

Solution to Question 2:

The structure marked in the given CT thorax image is the superior vena cava.

The image below depicts all the major structures labelled in the CT thorax image:



Solution to Question 3:

The given X-ray of the skull shows multiple radiolucent lytic lesions, typical of the salt and pepper appearance (pepperpot appearance). This is seen in hyperparathyroidism.

In hyperparathyroidism, there is resorption of the trabecular bone of the skull (which causes the radiolucent areas) in combination with cystic areas of osteopenia. There is also a loss of definition between the inner and outer tables of the skull.

Other options:

Option B: In multiple myeloma, multiple well-defined "punched out" lytic lesions exist.

Option C: Untreated hyperthyroidism for several years can lead to osteoporosis of bones.

Option D: In Paget's disease, the pelvis, spine, skull, and proximal long bones are most frequently affected. Radiological skull changes include:

• Osteoporosis circumscripta: Large, well-defined lytic lesion

• Cotton wool appearance: Mixed lytic and sclerotic lesions of the skull

• Diploic widening: Both inner and outer calvarial tables are involved, with the former usually more extensively affected

• Tam o'Shanter sign: Frontal bone enlargement, with the appearance of the skull falling over the facial bones, like a Tam o'Shanter hat

Solution to Question 4:

Iohexol is the contrast medium used for the diagnostic work-up for a suspected esophageal perforation.

Although gastrograffin (water-soluble) was used previously, it has a high chance of causing aspiration pneumonitis and pulmonary edema consequently. Hence it is not preferred nowadays. (Option A)

Barium contrast is not used in esophageal perforation as it causes barium-related inflammation of the mediastinum. (Option D)

Given below is a plain chest radiography with a water-soluble contrast swallow, showing contrast leakage in a patient with spontaneous rupture of the esophagus.

Contraindications to the use of barium contrast	Alternative dyes used
Tracheoesophageal fistula	Dianosil- water-soluble, non-i onic agent
Esophageal perforation	Iohexol
Perforation peritonitis	Gastrograffin / Iohexol



Solution to Question 5:

The given CT scan showing a double lumen and dilatation of the aorta is suggestive of an aortic dissection.

In this condition, blood enters the space below the intima of the aorta. This results in the formation of a true lumen and false lumen. The true lumen is often the smaller of the two because of being compressed by the false lumen, as shown below.



The beak sign in aortic dissection is a reliable sign to identify the false lumen. It refers to the acute angle formed at the edge of the false lumen in aortic dissection in axial cross-section. It is formed by the borders of the outer aortic wall and the intimal flap, and maybe partially thrombosed (blunted beak).



Solution to Question 6:

The image given is of hepatic adenoma. It is the most frequent hepatic tumour in young women on oral contraceptive pills.

Hepatic adenomas, also referred to as hepatocellular adenomas, are benign liver tumours. They are well-defined vascular lesions, which are generally solitary. They are almost always seen in women between the age of 25 and 50 years and are associated with the use of the oral contraceptive pill.

On CT, uncomplicated adenomas are usually homogeneous, with a well-defined margin. They enhance markedly and uniformly during the arterial phase and then merge with the surrounding liver during the portal phase.

Hepatic adenoma on the CT scan is shown below.



Other options:

Option D: On contrast-enhanced CT scan, hemangiomas show asymmetrical, nodular peripheral enhancement that is isodense with large vessels and progressive centripetal enhancement. Liver hemangioma on CT scan is shown below.

Option B: Most metastatic deposits appear as areas of low attenuation on unenhanced CT and portal-phase CT. Hypervascular tumours are often visible as low-attenuation lesions on unenhanced images and may enhance transiently in the arterial phase, some becoming invisible in the portal phase. Metastatic infiltrates in the liver in the portal phase on CT scan are shown below.



Option A: Hepatocellular carcinoma (HCC) typically shows a decreased signal on T1-weighted images and a moderately increased signal on T2-weighted images, with internal heterogenecity. MRI is regarded as the most sensitive imaging modality for detecting HCC. Given below is the MRI image of HCC:



Solution to Question 7:

The MR-Angiography shown is of Vein of Galen Malformation.

Vein of Galen aneurysmal malformations (VGAM's), also known as median prosencephalic arteriovenous fistula, are uncommon intracranial anomalies that tend to present in early childhood with features of a left-to-right shunt and high-output cardiac failure.

The vein of Galen, also known as the great cerebral vein or great vein of Galen, is a short trunk formed by the union of the two internal cerebral veins and basal veins of Rosenthal. It lies in the quadrigeminal cistern. It curves backwards and upward around the posterior border of the splenium of the corpus callosum and drains the anterior and central regions of the brain into the sinuses of the posterior cerebral fossa.

The following image shows the Vein of Galen:



Solution to Question 8:

The given image shows the hysterosalpingogram of a unicornuate uterus.

The endometrial cavity usually assumes a fusiform (banana type) shape, tapering at the apex and draining into a single fallopian tube. The uterus is generally shifted off the midline.

Solution to Question 9:

The Bragg peak effect is pronounced in protons.

The proton is a heavy and charged particle that gradually loses its speed as it interacts with human tissue. It is easily controlled and delivers its maximum dose at a precise depth and can go as far as 32 cm. The absorbed dose increases very gradually with greater depth and lower speed, suddenly rising to a peak when the proton is ultimately stopped. This is known as the Bragg peak.

The behavior of the proton can be precisely determined, and the beam can be directed, allowing the Bragg peak to occur within the tumor site. Immediately following this burst of energy, the protons cease radiating and come to a complete stop. This property of proton therapy enables targeting tumors within the body, delivering radiation directly to the tumor while sparing the surrounding healthy cells, offering a much less invasive alternative to treat cancer.

The Bragg curve shown below plots the energy loss of ionizing radiation during its travel through matter.



Solution to Question 10:

The image given is of the findings seen in hysterosalpingography. The given image shows bilateral retort-shaped fallopian tubes, indicating bilateral hydrosalpinx.

Hysterosalpingography (HSG) is a technique used for visualization of the uterine cavity and fallopian tubes by the injection of a radioopaque dye, and visualization of the dye by fluoroscopy as it passes through the uterine cavity and fallopian tubes.

It is best performed between the end of the menstrual period and ovulation (i.e. 9th or 10th day of the cycle). 10-15ml of the dye is injected using a Leech-Wilkinson cannula, Rubin's cannula, or Foley's catheter. If the tubes are patent, the dye will spill out of the tubal Ostia and smear the adjacent bowel.

It is used to detect:

• Tubal block: This can be due to strictures (fibrotic block), inspissated amorphous material, or spasms. The tubal spasm can occur in an apprehensive woman, and this is prevented by premedication with atropine and analgesia.

• Uterine anomalies

• Uterine polyps

• Uterine synechiae

• The feasibility of tuboplasty, by studying the location and extent of tubal pathology.

Contradictions:

• Presence of genital tract infection and bleeding

• Genital tuberculosis, due to the risk of spreading infection.

• Suspected pregnancy

• Allergy to the dye

• Premenstrual phase (there is a risk of embolism and endometriosis if HSG is done during this phase).

The image given below shows the appearance of a normal hysterosalpingogram.

Laparoscopic chromopertubation is a laparoscopic procedure for establishing the patency of fallopian tubes or verifying the findings of blocked tubes on HSG. Methylene blue is injected through the cervix to visualize the free spill. The absence of spill indicates blocked tubes.

Solution to Question 11:

The artery marked by the blue arrow is the superior mesenteric artery (SMA), which gives rise to various branches that supply the small intestines, cecum, ascending, and part of the transverse colon.

The CT also depicts nutcracker syndrome, a left renal vein entrapment syndrome characterized by extrinsic compression of the left renal vein (marked by a red arrow), most commonly between the abdominal aorta and superior mesenteric artery. This syndrome is one of the postulated reasons for left-sided varicoceles being more common than the right side.

Solution to Question 12:

The given MRI of the brain demonstrates a craniopharyngioma, which is a suprasellar tumour that arises from remnants of Rathke's pouch.

The two variants of craniopharyngioma are:

• Adamantinomatous type (common in children): Predominantly solid with cystic components. Diagnostic features include compact lamellar keratin ("wet keratin") and areas of dystrophic calcification.

• Papillary type (common in adults): Contains both solid sheets and papillae lined by well-differentiated non-keratinized squamous epithelium.

These tumours are usually large and heterogeneous, displaying solid and cystic components. The cysts contain a cholesterol-rich, thick brownish-yellow fluid that is compared to 'machine oil'.

Clinical features include headaches (most common), visual abnormalities - decreased acuity and bitemporal hemianopia, GH deficiency, growth failure, delayed sexual maturation due to pituitary hypofunction, and polyuria (central diabetes insipidus).

Radiologically:

• CT may show calcifications associated with the solid and cystic wall components

• MRI is the imaging modality of choice which shows:

Craniopharyngiomas tend not to expand the pituitary fossa unless they become huge, which is a differentiating feature from pituitary macroadenomas.

Other options:

Option A: Pituitary apoplexy is an acute clinical condition associated with the rapid expansion of the pituitary gland, which is evident on an MRI as enhancement around a non-enhancing infarcted centre.

Option B: Pituitary macroadenoma (with cystic degeneration or necrosis) shows indentation at the diaphragma sellae, giving rise to a snowman/figure of eight appearance/cottage loaf. Calcification is often absent. The image below shows the snowman appearance:

Cysts	
T1	Iso to hyperintense to the bra in (due to high protein conte nt "machinery oil cysts")
T2	Mostly hyperintense
Solid compon ent	
T1	Vivid enhancement
T2	Variable or mixed

Snowman appearance – Pituitary adenoma



Option C: Rathke cleft cyst shows no solid or enhancing component. Calcification is rare. It is unilocular and the majority are completely or mostly intrasellar.

Radiology NEET 2019

Question 1:

Match the following intracranial bleed:

A)Epidural hematoma	i) 
B)Subdural hematoma	ii) 
C)Subarchnoid haemorrhage	iii)
D)Intracerebral haemmorrhage	iv)

a) A -i B-ii C- iii D- iv

b) A -ii B-iv C-i D-iii

c) A-iv B-iii C -ii D-i

d) A -iv B-ii C-iii D-i

Question 2:

The following appearance in IVP is caused by:

a) Horse shoe kidney

b) Ureterocoele

c) Bladder tumour

d) Posterior urethral valve

Question 3:

Which of the following investigations is shown in the radiograph?

a) Barium swallow

b) Barium meal

c) Barium meal follow through

d) Barium enema

Question 4:

What is the best imaging study for earliest diagnosis of cerebral infarct?

a) FLAIR MRI

b) Non-contrast CT

c) CECT

d) Diffusion weighted MRI

Question 5:

During evaluation of PUO, a patient's chest X-ray is as seen below. What is your diagnosis?

a) Miliary tuberculosis

b) Pulmonary edema

c) Necrobiotic nodules

d) Tuberculoma

Question 6:

Identify the structure marked in the following MRI scan:

a) Midbrain

b) Cerebellum

c) Brainstem

d) Cerebrum

Question 7:

Which of the following is a water soluble contrast?

a) Iohexol

b) Barium

c) Bromium

d) Calcium

Question 8:

All of the following could be the cause of the following X-ray finding except:



a) Duodenal ulcer perforation

b) Post laparoscopy

c) Chilaiditi syndrome

d) Acute intestinal obstruction

Answer Key

Question No.	Correct Option
1	b
2	b
3	b
4	d
5	a
6	b
7	a
8	d

Detailed Explanations

Solution to Question 1:

Epidural hematoma:

Noncontrasted axial computed tomography scan showing a lenticular area of increased density in left parietal region.

• Typically occur in the groove of the middle meningeal artery associated with temporal bone fractures that have resulted in a tear of middle meningeal artery.

• They tend to not to cross suture lines of the skull.

Subdural hematoma:



A non contrasted computed tomography scan of acute subdural hematoma shows a cresentric area of increased density in the right posterior parietal region between brain and the skull. An area of intraparenchymal haemorrhage also seen. In addition, mass effect causes a midline shift to left.

• They can cross suture lines but they do not cross the tentorium or falx.

• Can be acute or chronic

Subarachnoid haemorrhage:



A non contrasted axial computed tomography scan shows the blood as areas of increased density near the base of the brain shows blood in Texaco star pattern formed blood radiating from suprasellar cistern into the sylvian fissures and anterior interhemispheric fissure.

• Most often accompanied by severe sudden onset headache.

• Can really be visualised in acute stage, when blood is radiographically dense (whiter) than the CSF.

Intracereberal haemorrhage:



Non contrast computed tomography shows a large area of fresh blood appearing white in the region of right thalamus. Blood is denser than CSF so settles dependently.

This settling is not seen with subarachnoid or interparenchymal blood.

Solution to Question 2:

The given intravenous pyelography (IVP) depicts the cobra head or adder head deformity or spring onion sign seen in bilateral ureterocele. These signs refer to a dilated, contrast-filled distal ureter surrounded by a thin rim of the ureteric wall and bladder mucosa.

Ureterocoele is a cystic enlargement of the intramural portion of the ureter, more commonly seen in females than males. It is thought to be caused due to ureteric orifice atresia. They are similar to ectopic ureters, which typically drain the upper pole of the kidney and are associated with dysplastic or non-functional renal tissue.

If they present clinically, ureteroceles often manifest with urinary tract infections (UTIs) in children, while in adults, they are more likely to present with stones in the lower ureter. Ureterocoeles can enlarge and obstruct the bladder neck or the contralateral ureteric orifice.

The classic sign of a ureterocele on intravenous urography is a ‘cobra head’, ‘adder head’, or 'spring onion sign'. On ultrasonography, it can be seen as a fluid-filled cystic mass in the bladder.

The treatment of simple ureteroceles is surgical excision with re-implantation of the ureter.

Solution to Question 3:

The given image is a double-contrast radiograph of a barium meal. It is used to examine the lower half of the esophagus, stomach, and entire duodenum. It is used in identifying ulcers, small fleshy lumps (polyps), tumours.

Barium meal follow-through is useful in visualising small bowel pathology and evaluating the function of the small bowel. It is performed following a barium meal examination where X-rays are taken 10-15 minutes after the intake of barium. This allows time for the barium to reach the small intestine. It is inferior to enteroclysis enema, where contrast is administered through a nasogastric tube.

GI Imaging	Preparations of barium	Structures best visualized
Barium swallow	Barium paste	Esophagus
Barium meal	95% $BaSO_4$	Stomach and duodenum
Barium meal follow-through	50% $BaSO_4$	Small bowel
Barium enema	25% $BaSO_4$	Large bowel

The below radiograph is a double-contrast barium enema with a normal large bowel structure. The contrast is administered through a rectal tube and a series of radiographs are obtained.

Solution to Question 4:

The best imaging modality for the early diagnosis of cerebral infarction is diffusion-weighted MRI (DWI), which can detect infarcts as early as 30 minutes of onset and identify cytotoxic oedema within a few hours of ischemic stroke.

However, the initial investigation in suspected stroke cases is non-contrast CT due to its rapid acquisition time, widespread availability, and cost-effectiveness. It helps quickly rule out hemorrhagic stroke and facilitates the timely administration of intravenous tPA in eligible patients.

Acute MCA territory infarct on CT is shown below:

The image below is of a DWI MRI of a patient clearly showing the extent of the infarct.



Solution to Question 5:

The given chest radiography image shows miliary tuberculosis.

It is the widespread dissemination of Mycobacterium tuberculosis, via hematogenous spread.

Classic miliary TB is defined as millet-like (mean, 2 mm; range, 1-5 mm) seeding of TB bacilli with uniform size and distribution in the lung, as seen on chest radiography. This particular sign is called the miliary shadow which is also seen in sarcoidosis and histoplasmosis.

Sputum-smear microscopy is negative in most cases.

Physical findings include hepatomegaly, splenomegaly, and lymphadenopathy

Bronchoalveolar lavage and transbronchial biopsy are used to provide bacteriologic confirmation

Solution to Question 6:

The given image shows an MRI of the sagittal section of the brain. The area marked with a red arrow is the cerebellum.

The other areas also visualized in the image are :



Solution to Question 7:

Iohexol is a nonionic, water-soluble radiographic contrast medium.

It is used to visualize the following structres:

• Arteries

• Veins

• Ventricles of the brain

• The urinary system

• Joints

It can be used during computer tomography (CT scan) as well. It can be given orally, intravenously, or can be injected into the body cavity.

Solution to Question 8:

The given X-ray shows air under the diaphragm, indicative of pneumoperitoneum, which is not caused by acute intestinal obstruction.

Causes of pneumoperitoneum include:

• Bowel perforation: Gastric and posterior duodenal perforations in long-standing cases of peptic ulcers., ischemic bowel, necrotizing enterocolitis

• Small bowel perforation secondary to trauma or penetrating injuries

• Post-surgery/laparoscopy

• Causes other than perforation are:

• Chilaiditi syndrome - the intestine is seen between the liver and diaphragm giving a false impression of free air

• Sub-phrenic abscess

• Postoperatively, air may be present in the abdomen for up to 7 days

An intestinal obstruction would show multiple dilated loops on X-ray with possible air-fluid levels. Only when the obstruction becomes severe and there is a perforation, pneumoperitoneum occurs.

The image given below is an abdominal radiograph showing multiple dilated loops with air-fluid levels suggestive of intestinal obstruction.



Radiology NEET 2020

Question 1:

A 56-year-old■ female patient presented with progressive dysphagia for solids, loss of weight, and loss of appetite. A barium swallow done is given below. What is the most probable diagnosis?



a) Carcinoma esophagus

b) Diffuse esophageal spasm

c) Esophageal leiomyoma

d) Esophageal web

Question 2:

A 30-year-old woman presents with persistent hypertension, chest pain and cramping in her legs, particularly during exertion. A chest radiograph was taken, which reveals the condition, as shown in the image below. What is the likely■ underlying cause of her hypertension?



a) Coarctation of aorta

b) Essential hypertension

c) Aortic aneurysm

d) Aortaarteritis

Question 3:

A 51-year-old man complains of abdominal pain and blood-stained stools. The barium study of his intestines shows the following finding. What is the most likely diagnosis?



a) Intussusception

b) Volvulus

c) Diverticulitis

d) Ulcerative colitis

Question 4:

A cystic lesion in the suprasellar region with calcification is seen on the MRI. What is the most likely diagnosis?



a) Craniopharyngioma

b) Pituitary adenoma

c) Meningioma

d) Oligodendroglioma

Question 5:

Personal monitoring of radiation is done by:

a) TLD badge

b) Collimators

c) Linear accelerator

d) Grid

Question 6:

The appearance of USG at saphenofemoral junction, showing common femoral vein, common femoral artery and the great saphenous vein is called as:

a) Mickey mouse sign

b) String sign

c) Tillaux sign

d) Stemmer sign

Question 7:

Plethoric lung fields are seen in:

a) Total Anomalous Pulmonary Venous Connection

b) Tricuspid atresia

c) Ebstein anomaly

d) Tetrology of Fallot

Question 8:

A patient complains of neck stiffness and severe headache. The CT image shown below is suggestive of which of the following conditions?



a) Meningitis

b) Subarachnoid hemorrhage

c) Intraparenchymal hemorrhage

d) Hemorrhagic stroke

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	a
5	a
6	a
7	a
8	b

Detailed Explanations

Solution to Question 1:

A patient with the given history and barium swallow findings of irregular narrowing, mucosal destruction, and shouldered margins of esophagus, are suggestive of carcinoma esophagus.

Esophageal carcinoma is more common in men, with tobacco use and alcohol being the strongest risk factors. The squamous cell carcinoma variant usually affects the upper 2/3rd, and the adenocarcinoma variant usually affects the lower 1/3rd of the esophagus. Clinical features include progressive dysphagia (most common), regurgitation of food, vomiting, odynophagia and weight loss.

On barium swallow, carcinoma esophagus presents with irregular narrowing and ulceration of esophagus. Flexible endoscopy with biopsy is preferred for confirmation of diagnosis. Contrast-enhanced CT and PET can be used to assess distant metastasis. Endoscopic ultrasonography (EUS) is the investigation of choice to assess tumor invasion and lymph node metastasis.

Other options:

Option B: Diffuse esophageal spasm results in the corkscrew appearance of the esophagus on barium study.

Option C: Esophageal leiomyoma appears as a smooth, semilunar filling defect on barium swallow. it is seen as a discrete ovoid mass that is well outlined by barium.

Option D: Esophageal webs occur in the upper and middle esophagus as a thin membrane that can be identified by the translucent part on contrast radiology.

Solution to Question 2:

This clinical scenario and the image showing the figure of 3 sign point towards coarctation of aorta as the cause of hypertension in this patient.

The figure of 3 sign refers to the indentation of the aorta at the coarctation site. There is pre and post-stenotic dilatation along with left paramediastinal shadow, as shown below.



Other chest X-ray features in coarctation of the aorta are dilated left subclavian artery, dilated ascending aorta and notching of ribs 4-8 due to rib erosion by dilated collateral vessels.

Solution to Question 3:

The given clinical scenario and the radiological images are suggestive of intussusception.

In intussusception, the occluding mass prolapsing into the lumen has been termed the coiled spring appearance (barium in the lumen of the intussusceptum and in the intraluminal space). In adults, it develops due to a tumor of the bowel.

The intussusception appears as a sausage-shaped mass in a longitudinal plane or a target mass in the transverse plane on CT and USG.

Ultrasound signs include:

• Target sign (also known as the doughnut sign)

• Pseudokidney sign- this is seen in the longitudinal section of the bowel

• the edematous bowel resembles renal parenchyma and

• the mesentery which is dragged into the intussusception resembles renal hilum

• Crescent in a doughnut sign.

Note: The colographic appearance of intussusception is also described as a claw sign, pincer defect, shouldering effect, and coiled-spring pattern.

Option B: Volvulus presents with constipation, abdominal bloating, nausea, and/or vomiting. It shows coffee bean sign

Option C: Diverticulitis presents with pain in the left iliac fossa

Option D: Ulcerative colitis causes backwash ileitis

Solution to Question 4:

A cystic lesion in the suprasellar region with calcification is suggestive of craniopharyngioma. It is derived from Rathke's pouch and arises near the pituitary stalk, commonly extending into the suprasellar cistern.

Other options:

Option B: Pituitary adenomas are seen in the sellar region. Microadenomas are best shown on contrast-enhanced images. Macroadenomas are seen as snowman or figure of eight appearance.

Snowman appearance – Pituitary adenoma



Option C: Meningioma is a supratentorial tumor, mainly in the parasagittal region. A classical dural tail may be seen in intracranial meningiomas but is not pathognomic. The MRI image below shows a well-defined, hyper-intense, and homogeneously enhanced mass close to the midline suggestive of meningioma.



Option D: Oligodendroglioma is exclusively found in the cerebral hemispheres, most commonly in the frontal lobes. Upto 90% of oligodendrogliomas contain visible calcification on CT. The CT image below shows a calcified cortical mass that suggests oligodendroglioma.

Solution to Question 5:

Personal monitoring of radiation is done by a TLD badge.

The TLD badge (thermoluminescent dosimeter) should be positioned according to local or national practices. In India, one TLD badge has to be worn at the chest level inside the lead apron if the apron is used. An additional wrist badge has to be worn for procedures requiring hands close to the primary beam.

The thermoluminescent material used is calcium sulfate doped with dysprosium or lithium fluoride. Radiation exposure is monitored every three months. The maximum dose limit for occupational radiation workers is 30 mSv in a year, provided the cumulative dose in a block of 5 years does not exceed 100 mSv.

The given image shows a TLD badge.

Solution to Question 6:

The normal anatomy of the femoral vessels and great saphenous vein at the saphenofemoral junction on USG is called the mickey mouse sign.

The given images below show the mickey mouse sign (FV- femoral vein; FA- femoral artery; GSV- great saphenous vein).



Other options:

Option B: String sign (of Kantor) on barium studies appears as a thin, thread-like stream of contrast passing through the stenosed bowel segment, particularly in Crohn's disease.

Option C: Tillaux sign is said to be positive if the mass/swelling is mobile perpendicular to the mesentery attachment line. Tillaux's sign is seen in a mesenteric cyst.

Option D: Stemmer's sign is the inability to tent the skin at the base of the toes due to skin thickening, as seen in lymphedema.

Solution to Question 7:

Plethoric lung fields are seen in Total Anomalous Pulmonary Venous Connection (TAPVC). In this condition, pulmonary venous obstruction results in pulmonary arterial hypertension, resulting in plethoric lung fields.

	Increasedpulmonary blood flow (pulmonary plethora)	Decreasedpulmonary blood flow
Acyanotic	Ventricular septal defect, Atrial septal defect, Patent ductus arteriosus	—
Cyanotic	TAPVC	Ebstein anomaly, Tetrology of Fallot, Tricuspid atresia

Solution to Question 8:

The clinical manifestations of neck stiffness and severe headache with CT revealing blood in the basal cistern points out to the diagnosis of subarachnoid hemorrhage (SAH).

Subarachnoid hemorrhage (SAH) is the bleeding into the subarachnoid space, which is the area between the arachnoid membrane and the pia mater surrounding the brain. The diagnosis is suspected when a hyperattenuating material is seen filling the subarachnoid space. Most commonly this is apparent around the circle of Willis, on account of the majority of berry aneurysms occurring in this region (65%), or in the Sylvian fissure (30%).

The modified Fisher scale is a method for grading SAH as seen on non-contrast CT (NCCT). The Hunt and Hess scale describes the severity of SAH and is used as a predictor of survival.

Xanthochromia on CSF is indicative of SAH in case of a negative CT scan.

Radiology INI-CET 2021

Question 1:

What is the structure marked in the CT scan ?



a) Anterior ethmoidal canal

b) Posterior ethmoidal canal

c) Optic nerve

d) Cribriform plate

Question 2:

What type of MRI image is given below?



a) Proton density image

b) Susceptibility weighted image

c) Gradient echo image

d) Inversion recovery image

Question 3:

During rounds, your senior was discussing the given image. The following is an image of which of the following investigations?



a) Contrast Dye study

b) CT scan

c) X-ray

d) Angiography

Question 4:

A 22-year-old male was brought to the emergency room following a road traffic accident. On examination, his GCS was E2M2V3. He had no fractures and no hydrocephalus or raised intracranial pressure. Non-contrast CT was performed and was found to be normal. The likely diagnosis is

a) Diffuse axonal injury

b) Post concussion syndrome

c) Extra dural hemorrhage

d) Herniation of brain

Question 5:

A 45-year-old patient presented with vague abdominal pain. On USG, he was found to have a renal cyst of Bosniak class III. CECT was done, as shown below. What is the diagnosis?



a) RCC

b) Renal oncocytoma

c) Renal angiomyolipoma

d) Renal cyst

Question 6:

A young child presented with mild intermittent upper abdominal pain. X-ray is given below. What is the diagnosis?



a) Bochdalek hernia

b) Morgagni hernia

c) Gastric volvulus

d) Eventration of diaphragm

Question 7:

A woman presented with confusion, constipation, and headache. Serum calcium was 13.1mg/dL and ALP was elevated. She was diagnosed with primary hyperparathyroidism. Which of the following is the imaging of choice?

a) MRI

b) FDG PET

c) CT

d) Sestamibi scan

Question 8:

Identify the imaging modality given below.



a) USG

b) Fluroscopy

c) X-Ray

d) MRI

Question 9:

Which of the following is true regarding CORADS?

a) CORADS- 0 indicates normal lung

b) CORADS-5 is typical of COVID-19

c) Pleural thickening is typical of COVID-19

d) Lobar consolidation is typical of COVID-19

Question 10:

Double bleb sign seen in early pregnancy is due to _____.



a) Twin pregnancy

b) Amnion and chorion

c) Yolk sac and amniotic sac

d) Heterotopic pregnancy

Question 11:

Which one of the following is the most radioresistant phase of the cell cycle?

a) S phase

b) M phase

c) G1 phase

d) G2 phase

Answer Key

Question No.	Correct Option
1	d
2	b
3	b
4	a
5	c

6	b
7	d
8	a
9	b
10	c
11	a

Detailed Explanations

Solution to Question 1:

The structure marked in the image is the cribriform plate of the ethmoid bone. The image shown is a coronal section of paranasal sinuses at the level of anterior ethmoidal cells.

The cribriform plate is identified as a bony structure that separates the nasal cavity from the anterior cranial fossa. It is part of the ethmoid bone and is a sieve-like structure as it is pierced by the olfactory nerve fibers.

Crista Galli is also a part of ethmoid bone that projects superiorly from the cribriform plate. It provides attachment for the falx cerebri.

The anterior ethmoidal canal is identified by its location below the roof of ethmoidal air cells. The anterior ethmoidal artery passes through this canal and it is important to identify this to avoid bleeding from this vessel during sinus surgeries.

The posterior ethmoid canal/foramen is an opening in the ethmoid bone through which the posterior ethmoidal artery, vein, and nerve traverse. It is not visualized in this image as it is a coronal section at the level of anterior ethmoidal cells.

The optic nerve can be identified as an oval intraorbital structure in the coronal sections.



Solution to Question 2:

The image shown is a susceptibility-weighted image(SWI) as it clearly demonstrates microhemorrhages as hypointense foci.

SWI is a form of MRI sequence that is especially sensitive to substances that distort the local magnetic field. Compounds with diamagnetic, paramagnetic, and ferromagnetic properties will interact with the local magnetic field and produce distraction. This would lead to a loss of signal in these areas.

Deoxyhemoglobin, hemosiderin, and ferritin are paramagnetic. Bone minerals and dystrophic calcifications are diamagnetic in nature.

The main clinical use is the identification of microhemorrhages/blood products or calcium that are not detected in other MRI sequences.

Other options:

Proton density image:

It is based on the number of nuclei in the area that is imaged. In PD-weighted image tissues with a higher density of protons produce stronger signals and appear brighter. PD weighted image can be identified by comparing fluid with the fat signal. Fluids in these images appear greyish white similar or the appearance of fat in the body.



Gradient echo sequences(GRE):

It uses gradient fields to generate transverse magnetization. In areas of hemorrhage, the iron that is present in the hemoglobin produces its own local magnetic field and dephases the spinning nuclei. This is useful to identify hemorrhagic contusions in the brain and pigmented villonodular synovitis.



Inversion recovery sequences:

This MRI sequence is used to selectively null certain tissues. It can be:

• Short tau inversion recovery(STIR) where fat is nulled. Fluids appear bright and fat appears dark in these images.

• Fluid attenuated inversion recovery(FLAIR) or double inversion recovery(DIR where fluid is nulled.- The fluid(CSF) normally appears dark and lesions appear to be brighter on these images.

FLAIR image-

Solution to Question 3:

The image in the question is a CT scan as it shows the bone as a high-intensity structure (white appearance).

In CT study X-rays are applied in a circular motion and body tissue slices are recreated mathematically. Different tissues are seen with different intensities depending on the proportion of attenuation of the X-rays that pass through them.

• Air and water are displayed as low intensities (dark appearance) because they produce only little attenuation.

• Bone is displayed as a high-intensity structure (bright appearance) because of its significant attenuation.

• High density is seen with hyperemia, freshly clotted blood, and when contrast is used.

• Edema and necrosis are seen as low-intensity lesions.

Other options:

Contrast dye study: In this imaging study, a special dye or contrast is given to evaluate structures that are not identified by conventional imaging. The contrast added would be visible in the image depending on the type of contrast and tissues targeted. Intravenous pyelogram, barium studies, angiography, etc are all examples of contrast studies.

X-ray: It shows different parts of the body in varying shades of black and white. The difference in the appearance of various tissues is due to the differences in the amount of radiation absorbed by them.

• Bone appears white because calcium has high x-ray absorption.

• Fat and soft tissues appear grey due to less x-ray absorption.

• The least absorption is seen with air which appears black.

Angiography: Here a special dye or contrast is injected to highlight the blood vessels are they are not clearly identified in normal imaging. Hence vessels would be highlighted in an angiogram.

Note: A contrast study is different from a contrast-enhanced CT scan, Intravenous pyelogram, barium studies, angiography, etc are all examples of contrast studies.

Solution to Question 4:

The given clinical scenario of an RTA with no imaging abnormality along with a disproportionate clinical manifestation suggests diffuse axonal injury.

It is caused due to stretching of axons in subcortical white matter due to sudden deceleration injuries. It is usually associated with mild nonspecific changes in NCCT. It is visualized on T2 MRI characterized by multiple lesions showing microbleeds or linear hypointensities. It may also show disruption of white matter tracts on tractography.

Other options

Post-concussion syndrome usually follows mild traumatic brain injury. It does not have a severe clinical status as seen in this patient.

Extradural hemorrhage is produced usually results secondary to trauma to head. The bleeding is usually arterial and blood collects between outer dura and inner table of the skull. It can very well be visualized on CT.

A cerebral herniation is defined as a shift of cerebral tissue in an adjacent space. Intracranial brain herniation can be classified into 3 types - subfalcine, transtentorial and tonsillar. A distinct clinical picture can be produced as a result of compression of neurovascular structures by the herniating part. CT is the imaging modality of choice because it can be done rapidly.

Solution to Question 5:

The CT image showing renal lesions that has hypoattenuating components consistent with macroscopic fat is highly suggestive of renal angiomyolipoma.

Imaging in renal angiomyolipoma: Demonstration of macroscopic fat is the main part of making a diagnosis. USG may show hyperechoic lesions in the cortex with posterior acoustic shadowing. In patients with tuberous sclerosis, the entire kidney is affected, producing an echogenic appearance with loss of normal corticomedullary junction.

CT findings:

• Cortical lesions that demonstrate macroscopic fat (less than -20 HU)

• Absence of ossification or calcification

• The presence of macroscopic fat is strongly suggestive of this condition, but not pathognomonic as RCC may rarely demonstrate macroscopic fat.

• Approximately 5% may be fat poor.

MRI: It is an excellent tool to demonstrate intramural fat components. Two sequences are used:

• Fat-saturated techniques

• Phase-in and phase-out imaging that shows India ink artifact at the junction between fat and non-fat components.

Digital subtraction angiography: They are hypervascular lesions showing 'sunburst appearance' in the arterial phase and 'onion peel appearance' in the venous phase.

Angiomyolipomas are benign tumors that occur sporadically or in association with tuberous sclerosis. The mean age of presentation is 43 years and there is a female predilection. It can present as spontaneous retroperitoneal hemorrhage, especially with larger tumors. The patient can present with flank pain, palpable mass, hematuria, urinary tract infections, hypertension, or renal failure. They usually do not require any intervention. mTOR inhibitors like everolimus can be used to decrease the size of the lesion.



Bosniak classification:

It classifies the renal cystic lesions mainly into four classes based on the CT characteristics.



Class I and II are benign and do not require follow-up imaging or surgery.

Class IIF requires follow-up for about 5 years to exclude malignancy.

Class III needs to be biopsied or surgically explored due to a higher risk of malignancy.

Class IV lesions are clearly malignant and need appropriate treatment.

Other options:

Renal cell carcinoma (RCC): It is the most common malignant tumor of the kidney. It usually occurs in the age group 50-70 years and macroscopic hematuria is seen in about 60% of the cases. It is a differential for angiomyolipoma but it is important to note that macroscopic fat is almost always associated with ossification or calcification.



Renal oncocytoma: They are relatively benign in nature and usually present in the 6th to 7th decade. An important point regarding this tumor is its similarity with RCC and hence it is very difficult to differentiate between them preoperatively. A sharp central stellate scar may be seen in one-third of the cases. But RCC may also show this stellate scar appearance in some cases.



Renal cyst: It is a broader term that includes the cystic lesions of the kidney. They can be simple cysts that are benign. Complex or complicated cystic lesions need evaluation and can be malignant. Bosniak classification of renal cysts is given above.

Image showing a simple renal cyst:



Solution to Question 6:

The clinical features and the chest X-ray showing an opacity in the right cardiophrenic angle are suggestive of Morgagni hernia.

Congenital diaphragmatic hernia(CDH) of anteromedial location is the Morgagni hernia. Here the defect is anterior and retrosternal. It is rarer than bochdalek hernia and accounts for only 2-5% of CDH. About 90% of the cases occur on the right side.

It is less symptomatic than bochdalek hernia and the diagnosis is usually delayed until childhood. The patient presents with respiratory or upper GI symptoms. Respiratory symptoms like distress, tachypnea, and recurrent pulmonary infections are common in children. Other symptoms are abdominal pain, dyspnea, flatulence, and indigestion.

Radiographic findings:

• Radiolucent paracardiac shadow may be seen retrosternally on lateral chest x-ray if there is bowel herniation.

• Rounded opacity in the right cardiophrenic angle may be seen if there is herniation of omentum or solid organs.

• A CT scan may be used to confirm the diagnosis.

Surgical repair is indicated in all cases either by a thoracic or abdominal approach.



Other options:

Bochdalek hernia: It is more common than Morgagni hernia. Here the defect is posterolateral in location. Respiratory distress is experienced by most infants. In the pediatric age group, symptoms are predominantly respiratory due to pulmonary hypoplasia. In adults, it is usually asymptomatic and is incidentally found. About 85% of the cases are left-sided.

Gastric volvulus: This is an uncommon condition and presents with the characteristic Borchardt triad which includes:

• Sudden-onset severe upper abdominal pain

• Recurrent retching with the production of little vomitus

• Inability to pass a nasogastric (NG) tube

It can be organoaxial or mesentroaxial. Chest X-ray may show intrathoracic upside-down stomach and retrosternal or mediastinal air-fluid level.

Image showing intrathoracic gastric volvulus with mediastinal gastric bubble-



Eventration of the diaphragm: It refers to the abnormal contour of the dome of the diaphragm without any disruption to the diaphragmatic continuity. It usually involves a segment of the hemidiaphragm. It can cause respiratory distress in newborns and is usually asymptomatic in adults.

Radiographic findings: The affected portion of the hemidiaphragm is elevated and is seen as a smooth hump, while the normal contour is seen in the remaining part. 'Double diaphragmatic contour' may be seen in frontal view and is confirmed with a lateral X-ray.

Image showing eventration of the right dome of the diaphragm-



Solution to Question 7:

Sestamibi scan is the imaging of choice to delineate parathyroid abnormalities.

Sestamibi is a monovalent lipophilic cation. It shows passive diffusion across cell membranes and accumulates in mitochondria. Because of its accumulation in the mitochondria, there is a difference in the rate at which it gets washed out depending upon the number of mitochondria in each tissue.

In parathyroid adenomas, there is a high concentration of oxyphilic cells with high mitochondrial content. The three protocols for sestamibi scanning are:

• Single-isotope dual-phase scan

• Dual-isotope subtraction imaging

• Single-photon emission computed tomography (SPECT).

SPECT is commonly used because of its improved spatial resolution. Sestamibi combined with CT can be used to detect smaller parathyroid tissues.



Other imaging modalities for parathyroid localization:

• Ultrasonography:

• It is inexpensive and noninvasive

• Mediastinal parathyroid tissue cannot be detected because USG cannot penetrate the sternum or clavicles.

• Four-dimensional CT:

• It has a very high sensitivity

• Increased radiation exposure than other routine modalities is a potential disadvantage

• Selective venous sampling for PTH can be used for the exact localization of adenomas

Solution to Question 8:

The image shown in the question is the USG of the neck.

Sonographic examination of the neck shows the thyroid gland, muscles, blood vessels, bony structures or cartilages, and nerves. The thyroid of a healthy patient appears homogeneously echogenic with a uniform echotexture on USG. Lateral to the thyroid, the large blood vessels (the common carotid artery and more laterally, the internal jugular vein) can be identified. In the above image, hypoechoic areas are suggestive of the lumen of vessels seen in the neck i.e common carotid artery and internal jugular vein.

Neck ultrasonography is the preferred modality for routine radiographic evaluation of the thyroid gland, as it offers the most accurate visualization of thyroid nodular disease. It also allows for the concomitant performance of percutaneous fine-needle aspiration biopsy of thyroid lesions.

Neck ultrasound can also be used to localize parathyroid disease.

Other options:

Option B: Fluoroscopy is an imaging modality that allows real-time X-ray viewing of a patient with high resolution. During this procedure, radiologists usually use contrast material (dye) to highlight the area being examined.

Below image shows the fluoroscopy of neck.



Option C: The below image shows the X-ray appearance of neck in AP view.

Option D: Relative to the neck musculature, the thyroid is hyperintense on MRI images.

Solution to Question 9:

Among the given options, CORADS-5 is typical of COVID-19 is the true statement.

COVID-19 Reporting and Data System (CO-RADS) is a categorical assessment for pulmonary involvement of COVID-19 on chest CT.

CO-RADS Category	Summary
CO-RADS 0	Scans are incomplete or of inferior quality
CO-RADS 1	Normal or non-infectious
CO-RADS 2	Typical features of other infections but not of COVID-19
CO-RADS 3	Features compatible with COVID-19 but also with other diseases
CO-RADS 4	Suspicious for COVID-19
CO-RADS 5	Typical for COVID-19
CO-RADS 6	Proven COVID-19 by positive RT-PCR test

CO-RADS categories:

• CO-RADS 0: Not interpretable - Technically insufficient to assign a score: Scans are incomplete or of inferior quality.

• CO-RADS 1: Very low level of suspicion for pulmonary involvement by COVID-19. Mild or severe emphysema, perifissural nodules, lung tumors, or fibrosis.

• CO-RADS 2: Low level of suspicion for pulmonary involvement by COVID-19. Includes infectious etiology that is not compatible with COVID-19 such as bronchitis, bronchiolitis, bronchopneumonia, lobar pneumonia, abscess.

• CO-RADS 3: Equivocal findings for pulmonary involvement by COVID-19. Findings overlap with other viral or infectious etiologies. Could include perihilar ground-glass, homogenous extensive ground-glass with or without sparing of secondary pulmonary lobules.

• CO-RADS 4: High level of suspicion for pulmonary involvement by COVID-19. Findings are similar to CO-RADS 5 but are not present in the typical locations. Findings may also be superimposed on severe pre-existing pulmonary abnormalities.

• CO-RADS 5 based on typical CT findings. It requires the presence of at least one confirmatory pattern (mentioned below).

• CO-RADS 6: Proven COVID-19 by positive RT-PCR test.

CT features typical of pulmonary involvement in COVID-19 include:

• Obligatory features

• Ground glass opacities, with or without consolidation in areas close to the visceral pleural surface, including the fissures.

• Confirmatory features

• Ground glass regions

• Crazy paving pattern

• Organizing pneumonia compatible patterns

• Thickened vessels with parenchymal abnormalities

Solution to Question 10:

The double bleb sign is due to the yolk sac and amniotic sac.

On ultrasonographic examination, a gestational sac may contain the yolk sac and amniotic sac, appearing like two small bubbles. This is known as the double bleb sign. The embryonic disc is located between the yolk sac and the amniotic sac. It is an important feature of an intrauterine pregnancy.

Other options:

In a twin pregnancy, the chorionicity of the placenta is best diagnosed with the help of an ultrasound examination.

• In dichorionic gestation, the twins are separated by amniotic membranes and an intervening chorion. So, a thicker twin peak sign or lambda (λ) sign is formed at the point at which the amnions meet the placenta.

Twin peak/ Lambda sign seen in dichorionic pregnancy



©MARROW

• In monochorionic diamniotic gestation, the twins are separated only by the two amniotic membranes with no intervening chorion. So, a thin T sign is formed at the point at which the amnions meet the placenta.

Heterotopic pregnancy is a condition in which a normal uterine pregnancy co-exists with an extrauterine (ectopic) pregnancy. Although heterotopic pregnancy can occur spontaneously, it is more likely after artificial reproductive technology (ART) and ovulation induction.

The below image shows heterotopic pregnancy with 2 sacs, one inside the uterus (white arrow) and the other in the cervix (black arrows).



Solution to Question 11:

S phase is the most radioresistant phase of the cell cycle.

The replication of the DNA genome occurs only at a specified time period during the cell cycle. This period is referred to as the synthetic or S phase. The S phase is separated from the mitotic, or M phase, by nonsynthetic periods referred to as gap 1 (G1) and gap 2 (G2) phases, occurring before and after the S phase, respectively. The cell prepares for DNA synthesis in G1, and for mitosis in G2.

The most radiosensitive phase of the cell cycle is the G2-M phase. Cells in the G2 or M phase do not have time to repair DNA damage that occurs before they start to divide. This affects mitosis, leading to catastrophe.

Cells become gradually less sensitive to radiation in the G1 phase to the early S phase and are least sensitive in the late S phase.

Radiology NEET 2021

Question 1:

A 15 year old patient with history of spinal tuberculosis was brought to the hospital with low grade fever, dull backache and persistent flexion of hip joint. Hip joint motion was limited and painful. Identify the muscle affected in the CT image.



a) 1

b) 2

c) 3

d) 4

Question 2:

A 12 week old child presented with fever, cyanosis and fatigue. His chest X ray is shown below. What is the most likely diagnosis?

a) Total anomalous pulmonary venous circulation

b) Tetrology of Fallot

c) Transposition of great arteries

d) Pericardial effusion

Question 3:

A patient following a skid resulting in a motor traffic accident was brought to the emergency room 2 hours later. On examination, he was stable with GCS 15/15. The pupil was reactive to light. Tenderness and bruising over the left lower chest wall with petechiae was seen. Severe tenderness was elicited in the left hypochondriac region and BP- 90/50 mm of Hg. What is the best investigation used in the ER?

a) FAST

b) X-ray

c) Diagnostic peritoneal lavage

d) CT scan

Question 4:

A patient presents with abdominal pain and distension. An X ray was taken which showed dilated bowel loops as given in the image below. Identify the segment involved.



a) Jejunum

b) Ileum

c) Duodenum

d) Transverse colon

Question 5:

A child with acute lymphoblastic leukemia undergoes prophylactic irradiation prior to autologous haematopoietic stem cell transplantation. Which of the following will be the least affected?

a) Spermatogonia

b) Intestinal epithelial cells

c) Neurons

d) Bone marrow/ erythroid precursor cells

Answer Key

Question No.	Correct Option
1	b
2	a
3	a

4	a
5	c

Detailed Explanations

Solution to Question 1:

The clinical vignette of a patient with low-grade fever, dull backache, and persistent flexion of the hip joint with a history of spinal tuberculosis is suggestive of psoas abscess involving the psoas muscle, marked 2 in the given image. CT scan shows a mass in the psoas muscle lateral to the vertebral column which is diagnostic of the same.

Psoas abscess occurs due to a collection of pus in the iliopsoas compartment. The compartment is retroperitoneal and hence, the abscess is also called a retroperitoneal abscess. The compartment contains iliacus and iliopsoas muscles. The psoas compartment muscles lie in close proximity to major abdominal organs and is richly supplied by iliolumbar, external iliac and common femoral arteries. Owing to these reasons, there is hematogenous spread of infections like tuberculosis leading to abscess.

Clinical presentation is with back pain, fever, and lassitude. The hip may be in fixed flexion or there may be pain on passive extension.

CT is performed for diagnosis. Treatment is with percutaneous CT-guided drainage and antibiotic therapy.

Other muscles marked in the image are depicted below.



Solution to Question 2:

The given X-ray shows figure of 8 or snowman appearance characteristic of total anomalous pulmonary venous circulation (TAPVC).

TAPVC refers to a condition in which pulmonary veins instead of draining into the left atrium drain into the right atrium directly or indirectly. This results in anomalous drainage of pulmonary venous blood flow into the systemic venous circulation. It is a cyanotic heart disease with features of congestive heart failure. The patients are irritable and have failure to thrive.

The following types are seen:

• Supracardiac (most common): Common pulmonary vein drains into superior vena cava or the left innominate vein

• Cardiac: Pulmonary veins drain into the right atrium or coronary sinus.

• Infracardiac (always obstructive and most dangerous): Pulmonary veins drain into portal vein or inferior vena cava.

Snowman appearance is classically seen in supracardiac variant of TAPVC. The dilated vertical vein on the left, brachiocephalic vein on top, and the superior vena cava on the right form the head of the snowman; the body of the snowman is formed by the enlarged right atrium. It is shown in the image below.



Other options:

Option B: Tetrology of Fallot comprises pulmonary stenosis, ventricular septal defect, overriding of the aorta over the ventricular septum, and right ventricular hypertrophy. On chest X-Ray, the cardiac silhouette appears like a boot ("coeur en sabot").

Option C: In the transposition of great arteries, the aorta arises from the right ventricle and the pulmonary trunk from the left ventricle. The X-ray shows cardiomegaly with cardiac contours which appear like an egg on a string.

Option D: Pericardial effusions occur when excessive fluid gets collected in the pericardial space. The X-ray shows globular enlargement of the cardiac shadow giving a water bottle configuration.

Solution to Question 3:

FAST (Focused Assessment with Sonography for Trauma) is the best investigation that can be performed in the emergency room (ER) for this patient with a blood pressure of 90/50 mm Hg who has sustained blunt abdominal trauma.

Focussed assessment with sonography for trauma is a non-invasive investigation that can be performed rapidly in the emergency room. It helps in identifying an abnormal collection of fluid or blood. FAST is indicated in:

• Haemodynamically unstable patients who had a blunt abdominal trauma

• Patients who sustained penetrating abdominal trauma with no other indications for immediate laparotomy (signs of haemodynamic instability, peritoneal irritation, peritoneal penetration and gunshot wound with transperitoneal trajectory).

FAST includes an examination of four regions:

• Pericardial sac

• Hepatorenal fossa

• Splenorenal fossa, and

• Pouch of Douglas

It is an operator-dependent procedure with limited visualization of the retroperitoneal structures.

Other options:

Option B - A chest X-Ray is used to exclude hemothorax or pneumothorax in a hemodynamically normal patient with penetrating trauma above the umbilicus or a suspected thoracoabdominal injury. Since the given patient is alert, awake, and with no mention of any pelvic pain or tenderness, a pelvic X-Ray is not indicated either.

Option C - Diagnostic peritoneal lavage is an invasive procedure and requires surgical expertise. For these reasons, it is done only in centres where FAST and/ CT are not available.

Option D - CT scan requires transporting the patient and is time-consuming. Hence, it cannot be performed in the ER in a haemodynamically unstable patient.

Solution to Question 4:

The given clinical scenario and X-ray suggest intestinal obstruction where the involved segment is the jejunum. The obstructed bowel loops are central in location and show valvulae connivente (mucosal folds that cross the entire diameter of the bowel) which is characteristic of jejunal involvement. The gas bubbles trapped between the valvulae connivente give a string of beads appearance.

Radiological features of bowel obstruction

• Small bowel - The involved dilated segments are straight and usually centrally located.

• Jejunum - It is characterized by the presence of valvulae connivente with gas bubbles trapped between them. This gives a string of beads/ ladder effect.

• Ileum - The distal ileal obstruction is usually described as featureless.

• Caecum - A rounded gas shadow in the right iliac fossa is suggestive of caecal obstruction.

• Large bowel - The presence of haustrations is a feature of large bowel obstruction. Haustral folds are irregular and do not cross the entire diameter of the bowel. These are absent in the caecum.

In intestinal obstruction, fluid levels appear as the obstruction progresses. When the fluid levels are prominent, it indicates that the obstruction is advanced. The number of fluid levels also increases if the lesion is more distal.

The most reliable criterion for the diagnosis of small bowel obstruction is the transition zone, seen on CT imaging. It identifies a definite point of obstruction. Dilated small bowel loops are seen proximal to the site of obstruction and collapsed loops are seen distal to the obstruction.

Valvulae connivente in jejenal obstruction

Large bowel obstruction -
X-Ray appearance



Solution to Question 5:

Neurons are least susceptible to radiation. The radiosensitivity of biological tissue is directly proportional to its mitotic activity and inversely proportional to its degree of differentiation. This is known as the law of Bergonie and Tribondeau.

Hence, hematopoietic cells are the most radiosensitive, and the central nervous system is the least sensitive to radiation.

Acute Radiation Syndromes (ARS) is an acute illness that is caused by exposure of the entire body to a high dose of penetrating radiation in a very short period of time. It is also known as radiation toxicity or radiation sickness. It consists of three syndromes:

Stages of ARS:

• Prodromal stage - This stage starts within minutes of exposure and may continue for a few days. Symptoms include nausea, vomiting, and anorexia.

• Latent stage - It begins within a few hours and continues up to a few weeks. The patient generally feels healthy.

• Manifest illness stage - This stage may continue up to months and symptoms vary according to the syndrome.

• Recovery/ death - The recovery process occurs over two years. Recovery cannot be expected in cardiovascular and central nervous system syndromes, which lead to death.

Syndrome	Radiation dose
Hematopoietic (Bone marrow)	>0.7 Gy
Gastrointestinal	>10 Gy
Cardiovascular/ Central nervous system	>50 Gy

Radiology INI-CET 2022

Question 1:

A patient presents to the casualty following blunt trauma to the chest. A chest X-ray was done. Among the following radiographs, in which case would you further evaluate the patient before putting a chest tube?



a) 4

b) 1

c) 2

d) 3

Question 2:

A 30-year-old patient presents with progressive difficulty in swallowing both solids and liquids for 8 months. A barium swallow was done and is shown in the image below. What other investigations should be done to confirm the diagnosis in this patient?



a) UGIE + manometry

b) UGIE + contrast enhanced CT

c) UGIE + PET

d) UGIE + 24 hr pH monitoring

Question 3:

A 28-year-old woman with a severe headache is scheduled to undergo a CT scan. During history taking, which of the following will you definitely ask for before the commencement of the scan?

a) Use of cochlear implant

b) LMP

c) History of fall

d) History of hypertension

Question 4:

Which of the following findings are seen in a high-resolution CT scan of fungal pneumonia?

a) 1, 2, 4

b) 1, 3, 4

c) 1, 2, 3

d) 2, 3, 4

Question 5:

An elderly lady presents with abdominal pain and distension. She has a history of scoliosis. Given below is the erect X-ray abdomen radiograph of the lady. What is the most likely diagnosis?



a) Large bowel obstruction due to sigmoid volvulus

b) Large bowel obstruction due to fecolith

c) Cecal volvulus

d) Small bowel obstruction due to bowel perforation

Question 6:

A child presents with cyanosis. His chest radiograph is shown below. What is the diagnosis?

a) Tetralogy of Fallot (TOF)

b) Ebstein's anomaly

c) Transposition of great arteries (TGA)

d) Total anomalous pulmonary venous return (TAPVC)

Question 7:

An HIV patient presented with a headache and altered sensorium. Consider the following statements and identify the essential criteria for the diagnosis of neurocysticercosis.



a) 1 and 2 only

b) 1 and 4 only

c) 1, 3, and 4

d) 1, 2, and 4

Question 8:

Identify the condition based on the non-contrast CT scan of a patient given below.



a) Hepatocellular carcinoma

b) Fibronodular hyperplasia

c) Liver abscess

d) Hydatid cyst

Question 9:

Match the following:

a) 1-b, 2-d, 3-a, 4-c

b) 1-c, 2-a, 3-b, 4-d

c) 1-d, 2-b, 3-a, 4-c

d) 1-d, 2-a, 3-b, 4-c

Question 10:

Whole-body CT scan protocol for a trauma patient includes all, except _____.

a) CT Head

b) CT Cervical spine

c) CT Limbs

d) CT Abdomen

Question 11:

A patient presents with hematuria and a mass in the right kidney on a CT scan, as shown below. What is the diagnosis?



a) Renal cell carcinoma

b) Renal oncocytoma

c) Renal angiomyolipoma

d) Renal cyst

Question 12:

Egg shell calcification is seen in all the following, except _____.

a) Silicosis

b) Sarcoidosis

c) Post-radiation therapy lymph nodes

d) Metastatic lymph nodes

Question 13:

A 55-year-old man presents with hematuria and a mass in the left kidney on a CECT scan, as shown below. What is the diagnosis?



a) Renal angiomyolipoma

b) Renal oncocytoma

c) Renal cell carcinoma

d) Renal cyst

Question 14:

A patient who was diagnosed with prostate cancer is being investigated. The bone scan is reported as a super scan with increased uptake in the bones and reduced activity in the kidney and soft tissue. What is the reason for this super scan appearance?

a) Increased uptake due to diffuse metastasis

b) Increased uptake by carcinoma prostate

c) Increased uptake by the bone

d) Decreased uptake by adrenal glands and kidney

Answer Key

Question No.	Correct Option
1	b
2	a
3	b
4	c
5	a
6	a
7	b
8	d
9	d
10	c
11	c
12	d
13	c
14	a

Detailed Explanations

Solution to Question 1:

Among the following radiographs, image 1 shows dilated bowel loops in the hemithorax, and is suggestive of diaphragmatic hernia and it needs to be evaluated further before putting a chest tube since it can cause injury to the intestinal contents in the thorax.

Blunt injury to the diaphragm can lead to rupture and therefore the abdominal contents can herniate into the chest. Nasogastric tube placement followed by chest radiography can be useful in case of herniation of the stomach into the chest. The most accurate investigation is video-assisted thoracoscopy or laparoscopy.

Operative repair should be done. Repair should be done via the abdomen in case of a diaphragmatic rupture.

Radiographs 2,3,4 are hemothorax, pneumothorax, and flail chest respectively. These conditions are life-threatening. Hence, the diagnosis and management have to be immediate. Rapid insertion of intercostal chest drains should be done. They are inserted into the fifth intercostal space. The safe landmark is identified as a triangle that lies anterior to the midaxillary line above the nipple, below, and lateral to the pectoralis major muscle.

The below images show the triangle of safety and technique of chest drain insertion.





Hemothorax can occur following blunt trauma. It occurs due to bleeding from the intercostal vessels or internal mammary artery secondary to rib fracture. Features include hemorrhagic shock, flat neck veins, unilateral absence of breath sounds, and dullness on percussion. Correction of hypovolemic shock, intercostal drain insertion, and intubation are useful in management.

Pneumothorax can be open or tension pneumothorax. In tension pneumothorax, the common etiology is penetrating or blunt trauma, and features such as tachypnea, dyspnea, distended neck veins, tracheal deviation, hyper resonance and decreased breath sounds are seen. Immediate decompression can be attained by large bore cannula insertion into the second intercostal space on the affected side in the midclavicular line after which a chest tube can be placed in the 5th intercostal space along the anterior axillary line. Chest radiograph shows mediastinal shift to the opposite side and diaphragmatic depression on the same side.

Flail chest occurs as a result of blunt trauma and is associated with rib fractures. Clinically, paradoxical movement of the chest can be present. Management includes oxygen administration, analgesia, chest drain, and surgery to stabilize the flail segment. Rib fractures can be seen on

chest radiograph.

Solution to Question 2:

The given clinical scenario and the barium swallow showing bird beak sign are suggestive of achalasia cardia. Upper gastrointestinal endoscopy (UGIE) and manometry should be done to confirm the diagnosis.

Achalasia is a motility disorder and is less common. It typically presents as dysphagia for solids and liquids. Other features such as regurgitation and chest pain are also present.

On fluoroscopy, a grossly dilated esophagus with tapering end towards the gastroesophageal junction is seen. This is called the bird beak sign.

Barium is retained in the esophagus for more than 5 minutes and is a significant finding.

Diagnosis is confirmed with endoscopy and manometry.

Solution to Question 3:

The most important information that should be ascertained during history taking in a 28-year-old woman of reproductive age group posted for a CT scan is the last menstrual period (LMP). CT scan is avoided during pregnancy due to the potential adverse effects of radiation on the fetus; hence, LMP is ascertained to rule out pregnancy.

CT is a high-resolution technique using x-rays to generate cross-sectional images of any area of the body. It is associated with high radiation doses.

Other options:

Option A: Use of cochlear implant is a contraindication for MRI and not CT. CT scan can be performed on a patient with a cochlear implant.

Option B: History of fall in a patient with a headache is asked for suspected head trauma. A CT scan in this patient can help in the diagnosis and assess extent of damage.

Option C: History of hypertension points to a possible intracranial bleed. A CT scan can be useful in ruling out the bleed.

Since it is imperative to rule out pregnancy before taking a CT scan, last menstrual period is the most appropriate answer to this question.

Solution to Question 4:

High-resolution computed tomography (HRCT) of fungal pneumonia shows interlobular septations, peripheral wedge-shaped consolidation, and pleural effusion. HRCT of fungal pneumonia shows a nodule (not a cavity) with surrounding ground-glass opacities.

A nodule surrounded by ground-glass attenuation on HRCT imaging is seen in aspergillosis, candidiasis, Pneumocystis jirovecii pneumonia, coccidioidomycosis, and paracoccidioidomycosis. Cavitation on HRCT is seen in cases of community-acquired pneumonia due to Mycobacterium tuberculosis, S. aureus, and gram-negative bacilli.

Aspergillosis is a fungal disease caused by Aspergillus sp. Pulmonary aspergillosis has been categorized into saprophytic, allergic, and invasive forms.

HRCT findings of various types of aspergillosis are as follows:

• Aspergilloma (saprophytic) shows a mass with soft-tissue attenuation, which is separated from the thoracic wall by airspace called the air crescent sign. Multiple nodules and bilateral involvement may be seen.

• Allergic bronchopulmonary aspergillosis (ABPA-allergic) shows hypodense mucous plugs.

• Angioinvasive aspergillosis shows multiple nodules surrounded by a halo of ground-glass attenuation ('halo sign') or pleural-based wedge-shaped areas of consolidation.

The following is an HRCT image of aspergilloma showing the air crescent sign:



The following CT image of aspergilloma shows the halo sign:



Candidiasis pneumonia caused by Candida sp. in immunocompromised patients shows multiple bilateral nodular opacities, areas of consolidation, and ground-glass opacities on HRCT. Pleural effusion, thickening of the bronchial walls, and cavitation are less commonly seen.

Paracoccidioidomycosis (PCM), is a disease caused by Paracoccidioides brasiliensis. On HRCT, ground-glass opacities, nodules, the 'halo' and 'reversed halo' signs, interlobular septal thickening, airspace consolidation, cavitation, and fibrosis.

Solution to Question 5:

The given clinical scenario with an old woman and the radiograph showing characteristic coffee bean appearance are highly suggestive of sigmoid volvulus causing a large bowel obstruction.

Sigmoid volvulus is a cause of large bowel obstruction and occurs when the sigmoid colon twists on the sigmoid mesocolon. Symptoms include constipation, abdominal bloating, nausea, and/or vomiting. The onset may be acute or chronic. It occurs more commonly in patients with a history of mental illness and chronic constipation.

X-ray abdomen shows a large, dilated loop of the colon, often with a few air-fluid levels. Specific signs include the coffee bean sign, Frimann-Dahl sign (three dense lines converge towards the site of obstruction), and the absence of rectal gas. CT may reveal a large gas-filled loop without haustral markings, forming a closed-loop obstruction.

Here are some pointers that help in differentiating the two volulus:

Other options:

Option B: In large bowel obstruction due to fecolith, the presence of haustrations is a feature. Haustral folds are irregular and do not cross the entire diameter of the bowel. These are absent in the cecum. The given image shows haustrations in large bowel obstruction.

The following image shows an x-ray radiograph of large bowel obstruction:

Sigmoid volvulus	Cecal volvulus
More common incidence	Less common incidence
Coffee bean sign is typical.	Coffee bean sign is occassion ally seen
Seen in old age, psychiatric p atients	Usually seen in those aged 3 0-60 years old
The top of the sigmoid volvul us usually lies very high in th e abdomen with its apex on t he left side.	In half the patients, the caec um twists and inverts so that the pole of the caecum and th e appendix occupy the left up per quadrant.In other patient s, the twist occurs in an axial plane without inversion and t he caecum still occupies the r ight half of the abdomen.
Haustra aren't usually seen	Haustra usually seen
Multiple air-fluid level seen	One air-fluid level seen
	Attached air-filled appendix c an be seen



Option C: Cecal volvulus shows characteristic radiological findings like cecal dilatation, single air-fluid level that may extend into the left upper quadrant, and absence of gas in the distal colon. The colonic haustrations are well maintained.

The following image shows an x-ray radiograph of cecal volvulus:



Option D: In small bowel obstructions, the obstructed bowel loops are central in location and lie transversely. The jejunal obstruction shows valvulae conniventes (mucosal folds that cross the entire diameter of the bowel) giving it a ladder effect. The distended cecum may show the presence of gas shadow in the right iliac fossa.

Common causes of small bowel obstruction are adhesions (~60%), neoplasms (~20%), hernias (~10%), Crohn's disease (~5%) and other miscellaneous conditions (~5%).

The following image shows an x-ray radiograph of small bowel obstruction:



Cases of bowel perforation have an erect abdominal radiograph showing free gas under the diaphragm. A supine radiograph may show dilated gas-filled loops of the bowel, suggestive of paralytic ileus.

The below radiograph image shows a hollow viscus perforation:



Solution to Question 6:

The given clinical scenario and chest X-ray showing a "coeur en sabot" or boot-shaped heart are suggestive of tetralogy of Fallot (TOF).

TOF is a cyanotic congenital heart disease with the following four components:

• Obstruction to right ventricular outflow (pulmonary stenosis)

• Malalignment type of ventricular septal defect (VSD)

• Dextroposition of the aorta so that it overrides the ventricular septum (overriding of the aorta), and

• Right ventricular hypertrophy (RVH)

In TOF, the hypertrophied right ventricle causes the rounded apical shadow to be uptilted so that it is situated higher above the diaphragm than normal. On x-ray, this appears as a boot-shaped heart or coeur en sabot, as shown below:

Boot-shaped heart



©MARROW

Option B: Ebstein anomaly is a rare cyanotic congenital heart disease with diminished pulmonary blood flow due to abnormality of the tricuspid valve.

In the Ebstein anomaly, there is severe right-sided cardiomegaly due to an elongated and enlarged right atrium, which leads to an elevated apex giving a box-shaped heart, as shown below:

Box-shaped heart seen in Ebstein anomaly



©MARROW

Option C: Transposition of great arteries (TGA) can present from birth to infancy with cyanosis, decreased exercise tolerance, and poor physical development TGA is the most specific heart disease in infants of diabetic mothers.

The following is a radiograph of TGA showing the egg-on-a-string appearance/egg-on-side appearance of the heart:



Option D: Total anomalous pulmonary venous return (TAPVC) occurs due to the anomalous connection of pulmonary veins instead of joining to the left atrium resulting in pulmonary venous blood reaching the right atrium.

In the supracardiac type of TAPVC, the configuration of the heart and the superior mediastinal borders resemble a snowman. It is also referred to as the figure of 8 sign, as shown below:



Solution to Question 7:

The MRI brain labeled as B shows multiple cysticerci within the brain suggesting neurocysticercosis. Among the given statements, the essential criteria for the diagnosis of neurocysticercosis include 1 and 4 (option B), i.e.,

- Brain biopsy showing the parasite.

• Neuroradiologic demonstration of cystic lesions with scolex (MRI labeled B in this case).

Neurocysticercosis is a condition caused by the larvae of Taenia solium.

Criteria for diagnosis of neurocysticercosis:

1. Absolute criteria

• Demonstration of cysticerci by histologic or microscopic examination of a biopsy taken from the brain or spinal cord lesion.

• Visualization of the parasite in the eye by funduscopy

• Neuroradiologic demonstration of cystic lesions containing a characteristic scolex

2. Neuroimaging criteria

• Major neuroimaging criteria: Cystic lesions without a discernible scolex, multilobulated cystic lesions in the subarachnoid space, typical small enhancing lesions, typical parenchymal brain calcifications

• Confirmative neuroimaging criteria: Resolution of cystic lesions spontaneously or after cysticidal drug therapy and migration of ventricular cysts documented on sequential neuroimaging studies

• Minor neuroimaging criteria: Obstructive hydrocephalus or abnormal enhancement of basal leptomeninges

3. Clinical/exposure criteria

• Major clinical/exposure criteria: Demonstration of antibodies to cysticerci or cysticerci antigen in cerebrospinal fluid by enzyme-linked immunosorbent assay; Household contact with an individual infected with T. solium; Evidence of cysticercosis outside the central nervous system (e.g., cigar-shaped soft-tissue calcifications)

• Minor clinical/exposure criteria: Clinical manifestations suggestive of neurocysticercosis; Residence in a cysticercosis-endemic area; Frequent travel to a cysticercosis-endemic area

Diagnosis is confirmed by:

• One absolute criterion

• Two major neuroimaging criteria plus any clinical/exposure criteria

• One major and one confirmative neuroimaging criterion plus any clinical/exposure criteria

• One major neuroimaging criteria plus two clinical/exposure criteria (including at least one major clinical/exposure criterion), with the exclusion of other pathologies producing similar neuroimaging findings

A probable diagnosis is supported by the fulfillment of

• One major neuroimaging criterion plus any two clinical/exposure criteria

• One minor neuroimaging criterion plus at least one major clinical/exposure criterion

Other options: The MRI labeled as A shows ventriculomegaly in brain atrophy.

Solution to Question 8:

The given CT image shows a cyst in the liver with the serpent sign and areas of calcification, which are suggestive of hydatid disease.

Hydatid disease or echinococcosis is a zoonotic infection caused by the Echinococcus species of flatworms. Its life cycle consists of dogs as definitive hosts and humans as accidental intermediate hosts.

Primary hydatid cysts most commonly occur in the liver. Most patients are asymptomatic but when symptomatic, the patient presents with hepatomegaly, pain, and obstructive jaundice.

The first/initial imaging test is USG, and CT scan can be used to diagnose the condition.
CT scan shows the presence of a space-occupying lesion with a smooth outline and septa. CT scan has an accuracy of 98% in the detection of hydatid cysts and high sensitivity to the detection of daughter cysts and extra-hepatic cysts.

On CT imaging, serpent sign is seen as wavy membranes within the cyst:

The USG of an active hydatid cyst shows multiple septations and a rosette or honeycomb appearance:



The ball of wool sign is seen on USG when the contents settle in the dependent portion of the cyst, as seen below:



On USG, the water-lily sign is seen in hydatid infections when there is detachment of the endocyst membrane, which results in floating membranes within the pericyst that mimic the appearance of a water lily.



The image below shows the structures present within a cyst.



Other options:

Option A: Hepatocellular carcinoma (HCC) is diagnosed with contrast-enhanced CT (CECT) or MRI. On a non-contrast CT, it appears as a low-attenuated lesion with occasional calcification. On CECT, HCC lesions show enhancement in the arterial phase and washout in the portal phase.

The following image shows the triphasic CECT of HCC:



Option B: Focal nodular hyperplasia is a benign solid lesion in the liver. Histologically, it is composed of normal liver elements, but the biliary elements are not connected to the normal biliary tree. On CECT, they exhibit homogeneous enhancement in the arterial phase of IV contrast.

The image below shows late arterial and portal venous phase CECT of focal nodular hyperplasia:



Option C: Liver abscess may be caused by various infections. It begins as a local inflammatory reaction and later develops liquefactive necrosis in the center, with an inflammatory margin.

On a non-contrast CT, it appears as an ill-defined lesion with low attenuation, as seen in the image below:



The given image shows a liver abscess with rim enhancement on CECT.

Solution to Question 9:

The correct match is 1-d, 2-a, 3-b, 4-c (option D).

Football sign is seen in cases of massive pneumoperitoneum in abdominal x-ray film taken in the supine position.

The following image shows football sign in a patient with pneumoperitoneum:

1. Football sign	d. Pneumoperitoneum
2. String sign	a. Hypertrophic pyloric steno sis

1. Football sign	d. Pneumoperitoneum
3. Rat tail sign	b. Achalasia cardia
4. Inverted umbrella sign	c. Ileocecal TB

Rat tail sign refers to dilatation of the esophagus with narrowing at the gastro-esophageal junction in barium swallow/ fluoroscopy studies. It is also known as bird beak sign and is typically seen in achalasia cardia. In the case of a rat tail sign with irregular contour, esophageal cancer should be suspected.

The following image shows rat tail sign or bird beak sign in a patient with achalasia cardia:



String sign refers to the string-like appearance of severely narrowed bowel lumen in fluoroscopy/barium studies. It is seen in hypertrophic pyloric stenosis, gastrointestinal tuberculosis, colon cancer, and carcinoid tumor. String sign of Kantor is classically described in Crohn's disease.



String Sign in Pyloric Stenosis.

Inverted umbrella sign, also known as Fleischner sign, is seen in barium studies of ileocecal tuberculosis. It refers to a widely gaping, thickened, patulous ileocecal valve and a narrowed, ulcerated terminal ileum.

Fleischner sign/Inverted umbrella sign



Solution to Question 10:

The whole-body CT of a trauma patient does not include CT limbs.

Whole-body CT or trauma CT or panscan is increasingly being used in the context of trauma patients with a dangerous mechanism of injury:

• Non-contrast CT of head

• Non-contrast CT of neck

• Contrast-enhanced CT of chest (arterial phase)

• Contrast-enhanced CT of abdomen (portal venous phase)

The purpose of the scan is to rapidly evaluate life-threatening injuries and accurately diagnose known and unknown injuries.

Solution to Question 11:

The CT image showing renal lesions that has hypo-attenuating components consistent with macroscopic fat is highly suggestive of renal angiomyolipoma.

Angiomyolipomas are benign tumors that occur sporadically or in association with tuberous sclerosis. The mean age of presentation is 43 years and there is a female predilection. It can present as spontaneous retroperitoneal hemorrhage, especially with larger tumors. The patient can present with flank pain, palpable mass, hematuria, urinary tract infections, hypertension, or renal failure.

Imaging in renal angiomyolipoma: Demonstration of macroscopic fat is the main part of making a diagnosis. USG may show hyperechoic lesions in the cortex with posterior acoustic shadowing. In patients with tuberous sclerosis, the entire kidney is affected and produces an echogenic appearance with loss of normal corticomedullary junction.

CT findings:

• Cortical lesions that demonstrate macroscopic fat (less than -20 HU)

• Absence of ossification or calcification

• The presence of macroscopic fat is strongly suggestive of this condition, but not pathognomonic as RCC may rarely demonstrate macroscopic fat.

• Approximately 5% may be fat-poor.

MRI: It is an excellent tool to demonstrate intramural fat components. Two sequences are used:

• Fat-saturated techniques

• Phase-in and phase-out imaging that shows India ink artifacts at the junction between fat and non-fat components.

Digital subtraction angiography: They are hypervascular lesions showing a 'sunburst appearance' in the arterial phase and an 'onion peel appearance' in the venous phase.

They usually do not require any intervention. mTOR inhibitors like everolimus can be used to decrease the size of the lesion.



Other options:

Renal cell carcinoma (RCC): It is the most common malignant tumor of the kidney. It usually occurs in the age group 50-70 years and macroscopic hematuria is seen in about 60% of the cases. It is a differential for angiomyolipoma but it is important to note that macroscopic fat is almost always associated with ossification or calcification.



Renal oncocytoma: They are relatively benign in nature and usually present in the 6th to 7th decade. An important point regarding this tumor is its similarity with RCC and hence it is very difficult to differentiate between them preoperatively. A sharp central stellate scar may be seen in one-third of the cases. But RCC may also show this stellate scar appearance in some cases.



Renal cyst: It is a broader term that includes the cystic lesions of the kidney. They can be simple cysts that are benign. Complex or complicated cystic lesions needs evaluation and can be malignant. Bosniak classification of renal cysts is given above.

Image showing a simple renal cyst:



Solution to Question 12:

Eggshell calcification is not seen in the case of metastatic lymph nodes.

A ring of calcification at the periphery of a lymph node is known as eggshell calcification.

Solution to Question 13:

The history of an elderly male presenting with painless hematuria, renal mass, and an axial renal cortical phase CECT image showing renal lesions with irregular enhancement is suggestive of renal cell carcinoma (RCC).

Renal cell carcinoma (RCC) is the most common malignant tumor of the kidney. It usually occurs in the age group 50–70 years and macroscopic hematuria is seen in about 60% of the cases. It usually presents with painless gross hematuria, flank pain, or painless flank mass.

Contrast-enhanced CT is the investigation of choice for RCC. On CT, small lesions enhance homogeneously, whereas larger lesions have irregular enhancement due to areas of necrosis. The clear cell subtype may show much stronger enhancement is a differential for angiomyolipoma. Still, it is essential to note that macroscopic fat is almost always associated with ossification or calcification.

T1-weighted MRI is often heterogeneous due to necrosis, hemorrhage, and solid components. T2-weighted MRI is hyperdense for clear cell RCC and hypodense for papillary RCC. Thus, a T2-weighted MRI can be used for histological diagnosis. Contrast-enhanced MRI is the investigation of choice for detecting vascular invasion of RCC.



Other options:

Renal angiomyolipoma: Angiomyolipomas are benign tumors that occur sporadically or in association with tuberous sclerosis. The mean age of presentation is 43 years, and there is a female predilection. CT findings show cortical lesions that demonstrate macroscopic fat (less than -20 HU) and absence of ossification or calcification.



Renal oncocytoma: They are relatively benign in nature and usually present in the 6th to 7th decade. An important point regarding this tumor is its similarity with RCC, and hence, it is very difficult to differentiate between them preoperatively. A sharp central stellate scar may be seen in one-third of the cases. But RCC may also show this stellate scar appearance in some cases.



Renal cyst: It is a broader term that includes the cystic lesions of the kidney. They can be simple cysts that are benign. Complex or complicated cystic lesions needs evaluation and can be malignant. Bosniak classification of renal cysts is given above.

Image showing a simple renal cyst:



Solution to Question 14:

The super scan appearance is due to increased uptake due to diffuse metastasis.

Diffuse osteoblastic metastatic diseases such as carcinoma breast and prostate carcinoma lead to a generalized increase in the skeletal uptake relative to soft tissue and absent or faint renal activity on a bone scan. This is known as a super scan. It may also be seen in generalized bone diseases such as hyperparathyroidism.

Carcinoma of the prostate is the most common malignant tumor in men over the age of 65 years. Most cancers are clinically silent and are detected on routine rectal examination or PSA measurements. Grossly, malignant tissue is gritty and firm in consistency (in contrast to the spongy appearance of benign tissue) and thus more readily palpable on digital rectal examination. They usually arise in the peripheral zone of the gland.

Local extension most commonly involves the periprostatic tissue, seminal vesicles, and the base of the urinary bladder. Metastasis spreads via the lymphatics to the obturator nodes and the para-aortic nodes. Hematogenous spread occurs chiefly through the bones to the axial skeleton and rarely to the viscera. The bony metastasis is blastic in origin and involves the lumbar spine, proximal femur, pelvis, thoracic spine, and ribs in decreasing order of frequency.

Prognosis and staging: The Gleason grading system correlates with the pathological stage, spread, and prognosis. Measurement of serum PSA levels may be useful to assist with the diagnosis and management.

Radiology NEET 2022

Question 1:

Identify the imaging modality given below.



a) Cerebral blood flow scan

b) PET CT

c) Sestamibi scan

d) Full body MDCT

Question 2:

A patient is brought to the casualty following a road traffic accident. On examination, the patient is conscious, BP is 90/60 mm Hg, respiratory rate is 40 breaths per minute, and pulse rate is 120 bpm. The X-ray is shown below. What is the next best step in management?



a) Chest tube insertion

b) Pericardiocentesis

c) Thoracotomy

d) Pleurodesis

Answer Key

Question No.	Correct Option
1	b
2	a

Detailed Explanations

Solution to Question 1:

The above imaging modality is a PET CT scan.

PET (positron-emission tomography) CT scan is a hybrid technique that combines PET and CT scan that is used for targeted imaging of tumor physiology and biology. It is based on the principle that tumor cells generally have high metabolic activity compared with normal tissues.

A small dose of the radiopharmaceutical agent 18F-fluorodeoxyglucose (FDG). This enters cells and is phosphorylated to FDG 6-phosphate which cannot be metabolized further. Since it is a highly polar molecule, it cannot easily diffuse back out of the cell. After intravenous injection of

FDG, it continuously accumulates in the metabolically active cell.

The main advantage of PET-CT is the identification of distant metastatic disease not seen on CT alone. The disadvantages of PET scans are radiation burden, poorer spatial resolution when used alone, and high cost.

Other options:

Option A: Cerebral blood flow scan refers to the perfusion MRI that uses gadolinium as contrast and provides information about the cerebral blood flow and volume.

Option B: The sestamibi scan is used for parathyroid localization and is now considered the most accurate and reliable method for imaging the parathyroid glands. Sestamibi accumulates in mitochondria. Parathyroid adenomas often have a high concentration of oxyphilic cells with high mitochondrial content. These retain sestamibi tracer and adenomas are therefore associated with a slow washout when compared with the thyroid gland.

Option D: The full body MDCT scan refers to the multidetector CT scan which is an advanced CT imaging technique that uses multiple detector elements that provide improvement in speed, quality, and scope.

Solution to Question 2:

The given X-ray features showing a well-demarcated area devoid of lung markings on the left side with tracheal deviation and the mediastinal shift to the right side, along with the clinical scenario is suggestive of tension pneumothorax. The next best step in the management is chest tube insertion.

Tension pneumothorax is a clinical diagnosis, and treatment should never be delayed while waiting for radiological confirmation. Imaging modalities used are ultrasound–extended focused assessment with sonar for trauma (eFAST), chest x-ray, and spiral CT.

A tension pneumothorax develops when there is a tear either in the parietal or visceral pleura. This tear acts as a one-way valve i.e., air enters from the lung or chest wall through it during inspiration but has no way to escape during expiration. This ultimately leads to the collapse of the affected lung.

It usually occurs following penetrating or blunt chest trauma, iatrogenic lung injury, or mechanical positive pressure ventilation.

It is characterized by:

• Chest pain

• Tachypnea and acute dyspnea

• Sudden onset hemodynamic compromise (tachycardia, hypotension, low O_2 saturation, loss of consciousness)

• Neck vein distention

• Tracheal deviation away from the side of the injury

• Decreased chest movements on the affected side

• Hyper-resonance on percussion on the affected side

• Absence or decreased breath sounds on the affected side

The X-ray below shows a left-sided tension pneumothorax. There is a large, well-demarcated area devoid of lung markings on the left side. The yellow arrows are pointing to the tracheal deviation and the mediastinal shift to the right side.



Management of tension pneumothorax:

• Immediate management- needle decompression or needle thoracostomy i.e. insertion of a wide-bore needle in the 5th ICS anterior to the midaxillary line

• Definitive management- tube thoracostomy or chest tube insertion in the 5th ICS in the triangle of safety, identified as a triangle that lies anterior to the midaxillary line above the nipple, below, and lateral to the pectoralis major muscle.

The images below show the triangle of safety and technique of chest drain insertion.



Other options:

• Pericardiocentesis is done in cases of cardiac tamponade which is characterized by the presence of hypotension, distended jugular veins, and muffled heart sounds (Beck's triad). The breath sounds are absent in tension pneumothorax while they are present in cardiac tamponade.

• Surgery for pneumothorax can be performed by video-assisted thoracoscopic surgery (VATS) or as an open procedure (thoracotomy) in cases of spontaneous pneumothorax failing to settle despite chest drainage. Pleural adhesion is achieved by chemical pleurodesis where talc is used and is insufflated into the chest cavity.

Radiology NEET 2023

Question 1:

A 2-month-old infant is brought to the OPD with a parietal swelling present since birth. The X-ray image is given below. What could be the probable diagnosis?



a) Subgaleal hematoma

b) Cephalhematoma

c) Caput succedaneum

d) Enecephalocele

Question 2:

A patient presented with abdominal pain and sterile pyuria, and the x-ray showed the following features. What is the likely diagnosis?

a) Putty kidney

b) Nephrocalcinosis

c) Staghorn calculus

d) Psoas calcification

Question 3:

A woman with endometrial carcinoma is undergoing radiotherapy. Which of the following is true?

a) Intensity is inversely proportional to the square of the distance from the source

b) Small blood vessels are most radioresistant

c) Rapidly proliferating cells are most radioresistant

d) Small intestinal mucosa is most radio-resistant

Question 4:

A patient presents with a history of fever and cough for the past 3-5 days. His examination findings reveal crepitations. Chest X-ray is given below. What is the probable diagnosis?

a) Apical segment of Right lower lobe consolidation

b) Right middle lobe consolidation

c) Posterior segment of right lower lobe consolidation

d) Loculated pleural effusion

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b

Detailed Explanations

Solution to Question 1:

In the given clinical picture, the swelling over the infant's head is likely to be a cephalhematoma.

Cephalohematoma is a collection of blood in between the pericranium and the flat bone (subperiosteal) of the skull, usually unilateral and over a parietal bone. It is never present at birth but gradually develops after 12–24 h. The swelling is limited by the suture lines of the skull as the pericranium is fixed to the margins of the bone.

The arrow mark in the image represents cephalohematoma.

Note: Cephalhematoma is more likely to be identified soon after birth, and its presence as a defined area of swelling that does not cross suture lines is more consistent with this diagnosis.



Subgaleal hematoma occurs between the galea aponeurotica and the periosteum. It is commonly caused by birth trauma, particularly forceps or vacuum-assisted delivery. Subgaleal hematoma can cross suture lines and typically presents as a soft, fluctuant swelling on the scalp.

Caput succedaneum is an edema of the scalp that can cross suture lines. It is usually present at birth or develops shortly after due to pressure on the presenting part of the head during labor. It generally resolves within a few days.

Encephalocele is a neural tube defect that results in a sac-like protrusion of the brain and its meninges through an opening in the skull. It is a congenital condition and is typically diagnosed at birth or via prenatal ultrasound.

Solution to Question 2:

A patient with abdominal pain and sterile pyuria, with the given x-ray showing diffuse alveolar calcification in the right renal region point towards a diagnosis of a putty kidney.

The putty kidney is a characteristic finding of genitourinary tuberculosis (TB), specifically renal TB. The given image shows a homogenous and ground-glass-like calcification of the kidney appearing like putty. It is caused by the dissemination of the organism through the bloodstream and is thus always a secondary TB. There is either reinfection or reactivation of old TB.

Genitourinary TB is a cause of sterile pyuria. Diagnosis is confirmed with a tuberculin test, or at least 3 consecutive early-morning specimens of urine are examined for acid-fast bacilli. They are stained with a Ziehl-Neelsen stain and subsequently cultured.

CT is the most sensitive modality and is the preferred means of visualizing renal calcifications. CT urography is done to assess the various stages of renal TB and its manifestations, which are as follows:

• Early:

• Papillary necrosis

• Progressive:

• Multifocal strictures

• Generalized or focal hydronephrosis

• Mural thickening and enhancement

• Poorly enhancing renal parenchyma

• End-stage:

• Progressive hydronephrosis

• Amorphous dystrophic calcification that involves the entire kidney - putty kidney

Treatment involves short-course therapy with antitubercular drugs like pyrazinamide, isoniazid, and rifampicin. Partial nephrectomy, nephrectomy, or reconstructive surgery is often done.

Option B: Nephrocalcinosis - This condition is characterized by the deposition of calcium in the kidney parenchyma or tubules, often seen in hypercalcemic conditions like hyperparathyroidism. However, it would not explain the clinical presentation of sterile pyuria.

Option C: Staghorn calculi are large kidney stones that fill the renal pelvis and extend into the renal calyces, often caused by urinary tract infections with organisms such as Proteus, Pseudomonas, and Staphylococcus. While these calculi can cause abdominal pain, they usually do not result in sterile pyuria, as they are associated with infection.

The image below shows a staghorn calculus of the left kidney.

Option D: Psoas calcification - This refers to calcification within the psoas muscle and is often seen after acute renal failure and rhabdomyolysis. It is not associated with sterile pyuria and is not the cause of the patient's symptoms.

Solution to Question 3:

The true statement is that the intensity of radiation is inversely proportional to the square of the distance from the source in brachytherapy.

Radiotherapy is a common treatment modality for various cancers, including endometrial carcinoma. It works by targeting rapidly dividing cancer cells with ionizing radiation, which can damage the DNA of these cells and lead to cell death. Radiotherapy can be delivered externally (external beam radiotherapy) or internally (brachytherapy). In brachytherapy, radioactive sources are placed within or adjacent to the tumor, leading to a high radiation dose delivered directly to the tumor while minimizing damage to surrounding healthy tissue.

Other options:

Option B: Larger blood vessels are more radioresistant than microvasculature.

Option C: Permanent non-dividing cells are more radioresistant than rapidly proliferating cells.

Option D: Small intestinal epithelial cells (intestinal crypts), which divide rapidly with no differentiation are the most radiosensitive.

Solution to Question 4:

The given clinical scenario of fever, cough, and crepitations suggests features of pneumonia. The chest X-ray showing opacities with obscuration of the right cardiac border is suggestive of right middle lobe consolidation.

This loss of clear demarcation between the normal lung and its surrounding soft structures (heart, diaphragm) is called the silhouette sign. It occurs due to loss of contrast between lungs and soft tissue when the lung parenchyma is no longer filled with air. This sign may be seen in consolidation, atelectasis, or mass lesions.

In the following images:

• Left image - red line indicates the normal demarcation between the right heart border and middle lobe of the right lung

• Right image - loss of demarcation i.e, positive silhouette sign due to right middle lobe consolidation



Positive silhouette sign and corresponding anatomical site of pathology:

• Right heart border - right middle lobe or medial right lower lobe

• Left heart border - lingula segments of the left upper lobe

• Left hemidiaphragm or descending aorta - left lower lobe

• Aortic knuckle - left upper lobe (posterior segment)

• Right hemidiaphragm - right lower lobe

• Right paratracheal stripe - right upper lobe

Other options:

Apical segment of Right lower lobe consolidation: In this case, the chest X-ray would show an opacity or consolidation in the apical segment of the right lower lobe. The consolidation would be located in the lower part of the right lung, towards the apex. The right heart border, however, would remain visible, as it is not adjacent to the apical segment of the right lower lobe.

Posterior segment of right lower lobe consolidation: A chest X-ray revealing consolidation in the posterior segment of the right lower lobe would show an opacity in the lower part of the right lung, more posteriorly. This would not affect the visibility of the right heart border, as it is not adjacent to the posterior segment of the right lower lobe. In a posteroanterior (PA) chest X-ray, this consolidation may be difficult to appreciate, but it would be more evident in a lateral view.

Loculated pleural effusion: In a chest X-ray showing a loculated pleural effusion, there would be a fluid collection within the pleural space. This fluid collection would appear as a well-defined opacity, typically with a convex border facing the lung parenchyma. It may appear as a lenticular shape, conforming to the space between the lung and chest wall or diaphragm. The loculated pleural effusion would not create a silhouette sign with the right heart border, as it is located in the pleural space rather than the lung parenchyma.

Radiology INI-CET 2023

Question 1:

A patient presented with dysuria and pain in the back. His lumbar spine X-ray shows the below finding. What is the likely diagnosis?

a) Wilson's disease

b) Addison's disease

c) Cystic fibrosis

d) Hyperparathyroidism

Question 2:

Identify the finding in the given ultrasound image.

a) Mercedes-Benz sign

b) Gall bladder polyp

c) Gall bladder cancer

d) Phrygian cap

Question 3:

Banana sign on USG is associated with which of the following conditions?

a) Spina bifida

b) Omphalocele

c) Downs syndrome

d) Klinefelter syndrome

Question 4:

Identify the fracture site in the X-ray shown below.



a) 5th, 6th rib

b) 5th, 6th & 7the rib

c) 7th and 8th rib

d) 6th and 7th rib

Question 5:

Which of the following is not an advantage of USG over Mammography?

a) Can be used for screening mammographically undetectable clinically non palpable breast mass

b) Useful in young females with dense breast

c) Differentiates between solid and Cystic lesions in the Breast

d) Guided biopsy can be done

Question 6:

An MRCP image is shown below. Which among the following is the most likely diagnosis?

a) Choledocholithiasis

b) Cholecystitis

c) Pancreatic duct stricture

d) Malignancy

Question 7:

The image given below shows the MRI of the knee. Identify the marked structure.

a) ACL

b) PCL

c) MCL

d) Patellar tendon

Question 8:

A child was brought to the emergency with drooling and choking following ingestion of a foreign body. Based on the given X-ray, what object was most likely swallowed by the child?



a) Cap of a cool drink bottle

b) Coin

c) Button battery

d) Metal ball

Answer Key

Question No.	Correct Option
1	d
2	d
3	a
4	d
5	a
6	a

7	b
8	c

Detailed Explanations

Solution to Question 1:

The given clinical scenario and X-ray showing diffuse calcification of both kidneys are suggestive of nephrocalcinosis which is seen in hyperparathyroidism.

Nephrocalcinosis is the deposition of calcium in the kidney parenchyma or tubules which is seen in conditions causing hypercalcemia like hyperparathyroidism.

Clinical features of hyperparathyroidism:

• Bones:

• Pathological fractures

• Subperiosteal bone resorption - 2nd and 3rd digits, most common in the radial aspect

• Salt and pepper skull

• Osteitis fibrosa cystica or VonReckling Hausen's disease of bone

• Stones

• Multiple and recurrent renal stones - most common

• Nephrocalcinosis

• Abdominal groans - colicky due to pancreatitis

• Psychiatric overtones - raised PTH causing CNS depression and psychiatric symptoms

• Band keratopathy

• Features of hypercalcemia -

• Vomiting

• Confusion

• ECG changes: increased PR interval, shortened QT interval

Investigations done in primary hyperparathyroidism:

• Ultrasonography of the neck is a safe and widely used technique for the localization of abnormal parathyroid glands.

• Nuclear medicine scanning with radiolabeled Sestamibi is also a widely used technique. Sestamibi accumulates in parathyroid adenomas. This radionuclide is concentrated in the thyroid and parathyroid tissue but usually washes out of normal thyroid tissue in under an hour. It persists in abnormal parathyroid tissue.

• Dual-energy radiographic absorptiometry is a useful tool to demonstrate skeletal involvement in primary hyperparathyroidism.

Solution to Question 2:

The ultrasound examination of the gallbladder shows a Phrygian cap. It is a normal anatomical variant.

Phrygian cap refers to an anatomical variant of the gallbladder, where, the fundus of the gallbladder folds on itself, or is separated from the body by a partial or complete septum. A true septum is uncommon and is usually an apparent septum due to the gallbladder wall folding. It is an incidental finding that may be seen on CT, MRI, or ultrasound. It causes no symptoms.

The Phrygian cap is named so, due to the resemblance of the ultrasonographic appearance to a hat or cap that is worn by the inhabitants of ancient Phrygia circa 1200 BC.



Solution to Question 3:

The banana sign on USG is seen in spina bifida.

Banana sign is an abnormal anterior curvature of the cerebellum along with effacement of the cisterna magna that produces this appearance. It is seen in spina bifida and is also associated with Arnold-Chiari malformation.

Spina bifida is a congenital anomaly occurring due to incomplete development of the neural tube. The most commonly cited signs present in spina bifida are the banana sign and the lemon sign.

The lemon sign occurs due to the overlapping of the frontal bones due to the posterior shift of intracranial contents as shown below.



Other options:

Option B: Omphalocele is a congenital midline anterior abdominal wall defect through which the solid abdominal viscera or bowel herniates. The herniated abdominal contents are covered by a

membranous sac as shown below.



The image below shows an ultrasound image of the fetus suggestive of an omphalocele (asterisk).

Option C: Nuchal translucency (NT) is used to antenatally screen Downs syndrome (Trisomy 21) during the first trimester. NT is the maximum thickness of the translucent area between the skin and soft tissue over the fetal spine at the back of the neck. When the NT measurement is increased, it is a marker for fetal aneuploidy and structural anomalies. The ultrasound image below shows increased NT thickness and absent nasal bone of a fetus with Trisomy 21 at 11 weeks gestation.



Solution to Question 4:

The given X-ray shows the fracture of the 6th and 7th rib

In the chest X-ray, ribs appear as light grey strips. The anterior part of the rib is more oblique and the medial part cannot be demonstrable as it ends with costal cartilage. The posterior part of the rib is more horizontal and the attachment of the rib with the vertebra can be easily made out.

Normal Chest X-ray Rib Identification



The anterior(yellow) and posterior(red) ribs are highlighted in the image given below.



Solution to Question 5:

USG cannot be used for screening mammographically undetectable clinically nonpalpable breast mass. Although ultrasound has its advantages in certain scenarios, mammography remains the primary screening tool for breast tumor detection.

Mammography is complementary to physical examination and can detect clinically non-palpable breast mass. The primary purpose of mammography is to detect small breast tumors. While mammography is typically the initial screening tool for breast cancer detection, there are specific scenarios where ultrasound may be preferred or used in conjunction with mammography. Advantages of USG over mammography include:

• USG is preferred in pregnant women, lactating women, and young women since breast tissue is dense (Option B) in these populations.

• USG can differentiate between solid and cystic lesions (Option C) of the breast and is able to detect abnormal axillary lymph nodes.

• Ultrasound can assist in guided breast biopsy (Option D) procedures.

Solution to Question 6:

The given MRCP image shows multiple round filling defects within the common bile duct indicative of choledocholithiasis.

Choledocholithiasis or common bile duct stones are classified based on their point of origin. Choledocholithiasis is mostly clinically silent but may present with biliary colic and even obstructive jaundice. Management of choledocholithiasis includes ERCP +/- sphincterotomy followed by laparoscopic cholecystectomy.

MRCP uses the fluid which is present in the biliary and pancreatic ducts as a contrast agent and does not require external contrast material administered into the ductal system. It is a non-invasive investigation for evaluating extrahepatic, intrahepatic bile duct, and pancreatic duct. No interventions are performed under MRCP.

MRCP uses T2 weighted MRI and it recognizes slow-flowing or stationary fluid within ducts, which appear brighter relative to adjacent solid tissues. MRCP can be used to evaluate various conditions of the pancreaticobiliary ductal system, examples include:

• Congenital anomalies of the cystic and hepatic ducts

• Post-surgical biliary anatomy

• Pancreas divisum

• Anomalous pancreaticobiliary junction

• Choledocholithiasis

• Biliary structures

• Chronic pancreatitis

• Pancreatic cystic lesions

• Trauma to the biliary system

MRCP in benign strictures would typically show smooth elongated tapering of the bile duct, while malignancy-associated strictures typically show abrupt shouldering.

Solution to Question 7:

In the above given image of an MRI of the knee joint, the arrow shows the posterior cruciate ligament.

The posterior cruciate ligament (PCL) is the only primary restraint to posterior translation throughout the entire knee range of motion. It prevents the internal rotation of the tibia on the femur when there is varus stress on the knee.

It acts as a secondary restraint to valgus/varus, external rotation.

Anterior cruciate ligament	Posterior cruciate ligament
Attached to the anterior part of the intercondylar area of t he tibia	Posterior part of an intercon dylar area of the tibia
Direction – Upwards, backwa rds & laterally & attached to posterior aspect of medial su rface of lateral condyle.	Upwards, forwards & mediall y & attached to the anterior a spect of the lateral surface of the medial condyle.
Prevents posterior dislocatio n of femur on tibia and forwa rds dislocation of tibia on fe mur.	Prevents anterior dislocation of femur on tibia and posteri or dislocation of tibia on fem ur

Anterior cruciate ligament	Posterior cruciate ligament
Limits hyperextension of kne e joint(taut during extension)	Limits hyperflexion of knee j oint. (taut during flexion)
Weaker	Stronger



Solution to Question 8:

The presence of a halo sign or double ring sign is indicative of the foreign body being a button battery. Furthermore, the presence of irregularities along the periphery of the battery indicates the possibility of corrosion.

Button batteries and sharp objects in the oesophagus and foreign bodies associated with respiratory symptoms all mandate urgent endoscopic removal.

Button batteries, in particular, are very dangerous foreign bodies, which must be expediently removed because they can cause:

• Mucosal injury in 1-hour contact time

• Involve all layers of the oesophagus in 4 hours

Injury is thought to be secondary to electrochemical burns from electrical discharge. Chemical burns from electrolyte alkali leakage and pressure necrosis also may contribute to mucosal injury.

Asymptomatic blunt objects and coins lodged in the oesophagus can be observed for up to 24 hr in anticipation of passage into the stomach.

Another AP and lateral X-ray demonstrating the presence of a button battery in the oesophagus is as shown below.


R
Supine
R
HRUMW

Radiology INI-CET 2024

Question 1:

Which of the following statements is correct about CT scan?

a) CT score of water is 0

b) CT consists of unfiltered x-ray beams

c) It is the investigation of choice for gallstones

d) 20mSV for NCCT head

Question 2:

'Mount Fuji sign' is seen in _____

a) Tension pneumomediastinum

b) Tension pneumothorax

c) Tension pneumoperitonium

d) Tension pneumocephalus

Question 3:

A patient presented with a stab wound over the right chest. The supine X-ray image of the patient is given below. What is the diagnosis?



a) Right pneumothorax

b) Pulmonary contusion

c) Right hemothorax

d) Right diaphragmatic rupture

Question 4:

Which of the following contrast agents is safe to use in renal failure patients?

a) Gadolinum contrast used in MRI

b) Iodinated ionic contrast agent

c) Iodinated non-ionic contrast agent

d) Ultrasound contrast agent

Question 5:

Which of the following structures is the arrow pointing to?



a) Anterior end of rib

b) Costal cartilage

c) Posterior end of rib

d) Right pulmonary artery

Question 6:

Which of the following radiological interventions are used in the management of cerebral aneurysm?

a) Coiling

b) Sclerotherapy

c) Stenting

d) Clipping

Answer Key

Question No.	Correct Option
1	a
2	d
3	c
4	d
5	c
6	a

Detailed Explanations

Solution to Question 1:

The correct statement is that the CT score of water is 0.

On CT, each image consists of a matrix of data points (pixels) that are assigned a number that represents the X-ray attenuation(μ) at that point in the body. CT number is a scaled version of μ that is normalised to the X-ray attenuation of water (μW).

CT = 1000 × (μ/μW – 1)

The resulting numbers are called 'Hounsfield units'(HU). Due to this normalisation, water is assigned 0 HU, and air is assigned a CT number of –1000 HU.

CT numbers:

• Water = 0 HU

• Solid soft tissues are centred around 0 HU

• Lungs = -1000 HU

• Air = –1000 HU

• Dense cortical bone = +1000 HU

• Acute Blood Clot = 60-90 HU

• Fat = -100 HU

Other options:

Option B: Computed tomography (CT) imaging involves the use of a narrow beam of X-rays that are emitted from a rotating X-ray tube.CT scanners typically use a combination of X-ray beams at different energies and filters to reduce scatter and improve image quality. Therefore, the X-ray beams used in CT imaging are often filtered to enhance image contrast and reduce radiation dose to the patient.

Option C: The investigation of choice for gallstones is an Abdominal ultrasound and not a CT Abdomen. An ultrasound can assess for complications of gallstones, such as inflammation of the gallbladder (cholecystitis) or obstruction of the bile ducts. In cases where ultrasound findings are inconclusive or further characterisation is needed, other imaging modalities such as CT scan or magnetic resonance cholangiopancreatography (MRCP) may be used.

Option D: The mSv (millisievert) value for a non-contrast computed tomography (NCCT) scan of the head typically ranges from around 1 to 2 mSv. This radiation dose can vary depending on factors such as the specific CT scanner used, the scanning protocol, and the individual patient characteristics. However, in general, NCCT head scans are associated with relatively low radiation doses compared to other medical imaging procedures.

Solution to Question 2:

'Mount Fuji sign' is seen in tension pneumocephalus.

Mount Fuji sign refers to the presence of air (pneumocephalus) between the tips of the frontal lobes giving the appearance of Mount Fuji.

Tension pneumocephalus occurs with subdural air causing a mass effect over the underlying brain parenchyma, often from a ball valve mechanism causing one-way entry of air into the subdural space.

The appearance of this sign often warrants immediate surgery to prevent permanent damage.

The CT image below shows the Mount Fuji sign.



Solution to Question 3:

The history of stab injury to the right chest and the X-ray showing opacity of the right hemithorax, with a slight tracheal deviation towards the opposite side, indicates a diagnosis of right hemothorax.

In a plain chest radiograph, it's difficult to differentiate an acute hemothorax from other collections of pleural fluid as they are indistinguishable. As blood clots, it often leads to localisation, occasionally forming a fibrin body. Pleural thickening and calcification are acknowledged as potential consequences.

The following are the findings commonly seen in a case of hemothorax:

• Blunting of Costophrenic Angle: Hemothorax results in the accumulation of blood in the pleural cavity, causing the lung to collapse partially. This can lead to the blunting or obliteration of the costophrenic angle on the affected side.

• Increased Opacity: The presence of blood in the pleural space leads to increased opacity or density in the affected hemithorax. This can manifest as a homogenous shadowing that obscures the lung markings.

• Mediastinal Shift: Depending on the volume of blood in the pleural cavity, there may be a mediastinal shift towards the opposite side. However, in mild to moderate hemothorax, this may not be readily apparent on supine X-rays.

• Air-Fluid Level: In cases where there is concurrent pneumothorax (air in the pleural cavity), an air-fluid level may be visible, with the blood layering dependently and the air layer superiorly.

• Unilateral Findings: The abnormalities will typically be limited to one side of the chest, corresponding to the side of the hemothorax.

In cases of suspected hemothorax, a lateral decubitus or chest X-ray in erect position may be more sensitive for detection. Additionally, other imaging modalities such as ultrasound or computed tomography (CT) scans can provide more detailed information about the presence and extent of a

hemothorax.

Other options:

The following image is characteristic of a pneumothorax (left-sided) with radiolucency over the left lung, lung collapse, and mediastinal shift on x-ray.



The given radiograph is a case of diaphragmatic hernia that shows elevation of the left hemidiaphragmatic line along with the bowel loops and stomach seen in the pleural cavity.



The following is an image of a right-sided pulmonary contusion post-blunt chest trauma. Note the increased opacity just below the horizontal fissure.



Solution to Question 4:

Ultrasound contrast agents are gas-filled microbubbles that are administered intravenously to the systemic circulation. Hence, considered as low risk in patients with renal impairment.

Ultrasound contrast agents are used to enhance the visibility of blood vessels, organs, and tissues during ultrasound imaging. These agents typically consist of tiny gas-filled microbubbles encapsulated within a shell. When injected intravenously, they flow through the bloodstream and resonate in response to ultrasound waves, producing strong echoes that enhance the ultrasound image.

Perflutren/Optison is an ultrasound contrast agent. Levovist (consisting of galactose microparticles, air bubbles, and palmitic acid) is another example of an ultrasound contrast agent. Sulfur hexafluoride with a phospholipid shell is also used as an ultrasound contrast agent.

Other options:

Option A: Gadolinium contrast used in MRI is predominantly excreted through the kidney. In renal failure, it gets accumulated in the body and releases gadolinium ions. Since the ion is toxic, the body tries to contain it by fibrosis resulting in multisystem painful fibrosis called nephrogenic systemic fibrosis.

Option B: Iodinated ionic contrast agents increase the risk of Contrast-induced nephropathy(CIN). Hence alternative imaging methods, without iodinated contrast media, should be chosen in patients with considerably impaired renal function (CKD 4 or 5).

Option C: Non-ionic contrast agents have a lower risk of causing adverse reactions compared to ionic contrast agents. They are generally better tolerated by patients, with fewer incidences of allergic reactions and nephrotoxicity. However, USG contrast agents are among the safest in renal failure patients.

Solution to Question 5:

The structure pointed by the arrow mark in the given chest X-ray PA view is the posterior end of the rib.

In the chest X-ray PA view, ribs appear as light grey strips. The anterior part of the rib is more oblique and the medial part cannot be demonstrable as it ends with costal cartilage. The posterior part of the rib is more horizontal, hyperdense and the attachment of the rib with the vertebra can be easily made out.

Normal Chest X-ray Rib Identification



The anterior(yellow) and posterior(red) ribs are highlighted in the image given below.



Solution to Question 6:

Coiling is the radiological intervention used in the management of cerebral aneurysm.

Aneurysms occur at the bifurcations of the large to medium-sized intracranial arteries. Rupture is into the subarachnoid space in the basal cisterns. Approximately 85% of the aneurysms occur in the anterior circulation, mainly on the circle of Willis.

Most unruptured intracranial aneurysms are completely asymptomatic. When the aneurysm ruptures, 45% of cases present with sudden onset severe headache, described as the worst headache of the patient's life. Other features include neck rigidity and transient loss of consciousness.

Diagnosis -

CT shows hyperattenuating material filling the subarachnoid space. CT angiography has a high sensitivity for aneurysms and AVMs, but digital subtraction angiography remains the gold standard.



Treatment -

Early aneurysm repair prevents re-rupture. At many centers, definitive repair is carried out within 24 h of the bleed in all patients who are stable enough to tolerate the procedure. An aneurysm can be “clipped” by a neurosurgeon or “coiled” by an endovascular approach. Surgical repair involves placing a metal clip across the aneurysm neck, thereby immediately eliminating the risk of rebleeding. This approach requires craniotomy (option D) and brain retraction, which is associated with neurologic morbidity.

Endovascular techniques involve placing platinum coils within the aneurysm via a catheter that is passed from the femoral artery. The aneurysm is packed tightly to enhance thrombosis and over time is walled off from the circulation.

Intracranial pressure should be measured and the cerebral perfusion pressure targeted at 60–70 mmHg.

Delayed cerebral ischemia remains the leading cause of morbidity and mortality following aneurysmal SAH. Treatment with the calcium channel antagonist nimodipine (60 mg PO every 4 h) improves outcome, perhaps by preventing ischemic injury rather than reducing the risk of vasospasm.

The image below is an arteriogram showing a coiled aneurysm (indicated by yellow arrows) of the posterior cerebral artery with a residual aneurysmal sac.



Other options :

Option B - Sclerotherapy is used in the management of venous insufficiency.

Option D - Stenting is employed mainly for stenotic vascular lesions.

Radiology NEET 2024

Question 1:

A patient presented with chest pain and hoarseness for one month, and on a CT angiogram, the following image is seen, what is the possible diagnosis?



a) Stenosis of aortic arch

b) Aortic dissection

c) Saccular aneurysm of distal aorta

d) Coarcatation of aorta

Question 2:

This is the radiograph of a child with a foreign body in the oesophagus. What is the sign seen in this?



a) Thumb sign

b) Steeple sign

c) Double ring sign

d) String sign

Question 3:

A patient presents with a one-month history of lethargy, headache, fever, and vomiting. Shortly after being admitted to the hospital, the patient tragically passes away. Radiological imaging reveals multiple cystic lesions in the cerebrum with invasion into the ventricles and subarachnoid space. What is the likely cause?



a) Neurocysticercosis

b) Astrocytoma

c) AV malformation

d) Brain metastasis

Question 4:

A patient complains of severe headache. NCCT and CT angiography images are as below. What is the likely etiology?



a) AV malformation

b) Subarachnoid hemorrhage

c) Intraparenchymal hemorrhage

d) Subdural hemorrhage

Question 5:

An elderly woman with a known history of Myasthenia Gravis presents with symptoms including fatigue, facial congestion, dilated neck veins, and breathlessness, which are indicative of superior vena cava (SVC) obstruction. What is the likely diagnosis?



a) Lipoma

b) Ectopic thyroid

c) Thymoma

d) Germ cell tumor

Question 6:

A 4-week-old infant presented to the ER with projectile vomiting. USG abdomen findings of the infant are given below. Probable diagnosis is _____

a) Sigmoid volvulus

b) Duodenal atresia

c) Congenital hypertrophic pyloric stenosis

d) Intussusception

Question 7:

An infant presents with respiratory distress. He has a previous history of recurrent respiratory infections. What is the diagnosis?



a) Lung abscess

b) Congenital malformations of the airway

c) Congenital diaphragmatic hernia

d) Pyopneumothorax

Question 8:

A young patient presents with nasal polyps and rhinosinusitis. X-ray and clinical images are as below. What is the diagnosis?





a) Young's syndrome

b) Kartagener's syndrome

c) Churg Strauss syndrome

d) Cystic fibrosis

Question 9:

Given below is a USG taken at approximately 8 weeks of gestation. Which anomaly can be observed here?

a) Anencephaly

b) Hydranencephaly

c) Meningocele

d) Encephalocele

Question 10:

Match the following using the radiological image given below:



a) 3-A, 2-B, 1-C, 4-D

b) 2-A, 3-B, 4-C, 1-D

c) 1-B, 2-C, 3-A, 4-D

d) 2-A, 1-B, 3-C, 4-D

Question 11:

Identify the diagnosis from the given CT scan.

a) PUJ obstruction

b) Renal cyst

c) Renal tumour

d) Multicystic kidney

Question 12:

What does this image below indicate?



a) Choledochal cyst

b) Gall bladder carcinoma

c) Cholangiocarcinoma

d) Choledocholithiasis

Question 13:

Identify the condition shown in the image:



a) Chronic calicific pancreatitis

b) Mesenteric calcification

c) Horseshoe kidney

d) Jejunal fecolith

Question 14:

A 6-year-old child presented with seizures which subsided with injection Midazolam. Blood investigations and lumbar puncture studies were normal. MRI is done and the image is given below. What is the next best investigation of choice for further assessment?



a) MR spectroscopy

b) Ocular B scan

c) CBNAAT of CSF

d) PET scan

Answer Key

Question No.	Correct Option
1	c
2	c
3	a
4	b
5	c
6	c
7	b
8	b
9	a
10	c
11	a
12	a
13	a
14	a

Detailed Explanations

Solution to Question 1:

The given CT shows saccular aneurysm of the distal aorta. The clinical presentation of chest pain and hoarseness supports the diagnosis.

Other options:

Option A: Stenosis of the aortic arch would show a narrowing rather than an aneurysmal dilation, and it is more likely to present with symptoms related to decreased blood flow to the upper body.

Option B: Aortic dissection would usually present with a tearing chest pain that radiates to the back and would show a double lumen or false lumen on imaging. The below image shows Stanford type A aortic dissection:



Option D: Coarctation of the aorta typically presents with hypertension in the upper limbs and weak or absent pulses in the lower limbs.

Solution to Question 2:

The presence of a halo sign or double ring sign is indicative of the foreign body being a button battery.

Button batteries and sharp objects in the oesophagus and foreign bodies associated with respiratory symptoms all mandate urgent endoscopic removal.

Button batteries, in particular, are very dangerous foreign bodies, which must be expediently removed because they can cause:

• Mucosal injury in 1-hour contact time

• Involve all layers of the oesophagus in 4 hours

Injury is thought to be secondary to electrochemical burns from electrical discharge. Chemical burns from electrolyte alkali leakage and pressure necrosis also may contribute to mucosal injury.

Asymptomatic blunt objects and coins lodged in the oesophagus can be observed for up to 24 hr in anticipation of passage into the stomach.

Another AP and lateral X-ray demonstrating the presence of a button battery in the oesophagus is as shown below.



Other options

Option A: Thumb sign is seen in acute epiglottitis.



Option B: A steeple sign is seen in croup or laryngotracheobronchitis.



Option D: String sign on the barium enema is in ileocecal tuberculosis.



The image below shows the string sign of Kantor associated with Crohn's disease.



Solution to Question 3:

The given clinical scenario along with CT showing cystic lesions in the brain, particularly within the cerebrum, ventricles, and subarachnoid space is clearly suggestive of neurocysticercosis.

Neurocysticercosis is a form of cysticercosis caused by Taenia solium.

It usually presents with seizures and symptoms of increased intracranial pressure, including headache, nausea, vomiting, vision changes, dizziness, ataxia, or confusion. Papilledema may develop secondary to hydrocephalus.

Diagnosis is done by demonstrating eggs of Taenia in faecal examination.

A biopsy of the brain or spinal cord lesion or fundoscopic visualization of subretinal cysticercus can also be done.

Imaging shows a scolex within a cystic lesion, cystic lesions with or without enhancement (e.g., ring enhancement), nodular calcifications, focal enhancing lesions, or multilobulated cystic lesions in the subarachnoid space.

Treatment of intestinal T. solium infection is by a single dose of praziquantel. Neurocysticercosis is treated with anti-epileptic drugs, anticysticidal drugs like albendazole or praziquantel, and steroids given before anticysticidal therapy.

Other options:

Option B: Astrocytoma is a type of glial cell tumour in the brain, ranging from low to high grade. They do not typically present as multiple cystic lesions and are more likely to appear as a single mass with variable enhancement on imaging.

Option C: Arteriovenous malformation (AVM) is a congenital vascular anomaly characterized by a tangle of abnormal blood vessels connecting arteries and veins. AVMs are more likely to cause hemorrhage or seizures and typically appear as highly vascular lesions on imaging, not as cystic lesions.

Option D: While metastases can present as multiple lesions on imaging, they usually appear solid rather than cystic masses and often enhance with contrast.

Solution to Question 4:

The clinical manifestations of severe headache with CT revealing blood in the basal cistern point to a diagnosis of subarachnoid haemorrhage (SAH).

Subarachnoid haemorrhage (SAH) is bleeding into the subarachnoid space, which is the area between the arachnoid membrane and the pia mater surrounding the brain. The diagnosis is suspected when a hyperattenuating material is seen filling the subarachnoid space. Most commonly this is apparent around the circle of Willis, on account of the majority of berry aneurysms occurring in this region (65%), or in the Sylvian fissure (30%).

The modified Fisher scale is a method for grading SAH as seen on non-contrast CT (NCCT). The Hunt and Hess scale describes the severity of SAH and is used as a predictor of survival. Xanthochromia on CSF is indicative of SAH in case of a negative CT scan.

Other options:

Option A: Arteriovenous malformation (AVM) is a congenital vascular anomaly characterized by a tangle of abnormal blood vessels connecting arteries and veins. AVMs are more likely to cause haemorrhage or seizures and typically appear as highly vascular lesions on imaging, not as cystic lesions.

Option C: Intraparenchymal hemorrhage results from damage to lenticulostriate arteries. Hyperdensity within the brain parenchyma is seen on CT.

Option D: Acute subdural haemorrhage results from the accumulation of blood between the arachnoid membrane and the dura. The tearing of a bridging vein from the cerebral cortex to the dural sinuses is the most common cause of acute SDH. On head CT scan, acute SDH appears as a hyperdense, crescent-shaped (concavo-convex).

Solution to Question 5:

The diagnosis of myasthenia gravis along with the CT scan image showing an anterior mediastinal mass, is suggestive of thymoma.

Thymoma is the most common mediastinal tumor and is derived from the thymus gland. Thymomas vary in behaviour from benign to aggressively invasive. Superior vena cava obstruction is a possible complication of mediastinal mass.

Masoaka classification system is used to stage thymomas.

Myasthenia gravis is often related to thymomas. Symptoms of myasthenia gravis can improve after excision of the thymus gland even in the absence of a thymoma.

Complete thymectomy is the treatment for thymoma. It can be performed by two approaches:

• A median sternotomy approach

• A transcervical approach with or without an additional VATS procedure (Video-Assisted Thoracoscopic Surgery) for smaller thymomas

Other options:

Option A: Lipomas are benign and slow-growing, less likely to cause significant compression of the SVC and produce the acute symptoms described.

Option B: Ectopic thyroid tissue can be found in the mediastinum, but it is typically asymptomatic and does not usually cause SVC obstruction.

Option D: Germ cell tumours can occur in the mediastinum and cause SVC obstruction, but they are less commonly associated with Myasthenia Gravis compared to thymomas.

Solution to Question 6:

The probable diagnosis is Congenital Hypertrophic Pyloric stenosis since the cervix sign and nipple sign are signs of Congenital Hypertrophic Pyloric stenosis on USG.

• Double-layered hypertrophic mucosa protrudes into the stomach, creating the so-called Nipple sign.

• The sonographic appearance of the hypertrophied pylorus resembles that of the uterine cervix and is sometimes referred to as the cervix sign.



Hypertrophic pyloric stenosis most presents with non-bilious projectile vomiting. It is common between 2 and 8 weeks of age. Hypertrophy of the circular muscle of the pylorus results in constriction and obstruction of the gastric outlet. Male infants are at a higher risk.

Clinically, an olive-shaped mass is palpable in the right upper quadrant or epigastrium. When the infant is given a test feed, visible peristalsis is usually observed moving from left to right across the abdomen.

Infants with hypertrophic pyloric stenosis develop a characteristic metabolic abnormality-hypochloraemic hypokalaemic metabolic alkalosis. Persistent vomiting results in loss of

hydrochloric acid leading to the development of metabolic alkalosis (due to loss of H+) and hypochloremia (due to loss of Cl-).

Ultrasound examination is the investigation of choice for this condition. The diagnostic criteria for pyloric stenosis are a muscle thickness of ≥4 mm and a pyloric length of ≥16 mm. The cervix sign and nipple sign are signs seen on USG.

As for the management, the child should be rehydrated and the metabolic abnormality has to be corrected before proceeding to surgery. The Ramstedt pyloromyotomy remains the standard surgical procedure for its treatment.

Solution to Question 7:

The given clinical scenario along with the image is suggestive of congenital malformations of the airway.

Congenital pulmonary airway malformation (CPAM) is a condition where more normal lung tissue is mixed with hamartomatous or dysplastic lung tissue, usually confined to one lobe. The lesion most likely originates from a maldevelopment of terminal bronchiolar structures caused by an embryologic injury that occurred prior to the 35th day of gestation.

Pneumothorax, recurrent respiratory infections, and respiratory distress can all manifest in patients during the newborn or early infancy period. One could mistake the lesion for a diaphragmatic hernia. Until mid-childhood, when episodes of recurrent or persistent pulmonary infection or chest pain occur, patients with smaller lesions typically are asymptomatic.

Treatment includes aspiration of macrocystic lesions, excision of the affected lobe for microcystic lesions, and infrequently, open foetal surgery for severely affected infants. For patients with symptoms in the postnatal period, surgery is recommended.

Other options:

Option A: Lung abscess represents necrosis and cavitation of the lung following microbial infection.

Chest X-ray features of lung abscess include a lung cavity with an air-fluid level and a thick and irregular cavity wall with surrounding pulmonary infiltrate. The image showing the lung abscess is given below:

Option C: Congenital diaphragmatic hernia (CDH) is a birth defect of the diaphragm. Malformation of the diaphragm allows the abdominal organs to push into the chest cavity, hindering proper lung formation and resulting in pulmonary hypoplasia.

The most common type of CDH is a Bochdalek hernia, which in most cases involves the left side. CDH is a life-threatening pathology in infants and a major cause of morbidity and mortality due to two complications: pulmonary hypoplasia and pulmonary hypertension. The image of Bochdalek hernia is given below:

Option D: Pyopneumothorax is a pleural collection of pus and gas, also referred to as empyemic hydropneumothorax or infected hydropneumothorax. It could be compared to a thoracic empyema variation that contains gas. Chest pain and fever are typically the patient's initial complaints.

X-ray shows gas-fluid level. It could have loculations. There appears to be a loss of silhouette around the diaphragm dome. The thick pleural lining is strongly indicative of the

pyopneumothorax.

Solution to Question 8:

The given clinical scenario along with an X-ray showing dextrocardia, CT showing maxillary polyp, and endoscopy showing nasal polyp is suggestive of Kartagener's syndrome.

Primary ciliary dyskinesia (also called immotile cilia syndrome) is an autosomal recessive disorder affecting ciliary motility. When a patient has bronchiectasis, chronic sinusitis, and situs inversus triad, he is known to have Kartagener syndrome. It occurs due to the absence of protein dynein in the A tubules of the respiratory mucosa.



The disease is usually diagnosed based on the clinical picture and should be evaluated in later stages when a child presents with recurrent lower and upper respiratory tract infections. Other features of primary ciliary dyskinesia include congenital heart diseases and neonatal hydrocephalus. Male patients with this condition tend to be infertile, due to asthenospermia (immotile spermatozoa due to impaired sperm flagella function).

Transmission electron microscopy is the gold standard investigation to analyze cilia. There is no specific treatment for this condition and is managed according to the symptoms.

Other options:

Option A: Churg Strauss syndrome clinically presents with asthma, fever and eosinophilia.

Option C: Young's syndrome clinically presents with infertility (azoospermia) and bronchiectasis.

Option D: Cystic fibrosis is an autosomal recessive condition due to a mutation of the CFTR gene on chromosome 7. Clinical features include recurrent sinopulmonary infection, intestinal obstruction, pancreatic insufficiency, and male infertility. Diagnosis is made by elevated sweat chloride levels, abnormal nasal potential difference, and genetic testing for CFTR mutation.

Solution to Question 9:

The given ultrasound image shows an open defect in the calvaria and skin. This is consistent with the diagnosis of anencephaly.

Anencephaly results from incomplete closure of the neural tube. It is one of the earliest detectable congenital defects that can be detected as early as 10 to 11 weeks of gestation. Screening can also be done by serum estimation of maternal serum alpha-fetoprotein (MSAFP). The following are the radiological signs seen in anencephaly.

• Frog eye sign - Absence of cranial calvaria, prominent orbits thereby resembling the face of a frog.

• Shower cap sign - The absence of the cranium could lead to the protrusion of a disorganized mass of brain tissue. This resembles a shower cap giving rise to a characteristic triangular facial appearance.

Neural-tube defects can be prevented with pre-conceptional supplementation of folic acid. A dose of 4 mg daily is recommended in subsequent pregnancies.

The frog eye and showercap signs are shown below:



Other options:

Option B: Hydranencephaly: This condition involves the absence of cerebral hemispheres, replaced by fluid-filled sacs.

Option C: Meningocele: It involves herniation of the meninges through a defect in the skull or spinal column, forming a sac filled with cerebrospinal fluid.

The image below shows meningocele:

Option D: Encephalocele is caused due to defects in the skull through which the meninges and/or neural tissue herniate. The squamous part of the occipital bone is the most frequently affected bone.

The image shows a newborn with encephalocele.

Solution to Question 10:

Here are the correctly matched structures:

1	B. Trachea
2	C. Esophagus
3	A. Arch of Aorta
4	D. Superior vena cava



Solution to Question 11:

The CT scan showing ballooning of the left renal pelvis with moderate hydronephrosis of the affected kidney indicating a likely diagnosis of ureteropelvic junction obstruction (UPJ)of the left side.

PUJ obstruction refers to a blockage or narrowing at the junction where the renal pelvis transitions into the ureter, leading to impaired drainage of urine from the kidney.

Indications for surgery in patients with UPJ obstruction include:

• On the diuretic renogram, UPJO with less than 40% in the split function of the afflicted kidney.

• Renal parenchymal atrophy in the presence of severe bilateral UPJO.

• Despite preventive medicines, recurring urinary tract infections.

• Obstructive UPJO that is symptomatic or linked with an abdominal mass

Hydronephrosis may be detected on antenatal ultrasound; it can, however, be transient or mild and resolves on its own. Re-evaluation should be done after birth and in cases presenting with symptoms of recurrent UTI, palpable flank mass or pain, urinary obstruction, and in cases of failure to thrive.

If the hydronephrosis is persistent, its severity is graded by the extent of the dilatation of the pelvicalyceal system and parenchymal thickness. It needs to be ascertained whether the other kidney is involved, the lower ureter is dilated, the bladder wall is thickened, and whether emptying of the bladder is incomplete. Based on these findings, a diagnosis can be made.

In cases of hydronephrosis, long-standing urine stasis and pressure may damage the kidneys.

The causes of UPJ obstruction are

• Intrinsic (due to an adynamic or stenotic segment of the proximal ureter) or

• Extrinsic (e.g. compression by accessory renal arteries). This needs to be corrected primarily.

The renal anatomy and function can be assessed using radioscintigraphy studies.
The investigation of choice for UPJ obstruction is 99mTc-MercaptoAcetyl-triGlycine (MAG3) scan. Micturating cystourethrogram is the investigation of choice in vesicoureteric reflux. 99mTc-dimercaptosuccinic acid (DMSA) scan is used for assessing renal scarring.

Treatment of intrinsic UPJ obstruction is done surgically if renal function is compromised or if there is bilateral involvement of the kidneys. In other cases, serial follow-up with ultrasonography is done to assess the spontaneous resolution of hydronephrosis.

Surgical options include pyeloplasty which entails the surgical division of the obstruction and the creation of a funnel-shaped anastomosis. Additionally, a temporary stent and a drain may be inserted. It is necessary to do further serial ultrasounds and MAG-3 imaging in order to ensure proper monitoring and evaluation. This procedure is now commonly being performed via laparoscopy and also with robotic assistance. Congenital PUJ obstruction is treated surgically by Anderson Hyen's pyeloplasty.

Below are CT scans of a left PUJ obstruction before(left) and after(right) surgery.



Other options:

Option B: A renal cyst would appear as a fluid-filled, well-demarcated structure within the kidney but does not involve the PUJ or cause obstruction.

Option C: A renal tumor would present as a solid, abnormal mass within the kidney, not typically involving the PUJ unless there is direct extension.

Option D: Multicystic kidney refers to the presence of multiple cysts within the kidney, often congenital, but is not specific to the PUJ and would not result in obstruction at the PUJ.

Solution to Question 12:

The given MRCP image showing saccular dilatation of the common bile duct suggests a diagnosis of type I choledochal cyst.

Todani classification:

• Type I- saccular dilatation of the whole (Ia) or a segment (Ib) of the common bile duct (CBD)

• Type II- diverticulum from the CBD

• Type III- choledochocele that opens into the duodenum

• Type IV- multiple cysts (both intrahepatic and extrahepatic)

• Type V- multiple cysts that are only intrahepatic (Caroli's disease)

The diagnosis is made initially with a right upper quadrant ultrasound, following which a confirmatory MRCP can be carried out to help in pre-operative planning.

Treatment is usually surgical with a Roux-en-Y hepaticojejunostomy with radical resection of all the biliary tissue. There is a risk of malignant transformation in long-standing disease, so surgical removal is usually recommended.

Other options:

Option B: Gall bladder carcinoma would show abnormal growths or masses in the gall bladder rather than cystic dilations of the bile ducts.

Option C: Cholangiocarcinoma is a malignancy of the bile ducts and would present as irregular masses or bile duct strictures rather than cystic structures.

Option D: Choledocholithiasis refers to the presence of stones in the bile duct, which would appear as filling defects within the bile ducts rather than cystic dilations.

Solution to Question 13:

As the given image shows, there are multiple calcifications within the region of the pancreas, which is characteristic of chronic calcific pancreatitis.

This condition results from long-standing inflammation of the pancreas, leading to the deposition of calcium. Chronic pancreatitis is often associated with alcohol abuse, genetic factors, or recurrent acute pancreatitis. The calcifications are a hallmark feature and help differentiate it from other abdominal pathologies.

It presents with symptoms like chronic abdominal pain with a history of alcoholism and bulky stools which indicates a lack of pancreatic lipases seen in chronic disease leading to impaired fat digestion and absorption.

Chronic pancreatitis is a progressive inflammatory disease in which there is an irreversible destruction of pancreatic tissue. It is more common in males, with a male: female ratio of 4:1.

The etiology of chronic pancreatitis and pancreatic exocrine insufficiency can be classified based on the TIGAR-O classification system:

• Toxic-metabolic -alcohol, smoking, hypercalcemia, hyperlipidemia

• Idiopathic

• Genetic- PRSS1, CFTR, and SPINK1 genes

• Autoimmune- autoimmune chronic pancreatitis, IgG4-related disease

• Recurrent and severe acute pancreatitis

• Obstructive- pancreas divisum, duct obstruction, malignancy

The diagnosis can be made using Rosemont criteria. It is based on endoscopic ultrasonographic (EUS) findings. It is explained in the pearl attached below.

ERCP is the most sensitive radiologic test for the diagnosis of chronic pancreatitis.

The below image shows the X-ray findings:



Other options:

Option B: Mesenteric calcification would present in a different anatomical region and is associated with conditions like ischemia.

Option C: Horseshoe kidney involves the kidneys and does not cause pancreatic calcifications.

Option D: Jejunal fecolith refers to hardened stool within the small intestine.

Solution to Question 14:

The given clinical history of a child along with an MRI showing a ring-enhancing intraaxial lesion of the brain could likely be a tuberculoma or neurocysticercosis and MR spectroscopy would be the next best investigation for further assessment.

Magnetic Resonance (MR) spectroscopy is a non-invasive test based on the principle of magnetic resonance used to analyse the metabolite distribution within the brain tissue. It contains N-acetyl aspartate (NAA), a neuronal viability signal; choline (a component of cell membranes and a marker of cellular turnover); creatine (present in metabolically active tissues); lipids (released with cell membrane disruption); and lactate (a hallmark of anaerobic metabolism).

The image below shows a normal MR spectroscopy.



In tuberculoma, MR spectroscopy has reduced NAA owing to the destruction of the normal brain parenchyma and increased levels of lactate or lipids. In neurocysticercosis, there is reduced NAA and creatine and elevated lactate, alanine, succinate, and choline levels.

Other options:

Option B- Ocular B- scan is done to rule out diseases in the posterior segment in the presence of media opacities which is not significant for this case.

Option C- CBNAAT of CSF will not be the next investigation as the lumbar puncture analysis was normal.

Option D- PET-CT is done to assess the stage of cancer and is hence not required in this case.

Medicine AIIMS 2017

Question 1:

An 18-year-old girl with the diagnosis of acute promyelocytic leukemia was treated medically. She developed fever and tachypnea and a chest X-ray showed pulmonary infiltrates. What drug should she be given next?

a) Cytarabine

b) Dexamethasone

c) Doxorubicin

d) Methotrexate

Question 2:

The definition of fever of unknown origin (FUO) includes all of the following, except:

a) Undiagnosed after one week of investigation

b) Immuno compromised status should be ruled out

c) Temperature more than 38.3 C

d) Fever persisting for more than 3 weeks

Question 3:

A 43-year-old patient presented with spikes of fever and difficulty in breathing. Transesophageal echo revealed vegetations in the heart. The culture was positive for Burkholderia cepacia. Which of the following is the first line of management?

a) Aminoglycosides and colistin

b) Cotrimoxazole with 3rd generation cephalosporin

c) Cefepime and tigecycline

d) Carbepenams and 3rd generation cephalosporins

Question 4:

The radio-iodine uptake scan of a patient evaluated for swelling of the thyroid is shown below. Which of the following is the most likely diagnosis?



a) Hypersecreting adenoma

b) Papillary Carcinoma thyroid

c) Graves disease

d) Lateral aberrant thyroid

Question 5:

Which of the following diseases primarily involves the marked structure?



a) Paralysis agitans

b) Depression

c) Alzheimers

d) Huntington chorea

Question 6:

All are components of CURB65 except

a) Confused state

b) Respiratory rate of > 30 /min

c) Blood urea nitrogen value greater than 7 mmol/L

d) Systolic blood pressure less than 100 mm Hg

Question 7:

A 22-year-old female patient came with complaints of vomiting and a decrease in movement of the right leg. In the past, she has had episodes of violent and aggressive behavior and abdominal pain. Which of the following is the most probable diagnosis?

a) AIDP with migraine

b) Right side paralysis

c) Acute intermittent porphyria

d) Conversion disorder

Question 8:

A 36 year old male came with complaints of ascending weakness in both lower limbs and upper body. There was no history of trauma. Which of the following investigations is done in the evaluation of this patient?

a) Serum potassium

b) Serum magnesium

c) Serum sodium

d) Serum calcium

Question 9:

On thyroid function test analysis, TSH value was much higher than normal with a low free T4. What is the diagnosis?

a) Subclinical hypothyroidism

b) Hyperthyroidism

c) Secondary hypothyroidism

d) Primary hypothyroidism

Question 10:

Which of the following is responsible for the elevated levels of TG and VLDL in diabetes mellitus?

a) Increased activity of lipoprotein lipase and decreased activity of hormone sensitive lipase

b) Increased activity of hormone sensitive lipase and decreased lipoprotein lipase activity

c) Increase in peripheral function of LDL receptors

d) Increased in activity of hepatic lipase

Question 11:

What is the window period for thrombolysis in case of stroke?

a) 1 ½ hours

b) 2 ½ hours

c) 3 ½ hours

d) 4 ½ hours

Question 12:

A 53-year-old man was admitted with a history of a cerebrovascular accident. On the second day, he becomes drowsy with minimal response. CT finding is shown in the image below. What will be your next line of management?



a) Aspirin and clopidogrel

b) Mannitol

c) Decompressive surgery

d) Mechanical thrombectomy

Question 13:

Disappearance of loud S1 in mitral stenosis occurs in all except:

a) First degree heart block

b) AR

c) Calcified valve

d) Mild MS

Question 14:

The given ECG depicts which of the following conditions:



a) Paroxysmal supraventricular tachycardia

b) Ventricular bigeminy

c) P pulmonale

d) Re-entrant atrial fibrillation

Question 15:

As an intern in the emergency ward, which of the following method would you consider as correct regarding nasogastric tube length estimation?

a) Tip of nose to ear to xiphisternum

b) Tip of nose to angle of ear to umbilicus

c) Mouth to ear to umbilicus

d) Mouth to ear to midway between xiphisternum and umbilicus

Question 16:

A morbidly obese diabetic woman is not responsive to metformin therapy. She has a history of pancreatitis and family history of bladder cancer. She does not want to take injections. Which of the following would be suitable to reduce her glucose levels?

a) Sitagliptin

b) Pioglitazone

c) Canagliflozin

d) Liraglutide

Question 17:

A patient has a right upper limb BP of 180/95 mm Hg and a left upper limb BP of 130/90 mm Hg. He also has an early diastolic murmur in right 2nd intercostal space. Which of the following is least likely to be associated with these findings?

a) Supravalvular aortic stenosis

b) Coarctation of aorta

c) Takayasu arteritis

d) Aortic dissection

Question 18:

A young donor came to the blood bank for the first time for platelet apheresis with platelet count of 1.9L. During the course he developed paresthesias and circumoral numbness during donation. His vitals remained stable though. ECG showed tachycardia with ST-T changes. What is the reason for his symptoms?

a) Hypovolemic shock

b) Hypocalcemia

c) Seizures

d) Allergic reaction

Question 19:

Which of the following anti-diabetic drugs should be stopped in a patient with a creatinine of 5.6 mg/dL?

a) Metformin

b) Linagliptin

c) Metoprolol

d) Insulin

Question 20:

A 53-year-old patient was admitted complains of dyspnea. On examination he has puffy face with engorged veins over the chest. SVC obstruction is suspected. Chest X-ray shows mediastinal enlargement. What is the next step?

a) Total blood count with peripheral smear

b) CT thorax

c) Start cyclophosphamide

d) Urgent referral to RT

Question 21:

A patient with ascites is found to have SAAG ≥1.1 . All the following can be the causes except

a) Cirrhosis

b) Liver failure

c) Metastasis to liver

d) TB peritonitis

Question 22:

A patient on anti-tubercular therapy develops tingling sensation in the limbs. Which of the following when substituted can result in improvement of symptoms?

a) Thiamine

b) Pyridoxine

c) Folic acid

d) Methylcobalamine

Question 23:

A patient comes to ED with fever and headache. On examination he has neck stiffness. CSF analysis was done and it shows the following findings. Most likely diagnosis is:

a) TB

b) Neisseria Gonorrhoea

c) Cryptococcus

d) Coxsackie

Question 24:

Metabolic acidosis with a normal anion gap is seen in all except _____.

a) Proximal RTA

b) Pancreatitis

c) Severe diarrhoea

d) Salicylate poisoning

Question 25:

What is the drug of choice for scrub typhus?

a) Doxycycline

b) Ciprofloxacin

c) Chloramphenicol

d) Azithromycin

Question 26:

A patient came with sudden onset palpitations. ECG showed a narrow complex, regular tachycardia without P waves, and a rate of 160 BPM. His blood pressure is 96/68 mmHg. What is the next step in management?

a) Carotid sinus massage

b) Adenosine

c) Beta-blocker

d) Defibrillator

Question 27:

Management of hyperkalaemia includes all except

a) Calcium gluconate

b) Insulin drip

c) Salbutamol nebulisation

d) Magnesium sulphate

Question 28:

Which of the following is not seen in tumor lysis syndrome?

a) Hyperphosphatemia

b) Hyperuricemia

c) Hypercalcemia

d) Hyperkalemia

Question 29:

Which of the following is a negative acute phase reactant?

a) Albumin

b) Haptoglobin

c) Ferritin

d) C reactive protein

Question 30:

A 3-month-old male child was brought with a history of fever and respiratory distress. Suspecting pneumonia, a chest X-ray was ordered. Which of the following is the most likely causative organism?



a) Klebsiella pneumoniae

b) Streptococcus pneumoniae

c) Mycoplasma pneumoniae

d) Staphylococcus aureus

Question 31:

What does the red cell distribution width deal with?

a) Hypochromia

b) Anisocytosis

c) Poikilocytosis

d) Anisochromia

Question 32:

A 25-year-old woman came to the OPD one year after the birth of her last child. She had been treated for iron deficiency anemia while pregnant. On examination, she is pale and investigations show a Hb of 5g/dl and a reticulocyte count of 9%. What is her corrected reticulocyte count? (Take normal Hb as 15g/dl)

a) 6

b) 4.5

c) 3

d) 1

Question 33:

A 44-year-old man is referred to your center for management of severe hyperglycemia. During examination, you note that the patient has enlarged feet, coarse facial features and prognathism. You suspect acromegaly. Which of the following tests can you use as a screening method for this condition?

a) Oral glucose challenge test with GH level

b) Insulin-like growth factor-1 binding protein-3 (IGFBP-3) level

c) Insulin like growth factor-1 levels

d) Random growth hormone levels

Question 34:

What is not a feature of myasthenia gravis among the following?

a) Muscle fatigue

b) Absent deep tendon reflexes

c) Normal pupillary reflex

d) Ptosis

Question 35:

Regarding the procedure done with the needle shown below, which of the following is false?

a) To be done in lateral recumbent position

b) Breath holding is not necessary

c) The bevel should face upwards while inserting the needle

d) Coagulopathy is not a contraindication

Question 36:

Pancreatic cancer has the highest association with _____.

a) FAMMM syndrome

b) FAP

c) Hereditary pancreatitis

d) Peutz-Jeghers syndrome

Question 37:

The flow-volume curve after spirometry in patient 1 is given below. The most likely diagnosis in this patient is:



a) Endobronchial neoplasm

b) Bronchial Asthma

c) Myasthenia gravis

d) Emphysema

Question 38:

A 45-year-old female patient has NYHA class III heart failure with dyspnea, pedal edema and K+ levels of 5.5 mg% and creatinine 1.1 mg%. Which of the following drugs can be used in this patient?

a) Digoxin

b) Spironolactone

c) Enalapril

d) Furosemide

Question 39:

Which of the following features of stroke need not be treated?

a) Dysphagia

b) Spasticity

c) Fever

d) Neglect

Question 40:

A 50-year-old man is brought to the emergency following an episode of syncope during his morning walk. His blood pressure is 80/60 mm Hg. He has a history of myocardial infarction 5 yrs back. His present ECG is given below. Which is the most appropriate treatment for the patient?



a) Lignocaine

b) Diazoxine

c) DC cardioversion

d) Lidocaine

Question 41:

A patient came with complaints of high-grade fever and altered sensorium. He was diagnosed to be suffering from meningococcal meningitis. Which of the following is the most appropriate empirical treatment option?

a) Penicillin G

b) Ceftriaxone

c) Cefotetan

d) Cefoxitin

Question 42:

A 60-year-old man post-MI had the following lab values. Which of the following must be his treatment plan?

a) Rosuvastatin + fenofibrate

b) Atorvastatin 80 mg

c) Fenofibrate

d) Rosuvastatin 10 mg

Question 43:

A 40-year-old smoker who is a known case of hypertension was on enalapril and hydrochlorothiazide. He had an episode of hemoptysis and on evaluation he was found to have bronchogenic carcinoma with brain metastasis. His lab values were Na -124 mg/dl, creatinine – 2.8 mg%, blood sugar 112 mg/dl, blood urea - 24 mg/dl, serum osmolality – 255mosm, urine osmolality-350mosm, 24 hour urinary sodium 110 and BP-150/90 mmHg. Which of the following is the most probable diagnosis of the low sodium values in him

a) SIADH

b) Diuretic induced hyponatremia

c) Cerebral salt wasting syndrome

d) Pseudohyponatremia

Question 44:

Non invasive test to detect hepatic fibrosis include:

a) Aspartate aminotransferase/Platelet ratio

b) FibroTest

c) Forns index

d) Serum laminin and serum hyaluronidase

Answer Key

Question No.	Correct Option
1	b
2	a
3	b
4	a
5	a
6	d
7	c
8	a
9	d
10	b
11	d
12	c
13	d
14	b
15	a
16	c
17	a
18	b
19	a
20	b
21	d
22	b
23	d
24	d
25	a
26	a
27	d
28	c
29	a
30	d
31	b
32	c

33	c
34	b
35	d
36	d
37	c
38	d
39	d
40	c
41	b
42	b
43	a
44	b

Detailed Explanations

Solution to Question 1:

The clinical scenario describes a patient of APL (acute promyelocytic leukemia) developing fever, tachypnea, and pulmonary infiltrates while on medical treatment. This is suggestive of APL differentiation syndrome, an adverse effect of all-trans-retinoic acid (ATRA). The management involves glucocorticoids like dexamethasone.

Along with glucocorticoids, the other management of APL differentiation syndrome includes chemotherapy for cytoreduction and supportive measures. In severe cases, temporary discontinuation of ATRA is necessary.

ATRA is an oral drug that induces the differentiation of leukemic cells bearing the t(15;17) chromosomal translocation. While ATRA reduces the incidence of DIC associated with cytarabine and daunorubicin used earlier in APL, it can cause APL differentiation syndrome due to the adhesion of differentiated neoplastic cells to the pulmonary vasculature endothelium.

In low-risk APL patients with a low leukocyte count at presentation, a combination of ATRA and arsenic trioxide (ATO) has been found to be superior and is the current standard of care.

Solution to Question 2:

The current definition of fever of unknown origin (FUO) does not include undiagnosed after one week of investigation. The criterion "diagnosis uncertain, after 1 week of evaluation" which was previously a part of the definition has been removed.

FUO is currently defined as:

• Fever ≥38.3°C (≥101°F) on at least two occasions

• Duration of ≥3 weeks

• No known immunocompromised state

• Diagnosis is uncertain after thorough history-taking, physical examination, and obligatory investigations.

Inflammation of Unknown Origin (IUO) is closely related to FUO. It has the same definition, except for the body temperature criterion. IUO is characterized by elevated inflammatory parameters (such as CRP or ESR) on multiple occasions for at least three weeks in an immunocompetent patient with normal body temperature. The causes and workups for IUO are similar to those for FUO.

Solution to Question 3:

The first-line agents used in the management of Burkholderia cepacia complex (BCC) infections are trimethoprim-sulfamethoxazole (cotrimoxazole), meropenem, minocycline, and ceftazidime (3rd generation cephalosporin).

Other options:

Option A: Burkholderia species are intrinsically resistant to aminoglycosides and polymyxins.

Option C: Tigecycline is less effective in the case of Burkholderia infections.

Option D: Carbapenems and 3rd generation cephalosporins have the same mechanism of action and compete for the same binding site. Hence, it is not a rational drug combination.

BCC is a group of bacteria known for causing severe respiratory distress and septicemia, particularly in patients with cystic fibrosis (CF). BCC organisms are also found in ICU patients and individuals with chronic granulomatous disease, leading to lung disease. BCC colonization in CF patients prior to lung transplantation increases the risk of death during the early transplant period.

Solution to Question 4:

The given image showing localized overactivity (increased uptake) in the left lobe of the thyroid i.e., hot nodule, with suppression of the remainder of the gland, is suggestive of a hypersecreting adenoma.

Other options:

Option B: The likelihood of malignancies like papillary Carcinoma thyroid is low (5%) in the case of a hot nodule.

Option C: Graves disease shows diffusely increased uptake on a thyroid scan.

Option D: Lateral aberrant thyroid presents with swelling of cervical lymph nodes and not thyroid. It is due to skip metastasis from papillary thyroid carcinoma to the ipsilateral cervical lymph nodes.

The thyroid scan appearance in different conditions is shown in the image below:



Diagnostic isotope studies, such as 99mTc-pertechnetate and iodine-123, are becoming less common in thyroid imaging because some warm swellings (5%) which are actually malignant could be mistaken as benign. Isotope scanning is most valuable when assessing a hyperthyroid patient presenting with a nodule or nodularity in the thyroid. It helps to identify overactivity in the gland, which can differentiate between a toxic nodule that suppresses the rest of the gland and a toxic multinodular goitre with multiple areas of increased uptake.

Solution to Question 5:

The structure marked in this transverse section of the brain at the level of superior colliculus is substantia nigra. Degeneration of substantia nigra is seen in Parkinson's disease, also known as paralysis agitans.

Substantia nigra is pigmented due to the presence of neuromelanin granules, which help in the synthesis of dopamine. In Parkinson's disease, there is an absence/reduced pigmentation of the substantia nigra.

Bradykinesia, rigidity, and resting tremor are classic features of Parkinson's disease. Because of motor disability, it is also referred to as paralysis agitans.

Other options:

Option B: Depression is often a comorbid finding with Parkinson's disease, but is not due to damage to the substantia nigra.

Option C: Alzheimer’s disease is the most common cause of dementia. It is due to atrophy of the entorhinal cortex, located in the medial temporal lobe.

Option D: Huntington’s chorea is an autosomal dominant disorder of movement and cognition primarily involving the striatum (caudate nucleus and putamen).



Solution to Question 6:

Among the options, systolic blood pressure less than 100 mm Hg (Option D) is not a component of CURB-65. CURB65 includes blood pressure, systolic ≤90 mmHg or diastolic ≤60 mmHg.

CURB-65 criteria is used to assess the severity of illness in a case community-acquired pneumonia. It includes confusion (C), urea >7 mmol/L (U), respiratory rate ≥30/min (R), blood pressure systolic ≤90 mmHg or diastolic ≤60 mmHg (B) and an age of ≥65 years.

Pneumonia severity index (PSI) is another index to assess the severity of illness and the risk of mortality in the case of pneumonia. It includes 20 variables and depending on the score, patients are assigned into one of the 5 classes.

Class, I, II, III - at low risk of death and can be treated on an outpatient basis

Class IV & V require hospitalization

Solution to Question 7:

The clinical scenario describing decrease in movement of right leg (neurological symptom) in a young female with history of episodic attacks of abdominal pain and aggressive behaviour (mental symptom) is consistent with the diagnosis of acute intermittent prophyria.

Conversion disorder (Option D) presents commonly in young women as acute, temporary loss of sensory or motor function preceded by psychological issues. Recurrent episodic attacks of neurovisceral symptoms make the diagnosis of conversion disorder unlikely.

Acute intermittent porphyria (AIP) is a hepatic porphyria inherited in an autosomal dominant pattern. It occurs due to a deficiency of porphobilinogen deaminase/hydroxymethylbilane synthase/uroporphyrinogen I synthase enzyme in heme synthesis. AIP is generally latent before puberty and is more common in women.

AIP presents with neurovisceral symptoms and no cutaneous manifestations are seen. Abdominal pain is the most common symptom. Nausea, vomiting, constipation, muscle weakness, sensory loss, dysuria, and urinary retention are seen. Tachycardia, hypertension, restlessness, tremors, and excess sweating are due to sympathetic overactivity. Peripheral neuropathy predominantly affecting motor neurons is due to axonal degeneration. During acute episodes, individuals may experience mental manifestations such as anxiety, insomnia, depression, hallucinations, and paranoia. Seizures may result from either neurological effects or hyponatremia.

Diagnosis can be made with the appearance of classic symptomology after exposure to an underlying precipitating factor. Plasma and urine ALA and PBG levels are significantly raised during an acute attack and remain high even after symptoms resolve.

Intravenous hematin as first-line treatment for acute attacks. Narcotic analgesia may be given for abdominal pain. Carbohydrate loading with intravenous glucose may be tried in milder attacks. Orthoptic liver transplant is curative in patients refractory to hemin therapy.

Givosiran is a recently approved hepatic-targeted RNA interference (RNAi) therapy for acute hepatic porphyrias.

Solution to Question 8:

The given scenario describes a patient with ascending paralysis. Among the given options, Serum potassium (Option A) is tested as disturbances in serum potassium levels (hypokalemia or hyperkalemia) present with ascending muscle weakness.

Option B: Hypomagnesemia can cause tetany. Hypermagnesemia can cause generalized weakness and paralysis. However, serum potassium is a better answer as disturbances in potassium levels are more likely to cause paralysis.

Option C: Alteration in serum sodium levels does not cause muscular paralysis.

Option D : Hypocalcemia presents with paresthesia, tetany, seizures and prolonged QT interval. Hypercalcemia presents with other clinical features like fatigue, depression, mental confusion, anorexia, nausea, vomiting, constipation, reversible renal tubular defects, increased urine output, short QT interval in the electrocardiogram, and, in some patients, cardiac arrhythmias.

Solution to Question 9:

An elevated TSH value with low free T4 is suggestive of primary hypothyroidism (Option D).

Option A: In subclinical hypothyroidism, free T4 levels remain normal despite elevated TSH.

Option B: TSH is low in hyperthyroidism.

Option C: The diagnosis of secondary hypothyroidism is made along with a deficiency of other pituitary hormones. TSH levels may be low, normal, or slightly increased. The elevation in TSH levels can be attributed to the secretion of immunoreactive but biologically inactive forms of TSH. To confirm the diagnosis, it is essential to detect a low level of unbound T4.

The diagnostic approach to hypothyroidism is given in the flowchart below:



Solution to Question 10:

The elevated levels of triglycerides (TG) and very low-density lipoproteins (VLDL) in diabetes mellitus (DM) are due to decreased activity of lipoprotein lipase (LpL) and increased activity of hormone-sensitive lipase (HSL).

Insulin normally has an inhibitory action on HSL and induces LpL on capillary walls. There is an absolute or relative deficiency of insulin in DM. In DM, the activity of LpL is decreased and HSL is increased.

The increased activity of HSL results in increased mobilisation of fat stores towards lipolysis releasing fatty acids. These fatty acids are partly cleared by the liver, resulting in increased synthesis of VLDL and TG, resulting in dyslipidemia. They can accumulate in the liver, resulting in steatosis (non-alcoholic fatty liver disease).

The reduced activity of LpL prevents the clearing of TG and VLDL in the bloodstream and their levels remain elevated.



Solution to Question 11:

The window period for thrombolysis in case of stroke is 4½ hours.

Intravenous Recombinant Tissue Plasminogen Activator (IV rtPA) 0.9 mg/kg upto a maximum of 90 mg, 10% as bolus, the remainder as infusion over 60 min is given. Tenecteplase 0.25 mg/kg IV bolus over 5 s is also used.

Blood pressure should be monitored frequently. No other antithrombotic therapy should be administered within a 24-hour period. In the case of a deterioration in neurological condition or uncontrolled high blood pressure, the infusion should be stopped, cryoprecipitate should be administered, and urgent imaging of the brain should be done. It is recommended to refrain from urethral catheterization for a minimum of 2 hours.

Solution to Question 12:

The clinical scenario describes a 53 year old patient admitted in view of a cerebrovascular accident developing neurologic deterioration (drowsy with minimal response) on the second day, and CT brain showing significant edema of MCA territory with midline shift. The next line of management is decompressive surgery.

Cerebral edema is a potential complication in 5-10% of patients with strokes, leading to obtundation and brain herniation. The severity of edema is highest on the second or third day after the stroke and can last for around 10 days.

Medical management may not be successful in case of significant edema or infarcts involving the cerebellum.

Decompressive surgery is recommended for patients who are less than 60 years old and experience deterioration ≤ 48 hours, despite receiving medical therapy. Hypodensity on head CT within the first 6 hours of stroke onset, involving ≥ one-third of the MCA territory, early midline

shift, and a diffusion-weighted imaging volume on MRI within 6 hours ≥ 80 mL are predictors of severe cerebral edema.

Solution to Question 13:

Loud S1 is heard in mild mitral stenosis (MS).

The intensity of S1 depends on the distance over which the anterior leaflet of the mitral valve moves, leaflet mobility, LV contractility, and PR interval. It is independent of the severity of MS.

Loud S1 is seen in early rheumatic MS, short PR interval, and hyperdynamic circulatory states.

S1 becomes soft in case of calcified leaflets, long PR interval, associated aortic regurgitation and LV dysfunction.

Solution to Question 14:

The given ECG shows a premature ventricular complex (PVC) that occurs every second beat (ventricular bigeminy).

PVCs exhibit premature timing by arising before the anticipated occurrence of the subsequent regular heartbeat. They display an aberrant QRS-T appearance, where the QRS complex appears unusually wide (0.12 sec or longer), and the T wave and QRS complex point in opposite orientations (discordant).

Frequent PVCs can manifest in different patterns. When two PVCs occur consecutively, they are termed a pair or couplet.

If three or more PVCs occur in succession (triplet), it is classified as ventricular tachycardia (VT). The repetitive pattern of one normal beat followed by one PVC is known as ventricular bigeminy.

When the repeating sequence involves two normal beats followed by a PVC, it is called ventricular trigeminy. Similarly, if three normal beats are followed by a PVC, it is referred to as ventricular quadrigeminy.

The ECG below shows ventricular bigeminy where every normal QRS (green underlines) is followed by a PVC (red arrows):



Solution to Question 15:

The commonly used method to determine the length of a nasogastric tube is the NEX method: the tip of the nose to the ear to the xiphisternum.

The approximate length is 120-125 cm. The markings indicate the traversing of the tube through various parts of the GIT:

• 40 cm: Cardia of the stomach

• 50 cm: Body of the stomach

• 60 cm: Pylorus

• 70 cm: Duodenum

A nasogastric tube or Ryle's tube is inserted in the sitting position with the neck slightly flexed. The tube is then gently inserted through the nostril and advanced until it reaches 10 cm. The patient is encouraged to swallow, allowing the tube to move down the esophagus with successive swallows until it reaches the measured distance.

Subjective confirmation involves insufflating air through the proximal opening of the NG tube and listening for the sound of air in the left upper quadrant where the stomach is located. Objective confirmation can be obtained through pH measurement, chest or abdominal radiographs, ultrasound, or calorimetric capnometry.

While widely practiced, blind NG tube insertion can lead to complications such as coiling of the tube in the nasopharynx, incorrect placement in the bronchial passage, or even esophageal perforation.

Solution to Question 16:

Among the given options, canagliflozin (Option C) is the most suitable treatment. It is an oral drug belonging to the class SGLT-2 inhibitors and causes weight loss. Although it increases the risk of urogenital infections, it has no risk of pancreatitis or bladder cancer.

Option A: Sitagliptin, a DPP-IV inhibitor, is associated with the risk of acute pancreatitis.

Option B: Pioglitazone, a thiazolidinedione, can increase the risk of bladder cancer and can cause weight gain and congestive cardiac failure.

Option D: Liraglutide is an injectable GLP-1 receptor agonist. Pancreatitis is a serious adverse effect of this drug. Hence, it is contraindicated in individuals with a history of pancreatitis and should be permanently discontinued if pancreatitis develops.

Solution to Question 17:

A significant difference in blood pressure measured between two arms (>10 mm Hg) may be seen in Takayasu arteritis (subclavian artery disease), supravalvular aortic stenosis, aortic coarctation, and aortic dissection (all four options).

An early diastolic murmur in the right second intercostal space (aortic area) is suggestive of aortic regurgitation. Supravalvular aortic stenosis (Option A), which involves the part of the aorta distal to the aortic valve, is unlikely to cause aortic regurgitation and hence the best answer.

Features	Supravalvular AS	Takayasu arteritis	Aortic dissection	Coarctation of aorta
Aortic regurgitation	Unlikely	Present in 25% of cases	Present in acute situations	Present
BP difference between arms	Present	Present	Present	May be present

So the single best answer would be Supravalvular AS > Coarctation of the aorta.

Solution to Question 18:

The clinical scenario of a young patient developing paresthesia and circumoral numbness and ECG showing ST-T changes during blood donation for platelet apheresis is consistent with hypocalcemia.

Hypocalcemia in this case is due to citrate which is used as an anticoagulant in apheresis machine. Citrate is a calcium chelator and bind to the calcium in the donor's circulation.

During a physical examination, Chvostek's sign (twitching of the muscles around the mouth when the facial nerve is gently tapped) and Trousseau's sign (carpal spasm induced by inflating a blood pressure cuff to 20 mmHg above the patient's systolic blood pressure for 3 minutes) may be seen. Severe hypocalcemia can lead to various complications including seizures, spasms in the hands and feet (carpopedal spasm), bronchospasm, laryngospasm, and QT prolongation.

For acute and symptomatic hypocalcemia, the initial treatment involves intravenous administration of calcium gluconate, usually 10 mL of a 10% solution given over 5 minutes.

Solution to Question 19:

Metformin should be stopped in case of elevated creatinine levels (renal insufficiency) > 1.5 mg/dl in men and >1.4 mg/dl in women.

Metformin acts by reducing hepatic glucose production and improves peripheral glucose utilization. Lactic acidosis and Vitamin B12 deficiency are specific adverse effects of metformin. It is not used in patients with a glomerular filtration rate <30 mL/min (moderate renal insufficiency), acidosis, unstable congestive cardiac failure, liver disease, severe hypoxemia, and patients undergoing contrast studies.

Solution to Question 20:

In a patient with suspected superior vena cava syndrome (SVCS), the next step is to perform a CT thorax to confirm the diagnosis and determine the underlying cause. CT reveals reduced or absent opacification of central venous structures along with noticeable collateral venous circulation.

The diagnosis of SVCS is primarily clinical, but imaging can aid the diagnosis. The chest radiographic finding commonly observed in SVCS is the widening of the superior mediastinum. MRI can be used to diagnose SVC obstruction with 100% sensitivity and specificity. Invasive procedures like bronchoscopy or biopsy may be performed if needed.

SVCS is caused by the obstruction of the superior vena cava, resulting in impaired blood flow from the head, neck, and upper extremities. Lung cancer is the most common cause of malignant SVCS cases. In young adults, malignant lymphoma followed by mediastinal germ cell tumors are the leading causes of SVCS. The use of intravascular devices such as central venous catheters has led to an increase in benign causes of SVCS.

Patients with SVCS present with swelling of the neck and face, dyspnea, cough, hoarseness, headaches, nasal congestion, and pain. Bending forward or lying down may aggravate the symptoms. Physical examination may reveal dilated neck veins, collateral veins on the chest wall, cyanosis, and edema of the face, arms, and chest.

Cerebral and laryngeal edema are rare but serious complications associated with SVCS, and they require urgent evaluation.

Treatment of SVCS depends on the underlying cause. Radiation therapy is the primary treatment for SVCS caused by non-small-cell lung cancer and metastatic solid tumors. Chemotherapy is effective for small-cell lung cancer, lymphoma, and germ cell tumors. Intravascular stenting is often used to provide rapid relief of symptoms.

Solution to Question 21:

The serum ascites albumin gradient (SAAG) in ascites due to TB peritonitis is < 1.1 g/dL as it is unrelated to portal hypertension.

SAAG helps differentiate ascites caused by portal hypertension from nonportal hypertensive ascites. A SAAG ≥1.1 g/dL indicates portal hypertension and suggests causes such as cirrhosis, cardiac ascites, hepatic vein thrombosis, sinusoidal obstruction syndrome, or liver metastases. A SAAG <1.1 g/dL suggests other causes like tuberculous peritonitis, peritoneal carcinomatosis, or pancreatic ascites. For high-SAAG ascites, the ascitic protein level provides additional information. A protein level of ≥2.5 g/dL suggests normal hepatic sinusoids, while <2.5 g/dL indicates damaged and scarred sinusoids. Pro-brain-type natriuretic peptide (BNP) levels may help identify heart failure as the cause of high-SAAG ascites.

Additional tests may be conducted in specific clinical situations, such as measuring ascitic glucose and lactate dehydrogenase (LDH) levels for suspected secondary peritonitis, ascitic amylase levels for suspected pancreatic ascites, and cytology for peritoneal carcinomatosis.

Tuberculous peritonitis may be associated with ascitic fluid lymphocytosis. The sensitivity of the smear for acid-fast bacilli and culture is low. Elevated ascitic adenosine deaminase > 30–45 U/L is highly sensitive.

Solution to Question 22:

In a patient on anti-tubercular therapy (ATT), the development of a tingling sensation is suggestive of isoniazid (INH) related peripheral neuropathy. In adults, pyridoxine (100 mg/day) is given until symptoms subside.

Alcoholics, malnourished persons, pregnant and lactating women, and patients with chronic renal failure, diabetes, and HIV infection are at higher risk of peripheral neuropathy. Patients not responding to pyridoxine may be treated with amitriptyline. Mothers receiving INH (10mg/day), their infants (5 mg/day), and all children receiving INH (10mg/day) should receive pyridoxine supplementation. Adults (50 mg/day) and children > 1 year (25 mg/day) on isoniazid preventive therapy are also supplemented with pyridoxine.

Drug-induced liver injury is another adverse effect of INH. Liver enzymes more than three times the upper limit with symptomatic jaundice or liver enzymes more than five times the upper limit in asymptomatic patients may warrant cessation of INH.

Ethionamide, a second-line anti-tubercular drug, also causes neurological adverse effects and may require pyridoxine supplementation.

Solution to Question 23:

The given symptoms cluster of fever, vomiting, and neck stiffness is suggestive of meningitis. A CSF analysis showing increased opening pressure, lymphocytic pleocytosis, mildly increased protein, and normal glucose is suggestive of viral meningitis. Among the given options, Coxsackie (Option D) is the only viral cause of meningitis.

Other options:

Option A: Tuberculous meningitis shows lymphocytic pleocytosis with low glucose.

Option B: Bacterial meningitis shows neutrophilic pleocytosis with low glucose.

Option C: Fungal meningitis shows lymphocytic/neutrophilic pleocytosis with low glucose.

Solution to Question 24:

Among the given options, salicylate poisoning (Option D) causes high anion gap metabolic acidosis. Proximal renal tubular acidosis, pancreatitis, and severe diarrhea are causes of normal anion gap acidosis.

Metabolic acidosis can be caused by increased production of endogenous acids, loss of bicarbonate, or impaired acid excretion by the kidneys. Blood pH decreases, leading to increased ventilation, particularly in tidal volume (Kussmaul respiration). It can be categorized into two main types: high-anion gap and non-anion gap acidosis.

Solution to Question 25:

The drug of choice for scrub typhus is doxycycline.

To treat scrub typhus, oral doxycycline 100 mg is given twice daily for 7–15 days. In the case of pregnant women, oral azithromycin 500 mg for 3 days may be used as doxycycline is contraindicated in pregnancy.

Scrub typhus is a rickettsial infection caused by Orientia tsutsugamushi. The vector is a trombiculid mite. Humans are infected when they are bitten by the mite larvae known as chiggers. The incubation period is 1- 3 weeks. Patients develop a characteristic eschar at the site of the mite bite, along with regional lymphadenopathy and maculopapular rash. It can be associated with fever, headache, myalgia, meningoencephalitis, myocarditis, and pneumonia.

Serologic assays are used for diagnosis. Weil-Felix reaction is a heterophile agglutination reaction used for diagnosis, wherein the sera are tested for agglutinins to the O antigens of Proteus strains OX19, OX2, and OXK. OXK alone is agglutinated in scrub typhus.

Solution to Question 26:

The given clinical scenario and the electrocardiogram (ECG) showing narrow QRS complex regular tachycardia without P waves are suggestive of paroxysmal supraventricular tachycardia (PSVT). The next step in the management of PSVT in a hemodynamically stable (systolic blood pressure >90 mm Hg) patient is vagotonic maneuvres such as carotid sinus massage or Valsalva maneuver.

Carotid sinus massage can be considered in the absence of carotid bruits and no history of stroke. If vagal maneuvers fail or cannot be performed, intravenous adenosine can be used.

Non-DHP calcium channel blockers and beta blockers are second-line pharmacological agents in the management of PSVT. They can cause hypotension before and after terminating the arrhythmia.

If the patient is hemodynamically unstable (systolic blood pressure <90 mmHg or mean arterial pressure <65 mm Hg), immediate QRS-synchronous direct current cardioversion is necessary.

The treatment algorithm for PSVT is described in the flowchart below:



Solution to Question 27:

Among the given options, magnesium sulphate has no role in the management of hyperkalemia.

Hyperkalemia is a medical emergency that requires urgent treatment. Even in the absence of ECG changes, patients with plasma K+ concentration ≥6.5 mEq/L should be aggressively managed.

Intravenous calcium gluconate (10 mL of 10% solution) is administered over 2-3 minutes with cardiac monitoring to protect the heart while other measures are taken to correct hyperkalemia. Calcium increases the threshold of action potential thereby reducing excitability.

Insulin-dextrose infusion is given to shift potassium into cells. The recommended dose is 10 units of regular insulin followed immediately by 50 mL of 50% dextrose. β2-agonists like salbutamol can also be used in conjunction with insulin.

Cation exchange resins, diuretics, and dialysis are used to eliminate potassium. Sodium polystyrene sulfonate is a cation exchange resin that exchanges sodium for potassium in the gastrointestinal tract, increasing fecal excretion of potassium. Newer intestinal potassium binders like patiromer and ZS-9 are preferred due to their lack of intestinal toxicity. Hemodialysis is the most effective method to reduce plasma potassium concentration.

In hypovolemic patients with oliguria, intravenous saline may be beneficial to improve distal delivery of sodium and enhancing renal potassium excretion. Loop and thiazide diuretics can be used in hypervolemic patients with sufficient renal function.

Solution to Question 28:

Hypocalcemia, not hypercalcemia, is seen in tumor lysis syndrome (TLS).

TLS is a potentially life-threatening condition characterized by the rapid destruction of a large number of tumor cells. The release of calcium phosphate from lysed cells contributes to hyperphosphatemia and subsequent reciprocal depression in serum calcium levels. The consumption of ionized calcium during the formation of calcium phosphate crystals leads to hypocalcemia.

TLS is commonly observed after initiation of chemotherapy in hematologic malignancies characterized by high cell turnover rates, such as Burkitt's lymphoma and acute lymphoblastic leukemia.

The destruction of neoplastic cells results in the release of nucleic acids, which are broken down into uric acid, leading to hyperuricemia. The acidic environment in the renal tubules can cause uric acid to precipitate, leading to acute renal failure. Potassium is released in large quantities during tumor cell lysis leading to hyperkalemia.

TLS is diagnosed using the Cairo-Bishop criteria which includes both laboratory and clinical criteria.

Prevention is the mainstay of the management of TLS, which involves includes allopurinol, which inhibits the production of uric acid, and aggressive hydration to promote urine flow. Urinary alkalization is not recommended. Febuxostat, a selective xanthine oxidase inhibitor, can be used as an alternative.

In severe cases or when renal failure is present, rasburicase, a recombinant urate oxidase, is used for treatment. Rasburicase rapidly lowers uric acid levels by converting it into a highly soluble compound called allantoin. Renal replacement therapy is often required and should be considered in case of acute renal failure.

Solution to Question 29:

Albumin is a negative acute phase reactant while haptoglobin, ferritin and C-reactive protein are positive acute phase reactants.

Acute phase reactants are classified as positive and negative based on whether their levels decrease or increase in inflammatory states.

Solution to Question 30:

The most likely causative agent of pneumonia in an infant with X-ray showing pneumatoceles (air-filled cysts within lung parenchyma) is Staphylococcus aureus.

In newborns and infants, S. aureus is a significant cause of severe respiratory tract infections, presenting with dyspnea, fever, and respiratory failure. Community-acquired infections are often

preceded by viral infections, most commonly influenza. Among adults, nosocomial S. aureus pulmonary infections are frequently observed in intubated patients within intensive care units.

S. aureus often causes necrotizing pneumonitis that may be associated with complications such as pneumothorax and empyema.

Pneumatocoele can also be caused by Streptococcus pneumoniae, Haemophilus influenzae, E. coli, group A streptococci, Klebsiella pneumoniae, etc. However, Staphylococcus aureus is most commonly implicated.

Other options:

Option A: Klebsiella pneumoniae causes lobar pneumonia in alcoholics and diabetics. Patients present with fever, dyspnea, and a productive cough with red currant jelly sputum. Classic chest X-ray findings include upper lobe consolidation, classically on the right side, cavitation, and bulging fissure sign.

The following image shows bulging fissure sign:

Option B: Streptococcus pneumoniae causes lobar pneumonia. Patients with pneumococcal pneumonia present with fever, dyspnea, and productive cough. On examination, the involved lobes are dull to percussion, bronchial breath sounds, and crackles are heard on auscultation.

The following image shows right lower lobe consolidation:

Option C: Mycoplasma pneumoniae causes atypical pneumonia. Patients present with dry cough, fever, chills, malaise, and upper respiratory tract symptoms. On auscultation, wheezes or rales are heard. Chest X-ray shows peribronchial pneumonia with prominent bronchial markings, interstitial infiltration, and atelectasis.

The below chest X-ray shows patchy consolidation at the hilum and fanning to the periphery:



Solution to Question 31:

Red cell distribution width (RDW) measures the range of red blood cell volumes and provides an assessment of their variability (anisocytosis).

The normal range of RDW is 11-14%. In the presence of uneven cell size (anisocytosis), the RDW increases to 15-18%.

RDW has clinical utility in two main scenarios:

• In cases of microcytic anemia (low MCV), it helps differentiate between iron deficiency and thalassemia. Thalassemia is characterized by uniformly small red blood cells with a normal RDW, while iron deficiency leads to greater size variability and an increased RDW.

• A high RDW can indicate dimorphic anemia caused by chronic atrophic gastritis. Vitamin B12 malabsorption results in macrocytic anemia and blood loss leads to iron deficiency (microcytic anemia).

Solution to Question 32:

The corrected reticulocyte count (CRC) of this patient is 3.

CRC = Reticulocyte % x (Actual Hb / Normal Hb)

Plugging in the values given in the question,

CRC = 9% x 5/15 = 3%

Alternatively, CRC can be calculated using hematocrit (PCV) as follows:

CRC = Reticulocyte % x (Actual PCV / Normal PCV)

For these calculations, take normal Hb as 15g/dL and normal PCV as 45%.

The patient's hemoglobin level (5 g/dL) is significantly lower than the normal hemoglobin level (15 g/dL). To obtain an accurate assessment of reticulocyte production relative to the reduced number of circulating red blood cells, a correction factor has to be applied. CRC accounts for the anemia and provides a more precise estimate of the reticulocyte count.

When polychromatophilic macrocytes (prematurely released reticulocytes/shift cells) are observed in the peripheral blood smear, a second correction is applied to account for the longer life of reticulocytes to yield reticulocyte production index (RPI).

In the classification of anemia, the patient's reticulocyte count is compared to the expected reticulocyte response. Reticulocytes can be identified by staining their ribosomal RNA with a supravital dye, resulting in the formation of blue or black punctate spots. They can be quantified manually or by using fluorescent dyes. The presence of reticulocytes indicates active red cell production.

The premature release of reticulocytes is typically associated with increased EPO stimulation. In hemolytic anemia, red cell production can increase six to seven times. If the RPI < 2 in a patient with anemia, a defect in erythroid marrow proliferation or maturation has to be suspected.

Solution to Question 33:

The best screening test for acromegaly is the measurement of serum insulin-like growth factor-1 (IGF-1) level.

Other options:

Options A and B: Oral glucose challenge test with growth hormone (GH) levels is used for further confirmation of the diagnosis. Increased insulin-like growth factor-1 binding protein-3 (IGFBP-3) levels can also be seen in patients with acromegaly. But they are not typically used as initial screening methods.

Option D: IGF-1 levels are more reliable than random GH levels because GH secretion is pulsatile. IGF-1 levels reflect long-term exposure to GH and provide a more stable indicator of excessive GH production.

Acromegaly is characterized by excessive GH secretion, usually caused by a somatotrope adenoma in the pituitary gland. It can also be due to ectopic GH secretion or hypothalamic tumors.

The clinical presentation of acromegaly includes frontal bossing, increased hand and foot size, coarse facial features, mandibular prognathism, widened space between lower incisors and other manifestations such as soft tissue swelling, deepening voice, carpal tunnel syndrome, arthropathy, and acanthosis nigricans.

It can also lead to cardiovascular complications (cardiomyopathy, left ventricular hypertrophy, and hypertension), obstructive sleep apnea, diabetes mellitus, an increased risk of colon polyps, and colonic malignancy.

The primary treatment for acromegaly is surgical resection of GH-secreting adenomas. Somatostatin receptor ligands like octreotide and lanreotide, which suppress GH and IGF-1 levels, are used as adjuvant treatment or when surgery is not possible or unsuccessful. Radiation therapy may be considered for patients who cannot tolerate or do not respond to medical therapy. Other treatment options include GH receptor antagonists like pegvisomant and dopamine agonists like cabergoline.

Solution to Question 34:

Absent deep tendon reflex is not a feature of myasthenia gravis (MG).

MG usually occurs in young females. Patients usually present with weakness and fatiguability of the muscles, which worsens on repeated use or late in the day and improves on rest. The cranial muscles, particularly the eyelids and extraocular muscles are involved early in the course of MG. Diplopia and ptosis are common initial complaints. When MG becomes generalized, the proximal muscles of the limbs are affected first. Despite the muscle weakness, the deep tendon reflexes remain normal. Snarling expression on smiling (facial weakness), nasal timbre of speech (weakness of palate and tongue), and difficulty swallowing may be seen. Bulbar weakness and respiratory depression are seen more frequently in MuSK antibody–positive MG.

MG is an autoimmune disorder most commonly caused by antibodies against the acetylcholine receptor (AChR). These antibodies cause damage to the postsynaptic muscle membrane and increased endocytosis of the AChR. This autoimmune response against the AChR seems to be mediated by the thymus. The myoid (muscle-like) cells present in the thymus express AChRs on their surface and are postulated to provide the autoantigen which triggers this autoimmune reaction. Thymic hyperplasia is seen in 65% of the patients, and thymomas are seen in 10% of the patients.

The tests which are used to confirm the diagnosis of MG include:

• Anti-AChR antibody immunoassay: 85% of myasthenic patients have detectable anti-AChR antibodies, but only 50% of patients with weakness limited to ocular muscles. The presence of anti-AChR antibodies is highly suggestive of MG, but a negative test does not exclude the disease. Antibodies to MuSK and LRP4 are found in some patients

• Recurrent nerve stimulation test: MG shows a decremental response > 10%, while Lambert-Eaton myasthenic syndrome (LEMS) shows an incremental response

• Block and jitter response on single-fiber electromyography

• Positive edrophonium test: improvement of ptosis on the administration of anticholinesterase such as edrophonium

• Positive ice-pack test: improvement of ptosis on the application of ice-pack on the affected eye

The first line of medical management in MG is anticholinesterase drugs. Pyridostigmine 30–60 mg three to four times daily is the most commonly used drug.

Patients are evaluated for associated thymic abnormalities using a CT chest. Thymectomy is considered in patients with thymoma and in generalized MG patients with anti-AChR antibodies.

Immunotherapy therapy can be initiated in patients with poor control on pyridostigmine after thymectomy. Glucocorticoids, cyclosporine, or tacrolimus are generally effective within 1-3 months. Azathioprine and mycophenolate mofetil may take several months to show beneficial effects and are used for the long-term management of MG. Rituximab has shown effectiveness in patients with MuSK antibodies. Intravenous eculizumab, which binds to terminal complement component 5, is also effective.

Myasthenic crisis is the life-threatening exacerbation of MG leading to respiratory failure. Treatment for myasthenic crisis should take place in intensive care units. It is important to rule out excessive anticholinesterase medication as a cause of deterioration by temporarily discontinuing these drugs. Plasmapheresis and intravenous immunoglobulin (IVIg) are beneficial in hastening recovery.

Solution to Question 35:

The given image shows a Quincke-type lumbar puncture needle. Coagulopathy is a contraindication to perform lumbar puncture (LP).

Contraindications for LP include:

• Patients receiving anticoagulant or antiplatelet medications and those with coagulation defects

• Possible raised intracranial pressure

• Suspected spinal epidural abscess

• Infection at the site of puncture

Before performing an LP, it is important to assess the patient's bleeding risk by checking their platelet count, international normalized ratio (INR), and partial thromboplastin time (PTT). LP is considered contraindicated if the platelet count is less than 20,000/µL, and some institutions recommend a platelet count of greater than 40,000/µL before proceeding with the procedure.

Proper positioning of the patient is crucial for an LP. The patient should lie on their side away from the examiner, with their knees bent and pulled up toward the abdomen (lateral decubitus). The neck should be gently bent forward, and the shoulders and pelvis should be aligned vertically. LP is typically performed at or below the L3-L4 interspace, which can be identified by palpating the spinous processes and using a line drawn between the posterior superior iliac crests as a guide.

Alternatively, the LP can be done with the patient seated upright, leaning forward to flex the back. This position is suitable for obese patients.

After sterile preparation of the insertion site and the injection of a local anesthetic, the LP needle is inserted midline between two spinous processes, with the bevel parallel to the dural fibers facing upwards. The needle is advanced until the subarachnoid space (SAS) is reached indicated by a loss of resistance. Once the SAS is reached, the opening pressure is measured, and CSF is allowed to drip into collection tubes for analysis.

A traumatic tap may cause a bloody tap, which can be differentiated from subarachnoid hemorrhage (SAH) by examining the supernatant after centrifugation. Prior to needle removal, the stylet is reinserted to prevent nerve entrapment and the development of a dural CSF leak. There is no need for prolonged recumbency after the procedure to prevent post-LP headache.

Neuroimaging is recommended before performing an LP in patients who have an altered level of consciousness, focal neurological deficits, new-onset seizures, or papilledema. This is because these patients are at a higher risk of developing cerebellar or tentorial herniation, a rare but potentially fatal complication. Imaging studies should also include the spine in patients with symptoms suggestive of spinal cord compression, such as back pain, leg weakness, urinary retention, or incontinence. In cases where acute meningitis is suspected and neuroimaging is necessary before the LP, it is recommended to administer antibiotics, preferably after blood culture, before conducting the imaging study.

The image below shows Quincke (dura cutting) and Whitacre (dura splitting) LP needles:

Solution to Question 36:

Pancreatic cancer has the highest association with Peutz–Jeghers syndrome.

The risk factors for pancreatic cancer include:

• Smoking: One to threefold increase in risk

• Chronic pancreatitis

• Occupational exposure

• Obesity: Up to threefold increase in risk

• New-onset diabetes associated with:

• Elderly patients

• Lower body mass index

• Weight loss

• No family history of diabetes

• Hereditary factors:

FAMMM: Familial atypical multiple mole melanoma

HNPCC: Hereditary nonpolyposis colorectal cancer

Syndrome	Germ-line mutation	Estimated increased risk (fold) of
Familial adenomatous polyposis	APC	4
HNPCC /Lynch syndrome	MLH1, MSH2, MSH6, PMS2	8
Hereditary breast and ovarian cancer	BRCA 2	10
Familial pancreatic cancer		18
FAMMM syndrome	p16/CDKN2A	20
Cystic fibrosis	CFTR	30
Hereditary pancreatitis	PRSS 1/SPINK1	53
Peutz–Jeghers syndrome	STKII (LKB1)	~132

Solution to Question 37:

The given flow-volume curve shows decreased lung volumes (total lung capacity, vital capacity, and residual volume) with normal shape, suggestive of a restrictive pattern. The curve is a miniature version of the normal curve with shift towards the right. It is seen in myasthenia gravis.

Myasthenia gravis presents with ventilatory restriction due to reduced respiratory muscle strength.

Asthma, emphysema, and endobronchial neoplasms present with obstructive flow patterns.



Solution to Question 38:

The clinical scenario describes a patient with symptomatic heart failure and K+ levels of 5.5 mg%. She requires a diuretic for symptom control. Furosemide can be given safely as it causes hypokalemia.

Other options:

Option A: Digoxin is not used as a first-line drug in the management of symptomatic heart failure.

Options B and C: Spironolactone (K+ sparing diuretic) and enalapril (ACE inhibitor) can cause hyperkalemia.

The first step in managing acute decompensated heart failure (ADHF) is addressing the factors that caused the decompensation. They include nonadherence to medications, dietary indiscretion, medication use that exacerbates heart failure, coronary ischemia, arrhythmias, valvular heart disease, infection, and pulmonary thromboembolism.

Volume management is crucial in ADHF. Intravenous loop diuretics are used to relieve congestion. The effectiveness of diuretic therapy can be assessed by changes in weight, improvement in symptoms, and clinical examination findings. Natriuretic peptide levels can also help assess the adequacy of therapy and predict risk.

Cardiorenal syndrome, characterized by the interplay between heart and kidney dysfunction, is a common complication of ADHF. Diuretic therapy should be carefully managed to avoid worsening renal function in patients with elevated right-sided filling pressures. In cases of severe low cardiac output, inotropic therapy or mechanical circulatory support may be necessary to preserve or improve renal function temporarily.

Vasodilators like nitroglycerin, sodium nitroprusside, and nesiritide (recombinant BNP) are used to reduce filling pressures and systemic vascular resistance.

Inotropic therapy with drugs like dopamine and milrinone is used to increase myocardial contractility and improve cardiac output in patients with low-output ADHF. Dopamine has

dose-dependent effects on different receptors, while milrinone is a phosphodiesterase-3 inhibitor.

Solution to Question 39:

Neglect is a feature of stroke that need not be treated. Although neglect indicates a poor prognosis, most patients recover without any treatment in 1-6 months.

Dysphagia(Option A) poses a high risk of aspiration and hence can't be left untreated. Spasticity(Option B) can be treated with baclofen, tizanidine. Untreated, it creates a psychosocial impact on the patient. Fever(Option C) is deleterious in stroke patients and must be treated.

Neglect can arise from injury to cortical-limbic-reticular network, that directs attention and integrates the localization and identification of a stimulus. Neglect is associated with poor scores and slower recovery, but severe neglect is rare beyond 6 months. Recovery is rapid in the first 2 weeks, regardless of the side of stroke. By 3 months, patients might have some residual visual neglect. The choice of an intervention depends on the mechanism of neglect, the patient can be treated for an underaroused right hemisphere which has difficulty processing sensory inputs.

Neurogenic dysphagia is an important cause of pulmonary infection in any patient with stroke, motor neuron disease, even transient dysphagia can lead to malnutrition, dehydration, aspiration pneumonia, and airway obstruction with asphyxia, it increases the risk of death and institutionalization. Early placement of nasogastric tube to feed patients after stroke has improved nutrition and reduced deaths.

Spasticity is seen in less than 20% of stroke patients. Patients with bilateral weakness from hypoxic injury are likely to have signs of spasticity. Exaggerated cutaneous and autonomic reflexes and involuntary spasms are disabling problems associated with spasticity. Physical modalities like slow stretching movements and daily passive exercises will reduce symptoms of spasticity and risk of contractures. Baclofen, tizanidine, and clonidine are used in reducing clonus and extensor spasms.

Solution to Question 40:

The clinical scenario describes a man with history of coronary artery disease developing syncope. The ECG shows wide QRS complex (> 0.12 s) tachycardia with a single QRS morphology suggestive of monomorphic ventricular tachycardia (VT). The most appropriate treatment of VT in a hemodynamically unstable (systolic blood pressure < 90 mm Hg) patient is DC cardioversion.

VT can be casued by structural heart disease, scar-related reentry, cardiomyopathies, or abnormalities in the conduction pathways. VT is said to be sustained if it lasts for at least 30 seconds or requires intervention for termination.

Rapid VT can lead to hypotension and syncope, especially in patients with ventricular dysfunction. Patients with normal cardiac function may tolerate sustained VT, presenting with palpitations. Monomorphic VT associated with structural heart disease can progress to ventricular fibrillation (VF) and sudden cardiac arrest.

QRS synchronous electrical cardioversion is performed for unstable patients. Intravenous amiodarone is the preferred drug for stable VT associated with heart disease.

After restoring sinus rhythm, further evaluation is necessary to determine the underlying heart disease. Treatment options include antiarrhythmic medications, catheter ablation, or implantable cardioverter-defibrillators (ICDs) for recurrent or high-risk VT. ICDs are commonly used for secondary prevention of sudden death in patients with sustained VT and structural heart disease.

Solution to Question 41:

Ceftriaxone or cefotaxime is the drug of choice for empirical treatment of meningococcal meningitis. However, Penicillin G remains the antibiotic of choice for susceptible strains.

Empirical antimicrobial therapy is initiated in patients with suspected bacterial meningitis before the results of CSF Gram stain and culture are known.

S. pneumoniae and N. meningitidis are the most common etiologic organisms of community-acquired bacterial meningitis. Due to the emergence of penicillin-resistant and cephalosporin-resistant S. pneumoniae, empirical therapy of community-acquired suspected bacterial meningitis should include a combination of:

• Dexamethasone

• Third- or fourth-generation cephalosporin (e.g., ceftriaxone, cefotaxime, or cefepime)

• Vancomycin

• Acyclovir - HSV encephalitis is the main differential

• Doxycycline - during tick season.

Ceftriaxone or cefotaxime provides better coverage for susceptible S. pneumoniae, H. influenzae and group B streptococci and good coverage for N. meningitidis. Cefepime is a broad-spectrum fourth-generation cephalosporin with activity similar to that of cefotaxime or ceftriaxone against S. pneumoniae and N. meningitidis with additional greater activity against Enterobacter species and Pseudomonas aeruginosa.

Solution to Question 42:

In this 60 yr old patient with post-myocardial infarction (Atherosclerotic cardiovascular disease patient), high-intensity statins should be given irrespective of cholesterol levels. So atorvastatin 80mg is the treatment plan in the given case scenario.

Examples of high-intensity statins include high-dose atorvastatin 40–80 mg/day and rosuvastatin 20–40 mg/day.

Note: Fenofibrate is not advocated unless TG is more than 500 mg/dl. It can be administered along with a moderate-intensity statin.



Solution to Question 43:

The clinical scenario of patient who is a known hypertensive, smoker and bronchogenic carcinoma with brain metastasis having low sodium, low serum osmolality and high urine osmolality, the most probable diagnosis is of SIADH (Syndrome of Inappropriate secretion of ADH).

From the given information, reduced serum osmolality rules out pseudohyponatremia(Option D) as such patients should be isotonic or have normal serum osmolality. The BP of the patient indicates there is no hypotension and patient is euvolemic, hence ruling out Cerebral Salt Wasting Syndrome(Option C) in which patient is hypovolemic and volume depleted. In thiazide induced SIADH(Option B), patients will be dehydrated and not euvolemic.

In this patient, lung cancer and thiazides may cause SIADH.

Bartter and Schwartz diagnostic criteria of SIADH

- Hyponatremia
- Clinically Euvolemic
- Serum osmolality less than 275 mOsm/Kg
- Urine Osmolality more than 100mOsm/Kg
- Urine sodium more than 40 mmol/L
- No use of diuretics

In SIADH, if the hyponatremia is mild and asymptomatic, restrict total water intake to 30 mL/kg/day less than urine output until the syndrome resolves spontaneously. If the hyponatremia is severe and symptomatic, the goal is to partially correct it by intravenous infusion of hypertonic (3%) saline or administration of an AVP antagonist such as tolvaptan or conivaptan. With hypertonic saline or vaptans, plasma osmolarity and sodium should be checked every 1–2 h, and water intake should be adjusted to keep osmolarity 270 mosm/L, at which point the treatment should be discontinued.

Solution to Question 44:

FibroTest is a sensitive test to detect hepatic fibrosis. It is a multiparameter blood test which incorporates haptoglobin, bilirubin, GGT, apolipoprotein A-I, and α2-macroglobulin.

Aspartate aminotransferase/Platelet ratio and Forns index are tests to estimate the risk of developing liver fibrosis.

Fibrotest has high positive and negative predictive values for diagnosing advanced fibrosis in patients with alcoholic liver disease, chronic hepatitis B, chronic hepatitis C, nonalcoholic fatty liver disease and psoriasis patients taking methotrexate.

Transient elastography (TE) known as FibroScan, and magnetic resonance elastography(MRE) are used in patients of liver disease. TE is a noninvasive procedure uses ultrasound waves to measure hepatic stiffness and is accurate for identifying advanced fibrosis in patients with chronic hepatitis C, primary biliary cholangitis, hemochromatosis and nonalcoholic fatty liver disease. MRE is superior to TE for staging liver fibrosis in patients with a variety of chronic liver diseases but is expensive.

Tests for estimating risk of developing hepatic fibrosis:

• AST/Platelet ratio

• Forns Index

• Fib-4

• Cirrhosis determinant score

Test for detecting hepatic fibrosis:

• FibroTest (also called Fibrosure)

Note: Serum laminin and serum hyaluronidase are non specific and is not preferred as a test to detect hepatic fibrosis.

Medicine AIIMS 2018

Question 1:

An 18-year-old girl with the diagnosis of acute promyelocytic leukemia was treated medically. She developed fever and tachypnea and a chest X-ray showed pulmonary infiltrates. What drug should she be given next?

a) Cytarabine

b) Dexamethasone

c) Doxorubicin

d) Methotrexate

Question 2:

A 50-year-old man was brought to the emergency in an unconscious state. He had a fever for the past two days and is a known case of severe COPD. His ECG is given below. What is the most likely diagnosis?



a) Ventricular tachycardia

b) Multifocal atrial tachycardia

c) Atrial fibrillation

d) Atrial tachycardia

Question 3:

Infertility in Kartagener's syndrome is due to which of the following?

a) Oligospermia

b) Asthenospermia

c) Undescended testes

d) Epididymis obstruction

Question 4:

Calculate the GCS Score in an intubated patient, with findings of eye movements to pain and abnormal flexion?

a) E2VTM4

b) E2VTM3

c) E2V1M3

d) E2VNTM3

Question 5:

Two brothers were arguing over a property dispute when the elder of the two complained of chest pain and collapsed and was later declared brought dead by the hospital. His family says he was previously healthy and there was no similar disease in the family members. What is the likely diagnosis?

a) Infective cardiomyopathy

b) Takotsubo cardiomyopathy

c) Dilated cardiomyopathy

d) Hypertrophic cardiomyopathy

Question 6:

Which of the following is used in the treatment of late cardiovascular syphilis?

a) Benzathine penicillin 2.4 million units weekly for three weeks

b) Benzathine penicillin 2.4 million units as single dose

c) Benzathine penicillin 12-24 million units for 21 days

d) Tetracycline 2g daily

Question 7:

Which of the following drugs is commonly used for treating community-acquired pneumonia in OPD?

a) Vancomycin

b) Ceftriaxone

c) Azithromycin

d) Streptomycin

Question 8:

Which of the following is not a component of Syndrome Z?

a) Obstructive sleep apnea

b) Blood pressure more than 130/85 mmHg

c) Fasting triglyceride more than150mg/dl

d) LDL more than 100mg/dl

Question 9:

A patient with a history of a backache for 10 days, has now presented with sudden onset difficulty in micturition and defaecation. There was no history of a cough or fever previously. What is the diagnosis?

a) Pott's Spine

b) Guillain Barre Syndrome

c) Cauda Equina Syndrome

d) Multiple Sclerosis

Question 10:

A 40-year-old male chronic smoker presents to the AIIMS OPD with fever, fatigue, yellow-colored urine, and clay-colored stools. For the past few days, he has developed an aversion to cigarette smoking. On examination icterus was present. What investigations

would you advise to rule out acute viral hepatitis? Liver function test results are given below:

a) HBsAg, IgM antiHBC, AntiHCV, AntiHEV

b) AntiHAV, HBsAg, IgM antiHBc, AntiHCV

c) HBsAg, IgM antiHBc, AntiHDV, AntiHCV, Anti HEV

d) AntiHAV, IgM antiHBc, AntiHCV, AntiHEV

Question 11:

A 22-year-old man presented with diarrhea and intolerance to dairy products. On investigation, he was found to have a lactase deficiency. Which of the following agents is least likely to cause symptoms of lactose intolerance?

a) Condensed Milk

b) Skimmed Milk

c) Yogurt

d) Ice Cream

Question 12:

What is the most likely causative organism of ventilator-associated pneumonia among the following?

a) Clostridium

b) Klebsiella

c) Acinetobacter

d) Hemophilus

Question 13:

The arterial blood gas findings of a patient are pH = 7.12, pCO_2 = 50, HCO_3 = 28. What is the diagnosis?

a) Respiratory acidosis with metabolic compensation

b) Metabolic acidosis with respiratory compensation

c) Respiratory alkalosis with metabolic compensation

d) Metabolic alkalosis with respiratory compensation

Question 14:

Which of the following drugs is not used in rheumatoid arthritis?

a) Febuxostat

b) Leflunomide

c) Etanercept

d) Methotrexate

Question 15:

A middle-aged woman with a history of constipation, dry skin and menorrhagia presents to the ER with altered sensorium, non-pitting edema, hypothermia, bradycardia, and hypotension. What is the most likely diagnosis?

a) Septic shock

b) Cardiogenic shock

c) Myxedema coma

d) Hypoglycemia

Question 16:

An elderly diabetic and hypertensive patient was carried to the emergency room in a comatose state. On examination, the blood pressure was 170/100 mmHg and pulse rate was >100/min. Plantar reflex was bilateral extensor. What is the next step to do?

a) Check blood sugar level

b) Give antihypertensive

c) IV mannitol

d) CT brain

Question 17:

A 60-year-old patient came with a history of hyponatremia. He was treated by a large volume of hypertonic fluids over 24 hours following which the patient developed quadriparesis. What is the most likely cause for this patient's condition?

a) Brain infarct

b) Brainstem injury

c) Central pontine myelinolysis

d) Rare cause of hypernatremia

Question 18:

An elderly woman presents with a chronic history of pain in the small joints of hands with stiffness of joints in the early hours of the day. The image of the patient's hands is given below. What is the most likely diagnosis?

a) Rheumatoid Arthritis

b) Osteoarthritis

c) Complex Regional Pain Syndrome

d) Villonodular synovitis

Question 19:

A triad of skin lesions, asymmetric mononeuritis multiplex and eosinophilia is seen in which of the following conditions?

a) Cryoglobulinemic vasculitis

b) Polyarteritis nodosa

c) Churg Strauss syndrome

d) Giant cell arteritis

Question 20:

'a' wave in JVP corresponds to?

a) Right atrial contraction

b) Closure of tricuspid valve

c) Onset of ventricular systole

d) Maximal atrial filling

Question 21:

Which among the following is wrong about DIC?

a) Increased schistocytes

b) Increased PT

c) Increased fibrinogen

d) Increased FDPs

Question 22:

A 16-year-old girl, who is taking antiepileptics, has had a seizure-free period of 6 months. She has no family history of epilepsy. Her EEG is now normal and she has a normal neurological exam and intelligence. What would your advice be?

a) Stop the treatment and follow up

b) Gradually taper the drug and stop treatment

c) Continue treatment for another 18 months

d) Continue lifelong treatment with antiepileptics

Question 23:

AKIN and RIFLE criteria are used to classify

a) Acute kidney injury

b) Chronic renal failure

c) Acute glomerulonephritis

d) Nephrotic syndrome

Question 24:

The most common bleeding manifestation seen in severe hemophilia is:

a) Recurrent hematomas

b) Recurrent hemarthrosis

c) Hematuria

d) Intracranial hemorrhage

Question 25:

A patient presents with left-sided facial paralysis and left-sided limb weakness for the past 1 hour. Her blood pressure is 160/100 mm Hg and her CT appears normal. What would be the next step of management?

a) Nothing, since CT was normal

b) Start on aspirin + clopidogrel

c) Intravenous thrombolysis

d) Advice BP control

Question 26:

Relative bradycardia occurs in:

a) Typhoid fever

b) Q fever

c) Leptospirosis

d) All of the above

Question 27:

Which of the following is the most specific marker for alcoholism?

a) ALT

b) GGT

c) ALP

d) LDH

Question 28:

Which of the following is a differentiating feature between cardiac tamponade and tension pneumothorax?

a) Breath sounds

b) Muffled heart sounds

c) Increased heart rate

d) Raised JVP

Question 29:

A patient presented with chronic diarrhea and steatorrhea. D-xylose test was normal and the Schilling test was abnormal. A duodenal biopsy was normal. What is the most likely diagnosis?

a) Celiac disease

b) Ulcerative colitis

c) Intestinal lymphangiectasia

d) Pancreatic insufficiency

Question 30:

Which of the following leads would show maximum R wave with the cardiac axis as +90 degrees?

a) Lead aVF

b) Lead aVL

c) Lead I

d) Lead II

Question 31:

A patient walks with a stomping gait. When asked to close his eyes and walk, he is unable to do so. Which of the following tracts is probably affected?

a) Spinocerebellar tract

b) Posterior column tract

c) Vestibulospinal tract

d) Rubrospinal tract

Question 32:

A patient with thalassemia has a history of multiple blood transfusions, iron overload, and cardiac arrhythmia. She has now come for blood transfusion and during the process, complains of backache and looks very anxious. What would you do next?

a) Stop the blood transfusion

b) Continue the transfusion but do an ECG

c) Stop the transfusion. Wait for patient to become normal and then start it again

d) Do clerical check and get ECG

Question 33:

A patient with chronic kidney disease came with complaints of vomiting and diarrhea. His blood gas reports show pH = 7.40, HCO_3^- = 23 mEq/L, Na^+ = 145 mEq/L, Cl^- = 100 mEq/L. What is your probable diagnosis?

a) No acid base abnormality

b) Respiratory acidosis

c) Metabolic alkalosis

d) High anion gap metabolic acidosis with metabolic alkalosis

Question 34:

Which of the following is not done before drawing blood for arterial blood gas analysis?

a) Allen's test

b) Rinse syringe with heparin

c) Flexion of wrist

d) Placing needle at 45 degree angle

Question 35:

Anti-nuclear antibodies are required for the diagnosis of which of the following?

a) Scleroderma

b) Systemic lupus erythematosus

c) Rheumatoid arthritis

d) Sjogren's syndrome

Question 36:

An intern is asked to measure the blood pressure of a patient with cardiac tamponade. He/she should ask the patient to:

a) Hold breath

b) Take rapid shallow breaths

c) Take long deep breaths

d) Breathe normally

Question 37:

In the sagittal section of the brain given below if the coloured area is affected, which of the following is not seen?



a) Urinary incontinence

b) Gait abnormality

c) Perianal anesthesia

d) Fecal incontinence

Question 38:

A hypertensive patient is on metoprolol. If verapamil is added, there is a risk of:

a) Atrial fibrillation

b) Bradycardia with AV block

c) Tachycardia

d) Torsades de pointes

Answer Key

Question No.	Correct Option
1	b
2	b
3	b
4	d
5	b
6	a
7	c
8	d
9	c
10	b
11	c
12	c
13	a
14	a
15	c
16	a
17	c
18	a
19	c
20	a
21	c
22	c

23	a
24	b
25	c
26	d
27	b
28	a
29	d
30	a
31	b
32	a
33	d
34	c
35	b
36	d
37	c
38	b

Detailed Explanations

Solution to Question 1:

The clinical scenario describes a patient of APL (acute promyelocytic leukemia) developing fever, tachypnea, and pulmonary infiltrates while on medical treatment. This is suggestive of APL differentiation syndrome, an adverse effect of all-trans-retinoic acid (ATRA). The management involves glucocorticoids like dexamethasone.

Along with glucocorticoids, the other management of APL differentiation syndrome includes chemotherapy for cytoreduction and supportive measures. In severe cases, temporary discontinuation of ATRA is necessary.

ATRA is an oral drug that induces the differentiation of leukemic cells bearing the t(15;17) chromosomal translocation. While ATRA reduces the incidence of DIC associated with cytarabine and daunorubicin used earlier in APL, it can cause APL differentiation syndrome due to the adhesion of differentiated neoplastic cells to the pulmonary vasculature endothelium.

In low-risk APL patients with a low leukocyte count at presentation, a combination of ATRA and arsenic trioxide (ATO) has been found to be superior and is the current standard of care.

Solution to Question 2:

The given clinical scenario of chronic pulmonary disease and the ECG showing different p wave morphologies are suggestive of multifocal atrial tachycardia.

Multifocal atrial tachycardia (MAT) occurs due to impulses arising from multiple abnormal foci in the atria. This leads to supraventricular tachycardia with p waves and QRS complexes of different morphologies. It is usually encountered in patients with chronic pulmonary disease and acute illness.

Therapy for MAT is directed at treating the underlying disease and correcting any metabolic abnormalities. Calcium-channel blockers and radiofrequency ablation of the AV node are the other treatment modalities.

The given ECG shows narrow QRS complexes with irregular RR intervals, hence it can be atrial fibrillation or multifocal atrial tachycardia. The next step is to look at the p waves. Multifocal atrial tachycardia (MAT) is characterized by at least 3 distinct P-wave morphologies (shown by arrows in the image). Whereas, in atrial fibrillation, there would be no distinct p waves seen.



Solution to Question 3:

Kartagener's syndrome is a type of primary ciliary dyskinesia characterized by the classical triad of

• Situs inversus

• Chronic sinusitis

• Bronchiectasis

Infertility in Kartagener's syndrome is due to asthenospermia (immotile spermatozoa due to impaired sperm flagella function), not azoospermia (complete absence of spermatozoa in ejaculate).

Primary ciliary dyskinesia (also called immotile cilia syndrome) is an autosomal recessive disorder affecting ciliary motility. It occurs due to ultrastructural defects in the dynein arm of cilia. This causes the retention of secretions due to a failed mucociliary clearance mechanism. Clinically, this manifests as chronic sinusitis and recurrent infections that ultimately lead to bronchiectasis. As

ciliary function is necessary to ensure proper rotation of the developing organs in the chest and abdomen, its absence will lead to situs inversus or a partial lateralizing abnormality. Approximately half of the patients with primary ciliary dyskinesia have Kartagener syndrome.

Solution to Question 4:

The GCS score is E2VNTM3. In this case:

• Eye-opening to pain: E2

• In case of endotracheal intubation or tracheostomy, the verbal component of the score is denoted as 'not testable/NT'. Earlier in intubated patients, the V score was denoted as VT. According to the current update, the V score should be VNT.

• Abnormal flexion: M3

Solution to Question 5:

The given clinical scenario of sudden heart failure in a previously healthy person under a stressful situation is suggestive of Takotsubo cardiomyopathy.

Takotsubo cardiomyopathy is also called broken heart syndrome, apical ballooning syndrome, or stress-induced cardiomyopathy. It is a transient myocardial stunning from microcirculation spams due to a catecholamine surge during stress. It is usually seen in middle-aged women. The patients can present with pulmonary edema, hypotension, and chest pain.

ECHO reveals apical dilatation with basal hypercontraction of the heart, resembling a Takotsubo (pot for catching octopus). The ECG changes can mimic myocardial infarction but changes extend beyond a particular coronary arterial territory. Coronary angiography, therefore, may be needed to rule out coronary occlusion. (Option B)

It is a self-limiting disorder resolving within days to weeks. The prognosis is generally good.

The image below shows left ventriculography during systole showing apical ballooning resembling a Takotsubo pot.



Other options:

Option A: Infective cardiomyopathy: The most common causes of infective myocarditis are viral infections and the protozoan Trypanosoma cruzi, which is associated with Chagas' disease. Acute viral myocarditis often presents with symptoms and signs of heart failure, although it can also manifest as chest pain and show electrocardiogram (ECG) changes resembling pericarditis or acute myocardial infarction. In Chagas' disease, specific features include abnormalities in the conduction system, particularly dysfunction of the sinus node and AV node, as well as the right bundle branch block.

Option C: Dilated cardiomyopathy: It is the most common cardiomyopathy and is predominantly characterised by systolic dysfunction. Though it may present acutely, it progresses chronically. The familial form is due to mutations in TTN gene which encode the sarcomeric protein titin. Other etiologocal agents include alcohol, infective myocarditis, doxorubicin, hemochromatosis and sarcoidosis.

Option D: Hypertrophic cardiomyopathy: Hypertrophic cardiomyopathy (HCM) is the most common genetic cardiovascular disorder and is a prominent cause of sudden death before the age of 35 years. Family history is commonly present, a sarcomere mutation is present in ~50% of patients. Sudden death can be due to polymorphic ventricular tachycardia/fibrillation. Risk factors for sudden death in this disease include young age, nonsustained ventricular tachycardia, failure of blood pressure to increase during exercise, recent (within 6 months) syncope, ventricular wall thickness >3 cm, and possibly the severity of LV outflow obstruction.

Solution to Question 6:

Benzathine penicillin 2.4 million units weekly for three weeks is the treatment of late cardiovascular syphilis.

The treatment of syphilis is based on the CDC's 2015 guidelines:

Note:

mU is a million units, not milli-units.

Stage of Syphilis	Patients without Penicillin Allergy	Patients with Confirmed Penicillin Allergy
Primary, Secondary and early latent	CSF Normal or not examined: Penicillin G benzathine (single dose 2.4mU IM)CSF Abnormal: Treat as Neurosyphilis	CSF normal or not examined: Tetracycline HCl (500 mg PO qid)ORDoxycycline(100 mg PO bid) for 2 weeks CSF abnormal: Treat as neurosyphilis
Late latent (or latent of uncertain duration),Cardiovascular or benign tertiary	CSF Normal or not examined: Penicillin G benzathine 2.4 mU IM weekly for 3 weeksCSF Abnormal: Treat as Neurosyphilis	CSF normal and patient not infected with HIV: Tetracycline HCl (500 mg PO qid)ORDoxycycline (100 mg PO bid) for 4 weeksCSF normal and patient infected with HIV: Desensitization and treatment with penicillin if compliance can not be ensuredCSF abnormal: Treat as neurosyphilis
Neurosyphilis (asymptomatic or symptomatic)	Aqueous crystalline penicillin G (18-24 mU/d IV), given as 3-4mU q4h or continuous infusion for 10-14 days. ORAqueous procaine penicillin G (2.4mU/d IM) + Oral probenecid(500mg qid), both for 10-14 days.	Desensitization and treatment with penicillin
Syphilis in Pregnancy	According to stage	Desensitization and treatment with penicillin

Solution to Question 7:

Azithromycin is the drug of choice for treatment of community-acquired pneumonia (CAP) in an OPD setup.

Outpatient management of CAP:

1. For previously healthy patients who have not taken antibiotics within the past 3 months:

• A macrolide – Azithromycin, 500 mg orally as a first dose and then 250 mg orally daily for 4 days, or 500 mg orally daily for 3 days, or

• Clarithromycin, 500 mg orally twice a day; or

• Doxycycline 100 mg orally twice a day

2. For patients with such comorbid medical conditions as chronic heart, lung, liver, or kidney disease, diabetes mellitus, alcoholism, malignancy, asplenia, immunosuppressant conditions or use of immunosuppressive drugs or use of antibiotics within the previous 3 months:

• A respiratory fluoroquinolone (moxifloxacin, 400 mg orally daily; gemifloxacin, 320 mg orally daily; levofloxacin, 750 mg orally daily) or

• A macrolide (as above) plus a beta-lactam (amoxicillin, 1 g orally three times a day; amoxicillin-clavulanate, 2 g orally twice a day; cefuroxime, 500 mg orally twice a day).

3. In regions with a high rate (> 25%) of infection with high level (MIC ≥ 16 mcg/mL) macrolide-resistant Streptococcus pneumoniae, consider the use of alternative agents listed in (2) for patients with comorbidities.

Other options: Vancomycin, Ceftriaxone, and Streptomycin are administered parenterally, and hence, not appropriate for outpatient management.

Solution to Question 8:

Syndrome Z is defined as a combination of obstructive sleep apnea syndrome with metabolic syndrome. LDL more than 100 mg/dL is not a criterion for the diagnosis of metabolic syndrome.

Metabolic syndrome is diagnosed based on The NCEP: ATP III Criteria. Three or more of the following must be present :

• Central obesity: waist circumference >102 cm (M), >88 cm (F)

• Hypertriglyceridemia: triglyceride level ≥150 mg/dL or specific medication

• Low HDL cholesterol: <40 mg/dL and <50 mg/dL for men and women, respectively, or specific medication

• Hypertension: blood pressure ≥130 mmHg systolic or ≥85 mmHg diastolic or specific medication

• Fasting plasma glucose level ≥100 mg/dL or specific medication or previously diagnosed type 2 diabetes

Solution to Question 9:

This is a case of cauda equina syndrome (CES).

CES signifies an injury of multiple lumbosacral nerve roots within the spinal canal distal to the termination of the spinal cord at L1-L2. It is commonly due to a ruptured lumbosacral intervertebral disk, lumbosacral spine fracture, hematoma within the spinal canal, compressive tumor, or other mass lesions.

Low back pain, weakness, and areflexia in the legs, saddle anesthesia, or loss of bladder function may occur.

Treatment options include surgical decompression, sometimes urgently in an attempt to restore or preserve motor or sphincter function, or radiotherapy for metastatic tumors.

Solution to Question 10:

The given clinical scenario is suggestive of jaundice. The tests that should be done in this patient to rule out acute viral hepatitis include:

• Anti-HAV

• HBsAg

• IgM anti-HBc

• Anti-HCV

We mainly test for hepatitis A, B, and C. Tests for hepatitis D and E are not used routinely.

Hepatitis D only occurs concurrently with Hepatitis B, either as a co-infection (when it is clinically indistinguishable from HBV infection) or superimposed on chronic HBV infection. In the latter scenario, it presents as a sudden worsening of the patient's condition. IgM anti-HDV, IgG anti-HDV, and HDV RNA assays are useful for diagnosis when hepatitis D is suspected.

Hepatitis E is not tested as the majority of patients who acquire hepatitis E virus (HEV) infection spontaneously clear the virus. Acute hepatic failure is more likely in those who are pregnant and in those who are malnourished or have preexisting liver disease. Chronic HEV has been described almost exclusively in immunocompromised hosts (e.g., organ transplantation). The diagnosis of acute HEV is typically based upon the detection of IgM antibodies to HEV. Both false positives and negatives are common with available assays and are thus, not very reliable. Thus, additional serologic testing or HEV RNA testing should be performed to confirm the diagnosis and not be useful as a screening test. [Please note the difference between anti-HBe and anti-HEV]

Condition Suspected	Targetted Test Performed
Acute Hepatitis A	IgM Anti-HAV
Acute Hepatitis B	HBsAg andIgM Anti-HBc*1 or both can be positive. A positive IgM AntiHBc with negative HBsAg is seen when HBsAg is below the detection threshold.
Acute Hepatitis C	Anti HCV

Solution to Question 11:

Yogurt is the least likely to cause symptoms of lactose intolerance. It contains lactobacillus that produces lactase and is generally well tolerated.

Lactase is a brush border enzyme that hydrolyzes lactose into glucose and galactose. In lactase deficiency, patients have great variability in clinical symptoms, depending on both the severity of lactase deficiency and the amount of lactose ingested.

The goal of treatment in patients with isolated lactase deficiency is achieving patient comfort through the elimination of dietary lactose.

Foods that are high in lactose include milk (12 g/cup), ice cream (9 g/cup), and cottage cheese (8 g/cup). Aged cheese has lower lactose content. Most people can tolerate up to 12g of lactose a day without significant symptoms.

Solution to Question 12:

Among the given options, Acinetobacter species is the most commonly isolated organism in cases of ventilator-associated pneumonia (VAP). VAP develops in a mechanically ventilated patient >48 hours after endotracheal intubation.

The organisms causing ventilator-associated pneumonia (VAP) significantly vary from one hospital to another. However, the organisms that frequently cause VAP in hospital settings are Staphylococcus aureus, Pseudomonas aeruginosa, Acinetobacter sp., Stenotrophomonas maltophilia, and gram-negative rods (Enterobacter species, K. pneumoniae, and Escherichia coli).

Three factors distinguish nosocomial pneumonia from community-acquired pneumonia:

• Different infectious agent

• Different antibiotic susceptibility patterns, specifically, a higher incidence of drug resistance

• Poorer underlying health status of patients putting them at risk for more severe infections

Within 48 hours of admission, 75% of seriously ill hospitalized patients have their upper airway colonized with organisms from the hospital environment. Impaired cellular and mechanical defense mechanisms in the lungs of hospitalized patients raise the risk of infection after aspiration has occurred.

VAP patients may be infected with Acinetobacter species and Stenotrophomonas maltophilia. Anaerobic organisms (Bacteroides, anaerobic streptococci, Fusobacterium) may also cause pneumonia in the hospitalized patient.

Solution to Question 13:

The above findings give the diagnosis of respiratory acidosis with metabolic compensation.

Interpretation of the ABG:

Step 1: pH is reduced (7.12), hence acidosis.

Step 2: Since pCO_2 is raised (50 mmHg), it is respiratory acidosis.

Step 3: HCO_3 is raised (28 mEq/L) suggesting there is metabolic compensation.

Solution to Question 14:

Febuxostat is not used in rheumatoid arthritis. It is used in the treatment of gout.

Solution to Question 15:

The given clinical history is typical of a case of hypothyroidism and the diagnosis is myxedema coma, which is a serious complication of hypothyroidism.

Myxedema coma is a part of a larger clinical picture of myxedema crisis. Myxedema crisis refers to severe, life-threatening manifestations of hypothyroidism. It is often precipitated by the administration of sedatives, antidepressants, hypnotics, anaesthetics, or opioids.

Clinical features of myxedema crisis :

• Impaired cognition, ranging from confusion to somnolence to coma (myxedema coma).

• Convulsions and abnormal CNS signs.

• Profound hypothermia (can reach 23°C), hypoventilation, hyponatremia, hypoglycemia, hypoxemia, hypercapnia, and hypotension.

• Rhabdomyolysis and acute kidney injury

• A high mortality rate

Other options:

Option A: Without a history of or focus of infection, septic shock is unlikely.

Option B: Cardiogenic shock would present with symptoms of left heart failure i.e. pitting pedal edema, abdominal distension, prominence of neck veins, and dyspnea.

Option D: Hypoglycemia usually presents with sweating, palpitations, and tachycardia

Solution to Question 16:

Ruling out glycemic extremes is the first step in the evaluation of this patient who is in a coma as most of the causes of coma are metabolic in origin and with appropriate treatment are completely reversible.

In this elderly patient, with no other relevant history, the most likely cause of coma is diabetic ketoacidosis, hyperglycemic hyperosmolar state, or even hypoglycemia (which may present with hypertension).

Other options :

The patient presents with tachycardia which is not a sign of raised intracranial pressure(bradycardia, altered respiratory pattern, raised systolic blood pressure) and hence starting iv mannitol (option 3) to reduce intracranial pressure or doing a CT (option 4) to rule out intracranial space-occupying lesion is not required.

The patient is a known case of hypertension which may cause hypertensive encephalopathy due to cerebral edema and would present with a similar history and signs of raised intracranial pressure giving antihypertensive (option 2) medications would treat the condition but it is not the first step as metabolic causes are more common and have to be addressed first.

Coma is defined as a deep sleep-like state from which the patient cannot be aroused.Coma could be due to:

• Metabolic and toxic encephalopathies, where brain imaging will be normal

• Meningitis syndromes

• Diseases with prominent focal signs like stroke, or hemorrhage

Urgent intervention is needed to prevent further brain damage. It is done by providing supplemental oxygen, intravenous thiamine (at least 100 mg), and intravenous 50% dextrose in water (25 g) empirically. It is crucial to measure baseline serum glucose levels before administering glucose.

The initial examination includes evaluating the general appearance,vitals, breath sounds, response to stimulation, pupil size and responsiveness, and any abnormal posturing or movements. In trauma cases, the neck should be stabilized until ruling out cervical spine fracture. Airway protection and intravenous access is necessary.

Immediate therapeutic intervention should be done for hypotension, significant hypertension, bradycardia, arrhythmias affecting blood pressure or marked hyperthermia. Signs of meningeal irritation or fever indicate the need for urgent lumbar puncture. Fundus examination for papilledema and a CT scan of the brain should be performed prior to lumbar puncture in comatose patients.

Solution to Question 17:

This is a typical history of sudden correction of hyponatremia leading to central pontine myelinolysis, a common complication associated with the treatment of hyponatremia.

Central pontine myelinolysis/ Osmotic demyelination syndrome is seen due to rapid correction of chronic hyponatremia (hyponatremia lasting for more than 48 hrs). It results in the efflux of organic osmolytes from brain cells, which reduces intracellular osmolality and osmotic gradient, and water entry.

Rapid correction of hyponatremia (>8–10 mM in 24 h or 18 mM in 48 h) causes hypertonic stress in astrocytes within the brain causing demyelination without inflammation at the base of the pons, with relative sparing of axons and nerve cells.

Clinically patients present one or more days after overcorrection of hyponatremia with paraparesis or quadriparesis, diplopia, a “locked-in syndrome,” dysphagia, dysarthria with/without loss of consciousness.

MRI is useful in establishing the diagnosis and may also identify partial forms that present as confusion, dysarthria, and/or disturbances of conjugate gaze without paraplegia.

Prevention can be done by restoration of severe hyponatremia gradually i.e. by ≤10 mmol/L (10 mEq/L) within 24 h and ≤20 mmol/L (20 mEq/L) within 48 h.



The image shows the MRI finding of central pontine myelinolysis with axial fat saturated T2-Weighted MRI showing hyperintense signal in the central pons.

Note: Brain infarct or brain stem injury usually do not result in a quadriparesis.

Solution to Question 18:

The above image shows flexion at the PIP joint and extension in the DIP joints which is known as boutonniere deformity which is seen in rheumatoid arthritis(RA).

If the patient gives a history of pain/stiffness in the joints in the early hours of the day which usually decreases as the day progresses suggesting inflammatory arthritis and if it presents with predominant involvement of smaller joints, it is most likely a case of RA(option A).

It occurs due to the accumulation of inflammatory mediators which on increased joint mobility are flushed out due to increased blood supply.

Pain and stiffness in osteoarthritis (Option B) usually increase as the day progresses due to an increase in the activity of the joint. They typically present with nodules around the distal interphalangeal joint (Heberden nodes) and proximal interphalangeal joint (Bouchard nodes).

Solution to Question 19:

A triad of skin lesions, asymmetric mononeuritis multiplex, and eosinophilia is seen in Churg–Strauss syndrome (eosinophilic granulomatosis with polyangiitis).

It is a necrotizing vasculitis of the small-to-medium-sized vessels and in order to be diagnosed with Churg-Strauss, a patient should have evidence of asthma, peripheral blood and tissue eosinophilia (>1000 cells/μL), extravascular granuloma formation and clinical features consistent with vasculitis.

Common presenting symptoms of this condition are lower respiratory tract symptoms such as severe attacks of asthma, upper respiratory tract symptoms such as allergic rhinitis, sinusitis, and nasal polyps, mono neuritis multiplex (a type of peripheral neuropathy affecting at least two nerves) which presents with a complaint of pain, numbness, or weakness in the affected areas and skin lesions like palpable purpura and subcutaneous nodules.

Renal involvement is less common and less severe than other small vessel vasculitis.

P-ANCA antibodies are present in Churg Strauss syndrome.

• Severe attacks of asthma

• Mononeuritis multiplex: a type of peripheral neuropathy affecting at least two nerves. Patients will complain of pain, numbness, or weakness in the affected areas.

• Skin lesions like palpable purpura and subcutaneous nodules.

Glucocorticoids show effective responses in these patients. Other drugs that have been tried are mepolizumab and rituximab.

For fulminant multisystem disease, daily cyclophosphamide with prednisolone followed by azathioprine or methotrexate is used.

Option A: Cryoglobulinemic vasculitis presents with palpable purpura, arthralgia, neuropathy, and glomerulonephritis. It is most commonly associated with Hepatitis C.

Option B: Polyarteritis nodosa can also present with mononeuritis multiplex and skin lesions, but they do not have eosinophilia, it has predominant neutrophilia.

Option D: Giant cell arteritis manifests with the cardinal symptom of headache localized to the temporal region. Common symptoms include headache, polymyalgia rheumatica , jaw claudication, fever, and decrease in body weight. Laboratory results show increased ESR, CRP levels, normochromic or mildly hypochromic anemia.

Solution to Question 20:

The "a" wave in jugular venous pressure (JVP) is a positive wave, which corresponds to right atrial contraction.

The waveforms seen in a normal JVP are:

• a wave corresponds to atrial contraction and occurs just after the P wave (on ECG) and just before the first heart sound.

• c wave represents the bulging of the tricuspid valve into the right atrium during ventricular systole.

• x descent corresponds to the atrial relaxation during atrial diastole.

• v wave represents atrial filling during ventricular systole and peaks at the second heart sound.

• y descent represents passive emptying of the right atrium into the right ventricle after tricuspid valve opening.



Solution to Question 21:

In disseminated intravascular coagulopathy (DIC), fibrinogen levels are decreased due to increased bleeding.

Disseminated intravascular coagulation (DIC) is an acquired syndrome characterized by the widespread activation of blood clotting factors within blood vessels, resulting in a loss of clotting control. The primary mechanism of DIC involves the uncontrolled generation of thrombin through multiple pathways which leads to sustained coagulation activation, the formation of fibrin, and the consumption of clotting factors and platelets.

Infections are the most frequent cause of DIC.

Laboratory investigations in DIC:

• Decreased platelet count

• Prolonged PT/TT

• Prolonged activated partial thromboplastin time (aPTT)

• Decreased fibrinogen

• Elevated fibrin degradation products (D-dimers)

• Peripheral smear: microangiopathic hemolytic anemia in DIC with schistocytes (fragmented red blood cells)

The treatment involves platelet transfusion when platelet counts are below 10,000-20,000/μL. Replacement of fibrinogen and coagulation factors is with fresh frozen plasma (FFP) along with cryoprecipitate or fibrinogen concentrate.

In cases of low-grade DIC associated with solid tumors or acute promyelocytic leukemia low doses of continuous-infusion heparin has been effective.

Solution to Question 22:

In the given clinical scenario of a girl on antiepileptics for 6 months with no further seizures, no family history, normal EEG, neurological examination, and intelligence, antiepileptics should be continued for another 18 months since she's already taken for 6 months (minimum 2 years) to prevent seizure recurrence.

Anti-epileptic drugs withdrawal is considered only when an adult patient has been seizure-free for ≥2 years along with the following criteria:

• Normal CNS examination including intelligence

• Normal EEG and normal MRI

• Single seizure type

• No family history of epilepsy.

It is preferable to reduce the dose of the drug gradually over 2–3 months.

Other options:

Option A: Sudden cessation of drug therapy may cause recurrence of seizure hence not advised.

Option B: The patient has a seizure-free interval of only 6 months and therefore, is not yet eligible for considering medication withdrawal.

Option D: Life-long treatment is not necessary in this case as the patient meets all the above-mentioned criteria for drug withdrawal after 2 years.

Note: Juvenile myoclonic epilepsy (JME), the most common idiopathic generalized epilepsy syndrome is known to have a high risk of seizure recurrence, and hence antiepileptics are continued lifelong in a majority of patients.

Solution to Question 23:

AKIN and RIFLE criteria are used to classify acute kidney injury (AKI).

The various classifications used in AKI are:

• RIFLE criteria (Risk, Injury, Failure, Loss, End-stage)

• AKIN (AKI Network) Classification

• KDIGO (Kidney Disease Improving Global Outcomes) Classification

RIFLE, AKIN, KDIGO definitions of AKI:

RIFLE, AKIN, KDIGO staging of AKI:

	RIFLE	AKIN	KDIGO
S. creatinine	>50% increase that develops in <7 days	>0.3 mg/dl increase or >50% increase that develops in <48 hr	>0.3 mg/dl increase or >50% increase that develops in <7 days

	RIFLE	AKIN	KDIGO
Urine output	<0.5ml/kg/hr for >6 hr	<0.5ml/kg/hr for >6 hr	<0.5ml/kg/hr for >6 hr

RIFLE	Increase in S.creatinine	AKIN	Increase in S.creatinine	KDIGO	Increase in S.creatinine	Urine output
Risk	≥50%	Stage 1	≥0.3mg/dl or ≥50%	Stage 1	≥0.3mg/dl or ≥50%	<0.5mL/kg/hr for >6h
Injury	≥100%	Stage 2	≥100%	Stage 2	≥100%	<0.5mL/kg/hr for >12 hr
Failure	≥200%	Stage 3	≥200%	Stage 3	≥200%	<0.3mL/kg/hr for >24hr or anuria >12hr
Loss	Need for renal replacement therapy (RRT)>4 weeks					
End-stage	Need RRT for >3 months					

Solution to Question 24:

Recurrent hemarthrosis is the most common bleeding manifestation of severe hemophilia. Hemophilia is said to be severe when the residual activity of factor VIII is <1%.

Hemophilia is an X-linked recessive hemorrhagic disorder and is caused by mutations in either the F8 gene (hemophilia A or classic hemophilia) or the F9 gene (hemophilia B). Clinically, it is impossible to distinguish between hemophilia A and hemophilia B as they present with similar symptoms. It can be classified based on residual activity of FVIII or FIX as severe (<1%), moderate (1–5%), or mild (6–30%)

Hemophilia presents early in life- bleeding after circumcision or when beginning to walk or crawl.

Clinical features:

• Acute hemarthrosis:

• Local pain, swelling

• Erythema

• Muscle contractures due to fixed positioning.

• Chronic hemarthrosis:

• Synovial thickening

• Synovitis (due to intraarticular blood)

The most common joint affected is the knee. One joint is affected more than the other. This is called the target joint. Muscle hematomas can lead to compartment syndromes.

Hematuria is common but self-limited. Life-threatening bleeds such as bleeding into oropharyngeal spaces, CNS, and retroperitoneum can occur.

Treatment: Without treatment, severe hemophilia may limit life expectancy. Factor replacement for hemophilia has been the mainstay of therapy.

• FVIII dose (IU) = Target FVIII levels – FVIII baseline levels × body weight (kg) × 0.5 unit/kg .

• FIX dose (IU) = Target FIX levels – FIX baseline levels × body weight (kg) × 1 unit/kg

The FVIII half-life of 8–12 h requires injections twice a day to maintain therapeutic levels, whereas the FIX half-life is longer, ~24 h so that once-a-day injection is sufficient.

Solution to Question 25:

Sudden onset hemiparesis with apparently normal CT indicates ischemic stroke in this patient. Intravenous thrombolysis(alteplase 0.9 mg/kg) should be immediately administered to patients with ischemic stroke who arrive in the window period i.e., within 3-4.5 hours of symptom onset.

Aspirin + clopidogrel(Option B) is used in secondary prevention of stroke and also to prevent stroke following TIA.

Transient ischemic attacks (TIAs) refer to episodes of stroke symptoms that have a brief duration, the standard definition of duration is <24 h, but most TIAs last <1 hr. TIAs may arise from emboli to the brain or from in situ thrombosis of an intracranial vessel. With a TIA, the occluded blood vessel reopens and neurologic function is restored. If a relevant brain infarction is identified on brain imaging, the clinical entity is now classified as stroke regardless of the duration of symptoms.

The risk of stroke following a transient ischemic attack (TIA) is approximately 10-15% in the first three months, majorly in the first two days. A reliable estimation of this risk can be obtained using the ABCD2 score.

Other options:

Option A: If CT is normal it only rules out hemorrhagic stroke and changes of ischemic stroke usually take time to appear on NCCT.

Option D: BP control is done when BP exceeds 220/120 mmHg, if there is malignant hypertension or concomitant myocardial ischemia, or if blood pressure is >185/110 mmHg and thrombolytic therapy is anticipated.

Solution to Question 26:

In the given options typhoid, Q fever, and leptospirosis are infectious causes of relative bradycardia.

Relative bradycardia, also known as Faget's sign, is observed when patients exhibit a lower heart rate than anticipated in relation to an increase in body temperature. Body temperature is commonly regarded as a factor that independently influences heart rate. It is estimated that for each 1°C (1.8°F) rise in temperature, the heart rate typically increases by approximately 15-20

beats per minute.

While this pulse-temperature dissociation is not highly sensitive or specific for establishing a diagnosis, it can still prove beneficial in resource-limited settings due to its easy accessibility and simplicity.

Infectious causes of relative bradycardia include:

• Salmonella typhi

• Francisella tularensis

• Coxiella burnetii (Q fever)

• Leptospira interrogans

• Legionella pneumophila

• Rickettsia spp.

• Orientia tsutsugamushi (scrub typhus)

• Corynebacterium diphtheriae

• Plasmodium spp. (malaria)

Noninfectious causes include:

• Drug fever

• Beta-blocker use

• Central nervous system lesions

• Malignant lymphoma

• Factitious fever

Solution to Question 27:

GGT (Gamma Glutamyl Transferase) is the most specific marker of alcoholism among the given options.

The markers with >60% sensitivity and specificity for heavy alcohol use are gamma-glutamyl transferase (>35 u) and carbohydrate-deficient transferrin (CDT), the combination of the two is likely to be more accurate than one. An isolated elevation in serum GGT or GGT elevation out of proportion to that of other enzymes (such as the ALT and ALP) may be an indicator of alcohol abuse or alcoholic liver disease.

ALT (option A) is usually elevated in acute hepatocellular conditions such as hepatitis. Alanine aminotransferase is predominantly found in the liver, hence a more specific marker of liver injury. A ratio of AST/ALT > 2:1 indicates the possibility of alcoholic liver disease and > 3:1 ratio is highly indicative of this condition. In alcoholic liver disease, ALT levels are often normal.

ALP (option C) is elevated in cholestasis, ■■is also elevated when there is rapid bone growth like in children and adolescents because of bone alkaline phosphatase and in normal pregnancies due to placental alkaline phosphatase.

LDH (option D) elevation is nonspecific.

Solution to Question 28:

The differentiating feature between cardiac tamponade and tension pneumothorax is breath sounds. The breath sounds are absent in tension pneumothorax while they are present in cardiac tamponade.

The presence of hypotension, distended jugular veins, and muffled heart sounds (Beck's triad) suggest a cardiac tamponade. These features can also be seen in tension pneumothorax. Other clinical features pointing towards tension pneumothorax are absent breath sounds, hyper resonance to percussion, and shift of the mediastinum to the contralateral side.

Heart sounds can be muffled by structures between the stethoscope and the heart, eg, adipose tissue deposition (obesity), fluid accumulation (pericardial effusion, pleural effusion), gas accumulation (pneumothorax, subcutaneous emphysema), soft tissue masses can lead to attenuation of heart sounds.

Solution to Question 29:

The clinical scenario describes a case of malabsorption syndrome (chronic diarrhea and steatorrhea), a normal xylose test and normal duodenal biopsy with abnormal schilling test points towards pancreatic insufficiency.

A normal xylose test and normal duodenal biopsy rules out small intestinal pathologies like celiac disease (option A) and intestinal lymphangiectasia (option C), presence of steatorrhea rules out ulcerative colitis (option B) as it affects the large bowel which is not the main site of absorption of fat.

Ileal disease is also characterized by normal xylose test, normal duodenal biopsy and steatorrhea and abnormal schilling test but among the given options the most appropriate answer would be pancreatic insufficiency due to chronic pancreatitis.

Pancreatic insufficiency is characterised by the deficiency of exocrine pancreatic enzymes that result in maldigestion and malabsorption.

The features in favor of the diagnosis:

• Steatorrhea - a deficiency of pancreatic lipases can lead to undigested fat in stools.

• D-xylose test - d-xylose is a monosaccharide that is directly absorbed by the intestinal mucosa without undergoing any enzymatic action. Normal d-xylose test denotes intact mucosa.

• Schilling test - it is abnormal as pancreatic enzymes are required for the absorption of vitamin B_{12}.

Diagnosis of various malabsorption syndromes:

Pathologies	D xylose test	Duodenal mucosal biopsy	Schilling test
Celiac disease	Decreased	Abnormal; Short or absen t villi; epithelial cell dama ge; crypts hypertrophy	Normal
Ileal disease	Normal	Normal	Abnormal
Chronic pancreatitis	Normal	Normal	Abnormal; normal after p ancreatic enzymes substit ution
Intestinal lymphangiectasi a	Normal	Abnormal; dilated lympha tics, clubbed villi	Normal
Bacterial overgrowth synd rome	Normal or may be abn ormal	Usually normal or can hav e diffuse non specific dam age to villi	Usually abnormal; normal after antibiotics administ ration

Solution to Question 30:

With the cardiac axis at +90 degrees, lead aVF would show the maximum R wave.

The cardiac axis represents the mean QRS axis (direction of mean depolarization vector).

The normal adult QRS axis is directed downward and to the left, which is between -30 degrees and +100 degrees.

The position of lead aVF is at +90 degrees. If the cardiac axis is also at +90 degrees, it would be in the same direction as lead aVF and hence aVF would show maximum R wave amplitude.

If the mean axis is:

• More negative than –30 degrees: left axis deviation

• More positive than +90 to +100 degrees: right axis deviation

• Lies between -90 to +180 degrees: extreme axis deviation



Solution to Question 31:

The given clinical scenario is suggestive of the involvement of the posterior column.

Posterior column carries proprioception which is essential for coordination of motor movements. Loss of this leads to the development of a stomping gait. This is seen in cases of vitamin B12 deficiency leading to subacute combined degeneration, tabes dorsalis or diabetic neuropathy.

Solution to Question 32:

The patient is being transfused blood and has complaints of backache, this is suggestive of an acute hemolytic transfusion reaction.

The first step is to stop transfusion immediately and then collect blood samples for testing.

Acute hemolytic transfusion reactions typically occur within the first 24 hours following a blood transfusion, primarily due to ABO incompatibility.

It presents with symptoms such as hypotension, tachypnea, tachycardia, fever (1-2°C increase in temperature), chills, chest and back pain, hemoglobinuria, and hemoglobinemia. In severe cases, disseminated intravascular coagulation (DIC), acute renal failure, shock, and death can occur.

Management of immune-mediated acute hemolytic transfusion reactions is to provide supportive care, administer vigorous hydration with isotonic saline and use diuretics to maintain urine output.

Solution to Question 33:

The probable diagnosis is a high anion gap metabolic acidosis (HAGMA) with metabolic alkalosis.

Interpretation of the ABG:

Step 1: The pH being 7.4 is within normal limits.

Step 2: Plasma HCO_3^- is 23 mEq/L and again is within normal limits.

At this juncture, we have ruled out options B and C as this particular clinical scenario is not suggestive of a simple acid-base disorder.

Step 3: Calculating the Anion gap:

Anion gap = $[Na^+] - ([Cl^-] + [HCO_3^-])$

= 145-(100+23)

= 22 mEq/L

The presence of a high anion gap indicates that there's indeed an acid-base disorder, thus ruling out option A.

A high anion gap indicates metabolic acidosis. However, HCO_3^- levels within normal limits suggest that there is also coexisting metabolic alkalosis in this patient.

This can be confirmed by calculating the delta gap.

Delta gap = $\Delta AG - \Delta HCO_3^-$

In this scenario: Delta gap = (22-10) - (24-23) = 11 mEq/L

Delta gap is > +6 suggesting the presence of metabolic alkalosis along with HAGMA.

If the delta gap is significantly positive (>+6), metabolic alkalosis is present because the rise in the anion gap is more than the fall in bicarbonate. If the delta gap is significantly negative (<6) then hyperchloremic acidosis is usually present because a rise in the anion gap is less than a fall in bicarbonate.

Alternatively, this can be confirmed by calculating the delta ratio.

Delta ratio = the increase in Anion Gap / the decrease in HCO_3^-

It is used in the presence of a high anion gap metabolic acidosis (HAGMA), to check it is a 'pure' HAGMA or if there is coexistant normal anion gap metabolic acidosis (NAGMA) or metabolic alkalosis.

Delta ratio values:

• 1-2 indicates HAGMA

• < 1 indicates NAGMA + HAGMA

• > 2 indicates HAGMA + Metabolic alkalosis

In this scenario,

Delta ratio = 12/1 = 12

Thus, the acid-base disorder for the above scenario is HAGMA with metabolic alkalosis.

Solution to Question 34:

Flexion of the wrist is not done just before drawing blood for ABG analysis.

During arterial blood gas analysis, the ideal position for drawing arterial blood from the radial artery is to maintain the wrist in extension. This provides stability to the artery and prevents it from slipping away during the puncture. To draw blood for ABG analysis, the skin is pierced at a 45° angle.



Allen's test is done prior to radial cannulation to identify the adequacy of radial and ulnar collateral blood flow to the hand.



The syringe used to draw blood for ABG analysis is rinsed with heparin prior to the procedure. Prior to collecting blood, 0.1 ml heparin is drawn into the syringe and then flushed out. The residue that lubricates the inner wall of the syringe is sufficient to prevent the blood from clotting.

Solution to Question 35:

Antinuclear antibodies (ANAs) are required for the diagnosis of systemic lupus erythematosus.

ANAs are autoantibodies that bind to the contents of the cell nucleus. They are present in > 95% of patients with SLE. A positive antinuclear antibody (ANA) test with a titer of at least 1:80

is the obligatory entry criterion along with the additive criteria in 7 clinical and 3 immunologic domains. If a total of >10 points achieved, it is indicative of SLE according to 2019 EULAR/ACR Classification Criteria for Systemic Lupus Erythematosus (SLE)

Most sensitive antibody for SLE: ANA

Most specific antibody for SLE: Anti smith > Anti dsDNA

ANA may also be positive in the other options but it is not a criterion for diagnosis of these conditions.

The following antibodies are specific for their respective autoimmune conditions.

• Scleroderma (option A): Anti-topoisomerase I(Scl-70) and anti-centromere antibodies highly specific for Systemic sclerosis.

• Rheumatoid arthritis (option C): RF and anti-CCP antibodies

• Sjogren syndrome (option D): Anti-Ro, Anti-La

Solution to Question 36:

The patient with cardiac tamponade is asked to breathe normally to elicit pulsus paradoxus.

In a patient with cardiac tamponade, both ventricles share an incompressible pericardial covering due to the fluid. As the right ventricle enlarges during inspiration, the interventricular septum bulges into the left ventricle, reducing the left ventricular volume.

This results in reduced stroke volume and hence reduced systolic arterial pressure.

As a result, there is a greater than normal (more than 10 mmHg) fall in systolic arterial blood pressure during inspiration, producing pulsus paradoxus.

This cannot be elicited if the patient is holding his/her breath, taking long or shallow breaths.

Pulsus paradoxus is also seen with effusive constrictive pericarditis, restrictive cardiomyopathy and right ventricular myocardial infarction.

Solution to Question 37:

Paracentral lobule (PCL) is the continuation of the precentral and postcentral gyri of the superolateral surface into the medial surface of cerebral hemisphere. It controls the motor and sensory innervations of the contralateral lower extremity. It is also responsible for cortical control of micturition and defecation.

The anterior two-thirds of PCL belong to Brodmann area 4, it reflects the primary motor cortex for the muscles of contralateral leg, foot, and perineum,it is functionally important in control of bladder and anal sphincters. Therefore, it's called the cortical center of defecation and micturition.

The posterior third of the PCL contains 3,2 and 1, it reflects the primary somatosensory representation of the leg and foot.

The A2 segment of the anterior cerebral artery supplies the paracentral lobule. Lesions in the paracentral lobule lead to:

• Weakness and loss of sensation in contralateral lower limb

• Urinary incontinence

• Fecal incontinence.

Solution to Question 38:

Metoprolol, when added with verapamil, leads to bradycardia with AV block.

Verapamil and β blockers have additive effect on inhibiting SA node and AV conduction, and the combination should be avoided to prevent severe bradycardia and conduction block.

These agents can also have additive effect on decreasing left ventricular function which could seriously impair cardiac output in a patient with heart failure.

Medicine AIIMS 2019 (May & Nov)

Question 1:

A 16-year-old girl, who is taking antiepileptics, has had a seizure-free period of 6 months. She has no family history of epilepsy. Her EEG is now normal and she has a normal neurological exam and intelligence. What would your advice be?

a) Stop the treatment and follow up

b) Gradually taper the drug and stop treatment

c) Continue treatment for another 18 months

d) Continue lifelong treatment with antiepileptics

Question 2:

AKIN and RIFLE criteria are used to classify

a) Acute kidney injury

b) Chronic renal failure

c) Acute glomerulonephritis

d) Nephrotic syndrome

Question 3:

Which of the following statements about polycythemia vera is true?

a) Risk of thrombosis strongly correlates with the degree of thrombocytosis

b) Generalized pruritus is a consequence of mast cell activation by JAK 2 mutation

c) It is protective against H.pylori infection

d) Acute myeloid leukemia is the most common cause of mortality

Question 4:

All of the following are the core clinical features for the diagnosis of Lewy body dementia except:

a) Fluctuating cognition

b) Recurrent visual hallucination

c) REM sleep behavior disorder

d) Orthostatic hypotension

Question 5:

A 68-year-old male admitted to the ICU for acute exacerbation of COPD develops a sudden onset of palpitations. The following ECG changes are seen. What is the most likely diagnosis?



a) Ventricular tachycardia

b) Multifocal atrial tachycardia

c) Atrial fibrillation

d) Atrial tachycardia

Question 6:

A patient has recurrent optic neuritis of both eyes with extensive transverse myelitis. Visual acuity in the right eye is 6/60 and visual acuity in the left eye 6/18. The patient showed a partial response to steroids. What is the likely diagnosis?

a) Neuromyelitis optica

b) Subacute combined degeneration of spinal cord

c) Posterior cerebral artery stroke

d) Neurosyphilis

Question 7:

Deep y descent in JVP is seen in all except:

a) Cardiac tamponade

b) Restrictive cardiomyopathy

c) Constrictive pericarditis

d) Tricuspid regurgitation

Question 8:

Tigroid white matter on MR imaging is seen in which of the following conditions?

a) Pantothenate kinase deficiency

b) Pelizaeus-Merzbacher disease

c) Neuroferritinopathy

d) Aceruloplasminemia

Question 9:

All of the following characteristics are found in the pleural effusion fluid of a rheumatoid arthritis patient except:

a) RA factor

b) High glucose

c) Cholesterol crystals

d) High LDH

Question 10:

What is the most common pulmonary manifestation of SLE?

a) Shrinking lung syndrome

b) Pleuritis

c) Intra alveolar hemorrhage

d) Interstitial inflammation

Question 11:

True about the definition of Postural Hypotension:

a) Decrease in Systolic blood pressure 20 mmHg after 6 mins of standing

b) Decrease in Systolic blood pressure 20 mmHg after 3 min of standing

c) Decrease in Diastolic blood pressure 20 mmHg after 6 mins of standing

d) Decrease in diastolic blood pressure 20 mmHg after 3 mins of standing

Question 12:

The given pattern of EEG is found in:

a) Hepatic encephalopathy

b) Creuzfeldt-Jakob Disease

c) Generalized tonic clonic seizures

d) Herpes simplex encephalitis

Question 13:

Calculate the anion gap from the following values: Na+ = 137 mmol/L K+ = 4 mmol/L Cl- = 100 mmol/L HCO_3^- = 15 mmol/L

a) 22 mmol/L

b) 16 mmol/L

c) 10 mmol/L

d) 12 mmol/L

Question 14:

The ECG of a patient who presented to the ER is as given below. The patient was already given a carotid massage and IV Adenosine but the BP is currently 60/30 mm Hg. What is the next best step of management?



a) Repeat Inj adenosine 6 mg

b) Inj amiodarone 300 mg

c) Synchronised cardioversion

d) Defibrillation

Question 15:

Identify the diagnosis.

a) Facial nerve palsy

b) Plexiform neurofibromatosis

c) Masticator space abscess

d) Fibrous dysplasia

Question 16:

What is the most common presentation of inhalational anthrax?

a) Atypical pneumonia

b) Hemorrhagic mediastinitis

c) Lung abscess

d) Broncho pulmonary pneumonia

Question 17:

A patient of hemophilia received multiple blood transfusions. Which of the following metabolic abnormalities can be seen in a patient?

a) Metabolic alkalosis

b) Respiratory alkalosis

c) Metabolic acidosis

d) Respiratory acidosis

Question 18:

A patient with renal failure develops vomiting in ICU. ABG analysis showed pH - 7.4, Na+ = 145 mEq/L, Cl- = 100 mEq/L, HCO_3^- = 24 mEq/L, $PaCO_2$ = 40 mmHg. What is the likely metabolic abnormality?

a) Normal ABG

b) Normal anion gap, metabolic acidosis

c) High anion gap metabolic acidosis with metabolic alkalosis

d) High anion gap metabolic alkalosis with normal metabolic acidosis

Question 19:

A 56-year-old woman presented with a gradually diminishing visual field. She was found to have a pituitary mass and underwent trans-sphenoidal hypophysectomy. Now she has low levels of ACTH, TSH, FSH, and LH. Which of the following hormone supplements is not needed in the patient postoperatively?

a) Mineralocorticoids

b) Thyroid hormones

c) Glucocorticoids

d) Estradiol

Question 20:

A patient with acute hepatitis B has recovered from the infection. Identify the serological marker marked in the image:

a) Anti-HBc Ab

b) Anti-HBs Ab

c) Anti-HBe Ab

d) HBsAg

Question 21:

The ECG of a pregnant lady having pre-eclampsia is shown below. Her vitals are stable. What is the best step in the management of the condition?



a) IV calcium

b) IV $MgSO_4$

c) DC Shock

d) Synchronised cardioversion

Question 22:

Which of the following is the most common cause of infection post-solid organ transplantation in an Indian setting?

a) Cytomegalovirus

b) Varicella zoster virus

c) Epstein-Barr virus

d) Herpes simplex virus

Question 23:

Addison's disease is commonly associated with ____

a) Autoimmune adrenalitis

b) Adrenocortical carcinoma

c) Hypernephroma

d) Medullary carcinoma thyroid

Question 24:

Which of the following is used in the treatment of hypercalcemia due to Vitamin D toxicity?

a) Dexamethasone

b) Hydroxychloroquine

c) Chloroquine

d) Ketoconazole

Answer Key

Question No.	Correct Option

1	c
2	a
3	b
4	d
5	b
6	a
7	a
8	b
9	b
10	b
11	b
12	b
13	a
14	c
15	b
16	b
17	a
18	c
19	a
20	a
21	b
22	a
23	a
24	a

Detailed Explanations

Solution to Question 1:

In the given clinical scenario of a girl on antiepileptics for 6 months with no further seizures, no family history, normal EEG, neurological examination, and intelligence, antiepileptics should be continued for another 18 months since she's already taken for 6 months (minimum 2 years) to prevent seizure recurrence.

Anti-epileptic drugs withdrawal is considered only when an adult patient has been seizure-free for ≥2 years along with the following criteria:

• Normal CNS examination including intelligence

• Normal EEG and normal MRI

• Single seizure type

• No family history of epilepsy.

It is preferable to reduce the dose of the drug gradually over 2–3 months.

Other options:

Option A: Sudden cessation of drug therapy may cause recurrence of seizure hence not advised.

Option B: The patient has a seizure-free interval of only 6 months and therefore, is not yet eligible for considering medication withdrawal.

Option D: Life-long treatment is not necessary in this case as the patient meets all the above-mentioned criteria for drug withdrawal after 2 years.

Note: Juvenile myoclonic epilepsy (JME), the most common idiopathic generalized epilepsy syndrome is known to have a high risk of seizure recurrence, and hence antiepileptics are continued lifelong in a majority of patients.

Solution to Question 2:

AKIN and RIFLE criteria are used to classify acute kidney injury (AKI).

The various classifications used in AKI are:

• RIFLE criteria (Risk, Injury, Failure, Loss, End-stage)

• AKIN (AKI Network) Classification

• KDIGO (Kidney Disease Improving Global Outcomes) Classification

RIFLE, AKIN, KDIGO definitions of AKI:

RIFLE, AKIN, KDIGO staging of AKI:

	RIFLE	AKIN	KDIGO
S. creatinine	>50% increase that develops in <7 days	>0.3 mg/dl increase or >50% increase that develops in <48 hr	>0.3 mg/dl increase or >50% increase that develops in <7 days
Urine output	<0.5ml/kg/hr for >6 hr	<0.5ml/kg/hr for >6 hr	<0.5ml/kg/hr for >6 hr

RIFLE	Increase in S.creatinine	AKIN	Increase in S.creatinine	KDIGO	Increase in S.creatinine	Urine output
Risk	≥50%	Stage 1	≥0.3mg/dl or ≥50%	Stage 1	≥0.3mg/dl or ≥50%	<0.5mL/kg/hr for >6h
Injury	≥100%	Stage 2	≥100%	Stage 2	≥100%	<0.5mL/kg/hr for >12 hr
Failure	≥200%	Stage 3	≥200%	Stage 3	≥200%	<0.3mL/kg/hr for >24hr or anuria >12hr

RIFLE	Increase in S.creatinine	AKIN	Increase in S.creatinine	KDIGO	Increase in S.creatinine	Urine output
Loss	Need for renal replacement therapy (RRT)>4 weeks					
End-stage	Need RRT for >3 months					

Solution to Question 3:

Generalized pruritus seen in polycythemia vera (PV) is a consequence of the activation of mast cells by JAK 2 mutation.

Other options:

Option A: The risk of thrombosis strongly correlates with erythrocytosis, not the degree of thrombocytosis.

Option C: In polycythemia vera, there is an increased risk of H.pylori infection.

Option D: Thrombosis is the major cause of mortality in polycythemia vera. The incidence of acute leukemia in patients not exposed to chemotherapy or radiation therapy is low.

PV is a clonal hematopoietic stem cell disorder in which red cells, granulocytes & platelets accumulate in the absence of a physiological stimulus. The most common mutation seen in PV is the JAK2 mutation.

Clinical features of polycythemia vera are:

• Thrombosis is the most common manifestation and is the major cause of mortality. Uncontrolled erythrocytosis causes increased viscosity of the blood causing thrombosis of the vessels. Patients may present with stroke, transient ischemic attacks, myocardial infarction, and Budd-Chiari syndrome.

• Aquagenic pruritis is generalized itching following a warm bath or a shower due to increased mast cells and histamine.

• Erythromelalgia is erythema and pain in the fingers, hands, and feet due to increased platelet stickiness.

• Splenomegaly due to increased destruction of RBCs in the spleen

• Gout is due to the excessive proliferation of the marrow cells leading to elevated uric acid levels.

Solution to Question 4:

Orthostatic hypotension is not a core clinical feature of Lewy body dementia (LBD) as per DSM-5.

LBD also known as dementia with Lewy bodies is the second most type of progressive dementia after Alzheimer's disease. Protein deposits, called Lewy bodies, deposit in nerve cells in the brain region involved in thinking, memory, and movement (motor control)

Core features of LBD as per DSM-5 are fluctuating cognition, attention, alertness, recurrent visual hallucination, and Parkinsonian symptoms. Suggestive features are REM sleep behavior disorder and severe neuroleptic sensitivity.

Orthostatic hypotension, erectile dysfunction, and constipation are other features of LBD.

Treatment of LBD includes rivastigmine and donepezil which may improve cognition, and reduce hallucinosis and delusion. Pimavanserin may be used to treat psychosis.

Solution to Question 5:

The given clinical scenario of a COPD patient in ICU developing palpitations and the ECG showing narrow QRS complex (<0.12 s) tachycardia with more than three different P wave morphologies, points to the diagnosis of multifocal atrial tachycardia (MAT).

Multifocal atrial tachycardia (MAT) is characterized by at least 3 distinct P wave morphologies (shown by arrows in the image) with rates typically between 100 and 150 beats/min.

It is usually encountered in patients with chronic pulmonary disease and acute illness. The clinical features are most commonly related to the underlying precipitating illness (COPD in this case). Unlike atrial fibrillation, there are clear isoelectric intervals between P waves.

Treating the underlying disease and correcting any metabolic abnormalities is the mainstay of treatment. Electrical cardioversion has no effect.

The calcium-channel blockers verapamil or diltiazem may slow the atrial and ventricular rate. MAT may respond to amiodarone, but long-term therapy with this agent is usually avoided due to its toxicities, particularly in pulmonary fibrosis.

Solution to Question 6:

The given clinical scenario of a patient with recurrent optic neuritis of both eyes with extensive transverse myelitis, showing partial response to steroids points to the diagnosis of neuromyelitis optica (NMO), also known as Devic's disease.

It is an aggressive inflammatory demyelinating disorder characterized by recurrent attacks of optic neuritis and myelitis. Attacks of optic neuritis can be bilateral or unilateral and myelitis can be severe and involve three or more contiguous vertebral segments.

The core clinical characteristics include optic neuritis, acute myelitis, area postrema syndrome with episodes of otherwise unexplained hiccups or nausea or vomiting, acute brainstem syndrome, symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions and symptomatic cerebral syndrome with NMOSD-typical brain lesions.

Acute attacks of NMO are treated with high-dose glucocorticoids. Plasma exchange can be used in acute attacks that do not respond to glucocorticoids.

Drugs used for prophylaxis against relapses include mycophenolate mofetil, rituximab, glucocorticoids, and azathioprine. Three monoclonal antibodies approved for attack prevention include eculizumab (terminal complement inhibitor), inebilizumab (B-cell depleter), and

satralizumab (IL-6 receptor blocker).

Solution to Question 7:

The y-descent in jugular venous pulse (JVP) corresponds to the fall in right atrial pressure after the tricuspid valve opening in the early diastolic phase of the right ventricle. Y-descent is attenuated or absent in cardiac tamponade because of resistance to ventricular filling in the early diastole.

Other options:

Options B and C: Restrictive cardiomyopathy and constrictive pericarditis can have deep y descent due to rapid filling of the right ventricle during early diastole.

Option D: In patients with tricuspid regurgitation, increased blood flow to the right atria through the incompetent tricuspid valve during ventricular systole results in a prominent v wave. When the tricuspid valve opens during ventricular diastole, this increased blood in the right atria enters rapidly into the right ventricle during early diastole resulting in deep y-descent.

Solution to Question 8:

Tigroid white matter on MR imaging is seen in Pelizaeus-Merzbacher disease.

It is a rare, X-linked recessive disorder characterized by central nervous system dysmyelination.

The proteolipid protein gene (PLP1) is affected, leading to the failure of axon myelination by oligodendrocytes in the central nervous system.

Clinical Features:

• Nystagmus

• Developmental delay

• Psychomotor delay

• Seizure

• Stridor

• Feeding difficulties

• Ataxia

• Hypotonia progressing to spasticity.

The degree of dysmyelination is correlated with the severity of the clinical manifestations.

Genetic analysis can confirm the diagnosis. MRI shows a lack of myelination and MR spectroscopy reveals aberrant N-acetyl aspartate (NAA) and choline concentrations that reflect axonal and myelination abnormalities.

Unlike other demyelinating leukodystrophies such as metachromatic leukodystrophy and adrenoleukodystrophy, where myelin is formed but subsequently destroyed, PMD shows a failure

of myelin formation.

Solution to Question 9:

Low glucose (<60 mg/dL) is seen in the pleural effusion fluid of a patient with rheumatoid arthritis (RA).

RA causes exudative pleural effusion and fulfills the Light's criteria. High LDH and RA factor are found in the pleural fluid of RA patients. Less commonly, patients with RA may have a cholesterol effusion. It is sterile and has a milky or opaque appearance. The milky appearance is due to the presence of cholesterol crystals and an elevated cholesterol level.

Light's criteria is used to distinguish between exudative and transudative pleural effusions. An exudative pleural effusion satisfies at least one of the following criteria:

• Pleural fluid protein/serum protein > 0.5

• Pleural fluid LDH / serum LDH > 0.6

• Pleural fluid LDH more than 2/3rds upper limit of normal for serum



Solution to Question 10:

The most common pulmonary manifestation of SLE is pleuritis, with or without pleural effusion.

The most common manifestations of different systems involved in SLE are:

• Musculoskeletal- arthritis and arthralgia

• Skin- photosensitivity

• Kidneys- proteinuria

• Neurologic- cognitive impairment

• Cardiac- pericarditis

• Pulmonary- pleuritis

• Hematologic- anemia

Solution to Question 11:

Orthostatic hypotension or postural hypotension is defined as a sustained drop in systolic BP of ≥20 mmHg or diastolic BP of ≥10 mmHg after 3 min of standing.

It can manifest as lightheadedness/dizziness, weakness or tiredness, difficulty thinking/concentrating, blurred vision, and anxiety.

In non-neurogenic causes of orthostatic hypotension such as hypovolemia, the drop in the blood pressure is accompanied by a compensatory increase in heart rate of >15 beats/min. In neurogenic causes, there is aggravation or precipitation of orthostatic hypotension by autonomic stressors like a meal, hot bath, or exercise.

Non-neurogenic causes of orthostatic hypotension include

• Cardiac pump failure - myocardial infarction, myocarditis, aortic stenosis

• Hypovolemia - dehydration, diarrhea, hemorrhage

• Metabolic - adrenocortical insufficiency, pheochromocytoma

• Venous pooling - sepsis, prolonged standing, or recumbency

• Medications - alpha-blockers, diuretics, phenothiazines, tricyclic antidepressants

Treatment includes removing reversible causes like drugs causing hypotension.
Non-pharmacological interventions include patient education regarding staged moves from supine to upright, raising the head of the bed, and increasing dietary fluid and salt. If these fail, drugs like fludrocortisone and midodrine can be given.

Solution to Question 12:

Periodic sharp-wave complexes seen as spikes are seen in Creutzfeldt-Jakob disease (CJD).

It is a spongiform encephalopathy that is characterized by rapidly progressive dementia, myoclonus, akinetic mutism, cerebellar/visual disturbances. Caused by Prions. It is fatal and death occurs usually within a year or less from the onset.

EEG shows characteristic periodic sharp wave complexes (bi/triphasic). CSF assay is positive for 14-3-3 protein.

Four types of Creutzfeldt-Jakob disease have been described:

• Sporadic (sCJD): Accounts for 85-90% of cases

• MRI shows cortical ribbon sign

• Variant (vCJD): mad cow disease, Kuru

• MRI shows hockey stick sign

• Familial (fCJD): 10% of cases (these individuals carry a PRPc mutation)

• Iatrogenic

Option A: Hepatic encephalopathy will show a triphasic wave pattern (negative-positive-negative wave). They are also seen in Hashimoto encephalopathy and CO2 narcosis.

Option C: GTCS (generalized tonic clonic seizures)will show high voltage bursts & high voltage spikes present in all the leads.

Option D: HSV encephalitis usually affects one side temporal lobe showing a characteristic periodic lateralised epiletptiform discharge.

Solution to Question 13:

The anion gap is the difference between measured cations and measured anions and is estimated as:

Anion gap = [Na+] – ([Cl-] + [HCO3-])

Substituting the given values, AG is 22 mmol/L. Normal AG is 6 to 12 mmol/L with an average of 10 mmol/L.

The concentrations of anions and cations in plasma must be equal to maintain electrical neutrality. Therefore, there is no real “anion gap” in the plasma.

However, only certain cations and anions are routinely measured in the clinical laboratory. The cation normally measured is Na+, and the anions are usually Cl– and HCO3–. The value of plasma potassium is omitted from the calculation of the anion gap.

Usually, the unmeasured anions exceed the unmeasured cations.

The anion gap will increase if unmeasured anions rise or if unmeasured cations fall.

The unmeasured anions include albumin, phosphate, sulfate, and other organic anions. The unmeasured cations include calcium, magnesium, and potassium.

Solution to Question 14:

The given ECG showing narrow QRS complex (<0.12 s) tachycardia with absent P wave is suggestive of paroxysmal ventricular tachycardia (PSVT). The next best step in the management of a patient with unstable vitals (systolic blood pressure < 90 mm Hg) is synchronized cardioversion.

Synchronized cardioversion is the treatment of choice in unstable PSVT patients, in which shocks are delivered with the defibrillator/monitor in synchronized mode with an initial energy level of 100J. Synchronization times the delivery so that it is given at the peak of the QRS complex.

If the shock occurs in the ST segment or the T wave (unsynchronized), it can precipitate a more serious arrhythmia such as ventricular fibrillation.

Synchronized cardioversion	Defibrillation
Used inunstable PSVT, atrial flutter, atrial fibrillation, and ventricular tachycardiawith a pulse	Used in ventricular tachycard iawithout pulseand ventricul ar fibrillation.
Shock is delivered in sync wit h the R wave (in absolute refr actory period)	Shock is delivered at any tim e during the cardiac cycle.
Monophasic: 100-200 JBipha sic: 50-100 J	Monophasic: 360 JBiphasic:1 20-200 J

Other options:

Option A: In the absence of hemodynamic instability, a repeat dose of injection adenosine (maximum of 12mg) could be attempted.

Option B: IV Amiodarone is rarely used in the acute management of PSVT. Instead, it is used as an alternative to calcium channel blockers in chronic suppressive therapy.

Option D: Defibrillation is only done for ventricular fibrillation and pulseless ventricular tachycardia.



Solution to Question 15:

The given image showing thickening of tissues in the temporal and orbital regions with coarse overlying skin suggests a diagnosis of plexiform neurofibroma.

Plexiform neurofibroma represents an uncommon variant (30%) of neurofibromatosis 1 (NF1). Plexiform neurofibromas are usually evident at birth and result from a diffuse thickening of nerve trunks and surrounding tissues that are usually located in the orbital or temporal region of the face. The skin overlying it may be coarse and associated with hyperpigmentation.

Plexiform neurofibromas may produce an overgrowth of an extremity and a deformity of the corresponding bone.

NF1 is clinically diagnosed when any 2 of the following 7 features are present:

• Six or more café-au-lait macules >5mm in prepubertal and >15mm in postpubertal

• Axillary or inguinal freckling

• Two or more iris lisch nodules

• Two or more neurofibromas or one plexiform neurofibroma

• A distinctive osseous lesion such as sphenoid dysplasia or cortical thinning of long bones with or without pseudoarthrosis.

• Optic gliomas

• A first-degree relative with NF-1

Solution to Question 16:

The most common presentation of inhalation anthrax is hemorrhagic mediastinitis, which is seen in 38% of the patients.

Inhalational anthrax also known as wool sorter's disease is caused by Bacillus anthracis. Infection usually occurs among workers handling contaminated animal wool, hair, and skin in textile and tanning industries.

Patients present with malaise, fever, cough, shortness of breath & headache. B. anthracis spores are phagocytosed by alveolar macrophages and transported to mediastinal lymph nodes. Hemorrhagic necrosis of the thoracic lymph nodes draining the lung results in hemorrhagic mediastinitis. Pulmonary infiltrates, mediastinal widening, and pleural effusion are other findings.

Treatment of inhalation anthrax includes a combination of intravenous ciprofloxacin and clindamycin for at least 14 days, antitoxins (Raxibacumab, Obiltoxaximab), anthrax immunoglobulins and steroids (if associated meningitis is present).

Prophylaxis with ciprofloxacin (500mg twice daily for 2 months) is recommended for anyone at high risk of inhalational exposure to anthrax spores.

Solution to Question 17:

Among the given options, the metabolic abnormality seen in a patient who has received multiple blood transfusions is metabolic alkalosis.

This is because the excess sodium citrate (anticoagulant) in the transfused blood gets converted to bicarbonate leading to metabolic alkalosis. Each mmol of citrate generates 3 mEq of bicarbonate (a total of 23 mEq of bicarbonate due to each unit of blood).

Metabolic acidosis can also be seen in multiple blood transfusions as citrate is acidic, but metabolic alkalosis is more common.

Solution to Question 18:

The likely metabolic abnormality in this patient is high anion gap metabolic acidosis with metabolic alkalosis.

Patients with renal failure may have metabolic acidosis and associated vomiting in this patient may cause metabolic alkalosis. However, the parameters of arterial blood gas analysis appear to be normal. This disparity raises the suspicion of a mixed acid-base disorder.

When metabolic acidosis and metabolic alkalosis coexist in the same patient, pH may be in the normal range. In such cases, the presence of an elevated anion gap denotes the presence of metabolic acidosis.

The anion gap (AG) is calculated as:

AG = [Na+] – ([Cl-] + [HCO3-])

= 145-(100+24)

= 21

AG is high, which points to high AG metabolic acidosis.

In high anion gap metabolic acidosis, excess H+ ions are buffered by bicarbonate and the bicarbonate level falls. The fall in HCO3- is expected to be equal to the rise in anion gap. The difference between the rise in anion gap and the fall in HCO3- (delta gap) is zero in simple HAGMA.

Delta gap = (Actual anion gap - Normal anion gap) - (Normal bicarbonate - Actual bicarbonate)

Associated vomiting in this patient results in loss of acid and hence the fall of bicarbonate level is less compared to the rise in the anion gap. A delta gap of more than +6 indicates the presence of additional metabolic alkalosis.

In this case, the delta gap is 10, which points to metabolic alkalosis in addition to high AG metabolic acidosis.

Solution to Question 19:

The given clinical scenario describes a woman presenting with hypopituitarism following trans-sphenoidal resection. Mineralocorticoid supplementation is not needed postoperatively as it is regulated by the renin-angiotensin-aldosterone system and not by ACTH. Therefore, mineralocorticoid levels would remain intact despite ACTH deficiency.

Thyroid hormone, estradiol, and glucocorticoid supplementation are needed in this patient as their secretion is regulated by anterior pituitary hormones.

Features of hypopituitarism include growth abnormalities in children with a growth hormone deficiency, insufficient levels of gonadotropins leading to menstrual irregularities, infertility in women and reduced sexual function, infertility and loss of secondary sexual characteristics in men. TSH deficiency leads to stunted growth in children and features of hypothyroidism in both

children and adults. ACTH deficiency leads to secondary adrenal insufficiency.

Polyuria and polydipsia may be seen and indicate a loss of vasopressin secretion in posterior pituitary lesions.

The following hormones must be replaced post-hypophysectomy:

Trophic hormone deficit	Hormone Replacement
ACTH	Hydrocortisone, cortisone acetate, prednisone
TSH	L-thyroxine (75–150 μg daily)
FSH/LH	Males: TestosteroneFemales: For fertility: menopausal gonadotropins, hCGPost-menopausal women: Conjugated estrogen, progesterone
GH	Somatotropin
Vasopressin	Intranasal desmopressin

Solution to Question 20:

The serological marker marked in the image which becomes detectable early in the infection and remains elevated for a long period after recovery from the infection is an anti-HBc antibody.

Anti-HBc is demonstrable in serum within the first 1–2 weeks after the appearance of HBsAg. It precedes detectable levels of anti-HBs by weeks to months.

IgM anti-HBcAg antibody is seen early in the infection for about 6 months and indicates a recent infection. IgG anti-HBcAg antibody is seen later, after 6 months, and indicates remote infection.

Solution to Question 21:

The given ECG showing a wide QRS complex tachycardia with waxing and waning QRS amplitude points to the diagnosis of torsades des pointes. As the patient is hemodynamically stable, intravenous $MgSO_4$ is the treatment of choice.

Polymorphic ventricular tachycardia (VT) has a continually changing QRS morphology. Polymorphic VT that occurs in individuals with congenital or acquired prolongation of QT interval is referred to as torsades des pointes. Patients present with near-syncope, syncope, or cardiac arrest.

IV $MgSO_4$ is highly effective for both treatment and the prevention of recurrence of long QT-related ventricular ectopic beats that trigger torsades des pointes.

If magnesium alone is ineffective, isoproterenol infusion or pacing, to a rate of 100-120 depolarization/minute usually suppresses VT recurrences.

If the patient is hemodynamically unstable, then immediate defibrillation should be done.

Synchronized cardioversion	Defibrillation
Used inunstable PSVT, atrial flutter, atrial fibrillation, and ventricular tachycardiawith a pulse	Used in ventricular tachycard iawithout a pulse, and ventric ular fibrillation.
Shock is delivered in sync wit h the R wave (in absolute refr actory period)	Shock is delivered at any tim e during the cardiac cycle.
Monophasic - 100 to 200J	Monophasic - 360J
Biphasic - 50 to 100J	Biphasic - 120 to 200J

Solution to Question 22:

The most common cause of infection post-solid organ transplantation in India is cytomegalovirus(CMV). Epstein-Barr virus (EBV), Herpes, and varicella are also known to cause infection post-solid organ transplantation but to a lesser extent.

CMV disease in recipients can present with fever, leukopenia, thrombocytopenia, malaise, and arthralgias. The diagnosis of CMV infection is established by nucleic acid testing (NAT) using the polymerase chain reaction (PCR).

Management of this includes use of anti-viral drugs like ganciclovir, valganciclovir, foscarnet, cidofovir and intravenous immunoglobulins.

Solution to Question 23:

Primary adrenal insufficiency, also known as Addison's disease, is most commonly caused by autoimmune adrenalitis.

Tuberculous adrenalitis is still a frequent cause of primary adrenal insufficiency in developing countries like India. It is characterized by the loss of glucocorticoid, mineralocorticoid, and adrenal androgen secretion.

Glucocorticoid deficiency results in fatigue, lack of energy, weight loss, fever, myalgia, normochromic anemia, and hypoglycemia. Mineralocorticoid deficiency is characterized by nausea, vomiting, low blood pressure, postural hypotension, salt craving, hyponatremia, and hyperkalemia. Adrenal androgen deficiency manifests as a lack of energy, dry itchy skin, loss of libido, and loss of axillary and pubic hair (in women).

Hyperpigmentation is seen in primary adrenal insufficiency due to excess proopiomelanocortin-derived peptides. Hyperpigmentation is not seen in secondary adrenal insufficiency.

The diagnosis of adrenal insufficiency can be made with the short cosyntropin test (cortisol level <450-500nmol/L sampled 30 to 60 minutes after ACTH stimulation is diagnostic of adrenal

insufficiency). Plasma ACTH, plasma renin, and serum aldosterone can help differentiate primary from secondary causes. Primary adrenal insufficiency will show elevated ACTH and a compensatory elevation in renin levels due to decreased aldosterone in the body. In secondary adrenal insufficiency, ACTH is low with normal renin and aldosterone.

Treatment includes glucocorticoid, mineralocorticoid, and adrenal androgen replacement.

Other options:

Option B: Patients with adrenocortical carcinoma present with features of glucocorticoid and adrenal androgen excess, and not adrenal insufficiency.

Options C and D: Hypernephroma and medullary carcinoma thyroid are not associated with Addison's disease.

Solution to Question 24:

In patients with hypercalcemia due to vitamin D toxicity, glucocorticoids like dexamethasone are the preferred therapy as they decrease 1,25 (OH2) vitamin D (calcitriol) production. This results in reduced intestinal calcium absorption and increased urinary excretion of calcium.

Hydroxychloroquine, chloroquine, and ketoconazole may also decrease calcitriol production and are used occasionally. But the preferred therapy is steroids.

Vitamin D toxicity presents acutely with features of hypercalcemia such as confusion, polyuria, polydipsia, anorexia, vomiting, and muscle weakness. Chronic intoxication may cause nephrocalcinosis, bone demineralization, and pain.

Lab features include elevated 25 (OH) vitamin D, high calcium, and suppressed PTH.

Dexamethasone can also be used in the treatment of hypercalcemia caused by the humoral hypercalcemia of malignancy (due to PTHr) and sarcoidosis.

Medicine AIIMS 2020 (May and Nov INI-CET)

Question 1:

Which of the following is the drug of choice for hairy cell leukemia?

a) Rituximab

b) Vemurafenib

c) Cladribine

d) Interferon-alpha

Question 2:

An intubated patient in the ICU had a chest X-ray taken to evaluate for fever spikes over the last 24-48 hours. The radiograph is shown below. What is the next best step in the management of this patient?



a) Suction the ET tube

b) Chest physiotherapy twice daily

c) Start antibiotics and send samples for culture

d) Ultrasound-guided aspiration

Question 3:

Following is a graphic representation of a patient admitted to the medicine ward with fever. What could be the possible diagnosis?



©MARROW

a) Cerebral malaria

b) Brucellosis

c) Leptospirosis

d) Typhoid

Question 4:

The laboratory investigations of a 65 year old woman revealed a hemoglobin level of 5.4 g/dl, creatinine level of 6 mg/dl, calcium level of 12 mg/dl, potassium level of 4.4 mmol/L. Her urine output in the last 24 hours was 50 ml. Bone marrow aspiration revealed more than 60% of plasma cells. What would be the next appropriate line of management for this patient?

a) Urgent consultation for dialysis

b) Lenalidomide + Bortezomib + Dexamethasone

c) Dexamethasone + fluids

d) Chemotherapy followed by autologous stem cell transplantation

Question 5:

A diabetic nephropathy patient is an ideal candidate for renal graft transplant. Which of the following statements is true?

a) The survival rate of graft is 95% in the first year

b) The transplantation is cost effective after the second transplant year

c) The life expectancy is doubled in a diabetic patient with renal transplant

d) The treatment of chronic rejection has improved over the last 10 years

Question 6:

A diabetic patient enters the OPD and greets you with a namaste. As the doctor, you notice a deformity in his hand which is shown in the image below. Based on your observation, what is the likely cause of this deformity?

a) Flexor tenosynovitis

b) Cheiroarthropathy

c) Dupuytrens contracture

d) Ankyloses

Question 7:

Dialysis will prevent all except?

a) Seizures

b) Peripheral neuropathy

c) Metabolic acidosis

d) Uremic pericarditis

Question 8:

Which of the following is not a treatment option in Guillain Barre syndrome?

a) Steroids

b) Plasmapheresis

c) IVIG

d) Ventilator support

Question 9:

Which part of the internal capsule is affected in hemiplegia due to ischaemic stroke?

a) Retrolentiform

b) Sublentiform

c) Anterior limb

d) Posterior limb

Question 10:

Which of the following is false about mycoplasma pneumonia?

a) Responds well to amoxiclav

b) Antibody are useful in diagnosis

c) Chest X-ray shows bilateral infiltrates

d) Can be cultured in a cell-free medium

Question 11:

A 50-year-old chronic smoker presented with mild recurrent hemoptysis without fever or weight loss. The chest X-ray appears normal and the sputum is negative for AFB. Which of the following is the next best diagnostic investigation?

a) MRI

b) CT

c) Fiberoptic bronchoscopy

d) Bronchography

Question 12:

Which of the factors would suggest a risk for a patient with TIA developing stroke in the future according to ABCD2 scoring?

a) Age < 60 years

b) SBP >130 mm Hg and DBP > 80 mm Hg

c) Duration of TIA >5 mins

d) Diabetes

Question 13:

The serum electrolyte values of a patient weighing 60kg are given below. Calculate the sodium deficit in the patient. pH: 7.42, Na+: 120 mEq/L, serum K+: 4 mEq/L, serum Cl-: 90 mEq/L.

a) 20 mEq

b) 200 mEq

c) 400 mEq

d) 720 mEq

Question 14:

In a patient with non-ST-elevation myocardial infarction (NSTEMI), all the following can be used for treatment except?

a) Aspirin

b) Clopidogrel

c) Streptokinase

d) Prasugrel

Question 15:

A patient presents with vertigo, diplopia, hoarseness, dysphagia, and left Horner's syndrome associated with numbness of the left face and right side limbs. Which artery is affected in this

patient?

a) Posterior inferior cerebellar artery

b) Superior cerebellar artery

c) Anterior inferior cerebellar artery

d) Basillar artery

Question 16:

Resistant hypertension is defined as?

a) BP >160/90 mmHg with one anti-hypertensive

b) BP >140/90 mmHg with one anti-hypertensive

c) BP >160/90 mmHg with three or more anti-hypertensives

d) BP >140/90 mmHg with three or more anti-hypertensives

Question 17:

Components of the MELD (2016) score are all except?

a) Bilirubin

b) INR

c) Albumin

d) Creatinine

Question 18:

A woman with Grave's disease was taking antithyroid medication throughout her pregnancy. The baby was born with aplasia cutis congenita. Which drug could have caused this condition?

a) Carbimazole

b) Levothyroxine

c) Methylthiouracil

d) Hydrouracil

Question 19:

In the given acid-base nomogram diagram, areas marked A and D indicate which of the following conditions?



a) Chronic respiratory acidosis, acute metabolic alkalosis

b) Acute Metabolic alkalosis, acute respiratory acidosis

c) Acute respiratory acidosis, acute respiratory alkalosis

d) Acute respiratory acidosis, acute metabolic alkalosis

Question 20:

A patient presented with the complaints of chest pain. Which among the following is the best investigation to diagnose MI within 2-4 hours?

a) LDH

b) CKMB

c) Troponin-T

d) BNP

Question 21:

A patient comes with palpitation to ER. His Pulse-180/min and BP-70mm of Hg systolic. ECG shows narrow QRS complex tachycardia with regular heart rate. What is the next step in management?

a) DC Cardioversion

b) Adenosine

c) Valsalva Maneuver

d) Verapamil

Question 22:

High anion gap acidosis with low bicarbonate levels is seen in all the following conditions except:

a) Renal tubular acidosis

b) Diabetic ketoacidosis

c) Lactic acidosis

d) Salicylate poisoning

Question 23:

A patient comes to the hospital with chest pain for the last 3 hours. Vitals are stable. ECG revealed ST depression and T wave inversion in anterior chest leads. Which of the following would be the most appropriate line of management in this patient?

a) PCI

b) Thrombolysis with alteplase

c) Prophylaxis for arrhythmia

d) Aspirin with heparin

Question 24:

Which among the following is the most life-threatening manifestation of CKD?

a) Decreased GFR

b) Increased creatinine

c) Hyperkalemia

d) Proteinuria

Question 25:

A patient presented with massive abdominal swelling. Paracentesis fluid was sent for lab analysis and results show serum protein to be 3g/dL and ascitic fluid protein to be 1g/dL. What is the likely diagnosis?

a) Congestive heart failure

b) Early Budd Chiari syndrome

c) Late Budd Chiari syndrome

d) Tuberculosis

Question 26:

Iritis is seen in all except:

a) SLE

b) Behcet's disease

c) Ulcerative colitis

d) Rheumatoid arthritis

Question 27:

A 13-year-old boy collapsed while playing football. ECG showed V5, V6 inversion and prominent inverted T-waves. 2D echocardiography showed systolic anterior motion of the mitral valve leaflet. Which of the following features will be seen on clinical examination and which of the following are true regarding the management?

a) 1 & 3, a, c, d

b) 2 & 4, b, c, e

c) 1 & 3, a, d

d) 2 & 4, b, e

Question 28:

You are posted as an intern in casualty. An elderly 60-year-old man is brought to you with complaints of chest pain. He has tested positive for COVID. Which of the following features will prompt you to get this patient admitted to dedicated COVID Health Centre?

a) A & B only

b) A, B, C, & E only

c) A, B & C only

d) A, B, C, D & E

Question 29:

Correct sequence after p wave: A- 'a' wave B- 1st heart sound C- Rapid filling of ventricles D- 't' wave

a) ABCD

b) ABDC

c) CABD

d) ACBD

Question 30:

A 23-year-old medical student diagnosed with dengue developed epistaxis that resolved spontaneously. He has been afebrile for the last 24 hours, and his platelet count is 14,000/µL. His BP is 110/70 mmHg and there is a 28% rise in hematocrit. What should be the management?

a) 1 and 2

b) 1, 2, and 4

c) 2, 3, and 4

d) 2 and 5

Question 31:

Which of the following are true regarding KF ring?

a) 1, 2 and 5 only

b) 1 and 5 only

c) 1, 2, 3, 4 and 5

d) 2, 3, 4 and 5 only

Answer Key

Question No.	Correct Option
1	c
2	d
3	c
4	c
5	b
6	b
7	b
8	a
9	d
10	a
11	b
12	d
13	d
14	c
15	a
16	d
17	c
18	a
19	c
20	c
21	a
22	a
23	d
24	c
25	c
26	d
27	a
28	b
29	b
30	c
31	b

Detailed Explanations

Solution to Question 1:

Cladribine is the drug of choice in hairy cell leukemia.

Cladribine is a purine analogue with potent and curative activity against hairy cell leukemia.

Other treatment options for hairy cell leukemia include deoxycoformycin, interferon-alpha, and splenectomy. Chemotherapy-refractory patients are treated with vemurafenib, a BRAF kinase inhibitor. The addition of rituximab results in more durable remissions.

Cladribine is also used to treat CLL, low-grade lymphomas, Langerhans cell histiocytosis, cutaneous T-cell lymphomas, and Waldenström macroglobulinemia.

The major dose-limiting toxicity of cladribine is myelosuppression.

Solution to Question 2:

The given clinical history of an intubated ICU patient and the X-ray features point toward a diagnosis of pleural effusion. Ultrasound-guided aspiration is the next step in the management of the patient. Antibiotics should be started once the pleural tap has been done and infective etiology is suspected.

Options A and B: ET tube suctioning and chest physiotherapy are done to remove thick secretions out of the upper and lower respiratory tract. It wouldn't help in removing the fluid of a pleural effusion.

Pleural effusion is a common finding in the ICU setting and in intubated patients. Many factors contribute to the impairment of pleural fluid turnover in such cases as prolonged recumbency, ventilator-produced positive pressure, and reduced plasma oncotic pressure.

The diagnosis of pleural effusion may be made by clinical examination (dull note on percussion) or by imaging. A pleural tap for confirmation of diagnosis and to relieve breathing difficulty is now done under ultrasound guidance.

The CXR features of pleural effusion include:

• Obliteration of costophrenic angle on the side of pleural effusion

• Air-fluid interface (meniscus sign)

• Large effusions show: Complete opacification of the lung, mediastinal shift, and tracheal deviation away from the effusion

Differential diagnosis in an intubated patient with opacities on CXR might be pneumonitis secondary to gastric acid or food aspiration. But the X-ray would show patchy infiltration unlike the homogenous density seen in pleural effusion.

Solution to Question 3:

The given graphical representation of a biphasic illness (acute and convalescent phases) and relevant investigations at different stages of the disease is consistent with a diagnosis of leptospirosis.

Malaria (Option A) is commonly diagnosed by demonstrating malarial parasites in a peripheral blood smear and not in CSF or urine.

Brucellosis (Option B) follows a characteristic pattern of undulating fever.

The mean incubation period in typhoid (Option D) usually ranges from 10 to 14 days and diagnosed based on examination of blood or stool.

Leptospirosis is a zoonotic disease caused by gram-negative Leptospira bacteria (especially L. interrogans). Direct transmission to humans occurs when broken skin and mucous membranes come into contact with the urine of infected reservoir hosts such as rodents.

Illness is classically biphasic after an incubation period of up to 1 month, though the distinction between the first and second phases is not always clear.

• Acute leptospiremic phase-organism can be cultured from blood (within 1 week of onset of symptoms)

• Resolving immune phase-organism can be cultured from urine (after 1 week)

Most cases are characterized by non-specific symptoms (e.g fever, headache, myalgia and conjunctival suffusion) and resolve spontaneously within a week but a few may progress to a severe form known as Weil's disease which typically presents with a triad of hemorrhage, jaundice, and acute kidney injury.

Diagnosis includes:

• Dark-field microscopy of urine or blood samples

• Serological tests

• Four-fold rise in the level of IgM titers within one month of the onset of symptoms

• Microscopic agglutination test (MAT)

• PCR

• Culture on EMJH or Fletchers media

Treatment:

• Mild disease: Oral doxycycline or azithromycin or ampicillin/ amoxicillin

• Severe disease: IV penicillin

Solution to Question 4:

The clinical scenario shows a patient with a bone marrow finding of plasma cells >60% which points to a diagnosis of multiple myeloma.

Anemia (Hb- 5.4 g/dl), hypercalcemia (serum calcium-12 mg/dl) and acute kidney injury (creatinine-6 mg/dl, reduced urine output) are myeloma-defining events suggesting end-organ damage.

The most appropriate next step in managing this patient would be to administer fluids to treat the acute kidney injury and hypercalcemia (hypercalcemia is the most common cause of renal failure in multiple myeloma), and prevent further renal impairment. Therefore, among the given options, dexamethasone and fuids would be the best answer.

Multiple myeloma is a malignant plasma cell dyscrasia and its diagnosis requires bone marrow biopsy finding of ≥ 10% clonal bone marrow plasma cells, a serum/urine M component, and at least one of the myeloma-defining events (hypercalcemia, renal insufficiency, anemia and bone lesions).

Dialysis (Option A) is indicated in the following conditions:

• Pulmonary edema not responding to diuretics

• Uremic pericarditis, uremic encephalopathy or uremic syndrome

• Metabolic acidosis not responding to treatment

• Hyperkalemia

Lenalidomide + Bortezomib + Dexamethasone (Option B) is the preferred treatment in multiple myeloma and can be initiated after initial management of acute kidney injury.

Chemotherapy followed by autologous stem cell transplantation (Option D) in transplant-eligible patients prolongs overall survival in multiple myeloma patients.



Solution to Question 5:

Transplantation becomes cost-effective after the second transplant year in a patient with diabetic nephropathy, provided there is no chronic rejection.

The cost of transplantation after the second year becomes lesser compared to thrice-weekly dialysis.

Other options:

Options A: Graft survival in the first year is not 95%. It is around 80%

Option C: The life expectancy is not doubled in a patient with diabetic nephropathy post-transplantation

Option D: As chronic rejection is due to so many variables, the treatment is very difficult. Risk factors that predispose a recipient to long-term graft rejection include HLA-mismatching, acute episodes of graft rejection, mismatch in donor-recipient age, and smoking. Better treatment options are still under research.

Solution to Question 6:

The image shows a positive namaste or prayer sign where the patient is unable to touch the palmar aspects of both hands and fingers together which is suggestive of diabetic cheiroarthropathy.

Diabetic cheiroarthropathy also known as diabetic stiff hand syndrome is a cutaneous condition characterized by waxy, thickened skin and limited joint mobility of the hands and fingers leading to flexion contractures (namaste or prayer sign). This occurs due to the non-enzymatic glycosylation of collagen. It is seen in both diabetes type 1 and type 2. Its prevalence increases with age and poor glycemic control.

Other options:

Option A: Flexor tenosynovitis refers to inflammation of the sheath surrounding the flexor tendons in the hand. Infectious flexor tenosynovitis is associated with the Kanavel's signs which include:

• Fusiform swelling along the affected finger

• Tenderness with palpation along the tendon sheath

• Flexed posture of the affected finger

• Pain with passive extension

Option C: Dupuytren's contracture is a fibroproliferative disorder affecting the palmar aponeurosis of mainly the 4th and 5th fingers. Skin puckering proximal to the flexor crease is the earliest sign, followed by the development of nodules and cords in the palmar fascia, leading to flexion contractures at the MCP and PIP joints as the disease progresses.

Option D: Ankylosis refers to the fusion of a joint, resulting in limited or no movement. It can occur due to various causes such as rheumatoid arthritis, osteoarthritis, trauma, or infection.

Solution to Question 7:

Dialysis does not prevent peripheral neuropathy. Complications like seizures, metabolic acidosis, and uremic pericarditis can be prevented by dialysis.

Peripheral neuropathy is postulated to be caused due to middle molecules (50-200kDa). Conventionally used dialysis membranes have pore sizes that prevent larger particles (such as middle molecules) from passing through them.

Suggested methods for the removal of middle molecules include:

• High flux dialysis

• Protein adsorptive membranes

• High-cutoff membranes

Aggressive removal of middle molecules or larger low-molecular-weight proteins (LMWPs) has been a growing concern since it causes massive albumin leakage.

β2-microglobulin (B2M),α1-microglobulin (A1M) and albumin leakage during a dialysis session are useful parameters for assessing middle-molecule removal.

Renal transplantation is the best treatment option available for peripheral neuropathy in diabetes.

Solution to Question 8:

Steroids are not used in the treatment of Guillain Barre syndrome (GBS).

GBS is an acute, fulminant, ascending polyradiculoneuropathy that is autoimmune in nature. GBS is caused when immune response/antibodies against external antigens like infections, vaccines, etc target similar proteins present in the person's nervous tissue. Hence the preferred treatment is to neutralize these autoantibodies using IVIg (I.V. immunoglobulins) or plasmapheresis (Option B and Option C).

GBS manifests as rapidly evolving ascending areflexic motor paralysis with or without sensory disturbance. GBS paralysis can also affect the muscles of respiration, possibly leading to death due to respiratory failure. In such cases, ventilatory support (Option D) is needed.

Note: Steroid therapy is used in Chronic inflammatory demyelinating polyneuropathy.

Solution to Question 9:

The posterior limb of the internal capsule is affected in hemiplegia due to ischaemic stroke.

Motor fibers (cortico-spinal tract) supplying the extremities are present in the posterior limb, hence a lesion in this region may cause hemiplegia.



Solution to Question 10:

Mycoplasma does not respond to amoxicillin-clavulanate (amoxiclav).

Mycoplasma is a bacteria that lacks a cell wall. Since beta-lactam antibiotics (amoxiclav) act on the cell wall of the bacterium, they can not be used in the treatment of mycoplasma infection.

The diagnosis of mycoplasma pneumonia (also known as walking pneumonia/ primary atypical pneumonia) can be established by:

• PCR detection of M. pneumoniae in respiratory secretions

• Serological tests for IgM and IgG antibodies to M.pneumoniae in paired serum samples

• Culture: Mycoplasma can be cultured on PPLO agar, a cell-free medium, and shows a typical fried egg appearance

Chest X-ray usually shows B/L peribronchial and perivascular interstitial infiltrates

Treatment options for mycoplasma pneumonia include:

- Macrolides (e.g. azithromycin)
- Tetracyclines (e.g. doxycycline)
- Respiratory fluoroquinolones. (e.g. levofloxacin)

Azithromycin is the drug of choice for the management of mycoplasma pneumonia.

Solution to Question 11:

In this case, considering the patient's age of 50 years, history of chronic smoking, and recurrent hemoptysis, all of which raise suspicion for lung malignancy, the next best diagnostic investigation would be a CT scan.

Any patient presenting with recurrent hemoptysis requires chest imaging. The initial imaging modality is a plain radiograph. However, patients with risk factors for malignancy need further evaluation with a CT scan.

CT can identify masses, bronchiectasis, and parenchymal lesions. Further management depends on the CT findings. Fiberoptic bronchoscopy, which is an invasive procedure, is done after a mass is seen on a CT scan to enable sample collection for biopsy. Also, bronchoscopy has an overall lower sensitivity than MDCT in detecting the underlying causes of bleeding.

Other options:

Option A: While MRI can be valuable in certain situations, such as evaluating the mediastinum or vascular structures, a chest CT scan is generally the preferred initial diagnostic investigation for assessing lung pathology, especially in cases of hemoptysis.

Option D: Bronchography is visualizing the tracheobronchial tree on plain radiography by instilling contrast material via the bronchus. It has been replaced by advancements in CT scans.

Risk factors for the development of lung cancer:

• Tobacco smoking: Associated with 90% of lung cancers.

• Occupational and environmental exposure to carcinogens: Includes passive smoking, environmental air pollution, radon, asbestos, arsenic, chromium, nickel, beryllium, and silica.

• Family history (genetic predisposition).

• Other risk factors: Pulmonary scarring, previous radiation, pulmonary fibrosis, and chronic infections (e.g., tuberculosis, HIV).

Solution to Question 12:

According to the ABCD2 scoring system, which is commonly used to assess the short-term risk of stroke following a transient ischemic attack (TIA), diabetes would suggest a higher risk factor.

Age more than 60 years, SBP more than 140 or DBP more than 90 mmHg, and duration of TIA of 10-59 minutes or more than 60 minutes are other criteria included in the ABCD2 scoring system.

TIA is defined as episodes of stroke symptoms that last for only a brief period of time (<24 hours). The ABCD2 scoring system is most frequently used to calculate the short-term risk of stroke after TIA.

Solution to Question 13:

The sodium deficit is calculated as (target sodium concentration-current sodium concentration) x 0.6 x Body weight (kg).

Substituting the values, sodium deficit = (140-120) x 0.6 x 60 = 720 mEq.

To calculate the sodium deficit in the patient, we first need to determine the target sodium concentration and compare it to the current sodium concentration.

The target sodium concentration can be calculated using the following formula:

Target Na concentration = (Lower limit of serum sodium concentration + upper limit of serum sodium concentration) / 2

In this case, we assume the lower limit for the serum sodium concentration is 135 mEq/L, and the upper limit is 145 mEq/L.

Target Na concentration = (135 + 145) / 2= 140 mEq/L

Next, the current sodium deficit can be calculated using the following formula:

Sodium Deficit = (Target sodium concentration-current sodium concentration) x Total Body Water (TBW)

Total Body Water (TBW) = Weight (in kg) × 0.6 (assuming 60% of body weight is water); TBW = 60 × 0.6= 36 liters

Sodium Deficit = (140 - 120) x 36 = 20 x 36 = 720 mEq

Therefore, the sodium deficit in the patient is 720 mEq.

Rapid sodium correction of 4–6 mEq/L should be achieved in the first 6 hours of therapy for patients with acute hyponatremia.

The rate of correction should be comparatively slow in chronic hyponatremia: <6-8 mEq/L in the first 24 h and <6 mEq/L each subsequent 24h, so as to avoid osmotic demyelination syndrome.

Solution to Question 14:

Streptokinase is not used in the management of non-ST segment elevation MI (NSTEMI). It is a fibrin non-specific plasminogen activator used as a fibrinolytic agent in the treatment of ST-segment elevation MI (STEMI). Aspirin, clopidogrel, and prasugrel are antiplatelet drugs commonly used in the management of NSTEMI.

Fibrinolytic therapy is not indicated in patients with unstable angina or NSTEMI, unlike in STEMI. This is because STEMI is almost always caused by plaque rupture leading to the formation of an occluding thrombus, whereas, NSTEMI can be caused by a stable non-occluding plaque with some amount of lumen still patent. Fibrinolytic therapy may lead to the rupture of the stable plaque causing more damage.

Acute coronary syndrome is classified into 2 groups:

• ST-segment Elevation Myocardial Infarction (STEMI)

• NSTE-ACS

NSTE-ACS further includes:

• NSTEMI (myocyte necrosis present and cardiac troponins elevated)

• Unstable Angina (myocyte necrosis absent and cardiac troponins not elevated)

The treatment of NSTEMI is as follows:

• Admission to a specialized cardiac unit

• Inhaled oxygen

• Bed rest with continuous ECG monitoring

• Monitoring cardiac biomarkers

• Morphine or opioid analgesics

• Nitrates

• Beta-blockers

• Antiplatelet drugs like aspirin, clopidogrel, prasugrel

Solution to Question 15:

These given features collectively indicate a diagnosis of Wallenberg or lateral medullary syndrome, which is typically caused by a lesion of the posterior inferior cerebellar artery.

The patient presents with hoarseness and dysphagia, suggesting the involvement of cranial nerves IX and X, which helps localize the lesion to the medulla. Additionally, the presence of vertigo and diplopia (involving the vestibular nerve), left-sided Horner's syndrome (involving the descending sympathetic tract), numbness on the left side of the face (involving the sensory tract of trigeminal nucleus), and numbness in the right side limbs (involving the lateral spinothalamic tract) further aids in localizing the lesion to the lateral part of the medulla.

The absence of motor symptoms suggests that the pyramidal tracts in the medial aspect of the medulla are not involved, ruling out medial medullary syndrome.

Wallenberg/lateral medullary syndrome affects:

• Descending tract and nucleus of CN V- loss of pain and sensation over the ipsilateral half of the face.

• Inferior cerebellar peduncle – ipsilateral ataxia

• Lateral spinothalamic tract – loss of pain and temperature sensation over the contralateral half of the body.

• CN IX and X - ipsilateral paralysis of palate, vocal cord, diminished gag reflex giving rise to dysphagia, dysarthria.

• Vestibular nucleus - Nystagmus, diplopia, vertigo, nausea

• Descending sympathetic tract – ipsilateral Horner's syndrome (miosis, ptosis, and anhidrosis).



Solution to Question 16:

Resistant hypertension is defined as blood pressure being persistently >140/90 mmHg despite taking three or more antihypertensive agents, including a diuretic.

Resistant hypertension is more common in patients aged >60 years than in younger patients.

Resistant hypertension may be related to pseudohypertension (high office blood pressure and lower home blood pressure measurements), nonadherence to therapy, and secondary causes of

hypertension including obesity, excessive alcohol intake, and primary aldosteronism.

Solution to Question 17:

Albumin is not a component of the MELD (2016) score.

The Model for End-Stage Liver Disease (MELD) is used as an indicator of short-term survival in patients with end-stage liver disease and to determine organ allocation priorities for liver transplantation. The components of the original MELD model were:

• Serum creatinine

• Serum bilirubin

• INR (international normalized ratio)

In 2016, it was modified to incorporate serum sodium (the MELD-Na score).

The latest MELD 3.0 scoring system has added two additional components:

• Serum albumin

• Female gender

Patients with high scores have poor prognoses without intervention and should therefore be prioritized for transplantation if eligible.

Solution to Question 18:

Carbimazole intake during the first trimester of pregnancy is associated with aplasia cutis congenita.

Carbimazole and methimazole are associated with teratogenic effects such as:

• Aplasia cutis congenita: Congenital absence of scalp hair and skin along with calvarial defects

• Choanal atresia

• Tracheoesophageal fistula

• Fetal goiter

The drug of choice for the treatment of hyperthyroidism during the first trimester of pregnancy is propylthiouracil (PTU). It is because it has lower teratogenic effects than carbimazole or methimazole.

However, propylthiouracil use is limited to the first trimester due to the risk of hepatotoxicity. It is replaced with methimazole (or carbimazole) in the second and third trimesters of the pregnancy.

Radioiodine therapy is contraindicated during pregnancy and lactation.

Solution to Question 19:

In the given illustrated acid-base nomogram, the letters A and D indicate acute respiratory acidosis and acute respiratory alkalosis.

An acid-base nomogram, also known as a Davenport diagram, is a graphical representation of acid-base balance to interpret arterial blood gas (ABG) results. It helps determine the primary acid-base disorder and identify compensatory mechanisms.



The y-axis of the acid-base nomogram represents the arterial bicarbonate (HCO3-) levels, and the x-axis represents the arterial pH values. The shaded areas are the range of values of the normal respiratory and metabolic compensations for primary acid-base disturbances.

Solution to Question 20:

Measurement of troponin is the best investigation to diagnose MI within 2-4 hours.



Troponin T/I:

• Highly specific for MI

• Troponin I is more important than troponin T.

• Troponins will remain elevated longer than creatine kinase (CK), for up to 14 days.

• This makes troponins a superior marker for diagnosing MI in the recent past (late diagnosis).

• Reinfarction: It is now clear that cardiac troponin (cTn) increases rapidly, even if it is from an abnormal baseline in patients with reinfarction.

• Sometimes cTn concentration is elevated, but stable or decreasing at the time of suspected reinfarction. In this case, the diagnosis of reinfarction requires a 20% or greater increase in the cTn measurement

Creatine kinase (CK):

• Less sensitive and specific than troponins

• When cTn is available, CK-MB should not be used for the initial diagnosis of acute MI

Lactate dehydrogenase (LDH):

• Normally, LDH2>LDH1; but, in MI, LDH1>LDH2. This is called flipping of the LDH ratio.

• Current recommendations suggest that LDH no longer has a role in the diagnosis of MI.

Myoglobin:

• Earliest enzyme to increase after MI.

• Not specific to cardiac muscle and can be elevated to any form of injury to skeletal muscle.

Heart-type fatty acid-binding protein (H-FABP), which mainly exists in cardiomyocytes, has recently emerged as a potential biomarker, which was found to be sensitive for the early diagnosis of acute myocardial infarction. H-FABP as an early marker and cTnI as a late marker would be the ideal combination to cover the complete diagnostic window for AMI

Solution to Question 21:

The given clinical scenario of the patient presenting with palpitations and the electrocardiogram (ECG) showing narrow QRS complex tachycardia with regular heart rate suggests a paroxysmal supraventricular tachycardia. As the patient is hemodynamically unstable (SBP of 70 mmHg), the next step in management is DC (direct current) cardioversion.

PSVT is a narrow QRS complex tachycardia with sudden onset and termination. It mainly presents with dizziness and palpitations. Evaluation of these patients includes monitoring of vitals and ECG.

The ECG below shows PSVT.



Management of PSVT is as follows:

1) When the patient is hemodynamically stable:

• Carotid sinus massage is attempted. It is avoided if carotid bruit is present and if a prior history of stroke is present

• Valsalva maneuver is done when the patient is cooperative

• Intravenous adenosine is useful in most cases. It is usually given in a bolus dose of 6 mg (IV bolus over 2-3 seconds followed by a rapid flush with 20 mL NS followed by a dose of 12 mg if required. However, its usage is contraindicated in patients with prior cardiac transplantation as it can aggravate bronchospasm and atrial fibrillation.

• Intravenous calcium channel blockers such as verapamil or diltiazem and beta-blockers are helpful but can cause hypotension.

• If there is no termination of the PSVT, then the next step is IV ibutilide + AV nodal–blocking agent (or) IV procainamide + AV nodal–blocking agent (or) cardioversion.

2) When the patient is hemodynamically unstable, i.e. when the patient has hypotension (BP<90/60) with unconsciousness or respiratory distress, QRS-synchronous DC cardioversion is done.



Solution to Question 22:

Renal tubular acidosis causes normal anion gap metabolic acidosis, not high anion gap acidosis.

Normally, in the serum, the total number of cations is equal to the total number of anions. But, a few of the anionic proteins (e.g., albumin), phosphates, sulfates, and organic anions are not measured and this is called an anion gap. It is the difference between measured cations and measured anions and is estimated as:

Plasma anion gap = [Na+] –([Cl-]+[HCO3-])

The normal anion gap ranges between 6-12 mEq/L.

Solution to Question 23:

Based on the given clinical scenario of a patient presenting with sudden onset of chest pain, with ECG findings of ST depression in the anterior chest leads and T wave inversion, the most likely diagnosis is a non-ST-segment elevation-acute coronary syndrome (NSTE-ACS). The most appropriate line of management in this patient would be to administer aspirin with heparin.

Percutaneous coronary intervention is indicated after coronary angiography in high-risk patients.

Fibrinolytic therapy is not indicated in patients with NSTE-ACS.

Arrhythmia prophylaxis is generally not routinely recommended in all patients with NSTE-ACS.

Acute coronary syndrome is classified into 2 groups:

• ST-segment Elevation Myocardial Infarction (STEMI)

• NSTE-ACS

NSTE-ACS further includes:

• NSTEMI (myocyte necrosis present and cardiac troponins elevated)

• Unstable Angina (myocyte necrosis absent and cardiac troponins not elevated)

The diagnosis of NSTE-ACS is based on the following:

Severe chest discomfort with at least one of 3 features:

• Occurring at rest or minimal exertion and lasting for more than 10 minutes

• Recent onset

• Crescendo pattern

ECG showing:

• ST segment depression

• T wave inversion

Elevated cardiac troponins in case of NSTEMI.

Management of NSTE-ACS:

To provide relief of pain and discomfort, initial treatment includes bed rest, nitrates, beta blockers, and inhaled oxygen (only if arterial O2 saturation <90%).

Antithrombotic therapy consisting of antiplatelet and anticoagulant drugs:

• Dual antiplatelet therapy (DAPT): Aspirin (162-325 mg) AND one ADP receptor inhibitor (Ticagrelor, clopidogrel or prasugrel)

• Anticoagulation: Enoxaparin, unfractionated heparin, fondaparinux or bivalirudin

• An early invasive strategy including diagnostic coronary angiography (within 2–72 hours of presentation) followed by revascularization (percutaneous coronary intervention/ coronary artery bypass grafting) is indicated for stabilized patients who are at high risk of coronary events (Based on TIMI, GRACE or HEART score).

Solution to Question 24:

Of the given options, hyperkalemia is the most life-threatening complication of chronic kidney disease (CKD) as it can result in cardiac arrhythmias and sudden death.

CKD is staged based on declining glomerular filtration rate (GFR) and albuminuria. The various clinical manifestations occur due to loss of kidney function and accumulation of various uremic toxins (urea, creatinine, and other products of protein metabolism).

Complications of declining kidney function include:

• Volume overload

• Hyperkalemia

• Metabolic acidosis

• Hyperphosphatemia

• Mineral and bone disorders

• Hypertension

• Anemia

• Dyslipidemia

Solution to Question 25:

The likely diagnosis in the given clinical scenario with serum-ascites albumin gradient (SAAG) >1.1g/dL [calculated as serum albumin level - ascitic fluid albumin level, which in this case is 3-1=2g/dL] and ascitic fluid protein <2.5 g/dL is late Budd Chiari syndrome.

Serum-ascites albumin gradient (SAAG) can be used to differentiate between ascites due to portal hypertension and non-portal hypertensive ascites. It represents the pressure within the hepatic sinusoids and correlates with the hepatic venous pressure gradient.

It is given by the formula: SAAG = Serum albumin level - Ascitic fluid albumin level

Interpretation of SAAG:

• SAAG ≥1.1 g/dL indicates that the ascites is due to increased pressure in the hepatic sinusoids. It reflects the presence of portal hypertension.

• SAAG <1.1 g/dL indicates that the ascites is not related to portal hypertension. It may be due to infection or malignancy.

In cases with high-SAAG (≥1.1), ascitic protein level analysis further helps in identifying the etiology:

• ≥2.5 g/dL indicates that the hepatic sinusoids are normal and are allowing the passage of protein into the ascites as in early Budd-Chiari syndrome.

• <2.5 g/dL indicates that the hepatic sinusoids have been damaged and scarred and no longer allow passage of protein as in late Budd-Chiari syndrome

Solution to Question 26:

Iritis is seen in all the above conditions except rheumatoid arthritis.

Iritis or iridocyclitis is also known as anterior uveitis.

Anterior uveitis is seen in:

• Ankylosing spondylitis

• Juvenile rheumatoid arthritis

• Psoriasis

• Reactive arthritis

• Systemic lupus erythematosus

• Leprosy

Posterior uveitis is seen in:

• Vogt-Koyanagi Harada syndrome

• Diseases like toxoplasmosis, cysticercosis, coccidioidomycosis, toxocariasis, histoplasmosis

• Infections- Candida, Pneumocystis carinii, Cryptococcus, Aspergillus, and Cytomegalovirus

• Cat-scratch disease, Whipple's disease, and brucellosis

Both anterior and posterior uveitis is seen in:

• Sarcoidosis

• Behcet's disease

• Inflammatory bowel disease

• Herpes infections, Syphilis, Lyme disease, TB, onchocerciasis

Intermediate uveitis/pars planitis is seen in multiple sclerosis.

Solution to Question 27:

The given clinical scenario of a young patient with a history of sudden collapse following physical activity along with ECG findings of V5, V6 inversion, and prominent inverted T-waves, and 2D echocardiography showing the systolic anterior motion of the mitral valve leaflet point toward a diagnosis of hypertrophic obstructive cardiomyopathy (HCM).

In HCM, clinical examination may reveal pulsus bisferiens and an ejection systolic murmur that increases on standing.

Beta-blockers and implantable cardioverter-defibrillators (ICD) are used in the management.

HCM is a leading cause of sudden death in young. It is inherited in an autosomal dominant pattern with variable penetrance. HCM is characterized by the thickening of the septal myocardium below the aortic valve leading to obstruction of the left ventricular outflow tract.

Symptoms:

• Exertional dyspnea

• Syncope

• Angina, palpitations, cardiac arrhythmias

• Sudden cardiac death (particularly during or after intense physical activity)

Examination:

• Systolic ejection murmur best heard at the apex or lower left sternal border

• Holosystolic murmur from mitral regurgitation

• Sustained apex beat

• S4 gallop

• Paradoxical split of S2

• Pulsus bisferiens: LV outflow obstruction causes a sudden quick rise of the pulse followed by a slower longer rise.

ECG findings:

• Left axis deviation

• P wave abnormalities

• Prominent abnormal Q waves

• Deeply inverted T waves

Echocardiography findings:

• LV wall thickness ≥15 mm

• Systolic anterior motion of the mitral valve leaflet

• Mitral regurgitation

Treatment:

• Beta-blockers and calcium channel blockers are the first-line treatment

• Implantable cardioverter-defibrillator (ICD) - indicated when the history of cardiac arrest or sustained ventricular tachycardia is present

Parameters	Murmur intensity increased	Murmur intensity decreased
Preload	ValsalvaStanding(↓preload = ↑obstruction)	SquattingPassive leg raising (↑preload = ↓obstruction)
Force of contraction	Exercise(↑force = ↑obstruction)	
Afterload	Vasodilators(↓afterload = ↑obstruction)	Isotonic handgripSquatting(↑ afterload = ↓obstruction)

Solution to Question 28:

A, B, C & E are clinical criteria of admission under COVID-19 ward.



Solution to Question 29:

The correct sequence is ABDC.

A - 'a' wave: The 'a' wave in the JVP waveform corresponds to atrial contraction and occurs just after the P wave on the ECG (which corresponds to atrial depolarization) and just before the first heart sound.

B- 1st heart sound: The closure of the mitral and tricuspid valves at the beginning of ventricular systole produces the 1st heart sound (S1) as the change in pressure across the AV valves causes them to close. This sound occurs soon after the QRS complex on the ECG waveform representing ventricular depolarization.

The x-descent of the JVP waveform is associated with the ventricular systole. The c-wave occurs due to the bulging of the tricuspid valve into the atrium until the semilunar valves open.

D - 'T' wave: Following ventricular depolarization, the ventricles start to repolarize represented by the 'T' wave on ECG. The 2nd heart sound (S2) occurs due to the closure of the semilunar valves.

C- Rapid filling of ventricles: Ventricular repolarization triggers the ventricles to enter diastole. During early diastole, there is a phase of rapid filling of the ventricles. This occurs as the atrial pressure exceeds the ventricular pressure, allowing blood to passively flow from the atria to the ventricles.

The 3rd heart sound (S3) may be heard here. This event is associated with the v-wave and the y-descent on the JVP waveform as the ventricle relaxes further.



Solution to Question 30:

The given clinical picture of a student diagnosed with dengue with history of an afebrile period, epistaxis, reduced platelet count and elevated hematocrit strongly suggests dengue hemorrhagic fever.

He should be kept under observation and given adequate oral and intravenous fluids (Statements 2,3 and 4).

Clinical criterion for dengue hemorrhagic fever is:

• Acute onset fever lasting 2-7 days.

• Hemorrhagic manifestations: positive tourniquet test, petechiae, ecchymoses, purpura, or bleeding from the mucosa, gastrointestinal tract, injection sites, or other locations.

• Platelet count ≤ 100,000 cells/mm.

• Objective evidence of plasma leakage: rising hematocrit/ haemoconcentration of ≥20% from baseline or signs of plasma leakage (e.g., pleural effusion, ascites, hypoproteinemia/ albuminemia).

Severe dengue symptoms often come after the fever has subsided, when the capillary permeability may increase along with rising hematocrit levels. Progressive leukopenia, followed by a rapid decrease in platelet count, often precedes plasma leakage.

The main pathology in dengue is plasma leakage, resulting in intravascular volume depletion and thereby shock. So, supportive management with oral or intravenous rehydration is the mainstay of treatment. If the patient is febrile, paracetamol is administered to maintain body temperature. NSAIDs should be avoided since they can cause gastritis, platelet dysfunction, and severe bleeding.

Patients should be kept under observation and vital parameters like blood pressure, urine output, platelet count, and rise or fall in hematocrit should be monitored.

Indications for platelet transfusion in dengue:

• Platelet count < 10,000/cu.mm.

• Prolonged shock with coagulopathy and abnormal coagulogram

• Systemic massive bleeding

Solution to Question 31:

Among the given options only statements 1 and 5 are true.

Kayser-Fleischer (KF) rings are seen in almost all patients of Wilson's disease with neurological involvement. It initially appears as a crescent in the superior pole of Descemet's membrane and gradually extends inferiorly, eventually becoming circumferential in shape.

KF rings occur due to copper deposition in the descemet's membrane. It is golden-brown to greenish-brown in color and can be best seen by slit-lamp examination.

The image below shows a KF ring :



While it is a characteristic sign of Wilson’s disease (statement 2), it can also be seen in:

• Primary biliary cholangitis

• Primary sclerosing cholangitis

• Cholestasis

KF rings are present in almost 95% of patients of Wilson's disease with neurological/psychiatric symptoms compared to around 65% of those with hepatic presentations (statement 4).

They disappear on treatment with copper-chelating agents like d-penicillamine (while desferrioxamine is an iron chelating agent) and reappear with disease progression (statement 3).

Medicine NEET 2018

Question 1:

Thrombosis of posterior inferior cerebellar artery causes _____.

a) Lateral medullary syndrome

b) Weber syndrome

c) Medial medullary syndrome

d) Millard Gubler syndrome

Question 2:

A patient presents with frequent urination, nocturia, and enuresis. 24-hour urine volume was measured and recorded to be 7 liters. The urine osmolarity was 260 mOsm/L. ADH assay was performed and the recorded values were reported as 0.8 pg/ml. An MRI of the brain was performed and T1 weighted indicated the absence of the bright spot. What is the most likely diagnosis?

a) Nephrogenic DI

b) Primary polydipsia

c) Pituitary DI

d) Mannitol infusion

Question 3:

What levels of prolactin are definitely suggestive of prolactinoma?

a) >50µg/L

b) >100µg/L

c) >150µg/L

d) >200µg/L

Question 4:

A patient develops prosthetic valve endocarditis 2 years after valve replacement surgery. Which of the following organism is the most likely cause?

a) Streptococci

b) Staphylococcus aureus

c) Coagulase negative staphylococci

d) HACEK organisms

Question 5:

What is the most common extra-articular manifestation of rheumatoid arthritis?

a) Subcutaneous nodule

b) Sjogren's syndrome

c) Felty's syndrome

d) Vasculitis

Question 6:

Periodic acid Schiff positive macrophages are seen in:

a) Agammaglobulinemia

b) Whipple's disease

c) Abetalipoproteinemia

d) Crohn's disease

Question 7:

According to the berlin definition, moderate ARDS is characterized by all, except

a) PaO2/FiO2 200 - 300mm/Hg

b) Bilateral interstitial infiltrates

c) Symptom onset within a week

d) No cardiac failure on echocardiography

Question 8:

Low serum copper levels attributed to ATP7A gene is due to:

a) Dubin-Johnson syndrome

b) Wilson's disease

c) Menke's disease

d) Gilbert's syndrome

Question 9:

A patient presents with cutaneous vasculitis, glomerulonephritis, and peripheral neuropathy. Which investigation should be performed next to help you diagnose the condition?

a) ANCA

b) RA factor

c) HbsAg

d) MIF

Question 10:

What is the approximate time interval between HIV infection & manifestation of AIDS?

a) 7.5 years

b) 10 years

c) 12 years

d) 5 years

Question 11:

Myasthenia gravis is associated with _____.

a) Decreased acetylcholine release at the nerve endings

b) Decreased myosin

c) Absence of troponin C

d) Decreased synaptic transmission at the myoneural junction

Question 12:

Which among the following is wrong regarding Weber's syndrome?

a) Contralateral hemiplegia

b) Ipsilateral oculomotor nerve palsy

c) Contralateral Parkinsonism

d) Ipsilateral paralysis of lower face

Question 13:

A 10-year-old boy complains of multiple episodes of acute severe pain in his fingers and toes which resolves on its own for the past 1 year. Which of the following conditions is he most likely suffering from?

a) Alpha-thalassemia

b) Beta-thalassemia

c) Sickle cell anemia

d) Von Willebrand disease type 1

Question 14:

Cryoglobulinemia is associated with _____.

a) Hepatitis C

b) Ovarian cancer

c) Diabetes

d) Leukemia

Question 15:

Which is the characteristic pattern seen in Brown-Sequard syndrome?

a) Contralateral loss of joint sense and position

b) Contralateral loss of pain sensation

c) Ipsilateral loss of complete sensory functions

d) Contralateral motor functions

Question 16:

A child presents with hepatitis and progressive neurological degeneration. A picture of his eye has been provided below. Among the following options, what would be the initial step regarding the investigations done for this child's diagnosis?



a) Serum ceruloplasmin levels

b) Serum copper levels

c) Enzyme assay

d) Karyotyping

Question 17:

Automatic implantable cardioverter-defibrillator (AICD) implantation is done for which of the following conditions?

a) Brugada syndrome

b) Ventricular fibrillation

c) Acute coronary syndrome with low ejection fraction

d) All of the above

Question 18:

Pseudo P pulmonale is seen in which electrolyte abnormality?

a) Hypokalemia

b) Hyponatremia

c) Hypocalcemia

d) Hypercalcemia

Question 19:

The Gold's criteria for very severe COPD is:

a) FEV1/FVC <0.7 and FEV1 < 30%

b) FEV1/FVC <0.7 and FEV1 < 70%

c) FEV1/FVC<0.7 and FEV1< 50%

d) Both A and C

Question 20:

A 35-year-old female patient has hypokalemia, hypertension and metabolic alkalosis. What is the most likely etiology?

a) Bartter syndrome

b) Gitelman's syndrome

c) Liddle's syndrome

d) Fanconi's syndrome

Question 21:

What is the diagnostic test of choice for HIV in a baby born to an HIV-infected mother?

a) HIV DNA PCR

b) Cord blood ELISA

c) Western blot

d) Third generation ELISA

Question 22:

Which of the following is not a common feature of allergic bronchopulmonary aspergillosis?

a) Distal bronchiectasis

b) Cough

c) Wheezing

d) Raised serum IgE levels

Question 23:

A 58-year-old male patient has been diagnosed with hypertension. On examination, there is no existing cardiovascular disease, no hypertension mediated organ damage (HMOD) and the renal function is normal. He should be immediately started with pharmacological management if his blood pressure is more than _____.

a) >130/80 mmHg

b) ≥160/100 mmHg

c) > 150/100 mmHg

d) ≥140/90 mmHg

Question 24:

A middle-aged woman who presented with tremors, palpitations, weight loss, and menstrual irregularities is diagnosed with Graves' disease. Which of the following statements would be correct with respect to her thyroid function test?

a) Low TSH levels

b) Low free T4 levels

c) Low serum T3 levels

d) Radioactive Iodine Uptake (RAIU) at 24 hours below normal

Question 25:

The observed myocardial stunning pattern does not match the ECG findings of a patient. What is the probable diagnosis?

a) Takotsubo cardiomyopathy

b) Restrictive cardiomyopathy

c) Brigade's cardiomyopathy

d) Pericardial tamponade

Question 26:

Which of the following antiretroviral drugs is not used in HIV & hepatitis B co-infection?

a) Tenofovir

b) Abacavir

c) Lamivudine

d) Emtricitabine

Question 27:

Who comes under category III non-heart-beating donor?

a) A patient who died during transportation to the hospital

b) A patient who died after failed resuscitation after reaching the hospital

c) A patient who was brought dead to the hospital

d) A patient who is awaiting death in the hospital

Question 28:

Which of the following conditions is associated with AV block?

a) Hypothyroidism

b) Cushing's syndrome

c) Hyperthyroidism

d) Pheochromocytoma

Question 29:

Which drug is used for mass chemoprophylaxis for meningococcal meningitis?

a) Ciprofloxacin

b) Chloramphenicol

c) Tetracycline

d) Penicillin

Answer Key

Question No.	Correct Option
1	a
2	c
3	d
4	a
5	a
6	b
7	a
8	c
9	a
10	b
11	d
12	d
13	c
14	a
15	b
16	a
17	d
18	a
19	a
20	c
21	a
22	a
23	d
24	a
25	a
26	b
27	d
28	a
29	a

Detailed Explanations

Solution to Question 1:

Thrombosis of the posterior inferior cerebellar artery (PICA) causes lateral medullary syndrome or Wallenberg syndrome.

PICA is a branch of the fourth part of the vertebral artery, which is a branch of the first part of subclavian artery.



Solution to Question 2:

The features of frequent urination, nocturia, with polyuria(>3L/day), low urine osmolarity(<280 mosm/L), low plasma ADH levels and MRI showing absent pituitary bright spot is suggestive of Pituitary Diabetes Insipidus.



Diabetes Insipidus(DI) :

Treatment:

• For Central DI, synthetic analog of ADH Desmopressin(DDAVP) is used.

• For Nephrogenic DI, a low sodium diet with thiazide diuretic and/or amiloride and prostaglandin synthesis inhibitor is used.

• For primary polydipsia, there are no effective drugs used but for Iatrogenic type patient education is helpful.

Other options:

Option A: In nephrogenic DI, there is polyuria, low urine osmolarity but the ADH levels will be >1pg/mL.

Option B: In primary polydipsia, there is polyuria, low urine osmolarity, ADH levels will be <1pg/mL but a pituitary bright spot present.

Option D: In solute loss (mannitol or hypertonic saline infusion), urine osmolality is >300 mOsm/L.

Types	Primary pathology	Caused by
Pituitary/ Central DI	Primary deficiency of ADH se cretion	AVP -Neurophysin II gene(m ost common gene defect)
Primary polydipsia	Suppression of ADH secretio n by excessive fluid intake	1. Dipsogenic DI : Following head trauma, TB meningitis, Multiple sclerosis.2. Psychog enic DI : Due to psychosis or OCD3. Iatrogenic polydipsia : Due to motivated health ben efits
Gestational DI	Deficiency due to Increased d egradation of ADH	N terminal aminopeptidase p roduced by placenta
Nephrogenic DI	Renal insensitivity to ADH or drugs like lithium or genetic mutation	defect in gene encoding V2 re ceptor on chXq28(most com mon gene defect)

Solution to Question 3:

The diagnosis of a prolactinoma is highly likely when prolactin levels are > 200 μg/L.

The normal range of serum prolactin (PRL) is 5-20 μg/L. Prolactin levels < 100 μg/L can be due to non-functioning pituitary adenomas or non-neoplastic causes of hyperprolactinemia. MRI brain must be done in all patients with hyperprolactinemia.

Prolactinoma are tumors arising from the lactotroph cells of the pituitary gland, they can be classified as microadenoma(<1cm in diameter, usually non invasive) or macroadenoma(>1cm, locally invasive to adjacent structures). Clinical features include amenorrhea, infertility, and galactorrhea in women, impotence, loss of libido, infertility in men and if tumor extends beyond sella, there can be visual field defects and other mass effects.

Treatment - In asymptomatic patients and those where fertility is not desired, they do not require treatment and can be monitored using regular PRL measurements and MRI. In symptomatic microadenoma cases, oral dopamine agonists (cabergoline and bromocriptine) form the main

treatment for hyperprolactinemia and reduction of tumor volume. For macroadenoma, before starting dopamine agonists visual field testing is to be done, and reassess it with MRI changes every 6-12 month intervals until maximum reduction is achieved, in unresponsive cases consider surgery.

Solution to Question 4:

Patient developing endocarditis 2 years after prosthetic valve surgery suggests late-onset endocarditis. In such cases, the most common causative organism is streptococci.

Solution to Question 5:

The most common extra-articular manifestation of rheumatoid arthritis(RA) is subcutaneous nodules.

The nodules are generally benign, non-tender, and firm, and occur in areas subjected to repeated trauma such as the forearm, sacral prominences, and Achilles tendon. They can also occur in the lungs, pleura, pericardium, and peritoneum.

Patients who have a history of cigarette smoking had early onset arthritic manifestations and tested positive for RA factor or ACPA are more likely to develop extra-articular manifestations. Other common extra-articular manifestations include secondary Sjogren's syndrome, interstitial lung disease, pulmonary nodules, and anemia. FELTY'S syndrome is a triad of neutropenia, splenomegaly, and nodular RA.

Solution to Question 6:

Periodic acid Schiff (PAS) positive macrophages are seen in Whipple's disease.

Whipple's disease is a chronic infection caused by Tropheryma whipplei, classically presents as chronic diarrhea, abdominal pain, intermittent fever, weight loss, arthralgia/arthritis with neurological(cognitive changes progressing to dementia, personality and mood alterations, hypothalamic involvement and supranuclear ophthalmoplegia) and cardiological involvement (endocarditis, congestive heart failure, embolic events, aneurysms).

Investigations- Intestinal biopsy showing PAS-positive diastase resistant inclusions in lamina propria, specific PCR-based diagnosis has higher sensitivity. Urine PCR is an important tool to assess the success of therapy.

Treatment- IV ceftriaxone (2 g q12h) or meropenem (2 g q8h) for 2–4 weeks and then treated with oral doxycycline, or minocycline plus hydroxy- chloroquine for at least 1 year. Trimethoprim/Sulfamethoxazole is used as an oral alternative, but high relapses.

Solution to Question 7:

Moderate ARDS is characterized by PaO2/FiO2: 101 - 200 mmHg. PaO2/FiO2: 200 - 300 mm/Hg is classified as mild ARDS.

Diagnostic criteria in ARDS—Berlin definition:(requires each of the following criteria to be present to diagnose ARDS)

• Onset - Acute: Respiratory symptoms must have begun < 1 week of a known clinical insult

• Bilateral alveolar or interstitial infiltrates: on a chest X-ray image or CT scan

• Respiratory failure must not be fully explained by cardiac failure or fluid overload. Objective assessment e.g Echo should be done in cases of no known cause.

• Impairment of oxygenation: The severity of hypoxemia defines the severity of ARDS.

Note: American-European Consensus Conference (AECC) classification of ARDS is no longer used.

Image: Alveolar/interstitial infiltrates in ARDS patient

Severity	PaO2/FiO2(On ventilator settings that include PEEP ≥5 cm H2O)
Mild	> 200 mmHgbut≤ 300 mmHg
Moderate	> 100 mmHgbut≤ 200 mmHg
Severe	≤ 100 mmHg



Solution to Question 8:

Low serum copper level attributed to the ATP7A gene is due to Menke's disease (Option C).

Menke's disease or Menke-Kinky hair syndrome is an X-linked recessive condition occurring due to a mutation in the ATP7A gene on chromosome X. This causes impaired transport of dietary copper from the small intestine to the liver leading to low levels of copper-containing enzymes. It presents within 3 years of age and is characterized by:

• Depigmented kinky hair

• Failure to thrive

• Neurodegenerative symptoms like loss of developmental milestones, truncal hypotonia, epilepsyLoose, wrinkled skin

• Multiple fractures

• Poor visual acuity

• Recurrent infections.

Lab investigations show low serum copper (<65 µg/dL) and low ceruloplasmin(<20 mg/dl). The disease is fatal as it has no curative treatment. But, administration of L-histidine and copper chloride injections may be beneficial, if started early.

Aceruloplasminemia is an autosomal recessive disease characterized by tissue iron overload, mental deterioration, microcytic anemia, and low serum iron and copper concentrations.

Option B: Low serum copper and low ceruloplasmin is also seen in Wilson's disease, but is associated with the gene defect ATP7B on chromosome 13. It is an autosomal recessive disorder of copper metabolism with features of hepatolenticular degeneration and Kayser Fleischer ring.

Option A: Dubin Johnson Syndrome is an autosomal recessive disorder characterized by impaired canalicular excretion of bilirubin due to mutation in MRP-2 protein leading to conjugated hyperbilirubinemia.

Option D: Gilbert's Syndrome is characterized by decreased UDP glucuronyl transferase activity leading to unconjugated hyperbilirubinemia, which can be autosomal recessive or dominant.

Solution to Question 9:

The presence of cutaneous vasculitis, glomerulonephritis, and peripheral neuropathy are suggestive of small-vessel vasculitis. Investigating for ANCA (antineutrophil cytoplasmic antibodies) should be the next step to aid in diagnosis.

Small Vessel Vasculitis/ ANCA- associated Vasculitis

Granulomatosis with Polyangiitis(Wegener's granulomatosis)	Microscopic Polyangiitis	Eosinophilic granulomatosis with Polyangiitis(Churg Strauss syndrome)
Characterized by a triad ofupper and lower respiratory tractinvolvement withglomerulonephritis.	Characterized bynecrotizing vasculitiswithglomerulonephritis(pauci immune, crescentic)	Characterized byasthma,peripheral and tissue eosinophilia,extravascular granulomaformation, andvasculitisof multiple organs.
Nodular cavitary infiltratesin lungs showing necrotizing granulomas, and vasculitis on lung biopsy	Diffuse alveolar hemorrhage, hemoptysis.No URTI, No granulomas, No pulmonary nodules/cavities.	Mononeuritis multiplex- most common and characteristic. Biopsy showing granulomas with eosinophilic infiltrates.
C-ANCA +	P-ANCA +	P-ANCA +

Solution to Question 10:

The time interval between primary HIV infection and the manifestation of AIDS is 10 years.



After exposure to the HIV virus, there occurs primary infection following an incubation period of 2-4 weeks, which is usually symptomatic in more than 50% of cases, following 3-6 weeks there is acute HIV syndrome due to wide dissemination of the virus, characterized by fever, sore throat, diarrhea, lymphadenopathy, and rash.

This peak viremia drops as the immune response develops, to reach a plateau about 3 months later. Following this there is a prolonged period of clinical latency, during which infected individuals are asymptomatic, lasting for a median of 10 years.

Following this period, chronic persistent infection develops which is the hallmark of HIV disease called Acquired immunodeficiency syndrome (AIDS).

Long-term non-progressors: a small part of untreated HIV-infected people with CD4 counts within the reference range for >10 years.

Elite controllers: Some long-term non-progressors have undetectable viral loads.

Solution to Question 11:

Myasthenia gravis is associated with a decrease in synaptic transmission at the myoneural junction.

Myasthenia gravis is a postsynaptic neuromuscular disorder that is characterized by weakness and fatigability of skeletal muscles. It is due to autoantibody-mediated destruction of the acetylcholine receptors (AChRs) at the neuromuscular junction (NMJ). This consequently leads to a decreased transmission at the neuromuscular (myoneural) junction.

Lambert Eaton Myasthenic Syndrome (LEMS) is a presynaptic disorder of the NMJ that can cause weakness similar to that of Myasthenia Gravis. It is due to autoantibodies directed against P/Q-type calcium channels at the motor nerve terminals.

Myasthenia gravis	Lambert Eaton Myasthenic S yndrome
Postsynapticcondition	Pre synapticcondition
Antibodies againstAchR,MuS K,LRP-4	Antibodies against fastP/Q ty pe calcium channel
Earliest manifestation -ptosis, diplopia	Earliest manifestation -proxi mal musclesoflower limbinvo lvement
DTR -normal	DTR -depressed/ absent
Autonomic changes -absent	Autonomic changes -presentl ike dry mouth and impotence
On exertion -increased weak nessseen	On exertion -decreased weak nessseen
EMG(repeated nerve stimulat ion) -decremental response	EMG(repeated nerve stimulat ion) -incremental response
Associated withthymicabnor mality	Associated withsmall cell car cinomaoflungin older adults

Solution to Question 12:

Ipsilateral paralysis of the lower face is not seen in Weber's syndrome. However, there is contralateral paralysis of the lower face.

Weber syndrome is caused by infarction in the distribution of the penetrating branches of the posterior cerebral artery (PCA). It affects the cerebral peduncle, especially medially, with damage to the fascicle of cranial nerve III and the pyramidal fibers.

The resultant clinical findings are contralateral hemiplegia of the face, arm, and leg secondary to corticospinal and corticobulbar tract involvement, and ipsilateral oculomotor paresis including a dilated pupil.

Structure damaged	Effect
Cranial nerve III fibres	Ipsilateral ptosisIpsilateral di vergent squintLoss of accom modation reflexLoss of light r eflex
Corticobulbar tract	UMN lesion leading to contra lateral paralysis of the lower f ace
Corticospinal tract	Contralateral hemiplegia



Solution to Question 13:

The features of multiple episodes of acute severe pain in fingers and toes with spontaneous resolution are suggestive of dactylitis (hand-foot syndrome), which usually occurs in sickle cell

anemia.

The sickled RBCs have a sticking tendency towards the endothelium of the small venules. These abnormalities provoke unpredictable episodes of vaso-occlusion. Dactylitis is characterized by severe pain affecting the bones of the hands, feet, or both. It is often the first symptom of sickle cell anemia in children.

Treatment - Administration of patient-controlled analgesia or a frequent fixed dose of opioids with rescue doses for breakthrough pain are preferred with frequent monitoring during the initial 48-72 hrs to prevent unexpected death due to arrhythmias or pulmonary embolism.

Other manifestations include:

• Splenic autoinfarction - repeated splenic microinfarcts

• Acute splenic sequestration - venous obstruction, needing emergency transfusion/splenectomy

• Painful ischemic crises - vaso-occlusion in connective and musculoskeletal tissue

• Renal papillary necrosis leading to isosthenuria

• Cerebrovascular strokes

• Chronic lower leg ulcers

• Priapism in males

α-thalassemia (Option A) and β-thalassemia (Option B) are characterized by pallor, jaundice, growth failure, hepatosplenomegaly, characteristic thalassemic facies, and hair-on-end appearance of the skull.

Option D: Von Willebrand Disease is characterized by predominantly mucosal bleeding symptoms like excessive bruising and epistaxis.

Solution to Question 14:

Cryoglobulinemia is associated with hepatitis C infection.

Following Hepatitis C infection, there is an aberrant immune response leading to an immune complex formation containing hepatitis C antigens, polyclonal IgG specific to Hepatitis C, and monoclonal IgM rheumatoid factor, which triggers an inflammatory cascade results in Cryoglobulinemic vasculitis.

Cryoglobulins are the immunoglobulins that precipitate at temperatures less than 37°C, the presence of these cryoglobulins in serum leads to Cryoglobulinemia, which is characterized by cutaneous vasculitis, palpable purpura, arthritis, peripheral neuropathy, and glomerulonephritis.

Important conditions associated with cryoglobulinemia are:

• Hepatitis C

• Multiple myeloma

• Connective tissue disorders

• Lymphoproliferative disorders

Solution to Question 15:

The Brown–Sequard syndrome presents with contralateral loss of pain sensation.

Lateral hemisection of the spinal cord produces the Brown–Séquard syndrome (also known as the hemicord syndrome). Knife or bullet injuries and demyelination are the most common causes of this condition.

Clinical features include:

• Ipsilateral weakness (corticospinal tract involvement)

• Ipsilateral loss of joint position and vibratory sense (posterior column involvement)

• Contralateral loss of pain and temperature sense (spinothalamic tract involvement) one or two levels below the lesion

• Ipsilateral segmental signs like radicular pain, muscle atrophy, or loss of a deep tendon reflex

Solution to Question 16:

The clinical features of hepatitis, progressive neurological degeneration, and the Kayser-Fleischer (KF) ring seen in the image are suggestive of Wilson's disease. The initial step regarding investigations done for the diagnosis is the measurement of serum ceruloplasmin levels.

Kayser-Fleischer (KF) ring is a golden to greenish-brown ring of deposition of copper in the Descemets membrane of the peripheral cornea. Patients with neurological manifestations manifest with KF ring in >95% of cases.

Lab investigations in Wilson's disease would show:

The gold standard technique for diagnosis of Wilson’s disease is percutaneous needle liver biopsy for measurement of hepatic copper levels >200 μg/g of dry weight.

Lab markers	Change in Wilson's disease	Reason
Serum ceruloplasmin	↓↓↓	ATP7B deficiency results in reduced bound copper
24h urinary copper	↑↑ (>100 μg/24h)	Excess free copper excreted in urine
Total serum copper	↓	Due to marked reduction in ceruloplasmin

Option B: The role of total serum copper in the diagnosis of Wilson's disease is limited. Measurement of serum ceruloplasmin and urinary copper is preferred.

Option C: ATP7B enzyme assays have shown promise in early studies as a neonatal screening test for Wilson's disease. However, it is not yet part of routine clinical practice.

Option D: Karyotyping is done for every asymptomatic 1st and 2nd degree relative of a patient with Wilson's disease. This helps differentiate the carriers (heterozygous) from the patients (homozygous) and can lead to early initiation of treatment before the onset of symptoms.

Solution to Question 17:

An Automated Implantable Cardioverter-Defibrillator (AICD) is a device placed in the chest or abdomen to detect arrhythmias and perform cardioversion, defibrillation, and pace the heart.

It is used in: -

• Brugada syndrome- refers to a sodium channelopathy with a high risk of arrhythmias that can lead to sudden cardiac death.

• Ventricular fibrillation and ventricular tachycardia.

• Acute coronary syndrome with a low ejection fraction.

ICDs are indicated for primary prevention in patients who had myocardial infarction >40 days ago (with Class II-III NYHA Heart failure with LVEF <35% and Class I NYHA with LVEF <30%). Cardiac resynchronization therapy for prevention of sudden cardiac death in heart failure. High-risk situations of sudden cardiac death in hypertrophic cardiomyopathy, non-ischemic dilated cardiomyopathy with low ejection fraction.

Solution to Question 18:

Pseudo P pulmonale refers to tall peaked P waves seen in hypokalemia.

P-pulmonale refers to P waves taller than >2.5mm in an ECG lead.



Hypocalcemia (Option C) prolongs the QT interval.

Hypercalcemia (Option D) shortens the QT interval.

Solution to Question 19:

Gold's criteria for very severe COPD (stage IV) are FEV1/FVC <0.7 and FEV1 < 30%.

The severity of COPD is assessed by Global Initiative for Chronic Obstructive Lung Disease (GOLD) classification:

Note: FEV1/FVC and FEV1 predicted are measured after the administration of a bronchodilator.

GOLD Stage	Severity	Spirometry
I	Mild	FEV1/FVC <0.7 and FEV1 ≥80% predicted
II	Moderate	FEV1/FVC <0.7 and FEV1 ≥50% but <80% predicted
III	Severe	FEV1/FVC <0.7 and FEV1 ≥30% but <50% predicted
IV	Very severe	FEV1/FVC <0.7 and FEV1 <30% predicted



Solution to Question 20:

Hypertension with hypokalemia and metabolic alkalosis is seen in Liddle's syndrome.

Liddle's syndrome (pseudo aldosteronism) is an autosomal dominant condition that is caused due to the increased activation of ENaC channels. This causes increased reabsorption of Na+ exchanged for K+ causing hypokalemia. Increased Na absorption leads to hypertension, and there

is also a loss of H+ ion with K+, leading to metabolic alkalosis. It is associated with low renin and low aldosterone levels.

Solution to Question 21:

In a baby born to an HIV-infected mother, HIV infection is diagnosed using HIV DNA PCR. NACO recommends the use of a qualitative HIV-1 DNA PCR (NAT) in children aged <18 months for Early Infant Diagnosis (EID).

Antibody assays (e.g., ELISA) are not reliable in infants and young children because of the persistence of transplacentally acquired maternal antibodies. For this reason, HIV virologic testing must be performed using tests that detect HIV DNA or RNA to diagnose HIV infection in children.

Scheduled repeated testing is necessary because test sensitivity increases with time from HIV virus exposure and very recent infection will not be detectable. The cord blood should not be used for testing because of the possibility of contamination of the sample with maternal blood.

Solution to Question 22:

Central bronchiectasis is a typical feature of allergic bronchopulmonary aspergillosis (ABPA), not distal bronchiectasis.

ABPA is an eosinophilic pulmonary disorder caused by a hypersensitivity reaction to the colonization of airways by the fungus Aspergillus fumigatus. It occurs almost exclusively in patients with asthma (~2.5% of patients) or cystic fibrosis (up to 15%). In chronic cases, repeated episodes of bronchial obstruction, inflammation, and mucoid impaction can lead to bronchiectasis, fibrosis, and respiratory compromise. While central bronchiectasis is characteristic, some patients can develop chronic cavitary lesions.

With the development of ABPA, asthma or cystic fibrosis typically worsens and may manifest with a new or worsening cough or an increase in sputum production or wheezing. Many patients report coughing up thick sputum casts. The presence of hyperattenuating mucus in the airways is highly specific.

ABPA is diagnosed serologically, based on the presence of:

Serum IgE > 1000 IU/ml

Positive skin prick test or IgE levels against A. fumigatus extract

Detection of Aspergillus-specific antibodies

Eosinophilia >500 cells/μL

Treatment - The use of systemic glucocorticoids is indicated for ABPA, but the course should be tapered by 3-6 months. Antifungal agents like itraconazole, voriconazole, and newer azoles can be used for over 4 months to control disease activity. Monoclonal antibodies against IgE, IL-4, and IL-5 receptors can be used for severe refractory cases.

Solution to Question 23:

In this patient with no markers of cardiovascular disease, no hypertension-mediated end-organ damage (HMOD), and normal renal function, immediate pharmacological treatment is recommended when the blood pressure (BP) is ≥140/90 mmHg according to the AHA and ESH on the management of hypertension.

Guidelines for Hypertension Management:

• Adults 18–79 y with SBP ≥140 mm Hg or DBP ≥90 mm Hg

• Adults ≥80 y with SBP ≥160 mm Hg but ≥140 mm Hg can be considered.

• In frail patients, individualize treatment.

• In patients with CVD, especially CAD, SBP ≥130 mm Hg, or DBP ≥80 mm Hg.

Treatment BP targets:

• Adults 18–79 y, primary SBP/DBP goal <140/90 mm Hg; if treatment well tolerated target SBP/DBP <130/80 mm Hg but not SBP <120 mm Hg or DBP <70 mm Hg. For adults with isolated systolic hypertension, target SBP lowering, albeit cautiously.

• In adults ≥80 y, target SBP/DBP <140/90 mm Hg, if well tolerated

AHA Hypertension grading:

BLOOD PRESSURE CATEGORY	SYSTOLIC BP	DIASTOLIC BP
Normal	<120 mm Hg	<80 mmHg
Elevated	120-129 mmHg	<80 mmHg
Stage 1	130-139 mmHg	80-89 mm Hg
Stage 2	≥140 mm Hg	≥90 mmHg
Hypertensive crisis	≥180 mm Hg	≥120 mm Hg

Solution to Question 24:

In Graves' disease, the level of serum TSH is low, free T4 is high and serum T3 is high. In patients with thyrotoxicosis, the radioactive iodine uptake (RAIU) at 24 hours is characteristically elevated.

Grave's disease is characterized by thyrotoxicosis caused by thyroid-stimulating immunoglobulins (TSIs) synthesized by lymphocytes in the thyroid gland, bone marrow, and lymph nodes. Thyroid peroxidase (TPO) and thyroglobulin (Tg) antibodies are also seen. This results in excess

production of free T3 and free T4 levels and in turn reduces serum TSH levels.

RAIU is used to differentiate Grave's from other thyroid disorders, here there is diffuse high uptake. Measurement of TRab and TPOs is also done to confirm the diagnosis if the diagnosis clinically is unclear.



Solution to Question 25:

Takotsubo cardiomyopathy is the probable diagnosis when the observed myocardial stunning pattern does not match ECG changes.

Myocardial stunning is a state where certain segments of the myocardium (usually corresponding to an area of major coronary occlusion) show forms of contractile abnormality (dysfunction) on echocardiography.

In Takotsubo cardiomyopathy, while the ECG changes mimic a myocardial infarction(MI), the left ventricular dysfunction extends well beyond the distribution of a specific coronary artery. Thus, the myocardial stunning pattern might not fully match the ECG. Ventriculography shows apical dilation with basal contraction.

Takotsubo cardiomyopathy occurs typically in older women and is associated with a sudden increase in emotional or physical stress, presenting with hypotension, chest pain with ECG changes of acute MI, and pulmonary edema. Its also known as apical ballooning syndrome or acute stress-induced cardiomyopathy.

Option B: Restrictive cardiomyopathy is characterized by diastolic dysfunction with reduced contractility and ejection fraction, most commonly seen in amyloidosis, and enlarged ventricles on imaging with low-voltage ECG complexes.

Option C: Arrhythmogenic right ventricular cardiomyopathy (ARVC) is a type of familial dilated cardiomyopathy (DCM), having defects in desmosomal proteins, affecting the ventricles showing thin ventricular walls on Echo, epsilon waves, and low voltage ECG complexes.

Option D: Pericardial tamponade is characterized by hypotension, muffled heart sounds and increased jugular venous pressure with muffled heart sounds with total electrical alternans and sinus tachycardia in ECG.

Solution to Question 26:

Abacavir is not used in HIV and hepatitis-B co-infection. It is used in the first-line ART regimen for patients with abnormal serum creatinine values.

Nucleoside reverse transcriptase inhibitor (NRTI) drugs like lamivudine, its closely related drug emtricitabine, and the nucleotide reverse transcriptase inhibitor (NRTI) tenofovir have activity against hepatitis B virus. These drugs are preferred in combination with other antiretrovirals for the treatment of HIV & HBV co-infections.

Solution to Question 27:

Category III donors are patients with non-survivable injuries who have their treatment withdrawn and if such patients wished in life to be organ donors, the transplant team can attend at the time of treatment withdrawal and retrieve organs after cardiac arrest has occurred.

Modified Maastricht classification- Categories of non-heart-beating donors:

Category	Description	Circumstances
Category I	Includesvictims of accident a nd suicidewho arefound dead at the sceneand resuscitation is deemed pointless (e.g. fata l cervical spine fracture).	Uncontrolled - transplant tea m not present or prepared
Category I I	victims of sudden cardiac or cerebral catastrophewho are brought to emergency depart ments while being resuscitate d by ambulance personnel or whodied in the department. Other sources include patient s suffering an isolated brain i njury, anoxia, and stroke, an d victims of major trauma wh o died soon after hospital ad mission.	Uncontrolled - transplant tea m not present or prepared

Category	Description	Circumstances
Category I II	Patients who are dying, often in an intensive care unit. Th ese are the patients awaiting cardiac arrest where the treat ing clinicians have decided to withdraw treatment and not commence resuscitation for v arious reasons.	Controlled - transplant team present and prepared
Category I V	Comprises patients who suffe runexpected cardiac arrestdu ring or after the determinatio n ofbrain death.	Controlled - transplant team present and prepared
Category V	Unexpected cardiac arrestdur ingintensive care-	Uncontrolled - transplant tea m not present or prepared

Solution to Question 28:

Hypothyroidism is associated with AV block. In hypothyroidism, there is reduced myocardial contractility and reduced pulse rate leading to a reduced stroke volume and bradycardia, AV block.

Causes of AV block:

Autonomic:

• Carotid sinus hypersensitivity

Metabolic:

• Hyperkalemia

• Hypermagnesemia

• Hypothyroidism

• Adrenal insufficiency

Drug related:

• Beta blockers

• Calcium channel blockers

• Adenosine

• Digitalis

• Lithium

• Class 1 and class 3 anti arrhythmics

Infections:

• Endocarditis

• Lyme disease

• Chaga's disease

• Syphilis

• Tuberculosis

• Diphtheria

• Toxoplasmosis

Inflammatory:

• SLE

• RA

• Scleroderma

Infiltrative:

• Amyloidosis

• Sarcoidosis

• Hemachromatosis

Solution to Question 29:

Ciprofloxacin is used for chemoprophylaxis of meningococcal meningitis for close contacts. Ceftriaxone and rifampicin are also used.

Antibiotic chemoprophylaxis among close contacts of a patient with invasive meningococcal disease is recommended to prevent secondary cases.

• Rifampin - was earlier used widely, but is not optimal due to the high risk of adverse events, emerging resistance, and the need for compliance for four doses.

• Ciprofloxacin - widely used as a single oral dose, but it cannot be given in pregnancy and those below 18 years of age.

• Ceftriaxone - can be given in pregnancy and all ages as a single IM or IV dose. However, reduced susceptibility of isolates to ceftriaxone has occasionally been reported.

Recommended chemoprophylaxis for high-risk close contacts for meningococcal meningitis:

Patient group	Drug	Age	Dose	Route and duration
Non-pregnant individuals	Rifampicin	<1 month	5 mg/kg	Oral every 12 hrs for 2 days
>1 month	10 mg/kg	Oral every 12 hrs for 2 days		
Ciprofloxacin	>18 years	500 mg	Oral single dose	
Children and pregnant women	Ceftriaxone	<15 years	125mg	IM single dose
> 15 years	250 mg	IM single dose		

Medicine NEET 2019

Question 1:

A 65-year-old female patient is brought to the emergency department in a state of unconsciousness. Her BP is 70/50 mm of Hg. Her ECG is shown below. What is the next best step in the management of her condition?



a) IV verapamil

b) IV adenosine

c) Carotid massage

d) DC cardioversion

Question 2:

Nasal polyps are commonly associated with:

a) Intrinsic asthma

b) Brittle asthma

c) Extrinsic asthma

d) Exercise-induced asthma

Question 3:

All of the following are AIDS defining illnesses except:

a) Encephalopathy attributed to HIV

b) Invasive cervical cancer

c) Mycobacterium tuberculosis of any site

d) Oral candidiasis

Question 4:

A 47-year-old man with a diagnosis of acute myeloid leukemia with a blood type O negative blood group presents to the transplant clinic to discuss proceeding with an allogeneic stem cell transplant. Which of the following would be an optimal donor?

a) His identical twin brother

b) Umbilical cord transplant

c) His HLA identical 50-year-old brother who is otherwise healthy and is blood type O+

d) An HLA identical matched unrelated donor who is blood type O–

Question 5:

A 30-year-old HIV-positive man presents with fever for 3 weeks, dry cough, and significant weight loss. He complains of retrosternal chest pain that is worse on inspiration. His chest X-ray is given below. What is the most likely diagnosis?



a) Staphylococcal pneumonia

b) Pneumococcal pneumonia

c) Tuberculosis

d) Pneumocystis jirovecii pneumonia

Question 6:

All are true about Guillain-Barre syndrome except:

a) Inflammatory condition

b) Descending paralysis is seen

c) Plasmapheresis is a treatment method

d) Demyelinating disorder

Question 7:

Which of the following is the best treatment for the relapse of immune thrombocytopenic purpura?

a) IV immunoglobulins

b) Steroids

c) Splenectomy

d) Blood transfusion

Question 8:

Which of the following cardiac chambers is enlarged in mitral stenosis on chest X-ray initially?

a) Left atrium

b) Right atrium

c) Left ventricle

d) Right ventricle

Question 9:

Which among the following is false?

a) Components of MELD (Model for end stage liver disease) scoring system are creatinine, bilirubin, International normalized ratio (INR), serum sodium (mEq/L).

b) Components of CTP(Child Turcotte Pugh) score are albumin, bilirubin, INR, ascites, encephalopathy

c) Components of CTP (Child Turcotte Pugh) are creatinine, bilirubin, International normalized ratio (INR)

d) MELD score is used to assess and prioritize patients awaiting liver transplantation

Question 10:

Identify the false statement about the bundle of Kent.

a) It is faster than AV nodal pathway

b) It is slower than AV nodal pathway

c) Leads to short PR interval

d) Leads to prolonged QRS duration

Question 11:

A 40-year-old male patient came to ED with complaints of weakness, paresthesia, and breathing difficulty. Relevant investigations were done. The ECG obtained is suggestive of :



a) Hypokalemia

b) Hyperkalemia

c) Hypocalcemia

d) Hypercalcemia

Question 12:

Osborn waves in ECG are seen in:

a) Hypothyroidism

b) Hypothermia

c) Hypocalcemia

d) Hypokalemia

Question 13:

The production of which of the following hormones is increased in obesity?

a) Insulin

b) Thyroxine

c) Growth hormone

d) Adiponectin

Question 14:

What is the diagnosis of this patient from the ECG shown below?



a) Normal ECG

b) Ventricular fibrillation

c) Ventricular tachycardia

d) Misplaced leads

Question 15:

All are true about patients with trigeminal neuralgia except:

a) More common in females

b) Pain along V2 and V3 division of trigeminal nerve

c) Deep seated pain

d) No objective signs of sensory loss

Question 16:

All of the following are features of Crohn's disease except:

a) Transmural involvement

b) Lead pipe appearance

c) Rectal sparing

d) Perianal fistula

Question 17:

Which of the following is the most common type of Guillian-Barre syndrome?

a) Acute motor axonal neuropathy

b) Acute inflammatory demyelinating polyneuropathy

c) Acute motor sensory axonal neuropathy

d) Miller Fisher syndrome

Question 18:

Cushing's ulcers are seen in

a) Burns

b) Stress

c) Head injury

d) Cell necrosis

Question 19:

Which of the following is true about paroxysmal nocturnal hemoglobinuria?

a) Inherited defect in PIG –A

b) Extravascular hemolysis

c) Deficiency of CD 55 and CD 59

d) Microcytic anaemia

Question 20:

A patient with rheumatoid arthritis is being treated with methotrexate and low-dose corticosteroids for the past 4 months. However, the disease is still progressing. What would be your recommendation for the further management of this patient?

a) Stop oral methotrexate and start parenteral methotrexate

b) Add sulfasalazine and hydroxychloroquine

c) Continue corticosteroids and methotrexate

d) Start only anti-TNF alpha agents

Question 21:

A lady presents with fever, oral ulcers, photosensitivity, and rashes on her face as shown below. What is the most likely diagnosis?

a) Dermatomyositis

b) Systemic lupus erythematosus

c) Rosacea

d) Melasma

Question 22:

Which of the following murmur increases on standing?

a) HOCM

b) MR

c) MS

d) VSD

Question 23:

On performing a pulmonary function test, reduction in FEV1/FVC is characteristic of:

a) Restrictive disease

b) Obstructive disease

c) Normal lung function

d) Interstitial lung disease

Question 24:

Calcitonin levels are increased in:

a) Hyperthyroidism

b) Hypoparathyroidism

c) Hyperparathyroidism

d) Cushing's syndrome

Question 25:

Which of the following does not lead to ARDS (Acute Respiratory Distress Syndrome) at the blood-lung interface?

a) Pneumonia

b) Fat embolism syndrome following femur shaft fracture

c) Sepsis

d) Transfusion-related lung injury

Answer Key

Question No.	Correct Option
1	d
2	a
3	d
4	c
5	d
6	b
7	c
8	a
9	c
10	b
11	a
12	b
13	a
14	c

15	c
16	b
17	b
18	c
19	c
20	b
21	b
22	a
23	b
24	c
25	a

Detailed Explanations

Solution to Question 1:

The given clinical scenario describes an unconscious patient, who is hemodynamically unstable (systolic BP <90 mmHg or mean arterial pressure <65 mmHg). The ECG shows a narrow QRS complex tachycardia (heart rate of 188 BPM) with regular RR interval and absent P waves, characteristic of paroxysmal supraventricular tachycardia (PSVT). The next step in management is QRS-synchronized DC cardioversion.

Shown below is the algorithm for the management of supraventricular tachycardia:



Options A, B, and C are all used in stable cases. Vagal maneuvers like carotid massage or Valsalva are tried first followed by IV Adenosine. If these are ineffective, IV Verapamil, Diltiazem, or Beta blockers are used.

Synchronized cardioversion	Defibrillation
Used in sustained VTUsed in SVT (unstable cases), atrial fl utter/fibrillation	Used in VT and VF without a pulse
The shock delivered insync w ith R wave(in absolute refract ory period)	The shock is delivered at any time during cardiac cycle
Characteristics of QRS compl ex:Narrow and regular - 50 t o 100JNarrow and irregular - 120 to 200JWide and regul ar - 100JWide and irregular - No synchronised cardiovers ion	Wide and irregular QRS com plexMonophasic: 360 JBipha sic:1 20-200 J

Solution to Question 2:

Nasal polyps are associated with intrinsic asthma.

Intrinsic asthma is characterized by the presence of asthma with negative skin prick tests for common antigens. The association of nasal polyps with asthma is described by Samter's triad (intrinsic asthma + aspirin hypersensitivity + nasal polyposis). Asthma is usually severe, persistent, and adult-onset.

Other options:

Option B: Brittle asthma refers to a severe variant of asthma where patients have very serious and often, life-threatening attacks despite appropriate therapy.

Option C: Extrinsic asthma is also referred to as atopic asthma. It is allergen mediated.

Option D: Exercise-induced asthma is airway narrowing due to strenuous physical activity. Symptoms typically start after the exercise has ended and last for around 30 minutes.

Solution to Question 3:

Oral candidiasis, although seen in AIDS patients, is not an AIDS-defining illness. Candidiasis of the esophagus, trachea, bronchi, and lungs are AIDS-defining illnesses.

AIDS-defining illnesses are specific opportunistic illnesses in patients with AIDS, which signify an advanced stage of illness (Stage 3).

A case of HIV can be classified into five stages as per current CDC guidelines (0, 1, 2, 3, and unknown).

Stage	Criteria	CD4+ count (>5 years)
0	A negative HIV test within 6 months of first diagnosis	NA
1	Based on CD4+ count	>=500
2	Based on CD4+ count	200-499
3	AIDS-defining opportunistic illness	<200
unknown	Unavailable CD4+ counts	unknown

Solution to Question 4:

In the case of acute myeloid leukemia (AML), allogeneic stem cell transplants are the treatment of choice. His HLA identical 50-year-old brother who is blood type O positive would be the best choice.

Other options:

Option A: An identical twin brother would not be the donor of choice. The therapeutic benefit of stem cell transplant in hematological malignancies is due to the graft-versus-leukemia (GvL). Donor T lymphocytes attack malignant hematopoietic stem cells of the recipient that remain after high-dose chemotherapy. Despite having a low rate of graft versus host disease (GvHD), stem cells from an identical twin do not have a desirable degree of GvL effect, leading to higher rates of recurrence post-transplant.

Option B: Autologous (umbilical cord transplant) is not recommended as higher rates of relapse have been reported.

Option D: An HLA identical sibling is preferred over an unrelated HLA identical donor because there is a greater chance of matching for minor histocompatibility complex (mHC). This reduces the risk of GvHD. ABO compatibility is also important to consider, but a minor ABO mismatch (Rh mismatch) doesn't have any impact on hemolysis, graft-versus-host disease, or survival.

Solution to Question 5:

The most likely diagnosis in a case of pneumonia in a HIV positive patient, along with the presence of fever, and non-productive cough with radiographic findings of diffuse interstitial infiltrates and symmetric perihilar involvement is Pneumocystis jirovecii pneumonia (PCP). A characteristic feature of PCP is retrosternal chest pain, which worsens on inspiration and is described as sharp or burning.

It is important to note that even though the incidence of PCP is the highest at CD4+ counts of <200/µL (80%), it can be seen in patients with higher counts.

Other options:

Option D: Tuberculosis (TB) should be considered as a possibility in cases of pneumonia in people living with HIV. However, the absence of hilar lymph node involvement, non-productive cough, and no extrapulmonary (pleural, pericardial, lymphatic) features are not consistent with tuberculosis.

TB can occur at any stage of HIV infection. In the early stages, pulmonary TB appears typically with upper-lobe infiltrates and cavitation. In advanced HIV infection, with CD4+ T-cell count below 200/µL, TB presents with diffuse interstitial infiltrates, minimal cavitation, pleural effusion, and lymphadenopathy. Extrapulmonary TB is fairly common in HIV, with it being documented in 40-60% of cases.

Option A: Staphylococcal pneumonia most commonly presents with a productive cough, and radiologically presents with pneumatoceles (cavitations)

Option B: Patients with pneumococcal pneumonia also present with fever, dyspnea, and productive cough. Imaging usually shows lobar involvement.



Solution to Question 6:

Ascending (not descending) paralysis is seen in Guillain-Barre syndrome (GBS). It generally begins in the bilateral lower limbs (rubbery legs) affecting both proximal and distal muscles and ascends to involve the trunk and upper limbs.

GBS is an acute, autoimmune, inflammatory, demyelinating polyneuropathy. The usual presentation is symmetrical, ascending paralysis evolving over hours to a few days. There is usually no progression beyond 4 weeks. The legs are usually more affected than the arms. It is associated with autonomic dysfunction with or without sensory involvement. The most commonly involved cranial nerve is the facial nerve (50%). Lower cranial nerves are also frequently involved, causing bulbar weakness and difficulty in maintaining an airway and handling secretions. Deep tendon reflexes diminish or disappear within the initial days of symptom onset. Bladder dysfunction may occur in severe cases but is typically transient and not a prominent feature.

70% of GBS cases occur within 1-3 weeks after an acute infectious process, commonly respiratory or gastrointestinal infections. Campylobacter jejuni, human herpes viruses, and Mycoplasma pneumoniae have been implicated.

The diagnosis is based on Brighton criteria. It includes a combination of clinical, electrophysiologic, and CSF findings. Cytoalbuminologic dissociation (elevated CSF protein level AND CSF total white cell count <50 cells/µL) is characteristic of GBS.

Treatment should ideally commence as early as possible. Intravenous immunoglobulin (2g/kg bodyweight over 5 days) and plasmapheresis are both viable treatment options, with similar efficacy for typical GBS. Approximately 30% of GBS patients require ventilatory assistance. Glucocorticoids have not shown efficacy in GBS management.

Solution to Question 7:

Splenectomy is the treatment of choice in a case of immune thrombocytopenic purpura (ITP) relapse after first line therapy with glucocorticoids. The relapse usually occurs once they are tapered. Vaccination against encapsulated organisms (pneumococcus, meningococcus, H. influenzae) is advised before splenectomy is performed.

It is important to note that in some patients ITP resolves by itself, and observation along with intermittent IV immunoglobulins and anti - Rh(D) administration or administration of a thrombopoietin receptor agonist (eltrombopag, romiplostim) can be a viable option before splenectomy is considered.

The treatment of ITP is as follows:

ITP is an acquired disorder characterized by the immune-mediated destruction of platelets and possibly inhibition of platelet release from the megakaryocyte. Secondary causes of ITP include systemic lupus erythematosus (SLE), HIV-AIDS, and Hepatitis C.

ITP usually presents with mucocutaneous bleeding, very low platelet count, and normal peripheral blood cells. Patients typically exhibit ecchymoses and petechiae or they may incidentally discover thrombocytopenia during a routine complete blood count. Rarely, life-threatening bleeding may occur. The presence of wet purpura (blood blisters in the mouth) and retinal hemorrhages can serve as warning signs of potentially life-threatening bleeding.

	Clinical features	Management
1.	Minor bleeding manifestations (and/or)Platelet count > 50000/ µL	Prednisone(OR)Anti-D immunoglobulin therapy- effective only in Rh-positive patients (OR)Intravenous immunoglobulin (IVIG)
2.	Severe ITP (any of the following):Significant GI / intracranial bleedingPlatelet count < 50000/µLSigns of impending bleeding (retinal or large oral mucosal bleeds)	Platelet transfusion (AND)High-dose glucocorticoids (AND)IVIG / Anti-D immunoglobulins + Rituximab
3.	Relapse after first-line therapy	Splenectomy
4.	Unresponsive to all other therapy or relapse after splenectomy	Thrombopoietin receptor agonists -romiplostim(subcutaneous),eltrombopag(oral)

Solution to Question 8:

Left atrium (LA) is the first cardiac chamber to get enlarged in mitral stenosis (MS). It is due to the pressure overload of LA. It can be appreciated on the lateral and left anterior oblique views.

Other radiographic changes seen in MS are straightening of the upper left border of the cardiac silhouette, posterior displacement of the esophagus by an enlarged LA, and Kerley B lines due to distention of interlobular septae and lymphatics with edema.

Severe MS causing pulmonary hypertension leads to enlargement of the pulmonary artery, right ventricle, and right atrium. Left ventricle remains unaffected in isolated MS.

Solution to Question 9:

The Child Turcotte Pugh (CTP) score does not include creatinine.

The components of the CTP score are:

• Albumin

• Bilirubin

• International normalized ratio (INR)

• Ascites

• Encephalopathy

The Model for End-stage liver disease (MELD 3.0) is used to prioritize patients awaiting liver transplantation and includes:

• Female sex

• Bilirubin

• Albumin

• Serum sodium (mEq/L)

• INR (International Normalized ratio)

• Creatinine

Solution to Question 10:

The bundle of Kent is the accessory pathway of conduction between the atria and ventricle in Wolff-Parkinson-White syndrome. It conducts impulses faster than the AV nodal pathway and avoids AV nodal delay, leading to shortened PR interval. QRS is prolonged because of the AV nodal impulse reaching the ventricles after the aberrant impulse.

Solution to Question 11:

The symptoms of weakness, paresthesia, and dyspnea along with the given ECG showing depression of the ST segment, a decrease in the amplitude of the T wave (flattening or inversion), and an increase in the amplitude of U waves (at the end of the T wave) are characteristic of hypokalemia.

In the given ECG, leads V1-V6 exhibit T wave inversion. U waves are well appreciated in leads V2-V5.



Hypokalemia leads to dynamic ECG changes. The image given shows the sequence of changes observed:



Hypokalemia is defined as plasma K+ concentration of <3.5meq/L. It can also cause premature atrial and ventricular beats, sinus bradycardia, paroxysmal atrial or junctional tachycardia, atrioventricular block, and ventricular tachycardia or fibrillation.

Other options:

Option B: Hyperkalemia causes tall peaked T waves with a shortened QT interval. There is a progressive lengthening of the PR interval and QRS duration with an increase in potassium levels, eventually leading to the disappearance of the P wave; and widening of the QRS complex to a sine wave pattern.



Option C: Hypocalcemia causes prolongation of the QT interval. There is a lengthening of the ST segment, while the T wave (which correlates with the time for repolarization) remains unaltered.

Option D: Hypercalcemia causes shortening of the QT interval and a decrease in the ST segment duration.



Solution to Question 12:

Osborn waves are seen in hypothermia.

Osborn waves can also be seen in severe hypothyroidism (Option A), but even in those cases, hypothermia is the reason for the appearance of Osborn waves. Therefore, hypothermia is the better answer here.

It consists of an abrupt ascent right at the J point (the junctional point between the QRS complex and the ST segment) and then an equally sudden plunge back to the baseline. It is also called the J wave. This is because systemic hypothermia prolongs repolarisation.



Osborn wave can also be seen in:

• Hypercalcemia

• Brain injury

• Subarachnoid hemorrhage

• Vasospastic angina

• Cardiopulmonary arrest from oversedation

• Idiopathic ventricular fibrillation

Solution to Question 13:

Insulin levels are increased in obesity. It stimulates the synthesis of fatty acids and TAG and decreases the rate of lipolysis. Thyroxine, growth hormone, and adiponectin have lipolytic effects, hence their levels are reduced in obesity.

Hyperinsulinemia and insulin resistance are consistent features of obesity. Insulin resistance is more strongly linked to intra-abdominal fat than to fat in other areas.

Adiponectin (Option D) is an anti-inflammatory cytokine that is produced by adipocytes. It performs vital functions that are reduced in obesity - like, increasing insulin sensitivity and the inhibition of gluconeogenic enzymes. It also enhances fatty acid oxidation.

Insulin resistance in obesity occurs as a result of the following:

• Insulin receptor downregulation

• Increased FFA (free fatty acids) and capable of impairing insulin action

• Intracellular lipid accumulation

• Circulating peptides produced by adipocytes such as TNF-α, IL-6, RBP4, and resistin modify insulin action.

Solution to Question 14:

The rate of the given ECG is around 188 BPM (>100 BPM) and it shows wide QRS complexes (≥ 0.12 s) with abnormal morphology (red arrows) and absence of P waves which suggests a diagnosis of ventricular tachycardia.

Ventricular fibrillation (Option B) would not show discreet QRS complexes, and hence is not the answer in this case.



Solution to Question 15:

Trigeminal neuralgia is associated with superficial pain and not deep-seated pain.

It is more common (~ 60% of cases) in females (Option A) over 50 years and is characterized by unilateral lancinating facial pain. This most commonly involves the second and/or third divisions (Option B) of the trigeminal nerve territory. Bilateral or early-onset pain is associated with multiple sclerosis. Unilateral trigeminal nerve compression can be due to the superior cerebellar artery.

Pain is repetitive, severe, and brief and can be triggered by touch, cold, wind, or eating. Physical signs are usually absent, although the spasms may make the patient wince and sit silently (tic douloureux). No signs of sensory loss are present (Option D)

The drug of choice is carbamazepine.

Note: It is important to note that trigeminal neuralgia and trigeminal autonomic cephalalgia (TAC) are different entities. TAC is a group of primary headaches that include - cluster headaches, paroxysmal hemicrania, short-lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT)/short-lasting unilateral neuralgiform headache attacks with cranial autonomic symptoms (SUNA), and hemicrania continua.

Solution to Question 16:

Lead pipe appearance is seen in ulcerative colitis (not Crohn's disease). This appearance is due to the loss of haustra in the diseased colonic segment.

The image below shows the lead pipe appearance of the colon on barium enema:



Solution to Question 17:

Acute inflammatory demyelinating polyneuropathy (AIDP) is the most common type of GBS.

There are several types of Guillain–Barre syndrome, which are recognized by electrodiagnostic studies and pathological distinctions. These include:

A. Demyelinating

• Acute inflammatory demyelinating polyneuropathy (AIDP) - most common

B. Axonal

• Acute motor axonal neuropathy (AMAN)

• Acute motor–sensory axonal neuropathy (AMSAN)

C. Regional GBS syndromes

• Miller–Fisher syndrome (MFS)

• Pure sensory forms

• Ophthalmoplegia with anti-GQ1b antibodies as part of severe motor-sensory GBS

• GBS with severe bulbar and facial paralysis

• Acute pandysautonomia

Solution to Question 18:

Cushing's ulcers are seen in patients with head injury. They can also be seen in cases of space-occupying lesions, such as intracranial tumors. The most common areas affected are the acid-producing areas of the stomach (fundus and body).

Stress ulcers observed in severe burns patients are called Curling's ulcers. They are usually seen in the duodenal mucosa and are related to reduced perfusion and ischemia.

Stress-related gastric ulcers are seen in patients with massive burns, severe shock, sepsis, or head injury. Inflammatory mediators released in these conditions result in gastric mucosal inflammation. The most common presentation is minimal GI bleeding.

Solution to Question 19:

Deficiency of CD55 and CD59 is seen in paroxysmal nocturnal hemoglobinuria (PNH).

It is due to an acquired mutation in the X-linked phosphatidylinositol class A PIG-A gene (Option A) causing a deficiency of GPI anchor proteins. GPI is a glycolipid that anchors membrane protective factors such as CD59 and CD55 to the membrane. The deficiency of CD55 and CD59 makes the RBC sensitive to complement-mediated intravascular hemolysis(Option B). The anemia is usually normo-macrocytic (Option D).

Hemolysis tends to occur at night due to a slight decrease in blood pH during sleep, which triggers complement activity (nocturnal hemoglobinuria). PNH is characterized by hemolytic anemia, pancytopenia, and venous thrombosis.

PNH is diagnosed using flow cytometry. The drugs of choice for treatment are Eculizumab or Ravulizumab which work against complement C5. Supportive management of PNH includes folic acid and iron supplementation. Bone marrow transplantation is another option for the treatment of this condition.

Solution to Question 20:

For further management of a patient with rheumatoid arthritis (RA) with worsening disease activity, the preferred therapy is a step-up to oral triple therapy with methotrexate, sulfasalazine, and hydroxychloroquine. The addition of leflunomide or a biological (anti - TNF) to methotrexate is another alternative.

Ideally, the approach to a patient with early RA is aggressive. Methotrexate is always the first DMARD that the patient is started on, and if an adequate response is not observed within 3-6 months, a step up in therapy is made by adding on drugs.

For assessment of progression, composite indices like Disease Activity Score - 28 (DAS-28) can used.

Other options:

Option A: Changing the route of administration of methotrexate from oral to parenteral is not a part of step up

Option C: Priority is given to minimising the duration the patient is on steroids.

Option D: Anti -TNF agents (adalimumab, etanercept) are not recommended as a monotherapy, and are reserved for treatment in combination with DMARDs. This is due to the fact that the efficacy of monotherapy is lesser than that of DMARDs, and the adverse effects include increased risk of serious infections, re-activation of latent TB and opportunistic fungal infections. They are reserved for cases where there is a lack of response with the first line agents.

Solution to Question 21:

The given clinical case describes a young lady with fever, oral ulcers, and photosensitivity. The image shows a well-demarcated erythematous butterfly-shaped rash over the malar eminences with sparing of the nasolabial folds (malar rash). The clinical presentation along with the image showing malar rash point to a diagnosis of systemic lupus erythematosus (SLE).

The diagnosis of SLE is based on the presence of clinical and immunological manifestations.

Clinical manifestations:

• Skin: Acute, subacute cutaneous manifestations (photosensitivity, malar or maculopapular rash) and Chronic cutaneous manifestations (discoid lupus, panniculitis, lichen planus-like)

• Oral or nasal ulcers

• Non-scarring alopecia

• Synovitis involving ≥2 joints

• Serositis (pleurisy, pericarditis)

• Renal: Protein/Creatinine ratio >0.5; RBC casts; > Class II lupus nephritis on renal biopsy

• Neurologic: seizures, psychosis, myelitis, neuropathy, confusional state

• Hemolytic anemia

• Leukopenia (<4,000/mcL)

• Thrombocytopenia (<1,00,000/mcL)

Immunological manifestations:

• Antinuclear antibodies >reference value (positive in >98% patients)

• Anti-dsDNA >reference value (specific)

• Anti-Smith antibody (specific)

• Antiphospholipid antibody (lupus anticoagulant, false-positive rapid plasma reagin)

• Low serum complement (C3, C4)

According to the Systemic Lupus International Collaborating Clinics (SLICC) criteria, at least 4 out of 11 of the above criteria must be positive with at least one clinical and one immunological criteria well documented to diagnose SLE. However, if a renal biopsy is suggestive of lupus nephritis and lupus autoantibodies are positive, other criteria are not required.

Life-threatening manifestations include - lupus nephritis leading to end-stage renal disease, myocarditis, venous and arterial thrombosis (including stroke and myocardial infarction), ARDS, and vasculitis involving the gastrointestinal vasculature - leading to ischemia, bleeding, and sepsis.

NOTE: There are two classifications systems used to diagnose SLE:

• SLICC criteria, which mentions the clinical features is easier to use for clinical diagnosis.

• EULAR/ACR criteria which apply weightage to each clinical manifestation. The patient must have a positive ANA antibody titre of >1:80 by immunofluorescence, and a score of 10 to be diagnosed. The EULAR criteria is more recent and is commonly used for clinical studies.

Solution to Question 22:

The murmur of hypertrophic cardiomyopathy (HOCM) increases with standing, as there is a reduction in preload. In conditions of reduced preload, there is lesser end diastolic volume in the left ventricle, leading to a narrower outflow tract, and a louder murmur is heard.

The murmur heard in HOCM is a mid-systolic murmur, which is loudest along the left sternal border. It increases in conditions of decreased preload and afterload (standing, valsalva, vagal maneuvers) or conditions of increased contractility (inotropic stimulation) and decreases when preload and afterload are increased (squatting, vasopressors).

Most murmurs decrease in intensity with reduced preload and afterload except HOCM and mitral valve prolapse (MVP).

Solution to Question 23:

Reduction in FEV1/FVC is characteristic of obstructive lung disease.

Forced vital capacity (FVC) is the volume of expired air recorded when a person inhales to the total lung capacity and exhales as hard and as completely as possible. The volume of air expired in the first second is called the forced expiratory volume 1 (FEV1).

Given below is a graphical representation of FEV and FEV1:



FVC is reduced in both obstructive and restrictive lung diseases. In obstructive diseases, the slope of FVC is steady, therefore the FEV1 value is low and the FEV1/FVC ratio decreases. In restrictive diseases, the airflow is fast in the beginning leading to a higher value of FEV1 and FEV1/FVC ratio increases.

Solution to Question 24:

Calcitonin levels are increased in hyperparathyroidism.

Due to elevated parathyroid hormone (PTH) levels in hyperparathyroidism, there is osteoclast activation. This leads to increased calcium levels in plasma. As a response to this, there is an increase in the level of calcitonin to regulate the high calcium levels.

Calcitonin is secreted by the parafollicular C-cells of the thyroid gland. Its principal effects are to lower serum calcium and phosphate by its action on bone and kidney. In bone, calcitonin inhibits osteoclasts and reduces bone resorption. In the kidney, it inhibits the reabsorption of both calcium and phosphate.

Other options

Option A: In hyperthyroidism, the direct effect of thyroid hormones can lead to bone resorption and mild hypercalcemia. However, hyperparathyroidism is a better answer.

Options B and D: Hypoparathyroidism and Cushing's syndrome will have low serum calcium and hence, will not have increased calcitonin levels.

Solution to Question 25:

Pneumonia leads to Acute Respiratory Distress Syndrome (ARDS) at the air–lung interface and not at the blood-lung interface.

ARDS can be caused by direct lung injury at the air-lung interface and indirect lung injury at the blood-lung interface.

Causes at the air-lung interface (direct injury):

• Aspiration

• Toxic gas or smoke inhalation

• Near-drowning

• Pneumonia

• Pulmonary contusion

Causes at the blood-lung interface (indirect injury):

• Sepsis

• Transfusion-related lung injury (TRALI)

• Pancreatitis

• Long bone fractures causing fat embolism syndrome

• Thromboembolism

ARDS is a clinical syndrome of rapid onset severe dyspnea within one week of known clinical insult, hypoxemia, and diffuse pulmonary infiltrates leading to respiratory failure.

The recommended therapies in ARDS with clinical or supportive evidence include:

• Low tidal volume ventilation (LTVV) strategy (4–6 mL/kg of ideal body weight) is recommended to maintain adequate ventilation and avoid ventilator-induced lung injury by barotrauma and volume trauma.

• High positive end-expiratory pressure (PEEP) or open lung mechanical ventilation improves arterial oxygenation by improving ventilation perfusion mismatch.

• Prone position

• Extracorporeal membrane oxygenation (ECMO)

• The routine use of glucocorticoids is not beneficial.

The following X-ray shows bilateral alveolar infiltrates as seen in ARDS:

PORT
ET/OG

Medicine NEET 2020

Question 1:

A patient came to the hospital with complaints of lethargy, increased sleep, and weight gain. Investigations revealed low plasma TSH concentration. However, on the ■administration of TRH, the TSH levels increased. Which of the following conditions is likely in this patient?

a) Hyperthyroidism due to disease in the pituitary

b) Hypothyroidism due to disease in the pituitary

c) Hypothyroidism due to disease in the hypothalamus

d) Hyperthyroidism due to disease in the hypothalamus

Question 2:

What is the Child-Turcotte-Pugh class for the patient who has a serum bilirubin 2.5 mg/dl, serum albumin 3 g/dl, prothrombin time increased by 5 seconds (INR = 2), no encephalopathy, and mild ascites?

a) Class A

b) Class B

c) Class C

d) Class D

Question 3:

An elderly woman is brought to the OPD with complaints of behavioral change, a history of multiple falls, urinary incontinence, and dementia. What is your diagnosis?

a) Normal pressure hydrocephalus

b) Frontotemporal dementia

c) Parkinson disease

d) Creutzfeldt-Jakob disease

Question 4:

A 60-year-old male patient who is a known case of COPD presented with acute exacerbation and is admitted to the ICU. Which of the following statements is correct with regards to the initial management of this patient?

a) Non-Invasive PPV should be given

b) Invasive PPV should be given

c) IV corticosteroids should be administered

d) Permissive hypercapnia is allowed

Question 5:

Thrombocytopenia, eczema and recurrent infections are seen in:

a) Wiskott-Aldrich syndrome

b) Chediak-Higashi syndrome

c) Thrombocytopenia-absent radius syndrome

d) Hermansky-Pudlak syndrome

Question 6:

Which among the following is more common in limited cutaneous systemic sclerosis when compared to diffuse cutaneous systemic sclerosis?

a) Esophageal dysmotility

b) Myopathy

c) Interstitial lung disease

d) Scleroderma renal crisis

Question 7:

Water hammer pulse is seen in which of the following conditions?

a) Aortic stenosis

b) Aortic regurgitation

c) Aortic stenosis and aortic regurgitation

d) Mitral regurgitation

Question 8:

The arterial blood gas analysis findings of a patient are as follows: pH= 7.30, HCO_3= 10 mEq/L, $paCO_2$= 30 mmHg, which is suggestive of a partially compensated metabolic acidosis. What is the mechanism for the secondary disorder?

a) Hypoventilation causing decreased CO_2 washout

b) Hyperventilation causing increased CO_2 washout

c) Increased tubular reabsorption of HCO_3-

d) Increased excretion of H+

Question 9:

Which of the following is seen in pituitary apoplexy?

a) Hypertension

b) Shock

c) Unconsciousness

d) Fatigue

Question 10:

A 20-year-old man presents to the emergency department with respiratory distress and hypotension following a trauma. He has subcutaneous emphysema and there is an absence of air entry into the right side of the lungs. What is the next best step?

a) Start IV fluids with large bore cannula

b) Needle decompression into the 5th intercostal space anterior to the midaxillary line

c) Take the patient to the ICU and intubate him

d) Start positive pressure ventilation

Question 11:

Which of the following is true about polyarteritis nodosa?

a) Necrotising inflammation of large vessels

b) Patient has hypogammaglobulinemia

c) 90% is associated with ANCA positivity

d) 30% is associated with Hepatitis B

Question 12:

Which type of aphasia is seen in the lesions of the posterior part of the superior temporal gyrus?

a) Fluent

b) Non fluent

c) Anomic

d) Conduction

Question 13:

A 70-year-old man with a known history of DM and HTN on regular medications presents with a history of progressively worsening dyspnoea on exertion. On examination, there is elevated JVP, positive hepatojugular reflux, crackles at the lung bases, hepatomegaly, ascites, and pedal oedema■. What is your provisional diagnosis?

a) Heart failure

b) Hypertrophic cardiomyopathy

c) Mitral regurgitation

d) Portal hypertension

Question 14:

The slowing of conduction in multiple sclerosis is due to which of the following?

a) Gliosis

b) Loss of myelin sheath

c) Defect at node of Ranvier

d) Leaky Na+ channel

Question 15:

A 15-year-old girl presented with fatigue, chronic diarrhoea, weight loss, bone pain, and abdominal distension. Investigations revealed iron deficiency anemia and osteoporosis.

Which of the following is the single best test to be done for her evaluation?

a) TSH levels

b) C-peptide levels

c) Urine sugar and ketone

d) IgA tissue transglutaminase antibody

Question 16:

A 55-year-old male presented with tachypnea and mental confusion. His blood sugar is 350mg/dl, pH- 7.2, HCO3-10, pCO2-30. What is the primarymetabolic abnormality?

a) Metabolic alkalosis

b) Metabolic acidosis

c) Respiratory alkalosis

d) Respiratory acidosis

Question 17:

A ten-year-old boy was brought to the OPD with fever and swelling of hands after playing football. He gives a history of previous episodes of swelling of hands. On imaging, his spleen was found to be shrunken. Which of the following is a likely diagnosis?

a) Sickle cell anaemia

b) Malaria

c) Pancreatitis

d) Measles

Question 18:

A 53-year-old woman underwent hip replacement surgery. A week after the surgery, the patient developed swelling of the legs associated with pain on palpation. Her heart rate is 70 beats per min. There is no history of hemoptysis or significant weight loss. There is no previous history of pulmonary embolism. What is the risk of developing pulmonary embolism in the patient based on Well's score?

a) Low

b) High

c) Moderate

d) Cannot comment without d-dimer values

Question 19:

The recommended temperature of water bath for a patient with frost bite is

a) 37 degree C

b) 42 degree C

c) 32 Degree C

d) 30 degree C

Question 20:

A mutation in the gene for aquaporin channel leads to which of the following conditions?

a) Liddle's syndrome

b) Nephrogenic diabetes insipidus

c) Cystic fibrosis

d) Bartter syndrome

Question 21:

Which of the following is not a radiological finding in a patient with left heart failure?

a) Kerley b lines

b) Focal oligemia

c) Increased venous blood in lung

d) Change in upper lobe circulation

Question 22:

A 60-year-old man presented with an episode of acute onset dizziness and loss of consciousness which lasted for a few seconds followed by a regain of full consciousness. There were no similar episodes in the past. Which of the following is a true statement regarding this scenario?

a) ECG to rule out atrial fibrillation

b) If ECG is normal, CT scan should be done

c) Tilt-table testing

d) Vestibular neuritis is a possible condition that can cause these symptoms

Question 23:

A 12-year-old child who is known to have type 1 diabetes mellitus presents with confusion and drowsiness. Her mother says that she seems to be breathing very fast. On examination, mucous membranes are dry and blood pressure is 70/50 mm Hg. Random blood glucose is 415 mg/dl and urine ketones are 4+. What is the next best step in management?

a) 2-3 L of normal saline over 1-3 hours

b) Insulin infusion at 0.1 units/kg/hour

c) Arterial blood gas

d) Insulin bolus of 0.1 units/kg given IV

Answer Key

Question No.	Correct Option
1	c
2	b
3	a
4	a
5	a
6	a
7	b
8	a
9	b
10	b
11	d
12	a
13	a
14	b
15	d
16	b

17	a
18	b
19	a
20	b
21	b
22	a
23	c

Detailed Explanations

Solution to Question 1:

The presenting symptoms of lethargy, increased sleep, and weight gain are suggestive of hypothyroidism. An increase in TSH on administration of TRH confirms the lack of stimulus for TSH secretion from the hypothalamus.

Although decreased TSH levels can be seen in hyperthyroidism, clinical features such as lethargy, increased sleep and weight gain are not. This pushes the diagnosis in favor of hypothyroidism.

If the disease was in the pituitary, there would be no increase in TSH secretion on administration of TRH (Option B).

Normally, the thyrotropin releasing hormone (TRH) is released by the hypothalamus. It stimulates the pituitary gland to release the thyroid-stimulating hormone (TSH), which in turn stimulates the thyroid gland to produce T3 and T4 hormones.

Low TSH levels may be seen in the following cases:

Disease	Pathogenesis	Presentation
Hyperthyroidism	Feedback inhibition of TSH r elease	Symptoms of hyperthyroidis m
Primary disease of the pituita ry	Decreased release of TSH, th ereby causing a decreased rel ease of thyroid hormones	Symptoms of hypothyroidism. Administration of TRH does not elevate the TSH levels in this case since the pituitary g land is abnormal
Primary disease of the hypot halamus	Decreased release of TRH lea ding to decreased TSH and h ypothyroidism	Administration of TRH cause s elevation of TSH levels sinc e the pituitary gland is norm al.



Solution to Question 2:

The given patient belongs to Child-Turcotte-Pugh class B.

The score can be calculated as follows:

Serum bilirubin-2.5mg/dl: 2

Serum albumin-3g/dl: 2

Prothrombin time-5 seconds or INR-2: 2

No encephalopathy: 1

Mild ascites: 2

Therefore, the Child-Turcotte-Pugh score is (2+2+2+1+2)= 9 which is included in Class B.

Solution to Question 3:

The given case scenario describes an elderly woman with urinary incontinence, dementia (cognitive decline), and a history of multiple falls (suggestive of a gait abnormality or ataxia). These form the classical triad of normal pressure hydrocephalus (NPH).

Both frontotemporal dementia (Option B) and Parkinson's disease (Option C) are common causes of dementia associated with falls, but urinary incontinence is not commonly appreciated. NPH is, therefore, the most likely diagnosis.

Normal pressure hydrocephalus refers to a disproportionate enlargement of the ventricles compared to the degree of cortical atrophy. It is a communicating hydrocephalus caused by delayed resorption of cerebrospinal fluid. Out of the classical triad of symptoms, most patients present with one or two. Hence, neuroimaging showing dilated ventricles is essential, as well as an L- dopa challenge test to rule out parkinson's disease is recommended.

A characteristic MRI of NPH is given below, showing dilated lateral ventricles which are out of proportion to sulcal deepening:



Solution to Question 4:

Non-invasive positive pressure ventilation (NIPPV) should be the first mode of ventilation used in COPD patients with acute respiratory failure provided there are no absolute contraindications. This approach improves gas exchange, reduces work of breathing and the need for intubation, decreases hospitalization duration and enhances overall survival outcomes.

Invasive positive pressure ventilation is only considered after failure of NIPPV, resulting in severe respiratory distress, life threatening hypoxemia, severe hypercarbia and/or acidosis, markedly impaired mental status, respiratory arrest or hemodynamic instability (Option B).

Apart from NIPPV, short-acting inhaled beta 2 agonists (albuterol, salbutamol) with or without short-acting anticholinergics (ipratropium) and Long-acting anticholinergics (tiotropium) are recommended as the initial bronchodilators. Maintenance therapy with long-acting beta 2 agonists (formoterol) must be started as soon as possible.

Systemic corticosteroids are useful in the treatment of acute exacerbations of COPD as they reduce airway inflammation and improve lung function over time. However, the effects are not immediate. NIPPV on the other hand provides immediate relief to the patient and reduces the work of breathing. Therefore, it would not be the best choice in this case. (Option C)

Antibiotics can also be considered, as they reduce duration of hospitalisation. Treatment is given for 5 days.

Permissive hypercapnia refers to a ventilation strategy where tidal volume is reduced in order to avoid hyperinflation-associated lung trauma. It is not preferred at the initial presentation. It is only permitted once invasive PPV is started following NIPPV failure (Option D)

An exacerbation of COPD is defined as an event characterized by dyspnea and/or cough and sputum that worsen over < 14 days. Exacerbations of COPD are often associated with increased

local and systemic inflammation caused by airway infection, pollution, or other insults to the lungs.

Solution to Question 5:

Thrombocytopenia, eczema and recurrent infections are characteristically seen in Wiskott-Aldrich syndrome.

Wiskott Aldrich syndrome is an X-linked recessive disorder due to mutations in the WASP gene. The characteristic triad of features is:

• Eczema

• Thrombocytopenia - along with decreased platelet size, impaired platelet aggregation response

• Immunodeficiency - The immunoglobulin pattern seen is low level of IgM, elevated IgA and IgE, and a normal or slightly low IgG. Defect in regulatory T cells results in recurrent bacterial infections.

Other options:

Option B: Chediak-Higashi syndrome (CHS) is an autosomal recessive disease due to mutations in CHS1/LYST gene (defective lysosomal transport). It leads to recurrent pyogenic infections, partial oculocutaneous albinism, neurologic abnormalities, and thrombocytopenia. Peripheral smear shows large azurophilic granules in neutrophils and eosinophils. It can lead to hemophagocytic lymphohistiocytosis.

Option C: Thrombocytopenia-absent radius syndrome is an autosomal recessive disorder with severe thrombocytopenia (diagnosed at birth) and bilateral absent radii (thumbs are always present).

Option D: Hermansky-Pudlak syndrome is an autosomal recessive disorder with oculocutaneous albinism, bleeding diathesis (deficient dense granules in platelets), pulmonary fibrosis, and granulomatous colitis.

Solution to Question 6:

Esophageal dysmotility is more commonly seen in limited cutaneous systemic sclerosis (lcSSc) compared to diffuse cutaneous systemic sclerosis.

Systemic sclerosis is divided into two based on the patterns of involvement - diffuse and limited.

Limited cutaneous systemic sclerosis is characteristically associated with CREST syndrome, which includes calcinosis cutis, Raynaud's phenomenon, esophageal dysmotility, sclerodactyly and telangiectasia.

Skin involvement, interstitial lung disease (Option C), myopathy (Option B), and scleroderma renal crisis (Option D) are more common in diffuse cutaneous systemic sclerosis (dcSSc) than limited cutaneous systemic sclerosis.

The incidence of ischemic digital ulcers, CREST features, and primary biliary cirrhosis is more in limited cutaneous systemic sclerosis. The frequency of pulmonary arterial hypertension (PAH) is equal in both types.

Solution to Question 7:

Water hammer pulse, also known as Corrigan's pulse, is characterized by a sharp rise and a rapid fall-off and is seen in aortic regurgitation (AR) due to the fact that arterial pressure falls rapidly in late systole and early diastole.

In AR, the pulse pressure is widened due to systolic hypertension and decreased diastolic pressure. The diastolic pressure is measured at Korotkoff phase IV. In certain patients with AR or a combination of aortic stenosis (AS) and AR, the carotid arterial pulse may exhibit a bisferiens pattern, characterized by two systolic waves separated by a trough.

The apical impulse is heaving and displaced inferiorly. A diastolic thrill may be palpable along the left sternal border.

A2 is usually absent in AR. An early diastolic murmur can be auscultated in chronic AR. It is best heard in the third intercostal space along the left sternal edge.



Severe AS (Option A) would show pulsus parvus et tardus, characterized by the carotid pulse rising slowly to a delayed peak.

Chronic mitral regurgitation (Option D) shows a sharp, low volume upstroke due to reduced cardiac output and acute mitral regurgitation demonstrates narrow pulse pressure.

Solution to Question 8:

The mechanism for the secondary disorder is hypoventilation causing decreased CO2 washout.

The given arterial blood gas findings are suggestive of a partially compensated metabolic acidosis. The compensation is partial because the fall in paCO2 much lower than what is expected for the given HCO3 levels. The fall in paCO2 is inadequate due to hypoventilation causing decreased CO2 washout. In this case, the secondary disorder is respiratory acidosis.

Hyperventilation causing increased CO2 washout (Option B) would allow for better compensation, with lower paCO2 levels than observed.

The expected paCO2 can be calculated using Winter's formula:

For metabolic acidosis, the expected paCO2 = [1 .5 X HCO3-] + 8 ± 2.

Substituting the values, expected PaCO2 = (1.5 x 10) +8 ± 2

= 15+8 ±2

= 21 to 25 mmHg

However, the actual PaCO2 is 30 mmHg, which is higher than the expected paCO2 indicating a secondary respiratory acidosis. A possible explanation for this could only be hypoventilation causing decreased CO2 washout.

Thus, the given ABG findings are suggestive of a mixed acid-base disorder characterized by metabolic acidosis with additional respiratory acidosis.

For example, in a patient with diabetic ketoacidosis and pneumonia, the lungs would fail to compensate, and the patient could develop hypoventilation leading to secondary respiratory acidosis.

Solution to Question 9:

Shock is seen in pituitary apoplexy.

Loss of consciousness (Option C) can also be seen but it occurs secondary to adrenal insufficiency and shock. Hence, shock is the most appropriate answer.

Pituitary apoplexy refers to acute hemorrhagic infarction of the pituitary gland. This results in the compression of surrounding structures and hypopituitarism leading to acute adrenal insufficiency. It may be seen spontaneously in a pre-existing pituitary adenoma, post-partum (Sheehan's syndrome), or be associated with hypertension, diabetes mellitus or sickle cell anemia or shock.

Pituitary apoplexy is an endocrine emergency that can lead to severe hypoglycemia, hypotension, shock, central nervous system (CNS) hemorrhage, and death. Acute clinical manifestations commonly encompass intense headaches with signs of meningeal irritation, bilateral visual changes, ophthalmoplegia, and even cardiovascular collapse.

The investigation of choice is MRI.

Patients who do not exhibit overt visual impairment or altered mental status can be managed conservatively with high-dose glucocorticoids. Progressive visual loss, cranial nerve palsy, or altered mental status necessitate expeditious surgical decompression.

Solution to Question 10:

The given clinical scenario of a young male developing respiratory distress with hypotension, subcutaneous emphysema, and unilaterally absent air entry is suggestive of tension pneumothorax. The immediate next step, in this case, would be needle decompression at the fifth intercostal space anterior to the mid-axillary line (as the patient is an adult).

In children, the second intercostal space in the midclavicular line is chosen as the site for needle decompression.

Needle decompression turns a tension pneumothorax into an open pneumothorax. It has to be followed by tube thoracostomy. An intercostal chest drain (ICD) tube is inserted in the triangle of safety, just anterior to the midaxillary line on the side of the lung injury. The site of insertion is the same in both children and adults.

The triangle of safety is bounded by the lateral border of the pectoralis major anteriorly, the anterior border of the latissimus dorsi posteriorly, and the fifth intercostal space inferiorly, with the apex below the axilla. This provides a safe landmark for the insertion of an ICD with minimal risk of tissue injury.

Tension pneumothorax is a condition where air accumulates in the pleural space due to a one-way valve air leak, leading to the collapse of the affected lung. This can cause a mediastinal shift, compression of the opposite lung, and decreased venous return, resulting in obstructive shock. It commonly occurs during positive-pressure mechanical ventilation, but can also happen after chest trauma, subclavian/internal jugular venous catheter insertion, or chest wall defects. The diagnosis is clinical, and immediate treatment is necessary without waiting for radiological confirmation.

Tension pneumothorax is characterized by chest pain, tachypnoea, tachycardia, hypotension, tracheal deviation away from the side of injury, absent breath sounds on the affected side, cyanosis, and neck vein distention. Patients can also develop subcutaneous emphysema as a complication secondary to the pneumothorax.

Solution to Question 11:

One-third of polyarteritis nodosa (PAN) cases are associated with Hepatitis B.

Other options:

Option A: Polyarteritis nodosa is a systemic vasculitis which is associated with necrotizing inflammation of small and medium-sized muscular arteries.

Option B: Patients show hypergammaglobulinemia.

Option C: PAN is rarely associated with ANCA positivity. There is no serological test for PAN. Most patients exhibit an elevated leukocyte count with a predominance of neutrophils.

PAN is usually seen in young adults, characterized by rapidly accelerating hypertension, abdominal pain, bloody stools, myalgias, and peripheral neuropathy.

Arterial inflammation is segmental, necrotizing, and transmural, leading to aneurysms and thrombosis.

The involved vessels are typically found at arterial bifurcations, spreading to veins but sparing venules. The most commonly affected organs are the kidney, heart, liver, and gastrointestinal tract, in that order. Pulmonary arteries are not involved. Inadequate blood flow leads to ischemia, ulcers, infarcts, atrophy, or hemorrhage.

Diagnosis relies on histopathological findings from organ biopsies or the demonstration of aneurysms in small to medium-sized arteries of the renal, hepatic, and visceral vasculature.

Untreated PAN has an extremely poor prognosis, with a 5-year survival rate of only 10 to 20%. Common causes of death include gastrointestinal complications, such as bowel infarcts and perforation, as well as cardiovascular issues. Timely treatment is vital for improved outcomes.

Combining prednisone and cyclophosphamide has shown significant benefits in survival rates. Glucocorticoids alone may be sufficient for milder cases. In PAN cases associated with hepatitis B or C, antiviral therapy is combined with glucocorticoids and plasma exchange. Managing hypertension effectively can help reduce vascular complications. Despite successful treatment, relapses occur in approximately 10-20% of patients.

A histopathological image of PAN showing necrotizing transmural inflammation is given below:



The image below shows aneurysms in renal vasculature:



Solution to Question 12:

Lesions of the posterior part of the superior temporal gyrus cause Wernicke's or Fluent aphasia due to the involvement of the Wernicke's area. The speech is full of jargon and neologisms that make little sense (area no 22).

Broca's (Option B) or motor/non-fluent aphasia is seen in lesions of the inferior frontal gyrus which contains the motor speech area. Speech is non-fluent and interrupted, with a paucity of functional words.

Comprehension of written language and the ability to name objects are compromised in anomic aphasia (Option C). This aphasia is caused by damage to the angular gyrus that disrupts visual information processing and transmission to the Wernicke area.

Conduction aphasia (Option D) is seen in lesions of the arcuate fasciculus, the bundle of nerve fibers connecting the Wernicke and Broca's areas. Repetition of words is impaired and paraphasias are observed, though not as much as in Wernicke aphasia.

Solution to Question 13:

The given symptoms and signs are suggestive of heart failure (Option A).

Portal hypertension (Option D) can be a result of right-sided heart failure, however, the presence of worsening dyspnea along with the other symptoms implies a primary cardiac condition.

Heart failure is a common endpoint for most chronic cardiac conditions such as hypertension, valvular heart disease, and coronary heart disease.

The symptoms of heart failure are commonly due to volume overload with pulmonary and/or systemic venous pressures.

RIGHT HEART FAILURE	LEFT HEART FAILURE
Symptoms occur due tosystemic congestion	Symptoms occur due topulmonary congestion
Raised pulsatile JVPHepatic congestionTender hepatomegalyPitting pedal edemaAscitesEarly satiety	Shortness of breathOrthopnea (dyspnea that occurs within 1-2 mins of lying down)Paroxysmal nocturnal dyspnea(shortness of breath that develops after lying down for 30 mins or longer)Pulmonary rales

Solution to Question 14:

In multiple sclerosis, the slowing of conduction is due to the loss of the myelin sheath.

Under normal circumstances, nerve conduction in myelinated axons occurs in a saltatory manner i.e from one node of Ranvier to another. But, when demyelination sets in, the nerve impulse are unable to travel as fast due to not being saltatory. This produces either a conduction block or delay.

Lesions due to multiple sclerosis begin with perivenular cuffing by inflammatory mononuclear cells. Myelin-specific autoantibodies cause degeneration of myelin sheaths. As lesions evolve, prominent astrocytic proliferation (gliosis) is observed.

Demyelination is the hallmark in the pathology of multiple sclerosis. But, it is not solely a disease affecting the myelin sheath. Partial or total axonal destruction (neuronal loss) can also occur.

Solution to Question 15:

The presence of fatigue, chronic diarrhea, weight loss and abdominal distention along with investigations showing features of malabsorption (iron deficiency) lead to the diagnosis of celiac disease or gluten sensitive enteropathy. The single best test for which is checking serum for the presence of IgA tissue transglutaminase (anti-TTG) antibodies (Option D).

Celiac disease is a disorder characterized by an immunological response to gluten causing damage to the proximal intestinal mucosa (villous atrophy) and malabsorption of nutrients.

Symptoms usually begin 6 to 24 months from the introduction of gluten into diet, although most patients go undiagnosed.

The common manifestations are - irritability, abdominal distention, chronic diarrhea, weight loss or muscle wasting and extra-intestinally, the symptoms observed are aphthous ulcers, delayed puberty, iron deficiency anemia and short stature.

Screening for celiac disease is by testing for tissue transglutaminase IgA, endomysial, and deamidated antigliadin antibodies. IgA levels in serum are measured to detect false-negative results because of IgA deficiency.

Small intestinal biopsy shows characteristic villus blunting, crypt hyperplasia, and increased intraepithelial lymphocytes.

The treatment involves the exclusion of gluten-containing food from diet.

Long term complications observed are osteoporosis, anemia, infertility and malignancy (enteropathy related-T cell lymphoma is the most common association)

The Marsh classification is used to categorize celiac disease into different types and to quantify severity of disease involvement.

Solution to Question 16:

The patient has a primary metabolic acidosis with partial respiratory compensation (Option B)

Step 1: Observe pH - Patient is in acidosis.

Step 2: Observe the HCO_3 value. The normal value of bicarbonate is 22 - 28 mEq/L, which means that this patient has low levels (10mEq/L) . This is indicative of a primary metabolic acidosis.

In case of a primary respiratory acidosis, the pCO_2 values would have been greater than normal (35 - 45 mmHg). Hence, the answer cannot be Option D.

Step 3: Calculate the expected compensatory fall in pCO_2 using Winter's formula :

$$\text{Expected } PaCO_2 = [1.5 \times HCO_3^-] + 8 \pm 2$$

$$= (1.5 \times 10) + 8 \pm 2$$

$$= 23 \pm 2 \text{ mEq/L}$$

This implies that the levels in the patient are higher than expected - suggestive of a partially compensated metabolic acidosis.

Note: The likely cause of the condition is diabetic ketoacidosis, as the patient has high blood sugar levels.

Solution to Question 17:

The findings of episodes of swelling of hands (dactylitis) along with the shrunken spleen (auto splenectomy or splenic infarction) are suggestive of sickle cell disease. Moreover, the symptoms were seen after playing football, which acts as a predisposing factor to sickling to which dehydration, infection and acidosis are the triggers.

Sickle cell anemia is an autosomal recessive disorder that occurs due to the point mutation in the β-globin gene, which results in the replacement of glutamate residue with valine. This causes the production of sickle-shaped red blood cells due to the polymerization of deoxygenated hemoglobin (HbS).

The cardinal clinical feature of sickle cell anemia is acute vaso-occlusive pain, due to ischemia of the tissues. Vaso-occlusive crisis can cause dactylitis, splenic infarction (shrunken spleen over time) and avascular necrosis (AVN).

Most commonly, the femoral head is affected in AVN. Other sites affected include the humeral head and mandible.

Dactylitis, referred to as a hand-foot syndrome, is often the first manifestation of pain in infants and young children with sickle cell anemia. Dactylitis often manifests with symmetric or unilateral swelling of the hands and/or feet.

Other manifestations include:

• Acute splenic sequestration due to venous obstruction, needing emergency transfusion/splenectomy

• Painful ischemic crises due to vaso-occlusion in connective and musculoskeletal tissue

• Renal papillary necrosis leading to isosthenuria

• Cerebrovascular strokes

• Chronic lower leg ulcers

• Priapism in males

Solution to Question 18:

The risk of developing pulmonary embolism in this patient is high (Option B) according to the three-tiered Well's clinical risk assessment score.

Calculation of Well's score:

• Clinical signs of DVT = 3; - swelling of legs, pain on palpation (Moses sign)

• No alternative diagnosis = 3

• Heart rate is <100 BPM = 0; - 70 BPM

• Surgery in previous 4 weeks = 1.5; - hip replacement surgery 1 week ago

• No previous pulmonary embolism = 0

• No Hemoptysis = 0

• No malignancy = 0; - no significant weight loss

Total score = 7.5

Risk stratification:

• High risk = > 6.0

• Moderate risk = 2.0 to 6.0

• Low risk = <2.0 (in patients with low risk, to further confirm the risk - pulmonary embolism rule-out criteria (PERC) may be referred)

Note: There is no other alternative diagnosis which is more likely in this condition, where there is a clear clinical presentation of a deep vein thrombosis, with the associated risk factors of a recent surgery and immobilization. Hence, 3 points can be added to the score.

Solution to Question 19:

The temperature of the water bath for a patient with frostbite should be maintained between 37-40°C (Option A). Circulating water is preferred for treatment.

In frostbite, tissue damage results from severe environmental cold exposure or from direct contact with a very cold object. Tissue injury results from both freezing and vasoconstriction. It usually affects the distal aspects of the extremities or exposed parts of the face, such as the ears, nose, chin, and cheeks.

The treatment includes - extraction from cold environment, rapid rewarming, analgesics for pain during thawing, antithrombotics or thrombolytics, cleaning of the injured area with soap or antiseptics, tetanus prophylaxis, monitoring for compartment syndrome and finally, elective late amputation after imaging in severe cases.

Solution to Question 20:

A mutation in the gene for aquaporin 2 channel leads to nephrogenic diabetes insipidus.

The antidiuretic hormone (ADH) binds to V2 receptors in the late distal tubules, collecting tubules, and collecting ducts, and stimulates the movement of an intracellular protein, called aquaporin-2 (AQP-2), to the luminal side of the cell membranes.

Diabetes insipidus (DI) is a condition in which there is the absence of or poor response to anti-diuretic hormone (ADH). This results in reduced permeability of the distal tubules and collecting ducts to water. This furthermore leads the kidneys to excrete large amounts of dilute urine that clinically manifests as polyuria and polydipsia.

DI can be of four types :

Note: Lithium - used to treat bipolar disorder, can cause acquired diabetes insipidus by decreasing the expression of the AQP2 gene.

Option A : Liddle's syndrome is due to an inherited gain of function mutation of genes regulating the Na channel in collecting duct ENaC, characterized by hypertension, hypokalemia, and metabolic alkalosis.

Option C : Cystic fibrosis is an autosomal recessive disorder due to mutation in CFTR gene that serves as a channel in the luminal plasma membranes of epithelial cells and regulates volume and composition of secretions in exocrine glands.

Option D : Bartter syndrome is due to loss-of-function mutations of transporters and ion channels in thick ascending loop of henle, most commonly, Na-K-2Cl transporter.

Types	Primary pathology	Caused by
Pituitary/ Central DI	Primary deficiency of ADH se cretion	Mc defect: AVP -Neurophysin II gene
Primary polydipsia	Suppression of ADH secretio n by excessive fluid intake	1. Dipsogenic DI : Following head trauma, TB meningitis, Multiple sclerosis.2. Psychog enic DI : Due to psychosis or OCD3. Iatrogenic polydipsia : Due to motivated health ben efits
Gestational DI	Deficiency due to Increased d egradation of ADH	N terminal aminopeptidase p roduced by placenta
Nephrogenic DI	Renal insensitivity to ADH or drugs or genetic mutation	Drugs like lithium - gene enc oding V2 receptor on chXq28

Solution to Question 21:

Focal oligemia is not a radiological feature of left heart failure. It is seen in pulmonary embolism and is also called Westermark's sign.



It is defined by a hyperlucent area with decreased vascularity due to oligemia of the involved part of the lung. It is caused by a cutoff of circulation due to an embolus, leading to the loss of visibility of a patch of pulmonary vasculature on an X- ray.

Radiological signs of left heart failure include :



• Cardiomegaly - enlargement of the cardiac silhouette (increased cardiothoracic ratio)

• Upper lobe pulmonary venous congestion (cephalisation of vessels)

• Interstitial edema.

• Kerley B lines - horizontal lines in the periphery of the lower posterior lung fields as shown below.



Other radiological signs of Pulmonary embolism :

• Hampton's hump - a peripheral wedged-shaped density usually located at the pleural base

• Palla's sign - an enlarged right descending pulmonary artery

Solution to Question 22:

The clinical scenario of transient loss of consciousness in this patient is suggestive of syncope. Hence, an ECG must be done to rule out cardiac causes for syncope like - arrhythmias (atrial fibrillation, sinus node dysfunction, supra ventricular tachycardias) or a structural cardiac abnormality.

A CT would only be required if a seizure or transient ischemic attack (TIA) is suspected. However, in this case - the episode only lasted for a few seconds, the patient has no history of similar episodes in the past and there are no lasting neurological deficits after the episode. Therefore, option B is incorrect.

Syncope is a transient loss of consciousness caused by global cerebral hypoperfusion. It can be due to multiple causes, but is broadly divided into - neurally mediated syncope, orthostatic hypotension and cardiac syncope.

The initial workup for a case of syncope includes:

• Detailed history of the episode and past history of similar episodes

• Physical examination mainly to check BP in standing and supine positions to

• rule out orthostatic hypotension.

• A resting ECG to look for cardiac arrhythmia

• Option B: If the ECG is normal, no further tests are needed. If the ECG is suggestive of cardiac arrhythmia, then further tests including continuous ECG monitoring and electrophysiological studies may be done.

Option C: Tilt-table testing is done in patients with recurrent syncopal attacks who are suspected to have vasovagal syncope. Psychiatric evaluation is done in patients with recurrent unexplained syncopal attacks.

Option D: Vestibular neuritis is associated with vertigo, where the patients typically complain of dizziness and a sense of spinning of the head.

Solution to Question 23:

The given clinical scenario is suggestive of diabetic ketoacidosis (DKA) and the first step is to assess the arterial blood gas (ABG) levels. This precedes correction of fluid balance, as assessment of the current status of the patient is necessary before beginning treatment. A baseline pH and electrolyte level must be obtained.

Diabetic ketoacidosis is the most serious complication of type 1 diabetes mellitus. The symptoms include dehydration, nausea, vomiting, confusion, drowsiness, and coma in severe cases.

In DKA, the insulin-to-glucagon ratio is decreased which promotes gluconeogenesis, glycogenolysis, and ketone body formation in the liver. Ketone bodies exist as ketoacids at physiological pH which are neutralized by bicarbonate. As bicarbonate stores are depleted, metabolic acidosis ensues. Increased lactic acid production also contributes to the acidosis.

The diagnostic biochemical criteria for DKA includes:

• Hyperglycemia (>250 mg/dL)

• Venous pH <7.3

• Plasma/serum bicarbonate <15-18 mmol/L

• Increased anion gap

• Ketonemia (beta-hydroxybutyrate [BOHB] >3 mmol/L) and ketonuria

Signs of ketoacidosis:

• Kussmaul breathing (deep, heavy, non-labored rapid breathing)

• Fruity breath odor (acetone)

• Prolonged Q-T interval

• Diminished neurocognitive function

• Coma


Management of diabetic ketacidosis
Hyperglycemia + Ketosis/Ketonuria
Check: Na, K, HCO_3, Cl, Mg, pH status
Start IV fluids: Normal saline (NS) at 1 L/h
IV fluids
Insulin
Potassium
Severe hypovolemia: NS at 1 L/h
Mild hypovolemia: Check corrected Na level
Low Na: Continue NS
High Na: Switch to 0.45% saline
When RBS < 250 mg/dl: Change to 5% glucose + 0.45% saline
IV bolus: 0.1 U/kg followed by IV infusion: 0.1 U/kg/h
If RBS does not fall by 50–70 mg/dl in 1st hour, double insulin dose
Glucose goal: 150–200 mg/dl
Reduce infusion to 0.02–0.05 U/kg/h Start SC insulin once DKA has resolved and patient is able to eat
Monitor every 2–4 hours until stable:
• Blood glucose
• Electrolytes
• BUN and Creatinine
• Venous pH
If < 3.3, do not start insulin until K^+ is corrected
If > 5.3, do not give KCl, but monitor every 2 hours
For serum K of 3.3 to 5.3, begin K correction at 10 mEq/h
©MARROW

Medicine INI-CET 2021

Question 1:

Identify the correct 'organism- drug to which it is intrinsically resistant' pair:

a) Aspergillus niger - Voriconazole

b) Aspergillus fumigatus - Micafungin

c) Candida albicans - Amphotericin B

d) Candida krusei - Fluconazole

Question 2:

A 45-year-old male patient presented with a history of bilateral lower limb weakness which progressed to his upper limbs in a year. On examination, he had weakness in both lower limbs and wasting in the left upper limb. Babinski sign was positive and the deep tendon reflexes were absent. He has no sensory loss or any autonomic dysfunction. What is the likely diagnosis?

a) MS

b) ALS

c) GBS

d) Tropical spastic paraparesis

Question 3:

In a PHC, which of the following initial screening tests are used to diagnose diabetes?

a) 1 and 2

b) 2 and 3

c) 2 and 4

d) 1, 2 and 3

Question 4:

A newly diagnosed HIV patient presented with fever, cough, and hemoptysis. His sputum sample showed acid-fast bacilli. What is the appropriate management for this patient?

a) Start ART if CD4<200

b) Start ART and then ATT after 2 weeks

c) Start ATT and then ART after 2 weeks

d) Start ATT and ART simultaneously

Question 5:

Identify the pathology from the given flow-volume loop:



a) Fixed distal airway obstruction

b) Fixed central airway obstruction

c) Variable intrathoracic obstruction

d) Variable extrathoracic obstruction

Question 6:

An elderly woman presented with symptoms of confusion, thirst, and abdominal pain. On examination, she had pallor and thoracic spine tenderness. X-ray spine showed osteolytic lesions. Her lab investigations showed the following findings. What is the most likely diagnosis?

a) Metastatic breast cancer

b) Multiple myeloma

c) Primary hyperparathyroidism

d) Milk alkali syndrome

Question 7:

A patient presented with sinusitis, ulcerative lesions on the nasopharynx, nodular lesions with cavitations of the lung, and renal failure. Which of the following is the most useful investigation in this patient?

a) Test for ANCA

b) Biopsy to show granuloma

c) AFB staining of sputum

d) BAL

Question 8:

Identify the urinary crystal shown below:



a) Calcium oxalate

b) Calcium phosphate

c) Uric acid

d) Cysteine

Question 9:

During the discharge of a COVID patient who was treated with steroids and remdesivir, which of the following will you inform him about?

a) 1, 3 and 4

b) 2, 3 and 4

c) 3, 4 and 5

d) 1, 2, 3, 4 and 5

Question 10:

A 50-year-old man suddenly developed right-sided weakness and aphasia within 2 hours. His BP recorded was 160/110mmHg and NCCT was clear. What is the next step in management?

a) CT angiography to look for large vessel occlusion

b) MRI to look for infarct

c) Tab labetalol 10 mg stat

d) Thrombolysis

Question 11:

In atrial fibrillation, choose the correct statements:

a) 1, 2, 3 and 4 correct

b) 1, 2, and 3 correct

c) 2 and 3 correct

d) 1 and 4 correct

Question 12:

Which of the following is not a major diagnostic criterion of neuromyelitis optica?

a) Area postrema syndrome

b) Acute Myelitis

c) Focal epilepsy

d) Optic neuritis

Question 13:

Acid-base imbalance is suspected in a patient. Which of the following would you use to determine it?

a) 1, 2

b) 1, 2, 3

c) 1, 2, 4

d) 1, 2, 3, 4

Question 14:

Which of the following does not show both hyperkinetic and hypokinetic movements?

a) Parkinson's disease

b) Wilson's disease

c) Neurodegeneration with Brain Iron Accumulation

d) Huntington's chorea

Question 15:

For the diagnosis of resistant hypertension, the patient should have uncontrolled hypertension despite taking three antihypertensive drugs. This should include at least one:

a) Calcium channel blocker

b) Beta blocker

c) Alpha blocker

d) Diuretic

Question 16:

Which of the following conditions is associated with hypokalemic metabolic alkalosis with hypertension?

a) Bartter syndrome

b) Liddle syndrome

c) Gitelman syndrome

d) Gordon syndrome

Question 17:

Which of the following is not true about syndrome of inappropriate antidiuretic hormone secretion (SIADH)?

a) Urinary sodium <20 mEq/L

b) Serum sodium <135 mEq/L

c) Urine osmolality >100 mOsm/kg

d) Patient can be clinically euvolemic to hypervolemic

Question 18:

A patient came with hypoxemia and a normal alveolar-arterial oxygen gradient. What can be the cause?

a) Right to left shunt

b) Ventilation/Perfusion mismatch

c) Hypoventilation

d) Alveolar membrane damage

Question 19:

Treatment of choice for an acute attack of cluster headache is :

a) Oral sumatriptan

b) Subcutaneous sumatriptan

c) 100% oxygen at 6 L/minute

d) 100% oxygen at 8 L/minute

Question 20:

All are true regarding heart failure except:

a) Regardless of ejection fraction and the type of heart failure, the 5 year mortality rate is around 50%

b) Non-cardiovascular death is more in HFrEF -30-40% as compared to HFpEF -15%

c) Angiotensin converting enzyme (ACE) inhibitors cause cough in 15% and angioedema in 1%

d) Atrial fibrillation is common in elderly and is the cause of stroke in quarter of the stroke cases

Question 21:

A 35-year-old female patient with class II pulmonary hypertension presents with a negative vasoreactivity test. Which of the following is used in the next step of management?

a) Iloprost

b) Ambrisentan

c) Nifedipine

d) Epoprostenol

Question 22:

A 30-year-old female patient came to the outpatient department with complaints of neck swelling. She gives a history of weight loss and palpitations. On examination, she has exophthalmos and a diffusely enlarged thyroid gland. What are the essential investigations that should be done for the patient?

a) 1, 2, 3, 4, and 5

b) 1, 3, 5 only

c) 3, 4, 5 only

d) 1, 3, 4 and 5 only

Question 23:

Which of the following drugs is preferred in the management of primary progressive multiple sclerosis ?

a) Natalizumab

b) Ocrelizumab

c) Fingolimod

d) Alemtuzumab

Question 24:

The blood investigation of a patient is given below. What is the probabale diagnosis?

a) Acute hepatitis B

b) Chronic hepatitis B

c) Remote infection with hepatitis B

d) Core window period

Question 25:

A 60-year-old male patient with critical mitral stenosis presented with atrial fibrillation. He has a history of multiple episodes of transient ischemic attacks. Which of the following are true regarding the prevention of stroke in this patient?

a) 1 only

b) 2 and 3

c) 2, 3 and 4

d) 1, 2, 3 and 4

Question 26:

True about electroencephalogram (EEG) is all except:

a) 1-5% of the population can have epileptiform discharges

b) Scalp EEG may be helpful in localizing frontal lobe epilepsy

c) Doing EEG is mandatory for diagnosis of seizures

d) Progressive multifocal leukoencephalopathy shows triphasic and slow waves on EEG

Question 27:

A 62-year-old male patient with heart failure is scheduled for a heart transplant. His renal function test is deranged and hemoglobin is 6 gm%. The physician ordered 2 units of whole blood. Four hours after transfusion, he developed severe respiratory distress. On examination, he is hypoxemic, has tachycardia and his mean arterial pressure is elevated. Which of the following are the best investigations for the above scenario?

a) 4 and 5

b) 3 and 5

c) 1 and 2

d) 2 only

Question 28:

A chronic smoker presented with shortness of breath, bilateral pitting pedal edema, and abdominal distension. On examination, jugular venous pulse was found to be elevated and liver was palpable 8 cm below the costal margin. An abnormality in which of the following structures is responsible for the patient's symptoms?



a) 1

b) 2

c) 3

d) 4

Answer Key

Question No.	Correct Option
1	d
2	b
3	c
4	c
5	b
6	b
7	a
8	a

9	c
10	a
11	c
12	c
13	a
14	d
15	d
16	b
17	a
18	c
19	b
20	b
21	b
22	d
23	b
24	c
25	b
26	c
27	c
28	b

Detailed Explanations

Solution to Question 1:

Among the given options, Candida krusei shows an intrinsic resistance to fluconazole. The primary or intrinsic resistance of C. krusei to azoles (fluconazole and itraconazole) is mediated through reduced susceptibility of the target enzyme (lanosterol 14α-demethylase) to inhibition by these drugs. However, improved susceptibility is seen with newer generation azoles such as voriconazole.

Life-threatening systemic infections caused by Candida and Aspergillus species are usually seen in severely ill and immunocompromised patients such as HIV, chemotherapy-induced neutropenia, and transplant recipients on immunosuppressants. C. albicans and C. dubliniensis are most susceptible to currently available azoles, C. glabrata is less susceptible, whereas C. krusei is intrinsically resistant to fluconazole. Resistance occurs due to overexpression or point mutation in the gene encoding the target enzyme lanosterol 14α-demethylase. Voriconazole is the preferred therapy for multi-drug resistant infection with C. krusei.

C. albicans showing resistance to Amphotericin B has been increasingly reported over the past several years. But this is a secondary resistance due to alteration or a decrease in the amount of

ergosterol in the cell membrane.

Aspergillus fumigatus, the most common species to cause Aspergillus infection worldwide, also shows an intrinsic resistance to fluconazole. It is susceptible to echinocandins like micafungin, which acts by inhibiting beta-(1,3)-D-glucan synthase, an enzyme necessary for the synthesis of the cell wall.

A. niger is a common cause of otomycosis and is susceptible to azoles like voriconazole, amphotericin B, and echinocandins.

Solution to Question 2:

The given clinical scenario with both UMN (Babinski positive) and LMN (absent deep tendon reflexes, wasting) findings in the absence of sensory loss and autonomic dysfunction points towards a diagnosis of amyotrophic lateral sclerosis (ALS).

ALS is a progressive motor neuron disease with mutations in the genes encoding:

• protein C9orf72 (open reading frame 72 on chromosome 9)

• cytosolic enzyme SOD1 (superoxide dismutase),

• RNA binding proteins TDP43 (encoded by the TAR DNA binding protein gene), and fused in sarcoma/translocated in liposarcoma (FUS/TLS).

The pathologic hallmark of ALS is the death of lower motor neurons (anterior horn cells in the spinal cord and cranial nerves innervating bulbar muscles) and upper motor neurons (corticospinal neurons originating in layer five of the motor cortex). At onset, ALS may involve only upper or lower motor neurons, but ultimately causes progressive loss of both categories of motor neurons. Clinical features of ALS include:

• LMN findings: fasciculations, muscle cramps, asymmetric distal weakness, atrophy, and bulbar palsy.

• UMN findings: spasticity, increased reflexes, flexor spasms, and pseudobulbar palsy.

• Pseudobulbar affect: degeneration of corticobulbar projections manifested as inappropriate laughing, crying, or yawning which is often mood incongruent or may be triggered by stimuli that would not have elicited such a response.

• Even in the late stages, sensory, bowel and bladder, ocular motility, and cognitive functions are preserved.

Treatment:

• Riluzole (100mg/day) reduces excitotoxicity by diminishing glutamate release and improves survival.

• Edaravone (IV infusion) works in the central nervous system as a potent scavenger of oxygen radicals that has the capability of slowing down ALS progression and improving survival.

• Tofersen is a newer drug proven to be effective for ALS associated with SOD1 mutation. It is an antisense oligonucleotide that targets SOD1 mRNA to reduce the synthesis of SOD1 protein.

Multiple sclerosis (MS) is an autoimmune demyelinating and inflammatory disease of the CNS characterized by gliosis and neuronal loss. It is caused by an autoimmune response directed against components of the myelin sheath. Genetic predisposition with HLA-DRB1, cigarette smoking, vitamin D deficiency, and exposure to Epstein Barr virus (EBV) are known risk factors. Clinical features of MS include:

• Visual impairment due to the involvement of the optic nerve (optic neuritis).

• Involvement of the brainstem causes cranial nerve signs such as ataxia, nystagmus, and internuclear ophthalmoplegia (INO).

• Weakness of limb (usually UMN type), facial weakness, spasticity, hyperreflexia, Babinski signs, sensory impairment of the trunk and limbs, and loss of bladder control.

Guillain-Barre syndrome (GBS) is an acute inflammatory demyelinating neuropathy causing rapidly evolving ascending paralysis. It is associated with diminished reflexes (areflexia) and autonomic involvement with or without sensory impairment.

Tropical spastic paraparesis also known as HTLV-1 associated myelopathy is a slowly progressive spastic syndrome with variable sensory and bladder disturbance but no LMN involvement. Diagnosis is by demonstration of HTLV-1 specific antibody in serum by ELISA and confirmation is done using radioimmunoprecipitation or Western blot analysis.

Solution to Question 3:

Fasting blood sugar >126 mg/dL and 2-hour postprandial blood sugar >200 mg/dl can be used as screening tests for diabetes in a primary health center (PHC).

HbA1c estimation is usually unavailable for screening at many PHCs due to low resources. In India, facilities for HbA1c estimation are usually available only at tertiary centers.

Random blood sugar >200 mg/dL can be used for diagnosis only if the patient presents with symptoms of diabetes, hence it is not a useful screening test.

Solution to Question 4:

In an HIV patient with TB, start antitubercular therapy (ATT) first and then start antiretroviral therapy (ART) as soon as ATT is tolerated (between 2 weeks and 2 months) by the patient. This is done to avoid immune reconstitution inflammatory syndrome (IRIS).

Among persons living with HIV with TB meningitis, ART should be delayed for at least 4 weeks (and initiated within 8 weeks) after treatment for TB meningitis is initiated.

Patients with HIV-related TB may experience a temporary paradoxical exacerbation of symptoms and radiographic manifestations of TB after starting ART (IRIS). This occurs as a result of immune reconstitution due to the simultaneous administration of ART and ATT.

It is characterized by high fever, lymphadenopathy, expanding intra-thoracic lesions, and worsening of chest X-ray findings. Diagnosis should be made only after ruling out treatment failure and other infections. For severe cases, prednisone 1-2 mg/kg is used for 1-2 weeks and then

gradually tapered.

Note: CBNAAT is the preferred diagnostic technique for testing TB in PLHIV (people living with HIV). Smear microscopy has poor sensitivity in detecting TB in PLHIV due to fewer organisms in sputum, and hence is not preferred.

Solution to Question 5:

The given flow-volume curve shows flattening of both the inspiratory and expiratory limbs, which is seen in fixed obstruction of the central or upper airway.

The curve denotes airflow limitation during both inspiration and expiration.

Examples of fixed central airway obstruction include tracheal stenosis (caused by intubation) and circular tracheal tumors.

Other options:

Option A: The image given below shows a concave upward, pattern in the expiratory limb (scooped-out or coved) pattern seen in lower airway obstruction. Examples include asthma, chronic obstructive pulmonary disease, bronchiectasis, and bronchiolitis.



Option C: The image below shows airflow limitation and flattening in the expiratory limb of the loop. It is seen in variable, nonfixed intrathoracic obstruction.

Examples: intrathoracic airway, bronchogenic cysts, or tracheal lesions.



Option D: The image below shows airflow limitation and flattening in the inspiratory limb of the loop. It is seen in nonfixed variable extrathoracic obstruction.

Examples: vocal cord paralysis, extrathoracic tracheomalacia, polychondritis, etc.





Solution to Question 6:

The given clinical scenario describes an elderly woman with confusion, thirst and abdominal pain along with elevated serum calcium levels (hypercalcemia), elevated creatinine suggestive of renal failure, anemia and osteolytic bone lesions (CRAB features); and hypoalbuminemia with elevated total proteins. The most likely diagnosis is multiple myeloma (MM).

MM is a malignant plasma cell dyscrasia with the proliferation of plasma cells derived from a single clone. It is associated with increased osteoclast activity with suppression of osteoblast activity, causing punched-out lytic lesions in the bone and hypercalcemia.

Signs and symptoms include bone pain (most common), pathological fractures, recurrent bacterial infections, normocytic normochromic anemia, elevated ESR, and normal ALP.

Electrophoresis shows monoclonal elevation of one immunoglobulin (M spike)

Urine analysis shows free kappa & lambda light chains (Bence Jones proteins)

Peripheral smear shows rouleaux formation.

The diagnosis of MM is made by using the Revised International Myeloma Working Group criteria.

Clonal bone marrow plasma cells ≥10% (or) Biopsy-proven bony or extramedullary plasmacytoma and any one or more of the following myeloma-defining events is required to meet the criteria:

Evidence of end-organ damage attributable to the underlying plasma cell proliferative disorder, specifically: (CRAB features)

• Hypercalcemia: Serum calcium level >1 mg/dL higher than the upper limit of normal or >11 mg/dL

• Renal insufficiency: Creatinine clearance <40 mL/min or serum creatinine >2 mg/dL

• Anemia: Hemoglobin level of >20 g/L (2gm/dl) below the lowest limit of normal, or Hb <10 g/dL

• Bone lesions: One or more osteolytic lesions on skeletal radiography, computed tomography (CT), or positron emission tomography (PET-CT).

Any one or more of the following biomarkers of malignancy:

• Clonal bone marrow plasma cells ≥60%

• Serum involved/uninvolved free light-chain ratio of 100 or greater

• >1 focal lesion on magnetic resonance imaging (MRI) that is at least 5 mm or greater in size.

Treatment includes a combination of lenalidomide/thalidomide (immunomodulatory agents), bortezomib (proteasome inhibitor), and dexamethasone.

International Staging System (ISS) is used for assessing prognosis in multiple myeloma patients. Serum β2-microglobulin is the most important prognostic marker.

Other options:

Option A: The bony metastasis in breast carcinoma is usually osteolytic. However, CRAB features, hypoalbuminemia, and elevated total protein suggest multiple myeloma.

Option C: Primary hyperparathyroidism is characterized by hypercalcemia and low phosphorus but is not associated with anemia and hypoalbuminemia.

Option D: Milk-alkali syndrome occurs after ingestion of large amounts of calcium and vitamin D supplementation. It results in hypercalcemia, alkalosis, nephrocalcinosis, and nephrolithiasis.

Solution to Question 7:

The given patient has the involvement of upper respiratory tract (chronic sinusitis, ulcerations on nasopharynx), lower respiratory tract (nodular lesions with lung cavitations), and kidneys (renal failure) which are suggestive of granulomatosis with polyangiitis (Wegener's granulomatosis) and testing for cytoplasmic ANCA (c-ANCA) would be the most helpful investigation in this condition.

Granulomatosis with polyangiitis is a granulomatous and necrotizing vasculitis of the small arteries and veins that mainly affects:

• Upper respiratory tract: chronic sinusitis, serous otitis media due to eustachian tube blockage, nasal septal perforation, and mucosal ulceration of the nasopharynx.

• Lower respiratory tract: presenting with cough, hemoptysis, dyspnea, and persistent pneumonitis. Lung involvement typically appears as multiple, bilateral nodular cavitary infiltrates.

• Kidneys: focal or segmental glomerulonephritis presenting with proteinuria, hematuria, and hypertension. If left untreated it can lead to rapidly progressive glomerulonephritis.

Apart from these organs, it can also affect the eyes and skin. Skin lesions present as palpable purpura or subcutaneous nodules.

Lab investigations will show anemia, leucocytosis, elevated ESR, and high titers of cytoplasmic ANCA (c-ANCA) targetted against anti-proteinase-3 (anti-PR-3). The specificity of a positive PR-3 ANCA for Wegener's is very high, especially in the presence of active glomerulonephritis. Following treatment, PR-3 ANCA titers fall and subsequent rise in titers is predictive of

relapse. Chest radiograph will show multiple, bilateral cavitating nodules and pulmonary infiltrates.

The diagnosis is confirmed by the presence of necrotizing granulomatous vasculitis on biopsy. This finding is commonly seen in the tissues obtained from the lungs, upper airways, skin, and rarely in the kidneys.

The radiograph below shows cavitating nodules in a patient with granulomatosis with polyangiitis (Wegener's).



Solution to Question 8:

The given urinary crystal is the bipyramidal or envelope-shaped crystals of calcium oxalate dihydrate. Risk factors for calcium oxalate stones include higher urine calcium, urine oxalate, and lower urine citrate.

Oxalate-rich foods like spinach, rhubarb, almonds, and potatoes should be avoided. Avoiding high-dose vitamin C supplements reduces endogenous oxalate production.

Chlorthalidone can lower urine calcium excretion and can reduce the recurrence of calcium oxalate stones. Consumption of fruits and vegetables or supplemental alkali is advised to increase urine citrate, which is a natural inhibitor of calcium oxalate and calcium phosphate stones.

Solution to Question 9:

The given clinical scenario describes a patient treated with steroids and remedesivir for COVID-19 infection being discharged. This patient should be advised to monitor blood glucose levels as steroids can worsen glycemic status. He should be advised to watch for headache, nasal discharge, and sinusitis-like symptoms as there is a higher incidence of mucormycosis among COVID patients.

Other options:

Option 1: Repeat RT-PCR after 7 days of discharge is not needed as per the updated discharge criteria.

Option 2: Anosmia may persist for weeks to months in those who recovered from COVID-19. Its persistence is not a dangerous sign.

Mucormycosis is most commonly caused by the fungus Rhizopus oryzae that is acquired through the inhalation of fungal spores from the air. It may occur during COVID infection or after a few weeks (usually around the 3rd week) of recovery from it.

Risk factors in COVID-19 patients include uncontrolled hyperglycemia, misuse and irrational use of steroids, leukopenia, and unhygienic humidifiers.

Symptoms of rhino-orbital-cerebral mucormycosis include facial pain, facial numbness, brownish or blood-tinged nasal discharge, conjunctival suffusion and swelling, black eschar over the face, and hard palate.

The presence of bilateral proptosis, chemosis, vision loss, and ophthalmoplegia suggests cavernous sinus thrombosis which is seen in advanced disease.

Diagnosis is done by biopsy with the histopathological examination which shows the presence of ribbon-like aseptate hyphae (5-15 μ thick) branching at right angles.

CT chest in pulmonary mucormycosis may show lobar consolidation, isolated masses, cavities, nodules, pleural effusion, or wedge-shaped infarct. MRI can be utilized to assess the degree of orbital and brain involvement.

Management includes strict glycemic control. Antifungal therapy started early on high clinical suspicion even prior to microbiological confirmation can reduce mortality.

Liposomal Amphotericin B (5-10 mg/kg) is the drug of choice and has to be continued for 3-6 weeks. It should be diluted in 5% dextrose and is incompatible with normal saline and Ringer lactate. Step-down therapy is to be done with oral posaconazole or isavuconazole. Renal function must be monitored during the entire treatment period.

Blood vessel thrombosis and tissue necrosis in mucormycosis can lead to limited penetration of antifungal drugs at the infection site. To address this, debridement of necrotic tissues is crucial in eradicating the disease.

Revised discharge policy for COVID-19:

Mild cases discharged after at least 7 days have passed from testing positive and with no fever for 3 consecutive days. No testing is needed prior to discharge.

For moderate cases, in patients whose signs and symptoms resolve and maintain saturation above 93% for 3 consecutive days, patient will be discharged as per the advice of the treating medical officer. No testing is needed prior to discharge.

In severe cases including immunocompromised, discharge criteria will be based on clinical recovery at the discretion of the treating medical officer.

Solution to Question 10:

The next step in management would be CT angiography. NCCT and CT angiography must be done concurrently during stroke evaluation. The given features of hemiparesis with aphasia and a clear NCCT are suggestive of an acute ischemic stroke with a proximal large vessel occlusion.

Though thrombolysis is the treatment of choice, occlusions in large vessels often fail to open up with thrombolysis alone. Hence, mechanical thrombectomy is also indicated and should be done ideally within 6 hours and not beyond 24 hours of symptom onset.

CT angiography determines if there is an emergent large vessel occlusion (ELVO), and can be used to triage patients for management by intra-arterial mechanical thrombectomy using a retriever device.

Other options:

Option D: Thrombolysis with alteplase is the first-line therapy for acute ischemic stroke and should be initiated within 4.5 hours of symptom onset. However, mechanical thrombectomy is indicated as well for acute ischemic stroke due to a large artery occlusion in the anterior circulation. Thus, CT angiography to look for large vessel occlusion would be the next best step in the management.

Option B: MRI with diffusion-weighted imaging (DWI) is superior to NCCT for detecting an acute infarct. However, MRI is not preferred to NCCT because of longer scan completion time, limited availability in emergency settings, and higher cost.

Option C: Labetalol, nicardipine, and clevidipine are the first-line anti-hypertensive agents used in the acute phase of stroke. The blood pressure goals in ischemic stroke are:

• If the patient is to receive thrombolytic therapy, maintain BP ≤185/110 mmHg.

• After thrombolytic therapy, maintain BP ≤180/105 mmHg for at least the next 24 hours.

• If not treated with thrombolytic therapy, BP should not be treated acutely unless it is more than 220/120 mmHg. BP should be lowered cautiously by only 15 percent in the first 24 hours in order to preserve perfusion to the ischemic penumbra.

Solution to Question 11:

Statements 2 and 3 are correct.

Novel Oral Anticoagulants (NOACs) also known as direct oral anticoagulants are a new class of anticoagulants that include oral thrombin inhibitor (dabigatran) and oral factor Xa inhibitors like rivaroxaban, apixaban, and edoxaban. They are easier to use, achieve reliable anticoagulation, and do not require dosage adjustment based on blood tests. They have a lower risk of intracranial hemorrhage compared to warfarin and are used as alternatives to warfarin for patients who have atrial fibrillation.

Anticoagulation with warfarin or NOACs should be given in patients with ischemic stroke due to atrial fibrillation. In patients with small or moderate-sized infarcts and no intracranial bleeding, warfarin can be started soon (after 24 hours) after admission. The risks of transformation to hemorrhagic stroke are minimal. NOACs should be started after 48 hours as they have a more rapid anticoagulant effect.

Statement 3: NOACs are contraindicated in patients with prosthetic mechanical heart valves due to a lack of sufficient evidence.

Statement 1: NOACs are not used in the treatment of moderate to severe mitral stenosis (MS) due to limited clinical evidence supporting their efficacy and safety in MS. Warfarin is the recommended anticoagulant for patients with MS and mechanical prosthetic valves.

Statement 4: Beta-blockers and calcium channel blockers are the first line drugs used to acheive rate control in atrial fibrillation.

Solution to Question 12:

Focal epilepsy is not considered a major diagnostic criterion of neuromyelitis optica (NMO).

Neuromyelitis optica (Devic's disease) is an aggressive inflammatory demyelinating disorder characterized by recurrent attacks of optic neuritis and myelitis.

It is associated with a highly specific autoantibody directed against aquaporin-4(AQP-4). AQP-4 is localized to the foot processes of astrocytes in close proximity to endothelial surfaces.

The core clinical characteristics include optic neuritis, acute myelitis, area postrema syndrome (unexplained hiccups/ nausea/ vomiting), acute brainstem syndrome, symptomatic narcolepsy or acute diencephalic clinical syndrome with Neuromyelitis Optica Spectrum Disorder (NMOSD)-typical diencephalic MRI lesions, and symptomatic cerebral syndrome with NMSOD-typical brain lesions.

Attacks of optic neuritis are bilateral and severe in NMO, whereas in multiple sclerosis (MS) optic neuritis is usually unilateral. Progressive symptoms are not typical of NMO, unlike in MS. CSF findings in NMO show higher pleocytosis, with neutrophils and eosinophils in acute cases, and rare occurrence of oligoclonal bands, distinguishing it from MS.

Acute attacks of NMO are treated with high-dose glucocorticoids. Plasma exchange can be used in acute attacks that do not respond to glucocorticoids.

Drugs used for prophylaxis against relapses include mycophenolate mofetil, rituximab, glucocorticoids, and azathioprine. Three monoclonal antibodies approved for attack prevention include eculizumab (terminal complement inhibitor), inebilizumab (B-cell depleter), and satralizumab (IL-6 receptor blocker).

NMO typically presents as a recurrent disease, with a monophasic course observed in less than 10% of patients. Untreated NMO can lead to disability, including respiratory failure, blindness, and limb paralysis.

Solution to Question 13:

Arterial and Venous pH can be used to determine acid-base imbalance.

Arterial blood gas (ABG) analysis is the gold standard method for the assessment of oxygen and acid-base status.

Venous blood gas (VBG) analysis is increasingly being used as a substitute for ABG as the difference between the pH obtained from VBG and ABG is clinically insignificant (arterial pH typically 0.03 higher than venous pH), thus making venous pH values acceptable.

Other options:

Option 3: The pO2 values also show a significant difference, with arterial pO2 (normal 95mmHg) values typically 55mmHg greater than the venous pO2 (normal 40mmHg), making venous pO2 values unacceptable.

Option 4: Venous pCO2 (normal range 41 to 51mmHg) values differ significantly from arterial pCO2 (normal range 35 to 45mmHg) by a range of 5 to 15mmHg which makes venous pCO2 values unacceptable.

Solution to Question 14:

Huntington's disease (formerly known as Huntington's chorea) is a movement disorder that shows predominantly only hyperkinetic movements like chorea.

Hyperkinetic movement disorders include tremor, dystonia, athetosis, chorea, myoclonus, and tics. Hypokinetic movements include rigidity, stiffness, and bradykinesia.

Huntington's disease is a degenerative disease of the caudate nucleus and putamen (striate neurons) caused by a trinucleotide repeat (CAG) in the Huntingtin gene on chromosome 4. It is inherited in an autosomal dominant pattern and is characterized by chorea, dysarthria, gait disturbance, behavioral disturbance and cognitive impairment with dementia. The clinical manifestations increase in severity with each successive generation (anticipation). Treatment includes haloperidol, and tetrabenazine.

With advancing disease, a reduction in chorea may be seen along with the emergence of dystonia, rigidity, myoclonus, and bradykinesia. It can also have an akinetic, rigid, or Parkinsonian presentation in younger patients (~10% of cases) known as the Westphal variant. However, the typical presentation of Huntington's disease is characterized by hyperkinetic movements like chorea.

Other options:

Option A: Parkinson's disease is characterized by both hyperkinetic (resting tremor) and hypokinetic (rigidity, stiffness, bradykinesia) movements.

Option B: The neurological manifestations in Wilson's disease occur due to damage of basal ganglia resulting in dystonia and tremor (hyperkinetic), parkinsonism with bradykinesia (hypokinetic), incoordination, dysarthria, and dysphagia. In advanced disease, the tremor progresses to a characteristic wing-beating flapping tremor.

Option C: Pantothenate kinase-associated neurodegeneration (PKAN) formerly known as Hallervorden-Spatz disease caused by PANK2 gene mutation is the most common form of Neurodegeneration with Brain Iron Accumulation (NBIA) and is manifested as a combination of dystonia (hyperkinetic), parkinsonism (hypokinetic), and spasticity. T2W-MRI shows the characteristic eye of the tiger sign due to iron accumulation in globus pallidus.

Solution to Question 15:

For the diagnosis of resistant hypertension, the patient should have uncontrolled hypertension despite taking three antihypertensive drugs, including at least one diuretic.

While a calcium channel blocker, beta blocker, or alpha-blocker can be part of the antihypertensive regimen, the inclusion of a diuretic is necessary for the diagnosis of resistant hypertension. The diuretic helps in addressing fluid retention which is often a significant contributor to hypertension.

Resistant hypertension is more common in patients aged >60 years than in younger patients. It may be related to nonadherence to therapy, obesity, and excessive alcohol intake.

In the absence of a specific identifiable cause, mineralocorticoid receptor antagonists like spironolactone have been found to be most effective add-on drugs for the treatment of resistant hypertension.

Solution to Question 16:

Liddle's syndrome is associated with hypokalemic metabolic alkalosis with hypertension.

It is an autosomal dominant condition due to a gain of function mutation in the ENaC gene, which codes for the epithelial sodium channels present in the kidney.

Hypertension is the result of increased sodium retention through ENaC channel and volume expansion. Volume expansion suppresses renin and aldosterone secretion, leading to hypokalemic metabolic alkalosis.

Treatment is with diuretics that directly block the ENaC channels, like amiloride and triamterene. Spironolactone is not effective as it is an aldosterone-receptor antagonist.

Other options:

Option A: Bartter syndrome is an autosomal recessive disorder occurring due to defects in various genes involved in the regulated Na+, K+, and Cl– transport by the thick ascending loop of Henle. There is extracellular fluid volume depletion with low or normal blood pressure. It is characterized by metabolic alkalosis, hypokalemia, hypochloremia, and hypercalciuria.

Option C: Gitelman syndrome is an autosomal recessive disorder due to a loss of function mutation in the thiazide-sensitive Na+-Cl– cotransporter of the distal convoluted tubule. It is characterized by hypokalemic metabolic alkalosis with hypomagnesemia and hypocalciuria with low blood pressure.

Option D- Gordon syndrome is a state of pseudohypoaldosteronism. It is characterized by hypertension, hyperkalemia, metabolic acidosis, and low-normal plasma renin activity and aldosterone concentrations.

Solution to Question 17:

In SIADH, the kidneys retain water, leading to volume expansion and natriuresis. Therefore, urinary sodium levels are typically higher than 20 mEq/L.

Other options:

Option B: SIADH is a condition where there is an inappropriate release of antidiuretic hormone (ADH), leading to impaired water excretion and dilutional hyponatremia.

Option C: ADH promotes the reabsorption of water and not the solutes resulting in concentrated urine. This is depicted as high urine osmolality.

Option D: Patients can be hypervolemic due to water retention. However, persistently elevated ADH levels can induce vasopressin escape mechanism, resulting in the typical euvolemic status.

Patients usually present with hyponatremia with a euvolemic status.

Acute hyponatremia may present with mild headache, confusion, anorexia, nausea, vomiting, coma, and convulsions.

Chronic hyponatremia allows cerebral adaptation, and the patients are asymptomatic despite the hyponatremia.

Laboratory findings in SIADH include:

• Hyponatremia: serum Na+<135 mEq/L

• Urinary Na+ >25 mEq/L

• Urine osmolality >100 mOsm/kg

• Blood urea nitrogen (BUN) levels are low, usually <10 mg/dL

• Serum urate is low

• Serum K+ and serum bicarbonate are normal

Mild and symptomatic hyponatremia can be managed by restricting fluid intake.

Severe, symptomatic hyponatremia can be corrected using hypertonic saline or vasopressin receptor-2 antagonists (tolvaptan and conivaptan).

Other drugs which are useful in the management are demeclocycline and fludrocortisone.

Solution to Question 18:

All of the given options can cause hypoxemia out of which the alveolar-arterial oxygen gradient (A-a gradient) remains normal in hypoventilation.

The A-a gradient or the alveolar-arterial oxygen gradient refers to the difference between the oxygen concentration in the alveoli (PAO2) and oxygen concentration in the arterial system (PaO2).

A-a gradient= PAO2 - PaO2

In hypoventilation, patients lack oxygen tension in the respiratory system as well as their arterial system. This results in a decrease in both PAO2 and PaO2 and A-a gradient remains normal. Hypoventilation can be seen in CNS depression, neuromuscular diseases such as myasthenia gravis, kyphoscoliosis, vertebral fractures, etc.

Other options:

Option A- In a right-to-left shunt, a portion of the pulmonary blood flow gets shunted away from the alveoli. This results in ventilation without perfusion which causes a higher V/Q ratio. Here, diffusion between capillaries and alveoli is unaffected, but the arterial PaO_2 is decreased due to the lack of oxygenation of the shunted blood. This results in hypoxemia and an increased A-a gradient. In the cases of the shunt, oxygenation does not improve hypoxemia.

Option B- Ventilation/Perfusion (V/Q) mismatch is the most common cause of hypoxemia.

Low V/Q mismatch occurs in chronic obstructive pulmonary disorder (COPD), asthma, interstitial lung disease, etc. In these cases, the A-a oxygen gradient is increased due to a defect in diffusion resulting in hypoxemia.

A high V/Q ratio is seen in pulmonary embolism (PE) where there is wasted ventilation. The alveolar oxygen is normal but there is decreased perfusion resulting in increased A-a gradient.

Inspiratory hypoxia, seen at high altitudes, can cause a V/Q mismatch. The decreased atmospheric pressure causes a decreased PAO_2. There is normal diffusion which suggests a normal A-a gradient while the arterial PaO_2 decreases, leading to hypoxemia.

Hypoventilation is the better answer as the cause of the V/Q mismatch is not specified.

Option D: Alveolar membrane damage affects diffusion between the alveoli and capillaries. The oxygen levels in the arterial blood decrease even though alveoli are being ventilated. This leads to an increased A-a gradient.

Solution to Question 19:

Among the given options, subcutaneous injection of sumatriptan (6 mg) is used in the treatment of an acute attack of cluster headache.

The treatment of choice for an acute attack of cluster headache is 100 % oxygen at 10-12 L /min. High flow and high oxygen content are important. 100 % oxygen at 6 or 8 L/min (Options C and D) is insufficient.

Oral sumatriptan (Option A) is not effective for the treatment of acute attack of cluster headaches.

Sumatriptan (20 mg) and zolmitriptan (5 mg) nasal spray and noninvasive vagus nerve stimulation (nVNS) can also be used for the treatment of acute episodes of cluster headaches.

Cluster headache is a rare primary headache characterized by daily attacks of short-duration unilateral headaches lasting 8-10 weeks a year. It presents as severe, brief (15-180 minutes) unilateral retro-orbital pain with autonomic features, such as unilateral lacrimation, nasal congestion, and conjunctival injection. Patients are usually perfectly well in between episodes.

Verapamil is the drug of choice for prophylaxis. Other drugs used are prednisone, topiramate, lithium, gabapentin, melatonin, and greater occipital nerve injection with lidocaine and glucocorticoids.

Solution to Question 20:

Non-cardiovascular deaths are more in heart failure with preserved ejection fraction (HFpEF) as compared to heart failure with reduced ejection fraction (HFrEF).

Non-cardiac causes accounted for 62% of deaths in HFpEF, 54% in HFmrEF (heart failure with mildly reduced ejection fraction), and 35% in HFrEF. Cancer was twice as frequent as a cause of death in HFpEF and HFmrEF versus HFrEF.

Heart failure is divided according to the ejection fraction:

• Heart failure with preserved ejection fraction (HFpEF): Left ventricular ejection fraction is >40-50%

• Heart failure with reduced ejection fraction (HFrPF): Left ventricular ejection fraction is <40%

Patients with a left ventricular ejection fraction between 40% and 50% are considered to have borderline or mid-range EF.

Other options:

Option A: 'Regardless of the EF and the type of heart failure, the 5-year mortality rate is around 50%' is a true statement. The development of symptomatic heart failure has a poor prognosis. 30-40% of patients die within 1 year of diagnosis. More than 50% (60-70%) die within 5 years.

Option C: Angiotensin-converting enzyme (ACE) inhibitors cause cough in 15% and rashes and angioedema in 1%. ACE inhibitors inhibit the angiotensin-converting enzyme and inhibit the conversion of angiotensin I to angiotensin II. ACE also splits bradykinin, substance P, enkephalins, etc. The bradykinin and substance P which are not broken down lead to cough, urticaria, and rarely angioedema.

Option D: The prevalence of atrial fibrillation increases with age, and >95% of AF patients are >60 years of age. Atrial fibrillation increases the risk of stroke by five-fold and is estimated to be the cause of 25% of strokes.

Solution to Question 21:

In a patient with class II pulmonary hypertension (low risk) with a negative vasoreactivity test, the next best step of management is to start on ambrisentan (endothelin receptor antagonist)

Pulmonary artery hypertension (PAH) is defined as mean pulmonary arterial pressure greater than 20 mmHg with a pulmonary vascular resistance of ≥3 wood units.

Exertional dyspnea, chest pain, fatigue, and lightheadedness are common early symptoms. With the progression of the disease, right ventricular dysfunction occurs which causes syncope, abdominal distention, ascites, and peripheral edema.

Vasoreactivity testing involves the administration of inhaled nitric oxide, IV epoprostenol, IV adenosine, or inhaled iloprost. The test is considered positive if the mean pulmonary arterial pressure (mPAH) decreases by ≥ 10 mmHg leading to mPAH ≤ 40 mmHg, without a decrease in cardiac output (CO).

In case of a positive vasoreactivity test, calcium channel blockers are indicated. Long-acting nifedipine, diltiazem or amlodipine can be used.

If the vasoreactivity test is negative,

• For patients with lower risk (class II, III) based on clinical assessment, oral therapy with endothelin receptor antagonists like ambrisentan, sitaxsentan, bosentan, or phosphodiesterase 5 inhibitors (PDE-5 I's) like sildenafil, tadalafil is the first line of therapy.

• For patients with higher risk (those with advanced heart failure or syncope, class IV) based on clinical assessment, continuous treatment with iv prostacyclins (epoprostenol or treprostinil) would be the first line of therapy.

Lung transplantation and/or atrial septostomy are reserved for patients who progress despite optimal medical treatment.

The treatment algorithm for pulmonary hypertension is given below:



Revised WHO classification of pulmonary hypertension:

• Group I: Pulmonary arterial hypertension (PAH)

• Group II: Pulmonary hypertension with left heart disease

• Group III: Pulmonary hypertension associated with lung diseases and/or hypoxemia

• Group IV: Pulmonary hypertension due to chronic thrombotic and/or embolic disease (CTEPH)

• Group V: Pulmonary hypertension due to miscellaneous causes

The table below provides guidance on the prognosis of PAH.

Determinants	Low risk	High risk
Clinical evidence of RV failure	No	Yes
Progression of symptoms	Gradual	Rapid
Echocardiography	Minimal RV dysfunction	Pericardial effusion, significant RV enlargement/dysfunction, right atrial enlargement
WHO functional class	II, III	IV
6min walk distance	Longer (>400 m)	Shorter (<300 m)

Determinants	Low risk	High risk
BNP	Minimally elevated	Significantly elevated

Solution to Question 22:

The given clinical scenario of a female patient presenting with weight loss, palpitations, exophthalmos, and diffusely enlarged thyroid gland is suggestive of Graves' disease.

The essential investigations in this patient would be thyroid function tests, ultrasound of the neck, thyroid scan, and anti-thyroid antibody. FNAC is not an essential investigation in this condition. FNAC should be done on nodules that are not classified as fully benign on ultrasound.

The hyperthyroidism of Graves' disease is caused by thyroid-stimulating immunoglobulin (TSI) synthesized by lymphocytes in the thyroid gland as well as in bone marrow and lymph nodes. TSI antibodies cause stimulation of the thyroid gland causing features of hyperthyroidism.

Symptoms include hyperactivity, heat intolerance, sweating, palpitations, weakness, weight loss, increased appetite, and diarrhea.

Signs include tachycardia, goiter, tremor, muscle weakness, and atrial fibrillation in the elderly.

Thyroid ophthalmopathy seen in one-third of patients with Grave's disease.

NO SPECS scoring system as describe below can be used to evaluate ophthalmopathy.

• No signs or symptoms

• Only signs (lid retraction, or lag)

• Soft tissue involvement (periorbital edema)

• Proptosis

• Extraocular muscle involvement

• Corneal involvement

• Sight loss

Thyroid dermopathy usually occurs in the presence of moderate or severe ophthalmopathy. It primarily manifests as noninflamed, indurated plaques with a deep pink or purple color and an "orange skin" appearance, commonly found on the anterior and lateral aspects of the lower leg (known as pretibial myxedema).

Investigations done in a patient with Graves' disease include:

• Thyroid function tests- TSH levels are decreased with or without elevated T3 and T4 levels.

• Anti-thyroid antibodies- Elevated levels of TSH receptor antibodies are diagnostic of Graves' disease. Other antibodies like anti-Tg and anti-TPO are elevated in most of the patients.

• Ultrasound of the neck is performed to evaluate vascularity, echogenicity, nodules, and clinical landmarks for surgical planning.

• A radioactive iodine uptake (RAIU) scan (thyroid scan) should be performed and a diffuse uptake confirms the diagnosis of Graves' disease. It is also useful to differentiate TSI-negative Graves

disease from toxic nodular disease based on diffuse or nodular uptake patterns.

• CT or MRI scans of the orbits are useful to evaluate Graves' ophthalmopathy.

Hyperthyroidism can be treated by antithyroid drugs, radioiodine treatment (I131), or thyroidectomy.

Solution to Question 23:

Ocrelizumab is preferred in the management of primary progressive multiple sclerosis (PPMS). Natalizumab, fingolimod, and alemtuzumab are used in the management of relapsing multiple sclerosis and not PPMS.

Ocrelizumab is a monoclonal antibody against CD20 molecules present on B cells. It reduces the progression of the disease and improves other clinical parameters.

Multiple sclerosis (MS) is an autoimmune disease characterized by chronic inflammation, demyelination, gliosis (scarring), and neuronal loss.

Based on clinical course, MS is classified as:

• Relapsing or bout onset MS (RMS)

• Secondary progressive MS (SPMS)

• Primary progressive MS (PPMS)

The drugs used in the management of RMS are listed in the table below:

Siponimod, a selective S1P1 S1P5 modulator, reduces the risk of progression of SPMS. Ocrelizumab, cladribine and high dose interferon beta can also be used.

Ocrelizumab is the first drug to alter the progression of disability in PPMS.

Highly effective	Moderately effective	Modestly effective
Anti CD20 antibodies: Ocrelizumab, Ofatumumab, RituximabAnti integrin antibody: NatalizumabAnti CD 52 antibody: AlemtuzumabMitoxantrone	Sphingosine-1-phosphate inhibitor: FingolimodDimethyl fumarateCladribine	Glatiramer acetateInterferon betaTeriflunomide

Solution to Question 24:

The most probable diagnosis is remote infection with hepatitis B virus (HBV).

A non-reactive HbsAg rules out active HBV infection (Options A and B).

Non-reactive IgM anti-HbcAg suggests that the patient is not in the window period (Option C).

Reactive IgG HBcAg is seen in the late convalescence stage and remains lifelong.

Option A: In acute hepatitis B, HBsAg is reactive, HBeAg can be reactive in highly infectious individuals, and IgM Anti HBcAg is reactive. Anti-Hbs is reactive in recovery.

Option B: In chronic hepatitis B, HBsAg is reactive, and IgG Anti HBcAg is reactive. Anti-HBs is non-reactive.

Option D: Core window period is a phase where even though hepatitis infection is present, HBsAg is non-reactive. Diagnosis is based on the presence of IgM Anti HBcAg.

Solution to Question 25:

Statements 2 and 3 are true regarding the prevention of stroke in this patient.

Statement 1: Clopidogrel combined with aspirin is better than aspirin alone for stroke prevention. Warfarin is superior to antiplatelets in stroke prevention.

Statement 2: Warfarin is the recommended anticoagulant for patients with mitral stenosis (MS) and mechanical prosthetic valves.

Statement 3: Direct oral anticoagulants (DOACs) like dabigatran, rivaroxaban, apixaban, and edoxaban are not used in the treatment of moderate to severe MS and in patients with prosthetic heart valves due to limited clinical evidence.

DOACs have a lower risk of intracranial hemorrhage compared to warfarin and are as effective as warfarin for stroke prevention in valvular heart disease (except moderate to severe mitral stenosis or a mechanical heart valve), ischemic stroke or TIA, and AF.

Statement 4: Mitral valvotomy is not recommended for the prevention of stroke in this patient. It is indicated in symptomatic (New York Heart Association Functional Class II-IV) patients with isolated severe mitral stenosis, whose effective orifice (valve area) is < 1.5 cm2 in normal-sized adults. It is contraindicated in cases of left atrial thrombus.

Solution to Question 26:

Doing EEG is not mandatory for the diagnosis of seizures. The diagnosis of a seizure in most cases is based only on clinical grounds when examination and laboratory studies are often normal.

All patients who are suspected to have a seizure disorder should be evaluated with an EEG. EEG may provide support to the clinical diagnosis, prognosis, and classification of the seizure disorder.

Other options:

Option A: Approximately 1–5% of the population will have epileptiform discharges on EEG. The presence of epileptiform activity on EEG is not specific to epilepsy. Even in known cases of epilepsy, the routine inter-ictal EEG can be normal in 60% of cases. Thus, the EEG cannot establish the diagnosis of epilepsy in many cases.

Option B: Scalp EEG recording is sometimes helpful in the localization of frontal lobe epilepsy but is usually normal or misleading. When focal seizures arise from regions distant from the scalp such as the inferior frontal lobe or medial temporal lobe, the EEG recorded during the seizure may be non-localizing.

Option D: Progressive multifocal leukoencephalopathy is a demyelinating disease of the central nervous system due to infection by the John Cunningham virus (JC) virus. Electroencephalography (EEG) is characterized by a slow background activity with triphasic waves and slow waves predominantly in the frontal regions.

Characteristic EEG findings in different conditions:

• Typical absence seizures: 3-Hz spike-and-slow-wave discharges

• Lennox-Gastaut syndrome: Slow (<3 Hz) spike-and-wave discharges

• Generalized tonic clonic seizure (GTCS) - in tonic phase, increasing generalized low-voltage fast activity, followed by high-amplitude, polyspike discharges & in clonic phase a spike-and-slow-wave pattern is seen

• Creutzfeldt-Jakob disease: generalized periodic sharp waves

• Hepatic encephalopathy: triphasic waves

Solution to Question 27:

The clinical scenario describes an elderly patient with heart failure who develops severe respiratory distress associated with hypoxia, tachycardia, and elevated mean arterial pressure following a blood transfusion. The most likely diagnosis is transfusion-associated circulatory overload (TACO). The recommended investigations for suspected TACO include chest X-ray and BNP levels.

Chest X-ray in TACO may show bilateral and predominant alveolar infiltrates. BNP levels are usually elevated in TACO secondary to volume overload.

Anti-leukocyte antibodies are present in transfusion-related acute lung injury (TRALI), not in TACO patients.

TACO is usually a clinical diagnosis suspected in any patient who has acute respiratory distress and/or hypertension during transfusion or within 6 hours of completion.

Risk factors include older age, pre-existing fluid overload, cardiac dysfunction, renal dysfunction, and transfusion of large volumes of blood components.

Management of TACO involves discontinuing the transfusion and providing oxygen and diuretics.

Identification of at-risk patients, a slow transfusion rate (1 PRBC over 3-4 hours), close monitoring, and the use of diuretics in hemodynamically stable patients with a history of TACO can prevent TACO.

Features	TRALI	TACO
Body temperature	Fever +	Fever +/-
Blood pressure	Hypotension	Hypertension
Respiratory sympto ms	Acute dyspnea	Acute dyspnea
Neck veins	Unchanged	Distended

Features	TRALI	TACO
Chest radiograph	Diffuse bilateral infiltrates	Diffuse bilateral infiltrates
Ejection fraction	Normal	Decreased
Pulmonary edema fl uid	Exudate	Transudate
Fluid balance	Neutral or negative	Positive
Response to diuretic s	Inconsistent	Significant improvement
White cell count	Transient leukopenia may be present	Unchanged
BNP	Maybe elevated in critically il l patients with TRALI	Elevated

Solution to Question 28:

The given clinical scenario of a patient with dyspnea, bilateral pitting pedal edema, abdominal distension, elevated jugular venous pulse (JVP) and a palpable liver are suggestive of right ventricular (RV) failure. The history of chronic smoking points towards chronic obstructive pulmonary disease (COPD) as the etiology for RV failure (cor pulmonale). The label 2 points to the right ventricle (RV), which is affected in this patient.

Cor pulmonale is defined as altered RV structure/function secondary to pulmonary hypertension in the context of chronic lung diseases.

The main pathophysiology behind cor pulmonale is pulmonary hypertension and increased right ventricular (RV) afterload which is sufficient to cause RV dilation with or without hypertrophy. Sustained pressure overload leads to RV dysfunction and failure.

Dyspnea is the most common symptom. Signs include elevated JVP with prominent v waves, hepatomegaly, pulsatile liver, ascites, pedal edema, S3 gallop and murmur of tricuspid regurgitation.

The ECG in severe pulmonary hypertension shows p-pulmonale, right axis deviation, and RV hypertrophy. Echo can be used to measure RV wall thickness and chamber dimensions, while doppler can be used to assess pulmonary artery pressures.

A specimen of the heart with the various parts labeled is given below:


Aorta
Right auricle
Pulmonary trunk
Right atrium
Left atrium
Left auricle
Right ventricle
Left ventricle

Medicine NEET 2021

Question 1:

A 37-year-old woman presents with headaches for 6 months. She has been taking analgesics regularly. The headache recently increased in severity for 3 days but was reduced on stopping the analgesic. What is the likely diagnosis?

a) Medication overuse headache

b) Tension headache

c) Chronic migraine

d) Cluster headache

Question 2:

A 20-year-old woman presents with breathlessness and chest pain. She is a known case of mitral stenosis. Her pulse is irregularly irregular. No thrombus is seen on echocardiography. What is the best agent to prevent future thrombotic events?

a) Dabigatran

b) Aspirin 150mg

c) Oral warfarin

d) Aspirin + Clopidogrel

Question 3:

Evalutaion of a patient revealed the presence of a mid diastolic murmur. JVP showed prominent a-waves. What is the likely diagnosis?

a) Mitral stenosis

b) Tricuspid stenosis

c) Mitral regurgitation

d) Tricuspid regurgitation

Question 4:

A chronic alcoholic male patient presented with abdominal distension, reduced urine output, and pedal edema. His serum creatinine was 1.6 mg/dL. What is the next line of management in this patient?

a) Methylprednisolone

b) Heparin

c) Torsemide

d) Octreotide plus albumin

Question 5:

A 68-year-old man presents with cough with yellowish sputum. Auscultation revealed bronchial breath sounds. He is hemodynamically stable and not confused. On examination, his respiratory rate is 20/min and blood pressure is 110/70 mmHg. Lab reports show urea levels of 44 mg/dL. What is the next best step in the management of this patient?

a) Give antibiotics and send the patient home

b) Admit to ICU without mechanical ventilation

c) Admit to ICU with invasive mechanical ventilation

d) Consider admission in a non-ICU setting

Question 6:

A 30-year-old man presented with excessive fatiguability towards the end of the day that improved with rest. He also gives a history of ptosis and difficulty in speech and swallowing. What is the most likely diagnosis?

a) Myasthenia gravis

b) Lambert-Eaton syndrome

c) Duchenne muscular dystrophy

d) Systemic lupus erythematosus

Question 7:

A woman with hysteria who was hyperventilating developed carpo-pedal spasm. What could be the likely cause for this?

a) Respiratory acidosis

b) Metabolic acidosis

c) Respiratory alkalosis

d) Metabolic alkalosis

Question 8:

A 10-year old child presented with recurrent pulmonary infections, and bulky greasy stools. Quantitative estimation of the stool fat was more than 10gm/day. Which of the following will be seen in this child?

a) Protein losing enteropathy

b) Distal intestinal obstruction

c) Hypernatremia

d) Rectal prolapse occurs after treatment

Question 9:

A 36-year-old woman presented with claudication in the forearm, transient loss of vision, and abdominal pain. Femoral pulses were weak. Fundus examination revealed retinal hemorrhages. What is the likely diagnosis?

a) Polyarteritis nodosa

b) Thromboangitis obliterans

c) Takayasu arteritis

d) Microscopic polyangitis

Question 10:

A diabetic woman is admitted with complaints of vomiting. She gives a history of consuming food from outside. On evaluation, her blood pressure is 90/60 mmHg. Her arterial blood gas (ABG) report shows the following findings. What is the cause of her ABG findings?

a) Diabetic ketoacidosis

b) Persistent vomiting

c) Septic shock

d) Renal tubular acidosis

Question 11:

A patient presents with fever, nocturnal cough, breathlessness, and wheezing for 4 weeks. Absolute eosinophil count >5000/µL. Chest X-ray shows a miliary pattern. What is the likely diagnosis?

a) Bronchial asthma

b) Miliary tuberculosis

c) Tropical pulmonary eosinophilia

d) Hypersensitivity pneumonitis

Question 12:

A middle-aged man with chronic renal failure on dialysis presents with sudden collapse to the casualty with labored breathing. ECG shows tall tented T waves. What is the most likely acid-base imbalance that can be seen in this patient?

a) pH- 7.14, pCO2- 20 mm Hg, HCO3- 5 mEq/L

b) pH- 7.14, pCO2- 20 mm Hg, HCO3- 34 mEq/L

c) pH - 7.54, HCO3 - 27 mEq/L, pCO2- 34 mm Hg

d) pH- 7.4, HCO3- 27 mEq/L, pCO2 - 40 mm Hg

Question 13:

A woman presents with numbness of her fingertips. On examination, her face is tightened. The antinuclear antibody(ANA) is found to be positive and immunofluorescence shows the nucleolar pattern. What is the likely diagnosis?

a) Systemic sclerosis

b) Sjogren's syndrome

c) Systemic lupus erythematosus

d) Rheumatoid arthiritis

Question 14:

A 45-year-old woman with a history of hypertension presented to the emergency room with loss of consciousness, chest pain, and diaphoresis. On examination, she was grossly unstable and the bilateral pulses were unequal. ECG showed nonspecific ST-T changes. What is the next best investigation?

a) MRI

b) Transesophageal echocardiography

c) Cardiac enzymes

d) X ray

Question 15:

A 10-year-old child with AIDS presented with fever and productive cough. On auscultation, bronchial breath sounds and crepitations were heard in the right infra-scapular region. Chest X-ray showed right lower lobe consolidation. The CD4 count was 55 cells/mm3. What is the most common causative organism for this condition?

a) Pneumocystis jirovecii

b) Streptococcus pnuemoniae

c) Staphylococcus aureus

d) Mycoplasma

Question 16:

A patient presents with a seizure. On evaluation, urine osmolality is 1000 mOsm/Kg and serum osmolality is 270 mOsm/Kg. Which electrolyte abnormality can be expected in this patient?

a) Hypernatremia

b) Hyponatremia

c) Hyperkalemia

d) Hypokalemia

Question 17:

A male patient had feeble femoral pulses with upper-limb blood pressure of 186/90 mmHg. Chest X-ray showed enlarged intercostal arteries. What is the likely diagnosis?

a) Coarctation of aorta

b) Atrial septal defect

c) Bicuspid aortic valve

d) Patent ductus arteriosus

Question 18:

A patient who is a known case of atherosclerosis underwent a circumflex artery bypass. The stent is now placed successfully. He is already taking lisinopril, verapamil, metoprolol. Which drug should be added?

a) PDE 3 inhibitor

b) Direct oral anticoagulant

c) P2Y12 receptor blocker

d) PDE 5 inhibitor

Question 19:

Which of the following statement is incorrect regarding pheochromocytoma?

a) Diagnosed by urine VMA & catecholamines

b) Surgical excision is the definitive treatment

c) Propranolol is given initially to manage hypertension

d) Can present as hypertension alone and sometimes with vomiting and pain abdomen

Question 20:

A patient with giant cell arteritis presented with headache, jaw claudication, polymyalgia rheumatica, and mononeuritis multiplex. What should be the first-line treatment for this patient?

a) Abatacept

b) Tocilizumab

c) Steroids

d) Aspirin

Question 21:

A 50-year-old man presents with a history of recurrent retrosternal chest pain with each episode lasting for 3-5min and subsiding with sublingual nitrate. ECG shows left ventricular hypertrophy and flat T-wave. He is a known case of hypertension, diabetes mellitus, hypercholesterolemia currently on aspirin, atenolol, metformin, and lovastatin. What is the next best step in management?

a) IV glyceryl trinitrate infusion

b) Injection enoxaparin

c) Add clopidogrel

d) Increase the dose of beta blocker

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	d
5	d
6	a
7	c
8	b
9	c
10	b
11	c
12	a
13	a
14	b
15	b
16	b
17	a
18	c
19	c
20	c
21	b

Detailed Explanations

Solution to Question 1:

The clinical features are suggestive of medication overuse headache.

Medication overuse headache:

It is also known as analgesic rebound headache, drug-induced headache, or medication-misuse headache. In this condition medications that are used for the treatment of headaches become the cause for headaches. It usually resolves when the overuse is stopped.

Diagnostic criteria: According to ICHD-3 (International Classification of Headache Disorders-3)

• Headache occurring on ≥15 days/month in a patient with a pre-existing headache disorder

• Regular overuse for >3 months of one or more drugs that can be taken for acute and/or symptomatic treatment of headache.

• Not better accounted for by another ICHD-3 diagnosis.

Management: It includes the following steps:

• Withdrawal of the overused medication

• Rescue therapy or the treatment of withdrawal symptoms

• Initiation of preventative therapy if required

• Alternative symptomatic medication for future headaches

Other options:

Option B: Tension headache is a chronic headache syndrome characterized by bilateral, tight band-like discomfort. It can be episodic or chronic in nature. It is usually not associated with nausea, vomiting, photophobia, photophobia, etc.

Option C: Chronic migraine - Diagnostic criteria for migraine headache is as follows: Repeated attacks of headache lasting 4-72 hours in patients with normal physical examination, no other reasonable cause for headache, and

Option D: Cluster headache is characterized by deep, excruciating, nonfluctuating pain which is usually retroorbital. The classical feature of this condition is its periodicity. At least one of the daily attacks of pain recurs at almost the same hour every day.

Typically the patient has daily bouts of one or two attacks of relatively short-duration unilateral pain for 8–10 weeks a year. In between the clusters, there is a pain-free interval.

At least 2 of the following features	Plus at least 1 of the following features
Unilateral pain	Nausea or vomiting
Throbbing pain	Photophobia and phonophobia
Aggravation by movement	
Moderate or severe intensity	

Solution to Question 2:

The clinical presentation of breathlessness, chest pain, and irregularly irregular pulse is suggestive of atrial fibrillation. Since the patient is having mitral stenosis oral warfarin is the best agent to reduce the risk of thrombotic events.

Anticoagulation in atrial fibrillation:

Anticoagulation is indicated for patients with mitral stenosis, hypertrophic cardiomyopathy, previous history of stroke. In nonvalvular atrial fibrillation, the risk of stroke and the need for anticoagulation is assessed with the CHA2DS2-VASc score.

Anticoagulants that can be used include:

• Vitamin K antagonist - Warfarin

• Antithrombin inhibitor - Dabigatran

• Factor Xa inhibitors - Rivaroxaban, Apixaban, and Edoxaban

In AF patients with rheumatic mitral stenosis or mechanical heart valves, the anticoagulant of choice is warfarin. The newer anticoagulants are not yet tested for rheumatic heart disease.

Warfarin is a less compliant drug because it requires routine PT/INR monitoring for dose adjustments and has many drug interactions. The newer oral anticoagulants are easy to use and do not require PT/INR monitoring for dose adjustments.

Bleeding is an important risk of anticoagulation. The newer oral agents have a lower risk of intracranial bleeding when compared to warfarin. The risk factors for bleeding include age >65–75 years, renal insufficiency, heart failure, prior bleeding, and excessive alcohol or nonsteroidal anti-inflammatory drug use.

Reversal agents for anticoagulants are as follows:

• Warfarin - Vitamin K and fresh frozen plasma

• Dabigatran - Idarucizumab

• Factor Xa inhibitors - Andexanet alpha and Ciraparantag

Solution to Question 3:

The presence of a mid-diastolic murmur and a prominent a-wave in JVP is suggestive of tricuspid stenosis.

Mid-diastolic murmurs result from obstruction and/or augmented flow at the level of the mitral or tricuspid valve. Thus, both mitral stenosis (MS) and tricuspid stenosis (TS) present with a mid-diastolic murmur. However, the presence of a prominent a-wave in JVP is pointing more towards tricuspid stenosis.

Causes for mid-diastolic murmur:

• Mitral stenosis

• Tricuspid stenosis

• Atrial myxomas

• Austin flint murmur in severe aortic regurgitation

• Carey-coombs murmur in acute rheumatic fever

Tricuspid stenosis:

It always occurs in association with MS and is rheumatic in origin.

Mid-diastolic murmur in TS is best heard along the left sternal border and over the xiphoid process. Murmur is augmented during inspiration and reduced during expiration.

The jugular veins are distended and a giant a-wave is seen in JVP. There is a slow y descent due to impedance of right atrial emptying during diastole.

ECG shows features of right atrial enlargement which includes tall, peaked P wave in lead II and a prominent, upright P wave in lead V1.



Other options:

Option A: Mid-diastolic murmur is seen in mitral stenosis (MS)

It occurs most commonly in rheumatic fever.

Loud S1 (first heart sound) with an opening snap following S2 (second heart sound) is heard. S2 is followed by a low-pitched, rumbling, mid-diastolic murmur which is best heard at the apex with the patient in a lateral recumbent position.

Option C: Holosystolic murmur is seen in mitral regurgitation (MR).

The major causes of MS are infective endocarditis, myocardial infarction, rheumatic fever, and cardiomyopathies.

Soft S1 with a widely split S2 and a low-pitched S3 (third heart sound) is heard in MR. In addition, a holosystolic murmur is prominent at the apex and radiates to the axilla.

Option D: Holosystolic murmur is seen in tricuspid regurgitation (TR).

TR is most commonly due to functional defects such as tricuspid annular dilatation and leaflet tethering caused by rheumatic heart disease, myocardial infarction, etc.

The jugular vein is distended. Prominent c-v waves and rapid y descents are seen in severe TR.

A blowing holosystolic murmur along the lower left sternal margin, which is augmented during inspiration (Carvallo's sign) and reduced during expiration is seen in TR.



Solution to Question 4:

The given features of abdominal distension in a chronic alcoholic are suggestive of liver failure and reduced urine output, pedal edema and elevated serum creatinine are suggestive of renal failure. This points towards the diagnosis of hepatorenal syndrome. Among the given options, octreotide plus albumin will be the next line of management.

Octreotide is a somatostatin analogue. It causes splanchnic vasoconstriction and improves renal perfusion. Intravenous albumin is a volume expander and is effective in the management of hepatorenal syndrome. It is used in combination with other vasoconstrictor agents like terlipressin or octreotide.

A hepatorenal syndrome is a form of functional renal failure without renal pathology that occurs in patients with advanced cirrhosis or acute liver failure.

There is an increased splanchnic production of vasodilators (nitric oxide) in patients with cirrhosis and portal hypertension. Splanchnic vasodilation leads to systemic arterial underfilling resulting in activation of the renin-angiotensin-aldosterone system (RAAS). RAAS activation causes renal vasoconstriction which leads to decreased renal perfusion resulting in hepatorenal syndrome.

International Ascites Club's Diagnostic Criteria of Hepatorenal Syndrome:

Major criteria

• Chronic or acute liver disease with advanced hepatic failure and portal hypertension

• Low GFR: serum creatinine >133 μmol/L or 24h creatinine clearance <40 mL/min

• Absence of shock

• Absence of gastrointestinal or renal fluid losses

• No sustained improvement in renal function following diuretic withdrawal and expansion of plasma volume with 1.5 L of isotonic saline

• Proteinuria <0.5 g/d and no ultrasonographic evidence of obstructive uropathy or parenchymal renal disease

Additional criteria

• Urine volume <0.5 L/d

• Urine sodium <10 mmol/L

• Urine osmolality > plasma osmolality

• Urine red blood cells <50 high power field

• Serum sodium concentration <130 mmol/L

Treatment:

The definitive treatment is liver transplantation.

Pharmacotherapy mainly consists of drugs that cause splanchnic vasoconstriction. They include vasopressin analogs (ornipressin, terlipressin), somatostatin analogs (octreotide), and alpha-adrenergic agonists (midodrine). Plasma volume expanders are also effective in the treatment of hepatorenal syndrome.

Other treatment options include dialysis and transjugular intrahepatic portosystemic shunt.

Option C: Torsemide is a diuretic. It may trigger hepatorenal syndrome because of intravascular volume depletion.

Solution to Question 5:

This patient should be considered for admission in a non-ICU setting.

Productive cough along with auscultatory findings of bronchial breath sounds in this patient is suggestive of community-acquired pneumonia.

The next step is to determine the need for hospitalization in community-acquired pneumonia based on the CURB-65 score.

CURB-65 scoring:

The table below shows the decision based on the CURB-65 score:

Calculating the CURB-65 score for the case given in the question:

• No confusion (score = 0)

• Urea is 44 mg/dL (score = 1)

• Respiratory rate is 20/min (score = 0)

• Blood pressure is 110/70 mmHg (score = 0)

• Age is 68 years (score = 1)

Therefore, the total score is 2, and the patient requires hospitalization in a non-ICU setting (medical ward).

Treatment in a non-ICU setting includes respiratory fluoroquinolones (moxifloxacin) or a beta lactam antibiotic (cephalosporins) plus a macrolide (azithromycin/clarithromycin).

Clinical factor	Score
Confusion	1
Bloodurea nitrogen > 20 mg/dL (> 7 mmol/L)	1
Respiratory rate ≥ 30 breaths per minute	1
Systolicblood pressure < 90 mm Hg orDiastolicblood pressure ≤ 60 mm Hg	1
Age ≥65years	1

Total Score	Decision
0	Treated as outpatient
1 or 2	Admission to the hospital
3 or more	Admission in ICU

Solution to Question 6:

This patient has symptoms of fatiguability that worsen with repetitive movements but resolve after rest, along with oculobulbar symptoms such as ptosis, dysarthria, and dysphagia. From the given clinical picture, myasthenia gravis is the most likely diagnosis.

Myasthenia gravis is an autoimmune neuromuscular junction disorder that is characterized by weakness and fatigability of skeletal muscles.

It is due to the autoantibody-mediated destruction of the acetylcholine receptors (AChRs) at the post-synaptic membrane. This consequentially leads to a decreased transmission at the neuromuscular junction. Failure of transmission results in muscle weakness.

Females are more commonly affected than males and the disease tends to occur in middle age.

Clinical features of myasthenia gravis:

• Muscle weakness and fatiguability increase with repeated use or during late in the day and may improve following rest or sleep. Limb weakness is proximal and asymmetrical

• Cranial muscles particularly the lids and extraocular muscles are involved early in the course - diplopia, ptosis

• Facial weakness - snarling expression on smiling

• Weakness in chewing

• Speech with nasal timbre due to the weakness of palate and tongue

• Difficulty in swallowing

• Deep tendon reflexes are intact

• Myasthenic crisis - acute exacerbation of muscle weakness especially intercostal and diaphragmatic muscles resulting in ventilatory failure. It is most commonly due to infection and anticholinesterase medication overuse.

Myasthenia gravis is associated with thymoma and other autoimmune diseases such as Hashimoto's thyroiditis, Graves' disease, neuromyelitis optics, and systemic lupus erythematosus.

Diagnosis:

• Ice-pack test - application of ice pack over the eyes improves ptosis (decreases activity of acetylcholinesterase, this increases the amount of acetylcholine in the synaptic cleft and increase transmission)

• Antibodies associated with myasthenia gravis: Anti-AChR antibodies, Anti-MuSK antibodies, Anti-LRP4 antibodies

• Repetitive nerve stimulation test shows a decremental response.

• Edrophonium chloride or Tensilon test - improvement of symptoms occurs on injecting edrophonium. Edrophonium is an acetylcholinesterase inhibitor. This increases the amount of acetylcholine in the synaptic cleft and improves transmission.

Treatment:

• Anticholinesterase drugs - Pyridostigmine

• Thymectomy

• Immunosuppression therapy with glucocorticoids, azathioprine, tacrolimus, rituximab, etc.

• Plasmapheresis and intravenous immunoglobulin

Other options:

Option B:

Lambert-Eaton myasthenic syndrome (LEMS) is an autoimmune disorder affecting the presynaptic membrane at the neuromuscular junction. It is due to the autoantibodies against the P/Q-type calcium channels at the motor nerve terminal which results in the impaired release of Ach from the presynaptic nerve terminal.

Patients present with proximal muscle weakness of the lower limbs, ptosis, and diplopia. However, symptoms might improve after brief exercise. Deep tendon reflexes are absent or reduced. They also experience autonomic dysfunction and present with dry mouth and impotence. It is most commonly associated with small cell lung cancer.

Repetitive nerve stimulation test shows an incremental response at higher rates.

Treatment involves plasmapheresis, immunotherapy, and drugs such as 3,4-diaminopyridine (3,4-DAP) and pyridostigmine.

Option C:

Duchenne muscular dystrophy (DMD) is X-linked muscular dystrophy due to mutations in the dystrophin gene. It most commonly affects boys in their childhood.

Patients initially present with proximal muscle weakness predominantly of the lower limbs. Calf muscle hypertrophy is a cardinal feature. Other features include mental retardation, cardiac failure, and myoglobinuria.

Elevated serum creatinine kinase levels and genetic testing are diagnostic of DMD.

Option D:

Systemic lupus erythematosus (SLE) is an autoimmune disease that most commonly presents with arthralgia with early morning stiffness. It predominantly occurs in middle-aged women. Other important features of SLE include malar rash, discoid rash, photosensitivity, oral ulcers, pleuritis, pericarditis, and lupus nephritis.

Anti-nuclear antibodies, Anti-dsDNA antibodies, anti-Smith antibodies are positive in SLE.

Solution to Question 7:

Respiratory alkalosis, developed secondary to hyperventilation, is the likely cause of the carpopedal spasm.

Hyperventilation leads to the washout of carbon dioxide, resulting in decreased levels of $PaCO_2$ (hypocapnia), which in turn increases the pH (alkalosis). Thus, acute respiratory alkalosis occurs during hyperventilation. An increase in extracellular pH increases the binding of free calcium to albumin, lowering the ionized calcium concentration. This leads to hypocalcemia.

The clinical features of hypocalcemia are:

• Tetany: Increased peripheral neuromuscular irritability. May include perioral paresthesias, muscle spasms, and cramps.

• Trosseau's sign: Induction of carpal spasm when the BP cuff is inflated 20 mm Hg over the systolic BP for three minutes or more.

• Chvostek's sign: Contraction of the ipsilateral circumoral facial muscles when the facial nerve is tapped just anterior to the ear.

• Severe hypocalcemia can cause seizures, carpopedal spasm, bronchospasm, and laryngospasm.

Treatment of the underlying condition is necessary. The patient should be reassured and rebreathing from a paper bag should be done during acute episodes.

The image below shows Trosseau's sign.



The image below shows Chvostek's sign.

Solution to Question 8:

The given clinical scenario showing a young child with recurrent pulmonary infections and steatorrhea with stool fat more than 10 gm/day is suggestive of cystic fibrosis (CF). A distal intestinal obstruction is a common complication seen in this age group.

Protein-losing enteropathy occurs due to exocrine pancreatic insufficiency and is seen most commonly in infancy (Option A). Rectal prolapse was one of the common complications in the past but occurs less frequently due to earlier diagnosis and initiation of pancreatic enzyme replacement therapy (Option D). Individuals with CF develop hypovolemic hyponatremia by sodium loss via sweat through a defective chloride ion transport channel, the CF transmembrane conductance regulator (CFTR), and hence hypernatremia is not seen in cystic fibrosis (Option C).

Cystic fibrosis is an autosomal recessive exocrinopathy affecting multiple epithelial tissues caused primarily due to a mutation in the CF transmembrane conductance regulator [CFTR] gene which serves as an anion channel in the apical membranes of epithelial cells and regulates the volume and composition of exocrine secretion. A defect in this gene results in thick, hyper-viscous exocrine secretions which lead to the primary pathologies seen in cystic fibrosis.

Disease manifestations include:

• Respiratory: Copious hyper-viscous and adherent pulmonary secretions obstruct small and medium-sized airways leading to respiratory compromise. Bacterial flora (including Staphylococcus aureus, Haemophilus influenzae, and Pseudomonas aeruginosa) is routinely cultured from CF sputum.

• Pancreas: Thick and adherent exocrine secretions obstruct pancreatic ducts and impair the production and flow of digestive enzymes. There is tissue destruction of the exocrine pancreas, with fibrotic scarring, cyst proliferation, and loss of acinar tissue.

• Hepatic: Obstruction of intrahepatic bile ducts and parenchymal fibrosis leading to hepatic insufficiency and multilobular cirrhosis.

• GIT: Contents of the intestinal lumen are often difficult to excrete, leading to meconium ileus in the newborns and distal intestinal obstructive syndrome in adults. With early diagnostic modalities and pancreatic enzyme replacement therapy, the occurrence of rectal prolapse is fairly rare.

• Infertility:

• Males- developmental abnormality secondary to improper secretion by the vas or associated structures leading to complete involution of the vas deferens and infertility. About 99% of males with CF are infertile

• Females- abnormalities of the reproductive tract secretions leads to an increased incidence of infertility

• Sinusitis: Present in most CF patients

Solution to Question 9:

The given clinical scenario of a patient presenting with claudication in the forearm, weak femoral pulses, abdominal pain and transient loss of vision is suggestive of Takayasu arteritis.

A patient with polyarteritis nodosa will present with abdominal pain, renal involvement, and hypertension. Musculoskeletal symptoms present as arthralgia, arthritis, or myalgia. The presence of large vessel symptoms such as claudication in the forearm and weak femoral pulses makes the diagnosis of polyarteritis nodosa unlikely (option A).

Takayasu arteritis is a chronic inflammatory medium- and large- vessel vasculitis that commonly affects adolescents and young females. It mainly involves the aortic arch and its branches. Depending on the artery occluded, patients present with the following symptoms:

• Subclavian artery (most common) - pain, numbness, claudication, and weak or absent radial pulses on the affected side

• Common carotid artery - visual symptoms, syncope, and stroke

• Abdominal aorta, superior mesenteric artery, and coeliac axis - abdominal pain, nausea, vomiting

• Renal artery - hypertension and renal failure

• Aortic arch or root - aortic insufficiency, heart failure

• Pulmonary artery - atypical chest pain, dyspnea

• Coronary artery - chest pain, myocardial infarction

• Iliac artery - Leg claudication, weak/absent pulses in the periphery

Patients also have constitutional symptoms such as malaise, fever, night sweats, weight loss, etc.

Investigations:

• Lab reports reveal an elevated ESR level, anemia, and elevated immunoglobulin levels.

• Arteriography is done as a confirmatory test.

Treatment includes glucocorticoid therapy, methotrexate, immunotherapy with anti-TNF agents, and surgery of the stenosed vessels.

Other options:

Option A: Polyarteritis nodosa (PAN) is necrotizing arteritis of the medium-sized blood vessels. It can affect the skin, peripheral nerves, renal vessels, mesenteric vessels, heart, and brain. Lungs are usually spared in PAN.The disease initially starts with non-specific symptoms like fever, malaise, arthralgia, and myalgia. Later, depending on the artery involved, patients may develop the following symptoms:

• Skin findings - ulcers, digital gangrene, subcutaneous nodules, and livedo reticularis

• Renal arteries - hypertension and renal failure

• Mesenteric arteries - abdominal pain, nausea, vomiting, bleeding, and a bowel perforation

Patients can have musculoskeletal symptoms (arthralgia, myalgia) and mononeuritis multiplex.

Option B: Thromboangiitis obliterans, also known as Buerger's disease, is an inflammatory occlusive vascular disorder involving medium- and small-sized vessels in the distal extremities. It is most frequently seen in male smokers less than 40 years of age.The patient presents with a triad of claudication of the distal extremities, Raynaud's phenomenon, and migratory superficial vein thrombophlebitis. Proximal pulses (brachial artery, femoral artery, popliteal artery) are normal. But distal pulses (radial, ulnar, tibial artery) are weak/absent.

Option D: Microscopic polyangiitis is a necrotizing vasculitis affecting small- and medium-sized vessels. It occurs most commonly in elderly males. Patients can present with non-constitutional symptoms such as fever, weight loss, and musculoskeletal pain. Renal involvement is very common (79%) in the form of glomerulonephritis, which can rapidly progress, causing renal failure. It is also associated with mononeuritis multiplex, alveolar hemorrhage (present with hemoptysis as 1st symptom), and cutaneous vasculitis. Lab findings include anemia, raised ESR and or CRP, leukocytosis, thrombocytopenia, and ANCA positive in 75% of patients,

Solution to Question 10:

This patient has hypovolemic, hypochloremic, hypokalemic, hyponatremic metabolic alkalosis with respiratory alkalosis. This is most probably due to persistent vomiting.

Inference from the given question stem:

Step 1: pH in this patient is 7.52 which is alkalemia.

Step 2: Bicarbonate level in this patient is 30 mEq/L which is increased. This signifies the presence of an excess of base. Hence, this is metabolic alkalosis.

Step 3: $PaCO_2$ level in this patient is 20 mmHg, which is less than normal. This signifies respiratory alkalosis.

Step 4: Electrolyte imbalance -

• Na - 123 mEq/L signifies hyponatremia

• K - 3.2 mEq/L signifies hypokalemia

• Cl - 67 mEq/L signifies hypochloremia

Step 5: Volume depletion - Blood pressure of 90/60 mmHg signifies hypovolemia due to persistent vomiting.

Vomiting leads to loss of Hcl (acid) from the stomach resulting in metabolic alkalosis and hypochloremia. This causes failure of secretion of HCO_3^- into the small intestine and HCO_3^- is added into the extracellular fluid.

Increased level of HCO_3^- will be excreted in the urine making the urine alkaline with increased potassium and sodium excretion in the urine. This leads to hypokalemia and hyponatremia.

Other options:

Option A: Diabetic ketoacidosis is associated with high anion gap metabolic acidosis along with hyponatremia and hypochloremia. It may initially present with hyperkalemia.

Option C: Septic shock is characterized by high anion gap metabolic acidosis.

Option D: Renal tubular acidosis is characterized by hyperchloremic, normal anion gap metabolic acidosis.

Solution to Question 11:

The clinical features along with elevated absolute eosinophil count and miliary pattern in chest x-ray are suggestive of tropical pulmonary eosinophilia.

Tropical pulmonary eosinophilia (TPE) is a hyperresponsive pulmonary syndrome that occurs in response to trapped microfilariae within the lung parenchyma. It is a type 1 hypersensitivity reaction.

Clinical features:

The patient has a history of residence in a filaria endemic area and usually presents with nocturnal paroxysmal wheeze and cough. Other manifestations include weight loss, low-grade fever, and lymphadenopathy. Hepatosplenomegaly may be seen due to the trapping of microfilariae in the reticuloendothelial system.

Investigations:

• Eosinophilia > 3000/µL, leucocytosis

• X-ray chest: Mottled opacities or military lesions in the middle and lower lung fields, increased bronchovascular markings

• PFT: Restrictive changes in most cases and obstructive defects in half

• Elevated levels of IgE and Antifilarial antibody titers

• Microfilaria are sequestrated in the lungs and hence not found in blood

Treatment: Diethylcarbamazine(DEC) at a daily dosage of 4-6 mg/kg for 14 days.

Differntial Diagnosis:

• Bronchial asthma

• Loffler syndrome

• Allergic bronchopulmonary aspergillosis

• Churg strauss syndrome

•

Chronic eosinophilic pneumonia

•

Hypereosinophilic syndromes

Other options:

Option A: The classical symptoms of bronchial asthma are wheezing, dyspnea, and cough which is worse at night and keeps the patient awake in the early morning hours. Chest auscultation reveals rhonchi throughout the chest. The chest x-ray is usually normal but hyperinflation may be seen in severe patients.

Option B: Miliary TB occurs due to the hematogenous spread of tubercle bacilli. Clinical features are nonspecific and depend on the site of involvement. Common manifestations include fever, night sweats, anorexia, weakness, and weight loss. Blood investigations may reveal anemia with leukopenia, lymphopenia, neutrophilic leukocytosis, leukemoid reactions, and polycythemia. Chest x-ray shows a military reticulonodular pattern in some cases.

Option D: Hypersensitivity pneumonitis, also known as extrinsic allergic alveolitis, is an inflammatory response to various inhaled antigens. The clinical presentation can be acute, subacute, and chronic. History of exposure to the offending antigen is present. Chest x-ray findings are nonspecific and can show ground-glass opacities or ill-defined micronodular opacities.

Solution to Question 12:

Metabolic acidosis is the most common acid-base abnormality seen in chronic kidney disease (CKD) patients due to the defective production of ammonia which impairs the excretion of H+ ions.

In metabolic acidosis, there will be low pH (<7.35), low bicarbonates (<22 mEq/L), and low pCO2 levels (<35 mm Hg). Since the pH in options C and D is >7.35, they can be ruled out. A pH of 7.14 indicates acidosis. The bicarbonate level of 34 mEq/L (in option B) is not consistent with metabolic acidosis, thus leaving us with option A.

During metabolic acidosis, there is respiratory compensation by hyperventilation resulting in loss of volatile acid causing a reduction in PCO2 levels. The decrease in PCO2 with a decrease in pH indicates that it is a compensatory mechanism. We can use Winter's formula to check the degree of respiratory compensation in metabolic acidosis:

$pCO2 = [1.5 \times HCO3^-] + 8 \pm 2$

$= [1.5 \times 5] + 8 \pm 2$

$= 13.5\text{-}17.5$

The expected pCO2 level is 13.5-17.5 mmHg. The pCO2 value of 20mmHg in option A is close to the expected values. Hence, this patient has a partially compensated metabolic acidosis.

In CKD, the kidneys are unable to produce enough ammonia (NH_3) in the proximal tubules to excrete the endogenous acid (H^+) into the urine in the form of ammonium (NH_4^+). Therefore, accumulation of H^+ occurs causing metabolic acidosis. In the late stages of CKD, the accumulation of phosphates, sulfates, and other organic anions cause an increase in the anion gap. Thus, non-anion gap metabolic acidosis is seen in the early stages of CKD which then progresses to high anion gap metabolic acidosis in later stages of CKD.

In CKD, the ability to excrete potassium is decreased. This causes hyperkalemia. Also, metabolic acidosis in CKD causes an extracellular shift of K^+ and aggravates hyperkalemia. This is responsible for the ECG findings of tall tented T waves in this patient.

Solution to Question 13:

Numbness of fingers associated with Raynaud's phenomenon, facial tightening, positive anti-nuclear antibodies (ANA) with nucleolar pattern in immunofluorescence is highly suggestive of systemic sclerosis.

Systemic sclerosis/ scleroderma:

Systemic sclerosis (SSc) is an autoimmune connective tissue disorder that results in fibrosis of the skin, vasculature, and internal organs. It is of two types, namely diffuse cutaneous systemic sclerosis and limited cutaneous systemic sclerosis.

Scleroderma is characterized by Raynaud's phenomenon, digital ischemia, sclerodactyly, calcinosis, telangiectasia, and cardiac, lung, gastrointestinal and renal disease. Skin thickening is the hallmark of SSc. Patients have a characteristic expressionless facial appearance with taut and shiny skin and microstomia.

Antinuclear antibodies (ANA) are positive in SSC. The presence of anti-topoisomerase 1 (Scl70) and anticentromere antibodies are seen in diffuse cutaneous SSC and limited cutaneous SSc respectively.

Nucleolar pattern of ANA staining in immunofluorescence is most commonly seen in systemic sclerosis. A centromeric pattern is also seen.

Image below shows nucleolar pattern in immunofluorescence seen in systemic sclerosis.



Other options:

Option B: Sjogren's syndrome is an autoimmune disease characterized by lymphocytic infiltration of salivary and lacrimal glands.

Patients present with dry eye (keratoconjunctivitis sicca) and dry mouth (xerostomia) due to fibrosis of the lacrimal and salivary gland respectively.

Anti-Ro and anti-La antibodies are positive in Sjogren's syndrome. ANA and RF (Rheumatoid factor) can also be seen. Centromeric pattern and speckled pattern of staining is seen in Sjogren's syndrome.

Image below shows speckled pattern (image 1) and centromeric pattern (image 2) of staining respectively.





Option C: Systemic lupus erythematosus (SLE) is an autoimmune disease that most commonly presents with arthralgia with early morning stiffness.

Other important features of SLE include malar rash, discoid rash, photosensitivity, oral ulcers, pleuritis, pericarditis, and lupus nephritis.

ANA, Anti-dsDNA antibodies, anti-smith antibodies are positive in SLE. Homogenous or diffuse staining and peripheral staining pattern are most commonly seen in SLE.

Image below shows a homogenous or diffuse stained immunofluorescence pattern seen in SLE.



Option D: Rheumatoid arthritis (RA) is a multisystem disease that classically presents with pain, joint swelling, and stiffness of the small joints of the hands, feet, and wrists in a symmetrical manner.

RF and anti-citrullinated peptide antibodies are positive in RA. Homogenous staining is the most common pattern seen in RA.



Solution to Question 14:

The clinical presentation and ECG showing nonspecific ST-T changes are highly suggestive of aortic dissection. As the patient is hemodynamically unstable the next best investigation is transesophageal ECHO.

Aortic dissection:

This occurs when a defect or flap occurs in the intima of the aorta. It results in blood tracking into the aortic tissues and splitting of the medial layer of the aorta.

Predisposing factors:

• Age

• Hypertension

• Marfan syndrome

• Pregnancy

• Other connective tissue disorders, for example, Ehlers–Danlos syndrome, giant cell arteritis, systemic lupus erythematosus

• Coarctation of the aorta

• Turner or Noonan syndromes

• Aortic cannulation site (iatrogenic)

Clinical features:

The most typical presentation of aortic dissection is a tearing interscapular pain. It is associated with diaphoresis. Other symptoms are syncope, weakness, and dyspnea. On examination, there can be hypertension or hypotension, loss of pulses, findings of aortic regurgitation, pulmonary edema, or neurological manifestations. Unequal pulses or blood pressure in the extremities may be seen due to changes of flow in the true and false lumens.

ECG findings: An ECG that shows no findings of acute myocardial infarction helps to differentiate aortic dissection from myocardial infarction.

Imaging:

In hemodynamically stable patients the diagnosis is established with a contrast-enhanced CT scan. The characteristic diagnostic finding is a double lumen aorta.

Transesophageal ECHO(TEE) is the investigation of choice in a hemodynamically unstable patient.



Treatment:

The initial management is with beta-blockers to control hypertension, along with pain management. The preferred beta-blocker is esmolol. These reduce cardiac contractility and blood pressure.

Isolated use of a direct vasodilator such as hydralazine is contraindicated in acute dissection because these agents can increase hydraulic shear and worsen the dissection. CCBs like verapamil and diltiazem may be used intravenously if nitroprusside or beta-blockers cannot be employed.

Once hypertension is controlled, definitive surgical repair of aortic dissection is done. Treatment options depend on the classification.

• Stanford Type A (DeBakey types I and II) requires surgical intervention once the patient's blood pressure is stabilized.

• Stanford Type B (DeBakey type III) is best managed by antihypertensive drugs. Surgical intervention is indicated in complicated cases if the pain increases (signaling impending rupture) or when there is evidence of malperfusion of organs or neurological symptoms.



Solution to Question 15:

The given clinical features of productive cough with fever, bronchial breath sounds and crepitations on auscultation along with lower lobe consolidation on chest X-ray point towards the diagnosis of lobar pneumonia.

The most common causative organism is Streptococcus pneumonia, irrespective of HIV status. In AIDS patients, opportunistic infection-causing bacterial pneumonia occurs at any clinical stage and at any CD4 cell count. Pneumocystis jirovecii pneumonia would present with dry cough and interstitial infiltrates on chest x-ray (option A).

Patients with pneumococcal (lobar) pneumonia present with fever, dyspnea, and productive cough. On examination, involved lobes are dull to percussion, bronchial breath sounds, and crackles are heard on auscultation.

A chest radiograph reveals focal, segmental, or lobar consolidation of involved lobes.



Other options:

Option A:

Pneumocystis jiroveci causes pneumocystis pneumonia (PCP) in AIDS patients with CD4 counts below 200 cells/mm3.

PCP presents with fever, dry cough, and dyspnea. Inspiratory crackles are heard on auscultation. Chest X-ray reveals bilateral, symmetric, interstitial infiltration. Initially, infiltrates are perihilar, as the disease progress, it becomes diffuse.

The chest radiographs given below show ground-glass opacities in PCP



Option C:

Staphylococcus aureus causes bacterial pneumonia with features similar to pneumococcal pneumonia. The differentiating feature is that chest X-ray in Staph. aureus infection shows pneumatoceles (cavitation).

The X-ray shows pneumatoceles (air-filled cysts within lung parenchyma).



Option D: Mycoplasma causes atypical pneumonia. Patients present with dry cough, fever, chills, malaise, and upper respiratory tract symptoms. On auscultation, wheezes or rales are heard. Chest X-ray shows peribronchial pneumonia with prominent bronchial markings, interstitial infiltration, and atelectasis.

The chest X-ray below shows patchy consolidation at the hilum and fanning to the periphery



Solution to Question 16:

This patient has high urine osmolality of 1000 mOsm/kg (normal 500-800 mOsm/kg) and low plasma osmolality of 270 mOsm/kg (normal 285-295 mOsm/kg). This is seen in the syndrome of inappropriate antidiuretic hormone synthesis (SIADH) and hyponatremia is the most common

electrolyte abnormality seen.

Syndrome of inappropriate antidiuretic hormone ADH release (SIADH) is a condition where there is an unsuppressed release of antidiuretic hormone (ADH) from the pituitary gland or nonpituitary sources or its continued action on vasopressin receptors.

ADH promotes the reabsorption of water from the collecting duct. Excessive secretion of ADH in SIADH results in enhanced water reabsorption. Therefore, only water is reabsorbed and not the solutes resulting in concentrated urine. This is depicted as high urine osmolality. The excess of water reabsorbed from the tubules expand and dilute the extracellular fluid leading to dilutional hyponatremia. This is depicted as low plasma osmolality.

Since there is an increase in body water with little or no change in body sodium in SIADH, this is known as euvolemic hyponatremia.

Patients may be asymptomatic or present with headache, confusion, nausea, vomiting, anorexia, seizures, and coma.

Solution to Question 17:

The given clinical features of hypertension in the upper limb, weak lower limb pulse, and enlarged intercostal arteries on chest X-ray are pointing towards the diagnosis of coarctation of the aorta.

Coarctation of the aorta is a congenital heart defect characterized by the narrowing of the thoracic aorta near the insertion of the ductus arteriosus. It is associated with the bicuspid aortic valve, atrial septal defect, ventricular septal defect, mitral regurgitation, berry aneurysm, and Turner syndrome. It is more common in males.

Depending on the site of constriction it may be-

• Preductal - constriction proximal to the ductus arteriosus. (infantile type)

• Postductal - constriction distal to the ductus arteriosus (adult type).

Physical examination findings include-

• Bounding upper limb pulses (radial, carotid) and weak lower limb pulses (femoral, popliteal, posterior tibial, and dorsalis pedis)

• Increased Radio-femoral delay

• Hypertension in upper limbs; hypotension in lower limbs

• Difference in blood pressure of more than 20 mmHg between upper and lower limbs

• Continuous machinery like murmur (due to collaterals) is heard over the scapula

• Systolic ejection click and systolic ejection murmur in the left upper sternal border (due to bicuspid aortic valve)

Radiologically, the chest X-ray may reveal-

• Figure 3 sign - Pathognomonic of coarctation of the aorta. Due to indentation of the aorta at the site of coarctation and pre and post-stenotic dilatation along the left paramediastinum.



• Notching of ribs (Roesler's sign) due to inferior rib erosion by dilated intercostal collateral vessels.

• Collateral blood supply occurs from the arch of the aorta to the descending thoracic aorta bypassing the narrowed aorta.

• It occurs via the arch of the aorta → internal thoracic artery → anterior intercostal artery → posterior intercostal artery → descending thoracic aorta.

• 4th to 8th ribs are most commonly involved. 1st and 2nd ribs are spared as the intercostal arteries in these ribs do not communicate with the descending aorta.



Solution to Question 18:

P2Y12 receptor blocker should be added to prevent stent thrombosis.

Clopidogrel, a P2Y12 receptor blocker, along with aspirin is called dual antiplatelet therapy (DAPT).

DAPT should be continued for at least 1 year following stent placement to prevent stent thrombosis. After the recommended duration of treatment with dual antiplatelet therapy, aspirin should be continued indefinitely.

Solution to Question 19:

Alpha-blockers like phenoxybenzamine are given initially to manage hypertension in pheochromocytoma. Beta-blockers such as propranolol are given only after adequate alpha blockade is achieved.

Pheochromocytoma is a catecholamine secreting tumor. High levels of circulating catecholamines stimulate the alpha receptors on the blood vessels and cause vasoconstriction, resulting in hypertension. If β-blockers are given initially, it will lead to unopposed α-stimulant action of catecholamines leading to hypertensive crisis. This is the rationale behind prescribing alpha-blockers initially and then followed by β-blockers in pheochromocytoma.

Pheochromocytoma:

Pheochromocytoma is a catecholamine secreting tumor derived from the chromaffin cells of the adrenal medulla.

It can be sporadical or hereditary. Hereditary pheochromocytoma occurs in association with syndromes such as MEN 2 (Multiple endocrine neoplasia type 2), familial paraganglioma syndrome, Von Hippel Lindau (VHL) syndrome, and neurofibromatosis (NF) type 1.

Clinical features:

It most commonly presents as a classic triad of intermittent headache, palpitation, and profuse sweating along with hypertension. The patient can also present with anxiety, panic attacks, nausea, abdominal pain, weight loss, and weakness.

Symptoms may be precipitated by surgery, pregnancy, position changes, exercise, and medications such as metoclopramide and tricyclic antidepressants.

Investigations:

• Biochemical screening test - measurement of urinary and plasma levels of catecholamines and metanephrines.

• MRI is the investigation of choice for localizing the tumor. It is preferred over a CT scan as the contrast used in CT scans can precipitate paroxysms.

• Other imaging modalities like MIBG scintigraphy, DOPA- or FDG- PET scan are used in identifying the primary tumor as well as the extra-adrenal tumors and metastases.

Treatment:

Surgical resection of the tumor is the definitive treatment for pheochromocytoma.

Preoperatively, hypertension is managed initially with alpha-blockers followed by beta-blockers.

Postoperatively, lifelong yearly follow-up should be done.

Solution to Question 20:

Steroids are the first-line management for giant cell arteritis.

Giant cell arteritis (GCA), also known as temporal arteritis is an inflammation of medium and large-sized vessels. It is more common in women and people aged >50 years.

The clinical features include headache associated with scalp pain and jaw claudication, fever, fatigue, malaise, and weight loss. Half of the patients have signs of superficial temporal artery inflammation, including erythema, pain on palpation, nodularity, thickening, or reduced pulsation on the affected side. Gentle pressure on the scalp may elicit focal or generalized tenderness.

GCA is very commonly associated with polymyalgia rheumatica characterized by morning stiffness and pain around the shoulders and hip. It is also associated with mononeuritis multiplex which is characterized by peripheral neuropathy in two or more separate nerve areas.

A dreaded complication is ischemic optic neuropathy, which can lead to sudden blindness. Therefore, patients with a clinical diagnosis of GCA, especially with headaches or visual symptoms should immediately be started on steroids. If treatment is delayed beyond 48 hours, there may be irreversible damage.

Lab reports reveal an elevated ESR level, hypo-/normo-chromic anemia, elevated alkaline phosphatase levels, and increased levels of IgG and complement. The diagnosis can be confirmed by a temporal artery biopsy. It may not be positive in all patients, due to patchy involvement of the vessels. Obtaining a biopsy segment of 3-5 cm with serial sectioning may yield a positive result. A color doppler ultrasound can be used as an alternative.

The mainstay of treatment in GCA is steroids. Prednisolone 40-60 mg is given for one month followed by tapering doses. The duration of treatment is approximately 2 years or more. Methylprednisolone 1000 mg for 3 days should be given for patients presenting with visual symptoms.

Aspirin reduces the occurrence of visual loss and can be given in addition to steroids. Newer drugs used in the treatment of giant cell arteritis are tocilizumab (anti-IL-6 receptor) and abatacept (CTLA4-Ig). Both these drugs are used adjunctively to steroid therapy.

Histopathology of giant cell arteritis shows panarteritis with infiltration of inflammatory mononuclear cells with giant cell formation. Intimal thickening and fragmentation of the internal elastic lamina is also seen.

The image below shows internal elastic lamina fragmentation (black arrow) and giant cells (yellow arrow) in a cerebral artery.

Solution to Question 21:

The given clinical scenario of recurrent chest pain relieved on nitrates in a patient with hypertension, diabetes mellitus and hypercholesterolemia are suggestive of acute coronary syndrome. ECG showing flat T wave and left ventricular hypertrophy favors the diagnosis of non-ST segment elevated acute coronary syndrome (NSTE-ACS). Injection enoxaparin is the next best step in management.

Enoxaparin, a low molecular weight heparin, is used as a first-line anticoagulant therapy to prevent thrombosis in patients with NSTE-ACS.

Patients with acute coronary syndrome are classified into ST-segment elevated myocardial infarction (STEMI) and NSTE-ACS.

Management of NSTE-ACS:

Supportive therapy - supplemental oxygen

Anti-ischemic drugs:

• Sublingual or IV nitrates

• Beta-blockers

Antithrombotic treatment consists of anticoagulants and antiplatelet therapy.

The anticoagulants used are low molecular weight heparin such as enoxaparin, fondaparinux. Other anticoagulants used are unfractionated heparin (UFH), bivalirudin, and direct oral anticoagulants such as dabigatran, apixaban, and rivaroxaban. UFH is the mainstay of therapy.

Antiplatelet drug, aspirin, is the preferred drug in NSTE-ACS treatment. P2Y12 receptor inhibitors like prasugrel, ticagrelor, cangrelor, clopidogrel are also used. Clopidogrel is used only when prasugrel or ticagrelor are not available/ contraindicated.

GPIIb/IIIa inhibitors - eptifibatide and tirofiban - are also used as alternatives.

Invasive treatment - percutaneous coronary intervention- is done after stabilizing the patient with the above-mentioned drugs.

Other options:

Option A: Intravenous glyceryl trinitrate infusion is only given when chest pain is not relieved by sublingual nitrates. This patient has chest pain controlled by sublingual nitrates. Hence, IV GTN infusion is not recommended.

Option C: Clopidogrel is used in combination with aspirin or when aspirin is intolerable/contraindicated. However, prasugrel or ticagrelor is preferred over clopidogrel and is initiated after anticoagulant therapy.

Option D: Beta-blockers are used to reduce pain. This patient does not present with pain and also mentions that the pain is relieved on sublingual nitrates. Increasing the dose of beta-blocker has no benefit in this situation.

Medicine INI-CET 2022

Question 1:

A diabetic patient's fasting blood glucose level is found to be 160 mg/dL. What will you advise the patient regarding non-pharmacological management?

a) At least 80 mg dietary fibre

b) <5 g sodium intake everyday

c) <30% of the calories should come from fat

d) Cholesterol <100 mg

Question 2:

Which of the following is the most characteristic feature of Kallmann syndrome?

a) Anosmia

b) Syndactyly in males

c) Precocious puberty in females

d) White forelock

Question 3:

A chronic cigarette smoker now joined a construction company. His pulmonary function test results are given below. What is the most likely diagnosis for this patient?

a) Vascular disease with bronchodilator reversibility

b) Restrictive lung disease with bronchodilator reversibility

c) Restrictive lung disease without bronchodilator reversibility

d) Obstructive disease with bronchodilator reversibility

Question 4:

Which of the following is not a first-line drug for the management of a patient with rheumatoid arthritis?

a) Sulfasalazine

b) Hydroxychloroquine

c) Methotrexate

d) Azathioprine

Question 5:

AIDS-defining malignancies include all except :

a) Kaposi's sarcoma

b) Melanoma

c) Invasive cervical carcinoma

d) Non hodgkins lymphoma

Question 6:

A 40-year-old chronic alcoholic presented with withdrawal symptoms. His liver function tests showed AST 140 IU/L, ALT 110 IU/L, and GGT 500 IU/L. Which of the following benzodiazepines can be safely administered to this patient?

a) Diazepam

b) Alprazolam

c) Clonazepam

d) Lorazepam

Question 7:

Which of the following does not help in the localization of lesions in the spinal cord?

a) Contralateral hemiplegia

b) Fasciculation at the level of lesion

c) Upper motor neuron lesion and lower motor neuron lesion

d) Bladder involvement

Question 8:

Mark the correct statement regarding inflammatory bowel disease.

a) Skip lesions are present in Crohn's disease

b) Mucosal layers are involved in Crohn's while transmural involvement seen in ulcerative colitis

c) Inflammatory bowel disease has a strong genetic predisposition

d) Crohn's is curable through surgical resection of the affected segment

Question 9:

A 40-year-old man presents with 15 days of fever and altered sensorium for 1 day. Rapid diagnostic test was done and Plasmodium falciparum was diagnosed. Which of the following is not a complication?

a) Blood glucose <40 mg/dL

b) Arterial pH >7.2

c) Serum creatinine - 5.2

d) Unarousable coma

Question 10:

A person presents to the hospital with fever and chills. Fever profile is ordered and is found to be negative for malaria and dengue. rk39 test is found to be positive. What is the treatment of choice?

a) Amphotericin B

b) Griseofulvin

c) Dapsone

d) Hydroxychloroquine

Question 11:

Incorrect statement regarding the management of frostbite:

a) Amputation in severe cases

b) Rewarming is done

c) Antibiotics and analgesics not used

d) The area is dried and cleaned

Question 12:

Which of the following is not a cause of central cyanosis?

a) Methemoglobinemia

b) Pulmonary arteriovenous fistula

c) High altitude

d) Hypothermia

Question 13:

A female patient presents to the hospital with palpitations. Her vitals are stable and the ECG shows the following. Which of the following is used as first-line management to treat the condition?



a) Amiodarone

b) Adenosine

c) DC cardioversion

d) Primary PCI

Question 14:

Which of the following is not a feature of Brown Sequard syndrome?

a) Complete transection of spinal cord

b) Ipsilateral loss of vibration & touch

c) Contralateral loss of pain & temperature

d) Ipsilateral loss of proprioception

Question 15:

Which of the following drug is not used for the emergency (immediate) management of hyperkalemia?

a) 10% calcium gluconate over 10 min

b) Salbutamol nebulisation

c) Insulin-dextrose

d) Injection $MgSO_4$

Question 16:

Chronic HBV infection would be consistent with which of the following serology?

a) 1, 3, 4

b) 1, 3

c) 1, 2

d) 1, 2, 4

Question 17:

Tongue fasciculations are seen in:

a) Guillain-Barre syndrome

b) Spinal muscular atrophy

c) Duchenne muscular dystrophy

d) Myasthenia gravis

Question 18:

In a patient with HIV, the incorrect statement regarding prophylaxis for various organisms is ____.

a) Prophylaxis for PCP is indicated when CD count is <200 cells/µL

b) Stop prophylaxis for Coccidioides if CD count is >250 cells/µL for 6 months

c) Prophylaxis for MAC is indicated when CD count is <50 cells/µL

d) Prophylaxis for Cryptococcus is indicated when CD count is >150 cells/µL

Question 19:

A patient presents with fever and nuchal rigidity. The resident was instructed to perform a lumbar puncture on this patient. What is the correct order of managing this patient?

a) 1,2,4,3

b) 1,4,3,2

c) 4,3,2,1

d) 2,4,3,1

Question 20:

A female patient was clinically diagnosed to have hypothyroidism. Measurements of the following parameters were obtained: TSH; combined protein levels; and total and free T3, T4, and rT3. Which one of the following options represents her condition?

a) 1

b) 2

c) 3

d) 4

Question 21:

A patient, who is a known case of HIV with a CD4 count of 200 cells/cu.mm, presents with 5 days of cough and high-grade fever without chills and rigors. There is no history of diarrhea, vomiting, or nuchal rigidity. Chest x-ray is normal. What treatment will you give?

a) Amoxicillin-clavulanic acid + Azithromycin

b) Co-trimoxazole + steroids

c) Co-trimoxazole only

d) Antitubercular treatment

Question 22:

The following clinical examination was performed. What lesion would cause an exaggerated reflex?



a) Polyneuropathy

b) Radiculopathy

c) Upper motor neuron

d) Lower motor neuron

Question 23:

A 65-year-old female patient weighing 60 kg is on mechanical ventilation for ARDS secondary to urosepsis. The respiratory parameters are as follows: tidal volume- 360 mL; frequency- 30 breaths/min; PEEP- 5 cm of H_2O; and FiO_2- 90%. The arterial blood gas findings are as follows: paO_2- 50 mmHg; $paCO_2$- 38 mmHg; and pH- 7.38. What is the next step?

a) Reduce FiO_2

b) Increase tidal volume

c) Increase respiratory rate

d) Increase PEEP

Question 24:

Identify the conduction abnormality from the ECG given below.

a) First degree heart block

b) Ventricular tachycardia

c) Third degree heart block

d) Second degree heart block

Question 25:

An elderly male patient was brought by his son because he was not using his left arm for day-to-day functioning. He has an unkempt appearance on the left side of his body and has a history of bumping into door frames while walking through them. The evaluation performed is shown below. Where does a lesion in the brain cause this presentation?

©MARROW

a) Right parietotemporal

b) Left parietal

c) Right parietal

d) Right premotor cortex

Question 26:

A young man is brought to the hospital with high-grade fever and altered consciousness. On examination, he had neck rigidity and pain when bending the neck. A lumbar puncture was performed, which showed a WBC count of 45 cells/µL, majorly being lymphocytes, elevated pressure with protein of 120 mg/dL, and glucose level of 70 mg/dL. How will you manage the index case?

a) Piperacillin + tazobactam

b) Vancomycin + ceftriaxone

c) Amphotericin B + flucytosine

d) Antitubercular therapy

Question 27:

The absence of loud S1 in mitral stenosis would indicate all of the following, except _____.

a) Calcified valve

b) Aortic regurgitation

c) First degree heart block

d) Mild mitral stenosis

Question 28:

A patient presents with perioral numbness, tingling, and tetany. All the following conditions can lead to this, except _____.

a) Vitamin D toxicity

b) Acute pancreatitis

c) Chronic kidney disease

d) Hypoparathyroidism

Question 29:

Thrombolysis can be considered in all of these conditions, except _____.

a) MRI showing density in less than 1/3rd of the area supplied by MCA

b) Blood pressure of more than 185/110 mmHg

c) Ischemic stroke within 2 hours

d) Onset of symptoms <4 hours

Question 30:

Serologic findings of a patient with hepatitis are given below.

a) High HDV viral load

b) High HBV viral load

c) Presence of IgM anti HDV antibodies

d) Absence of anti HBc IgG antibodies

Question 31:

A patient was brought following an RTA. He was found to have an incomplete lesion of the spinal cord. What would be seen if the central part of the cord was involved?

a) 1 and 4

b) 2 and 3

c) 2 and 4

d) 1 and 3

Question 32:

A patient with heart disease has breathlessness on going to the bathroom. What grade does he belong to?

a) NYHA 3

b) NYHA 4

c) MMRC 4

d) MMRC 5

Question 33:

A 50-year-old male patient presented with left-sided hemiparesis. Damage to which part of the internal capsule leads to this presentation?

a) Retrolentiform

b) Sublentiform

c) Anterior limb

d) Posterior limb

Question 34:

An increased anion gap is seen in _____.

a) Respiratory acidosis

b) Respiratory alkalosis

c) Metabolic acidosis

d) Metabolic alkalosis

Question 35:

A young patient presents to the clinic with erythematous lesions over the exposed areas of the skin like hands, arms, chest, etc. She also complains of arthralgia and breathlessness. Which among the following antibodies will be useful in diagnosing this condition?

a) Antihistone antibodies

b) Anti dsDNA antibodies

c) Anticentromere antibodies

d) Antinuclear antibodies

Question 36:

Based on the ECG given below, which of the following drugs is not used in the management of the condition?



a) Metoprolol

b) Amiodarone

c) Adenosine

d) Diltiazem

Question 37:

A patient presented with violent, flinging movements. Where is the lesion causing the hemiballismus seen?

a) Putamen

b) Subthalamic nucleus

c) Caudate nucleus

d) Globus pallidus

Question 38:

An elderly female patient with a history of severe vomiting was treated aggressively for severe dehydration at an outside hospital. One week later, she became mute, quadriplegic, and rigid. What is the most likely reason?

a) Malignant hyperthermia

b) Severe catatonia

c) Rapid sodium correction

d) Neuroleptic malignant syndrome

Question 39:

Which of the following is seen in MEN 2B syndrome?

a) Cafe au lait spot

b) Medullary thyroid carcinoma

c) Parathyroid adenoma

d) Optic nerve glioma

Question 40:

A patient presents with severe abdominal pain. Identify the false statement given about the examination of the abdomen.

a) Hemoperitoneum leads to reddish discoloration of the flanks of the abdomen

b) Cullen's sign refers to discoloration around the umbilicus

c) Palpation begins at the site of pain

d) Guarding and rigidity indicate peritoneal inflammation

Question 41:

A patient with diabetes develops a UTI, which gets complicated with hypotension that is resistant to IV fluids. Which of the following antibiotics can be used?

a) Piperacillin-Tazobactam

b) Amoxicillin-Clavulanate

c) Ceftriaxone

d) Nitrofurantoin

Question 42:

A patient with diabetes, hypertension, and chronic kidney disease has elevated serum creatinine and urea levels. Which of the following oral hypoglycemic agents is safe to use?

a) Glimepiride

b) Exenatide

c) Vildagliptin

d) Linagliptin

Question 43:

A hypertensive patient who is on losartan develops a HBA1c level of 8.1%. Which of the following drugs would you replace losartan with?

a) Amlodipine

b) Telmisartan

c) Metoprolol

d) Olmesartan

Question 44:

Which of the following is true regarding the placement of ECG leads?

a) The earth to all leads is in the right leg.

b) Lead I: Positive pole to the right and negative pole towards the left

c) Lead III: Positive pole on the left and negative pole towards the right

d) Lead II: Positive pole on the right and negative pole towards the left

Question 45:

What is depicted in the following image?



a) Babinski sign

b) Withdrawl reflex

c) Grasp reflex

d) Plantar reflex

Question 46:

What is the most common site of abdominal tuberculosis?

a) Rectum

b) Colon

c) Small intestine

d) Ileocecal junction

Question 47:

An 11-year-old child diagnosed with acute rheumatic fever and having no valvular pathology comes to your clinic. What is the duration of prophylaxis given to this child?

a) 5 years

b) 40 years

c) 10 years

d) 20 years

Question 48:

Which of the following electrolyte abnormalities is seen in methanol intoxication?

a) Metabolic alkalosis with high anion gap

b) Metabolic acidosis with high anion gap

c) Metabolic acidosis with normal anion gap

d) Metabolic alkalosis with normal anion gap

Question 49:

A woman presents with altered sensorium, breathlessness, hypotension, and bradycardia. Examination revealed non-pitting edema of the extremities. She has a long-standing history of weight gain, constipation, cold intolerance, and menorrhagia. What is the most likely diagnosis?

a) Septic shock

b) Cardiogenic shock

c) Myxedema coma

d) Hyperthyroidism

Question 50:

Which of the following types of headaches requires further evaluation?

a) 2, 3

b) 2, 4

c) 2, 3, 4

d) 1, 2, 3, 4

Question 51:

A patient presents with cough, cold, fever, and malaise associated with weight loss and ascites. A radiograph of the chest is shown below. Ascitic fluid analysis showed elevated adenosine deaminase. Which of the following is the next step in management?

a) Biopsy

b) Start ATT after laparotomy and stricture removal

c) Start anti-tubercular therapy

d) Conservative management

Question 52:

A female patient presents to the AIIMS emergency department with severe fatigue and prostration. She is a known case of chronic stable angina. While collecting a blood sample, you notice the blood has a brownish hue. What is the underlying diagnosis?

a) Carboxy-hemoglobinemia

b) Sideroblastic anemia

c) Methemoglobinemia

d) Sulfhemoglobinemia

Question 53:

A patient presented with ipsilateral Horner's syndrome, ipsilateral loss of pain and temperature sensations in the face, vertigo with numbness and loss of sweating and dysarthria on the contralateral side. All these symptoms are caused due to a lesion in:

a) B and D

b) B, C, D

c) A, B, C

d) A and B

Question 54:

A female presented with loss of pain and temperature, but her touch sensation was intact. Imaging showed cavitation around the central canal. What is the probable diagnosis?

a) Brown-Sequard syndrome

b) Tabes dorsalis

c) Syringobulbia

d) Syringomyelia

Question 55:

A patient was treated for peptic ulcer with an H.pylori regimen. Which of the following is used to assess the success of treatment?

a) Endoscopy

b) IgG antibody study

c) Urea breath test

d) Urease test of biopsy

Question 56:

The patient has a fever, cough, and cold. The chest x-ray is given below. What will be the examination finding in the infrascapular region?



a) Hyperresonance on percussion

b) Coarse crackles

c) Dull note on percussion

d) Stridor

Question 57:

What does the given ECG show?



a) P-pulmonale

b) Ventricular bigeminy

c) Electrical alternans

d) Improper calibration

Question 58:

Hamman sign is seen in:

a) Pneumoperitoneum

b) Pneumopericardium

c) Pneumomediastinum

d) Hydropneumothorax

Question 59:

Serologic findings of a patient with hepatitis are given below.

a) HDV RNA copies

b) Anti-HBc IgM positivity

c) Anti-HBs IgM positivity

d) Anti-HBc IgG positivity

Question 60:

Match the following:

a) 1-D,2-C,3-A,4-B

b) 1-C,2-D,3-A,4-B

c) 1-A,2-B,3-D,4-C

d) 1-B,2-A,3-C,4-D

Question 61:

The given image shows a normal graph on the left and the patient's graph on the right. Which of the following diagnoses can be inferred from the graph?



a) Chest wall neuromuscular disease

b) Sarcoidosis

c) Idiopathic pulmonary fibrosis

d) Bronchiectasis

Question 62:

Which of the following are signs of heart failure?

a) 1 and 2

b) 2, 3, 4

c) 1, 2, 3

d) 1, 3, 4

Answer Key

Question No.	Correct Option
1	c
2	a
3	d
4	d
5	b

6	d
7	a
8	a
9	b
10	a
11	c
12	d
13	b
14	a
15	d
16	a
17	b
18	d
19	c
20	d
21	c
22	c
23	d
24	d
25	c
26	c
27	d
28	a
29	b
30	d
31	d
32	a
33	d
34	c
35	b
36	c
37	b
38	c
39	b
40	c
41	a
42	d

43	b
44	a
45	a
46	d
47	c
48	b
49	c
50	c
51	c
52	c
53	a
54	d
55	c
56	c
57	a
58	c
59	d
60	a
61	d
62	b

Detailed Explanations

Solution to Question 1:

The advice that can be given to the patient regarding non-pharmacological management is that <30% of the calories should come from fat.

Medical nutrition therapy (MNT) refers to the optimal coordination of caloric intake along with other aspects of diabetes therapy (insulin, exercise, and weight loss). According to the Indian Research Society for Diabetes, these are the non-pharmacological recommendations:

• Carbohydrate content should be limited to 50%-60% of total calorie intake. Complex carbohydrates and low glycemic index foods should be preferred. Fiber intake should be around 25-40g per day.

• Protein intake should be 15% of the total calorie intake. Intake of red meat should be avoided.

• Fat intake should be limited to less than 30% of the total calorie intake. Oils with high monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA) should be used. Saturated fatty acid intake should be less than 10% of the total calories per day. Non vegetarians should consume 100■200 g of fish/week as these are rich sources of PUFA. Vegetarians should

consume vegetable oils (soybean/ safflower/sunflower), walnuts, and flaxseeds.

• Patients should consume a diet rich in fruits, leafy vegetables, nuts, fibers, whole grains, unsaturated fat, legumes, unprocessed vegetables, and low-fat dairy.

• Salt consumption should be less than 5g per day (not sodium). Sodium consumption should be less than 2300 mg per day.

• Patients should be advised to avoid smoking, alcohol intake, sugars, and sweetened beverages.

A minimum of 150 min/week of physical activity is recommended for healthy Indians to prevent type 2 diabetes mellitus and coronary artery disease. This should involve:

• ≥30 min of moderate■intensity aerobic activity each day

• 15■30 min of work■related activity

• 15 min of muscle■strengthening exercises (at least 3 times/week)

Solution to Question 2:

Anosmia is the most characteristic feature of Kallman's syndrome.

Kallman's syndrome is a hypothalamic disorder that occurs due to defective gonadotropin-releasing hormone (GnRH).

It is mainly associated with anosmia or hyposmia or hypoplasia. Other associated features such as colour blindness, optic atrophy, deafness, cleft palate, renal abnormalities, and cryptorchidism may be present.

The presenting features in males are delayed puberty and micropenis while females present with primary amenorrhoea and failure of secondary sexual development.

The condition is characterised by low luteinizing hormone (LH), follicle-stimulating hormone (FSH), testosterone and estradiol.

Treatment includes human chorionic gonadotropin (hCG) or testosterone in males for restoration of puberty and secondary sexual characteristics. In females, cyclic estrogen and progestin can be given. Fertility is restored by giving gonadotropins or pulsatile GnRH

Solution to Question 3:

In the given clinical scenario, FEV1/FVC is less than 0.7, hence it is obstructive. After bronchodilation, FEV1 has increased from 0.9L to 1.9L, which is more than 12% implying its reversible. Therefore, the most likely diagnosis is an obstructive disease with bronchodilator reversibility.

In obstructive pulmonary disease, FEV1/FVC is reduced to <0.7 and FEV1 is also reduced <80%. Reversibility is defined as a ≥12% increase and absolute increase of 200mL in the FEV1 at least 15 min after administration of a β2 -agonist. Examples of obstructive pulmonary disease include COPD, asthma, bronchiectasis, etc.

In restrictive pulmonary disease, FEV1/FVC is normal or increased (>0.7) and FEV1 and FVC are also reduced. Examples of restrictive pulmonary disease include asbestosis, sarcoidosis, idiopathic pulmonary fibrosis, etc.

Solution to Question 4:

Azathioprine is not a first-line drug used in the management of rheumatoid arthritis (RA). It is a purine synthesis inhibitor and has been used for the management of RA earlier, but it is not preferred now due to inconsistent efficacy, myelosuppression, high hepatic toxicity, and renal damage.

The drugs used for the treatment of RA can be classified into broad categories:

• NSAIDs for pain relief.

• Glucocorticoids are given before initiating DMARD (disease-modifying anti-rheumatoid drug) therapy.

• Biological agents such as TNF-α inhibitors, abatacept, anakinra, rituximab and interleukin -6 inhibitors are useful in RA.

• DMARDs

DMARDs as follows:

• Methotrexate is the first choice, supplemented with folic acid to prevent side effects.

• Hydroxychloroquine (HCQ) is given as an adjuvant in combination with other DMARDs.

• Sulfasalazine helps to reduce the radiographic progression of the disease.

• Leflunomide

Rheumatoid arthritis (RA) is a chronic inflammatory disease, presenting with early morning joint pain and stiffness that relieves with activity, mainly involving the metacarpophalangeal (MCP) and proximal interphalangeal (PIP) joints. Other hallmark features include flexor tendon tenosynovitis, swan neck deformity, boutonniere deformity, and Z-line deformity.

Extra Articular features like subcutaneous nodules (most common), secondary Sjogren's syndrome, interstitial lung disease, pulmonary nodules, and anemia may be present.

Patients may have elevated ESR (Erythrocyte sedimentation rate) or C reactive protein (CRP). Rheumatoid factor and anti-CCP (cyclic citrullinated peptide) antibodies are helpful in differentiating RA from other polyarticular conditions and also have prognostic significance.

Solution to Question 5:

Melanoma is not an AIDS-defining malignancy.

AIDS-defining malignancies are those cancers that a human immunodeficiency virus (HIV) infected person is at a high risk of developing. When a HIV-infected person develops one of the AIDS-defining malignancies it suggests that he has AIDS.

AIDS-defining malignancies include the following:

• Kaposi sarcoma - It is a multicentric neoplasm of vascular origin and has a strong association with HHV - 8.

• Non-Hodgkin's lymphoma

• Invasive cervical cancer

Given below is the image of kaposi sarcoma:

Note : The FDA has approved pomalidomide, a thalidomide analogue to be included in treating adult patients with AIDS-related Kaposi's sarcoma after failure of highly active antiretroviral therapy and Kaposi's sarcoma in adult patients who are HIV-negative.

Solution to Question 6:

In the given scenario the patient has alcohol withdrawal symptoms along with deranged liver function test findings. So the benzodiazepine that can be safely administered to this patient is lorazepam.

Lorazepam has a short half-life and it has to be considered in patients with liver impairment or brain damage. Lorazepam, Oxazepam, and Temazepam are metabolized by conjugation and are less dependent on liver function. The rest of the benzodiazepines require hepatic CYP-mediated oxidation to metabolize and are hence not recommended for patients with elevated transaminases.

Alcohol withdrawal syndrome is characterized by withdrawal symptoms in an alcoholic patient who suddenly stops drinking. Features include tremors of the hands, agitation, tachycardia, tachypnoea, sweating, and insomnia. These symptoms begin after 5-10 hours of abstinence and peak on day 2 or 3 and improve by the fourth or fifth day. If these features are associated with delirium which includes confusion, agitation, and fluctuating consciousness, then it is known as delirium tremens.

Management is mainly by benzodiazepines such as chlordiazepoxide, lorazepam, and diazepam.

Solution to Question 7:

Contralateral hemiplegia does not help in the localization of the lesion in the spinal cord.

Contralateral hemiplegia is a feature of involvement of the corticospinal tract from its origin in the cerebrum (motor homunculus) till it crosses at the level of the lower medulla. At the level of spinal cord lesion, it leads to ipsilateral hemiplegia (eg: Brown Sequard syndrome)

Fasciculation at the level of lesion, upper motor neuron lesions (UMNL) and lower motor neuron lesions (LMNL), and bladder involvement are features of spinal cord lesions.

The presence of fasciculation (option B) indicates segmental signs corresponding to the disturbed individual cord segment. Other features include altered sensation, muscle atrophy, or absent deep tendon reflex.

Upper motor neuron (UMN) lesions and lower motor neuron (LMN) lesions (option C) can help localize lesions in the spinal cord. UMN signs include paraplegia or quadriplegia with exaggerated deep tendon reflexes, positive Babinski sign, and spasticity. LMN lesions include muscle atrophy, fasciculations, decreased reflexes, and flaccid paralysis.

Autonomic disturbances such as absent sweating below the implicated cord level and bladder involvement (option D) are seen when there is transverse damage to the spinal cord.

Solution to Question 8:

The correct statement regarding inflammatory bowel disease (IBD) is that skip lesions are present in Crohn's disease.

Inflammatory bowel disease (IBD) is a chronic inflammatory condition of the gastrointestinal tract and is mainly idiopathic. It includes ulcerative colitis (UC) and Crohn's disease (CD). Genetic predisposition has been identified for IBD. But its association is weak for both CD and UC (option C)

Ulcerative colitis mainly involves the mucosa of the rectum (Option B) and extends proximally to the colon. Microscopically, crypt architecture is distorted. Neutrophils invade the epithelium and the crypts leading to cryptitis and crypt abscess. Clinically, UC presents with diarrhea, rectal bleeding, tenesmus, and abdominal cramps.

Crohn's disease can affect any part of the gastrointestinal tract from the mouth to the anus but mostly involves the terminal ileum. It is characterized by transmural inflammation. The characteristic lesions include cobblestone appearance, skip lesions, and transmural inflammation (Option B). CD presents as ileocolitis characterized by right lower quadrant pain, palpable mass, fever, and leukocytosis. Bowel obstruction can also be seen. They also present as jejunocolitis characterized by malabsorption and steatorrhea.

Management can be:

• Medical management with sulfasalazine, glucocorticoids, and immunosuppressants like azathioprine and mercaptopurine are useful

• Surgical management is indicated in ulcerative colitis. It does not cure Crohn's disease (Option D)

Solution to Question 9:

The clinical scenario of a patient presenting with prolonged fever, altered sensorium, and test showing Plasmodium falciparum positive is suggestive of severe falciparum malaria, Arterial pH >7.2 is not a complication of falciparum malaria.

Malaria is a protozoal disease transmitted by the bite of a female anopheles mosquito that has been infected. Five different species of Plasmodium are responsible for disease in humans. They are P. falciparum, P.vivax, P.ovale, P.malariae, and P.knowlesi.

Features of severe falciparum malaria are as follows:

• Unarousable coma or cerebral malaria

• Acidosis is characterized by pH <7.25, base deficit >8 meq/L or plasma bicarbonate <15 mmol/L, venous lactate >5 mmol/L, and labored deep breathing.

• Anemia is severe with a normocytic and normochromic picture with hemoglobin <5g/dL or hematocrit <15%

• Renal failure is characterized by serum creatinine >3 mg/dL, urine output <400 mL, and there is no improvement with rehydration.

• Pulmonary edema or adult respiratory distress syndrome

• Hypoglycemia with plasma glucose <40 mg/dL

• Hypotension or shock is characterized by systolic blood pressure < 50 mmHg in children and < 80 mm Hg in adults, and a prolonged capillary refilling time of >2s

• Bleeding or disseminated intravascular coagulation

• Convulsions, weakness

• Hemoglobinuria

• Jaundice

Solution to Question 10:

The given scenario of a person with fever with chills and a positive rk39 test is suggestive of kala-azar. The treatment of choice of this condition is amphotericin B.

Kala-azar or visceral leishmaniasis is a protozoal infection mainly affecting the reticuloendothelial system. It is caused by the Leishmania donovani complex that includes L. donovani and L. infantum. These organisms are transmitted by sandflies.

Clinical features include:

• Moderate to high-grade fever with chills and rigors

• Splenomegaly and hepatomegaly

• Anemia

• Lymphadenopathy

• Pedal edema, ascites - due to hypoalbuminemia

• Epistaxis, gastrointestinal bleed, and retinal hemorrhages - due to thrombocytopenia

The gold standard test for the diagnosis is the demonstration of amastigotes in the smears of tissue aspirates. Another commonly used test in the detection of antibodies to a recombinant antigen called rK39 by rapid immunochromatographic test.

Amphotericin B is the first-line drug used in India. Other drugs useful in treating the condition are sodium stibogluconate, meglumine antimoniate, paromomycin, and miltefosine.

Solution to Question 11:

The incorrect statement regarding management of frost bite is statement 3, since antibiotics and analgesics are useful for a patient with frost bite.

In frostbite, tissue damage results from severe environmental cold exposure or from direct contact with a very cold object. Tissue injury results from both freezing and vascular stasis. It usually affects the distal aspects of the extremities or exposed parts of the face, such as the ears, nose, chin, and cheeks.

Clinical features include decreased sensation (especially for pain, and temperature sensations) in the acral tips and extremities. Affected area appears waxy, mottled, or violaceous. Patients may have a wood-like sensation in the affected area.

Treatment:

• Initial treatment is by rapid rewarming (or thawing), which is accomplished by immersion of the affected part in a water bath at temperatures of 37°–40°C (99°–104°F) for 30–60 min. Rewarming by exercise, rubbing, or friction is not advised.

• Analgesics such as ibuprofen are often required during rewarming.

• Antithrombotics, thrombolytics and vasodilators such as prostacyclin can be given to improve circulation after rewarming.

• Frost bites are associated with an increased risk of infections and tetanus. Therefore, the injured area should be cleaned with soap or antiseptic, and sterile dressings should be applied. Tetanus prophylaxis must be given, but, empirical antibiotics are not recommended.

• The patient must be monitored for the development of compartment syndrome.

• Elective late amputation after imaging may be required in severe cases.

Image showing frost bite of the toes

Solution to Question 12:

Causes of central cyanosis are high altitude, methemoglobinemia, pulmonary arteriovenous fistula, certain congenital heart diseases etc.

Cyanosis appears when the reduced haemoglobin concentration of the blood in the capillaries is more than 5 g/dL. It depends on the total amount of haemoglobin in the blood and the degree of haemoglobin unsaturation. The state of the capillary circulation is also a contributing factor. Cyanosis is examined in the nail beds and mucous membranes, earlobes, lips, and fingers.

Cyanosis is divided into central and peripheral types. If the oxygen saturation is reduced along with an abnormal haemoglobin derivative it is called central cyanosis. Both the mucous membranes and skin will be affected. Peripheral cyanosis occurs as a result of slow blood flow and increased extraction of O2 from normally saturated arterial blood.The mucous membrane of the oral cavity and sublingual mucosa are spared. The causes are reduced cardiac output, cold exposure, arterial and venous obstruction etc.

Solution to Question 13:

The given clinical scenario and the ECG findings of narrow QRS complexes without P waves and pseudo R waves in aVR lead suggest paroxysmal supraventricular tachycardia (PSVT) and the first line of management to treat the condition is adenosine.

PSVT is a narrow QRS complex tachycardia with sudden onset and termination. They mainly present with dizziness and palpitations. Evaluation of these patients includes monitoring of vitals and ECG.

Management is as follows:

1) When the patient is hemodynamically stable:

• A carotid sinus massage can be done. It is avoided if carotid bruit is present and if a prior history of stroke is seen.

• Valsalva maneuver is done if the patient is cooperative

• Intravenous adenosine is useful. It is avoided in patients with atrial fibrillation.

• Intravenous calcium channel blockers and beta-blockers are helpful. It can cause hypotension.

2) When the patient is hemodynamically unstable, direct current cardioversion (DC cardioversion) should be done.

The below flowchart shows the treatment algorithm for PSVT.



Solution to Question 14:

Complete transection of the spinal cord is not a feature of Brown Sequard syndrome.

Brown sequard syndrome occurs due to transection of the spinal cord only on one side. This results in blocking of all motor functions on the side of the transection in all segments below the transection level.

Features include:

• Ipsilateral weakness - due to corticospinal tract

• Ipsilateral loss of joint position and vibration sensation - due to posterior column

• Contralateral loss of pain and temperature sensation - due to the spinothalamic tract involvement

• Unilateral radicular pain

• Muscle atrophy or loss of deep tendon reflex may be present

Solution to Question 15:

Intravenous magnesium sulfate (MgSO4) is not used in the management of hyperkalemia.

Hyperkalemia is an increase in the levels of serum potassium >5.5 mM. It is a medical emergency and has to be treated when ECG changes are present or severe hyperkalemia (serum K >6.5) without ECG changes.

Features include cardiac arrhythmias which include sinus bradycardia, sinus arrest, ventricular tachycardia, ventricular fibrillation and asystole.

ECG shows tall and peaked T waves, widening of QRS complexes, loss of P waves and sine wave pattern.



Management is as follows

• Intravenous infusion of 10 ml of 10% calcium gluconate for 2-3 minutes with cardiac monitoring. It helps to antagonise the cardiac effects.

• Intravenous regular insulin 10 units followed by 50mL of 50 % dextrose. This helps in shifting the plasma K+ into the cells

• Nebulisation with β2 agonists such as albuterol or salbutamol are effective.

• Cation exchange resins, diuretics and or dialysis are useful to remove the potassium.

Solution to Question 16:

Chronic HBV infection would be consistent with HBs Ag, Anti HBc Ag, and HBV DNA.

HBsAg (Hepatitis B surface antigen) is the first marker to appear in the serum and it indicates the presence of infection. If it is persistent for more than 6 months, it is suggestive of chronic infection.

Anti HBs are the antibodies to HBs and are seen after clearance of HBsAg or post-vaccination. Thus, it is suggestive of recovery, non-infectivity and immunity.

Anti HBc (antibodies to hepatitis B core antigen) has IgM and IgG. IgM anti-HBc is diagnostic of acute hepatitis and it appears soon after HBs Ag. It persists for 3-6 months and can re-appear in flare-ups. IgG anti-HBc appears during acute infection and persists indefinitely irrespective of the recovery or progression to chronic disease.

HBV DNA is a sensitive indicator for viral replication and infectivity. HBeAg also indicates viral replication and infectivity. It appears shortly after HBs Ag and if it is persisting beyond 3 months, it is suggestive of chronic hepatitis.

Hepatitis B virus is mainly transmitted by percutaneous route followed by perinatal and sexual transmission. Chronic hepatitis B is a progression of acute hepatitis. Chronic hepatitis B infection has five phases - immune tolerant phase, immune active phase, inactive HBsAg carrier state, reactivated chronic hepatitis B phase, and HBsAg negative phase.

Treatment options include interferon, lamivudine, adefovir, entecavir, telbivudine, and tenofovir.

Prophylaxis is by hepatitis B immunoglobulin (HBIG) and recombinant vaccine.



Solution to Question 17:

Tongue fasciculations are seen in spinal muscle atrophy (SMA).

Spinal muscle atrophy is a lower motor neuron disease with early-onset and occurs as a result of the loss of a protein called survival motor neuron (SMN).

Features of lower motor neuron disease include flaccid paralysis, muscle atrophy, fasciculations characterized by visible muscle twitches appearing as flickers under the skin, hypotonia (decreased muscle tone), hyporeflexia, or areflexia.

Clinical forms of SMA are

• In infantile SMA, the onset is very early and rapid. It is characterized by decreased fetal movements in the third trimester. In infancy, features include weak and floppy infant (hypotonic) and they lack muscle stretch reflex. Tongue fasciculations are typically seen in a severe form of SMA known as Werdnig-Hoffman disease. It can affect infants between 0-6 months.

• Chronic childhood SMA has a slow onset and a progressive course.

• Juvenile SMA is seen in late childhood with a slow course.

Nusinersen is approved for therapy and another treatment is systemic administration of adeno-associated virus to deliver the missing SMN gene.

Other options:

Option A: GBS or Guillain-Barre syndrome is an autoimmune polyradiculopathy with an acute onset. It is characterized mainly by motor paralysis with or without sensory disturbances. Weakness is seen more in the legs than arms. Other features include pain, bladder dysfunction, affected deep tendon reflexes, and proprioception. Hypotension and arrhythmias can also be seen

Option C: Duchenne muscular dystrophy is an x-linked recessive disorder due to mutation in the dystrophin gene. Typical features include proximal muscle weakness in lower extremities, calf muscle hypertrophy, intellectual disability, and heart failure. Myalgia without weakness and myoglobinuria can also be present.

Option D: Myasthenia gravis is a neuromuscular disorder due to autoimmune damage to the acetylcholine receptors. It is characterized by weakness and muscle fatiguability. Weakness increases with usage. Ocular features like ptosis and diplopia can be seen. Proximal limb weakness is also present.

Solution to Question 18:

The incorrect statement regarding prophylaxis in a patient with HIV is that prophylaxis for Cryptococcus is indicated when the CD count is >150 cells/µL.

In cryptococcal infection, the indication for prophylaxis is a prior documented disease. The drugs used in prophylaxis can be oral fluconazole 200 mg/day or itraconazole 200 mg/day.

Prophylaxis can be stopped when CD4+ T-cell count is >100/µL and when there is no evidence of infection.

Solution to Question 19:

The correct order of events is 4, 3, 2, 1.

The classic clinical symptoms of fever and neck rigidity are suggestive of meningitis. Other common complaints are nausea, vomiting, and photophobia. Nuchal rigidity (“stiff neck”) is the pathognomonic sign of meningitis.

Bacterial meningitis in adults must be considered a medical emergency and requires urgent administration of empirical antibiotic therapy. However, as part of pre-treatment evaluation, lumbar puncture and blood cultures must be sent for testing prior to initiation of antibiotics.

Therefore a guarded lumbar puncture must be performed as the first step.

Once the cerebrospinal fluid samples are sent for culture and sensitivity testing, ceftriaxone injection must be administered as the emergent next step as part of empirical antibiotic therapy. It is recommended that antibiotic therapy is initiated within 60 minutes of the patient's arrival to the emergency room.

IV cannulation must be performed to attain intravenous access to administer fluids. Studies have shown that the administration of maintenance fluids in the initial 48 hrs of admission has reduced the incidence of adverse sequelae. Since IV cannulation is mentioned to be done for administering fluids, it can be done after lumbar puncture and antibiotics. Also, caution must be taken before administering fluids because there might be altered intracranial pressures.

Solution to Question 20:

Laboratory investigations in a patient with hypothyroidism would show low levels of total and free T3, T4, and rT3, with high TSH levels and normal combined protein levels.

Normally, the thyrotropin-releasing hormone (TRH) is released by the hypothalamus. It stimulates the pituitary gland to release the thyroid-stimulating hormone (TSH), which in turn stimulates the thyroid gland to produce T3 and T4 hormones.

Elevated TSH with a low level of free T4 is suggestive of primary hypothyroidism.

In the case of secondary hypothyroidism, TSH levels may be low, normal, or even slightly increased. The latter is due to the secretion of immunoactive but bio-inactive forms of TSH. The diagnosis is confirmed by the detection of a low unbound T4 level.

Patients with hypothyroidism often present with tiredness, weakness, dry skin, hair loss, difficulty in concentrating, poor memory, constipation, and weight gain with poor appetite. On examination, they may have cool peripheral extremities, puffy faces, bradycardia, and peripheral edema.

Patients with hypothyroidism who have no evidence of heart disease are managed with levothyroxine 50 to 100 mcg daily. The goal of the treatment is to maintain TSH in the lower half of the reference range.

Solution to Question 21:

The clinical scenario of an HIV-infected patient with a CD4 count of 200/mm3 with fever, cough, and a normal chest x-ray is indicative of underlying Pneumocystis pneumonia. Pneumocystis jirovecii pneumonia (PCP) is an opportunistic infection in immunocompromised patients. It is treated with co-trimoxazole.

Patients usually present with fever and nonproductive cough. It can also affect other organs like lymph nodes, spleen, and liver.

The classic radiographic appearance is likely normal or could consist of faint bilateral interstitial infiltrates. Perihilar infiltrates are less common in patients with AIDS. CT will show diffuse ground-glass opacities.

Histopathology of the tissue will show foamy alveolar infiltrate and mononuclear interstitial infiltrates, which is pathognomonic of PCP. Specific stains like methenamine silver, toluidine blue O, or Giemsa can show the cysts. Other tests that are used for the diagnosis of PCP are polymerase chain reaction, (1→3)-β-d-glucan, etc.

The treatment of PCP is with cotrimoxazole (trimethoprim-sulfamethoxazole).

For patients with a PaO2 <70 mmHg or with an arterial-alveolar (a–A) gradient >35 mmHg, adjunct glucocorticoid therapy should be used in addition to cotrimoxazole. As the clinical stem does not indicate poor oxygenation, steroid therapy does not have to be initiated. (Option B)

Atovaquone, pentamidine, and primaquine + clindamycin are alternative agents for the treatment of PCP.

Other options

Option A - Amoxicillin with azithromycin is used for the empirical management of bacterial pneumonia on an outpatient basis. Bacterial pneumonia, however, tends to present with consolidation on chest x-ray, which is not seen in this patient.

Below is an image of bacterial pneumonia showing consolidation on x-ray:



Option D: Antitubercular therapy would be indicated if tuberculosis was suspected in the patient. Tuberculosis in HIV-infected patients can be seen in relatively high CD4 counts and typically presents with fever, cough, weight loss, night sweats, and an abnormal chest x-ray revealing upper lobe cavitary disease.

Solution to Question 22:

The examination being performed is the ankle jerk reflex, also known as the Achilles reflex. Upper motor neuron (UMN) lesions are associated with exaggerated deep tendon reflexes.

Hyperreflexia is a feature of UMN nerve injury. There may be an acute decrease in reflexes, but this is normally followed by hyperreflexia. LMN lesions on the other hand will present with

depressed reflexes.

UMNs refer to corticospinal tract neurons and brainstem neurons that control spinal motor neurons. Injury to UMNs leads to:

• Spasticity

• Hypertonia

• Hyperreflexia

• Abnormal plantar extensor reflex (positive Babinski sign).

Other options:

Options A, B and D: Polyneuropathy/ neuropathic weakness and radiculopathy are types of lower motor neuron weakness and will not cause exaggerated reflexes.

The following image shows the Babinski reflex:



Solution to Question 23:

Increasing PEEP in this patient would be the next step in ensuring adequate oxygenation. Applying adequate PEEP can improve arterial oxygenation by reducing ventilation-perfusion (V/Q) mismatch in areas of atelectasis.

This patient's low paO2 and low paO2/FiO2 ratio are suggestive of impaired oxygenation. The ratio of arterial oxygen partial pressure to the fraction of inspired oxygen (PaO2/FiO2) is ≤300 mmHg as assessed on 5 cm H2O of positive end-expiratory pressure (PEEP) or continuous positive airway pressure (CPAP). In this patient the ratio is ≤100 mmHg, implying that the PEEP of 5 cm H2O is insufficient and must be increased.

Acute respiratory distress syndrome (ARDS) is the most severe type of acute lung injury characterized by poor oxygenation, pulmonary infiltrates, and acuity of onset. It is associated with capillary endothelial injury and diffuse alveolar damage.

Etiology for ARDS:

• Direct lung injury- pneumonia, aspiration of gastric contents, pulmonary contusion, near-drowning, toxic inhalational injury

• Indirect lung injury- sepsis, severe trauma like multiple bone fractures, flail chest, head trauma, burns, multiple transfusions, drug overdose, pancreatitis, post-cardiopulmonary bypass

ARDS is characterized by the development of dyspnea and hypoxemia, which progressively worsens within 6 to 27 hours of the initial provoking event.

Recommended therapies in ARDS with clinical or supportive evidence:

• The low tidal volume ventilation (LTVV) strategy (4–6 mL/kg of ideal body weight) is recommended to maintain adequate ventilation and avoid ventilator-induced lung injury by barotrauma and volume trauma.

• High positive end-expiratory pressure (PEEP) or open lung mechanical ventilation

• Prone position

• Extracorporeal membrane oxygenation (ECMO)

Other options:

Option A: Reducing FiO_2 will further worsen the patient's arterial oxygen as PaO_2/FiO_2 is ≤100 mmHg.

Option B: The LTVV strategy recommends a goal of 4–6 mL/kg of ideal body weight. The tidal volume setting in this patient is already at a goal of 6 mL/kg and further increasing it without first recruiting atelectatic lung tissue, would increase the risk of volume trauma.

Option C: Increasing the respiratory rate is an acceptable measure in the event of acute respiratory acidosis. The results of arterial blood gas analysis in this patient do not reflect acidosis. Furthermore, since the respiratory rate is already set at 30 breaths/min, increasing it further can worsen the risk of barotrauma and hyperinflation.

Solution to Question 24:

The ECG shown above is the second-degree atrioventricular block type 1 - Wenckebach phenomenon.

Second Degree Heart Block: It occurs due to the intermittent failure of electrical impulse conduction from the atrium to the ventricle. It is characterized by a missed QRS. It is of two types:

• Mobitz type 1/ Wenckebach occurs due to the decremental conduction of electrical impulses in the AV node. It is characterized by a progressive PR elongation leading to an eventual drop in the QRS complex. This may be seen in young athletes.



• Mobitz type 2 occurs in the distal or infra-His conduction system and is often associated with intraventricular conduction delays. It is characterized by an intermittent failure of conduction of the p wave without any change in the preceding impulses. On ECG there is a fixed PR interval and missed QRS.



Solution to Question 25:

The presentation shown in the clinical stem and image is suggestive of hemispatial neglect. Hemispatial neglect is the inability to report, respond, or orient to stimuli in one-half of the space despite having no motor or sensory deficits. It is primarily associated with lesions in the non-dominant parietal lobe i.e. right parietal lobe.

Contralateral hemispatial neglect is due to the damage of cortical or subcortical components of the parieto-frontal network for spatial orientation.

The right hemisphere directs attention within the entire extrapersonal space - both the right and left side, while the left hemisphere directs attention mostly within the contralateral right hemispace.

Thus, left hemisphere lesions do not give rise to much neglect because of the intact right hemisphere. Severe neglect for the right hemispace is rare, even in left-handers with left hemisphere lesions.

Right hemisphere lesions, however, give rise to severe left hemispatial neglect because the unaffected left hemisphere does not contain ipsilateral attentional mechanisms and cannot compensate.

The features of parietal lobe lesions are as follows:

• Apraxia

• Acalculia

• Alexia

• Agnosia

• Neglect

• Inferior quadrantic homonymous hemianopia.

Two bedside tests can be used to evaluate neglect: simultaneous bilateral stimulation and visual target cancellation.

• In simultaneous bilateral stimulation, when the examiner provides unilateral or simultaneous bilateral stimulation in the visual, auditory, or tactile modalities, patients have no difficulty detecting unilateral stimuli on either side but perceive the bilaterally presented stimulus as coming only from one side. This phenomenon is known as extinction.

• In visual target cancellation, the patient fails to detect the targets on one-half of the field in target-detection tasks.

Solution to Question 26:

The clinical symptoms along with the cerebrospinal fluid analysis of elevated pressure and proteins, normal glucose levels, and lymphocyte predominance suggests a diagnosis of fungal meningitis. Cryptococcal meningitis, one of the most common causes of fungal meningitis is managed with amphotericin B + flucytosine.

In cryptococcal meningitis, there is lymphocytic predominance in the CSF analysis of white blood cells.

Cryptococcus neoformans is an opportunistic fungus that is found worldwide as a saprophyte in soil enriched with pigeon droppings. It may cause infection in immunocompetent individuals, resulting in a flu-like illness with complete recovery.

Infection in immunocompromised people like AIDS patients or patients with hematological malignancies results in meningoencephalitis.

Pathogenic varieties of Cryptococcus, including C. neoformans and Cryptococcus gattii, are laccase positive and have the ability to grow at 37°C. These factors are used to differentiate them from non-pathogenic cryptococci.



The primary pulmonary infection may be asymptomatic or may mimic an influenza-like respiratory infection, often resolving spontaneously. Cutaneous infection results in ulcers and granulomata.

In immunocompromised individuals, the yeasts may multiply and disseminate to other parts of the body, but preferentially to the CNS, causing cryptococcal meningoencephalitis, which presents with headache and nuchal rigidity.

Diagnosis is established by the microscopic examination of cerebrospinal fluid, which shows gram-positive round budding yeast cells on gram staining (option B). Also, India ink preparation reveals capsulated budding yeast cells.

The image given below shows Cryptococcus neoformans stained with India ink.



Solution to Question 27:

The early phases of mitral stenosis typically show a loud S1 (first heart sound).

The loud S1 is due to the greater distance covered by the leaflets during closure since elevated left atrial pressure has kept the leaflets relatively wide apart. The S1 becomes softer in the later stages of MS when the leaflets are rigid and get calcified (Option A)

The first heart sound (S1) includes mitral and tricuspid valve closure. The intensity of S1 is determined by the

• distance over which the leaflet of the valve must travel to return to its annular plane

• LV contractility

• PR interval

Causes of loud S1

• Early mitral stenosis

• Hyperadrenergic states (thyrotoxicosis, anxiety, anemia, pregnancy, exercise)

• Short PR interval

Soft/ low intensity of S1 is caused by states that cause the AV valves to close prior to ventricular systole or due to a reduction in rate of development of pressure in the ventricles. The causes of soft S1 are:

• First-degree heart block (Option C)

• Mitral regurgitation

• Aortic regurgitation (Option B)

• Attenuation due to increased tissue between the heart and chest wall (obesity, obstructive lung disease, pericardial effusion)

• Calcified Mitral stenosis

• Left ventricular contractile dysfunction

• Beta-receptor blockers

Solution to Question 28:

The above clinical scenario of a patient presenting with perioral paresthesias and tetany is suggestive of hypocalcemia. Vitamin D toxicity does not cause hypocalcemia but rather leads to hypercalcemia.

Vitamin D toxicity occurs due to excessive intake of vitamin D supplements, usually in amounts of >40,000–100,000 IU/day. The symptoms are mainly due to hypercalcemia. Laboratory investigations will show elevated serum vitamin-D levels (>100 ng/mL) and calcium.

In such patients, vitamin D supplements should be discontinued immediately, along with restricted exposure to the sun and avoiding calcium supplements. Since vitamin D is stored in the body for a long time, the symptoms may persist for some days even after stopping the intake.

Causes of hypocalcemia

• Pancreatitis (sequestration of the calcium via chelation with free fatty acids)

• Renal failure/chronic kidney disease (impaired 1,25(OH)2D production)

• Pancreatic and small bowel fistulas

• Hypoparathyroidism (PTH is the main defense against hypocalcemia)

• Abnormalities in magnesium levels

• Tumor lysis syndrome

Symptoms and signs of hypocalcemia may be circumoral or digital numbness, paresthesia, laryngeal spasms, and cardiac arrhythmias. Given below are two named signs of hypocalcemia.

• Chvostek sign: This is elicited by gently tapping over the facial nerve in front of the tragus of the ear. There is hyperexcitability of the facial muscles, manifesting as a contraction of the muscles.

• Trousseau sign: This is elicited by making the forearm ischemic by applying a sphygmomanometer cuff to the arm and keeping the pressure above systolic blood pressure (SBP) for 5 minutes. The hand goes into a painful carpopedal spasm and assumes the accoucheur's hand position (shown in the image below).

The below image shows Trousseau's sign:



Acute symptomatic hypocalcemia is an emergency and must be treated with:

• 10 mL of 10% calcium gluconate is given slowly through the intravenous route.

• Calcium and vitamin D replacement is given orally.

• Magnesium supplementation may also be required due to the synergistic actions of calcium and magnesium.

Calcitonin and bisphosphonates are used in cases of severe toxicity causing hypercalcemia. Some patients might require glucocorticoids, which lower plasma calcium levels by reducing calcium absorption from the intestines and increasing its urinary excretion.

Solution to Question 29:

Thrombolysis is contraindicated if blood pressure is greater than 185/110 mmHg. Blood pressure must be lowered before considering thrombolysis.

Thrombolysis refers to the dissolution of the intravascular clots to prevent ischemic damage and improve blood flow.

Intravenous thrombolysis with recombinant tissue plasminogen activator (rTPA), alteplase (dose 0.9 mg/kg) should be immediately administered to patients with ischemic stroke who arrive in the window period within 4.5 hours of symptom onset.

Solution to Question 30:

Anti-HBc IgG antibodies are markers of a chronic hepatitis B infection. Its presence along with Anti HDV antibodies suggests a superinfection of HDV on chronic HBV. Therefore the absence of Anti HBc IgG antibodies is an indication of an acute episode of hepatitis which is associated with HBV/HDV (hepatitis B/hepatitis D) coinfection.

HBV/HDV coinfection occurs when a person simultaneously becomes infected with both HBV and HDV. In acute hepatitis, simultaneous infection with HBV and HDV can lead to mild-to-severe hepatitis. The signs and symptoms typically appear 3–7 weeks after the initial infection and include fever, fatigue, loss of appetite, nausea, vomiting, dark urine, pale-coloured stools, jaundice (yellow eyes), and even fulminant hepatitis. However, recovery is usually complete. Co-infection usually ends with recovery and viral eradication (<5% chronicity).

HDV superinfection occurs when a person who is already chronically infected with HBV acquires HDV. The infection of HDV on chronic hepatitis B can accelerate the progression to severe disease in all ages with progression to cirrhosis almost a decade earlier than in HBV mono-infected persons. The outcome is usually persistent infection with 80% progressing to chronicity.

Diagnosis of HDV infection is by detecting high levels of anti-HDV immunoglobulin G (IgG) and immunoglobulin M (IgM). It is confirmed by the detection of HDV RNA in serum and their levels are not helpful in differentiating between acute and chronic infections. There is no specific treatment for HDV infection. Pegylated interferon alpha may be used (with limited efficacy).

Solution to Question 31:

In this patient with central cord syndrome, LMN lesion of upper limbs and trunk, UMN of lower limbs, bladder and bowel involvement will be seen.

Central cord syndrome occurs due to damage to the central region of the spinal cord. It results from selective damage to the grey matter nerve cells and the decussating spinothalamic tracts surrounding the central canal. These decussating fibers carry crude touch, pain, and temperature sensations.

It commonly occurs due to hyperextension of the cervical spine, causing the spinal cord to compress between the vertebral body and ligamentum flavum. Spinal trauma, syringomyelia, and intrinsic cord tumors are the other causes.

This condition affects the cervical cord producing arm weakness > leg weakness.

There is dissociated sensory loss, i.e, loss of crude touch, pain, and temperature senses with a preserved light touch, proprioception, and vibration sense over the shoulders, lower neck, and upper trunk (cape distribution). There is bilateral lower motor neuron paralysis and muscular atrophy in the segment of the lesion.

Eventually, expansion of the cavity may lead to compression of the long tracts. This results in spasticity and weakness with loss of pain, temperature, light touch, and pressure sensations occurring below the level of the lesion. There is a characteristic sacral sparing.

Bowel and bladder dysfunction and Horner's syndrome may also be seen in cases of central cord syndrome.

The following image shows incomplete lesions of the spinal cord:



Solution to Question 32:

This patient with heart disease with dyspnea on going to the bathroom, which is an activity of daily living/ ordinary activity, may be classified as NYHA class 3.

The New York Heart Association (NYHA) has outlined criteria that may be used to classify patients with heart disease based on their functional disability.

NYHA Classification of Heart Failure:

The Modified Medical Research Council (mMRC) Dyspnea Scale is used to classify the severity of symptoms in respiratory diseases, specifically COPD.

Modified MRC (mMRC) Dyspnea Scale:

• 0- Breathless only with strenuous exercise

• 1- Breathless when hurrying on a level or walking up a hill

• 2- Walks slower than peers or has to stop for breath when walking on level

• 3- Stops for breath after walking about 100m or for a few minutes

• 4- Breathless at rest

NYHA class	Description
I	No limitation of physical activityOrdinary physical activity does not cause symptoms
II	Slight limitation of physical activityComfortable at rest, but ordinary physical activity causes symptoms
III	Marked limitation of physical activityComfortable at rest, butless than ordinary physical activitycauses symptoms
IV	Unable to carry out any physical activity without symptoms or symptoms at rest

Solution to Question 33:

The posterior limb of the internal capsule is affected in hemiplegia/ hemiparesis due to ischaemic stroke.

Motor fibers (cortico-spinal tract) supplying the extremities are present in the posterior limb, hence a lesion in this region may cause hemiplegia.

The types of lacunar syndromes are as follows:

• Pure motor hemiparesis - infarct in the posterior limb of the internal capsule

• Pure sensory stroke - infarct in the thalamus

• Ataxic hemiparesis - infarct in the pons or internal capsule

• Dysarthria and clumsy hand syndrome - infarction in the pons or internal capsule.



Solution to Question 34:

An increased anion gap is seen in metabolic acidosis.

Normally, in the serum, the total number of cations is equal to the total number of anions. But, a few of the anions (lactate, phosphate, and sulfate) cannot be measured, and this is called an anion gap.

Anion gap (AG) = [Na+] – [Cl-+HCO3-]

The normal anion gap is 6-12 mEq/L.

When acid anions, such as acetoacetate and lactate, accumulate in extracellular fluid, the AG increases, causing a high-AG acidosis. Metabolic acidosis with an increased serum anion gap is seen if the anion gap is >12 mEq/L. It occurs in conditions such as methanol poisoning, uremia, diabetic ketoacidosis, paracetamol poisoning, lactic acidosis, salicylate poisoning, etc.

A high anion gap indicates metabolic acidosis. When metabolic acidosis and metabolic alkalosis coexist in the same patient, the pH may be in the normal range. In this circumstance, the presence of a high AG helps identify the presence of metabolic acidosis.

Solution to Question 35:

The clinical scenario of a patient with a photosensitive rash, arthralgias, and shortness of breath is suggestive of systemic lupus erythematosus (SLE). Anti-dsDNA antibodies are specific for the diagnosis of SLE.

SLE is an autoimmune disorder that results in damage to multiple organs and organ systems in the body most commonly seen in women of reproductive age.

Clinical and immunological manifestations of SLE:

• Skin

• acute, subacute cutaneous manifestations (photosensitivity, malar or maculopapular rash)

• chronic cutaneous manifestations (discoid lupus, panniculitis, lichen planus-like)

• Oral or nasal ulcers

• Non-scarring alopecia

• Synovitis involving ≥2 joints

• Serositis (pleurisy, pericarditis)

• Renal:

• Protein/Creatinine ratio >0.5

• RBC casts

• Neurologic: seizures, psychosis, myelitis, neuropathy, confusional state

• Hemolytic anemia

• Leukopenia

• Thrombocytopenia

• Immunological manifestations

• Antinuclear antibodies >reference value (positive in >98% patients)

• Anti-dsDNA >reference value (specific)

• Anti-Smith antibody (specific)

• Antiphospholipid antibody (lupus anticoagulant, false-positive rapid plasma reagin)

• Low serum complement

To diagnose SLE, at least 4 out of 11 of the above criteria must be positive. However, if a renal biopsy is suggestive of lupus nephritis and lupus autoantibodies are positive then four criteria are not required.

Antinuclear antibody detection is sensitive for the screening of SLE but not specific to diagnose it.

Solution to Question 36:

The given ECG findings of an irregularly irregular rhythm, absent p waves, and narrow QRS complexes are suggestive of atrial fibrillation(AF). Adenosine is not used. Intravenous adenosine administration has been found to provoke spontaneous AF in 12-16% of patients.

Atrial fibrillation is the most common sustained arrhythmia. There is rapid and irregular atrial activation which is transmitted through the atrioventricular node. The conducted ventricular rate is usually rapid and irregularly irregular. It can vary from 110-160 beats/minute. A pulse deficit is an important clinical finding associated with the condition. This arises due to preload reduction.

Symptoms include palpitations, fatigue, presyncope, and dyspnea. An ECG shows the absence of P waves.

In acute atrial fibrillation with hemodynamic instability, electrical or chemical cardioversion is the first step. The latter is done by intravenous administration of class III antiarrhythmic - ibutilide. In a stable patient, ventricular rate control is the primary aim. This is done by intravenous or oral administration of beta-blockers (metoprolol). Calcium channel blockers such as verapamil and diltiazem can also be used.

If the current episode duration is >48 hours, cardioversion can precipitate thromboembolic complications. In such cases, cardioversion should be carried out only after 3 weeks of anticoagulation. It has to be continued for 4 weeks after cardioversion. Another option is to evaluate the presence of a thrombus in the left atrial appendage. This can be done either by transesophageal echocardiography (TEE) or high-resolution CT.

If the duration is <48 hours and there are no high-risk factors, cardioversion can be done. The CHA2DS2-VASc scoring system is used to assess stroke risk.

Warfarin is recommended in patients with rheumatic mitral stenosis or mechanical heart valves who present with atrial fibrillation. In mitral stenosis, thrombi arise frequently from the dilated left atrium. Atrial fibrillation in the absence of mitral stenosis is known as nonvalvular atrial fibrillation. Oral factor Xa inhibitors such as apixaban, edoxaban, or rivaroxaban can be used in such cases.



Solution to Question 37:

The above clinical scenario is suggestive of hemiballismus. A lesion in the subthalamic nuclei is responsible for this condition.

A lesion in the subthalamic nucleus causes sudden, violent, involuntary, and flinging movements confined to one side of the body, commonly affecting the proximal muscles of the limbs, a condition called hemiballismus. The movements are seen contralateral to the side of the lesion, i.e., involuntary movements of the right arm and leg are seen in the lesion of the left subthalamic nucleus and vice versa.



Basal ganglia circuits have both direct and indirect pathways. The direct pathway stimulates the cortex to produce movement, whereas the indirect pathway inhibits it. The striatal activity in the direct pathway is mediated by D1 dopamine receptors and in the indirect pathway by D2 dopamine receptors.

The subthalamic nucleus is a part of the indirect pathway. Hence, a lesion at subthalamic nuclei leads to loss of inhibitory effect of the indirect pathway resulting in dopaminergic overactivity. This is responsible for features of hemiballismus.

This movement disorder is usually self-limiting and tends to resolve spontaneously after weeks or months. Dopamine antagonistic agents can be given to manage it as well.

Other options:

Options A and C: A lesion in the putamen and caudate nucleus can give rise to chorea. Chorea is defined as rapid, semi-purposeful, graceful, dance-like nonpatterned involuntary movements involving distal or proximal muscle groups.

Option D: A lesion in the globus pallidus can give rise to athetosis. This involves spontaneous and continuous writhing movements of the hand, arm, neck, or face.

Solution to Question 38:

Severe/ intractable vomiting is a common cause of hyponatremia. The development of quadriplegia, mutism, and rigidity after treatment is suggestive of rapid correction of hyponatremia, leading to central pontine myelinolysis (CPM). It is a complication associated with the treatment of hyponatremia and can present a few days after the initial correction, as seen in this patient.

Rapid correction of hyponatremia (at a rate of >8–10 mM in 24 hours or 18 mM in 48 hours) causes hypertonic stress, leading to apoptosis and cell death. Demyelination without inflammation occurs at the base of the pons with the relative sparing of axons and nerve cells.

CPM, better known now as osmotic demyelination syndrome, commonly involves the pons; but it may also cause extrapontine myelinolysis, affecting the cerebellum, putamen, thalamus, or lateral geniculate bodies. This can result in quadriparesis, dysphagia, dysarthria, diplopia, mutism, ataxia, parkinsonism, and catatonia.

MRI is useful in establishing the diagnosis and may also identify partial forms that present as confusion, dysarthria, and/or disturbances of conjugate gaze without paraplegia.

Prevention of CPM depends on therapeutic guidelines for the restoration of severe hyponatremia which should aim for gradual correction i.e. at a rate of ≤10 mmol/L (10 mEq/L) within 24 h and ≤20 mmol/L (20 mEq/L) within 48 h.

The image shows the MRI finding of central pontine myelinolysis with axial fat-saturated T2-weighted MRI showing a hyperintense signal in the central pons.



Option A: Malignant hyperthermia is a condition in which the patient develops sustained muscle contraction and hyperthermia. It occurs following the administration of succinylcholine and other inhaled anesthetics like halothane, isoflurane, sevoflurane, desflurane, etc.

Option B: Catatonia is a clinical syndrome characterized by striking behavioral abnormalities such as stupor, mutism, and posturing.

Option D: Neuroleptic malignant syndrome (NMS) is a life-threatening adverse effect of antipsychotics characterized by increased pulse and blood pressure; motor and behavioral symptoms such as muscular rigidity and dystonia, akinesia, and agitation; and autonomic symptoms like hyperthermia. Laboratory findings include an increased white blood cell count and increased levels of creatine phosphokinase, plasma myoglobin, and myoglobinuria.

Solution to Question 39:

Medullary thyroid carcinoma is seen in multiple endocrine neoplasia 2B (MEN 2B) syndrome.

Multiple endocrine neoplasia (MEN) describes conditions with tumors involving two or more endocrine glands. There are four types - MEN 1-4. They have an autosomal dominant pattern of

inheritance and can also be sporadic.

MEN type 2 comprises groups of disorders caused by gain-of-function mutations of the RET proto-oncogene at chromosomal locus 10q11.2. There are three variants:

• MEN 2A, also known as MEN 2 (most common type)

• MEN 2B, also known as MEN 3

• MTC only

MEN 2A is also known as Sipple's syndrome. It is characterized by medullary thyroid carcinoma (MTC), pheochromocytomas, and parathyroid tumors.

In MEN 2B along with MTC and pheochromocytoma, it is associated with other abnormalities such as mucosal neuromas, marfanoid habitus, medullated corneal nerve fibers, and megacolon. Parathyroid tumors are generally not seen here, in contrast to the MEN 2A type.

MTC presents early and is usually aggressive. MTC is the most common feature, presents as a palpable mass in the neck, and can be associated with pressure symptoms. Ectopic adrenocorticotropic hormone (ACTH) secreted by the tumor can cause Cushing's syndrome.

Pheochromocytoma is frequently bilateral. Palpitations, sweating, and headaches are the features of pheochromocytoma.

Mucosal neuromas can present with bumpy and enlarged lips, tongue, and eyelids. Intestinal autonomic ganglion dysfunction can lead to megacolon and diverticula. Medullated corneal fibers and delayed puberty can also be seen. Marfanoid habitus marked by tall stature, long limbs, arachnodactyly, and joint hypermobility is an associated feature.

Other options:

Cafe au lait spots and optic nerve gliomas are seen in neurofibromatosis 1 (NF1).

Solution to Question 40:

The wrong statement regarding the examination of the abdomen is statement 3. Palpation should ideally begin away from the site of pain. Palpating from the least to the most painful area allows one to appreciate and compare normal tissue with the painful region.

Other statements:

Cullen sign refers to periumbilical bruising in cases of hemoperitoneum, especially those of pancreatic origin.

Grey-Turner sign refers to bluish discoloration around flanks in cases of hemoperitoneum, especially acute hemorrhagic pancreatitis.

Guarding refers to the contraction of the abdominal muscles. It can either be voluntary or involuntary (reflex contraction). It indicates inflammation of the parietal peritoneum.

Rigidity of the anterior abdominal wall muscles is seen when there is generalized peritonitis.

Discoloration of flanks: Grey Turner sign

Cullen's sign: Periumbilical bruising seen in pancreatitis:

Solution to Question 41:

A diabetic patient with a urinary tract infection, now presenting with hypotension unresponsive to IV fluids is suggestive of urosepsis (septic shock). Hence, the patient should be treated with Piperacillin-Tazobactam. In septic shock, there is peripheral vasodilation due to toxins from the infectious agent causing endothelial injury and fluid leak, leading to hypotension which can be refractory to fluid therapy in the early stages.

Urinary tract infections (UTIs) are more common in diabetic patients due to poor metabolic control, altered immune status, poor bladder control and incomplete voiding. E.coli is the most common organism to cause urinary tract infections. Other organisms causing infections include Klebsiella spp, Proteus spp, Enterobacter spp and Enterococci. They are more prone to resistance

to common antibiotics used in uncomplicated urinary tract infections. UTIs can range from asymptomatic bacteriuria, cystitis, pyelonephritis and complicated urosepsis.

Other options:

Option B: Amoxiclav is used for the treatment of upper respiratory tract infections and skin and soft tissue infections. Though most cases of UTI are caused by E.coli, due to increased resistance (ESBL-producing strains), it can be challenging to treat with oral beta-lactams and generally, they have lower cure rates compared to other classes of drugs such as fluoroquinolones or trimethoprim-sulfamethoxazole.

Option C: When a patient with uncomplicated pyelonephritis is treated with trimethoprim-sulfamethoxazole and organism susceptibility is unknown, ceftriaxone 1 gram IV can be used initially.

Option D: Nitrofurantoin is used for the treatment of uncomplicated cystitis.

Factors that suggest a complicated urinary tract infection:

• Functional or anatomic abnormality of the urinary tract

• Male gender

• Diabetes

• Elderly

• Immunosuppression

• Childhood urinary tract infection

• Recent antimicrobial usage

• Indwelling urinary catheter

• Urinary tract instrumentation

• Hospital-acquired infection

• Symptoms more than 7 days at presentation.

Solution to Question 42:

Linagliptin does not require dose adjustment with renal failure as it is cleared primarily by the hepatobiliary system and hence no adjustment is needed in renal failure.

Glimepiride (Option A) is a second-generation sulfonylurea which is excreted by the kidneys and hence cannot be used in renal failure. Vildagliptin (Option C) and other DPP-4 inhibitors (saxagliptin, sitagliptin, and alogliptin) are largely excreted by the kidneys and cannot be used in chronic renal failure. Exenatide (Option B) is a GLP-1 agonist which is excreted by the kidneys and hence cannot be used in renal failure.

Linagliptin is a dipeptidyl-peptidase-4 (DPP-4) inhibitor used in type 2 diabetes as monotherapy or in combination therapy when metformin is not tolerated or contraindicated. It binds extensively to plasma proteins and is cleared primarily by the hepatobiliary system. Hence, no dose adjustment is needed in renal failure.

Oral semaglutide is the first oral GLP-1 agonist approved for the treatment of type 2 diabetes mellitus, available as 3 mg, 7 mg and 14 mg tablets. Unlike most DPP-4 inhibitors, it is not eliminated by renal clearance and can be used in patients with renal impairment. Patients should be advised to take the tablet 30 minutes before food intake. Side effects include nausea, vomiting and abdominal pain. Some trials have shown semaglutide to be associated with an increased risk of gallbladder disease.

Solution to Question 43:

A hypertensive patient developing diabetes should be switched to telmisartan.

Telmisartan, an angiotensin receptor blocker (ARB), has shown evidence of having PPAR-gamma agonist action. It is very useful in hypertensive patients with diabetes. PPAR-gamma agonists are the thiazolidinedione group of drugs that act by inducing several insulins sensitizing genes, thereby reducing overall insulin resistance.

ACE inhibitors or ARBs are the first-line treatment of choice for hypertension in a diabetic patient. They reduce proteinuria and delay diabetic nephropathy, a benefit that is not seen with beta-blockers or calcium channel blockers.

Solution to Question 44:

The earth to all leads is in the right leg is the correct statement about the placement of ECG leads.

Bipolar limb leads for ECG recording

- LA → left arm
- RA → right arm
- LL → left foot

Lead type	Positive input (+)	Negative input (-)
Lead I	LA	RA
Lead II	LL	RA
Lead III	LL	LA



Solution to Question 45:

The image shows extensor plantar response, which is a positive Babinski sign.

Babinski sign refers to dorsiflexion of the great toe and fanning of other toes when the lateral aspect of the sole of the foot is scratched. It is a sign of an upper motor neuron lesion. In normal adults, the response is plantar flexion of all the toes.

UMNs refer to corticospinal tract neurons and brainstem neurons that control spinal motor neurons.

Injury to UMNs leads to

• Spasticity

• Hypertonia

• Hyperreflexia

• Abnormal plantar extensor reflex (positive Babinski sign)

The below image shows the positive and negative Babinski signs.



The image below shows a positive Babinski sign in an adult, which is abnormal and indicative of a UMN lesion.

In infants whose corticospinal tracts are still underdeveloped, dorsiflexion of the great toe and fanning of other toes is a normal response to scratching of the sole of the foot. The image below shows an extensor response in an infant, which is normal.

Solution to Question 46:

The most common site of abdominal tuberculosis is the ileocaecal junction.

Amongst extrapulmonary tuberculosis (TB), abdominal TB is a relatively uncommon form that may occur due to direct seeding by swallowing of sputum, hematogenous spread, or ingestion of milk from cows with bovine TB. It most commonly involves the terminal ileum and caecal regions, though any part of the gastrointestinal tract may be involved.

Clinical features include abdominal pain, palpable mass, hematochezia, and obstruction, along with symptoms characteristic of TB such as fever, weight loss, and night sweats.

The characteristic findings on barium meal and follow-through are strictures, narrow ileum, and high subhepatic cecum.

The image below shows string sign on the barium enema. This finding is seen in ileocaecal tuberculosis with narrowing of terminal ileum, ileocaecal junction, and dilated proximal ileum:



Solution to Question 47:

The duration of prophylaxis given to this child is 10 years.

The given scenario is a case of acute rheumatic fever without carditis and no valvular pathology. The secondary prophylaxis for this child is given for 5 years or till the age of 21, whichever is longer. Since the boy is 11 years old, prophylaxis for 10 years till the age of 21 years is longer than 5 years.

ARF is an autoimmune disease that develops in response to group A streptococcal infection. It is commonly seen in children between the age of 5-14 years. An upper respiratory tract infection with some M serotypes of the organism is the trigger. The concept of molecular mimicry is used to explain the pathogenesis of the condition.

Since one attack of ARF increases the risk for further episodes and development of RHD, secondary prophylaxis is recommended by the American heart association (AHA) as follows:

• Rheumatic fever without carditis: For 5 years after the last attack or till 21 years of age (whichever is longer).

• Rheumatic fever with carditis but no residual valvular disease: For 10 years after the last attack or till 21 years of age (whichever is longer).

• Rheumatic fever with persistent valvular disease, evident clinically or on echocardiography: For 10 years after the last attack or till the age of 40 (whichever is longer). Sometimes lifelong prophylaxis can be given.

Symptoms usually present after a latent period of about three weeks. Indolent carditis and chorea are the exceptions that can present even six months later. Migratory asymmetric polyarthritis which affects large joints and high-grade fever are common features. Cutaneous manifestations include erythema marginatum and subcutaneous nodules. Sydenham's chorea is commonly found in females and frequently affects the head and upper limbs. Acute rheumatic carditis can be seen as the involvement of the endocardium, myocardium, or pericardium.

Histopathological examination shows Aschoff bodies and activated macrophages called Anitschkow cells.

The image below shows Aschoff bodies:



In chronic rheumatic heart disease, there is a leaflet and tendinous cord thickening along with commissural fusion and shortening of the mitral valve. Ultimately the valve can undergo stenotic changes giving it a fish-mouth appearance as shown below.



The modified Jones criteria is used for diagnosis. Management of ARF is mainly symptomatic by the use of antibiotics and NSAIDs.

Solution to Question 48:

Methanol intoxication is characterized by metabolic acidosis with a high anion gap.

Methanol is metabolized in the body as follows:

Methanol -Alcohol Dehydrogenase (ADH)→ Formaldehyde -Aldehyde Dehydrogenase→ Formic acid

Formic acid is an unmeasured anion causing a high-anion gap and metabolic acidosis. It accumulates in the CNS and retina and causes blurring vision/loss of vision.

Other causes of high-anion gap metabolic acidosis:

• Lactic acidosis

• Ketoacidosis: alcohol, starvation, diabetes

• Toxins: Ethylene glycol, methanol, salicylate

• Renal failure (acute and chronic)

Solution to Question 49:

The given clinical history suggests a typical case of hypothyroidism, and the likely diagnosis is myxedema coma.

Myxedema coma is a severe neurological complication of hypothyroidism. It is primarily seen in the elderly and has a high mortality rate of 20–40%. It is often precipitated by the following factors:

• Drugs like sedatives, antidepressants, and anesthetics

• Infections like pneumonia

• Congestive heart failure, myocardial ischemia

• Cerebrovascular accidents

• Gastrointestinal bleeding

• Exposure to cold

All these causes impair respiration. Hypoventilation and the ensuing hypoxia and hypercapnia play a major role in the pathogenesis of myxedema coma.

Clinical features include altered consciousness, seizures, hypothermia (can reach 23°C), metabolic disturbances, and symptoms and signs of hypothyroidism (fatiguability, cold intolerance, non-pitting edema, weight gain, constipation, menorrhagia, etc).

Management of myxedema coma involves:

• Pharmacotherapy:

• Levothyroxine (T4): Single loading IV dose (200–400 mcg) and maintenance oral dose (1.6 mcg/kg/day)

• Liothyronine (T3): Added along with T4 either via nasogastric tube or IV

• Parenteral hydrocortisone (50 mg every 6 h): Given to correct the impaired levels of corticoid hormones

• Broad-spectrum IV antibiotics: Given in case of infections

• Supportive therapy is by correction of associated metabolic imbalance, external warming (if body temperature is <30 degrees) to prevent circulatory collapse by using space blankets, and ventilatory support with regular blood-gas analysis in the first 48 hrs.

Other options:

Option A: Septic shock occurs due to systemic inflammation like bacterial and fungal infections, toxic shock syndrome, pancreatitis, massive trauma, and severe burns, resulting in organ failure. A massive release of inflammatory cytokines causes peripheral vasodilation and pooling, which causes the skin to look flushed. Vascular damage, DIC, metabolic derangements, and diffuse alveolar damage (shock lung) can be seen.

Option B: Cardiogenic shock occurs due to myocardial pump failure, resulting in low cardiac output. It is associated with conditions like myocardial infarction, arrhythmias, cardiac tamponade, and pulmonary embolism. Patients present with hypotension, a weak rapid pulse, tachypnea, and cool clammy skin.

Option D: Hyperthyroidism presents with clinical features that include heat intolerance and sweating, tachycardia and palpitations, weight loss with increased appetite, diarrhea, and oligomenorrhea. Clinical findings include exophthalmos, tremors, and myopathy.

Solution to Question 50:

The headaches that need further evaluation are 2, 3, and 4.

Statement 1: Headache for >4 hours can be a chronic tension-type headache or migraine, which does not require further evaluation unless associated with other warning symptoms.

Statement 2: Worst headache of life - Patients with subarachnoid hemorrhage often describe their headache as the worst headache of their life. This condition requires further evaluation, usually a non-contrast CT scan.

Statement 3: New-onset progressive headache has a higher chance of being caused by a serious underlying cause as compared to headaches that have a similar intensity and a longstanding history. Underlying causes may include meningitis, subarachnoid hemorrhage, subdural or epidural hematoma, and purulent sinusitis. Hence, these headaches need further evaluation.

Statement 4: Headache with blurring of vision could be due to a variety of causes such as migraine, increased intracranial pressure, acute angle closure glaucoma, etc. Hence, these headaches need to be evaluated to find the exact cause and rule out serious underlying conditions.

Solution to Question 51:

This clinical vignette is suggestive of tuberculous peritonitis secondary to pulmonary tuberculosis. Anti-tubercular therapy should be started as the next step in the management of this condition.

This patient presents with cough, cold, fever, and malaise associated with weight loss, and a radiograph showing a homogenous dense infiltrate in the right lung is consistent with pulmonary tuberculosis (TB). The presence of ascites along with the above-mentioned signs and symptoms is suggestive of tuberculous peritonitis.

The various pathogenic mechanisms of gastrointestinal TB include direct seeding into the abdomen due to swallowing of the sputum, hematogenous seeding of tubercle bacilli from rupture of lymph nodes or intraabdominal organs, and ingestion of milk from cows infected with bovine TB. It presents with abdominal pain, swelling, obstruction, hematochezia, and a palpable abdominal mass. Constitutional symptoms such as fever, weight loss, night sweats, and anorexia are also present. The terminal ileum and cecum are the most commonly involved sites.

The presence of fever, abdominal pain, and ascites should raise the suspicion of tuberculous peritonitis. The clinical presentation may be similar to Crohn's disease if the intestinal wall is involved, leading to ulcerations and fistulae.

Paracentesis reveals an exudate with high protein content, lymphocytosis, and elevated levels of adenosine deaminase(ADA). Elevated ADA levels have very high sensitivity for tuberculous peritonitis. Standard anti-tubercular therapy is recommended for ascites caused by tuberculous peritonitis. If the etiology of the ascites is uncertain, laparoscopy with peritoneal biopsy for culture and histology remains the gold standard.

Solution to Question 52:

The given clinical scenario of a patient with chronic stable angina, presenting with fatigue and brownish-colored blood is suggestive of methemoglobinemia. For the management of chronic stable angina, she must be taking nitrates and methemoglobinemia is a rare complication of nitrate (isosorbide) consumption.

Nitrate ions released during the metabolism of isosorbide oxidize hemoglobin (Fe^{2+}) to the brown methemoglobin (Fe^{3+}), thereby producing methemoglobinemia. The risk of developing clinically significant methemoglobinemia is higher in patients with coronary ischemia and anemia.

Interestingly, pulse oximetry in methemoglobinemia shows a plateau of 80% to 85% irrespective of the arterial oxygen saturation and irrespective of the methemoglobin levels. Co-oximeters that can differentiate methemoglobin from carboxyhemoglobin, oxyhemoglobin, and deoxyhemoglobin are thus preferred for the diagnosis of methemoglobinemia.

Treatment of methemoglobinemia associated with isosorbide toxicity involves supportive care, cessation of isosorbide consumption, and reduction of methemoglobin to hemoglobin by intravenous methylene blue (1–2 mg/kg).

Other options:

Option A: Carboxy-hemoglobinemia occurs due to acute and chronic carbon monoxide intoxication through occupational exposure and exposure to incompletely burned hydrocarbons. It presents with headache, altered mental status, and other constitutional symptoms.

Option B: Sideroblastic anemia is a type of anemia that occurs in the presence of normal to high levels of iron. The characteristic ring sideroblasts are seen in the peripheral smear. Causes of sideroblastic anemia include pyridoxine deficiency, lead poisoning, and alcoholism.

Option D: Sulfhemoglobinemia is a rare irreversible condition that can occur with the ingestion of drugs such as dapsone and sulphonamides.

Solution to Question 53:

The given clinical scenario is suggestive of lateral medullary syndrome/Wallenberg syndrome. This may result from a stroke affecting the ipsilateral vertebral artery.

Lateral medullary syndrome: The clinical findings correspond to the neuroanatomical structures affected in the lateral portion of the medulla:

• Descending tract and nucleus of CN V (trigeminal)- loss of pain and sensation over the ipsilateral half of the face.

• Inferior cerebellar peduncle – ipsilateral ataxia (gait and limb)

• Lateral spinothalamic tract – loss of pain and temperature sensation over the contralateral half of the body.

• IX and X CN - ipsilateral paralysis of palate, vocal cord, diminished gag reflex giving rise to dysphagia, dysarthria.

• Vestibular nucleus - Nystagmus, diplopia, vertigo, nausea

• Descending sympathetic tract – ipsilateral Horner's syndrome (miosis, ptosis, and anhidrosis).



Solution to Question 54:

The probable diagnosis based on the given clinical scenario and the image showing cavitation around the central canal is syringomyelia.

Syringomyelia is a progressive disease of the spinal cord in which there is an enlarging cavity in the cord. Syrinx cavities are acquired cavities of the cord in necrosed areas secondary to trauma, myelitis, and spinal cord tumors. It usually presents in adolescence or early adulthood with central cord symptoms.

Most patients present with regional dissociated sensory loss of pain and temperature of the upper limbs. Touch and vibration senses are intact. The loss of sensation has a 'cape distribution'

affecting the nape of the neck, shoulders, and upper arms. Eventually, expansion of the cavity may lead to compression of the long tracts. This results in spasticity and weakness of the legs, bowel and bladder dysfunction, and Horner's syndrome.

MRI is used for the diagnosis. There is no satisfactory treatment for this condition. Symptomatic syrinx cavities secondary to trauma or infection, are treated with a decompression and drainage procedure.

Other options:

Option A: Brown-Sequard syndrome occurs due to the transection of the spinal cord only on one side. This results in ipsilateral weakness and loss of joint position and vibration sensation with contralateral loss of pain and temperature sensation.

Option B: Tabes dorsalis is a manifestation of neurosyphilis, which occur within a year or up to 10-35 years after primary infection. It causes demyelination of the dorsal column of the spinal cord and presents with bladder disturbances, foot drop, impotence, and ataxia. Loss of position and vibration sense result in a positive Romberg test.

Option C: Syringobulbia occurs when the syrinx of syringomyelia extends into the medulla. It is characterized by palatal or vocal cord paralysis, episodic vertigo, tongue weakness with atrophy, and horizontal nystagmus.

Solution to Question 55:

The success of treatment after H.pylori can be assessed by the urea breath test.

Peptic ulcer disease is most likely caused by Helicobacter pylori. H. pylori is a gram-negative motile bacillus. It has a unique feature of producing urease which is helpful in diagnosis. The sole source of H. pylori is the human gastric mucus. H.pylori causes chronic superficial gastritis on long-term colonization. Only a few people develop peptic ulceration, gastric adenocarcinoma, and mucosa-associated lymphoid tissue (MALT) lymphomas.

Diagnosis can be made by tests based on endoscopic biopsy:

• Biopsy urease test, which is quick and simple (Option D)

• Histology can give additional information.

• Culture can determine antibiotic susceptibility.

Non-invasive tests:

• Serology tests (Option B: IgG antibody study) are inexpensive and convenient but cannot be used to monitor treatment success.

• 13C Urea breath test requires fasting but is useful for follow-up.

• Stool antigen test is useful, particularly in children, and is also useful for follow-up.

The urea breath test is used to assess the success of treatment as it is non-invasive and less expensive than endoscopy. However, because this test is dependent on H. pylori load, use in less than 4 weeks after treatment may yield false-negative results. Endoscopy and a biopsy-based test are also used in follow-up after treatment. The main indication for these invasive tests is to exclude underlying gastric adenocarcinoma in a case of gastric ulcer.

Current treatment regimens consist of a proton-pump inhibitor (omeprazole) and two or three antimicrobial agents given for 10–14 days.

Solution to Question 56:

The above X-ray and symptoms are suggestive of pleural effusion secondary to a lower respiratory tract infection. On examination, dullness to percussion will be observed.

Pleural effusion refers to the pathological accumulation of fluid in the pleural space due to

• excess fluid formation from the interstitial spaces of the lung, parietal pleura, or peritoneal cavity

• decreased fluid removal by the lymphatics

Pleural effusion may present with dyspnea, pleuritic pain, dry cough, and associated symptoms like fever, weight loss, edema, and ascites depending on the etiology. On examination, there may be tachypnea, dullness on percussion and decreased/absent breath sounds at the base(s).

The etiology of pleural effusion can be of two types:

• Exudative: disruption in pleural fluid formation and drainage because of local factors. Examples include malignancy, infections, and pulmonary embolism.

• Transudative: disruption in pleural fluid formation and drainage due to systemic factors. Examples include congestive heart failure, hepatic cirrhosis, and nephrotic syndrome.

Option A: Hyperresonance is a sign of pneumothorax or emphysema.

Option B: Crackles are heard in cases of pulmonary edema or pneumonia.

Option D: Stridor is due to a lesion in the upper airways, usually tracheal or laryngeal.

Solution to Question 57:

The given ECG shows P-pulmonale.

P-pulmonale is characterized by an increase in P-wave amplitude ≥2.5 mm. Occasionally, a prominent positive wave in the biphasic P wave of lead V1 is seen. It is caused by overload in the right atrium and often occurs with severe pulmonary disease.

It is of limited sensitivity and specificity in the diagnosis of right atrial enlargement.

Causes of right atrial enlargement are:

• Pulmonary diseases, such as pulmonary embolism, asthma, COPD, and chronic thromboembolic pulmonary hypertension.

• Congenital heart disease, such as pulmonary valve stenosis, atrial septal defect, Ebstein's abnormality and Fallot's tetralogy.

• Tricuspid stenosis or regurgitation.

• Cardiomyopathies, such as arrhythmogenic right ventricular cardiomyopathy/dysplasia (ARVC/D).

The image below shows the ECG changes in atrial enlargement.



Solution to Question 58:

Hamman's sign can be elicited in pneumomediastinum. It is a crunching or clicking sound that is synchronous with the heartbeat and best heard in the left lateral decubitus position.

Pneumomediastinum:

The main causes of gas/air in the interstices of the mediastinum are

• Alveolar rupture, perforation or

• Rupture of esophagus/trachea/bronchi,

• Dissection of air from the neck or abdomen into the mediastinum.

Clinical features include severe substernal chest pain (may or may not radiate to the neck and arms) and subcutaneous emphysema over the suprasternal notch.

The examination may reveal subcutaneous emphysema in the suprasternal notch with Hamman's sign.

Chest x-ray confirms the diagnosis. No treatment is usually required, but inhaling high concentrations of oxygen will help absorb the mediastinal air. Needle aspiration can be performed if mediastinal structures are compressed.

Solution to Question 59:

As anti-HBc IgG antibodies are representative of a chronic hepatitis B infection, their presence may be indicative of an HBV/HDV (hepatitis B/hepatitis D) superinfection.

HDV superinfection occurs when a person who is already chronically infected with HBV acquires HDV. The infection of HDV on chronic hepatitis B can accelerate the progression to severe disease in all ages. HDV superinfection accelerates progression to cirrhosis almost a decade earlier than HBV mono-infected persons.

HBV/HDV coinfection occurs when a person simultaneously becomes infected with both HBV and HDV. In acute hepatitis, simultaneous infection with HBV and HDV can lead to mild-to-severe hepatitis. The signs and symptoms typically appear 3–7 weeks after the initial infection and include fever, fatigue, loss of appetite, nausea, vomiting, dark urine, pale stools, jaundice (yellow eyes), and even fulminant hepatitis. However, recovery is usually complete.

Diagnosis of HDV infection is by detecting high levels of anti-HDV immunoglobulin G (IgG) and immunoglobulin M (IgM). It is confirmed by the detection of HDV RNA in serum, and their levels are not helpful in differentiating between acute and chronic infections. There is no specific treatment for HDV infection. Pegylated interferon alpha may be used (with limited efficacy).

Anti-HBc IgM and anti-HBs IgM are indicative of acute phases of hepatitis B and, therefore, more likely to be associated with coinfection of HDV.

Solution to Question 60:

The correctly matched pair is 1-D,2-C,3-A,4-B.

Acid-base disorders can be divided into metabolic and respiratory causes. A disturbance in one leads to compensatory changes in the other. The best and quickest method to answer this question is to match the given values with the condition without an elaborate calculation.

• Increased pH with increased bicarbonate is metabolic alkalosis. The lungs are trying to compensate by retaining carbon dioxide via hypoventilation. Hence this is a case of metabolic alkalosis which is undergoing compensation.

• Decreased pH with decreased bicarbonate suggests metabolic acidosis. But PCO_2 levels are normal, suggesting no compensatory changes occurring in the lungs. This is uncompensated metabolic acidosis

• The pH is normal but PCO_2 is low suggesting respiratory alkalosis from hyperventilation. The kidneys have compensated by increased excretion of bicarbonate turning pH to normal range. As the pH has come to normal, it is a fully compensated respiratory alkalosis

• The pH is normal here, but an increased PCO_2 suggests respiratory acidosis due to hypoventilation. Here bicarbonate is increased suggesting a metabolic compensation. This is a case of fully compensated respiratory acidosis.

Solution to Question 61:

The patient's graph shows a reduction in both FVC & FEV1/FVC which is < 70%, indicating an obstructive lung disorder. Therefore, the most likely diagnosis is Bronchiectasis.

Chest neuromuscular disease (option A), sarcoidosis (option B) & idiopathic pulmonary fibrosis (option C) are restrictive lung disorders, where there is a reduction of FVC to < 4L, Without a loss in FEV1/FVC ratio (normal or >70%).

These values are measured as a part of the pulmonary function test known as spirometry. At a given lung volume, when a person inhales to the total lung capacity and exhales as hard and completely as possible, the volume of expired air is recorded as forced vital capacity (FVC), and the volume of air expired in the first second is called the forced expiratory volume (FEV1).



Solution to Question 62:

Orthopnea, dyspnea after 2 hours of sleep, and right upper quadrant pain/right hypochondrial pain are signs of heart failure.

Heart failure is a manifestation of most chronic cardiac conditions such as hypertension, valvular heart disease, and coronary artery disease. Symptoms such as shortness of breath, orthopnea, and paroxysmal nocturnal dyspnea (shortness of breath that develops after lying down for 30min or longer) point to left heart failure due to pulmonary congestion. In contrast, symptoms such as pedal edema, weight gain, early satiety, and hepatic congestion, tender hepatomegaly are characteristic of right heart failure.

Raised non-pulsatile JVP is seen in cardiac tamponade. Tamponade results in elevated right atrial pressure and elevated right ventricular pressures. This results in a prominent x descent and a diminished or absent y descent in the JVP. The resultant JVP is elevated but nonpulsatile in nature due to the persistently raised pressures in the right heart. (Statement 1)

Right heart failure	Left heart failure
Due tocongestionin thesyste mic circulation	Due tocongestionin thelungs.

Right heart failure	Left heart failure
Symptoms and signs:Raised pulsatile JVPCongestive hepatomegalyPositive hepatojugular refluxAscitesBilateral pitting edemaWeight gain- due to edemaFatigue- due to ↓pumping of blood to lungs leading to ↓oxygenation of the blood	Symptoms and signs:Dyspnea on exertionOrthopneaParoxysmal nocturnal dyspnea(characteristic of LVF)Pulmonary ralesIncreased PCWPFatigue- due to ↓cardiac output

Medicine NEET 2022

Question 1:

A patient with diabetes mellitus for the past 5 years presents with vomiting and abdominal pain. She is non-compliant with medication and appears dehydrated. Investigations revealed a blood sugar value of 500 mg/dL and the presence of ketone bodies. What is the next best step in management?

a) Intravenous fluids with long acting insulin

b) Intravenous fluids

c) Intravenous insulin

d) Intravenous fluids with regular insulin

Question 2:

The cerebrospinal fluid (CSF) specimen of a patient is shown below along with the microscopy. The report shows mononuclear cytosis, elevated proteins, and low sugars. Which of the following is the likely etiology?



a) Tuberculous meningitis

b) Aseptic meningitis

c) Bacterial meningitis

d) Chemical meningitis

Question 3:

A patient diagnosed to be retro-positive was started on highly active antiretroviral therapy (HAART). Which of the following can be used to monitor treatment efficacy?

a) CD4+ T cell count

b) Viral load

c) p24 antigen

d) Viral serotype

Question 4:

A female patient presents to you with a unilateral headache. It is associated with nausea, photophobia, and phonophobia. What is the drug of choice for acute management?

a) Flunarizine

b) Sumatriptan

c) Propranolol

d) Topiramate

Question 5:

A hypertensive patient who is non-compliant with medication presents to you with sudden onset breathlessness. A chest X-ray was done, which is shown below. How will you manage this patient?



a) Intravenous salbutamol

b) Intravenous nitroglycerine

c) Nebulization with salbutamol

d) Oxygen and antibiotics

Question 6:

A patient on anti-depressants presented to you with hypotension. An ECG was done, which showed wide QRS complexes and right axis deviation. How will you manage this patient?

a) Antiarrhythmics

b) Intravenous sodium bicarbonate

c) Propranolol

d) Phenytoin

Question 7:

A patient presents to you with fever, night sweats, ptosis, and bilateral facial nerve palsy. Investigations showed leukocytosis and bilateral hilar lymphadenopathy. Which of the following is the most likely diagnosis?

a) Sarcoidosis

b) Tuberculosis

c) Lymphoma

d) Hypersensitive pneumonitis

Question 8:

A 35-year-old female patient presents to you with fever, breathlessness, and cough with expectoration. A CT scan was done which is shown below. What is the most likely diagnosis?



a) Consolidation with air bronchogram

b) Mediastinal mass

c) Pleural effusion

d) Diaphragmatic hernia

Question 9:

A 25-year-old patient is undergoing tooth extraction for dental caries. Which of the following does not require prophylaxis against infective endocarditis?

a) Prior history of endocarditis

b) Atrial septal defect

c) Unrepaired cyanotic heart disease

d) Prosthetic heart valves

Question 10:

A patient presents to the emergency department with a history of ingestion of ten tablets of paracetamol. He has developed oliguria and liver function tests show deranged values. Which of the following can be used in the management of this condition?

a) N-acetylcysteine

b) Dopamine

c) Ursodeoxycholic acid

d) Furosemide

Question 11:

A female patient with a negative urine pregnancy test presents to you with galactorrhea. An MRI was done which revealed a large pituitary tumor. If the patient is not willing for surgery, which of the following is the best drug for treatment?

a) Bromocriptine

b) Promethazine

c) Octreotide

d) Clozapine

Question 12:

A patient presents to you with an irregularly irregular pulse of 120/minute and a pulse deficit of 20. Which of the following would be the jugular venous pressure (JVP) finding?

a) Absent p wave

b) Absent a wave

c) Cannon a wave

d) Raised JVP with normal waveform

Question 13:

A patient with a history of chronic liver disease presents with abdominal distension, jaundice, and pruritis. Ascitic fluid analysis revealed neutrophil count > 650 per cubic mm. What is the most likely diagnosis?



a) Spontaneous bacterial peritonitis

b) Malignant ascites

c) Tubercular ascites

d) Intestinal obstruction

Question 14:

Laboratory investigations of a patient being evaluated for jaundice show elevated bilirubin and alkaline phosphatase levels. Levels of the remaining liver enzymes are normal. What is the likely diagnosis?

a) Obstructive jaundice

b) Hemolytic jaundice

c) Hepatic jaundice

d) Prehepatic jaundice

Question 15:

A woman presents to you with fever, arthralgia, ulcers, fatigue for the past six months, and new-onset hematuria. Urine examination reveals RBC casts and proteinuria. What is the likely diagnosis?

a) Acute interstitial nephritis

b) Poststreptococcal glomerulonephritis

c) Lupus nephritis

d) IgA nephropathy

Question 16:

An 11-year-old child with a history of streptococcal pharyngitis presents you with fever and arthralgia. There is no past history of rheumatic heart disease or features of carditis or valvular disease. How often is 6,00,000 IU of benzathine penicillin recommended for prophylaxis of rheumatic heart disease?

a) Immediately

b) Thrice weekly lifelong

c) Once in three weeks for 5 years or till the age of 18, whichever is longer

d) Once in three weeks for 10 years or till the age of 25, whichever is longer

Question 17:

A man on diuretics presents with weakness. An ECG was done which showed flat T waves and prominent U waves. What is the most likely diagnosis?

a) Hypokalemia

b) Hyperkalemia

c) Hypomagnesemia

d) Hypernatremia

Question 18:

A male patient presents to the emergency department. The arterial blood gas report is as follows: pH, 7.2; pCO_2, 81mm Hg; and HCO_3, 40 meq/L. Which of the following is the most likely diagnosis?

a) Respiratory alkalosis

b) Metabolic acidosis

c) Respiratory acidosis

d) Metabolic alkalosis

Question 19:

Which of the following is not seen in MEN 2B syndrome?

a) Megacolon

b) Parathyroid adenoma

c) Mucosal neuroma

d) Marfanoid habitus

Question 20:

A patient with coronary artery disease presents to you with chest pain and palpitations. The ECG is shown below. Which of the following can be used in the management?



a) Oral metoprolol

b) Oral amiodarone

c) Intravenous amiodarone

d) Intravenous metoprolol

Question 21:

Multidrug-resistant (MDR) tuberculosis shows resistance to which of the following drugs?

a) Isoniazid, rifampicin, and fluoroquinolone

b) Fluoroquinolones

c) Isoniazid and rifampicin

d) Isoniazid, rifampicin, and kanamycin

Question 22:

A patient presents to you with fever, jaundice, and malaise. What is the most likely diagnosis based on the serology reports given below?

a) Acute hepatitis B

b) Acute hepatitis C

c) Chronic hepatitis B

d) Chronic hepatitis C

Question 23:

A child presents to the emergency department with a history of ingestion of 10-20 ferrous sulfate tablets. Arterial blood gas revealed acidosis. Which of the following can be used in the management of this condition?

a) Deferoxamine

b) Activated charcoal

c) Dimercaprol

d) Penicillamine

Answer Key

Question No.	Correct Option
1	d
2	a
3	b
4	b
5	b
6	b
7	a
8	a
9	b
10	a

11	a
12	b
13	a
14	a
15	c
16	c
17	a
18	c
19	b
20	c
21	c
22	a
23	a

Detailed Explanations

Solution to Question 1:

The clinical vignette points to a diagnosis of diabetic ketoacidosis (DKA) for which the next best step in management would be intravenous fluids with regular insulin.

DKA is a complication of diabetes mellitus that arises due to relative or absolute insulin deficiency, seen more commonly in patients with diabetes mellitus type 1. Clinical features include nausea, vomiting, abdominal pain, and tachycardia. Hyperglycemia leads to volume depletion and subsequent hypotension. Kussmaul respiration and fruity odor on the breath are characteristic of the condition.

Investigation shows elevated glucose levels (250-600 mg/dL), ketonemia, and metabolic acidosis with a high anion gap.

Patients with DKA are also at an increased risk of developing rhinocerebral mucormycosis. Coma and cerebral edema are extremely serious complications that can arise.

Once the diagnosis is confirmed intravenous fluids should be given along with a bolus of short-acting regular insulin. Then, intravenous administration of 0.1 units/kg/hour of regular insulin is administered until acidosis resolves and the patient is stable.

Insulin treatment and resolution of acidosis can lead to potassium store depletion in DKA but it might be normal or slightly elevated at presentation. This is secondary to acidosis and volume depletion. If the initial potassium level is <3.3 meq/L, insulin should not be administered till it is corrected.

Other options:

Option A: Long-acting insulin is started only when the patient resumes eating.

Option B: Intravenous insulin (bolus) is one of the initial steps along with intravenous fluids.

Option C: Intravenous fluids are given as the first step in management. A bolus of intravenous insulin is also required.

Solution to Question 2:

The given clinical scenario along with the cerebrospinal fluid (CSF) sample with cobweb-like clot and slide showing lymphocytic predominance points to a diagnosis of tubercular meningitis.

Tuberculous meningitis caused by Mycobacterium tuberculosis is caused by the formation of tubercles in the brain parenchyma. This happens during the hematogenous spread of the bacilli during primary infection. It can also be due to the rupture of a subependymal tubercle into the subarachnoid space.

Tuberculous meningitis presents as a subacute infection and has a longer course than bacterial meningitis. Ocular nerve palsy is a common association as there is marked meningeal involvement at the base of the brain. The patient can also develop hydrocephalus.

The CSF sample shows a pellicle or cobweb-like clots. Acid-fast bacilli can be best isolated from these. There is usually lymphocytic leukocytosis, low glucose concentration, and elevated proteins. CSF culture remains the gold standard for diagnosis.

The clinical triad of fever, headache, and nuchal rigidity is common to all types of meningitis. Nuchal rigidity is the resistance of the neck to passive flexion and is pathognomonic of meningeal irritation. Kernig's and Brudzinski's signs are other features of meningeal irritation. Raised intracranial pressure (ICP) is a complication that presents as papilledema, disorientation, or dilated poorly reactive pupils.

Other options:

Option B - Aseptic meningitis is commonly caused by enteroviruses such as coxsackieviruses, and varicella-zoster virus. It is more benign and usually self-limiting compared to pyogenic meningitis. Gram's staining of the CSF sample shows lymphocytosis but not the organism.

Option C - Bacterial meningitis is most commonly caused by Streptococcus pneumoniae and Neisseria meningitidis. Blood culture can be used for diagnosis. CSF findings are described in the pearl below. Antibiotic prophylaxis is provided to the close contacts of persons with confirmed meningococcal disease.

Option D - Chemical meningitis is a common cause of chronic meningitis. It can arise due to the rupture of tumor contents from a cystic glioma, dermoid cyst, or craniopharyngioma into the CSF.

Solution to Question 3:

The viral load can be assessed in a retro-positive patient started on highly active antiretroviral therapy (HAART) to monitor treatment efficacy.

HAART or combination antiretroviral therapy is the mainstay in the management of retro positive patients. It is initiated soon after diagnosis except in cases of cryptococcal or tuberculous meningitis.

After initiation of therapy, a ten-fold reduction in plasma human immunodeficiency virus (HIV) RNA levels in 1-2 months can be expected. Within six months, a slower decline to <50 copies/mL is anticipated. During the first month of therapy, a rise in CD4+ T cell count is also predicted. Out of these two parameters, viral load is the better choice to monitor treatment efficacy as it is more cost-effective.

Plasma HIV RNA levels signify the rate of CD4+ T-cell destruction and the magnitude of HIV replication. It should be monitored within 2-4 weeks of initiation of therapy and after any change in the treatment regimen. It should be repeated every 4-8 weeks till the RNA levels drop to <200 copies/mL and then every 3-6 months subsequently.

The indications for change in the treatment regimen are:

• < 10-fold drop in HIV RNA levels within 4 weeks of initiation

• ≥ 3-fold increase from the nadir of plasma HIV RNA level not attributable to other causes

• Persistent decline in CD4+ T-cell counts

• Adverse effects

• Clinical deterioration

Other options:

Option C: p24 antigen is a core protein present in the capsid of the virus.

Option D: Viral serotype does not play a role in monitoring treatment efficacy. Most cases of HIV are due to HIV-1. HIV-2, another human immunodeficiency virus can also cause disease in humans.

Solution to Question 4:

The clinical vignette is suggestive of an acute attack of migraine and the drug of choice for acute management is sumatriptan.

Migraine is the second most common cause of headaches and is seen more commonly in women. It presents as an episodic headache associated with nausea, vomiting, photosensitivity, and sensitivity to sound (phonophobia). Common triggers include bright lights, sounds, and hunger.

The release of vasoactive peptides such as calcitonin gene-related peptide (CGRP) by the trigeminal nucleus cells is the theory used to explain the pathogenesis of the condition. The neurotransmitters serotonin and dopamine also play a part.

Triptans are 5-HT1B and 5-HT1D receptor agonists and they act by arresting the nociceptive nerve signaling in the trigeminovascular system. In addition, they cause cranial vasoconstriction. Cardiac side effects such as transient myocardial ischemia, coronary artery vasospasm, and myocardial infarction can occur in patients with risk factors for coronary artery disease. Other side effects include sweating, nausea, and serotonin syndrome when used in combination with selective serotonin reuptake inhibitors (SSRIs) or tricyclic antidepressants (TCAs).

Ergotamine, an ergot alkaloid are serotonin receptor agonists which can be used to treat an acute attack. They produce long-lasting vasoconstriction and can result in gangrene of fingers and toes (ergotism). These drugs are contraindicated in sepsis, ischemic heart disease, and peripheral

vascular disease.

Migraine can broadly be classified into:

• Migraine with aura

• Migraine without aura

• Chronic migraine

The migraine disability assessment score (MIDAS) can be used to assess the severity of the condition. Avoidance of recognized triggers is helpful to avoid symptoms. The various drugs used to treat an acute attack of migraine are non-steroidal anti-inflammatory drugs (NSAIDs), triptans, gepants (rimegepant), ditans (lasmiditan), and dopamine receptor antagonists (domperidone).

In patients with ≥4 attacks in a month, prophylactic medications can be used. The various FDA-approved drugs include propranolol, timolol, flunarizine, and topiramate.

Other options:

Option A - Flunarizine is a calcium channel blocker that can be used in migraine prophylaxis. Side effects include weight gain, depression, and parkinsonism.

Option C - Propranolol is a non-selective β-receptor antagonist that can be used to treat hypertension, arrhythmias, and migraine prophylaxis. It is contraindicated in patients with chronic obstructive pulmonary disease (COPD) as it can cause bronchoconstriction. Side effects include fatigue, depression, and insomnia.

Option D - Topiramate is an FDA-approved drug for generalized tonic-clonic seizures, Lennox-Gastaut syndrome, and for migraine prophylaxis in adults. Adverse effects include somnolence, weight loss, and nervousness.

Solution to Question 5:

The clinical vignette along with the X-ray given points to a diagnosis of acute cardiogenic pulmonary edema. It can be managed by the administration of intravenous nitroglycerin.

Acute severe hypertension, acute myocardial infarction, or exacerbation of chronic heart failure are some of the common causes of cardiogenic pulmonary edema. It can be precipitated by the discontinuation of medications (non-compliant patient), excessive salt intake, or tachyarrhythmias.

It presents with severe dyspnea, pink frothy sputum production, diaphoresis, and cyanosis. Auscultation reveals rales and rhonchi in all lung fields. Chest X-ray shows increased interstitial markings, vascular redistribution, and alveolar edema in a butterfly pattern. Cardiomegaly might be seen in cases of chronic heart failure.

Management of pulmonary edema includes the use of diuretics, nitrates, and morphine to decrease the preload. Nitroglycerin and isosorbide dinitrate have a rapid onset of action and can be given sublingually. If there is persistence of pulmonary edema and no hypotension, it can be administered intravenously.

Oxygen therapy to maintain a saturation >92% is recommended. Positive-pressure ventilation through a face mask or endotracheal intubation is required if saturation is not maintained even

with supplemental oxygen. A sitting position with legs hanging by the side of the bed can be used to reduce venous return in patients without hypotension.

Non-cardiac causes of pulmonary edema are high altitude, sepsis, and disseminated intravascular coagulation.

Heart failure is a manifestation of most chronic cardiac conditions such as hypertension, valvular heart disease, and coronary artery disease. Symptoms such as shortness of breath, orthopnea, and paroxysmal nocturnal dyspnea point to left heart failure due to pulmonary congestion. In contrast, symptoms such as pedal edema, weight gain, early satiety, and hepatic congestion are characteristic of right heart failure.

Other options:

Options A and C - Salbutamol is a short-acting β2-agonist that is used to treat acute severe asthma. Salbutamol nebulization is used to treat hyperkalemia. Side effects include tachycardia, tremors, hypokalemia, and raised glucose levels.

Option D - Oxygen supplementation is recommended for pulmonary edema but antibiotics have no role in cardiogenic pulmonary edema.

Solution to Question 6:

The given clinical scenario along the ECG changes described suggests a diagnosis of tricyclic antidepressant toxicity. This can be managed with intravenous sodium bicarbonate.

Tricyclic antidepressants (TCAs) are used in the treatment of psychotic depression, insomnia, and several pain conditions. Their mechanism of action is inhibition of norepinephrine and serotonin reuptake. They have a quinidine-like sodium channel blockade effect on cardiac conduction. Hence, they cause QT prolongation. Other cardiotoxic effects include atrioventricular (AV) block and hypotension. Rightward-axis deviation is also an associated finding.

TCA toxicity can result in ventricular tachyarrhythmias, status epilepticus, hyperthermia, and coma. It can also result in anticholinergic syndrome due to muscarinic receptor antagonism. Symptoms include dilated pupils, high temperature, dry mouth, and myoclonic movements. It is managed with intravenous sodium bicarbonate bolus (50-100 mEq). This helps in the reversal of sodium channel blockade. Gastric lavage and administration of activated charcoal can be carried out in cases of recent ingestion.

Other side effects of TCAs include urinary retention, tachycardia, and postural hypotension. ECG changes such as T wave inversion or suppression are also common.

In some cases, it can result in polymorphic ventricular tachycardias with QT prolongation known as torsades de pointes. Recurrent episodes can be treated with intravenous magnesium sulfate.

Other options:

Option A: Class IA, IC, and III antiarrhythmic drugs are contraindicated as they potentiate the action of TCAs.

Option B: Propranolol is a class II antiarrhythmic that can cause wide QRS complexes. It is used for rate control in atrial fibrillation, angina, and hypertension.

Option D: Phenytoin is an antiepileptic used for the treatment of generalized tonic-clonic seizures and focal seizures. It inactivates sodium channels and stabilizes neuronal membranes. It can displace TCAs from their protein binding sites which can result in transient overdose symptoms.

Solution to Question 7:

The given clinical vignette points to a diagnosis of sarcoidosis.

Sarcoidosis is a granulomatous inflammatory condition with multisystem involvement. The most common organs affected are the lungs, liver, eye, and skin. The most widely accepted etiological hypothesis is an inflammatory response to an infectious or non-infectious trigger in a susceptible host.

There are two syndromes that are associated with the condition.

• Lofgren's syndrome is characterized by erythema nodosum and hilar adenopathy. Periarticular arthritis can also be seen.

• Mikulicz syndrome is bilateral involvement of parotid, submaxillary, and sublingual glands.

The most common presenting features are cough, dyspnea, fever, night sweats, and weight loss. The eyes can be involved in the form of anterior uveitis, photophobia, dry eyes, and blurred vision. Neurological involvement in the form of myelopathy, basilar meningitis, and facial nerve paralysis. Skin involvement can be seen as maculopapular lesions, keloid formation, ad subcutaneous nodules. Lupus pernio refers to the skin involvement of the bridge of the nose, periocular region, and cheeks and points to a chronic state.



Histopathological examination reveals non-necrotizing granulomas made of macrophages and giant cells. Calcium and protein concretions known as Schaumann bodies and stellate inclusions called asteroid bodies may be seen within the giant cells.



Schaumann body within a giant cell



Chest X-ray is preferred to stage lung involvement as follows.



CT features include peribronchial thickening and reticular nodular changes. Glucocorticoids are the drug of choice for management.

Other options:

Option B - Tuberculosis (TB) is a chronic granulomatous systemic disease. The underlying pathophysiology is type IV hypersensitivity reaction. It can present as pulmonary or extrapulmonary TB. Symptoms include fever, night sweats, weight loss, cough, and hemoptysis. Histopathological examination reveals caseating tubercles with multinucleated giant cells and epitheloid cells.

Option C - Lymphomas are broadly classified into Non-Hodgkin's lymphoma (NHL) and Hodgkin's lymphoma. Symptoms include fever, weight loss, night sweats, lymphadenopathy, and splenomegaly.

Option D - Hypersensitivity pneumonitis arises due to inhalation of antigenic triggers which result in an inflammatory response. Symptoms include fever, chills, malaise, and dyspnea. Cough, weight loss, and fatigue can be seen in chronic cases.

Solution to Question 8:

The given history and the CT scan image are highly suggestive of pneumonia in which, consolidation with an air bronchogram can be seen.

A patient presenting with fever, breathlessness, and cough with expectoration is most likely suffering from pneumonia. Other symptoms include nausea, vomiting, diarrhea, fatigue, headache, myalgias, arthralgias, etc. The CT image shows a sharply demarcated homogenous consolidation that crosses segmental boundaries and involves only a single lobe. It is suggestive of lobar pneumonia.

In lobar pneumonia, the infection develops within the distal airspaces. It spreads via the collateral air drift and produces homogenous opacification of complete or partial lung segments. The airways remain patent as they are not primarily involved. Patent bronchi may be seen within the

area of consolidation called air bronchograms. Total resolution of the opacification occurs within 4-6 weeks.

The most common organism implicated in community-acquired adult bacterial pneumonia is Streptococcus pneumoniae. Some of the predisposing factors include:

• Chronic illness

• Splenectomy

• Alcoholism

• Sickle cell disease

Other options:

Option B: A mediastinal mass can be in one of the three mediastinal compartments namely the anterior, middle, or posterior compartment. CT scan is the only imaging modality that should be done for the evaluation in most instances. Barium studies are indicated in patients with posterior mediastinal masses. Given below is a CT image of a large mass in the left anterior mediastinum, most likely a thymoma.



Option C: Pleural effusion is the accumulation of fluid within the pleural space. It can be either transudative (seen in cardiac failure) or exudative (seen in malignancies, infections, pulmonary embolism, etc.). On a CT image, pleural effusion appears as a dependent sickle-shaped opacity of low attenuation. The image below shows a CT image of right-sided pleural effusion.



Option D: Diaphragmatic hernia occurs due to intrathoracic movement of the abdominal contents through a diaphragmatic defect. The types include:

• Bochdalek hernia (posterior): Most common type. Herniation occurs through the pleuroperitoneal foramen.

• Morgagni hernia (anterior): Herniation occurs through the foramen of Morgagni.

Given below is a CT image showing Bochdalek hernia.

Solution to Question 9:

Atrial septal defect is not considered a high-risk condition that warrants prophylaxis against infective endocarditis.

Bacterial or fungal infection of the valvular or endocardial surface of the heart is known as infective endocarditis. Large friable vegetations on the heart valves composed of fibrin, inflammatory cells, and organisms are characteristic of the condition. Predisposing conditions include mitral valve prolapse, ventricular septal defect, rheumatic heart disease, intravenous drug use, and prosthetic valves.

Prophylaxis against infective endocarditis is recommended during certain dental procedures and respiratory tract procedures. In patients not allergic to penicillin, 2g of amoxicillin can be given orally one hour before the procedure. Another option is to administer 2g of ampicillin intravenously or intramuscularly 30 minutes before the procedure. Though prophylaxis is not advised, it is recommended that skin and genitourinary tract infections be treated before undergoing the procedure at these sites.

The most common organisms responsible are Staphylococcus aureus, streptococci, and pneumococci. The oral cavity is the portal for viridans streptococci. Staphylococci are responsible for early prosthetic valve endocarditis. In late prosthetic valve endocarditis, streptococci are commonly implicated.

Clinical features include subungual splinter hemorrhages, Osler nodes, Janeway lesions, and Roth spots.

The image below shows Osler nodes which are painful raised violaceous lesions commonly affecting the fingers, toes, and feet. It is immunologically mediated.

The image below depicts Roth's spots which are exudative lesions in the retina. This is an immunological phenomenon.



The image given below shows Janeway lesions which are vascular lesions and not immunologically mediated.

Symptoms such as dyspnea, cough, arthralgias, and flank pain can also arise due to embolization or immunologically mediated phenomena.

Antibiotics are the mainstay in management. In certain cases, surgical management is necessary.

Other options:

Options A, B, and C - According to American Heart Association (AHA), prior history of endocarditis, unrepaired cyanotic heart disease, and prosthetic heart valves are all high-risk cardiac lesions that necessitate antibiotic prophylaxis.

Solution to Question 10:

The clinical vignette suggests acetaminophen (Paracetamol) toxicity, which can be managed with N-acetylcysteine.

Early manifestations (within 4-12 hours) of acetaminophen toxicity include nausea, vomiting, diarrhea, and abdominal pain. Later, renal failure and myocardial injury may be seen. Low bilirubin levels along with raised aminotransferase levels are characteristic of the condition.

Acetaminophen toxicity is an important cause of acute liver failure. Ingestion of ≥25g can be fatal. Cytochrome P450 CYP2E1 metabolizes acetaminophen to N-acetyl-p-benzoquinone-imine (NAPQI). Glutathione is responsible for its detoxification, after which it is excreted in the urine. When glutathione reserves are depleted or large amounts of NAPQI are formed, it binds to hepatocyte macromolecules. This can then lead to hepatocyte necrosis.

Initial management includes gastric lavage and oral charcoal or cholestyramine. It should be carried out within thirty minutes of ingestion. If they are used, gastric lavage must be done afterward before using another agent. Acetaminophen blood levels can be estimated from a nomogram plot. In patients with high acetaminophen levels, intravenous N-acetylcysteine administration within 8 hours of ingestion is recommended. It helps in the replenishment of glutathione. It is partially effective even up to 36 hours. Liver transplantation is necessary if signs of hepatic failure continue despite therapy.

Other options:

Option B: Dopamine is a neurotransmitter that is used in severe congestive cardiac failure. It acts on β-1 receptors at a dose of 2-10 mcg/kg/minute.

Option C: Ursodeoxycholic acid is a bile acid that is the first-line treatment for primary biliary cholangitis. It is useful in cholestatic drug hepatotoxicity, but is not recommended.

Option D: Furosemide is a loop diuretic commonly used in pulmonary edema and congestive heart failure. Side effects include hyponatremia, hypotension, and ototoxicity.

Solution to Question 11:

The clinical vignette points to a diagnosis of prolactinoma and the best drug for the treatment of the condition is bromocriptine.

Prolactinomas are pituitary tumors that can present as microadenomas or macroadenomas. Tumors <1cm in diameter and no regional invasion are known as microadenomas. It is more commonly seen in women. Tumors >1cm in size are known as macroadenomas. It can be locally invasive. Prolactin levels >200µg/L are suggestive of prolactinomas.

Women present with features such as amenorrhea, infertility, and galactorrhea. Visual field defects can also be seen. Symptoms in males include impotence, infertility, and loss of libido. An MRI is recommended in all patients who present with hyperprolactinemia.

Microadenomas can be managed with regular monitoring of prolactin levels and MRI scans, if fertility is not a requisite. Oral dopamine agonists such as bromocriptine and cabergoline are used for medical management. They are agonists at D2 receptors. Side effects include constipation, nasal stuffiness, and dry mouth. Resistance or intolerance to these drugs requires surgical debulking. Cases that do not respond to medical or surgical management can be treated with radiotherapy.

Other options:

Option B: Promethazine is an antihistamine used to treat motion sickness, pruritus, and oculogyric crisis.

Option C: Octreotide is a somatostatin analog that is used in acromegaly, variceal bleeding, and hepatorenal syndrome.

Option D: Clozapine is an atypical antipsychotic used in refractory schizophrenia and mania.

Solution to Question 12:

The clinical scenario points to a diagnosis of atrial fibrillation and the JVP finding seen would be absent a wave.

Atrial fibrillation is the most common sustained arrhythmia. There is rapid and irregular atrial activation which is transmitted through the atrioventricular node. The conducted ventricular rate is usually rapid and irregularly irregular. It can vary from 110-160 beats/minute. A pulse deficit is an important clinical finding associated with the condition. This arises due to preload reduction.

Symptoms include palpitations, fatigue, presyncope, and dyspnea. An ECG shows the absence of P waves.



In acute atrial fibrillation with hemodynamic instability, electrical or chemical cardioversion is the first step. The latter is done by intravenous administration of class III antiarrhythmic - ibutilide. In a stable patient, ventricular rate control is the primary aim. This is done by intravenous or oral administration of beta-blockers. Calcium channel blockers such as verapamil and diltiazem can also be used.

If the current episode duration is >48 hours, cardioversion can precipitate thromboembolic complications. In such cases, cardioversion should be carried out only after 3 weeks of anticoagulation. It has to be continued for 4 weeks after cardioversion. Another option is to evaluate the presence of a thrombus in the left atrial appendage. This can be done either by transesophageal echocardiography (TEE) or high-resolution CT.

If the duration is <48 hours and there are no high-risk factors, cardioversion can be done. The CHA2DS2-VASc scoring system is used to assess stroke risk.

Warfarin is recommended in patients with rheumatic mitral stenosis or mechanical heart valves who present with atrial fibrillation. In mitral stenosis, thrombi arise frequently from the dilated left atrium. Atrial fibrillation in the absence of mitral stenosis is known as nonvalvular atrial fibrillation. Oral factor Xa inhibitors such as apixaban, edoxaban, or rivaroxaban can be used in such cases.

Other options:

Option A: Absent p wave is an ECG finding seen in sinoatrial nodal disease and atrial fibrillation.

Option C: Cannon a wave is seen in atrioventricular dissociation and contraction of the right atria against a closed tricuspid valve.

Option D: Raised JVP with normal waveform is seen in conditions such as congestive heart failure, right ventricular infarction, and superior vena cava obstruction.

Solution to Question 13:

In the given scenario and image of a patient with abdominal distension, jaundice, pruritus, and neutrophil count >650 per mm3, the likely diagnosis is spontaneous bacterial peritonitis.

Spontaneous bacterial peritonitis (SBP) is an acute bacterial infection of the ascitic fluid. It is commonly seen in adults in association with liver cirrhosis. The pathogens involved are Escherichia coli, streptococci, enterococci, and pneumococci.

In a patient with ascites, SBP presents with fever, and acute abdominal pain. On examination, tachycardia, tachypnea, and signs of peritonitis are seen. Hepatic encephalopathy and gastrointestinal bleeding may also occur.

Diagnosis is by paracentesis, and an ascitic fluid neutrophil count of >250/mm3 is suggestive of spontaneous bacterial peritonitis. Ascitic fluid culture may reveal the pathogenic organisms.

Management is mainly by third-generation cephalosporins such as cefotaxime. Albumin infusion can be given as it decreases the risk of renal failure and improves patient survival. Early diagnosis and treatment improve the prognosis.

Other options:

Option B: Malignant ascites occurs as a complication of intrabdominal and extraabdominal malignancies. Patients can present with increasing abdominal girth, generalized abdominal pain, and shortness of breath. Weight loss may be more common. Ascitic fluid analysis reveals a milky or bloody appearance with elevated levels of tumor markers and VEGF (vascular endothelial growth factor). Treatment is by paracentesis and peritoneovenous shunt.

Option C: Ascites are often seen in patients with tubercular peritonitis. Multiple tubercles are seen on the peritoneum. Ascitic fluid analysis reveals a straw-colored fluid with a white blood cell count >500/mm3. Laparoscopy and peritoneal biopsy may be helpful in diagnosis. Management is by supportive and anti-tubercular therapy.

Option D: Intestinal obstruction can present as pain, abdominal distension, vomiting, and absolute constipation. Abdominal radiographs taken in a supine position are useful in diagnosis.

Abdominal radiograph in erect posture shows fluid levels. Management is mainly by insertion of a nasogastric tube, replacement of fluid and electrolytes, and surgery.

Solution to Question 14:

The given laboratory reports showing elevated alkaline phosphatase and bilirubin points to a likely diagnosis of obstructive jaundice.

Obstructive jaundice arises due to an obstruction in the biliary outflow due to choledocholithiasis, cholelithiasis, biliary strictures, etc. The patient presents with jaundice, right upper quadrant pain, pruritus, dark urine, and light-colored stools. The symptoms can be intermittent in cases of periampullary carcinoma and cholangiocarcinoma.

Elevated conjugated bilirubin and alkaline phosphatase (ALP) are characteristic of the condition. An associated rise in gamma-glutamyl transpeptidase (GGT) levels confirms the hepatic origin of ALP.

An ultrasound is done initially in suspected cases of obstructive jaundice. A CT or magnetic resonance cholangiopancreatography (MRCP) is recommended next. Endoscopic retrograde cholangiopancreatography (ERCP) is considered the gold standard for choledocholithiasis.

Jaundice arises due to the deposition of bilirubin in the body tissues. Bilirubin is a breakdown product of heme that circulates in the blood in the unconjugated/ indirect form. In the hepatocytes, it is conjugated with glucuronic acid to form conjugated/ direct bilirubin. Direct and indirect bilirubin can be assessed using Van den Bergh reaction.

Jaundice is broadly classified into:

• Pre-hepatic jaundice

• Hepatic jaundice

• Posthepatic jaundice

Other options:

Options B and D: Hemolytic jaundice is one of the common causes of prehepatic jaundice. These include sickle cell anemia, spherocytosis, and thalassemia. Inherited enzyme disorders, such as Gilbert's syndrome, and Crigler-Najjar types I and II, are other causes. In these cases, there is an isolated elevation of unconjugated bilirubin.

Option C: Hepatic jaundice shows an elevation of aspartate transaminase (AST) and alaline transaminase (ALT) out of proportion to ALP. An increased bilirubin level is also seen. Common causes include viral hepatitis, alcoholic liver disease, and autoimmune hepatitis. An AST/ALT ratio ≥ 2 is highly suggestive of alcoholic hepatitis.

Solution to Question 15:

The above clinical scenario points to a diagnosis of lupus nephritis.

Lupus nephritis is a common complication of systemic lupus erythematosus (SLE). It results from immune complex deposition containing DNA and anti-DNA. Clinical features include proteinuria, hematuria, and hypertension. The disease is divided into 6 classes based on the severity. Patients with class V disease are at an increased risk of thrombotic complications. Lupus nephritis can result in end-stage disease warranting dialysis or transplantation.

SLE is an autoimmune condition more commonly seen in women. Antinuclear antibodies, anti-dsDNA, and anti-Sm are the autoantibodies implicated. Symptoms include maculopapular butterfly rash, photosensitivity, oral or nasal ulcers, fever, and intermittent polyarthritis. Pleuritis and pericarditis can also be associated. Systemic glucocorticoids, cyclophosphamide, or mycophenolate mofetil are used for management. Due to the possible teratogenic effects of these drugs, hydroxychloroquine is the mainstay in pregnancy.

Other options:

Option A - Acute interstitial nephritis is a type of acute kidney injury. It is marked by oliguria and raised creatinine. Various causes of interstitial injury include the use of non-steroidal anti-inflammatory drugs (NSAIDs), pyelonephritis, leukemia, and sepsis. White blood cell (WBC) casts and granular casts may be seen.

Option B - Post-streptococcal glomerulonephritis usually affects male children between the age of 2-14. Skin or pharyngeal infection with M types of the organism is implicated. Symptoms include hematuria, pyuria, and hypertension. Flank pain and headache are also associated symptoms. Red blood cell (RBC) casts are seen in the urine.

Option D - IgA nephropathy arises due to mesangial IgA deposition. It is characterized by recurrent macroscopic hematuria after an upper respiratory infection associated with proteinuria. Mesangial proliferation is a distinctive feature. Angiotensin-converting enzyme (ACE) inhibitors and steroids can be used for management.

Solution to Question 16:

The given clinical scenario points to a diagnosis of acute rheumatic fever (ARF). The secondary prophylaxis after an episode for a patient weighing <27 kilograms is 6,00,000 IU of benzathine penicillin injection once every three weeks for 5 years or till the age of 18, whichever is longer.

ARF is an autoimmune disease that develops in response to group A streptococcal infection. It is commonly seen in children between the age of 5-14 years. An upper respiratory tract infection with some M serotypes of the organism is the trigger. The concept of molecular mimicry is used to explain the pathogenesis of the condition.

Since one attack of ARF increases the risk for further episodes and development of RHD, secondary prophylaxis is recommended by the American heart association (AHA) as follows:

• Rheumatic fever without carditis: For 5 years after the last attack or till 18/21 years of age (whichever is longer).

• Rheumatic fever with carditis but no residual valvular disease: For 10 years after the last attack or till 21/25 years of age (whichever is longer).

• Rheumatic fever with persistent valvular disease, evident clinically or on echocardiography: For 10 years after the last attack or till the age of 40 (whichever is longer). Sometimes lifelong

prophylaxis can be given.

Symptoms usually present after a latent period of about three weeks. Indolent carditis and chorea are the exceptions that can present even six months later. Migratory asymmetric polyarthritis which affects large joints and high-grade fever are common features. Cutaneous manifestations include erythema marginatum and subcutaneous nodules. Sydenham's chorea is commonly found in females and frequently affects the head and upper limbs. Acute rheumatic carditis can be seen as involvement of the endocardium, myocardium, or pericardium. Histopathological examination shows Aschoff bodies and activated macrophages called Anitschkow cells.

The image below shows Aschoff bodies.



In chronic rheumatic heart disease, there is a leaflet and tendinous cord thickening along with commissural fusion and shortening of the mitral valve. Ultimately the valve can undergo stenotic changes giving it a fish-mouth appearance as shown below.



The modified Jones criteria is used for diagnosis. Management of ARF is mainly symptomatic by the use of antibiotics and NSAIDs.

Solution to Question 17:

Based on the clinical scenario, the most likely diagnosis is hypokalemia.

Hypokalemia is defined as a plasma concentration of K+ <3.5 meq/L. Common causes include metabolic alkalosis, diarrhea, use of diuretics, starvation, and β2-agonists. It can also be seen after massive blood transfusion and in cystic fibrosis. It affects the cardiac, skeletal, and intestinal cells. It can lead to arrhythmias and intestinal ileus and increases the risk for rhabdomyolysis.

ECG changes include ST depression, prominent U waves, and prolonged repolarization. Pseudo P pulmonale waves can also be seen.





Oral potassium chloride is the mainstay in management. In patients unable to use the enteral route, intravenous potassium chloride can be given in saline.

Other options:

Option B: Hyperkalemia is a plasma K+ concentration >5.5 meq/L.Common causes include acidosis, the use of angiotensin-converting enzyme (ACE) inhibitors, and β2-antagonists. It can be seen as a manifestation of chronic kidney disease, and after a massive blood transfusion. It leads to increased aldosterone secretion. ECG changes include tall T waves, prolonged PR intervals, and small P waves. In severe cases, wide QRS complexes which progress to an undulating pattern (sine wave) and asystole can be seen.



Option C: Hypomagnesemia can arise due to prolonged use of proton pump inhibitors. It presents as delirium, neuromuscular dysfunction, reduced parathyroid function, and treatment-resistant hypokalemia. ECG might reveal hypokalemia potentiated by the condition. It has been implicated in ventricular arrhythmias with acute myocardial infarction and torsades de pointes. It can also enhance digitalis toxicity.

Option D: Hypernatremia is defined as plasma Na+ concentration >145 meq/L. Common causes include diarrhea, hyperglycemia, and post obstructive diuresis. Altered mental status and generalized weakness are the common manifestations. It can also cause neonatal seizures.

Solution to Question 18:

The arterial blood gas values suggest a likely diagnosis of respiratory acidosis.

The various buffering systems present in the body maintain the systemic arterial pH between 7.35-7.45. The most common acid-base disorders are metabolic acidosis or alkalosis and respiratory acidosis or alkalosis.

Normal values in an arterial blood gas are as follows:

• pH: 7.35-7.45

- PaCO2: 35-45 mmHg
- HCO3: 22-26 mEq/L
- PaO2: 75-100 mmHg

Respiratory changes (changes in PCO2) promote a corresponding change in metabolic response (changes in HCO3-) and vice versa as a compensatory mechanism. In general, PCO2 and HCO3- move in the same direction for compensation in metabolic and respiratory acid-base disorders respectively. A change in the two parameters in opposite directions points to a mixed acid-base disturbance.

Respiratory acidosis can be secondary to severe pulmonary disease, respiratory muscle fatigue, or ventilatory control abnormalities. It is marked by an increase in PaCO2 and a decrease in pH. A compensatory rise of HCO3- can also be seen.

A rapid rise in PCO2 (acute hypercapnia) can present with dyspnea, confusion, and psychosis. It can be seen in severe asthma secondary to bronchospasm, anaphylaxis, or inhalational burns. Chronic hypercapnia causes sleep disorders, memory loss, and daytime somnolence. It is commonly seen in end-stage obstructive lung disease.

Acute respiratory acidosis may need tracheal intubation and assisted mechanical ventilation. Rapid correction of hypercapnia should be avoided as it can trigger cardiac arrhythmias and seizures. Chronic respiratory acidosis is difficult to manage and improvement in lung function is the primary aim.

Other options:

Option A: Respiratory alkalosis arises due to alveolar hyperventilation and can be due to salicylates, high altitude states, cardiac failure, and anemia.

Option B: Metabolic acidosis arises due to an accumulation of endogenous acids. Examples include lactic acidosis, diabetic ketoacidosis, and starvation. It can also be seen in renal failure.

Option D: Metabolic alkalosis can be seen in Liddle's syndrome, hypertrophic pyloric stenosis, massive blood transfusion, and cystic fibrosis. It is marked by elevated pH and HCO3- values. A rise in PaCO2 indicates a compensatory respiratory response.



Solution to Question 19:

Parathyroid adenoma is not seen in multiple endocrine neoplasia 2B (MEN 2B) syndrome.

Multiple endocrine neoplasia (MEN) describes conditions with tumors involving two or more endocrine glands. There are four types - MEN 1-4. They have an autosomal dominant pattern of inheritance and can also be sporadic.

MEN type 2 comprises groups of disorders caused by gain-of-function mutations of the RET proto-oncogene at chromosomal locus 10q11.2. There are three variants:

• MEN 2A, also known as MEN 2 (most common type)

• MEN 2B, also known as MEN 3

• MTC only

MEN 2A is also known as Sipple's syndrome. It is characterized by medullary thyroid carcinoma (MTC), pheochromocytomas, and parathyroid tumors.

In MEN 2B along with MTC and pheochromocytoma, it is associated with other abnormalities such as mucosal neuromas, marfanoid habitus, medullated corneal nerve fibers, and megacolon. Parathyroid tumors are generally not seen here, in contrast to the MEN 2A type.

MTC presents early and is usually aggressive. MTC is the most common feature, presents as a palpable mass in the neck, and can be associated with pressure symptoms. Ectopic adrenocorticotropic hormone (ACTH) secreted by the tumor can cause Cushing's syndrome.

Pheochromocytoma is frequently bilateral. Palpitations, sweating, and headaches are the features of pheochromocytoma.

Mucosal neuromas can present with bumpy and enlarged lips, tongue, and eyelids. Intestinal autonomic ganglion dysfunction can lead to megacolon and diverticula. Medullated corneal fibers and delayed puberty can also be seen. Marfanoid habitus marked by tall stature, long limbs, arachnodactyly, and joint hypermobility is an associated feature.

Solution to Question 20:

The clinical scenario along with the ECG points to a diagnosis of ventricular tachycardia. Intravenous amiodarone (class III antiarrhythmic) can be used in the management of this condition.

The patient here complaining of chest pain, and palpitations and has ECG changes showing wide QRS complex tachycardia with absent p waves is likely to have acute ventricular tachycardia. Since the patient is stable and has an underlying coronary artery disease the drug of choice is intravenous amiodarone.

Ventricular tachycardias can be defined as three or more pre-ventricular contractions at a rate >100/minute.

ECG changes include wide QRS complexes, absence of P waves, and ventriculoatrial dissociation. It can be broadly classified as sustained VT or non-sustained VT. Nonsustained VT has QRS complexes lasting <30 seconds while sustained VT lasts for >30 seconds. In monomorphic VT, all QRS complexes look the same. Polymorphic VT has QRS complexes varying in shape or direction in the same lead.

Most monomorphic VT have an underlying cardiac disease. Symptoms of ventricular tachycardia include palpitations, syncope, and shortness of breath.

In a hemodynamically stable patient, amiodarone, procainamide, and lidocaine can be used to treat the condition. In the presence of heart disease, intravenous amiodarone is the drug of choice. In hemodynamically unstable patients, electrical cardioversion is done. Further management depends on the underlying heart condition and frequency of VT. Frequent episodes warrant the use of antiarrhythmics or catheter ablation.

Torsades de Pointes is a type of polymorphic VT occurring with QT prolongation. It can progress to ventricular fibrillation which is treated with defibrillation. Recurrent episodes can be managed with intravenous magnesium sulfate.

Other causes of wide QRS complexes include supraventricular tachycardias with bundle branch block and Wolff-Parkinson-White syndrome.

Other options:

Options A and D: Metoprolol is a selective β1-antagonist and a class II antiarrhythmic. It can be used in hypertension, angina, and heart failure. The intravenous form can be used for initial management in acute myocardial infarction.

Option B: Oral amiodarone can be used in the chronic treatment of atrial fibrillation, ventricular tachycardia, and ventricular fibrillation. It has shown a modest beneficial effect on mortality after acute myocardial infarction.

Solution to Question 21:

Muti-drug resistant (MDR) tuberculosis (TB) shows resistance to isoniazid and rifampicin.

MDR-TB poses a serious problem in the treatment of TB as isoniazid and rifampicin are the most potent first-line drugs.

The treatment of MDR-TB is curated with drugs grouped into different classes by the world health organization (WHO).

• Group A: Levofloxacin, bedaquiline, linezolid

• Group B: Clofazimine, cycloserine

• Group C: Ethambutol, pyrazinamide, delamanid, ethionamide

In patients treated previously or with additional drug resistance, a longer MDR-TB regimen of 18-20 months is recommended. Drugs belonging to group A are preferred along with one drug from group B whenever possible. A shorter regimen of 6 months including bedaquiline with other drugs can be used in patients not treated before and without extensive disease. In pregnant women with MRD-TB, drugs such as bedaquiline and delamanid should be avoided.

The nucleic acid amplification test (NAAT)/ gene Xpert is used to detect rpoB gene polymorphisms, which are responsible for rifampicin resistance.

The four drugs considered first-line agents in the treatment of TB are isoniazid, rifampicin, pyrazinamide, and ethambutol.

Isoniazid is an antibiotic that works by inhibiting the synthesis of mycolic acid, an essential component of the mycobacterial cell wall. It is a bactericidal drug. It acts as a vitamin B6 antagonist, so oral pyridoxine supplementation is recommended during treatment. Side effects include peripheral neuritis, seizures, and ataxia.

Rifampicin is a macrocyclic antibiotic that inhibits RNA synthesis and is bactericidal. It is also used in the prophylaxis of meningococcal infections. Side effects include rash, vomiting, and hepatitis. It is a microsomal enzyme inducer that can result in contraceptive failure.

Ethambutol inhibits the mycobacterial cell wall assembly. Side effects include optic neuritis, pruritus, and joint pain.

Pyrazinamide is a nicotinamide analogue activated in acidic conditions. Side effects include liver injury and hyperuricemia.

Other options: Fluoroquinolones are considered second-line agents used in the treatment for drug-resistant TB. Kanamycin, an injectable aminoglycoside is excluded from the new WHO classification of second-line drugs.

Solution to Question 22:

The given clinical scenario along with the serology reports suggest a diagnosis of acute hepatitis B.

Acute hepatitis B presents after an incubation period of 30-180 days with symptoms such as anorexia, malaise, and myalgias. A low-grade fever can also be associated. These symptoms can precede jaundice by 1-2 weeks. Mild weight loss and tender hepatomegaly are other features.

The presence of HBsAg (hepatitis B surface antigen), also known as Australia antigen, IgM anti-HBc antibodies (antibodies against hepatitis B core antigen), and the absence of anti-HBs is the characteristic picture seen in acute hepatitis B with high infectivity. The absence of anti-HCV

antibodies excludes a diagnosis of hepatitis C.



Aspartate transaminase (AST) and alanine transaminase (ALT) values are usually elevated along with bilirubin levels. Serological markers vary according to the phase of the disease process as shown below.

Most cases undergo spontaneous recovery. Prophylaxis can be carried out with the use of vaccines containing HbsAg. Anti-HBs titres ≥ 10mIU/mL are protective.

Acute viral hepatitis can be due to due to any of the five agents - hepatitis A virus (HAV), hepatitis B virus (HBV), hepatitis C virus (HCV), hepatitis D virus (HDV), and hepatitis E virus (HEV). Clinical features of all acute viral hepatitis are similar. A patient suspected to have acute hepatitis should be tested for HBsAg, IgM anti-HAV, IgM anti-HBc, and anti-HCV.

Other options:

Option B: Acute hepatitis C is caused by HCV which is an RNA virus. It is a blood-borne infection. Detection of anti-HCV and HCV RNA is used for diagnosis. HCV-RNA appears before anti-HCV and persists for the entire duration of the disease.

Option C: Chronic hepatitis B is a late complication of acute hepatitis B. Persistence of HBeAg for >3 months or HBsAg for >6 months is suggestive of the development of the condition. Investigations show the presence of HBsAg, IgG anti-HBc, HBeAg, and HBV DNA. Fatigue and intermittent jaundice are the main features. It can progress to cirrhosis or hepatocellular carcinoma. Hepatitis D superinfection can accelerate this progression.

Option D: Chronic hepatitis C is a common complication of acute hepatitis C. It can progress to cirrhosis or hepatocellular carcinoma HCV-RNA and anti-HCV can be detected on serology.

Solution to Question 23:

The clinical scenario points to iron toxicity, which is managed with deferoxamine (also known as desferrioxamine).

Large amounts of ferrous salts can be fatal in children. Initial symptoms include abdominal pain, diarrhea, vomiting, and gastrointestinal bleeding. In severe cases, cyanosis and hyperventilation due to acidosis can develop. This can sometimes be followed by an asymptomatic phase. Later, hypoglycemia and hepatic and renal failure can develop. Pyloric stenosis and gastric scarring can develop due to its corrosive nature. Ultimately, if left untreated, the child can die.

The gastric contents can be used for an iron color test to estimate the iron concentration in plasma. As iron tablets are radiopaque, an X-ray can be used to assess the number of pills taken. Deferoxamine, an iron chelator has to be administered intramuscularly. Side effects include pruritus, wheals, rash, and anaphylaxis.

Deferiprone and deferasirox are other oral iron chelators used to treat iron overload in thalassemia patients.

Other options:

Option B: Activated charcoal can be used in the initial management of acetaminophen and opioid overdose. It is used for gastrointestinal decontamination to prevent further absorption of the poison.

Option C: Dimercaprol, also known as BAL (British anti-Lewisite), is used to treat arsenic, mercury poisoning, and Wilson's disease.

Option D: D-penicillamine (Cuprimine) can be used to treat scleroderma, Wilson's disease, and rheumatoid arthritis. Side effects include nephrotoxicity, hematologic derangements, and a characteristic rash that involves the neck and axillae—elastosis perforans serpiginosa. It is associated with membranous glomerulonephritis, drug-induced lupus, and myasthenia gravis.

Medicine NEET 2023

Question 1:

Identify the condition associated with the murmur depicted in the image below.



a) Mitral regurgitation

b) Aortic regurgitation

c) Aortic stenosis

d) Mitral stenosis

Question 2:

A young man came to the medical OPD with complaints of early morning backache and stiffness, which improves on exercise, and persistent red eyes. On examination, lung expansion was less than 3 cm. X-ray is shown in the image given below. What is the most probable diagnosis?

a) Ankylosing spondylitis

b) Paget's disease

c) Healed tuberculosis

d) Osteopetrosis

Question 3:

A patient with hyperkalemia and elevated urea levels underwent dialysis. Towards the end of the session, she became drowsy and had a sudden seizure episode. On examination, the patient was hypotensive. What is the treatment for this condition?

a) Bumetanide

b) Ethacrynic acid

c) Nesiritide

d) IV Mannitol

Question 4:

A female patient presents with severe restlessness, palpitations, and tremors to the emergency department. She is a known case of bronchial asthma. On examination, the neck looks swollen. Blood pressure is elevated, and tachycardia is noted. ECG shows atrial fibrillation. Which of the following drugs is used for immediate management in this patient?

a) Diltiazem

b) Propranolol

c) Esmolol

d) Propylthiouracil

Question 5:

A 30-year-old male is found to be positive for HBsAg and HBeAg and is diagnosed with chronic hepatitis B. The viral load of the patient was 2 x 100000 and SGPT is found to be doubled. What is the appropriate management in this patient?

a) Lamivudine for 30+ weeks

b) Tenofovir for > 40 weeks

c) Pegylated interferon for 52 weeks

d) Combined pegylated interferon with lamivudine

Question 6:

A 25-year-old male patient came with complaints of palpitations, sweating, and restlessness. He has a complaint of sweaty palms. Clinical findings are depicted in the image given below. What is the diagnostic finding seen in this patient?

a) Anti-thyroglobulin antibody

b) Anti-thyroid peroxidase antibody

c) Thyroid receptor antibody

d) Elevated ultrasensitive thyrotropin levels

Question 7:

A 50-year-old HIV patient presented with a painful lesion, as shown in the image. What is the most likely diagnosis?



a) Basal cell carcinoma

b) Kaposi sarcoma

c) Malignant melanoma

d) Squamous cell carcinoma

Question 8:

A patient is brought to the OPD by his wife complaining about difficulty in expressing emotions and not taking part in daily activities. On examination, resting tremors and rigidity are noted. Given the possible diagnosis, which part of the brain is affected in this patient?

a) Basal ganglia

b) Hippocampus

c) Cerebellum

d) Premotor cortex

Question 9:

A 45-year-old HIV-positive male came with complaints of persistent cough and weight loss. He has skin lesions that appear as umbilicated papules and nodules predominantly on the

face, trunk, and upper extremities. Chest x-ray showed multiple bilateral nodular infiltrates. His sputum CBNAAT for tuberculosis was negative and he has a low CD4 count. What is the probable diagnosis?

a) HIV with disseminated histoplasmosis

b) HIV with disseminated cryptococcosis

c) HIV with molluscum contagiosum

d) HIV with tuberculosis

Question 10:

A male patient presented to the emergency room with seizures. He has a history of fever, headache, and confusion. An MRI brain was done, and it showed inflammation involving the bitemporal lobe. What is the most likely etiology for this presentation?

a) Cytomegalovirus

b) Toxoplasma gondii

c) Herpes simplex virus

d) Mycobacterium tuberculosis

Question 11:

A male patient presents with sensory loss and weakness of limbs for 3 months. He also has angular stomatitis. On examination, there is loss of proprioception, vibration sensations, UMN type of lower limb weakness, and absent ankle reflex. What is the most probable diagnosis?

a) Extradural cord compression

b) Amyotrophic lateral sclerosis

c) Multiple sclerosis

d) Subacute combined degeneration of cord

Question 12:

A 40-year-old female patient came with complaints of chest pain, palpitation, and shortness of breath. On examination, a mid-diastolic murmur was heard, and a prominent 'a' wave was found on JVP. What is the most likely diagnosis?

a) Mitral stenosis

b) Tricuspid stenosis

c) Mitral regurgitation

d) Tricuspid regurgitation

Question 13:

A patient who is a known case of hypertension on multiple anti-hypertensive medications came to OPD. His ECG finding is given below. Which of the following drugs is responsible for the ECG finding?



a) Prazosin

b) Metoprolol

c) Hydrochlorothiazide

d) Spironolactone

Question 14:

A 78-year-old woman presents with a progressive decline in daily activity. She gives a history of convulsions and visual hallucinations. She does not talk to anyone and keeps looking at the sky. Pathological examination shows the presence of Lewy bodies within the neurons. What is the most probable diagnosis?

a) Parkinson's disease

b) Prion disease

c) Huntington's chorea

d) Alzheimer's disease

Question 15:

A patient came to medical OPD with complaints of polyuria. He has a history of undergoing total hypophysectomy. His Na+ levels are found to be 155 mEq/L, urine osmolarity was 200 mOsm/L. What is the definitive management in this patient?

a) DDAVP for 2 weeks and then discontinue

b) DDAVP supplementation for lifelong

c) Upsetting of receptors so no treatment is required

d) Thiazides for 2 weeks

Question 16:

A 65-year-old chronic smoker came to the medicine outpatient department with complaints of upper chest discomfort and drooping of an eyelid. He also complained of pain radiating to the upper arm and a tingling sensation in the 4th and 5th digits of his right hand. The chest X-ray is given below. Which of the following is the most likely diagnosis?



a) Pancoast tumor

b) Upper lobe pneumonia

c) Superior vena cava obstruction

d) Aspergilloma

Question 17:

A chronic alcoholic patient presents with acute pain and swelling of the left great toe. There is no history of trauma. Synovial fluid analysis shows raised leukocytes. Lab investigations show normal serum uric acid levels. What is the most likely diagnosis?

a) Pseudogout

b) Acute gout

c) Reactive arthritis

d) Septic arthritis

Answer Key

Question No.	Correct Option
1	a
2	a
3	d
4	a
5	b
6	c
7	b
8	a
9	a
10	c
11	d
12	b
13	d
14	a
15	b
16	a
17	b

Detailed Explanations

Solution to Question 1:

The given image shows pansystolic murmur, and it is associated with mitral regurgitation.

Mitral regurgitation may result from an abnormality or disease process that affects functional components of the mitral valve apparatus.

Causes of mitral regurgitation include

• Acute: Blunt trauma, papillary muscle rupture, infective endocarditis

• Chronic: Mitral valve prolapse, rheumatic fever, healed infective endocarditis, ischemic cardiomyopathy, HOCM, etc.

A pansystolic murmur of at least grade III/VI intensity can be auscultated in chronic severe mitral regurgitation. It is heard best at the apex and radiates to the axilla. Ruptured chordae tendineae may produce a murmur with a cooing or "seagull" quality, whereas a flail leaflet may produce a murmur with a musical quality.

S1 is generally absent or soft. Wide splitting of S2 may occur due to premature closure of the aortic valve. A low-pitched third heart sound (S3) may occur following A2. S4 can also be audible in patients with acute severe MR.

The image below depicts the pansystolic murmur in mitral regurgitation.



Other options:

Option B: An early diastolic murmur can be auscultated in chronic aortic regurgitation. It is high-pitched and blowing in character. It is best heard in the third intercostal space along the left sternal edge. It may be accompanied by a mid-systolic ejection murmur in isolated AR or the Austin Flint murmur in severe AR. The latter is a soft, low-pitched, rumbling, mid-to-late diastolic murmur produced by the displacement of the anterior leaflet of the mitral valve by the AR stream.

A2 is usually absent, and occasionally, S4 also may be heard.



Option C: In aortic stenosis, an ejection mid-systolic murmur can be auscultated. It begins after the first heart sound (S1), increases in intensity toward the middle of the ejection, and ends just before aortic valve closure. It is low-pitched, rough, and rasping in character. It is best heard at the base of the heart, most commonly in the second right intercostal space.

A systolic ejection click can be heard in young patients with a bicuspid aortic valve, but disappears as the valves become calcified. A2 and P2 may be heard at the same time, or A2 may follow P2, resulting in the paradoxical splitting of the second heart sound (S2). The fourth heart sound (S4) is often heard at the apex, indicating the presence of left ventricular hypertrophy.



Option D: In mitral stenosis, a diastolic murmur can be heard during auscultation.



Solution to Question 2:

The given clinical vignette points towards the diagnosis of ankylosing spondylitis.

Ankylosing spondylitis is a chronic inflammatory disease affecting the spine and sacroiliac joint. It is more common in young males and has a strong association with HLAB27.

Clinically, patients present with recurring backache which worsens in the early morning or with inactivity and improves with exercise. Later, they develop fatigue, pain, joint swellings, and heel pain. Extra-articular manifestations include anterior uveitis, aortic incompetence, carditis, and apical lobe fibrosis. Examination findings include loss of lumbar lordosis, increased thoracic kyphosis, diminished spinal movements, and decreased chest expansion.

X-ray findings:

• Erosion of sacroiliac joints - earliest sign

• Periarticular sclerosis in later stages

• Squaring of vertebrae due to the flattening of normal anterior concavity of the vertebral body is the earliest vertebral change

• Bamboo spine appearance due to bridging of adjacent vertebrae

• Wedging of the vertebra, leading to hyperkyphosis

Complications of ankylosing spondylitis include

• Spinal fractures, the most common site is C5–7

• Hyperkyphosis

• Spinal cord compression

• Lumbosacral nerve root compression

Treatment consists of maintaining posture and preserving mobility. Pain and stiffness can be controlled with anti-inflammatory drugs. In severe cases, TNF inhibitors are used. Surgery can be considered to correct the deformity.

The image below shows the bamboo spine's appearance:



Solution to Question 3:

The given scenario is suggestive of the dialysis disequilibrium syndrome (DDS) or the reverse urea effect. It is an acute complication that occurs especially in patients undergoing dialysis for the first time.

During dialysis, urea in blood vessels is rapidly removed as compared to that in the interstitium. This decreases the osmolality of blood. This results in a fluid shift from the blood vessels into the interstitium and brain parenchyma, causing cerebral edema.

DDS can present as headache, nausea, blurred vision, restlessness, confusion, and severe manifestations like coma and seizures in rare cases.

The most common risk factors include

• First hemodialysis treatment

• High blood urea nitrogen (BUN) (above 175 mg/dL or 60 mmol/L) before initiation of dialysis

• Children and elderly

• Sudden change in dialysis regimen

• Hyponatremia, hepatic encephalopathy

• Sepsis, meningitis, encephalitis, hemolytic uremic syndrome, vasculitis

•

History of stroke, malignant hypertension, head trauma, or seizure disorder

Management includes changing the sodium dialysate bath or engaging the changed prescription on the dialysis machine. Intravenous mannitol can be used to reduce intracerebral pressure in severe symptoms of DDS.

DDS can be prevented by

• Slow and gentle initial hemodialysis

• Limiting the clearance of urea to prevent the development of an osmotic gradient

• Increasing dialysate sodium level

• Administration of osmotically active substance

Solution to Question 4:

The given clinical scenario suggests the diagnosis of thyroid storm/thyroid crisis presenting with atrial fibrillation, and the immediate management in this patient is to rate control the arrhythmia with diltiazem (calcium channel blocker).

Thyroid storm/Thyrotoxic crisis is a rare but severe life-threatening complication of hyperthyroidism induced by the release of excessive thyroid hormones.

It is caused by abrupt cessation of antithyroid medications, infection, inadequately prepared hyperthyroid patient for thyroidectomy, amiodarone administration, exposure to iodinated contrast agents, and following radioactive iodine (RAI) therapy.

Features:

• Fever and severe dehydration

• Cardiovascular: Circulatory collapse, palpitations, tachycardia, cardiac arrhythmias, and cardiac failure

• GI symptoms: Vomiting, diarrhea, jaundice

• Neurological symptoms: Restlessness, irritability, delirium, tremor, convulsions, and coma

Treatment:

• If the patient presents with atrial fibrillation, the first step in the management is to do rate control using beta-blockers, calcium channel blockers, digoxin, or amiodarone. Since this patient is a known asthmatic, beta-blockers (propranolol and esmolol) are contraindicated. So, a calcium channel blocker (diltiazem) is the preferred drug for immediate management in this patient.

• Supportive measures – rehydration by proper fluid therapy, tepid sponging, and cooling blankets.

• Thionamides like propylthiouracil (PTU) or methimazole or carbimazole can be used. PTU (Option D) is the drug of choice in thyroid storm, but it is given only after controlling atrial fibrillation.

• Lugol's iodide is given one hour after intake of thionamides. They inhibit thyroid hormone synthesis by Wolff- Chaikoff effect.

• Glucocorticoids – reduce the peripheral conversion of T4 to T3 and controls the shock.

• Digoxin and cardiac monitoring are crucial.

Other options:

Options B & C: Beta-blockers like propranolol and esmolol are contraindicated in bronchial asthma patients.

Solution to Question 5:

The appropriate treatment for this patient based on the given clinical scenario is to administer tenofovir for more than 40 weeks.

The serological markers present in the case of chronic hepatitis B are

• HBsAg (Remains detectable beyond six months)

• Anti-HBc (Primarily of IgG class)

• HBeAg (Qualitative marker of replicative phase)

• HBV DNA (Quantitative marker of replicative phase)

Treatment of chronic hepatitis B is focused on suppressing the level of virus replication.

Tenofovir disoproxil fumarate (TDF) is an acyclic nucleotide analog that is used as first-line therapy for the treatment of chronic hepatitis B and lamivudine-resistant chronic hepatitis B. It is a potent antiretroviral agent used in the treatment of HIV infection. It is more potent in suppressing HBV DNA and is active against both wild-type and lamivudine-resistant HBV. It is given as an oral once-daily dose of 300 mg for 48 weeks. Tenofovir therapy provided histologic improvement, reduction in fibrosis score, and regression of cirrhosis in patients. Side effects are acute renal failure and low blood phosphate levels.

Other options:

Option A: Lamivudine is a nucleoside analog that inhibits reverse transcriptase activity of both HIV and HBV and is an effective agent for chronic hepatitis B. It is given at a daily dose of 100 mg for 48-52 weeks. It is no longer preferred.

Option C: Pegylated interferon is recommended as one of the first-line agents for hepatitis B. It is given as a subcutaneous injection at a dose of 180 μg/week for 48-52 weeks. It is not preferred over oral agents because of its poor tolerability.

Option D: The combination of lamivudine and pegylated interferon is found to suppress HBV DNA more profoundly during therapy than monotherapy. However, usage of these drugs for a year provides results similar to drugs given as monotherapy. Hence, this options is not preferred.

Solution to Question 6:

Based on the clinical findings the most likely diagnosis is Graves' disease. Elevated levels of thyroid receptor–stimulating antibodies are diagnostic of Graves' disease.

The hyperthyroidism of Graves' disease is caused by thyroid-stimulating immunoglobulins (TSI) that are synthesized in the thyroid gland as well as in bone marrow and lymph nodes. TSI cause stimulation of the thyroid gland causing features of hyperthyroidism in the presence of low TSH levels.

Clinical features of Graves' disease:

• Hyperthyroid features like hyperactivity, irritability, dysphoria, heat intolerance and sweating palpitations, fatigue, and weakness, weight loss with increased appetite, diarrhea, polyuria, oligomenorrhea, loss of libido, tachycardia

• Eye signs as mentioned in Graves' ophthalmopathy.

• Graves ophthalmopathy: It is due to increased volume of the retroorbital connective tissues and extraocular muscles because of marked lymphocytic infiltration of the retro-orbital space, accumulation of hydrophilic glycosaminoglycans such as hyaluronic acid and chondroitin sulfate, and increased numbers of adipocytes (fatty infiltration). Corneal exposure, lid retraction, lid lag on downgaze, conjunctival injection, restriction of gaze, diplopia, and visual loss from optic nerve compression are cardinal symptoms.

Histopathology shows follicular hyperplasia with scalloped margins on biopsy.

Management involves

• Propylthiouracil is the mainstay of treatment for Graves' disease but it does not affect Graves' ophthalmopathy.

• Radioactive Iodine is also used as a treatment modality for hyperthyroidism in suitable patients but does not affect Grave's ophthalmopathy.

Other options:

Levels of anti-thyroid peroxidase (Option B) and anti-thyroglobulin (Option A) are elevated in autoimmune hypothyroidism which is characterized by features of dry skin, cold intolerance, tiredness, weakness, feeling tired, weight gain, etc.

Option D: Ultrasensitive thyrotropin level (TSH) is elevated in primary hypothyroidism.

Solution to Question 7:

The above image showing a violaceous growth on the hard palate in an HIV patient is diagnostic of Kaposi sarcoma.

Kaposi sarcoma (KS) is a vascular neoplasm caused by human herpesvirus 8 (HHV8) and is very common in patients suffering from acquired immunodeficiency syndrome (AIDS). KS is considered to be an AIDS-defining disease.

AIDS-associated KS is an AIDS-defining illness and is the most common malignancy associated with HIV infection. Involvement of the oral cavity is most common in AIDS-related KS, but it can be seen in other variants also. Oral KS (OKS) most often involves the hard and soft palate, gingiva, and dorsal tongue with plaques or tumors. The color of the lesions may range from non-pigmented to brownish-red or violaceous.

Other forms of KS include

• Classic Kaposi sarcoma: It has the following three clinical stages patches, plaques, and nodules.

• Endemic African KS

• Transplant-associated KS is more common after solid organ transplant patients receive immunosuppression as compared to the marrow or stem-cell transplant patients.

The given images show coalescent red-purple macules and plaques of the skin and the histologic appearance of Kaposi sarcoma, demonstrating sheets of proliferating spindle cells:



There is no definitive cure but KS is highly radiosensitive and achieves complete response in most cases. In AIDS■associated KS, instituting highly active antiretroviral therapy (HAART) often (but not always) results in regression of KS. In localized cutaneous KS, excision and cryotherapy can be used. Intra-lesional administration of vinblastine, interferon■α2b, and imiquimod is also effective.

Other options:

Option A: Basal cell carcinoma (BCC) most commonly tends to occur in sun-exposed areas of the face like the nose, eyelid, and other parts of the face. BCC can vary from red, flesh-colored, or white macules or papules, to nodules and ulcerated lesions. The nodular variant is the most common subtype characterized by raised, pearly pink papules with telangiectasias and occasionally a depressed tumor center with raised borders giving the classic “rodent ulcer” appearance.

The image below shows a case of basal cell carcinoma.

Option C: Malignant melanoma primarily arises from melanocytes at the epidermal-dermal junction but may also originate from mucosal surfaces of the oropharynx, nasopharynx, eyes, proximal esophagus, anorectal, and female genitalia. Lentigo maligna, superficial spreading, acral lentiginous, and nodular are the variants of malignant melanoma. It includes pigmented lesion that has enlarged, ulcerated, or bled. Amelanotic lesions appear as raised pink, purple, or flesh-colored skin papules. The diagnostic clue for melanoma is the ABCDE criteria.

It comprises of

• Asymmetry

• Border irregularity

• Color variegation

• Diameter of more than 6 mm

• Evolution or change

The image below shows a case of malignant melanoma.

Option D: Squamous cell carcinoma (SCC) is the second most common skin cancer. UV radiation exposure is the primary risk factor for SCC. Classically, it appears as a scaly or ulcerated papule or plaque that can bleed with minimal trauma, but the pain is rare. It can exhibit in situ (confined to the epidermis) or invasive subtypes. Wide surgical excision including subcutaneous fat is the treatment of choice for SCC.

Solution to Question 8:

Based on the clinical findings given above the most likely diagnosis is Parkinson's disease and it is due to degeneration of the substantia nigra in the basal ganglia.

Parkinson’s disease, also known as paralysis agitans, results from widespread degeneration of the portion of the substantia nigra that sends dopamine-secreting nerve fibers to the caudate nucleus and putamen.

It is characterized by rigidity, resting tremors, akinesia, bradykinesia, postural instability, dysphagia, speech disorders, gait disturbances, and fatigue. Other non-motor features include anosmia, sensory disturbances, mood disorders (e.g., depression), sleep disturbances, autonomic disturbances, and cognitive impairment/dementia.

Other options:

Option B: Lesions of the hippocampus mainly affect learning, memory, emotional behavior, regulation of hypothalamic functions, and spatial navigation.

Option C: Features of cerebellar lesions are ataxia, intention tremors, nystagmus, speech defect (slurring speech), hypotonia or atonia, dysmetria, dysdiadochokinesia (failure to perform rapid alternating movements), decomposition of movement (lack of smooth flow of movement), rebound phenomenon

Option D: Premotor cortex is concerned with setting posture at the start of a planned movement and with getting the individual prepared to move. It is most involved in the control of proximal limb muscles needed to orient the body for movement.

Solution to Question 9:

The diagnosis based on the given clinical scenario is HIV with disseminated histoplasmosis.

Progressive disseminated histoplasmosis (PDH) is seen in immunocompromised patients. Common risk factors are AIDS (CD4+ T-cell count, <200/μL), extremes of age, the administration of immunosuppressive drugs (e.g., prednisone, mycophenolate, calcineurin inhibitors), and use of drugs like methotrexate, anti-TNF-α agents, and other biologic response modifiers for autoimmune disorders. It most commonly involves the lungs and bone marrow.

Patients with PDH can present with diffuse interstitial or reticulonodular lung infiltrates that can cause respiratory failure, shock, coagulopathy, and multiorgan failure. Clinical features are fever, weight loss, hepatosplenomegaly, and thrombocytopenia. They may also present with meningitis or focal brain lesions, ulcerations of the oral mucosa, gastrointestinal ulcerations and bleeding, and adrenal insufficiency.

Cutaneous involvement is noted in about 10% of cases of PDH. Skin lesions can be caused by the fungus itself resulting in the formation of papules, plaques, nodules, or ulcers, or by an immune response to the infection causing erythema nodosum or erythema multiforme.

Laboratory investigations include looking for Histoplasma antigen in bronchoalveolar lavage (BAL) fluid, serum, and urine, IgG and IgM antibodies in serology, a fungal culture of blood or bone marrow aspirate, and cytopathology on biopsy of the affected organ.

Histoplasmosis caused by Histoplasma capsulatum is a thermally dimorphic fungus. The yeast form occurs mainly in tissues at a temperature of 37 °C and the mycelial form occurs in the soil at 25 °C. At room temperature, white, cottony, mycelial growth appears, with large (8-20 μ) thick-walled, spherical spores with tubercles (finger-like projections called tuberculate spores). Transmission is mainly via inhalation of microconidia.

Treatment involves liposomal amphotericin B, followed by itraconazole for at least 12 months.

Other options:

Option B: If the disease involves skin with signs of meningitis, then HIV with disseminated cryptococcosis can be considered. Since the patient has no signs of meningitis, this option can be ruled out.

Option C: If the disease is restricted to the skin alone, it can be HIV with disseminated molluscum contagiosum. Since the question mentions lung involvement, the option is ruled out.

Option D: The question mentions that CBNAAT for tuberculosis was negative. Hence, this option is ruled out.

Solution to Question 10:

The given clinical vignette suggests the diagnosis of encephalitis caused by the herpes simplex virus (HSV).

HSV encephalitis is defined as inflammation of brain parenchyma caused by the Herpes virus and it typically presents as an acute febrile illness.

Patients typically present with acute onset of fever, altered levels of consciousness, personality changes, and evidence of either focal or diffuse neurologic signs and symptoms, especially in the temporal lobe. Focal or generalized seizures occur in many patients with HSV encephalitis. Other most commonly encountered focal findings include aphasia, ataxia, upper motor neuron or lower motor neuron type lesions, and cranial nerve deficits.

HSV-DNA demonstration in CSF by PCR is the investigation of choice for HSV encephalitis. Other investigations that can help include CSF examination, CSF culture, serological studies, MRI, CT, EEG, and brain biopsy.

The T2-weighted MRI of the brain will show hyperintensity in the temporal lobe in most of the patients. Bilateral involvement of the temporal lobe is more common in HSV encephalitis.

The image below is a coronal T2-weighted MR image showing a high signal intensity in the temporal lobes in a patient with HSV encephalitis.



Management includes supportive therapy(monitoring of intracranial pressure, fluid restriction, avoidance of hypotonic IV fluids), anticonvulsants for seizures, and intravenous acyclovir is beneficial in the treatment of HSV encephalitis.

Other options:

Option A: Cytomegalovirus (CMV) encephalitis in adult immunocompromised adult patients often have enlarged ventricles with areas of increased T2 signal on MRI outlining the ventricles (periventricular calcifications).

Option B: In immunocompetent patients, acute toxoplasmosis is usually asymptomatic and self-limited. Immunocompromised patients with Toxoplasma encephalopathy usually have multiple lesions located in both hemispheres, with the basal ganglia and corticomedullary junction most commonly involved in the MRI.

Option D: Mycobacterium tuberculosis causes tuberculous (TB) encephalopathy and it is exclusively seen in children and infants with pulmonary TB. It is characterized by cerebral edema sometimes with features similar to acute disseminated encephalopathy and may manifest with a variety of symptoms ranging from focal neurological deficits to convulsions and decreased conscious state.

Solution to Question 11:

The given case scenario is suggestive of subacute combined degeneration (SACD) due to vitamin B12 deficiency.

Subacute combined degeneration (SACD) is due to the deficiency of vitamin B12. Pernicious anemia, partial gastrectomy, and Diphyllobothrium latum infestation can lead to this condition. The most severely affected regions in SACD are the posterior columns at the cervical and upper thoracic levels.

Patients with SACD present with the following features:

• Loss of vibration sense and proprioception

• Paresthesia - upper limb paresthesias occur before lower limbs are affected.

• Unsteady gait - due to sensory ataxia

• Loss of deep tendon reflexes

• Peripheral neuropathy

• Optic neuropathy - can present as blurring of vision

• Psychiatric manifestations such as:

• Irritated behavior

• Severe dementia

• Frank psychosis

MRI spine shows myelopathy and areas of demyelination. Myelopathy of SACD tends to be diffuse rather than focal.

Treatment is replacement therapy with 1000 μg of intramuscular vitamin B12 repeated at regular intervals or by oral vitamin B12.

Other options:

Option A: Patients with extradural spinal cord compression usually present with back pain that is aggravated in the supine position, tingling, or electric sensation along the back and upper and lower limbs upon flexion or extending the neck. Loss of pain, vibration, and proprioception is common. Extensor plantar reflex may be seen in significant compression. Deep tendon reflexes may be brisk.

Option B: Patients with amyotrophic lateral sclerosis present with, fasciculations, muscle cramps, distal weakness and atrophy, bulbar palsy - dysarthria, dysphagia, respiratory compromise, spasticity, increased reflexes, flexor spasms, and pseudobulbar effects - inappropriate laughing, crying, yawning.

Option C: Multiple sclerosis is an autoimmune demyelinating disease of the central nervous system. The common symptoms of multiple sclerosis include paresthesia (tingling sensations, pins, and needle sensation), hypesthesia (numbness), optic neuritis - decreased visual acuity and color perception, unilateral/ bilateral, limb weakness - spastic paralysis, gait disturbances, Babinski's sign positive, facial weakness, vertigo, hearing loss, bladder dysfunction, Lhermitte's sign (shooting pain down the back on neck movement), Uhthoff's sign (worsening of neurological symptoms on exposure to elevated temperatures eg. hot shower)

Solution to Question 12:

The presence of a mid-diastolic murmur and a prominent a-wave in JVP is suggestive of tricuspid stenosis.

Mid-diastolic murmurs result from obstruction and/or augmented flow at the level of the mitral or tricuspid valve. Thus, both mitral stenosis (MS) and tricuspid stenosis (TS) present with a mid-diastolic murmur. However, the presence of a prominent a-wave in JVP points toward tricuspid stenosis.

Causes for mid-diastolic murmur:

• Mitral stenosis

• Tricuspid stenosis

• Atrial myxomas

• Austin Flint murmur in severe aortic regurgitation

• Carey-Coombs murmur in acute rheumatic fever

Tricuspid stenosis:

It always occurs in association with MS and is rheumatic in origin. Mid-diastolic murmur in TS is best heard along the left sternal border and over the xiphoid process. Murmur is augmented during inspiration and reduced during expiration. The jugular veins are distended, and a giant a-wave is seen in JVP. There is a slow y descent due to impedance of right atrial emptying during diastole. ECG shows features of right atrial enlargement, which includes tall, peaked P wave in lead II and a prominent, upright P wave in lead V1.



Other options:

Option A: Mid-diastolic murmur is seen in mitral stenosis (MS). It occurs most commonly in rheumatic fever. Loud S1 (first heart sound) with an opening snap and a following S2 (second heart sound) is heard. S2 is followed by a low-pitched, rumbling, mid-diastolic murmur, which is best heard at the apex with the patient in a lateral recumbent position.



Option C: Holosystolic murmur is seen in mitral regurgitation (MR). The major causes of MR are infective endocarditis, myocardial infarction, rheumatic fever, and cardiomyopathies. Soft S1 with a widely split S2 and a low-pitched S3 (third heart sound) is heard in MR. In addition, a holosystolic murmur is prominent at the apex and radiates to the axilla.



Option D: Holosystolic murmur is seen in tricuspid regurgitation (TR). TR is most commonly due to functional defects such as tricuspid annular dilatation and leaflet tethering caused by rheumatic heart disease, myocardial infarction, etc. The jugular vein is distended. Prominent c-v waves and rapid y descents are seen in severe TR. A blowing holosystolic murmur along the lower left sternal margin, which is augmented during inspiration (Carvallo's sign) and reduced during expiration is seen in TR.

Solution to Question 13:

Based on the ECG findings (tall T waves), the most likely diagnosis is hyperkalemia, and among the given drugs, spironolactone is the drug that is most likely to cause hyperkalemia.

Spironolactone is a potasium-sparing diuretic. It is a competitive inhibitor of the mineralocorticoid receptor in the late distal tubule and collecting duct of the kidneys. It causes a decrease in sodium reabsorption and potassium excretion in the distal tubule. As a result, spironolactone promotes sodium diuresis but maintains body potassium levels.

Hyperkalemia is defined as a serum potassium level of more than 5.5 mEq/L.

Hyperkalemia can cause drastic conduction changes. It can be diagnosed by a sequence of changes affecting the ST-T segments and QRS complex. ST-T segment changes reflect a change in repolarization while QRS complex changes reflect a change in depolarization.

The ECG changes that can be seen in hyperkalemia depend on the extent of hyperkalemia.

S. No.	Serum potassium level	ECG change
1	5.5-6.5 mEq/L	Tall, peaked T waves with 'tented' or 'pinched' shape
2	6.5-7.5 mEq/L	loss of P waves
3	7.0-8.0 mEq/L	widened QRS complex

S. No.	Serum potassium level	ECG change
4	> 8.0 mEq/L	Sine wave



•

Drugs that cause hyperkalemia include:

• Angiotensin-converting enzyme inhibitors

• Angiotensin receptor blockers eg. Losartan

• Mineralocorticoid receptor blockers like spironolactone, eplerenone, drospirenone

• Epithelial sodium channel (ENaC) blockers like amiloride and triamterene

• Others: Trimethoprim, pentamidine, nafamostat

Treatment of hyperkalemia involves the following:

• Administer intravenous calcium gluconate, which raises the threshold and reduces excitability. It is cardioprotective

• Administer regular insulin with dextrose. The insulin will reduce the serum potassium concentration by causing an intracellular shift. The dextrose will counteract the hypoglycemic effects of insulin.

• Salbutamol nebulization will also shift potassium inside the cell.

• Remove potassium with calcium-based resins, diuretics, or hemodialysis.

Other options:

Option A: The major potential side effect of prazosin is postural hypotension and syncope.

Option B: Metoprolol is a cardioselective beta blocker. Cardioselective beta blockers are less likely to cause hyperkalemia as an adverse effect. The risk of metoprolol-induced hyperkalemia is increased if the patient is also diabetic.

Option C: Hydrochlorothiazides increase the renal excretion of potassium and thus cause hypokalemia if the dietary intake of potassium is less.

Solution to Question 14:

Based on the given clinical scenario and the pathological examination showing Lewy bodies within the neurons the most likely diagnosis, among the given options is Parkinson's disease.

Parkinson's disease is due to widespread degeneration of the portion of the substantia nigra that sends dopamine-secreting nerve fibers to the caudate nucleus and putamen. On pathological examination, intraneuronal proteinaceous inclusions called Lewy bodies that stain for α-synuclein are seen. Mutations in α-synuclein (SNCA), GBA, LRRK2, and PINK1/Parkin are found to be responsible for this condition.

It is characterized by rigidity, resting tremors, akinesia, bradykinesia, and postural instability.

Other motor features are masked facies, reduced eye blinking, dysphagia, speech disorders, freezing, and falling. Other non-motor features include anosmia, sensory disturbances, mood disorders (e.g., depression), sleep disturbances, autonomic disturbances, and cognitive impairment/dementia.

Levodopa combined with peripheral decarboxylase inhibitor (carbidopa/benserazide) is considered the gold standard in the treatment of this disease.

Other drugs used in the treatment:

• Dopamine agonists- Pramipexole, ropinirole, rotigotine

• Monoamine oxidase (MAO-B) inhibitors - Selegiline and rasagiline

• Catechol-O-methyl transferase (COMT) inhibitors- Tolcapone and entacapone

Other options:

Option B: Prion disease is a rapidly progressing neurodegenerative disorder that occurs due to the abnormal folding of prion proteins. These proteins can undergo a structural change from an alpha helix to beta-pleated sheets (PrPC to PrPSC). These proteins can rapidly spread, invading the central nervous system and inducing neuronal cell death. Prion diseases can be sporadic in nature, but more often they have a familial mode of spread due to mutations in the PRNP gene that codes for PrP. The brain classically undergoes spongiform degeneration due to vacuole formation and rapid death of the neurons. Histopathological examination during post-mortem can reveal inclusion such as PrPSC deposits and Kuru plaques.

Option C: Huntington's disease (HD) is an autosomal dominant progressive brain disorder that causes uncontrolled movements, emotional problems, and loss of thinking ability. It occurs due to atrophy, with loss of neurons and gliosis in the caudate lobe, putamen, and globus pallidus. There is an abnormal expansion of the trinucleotide CAG in the Huntington gene on chromosome 4.

Option D: Alzheimer's disease is the most common cause of dementia in the elderly. It has an insidious onset with deficits developing over the course of 5-10 years. Initially, the patient presents with forgetfulness but then progresses to language deficits, loss of mathematical skills, and motor skills. In the end stages, the patient becomes mute, incontinent, and unable to walk. The pathogenesis of Alzheimer's disease involves the accumulation of Aβ amyloid and tau in the

brain. The hallmarks of this condition are amyloid plaques comprised of aggregated Aβ amyloid and neurofibrillary tangles.

Note: The possible diagnosis based on the given clinical scenario points to Lewy-body dementia. Since the option was not given, the second best answer would be Parkinson's disease.

Dementia with Lewy bodies is a progressive neurodegenerative associated with dementia, psychosis, and features of parkinsonism. It is characterized by the protein deposition of Lewy bodies in the substantia nigra, limbic system, and cerebral cortex, comprising of α-synuclein and ubiquitin. It is the third most common type of dementia after Alzheimer's disease and vascular dementia.

The clinical features of dementia with Lewy bodies are characterized by visual hallucinations, features of Parkinsonism, neuroleptic sensitivity, rapid eye movement (REM) sleep, behavior disorder, and excessive daytime sleepiness. They can also present with delirium and orthostatic hypotension. Memory is usually preserved in patients with Lewy body dementia. If a typical antipsychotic is prescribed for hallucinations and mild cognitive deficits, it can result in profound extrapyramidal adverse effects due to the sensitivity to neuroleptic agents. The features of Parkinsonism would include bradykinesia, resting tremor, cog-wheel rigidity, and festinating gait.

Dementia or neuropsychiatric conditions emerge concurrently with parkinsonism features in dementia with Lewy bodies whereas Parkinson's disease dementia occurs in long-standing cases of Parkinson's disease. Recently it has been decided that if dementia or any cognitive disability occurs after 12 months of Parkinson's disease, then the disorder is most likely Parkinson's disease dementia. If it is less than 12 months, then it is dementia with Lewy bodies.

Treatment: The pharmacotherapy would include cholinesterase inhibitors like rivastigmine, galantamine, and donepezil to treat the cognitive impairments. Other medications include antidepressants, atypical antipsychotics to treat depression and hallucinations respectively. Supportive therapies include physiotherapy, exercise, and family therapies.

Solution to Question 15:

Based on the given clinical vignette, the most likely diagnosis is central diabetes insipidus. Lifelong supplementation of desmopressin (DDAVP), a synthetic analog of AVP, is the definitive management in this scenario.

Diabetes insipidus (DI) is a syndrome characterized by the excretion of abnormally large volumes of dilute urine. The 24-hour urine volume exceeds 40 mL/kg body weight, and the 24-hour urine osmolarity is <280 mOsm/L. There will be a slight increase in plasma osmolarity/sodium that stimulates thirst (polydipsia). Hence, clinical symptoms and signs of dehydration are uncommon unless thirst and/or water intake are also impaired.

Causes of central diabetes insipidus include

• Idiopathic

• Familial and congenital diseases - familial neurohypophyseal DI, Wolfram syndrome, congenital hypopituitarism

• Neurosurgery and trauma - Trauma to the hypothalamus and posterior pituitary or neurosurgery with a transsphenoidal approach.

• Neoplasms

• Hypoxic encephalopathy

• Infiltrative disorders- Langerhans cell histiocytosis, granulomatosis with polyangiitis, autoimmune lymphocytic hypophysitis, and sarcoidosis

Treatment of central diabetes insipidus due to resection of pituitary tumor includes lifelong supplementation with vasopressin analogs desmopressin (DDAVP) (1-deamino-8-D-arginine vasopressin).

Note: If partial hypophysectomy, which is the usual procedure, is done, DDAVP supplementation is usually required only for a short duration as there is only a transient AVP deficiency. However, in complete hypophysectomy which results in permanent diabetes insipidus, life-long administration of desmopressin may be required.

Solution to Question 16:

Pancoast's tumor is the most likely diagnosis in this patient.

Pancoast syndrome/superior sulcus tumor results from a local extension of a tumor growing in the apex of the lung, with the involvement of C8, T1, and T2 nerve roots, stellate ganglion, and the chest wall. It presents with pain that radiates to the ulnar distribution of the arm, often with the destruction of the 1st and 2nd ribs. Patients with the central or endobronchial growth of the primary tumor may present with a cough, hemoptysis, wheeze, stridor, dyspnea, or post-obstructive pneumonitis.

Regional spread can cause

• Tracheal obstruction

• Esophageal compression with dysphagia

• Recurrent laryngeal paralysis with hoarseness

• Phrenic nerve palsy with the elevation of the hemidiaphragm and dyspnea

• Sympathetic nerve paralysis with Horner’s syndrome (enophthalmos, ptosis, miosis, and anhydrosis)

Investigations in Pancoast tumor:

• MRI > CT is the investigation of choice for superior sulcus tumor/Pancoast tumor as the associated soft tissue infiltration, the extent of which can be best visualized using an MRI.

• FDG-PET scan shows high uptake by the tumor cells. FDG is taken up by all metabolizing cells but more avidly by cancer cells.

• Bronchoscopy can be done in endobronchial tumors to obtain a tissue diagnosis.

The image below shows the management of the Pancoast tumor.



Other options:

Option B: Pneumonia patients usually present with fever, cough, chills, tachycardia, and/or sweats. Cough may be nonproductive or productive of mucoid, purulent, or blood-tinged sputum. Other symptoms may include fatigue, headache, myalgias, and arthralgias.

Option C: Patients with superior vena cava obstruction usually present with neck and facial swelling, dyspnea, and cough. Symptoms may aggravate while bending forward or lying. The characteristic physical findings are dilated neck veins, cyanosis, and edema of the face, arms, and chest. Facial swelling and plethora are typically exacerbated when the patient is supine.

Option D: Aspergilloma (fungal ball) is a late manifestation of chronic cavitary pulmonary aspergillosis, but some patients are asymptomatic. Signs and symptoms include cough (sometimes productive), hemoptysis, wheezing, and mild fatigue. More significant signs and symptoms are associated with chronic cavitary pulmonary aspergillosis.

Solution to Question 17:

The given clinical vignette points towards the diagnosis of acute gout.

Gout is a hyperuricemic metabolic condition, typically manifested by episodic inflammatory arthritis due to crystal formation and deposition of monosodium urate (MSU) crystals within the joints and connective tissue.

Acute recurrent gout flare is characterized by painful, warm, red, tender, and substantially swollen joints. The metatarsophalangeal joint of the first toe is involved in 70–90% of cases (podagra), followed by tarsal joints, ankles, and knees. Triggers of gout flares include purine-rich food, alcohol, diuretic use, local trauma, and medical illnesses such as congestive heart failure and respiratory hypoxic conditions.

Diagnosis is made by synovial fluid examination, which may show needle-shaped monosodium urate crystals. These crystals show bright, negative birefringence under polarized light. WBC count may be increased in synovial fluid. Serum uric acid level is usually elevated in chronic gout

patients, but it may remain normal or low during acute attacks, because of the uricosuric property of inflammatory cytokines.

Management of acute gout includes the application of ice packs, rest, NSAIDS, colchicine, and glucocorticoids. In patients with chronic hyperuricemia urate-lowering agents like xanthine oxidase inhibitors(eg. allopurinol and febuxostat) or uricosuric drugs (e.g. probenecid) can be used.

The negatively birefringent needle-shaped monosodium urate crystals under polarized microscopy are shown below.



Other options:

Option A: Pseudogout occurs due to the deposition of calcium pyrophosphate crystal deposition (CPPD). It presents with features similar to acute gout and predominantly affects the elderly. Acute attacks of CPPD present with joint swelling and pain, affecting the knee (most common), wrist, shoulders, elbows, and hands. It also has systemic features of inflammation including fever, chills, and elevated CRP.

Chronic CPPD commonly presents as polyarticular arthritis resembling osteoarthritis (pseudo osteoarthritis). It affects atypical joints such as metacarpophalangeal, wrist, elbow, and ankle joints causing severe damage.

X-ray or ultrasonography of the joint shows chondrocalcinosis, i.e, punctate or linear radiodense deposits in the joint menisci or articular cartilage. Detection of rod-like or rhomboid crystals in synovial fluid or articular cartilage is diagnostic of pseudogout.

Option C: Reactive arthritis, also called Reiter syndrome, is inflammatory arthritis that manifests several days to weeks after a gastrointestinal or genitourinary infection. It is also described as a classic triad of arthritis, urethritis, and conjunctivitis.

Option D: Septic arthritis is more common in children. It is the inflammation of joints secondary to an infection, usually bacteria. Hematogenous, secondary to nearby osteomyelitis, iatrogenic, penetrating wounds, and umbilical cord sepsis are the common routes of the spread of infection. The knee, hip, shoulder, and elbow are the common joints involved. Synovial fluid may look cloudy and WBC count is usually elevated predominantly with polymorphonuclear cells. The

causative organism can be seen under microscopic examination and the same can be cultured.

Medicine INI-CET 2023

Question 1:

A patient in emergency has the following lab parameters: creatinine 1.5 and BUN 45. The chest x-ray is shown below. In gram staining and culture, only rhinovirus was isolated. The patient is shifted to ICU. What is the treatment of choice in this patient?



a) Azithromycin

b) Ceftriaxone plus azithromycin

c) Amoxicillin plus clavulanic acid

d) Piptaz plus ciprofloxacin

Question 2:

A patient presents with severe vomiting and diarrhea and has orthostatic hypotension. What metabolic abnormalities would you expect in this patient?

a) 2, 4 only

b) 1, 3 only

c) 1, 2, 3, 4

d) 1, 2, 3

Question 3:

Order in which the following heart sounds are heard from superior to inferior on the left side chest wall?

a) $3 \rightarrow 1 \rightarrow 2$

b) $3 \rightarrow 2 \rightarrow 1$

c) $2 \rightarrow 1 \rightarrow 3$

d) $1 \rightarrow 2 \rightarrow 3$

Question 4:

A 35-year-old female presents with skin thickening, muscle weakness, and the peripheries becoming pale on exposure to cold. She is positive for ANA and anti-Scl-70 antibodies. Her creatinine kinase is elevated and muscle biopsy shows perifascicular infiltration. What is the antibody associated with this?

a) Anti-U3 RNP

b) Anti-centromere antibody

c) Anti-PM-Scl antibody

d) Anti-Jo-1 antibody

Question 5:

During which phase of the cardiac cycle is an AR murmur likely to be heard?

a) Ventricular ejection

b) Throughout cardiac cycle

c) Isovolumetric contraction

d) Ventricular diastole

Question 6:

Identify the given ECG.



a) Torsades de pointes

b) Ventricular fibrillation

c) Atrial flutter

d) Monomorphic ventricular tachycardia

Question 7:

Total plasma exchange is done in thrombotic thrombocytopenic purpura. How does TPE help in the management of TTP?

a) TPE removes the von Willebrand factor

b) TPE removes the antiplatelet antibodies

c) TPE replenishes the coagulation factors

d) TPE removes anti-ADAMTS13 autoantibodies and thereby increases the ADAMTS protease activity

Question 8:

Identify the neurological test being performed.

a) Babinski sign

b) Ankle jerk

c) Jendrassik

d) Normal plantar response

Question 9:

Which among the following indicates severe COVID?

a) RR >24/min

b) MAP 70 mmHg

c) Lung infiltrates >42% on CT

d) SpO2 < 85% on room air

Question 10:

Which of the following are seen in iron deficiency anemia?

a) 1, 2

b) 3, 4

c) 1, 2, 3, 4

d) 1, 3, 4

Question 11:

A patient recovered from hepatitis B infection. Which of the following curve represents antibodies seen 4 weeks post-recovery?



a) Anti-HBs

b) Anti-HBc

c) HBeAg

d) HBsAg

Question 12:

Which of the following is not a cause of reversible dementia?

a) B12 deficiency

b) Normal pressure hydrocephalus

c) Lewy body dementia

d) Hypothyroidism

Question 13:

A female presented with excessive daytime sleepiness and impaired concentration and memory for one year. On examination, the BMI was 43 kg/m2 and BP was 170/98 mmHg. ABG was done in awake state and the findings are given below. Which of the following is the likely diagnosis?

a) Obesity hypoventilation syndrome

b) Obstructive sleep apnea

c) Central sleep apnea

d) Narcolepsy

Question 14:

A patient with chronic kidney disease has the following laboratory reports. Calculate the anion gap in this patient.

a) 25

b) 68

c) 35

d) 43

Question 15:

A person comes to the OPD with symptoms of not using his left side and not washing the left side of his body while bathing. On examination, he wasn't able to draw the left side of the images. Which of the following is the most likely location of the lesion?



©MARROW

a) Left motor cortex

b) Left posterior parietal area

c) Right posterior parietal area

d) Right motor cortex

Question 16:

Identify the type of ANA staining pattern shown in the image below.



a) Centromeric

b) Homogenous

c) Nucleolar

d) Speckled

Question 17:

A chronic cigarette smoker now joined a construction company. His pulmonary function test results are given below. What is the most likely diagnosis for this patient?

a) Vascular disease with bronchodilator reversibility

b) Restrictive lung disease with bronchodilator reversibility

c) Restrictive lung disease without bronchodilator reversibility

d) Obstructive disease with bronchodilator reversibility

Question 18:

Which of the following are true about subclinical myocarditis?

a) 1, 2

b) 3, 4

c) 1, 2, 3, 4, 5

d) 2, 3, 4, 5

Question 19:

A female with nephrotic syndrome underwent a renal biopsy. Which of the following statements is not correct?

a) 1, 2, 3, 4

b) 4

c) 1, 4

d) 2, 3, 4

Question 20:

Which of the following will not be measured to diagnose metabolic syndrome?

a) HDL levels

b) Hip circumference

c) Blood glucose

d) Blood pressure

Question 21:

An old man presents with dyspnea and swelling of his feet. On examination, rales are heard and he is found to have an enlarged heart on chest x-ray. Which of the following will delay the progression?

a) Furosemide

b) Carvedilol

c) Digoxin

d) Captopril

Question 22:

A 27-year-old female patient presents with weight loss, anxiety, tachycardia, and restlessness. The resting pulse was 110 beats per minute. Which of the following tests should be done to evaluate her condition further?

a) Total T3

b) TPO

c) Thyroid Scan

d) TSH and free T4

Question 23:

A 65-year-old man was brought after a fall. He has a fever and confusion. On examination, he had a cough with rales. The patient was also tachypneic. How will you manage this patient?

a) Admit and give Oral levofloxacin

b) Treat at home and oral augmentin

c) Admit and iv ceftriaxone

d) Admit and give IV levofloxacin

Question 24:

A 60-year-old with a history of COPD is brought to the ER with worsening dyspnea. ABG was done and pH was 7.3, pCO2 was 60mmHg with HCO3- being 28 mEq/L. Which of the following is seen in this patient?

a) Respiratory acidosis, hyperventilation, inadequate metabolic compensation

b) Respiratory alkalosis, hypoventilation, inadequate metabolic compensation

c) Respiratory acidosis, hypoventilation, inadequate metabolic compensation

d) Respiratory acidosis, hypoventilation, adequate metabolic compensation

Question 25:

A 60-year-old man presented with acute onset of chest pain radiating to the left arm and diaphoresis. His ECG and cardiac biomarkers show non-ST elevation myocardial infarction. Which of the following will be used in the treatment of this patient?

a) 2, 3

b) 4, 5

c) 1, 2, 3, 4, 5

d) 1, 2, 3, 4

Question 26:

A patient with a history of consumption of 2-3 drinks per week comes with complaints of right upper quadrant discomfort. He has grade 2 steatosis. His BMI was 29kg/m2 and AST and ALT levels were slightly elevated. What would you advise him?

a) Lifestyle modification and vitamin E

b) Ursodeoxycholic acid and lifestyle modification

c) Lifestyle modifications and weight loss

d) Pioglitazone and lifestyle modifications

Question 27:

The blood investigation of a patient is given below. What is the probabale diagnosis?

a) Acute HBV infection in window period

b) Immune with recombinant HBV Vaccine

c) HBV infection in the remote past, completely recovered

d) Chronic HBV infection - inactive carrier

Question 28:

The given image shows a normal graph on the left and the patient's graph on the right. Which of the following diagnoses can be inferred from the graph?



a) Chest wall neuromuscular disease

b) Sarcoidosis

c) Idiopathic pulmonary fibrosis

d) Bronchiectasis

Question 29:

In which of the following malignancies, a liver transplantation is not contraindicated?

a) Renal carcinoma

b) Breast carcinoma

c) Melanoma

d) Testicular cancer

Question 30:

A patient with CKD with refractory hypertension is brought to the casualty with dizziness. His BP was found to be 200/120mmHg. His BUN was 50mg/dl and creatinine was 3mg/dl. Potassium was 6mEq/L. Which of the following is an indication of emergency hemodialysis?

a) Severe acidosis with elevated BUN and creatinine

b) Hypertension

c) Potassium level of 6mEq/L

d) Metabolic alkalosis

Question 31:

What is the diagnosis in the ECG shown below?



a) First degree heart block

b) Third degree heart block

c) Wenckeback heart block

d) Mobitz type 2 block

Question 32:

A 60-year-old man with a known history of diabetes mellitus presents with a BP of 150/90 and dizziness. What is the best way to manage this patient?

a) Diet, Exercise, Lifestyle modification + Metoprolol

b) Diet, Exercise, Lifestyle modification + amlodipine

c) Diet + prazosin

d) Diet, Exercise, Lifestyle modification

Question 33:

A 40-year-old male presents with increased thirst and increased frequency and volume of urination. He has no other symptoms. His random blood sugar was 205mg/dl. What

appropriate investigations should be done?

a) i, ii, iii, v

b) i, ii, iv, vi

c) i, ii, iii, iv

d) i, ii, iv, v

Question 34:

A young child presented with nocturnal chest pain and a history of syncope at school. ECG revealed a cove pattern of ST elevation in V2 and V3. What is the most probable diagnosis?

a) Myocardial infarction

b) HOCM

c) Brugada syndrome

d) DCM

Question 35:

A 26-year-old female presents to the ER with menorrhagia. She mentions that her father and sister also suffer from bleeding disorders. Which of the following is an autosomal dominant bleeding disorder?

a) Factor X deficiency

b) Factor VIII deficiency

c) Factor IX deficiency

d) Von Willebrand Disease

Question 36:

A woman comes with complaints of hyperpigmentation of her palms after an adrenalectomy. She also has new onset vision deficits. What is the likely diagnosis?

a) Nelson syndrome

b) Sheehan syndrome

c) Conn's syndrome

d) Addisons disease

Question 37:

These findings are suggestive of which of the following conditions?

a) Addison's disease

b) Gigantism

c) Acromegaly

d) Cushings syndrome

Question 38:

A 60-year-old lady with diabetes mellitus comes with dizziness. Which of the following would be seen in a patient with hyperinsulinemia?

a) Hyperketonemia

b) Increased amino acids

c) Raised glucose

d) Increased fatty acid synthesis

Answer Key

Question No.	Correct Option
1	b

2	d
3	d
4	c
5	d
6	a
7	d
8	a
9	d
10	c
11	a
12	c
13	a
14	c
15	c
16	a
17	d
18	d
19	c
20	b
21	d
22	d
23	d
24	c
25	d
26	a
27	c
28	d
29	d
30	a
31	d
32	b
33	d
34	c
35	d
36	a
37	c
38	d

Detailed Explanations

Solution to Question 1:

The chest radiograph is suggestive of pneumonia. Since this patient requires treatment in an ICU setting, the drugs of choice would be ceftriaxone plus azithromycin.

Initial treatment in inpatients with pneumonia:

Appropriate antibiotics for Pseudomonas aeruginosa or MRSA are to be added if there is a history of prior isolation from respiratory samples.

Non-severe disease	A β-lactam + a macrolide (ceftriaxone + azithromycin)orA respiratory fluoroquinolone
Severe disease	A β-lactam + a macrolide (ceftriaxone + azithromycin)orA β-lactam + respiratory fluoroquinolone

Solution to Question 2:

In the given clinical scenario, the patient is likely to have hypokalemia, hypochloremia and metabolic alkalosis.

The given clinical scenario describes a patient who has a history of vomiting and diarrhea and now has orthostatic hypotension. This indicates that the patient is dehydrated and there is a contraction of the extracellular fluid volume. This would stimulate the renin-angiotensin-aldosterone system (RAAS).

Elevated aldosterone causes net H^+ and K^+ excretion from the distal tubule of the kidneys. This causes hypokalemia and metabolic alkalosis. Alkalosis is worsened because angiotensin II stimulates the Na^+-H^+ exchanger in the renal proximal convoluted tubule and increases the reabsorption of Na^+ and HCO_3^-. Hypochloremia occurs because of loss of HCl during vomiting, and is maintained due to loss of Cl^- in the kidney which accompanies increased HCO_3^- reabsorption.

In a patient with metabolic alkalosis, the compensatory mechanism should be respiratory acidosis, not respiratory alkalosis (statement 4). This occurs in the form of hypoventilation, which increases $PaCO_2$.

Solution to Question 3:

The order of heart sounds as they are heard on the left chest wall from superior to inferior is as follows:

• Pulmonary — second intercostal space

• Aortic — can be auscultated in the neoaortic area or the Erb's point, in the third intercostal space

• Mitral — fifth intercostal space

The following picture shows the areas of auscultation:



• Aortic area, second right intercostal space: this area is primarily used for auscultation of the aortic valve. The murmurs heard best here are those of aortic stenosis, aortic regurgitation, and hypertrophic cardiomyopathy.

• Pulmonic area, second left intercostal space: this area is primarily used for the auscultation of the pulmonic valve.

• Erb's point, third left intercostal space: this area lies close to both the semilunar valves. Pathologies of both aortic and pulmonic valves can be heard at this point. Murmurs in this region are more audible if the patient leans forward.

• Tricuspid area, fourth left intercostal space: murmurs of tricuspid stenosis and tricuspid regurgitation, and ventricular septal defects are heard in this area.

• Mitral area, fifth left intercostal space in the midclavicular line: murmurs of mitral stenosis, mitral regurgitation, and mitral valve prolapse are heard here.

Solution to Question 4:

The given patient with a history and findings suggestive of diffuse systemic sclerosis and dermatomyositis/polymyositis is most likely to be suffering from overlap syndrome. The presence of skin thickening, Reynaud's phenomenon, and anti-Scl-70 antibodies are suggestive of diffuse systemic sclerosis, while muscle weakness and perifascicular infiltration are suggestive of

dermatomyositis/polymyositis.

In overlap syndromes involving polymyositis/dermatomyositis and systemic sclerosis, anti-PM-Scl antibodies are found in close to 50% of the patients, while anti-Jo-1 antibodies were found to be positive in 20-30% of the patients.

Anti-centromere antibodies (option B) are seen in limited-cutaneous systemic sclerosis. Anti-U3-RNP antibodies (also known as anti-fibrillin antibodies) are seen in both limited-cutaneous and diffuse systemic sclerosis. They signify an increased risk of the patient developing pulmonary arterial hypertension, interstitial lung disease, scleroderma renal crisis, and myositis.

Solution to Question 5:

Aortic regurgitation (AR) murmur is most likely to be heard during the ventricular diastole.

On auscultation of patients with chronic AR, a high-pitched decrescendo diastolic murmur is heard. It can be best noted in the third intercostal space on the left sternal border. In patients with mild AR, the murmur is brief. But as the disease progresses, it becomes longer in duration, and can indeed become holodiastolic. In patients in whom AR is due to primary valvular disease, the diastolic murmur is heard louder on the left sternal border. However, if the murmur is louder on the right sternal border, it indicates that AR is due to aneurysmal dilation of the aortic root. The auscultatory features of AR are intensified by handgrip, which augments systemic vascular resistance and increases LV afterload.

The heart murmurs heard in systole are:

• Aortic and pulmonic stenosis: ejection systolic murmur

• Chronic mitral and tricuspid regurgitation: holosystolic murmur

• Mitral valve prolapse: midsystolic click followed by a late systolic murmur

• Ventricular septal defect: holosystolic murmur

The heart murmurs heard in diastole are:

• Aortic regurgitation: early diastolic, decrescendo murmur

• Mitral stenosis: opening snap following by mid to late diastolic murmur

Solution to Question 6:

The given ECG shows broad QRS tachycardia with changing QRS morphology with every beat. The most likely diagnosis in this case is polymorphic ventricular tachycardia or Torsades de pointes.

The causes of torsades de pointes include congenital long QT syndrome, electrolyte abnormalities (hypokalemia, hypomagnesemia, and hypocalcemia), certain drugs, cardiac conditions like myocardial infarction and myocarditis, and endocrine disorders like hypothyroidism and pheochromocytoma.

Other options:

Ventricular fibrillation (option B) shows a completely disorganized rhythm with no identifiable waves on ECG, as shown in the image below.



Atrial flutter (option C) shows sawtooth appearance of the P-waves. This is due to rapid and successive atrial depolarization. The arrhythmia usually originates in the area of the tricuspid annulus in the right atrium. The ECG given below depicts atrial flutter, with the sawtooth P-waves most apparent in leads V1 and V2.



Monomorphic ventricular tachycardia (option D) shows wide QRS complex tachycardia but with the same QRS morphology in each beat. It occurs most commonly due to structural heart diseases (cardiomyopathy, or scarring after myocardial infarction). The patients are at high risk for sudden cardiac death. The ECG given below shows monomorphic ventricular tachycardia.



Solution to Question 7:

Therapeutic plasma exchange (TPE) removes autoantibodies to ADAMTS13. This makes ADAMTS13 enzyme functional and it is able to degrade multimers of von Willebrand factor.

Thrombotic thrombocytopenic purpura (TTP) is thrombotic microangiopathy that can be inherited or spontaneously acquired. It results from a deficiency of or antibodies to the metalloprotease ADAMTS13, which normally cleaves multimers of vWF. Failure to cleave these multimers results in excessive platelet adhesion and aggregation. This causes platelet thrombi and fibrin deposits to form at various sites in the circulation. These thrombi result in the classic pentad of clinical features:

- Microangiopathic hemolytic anemia (MAHA)
- Thrombocytopenia
- Renal failure
- Neurologic findings
- Fever

The platelet thrombi narrow the vessel lumen, causing shearing stress on the red blood cells that are passing through. This results in microangiopathic hemolytic anemia. The same platelet thrombi cause microvascular occlusion and result in features like renal failure and neurological manifestations. On peripheral smear, these fragmented red cells appear as schistocytes or helmet cells, as shown in the image below:



The mainstay of treatment of TTP is with therapeutic plasma exchange (TPE). TPE is continued till the platelet count becomes normal and signs of hemolysis are absent for at least 2 days. Glucocorticoids have been used as an adjunct to TPE. Addition of rituximab as an initial therapy also decreases the duration of TTP and relapses.

Solution to Question 8:

The image shows extensor plantar response, which is a positive Babinski sign.

The normal response to uncomfortable or painful cutaneous stimulation of the file of the foot includes plantar flexion of the toes. This can be accompanied by flexion withdrawal of the limb. If there is an upper motor neuron lesion affecting the leg being tested, there will be an extensor plantar response, which is a positive Babinski sign. The minimum extensor response to be looked for is at least the extension of the great toe. It can be accompanied by fanning of other tows and contraction of the tensor fascia lata. Other features which may be seen in UMN lesions are spasticity of the leg, pathologically brisk reflexes, and ankle clonus.

The below image shows the positive and negative Babinski signs.



The image below shows a positive Babinski sign in an adult, which is abnormal and indicative of a UMN lesion.

In infants whose corticospinal tracts are still underdeveloped, dorsiflexion of the great toe and fanning of other toes is a normal response to scratching of the sole of the foot. The image below shows an extensor response in an infant, which is normal.

Solution to Question 9:

SpO2 < 90% on room air indicates severe COVID.

According to the clinical guidelines for the management of COVID-19 patients released by the AIIMS/ICMR task force, a respiratory rate of >24/min constitutes moderate COVID, not severe COVID.

According to the clinical management protocol given by the Ministry of Health and Family Welfare, the clinical severity of COVID-19 is given as:

Severity	Presentation	Clinical parameters	Management
Mild	Uncomplicated URTI	Mild symptomssuch as fever, cough, sore throat and head acheNo shortness of breath or hypoxia:SpO2≥ 94%on room air	Managed at home or COVID care center.
Moderate	Pneumonia with no signs of severe disease	Features of pneumoniaDyspnea or hypoxia present:SpO290 to ≤ 93%on room airRespiratory rate ≥ 24/min	Managed at a dedicated COVID health center.
Severe	Severe pneumonia	Features of pneumoniaSpO2 < 90%on room airRespiratory rate >30 breaths/min	Managed at a dedicated COVID hospital.
	ARDS	As per ARDS diagnostic criteria	

Severity	Presentation	Clinical parameters	Management
	Sepsis	Life-threateningorgan dysfunctionAltered mental status, reduced urine output, weak pulse, cold extremities,weak blood pressure, coagulopathy	
	Septic shock	Persistent hypotension despite volume resuscitation, requiring vasopressors to maintain MAP ≥ 65 mmHg	



Solution to Question 10:

Iron deficiency anemia (IDA) is characterized by reduced serum iron level, reduced transferrin saturation, increased total iron-binding capacity (TIBC), and reduced ferritin level.

Serum iron level represents the amount of circulating iron bound to transferrin. When the iron stores become depleted in IDA, the serum iron level begins to fall.

TIBC is an indirect measure of the circulating levels of the iron-transporting protein, transferrin. IDA is characterized by more free transferrin (not bound to iron) available in circulation. This increases the TIBC. Transferrin saturation is calculated as: (serum iron ÷ TIBC) x 100. Normal transferrin saturation is 25-50%. Iron deficiency states show low transferrin saturation. A transferrin saturation >50% indicates a large amount of iron being delivered to the tissues, and can cause tissue iron overload. This is seen in hemochromatosis.

Serum ferritin is the most useful laboratory test to assess the levels of iron stores in the body. As iron stores are depleted, serum ferritin levels fall. Levels <15 µg/L are diagnostic of completely absent iron stores.

Solution to Question 11:

The serological marker marked in the image is anti-HBs antibody. In typical cases of resolving hepatitis B, HBsAg becomes undetectable 1-2 months after the appearance of clinical features. After this, anti-HBs becomes detectable in the serum.

After a person is infected with the hepatitis B virus (HBV), the first serologic marker to increase in level is HBsAg. It becomes detectable in the serum between 1-12 weeks. It becomes undetectable 1–2 months after the onset of jaundice and rarely persists beyond 6 months. Once the HBsAg becomes undetectable, anti-HBs antibody starts rising in serum and remains detectable indefinitely thereafter. The gap between the disappearance of HBsAg and the appearance of anti-HBs antibodies is called the window period.

HBcAg is located inside the cells and is usually enclosed within an HBsAg coat when present in the serum. As a result, HBcAg is not commonly detectable in the serum of patients with HBV infection since naked core particles are not found in the bloodstream. Conversely, anti-HBc can be detected in the serum shortly after the emergence of HBsAg, usually within 1-2 weeks. Anti-HBc may be the only serologic marker useful for the identification of HBV infection during the window period.

HBeAg appears shortly after the appearance of HBsAg. Its appearance correlates with high levels of virus replication. Anti-HBe correlates with a period of lower infectivity.



Solution to Question 12:

Lewy body dementia causes irreversible dementia.

Dementia is an acquired deterioration in cognitive abilities that impairs the successful performance of activities of daily living. The most commonly lost cognitive function is episodic memory, that is, the ability to recall events specific in time and place.

Causes of dementia are:

Reversible

• Hypothyroidism

• Thiamine deficiency, vitamin B12 deficiency

• Normal pressure hydrocephalus

• Subdural hematoma

• Brain tumor

• Chronic infection

• Autoimmune encephalopathy

Irreversible

• Alzheimer's disease

• Huntington's disease

• Lewy body dementia

• Vascular dementia

• Creutzfeld-Jacob disease

• Frontotemporal dementia

• Parkinson's disease

Solution to Question 13:

The given clinical scenario of an obese patient with daytime sleepiness and hypercapnia on ABG is suggestive of obesity hypoventilation syndrome.

Obesity hypoventilation syndrome (OHS) occurs due to increased work of breathing, respiratory muscle impairment related to the increased load, and depressed central ventilatory response to hypoxemia and hypercapnia. The diagnosis of OHS requires:

• BMI ≥30 kg/m2

• $PaCO_2$ ≥45 mmHg which signifies daytime alveolar hypoventilation

• Evidence of sleep-disordered breathing, which in most cases is due to obstructive sleep apnea (OSA)

Obstructive sleep apnea (OSA) (option B) is diagnosed if the patient has either symptoms of nocturnal breathing disturbances or daytime sleepiness, and ≥5 episodes of obstructive apnea or hypopnea per hour of sleep. Each such episode should last for at least 10 seconds, resulting in ≥3% drop in oxygen saturation or a brain cortical arousal. OSA is diagnosed by overnight polysomnogram (PSG). This includes measurement of breathing, oxygenation, body position, cardiac rhythm, sleep stages, and snoring intensity. Waking hypoxemia or hypercarbia in a patient with OSA suggests cardiopulmonary disease or obesity hypoventilation syndrome.

Central sleep apnea (CSA) (option C) is caused due to increased sensitivity to PCO_2. This causes an unstable breathing pattern which manifests as hyperventilation alternating with apnea. Prolonged circulation delay between pulmonary capillaries and carotid chemoreceptors is also a contributing cause. Hence, patients with heart failure may show CSA. They can present with

Cheyne-Stokes breathing.

Narcolepsy (option D) presents with difficulty sustaining wakefulness and disturbed nocturnal sleep. However, unlike sleep apnea, patients with narcolepsy usually feel well-rested on awakening and then feel tired throughout the day. One of the most characteristic clinical features which is seen is cataplexy – sudden muscle weakness without a loss of consciousness, which is usually triggered by strong emotions.

Solution to Question 14:

The anion gap in this patient is 35 mEq/L.

The anion gap is calculated as $Na^+ – [Cl^- + HCO_3^-]$. In this patient, the anion gap will be calculated as: 145 – (90 + 20) = 145 – 110 = 35 mEq/L.

Anion gap is representative of unmeasured anions which are normally present in the plasma, such as anionic proteins (albumin), phosphate, sulfate, and organic anions. When acid anions such as acetoacetate and lactate accumulate in the extracellular fluid, the anion gap increases, causing a high anion gap metabolic acidosis. A decrease in anion gap can occur due to an increase in abnormal cations (such as lithium, immunoglobulins), reduction in plasma albumin (nephrotic syndrome) or hyperviscosity and hyperlipidemia.

Solution to Question 15:

The given presentation and image suggest hemispatial neglect on the left side. This is likely due to a contralateral (right) parietal lobe lesion.

The right hemisphere of the brain directs attention within the entire extrapersonal space – both right and left side. If there is a lesion in the left hemisphere, there is not much contralesional neglect because the right hemisphere can compensate for both sides. The left hemisphere directs attention only in the contralateral right hemispace. If there is a lesion in the right hemisphere, it leads to severe contralesional hemineglect because the left hemisphere does not contain ipsilateral attention functions.

Patients with neglect may not be able to dress or groom the left side of the body. They may fail to copy the details on the left side of a drawing. Two bedside tests can be used to evaluate neglect:

• In simultaneous bilateral stimulation, when the examiner provides unilateral or simultaneous bilateral stimulation in the visual, auditory, or tactile modalities, patients have no difficulty detecting unilateral stimuli on either side but perceive the bilaterally presented stimulus as coming only from one side. This phenomenon is known as extinction.

• In visual target cancellation, the patient fails to detect the targets on one-half of the field in target-detection tasks.

Other features of non-dominant pareital lobe lesions include:

• Constructional apraxia

• Visuo-spatial disorientation

• Dressing apraxia

Solution to Question 16:

The given picture shows discrete and large speckles in the resting cells, and the alignment of the speckles in the chromosome region in the dividing cells. This is suggestive of centromeric pattern of ANA staining. Centromeric pattern is seen in limited cutaneous systemic sclerosis (CREST syndrome).

The main patterns of ANA staining are as follows:

Homogenous pattern refers to diffuse staining of the nucleus in the resting cells. There is also staining of the chromosome region in the dividing cells. The usual antibodies which show this pattern are anti-histone antibodies and anti-dsDNA antibodies. This type of pattern is also seen in chronic autoimmune hepatitis and juvenile idiopathic arthritis. Homogenous staining is shown in the image below.



Fine speckled pattern refers to hundreds of dots present throughout the nucleus. However, there is no staining of the chromosome region of the dividing cells as shown in the image below. This pattern is shown by anti-SS-A/Ro and anti-SS-B/La antibodies.



Coarse speckled pattern refers to hundreds of dots present in the nucleus, but generally larger than the dots seen in fine speckled pattern. There is no staining of nucleoli or dividing chromosomes. This type of pattern is seen in anti-Sm and anti-U1 RNP antibodies.

Dense fine speckled pattern shows the speckles distributed throughout the nucleoli. This pattern differs from the fine and coarse speckled patterns in that the speckles are associated with the chromosomes in the dividing cells. This type of pattern is not associated with systemic rheumatic disease.



Solution to Question 17:

In the given clinical scenario, FEV1/FVC is less than 0.7, hence it is obstructive. After bronchodilation, FEV1 has increased from 0.9L to 1.9L, which is more than 12% implying its reversible. Therefore, the most likely diagnosis is an obstructive disease with bronchodilator

reversibility.

In obstructive pulmonary disease, FEV1/FVC is reduced to <0.7 and FEV1 is also reduced <80%. Reversibility is defined as a ≥12% increase and absolute increase of 200mL in the FEV1 at least 15 min after administration of a β2 -agonist. Examples of obstructive pulmonary disease include COPD, asthma, bronchiectasis, etc.

In restrictive pulmonary disease, FEV1/FVC is normal or increased (>0.7) and FEV1 and FVC are also reduced. Examples of restrictive pulmonary disease include asbestosis, sarcoidosis, idiopathic pulmonary fibrosis, etc.

Solution to Question 18:

A patient is diagnosed with possible subclinical acute myocarditis when there is a typical syndrome but without any cardiac symptoms such as palpitations. Biomarkers of cardiac injury are elevated (statement 2), ECG is abnormal (statement 4), and there is a reduced LV ejection fraction (statement 5) and regional wall motion abnormality on 2D echo (statement 3).

Probable acute myocarditis is diagnosed when the aforementioned criteria are accompanied by cardiac symptoms such as palpitations, shortness of breath, and chest pain. When there are signs of pericarditis along with elevated CK-MB or troponins and regional wall motion abnormalities, the term myopericarditis or perimyocarditis is used.

Definite myocarditis is diagnosed when there is histological evidence of inflammation on endomyocardial biopsy, and does not require any other laboratory or clinical criteria.

Solution to Question 19:

Patients undergoing a percutaneous kidney biopsy are observed for a 24-hour period (statement 2) in the hospital. If any major complication occurs, it is likely to happen during this period, and thus likely to be identified and treated early. Bleeding and hematoma formation are relatively common complications of kidney biopsy and can present as frank hematuria or dark red clots in the urine (statement 3).

Focal segmental glomerulosclerosis (FSGS) is a pathological diagnosis, so obtaining a tissue sample from kidney biopsy is necessary (statement 1). On light microscopy, there are focal (affecting only some of the glomeruli) and segmental (affecting only part of the glomerulus) lesions. There is a collapse of capillary loops, an increase in the matrix, and deposition of plasma proteins.

In patients with features of systemic amyloidosis (statement 4), rectal or gingival tissues can be biopsied for the confirmation of the diagnosis. Examination of abdominal fat pat aspirates can also be used for confirmation of systemic amyloidosis.

Solution to Question 20:

Hip circumference is not used in the diagnosis of metabolic syndrome. Waist circumference is the parameter used.

Metabolic syndrome is present if three or more of the following five criteria are met, as defined by National Cholesterol Education Program and Adult Treatment Panel III (NCEP-ATP III):

• Central obesity: Waist circumference of >102 cm in males and >88 cm in females

• Hypertriglyceridemia: Triglyceride level of ≥ 150 mg/dL

• Low HDL cholesterol: <40 mg/dL and <50 mg/dL for men and women, respectively

• Hypertension: Systolic BP of ≥130 mmHg or diastolic BP of ≥ 85 mmHg

• Fasting plasma glucose level: ≥100 mg/dL or previously diagnosed type 2 diabetes

BMI and LDL cholesterol levels are not part of NCEP-ATP III.

Metabolic syndrome (syndrome X) is a group of conditions that increases the risk for cardiovascular disease. These include hypertension, hyperglycemia, hypertriglyceridemia, and central obesity. Increasing age, genetics, and sedentary lifestyle are considered risk factors.

Insulin resistance is the most accepted hypothesis used to explain the pathophysiology of the condition. Elevated circulating fatty acids are considered a major factor in developing insulin resistance. It is associated with acanthosis nigricans which are velvety hyperpigmented lesions in skin flexures such as the axilla, groin, and back of the neck.

Solution to Question 21:

The given scenario is suggestive of heart failure. ACE inhibitors like captopril are the first-line drugs to prevent further cardiac remodeling and prevent the progression of disease.

Heart failure is a complex clinical syndrome that occurs because the ventricle is unable to fill with or eject blood. The important etiologies of heart failure include hypertension, diabetes mellitus, coronary artery disease, myocardial infarction, high output states, such as beriberi, thyrotoxicosis, and severe anemia.

Clinical features of heart failure include dyspnea on exertion, orthopnea, pedal edema, anorexia, pallor, elevated JVP, ascites, and hepatomegaly.

Heart failure is classified based on the ejection fraction:

• Heart failure with preserved ejection fraction (HFpEF) - left ventricular ejection fraction is ≥50%

• Heart failure with mid-range ejection fraction (HFmEF) - left ventricular ejection fraction 40-50%

• Heart failure with reduced ejection fraction (HFrEF) - left ventricular ejection fraction is ≤40%

The mainstay management of heart failure is medical therapy and device therapy is reserved for refractory cases. The type of medications used in medical therapy are:

• Drugs with symptomatic and mortality benefits provide symptomatic relief and reduce mortality risk by preventing cardiac remodeling. It includes -

• Angiotensin receptor–neprilysin inhibitor (ARNI) - a combination of sacubitril and valsartan

• Angiotensin-converting enzyme (ACE) inhibitors - ramipril, enalapril

• β-blockers - metoprolol, carvedilol, bisoprolol

• Mineralocorticoid receptor antagonist - spironolactone, eplerenone

• Sodium-glucose cotransporter 2 inhibitors (SGLT2 inhibitor) - dapagliflozin

• Drugs with only symptomatic benefits provide only symptomatic relief and have no effect on reducing mortality risk. It includes -

• Loop diuretics - furosemide

• Digoxin

Unfortunately, evidence of the clinical benefits of most therapies is restricted to HFrEF. Pharmacological therapy used in HFpEF is mainly diuretics.

Non-pharmacological therapy includes implantable cardioverter defibrillators (ICD), cardiac resynchronization using biventricular pacing, etc. These treatments are not found to be effective in HFpEF.

Solution to Question 22:

This patient is presenting with symptoms of hyperthyroidism. The next step in diagnosis would be to obtain a serum TSH level and Free T4 level.

Investigations in a patient with Graves' disease

• Thyroid function tests- In Graves disease, the thyroid-stimulating hormone (TSH) levels are decreased with or without elevated T3 and T4 levels. It is the first step in evaluating suspected thyroid disease.

• Anti-thyroid antibodies- Elevated levels of TSH receptor antibodies are diagnostic of Graves' disease. Other antibodies like anti-Tg and anti-TPO are elevated in most of the patients.

• Ultrasound of the neck is performed to evaluate vascularity, echogenicity, nodules, and clinical landmarks for surgical planning.

• CT or MRI scans of the orbits are useful to evaluate Graves' ophthalmopathy.

The hyperthyroidism of Graves' disease is caused by thyroid-stimulating immunoglobulin (TSI) that are synthesized in the thyroid gland as well as in bone marrow and lymph nodes. TSI antibodies cause stimulation of the thyroid gland causing features of hyperthyroidism in the presence of low TSH levels. Histopathology shows follicular and papillary hyperplasia with lymphoid infiltration.

A patient with Graves' disease will present with complaints of:

• Hyperactivity, irritability

• Heat intolerance and sweating

• Palpitations

• Weakness

• Weight loss with increased appetite

• Diarrhea

• Polyuria, oligomenorrhea, and loss of libido.

On examination, they may have

• Tachycardia

• Exophthalmos

• Lid retraction or lag

• Goiter

• Tremor

• Gynecomastia

• Myopathy.

Other options: TPO, Serum T3 and thyroid scans are obtained only after serum TSH is obtained to confirm the diagnosis. But the first step remains measuring TSH levels.

Solution to Question 23:

The given clinical scenario of a patient with fever and productive cough point to a diagnosis of pneumonia. Since this patient's CURB-65 score is at least 3 (1 point each for confusion, tachyapnea, and Age 65), the most appropriate management would be ICU admission and starting IV antibiotics. This patient with a score of 3 will need ICU admission and treatment with a respiratory quinolone like levofloxacin.

Once the diagnosis of pneumonia is made, the next step in management is to determine the site of care for the patient. To identify the patients that require in-patient and ICU management, CURB-65 score is used. It includes:

• Confusion

• Urea ≥ 20 mg/dL

• Respiratory rate ≥30/min

• Blood pressure: SBP ≤90mmHg or DBP ≤60 mmHg

• Age ≥ 65 years

1 point is given for the presence of each of the above. The total score determines the site of care, as shown below:

Note: Treatment with a combination of Beta-lactam and a macrolide or a respiratory fluoroquinolone alone results in lower mortality than monotherapy with a beta-lactam antibiotic. Therefore, out of the given options, D is the best answer.

Total Score	Decision
0	Outpatient treatment

Total Score	Decision
1 or 2	Inpatient treatment in non-ICU setting
≥3	Inpatient treatment in ICU setting

Solution to Question 24:

The arterial blood gas values suggest a likely diagnosis of respiratory acidosis (pH of 7.3 and elevated $PaCO_2$ of 60mmHg indicating inadequate respirations). Normal serum bicarbonate levels are 22-26mEq/L indicating that this patient's acidosis is inadequately compensated.

The various buffering systems present in the body maintain the systemic arterial pH between 7.35-7.45. The most common acid-base disorders are metabolic acidosis or alkalosis and respiratory acidosis or alkalosis.

Normal values in an arterial blood gas are as follows:

• pH: 7.35-7.45

•

$PaCO_2$: 35-45 mmHg

•

HCO_3: 22-26 mEq/L

•

PaO_2: 75-100 mmHg

Respiratory changes (changes in PCO_2) promote a corresponding change in metabolic response (changes in HCO_3-) and vice versa as a compensatory mechanism. In general, PCO_2 and HCO_3- move in the same direction for compensation in metabolic and respiratory acid-base disorders respectively. A change in the two parameters in opposite directions points to a mixed acid-base disturbance.

Respiratory acidosis can be secondary to severe pulmonary disease, respiratory muscle fatigue, or ventilatory control abnormalities. It is marked by an increase in $PaCO_2$ and a decrease in pH. A compensatory rise of HCO_3- can also be seen.

A rapid rise in PCO_2 (acute hypercapnia) can present with dyspnea, confusion, and psychosis. It can be seen in severe asthma secondary to bronchospasm, anaphylaxis, or inhalational burns. Chronic hypercapnia causes sleep disorders, memory loss, and daytime somnolence. It is commonly seen in end-stage obstructive lung disease.

Acute respiratory acidosis may need tracheal intubation and assisted mechanical ventilation. Rapid correction of hypercapnia should be avoided as it can trigger cardiac arrhythmias and seizures. Chronic respiratory acidosis is difficult to manage and improvement in lung function is the primary aim.



Solution to Question 25:

While atorvastatin, aspirin, clopidogrel, and low molecular weight heparin all have a role in the management of a patient with NSTEMI, there is no role of alteplase.

Initial management of a patient with NSTEMI includes bed rest to decrease the load on the heart. Oxygen is administered only if the patient has hypoxemia (SpO2 <90%) or heart failure. Further management includes the following medical therapy:

Nitrates (such as nitroglycerine given sublingually) reduce the venous return to the heart, thus reducing the cardiac output and the oxygen demand. Nitrates are contraindicated if the patient has hypotension.

β-blockers also reduce the myocardial oxygen demand predominantly by their negative chronotropic effect. Diltiazem or verapamil are given to reduce the heart rate if the patient doesn't respond to β-blockers.

Morphine is given if the patient continues to have severe chest pain despite the administration of the abovementioned anti-ischemic drugs.

Antithrombotic therapy, which includes:

• Dual antiplatelet therapy (DAPT) with aspirin and a P2Y12 blocker such as clopidogrel. DAPT is continued for a minimum of 3 months, but preferably for 12 months.

• Anticoagulants such as low molecular weight heparin and bivalirudin

Clot formation and fibrinolysis is a dynamic process occurring in the coronary arteries. Antithrombotic therapy is given to prevent new clot formation while fibrinolysis is occurring by intrinsic or extrinsic mechanisms.

Statins, such as atorvastatin are started because they reduce the risk of recurrence of NSTEMI and stabilize the atherosclerotic plaque in the coronary arteries.

Solution to Question 26:

This patient with a history of consumption of 2-3 alcoholic drinks per week(acceptable limit is <5 per week for men)presents with right upper quadrant discomfort with a BMI of 29kg/m2 and slightly elevated AST and ALT most likely has nonalcoholic fatty liver disease(NAFLD). The first line for management of this condition is to advise him on lifestyle modifications and vitamin E. All the other options include drugs that are used off-label but are not FDA-approved for use in NAFLD.

Non-alcoholic fatty liver disease (NAFLD) is associated with diabetes and obesity. The fatty change may also be caused due to the effect of toxins (including certain medications), protein malnutrition, dyslipidemia, and anoxia. Bleomycin, methotrexate, doxycycline, and l-Asparaginase are other drugs causing hepatic steatosis.

Most of the patients with NAFLD and NASH are asymptomatic. They are usually diagnosed incidentally on routine blood work. Some patients with NASH can present with right upper quadrant pain due to stretching of the liver capsule from hepatomegaly, or acanthosis nigricans due to insulin resistance or with signs like palmar erythema, spider telangiectasia, jaundice, splenomegaly, and ascites if they have liver cirrhosis. Currently, a biopsy is the gold standard investigation to establish the diagnosis.

The current approach to NAFLD management focuses on treatment to avoid or decrease the risk factors for NASH (like obesity, insulin resistance, metabolic syndrome, and dyslipidemia). For patients with morbid obesity, bariatric surgery can be done. In patients with decompensated cirrhosis, liver transplantation is can be considered.

Note: The FDA in 2024 has approved Resmetirom, a thyroid hormone receptor-beta (THR-beta) agonist for noncirrhotic NASH with moderate to advanced fibrosis.

Solution to Question 27:

The most probable diagnosis is remote infection with hepatitis B virus (HBV).

A non-reactive HbsAg rules out active HBV infection (Options D).

Non-reactive IgM anti-HbcAg suggests that the patient is not in the window period (Option A).

The absence of Anti HbS makes vaccination unlikely.(Option B)

Solution to Question 28:

The patient’s graph shows a reduction in both FVC & FEV1/FVC which is < 70%, indicating an obstructive lung disorder. Therefore, the most likely diagnosis is Bronchiectasis.

Chest neuromuscular disease (option A), sarcoidosis (option B) & idiopathic pulmonary fibrosis (option C) are restrictive lung disorders, where there is a reduction of FVC to < 4L, Without a loss in FEV1/FVC ratio (normal or >70%).

These values are measured as a part of the pulmonary function test known as spirometry. At a given lung volume, when a person inhales to the total lung capacity and exhales as hard and

completely as possible, the volume of expired air is recorded as forced vital capacity (FVC), and the volume of air expired in the first second is called the forced expiratory volume (FEV1).



Solution to Question 29:

Liver transplantation is relatively not contraindicated in testicular cancer. Past history of malignancy is a relative contraindication to transplantation, because the effect of surgery and resulting immunosuppression may predispose the patient to early recurrence of the tumor.

Testicular carcinoma has a relatively low risk of recurrence (0%-10%) following transplantation. In contrast, renal carcinoma, breast carcinoma, and melanoma pose a higher risk of recurrence, exceeding 10%.

Solution to Question 30:

Severe acidosis with elevated BUN and creatinine with symptoms like dizziness and altered mental status is an indication for hemodialysis.

The emergent indications of dialysis or renal replacement therapy can be remembered by the mnemonic - 'AEIOU' (vowels in the English alphabet). These include:

• Severe metabolic Acidosis (pH < 7.1) with concomitant renal injury (Option D).

• Electrolyte abnormalities - Severe hyperkalemia (>6.5 mmol/L) or rapidly rising potassium levels (Option C).

• Intoxication - In cases of severe poisoning of toxic substances like methanol or ethylene glycol, and in cases of overdose with a dialyzable drug.

• Fluid Overload that is not responsive to diuretic therapy.

• Uremia - elevated BUN with signs and symptoms of uremia like bleeding, pericarditis, and altered mental status.

Solution to Question 31:

The ECG shown above shows regular PR intervals with a sudden dropped beat. This indicates a second-degree atrioventricular block, Mobitz type 2.

Second Degree Heart Block: It occurs due to the intermittent failure of electrical impulse conduction from the atrium to the ventricle. It is characterized by a missed QRS. It is of two types:

• Mobitz type 1/Wenckebach occurs due to the decremental conduction of electrical impulses in the AV node. It is characterized by a progressive PR elongation leading to an eventual drop in the QRS complex. This may be seen in young athletes.



• Mobitz type 2 occurs in the distal or infra-His conduction system and is often associated with intraventricular conduction delays. It is characterized by an intermittent failure of conduction of the P wave without any change in the preceding impulses. On ECG there is a fixed PR interval and missed QRS.



Other options:

Option A: In a first-degree AV block, a prolonged PR interval (of over 200 ms) is seen which indicates a slowing of conduction between the atria and ventricles, usually due to slow conduction through the atrioventricular node (AV node).

Option B: Third-degree heart block presents as a complete dissociation between the P wave and the QRS complex and occurs due to complete blockage of transmission of impulses from the AV node to the ventricles. Here, the ventricles assume an idioventricular rhythm and do not correlate to the P waves.

Solution to Question 32:

In this patient with diabetes mellitus with new-onset hypertension, he should be advised conservative measures along with amlodipine for controlling his blood pressure.

Patients with co-existing hypertension and diabetes are at increased risk of cardiovascular events. Thus blood pressure should be adequately controlled in these patients regardless of ASCVD 10 year risk.

Lifestyle modifications to manage hypertension:

• Attain and maintain a BMI of <25 kg/sq m

• A diet rich in fruits, vegetables, and low-fat dairy products with reduced content of saturated and total fat. Diet rich in potassium, calcium and magnesium.

• Dietary salt <6 grams/day

• For those who drink alcohol, 2 or less drinks/day for men and 1 or less drink/day for women

• Physical activity: Regular aerobic activity, eg, brisk waking for 30 min/day.

Prazosin would be contraindicated in this patient due to his history of dizziness as it can cause orthostatic hypotension.

Solution to Question 33:

This patient presenting with increased thirst(polydipsia), increased urination(polyuria) and a random blood sugar of 205 makes a diagnosis of type 2 diabetes mellitus likely. For a newly diagnosed case of diabetes, patients need to undergo other investigations to check for systemic progression of disease. These investigations include HbA1c, Urine albumin creatinine ratio, Fundoscopy, and ECG.

Diabetes can cause many microvascular and macrovascular complications and requires screening for these conditions.

• HbA1c is done to estimate the glycemic control over the last 3 months and helps guide further therapy. It should be done 2-4 times per year based on the previous values.

• Urine albumin creatinine ratio is done to check if there is progression to diabetic kidney disease. It should ideally be done annually to check for kidney disease.

• Dilated fundoscopy should be done annually in diabetics to rule out diabetic retinopathy. For a newly diagnosed case of T2DM, fundoscopy should be done at the time of diagnosis and then every year till there is no diabetic retinopathy.

• ECG should be done to check for resting electrolyte abnormalities in diabetics. Additionally, many diabetic patients also have co-existing hypertension so ECG is done to rule out left ventricular hypertrophy.

• Testing for diabetic neuropathy should be done annually with the monofilament test.

• BP screening 2-4 times a year

Solution to Question 34:

A given clinical scenario of nocturnal chest pain and a history of syncope in a young kid along with ECG findings of coved ST-segment elevation of V1-V3, points towards the diagnosis of Brugada syndrome.

Brugada syndrome is an inherited sodium channelopathy (SCN5A mutation) and is most commonly seen in Southeast Asian males. The age of presentation is usually 40 years. This syndrome is characterized by an increased risk for sudden cardiac death due to polymorphic ventricular tachyarrhythmias. Implantable cardioverter-defibrillator is the treatment of choice.

Brugada syndrome is associated with right bundle branch block. It is an inherited disease that predisposes patients to potentially life threatening arrythmias.

The ECG below is depictive of Brugada syndrome with concave ST segments and negative T waves (to be seen in more than one anterior precordial leads, i.e., V1-V3)



Solution to Question 35:

This patient's clinical features are suggestive of von-Willebrand disease. It is caused by a deficiency of the von-Willebrand factor.

They are present within cytoplasmic vesicles called Weibel-Palade bodies.

Normal Platelet count, Prolonged BT, Normal PT, and Prolonged aPTT are seen in von-Willebrand disease. Factor VIII is found bound to the Von-Willebrand factor (vWF) in the blood. Deficiency of vWF results in Factor VIII deficiency and prolongation of aPTT in this disease. Prolonged BT is due to abnormal platelet-mediated hemostasis.

vWF is also found in

• Platelet granules (Less as compared to endothelial cells)

• Subendothelium, where it binds to collagen

Von Willebrand disease is the most common inherited bleeding disorder in humans. The bleeding tendency is usually mild and manifests during surgical or dental procedures. The most common presenting symptoms are spontaneous bleeding from mucous membranes (epistaxis), excessive bleeding from wounds, or menorrhagia.

It is usually transmitted as an autosomal dominant disorder, but rare autosomal recessive variants can be seen.

Three types are recognized, with phenotypic variations:

Type 1 and type 3 von-Willebrand disease are associated with quantitative defects in vWF.

Type 1 is an autosomal dominant disorder characterized by a mild to moderate vWF deficiency, accounting for about 70% of all cases. It is associated with mild disease.

Type 3 disease is an autosomal disorder resulting in little to no vWF synthesis. Because vWF stabilizes factor VIII in the circulation, factor VIII levels are also reduced in type 3 disease and the associated bleeding disorder is often severe.

Type 2 von-Willebrand disease is characterized by qualitative defects in vWF. Type 2A is the most common. It is inherited as an autosomal dominant disorder.

Other options:

Factor X, VIII and IX deficiencies are all causes of hemophilias. These conditions manifest as prolonged bleeding after injuries, dental treatment and hemarthrosis. They are also additionally X-linked conditions so the presence of this condition in her father and sister makes X-linked transmission impossible.

Solution to Question 36:

The clinical scenario of hyperpigmentation, history of adrenalectomy and visual deficits indicating a probable pituitary adenoma points to the diagnosis of Nelson's syndrome.

Nelson's syndrome is a rare complication of total bilateral adrenalectomy (TBA). TBA is a procedure used to control hypercortisolism in Cushing's disease (due to ACTH-secreting pituitary adenoma) when resection of the primary tumor is not possible.

The loss of feedback inhibition of the hypothalamic-pituitary-adrenal axis following TBA increases CRH. This leads to stimulation of ACTH-secreting corticotroph cells, causing enlargement of the original pituitary tumor or formation of a new one.

Nelson's syndrome is associated with a clinical triad of:

• Hyperpigmentation

• Excessive ACTH secretion

• Corticotroph adenoma

Screening is done with fasting plasma ACTH levels and MRI brain performed at regular intervals. Prophylactic pituitary irradiation may be recommended to prevent its incidence. The primary treatment is transsphenoidal surgery. Patients will also require life-long glucocorticoid replacement, as they do not produce cortisol.

The use of steroidogenic inhibitors in the treatment of Cushing's disease has decreased the need for bilateral adrenalectomy these days.

Option B: Sheehan syndrome refers to panhypopituitarism. It classically follows massive postpartum hemorrhage and associated hypotension. The abrupt, severe hypotension leads to pituitary ischemia and necrosis. Patients with the most severe form develop shock due to pituitary apoplexy. Pituitary apoplexy is characterized by a sudden onset of headache, nausea, visual deficits, and hormonal dysfunction due to acute hemorrhage or infarction within the pituitary.

Option C: Conn's syndrome or primary aldosteronism occurs due to aldosterone-secreting adrenal adenoma leading to mineralocorticoid excess. It is characterized by hypokalemia, hypomagnesemia, muscle weakness, proximal myopathy and muscle cramps.

Option D: Addison's disease (primary adrenal insufficiency), most commonly caused by autoimmune adrenalitis presents with hyperpigmentation of the skin, especially in areas exposed to sunlight or prone to friction. This is due to excessive stimulation of melanocytes by the raised adrenocorticotropic hormone (ACTH). This is seen in chronic cases. However, there are no visual symptoms in Addison's disease.

Solution to Question 37:

The clinical image of a man with mandibular enlargement, prognathism (protrusion of jaw), and jaw malocclusion suggests the diagnosis of acromegaly.

Acromegaly is a result of growth hormone hypersecretion due to a pituitary adenoma. It can also occur due to extra pituitary lesions. If growth hormone hypersecretion occurs before the fusion of the growth plates at puberty, it leads to gigantism.

75g of oral glucose load is administered to the patient. The growth hormone levels are measured within 1-2 hours after administration. Normally, GH should be suppressed to less than 0.4 ug/L after the oral glucose load. A failure of suppression of growth hormone confirms the diagnosis of acromegaly. In some patients, growth hormone levels may demonstrate a paradoxical rise.

Acral bony overgrowth in acromegaly manifests as frontal bossing, increased hand and foot size, mandibular enlargement, prognathism, and widened space between incisors. Soft tissue swelling results in increased heel pad thickness, coarse facial features, and a large fleshy nose.

Generalized visceromegaly is seen including cardiomegaly, macroglossia, thyroid enlargement, etc. Serious complications include diabetes mellitus, obstructive sleep apnoea, and increased risk of colonic polyps and cancer.

Acromegaly can be treated by transsphenoidal resection of the tumor. It can also be managed medically by using octreotide (a somatostatin analog), pegvisomant (Growth Hormone receptor antagonist), dopamine agonists or radiation.

Other options:

Option A: Addison's disease is a primary adrenal insufficiency in which the adrenal is unable to secrete adequate amounts of cortisol (glucocorticoid), and aldosterone (mineralocorticoid). The features are fatigue, weight loss, anorexia, low blood pressure, postural hypotension, hypoglycemia, nausea, vomiting, abdominal pain, salt craving, hyponatremia and hyperkalemia.

Option B: Pituitary gigantism is caused by the overproduction of growth hormone, before the closure of epiphysis. It usually does not present with prognathism as this occurs after the closure of the growth plates. It presents as longitudinal growth acceleration. It is a rare condition, caused mostly by pituitary adenoma

Option D: Cushing syndrome results from chronic exposure to excess glucocorticoids. It presents with weight gain, central obesity, hypertension, and hypokalemia. Other clinical features include thin brittle skin with easy bruisability, purple stretch marks, diabetes, dyslipidemia, and osteopenia.

Solution to Question 38:

Insulin increases fatty acid synthesis in adipose tissue.

Effects of insulin on various tissues:

Tissue	Increases	Decreases
Adipose tissue	Increased glucose entryIncreased fatty acid synthesisIncreased glycerol phosphate synthesisIncreased triglycerides depositionActivation of lipoprotein lipaseIncreased K+ uptake	Inhibition of hormone-sensitive lipase
Muscle	Increased glucose entryIncreased glycogen synthesisIncreased amino acid uptakeIncreased protein synthesis in ribosomeIncrease ketone uptakeIncrease K+ uptake	Decrease protein catabolism Decrease release of glucogenic amino acids
Liver	Increased protein synthesisIncreased lipid synthesisIncreased glycogen synthesis & glycolysis	Decrease ketogenesisDecreased glucose output due to decreased gluconeogenesis
General	Increased cell growth	

Medicine INI-CET 2024

Question 1:

A patient is admitted with gastroenteritis. His serum K+ is 2.8 mEq/L. Which is not a possible complication of this condition?

a) Diabetes insipidus

b) Cerebral edema

c) Quadriparesis

d) Ventricular tachycardia

Question 2:

A man was brought to the emergency department unconscious and hypotensive. His ECG revealed narrow complex tachycardia. What is the next best step in management?

a) Synchronised cardioversion

b) Carotid massage

c) IV adenosine 6mg immediately

d) Amiodarone bolus followed by saline infusion

Question 3:

A 40 year old female patient presents with chest pain and breathlessness. She had a brief viral illness 1 week ago. Her chest X-ray is given below. What is not true about this condition?



a) Loud S1

b) Pulsus paradoxus may be seen

c) Emergency pericardiocentesis may be required

d) ECHO showing diastolic collapse of RA and RV wall during expiration requires immediate intervention

Question 4:

A patient is diagnosed with enterococcal infective endocarditis. He is allergic to penicillin. Which of the following is the antibiotic of choice for empiric therapy?

a) Piperacillin

b) Vancomycin

c) Aztreonam

d) Ceftriaxone

Question 5:

Which is not used as a diagnostic criterion for diabetic ketoacidosis?

a) Serum glucose>250mg/dl

b) Serum potassium <3.5mEq/L

c) Serum bicarbonate <15mEq/L

d) ph <7.3

Question 6:

A patient presents with complaints of recurrent kidney stones, abdominal pain, and altered mentation. Decreased serum Ca2+ and elevated parathyroid hormone are seen on further investigation. Which of the following disorders explains this patient's condition?

a) Bone - Metabolic bone disorder

b) Parathyroid - Hyperparathyroidism

c) Thyroid - Hypothyroidism

d) Adrenal gland - Pheochromocytoma

Question 7:

Interpret the ABG findings

a) Respiratory acidosis

b) Respiratory alkalosis

c) Metabolic acidosis

d) Metabolic alkalosis

Question 8:

A patient presents with complaints of chest pain for 1 hour and perspiration. He also appears anxious. ECG image is given below. What is the possible diagnosis?



a) NSTEMI

b) Unstable angina

c) Posterior wall MI

d) Acute Myocarditis

Question 9:

Which of the following is the least common cause of bipedal edema?

a) Chronic venous insufficiency

b) Lupus nephritis

c) Chronic liver disease

d) Heart failure

Question 10:

A patient presents with sudden onset lower motor neuron facial paralysis on his left side. Radiological tests showed no deficits. What is the next best step in the management?

a) Intravenous antibiotics, and mastoidectomy

b) Acyclovir, eye patching and eye ointment

c) Electrical stimulation of facial nerve

d) Facial nerve decompression

Question 11:

A 50year old diabetic male presents with complaints of breathlessness on lying down and walking uphill. His ECHO shows ejection fraction of 40%. Fasting blood glucose is 135mg/dl, BP -164/93 mmHg, HbA1c is 7.8%, Hb- 13mg/dl. Which of the following anti-diabetic drugs can be used for this patient?

a) Dapagliflozin

b) Liraglutide

c) Glimepiride

d) Pramlintide

Question 12:

Which of the following are features of Korsakoff's psychosis?

a) a,b,c,d

b) a,b,c

c) a,c,d

d) a,b,d

Question 13:

Which of the following is an indication to start corticosteroids in a patient with IgA vasculitis?

a) Gastrointestinal bleeds

b) Arthritis

c) Mild pain abdomen

d) Severe skin rash

Question 14:

A 35-year-old female patient presents with dyspnea and palpitations. ECG is given below. Which of the following is false about her condition?



a) Is a rare finding in a morphologically normal heart

b) Treated with intravenous digoxin to control ventricular rate

c) Should be treated with cardioversion in all cases

d) Difference in PR and HR can be used to diagnose

Question 15:

A patient presents with cough and hemoptysis and is diagnosed with pulmonary aspergillosis. He was treated with voriconazole. Which of the following is false about voriconazole?

a) It should be taken on an empty stomach

b) Therapeutic drug monitoring is required

c) It can be combined with pyrazinamide

d) A loading dose should be given

Question 16:

A patient with a history of sore throat was treated with antibiotics. He now presents with facial swelling. Urine examination shows eosinophilia and WBC casts. What is the likely diagnosis?

a) Acute interstitial nephritis

b) Lupus nephritis

c) Pyelonephritis

d) Post streptococcal glomerulonephritis

Question 17:

Which of the following is a worrisome feature in a patient presenting with headache?

a) 1,2,3

b) 1,2,3,4

c) 1,2,4

d) 3,4

Question 18:

Which of the following conditions is cannabinoid oil not indicated in?

a) Lennoux gustaut

b) Dravet syndrome

c) Tuberous sclerosis

d) Rett syndrome

Question 19:

ABG findings of a person presenting with abdominal pain and vomiting are as follows. pH 7.31, HCO3-20,Na 135, K+ 5, Cl- 92. Blood glucose is 450mg/dL. What is the interpretation?

a) High anion gap respiratory acidosis

b) Normal anion gap metabolic acidosis

c) High anion gap metabolic acidosis

d) Normal anion gap respiratory acidosis

Question 20:

A farmer presents with high grade fever, rash, vomiting, retro orbital pain and myalgia since 5 days is suspected to have dengue hemorrhagic fever. Which of the following investigations can be done for the diagnosis?

a) RTPCR for viral detection

b) NS1 antigen

c) IgM ELISA

d) Viral isolation of Aedes albopictus ■(C6/36) culture

Question 21:

Which of the following are features of upper motor neuron lesions?

a) 1, 2, 3

b) 1, 4

c) 3, 4, 5

d) 2, 5

Question 22:

Which of the following is not associated with celiac disease?

a) IgA transglutaminase positive

b) IgG antigliadin positive

c) Modified Marsh classification

d) HLA DQ6 and HLA DQ11

Question 23:

All of the following help to prevent the spread of COVID-19 except

a) Cough etiquette

b) Mask

c) Ivermectin

d) Vaccine

Question 24:

Which among the following viruses is primarily water-borne?

a) Viral hepatitis A

b) Viral hepatitis B

c) Viral hepatitis C

d) Viral hepatitis D

Question 25:

A patient presented with a blood pressure of 220/120 mmHg in an altered sensorium. He also had an episode of vomiting and a history of weakness on the left side of his body. Which of the following is incorrect regarding hypertensive management in this case?

a) In hemorrhagic stroke, reduce the BP to less than 160/100 mmHg with antihypertensives to prevent rebleed

b) In hemorrhagic stroke, aggressive use of IV antihypertensive if SBP > 220

c) In ER multiple blood pressure readings should be taken before starting anti-hypertensive therapy

d) In ischemic stroke, BP should be reduced to <140/90 mmHg immediately

Question 26:

Which of the following is not a feature of thrombotic thrombocytopenic purpura?

a) Thrombocytopenia

b) Thrombosis

c) Microangiopathic hemolytic anemia

d) Renal failure

Question 27:

Which of the following cannot help in differentiating psoriatic arthritis from other forms of arthritis?

a) Dactylitis

b) Presence of diarrhea

c) Enthesitis

d) Nail pitting

Question 28:

What is the mutation seen in Sickle cell anemia?

a) Glu---> Val(6)

b) Lys---> Arg

c) Glu--> lysine

d) Ala--> Gly (9)

Question 29:

A female patient with type 1 DM and hypothyroidism presented with chronic oral mucocutaneous candidiasis for 5 years. She is diagnosed with autoimmune polyendocrinopathy syndrome type 1. Which of the following is incorrect about this condition?

a) DM type-1 association is high

b) Starts in infancy

c) Hypoparathyroidism

d) High association with candidiasis

Question 30:

All of the following are examples of HAGMA except?

a) DKA

b) Diarrhoea

c) AKI

d) Inborn error of metabolism

Question 31:

LMN lesions are associated with?

a) 1,2

b) 1,2,3

c) 1,2,3,4

d) 1,4

Question 32:

In hemolytic anemia

a) 1, 3, 5

b) 1, 4

c) 2, 5

d) 1, 4, 5

Question 33:

A patient has undergone an acute renal allograft transplantation. What would indicate its rejection?

a) Any rise in serum creatinine above pre-transplant values is s/o acute renal allograft rejection

b) Rise of creatinine by more than 5% from the baseline is s/o acute renal allograft rejection

c) Serum creatinine does not rise in acute renal allograft rejection

d) Rise of creatinine by more than 10% from the baseline is s/o acute renal allograft rejection

Question 34:

A lady presents with recurrent bluing of her hand and stretching of her skin. She also has difficulty in swallowing and lung lesions. What is the pattern of ANA and the diagnosis?



a) Speckled pattern: Systemic sclerosis

b) Anti-centromere pattern: Systemic sclerosis

c) Nucleolar pattern: SLE

d) Homogenous pattern: SLE

Question 35:

A 60-year-old male patient complains of numbness in both lower limbs, On examination Rhomberg test is positive with broad-based gait and loss of proprioception of both lower limbs. He also has a history of low BP, hypothyroidism, and gastrectomy for gastric cancer a year back. Which of the following deficiencies can cause the current condition?

a) Vitamin B12 deficiency

b) Folic acid deficiency

c) Thiamine deficiency

d) Transferrin deficiency

Question 36:

A known case of ulcerative colitis is not responding to the anti-TNF alpha agent. Which JAK inhibitor is preferred for moderate to severe ulcerative colitis?

a) Vedolizumab

b) Tofacitinib

c) Ozanimod

d) Leflunomide

Question 37:

In children with GBS, all are true except?

a) Areflexia of the affected limb is a mandatory diagnostic criteria

b) Plasmapheresis is not preferred in refractory cases

c) Intravenous immunoglobulin is the first line of management

d) Albuminocytological dissociation in CSF is characteristic

Question 38:

As per KDIGO, which of the following falls under G3b?

a) 45-59

b) 60-89

c) 30-44

d) 15-29

Question 39:

A 46-year-old diabetic patient presented with fever, malaise, and a productive cough. His chest X-ray is given below. Which of the following is incorrect?



a) Patchy infiltrates as seen in pyogenic pneumonia

b) Interstitial infiltrates as seen in mycoplasma pneumonia

c) Hilar lymphadenopathy with infiltrates as seen in malignancy

d) Aspirational pneumonia is more common in the upper segment of the lower lobe and posterior segment of the upper lobe

Question 40:

A patient presented with uveitis, enthesitis, and sacroiliitis. Which of the following is most likely associated with this condition?

a) HLA-B27

b) HLA-B57

c) HLA-B51

d) HLA-B5

Question 41:

A 25-year-old male comes with hematuria. Which of the following are seen in glomerular hematuria?

a) 1 only

b) 1, 2

c) 1, 2, 3

d) 1, 2, 4

Question 42:

A male patient presented with oral ulcers and ulcers in the anal canal. He was diagnosed with Behcet's disease. Which of the following is not seen in this condition?

a) Ulcer over the dorsum of the penis

b) It is similar to Crohn's disease than ulcerative colitis

c) Bilateral panuveitis

d) Pathergy test positive is a criterion of ISG diagnostic criteria for Behcet's

Question 43:

The female patient comes to the ER with palpitations and syncopal attacks following exercise. She has had similar episodes in the past. An ECG was done as follows. What is the most probable diagnosis?

a) Uhl's disease

b) Arrhythmogenic Right Ventricular Dysplasia

c) Hypertrophic Cardiomyopathy

d) Brugada's syndrome

Question 44:

A 20-year-old male patient presented with a history of occasional palpitation, high BP, and headache. The patient was diagnosed with pheochromocytoma. Which of the following will be the preferred first investigation?

a) USG-abdomen

b) MIBG

c) MRI- abdomen

d) 24-hour urine metanephrines

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	b
6	a
7	d
8	c
9	a
10	b
11	a
12	d
13	a
14	c
15	c
16	a
17	a
18	d
19	c
20	c
21	d
22	d
23	c
24	a

25	d
26	d
27	b
28	a
29	a
30	b
31	d
32	b
33	d
34	b
35	a
36	b
37	b
38	c
39	a
40	a
41	d
42	a
43	b
44	d

Detailed Explanations

Solution to Question 1:

This patient's potassium is 2.8mEq/L indicating a hypokalemia. Cerebral edema is not a complication associated with hypokalemia, rather it is seen with hyponatremia.

Hypokalemia is defined as a plasma concentration of K+ <3.5 meq/L. Common causes include metabolic alkalosis, insulin infusions, diarrhea, use of diuretics, starvation, β2-agonists,magnesium deficiency, and primary hyperaldosteronism.

Hypokalemia manifests as systemic effects primarily targeting cardiac, skeletal, and intestinal muscles.

• In the cardiovascular system, hypokalemia poses a significant risk for arrhythmias, both ventricular and atrial. It can lead to ventricular tachycardia which can be fatal. (Option D)

• Electrocardiographic alterations characteristic of hypokalemia include the presence of broad flat T waves, ST segment depression, and QT interval prolongation, with pronounced changes observable when serum potassium levels drop below 2.7 mmol/L.

• Skeletal muscle function is also profoundly affected, as hypokalemia induces hyperpolarization, compromising the muscle's ability to depolarize and contract effectively. Consequently, individuals may experience weakness and paralysis. (Option C)

• Intestinal smooth muscle paralysis is another consequence of hypokalemia, potentially leading to intestinal ileus.

• Prolonged hypokalemia can precipitate acute kidney injury and, in chronic cases related to underlying conditions such as eating disorders or laxative abuse, contribute to the progression of end-stage renal disease. Hypokalemic polyuria occurs due to a combination of central polydipsia and an AVP-resistant renal concentrating defect, mimicking nephrogenic diabetes insipidus. This condition leads to the production of large volumes of dilute urine because the kidneys are unable to concentrate urine effectively in response to antidiuretic hormone (ADH).(Option A).

Solution to Question 2:

The given clinical scenario and the electrocardiogram (ECG) showing narrow QRS complexes suggest paroxysmal supraventricular tachycardia (PSVT). As the patient is hemodynamically unstable, the next step in management is DC (direct current) cardioversion.

PSVT is a narrow QRS complex tachycardia with sudden onset and termination. They mainly present with dizziness and palpitations. The evaluation of these patients includes monitoring their vitals and ECG.

Management is as follows:

1) When the patient is hemodynamically stable:

• Carotid sinus massage can be done. It is avoided if carotid bruit is present and if a prior history of stroke is seen

• Valsalva maneuver is done if the patient is cooperative

• Intravenous adenosine is useful in most cases. However, its usage is contraindicated in patients with prior cardiac transplantation. It can aggravate bronchospasm and atrial fibrillation, hence, it must be used with great caution in such patients.

• Intravenous calcium channel blockers such as verapamil or diltiazem and beta-blockers are helpful but can cause hypotension.



2) When the patient is hemodynamically unstable, i.e., hypotension with unconsciousness or respiratory distress, QRS-synchronous direct current cardioversion is done.



Solution to Question 3:

The patient presents with chest pain and breathlessness following a recent viral illness, and her chest X-ray shows a "water bottle" sign, indicating a pericardial effusion. In cases of significant pericardial effusion or cardiac tamponade, heart sounds, including the first heart sound S1, are typically muffled due to the insulating effect of the accumulated fluid around the heart. This makes it difficult to hear the heart sounds clearly, hence there won't be a loud S1.

Pericardial effusion is usually associated with chest pain and/or the ECG changes of pericarditis (widespread ST elevation, often with upward concavity). If the effusion is large, the ECG may show electrical alternans. Heart sounds may be faint, friction rub, and the apical impulse may not

be audible.

Echocardiography is a widely used imaging technique that allows localization and estimation of the quantity of pericardial fluid.

The chest X-ray shows an enlargement of the cardiac silhouette. It may be normal in small effusions. The lateral radiograph may show the oreo cookie sign. It is a vertical opaque line (pericardial fluid) separating a vertical lucent line (paracardial fat) directly behind the sternum anteriorly from a similar lucent vertical line (epicardial fat) posteriorly.

CT and MRI are indicated when echocardiography is inconclusive, when the effusion is loculated or hemorrhagic or if there is pericardial thickening.

Other options:

Option B: Pulsus paradoxus may be seen: This statement is true. Pulsus paradoxus, an exaggerated decrease in systolic blood pressure during inspiration, is commonly seen in patients with massive pericardial effusions.

Option C: Emergency pericardiocentesis may be required: This statement is true. In cases of significant pericardial effusion or cardiac tamponade, emergency pericardiocentesis is often necessary to relieve pressure on the heart and prevent hemodynamic compromise.

Option D: ECHO showing diastolic collapse of RA and RV wall during expiration requires immediate intervention: This statement is true. Echocardiographic evidence of diastolic collapse of the RA and RV is a sign of cardiac tamponade, which necessitates immediate intervention.

Solution to Question 4:

This patient with a history of allergy to penicillin and a diagnosis of enterococcal endocarditis should receive vancomycin for the treatment of his endocarditis.

Enterococci are one of the major hospital-acquired pathogens, producing various infections like:

• Urinary tract infections

• Bacteremia and mitral valve endocarditis (in intravenous drug abusers)

• Intrabdominal, pelvic and soft tissue infections

• Late-onset neonatal meningitis and sepsis

Strains resistant to penicillin and other antibiotics occur frequently, so it is essential to perform antibiotic sensitivity for proper therapy. Enterococci are intrinsically resistant to cephalosporins(Option D) and offer low-level resistance to some aminoglycosides.

In penicillin-sensitive strains, synergism occurs with combination treatment with penicillin and aminoglycosides. In case of high-level resistance, the choice of drug is vancomycin. However, in infections caused by E. faecalis ampicillin is used as a first-line antibiotic as resistance is rare.

Recently vancomycin-resistant enterococci (VRE) have begun to emerge. These are treated with linezolid, streptogramins (active against E.faecium, but not to E.faecalis) and daptomycin.

Other options:

Option B: Piperacillin would be contraindicated in a patient with a penicillin allergy as there is cross-reactivity between the two drugs.

Option C: Aztreonam can be used in patients who have enterococci resistant to vancomycin. However, vancomycin is preferred for empiric therapy due to a better side effect profile.

Solution to Question 5:

Serum potassium <3.5mEq/L is not a diagnostic criterion of diabetic ketoacidosis.

DKA is an acute complication of type 1 diabetes mellitus. It is usually precipitated by absolute or relative insulin deficiency caused by increased metabolic demand (infections) or insulin noncompliance.

The common symptoms include nausea, vomiting, and severe abdominal pain. The common signs include Kussmaul breathing, fruity odor of the breath, dehydration, tachycardia, hypotension, and delirium.

The diagnosis of DKA is made based on the following parameters:

• Hyperglycemia > 250 mg/dl

• Ketosis or ketones in urine

• Metabolic acidosis with pH < 7.3

• HCO3 < 15-18 mEq/L

• High anion gap due to ketoacids and lactic acidosis

Hyponatremia is seen as a result of hyperglycemia (osmotic shift causes dilutional hyponatremia). Normal sodium levels on admission indicate severe dehydration. Potassium levels at presentation may be normal or elevated despite total body deficit (due to intracellular shift).

Solution to Question 6:

The patient presents with recurrent kidney stones, abdominal pain, and altered mentation, along with laboratory findings of decreased serum calcium (Ca^{2+}) and elevated parathyroid hormone (PTH). These symptoms and laboratory results are indicative of renal osteodystrophy, a metabolic bone disorder associated with chronic kidney disease (CKD).

Renal osteodystrophy is a complex condition that results from the kidneys' inability to maintain the proper levels of calcium and phosphate in the blood, leading to secondary hyperparathyroidism. In CKD, the kidneys are not able to convert vitamin D to its active form, calcitriol, which is necessary for calcium absorption in the intestines. The reduced levels of calcitriol lead to decreased calcium absorption, resulting in hypocalcemia. The body responds to hypocalcemia by increasing PTH secretion from the parathyroid glands in an attempt to normalize serum calcium levels.

The elevated PTH, known as secondary hyperparathyroidism, causes increased bone resorption to release calcium and phosphate into the bloodstream, leading to bone pain and fragility, and

contributing to the formation of kidney stones due to the elevated phosphate and calcium levels in the urine. Additionally, the ongoing calcium and phosphate imbalance can cause vascular calcifications and other systemic complications leading to hypertensive crises, contributing to the patient's altered mentation and other symptoms.

Therefore, the constellation of symptoms and laboratory findings in this patient points to renal osteodystrophy, a metabolic bone disorder secondary to chronic kidney disease, rather than primary hyperparathyroidism, hypothyroidism, or pheochromocytoma.

Other options:

Option B: Primary hyperparathyroidism would typically present with hypercalcemia due to autonomous overproduction of parathyroid hormone. In this case, the patient has decreased serum calcium, which is more indicative of secondary hyperparathyroidism due to chronic kidney disease.

Option C: Hypothyroidism generally presents with symptoms such as fatigue, weight gain, cold intolerance, and dry skin. It does not typically cause elevated parathyroid hormone levels or the specific constellation of symptoms seen in this patient (recurrent kidney stones, abdominal pain, and altered mentation).

Option D: Pheochromocytoma is a tumor of the adrenal gland that secretes excessive catecholamines, leading to symptoms such as hypertension, palpitations, and episodic headaches. It does not cause hypocalcemia or elevated parathyroid hormone levels, and it is not associated with kidney stones or the other symptoms described in this patient.

Solution to Question 7:

The ABG values indicate a metabolic alkalosis because of the significantly elevated bicarbonate level (HCO_3^- 44 mEq/L, normal 22-28mEq/L) and a secondary respiratory compensation (increased pCO_2 to 50 mmHg, normal 35-45 mmHg). The elevated pH (7.51, normal 7.35-7.45) confirms the alkaline state.

The compensation is not complete, as the pH remains above normal, indicating a partially compensated metabolic alkalosis.

Solution to Question 8:

The patient's presentation of chest pain for 1 hour, perspiration, and anxiety, combined with ECG findings showing ST depressions in leads V2 and V3, suggests a possible diagnosis of posterior wall MI. Posterior wall MI can present with ST depressions in the anterior leads (V1-V3) due to the reciprocal changes associated with ischemia in the posterior wall of the heart. These depressions indicate that the infarction is not directly visible in the standard ECG leads but is affecting the posterior part of the heart.

Posterior wall MI is typically caused by occlusion of the circumflex artery or the posterior descending artery, which supplies the posterior aspect of the heart. Management of posterior wall MI involves immediate medical intervention, including the administration of antiplatelet agents, anticoagulants, nitrates, beta-blockers, and, in many cases, reperfusion therapy through

percutaneous coronary intervention (PCI) or thrombolysis to restore blood flow to the affected area of the heart.

Other options:

Option A and B: NSTEMI (Non-ST segment elevation myocardial infarction) could present with ST depression, but it typically affects a wider range of leads and is less likely to present with such a specific pattern of depression isolated to V2 and V3. Unstable angina, characterized by chest pain at rest or with minimal exertion, can cause ST depression, but it usually does not present with the level of severity and specific pattern seen in this case, particularly without elevation in cardiac biomarkers.

Option D: Acute myocarditis, while it can present with chest pain and ECG changes, usually shows diffuse changes, not isolated to specific leads like V2 and V3. It typically involves diffuse ST elevation or PR depression across multiple leads. Therefore, the combination of acute symptoms and specific ECG changes in leads V2 and V3 strongly suggests that a posterior wall MI is the most likely diagnosis for this patient.

Solution to Question 9:

Chronic venous insufficiency is the least common cause of bipedal edema. This is because, the other conditions listed cause edema that can affect both limbs at the same time due to decreased albumin concentration or fluid retention. However, venous insufficiency is not as commonly associated with bilateral lower limbs and is more likely to present as unilateral lower extremity edema.

Lower extremity edema is the accumulation of fluid in the tissues of the legs, leading to swelling. It can result from various conditions affecting the circulatory, lymphatic, or renal systems.

Common causes include:

• Venous Insufficiency: Chronic venous insufficiency (CVI) occurs when veins in the legs do not efficiently return blood to the heart, leading to pooling of blood and fluid leakage into surrounding tissues. However, this edema is usually unilateral depending on the limb that is worse effected.

• Heart Failure: When the heart is unable to pump blood effectively, it can cause fluid buildup in the legs.

• Kidney Disease: Impaired kidney function can lead to fluid retention and swelling in the lower extremities.

• Liver Disease: Conditions like cirrhosis can cause decreased production of proteins like albumin, leading to fluid leakage into tissues.

• Lymphedema: Blockage or damage to the lymphatic system can prevent proper drainage of lymphatic fluid, causing swelling.

• Medications: Certain drugs, such as calcium channel blockers, nonsteroidal anti-inflammatory drugs (NSAIDs), and corticosteroids, can cause fluid retention.

• Infections: Conditions like cellulitis can cause localized swelling and inflammation.

• Pregnancy: Increased pressure on the veins and hormonal changes can lead to edema.

• Prolonged Standing or Sitting: Lack of movement can cause fluid to accumulate in the legs due to gravity.

Treatment: The treatment of lower extremity edema focuses on addressing the underlying cause and alleviating symptoms. This can include lifestyle modifications such as elevating the legs above heart level to reduce swelling, wearing compression stockings to improve venous return and reduce fluid accumulation, and regular exercise to promote circulation.

In systemic conditions, the underlying cause should be treated to improve the edema.

Solution to Question 10:

The given clinical features are suggestive of paralysis of the facial nerve also known as Bell's palsy. Treatment for this condition is with acyclovir, eye patching and eye ointments to prevent desiccation.

Bell's palsy is an acute onset idiopathic, peripheral facial paralysis. While in most cases the cause is unknown, there can be exposure to the herpes virus in some patients. It is a lower motor neuron lesion of the facial nerve and involves the ipsilateral side of the upper and lower halves of the face.

It presents with an inability to close the eye, and if the patient attempts to close the eye, it turns up and out. This is known as the Bell's phenomenon. Other features include drooling of saliva and facial asymmetry. There may be noise intolerance and a loss of taste sensation.

Bell's palsy is a diagnosis of exclusion. All other possible causes of facial paralysis should be ruled out by investigations.

Treatment of Bell's palsy involves prednisone 60-80 mg daily during the first 5 days and then tapered over the next 5 days. Acyclovir can also be used in patients who are suspected to have exposure to the herpes virus. Tape or eye masks to depress the upper eyelid during sleep and artificial tears can be used to prevent corneal drying. Most patients recover within a few weeks or months.

Solution to Question 11:

This patient has elevated fasting glucose values indicating a diagnosis of diabetes mellitus. He also has reduced ejection fraction, elevated blood pressure with signs of orthopnea(breathlessness on lying down) indicating heart failure. Dapagliflozin, which is a sodium-glucose transporter-2 (SGLT-2) inhibitor used for diabetes has been recently approved for use in patients with heart failure with reduced ejection fraction and is the preferred drug in this patient as it can help manage both his conditions.

Dapagliflozin works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) in the proximal renal tubules resulting in reduced reabsorption and increased urinary excretion of glucose. The glucose-lowering effect is independent of insulin. Dapagliflozin acts as a diuretic and natriuretic by reducing sodium and volume load, causing intravascular contraction. Dapagliflozin is also associated with weight loss, with reductions in blood pressure without increasing heart rate. Dapagliflozin has been found to decrease the risk of mortality from cardiovascular causes in

diabetic patients. SGLT-2 inhibitors have a cardioprotective effect resulting from improved cardiac cell metabolism, improvement of ventricular loading mechanism, reduction of cardiac cell necrosis and fibrosis. It also has a renoprotective property.

Side effects of SGLT-2 inhibitors include dehydration, hyperkalemia, worsening of diabetic ketoacidosis, increased risk of fractures, leg and foot amputation with canagliflozin, and a possible increased risk of bladder cancer with dapagliflozin. Dapagliflozin has also been associated with an increased risk of breast cancer.

Other options :

Option B: Liraglutide is a GLP 1 receptor agonist used for the management of diabetes mellitus. It has recently been approved for use in obesity. However, it has no role in managing heart failure.

Option C: Glimepiride is a sulfonylurea that is used in patients who are not sufficiently controlled with metformin for diabetes mellitus. However, it has no role in managing heart failure.

Option D: Pramlintide is a subcutaneous amylin analogue used in the management of diabetes mellitus. It has no role in the management of heart failure.

Solution to Question 12:

Amnesia, polyneuropathy and confabulation are seen in Korsakoff's psychosis. Ophthalmoplegia is seen in Wernicke's disease. Peripheral neuropathy is a consequence of B1 deficiency which is the vitamin deficit in Korsakoff psychosis.

Korsakoff's psychosis is a chronic amnestic syndrome that can follow Wernicke's encephalopathy due to thiamine deficiency (secondary to alcoholism). The cardinal features are impaired mental syndrome and anterograde amnesia. Only 20% of patients recover with treatment (whereas in Wernicke's encephalopathy, there is complete reversal with treatment).

Clinical features:

• Impaired mental syndrome - especially recent memory

• Striking inability to form new memories - severe anterograde amnesia

• Retrograde amnesia is less severe.

• Present with confabulation as an attempt to cover the memory deficit

• There is a lack of insight and no intellectual functional loss.

• Overall intact intelligence

• Implicit learning (e.g., learning procedures) is preserved.

• Long-term and immediate memory is normal.

Treatment: Thiamine is given 100mg orally two to three times daily for 3 to 12 months.

Solution to Question 13:

The presence of gastrointestinal bleeds is an indication for corticosteroid therapy in IgA vasculitis/Henoch Schonlein Purpura.

Corticosteroids have improved outcomes in patients who develop severe GI complications including GI bleeds and intractable abdominal pain in patients with HSP. This is because it can reduce the intensity of disease and improve outcomes by helping resolve the inflammation. The other indication for the use of corticosteroids is when the patient develops crescentic glomerulonephritis, ie rapidly progressive glomerulonephritis. While corticosteroids can be used for arthritis and abdominal pain, there is limited evidence of their efficacy. Thus it is strictly reserved for severe GI and renal involvement.

Henoch-Schönlein purpura (HSP), also known as IgA vasculitis, is a common small vessel vasculitis. It is a type III hypersensitivity reaction characterized by the deposition of IgA in the small vessels of the skin, joints, gastrointestinal tract, and kidney resulting in vasculitis. The exact etiology of HSP remains unknown. But, it usually occurs following an upper respiratory tract infection caused by group A Streptococcus, Staphylococcus aureus, Mycoplasma, and Adenovirus.

Patients usually present with the following features:

• Palpable purpura is symmetric and occurs in the lower extremities, extensor aspect of the upper limbs, and buttocks.

• Arthralgia and arthritis- commonly affecting hip, knee, or ankle joints.

• Gastrointestinal manifestations like abdominal pain, vomiting, diarrhea, intestinal perforation and intussusception.

• About 30% of the patients with HSP develop renal manifestations, presenting as microscopic hematuria, hypertension, and progressive glomerulonephritis.

Nephritis due to IgA vasculitis should be treated with immunosuppressants like prednisolone and azathioprine.

The image given below shows palpable purpura in the lower limbs seen in HSP.

Solution to Question 14:

In this patient presenting with palpitations, and irregular RR intervals with absent P waves on ECG, the most likely diagnosis is atrial fibrillation. The false statement is that atrial fibrillation should be treated with cardioversion in all cases.

Cardioversion, whether electrical or pharmacological, is not always appropriate for every patient with AF. The decision to perform cardioversion depends on several factors, including the duration of the AF episode, the presence of underlying heart disease, the patient's symptoms, and the risk of thromboembolism. In some cases, rate control and anticoagulation may be the preferred treatment approach rather than attempting to restore sinus rhythm immediately. Therefore, while cardioversion is a treatment option for some patients with AF, it is not universally indicated for all cases.

Atrial fibrillation (AF) is a supraventricular tachycardia caused by ectopic foci or small reentry circuits within the atria. It has an atrial rate of 400-600/min and a ventricular rate of 100-160/min.

Risk factors for developing AF include:

• Age- usually above 60 years of age

• Underlying heart disease

• Hypertension

• Diabetes mellitus

• Obesity

• Sleep apnea

• Post-cardiac surgery

• Inflammatory pericarditis

• Acute alcohol consumption

Features include heart rate typically ranges from 110-160 beats/minute. It is irregular and rapid. Presyncope and syncope can be seen. Thrombus formation can occur and thromboembolic events can take place.

Typically, ECG shows an absent P wave and irregular RR intervals.

Management is mainly by control of rate and rhythm, avoiding the thromboembolism risk and modifiable risk factors.

Other options:

Option A: Atrial fibrillation is usually seen in a patient with underlying heart disease and is a rare occurrence in morphologically normal hearts.

Option B: Intravenous digoxin can be used to control the ventricular rate in atrial fibrillation, especially in patients with heart failure or those who are not suitable for beta-blockers or calcium channel blockers. However, digoxin is not always the first-line treatment for rate control.

Option D: The difference between the pulse rate (PR) and the heart rate (HR) can indeed be used to diagnose atrial fibrillation. This discrepancy, known as a pulse deficit, occurs because not all heartbeats result in a palpable pulse due to the irregular and often rapid ventricular response.

Solution to Question 15:

Voriconazole should not be combined with pyrazinamide. This is because, both these drugs are hepatotoxic and can cause fulminant liver failure if combined together. Thus, this is false with regards to voriconazole.

Voriconazole is an azole antifungal used in the management of pulmonary aspergillosis. It is also the drug of choice for invasive aspergillosis, disseminated infections caused by fluconazole-resistant Candida, Fusarium infections, and febrile neutropenia not responding to antibacterial therapy.

Patients who are started on voriconazole should initially receive a loading dose of 6mg/kg body weight for 24 hours followed by 3-4mg/kg body weight for 12 hours. After this, patients can be shifted to oral administration of the drug. Voriconazole is metabolized by the cytochrome p450 group of enzymes and thus its blood levels should be monitored. This is especially indicated in patients with liver disease to adequately titrate the drug dosage.(Option B). Patients are advised to take the drug on an empty stomach as fatty meals reduce the absorption of the drug(Option A)

Other adverse effects of voriconazole include rashes, photosensitivity, transient visual or auditory hallucinations and anaphylactoid reactions. Intravenous voriconazole should be used with caution in patients with renal failure.

Solution to Question 16:

In this patient with a history of antibiotic use, after a sore throat presenting with facial swelling and WBC casts in the urine, the likely diagnosis is acute interstitial nephritis(AIN) likely secondary to penicillin use.

AIN usually occurs as an allergic response to certain medications. Offending agents include beta-lactam antibiotics like methicillin, nonsteroidal anti-inflammatory drugs, proton pump inhibitors, sulfonamide derivatives, diuretics, antiepileptics (phenytoin, valproate), and sulfasalazine.

Individuals with AIN are usually asymptomatic and are incidentally found to have elevated creatinine. However, in some instances, they can develop fever and rash. In severe cases, renal failure with proteinuria can occur.

Laboratory studies may show peripheral eosinophilia and urinary eosinophilia. However, these are not specific or sensitive enough to serve as diagnostic criteria.

Treatment depends on how severe AIN is. Milder cases can be treated by stopping the offending drug. More severe cases require hospitalization and follow-up.

Solution to Question 17:

Statements 1,2,3 are indications for further evaluation of a headache

In a patient presenting with a headache, the following features are worrisome:

• Tenderness over the temporal area: This is a worrisome feature because it may indicate temporal arteritis (giant cell arteritis), a condition that requires prompt diagnosis and treatment to prevent complications such as vision loss.

• Subacute onset, worsening over 2 weeks: This is concerning as it suggests a progressive underlying condition, which could be indicative of serious issues such as a tumor, an expanding subdural hematoma, infection, or other intracranial processes that require further investigation.

• Age of 50 years: Age is a significant factor because the incidence of secondary headaches due to serious underlying causes increases with age. Conditions such as temporal arteritis and intracranial neoplasms are more common in older adults.

In contrast, a history of cocaine abuse is not specifically a worrisome feature in the context of the headache itself. While cocaine abuse can lead to headaches due to its vasoconstrictive properties and potential to cause hypertensive emergencies or strokes, it is not a primary red flag feature for serious underlying causes of headaches in the same way as the other listed factors. However, a history of cocaine abuse does warrant attention for other potential health risks and complications associated with drug use.

Solution to Question 18:

Cannabinoid oil is not indicated in Rett syndrome.

The FDA recently approved cannabidiol oral solution for the treatment of seizures associated with tuberous sclerosis complex in patients one year of age and older. It was previously approved for the treatment of seizures associated with Lennox-Gastaut syndrome and Dravet syndrome.

Cannabinoid oil is an oral derivative of the cannabis plant. The mechanism by which it improves seizure threshold is not understood yet and there are various studies ongoing to survey its use. The cannabinoid receptors are G protein-coupled receptors located predominantly in the presynaptic nerve terminals.

The CB1 receptor is located in the central nervous system. Activation of CB1 receptor results in euphoria and alleviation of pain (anti-nociceptive effect). The CB2 receptor is located mainly in the periphery and their activation does not have a euphoric effect.

The endogenous cannabinoid system is a potential target for use in the treatment of chronic pain. The most common adverse effects of CBD at doses used for treating epilepsy include somnolence, fatigue, decreased appetite, diarrhea, and elevated hepatic transaminases. The elevations in liver transaminases are dose-related.

Solution to Question 19:

The ABG findings of a person presenting with abdominal pain and vomiting, showing a pH of 7.31, HCO_3^- of 20 mEq/L, Na^+ of 135 mEq/L, K^+ of 5 mEq/L, Cl^- of 92 mEq/L, and a blood glucose level of 450 mg/dL, with an anion gap of 23 indicate a high anion gap metabolic acidosis likely secondary to a diabetic ketoacidosis state.

Anion Gap=Na+ – (Cl- + HCO3-), so Anion Gap=135–(92+20)=135–112=23

The low pH and low bicarbonate confirm acidemia with a metabolic component. The significantly elevated blood glucose level, along with symptoms of abdominal pain and vomiting, strongly suggests diabetic ketoacidosis (DKA) as the underlying cause.

Solution to Question 20:

The given clinical scenario is of a farmer with suspected dengue hemorrhagic fever. As he has presented 5 days after the onset of symptoms, the most appropriate test for him would be IgM ELISA. NS1 antigen is the preferred test had he presented before 5 days.

Dengue is an acute febrile illness caused by the dengue virus that is transmitted by the Aedes aegypti mosquito.

It presents as sudden-onset fever, retroorbital pain, myalgia, arthralgia, and a rash. The initial rash in dengue may appear as flushing erythema of the face in the first 2 days of symptoms. A second rash occurs 3-6 days after the onset of fever and appears maculopapular or morbilliform. Sometimes, lesions may coalesce and appear as 'white islands in a red sea' due to generalized erythema with petechiae and islands of sparing.

Before day 5 of illness, during the febrile period, dengue infections may be diagnosed by detection of viral antigens like non-structural viral protein antigen(NS 1) by ELISA or rapid tests. This is considered the mainstay of diagnosis.

Virus isolation in cell culture and viral RNA detection by nucleic acid amplification tests (NAAT) can also be done.

Serological tests, such as IgM ELISA, are performed after day 5 when dengue viruses and antigens are no longer present in the blood due to the appearance of specific antibodies.

Solution to Question 21:

Spasticity and exaggerated reflexes are features of an upper motor neuron lesion.

Fasciculations and severe muscle atrophy are features of lower motor neuron lesions.

Rigidity is a feature of extrapyramidal lesions.

Solution to Question 22:

HLA-DQ2 and DQ8 are associated with celiac disease, not HLA DQ6 and HLA DQ11.

Celiac disease or celiac sprue or gluten-sensitive enteropathy is triggered by the ingestion of gluten-containing foods such as barley, rye, oats, and wheat (mnemonic: BROW) in genetically predisposed individuals. It usually affects the 2nd part of the duodenum or proximal jejunum.

Diagnosis:

• Tissue Transglutaminase Ab (tTG-IgA) - single best screening test (Option A)

• IgG Anti-gliadin antibodies (Option B) are produced in response to gliadin, a component found in wheat. These are associated with celiac disease along with other antibodies like IgA Anti-tissue transglutaminase antibodies and Anti-endomysial antibodies.

• HLA-DQ2 or DQ8- HLA testing is useful for ruling out celiac disease.

• Endoscopy with small bowel biopsy is used as a confirmatory test to diagnose celiac disease and not as an initial test in suspected cases.

• The histopathology is characterized by villous atrophy, crypt hyperplasia, and increased intraepithelial CD8+T lymphocytes. An increase in the number of intraepithelial lymphocytes is a sensitive marker of celiac disease, even in the absence of epithelial damage and villous atrophy.

• Modified MARSH's Score (Option C) is used to assess celiac disease histologically.



Solution to Question 23:

Ivermectin does not help prevent the spread of COVID-19.

Basic protective measures to prevent the spread of COVID-19:

• Wash your hands frequently for at least 20 seconds.

• Maintain social distancing- Maintain at least 1 metre distance between yourself and anyone who is coughing or sneezing wearing a face mask (Option B)

• Avoid touching eyes, nose and mouth

• Practice respiratory hygiene (cough etiquette) - cover your mouth and nose with your bent elbow or tissue when you cough or sneeze, then dispose of the used tissue immediately. (Option A)

• If you have a fever, cough, and difficulty breathing, seek medical care early

• Stay informed and follow the advice given by your healthcare provider

• Vaccination (Option D)

COVID-19 vaccines approved and in development:

Vaccines	Mechanism
Pfizer/BioNtech	mRNA based vaccine
Moderna	mRNA based vaccine
Oxford University/ Astra Zeneca/ Serum Institute of India (Covishield)	Non-replicating viral vector
Gamaleya (Sputnik V)/ Dr.Reddys	Non-replicating viral vector
Johnson and Johnson	Non-replicating viral vector
Bharat biotech/NIV (Covaxin)	Inactivated vaccine
Zydus Cadila (ZyCoV-D)	Plasmid DNA vaccine
Novavax	Protein subunit

Solution to Question 24:

Viral hepatitis A is primarily water-borne.

Hepatitis A and E are transmitted by fecal-oral route (waterborne). Overcrowding, poor hygiene, and faecal contamination of water supplies, such as after monsoon flooding, can lead to the spreading of the disease.

Hepatitis B, C, and D are transmitted by parenteral routes, e.g. blood transfusions.

Solution to Question 25:

Option D is incorrect regarding hypertensive management in this case. In ischemic stroke, BP should be reduced to <185/110 mmHg immediately before doing thrombolysis.

In ischemic stroke, blood pressure is generally not aggressively lowered immediately unless it is extremely high (usually >220/120 mmHg), because maintaining higher blood pressure can help ensure adequate perfusion of the brain tissue and reduce the risk of worsening ischemia. Before performing thrombolysis in ischemic stroke, BP must be reduced to <185/110mm Hg immediately, and not less than <140/90 mmHg. If the BP is reduced to <140/90 mmHg, it would further compromise the cerebral perfusion and can increase the infarct size. The goal is typically to avoid lowering BP too quickly, and a more gradual reduction is preferred.

Other options

Option A: In hemorrhagic stroke, BP should be reduced to less than 160/100 mmHg with antihypertensives to prevent rebleed.

Option B: In the case of hemorrhagic stroke, if the systolic blood pressure (SBP) is greater than 220 mmHg, aggressive lowering of blood pressure with IV antihypertensives is recommended to reduce the risk of further bleeding.

Option C: In the emergency room, multiple blood pressure readings should be taken to accurately assess the patient's blood pressure before initiating any antihypertensive therapy, especially in a setting like hypertensive emergencies.

Solution to Question 26:

Renal failure is not a feature of thrombotic thrombocytopenic purpura

Renal injury can occur in TTP due to microvascular injury, but it is not a defining feature of TTP, whereas the other three features (thrombocytopenia, thrombosis, microangiopathic hemolytic anaemia) are key components of the disease.

Thrombotic thrombocytopenic purpura (TTP) is a rare hematological disorder primarily characterized by thrombocytopenia, microangiopathic hemolytic anaemia, neurological symptoms, and sometimes fever. It is typically associated with the following:

• Thrombocytopenia

• Thrombosis

• Microangiopathic hemolytic anaemia

Other options

Option A: Thrombocytopenia is a hallmark feature of TTP. Platelets are consumed in the formation of microthrombi, leading to a low platelet count.

Option B: Thrombosis: TTP involves the formation of microthrombi (not large thrombosis) in small vessels, particularly in organs like the kidneys, brain, and heart. These microthrombi lead to other characteristic features, including hemolysis and thrombocytopenia.

Option C: Microangiopathic hemolytic anemia is another key feature of TTP. The microthrombi damage red blood cells as they pass through small blood vessels, resulting in hemolysis (destruction of red blood cells).

Solution to Question 27:

The presence of diarrhea is not a characteristic symptom of psoriatic arthritis, therefore, it cannot help in differentiating the condition from other forms of arthritis.

Psoriatic arthritis can sometimes be associated with inflammatory bowel disease causing diarrhea. However, diarrhea is not specific to psoriatic arthritis and can occur in other types of arthritis such as reactive arthritis and certain types of infectious arthritis.

Psoriatic arthritis is an inflammatory type of arthritis mainly affecting the distal interphalangeal joints. In most cases, psoriasis precedes arthritis. Other features include nail changes (pitting, horizontal ridging, onycholysis, yellowish discoloration, hyperkeratosis), dactylitis, enthesitis, and

tenosynovitis. The Classification of Psoriatic Arthritis (CASPAR) criteria are a useful tool for early diagnosis of psoriatic arthritis.

Other options:

Option A: Dactylitis is a characteristic feature of psoriatic arthritis, referring to the inflammation of an entire finger or toe. It is not commonly seen in other types of arthritis.

Option C: Enthesitis (inflammation of the entheses) is a hallmark of psoriatic arthritis and is less common in other forms of arthritis like rheumatoid arthritis.

Option D: Nail pitting (small depressions on the surface of the nails) is a feature strongly associated with psoriatic arthritis and is rarely seen in other forms of arthritis.

Solution to Question 28:

Glu---> Val(6) mutation is seen in Sickle cell anemia.

Sickle cell anemia is an autosomal recessive disorder that occurs due to a missense mutation (a type of point mutation) in the β-globin gene of HbS, which results in the replacement of polar glutamate residue with non-polar valine at the 6th position, generating a hydrophobic "sticky patch" on the surface of the β subunit of deoxygenated HbS.



Both HbA and HbS contain a complementary sticky patch on their surfaces that is exposed only in the deoxygenated state. Thus, at low Po2, deoxyHbS polymerizes to form long, twisted helical fibers. This distorts the biconcave RBC into a sickle shape. The precipitated HbS also damages the RBC membrane, so the cells become highly fragile and get easily hemolyzed leading to anemia.



The major pathophysiology involves chronic hemolysis and microvascular occlusion. Hypoxia increases the concentration of deoxygenated HbS promoting RBC sickling. These sickle cells can disrupt circulation causing vascular occlusion and tissue infarction. Vaso-occlusive crises, also known as pain crises, are a result of this hypoxic injury and infarction. These crises predominantly affect the bones, lungs, liver, brain, spleen, and penis.

Solution to Question 29:

Diabetes mellitus type 1 usually occurs in APS-2 and is less commonly associated with APS-1.

This clinical scenario is suggestive of autoimmune polyendocrinopathy (APS) type 1. It is also called APECED (autoimmune polyendocrinopathy, candidiasis, and ectodermal dystrophy).

It is caused by mutations in the autoimmune regulator (AIRE) gene. This gene usually increases the tolerance of central T-cells to peripheral tissue antigens. When this function is lost, the T-cells become more prone to mount immune reactions against tissue antigens, promoting autoimmunity. These patients develop autoantibodies against IL-17 and IL-22, which are crucial for defence against fungal infections. APS type 1 usually presents in childhood. It can also start in infancy (option B) and the presentation can evolve over time.

APS type 1 presents in childhood with the following features:

• Chronic mucocutaneous candidiasis (option D)

• Ectodermal dystrophy (abnormalities of the skin, dental enamel, and nails)

• Organ-specific autoimmune disorders such as -

• Autoimmune adrenalitis

• Autoimmune hypoparathyroidism (option C)

• Idiopathic hypogonadism

• Pernicious anemia

Autoimmune polyendocrine syndrome (APS) type 2 usually presents in early adulthood as a combination of the following:

• Adrenal insufficiency

• Autoimmune thyroiditis or type 1 diabetes

Solution to Question 30:

Diarrhoea is an example of NAGMA (Normal anion gap metabolic acidosis), not HAGMA (High anion gap metabolic acidosis).

Other three options, i.e., DKA, AKI and inborn error of metabolism are examples of HAGMA.

Solution to Question 31:

Lower Motor Neuron lesions result in flaccid paralysis, characterized by decreased muscle tone (hypotonia) and fasciculations.

The hallmark signs of Lower Motor Neuron lesions include:

• Flaccid paralysis: Decreased muscle tone (hypotonia) and reflexes (hyporeflexia).

• Muscle atrophy: Due to denervation.

• Fasciculations: Due to spontaneous activity in motor neurons.

• Absent Babinski reflex.

Sign	Upper Motor Neuron	Lower Motor Neuron
Atrophy	None	Severe
Fasciculations	None	Common
Tone	Spastic	Decreased
Distribution of weakness	Pyramidal/ regional	Distal/ segmental
Muscle stretch reflexes	Hyperactive	Hypoactive/ absent
Babinski sign	Present	Absent

Solution to Question 32:

Statements 1 and 4 are true regarding hemolytic anemia.

In hemolytic anaemia, the following features are typically observed:

• Indirect Van den Bergh test is positive (statement 1). This indicates an increase in unconjugated (indirect) bilirubin, which is a hallmark of hemolysis.

• Urobilinogen is high (statement 4). Increased breakdown of haemoglobin leads to more bilirubin being converted to urobilinogen in the intestines, which is then absorbed and excreted in urine.

Van den Bergh test can be used to determine the serum bilirubin level in a patient with jaundice. Van den Bergh's reaction is a colourimetric method to measure serum bilirubin. It is based on the formation of red-purple azodipyrroles when diazotised sulfanilic acid reacts with bilirubin.

Other options

Statement 2: Unconjugated (indirect) bilirubin is elevated in hemolytic anaemia,

Statement 3: Raised serum transaminases are usually associated with liver damage, not hemolysis.

Statement 5: Gamma-glutamyl transferase (GGT) elevation is related to liver or biliary tract pathology, not hemolytic anaemia.

Clinical condition	Reaction with reagent	Interpretation
Normal serum	No reaction	Negative
High conjugated bilirubineg: Obstructive jaundice	Reacts in the absence of met hanol	Direct positive
High unconjugated bilirubine g:Hemolytic anaemia	Reacts in the presence of met hanol to give total bilirubin	Indirect positive
High-conjugated and unconj ugated bilirubineg: Hepatic j aundice	Reacts strongly both with an d without methanol	Biphasic reactio n

Solution to Question 33:

A rise in serum creatinine by more than 10% from the baseline is suggestive of acute renal allograft rejection.

Acute renal allograft dysfunction is defined as a rise in serum creatinine of >10% from baseline or an absolute rise of ≥20 μmol/L which requires detailed investigations. The causes of early graft dysfunction include:

• Acute rejection (antibody mediated or cell mediated)

• Calcineurin inhibitor toxicity

• Dehydration

• Urinary tract infection or pyelonephritis

• Sepsis

• Renal vein or renal artery thrombosis

• Ureteric obstruction or urine leak

If graft dysfunction's cause is unclear, a percutaneous needle-core biopsy should be performed to confirm allograft rejection.

Solution to Question 34:

Clinical and immunofluorescence patterns are suggestive of a diagnosis of Systemic sclerosis and Anti-centromere pattern in ANA.

The given clinical scenario featuring the recurrent bluing of hands(Raynaud phenomenon), difficulty swallowing (GERD), and tightening of the skin in the hands is suggestive of CREST syndrome/ limited cutaneous systemic sclerosis, and anti-centromere antibodies are most specific for this disease.

CREST syndrome consists of the following:

• Calcinosis cutis

• Raynaud's phenomenon

• Esophageal dysmotility

• Sclerodactyly

• Telangiectasia

Systemic sclerosis (SSc) is a chronic autoimmune disease. Progressive fibrosis due to the deposition of extracellular matrix components in different tissues and organs is the prominent feature of SSc.

The characteristic features of SSc are vascular damage, inflammation, and the presence of specific autoantibodies.

SSc is classified into two based on the extent of skin involvement.

• Limited cutaneous systemic sclerosis / CREST syndrome

• Diffuse cutaneous systemic sclerosis

In limited cutaneous systemic sclerosis (lcSSc), fibrosis in the skin of acral parts of the body, like the face and limbs (distal to elbow and knee), takes place. Skin fibrosis in limited type has a slow progression, but microcirculation disturbances like Raynaud's phenomenon start long before the skin symptoms and are associated with the involvement of the lungs and esophagus.

Since it has late and slow organ involvement, it is associated with a good prognosis with a 10-year survival of over 90%.

Antinuclear antibodies (ANA) are present in 90 - 95% of cases with SSc.

Autoantibodies specific to limited cutaneous systemic sclerosis include:

• Anti-centromere

• Anti-Th/To

• Anti-U1-RNP

• Anti-hUBF

Other options:

Option A: A speckled/granular pattern of staining is seen in systemic sclerosis due to antibodies against Scl-70, SLE, and Sjogren's syndrome. The image given below shows the speckled pattern of immunofluorescence.



Options C and D: Systemic lupus erythematosus (SLE) is an autoimmune disease that most commonly presents with arthralgia with early morning stiffness. Other important features of SLE include malar rash, discoid rash, photosensitivity, oral ulcers, pleuritis, pericarditis, and lupus nephritis.ANA, anti-dsDNA antibodies, and anti-smith antibodies are positive in SLE. Homogenous or diffuse staining and peripheral staining patterns are most commonly seen in SLE.

Solution to Question 35:

The given clinical presentations of numbness in the lower limbs, positive Romberg test, broad-based gait, and loss of proprioception are suggestive of dorsal column involvement of the spinal cord due to underlying vitamin B12 deficiency.

Vitamin B12 deficiency causes subacute combined degeneration of the spinal cord (SACD). It affects the following structures:

• Dorsal column - sensory loss, paresthesias, incontinence, sensory ataxia (positive Romberg sign with eyes closed)

• Lateral column - spasticity and paraplegias in later stages

The history of low BP, hypothyroidism, and recent gastrectomy further supports the diagnosis. Gastrectomy can lead to vitamin B12 malabsorption because the intrinsic factor(required for vitamin B12 absorption) is reduced after stomach removal leading to vitamin B12 deficiency. Hypothyroidism can sometimes coexist with B12 deficiency.

Folate deficiency (Option B) is typically not associated with neurological manifestations. In patients with a combined deficiency of B12 and folate, B12 deficiency must always be corrected first. This is because the administration of folate supplements may improve hematologic

abnormalities but worsen neurological manifestations.

Treatment of cobalamin deficiency is as follows:

• Replenishment of body stores: Six 1000-µg IM injections of hydroxocobalamin given at 3- to 7-day intervals.

• Maintenance therapy: 1000 µg hydroxocobalamin IM once every 3 months

Treatment of folate deficiency is as follows:

• 5–15 mg oral folic acid daily for a minimum of 4 months.

Other options

Option C: Thiamine deficiency causes Wernicke's syndrome with ophthalmoplegia, ataxia, and nystagmus. As this is due to cerebellar damage, Romberg's sign would be positive even with eyes open.

Option D: Transferrin deficiency, which affects iron transport, is rare and would not cause any neurological symptoms. Iron deficiency anaemia, which can cause fatigue and weakness, is more common, but it does not lead to proprioception loss or ataxia.

Solution to Question 36:

Tofacitinib is the JAK inhibitor that is preferred for moderate to severe ulcerative colitis.

Tofacitinib is a novel, oral JAK 1/3 - Inhibitor that will inhibit the JAK-STAT pathway responsible for the production of cytokines and chemokines. It has been approved for the treatment of ulcerative colitis, psoriatic arthritis and rheumatoid arthritis.

Other options:

Option A: Vedolizumab is a humanised anti-alpha-4-beta-7 integrin monoclonal antibody that prevents lymphocyte recruitment to the intestinal mucosa. It is approved for use in moderate-severe Crohn's disease and ulcerative colitis.

Option C: Ozanimod was recently FDA-approved for treating moderate to severe ulcerative colitis. Ozanimod is a potent sphingosine-1-phosphate (S1P1) receptor modulator. By preventing the disease-exacerbating lymphocytes from reaching the gut, ozanimod provides immunomodulatory effects and moderates disease processes.

Option D: Leflunomide is a pyrimidine synthesis inhibitor indicated for the management of rheumatoid arthritis in adults.

Solution to Question 37:

Plasmapheresis is preferred in refractory cases of GBS.

Guillain-Barré Syndrome (GBS) is an acute immune-mediated polyneuropathy characterized by weakness, areflexia, and sensory symptoms. It often follows an infection, vaccination, or other immune trigger. The autoimmune mechanism attacks the peripheral nervous system, leading to demyelination or axonal damage.

Mandatory diagnostic criteria for GBS:

• Progressive, symmetric muscle weakness of more than one limb: Weakness evolves over a period of hours to weeks. Typically starts in the legs and ascends (ascending paralysis).

• Areflexia or hyporeflexia: Deep tendon reflexes are reduced or absent in affected limbs. Reflex loss is a hallmark clinical feature.

Symptoms:

• Progressive weakness, starting in the legs and moving upwards (ascending paralysis).

• Areflexia (loss of reflexes) or hyporeflexia in the affected limbs. (Option A)

• Sensory symptoms like tingling or numbness may occur but are less prominent.

• Autonomic dysfunction, such as fluctuations in heart rate or blood pressure, in severe cases.

CSF Findings:

• Albuminocytological dissociation: High protein levels in cerebrospinal fluid (CSF) with normal or low white blood cell counts. (Option D)

Treatment:

• Intravenous Immunoglobulin (IVIG): First-line therapy. (Option C)

• Plasmapheresis (plasma exchange): in severe or refractory cases. (Option B)

• Supportive care for respiratory or autonomic complications.

Solution to Question 38:

As per KDIGO, 30-44 ml/min falls under G3b.

Kidney Disease Improving Global Outcomes (KDIGO) stages of chronic kidney disease (CKD):

Stage	GFR
Stage 1	> 90 ml/min/1.73 m^2
Stage 2	60 to 89 ml/min/1.73 m^2
Stage 3a	45 to 59 ml/min/1.73 m^2
Stage 3b	30 to 44 ml/min/1.73 m^2
Stage 4	15 to 29 ml/min/1.73 m^2
Stage 5 (end-stage renal disease)	< 15 ml/min/1.73 m^2



Solution to Question 39:

The clinical presentation and the chest X-ray findings of patchy infiltrates suggest pneumonia. However, patchy infiltrates are not commonly seen in pyogenic pneumonia. Patchy infiltrates are more characteristic of atypical pneumonia, such as Mycoplasma or viral infections.

Pyogenic pneumonia (community-acquired pneumonia) often shows lobar consolidation rather than patchy infiltrates.

Other options:

Option B: Mycoplasma pneumonia classically presents with interstitial infiltrates.

Option C: Hilar lymphadenopathy is indeed a feature of malignancy but it is unrelated to the acute presentation in this case.

Option D: Aspirational pneumonia is more common in the upper segment of the lower lobe and posterior segment of the upper lobe. Aspiration pneumonia typically affects these dependent lung segments.

Solution to Question 40:

The given clinical presentations such as uveitis, enthesitis, and sacroiliitis are suggestive of ankylosing spondylitis. HLAB27 is most likely associated with this condition.

The presence of HLA-B27 is strongly associated with ankylosing spondylitis (AS) and other spondyloarthropathies, such as reactive arthritis, psoriatic arthritis, and enteropathic arthritis, which often present with uveitis, enthesitis, and sacroiliitis. HLA-B27 is a genetic marker that increases the risk for these conditions, especially in the context of inflammatory back pain and other characteristic findings.

Other options

Option B: HLA-B57 is associated with HIV and some autoimmune conditions, but it is not a characteristic marker for spondyloarthropathies.

Option C: HLA-B51 is associated with Behçet's disease, which can present with uveitis but does not typically involve sacroiliitis or enthesitis in the same pattern as spondyloarthropathies.

Option D: HLA-B5 is associated with certain forms of autoimmune disease but is not typically associated with the spondyloarthropathies that present with sacroiliitis, enthesitis, and uveitis.

Solution to Question 41:

Brown-coloured urine is seen in glomerular hematuria, dysuria is absent and rare edema is present.

Glomerular hematuria is the presence of red blood cells (RBCs) in the urine that originate from the glomeruli. It can be identified by :

• Dysmorphic RBCs: (>40%)

RBCs that are misshapen or fragmented, often due to mechanical damage while passing through the damaged glomerular filtration barrier. These deformed RBCs are a hallmark of glomerular hematuria.

• Acanthocytes: (>5%)

These are abnormally shaped RBCs with a spiky or irregular surface, which suggest that the RBCs have been damaged as they pass through the glomerulus.

• RBC Casts: (>1)

The presence of RBC casts in the urine indicates that the blood is originating from the glomeruli, as casts are formed in the renal tubules.

These findings are indicative of glomerular injury, distinguishing glomerular hematuria from non-glomerular hematuria, where RBCs are typically normal and do not exhibit these characteristics.

Hemoglobin from red blood cells that have passed through the damaged glomerular basement membrane gets oxidized to methemoglobin, giving urine a dark or "cola-brown" appearance. This is a hallmark feature of glomerular hematuria. (A)

Dysuria (painful urination) is typically not a feature of glomerular hematuria, as this symptom is more common in lower urinary tract infections or bladder pathologies. (B)

Crystals in urine suggest kidney stones or metabolic disorders, not glomerular hematuria. (C)

Edema is a feature in nephritic syndrome or glomerulonephritis, which are associated with glomerular hematuria, due to sodium retention or hypoalbuminemia. Therefore, edema can be present in conditions causing glomerular hematuria. (D)

Solution to Question 42:

Ulcer over the dorsum of the penis is not seen in Behcet’s disease.

Behçet's disease is a systemic vasculitis that can cause recurrent oral ulcers, genital ulcers, skin lesions, and uveitis, among other manifestations. Genital ulcers in Behcet's disease typically occur on the scrotum in men and on the vulva in women, but not specifically on the dorsum of the penis.

Other options:

Option B: It is similar to Crohn's disease compared to ulcerative colitis. Behcet's disease shares some features with Crohn's disease, such as gastrointestinal involvement and the potential for both having systemic vasculitis. Ulcerative colitis, on the other hand, typically affects only the colon and does not present with the systemic symptoms seen in Behcet's disease.

Option C: Bilateral panuveitis is a common and serious manifestation of Behcet's disease and can lead to vision loss if untreated.

Option D: Pathergy test positive is a criterion of ISG diagnostic criteria for Behcet's. The pathergy test, which involves pricking the skin and observing for a hyper-inflammatory response (a red papule or pustule), is indeed one of the diagnostic criteria for Behcet's disease, though it is not always present in all patients.

Solution to Question 43:

The ECG provided shows epsilon waves and T wave inversions, which is a hallmark finding suggestive of Arrhythmogenic Right Ventricular Dysplasia (ARVD).

The clinical presentation of young females with palpitations and syncope, especially post-exercise, are common in ARVD due to ventricular arrhythmias.

The changes marked in ECG show a notched deflection at the end of the QRS complex due to delayed ventricular activation known as an epsilon wave. It is commonly seen in patients with ARVD.

ARVD is autosomal dominant cardiomyopathy characterized by fibrofatty replacement in the myocardium of the ventricles. It most commonly affects the right ventricle. It is due to mutations in the gene encoding desmosome-associated protein plakophilin 2. This leads to defective cell-adhesion proteins in the desmosomes that link adjacent cardiac myocytes.

The patients usually present with rhythm disturbances, right-sided heart failure, and sudden cardiac death. ARVD is associated with Naxos syndrome. This syndrome is characterised by arrhythmogenic cardiomyopathy and hyperkeratosis of the palmar and the plantar surfaces.

Morphology shows severe thinning of the right ventricular wall due to loss of myocytes, accompanied by extensive fatty infiltration and fibrosis. In ARVD, there is T inversion and delayed ventricular activation manifesting as epsilon waves. This reflects the involvement of the right ventricular myocardium.



Other options

Option A: Uhl's disease is rare and typically associated with the absence of RV muscle. It does not have specific ECG patterns like those in ARVD.

Option C: Hypertrophic Cardiomyopathy (HCM) usually shows LV hypertrophy with left-axis deviation and repolarization abnormalities, particularly in lateral leads.

Option D: Brugada syndrome is characterized by coved-type ST elevation in leads V1-V3, not T wave inversions as seen here.

Solution to Question 44:

The preferred first investigation for a patient diagnosed with pheochromocytoma would be 24-hour urine metanephrines.

This test (24-hour urine metanephrines) is highly sensitive and specific for diagnosing pheochromocytoma. Elevated levels of metanephrines (products of catecholamine metabolism) in the urine indicate the condition. After biochemical confirmation, other imaging studies, like MRI or MIBG scans, are typically used to localise the tumour.

Other options

Option A: USG-abdomen can help detect abdominal masses, but it is not sensitive enough for pheochromocytoma, as it is not accurate in detecting it.

Option B: MIBG scan is specific for detecting pheochromocytoma, but it is not typically used as a first-line investigation. It is used only after biochemical confirmation.

Option C: MRI-abdomen can be useful for tumour localisation, but biochemical testing (like 24-hour urine metanephrines) is preferred first for diagnosis.

Medicine NEET 2024

Question 1:

An elderly patient who is a known case of scleroderma presented with shortness of breath. On examination, her right ventricular pressure was 70-80 mmHg and later on catheterization, pulmonary capillary wedge pressure was found to be 35 mmHg. Which of the following is true regarding her condition?

a) Calcium channel blockers given for all the patients

b) Lung transplant is the best treatment, if identified early

c) Endothelin receptor antagonists improves symptoms

d) Lifestyle modification is the only possible approach

Question 2:

A 26-year-old patient with bronchial asthma who has two episodes in the daytime and one episode at night time per week is prescribed albuterol to use as and when required. MDI FEV1 improved from 76% to 83% after taking bronchodilator. What is the further treatment plan?

a) Add oral prednisolone 10 mg daily

b) Continue albuterol whenever required

c) Add fluticasone propionate 40 mg BD

d) Change to salbutamol MDI twice daily

Question 3:

A 30-year-old patient came with chest pain shortness of breath, and pedal edema. JVP showed rapid y descent. On auscultation, he had high-pitched diastolic sound. What is the diagnosis?

a) Chronic constrictive pericarditis

b) Cardiac tamponade

c) RHD with Mitral stenosis

d) Bicuspid aortic valve with Aortic regurgitation

Question 4:

A 30-year-old female patient comes to the outpatient department with complaints of neck swelling. She gives a history of weight loss and palpitations. On examination, she has exophthalmos and a diffusely enlarged thyroid gland. Which of the following is not true?

a) Severe disease can cause vision loss

b) NO SPECS criteria is used for this condition

c) 10% of isolated cases is present without hyperthyroidism

d) Condition will improve simultaneously with improvement in thyrotoxic state

Question 5:

A 56-year-old presented with fever chills & cough with sputum. Blood pressure was measured to be 150/88mmhg, pulse rate - 110bpm and respiratory rate - 18, spo2 - 88% at room air. An X-ray was done and shown below. What is the probable diagnosis?



a) Tubercular pleural effusion

b) Community acquired pneumonia

c) Coal worker Pneumoconiosis

d) Silicosis

Question 6:

A 45-year-old man presented with fatigue and hemoptysis. His current hemoglobin is 8g/dL (previous hemoglobin one week ago was 12g/dL). His bronchoalveolar lavage is reddish and his serum creatinine was 1.7mg/dL. He is diagnosed with ANCA-positive glomerulonephritis. Biopsy of the kidney revealed pauci-immune deposits and the absence of granulomas. Which of the following conditions is he likely suffering from?

a) Churg-Strauss syndrome

b) Microscopic polyangiitis

c) SLE

d) Henoch–Schonlein purpura

Question 7:

A patient with an acute exacerbation of COPD required mechanical ventilation and experienced a consistent drop in oxygen saturation. Auscultation revealed vesicular breath sounds on the right side. Based on the chest radiography findings, what would you expect to hear on auscultation of the left side?

a) Absent breath sounds

b) Bronchial breath sounds

c) Increased vesicular breath sounds

d) Increased vocal fremitus

Question 8:

A patient with salicylic acid poisoning has the following arterial blood gas analysis report: pH = 7.3, pCO2 =16mmHg, HCO3-=6.3mmol/L, Na+- 135, K+- 4.5 Cl- 113.7, lactate was normal. Which of the following options correctly describes the condition?

a) Metabolic acidosis with partial compensated respiratory alkolosis with high anion gap

b) Respiratory acidosis with compensatory metabolic alkalosis

c) Metabolic acidosis with uncompensated respiratory alkalosis

d) Respiratory acidosis with metabolic alkalosis

Question 9:

Match the following ECG waves with condition: atrial flutter, atrial fibrillation, SVT, ventricular tachycardia



a) 1-■, 2-■, 3-■, 4-■

b) 1-■, 2-■, 3-■, 4-■

c) 1-■, 2-■, 3-■, 4-■

d) 1-■, 2-■, 3-■, 4-■

Question 10:

A 34-year-old patient presents with lethargy and tiredness. On examination, he has perioral hyperpigmentation. His blood pressure is 90/60 mmHg and his heart rate is 110 beats/minute. Laboratory investigations revealed Na+ of 125mEq/L and K+ of 5.5mEq/L. What is the most appropriate treatment given to this patient?

a) ACTH

b) Hydrocortisone + Fludrocortisone

c) Dexamethasone

d) Saline infusion

Question 11:

A young woman comes with weight loss, tachycardia, heat intolerance, and tremors. She has exophthalmus on examination. What is the diagnosis?

a) Addison's disease

b) Parkinson's disease

c) Graves disease

d) Alzheimer's disease

Question 12:

For MDR-RR TB, what is the indication of short course bedaquiline?

a) Resistance to rifampicin and sensitive to fluroquinolones

b) Mutation only to INH A and kat G gene

c) Rifampicin sensitive

d) Extrapulmonary TB like tubercular meningitis

Question 13:

In a patient with type 1 diabetes with 3rd stage beta cell damage, which of the following situations is associated with the same?

a) Autoimmune positive with dysglycemia

b) Autoimmune positive with normoglycemia

c) Autoimmune negative with normoglycemia

d) Autoimmune negative with dysglycemia

Question 14:

An alcoholic patient presented to the emergency in an unconscious state. Relatives stated that he had a similar episode of binge drinking one week back. On examination, his glucose was 49mg/dl. What is the appropriate management?

a) IV thiamine followed by dextrose infusion

b) Injection vitamin K followed by dextrose infusion

c) IV 25% dextrose

d) NS infusion

Question 15:

A patient came to the emergency department with dyspnea and palpitations. He is a known case of chronic kidney disease. ECG of the patient is given below. What is the diagnosis?



a) Hyperkalemia

b) Hypokalemia

c) Hypocalcemia

d) Hypercalcemia

Question 16:

A 50-year-old female presented with dyspnea and angina episodes. On workup, she is found to have aortic stenosis. Which of the following findings favor the diagnosis of aortic stenosis over aortic regurgitation?

a) Increase in preload causes more reduction in myocardial oxygen demand than increase in afterload

b) Workload has nothing to do with myocardial O2 consumption

c) Aortic stenosis causes reduced pressure than aortic valve

d) Increase in myocardial O2 consumption is seen with increased pressure work than volume

Question 17:

A patient presented with fatigue and tingling sensation in the arms, weight loss and h/o polyuria. The following findings were detected on bone marrow biopsy. What is probable cause ?



a) Subacute combined degeneration due to cobalamin deficiency

b) Multiple myeloma

c) Subacute degeneration due to diabetes

d) Megaloblastic anemia

Question 18:

A 60-year-old patient presents with dyspnea and chest pain. The pleural fluid analysis shows pleural protein/ serum protein > 0.5, and pleural LDH/ serum LDH > 0.4. What is the diagnosis?

a) Congestive heart failure

b) Chronic kidney disease

c) Liver cirrhosis

d) Tuberculosis

Question 19:

An adolescent presented with fever, purpura on the thigh, swelling of the right knee and left ankle, and abnormal and diastolic murmur in the cardiac apical area. He was seronegative for rheumatoid factor. What is the probable diagnosis?

a) Acute rheumatic fever

b) Juvenile idiopathic arthritis

c) Rheumatoid arthritis

d) Seronegative arthropathy

Question 20:

A patient came with complaints of easy fatiguability, anemia, hyperpigmentation of knuckles, oral mucosa, palm and hands, and tachycardia. The blood pressure of the patient is 80/60 mm Hg and Na+ 130 meq/L. What is the most probable diagnosis?

a) Conn's syndrome

b) Primary adrenal insufficiency

c) Vitamin B12 deficiency

d) Cushing syndrome

Question 21:

A female presents with choreo-athetoid movements, dysarthria, and dystonia with the given finding. Which of the following is the best investigation to confirm the same?

a) Serum ceruloplasmin levels

b) Hepatic copper content

c) 24-hour urinary excretion of copper

d) Blood copper levels

Question 22:

A 27-year-old footballer collapsed on the field and could not be revived. An autopsy was conducted and on cardiac biopsy, the left ventricle was hypertrophied. Identify the possible cause for the same.



a) Ischemic cardiomyopathy

b) Arrhythmogenic right ventricular dysplasia

c) Hypertrophic cardiomyopathy

d) Takotsubo cardiomyopathy

Question 23:

A patient suffered from RTA. After a few days, he had exaggerated right-sided extensor plantar reflex and knee jerk. What other finding would be seen?

a) Ankle clonus in right foot

b) Proprioception lost in left leg

c) Pain and temperature was lost in right leg

d) Fasciculations in right foot

Question 24:

A patient after RTA underwent ileal resection. He presents with fatigue, which of the following will not be seen?

a) Cyto-nuclear dys-synchrony

b) Microcytic hypochromic anemia

c) Romberg's sign positive

d) Subacute combined degeneration

Question 25:

A patient came with a history of dyspnea. His CT scan is shown below. What is the likely diagnosis?



a) Mediastinal mass invading vascular structures

b) Thrombosed arch of aorta

c) Mediastinal mass with fat and calcification

d) This is characteristic of mediastinal Hydatid cyst

Question 26:

Which of the following is most likely based on the lab values given?

a) Type 1 renal tubular acidosis

b) Type 2 renal tubular acidosis

c) Type 3 renal tubular acidosis

d) High Anion Gap Metabolic Acidosis

Question 27:

A 36-year-old girl presented with tremors and weight loss. Her ophthalmic examination shows the following findings. Identify the true statement regarding the following condition.



a) No association with HLA

b) Medial rectus is the first muscle to be involved.

c) The muscles and connective tissue around the eye shares similar antigen with diseased organ

d) More common in males

Question 28:

A patient presented to the casualty with chest pain and palpitations. His blood pressure was 130/90mmHg and pulse rate- 101bpm. Echocardiography showed a collapse of the right atrium during diastole. What is the most appropriate management for this patient?

a) NSAIDs

b) Pericardiocentesis

c) Chest tube insertion or thoracotomy

d) Steroids

Question 29:

A patient gives a history of epigastric pain radiating to the back for five days. Lab investigations show that amylase and lipase are elevated five times the normal values. Two litres of saline was administered in 24 hours to the patient. He also gave a history of consumption of alcohol. Given below is the chest X-ray of the patient. What is the most probable diagnosis?

a) Aspiration pneumonia

b) Non-cardiogenic pulmonary edema

c) Miliary tuberculosis

d) Bilateral pulmonary embolism

Question 30:

What is the most likely cause of death in a patient suffering from Idiopathic pulmonary fibrosis?

a) Pulmonary hypertension

b) Pulmonary embolism

c) Respiratory failure

d) Lung carcinoma

Question 31:

A 35-year-old female presented with complaints of fever for 5 days and there were no signs of meningitis. MRI with diffusion-weighted and contrast are shown. What is the most likely diagnosis?



a) Tuberculoma

b) Toxoplasmosis

c) Glioma

d) Brain abscess

Question 32:

A patient presented to the OPD with a temperature of 102°F, and a cough with foul-smelling sputum. On examination, clubbing is present. His chest X-ray is given below. What is the probable diagnosis?

a) Pneumothorax

b) Pneumatocele

c) Empyema

d) Lung abscess

Question 33:

A young athlete presented with chest pain and dyspnea. His brother passed away due to sudden cardiac death. He is diagnosed with HCM. A harsh systolic murmur is heard in the left sternal border. Which of the following will increase the murmur?

a) Squatting

b) Leaning forward with head down/up

c) Hand grip

d) Valsalva

Question 34:

A patient presented with right upper and lower limb weakness with left sided facial nerve palsy and left sided horizontal gaze palsy. What is the probable diagnosis?

a) Millard- Gubler syndrome

b) Benedikt's syndrome

c) Foville syndrome

d) Locked in syndrome

Question 35:

A male patient presents with headaches and visual disturbance. The patient had frontal bossing, thick palms, and soles, and gaping of central incisors. He is also hypertensive and diabetic. Which of the following statements is true about screening investigations done on this patient?

a) GH levels <0.4 after 1 hour of 75 gm glucose tolerance test.

b) Single early morning fasting level of GH is used to establish the diagnosis.

c) Age specific IGF-1 levels are useful for screening.

d) GH levels are used to assess the severity of the condition.

Question 36:

A person goes through routine tests while applying for a job. His ECG showed a PR interval of 0.21 sec. What is the diagnosis?

a) 1st degree block

b) Normal ECG

c) 2nd degree block

d) Complete heart block

Question 37:

A woman presented with an irregularly irregular rhythm with a heart rate of 150 bpm. On auscultation P2 was loud and no murmur was heard. What would be the JVP finding?

a) Rapid x descent

b) Absent a wave

c) Cannon a wave

d) Rapid y descent

Question 38:

A young man had a history of jaundice and raised HBsAg levels a year back. On follow-up now, he has normal levels of liver enzymes. His current profile is shown in the image. What is the diagnosis for his current condition?



a) Chronic HepB with HBe Ag negative

b) Resolved infection

c) Chronic HepB with HBe Ag positive

d) Acute HepB

Question 39:

A 40-year-old female patient with a known case of myasthenia gravis on mycophenolate mofetil and pyridostigmine now presents with breathing difficulties. Her calcium levels were 13.5g/dl. What is the most likely cause for the increase in calcium?

a) Small cell carcinoma of lung

b) Drug induced hypercalcemia

c) Parathyroid adenoma

d) Adenocarcinoma lung

Question 40:

A 35-year-old female with a history of hypertension presents with palpitations, dyspnea, bilateral basal crackles and irregularly irregular pulse. ECG is given below. Which among the following management is least useful?



a) Cardioversion in a haemodynamically unstable patient

b) Direct oral anticoagulants

c) Beta blockers is to be given for rate control if patient is intolerant to Calcium channel blockers

d) IV digoxin for ventricular rate control

Question 41:

A 25-year-old male presents with hyperthermia, autonomic instability, confusion, hyperreflexia, muscle rigidity, and dilated pupils. The patient takes sertraline and was recently prescribed amitryptiline and sumatriptan for depression and migraines, respectively. Which medication is most likely to be effective in treating this patient's current condition?

a) Cyproheptadine

b) Propranolol

c) Diphenhydramine

d) Nitroprusside

Answer Key

Question No.	Correct Option
1	c
2	c
3	a
4	d
5	b
6	b
7	a
8	a
9	a
10	b
11	c
12	a
13	a
14	a
15	a
16	d
17	b
18	d

19	a
20	b
21	b
22	c
23	a
24	b
25	c
26	a
27	c
28	b
29	b
30	c
31	d
32	d
33	d
34	c
35	c
36	a
37	b
38	c
39	c
40	c
41	a

Detailed Explanations

Solution to Question 1:

In this elderly patient with scleroderma presenting with elevated right ventricular pressure and pulmonary capillary wedge pressure, is suggestive of scleroderma/systemic sclerosis(SSc)-associated pulmonary hypertension, and endothelin receptor antagonists improve the symptoms.

Right heart catheterization, the gold standard for diagnosing pulmonary hypertension, confirms the condition when the mean pulmonary arterial pressure (mPAP) exceeds 20 mmHg. In this elderly patient with scleroderma, elevated right ventricular pressure (70-80 mmHg) and pulmonary capillary wedge pressure (35 mmHg) indicate severe pulmonary hypertension.

This condition involves intimal thickening and narrowing of pulmonary arteries, leading to increased blood flow resistance and symptoms such as chest pain, dyspnea, and hemoptysis. As

the disease progresses, it causes right ventricular dysfunction, presenting as syncope, abdominal distension, ascites, and peripheral edema. The disease leads to vascular changes like intimal proliferation, fibrosis, and thrombosis, which increase right ventricular afterload and total pulmonary vascular resistance, ultimately resulting in decreased cardiac output, irreversible damage, and potential right heart failure and death.

An echocardiogram with a bubble study is an important screening test that will reveal a dilated and hypertrophied right ventricle. On chest x-ray, 'vascular pruning' may be seen as well as enlarged pulmonary arteries.

Management for pulmonary hypertension:



Note: Early identification of PAH(option B) allows for effective management with available medical therapies, making lung transplant a later option rather than the immediate treatment of choice.

Vasoreactivity testing involves the administration of inhaled nitric oxide, IV epoprostenol, IV adenosine, or inhaled iloprost. The test is considered positive if the mean pulmonary arterial pressure (mPAH) decreases by ≥ 10 mmHg leading to mPAH ≤ 40 mmHg, without a decrease in cardiac output (CO).

Solution to Question 2:

In the given clinical scenario of a patient with mild persistent asthma taking albuterol and FEV1 improving from 76% to 83% after taking bronchodilator, the patient should be prescribed fluticasone propionate 40mg BD.

Treatment guidelines for different types of asthma are given below:

LABA - Long-acting Beta 2 Agonists, ICS - Inhaled corticosteroids

Add-on therapy includes long-acting anticholinergics, leukotriene antagonists, mast cell stabilizers, methylxanthines, and antibodies.

Drugs used in the treatment of bronchial asthma

Bronchodilators

• Beta-2 sympathomimetics: Salmeterol, Formoterol

• Methylxanthines: Theophylline, Aminophylline

• Anticholinergics: Ipratropium bromide, Tiotropium bromide

Anti-inflammatory

• Corticosteroids (Inhalational): Beclomethasone, Budesonide, Fluticasone

• Corticosteroids (Systemic): Prednisolone, Hydrocortisone

• Mast cell stabilizers: Sodium cromoglycate, Ketotifen

• Leukotriene antagonists: Montelukast, Zafirlukast

• anti-IgE - Omalizumab

• anti-IL5 - Mepolizumab, Reslizumab

• anti-IL4 receptor - Dupilumab

Important Update - According to GINA 2019, Short-acting beta 2 agonists (SABA) have no role in treating acute exacerbations. Even though SABA provides short-term relief of symptoms, there is strong evidence that SABA only treatment increases the risk of exacerbations.

GINA now recommends that symptom-driven (in mild asthma), a daily dose of ICS (inhaled corticosteroids) containing treatment to reduce the risk of serious exacerbations. This is said to be the most important change in asthma management in 30 years

Step 1 (intermittent)	As needed low dose ICS + Fo rmoterol
Step 2 (mild persistent)	Daily low-dose ICS
Step 3 (moderate persistent)	Daily low dose ICS + LABA
Step 4 (severe persistent)	Medium dose ICS + LABA +/- add-on therapy
Step 5	High dose ICS + LABA + oral steroids +/- add-on therapy

Solution to Question 3:

The combination of the given clinical history and JVP showing rapid y descent is highly suggestive of chronic constrictive pericarditis.

Constrictive pericarditis is a chronic type of pericarditis that results when there is healing of acute fibrinous pericarditis or absorption of chronic pericardial effusion. The most common cause in developing countries is tuberculosis, following viral pericarditis, cardiac surgery, mediastinal irradiation, or as a manifestation of rheumatoid arthritis and SLE. The main abnormality that occurs in constrictive pericarditis is the inability of the ventricles to fill because of the rigid and

thickened pericardium. The restriction is not uniform through the diastole as seen in tamponade. Ventricular filling is normal early in the diastole but stops suddenly when the elastic limit of the pericardium is reached.

Y-descent on the JVP curve is prominent in constrictive pericarditis because it represents the rapid early filling of the ventricles. A sudden stop in ventricular filling and subsequent rising pressure is seen as a square root sign in the JVP curve.

The patient presents with exertional dyspnea and orthopnea. There may be a paradoxical rise in venous pressure during inspiration (Kussmaul's sign) reflecting impaired right ventricular compliance. The ECG often shows low-voltage QRS complexes and diffuse flattening of T-waves. A chest radiograph may show pericardial calcification, especially in tuberculous pericarditis.



The waveforms seen in a normal JVP are:

• a wave corresponds to atrial contraction and occurs just after the P wave (on ECG) and just before the first heart sound.

• c wave represents the bulging of the tricuspid valve into the right atrium during ventricular systole.

• x descent corresponds to the atrial relaxation during atrial diastole.

• v wave represents atrial filling during ventricular systole and peaks at the second heart sound.

• y descent represents passive emptying of the right atrium into the right ventricle after tricuspid valve opening.

A normal tracing of the jugular venous pulse is given below:



Other options:

Option B: Cardiac tamponade: On JVP tracing, cardiac tamponade characteristically shows a prominent x-descent and absent y-descent.

The image of JVP in cardiac tamponade is given below:



Option C: Mid-diastolic murmur is seen in mitral stenosis (MS). It occurs most commonly in rheumatic fever. A loud S1 (first heart sound) with an opening snap following S2 (second heart sound) is heard. S2 is followed by a low-pitched, rumbling, mid-diastolic murmur, which is best heard at the apex with the patient in a lateral recumbent position.

Option D: Aortic regurgitation can occur due to bicuspid aortic valve, rheumatic fever, degeneration of the valve, etc. Auscultation will have a high-pitched, blowing, decrescendo diastolic murmur, heard best in the third intercostal space along the left sternal border. Mid systolic ejection murmur can also be heard.

Solution to Question 4:

The given clinical scenario of a female patient presenting with neck swelling weight loss, palpitations, exophthalmos, and diffusely enlarged thyroid gland is suggestive of Graves' disease with thyroid eye disease. Thyroid eye disease (TED) does not improve simultaneously with improvement in the thyrotoxic state.

Graves' disease causes hyperthyroidism through thyroid-stimulating immunoglobulin (TSI) synthesized by lymphocytes in the thyroid gland, bone marrow, and lymph nodes. TSI antibodies stimulate the thyroid gland, causing hyperthyroidism.

Symptoms include hyperactivity, heat intolerance, sweating, palpitations, weakness, weight loss, increased appetite, and diarrhoea. Signs include tachycardia, goitre, tremor, muscle weakness, and atrial fibrillation in the elderly.

Thyroid ophthalmopathy is seen in one-third of patients with Grave's disease. Thyroid dermopathy usually occurs in the presence of moderate or severe ophthalmopathy. It primarily manifests as noninflamed, indurated plaques with a deep pink or purple colour and an "orange skin" appearance, commonly found on the anterior and lateral aspects of the lower leg (pretibial myxedema).

Investigations done on a patient with Graves' disease include:

• Thyroid function tests- TSH levels are decreased with or without elevated T3 and T4 levels.

• Anti-thyroid antibodies—Elevated levels of TSH receptor antibodies are diagnostic of Graves' disease. Other antibodies, like anti-Tg and anti-TPO, are elevated in most patients.

• Ultrasound of the neck is performed to evaluate vascularity, echogenicity, nodules, and clinical landmarks for surgical planning.

• A radioactive iodine uptake (RAIU) scan (thyroid scan) should be performed and a diffuse uptake confirms the diagnosis of Graves' disease. It is also useful to differentiate TSI-negative Graves disease from toxic nodular disease based on diffuse or nodular uptake patterns.

• CT or MRI scans of the orbits are useful to evaluate Graves' ophthalmopathy.

Hyperthyroidism can be treated by antithyroid drugs, radioiodine treatment (I131), or thyroidectomy.

Other options:

Option A: Severe disease can cause vision loss mainly due to compression of the optic nerve at the orbit's apex, which can result in peripheral field defects, papilledema, and, in the worst case scenario, irreversible blindness.

Option B: NO SPECS scoring system as described below can be used to evaluate ophthalmopathy.

• No signs or symptoms

• Only signs (lid retraction, or lag)

• Soft tissue involvement (periorbital oedema)

• Proptosis

• Extraocular muscle involvement

• Corneal involvement

• Sight loss

Option C: 10% of isolated cases of thyroid eye disease are present without hyperthyroidism as they may also present as euthyroid or hypothyroid.

Solution to Question 5:

The clinical vignette describes a case of community-acquired pneumonia. The findings of fever, chills, cough with expectoration, and chest X-ray findings of lobar involvement support the diagnosis.

Clinical features include acute onset fever, cough (with or without sputum production), and shortness of breath. In some cases, pleuritic chest pain, tachycardia, tachypnea, hypoxemia, or increased work of breathing may be present on physical examination.

Diagnosis can be made using a chest X-ray. The presence of an infiltrate on a plain chest radiograph is considered the gold standard for diagnosing pneumonia. Radiographic findings consistent with the diagnosis of CAP include lobar consolidations, interstitial infiltrates, and/or cavitations. CT chest is used for better resolution or if chest X-ray was non-diagnostic. Blood and sputum cultures are obtained to guide the antibiotic regimen.

Other options:

Option A: Pleural effusion may present with dyspnea, pleuritic pain, dry cough, and associated symptoms like fever, weight loss, oedema, and ascites depending on the aetiology. On examination, there may be tachypnea, dullness on percussion and decreased/absent breath sounds at the base(s).



Option C: Coal worker's pneumoconiosis (CWP) is an occupational disease due to chronic exposure to coal dust. It is seen in people working in mines. It presents small opacities in the

upper lobe of the lungs. Complicated CWP has nodules ≥1 cm in diameter generally confined to the upper half of the lungs.

Option D: Silicosis is an occupational disease and is caused by exposure to silica. It is seen in people who work in mining, stone cutting, sandblasting, and quarrying. It presents as small rounded opacities in the upper lobe. There may be calcification of hilar lymph nodes which leads to the appearance of eggshell calcification. HRCT will show a characteristic crazy paving pattern. The image is given below



Solution to Question 6:

The given clinical scenario is suggestive of microscopic polyangiitis. The absence of granulomatous inflammation differentiates microscopic polyangiitis from other forms of ANCA-associated vasculitis.

Anti-neutrophil cytoplasmic antibodies (ANCA)-associated vasculitis is categorized into 3 distinct forms:

• Granulomatosis with polyangiitis (GPA; formerly Wegener granulomatosis)

• Microscopic polyangiitis (MPA)

• Eosinophilic granulomatosis with polyangiitis, formerly called Churg–Strauss syndrome (CSS)

Necrotizing vasculitis is the cardinal histologic feature in both GPA and MPA. Kidney biopsies typically demonstrate crescentic glomerulonephritis with little or no immune complex deposition. hence the name “pauci-immune vasculitis.”

Granulomatous inflammation is typically absent in MPA but is common in both GPA and CSS.

ANCA-associated vasculitis (AAV) comprises a group of multi-systemic diseases affecting small-to-medium-sized vessels in organs like kidneys and lungs. It is characterized by the presence of ANCA with specificity for either proteinase-3 (PR3) or myeloperoxidase (MPO).

Other options:

Option A: Churg-Strauss syndrome is typically characterized by a necrotizing granulomatous small vessel vasculitis with a history of bronchial asthma and peripheral eosinophilia.

Option C: SLE is an autoimmune disease involving multiple organs, with a vast array of autoantibodies, particularly antinuclear antibodies (ANA). The lesion occurs mainly by deposition of immune complexes and binding of antibodies to various cells and tissues

Option D: Henoch–Schonlein purpura is the most common vasculitis in childhood. It is characterized by leukocytoclastic vasculitis and IgA deposition in the skin, kidney, joints, and GIT. The clinical manifestations include palpable purpura, joint involvement in the form of arthritis or arthralgias, diarrhoea, vomiting, renal involvement in the form of microscopic hematuria, proteinuria, etc.

Solution to Question 7:

Absent breath sounds on the left side are expected in this case of left pneumothorax.

A pneumothorax occurs when air enters the pleural space, causing lung collapse and preventing normal lung expansion, which leads to absent or diminished breath sounds over the affected area. In patients with COPD, particularly during an acute exacerbation, the already weakened and hyperinflated lung tissue can rupture, leading to a pneumothorax. This is a serious complication, especially when mechanical ventilation is required, as the pressure from the ventilator can exacerbate the pneumothorax.

Other options:

Option B: Bronchial breath sounds: Typically heard in conditions like pneumonia, where there is lung consolidation, not in a pneumothorax.

Option C: Increased vesicular breath sounds: These indicate hyperinflation or normal lung function, which is not possible in a collapsed lung.

Option D: Increased vocal fremitus: Fremitus is usually decreased or absent in pneumothorax due to the insulating effect of air in the pleural space.

Solution to Question 8:

The acid-base disorder here is a mixed disorder of metabolic acidosis with partially compensated respiratory alkalosis with high anion gap.

It should not be confused with metabolic acidosis with uncompensated respiratory alkalosis. This can be understood by the following calculation:

Step 1: The pH in this patient is 7.3 (pH <7.35) which indicates acidosis

Step 2: HCO_3^- is 6.3 mmol/L (Normal HCO_3^- - 22–30 mmol/L). In metabolic acidosis, HCO_3^- levels decrease.

Hence, this is a case of metabolic acidosis.

Step 3: Calculate the anion gap.

Anion gap = Na+-(Cl-+HCO3-)

= 135 – (113.7 +6.3)

= 15, which is a high anion gap (normal anion gap is between 6-12 mEq/L)

Step 4: As per Winter's formula, the expected respiratory compensation in metabolic acidosis can be calculated using,

$pCO_2 = (1.5 \times [HCO_3^-]) + 8 \pm 2$

In this case, Expected $pCO_2 = (1.5 \times [6.3]) + 8 \pm 2$

$= 17.5 \pm 2$

$= 15.45 - 19.45$ mmHg

The pCO_2 in this patient is 16 mmHg, which implies there is an overcompensation of pCO_2. This overcompensation can happen only if there is an additional component of respiratory alkalosis contributing to reduced pCO_2.

Hence, this patient has a mixed disorder - metabolic acidosis and respiratory alkalosis.

Solution to Question 9:

The matched option is option A, i.e., 1-■, 2-■, 3-■, 4-■.

Atrial flutter rhythm is regular. The rate is 250-300 bpm. Normal P waves are replaced by saw tooth-shaped waves known as flutter waves.



Atrial fibrillation rhythm is irregularly irregular. P waves are absent, these irregular waves are known as fibrillation waves, characteristic of atrial fibrillation.



Paroxysmal ventricular tachycardia (PSVT) ECG shows narrow QRS complex (<0.12 s) tachycardia with an absent P wave.



Ventricular tachycardia ECG changes include wide QRS complexes, absence of P waves, and ventriculoatrial dissociation. It can be broadly classified as sustained VT or non-sustained VT. Nonsustained VT has QRS complexes lasting <30 seconds while sustained VT lasts for >30 seconds. In monomorphic VT, all QRS complexes look the same. Polymorphic VT has QRS complexes varying in shape or direction in the same lead.



Solution to Question 10:

This clinical scenario is suggestive of primary adrenal insufficiency (Addisonian crisis) and IV hydrocortisone+fludrocortisone is the most appropriate treatment for this condition.

Adrenal insufficiency can be primary (due to adrenal pathology), secondary (due to pituitary pathology- lack of ACTH), or tertiary (due to hypothalamus pathology- lack of CRH). Autoimmune adrenalitis is the most common cause of primary adrenal insufficiency worldwide. Tuberculosis is the most common cause of primary adrenal insufficiency in India.

Acute adrenal insufficiency (adrenal or addisonian's crisis) is a medical emergency caused due to a precipitating illness unmasking insufficiency or trauma or severe sepsis causing adrenal hemorrhage (Waterhouse-Friderichsen syndrome) or surgical removal of the glands. It presents with non-specific symptoms such as shock with fever, nausea, vomiting, abdominal pain, hypocalcemia, and electrolyte imbalances. It may be often misdiagnosed as an acute abdominal emergency.

Acute adrenal insufficiency requires the following therapeutic strategies:

• Rehydration - saline infusion at an initial rate of 1 L/h with continuous cardiac monitoring.

• Glucocorticoid replacement - bolus injection of 100 mg hydrocortisone, followed by the maintenance dose of 100–200 mg over 24h.

• Mineralocorticoid replacement - initiated once the daily hydrocortisone dose has been reduced to <50 mg as higher doses of hydrocortisone have sufficient mineralocorticoid action.

Solution to Question 11:

Based on the clinical findings the most likely diagnosis is Graves' disease.

The hyperthyroidism of Graves' disease is caused by thyroid-stimulating immunoglobulins (TSI) that are synthesized in the thyroid gland as well as in bone marrow and lymph nodes. TSI cause stimulation of the thyroid gland causing features of hyperthyroidism in the presence of low TSH levels.

Clinical features of Graves' disease include hyperthyroid features like hyperactivity, irritability, dysphoria, heat intolerance and sweating palpitations, fatigue, and weakness, weight loss with increased appetite, diarrhea, polyuria, oligomenorrhea, loss of libido, tachycardia.

Graves ophthalmopathy is due to increased volume of the retroorbital connective tissues and extraocular muscles because of marked lymphocytic infiltration of the retro-orbital space, accumulation of hydrophilic glycosaminoglycans such as hyaluronic acid and chondroitin sulfate, and increased numbers of adipocytes (fatty infiltration). Corneal exposure, lid retraction, lid lag on downgaze, conjunctival injection, restriction of gaze, diplopia, and visual loss from optic nerve compression are cardinal symptoms.

Histopathology shows follicular hyperplasia with scalloped margins on biopsy.

Management involves propylthiouracil is the mainstay of treatment for Graves' disease. Radioactive iodine is also used as a treatment modality for hyperthyroidism in suitable patients but can aggravate Grave's ophthalmopathy.

Other options:

Option A: Patients with Addison's disease usually present with complaints of fatigue, myalgias, weight loss, and joint pain. Mineralocorticoid deficiency manifests as salt craving, dizziness due to hypotension, and abdominal pain. Excess ACTH stimulation results in hyperpigmentation of exposed areas of the skin such as palmar creases and dorsal foot. Hyponatremia is a characteristic feature and is often associated with hyperkalemia.

Option B: Parkinson's disease is a degenerative condition of the dopamine-secreting neurons of the substantia nigra. Patients present with muscular rigidity, involuntary resting tremors of the hands, akinesia - difficulty in initiating movement - and postural instability. On examination, cog-wheel rigidity and a shuffling gait are seen.

Option D: Alzheimer's disease is the most common cause of dementia in the elderly. It begins with memory impairment, which later progresses to language and visuospatial deficits. Later, the patient has difficulty in keeping track of finances, following instructions, driving, shopping, and housekeeping. In severe impairment, the patient easily gets lost and confused disturbances and cognitive impairment are also seen.

Solution to Question 12:

Resistance to rifampicin and sensitivity to fluoroquinolones is an indication of a short course of bedaquiline.

A shorter MDR-TB regimen of 6 months, BPaLM, is the treatment of choice for MDR/ RR TB as per the latest WHO guidelines.

A longer TB regimen has to be used instead of the shorter MDR TB regimen if:

•

• XDR TB is suspected

• Pregnancy and lactation (Pretomanid is not safe in pregnancy or breastfeeding)

• Patients aged < 14 years

• Resistance to the components of BPaLM regimen

Regimen class	Drugs	Comments
BPaLM regimen6 months M DR/RR TB regimen	Bedaquilline, pretomanid, lin ezolid, moxifloxacin	6 months long
9- 11 months MDR TB regim en	Intensive Phase:Mfx Km/Am Eto Cfz Z H E for 4-6 months Continuation phase:Mfx Cfz Z E for 5 months	9-11 months longInjectables used in intensive phase.
Longer MDR TB regimen	Bdq(6 months) Lfx Lzd Cfz C s for 18-20 months	18-20 months longOral drugs onlyThis regimen is used for XDR TB patients for 20 mon ths.

Solution to Question 13:

In type 1 diabetes mellitus, autoimmune positivity with dysglycemia indicates Stage 3, where clinical diabetes has become evident. This advanced stage is characterized by significant beta cell destruction and insulin deficiency, resulting in symptomatic diabetes.

Individuals with a genetic predisposition are exposed to a trigger that initiates an autoimmune process, leading to the production of islet autoantibodies and a progressive decline in beta cell function.

Stages of type 1 diabetes mellitus are:

Stage 1: It is marked by the presence of two or more islet cell autoantibodies, such as antibodies against zinc transporter (ZnT-8 antibodies), islet cell antibodies (ICA), anti-glutamic acid decarboxylase antibodies (GAD) and insulin autoantibodies (IAA) while glucose levels remain normal.

Stage 2: It involves ongoing autoimmunity and the onset of dysglycemia, where glucose regulation becomes impaired but does not yet meet the criteria for diabetes. This stage reflects a decline in beta cell function but is not yet characterised by the full-blown clinical symptoms of diabetes.

Stage 3: It is defined by the development of hyperglycemia that exceeds the diagnostic criteria for diabetes, indicating significant beta cell destruction and the onset of symptomatic diabetes. At this stage, patients have both autoimmune markers and impaired glucose regulation, confirming advanced disease progression.

Solution to Question 14:

The above clinical history is suggestive of Wernicke's encephalopathy. The current standard of treatment for patients with symptoms of Wernicke's encephalopathy is to give thiamine 100 mg intravenously before administering glucose-containing IV fluids like IV dextrose.

Thiamine deficiency leads to the following:

• Wernicke's encephalopathy: It is more commonly seen in alcoholics with chronic thiamine deficiency. It is characterized by:

• Global confusion

• Truncal ataxia

• Ophthalmoplegia

• Horizontal nystagmus

• Korsakoff psychosis: symptoms in addition to those of Wernicke's encephalopathy include:

• Memory loss

• Confabulation

• Beriberi: It can present with:

• Dry beriberi: sensorimotor peripheral neuropathy, muscle weakness

• Wet beriberi: tachycardia, cardiomegaly, high-output cardiac failure

Solution to Question 15:

The given clinical presentation and the tall T waves on ECG indicate hyperkalemia, which is commonly seen in a patient with chronic kidney disease.

The causes of hyperkalemia are given below -

Intra- to extracellular shift -

• Acidosis

• Hyperosmolality; radiocontrast, hypertonic dextrose, mannitol

• β2-Adrenergic antagonists (non-cardioselective agents)

• Digoxin and related glycosides (yellow oleander, foxglove, bufadienolide)

• Hyperkalemic periodic paralysis

• Rapid tumour lysis

Inadequate excretion

• Inhibition of the renin-angiotensin-aldosterone axis; which increases the risk of hyperkalemia when used in combination with angiotensin-converting enzyme (ACE) inhibitors, renin-inhibitors, angiotensin receptor blockers (ARBs), blockade of the mineralocorticoid receptor(spironolactone, eplerenone, drospirenone), blockade of the epithelial sodium channel (ENaC)(amiloride, triamterene, trimethoprim, pentamidine, nafamostat)

• Advanced renal insufficiency

•

Primary adrenal insufficiency

•

Renal resistance to mineralocorticoid

The steps of management are as follows -

• Immediate antagonism of the cardiac effects of hyperkalemia: Calcium raises the action potential threshold and reduces excitability, without changing the resting membrane potential, thereby reversing the depolarizing blockade due to potassium. 10 mL of 10% calcium gluconate should be infused intravenously over 2-3 mins with cardiac monitoring. Effects start in 1-3 mins.

• Rapid reduction of plasma K+ concentration by redistribution into cells: Insulin shifts K+ into the cells; 10 units of intravenous regular insulin followed by 50 mL of 50% dextrose. Effects start in 10-20 mins.

β2-agonists with insulin have additive effects on potassium concentration. 10-20 mg of nebulized albuterol in 4 mL of normal saline, inhaled over 10 mins. Effects start at 30 mins

• Removal of potassium: Cation exchange resins such as sodium polysterone sulfonate (SPS) exchange Na+ for K+ in the GI tract and increase fecal excretion of K+. 15-30 g of powder in a suspension with 33% sorbitol

Hemodialysis is the most reliable and effective method

Other options:

Option B: Hypokalemia is defined as a plasma concentration of K+ <3.5 meq/L. Common causes include metabolic alkalosis, insulin infusions, diarrhoea, use of diuretics, starvation, and β2-agonists.ECG changes include ST depression, prominent U waves, and prolonged repolarization.



Option C: Hypocalcemia causes prolongation of the QT interval. There is a lengthening of the ST segment, while the T wave (which correlates with the time for repolarization) remains unaltered.

Option D: Hypercalcemia causes shortening of the QT interval and a decrease in the ST segment duration.



Solution to Question 16:

In evaluating the case of a 50-year-old female presenting with dyspnea and angina episodes, it is essential to differentiate between aortic stenosis (AS) and aortic regurgitation (AR). The key factor to consider in aortic stenosis(AS) is that increased myocardial oxygen consumption is more closely associated with increased pressure work rather than increased volume.

AS is the narrowing of the aortic valve. It may be caused due to the calcification of the normal aortic valve, calcification of the congenital bicuspid valve, and rheumatic heart disease. In AS, the heart works harder to overcome the elevated pressure gradient between the left ventricle (LV) and the aorta, leading to increased myocardial oxygen demand. This is in contrast to AR, where the primary issue is volume overload rather than pressure overload.

Three cardinal clinical features of aortic stenosis are:

• Exertional dyspnea - this occurs due to left ventricular (LV) diastolic dysfunction. Raised end-diastolic LV volume and reduced LV compliance lead to increased LV pressure, leading to elevation of pulmonary capillary wedge pressure.

• Angina pectoris - it occurs due to the combination of the increased O2 needs of hypertrophied myocardium and reduction of O2 delivery.

• Syncope - it occurs due to the combination of decreased cardiac output and systemic arterial dilation which lead to a reduction of cerebral perfusion.

The progression of AS involves a gradual narrowing of the valve orifice, which increases pressure work for the LV and leads to concentric hypertrophy as an adaptive response. This process, akin to vascular atherosclerosis, involves endothelial dysfunction, lipid accumulation, and inflammation, contributing to the valve's calcification and eventual stenosis. The pathophysiology of AS results in a significant systolic pressure gradient across the aortic valve. Initially, this

pressure gradient leads to LV hypertrophy as the heart compensates for the increased afterload. Over time, however, this compensatory mechanism becomes maladaptive, leading to decreased LV function and heart failure symptoms. In severe cases, symptoms such as exertional dyspnea, angina, and syncope may develop, reflecting the increased myocardial oxygen demand due to the elevated pressure work.

Solution to Question 17:

The given clinical scenario and marrow aspirate with plasma cells suggest multiple myeloma. The clinical features of fatigue, tingling sensation in the arms, weight loss, and polyuria are due to hypercalcemia.

Plasma cell dyscrasias are B-cell proliferations that contain neoplastic plasma cells that secrete a monoclonal Ig or Ig fragment. Multiple myeloma is a malignant neoplasm of plasma cells characterized by monoclonal immunoglobulin or light chains in the serum and urine. It is a disease of older adults, with a peak age of incidence of 65 to 70 years.

Signs and symptoms include hypercalcemia, fractures/ lytic bone lesions, renal failure and anaemia.

Investigations reveal:

• Radiography - Punched-out lytic lesions on X-ray

• Electrophoresis - Monoclonal elevation of one immunoglobulin

• Urinalysis - Free kappa and lambda light chains (Bence Jones proteins)

• Bone marrow biopsy - 10 to 20% plasma cells (normal is < 5%)

• Peripheral smear - Rouleaux formation. A high level of M proteins in myeloma causes red cells in the peripheral blood smears to stick to one another in linear arrays, a finding referred to as rouleaux formation.

Plasma cell tumours are positive for CD138.

Morphology:

Myeloma cell size varies from small differentiated cells to large immature and undifferentiated cells. Plasmablasts with vesicular nuclear chromatin and a prominent single nucleolus, or bizarre, multinucleated cells may predominate. Other cytologic variants include flame cells and Mott cells.

Flame cells have fiery red cytoplasm, and Mott cells are with multiple grape-like cytoplasmic droplets. Other inclusions include fibrils, crystalline rods, and globules.

Image showing Mott cell:



The globular inclusions are Russell bodies (if cytoplasmic) or Dutcher bodies (if nuclear). Rouleaux formation is characteristic but not specific.

Image showing Russell and Dutcher bodies:



Other options:

Options A and D: Megaloblastic anaemia and subacute combined degeneration(SACD) occur due to vitamin B12 deficiency. Patients with SACD present with loss of vibration sense and proprioception, paresthesia, unsteady gait, loss of deep tendon reflexes, peripheral neuropathy, and optic neuropathy.

Option C: Diabetes does not lead to subacute degeneration.

Solution to Question 18:

The given clinical scenario is suggestive of an exudative pleural effusion according to Light's criteria. Among the given options, tuberculosis can lead to exudative pleural effusion.

In any patient diagnosed with a pleural effusion, an effort should be made to find the cause. This requires a diagnostic thoracocentesis and analysis of the pleural fluid in terms of whether it is a transudate or an exudate based on Light's criteria:

• Pleural fluid protein/serum protein >0.5

• Pleural fluid LDH/serum LDH >0.6

• Pleural fluid LDH is more than two-thirds the upper limit of normal for serum

If the pleural fluid meets one of these criteria, it is considered exudative pleural effusion. Sometimes, special investigations like ADA and cytology might be required to rule out tuberculosis and malignancy respectively.

For thoracocentesis, a posterior approach is preferred. The patient should sit on the edge of the bed, leaning forward with the arms abducted onto a pillow on a bedside stand. The patient might have severe dyspnea, so it is important to assess whether they can maintain this position for 10 minutes before performing the procedure. The entry site for the thoracocentesis is at the superior aspect of the rib. This prevents any injury to the intercostal nerve and vessels.

While maintaining gentle negative pressure, the needle should be slowly advanced into the pleural space. If a diagnostic tap is being performed, 30-50 mL of fluid will be adequate. If a therapeutic thoracocentesis is being performed, a maximum of 1 litre of pleural fluid should be withdrawn at a given time. Withdrawal of quantities more than this can cause re-expansion pulmonary edema. A chest radiograph should be obtained post-procedure to evaluate for a pneumothorax. The patient should be instructed to inform the doctors if new shortness of breath develops.

Other options

Option A and C: Congestive heart failure and liver cirrhosis lead to transudative pleural effusion.

Solution to Question 19:

The adolescent's symptoms, which include fever, purpura, swelling of joints (right knee and left ankle), and abnormal heart sounds, are indicative of acute rheumatic fever (ARF)

Acute rheumatic fever (ARF) is an inflammatory disease that can develop after a Group A Streptococcal infection, such as strep throat. It commonly affects the heart, joints, skin, and central nervous system. Symptoms include fever, migratory arthritis (joint swelling and pain), and carditis. The presence of abnormal heart sounds, such as diastolic murmurs, suggests valvular involvement, typically of the mitral valve. ARF can also present with skin manifestations like purpura.

Other Options:

Option B: Juvenile idiopathic arthritis typically presents with persistent joint swelling and stiffness, not with the acute symptoms or heart involvement seen in ARF.

Option C: Rheumatoid Arthritis is a chronic autoimmune disease that primarily affects adults, though juvenile forms exist. It usually presents with symmetrical joint involvement and positive rheumatoid factor in many cases, which is absent in this patient.

Option D: Seronegative Arthropathy is a group of disorders that includes conditions like ankylosing spondylitis and psoriatic arthritis, which typically involve the spine and entheses rather than the acute joint swelling, purpura, and heart involvement seen in ARF.

Solution to Question 20:

The given clinical scenario of a patient with easy fatiguability, skin and soft tissue hyperpigmentation, and a low blood pressure and hyponatremia is suggestive of primary adrenal insufficiency(Addison's disease).

Primary adrenal insufficiency (Addison's disease) is most commonly due to autoimmune adrenalitis. These patients are likely to develop other autoimmune diseases such as vitiligo, premature ovarian failure, and autoimmune thyroid disease. Other causes of Addison's disease include destruction of the adrenal gland due to infections such as TB, hemorrhage (Waterhouse-Friderichsen syndrome due to meningococcemia) or infiltration.

Secondary adrenal insufficiency occurs due to hypothalamic-pituitary axis dysfunction. This can include iatrogenic suppression due to chronic therapy with glucocorticoids, or because of pituitary and hypothalamic tumors. Pituitary apoplexy because of hypovolemic shock during parturition (Sheehan syndrome) can also cause secondary adrenal insufficiency.

The patients present with fatigue and loss of energy. Primary adrenal insufficiency causes hyperpigmentation due to excess ACTH, which stimulates the melanocytes to produce more pigment. Hyperpigmentation is maximally seen in areas exposed to sunlight or that which undergo friction. However, secondary adrenal insufficiency presents with skin paleness due to the complete absence of ACTH. The diagnosis of adrenal insufficiency is established by the ACTH stimulation test.

ACTH stimulation test (also known as the short cosyntropin test or short Synacthen test) involves the administration of an exogenous, synthetic peptide that has an activity similar to ACTH on the adrenal glands. Serum cortisol levels are measured prior to, and after the administration of the synthetic peptide. Cortisol levels lower than the cut-off value indicate a failure of the adrenal glands to increase the production of cortisol in response to synthetic ACTH — thus confirming the diagnosis of adrenal insufficiency. Note that this will be seen in both primary and secondary adrenal insufficiency.

Primary and secondary adrenal insufficiencies can be differentiated biochemically:

Treatment is with oral hydrocortisone 15-25 mg/day in 2-3 divided doses. Long-acting agents are not preferred because they cause receptor activation even when the cortisol levels are supposed to be physiologically low in the body. Mineralocorticoid replacement is done with fludrocortisone 100-150 μg.

Other options

Option A: Primary hyperaldosteronism (Conn syndrome), is characterized by excessive production of aldosterone by the adrenal glands. Patients commonly present with fatigue, muscle weakness, headaches, palpitations, and high blood pressure. Hypokalemia can cause muscle cramps, weakness, and even paralysis in severe cases. The diagnosis is suspected based on clinical presentation and laboratory findings of elevated plasma aldosterone and suppressed plasma renin activity.

Option C: Vitamin B12 deficiency can cause subacute combined degeneration. Patients present with sensory ataxia (involvement of the dorsal column) and UMN type of motor symptoms such as spasticity and positive Babinski sign (involvement of corticospinal tract). Because of coexisting peripheral neuropathy, deep tendon reflexes can be absent. Some patients also develop cortical defects such as dementia. These patients can also have skin hyperpigmentation, but the presence of hyponatremia and low blood pressure makes this unlikely.

Option D: Cushing's syndrome is characterized by clinical features that result from chronic exposure to excess glucocorticoids that can be due to prolonged exogenous administration of glucocorticoids, excess endogenous cortisol secretion, and ectopic ACTH production. Clinical features include moon facies, purple skin striae, abnormal masculinization (acne, bruising, enlargement of the clitoris in girls), pubertal developmental delay and amenorrhea in girls, proximal muscle weakness, osteoporosis, hypertension, hyperpigmentation, hypokalemic alkalosis, hyperglycemia.

Primary adrenal insufficiency	Secondary adrenal insufficiency
Serum cortisol after ACTH stimulation: low	Serum cortisol after ACTH stimulation: low
Serum ACTH:high	Serum ACTH:low
Plasma renin:high(minerocorticoid deficiency also present, since adrenal gland itself is not functioning)	Plasma renin:normal(adrenal gland itself is functional and can be regulated by renin-angiotensin system)

Solution to Question 21:

The given clinical scenario along with the image showing the Kayser–Fleischer ring, is suggestive of Wilson's disease and the best investigation to confirm the same is by measuring the hepatic copper content.

The diagnosis of Wilson's disease is based on the following:

• Increased 24-hour urinary excretion of copper (the most specific screening test)(Option C)

• Increase in hepatic copper content (the most sensitive and accurate test)

• A decrease in serum ceruloplasmin(Option A)

• Kayser–Fleischer rings

• Genetic testing for mutations in the ATP7B gene

Serum copper levels are of no diagnostic use since they may be low, normal, or elevated depending on the stage of the disease. (Option D)

Wilson's disease is an autosomal recessive disorder caused by loss of function mutation of the ATP7B gene, which codes for the enzyme copper-transporting ATPase present on the hepatocyte canalicular membrane.

In general, the free copper is taken up by hepatocytes and incorporated to form ceruloplasmin gets released into the bloodstream, and accounts for 90% to 95% of plasma copper. Excess hepatic copper gets transported into the bile and eventually excreted in the faeces.

The defect in the ATP7B gene impairs copper transport into the bile, its incorporation into ceruloplasmin, and secretion of ceruloplasmin into the blood. This results in a decrease in circulating ceruloplasmin levels and increased levels of hepatic copper. Total serum copper level may be decreased due to the deficiency in ceruloplasmin.

The excessive accumulation of copper injures the hepatocytes. This in turn releases the non-ceruloplasmin-bound copper into the bloodstream, which later gets deposited in tissues like the basal ganglia of the brain and the cornea. This results in behavioural disturbances as well as extrapyramidal symptoms and the characteristic KF ring that is seen on slit-lamp examination. There is a marked compensatory urinary copper excretion. However, these levels do not suffice to prevent the deposition of copper in tissues.

Solution to Question 22:

The given clinical scenario and autopsy findings are suggestive of hypertrophic cardiomyopathy (HCM). The image given is a histologic section of cardiac muscle showing myocyte disarray and extreme hypertrophy as seen in HCM.

HCM is the leading cause of sudden cardiac death in young adults. It is defined as left ventricular hypertrophy of >15mm on ECHO. The left ventricular myocardium is poorly compliant. This leads to abnormal diastolic filling and intermittent ventricular outflow obstruction, i.e., primarily diastolic dysfunction.

The gross features of HCM are:

• Massive myocardial hypertrophy usually without ventricular dilation

• Asymmetric septal hypertrophy- disproportionate thickening of the ventricular septum relative to the left ventricle free wall

• Banana-shaped left ventricle- due to bulging of the ventricular septum into the lumen

The image given below shows changes in hypertrophic cardiomyopathy compared to a normal heart. HCM shows asymmetric septal hypertrophy and a thickened left ventricle wall.



Histologic features of the HCM are:

• Massive myocyte hypertrophy- with transverse myocyte diameters frequently greater than 40 µm, normal is approximately 15 µm.

• Myofiber disarray- haphazard disarray of bundles of myocytes, individual myocytes, and contractile elements in sarcomeres within cells

• Interstitial fibrosis and replacement fibrosis

The image given below is a histologic section of cardiac muscle showing myocyte disarray and extreme hypertrophy seen in an abnormal HCM septum.



Solution to Question 23:

The given clinical scenario is suggestive of an upper motor lesion secondary to trauma and ankle clonus in the right foot will be seen in this patient along with other clinical findings.

An upper motor neuron (UMN) refers to corticospinal tract neurons and brainstem neurons that control spinal motor neurons. UMN lesion is characterized by signs such as spasticity, hypertonia, and hyperreflexia, indicating a disruption in the neural pathways that control voluntary movement. These lesions can result from various conditions, including stroke, trauma, or degenerative diseases.

In a normal response to uncomfortable or painful stimulation of the foot, there is plantar flexion of the toes, often accompanied by flexion withdrawal of the limb. However, with a UMN lesion, the response will be an extensor plantar response, known as a positive Babinski sign, characterized by extension of the great toe and possibly fanning of other toes, along with contraction of the tensor fascia lata.

Clonus is a rhythmic oscillatory series of muscle contractions that is induced by stretching the tendon. It can be elicited in many joints but is most commonly seen in the ankle joint by suddenly dorsiflexing the foot and maintaining light upward pressure. It is seen in UMN lesions and is associated with other signs of UMN lesions like spasticity.

Solution to Question 24:

After complete ileal resection, a patient with fatigue is unlikely to have microcytic anemia. In contrast, the patient is likely to have macrocytic anemia due to B12 deficiency.

Vitamin B12 is absorbed in the ileum. Vitamin B12 binds to intrinsic factor, a protein secreted by the parietal cells of the stomach, and the complex is absorbed across the ileal mucosa. Hence after ileal resection, there is impaired absorption of vitamin B12.

Vitamin B12 deficiency can cause subacute combined degeneration(option D). Patients present with sensory ataxia (involvement of the dorsal column) and UMN-type motor symptoms such as spasticity and positive Babinski sign (involvement of corticospinal tract). Because of coexisting peripheral neuropathy, deep tendon reflexes can be absent. Some patients also develop cortical defects such as dementia.

Its deficiency can manifest as pernicious anemia, dementia, spinal cord degeneration, and homocysteinemia.

Other options:

Option A: Vitamin B12 deficiency shows hypersegmented neutrophils (>5 lobes), ovalocytes, and some amount of cyto-nuclear dyssynchrony.

Option C: In the Romberg test, the standing patient is asked to close their eyes. An increased loss of balance is interpreted as a positive Romberg's test. It tests the proprioception, which requires dorsal columns of the spinal cord. Hence, with the involvement of the dorsal column due to Vitamin B12 deficiency, the Romberg's sign is seen.

Solution to Question 25:

In a patient with dyspnea, a CT scan shows a mediastinal mass with an attenuation of -62 HU, indicative of fat content, along with areas of calcification. This suggests the presence of a mediastinal mass comprised of fat and calcified tissue.

Thymic neoplasms usually are detected in the anterior mediastinum. Fewer than 4% of thymic tumors occur in the middle mediastinum, defined as mediastinal space surrounded by the left brachiocephalic vein, superior vena cava, esophagus, trachea, and main bronchus up to the

cross-section of the left brachiocephalic vein and the center of the trachea. Thymic carcinoma is a rare thymic neoplasm. Moreover, few reports of middle mediastinal thymic carcinomas are histopathologically diagnosed as having immunohistochemical features.

Other options:

Option A: The CT shows a mediastinal mass but no clear evidence of vascular invasion.

Option B: It shows a picture of a mediastinal mass rather than aorta.

Option D: A mediastinal hydatid cyst on a chest CT can confirm the diagnosis by objectifying a mass of fluid density, often very limited, unmodified by the injection of contrast, however, contrast uptake by a pericyst can be observed. The presence of peripheral calcifications supports the diagnosis. A predilection for the posterior compartment of the mediastinum has been reported in the literature.

Solution to Question 26:

The given laboratory values of low pH, low bicarbonate, low potassium and urinary pH > 5.5 are suggestive of metabolic acidosis due to type 1 renal tubular acidosis (RTA).

Metabolic acidosis associated with RTA I has a normal serum anion gap due to impaired urinary acidification. It is also associated with hypercalciuria. The hypercalciuria leads to a tendency to form calcium oxalate and calcium phosphate stones. The patients can also have recurrent UTIs and hematuria. The urinary anion gap is positive in these cases, and urine pH is greater than 5.5.

Solution to Question 27:

The clinical scenario of tremors, weight loss, and the image showing exophthalmos are suggestive of Graves ophthalmopathy. The muscles and connective tissue around the eye share similar antigens with diseased organ i.e. Thyroid.

Thyroid-stimulating hormone (TSH) receptors have been found in the orbit's connective tissues and fibroblasts. TSH stimulates the production of inflammatory cytokines. Thyroid eye disease is considered an autoimmune disease where orbital fibroblasts are the primary target and extraocular muscles are secondarily involved. Thyroid follicular cells and orbital fibroblasts likely share a common antigen.

TAO is also linked to increased glycosaminoglycan synthesis, which results in restrictive myopathy. Fat accumulation is caused by orbital fibroblast proliferation and differentiation into adipocytes.

Graves ophthalmopathy, also known as dysthyroid eye illness or thyroid-associated ophthalmopathy (TAO), is one of the most serious ocular manifestations of Graves disease. Between the ages of 25 and 50, it's the leading cause of unilateral and bilateral proptosis in adults. It is seen in 10% of cases without hyperthyroidism.

Proptosis, eyelid retraction, lid retraction, lid lag on downgaze, restricted myopathy, and perhaps compressive optic neuropathy are clinical manifestations that could potentially be used for diagnosis.

American Thyroid Association (ATA) has classified Graves' ophthalmopathy, irrespective of the hormonal status into the following classes. This classification is also known as Werner classification which reflects the severity of the ophthalmopathy and is well known by the acronym NO SPECS.

Treatment of thyroid ophthalmopathy:

Other options

Option A: Thyroid eye disease is associated with HLA-DR3 and HLA-B8.

Option B: The most common ocular motility defect is unilateral elevator palsy due to involvement of the inferior rectus followed by limitation of abduction due to medial rectus palsy.

Option D: Thyroid eye disease is 4-6 times more common in females.

Grade	Feature	Acronym
0	No signs or symptoms	N
1	Only signs (lid retraction), No symptom	O
2	Soft tissue involvement along with signs and symptoms(chemosis, lacrimation, photophobia)	S
3	Proptosis is well-established	P
4	Extraocular muscle involvement (limitation of movement and diplopia)	E
5	Corneal involvement (exposure keratitis).	C
6	Loss ofSight due to optic nerve involvement with disc pallor or papilloedema and visual field defects.	S

Non-surgical management	Surgical management
Systemic steroids	Orbital decompression
Radiotherapy	Extraocular muscle surgery
Teprotumumab (IGF-1R inhibitor)	Eyelid surgery

Solution to Question 28:

The patient in the clinical vignette with chest pain and palpitations, stable vitals and echocardiography findings of the diastolic collapse of the right atrium is suggestive of cardiac tamponade, and the most appropriate management will be pericardiocentesis.

Cardiac tamponade is characterised by the accumulation of fluid in the pericardial space, leading to impaired diastolic relaxation of the cardiac chambers. This leads to a decline in cardiac output.

Pericardiocentesis, also known as pericardial tap, is performed with sub-xiphoid, parasternal, or apical approaches. The most common approach is a sub-xiphoid approach under echocardiographic guidance. As a preparatory step, IV saline should be administered to the patient, and intrapericardial pressure should be measured without delaying the treatment, as the condition can be fatal. The fluid should be completely drained, and a multiholed catheter is left in situ to drain the fluid which re-accumulates. The fluid obtained should be sent for analysis.

The echocardiography shows a swinging heart sign, which is characteristic of cardiac tamponade. The pericardial pressures exceed the left ventricular diastolic filling pressures, causing the collapse of heart chambers, IVC plethora, and respiratory variation in volumes and flows.



Other options:

Option A - NSAIDs can be given for pain management, but the primary aim is to reduce the pressure on the heart by urgent pericardiocentesis or surgical drainage. They can, however, be used in uremic pericarditis along with dialysis.

Option C- Intercostal chest tube drainage is done in tension pneumothorax, which is a close differential diagnosis for cardiac tamponade where there will be reduced air entry and mediastinal shift.

Option D: Steroids are not used in the management of acute cardiac tamponade. They can be used in pericardial effusion secondary to rheumatological and inflammatory diseases like SLE, rheumatoid arthritis, scleroderma, and polyarteritis nodosa to prevent inflammation and reaccumulation of fluid.

Solution to Question 29:

The given clinical scenario of an alcoholic patient with epigastric pain radiating to the back, elevated amylase and lipase with the chest X-ray findings of diffuse bilateral lung opacities is suggestive of non-cardiogenic pulmonary oedema or ARDS (acute respiratory distress syndrome) secondary to acute pancreatitis.

Acute pancreatitis gives rise to inflammatory processes and endothelial damage that result in proteinaceous interstitial oedema. Unlike cardiogenic pulmonary oedema, this does not clear with diuretic therapy and persists for days to weeks. Early findings on a chest X-ray may be normal or may have diffuse bilateral interstitial opacities, obscuring the pulmonary vascular markings which progress later to a more diffuse and extensive consolidation.

Clinical disorders commonly associated with ARDS:

• Direct lung injury: Pneumonia, aspiration of gastric contents, pulmonary contusion, near-drowning, toxic inhalation injury.

• Indirect lung injury: Sepsis, severe trauma- multiple fractures, flail chest, head trauma, burns, multiple transfusions, drug overdose, pancreatitis, post-cardiopulmonary bypass

Other options:

Option A: Aspiration pneumonia results from aspiration of oropharyngeal or gastric contents into the lower respiratory tract. Although it is more common in alcoholics, the clinical history of the patient suggests pancreatitis, making ARDS a more likely diagnosis. Below is a chest X-ray showing diffuse opacities in the right middle and lower lobes.



Option C: The chest radiograph given below shows extensive bilateral reticulonodular infiltrates seen in miliary tuberculosis, which is the widespread dissemination of tuberculosis via the hematogenous route.



Option D: Pulmonary embolism is an embolic occlusion of the pulmonary arterial system in the presence of medical or surgical risk factors. CT pulmonary angiography is the investigation of choice in pulmonary embolism.

Given below are the chest X-rays of pulmonary embolism showing Westermark's sign (hyperlucent area with decreased vascularity due to oligemia of the involved part of the lung) and Hampton's hump (peripheral wedge-shaped opacity with apex pointing towards the occluding vessel), respectively.





Solution to Question 30:

The most likely cause of death in a patient suffering from Idiopathic pulmonary fibrosis (IPF) is respiratory failure.

The natural history is usually of steady decline, however, some patients present with acute exacerbations manifesting as dyspnea, impaired gas exchange, or a new consolidation on HRCT(High-Resolution CT).In advanced cases, patients may develop pulmonary hypertension and features of right heart failure. The finding of high numbers of fibroblastic foci on biopsy is suggestive of more rapid deterioration.

IPF is the most common interstitial lung disease occurring due to an unknown cause. The patients are usually over 60 years of age and present with progressive exertional dyspnea and dry cough. On examination, fine inspiratory, velcro crackles may be heard. Digital clubbing and cyanosis are signs of advanced disease.

Pulmonary function tests (PFT) will show a restrictive deficit and reduction of TLC, FEV1, and FVC. HRCT shows predominant subpleural and basal involvement in the form of reticulation, honeycombing, and traction bronchiectasis. Pirfenidone inhibits TGF-beta1-stimulated collagen synthesis and has anti-fibrotic, anti-inflammatory & anti-oxidant effects. It is also used to manage acute exacerbations of pulmonary hypertension along with slowing the progression of IPF. Nintedanib is another drug also used for IPF based on the mechanism of tyrosine kinase inhibition. However, it carries a poor prognosis.

Other options:

Option A: Patients may develop pulmonary hypertension and features of right heart failure in the advanced stage only.

Option B and D: Pulmonary embolism and Lung carcinoma are not directly related to IPF.

Solution to Question 31:

The given clinical scenario of a patient presenting with fever without evidence of meningitis along with an image showing a ring-enhancing lesion with enhancement on diffusion-weighted MRI (Diffusion restriction) is suggestive of Brain abscess.

Ring-enhancing brain lesions can be caused by the following conditions:

• Neurocysticercosis

• Tuberculoma

• Cerebral abscess

• Metastasis

• Toxoplasmosis

• Glioblastoma

• Subacute infarct /hemorrhage/contusion

Diffusion restriction is seen in

• Brain abscess

• Acute Infarcts

• Epidermoid cysts

• Subacute bleeds/ Diffuse axonal injury

• Lymphoma

The combination of ring-enhancing lesion along with diffusion restriction is suggestive of Brain abscess.

The image below shows ring enhancement in glioblastoma:



Solution to Question 32:

The clinical scenario of the patient with fever, cough with foul-smelling sputum, clubbing and a chest X-ray showing air-fluid level is suggestive of a lung abscess.

A lung abscess is caused by microbial infection, leading to necrosis and cavitation of the lung. These could either occur following aspiration or secondary to bronchial obstruction by a foreign body or malignancies, often in the background of immunosuppression. The patients usually present with complaints of fever, cough with sputum, and chest pain. If the lung abscess is putrid, it may produce discoloured and foul-smelling sputum. Chronic cases may present with anaemia, night sweats and fatigue. Patients usually have poor dental hygiene and gingival disease. Examination findings include clubbing, absent gag reflex, and cavernous breath sounds.

Radiological features of lung abscess:

• A lung cavity with an air-fluid level

• The cavity wall is thick and irregular with a surrounding pulmonary infiltrate and acute angle to the chest wall.

• Computed tomography (CT) is more sensitive than chest radiography and is used to detect small cavities

The X-ray image given below shows thick-walled cavities with air-fluid levels on lateral view.



Other options:

Option A: In pneumothorax, the patient will present with a history of breathlessness, and chest X-ray will show deep costophrenic angle, hyperlucency (darker area without lung markings) with inward shifting of the visceral pleural line (sharp white line). In tension pneumothorax, there will be a mediastinal shift to the opposite side.

The image below shows a left pneumothorax with total left lung collapse and tracheal deviation to the right.



Option B: Pneumatoceles are thin-walled cavities in the lung parenchyma that are seen in respiratory tract infections, most often caused by Staphylococcus aureus in children and Pneumocystis jirovecii in immunocompromised adults. Given below is the chest X-ray showing pneumatocele.



Option C: Empyema is the collection of pus in the pleural cavity. It is usually associated with pneumonia but may also develop after thoracic surgery or thoracic trauma. At the margins of the empyema, the pleura can be seen dividing into parietal and visceral layers, the split pleura sign. It is the most sensitive and specific sign on CT and helps distinguish an empyema from a peripheral lung abscess.

The image given below shows a split pleura sign of empyema.



Solution to Question 33:

The clinical vignette of an athlete with chest pain and dyspnea whose family history is suggestive of sudden cardiac death due to HCM and a harsh systolic murmur at the left sternal border are all in favour of HCM murmur. It increases in intensity while performing the Valsalva manoeuvre.

The murmur of HCM is an ejection systolic murmur best heard at the lower left sternal border.

The intensity of the mid-systolic murmur is

Increased by -Valsalva strain phase, Standing posture, and exercise

Decreased by - Squatting, passive leg raising, and handgrip.

Valsalva manoeuvre involves the performance of forced expiration against a closed glottis.

HCM is the leading cause of sudden cardiac death in young adults. It is defined as left ventricular hypertrophy of >15mm on ECHO. The left ventricular myocardium is poorly compliant. This leads to abnormal diastolic filling and intermittent ventricular outflow obstruction, i.e., primarily diastolic dysfunction.

Gross features of HCM:

• Massive myocardial hypertrophy usually without ventricular dilation

• Asymmetric septal hypertrophy- disproportionate thickening of the ventricular septum relative to the left ventricle free wall

• Banana-shaped left ventricle- due to bulging of the ventricular septum into the lumen

The image given below shows changes in hypertrophic cardiomyopathy compared to a normal heart. HCM shows asymmetric septal hypertrophy and a thickened left ventricle wall.



Histologic features of the HCM:

Massive myocyte hypertrophy- with transverse myocyte diameters frequently greater than 40 µm, normal is approximately 15 µm.

Myofiber disarray- haphazard disarray of bundles of myocytes, individual myocytes, and contractile elements in sarcomeres within cells

Interstitial fibrosis and replacement fibrosis

The image given below is a histologic section of cardiac muscle showing myocyte disarray and extreme hypertrophy seen in an abnormal HCM septum.



Pathophysiology:

HCM is characterized by reduced stroke volume due to impaired diastolic filling. Approximately 25% of patients with HCM have dynamic obstruction to the left ventricular outflow as the anterior mitral leaflet moves toward the ventricular septum during systole. The compromised cardiac

output in conjunction with a secondary increase in pulmonary venous pressure explains the exertional dyspnea seen in these patients.

Major clinical problems in HCM are atrial fibrillation, mural thrombus formation leading to embolization and stroke, cardiac failure, ventricular arrhythmias, and sudden death.

ECG - findings:

Prominent abnormal Q waves

P wave abnormalities

Left axis deviation

Deeply inverted T waves

Echocardiography:

LV wall thickness ≥15 mm

Systolic anterior motion of the mitral valve

LVOT obstruction

The management options in patients with HCM:

Beta-blockers

Non-dihydropyridine calcium channel blockers

Disopyramide

Drugs to be avoided in patients with significant LVOT obstruction:

Dihydropyridine calcium channel blockers

Nitroglycerin

Angiotensin-converting enzyme inhibitors, ARBs

Diuretics

Preventive intra-cardiac defibrillator is indicated for patients with HCM and documented cardiac arrest or sustained ventricular tachycardia.

Note: Valsalva maneuver decreases the intensity of all murmurs except MVP (mitral valve prolapse) and HCM.

Solution to Question 34:

The given clinical scenario of a patient with right hemiplegia, left-sided facial nerve palsy, and left horizontal gaze palsy is suggestive of Foville syndrome or dorsal pontine syndrome.

Brainstem syndromes cause crossed hemiplegia with ipsilateral cranial nerve palsies and contralateral hemiplegia. Pontine syndromes occur due to the involvement of pontine branches of the basilar artery and anterior inferior cerebellar artery.

In Foville's syndrome, there is a dorsal pontine injury (lesion in the abducens nucleus) that causes lateral/ horizontal gaze palsy. In Millard-Gubler syndrome, there is a ventral pontine injury (lesion

in the abducens fascicle) that causes lateral rectus weakness.

Locked-in syndrome occurs when a pontine stroke causes quadriplegia and paralysis of all voluntary muscles except vertical movements of the eyes and blinking. The vertical gaze nucleus, the rostral interstitial nucleus of medial longitudinal fasciculus, is located in the midbrain and is hence preserved.

CNS lesion	Effects
Foville's Syndrome/ Dorsal pontine syndrome	Lateral gaze palsyIpsilateral facial nerve palsyContralateral hemiparesis
Millard–Gubler Syndrome/Ventral pontine syndrome	Ipsilateral abducens nerve weaknessIpsilateral facial nerve palsyContralateral hemiplegia

Solution to Question 35:

The given clinical scenario of a patient with hypertension and diabetes, and frontal bossing, thick palms and soles, and gaping of central incisors is suggestive of acromegaly. Age-specific IGF-1 levels are used as a screening test in this condition.

The confirmation of acromegaly is done by administering 75g of oral glucose load to the patient. Normally, GH should be suppressed to less than 0.4 ug/L after the oral glucose load. A failure of suppression of growth hormone confirms the diagnosis of acromegaly.

Acromegaly is a result of growth hormone hypersecretion due to a pituitary adenoma. It can also occur due to extra pituitary lesions. If growth hormone hypersecretion occurs before the fusion of the growth plates at puberty, it leads to gigantism.

Acral bony overgrowth in acromegaly manifests as frontal bossing, increased hand and foot size, mandibular enlargement, prognathism, and widened space between incisors. Soft tissue swelling results in increased heel pad thickness, coarse facial features, and a large fleshy nose. Generalized visceromegaly is seen including cardiomegaly, macroglossia, thyroid enlargement, etc. Serious complications include diabetes mellitus, obstructive sleep apnoea, and increased risk of colonic polyps and cancer.

The physiological actions of the growth hormone are mostly mediated through IGF-1, which is synthesized in the liver. Since the growth hormone is released in a pulsatile manner, serum IGF-1 levels are usually measured to assess growth hormone activity. Serum IGF-1 level is decreased in pituitary failure, defect or blockade of GH receptor. Since the low-calorie state is associated with growth hormone resistance, IGF-1 is also decreased in conditions such as cachexia, malnutrition, and sepsis. Serum IGF-1 level is increased in acromegaly.

Acromegaly can be treated by transsphenoidal resection of the tumor. It can also be managed medically by using octreotide (a somatostatin analog), pegvisomant (Growth Hormone receptor antagonist), dopamine agonists, or radiation.

Other options:

Option A: A failure of suppression to <0.4ng/ml using ultrasensitive assays or to <1ng/ml using standard assays after an oral glucose load is used as a confirmatory test.

Option B: Since the growth hormone is released in a pulsatile manner, measurements of a single GH level are not used in the diagnosis of acromegaly.

Option D: Though, GH levels > 40 ng/ml indicate poor prognosis but it is not routinely done to assess the severity of acromegaly.

Solution to Question 36:

Normal PR interval ranges between 0.12 and 0.20 seconds since this patient's ECG has a PR interval of 0.21 s(prolonged), it is suggestive of a first-degree heart block.

This type of block is characterized by a fixed, prolonged PR interval without any dropped beats. It indicates a delay in conduction through the AV node, but all atrial impulses still reach the ventricles. It is most often asymptomatic. Canon waves are observed in JVP. It carries a very good prognosis and no treatment is needed.

A heart block is caused by an interruption of impulse conduction within the conducting system of the heart. It may be sinoatrial, atrioventricular, or intraventricular. Atrioventricular blocks are further divided into type I, type II, and type III blocks. The type II blocks are subdivided into Mobitz type I and Mobitz type II blocks.

Other options:

Option B: Normal ECG will show a PR interval between 0.12 and 0.20 seconds

Option C: Second-Degree AV Block Mobitz Type I (Wenckebach). is characterized by a progressive prolongation of the PR interval until a beat is eventually dropped. This type of block is usually benign and often does not require treatment.

Second-degree AV Block Mobitz Type II involves intermittent dropped beats without the progressive prolongation of the PR interval. It is often associated with a fixed PR interval and is more serious than Mobitz Type I because it has a higher risk of progressing to a complete heart block.

Option D: Third-degree AV Block also known as complete heart block, involves complete dissociation between atrial and ventricular activity. The atria and ventricles beat independently of each other, which can result in severe bradycardia and require immediate medical intervention.

Solution to Question 37:

The clinical scenario points to a diagnosis of atrial fibrillation and the JVP finding seen would be absent a wave.

Atrial fibrillation is the most common sustained arrhythmia. There is rapid and irregular atrial activation which is transmitted through the atrioventricular node. The conducted ventricular rate is usually rapid and irregularly irregular. It can vary from 110-160 beats/minute. A pulse deficit is an important clinical finding associated with the condition. This arises due to preload reduction.

Symptoms include palpitations, fatigue, presyncope, and dyspnea. An ECG shows the absence of P waves.



In acute atrial fibrillation with hemodynamic instability, electrical or chemical cardioversion is the first step. The latter is done by intravenous administration of class III antiarrhythmic - ibutilide. In a stable patient, ventricular rate control is the primary aim. This is done by intravenous or oral administration of beta-blockers. Calcium channel blockers such as verapamil and diltiazem can also be used.

If the current episode duration is >48 hours, cardioversion can precipitate thromboembolic complications. In such cases, cardioversion should be carried out only after 3 weeks of anticoagulation. It has to be continued for 4 weeks after cardioversion. Another option is to evaluate the presence of a thrombus in the left atrial appendage. This can be done either by transesophageal echocardiography (TEE) or high-resolution CT.

If the duration is <48 hours and there are no high-risk factors, cardioversion can be done. The CHA2DS2-VASc scoring system is used to assess stroke risk.

Warfarin is recommended in patients with rheumatic mitral stenosis or mechanical heart valves who present with atrial fibrillation. In mitral stenosis, thrombi arise frequently from the dilated left atrium. Atrial fibrillation in the absence of mitral stenosis is known as nonvalvular atrial fibrillation. Oral factor Xa inhibitors such as apixaban, edoxaban, or rivaroxaban can be used in such cases.

Other options:

Option A: Rapid x-descent is seen in Cardiac tamponade.

Option C: Cannon a wave is seen in atrioventricular dissociation and contraction of the right atria against a closed tricuspid valve.

Option D: Rapid y-descent is seen in Constrictive pericarditis and isolated severe tricuspid regurgitation.

Solution to Question 38:

Chronic hepatitis B with active viral replication has the following serological findings: HBsAg positive, Anti-HBs negative, Anti-HBc IgG positive, HBeAg positive.

After a person is infected with the hepatitis B virus (HBV), the first serologic marker to increase in level is HBsAg. It becomes detectable in the serum between 1-12 weeks. It becomes undetectable 1–2 months after the onset of jaundice and rarely persists beyond 6 months. Once the HBsAg becomes undetectable, the anti-HBs antibody starts rising in serum and remains detectable indefinitely thereafter. The gap between the disappearance of HBsAg and the appearance of anti-HBs antibodies is called the window period.

However, if HBsAg persists beyond 6 months, it is likely to be chronic hepatitis B, marked as an orange line in the image.

HBcAg is located inside the cells and is usually enclosed within an HBsAg coat when present in the serum. As a result, HBcAg is not commonly detectable in the serum of patients with HBV infection since naked core particles are not found in the bloodstream.

Conversely, anti-HBc (red line) can be detected in the serum shortly after the emergence of HBsAg, usually within 1-2 weeks. Anti-HBc may be the only serologic marker useful for the identification of HBV infection during the window period. Initially, the anti-HBc is of the IgM variety, but in chronic infections, it is of the IgG class.

HBeAg (green line) appears shortly after the appearance of HBsAg. Its appearance correlates with high levels of virus replication. Anti-HBe correlates with a period of lower infectivity or recovery from the infection.

Other options:

Option A: Chronic HepB with HBe Ag negative could be possible if there had not been a green line in the graph marked as HBeAg, which is persisting for more than 52 weeks.

Option B: Persistence of Hepatitis B surface antigen (HBsAg) beyond 6 months indicates chronic infection (orange line), ruling out the possibility of a resolved infection.

Option D: Acute Hepatitis B patient is unlikely to have a history dating back a year ago and persistent surface antigen.

Solution to Question 39:

The cause of hypercalcemia in the female with myasthenia gravis on mycophenolate mofetil and pyridostigmine is likely to be a parathyroid adenoma.

Parathyroid adenomas cause primary hyperparathyroidism and may present as a solitary adenoma or as a part of genetic syndromes like MEN syndromes(parathyroid hyperplasia). Elevated parathyroid hormone increases blood calcium via bone resorption, intestinal absorption and renal reabsorption.

Primary hyperparathyroidism is mostly asymptomatic. However, it may also present as painful bones (subperiosteal resorption, osteitis fibrosa cystica), renal stones, abdominal groans (pain, nausea, vomiting, heartburn), neuropsychiatric (irritability, paranoid delusion, depression), fatigue, and proximal muscle weakness. Excess calcium can also deposit in the cornea

(band-shaped keratopathy), blood vessels (calciphylaxis), and extravascular tissues (calcinosis).

Other options:

Options A and D: Small cell carcinoma and adenocarcinoma of the lung are not associated with hypercalcemia. Squamous cell carcinoma is associated with hypercalcemia due to PTHrP.

Option B: Mycophenolate mofetil or pyridostigmine are not known to cause hypercalcemia.

Solution to Question 40:

In this patient presenting with palpitations, dyspnea bilateral basal crackles and irregular RR intervals with absent P waves on ECG, the most likely diagnosis is acute decompensated heart failure with atrial fibrillation. Beta-blockers are not employed for rate control in patients with acute heart failure.

Beta-blockers and calcium channel blockers are negative ionotropic drugs. They cause decreased cardiac output, decreased blood pressure, and increased left ventricular filling pressure thereby worsening acute heart failure. Beta-blockers are first-line therapy for rate control for Atrial fibrillation in patients without acute heart failure.

Atrial fibrillation (AF) is a supraventricular tachycardia caused by ectopic foci or small reentry circuits within the atria. It has an atrial rate of 400-600/min and a ventricular rate of 100-160/min.

Risk factors for developing AF include:

• Age- usually above 60 years of age

• Underlying heart disease

• Hypertension

• Diabetes mellitus

• Obesity

• Sleep apnea

• Post-cardiac surgery

• Inflammatory pericarditis

• Acute alcohol consumption.

Features include heart rate typically ranging from 110-160 beats/minute. It is irregular and rapid. Presyncope and syncope can be seen. Thrombus formation can occur and thromboembolic events can take place.

Typically, ECG shows an absent P wave and irregular RR intervals.

Management is mainly by controlling the rate by using drugs like β blockers (such as metoprolol) and calcium channel blockers (such as diltiazem and verapamil). Reversion to sinus rhythm can be done electrically or using medications. Pharmacological cardioversion can be done using ibutilide, dofetilide and amiodarone (class III antiarryhthmics) or flecainide and propafenone (class Ic drugs).

Other options:

Option A: Synchronized cardioversion is the treatment of choice in haemodynamically unstable patients.

Option B: Atrial fibrillation predisposes risk for thrombus formation, therefore appropriate anticoagulation has to be started. Direct-acting oral anticoagulants (DOACs) are preferred over warfarin for anticoagulation. However, warfarin is the preferred drug if the patient has a mechanical prosthetic valve, advanced CKD or moderate-severe mitral stenosis.

Option D: Intravenous digoxin can be used to control the ventricular rate in atrial fibrillation, especially in patients with heart failure or those who are not suitable for beta-blockers or calcium channel blockers. However, digoxin is not always the first-line treatment for rate control.





Solution to Question 41:

This patient with a history of depression treated with sertraline and now amitryptiline most likely has developed serotonin syndrome. Serotonin syndrome is treated with cyproheptadine (5-HT2 receptor antagonist). Cyproheptadine can effectively treat serotonin syndrome by blocking the effects of excess serotonin.

Serotonin syndrome is characterized by a triad of:

• Autonomic instability - hyperthermia, hypertension, tachycardia, diaphoresis, vomiting, diarrhea.

• Altered mental status - agitation, confusion, delirium.

• Neuromuscular hyperactivity - tremor, hyperreflexia, myoclonus.

Serotonin syndrome rarely occurs with a single serotonergic drug used at therapeutic doses. More commonly, it occurs due to the combined side effects with any drug that increases serum serotonin levels like MAOIs (monoamine oxidase inhibitors), SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), TCAs (tricyclic antidepressants), tramadol, ondansetron, triptans, linezolid, MDMA (amphetamine), dextromethorphan, etc.

Surgery AIIMS 2017

Question 1:

A man met with a road traffic accident, which resulted in paralysis of both his upper and lower limbs. He has not passed urine and tenderness is elicited in the cervical region. What will you advise regarding radiography?

a) The patient should not be shifted and portable x-ray machine should be used after neck stabilization.

b) Instruct the radiographer to take cervical and chest x-ray.

c) Instruct the radiographer to take cervical x-ray AP and lateral view without any cervical support.

d) Order a cervical x-ray and shift the patient from the trolley by himself.

Question 2:

A 42-year-old woman presented with a history of painless breast lump measuring 6.2 X 4.5 cm in the left upper outer quadrant and no axillary nodes. Core cut biopsy showed ductal carcinoma in situ. She underwent tumor resection with adequate margins and postoperative HPE showed DCIS with intermediate-grade necrosis and with 6-mm clearance on margins. What is the next step of treatment?

a) No additional treatment

b) Adjuvant radiotherapy

c) Adjuvant chemotherapy

d) Adjuvant chemoradiotherapy

Question 3:

Which of the following is false regarding imaging in the primary survey of a trauma patient?

a) Hemodynamically unstable patient should not be sent for CT scan

b) All patients should have chest X-ray PA view only

c) X-ray of the pelvis is essential in primary survery

d) X-rays in primary survey must also be done in pregnant women

Question 4:

A patient presents with renal calculus of > 4 cm of the staghorn type. What is the best management?

a) Extracorporeal shockwave lithotripsy

b) Percutaneous nephrolithotomy

c) Ureteroscopic removal

d) Retrorenal surgery

Question 5:

All of the following can be done to transport a polytrauma patient who is conscious except?

a) Strap their thorax, abdomen and lower limbs on a stretcher

b) Place in prone position to prevent aspiration

c) Put patient on lateral position and shift him immediately

d) Roll the patient onto a hard board

Question 6:

Identify the condition.



a) Cholesterolosis

b) Strawberry gallbladder

c) Gallbladder polyps

d) Gallbladder carcinoma

Question 7:

Colour of 22G IV Cannula is?

a) Pink

b) Green

c) Blue

d) Yellow

Question 8:

Fr in size of Foley's catheter refers to:

a) Outer Diameter

b) Inner Diameter

c) Inner Circumference

d) Radius

Question 9:

A school bus full of children met with an accident. As the doctor in charge of triage, which of the following will you attend to first?

a) Airway Obstruction

b) Grade 4 Shock

c) Flail Chest

d) Severe head injury

Question 10:

Which parameter is the best guide to monitor the adequacy of fluid resuscitation in a patient?

a) Central venous pressure

b) Urine output

c) Blood pressure

d) Oxygen saturation

Question 11:

Which of the following is the most common presentation of abdominal desmoid tumor?

a) Abdominal pain

b) Abdominal mass

c) Fever

d) Rectal prolapse

Question 12:

A 10-year-old child came with complaints of pain over the right lower quadrant and lower back pain. On examination, the patient had pain over the right side of the hip on passive extension with a mass in the right lumbar region pointing towards the groin. What is the most probable diagnosis?

a) Pyelonephritis

b) Psoas abscess

c) Appendicitis

d) Torsion of right undescended testis

Question 13:

Which layer is involved in blister formation in a superficial partial thickness burn?

a) Epidermis

b) Subcutaneous tissue

c) Papillary dermis

d) Reticular dermis

Question 14:

Proper functioning of ICD tube is indicated by _____.

a) Continuous air bubbles coming out of the underwater drain

b) Oscillation of water column in the tube

c) Chest X-ray

d) Auscultation

Question 15:

In a preoperative surgical safety checklist, which of the following is not present?

a) Signature of the doctor

b) Consent

c) Functioning pulse oximeter

d) Sterility of instruments

Question 16:

A trauma patient suffers a renal laceration with urine extravasation. He is managed conservatively. However, after a few days, he presents with a urinoma. The next best management for this patient would be?

a) Ureteral stenting

b) Surgical repair and exploration

c) Observation

d) Percutaneous drainage

Question 17:

Which of the following is considered balanced resuscitation?

a) Maintaining airway, breathing and circulation

b) Allow permissible hypotension

c) Maintain ph by ensuring acid-base balance

d) Fluid resuscitation by giving colloids and crystalloids

Question 18:

The flap shown in the picture is which of the following?



a) Serratus anterior flap

b) Pectoral myocutaneous flap

c) TRAM flap

d) VRAM flap

Question 19:

A 25-year-old patient presented with neck swelling which moves with deglutition as shown below. What is your diagnosis?

a) Branchial cyst

b) Cystic hygroma

c) Thyroglossal cyst

d) Cervical lymphadenopathy

Question 20:

A 28-year-old patient was brought to the casualty ward with high fever and pain abdomen. He had a history of bloody diarrhea for several months. His past history was negative for long-term medications and previous surgeries. The abdominal X-ray image obtained is shown below. What is the most probable diagnosis?



a) Toxic megacolon

b) Volvulus

c) Ogilvie syndrome

d) Pneumatosis intestinalis

Question 21:

A USG can detect lymph node metastases in a patient with breast cancer and clinically negative lymph nodes by what percentage?

a) 50 – 60%

b) 10 – 20%

c) 20 – 30%

d) 30 – 50%

Question 22:

A young boy was brought to the casualty with a history of fever and abdominal pain. On examination, he was febrile and his pulse rate was 104 beats/min. The CT scan image is shown below. Which of the following signs is not seen in this disease?

a) McBurney's tenderness

b) Rovsing's sign

c) Psoas sign

d) Ballance sign

Question 23:

A 12-year-old boy presented to the OPD with a scald injury of his arm. The lesion was pink, oozing, and painful to moving air and touch. Which of the following is the most appropriate management?

a) Silver sulphadiazine cream dressing

b) Excision and grafting

c) Paraffin gauze dressing

d) Collagen dressing

Question 24:

A 25-year-old pregnant women came with a history of thyroid swelling. FNAC report was suggestive of papillary carcinoma thyroid and the nodule was “cold” on scan. Which of the following will not be a treatment option for her?

a) Total thyroidectomy

b) Total thyroidectomy plus lymph node dissection

c) Radioiodine ablation of thyroid

d) Right hemithyroidectomy

Question 25:

A 41-year-old woman underwent treatment for breast cancer of 1.2 X 1 cm with positive axillary nodes. She completed radiotherapy and chemotherapy and is now receiving tamoxifen. How will you follow up the patient?

a) Annual mammogram

b) Regular checking of LFT once in 6 months

c) Follow-up with tumour markers once in 6 months

d) Yearly bone scans

Question 26:

A patient underwent laparoscopic cholecystectomy and was discharged after removing the drain tube on the first postoperative day. On day three, the patient came back to the hospital and was febrile. On ultrasound examination, there was a 5 x 5 cm collection in the subhepatic space. Next best step is _____.

a) To give oral antibiotics since the given accumulation is normal

b) To re-explore the wound and insert a T-tube

c) Insert pigtail catheter for drainage

d) ERCP

Question 27:

A 22-year-old boy was admitted to the casualty with a history of breathlessness following massage by his mother. He had a history of hairline ankle fracture and was on a below-knee cast which was removed recently. The most probable diagnosis is

a) Pneumonitis

b) Septicemia

c) Acute MI

d) Pulmonary thromboembolism

Question 28:

A 65-year-old man has the following lesion over the face. He complains of bleeding while itching. The most likely diagnosis is

a) Dermatofibrosarcoma

b) Marjolin's ulcer

c) BCC

d) Naevus

Question 29:

A 54-year-old comes to the casualty with a history of massive hemoptysis. His routine investigation and chest X-ray were normal. Which of the following you will not do to prevent hemoptysis?

a) Bronchial artery embolization

b) Pulmonary artery embolization

c) Procoagulant drugs

d) Surgical removal of lobe

Question 30:

A 30-year-old male met with an accident and came to the casualty with a contusion on the left precordium. His vitals are BP 120/80 mmHg, RR – 16/min. Normal heart sounds were heard, but decreased breath sounds on the left side, and the trachea deviated to the right. Which of the following is the first line of management?

a) Needle thoracocentesis

b) Pericardiocentesis

c) Chest tube insertion

d) Open surgery

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	b
5	c
6	c
7	c
8	a
9	a
10	b
11	b
12	b
13	c
14	b
15	a
16	a
17	b
18	a
19	c

20	a
21	a
22	d
23	d
24	c
25	a
26	c
27	d
28	c
29	c
30	c

Detailed Explanations

Solution to Question 1:

The given clinical situation suggests a cervical spine injury due to trauma. The patient should not be shifted and a portable X-ray machine should be used for radiography after neck stabilization.

Management in a trauma patient with a suspected cervical fracture involves the following steps:

• Spinal immobilization with a rigid cervical collar and a long spine board is the immediate step in the management of blunt/penetrating cervical trauma cases.

• It is followed by a gross assessment of spinal cord function by eliciting extremity movement. Tenderness over the injured vertebrae or the presence of a deformity is indicative of an associated fracture.

• Imaging is done by X-ray or CT. Portable X-ray machines are preferred to avoid further injury while mobilizing the patient. They are useful in visualizing thoracic and lumbar vertebrae but not the cervicothoracic junction. Cervical CT has been made the modality of choice in most emergency departments owing to the high quality of images that can be obtained.

Solution to Question 2:

According to Silverstein Van Nuys prognostic index for DCIS, the total score is 9. Hence, the patient would require adjuvant radiotherapy.

Age: 42 years (score 2), size: 62mm (score 3), clearance margin: 6mm (score 2) with intermediate-grade necrotic features (score 2).

Silverstein Van Nuys prognostic index for DCIS:

Treatment:

• Scores 4-6 with >3mm margin: Excision alone

• Score 7 with < 3 mm margin, score 8 with > 3mm margin, score 9 with >5mm margin: Excision with RT

• Score 9 with < 5mm margin & score 10-12: Mastectomy

If the patient is ER/PR+, adjuvant hormonal therapy with tamoxifen or an aromatase inhibitor is indicated. Current data does not support the use of HER-2-directed therapy in DCIS.

Indications for adjuvant radiotherapy in breast cancer:

• Locally advanced breast cancers (T3, T4, N1, N2, N3)

• Following breast conservation surgery

• After mastectomy if tumor size ≥5 cm, skin or chest wall involvement, lymphovascular invasion and axillary node-positive for metastasis.

Post-treatment surveillance after treatment for DCIS includes history, physical examination, and routine mammography (if applicable).

Parameters	Score 1	Score 2	Score 3
Age (years)	≥61	40-60	<40
Size (mm)	≤15 mm	16-40 mm	>40 mm
Margin (mm)	≥10 mm	1-9 mm	<1 mm
Class (grade and necrosis)	Grades 1 and 2no necrosis	Grades 1 or 2with necrosis	Grade 3 with or without necrosis

Solution to Question 3:

Chest X-ray and X-ray pelvis are taken in AP view in the adjuncts to the primary survey of a trauma patient.

Solution to Question 4:

The best management for staghorn renal calculus of >4 cm is percutaneous nephrolithotomy (PCNL).

The initial management of renal stones depends on the stone size and includes conservative management, extracorporeal shockwave lithotripsy (ESWL), and PCNL. PCNL is preferred for stones > 2cm in size and for staghorn calculi, while ESWL is used to treat renal stones of size 0.6 to 1.5 cm. PCNL is indicated in

• Renal stones >2 cm

• Staghorn calculus

• Failed shockwave lithotripsy (SWL) or retrograde intrarenal surgery (RIRS) for renal calculi

• Lower pole renal stones with anatomy that is unfavorable for ESWL

Other options:

Option C: Ureteroscopic removal is attempted for failed SWLs or PCNLs, and is not the first choice in the surgical management of renal stones.

Option D: RIRS is a minimally invasive procedure similar to ureteroscopic removal, where the kidney is accessed using a slim and flexible ureteroscope in a retrograde manner via the ureteric orifice. It is indicated for lower pole renal stones < 2 cm.

Solution to Question 5:

Patients with polytrauma are shifted into a supine or prone position (not in a lateral position) by laying them over a firm surface to restrict spinal movement.

As per ATLS guidelines, in patients with trauma, airway management is first done along with cervical spine stabilization. All trauma patients until proven otherwise are considered to have an injured vertebral column.

The patients are placed supine on a firm surface without rotating or bending the spinal column and a rigid cervical collar (philadelphia collar) is applied. Aspiration can be prevented by putting the patient in the prone position and strapping can be done to provide further stabilization.

Long spine boards with straps are helpful in the rapid transport of the patient from one place to another. The patient can be put on the long spine board by log-rolling (a technique where four people hold the patient and roll the patient like a log without allowing rotation or bending of the spine).

Conscious and alert patient indirectly means that they don't need intubation and their airway is patent. If the patient is unconscious always check the airway and if needed, endotracheal intubation must be done immediately by various methods. Preferably by direct laryngoscopy and intubation.

Solution to Question 6:

The given image shows a resected specimen of a gallbladder with a polyp on the inner surface.

Specimen of gallbladder carcinoma would include an en-bloc dissection of the gall bladder, surrounding liver tissue and regional lymph nodes (option D). In specimens of cholesterolosis, also known as strawberry gallbladder, the gall bladder would appear yellow and coarsely granular due to the presence lipid nodules (options A and B).

Gall bladder polyps are usually incidental findings during abdominal imaging. Most of them are benign and comprise cholesterol polyps, adenomyomas or inflammatory polyps. Adenomatous polyps are the most common neoplastic polyps. The majority of patients with polyps are stable and smaller polyps are kept under observation. Cholecystectomy is done for symptomatic patients (biliary colic, pancreatitis), patients >50 years, sessile polyps with thickening >4 mm and polyps >10 mm.

The risk factors for malignancy in an adenomatous polyp include age >50 years, size >10 mm, sessile polyp, documented increase in size, and coexistence with gallstones.

Other options:

Option D: The gallbladder in a patient with carcinoma is removed via an en-bloc dissection with surrounding liver tissue along with regional lymph nodes. The given image below shows a cut surface of the resected gallbladder with a tumor at the neck along with liver bed involvement (marked by the red arrow).

Options A and B: Cholesterolosis (strawberry gallbladder) is characterized by the accumulation of lipids (triglycerides, cholesterol precursors and cholesterol esters) in the lamina propria of the gallbladder wall. The nodules are less than 1 mm in diameter and there is mucous villous hyperplasia, which gives it a coarse, yellow and granular appearance, giving it a strawberry-like appearance. The given images below show cholesterolosis.

Solution to Question 7:

The 22G IV cannula is blue in colour.

Solution to Question 8:

The French (Fr) measurement unit refers to the outer diameter.

As per the French system, the outer circumference equals 3 times the outer diameter. One French unit (1 Fr) equals 1 mm of the outer circumference and 0.3 mm of the outer diameter. Therefore, the outer diameter in mm can be determined by dividing the Fr size by 3.

Example: An 18 Fr would have a circumference of 18 mm and an outer diameter of 6 mm.

Solution to Question 9:

The patient with airway obstruction, categorized as red, should be attended to first.

Triage is used to classify the priority of treatment. It is carried out at the site of the disaster to determine the priority for immediate medical attention, transportation, and admission to the hospital. Triage decisions are made by deciding which injuries constitute the greatest immediate threat to life. In general, airway problems are more rapidly lethal than breathing problems > circulation problems > neurologic injuries.

The internationally accepted four-color coded system is commonly used. It includes:

• Red - High-priority attention or transfer

• Yellow - Medium priority

• Green - Ambulatory patients

• Black - Dead or moribund patients

Solution to Question 10:

Urine output is the best measure of organ perfusion and the best monitor for adequacy of fluid resuscitation.

Adequate end-organ perfusion is best indicated by urine output of >0.5 to 1 mL/kg/h. However, it does not give a minute-to-minute value.

The level of consciousness is an important marker for cerebral perfusion.

The only clinical indicators of perfusion of the gastrointestinal tract and muscular beds are measures of lactic acidosis (lactate and base deficit) and mixed venous oxygen saturation.

Other options:

Option A: Central venous pressure (CVP) is the mean pressure in the superior vena cava, used for estimation of preload and right atrial pressure. Normal CVP ranges from 2 to 7 mm Hg (3 to 9 cm H2O). A sick or injured patient with a CVP <3 mm Hg is presumed to be volume depleted and may be given fluids with relative safety. However, CVP is unreliable in assessing volume status or left ventricular function with respect to resusitation.

Options C and D: Blood pressure and oxygen saturation do not reflect body fluid levels.

Solution to Question 11:

Explanation:

The most common presentation of an abdominal desmoid tumor is an abdominal mass.

Desmoid tumors, also known as fibromatosis, aggressive fibromatosis or desmoid-type fibromatosis are locally aggressive tumors that lack metastatic potential. Desmoid tumors can be sporadic in origin or associated with familial adenomatous polyposis.

These tumors arise from fibro-aponeurotic tissue and typically are manifested as a slowly growing mass. They present as firm, non-painful, non-tender masses in the abdominal wall, shoulder, hip and limbs, mesentery, and pelvis. They are also associated with trauma, pregnancy, and oral contraceptive usage.

The investigation of choice for a desmoid tumor is MRI. It is superior to CT in defining the pattern and the extent of involvement and can determine recurrence after surgery. Diagnosis is established by a biopsy with immunohistochemistry, which stains positive for beta-catenin, actin and vimentin, and negative for desmin, cytokeratin and S-100.

Treatment:

• An initial observation is preferred for stable, asymptomatic, primary or recurrent desmoids, especially if surgery can cause high mortality.

• Surgical resection with negative margin clearance is reserved for patients with small abdominal wall tumors associated with pain or an increase in size.

• Tamoxifen and NSAIDs are used as first-line medical therapy. Other systemic treatment options are low-dose chemotherapy, tyrosinase kinase inhibitors, and full-dose chemotherapy

• Radiotherapy is preferred for candidates unfit for surgery, patients refusing surgery and when surgical resection is associated with high morbidity.

Solution to Question 12:

The given clinical history of lower back pain, mass in the right lumbar region pointing towards the groin and pain on passive extension of the ipsilateral hip is suggestive of a psoas abscess.

Psoas abscess is a retroperitoneal abscess, caused due to various etiologies. Previously, tuberculosis of the spine was the most common cause secondary to spreading infection from the digestive or urinary tract, which can cause psoas abscess. Nowadays, Crohn's disease is the most common cause. Patients present with flank pain, fever and chills, malaise and weight loss. They can have referred pain to the hip, groin or knee. When the swelling extends across the groin and below the inguinal ligament along the course of the iliopsoas, it becomes cross fluctuant across the inguinal ligament.

Imaging is done by a CT scan, which shows a low-density mass with surrounding inflammation and infiltration of surrounding fat and gas or air-fluid levels.



Treatment is done by percutaneous CT-guided drainage and appropriate antibiotics.

Other options:

Option A: Pyelonephritis is usually associated with a history of urinary tract infection and sudden onset of pain in the flank and hypochondrium, not the lumbar region.

Option C: Appendicitis (retrocaecal) can present with pain on passive extension of the hip (psoas sign), but the presence of a mass in the right lumbar region pointing towards the groin makes it an unlikely diagnosis.

Option D: Testicular torsion is a surgical emergency condition presenting with sudden onset pain in the scrotum, groin and lower abdomen, tenderness, redness and edema of the scrotal skin.

Solution to Question 13:

The papillary dermis is involved in blister formation in a superficial partial-thickness burn.

First-degree burns involve only the epidermis, with pain, redness, and mild swelling.

Second-degree (partial thickness) burns involve the epidermis and some parts of the dermis (also known as partial-thickness burns). Further divided into:

• Superficial second-degree burns - involves the upper part of the dermis (papillary) and shows pain, blisters, and severe swelling.

• Deep second-degree burns - extends up to the reticular layer of the dermis and appears more pale and mottled, does not blanch to touch, but remains painful to pinprick.

Third-degree or full-thickness burns involve the full thickness of the dermis and subcutaneous tissue.

Fourth-degree burns involve deep structures such as muscle and bone.

Solution to Question 14:

Oscillation of the water column level (moving up and down during respiration) within the tube in the water seal chamber is indicative of a functioning (patent) ICD tube.

During spontaneous respiration, the water level rises with inspiration and falls with expiration. Whereas during positive pressure ventilation, the opposite occurs. This oscillation is also known as tidaling.

Other options:

Option A: Continous bubbling indicates an air leak in the drainage system, which should be corrected

Option B: Chest X-ray is useful for assessing the position of the ICD tube inside the chest, not its functioning.

Option C: Auscultation is not useful for assessing the functioning of an ICD tube.

Solution to Question 15:

In a preoperative patient surgical safety checklist, doctor's signature is not required.

WHO Guidelines for Surgical Safety Checklist



FIGURE 10-4 Surgical safety checklist published by the World Health Organization. www.who.int/patientsafety/safesurgery/checklist/en.

Solution to Question 16:

The patient with a history of grade IV renal injury (renal laceration with urine extravasation) who develops a persistent urinoma (a mass formed by encapsulated extravasated urine) should be managed with the placement of an internal ureteral stent.

In patients not cured by ureteral stenting, percutaneous nephrostomy or transcutaneous drain may be required.

Management of renal injuries

Blunt trauma is the main cause of renal injuries and is self-limiting. Blunt trauma is mostly treated conservatively. Most penetrating injuries require surgical renal exploration.

• Grading of injury through CECT

• If hemodynamically stable

• Conservative management - Bedrest & hemodynamic monitoring

• In grade IV, a repeat CT scan to ensure the extravasation of urine has resolved.

• If hemodynamically unstable

• Surgical exploration and repair

• In grade V, nephrectomy.

Solution to Question 17:

Permissive hypotension is a part of balanced resuscitation in trauma management to avoid bleeding.

Damage control resuscitation (DCR) i.e. balanced resuscitation is used in the settings of major trauma. It aims to control hemorrhage and prevent coagulopathy prior to the development of the trauma triad of death (hypothermia, acidosis, and coagulopathy). The components of DCR include:

• Permissive hypotension until definitive surgical control

• Minimize crystalloid based resuscitation

• Early use of blood products (PRBCs, FFP, platelets, cryoprecipitate) in the ratio of 1:1:1

• Use of hemostatic agents such as tranexamic acid, recombinant factor 7a, prothrombin complex concentrates

DCR has the following advantages:

• Minimizes the impact of trauma-induced coagulopathy

• Limits blood product waste

• Reduces the complications that occur with aggressive crystalloid resuscitation.

Solution to Question 18:

The given image shows a serratus anterior flap. The serratus anterior flap is a highly reliable flap with a consistently long pedicle. Its anatomy permits coverage of complex wounds. The underlying rib can be included as a myoosseous flap to expand the versatility of this flap.

Classification of muscle flaps		
Based on blood supply	Random flapUnnamed vessel supplying flap	Axial flapNamed vessel supplying flap.
Method of movement	Local transferRotation flapTransposition flapAdvancement flap	Distant transferPedicle flapFree flap
Based on composition	Simple FlapCutaneous flapMusculocutaneousFasciocutaneous	Composite flapOsseocutaneous flapSensory flap

Solution to Question 19:

A thyroglossal cyst is the only midline neck swelling that moves with deglutition. This is because the thyroid gland is fixed to the hyoid.

Thyroglossal duct cyst:

It is a congenital midline neck swelling that typically presents during childhood often noticed after an upper respiratory infection. It represents the persistence of the track of the thyroid gland and can be found adjacent to the midline. The thyroid gland descends early in fetal life from the base of the tongue towards its position in the lower neck. At the time of descent, the hyoid bone has not been formed and the track of the thyroid gland is variable, passing in front, through, or behind the eventual position of the hyoid body.



Clinical features of thyroglossal cyst:

• Classically the cyst moves upwards on swallowing and on tongue protrusion. This indicates attachment to the hyoid bone.

• Swelling is smooth, soft, fluctuant (cystic), nontender, mobile, often transilluminant.

• Thyroid fossa is empty, if there is no thyroid in the normal location.

Complications of thyroglossal cyst:

• The most common complication is infection.

• Thyroglossal fistula is an acquired condition that either follows infection of the cyst which bursts open or due to inadequate removal of the cyst.

• 1% of thyroglossal duct cysts are found to be malignant of which the most common type of cancer is papillary (85%).

Neck swellings that move with deglutition are:

• Thyroid swellings

• Subhyoid bursitis

• Dermoid cysts attached to the hyoid bone

However, only the thyroglossal cyst moves with tongue protrusion. This is because the duct extends from the foramen caecum of the tongue to the isthmus of the gland.

Sistrunk's operation is performed to excise the cyst and prevent a recurrence. It involves the excision of the whole thyroglossal tract, involving removal of the body of the hyoid bone, and suprahyoid tract through the tongue base to the vallecula at the site of the primitive foramen caecum along with the core of tissue on either side.

Solution to Question 20:

The clinical scenario and the radiograph are suggestive of toxic megacolon.

Toxic megacolon may present with fever, abdominal pain, and abdominal distension. Abdominal radiography shows transverse or right colonic dilation of >6 cm and loss of haustrations. It is usually associated with ulcerative colitis as indicated by history of chronic bloody diarrhoea.

It can be triggered by electrolyte imbalance and narcotics which interfere with gut motility. Features of secondary bacterial infecton and sepsis may be present. It can be complicated by intestinal perforation.

It usually resolves with supportive treatment (correcting electrolyte imbalance, fluids, and bowel rest) and medical therapy which includes glucocorticoids, anti-inflammatory drugs, and antibiotics. However, an urgent colectomy is indicated in patients who do not show signs of improvement.

Indications for urgent surgery include:

• Frank intraperitoneal peritonitis, secondary to perforation

• Life-threatening hemorrhage or increasing transfusion requirements

• Worsening systemic toxicity

• Worsening colonic dilatation

Other options:

Option B: Volvulus occurs when the air-filled segment of the colon twists around the mesentery. The sigmoid colon is the most commonly involved part of the large intestine (90%). The patient

can present with abdominal distension, nausea and vomiting. X-ray abdomen shows a bent inner tube or coffee bean appearance.



Option C: Ogilvie syndrome, also known as colonic pseudo-obstruction is a functional disorder with colonic dilatation without mechanical obstruction. It occurs commonly in hospitalized patients and is associated with narcotics, bed rest and comorbid disease. X-ray shows dilatation of the colon (predominantly transverse colon) with absent mechanical obstruction.

Option D: Pneumatosis intestinalis is the presence of gas within the bowel wall seen in any part of the GI tract but most commonly in the jejunum. It is a pathognomic feature of necrotizing enterocolitis. X-ray of the abdomen will show intramural gas in the bowel.



Solution to Question 21:

A USG can detect lymph node metastasis in a patient with clinically negative lymph nodes by 50-60%.

The clinical determination of axillary lymph node metastases has an accuracy of only 33%.

Ultrasound (USG) is more sensitive than physical examination alone in determining axillary lymph node involvement during the preliminary staging of breast carcinoma.

Fine-needle aspiration cytology (FNAC) or core biopsy of sonographically indeterminate or suspicious lymph nodes can provide a more definitive diagnosis than USG alone.

Ultrasonography can also be utilized to image the regional lymph nodes in patients with breast cancer. The sensitivity for the status of axillary nodes ranges from 35% to 82% and specificity ranges from 73% to 97%.

The features of a lymph node involved with cancer on USG include:

• Cortical thickening

• Change in shape of the node to a more circular appearance

• Size larger than 10 mm

• Absence of a fatty hilum

• Hypoechoic internal echoes

Solution to Question 22:

The patient's history and radiological findings are indicative of acute appendicitis. Ballance sign is not seen in acute appendicitis.

Findings on CT that are suggestive of acute appendicitis include:

• Appendiceal dilatation (>6 mm in diameter)

• Appendiceal wall thickening (>3 mm) and enhancement

Ballance sign is seen due to a ruptured spleen. The ballance sign is positive if there is a fixed dullness to percussion in the left flank and shifting dullness to percussion in the right flank. The fixed dullness is due to the presence of coagulated blood near the lacerated spleen while the shifting dullness is due to the presence of blood in the peritoneal cavity.

Option A: Deep tenderness at McBurney's point, known as McBurney's sign, is a sign of acute appendicitis. McBurney's point is a point at one-third of the distance from the anterior superior iliac spine and umbilicus.

Option B: Rovsing sign is seen in appendicitis. Deep palpation of the left iliac fossa may cause pain in the right iliac fossa.

Option C: Psoas sign is seen in appendicitis. Occasionally, an in■amed appendix lies on the psoas muscle, and the patient will lie with the right hip ■exed, for pain relief.

Solution to Question 23:

Scald injury (damage from contact with hot liquids) that appears pink in colour, has oozing, and pain is suggestive of a 2nd-degree burn (i.e. partial-thickness burn). Collagen dressing is the most appropriate management for this injury.

The best dressing for burn wounds is skin; collagen is a good substitute for skin, especially in partial-thickness burns. For full-thickness burns, sequential excision and grafting should be considered.

Initial treatment for burns includes cleaning burnt areas and removing loose dead skin. Blisters are de-roofed and an appropriate dressing is applied.

Biological dressing (collagen dressings, amniotic membrane, etc.) is the preferred dressing material, especially in partial thickness burns.

If this is not available then a non-adhesive dressing like paraffin gauze impregnated with silver sulfadiazine (SSD) cream, colloidal silver, ionic silver cream, or silver nitrate cream can be applied. Since increased resistance to SSD has recently emerged, it should be judiciously used and reserved for deep dermal, full thickness, infected, and old neglected burns.



Solution to Question 24:

Radioactive ablation of the thyroid is an absolute contraindication in pregnancy as it can cause fetal goiter and hypothyroidism.

Radioactive ablation of the thyroid is done using the radioactive isotope of iodine (I-131), which has a half-life of 8 days. I-131 emits gamma rays (useful in imaging) and beta particles (useful in therapy). I-131 is concentrated within the thyroid follicles and emits radiation from within the tissue. The beta particles penetrate only 0.5-2 mm of the tissue. The thyroid follicular cells are affected from within and undergo pyknosis and necrosis followed by fibrosis without damage to neighboring tissues.

Radioactive iodine is administered as the sodium salt of I-131 dissolved in water and taken orally.

Therapeutic uses of radioactive iodine:

• Distant metastasis at diagnosis (papillary or follicular carcinoma)

• Incomplete tumor resection

• Patients at high risk for mortality or recurrence

• Graves disease

• Toxic multinodular goitre

The average therapeutic dose is 3-6 m curie, calculated based on previous tracer studies and thyroid size.

Contraindications:

• Pregnancy

• Lactation

• Young age (<25 years)

• Moderate to severe Graves ophthalmopathy

Management of differentiated thyroid cancer in pregnancy:

• Thyroidectomy after delivery for patients with no evidence of advanced disease or no rapid progression and in the second trimester for patients with rapid progression or signs of advanced disease.

• Radioactive iodine is contraindicated in pregnancy and lactation. If it is needed after delivery, breastfeeding shall be stopped at least 6 weeks before ablative therapy and mothers should be advised against nursing.

• Levothyroxine supplementation: as a suppressive treatment for those who decided to postpone surgery till the second trimester or after delivery, as a replacement therapy after thyroidectomy, and as a suppressive therapy for patients with residual disease.

• Differentiated thyroid cancer not responding to radioactive iodine can be treated with tyrosine kinase inhibitors.

Diagnostic radioiodine scan: 25–100 μ curie of I-131 is given; counting or scanning is done at intervals.

Solution to Question 25:

Follow-up of patients after treatment of breast cancer consists of a physical examination once in 3-6 months for the first 3 years and an annual mammogram.

Mammography is the only imaging modality necessary for follow-up after breast cancer surgery. There is currently no role for tumor markers, serum investigations, or any other imaging studies like bone scans in routine follow-ups.

Guidelines for follow-up after primary treatment:

• Physical examination: every 3-6 months for the first 3 years, every 6-12 months for years 4 and 5, and annually thereafter.

• Mammography: initially, 6 months after surgery, and thereafter, annually.

• Breast self-examination: Monthly

• Pelvic examination: Annually in patients taking tamoxifen

Solution to Question 26:

The given clinical situation suggests a bile duct injury identified on post-op day 3 following laparoscopic cholecystectomy. The next best step after imaging would be to control inflammation by drainage of fluid collection (pigtail insertion and drainage) and administration of parenteral broad-spectrum antibiotics.

Bile duct injury

• Usually is manifested in the postoperative period

• Clinical features include fever, increasing abdominal pain, jaundice and bile leakage from an incision.

Imaging: Ultrasonography is done to assess for fluid collection and to evaluate the biliary tree.

Treatment Strategy

• Control of infection, limiting inflammation

• Percutaneous drainage of periportal fluid collections

• Parenteral antibiotics (option A)

• Clear and thorough delineation of entire biliary anatomy

• Magnetic resonance cholangiopancreatography (MRCP) or percutaneous transhepatic cholangiography (PTC)

• Endoscopic retrograde cholangiopancreatography (ERCP), especially if a cystic duct stump leak is suspected

• Re-establishment of biliary-enteric continuity (surgery)

• Tension-free, mucosa-to-mucosa anastomosis

• Roux-en-Y hepaticojejunostomy

Other options:

Option B: Re-exploration and repair in indicated for biliary leaks identified in within 48 hours post cholecystectomy.

Option D: ERCP would be indicated if drainage using a pigtail catheter fails.



Solution to Question 27:

The given clinical history of immobilization after the application of the cast is a risk factor for deep vein thrombosis (DVT). Massaging the leg may dislodge the thrombus causing pulmonary

embolism (PE). It is the commonest complication of DVT leading to death.

PE and DVT are part of the same spectrum of venous thromboembolism. PE occurs due to obstruction of the pulmonary arterial circulation by deep venous thrombi that detach from their site of formation and embolize to the vena cava, right atrium, and right ventricle. Virchow's triad of venous stasis, hypercoagulability, and endothelial injury, leading to recruitment of activated platelets, provides the ideal environment for thrombus formation.

DVT presents with pain and swelling of the calf (the most common symptom), tenderness and edema. Clinical examination is unreliable and the elicitation of signs such as Homan's or Mose's signs is not recommended owing to the risk of clot dislodgement. Patients with symptoms of PE present with fever, dyspnea, pleuritic chest pain, and hemoptysis. On examination, they can have cyanosis, jugular venous distension, fixed split-second heart sound and pleural rub, but may be subtle or absent. Some patients are asymptomatic.

Depending on the clinical presentation, patients with PE can be stratified as:

• High-risk (massive) PE is characterized by systemic hypotension. Patients can have dyspnea and syncope and may present in cardiogenic shock.

• Intermediate-risk PE is characterized by right ventricle (RV) dysfunction but normal blood pressure. RV dysfunction can be diagnosed with echocardiography and ECG.

• Low-risk PE is characterized by normal blood pressure and the absence of any RV dysfunction.

Investigations:

• Venous duplex ultrasound to look for thrombi in the leg.

• Chest x-ray in PE shows hyper lucency in an area of oligemia (Westermark sign). D-dimer levels are elevated and ECG will show sinus tachycardia (the most common ECG abnormality) and an S1Q3T3 sign (an S wave in lead I, a Q wave in lead III, and an inverted T wave in lead III).

• Definitive diagnosis of PE is by a CT-Pulmonary Angiography.



The management of PE will vary according to the risk stratification:

• A patient with high-risk PE will primarily be treated with anticoagulation and thrombolysis or catheter/surgical embolectomy.

• A patient with low-risk PE will be treated with anticoagulation alone, or an IVC filter as secondary prophylaxis.

• Patients with intermediate-risk PE require individualized treatment.

Anticoagulation can be done with the classical strategy of low-molecular-weight heparin followed by warfarin, or by low-molecular-weight heparin for 5 days, switched to novel oral anticoagulant like Dabigatran after that. Fibrinolysis, if indicated, is done with recombinant TPA administered over 2 hours. It can be done up to 14 days after the PE has occurred.

Management of DVT:

• Anticoagulation with low molecular weight heparin (LMWH) and unfractionated heparin (UFH) if renal impairment is present.

• After treatment with heparin, patients should be commenced on oral anticoagulants for at least 3 months with either factor Xa inhibitors (rivaroxaban and apixaban)
or direct thrombin inhibitors (dabigatran).

• Other measures such as rest, limb elevation and compression stockings are also used.

Solution to Question 28:

The given history with the image showing an ulcer over the face with pearly, rolled-out borders points to a diagnosis of basal cell carcinoma.

Basal cell carcinoma (BCC) is an epithelial tumor arising from basal cells. It is the most common type of skin cancer most commonly seen over the face, above the line joining corner of the mouth, and lobule of the ear. Risk factors for BCC include ultraviolet radiation (strongest risk factor), age 40-80 years with excessive sun exposure, light-skinned people, and exposure to arsenic compounds, coal tar, aromatic hydrocarbons, and infrared light. It is more common in males than females. It is also associated with syndromes such as xeroderma pigmentosum and Gorlin's syndrome.

Features of BCC include:

• Pearly nodule with rolled border

• Tortuous telangiectatic vessels

• Surface which can get ulcerated with raised and beaded edge, often with a central depression or umbilication

• Bleeding on minor trauma

BCC has various subtypes such as nodular (most common type), pigmented, nodulocystic, morphoeic, infiltrative, and fibroepithelioma of Pinkus.

It is locally aggressive and can erode the bone or facial sinuses. It rarely metastasizes via lymph nodes.

The occasional presence of melanin in the nodular variant of BCC (pigmented BCC) may lead to confusion with melanoma.

Depending upon the type of the lesion, a shave, punch or excisional biopsy may be employed for diagnosis. X-ray or CT may be needed if there is a local invasion. Histology shows peripheral palisading of cells.

Treatment:

• Surgical: Wide excision (4-6 mm clearance) with a skin graft or flap or primary suturing

• Mohs' surgery for BCC close to the eyes, nose, or ear to preserve more tissue.

• Non-surgical: Radiotherapy, vismodegib and sonedegib.

Other options:

Option A: Dermatofibrosarcoma, also known as dermatofibrosarcoma protuberans (DFSP) is a rare, slow-growing, well-differentiated primary fibrosarcoma of the skin, associated with increased production of platelet-derived growth factor-beta. It presents as a protuberant nodule within a firm, indurated plaque that can ulcerate. They are locally aggressive tumors with high recurrence rates. Treatment is wide excision and Mohs microscopic surgery. Imatinib is used for unresectable, metastatic, and recurrent tumors. The given image below shows a DFSP near the umbilicus.

Option B: Marjolin's ulcer is a painless, slow-growing, locally malignant, well-differentiated squamous cell carcinoma that occurs in a chronic scar (burns scar, scar of venous ulcer). It presents as an indurated, non-tender ulcer with raised and everted edges with scarring in the surrounding area with marked fibrosis. The ulcer can spread to the lymph nodes when it reaches the normal skin. A wedge biopsy and MRI of the region involved confirm the diagnosis. Management is by wide excision, for larger ulcers, amputation is done. The given image below shows a marjolin's ulcer of the leg.

Option D: Naevus is a common benign neoplasm caused due to mutations in the components of the RAS signaling pathway. It has various types, such as congenital naevus, blue naevus, Spitz naevus, halo naevus, and dysplastic naevus. The given image below shows a patient with multiple dysplastic nevi.

Solution to Question 29:

Procoagulant drugs are not used to prevent hemoptysis. Procoagulant drugs like vasopressin are indicated only if there is an abnormal coagulation profile.

Massive hemoptysis is generally defined as the expectoration of over 600 mL of blood within a 24-h period.

Treatment priorities in the management of massive hemoptysis:

• First priorities are to secure the airway, optimize oxygenation and stabilize the hemodynamic status by administering IV fluids.

• Chest X-ray is the first investigation of choice. CT scan is done only in hemodynamically stable patients.

• If the bleeding site is known, the patient should be placed in a lateral decubitus position with the bleeding side down to prevent aspiration into the unaffected lung. Supplemental oxygen should be given.

• If large-volume bleeding continues or the airway is compromised, the patient should be intubated and undergo an emergency bronchoscopy.

• The bleeding site is localized using flexible/rigid bronchoscopy.

• Control the hemorrhage:

• Endobronchially with Nd: YAG laser or bronchial occlusion

• Endovascularly with bronchial and/or pulmonary artery embolization

• Electrocauterization

• Surgically with resection of the involved lobe

The cause of the hemoptysis must be determined to definitively prevent a recurrence.

Solution to Question 30:

The given clinical scenario of trauma, decreased breath sounds, and tracheal deviation with stable vitals are suggestive of a simple pneumothorax. Management involves inserting a chest tube.

Traumatic pneumothorax can be:

• Simple pneumothorax - with hemodynamic stability

• Tension pneumothorax - without hemodynamic stability

Initial management of simple pneumothorax with open sucking wound involves prompt closing of the defect with a sterile occlusive plastic dressing (e.g., Opsite), taped on three sides to act as a flutter-type valve followed by chest tube insertion as soon as possible in a site remote from the injury site.

Other options:

Option A: Features of tension pneumothorax include hypotension, dyspnea, distended neck veins, tracheal deviation, absent or decreased breath sounds, hyper-resonant note on percussion and normal heart sounds. The first line of management is needle thoracocentesis in the 5th intercostal space in the midaxillary line.

Definitive management - insertion of an intercostal tube in the triangle of safety. If a sucking wound is present, then it is covered with gauze pieces and taped on 3 sides.

Option B and D: Pericardial tamponade resembles tension pneumothorax but there is no tracheal deviation, normal breath sounds, normal percussion note and muffled heart sounds. Initial management is by needle pericardiocentesis followed by definitive sternotomy or lateral thoracotomy.

REMEMBER RECENT UPDATE :(Acc to ATLS 10th edition):

•

• In adults, the needle thoracotomy location changed to the 5th ICS mid-axillary line from the 2nd ICS midclavicular line.

• In the pediatric age group, the thoracotomy location remains at the 2nd ICS mid-clavicular line.

Surgery AIIMS 2018

Question 1:

All are true about intestinal transplantation except _____.

a) Most of the grafts are multi-visceral grafts.

b) It is not widely used due to vigorous rejection reactions.

c) The most dangerous complication is sepsis.

d) The most common indication is the ischemia

Question 2:

All of the following are boundaries of the triangle of doom except?

a) Vas deferens

b) Peritoneal Reflection

c) Iliopubic tract

d) Spermatic vessels

Question 3:

All of the following are true about superior mesenteric artery syndrome except?

a) Third part of duodenum is compressed

b) Angle between SMA and aorta is more than 70 degrees

c) Surgical management is by dudenojejunostomy

d) Also known as Wilkie's Syndrome

Question 4:

Which of the following type of renal calculi is resistant to the extracorporeal shockwave lithotripsy?

a) Struvite

b) Cystine

c) Calcium oxalate

d) Uric acid

Question 5:

An abdominal aortic aneurysm is operated when its size is greater than _____

a) 5 cm

b) 6 cm

c) 5.5 cm

d) 6.5 cm

Question 6:

A 3-day-old neonate presented with recurrent bilious vomiting. An X-ray erect abdomen is given below. What is the next best step in managing this condition?

a) CT abdomen

b) Gastrograffin follow-through

c) USG abdomen

d) Explorative laparotomy

Question 7:

The image shows methylene blue dye being injected in the peritumoral region. What is this procedure useful for?



a) Localizing the tumor

b) Identifying the margins

c) Sentinel lymph node biopsy

d) Identifying metastasis

Question 8:

A 3-month-old child presented with umbilical discharge. Abdomen examination was normal except for the below finding. The next step of management is?

a) MRI Abdomen

b) USG Abdomen

c) CECT

d) Exploratory laparotomy

Question 9:

Which of the following cancers rarely metastasizes to the liver?

a) Colon

b) Breast

c) Prostate

d) Melanoma

Question 10:

A 40-year-old female presents to the ER with a history of RTA. On examination, GCS is 15/15, heart rate = 110/min, BP = 90/60 mmHg, and there is elevated JVP. The right chest is hyperresonant on percussion with decreased breath sounds. The left chest appears normal. But the patient is dyspneic and ecchymosis is noted over the chest wall. The abdomen was soft. The probable diagnosis is:

a) Cardiac tamponade

b) Tension pneumothorax

c) Pericardial effusion

d) Massive hemothorax

Question 11:

A 50-year-old male presents to the emergency with a BP of 160/100 mmHg and a heart rate of 120. CECT done reveals a Stanford A aortic dissection. What is the initial management of this given condition?

a) Beta blocker medication

b) LMWH

c) Surgical repair

d) Vitamin K inhibitors

Question 12:

A 22-year-old male presented with a hard painless testicular swelling. There were no inguinal, intra-abdominal or mediastinal lymph nodes on CT Scan. The blood investigations showed AFP levels of > 3000. USG shows uniform echotexture and small areas of necrosis. The next line of management is _____.

a) High inguinal orchidectomy

b) Core needle trucut biopsy

c) FNAC

d) Whole body PET/CT

Question 13:

A condition requires the surgery depicted in the image shown below. All the following are usually seen associated with this condition except?



a) Absent kidney

b) Hepatic artery anomalies

c) Malrotation of gut

d) Polysplenia

Question 14:

You are doing an emergency laparotomy for a case of intestinal obstruction. Which organ will you first visualize to know if this is a case of small bowel or large bowel obstruction?

a) Ileum

b) Caecum

c) Jejunum

d) Sigmoid colon

Question 15:

What is the best position for a conscious patient to be in during the insertion of a Ryle's tube?

a) Supine with neck flexion

b) Supine with neck extension

c) Sitting with neck flexion

d) Sitting with neck extension

Question 16:

Pick from the following the usual cause of thyroid storm in thyroidectomy:

a) Rough handling of thyroid during surgery

b) Due to parathyroid gland

c) Inadequate patient preparation

d) Infections

Question 17:

Which of the following colored cannula will you use to achieve maximum flow rate, in a patient who needs a large amount of fluids?

a) Grey

b) Green

c) Blue

d) Pink

Question 18:

A patient who was admitted after a road traffic accident is put on mechanical ventilation. He opens his eyes on verbal command and moves all four limbs spontaneously. Calculate his GCS.

a) Eyes-2, Verbal-1, Motor-5

b) Eyes-2, Verbal-NT, Motor-5

c) Eyes-3, Verbal-1, Motor-6

d) Eyes-3, Verbal-NT, Motor-6

Question 19:

A patient was brought to the casualty after a road traffic accident (RTA) with chest trauma. He is conscious and speaks single words. He has a respiratory rate of 44 cycles per min and a BP of 90/40 mmHg. There is a hyper-resonance on percussing the right side of his chest. What will be your next, immediate step?

a) Intubate the patient

b) Insert needle in appropriate intercostal space

c) Start intravenous fluids urgently

d) Take a chest X-ray

Question 20:

Heller's operation is done for _____.

a) Achalasia cardia

b) Carcinoma esophagus

c) Esophageal stricture

d) Hypertrophic pyloric stenosis

Answer Key

Question No.	Correct Option
1	a
2	c
3	b

4	b
5	c
6	b
7	c
8	b
9	c
10	b
11	a
12	a
13	a
14	b
15	c
16	c
17	a
18	d
19	b
20	a

Detailed Explanations

Solution to Question 1:

Multivisceral grafts are rarely used in intestinal transplantation. The most common intestinal graft is small bowel graft, with or without the colon or a combined liver and small bowel graft.

All the other statements are correct.

Small bowel transplantation is the final treatment option after the outcome of reconstruction and lengthening procedures are unsatisfactory.

The small bowel transplant is highly immunogenic since it contains a large amount of gut-associated lymphoid tissue, donor dendritic cells, and macrophages. Acute rejection has been reported in 56–75% of patients, with small bowel transplant limiting the overall survival to only one-third of recipients.

Solution to Question 2:

Iliopubic tract does not form a boundary of the triangle of doom.

The triangle of doom is bounded

• Medially by the vas deferens

• Laterally by the vessels of the spermatic cord

• Inferiorly by peritoneal reflection

Apex oriented superiorly at the internal ring

The contents of the space include

• External iliac vessels

• Deep circumflex iliac vein

• Femoral nerve

• Genital branch of the genitofemoral nerve

Note: Staples are avoided in this region during hernia repair due to the risk of injury to the vessels which might result in severe hemorrhage.



Solution to Question 3:

Option B is the false statement. In superior mesenteric artery (SMA) syndrome, the aortomesenteric angle is reduced to less than 25 degrees. The normal aortomesenteric angle is approximately between 38-65 degrees.

SMA syndrome, also called Wilkie's syndrome (option D), is characterized by an obstruction of the third part of the duodenum by the SMA (option A) due to compression against fixed retroperitoneal structures.

Symptoms include episodic epigastric pain associated with vomiting. Risk factors include rapid childhood growth without compensatory weight gain, profound weight loss, abdominal surgery, inflammatory diseases of the abdomen, and immobilization in a cast that increases lordosis and accentuates the angle of the SMA with the aorta.

Abdominal CT angiography is the investigation of choice. The CT image below shows narrowed aortomesenteric angle in SMA syndrome:



Lifestyle therapy includes frequent small feedings and patients may benefit from lying prone or on their left side postprandially.

In some cases, weight gain or removal of a body cast improves symptoms. Laparoscopic lysis of the ligament of Treitz is beneficial in few cases. The preferred surgical treatment is Strong's procedure involving duodenojejunostomy (option C).



Solution to Question 4:

Cystine stones are resistant to extracorporeal shockwave lithotripsy (ESWL) due to their hardness. Cystine, brushite, and calcium oxalate monohydrate stones are the most refractory to ESWL fragmentation, with cystine and brushite being the most resistant.

ESWL is a non-invasive method that generates shockwaves outside the body that focus on the stones. It is the most common method of treating urinary tract stones nowadays. Stones up to approximately 1.5 cm in size are suitable for this form of treatment.

Complications: Infection, haematuria, parenchymal hemorrhage, and perirenal hematoma.

Contraindications: Obese patients, pregnant patients, and patients taking oral anticoagulants.

Note: Stone fragments collecting in the distal ureter following ESWL are known as steinstrasse.

Solution to Question 5:

An asymptomatic abdominal aortic aneurysm in an otherwise fit patient should be considered for repair if >5.5 cm in diameter (measured by ultrasonography). The annual incidence of rupture rises from 1% or less in aneurysms that are <5.5 cm in diameter to a significant level of nearly 25%, in those that are 7 cm in diameter.

As open elective surgery (transabdominal) carries a 5% mortality rate, the balance is in favor of elective operation once the maximum diameter is >5.5 cm.

6 monthly follow-up ultrasonographic assessment is indicated for asymptomatic aneurysms <5.5cm in diameter.

Solution to Question 6:

Explanation:

The given clinical history with an x-ray showing a triple bubble sign is suggestive of jejunal atresia. The next best step is to perform a contrast study to localize the site of obstruction.

Small bowel atresias present with intestinal obstruction soon after birth.

• Bilious vomiting is the dominant feature in jejunal atresia

• Abdominal distension is more prominent in ileal atresia

A small amount of pale meconium may be passed, despite the atresia.

Plain abdomen radiographs in a jejunal/ileal atresia case will show multiple air-fluid levels and dilated loops. They don't reliably distinguish small and large bowels in neonates and hence contrast studies are often required to localize the obstruction.

Water-soluble enemas (Gastrogaffin, Hypaque) are particularly useful in differentiating atresia from meconium ileus and Hirschsprung disease. Contrast studies also help to rule out distal intestinal obstruction and occasionally may act therapeutically to relieve obstructions.

In most cases of jejunal/ileal atresia, the distal end of the dilated proximal small bowel is resected and a primary end-to-end anastomosis is done. If the proximal bowel is extremely dilated, it may need to be tapered to the distal bowel before anastomosis.

Other options:

Option A: CT is not recommended for neonates owing to the harmful effects of radiation.

Option C: USG abdomen does not help to localize the site of obstruction and is more useful in the diagnosis of intussusception, where there is an abdominal mass.

Option D: Explorative laparotomy is not done for diagnosing intestinal atresias.

Solution to Question 7:

The utility of injection of methylene blue in the region around the tumor is done for sentinel lymph node biopsy (SLN) in patients without clinically positive lymph nodes.

SLN biopsy rationale:

• Afferent lymphatics from a primary tumor first drain into the sentinel node before draining into the other nodes. So, if a sentinel lymph node is positive for metastsis, it is possible that the other axillary nodes are also involved. Hence, axillary dissection would be necessary.

• If the sentinel node is tumor-free, then regional nodal metastasis is unlikely.

The nodes can be identified using fluorescein dye, methylene blue, or radioactive sulfur colloid-labeled albumin. The dye can be injected into the peritumoral tissue, subareolar, periareolar, or intradermal plane.

Contraindicated in patients with inflammatory breast cancer, stage T4 disease, a history of previous breast cancer or chest wall surgery, breast scarring, and radiotherapy.

Solution to Question 8:

The given clinical image is that of a congenital umbilical anomaly such as patent urachus, urachal cyst, and vitelline cyst (resulting from the persistence of omphalomesenteric duct). The next best step is the USG abdomen to diagnose if the anomaly is cystic, fibrous or fistula and it's connections.

Omphalomesenteric duct abnormalities: Vitello intestinal duct or omphalomesenteric duct is the connection between the yolk sac and the midgut during embryonic life, which eventually gets obliterated. The persistence of the vitellointestinal duct leads to various abnormalities depending on which part of the duct remains patent.

• Persistence of the proximal intestinal end of the duct leads to Meckel's diverticula.

• Persistence of the distal end near the umbilicus leads to the umbilical polyp or sinus.

• Persistence of the distal end with mucus retention leads to the cyst, which may be associated with the intestine or the umbilicus by a fibrous band.

• A fully patent vitellointestinal duct can lead to an enterocutaneous fistula.



Patent urachus occurs due to the failure of urachus to obliterate, resulting in a connection between the bladder and the umbilicus.

Urachal cyst occurs when the urachus fails to obliterate but there is no connection between the bladder and the umbilicus.

The given image below shows a urachal cyst (UC) attached to Meckel's diverticulum (MD) on one side and urinary bladder (UB) on the other side with a patent tract- urachus (U).



Solution to Question 9:

Prostate cancers rarely metastasize to the liver.

The most common malignant tumors of the liver are metastatic lesions. The liver is a common site of metastases from gastrointestinal tumors because of dissemination through the portal

venous system. The most relevant metastatic tumor of the liver to the surgeon is colorectal cancer because of the well-documented potential for long-term survival after complete resection.

The more frequent primary sites include carcinomas of the colon, pancreas, lung, and breast in adults, and neuroblastoma, Wilms tumor, and rhabdomyosarcoma in pediatric patients. Although the origin of the tumor is often known when metastases are biopsied, carcinomas of the pancreas, stomach, and lung are particularly notorious for being occult primary neoplasms.

Solution to Question 10:

The given clinical scenario of trauma with decreased breath sounds and a hyper-resonant note on percussion indicates pneumothorax. The presence of tachycardia, hypotension and elevated JVP is suggestive of tension pneumothorax.

Tension pneumothorax develops when there is a parietal or visceral pleural tear. This tear acts as a one-way valve i.e. air enters the pleural space from the lung or chest wall during inspiration but has no way to escape during expiration, leading to the collapse of the affected lung.

It is characterized by chest pain, tachypnoea, acute dyspnea, sudden onset hemodynamic compromise (tachycardia, hypotension, low O2 saturation, loss of consciousness), neck vein distention, tracheal deviation away from the side of injury, decreased chest movements on the affected side, hyper-resonance on percussion on the affected side and absent or decreased breath sounds on the affected side.

Tension pneumothorax is a clinical diagnosis, and treatment should never be delayed awaiting radiological confirmation.

Management includes:

• Immediate management: Needle decompression or needle thoracostomy with a wide bore needle in the 5th intercostal space anterior to the mid-axillary line. In the pediatric age group, the needle thoracostomy location is at the 2nd intercostal space in the mid-clavicular line.

• Definitive management: Tube thoracostomy or chest tube insertion in the 5th intercostal space in the triangle of safety.

The triangle of safety is bounded anteriorly by the lateral border of the pectoralis major, posteriorly by latissmus dorsi and inferiorly by a horizontal line at the level of the nipple.

The X-ray below shows a left-sided tension pneumothorax. There is a large, well-demarcated area devoid of lung markings on the left side. The yellow arrows are pointing to the tracheal deviation and the mediastinal shift to the right side.



Other options:

Options A: Pericardial tamponade can mimic tension pneumothorax, but the absence of a tympanic note on percussion rules out its possibility. A patient with pericardial tamponade can present with hypotension, raised JVP, and muffled heart sounds (Beck's triad), with normal breath sounds and no tracheal shift.

Option C: Dullness in percussion at the base of the left inferior angle of scapula in conjunction with tubular breath sounds (Ewart's sign), chest pain that improves while sitting upright and worsens when lying down and pulsus paradoxus are indicative of pericardial effusion. Pericardial effusion develops into tamponade when the effusion leads to increased pressure, impairing ventricular filling and reducing cardiac output.

Option D: Massive hemothorax is the accumulation of >1500 ml of blood or one-third or more of the patient's blood volume in the chest cavity. The patient will have collapsed neck veins, decreased breath sounds and a dull note on percussion.

Solution to Question 11:

The initial management of patients with all types of aortic dissection is beta blocker medication to control hypertension along with pain management. Patients with Stanford Type A (DeBakey Types I and II) will require surgical intervention once the patient's blood pressure is stabilized.

Aortic dissection occurs when a defect or flap occurs in the intima of the aorta.

Predisposing factors:

• Age

• Hypertension

• Marfan syndrome

• Pregnancy

• Other connective tissue disorders, for example, Ehlers–Danlos syndrome, giant cell arteritis, systemic lupus erythematosus

• Coarctation of the aorta

• Turner or Noonan syndromes

• Aortic cannulation site (iatrogenic).

The most typical presentation of aortic dissection is a tearing interscapular pain. The dissection may also result in aortic wall rupture into the pericardium (cardiac tamponade) or mediastinum (left haemothorax).

Contrast-enhanced CT is currently the most preferred imaging modality for aortic dissection. Double-lumen aorta is seen on CECT. Transesophageal echocardiography is preferred for patients who are unstable for transportation.

Acute aortic dissection can be classified by the original DeBakey system, based on the location of the proximal tear and distal extent of the dissection flap, or the Stanford system which uses the location of the proximal tear alone.

Site of dissection	DeBakey	Stanford
Both ascending and descending aorta	Type I	Type A
Only ascending aorta and arch of aorta	Type II	Type A
Descending the aorta distal to the subclavian artery	Type IIIaLimited to the thoracic aorta	Type B
Type IIIbExtends to the abdominal aorta		



Management of aortic dissection:

The initial management of patients with all types of aortic dissection is beta-blocker medication to control hypertension along with pain management. The preferred beta-blocker drug is esmolol. Hemodynamic stability is an important consideration.

• Stanford Type A (DeBakey Types I and II) requires surgical intervention once the patient's blood pressure is stabilized.

• Stanford Type B (DeBakey type III) is best managed by antihypertensive drugs. For patients with complicated type B dissection, percutaneously placed endovascular stents are currently preferred. Surgery is reserved for the rare patients not suitable for stenting.

Other options:

Options B and D: Low molecular weight heparin and vitamin K inhibitors are contraindicated in aortic dissection as they can worsen the risk of bleeding.



Solution to Question 12:

The given clinical features of painless testicular swelling with raised AFP levels without distant metastasis point towards the likely diagnosis of stage 1 non-seminomatous germ cell tumour (NSGCT) of the testis. The next step in management is a high inguinal orchidectomy.

Core needle trucut biopsy and FNAC should be never done when there is a suspicion of testicular neoplasm as it will result in the seeding of tumor cells into the skin and lymphatics, leading to upstaging of the tumor (Options B and C). Currently, there is no role for PET-CT in the routine evaluation of non-seminomatous germ cell tumor and seminoma at the time of diagnosis (Option D).

Testicular tumors are classified according to their predominant cell type:

• Germ cell tumors (90-95%): seminomatous (seminoma) or NSGCT (embryonal carcinoma, yolk sac tumor, teratoma and choriocarcinoma).

• Sex cord-stromal tumors (1-2%): Leydig cell tumors, Sertoli cell tumors

• Lymphoma (3-7%)

• Other tumors (1-2%)

The most common presentation of testicular cancer is painless testicular mass. Pain is more commonly associated with NSGCT and two-thirds of patients present with metastatic disease at the time of diagnosis. The symptoms of metastatic disease include abdominal pain, dyspnea, chest pain, cough or hemoptysis.

Management of testicular tumours:

• Ultrasound of the testis should be done in all suspected cases of a testicular tumour to confirm the diagnosis.

• Tumor markers (AFP, hCG, LDH) are measured for the staging of cancer.

• CT- chest, abdomen, and pelvis are done to detect metastasis.

• Orchidectomy via an inguinal incision (high inguinal orchidectomy) is the best line of treatment in case of a suspected testicular tumour.

• Further management is based on staging. The older staging is preferred over the TNM staging for treatment and is given below:

• Stage I: Tumour is confined to testis and epididymis

• Stage II: Nodal tissue is present but is confined to nodes below the diaphragm

• Stage III: Nodal involvement above the diaphragm

• Stage IV: Non-lymphatic metastatic spread (mostly in lungs)

• Stage I tumours:

• Combination chemotherapy with bleomycin, etoposide, and cis-platinum (BEP chemotherapy) is given for NSGCTs, as they are radio-resistant.

• Chemotherapy with cisplatinum and radiotherapy to para-aortic lymph nodes can be given in seminomas as they are radiosensitive.

• Stage II-IV tumours (seminomas and NSGCTs) are treated with BEP chemotherapy and retroperitoneal lymph node dissection if needed.

Solution to Question 13:

The given image shows a Roux-en-Y loop of the small intestine (jejunum) attached to an area of the liver capsule (Roux-en-Y hepaticojejunostomy), which is suggestive of Kasai hepatoportoenterostomy procedure done for biliary atresia in infants. An absent kidney is not associated with this condition.

Extrahepatic biliary atresia (EHBA) is defined as a partial or total absence of permeable bile duct between porta hepatis and the duodenum. It is due to inflammatory fibrosis of the biliary tree. It leads to obstruction to bile flow thereby causing leaking of conjugated bilirubin into the circulation.

Associated anomalies include cardiac lesions, polysplenia, situs inversus, absent vena cava, and a preduodenal portal vein.

In the Kasai procedure, the extrahepatic biliary tree is dissected proximally till the liver capsule, the porta hepatis is transected and a jejunal loop is anastomosed in order to provide a drainage pathway for bile.

Infants with biliary atresia, who do not achieve successful drainage after Kasai's procedure, require liver transplantation within the first year of life. This is the most common reason for liver transplantation in children.

Management of extrahepatic biliary atresia		
Type I	Atresia restricted to the common bile duct	Roux-en-Y hepaticojejunostomy
Type II	Atresia of the common hepatic duct	Kasai procedure(Roux-en-Y hepatoportoenterostomy)
Type III	Atresia of the right and left hepatic duct	

Solution to Question 14:

The cecum is the first organ to be visualized to differentiate between the small and large bowel in cases of intestinal obstruction.

Distension of the cecum will confirm large bowel involvement. The identification of a collapsed distal segment of the large bowel and its sequential proximal assessment will lead to the identification of the cause.

Solution to Question 15:

The best position for a conscious patient to be in during the insertion of Ryle's tube (RT) is sitting with neck flexion. The described position helps in directing the RT towards the esophagus and also prevents it from coiling in the oropharynx or entering the trachea.

RT, also known as nasogastric (NG) tube, measures 120-125 cm. It has 4 markings at 40cm (cardia), 50cm (body), 60cm (pylorus), and 70 cm (duodenum) to indicate the approximate position after the placement.

Indications for NG tube:

• Feeding in patients with dysphagia, decreased appetite, or those who are nil per os (NPO) for medical or surgical reasons.

• Aspiration of gastric fluid

• Gastric decompression in cases of intestinal obstruction and paralytic ileus.

• Zollinger-Ellison syndrome

• Bowel irrigation

The absolute contraindications for an NG tube are severe midface trauma, recent nasal surgery, and basilar skull fractures

Steps for the correct insertion of RT in an adult:

• Examination of the patient's nostril for any anatomical abnormalities and then instillation with 10 ml of 1% lignocaine

• NEX method is used to measure the length of the RT: Place the exit port of the tube at the tip of the nose. Extend the tube to the earlobe, and then to the xiphisternum.

• The patient is asked to sit upright with partial neck flexion. The tip of the Ryle's tube is lubricated and passed along the floor of the nasal passage until a resistance is felt(corresponds to nasopharynx)

• The patient is subsequently asked to swallow water to guide the tube through the esophagus into the stomach until the marked length (based on the NEX method).

• NG tube is finally taped to secure it in place.

Confirmation of the correct position is done by:

• Auscultation in the epigastric area while injecting a bolus of air

• X-ray chest to locate the lead markers

• Aspiration of contents and subsequently confirming gastric origin (pH <6 indicates gastric placement)

Solution to Question 16:

The usual cause of thyroid storm in thyroidectomy is inadequate patient preparation before surgery. Patient preparation pre-operatively involves ensuring a euthyroid state.

Rough handling of the thyroid in thyroidectomy does not cause a thyroid storm. Though a significant amount of thyroglobulin is stored in the acinar cells, they are inactive unless converted

to thyroxine (T_3 and T_4).

Solution to Question 17:

Grey colored cannula is used to achieve maximum flow rate in a patient who needs a large amount of fluids.

Solution to Question 18:

The patients GCS is Eyes-3, Verbal-NT, Motor-6.

It is calculated as follows:

• Eye opening: Opening to verbal command - 3

• Verbal response: In the case of endotracheal intubation, the verbal response score is noted as NT (non-testable)

• Motor respone: Movement of all 4 limbs spontaneously - 6

Total GCS score is not recorded since one of the components is non-testable.

Solution to Question 19:

The given clinical scenario of chest trauma with a hyper resonant note on the right side suggests pneumothorax. Hemodynamic instability in this patient suggests tension pneumothorax and this is managed by immediate decompression by insertion of a needle in the appropriate place.

Pneumothorax refers to the presence of excess air in the pleural space. Entrapment of air occurs by:

• Primary spontaneous pneumothorax - Due to the spontaneous rupture of apical pleural blebs or cystic spaces immediately below the visceral pleura.

• Secondary pneumothorax - Due to the rupture of dilated cystic spaces in lung diseases like chronic obstructive pulmonary disease (COPD)

• Tension pneumothorax - Due to a one-way valve mechanism that directs airflow into the pleural space leading to a higher than atmospheric pressure environment. This leads to the affected lung being unable to expand appropriately and leads to hemodynamic instability.

Tension pneumothorax is a surgical emergency and the excessive pressure has to be immediately relieved. A chest X-ray is not required before emergency decompression as it is a clinical diagnosis, but an X-ray should be done following chest tube insertion (option D). Tension pneumothorax is managed by immediate needle decompression in the 5th ICS (adults), followed by definitive management with a chest tube inserted in the same space.

Other options:

Option A: Intubation is performed when the patient is unresponsive post chest tube insertion.

Option C: While IV fluids are a part of the resuscitation protocol, the management of tension pneumothorax takes higher precedence as the next immediate step.

Note: According to ATLS 10th edition

• In adults, needle thoracostomy or needle decompression location is changed to the 5th ICS anterior to the midaxillary line from the 2nd ICS midclavicular line.

• In the pediatric age group, the needle thoracostomy location remains at the 2nd ICS midclavicular line.

Solution to Question 20:

The most common surgical procedure for achalasia cardia is laproscopic Heller myotomy in conjugation with partial fundoplication.

Surgery AIIMS 2019 (May & Nov)

Question 1:

A patient presents with renal calculus of > 4 cm of the staghorn type. What is the best management?

a) Extracorporeal shockwave lithotripsy

b) Percutaneous nephrolithotomy

c) Ureteroscopic removal

d) Retrorenal surgery

Question 2:

You are doing an emergency laparotomy for a case of intestinal obstruction. Which organ will you first visualize to know if this is a case of small bowel or large bowel obstruction?

a) Ileum

b) Caecum

c) Jejunum

d) Sigmoid colon

Question 3:

What is the best position for a conscious patient to be in during the insertion of a Ryle's tube?

a) Supine with neck flexion

b) Supine with neck extension

c) Sitting with neck flexion

d) Sitting with neck extension

Question 4:

Pick from the following the usual cause of thyroid storm in thyroidectomy:

a) Rough handling of thyroid during surgery

b) Due to parathyroid gland

c) Inadequate patient preparation

d) Infections

Question 5:

Which of the following colored cannula will you use to achieve maximum flow rate, in a patient who needs a large amount of fluids?

a) Grey

b) Green

c) Blue

d) Pink

Question 6:

A patient who was admitted after a road traffic accident is put on mechanical ventilation. He opens his eyes on verbal command and moves all four limbs spontaneously. Calculate his GCS.

a) Eyes-2, Verbal-1, Motor-5

b) Eyes-2, Verbal-NT, Motor-5

c) Eyes-3, Verbal-1, Motor-6

d) Eyes-3, Verbal-NT, Motor-6

Question 7:

A patient was brought to the casualty after a road traffic accident (RTA) with chest trauma. He is conscious and speaks single words. He has a respiratory rate of 44 cycles per min and a BP of 90/40 mmHg. There is a hyper-resonance on percussing the right side of his chest. What will be your next, immediate step?

a) Intubate the patient

b) Insert needle in appropriate intercostal space

c) Start intravenous fluids urgently

d) Take a chest X-ray

Question 8:

Which of the following is used for wound infection grading?

a) ASA classification

b) SIRS criteria

c) Southampton grading

d) APACHE II score

Question 9:

The most common mediastinal tumor is _____.

a) Thymoma

b) Neurogenic

c) Lymphoma

d) Bronchial cyst

Question 10:

On the 5th postoperative day post CABG, the patient develops tachypnea and hypotension. What is the most probable cause?

a) Sepsis

b) Acute kidney failure

c) Acute respiratory failure

d) Acute cardiac failure

Question 11:

The probable diagnosis based on the below clinical image of the breast is?

a) Paget's disease

b) Mondor's disease

c) Zuska's disease

d) Tietze syndrome

Question 12:

The most commonly involved artery in pseudoaneurysm formation:

a) Aorta

b) Femoral artery

c) Popliteal artery

d) None of the above

Question 13:

A chronic alcoholic patient came to the emergency with pain in the epigastrium radiating to the back and recurrent vomiting. On examination, guarding was present in the upper epigastrium. The erect chest X-ray was normal. What is the next step?

a) Upper GI endoscopy

b) Alcohol breath test

c) Serum lipase

d) CECT

Question 14:

The best way to manage the wound shown in the image is:

a) Cleaning and dressing

b) Debridement and dressing

c) Cleaning and skin grafting

d) Cleaning and use of allograft

Question 15:

Device shown in the image is used for which purpose?



a) Prevent viral infections

b) Prevent transfusion related reactions

c) Prevent infections

d) All of the above

Question 16:

In a child with a unilateral undescended testis, a diagnostic laparoscopy is performed which showed the blind-ending testicular vessels on the right side. What is the next best step?

a) Inguinal exploration

b) Abdominal exploration

c) Scrotal exporation

d) No further exploration

Question 17:

What is the surgical site infection rate expected in a patient with a clean-contaminated wound that is operated on after antibiotics prophylaxis?

a) 1-2%

b) 3%

c) 13-20%

d) 20-30%

Question 18:

Which of the following is not a component of qSOFA?

a) Respiratory rate ≥22/min

b) Systolic BP ≤100 mm Hg

c) Altered mental status

d) Bilateral pupillary reflex

Question 19:

Which of the following does not classify as locally advanced breast cancer?

a) Tumor more than 4 cm

b) Inflammatory breast cancer

c) Chest wall involvement

d) Skin involvement

Question 20:

All of the following are features of SIRS except:

a) Temperature > 38 degrees Celsius

b) Respiratory Rate > 20/minute and PaCO2 <35 mmHg

c) Pulse rate > 90/minute

d) WBC > 12000/mm3 or < 4000/mm3

Question 21:

The most common neuroendocrine tumor of the pancreas is

a) Gastrinoma

b) Somatostatinoma

c) Insulinoma

d) VIPoma

Question 22:

After parotidectomy patients complains of loss of sensation while shaving. Which nerve is most likely affected?

a) Auriculotemporal nerve

b) Great auricular nerve

c) Facial nerve

d) Mandibular nerve

Question 23:

ABCDE rule applies for:

a) Malignant melanoma

b) BCC

c) SCC

d) Actinic keratosis

Question 24:

Identify the following instrument?

a) Unipolar cautery

b) Bipolar cautery

c) Harmonic scalpel

d) Ligasure

Question 25:

Radical neck dissection spares which of the following structures?

a) Digastric muscle

b) Sternocleidomastoid muscle

c) Spinal accessory nerve

d) Ipsilateral cervical lymph nodes

Question 26:

A patient comes with active variceal bleeding. Which of the following is not done?

a) Platelet transfusion

b) Arrange pRBC

c) IV Octreotide

d) Urgent Endoscopy

Question 27:

A child presents with pain and swelling in the right side of his scrotum and testicular torsion is suspected. Which is the most appropriate management?

a) Immediate exploration on both symptomatic and asymptomatic side

b) Immediate exploration on symptomatic side only

c) Delayed exploration on both symptomatic and asymptomatic side

d) Delayed exploration on symptomatic side only

Question 28:

What is the instrument shown below that is used to create a pneumoperitoneum ?

a) Veress needle

b) Salah's needle

c) Hasson Trocar

d) Von Graefe needle

Question 29:

Which of the following is the principle behind the wound-healing procedure shown in the image below?

a) Negative pressure wound therapy

b) Positive pressure wound therapy

c) Foam therapy

d) Sponge therapy

Question 30:

Stage the bedsore in the following image:

a) 4

b) 3

c) 2

d) 1

Question 31:

A patient with a road traffic accident presented to the emergency room and requires a large amount of fluids to be transfused. Which of the following IV cannula is preferred?

a) Green

b) Blue

c) Pink

d) Grey

Question 32:

A patient with a nodule of the neck presents to the surgery OPD. A junior resident performs the following method to palpate it. Identify the procedure shown in the video below.

a) Lahey's method

b) Crile's method

c) Kocher's method

d) Pemberton's method

Question 33:

The instrument given in the image is:

a) Hemovac

b) Jackson-Pratt drain

c) Minivac

d) Corrugated drain

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	c
5	a
6	d
7	b
8	c
9	a
10	c
11	b
12	b
13	c
14	a

15	b
16	d
17	b
18	d
19	a
20	b
21	c
22	b
23	a
24	a
25	a
26	a
27	a
28	a
29	a
30	a
31	d
32	a
33	b

Detailed Explanations

Solution to Question 1:

The best management for staghorn renal calculus of >4 cm is percutaneous nephrolithotomy (PCNL).

The initial management of renal stones depends on the stone size and includes conservative management, extracorporeal shockwave lithotripsy (ESWL), and PCNL. PCNL is preferred for stones > 2cm in size and for staghorn calculi, while ESWL is used to treat renal stones of size 0.6 to 1.5 cm. PCNL is indicated in

• Renal stones >2 cm

• Staghorn calculus

• Failed shockwave lithotripsy (SWL) or retrograde intrarenal surgery (RIRS) for renal calculi

• Lower pole renal stones with anatomy that is unfavorable for ESWL

Other options:

Option C: Ureteroscopic removal is attempted for failed SWLs or PCNLs, and is not the first choice in the surgical management of renal stones.

Option D: RIRS is a minimally invasive procedure similar to ureteroscopic removal, where the kidney is accessed using a slim and flexible ureteroscope in a retrograde manner via the ureteric orifice. It is indicated for lower pole renal stones < 2 cm.

Solution to Question 2:

The cecum is the first organ to be visualized to differentiate between the small and large bowel in cases of intestinal obstruction.

Distension of the cecum will confirm large bowel involvement. The identification of a collapsed distal segment of the large bowel and its sequential proximal assessment will lead to the identification of the cause.

Solution to Question 3:

The best position for a conscious patient to be in during the insertion of Ryle's tube (RT) is sitting with neck flexion. The described position helps in directing the RT towards the esophagus and also prevents it from coiling in the oropharynx or entering the trachea.

RT, also known as nasogastric (NG) tube, measures 120-125 cm. It has 4 markings at 40cm (cardia), 50cm (body), 60cm (pylorus), and 70 cm (duodenum) to indicate the approximate position after the placement.

Indications for NG tube:

• Feeding in patients with dysphagia, decreased appetite, or those who are nil per os (NPO) for medical or surgical reasons.

• Aspiration of gastric fluid

• Gastric decompression in cases of intestinal obstruction and paralytic ileus.

• Zollinger-Ellison syndrome

• Bowel irrigation

The absolute contraindications for an NG tube are severe midface trauma, recent nasal surgery, and basilar skull fractures

Steps for the correct insertion of RT in an adult:

• Examination of the patient's nostril for any anatomical abnormalities and then instillation with 10 ml of 1% lignocaine

• NEX method is used to measure the length of the RT: Place the exit port of the tube at the tip of the nose. Extend the tube to the earlobe, and then to the xiphisternum.

• The patient is asked to sit upright with partial neck flexion. The tip of the Ryle's tube is lubricated and passed along the floor of the nasal passage until a resistance is felt(corresponds to nasopharynx)

• The patient is subsequently asked to swallow water to guide the tube through the esophagus into the stomach until the marked length (based on the NEX method).

• NG tube is finally taped to secure it in place.

Confirmation of the correct position is done by:

• Auscultation in the epigastric area while injecting a bolus of air

• X-ray chest to locate the lead markers

• Aspiration of contents and subsequently confirming gastric origin (pH <6 indicates gastric placement)

Solution to Question 4:

The usual cause of thyroid storm in thyroidectomy is inadequate patient preparation before surgery. Patient preparation pre-operatively involves ensuring a euthyroid state.

Rough handling of the thyroid in thyroidectomy does not cause a thyroid storm. Though a significant amount of thyroglobulin is stored in the acinar cells, they are inactive unless converted to thyroxine (T3 and T4).

Solution to Question 5:

Grey colored cannula is used to achieve maximum flow rate in a patient who needs a large amount of fluids.

Solution to Question 6:

The patients GCS is Eyes-3, Verbal-NT, Motor-6.

It is calculated as follows:

• Eye opening: Opening to verbal command - 3

• Verbal response: In the case of endotracheal intubation, the verbal response score is noted as NT (non-testable)

• Motor respone: Movement of all 4 limbs spontaneously - 6

Total GCS score is not recorded since one of the components is non-testable.

Solution to Question 7:

The given clinical scenario of chest trauma with a hyper resonant note on the right side suggests pneumothorax. Hemodynamic instability in this patient suggests tension pneumothorax and this is managed by immediate decompression by insertion of a needle in the appropriate place.

Pneumothorax refers to the presence of excess air in the pleural space. Entrapment of air occurs by:

• Primary spontaneous pneumothorax - Due to the spontaneous rupture of apical pleural blebs or cystic spaces immediately below the visceral pleura.

• Secondary pneumothorax - Due to the rupture of dilated cystic spaces in lung diseases like chronic obstructive pulmonary disease (COPD)

• Tension pneumothorax - Due to a one-way valve mechanism that directs airflow into the pleural space leading to a higher than atmospheric pressure environment. This leads to the affected lung being unable to expand appropriately and leads to hemodynamic instability.

Tension pneumothorax is a surgical emergency and the excessive pressure has to be immediately relieved. A chest X-ray is not required before emergency decompression as it is a clinical diagnosis, but an X-ray should be done following chest tube insertion (option D). Tension pneumothorax is managed by immediate needle decompression in the 5th ICS (adults), followed by definitive management with a chest tube inserted in the same space.

Other options:

Option A: Intubation is performed when the patient is unresponsive post chest tube insertion.

Option C: While IV fluids are a part of the resuscitation protocol, the management of tension pneumothorax takes higher precedence as the next immediate step.

Note: According to ATLS 10th edition

• In adults, needle thoracostomy or needle decompression location is changed to the 5th ICS anterior to the midaxillary line from the 2nd ICS midclavicular line.

• In the pediatric age group, the needle thoracostomy location remains at the 2nd ICS midclavicular line.

Solution to Question 8:

Southampton wound grading system is used to score the severity of wound site infection.

Southampton wound grading system:

• Grade 0 - Normal healing

• Grade I - Normal healing with different grades of erythema

• Grade II - Erythema with other signs of inflammation at different parts of the wound.

• Grade III - Clear or serosanguinous discharge at different sites

• Grade IV - Pus discharge

Other options:

Option A: American Society of Anesthesiologists (ASA) grading is used in pre-anesthetic checkups to assess a patient's physical health and fitness for anesthesia and surgery.

Option B: Systemic Inflammatory Response Syndrome (SIRS) is a set of clinical criteria used to identify and monitor critically ill patients with high-grade burns, severe trauma, and other major surgeries.

Option D: Acute Physiology and Chronic Health Evaluation II (APACHE II) is a scoring system used to assess the severity and predict the outcome of pancreatitis.

Solution to Question 9:

The most common mediastinal tumors in descending order of incidence are thymoma (25%), germ cell tumor (13%), lymphoma, mesenchymal, and thyroid.

The Masaoka classification and TNM staging are used in the staging of thymoma.

The most common mediastinal tumor in children is the neurogenic tumor.

Solution to Question 10:

The clinical presentation of tachypnoea and hypotension in a post-op day 5 patient who underwent coronary artery bypass grafting (CABG) points towards lung collapse (atelectasis) - type 3 respiratory failure.

Atelectasis is the most common respiratory complication in a post-operative patient.

The factors predisposing to atelectasis are general anesthesia, upper abdominal surgery, and thoracic surgery. They commonly present with tachypnoea and decreased oxygen saturation. On examination, the breath sounds are bronchial in nature and reduced in intensity.

Lung atelectasis can be reversed (within 24h-48h) with early mobilization, deep breathing (5 sequential breaths for 5s), incentive spirometry, and chest physiotherapy.

Other postoperative systemic complications include:

• Pneumonia and myocardial infarction : First few days after surgery

• Venous thromboembolism, kidney failure, and urinary tract infection: Days 4 to 7

• Superficial and deep surgical site infections: 2nd post-op week

Other options:

Option A: For a diagnosis of sepsis, SEPSIS 3 criteria needs to be satisfied:

• A documented infection

• An acute increase in >2 SOFA points

Option B: Acute kidney failure is defined as a decrease in urine output (<0.5 mL/kg/h for 6-12h) and a rise in serum creatinine values (>1.5x baseline or >0.3 mg/dL).

Option D: Acute cardiac failure has features of hypotension and tachycardia, with peripheral signs of hypoperfusion characterized by cold clammy extremities.

Solution to Question 11:

The image shows a thrombosed subcutaneous cord attached to the skin of the breast which is the pathognomic feature of Mondor's Disease.

Mondor's disease is the thrombophlebitis of the superficial veins of the breast and anterior chest wall. It may also occur in the arm or penis. On raising the arm, the skin over the chest gets stretched and a narrow, shallow groove alongside the cord is seen. It is treated by restricting arm movements and usually subsides within a few months without recurrence, complications, or deformity.

The differential diagnosis to be considered is lymphatic permeation from an occult carcinoma of the breast.

Other options:

Option A: Paget's disease is a type of ductal carcinoma in situ arising from the nipple and commonly presents as nipple erosion.

Option B: Zuska's disease, also called recurrent periductal mastitis, is a condition of recurrent retroareolar infections and abscesses. Smoking has been implicated as a risk factor for this condition. Managed symptomatically by antibiotics coupled with incision and drainage as necessary.

Option D: Tietze's syndrome refers to the benign inflammation of one or more costal cartilages.

Solution to Question 12:

The most commonly involved artery in pseudoaneurysm is the femoral artery.

A true aneurysm involves all three vessel wall layers, while a pseudoaneurysm involves only the intimal and medial layers. Pseudoaneurysms are most commonly iatrogenic due to repeated arterial puncture for blood sampling or catheterization procedures which injure the arterial wall. The most commonly used artery in endovascular surgeries is the femoral artery, and hence, also the most commonly involved in the formation of pseudo aneurysms.

Aneurysms are usually pulsatile on palpation and a systolic bruit can be heard on auscultation.

Doppler ultrasound reveals a bidirectional, 'to-and-fro' flow of blood which is known as the yin-yang sign.

Treatment options include endovascular stent insertion, ultrasound-guided thrombin injection, and surgical repair.



Other options:

Option A: Aortic aneurysm can be types - thoracic or abdominal. They are usually seen in heritable conditions like Marfan syndrome or vascular-type Ehlers-Danlos.

Option C: The most common peripheral aneurysm is the popliteal aneurysm (80% of the cases) followed by the femoral aneurysm.

Solution to Question 13:

The clinical scenario of the patient presenting with abdominal pain, recurrent vomiting, and upper epigastric guarding in a chronic alcoholic is highly suggestive of acute pancreatitis. The normal erect chest X-ray indicates that there is no pneumoperitoneum (effectively ruling out perforation). The next step is to send blood samples for serum lipase levels.

While contrast-enhanced computed tomography (CECT) of the abdomen (option D) is the best investigation to rule out a case of acute pancreatitis, it is not the next step after an abdominal X-ray.

Acute pancreatitis is often caused by gallstones (70%) or alcoholism (25%). The cardinal symptom is acute-onset severe pain in the upper quadrant of the abdomen. The patient usually presents in shock (hypotension and tachycardia) and tachypnoea. Some may also present with signs of bleeding into the fascial planes leading to discoloration of the sites involved:

• Cullen's sign: Around the umbilicus

• Grey Turner's sign: Flanks

• Fox sign: Inguinal region

• Bryant's sign: Scrotum

The diagnosis is often clinical with a laboratory diagnosis of serum lipase preferred over serum amylase. The severity of the acute pancreatitis is assessed using the Ranson and Glasgow scoring system (≥3 at 48h of onset of pain). APACHE, SOFA, SAPS, and MODS can also be used to assess the severity of pancreatitis in intensive care (ICU) settings.

Pancreatitis is managed conservatively with initial nasogastric drainage followed by intravenous fluids, analgesia, and frequent monitoring of vital signs, urine output, and blood gases.

Other options:

Option A: Upper gastrointestinal intestinal (GI) endoscopy is performed by passing a flexible endoscope and is the best method for examining the upper gastrointestinal mucosa. It doesn't help us diagnose pancreatitis.

Option B: The alcohol breath test is used as a screening test to measure the concentration of alcohol in a person's breath, but doesn't help us diagnose pancreatitis.

Solution to Question 14:

The given image shows a superficial injury to the skin, an abrasion. Cleaning and dressing is the appropriate management of an abrasion.

Wound cleaning is the removal of bacteria and particles to decrease the chances of infection. It includes the "TIME" acronym:

• T - Tissue : Assessment and debridement of non-viable tissue or foreign material

• I - Inflammation/Infection : Assess the etiology of each wound

• M - Moisture Balance : Assess the etiology of wound exudate

• E - Edge of the wound : Assess the edge and surrounding skin

Wound irrigation or toilet is washing the wound with normal saline.

Wound debridement is performed using a surgical blade or fine scissors to remove any devitalized, nonviable tissue or any embedded foreign body (option B).

Skin grafting is required only in situations of wonuds with larger areas soft tissue loss (options C and D).

Solution to Question 15:

The image shows a leukoreduction filter which is used for filtering out donor WBCs. This helps in reducing the incidence of transfusion-related reactions by removing WBC microaggregates. The optimum time for performing leukoreduction is before the storage of blood.

Potential benefits include reduced risk of:

• Sensitization to HLA antigens (important for transplant patients & preventing platelet refractoriness)

• Febrile non-hemolytic transfusion reactions (FNHTR)

• CMV transmission

Note: CMV also has an affinity for monocytes and macrophages (types of WBCs). Reduction in WBCs is the cause for the decrease in CMV transmission (accidental benefit), and thus, the main purpose of a leukoreduction filter is not the prevention of CMV transmission (options A and C).

Possible reduced risk of:

• Transfusion-associated graft versus host disease (TA- GVHD)

• Transfusion-related immuno-modulatory (TRIM) effects

• Transmitting variant Creutzfeldt-Jakob Disease

Other Methods of Leucodepletion are:

• Specific leucodepletion filters (Most Efficient)

• Washing

• Freezing

• Buffy coat removal

• Micro-aggregate filtration

• Low leucocyte apheresis devices

Solution to Question 16:

In non-palpable testis cases, laparoscopically encountering blind-ending vas deferens and testicular vessels prior to entering the internal ring is sufficient to diagnose a vanishing testis. In such cases, no further exploration is required.

Cryptorchidism is the failure of the descent of the intra-abdominal testes into the scrotal sac. It is the most common congenital abnormality of the genitourinary tract. The descent of the testes can arrest anywhere, and the most common site is the inguinal region.

Phases of testicular descent:

• Transabdominal phase: Testes come to lie in the lower abdomen or pelvis, and this descent is regulated by Mullerian-inhibiting substances.

• Inguinoscrotal phase: Testes descend along the inguinal canal and reach the scrotal sac, and this phase of the descent is regulated by androgen-induced calcitonin gene-related peptide from the genitofemoral nerve.

The exact pathogenesis remains unknown; however, deficiency in gonadotropins and Mullerian-inhibiting substance (MIS) and increased expression of estradiol in the placenta are thought to play a role.

Cryptorchidism is histopathologically characterized by the following:

• Testicular atrophy: thickening of the basement membrane, arrested germ cell development, marked hyalinization

• Tubules showing Sertoli cells but no spermatogenesis

• Increase in the interstitial Leydig cells

It is associated with testicular atrophy and dysfunction, which result in sterility. Cryptorchid testes carry a 3–5-fold higher risk for testicular cancer (usually seminoma).

Solution to Question 17:

The surgical site infection rate expected in a patient with a clean contaminated wound that is operated upon after antibiotics prophylaxis is 3%.

SSI rates relating to wound contamination with and without using antibiotic prophylaxis:

Type of surgery	Infection rate without prophylaxis (%)	Infection rate with prophylaxis (%)
Clean(no viscus opened)	1-2	1-2
Clean-contaminated(viscus opened, minimal spillage)	6-9	3
Contaminated(open viscus with spillage or inflammatory disease)	13-20	6
Dirty(pus or perforation, or incision through an abscess)	40	7

Solution to Question 18:

qSOFA score (quickSOFA) is a bedside prompt that may identify patients with suspected infection who are at greater risk for a poor outcome outside the intensive care unit (ICU).

It uses three criteria, assigning one point each for

• Low blood pressure (SBP≤100 mmHg),

• High respiratory rate (≥22 breaths per min)

• Altered mentation (Glasgow coma scale, <15).

The score ranges from 0 to 3 points. The presence of 2 or more qSOFA points is a simple prompt to identify infected patients outside the ICU who are likely to develop sepsis.

Solution to Question 19:

A breast tumor of size > 5 cm (not 4 cm) is classified as locally advanced breast carcinoma (LABC).

Invasive breast carcinoma is of 3 types:

• Early breast carcinoma (EBC) - TNM combination of T1 to T2, N0 to N1, and M0

• Locally advanced breast carcinoma (LABC) - T4N0M0, T3N1M0, TanyN2-3M0

• Advanced breast carcinoma (ABC) - TanyNanyM1

Locally advanced breast cancer (LABC) includes:

• Breast tumors more than 5 cm in diameter

• Cancers that involve the skin or the underlying muscles of the chest (T4 of TNM staging)

• Cancers that involve multiple local lymph nodes (axillary or sub clavicular)

• Inflammatory breast cancer - a rapidly growing type of cancer that makes the breast appear red and swollen

Solution to Question 20:

Respiratory Rate > 20/minute or (Not and) PaCO2 < 32 mmHg (Not <35 mmHg) would be a feature of SIRS (Systemic inflammatory response syndrome).

Systemic inflammatory response syndrome (SIRS) is an exaggerated defense response of the body to a noxious stressor (infection, trauma, surgery, acute inflammation, ischemia, or reperfusion) to localize and then eliminate the endogenous or exogenous source of the insult.

SIRS is diagnosed when any two of the following are present:

• Hyperthermia (>38 Degree Celsius) or Hypothermia (<36 Degree Celsius)

• Tachypnoea (>20/min) or PaCO2 < 32 mmHg

• Tachycardia (>90/min)

• White cell count >12 x 109/liter or <4 x 109/liter or >10% Band forms

Sepsis is SIRS with a documented infection.

Solution to Question 21:

The most common neuroendocrine tumor of the pancreas is insulinoma (4 per million).

Insulinoma is an insulin-producing tumor of the pancreas, with more than 90% of them being benign. It occurs in >40 years in patients without MEN 1 syndrome and <40 years in patients with MEN 1. It presents with symptoms of hypoglycemia such as weakness, headaches, sweating, faintness, seizures, and altered behavior. These occur after fasting or exertion and improve after glucose intake.

The investigation of choice for insulinoma is the determination of 72-hour fasting glucose and serum insulin levels. C-peptide and proinsulin levels are also elevated.

Medical management includes:

• Frequent carbohydrate-rich meals

• Somatostatin analogs like octreotide are effective in controlling symptoms due to excess insulin secretion. They should be used with caution as they can suppress counterregulatory hormones, such as growth hormone (GH), glucagon, and catecholamines, and precipitously worsen hypoglycemia.

• Diazoxide can be used in the initial management of insulinoma as it inhibits insulin release. Its side effects include sodium retention and nausea.

• Everolimus has been found to be highly effective in improving glycemic control in patients with insulinoma owing to its antitumor effect and induction of insulin resistance.

Enucleation and pancreatectomy are the definitive surgical management of insulinoma.

Other options:

Option A: Gastrinoma has an incidence of 0.5 to 4 in a million.

Options B & D: Somatostatinoma and VIPoma are much rarer than gastrinoma.

Solution to Question 22:

After parotid surgery, the nerve involved in the loss of sensation while shaving is the great auricular nerve.

The great auricular nerve (GAN) originates from the second and third nerve roots of the cervical plexus (C2-C3). It supplies the skin over the lower pinna, pre-auricular region, and angle of the mandible (image below). The main trunk passes over the sternocleidomastoid and into the anterosuperior portion of the parotid glands and branches into the anterior and posterior GAN. The anterior branch of GAN is often injured during parotid surgery during the initial elevation of the skin flap.

When GAN is injured, it can cause reduced sensations in the earlobe, pre-auricular skin, and the angle of the mandible that can interfere with habits such as shaving, wearing earrings, etc.



Other options:

Option A: The auriculotemporal nerve, a branch of the mandibular nerve (V3 of trigeminal) supplies the posterior part of the temple, tragus, and adjoining auricle. It does not supply the area where a person is likely to shave. However, the injury to the auriculotemporal nerve resulting in aberrant regeneration would result in Frey's Syndrome (gustatory sweating).

Option C: Though the facial nerve is the most common nerve affected during parotidectomy, any damage would result in facial paralysis and not loss of sensation of the face.

Option D: The mandibular nerve is a branch of the trigeminal nerve (cranial nerve V) that supplies the skin over the mandible, lower lip, and cheek. However, they are not injured during parotidectomy.

Solution to Question 23:

ABCDE algorithm is used to diagnose melanoma.

It comprises of:

• Asymmetry

• Border irregularity

• Color variegation

• Diameter of more than 6 mm

• Evolution or change

Melanoma is a malignant tumor arising from melanocytes at the epidermal-dermal junction. Though it represents only 4% of skin cancers, it accounts for 80% of deaths (high mortality rate). It presents as a pigmented lesion that has enlarged, ulcerated, or bled.

Histopathological confirmation is the gold standard in the diagnosis of melanoma. It is divided into four main types: nodular melanoma (NM), superficial spreading melanoma (SSM), lentigo maligna melanoma (LMM), and acral lentiginous melanoma (ALM).

The treatment consists of wide local excision of the tumor and follow-up to detect any local recurrence around the scar and excluding nodal spread.

Solution to Question 24:

The given instrument is a unipolar cautery/monopolar cautery.

Other options:

Option B: The bipolar cautery contains two tips, unlike unipolar cautery which has only one tip. The current flows between these two tips and hence it does not need a return electrode, unlike the unipolar cautery.

Option C: The harmonic scalpel (shown below) is a surgical instrument used to simultaneously cut and cauterize tissue. Ultrasonic energy is used in Harmonic scalpel where ultrasonic energy is converted to mechanical energy at the active blade.

Option D: LigaSure vessel sealing system (shown below) is a bipolar apparatus for sealing vascular tissue. It applies a high coaptive pressure during the generation of tissue temperature under 1000°C, hydrogen cross-links are first ruptured and then renatured, resulting in a vascular seal that has high tensile strength.

Solution to Question 25:

Digastric muscle is spared in radical neck dissection (RND). RND is used in the treatment of malignant lesions of the neck with local spread or metastasis.

In 1905, Crile described RND where Level I-V LN, sternocleidomastoid (SCM), internal jugular vein (IJV), and XI cranial nerve (spinal accessory nerve) are removed.

In 1962, Suarez described modified RND (MRND) where Level I-V LNs are removed and the following are preserved:

• MRND I - XI cranial nerve

• MRND II - XI cranial nerve and IJV

• MRND III - XI cranial nerve, IJV, and SCM

Note: RND with an additional LN/muscle/nerve is called extended RND.

Solution to Question 26:

Platelet transfusion is not done in the case of active variceal bleeding unless coagulopathy is confirmed as the cause. Platelet transfusions are indicated only in patients with a platelet count of <50,000/microL or in patients with evidence of microvascular bleeding.

Variceal hemorrhage is a medical emergency. Varices usually present with the acute onset large-volume haematemesis, the lower esophagus being the most common site for variceal bleeding.

The sites for portosystemic anastomosis are:

• Left gastric vein to distal oesophageal veins: oesophageal and gastric variceal bleeding

• Superior rectal veins to Middle and inferior rectal veins: hemorrhoids and rectal bleeding

• Left branch of the portal vein to Periumbilical veins: caput-medusae

• The right branch of the portal vein to retroperitoneal veins draining into azygos, hemiazygos, lumbar, intercostal, and phrenic veins

• Omental and colonic veins near hepatic and splenic ■exures to Retroperitoneal veins near hepatic and splenic ■exures

Medical management of variceal bleeding includes blood transfusion (option B) and drug therapy with somatostatin analogs like octreotide (option C). Short-term antibiotics like ceftriaxone are known to decrease bacterial infection and increase survival.

Surgical measures are preferred if medical management fails. They include balloon tamponade, endoscopic banding (option D), transjugular intrahepatic portosystemic shunt (TIPS), portosystemic shunts, splenectomy, and gastric revascularization

Note: Sugiura procedure is the non-shunt surgical management of refractory oesophageal variceal bleeding. It is a procedure of de-vascularizing the distal esophagus and stomach.



Solution to Question 27:

In a case of suspected testicular torsion, immediate exploration needs to be done on both the symptomatic as well as the asymptomatic side. After the de-torsion, the adequacy of the blood flow has to be assessed by Doppler.

In testicular torsion, the blood supply to the testicle is compromised due to the twisting of the spermatic cord within the tunica vaginalis, resulting in ischemia of the epididymis and the testis. The patient usually presents with acute onset pain and swelling of the scrotum. On examination, the cremasteric reflex is lost, and the scrotum looks asymmetrical.

Prehn's sign is used to differentiate epididymitis from testicular torsion. Prehn's sign is said to be positive when the pain gets relieved on lifting the testis, and it is seen in epididymo-orchitis.

At the time of surgical exploration, contralateral testes also must be explored and fixed to the dartos fascia in order to prevent possible torsion occurring in this testes.

More than 80% of the testis can be salvaged if surgery is performed within 6 hours. If the testis is found to be necrosed on scrotal exploration, orchidectomy of the necrosed testis and orchidopexy of the contralateral testis is done.

Solution to Question 28:

The given image showing a long, slender laparoscopic instrument is Veress Needle. It is used in direct puncture laparoscopy to produce a pneumoperitoneum.

Direct puncture laparoscopy (closed method):

• The abdominal wall is grasped with towel clips or fingers to elevate the abdominal wall away from the underlying structures.

• A small incision is made in the umbilicus and the Veress needle is introduced into the abdomen. Two distinct pops can be felt as the needle pierces the abdominal wall fascia and the peritoneum.

The umbilicus is the preferred location for introducing a Vereess needle because the abdominal wall is thin at this location.

Carbon dioxide (CO_2) gas is usually used and the maximum pressure is in the range of 14-15 mmHg. But N_2O is used if the procedure is done under local anesthesia. Insufflation with N_2O is stopped at a pressure of 10 mmHg or when 2L of gas is insufflated.



The Hasson Trocar is used to gain access to the peritoneal cavity and generate pneumoperitoneum by the open direct peritoneal access technique (option C). This is known as the Hasson technique and is done only occasionally. It is preferred for patients who have undergone previous abdominal surgeries as the small bowel may be adherent to the undersurface of the abdominal wound in such cases.



Other options:

Option B: Salah's needle is a type of reusable bone marrow biopsy needle that has a guard situated 2 cm from the tip and a side screw.

Option D: Von Graefe needle (also known as Graefe's knife) is primarily used in ophthalmic surgery.

Solution to Question 29:

The given image shows negative pressure wound therapy using a vacuum-assisted closure (VAC) device to promote healing in chronic wounds.

Principle: An intermittent negative pressure of -125 mmHg hastens debridement and formation of granulation tissue in chronic wounds.

Indications: Chronic wounds (such as diabetic foot ulcers, stage III and IV pressure ulcers), traumatic wounds, flaps and grafts, and dehisced incisions.

Contraindications:

• Exposed vital structures (organs, blood vessels, or vascular grafts)

• Preexisting infection

• Inadequate debridement of devitalized soft tissue or bone increases the risk of infection.

• The presence of malignancy (Marjolin's ulcer).

• An allergic reaction towards the adhesive used to maintain the vacuum seal.

Though not an absolute contraindication, there is no appreciable benefit for ischemic wounds.

Solution to Question 30:

The given image shows complete skin loss that involves muscles and shows exposed bone indicating a stage 4 pressure sore.

Pressure sores are skin or soft tissue injuries that occur due to prolonged pressure over specific areas of the body. An external pressure of >30 mmHg leads to the formation of pressure sores. The common sites of pressure sores in decreasing frequency are the ischium, greater trochanter, sacrum, heel, malleolus (lateral followed by medial), and occiput.

Pressure sores can be prevented with good skincare, the use of air-fluidized beds, special pressure-dispersion cushions, or foams. Repositioning the patient every two hours can lessen the pressure on the area.

Ulcers of stages 1 and 2 often do not require operative measures. They are managed with regular wound cleaning and dressing with a good antibiotic cover.

Stage 3 and 4 ulcers are adequately debrided, and vacuum-assisted closure (VAC) is used to promote the growth of granulation tissue required for surgical closure. Large skin flaps that have muscle with sensory innervation are best for reconstruction.

Solution to Question 31:

Grey coloured IV cannula is required for this patient in order to transfuse large amount of fluids.

The ATLS 10th edition guidelines state that 2 cannulae of atleast 18G should be used to gain vascular access in trauma patients. In this scenario requiring large fluid transfusions, the grey (16G) cannula would be preferable to the green (18G) cannula due to the higher flow rate.

Solution to Question 32:

The video depicts Lahey's method of palpation of the left lobe of the thyroid gland.

In Lahey's method, for palpating the left lobe, the examiner initially pushes the right lobe medially with the left hand, and then using the right hand, the border and consistency of the left lobe is palpated.

Other options:

Option B: In Crile's method the patient is asked to swallow, while the examiner palpates the gland with the thumb. This gives a clear indication regarding the slight enlargement of the thyroid gland or small nodules.



Option C: Kocher's method is done to demonstrate the pressure effect of the thyroid gland on the trachea or larynx. Lateral compression will result in stridor.

Option D: Pemberton's sign (not method), is engorgement of neck veins when the patient is asked to raise their hands above the head and the arms should touch the ears. It is seen in patients with retrosternal goiter.

Solution to Question 33:

The given image is of a Jackson-Pratt drain.

Drains are placed at surgical sites to freely drain the air or fluids that may accumulate. They promote wound healing by preventing the formation of unnecessary cavities. Drains can be:

• Open : They are based on the principle of gravity

• Semi-open : Their action is based on capillary effect

• Closed : They carry out their actions through suction

Jackson-pratt drain is a closed, air-tight drainage system that operates by self-suction. The drain is sutured in place on the skin at the site of insertion to promote stability. There are two main parts to a JP drain:

• Hollow tube

• Collection bulb

The drain should be emptied as often as possible to allow the bulb to be compressed fully in order to maintain suction. This is usually done every 4-6 hours the first few days.



Other options:

Option A: Hemovac is also a closed type of suction drain which can typically handle larger amounts of fluid than the JP drain.

Option C: Minivac is a closed wound drainage system under negative pressure used in minor surgeries.

Option D: Corrugated drain is an open drainage system to remove excess fluid from the operated site.

Surgery AIIMS 2020 (May and Nov INI-CET)

Question 1:

A 65-year-old man presents with abdominal pain for the past 6 hours, BP- 89/50mmHg, RR-24/min, saturation 92% with 6 L of oxygen. Radiograph shows extraluminal air in the abdomen. Which of the following must be done before shifting this patient into the OT?

a) Intubate the patient

b) Insert central venous catheter as initial IV access of choice

c) 1 litre of crystalloid infusion

d) Confirm the diagnosis with CECT

Question 2:

A patient with recurrent pain in the right hypochondrium, nausea, and vomiting is diagnosed with biliary dyskinesia. Which of the following is least useful in the diagnosis of this condition?

a) ERCP

b) HIDA scan

c) USG

d) Post surgery follow-up

Question 3:

A 32-year-old woman was declared brain dead. Her family has decided to donate her organs. Who is the ideal/best recipient?

a) 14 year old child with multi organ dysfunction syndrome

b) 50 year alcoholic man with liver cirrhosis with 1 month abstinence

c) 70 year old woman with end stage renal disease

d) 32 year old woman with renal failure

Question 4:

ERCP is least useful in the diagnosis of _____

a) Recurrent pancreatitis

b) Unexplained jaundice

c) Periampullary mass

d) Acute pancreatitis without choledocholithiaisis

Question 5:

Which of the following is a correct statement regarding retroperitoneal trauma?

a) All zone 1 injury to be treated conservatively

b) Zone 3 injury whether blunt or penetrative should always be explored

c) Zone 2 injury with expanding hematoma should be treated surgically

d) Mattox maneuver is when the ascending colon is mobilized to the right to visualise the aorta

Question 6:

Which of the following nerve is least injured during thyroid surgery?

a) Recurrent laryngeal nerve

b) Ansa cervicalis

c) External laryngeal nerve

d) Marginal mandibular nerve

Question 7:

eFAST is least helpful in diagnosing which of the following conditions?

a) Pericardial Effusion

b) Pneumothorax

c) Retroperitoneal hematoma

d) Renal injury

Question 8:

A 23-year-old man presented with burn injury. He is covered with burns on his head & neck, both upper limbs, front and back of the chest. Calculate the percentage of burns.

a) 27%

b) 36%

c) 45%

d) 18%

Question 9:

FAST stands for _____

a) First assessment with sonography in trauma

b) Focussed assessment with sonography in trauma

c) First abdominal sonography in trauma

d) Focussed abdominal sonography in trauma

Question 10:

A patient with unilateral breast cancer is started on tamoxifen. This will increase the risk of?

a) Ovarian cancer

b) Endometrial cancer

c) Breast cancer on other side

d) Cervical cancer

Question 11:

Which of the following is the most appropriate mesh for use in hernioplasty?

a) Heavyweight with small pores

b) Heavyweight mesh with large pores

c) Lightweight mesh with small pores

d) Lightweight mesh with large pores

Question 12:

A 30-year-old teacher underwent parotidectomy following which she developed lower lip paralysis. This probably occurred because of an injury to which of the following structures?

a) Temporal branch of facial nerve

b) Cervical branch of facial nerve

c) Buccal branch of the facial nerve

d) Mandibular branch of facial nerve

Question 13:

Which of the following is not an adverse effect seen after a massive blood transfusion?

a) Hypocalcemia

b) Hypercalcemia

c) Hypokalemia

d) Hyperkalemia

Question 14:

Which of the following gene mutations are involved in carcinoma of the colon?

a) APC, BRCA, k-ras

b) APC, k-ras, p53

c) MYh, k-ras, APC

d) MYH, p53, DCC

Question 15:

Which among the following is correct regarding adequate thyroid FNAC sampling?

a) 6 follicular clusters with 10 cells each

b) 3 follicular clusters with 10-15 cells each

c) 10 follicular clusters with 6-8 cells each

d) 12 follicular clusters with 10-12 cells each

Question 16:

Which of the following are used to stop esophageal variceal bleed?



a) A, B, C and D

b) A and B only

c) A, B and C

d) A, B and D

Question 17:

A 25-year-old male patient was brought to the ER after RTA, his RR is 25/min, BP is 80/45mmHg. On examination, hyper-resonance is heard over the left side of the chest. The chest X-ray performed is given below. What would be the appropriate management in this case?



a) Wide bore needle insertion in the 5th intercostal space (ICS)

b) Wide bore needle insertion in the 5th ICS followed by an intercostal drain in the 5th ICS

c) Aspiration of the pleural fluid

d) No intervention needed

Question 18:

A patient presents with pain in the right upper quadrant on the second postoperative day following a cholecystectomy. Evaluation reveals a collection in the subhepatic space. A bile leak was suspected. Which of the following is the best investigation for this patient?

a) MRCP

b) ERCP

c) HIDA

d) CECT

Question 19:

What is the most likely diagnosis of the lesion shown in the image below?

a) Arterial insufficiency

b) Neuropathic ulcer

c) Venous ulcer

d) Tuberculous ulcer

Question 20:

A woman was diagnosed to have metastatic liver adenocarcinoma with CK 7 (-), CK20 (+), and CDX-1 (+) IHC markers. Which of the following is the most likely primary site of origin for this metastasis?

a) Colon carcinoma

b) Breast Carcinoma

c) Pancreatic carcinoma

d) Ovarian carcinoma

Question 21:

A 35-year-old chronic smoker presents with bluish discoloration of the toe along with intermittent claudication. Which of the following will be the next appropriate investigation in this patient?

a) CT angiography

b) MR angiography

c) Digital subtraction angiography

d) Duplex scan

Question 22:

A 40-year-old woman presents with complaints of swelling of the right breast over the past 15 days. On examination, there was local rise of temperature, and the swelling was mobile over the chest wall. What is the likely diagnosis?

a) Inflammatory carcinoma

b) Giant fibroadenoma

c) Phyllodes tumor

d) Galactocele

Question 23:

A middle-aged man presents with progressive difficulty in swallowing for both solids and liquids for 8 months. Barium swallow done is shown below. What other investigations need to be done in this patient?



a) Upper GI endoscopy & Manometry

b) Upper GI endoscopy & PET scan

c) Upper GI endoscopy & CECT

d) Upper GI endoscopy and 24 hour PH monitoring

Question 24:

A 55-year-old man with IHD presents with sudden, severe, generalized abdominal pain and multiple episodes of vomiting. On examination, the pulse rate is 118bpm, rhythm is irregularly irregular, BP is 90/60 mmHg. The abdomen appears soft, non-distended, and bowel sounds are absent. Which of the following could be a likely diagnosis in this case?

a) Acute severe pancreatitis

b) Ruptured aortic aneurysm

c) Acute mesenteric ischemia

d) Peptic ulcer perforation

Question 25:

A 15-year-old girl presented with a slow-growing, rubbery, non-tender swelling on the side of the neck as shown in the image. Examination shows the presence of freckles and skin patches over the face and trunk. What is the most likely diagnosis?

a) Cystic hygroma

b) Chemodectoma

c) Neural tumor

d) Lipoma

Question 26:

What is the size of the cannula shown in the below image?



a) 16 G

b) 18 G

c) 20 G

d) 22 G

Question 27:

The position of the patient in the image shown below is known as:

a) Lloyd Davis position

b) Jack knife position

c) Trendelenberg position

d) Reverse trendelenberg position

Question 28:

What will be the most likely associated condition with increased TSH and low T4 levels?

a) Hypoparathyroidism

b) Pituitary Adenoma

c) Grave's disease

d) Hashimoto's thyroiditis

Question 29:

In e-FAST which of the following body part is included that is not included in FAST?

a) Pelvic cavity

b) Retroperitoneum

c) Thoracic cavity

d) Dural and subdural spaces

Question 30:

According to Strasberg's classification of bile duct injuries, injury to the right aberrant posterior sectoral duct with bile leak is:

a) Type A

b) Type B

c) Type C

d) Type D

Question 31:

A bus conductor was diagnosed with varicose veins. On examination, skin eczema and lipodermatosclerosis were noted. According to the CEAP classification system, He would be classified as

a) C1

b) C2

c) C3

d) C4

Question 32:

Which of the following is the most commonly seen tumor in the retroperitoneal space?

a) Liposarcoma

b) Rhabdomyosarcoma

c) Synovial sarcoma

d) Neurofibroma

Question 33:

A 50-year-old female patient presents with a 4*3cm irregular mass in the upper outer quadrant of the right breast, which is not fixed to the skin or the chest wall. A lymph node of 2cm is palpated in the right axilla. No mass is palpated in the opposite breast. Which of the following investigations would you perform on this patient?

a) 1, 2, 3 & 4

b) 1, 2 & 3

c) 1, 3 & 4

d) 1 & 3

Question 34:

According to the European Hernia Society classification, M3 incisional hernia refers to:

a) Epigastric

b) Infraumbilical

c) Umbilical

d) Suprapubic

Question 35:

Which of the following is true about the Tillaux sign?

a) Mass is felt in the periumbilical region

b) Mass is brilliantly transilluminant

c) Mass is mobile perpendicular to mesentery attachment line

d) Fluctuation is positive

Question 36:

Match the correct answer related to the nerve injuries in various surgeries:

a) A-1, B-2, C-3, D-4

b) A-2, B-3, C-4, D-1

c) A-3, B-4, C-1, D-2

d) A-4, B-1, C-2, D-3

Question 37:

A child is diagnosed with congenital hypertrophic pyloric stenosis, which among the following can be seen in this child?

a) 2, 3 and 4

b) 3 and 4 only

c) 1, 2 and 4

d) 1, 2, 3 and 4

Question 38:

The components of qSOFA scoring include:

a) 1 & 2

b) 3 & 4

c) 1, 2 & 3

d) 1, 2, 3 & 4

Question 39:

Which of the following is true regarding inguinal hernia?

a) 1 and 3

b) 1 and 4

c) 2 and 3

d) 2 and 4

Question 40:

Match the following:



a) A-3, B-4, C-2, D-1

b) A-3, B-1, C-2, D-4

c) A-2, B-4, C-3, D-1

d) A-2, B-1, C-3, D-4

Question 41:

A 40-year-old man presents to the ER with severe abdominal pain. On examination, his HR is 110bpm, BP- 88/60mmHg. On per abdomen examination, guarding is present. The erect abdominal X-ray is shown below. Which of the following would you proceed with, for the management of this case?

a) 2 & 3

b) 2 & 4

c) 1, 2 & 3

d) 1, 2 & 5

Answer Key

Question No.	Correct Option
1	c
2	c
3	d
4	d
5	c
6	d
7	c
8	c
9	b
10	b
11	d
12	d
13	b
14	b
15	a
16	a
17	b
18	b
19	b
20	a
21	d
22	a
23	a
24	c
25	c
26	c
27	d

28	d
29	c
30	c
31	d
32	a
33	c
34	c
35	c
36	a
37	b
38	c
39	c
40	b
41	a

Detailed Explanations

Solution to Question 1:

Volume resuscitation with 1 liter of crystalloid infusion should be done before shifting this patient to the OT for an emergency exploratory laparotomy.

The given clinical scenario in a patient with hypotension and tachypnea with extraluminal gas in the abdomen is suggestive of perforation peritonitis with hemodynamic instability. The immediate step is volume resuscitation to maintain circulating volume and organ perfusion.

Peritonitis is the inflammation of the peritoneum and can be categorized as local or diffuse and acute or chronic.

Causes:

• Gastrointestinal perforation e.g. perforated ulcer, appendix, diverticulum

• Transmural translocation (no perforation), e.g. pancreatitis, ischaemic bowel, primary bacterial peritonitis

• Exogenous contamination, e.g. drains, open surgery, trauma, peritoneal dialysis

• Female genital tract infection, e.g. pelvic inflammatory disease

• Haematogenous spread (rare), e.g. septicaemia

• Miscellaneous- e.g. familial Mediterranean fever

Clinical features:

• Abdominal pain, worse on movement, coughing and deep respiration

• Constitutional symptoms: anorexia, malaise, fever, lassitude

• nausea +/– vomiting

• Pyrexia (may be absent)

• Tachycardia

• Tenderness +/– guarding/rigidity/rebound of the abdominal wall

• Pain/tenderness on rectal/vaginal examination (pelvic peritonitis)

• Absent or reduced bowel sounds

• Septic shock (systemic inflammatory response syndrome [SIRS] and multiorgan dysfunction syndrome [MODS]) in later stages

Management:

General care of the patient:

• Correction of fluid and electrolyte imbalance

• Insertion of a nasogastric tube and urinary catheter

• Broad-spectrum antibiotic therapy

• Aprropriate analgesia

Surgical treatment: Exploratory laparotomy and repair of perforation is the definitive management. Presence of peritoneal signs and hemodynamic instability are indications for urgent surgery.

Other options

Option A: Intubation is not indicated by the patient's current vital signs.

Option B: Initial volume resuscitation can be done with the peripheral large-bore cannula. A central venous catheter may be needed if peripheral access is dificult, vasopressors are administered, patient requires TPN.

Option D: A CT scan, while helpful for locating lesion and confirming the diagnosis, is not indicated for a hemodynamically unstable patient.

Note:

According to recent ATLS guidelines, crystalloid infusion should be limited to a maximum of 1.5 litres as crystalloid resuscitation beyond 1.5 liters increases the risk of death.

Solution to Question 2:

USG is the least useful investigation in the diagnosis of biliary dyskinesia or sphincter of Oddi dysfunction since the findings can be normal or show non-specific changes such as dilatation of the common bile duct.

Sphincter of Oddi dysfunction (SOD) is caused due to a structurally or physiologically abnormal sphincter with a higher tone, which fails to relax, causing pain and recurrent episodes of pancreatitis with normal liver function tests. There are two clinical types: biliary SOD and pancreatic SOD. Each type is further classified into types 1,2 and 3.

Other options:

Option A: The best investigation for the diagnosis of SOD is ERCP with manometry of the biliary and pancreatic sphincters.

Option B: HIDA scan will show a reduced gallbladder ejection fraction (EF). A gallbladder EF <35% is considered diagnostic of biliary dyskinesia.

Option D: Biliary-type SOD should be considered and excluded in post-cholecystectomy syndrome; hence, post-surgery follow-up is essential.

CT and MRCP can demonstrate dilatation of the biliary and pancreatic ducts.

Solution to Question 3:

Among the given options, a 32-year-old woman with renal failure is the most suitable recipient for transplantation.

This is because the prognosis for the other patients mentioned above is poor. Prioritization of organ transplant recepients should be done based on post-transplant outcomes.

Option A - 14-year-old with multi-organ dysfunction syndrome is not suitable because there is a poor prognosis for the ongoing organ dysfunction.

Option B - Alcoholic patients with just 1-month of abstinence are not considered for a transplant.

Option C - A 70-year-old woman with end-stage renal disease is not the right candidate because the chances of long-term survival and a successful outcome are very less.

Solution to Question 4:

ERCP is least useful in the diagnosis of acute pancreatitis without choledocholithiasis.

ERCP stands for Endoscopic Retrograde Cholangio-Pancreatography. It involves the use of a side-viewing endoscope which is passed through the pylorus of the stomach into the second part of the duodenum to visualise the papilla. This is then cannulated, and dye is injected into the pancreatic or bile duct to visualise the pancreatic or biliary tree. It can be therapeutic or diagnostic. It is the best diagnostic modality for viewing the distal biliary tree, but MRCP is preferred as the diagnostic modality unless a therapeutic intervention is also needed.

Indications of ERCP:

• Choledocholithiasis

• Acute cholangitis

• Biliary stricture, biliary leak

• Pancreatic duct stricture, pancreatic stones

• Extrahepatic biliary strictures related to primary sclerosing cholangitis

• Endoscopic therapy for some patients with sphincter of Oddi dysfunction

• Biliary or pancreatic ductal system tissue sampling

• Balloon dilatation of the duodenal papilla or ductal strictures

• Sphincterotomy

The most common therapeutic indication for ERCP is the relief of biliary obstruction due to gallstone disease and benign or malignant biliary strictures.

Contraindications of ERCP include patients who cannot tolerate anesthesia, patients with type 3 sphincter of Oddi dysfunction, luminal gastrointestinal obstruction (examination limited to an area proximal to the level of obstruction and untreated hemostatic disorder (high risk of bleeding).

Complications of ERCP include duodenal perforation, hemorrhage, pancreatitis and sepsis.

Other options:

Option A: ERCP can help in ruling out the etiology in cases of recurrent pancreatitis.

Option B: ERCP can act as both diagnostic and therapeutic in cases of unexplained jaundice.

Option C: ERCP can be useful in case of a periampullary mass to take a tissue biopsy and cytology.

Solution to Question 5:

A zone 2 retroperitoneal injury with expanding hematoma should be treated surgically.

Zone of retroperitoneal injury	Type of viscera	Differential management
Zone 1	Central	Always explored, after proximo-distal vascular control
Zone 2	Lateral	Only explored if the hematoma is pulsatile or expanding, else conservative/ angioembolization
Zone 3	Pelvic	Only explored if the hematoma is pulsatile or expanding, else conservative/ angioembolization



Surgical exploration is done in cases of penetrating injuries to any zone.

Other options:

Option A: Zone 1 injuries are always explored.

Option B: Depending on the situation, zone 3 injuries can be managed with surgical exploration or conservatively.

Option D: Mattox manoeuvre is left medial visceral rotation to expose the aorta, while the Cattell-Braasch manoeuvre is done by right medial visceral rotation to expose IVC.

Solution to Question 6:

The marginal mandibular nerve is least likely to be injured during thyroid surgery as it is high up (mandibular margin) than the dissection plane of the approach to thyroid surgery.

Other nerves that might be at risk of injury include:

• Recurrent laryngeal nerve may be injured if dissection is done below the cricoid cartilage or near the tracheoesophageal groove (option A).

• Ansa cervicalis may be injured if strap muscles are not cut high enough. (option B)

• External laryngeal nerve may be injured during ligation of superior thyroid vessels as it lies very close to the superior thyroid artery (option C).

Complications of thyroidectomy:

• Hemorrhage - most common complication overall; due to slipping of ligature

• Nerve injury

• External laryngeal nerve - hoarseness (more common)

• Recurrent laryngeal nerve - hoarseness if unilateral, life-threatening aspiration and aphonia if bilateral

• Respiratory obstruction

• Laryngeal edema - the most common cause of respiratory obstruction

• Bilateral recurrent laryngeal nerve palsy

• Laryngomalacia

• Tension hematoma

• Hypoparathyroidism - infarction of parathyroids

• Thyroid storm - inadequately prepared for surgery

• Hypothyroidism

• Recurrent thyrotoxicosis

• Wound infection, stitch granuloma, keloid.

Solution to Question 7:

Among the given options, eFAST (focussed assessment with sonography in trauma) is the least useful in diagnosing retroperitoneal hematoma as the sensitivity of ultrasound for the detection of retroperitoneal hematomas is extremely poor.

Computed Tomography (CT Scan) is the investigation of choice for retroperitoneal injury, but it can only be done in a hemodynamically stable patient.

FAST stands for Focussed Assessment with Sonography for Trauma. It is a rapid, non-invasive bedside test for detecting free fluid using sonography used to assess patients for potential thoracoabdominal injuries.

Technique:

• The patient is placed in a supine position and the following regions are scanned:

• Subxiphoid transverse view: assess for pericardial effusion and left lobe liver injuries

• Longitudinal view of the right upper quadrant: assess for right liver injuries, right kidney injury, and Morison pouch

• Longitudinal view of the left upper quadrant: assess for splenic injury and left kidney injury

• Transverse and longitudinal views of the suprapubic region: assess the bladder and pouch of Douglas

eFAST stands for extended FAST and incorporates left and right thoracic views to assess for pneumothorax and haemothorax.

Limitations of eFAST are:

• Operator-dependent

• Bowel gas and subcutaneous air distort images.

• Can miss diaphragm, bowel, and pancreatic injuries

• Unreliable for assessment of the retroperitoneum

• Cannot reliably exclude injury in penetrating trauma

• May need repeating or supplementing with other investigations



Solution to Question 8:

The percentage of burns calculated in the given patient is 45% according to Wallace's rule of nine.

Calculation:

• Burns on the head&neck would account for 9% of burns.

• Burns on both upper limbs (9% for each limb) would add to 18%

• Burns on both sides of the chest would account for 18%

Therefore the total percentage of burns in the given patient is 9+18+18 = 45%

Solution to Question 9:

FAST stands for Focussed Assessment with Sonography in Trauma.

It is a rapid, non-invasive bedside test for detecting free fluid using sonography to assess patients for potential thoracoabdominal injuries. It can be performed within 2-4 minutes.

Technique:

• The patient is placed in a supine position and the following regions are scanned:

• Subxiphoid transverse view: assess for pericardial effusion and left lobe liver injuries

• Longitudinal view of the right upper quadrant: assess for right liver injuries, right kidney injury, and Morison pouch

• Longitudinal view of the left upper quadrant: assess for splenic injury and left kidney injury

• Transverse and longitudinal views of the suprapubic region: assess the bladder and pouch of Douglas

eFAST stands for extended FAST and incorporates left and right thoracic views to assess for pneumothorax and haemothorax.

Limitations of eFAST are:

• Operator-dependent

• Bowel gas and subcutaneous air distort images.

• Can miss diaphragm, bowel, and pancreatic injuries

• Unreliable for assessment of the retroperitoneum

• Cannot reliably exclude injury in penetrating trauma

• May need repeating or supplementing with other investigations



Solution to Question 10:

Tamoxifen increases the risk of endometrial cancer by 1-2%.

The benefit of tamoxifen in breast cancer outweighs the risk of endometrial cancer.

Tamoxifen is a selective estrogen receptor modulator, approved for the treatment of estrogen receptor-positive (ER) breast cancer. It is used only in premenopausal women for 5 years in low-risk patients and for 10 years in patients with a high risk of relapse (node-positive, tumor >5 cm, lymphovascular invasion). It is also known to reduce the incidence of second primary breast cancer in the contralateral breast.

Tamoxifen increases the risk of endometrial cancer in postmenopausal women; hence aromatase inhibitors are used in postmenopausal women. Tamoxifen is also not recommended in patients who have had bilateral mastectomies with ER-positive ductal carcinoma in situ. Other side effects of tamoxifen include bone pain, hot flashes, nausea, vomiting, fluid retention and rarely

thrombosis and cataract.

Solution to Question 11:

The most appropriate mesh used in hernioplasty is a lightweight mesh with large pores.

Lightweight meshes generally contain less material and induce a less-pronounced foreign body reaction and decreased inflammatory response, which results in better tissue incorporation, increased prosthesis compliance, and decreased patient discomfort and pain.

Meshes with large pores exhibit less inflammatory infiltrate, connective tissue, and scar bridging, which allows increased soft tissue ingrowth. They have greater elasticity and flexibility and also shrink less resulting in lesser adhesions, better tissue incorporation, and decreased patient discomfort and pain. So the best mesh type among the given options is lightweight mesh with large pores.

Meshes used in hernioplasty can be synthetic or biological depending on their composition.

Synthetic meshes are made from polypropylene, polyester, or PTFE (polytetrafluoroethylene).

PTFE meshes are hydrophobic with a lower risk of adhesion.

Biological meshes are acellular matrix materials derived from humans or animals and have lesser complications. But due to poor collagen deposition and absorption, they rapidly lose mechanical strength and hence their use is restricted to infected environments.

Note: To promote better tissue integration, a composite mesh was developed which combines the attributes of polypropylene and PTFE by layering the two substances on top of one another.

POLYPROPYLENE	POLYESTER	PTFE
monofilament	multifilament	solid
hydrophobic	hydrophilic	hydrophobic
rigid	elastic	rigid

Solution to Question 12:

Lower lip paralysis occurs when there is an injury to the marginal mandibular branch of the facial nerve.

Branches of the facial nerve are:

• Temporal

• Zygomatic

• Buccal

• Marginal mandibular

• Cervical

The facial nerve passes through the parotid gland, hence it might be injured during parotidectomy.

In contrast to this, the upper lip is supplied by buccal branches of the facial nerve.

Complications of parotidectomy include hematoma formation, infection, deformity (unsightly scarring and retromandibular hollowing), temporary/permanent facial weakness, sialocele, facial numbness, permanent numbness of the earlobe due to great auricular nerve transection and Frey's syndrome.



Anatomy of Facial Nerve & Topographical Tests to Detect Lesions at Various Levels

Solution to Question 13:

Hypercalcemia is not an adverse effect of massive blood transfusion.

Massive blood transfusion instead leads to hypocalcemia as the anticoagulant citrate binds to calcium. The lethal triad of massive blood transfusion is acidosis, hypothermia, and coagulopathy.

Solution to Question 14:

APC, k-ras, and p53 gene mutations are implicated in carcinoma of the colon.

In the adenoma to carcinoma sequence, the loss of one normal copy of the tumor suppressor gene APC is the first hit, followed by the loss of the intact second copy of the APC gene (second hit). Other changes in the sequence include the mutation of k-ras and the inactivation of the tumor suppressor gene p53, leading to the emergence of carcinoma. The accumulation of these mutations is more important than their occurrence in a specific order. This is known as the Fearon-Vogelstein adenoma-carcinoma multistep model of colorectal neoplasia.

A number of other genes on chromosome 18q are implicated in colorectal cancer, including SMAD2, SMAD4, and DCC.

Solution to Question 15:

6 follicular clusters each with 10 cells are considered adequate thyroid FNAC sampling.

There are different criteria for the adequacy of thyroid FNAC:

• Six groups with at least 10 cells in each group, or

• Ten groups with at least 20 cells in each group, are considered adequate.

Cystic goitre may not yield thyroid follicular cells, hence, the aspiration should be repeated under USG guidance.

FNAC is the investigation of choice in discrete thyroid swellings; it has excellent patient compliance, simple and quick to perform in the outpatient department and is readily repeated.

Bethesda criteria for reporting thyroid cytopathology:



The 2023 Bethesda System for Reporting Thyroid Cytopathology

Bethesda class	Diagnostic category	Management
I	Nondiagnostic	Repeat FNA with ultrasound guidance
II	Benign	Clinical and ultrasound follow up
III	Atypia of undetermined significance Specify if AUS-nuclear atypia or AUS-other	Repeat FNA, molecular testing, diagnostic lobectomy, or surveillance
IV	Follicular neoplasm	Molecular testing+, hemithyroidectomy or lobectomy
	Follicular neoplasm-oncocytic follicular neoplasm*	
V	Suspicious for malignancy	Molecular testing+, lobectomy or near total thyroidectomy
VI	Malignant	Lobectomy or near total thyroidectomy

* Previously called Hurthle cell neoplasm
\+ Molecular testing can be performed to assess the type of surgery to be performed (lobectomy or total thyroidectomy)

©MARROW



Solution to Question 16:

All the given options can be used to stop esophageal variceal bleeding.

Image A shows the Sengstaken-Blakemore tube. It has 3 lumens - one aspiration port and 2 inflation balloons. They are:

• Gastric aspiration port

• Gastric balloon port

• Esophageal balloon port

The gastric balloon is inflated and tension is applied to the gastroesophageal junction. If this does not control the haemorrhage, the esophageal balloon is inflated as well, compressing the venous plexus between them. Only gastric contents can be aspirated.



Image B shows the Minnesota tube. It has 4 openings - 2 inflation tubes and 2 suction tubes i.e. both gastric and oesophageal aspiration ports are present, which allows the aspiration of both gastric and esophageal contents.

Images C and D both show an upper GI endoscope. They are used to identify the source of bleeding and once identified, sclerotherapy and endoscopic variceal banding are performed, which have been shown to control hemorrhage effectively.

Banding has a lower rate of complications. If bleeding does not stop by banding/sclerotherapy, balloon tamponade should be tried.

The protocol for the management of esophageal varices is given below.



Solution to Question 17:

The given clinical scenario of hemodynamic instability, hyper-resonance on the left side of the chest, and the X-ray findings are suggestive of tension pneumothorax. It is managed by inserting a wide-bore needle in the 5th intercostal space (ICS) followed by placing an intercostal drain in the 5th ICS.

Traumatic pneumothorax can be:

• Simple pneumothorax: Without hemodynamic instability

• Tension pneumothorax: With hemodynamic instability

The X-ray given below is showing a left-sided tension pneumothorax. A large, well-demarcated area is devoid of lung markings on the left side. The yellow arrows are pointing to the tracheal deviation and the mediastinal shift to the right side.



Management:

• Simple pneumothorax: Tube thoracostomy or chest tube insertion in the 5th ICS in the triangle of safety

• Tension pneumothorax:

• Immediate management: Urgent needle decompression or needle thoracostomy i.e. insertion of a wide-bore needle in the 5th ICS, anterior to the midaxillary line

• Definitive management: Tube thoracostomy or chest tube insertion in the 5th ICS in the triangle of safety

Note: According to ATLS 10th edition

• In adults, needle thoracostomy or needle decompression location is changed to 5th ICS anterior to the midaxillary line from 2nd ICS midclavicular line.

• In the pediatric age group, the needle thoracostomy location remains at the 2nd ICS midclavicular line.

Solution to Question 18:

Endoscopic retrograde cholangiopancreatography (ERCP) is the best investigation in a patient suspected of post-cholecystectomy biliary leaks.

Patients with post-cholecystectomy biliary leak usually present with fever, jaundice and abdominal pain. USG is used for an initial evaluation in such cases and can show a collection in the sub-hepatic space.

The most common bile leak following cholecystectomy is from the cystic duct. This can be treated by placing a biliary endoprosthesis (stent) in the common bile duct across the origin of the cystic duct.

MRCP shows fluid collections arising from bile leaks but cannot differentiate these from other causes of fluid collections.

ERCP is the best investigation as it has the advantage of being both diagnostic and therapeutic. ERCP is useful post-cholecystectomy as there is a high likelihood of a retained stone and stone extraction and/or temporary stenting may be required.

HIDA scan is the most sensitive investigation for biliary leakage.

CT scan can show focal fluid collections, ascites, biliary obstruction, hepatic atrophy, signs of secondary biliary cirrhosis and any vascular injury.



Solution to Question 19:

The given image which shows a punched-out ulcer with regular margins on the plantar surface of the dry, crackled foot is suggestive of a neuropathic ulcer.

Neuropathic ulcers occur as a result of repeated trauma to the insensitive part due to peripheral neuropathy. There is a lack of sensation over the pressure points, leading to unnoticed microtrauma, wounds or cuts which develop into an ulcer. Causes of neuropathic ulcers include diabetic neuropathy, B12 deficiency, alcoholism, hypothyroidism, leprosy, peripheral nerve injury, transverse myelitis and myelomeningocele.

Other options:

Option D: Tuberculous ulcers occur when a cold abscess from bone and joint tuberculosis breaks out on the surface. The most characteristic feature of this ulcer is its edge which is thin and undermined. The following is a picture of a tuberculous ulcer:

Type of Ulcer	Arterial(option A)	Venous(option C)	Neuropathic	Pressure
Underlying co ndition	Peripheral vascular d isease, hypertension	Varicose veins, DVT, obesity, previous thr ombophlebitis	Diabetes, peripheral neuropathy	Limited mobility

Type of Ulcer	Arterial(option A)	Venous(option C)	Neuropathic	Pressure
Location	Lateral malleolus, toes, and dorsum of the foot	Lower leg, above the medial malleolus	The plantar surface of the foot, toes	Bony prominences (e.g.heel)
Characteristics	Deep, punched-out, irregular margins. The wound bed has unhealthy necrotic tissue	Shallow, flat with exudate	Punched-out with regular calloused margins.	Deep, macerated
Condition of leg/foot	Cool thin shiny skin, pallor on leg elevation, absent peripheral pulses, and delayed capillary refill	Excematous and pigmented. Limb edema may be present	Dry, cracked, insensate	Atrophic skin, loss of muscle

The following is a picture of an arterial ulcer.

The following is a picture of a venous ulcer.

The following is a picture of a pressure ulcer.

Solution to Question 20:

Colorectal carcinoma is the most likely primary tumor for the given metastases. Among the given options, cytokeratin 7 is negative only in colon carcinoma.

The most common site of distant metastasis of colorectal cancer is the liver. These metastases arise from hematogenous spread via the portal venous system. The lung is the next most common site for metastasis.

Cytokeratin profiles of various primary tumors:

CK 7 positiveCK20 positive	CholangiocarcinomaOvarian mucinous carcinoma
CK 7positiveCK 20 negative	Breast carcinomaOvarian serous adenocarcinomaThyroid carcinomaPancreatic adenocarcinomaLung carcinoma
CK 7negativeCK 20positive	Colorectal adenocarcinomaMerkel cell carcinoma
CK 7 negativeCK 20 negative	Hepatocellular carcinomaRenal cell carcinomaProstatic adenocarcinomaSquamous cell carcinoma

Solution to Question 21:

Chronic smoker presenting with bluish discoloration of the toe along with intermittent claudication is suggestive of peripheral arterial disease and duplex scanning will be the next appropriate investigation.

Duplex Scanning is a major non-invasive technique that uses B-mode ultrasound to provide an image of the vessel. Most scanners now have color coding, which allows detailed visualization of blood flow in a vessel. Different colors indicate changes in the direction and velocity of flow, with areas of high flow usually indicating stenotic lesions.

Other options:

Option A: CT Angiography (CTA) is more useful in the evaluation of visceral arterial disease (eg. mesenteric ischemia). It can be beneficial in peripheral artery diseases where a duplex scan shows poor images (aortoiliac segment involvement). CTA is not routinely recommended in peripheral artery disease because of concomitant renal dysfunction in these patients which can lead to contrast-induced nephropathy.

Option B: MR Angiography is time-consuming and expensive, hence not done as the next investigation. It is contraindicated in patients with claustrophobia and metallic implants.

Option C: Digital subtraction angiography (DSA) is an invasive and expensive procedure compared to other investigations. The benefits of DSA are that it provides dynamic arterial flow information and can be combined with a definitive endovascular intervention when indicated. DSA is indicated only when there is a discrepancy in diagnosis or intervention is planned. Complications associated with DSA include bleeding, hematoma, false aneurysm formation, thrombosis, arterial dissection, distal embolisation, renal dysfunction, and allergic reaction.



Solution to Question 22:

In the given scenario, the rapid onset of the disease and the examination findings are suggestive of inflammatory carcinoma of the breast.

Inflammatory breast cancer is the most aggressive subtype of breast cancer. It mimics acute mastitis. The hallmark of inflammatory breast carcinoma is the diffuse tumor involvement of the dermal lymphatics within the breast and the overlying skin, often without a discrete underlying tumor mass (peau d' orange appearance). The patient presents with rapidly progressive swelling of the breast, warmth, erythema and edema. A breast lump may or may not be present.

A diagnosis of inflammatory breast carcinoma should be suspected when a patient with rapidly progressive swelling with inflammatory skin changes does not respond to antibiotics. The imaging features with ultrasound and a mammograph is non-specific due to the underlying erythema and edema. A biopsy should be done to confirm the diagnosis.

Treatment includes combined-modality treatment:

• Neoadjuvant chemotherapy

• Mastectomy

• Radiation therapy

• Hormonal therapy- for ER-positive tumors

• Trastuzumab for HER-2-positive tumors

Option B: Giant fibroadenomas occasionally occur during puberty. These are benign solid encapsulated slow-growing tumors over 5 cm in diameter. It presents as a painless lump in the breast. They are managed by excision via a submammary incision.

Option C: Phyllodes tumors usually occur in women over the age of 40 years but can appear in younger women. Phyllodes tumor is mostly benign even though it is also called cystosarcoma phyllodes. It presents as a large tumor with an unevenly bosselated surface. They are usually

mobile over the chest wall despite their large size. This tumor does not infiltrate the skin. Nipple retraction is not seen in this condition. Benign lesions can be treated with enucleation or wide local excision. Massive tumours, recurrent tumours and malignant lesions are treated with a mastectomy.

Option D: Galactocele are the most common benign breast lesions in lactating women. It is a cystic enlargement of the mammary gland containing milk. They are mostly detected during lactation or in the third trimester of pregnancy but rarely, may also present after stopping breastfeeding. Clinically, the mass is usually firm and non-tender and presents as a solid tumour-like lesion on physical examination. The patient usually notes the lesion during lactation or some period after lactation. Needle aspiration, followed by excision should it recur, is the standard method of treating galactocele.

Solution to Question 23:

The given clinical scenario and the barium swallow showing a bird beak appearance are suggestive of achalasia cardia.

Achalasia is a primary esophageal motility disorder characterized by the absence of esophageal peristalsis and impaired relaxation of the lower esophageal sphincter (LES) in response to swallowing.

Manometry is considered the most sensitive investigation to diagnose achalasia as it identifies early disease before esophageal dilatation and food retention. The diagnostic criteria for achalasia with manometry are impaired lower esophageal sphincter (LES) relaxation and absent peristalsis with an elevated LES pressure.

Endoscopy is useful for ruling out pseudo-achalasia.

Note: 24-hour pH monitoring is ideal and most accurate in patients with the features of gastroesophageal reflux disease.

Solution to Question 24:

The given clinical scenario of a patient with sudden onset severe abdominal pain which is out-of-proportion to the physical examination findings along with atrial fibrillation is suggestive of acute mesenteric ischemia.

Mesenteric ischemia may be acute or chronic. The superior mesenteric artery (SMA) is most likely affected. An embolus is the most common cause of SMA occlusion. Possible sources for SMA emboisation include:

• Left atrium in atrial fibrillation

• Previous history of ischemic heart disease/myocardial infarction

• Vegetations on mitral and aortic valves associated with endocarditis

• An atheromatous plaque from an aortic aneurysm.

Other causes of mesenteric ischemia include:

• Atherosclerosis of mesenteric vessels

• Vasculitides

• Venous thrombosis in factor V Leiden mutation

• Non-occlusive mesenteric ischemia due to alterations in splanchnic blood flow.

The most important clue to an early diagnosis of acute mesenteric ischemia is the sudden onset of severe abdominal pain in a patient with atrial fibrillation or atherosclerosis. The pain is typically in the central abdomen and out of proportion to the physical findings.

Associated clinical features include nausea, vomiting, diarrhea, abdominal distension, peritonitis, and the passage of bloody stools.

The gold standard for the diagnosis of acute mesenteric ischemia is angiography, and management is laparotomy. The goal of operative exploration is to resect the compromised bowel and restore blood supply via embolectomy.

Other options:

Option B: A pulsatile abdominal mass is seen in the case of an abdominal aortic aneurysm. Acute pain and hypotension occur with rupture of the aneurysm, which requires an emergency operation.

Options A and D: Acute severe pancreatitis and peptic ulcer perforation have no association with atrial fibrillation. They usually present with pain over the upper abdomen associated with guarding and rigidity on examination.

Solution to Question 25:

A lateral neck mass in a pediatric patient, which is non-tender and present in association with freckles, and hyperpigmented skin patches (cafe-au-lait spots) is suggestive of neural tumor i.e. neurofibroma.

Neurofibroma is a benign tumor arising from the connective tissue of the nerve containing ectodermal, neural and mesodermal connective tissue components. It can be single or multiple. It is the most common intradural extramedullary spinal tumor. Neurofibromas can be local, generalized (von Recklinghausen's disease) or plexiform types. Patients present with mild pain or a painless swelling usually in the subcutaneous or cutaneous plane with tingling, numbness and paraesthesia along the distribution of the nerve. Complications of neurofibroma include cystic degeneration, erosion into deeper planes, cord compression and paralysis (spinal neurofibroma) and sarcomatous changes.

Treatment is excision under general or regional anesthesia. Indications for surgery in neurofibroma include symptomatic patients (pain and pressure symptoms), cosmetic problems and malignant transformation (5%).

Option A: Cystic hygroma is a type of congenital lymphangioma. This swelling is soft and partially compressible, which shows positive fluctuation and translucency test.

Option B: Chemodectoma or carotid body tumor is a rare tumor, most commonly present in the fifth decade and there is an association with pheochromocytoma. These patients usually present with a slowly enlarging, painless lump at the carotid bifurcation.

Option D: A lipoma is a soft painless swelling. It shows the characteristic slip sign and is freely mobile.

Solution to Question 26:

The given cannula is 20 gauge IV cannula as its color code is pink.

Solution to Question 27:

The above image shows a reverse Trendelenburg position, where the patient's head is elevated 15 - 30 degrees higher than the feet. It is named so, as the position is reverse of the Trendelenburg position, where the head end is lower and the feet are at a higher level.

The Trendelenburg and reverse Trendelenburg positions are used in surgeries that are required to move the abdominal viscera, such as the small bowel, away from the operative field. Restraints and sideboards are placed to prevent the patient from sliding from the bed.

In laparoscopic cholecystectomy, the patient is placed in a reverse Trendelenburg position slightly rotated to the left. This exposes the fundus of the gallbladder which is retracted towards the diaphragm.

Other options:

Option A: Lloyd Davies's position is a 30-degree Trendelenburg position with the hips flexed at 15 degrees. It is used for rectal and anal canal surgeries like low anterior resection and abdominoperineal resection. It can cause compartment syndrome of the calves.



Option B: Jackknife position is a modified prone position where the patient's hips are supported on a pillow and the table is flexed at a 90-degree angle with the head and feet at a lower level. It is not used anymore due to the chest compression causing impaired ventilation leading to positional asphyxia. It was used for hemorrhoids and fissure surgery.



The image below shows commonly used patient positions in the operating room:



Solution to Question 28:

Increased TSH and low T4 levels are seen in primary hypothyroidism. Hashimoto's thyroiditis is the only cause of primary hypothyroidism in the given options.

Hashimoto's thyroiditis is an autoimmune disease that results in the destruction of the thyroid gland and gradual & progressive thyroid failure. It is the most common cause of primary hypothyroidism in iodine-sufficient areas. They also have autoantibodies against thyroid receptors, thyroid peroxidase and thyroglobulin.

Other options:

Option A: Hypoparathyroidism is generally not associated with hypothyroidism, except in certain congenital multisystem disorders such as DiGeorge's syndrome.

Option C: Grave's disease is the most common cause of hyperthyroidism. It is an autoimmune disorder caused due to autoantibodies against the TSH receptor. There will be elevated T3 and T4 levels with low TSH levels.

Option B: A TSH-producing pituitary adenoma can have increased TSH levels, but there will be elevated T3, T4 levels as well.

Solution to Question 29:

e-FAST is used to additionally assess the presence of free fluid in the thoracic cavity.

FAST stands for focused assessment with sonography for trauma. It is a rapid diagnostic examination to assess patients with potential thoracoabdominal injuries. It can be performed within 2-4 minutes.

The sequential order in which FAST is performed: pericardial sac, the right upper quadrant (RUQ), left upper quadrant (LUQ), and pelvis. 4 'Ps' are evaluated in sequence: Pericardial sac → Perihepatic region → Perisplenic region → Pelvis.

Technique:

• Following regions are scanned:

• Subxiphoid: assess for pericardial effusion and left lobe liver injuries

• Right upper quadrant: assess for right liver injuries, right kidney injury, and Morison pouch

• Left upper quadrant: assess for splenic injury and left kidney injury

• Suprapubic region: assess the bladder and pouch of Douglas

An 'extended FAST or eFAST' scan may be performed: left and right thoracic views to assess for pneumothorax and hemothorax.

Solution to Question 30:

Injury to the right aberrant posterior sectoral duct associated with bile leak is classified as Strasberg's type C injury.

Biliary injury may occur post-laparoscopic cholecystectomy. The most common site of bile leak following cholecystectomy is the cystic duct.

MRCP is the gold standard for complete morphological evaluation of the biliary tree. MRCP with magnetic resonance angiography is more useful in cases of associated vascular injury. A HIDA scan can confirm the presence of a bile leak or biliary obstruction.

The Strasberg classification of bile duct injury is widely used to anatomically define bile duct injuries.

Classification

• Type A: Bile leak from cystic duct stump or a minor biliary radical in the gallbladder fossa
• Type B: Occluded right posterior sectoral duct.
• Type C: Bile leak from the divided right posterior sectoral duct.
• Type D: Bile leak from the main bile duct without major tissue loss.
• Type E: Injury to the main hepatic duct; classified according to the level of injury
• E1: Transected main bile duct with a stricture more than 2cm from the hilus.
• E2: Transected main bile duct with a stricture less than 2cm from the hilus
• E3: Stricture of the hilus with right and left ducts in communication.
• E4: Stricture of the hilus with separation of right and left hepatic ducts.
• E5: Stricture of the right aberrant sectoral and main bile ducts.



Management of bile duct injuries is shown below:



Solution to Question 31:

According to CEAP classification, varicose veins with skin eczema and lipodermatosclerosis come under the clinical class of C4.

The clinical class of C4 under CEAP includes changes in skin and subcutaneous tissue secondary to chronic venous disease. It is further classified as,

• C4a- pigmentation or eczema

• C4b- lipodermatosclerosis or atrophie blanche

• C4c- corona phlebectatica

Recent 2020 update on the CEAP system:

• C2r for recurrent varicose veins

• C4c for Corona phlebectatica

• C6r for recurrent active venous ulcer.

Solution to Question 32:

Liposarcoma is the most commonly seen tumor in retroperitoneal space.

Retroperitoneal tumors can be benign or malignant. Malignant retroperitoneal tumors may be mesodermal (75%), neuroectodermal (24%), or embryonic remnant (1%) in origin. Rarely it can be urologic or metastatic also. The order of frequency of primary retroperitoneal malignancy is as follows:

• Liposarcoma

• Leiomyosarcoma

• Rhabdomyosarcoma

• Fibrosarcoma

• Malignant nerve sheath tumors

Liposarcoma is composed of three histologic varieties (listed in order of decreasing frequency):

• Well-differentiated and undifferentiated liposarcoma

• Pleomorphic liposarcoma

• Myxoid/round cell liposarcoma

Among the three subtypes, pleomorphic liposarcoma is the most malignant and often metastasizes, while the well-differentiated is the least malignant. Myxoid liposarcoma has intermediate malignant behavior.

Diagnosis is confirmed by contrast-enhanced CT with CT of the chest to look for metastases.

Surgery is the main modality of treatment. Wide excision with complete en bloc dissection is done. Post-operative radiotherapy and chemotherapy are used as adjuvant methods. Radiotherapy is given at a dose of 5000 cGY, it can be external beam or intra-operative radiotherapy. Chemotherapeutic agents used are mesna, adriamycin, ifosfamide and dacarbazine (MAID regimen).

Solution to Question 33:

The given clinical scenario is suggestive of breast malignancy with locoregional spread with unlikely distant metastasis. For further evaluation of this lesion, bilateral mammography, a biopsy of the breast mass and a USG-guided biopsy of the suspicious lymph node need to be performed in order to determine staging and decide on further management.

Statement 2: PET Scan is usually performed in advanced cases of breast cancer. It is not routinely used in the assessment of locoregional disease.

Statement 1: Diagnostic mammography is used to evaluate women with abnormal findings such as a breast mass or nipple discharge. Mammography is more accurate than clinical examination for the detection of early breast cancers. Mammographic features suggestive of breast cancer include a solid mass with or without stellate features, microcalcifications, irregular soft tissue shadow, spiculations and duct distortion.

Statement 3: Pathological diagnosis of carcinoma is made by a core needle biopsy (tru-cut biopsy) or fine-needle aspiration before any type of treatment is started.

Statement 4: The presence or absence of axillary lymph node involvement in breast cancer is one of the most important prognostic factors for determining staging and treatment. Ultrasound-guided lymph node biopsy should be done in patients with suspicious or clinically palpable lymph nodes.

Ultrasound features suggestive of breast cancer include a hypoechoic, irregular mass with indistinct, angulated or spiculated margins, acoustic shadowing, ductal extension and a branched pattern of mass.

Solution to Question 34:

According to the European Hernia Society classification, M3 incisional hernia refers to umbilical hernia.

According to the European Hernia Society classification of incisional hernias:

Midline incisional hernias are classified as-

Lateral incisional hernias are classified as-

M1	Subxiphoidal
M2	Epigastric

M1	Subxiphoidal
M3	Umbilical
M4	Infraumbilical
M5	Suprapubic

L1	Subcostal
L2	Flank
L3	Iliac
L4	Lumbar

Solution to Question 35:

Tillaux's sign is said to be positive if the mass/swelling is mobile perpendicular to the mesentery attachment line. Tillaux’s sign is seen in a mesenteric cyst.

Mesenteric cysts occur either in the small intestine (60%) or the colon (40%) and can be classified as the following:

• Chylolymphatic (most common type)

• Enterogenous

• Urogenital remnant

• Dermoid

• Other causes: traumatic hematoma and cyst formation, tuberculous cold abscess of mesentery and hydatid cyst of the mesentery

They most commonly occur in adults with a mean age of 45 years. It is more common in females than males (2:1). The most common presentation is a painless abdominal swelling near the umbilicus, which is smooth and not moving with respiration with Tillaux's triad:

• Soft, smooth swelling in the umbilical region

• Freely mobile in a direction perpendicular to the mesentery (Tillaux's sign)

• Zone of resonance around the cyst

Other presentations include recurrent attacks of abdominal pain due to temporary impaction of a food bolus in a segment of the bowel narrowed by the cyst and vomiting.

Complications of mesenteric cyst include torsion, rupture, infection and hemorrhage into the cyst.

Ultrasonography and CT are used to demonstrate mesenteric cysts.

Treatment:

• Chlyolymphatic cyst is managed by enucleation.

• Enterogenous cyst is managed by removal of the cyst with resection of the adjacent bowel as it arises as a diverticulum or duplication from the adjacent bowel.

In cases where complete excision is not possible due to the close proximity to vital structures, partial excision or marsupialization should be performed.



Solution to Question 36:

A-1, B-2, C-3, and D-4 is the correct option.

A: During breast surgery, removal of the axillary lymph nodes is commonly done. This carries a risk of injury to the long thoracic nerve, which originates from the C5-C7 roots of the brachial plexus. It innervates the serratus anterior muscle. Injury to this nerve during surgery results in paralysis of the serratus anterior muscle leading to winging of the scapula.

B: The external laryngeal nerve is a motor nerve that supplies the cricothyroid muscle. The close relationship of the external laryngeal nerve to the superior thyroid artery puts the nerve at potential risk when the artery is clamped during thyroid surgeries.

C: Excision of the submandibular gland results in potential risk to 3 cranial nerves:

• Marginal mandibular branch of the facial nerve

• Lingual nerve

• Hypoglossal nerve

D: The vagus nerve is at greater risk of injury during surgeries like thymectomy.

Solution to Question 37:

Patients with congenital hypertrophic pyloric stenosis (CHPS) will present with non-bilious projectile vomiting and visible peristalsis observed moving from left to right (not from right to left). Metabolic abnormalities include hypokalemia, hypochloremia and metabolic alkalosis.

CHPS occurs due to hypertrophy of the circular muscle of the pylorus, which results in constriction and obstruction of the gastric outlet.

• Patients commonly present between 2 and 8 weeks of age.

• First-born male infants are at the highest risk.

• Infants usually present with projectile, non-bilious vomiting.

• Clinically, an olive-shaped mass is palpable in the right upper quadrant or epigastrium. When the infant is given a test feed, visible peristalsis is usually observed moving from left to right (not from right to left), followed by projectile vomiting.

Metabolic abnormality:

• Infants with hypertrophic pyloric stenosis develop a characteristic metabolic abnormality i.e. hypokalemic hypochloremic metabolic alkalosis with paradoxical aciduria.

• Persistent vomiting results in loss of hydrochloric acid, leading to metabolic alkalosis (due to loss of H^+) and hypochloremia (due to loss of Cl^-).

• Kidneys try to compensate for alkalosis by secreting Na^+ and HCO_3^-. However, sodium loss stimulates aldosterone secretion through the RAAS pathway.

• Aldosterone causes active reabsorption of sodium in exchange for K^+ and H^+ ions leading to the development of hypokalemia and paradoxical aciduria, since the body initially loses H^+ ions due to the loss of HCl, and instead of compensating for that loss, the body is losing more H^+ ions through urine.

Management:

• The child should be rehydrated and the metabolic abnormality must be corrected before surgery.

• Ramstedt pyloromyotomy remains the standard surgical procedure for its treatment.

Solution to Question 38:

The quick Sequential Organ Failure Assessment (qSOFA) scoring system does not rely on the heart rate.

It is a bedside scoring system that may identify patients with suspected infection, who are at greater risk for sepsis outside the intensive care unit (ICU).

It uses three criteria, assigning one point each:

• Low blood pressure (SBP≤100 mmHg)

• High respiratory rate (≥22 breaths per min)

• Altered mental status

The qSOFA suggests potentially life-threatening sepsis when at least two of the parameters are met.

Note :

The primary method to evaluate the risk of organ dysfunction (sepsis) in patients with suspected infection is by determining the sequential (sepsis-related) organ failure assessment (SOFA) score; which includes an assessment of:

• Circulation - Systolic BP & need for vasopressors; platelet count

• Respiration - PaO2/FiO2 & need for mechanical ventilation

• Mentation - Glasgow Coma Score

• Hepatic function - Bilirubin

• Renal function - Creatinine

Organ dysfunction is identified by a SOFA score of ≥ 2

The SOFA score is limited by its requirement of multiple laboratory tests which makes it complex and time-consuming. In order to simplify the scoring process, the "Sepsis -3" task force recommended the quick sepsis-related organ dysfunction assessment "qSOFA" score.

Solution to Question 39:

The medial border of the triangle of doom is bounded by vas deferens and the medial border of the myopectineal orifice is bounded by the lateral border of the rectus abdominis.

The triangle of doom is bounded

• Medially by the vas deferens

• Laterally by the vessels of the spermatic cord (gonadal vessels)

• Inferiorly by peritoneal reflection

The myopectineal orifice of Fruchaud is an osseo-myo-aponeurotic tunnel through which all groin hernias occur.

It is bounded:

• Medially by the lateral border of the rectus sheath.

• Laterally by the iliopsoas muscle.

• Above by the arched fibers of the internal oblique and transversus abdominis muscle.

• Below by the pectin pubis and fascia covering it.



Solution to Question 40:

A-3, B-1, C-2, and D-4 is the correct option.

A: Uro bags are used for the collection of urine. They are usually attached to Foley's catheter, which is placed to drain the bladder (commonly in postoperative patients).

B: Nasogastric tube/Ryle's tube is used for nasogastric feeding/stomach wash/aspiration.

C: An underwater seal bag is used for the collection of drainage fluid from the thoracic cavity. It acts as a one-way valve allowing air to leave the pleural cavity, but it cannot be drawn back in by the negative pressure that is created in the intrathoracic cavity.

D: A metallic suction cannula is used to clear the operative field by suctioning secretions/blood and aspirating the fluid from various cysts and in a procedure like a hydrocele.

Solution to Question 41:

The given clinical scenario and erect X-ray abdomen showing gas under the diaphragm are suggestive of pneumoperitoneum, which is managed by initial resuscitation with IV fluids followed by exploratory laparotomy.

Pneumoperitoneum is the presence of air or gas in the peritoneal cavity. The most common cause of pneumoperitoneum is a perforation of a hollow viscus. Other causes of pneumoperitoneum in adults include peritoneal dialysis, and mechanical ventilation, after procedures like colonoscopy and endoscopy. In children, it can be caused by necrotizing enterocolitis, hirschsprung's disease, meconium ileus and mechanical ventilation.

The patient presents with abdominal pain and distension, vomiting, fever, constipation, tachycardia, tachypnea, hypotension, and reduced urine output. Per-abdomen examination can show guarding and rigidity.

An erect X-ray abdomen is highly sensitive for the detection of free intraperitoneal gas. Some of the signs seen in the X-ray include:

• Cupola sign is the presence of air underneath the diaphragm, giving a dome-like appearance.

• Rigler's sign is where both bowel surfaces are seen together as a sharp margin.

• Falciform ligament sign is the outlining of the falciform ligament with free abdominal gas.

• Liver/gallbladder edge sign is where the sharp margins of the liver/gallbladder are visible due to the collection of air.

Management includes,

• Adequate preoperative resuscitation with intravenous fluids followed by correction of electrolyte imbalance, instituting broad-spectrum intravenous antibiotics and

• Early surgical intervention/laparotomy to address the cause of pneumoperitoneum and prevent further contamination of the peritoneal cavity.

Surgery NEET 2018

Question 1:

Which of the following are indications for surgical intervention in abdominal aortic aneurysms?

a) Size more than 5.5 cm in males

b) Size more than 5 cm in females

c) Expansion of >0.5 cm within 6 months

d) All the above

Question 2:

The following image shows the retrograde ureteropyelogram of a patient with loin pain. What is the most probable clinical diagnosis?



a) Hydronephrosis

b) Duplex kidney

c) Renal carcinoma

d) Renal stone

Question 3:

Which of the following types of renal stones are produced by laxative abuse?

a) Uric acid

b) Ammonium urate

c) Struvite

d) Calcium oxalate

Question 4:

A patient is confused, opens his eyes on painful stimuli, flexes one limb on painful stimuli, and can localize the stimulus with the other limb. His Glasgow Coma Score is _____

a) 8

b) 9

c) 10

d) 11

Question 5:

A patient presents with a painless swelling of the neck. The histology of the tissue is as given in the image below. What is the diagnosis?



a) Medullary carcinoma

b) Non Hodgkin's lymphoma

c) Grave's disease

d) Hashimoto's thyroiditis

Question 6:

A 45-year-old male patient who presented with a scrotal swelling underwent an orchidectomy. The image below shows the histopathology findings. What is your diagnosis?



a) Seminoma

b) Yolk-sac tumor

c) Embryonal cell carcinoma

d) Teratoma

Question 7:

A surgeon sees a necrotic bowel when operating on a child for an inguinal hernia. What is true about the resection and anastomosis of the bowel?

a) The surgeon must use chromic catgut

b) The surgeon must not include the submucosa in the sutures

c) The surgeon must use the single-layer, extramucosal suture technique

d) The surgeon must use the single-layer seromuscular suture technique

Question 8:

A lady comes with a reddish firm nodule on her chest which hasn't grown in size. Its histolopathological image is shown below. What is the diagnosis?



a) Lipoma

b) Fibroadenoma

c) Nevus

d) Hemangioma

Question 9:

Which is the most common organism causing acute bacterial prostatitis?

a) Enterococcus

b) Proteus

c) Streptococcus Agalacticia

d) E.coli

Question 10:

A 35-year-old man came with complaints of dark-colored stools to the OPD. After undergoing endoscopy, the patient was diagnosed with an upper GI bleed. Which of the following structures is used as a demarcation point to differentiate it from a lower GI bleed?

a) Ampulla of Vater

b) Ligament of Treitz

c) Superior duodenal flexure

d) Ileocaecal junction

Question 11:

Which is the best graft for a femoropopliteal bypass?

a) Dacron

b) Great saphenous vein

c) Polytetrafluoroethylene

d) Short saphenous vein

Question 12:

What is the most common cause of chronic pancreatitis?

a) Gallstones

b) Alcohol

c) Autoimmune

d) Drugs

Question 13:

A probe is inserted into the duct from which the nipple discharge is emerging. A tennis racquet incision is made and the flap is dissected to reach the duct. The duct is then excised. This procedure is known as?

a) Macrodochectomy

b) Microdochectomy

c) Cone excision

d) Wide excision of the lactiferous duct.

Question 14:

Which of the following is true about chromic catgut?

a) It is made from cat gut

b) It is made from rabbit gut

c) It is absorbed by enzymatic degradation and macrophage phagocytosis

d) It is tanned with nickel salts to improve handling and to resist degradation in tissue

Question 15:

Identify the instrument given in the image below:

a) Deaver's retractor

b) Morris retractor

c) Czerney's retractor

d) Double hook retractor

Question 16:

Extra mammary Paget's disease is commonly seen in the:

a) Vulva

b) Vagina

c) Uterus

d) Cervix

Question 17:

The best method to obtain tissue for the diagnosis of pancreatic carcinoma is:

a) Transgastric FNAC guided by endoscopic ultrasound

b) MRI-guided biopsy

c) Laparoscopic-guided biopsy

d) Percutaneous transperitoneal biopsy

Question 18:

An abdominal mass in a patient with congenital hypertrophic pyloric stenosis is best seen _____.

a) During feeding

b) Soon after birth

c) During palpation over the left hypochondrium

d) During palpation over the epigastrium

Question 19:

While performing a fasciotomy for compartment syndrome, the following layers are incised:

a) Skin and Subcutaneous tissue

b) Skin, Subcutaneous tissue, Superficial fascia

c) Skin, Subcutaneous tissue, Superficial fascia and Deep fascia

d) Skin, Subcutaneous tissue, Superficial fascia, Deep fascia and a few muscular fibres

Question 20:

Which of the following statements is true about Crohn's disease?

a) For effective treatment of a Crohn's mass, only antibiotics are prescribed.

b) Symptomatic relapse after surgery is not related to the presence of disease during resection.

c) The proximal ileum is most commonly involved in Crohn's disease.

d) On endoscopic examination, continuous lesions are visualised, which are diagnostic of Crohn's disease.

Question 21:

Which of the following is the earliest sign in Crohn's disease?

a) Cobblestone appearance

b) Deep fissured ulcers

c) Rose-thorn appearance

d) Aphthous ulcers

Question 22:

The Van Nuys grading system for ductal carcinoma in situ does not include:

a) Age of the patient

b) Size of the tumor

c) Hormone receptor staining

d) Presence of microcalcification

Question 23:

Which of the following statements are true with regards to a keloid scar?

a) The extent does not cross the wound margins

b) Wide excision of the keloid scar prevents recurrence

c) Histopathology of a keloid scar shows reduced collagen and increased vascularity

d) A keloid scar forms due to increase in the level of growth factors

Question 24:

Heller's operation is done for _____.

a) Achalasia cardia

b) Carcinoma esophagus

c) Esophageal stricture

d) Hypertrophic pyloric stenosis

Question 25:

Which of the following is not true about Barrett's esophagus?

a) It is the intestinal metaplasia of esophagus

b) Mucus secreting foveolar cells are characteristic

c) Alician blue is used in the staining of biopsy specimen

d) It is a reversible cell adaptation

Question 26:

Modified shock index is calculated using?

a) HR/SBP

b) HR/DBP

c) HR/MAP

d) PR/SBP

Answer Key

Question No.	Correct Option
1	d
2	a
3	b
4	d
5	d
6	a
7	c
8	d
9	d
10	b
11	b
12	b
13	b
14	c
15	b
16	a
17	a
18	a

19	c
20	b
21	d
22	c
23	d
24	a
25	b
26	c

Detailed Explanations

Solution to Question 1:

All the given are indications for surgical intervention in abdominal aortic aneurysm (AAA). The indications include:

• Size >5.5 cm in males (option A)

• Size >4.5-5 cm in females (option B)

• Expansion of >0.5 cm within 6 months or >1 cm in one year (option C)

• Painful, tender aneurysm

• Thrombosed aneurysm, aneurysm with distal emboli

The rupture risk is high for aneurysms >5.5 cm and begins to rise exponentially with an increase in size. Women are at higher risk for rupture than men, with a lower threshold of 4.5 to 5.0 cm (depending on the reference used).

Screening is done for:

• Cardiovascular patients in men and women (60-85 years)

• Men and women >50 years with positive family history

• Annual screening in asymptomatic patients with 4-4.5 cm size, size >4.5 cm once in every 6 months.

Management of AAA can be done by open repair or endovascular aneurysm repair (EVAR). About 75% of infrarenal aneurysms are suitable for endovascular aneurysm repair (EVAR), depending on the morphology of the aneurysm assessed by a CT scan.

Aneurysms that fall below the mentioned indications may safely be followed with CT or ultrasound at 6-month intervals since long-term outcomes to routine follow-up are equivalent to earlier surgical repair of the aneurysm. Smoking cessation is very important and reduces the mortality rate.

The technique of EVAR:



Solution to Question 2:

The given image showing dilated clubbed calyces with stricture of the left ureteropelvic junction points to a diagnosis of hydronephrosis.

Hydronephrosis refers to the dilatation of the pelvicalyceal system caused by a partial or complete blockage of urinary flow.

Investigations:

• Ultrasound is the least invasive technique for diagnosing hydronephrosis.

• Retrograde pyelography is rarely indicated but will confirm the site of obstruction before corrective surgery.

• Intravenous urogram (IVU) is helpful only if the obstructed kidney is functional. In IVU, the extrarenal pelvis is dilated and the minor calyces lose their normal cupping. In early stages, there is blunting of calyces followed by its clubbing in the late stages.

• A renal isotope scan (DTPA or MAG-3) is the best modality to confirm the obstructive dilatation of the collecting system.

Solution to Question 3:

Ammonium urate stones are most likely to be seen in patients with laxative abuse.

Loss of potassium and sodium in laxative abuse causes intracellular acidosis, which increases the production of ammonia in urine. Dehydration and sodium loss further exacerbate hypokalemia and reduces the urinary volume, resulting in hyperuricemia. Highly concentrated uric acid and ammonia promote the formation of ammonium urate stones.

These are also associated with inflammatory bowel disease, chronic diarrhea, ileostomy, recurrent urinary tract infection and recurrent uric acid stone formation.

The below image shows an ammonium urate stone.



Solution to Question 4:

The GCS in this patient is 11.

• The score for verbal response: 4, because he is confused.

• The score for eye-opening: 2, because his eyes are opening to only painful stimuli.

• The score for best motor response: 5, because he can localize pain.

Therefore, the GCS score is (4+2+5) = 11.

Solution to Question 5:

The slide shows the histopathology of Hashimoto thyroiditis, with:

• Follicular destruction

• Reduced colloid with increased interstitial connective tissue

• Small lymphocyte and plasma cell infiltration

• A well-developed germinal centre may be seen

• Hurthle (Askanazy) cells that have abundant eosinophilic granular cytoplasm lining the follicles



Hashimoto's thyroiditis is the most common cause of hypothyroidism in iodine-sufficient areas. It is also known as struma lymphomatosa or lymphocytic thyroiditis. It is an autoimmune disease that results in the destruction of the thyroid gland and gradual & progressive thyroid failure.

Clinical features:

• Most prevalent between 45-65 years of age

• More common in females

• Painless enlargement of the thyroid

• Hashitoxicosis: When the auto-antibodies attack and damage the follicles, there is a sudden release of thyroid hormones from these follicles. This leads to a transient phase of thyrotoxicosis.

• Hypothyroidism: Since the follicles are damaged and cannot form the hormones anymore, these patients tend to be in a state of hypothyroidism.

Diagnosis:

• T3 and T4 are low

• TSH is raised

• Autoantibodies against:

• Thyroid receptors

• Thyroid peroxidase (TPO)

• Thyroglobulin

These patients are managed with thyroid hormone replacement therapy.

In patients with long-standing goiter, there is a risk of compressive features and the development of thyroid lymphoma. Such cases can be treated with total thyroidectomy. It may also predispose to papillary thyroid carcinoma.

Other options:

Option A: Histological features of medullary carcinoma of the thyroid include polygonal to spindle-shaped cells in the form of nests, trabeculae or follicles. Small, round anaplastic cells may be found in some cases and amyloid deposits derived from calcitonin polypeptides are present in the stroma.

The given slide below shows medullary carcinoma of thyroid:



Option C: Grave's disease is an autoimmune disorder characterized by hyperthyroidism due to circulating autoantibodies. The thyroid-stimulating immunoglobulins bind to the TSH receptors, causing overgrowth of the thyroid gland and increased thyroid hormone synthesis. Histologically, there is acinar cell hypertrophy and hyperplasia with the absence of normal colloid in the tall columnar epithelium (normally the epithelium is flat with colloid). Lymphoid infiltrates, consisting predominantly of T cells, along with scattered B cells and mature plasma cells, are present throughout the interstitium. Germinal centers are common.

The given slide below shows Grave's disease:



Solution to Question 6:

Based on the gross and microscopic appearance of the specimen, the diagnosis is seminoma.

The gross specimen shows a well-circumscribed, pale, gray-white, fleshy, homogenous mass, with a lobulated cut surface, usually devoid of hemorrhage or necrosis. Histopathology shows uniform tumor cells with abundant clear cytoplasm, distinct cell borders, and large central nuclei. There is a sparse lymphocytic infiltrate. Classically, they have a fried egg appearance.

Other options:

Option B: Yolk-sac tumor has a yellowish-white, mucinous appearance. Histologically, they are typically composed of a lace-like (reticular) network of medium-sized cuboidal, flattened, or spindled cells, which generally show less cytologic atypia than embryonal carcinoma. In approximately 50% of tumors, structures resembling endodermal sinuses (Schiller-Duval bodies) may be seen; these consist of a mesodermal core with a central capillary and a visceral and parietal layer of cells resembling primitive glomeruli.



Option C: Embryonal cell carcinoma appears variegated with foci of hemorrhage or necrosis. Histologically, the cells grow in alveolar or tubular patterns, sometimes with papillary folds. More undifferentiated lesions may display sheets of cells with cleft-like spaces. The neoplastic cells have an epithelial appearance, are large and anaplastic, and have hyperchromatic nuclei with prominent nucleoli. The cell borders are usually indistinct, and there is considerable variation in cell and nuclear size and shape (pleomorphism). Mitotic figures and tumor giant cells are frequently seen.



Option D: Teratomas also have a variegated appearance but have cysts reflecting the multiplicity of tissue types. Histologically, collections of differentiated cells or organoid structures originating in all 3 germ layers can be seen. Elements may be mature (resembling various adult tissues) or immature (sharing histologic features with fetal or embryonic tissue).



Solution to Question 7:

The standard technique used currently for resection anastomosis of the bowel is a one-layer extra mucosal suture technique since it causes the least tissue necrosis and luminal narrowing. Interrupted and continuous single-layer suture techniques are adequate and safe.

The extra mucosal suture must also include the submucosa because this layer has a high collagen content and it is the most stable suture layer in all sections of the gastrointestinal tract.

Catgut has now been replaced with absorbable synthetic polymers. Stapling devices can be used when speed is required.

Solution to Question 8:

The given clinical scenario of a woman with reddish firm nodule on her chest and histology showing multiple dilated vascular structures with a single layer of epithelial cells is suggestive of a hemangioma.

Hemangioma is a benign tumor caused by the abnormal growth of blood vessels. Types of hemangioma are capillary hemangioma (most common), juvenile/ strawberry hemangioma, cavernous hemangioma and pyogenic hemangioma.

Indications for immediate treatment are complications such as ulceration or those pertaining to the location of the mass (e.g. visual issues due to a large hemangioma over the eyelid).

Treatment options include:

• Beta-blockers - oral propranolol

• Steroids - intralesional triamcinolone, topical clobetasol, systemic steroids

• Interferon alfa-2a and vincristine - for steroid-resistant lesions

• Local resection with cutting cautery or carbon dioxide laser

The given image shows a flat to elevated, irregular, pinkish-red mass over the skin of the abdomen suggestive of a strawberry hemangioma.



The given image shows a capillary hemangioma in a child

Solution to Question 9:

The most common organism causing acute bacterial prostatitis (ABP) is Escherichia coli.

It is commonly caused by an ascending urethral infection or reflux of infected urine into prostatic ducts. Sexually transmissible infections and other urinary pathogens, such as Staphylococcus aureus and Streptococcus faecalis may also be responsible.

The patient presents with fever, perineal pain, chills and rigor, and dysuria. Per rectal examination will reveal a tender, boggy prostate. Care should be taken to avoid aggressive palpation or prostatic massage as it can lead to bacterial dissemination and sepsis.

Urine examination and culture should be done. Urine examination reveals threads in urine.

Management:

Patients with systemic signs of infection should be admitted for IV antibiotics, hydration, and monitoring of laboratory parameters.

Patients who have no signs of systemic infection, who can tolerate oral intake, and who have no urinary retention can be treated with oral antibiotics. Antibiotics are given for 2-6 weeks to ensure the eradication of the infection as antibiotics don't penetrate the prostate capsule easily.

For outpatient therapy: Ciprofloxacin 500 mg oral every 12 hours or levofloxacin 500 mg OD or trimethoprim-sulfamethoxazole double-strength tablet oral every 12 hours can be used.

For hospitalized patients: Intravenous ciprofloxacin or levofloxacin with or without an aminoglycoside (gentamicin or tobramycin) or intravenous beta-lactam with or without an aminoglycoside. Gram-positive infections are treated with cephalosporins (cephalexin, cefazolin) or dicloxacillin and vancomycin.

Solution to Question 10:

The ligament of Treitz is used as the demarcation point to differentiate between upper gastrointestinal bleeding and lower gastrointestinal bleeding. It is a part of the duodenojejunal junction.

Upper GI bleeding refers to bleeding proximal to the ligament of Treitz, and lower GI bleeding refers to bleeding distal to the ligament of Treitz.



Solution to Question 11:

The great saphenous vein (GSV) is the best as it lasts longer than polytetrafluoroethylene (PTFE), dacron, and human umbilical vein as a graft in a femoropopliteal bypass.

The GSV is the conduit of choice for arterial bypass grafts below the inguinal ligament.

For bypass above the inguinal ligament, while both GSV and PTFE grafts have similar patency rates in the first 2 years, the GSV is superior thereafter and hence preferred over PTFE or any other grafts.

The ipsilateral GSV is the conduit of choice for femoropopliteal bypass. When it is not available, contralateral GSV should be used. Other veins are preferred when both ipsilateral and contralateral GSV is not available and they include a small saphenous vein, cephalic vein, and basilic vein.

Non-autogenous grafts are used for patients who lack a suitable venous conduit. The most commonly used prosthetic conduit is PTFE. Recent studies have shown that heparin-bonded PTFE is superior to compared to PTFE without heparin.

Solution to Question 12:

Alcohol consumption is the most common cause of chronic pancreatitis, accounting for 60–70% of the cases. Smoking increases the risk of alcohol-induced chronic pancreatitis.

Cystic fibrosis is the most common cause of chronic pancreatitis in the pediatric age group.

Other causes of chronic pancreatitis include:

• Pancreatic divisum

• Annular pancreas

• Hyperlipidemia

• Hypercalcemia

• Hyperparathyroidism

• Chronic duct obstruction

• Autoimmune pancreatitis

• Cystic fibrosis

• Tropical pancreatitis

• Hereditary pancreatitis

Solution to Question 13:

Microdochectomy is the excision of the entire lactiferous duct (single duct disease). It is done for duct papilloma.

Microdochectomy:

• In a duct papilloma where no palpable mass is present, the duct is expressed to observe the discharge.

• A fine probe or stiff nylon suture is gently inserted into the duct, and the probe is secured to the nipple skin using silk sutures to prevent displacement during the procedure.

• A tennis racquet shaped incision is made to elevate the skin flaps, and the duct with the probe is excised along with the skin flap for 5 cm until the papilloma is identified (The papilloma will always be located within 5 cm from the duct orifice).

• The wound is closed without a drain, and a compression bandage is applied.

It is ideal to rule out DCIS by frozen section biopsy.

Cone excision and wide excision are done under major duct excision (Adair-Haldfield operation). It is done for duct ectasias with multiple duct involvement and multiple papillomas.

Solution to Question 14:

Chromic catgut is absorbed through macrophage phagocytosis and enzymatic degradation within 90 days.

It is made from collagen derived from healthy sheep or cattle. It is then tanned with chromium salts to improve handling and resist degradation in tissue.

The tensile strength lasts for 21-28 days. It has a moderate tissue reaction as compared to plain catgut, which has a high tissue reactivity.

It is used to ligate superficial vessels, suture subcutaneous tissues, and is used for stomas and other tissues that heal rapidly. It is not used in tissues that heal slowly and require prolonged support.

Solution to Question 15:

The given image shows a Morris retractor with a single, flat transversely curved blade and a blunt projecting ridge with backward projection.

Morris retractor can be single or double-bladed. It is used to retract abdominal wall incisions, loin incisions, subcostal wounds or any wounds in depth.

It can be used to improve visibility on one side of an incision. Thus, it is valuable during the initial phase of a laparotomy.

Other options:

Option A: Deaver's retractor is a question mark-shaped stout, atraumatic retractor with a broad, gently curved long blade. It is used to retract the abdominal wall and viscera. Moist mops should be kept underneath prior to placing this retractor. It is used for major abdominal surgeries, bladder surgeries and lumbar sympathectomy.

Option C: Czerny's retractor is a superficial retractor with a thick and small blade on one side and a biflanged hook on the other side in opposite directions. It is used in hernia surgery and laparotomy, especially during closure.



Option D: Double-hook retractor consists of two hooks with a handle. It is used in excision of swellings like lipoma, neurofibroma and cyst. It is also used to hold the skin flaps when raising them.



Solution to Question 16:

Extramammary Paget's disease arises most frequently on the vulva. Other sites include the perineum, scrotum, perianal skin, and penis.

Extramammary Paget's disease (EMPD) can be primary or secondary.

• Primary EMPD is more common and occurs in the absence of an underlying malignancy, thought to arise from intraepidermal parts of apocrine glands or from pluripotent cells from the epidermis.

• Secondary EMPD arises from an underlying adnexal carcinoma or internal malignancy due to the epidermotropic spread of malignant cells from the underlying tumor.

Visceral malignancies associated with EMPD are carcinomas of the colon, rectum, bladder, urethra, cervix, and prostate. The patient presents with well-defined, moist, erythematous scaly plaques and severe pruritis. Lichenification and excoriations may also be seen.

Diagnosis is confirmed by punch, wedge, or excisional skin biopsy. A systemic workup should also be done to identify any underlying malignancy and they include colonoscopy, cystoscopy, colposcopy, and pelvic examination with pap smear in females.

Treatment:

• Mohs microscopic surgery with 5 cm margin clearance

• Radiotherapy

• Topical 5-fluorouracil and imiquimod

• Chemotherapy

• Photodynamic therapy

The given image below shows extramammary Paget's disease of the vulva.

Solution to Question 17:

Transgastric or transduodenal FNAC or trucut biopsy performed under endoscopic ultrasound (EUS) guidance is the best method to obtain tissue for the diagnosis of pancreatic carcinoma. This is because it avoids the spillage of tumor cells into the peritoneal cavity.

Endoscopic ultrasonography with a fine needle aspiration (EUS-FNA) is done if CT has failed to demonstrate a tumor. When a tissue diagnosis is required prior to surgery to distinguish between inflammation and neoplasia, EUS-FNA is helpful.

Diagnostic laparoscopy prior to resection can spare an unnecessary laparotomy by identifying small peritoneal and liver metastases.

Percutaneous transperitoneal biopsy of potentially resectable pancreatic tumors is avoided.

Solution to Question 18:

In a child with congenital hypertrophic pyloric stenosis (CHPS), the abdominal mass is best seen during test feed in the right upper quadrant.

Hypertrophic pyloric stenosis most presents with non-bilious projectile vomiting. It is common between 2 and 8 weeks of age. Hypertrophy of the circular muscle of the pylorus results in constriction and obstruction of the gastric outlet. Male infants are at a higher risk.

Clinically, when the infant is given a test feed, an olive-shaped mass is palpable in the right upper quadrant and visible peristalsis is usually observed moving from left to right across the abdomen.

Infants with hypertrophic pyloric stenosis develop a characteristic metabolic abnormality- hypochloraemic hypokalaemic metabolic alkalosis. Persistent vomiting results in loss of hydrochloric acid leading to the development of metabolic alkalosis (due to loss of H^+) and hypochloremia (due to loss of Cl^-).

All infants with congenital hypertrophic pyloric stenosis are evaluated with an ultrasound (investigation of choice). It confirms the diagnosis in the majority of cases.

Management: Rehydration and metabolic correction prior to definitive surgery - Ramstedt's pyloromyotomy.

Solution to Question 19:

The incision for a fasciotomy involves the skin, subcutaneous tissue, superficial fascia, and deep fascia.

Fasciotomy involves incising the deep muscle fascia and is best carried out via longitudinal incisions of skin, subcutaneous fat, and superficial and deep fascia. The muscle will be seen bulging out through the fasciotomy opening.

The lower limb can be decompressed via two incisions, each being lateral to the subcutaneous border of the tibia. This gives access to the two posterior compartments and the leg's peroneal and anterior compartments.



Solution to Question 20:

Symptomatic relapse of Crohn's disease after surgery is not related to the presence of the disease during resection.

The course of the disease after surgery is unpredictable since surgery does not cure Crohn's disease. Instead, the probability of relapse increases with the time since resection.

Other options:

Option A: Antibiotics alone cannot treat a Crohn's mass effectively. Abscesses can be treated with metronidazole and ciprofloxacin and should also undergo drainage (and resection if needed).

Option C: The terminal ileum is most commonly involved in Crohn's disease.

Option D: On endoscopic examination, discontinuous "skip" lesions are visualized. These are irregular areas of inflamed mucosa that are ulcerated with mucopurulent exudate separated by intervening, normal-appearing intestinal mucosa.

Solution to Question 21:

Aphthous ulcers, surrounded by a rim of erythematous mucosa, are the earliest lesions seen in a patient with Crohn's disease during endoscopy.

Crohn's colitis usually spares the rectum and is most severe in the ileocaecal region.

• In early Crohn's disease, small, punched-out aphthous ulcers are often seen (the result of submucosal lymphoid follicle expansion).

• In moderate disease, the aphthous ulcers coalesce into larger ulcers, which may take on the appearance of a star (stellate ulcers).

• As disease severity increases, submucosal edema and injury can result in cobblestoning of the mucosa (option A) and eventually the formation of serpiginous ulcers, which are typical findings in Crohn's disease.

Thorn-like extraluminal projections in barium studies are known as rose-thorn appearance, and are a typical sign of Crohn's disease. They represent deep penetrating linear ulcers or fissurings typically seen within stenosed terminal ileum with a thickened wall. (Options C and D)

Solution to Question 22:

Staining for estrogen and progesterone receptors is not part of the Van Nuys system used for grading DCIS.

The components of the Van Nuys prognostic index (VNPI) are as follows:

• Size of the tumor

• Extent of resection margin

• Histological variant of DCIS - based on nuclear grade, presence of necrosis, presence of microcalcification

• Age of the patient

Patients with a high score in this grading system benefit from radiotherapy after excision, whereas those with low-grade and completely excised tumors, need no further treatment.

A score ranging from 1 (lesions with the best prognosis) to 3 (lesions with the worst prognosis) is given for each of the four prognostic predictors. Total scores range from 4 (least likely to recur) to 12 (most likely to recur).

Note: The type of microcalcification seen on imaging, helps in determining the nuclear grade of DCIS. While the linear branching type of microcalcification is associated with high nuclear grade, fine granular type is associated with low nuclear grade.

Silverstein Van Nuys prognostic index for DCIS

Parameters	Score 1	Score 2	Score 3
Size(mm)	≤15	16-40	≥41
Margin(mm)	≥10	1-9	<1
Class	Grade 1 or 2 without necrosis	Grade 1 or 2 with necrosis	Grade 3 with or without necrosis
Age(years)	>60	40-60	<40

Scores 4-7 with >3mm margin:Excision aloneScore 7 with < 3 mm margin, score 8 with > 3mm margin, score 9 with >5mm margin:Excision with RTScore 9 with < 5mm margin & score 10,11,12:Mastectomy

Solution to Question 23:

Keloid scars are associated with elevated levels of growth factor leading to increased proliferation of fibrous tissue.

A keloid scar is defined as excessive scar tissue that extends beyond the boundaries of the original incision or wound (option A) and rarely regresses on its own. It is more common in darker-pigmented ethnicities.

It can arise from surgery, burns, skin inflammation, infections, ear piercing, and even spontaneously. It occurs 3 months to years after the initial insult with even minor injuries causing keloid formation.

It most commonly occurs on the skin of the earlobe, deltoid, presternal, upper back, and rarely extends into the subcutaneous tissue.

The histology of keloid scars shows excess collagen and hypervascularity with collagen fibres arranged haphazardly in loose sheets (option C).

Treatment: Keloid scars do not improve spontaneously with time and excision alone has a high rate of recurrence (option B). The following can be done:

• Pressure - local moulds or elastic garments.

• Silicone gel sheeting.

• Intralesional triamcinolone injection.

• Excision and steroid injections.

• Excision and postoperative radiation (external beam or brachytherapy).

• Intralesional excision.

• Laser to reduce redness.

• Vitamin E and palm oil massage.

Differences between Keloids and Hypertrophic scars:

	Keloid	Hypertrophic scar
Incidence	Rare	Frequent
Ethnic groups	African, American, Asian, Hispanic	No predilection
Prior injury	Yes	Yes
Site predilection	Neck, chest, earlobes, shoulders, upper back	Anywhere
Genetics	Autosomal dominantwith incomplete penetration	No
Timing	Symptom-free interval; may appear years after injury	4-6 weeks post-injury

	Keloid	Hypertrophic scar
Symptoms	Pain, pruritus, hyperesthesia, growth beyond wound margins	Raised, some pruritus, respects wound confines
Regression and contracture	No, and rare	Frequent
Histology	Hypocellular, thick, wavy collagen fibres inrandom orientation	Parallel orientationof collagen fibres

Solution to Question 24:

The most common surgical procedure for achalasia cardia is laproscopic Heller myotomy in conjugation with partial fundoplication.

Solution to Question 25:

Mucus-secreting goblet cells (not foveolar cells) are characteristic of Barrett's esophagus, as there is intestinal metaplasia of the esophagus. Foveolar cells are the mucus-secreting cells in the stomach.

Barrett's esophagus is a complication of chronic gastroesophageal reflux disease (GERD), and it is estimated to occur in about 10% of individuals with symptomatic GERD. Symptoms include heartburn, regurgitation and sometimes dysphagia. Obese white men over the age of 60 years represent the highest-risk group.

Intestinal metaplasia in the esophagus occurs due to chronic irritation of the esophagus in GERD (option A) since intestinal epithelium is better suited to handling gastric acid exposure. Hence, mucus-secreting goblet cells are characteristic of Barrett's esophagus. Since all metaplasia are reversible, Barrett's esophagus is also reversible (option D). Only 0.2 to 1% of Barrett's esophagus leads to dysplasia and adenocarcinoma.



Alician blue is used in the staining of the biopsy specimen, as it stains the goblet cells (option C). Mucicarmine is the specific stain.

Seattle protocol is used for the management of Barrett's esophagus.



Solution to Question 26:

• Modified shock index = HR/MAP

• Original shock index = HR/SBP

The shock index is a hemodynamic stability indicator. In patients with shock, it is a better predictor of mortality than blood pressure or heart rate alone. A shock index >0.9 is indicative of higher mortality.

The modified shock index is a more sensitive indicator of tissue perfusion than the original shock index. In critically ill patients, the DBP decreases earlier than SBP, hence, the modified shock index using MAP is better suited to assess the severity of shock.

Surgery NEET 2019

Question 1:

The type of knot seen in the image is:



a) Half hitch knot

b) Reef knot

c) Surgeon's knot

d) Crossed half hitch knot

Question 2:

Which of the following drugs causes carcinoma bladder?

a) Cyclophosphamide

b) Cisplatin

c) Taxane

d) Tamoxifen

Question 3:

The most common site of branchial cyst is _____.

a) Junction of upper and middle third of the sternomastoid muscle

b) Junction of middle and lower third of the sternomastoid muscle

c) Middle of the sternomastoid muscle

d) Between the two heads of the sternomastoid muscle

Question 4:

Dohlman's procedure is for:

a) Meckel's diverticulum

b) Zenker's diverticulum

c) Bochadlek hernia

d) Menetrier's disease

Question 5:

Parathyroid gland post surgery is commonly transferred to:

a) Triceps

b) Biceps

c) Sartorius

d) Brachioradialis

Question 6:

What is the drug used for estrogen dependant breast cancer?

a) Estrogen

b) Clomiphene citrate

c) Tamoxifen

d) Adriamycin

Question 7:

The commonest surgical approach to a retrosternal goiter is via:

a) Neck

b) Chest

c) Median sternotomy

d) None of the above

Question 8:

An intern placed an intercostal drainage tube in a patient. Which of the following steps is incorrect?

a) Identified 5th intercostal space anterior to mid axillary line to place the tube

b) Inserted tube along the lower border of upper rib

c) Directed tube posteriorly to prevent injury

d) Incised and digitally explored

Question 9:

What is the size of the given blade used for incision and drainage of a superficial abscess?

a) 10

b) 11

c) 22

d) 15

Question 10:

In a blast injury, which of the following organs is least vulnerable to the blast wave?

a) GI tract

b) Lungs

c) Liver

d) Ear drum

Question 11:

A man presents with a slowly progressive swelling over the past 1 and 1/2 years. On examination, it has a variable consistency and appears as shown in the image. What is the probable cause?

a) Adamantinoma

b) Dermoid cyst

c) Pleomorphic adenoma

d) Sebaceous cyst

Question 12:

What is shown in the given image?



a) Dermoid cyst

b) Sebaceous cyst

c) Meningioma

d) Post auricular fistula

Question 13:

A female patient presents with a breast abscess. An aspiration was attempted under antibiotic cover which failed to reveal any aspirate. What is the next step in management?

a) Incision and Drainage

b) Continue antibiotics only

c) A second attempt at aspiration

d) Stop antibiotics as no aspirate was collected

Question 14:

Which nerve is involved in Frey's syndrome?

a) Facial Nerve

b) Oculomotor nerve

c) Auriculotemporal nerve

d) Glossopharyngeal nerve

Question 15:

Which of the following statements is false regarding deep vein thrombosis (DVT)?

a) Clinical assessment is unreliable

b) Mostly bilateral

c) Most common clinical presentation is pain and tenderness in calf

d) Some cases may directly present as pulmonary thromboembolism

Question 16:

A truck driver presents with dilated, tortuous veins of 4 mm in the foot. What is the stage of clinical classification according to CEAP(Clinical-etiology-anatomy-pathophysiology for chronic venous disorders)?

a) C0

b) C1

c) C2

d) C3

Question 17:

Which of the following statements is false regarding rectal carcinoma?

a) Hartmann's operation is done in elderly debilitated patients

b) Adenocarcinoma is a rare variant of rectal carcinoma

c) Early morning spurious diarrhoea and tenesmus can occur

d) Growth confined to the rectal mucosa is stage A of modified Duke's staging

Question 18:

A gasless abdomen is seen in:

a) Ulcerative colitis

b) Intussusception

c) Acute pancreatitis

d) Crohn's disease

Question 19:

Gall stone chemistry most commonly seen amongst the following is:

a) Crystalline cholesterol monohydrate

b) Crystalline cholesterol dihydrate

c) Amorphous cholesterol dihydrate

d) Amorphous cholesterol monohydrate

Question 20:

A 56-year-old patient suffering from alcoholic cirrhosis receives a healthy liver from his biological son. What is this type of transplant known as?

a) Autograft

b) Allograft

c) Xenograft

d) Isograft

Question 21:

Esophageal endoscopy of a patient reveals the presence of serpiginous ulcers. This finding indicates:

a) CMV esophagitis

b) Candida esophagitis

c) Herpetic esophagitis

d) Radiation esophagitis

Question 22:

Which of the following is not a part of revised trauma score (RTS)?

a) Glasgow coma scale

b) Systolic blood pressure

c) Pulse rate

d) Respiratory rate

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	d
6	c
7	a
8	b
9	b
10	c
11	c
12	b
13	c
14	c
15	b
16	c
17	b
18	c
19	a
20	b
21	a
22	c

Detailed Explanations

Solution to Question 1:

The given image shows the suture material exiting the knot is on the same side as the entry i.e no cross-over is seen. This is seen in a reef knot.

A knot is a fastening made of threads intertwined together or on themselves. Each knot consists of- 2 half-hitches placed in sequential throws. Squaring refers to tightening the knot/half-hitch in a manner that the threads lay flat due to the alternating direction of half-hitches in adjacent layers.

Types of knots:

• Granny knot: half-hitch over a half-hitch in the same direction. It has better-holding power than half-hitch

• Reef knot: half-hitch followed by a second half-hitch after squaring.

• Triple throw knot: Reef knot with another half-hitch knot for added security. Knots made with prolene sutures require 6-8 throws to prevent slippage.

• Surgeon's knot: Modification of the reef knot with two half-hitches in the first layer.

• Slip knot: Modification of either granny/reef knot formed when one end of the thread is kept taut while tightening the knot.



Solution to Question 2:

Cyclophosphamide (an alkylating agent) causes bladder cancer

The risk of bladder cancer with cyclophosphamide is directly proportional to the duration and intensity of cyclophosphamide treatment. Phosphoramide mustard is the primary mutagenic metabolite that causes bladder cancer in patients exposed to cyclophosphamide.

Painless, gross hematuria is the most common symptom that is indicative of a bladder carcinoma until proven otherwise.

Depending upon the depth of invasion, bladder carcinoma can be divided into

• Flat, non-invasive carcinoma in situ (CIS)

• Non-muscle invasive bladder cancer (70% of cases) - does not invade detrusor muscle but can invade lamina propria.

• Muscle-invasive bladder cancer - invades the detrusor muscle.

Diagnosis is by cystourethroscopy (directly visualizing the growth) and biopsy. Management is as follows:

• Non-muscle invasive bladder cancer:

• Transurethral resection of bladder tumor (TURBT) followed by intravesical chemotherapy with mitomycin C or epirubicin

• Immunotherapy with regular intravesical instillations of Bacille Calmette Guerin (BCG) in high-risk tumors.

• Muscle invasive bladder cancer: Radical cystectomy along with urinary diversion procedure and lymph node dissection +/- neoadjuvant chemotherapy

Other options:

Option B: Cisplatin is an anticancer drug used to treat testicular cancer. It is associated with an increased risk for acute myeloid leukemia (AML).

Option C: Taxane is an anticancer drug and is the main component in the treatment regimens of metastatic cancers. Taxane's adverse effects are neutropenia and fluid retention (pulmonary edema).

Option D: Tamoxifen is a selective estrogen receptor modulator (SERM) and is used in the treatment of estrogen-dependent breast cancer. Due to its partial agonistic activity, it is known to increase endometrial cancer in post-menopausal women.

Solution to Question 3:

The most common site of a branchial cyst is at the junction of the upper and middle third of the sternomastoid muscle at its anterior border.

The below image shows a large right-sided branchial cyst.

Branchial cyst develops due to the persistence of the cervical sinus. The cervical sinus is formed when the 2nd arch mesoderm overgrows the succeeding arches and overhangs them as shown in the image. The space between the overhanging 2nd pharyngeal arch and the 3rd, 4th, and 6th arches is called the cervical sinus. The cavity of the cervical sinus is lined by ectoderm and

normally gets obliterated.

Branchial cysts are usually lined by squamous epithelium and contain thick, turbid fluid full of cholesterol crystals. It is fluctuant and may transilluminate when it contains clear fluid.

A branchial fistula may be acquired or congenital. The congenital fistula is seen at the junction of the middle and lower third of the sternomastoid (option B). An acquired fistula is usually seen in cases of incomplete surgery for the branchial cyst or spontaneous rupture, and it is seen at the same site as the cyst.



Solution to Question 4:

Dohlman's procedure is done for Zenker's diverticulum.

It is a transoral endoscopic approach, that ablates the septum between the diverticulum and the cervical esophagus. It can be used for diverticula between 2 and 5 cm in size.

Other options:

Option A: Meckel's diverticulum is a remnant of the vitellointestinal duct and is treated by surgical resection.

Option C: Bochadlek's hernia is a congenital posterolateral defect on the left side of the diaphragm due to defective embryological development.

Option D: Menetrier's disease is a premalignant condition with gross hypertrophy of gastric mucosal folds and is treated with gastrectomy.

Solution to Question 5:

The parathyroid is commonly transferred to the brachioradialis muscle post-surgery. This will facilitate easy removal if there is disease recurrence after surgery.

Parathyroidectomy is indicated in:

• Symptomatic primary hyperparathyroidism (HPT):

• Asymptomatic primary HPT with:

• raise in serum calcium >1mg/dl of the normal range

• >400 mg urinary calcium in 24-hour urine test

• < -2.5 Z-score of bone mineral density

• Age <50yr and failed medical treatment

• Severe cases of secondary HPT

Parathyroidectomy is performed in neck hyperextension (semi-Fowler position). The thyroid gland is mobilized, and the parathyroid gland with adenoma is identified and resected. The parathyroid tissue is transplanted to the non-dominant forearm.

Pockets are made in the belly of the brachioradialis muscle, and one to two pieces of parathyroid tissue measuring 1 mm each are placed into each pocket. A total of 12 to 14 pieces are transplanted. This is a type of heterotopic transplant.

Miami criteria (confirm the absence of another hyperactive parathyroid): A drop of parathyroid hormone intraoperatively, to more than half the maximum value at 10 minutes postresection is considered an adequate resection in a parathyroid adenoma. However, if there is no drop in PTH level, then a multi-gland disease is suspected and the surgery is converted to a bilateral neck exploration.

Complications of parathyroidectomy include:

• Persistent or recurrent hyperparathyroidism

• Hypoparathyroidism (presenting as hypocalcemia)

• Hungry bone syndrome

• Injury to the recurrent laryngeal nerve

Solution to Question 6:

Tamoxifen is a selective estrogen receptor modulator (SERM) that is mainly used for estrogen receptor-positive breast cancers in pre-menopausal women.

It exhibits selective agonistic and antagonistic activity for ER receptors (ERα and ERβ):

• Agonist activity at the uterus, bone, liver, and pituitary

• Antagonist activity in breast and blood vessels

Uses:

• Part of the primary treatment regimen for estrogen receptor-positive (ER+) breast cancers in pre-menopausal women

• Adjuvant therapy after breast conservative surgery to prevent a recurrence

• Tamoxifen also is effective (a 40%–50% reduction in tumor incidence) in preventing ER+ breast cancer in women at increased risk.

Adverse effects:

• Vasomotor symptoms (hot flashes) and sweating

• Endometrial cancer

• Increased risk of thromboembolic events

• Fatty liver disease

Solution to Question 7:

The commonest surgical approach to a retrosternal goiter is via the neck and the most common type of goitre approached is the secondary retrosternal goiter.

A goiter is considered to be retrosternal if at least 50% of the thyroid tissue is located intrathoracically. They can be primary or secondary.

Retrosternal goiter presents with dyspnoea, stridor, and dilated neck veins. Pemberton's sign (venous congestion of the neck on the raising of hands above the head) is seen.

Contrast-enhanced computed tomography (CECT) of the neck is performed to assess the location of the thyroid. The definitive treatment is surgical excision of the ectopic thyroid.

Other options:

Option B & C: Median sternotomy (via chest) is indicated in:

• Primary mediastinal goiter

• Malignant retrosternal goiter

• Very large retrosternal goiter

• Recurrence in the mediastinum

	Primary retrosternal goiter	Secondaryretrosternal goiter
Occurrence	1%	Majorityof retrosternal goiters
Origin	Arise fromectopic thyroid tissuein the chestNo connection with the thyroid tissue in the neck	Arise from thedownward extension of cervical thyroid tissuealong the fascial planes of the neck
Blood supply	Intrathoracicblood vessels	Superior and inferior thyroid arteries

Solution to Question 8:

An intercostal tube (ICD) is placed along the upper border of the lower rib at the 5th intercostal space (ICS) to prevent damage to the neurovascular bundle. Hence, option B is incorrect.

Chest tube insertion (thoracostomy) is indicated in pneumothorax, hemothorax, chylothorax, and pleural effusion in order to drain air/fluid from the pleural space.

The triangle of safety, usually in the 5th intercostal space, is the safest site for chest tube insertion, it is bound by:

• anterior to the midaxillary line

• above the horizontal line marked by the level of the nipple

• below and lateral to the pectoralis major muscle



Contraindications include pulmonary adhesions, pulmonary trauma, coagulopathy, and diaphragmatic hernia.

Complications of this procedure:

• Infection at the site of insertion

• Unintentional injury to the lung - hemothorax

• Intercostal nerve/ vessel injury

• Subcutaneous hematoma/ emphysema

Solution to Question 9:

The size of the surgical blade shown is the 11-size blade (stab knife) that has a triangular shape with an oblique edge and a sharp-pointed tip.

Uses:

• Incision and drainage (I & D) of an abscess

• Insertion of drains and arteriotomy.

Other options:

Option A: The 10-size blade is used for skin incisions. It has a curved cutting edge that tapers to a pointed tip.

Option C: The 22-size blade is often used for abdominal incisions. It has a flat, unsharpened back edge and a curved front cutting edge.

Option D: The 15-size blade is used for minor surgical procedures (lipoma excision) and precise skin incisions. It is a smaller blade with a small cutting edge.

Solution to Question 10:

In a blast injury, the liver is the least vulnerable to blast wave (primary blast) injury.

Blast injury is a complex type of physical trauma resulting from direct or indirect exposure to an explosion. Primary blast injuries result from the rapid overpressure or shock waves produced by an explosion. These injuries result from the dramatic changes in barometric pressure projecting outwards from the point of detonation. They predominantly cause damage to air-filled hollow organs of the body from rapid pressure change (barotrauma). This includes the middle ear, lungs, and GIT.

The most sensitive and most frequently injured hollow organs are the tympanic membranes > lungs. Blast damage to the lungs is the most common cause of life-threatening injury following an explosion.

Solid organs are the most resistant to blast injuries as the parenchymal disruption requires very high levels of overpressure.

Classification of blast injuries:

Classification	Explanation	Injury type	Examples
Primary blast	Due to air pressure from the blast	Shockwave (overpressure)	Tympanic membraneinjury, blast lung, intestinal blast inj ury
Secondary blas t	Fragments from the blast, fo llowing a detonation	Penetrating/fragmentation	Allpenetratinginjuries
Tertiary blast	Gross movements of persons, objects, or infrastructure by the blast wind	Blunt trauma	Blunt and crushinjuries. Tra umatic amputation.
Quaternary bl ast	Miscellaneous causes	Exposure to explosive produ cts or toxic substances	Burns, inhalation injury
Quinary blast	Theintentional additionof bi ological or radioactive mater ial to the explosive device	Environmental contaminant s	Radiationsickness,infection.

Solution to Question 11:

A slowly progressive swelling, with a variable consistency, lifting the ear lobe, is suggestive of pleomorphic adenoma of the parotid gland.

Pleomorphic adenoma, the most common benign salivary gland tumor, primarily affects the parotid gland, particularly its superficial lobe. It is also referred to as a mixed tumor due to its combination of epithelial and mesenchymal components. This tumor most commonly occurs in females around the age of 40.

Clinical features:

• Painless, slow-growing swelling

• Lobulated, firm, variable in consistency

• Lifts the earlobe

• No facial nerve involvement

• Curtain sign positive - cannot be moved above the zygomatic bone.

• In case of deep lobe involvement, tonsils, and faucial pillars will be pushed medially.

• Features of malignancy - a sudden increase in size, ulceration of the overlying skin, facial nerve palsy (involvement of facial nerve), painful swelling, and restriction of jaw movements.

Fine needle aspiration cytology (FNAC) is the investigation of choice in diagnosing pleomorphic adenoma. Open biopsy is contraindicated due to the chances of injury to the facial nerve.

All tumors of the superficial lobe of the parotid gland should be managed by superficial parotidectomy. Radical parotidectomy is performed for malignant pleomorphic adenoma.

Other options:

Option A: Adamantinoma is a low-grade malignant tumor that usually presents with pre-tibial pain and swelling since it most commonly affects the tibial diaphysis.

Option B: Dermoid cysts present as firm painless subcutaneous nodules and are of 4 types:

• Sequestration dermoid - sequestration of epithelium during embryological development.

• Tubulodermoid - accumulation of secretions from ectodermal tube/duct

• Implantation dermoid - implantation of the epidermis due to trauma

• Teratomatous dermoid - having all 3 germ layers (ecto, meso, and endoderm)

Option D: Sebaceous cysts are retention cysts of the sebaceous glands that present as smooth, non-tender, non-transillumination swelling with a punctum (in 70% of the patients).

Solution to Question 12:

The image shows a dark-colored summit over a swelling, usually seen in a sebaceous cyst.

Sebaceous cysts are the retention cysts of the sebaceous gland that are usually present on the face, scalp, or scrotum. They contain yellowish-white cheesy material with fat and epithelium.

Clinical features:

• The swelling is painless, soft, and freely mobile.

• Though non-transilluminating, they are Paget's test positive (fluctuant).

• Punctum is seen in 70% of the patients with sebaceous cysts. They are dark in color due to the accumulation of denuded epithelium.

• They also have an unpleasant sebum odor.

• The swelling can ulcerate and become painful (Cock's peculiar tumor).

• Some cysts over the scalp may also harden over time (Sebaceous horn).

• Many may get infected and lead to abscess formation.

Treatment:

• Excision including skin adjacent to punctum using elliptical incision-dissection method.

• Incision and avulsion of cyst wall.

• In the presence of an abscess, an initial treatment with an incision later followed by excision of the cyst

Other options:

Option A: Dermoid cysts present as firm painless subcutaneous nodules and are of 4 types:

• Sequestration dermoid - sequestration of epithelium during embryological development

• Tubulodermoid - accumulation of secretions from ectodermal tube/duct

• Implantation dermoid - implantation of the epidermis due to trauma

• Teratomatous dermoid - having all 3 germ layers (ecto, meso, and endoderm)

Option C: Meningiomas are benign tumors arising from the meningothelial cells of the dura. They present with headaches, seizures, or focal neurological deficits.

Option D: Postauricular cutaneous mastoid fistula is a rare complication of chronic otitis media that heals very slowly and needs surgical closure.

Solution to Question 13:

In a patient with a breast abscess, the next step after a failed aspiration is to attempt aspiration once again under antibiotic cover.

Incision and drainage is not preferred, as it can potentially interfere with breastfeeding, and delicate structures such as ducts, lobules, and tubules may be injured, possibly leading to a non-healing milk fistula.

A breast abscess is a complication of acute mastitis, which is more common in new mothers at the beginning of lactation (lactational mastitis). Non-lactational mastitis is more common in diabetic, immunosuppressed people and people predisposed by duct ectasia and periareolar infections.

Clinical features of a breast abscess include:

• Fever and malaise

• Localized pain and tenderness

• Redness and swelling of the affected breast

• Positive fluctuation test.

An ultrasonogram of the affected breast is performed to confirm the abscess, and treatment proceeds as follows:

• Analgesics, antipyretics, and breast support for pain and fever relief.

• Aspiration is performed under antibiotic cover, and the pus is cultured. If aspiration proves unsuccessful, it should be reattempted after a few days.

• Incision and drainage should be considered in cases of:

• Mastitis unresponsive to antibiotics for 48 hours.

• Brawny induration.

• Progressive mastitis with persistent fever

Note:

	Bacterial mastitis	Breast abscess
Condition	Localized inflammation due t o ascending infection or infec tion due to milk stasis	Occurs as a complication of mastitisPus formation
Symptoms	Dullpain	Throbbingpain
Signs (if any)	TendernessErythema	Point tendernessErythemaIn duration of skin(if chronic)Ly mphadenopathyFluctuation t est +vePus pointmay be prese nt
Managemen t	AnalgesicsAntibiotics (Diclox acillin500 mg every 6 hours orally; Erythromycin if allergi c to penicillin)Adequate brea st support	AnalgesicsAdequate breast su pportAspiration of Pus under antibiotic cover; repeated as pirations if requiredIncision & drainage

Solution to Question 14:

The auriculotemporal nerve (V3 nerve) is involved in Frey's syndrome (gustatory sweating).

Frey's syndrome is also known as Dupuy syndrome or auriculotemporal syndrome. It commonly develops after parotid surgery. It is caused by an aberrant regeneration of post-ganglionic parasympathetic nerve fibers crossed along the sympathetic pathways to the sweat glands near the parotid region.

Clinical features are due to the result of salivary stimulation (by the smell or taste of food) leading to the sweating and flushing of the periauricular region on the involved side of the face. A starch iodine test is used for diagnosis.

Treatment ranges from antiperspirants (in mild cases) to tympanic neurectomy or botulinum toxin injection (in severe cases). It can be prevented by incorporating careful extracapsular dissection and closure of the superficial musculoaponeurotic layer during parotidectomy.



Other options:

Option A: Facial nerve is the 7th cranial nerve that supplies the muscles of the face, and results in facial paralysis if injured during parotidectomy

Option B: Oculomotor is the 3rd cranial nerve which supplies the extraocular muscles. If injured during ophthalmic surgeries, it results in ocular paralysis and the eye pointing down and out.

Option D: Glossopharyngeal nerve is the 9th cranial nerve and its injury causes loss of taste sensation from the posterior 1/3rd of the tongue.

Solution to Question 15:

Most cases of deep vein thrombosis (DVT) are unilateral. It is bilateral in only 30% of the individuals.

DVT refers to thrombosis of the deep venous system.

Caused by:

• Postoperative thrombosis (most common cause)

• Oral contraceptive pills (OCPs)

• After childbirth

• Immobility

Sites:

• Pelvic veins—common

• Leg veins—common in femoral and popliteal veins (common on the left side).

Fever is the earliest symptom, which is also most commonly accompanied by pain and tenderness in the calf and thigh (option C).

The following signs could be elicited:

• Positive Homan's sign: Resistance of calf muscles on passive foreceful dorsiflexion of the foot.

• Mose's sign: Gentle squeezing of the lower part of the calf from side to side is painful.

• Neuhof's sign: Thickening and deep tenderness seen on deep palpation of the calf muscles.

Sometimes, the thrombus may get dislodged and present as pulmonary thromboembolism (option D) with features of dyspnoea, tachypnoea, and chest pain.

Clinical features are unreliable for making a definitive diagnosis (option A), and a venous Doppler is the best modality for confirming DVT. It reveals a non-compressible dilated vein. The likelihood of pulmonary embolism can be predicted using modified Wells criteria.

Patients with DVT should be immediately started with subcutaneous low molecular weight heparin (LMWH). If contraindicated due to renal insufficiency or thrombocytopenia-related conditions, they should be started on fondaparinux (indirect factor Xa inhibitor) or bivalirudin (direct thrombin inhibitor). Anticoagulants should be administered for a minimum period of 3 months. Vena cava filters are used in patients with bleeding risks (like hemophilia).

Solution to Question 16:

Dilated tortuous veins of 4 mm indicate varicose veins, so the stage is C2.

Solution to Question 17:

Adenocarcinoma is the most common variant of rectal carcinoma.

Colorectal cancer is:

• The second most common malignancy globally

• The second most common cancer in women

• The third most common cancer in men

• The fourth most common cause of death due to cancer.

Symptoms of rectal carcinoma include:

• Painless bleeding - earliest and most common

• Alteration of bowel habits - change in frequency and consistency of the stool

• Tenesmus - unable to pass a motion even when there is the sensation of needing to evacuate the bowel. It is suggestive of a tumor of the lower half of the rectum

• Early morning bloody diarrhea (option C) - passage of blood and mucus in addition to feces. Pain and weight loss are usually late symptoms

Abdominal examination and digital rectal examination are used for the primary evaluation. Rigid sigmoidoscopy is performed to obtain a biopsy. A synchronous tumor (primary tumor with a different origin than the one currently investigated) is excluded by performing a colonoscopy.

Modified Duke's classification:

A	Growth limited up to the rect al mucosa (option D)
B	B1: Tumor invading muscular is propria, no lymph node in vasion.B2: Tumors invading i nto or through serosa, no ly mph node invasion.
C	C1: The tumour did not excee d the bowel wall, lymph node metastasis.C2: Tumour exce eded the intestinal wall and l ymph node metastasis.
D	Distant metastasis



Surgery is the mainstay of management for rectal carcinoma. Non-operative management is offered to only those patients who have a complete clinical response to neoadjuvant.

The type of surgery depends on multiple factors (size, location, age of the patient, etc). However, the general consensus is:

• Early cancer (T1 and a few T2) is locally excised via the transanal approach.

• Most rectal cancer is treated by anterior resection and colorectal/coloanal anastomosis.

• Low extensive tumors are treated by laparoscopic abdominoperineal resection(APR) and permanent colostomy.

• Hartmann's operation (option A) is preferred in old and frail patients due to poor anal sphincter and postoperative incontinence.

Solution to Question 18:

A gasless abdomen is an X-ray feature that can be seen in acute pancreatitis. Patients with acute pancreatitis can have significant vomiting and diarrhea that may produce a gasless abdomen.

Common causes of gasless abdomen include:

• High obstruction: gastric outlet obstruction, congenital atresia

• Small bowel obstruction

• Bowel ischemia

• Ascites

• Acute pancreatitis

• Gastroenteritis

• Large abdominal mass - due to displacement

• Post-surgery following total colectomy, esophagogastrectomy, gastrectomy, and low anterior resection

• A paucity of small bowel gas may be a normal finding in some adults

Note: In intussusception, an X-ray shows small bowel obstruction, a mass, and a paucity of gas in the right iliac fossa.

Solution to Question 19:

The most common gallstone chemistry seen is crystalline cholesterol monohydrate.

Types of gallstones:

• Cholesterol stones (75% of gallstones) have >70% cholesterol by weight and are composed of crystalline cholesterol monohydrate. The majority of cholesterol stones have a mixed composition with bile pigments and calcium, in addition to the cholesterol. They are the most common type of gall stones worldwide and in India.

• Pigment stones have <20% cholesterol and are dark due to the presence of calcium bilirubinate as the major constituent.

• Black pigment stones (20% of gallstones) - are formed due to supersaturation of unconjugated bilirubin (seen in hereditary hemolytic anemia (like sickle cell and hereditary spherocytosis)

• Brown pigment stones (5% of gallstones) - formed secondary to infections (like E. coli, and Ascaris lumbricoides)

Note: Pure cholesterol stones are uncommon and account for <10% of all stones.

Solution to Question 20:

Transplant of an organ/tissue from a son to a parent is an example of an allograft.

An allograft is transplant of cells/tissues/organs from a genetically non-identical donor of the same species.

Isografts are allografts in which cells/tissues/organs are transplanted from a donor to a genetically identical recipient. (monozygotic twins)

Solution to Question 21:

The image showing serpiginous ulcers in the otherwise normal mucosa of the distal esophagus is suggestive of cytomegaloviral esophagitis.



Solution to Question 22:

Pulse rate is NOT a part of revised trauma score.

The Revised Trauma Score is made up of 3 categories: Glasgow Coma Scale, systolic blood pressure, and respiratory rate.

The score range is 0–12.

A patient with an RTS score of 12 is labelled delayed, 11 is urgent, and 3–10 is immediate. Those who have an RTS below 3 are declared dead.

Surgery NEET 2020

Question 1:

Which of the following is true about the flap given in the image.



a) Pedicled TRAM based on superior epigastric artery

b) Free TRAM based on superior and inferior epigastric artery

c) DIEP based on deep inferior epigastric artery

d) Lower Abdominal Fat based inferior epigastric vessels

Question 2:

What is the most common site of gastrinoma in MEN 1 syndrome?

a) Jejunum

b) Ileum

c) Duodenum

d) Stomach

Question 3:

A 50-year-old male who is a chronic smoker presented to the hospital with intermittent claudication pain of both thigh and buttock region on walking about 500 m. Which of the following is a likely diagnosis?

a) Arterial disease involving the superficial femoral artery

b) Arterial disease with aortoiliac involvement

c) Femoral venous insufficiency

d) Arterial disease involving the profunda femoris artery

Question 4:

Which of the following statements is true about retrosternal goitre?

a) Operated in all patients regardless of symptoms

b) Sternal incision is always required to operate

c) It receives blood supply from the thoracodorsal artery

d) Most of the retrosternal goitres can be removed by a neck incision

Question 5:

While performing surgery on a patient who presented with clinical features suggestive of peritonitis, a diverticular perforation is seen along with fecal peritonitis. What is the Hinchey's stage?

a) Stage 1

b) Stage 2

c) Stage 3

d) Stage 4

Question 6:

An 80-year-old woman presented with complaints of fatigue. On imaging, she was found to have a scleroatrophic gall bladder with impacted stones in it. A large stone was also present in the common bile duct and the biliary tree is dilated. What is the next step in management?

a) Immediately take up patient for cholecystectomy

b) Endoscopic retrograde cholangiopancreatography

c) Wait and watch

d) CT scan

Question 7:

A 35-year-old man presented with recurrent episodes of fever with chills. MRCP image is shown below. What is the most likely diagnosis?



a) Primary sclerosing cholangitis

b) Caroli disease

c) Oriental cholangitis

d) Primary biliary carcinoma

Question 8:

A 5-year-old child with Wilson's disease is being considered for liver transplantation. Which of the following is not a parameter in the selection criteria for transplantation?

a) Age <10years

b) PT >100 s

c) Bilirubin >300 µmol/L

d) Jaundice less than 7 days before development of encephalopathy

Question 9:

The most common neuroendocrine tumor of the pancreas is

a) Gastrinoma

b) Somatostatinoma

c) Insulinoma

d) VIPoma

Question 10:

ASEPSIS wound scoring takes all of the following into consideration except:

a) Wound swab from the site

b) Serous discharge

c) Erythema

d) Induration

Question 11:

A patient comes with long-standing non-pitting edema in the legs. What is the most probable cause?

a) Lymphedema

b) Arterial Disorders

c) Venous Disorders

d) Coronary heart disease

Question 12:

A 41-year-old female patient presented with a tortuous swelling over the left side of the scalp and face as shown in the image given below. On local examination, the swelling was pulsatile. On auscultation, a bruit was heard. Which of the following is the most likely diagnosis?

a) Neurofibromatosis

b) Liposarcoma

c) Cirsoid Aneurysm

d) Varicocele

Question 13:

What does the given barium swallow film show?



a) Aortic aneurysm

b) Pharyngeal pouch

c) Carotid body tumor

d) Esophageal carcinoma

Question 14:

True statement about intermittent claudication:

a) Felt at rest

b) Most commonly felt in the calf muscles

c) Day to day distance is variable

d) Relived after getting out of bed

Question 15:

Which of the following is not a part of the primary survey?

a) Measuring blood pressure

b) Removing clothes to inspect for wounds

c) Taking CT scan of the patient

d) Checking airway patency

Question 16:

A 50-year-old male patient, during his routine annual checkup, is found to have an abdominal aortic aneurysm measuring 40 mm in diameter on USG. Which of the following is true about the management of this patient?

a) Urgent endovascular aneurysm repair

b) Surgery is indicated if aneurysmal diameter is >55mm

c) Urgently wheel the patient to OT for surgery

d) Surgery is indicated only if aneurysm is >77mm

Question 17:

Which of the following is the Miami criteria for intraoperative parathyroid hormone (PTH) measurement?

a) Quarter fall in 5 min

b) Quarter fall in 10 min

c) Half fall in 5 min

d) Half fall in 10 min

Question 18:

A 15-year-old boy undergoes bilateral adrenalectomy for phaeochromocytoma. On the third postoperative day, the boy feels lethargic and tired. Blood pressure is 80/60 mm Hg, there is no sign of bleed. What could be the cause of this?

a) Addisonian crisis

b) SIADH

c) Diabetes insipidus

d) Cerebral salt wasting disease

Question 19:

Which of the following is not a component of Thoracoscore?

a) Performance status

b) Complication of surgery

c) ASA grading

d) Priority of surgery

Question 20:

A middle-aged man presented to the emergency department with upper abdominal pain after having a heavy meal. On clinical examination, upper abdominal tenderness was noted. On chest radiography widening of the mediastinum with pneumomediastinum was seen. What is the probable diagnosis?

a) Spontaneous perforation of esophagus

b) Perforation of peptic ulcer

c) Foreign body in esophagus

d) Rupture of emphysematous bulla of lung

Question 21:

A 32-year-old male was diagnosed with achalasia cardia. High-resolution manometry revealed massive, simultaneous, high-pressure contractions of the distal oesophagus. According to the ■Chicago classification, the patient has which type of achalasia?

a) Type II

b) Type I

c) Type IV

d) Type III

Question 22:

Identify the investigation and the lesion depicted in the image given below.



a) Retrograde cystourethrogram with membranous urethra stricture

b) Retrograde cystourethrogram with penile urethral stricture

c) Micturating cystourethrography with bulbar stricture

d) Micturating cystourethrography with prostratic urethra stricture

Question 23:

According to the TNM staging, a patient with lung cancer of size 2.5 cm is staged as _____

a) T1b

b) T1c

c) T2a

d) T1a

Question 24:

Which of the following should be the treatment of choice for renal cell carcinoma with tumor less than 4 cm in size?

a) Partial nephrectomy

b) Partial nephrectomy + neoadjuvant chemotherapy

c) Radical nephrectomy

d) Radical nephrectomy + postoperative radiotherapy

Question 25:

A 38-year-old man was brought to the emergency department after suffering a stab injury to the lower chest. On presentation, he had bradycardia and hypotension which did not improve despite IV fluid administration. Chest radiography showed clear lung fields. What is the next step in management?

a) eFAST

b) Tube thoracostomy

c) CECT chest

d) CECT abdomen

Question 26:

A man was brought to the emergency department after he fell into a manhole and injured his perineum. He gets an urge to micturate but is unable to urinate. On examination, blood was seen at the tip of the urethra, and swelling of the penis and scrotum was seen. What is the site of injury?

a) Membranous urethra rupture

b) Bulbar urethra injury

c) Prostatic urethra injury

d) Urinary bladder rupture

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	d
5	d
6	b
7	a
8	d
9	c
10	d
11	a
12	c
13	b
14	b
15	c
16	b
17	d
18	a
19	b
20	a
21	d
22	b
23	b
24	a
25	a
26	b

Detailed Explanations

Solution to Question 1:

The given image showing only skin and fat moved to the chest is suggestive of a deep inferior epigastric artery perforator flap (DIEP), based on the deep inferior epigastric artery.

DIEP is a variant of TRAM flap and free flap where only the skin and fat are moved to the chest. A microvascular technique is used in which no muscle is taken and the perforating vessels are dissected out.

As the muscle is spared in DIEP, there is a reduced risk of abdominal wall weakness and its associated complications.

Other options:

Options A and B: The pedicled TRAM flap is based on the superior epigastric artery and the free TRAM flap is based on superior and inferior epigastric arteries (supercharged TRAM flap); however, the given image depicts a deep inferior epigastric artery flap. In TRAM flaps:

• The skin, fat, and muscle are moved to the chest.

• It provides good cosmesis and bulk since it has a soft tissue consistency similar to the breast.

• Disadvantages include epigastric bulge, partial flap loss, fat necrosis, abdominal wall weakness, and hernias.

The given image below shows a TRAM flap:



Solution to Question 2:

The most common site of gastrinoma in MEN 1 syndrome is the duodenum.

Gastrinoma is the most common pancreatic islet tumor in MEN 1 syndrome, arising from the G cells of the pancreas. Gastrinoma, associated with marked gastric acid production and recurrent peptic ulcers, is called Zollinger-Ellison syndrome. The majority of gastrinomas occur in the gastrinoma (Passaro's) triangle.

In MEN 1, they often occur in patients <30 years. Patients present with abdominal pain (the most common symptom), diarrhea and steatorrhea.

Management:

• Medical management for tumors <2 cm with proton-pump inhibitors.

• Surgery for tumors >2 cm.

• Proton-pump inhibitors with somatostatin analogues for advanced locoregional disease or metastasis.

Solution to Question 3:

The given clinical scenario is suggestive of arterial disease with aortoiliac involvement. Classically, the muscle group affected by claudication is one anatomical level below the level of arterial disease.

Intermittent claudication is a feature of chronic arterial disease or occlusion of the artery. Aortoiliac disease affects the infrarenal aorta and iliac arteries.

Management involves smoking cessation, lifestyle modifications, and treatment of underlying diabetes or hyperlipidemia if present.

For claudication, cilostazol or pentoxyfilline is used. Cilostazol is superior to pentoxyfilline.

Surgical options include aortoiliac bypass, aortobifemoral bypass, percutaneous transluminal angioplasty (PTA), which can be with or without stent placement, and thromboendarterectomy.

Correlation of clinical findings to the Site of Disease	
Aortoiliac obstruction	Claudication inbuttocks, thighs, and calvesBruit over the aortoiliac regionImpotence(Leriche's syndrome)
Iliac obstruction	Claudication ofthigh and calf, unilaterallyAbsence of femoral and distal pulsesBruit over iliac region
Femoropopliteal obstruction	Claudication incalfFemoral pulse felt but distal pulses absent
Distal obstruction	Ankle pulses absentClaudication incalf and footFemoral and popliteal pulses palpable

Solution to Question 4:

Most of the retrosternal goiters can be removed by neck incision (transcervical approach).

Retrosternal goitre is defined as having >50% of the goiter below the suprasternal notch. It can be either primary or secondary.

• Primary: rare and arises from ectopic thyroid tissue in the mediastinum; receives blood supply from intrathoracic blood vessels. It is not related to thyroid tissue in the neck.

• Secondary: they are the most common type; an extension of enlarged thyroid from the neck, usually arising from the lower pole of nodular goitre. It receives its blood supply from superior and inferior thyroid vessels.

The majority of patients are asymptomatic or have minimal symptoms. If the gland enlarges within the thoracic inlet, the pressure may lead to dysphagia, tracheal compression, and eventually airway symptoms. Pemberton sign can be positive (facial flushing and dilatation of cervical veins upon raising the arms above the head).

CECT neck and thorax is the investigation of choice.

Over 95% of retrosternal goiters can be removed via a transcervical incision since they are an extension from thyroid tissue in the neck.

Other options:

Option A: Surgical removal of the retrosternal goiter is indicated in patients with significant airway compression, and in young patients who have symptoms such as dyspnoea or dysphagia.

Option B: Median sternotomy approach is indicated in:

• Primary mediastinal goiter

• Malignant retrosternal goitre

• Very large retrosternal goiter

• Recurrence in the mediastinum

For a primary mediastinal goiter whose boundaries are completely within the chest, resection is often achieved through median sternotomy or thoracotomy to ensure that the vessels supplying the tumor are properly ligated. Removing the tumor through a cervical approach can result in uncontrolled bleeding and is generally avoided.

Option C: Blood supply to the retrosternal goiter is from the thyroid vessels (secondary retrosternal goiter).

Solution to Question 5:

Fecal peritonitis belongs to grade 4 in the Hinchey classification.

Diverticulitis refers to the inflammation and infection associated with the diverticula. Hinchey classification is the most commonly used classification for the severity and complication of acute diverticulitis. It is used additionally to guide the overall management of acute diverticulitis.

Hinchey classification:	
Stage 1	Mesenteric or pericolic abscess
Stage 2	Pelvic abscess
Stage 3	Purulent peritonitis

Hinchey classification:	
Stage 4	Fecal peritonitis

Solution to Question 6:

An 80-year-old woman with imaging showing a scleroatrophic gall bladder with impacted stones and a stone in the bile duct is suggestive of chronic cholelithiasis with choledocholithiasis. ERCP is the next line of management for this patient.

Choledocholithiasis is the presence of stone in the common bile duct (CBD) or biliary tree. It can be primary or secondary:

• Primary stones are formed in the CBD or biliary tree itself and are rare.

• Secondary stones are passed from the gallbladder to the CBD. They are easier to manage than primary stones.

It can be an asymptomatic and incidental finding or present with features such as jaundice (most common), abdominal pain, and fatigue. When there is inflammation of the biliary tree (cholangitis), the patient will have intermittent pain, fever, and jaundice (Charcot's triad).

Investigations:

• Liver function tests will show elevated direct bilirubin and alkaline phosphatase levels.

• Abdominal ultrasound is the first imaging modality done and may show gallstones and dilated bile duct (>8 mm).

• CT is sensitive for CBD stones. It shows stones, location, ductal stricture or block, ductal dilatation, and intrahepatic biliary changes.

• MRCP is highly sensitive and specific for CBD stones, but it is only diagnostic and intervention cannot be performed.

Management:

• ERCP with sphincterotomy is the procedure of choice, followed by the removal of stones using a dormia basket or placement of a stent.

• Despite ERCP being an invasive procedure, it has the advantage of being both a diagnostic and therapeutic procedure and is more suited for elderly patients who cannot tolerate anesthesia and surgery.

Other options:

Option A: Cholecystectomy is not recommended for elderly patients, owing to the risk of old age and associated anesthetic and surgical complications.

Option C: Wait and watch approach can be done for asymptomatic patients, and patients with smaller stones.

Option D: CT is sensitive for CBD stones, but it is only diagnostic and intervention cannot be performed.

Solution to Question 7:

The presence of fever and chills, along with the MRCP showing multiple areas of intrahepatic beading is typical of primary sclerosing cholangitis.

Option B: Caroli's disease is a rare congenital condition characterized by irregular dilatation of the intrahepatic bile ducts with a normal extrahepatic duct system. It is common in both sexes and mostly presents before the age of 30 years with abdominal pain, cholangitis, or end-stage liver disease. MRCP shows irregular, cystic dilation of the large proximal intrahepatic bile ducts with a normal common bile duct.

Primary sclerosing cholangitis (PSC) is an autoimmune condition characterized by inflammatory strictures involving the intra and extra-hepatic biliary tree. It is a progressive condition that leads to secondary biliary cirrhosis.

MRCP or ERCP will show demonstrate multifocal, diffuse dilatation and strictures of the intrahepatic and extrahepatic biliary trees, this pattern is a beading or chain of lakes appearance. Pseudodiverticula of extrahepatic bile ducts are pathognomic of PSC. MRCP is useful in diagnosis and surveillance, while ERCP is reserved for the treatment of dominant strictures and to exclude malignancy.

The below image shows a cholangiogram of primary sclerosing cholangitis:



The common age group of primary sclerosing cholangitis is 30-45 years. Men are affected twice as commonly as women. Two-thirds of these cases are associated with ulcerative colitis. It is also associated with Riedel’s thyroiditis and retroperitoneal fibrosis. 10–20% of patients develop cholangiocarcinoma.

Clinical features of a patient with PSC include pruritis, fever, chills, right upper quadrant discomfort, jaundice, and weight loss.

Liver function tests show a cholestatic pattern with an elevation of ALP and GGT and smaller rises in the aminotransferases. Bilirubin values may be variable.

Liver biopsy excludes other causes of cholestatic liver disease and defines the stage of PSC. It shows onion skin concentric periductal fibrosis. With disease progression, periportal fibrosis

occurs, progressing to bridging necrosis, and eventually biliary cirrhosis.

Currently, there is no known effective medical therapy. Surgical management involving resection of extrahepatic biliary tree and hepaticojejunostomy has shown reasonable effects. For patients with advanced liver disease, liver transplantation is the only available option.

Other options:

Option C: Oriental cholangitis, also known as recurrent pyogenic cholangitis is characterized by recurrent attacks of cholangitis due to bile stasis and stone formation. Patients between 30-40 years and both sexes are equally affected. It is caused due to infections (liver flukes, E.coli, Klebsiella, Pseudomonas, and Proteus). Patients present with Charcot's triad (intermittent fever, pain, and jaundice). MRCP shows intra or extrahepatic calculi, dilatation of ducts, and strictures.

Option D: Biliary carcinoma (cholangiocarcinoma) most commonly affects patients >60 years and presents with fever, abdominal pain, weight loss, and features of obstructive jaundice. MRCP shows irregular narrowing of the bile duct involved by the tumor, with asymmetric upstream dilatation of the intrahepatic ducts.

Solution to Question 8:

A 5-year-old child with Wilson's disease is a case of non-paracetamol-induced acute liver failure and jaundice for more than 7 days before the development of encephalopathy is one of the parameters in the King's College criteria for liver transplantation.

It includes the following parameters:

Non-paracetamol-induced acute liver failure:

• PT >100 s (INR >6.5)

or/and

• Any three of the following:

• Age <10 years or >40 years

• Etiology non-A, non-B hepatitis, or idiosyncratic drug reaction

• Jaundice more than 7 days before the development of encephalopathy i.e., not hyperacute

• PT >50 s (INR > 3.5)

• Bilirubin > 18 mg/dl (300 μmol/L)

Paracetamol-induced acute liver failure:

• pH <7.30 (irrespective of the grade of encephalopathy)

or/and

• All three of the following:

• Prothrombin time >100 s (INR >6.5)

• Serum creatinine >3.4 mg/dl (300 μmol/L)

• Grade 3 or 4 hepatic encephalopathy

Solution to Question 9:

The most common neuroendocrine tumor of the pancreas is insulinoma (4 per million).

Insulinoma is an insulin-producing tumor of the pancreas, with more than 90% of them being benign. It occurs in >40 years in patients without MEN 1 syndrome and <40 years in patients with MEN 1. It presents with symptoms of hypoglycemia such as weakness, headaches, sweating, faintness, seizures, and altered behavior. These occur after fasting or exertion and improve after glucose intake.

The investigation of choice for insulinoma is the determination of 72-hour fasting glucose and serum insulin levels. C-peptide and proinsulin levels are also elevated.

Medical management includes:

• Frequent carbohydrate-rich meals

• Somatostatin analogs like octreotide are effective in controlling symptoms due to excess insulin secretion. They should be used with caution as they can suppress counterregulatory hormones, such as growth hormone (GH), glucagon, and catecholamines, and precipitously worsen hypoglycemia.

• Diazoxide can be used in the initial management of insulinoma as it inhibits insulin release. Its side effects include sodium retention and nausea.

• Everolimus has been found to be highly effective in improving glycemic control in patients with insulinoma owing to its antitumor effect and induction of insulin resistance.

Enucleation and pancreatectomy are the definitive surgical management of insulinoma.

Other options:

Option A: Gastrinoma has an incidence of 0.5 to 4 in a million.

Options B & D: Somatostatinoma and VIPoma are much rarer than gastrinoma.

Solution to Question 10:

ASEPSIS wound score does not take induration into consideration. Instead, the I in ASEPSIS stands for the isolation of bacteria from the wound.

ASEPSIS wound score and Southampton wound score are used to determine the severity of the surgical site wound infection (SSI).

ASEPSIS score includes:

• Additional treatment

• Antibiotics for wound infection

• Drainage of pus under local anesthesia

• Debridement of the wound under general anesthesia

• Serous discharge

• Erythema

• Purulent exudate

• Separation of deep tissue

• Isolation of bacteria from the wound

• Stay as inpatient prolonged over 14 days as a result of wound infection

Solution to Question 11:

Lymphedema is the most probable cause of long-standing non-pitting edema in the limbs.

Lymphedema is the abnormal accumulation of protein-rich fluid in the interstitial spaces of the tissues.

Lymphedema can be primary or secondary.

Primary lymphedema includes:

• Lymphedema congenita (0-2 years): more common in males, involves multiple limbs, face, and genitalia. It can be familial or sporadic. The familial type is called Nonne Milroy syndrome.

• Lymphedema praecox (2-35 years): more common in females, the most common type of primary lymphedema with a peak incidence in puberty. It can be familial or sporadic. The familial type is called Meig's disease.

• Lymphedema tarda (>35 years): Rare and uncommon.

Secondary lymphedema is more common than primary and is the most common cause of lymphedema overall. Causes include lymph node dissection (axillary or inguinal), post-mastectomy, postoperative radiotherapy, obesity, varicose vein surgery, and trauma.

Complications of lymphedema include infections, skin changes, recurent cellulitis, non-healing ulcers, and lymphangiosarcoma.

Treatment mostly includes patient education and nonsurgical interventions, which include limb elevation, and external compressive garments. When the above methods fail, surgery can be considered.

Solution to Question 12:

The presence of a tortuous, pulsatile swelling over the scalp and face is suggestive of a cirsoid aneurysm.

Cirsoid aneurysm is a slow-growing, tortuous swelling that develops due to abnormal arteriovenous connections among the subcutaneous blood vessels. They are usually congenital. However, in some cases, they may be traumatic. The superficial temporal artery is most commonly involved.

Patients present with a tortuous, pulsatile swelling over the scalp or face. They are usually asymptomatic and present to the clinic due to cosmetic deformity. However, some patients may have headache, tinnitus, and/or bleeding. A loud bruit is heard on auscultation.

Angiography is the investigation of choice.

Treatment includes surgical resection, endovascular occlusion or injection of percutaneous sclerosing agents.

Solution to Question 13:

The given barium swallow film shows an outpouching from the posterior wall of the pharynx suggestive of a Zenker's diverticulum (pharyngeal pouch).

Solution to Question 14:

Intermittent claudication is most commonly felt in the calf muscles because of the involvement of the superficial femoral artery in chronic vascular occlusive disease.

Intermittent claudication:

• Intermittent claudication is the debilitating cramp-like pain felt in the leg muscles due to reduced blood flow and anaerobic metabolism brought on by peripheral arterial disease.

• It usually signifies mild to moderate chronic vascular occlusive disease

• The pain is brought on by walking (option D)

• Not present on taking the first step

• Relieved by rest in both the standing and sitting positions, usually within 5 minutes (option A).

• The distance that a patient is able to walk without stopping is known as claudication distance. It varies only slightly from day to day (option C).

• The superficial femoral artery (70%) is the most commonly affected artery by claudication and manifests as pain in the calf muscles.

• Aorto-iliac occlusion may cause thigh or buttock claudication. Leriche's syndrome occurs due to aortoiliac occlusion and presents with a triad of claudication, impotence, and absent femoral pulses.

Boyd's classification of claudication:

• Grade I: Patient complains of pain after walking. If the patient continues to walk, due to increased blood flow in muscle and opening of collaterals metabolites causing pain are washed away and the pain subsides.

• Grade II: Pain still persists on continuing to walk; but can walk with effort.

• Grade III: The patient has to take rest to relieve the pain.

Solution to Question 15:

Taking a CT scan of the patient is a part of secondary survey.

Primary survey identifies and manages the most immediately life-threatening pathologies first.

The secondary survey includes investigations that help identify all other injuries, excluding the apparent ones. This includes X-rays of the spine and extremities and a CT of the abdomen, head, chest, and spine.

Solution to Question 16:

Surgery is indicated in an asymptomatic abdominal aortic aneurysm (AAA) only if its diameter is more than 55 mm in a male patient.

If the diameter is less than 55 mm, regular ultrasound monitoring is required. This is because elective open surgery has a higher rate of mortality when compared to conservative management for aneurysms of size < 55 mm.

Indications for surgery in abdominal aortic aneurysm:

• Size >5.5 cm in males

• Size >5 cm in females

• Expansion of >1 cm in one year

• Painful tender aneurysm

• Thrombosed aneurysm; aneurysm with distal emboli

Management of aortic aneurysm: AAA is surgically managed by dacron graft repair. It can be either open or EVAR (endovascular aneurysm repair)

• Open surgical repair is done by exposing the aorta through a mid-sternal incision and graft replacement. To expose the aorta the descending colon should be mobilized medially and this is known as left visceral rotation or Mattox procedure.

• EVAR is a minimally invasive procedure in which an aortic stent graft is placed on the nonaneurysmal aortoiliac segment.

Complications following an AAA repair are:

• Cardiovascular (ischemia and infarction)

• Respiratory (atelectasis or consolidation).

• Colonic ischemia.

• Renal failure.

• Aortoduodenal fistula

• spinal cord ischemia

The given image below shows the technique of EVAR:



Solution to Question 17:

A drop in the level of PTH to less than half the maximum value (corresponding to the preoperative levels) at 10 min postresection is the Miami criteria.

It is used to determine the extent of resection of the parathyroid gland which is done in a case of primary hyperparathyroidism due to adenoma.

Intraoperative parathyroid hormone monitoring is a commonly used adjunct, which guides the surgeon on when to stop parathyroid exploration and helps in confirming there is no residual hyperfunctioning parathyroid tissue. The PTH levels are expected to drop after the gland is excised due to its short half-life (3-5 minutes). Blood is drawn at various time points and the hormone levels are assessed.

Based on the Miami Criteria, a more than 50% decline in PTH at 10 minutes following parathyroid gland removal compared to the highest baseline level, either the pre-incision or pre-excision, suggests a cure, and the procedure can be terminated.

Indications for parathyroidectomy:

All symptomatic patients with hyperparathyroidism should be offered surgery. Indications for surgery in asymptomatic patients include:

*BMD - bone mineral density

Serum calcium levels	0.25 mmol/L (1.0 mg/dL)above the upper limit serum calcium of normal
Skeletal	BMD by DEXA:T score –2.5at the lumbar spine, total hip, femoral neck, or distal one-third of radiusVertebral fracture
Renal	Creatinine clearance<60 mL/min24-hour urinary calcium >10 mmol/dL (>400 mg/day)Increasedrisk of stone formationby risk analysis
Age	<50years

Solution to Question 18:

The given clinical scenario of hypotension with no presence of bleeding and bilateral adrenalectomy is suggestive of an Addisonian crisis. It is a life-threatening condition caused due to sudden insufficiency of adrenal hormones.

Acute adrenal insufficiency (adrenal or addisonian's crisis) is a medical emergency caused due to a precipitating illness unmasking insufficiency or trauma or severe sepsis causing adrenal hemorrhage (Waterhouse-Friderichsen syndrome) or surgical removal of the glands. It presents with non-specific symptoms such as shock with fever, nausea, vomiting, abdominal pain, hypocalcemia and electrolyte imbalances. It may be often misdiagnosed as an acute abdominal emergency.

Adrenal insufficiency can be primary (due to adrenal pathology), secondary (due to pituitary pathology- lack of ACTH) or tertiary (due to hypothalamus pathology- lack of CRH).

Autoimmune adrenalitis is the most common cause of primary adrenal insufficiency worldwide.

Tuberculosis is the most common cause of primary adrenal insufficiency in India.

Adrenal crisis requires immediate intervention owing to its high mortality rate. Immediate rehydration is initiated with a saline infusion at 1 litre/hr with continuous cardiac monitoring. Glucocorticoid replacement is initiated with an initial bolus injection of hydrocortisone 100 mg followed by 200 mg over 24 hours, preferably by continuous infusion or alternatively by IV or IM

injections. Mineralocorticoid replacement is initiated once the daily dose of hydrocortisone has been reduced to <50 mg, since hydrocortisone at higher doses provides sufficient stimulation of mineralocorticoids.

Solution to Question 19:

The complication of surgery is not a component of thoracoscore.

Thoracoscore is used to assess the risk of operative mortality in a patient undergoing thoracic surgery. It includes 9 variables:

- Age
- Sex
- American Society of Anesthesiologists score (ASA)
- Performance status
- Dyspnoea score
- Priority of surgery
- Extent of surgery
- Malignant diagnosis
- Composite comorbidity score

Solution to Question 20:

The clinical scenario points to a diagnosis of spontaneous perforation of the esophagus.

In a perforated peptic ulcer, patients develop sudden onset severe generalized abdominal pain, tenderness, and board-like rigidity and an erect plain chest X-ray shows gas under the diaphragm (Option B). The foreign body in the esophagus can be asymptomatic or may present with perforation after ingestion. The radiopaque foreign body can be seen with a chest X-ray (Option C). Rupture of emphysematous bullae of the lung will lead to secondary pneumothorax, which manifests as chest pain and dyspnoea. A plain chest X-ray shows air in the thorax, and the mediastinum is pushed to the opposite side (Option D).

Spontaneous perforation of the esophagus needs immediate diagnosis and intervention due to the higher mortality associated with it. The most common site of perforation is the lower esophagus towards the left esophageal cavity.

Causes:

- Iatrogenic perforation (most common)
- Boerhaave's syndrome (spontaneous emetogenic perforation)
- Sharp foreign body
- Corrosive ingestion

• Esophageal carcinoma

It usually manifests as severe pain in the chest or abdomen after a meal or after a bout of drinking, commonly associated with dyspnoea. On clinical examination, there is rigidity and tenderness in the upper abdomen. The diagnosis is usually confirmed by a chest radiograph which shows pneumomediastinum, pneumoperitoneum, and pneumothorax. Subcutaneous emphysema can be seen in Boerhaave's syndrome. Patients with sepsis will have a fever, tachycardia and tachypnea.

Treatment includes adequate drainage, antibiotics, closure of the perforation site if possible and supportive measures.

Solution to Question 21:

The given features are suggestive of type III achalasia cardia.

Type III achalasia cardia is characterised by massive, simultaneous, high-pressure contractions of the distal esophagus.

Achalasia cardia occurs due to the loss of inhibitory ganglion cells in the myenteric plexus of GIT. During swallowing, the food bolus normally triggers a primary peristaltic wave, which is responsible for carrying the bolus to the stomach, accompanied by the relaxation of the lower esophageal sphincter (LES) to allow esophageal emptying into the stomach.

Achalasia is characterized by a disruption of this normal process. Failure of relaxation of the LES causes an elevated median integrated relaxation pressure and a complete failure of peristalsis. The esophagus becomes progressively more tortuous, leading to a megaesophagus.

Solution to Question 22:

The given image shows retrograde cystourethrogram with findings consistent with penile urethral stricture.

The male urethra can be visualized by retrograde urethrography (RGU) or voiding cystourethrogram (VCU), also known as micturating cystourethrogram (MCU).

In RGU, the urethra is visualized with the instillation of contrast into the anterior urethra (penile urethra and bulbar urethra). It is helpful for diagnosing anterior urethral abnormalities.

The following image shows an RGU with bulbar urethral stricture:



In VCU, after injecting contrast into the bladder using a Foleys catheter, the posterior urethra (membranous urethra and prostatic urethra) is visualized during micturition. It is helpful for diagnosing posterior urethral abnormalities and bladder abnormalities. It is the investigation of choice for vesicoureteric reflux (VUR) and posterior urethral valve.



Solution to Question 23:

According to TNM staging, a patient with lung carcinoma of size 2.5cm is staged as T1c.

Stage T1-3/N0-1/M0 are amenable to resection.

TNM staging system of Lung carcinoma	
T: Primary tumor	
Tx	Primary tumor cannot be assessed or tumor proven by the presence of malignant cells in sputum or bronchial washings but not visualized by imaging or bronchoscopy
T0	No evidence of primary tumor
Tis	Carcinoma in situ
T1	Tumour ≤3 cm in the greatest dimension surrounded by lung or visceral pleura without bronchoscopic evidence of invasion more proximal than the lobar bronchus (i.e. not in the main bronchus)
T1a(mi)	Minimally invasive adenocarcinoma
T1a	Tumour≤1 cmin greatest dimension
T1b	Tumor>1 cm but ≤2 cmin greatest dimension
T1c	Tumor >2 cm but≤3 cmin the greatest dimension
T2	Tumor >3 cm but≤5 cmor tumor with any of the following features:– involves themain bronchus regardless of distancefrom thecarinabutwithoutthe involvement of the carina – invades visceral pleura– associated withatelectasisorobstructive pneumonitisthat extends to thehilar region, involving part or all of the lung
T2a	Tumor>3 cm but ≤4 cmin greatest dimension
T2b	Tumor>4 cm but ≤5 cmin greatest dimension

TNM staging system of Lung carcinoma	
T3	Tumor>5 cm but ≤7 cmin greatest dimension or associated with separate tumor nodule(s) in the same lobe as the primary tumorordirectly invades any of the following structures:chest wall (including the parietal pleura and superior sulcus tumors), phrenic nerve, the parietal pericardium
T4	Tumor>7 cmin greatest dimension or associated with separate tumor nodule(s) in a different ipsilateral lobe than that of the primary tumor or invades any of the following structures:diaphragm, mediastinum, heart, great vessels, trachea, recurrent laryngeal nerve, esophagus, vertebral body, and carina
N: Lymph node involvement	
Nx	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in ipsilateral peribronchial and/or ipsilateral hilar lymph nodes and intrapulmonary nodes, including involvement by direct extension
N2	Metastasis in ipsilateral mediastinal and/or subcarinal lymph node(s)
N3	Metastasis in contralateral mediastinal, contralateral hilar, ipsilateral or contralateral scalene, or supraclavicular lymph node(s)
M: Distant metastasis	
M0	No distant metastasis
M1	Distant metastasis present
M1a	Separate tumor nodule(s) in a contralateral lobe; tumor with pleural or pericardial nodule(s) or malignant pleural or pericardial effusion

TNM staging system of Lung carcinoma	
M1b	Single extrathoracic metastasis
M1c	Multiple extrathoracic metastases in one or more organs

Solution to Question 24:

Partial nephrectomy (also known as nephron-sparing surgery) is the treatment of choice for tumors <4 cm.

Indications for partial nephrectomy:

• Tumors < 4cm

• Well-selected tumors between 4-7 cm

• Bilateral tumors

• Tumor in solitary kidneys

• Patients with pre-existing renal dysfunction.

Recently, partial nephrectomy is also increasingly being performed for stage T1b tumors.

Treatment of renal cell carcinoma:

RCC is a radio and chemoresistant tumor.

Radical nephrectomy consists of the removal of the kidney, perirenal fat, adrenal gland, and regional lymph nodes. The adrenal gland is no longer removed except in cases of obvious tumor involvement. Lymphadenectomy is indicated only in high-risk patients with large primary tumors and enlarged lymph nodes.

Cryoablation is indicated in the T1a stage of RCC if surgery cannot be performed. It is also indicated in advanced/metastatic tumors.

Note: Sunitinib and sorafenib are also used in GIST and hepatocellular carcinoma.

Staging	Description	Management
Stage 1	Tumor <7 cm, confined to the kidney	Partial nephrectomy
Stage 2	Tumor >7 cm, confined to the kidney	Radical nephrectomy
Stage 3	Tumor extends into the major veins or adrenal gland with intact Gerota's fascia or regional lymph nodes involved.	Radical nephrectomy

Staging	Description	Management
Stage 4	Tumor invades beyond Gerota fascia (including contiguous extension into the ipsilateral adrenal gland)	Debulking surgerymTOR inhibitors such as everolimus, sirolimusTyrosine kinase inhibitors: sunitinib, sorafenib

Solution to Question 25:

The given clinical scenario of a patient with a stab injury and hypotension not responding to IV fluids indicates persistent hypotension due to ongoing hemorrhage and the next step in the management is to perform an eFAST.

Cardiogenic shock or hemorrhagic shock are the most likely causes of persistent hypotension in a trauma patient. Ultrasound evaluation using eFAST to look for pericardium, pleural cavities, and abdomen in combination with plain radiographs of the chest and pelvis will usually identify the source of the bleed.

In patients with persistent hypotension due to blunt trauma, the pelvis should be wrapped in a sheet for stabilization until radiography can be done.

Evaluation of CVP or ultrasound of the IVC may further assist in distinguishing between cardiogenic and hypovolemic shock.

It is mostly fatal if not controlled, and requires immediate management. Blood and plasma transfusions are provided in a ratio of 2:1. Patients with persistent hypotension requiring surgical intervention include:

• Blunt trauma and massive hemothorax

• Penetrating trauma

• Hemoperitoneum

• Gunshot injuries to the chest or abdomen

• Penetrating weapon remaining in place.

Other options:

Options C and D: CECT can be done only when the patient is hemodynamically stable.

Option B: Tube thoracostomy has no role in this patient's management since the X-ray shows clear lung fields.

Solution to Question 26:

The given clinical scenario with a history of a blow to the perineum, blood at the tip of the meatus, swelling of the penis and scrotum is suggestive of an injury to the bulbar urethra.

Urethral injuries can be classified into:

Note: Urinary bladder rupture occurs due to an injury to the suprapubic region. Extraperitoneal bladder rupture is more common and associated with pelvic fracture. Intraperitoneal bladder rupture is caused by a blow or a fall on a distended bladder. It manifests as sudden severe pain in the hypogastrium and syncope.

Type	Proximal urethral injury	Distal urethral injury
Site of injury	Prostatic urethraMembranous urethra	Bulbar urethraPenile urethra
Mechanism	Secondary topelvic fracture	Occurs due to a blow to the perineum by:Straddle injuryLoose manhole coveringGymnasium accidents
Clinical features	Urinary retentionBlood at the urethral meatusHigh-riding prostate on digital rectal examination (Vermooten sign)Deep perineal hematoma	Urinary retentionBlood at the urethral meatusPerineal hematoma (butterfly hematoma)

Surgery INI-CET 2021

Question 1:

What is triage used for?

a) To prepare for a disaster

b) To asess the impact of a disaster

c) To classify priority of treatment

d) To rehabilitate following a disaster

Question 2:

Match the following

a) 1-C,2-D,3-B,4-A

b) 1-A,2-D,3-B,4-C

c) 1-C,2-D,3-A, 4-B

d) 1-D,2-C,3-A,4-B

Question 3:

A mother notices a swelling in the abdomen of her 3-year-old child while bathing him. He had a history of hematuria two weeks back which resolved spontaneously. On examination, a right-sided reniform ballotable mass was found. What all investigations will you do?

a) 1 and 2

b) 2 and 4

c) 1, 2, and 3

d) 1, 2, 3 and 4

Question 4:

A 26-year-old male presented with a large, round, firm, painless, right-sided testicular swelling associated with a dragging sensation. The left testis was normal. AFP was around 3000U. What is the next best step in the management?

a) FNAC

b) Scrotal trucut biopsy

c) High inguinal orchidectomy

d) Repeat tumor markers

Question 5:

A woman who had undergone cholecystectomy presented three days after the surgery with abdominal pain and tachycardia. WBC count is 11000 cells/mm3. USG showed fluid collection 5x5 cm in right sub-hepatic space. What is the next step in management?

a) Antibiotics and send home

b) USG guided pigtail catheter drainage

c) Laparoscopy and re-exploration

d) ERCP and T-tube placement

Question 6:

The instrument shown in the image is used to insufflate the abdomen for laparoscopic surgery. What is its name?



a) Veress needle

b) Lind's needle

c) Jacob needle

d) Mathew needle

Question 7:

A 20-year-old was brought to the ER after falling off his bicycle. His vitals were stable, but he had a bruise over the perineum and he had not passed urine yet. Blood was seen at the meatus and the bladder was not palpable. How do you manage him?

a) Wait and send him home

b) Immediate insertion of foleys catheter

c) Suprapubic aspiration of urine

d) Wait for bladder to fill and urge to urinate

Question 8:

A patient who was admitted after a road traffic accident is put on mechanical ventilation. He opens his eyes on verbal command and moves all four limbs spontaneously. Calculate his GCS.

a) Eyes-2, Verbal-1, Motor-5

b) Eyes-2, Verbal-NT, Motor-5

c) Eyes-3, Verbal-1, Motor-6

d) Eyes-3, Verbal-NT, Motor-6

Question 9:

What is the gauge size of the cannula used here?

a) 14

b) 16

c) 18

d) 20

Question 10:

A 55-year-old female patient presented with a 4x3 cm lump in the right upper outer quadrant, with no axillary lymph node involvement. Mammography revealed BIRADS 4b staging. She underwent breast conservation surgery, and the final HPE report showed high nuclear grade DCIS with necrosis and 10 mm margin clearance. What is the further management?

a) Adjuvant radiotherapy

b) Adjuvant chemotherapy

c) Follow up 6 monthly for 2 years and then yearly follow up

d) Trastuzumab therapy

Question 11:

A lady presents with abdominal pain. USG reveals multiple gall stones with no wall thickening. CBD diameter is 12 mm and serum bilirubin was 0.8. Alkaline phosphatase is 380 and GGT is raised by 5 times. What is the next best step of management?

a) MRCP

b) ERCP

c) Endoscopic ultrasound

d) Laparoscopic cholecystectomy

Question 12:

You are suturing a laceration in the ER using the interrupted suturing technique. What is the angle of needle placement?

a) 60 degrees

b) 70 degrees

c) 80 degrees

d) 90 degrees

Question 13:

A 26-year-old male patient was brought to the emergency department with abdominal pain and obstipation for 3 days. He gives a history of bull gore to the abdomen 3 days back. His X-ray is given below. What is the probable diagnosis?



a) Hemothorax

b) Hollow viscus perforation

c) Pneumothorax

d) Intestinal obstruction

Question 14:

A man presented with a history of stab injury over the arm. On examination, there was a hematoma present, and radial and ulnar pulses were not felt. A 3.5cm longitudinal tear is noted in the brachial artery. Which among the following is the best treatment option?

a) Primary repair with end to end anastomosis

b) Repair with great saphenous vein graft

c) Repair with prosthetic graft

d) Repair with saphenous vein patch

Question 15:

A 24-year old male patient was brought to the emergency department following a road traffic accident. His heart rate was 100/min, respiratory rate was 24/min, and his chest X-ray is given below. Which of the following is contraindicated in him?



a) Nasogastric tube insertion

b) Chest tube insertion

c) Log roll

d) Epidural anaesthesia

Question 16:

A patient was received in the casualty following a road traffic accident. His BP was 90/60 mmHg and his pulse rate was 120/min. The emergency room physician asked you to replace the fluid loss in this patient. What is the maximum flow rate of fluid with the IV canula given in the image?

a) 86ml/ min

b) 96ml/ min

c) 103 ml/ min

d) 63 ml/ min

Question 17:

The most common cause of bladder outlet obstruction in a male child is

a) Posterior urethral valve

b) Urethral atresia

c) Anterior urethral valve

d) Ureterocele

Question 18:

Which of the following statements are true about colon cancer?

a) 1, 2, and 3 only

b) 2, 3, and 4 only

c) 2 and 4 only

d) 1, 2, 3 and 4

Question 19:

Frozen section was done during the surgery of carcinoma breast. This procedure will be useful for all except

a) To know sentinel lymph node

b) To know margin positivity

c) To know metastasis

d) To confirm definitive diagnosis immediately in all cases

Question 20:

A 40-year-old woman presented with jaundice, fever, and abdominal pain. She is suspected to have cholangitis. Which of the following apart from abdominal pain makes up Reynold's pentad?

a) 1,2,3,4

b) 2,3,4,5

c) 1,3,4,5

d) 1,2,4,5

Question 21:

The following drain is placed after mastectomy. Identify the type of drain.

a) Open

b) Closed

c) Semi open

d) Semi closed

Question 22:

Consider the following about hepatocellular carcinoma (HCC) and choose the correct statements.

a) 1, 2, 3

b) 2, 3, 4

c) 1, 3, 4

d) 1 and 2 only

Question 23:

A 56-year-old patient with obstructive jaundice has a distended gall bladder and a dilated common bile duct and intrahepatic biliary radicles on ultrasound. Contrast-enhanced CT confirmed the same findings with no additional findings. Which of the following will be most useful to localize the cause?

a) Magnetic resonance imaging

b) Endoscopic ultrasound

c) Percutaneous transhepatic cholangiography

d) Endoscopic retrograde cholangiopancreatography

Question 24:

All of the following are lower urinary tract symptoms except

a) Sudden urge to urinate

b) Incontinence while asleep

c) Incontinence when patient lies down and coughs

d) Incontinence during intercourse

Question 25:

Fluorescence in-situ hybridization(FISH) is required in which of the following IHC interpretations of Her2/neu?

a) 1+

b) 2+

c) 3+

d) All of the above

Question 26:

A 50-year-old male patient came to the outpatient department with complaints of hematuria. A 2x2 cm bladder mass is seen which is a low-grade transitional cell carcinoma. Which among the following is the ideal management?

a) Transurethral resection of the tumour

b) Resection with ileal conduit

c) Partial cystectomy with bladder reconstruction

d) Neo-adjuvant chemotherapy

Question 27:

Which of the following statements are true?

a) 2, 3, 4

b) 1, 2, 3, 4

c) 4 only

d) 1, 3, 4

Question 28:

Which of the following is the correct method for measurement of nasogastric tube?

a) Tip of nose to ear lobule to xiphisternum

b) Angle of mouth to tragus to xiphisternum

c) Ala of nose to angle of mandible to xiphisternum

d) Angle of mouth to ear lobule to umbilicus

Question 29:

A surgeon performs an appendicectomy with a grid iron incision. He later converts the incision to Rutherford Morrison incision. Which of the following are cut?

a) 1, 3, 4

b) 2, 3, 4

c) 1, 2, 3

d) 1, 2, 4

Answer Key

Question No.	Correct Option
1	c
2	c
3	c
4	c
5	b
6	a
7	d
8	d
9	c
10	a

11	a
12	d
13	b
14	b
15	b
16	c
17	a
18	d
19	d
20	a
21	b
22	c
23	a
24	d
25	b
26	a
27	d
28	a
29	c

Detailed Explanations

Solution to Question 1:

Triage is used to classify the priority of treatment.

Triage is carried out at the site of a disaster to determine the priority for immediate medical attention, transportation, and hospital admission. This process involves the rapid classification of the injured based on the severity of their injuries and the likelihood of survival with timely medical intervention.

Its objective is to provide maximum benefit to the largest number of injured individuals in a disaster situation. Victims whose prognosis would be favorable with simple medical care are given higher priority, while patients in critical condition, who may not benefit significantly from medical attention, are given lower priority. An internationally accepted four-colour coded is commonly used, which includes:

• Red - High priority attention or tranfer

• Yellow - Medium priority

• Green - Ambulatory patients

• Black - Dead or moribund patients

Assessment of impact, rehabilatation and preparedness following a disaster are included in other phases of the disaster management cycle (options A, B and D).



Solution to Question 2:

The correct option is 1-C,2-D,3-A, 4-B

• Hirschsprung's disease - Rectal biopsy

• Posterior urethral valve - Micturating cystourethrogram

• Choledochal cyst - MRCP

• Intussusception - CT scan

Hirschsprung's disease:

• Rectal biopsy is the gold standard investigation for the diagnosis of Hirschsprung's disease. The affected part shows the absence of ganglion cells, hypertrophy of nerve trunks, and positive immunostaining for acetylcholinesterase.

• Water-soluble contrast enema: It shows the presence of a contracted aganglionic rectum, transition zone, and a dilated normally innervated colon.

• Anorectal manometry: It shows the failure of the relaxation of the internal anal sphincter.

Surgical intervention is the only definitive treatment available, and includes the Swenson, Duhamel, and Soave procedures.

Contrast enema image in Hirschsprung's disease-



Posterior urethral valve (PUV):

• Diagnosis of this condition is commonly made during the antenatal period with an ultrasound, which demonstrates bilateral hydronephrosis above a distended bladder. Pulmonary hypoplasia secondary to oligohydramnios may be detected in some complicated cases. In cases that are not detected during the antenatal period, these children usually present with urinary infection in the neonatal period or with uremia and renal failure.

• The confirmatory investigation for PUV is a micturating/voiding cystourethrogram. It demonstrates poor bladder emptying and dilated posterior urethra.

• It is treated by cystoscopic resection or ablation of the valve.

Preoperative(a) and postoperative(b) micturating cystourethrogram images in a patient with PUV is shown below. The dilated posterior urethra seen in the first image is resolved in the second image after surgery.



Choledochal cyst:

• MRI/MRCP is highly useful to access the biliary anatomy especially the relation between the lower end of the bile duct and the pancreatic duct.

• Ultrasound is helpful to identify the presence of an abnormal cyst.

• CT scan can be used to delineate the extent of intrahepatic or extrahepatic dilatation

MRCP image showing multiple intrahepatic biliary duct cystic dilatations-



Intussusception:

CT scan is the investigation of choice. It may show the characteristic target sign.

USG findings in intussusception include:

• Target sign (also known as the doughnut sign)

• Pseudokidney sign

• Crescent in a doughnut sign

Barium enema shows the characteristic coiled spring appearance.

USG image showing the target sign:



Solution to Question 3:

The clinical features of hematuria and a ballotable renal mass in a 3-year-old child are suggestive of Wilms tumor (Nephroblastoma). CECT abdomen is used to identify the extent of the tumor, integrity of the other kidney, and metastasis. Wilms tumor is metabolically active and can concentrate fluorodeoxyglucose therefore PET/CT is also helpful to identify regional spread and metastatic lesions. A biopsy is discouraged as it may lead to disease upstaging. 24-hour urinary vanillyl mandelic acid is measured to exclude neuroblastoma, which is an important differential diagnosis of Wilms tumor.

Wilms tumor is the most common primary malignant renal tumor in children, typically affecting children less than 5 years. After neuroblastoma, it is the second most common malignant abdominal tumor in childhood. It most commonly presents as an asymptomatic abdominal mass. Other findings include hypertension, abdominal pain, gross painless hematuria, and constitutional symptoms such as fever and weight loss.

Imaging studies done to diagnose Wilms tumor include plain radiography, USG, and CT of the abdomen to demonstrate the renal mass. USG with doppler study of renal veins is useful in the evaluation of collecting system and to look for tumor thrombi in renal veins and inferior vena cava. A chest CT is done preoperatively to screen for pulmonary metastasis. A positron emission tomography (PET) scan is useful in identifying regional spread and metastatic lesions. A biopsy is discouraged in this condition as it results in disease upstaging, and is considered only in cases of unusual presentation or imaging.

Neuroblastoma is an important differential diagnosis to consider, as it can present similarly to Wilms tumor. It is usually seen in preschool-aged children and commonly presents as a palpable abdominal mass. The retroperitoneum is the most common site for this tumor. Therefore, it is essential to test for elevated vanillyl mandelic acid (VMA) in the urine in this patient to rule out neuroblastoma.

Children's oncology group (previously known as National Wilms Tumor study group) staging system for Wilms tumor:

Treatment: Surgery and chemotherapy, with or without radiotherapy, are the mainstays of treatment.

• Radical nephrectomy is the standard surgical procedure done. Routine sampling of the para-aortic, hilar, and paracaval lymph nodes should be done.

• Nephron-sparing nephrectomy is done for children with bilateral disease or with a solitary kidney. Preoperative chemotherapy can be used in these cases to induce tumor shrinkage.

• The classical chemotherapy includes vincristine and dactinomycin with the addition of Adriamycin(Doxorubicin).

Stage I	Tumour confined to the kidney and completely resected with negative lymph nodes.
Stage II	Tumour extends beyond the kidney but is completely resected with negative margins and lymph nodes.
Stage III	Residual tumour present following surgery confined to the abdomen, including gross or microscopic tumour.
Stage IV	Hematogenous metastases (lung, liver, bone, brain, etc.) or lymph node metastases outside the abdominopelvic region.
Stage V	Metastasis to opposite kidney (bilateralrenal involvement).

Solution to Question 4:

The clinical features of a male patient presenting with a large, round, firm, and painless testicular mass with elevated AFP levels are suggestive of a non-seminomatous germ cell tumor (NSGCT). Among the given options, a high inguinal orchidectomy would be the next best step in the management.

An ultrasound scan (USG) of the testis is mandatory to confirm the diagnosis of a testicular tumor and assess the contralateral testis. Since USG is not given as an option, the next best step in this scenario would be a high inguinal orchidectomy. FNAC and trucut biopsy are contraindicated due to risk of seeding of the scrotrum and upstaging of the cancer.

Radical inguinal orchiectomy is the gold standard treatment for excision of the primary tumor. Sperm cryopreservation should be offered before surgery.

After removal of the affected testis, a metastatic workup is done with CT of the chest, abdomen, and pelvis to stage the disease as follows:

• Stage I: Localized tumor with no lymph node involvement or distant metastasis

• Stage II: Regional lymph node metastasis

• Stage III: Distant metastasis or nodal metastasis with elevated tumor markers

Further management is based on staging with treatment options including active surveillance, combination chemotherapy with bleomycin-etoposide-cisplatin and retroperitoneal lymph node dissection.

Solution to Question 5:

This clinical scenario involves a patient presenting with abdominal pain and tachycardia in the post-operative period following cholecystectomy. An ultrasound scan (USG) reveals a fluid collection in the right sub-hepatic space, strongly suggesting a bile duct injury following cholecystectomy. As the patient presents more than 48 hours after the surgery, the next best step in managament would be to drain the fluid collection using a USG-guided pigtail catheter.

More than 80% of bile duct injuries occur during cholecystectomy. A bile duct injury can result in either bile leakage or stricture. Patients with bile leakage may present with fever, increasing abdominal pain, jaundice or bile leakage from an incision.



Management strategy:

• Imaging: Ultrasonography to assess for a fluid collection and to evaluate the biliary tree

• Control of infection, limiting inflammation

• Percutaneous drainage of periportal fluid collections

• Parenteral antibiotics

• Clear and thorough delineation of entire biliary anatomy

• Endoscopic retrograde cholangiopancreatography (ERCP), especially if cystic-duct stump leak is suspected. It is both diagnostic and therapeutic.

• Magnetic resonance cholangiopancreatography (MRCP)

- HIDA scanning is also helpful to identify and quantitate bile leaks.
- Definitive surgical correction
- Roux-en-Y choledochojejunostomy can be done for benign strictures or duct transections.
- Balloon dilatation with placement of a stent can be done for strictures of recent onset.



Solution to Question 6:

The instrument shown in the image is the Veress needle which is used to insufflate the abdomen for laparoscopic surgery.

The image below shows the Veress needle:



There are two methods for establishing abdominal access during laparoscopic surgeries:

• Direct puncture using a Veress needle

• Direct peritoneal access (Hasson's technique)

Direct puncture laparoscopy:

In this procedure, first, the abdominal wall is grasped with towel clips or fingers to elevate it away from the underlying structures. Then, a small incision is made in the umbilicus, and insufflation is performed using a Veress needle. When introducing the Veress needle, two distinct pops can be felt as the needle pierces the abdominal wall fascia first and then the peritoneum. The umbilicus is the preferred location for introducing a Veress needle because the abdominal wall is thinner at this site.

Laparoscopic surgeries can be performed under general anesthesia or local anesthesia; although, general anesthesia is preferred. Carbon dioxide gas is usually used for insufflation, and the maximum pressure is typically in the range of 14-15 mm Hg. However, if the procedure is done under local anesthesia, nitrous oxide is used. Insufflation with nitrous oxide is stopped at a pressure of 10 mmHg or when 2L of gas has been insufflated.

Direct peritoneal access technique (Hasson technique): The surgeon makes a small umbilical incision and gains access to the peritoneal cavity under direct vision by dissecting through the abdominal fascia and peritoneum. It is preferred in patients who have undergone previous abdominal operations where the small bowel may be adherent to the undersurface of the abdominal wound.

Solution to Question 7:

The clinical features of a perineal bruise and blood at the urethral meatus following a fall from a bicycle are suggestive of a bulbar urethral injury. In this case, since the bladder is not palpable, the best option would be to wait for the bladder to fill and for the urge to urinate to occur. Once the bladder is full, the urine should be drained using a suprapubic catheter.

The image given below shows the various parts of the urethra:



Urethral injuries can be classified into:

A retrograde urethrogram with water-soluble contrast is used for confirmation of the diagnosis.

Type	Proximal urethral injury	Distal urethral injury
Site of injury	Prostatic urethraMembranous urethra	Bulbar urethraPenile urethra
Mechanism	Secondary topelvic fracture	Occurs due to a blow to the perineum by:Straddle injuryLoose manhole coveringGymnasium accidents
Clinical features	Urinary retentionBlood at the urethral meatusHigh-riding prostate on digital rectal examination (Vermooten sign)Deep perineal hematomaInjury to themembranous urethraleads to accumulation of urine in thedeep perineal pouch.	Urinary retentionBlood at the urethral meatusPerineal hematoma (Butterfly hematoma)



Solution to Question 8:

The patients GCS is Eyes-3, Verbal-NT, Motor-6.

It is calculated as follows:

• Eye opening: Opening to verbal command - 3

• Verbal response: In the case of endotracheal intubation, the verbal response score is noted as NT (non-testable)

• Motor respone: Movement of all 4 limbs spontaneously - 6

Total GCS score is not recorded since one of the components is non-testable.

Solution to Question 9:

The green IV cannula shown in the image corresponds to a gauge size of 18G.

Solution to Question 10:

The next step in the management in this patient with ductal carcinoma in situ (DCIS) would be adjuvant radiotherapy, according to the University of Southern California/Van Nuys Prognostic Index (USC/VNPI).

The USC/VNPI is a scoring system that evaluates the following four factors to predict recurrence in patients with DCIS who are undergoing conservative treatment:

• Tumor size

• Margin width

• Pathological classification (based on nuclear grade, and presence or absence of comedonecrosis)

• Age

This index also provides recommendations for managing patients based on the calculated score.

Scoring system for USC/VNPI:

The current recommendations for the management of DCIS according to this index are:

• Excision alone for scores 4,5, and 6

• Excision plus radiotherapy for scores 7,8, and 9

• Mastectomy for scores 10,11, and 12

In this case,

• Tumor size of 4cm (40mm) - score 2

• 10 mm margin clearance - score 1

• High nuclear grade with necrosis - score 3

• Age of 55 years - score 2

The total score is 8; hence radiotherapy is indicated after breast conservation surgery.

Parameters	Score 1	Score 2	Score 3
Size(mm)	≤15	16-40	≥41
Margin(mm)	≥10	1 - 9	<1
Class	Grade 1 or 2 without necrosis	Grade 1 or 2 with necrosis	Grade 3 with or without necrosis
Age(years	>60	40-60	<40

Solution to Question 11:

In this clinical scenario, a patient with gallstones, a dilated common bile duct, and an elevated alkaline phosphatase level raises suspicion for common bile duct stones. The next best step would be to perform magnetic resonance cholangiopancreatography (MRCP) to confirm their presence.

A dilated common bile duct (>8 mm) detected on ultrasound in a patient with gallstones, jaundice, biliary pain and an abnormal liver function panel (elevated bilirubin, alkaline phosphatase, and transaminases; but these can be normal in some cases) strongly indicates the presence of common bile duct stones (choledocholithiasis).

If there is uncertainty regarding the presence of bile duct stones, as in this case where bilirubin levels are normal, MRCP offers precise anatomical visualization and exhibits high sensitivity and

specificity for diagnosing choledocholithiasis.

While endoscopic retrograde cholangiopancreatography (ERCP) is also highly effective in diagnosing choledocholithiasis, due to its invasive nature and the risks associated with the procedure, it is rarely used as a purely diagnostic modality. Instead, it is reserved for cases in which a therapeutic intervention, such as stone extraction or sphincterotomy, is planned.

Treatment options:

• ERCP along with sphincterotomy and stone removal.

• Laparoscopic common bile duct exploration

• Open common bile duct exploration is only done when endoscopic and laparoscopic methods are not feasible or when concomitant biliary drainage is needed.

Solution to Question 12:

In the interrupted suture technique, the needle is placed at an angle of 90 degrees (at right angles).

Interrupted sutures:

The needle should be inserted and exit the skin at right angles. It should be rotated through the tissue rather than dragged to prevent unnecessary enlargement of the needle hole. The distance from the suture entry point to the wound edge should be approximately equal to the depth of the sutured tissue. There should be twice this distance between two consecutive sutures.

The image below shows interrupted suture:



Other suturing techniques:

Continuous Suture: The first suture is inserted in a manner identical to that in an interrupted suture. The remaining sutures are inserted in a continuous manner until the far end of the wound is reached. Care should be taken to maintain the tension of the suture to avoid excessive tension

or leaving it too slack; hence, the use of an assistant is warranted. An Aberdeen knot is used to secure the end of the suture.



Mattress sutures - They can be vertical or horizontal and are used to create either an eversion or inversion of the wound edges. The initial suture is inserted in the same way as an interrupted suture, but then, the needle either moves horizontally or vertically and traverses both edges of the wound once again, as seen in the image below. The main advantage of the mattress suture is that it produces an accurate approximation of wound edges, especially in settings where the edges are not easily approximated.



Subcuticular sutures: It is used for a better cosmetic appearance. It is also employed in situations where the edges of the wound are easily approximated. In this technique, small bites of the subcuticular tissues are taken on alternate sites of the wound, and then, they are gently pulled together. This causes an approximation of the wound edges without the risk of the cross-hatched markings of interrupted sutures.



Solution to Question 13:

The given clinical scenario involves a patient with a history of penetrating abdominal trauma and presenting symptoms of abdominal pain and obstipation. The erect radiograph shows air under the diaphragm which strongly points to a diagnosis of hollow viscous perforation.

Perforation of the alimentary tract is a life-threatening condition. This leads to the leakage of the enteric contents, thereby leading to acute bacterial peritonitis. Peritonitis is the generalized or localized inflammation of the peritoneum either due to the free spillage of fluids into the peritoneal cavity or secondary bacterial infection.

Clinical features include abdominal pain and distension, nausea, vomiting, tachycardia, tenderness, rebound tenderness, decreased or absent bowel sounds, and guarding on palpation. Fever may be present. Features of septic shock can be seen at later stages.

Investigations include estimation of the white blood cell count, urea, serum electrolytes, and an erect abdominal radiograph showing free gas under the diaphragm. A supine radiograph may show dilated gas-filled loops of the bowel suggestive of paralytic ileus. However, CECT is the most sensitive investigation to detect minimal/ suspected pneumoperitoneum.

Hollow viscus perforation



Management of a hollow viscous perforation mainly includes resuscitation, analgesics, intravenous broad-spectrum antibiotics, and emergency exploratory laparotomy.

Other options:

Option A- Hemothorax appears as a homogenous increase in opacity of the hemithorax.

Right Hemothorax

Option C- Pneumothorax is the collection of air in the pleural space. On radiography, there may be increased lucency on the affected side, depression of the affected hemidiaphragm, and mediastinal shift to the opposite side. Associated rib fractures might also be visible.

Left Pneumothorax



Option D- Intestinal obstruction

• In small bowel obstruction, the obstructed bowel loops are centrally located and lie transversely. In cases of jejunal obstruction, valvulae conniventes, which are mucosal folds that span the entire diameter of the bowel, create a ladder effect. Additionally, a distended cecum may exhibit a gas shadow in the right iliac fossa.

Small bowel obstruction



• In large bowel obstruction, the presence of haustrations is a characteristic feature. Haustral folds are irregular and do not extend across the entire diameter of the bowel. However, it is important to note that haustral folds are absent in the caecum.



Solution to Question 14:

In this patient, who has sustained a long tear to the brachial artery due to a stab injury, the best treatment option is to repair it with a great saphenous vein interposition graft harvested from the patient's uninjured leg.

The most desirable graft used is the great saphenous vein . A native vein graft is preferable to a prosthetic graft because:

• Its elastic properties are comparable to that of an artery and enable pulsatile flow

• The diameter is similar to that of an extremity artery

• It is not thrombogenic and has long term patency

Other grafts that can be used are cephalic vein, lesser saphenous vein, and prosthetic conduits. Cephalic vein is not the first choice because it is less muscular than greater saphenous vein, and might also compromise the upper limb venous access in the already injured patient. Both the cephalic and the lesser saphenous veins present problems while harvesting in a trauma patient.

Other options:

Option A: Primary arterial repair with end-to-end anastomosis can be done only if the repair is tension-free. If the length of the injured area is long, vessel injuries cannot be repaired by primary end-to-end anastomoses and instead require an interposition graft.

Option C: Prosthetic grafts are only preferred when there has been a trauma to the native saphenous vein or when there is a size mis-match. Polytetrafluoroethylene (PTFE) graft is commonly used.

Option D: Saphenous vein patch is used in the closure of the arteriotomy site during carotid endarterectomies.

Solution to Question 15:

The given radiograph shows elevation of the left hemidiaphragmatic line along with the bowel loops and stomach seen in the pleural cavity. The probable cause would be a diaphragmatic hernia following blunt injury due to a road traffic accident. Chest tube insertion is contraindicated in diaphragmatic hernia due to risk of injury to bowel.

Chest tube insertion/thoracostomy is indicated cases of pneumothorax, hemothorax, chylothorax, pleural effusion, and pleural empyema. Contraindications include pulmonary adhesions, pulmonary trauma, coagulopathy, and diaphragmatic hernia.

Chest radiography performed after the placement of a nasogastric tube can be helpful in identifying any herniation of the stomach into the chest. Log roll and epidural anesthesia are not contraindicated in diaphragmatic hernia.

Blunt trauma to the diaphragm results in a linear diaphragmatic rupture, which is typically large and accompanied by the herniation of abdominal contents into the chest. Acute diaphragmatic injuries require repair through the abdominal approach to address associated visceral injuries. The diaphragmatic edges are approximated using Allis clamps, and the defect is closed with a running 1-0 polypropylene suture.

Solution to Question 16:

The green IV cannula used in this hemodynamically unstable patient is an 18G cannula, which has a maximum flow rate of 103 ml/ min.

Solution to Question 17:

The most common cause of bladder outflow obstruction in a male child is the posterior urethral valve (PUV).

A PUV is a membranous fold in the posterior urethra which results from the persistence of the urogenital membrane. It is typically located just distal to the verumontanum and is the primary cause of bladder outlet obstruction in newborn males.

Diagnosis is most commonly made antenatally with ultrasound which demonstrates bilateral hydronephrosis, a distended bladder, and a dilated posterior urethra which appears as the keyhole sign on imaging. In severe cases, there is oligohydramnios and pulmonary hypoplasia during the antenatal period.

Posterior Urethral Valve - Keyhole Sign



If the diagnosis is not made antenatally, then babies typically present with urinary tract infections in the neonatal period or with uraemia and renal failure. Clinically, a poor urinary stream, dribbling of urine, and palpable bladder (due to diverticula) can be present.

The investigation of choice is micturating cystourethrography.

The diagnosis is best made during the micturition phase of micturating cystourethrography in lateral or oblique views so that the posterior urethra can be imaged adequately. The following findings are seen:

• Dilatation and elongation of the posterior urethra - equivalent of the ultrasonographic keyhole sign

• Linear radiolucent band corresponding to the valve - seen occasionally

• Vesicoureteral reflux - seen in 50% of patients

• Bladder trabeculation/diverticula

Posterior urethral valve - Micturating cystourethrogram



Treatment is initially supportive by inserting a urinary catheter to relieve symptoms of obstruction. Definitive treatment is by urethroscopic destruction of the valves with continuing lifelong supportive treatment of the dilated urinary tract, recurrent urinary infections and uremia.

Other options:

Options B and C- Urethral atresia and anterior urethral valve are rare congenital anomalies that can also lead to urinary obstruction.

Option D- Ureterocele is the cystic enlargement of the intramural ureter. It is common in females. They can present with features of urinary tract infection, bladder obstruction, or ureteral stones. Excretion urography will show cobra head appearance.

The image shown below depicts the "cobra head" sign in a patient with bilateral ureterocele.

Solution to Question 18:

All the 4 given statements on colon cancer are true:

• Right and left colon cancers present differently. Left-sided tumors commonly cause obstruction and present with changes in bowel habits or rectal bleeding. On the other hand, right-sided tumors typically present with iron deficiency anemia or a mass.

• While the majority of colorectal cancers occur sporadically, hereditary cases are also seen in patients with a family history of colorectal cancer. Genetic testing aids in early diagnosis for individuals with a positive family history.

• Metformin, a commonly used biguanide compound for diabetes mellitus, reduces the risk of cancer, including colorectal cancer. It achieves this by disrupting cellular energy supply through the inhibition of aerobic glycolysis.

• Liver is the most frequent site of distant metastasis for colorectal cancer, with metastases occurring via the portal venous system. The risk of hepatic metastasis increases with tumor size and grade. The lung is the second most common site of distant metastasis, also due to hematogenous spread. Carcinomatosis (diffuse peritoneal metastases) results from peritoneal

seeding.

Note: Regional lymph node involvement is the most common form of spread of colorectal carcinoma and usually precedes distant metastasis.

Solution to Question 19:

Frozen section analysis performed during breast carcinoma surgery is not suitable for definitive diagnosis due to technical challenges, potential quality issues in processing and staining, and the difficulty of obtaining accurate results.

Frozen section analysis is performed when a prompt biopsy report is required, allowing the surgeon to make immediate decisions regarding the further steps of management.

• Frozen section analysis plays a valuable role in assessing involvement of sentinel lymph nodes during surgery (option A). When a positive node is identified intraoperatively, the surgeon can proceed with immediate axillary lymph node clearance. On the other hand, if the sentinel node does not show tumor involvement, further axillary dissection can be avoided.

• It also helps in diagnosing nodal metastasis in clinically identified lymph nodes (option C). However, it may not detect subcapsular micrometastases.

• While performing breast-conserving surgeries, assessment of tumor margins (option B) using frozen sections is helpful in ensuring complete tumor excision.

Solution to Question 20:

Apart from abdominal pain, 1, 2, 3, and 4 form the Reynold's pentad which is seen in severe cholangitis.

The components of Reynold's pentad are:

• Right upper quadrant abdominal pain

• Fever

• Jaundice

• Hypotension

• Altered sensorium

Cholangitis is the infection of the common bile duct (CBD) secondary to obstruction and increased intraluminal pressure due to CBD stones. Cholangitis presents with the classical triad of fever, right upper quadrant pain, and jaundice, referred to as Charcot's triad. It is a medical emergency and if left untreated, it may progress in severity and presents with Reynold's pentad - Charcot's triad with hypotension and altered mental status due to sepsis.

Abdominal ultrasound is performed to check for the presence of biliary stones and bile duct dilatations. The patient should be started immediately on a first-line broad-spectrum antibiotic and rehydration, along with biliary drainage. Biliary drainage is done using endoscopic retrograde cholangiopancreatography (ERCP). Biliary stones are the cause of cholangitis the majority of the

time, which can be removed by endoscopic sphincterotomy during ERCP.

Solution to Question 21:

The Romo Vac drain, depicted in the image, is a closed type of drain commonly used after mastectomy.

Drains are placed at surgical sites to freely collect the air or fluids that may accumulate. This enhances wound healing by preventing the formation of unnecessary cavities. Drains are prophylactic in elective surgeries and therapeutic in emergency surgeries. Drains can be -

• Open - They are based on the principle of gravity

• Semi-open - Their action is based on capillary effect

• Closed - They carry out their actions through suction

Suction drains are commonly used in chest, breast, abdominal, thyroid, and plastic surgeries. They remove intraperitoneal secretions, anastomotic leaks, and blood from hemorrhage at the surgical site. Romo Vac drain is a negative suction drain closed drain kept in the subcutaneous tissue. Romo Vac drain is indicated in modified radical mastectomy, subcutaneous swelling excision, and amputations.

Special types of drains:

• Chest drains are connected to an underwater seal to aid the exit of air from the pleural cavity. They are used in pneumothorax, pleural effusion, and hemothorax.

• After surgical exploration of the bile duct, the sphincter of Oddi goes into spasm. To facilitate drainage of bile, a T tube may be used.

Drain insertion site care: The site of insertion of the drain is kept clean by drying and patting the drain insertion site with a clean towel after bath. The dressing is changed daily with folded gauze pieces placed below and above the drain site.

Removal of drains: The catheter at the site is milked for residual collections. It is then clamped and the suture holding the catheter is cut. The tube is then gently pulled out and any clots at the perforation end are removed. The time of removal of drains for various conditions is as given below.

• Drains used in surgeries like thyroidectomy are usually removed within 24 hours. Their main purpose is to cover postoperative bleeding.

• In a mastectomy, a drain is placed for 5 days to collect serous collections.

• Drains used to cover the colorectal anastomoses are placed for 5-7 days.

• Common bile duct T-tube drains are usually placed for 10 days.

• Drains for infections are removed when the infection subsides or when the content of the drain is minimal.

After the drain is removed, a folded gauze is placed over the drain insertion area. The dressing can be removed after 24 hours. The patient may be given an antibiotic ointment to apply over the area for a week.

Solution to Question 22:

The correct statements about hepatocellular carcinoma (HCC) are 1,3 and 4. Lenvatinib is used in the advanced stages of hepatocellular carcinoma (HCC).

HCC is the most common cause of primary liver cancer. It is more commonly seen in males and the peak age group is between 65-70 years. The main risk factor for HCC is cirrhosis and associated chronic liver damage due to chronic infection by hepatitis B or C virus (HBV or HCV) infection, alcohol abuse, metabolic syndrome, or hemochromatosis.

The incidence of HCC is increasing (statement 3) worldwide. This is contributed by the increase in HCV and HBV infections, along with lifestyle disorders like obesity. Obesity is associated with non-alcoholic fatty liver disease (NAFLD). This progresses on to non-alcoholic steatohepatitis (NASH) causing cirrhosis and hepatocellular carcinoma. Thus, NASH and NAFLD are risk factors of HCC (statement 4).

Barcelona-Clinic-Liver Cancer (BCLC) Classification is the commonly used staging system. It stages the tumors as follows:

• Very early HCC - Single, well-differentiated tumor <2cm

• Early HCC - Single tumor or a maximum of 3 nodules <3cm in diameter

• Intermediate HCC - Any other size or number confined to the liver

• Advanced HCC - Presence of cancer-related symptoms or tumors with macrovascular invasion (of any type, including branch, hepatic, or portal vein), lymph node involvement, or extrahepatic spread

• End-stage disease - Severe liver dysfunction with impaired quality of life

Treatment

Very early HCC is treated by ablation. Early HCC with a single nodule is resected but when there are multiple nodules over 3 cm, ablation or transplantation are considered.

Intermediate stage HCC is treated by chemoembolization. Transarterial chemoembolization(TACE) is used for multinodular lesions (statement 1).

Advanced stages of the disease with macrovascular invasion are treated with systemic therapies. Lenvatinib and sorafenib are first-line drugs. Supportive care is given at the terminal stage of the disease.

Solution to Question 23:

The given clinical scenario with USG and CECT showing a distended gall bladder, dilated common bile duct, and intrahepatic biliary radicles raises suspicion of common bile duct stones (CBD stones). Among the given options, magnetic resonance imaging (MRI/MRCP) will be next most useful modality to localize the cause and level of obstruction.

Magnetic resonance cholangiopancreatography (MRCP) uses T2-weighted MRI sequences and it is highly sensitive and specific in identifying CBD stones. MRCP provides visualization of dilated bile ducts and often enables accurate assessment of the level of occlusion in the biliary tree. It is a non-invasive test and is preferred by many surgeons to assess the need for endoscopic retrograde cholangiopancreatography (ERCP). ERCP is also highly effective at diagnosing CBD stones. However, due to the risks associated with the procedure, it is rarely used as a purely diagnostic modality, rather it is now used only as a therapeutic modality such as when stone extraction or sphincterotomy is planned (option D).

Ultrasonography (USG) can detect gallstones and assess the size of the common bile duct. If a patient with gallstones, jaundice, and biliary pain shows a dilated common bile duct (>8 mm in diameter) on USG, it strongly suggests the presence of CBD stones. CT scan is also a sensitive diagnostic tool for CBD stones. USG and CT will generally be the first line modalities due to their easier availability, and lower cost and time requirements.



Other options:

Option B- Endoscopic ultrasound is used for detecting CBD stones. It is considered as good as ERCP but it is invasive and requires expertise in endoscopic procedure and hence it is not done routinely.

Option C- Percutaneous transhepatic cholangiography (PTC) is rarely performed in patients with CBD stones. In patients where endoscopic or surgical procedures are contraindicated, it may be used for both diagnostic and therapeutic purposes.



Solution to Question 24:

Incontinence during intercourse is not a lower urinary tract symptom (LUTS).

The lower urinary tract symptoms (LUTS) can be classified as:

Storage LUTS: This occurs due to the failure of the bladder to act as a urine reservoir. It is commonly seen in the overactive bladder or bladder neuropathy. Symptoms related to storage LUTS are:

• Frequency - Increased frequency of voiding

• Nocturia - a patient has to wake up at least once at night to void

• Urgency (option A) - the sudden uncontrollable desire to pass urine

• Urge incontinence - a large volume of involuntary urine leakage preceded by urgency

• Stress incontinence (option C) - dribbling of urine during coughing, laughing, sneezing, or exercising. This occurs in women who have undergone vaginal deliveries.

• Nocturnal enuresis (option B) - involuntary leakage of urine during sleep. It is commonly seen in elderly men having chronic retention of urine with overflow incontinence.

• Strangury - is a sensation of constantly needing to void. Commonly occurs due to a lower urinary tract infection.

Voiding LUTS: This is commonly seen in bladder outlet obstruction (BOO). Symptoms related to voiding LUTS are:

• Hesitancy - difficulty in initiating micturition

• Reduced stream

• Straining - muscular effort is needed to initiate or to improve the urinary stream

Post micturitional LUTS: This is also caused due to BOO. Symptoms related to post micturitional LUTS are:

• Incomplete emptying - a sensation that the bladder is full still remains after micturition

• Post micturition dribble - Involuntary passage of urine after the patient has finished passing urine.

Solution to Question 25:

Fluorescence in situ hybridization (FISH) for gene amplification should be performed in all cases of breast tumors with an immunohistochemistry (IHC) score of 2+.

HER2/neu is a protein promoting rapid growth of breast cancer cells. All invasive breast cancers show undergo HER2/neu testing on biopsy or post-surgical removal. HER2 status is crucial for guiding breast cancer treatment. Two tests, immunohistochemistry (IHC) and fluorescence in situ hybridization (FISH), determine the presence of high HER2 protein levels. As FISH is costlier and more time-consuming, IHC is often performed as the initial test.

Her-2/neu IHC interpretation:

• 0 to 1+: Negative for HER2. These cancers do not respond to drugs targetting HER2 receptors.

• 2+: Equivocal. Hence, further testing with FISH is required to clarify the HER2 status.

• 3+: Positive for HER2. These cancers respond to drugs targetting HER2 receptors.

Transtuzumab and pertuzumab are monoclonal antibodies specifically used for HER2/neu-positive tumors.

Solution to Question 26:

This patient with low-grade 2x2 cm bladder cancer is likely to have non-muscle invasive bladder carcinoma (NMIBC) should undergo transurethral resection of the bladder tumor (TURBT) for accurate staging purposes.

Bladder cancer is graded (G1/2/3) based on level of differentiation. NMIBC includes stages Tis (in-situ), Ta (non-invasive papillary carcinoma), and T1 (invades lamina propria) while muscle invasive bladder cancer (MIBC) includes stages T2 (invades muscularis propria), T3 (invades perivesicular soft tissue), and T4 (invades adfjacent organs). NMIBC accounts for almost 70% of cases at presentation.

Treatment:

• NMIBC - The initial management of bladder tumors consists of transurethral resection of bladder tumor (TURBT) for accurate staging, which aids in determining the subsequent treatment plan. Intravesical therapy, involving chemotherapy (with mitomycin, doxorubicin, or epirubicin) or immunotherapy (BCG), is administered after TURBT to eradicate any remaining cancer cells. This therapy primarily targets the bladder lining and is not effective for cancers that have invaded deeper layers.

• MIBC - The standard treatment for muscle-invasive tumors is radical cystectomy and pelvic lymphadenectomy with urinary diversion (ileal conduit) (option B). Systemic adjuvant and neoadjuvant chemotherapy with M-VAC (Methotrexate, vinblastine, doxorubicin, and cisplatin) has shown to be beneficial (option D).

Option C: Partial cystectomy can be attempted in patients with focal bladder cancer, but the risk of recurrence remains high.

Solution to Question 27:

Statements 1, 3 and 4 are true.

Screening with noninvasive tests for common cancers aims to achieve early diagnosis, enable less invasive surgical treatments with reduced complications, and potentially improve surgical cure and overall survival rates.

Statement 1: Mammography as a routine screening tool has been found to lower breast cancer mortality at an acceptable cost. In the United Kingdom, an expert panel reported a 20% reduction in breast cancer mortality due to screening, despite the occurrence of overdiagnosis. The panel recommended continuing breast screening due to its significant benefits.

Statement 3: Visual inspection of the oral cavity is widely used for screening oral cancer. The target population includes individuals aged 30 and above who use tobacco and/or alcohol. Visual inspection is advantageous due to its feasibility, safety, acceptability, accuracy in detecting precancerous and cancerous lesions, and cost-effectiveness in reducing oral cancer mortality.

Statement 4: Pap smear is a screening test for cervical cancer. Cervical cancer screening should start at age 21. Women aged 21-29 should have Pap smear every 3 years, while women aged 30-65 should have both HPV and Pap co-test every 5 years.

Other statements:

Statement 2: American Cancer Society guidelines recommend colon cancer screening for average-risk individuals starting at age 50. Screening options include:

• Yearly fecal occult blood testing (FOBT) / fecal immunohistochemical testing (FIT)

• Flexible sigmoidoscopy every 5 years,

• FOBT/FIT and flexible sigmoidoscopy in combination every 5 years, or

• Colonoscopy every 10 years.

Solution to Question 28:

The correct method for measuring a nasogastric tube (Ryle's tube) is from the tip of the nose to the ear lobule to the xiphisternum.

Nasogastric (NG) tubes play a role in treating intestinal obstruction and can also be used to provide nutritional support. The commonly used method to determine the length of a nasogastric tube (NG tube) is the NEX method, which involves measuring from the tip of the nose to ear to

xiphisternum. The approximate length to be inserted is 55cms.

Ideally, Ryle's tube is inserted in the sitting position with the neck slightly flexed. Confirmation of the correct position is done by the following:

• Auscultation in the epigastric area while injecting a bolus of air

• An abdominal radiograph is the best way to confirm the location of the tube

An NG tube placed for suction should ideally be located within the stomach to avoid potential damage to the duodenum. For decreased risk of aspiration, the ideal location for an NG tube is postpyloric (55-60cm).

Indications for NG tube :

Drainage purposes:

• Conservative management of bowel obstruction and postop paralytic ileus

• Gastric decompression prior to emergency surgery

• Aspiration of gastric fluid

Feeding purposes:

• Nutritional support in patients in motor neuron disease, stroke, and following upper aerodigestive tract procedures

• Administration of medication

Contraindications:

Absolute:

• Severe midface trauma

• Recent nasal surgery

Relative:

• Coagulation abnormality

• Esophageal varices

• Recent banding of esophageal varices

• Anastomoses in the stomach and esophagus

Instruments tray used for nasogastric tube insertion:

Solution to Question 29:

The structures cut when the initial incision gridiron incision (McBurney's incision) is converted to the Rutherford Morrison incision are the external oblique, internal oblique, and transverse abdominis muscles.

When the preoperative diagnosis is reasonably certain, the widely used incision for appendectomy is known as the gridiron incision. It is made perpendicular to a line connecting the anterior superior iliac spine to the umbilicus, with its center at McBurney's point. When better access is needed, the gridiron incision can be converted to a Rutherford Morison incision by extending the incision obliquely upwards and laterally.

Rutherford Morison's incision is suitable for a para- or retrocecal appendix that is fixed. It is an oblique muscle-cutting incision starting from McBurney's point and extending upward and laterally as needed. All layers (skin, subcutaneous tissues, external oblique, internal oblique, transversus abdominis, and the peritoneum) are divided along the incision line.

In recent years, the transverse skin crease (Lanz) incision has gained popularity due to improved exposure and ease of extension when necessary. This incision is placed approximately 2 cm below the umbilicus along the midclavicular-midinguinal line, with its length adjusted according to the patient's size and obesity. If required, the incision can be extended medially by retracting or dividing the rectus abdominis muscle (statement 4).

When there is uncertainty in the diagnosis, especially in the presence of intestinal obstruction, a lower midline abdominal incision is preferred over a right lower paramedian incision.


Incisions used in appendicectomy
Grid iron incision
Lanz incision
Rutherford Morrison's incision
Lower midline abdominal incision
©MARROW

Surgery NEET 2021

Question 1:

A patient underwent surgery for breast carcinoma. She now presents with painless swelling involving the arm as shown in the image below. What is the most likely diagnosis?

a) Lymphedema

b) Lymphangiosarcoma

c) Venous thrombosis

d) Cellulitis

Question 2:

Which of the following statements is true regarding the condition given below?

a) Patient must be started on anti-gas gangrene serum therapy

b) It is a mixed flora aerobic and anaerobic infection

c) Bilateral orchidectomy is generally performed

d) Compulsory urinary diversion is performed

Question 3:

A baby playing unsupervised was brought with complaints of difficulty in swallowing for the last few hours. The X-ray image is given below. What is the diagnosis?



a) Foreign body in the trachea

b) Foreign body in the esophagus

c) Artifact

d) Soft tissue calcification in the neck

Question 4:

A 75-year-old man with prostate carcinoma presents to you with a PSA of 9 ng/mL with a small tumor focus. His Gleason score is 6. What will be your most likely management?

a) Radical prostatectomy

b) External beam radiation

c) Brachytherapy

d) Active surveillance

Question 5:

Identify the gross specimen obtained from a cystoprostatectomy.

a) Malakoplakia

b) Bladder carcinoma

c) Prostate carcinoma

d) Schistosomiasis

Question 6:

A patient after a road traffic accident presented with pain abdomen. The resident examines the child and found that vitals were stable and tenderness was present in the left lumbar region. Which is the best investigation of choice?

a) Contrast enhanced CT scan

b) Retrograde urethrogram

c) Wait and watch

d) Emergency laparotomy

Question 7:

A patient with rectal prolapse underwent surgery. The postoperative image is given below. Identify the procedure.



a) Stapled hemorroidopexy

b) Well's procedure

c) Thiersch wiring

d) Altemeier repair

Question 8:

A patient presents after a motorcycle road traffic accident with no reported pain or injury. During examination, blood is observed at the tip of the urethral meatus, and the bladder is

non-palpable. What would be your next step in management?

a) Foley's catheterization

b) Wait for bladder to fill

c) Nephrogram

d) Suprapubic cystostomy

Question 9:

A patient presents with a hernia medial to the inferior epigastric artery. What is the most likely diagnosis and appropriate management for the same?

a) Direct hernia and Bassini repair

b) Direct hernia and Lichtenstein mesh repair

c) Indirect hernia and Bassini repair

d) Indirect hernia and Lichtenstein mesh repair

Question 10:

A patient following a road traffic accident presented with subcutaneous emphysema. On auscultation, there was no air entry on the right side. Vitals were unstable. What will be your immediate management?

a) Insertion of wide bore needle in the 5th intercostal space (ICS)

b) Wide bore needle decompression and IV fluids

c) Intubation and positive pressure ventilation

d) eFAST

Question 11:

A 54-year-old female patient presents with abdominal pain and bilious vomiting. She had a history of abdominal hysterectomy 2 years back. On examination, abdomen auscultation revealed high-pitched bowel sounds. Investigations showed air within the biliary tree. What is the most likely diagnosis?

a) Gallstone ileus

b) Adhesive intestinal obstruction

c) Ischemic enterocolitis

d) Diverticulitis

Question 12:

In a patient with abdominal pain and sterile pyuria, what is the finding shown in the image given?

a) Putty kidney

b) Nephrocalcinosis

c) Staghorn calculus

d) Psoas calcification

Question 13:

A patient was found to have a mass arising from the antrum of the stomach which involves the head of the pancreas and metastasis to the right lobe of the liver. How will you manage this patient?

a) Palliative gastrojejunostomy followed by chemotherapy

b) Radical gastrectomy

c) Whipple's procedure

d) Gastrectomy with right hepatectomy

Question 14:

A patient underwent cholecystectomy for gallstones over 2 years ago. She now presents with symptoms of obstructive jaundice. Investigation reveals a stone in the common bile duct. What is the stone most likely to be?

a) Primary

b) Secondary

c) Tertiary

d) Retained

Question 15:

A 70-year-old female patient presents with pain in the thigh during walking. Investigation reveals the following finding as given below. What will be the best surgical management?



a) PTA with stenting

b) Iliofemoral bypass

c) Aortofemoral bypass

d) Aorto-iliac bypass

Question 16:

A young air-hostess presented to the outpatient department with dilated veins on her left leg (image). According to the CEAP classification for varicose veins, what will be the C stage in this patient?

a) C2A

b) C2S

c) C4a

d) C1

Question 17:

An alcoholic patient presented with severe abdominal pain. Pancreatitis was suspected and a CT scan revealed lesions with fluid collected near the body of the pancreas. Which of the following enzymes is most likely to be elevated in this scenario?

a) Serum GGT

b) CEA

c) Serum lipase

d) Serum bilirubin

Question 18:

A 25-year-old patient presents with abdominal pain and fever for 3 days duration. The chest X-ray in the erect position is given below. What will be the most likely diagnosis?



a) Gastric volvulus

b) Hollow viscus perforation

c) Liver abscess

d) Empyema thoracis

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	d
5	b
6	a
7	c
8	b
9	b
10	a
11	a
12	a
13	a
14	a

15	c
16	a
17	c
18	b

Detailed Explanations

Solution to Question 1:

The generalized swelling of the left arm of the patient post-surgery for breast carcinoma is suggestive of post-mastectomy lymphedema.

Lymphoedema occurs when interstitial fluid accumulates due to defective lymphatic drainage. It is a common complication of breast cancer treatment, involving axillary lymph node dissection and radiotherapy. Sentinel lymph node dissection (SLND) reduces the risk of lymphedema significantly.

Clinical signs and symptoms are observed in the ipsilateral arm. A significant increase in limb size due to swelling is seen. Other symptoms are heaviness, tightness, numbness, tingling, limb fatigue, and reduced limb mobility.

Diagnosis is predominantly based on the clinical swelling of the arm. The girth of the affected arm >2 centimetres than the normal limb, a 200-mL or more increase in limb volume, or a 5% or greater limb volume change are other diagnostic criteria.

Management:

• Limb elevation, elastic arm stockings, and pneumatic compression devices to reduce swelling.

• Using a low-pH water-based moisturizer to prevent dry skin caused by excess fluid accumulation.

• Manual lymphatic drainage (MLD) by a therapist followed by simple lymphatic drainage (SLD) by the patient daily aids interstitial fluid evacuation.

• Surgical options include lymphaticovenular anastomosis, liposuction, and debulking.

Other options:

Option B: Stewart-Treves syndrome refers to lymphangiosarcoma seen in chronic long-standing lymphedema following mastectomy (more than 10 years). They present as single/multiple bluish-red skin and subcutaneous nodules. They can ulcerate and hemorrhage.

Option C: Axillary venous thrombosis or Paget-Schrotter disease is seen in patients doing rigorous exercise and in weightlifters. The compression of the subclavian vein at the thoracic outlet causes painful swelling of the arm.

Option D: Cellulitis (lymphangitis) is a bacterial infection of the skin and subcutaneous tissue. It is characterized by a painful area of erythematous and edematous tissue.

Solution to Question 2:

The given image of extensive perineal soft tissue destruction with intact testis is suggestive of Fournier's gangrene, which is a mixed flora of aerobic and anaerobic infection.

Fournier's gangrene is a type of necrotizing fasciitis (Dartos, Scarpa, and Colle's fascia) of the perineum, external genitalia, and perianal region. Risk factors include:

• Immunocompromised states such as HIV, alcoholism, diabetes, etc.

• Minor procedures in the perineal area (opening of a periurethral abscess).

Pathogenesis is due to the obliteration of arterioles supplying the scrotal skin caused by fulminating subcutaneous polybacterial infection. After a while, the blackish discolouration of the scrotum (gangrene) with crepitus can be seen (image). The testis is usually not affected.

Clinical features of Fournier's gangrene are scrotal pain, fever, prostration, and pallor. Cellulitis spreads rapidly and may be associated with eschars, necrosis, flaking skin, and crepitus.

Diagnosis of Fournier's gangrene is majorly clinical and is a surgical emergency. It should be aggressively managed by urgent surgical debridement and skin grafting in later stages. The infection is controlled by the administration of broad-spectrum antibiotics.

Other options:

Option A: Anti-gas gangrene serum refers to refined equine antitoxin against Clostridium species. Hence, it has no role in Fournier's gangrene.

Option C: Bilateral orchidectomy is not performed as testes are not affected and need not be removed.

Option D: Urinary diversion may be necessary in a few cases, and is not compulsory.

Solution to Question 3:

The clinical picture of the child presenting with difficulty in swallowing and the chest X-rays (AP and lateral) showing a circular homogenous opacity in the neck region is suggestive of a foreign body (coin) lodged in the esophagus.

Foreign bodies commonly ingested by children are coins, disk batteries, and small toy items. In adults, it is usually a food bolus (mostly meat) that gets lodged.

The flat surface of a coin is seen on the anteroposterior view (AP view) and the edge is seen on the lateral view. This indicates that it is oriented in the coronal plane, which is seen when it is lodged in the esophagus (and not in the trachea).

Initial symptoms are choking, gagging, and coughing followed by excessive salivation, dysphagia, food refusal, vomiting, or pain in the neck and throat. Respiratory symptoms such as stridor, wheezing, cyanosis, or dyspnea are seen if the esophageal foreign body impinges on the larynx or tracheal wall.

Imaging with chest and abdominal X-rays should be performed in AP and lateral positions. Chest CT should always be performed if oesophageal perforation is suspected.

Endoscopic removal of the esophageal foreign body is the procedure of choice. It should be performed

• Immediately in esophageal obstruction, disk batteries, and sharp pointed objects.

• Within 12-24h in non-pointed objects, pointed objects in the stomach, and coins in the esophagus.

• Non-urgent removal in objects in stomach >2.5cm in diameter, and disk battery in the stomach (up to 48h)

Other options:

Option A: For coins lodged in the trachea, the edge is seen anteroposteriorly view and the flat side is seen in the lateral view of the X-rays (oriented in the sagittal plane).

Option C: Artifacts refer to an appearance or unintended structure that is not normally seen on the radiograph and is produced by some artificial means. It can occur due to technical or processing errors while taking a radiograph.

Option D: Soft tissue calcification in the neck is due to physiologic or pathological conditions. Commonly encountered soft tissue calcifications in the head and neck region include tonsilloliths, carotid artery calcifications, and calcified stylohyoid ligaments.

Solution to Question 4:

The given clinical vignette of a 75-year-old man, with a low PSA (<10 ng/ml), Gleason score of 6, and a small tumor focus point to a diagnosis of low-risk prostatic carcinoma. The most likely management for this patient would be active surveillance only.

Active surveillance is done when there is:

• Gleason score of 6

• Early-stage disease (T1c)

• Small volume disease on biopsy

• Low PSA (< 10 ng/ml)

Active surveillance protocols include a digital rectal examination and PSA monitoring every 3 to 6 months, with TRUS-guided prostate biopsies every 1 to 3 years.

Gleason grading system evaluates the abnormalities in the glandular architecture of the prostate. It is based on a scale of 1 to 5, with 1 being the most differentiated grade and 5 being the least differentiated. The grades are added to give the Gleason score.

The classification and management of carcinoma prostate are outlined below:

EBRT: External beam radiation therapy, TURP: Transurethral resection of the prostate

Grade	Treatment
Low Risk	Radical prostatectomy (<70 years),Active surveillance (>70 years)
Intermediate Risk	Radical prostatectomy and/or Radical radiotherapy(EBRT) (<70 years)Active surveillance +/- TURP and hormone therapy (>70 years)
High Risk	Early androgen ablation + Radiotherapy + Surgical treatment (< 70 years)
Metastatic disease	Systemic chemotherapy with palliative treatment (<70 years)Androgen ablation therapy and palliative treatment for symptomatic patients

Solution to Question 5:

The given image of the cystoprostatectomy specimen shows a large, fungating growth from the bladder wall involving the entire bladder. The two lobes of the prostate are not affected. The given specimen is that of bladder carcinoma.

Clinical features of bladder carcinoma include painless, gross hematuria (most common symptom), which may cause clots and urinary retention subsequently. Increased frequency and discomfort during urination may be seen.

Depending upon the depth of invasion, bladder carcinoma is divided into 3 groups:

• Non-muscle invasive bladder cancer - does not invade detrusor muscle but can invade lamina propria. It accounts for 70% of cases.

• Muscle-invasive bladder cancer - invades the detrusor muscle

• Flat, non-invasive carcinoma in situ (CIS)

Staging and diagnosis:

It is done by cystoscopy and resection of the tumor. A bimanual examination should be performed after resection of the tumor. Complete urinary tract imaging is required using contrast computed tomography (CT) or magnetic resonance (MR) urography.

Management:

• Non-muscle invasive bladder cancer:

• Transurethral resection of bladder tumor (TURBT) followed by intravesical chemotherapy with mitomycin C or epirubicin

• Immunotherapy with regular intravesical instillations of Bacille Calmette Guerin (BCG) in high-risk tumors.

• Muscle invasive bladder cancer: Radical cystectomy along with urinary diversion procedure and lymph node dissection +/- neoadjuvant chemotherapy

Other options:

Option A: Malakoplakia is a rare inflammatory condition that is typically seen in immunocompromised individuals. The term 'malakoplakia' means 'soft plaques'. General principles of treatment include systemic antibiotics, surgical excision, and limiting immunosuppression. Surgical excision is done for malakoplakia lesions based on the site affected, complications, and if medical treatment fails.

Option B: Prostatic carcinoma can be surgically managed by radical prostatectomy, which does not involve cystectomy. In prostatic carcinoma usually, one or more of the lobes are predominantly involved unlike in the given gross specimen.

Option D: Schistosomiasis of the bladder is a parasitic infection by Schistosoma hematobium. On cystoscopy, it can look like nodules, pseudo tubercles, ulceration, sandy patches, and in the end stage can cause squamous cell carcinoma of the bladder.

Solution to Question 6:

The given clinical vignette with abdominal pain and tenderness in the left lumbar region following a road traffic accident is suggestive of blunt renal trauma. The investigation of choice is CECT in a hemodynamically stable patient.

Blunt trauma is the main cause of renal injuries and is generally self-limiting. Radiological investigations are indicated when there is gross haematuria, shock in combination with microscopic haematuria, pediatric renal trauma, and penetrating injuries.

CECT is the investigation of choice. It assesses the extent of the injury, showing laceration, extravasation, surrounding hemorrhage, and vessel injury. It also shows non-renal injuries and effectively stages renal pedicle injuries. Arterial occlusion will manifest as rim enhancement of the normal renal contour.

Blunt trauma is mostly treated conservatively. Most of the penetrating injuries require surgical renal exploration.

Absolute indications for surgical intervention in renal trauma are:

• Massive hemorrhage

• Renal pedicle avulsion (grade 5 injury)

• Expanding, pulsatile, or uncontained retroperitoneal hematoma

Relative indications are:

• Large laceration of the renal pelvis or avulsion of the ureteropelvic junction

• Bowel or pancreatic injuries

• Persistent urinary leakage, postinjury urinoma, or perinephric abscess

• Complete renal artery thrombosis of both kidneys

• Renal hypertension

Other options:

Option D: Exploratory laparotomy is recommended in unstable patients and those with penetrating renal injuries.

Solution to Question 7:

The postoperative image shows a purse-string suture which is suggestive of Thiersch wiring for rectal prolapse.

In Thiersch's wiring, a steel wire or a silastic/nylon tape is placed around the anal canal. It is rarely done now due to post-operative problems like chronic perineal sepsis, anal stenosis, and obstructed defecation.

Rectal prolapse is of two types:

• Partial rectal prolapse - only the mucosal layer of the rectum is prolapsed. It is seen in children and is managed by digital repositioning. If there is a recurrence, then Theirsch's wiring or sclerotherapy is performed.

• Complete rectal prolapse - all the layers of the rectum are involved. It is seen in adults due to the weakening of the pelvic floor. It is managed by using the following approaches:

• The perineal approach - easy to perform, and has fewer complications but a high recurrence rate. It is preferred in elderly patients with comorbidities. The surgeries done include:

• Thiersch's wiring

• Delorme's mucosectomy - In this procedure, the rectal mucosa is stripped circumferentially from the rectum over the length of the prolapse. The underlying muscle is plicated with a series of sutures. The excess rectal mucosa is excised and a mucosal anastomosis is performed.

• Altemeier's recto sigmoidectomy (option D) - In this procedure, full-thickness resection of the prolapsed rectum is performed, incorporating any associated colonic prolapse. Restoration of colorectal continuity can be performed by hand-sewn/stapled anastomosis. This is the procedure of choice in patients presenting with incarcerated and strangulated prolapse.

• The abdominal approach - It is difficult to perform and has more complications, but a lower recurrence rate. The surgeries done include:

• Ripstein's procedure - It involves the placement of a prosthetic mesh around the mobilized rectum, with the attachment of the mesh to the presacral fascia below the sacral promontory.

• Well's procedure (option B) - It involves posterior rectopexy.

Other options:

Option A: Stapled hemorroidopexy is used in the treatment of internal hemorrhoids, where a circular stapling device is used to excise a cylinder of mucosa and submucosa and simultaneously staple the mucosal ends together.

Solution to Question 8:

The given clinical scenario showing blood at the tip of the urethral meatus following a road traffic accident (RTA) is a clinical hallmark of a urethral injury. The next best step in this scenario is to wait and watch.

Injuries to the urethra are more common in males. They are of:

Part of the urethra	Injury type	Clinical features
Bulbarurethra(distal urethral injury)	Blunt traumato the perineum	Butterfly hematoma(perineal bruising), urinary retention, and bleeding from the urethral tip
Bulbomembranousjunction(proximal urethral injury)	Road traffic accidents	Blood at the tip of the meatus and urinary retentionHigh-riding prostateon digital rectal examination (Vermooten sign)



Retrograde urethrography using water-soluble contrast is used for localizing the site of urethral injury.

The patient should be managed with analgesia and antibiotics. If the bladder is full, it should be drained with a catheter placed by percutaneous suprapubic puncture using a Seldinger technique.

Other options:

Option A: Foley catheterization should not be done since it might convert a partial tear into a complete tear.

Option C: Nephrogram is not done but retrograde urethrogram is performed in urethral injury.

Option D: Suprapubic Cystostomy is done only in cases where the bladder is full or the patient is unable to urinate following a pelvic fracture/ RTA.

Solution to Question 9:

The given clinical scenario of a hernia medial to the inferior epigastric artery is suggestive of a direct hernia, for which Lichtenstein mesh repair is the appropriate surgical method for its management.

The contents of a direct hernia protrude outward, forward, and medial to the inferior epigastric artery and the deep inguinal ring. Whereas, the contents of an indirect hernia protrude lateral to the inferior epigastric artery. The sac of an indirect inguinal hernia passes from the deep inguinal ring obliquely towards the superficial inguinal ring and then to the scrotum.

A direct hernia protrudes through the Hesselbach triangle, which is a zone of weakness for direct hernias.

A Lichtenstein mesh repair is currently one of the most frequently used methods for treating an inguinal hernia. It is done for the reconstruction of the inguinal floor in adults. It is an anterior tension-free mesh repair that reduces the chances of recurrence of a hernia. A piece of prosthetic nonabsorbable mesh is used to bridge the defect in this repair.



Other surgical methods used for the repair of an inguinal hernia are:

• Bassini repair - It is done for the reconstruction of the inguinal floor in adults. In this method, a mesh is not used. It is associated with a high recurrence and is not used frequently now. It is used only in cases of hernial strangulation and peritonitis where the use of a mesh is not preferred. It is performed by suturing the transversus abdominis and internal oblique musculoaponeurotic arches/conjoined tendon to the inguinal ligament.

• Shouldice repair - It is a multilayered repair, where the posterior wall of the inguinal canal is strengthened by continuous sutures. It involves double-breasting of the transversalis fascia. It is associated with a very low recurrence rate.

• Gilbert repair (plug and patch repair) - It is an extension of Lichtenstein's original mesh repair and has now become the most commonly performed primary anterior inguinal hernia repair. In this technique, a cone-shaped plug of polypropylene mesh is used. It is deployed like an upside-down umbrella when inserted into the internal inguinal ring to occlude the hernia

• McVay repair (Cooper's ligament repair) is used for the correction of direct inguinal hernia, large indirect hernia, recurrent hernia, femoral hernias (particularly suited for strangulated femoral hernias). In this method, the transversus abdominis aponeurosis is approximated to Cooper's ligament.

• Stoppa-Rives repair - It is used for the repair of large, recurrent, or bilateral hernias. In this method, a sub-umbilical midline incision is given to place a large mesh into the preperitoneal space.

• Herniotomy - It is used to treat inguinal hernia in a child which occurs due to the presence of a patent processus vaginalis. Therefore in herniotomy, the repair of the inguinal ring is not done.

Solution to Question 10:

The given clinical scenario of a patient with h/o road traffic accident (RTA) with no air entry on the right side and hemodynamic instability suggests a diagnosis of tension pneumothorax. This is managed by immediate decompression by insertion of a needle in the appropriate place.

Pneumothorax refers to the presence of excess air in the pleural space. Entrapment of air occurs by:

• Primary spontaneous pneumothorax - due to the spontaneous rupture of apical pleural blebs or cystic spaces immediately below the visceral pleura.

• Secondary pneumothorax - due to the rupture of dilated cystic spaces in lung diseases like chronic obstructive pulmonary disease (COPD)

• Tension pneumothorax - a one-way valve mechanism that directs airflow into the pleural space leading to a higher than atmospheric pressure environment, and prevents expansion of the lung.

Tension pneumothorax is characterized by chest pain, tachypnoea, tachycardia, and hypotension. On examination, the trachea is deviated away from the side of the injury, and breath sounds are absent on the affected lung. A patient can develop subcutaneous emphysema as a complication, secondary to the pneumothorax.

Tension pneumothorax is a surgical emergency and the excessive pressure has to be immediately relieved. Tension pneumothorax is managed by immediate needle decompression in the 5th intercostal space (ICS), followed by definitive management with a chest tube inserted in the same

space.



Other options:

Option B: While IV fluids are a part of the resuscitation protocol, hypotension in a tension pneumothorax is due to an obstructive type of shock, increasing circulating volume will have minimal effect on improving blood pressure. The management of the primary cause, tension pneumothorax, takes higher precedence as the next immediate step.

Option C: Intubation is performed when the patient is unresponsive post chest tube insertion.

Option D: Extended Focused Assessment with Sonography in Trauma (eFAST) is not required before emergency decompression as tension pneumothorax is a clinical diagnosis, but it can be performed following chest tube insertion.

Solution to Question 11:

The given scenario shows abdominal pain with bilious vomiting and high-pitched bowel sounds which are suggestive of intestinal obstruction along with radiographic features of pneumobilia (air within the biliary tree) pointing to the diagnosis of gallstone ileus.

Gallstone ileus is a condition that occurs secondary to the erosion of the gallbladder wall, which results in a fistula to the duodenum. Through this, a large gallstone can pass from the gallbladder to the duodenum and usually gets impacted in the distal ileum (the most common site), where it may cause intestinal obstruction.

Rigler's triad is a characteristic radiological feature seen in gallstone ileus and consists of:

• Pneumobilia - air present in the intestine passes into the biliary tree through the fistula

• Features of small bowel obstruction (abdominal pain, bilious vomiting, hyperechoic bowel sounds)

• Ectopic gallstone - usually in the right iliac fossa



Plain X-rays show air-fluid levels consistent with a small bowel obstruction and pneumobilia can also be visualized on CT imaging.

Gallstone ileus is managed by an explorative laparotomy to relieve the bowel obstruction. The stone is milked away from the site of impacted and then crushed or removed.

Other complications of gallstones include biliary colic, acute and chronic cholecystitis, empyema of the gallbladder, mucocele, perforation, acute cholangitis, and acute pancreatitis.

Other options:

Option B: Adhesive intestinal obstruction is seen after any abdominal surgery or any cause for peritoneal irritation, the features are consistent with bowel obstruction (air-fluid levels, featureless distal ileum) but there is no pneumobilia.

Option C: Ischaemic colitis on contrast-enhanced CT of the abdomen (CECT) shows bowel wall thickening, and signs of vascular insufficiency.

Option D: Diverticulitis on CT abdomen shows bright bowel outpouching, which may be complicated by infection and abscess formation.

Solution to Question 12:

A patient with abdominal pain with sterile pyuria and the abdominal X-ray showing homogenous ground glass-like calcification is suggestive of a putty kidney.

The putty kidney is a characteristic finding of genitourinary tuberculosis (GUTB). It is caused by the dissemination of the organism through the bloodstream and is always a secondary TB (reinfection or reactivation of old TB). Genitourinary TB usually presents as sterile pyuria.

Computed tomography (CT) of the abdomen is the most sensitive modality and is the preferred means of visualizing renal calcifications. Microbiological diagnosis is confirmed by at least 3 consecutive early-morning urine samples showing acid-fast bacilli using Ziehl–Neelsen (ZN) stain and subsequently cultured (gold standard).

GUTB is treated with a short course of anti-tubercular therapy (ATT). Ureteric strictures require double J stenting or percutaneous nephrostomy. Definitive treatment is after 3-6mn of ATT, where a reconstructive procedure (open surgical repair > balloon dilatation) is performed. In case the kidney is poorly functioning, a nephrectomy is done.



Option B: Nephrocalcinosis is the deposition of calcium in the kidney parenchyma or tubules, seen in hypercalcemic conditions like hyperparathyroidism.



Option C: Staghorn calculus (a triple phosphate stone) is found in urinary tract infections (UTI) by urease-producing pathogens like proteus and is not a cause of sterile pyuria.

Option D: Psoas calcification is a muscle calcification that develops after acute renal failure and rhabdomyolysis. It is not associated with sterile pyuria.



Solution to Question 13:

The given clinical scenario of a tumour arising from the pyloric antrum of the stomach which involves the head of the pancreas with liver metastasis is suggestive of metastatic carcinoma of the stomach and is managed by palliative gastrojejunostomy followed by chemotherapy.

Gastric carcinoma has a higher incidence in Japan and affects men more than women. H.pylori is a significant association, and other risk factors include smoking and industrial dust ingestion. The most common sites are - the proximal stomach (Western countries) and distal stomach (Japan and the rest of the world).

In the early stages, it usually presents as dyspepsia (non-specific). Vomiting and profound weight loss is seen in advanced cases. Endoscopy and biopsy of the lesion (gold standard) are performed to diagnose gastric carcinoma. Pre-operative staging is done with contrast-enhanced computed tomography (CECT) of the abdominal/pelvis. The most important prognostic indicators in gastric cancer are - lymph node involvement and depth of tumor invasion.

Gastric carcinoma is treated with surgical resection with lymphadenectomy. Based on the tumor location, different types of gastrectomy and anastomoses can be performed.

• Billroth I gastrectomy - the lower half of the stomach is removed + cut end of the stomach is anastomosed to the first part of the duodenum.

• Billroth II gastrectomy - two-thirds of the stomach is removed, the duodenal stump is closed and the stomach is anastomosed to the jejunum.

• Roux-en Y gastrojejunostomy - a loop of jejunum is anastomosed to a proximal gastric pouch







Other options:

Option B: Radical gastrectomy is done in proximal gastric cancer and linitis plastica.

Option C: Whipple's procedure is done for gastric carcinoma that has spread to the head of the pancreas. Since the patient already has metastases to the liver, this will not be the preferred option.

Option D: Gastrectomy with right hepatectomy is done in gastric cancer that has hepatic metastasis.

Solution to Question 14:

This patient has a history of cholecystectomy done for gallstones 2 years ago. Thus, in this patient with choledocholithiasis, the stones found in the common bile duct would most likely be primary stones.

Choledocholithiasis or common bile duct stones are classified based on their point of origin:

• Primary stones are brown pigment stones that arise de novo in the bile duct and are composed of bile pigments and cholesterol.

• Secondary stones (option B) are usually cholesterol stones, which originate in the gall bladder and pass down into the common bile duct. Since this patient has already undergone cholecystectomy more than 2 years ago, secondary stones are unlikely.

• When stones are found in the common bile duct within 2 years of history of cholecystectomy, they are termed retained stones (option D).

• If they are found more than 2 years after cholecystectomy, they are known as recurrent stones (which are mostly primary).

Clinical features: Mostly clinically silent, but may present with biliary colic and even obstructive jaundice. When there is fever, jaundice, and associated right upper quadrant abdominal pain, it is known as Charcot's triad, seen in cholangitis.

Management of choledocholithiasis:

ERCP +/- sphincterotomy followed by laparoscopic cholecystectomy is the most preferred management of choledocholithiasis.

If stones cannot be removed through ERCP and sphincterotomy, then common bile duct exploration is carried out intraoperatively either in an open procedure or via laparoscopy. A T-tube is left in place after choledochotomy.

Solution to Question 15:

The symptom of pain in the thigh while walking and the given angiography image showing common iliac artery obstruction is suggestive of peripheral obstructive vascular disease with aortoiliac obstruction. The best surgical course, in this case, is to perform an aortofemoral bypass.

Clinical findings of aortoiliac obstruction are :

• Intermittent claudication is the debilitating cramp-like pain felt in the buttocks, leg, and thigh muscles due to reduced blood flow.

• Femoral and distal pulses are absent in both limbs.

• Arterial bruit is heard over the aortoiliac region.

• Impotence (Leriche syndrome) may occur. Buttock claudication is present.

As the disease progresses, rest pain, ulceration, gangrene, and slow capillary refilling may be seen.

CT angiography (CTA) and MR angiography (MRA) are specific investigations. Doppler ultrasound, duplex scanning, and digital subtraction percutaneous angiography (DSA) are the other available investigations.

An aortofemoral bypass is the choice of surgical intervention in aortoiliac obstruction. Dacron graft is a collagen-impregnated, knitted graft that is used to perform the proximal aortic anastomosis in aortofemoral bypass. It can also be used in aortic valve repair. In unfit patients, axillofemoral bypass may be done.

Other options:

Option A: Percutaneous Transluminal Angioplasty (PTA) is most useful in the treatment of isolated iliac stenoses (<4cm in length) than in occlusive disease.

Option B: Iliofemoral bypass is done in patients with unilateral occlusion of the distal common iliac or external iliac arteries.

Option D: Aortoiliac bypass more commonly known as the aortoiliac endarterectomy is rarely performed now a days because of greater blood loss associated with this surgery and greater sexual dysfunction caused, additionally it is difficult to perform.

Note: Aortofemoral bypass surgery has long-term patency of 70-80% at 10years and is often done bilaterally as the occlusion is more likely to be bilateral with one side affected more severely than the other. Bilateral aortofemoral bypass does not add to physiological stress and has equal benefits comparatively.

Solution to Question 16:

The image above shows dilated, tortuous veins of the lower limb. In asymptomatic patients, they are classified as C2A according to the Clinical-Etiology-Anatomy-Pathophysiology (CEAP) classification for chronic venous disorders.

Varicose veins are characterized by the tortuous dilatation of subcutaneous veins resulting from venous insufficiency. Patients have heaviness, burning, and aching over the affected area or whole limb, which increases throughout the day or on prolonged standing. Compression and elevation can relieve the symptoms.

Diagnosed by duplex ultrasound and is treated by modalities such as compression, laser ablation, radiofrequency ablation, endovenous glue, ultrasound-guided foam sclerotherapy, and open surgery.

Image A shows tortuous dilated veins (C2), Image B is atrophie blanche (C4b), and Image C shows fan-shaped telangiectasia (C4c).



The images below show C5 and C6 - healed and active venous ulcers respectively.

Solution to Question 17:

In an alcoholic patient with acute pancreatitis, serum lipase is the enzyme most likely to be elevated.

Acute pancreatitis is often caused by gallstones (70%) or alcoholism (25%). The cardinal symptom is acute-onset severe pain in the upper quadrant of the abdomen. The patient usually presents in shock (hypotension and tachycardia) and tachypnoea. Some may also present with signs of bleeding into the fascial planes leading to discoloration of the sites involved:

• Cullen's sign: Around the umbilicus

• Grey Turner's sign: Flanks

• Fox sign: Inguinal region

• Bryant's sign: Scrotum

The diagnosis is often clinical with a laboratory diagnosis. Though serum amylase (>3 times) is indicative of pancreatitis, serum lipase is preferred over serum amylase as it is more sensitive and specific. Contrast-enhanced tomography (CECT) of the abdomen can be used to exclude pancreatitis if in doubt.

The severity of the acute pancreatitis is assessed using the Ranson and Glasgow scoring system (≥3 at 48h of onset of pain). APACHE, SOFA, SAPS, and MODS can also be used to assess the severity of pancreatitis in intensive care (ICU) settings. Pancreatitis is managed conservatively with initial nasogastric drainage followed by intravenous fluids, analgesia, and frequent monitoring of vital signs, urine output, and blood gases.

Other options:

Option A: Serum gamma-glutamyl transferase (GGT) can also be elevated. It is a biomarker for pancreatic disease and alcohol consumption but is not sensitive and specific to acute pancreatitis.

Option B: Carcinoembryonic antigen (CEA), a tumor marker that is elevated in the adenocarcinoma of the pancreas.

Option D: Elevated serum bilirubin is a result of (i) overproduction of bilirubin (E.g: hemolytic anemia) or (ii) impaired uptake by the liver (liver cirrhosis) or (iii) regurgitation of bilirubin (obstructive jaundice). It is not seen in acute pancreatitis.

Solution to Question 18:

The given clinical scenario of acute abdominal pain and fever with chest X-ray in an erect position showing air under the diaphragm (pneumoperitoneum) is a characteristic of hollow viscus perforation.

The most common cause of pneumoperitoneum is bowel perforation (>90% of the cases). Gastric and posterior duodenal perforations are seen in long-standing cases of peptic ulcers. Intestinal perforations are seen with ischemia, inflammation, diverticula, or obstruction (E.g:- colonic carcinoma). Other causes also include iatrogenic, penetrating injuries, prolonged mechanical ventilation, peritoneal dialysis, etc.

Perforation of a viscus is a life-threatening condition due to leakage of the enteric contents which can lead to acute bacterial peritonitis.

Clinical features of hollow viscous perforation are due to the developed peritonitis. They include abdominal pain, nausea, and vomiting. On examination, the patient has tachycardia, tenderness, rebound tenderness, decreased or absent bowel sounds, and guarding on palpation. Fever may be present.

The patient is initially investigated with an upright chest X-ray which typically shows air beneath the diaphragm. Various other signs for pneumoperitoneum in the supine abdominal X-rays are Rigler's sign, Cupola sign, falciform ligament sign, football sign, and Doge's cap sign. However, CECT is the most sensitive investigation to detect minimal/ suspected pneumoperitoneum along with the cause and site of perforation.



Other options:

Option A: Gastric volvulus is a twisting or axial rotation of the stomach along its longitudinal axis (organo-axial) or vertical axis (mesentero-axial). The sudden onset of constant and severe upper abdominal pain, recurrent retching with the production of little vomitus, and the inability to pass an NG tube constitute the Borchardt triad. Plain films of the abdomen reveal a gas-filled viscus in the chest or upper abdomen.

Option C: Liver abscess will mostly have the classic presenting symptoms of fever, jaundice, and right upper quadrant pain, with a tenderness to palpation. Chest X-ray findings generally reflect subdiaphragmatic diseases, such as an elevated right hemidiaphragm, right pleural effusion, or atelectasis.

A chest radiograph shows an air-fluid level in the right upper abdomen (arrow) and multiple nodular opacities in the bilateral lungs (arrowheads)



Option D: Empyema thoracis is an infection of the pleural space. An empyema thoracis on a chest X-ray can mimic a pleural effusion or a lung abscess.The fluid is commonly drained completely with a chest tube or small-bore catheter.

The chest radiograph showing pleural empyema.



Surgery INI-CET 2022

Question 1:

Which of the following is not a perineal approach for the condition shown in the image?

a) Ripstein's procedure

b) Delorme's operation

c) Thiersch operation

d) Altemeier's procedure

Question 2:

Identify the true statement regarding split skin and full-thickness skin grafts:

a) 2,3

b) 2,4

c) 1,3

d) 1,4

Question 3:

A 37-year-old unmarried nulliparous woman, having regular intercourse is on oral contraceptive pills. Her mother was diagnosed with carcinoma breast at 60 years of age and her elder sister was diagnosed with carcinoma ovary at 40 years of age. What is the next line of management?

a) Genetic counselling and screening for BRCA

b) Routine mammography

c) Stop taking oral contraceptive pills

d) Prophylactic surgery

Question 4:

Which of the following causes maximum malnutrition?

a) Pancreatic fistula

b) Duodenal fistula

c) Distal ileal fistula

d) Colonic fistula

Question 5:

A 50-year-old alcoholic patient presents with a history of waxing and waning jaundice for the past 2 months. His CT examination reveals, dilatation of the common bile duct and pancreatic duct. What is the most probable diagnosis in this patient?

a) Cholangiocarcinoma

b) Periampullary carcinoma

c) Common bile duct stone

d) Head of pancreas carcinoma

Question 6:

Which investigation is not necessary for the following condition?



a) Timed barium study

b) Endoscopy

c) Esophageal manometry

d) 24-h pH monitoring

Question 7:

The following procedure is performed for the management of:

a) Advanced gastric carcinoma

b) Gallbladder carcinoma

c) Distal cholangiocarcinoma

d) Chronic calcific pancreatitis

Question 8:

A patient developed a loss of sensations over the root of the penis following open hernia surgery. Which of the following nerves would you implicate?

a) Iliohypogastric nerve

b) Ilioinguinal nerve

c) Genitofemoral nerve

d) Medial cutaneous nerve of thigh

Question 9:

Which of the following statements about Psoas abscess are true?

a) 1 and 2 only

b) 1, 3, and 4

c) 1, 2, 3, and 4

d) 2, 3, and 4

Question 10:

A patient who had grade 3 splenic trauma was managed conservatively in the ICU. Which of the following signs on day 2 would indicate a need for surgery?

a) Haemoglobin fall from 12g/dL to 10g/dL

b) Extraperitoneal bladder rupture

c) Pneumoperitoneum

d) Air in gallbladder

Question 11:

A 34-year-old woman was brought to the AIIMS emergency following a road traffic accident. She was hemodynamically stable and her GCS score was 3. She was intubated and sent for non-contrast CT which was found to be normal. What is the most likely diagnosis?

a) Subdural hemorrhage

b) Extradural hemorrhage

c) Diffuse axonal injury

d) Cerebral contusion

Question 12:

Which of the following findings is likely to be seen in this patient?

a) Decreased posterior tibial & dorsalis pedis pulsations

b) Loss of sensation over foot

c) Positive trendelenberg sign

d) Pain on compression of calf

Question 13:

A 30-year-old male patient presents with a pelvic fracture following blunt abdominal trauma. No blood is seen at the meatus. Catheterization does not produce significant urine output. The radiological image is given below. What is the next step in management?

a) Suprapubic cystostomy

b) Perurethral catheterization

c) Laparotomy

d) Pelvic packing along with B/L percutaneous nephrostomy

Question 14:

Which of the following would be the most commonly encountered ectopic tissue in Meckel's diverticulum?

a) Gastric and pancreatic

b) Colonic and pancreatic

c) Gastric and colonic

d) Thyroid and lung

Question 15:

What is true about gall bladder carcinoma?

a) Palpable abdominal lump is the earliest sign

b) Earliest sign is jaundice

c) Clinical presentation of gall stones can be distinguished from gall bladder cancer

d) Median survival is 6 months

Question 16:

Neurogenic shock is associated with _____.

a) Hypertension + tachycardia

b) Hypotension + tachycardia

c) Hypotension + bradycardia

d) Hypertension + bradycardia

Question 17:

An old lady sustained a fall and developed a headache and neurological symptoms a few days later. What is your diagnosis?

a) Multi-infarct dementia

b) Frontal lobe meningioma

c) Korsakoff disease

d) Subdural hematoma

Question 18:

Which of the following is not a part of damage control surgery?

a) Control of hemorrhage

b) Resuscitation

c) Decontamination

d) Vascular anastomosis

Question 19:

All the following are criteria for SIRS, except _____.

a) Systolic BP <90 mmHg

b) Temperature >38 degree celsius or <36 degree celsius

c) Respiratory rate >20 bpm

d) Heart rate >90/ min

Question 20:

Which is the correct sequence for the correction of hypospadias?

a) 1->3->2->4

b) 3->1->2->4

c) 3->1->4->2

d) 1->3->4->2

Question 21:

A female patient presents with a lump in her breast. All the following can be used to rule out malignancy, except _____.

a) Radiological investigations

b) PET scan

c) Clinical examination

d) Histology and cytology

Question 22:

An elderly woman underwent surgical resection of her bowel. After a few months, she presents with easy fatiguability and tiredness. Blood smear evaluation shows macrocytic anemia. Resection of all the following can cause this, except _____.

a) Jejunum

b) Duodenum

c) Ileum

d) Stomach

Question 23:

Match the following:

a) A-1, B-2, C-3, D-4

b) A-4, B-1, C-2, D-3

c) A-3, B-2, C-1, D-4

d) A-4, B-3, C-2, D-1

Question 24:

A female patient was diagnosed with gastric cancer. On immunohistopathology, the biopsy sample was found to be positive for CD117 and negative for LCA, cytokeratin, and S100. What is the diagnosis?

a) GIST

b) Adenocarcinoma

c) Hodgkins lymphoma

d) SCC

Question 25:

Which of these is the most life-threatening injury that can be identified by assessing the breathing component of the patient?

a) Cervical spine injury

b) Tension pneumothorax

c) Blunt cardiac injury

d) Laryngotracheal injury

Question 26:

A 32-year-old female patient with Graves' disease with eye signs and enlarged thyroid is planned for a total thyroidectomy. What can be given in the preoperative period to reduce intraoperative bleeding in the patient?

a) Potassium iodide

b) Propranolol

c) Propylthiouracil

d) Betamethasone

Question 27:

In all the following places, hand rub can be used according to standard hand hygiene protocol, except _____.

a) If the hands are visibly soiled

b) While moving from a contaminated site to a clean site during patient care

c) Before donning on gloves

d) During direct patient contact

Question 28:

Which of the following is true regarding all types of shock?

a) Anaerobic respiration produces carbon dioxide

b) Low pH mainly activates lysosomal enzymes

c) Peripheral vasoconstriction

d) Renal perfusion is maintained

Question 29:

A patient presents with multiple stab injuries to the chest. On examination, there is reduced chest movement on the right side and tracheal deviation to the left side. Additionally, the neck veins are distended. What is the next best step in the management of this patient?

a) Resuscitative thoracotomy

b) ICD placement

c) Needle thoracocentesis

d) Finger thoracotomy

Question 30:

Match the following:

a) A-4, B-1, C-4, D-2

b) A-3, B-1, C-4, D-2

c) A-3, B-1, C-2, D-4

d) A-1, B-4, C-3, D-4

Question 31:

What is the correct sequence of management in a patient who presents to the casualty with an RTA?

a) 1, 3, 2, 4

b) 1, 2, 3, 4

c) 2, 1, 3, 4

d) 2, 1, 4, 3

Question 32:

A 32-year-old male patient presents to the surgical emergency with a history of peptic ulcer disease and pain abdomen, bloating, and obstipation for 2 days. What is the next line in management?



a) A and B

b) A and C

c) A and D

d) A, B, and C

Question 33:

A female patient is planned for cholecystectomy. Which of the following is not an advantage of laparoscopic surgery?

a) Lesser post-operative pain

b) Cosmetically better

c) Earlier return to normal physiological function

d) Longer duration of stay in hospital

Question 34:

What does the magnified number marked on the Foley catheter correspond to?



a) Inner lumen diameter

b) Length of the tube

c) Outer tube diameter

d) Bulb size

Question 35:

A 49-year-old female presents with a breast lump. Which of the following findings is in accordance with basal-like breast cancer?

a) ER +, PR +, HER2 -

b) ER +, PR -, HER2 -

c) ER -, PR -, HER2 -

d) ER -, PR -, HER2 +

Question 36:

A patient presented with large intra-abdominal fluid accumulation and was found to have pancreatic necrosis. Which of the following are included in the management of the condition?

a) 1, 2, 3, 4, and 5

b) 2, 4, 5 and 6

c) 1, 2, 3, and 5

d) 1, 2, 3, 4, 5 and 6

Question 37:

A patient presents with a chest x-ray finding of pneumothorax. Which of the following is not a boundary of the triangle of safety for ICD insertion?

a) Lateral edge of pectoralis major

b) Mid-clavicular line

c) Base of axilla

d) Lateral border of latissimus dorsi

Question 38:

Which of the following cancers are correctly matched with the criteria for the minimum number of lymph nodes required for pathological staging?

a) B, C

b) A, B, C

c) A, C, D

d) A, B, C, D

Question 39:

Severe intractable pain in the leg due to sciatica in a patient with carcinoma rectum is due to:

a) Anterior infiltration in females

b) Anterior infiltration in males

c) Lateral infiltration

d) Posterior infiltration

Question 40:

Identify the correct statement regarding gastric carcinoma.

a) 1,3,4

b) 2, 3

c) 2,3,4

d) 1,4

Question 41:

All the following are true about acute cholecystitis, except _____.

a) Gall bladder thickness >3 mm on USG

b) Murphy's sign positive

c) Leukocytosis

d) Preferential visualization of gall bladder in HIDA scan

Question 42:

Lymphatic drainage from the breast is mostly drained into the axillary lymph nodes. Which of the following is considered the location of the apical lymph nodes?

a) In the axillary pad of fat at the apex of the axilla

b) Along the margin of the pectoralis minor

c) Along the subscapular veins

d) Anterior in the center

Question 43:

A 10-year-old child with hereditary spherocytosis and hypersplenism presents to the hospital. A splenectomy is planned for this child. All the following vaccines are recommended before splenectomy, except _____.

a) Pneumococcal vaccine

b) Typhoid vaccine

c) Haemophilus influenzae B vaccine

d) Meningococcal vaccine

Question 44:

In proximal transverse colon carcinoma, which of the following arteries are resected to prevent the spread of cancer?

a) Ileocolic, right colic and branch of middle colic

b) Ileocolic, left colic, and stem of middle colic

c) Ileocolic, right colic, and trunk of middle colic

d) Middle colic only

Question 45:

Which of the following statements are true regarding central venous access and peripheral venous access?

a) 1,3,4

b) 2,3,4

c) 1,2,3,4

d) 1,2,3

Question 46:

A patient who had undergone appendicectomy 2 days ago presents with severe abdominal pain. On exploration, there is a collection of blood in the abdomen. Which of the following arteries should have been ligated during surgery?

a) Ileocolic artery

b) Right colic artery

c) Superior rectal artery

d) Superior mesenteric artery

Question 47:

A 25-year-old woman presents with a breast lump with no skin changes or induration. On histology, there was a fibroadenoma appearance. Which of the following categories does it fall

under?

a) BIRADS 1 and 2

b) BIRADS 2 or 4

c) BIRADS 2, 3, 4

d) BIRADS 2

Question 48:

Which of the following is not a contraindication for breast conservative surgery?

a) Tumor size < 4cm, with N1 axillary metastasis

b) Tumor more than the size of the breast

c) Prior chest irradiation

d) Multifocal tumors

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	b
5	b
6	d
7	c
8	b
9	c
10	c
11	c
12	b
13	c
14	a
15	d
16	c

17	d
18	d
19	a
20	c
21	b
22	b
23	b
24	a
25	b
26	a
27	a
28	b
29	c
30	c
31	b
32	b
33	d
34	c
35	c
36	c
37	b
38	a
39	a
40	b
41	d
42	b
43	b
44	c
45	c
46	a
47	d
48	a

Detailed Explanations

Solution to Question 1:

The condition shown in the image is a rectal prolapse and Ripstein's procedure is done using the abdominal approach.

Rectal prolapse occurs when a mucosal or full-thickness layer of rectal tissue protrudes through the anal orifice. Rectal prolapse is of two types:

• Partial rectal prolapse: Only the mucosal layer of the rectum is prolapsed. It is seen in children.

• Complete rectal prolapse: All the layers of the rectum are prolapsed. It is seen in adults due to the weakening of the pelvic floor.

Management:

• The first episode of a partial rectal prolapse is managed by digital repositioning. On recurrence, Theirsch's wiring or sclerotherapy are considered.

• Complete rectal prolapse is managed surgically, either by perineal or abdominal approaches:

1. Perineal approach: Often preferred in patients with high operative risk and limited life expectancy, i.e. in the elderly. It is associated with less morbidity but has a higher recurrence rate.

• Thiersch's wiring- A steel wire/nylon tape is placed around the anal canal. Largely obsolete now due to problems with chronic perineal sepsis, anal stenosis, and obstructed defecation.

• Delorme's mucosectomy- The rectal mucosa is stripped circumferentially from the rectum over the length of the prolapse. The underlying muscle is plicated with a series of sutures. The excess rectal mucosa is excised and a mucosal anastomosis is performed.

• Altemeier's rectosigmoidectomy- The rectum is prolapsed through the anal canal and a full-thickness resection is performed, incorporating any associated colonic prolapse. Restoration of colorectal continuity can be performed by hand-sewn/stapled anastomosis. This is the procedure of choice in patients presenting with incarcerated and strangulated prolapse.

2. Abdominal approach: Difficult to perform, more complications but a lower recurrence rate.

• Ripstein's procedure- involves the placement of a prosthetic mesh around the mobilized rectum, with the attachment of the mesh to the presacral fascia (anterior) below the sacral promontory.

• Well's procedure- involves posterior rectopexy.

• Suture rectopexy

Solution to Question 2:

Statements 2 and 3 are correct.

Split skin grafts (Thiersch grafts) are partial-thickness skin grafts consisting of the epidermis and a part of the dermis. Whereas, full-thickness grafts (Wolfe grafts) consist of the epidermis along with the entire dermis, with portions of the sweat glands, sebaceous glands, and hair follicles.

Primary contracture of the skin graft occurs due to passive recoil of the elastin fibers of the dermis. It occurs soon after the graft has been harvested.

• Since full-thickness grafts have more dermis (implying more elastin fibers), they undergo more primary contracture.

• Whereas, split skin grafts have less dermis, therefore, undergo less primary contracture.

Secondary contracture of the skin graft occurs due to the pull of myofibroblasts. It occurs over time, when the transplanted graft starts healing.

• Full-thickness grafts have more ability to resist the pull of myofibroblasts, hence they have less secondary contracture.

• Whereas, split skin grafts yield to the pull and develop more secondary contractures.

Advantages of split over full-thickness skin graft:

• Large grafts can be harvested.

• In the case of thinner split-thickness grafts, the donor site heals more rapidly and can be reused in severe cases such as burns.

• Uptake is good. Meshing can be done, i.e. small splits are done in graft, allowing exudates to drain out, thus improving graft take up.

• Meshing also allows the graft to expand.

Advantages of full-thickness over split skin graft:

• It has a lesser degree of post-graft contraction

• Best cosmetic appearance as they include the highest number of skin appendage elements.

Primary contracture	Secondary contracture
Elastic recoil	Pull of myofibroblasts
Less in split-skin grafts	More in split-skin grafts
More in full-thickness grafts	Less in full-thickness grafts

Solution to Question 3:

The given family history of two primary relatives with breast and ovarian cancer with one of the relatives with onset <50 years is suggestive of a woman with a high risk of breast cancer. The next line of management is genetic counselling and screening for BRCA.

Genetic counselling is the process of discussing various genetic tests such as BRCA testing and multiple gene panel testing.

Genetic counselling and testing for genetic mutations (BRCA 1, BRCA 2 and others such as TP53, PTEN) are recommended for high-risk women with:

• Breast cancer diagnosed before age 50

• Triple-negative breast cancer

• Bilateral breast cancer

• Breast and ovarian cancer in the same individual

• Breast cancer in men

• Family history (maternal or paternal) of two or more individuals with breast and ovarian cancer

• A close male relative with breast cancer

• A close relative with early-onset (<50 years) breast or ovarian cancer

• Known BRCA1 or BRCA2 mutation in the family

Other options:

Option B: Routine mammography is not the initial investigation of choice, but is used for surveillance after the genetic screening. Surveillance strategies for women with a family history of breast cancer or ovarian cancer include

• Self-examination of breasts starting at age 18 to 20 years - Each month

• Clinical breast examination starting at age 25 years - Every 6 months

• Mammography starting at the age of 25 years or 10 years earlier to breast cancer onset in a family member - Every year

Option C: Oral contraceptives can be continued in this patient as they don't increase the risk of breast cancer in a woman with a positive family history of breast cancer. It is also useful in reducing the risk of ovarian and endometrial cancers, which are associated with BRCA mutations.

Option D: Prophylactic surgery is advised after the identification of BRCA positivity in high-risk women. It is proven to be more effective than regular surveillance by mammography and MRI. It is usually combined with a prophylactic risk-reducing salpingo-oophorectomy that reduces the risk of ovarian and breast cancers.

Solution to Question 4:

Among the given conditions, maximum malnutrition is caused by duodenal fistulas.

Enterocutaneous fistulas are classified according to their location and volume of daily output.

• Proximal fistulas are associated with higher output, greater fluid and electrolyte loss, and greater loss of digestive capacity.

• Distal fistulas tend to have lower output, making them easier to manage and more likely to close spontaneously.

Although both gastric and duodenal fistulas are proximal fistulas, duodenal fistulas receive an additional 1 to 2 liters of enzymes, electrolytes, and water from the pancreatic and bile ducts. Hence duodenal fistulas are at greater risk for more electrolyte imbalance than gastric fistulas.

The management of enterocutaneous fistula is based on well-established principles known as SNAP which include:

• S - Sepsis elimination and skin protection

• N - A period of parenteral nutrition may well be required

• A - Anatomical assessment

• P - Definitive planned surgery

Solution to Question 5:

In the given scenario, a characteristic history of waxing and waning jaundice along with a dilated common bile duct and pancreatic duct on CT scan, points to a diagnosis of periampullary carcinoma.

Periampullary carcinoma arises from the vicinity of the ampulla of Vater. Patients with hereditary polyposis syndromes, such as familial adenomatous polyposis (FAP) and hereditary nonpolyposis colorectal cancer (HNPCC) have a higher risk of developing it.

Clinical features:

• Jaundice: It intermittently waxes and wanes because, whenever the tumor undergoes necrosis and sloughing, obstruction to bile ducts is partially relieved.

• Weight loss

• Pain: It depends on the location of the tumor and may involve back pain as well due to the involvement of retropancreatic nerves, pancreatic duct obstruction or stasis, and disruption of nerve sheath by the tumor.

• Nausea/vomiting

• Pruritus

• Fever

• Gastrointestinal bleeding

In suspected periampullary disease, a three-phase CT scan should be done routinely.

Management includes resection, along with chemotherapy/ chemoradiotherapy.

Solution to Question 6:

The given barium swallow shows a bird beak appearance and is suggestive of achalasia cardia. 24-h pH monitoring is not necessary for achalasia cardia. It is used in the diagnosis of gastroesophageal reflux disease (GERD).

Achalasia is a primary esophageal motility disorder. It is characterized by the absence of esophageal peristalsis and impaired lower esophageal sphincter (LES) relaxation in response to swallowing. This is due to the loss of ganglion cells in the myenteric (Auerbach's) plexus. These abnormalities cause a functional obstruction at the gastroesophageal junction.

Option A: On performing a timed barium study, barium will be retained for more than five minutes. The fluoroscopic examination will reveal a dilated esophagus that tapers to form a 'bird's beak' at the gastroesophageal junction as seen in the image above.

Option B: Endoscopy is an important investigation carried out to rule out cancer of the gastric cardia.

Option C: Esophageal manometry is the investigation of choice and shows:

• Incomplete LES relaxation in response to swallowing

• High resting LES pressure

• Absent esophageal peristalsis

Solution to Question 7:

The given image shows the Whipple procedure.

The Whipple procedure is a complex surgery done for carcinomas involving the head of the pancreas and distal cholangiocarcinoma. It is done only in cases when:

• The entire duodenum has to be removed e.g. in familial adenomatous polyposis (FAP)

• Tumor encroaches in the first part of the duodenum or the stomach

• A clear resection margin is not achieved by pylorus-preserving pancreatoduodenectomy

The following structures are removed during Whipple's procedure:

• Distal stomach

• Duodenum

• Head of the pancreas

• Gallbladder

• Common bile duct (CBD)

• Regional lymphatics



Solution to Question 8:

Loss of sensation in the root of the penis and adjacent part of the scrotum is suggestive of an injury to the ilioinguinal nerve. It is most commonly injured within the inguinal canal in open hernia repair due to traction or thermal injury.

Other options:

Option A: The iliohypogastric nerve carries the sensory supply from the suprapubic area. Injury to this nerve in the right iliac fossa rarely causes sensory disturbances, as the suprapubic area has a rich nerve supply from other sources too. If the nerve is severed above the anterior superior iliac spine, it can lead to the weakened posterior wall of the inguinal canal resulting in a direct inguinal hernia. The iliohypogastric nerve can also get injured following appendicectomy.

Option C: Injury to the genital branch of the genitofemoral nerve results in loss of sensation in the skin of the scrotum of males, and loss of sensation of the mons pubis and labia majora in females.

Injury to the femoral branch of the genitofemoral nerve causes hypoesthesia over the anterior thigh below the inguinal ligament.

Option D: The medial cutaneous nerve of thigh is a sensory branch of the femoral nerve. Injury to this nerve leads to loss of sensations over the areas supplied by this nerve, which include anterior-medial aspect of the thigh and knee.



Solution to Question 9:

The correct statements regarding psoas abscess are 1, 2, 3, and 4.

Psoas abscess occurs due to a collection of pus due to an infection in the iliopsoas compartment. The compartment is retroperitoneal and hence, the abscess is also called a retroperitoneal abscess. The compartment contains iliacus and iliopsoas muscles.

Psoas abscess is either primary or secondary.

• Tuberculosis of the spine leading to pott's abscess was the leading cause of psoas abscess (statement 2).

• Recently, hematogenous spread of infection from an occult primary site (gastrointestinal or urinary tract) has become common, especially in immunocompromised patients and the elderly (statement 4).

• Due to contiguous extension of infection from neighbouring structures.

Features include

• Back pain and fever.

• Soft, fluctuant swelling pointing towards the groin (hip) (statement 1).

• Pain on passive hip extension or active flexion of the psoas (statement 3).

CT scanning is useful in diagnosis. Percutaneous CT-guided drainage and antibiotic therapy are useful in the treatment of the condition.

Radiological images showing psoas abscess (mass lateral to the vertebral column):



Solution to Question 10:

The presence of pneumoperitoneum on imaging would warrant a laparotomy as it indicates visceral perforation which can lead to secondary peritonitis.

Secondary peritonitis usually requires surgery along with antibiotics.

The spleen is one of the most commonly injured abdominal organs. The severity of the injury is graded by the Americian association for the surgery of trauma (AAST) spleen injury scale. The management is based on the grade of the injury. Abdominal CT with intravenous contrast is the investigation of choice.

Conservative management is recommended in most cases according to current guidelines. There must be no evidence of active blood loss. Splenic angiography and embolization are also used for non-surgical management.

Surgical management is recommended when there is hemodynamic instability or failure to respond to conservative management. It can be in the form of splenic repair, partial splenectomy, or total splenectomy. Splenectomy is recommended in significant injuries to the hilum, pulverized splenic parenchyma, or any injury >grade 2 with coagulopathy.

Vaccination against encapsulated bacteria must be provided to prevent infection if a splenectomy is done.

Other options:

Option A - The initial hemoglobin level should be carefully assessed to monitor blood loss. Patients who are stabilized after crystalloid infusion can also be managed conservatively but a lower threshold for surgery must be kept. But hemoglobin fall from 12g/dL to 10g/dL is not an indication for laparotomy as the fall in hemoglobin is not marked.

Option B - Extraperitoneal bladder rupture can be managed by decompression with a urinary catheter. Later a cystography can be done to confirm healing.

Option D - Air in the gallbladder can be seen in acute cholecystitis if gas-forming organisms cause secondary bacterial infection (emphysematous cholecystitis). It is managed with antibiotics and intravenous fluids. Laparoscopic cholecystectomy is the surgery of choice.

Solution to Question 11:

The clinical scenario of a patient with a poor GCS score and a normal NCCT brain scan following a road traffic accident is suggestive of diffuse axonal injury.

Diffuse axonal injury (white matter shearing) results from shearing forces, typically from a rotational acceleration of the brain within the rigid cranium. This causes damage to microvasculature and disruption of axons in the subcortical parasagittal white matter, cerebellum, fornix, and internal and external capsule.

Most commonly it occurs in automobile accidents. Patients are usually unconscious from the moment of impact and remain the same until death.

Diffuse axonal injury is characterized by multiple focal lesions with a characteristic distribution:

• CT may be normal.

• MRI is the modality of choice for assessing suspected diffuse axonal injury, even in patients with entirely normal CT of the brain. MRI, especially SWI (susceptibility-weighted imaging) demonstrates small regions of susceptibility artifact or hemorrhage.

Other options:

Option A: Subdural hemorrhage is often due to the rupture of veins that cross the subdural space. In subdural hemorrhage, the CT shows concavo-convex opacity.

Option B: Extradural hemorrhage is mostly due to the rupture of the middle meningeal artery. Since the dura matter tends to adhere to the skull the hemorrhage is seen in CT as a biconvex dense area. They usually do not cross cranial sutures.

Option D: Cerebral contusion occurs due to traumatic brain injury, at the sites that come in direct contact with the skull. Imaging may show microhemorrhages and edema. Usually, no treatment is required.

Solution to Question 12:

In the given image, the lesion present over the plantar surface of the foot is likely to be a neuropathic ulcer. Neuropathic ulcers can lead to loss of sensation over the foot.

A neuropathic ulcer is one that occurs as a result of peripheral neuropathy. In peripheral neuropathy, lack of sensation over the pressure points leads to cumulative microtrauma, minor wounds, or cuts that go unnoticed eventually developing into ulcers. They are painless ulcers that are punched out with regular margins. They are often located on the plantar surface of the foot. Ulcer margins show the presence of calluses.

Other options:

Option A: Decreased posterior tibial and dorsalis pedis pulsations are usually seen in peripheral arterial disease, which causes arterial ulcers that are usually found on the dorsal surface of the foot.

Option C: Positive Brodie-Trendelenburg test is seen in varicose veins. This condition leads to venous ulcers which are usually seen on the medial malleolus of the foot.

Option D: Pain on compression of calf is a sign of deep venous thrombosis.

Solution to Question 13:

The given clinical scenario of blunt abdominal trauma, minimal urine output on catheterization, and the pattern of contrast extravasation in the given image indicate intraperitoneal bladder rupture. The absence of blood at the meatus rules out urethral injury. The preferred next step of management, in this case, is laparotomy.

Bladder rupture may be extraperitoneal or intraperitoneal.

• The bladder is related to the peritoneum only on its superior surface. Therefore, if the bladder rupture is towards the superior surface, the urine leaks into the peritoneal cavity, i.e. urine extravasation will be intraperitoneal.

• Whereas, bladder rupture due to injury to the anterior or lateral surfaces does not rupture the peritoneum. In such cases, the urine leaks into the perivesical space but does not enter the peritoneum, i.e. urine extravasation will be extraperitoneal.

Intraperitoneal bladder rupture is caused by a blow or fall on the distended bladder. Intraperitoneal bladder rupture manifests as sudden severe pain in the hypogastrium and syncope. There will be an associated increase in BUN and creatinine due to systemic absorption of the extravasated urine and it may also lead to chemical peritonitis.



Investigations :

• CT scan can be done to diagnose bladder rupture.

• A plain erect X-ray may show a ground-glass appearance.

• Intravenous urography (IVU) may confirm a leak.

• Retrograde cystography will confirm the diagnosis.

Intraperitoneal bladder rupture leads to extravasation of the contrast material into the peritoneal cavity, around the bowel loops and paracolic gutters. Hence, an outline of bowel loops can be seen. The image below shows intraperitoneal bladder rupture, with arrows showing contrast material in the peritoneal cavity:



Extraperitoneal bladder rupture leads to extravasation of the contrast material into the perivesical space and the fascia planes anterior to the peritoneum. The image below shows extraperitoneal bladder rupture, with contrast material in the retro-pubic space (no contrast can be seen within the peritoneal cavity).





Solution to Question 14:

The ectopic tissue encountered in Meckel's diverticulum includes gastric and pancreatic tissue.

Meckel's diverticulum is caused by a failure of normal regression of the vitelline duct that develops during weeks 5 to 7 of gestation.

The most common presentation is painless, massive, lower GI bleeds in children younger than 5 years. A Meckel's can be the lead point for ileoileal or ileocolic intussusception. The bleeding Meckel's diverticulum may be confirmed by a 99 mTc-pertechnetate isotope scan to detect gastric mucosa and pancreatic tissue.

Surgical treatment is the definitive therapy for Meckel's diverticulum. A V-shaped diverticulectomy with transverse closure of the ileum can be done.

Solution to Question 15:

Gallbladder carcinoma is a rare malignancy that is seen in the elderly. It is associated with a poor prognosis and median survival is less than 6 months.

Gallbladder carcinoma is indistinguishable from cholecystitis or cholelithiasis (option C). It is usually diagnosed when patients undergo cholecystectomy for gallstones. Its clinical features include right upper quadrant pain, nausea, vomiting, and anorexia. Jaundice, palpable mass, and anorexia are late features (options A and B). CA 19-9 is elevated in the carcinoma gallbladder and is used for monitoring.

The preoperative diagnosis is often made on ultrasonography and followed up by a CT scan or MRI/ Magnetic resonance cholangiopancreatography.

A percutaneous biopsy under radiological guidance is done for pathological examination. PET scanning helps in detecting metastatic disease.

In select cases, a laparoscopic examination is useful in staging the disease. Laparoscopy can also detect peritoneal or liver metastases. The most important prognostic factor is the depth of invasion (T-stage).

Solution to Question 16:

Neurogenic shock is associated with Hypotension + Bradycardia.

Neurogenic shock refers to reduced tissue perfusion due to loss of vasomotor tone to peripheral arterial beds. It is a type of distributive shock.

Neurogenic shock is usually secondary to spinal cord injuries due to vertebral body fractures of the cervical or high thoracic region that disrupts the sympathetic regulation of peripheral vascular tone. Loss of vasoconstrictor impulses results in increased vascular capacitance decreased venous return and decreased cardiac output.



The classic description of neurogenic shock consists of:

• Hypotension associated with bradycardia (absence of reflexive tachycardia due to disrupted sympathetic discharge)

• Warm extremities (due to loss of peripheral vasoconstriction)

• Motor and sensory deficits indicative of a spinal cord injury

• Radiographic evidence of a vertebral column fracture

Solution to Question 17:

The given clinical scenario of an elderly suffering from headache and neurological symptoms following trauma is suggestive of subdural hematoma (SDH).

SDH (acute/chronic) is the accumulation of blood between the arachnoid membrane and the dural layers of the meninges. The source of the bleeding is usually from the rupture of bridging veins running from the cerebral cortex to the dural sinuses. Elderly, alcoholics & patients on anticoagulation are at a higher risk of developing SDH, especially after head trauma including trivial incidents.

Non-contrast CT is the investigation of choice.

Radiographic features of subdural hemorrhage are:

• Typically unilateral (85%).

• Crescent-like distribution (concavo-convex) around the periphery of the brain.

• Not limited by sutures (vs extradural hemorrhage which is limited by sutures)

• Hyperdense lesion on CT: Acute SDH

• Hypodense lesion on CT: Chronic SDH

The given image shows the hyperdense collection seen in acute SDH.

Open craniotomy for the evacuation of acute SDH is indicated for the following:

• Thickness >1 cm

• Midline shift >5 mm

• Or GCS drop by two or more points from the time of injury to hospitalization.

Other options:

Option A: Multi-infarct dementia (MID) is a type of vascular dementia occuring due to multiple small infarcts leading to brain tissue injury. It commonly causes memory loss in the elderly.

Option B: Frontal lobe meningiomas are benign brain tumors that develop from the meninges. They cause personality changes that may be mistaken for depression.

Option C: Korsakoff syndrome is a memory disorder that results from vitamin B1 deficiency and is commonly seen in alcoholics. These patients present with impairment in storing new memories, disorientation, and confabulation. Consciousness is preserved.

Solution to Question 18:

Vascular anastomosis is undertaken as a part of definitive surgery, not damage control surgery (DCS).

DCS is minimal intervention done in order to stabilize the patient and followed by definite surgery later. It aims to coordinate staged operative interventions with periods of aggressive resuscitation to salvage trauma patients sustaining major injuries.

The initial focus is hemorrhage control, resuscitation, followed by control and limitation of contamination. It includes an abbreviated laparotomy, temporary packing, and closure of the abdomen in an e■ort to blunt the physiologic response to prolonged shock and massive hemorrhage.

After the initial steps of DCS, definitive surgery involving vascular anastomosis may be done. This is individualized to the patient and preferred to be completed in the first 24–72 hours.

Note: Vascular anastomosis is the option least likely to be part of the initial stages of DCS. Hence, it is the best option out of the four for this testing point.

Solution to Question 19:

Systolic BP <90 mmHg is not a criterion for SIRS (systemic inflammatory response syndrome).

SIRS is diagnosed when two of the following are present:

• Hyperthermia (>38 degrees celsius) or hypothermia (<36 degrees celsius)

• Tachypnoea (>20/min) or arterial carbon dioxide tension (PaCO2) < 32 mmHg

• Tachycardia (>90/min)

• White cell count >12 x 109/litre or <4 x 109/litre or >10% Band forms

Sepsis is SIRS with a documented infection.

Solution to Question 20:

The correct sequence for the correction of hypospadias is option C: 3->1->4->2

The main principles of surgery in hypospadias, in sequence, include:

• Orthoplasty - Chordee correction

• Urethroplasty - To reposition the urethral opening and tubularize urethra

• Glanuloplasty - Reconstruction of glans

• Meatoplasty - Reconstruction of distal urethral meatus

• Skin cover and scrotoplasty

Hypospadias is the most common congenital abnormality of the urethra. It is characterized by the combination of a ventrally (underside of penis or perineum) placed urethral meatus, a hooded foreskin, and ventral chordee (bending of the penis).

The image given below shows the classification of hypospadias:



Perineal hypospadias is the most severe type of deformity.

Operations for hypospadias are performed for sexual, urinary stream, or cosmetic reasons. It is done between 6 and 18 months of age.

Circumcision should not be done in hypospadias as the foreskin will be used for surgical reconstruction.

Tubularized incised plate (TIP) urethroplasty is the most commonly done procedure. It is done for mid-distal hypospadias.

Solution to Question 21:

In a patient presenting with breast lump or other suspicions of breast carcinoma a triple assessment method is used to rule out malignancy. PET scan is used to assess the spread and sites of distant metastasis, it is not used to rule out malignancy. It is also not a part of the triple assessment.

The triple assessment in the examination of breast lump includes:

• Clinical assessment - Clinical history and physical examination of the lump.

• Radiological examination - USG/ mammography/ both.

• Cytological or histological examination - FNAC (Fine needle aspiration cytology)/ core biopsy of the lump.

Features of malignancy seen on clinical examination can be enlarged axillary lymph nodes, circumferential retraction of the nipple, peau'd orange appearance of the skin and ulcerations over

the breast. Benign features include non-progressive, smooth, circumscribed, soft, mobile lumps without skin changes.

Microcalcifications seen on breast radiography are also highly suggestive of breast carcinoma.

FNAC and biopsy can detect malignant and atypical cells, suggestive of malignancy.

The triple assessment method has a positive predictive value of about 99.99%. In other words, if all three modalities give negative results, breast cancer can be excluded.

Solution to Question 22:

The given clinical vignette indicates a diagnosis of macrocytic anemia due to either vitamin B12 (cyanocobalamin) and/or vitamin B9 (folate) deficiency. The resection of the duodenum is least likely to cause this condition.

The duodenum is involved in iron absorption and hence its resection causes iron deficiency anemia which shows a microcytic hypochromic picture on peripheral blood smear.

The stomach and ileum are involved in the absorption of vitamin B12, and the jejunum is mainly used in folic acid absorption. Hence, a resection of these segments will result in a deficiency of the vitamins which in turn causes megaloblastic anemia with a peripheral smear showing macro-ovalocytes and a hypersegmented neutrophil.

Vitamin B12 (cobalamin), is a water-soluble vitamin derived from animal products such as red meat, dairy, and eggs. Absorption of vitamin B12 begins when pepsin in the stomach releases vitamin B12 from the binding proteins in the food. The released B12 binds to haptocorrin (R-factor), which is a salivary protein. In the duodenum, pancreatic proteases release B12 from haptocorrin, which then binds to intrinsic factors to form the IF- B12 complex. Intrinsic factor (IF) is a glycoprotein secreted by the parietal cells of the stomach. In the terminal ileum, this complex is absorbed by the enterocytes.

Pernicious anemia is a specific form of megaloblastic anemia caused by autoimmune gastritis. It is due to impaired production of the intrinsic factor from the parietal cells.It is a common cause of vitamin B12 deficiency. The other causes of vitamin B12 deficiency are gastrectomy, gastric bypass surgery, inflammatory bowel disease, pancreatic insufficiency and so on.



Vitamin B9 (folate) is a water-soluble vitamin, and its active form is tetrahydrofolate (THFA). They are mainly derived from liver, yeast, spinach, other greens, and nuts. Its deficiency results in megaloblastic anemia, homocysteinemia, and neural tube defects (spina bifida, Anencephaly) in the fetus.

Absorption of vitamin B9 is mainly from the upper small intestine, specifically the proximal jejunum.

Solution to Question 23:

The correct answer is option B: A-4, B-1, C-2, D-3

Marjolin ulcer is a rare and aggressive form of squamous cell carcinoma that grows from long-standing venous ulcers, burns, and scars. A wedge biopsy is taken from the edge of the Marjolin's ulcer for diagnosis.

The image below shows a Marjolin's ulcer in a burn scar:

Marjolin ulcer over the dorsum of foot measuring3 cm x 2 cm	Wedge biopsy
1.5 cm cervical lymph node in the neck	Excision biopsy
Soft tissue sarcoma over mid-thigh	Core needle biopsy
Solitary thyroid nodule	USG guided FNAC

In a patient with lymph node pathology, an excisional biopsy is performed as it provides the entire node for histopathological examinations.

Core needle biopsy is performed for:

• Soft tissue sarcomas.

• Breast lumps (Triple assessment).

USG-guided fine-needle aspiration cytology (FNAC) for a solitary thyroid nodule helps to diagnose various benign & malignant pathologies of the thyroid gland. However, this technique cannot differentiate between follicular adenoma and follicular carcinoma of the thyroid.

Solution to Question 24:

The given clinical scenario, along with the immunohistopathology findings showing positivity for CD117 and negativity for LCA, cytokeratin, and S100, is suggestive of gastrointestinal stromal tumors (GIST).

Other IHC markers mentioned are seen in:

• S-100: In melanoma (S-100,HMB-45,Melan-A).

• Cytokeratin: In squamous cell carcinoma (SCC).

• Leukocyte common antigen: In lymphomas.

Solution to Question 25:

Tension pneumothorax should be identified as the most life-threatening injury during the assessment of the breathing component of the primary survey (ABCDE protocol).

Both breathing and ventilation are assessed. Injuries that significantly impair ventilation in the short term include tension pneumothorax, massive hemothorax, open pneumothorax, and flail chest. These injuries should be identified during the primary survey and immediate intervention must be carried out.

Other options:

Options A and D: Cervical spine injury and laryngotracheal injury are identified during the airway assessment.

Option C: Blunt cardiac injury (and dissection of the thoracic aorta) are life-threatening injuries identified during circulation assessment.

Solution to Question 26:

Potassium iodide can be given in the 7-10 days preoperative period to reduce intraoperative bleeding during total thyroidectomy.

Preoperative preparation of a hyperthyroid patient (such as in Graves disease) includes:

• Potassium iodide/ Lugol's iodine reduces the vascularity of the thyroid gland and produces transient remission in a hyperthyroid patient.

• Anti-thyroid medications such as carbimazole/propylthiouracil (Option C) are started 6-8 weeks before the surgery. They help achieve a euthyroid state by decreasing the synthesis of thyroid hormones. The last dose of anti-thyroid medication is given in the evening before the day of surgery.

• Beta blockers such as propranolol/nadolol (Option B) reduce the risk of thyroid storm by blocking the adrenergic receptors. They should be continued for 7 days post-operatively as well. The appropriate dosages are propranolol 40 mg TDS or the longer-acting nadolol 160 mg once daily. They also inhibit the peripheral conversion of T4 to T3.

Option D: Betamethasone is a glucocorticoid that reduces the peripheral conversion of T4 to T3. It is not used preoperatively in thyroidectomy.

Solution to Question 27:

According to standard hand hygiene protocol, hand rub cannot be used if the hands are visibly soiled.

Hand hygiene is the most important way of preventing infection transmission, nosocomial infections and it must be performed by all individuals who are involved in patient care.

Routinely, alcohol-based hand rubs are preferred to soap and water handwash because they are faster, more efficient, and better tolerated by the hands. But, these hand rubs cannot be used in case of visibly soiled hands.

Soap and water hand hygiene method is preferred in cases of soiled hands and while dealing with spore-forming pathogens, including outbreaks of Clostridium difficile.

Hand hygiene must be performed in the following cases [WHEN] : (Options B,C and D)



The following infographic gives the steps for using hand rub [HOW] :

The following infographic gives the steps for washing hands with soap and water [HOW] :



Solution to Question 28:

The true statement among the given options is that low pH mainly activates lysosomal enzymes.

Lysosomes are membraned organelles that enclose all the hydrolyzing enzymes capable of breaking down biological polymers. The lysosomal enzymes are only active in acidic pH (low pH value) and this pH is maintained within the organelle. Hypoxic tissue injury during shock causes membrane damage, and extravasation of the enzymes from lysosomes. Thus, leading to tissue death.

Other options:

Option A: Anaerobic respiration produces lactic acid. Accumulation of lactic acid causes metabolic acidosis.

Option C: Peripheral vasoconstriction is seen in all types of shock, except septic shock.

Option D: Renal perfusion is maintained during the compensated shock stages. There is renal decompensation as shock progresses causing oliguria, and eventually anuria.

Shock is a state of circulatory failure resulting in tissue hypoperfusion and hypoxia. Prolonged shock eventually leads to irreversible tissue damage.

In cases of shock, there is inadequate oxygenation. Thus cells switch from aerobic to anaerobic respiration where lactic acid, not carbon dioxide, is produced. The accumulation of lactic acid in the blood leads to systemic acidosis.

Shock progresses through a series of well-defined stages:

• Initial nonprogressive stage: Compensatory neurohumoral mechanisms come into play to preserve tissue perfusion.

• Progressive stage: Tissue hypoperfusion leads to circulatory and metabolic disturbances. Acidosis occurs due to intracellular anaerobic glycolysis. This causes peripheral pooling, thereby leading to reduced cardiac output and disseminated intravascular coagulation (DIC).

• Irreversible stage: Tissue injury cannot be reversed even with hemodynamic correction. There is worsening myocardial contractility, renal function impairment, and bacteremia (due to the release of intestinal flora into the circulation).

The various types of shock are shown in the images below:





Hypovolemic shock



©MARROW

Obstructive shock



©MARROW

Neurogenic shock



©MARROW

Solution to Question 29:

Multiple stab injuries to the chest, reduced chest movement on the right side, tracheal deviation to the left side, and distension of neck veins suggest a diagnosis of tension pneumothorax on the right side. The next best step in the management would be needle thoracocentesis (emergent management).

ICD placement (Option B) is the definitive management of the tension pneumothorax, inserted after performing the initial emergency procedure.

Tension pneumothorax develops when a "one-way valve" air leak occurs causing air to get sucked in and get collected between the lung and chest wall causing the ipsilateral lung to collapse.

It is characterized by chest pain, tachypnea, tachycardia, hypotension, tracheal deviation away from the side of injury, unilateral absence of breath sound, cyanosis, and neck vein distension.

Tension pneumothorax is a clinical diagnosis and treatment is started without waiting for radiological confirmation.

The next step/ immediate management is chest decompression by rapid insertion of a large-bore cannula on the affected side (needle thoracocentesis) in the appropriate space.

If it is not successful, finger thoracostomy can be tried.

ICD placement is mandatory after needle or finger decompression of the chest and is the definitive management. This should be followed by confirmation of position by chest X-ray.

Recent update (Acc. to ATLS 10th edition):

• In adults, the needle thoracotomy location was changed from the 2nd ICS midclavicular line to the 5th intercostal space (ICS) mid-axillary line.

• In the pediatric age group, thoracotomy location remains the same, at the 2nd ICS midclavicular (MCL) line.

Solution to Question 30:

The correct option is A-3, B-1, C-2, D-4.

Nylon is a monofilament or braided multifilament type non-absorbable suture material. It is a polyamide polymer with low tissue reaction and is used in skin closure, abdominal wall mass closure, hernia repair, plastic surgery, neurosurgery, microsurgery, ophthalmic surgery etc. However, it also loses 15-20% of its tensile strength every year and degrades at the rate of 15-20% every year.

Polyglactin is a braided multifilament of type absorbable suture material. It is a copolymer of lactide and glycolide in a ratio of 90:10, coated with polyglactin and calcium stearate. It is a 'workhorse' suture used in general surgical practices gut anastomoses, vascular ligatures, subcuticular wound closures, and ophthalmic surgery.

Polypropylene is a monofilament, non-absorbable suture material. It is a propylene polymer. It is used in plastic surgery, cardiovascular surgery, hernia repair, ophthalmic surgery, and general surgical subcuticular skin closure. Polypropylene mesh is used for hernia repair.

A. Nylon 3-0	3. Midline laparotomy skin w ound closure
B. Polyglactin	1. Bowel anastomosis
C. Polypropylene 1-0	2. Suturing linea alba in the midline
D. Polypropylene 2-0	4. Fixing mesh in hernia surg ery

Solution to Question 31:

The correct sequence of management in a patient who presents to the casualty with an RTA is Option B: Cervical spine stabilization, Intubation, IV cannulation, CECT as per the ABCDE protocol.

Solution to Question 32:

The given clinical scenario and the chest X-ray showing air under the diaphragm is suggestive of a hollow viscus perforation occuring secondary to the peptic ulcer disease(PUD) in this patient. The next line of management in this condition is administration of IV fluids (option A) and exploratory laparotomy(option C).

Hollow viscus perforation is a life-threatening condition. In PUD, the perforation most commonly occurs as a complication of anterior duodenal ulcers and gastric ulcers.This results in leakage of gastric contents into the abdomen resulting in acute chemical peritonitis.

Clinical features of perforation include abdominal pain and distension, nausea, vomiting, tachycardia, tenderness, guarding & rigidity, rebound tenderness and decreased or absent bowel sounds. Fever may be present. Sometimes, features of septic shock can be seen at later stages.

An erect xray abdomen shows gas under the diaphragm. If the patient can't stand an x-ray can be taken in lateral decubitus position and may show dilated gas-filled loops of the bowel, suggestive of paralytic ileus.

The below image shows an x-ray radiograph of a hollow viscus perforation:



CECT (option B) is performed in hemodynamically stable patients and is the most sensitive investigation to detect minimal/ suspected pneumoperitoneum.

Management of a hollow viscous perforation is mainly resuscitation, IV fluids, analgesics, IV broad-spectrum antibiotics, and emergency laparotomy.

Upper GI endoscopy is done in a patient with upper GI hemorrhage (m/c variceal veins rupture) to diagnose and manage the source of bleeding with techniques such as banding and sclerotherapy (option D).

Solution to Question 33:

Longer duration of stay in the hospital is not an advantage of laparoscopic surgeries.

Laparoscopic surgeries are performed by introducing a rigid laparoscope through a port into the peritoneal cavity. Carbon dioxide is insufflated into the abdominal cavity to create a pneumoperitoneum. The other laparoscopic instruments used for dissection are introduced via other surgically created ports.

Evidence shows that laparoscopic surgery has short-term benefits over open surgery. Laparoscopic surgery has a better outcome with regard to blood loss, postoperative pain, length of stay, better cosmetic appearance, fewer adhesions, better anatomical perception, and earlier recovery.

The disadvantage of laparoscopic over open surgery is not knowing the extent of the thermal burns due to the relative lack of visibility of the entire instrument.

Solution to Question 34:

The number shown in the above image of the Foley catheter indicates the outer diameter.

"Fr" stands for French scale and refers to the external diameter of the catheter. 1Fr is 0.3 mm in diameter.

An 18 Fr. catheter will have a circumference of approximately 18 mm and a diameter of 6 mm.

Solution to Question 35:

Basal-like or triple-negative breast cancer on immunohistochemical staining is ER-, PR- and HER2-.

Immunohistochemistry staining:

Positive staining is indicated by the presence of brown-colored tissue.

• The estrogen receptor (ER) and progesterone receptor (PR) are both nuclear receptors, so the nucleus is stained brown if positive.

• The human epidermal growth factor receptor type 2 (HER2) is a membrane receptor, so the cell membrane is stained brown if positive.

The image below shows positive and negative staining for ER, PR, and HER2.



Molecular subtypes	ER	PR	HER 2	Ki-67	Cytokeratin 5 and 6
Luminal A(Most common type and has the best prognosis)	+ve	+ve	-ve	Low	-ve
Luminal B	+ve	+ve	-ve	High	-ve
	+ve	+ve	+ve	Any	-ve
HER2 enriched	-ve	-ve	+ve	Any	-ve

Molecular subtypes	ER	PR	HER 2	Ki-67	Cytokeratin 5 and 6
Basal-like or Triple-negative breast cancer(Aggressive, has worst prognosis but shows g ood response to chemotherap y, high PCR rates)	-ve	-ve	-ve	Any	+ve
Unclassified, now also known asClaudin-low	-ve	-ve	-ve	Any	-ve

Solution to Question 36:

The management of acute necrotizing pancreatitis (ANC) includes administering intravenous (IV) fluids, IV antibiotics, culture, and assisted drainage.

ANC is a complication of acute pancreatitis. Necrosis is the presence of nonviable pancreatic parenchyma or peripancreatic fat and can manifest as a focal area or diffuse involvement of the gland.

Steps in the management of ANC:

• Aggressive fluid resuscitation with isotonic crystalloid solution, pain control, and early nutrition, similar to acute pancreatitis

• CECT is done to diagnose ANC. The Balthazar Score is used to grade severity based on CT findings.

• Infection screening:

• One of the main complications of ANC is infection. If infected necrosis is suspected, fine-needle aspiration (FNA) is done, and the aspirate is subjected to Gram staining and culture sensitivity.

• Once the infection is confirmed, IV antibiotics are started. Carbapenem is the first-choice drug. The other drugs that can be used include quinolones, metronidazole, third-generation cephalosporins, and piperacillin.

• Definitive management:

• Traditionally, surgical debridement with necrosectomy, close continuous irrigation, or open packaging. Due to the high mortality, these surgeries are no longer performed, instead, a "step-up" approach has been adopted.

• Step-up approach: Percutaneous drainage followed by minimally invasive video-assisted retroperitoneal debridement.

Note: Pancreatic necrosectomy via midline laparotomy is considered only if sepsis persists despite conservative measures.

Solution to Question 37:

The mid-clavicular line is not a boundary of the triangle of safety for chest tube insertion.

Chest tube insertion is carried out in the 5th intercostal space (ICS) anterior to the midaxillary line in the triangle of safety (image below).

Boundaries of Triangle of Safety:

• Anteriorly (red line) : Below and lateral to the pectoralis major muscle

• Posteriorly(yellow line) : Anterior to the mid-axillary line (some references mention the lateral border of latissimus dorsi)

• Inferiorly (blue line) : Above the horizontal level of the nipple at the level of the 5th ICS

Chest tube insertion/ thoracostomy is indicated in pneumothorax, hemothorax, chylothorax, pleural effusion, etc. Contraindications include pulmonary adhesions, pulmonary trauma, coagulopathy, and diaphragmatic hernia.

Possible complications of this procedure include:

• Clogging of the tube (most common)

• Incorrect placement

• Intercostal nerve/ vessel injury

• Subcutaneous hematoma/ emphysema

• Pain

The image below depicts chest tube insertion in a step-wise manner:



Solution to Question 38:

The minimum number of lymph nodes to be examined for accurate staging of colon carcinoma and gall bladder carcinoma are correctly matched.

The minimum number of lymph nodes to be examined for effective staging are:

Type of Carcino ma	Minimum no. of Lymph n odes
Stomach CA	16
Colon CA	12
Gall Bladder CA	6
Breast CA	10

Solution to Question 39:

The most appropriate and specific answer, in this case, is that sciatica in patients with rectal carcinoma is due to anterior infiltration in females.

Sciatica in rectal carcinoma occurs due to perineural invasion of the inferior hypogastric plexus, from which the sciatic nerve arises. In females, the inferior hypogastric plexus lies anterolateral to the rectum in females, and in males, it lies lateral to the rectum.

With the given options, anterior infiltration in females is the most specific.

Solution to Question 40:

The correct statements regarding gastric carcinoma are 2 and 3.

Most cases of gastric cancer are sporadic (statement 1).

Lauren's system classifies gastric cancer into Type I - intestinal and Type II - diffuse.

At the molecular level,

• the diffuse-type is characterized by decreased E-cadherin mainly due to loss-of-function mutations in the CDH1 gene (statement 2).

• the intestinal type of cancer is characterized by mutations in the APC gene or β-catenin causes increased signaling via the Wnt pathway (statement 3).

Symptoms of both types include nausea, dyspepsia, dysphagia, weight loss, anorexia, early satiety, and anemia.

The image below shows the histopathological appearances of intestinal and diffuse types of gastric carcinoma:



Endoscopy and biopsy of the lesion (gold standard) are performed to diagnose gastric carcinoma.

When possible, surgery remains the preferred treatment approach. Local invasion into the duodenum, pancreas, and retroperitoneum is common, where chemo or radiation therapy with palliative care is the treatment.

The TNM staging is most widely used. It uses depth of invasion along with nodal and distant metastases as prognostic indicators (statement 4).



Solution to Question 41:

Preferential visualization of the gall bladder in a hepatobiliary iminodiacetic acid (HIDA) scan is not a feature of acute cholecystitis. On a HIDA scan, non-visualization of the gall bladder even after 3 hours of radioisotope administration suggests the diagnosis of acute cholecystitis.

Acute cholecystitis usually occurs due to the presence of gallstones (90%) leading obstruction of the cystic duct. This leads to inflammation, distension, and edema of the gallbladder. Acalculous cholecystitis and tumor obstructing the cystic duct are rare causes.

Clinical features of acute calculous cholecystitis: Fever, pain and right upper quadrant tenderness. A positive Murphy's sign (Option B) is right upper quadrant tenderness on palpation.

Diagnosis: An ultrasound scan is sensitive in acute cholecystitis and demonstrates a distended gallbladder with a thickened wall of >3 mm (Option A), pericholecystic fluid, the presence of gallstone(s), and the sonographic Murphy's sign. CT scan findings are similar to USG. CBC will show leukocytosis (Option C).

The following image depicts the confirmatory HIDA (Technetium-99m radioisotope) scan in which the gallbladder is not visualized even after 4 hours.



Management: Patients with acute cholecystitis are kept NPO and started on parenteral antibiotics. Cholecystectomy is the treatment of choice for acute cholecystitis. Treatment including cholecystectomy done within 72 hours has better outcomes that interval surgery.

Solution to Question 42:

The apical group of the axillary lymph nodes are present partly above the superior margin of the pectoralis minor and partly posterior to the superior part of the muscle. They are present along the axillary vessels.

The central group of axillary lymph nodes are present in the axillary pad of fat at the apex of the axilla.

Axillary lymph nodes collectively drain the entire upper limb, breast, and trunk above the umbilicus and can be divided into 5 groups namely:

- Lateral/ humeral
- Anterior/ pectoral
- Posterior/ subscapular
- Central
- Apical

Given below is a diagrammatic representation and details about the individual axillary groups of lymph nodes.



	Location	Afferents	Efferents
Humeral/ Lateral group	Posteromedial toaxillary vein	Whole limb except vessels accompanying cephalic veins	• Central nodes• Apical nodes• Inferior deep cervical nodes
Pectoral/ Anterior group	Alonglateral thoracic vessels	Skin and muscles of supra-umbilical anterolateral body wall and breast	• Central nodes• Apical nodes
Subscapular/ Posterior group	Along thesubscapular vessels	Skin and muscles of the inferior posterior region of the neck, dorsal aspect of the trunk up to iliac crest	• Central nodes• Apical nodes
Central group	Embedded in theaxillary padoffat	All preceding groups	• Apical group
Apical group	Deep to clavipectoral fascia, along theaxillary vessels	Those accompanying cephalic vein	Unite as subclavian trunk, right trunk drains into Jugulo-subclavian venous junction and left trunk ends in thoracic duct

Solution to Question 43:

The typhoid vaccine is not part of recommendations for a patient undergoing splenectomy.

Splenic insufficiency following splenectomy leads to increased susceptibility to sepsis caused by encapsulated bacteria such as pneumococci, meningococci, and H.influenzae. Such patients must be vaccinated against these agents to reduce the risk of sepsis.

Solution to Question 44:

Transverse colonic carcinoma is treated with an extended right colectomy. The vessels to be ligated are ileocolic, right colic, and trunk of the middle colic vessels.

Procedures performed for tumors of the colon are summarized below:

Tumor location	Procedure of choice
Cecum/ Ascending colon	Right hemicolectomyLigation and division of the ileocolic, right colic, and right branch of the middle colic vesselsResection of cecum + approx 10 cm of the terminal ileum with ileocecal junction + ascending colon + hepatic flexure
Hepatic flexure/Transverse colon	Extended right hemicolectomyLigation and division of theileocolic, right colic,and trunk ofmiddle colic vesselsResection of the cecum + approx 10 cm of the terminal ileum with ileocecal junction + ascending colon + hepatic flexure +proximal transverse colon(distal transverse colonandsplenic flexureare also sometimes removed)
Splenic flexure/ Descending colon	Left hemicolectomyLigation and division of the left branch of the middle colic vessels, the left colic vessel, and the first branches of the sigmoid vesselsResection of the transverse colon (left MCA territory) + Splenic flexure + Descending colon up to the sigmoid colon
Sigmoid colon	Sigmoid colectomy, or LAR (Low anterior resection)Ligation and division of the sigmoid branches of the inferior mesenteric artery.Removal of the entire sigmoid colon and rectum



Solution to Question 45:

All the given statements are true regarding central and peripheral venous catheters.

Peripheral intravenous cannulation is the insertion of a single-lumen plastic conduit across the skin into a peripheral vein. They are used in the transfusion of fluids, medications, and other therapies such as blood products. Phlebitis is the most common complication in peripheral venous access (Statement 1). Relative contraindications are coagulopathy, local infection, burns, arteriovenous fistula (Statement 3), or deep venous thrombosis on the affected limb.

Central venous catheter insertion is used in critically ill patients. It is inserted into a large, central vein. The most common veins are the internal jugular and subclavian veins. It is used in case of multiple infusions incompatible with peripheral intravenous access, and hemodynamic monitoring. Vasopressors are an integral component in the management of septic shock and are given in the central line (Statement 4).

To prevent or avoid the formation of thrombus in venous catheters, the solutions used include 0.9% sodium chloride, heparin (Statement 2), vitamin C, and sodium citrate.

Solution to Question 46:

In the given scenario, the ileocolic artery should have been ligated.

The right colic, superior rectal and superior mesentric arteries are not ligated during an appendicectomy.

When an appendicectomy is performed, the inferior branch of the ileocolic artery needs to be ligated at the base of the appendix. Otherwise, postoperative hemorrhage can occur.

The appendicular artery usually arises directly from the ileocolic artery and descends posterior to the terminal ileum to enter the mesoappendix. The artery runs very close to the free margin of the

mesoappendix and terminates in multiple small branches that supply the appendix.

The terminal part of the artery lies on the wall of the appendix and may become thrombosed in appendicitis, resulting in distal gangrene or necrosis.

The blood supply of the appendix is shown in the image below.





Solution to Question 47:

Fibroadenoma is a benign tumor in a 25-year-old woman with no skin changes and no indurations and is categorized as BIRADS 2.

Fibroadenoma is a form of aberrations of normal development and involution of the breast (ANDI), and it is the most common cause of a breast lump in women aged 15–25 years.

It arises from hyperplasia of a single lobule and presents as a well-circumscribed, smooth or lobular, firm, non-tender, freely mobile mass (mouse in the breast). The histopathology shows the proliferation of intralobular stroma, and the epithelium may be surrounded by stroma (pericanalicular pattern) or compressed and distorted (intracanalicular pattern).

USG is done to confirm the diagnosis. A biopsy is needed if USG shows atypical features or the age of presentation is above 25 years.

The image below shows the histology of fibroadenoma:



Solution to Question 48:

Tumor size < 4cm, with N1 axillary metastasis is not a contraindication for breast conservative surgery.

Breast conservative therapy (BCT) involves resection of primary breast cancer with adequate normal breast tissue, adjuvant radiotherapy, and assessment of regional lymph nodes via a separate incision in the axilla. Removal of primary breast cancer is also called lumpectomy/ tylectomy/ partial mastectomy/ wide local excision.

Absolute contraindications of BCT:

• Pregnancy

• Persistently positive margin (2 microscopy shows positive margins)

• Multicentric lumps

• Diffuse microcalcification

•

Extensive ductal carcinoma in situ (DCIS)

•

Inflammatory breast cancer

Relative contraindications of BCT:

• Active SLE (systemic lupus erythematosus), scleroderma

• Large lump-to-breast ratio (>5 cm size lump)

• Known or suspected genetic predisposition to breast cancer (i.e., BRCA1, BRCA2), Li-Fraumeni syndrome (p53 mutation)

• Multifocal disease

• Prior radiation therapy to the chest wall or breast

Surgery NEET 2022

Question 1:

A 6-year-old boy came with a history of recurrent urinary tract infections. Imaging was done and is shown below. What is the diagnosis?



a) Vesicoureteric reflux

b) Urinary bladder diverticulum

c) Urinary bladder hernia

d) Vesicocolic fistula

Question 2:

A 45-year-old female patient underwent thyroidectomy. Three days after the surgery, she developed perioral numbness. Which of the following investigations need to be done for her?

a) Free T_3, T_4

b) T_3, T_4, Thyroid-stimulating hormone

c) Radioiodine scan

d) Calcium, phosphate and parathormone levels

Question 3:

A patient presented with blunt trauma to the abdomen. On evaluation, liver injury was noted for which primary repair was done. Coagulation function was monitored intraoperatively, using the method shown below. What is the method used?



a) Thromboelastography

b) Plethysmography

c) Sonography

d) Elastography

Question 4:

A 59-year-old lady presents with a progressive painless lump in the breast. What is the cause for the following skin change?

a) Infiltration of subdermal lymphatics

b) Infiltration of lactiferous duct

c) Involvement of Cooper's ligament

d) Spread of tumor to anterior chest wall

Question 5:

A child had a history of stab injury on the anterior abdominal wall and the image is shown below. The child is hemodynamically stable. Which will be the next course of treatment?

a) Emergency laparotomy

b) Observation

c) Intravenous hydration

d) Wait and watch

Question 6:

Which of the following is most likely to be seen due to rupture of a saccular aneurysm?

a) Subdural hemorrhage

b) Subarachnoid hemorrhage

c) Intracerebral hemorrhage

d) Hydrocephalus

Question 7:

A patient presents with sudden-onset right leg pain. An investigation was done, and the obtained image is shown below. What is the investigation?



a) Digital subtraction angiography

b) Ultrasound Doppler

c) MR angiography

d) Plethysmography

Question 8:

A male patient presented with a 0.3-cm nodule on the left nasolabial fold. It was excised and a pathological examination was done. The image is given below. What is the diagnosis?



a) Basal cell carcinoma

b) Melanoma

c) Squamous cell carcinoma

d) Nevus

Question 9:

A 45-year-old patient complained of pain in one side of the neck. She is afraid of eating food as it worsens the pain. Ultrasound imaging of the salivary glands is done and shown below. What is the most likely diagnosis?



a) Sialolithiasis

b) Foreign body

c) Osteoma of floor of mouth

d) Cervical lymphadenopathy

Question 10:

A delayed intravenous urogram of a patient is given below. What is the likely diagnosis?

a) Pelviureteric junction obstruction

b) Putty kidney

c) Staghorn calculus

d) Cystic kidney

Question 11:

A young male patient with a history of a motor vehicle accident is unable to pass urine. Blood is seen at the meatus. What is the most likely site of urethral injury?

a) Bulbar urethra

b) Spongy urethra

c) Membranous urethra

d) Penile urethra

Question 12:

A male patient presented with a midline neck swelling. He later developed cervical node enlargement. The histopathology of the lesion is shown below. Which of the following statements is false about this condition?



a) Excellent prognosis is associated with this condition

b) It spreads quickly via lymphatics

c) Nuclear characteristics are used for identification

d) Fine needle aspiration cytology (FNAC) is not diagnostic

Question 13:

A 40-year-old man who is a known case of hypertension presented with multiple episodes of hematuria and loin pain. His elder brother passed away due to stroke at the age of 40. The ultrasound abdomen is shown below. What is the probable diagnosis?



a) Renal cell carcinoma

b) Autosomal dominant polycystic kidney disease

c) Tuberculosis of the kidney

d) Autosomal recessive polycystic kidney disease

Question 14:

A patient presents with acute-onset, severe abdominal pain. He is hemodynamically stable. A chest x-ray is shown below. What is the next step in management?



a) Gastric lavage

b) Emergency laparotomy

c) Ultrasound

d) Chest tube insertion

Question 15:

Calculate the percentage of burns in a preschool child shown in the image below.



a) 15-20%

b) 25-30%

c) 35-40%

d) 10-15%

Question 16:

A 32-year-old patient who is a chronic alcoholic presents with oral ulcers and a burning sensation. A picture of the oral cavity is given below. What is the most likely diagnosis?

a) Leukoplakia

b) Erythroplakia

c) Submucosal fibrosis

d) Malakoplakia

Question 17:

A patient presents with a tender pulsatile mass as shown in the image below. What will be done next for this patient?

a) Ultrasound doppler

b) Needle aspiration

c) CT angiogram with percutaneous management

d) Intravenous antibiotics for 7 days

Question 18:

A patient presents with carcinoma of the prostate, which has now spread to the vertebra. What is the route of spread to the lumbar vertebra?

a) Prostatic venous plexus

b) Transcoelomic spread

c) Inferior vesical vein

d) Internal iliac vein

Question 19:

A male patient presents with impotence and bilateral resting pain. The following finding is seen in the legs. At which level has the pathology occurred?

a) Bilateral popliteal arteries

b) Bilateral internal iliac arteries

c) Aortoiliac bifurcation

d) Bilateral femoral arteries

Question 20:

An elderly male patient presented with a sudden onset of scrotal pain and discharge. Based on the image below, what is the likely diagnosis?

a) Torsion of testis

b) Acute epididymo-orchitis

c) Scrotal carcinoma

d) Fournier's gangrene

Question 21:

Identify the fluid shown below.



a) Haemaccel

b) Ringer lactate

c) Isolyte P

d) Isolyte M

Question 22:

The image below shows a pressure sore. Which stage does this belong to ?



a) Stage 1

b) Stage 2

c) Stage 3

d) Stage 4

Question 23:

Identify the most common site of an intra-abdominal abscess.



a) 1

b) 2

c) 3

d) 4

Question 24:

A man came with complaints of recurrent discharge and pain due to lesions around the anus for 3 years. What is the diagnosis?

a) Pilonidal sinus

b) Carbuncle

c) Boil

d) Fistula in ano

Question 25:

According to triage, which of the following categories of patients comes under green?

a) Ambulatory patients

b) Medium risk patients

c) High risk patients

d) Dead patients

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	a
5	a
6	b
7	b
8	a
9	a
10	a
11	c
12	d
13	b
14	b
15	b
16	a
17	a
18	a
19	c

20	d
21	b
22	d
23	d
24	d
25	a

Detailed Explanations

Solution to Question 1:

In the given scenario of a 6-year-old boy with recurrent urinary tract infections and the micturating cystourethrogram showing a dilated ureter, the likely diagnosis is vesicoureteric reflux.

Vesicoureteric or vesicoureteral reflux (VUR) is a condition in which there is a retrograde urinary flow from the bladder to the ureter and kidney. It is usually congenital.

VUR can cause urinary stasis and lead to urinary tract infection (UTI). It also predisposes to pyelonephritis, which can lead to renal scarring due to parenchymal inflammation. This is known as reflux nephropathy.

Micturating cystogram is used to confirm the VUR. VUR can be graded based on the appearance of the urinary tract on the micturating cystogram. It is as follows:

Grade 1- reflux into the ureter

Grade 2- reflux into the ureter and renal pelvis

Grade 3- mild to moderate dilatation on IVU along with reflux and mild blunting of calices

Grade 4- blunting of fornices

Grade 5- absent papillary impressions



Grades 1-3 resolve spontaneously. Surgical management is done when acute pyelonephritis recurs in spite of antibiotic therapy or if severe reflux is associated with a defect that can be corrected surgically.

Other options:

Option B: Urinary bladder diverticula occurs due to pressure changes in the bladder leading to the formation of saccules that may protrude through the bladder wall. It is more common in elderly males >50 years and is usually asymptomatic or not infected. Symptoms may occur due to associated urinary obstruction, recurrent infection, or pyelonephritis. They are diagnosed incidentally on cystoscopy or ultrasonography.

Option C: Urinary bladder hernia is an uncommon condition characterized by herniation of the urinary bladder or ureter into the inguinal canal, scrotum, or femoral canal. Urinary symptoms such as dysuria, frequency, urgency, nocturia, and hematuria can be seen. Retrograde cystography is the best technique to image a bladder hernia.

Option D: Vesicocolic fistula is an abnormal communication between the bladder and the colon. As a result of this, the fecal matter may enter the bladder. Diverticulitis is the most common cause. Bubbles in the urine, pain, hematuria, and recurrent UTI can occur. Cystoscopy reveals edema and gas bubbles. The fistula can be seen in a cystogram.

Solution to Question 2:

Calcium, phosphate, and parathyroid hormone (PTH) levels should be done in this patient to evaluate hypoparathyroidism secondary to thyroidectomy.

Hypoparathyroidism is a common complication of thyroid surgery. It is temporary and usually lasts for 4-6 weeks. Hypoparathyroidism following thyroid surgery can be diagnosed when there are decreased serum calcium and/ or PTH levels, or if there is symptomatic hypocalcemia.

The features of hypocalcemia are circumoral paresthesia, and paresthesia of the neck, fingers, and toes. Muscle twitching, weakness of the forearm, hand, foot, and carpopedal spasms can be

seen. A carpal spasm can be induced by inflating a blood pressure cuff to 20 mmHg above the systolic blood pressure for 3 minutes. This is known as Trousseau's sign. Chvostek sign wherein twitching of facial muscles is demonstrated by tapping above the angle of the jaw. It is due to the stimulation of the branches of the facial nerve.

ECG shows prolonged QT interval.

Management of acute and symptomatic hypocalcemia is by giving 10 mL of intravenous 10% calcium gluconate over 5 minutes.

Solution to Question 3:

In the case of liver injury, intraoperative monitoring of coagulation function is done by thromboelastography which is shown in the given image.

Thromboelastography (TEG) is a non-invasive test that quantifies the ability of whole blood to form a clot. It is useful in intraoperative coagulation status assessment. The dynamic changes in the viscoelastic properties of a blood sample during the clotting can be detected and quantified by thromboelastography. Underlying coagulation deficiencies show different shapes in the TEG. They are shown in the image as follows



Advantages of thromboelastography:

-

Blood coagulability, including coagulation cascade, platelet function, and fibrinolysis can be assessed.

-

It is a simple bedside test

-

Can diagnose coagulopathic bleeding

- Can be used as a guide in transfusion therapy and decrease the use of blood products
- Can detect dynamic changes in blood coagulation during resuscitation
- Predict the clinical efficacy of therapeutic agents affecting blood coagulability

Other options:

Option B: Plethysmography is a semiquantitative method to measure segmental blood flow and velocity in the carotid and peripheral vascular systems. It is safe, inexpensive and easy to use. It is useful in the diagnosis of vascular disease.

Option C: Sonography or ultrasound is a cheap and easily available imaging method. It is used as a first-line imaging method to assess liver, biliary, renal tract, obstetric, and gynecological disease. It is also useful in guiding needle placement during interventional procedures. Ligaments, tendon and muscle injuries can also be studied by an ultrasound.

Option D: Elastography is a newer imaging modality, which detects the changes in elasticity of tissues. It is based on the concept that the pathological process alters the elastic properties of tissue. It is useful in differentiating malignant and benign breast neoplasms, identifying traumatic changes in the muscles and tendons, helping to decide the biopsy site accurately, assess liver fibrosis and steatosis.

Solution to Question 4:

The image shows the peau d'orange appearance and it occurs due to the infiltration of the subdermal lymphatics.

Peau d'orange appearance is a skin change in which the skin of the breast appears like that of an orange. It is due to cutaneous lymphatic edema caused by dermal lymphatic infiltration by malignant cells. It is seen in several conditions such as:

- inflammatory carcinoma of the breast
- mastitis
- chronic abscess
- following breast surgery or radiation

Other options:

Option B: Infiltration of lactiferous duct leads to nipple retraction.

Option C: Involvement of Cooper's ligament leads to dimpling of the skin over the breast.

Option D: In case of spread of the tumor to the anterior chest wall, the breast will appear fixed and will not fall forwards when bending. On lifting the arm above the shoulder, the breast will not move upward.

Solution to Question 5:

The next course of treatment in the given scenario of a stab injury to the abdomen with the image showing evisceration of the omentum is emergency laparotomy.

Emergency laparotomy is performed to explore and repair abdominal injuries in case of

• Hemodynamic instability

• Peritonitis

• Evisceration

• Gunshot wounds

The algorithm for management of stab wounds of the abdomen is as follows



Solution to Question 6:

Subarachnoid hemorrhage is most likely to be seen due to the rupture of a saccular aneurysm.

Saccular aneurysms are thin-walled outpouchings measuring 2 or 3 cm in diameter with a bright, red, translucent wall. These are mainly seen at the arterial branching points along the circle of Willis.

The absence of smooth muscle and internal elastic lamina is the main pathogenesis in saccular aneurysms. Female sex, smoking, cocaine abuse, and family history are some other risk factors.

Genetic conditions predisposing to subarachnoid hemorrhage are:

• Adult polycystic kidney disease

• Fibromuscular dysplasia

• Neurofibromatosis type 1

• Ehler Danlos syndrome and Marfan's syndrome.

Clinically, subarachnoid hemorrhage presents with a sudden and severe headache commonly referred to as thunderclap headache. It may be associated with seizures, unresponsiveness, and vomiting.

CT scan is the imaging method of choice. It shows blood in basal cisterns and aneurysms appear as round enhancing lesions. Management includes craniotomy and clipping or coiling by endovascular embolization. Medical management includes fluid replacement with normal saline, oral nimodipine 60 mg every 4 hours, and supportive measures.

Hyponatremia can be a frequent complication in subarachnoid hemorrhage. The below image shows a CT of subarachnoid hemorrhage with blood in basal cisterns.



Other options:

Option A: Subdural hemorrhage or subdural hematoma occurs mainly due to rupture of the cortical bridging veins. Symptoms of raised intracranial pressure (ICP) such as headache, vomiting, blurring of vision, as well as focal deficits are seen.

Option C: Intracerebral hemorrhages may spontaneously occur in patients with hypertension, amyloid angiopathy, coagulation disorders, or as a complication of ischemic stroke. Headache, raised ICP, and focal neurological deficits can be present.

Option D: Hydrocephalus is a condition where the cerebrospinal fluid (CSF) circulation is obstructed or there is a failure to resorb thereby leading to an increase in CSF volume with enlargement of the ventricles. Symptoms of raised ICP are seen.

Solution to Question 7:

The image shows a triphasic pattern of the arterial waveform, which is seen in the ultrasound doppler.

Doppler ultrasound is used in diagnosing peripheral arterial disease. It uses a handheld doppler ultrasound probe placed at an arterial site. This transmits a continuous wave. A receiver inside the probe picks up the reflected beam. The change in frequency between the reflected beam and the

transmitted beam gets converted to a pulsatile audio signal.

A normal arterial wave has a triphasic signal. A triphasic wave has three waveforms consisting of a sharp systolic wave with a forward uprise and fall, a second wave with reverse flow occurring in diastole, and the last wave with a forward flow taking place during diastole.

Doppler ultrasound can detect moving blood. However, it does not provide information on whether the blood flow is sufficient for the maintenance of limb viability.

Other options:

Option A: Digital subtraction percutaneous angiography (DSA) is a procedure wherein a radiopaque dye is usually injected percutaneously into the femoral artery. Images obtained are digitalized and the background may be removed for a clear image. It provides information on dynamic arterial flow and can be combined with endovascular intervention if needed.

Option C: Plethysmography is a semi-quantitative method to measure segmental blood flow and velocity in the carotid and peripheral vascular systems.

Option D: Magnetic resonance angiography (MRA), or MR angiography, is an imaging modality that can be used when duplex ultrasonography cannot be done or if it cannot produce a clear image. MRA avoids ionizing radiation.

Solution to Question 8:

The given clinical scenario of a nodule on the nasolabial fold and the histopathological image showing peripheral palisading is likely to be basal cell carcinoma (BCC).

Basal cell carcinoma is a slow-growing malignant tumor, with locally invasive characteristics. It affects the pilosebaceous skin.

Ultraviolet radiation exposure is the strongest predisposing factor. Other risk factors include exposure to arsenic, coal tar, aromatic hydrocarbons, and ionizing radiation.

It mainly affects the elderly or middle-aged men who are excessively exposed to sun, common in the age group 40-80 and white-skinned people.

Macroscopically, it is categorized as localized and generalized. Nodular, nodulocystic (nodular and nodulocystic together account for 90% of BCC), pigmented, and nevoid are the localized cancers while superficial multifocal and superficial spreading, infiltrative, and cicatrizing are the generalized types. Nodular type is described as pearly waxy nodules at the margins. A central depression with umbilication or ulceration and rolled-out edges are characteristic of BCC.

The below gross image shows pearly, rolled-out borders suggestive of basal cell carcinoma.

Histopathologic features show ovoid cells in nests with a single palisading layer. The outer layer of cells is in the active division and thus incomplete excision can lead to being more aggressive. The below image shows the histopathology of BCC.



Sites near the eye, nose, and ear and size >2 cm are considered to be high risk. Direct invasion at these sites can reach the cranium.

Management is by Moh's microscopic surgery. Non-surgical measures such as topical 5-fluorouracil, and imiquimod are useful. Metastasis is extremely rare.

Other options:

Option B: Melanoma is a malignancy, commonly seen in young adults, arising from melanocytes and involves the skin, mucosa, retina, and leptomeninges. Microscopically, malignant changes are seen in the basal epidermis with breach and invasion of the dermis as shown below.



Option C: Squamous cell carcinoma is the second most common malignant tumor, seen more commonly in men. Microscopically, the proliferation of irregular masses of squamous epithelium is seen. The image is shown below.



Option D: Nevus are aggregates of melanocytes at the dermis or dermo-epidermal junction. It can get transformed into malignant melanoma and the features suggestive of malignant transformation are a change in size, shape, color, thickness, satellite lesions, and tingling/ itching/serosanguinous discharge.

Solution to Question 9:

The given clinical scenario of pain on one side of the neck that worsens with food and the ultrasound image showing a stone within Wharton's duct is suggestive of sialolithiasis (salivary

gland stone).

The submandibular gland has highly viscous secretions and hence forms the most common site for salivary gland stone formation. The two most common sites of stone impaction are near the punctum and the hilum of the gland where the duct bends over the mylohyoid muscle.

The patient usually presents with acute painful swelling in the submandibular region, which is aggravated with food intake. Symptoms resolve 1-2 hours after the meal is completed. On examination, a tender and enlarged submandibular gland can be noted on bimanual examination. Pus may be seen draining from the sublingual papilla or can be expressed on bimanual palpation of the gland.

Most submandibular stones are radio-opaque and visible in radiographs. Stones of size <4 mm are removed via a dormia basket using endoscopic or ultrasound-guided methods under local anesthesia. Stones of size >4 mm are broken first using intracorporeal or extracorporeal lithotripsy and then removed using a dormia basket.

Submandibular gland excision is done only when minimally invasive techniques have failed or there is evidence of a salivary gland tumor.

Nerves at risk during the removal of the submandibular gland:

• the marginal mandibular branch of the facial nerve

• the lingual nerve

• the hypoglossal nerve





Solution to Question 10:

The delayed intravenous urogram (IVU) of the patient showing hydronephrosis with an absence of contrast in the ureter makes the likely diagnosis a pelviureteric junction obstruction.

Pelviureteric junction obstruction is the commonest type of urinary obstruction in childhood, but can also be seen in young adults. It is usually caused by intrinsic stenosis. It manifests as abdominal flank or back pain, and hematuria. In newborns, a renal mass is palpable.

Antenatal sonography is the least invasive technique to diagnose PUJ obstruction in utero and reveals fetal hydronephrosis. intravenous urography is useful if the kidney is functioning. Extrarenal pelvis dilatation, clubbing of the minor calyces, and slow diffusion of contrast down the blockage can be seen. A renal isotope scan using intravenous DTPA or MAG-3 is the best modality to confirm the obstructive dilatation of the collecting system.

The treatment is surgical if there are bouts of renal pain, increasing hydronephrosis, evidence of parenchymal damage, and infection.

Other options:

Option B: Putty kidney is an end-stage manifestation of renal tuberculosis where there is amorphous dystrophic calcification involving the entire kidney. It is shown in the image below.

Option C: Staghorn calculus is seen following urinary infections with Proteus, Pseudomonas, and Staphylococcus. These organisms can break down urea to produce ammonia and carbon dioxide. The urine becomes alkaline and forms struvite calculi which grow into staghorn calculi. A staghorn calculus in the left kidney is shown in the image below.

Option D: Cystic kidney is seen in autosomal dominant polycystic kidney disease (ADPKD) due to mutation in genes PKD1 and PKD2. It is associated with hypertension. Multiple renal bilateral cysts will be seen, along with cysts in other organs as well such as the liver, pancreas, etc.

Solution to Question 11:

In the given scenario of a young patient with urinary retention, blood at the meatus, and a history of a motor vehicle accident, the most likely site of injury is the membranous urethra. The given

retrograde urethrogram shows contrast leakage at the membranous urethra.

Urethral injuries in men usually occur after catheterization or during cystoscopy. Injuries following external trauma are rare, but mainly affect the bulbar or the membranous urethra.

Membranous urethral rupture usually occurs following a road traffic accident, crush injury, or a fall. It is associated with pelvic fracture and extraperitoneal bladder rupture. Clinical features include urinary retention, blood at the meatus, and a high riding prostate on digital examination. Bruising is seen over the pubic area, scrotum, and penis.

Management is by inserting a suprapubic catheter. If there is an associated bladder rupture, the bladder will not be seen on the ultrasound, and management in such a case is by exploration, repair, and placement of a suprapubic catheter.

Other options:

Option A: Bulbar urethral injury occurs following a blow to the perineum with a fall astride injury during cycling or gymnasium accidents. A typical feature is a butterfly-shaped hematoma in the perineum. Other features such as urinary retention and meatal bleeding can also be present. Analgesia, antibiotics, and suprapubic catheterization are needed. The patient should be discouraged from passing urine. Definitive management is delayed urethroplasty after 8-12 weeks.

Options B and D: Spongy and penile urethral rupture can occur following a penile fracture during coitus. Hematuria and difficulty in voiding can be present in spongy urethral injury. A pop sound followed by severe penile pain and penile deformity is characteristic of penile urethral injury. Hematoma, blood at the meatus, and voiding difficulties are also seen.

Retrograde urethrogram shows loss of smooth contour of the urethra with extravasation of the dye in case of a spongy urethral rupture. Surgical repair is needed in both.

Solution to Question 12:

The given scenario of a male patient with a midline swelling, cervical lymph node enlargement, and the histopathology showing branched papillae with orphan Annie eye nuclei is suggestive of papillary thyroid carcinoma. The false statement regarding the condition is that FNAC is not diagnostic.

Papillary thyroid carcinoma is the most common thyroid malignancy. It is more commonly seen in women between the ages of 30 and 50. Radiation exposure and family history of thyroid malignancies are the common etiologies. Genetic abnormalities in either the RET gene or the BRAF gene can be seen in papillary thyroid carcinoma.

Clinical features include an asymptomatic thyroid swelling with cervical lymph node enlargement. In advanced disease, there may be features like hoarseness, dysphagia, cough, or dyspnea.

An ultrasound scan accurately assesses the size and number of nodules, as well as categorizes the lesions as benign, indeterminate, or malignant. Fine needle aspiration cytology (FNAC) is a diagnostic test, helpful in investigating an indeterminate or malignant lesion.

Histopathology shows the following features:

• Branching papillae

• Orphan Annie eye nuclei: optically clear nuclei with fine nuclear chromatin

• Pseudo inclusions: invaginations of the nuclear membrane

• Psammoma bodies: concentric calcifications in the core of the papillae.

Papillary thyroid carcinoma is known to spread via lymphatics, hence lymph node metastasis is commonly seen.

Management is by thyroidectomy with neck dissection (central or local depending upon the site involved) followed by lifelong levothyroxine supplementation. Radioactive iodine is also useful.

The disease has an excellent prognosis, with a high survival rate.

Solution to Question 13:

This clinical scenario with a family history of stroke and USG showing a cystic appearance of the kidney is suggestive of autosomal dominant polycystic kidney disease (ADPKD).

ADPKD is inherited as an autosomal dominant trait with mutations in the PKD1(chromosome 16) and PKD2 genes (chromosome 4).

The disease does not usually manifest before 30 years of age, and in some patients, it is never diagnosed. The progression of the disease is faster in males.

Hypertension is the most common presentation. Renal manifestations of ADPKD include:

• Flank/loin pain - either due to hemorrhage inside cysts or due to renal stones (most commonly uric acid stones)

• Irregular upper quadrant renal mass

• Hematuria

• Uremia

• Renal failure (seen in 10–15% of the patients who receive hemodialysis)

The most common extrarenal manifestation is hepatic cysts, which are derived from the biliary epithelium. Other extra-renal manifestations of ADPKD:

• Pancreatic cysts

• Berry aneurysms (Rupture of cerebral aneurysms is an important cause of morbidity and The prevalence of intracranial aneurysms in patients with PKD is 40%.)

• Colonic diverticula

• Abdominal hernias

• Mitral valve prolapse

Screening cerebral angiography is strongly considered in all patients with PKD and a family history of cerebral aneurysm or subarachnoid hemorrhage.

Intravenous pyelography shows characteristic spider leg sign, as a result of stretched calyces due to multiple cysts. The nephrogram shows a swiss cheese appearance. On prenatal USG, the presence of either > 3 cysts in unilateral kidneys or > 2 cysts in each kidney is suggestive of PKD.

Treatment includes

• Surgical decompression is not a routine procedure performed in these patients. It may be required only in cases of very large cysts causing persistent pain and infected cysts.

• Analgesics

• Increased fluid intake

• Prevents renal stone formation

• Reduces growth of cysts by suppressing ADH release

• ACE inhibitors and ARBs

• Blood pressure control

• Slowing of cyst growth

• Tolvaptan (V2 receptor antagonist)

• Reduce pain and slow the decline in renal function

• Long-term dialysis

• Renal transplant

• Definitive management

• Pre-transplant nephrectomy may be required in patients with very large polycystic kidneys

Autosomal recessive polycystic kidney disease (ARPKD) occurs secondary to a mutation of the PKHD1 gene. The severe form manifests in-utero or during infancy; milder cases can manifest later in childhood. 50% of affected newborns die in the first few days of life.

Renal cell carcinoma (RCC) is a relatively rare complication of ADPKD. There does not seem to be an increased frequency of RCC in these patients compared to the general population. However, when it occurs, it tends to be bilateral, multicentric, and of sarcomatoid type.

Tuberculosis kidney presents as putty kidney characterized by amorphous dystrophic calcification.

Solution to Question 14:

The given clinical scenario and the chest radiograph showing air under the diaphragm are suggestive of a hollow viscus perforation. The next best step in management among the given options is emergency laparotomy.

Perforation of a viscus is a life-threatening condition that occurs following the rupture of the alimentary tract. This leads to the leakage of the enteric contents, thereby leading to acute bacterial peritonitis. Peritonitis is the generalized or localized inflammation of the peritoneum either due to the free spillage of fluids into the peritoneal cavity or secondary bacterial infection.

Clinical features of hollow viscous perforation occur due to the developed peritonitis and the features include abdominal pain and distension, nausea, vomiting, tachycardia, tenderness, rebound tenderness, decreased or absent bowel sounds, and guarding on palpation. Fever may be present. Sometimes, features of septic shock can be seen at later stages.

Investigations include estimating the white blood cell count, urea, serum electrolytes, and an erect abdominal radiograph showing free gas under the diaphragm. A supine radiograph may show dilated gas-filled loops of the bowel, suggestive of paralytic ileus. The below image shows a hollow viscus perforation.



Management of a hollow viscous perforation is mainly resuscitation, analgesics, intravenous broad-spectrum antibiotics, and emergency laparotomy. Peritonitis is managed by nutritional support, pain relief, correction of fluid loss and circulating volume, urinary catheterization with or without gastrointestinal compression, antibiotics, analgesics, and specific surgical treatment of the cause.

Solution to Question 15:

From the given image, the percentage of burns in the child is 25-30% calculated based on Lund and Browder chart.

In the given image, the right arm, chest, abdomen, and both thighs are involved in burns. According to Lund and Browder chart, 2% is for the right arm, 13% for chest and abdomen involvement, and 4% for each thigh. Hence, the total involvement is 2+13+4+4 = 23%. The closest value to 23% among the given options is 25-30% (option B).

Lund and Browder chart is an accurate method to measure burns if it is used correctly. In this, different proportions of body surface areas in children are taken into consideration. This chart compensates for the changes in the shape of the body with age and can accurately assess the burns area in children.

The below image shows the Lund and Browder chart.



Relative percentages of areas affected by growth (age in years)

	0	1	5	10	15	Adult
A: half of head	9½	8½	6½	5½	4½	3½
B: half of thigh	2¾	3¼	4	4¼	4½	4¾
C: half of leg	2½	2½	2¾	3	3¼	3½

©MARROW

Note: Wallace's rule of nines is used in estimating medium to large burns in adults. It helps in the quick estimation of the burns. In this, the total burn is calculated by dividing the body into areas of 9%. It is not accurate in children.

Solution to Question 16:

The most likely diagnosis in the given scenario of a 32-year-old chronic alcoholic with oral ulcers and a burning sensation and the image showing white lesions on the oral cavity is leukoplakia.

Leukoplakia is a premalignant condition of the oral cavity characterized by whitish patches or plaques that cannot be rubbed off.

They appear as small, well-circumscribed lesions and can extensively involve a large area of the oral mucosa.

Variants of leukoplakia are

• Speckled leukoplakia is whitish lesions on an erythematous base. They have the highest risk of malignant transformation.

• Proliferative verrucous leukoplakia is rare and has a high risk for malignant transformation.

Management includes smoking cessation and reduction in alcohol consumption. A biopsy should be done and a photographic record of the lesion is used for follow-up.

Other premalignant conditions of the oral cavity are erythroplakia and oral submucosal fibrosis.

Erthroplakia (option B) refers to irregular, reddish velvety plaques on the oral cavity. The surface may be nodular and may bleed. It has a high risk of malignant transformation. Biopsy should be done followed by surgical ablation.

Oral submucosal fibrosis (option C) is characterized by the formation of fibrous bands beneath the oral cavity. The disease is progressive and causes limitation of mouth opening and restricted tongue movements. Intralesional steroids and surgical excision of fibrous bands are recommended.

Malakoplakia (Option D) is a rare inflammatory condition affecting immunocompromised individuals. It mainly affects the urinary tract but can involve any organ system. The clinical presentation includes tender ulcerations, nodules, papules, and plaques or fungating masses with perianal and genital skin involvement. Systemic antibiotics and surgical excision are recommended.

Solution to Question 17:

The given scenario of a patient with tender pulsatile mass, and the image showing an erythematous swelling at the base of the neck on the left side, is likely to be a vascular anomaly. Ultrasound Doppler should be done next in this patient.

The diagnosis of vascular anomalies is done clinically and with imaging like ultrasound Doppler. MRI plays an important role in diagnosing the type and extent of the lesion. CT angiogram can be used to find out the feeding vessel (option C).

Vascular anomalies include malformations and tumors. Vascular malformations occur as a result of developmental abnormality of an artery, vein, capillary or lymphatic system. They can be pulsatile or have a thrill when the underlying component is an artery.

Hemangiomas are the most common vascular tumors. They are seen in infants and present as bright red lesions. Common sites of a hemangioma include the head and neck. It can be of two types capillary and cavernous hemangioma. Compressibility and bluish surface are some clinical features of hemangiomas.

These lesions can be treated by complete resection. Debulking is done when complete resection is not possible. Sclerotherapy is mainly useful in venous malformation. Embolization can be done after surgical resection of the lesion.

Note: Cirsoid aneurysm is a slow-growing, tortuous swelling over the scalp or face that develops due to abnormal arteriovenous connections in the scalp and extremities. They are usually congenital. However, in some cases, they may be traumatic. The superficial temporal artery is commonly involved in cirsoid aneurysms of the scalp.

Solution to Question 18:

The route of the spread of prostate carcinoma to the lumbar vertebra is by prostatic venous plexus or Batson plexus.

Batson venous plexus is a valveless vertebral venous plexus communicating with prostatic venous plexus.

Carcinoma of the prostate metastasizes to the bone usually by the bloodstream through this Batson plexus. Long bones and the lower lumbar vertebrae are usually involved. Other routes of spread of prostate carcinoma are local spread to the seminal vesicles, bladder neck, trigone, and lymphatic spread.

Carcinoma of the prostate is the most common malignancy in men above 65 years. The clinical features of prostate cancer are bladder outlet obstruction, pelvic pain, hematuria, arthritis,

anemia, pancytopenia, and renal failure. Per rectal examination reveals a hard, nodular, irregularly indurated gland.

Prostate-specific antigen (PSA) is useful in assessing the follow-up. PSA >10 ng/mL is suggestive of cancer and levels >35 ng/mL are diagnostic of advanced prostate cancer. The prostatic biopsy is indicated when there is a raised PSA level or suspicion of prostate cancer.

Localized cancer is treated by radical prostatectomy and radiation therapy, while in advanced cancer palliative and hormone ablation should be carried out.

Solution to Question 19:

The given scenario of a male patient with impotence, bilateral resting pain, and the image showing a gangrenous foot with amputated toes is suggestive of Leriche's syndrome and it indicates that the level of occlusion is at the aortoiliac bifurcation.

Peripheral arterial diseases occur mainly due to atherosclerosis or thromboembolism. Lerich's syndrome is a rare condition in which aortoiliac obstruction causes impotence and claudication (crampy pain) in the thigh or buttocks. Other features of aortoiliac arterial occlusion include absent femoral and distal pulses and a bruit over the aortoiliac region.

In general, features of chronic lower limb arterial occlusion are intermittent claudication, rest pain, ulcers, gangrene, absence of arterial pulses, bruit, and delayed capillary refilling. Rest pain occurs when limb perfusion is so severely compromised that anaerobic respiration occurs even at rest. This causes pain that increases on lying down or elevation of the foot. It reduces when the foot hangs below.

Investigations that are useful include Doppler ultrasound, duplex scanning, ankle-brachial pressure index, and digital subtraction percutaneous angiography (DSA). Through a DSA, endovascular intervention can be performed along with obtaining information on the dynamic arterial flow.

Management is by exercise, smoking cessation, control of diabetes, and weight loss if obese. Drugs such as HMG CoA reductase inhibitors and 75 mg of clopidogrel or aspirin should be given. Transluminal angioplasty and stenting are useful.

Other options:

Options A and D: Femoropopliteal occlusion can lead to claudication in the calf with a palpable femoral pulse but absent distal pulses.

Option B: Iliac obstruction causes thigh and calf claudication with a bruit over the aortoiliac region. The femoral and distal pulses are absent.

Solution to Question 20:

The given scenario of an elderly male patient with sudden-onset scrotal pain and discharge, with the image showing a blackish discoloration of the scrotum, is likely to be Fournier's gangrene.

Fournier's gangrene is an uncommon polymicrobial infection involving the perineal soft tissue, external genitalia, and perianal region. There is subcutaneous tissue inflammation leading to obliterative arteritis and gangrene. It typically follows minor injuries or procedures in the perineal area, urethral dilatation, and injection for hemorrhoids are some of the etiological factors.

Clinically, sudden scrotal pain with associated prostration, pallor, and pyrexia can be seen. Other features such as crepitus and foul-smelling exudate can be present.

Emergency surgical debridement is needed after initial measures like fluid resuscitation and broad-spectrum intravenous antibiotics are undertaken. Urinary diversion and other supportive measures are useful in the management of the condition.

Other options:

Option A: Torsion of testis is a surgical emergency typically seen in the age group of 10-25 years. It is characterized by sudden groin and lower abdominal pain, nausea, and vomiting. Examination findings reveal a swollen, tender scrotum and testis with a twisted tender cord above it and loss of cremasteric reflex.

Option B: Acute epididymo-orchitis is an infection of the epididymis and testis. It is seen in sexually-active young men. In older men, it is seen following a urinary tract infection, secondary to an indwelling catheter. The features include groin ache, fever, swollen and painful epididymis and testis.

Option C: Scrotal carcinoma is a rare carcinoma that is seen in workers exposed to oil, coal products, and aromatic hydrocarbons. It starts as a wart or ulcer in the scrotum and can extend to the testis.

Solution to Question 21:

The fluid shown in the image is ringer lactate or Hartmann's solution.

Ringer lactate, or Hartmann's solution, is a crystalloid made up of sodium 131 mEq, chloride 112 mEq, potassium 5 mEq, bicarbonate (as lactate) 29 mEq, and 2 mEq of calcium. It is devoid of glucose.

It is useful in the following conditions:

• Correction of severe hypovolemia

• Metabolic acidosis

• Postoperative period, burns, diarrhea

• Correction of hypokalemia

• Diabetic ketoacidosis.

It should be avoided in severe shock, liver diseases, vomiting, nasogastric aspiration, and metabolic alkalosis. It cannot be used along with blood transfusion since calcium in the ringer lactate can bind with citrate in the transfused blood leading to clot formation.

Other options:

Option A: Haemaccel is a colloid and contains 145 mmol of sodium, 6.25 mmol of calcium, and 5.1 mmol of potassium. It is mainly used for rapid volume replacement in circulatory shock, burns, and trauma. These release histamine and hence, H2 blocker has to be supplemented. It is also used in heart-lung machines.

Options C and D: Isolyte P and Isolyte M are crystalloids. Isolyte P is commonly used in fluid therapy for pediatrics. One liter contains 50 grams of glucose, 25 mEq of sodium, 22 mEq of potassium, 3 mEq of magnesium, and 23 mEq of acetate. Hyponatremia, renal failure, hyperkalemia, and hypovolemic shock are some of the contraindications.

Isolyte M is used for maintenance. It mainly contains 50 grams of glucose, 40 mEq sodium, 38 mEq chloride, 35 mEq potassium, 15 mEq phosphate, and 20 mEq acetate. It is useful in correcting hypokalemia in patients with good renal function, and acidosis. It is contraindicated in hyponatremia.

Solution to Question 22:

The given image shows a pressure sore with extensive tissue destruction, including damage to the muscle layer and thus it belongs to stage 4.

Pressure sores also referred known as bed sores, pressure ulcers and decubitus ulcers are ulcers that occur due to prolonged pressure on certain sites thereby leading to tissue necrosis. The commonest sites involved are ischium, greater trochanter, sacrum, heel, malleolus (lateral >medial), and occiput. Paraplegic patients, elderly patients, and severely ill patients are at a higher risk of developing pressure sores.

They are staged as:

• Stage 1 - Erythema with no blanching and no breach in the epidermis.

• Stage 2- Partial-thickness skin loss with epidermal and dermal involvement.

• Stage 3- Full-thickness skin loss with intact fascia but the involvement of the subcutaneous tissue.

• Stage 4- Full-thickness skin loss extending to the fascia with massive tissue destruction and involvement of muscle, bone, tendon, or joint.

Management is primarily proper skin care with pressure dispersion cushions, air-fluidized beds, and urinary or fecal diversion if required. Patients who are bed-bound should be turned once in 2 hours. Those in a wheelchair should lift themselves off the seat for 10 seconds every 10 minutes.

Surgery is the last resort and is done after a proper initial debridement and vacuum-assisted closure. Vacuum-assisted closure is a wound closure technique in which negative pressure is applied to the wound which leads to a decrease in edema. This hastens the debridement and formation of granulation tissue. Skin flaps with a muscle are also used for closure.

Solution to Question 23:

Among the marked parts, 4 represents the right subhepatic space. It is the most common site of an intra-abdominal abscess.

The right subhepatic space is a deep space situated in Rutherford Morison's pouch. It is located between the liver above and the transverse colon and the hepatic flexure below. The relations of the right subhepatic space are:

• the right lobe of the liver and diaphragm on the right side

• foramen of Winslow on the left

• duodenum below

• liver and the gall bladder in the front

• right kidney and the diaphragm behind.

Abscesses here usually arise from appendicitis, cholecystitis, perforated duodenal ulcer, or as sequelae of an upper abdominal surgery.

Intraperitoneal abscesses may develop following intraperitoneal sepsis in specific sites in the abdomen (intra-abdominal abscess) or pelvis (pelvic abscess). The intra-abdominal abscess may occur in subphrenic spaces, or the subhepatic spaces, on either the right or left side.



An intra-abdominal abscess is a localized collection of infected fluid within the abdominal cavity due to failure of the host's intra-abdominal defense mechanisms. Patients usually present with acute abdominal pain, fever, rigors, tachycardia, tachypnea, and leukocytosis. Other features such as nausea, anorexia, emesis, constipation, and diarrhea can occur.

CT scan with IV contrast is the imaging method of choice for an intra-abdominal abscess. Ultrasonography can also be useful.

Initial management is by early intravenous antibiotic therapy, drainage of the abscess, and fluid resuscitation with crystalloids. Vasopressors, inotropes, and packed red blood cell transfusion are given when needed.

Other measures include ultrasound or CT-guided percutaneous drainage (non-surgical) and operative drainage. Resection and debridement can also be done. It can be done through a

minimally invasive technique or laparotomy.

Solution to Question 24:

The given clinical scenario of recurrent pain and discharge due to lesions around the anus, shown in the image is likely to be a fistula-in-ano.

A fistula-in-ano is an abnormal connection between the anal canal and the perianal skin, lined with granulation tissue. It may follow the drainage of a perianal abscess or may be seen in association with diseases like tuberculosis or Crohn's disease.

Clinical features are perianal pain, swelling, and intermittent purulent discharge, which might be blood-stained. On examination, one or more external openings may be visible.

Goodsall's rule can be used as a guide in determining the location of the internal opening of the fistula. Fistulas with an external opening anteriorly connect to the internal opening by a short, radial tract. Fistulas with an external opening posteriorly track in a curvilinear fashion to the posterior midline.

However, exceptions to this rule often occur. Fistulas with an anterior external opening greater than 3 cm from the anal margin usually track to the posterior midline.



The investigations done for fistula-in-ano can include a fistulogram, endoanal ultrasound with hydrogen peroxide, MRI, and computed tomography (CT) scan. An MRI is the gold standard investigation for fistula imaging. It helps us to identify the secondary extensions of the track, which when missed may cause persistence of the fistula.

Surgical management includes fistulotomy, fistulectomy, ligation of the intersphincteric fistula tract (LIFT), setons, or advancement flaps.

Solution to Question 25:

Ambulatory patients that are stable are likely to survive even if treatment is delayed. Their medical need is not urgent and they come under green according to triage.

Triage is a method employed to identify the patients in whom treatment at the earliest produces the maximum benefit to the patient. It is used in the management of mass casualties.

Triage categories are based on the vital signs, patient's ability to walk, mental status, and presence or absence of ventilation/capillary perfusion. It is as follows:

Priority	Color	Medical need	Clinical status
First	Red	Immediate	Critical but can survive with early treatment
Second	Yellow	Urgent	Critical but can survive when treatment is given within hours
Third	Green	Non-urgent	Patients arestableand cansurvivewithdelayed treatment
Last	Black	Unsalvageable	Not breathing, no pulse, severely injured and medical care is unlikely to help.

Surgery NEET 2023

Question 1:

A 55-year-old male presented with verrucous carcinoma around the glans of the penis. Examination reveals that the inguinal lymph nodes are not enlarged. What is the appropriate management for this patient?

a) Total penectomy.

b) CO_2 laser excision

c) Topical 5-fluorouracil

d) Partial penectomy.

Question 2:

Identify the procedure being performed in the image given below.



a) Intraosseous cannula for pain relief

b) Bone marrow aspiration

c) Intraosseous route access for giving IV fluids

d) Intraosseous abscess drainage

Question 3:

A diabetic patient presents with sudden-onset perineal pain. On examination, foul-smelling discharge, and necrotic tissue as seen in the image is noted. Which of the following is true about the given condition?

a) Anti-gas gangrene serum indicated for all cases

b) Urinary diversion is the next step

c) Bilateral orchidectomy must be done

d) Mixed aerobic and anaerobic infection

Question 4:

Which of the following is the most common complication following ligation of the first vessel during abdominoperineal resection for rectal carcinoma?

a) Parasympathetic - bladder dysfunction and retrograde ejaculation.

b) Sympathetic - bladder dysfunction and impotence.

c) Sympathetic - retrograde ejaculation and bladder dysfunction.

d) Sympathetic - Impotence and loss of cutaneous sensation in perineal region

Question 5:

A patient presents to the clinic with the lesion given in the image. He had a traumatic injury to the chest one year ago. What is the most likely diagnosis?

a) Hemangioma

b) Hypertrophic scar

c) Keloid

d) Neurofibroma

Question 6:

A 45-year-old male underwent bilateral laparoscopic hernia repair for inguinal hernia. Postoperatively, he complained of pain in the right thigh. Which of the following nerve entrapment leads to this symptom?

a) Ilioinguinal nerve

b) Iliohypogastric nerve

c) Femoral nerve

d) Lateral cutaneous nerve of thigh

Question 7:

A patient was brought to the ER following a road traffic accident. On examination, the patient opens his eyes to a painful stimulus, speaks inappropriately, and withdraws his limbs to a painful stimulus. What is his GCS score?

a) E2V2M3

b) E3V3M3

c) E2V3M4

d) E3V2M2

Question 8:

A patient presented with fever and abdominal pain with jaundice. Investigations showed the given findings. Which of the following statements is true regarding this condition?

a) Surgery is mandatory

b) Fine-needle aspiration cytology is diagnostic

c) Angioembolization is the treatment of choice

d) Echinococcus species is involved in etiology

Question 9:

A 40-year-old male patient presents with a swelling that has been slowly growing in the past 2 years. On examination, it was variable in consistency and fully mobile. What is the most likely diagnosis?

a) Dermoid cyst

b) Parotid tumor

c) Sebaceous cyst

d) Cervical lymph node

Question 10:

A patient who is a known case of alcohol dependence syndrome presents with sudden and unintentional weight loss. What is the most likely diagnosis?

a) Hepatic adenoma

b) Cholangiocarcinoma

c) Hepatocellular carcinoma

d) Alcoholic hepatitis

Question 11:

A patient presented with features of chronic pancreatitis with recurrent attacks and has a 10 mm dilatation of the pancreatic duct with intraductal calculi present. Which of the following is the surgery of choice?

a) Pancreaticoduodenectomy

b) Longitudinal pancreaticojejunostomy

c) ERCP and sphincterotomy

d) Coring of pancreas head

Question 12:

A woman presented with a BIRADS-5 breast lesion. Which of the following is a good prognostic factor for this lesion?

a) BRCA-1 positive

b) p53 positive

c) ER positive

d) High Ki-67

Question 13:

A patient presents with sudden onset of chest pain shooting to the neck and interscapular region. X-ray shows widened mediastinum. BP is 110/90 mmHg in the right upper limb and 160/100 mmHg in the left upper limb. What is the most likely diagnosis?

a) Acute coronary syndrome

b) Acute pulmonary embolism

c) Acute aortic dissection

d) Esophageal rupture

Question 14:

A patient presented with right hypochondriac pain. He had an episode of diarrhea 1 week prior. CT scan of the abdomen reveals a liver abscess of around 25 ccs. What is the next step in management?

a) PAIR

b) Surgery

c) Medical therapy

d) Percutaneous drainage

Question 15:

A 50-year-old patient with a history of trauma one week back presents with confusion and ataxia. He is on treatment for alcohol dependence and his last intake was 2 weeks back. CT scan reveals the following finding. What is the diagnosis?

a) Extradural hemorrhage

b) Subdural hemorrhage

c) Intra-parenchymal hemorrhage

d) Subarachnoid hemorrhage

Question 16:

Which of the following is false about the given condition?

a) Repeated radical surgery every 24 hours may be required to adequately manage this condition

b) This is a case of Meleney's gangrene

c) The condition is caused by beta hemolytic streptococcus and may sometimes be polymicrobial

d) Hyperbaric oxygen has no role in the treatment of this condition

Question 17:

A patient presents with a swelling in the groin region and it is diagnosed as a direct inguinal hernia. Weakness in which of the following structures is most likely to cause this condition?

a) Pectineal ligament

b) Conjoint tendon

c) Reflected part of inguinal ligament

d) Lacunar ligament

Question 18:

A patient presents with sudden onset of hematemesis. He has been taking aspirin for his arthritis and drinks alcohol occasionally. He gives a history of occasional abdominal pain. On examination, there is no abdominal mass or tenderness. What is the likely diagnosis?

a) Mallory-Weiss tear

b) Esophagitis

c) Peptic ulcer

d) Esophageal varices

Question 19:

A construction worker came to the OPD complaining of white fingers. He has been working in the cement and concrete industry and has been working on heavy machinery drills, wood, and furniture polishing for 20 years. What is the most likely etiology for this condition?

a) Candidal infection of the fingers due to continuous exposure to water

b) Exposure to thinners and paints

c) Continuous exposure to cement and concrete

d) Continuous exposure to drills and machines

Question 20:

A 5-year-old child was admitted to the hospital for prolapsing rectal mass and painless rectal bleeding. Histopathological examination reveals enlarged and inflamed glands filled with mucin. What is the likely diagnosis?

a) Adenoma

b) Choristoma

c) Hamartoma

d) Carcinoma

Question 21:

Identify the given condition.

a) Venous ulcer

b) Malignant ulcer

c) Arterial ulcer

d) Trophic ulcer

Question 22:

A patient with varicose veins comes with complaints of eczema over the same region. What is the stage of the clinical classification according to the CEAP guidelines?



a) C2

b) C3

c) C4a

d) C4b

Question 23:

A patient comes to the casualty with a severe headache. His BP was found to be 160/100 mmHg. CT scan revealed a subarachnoid hemorrhage. What is the next best step in the management of this patient?

a) Nimodipine

b) Angiography

c) Surgery

d) Fibrinolytic therapy

Question 24:

A 20-year-old female patient consumed a cleaning product containing 90% sodium hydroxide. She is having complete dysphagia. What is the best step in management?

a) Esophagojejunostomy

b) Stent placement

c) Feeding jejunostomy

d) Gastrojejunostomy

Question 25:

A 23-year-old male patient presents with midline swelling in the neck. The swelling moves with deglutition and protrusion of the tongue. What is the likely diagnosis?

a) Brachial cyst

b) Thyroglossal cyst

c) Plunging ranula

d) Dermoid cyst

Question 26:

A 54-year-old woman presents with cervical cancer stage 2A. It is decided to give neoadjuvant chemotherapy. What does neoadjuvant chemotherapy mean?

a) Chemotherapy given along with radiation

b) Chemotherapy given during surgery

c) Chemotherapy given before radical surgery to reduce bulk of tumor

d) Chemotherapy given after radical surgery for micrometastases

Question 27:

A young patient is admitted with a history of fever for 5 days. BP is 90/60 mmHg, PR-120 bpm, RR-24 breaths/min, GCS score-10; laboratory tests show leucocytosis with neutrophilia and serum creatinine level of 2.6 mg/dL. An intern doctor wants to calculate the qSOFA score to predict prognosis and hospitalization duration. Which of the following comprises the qSOFA score?

a) BP, RR and CBC

b) Creatinine, PR, BP

c) RR, Body temperature, PR

d) SBP, RR, GCS score

Answer Key

Question No.	Correct Option
1	d
2	c
3	d
4	c
5	c
6	d
7	c
8	d
9	b
10	c
11	b
12	c
13	c
14	c
15	b
16	d
17	b
18	c
19	d
20	c
21	d
22	c
23	b
24	c
25	b
26	c
27	d

Detailed Explanations

Solution to Question 1:

The most appropriate management for this patient with verrucous carcinoma around the glans of the penis is partial penectomy.

Carcinoma of the penis is commonly a squamous cell carcinoma. It arises from the skin of the glans penis or the prepuce. The lesion can be flat or warty in appearance.

• T1- confined to the skin

• T2- invades the corpus spongiosum or cavernosum.

• T3- invades the urethra

• T4- invades adjacent structures.

The first lymph node to be involved is the superficial inguinal lymph node and thus, the sentinel lymph node biopsy (SLNB) is helpful for evaluating the spread of penile cancer.

Tumors of low-grade and low-stage are managed by Mohs' surgery. Tumors involving the glans penis require glansectomy, and in advanced cases require partial penectomy.

Other options:

Option A: In penile carcinoma, total amputation of the penis is done if the stump is < 2cm after resection of the tumor.

Options B, and C: Topical 5-fluorouracil cream and CO_2 laser ablation are preferred in case of penile intraepithelial neoplasia.



Solution to Question 2:

The procedure being performed in the above image is intraosseous route access for transfusing IV fluids in the anterior proximal tibia.

The intraosseous route is an alternative to the intravenous (IV) route in the transfusion of fluids in shock. The marrow of long bones has a rich network of vessels that drain into a central venous canal, emissary veins, and ultimately the central circulation. So, the bone marrow functions as a non-collapsible venous access route.

Intraosseous cannulation is recommended for patients in whom IV access proves difficult or unattainable. (e.g., in cardiopulmonary arrest). In such a case, if venous access is not available within 1 min, an intraosseous needle should be placed in the anterior proximal tibia. During a resuscitation procedure, all anesthetic drugs and fluids can be administered through intraosseous cannulae.

Insertion sites for intraosseous infusion:

• In the proximal tibia, 1-2 cm anteromedial from the tibial tuberosity

• Distal tibia at the junction of the medial malleolus and the shaft of the tibia

• Distal one-third of the femur

The response time to drug administration in intraosseous infusion is close to the IV route. Fluids should be given as boluses using a 50 mL syringe.

The image below shows the procedure of intraosseous cannulation.



Solution to Question 3:

The given condition is likely to be Fournier gangrene. It is a mixed flora aerobic and anaerobic infection.

Fournier's gangrene is a polymicrobial infection and a type of necrotizing fasciitis of the perineum, external genitalia, and perianal region. Risk factors include immunocompromised states such as HIV, alcoholism, and diabetes. The infection may be precipitated by minor

procedures in the perineal area, such as a bruise, scratch, urethral dilatation, injection of hemorrhoids, or opening of a periurethral abscess. The infection spreads along the dartos, Scarpa, and Colles fascia.

The sequence of events that lead to the development of gangrene are as follows: A mixed infection of the aerobic and anaerobic bacteria → fulminating inflammation of subcutaneous tissue → obliterative arteritis of the arterioles that supply the scrotal skin → gangrene.

Clinical features of Fournier gangrene are scrotal pain, fever, prostration, and pallor. Cellulitis spreads rapidly and may be associated with eschars, necrosis, flaking skin, and crepitus. The testis is usually not affected.

Diagnosis is majorly clinical and very rarely made by laboratory or imaging findings.

The steps in the management of this condition:

• IV fluids

• IV antibiotics

• Urgent surgical excision followed by skin grafting in later stages. Wide surgical excision is often necessary.

When the infection has been controlled, a vacuum-assisted dressing may be helpful. Skin grafting is essential for a patient surviving an acute episode. Despite the best therapy, mortality rates are around 50%.

Other options:

Option A: Anti-gas gangrene serum is a treatment for gas gangrene that is of historical importance and is no longer used. Treatment of gas gangrene is now done with antibiotics.

Option B: While urinary diversion and other supportive measures are useful in the management of the condition, it is not the next best step.

Option C: The blood supply of the testes is separate from that of the overlying skin, and hence, testes can be preserved during debridement.

Solution to Question 4:

The first vessel that is usually ligated during abdominoperineal resection (APR) is the inferior mesenteric artery (IMA). A high inferior mesenteric artery ligation during abdominoperineal resection can injure the superior hypogastric nerve, resulting in sympathetic dysfunction and leading to retrograde ejaculation and bladder dysfunction.

Abdominoperineal resection is the complete excision of the rectum, anal canal, and sphincter apparatus. This is done with the creation of a permanent colostomy. This is commonly preferred in elderly patients with distal rectal cancer with poor sphincter function. In these patients, the end colostomy offers a better quality of life.

Complications:

• Higher recurrence rates compared to low anterior resection.

• Anastomotic leaks or dehiscences

• Necrosis of the transposed colon

• Bleeding

• Strictures

The sympathetic innervation of the rectum is from L1, L2, and L3, which together form the superior hypogastric plexus. They divide into left and right hypogastric nerves at the level of the sacral promontory supplying the rectum and the genitourinary system. A high IMA ligation during surgery can result in sympathetic dysfunction causing retrograde ejaculation.

Injury to the pelvic plexus and nervi erigentes, during the division along the pelvic side wall, can cause erectile dysfunction, impotence, and atonic bladder. Injury to the periprostatic plexus can lead to sexual and bladder dysfunction.

Solution to Question 5:

The image showing a large area of scar tissue is suggestive of a keloid.

A keloid refers to excess accumulation of collagen at the scar site with growth of the scar tissue beyond the boundaries of the original wound. It does not regress. The patient's history of traumatic injury to the chest one year ago, followed by the development of an expanding scar, is suggestive of a keloid. The keloid is often glossy in appearance and rubbery in texture. They are composed of thick bundles of collagen.

Treatment of keloids usually involves intralesional injections of triamcinolone. Multiple doses are required at intervals over several weeks and may result in flattening but rarely does the wound narrow.

Option A: A hemangioma is a vascular tumor that arises from the endothelium. They typically present as red or pinkish macules or raised lesions.

Option B: Hypertrophic scars, unlike keloids, remain within the boundaries of the original wound and tend to regress over time. Since this scar has failed to regress over 1 year and has grown beyond the original wound margins, it is more likely to be a keloid.

Option D: Neurofibromas are benign tumors arising from Schwann cells. They usually present as soft, flesh-colored nodules, often in association with neurofibromatosis type 1, a genetic disorder. The patient's history of chest trauma and the appearance of the lesion make neurofibroma an unlikely diagnosis in this case.

Solution to Question 6:

Entrapment of the lateral cutaneous nerve of the thigh is the most likely cause of pain in this patient with history of laparoscopic hernia repair.

The laparoscopic approach is preferred for bilateral or recurrent inguinal hernias. Transabdominal preperitoneal (TAPP) repair and total extraperitoneal (TEP) repair are the methods commonly used.

The nerves commonly injured in laparoscopic repair are the genitofemoral and lateral femoral cutaneous nerves of the thigh. Patients present with acute neuropathic pain along the dermatomal distribution. Injury to the lateral femoral cutaneous nerve results in meralgia paresthetica.

Treatment consists of rest, ice, NSAIDs, physical therapy, and possible local corticosteroid and anesthetic injection. Drugs like gabapentin can be used.

In laparoscopic TAPP repair, to reduce the chances of nerve injury during mesh placement, tacks are ideally placed above the iliopubic tract to avoid nerve injury.

Other options:

Option A: Ilioinguinal nerve may be commonly injured during open hernia repair. It leads to sensory disturbances and pain over the skin of the genitalia and upper medial thigh.

Option B: Iliohypogastric nerve can be injured while making an incision in the right iliac fossa during hernia surgery.

Option C: Femoral nerve can be compressed by retroperitoneal tumors or hemorrhage in patients on anticoagulants or bleeding diathesis. It presents as wasting and weakness of the quadriceps femoris muscle making the patient difficult to walk. Pain and paraesthesia are seen along the region of the distribution of the nerve.

Solution to Question 7:

The GCS score of this patient: E2V3M4

- He opens his eyes to a painful stimulus - E2
- He produces inappropriate words only - V3
- He withdraws his limbs - M4

Solution to Question 8:

The given CT image shows a cyst in the liver with serpentine linear membranes seen as wavy membranes within the cyst is suggestive of hydatid disease.

Hydatid disease or echinococcosis is a zoonotic infection caused by the Echinococcus species of flatworms. Its life cycle consists of dogs as definitive hosts and humans as abberant intermediate hosts. Primary hydatid cysts most commonly occur in the liver. Most patients are asymptomatic but when symptomatic, the patient presents with hepatomegaly, pain, and obstructive jaundice.

The first/initial imaging test is USG, and a CT scan can be used to diagnose the condition. CT scan shows the presence of a space-occupying lesion with a smooth outline and septa. CT scan has an accuracy of 98% in the detection of hydatid cysts and high sensitivity to the detection of daughter cysts and extra-hepatic cysts. On CT imaging, the serpent sign is seen as wavy membranes within the cyst.

The USG of an active hydatid cyst shows multiple septations and a rosette or honeycomb appearance:



The ball of wool sign is seen on USG when the contents settle in the dependent portion of the cyst, as seen below:



On USG, the water-lily sign is seen in hydatid infections when there is detachment of the endocyst membrane, which results in floating membranes within the pericyst that mimic the appearance of a water lily.



Other options:

Option A: While surgical treatment of hydatid cyst is useful and effective, alternatives like percutaneous aspiration, injection, re-aspiration (PAIR) are equally effective and are currently used.

Option B: IgG ELISA and indirect hemagglutinin assay are used for the diagnosis of hydatid cyst.

Option C: Angioembolization has no role in the treatment of hydatid cyst.

Solution to Question 9:

The given clinical scenario and image depict a likely case of parotid tumor, likely to be a pleomorphic adenoma.

Pleomorphic adenoma is the most common benign salivary gland tumor. The most commonly affected gland is the parotid gland. The superficial lobe is most commonly affected. It is also called as mixed tumor since it has both epithelial and mesenchymal components. It most commonly affects females around 40 years of age.

The common presenting feature is a painless swelling below, in front, or behind the ear. The involvement of the deep lobe leads to dumbbell tumor – swelling in both the neck and oral cavity.

Specific features:

• Slowly growing benign tumor

• Unicentric in origin

• Potentially malignant

• Possibility of protrusion of fringes of the tumor mass through the capsule

• Radioresistant

• Usually affects adults, females > males

• Painless swelling

• Well-defined margins

• Lobulated, firm, elastic-to-hard consistency

• Free from deeper structures

• No facial nerve involvement

• The deep lobe pushes tonsils and faucial pillars

Diagnosis is made by FNAC, which is considered the investigation of choice. Open biopsy contraindicated due to chances of injury to the facial nerve and seedling of the tumor

Surgery is the mainstay of treatment - Superficial parotidectomy (Patey's operation). Radiotherapy is considered in high-grade malignancies demonstrating extra-glandular disease, perineural invasion, direct invasion of surrounding tissue, or regional metastases.

Pathology:

Gross features:

• The tumors are rounded, well-demarcated, measuring <6 cm in greatest dimension.

• They are usually encapsulated, except in some areas such as the palate.

• The cut surface is gray-white with myxoid and blue translucent areas of the chondroid (cartilage-like) stroma.



Microscopic features:

• The epithelial elements resemble ductal cells or myoepithelial cells. They are arranged as ducts, acini, irregular tubules, strands, or sheets of cells.

• These epithelial elements are dispersed within a background of loose myxoid and hyaline tissue containing islands of cartilage and, rarely, foci of bone.

The image below shows pleomorphic adenoma of the salivary gland consisting of mixed epithelial (left) and mesenchymal cell components (right).



Dermoid cysts are slow-growing benign lesions. They are mostly seen on the scalp and skull. It is made of hair, teeth, cartilage, and skin glands. Other common sites are the forehead, postauricular, and lateral part of the eyebrow.

A sebaceous cyst is derived from hair follicle infundibulum or traumatic inclusion. The most common site of a sebaceous cyst is the postauricular sulcus or below and behind the ear lobule. Malignancy is rare.

Clinical features of a sebaceous cyst include:

• Benign swelling that is mobile, fluctuant, and fixed to the skin. It presents with a central punctum in about 70% of cases.

• It is generally an asymptomatic, non-tender swelling. It can be painful and red if inflamed.

• It may ooze out a foul-smelling, cheese-like discharge.

Solution to Question 10:

The most likely diagnosis in this patient with a history of alcohol dependence, sudden unintentional weight loss, and raised AFP is hepatocellular carcinoma (HCC).

HCC is the most common cause of primary liver cancer. It is more commonly seen in males, and the peak age group is between 65-70 years. The main risk factor for HCC is cirrhosis and associated chronic liver damage due to chronic infection by hepatitis B or C virus (HBV or HCV), alcohol abuse, metabolic syndrome, or hemochromatosis. The incidence of HCC is increasing worldwide. This is contributed by the increase in HCV and HBV infections, along with lifestyle disorders like obesity. Obesity is associated with non-alcoholic fatty liver disease (NAFLD). This progresses on to non-alcoholic steatohepatitis (NASH) causing cirrhosis and hepatocellular carcinoma. Thus, NASH and NAFLD are risk factors for HCC.

Regular surveillance, including liver imaging and monitoring of alpha-fetoprotein levels, is recommended for high-risk populations, such as those with liver cirrhosis, chronic hepatitis B or C infection, or significant alcohol consumption. Early detection of HCC can improve treatment

outcomes and overall prognosis.

Very early HCC is treated by ablation. Early HCC with a single nodule is resected, but when there are multiple nodules over 3 cm, ablation or transplantation are considered. Intermediate-stage HCC is treated by chemoembolization. Transarterial chemoembolization (TACE) is used for multinodular lesions.

Other Options:

Option A: Hepatic adenomas are benign liver tumors that are often associated with oral contraceptive use and anabolic steroid use. The patient's history of alcohol dependence and high alpha-fetoprotein levels are not consistent with hepatic adenoma. Additionally, hepatic adenomas are usually asymptomatic and do not present with sudden unintentional weight loss.

Option B: Cholangiocarcinoma is a cancer that arises from the bile ducts within the liver. While patients with alcohol dependence may have an increased risk of cholangiocarcinoma, the elevated alpha-fetoprotein levels are not typical for this type of cancer. Cholangiocarcinomas usually present with jaundice, pruritus, and abdominal pain, but sudden weight loss is less characteristic.

Option D: Alcoholic hepatitis is an inflammation of the liver caused by excessive alcohol consumption. It usually presents with jaundice, fever, and abdominal pain. While the patient's history of alcohol dependence might suggest alcoholic hepatitis, elevated alpha-fetoprotein levels and unintentional weight loss are not typical for this condition. Moreover, the AST/ALT ratio is usually greater than 2 in alcoholic hepatitis, but it is 0.5 in this case.

Solution to Question 11:

In this case of chronic pancreatitis with recurrent attacks, 10 mm dilatation of the pancreatic duct, and intraductal calculi, longitudinal pancreaticojejunostomy (also known as Puestow procedure or lateral pancreaticojejunostomy) is the surgery of choice. This procedure involves opening the pancreatic duct along its length and connecting it to a loop of the jejunum, which helps to drain the obstructed pancreatic duct and alleviate pain.

Patients with chronic pancreatitis mainly experience pain due to ductal hypertension or parenchymal inflammation. Acute exacerbations in duct pressure lead to recurrent acute attacks. The pancreatic duct dilation observed on pancreatography for chronic pancreatitis is classically described as a chain of lakes, which reflects the presence of multiple dilations and stenoses. Response to treatment has been found to be dependent upon the degree of preoperative pancreatic fibrosis.

Longitudinal pancreaticojejunostomy, also known as the Puestow procedure or lateral pancreaticojejunostomy, is a surgical treatment primarily used for chronic pancreatitis with a dilated main pancreatic duct and intraductal calculi.

In longitudinal pancreaticojejunostomy, a longitudinal incision is made along the length of the main pancreatic duct, typically from the tail to the head of the pancreas. The dilated pancreatic duct is then opened, and any intraductal calculi or obstructions are removed. The opened duct is then connected to a loop of the jejunum to create a side-to-side anastomosis. This allows for the drainage of pancreatic secretions directly into the jejunum, bypassing any narrowed or obstructed areas of the duct.

This procedure can effectively relieve pain and improve the quality of life for patients with chronic pancreatitis. However, it is important to note that it may not reverse the underlying pathology of chronic pancreatitis or prevent the progression of the disease. It is also not suitable for all patients with chronic pancreatitis, particularly those with a non-dilated pancreatic duct or isolated head involvement.

Other Options:

Pancreaticoduodenectomy: This procedure, also known as Whipple procedure, involves removing the head of the pancreas, duodenum, gallbladder, and part of the bile duct. It is typically performed for pancreatic cancer or tumors in the head of the pancreas, not for chronic pancreatitis with ductal obstruction and intraductal calculi.

ERCP and sphincterotomy: Endoscopic retrograde cholangiopancreatography (ERCP) with sphincterotomy is a procedure that involves opening the sphincter of Oddi to facilitate the drainage of bile and pancreatic secretions. Although it may provide temporary relief, it is not a definitive treatment for chronic pancreatitis with large ductal dilatation and intraductal calculi. ERCP with sphincterotomy may be useful in some cases of chronic pancreatitis for the removal of small pancreatic duct stones, but for large ductal dilatation and intraductal calculi, longitudinal pancreaticojejunostomy is more appropriate.

Coring of pancreas head: Coring of the pancreas head, also known as pancreatic head resection or Beger procedure, is a surgical technique used to treat chronic pancreatitis, particularly when the inflammation and fibrosis predominantly affect the head of the pancreas. The procedure involves removing the inflamed and fibrotic tissue from the head of the pancreas while preserving the duodenum, bile duct, and the main pancreatic duct. Longitudinal pancreaticojejunostomy is a more targeted and effective treatment for this patient's condition.

Solution to Question 12:

ER positivity is a good prognostic factor in breast cancer.

The ER and PR are two steroid hormone receptors present in breast cancer. The expression of these receptors indicates a favorable prognosis and response to hormonal therapy.

Other options:

Option A: The mutations in BRCA1 gene account for up to 40% of familial breast cancers. These tumors are high-grade, hormone receptor–negative, and aneuploid and do not have a good prognosis.

Option B: p53 is a tumor suppressor gene, which on mutation acts as an oncogene. p53 positivity is associated with highly malignant breast cancers and has poor survival.

Option D: Ki-67 is used to access the proliferative index (activity) of the tumor. High Ki-67 indicates a poor prognosis.

Solution to Question 13:

Sudden-onset severe chest pain, with a wide discrepancy in blood pressure between the two limbs, and a widened mediastinum on x-ray point towards a diagnosis of aortic dissection.

It is initiated by an intimal tear, which allows blood to penetrate into the medial layer, producing a cleavage plane (false lumen) between the inner two-thirds and outer one-third of the media. The true lumen is separated from the false lumen by an intimomedial flap.

Classification of aortic dissection:

Acute aortic dissection can be classified by the original DeBakey system, which is based on the location of the proximal tear and distal extent of the dissection flap, or the Stanford system, which is based only on the location of the proximal tear.

	DeBakey	Stanford
Site of dissection		
Both ascending and descending aorta	Type I	Type A
Only ascending aorta and arch of the aorta	Type II	Type A
Descending the aorta distal to the subclavian artery	Type IIIaLimited to the thoracic aorta	Type B
Type IIIbExtends to the abdominal aorta		



Clinically, patients present with tearing chest/back pain associated with diaphoresis, syncope, weakness, and dyspnea or neurological manifestations.

On examination, unequal pulses or blood pressure in the extremities may be seen due to changes of flow in the true and false lumens. There can be hypertension or hypotension, loss of pulses, findings of aortic regurgitation, and pulmonary edema.

Imaging:

Transesophageal echocardiography (TEE) is the investigation of choice in a hemodynamically unstable patient. In hemodynamically stable patients, the diagnosis is established with a contrast-enhanced CT scan. The characteristic diagnostic finding is a double-lumen aorta. The image below shows aortic dissection in the thoracic aorta with the true lumen, as well as the false lumen. The true lumen is often the smaller of the two because of being compressed by the false lumen.

Acute coronary syndrome and acute pulmonary embolism may both present with sudden onset chest pain but widened mediastinum and blood pressure discrepancy make aortic dissection more likely. (Options A and B)

Both sudden onset chest pain and widened mediastinum may be seen in esophageal rupture but blood pressure discrepancy between the upper limbs makes aortic dissection more likely. (Option D)

Chest x-ray showing widened mediastinum in a patient with aortic dissection



The illustration given below shows the flowchart for the management algorithm for aortic dissection:



Solution to Question 14:

This patient with an abscess in the liver with a history of diarrhea most likely has an amebic liver abscess. The initial treatment of choice for an amebic liver abscess is medical therapy with metronidazole.

Ingestion of Entamoeba histolytica cysts causes intestinal diseases like diarrhea. Following this, the trophozoites reach the liver mostly through the portal venous system and cause amebic liver abscess. The common presenting features are right upper quadrant pain, tenderness, fever, hepatomegaly, and diarrhea.

CT abdomen is helpful in differentiating amebic from a pyogenic abscess. A rim enhancement is noted in case of a pyogenic abscess.

The mainstay of treatment for amebic abscesses is metronidazole, which is curative in more than 90% of patients. Luminal agents such as iodoquinol, paromomycin, or diloxanide furoate can be administered to treat the carrier state.

Aspiration of abscess is recommended in case of

• Diagnostic uncertainty

• Failure to respond to metronidazole therapy

• Abscesses larger than 5 cm in diameter and in the left liver, since they are thought to carry a high risk of rupture

Solution to Question 15:

The given clinical scenario and the plain CT scan showing a concavo-convex-shaped hematoma on the left side with a midline shift to the right is suggestive of subdural hematoma (SDH).

An acute SDH is the result of the accumulation of blood between the arachnoid membrane and the dura. It is usually due to the rupture of bridging veins running from the cerebral cortex to the dural sinuses. The level of consciousness is usually impaired at presentation and may deteriorate further as the hematoma expands. Elderly and alcoholic patients are at a higher risk of acute SDH formation after head trauma due to brain atrophy. On CT scan, acute SDH appears as a hyperdense, crescent-shaped (concavo-convex) lesion.

Open craniotomy for the evacuation of acute SDH is indicated for the following:

• Thickness >1 cm

• Midline shift >5 mm

• Glasgow Coma Scale (GCS) score drops by two or more points from the time of injury to hospitalization.

Chronic subdural hematoma results from a hematoma that expands over days or weeks by osmosis, ultimately producing symptoms of raised ICP or focal deficits. On a CT scan, diffuse hypodensity overlying the brain surface would be seen.

Solution to Question 16:

The given image is likely that of Meleney's gangrene. Hyperbaric oxygen has been shown to reduce mortality in patients with Meleney's gangrene and therefore plays a significant role in the management of this condition.

Necrotizing fasciitis of the abdominal wall is known as Meleney's gangrene. It presents with extensive necrosis of the abdominal wall, thrombosis of the microvasculature, and an acute inflammatory infiltrate. Bullae develop on the abdominal wall, which progress to overt cutaneous gangrene. The affected area is often painful, erythematous, and edematous, with possible systemic signs of infection such as fever and malaise. Early recognition and aggressive treatment are essential for a favorable outcome.

Management includes prompt surgical debridement of the necrotic tissue, broad-spectrum antibiotics to cover the polymicrobial nature of the infection, and supportive measures such as fluid resuscitation and correction of electrolyte imbalances. In some cases, hyperbaric oxygen therapy may be utilized as an adjunctive treatment to improve tissue oxygenation and enhance the healing process.

Other options:

Option A: Surgical excision of necrotic tissue is often required daily as the infection tends to spread beyond the borders of the debridement wound.

Option C: Meleney's gangrene is caused by beta-hemolytic streptococci or Staphylococcus aureus and may also be polymicrobial.

Solution to Question 17:

A direct inguinal hernia is caused due to weakness in the conjoint tendon.

The innermost layers of the lateral abdominal wall contain two muscles, namely internal oblique and transverse abdominis. These muscle fibers arch over the deep inguinal ring from lateral to medial, fuse, and become tendinous forming the conjoint tendon. The area below this arch lacks muscle and contains external oblique aponeurosis and fascia transversalis, making it weak.

Direct inguinal hernia is common in elderly patients. It is seen medial to the inferior epigastric vessels, in Hesselbach's triangle. This area lacks muscle and contains only fascia.

Other options:

Option A: Pectineal ligament also known as Cooper's ligament is a strong fibrous band that extends laterally along the pectineal line from the pectineal attachment of the lacunar ligament.

Option C: The inguinal ligament is external oblique aponeurosis that extends between the anterior superior iliac spine and the pubic tubercle. On the point of attachment to the pubic tubercle, some fibers pass upwards behind the superficial inguinal ring and external oblique aponeurosis to join the rectus sheath and linea alba. These form the reflected part of the inguinal ligament.

Option D: Lacunar ligament is formed from the fibers of the medial end of the inguinal ligament and the fibers from the fascia lata of the thigh. It forms the medial wall of the femoral canal.



Solution to Question 18:

This patient with a history of aspirin use, now presenting with abdominal pain and a sudden onset of hematemesis most likely has a bleeding peptic ulcer.

Peptic ulcer disease (PUD) accounts for around 2/3rd of upper gastrointestinal bleeding. The major causative factors are Helicobacter pylori and NSAIDs. Prostaglandins protect gastric and duodenal mucosa from acid and pepsin. They increase mucin and bicarbonate secretion and increase mucosal blood flow. They also promote epithelial cell proliferation. NSAIDs like aspirin disrupt these naturally protective mechanisms and increase the risk of peptic ulcer formation in the stomach and the duodenum.

They present with abdominal pain, bleeding, or signs of peritonitis due to perforation. Significant hemorrhage occurs when duodenal or gastric ulcers penetrate branches of the gastroduodenal or left gastric arteries, respectively.

Other options:

Option A: Mallory-Weiss tears are partial-thickness tears of the mucosa and submucosa occurring near the gastroesophageal junction. They occur in patients with a history of repeated emesis and are classical in alcoholic patients after a period of intense retching and vomiting following binge drinking.

Option B: Esophagitis commonly occurs due to repeated exposure of mucosa to the acidic gastric secretions in gastroesophageal reflux disease. It results in chronic blood loss and commonly presents as anemia or guaiac-positive stools. Ulcerations of superficial mucosa due to esophagitis generally do not bleed acutely.

Option D: Esophageal varices commonly occur due to portal hypertension in patients with cirrhosis. A history of liver disease or the stigmata of portal hypertension and cirrhosis would suggest variceal bleeding.

Solution to Question 19:

The clinical vignette suggests the diagnosis of Raynaud's syndrome due to chronic exposure to drills and machines (vibrating tools).

Raynaud's syndrome occurs due to peripheral vasospasm, and it is more common in the upper extremities. It may occur due to exposure to various stimuli, including cold temperatures, tobacco, or emotional stress, and certain occupational groups, such as those who use vibrating tools, or it can be secondary to scleroderma, systemic lupus erythematosus, Sjogren's syndrome, and antiphospholipid syndrome.

The pathophysiology behind Raynaud's syndrome is believed to be due to increased levels of $\alpha 2$-adrenergic receptors and their hypersensitivity and also due to an abnormal thermoregulatory response by the sympathetic nervous system.

A Raynaud's phenomenon attack is sudden, and it ranges from intense pallor (white attack) to cyanosis (blue attack), followed by reactive hyperemia due to subsequent relaxation of blood vessels. It typically affects fingers (distal areas), and the thumb is usually spared. If the thumb is involved, it may indicate underlying secondary etiology. Other common complaints include pin and needles sensation, numbness, or finger aches or pain. In severe attacks, digital ulcerations and gangrene may develop.

Diagnosis is primarily based on the clinical history and physical examination. Nail fold capillary microscopy can be helpful, particularly for identifying secondary Raynaud's phenomenon associated with connective tissue diseases. Angiography can be done for patients with digital ulcerations and gangrene.

Management is mainly conservative, and it includes wearing gloves, use of electric or chemically activated hand warmers, avoiding occupational exposure to vibratory tools, abstinence from tobacco, and relocating to a warmer, dryer climate.

Other management options:

• Calcium channel–blockers such as diltiazem and nifedipine are the drugs of choice.

• Fluoxetine has been shown to reduce the frequency and duration of vasospastic episodes.

• Prostaglandins can be used in patients with severe symptoms.

Complications, including digital ulcerations and gangrene, can be surgically treated by debridement and amputation, respectively.

Solution to Question 20:

The given features are suggestive of a juvenile polyp, which are a type of hamartomatous polyp. They are benign growths that are composed of a disorganized mixture of different types of tissues.

Juvenile polyps are malformations of epithelium and lamina propria commonly located in the rectum. It is seen in children <5 years of age. The typical presentation is painless rectal bleeding. Sometimes, there may be a proliferating rectal mass.

The disease may be sporadic or syndromic.

• Sporadic juvenile polyp occurs as a single mass and is called a retention polyp.

• Juvenile polyposis syndrome is characterized by multiple hamartomatous polyps, which may be premalignant.

Morphologically, both of these types are indistinguishable.

• Gross appearance: They appear as pedunculated, smooth reddish lesions, and with cystic spaces.

• Microscopic appearance: The cysts appear as dilated glands filled with mucin and inflammatory debris.



Adenoma: Adenomas are benign tumors that arise from glandular tissue. While they may present with rectal bleeding, they are not commonly seen in young children and are not typically pedunculated like juvenile polyps.

Choristoma: Choristomas are non-neoplastic masses that arise from heterotopic tissue, i.e., tissues found in abnormal locations. They are not associated with juvenile polyps.

Solution to Question 21:

The given picture showing an ulcer with punched-out edges over a pressure point is most likely that of a trophic ulcer/neuropathic ulcer.

A neuropathic ulcer is one that occurs as a result of peripheral neuropathy. In peripheral neuropathy, lack of sensation over the pressure points leads to cumulative microtrauma, minor wounds, or cuts that remain unnoticed but eventually develop into ulcers.

Features of a neuropathic ulcer include

• Painless ulcer

• Punched out ulcer with regular margins

• Often located on the plantar surface of the foot

• Ulcer margins show the presence of calluses

	Ischemic/Arterial Ulcer	Venous Ulcer	Neuropathic Ulcer	Tuberculous Ulcer
Underlying condition	Peripheral vascular disease- mainly atherosclerosis, thromboangiitis obliterans, cervical rib, Raynaud's phenomenon	Varicose veins, previous deep vein thrombosis	Diabetes, peripheral neuropathy	Tuberculosis lymph nodes, cold abscess from bone and joint tuberculosis
Location	Toes, feet, legs, or even upper limb digits	Lower limb, commonly on the medial side of the ankle, never above the middle third of the leg	The plantar surface of the foot, toes	Neck, axilla, and groin
Characteristics	Deep, penetrates the deep fascia exposing the tendons, muscles, and underlying bone. The ulcer is painful, tender, and hyperaesthetic.	Vertically oval, painless, the floor may have pale to no granulation tissue, induration, and tenderness at the base.	Punched-out with regular calloused margins.	Edges are thin, reddish blue, undermined; pale granulation tissue, scanty discharge on the floor, and slight induration at the base
Other features	Features of ischemia in adjacent areas, like pallor, dry skin, brittle nails, loss of hair	Inguinal lymph nodes are enlarged, usually the vertical group, and the ulcer is recurrent; forms again after healing	Dry, cracked, insensate	Regional lymph nodes are enlarged, non-tender, matted

Option A) The following is an image of a venous ulcer

Option B) The following is an image of a malignant ulcer (squamous cell carcinoma)

Option C) The following is an image of an arterial ulcer

Solution to Question 22:

A patient with varicose veins and overlying eczema is classified as C4a under CEAP guidelines.

The Clinical-Etiology-Anatomy-Pathophysiology (CEAP) classification system is used to classify chronic lower extremity venous disease.

The clinical class of C4 under CEAP includes changes in skin and subcutaneous tissue secondary to chronic venous disease. It is further classified as follows:

• C4a - Pigmentation or eczema

• C4b - Lipodermatosclerosis or atrophie blanche

• C4c - Corona phlebectatica

Image A shows tortuous dilated veins - C2, image B shows atrophie blanche - C4b, and image C shows fan-shaped telangiectasia - C4c.



Solution to Question 23:

In this patient with subarachnoid hemorrhage (SAH), the next step is to do an angiography to see the vascular anatomy and identify the site of bleed.

Trauma is the commonest cause of subarachnoid hemorrhage. Spontaneous SAH usually results from the rupture of an aneurysm or an arterio-venous malformation (AVM). Patients present with a sudden and severe headache which is different from any headache they have experienced before. This is described as a thunderclap headache. Other symptoms include meningitic features like neck rigidity and photophobia, impaired consciousness, intraocular hemorrhages, with or without focal neurological deficits. Compression from a posterior communicating artery aneurysm would result in third nerve palsy.

Non-contrast CT head is the first imaging to be done in a suspected case, which would confirm bleeding in more than 98% of cases. In patients with high suspicion and a negative CT, lumbar puncture is done to detect hemoglobin breakdown products like oxyhemoglobin and bilirubin. After confirming the diagnosis, CT or MR angiography is done to locate the aneurysms or AVM. Digital subtraction angiography remains the gold standard investigation.

The definitive management of subarachnoid hemorrhage (SAH) is done surgically by craniotomy and clipping or by endovascular embolization known as coiling. This prevents rebleeding of the aneurysm.

Patients with subarachnoid hemorrhage are at risk of developing complications such as rebleeding, cerebral vasospasm, hydrocephalus, seizures, and hyponatremia. Early recognition and management of these complications are essential to improve outcomes. In addition to surgery and nimodipine, other management strategies for SAH include strict blood pressure control, analgesia for headache, antiemetics, and anticonvulsants if seizures occur. Maintaining a close monitoring of the patient's neurological status, intracranial pressure, and electrolytes is also crucial.

Other Options

Nimodipine: Although it is used to prevent cerebral vasospasm and delayed ischemic neurological deficits, it is not the immediate next step in the management of SAH. Angiography is needed first to locate the aneurysms or AVM. It is administered after the procedure.

Surgery: Although surgery is an important management option for SAH, it is not the immediate next step. Angiography must be done first to identify the site of the bleed and plan for the appropriate surgical or endovascular intervention.

Fibrinolytic therapy: This option is not appropriate for the management of SAH as it may exacerbate the bleeding. Fibrinolytic therapy is used in the treatment of ischemic stroke, not hemorrhagic stroke.

Solution to Question 24:

This patient with a corrosive esophageal injury with complete dysphagia requires a feeding jejunostomy for enteral feeding.

The injury caused by alkali ingestion is deeper due to the process of saponification of fats and dehydration. The injury caused by acids is more superficial because they cause coagulative necrosis with eschar formation, which prevents the deeper penetration of the acid.

Severe corrosive injury may lead to esophageal perforation, bleeding, stricture, and death. Patients can develop dysphagia or odynophagia, which implies esophagal damage.

Minor injuries cause mucosal edema that resolves rapidly with no late sequelae and these patients can be safely fed. In case of severe injuries, a feeding jejunostomy may be appropriate until the patient can swallow saliva satisfactorily.

Management of corrosive esophagitis:

• Hemodynamic stabilization and management of patient's airway

• Endoscopic evaluation of injuries

• Surgical management includes laparotomy for bleeding or perforation. Esophageal replacement is done for very long or multiple strictures.

Stenting is used in patients who do not respond to repeated dilatations for esophageal strictures. They are not used as a treatment in acute settings.

Solution to Question 25:

The given clinical scenario is suggestive of a thyroglossal cyst. It moves with deglutition, protrusion of the tongue, and side-to-side movement is also possible.

A thyroglossal duct cyst is a congenital midline neck swelling that typically presents during childhood often noticed after an upper respiratory infection.

A thyroglossal cyst represents the persistence of the track of the thyroid gland and can be found adjacent to the midline. The thyroid gland descends early in fetal life from the base of the tongue towards its position in the lower neck. At the time of descent, the hyoid bone has not been formed

and the track of the thyroid gland is variable, passing in front, through, or behind the eventual position of the hyoid body.

It can occur anywhere along the migratory course of the duct:

• Infrahyoid (most common site)

• Region of thyroid cartilage

• Suprahyoid

• Floor of mouth

• Beneath foramen cecum

Clinical features of thyroglossal cyst:

• Classically the cyst moves upwards on swallowing and on tongue protrusion. This indicates attachment to the hyoid bone.

• Swelling is smooth, soft, fluctuant (cystic), nontender, mobile, and shows transillumination.

• Thyroid fossa is empty, if there is no thyroid in the normal location.

Complications of thyroglossal cyst:

• The most common complication is infection.

• Thyroglossal fistula is an acquired condition that either follows infection of the cyst which bursts open or due to inadequate removal of the cyst.

• 1% of thyroglossal duct cysts are found to be malignant of which the most common type of cancer is papillary (85%).

However, only the thyroglossal cyst moves with tongue protrusion. This is because the duct extends from the foramen cecum of the tongue to the isthmus of the gland.

Sistrunk's operation is performed to excise the cyst and prevent a recurrence. It involves excision of the whole thyroglossal tract, involving removal of the body of the hyoid bone, and suprahyoid tract through the tongue base to the vallecula at the site of the primitive foramen cecum along with the core of tissue on either side.

Features of hoarseness of voice, cough, neck swelling that increases in size on coughing/Valsalva maneuver, and a gurgling sound heard on pressing the swelling are suggestive of laryngocele.

Solution to Question 26:

Neoadjuvant chemotherapy is defined as chemotherapy given before the definitive treatment (usually surgery) to reduce the bulk of the tumor.

Neoadjuvant therapy is a type of induction therapy that is given before the definitive surgery to downstage the tumor, reduce the morbidity of treatment, know the in vivo response of the tumor to therapy, and increase its chances of success. Chemotherapy, radiotherapy, or hormonal therapy can be given as neoadjuvant therapy.

Option A: Treatment that involves the administration of chemotherapy along with radiation is called chemoradiation.

Option B: Hyperthermic intraperitoneal chemotherapy (HIPEC) is used during surgery in which a hyperthermic solution of anticancer drugs is infused directly into the abdominal cavity after the definitive surgery to kill any residual tumor cells. It is often used to treat cancers that have spread to the peritoneum.

Option D: Treatment given after surgical resection of the tumor to reduce the risk of recurrence and micrometastasis is called adjuvant therapy. Adjuvant therapy can be chemotherapy, radiation therapy, hormone therapy, targeted therapy, or biological therapy.

Solution to Question 27:

qSOFA score comprises of SBP, RR, and GCS.

qSOFA score (quickSOFA) is a bedside prompt that may identify patients with suspected infection who are at greater risk for a poor outcome outside the intensive care unit (ICU).

It uses three criteria, assigning one point each for

• low blood pressure (SBP≤100 mmHg),

• high respiratory rate (≥22 breaths/min),

• altered mentation (GCS score, <15).

The score ranges from 0 to 3 points. The presence of 2 or more qSOFA points is a simple prompt to identify infected patients outside the ICU who are likely to develop sepsis. This is a simplified derivative of SOFA score. The primary method to evaluate the risk of organ dysfunction (sepsis) in patients with suspected infection is by determining the Sequential (sepsis-related) Organ Failure Assessment (SOFA) score, which includes assessment of:

• Circulation - Systolic BP and need for vasopressors; platelet count

• Respiration - PaO_2/FiO_2 and need for mechanical ventilation

• Mentation - Glasgow Coma Scale score

• Hepatic function - Bilirubin level

• Renal function - Creatinine level

Organ dysfunction is defined by a SOFA score of ≥2.

The SOFA score is limited by its requirement of multiple laboratory tests which makes it complex and time-consuming. In order to simplify the scoring process, the "Sepsis - 3" task force recommended the quick sepsis-related organ dysfunction assessment "qSOFA" score.

SOFA vs. qSOFA:

• SOFA is a more comprehensive and accurate scoring system, which is particularly superior to qSOFA in the ICU setting.

• qSOFA is a simplified derivative of SOFA for bedside assessment of individuals at risk for sepsis, particularly useful outside the ICU setting (i.e. ward/emergency department). However, note that it was originally designed as a predictive tool that calculates the risk of death from sepsis rather than a diagnostic tool.

Thereby, the best description of sepsis based on "Sepsis-3" task force recommendations would be as follows:

• Suspected or documented infection with SOFA score of ≥2 or

• Suspected or documented infection with qSOFA score of ≥2

Another score that has been used is the National Early Warning Score (NEWS). It ranges from 0 to 20 and is based on respiratory rate, oxygen saturation, use of supplemental O2, temperature, systolic blood pressure, pulse rate, and level of consciousness. ≥5 out of 20 is considered a positive score.

Other scoring systems for sepsis:

The Systemic Inflammatory Response Syndrome (SIRS) score is no longer recommended as is it not specific to infection.

The Logistic Organ Dysfunction System (LODS) score is not recommended as it is complex and has too many variables.

Surgery INI-CET 2023

Question 1:

A 25-year-old patient presents with fever and weight loss. He gives a history of blood in the stools and vomiting 2 weeks ago. On examination, there is right hypochondriac tenderness and hepatomegaly. Abdominal CT a lesion in the right lobe of the liver with ragged edges, multiple septations, and heterogenous densities. What is the most likely diagnosis?

a) Pyogenic liver abscess

b) Hydatid cyst

c) Amoebic liver abscess

d) Hemangioma

Question 2:

Which of the following tests is done before performing an A-V fistula procedure? The image of the test is given below.

a) Allen's test

b) Adson's test

c) Elevated arm stress test

d) Nicoladoni's test

Question 3:

Match the following positions of the appendix:



a) 1-b, 2-a, 3-d, 4-c

b) 1-c, 2-a, 3-b, 4-d

c) 1-b, 2-d, 3-a, 4-c

d) 1-c, 2-b, 3-a, 4-d

Question 4:

A patient presents with dull aching pain in both legs. He gives a history of prolonged standing and tortuous veins in the legs. Which of the following tests are done for saphenofemoral incompetence?

a) 1, 2, 3 and 4

b) 1, 2 and 3

c) 1, 2

d) 3, 4

Question 5:

Which of the following statements is true about thymoma?

a) Most common neoplasm of thymus

b) Most common posterior mediastinal tumor .

c) Usually symptomatic with dyspnoea, dysphagia and SVC syndrome

d) Chest Xray is the investigation of choice

Question 6:

In a patient who underwent laparoscopic cholecystectomy, what is the ideal urine output?

a) 500 ml/day

b) 0.5 ml/minute

c) 0.5-1.5 ml/kg/hour

d) 1 ml/kg

Question 7:

In a patient presenting with rectal prolapse, which of the following procedures is not done via the perineal approach?

a) Ripstein operation

b) Thiersch operation

c) Altemeier procedure

d) Delorme operation

Question 8:

In a patient with an inguinal hernia, the surgeon reduces the entire hernial sac into the abdominal cavity without reducing the contents. What is this known as?

a) Maydl's hernia

b) Reduction en masse

c) Sliding hernia

d) Incarcerated hernia

Question 9:

Which of the following statements about claudication pain is not true?

a) Rest pain is seen more in the night

b) Pain in intermittent claudication comes at first step of walking.

c) Most common cause is atheroma

d) The clinical stem does not reflect the site of peripheral arterial obstruction

Question 10:

A 73-year-old patient presents with painless rectal bleeding. Which of the below-mentioned investigations are done for this patient?

a) 3 and 4

b) 2, 3 and 4

c) 1 and 2

d) 1, 2 and 3

Question 11:

Smoking increases the risk of which of the following diseases?

a) 2 & 3

b) 1 & 4

c) 2, 3, 4

d) 1, 3, 4

Question 12:

Identify the following image:



a) Hemangioma

b) Cystic hygroma

c) Branchial cyst

d) Ranula

Question 13:

Feeding is done through the given method. Which of the following is not a complication of this?



a) Refeeding syndrome

b) Osteoporosis

c) Pneumothorax

d) Aspiration pneumonia

Question 14:

Identify the surgical instrument shown in the image.

a) Adson's forceps

b) Kocher's forceps

c) Mixter forceps

d) Rampley forceps

Question 15:

A 25-year-old male patient was brought to the casualty after a road traffic accident with a suspected pelvic fracture. On examination, he is found to be in hypovolemic shock. What is the earliest treatment for the control of hemorrhage in this case?

a) Immediate internal fixation

b) Multiple blood transfusions

c) Use bedsheets to compress the pelvis

d) Emergency external fixation

Question 16:

Which of the following are done in Breast Conservation Surgery?

a) 1, 2, 3

b) 1, 3, 4

c) 1 , 2

d) 2, 3

Question 17:

Match the features seen due to compression of the following structures in case of thyroid swelling.

a) 1-d, 2-b, 3-c, 4-a

b) 1-c, 2-d, 3-b, 4-a

c) 1-c, 2-d, 3-a, 4-b

d) 1-a, 2-b, 3-d, 4-c

Question 18:

A 63-year-old female presented to the OPD with abdominal pain and a tense and tender swelling below and lateral to the pubic tubercle. What is the most likely diagnosis of this condition?

a) Strangulated femoral hernia

b) Strangulated inguinal hernia

c) Intussusception

d) Adhesions

Question 19:

A patient presented with a Type 1 Choledochal cyst. All of the following statements are correct except?

a) Increased risk of cholangiocarcinoma

b) Cyst opened and drainage is done

c) Todani classification is used

d) Due to anomalous pancreaticobiliary junction

Question 20:

A patient underwent laparoscopic cholecystectomy for cholelithiasis. Specimen revealed that she had gall bladder carcinoma infiltrating the muscularis layer. What is the management?

a) Radical cholecystectomy with adjuvant chemotherapy

b) Adjuvant chemotherapy

c) Extended cholecystectomy

d) Surveillance

Question 21:

A man presented with a history of stab injury over the arm. On examination, there was a hematoma present, and radial and ulnar pulses were not felt. A 3.5cm longitudinal tear is noted in the brachial artery. Which among the following is the best treatment option?

a) Primary repair with end to end anastomosis

b) Repair with great saphenous vein graft

c) Repair with prosthetic graft

d) Repair with saphenous vein patch

Question 22:

Which of the following drugs are not used in the management of a thyrotoxic crisis?

a) Acetaminophen

b) Aspirin

c) Lugol's iodine

d) Propranolol

Question 23:

Identify the incision.



a) Battle incision

b) Gibson incision

c) Robson incision

d) Lanz incision

Question 24:

Which of the following is the best method to conclusively localize a parathyroid adenoma?

a) SPECT sestamibi

b) Tc99 sestamibi scan

c) USG neck

d) MRI

Question 25:

Which classification of bile duct injury takes into account both vascular as well as the location of the injury?

a) Strassberg

b) Hannover

c) Bismuth

d) McMahon

Question 26:

If open cardiac massage were to be performed on a patient, what category would it come under?

a) Clean

b) Contaminated

c) Clean-contaminated

d) Dirty

Question 27:

A patient who was a known case of rheumatoid arthritis is brought to the ER due to dizziness. Her BP was 90/60mmHg. She was prescribed a medication by her local doctor which she stopped taking suddenly. Her blood sugar was low and serum potassium was 6mEq/L. Which of the following drugs should be given to her next?

a) IV Saline

b) Hydrocortisone

c) Dopamine infusion

d) Sodium bicarbonate

Question 28:

What is the most common cause of hypercalcemia in hospitalized patients?

a) Hypervitaminosis D

b) Malignancy

c) Parathyroid adenoma

d) Parathyroid hyperplasia

Question 29:

An elderly lady was diagnosed to have a gallbladder polyp. She was told that she has a risk for gallbladder cancer. Which of the following is not a risk factor for gallbladder cancer?

a) Porcelain GB

b) Primary sclerosing cholangitis

c) A single gall bladder polyp < 1 cm

d) Gall stone > 3cm

Question 30:

Which of the following is the most important prognostic factor of breast carcinoma?

a) Axillary lymph node status

b) Size

c) Distant metastasis

d) Hormonal status

Answer Key

Question No.	Correct Option
1	c
2	a
3	c
4	c
5	a
6	c
7	a
8	b
9	b
10	d
11	d
12	d
13	d
14	c
15	c
16	a
17	b
18	a
19	b
20	c

21	b
22	b
23	b
24	a
25	b
26	b
27	b
28	b
29	c
30	a

Detailed Explanations

Solution to Question 1:

Features of fever, hepatomegaly, and hypochondriac tenderness are suggestive of a liver abscess. A liver abscess in a patient with a history of blood in the stools and vomiting points to a diagnosis of amoebic liver abscess.

Amoebic liver abscess is more common in males in the age group of 20 - 40 years with a history of travel to endemic areas. Ingestion of Entamoeba histolytica cysts causes diarrhea initially. Following this, the trophozoites reach the liver through the portal venous system and cause an amebic liver abscess. 80% of the cases involve the right lobe. The clinical features of amoebic liver abscess are right upper quadrant pain, tenderness, fever, hepatomegaly, and diarrhea.

Enzyme immunoassays detecting the presence of antibodies against the parasite have high sensitivity and specificity. Ultrasonography and CT scans can visualize the liver abscess without a peripheral rim of enhancement.



The mainstay of treatment for amebic abscesses is metronidazole, which is curative in more than 90% of patients. Luminal agents such as diloxanide furoate can be administered to treat the carrier state. Aspiration of abscess is recommended in case of a high risk of rupture. Pigtail catheter drainage is done in patients not responding to intravenous metronidazole in the first 48–72 hours to improve antibiotic penetration.

Option A: Pyogenic liver abscess is usually seen after 50 years of age and is not associated with a history of travel or diarrhea. Jaundice, bacteremia, and positive blood cultures are common in pyogenic liver abscesses compared to amoebic liver abscesses. Serology testing for Entamoeba is negative. A characteristic rim enhancement is noted in the case of a pyogenic abscess on a CECT abdomen and can differentiate it from an amoebic liver abscess.



Option B: Hydatid cyst a zoonotic infection caused by the Echinococcus species of flatworms. Primary hydatid cysts most commonly occur in the liver. Most patients are asymptomatic but when symptomatic, the patient presents with hepatomegaly, pain, and obstructive jaundice. The rupture of a hydatic cyst can result in fever, pruritus, or anaphylaxis. Imaging findings are cystic lesion with multiple daughter cysts within, detachment of the endocyst membrane (water-lily sign on USG shown below), and areas of calcification.



Option D: Hemangiomas are more common in women and are usually asymptomatic. Larger hemangiomas may present with vague upper abdominal symptoms. On contrast-enhanced CT scan, hemangiomas show asymmetrical, nodular peripheral enhancement that is isodense, with large vessels and progressive centripetal enhancement.

Solution to Question 2:

The image is suggestive of Allen's Test. It is usually done prior to radial artery cannulation and performing an A-V fistula procedure to test the adequacy of ulnar collateral blood flow to the hand.

Allen's test is performed by occluding the radial and ulnar arteries. The patient is asked to make a tight fist so as to exsanguinate the palm. Then the patient is asked to open the hand and the occlusion over the ulnar artery is released. As occlusion of the ulnar artery is released, the color of

the open palm is observed. Normally, the color will return to the palm within 5 – 6 seconds. If the palm remains pale for more than 6-10 seconds, it indicates reduced ulnar collateral flow. The modified Allen's test considers a threshold of 5-15 seconds.



In recent years, studies have shown that, although the Allen test is often used to identify patients at increased risk for ischemic complications from radial artery cannulation, the predictive value of this test is poor. It is also used for identifying patients acceptable for radial artery harvest or use for coronary angiography.

Other options:

Option B: Adson's test is a provocative maneuver for thoracic outlet syndrome. The test is performed by extending the patient's neck and turning the head towards the affected side while they breathe in deeply. This compresses the interscalene space and will result in paraesthesia and obliteration of the radial pulse.

Option C: Elevated arm stress test or Roos test is a provocative test for the diagnosis of thoracic outlet syndrome. For this test, the patient abducts the shoulders to 90 degrees and flexes the elbow to 90 degrees. The patient is made to open and close the hand slowly for three minutes. The test is positive if the patient is unable to complete the test or experiences cramping, pain, or numbness on the affected side.

Option D: The reflex bradycardia and increase in blood pressure that immediately follows the sudden occlusion of an arteriovenous (A-V) fistula is known as the Nicoladoni-Branham sign.



Solution to Question 3:

The correctly matched positions of the appendix are:

1 - b: Retrocaecal - most common position, seen in 74%

2 - d: Paracaecal - seen in 2%

3 - a: Subcaecal - seen in 1.5%

4 - c: Inside the pelvis - seen in 21%

The appendix is a short and broad blind muscular tube that becomes tubular with the differential growth of the caecum. This continued growth of the caecum rotates the appendix into a retrocaecal but intraperitoneal position. In some cases, rotation of the appendix does not occur, resulting in a pelvic, subcaecal, or paracaecal position. During development, if the caecum does not migrate to its normal position in the right lower quadrant of the abdomen, the appendix can be found near the gallbladder. In cases of intestinal malrotation, the appendix may be found in the left iliac fossa.

Various positions of the appendix are shown in the image below.



Solution to Question 4:

Brodie Trendelenburg test and Morrissey's test are performed to test the incompetence of the saphenofemoral valve. Fegan's method is used to detect perforator incompetence while Modified Perthe's test is a test for deep vein thrombosis (DVT).

Brodie Trendelenburg test - Lay the patient down and elevate the leg. Empty the vessels and tie a tourniquet below the SFJ. Ask the patient to stand up:

• Trendelenburg's test part l - Here the tourniquet is released. A rapid filling from above shows saphenofemoral incompetence.

• Trendelenburg's test part ll - Here the tourniquet is not released. A gradual filling from below to upwards indicates perforator incompetence.

Morrissey's cough impulse test - The limb is elevated to empty the varicose veins. Now it is put to bed and the patient is asked to cough forcibly. An expansile cough impulse is felt at the saphenous opening if the saphenofemoral valve is incompetent. Similarly, a bruit may be heard on auscultation. The patient is asked to cough. A cough impulse at the saphenofemoral junction indicates SFJ incompetence.

Fegan's method - Mark the places of excessive bulges within the varicosities. Ask the patient to lie down and empty the varicose vessels. Palpate along the marked area to look for gaps and pits in the deep fascia, which indicates perforator incompetence.

Modified Perthe's test - A tourniquet is tied around the upper part of the thigh to prevent reflux down the vein and the patient is asked to walk around. In the presence of normal communicating and deep veins, the varicose veins will shrink. If there is any block, the veins will get more distended. An increase in pain and prominence of dilated veins indicate DVT.

Solution to Question 5:

The true statement is that thymoma is the most common neoplasm of the thymus.

Thymoma is the most frequently encountered neoplasm of the anterior mediastinum (Option B) in adults and is rare in children. It is seen most frequently in the age group of 40 and 60 years. Most patients with thymoma are asymptomatic (Option C). Large thymic tumors may present with symptoms related to a mass effect, which may include cough, chest pain, dyspnea, or SVC syndrome. Around 10% to 50% of patients with thymoma may have symptoms suggestive of myasthenia gravis or have circulating antibodies to the acetylcholine receptor. Other paraneoplastic syndromes, including red cell aplasia, hypogammaglobulinemia, systemic lupus erythematosus, Cushing's syndrome, or SIADH, may be present.

While the mediastinal mass can be identified on screening chest X-ray, imaging with contrast-enhanced CT and MRI (Option D) have a critical role in characterizing the mediastinal masses. The diagnosis may be suspected based on history and CT imaging but imaging alone is not diagnostic. The diagnosis is usually made after surgical resection due to the relative difficulty of obtaining a needle biopsy. Biopsy should be avoided in cases where imaging is highly suggestive of thymoma. Masaoka staging system is used for thymomas and is based on the gross or microscopic invasion of the capsule and of surrounding structures, as well as on the presence of metastases.

The definitive treatment for thymoma is complete surgical removal. Thymectomy leads to improvement or resolution of symptoms of myasthenia gravis in only about 25% of patients with thymomas.

Solution to Question 6:

The ideal urine output in that patient following laparoscopic cholecystectomy is 0.5-1.5 ml/kg/hour.

Normal urine output is expected in the postoperative period of a patient who underwent laparoscopic cholecystectomy. The average normal urine output in an adult is 0.5-1.5 ml/kg/hour (Option B) or approximately 1500 mL/day (Option A). A minimum urine output of 400 mL/day is essential to excrete the end products of protein metabolism.

Option D: Normal urine output in ml/kg would be 0.5 - 1.5 ml/kg/hour multiplied by 24 which is 12-36 ml/kg per day.

Solution to Question 7:

Among the given options, the Ripstein operation is the abdominal approach for rectal prolapse.

It is managed by using the following approaches:

• The perineal approach is easy to perform with fewer complications but a high recurrence rate. The surgeries done using this approach for rectal prolapse are the following:

• Thiersch's wiring

• Delorme's mucosectomy

• Altemeier's procedure or perineal rectosigmoidectomy

• The abdominal approach is difficult to perform with more complications but a lower recurrence rate. The surgeries done using this approach are the following:

• Moschcowitz repair

• Ripstein and Wells rectopexy; ventral rectopexy or suture rectopexy

• Resection rectopexy

If the patient is fit enough for an abdominal procedure, it is preferred over the perineal procedure, since the risk of recurrence is lower. The perineal approach is preferred in elderly patients with comorbidities and in young men since the abdominal approach may injure the pelvic autonomic nerves which may result in sexual dysfunction.

Solution to Question 8:

The surgeon reducing the entire hernial sac into the abdominal cavity without reducing the contents is referred to as reduction en masse of the hernia.

Reduction en-masse of inguinal hernia refers to the reduction or migration of the hernial sac along with the incarcerated bowel into the properitoneal space. The properitoneal space lies between the parietal peritoneum and the anterior abdominal wall. This is likely to occur following forcible attempts to reduce the hernia but may occur spontaneously. Though the hernial sac no longer lies in the inguinal or femoral canal, it contains an obstructed bowel loop that causes small bowel obstruction. This is a rare condition as forcible reduction of hernias is not done and early repair is carried out.

The patient may give a history of difficult reduction and the symptoms of bowel obstruction persist even after the hernia is reduced. On a CT scan, a hernial sac in properitoneal space with a fibrous constriction at the neck may be visualized. This hernial sac will contain the bowel loop with wall enhancement and fluid that suggest strangulation. Early surgical intervention is required.

Other options:

Option A: Maydl's hernias are defined as the presence of two small bowel loops within a single hernial sac, that is, there are two efferent and two afferent loops of the bowel, forming a "W" shape.



Option B: Sliding hernia occurs when an internal organ composes a portion of the wall of the hernia sac. The most common viscus involved is the sigmoid colon and its mesentery on the left side and the caecum on the right side. It is more common on the left side. They are usually variants of indirect inguinal hernias. The failure to recognize the visceral component of the hernia sac resulting in injury to the bowel or bladder is the major concern in sliding hernia.

Option D: Hernias which cannot be reduced are referred to as incarcerated hernias while hernias with compromised blood supply to its contents are known as strangulated hernias.

Solution to Question 9:

Pain in intermittent claudication is brought on by walking and doesnot start with the first step of walking.

Peripheral artery occlusion or stenosis is predominantly caused by atherosclerosis or atheroma (Option C) and thromboembolic disease. The resulting stenosis or occlusion produces signs and symptoms related to the organ that is supplied by the affected artery. Such occlusion in the lower limbs presents with claudication pain, rest pain, and gangrene. Though the muscle groups affected by claudication are present at one anatomical level below the level of arterial disease, the clinical stem does not reflect the exact site of the peripheral arterial obstruction. (Option D)

Intermittent claudication pain is a feature of arterial stenosis or occlusion that occurs due to the anaerobic metabolism occurring in the muscle. Claudication pain may be of different types and can be differentiated as follows:

• Vascular (intermittent) claudication pain is brought on by walking and is relieved by rest within 5 minutes. (Option B)

• Claudication pain present on taking the first step of walking is classical for osteoarthritis.

• Neurogenic claudication occurs with changes in the position of the back such as bending forward and straightening. It takes more time to be relieved by rest (more than 5 minutes). It may be seen in cases of nerve compression from lumbar intervertebral disc prolapse or spinal stenosis.

Rest pain occurs when the occlusive disease progresses in severity and anaerobic respiration occurs even at rest. Though this pain is present during the day, it is exacerbated on lying down at night (Option A) due to loss of gravity dependant perfusion of the foot. This may be relieved by hanging the foot out of the bed to restore the gravitational effects.

Solution to Question 10:

Colonoscopy, CECT and CEA (Carcinoembryonic antigen) can be done in an old patient presenting with painless rectal bleeding. Serum CA 19-9 is a pancreatic cancer biomarker and has no role in this case.

Rectal bleeding occurs due to pathology of the lower gastrointestinal tract below the ligament of Treitz. The differentials that can be considered in a case of rectal bleeding are:

• Colon cancer

• Inflammatory bowel disease

• Diverticular diseases

• Hemorrhoids

• Angiodysplasia

Colonoscopy with biopsy has a key role in the diagnosis of inflammatory bowel disease, distinguishing between ulcerative colitis and Crohn's colitis and monitoring the response to treatment. Colonoscopy is also the investigation of choice if colorectal cancer is suspected, especially in old patients. Angiodysplasia is suspected in patients above the age of 50 and can be identified by a colonoscopy.

Computed tomography (CT) colonography is sensitive in picking up polyps and has the advantage of being less invasive than colonoscopy. Spiral CT can aid in the diagnosis of complicated diverticular disease.

Carcinoembryonic antigen (CEA) is a non-specific serum biomarker that is elevated in colorectal cancer and hence may be done in this case.

Solution to Question 11:

Smoking increases the risk of which of duct ectasia, Mondor's disease, and breast carcinoma.

Duct ectasia is defined as dilated major milk ducts. It is considered a disorder of involution as part of ANDI. Abnormally dilated ducts are filled with debris. This acts as an irritant and can lead to periductal inflammation and subsequent fibrosis, leading to nipple retraction. Duct ectasia appears to be associated with smoking and diabetes.

The thrombophlebitis of the superficial veins of the breast and anterior chest wall is called Mondor's disease. The pathognomonic feature is a tender thrombosed subcutaneous cord, usually attached to the skin. When the skin over the breast is stretched by raising the arm, a narrow, shallow, subcutaneous groove alongside the cord becomes apparent. Smoking was found to be a potential factor involved in the pathogenesis of Mondor's disease. Other risk factors include breast

surgery, breast biopsy, an inflammatory process, breast cancer, and trauma.

The risk factors of breast cancer can be divided into modifiable (those that can be modified by adopting a healthy diet and lifestyle) and non-modifiable risk factors. The events increasing the oestrogenic exposure of the breast are said to be risk factors for breast cancer, such as early menarche, late menopause, nulliparity, late first pregnancy, and hormone replacement with high estrogen therapy. Smoking is a modifiable risk factor with a relative risk (RR) = 1.14 for smoking 25 or more cigarettes/day and RR = 1.07 for smoking for 20 years or more.

Fibroadenoma is the most common cause of a breast lump in women aged 15–25 years. It arises from the hyperplasia of a lobule and is surrounded by a well-defined capsule. Smoking is not a risk factor for fibroadenomas.

Solution to Question 12:

The given scenario and the image of a cystic swelling on the floor of the mouth are suggestive of a ranula.

Ranula is a mucus extravasation cyst arising from the sublingual gland. It produces a characteristic translucent swelling that gives the appearance of a frog's belly. A ranula usually resolves spontaneously.

A plunging ranula is a mucus retention cyst. It has mucus collecting below the gland and perforating through the mylohyoid muscle.

Treatment should include removal of the sublingual gland as this gland has multiple ducts and ranulas can recur if the gland is left behind. Incision, drainage and marsupialisation have low success rates.

Other options:

Options A: Hemangioma is the most common vascular tumor and is characterized by increased abnormal proliferation of endothelium. Hemangiomas develop in infancy and are usually not visible at birth. They appear red or maroon if superficial, and bluish if deep.

Option B: Cystic hygroma is usually seen in neonates and early childhood. They are cystic swellings filled with lymph fluid and are characteristically brilliantly translucent. They occur in the neck and can involve the parotid, submandibular, tongue, etc. The swelling is soft and partially compressible. Definitive treatment is the excision of the swelling.

Option C: The branchial cyst develops from the vestigial remnants of the second branchial cleft. Swelling in the neck beneath the anterior border of the upper third of the sternomastoid muscle. It is smooth, soft, fluctuant, and often transilluminate. The below image shows a large right-sided branchial cyst.

Solution to Question 13:

The given image shows a central line. Pneumothorax, osteoporosis, and refeeding syndrome are likely complications in patients receiving total parenteral nutrition through a central line.

Aspiration pneumonia is more common in enteral feeding and can be prevented by placing the patient in a semi-recumbent position while feeding.

Complications of enteral nutrition are as follows:

• Tube-related (most common): Malposition, displacement, blockage, breakage/leakage, local complications (e.g. erosion of skin/mucosa)

• Gastrointestinal: Diarrhoea, bloating, nausea, vomiting, abdominal cramps, aspiration, constipation

• Metabolic/biochemical: Electrolyte disorders, vitamin, mineral, and trace element deficiencies, drug interactions

• Infective: Exogenous (handling contamination) and endogenous (patient)

Other options:

Option A: Refeeding syndrome is a significant metabolic complication of both parenteral and enteral feeding. It occurs days after feeding is commenced in patients who have been severely malnourished. The major pathological process is hypophosphatemia which causes fluid and electrolyte shifts between the intra- and extracellular compartments.

Option B: Osteoporosis and osteomalacia are complications of long-term parenteral nutrition and may lead to fractures or kidney stones. Calcium, phosphate, vitamin D supplementation, and bisphosphonates can help prevent as well as treat this complication.

Option C: Pneumothorax is the most common complication of line insertion and most commonly occurs during the insertion of subclavian lines. It is managed by insertion of a chest drain.

Solution to Question 14:

The given surgical instrument is a Mixter's right angle forceps that can be used to pass ligatures and dissect pedicles.

Other options:

Option A: Adson's forceps or Cerci-Adson-Brown forceps have a double grasping tip that ensures grasping with better stabilization. It is the most appropriate forceps for grasping cartilage in rhinoplasty. Its multi-teeth tip prevents cartilage damage or bounce.

Option B: Kocher forceps have an opposing tooth in the tip and serrations in the distal blades. It is used to hold pedicles, tough structures, and cut ends of muscles. It is also used for holding ribs during rib dissection and for holding the resected bowel.

Option D: Rampley's sponge-holding forceps have a fenestrated, serrated, and flat end. The uses are:

• to clean the operative field and swab the cavities

• to mop the oozing area

• to hold the gallbladder and cervix during surgeries

• for blunt dissections

Solution to Question 15:

The earliest treatment to control hemorrhage in a case of suspected pelvic fracture is to use bedsheets to compress the pelvis similar to a compression binder.

The most likely source of bleeding in hemodynamically unstable patients following blunt trauma is abdominal or pelvic bleeding. A pelvic binder or a sheet should be applied around the true pelvis of all hemodynamically unstable patients and not removed until after a pelvic fracture has been excluded. It must be correctly positioned at the level of the greater trochanters. This is a potentially life-saving procedure that has to be done in the emergency department. Following this, they require an immediate pelvic radiograph. Mechanical testing of the pelvis should never be performed. A pelvic fracture should always be diagnosed radiographically.

In fractures of the pelvic ring, there is a risk of injury to the major vessels (internal iliac and external iliac vessels) and to the venous plexus around the sacroiliac joint. Bleeding from the venous plexus is the major cause of pelvic hemorrhage. Pelvic fractures can have a high mortality rate of more than 40% in some open pelvic fractures. Early mortality is mostly the result of hemorrhage.

Bleeding of venous origin can be controlled by compression either by binding/ external fixator or by extraperitoneal pelvic packing to compress the pelvic veins. Persistent bleeding after packing or bleeding of arterial origin may require angioembolisation. Severe pelvic injuries require a multidisciplinary team approach. An exploratory laparotomy may be considered if

• the source of the bleeding is in doubt

• FAST/CT results are positive, showing a significant amount of blood in the peritoneal cavity

• concurrent intra-abdominal injury cannot be excluded

Solution to Question 16:

Wide local excision, axillary node dissection, and frozen section (biopsy) during surgery are done as a part of Breast Conservation Surgery (BCS). All patients with BCS must receive radiotherapy (Statement 4) and are together known as Breast Conservation Therapy (BCT).

Breast conservation surgery involves removing the tumor with a 1mm margin of normal breast tissue. Removal of primary breast cancer is also called lumpectomy/ tylectomy/ partial mastectomy/ wide local excision (Statement 1). Histological examination via intraoperative frozen section (Statement 3) is done to examine for positive margins. Patients with positive margins should have a revision of margins known as 'cavity shave'. Assessment of regional lymph nodes is done via a separate incision in the axilla (Statement 2).

Absolute contraindications of BCT:

• 1st trimester of pregnancy (may be done for some 2nd and many 3rd trimester patients who can receive radiation after delivery)

• Persistently positive margin (2 microscopy shows positive margins)

• Multicentric lumps

• Diffuse microcalcification

•

Extensive ductal carcinoma in situ (DCIS)

•

Inflammatory breast cancer

Relative contraindications of BCT:

• Active SLE (systemic lupus erythematosus), scleroderma

• Large lump-to-breast ratio (>5 cm size lump)

• Known or suspected genetic predisposition to breast cancer (i.e., BRCA1, BRCA2), Li-Fraumeni syndrome (p53 mutation)

• Multifocal disease

• Prior radiation therapy to the chest wall or breast

Solution to Question 17:

The features seen due to compression of the surrounding structures in case of thyroid swelling are as follows:

The normal thyroid gland is impalpable. Thyroid swellings may be uniform enlargement of the gland, isolated solitary nodules, or multinodular goiter. Large thyroid swelling may compress the surrounding structures and present as pressure symptoms.

• Compression of the trachea can cause respiratory symptoms like dyspnea and stridor.

• Pressure on the esophagus causes dysphagia.

• Compression of the sympathetic chain leads to Horner's syndrome.

• Unilateral paralysis of the recurrent laryngeal nerve produces hoarseness. Vocal cord paralysis from RLN compression is extremely rare and should raise concern for malignant change.

1. Trachea	c. Dyspnea
2. Esophagus	d. Dysphagia
3. Sympathetic chain	b. Horner's Syndrome
4. Recurrent Laryngeal Nerve	a. Hoarseness

Solution to Question 18:

The above clinical scenario of a tense and tender swelling present lateral and below the pubic tubercle is suggestive of a femoral hernia. The femoral hernia traverses the femoral canal, which lies medial to the femoral vein.

A femoral hernia occurs in both males and females, but it is more common in females. It will pass through the femoral ring proximally to enter the femoral canal. The femoral canal is bound by the femoral vein laterally, the inguinal ligament anteriorly, and the pelvic bone with iliopectineal ligament (Cooper's) and lacunar ligament (Gimbernat's) medially. Lacunar ligament has a sharp unyielding edge that prevents the reduction of the femoral hernia. The incidence of strangulation in femoral hernias is high because of the narrowness of the neck of the sac and the rigidity of the femoral ring.

All patients with unexplained bowel obstruction should be examined properly for a femoral hernia. Surgery is the only treatment for femoral hernia, and these patients should be treated on priority because of the high risk of strangulation. The surgical approaches which are used are:

• Low approach (Lockwood): suitable when there is no risk of strangulation; bowel can't be inspected and resected in this approach

• Inguinal approach (Lotheissen)

• High approach (McEvedy): suitable for emergency situations when the risk of bowel strangulation is high

• Laparoscopic approach: both TEP and TAPP approaches can be used; the laparoscopic approach is recommended in women.

Other Options:

Option B: Inguinal hernia is above and medial to the pubic tubercle. Another structure that is useful in differentiating between inguinal and femoral hernia is inguinal ligament.
An inguinal hernia leaves the abdominal cavity just above the inguinal ligament and a femoral hernia leaves just below the inguinal ligament.

Option C: In case of Intussusception, patients present with cramping abdominal pain, palpable sausage-shaped abdominal mass, vomiting, and passage of red currant jelly stools.

Option D: Adhesions can lead to intestinal obstruction, but there won't be any swelling below and lateral to the pubic tubercle,.

Solution to Question 19:

In the given clinical scenario, a patient presented with a Type 1 Choledochal cyst. Cyst opening and drainage will not be done in this patient. the preferred course of action involves the complete surgical excision of the cyst. Complete resection of the cyst is essential because there is a risk of malignant transformation (cholangiocarcinoma). (Option A)

Choledochal cysts are congenital cystic dilations of the intrahepatic and/or extrahepatic biliary tree. They present with a triad of jaundice, a palpable upper quadrant mass, and abdominal pain.

The clinical triad (abdominal pain, mass, and jaundice) and the MRCP (Magnetic Resonance Cholangiopancreatography) findings suggest choledochal cyst. According to the Todani classification, the type III choledochal cyst opens into the duodenum. The treatment of choice for type III choledochal cyst is Endoscopic sphincterotomy followed by surveillance.

Choledochal cysts are congenital dilations of the intra and/or extrahepatic biliary system.

Pathogenesis involves an Anomalous Pancreaticobiliary Junction (APBJ) which may allow reflux into the biliary tract, leading to inflammation, increased biliary pressure, and cyst formation. (Option D)

Clinical features of choledochal cysts are as follows

• 60% of cases are diagnosed before the age of 10 years

• more common in females than in males (4:1)

• The classic symptom triad includes abdominal pain, right upper quadrant mass, and jaundice (most consistent symptom)

• Other symptoms- nausea, weight loss, fever, pruritus, etc

Complications include

• Cholangitis (may lead to cirrhosis and portal hypertension)

• Pancreatitis

• Bile peritonitis (due to cyst rupture)

• Cholangiocarcinoma

Diagnostic investigations for choledochal cysts are as follows:

• Ultrasonography:

• Confirms the presence of an abnormal cyst

• Define the dimensions of the cyst

• Define its relationship to the vascular structures in the porta hepatis, as well as the intrahepatic ductal configuration

• Magnetic Resonance Imaging (MRI/MRCP) is the investigation of choice. It reveals the anatomy of the ducts

• CT - delineates the extent of the intra- or extrahepatic dilatation

• ERCP - defines the distal biliary tree

Todani classification (Option C) of the choledochal cysts along with their management is as shown below,

Biliary-enteric reconstruction is not required in Type III choledochal cyst as it has the lowest risk of malignancy, and it may not be associated with ABPJ (anomalous pancreaticobiliary junction). Therefore, endoscopic drainage is sufficient.

Type of choledochal cyst	Surgical management	
Type I - fusiform dilatation of the bile duct (most common - seen in 75% of patients)	complete surgical excision, cholecystectomy, and Roux-en-Y hepaticojejunostomy	
Type II - saccular diverticulum from the common bile duct	Diverticulectomy, cholecystectomy, and common bile duct repair	
Type III- diverticulum within the pancreas, also known as choledochoceles	ERCP + sphincterotomy (Only cyst in which reconstruction is not required)	
Type IV	a - multiple cysts (both intrahepatic and extrahepatic)	Liver transplant
b - multiple cysts limited to the extrahepatic biliary tree	Kasai procedure (Portoenterostomy)	
Type V - multiple cysts involving only intrahepatic ducts (Caroli's disease)	Liver transplant	

Solution to Question 20:

In the given clinical scenario, gallbladder carcinoma with T1b lesions is detected incidentally after cholecystectomy. Treatment guidelines for the same are as follows:

• Up to stage Ia and stage Ib with negative margins, a standard cholecystectomy would suffice.

• Stage Ib with perineural, lymphovascular invasion, positive cystic duct margins, or poor differentiation, and stage II, extended cholecystectomy should be performed. Extended cholecystectomy involves the removal of the gall bladder, segments IVB and V of the liver, and lymph nodes along the hepatoduodenal ligament, with or without removal of the common bile duct (CBD).

However, the likelihood of nodal disease increases significantly with T1b lesions and perineural, lymphatic, or vascular invasion, therefore, extended cholecystectomy is generally recommended for all patients who are medically fit with T1b or greater level of invasion.

If the initial procedure was performed laparoscopically, the surgeon should examine the laparoscopic port sites. The finding of disease at the port sites is a sign of generalized peritoneal disease and carries a very poor prognosis. However, resection of port sites is no longer

recommended.

Chemotherapy is not required in early gallbladder cancer. Radiation has no role.

Treatment of gallbladder carcinoma:

TNM staging	Treatment
T1a	Simple cholecystectomy
T1b, T2, T3	Radical cholecystectomy
T4	Unresectable, chemotherapy and palliative nonsurgical therapy are provided
N1	Regional lymphadenectomy
N2	Radical cholecystectomy plus hepatopancreatic duodenectomy

Solution to Question 21:

In this patient, who has sustained a long tear to the brachial artery due to a stab injury, the best treatment option is to repair it with a great saphenous vein interposition graft harvested from the patient's uninjured leg.

The most desirable graft used is the great saphenous vein . A native vein graft is preferable to a prosthetic graft because:

• Its elastic properties are comparable to that of an artery and enable pulsatile flow

• The diameter is similar to that of an extremity artery

• It is not thrombogenic and has long term patency

Other grafts that can be used are cephalic vein, lesser saphenous vein, and prosthetic conduits. Cephalic vein is not the first choice because it is less muscular than greater saphenous vein, and might also compromise the upper limb venous access in the already injured patient. Both the cephalic and the lesser saphenous veins present problems while harvesting in a trauma patient.

Other options:

Option A: Primary arterial repair with end-to-end anastomosis can be done only if the repair is tension-free. If the length of the injured area is long, vessel injuries cannot be repaired by primary end-to-end anastomoses and instead require an interposition graft.

Option C: Prosthetic grafts are only preferred when there has been a trauma to the native saphenous vein or when there is a size mis-match. Polytetrafluoroethylene (PTFE) graft is commonly used.

Option D: Saphenous vein patch is used in the closure of the arteriotomy site during carotid endarterectomies.

Solution to Question 22:

Aspirin has no role in the management of a thyrotoxic crisis. All of the other options listed are used in managing the symptoms. Aspirin and other NSAIDs are to be avoided as they can interfere with protein binding and further exacerbate the condition.

Thyroid storm/ Thyrotoxic crisis:

It is a rare but severe life-threatening complication of hyperthyroidism induced by the release of excessive thyroid hormones.

Causes:

• Abrupt cessation of antithyroid medications

• Infection

• When a thyrotoxic patient has been inadequately prepared for thyroidectomy.

• Thyroid or non-thyroid surgery

• Trauma in patients with untreated thyrotoxicosis

• Amiodarone administration

• Exposure to iodinated contrast agents

• Following RAI (radioactive iodine) therapy

Features:

• They present 12–24 hours after surgery with hyperpyrexia and severe dehydration

• Cardiovascular: Circulatory collapse, hypotension, palpitations, tachycardia, cardiac arrhythmias, cardiac failure

• GI symptoms: vomiting, diarrhea, jaundice.

• Neurological symptoms: restlessness, irritability, delirium, tremor, convulsions, and coma

Treatment:

• Supportive measures – rehydration by proper fluid therapy, tepid sponging, cooling blankets

• Acetaminophen for fever.

• Thionamides like Propylthiouracil (PTU) or methimazole or carbimazole are used. They inhibit thyroid hormone synthesis. Propylthiouracil also inhibits the peripheral conversion of T4 to T3. Hence, PTU is the preferred drug in thyroid storm. However, the risk for severe liver injury and acute liver failure associated with PTU therapy makes either carbimazole or methimazole first-choice in patients with deranged liver function tests.

• Lugol's iodide is given one hour after intake of thionamides. They inhibit thyroid hormone synthesis by Wolff- Chaikoff effect. Large doses of iodide inhibit organic binding of the iodide and hence decrease thyroid hormone synthesis.

• Propranolol is used to reduce tachycardia and other adrenergic manifestations. It also reduces the peripheral conversion of T4 to T3.

• Glucocorticoids – reduce the peripheral conversion of T4 to T3 and controls the shock.

• Digitoxin, cardiac monitoring is crucial.

Solution to Question 23:

The image above shows a Gibson incision. Gibson incision is used for kidney transplantation or as an extraperitoneal approach to the distal ureter with low morbidity.

The traditional pelvic Gibson incision is an oblique or curvilinear incision that starts 3 cm above and parallel to the inguinal ligament and extends vertically 3 centimeters medial to the anterior superior iliac spine (ASIS) extending down toward the inguinal fold and terminating just lateral to the rectus muscle or continuing to above the symphysis pubis.

The Gibson incision is the traditional approach for renal transplant and transplant nephrectomy. This muscle-splitting incision provides great extraperitoneal access to pelvic vessels, the lower ureter, and the bladder for benign and malignant disease and is also utilized as a part of minimally invasive nephrectomy for specimen extraction or use of a hand port.

Other Options:

Option A: battle incision a lower right paramedian incision but placed more laterally than the standard paramedian incision. It is suitable for dealing with acute appendicitis and pathologies in the right lower quadrant of the abdomen.

Option C: Mayo Robson's incision is a right upper paramedian incision extended like a hockey stick to the midline. It is indicated in cholecystectomy.

Option D: Lanz incision, a transverse or skin crease incision for appendicitis, 2 cm below the umbilicus, centered on the midclavicular–mid-inguinal line. Due to its continuation with Langer's lines, the Lanz incision produces much more aesthetically pleasing results with reduced scarring.

An image depicting other abdominal incisions is shown below:



Solution to Question 24:

The best method to conclusively localise a parathyroid adenoma is SPECT sestamibi. Single-photon emission computed tomography (SPECT) is commonly used because of its improved spatial resolution. Sestamibi combined with CT can be used to detect smaller parathyroid tissues. It can also detect ectopic glands in the mediastinum or in the deeper cervical region.



Ultrasonography:

• It is inexpensive and noninvasive.

• Mediastinal parathyroid tissue cannot be detected because USG cannot penetrate the sternum or clavicle.

Nuclear medicine scanning with radiolabeled Sestamibi (Tc99 sestamibi scan) is also a widely used technique. Sestamibi accumulates in parathyroid adenomas. This radionuclide is concentrated in the thyroid and parathyroid tissue but usually washes out of normal thyroid tissue in under an hour. It persists in abnormal parathyroid tissue. Tc99 sestamibi scan is a 2D study, so SPECT sestamibi, which is a 3D study is preferred over Tc99 sestamibi. scan. SPECT sestamibi has high specificity and sensitivity as compared to Tc99 sestamibi.

MRI is rarely used because of its low sensitivity. Occasionally, MRI may be indicated in pregnant patients in whom ultrasound is noninformative because MRI avoids radiation exposure.

Solution to Question 25:

Hannover classification of bile duct injury takes into account both vascular as well as the location of the injury.

Hannover, bile duct injuries were divided into five types from A to E. Vascular injuries are included in Type C and Type D.

• Type A is peripheral bile leakage.

• Type B is stricture of CHD or CBD without injury.

• Type C is lateral CHD or CBD injury.

• Type D is the total transection of CHD.

• Type E is bile duct stricture of the main bile duct without bile leakage at the postoperative state.

Other Options:

Option A: The Strasberg classification of bile duct injury is a widely used system to anatomically define bile duct injuries.

Option C: The Bismuth–Corlette classification system provides an anatomic description of the tumor location and longitudinal extension in the biliary tree.

Option D: The McMahon classification separated the injuries by the width of bile duct injury

Solution to Question 26:

Open cardiac massage is classified as a contaminated wound.

Contaminated wounds are class III surgical wounds. They include fresh open accidental wounds, surgeries with a major break in sterile technique, gross spillage of the gastrointestinal tract, or incision through an inflamed nonpurulent tissue. Examples include:

• Open cardiac massage

• Penetrating abdominal trauma

Surgical wounds are classified based on the expected amount of bacterial load at the time of surgery.

Open cardiac massage is performed when there is inadequate cardiac output, after emergency thoracotomy for penetrating chest trauma, during pericardial tamponade.

Class I surgical wounds are clean wounds (option A). They are noninfected surgical wounds without any inflammation. Examples include:

• Hernia repair

• Breast biopsy specimen

Class II surgical wounds are clean/contaminated wounds (option C). They are surgical wounds that include an entry hollow viscus such as the respiratory tract or gastrointestinal tract under controlled conditions without significant spillage. Examples include:

• Cholecystectomy

• Elective gastrointestinal surgery

Class IV surgical wounds are dirty wounds (option D). They are old traumatic wounds with the presence of necrotic tissue and perforated viscus with a heavy load of contamination. Examples

include:

• Perforated diverticulitis

• Necrotizing soft tissue infection

Solution to Question 27:

The given clinical scenario is suggestive of an Addisonian crisis. It is a life-threatening condition caused due to sudden insufficiency of adrenal hormones. If a patient displays features of adrenal insufficiency, treatment must immediately be commenced, before awaiting the biochemical diagnosis. Intravenous administration of hydrocortisone 100 mg every 6 hours, >2 litres of saline is given in 6 hours under careful cardiovascular monitoring.

The symptoms of adrenal insufficiency are as follows:

• Glucocorticoid deficiency leads to:

• Fatigue, lack of energy

• Weight loss, anorexia

• Myalgia

• Hypoglycemia

• Low blood pressure, postural hypotension

• Mineralocorticoid deficiency leads to:

• Abdominal pain, nausea, vomiting

• Dizziness, postural hypotension

• Low blood pressure, postural hypotension

• Hyponatremia

• Hyperkalemia

• Adrenal androgen deficiency leads to dry and itchy skin, loss of libido

• The hyperpigmentation of sun-exposed areas results from excess ACTH (adrenocorticotropic hormone) secretion which occurs in response to decreased adrenal cortical hormones.

Primary causes of the Addisonian crisis include:

• Adrenal haemorrhage

• Acute infection

Secondary causes of the Addisonian crisis include:

• Surgery

• Cessation of steroid therapy

• Postpartum pituitary necrosis.

Solution to Question 28:

Cancer is the most common cause of hypercalcemia in hospitalized patients, whereas hyperparathyroidism is the most common cause of hypercalcemia detected incidentally in outpatients. The causes of hypercalcemia in cancer :

• Cancers affecting bone like multiple myeloma, leukemia, bony metastasis - cause the bone to break down which releases excess calcium into the blood

• Pseudohyperparathyroidism - tumours producing parathyroid hormone-related peptide (PTHrP) cause the bone to release calcium into the blood. The most common carcinoma associated is squamous cell carcinoma of lung.

• Cancers affecting kidneys - failure to remove excess calcium from the blood

• Dehydration caused by nausea and vomiting - difficult for the kidneys to remove calcium from the blood properly

• Lack of physical activity - causes bone breakdown, releasing calcium into the blood

Solution to Question 29:

A single gallbladder polyp <1 cm is not a risk factor for carcinoma gallbladder. Gallbladder polyps >10mm have malignant potential.

Risk factors for carcinoma gallbladder:

• Gallstones of size >3 cm (most important risk factor)

• Porcelain gallbladder

• Anomalous pancreato-biliary junction

• Choledochal cyst

• Adenomatous polyps

• Primary sclerosing cholangitis

• Obesity

• Salmonella typhii infection.

Gallbladder carcinoma is a rare malignancy that is seen in the elderly. It is associated with a poor prognosis and is indistinguishable from cholecystitis or cholelithiasis. It is usually diagnosed when patients undergo cholecystectomy for gallstones.

Its clinical features include right upper quadrant pain, nausea, vomiting, and anorexia. Jaundice is a late feature. CA 19-9 is elevated in the carcinoma gallbladder and is used for monitoring.

The preoperative diagnosis is often made on ultrasonography and followed up by a CT scan or MRI/Magnetic resonance cholangiopancreatography.

A percutaneous biopsy under radiological guidance is done for pathological examination. PET scanning helps in detecting metastatic disease.

In select cases, a laparoscopic examination is useful in staging the disease. Laparoscopy can also detect peritoneal or liver metastases. The most important prognostic factor is the depth of invasion (T-stage).

Solution to Question 30:

Axillary lymph node status is the most important prognostic factor in patients with breast cancer (non-metastatic). Women with node-negative disease have less than a 30% risk of recurrence, compared with as much as a 75% risk for women with node-positive disease. The presence of lymph node involvement at initial diagnosis also serves as a prognostic factor in recurrent breast cancer.

Prognostic factors in metastatic breast cancer include:

- Relapse-free intervals
- Site of metastasis
- Estrogen receptor (ER) or progesterone receptor (PR) status

Hormone receptor status (estrogen receptor [ER]/progesterone receptor [PR]) is the most important prognostic factor in patients with metastatic breast cancer. Long relapse-free intervals, absence of brain or visceral metastases, and ER/PR positivity are associated with better prognosis in metastatic breast cancer.

Surgery INI-CET 2024

Question 1:

Which of the following are extra-esophageal complications associated with GERD?

a) 1,2

b) 2,3,4

c) 1,2,3,4

d) 3,4

Question 2:

A 25 year old male presented with difficulty in swallowing food and weight loss for past few months. He also complains of regurgitation of undigested food and halitosis. Barium swallow was performed. What is the diagnosis?



a) Achalasia cardia

b) Esophageal carcinoma

c) Barrett's esophagus

d) Esophageal stricture

Question 3:

The instruments shown below are used for which of the following procedures?



a) Orogastric tube insertion

b) Chest tube drainage

c) Percutaneous endoscopic gastrostomy

d) Flatus tube insertion

Question 4:

Patients diagnosed with breast cancer are usually advised to undergo ER screening test. What is its significance?

a) Diagnostic marker

b) Prognostic marker

c) Monitoring biomarker

d) Surrogate molecular marker

Question 5:

A 45-year-old female presents with recurrent episodes of right flank pain, hematuria, and a history of urinary tract infections. Urine analysis shows alkaline urine with a pH of 8.0. Which of the following is the most likely underlying cause of this patient's condition?



a) Triple phosphate stones

b) Calcium oxalate

c) Uric acid stones

d) Ammonium urate stones

Question 6:

All of the following statements are true regarding Wilm's tumour except?

a) It presents most commonly as abdominal mass

b) Most common age of presentation is within 5 years

c) 95% of patients live long term with radiotherapy and chemotherapy

d) Metastasis to lungs is rare and occurs late

Question 7:

What is the colour code in trauma triage for patients with life threatening conditions?

a) Red

b) Yellow

c) Green

d) Black

Question 8:

Which of the following conditions is not a risk factor for the development of cholangiocarcinoma?

a) Hepatitis B

b) Caroli disesase

c) Primary sclerosing cholangitis

d) Infection with Opisthorchis viverrini

Question 9:

A woman presented with ulcerative lesion over the breast associated with erythema of surrounding skin and history of retraction of nipple. Biopsy revealed invasive ductal carcinoma. According to 8th edition of AJCC classification, the patient can be classified under which stage?

a) Stage cant be M1

b) Stage IIIb

c) Stage 4, if N1-N3

d) T4b

Question 10:

A 35-year-old male had an episode of loss of consciousness subsequent to a head injury from a bike accident. Upon awakening, he began experiencing severe headache and nausea. He appears drowsy with a Glasgow Coma Scale (GCS) score of 10. One of his pupils is dilated and unresponsive. CT head given. What is the management?



a) Anticoagulants to prevent further clotting of blood

b) Observe and serial monitoring in the ICU

c) Osmotic diuresis to decrease intracranial pressure

d) Urgent surgical evacuation

Question 11:

A patient with papillary carcinoma thyroid underwent thyroidectomy. She developed hoarseness of voice after surgery. Which of the following steps in surgery could not have resulted in this complication?

a) Division of strap muscles

b) Dissection close to Zuckerkandl tubercle

c) Dissection along triangle of Beahr

d) Ligation of superior thyroid vessels

Question 12:

Which is the most common hernia in females?

a) Indirect inguinal

b) Direct inguinal

c) Femoral

d) Obturator

Question 13:

Which of the following is not done during the primary survey in trauma management?

a) 1, 3 and 4

b) 3 and 4

c) 1 only

d) 1 and 3

Question 14:

A 45-year-old male arrived at the ER with altered mental status and hematemesis that began a day ago. He has a 20-year history of alcoholism, with his most recent binge occurring a week prior. An ultrasound revealed altered echo texture of liver and mild portal vein dilation. CT scan brain was normal. He was started on intravenous pantoprazole and antiemetics. What should be the next step in his management?

a) Start a course of broad spectrum antibiotics

b) Order a liver biopsy to asses for fibrosis

c) Perform an endoscopy to look for varices

d) Administer intravenous fluids and electrolyte replacement

Question 15:

A 23-year-old female presented with complaints of neck swelling, difficulty swallowing, and hoarseness of voice. Radiological evaluation revealed a mass compressing the trachea. Biopsy showed cells arranged in nests with amyloid deposits. What is the origin of these cells?

a) Follicular cells

b) Parafollicular cells

c) Hurthle cells

d) Ashkanazy cells

Question 16:

A 34-year-old male, on investigation serum calcium level of 12.8 mg/dl (N- 8.5-10.25mg/dl) with raised PTH. On ultrasonography, a 1.5x1.5 mass was detected on the right lower lobe of

the thyroid gland. Which of the following is true regarding this condition?

a) A normal DEXA scan rules out the need for surgery

b) MIBI followed by focussed parathyroidectomy to be done

c) No surgery is required, only surveillance

d) Creatinine clearance >80ml/min mandates surgery

Question 17:

A middle-aged woman presents with flank pain and painful micturition. The pain was relieved by IV analgesics given at a nearby primary healthcare centre. The patient is diagnosed with ureteric colic due to ureteric calculi. What is the best diagnostic modality in this case?

a) NCCT KUB

b) CECT

c) USG KUB

d) Ureteroscopy

Question 18:

Identify the marked structure.



a) Corpora amylacea

b) Corpora arenacea

c) Lewy body

d) Psammoma bodies

Question 19:

A 25-year-old patient sustains a fall from the 2nd floor of a building and is brought to the emergency department. Which of the following is not an adjunct to the primary survey?

a) X-ray pelvis AP

b) X-ray chest AP

c) Extended FAST

d) CT abdomen and pelvis

Question 20:

A 20-year-old male presents with acute epigastric pain and fever. He has a past history of cholelithiasis. On examination, abdominal guarding and rigidity present. His serum amylase, lipase are raised and patient has leucocytosis. What is true regarding his condition?

a) Serum amylase remains elevated even after 30 days of resolution of acute pancreatitis

b) Serum lipase is more specific than serum amylase

c) > 75 percent patients present with hypercalcemia

d) Increase in the level of serum lipase is directly proportional to the severity

Question 21:

A 34-year-old female presents with a lump in her left breast for 2 years. On examination, a 6 x 4 cm mass is palpated. Dimpling of the skin and bloody discharge from the nipple is present. The lesion is invading the pectoralis major muscle. Ipsilateral Level I and Level II matted axillary lymph nodes present. There was no distant metastasis. What is the TNM staging (AJCC 8th)?

a) T4bN1M0

b) T3N2M0

c) T4aN1M0

d) T2N1M1

Question 22:

An 84-year-old man with a history of stroke 2 years ago, who is on prolonged bed rest, develops a pressure ulcer with partial thickness skin loss and a clean base. What is the grade of his pressure sore?

a) Grade 1

b) Grade 4

c) Grade 3

d) Grade 2

Question 23:

During ward rounds, you see a patient who is admitted with a ca esophagus. What is the most likely presentation in this patient?

a) Dysphagia of solids followed by liquids

b) Dysphagia of liquids followed by solids

c) Chest pain

d) Neck mass

Question 24:

What is the T stage of this oral cavity tumour?

a) T4a

b) T4b

c) T3

d) T2

Question 25:

A 30-year-old patient with peptic ulcer disease presents with abdominal pain that is not responding despite treatment in multiple hospitals. He was found to have ulcers in the jejunum and ileum. Gastric pH < 2, and gastrin levels after adequate cessation of PPIs are 10 times the normal value. Identify the lesion present and the boundaries of the triangle containing it.

a) Gastrinoma; triangle formed by common bile duct, junction of D2-D3, neck and body of pancreas

b) Insulinoma; triangle formed by cystic duct, junction of D2-D3, neck and body of pancreas

c) Gastrinoma; triangle formed by cystic duct, junction of D2-D3, neck and body of pancreas

d) Insulinoma; triangle formed by common bile duct, junction of D2-D3, neck and body of pancreas

Question 26:

An intern is assisting with laparoscopic surgery, and the resident is orienting the intern to the instruments and the operating ports. Your surgeon asks you what the angle between the camera port and the working port is.

a) Azimuth's angle

b) Manipulation angle

c) Crossing angle

d) Elevation angle

Question 27:

A 35-year-old male underwent surgery for an ileal perforation. Due to heavy contamination, only the sheath was closed, and the wound was left open. Daily wound washing was performed, and the wound was closed on postoperative day 5. By which mechanism does wound healing occur in this case?

a) Primary intention

b) Secondary intention

c) Tertiary intention

d) Quarternary intention

Question 28:

A 40-year-old patient who is a known case of Hodgkin's lymphoma with a previous history of irradiation presents with right-sided thyroid swelling. FNAC reveals papillary carcinoma of the thyroid and there is no lymph node involvement. What is the next step in management?

a) Hemi-thyroidectomy

b) Hemi-thyroidectomy with modified radical neck dissection

c) Total thyroidectomy

d) Total thyroidectomy with modified radical neck dissection

Question 29:

One of your friends is teaching you about enhanced recovery after surgery. Which of the following he states is incorrect?

a) Prolonged fasting is necessary

b) Epidural anaesthesia is recommended in laparoscopic surgery

c) Reduction of the hospital stay by 2-3 weeks

d) Beta-blockers are given after major surgery done under general anaesthesia to improve outcome

Question 30:

A chronic smoker comes with complaints of pain in his legs after walking 100 meters. On examination, he has an absent pulse in the bilateral dorsalis pedis, anterior tibial, and posterior tibial arteries, but no gangrene is present. Intermittent claudication is implicated. Which of the following statements is true regarding this condition?

a) Pain is more at night and sleep

b) Development of pain on the 1st step of walking

c) Pain aggravated by standing and relieved by bending forward

d) Pain develops after walking some distance and is relieved by rest

Question 31:

A 25-year-old male with a history of a road traffic accident is brought to the emergency department. His GCS is 8/15, oxygen saturation is 85%, BP is 100/ 60 mmHg, and heart rate is 60 beats per minute. On auscultation his right side breath sounds were absent. What is the next step in management?

a) Secure airway, give oxygen, insert ICD, secure IV line

b) Administer IV dextrose followed by NCCT head

c) Start IV fluids, give atropine and secure airway

d) Evaluate for head injury and do CT-head

Question 32:

An elderly male patient comes with foul-smelling breath and dysphagia. A barium swallow X-ray is attached. What is the most likely diagnosis?



a) Achalasia cardia

b) Zenker's diverticulum

c) Tracheoesophageal fistula

d) Carcinoma esophagus

Question 33:

Truelove and Witts classification is used for?

a) Acute encephalitis

b) Acute pancreatitis

c) ARDS

d) Acute severe ulcerative colitis

Answer Key

Question No.	Correct Option
1	c
2	a
3	c
4	b
5	a
6	c
7	a
8	a
9	b
10	d
11	a
12	a
13	c
14	d
15	b
16	b
17	a
18	a
19	d
20	b
21	b
22	d
23	a
24	a
25	c
26	a
27	c

28	c
29	a
30	d
31	a
32	b
33	d

Detailed Explanations

Solution to Question 1:

Asthma, laryngitis, pulmonary fibrosis, and cough are all extra esophageal complications in gastroesophageal reflux disease(GERD)

Extraoesophageal symptoms of GERD are:

• Chronic cough

• Laryngitis,

• Asthma

• Dental erosions (especially on the lingual and palatal tooth

surfaces).

• Pulmonary fibrosis

• Sinusitis

• Pharyngitis

• Recurrent otitis media

Extraesophageal symptoms of GERD originate from the respiratory tract and include both laryngeal and pulmonary issues These symptoms can arise through two mechanisms. The first and most common mechanism involves proximal esophageal reflux and microaspiration of gastroduodenal contents, which cause direct injury to the larynx and lower respiratory tract. The second mechanism involves distal esophageal acid exposure triggering a vagal nerve reflex, leading to bronchospasm and cough, due to the shared vagal innervation of the trachea and esophagus.

The incidence of Barrett's esophagus is also more in patients with long term GERD, which in turn increases the risk of adenocarcinoma of esophagus.

Solution to Question 2:

The given complaints of difficulty in swallowing, weight loss, regurgitation of undigested food and halitosis along with the barium swallow showing bird beak appearance is suggestive of achalasia

cardia.

Achalasia is a primary esophageal motility disorder caused due to the degeneration of neurons in the myenteric plexus of the esophagus. It is characterized by the absence of esophageal peristalsis and impaired relaxation of the lower esophageal sphincter (LES), thereby causing a functional obstruction at the gastroesophageal junction.

The patients present with a triad of dysphagia, regurgitation, and weight loss. The dysphagia initially begins with liquid and then progresses to solid foods, halitosis is present due to the accumulation of food. Patients may develop pneumonia, lung abscess, etc. at a later stage.

The initial diagnosis is made by barium swallow or endoscopy, which shows the presence of dilated esophagus due to the constriction of the lower esophageal sphincter (bird beak appearance). The gold standard test to confirm achalasia cardia is esophageal manometry. The findings are impaired lower esophageal sphincter (LES) relaxation and absent peristalsis with elevated LES pressure.

Management includes:

• Surgical esophagomyotomy.

• Non-surgical measures like sublingual nitroglycerin, nitrates, or calcium channel blockers.

• Injection of botulinum toxin

• Peroral endoscopic myotomy (POEM)

Solution to Question 3:

The instruments shown in the image is used to perform percutaneous endoscopic gastrostomy.

A standard endoscopy is performed to ensure that there are no contraindications to gastrostomy insertion. The stomach is insufflated with air and direct percutaneous needle puncture made at a point where the stomach abuts the abdominal wall. Lignocaine is infused on withdrawal. A trocar is inserted and a wire is passed into the stomach, which can be caught with a snare. The scope is withdrawn, pulling the wire out through the mouth,at which point it is attached to the gastrostomy tube. The gastrostomy is pulled through into the stomach and out through the track created by the trocar insertion.

The insertion of a percutaneous endoscopic gastrostomy (PEG) tube improves nutritional and functional outcomes in patients who cannot maintain oral nutritional intake. PEG insertion is often a preliminary step in the treatment of complex orofacial malignancies and can also be used to support nutrition in patients with various malignant, degenerative, or inflammatory diseases.

Procedure risks include bleeding, perforation,aspiration risk, dislodgment with peritoneal contamination, wound site infection and granulation.

Solution to Question 4:

Estrogen receptor (ER) is used as a prognostic marker.

The term prognostic marker generally is used to describe molecular markers that predict disease-free survival, disease- specific survival, and overall survival, whereas the term predictive marker often is used in the context of predicting response to certain therapies.

Other options

Option A: A diagnostic biomarker detects or confirms the presence of a disease or condition of interest, or identifies an individual with a subtype of the disease.

Option C:When a biomarker can be measured serially to assess the status of a disease or medical condition for evidence of exposure to a medical product or environmental agent, or to detect an effect of a medical product or biological agent, it is a monitoring biomarker

Option D: The most common and serious error in evaluating biomarkers is assuming that a correlation between a biomarker's level and a clinical outcome means the biomarker is a valid surrogate. However, for a biomarker to be a valid surrogate, it must not only be correlated with the outcome but also explain the change in the clinical outcome.

Solution to Question 5:

The given clinical scenario and microscopic image showing coffin lid shaped crystals are suggestive of triple phosphate stones. They are also known as staghorn or struvite stones.

This is caused due to infection with urea-splitting organisms, such as Proteus, Klebsiella, Serratia, or Enterobacter species.They are usually formed in alkaline urine.

Staghorn calculi are radio-opaque stones that grow in a branching pattern along the pelvicalyceal system. The above organisms metabolize the urea into ammonium and bicarbonate. Increased metabolism of urea results in alkalinity of the urine, and precipitation of magnesium, ammonium, and phosphate - resulting in the formation of struvite stones.

Complete removal of a staghorn calculus is essential, as any remaining fragments after treatment can lead to a rapid recurrence and continued presence of bacteriuria. Following the successful removal of the infection stone, long-term chemoprophylaxis is required for several months.

Solution to Question 6:

Option C is not true regarding Wilms tumour. The standard treatment of Wilm's tumour involves a combination of chemotherapy and surgery.

Wilms tumor is the most common primary malignant renal tumor in children, typically affecting children less than 5 years (OptionB). After neuroblastoma, it is the second most common malignant abdominal tumor in childhood. It most commonly presents as an asymptomatic abdominal mass. (Option A) Other findings include hypertension, abdominal pain, gross painless hematuria, and constitutional symptoms such as fever and weight loss.

CECT abdomen is used to identify the extent of the tumor, integrity of the other kidney, and metastasis. Wilms tumor is metabolically active and can concentrate fluorodeoxyglucose therefore PET/CT is also helpful to identify regional spread and metastatic lesions. A biopsy is discouraged as it may lead to disease upstaging.

Treatment: Surgery and chemotherapy, with or without radiotherapy, are the mainstays of treatment.

• Radical nephrectomy is the standard surgical procedure done. Routine sampling of the para-aortic, hilar, and paracaval lymph nodes should be done.

• Nephron-sparing nephrectomy is done for children with bilateral disease or with a solitary kidney. Preoperative chemotherapy can be used in these cases to induce tumor shrinkage.

• The classical chemotherapy includes vincristine and dactinomycin with the addition of Adriamycin(Doxorubicin).

Despite certain adverse risk factors such as metastases, unfavorable histology and recurrent disease, the prognosis for most children with Wilms' tumor remains very positive. The overall survival rate for children with Wilms' tumor is over 90%.

The tumour can extend into the renal vein and vena cava and metastasises to lymph nodes and lung. But metastasis to lungs occur late in the disease. (Option D)

Solution to Question 7:

Red is the colour code for patients with life threatening conditions who need immediate medical care.

Triage includes rapid classification of the injured based on their injuries and the chances of survival with prompt medical intervention. It provides maximum benefit to the greatest number of injured in a disaster situation. It lessens the burden on medical facilities.

The internationally accepted four-color coded system is used commonly. It includes:

• Red - High priority attention or transfer; Critical, but likely to survive if treatment is given early

• Yellow - Medium priority; Critical, likely to survive if treatment is given within hours

• Green - Ambulatory patients; Stable, likely to survive even if treatment is delayed for hours to days

• Black - Dead or moribund patients; Not breathing, pulseless, so severely injured that no medical care is likely to help

Solution to Question 8:

While all the given options are associated with the causation of cholangiocarcinoma, hepatitis B in particular has an association with intrahepatic cholangiocarcinoma and rarely extrahepatic while all the other options are associated with both intra and extrahepatic cholangiocarcinoma. Hence it is not as significantly associated with cholangiocarcinoma compared to the other options

Cholangiocellular carcinomas (CCCs) or cholangiocarcinomas are mucin-producing adenocarcinomas of the biliary duct system that may originate in the liver and extrahepatic bile ducts. Patients present with signs of obstructive jaundice associated with weight loss and anorexia. Initial workup by ultrasound shows dilation of the intrahepatic biliary tree, but

definitive diagnosis requires visualisation of the biliary tree and MRCP or ERCP (biopsy possible) is preferred

Risk factors for cholangiocarcinoma:

• Chronic inflammatory conditions- PSC, Oriental cholangiohepatitis,Hepatitis C infection

• Parasitic infections- Opisthorchis viverrini, C. sinensis

• Congenital - Choledochal cysts, Caroli's disease

• Chemical agents- Thorium dioxide (Thorotrast), Vinyl chloride, Dioxin, Asbestos

Solution to Question 9:

The patient presenting with ulcerative lesion with inflammatory signs can be classified as Stage IIIb. Inflammatory breast carcinoma (stage T4d) is characterized by the skin changes of brawny induration, erythema with a raised edge, and edema (peaud'orange) involving >1/3rd of the breast.

T4d in the TNM staging of carcinoma breast points toward a diagnosis of inflammatory breast carcinoma. It shows extensive invasion and proliferation of tumor cells. Inflammatory breast cancer is the most aggressive subtype of breast cancer. The hallmark of this condition is the overexpression of e-cadherin. This results in increased tumor cohesion and the formation of tumoral clumps which can result in emboli. It has a particularly poor prognosis.

In addition to anatomical staging, the eighth edition of the AJCC TNM staging system includes the histological grade, ER, PR, HER2/neu and Ki-67 assessment, multigene testingwith Oncotype DXand the response to neoadjuvant chemotherapy to refine the prognostic information.

Solution to Question 10:

The given scenario of a man with a head injury, unconsciousness, nausea, dilated and fixed pupils, and the CT finding of a lenticular hemorrhage also showing a mid-line shift is characteristic of an extradural or epidural hemorrhage. Urgent surgical evacuation of heamatoma by craniotomy is the management.

Epidural hemorrhage/hematoma is an intracranial hemorrhage in which the hemorrhage is between the dura and the skull. It classically occurs due to a tear of the middle meningeal artery, following a fracture of the temporal bone. Management is by surgical evacuation of the hematoma or by ligation of the damaged middle meningeal artery

Clinically, patients with extradural hemorrhage typically present with an initial transient loss of consciousness. A lucid interval is then seen, which may be minutes to hours during which the patient usually presents. Neurological deterioration then occurs following the lucid interval as the hematoma expands.

As the hematoma expands, compensation is exhausted with rapid deterioration. There is contralateral hemiparesis, a reduced conscious level, and ipsilateral pupillary dilatation, the cardinal signs of brain compression and herniation. The prognosis for promptly evacuated

extradural hematoma, without associated primary brain injury is excellent.

Solution to Question 11:

The patient presenting with hoarseness of voice post thyroidectomy can be due to injury to external laryngeal nerve or unilateral recurrent laryngeal nerve. The step in thyroidectomy where it is least to happen is during division of strap muscles.

Recurrent laryngeal nerve is related to the posterior branch of the inferior thyroid artery near the lobes of the thyroid gland. Therefore, during surgery, Inferior thyroid artery is ligated close to the gland to avoid injury to the recurrent laryngeal nerve.

Thyroid surgery may cause injury to both external and recurrent laryngeal nerves (RLN). During thyroidectomy surgeries, there might be unilateral or bilateral recurrent laryngeal nerve injury. Unilateral RLN injury may be asymptomatic but in a few may cause hoarseness of voice.

Bilateral recurrent laryngeal nerve palsy can also occur postoperatively. It causes dyspnea and stridor but the voice is good. These cases required emergency tracheostomy. External laryngeal nerve injury is more common and could result in hoarseness of voice.

Other options

Option B: The RLN frequently passes immediately medial to the tubercle of Zuckerkandl. In order to find and protect RLN during thyroid surgery, a careful, bloodless, and meticulous dissection should be carried out around the tubercle.

Option C: Beahrs' triangle is bounded by common carotid artery as base, inferior thyroid artery superiorly and RLN as the lower arm of triangle. Recognizing the boundaries of the triangle posterior to thyroid after retracting the ipsilateral thyroid lobe medially, helps in avoiding RLN injury.

Option D: The external branch of the superior laryngeal nerve courses along the cricothyroid muscle just medial to the superior pole vessels. To avoid injury to this nerve, the superior pole vessels are divided individually as close as possible to the point where they enter the thyroid gland

Solution to Question 12:

The most common hernia in females is an indirect inguinal hernia.

Inguinal hernia is the most common hernia in men and women. It is classified into :

• Indirect hernia - protrusion of the contents occurs through the deep inguinal ring, along with the spermatic cord, lateral to the inferior epigastric artery. It is the most common subtype of groin hernia in both males and females.

• Direct inguinal hernia - Occurs through the posterior wall of the inguinal canal through the Hasselbach's triangle. The sac is medial to the inferior epigastric artery.

Other options:

Option C: Femoral hernia occurs when the hernial sac descends vertically through the femoral canal up to the saphenous opening and then escapes out into the loose areolar tissue. It is more common in women than in men; however, it is less common than inguinal hernia.

Option D: In an obturator hernia, the contents pass through the obturator canal. It is more commonly seen in females than males; however, it is less common than inguinal hernia.

Note: Inguinal hernias are five times more common than femoral hernias.

Solution to Question 13:

NCCT brain is not done during the primary survey in trauma management. (Option C)

Endotracheal intubation is part of airway management in the primary survey. Intercostal drainage for hemothorax or pneumothorax and chest X-ray especially AP view is part of breathing component assessment of the primary survey in trauma management.

NCCT brain is a part of the secondary survey of trauma management and is performed after initial resuscitation and stabilisation. A CT scan of the head is performed within 1 hour if the GCS is <14 at the initial assessment. Other indications include a tense fontanelle, suspicion of an open/depressed or base of skull fracture, abnormal pupillary response, abnormal posturing, and a focal neurological defect. A head CT scan should also be performed if there are three or more vomiting episodes, a witnessed loss of consciousness or amnesia for >5 minutes.

Solution to Question 14:

The next immediate step in managament of upper gastrointestinal bleeding is administration of intravenous fluids and electrolyte replacement.

USG findings and history of alcoholism are suggestive of cirrhosis leading to portal hypertension. The upper GI endoscopy provides a definitive diagnosis and aids in predicting the risk of bleeding with an option to treat via band ligation or sclerotherapy. (Option C)

The management of esophageal variceal bleeding involves:

• Resuscitation with IV fluids, airway protection(intubation to protect airway and prevent aspiration), and oxygen

• Blood transfusion to maintain Hb between 7-9g/dL

• Antibiotic prophylaxis with ceftriaxone before endoscopy (Option A)

• An attempt at hemostasis with octreotide or terlipressin or vasopressin

• Upper GI endoscopy within 12 hours after fluid resuscitation to confirm the source of the bleed

• Endoscopic variceal band ligation or sclerotherapy

• In case, of failure or rebleeding - Transjugular intrahepatic portosystemic shunt (TIPS)



Solution to Question 15:

The clinical scenario along with biopsy showing cells arranged in nests along with amyloid deposits is suggestive of medullary carcinoma of the thyroid. Medullary carcinoma of the thyroid (MTC) arises from the parafollicular C cells.

Histopathology shows polygonal to spindle-shaped cells which form nests, trabeculae, or follicles. Amyloid deposits that are derived from the altered calcitonin molecules are present in the stroma. Calcitonin in the tumor cells and stroma is easily demonstratable by immunohistochemistry.

Features of MTC:

• Painful neck mass with cervical lymphadenopathy

• Dysphagia, dysphonia, or dyspnea if there is local invasion

• Diarrhea due to increased intestinal motility

• Cushingoid habitus due to ectopic ACTH production

• Blood-borne metastasis

Diagnosis is by careful family history, physical examination, raised calcitonin and CEA levels, and fine-needle aspiration biopsy of the mass.

Other options:

Option C and D: A Hurthle cell is a cell in the thyroid that is often associated with Hashimoto's thyroiditis as well as benign and malignant tumors (Hürthle cell adenoma and Hürthle cell carcinoma). Ashkanazy cells are another name given to the cells with Hurthle bodies in Hashimoto's thyroiditis.



Solution to Question 16:

The lab reports of this patient, associated with a thyroid mass indicate primary hyperparathyroidism due to parathyroid adenoma. MIBI followed by focused parathyroidectomy to be done in this condition.

The patient presents with hypercalcemia (serum calcium 12.8 mg/dL) and elevated parathyroid hormone (PTH), which strongly indicates primary hyperparathyroidism (PHPT).

The ultrasonographic finding of a mass in the right lower lobe of the thyroid is consistent with a parathyroid adenoma, the most common cause of PHPT.

99mTc-Sestamibi (MIBI) scan is commonly used in conjunction with ultrasonography to localise the hyperfunctioning parathyroid gland before surgery.

Focused parathyroidectomy is the treatment of choice in such cases when a single adenoma is identified. This approach is minimally invasive and guided by the imaging findings.

Other options

Option A: A normal DEXA scan does not negate the need for surgery in a patient with severe hypercalcemia like this. Severe hypercalcemia (>12 mg/dL) is an absolute indication for surgery, irrespective of the results of the DEXA scan.

Option C: Observation is only appropriate for patients with mild hypercalcemia, no symptoms, and no target organ damage. Severe hypercalcemia (>12 mg/dL) is an absolute indication for surgery.

Option D: Creatinine clearance < 60 ml/min mandates surgery.

Solution to Question 17:

NCCT KUB (Non-contrast CT of the kidney, ureters, and bladder) is the best diagnostic modality for the clinical evaluation of ureteric stones.

NCCT is the preferred investigation for evaluation of:

• Renal stones except for indinavir stones

• Ureteric stones

• Head injury

• Salivary gland stones

Solution to Question 18:

The marked structure represents Corpora amylacea.

Corpora Amylacea are small, rounded, eosinophilic structures composed of glycoproteins and carbohydrates, commonly found in glandular tissues such as:

• Prostate: Seen in benign prostatic hyperplasia (BPH), within prostatic acini.

• Astrocytic processes of the CNS, associated with aging or neurodegenerative diseases.

Clinical Significance:

• Generally benign and age-related in both the prostate and CNS.

• In CNS, linked to diseases like multiple sclerosis or Alzheimer's.

Other Options:

Option B: Corpora arenacea: The pineal gland has a structure called the Corpora arenacea, located within the gland. These bodies are prone to an increase in calcification with advancing age. They are visible on x-ray and can be used as landmarks.



Option C: Lewy body: aggregates of insoluble fibrillar proteins called alpha-synuclein (specific) and ubiquitin in the neural cells. They are typically seen in Parkinson's disease, Lewy body

dementia, and a few variants of Alzheimer's disease.

Option D: Psammoma bodies : calcified structures seen in areas of cellular degeneration. They have a concentric laminated, or onion-like, appearance. Diagnostic feature of papillary thyroid carcinoma (PTC), the most common form of thyroid cancer.



Solution to Question 19:

CT abdomen and pelvis are not an adjunct to the primary survey. CT scan is a part of a secondary survey.

The primary survey identifies and manages the most immediately life-threatening pathologies first.

The secondary survey includes investigations that help identify all other injuries, excluding the apparent ones. This includes X-rays of the spine and extremities(limbs) and a CT of the abdomen, head, chest, pelvis and spine.

Solution to Question 20:

The patient presents with features suggestive of acute pancreatitis (epigastric pain, fever, history of cholelithiasis, guarding/rigidity, raised serum amylase and lipase, and leukocytosis). For acute pancreatitis, serum lipase is more specific than serum amylase.

Lipase has a higher specificity for diagnosing acute pancreatitis, as elevated amylase can occur in other conditions such as salivary gland disorders, intestinal obstruction, or renal failure. Lipase also remains elevated longer than amylase.

The diagnostic criteria for acute pancreatitis require at least two of the following three features:

• Characteristic Abdominal Pain:

Acute onset of severe epigastric pain, often radiating to the back.

Pain may worsen after eating and is associated with nausea and vomiting.

• Serum Lipase and/or Amylase Elevation:

At least 3 times the upper limit of normal (ULN).

Lipase is preferred due to its higher specificity and longer half-life.

• Imaging Findings:

Characteristic findings of pancreatitis on imaging (CT scan, MRI, or ultrasound), such as:

Pancreatic enlargement

Peripancreatic fluid collections

Inflammatory changes in and around the pancreas

Other Options:

Option A: Amylase levels normalize within 3-5 days after an episode of acute pancreatitis.

Option C: Hypocalcemia is more commonly observed, due to fat saponification in severe cases.

Option D: The degree of elevation of serum lipase does not correlate with the severity of acute pancreatitis. Severity is assessed using clinical scores like Ranson's criteria, APACHE II, or CT severity index.



Solution to Question 21:

TNM staging of the tumour is T3 N2 M0.

In the given scenario:

• 6 x 4 cm mass - T3

• Ipsilateral Level I and Level II matted axillary lymph nodes- N2

• No distant metastasis - M0.

In tumour size assessment, the greatest dimension is taken into consideration. For example, in a 6 x 4 cm lump, 6 cm is taken into account, and the tumour is classified as T3 (>5 cm). If 2 lumps are present in the same breast, staging is according to the bigger lump. If lumps are present in bilateral breasts, separate staging is done for both lumps.

Any contralateral lymph node involvement is considered as distant metastasis.

Solution to Question 22:

The pressure sore is of grade 2 in this case.

Grade 2 of bedsore corresponds to partial-thickness skin loss involving the epidermis and dermis.

Pressure sores, also known as bedsores, pressure ulcers, or decubitus ulcers, are defined as tissue necrosis with ulceration due to prolonged pressure. When the external pressure exceeds the capillary occlusive pressure, blood flow ceases leading to tissue anoxia, necrosis, and ulceration. An external pressure of >30 mmHg leads to the formation of pressure sores.

The common sites of pressure sores in decreasing frequency are the ischium, greater trochanter, sacrum, heel, malleolus (lateral followed by medial), and occiput.

Given below are images showing the stages of pressure sore.

Prevention is the best treatment for pressure sores. They can be prevented with good skincare, the use of air-fluidized beds, special pressure-dispersion cushions, or foams. Fecal or urinary diversion can be considered in special cases.

The position of the bedridden patients should be changed once every 2 hours and wheel-chair bound patients should be taught to get off their chairs for 10 seconds every 10 minutes.

Surgical management of pressure sores follows the same principles involved in acute wound treatment. Prior to the operation, the pressure sores should be adequately debrided, and vacuum-assisted closure (VAC) can be used to promote the growth of granulation tissue required

for surgical closure. Large skin flaps that have muscle with sensory innervation are best for reconstruction.

VAC is done by applying intermittent negative pressure of approximately -125 mmHg to hasten debridement and promote granulation tissue formation. Negative pressure decreases edema, removes interstitial fluid, and increases blood flow. This decreases bacterial growth and increases cell proliferation.

Solution to Question 23:

Dysphagia of solids followed by liquids is the most likely presentation in a patient with ca esophagus.

Initially, patients have difficulty swallowing solids. Later, as the tumor grows and narrows the esophageal lumen, dysphagia progresses to liquids. This progression occurs because solid foods require a wider lumen to pass through, and the narrowing caused by the tumor becomes more critical as it advances.

Carcinoma esophagus is one of the most common malignancies manifesting in mid to late adulthood. It can be a squamous cell or adenocarcinoma.

Squamous cell carcinoma (SCC) affects the upper two-thirds of the esophagus while adenocarcinoma affects the lower third. Tobacco smoking and alcohol are the risk factors for SCC. Obesity and gastroesophageal reflux disease increase the risk of adenocarcinoma.

Dysphagia can be the most common presenting feature but may present late. Other features such as regurgitation, weight loss, vomiting, and odynophagia can be seen. Lymph node involvement indicates a poor prognosis.

Endoscopy is the first line of investigation done. CT is useful to rule out metastasis. Endoscopic ultrasonography is used to assess the depth of tumor invasion through the wall of the esophagus, adjacent organs, and lymph node metastasis.

Management options include radical esophagectomy, neoadjuvant chemotherapy, and palliative chemoradiation or endoscopic treatments.

Other options

Option B: Dysphagia of liquids followed by solids is typically seen in motility disorders, such as achalasia cardia, where there is functional obstruction rather than a mechanical obstruction.

Option C: Chest pain may occur in the CA esophagus, but it is not the most common presenting symptom. It usually arises in advanced cases or due to tumor invasion into surrounding structures.

Option D: A neck mass might occur in metastatic disease (e.g., supraclavicular lymphadenopathy), but not the typical initial presentation.

Solution to Question 24:

This oral cavity tumour is stage T4a.

In the given image, the oral cancer involves the floor of the mouth as well as the lip (adjacent structures). When adjacent structures are involved, the cancer meets the criteria for T4a disease. (resectable)

T4a is defined as "moderately advanced local disease" and includes tumors that invade floor of the mouth or the lip, and is resectable. These structures involve:

• Cortical bone (e.g., mandible or maxilla)

• Inferior alveolar nerve

• Floor of the mouth

• Skin of the face (chin or nose)

The T staging for oral cavity tumors, based on the AJCC 8th edition, is as follows:

• T1: Tumor ≤ 2 cm in greatest dimension, no invasion of adjacent structures.

• T2: Tumor > 2 cm but ≤ 4 cm in greatest dimension, no invasion of adjacent structures. (Option A)

• T3: Tumor > 4 cm in greatest dimension or minor invasion into structures such as the deep tongue muscle. (Option C)

• T4a (Moderately advanced local disease): Tumor invades adjacent structures like cortical bone, inferior alveolar nerve, floor of the mouth, or skin (chin or nose).

• T4b (Very advanced local disease): Tumor invades masticator space, pterygoid plates, skull base, or encases the internal carotid artery. (Option B)

The gold standard investigation for oral cancer includes an incisional biopsy. An MRI is the imaging modality of choice for assessing primary extent. Surgical management with chemotherapy and adjuvant radiotherapy is the mainstay of management.

Solution to Question 25:

The above clinical scenario of abdominal pain, ulcers in the jejunum and ileum, gastric pH < 2, and elevated gastrin levels are suggestive of gastrinoma. It is commonly located in a triangle formed by the cystic duct, junction of D2-D3, neck and body of pancreas.

Gastrinoma is a neuroendocrinal tumour characterized by gastrin hypersecretion, resulting in gastrointestinal tract ulcers. It occurs more commonly in a sporadic pattern but can be associated with MEN-1 (multiple endocrine neoplasia-1) syndromes in the familial pattern.

Gastrinoma is most commonly located in the gastrinoma triangle (Passaro's triangle) bounded by the lines connecting the:

• cystic duct

• the junction between the second and third parts of the duodenum

• the junction between the neck and the body of the pancreas



The most common presentations of gastrinoma are epigastric pain, diarrhoea, and peptic ulcers. The ulcers occur at typical locations (proximal duodenum) and atypical locations (distal duodenum, jejunum). Duodenal ulcers are the most common. Gastrinoma if associated with marked gastric acid production and recurrent peptic ulcer is called Zollinger-Ellison syndrome.

Diagnosis:

The gastric pH <2, basal acid output >15 mEq/hr and serum gastrin concentration more than 1000 pg/ml (normal is <100 pg/ml) confirm the diagnosis of gastrinoma. When the diagnosis is in doubt (like gastrin levels 100-1000 pg/ml), then a secretin test is performed. A secretin test is considered positive when serum gastrin levels are more than 200 pg/ml on secretin administration and confirms the diagnosis of gastrinoma.

For localization of the tumour for surgical management, CT or MRI scans should be done first. If the tumour is not localized on these scans, then somatostatin receptor scintigraphy or endoscopic ultrasound is done for better localization.

Solution to Question 26:

The angle between the camera port and the working port is known as Azimuth's angle. The angle must be a minimum of 30° to avoid any clashing between these instruments.



Other options

Option B: Manipulation angle is the angle between two working ports. The minimum angle should be 60°.

Option C: Crossing angle refers to the intersection of instruments inside the abdominal cavity.

Option D: The elevation angle is the angle between the working port and the horizontal line. The minimum angle must be 60° to avoid colliding the instrument with the abdominal wall.

Solution to Question 27:

Wound healing occurs by tertiary intention in this case. The wound is cleaned and left open for a few days. This allows soft tissues to drain and also ensures no infection is present before surgical closure can be done.

Wound closure is categorized into primary, secondary, or tertiary:

Primary closure is done for clean wounds with minimal tissue loss that are amenable to approximation of wound edges. Healing occurs by primary intention. Primary closure is ideally performed within 6 to 8 hrs of trauma, but wounds in vascular areas (e.g. face) may be delayed to up to 24 hours after trauma.

Secondary closure allows healing by secondary intention. The wound is left open to heal largely by the formation of granulation tissue and contraction. These wounds often have significant tissue loss precluding tension-free approximation of edges, with devitalized edges. Wound dressings are changed daily to aid in the formation of granulation tissue and subsequent contraction.

Tertiary closure, also known as delayed primary closure, results in a combination of healing by primary and secondary intention. The wound is thoroughly irrigated, debrided, and left open for 2 to 5 days before surgical closure. It is used for wounds that are grossly contaminated without significant tissue loss.

Solution to Question 28:

Total thyroidectomy is performed in patients with radiation-induced differentiated thyroid cancers irrespective of their size.

Management of differentiated thyroid cancers (DTC)

Differentiated thyroid cancers include papillary thyroid cancer, follicular thyroid cancer, and hurtle cell cancer.

Hemithyroidectomy is performed for DTCs between 1-4 cm without extrathyroidal extension or evidence of metastatic disease.

However, total thyroidectomy is performed in the following scenarios

• Tumor at least 4 cm

• Gross extrathyroidal extension

• Evidence of metastatic disease

• Radiation-induced DTC

• Familial nonmedullary thyroid cancer

• Multifocal bilateral DTC.

Central neck dissection is done only in case of involvement of level VI nodes.

If level VI plus other lymph nodes are enlarged - then total thyroidectomy + central neck dissection + modified radical neck dissection is done.

In case of tumor size >4cm (T3 &T4), prophylactic central neck dissection is done.

Below is the TNM staging of differentiated thyroid carcinoma.

Papillary carcinoma is the most common thyroid carcinoma. It is most common in iodine-sufficient areas, among children, and in those exposed to ionizing radiation. It is more common in women than men (2:1). Dysphagia, dysphonia, and dyspnea are associated with local invasion. It usually presents as a cold nodule on the scan.

Lymph node metastasis is common. Lateral aberrant thyroid denotes a cervical lymph node invaded by papillary carcinoma. The most common sites of metastasis: lung > liver > bone > brain. It has an excellent prognosis, with a 10-year survival rate in excess of 95%.

Primary Tumor (T)TX Primary cannot be assessedT0 No primary tumorT1a ≤1 cm, intrathyroidalT1b 1-2 cm, intrathyroidalT2 2-4 cm, intrathyroidalT3 >4 cmlimited to the thyroid or any size withminimal extrathyroidal extension(muscle, soft tissue)T3a- >4cm limited to the thyroidT3b- gross extrathyroidal extension involving only the strap muscle.T4a Invades subcutaneous tissue, larynx, trachea, esophagus, or RLN T4b Invades prevertebral fascia or encases carotid or mediastinal vessels	Regional Lymph Nodes (N)NX Nodes cannot be assessed N0 Negative nodesN1aPositive in levelsVI & VIIN1b Positive unilateral, bilateral, contralateral lateral neck lymph nodes.Distant Metastasis (M)M0 No distant metastasis M1 Distant metastasis

Solution to Question 29:

Prolonged fasting is not necessary for enhanced recovery after surgery.

Early oral hydration and nourishment and avoidance of unnecessary IV fluid administration favors the Enhanced Recovery After Surgery (ERAS) protocol.

ERAS is an approach designed to speed the clinical recovery of the patient and reduce the cost and length of stay in the patient. It is achieved by optimizing the patient's health before surgery and then delivering evidence-based best care in the perioperative period.

It includes:

• Preference of minimally invasive procedures (e.g. laparoscopic)over open procedures.

• Minimal perioperative starvation.

• Preference for local or regional anesthesia, and adequate pain control with patient-controlled analgesia.

• Early removal of Ryles tube and early feeding techniques, rather than IV fluid administration.

• Early mobilisation planned physiotherapy.

• Discharge planning is started before the patient is admitted to hospital and involves support from stoma care nurses, physiotherapists etc.

Other Options:

Option B: Epidural anaesthesia is recommended in laparoscopic surgery for pain management.

Option C: Reduction of the hospital stay by 2-3 weeks: ERAS protocols do reduce hospital stay so as to plan for early discharge.

Option D: Beta-blockers are given after major surgery done under general anaesthesia to improve outcome: Beta-blockers may be used postoperatively in certain cases to control heart rate, manage perioperative stress, and improve outcomes, especially in high-risk cardiac patients.

Solution to Question 30:

In intermittent claudication, pain develops after walking some distance and is relieved by rest.

Intermittent(vascular) claudication is most commonly felt in the calf muscles because of the involvement of the superficial femoral artery in chronic vascular occlusive disease.

Intermittent claudication is the debilitating cramp-like pain felt in the leg muscles due to reduced blood flow and anaerobic metabolism brought on by peripheral arterial disease. It usually signifies mild to moderate chronic vascular occlusive disease. The pain is brought on by walking, not present on taking the first step. Relieved by rest in both the standing and sitting positions, usually within 5 minutes (option A). The distance that a patient can walk without stopping is known as claudication distance. It varies only slightly from day to day. The superficial femoral artery (70%) is the most commonly affected artery by claudication and manifests as pain in the calf muscles. Aorto-iliac occlusion may cause thigh or buttock claudication. Leriche's syndrome occurs due to aortoiliac occlusion and presents with a triad of claudication, impotence, and absent femoral pulses.

Boyd's classification of claudication:

• Grade I: Patient complains of pain after walking. If the patient continues to walk, due to increased blood flow in muscle and opening of collaterals metabolites causing pain are washed away and the pain subsides.

• Grade II: Pain persists on continuing to walk; but can walk with effort.

• Grade III: The patient has to take rest to relieve the pain.

Other options

Option A: Rest pain occurs when the occlusive disease progresses in severity. Though this pain is present during the day, it is exacerbated on lying down at night due to loss of gravity-dependent perfusion of the foot. This may be relieved by hanging the foot out of the bed to restore the gravitational effects.

Option B: The development of pain on taking the first step of walking is a classic sign of osteoarthritis.

Option C: Pain aggravated by standing and relieved by bending forward is seen in neurogenic claudication. It may be seen in cases of nerve compression from lumbar intervertebral disc prolapse or spinal stenosis.

Solution to Question 31:

The given clinical scenario of hemodynamic instability, decreased oxygen saturation (85%), GCS of 8/15 and absent breath sounds on the right side of the chest, with a history of a road traffic accident are suggestive of tension pneumothorax. The next step of management in this case is to secure the airway, give oxygen, insert the ICD, and secure the IV line.

According to the latest ATLS guidelines, the order of primary survey in a trauma patient is ABCDEF following the principle of ‘what kills first should be managed first’ -

• Airway

• Breathing

• Circulation with control of haemorrhage

• Disability(neurological)

• Exposure to environmental control

• Fluid resuscitation

After securing the airway, breathing can be assessed by auscultating for breath sounds and a chest X-ray to look for a pneumothorax or hemothorax. Administering oxygen through a bag and mask is useful during desaturation only if the airway is patent. Therefore, securing the airway is the most important next step for this patient.

Tension pneumothorax is mainly managed by inserting a wide-bore needle in the 5th intercostal space (ICS) followed by placing an intercostal drain in the 5th ICS. Once the airway and breathing are secured, hemodynamic instability must be corrected by securing an IV line to maintain circulation.

Other investigations such as NCCT head and CT head are done later on during further evaluation in the secondary survey.

Solution to Question 32:

The given clinical scenario of foul-smelling breath and dysphagia, and the given barium swallow X-ray showing the collection of contrast in a posterior pharyngeal pouch-like structure, are suggestive of Zenker's diverticulum.

It often presents with the following:

• Dysphagia

• Regurgitation

• Halitosis

• Gurgling sound on swallowing

• Aspiration

A barium swallow study may show a diverticulum arising from the midline of the posterior wall of the distal pharynx near the pharyngoesophageal junction. The diverticulum is typically noted at the C5-6 level. The outpouching may be transient, sometimes referred to as a pharyngeal pouch. The pouch is best identified during swallowing and is best visualized on the lateral view.

Solution to Question 33:

Truelove and Witts classification is used for acute severe ulcerative colitis.

Given below is the Truelove and Witts severity index for ulcerative colitis.

	Mild	Moderate	Severe
Bowel movements	< 4 per day	4 – 6 per day	> 6 per day
Blood in stool	Small	Moderate	Severe
Fever	None	< 37.5°C	> 37.5°C
Tachycardia	None	< 90 mean pulse	>90 mean pulse
Anemia	Mild	Hb > 75% of normal Hb	Hb < 75% of normal Hb
ESR	< 30mm	< 30mm	> 30mm
Endoscopic appearance	ErythemaDecreased vascular markingsFine granularity	Marked erythemaAbsent vascular markingsCoarse granularityContact bleedingNo ulcerations	Spontaneous bleedingUlcerations

Surgery NEET 2024

Question 1:

A male presented with discharge as shown in the image. What is the diagnosis?



a) Fistula in ano

b) Fissure in ano

c) External Haemorrhoids

d) Pilonidal sinus

Question 2:

A man presented to the emergency department after RTA with complaints of chest pain and breathing difficulty. An X-ray was done and is given below. What type of breathing pattern is observed in this patient?



a) Cheyne stokes

b) Paradoxical

c) Biot

d) Kussmaul

Question 3:

A mother brought her 3-month-old male child to the hospital, who is currently unable to pass urine. A similar issue was reported 15 days ago. The catheter could be passed without difficulty. The image of the micturating cystourethrogram is given below. What is the most appropriate management for this condition?



a) Partial resection of verumontanum

b) Endoscopic dilation of bladder neck

c) Indwelling catheter

d) Destruction of valves

Question 4:

What is the condition seen here?

a) Gastric Ulcers

b) Oesophageal varices

c) Candidiasis

d) Barret's esophagus

Question 5:

The baby was born with this deformity. What is the earliest age to correct this defect?

a) 3 months

b) 1 year

c) 6 months

d) 18 months

Question 6:

A patient with a history of tuberculosis presents with dysuria and sterile pyuria. The cystoscopy image of the patient is provided below. Based on the patient's history and the cystoscopic findings, which of the following conditions is most likely associated?



a) Linear ulcer of the bladder

b) Golf hole ureter

c) Ureterocele

d) Urethral stricture

Question 7:

What instrument is required for the following procedure?



a) Ureteroscope

b) Cystourethroscope

c) Nephroscope

d) Ultrasound guided scope

Question 8:

A patient with a history of umbilical swelling, initially asymptomatic for many years, now complains of pain over the swelling. The swelling was reducible initially but is now non-reducible. What is the appropriate management?

a) Umbilectomy with herniorrhaphy

b) Umbilectomy

c) Conservative management with antibiotics

d) Incision and drainage

Question 9:

A baby was brought in with complaints of intermittent feculent discharge from the umbilicus. What is the likely diagnosis?



a) Umbilical polyp

b) Ileal diverticulum

c) Urachus

d) Urachal cyst

Question 10:

A patient presents with renal colic and recurrent urinary tract infections. A CT scan reveals no significant findings. The patient undergoes a percutaneous nephrolithotomy and his urine analysis is as shown. Based on the clinical presentation and the urine analysis findings, what are these crystals made up of?

a) Struvite

b) Oxalate

c) Uric acid

d) Cystine

Question 11:

What is the treatment plan for this condition?

a) Full thickness skin grafting

b) Split thickness skin grafting

c) Vaccum-assisted closure

d) Flap

Question 12:

A patient with oral cancer underwent radical surgery and PMMC flap reconstruction as shown. What is the most important indicator for considering adjuvant radiotherapy in this patient?

a) Close margin < 5 mm

b) Extra nodal extension

c) Multiple lymph nodes

d) Perineural invasion

Question 13:

Identify the stage of bed sore in the sacral region.



a) Stage 1

b) Stage 2

c) Stage 3

d) Stage 4

Question 14:

Based on the constituents, identify the fluid given to post-operative patients.



a) Haemaccel

b) Ringer lactate

c) Isolyte P

d) Isolyte M

Question 15:

A female patient presented with a breast mass, as shown in the image below, with concomitant liver metastasis. What is the recommended management plan for this patient?

a) Simple mastectomy

b) Neo adjuvant chemotherapy followed by surgery

c) Modified radical mastectomy

d) Radical mastectomy

Question 16:

An elderly male who presented with the following lesion as shown, is diagnosed with stage T3 carcinoma with no clinically palpable inguinal lymph nodes. What is the management strategy for this patient?

a) Penectomy alone

b) Penectomy + Orchidectomy

c) Penectomy + Inguinal lymph node dissection

d) Penectomy + Superficial inguinal lymph node dissection

Question 17:

The most commonly occurring complication of TPN in a sepsis patient in ICU:

a) Hyperlipidemia

b) Hypertriglyceridemia

c) Reactive Hypoglycemia

d) Hyperglycemia

Question 18:

A CT scan of a patient with a hydatid cyst is shown below. What is the type of hydatid cyst as per the Gharbi classification?

a) Type 4

b) Type 3

c) Type 2

d) Type 1

Question 19:

Identify the image:

a) Ileal diverticulum

b) Bladder exstrophy

c) Omphalocele

d) Gastroschisis

Question 20:

Identify the following:



a) Fluoroscopy

b) X-ray of alkali ingestion

c) Barium swallow

d) X-ray of acid ingestion

Question 21:

Identify pseudomembranous colitis from the given image.



a) 3

b) 4

c) 1

d) 2

Question 22:

What is the structure marked with the yellow arrow in the intraoperative image of spermatic cord of a patient with a congenital inguinal hernia?



a) Obturator vein

b) Femoral vein

c) Inferior epigastric vein

d) Testicular vein

Question 23:

A chronic tobacco chewer presented with an ulcerative-nodular lesion of 4x5 cm over the lip, as shown below. Which investigation would most likely confirm his diagnosis?

a) Superficial incisional biopsy along with normal tissue

b) Deep incisional biopsy of Ulcer along with normal tissue

c) Excisional biopsy along with normal tissue

d) Brush biopsy

Question 24:

A patient presented with recurrent swelling in his lower lip for the past 6 months as shown in the image. What is the diagnosis and treatment option for the patient?

a) Plunging ranula ; Incision and drainage

b) Mucous retention cyst ; Excision

c) Ranula ; Incision and Drainage

d) Mucocele ; Aspiration

Question 25:

A male child experienced acute scrotal pain while playing. He was taken to the emergency department and a USG doppler was performed, followed by immediate surgical exploration. The following finding was seen. What is the probable diagnosis?



a) Testicular hematoma

b) Strangulated inguinal hernia

c) Torsion of testicular appendage with Gangrene

d) Testicular torsion with Gangrene

Question 26:

Identify the procedure done in the given image.

a) Bilateral Orchidectomy with bilateral inguinal node dissection

b) Hernia repair

c) Sentinel Lymph node biopsy

d) Total penectomy with inguinal node dissection

Question 27:

The patient presented with headaches, repeated episodes of elevated blood pressure, and tachycardia. Urine shows elevated levels of vanillylmandelic acid. Her CT scan revealed a retroperitoneal tumor in the right lumbar region. Which among the following statements is true regarding the condition?

a) It can be associated with MEN 2A syndrome

b) It is commonly seen as as a bilateral condition

c) It is most commonly malignant

d) It is more commonly seen in children

Question 28:

What is the ideal pressure exerted by the method of dressing for a non-healing leg ulcer shown below?

a) 40-50 mmHg

b) 60-70 mmHg

c) 110-130 mmHg

d) 90-100 mmHg

Question 29:

A patient is diagnosed with Stanford type B aortic dissection. What is the most appropriate initial management of this patient?



a) Administer beta blocker and intensive blood pressure monitoring

b) Bypass graft

c) Percutaneous endovascular stent

d) Balloon dilation

Question 30:

A patient presented with dysuria and increased frequency of micturition. He has a history of recurrent urinary tract infections with Proteus species. The radiographic image is given below. Identify the calculi shown.

a) Cystine

b) Calcium Phosphate

c) Xanthine

d) Calcium oxalate

Question 31:

Identify the given drain.

a) Underwater seal drain

b) Closed suction drain

c) Penrose drain

d) Nelaton drain

Question 32:

Identify the ulcer shown in the image.

a) Arterial ulcer

b) Venous ulcer

c) Callous ulcer

d) Neuropathic ulcer

Question 33:

A patient underwent surgery for a leg ulcer due to burns. What is the mechanism by which a graft receives its nutritional supply on postoperative day 3?

a) Inosculation

b) Imbibition

c) Diffusion

d) Neovascularisation

Question 34:

A woman noticed a neck swelling 2 months back. Clinical examination reveals a solitary nodule on the right side as shown. What is the most appropriate initial investigation for this patient?

a) Radioisotope I-131 assay

b) PET Scan

c) Radioisotope I-123 assay

d) Serum TSH, T3 and T4

Question 35:

A 76 year old elderly man presents with confusion. He is on antihypertensive medication, and on aspirin after a recent MI. He recalls a trivial fall from chair 3 weeks ago. Other neurological examination was normal. What is the likely diagnosis?



a) Epidural hemorrhage

b) Subarachnoid hemorrhage

c) Subdural hemorrhage

d) Normal

Question 36:

A 45-year-old male presents with acute retention of urine which is drained via the suprapubic route. A retrograde urethrogram was done and is as shown below. What is the most appropriate management?

a) Serial dilatation

b) Urethral repair with graft from buccal mucosa

c) Metallic catheter placement

d) Endoscopic urethrotomy

Question 37:

A patient diagnosed with GERD has no improvement after long-term medical therapy. He underwent surgical treatment for GERD. Identify the procedure.

a) Toupet's fundoplication

b) Dor's fundoplication

c) Watson's fundoplication

d) Nissen's fundoplication

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	b
5	c
6	b
7	b
8	a
9	b
10	d
11	b
12	b
13	d
14	b
15	a
16	d
17	d
18	c
19	b
20	c
21	a
22	d
23	b
24	b
25	d
26	d
27	a
28	c

29	a
30	b
31	b
32	b
33	a
34	d
35	c
36	b
37	d

Detailed Explanations

Solution to Question 1:

The diagnosis, considering the purulent discharge shown in the image, is a fistula in ano.

A fistula-in-ano is an abnormal hollow tract or cavity that is lined with granulation tissue and that connects a primary opening inside the anal canal to a secondary opening in the perianal skin. The most common cause is a perianal abscess. Other causes include tuberculosis or Crohn's disease.

Symptoms:

• Perianal pain

• Discharge- intermittent purulent discharge (which may be bloody)

Goodsall's rule can be used as a guide in determining the location of the internal opening of the fistula.

• Fistulas with an external opening anteriorly connect to the internal opening by a short, radial tract.

• Fistulas with an external opening posteriorly track in a curvilinear fashion to the posterior midline.



Exceptions to Goodsall's Rule

However, exceptions to this rule often occur: Fistulas with an anterior external opening greater than 3 cm from the anal margin usually track to the posterior midline.

Investigations:

The investigations done for fistula-in-ano are as follows:

• Fistulogram

• Endoanal ultrasound

• MRI

• Computed tomography (CT) scan

• Endocoil MRI

• Hydrogen peroxide–enhanced 3-D endoanal ultrasonography

MRI is the gold standard investigation for fistula imaging. It helps us to identify the secondary extensions of the track, which when missed may cause persistence of the fistula.

Treatment:

• Control of constipation

• Sitz bath

• Fistulotomy

• Fistulectomy

• Ligation of intersphincteric fistula tract (LIFT)

• Setons

• Advancement flaps

Other options:

Option B: A fissure in ano is a linear ulcer usually found in the midline, distal to the dentate line. It is one of the most common causes of anal pain and bleeding. They typically start with a tear in the anoderm within the distal half of the anal canal which then triggers cycles of recurring anal pain and bleeding, which lead to the development of a chronic anal fissure. The exposed internal sphincter muscle within the bed of the fissure frequently spasms, which not only contributes to severe pain but also can restrict blood flow to the fissure, preventing its healing.

Option C: Hemorrhoids are cushions of submucosal tissue containing venules, arterioles, and smooth muscle fibers located in the anal canal. External hemorrhoids are distal to the dentate line and are covered with anoderm; these may engorge, causing pain. Thrombosis of these external hemorrhoids causes severe pain.

Option D: Pilonidal sinus disease is commonly seen in the natal cleft, more in men, usually after puberty. Drivers and hairdressers are more commonly affected. The pilonidal sinus opening is usually seen in the midline between the sacrococcygeal joint's level and the coccyx's tip. The other sites that can be affected include the interdigital cleft and axilla.

Solution to Question 2:

The type of breathing pattern observed in this patient with the X-ray showing contiguous rib fractures on the right side suggestive of a flail chest is paradoxical respiration.

A flail chest usually results from blunt trauma associated with 3 or more rib fractures in 2 or more places. It arises when these injuries cause a part of the chest wall to move independently of the rest of the chest wall. It is diagnosed clinically by observing paradoxical respiration. On inspiration, the loose segment of the chest wall gets displaced inwards, and on expiration, it moves outwards. They are at higher risk of developing pneumothorax/hemothorax. CT scan is the gold standard investigation.

Management includes oxygen administration, adequate analgesia (including opiates) or epidural analgesia, and physiotherapy. Intermittent positive pressure ventilation (IPPV) is reserved for patients developing respiratory failure despite adequate analgesia and oxygen

Treatment algorithm for treatment of flail chest with respiratory failure:



Different patterns of breathing observed are shown below:



Solution to Question 3:

The given image of a micturating cystourethrogram (MCU) shows dilated prostatic urethra and transverse filling defects corresponding with the posterior urethral valves and the most appropriate management for this condition is destruction of valves.

Posterior urethral valves are the most common cause of severe obstructive uropathy in children. The pathological valves are membranes that have a small posterior slit within them, which typically lie just distal to the verumontanum. They act as flap valves and are obstructive to antegrade urinary flow.

Diagnosis is most commonly made antenatally with ultrasound, which demonstrates bilateral hydronephrosis above a distended bladder in fetus along with oligohydramnios and pulmonary hypoplasia.

If the diagnosis is not made antenatally, then babies typically present with urinary infections in the neonatal period or with uremia and renal failure. Clinically a poor urinary stream, dribbling, and palpable bladder (due to diverticula) can be present. The investigation of choice is micturating cystourethrography.

The initial step in treatment is to insert a urinary catheter to drain the urine and decompress the bladder and upper urinary tracts. A small polyethene feeding tube (No. 5 or No. 8 French) is inserted into the bladder, which can be difficult because the tube's tip can coil in the prostatic urethra. A Foley catheter should not be used because the balloon can cause severe bladder spasms, which can result in severe ureteral obstruction.

Endoscopic valve ablation and symptomatic management of urinary tract infections are the definitive treatments.

The below image shows the micturating cystourethrogram (MCU) with dilated prostatic urethra and transverse filling defects (constriction) corresponding with the posterior urethral valves.



The ultrasound shows the typical 'key-hole' sign in which the bladder and the proximal urethra are dilated.



Solution to Question 4:

The given endoscopic image of a patient suggests esophageal varices.

Esophageal varices are tortuous dilated veins within the mucosa and submucosa of the distal esophagus and proximal stomach. Variceal hemorrhage is an emergency that can be treated medically by inducing splanchnic vasoconstriction or endoscopically by sclerotherapy, balloon tamponade, or variceal ligation.

Other options:

Option A: Gastric ulcers are often found near the lesser curvature of the stomach, especially in the antral area. Although duodenal ulcers are more common overall, gastric ulcers are frequently located near the lesser curvature, within a few centimetres of the pylorus. A significant cause of these ulcers is infection with Helicobacter pylori.

The following is an endoscopic image of Gastric ulcer:



Option C: Esophageal candidiasis is a fungal infection of the esophagus caused by the overgrowth of yeast, commonly Candida albicans. This condition is often seen in individuals with weakened immune systems, such as those with HIV/AIDS, diabetes, or those undergoing immunosuppressive treatments.

The following is an endoscopic image of esophageal candidiasis:



Option D: Barrett's esophagus is a type of metaplasia at the gastro-esophagal junction due to chronic gastroesophageal reflux disease (GERD). The squamous esophageal mucosa is converted to a columnal/intestinal type that is better suited to handle acid reflux. It is associated with an increased risk of esophageal adenocarcinoma.

The following shows an endoscopic image of Barrett's esophagus:

Solution to Question 5:

The earliest age to correct this defect in a child with bilateral cleft lip and palate is 6 months.

Cleft lip repair is done between 3-6 months of age and palate repair is done between 6-12 months of age. As this child's deformity needs to be corrected by palatal repair, the best answer in this scenario is 6 months.

A cleft lip is due to the failure of the fusion of maxillary prominence with medial nasal prominence. It can be unilateral or bilateral. It can vary in severity from a small notch in the upper lip to a double cleft extending into both nostrils. There is also associated underdevelopment of the part of the maxilla that bears the incisors. Millard cleft lip repair is done by rotating the local nasolabial flaps. Cleft lip repair is preferably done between 3-6 months of age.

A cleft palate can occur in the primary palate (anterior cleft) or the secondary palate (posterior palate). A Complete cleft palate involves both the primary and secondary palate. An incomplete cleft palate involves only the secondary palate.

In the bilateral complete cleft palate, a Y-shaped cleft is seen with gaps between the primary and secondary palate and between the two halves of the secondary palate. In addition, there is a bilateral cleft upper lip.

The different types of facial clefts are shown below:



- Unilateral cleft lip with alveolar involvement
- Bilateral cleft lip with alveolar involvement
- Unilateral cleft lip associated with cleft palate
- Bilateral cleft lip and palate
- Cleft palate only

Note: Palate repair is done between 6-12 months of age. The palatal repair is done maintaining the balance between speech outcomes and mid-facial growth. The generally agreed timeline is before speech development. However, the exact timeline is still debated and no uniform consensus is available.

Solution to Question 6:

The cystoscopic findings are suggestive of a "golf hole ureter", which is most likely associated with genitourinary tuberculosis(TB).

Genitourinary TB is caused by the dissemination of the organism through the bloodstream and is thus always a secondary TB. There is either reinfection or reactivation of old TB. It can affect the kidneys, ureters, urinary bladder, prostate, seminal vesicles, epididymis, and vas deferens.

Lesions caused by tuberculosis in the kidney include:

- Golf hole ureteric orifice - This is due to fibrosis causing the rigid withdrawn dilated ureteric orifice to look like a golf hole (cystoscopic finding).
- Papillary ulcer - earliest lesion
- Pyonephrosis - caused by multiple scattered papillary ulcers undergoing caseous necrosis, which later become filled with pus
- Perinephric abscess - by extension of pyonephrosis or tuberculous renal abscess

• Pseudocalculi - occurs when calcium gets deposited in a papillary ulcer healed with fibrosis

• Ghost calyces - caused by the destruction of the calyces

• Thimble bladder -the entire urinary bladder gets fibrosed, stiff, and unable to dilate and accommodate urine (IV urogram finding).

• Putty/cement kidney - amorphous dystrophic calcification eventually involves the entire kidney (seen in X-ray and CT)

The following X-ray KUB shows a thimble bladder:



Ureteric lesions can also be seen in genitourinary tuberculosis. They include:

• Kerr's kink - kinking of the ureteropelvic junction (UPJ)

• Ureteral stricture

• Beaded ureter

• Shortening of the ureter

• Golf hole ureter - Longitudinal fibrosis leads to the shortening of the ureter and pulling up the orifice as a gaping hole. Reimplantation of ureter is done.



Solution to Question 7:

The instrument required for placing the Double J stent (DJ stent), as shown in the radiograph is a cystourethroscope.

Cystourethroscopy allows for the examination and access of the bladder and urethra, which is crucial for guiding the DJ stent from the bladder into the ureter. This procedure involves inserting a flexible or rigid endoscope through the urethra and into the bladder, where the stent can be manoeuvred into the ureter.

While ureteroscopy is typically used to visualize and treat issues directly within the ureter, cystourethroscopy provides the necessary access and visualization to position the DJ stent correctly. This approach is effective for ensuring the stent is placed accurately, particularly in cases where access to the bladder is essential.

Other options:

Option A: A ureteroscope is used to examine and treat the ureters (the tubes connecting the kidneys to the bladder). It is commonly used for removing kidney stones or evaluating issues in the ureter.

Option C: A nephroscope is a type of scope used to examine the kidneys directly. It is often used in procedures like percutaneous nephrolithotomy (PCNL) to remove large kidney stones.

Option D: An ultrasound-guided scope refers to an endoscope or medical device used in conjunction with ultrasound imaging to guide procedures. It's often used in various parts of the body, including the gallbladder, liver, or kidney to help visualize structures during minimally invasive surgeries or biopsies.

Solution to Question 8:

The appropriate management for this patient with a diagnosis of umbilical hernia with painful, non-reducible swelling is umbilectomy with umbilical herniorrhaphy.

An umbilical hernia occurs when a defect in the umbilical or median raphe area allows abdominal contents to protrude. In adults, these hernias often present as a swelling near the umbilicus that can become painful and non-reducible over time. Conditions that cause stretching and thinning of the midline raphe, such as obesity, pregnancy, or liver cirrhosis with ascites, increase the risk of such complications.

The management of painful, non-reducible umbilical hernia typically involves surgical intervention due to the risk of strangulation and bowel obstruction. Umbilical herniorrhaphy with umbilectomy is the standard approach, addressing both the hernia and the associated defect.

In children, the umbilical defect is usually present at birth and typically closes as the umbilical cord stump heals within the first week. If this closure process is delayed, it can result in herniation during the neonatal period. Additionally, in adults, the umbilical ring can stretch and reopen, leading to similar herniation.

Solution to Question 9:

The presence of feculent discharge at the umbilicus with the persistence of the vitellointestinal duct is suggestive of vitelline fistula which occurs due to an ileal diverticulum.

The given clinical image is that of a congenital umbilical anomaly such as patent urachus, urachal cyst, and vitelline cyst (resulting from the persistence of omphalomesenteric duct). The next best step is the USG abdomen to diagnose if the anomaly is cystic, fibrous or fistula and its connections.

Vitello intestinal duct or omphalomesenteric duct is the connection between the yolk sac and the midgut during embryonic life, which eventually gets obliterated. The persistence of the vitellointestinal duct leads to various abnormalities depending on which part of the duct remains patent.

• Persistence of the proximal intestinal end of the duct leads to the Ileal diverticula/Meckel's diverticula.

• Persistence of the distal end near the umbilicus leads to the umbilical polyp or sinus.

• Persistence of the distal end with mucus retention leads to the cyst, which may be associated with the intestine or the umbilicus by a fibrous band.

• A fully patent vitellointestinal duct can lead to an enterocutaneous fistula.



Patent urachus occurs due to the failure of urachus to obliterate, resulting in a connection between the bladder and the umbilicus.

Urachal cyst occurs when the urachus fails to obliterate but there is no connection between the bladder and the umbilicus.

The image below shows a urachal cyst (UC) attached to Meckel's diverticulum (MD) on one side and a urinary bladder (UB) on the other side with a patent tract- urachus (U).



Solution to Question 10:

The image shows a hexagonal crystal, suggesting cystine stones. These stones are a recurrent feature in patients with cystinuria.

Cystinuria is the most common inborn error of amino acid transport. It occurs due to defective basic amino acid transporter in the proximal tubules of the kidney. This leads to the excretion of cystine(dimer of cysteine), ornithine, lysine and arginine, abbreviated as COLA.

The disease manifests as precipitation of cystine stones in the urine leading to renal colic. Oral hydration is the mainstay of treatment.

Crystals in urine are usually associated with kidney stones. Crystals form in the urine depending upon a variety of factors including:

• The degree of concentration of constituent molecules

• The urine pH

• The presence of inhibitors of crystallisation

Calcium oxalate and uric acid crystals can sometimes be found in normal urine that has been left to stand, due to crystal formation ex vivo.

Solution to Question 11:

The given clinical image is suggestive of partial thickness burns and the treatment plan for this condition is split thickness skin grafting.

Split-thickness skin graft (Thiersch graft): Partial-thickness skin grafts consisting of the epidermis and a portion of the dermis are called split-thickness skin grafts. Less dermis in the split-thickness skin graft translates into less primary contraction (the degree to which a graft shrinks in surface area after harvesting and before grafting) and more secondary contraction (the degree to which a graft shrinks during healing). Humby knife is used to harvest split-thickness skin grafts.

Advantages of split skin graft over full-thickness skin graft:

• Large grafts can be harvested.

• In the case of thinner split-thickness grafts, the donor site heals more rapidly and can be reused in severe cases such

as burns.

• Uptake is good.

• Meshing can be done, i.e., small splits are made in graft and any exudates collected can come out through this, improving graft takeup. It also allows the graft to expand.

Other options

Option A: Full-thickness skin grafts (Wolfe graft) include the epidermis and entire dermis, with portions of the sweat glands, sebaceous glands, and hair follicles. More dermis in the full-thickness skin graft translates into more primary contraction and less secondary contraction.

Option C: VAC is a negative pressure dressing. An occlusive dressing is placed over the sore and it is connected to VAC that generates -125mmHg of pressure. It is used in chronic non-healing ulcers, venous ulcers, diabetic ulcers, burn wounds, and bed sores.

Venous ulcers are due to chronic venous insufficiency and are particularly seen in the lower ■ of the leg above the ankle on the medial aspect, known as the gaiter zone. Ambulatory venous

hypertension is the most acceptable theory in the pathophysiology of venous ulcers. Venous ulcers have a sloping edge and the floor contains granulation tissue covered by slough and exudate. Marjolin's ulcer is a type of squamous cell carcinoma seen in chronic long-standing venous ulcers.

Option D: Flaps are tissues with a blood supply, providing vascularity to a new area. Types include:

• Random Flaps: Rectangular, with no specific blood supply, length-to-breadth ratio ≤ 1.5:1.

• Axial Flaps: Long flaps based on known blood vessels, suitable for large distances.

• Pedicled/Isolated Flaps: Axial supply allows them to be moved on a stalk or fully isolated.

• Free Flaps: Blood supply disconnected and reconnected via microsurgery.

• Composite Flaps: Combine tissues like skin with bone or muscle.

• Perforator Flaps: Use small vessels from major blood vessels to supply the surface.

Solution to Question 12:

The most important indicator for considering adjuvant radiotherapy in this patient is extranodal extension (ENE).

For patients with oral cavity cancer, ENE is a key indicator for adjuvant radiotherapy. Adjuvant therapy, typically involving external beam radiotherapy, is recommended based on pathological features such as ENE, close margins (1–4.9 mm), or involved margins (≤ 1 mm), lymphovascular invasion (LVI), and perineural invasion (PNI).

The presence of one major criterion - ENE or margins ≤1 mm or two minor criteria - close margins (1–4.9 mm), multiple involved nodes, largest node >3cm, LVI/PNI or T3/4, guides the decision for additional treatment.

Major criteria

• Extranodal invasion

• Lymphovascular invasion

• Perineural invasion

• T3/T4

Minor criteria

• Close margins (1–4.9 mm)

• Multiple involved nodes

• Largest node >3cm

Landmark trials have demonstrated that high-dose cisplatin added to radiotherapy enhances outcomes in high-risk patients, particularly those with ENE and positive margins.

Note: The PMMC flap is the most commonly used flap for soft tissue reconstruction in head and neck surgery. The pectoralis flap can be harvested as a musculocutaneous flap or muscle-only flap and is based on the pectoral branch of the thoracoacromial artery. Following flap harvest, a subcutaneous tunnel is created from the donor site, over the clavicles, to the defect. Division of the pectoral nerve branches ensures atrophy of the muscle to reduce the bulge over the clavicle over time.

Solution to Question 13:

The given image shows a pressure sore with extensive tissue destruction, including damage to the muscle layer and thus it belongs to stage 4.

Pressure sores also known as bed sores, pressure ulcers and decubitus ulcers are ulcers that occur due to prolonged pressure on certain sites thereby leading to tissue necrosis. The commonest sites involved are ischium, greater trochanter, sacrum, heel, malleolus (lateral >medial), and occiput. Paraplegic patients, elderly patients, and severely ill patients are at a higher risk of developing pressure sores.

Management is primarily proper skin care with pressure dispersion cushions, air-fluidized beds, and urinary or fecal diversion if required. Patients who are bed-bound should be turned once in 2 hours. Those in a wheelchair should lift themselves off the seat for 10 seconds every 10 minutes.

Surgery is the last resort and is done after a proper initial debridement and vacuum-assisted closure. Vacuum-assisted closure is a wound closure technique in which negative pressure is applied to the wound which leads to a decrease in edema. This hastens the debridement and formation of granulation tissue. Skin flaps with a muscle are also used for closure.

Solution to Question 14:

The fluid shown in the image is ringer lactate or Hartmann's solution.

Ringer lactate, or Hartmann's solution, is a crystalloid made up of sodium 131 mEq, chloride 112 mEq, potassium 5 mEq, bicarbonate (as lactate) 29 mEq, and 2 mEq of calcium. It is devoid of glucose.

It is useful in the following conditions:

• Correction of severe hypovolemia

• Metabolic acidosis

• Postoperative period, burns, diarrhoea

• Correction of hypokalemia

• Diabetic ketoacidosis.

It should be avoided in severe shock, liver diseases, vomiting, nasogastric aspiration, and metabolic alkalosis. It cannot be used along with blood transfusion since calcium in the ringer lactate can bind with citrate in the transfused blood leading to clot formation.

Other options:

Option A: Haemaccel is a colloid and contains 145 mmol of sodium, 6.25 mmol of calcium, and 5.1 mmol of potassium. It is mainly used for rapid volume replacement in circulatory shock, burns, and trauma. These release histamine and hence, an H2 blocker has to be supplemented. It is also used in heart-lung machines.

Options C: Isolyte P is a crystalloid. Isolyte P is commonly used in fluid therapy for paediatrics. One litre contains 50 grams of glucose, 25 mEq of sodium, 22 mEq of potassium, 3 mEq of magnesium, and 23 mEq of acetate. Hyponatremia, renal failure, hyperkalemia, and hypovolemic shock are some of the contraindications.

Option D: Isolyte M is a crystalloid. Isolyte M is used as a maintenance fluid. It mainly contains 50 grams of glucose, 40 mEq sodium, 38 mEq chloride, 35 mEq potassium, 15 mEq phosphate, and 20 mEq acetate. It is useful in correcting hypokalemia in patients with good renal function, and acidosis. It is contraindicated in hyponatremia.

Solution to Question 15:

The image of a large breast mass with an unevenly bosselated surface, overlying skin ulceration, and concomitant liver metastasis is indicative of a malignant phyllodes tumor which is managed by simple mastectomy.

Phyllodes tumors spread primarily through the bloodstream (hematogenous). Lymph node metastases are uncommon. Large or malignant phyllodes tumors may require a simple mastectomy, but axillary dissection is not recommended (Option C and D) due to the rare involvement of axillary lymph nodes. The role of adjuvant chemotherapy (Option B) in malignant phyllodes tumors remains uncertain and is generally not recommended as a standard treatment.

Phyllodes tumor also known as a sero-cystic disease of Brodie or cystosarcoma phyllodes, originates from hormonally responsive stromal cells of the breast. It is usually a benign condition and affects women > 40 years of age, but can be seen in younger women. Sometimes, may develop malignant transformation.

The clinical features include large, sometimes massive, with an unevenly bosselated surface, ulceration of overlying skin occurs because of pressure necrosis and it remains mobile on the chest wall.

There is no role of FNAC in phyllodes tumor. Histopathology of the lump biopsy shows increased stromal cellularity, overgrowth, and typical leaflike architecture. Tumors are treated by excision with margins of at least 2 cm to prevent local recurrence.

Solution to Question 16:

The most appropriate management for this patient with stage T3 carcinoma of the penis with no clinically palpable inguinal lymph nodes is penectomy with superficial inguinal lymph node dissection.

In penile carcinoma, total amputation of the penis is done if the stump is < 2cm after resection of the tumour and partial penectomy is done if the stump is >2cm. Carcinoma of the penis is commonly a squamous cell carcinoma. It arises from the skin of the glans penis or the prepuce. The lesion can be flat or warty in appearance.

• T1- confined to the skin

• T2- invades the corpus spongiosum or cavernosum.

• T3- invades the urethra

• T4- invades adjacent structures.

The first lymph node to be involved is the superficial inguinal lymph node and thus, superficial inguinal lymph node dissection or sentinel lymph node biopsy (SLNB) helps evaluate the spread of penile cancer.

Tumours of low-grade and low-stage are managed by Mohs' surgery. Tumors involving the glans penis require glansectomy, and in advanced cases require partial penectomy.



Solution to Question 17:

The most commonly occurring complication of TPN in a sepsis patient in the ICU is hyperglycemia.

To monitor or prevent complications in patients on TPN, the following steps may be taken:

• Infections

• Monitor the development of glucose intolerance as it is the earliest sign of systemic sepsis. (Note: glucose intolerance- glucose not processed effectively, leading to hyperglycemia)

• Use subclavian vein preferably

• Metabolic complications

• BUN and LFTs weekly

• Serum electrolytes every 2-3 days

• Intestinal atrophy

• Provide at least some nutrients enterally if feasible

TPN also referred to as central parenteral nutrition is used for patients who can't absorb nutrients through their digestive tract, often due to conditions like short bowel syndrome from extensive intestinal resection or fistulas. It can also be temporary until other feeding methods, like a nasojejunal tube, are established.

TPN typically comes in a 3-liter bag with a lipid emulsion, amino acids, glucose, electrolytes, and vitamins. The energy provided is about 150–250 kcal per gram of protein nitrogen, with 30–50% from fat. Folic acid is added weekly, with other vitamins given daily, and long-term TPN may require vitamin B12 supplementation.

Solution to Question 18:

The given CT scan showing the hydatid cyst with detached membranes(Water Lily sign) is classified as CE-3 according to the WHO classification and Type 2 according to the Gharbi classification.



Hydatid Cyst is a zoonotic infection caused by the Echinococcus species(dog tapeworm). Dogs are the definitive hosts of E. granulosus. Sheep are the usual intermediate hosts, but humans are accidental intermediate hosts. Humans are end-stage hosts of the parasite since there is no human-to-human transmission.

The clinical presentation of a hydatid cyst is largely asymptomatic until complications occur. The most common presenting symptoms are abdominal pain, dyspepsia, and vomiting. The most frequent sign is hepatomegaly.

Approximately 75% of hydatid cysts are located in the right lobe of the liver and are solitary.

CT scan is the best imaging modality for hydatid cysts. It shows the presence of a space-occupying lesion with a smooth outline with septa. Diagnosis is suggested by the finding of a multiloculated cyst on ultrasound and is further supported by the finding of a floating membrane within the cysts on a CT scan. Active cysts contain a large number of smaller daughter cysts (multi-vesicular rosette-like cystic lesions with daughter cysts) which is characteristic of hydatid cysts of the liver.



If there is a disruption of the inner layers of the cyst, the following signs may be noted:

• Water lily sign/ camalote sign - air-fluid level with a floating membrane (as shown below)

• Rising sun sign/ serpent sign - a double-wall, dry cyst with crumpled membranes lying at its bottom

• Empty cyst sign - when all the contents are removed



The ultrasound of an active hydatid cyst shows multiple septations and a rosette or honeycomb appearance (as shown below)



The serpent sign is seen as wavy membranes within the cyst, on CT imaging.

A ball of wool sign is seen on ultrasound when the contents settle in the dependent portion of the cyst, as seen below.



The water-lily sign is seen in hydatid infections when there is detachment of the endocyst membrane which results in floating membranes within the pericyst that mimic the appearance of a water lily, on USG.



Solution to Question 19:

The given image showing a midline defect situated in the lower abdominal region below the umbilicus with an exposed bladder is suggestive of ectopia vesicae/ bladder exstrophy.

Ectopia vesicae/ bladder exstrophy is a congenital abdominal wall defect wherein the bladder protrudes from the lower abdominal wall and its mucosa is exposed. This results in the separation of the pubic rami and the rectus muscles.

The condition can be associated with complete epispadias with dorsal chordee, undescended testes, and inguinal hernias in males. Females may also have epispadias and separation of the two halves of the clitoris (bifid clitoris). The condition can be associated with rectal prolapse and short stature in both sexes.

Untreated exstrophy can cause urinary incontinence and is associated with an increased incidence of bladder cancer. Management involves a staged surgical reconstruction of the bladder and genitalia.

Other options:

Option A: The primitive gut is connected to the definitive yolk sac through a narrow stalk-like communication known as the vitellointestinal duct. In about 2% of the population, a small section of the vitellointestinal duct persists to form an outpouching of the ileum called the Meckel diverticulum or ileal diverticulum.

Option C: Gastroschisis is a condition where the intestinal loops herniate through the defect in the abdominal wall. The defect usually lies on one side of the umbilicus (Right side commonly). The loops of the bowel are not covered by amnion. It is associated with elevated alpha-fetoprotein levels.

Option D: Omphalocele is another congenital midline abdominal wall defect at the umbilicus causing herniation of the abdominal contents into the base of the umbilical cord. The herniated viscera is covered with peritoneum without any overlying skin. This condition can be associated

with Beckwith-Wiedemann syndrome which is associated with embryonal tumors such as Wilms tumor.

Solution to Question 20:

The given image shows a barium swallow. It is a type of imaging where the patient ingests barium, a contrast agent, to visualize the esophagus and detect abnormalities. The described indentations in the esophageal lumen caused by anterior osteophytes (bony outgrowths) are characteristic findings that can be seen with a barium swallow.

Fluoroscopy (option A) is a technique that provides real-time X-ray images and is often used during procedures like a barium swallow, but the term is more general. X-ray of alkali ingestion (Option B) and X-ray of acid ingestion (option D) might show damage to the esophagus in barium swallow which is not seen in the image.

Solution to Question 21:

The image 3 given above shows yellow plaques and nodules on the mucosa are suggestive of pseudomembranous colitis. It is usually caused by Clostridium difficile. The pathogenesis is mediated by the enterotoxins - toxin A and toxin B.

Clostridium difficile is a gram-positive anaerobic bacillus that causes Clostridioides (formerly Clostridium) difficile infection after ingestion of spores of the organism. The more severe form of the condition is known as pseudomembranous colitis (PMC). Enzyme immunoassay to detect glutamate dehydrogenase is less specific compared to toxin detection for diagnosis. So a combination of the two can be used for diagnostic confirmation.

The condition is most commonly associated with prolonged antimicrobial use, which interferes with the commensal colonic microflora. The most commonly implicated antibiotics are cephalosporins, ampicillin, and clindamycin. The spores are spread by contact with the feces of affected individuals or asymptomatic carriers. It can persist on surfaces for months. Risk factors include poor hand hygiene, old age, and gastrointestinal surgery.

The spores germinate in the small intestine and produce two toxins—toxin A (enterotoxin) and toxin B (cytotoxin). These interfere with the epithelial-cell barrier function and result in diarrhea and pseudomembrane formation. The pseudomembranes appear as small whitish-yellow plaques initially which later coalesce to involve the entire colon wall. It contains necrotic leukocytes, fibrin, and cellular debris. The anamnestic IgG response determines which patients develop the disease or remain asymptomatic.

Symptoms include diarrhea, fever, and abdominal pain. The diagnosis is based on a combination of clinical criteria and detection of toxin or the organism in stool samples. This can be done by nucleic acid amplification testing (NAAT) methods such as polymerase chain reaction (PCR) or by culture.

Discontinuation of any ongoing antimicrobials and adequate hydration are general measures that are recommended. Oral fidaxomicin or vancomycin is the first-line treatment. Oral metronidazole can be used when these are not available.

Other options:

Option C: Familial Adenomatosis Polyposis (image 1)is caused due to a germline loss of function mutation of the tumor suppressor gene, APC(adenomatous polyposis coli) on the 5th chromosome leading to multiple adenomas in the GI tract with 100% risk of malignant conversion associated with congenital retinal pigment epithelium hypertrophy. Variants of FAP are Gardner and Turcot syndromes. Its features are described in the pearl attached.

Option B: Esophageal varices (image 4) are tortuous dilated veins within the mucosa and submucosa of the distal esophagus and proximal stomach. Variceal hemorrhage is an emergency that can be treated medically by inducing splanchnic vasoconstriction or endoscopically by sclerotherapy, balloon tamponade, or variceal ligation. It commonly occurs due to portal hypertension in patients with cirrhosis.

Option D: Ulcerative colitis (UC) (image 2) which is characterized by continuous inflammation of the colon, pseudopolyps, biopsy findings of crypt distortion, ulceration, and infiltration of inflammatory cells until the submucosal layer of the intestinal wall. Diagnosis can be made by colonoscopy followed by a biopsy.

UC is a type of inflammatory bowel disease that primarily affects the rectum and extends proximally to involve the colon. Patients typically present with diarrhea, rectal bleeding, tenesmus, crampy abdominal pain, and mucus discharge. They may also present with anorexia, nausea, vomiting, fever, and weight loss.

The presentation of UC differs from that of Crohn's disease, which is characterized by transmural inflammation of the intestine. Differences are mentioned in the pearl.

Solution to Question 22:

The given clinical image is an open inguinal hernia repair of a child with a structure marked as the spermatic cord whose content includes the testicular vein.

Normally the funicular process of the peritoneum becomes obliterated after the testis has reached the scrotum. The scrotal part of the process remains patent and acquires the name 'tunica vaginalis'. In case of congenital hernia, the whole process remains patent. With the increase in abdominal pressure, abdominal contents come out through the patent peritoneal process. So it reaches the bottom of the scrotum very quickly. It may so happen that the funicular process remains patent up to the top of the testis. So the hernia stops at the top of the testis and is known as a congenital funicular hernia.

Indirect hernia comprises more than 80% of inguinal hernias which are common in women and children. It occurs earlier than a direct hernia and is often complete. The two forms often seen under indirect are congenital and acquired. Unlike congenital, acquired hernias do not become complete at once. They progress gradually. Clinically a congenital hernia may present at a later period.

The pearls attached below describe the inguinal canal and the differences between direct and indirect inguinal hernia.

Other options are not the contents of the spermatic cord. Contents are mentioned in the pearl below.

Solution to Question 23:

The given clinical scenario along with the image showing ulcero-proliferative mass is suggestive of lip cancer which is a type of oral carcinoma. Histopathological diagnosis in the form of a deep incisional biopsy of an ulcer along with normal tissue is the gold standard for diagnosis.

Incisional biopsy is taken from the edge of the ulcer along with normal tissue as the centre of the ulcer is usually necrotic. A narrow deep biopsy is better than a shallow broad biopsy. (Option A)

Excisional biopsy is done for smaller benign lesions. It is not done in oral cancers as they are usually bigger in size. An excisional biopsy may increase the spread of cancer or cause functional impairment. (Option C)

Techniques like brush biopsy and exfoliative biopsy are less sensitive. Hence, they are not used for diagnosis. (Option D)

Solution to Question 24:

The given scenario of recurrent swelling on the lip for 6 months and the image showing soft, and transparent cystic swelling points to a diagnosis of a mucous retention cyst. It has to be managed by an excision.

Mucous retention cysts result from duct obstruction by periductal scars or sialolithiasis (salivary gland stone formation) These cysts are mainly found in the ductal system of a minor salivary gland. The cyst cavity may contain fragments of a sialolith or mucous material. Some may regress spontaneously but best management is done by an excision.

Other Options:

Option A: Cervical or plunging ranula- is a rare type of retention cyst arising from the sublingual gland. The mucous passes through the mylohyoid muscle diaphragm to enter the neck. Clinically present as intraoral and cervical swelling. USG or MRI makes the diagnosis. Management includes excision of the sublingual gland + aspiration of saliva out of the sac.

Option C: Oral ranula - It is a benign cystic lesion formed by extravasation of mucus, involving the sublingual gland. Clinically present as cystic swelling in the floor of the mouth. On examination fluctuation test and transillumination test is positive. It is conservatively managed by marsupialisation. Definitive management is the excision of the cyst. Some may resolve spontaneously.

Option D: Mucocele occurs when an obstruction in the ducts of minor salivary glands, such as those in the labial or buccal areas, leads to swelling and aspiration is not a definitive treatment as it only provides temporary relief and does not prevent recurrence.

Solution to Question 25:

The given clinical scenario along with the intraoperative specimen is suggestive of testicular torsion with gangrene.

Testicular torsion is the twisting of the spermatic cord and its contents such that the testicular blood supply becomes compromised. It is a surgical emergency and if left untreated, the testes may become gangrenous. Early surgery undertaken to untwist the testis gives a better outcome.

The following are anatomical predispositions to testicular torsion:

• High investment in the tunica vaginalis causes the testis to hang within the tunica like a clapper in a bell. This is the most common cause in adolescents and is typically a bilateral abnormality

• Inversion of the testis. The testis is rotated so that it lies transversely or upside down.

• Separation of the epididymis from the body of the testis. This allows torsion of the testis on the pedicle that connects the testis with the epididymis.

Males in the age group of 10-25 years present with acute scrotal or groin pain with associated nausea and vomiting. On examination, the scrotum appears swollen and tender.

Torsion Testis has to be differentiated from Acute epididymo-orchitis. This is done by Prehn's sign. In the case of acute epididymo-orchitis, the pain gets relieved on the elevation of the testis but not in the torsion of the testes.

Other Signs seen in torsion of the testis are:

• Deming's sign: Spasm of the cremaster muscle and the twisting of the cord cause the affected testis to be positioned higher

• Angell's sign: The testis on the opposite side lies horizontally because of the bilaterality of the mesorchium between the testis and epididymis.

• Cremastric reflex- absent on the side of the affected testis

The morphologic changes range from intense congestion to widespread hemorrhage to testicular infarction depending on the duration of torsion. In advanced stages, the testis is markedly enlarged and consists entirely of soft, necrotic, hemorrhagic tissue.

A Doppler ultrasound may be taken to assess the blood supply but surgical exploration is still indicated as a clinical decision. If the testis is infarcted after the cord is untwisted, an orchidectomy must be undertaken and the opposite testis must be fixed to the scrotum to preserve fertility and prevent torsion. If the affected testis is viable after the cord is untwisted, both the affected and the contralateral testes are fixed to the scrotum (bilateral orchiopexy) using non-absorbable sutures.

Other Options :

Option A: Testicular hematoma will usually be associated with trauma to the testis.

Option B: Strangulated hernia will have a history of inguinal swelling and the presence of cough impulse over the swelling. the following image shows ischemic and necrotic bowel herniated into the sack.

Option C: Testicular appendage torsion is torsion of vestigial appendage present along the testis. It has no function but torsion of appendage can be painful. It presents with a blue dot sign on clinical examination. It is a self-limiting condition and conservative management is done.



Solution to Question 26:

The given post-operative image suggests that this patient underwent Total penectomy with inguinal lymph node dissection.

Total penectomy is done in cases of penile carcinoma if the stump is < 2cm after resection of the tumour and partial penectomy is done if the stump is >2cm. Carcinoma of the penis is commonly a squamous cell carcinoma. It arises from the skin of the glans penis or the prepuce. The lesion can be flat or warty in appearance.

The first lymph node to be involved is the superficial inguinal lymph node and thus, superficial inguinal lymph node dissection or sentinel lymph node biopsy (SLNB) helps evaluate the spread of penile cancer.

Tumours of low-grade and low-stage are managed by Mohs' surgery. Tumors involving the glans penis require glansectomy, and in advanced cases require partial penectomy.



Other options:

Option A: Testis is visible in the given image, therefore it is unlikely to be a specimen of Bilateral Orchidectomy with bilateral inguinal node dissection.

Option B: Hernia repair would be done higher up at the inguinal region.

Option C: SLNB will be done through a smaller incision.

Solution to Question 27:

The given clinical features along with elevated urinary levels of vanillylmandelic acid, and her CT scan showing a retroperitoneal tumor in the right lumbar region is suggestive of pheochromocytoma whose familial forms are associated with MEN2A syndrome.

Patients commonly present with a classical triad of palpitation, headache, and profuse sweating.

Pheochromocytoma is associated with:

• MEN-2: caused by germline mutations of the RET proto-oncogene

• Familial paraganglioma syndrome: caused by germline mutations within the succinate dehydrogenase complex subunit B (SDHB), SDHD, and SDHC genes.

• Von Hippel-Lindau syndrome: caused by a germline mutation in the VHL gene.

• Neurofibromatosis type 1: caused by a germline mutation in the NF1 gene.

Histopathology: Cells with salt and pepper chromatin. Other tumors with similar findings include medullary thyroid carcinoma, small cell cancer lung, and extra-adrenal counterpart of pheochromocytoma- paragangliomas. Given below is the histopathological image of pheochromocytoma.



Other Options:

Option B: Only 10% of the cases are bilateral.

Option C: Only 10% of the cases are malignant. Pheochromocytomas associated with MEN2A are rarely malignant(3-5%), but often malignant with SDHB gene mutation(50%).

Option D: It is a tumor of adulthood.

Solution to Question 28:

The pressure used in vacuum-assisted closure for wound healing is around –125 mmHg. That lies in the range of 110-130 mmHg

Applying negative pressure appears to hasten debridement and the formation of granulation tissue in chronic wounds and ulcers.

It acts by decreasing edema, by removing interstitial fluid, and as a result, helps in decreasing bacterial counts and cell proliferation increases.

Solution to Question 29:

The initial management of patients with all types of aortic dissection is beta blocker medication to control hypertension along with intense blood pressure monitoring and pain management.

Aortic dissection occurs when a defect or flap occurs in the intima of the aorta. Predisposing factors include mainly age, hypertension, Marfan syndrome, and pregnancy. The most typical presentation of aortic dissection is a tearing interscapular pain. The dissection may also result in aortic wall rupture into the pericardium (cardiac tamponade) or mediastinum (left haemothorax).

Contrast-enhanced CT is currently the most preferred imaging modality for aortic dissection. Double-lumen aorta is seen on CECT. Transesophageal echocardiography is preferred for patients

who are unstable for transportation.

Acute aortic dissection can be classified by the original DeBakey system, based on the location of the proximal tear and distal extent of the dissection flap, or the Stanford system which uses the location of the proximal tear alone.

Site of dissection	DeBakey	Stanford
Both ascending and descending aorta	Type I	Type A
Only the ascending aorta and arch of the aorta	Type II	Type A
Descending aorta distal to the subclavian artery	Type IIIaLimited to the thoracic aorta	Type B
Type IIIbExtends to the abdominal aorta		



Management of aortic dissection:

The initial management of patients with all types of aortic dissection is beta-blocker along with pain management. The preferred drug is esmolol. Hemodynamic stability is an important consideration.

• Stanford Type A (DeBakey Types I and II) requires surgical intervention once the patient's blood pressure is stabilized.

• Stanford Type B (DeBakey type III) is best managed by antihypertensive drugs. For patients with complicated type B dissection, percutaneously placed endovascular stents(option C)are currently preferred. Surgery (options B and D) is reserved for the rare patients not suitable for stenting.



Solution to Question 30:

The given clinical scenario and X-ray KUB showing a homogenous huge density with the smooth edge seen in the bladder is suggestive of struvite calculi, which is formed due to the precipitation of calcium, magnesium, and ammonium phosphate.

Struvite stones (triple phosphate stones) are usually seen in a setting of infection with urease-producing bacteria like Proteus, Pseudomonas, and Staphylococcus. They can grow very large and form a cast of the renal pelvis and calyces resulting in staghorn calculi. Production of urease enzyme by Proteus, which hydrolyses urea to ammonia causes alkalinization of urine. Alkalinization of urine leads to precipitation of organic and inorganic compounds which contributes to the formation of Struvite stones.

Almost 90% of the renal stones are radio-opaque and the remaining 10% are radiolucent. The investigation of choice for renal stones is NCCT.

Struvite stones are usually managed with percutaneous nephrolithotomy (PCNL) as it is usually larger in size and associated with Urinary tract infections.

Extracorporeal shock wave lithotripsy (ESWL) is a non-invasive method that generates shockwaves outside the body, which are focused on the stones. It is the most common method of treating urinary tract stones nowadays. Stones up to approximately 1.5 cm in size are suitable for this form of treatment.

PCNL is indicated in:

• Staghorn calculi - Stone >2 cm

• Multiple stones - >1 cm

• Proximal ureteral stone - >1 cm

Other options:

Option A: Patients with cystinuria (an autosomal recessive disorder) are at risk for Cystine stones. These stones are hexagonal.

Option C: Xanthine stones are seen in inborn errors of metabolism like Lesch Nyhan syndrome and cases of hyperuricemia. It is a radiolucent stone.

Option D: Calcium oxalate stones are seen in conditions causing high urine calcium such as primary hyperparathyroidism and chronic acidemia from distal renal tubular acidosis. Also seen in the condition of high urine oxalate which is seen during a low-calcium diet and in malabsorption syndromes(small bowel disease (eg, Crohn's disease), including a jejunoileal bypass for obesity (bariatric surgery), surgical bowel resection or diversion, or cystic fibrosis). Calcium oxalate monohydrate stones are usually dumbbell-shaped and Calcium oxalate dihydrate stones are usually envelope-shaped.

Solution to Question 31:

The Romo Vac drain, depicted in the image, is a closed suction drain commonly used after mastectomy.

Drains are placed at surgical sites to freely collect the air or fluids that may accumulate. This enhances wound healing by preventing the formation of unnecessary cavities. Drains are prophylactic in elective surgeries and therapeutic in emergency surgeries. Drains can be -

• Open - They are based on the principle of gravity

• Semi-open - Their action is based on capillary effect

• Closed - They carry out their actions through suction

Suction drains are commonly used in chest, breast, abdominal, thyroid, and plastic surgeries. They remove intraperitoneal secretions, anastomotic leaks, and blood from hemorrhage at the surgical site. Romo Vac drain is a negative suction drain closed drain kept in the subcutaneous tissue. Romo Vac drain is indicated in modified radical mastectomy, subcutaneous swelling excision, and amputations.

Special types of drains:

• Chest drains (option A) are connected to an underwater seal to aid the exit of air from the pleural cavity. They are used in pneumothorax, pleural effusion, and hemothorax.

• After surgical exploration of the bile duct, the sphincter of Oddi goes into spasm. To facilitate drainage of bile, a T tube may be used.

Drain insertion site care: The site of insertion of the drain is kept clean by drying and patting the drain insertion site with a clean towel after bath. The dressing is changed daily with folded gauze pieces placed below and above the drain site.

Removal of drains: The catheter at the site is milked for residual collections. It is then clamped and the suture holding the catheter is cut. The tube is then gently pulled out and any clots at the perforation end are removed. The time of removal of drains for various conditions is given below.

• Drains used in surgeries like thyroidectomy are usually removed within 24 hours. Their main purpose is to cover postoperative bleeding.

• In a mastectomy, a drain is placed for 5 days to collect serous collections.

• Drains used to cover the colorectal anastomoses are placed for 5-7 days.

• Common bile duct T-tube drains are usually placed for 10 days.

• Drains for infections are removed when the infection subsides or when the content of the drain is minimal.

After the drain is removed, a folded gauze is placed over the drain insertion area. The dressing can be removed after 24 hours. The patient may be given an antibiotic ointment to apply over the area for a week.

Other options:

Option A: The following image shows an underwater seal drain.



Option C: A Penrose drain is a flexible, soft, latex tube used to drain fluid from a surgical site after surgery. It's a type of passive drain.

Option D: Nelaton drain/catheter is commonly used as an intermittent urinary catheter. Like other intermittent catheters, it is also attached temporarily for draining the bladder and then removed.



Solution to Question 32:

The image of a healing ulcer with a sloping edge in the Gaiter's area indicates a venous ulcer.

Venous ulcers due to chronic venous insufficiency are most commonly seen in the lower ■ of the leg and foot. They usually lie just proximal to the medial or lateral malleolus.

Ambulatory venous hypertension is the most acceptable theory in the pathophysiology. They have a sloping edge and the floor contains granulation tissue covered by slough and exudate. Marjolin's ulcer is a type of squamous cell carcinoma seen in chronic long-standing venous ulcers. These are accompanied by lipodermatosclerosis and hemosiderosis.

Management of venous ulcers:

1) Bisgaard's regimen (4Es)

• Education of the patient

• Elevation of the affected limb to reduce the edema

• Elastic compression stockings - The ideal interface pressure in venous ulceration is 35-40 mmHg.

• Evaluation

2) 4-layer bandaging system comprising of:

• Orthopaedic wool

• Cotton crepe

• Elastic bandage - first compressive layer

• Cohesive bandage - second compressive layer

3) Antibiotics are given only if the wound is infected.

4) Regular dressing.

5) Medical management by oral pentoxifylline is considered which helps in reducing the viscosity and improving circulation.

6) Surgical management by ulcer excision with pinch mesh grafting has a good outcome and helps in preventing recurrence.

Note: Vacuum-assisted closure (VAC) is a negative pressure dressing. An occlusive dressing is placed over the sore and it is connected to VAC that generates -125mmHg of pressure. It is used in chronic non-healing ulcers, venous ulcers, diabetic ulcers, burn wounds, and bed sores.

Other options:

Option A: The following image depicts an arterial ulcer

Type of Ulcer	Ischaemic/Arterial	Venous	Neuropathic	Tuberculous
Underlying condition	Peripheral vascular disease mainly atherosclerosis, thromboangitis obliterans, and Raynaud's phenomenon	Varicose veins, previous deep vein thrombosis	Diabetes, peripheral neuropathy	Tuberculosis lymph nodes, cold abscess from bone and joint tuberculosis
Location	Toes, feet, legs, or even upper limb digits	Lower limb, commonly on the medial side of the ankle, never above the middle third of the leg	The plantar surface of the foot and toes	Neck, axilla, and groin
Characteristics	Deep, penetrates the deep fascia exposing the tendons, muscles, and underlying bone. The ulcer is painful, tender, and hyperaesthetic.	Vertically oval, and painless.The floor may have pale to no granulation tissue, induration, and tenderness at the base.	Punched-out with regular calloused margins.	Edges are thin, reddish blue, and undermined. Pale granulation tissue, scanty discharge on the floor, and slight induration at the base
Other features	Features of ischemia in adjacent areas, like pallor, dry skin, brittle nails, loss of hair	Inguinal lymphnodes are enlarged usually the vertical group. The ulcer is recurrent.	Dry, cracked, insensate	Regional lymph nodes are enlarged, non-tender, and matted

Option C: A callous ulcer is a type of foot ulcer that develops from a pre-ulcerative callus. It's a chronic, non-healing wound that can last for months or years. Callous ulcers are characterized by a hard base, rigid walls, pale tissue inside the ulcer, and a variety of shapes and sizes.

Option D: The following image is that of a neuropathic ulcer

Solution to Question 33:

The phase of skin graft uptake in which the graft receives its nutritional supply on postoperative day 3 is known as inosculation. It is the period during which the graft is most at risk for failure.

Split skin graft survives the transplantation surgery by the following process:

• The graft survives the first 48 hours because of the imbibition (option B) of plasma from the wound bed.

• After 48 hours, fine anastomotic connections are made, which leads to the inosculation of blood.

• Neovascularization (option D) is the process by which new blood vessels form, directly invading the graft and restoring the pink color of the skin. There is no step of diffusion (option C) in the process of graft transplantation.

Skin grafts can be broadly divided into two categories:

Split-thickness skin graft (Thiersch graft): Partial-thickness skin grafts consisting of the epidermis and a portion of the dermis are called split-thickness skin grafts. Less dermis in the split-thickness skin graft translates into less primary contraction (the degree to which a graft shrinks in surface area after harvesting and before grafting) and more secondary contraction (the degree to which a graft shrinks during healing).

Full-thickness skin grafts (Wolfe graft) include the epidermis and entire dermis, with portions of the sweat glands, sebaceous glands, and hair follicles. More dermis in the full-thickness skin graft translates into more primary contraction and less secondary contraction.

Advantages of split skin graft over full-thickness skin graft:

• Large grafts can be harvested.

• In the case of thinner split-thickness grafts, the donor site heals more rapidly and can be reused in severe cases such

• as burns.

• Uptake is good.

• Meshing can be done, i.e., small splits are made in graft and any exudates collected can come out through this, improving graft takeup. It also allows the graft to expand.

Advantages of full-thickness skin graft over split skin graft:

• It has a lesser degree of post-graft contraction

• Best cosmetic appearance as they include the highest number of skin appendage elements.

Solution to Question 34:

This patient is presenting with thyroid swelling along with image given is suggestive of Solitary Thyroid Nodule. Assessing the biochemical function of the thyroid, i.e. thyroid function test (TFT) which includes serum TSH, T3, and T4 is the first investigation done on any thyroid condition.



Investigations in a patient with Solitary thyroid nodule

• Thyroid function tests- Thyroid stimulating hormone (TSH) levels are the first step in evaluating suspected thyroid disease. Based on TSH levels, further investigations is decided.

• Ultrasonography is a non-invasive and inexpensive way to stratify the risk of the nodule into benign or malignant tissue.

• FNAC: Although other investigations can assist in evaluating a nodule, FNA cytology provides a definitive diagnosis.

• Radioiodine Thyroid scan: It helps to differentiate between a hot nodule and a cold nodule. Thyroid nodules that absorb radioiodine are called hot nodules. These nodules produce excess thyroid hormones and are functional nodules. Whereas, the thyroid nodules that do not absorb radioiodine are called cold nodules and they indicate non-functional nodules. It is mostly done by I-123 radioisotope. (Option C)

Other options:

Option A: I-131 radioisotope is more commonly used to radio-ablate the thyroid tissue.

Option B: PET scan has no role in the evaluation of Solitary thyroid nodules.

Solution to Question 35:

The given clinical vignette along with the CT image showing a crescent-shaped hyperdense lesion that crosses the suture lines is indicative of chronic subdural hematoma/hemorrhage.

Subdural hemorrhage (SDH) is a collection of blood between the dura and the arachnoid layers of the meninges. Most commonly, it results from the rupture of cortical bridging veins. Non-contrast CT is the investigation of choice.

Radiographic features of subdural hemorrhages:

• Typically unilateral (85%)

• Crescent-like distribution (concavo-convex) around the periphery of the brain.

• They are not limited by sutures and can fill dural reflection

• Crescent-shaped hypodense lesion on axial CT: Chronic subdural hemorrhage

• Hyperdense collections on CT: Acute subdural hemorrhage.

CT image showing acute subdural hemorrhage

CT image showing acute on chronic subdural hemorrhage.

Other Options:

Option A: Epidural haemorrhage will be usually lens-shaped and is limited by sutures.

Option B: Subarachnoid haemorrhage will show Hyperdensity in the subarachnoid space.

Solution to Question 36:

The given clinical scenario with a long-segment urethral stricture which will be best managed by urethral repair with a graft from buccal mucosa.

The management of a urethral stricture depends on the site, the length, and the cause of the stricture. They include:

• Endoscopic urethrotomy - The stricture is released using a knife passed through a rigid urethroscope. This is done for a short and incomplete stricture. (Option D)

• Urethroplasty - Excision of the stenosed length of the urethra and re-anastomosis is done. End-end anastomoses are done for strictures that are short and complete, while for a long complete stricture, a buccal mucosa graft can be used to bridge the gap. It is considered when a stricture is due to trauma or recurred following endoscopic treatment.

• Urethral dilation - The urethra is stretched using urethral dilators. This is a blind procedure and it can cause injury. Hence, it is not done frequently. (Option A)

• Intermittent self-dilatation - In patients with recurrent strictures who cannot undergo urethroplasty, intermittent self-dilatation with soft hydrophilic catheters is done daily.

• Metallic catheter placement - This is usually done for palliative care in end stage malignant diseases and recurrent cases. (Option C)

Solution to Question 37:

The procedure in the given image shows a 360-degree wrap, which is known as Nissen's fundoplication; this procedure is done for GERD. Nissen's fundoplication is a complete fundoplication procedure.

Fundoplication is the procedure in which the fundus of the stomach is wrapped around the lower end of the esophagus. In the Nissen's fundoplication procedure, the fundus is wrapped the entire 360 degrees around the esophagus.



The most common complication of Nissen's fundoplication is the gas-bloat syndrome. In this condition, patients are unable to belch gas from the stomach into the oesophagus. This causes increased gas in the stomach, resulting in a bloating sensation, abdominal discomfort, and flatulence.

Other options:

Option A: The toupet procedure is the re-creation of a sphincter around the EGJ done by wrapping the fundus around the oesophagus. A posterior 180-270-degree wrap is used.

Option B: The Dor procedure is fundoplication, most commonly used in the setting of the esophagal myotomy. It consists of an anterior 180-degree wrap.

Option C: Watson's fundoplication is a 90-degree anterior fundoplication. It is reported to achieve good reflux control while permitting burping and active vomiting and reducing gas bloat.



Orthopedics AIIMS 2017

Question 1:

A 5 year old boy fell on his outstretched hand. X-ray is shown below. Which of the following vessel is most commonly affected?

a) Radial artery

b) Brachial artery

c) Ulnar artery

d) Cubital vein

Question 2:

True regarding Galeazzi fracture-dislocation:

a) Radial collateral ligament tear with interosseous membrane tear with radial shaft fracture

b) Interosseous membrane tear with ulnar shaft fracture

c) Interosseous membrane tear with triangular fibrocartilage complex (TFCC) tear and radial shaft fracture

d) Interosseous membrane tear with triangular fibrocartilage complex (TFCC) tear and ulnar shaft fracture

Question 3:

A 25-year-old male presented to the emergency department with pain and swelling of the right knee. The radiograph showed the following finding. What is the best treatment for the given condition?

a) Tension band wiring with K-wires

b) Tension band wiring with cancellous screws

c) Cylinder cast

d) Patellectomy

Question 4:

Which of the following movements is being tested in the picture given below?

a) Internal rotation of hip

b) Abduction

c) Flexion

d) External rotation of hip

Question 5:

Which is the classification system used for the fracture shown in the x-ray given below?



a) Schatzker classification

b) Ideberg classification

c) Neer classification

d) Gartland classification

Question 6:

An elderly male presented with a history of fall on outstretched hand following which he developed swelling of the wrist. His X-ray is shown below. What is your diagnosis?



a) Fracture scaphoid

b) Fracture scaphoid with perilunate dislocation

c) Fracture hamate

d) Fracture capitate

Question 7:

Gustillo anderson classification type IIIb is?

a) Open fracture, wound more than 1 cm , less than 10 cm in length without extensive soft tissue damage, flaps, avulsions

b) Open fracture, wound less than 1 cm in length without extensive soft tissue damage, flaps, avulsions

c) Open fracture, wound more than 10 cm in length with extensive soft tissue loss and periosteal stripping and bone damage

d) Open facture, wound more than 10 cm in length associated with an arterial injury

Question 8:

An overweight man (90 kg) presented with complaints of not being able to squat and rest pain of left hip for 6 months duration. He had a past history of steroid and creatine use. His X-ray revealed destruction of the articular surface of femoral head with the presence of subchondral cysts and positive crescent sign. His MRI is suggestive of marrow edema. What is your diagnosis?

a) Tuberculosis of hip

b) Fracture neck of femur

c) Osteochondroma of femur

d) Osteonecrosis of femoral head

Question 9:

All of the following are associated with Sprengel shoulder except?

a) Diastematomyelia

b) Klippelfeil syndrome

c) Dextrocardia

d) Congenital scoliosis

Question 10:

A man with a POP cast for forearm fracture is prescribed analgesics. The earliest way to check for compartment syndrome is to look for:

a) Disappearance of pulse by displacing the cast

b) Discolouration of fingers

c) Decreased response to analgesics

d) Odour and discharge

Question 11:

Which of the following is not true about the posterior cruciate ligament:

a) Extra synovial

b) Prevent the posterior displacement of tibia

c) Primary action is to prevent the internal rotation of knee joint

d) Attached to anterolateral aspect of medial condyle

Question 12:

A patient presented with lower limb weakness. The examiner places one hand under the patient's heel and the patient is then asked to raise his other leg against downward resistance. What is the test being performed?

a) Hoover's Test

b) Mc Bride Test

c) Waddell's Test

d) O'Donoghue's Test

Question 13:

March fracture is seen in the ____ of second metatarsal bone.

a) Head

b) Neck

c) Shaft

d) Base

Question 14:

Identify the name of the instrument given below:



a) Bent Hohmann retractor

b) Bone holding forceps

c) Bone distractor

d) Bone curette

Question 15:

A patient came to the casualty after a RTA with fracture of femur. On 3rd day of admission, he developed sudden breathlessness. Which of the following is the most common cause?

a) Hypovolemic shock

b) Pulmonary hypertension

c) Post trauma MI

d) Fat embolism

Question 16:

Hand, knee gait in polio paralysis is due to weakness of the _____.

a) Popliteus

b) Quadriceps

c) Gastro-soleus

d) Gluteus maximus

Question 17:

The most common sequela of traumatic dislocation of the shoulder in young adults is_____.

a) Subscapular tendinitis

b) Rotator cuff injury

c) Recurrent dislocation of shoulder

d) Frozen shoulder

Question 18:

After a fracture, the growth of callus will be increased by which of the following?

a) Strict immobilization

b) Repeated tiny movements at the fracture site

c) Intermedullary nail

d) K wire fixation

Question 19:

Haglund deformity is seen around which of the following joints?

a) Knee

b) Ankle

c) Wrist

d) Elbow

Question 20:

An elderly patient slipped in the bathroom and sustained injury over the left hip. The radiograph is shown below. His attitude of the leg will be:

a) Shortened and abducted

b) Lengthened and internally rotated

c) Shortened and externally rotated

d) Flexed, adducted and internally rotated

Answer Key

Question No.	Correct Option
1	b
2	c
3	a
4	d
5	c
6	a
7	c
8	d
9	c
10	c
11	c
12	a
13	b
14	b

15	d
16	b
17	c
18	b
19	b
20	c

Detailed Explanations

Solution to Question 1:

The given X-ray shows a supracondylar fracture of the humerus and injury to the brachial artery is the most common vascular complication.

The fracture most commonly caused by a fall on an outstretched hand in a child is the supracondylar fracture of the humerus. As the hand strikes the ground, the elbow is forced into hyperextension resulting in a fracture of the humerus above the condyles.

• The brachial artery is usually injured by the sharp edge of the proximal fragment.

• Sometimes flexor muscles of the forearm - flexor pollicis longus and flexor digitorum profundus (medial-half) may suffer ischemic damage leading to Volkmann's ischemia leading to compartment syndrome.

While injury to the brachial artery is the commonest vascular complication, however, malunion is the most common complication of supracondylar fracture of the humerus.



Other delayed complications:

• Malunion - Cubitus varus / Gunstock deformity

• Volkmann's ischaemic contracture

• Myositis ossificans

Solution to Question 2:

Galeazzi fractures consist of a fracture of the radial shaft (lower one-third) with dislocation of the distal radioulnar joint (DRUJ).

Triangular Fibrocartilage Complex (TFCC)

The TFCC serves as the medial continuation of the distal articular surface of the radius as well as a static stabilizer of the DRUJ.

It consists of:

• Articular disc

• Dorsal radioulnar ligament - primary stabilizer

• Palmar radioulnar ligament - primary stabilizer

• The meniscus homologue

• The ulnar collateral ligament

• The sheath of the extensor carpi ulnaris

Other options:

Option A: Mentions radial collateral ligament but TFCC consists of the ulnar collateral ligament.

Options B and D: Mention ulnar shaft fracture when Galeazzi fracture is a radial shaft fracture.

Note: Galeazzi # are more common than Monteggia # and may be associated with ulnar nerve damage.

Solution to Question 3:

The radiograph shows a displaced patellar fracture. The best treatment for the given condition is tension band wiring with K- wires. In this method, two K-wires are used to transfix the reduced patellar fragments and a flexible wire is looped around them tightly.

Treatment options for patellar fracture are:

• Undisplaced patellar fractures with minimal joint incongruity and intact extensor retinaculum - cylinder cast

• Displaced patellar fractures - tension band wiring

• Severely comminuted fractures with difficulty in reducing fragments - patellectomy

Note: Tension band wiring with cancellous screw is done when compressive forces are also needed along with tensile forces to resist the displacement of fractures e.g., olecranon fractures.

Solution to Question 4:

External rotation of hip is being tested in the given image.

Rotatory movements of the hip joint can be measured with the knee flexed or extended. However, more precise measurements are obtained when the rotation is tested with the flexed knee. The procedure involves the following steps.

• The patient is made to lie supine.

• The examiner holds the knee with one hand and ankle with the other hand.

• The knee and hip are flexed to 90 degrees.

• Rotation movement is produced at the hip joint by stabilising the knee and moving the leg as the lever.

• If the hip is rotated such that the leg moves medially, external rotation of the hip occurs.

• If the hip is rotated such that the leg moves laterally, internal rotation of the hip occurs.



Solution to Question 5:

The given image is of the fracture of the proximal humerus. The most widely accepted classification is the Neer classification, which involves 4 major segments in injuries:

- Head of humerus
- Greater tuberosity
- Lesser tuberosity
- Shaft of humerus

Neer classi■cation distinguishes between the number of displaced fragments, with displacement de■ned as greater than 45 degrees of angulation or 1 cm of separation.



Other options

Option A: The Schatzker classification is a method of classifying tibial plateau fractures.

Option B: The Ideberg classification is a method of classification of intra-articular glenoid fossa fractures.

Option D: The Gartland classification is a method of classification of supracondylar fractures of the humerus.

Solution to Question 6:

The above X-ray shows fracture scaphoid.

Scaphoid fractures account for almost 75 percent of all carpal fractures although they are rare in the elderly and in children.

Mechanism of injury- The combination of forced carpal movement and compression, as in a fall on the dorsiflexed hand.

Diagnostic sign- precisely localized tenderness in the anatomical snuff box.

X-ray-

• Anteroposterior, lateral and oblique views are all essential

• often a recent fracture shows only in the oblique view.

• Usually, the fracture line is transverse, and through the narrowest part of the bone (waist), but it may be more proximally situated (proximal pole fracture).

Old, un-united fractures have hard borders, making it seem as if there is an extra carpal bone.

• Relative sclerosis of the proximal fragment is pathognomonic of avascular necrosis.

NOTE: The distal fragment, unrestrained by the scapholunate ligament, flexes and the proximal fragment tilt dorsally with the lunate (a DISI deformity). The hump-backed deformity of the scaphoid is permanent.

Solution to Question 7:

An open fracture with a wound with extensive soft tissue loss and periosteal stripping and bone damage belongs to classification IIIB.

Solution to Question 8:

The complaints of hip pain in an overweight man with a history of long-term steroid use along with the radiological findings of subchondral cysts and positive crescent sign in the femoral head are characteristic of osteonecrosis of the femoral head.

Avascular necrosis or osteonecrosis is a painful bone condition that results from diminished blood supply mainly due to trauma or non-traumatic etiologies. Alcohol, corticosteroids, immunosuppressives, and cytotoxic drugs, either singly or in combination, are the most common causes of non-traumatic osteonecrosis. The typical features are pain with movements in or near the joint, click in the joint, joint stiffness, limping gait, and deformity. Restriction of movements is also a feature of this condition.

An x-ray shows reactive new bone formation, increased radiolucency in the subchondral bone, and a distorted articular surface. The crescent sign is a subchondral fracture overlying the necrotic segment of the femoral head seen on radiography. MRI is the investigation of choice in early disease. It allows the identification of AVN of the femoral head and also determines the exact stage and extent of the pathologic process.



Management is based on Ficat and Arlet Classification:

Other options:

Option A: Tuberculosis of the hip is characterized by painful and restricted movements of the hip, especially at night. The radiographic findings are not appreciable in the early stages. Later stages show periarticular osteoporosis, hazy and irregular joint margins with a reduction in the joint space, and wandering acetabulum.

Stage	Findings	Management	Method
1	X-ray normal, MRI +ve	Conservative	Bisphosphonates
2a	X-ray: sclerosis, cysts, no collapse	Conservative	Bisphosphonates
2b	X-ray: crescent sign, no collapse	Surgical decompression	Core decompression
3	Collapse+loss of sphericity	Surgical	Osteotomy
4	Advanced arthritis	Reconstructive	Total hip replacement

Option B: Fracture neck of femur is common in elderly people after a trivial fall due to osteoporosis. Examination reveals tenderness in the groin along with shortening and external rotation of the leg. Xrays show a break in Shenton's line and trabecular stream, overriding of greater trochanter so that it lies at the level of the head of the femur, and external rotation of the femur is evident by prominent lesser trochanter.



Option C: Osteochondromas are common benign bone tumors commonly present in adolescents. The lesions consist of a bony mass, often in the form of a stalk, produced by endochondral ossification of a growing cartilaginous cap. They are usually painless. The common sites of involvement are the distal femur, proximal tibia, and proximal humerus.



Solution to Question 9:

Dextrocardia is not associated with Sprengel deformity.

Sprengel deformity is the congenital elevation of the scapula. It is the most common congenital shoulder abnormality.

The scapula is abnormally small and too high. Sometimes the clavicle is affected as well. Shoulder movements may be restricted and on abduction or elevation, the scapula moves very little or not at all.

Sprengel's deformity may be associated with other defects of the cervical spine (e.g. Klippel–Feil syndrome), and high thoracic kyphosis or scoliosis is quite common.

The high scapula may still be attached to the spine by a tough ■brous band or a cartilaginous bar (the omovertebral bar). Associated vertebral or rib anomalies are quite common.

An unappreciated association between Sprengel anomaly and diastematomyelia (congenital splitting of spinal cord) of the lumbar spine also raises questions about the embryologic origin of the scapula.

Associations of Sprengel deformity of Shoulder

• Klippel-Feil Syndrome

• Spina Bifida and Diastematomyelia

• Kyphosis or Scoliosis

• Torticollis

• Underdevelopment of clavicle or humerus

Solution to Question 10:

Decreased response to analgesia → suspect compartment syndrome.

Pain is the most common initial symptom in case of compartment syndrome. Red flags for compartment syndrome are pain out of proportion with the injury, and pain on passive stretch of the digits.

Later signs include pallor, paresthesia, paralysis, and pulselessness.

As the ischemia occurs at capillary level in compartment syndrome, pulses may still be felt and skin may not appear pale.

Solution to Question 11:

The lateral collateral ligament, not the posterior, is responsible for preventing the internal rotation of the knee joint

The posterior cruciate ligament (PCL) is an intra-articular but extra-synovial structure because of the presence of a synovial sheath that lines the ligament. It is the only primary restraint to posterior translation throughout the entire knee range of motion. It prevents the internal rotation of the tibia on the femur when there is varus stress on the knee.

It acts as a secondary restraint to valgus/varus, external rotation.

Anterior cruciate ligament	Posterior cruciate ligament
Attached to anterior part of t he intercondylar area of tibia	Posterior part of intercondyla r area of tibia
Direction – Upwards, backwa rds & laterally & attached to posterior aspect of medial su rface of lateral condyle.	Upwards, forwards & mediall y & attached to the anterior a spect of the lateral surface of the medial condyle.
Prevents posterior dislocatio n of femur on tibia and forwa rds dislocation of tibia on fe mur.	Prevents anterior dislocation of femur on tibia and posteri or dislocation of tibia on fem ur
Limits hyperextension of kne e joint(taut during extension)	Limits hyperflexion of knee j oint. (taut during flexion)
Weaker	Stronger



Solution to Question 12:

The test being performed by the examiner is Hoover's test. This test is useful in differentiating between the organic and functional causes of limb weakness.

Hoover's test involves the following steps:

• The examiner places both their hands under the heels of the patient.

• The patient is asked to flex the affected hip while keeping the knee extended.

• Downward pressure in the healthy limb is felt by the examiner in an organic cause of limb weakness and in healthy individuals. This pressure is not felt in the functional cause of limb weakness and malingering.

In a normal individual, when raising one leg by flexing at the hip, the other limb exerts a downward pressure which can be felt at the ankle.



The McBride's test, Waddell's test, and O'Donoghue's test are other examination techniques that help in distinguishing between organic and non-organic causes of lower back pain.

Solution to Question 13:

March fracture/stress fracture is seen most commonly in the neck of the second metatarsal bone.

It results from long, repeated stresses like prolonged walking or running, usually seen in military men, athletes, and dancers. They may complain of gradual onset of pain over the 2nd metatarsal neck starting 2-4 weeks after the beginning of stressors. On examination, a tender lump may be noted. The x-ray may be normal initially but later show a hairline crack. After a period of 4-6 weeks, a mass of callus can be identified.

Management: Reduction and splinting are not necessary as it is an undisplaced fracture. Normal walking is permitted while using an elastic bandage on the forefoot.

The below image shows a March fracture of the second metatarsal bone.

Note: According to Campbell orthopedics (14th edition) in March fractures, patients often note the gradual onset of pain directly over the second metatarsal neck region. In Uptodate, it is mentioned that many of the stress fractures occur in the second metatarsal shaft, especially at the neck. Hence, neck is considered a more correct answer than shaft.

Solution to Question 14:

The image shows a bone holding forceps.

Solution to Question 15:

History of sudden onset breathlessness on the 3rd day of admission after femur fracture should raise suspicion of fat embolism.

Fat embolism is the presence of fat globules in vital organs and peripheral circulation after a fracture of the long bone or other major trauma. Its pathogenesis can be related to 2 factors: the release of free fatty acids by the action of lipases on the neutral fats and the actual obstruction of small pulmonary vessels by fat globules.

Clinical features:

• Early signs are fever and tachycardia

• Presenting features are of 2 types -

• Cerebral type - drowsiness, restlessness, disorientation

• Pulmonary type - tachypnoea, tachycardia

• Classical triad -

• Respiratory distress

- Neurological abnormalities
- Petechial hemorrhages
- Pathognomonic signs are petechiae on the trunk, axillae, conjunctival folds, and retina.

Treatment of fat embolism is mainly supportive as given below:

- Respiratory support with oxygen
- Heparinization increases lipoprotein lipase activity
- Intravenous low molecular weight dextran
- Corticosteroids to avoid pneumonitis
- Prompt stabilization of long-bone fractures

Solution to Question 16:

Polio commonly affects the quadriceps. The weakness of the quadriceps leads to the buckling (bending) of the knee while walking or bearing weight on the leg. So, in order to walk, the person has to prevent the buckling of the knee by supporting with the hand, producing a hand-knee gait.

Few important examination findings in paralytic polio:

- Asymmetric: involvement of affected muscles is haphazard.
- Most common muscle affected: Quadriceps, often partially paralyzed.
- Most common muscle to undergo complete paralysis: Tibialis anterior
- Most common muscle in hand: Opponens pollicis
- Motor paralysis is not associated with sensory loss.
- In the knee, triple deformity in severe cases: flexion, posterior subluxation, and external rotation

Solution to Question 17:

The most common sequela of traumatic shoulder dislocation in young adults is recurrent dislocations of the shoulder joint.

Traumatic shoulder dislocation is associated with tear of the glenoid labrum (Bankart lesion). This results in loss of attachment of the inferior glenohumeral ligament, which is the primary static stabilizer of the shoulder.

In addition, there is a loss of concavity due to damage to the labrum. Therefore, the shoulder joint becomes unstable. This predisposes the person to repeated shoulder dislocations. Shoulder instability/recurrent dislocation is an indication for surgical intervention.

Solution to Question 18:

Micro-movements at the fracture site stimulate callus formation.

Other options:

Option A: It is not necessary to immobilize all fractures (e.g., fracture ribs, scapula, etc) as they heal anyway. Some other fractures need strict immobilization (e.g., fracture of the neck of the femur) and may still not heal.

Options C, D: Kirschner wires (K-wire) and intra-medullary nails are used when a fracture is so unstable that it is difficult to maintain it in an acceptable position by nonoperative means. They are used in order to secure rigid immobilization. They do not affect the growth of the callus directly.

Solution to Question 19:

Haglund deformity, pump bump, or Mulholland deformity is seen around the ankle joint. It refers to retrocalcaneal exostosis formed at the posterosuperior aspect of the calcaneum.

retrocalcaneal bursa usually protects the Achilles tendon from the calcaneal tuberosity. But in Haglund deformity, there is chronic inflammation of this retrocalcaneal bursa due to increased or repetitive chafing between the tuberosity and Achilles tendon. Due to prolonged inflammation, degenerative changes occur and osteophytes form within the tendon leading to calcification at the site of the tendon insertion. This leads to deformity and pain symptoms.

Management is usually conservative, using a heel lift, silicone pads in closed shoes, ice application, anti-inflammatory drugs, and stretching exercises of the calf.

The image below shows the Haglund deformity.

Solution to Question 20:

The history of fall and hip injury in an elderly patient along with the radiological findings of a broken Shenton's line is suggestive of a neck of femur fracture. In this condition, the attitude of the leg will be shortened and externally rotated and movements of the hip joint will be painful.

A plain radiograph may show the following features:

• Fracture line in the neck of femur

• External rotation of the femur with the lesser trochanter appearing more prominent

• Displacement of the greater trochanter

• Break in the trabecular stream

• Break in Shenton's line



Orthopedics AIIMS 2018

Question 1:

The X-ray image of a patient that presented to the casualty is given below. How will you manage this case?

a) Nailing

b) Plating

c) Tension band wiring

d) Knee splint

Question 2:

A 55-year-old female patient presented with a flexion contracture of the hip. Which of the following tests is useful in diagnosis?

a) Allen's test

b) Trendelenburg test

c) Thomas test

d) Ober test

Question 3:

A 10-year-old came to the OPD with the following deformity of his left upper limb. What is the most likely cause of the fracture?



a) Supracondylar Fracture

b) Olecranon Fracture

c) Radial Head fracture

d) Triad of elbow

Question 4:

A 4-year-old child was brought to the OPD with a painless deformity. A radiograph of the child's legs is shown below. The most common cause of this deformity is _____.



a) Skeletal dysplasia

b) Rickets

c) Scurvy

d) Blount disease

Question 5:

A 25 year old man involved in a fight, injured his finger and the deformity is shown in the following image. X-ray is normal. The most appropriate management at this stage is:

a) Splint the finger in flexion

b) Buddy strapping

c) Surgical repair of the flexor tendon

d) Splint the finger in hyperextension

Question 6:

What is your diagnosis from the below X-ray images?

a) Solitary bone cyst

b) Aneurysmal bone cyst

c) Giant cell tumor

d) Osteoblastoma

Question 7:

Which of the following is true regarding the scar tissue in patellar clunk syndrome?

a) Superior pole of patella, impinging during flexion

b) Superior pole of patella, impinging during extension

c) Inferior pole of patella, impinging during flexion

d) Inferior pole of patella, impinging during extension

Question 8:

The earliest symptom of acute compartment syndrome is _____

a) Pain

b) Pallor

c) Paresthesia

d) Paralysis

Question 9:

Identify the following marked structures.



a) 1 - Sequestrum, 2 - Involucrum, 3 - Cloacae

b) 1 - Sequestrum, 2 - Cloacae, 3 - Involucrum

c) 1 - Involucrum, 2 - Cloacae, 3 - Sequestrum

d) 1- Cloacae, 2 - Sequestrum, 3 - Involucrum

Question 10:

Which tumor does not cause osteomalacia and phosphaturia?

a) Osteosarcoma

b) Hemangiopericytoma

c) Fibrosarcoma

d) Breast carcinoma

Question 11:

A polytrauma patient with femur fracture and massive hemorrhage presents to the emergency department after a road traffic accident. Which of the following is not an immediate treatment consideration in this patient?

a) Administration of IV crystalloids

b) Administration of tranexamic acid

c) Check coagulation with thromboelastography

d) Internal fixation of femur fracture

Question 12:

What is the most common form of arthritis?

a) Rheumatoid arthritis

b) Psoriatic arthritis

c) Seronegative arthritis

d) Osteoarthritis

Answer Key

Question No.	Correct Option
1	c
2	c
3	a
4	b
5	d
6	c
7	b
8	a
9	c
10	d
11	d
12	d

Detailed Explanations

Solution to Question 1:

The given radiograph shows a transverse fracture of the patella which should be managed by tension band wiring.

Patellar fractures can be undisplaced, transverse, comminuted, or longitudinal. The radiological examination includes AP and lateral views. A skyline view of the patella is required in some cases of undisplaced fractures.

Management of patellar fractures:

• Undisplaced fractures: Treated with a cylinder cast extending from the groin to just above the malleoli, with the knee in full extension, left for at least 3 weeks.

• Transverse fractures: Surgical reduction using tension-band wiring and K-wire fixation, followed by extensor retinaculum repair and physiotherapy.

• Comminuted fractures: Open reduction and fixation with patellectomy. Patella-saving operations are also being performed with improved fixation techniques.

• Longitudinal fractures: As the extensor apparatus is usually intact, they can be treated with a cylinder cast or extension brace initially, followed by range-of-motion exercises.

Solution to Question 2:

Thomas test is useful in diagnosing flexion contracture of the hip.

Flexion deformity is a common hip deformity, often caused by stronger flexor muscles overpowering the extensor muscles. When there is a spasm of the muscles, the stronger flexors pull the hip in flexion. Soft tissue contractures can convert this into a fixed flexion deformity, making locomotion impossible.

The Thomas test is used to evaluate the degree of flexion deformity. The aim is to eliminate compensatory lumbar lordosis and reveal the obvious flexion deformity for measurement.

During the test, the patient lies supine on a hard surface with legs straight. The patient may compensate for the flexion deformity by arching the lower back, creating excessive lumbar lordosis. The easy passage of the examiner's hand between the bed and the back of the patient is considered a positive test as normally this is not possible.

By gradually flexing the unaffected hip, the pelvis tilts and eliminates lumbar lordosis, causing the affected hip to assume the flexed position. The angle between the affected thigh and the bed indicates the degree of flexion deformity.



Other options:

Option A: Allen's Test is mainly used to test the adequacy of ulnar collateral blood flow to the hand, which is usually done prior to radial artery cannulation.



Option B: Trendelenberg test is used to assess the weakness or paralysis of the gluteus medius muscle.



Option D: Ober's test is used in physical examination to identify the tightness of the iliotibial band.

Solution to Question 3:

The given image shows cubitus varus deformity, where the forearm has deviated medially in relation to the upper arm, which is a common complication of supracondylar humeral fracture.

Malunion is the commonest complication of a supracondylar fracture and results in a cubitus varus deformity, also known as gunstock deformity. This occurs when the fracture unites with the distal fragment tilted medially and in internal rotation.

The image below shows gunstock deformity:

Cubitus varus is primarily a cosmetic concern with minimal functional impairment. Mild cases may not necessitate treatment, while severe deformities should be corrected through a lateral closed wedge osteotomy (French osteotomy).

Other options:

Option B: Olecranon fracture results from a direct injury as in a fall onto the point of the elbow. Complications include non-union, elbow stiffness, and osteoarthritis

Option C: Radial head fracture results from a fall on an outstretched hand and presents with pain over the lateral aspect of the forearm near the elbow along with restricted elbow movements, pronation, and supination. Complications include posterior interosseous nerve injury, myositis ossificans, and osteoarthritis.

Option D: Terrible triad of elbow refers to elbow dislocation with fracture of the radial head, coronoid process, and medial collateral ligament rupture.

Solution to Question 4:

The given X-ray shows bow legs or genu varum which is most commonly seen in rickets.

Rickets is a bone mineralization disorder that primarily affects children with open growth plates, resulting from a deficiency of vitamin D, calcium, or phosphate.

In toddlers, genu varum is one of the most common initial sign of rickets. In older children, genu valgum or coxa vara might be present.



Solution to Question 5:

Based on the mechanism of injury and the deformity characterized by flexion at the distal interphalangeal joint, along with a normal radiograph, the likely diagnosis is mallet finger, which is managed by splinting the finger in hyperextension.

Mallet finger, also known as baseball finger, occurs when there is a sudden passive flexion of the distal interphalangeal joint, causing avulsion of the extensor tendon from its insertion at the base of the distal phalanx.



Surgery is considered only if there is a large fragment (>50%) and subluxation of the DIP joint.

Solution to Question 6:

The given radiographs show the anteroposterior and lateral views of the wrist, demonstrating a solitary lytic lesion with cortical expansion and a soap-bubble appearance in the epiphysis of the distal radius, suggestive of a giant cell tumor (GCT).

GCT usually affects young adults aged 20 to 40 years, with slight female predominance. It is a benign but locally aggressive tumor. It is exclusively seen in adults and affects the ends of long

bones, typically involving the epiphysis. The most common location of the tumor is the distal femur followed by the proximal tibia. Involvement of the distal end of the radius is also seen and the tumor is more aggressive in this location.

Important X-ray features of a GCT:

• A solitary, possibly loculated, lytic lesion in the epiphysis

• Eccentric location, most often sub-chondral

• Expansion of the overlying cortex

• Soap bubble appearance (the tumor is homogenously lytic with trabeculae of the remnants of bone traversing it, giving it a loculated appearance).

• No calcification within the tumor

• No or minimal reactive sclerosis around the tumor

• Thinned out cortex, may be perforated

The classical "soap bubble" appearance of a giant cell tumor (GCT) is shown below:

Other options:

Option A: A solitary bone cyst is most commonly occurs in the metaphysis of the bone. On an X-ray, a bone fracture fragment may be observed resting in a dependent portion of the cyst, referred to as the fallen fragment or fallen leaf sign. This sign is considered pathognomonic of a simple bone cyst.

The image below shows a bone cyst in the humerus along with a pathological fracture and fallen leaf sign:

Option B: On an X-ray, an aneurysmal bone cyst is seen as an eccentric, expansile, lytic lesion with well-defined margins in the metaphysis.

Option D: Osteoblastoma is a benign bone tumor usually occurring in the jaw and spine. On X-ray, a radiopaque central nidus with a surrounding radiolucent halo is seen. If occurring in the long bones, it involves the diaphysis or metaphysis, but never in the epiphysis.

The image below shows an osteoblastoma in C7 vertebrae:

Solution to Question 7:

In patellar clunk syndrome, the scar tissue which is formed at the superior pole of the patella becomes impinged during knee extension.

Patellar clunk syndrome is seen after posterior-stabilized knee arthroplasty. A fibrous nodule forms on the posterior surface of the quadriceps tendon just above the superior pole of the patella. This can become entrapped in the intercondylar notch of the femoral prosthesis and cause the knee to "clunk" at 30- 45 degrees to the femoral axis while the patient actively extends the knee.

The recommended treatment is arthroscopic debridement of the nodule.

Solution to Question 8:

The earliest symptom of acute compartment syndrome is pain.

Acute compartment syndrome is a surgical emergency caused by increased pressure within a fascial compartment, leading to tissue ischemia. It is characterized by rapid progression of pain and swelling in an extremity, commonly precipitated by traumatic injury.

The classic features of ischemia in acute compartment syndrome include the 5 P's:

• Pain - It is the first symptom of acute compartment syndrome. Ischemic muscle is highly sensitive to touch. On passive stretching of the muscles, severe pain can be elicited.

• Paresthesia - It is the first sign of nerve ischemia.

• Pallor - It may not be seen always. The skin outside the compartment may be normally perfused, as only the intra-compartmental perfusion may be affected.

• Paralysis - Paralysis of muscle groups affected in that compartment is a late sign.

• Pulselessness - Usually, the peripheral pulses and capillary return are intact in the early stages, unless there is a major arterial injury. If pulses are absent, arteriography is indicated.

The diagnosis can be confirmed by measuring the intra-compartmental pressure using a split catheter introduced into the compartment. The difference between diastolic pressure and compartment pressure is known as differential pressure (ΔP). A differential pressure (ΔP) of less than 30 mm Hg (4.00 kilopascals) is an indication for immediate compartment decompression by performing a fasciotomy.

Solution to Question 9:

In the given radiographs showing features of chronic osteomyelitis, the marked structures are as follows:

• Involucrum is the thick sheath of periosteal new bone that surrounds the sequestrum.

• Cloacae are openings in the involucrum that allow drainage of purulent and necrotic material out of the dead bone.

• Sequestrum is a piece of devitalized bone that has been separated from its surrounding bone during the process of necrosis.

Solution to Question 10:

Breast carcinoma does not cause osteomalacia and phosphaturia.

Tumor-induced osteomalacia (TIO) is a rare paraneoplastic syndrome which presents with hypophosphatemia, osteomalacia and renal phosphate wasting. It is most commonly seen with mesenchymal tumors.

Tumors associated with osteomalacia are:

- Hemangiopericytoma (option B)
- Non-ossifying fibroma
- Osteosarcoma (option A)
- Osteoblastoma
- Fibrosarcoma (option C)

Solution to Question 11:

The internal fixation of a femur fracture is not an immediate concern in a patient with polytrauma and hemodynamic instability secondary to severe blood loss. Instead, splinting is preferred until the patient has been stabilized.

Damage Control Orthopaedics (DCO) in unstable/borderline polytrauma patients involves 3 components:

- Stage 1: Resuscitative procedures for rapid hemorrhage control
- Stage 2: Restoration of normal physiologic parameters
- Stage 3: Definitive surgical management

The first stage is early temporary stabilization of unstable fractures and the control of the bleeding. In femur fracture, the most suited approach is temporary stabilization using an external fixator.

The second stage involves resuscitation and stabilization of the patients in the ICU.

Finally, the third stage involves delayed definitive fracture management after the patient's condition becomes suitable. In femur fracture, the definitive procedure is intramedullary nailing which is performed after the patient becomes stable.

Solution to Question 12:

Osteoarthritis is the most common type of arthritis.

Osteoarthritis is strongly associated with age and is highly prevalent among older individuals. Studies suggest that over 80% of people aged 55 and above have osteoarthritis in at least one joint.

Orthopedics AIIMS 2019 (May & Nov)

Question 1:

Following a fall in the bathroom, an elderly lady is diagnosed with a fractured neck of the femur. Which of the following coexisting conditions is most likely to have predisposed to this injury?

a) Osteoporosis

b) Osteopetrosis

c) Osteoarthritis

d) Osteomalacia

Question 2:

At what age in women is a DEXA scan done routinely?

a) At 50 years

b) At 55 years

c) At 60 years

d) At 65 years

Question 3:

Most specific feature for ankylosing spondylitis:

a) HLA-B27 positivity

b) Bilateral sacroiliitis

c) Elevated ESR

d) Lumbar laxity

Question 4:

The treatment for the condition given below is:

a) Extended curettage with allograft

b) Bone biopsy

c) Curettage

d) Extended curettage with autograft

Question 5:

Identify the instrument given below.

a) Bone nibbler

b) Bone curette

c) Plate holding forceps

d) Bone holding forceps

Question 6:

A 30-year-old man presents with numbness of both hands. He was asked to perform the test demonstrated below. What is this test?

a) Adson's test

b) Pemberton test

c) Roos test

d) Wright test

Question 7:

The test performed in the following video is



a) Allen's test

b) Roos test

c) Adson's test

d) Wright test

Question 8:

A child presents with difficulty in walking. On biopsy of the hip, some cells with grooved nuclei and eosinophilic cytoplasm are present. Which investigation would you order next?

a) MRI

b) Serum calcium

c) Serum PTH

d) CD1a marker

Question 9:

A 28-yr old man has recently finished a half- marathon. He complains of pain along the anterior and medial aspect of the tibia which increases on jogging & walking for a long time. Plain X-Ray radiograph of the leg was found to be normal. What is the probable diagnosis?

a) Jones fracture

b) Shin splint

c) Lisfranc fracture

d) Nutcracker fracture

Question 10:

The following X-ray shows a lytic lesion of the lower limb. The biopsy shows fibroblastic proliferation, osteoclast and inflammatory cells. From the given options, the most likely diagnosis that can be inferred from the above clinical picture is_____.

a) Aneurysmal bone cyst

b) Giant cell tumor

c) Chondroblastoma

d) Langerhan cell histiocytosis

Question 11:

Which of the following is used to stabilise a fracture of the distal femur as shown in the image below?



a) Rail fixator

b) Ilizarov fixator

c) K wire binding

d) Interlocking nail

Question 12:

What is the type of scoliosis shown in the X-ray given below?



a) Congenital scoliosis

b) Postural scoliosis

c) Infantile scoliosis

d) Degenerative scoliosis

Question 13:

The following instrument is used for:

a) Nibbling the ends of the bones

b) Holding the bone and plate

c) Controlled traction of fracture segments

d) Cutting the K- wire

Question 14:

A patient presents to the emergency department after he fell on his outstretched hand. An X-ray was performed and is given below. What is the diagnosis?

a) Colle's fracture

b) Rolando's fracture

c) Galeazzi fracture

d) Montegia fracture

Question 15:

Identify the following device.

a) Cock-up splint

b) Colles' cast

c) Knuckle bender splint

d) Dynamic finger splint

Answer Key

Question No.	Correct Option
1	a
2	d
3	b
4	d
5	a
6	c
7	a
8	d
9	b
10	b
11	a
12	a
13	b
14	c
15	a

Detailed Explanations

Solution to Question 1:

Given the clinical history of an elderly woman experiencing a fracture following a trivial fall, which indicates a pathological fracture, the most probable predisposing condition would be osteoporosis.

Osteoporosis is the most common metabolic bone disorder characterized by abnormally low bone mass, which makes the bone unusually fragile and increases the risk of fractures.

A DEXA scan is performed in the following patients to assess bone mineral density and screen for osteoporosis, and the results are reported in terms of T-score or Z-score:

• Females aged 65 years or more

• Males aged 70 years or more

• Post-menopausal females under the age of 65 years with risk factors

• Fracture after the age of 50 years

• Males aged 50-69 with risk factors

Solution to Question 2:

Dual-energy x-ray absorptiometry (DEXA scan) is routinely done in women at 65 years of age to screen for osteoporosis.

Osteoporosis is the most common metabolic bone disorder characterized by abnormally low bone mass, which makes the bone unusually fragile and increases the risk of fractures.

The criteria to screen for osteoporosis according to the national osteoporosis foundation is as follows:

• Females aged 65 years or more

• Males aged 70 years or more

• Post-menopausal females under the age of 65 years with risk factors

• Fracture after the age of 50 years

• Males aged 50-69 with risk factors

A DEXA scan is performed in the above patients to assess bone mineral density, and the results are reported in terms of T-score or Z-score.

Solution to Question 3:

Bilateral sacroiliitis is the most specific sign of ankylosing spondylitis (AS).

While HLA-B27 is positive in up to 90% of patients with AS, it is not specific to AS (option A). Erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) are often, but not always, elevated and are not specific to AS (option C). Lumbar laxity is not a feature of AS (option D).

The Assessment of Spondyloarthritis International Society classification criteria for axial spondyloarthritis includes lower back pain for more than 3 months in patients less than 45 years of age and one of the following:

• Sacroiliitis confirmed on x-ray or MRI and at least one typical clinical or laboratory finding

• A positive HLA-B27 test and at least two typical clinical or laboratory findings

Clinical and laboratory findings used in the diagnostic criteria for AS include:

• Articular manifestations: inflammatory back pain, dactylitis, arthritis

• Extraarticular manifestations: enthesitis, uveitis, inflammatory bowel disease, psoriasis

• Patient history: positive family history, good response to NSAIDs

• Laboratory findings: positive for HLA-B27; elevated CRP and/or ESR

HLA-B27 is strongly associated with the following conditions:

• Ankylosing spondylitis

• Psoriatic arthritis

• Acute anterior uveitis

• IBD-associated ankylosing spondylitis

• Reactive arthritis

Solution to Question 4:

The provided radiograph of the knee joint shows an expansile, eccentric, subarticular, lytic lesion in the epiphysis of the distal femur with a soap bubble appearance, which suggests a giant cell tumor (GCT). The recommended approach for management is extended curettage and autograft.

GCT or osteoclastoma usually affects young adults in the age group of 20-40 years. The most common location of the tumor is the distal femur followed by the proximal tibia.

Important X-ray features of a GCT:

• A solitary, possibly loculated, lytic lesion in the epiphysis.

• Eccentric location, most often subchondral.

• Expansion of the overlying cortex.

• Soap bubble appearance - The tumor is homogeneously lytic with trabeculae of the remnants of bone traversing it, giving it a loculated appearance called soap bubble appearance.

• No calcification within the tumor.

• No or minimal reactive sclerosis around the tumor.

The image below shows an X-ray of a giant cell tumor showing the typical soap bubble appearance:

The treatment of choice for GCT is extended curettage and bone grafting.

• GCT in the relatively insignificant bone like the distal ulna, clavicle, and proximal fibula is treated by resection without reconstruction.

• Bisphosphonates have been shown to reduce the proliferation of osteoclasts and hence have been shown to stabilize local and metastatic disease.

• Radiation/embolization can be used for inoperable spinal or sacral tumors.

• Denosumab is approved for use in unresectable tumors in skeletally mature patients.



Solution to Question 5:

The instrument given is a bone nibbler, which is designed to remove or shape small sections of bone by gently nibbling or biting away at the bone tissue.

It is available in various sizes and with different angles of the nose:

• Straight nibbler – for general use.

• Curved nibbler – for spinal surgery.

• Double-action nibbler – straight or curved. The double-action reduces the force required for cutting the bone and is mechanically superior.

Solution to Question 6:

The test performed in the image is Roos test. It is also referred to as the elevated arm stress test (EAST). It is a provocative test for the diagnosis of thoracic outlet syndrome.

For this test, the patient abducts the shoulders to 90 degrees and flexes the elbow to 90 degrees. The patient is made to open and close the hand repeatedly for three minutes. The test is positive if

the patient is unable to complete the test or experiences cramping, pain, numbness, on the affected side.

Solution to Question 7:

The test performed in the given video is Allen's test.

Allen's test is mainly used to test the adequacy of ulnar collateral blood flow to the hand, which is usually done prior to radial artery cannulation.

Procedure:

• The radial and ulnar arteries are occluded as the patient makes a tight fist so as to exsanguinate the palm.

• Then the patient is asked to open the hand and the occlusion over the ulnar artery is released.



Interpretation:

• As occlusion of the ulnar artery is released, the color of the open palm is observed. Normally, the color will return to the palm within 5 – 6 seconds.

• If the palm remains pale for more than 6-10 seconds, it indicates reduced ulnar collateral flow.

• The modified Allen's test considers a threshold of 5 to 15 seconds.

Clinical significance:

• In recent years, studies have shown that, although the Allen test is often used to identify patients at increased risk for ischemic complications from radial artery cannulation, the predictive value of this test is poor.

• It is also used for identifying patients acceptable for radial artery harvest or use for coronary angiography.

Solution to Question 8:

Histopathologically, cells with grooved nuclei and eosinophilic cytoplasm can be observed in both Langerhans cell histiocytosis (LCH) and chondroblastoma. Therefore, the next step to differentiate between them would be to obtain a CD1a marker, which is typically positive in LCH.

The specific markers for LCH are CD1a and langerin (CD207)

It is worth noting that while an MRI can provide valuable information for evaluating the pathology, it is ideally performed before the biopsy.

Solution to Question 9:

The given clinical scenario, with a history of lower limb overuse (marathon running) and pain along the medial tibial border which increases with activity (jogging and long walks), along with a normal radiograph, suggests medial tibial stress syndrome (MTSS), also known as shin splints.

MTSS is a stress injury of the tibia which occurs due to the frequent overuse of lower limbs. It is commonly seen in athletes and military personnel.

Exercise-induced pain along the distal two-thirds of the medial tibial border is typical. Since it is a stress injury, the radiograph of the leg appears normal.

Management is mainly conservative with rest and activity modification with less repetitive and load-bearing exercises.

Solution to Question 10:

The provided X-ray reveals a lytic lesion in the epiphysis of the distal tibia, and the biopsy, which shows osteoclast cells, suggests a diagnosis of a giant cell tumor (GCT).

Important X-ray features of a GCT:

• A solitary, possibly loculated, lytic lesion in the epiphysis.

• Eccentric location, most often subchondral.

• Expansion of the overlying cortex.

• Soap bubble appearance - The tumor is homogeneously lytic with trabeculae of the remnants of bone traversing it, giving it a loculated appearance called soap bubble appearance.

• No calcification within the tumor.

• No or minimal reactive sclerosis around the tumor.

The image below shows an X-ray of a giant cell tumor showing the typical soap bubble appearance:

Histologically, GCT is characterized by the presence of a large number of osteoclast-like giant cells among scattered round or spindle-shaped mononuclear cells. The tumor stroma is well-vascularized with bands of cellular or collagenous fibrous tissue.

The image below shows the histopathology of GCT:



Other options:

Option A: Aneurysmal bone cyst usually occurs in the metaphysis of long bones and the posterior elements of vertebral bodies. It is characterized by multiloculated blood-filled cystic spaces separated by septae, composed of multinucleated osteoclast-like giant cells and reactive woven bone. Radiographically, an aneurysmal bone cyst is seen as an eccentric, expansile, lesion with well-defined margins. Most lesions are completely lytic and often contain a thin shell of reactive bone at the periphery.

Option C: Chondroblastoma is a benign cartilaginous tumor that occurs mainly in bones around the knee joint. On x-ray, the tumor appears as a well-circumscribed, round, or oval lytic lesion in the center of the epiphysis of long bones, with areas of calcification within the tumor, giving the characteristic mottled appearance. On histology presence of "chicken wire calcification" is pathognomic.

Option D: Langerhans cell histiocytosis commonly involves the skull, pelvis, femur, and mandible. Radiographs show a lytic lesion with periosteal new bone formation. In the skull, lesions appear as a hole within a hole, and in the vertebra, vertebra plana (flattening of the vertebral body) can be seen. Histopathologically, Langerhans cells have a single large nucleus with a characteristic folded or grooved nucleus (“coffee-bean” appearance) with small inconspicuous nucleoli. On electron microscopy, the presence of Birbeck granules in the cytoplasm is characteristic.

Solution to Question 11:

The above image shows a rail fixator.

In this type of external fixator, multiple pins are inserted into the bone. These pins are then attached to different blocks which can move over a track or rail.

Given below is an Ilizarov ring fixator. Here, pins are attached to an adjustable ring system.

Tension band wiring is used in the fixation of transverse patellar fractures. In this method, two K-wires are used to transfix the reduced patellar fragments and a flexible wire is looped around them tightly.

The following X-ray image shows tibial intramedullary interlocking nailing.



Solution to Question 12:

In the given radiograph, the presence of a single hemivertebra at T9 and concavity to the right in the lower thoracic region is suggestive of congenital scoliosis.

Congenital scoliosis is a lateral curvature of the spine which is always associated with a radiologically demonstrable anomaly of the vertebral bodies, which includes:

• Hemivertebrae (where only one-half of the vertebra grows)

• Block vertebrae (where two vertebral bodies fuse)

• An unsegmented bar (a bone bar joining two adjacent vertebrae on one side, which hinders growth on that side)

The image below shows a unilateral and unsegmented bar with contralateral hemivertebra:

In infantile scoliosis (option C), most curves are thoracic with convexity to the left and the anomalies described above are absent.

Note: The most progressive type of congenital scoliosis is a concave, unilateral unsegmented bar with a convex hemivertebra.

Solution to Question 13:

The image shown is of bone holding forceps (Verbrugge forceps). It is used to hold the bone and plate.

It consists of a ratcheted clamp with a curved distal end.

Solution to Question 14:

The provided history of falling on an outstretched arm and the X-ray revealing an oblique fracture in the lower third of the radius with dislocation of the distal radio-ulnar joint point to the diagnosis of Galeazzi's fracture.

Galeazzi's fracture is the fracture of the lower third of the radius and the subluxation or dislocation of the distal radioulnar joint. This injury usually happens when one falls on an outstretched hand.

Patients present with pain, swelling, and deformity at the lower end of the radius. On examination, the disruption of the radio-ulnar joint can be elicited by pressing the lower end of the ulna. The ulna moves down and comes back like the key of a piano (piano-key sign).

The aim of treatment for this fracture is to preserve supination and pronation of the forearm. In children, closed reduction is usually enough to achieve perfect reduction. However, in adults,

open reduction using a k-wire fixation or compression plating is usually required. It is also important to look for ulnar nerve damage.

Galeazzi's fracture can cause complications such as malunion due to displacement of the fragment. This often results in deformity and limitation in supination and pronation.

Solution to Question 15:

The device shown in the image is a cock-up splint. It is used to manage radial nerve palsy.

Radial nerve palsy manifests as an inability to dorsiflex the wrist and digits (wrist drop and finger drop). Numbness occurs on the dorsoradial aspect of the hand and the dorsal aspect of the radial 3 1/2 digits. The cock-up splint is designed to counteract the wrist drop position by holding the wrist in a slightly extended or neutral position, which helps prevent further contracture and allows for functional use of the hand.

Other options:

Option B: Undisplaced Colles' fracture is managed by immobilization in a below-elbow plaster cast for six weeks. Displaced fractures are managed by manipulative reduction followed by immobilization in Colles' cast in palmar flexion, pronation, and ulnar deviation.

Option C: Knuckle bender splint is used in cases of ulnar claw hand where ulnar nerve recovery is anticipated. The splint can be used to correct and prevent metacarpophalangeal hyperextension by maintaining flexion as shown below.

Option D: Dynamic finger splint can be used for:

• Immobilization in case of fracture of the proximal phalanx

• Extensor tendon injuries of the finger

• Prevention of deformities in rheumatoid arthritis.

• Post-operative immobilization.



Orthopedics AIIMS 2020 (May and Nov INI-CET)

Question 1:

All of the following are indications of surgery in a patient of Potts' spine except?

a) Cold abscess without neurological complication

b) Doubtful diagnosis

c) Drug resistant TB

d) Progressive cauda equina syndrome/conus medullaris syndrome

Question 2:

In case of a meniscal tear repair, which of the following zones heals best?

a) Red zone

b) White zone

c) Gray zone

d) Red+White zone

Question 3:

A 22-year-old male patient presents with a history of RTA. His leg is in abduction and external rotation. What could be the cause?

a) Anterior dislocation of the hip

b) Posterior dislocation of the hip

c) Intertrochanteric fracture

d) Fracture neck of femur

Question 4:

The most specific test for carpal tunnel syndrome is?

a) Durkan's test

b) Phalen test

c) Tinel test

d) Two point discrimination

Question 5:

A female patient with a history of fall few months back presents with the deformity shown in the image. Probing into history reveals it was managed with traditional treatment. This deformity is likely to be due to _____

a) Intra-articular fracture of wrist

b) Extra-articular fracture of wrist

c) Boxer's fracture

d) Barton's fracture

Question 6:

Scoliosis screening in school children is done by?

a) Straight leg raising test

b) Forward bending test

c) Backward bending test

d) Single leg standing test

Question 7:

Fishtail deformity is seen in the fracture of _____

a) Distal end of radius

b) Distal end of humerus

c) Distal end of tibia

d) Distal end of femur

Question 8:

Ring sequestrum is seen in?

a) Salmonella infection

b) TB osteomyelitis

c) Pin tract infection

d) Chronic osteomyelitis

Question 9:

A counterforce brace is used in?

a) Tennis elbow

b) Ankle sprain

c) Ulnar claw hand

d) Foot drop

Question 10:

A 10-year-old boy presents with pain around the left knee joint. On X-ray, there is a lesion at the distal femur extending up to the terminal end. The gross image of the condition is given below. What can be the most probable diagnosis?



a) Giant cell tumour

b) Chondroblastoma

c) Osteoid osteoma

d) Osteosarcoma

Question 11:

Which of the following substances is used for bone formation in a patient with non-union?

a) PMMA

b) BMP

c) Ca PO_4

d) $CaSO_4$

Question 12:

The brace shown in the given image is _____.

a) Boston brace

b) Milwaukee brace

c) Taylor brace

d) ASHE brace

Question 13:

A 40-year-old man is suffering from increasing difficulty in bending forward with increasing stiffness of the spine and reduced chest movements. A CT was performed and shown below, the likely diagnosis is?

a) Ankylosing spondylitis

b) Rheumatoid arthritis

c) Reiter's syndrome

d) Fluorosis

Question 14:

In spine fusion surgery, the screws are fixed to which part of the vertebra?

a) Spinous process

b) Lamina

c) Pedicle

d) Facet joint

Question 15:

Which of the following tests are not commonly used in osteoporosis:

a) 1, 2 & 3

b) 2 & 4

c) 2 & 3

d) 1, 3, 4 & 5

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	a
5	b
6	b
7	b
8	c
9	a
10	d

11	b
12	b
13	a
14	c
15	c

Detailed Explanations

Solution to Question 1:

A cold abscess without neurological complications is not an indication for surgery in Pott's spine/TB spine.

Indications of surgery in TB spine are:

• Progressive neurologic deficit such as progressive cauda equina syndrome/conus medullaris syndrome (Option D)

• Progressive increase in spinal deformity

• Resistance to anti-tubercular therapy (Option C)

• Recurrence of disease

• Cord compression

• Severe pain due to abscess or spinal instability

• Progressive impairment of pulmonary function

• Uncertain diagnosis (Option B): Inability to obtain a microbiological diagnosis through microscopy, culture, or PCR techniques may occur. In cases where there is no accessible source for sampling (such as discharging sinus or superficial abscess) or insufficient specimen obtained from image-guided percutaneous biopsy, an open surgical biopsy is the only option to obtain a specific culture and sensitivity report.

Antero-lateral decompression is the surgery of choice for Pott's paraplegia as it helps in the removal of both solid and liquid debris.

Solution to Question 2:

The red zone heals best in case of a meniscal tear due to its rich blood supply.

The menisci are divided into three zones depending on the perfusion of blood. The menisci are supplied by perimeniscal blood vessels from superior and inferior branches of medial and lateral genicular vessels.

Following birth, the menisci become increasingly avascular. By adulthood, only the peripheral 10% to 25% of the tissue is perfused. This gives rise to 3 distinct regions of the menisci:

• The peripheral relatively vascularised region called the red-red zone (Zone 1)

• A zone of transition between the 2 zones, which is called the red-white zone (Zone 2)

• The completely avascular inner zone known as the white-white zone (Zone 3)

There is a direct relationship between the vascularization and capacity of the tissue to heal, predisposing the white-white zone to permanent post-traumatic and degenerative lesions.



Solution to Question 3:

The patient presents with abduction and external rotation of the lower limb following a road traffic accident, which is suggestive of anterior dislocation of the hip.

Anterior dislocation of the hip is usually sustained following a high-energy accident. On physical examination, the affected limb will show abduction, external rotation, and slight flexion. Lengthening of the limb is observed, along with a prominent anterior bulge in either the pubic or groin region.

In posterior dislocation of the hip, the lower limb is flexed, adducted, and internally rotated, and this condition is associated with limb shortening. The neck of femur fracture and intertrochanteric fracture present with external rotation and shortening of the lower limb.

The given image shows a patient with an externally rotated, abducted, and slightly flexed left leg due to anterior dislocation of the hip:

According to the position assumed by the femoral head, anterior dislocation of the hip can further be classified as:

• Type I - Superior or pubic

• Type II - Inferior or obturator

A lateral view X-ray will be confirmatory. Radiological findings include:

• The displaced femoral head will appear larger than the normal side

• As the thigh is externally rotated lesser trochanter is more prominent

X-RAY showing anterior dislocation of the hip:



Treatment of anterior dislocation includes a closed reduction under sedation or general anesthesia. If closed reduction fails, an open procedure may be planned.

Solution to Question 4:

Among the given options, Durkan's test is the most specific test for the diagnosis of carpal tunnel syndrome (CTS).

CTS is a common entrapment mononeuropathy caused by median nerve compression as it passes through the fibro-osseous carpal tunnel beneath the flexor retinaculum. Patients present with pain and paresthesia over the radial 3 and a half digits. There may be a weakness of thumb opposition. In chronic cases, the thenar eminence is wasted. The diagnosis is usually made on clinical grounds but nerve conduction studies can confirm the diagnosis.

Durkan's test is the most specific test for the diagnosis of CTS. In Durkan's test, direct median nerve compression for about 30 seconds, elicits paresthesias and pain.

Other tests for CTS:

• Phalen's test - Elicited by holding the wrist flexed fully. The appearance of sensory symptoms within 60 seconds implies a positive test.



• Tinel's test - Sensory symptoms reproduced by percussing over the median nerve imply a positive test.

Two-point discrimination is done to assess sensory loss and is not specific to carpal tunnel syndrome.

Solution to Question 5:

The given image shows the dinner-fork deformity seen in Colle's fracture which is an extra-articular fracture of the wrist.

Colle's fracture is a fracture of corticocancellous junction of the distal end of the radius. The dinner fork deformity is the prominence on the back of the wrist and depression in the front. It is due to the dorsal angulation of the distal fracture fragment. It is treated by a Colles' cast, which immobilizes the wrist in palmar flexion, pronation, and ulnar deviation.

Displacements that may be seen in Colles' fracture:

• Dorsal tilt

• Dorsal shift

• Lateral tilt (Radial tilt)

• Lateral shift (Radial shift)

• Supination

• Impaction of fragments

The given radiograph shows the dorsal tilt of the distal fragment leading to dinner fork deformity.



Solution to Question 6:

The forward bending test is used as a screening tool in scoliosis.

Adam's forward bending test is the best clinical test for evaluating spinal curvature in scoliosis.

Steps of Adam's test :

• The patient is asked to bend forward at the waist until the spine is at the horizontal level

• Knees must be maintained straight, with the feet together, the arms dependent, and the palms in opposition

• The examiner observes the trunk from behind (to assess midthoracic and lumbar rotation) and from the front (to assess upper thoracic rotation)

It is used to differentiate between structural and non-structural curves in scoliosis. On bending forward, non-structural scoliosis disappears, while structural scoliosis persists.



A straight leg raising test is used to find nerve compression in the lumbosacral area.

Solution to Question 7:

Fishtail deformity is seen following a fracture of the distal end of the humerus.

Fishtail deformity of the elbow is a rare complication that can occur in childhood after a distal humeral fracture. It is characterized by a contour abnormality of the distal humerus due to the absence or resorption of the lateral trochlear ossification centers.

Fishtail deformity of the distal humerus can be of 2 types:

• The first type is a sharp-angled wedge, which commonly occurs after fractures of the lateral condyle. It is caused by the persistence of a gap between the lateral condylar physis ossification center and the medial ossification of the trochlea. Because of this gap, the lateral crista of the trochlea may be underdeveloped, which may represent a small “bony bar” in the distal humeral physis

• The second type is believed to be associated with osteonecrosis of the lateral part of the medial crista of the trochlea.

Solution to Question 8:

Ring sequestrum is seen in a pin tract infection.

Sequestrum is a fragment of dead bone lying within a cavity formed by necrosis. Infection at a pin site in an external fixation construct leads to a ring sequestrum.

Types of sequestra:

Type	Seen in
Tubular	Pyogenic infections like osteo myelitis
Ring	Around pin tracts in external fixators
Black	Actinomycosis
Corallifor m	Perthes disease
Coke	Tuberculosis (cancellous bon e)
Feathery	Syphilis, Tuberculosis
Sandy	Tuberculosis

Solution to Question 9:

A counterforce brace is used in tennis elbow.

Tennis elbow, also known as lateral epicondylitis, is a condition involving the lateral epicondyle caused by repetitive contraction of the extensor muscles during activities that require forceful wrist extension and supination, such as playing tennis or using a screwdriver.

A counterforce brace is a treatment option for tennis elbow (shown in the image below). It works by applying compression to the extensor muscles, specifically the extensor carpi radialis brevis (ECRB) region, which helps release the forces exerted on the lateral epicondyle. This compression prevents the transmission of tension up to the lateral epicondyle, providing relief and promoting healing in the affected area.

Management options in tennis elbow:

• Conservative management: Rest and physiotherapy

• Medical intervention: Injection of local anesthetic and steroids

• Surgical methods: Percutaneous release of epicondylar muscles

• Other management methods: Extracorporeal shock wave therapy, arthroscopic release, autologous blood injections, counterforce bracing, rehabilitative exercises, and ultrasound-guided percutaneous needle therapy.

Solution to Question 10:

The given clinical scenario, involving a child and a gross specimen revealing a metaphyseal tumor in the distal femur that has infiltrated through the cortex, lifted the periosteum, and extended through the epiphyseal plate, supports the diagnosis of osteosarcoma.

A giant cell tumor (option A) is primarily an epiphyseal tumor that usually affects adults between ages 20 and 40 years when skeletal maturity is complete. Chondroblastoma (option B) is a benign tumor of childhood most commonly seen in the epiphysis. Osteoid osteoma (option C) is a benign tumor most commonly seen in the diaphysis of the proximal femur.

Osteosarcoma is the most common non-hematologic primary malignancy of bone. It occurs most commonly in the second decade of life. It is typically metaphyseal. The distal femur, proximal tibia, and proximal humerus are common sites.

Morphologically, osteosarcomas are bulky, gritty, gray-white tumors, often with hemorrhage and cystic degeneration. They destroy cortices, form soft tissue masses, and spread extensively in the medullary canal, replacing hematopoietic marrow. Although infrequent, they may also penetrate the epiphyseal plate or enter the joint.

Radiological features:

• Irregular destruction in the metaphysis overshadowed by the formation of new bone

• Lifting of the periosteum (Codman triangle) due to the formation of new bone inciting a periosteal reaction

• Bone formation along the blood vessels within the tumor centrifugally, giving rise to a sunburst (sunray) appearance



Solution to Question 11:

Among the given options, bone morphogenic protein (BMP) is the substance used for bone formation in a patient with non-union.

BMP is a bone-stimulating protein used as a bone graft in the treatment of non-union and open tibial fractures.

Bone grafts can be:

• Osteoinductive: They stimulate osteogenesis through the differentiation of mesenchymal cells into osteoprogenitor cells. BMPs are osteoinductive.

• Osteoconductive: They provide linkage across defects and a scaffold upon which new bone can form. Calcium phosphate (Option C), calcium hydroxyapatite, and calcium sulfate (Option D) are osteoconductive.

Polymethylmethacrylate (PMMA), also known as bone cement, holds bone and metal implants together by forming an interlocking network between the irregularities.

Solution to Question 12:

The brace shown in the image is the Milwaukee brace.

The Milwaukee brace is an active corrective spinal orthosis primarily used in the ambulatory treatment of structural scoliosis. Designed as a cervico-thoracic-lumbar-sacral orthosis (CTLSO), the main objective of the Milwaukee brace is to delay or eliminate the need for surgery. However, in recent years, the use of the Milwaukee brace has declined due to the availability and convenience of modern braces. Nonetheless, it is still employed for scoliosis cases characterized by high thoracic and cervical spine curves.

Option A: Boston brace is a type of thoraco-lumbo-sacral-orthosis (TLSO brace). It has a hard polypropylene outer shell with a soft foam lining inside. It is used in patients whose scoliosis curve apex is at T7 or lower. The below image shows a Boston brace.



Option C: Taylor brace is designed to control the upper and lower regions of the spinal column and support your spine whilst standing, sitting, and walking after spinal surgery.



Option D: ASHE brace (Anterior spinal hyperextension brace) is used in dorso lumbar spinal injury.

Solution to Question 13:

The given clinical scenario of a 40-year-old man with increasing stiffness in the spine, along with CT spine showing ossification of the anterior longitudinal ligament and ankylosis involving the thoracolumbar vertebral bodies points to a diagnosis of ankylosing spondylitis (AS).

AS is a chronic inflammatory disease affecting the spine and sacroiliac joint. It is more common in young males and has a strong association with HLA-B27.

Features of lower back pain in AS:

• Age at onset < 40 years

• Insidious onset

• Morning stiffness > 30 minutes

• Improvement with exercise

• Pain persists with rest and is also present at night.

• Alternating gluteal pain

• Persistence for more than 3 months

Examination findings in AS:

• Loss of lumbar lordosis

• Increased thoracic kyphosis

• Diminished spinal movements

• Decreased chest expansion (<7 cm)

• Tenderness over sacroiliac joints

Important X-ray features of ankylosing spondylitis:

• Sacro-iliac joints:

• Cardinal sign/earliest sign is the erosion and fuzziness of the sacroiliac joints

• Later- periarticular sclerosis and bony ankylosis of the joint.

• Vertebral changes:

• Early spondylitis is characterized by small erosions at the corners of vertebral bodies with reactive sclerosis: Romanus lesions of the spine (Shiny corner sign)

• Squaring of vertebral bodies or flattening of the normal anterior concavity of the spine due to inflammatory resorption of the bone.

• Later signs are ossification of the ligaments around the intervertebral discs producing syndesmophytes (bridges) between adjacent vertebrae

• Bamboo spine appearance is due to bridging at several vertebral levels

• Linear ossification along the central spine, representing interspinous ligament ossification can give a Dagger sign appearance on frontal radiographs.

CT imaging in AS helps to visualize bone erosions, osteoporosis/osteosclerosis, and new bone formation in the form of syndesmophytes, ligamentous ossification, and ankylosis.

The X-ray below shows the bamboo spine appearance in AS:





Solution to Question 14:

In spine fusion surgeries, screws are placed in the pedicle of the vertebrae.

Spinal fusion or spondylodesis is a surgical technique that joins two or more vertebrae. This procedure prevents any movement between the fused vertebrae.

Pedicle screws are placed above and below the vertebrae that are to be fused. These are used in a spinal fusion to add extra support and strength to the fusion while it heals.

Rods are used to connect the screws as shown below. These prevent movement and allow the bone to heal.



The post-operative X-ray given below shows fusion of L5 and S1 vertebrae.



Solution to Question 15:

Among the given options, bone scan and quantitative CT are not commonly used in the diagnosis of osteoporosis.

Osteoporosis is a common metabolic bone disease characterised by a diffuse reduction in bone density due to a decrease in bone mass.

DEXA, chemical tests and X-ray are commonly used in the diagnosis of osteoporosis. While quantitative CT also helps to measure bone mass density at the spine and hip, it is rarely used clinically because the radiation exposure and cost are both much higher than with DXA. A bone scan uses a radioactive tracer injected into the bloodstream to detect altered metabolic activity in the bone with the aid of a gamma camera. It is not used in the diagnosis of osteoporosis.

DEXA - dual-energy X-ray absorptiometry, is the investigation of choice for the diagnosis of osteoporosis. T-score less than or equal to 2.5 in the lumbar spine, femoral neck, or total hip indicates osteoporosis.

Chemical tests like serum calcium, phosphorus and alkaline phosphatase levels are usually normal in primary osteoporosis and help to rule out other conditions before establishing a diagnosis of osteoporosis.

X-ray features of osteoporosis:

• Loss of vertical height of a vertebra occurs due to collapse

• Cod fish appearance characterized by the disc bulging into the adjacent vertebral bodies, resulting in a biconvex shape

• Ground glass appearance of the bones

Orthopedics NEET 2018

Question 1:

De Quervain's disease classically affects :

a) Extensor pollicis longus and abductor pollicis longus

b) Extensor pollicis longus and abductor pollicis brevis

c) Extensor pollicis brevis and abductor pollicis brevis

d) Extensor pollicis brevis and abductor pollicis longus

Question 2:

You are an intern in the casualty. For which of the following patients will you need to urgently call the orthopedic resident on call?

a) Patient with recurrent shoulder dislocation

b) Patient with a fractured arm with capillary refill time of less than 3 seconds in his fingers

c) Patient with a fractured arm with a 10 cm long incision over the arm

d) Patient with a fractured arm with capillary refill time of 5 seconds in his fingers

Question 3:

Which of the following is not associated with Sprengel deformity?

a) Diastematomyelia

b) Klippel-Feil syndrome

c) Dextrocardia

d) Congenital scoliosis

Question 4:

What is the most common bone to be fractured in children?

a) Distal forearm bones

b) Humerus

c) Clavicle

d) Carpals

Question 5:

A patient has a history of a road traffic accident 2 years back, and has developed pain and swelling now at the same site. His X-ray image is shown in the image given below. What is the probable diagnosis?



a) Osteogenic sarcoma

b) Ewing's sarcoma

c) Chronic osteomyelitis

d) Multiple myeloma

Question 6:

A 35-year-old man comes with a history of musculoskeletal pain. Which of the following suggests that the pain is non-articular?

a) Swelling

b) Pain in both movement and at rest

c) Painless on passive movement

d) Presence of crepitation

Question 7:

The pathognomonic sign of a simple bone cyst is _____

a) Fallen fragment sign

b) Scalloping of cortex

c) Never breaches physis

d) Central radiolucent lesion

Question 8:

Which of the following is incorrect about perilunate dislocation?

a) Lunate is dislocated anteriorly but the rest of the carpals remains in position

b) Lunate remains in position but rest other dislocate dorsally

c) Involves median nerve injury

d) Tavernier's manoeuvre is used for closed reduction

Question 9:

Identify the carpal bone that is most commonly fractured after fall on an outstretched hand.



a) A

b) C

c) B

d) D

Question 10:

For hypertrophic non-union following a fracture, the most appropriate treatment would be _____

a) Stabilization

b) Bone grafting

c) Stabilization and bone grafting

d) None of the above

Answer Key

Question No.	Correct Option
1	d
2	d
3	c
4	a
5	c
6	c
7	a
8	a
9	a
10	a

Detailed Explanations

Solution to Question 1:

De Quervain's disease classically affects extensor pollicis brevis and abductor pollicis longus.

De Quervain's disease is a stenosing tenosynovitis of the extensor tendons of the first dorsal compartment, namely extensor pollicis brevis (EPB) and abductor pollicis longus (APL). It involves noninflammatory thickening of both the tendons and the tunnel or sheath through which they pass. This condition is prominent among females, especially women of childbearing age.

Key diagnostic factors:

• History of joint overuse at home or at work.

• Pain and tenderness at the tip of the radial styloid.

• Positive Finkelstein test: The examiner grasps the thumb of the patient and abducts the hand towards the ulnar side. This causes severe pain over the tip of the styloid process



Treatment in most patients includes conservative management with splinting and steroid injections. Only a few patients require surgical release of the 1st dorsal compartment (housing the APL and EPB tendons).

Solution to Question 2:

A patient with a fractured arm and a prolonged capillary refill time of 5 seconds in his fingers requires urgent evaluation and intervention by an orthopedic resident.

The capillary refill time is a simple clinical test used to evaluate peripheral circulation and assess blood flow to tissues. It can be assessed by applying pressure to the skin of the heel or toe pulp for approximately 5 seconds, causing blanching, and then releasing the pressure to allow the color to return. A capillary refill time of less than 3 seconds is considered normal, indicating good blood perfusion and adequate circulation.

In this patient with a fractured arm, a capillary refill time of 5 seconds suggests compromised blood flow to the fingers, which could be indicative of a potential vascular injury or compartment syndrome. Both of these conditions are surgical emergencies and require immediate attention to prevent further damage and ensure proper blood supply to the affected limb.

While the other patients should also be evaluated by an orthopedic resident, they do not present an immediate life-threatening situation.

Solution to Question 3:

Dextrocardia is not associated with Sprengel deformity.

Sprengel deformity is the congenital elevation of the scapula. It is characterized by an elevated shoulder on the affected side, with an abnormally high, smaller, and prominent scapula. The neck appears shorter, and there may be kyphosis or scoliosis of the upper thoracic spine. While shoulder movements are painless, abduction and elevation may be limited due to scapular fixation.

Abnormalities associated with Sprengel deformity include:

• Klippel-Feil Syndrome

• Kyphosis or Scoliosis

• Cervical rib

• Omovertebral bar (a bony bridge between the scapula and the cervical spine)

• Spina bifida

• Diastematomyelia (congenitally split spinal cord)

• Torticollis

• Renal abnormalities

The image below shows Sprengel deformity:

Solution to Question 4:

The most common bones to be fractured in children are the distal forearm bones.

The distal radius is the most commonly fractured bone in childhood. It is more commonly fractured than the clavicle.

The clavicle is most commonly fractured during birth.

Solution to Question 5:

The given clinical scenario of a patient with a history of an accident 2 years ago, experiencing symptoms of pain and swelling at the site of injury, and an X-ray of the lower limb revealing sequestrum and involucrum in the tibia, strongly suggests a diagnosis of chronic osteomyelitis.



Chronic osteomyelitis may occur by direct inoculation of the infective organism during trauma (like road traffic accidents). The infection continues to thrive within the bone leading to alternating quiescent and flare-up stages of chronic osteomyelitis.

Sequestrum is a piece of dead bone. It is formed when a fragment of bone gets separated from the healthy bone due to necrosis. It varies in size from a spicule to a large fragment and acts as a nidus for infection. It leads to abscess and persistent pus discharge in the bone. Sinus tracts are formed when the abscess from the infected bone drains out through the skin.

The involucrum is the dense new sclerotic bone surrounding the sequestrum. It is formed from deep layers of stripped periosteum. Cloacae are the perforations in the involucrum. Pus and tiny spicules of bone may be discharged through them.

The radiograph below shows sequestrum, involucrum and cloacae:

Option A: Osteogenic sarcoma characteristically shows the sun-burst appearance and Codman's triangle, as shown below.

Option B: Ewing's sarcoma shows an onion-skin appearance of the periosteum, as shown below.

Option D: Multiple myeloma shows multiple punched-out lesions of the bone, as shown below.

Solution to Question 6:

Non-articular disorders tend to be painless on passive movement.

Feature	Articular	Non-articular
Structures involved	Synovium, synovial fluid, articular cartilage, intraarticular ligaments, joint capsule, and juxta-articular bone	Extraarticular ligaments, tendons, bursae, muscles, fascia, bone, nerves, and overlying skin
Pain	Deep or diffuse	Localized

Feature	Articular	Non-articular
Pain with movement	Both on active and passive movement	Only on active movement
Swelling	Present	Absent
Crepitation	Present	Absent
Instability	Present	Absent
Deformity	Present	Absent
Examples	Osteoarthritis, infectious arthritis, ankylosing spondylitis	Fibromyalgia, myofascial pain, acute muscle strain, myotonic dystrophy

Solution to Question 7:

The fallen fragment sign is the pathognomonic radiographic sign of a simple bone cyst.

Simple bone cysts typically occur during the first two decades of life, predominantly between the ages of 4 and 10, with a 2:1 male preponderance. They are benign unilocular cysts filled with straw-colored serosanguinous fluid. The most common site for these cysts is the proximal humerus. Although they are usually asymptomatic, they can lead to pathological fractures through the cyst wall, resulting in pain and swelling. Additionally, due to their proximity to the growth plate, the cysts may cause growth disturbances.

On an X-ray, a bone fragment may be observed resting in a dependent portion of the cyst. This finding, known as the fallen fragment or fallen leaf sign, is considered pathognomonic of a simple bone cyst. Another variant of this sign is the trapdoor sign, in which a periosteal hinge prevents the fragment from falling down and enables it to change position with the patient's movement.

The radiograph below shows the fallen fragment sign:



Solution to Question 8:

Among the given options, statement A is incorrect.

A fall on a dorsiflexed hand may tear the tough ligaments that normally bind the carpal bones, leading to dislocations. In peri-lunate dislocation, the lunate remains in position in the lunate fossa of the distal radius but the rest of the carpals dislocate dorsally. It is frequently associated with compression of the median nerve.

Prompt reduction reduces swelling and potential damage to the median nerve. The Tavernier maneuver is the most commonly used method of closed reduction. It involves locking the capitate into the distal concavity of the lunate. If the injury cannot be reduced using this method, urgent reduction in the theater should be performed.

In lunate dislocation, the lunate dislocates anteriorly while the rest of the carpal bones remain in place.



Solution to Question 9:

The scaphoid (labelled as A) is the most commonly fractured carpal bone after a fall on an outstretched hand.

It is more susceptible to fracture because it lies obliquely across the two rows of carpal bones. It also lies along the line of loading (i.e, in the line of weight transmission) between the forearm and thumb. A combination of forced carpal movement and compression, such as in a fall on the dorsiflexed (outstretched) hand, exerts severe stress on the scaphoid making it liable to fracture.

Due to its blood supply originating from the dorsal distal pole, the scaphoid's proximal pole possesses a poor blood flow, making it less prone to heal and resulting in non-union.



Solution to Question 10:

In cases of hypertrophic non-union following a fracture, stabilization alone has the potential to induce union.

Non-union following a fracture occurs when the healing process fails to progress, resulting in a failure to unite. There are two primary types of non-union:

• Hypertrophic - where callus formation is present but does not bridge the fracture site

• Atrophic - characterized by minimal or no attempt at callus formation

For hypertrophic non-union without deformity, conservative treatment using functional bracing or rigid fixation alone (internally or externally) can promote union. However, in cases of atrophic non-union, fixation alone is insufficient, requiring the excision of fibrous tissues in the fracture gap and sclerotic bone ends, followed by the application of bone grafts around the fracture.

Orthopedics NEET 2019

Question 1:

A 40-year-old patient presents to the emergency department following a road traffic accident. The radiograph of his pelvis is shown below. What is the most likely diagnosis?

a) Central dislocation of the hip

b) Fracture shaft of femur

c) Posterior dislocation of the hip

d) Anterior dislocation of the hip

Question 2:

A 12-year-old boy presents with swelling and pain for 6 months over the upper tibia. His radiographic image is shown below. The most likely diagnosis is:



a) Osteosarcoma

b) Osteoclastoma

c) Brodie's abscess

d) Ewing's sarcoma

Question 3:

Which of the following fractures is most prone to non-union?

a) Proximal scaphoid

b) Inter-trochanteric

c) Distal radius

d) Olecranon fracture

Question 4:

Painful arc syndrome is seen in all except:

a) Complete tear of supraspinatus

b) Fracture greater tuberosity

c) Subacromial bursitis

d) Supraspinatus tendinitis

Question 5:

De-gloving injury refers to _____

a) Skin and subcutaneous fat are stripped from the underlying fascia

b) Skin, subcutaneous fat and fascia are stripped from tendons

c) Skin, subcutaneous fat, fascia and tendons are stripped from bone

d) Only skin is stripped off

Question 6:

Rheumatoid arthritis management in a patient with deformity is:

a) Steroids

b) Methotrexate with steroids

c) Methotrexate with anti-TNF

d) Steroids only after NSAIDs fail

Question 7:

A 60 year old woman presents with low backache and on a lumbar CT the following is seen. What is your diagnosis?



a) Spondylolisthesis

b) Discitis

c) Osteoporosis

d) Pott's spine

Question 8:

An 18-year-old boy presents to the orthopedician with complaints of painful swelling below the knee joint. An X-Ray was obtained which is given below. Which of the following is the preferred treatment modality for this patient?

a) Extended curettage and bone grafting

b) Simple curettage and bone grafting

c) Radiofrequency ablation

d) Radical excision

Question 9:

The following images A and B show:



a) Mallet finger & Jersey finger

b) Boxer fracture & march fracture

c) Mallet finger & boxer fracture

d) Jersey finger & baseball finger

Answer Key

Question No.	Correct Option
1	c
2	c
3	a
4	a
5	a
6	c
7	c
8	a
9	d

Detailed Explanations

Solution to Question 1:

The radiograph of the patient's pelvis following a road traffic accident reveals upward and lateral dislocation of the left femoral head, accompanied by an obscured lesser trochanter, indicating a posterior dislocation of the left hip.

Posterior hip dislocation is commonly seen following dashboard injuries and the patient presents with flexion, adduction, and internal rotation deformity of the lower limb.

Radiological findings:

• Loss of continuity of acetabulum with the femoral head (break in Shenton's line)

• Upward and lateral displacement of the femoral head

• The femoral head appears smaller than normal

• The lesser trochanter is not visualized as the thigh is internally rotated

• Femoral head and posterior acetabular wall fractures may also be seen

The image below shows posterior dislocation with disruption of Shenton's line:



Treatment:

An early reduction should be done as the risk of osteonecrosis and subsequent osteoarthritis increases with time.

Closed reduction under sedation or general anaesthesia is done in most cases. If there is an associated femoral neck fracture, a closed reduction should not be attempted as it can lead to the displacement of the femoral neck and disrupt the blood supply to the femoral head.

In closed reduction the following steps are done:

The pelvis is kept in a steady position and traction is applied in line of the femur with the knee flexed at 90 degrees. The hip is then flexed maintaining the traction along with internal rotation and adduction. This is called Allis' maneuver. The below image describes this technique:



Other methods of closed reduction include - Stimson's maneuver, East Baltimore lift maneuver, and Bigelow's maneuver.

After reduction, the stability of the hip is tested by flexing the hip to 90 degrees, and a longitudinal and posteriorly directed force is applied. While performing this the joint is monitored in an image intensifier to look for any signs of subluxation. In case of evidence of subluxation or any significant fractures then surgical repair should be done. If there is a delay in surgical correction, skeletal traction with a distal femoral traction pin can be done.

If there is damage to the acetabulum, or intervening structures between the femoral head and the acetabular surface are present - open reduction can be considered.

Early complications:

• Sciatic nerve injury - It can be damaged in 10-20% of the cases

• Vascular injury - Superior gluteal artery tear can occur

• Concurrent femoral shaft fracture - When both occur together hip dislocation may be missed

Late complications:

• Osteonecrosis of the femoral head

• Myositis ossificans

• Secondary osteoarthritis

• Recurrent instability

Solution to Question 2:

The given clinical scenario of a child with prolonged swelling and pain over the upper tibia, along with a radiograph showing a lytic lesion with a dense sclerotic rim and minimal periosteal reaction, suggests the diagnosis of Brodie's abscess.

Brodie's abscess is a localized form of subacute osteomyelitis that occurs when organisms of low virulence infect an immunocompetent host. The abscess is walled off by the thickening of the surrounding bone trabeculae. It most commonly involves the metaphysis of the proximal tibia and distal femur.

The patient is typically a child or adolescent experiencing prolonged pain near a large joint for several weeks or months. Symptoms may include a limp, slight swelling, muscle wasting, and localized tenderness.

The typical radiographic finding is a circumscribed, round or oval radiolucent cavity measuring 1-2 cm in diameter, surrounded by a halo of sclerosis, with minimal or no periosteal reaction. The radioisotope scan shows markedly increased activity. MRI may be helpful in diagnosis. Open biopsy with curettage is indicated if there is any doubt in the diagnosis.

Differential diagnoses include:

• Cystic tuberculosis

• Eosinophilic granuloma

• Osteoid osteoma

• Ewing's sarcoma

Treatment:

• Immobilization and antibiotics (flucloxacillin and fusidic acid) intravenous for 4-5 days, followed by oral administration for another 6 weeks; healing may take up to 12 months

• Curettage is needed if the x-rays show no healing after conservative management.

The following X-ray and MRI images show Brodie's abscess of the radius in a patient.



The differentiating features of each of the given conditions are as follows:

Condition	Age group	Radiographic findings
Brodie's abscess	Children, adolescents	Lytic lesion with dense sclero tic rim, minimal periosteal re action
Osteosarcoma (Option A)	Bimodal: Children and adoles cents, 70-80 yearsRapidly pr ogressivetumor	Periosteal reaction leading to sunray appearance and Cod man triangle
Osteoclastoma (Option B)	20-40 years (unlikely in child ren)	Lytic lesion withthin sclerotic rim, minimal periosteal reac tion
Ewing's sarcoma (Option D)	10-20 years	Diaphysealmass and perioste al reaction with onion-peel a ppearance

Solution to Question 3:

Among the given fractures, proximal scaphoid fractures are most prone to non-union.

Since the nutrient artery enters the scaphoid near its distal end, proximal fractures are associated with avascular necrosis and non-union. The more proximal the fracture in the scaphoid, the higher the chances of avascular necrosis and non-union.

Certain bones have a relatively limited or watershed blood supply that can potentially get interrupted by fracture. These include:

• Head of the femur: after fracture of the neck or posterior dislocation of the hip.

• Proximal scaphoid: due to fracture through the waist of the scaphoid.

• Lunate: following a dislocation.

• Body of talus: after fracture of its neck.

Fractures in these sites have a higher risk for non-union or development of osteonecrosis.

Other options:

Option B: Inter-trochanteric fractures involve either the greater or lesser trochanters or both, more common in the elderly following a history of a fall. Malunion is the most frequent complication. The vast metaphyseal region has an abundant blood supply, contributing to a higher union rate and fewer chances of avascular necrosis.

Option C: Distal radius fractures occur by fall on an outstretched hand and they include Colles' fracture (most common), Smith's fracture, Barton's fracture, and Chauffer's fracture. The complications include joint stiffness, malunion (more common), carpal tunnel syndrome, subluxation of the inferior radio-ulnar joint, Sudeck's dystrophy, and rupture of the extensor pollicis longus tendon. Avascular necrosis or non-union is rare.

Option D: Olecranon fracture results from a direct injury as in a fall onto the point of the elbow. Complications include non-union, elbow stiffness, and osteoarthritis.



Solution to Question 4:

The partial tear of the supraspinatus tendon is responsible for painful arc syndrome rather than a complete tear.

Painful arc syndrome, also known as impingement syndrome is a condition where there is pain in the shoulder and upper arm during the mid-range of shoulder abduction (between 60-120 degrees), arising due to repetitive rubbing of the supraspinatus tendon under the coracoacromial arch.

Causes include:

• Minor tears of the supraspinatus tendon

• Supraspinatus tendinitis

• Calcification of supraspinatus tendon

• Fracture of greater tuberosity

• Subacromial bursitis

Important tests for impingement are:

• Painful arc test

• Hawkin-Kennedy's test

• Neer's impingement sign

• Jobe's test

Investigations:

• An x-ray of the shoulder helps to detect bony avulsions, spurs, calcific deposits, sclerotic areas etc.

• Ultrasound of the shoulder joint is highly reliable with a sensitivity of 98%.

• Single contrast arthrogram is considered the gold standard in diagnosing rotator cuff tears

Management:

• Conservative: heat massage, exercises, NSAIDs, subacromial steroid injections, and temporary immobilization (90% of patients recover with these measures).

• Indications for surgery include failure of conservative measures for 3 months, increasing loss of shoulder function, and patients who are young and active.

• Surgical techniques include excision of adhesions and manipulation of the shoulder, excision of calcium deposits, repair of incomplete tear, acromioplasty, acromionectomy for more disabling pain with normal range of movements, direct suture for complete rupture of the rotator cuff, rotation and transposition of the flap, and free graft.



Hawkins–Kennedy test



The affected arm is elevated 90 degrees in the scapular plane with the elbow in 90 degrees flexion. The upper arm is then stabilized and rotated internally.

©MARROW

Neer's impingement sign

Keeping the scapula stable, the affected arm is raised fully in passive flexion, abduction, and internal rotation.

©MARROW

Jobe's test

Arms elevated in the scapular plane, elbows extended with the thumb pointing downwards, while the examiner applies downward pressure.

©MARROW

Solution to Question 5:

Degloving is the avulsion of skin and subcutaneous fat from the underlying fascia, muscle or bone.

Degloving injuries can be open or closed. An example of open degloving injury is finger avulsion injury with loss of skin. Closed degloving injuries result from shearing forces (e.g. motor vehicle collisions).

The extent of degloving injuries are often underappreciated and much of the skin may be non-viable. Disruption of perforating vascular and lymphatic vessels may result in a characteristic haemolymphatic collection between the fascial planes called a Morel-Lavallée lesion.

Patterns of degloving injury:

• Limited degloving with abrasion or avulsion

• Non-circumferential degloving

• Circumferential single plane degloving

• Circumferential multiplanar degloving

Assessing the viability of degloved tissue can be difficult and may therefore require more than one surgical exploration and debridement before definitive reconstruction. Non-viable skin may show fixed staining and thrombosis of subcutaneous veins. Most surgeons serially excise the degloved skin until punctate dermal bleeding is seen from viable tissue.

Intravenous fluorescein can help in delineating non-viable tissue, but it requires specialist equipment and there is a small risk of anaphylaxis. Indocyanine green has been found to have less risk of allergy.

Open degloving injuries are managed by reattaching the viable skin and grafting. Closed degloving injuries are managed by draining the accumulated fluid, removing the dead tissue and sclerotherapy. For severe, multiplanar degloving, amputation is done.

The given image below shows a degloving injury of the lower limb:

Solution to Question 6:

Methotrexate with anti-TNF is superior to any other treatment in limiting progression of joint destruction and disability.

In rheumatoid arthritis (RA) joint inflammation is the main cause of joint damage and is the most important factor of joint disability and deformities.

Methotrexate belongs to the category of drugs known as disease-modifying antirheumatic drugs (DMARDs). They slow down or prevent the structural progression of RA and hence are used in most combination therapies. Among the DMARDs, methotrexate is the preferred first-line drug.

Biologic DMARDs such as anti-TNF agents target cytokines and cell-surface molecules to slow the progression of joint inflammation. These drugs are typically used in combination with background methotrexate therapy.

Combination therapy with methotrexate and an anti-TNF drug is the most commonly used next step for the treatment of patients with deformities, having an inadequate response to methotrexate therapy.

Steroids are used when rapid control of the disease is required. They have a rapid onset of action, while the DMARDs take anywhere from 6-12 weeks to come into effect. They can be used as a cover during this period. Steroids can also be used in short courses for acute flare-up episodes of RA. Long-term use of steroids is not advised due to their adverse effects such as osteoporosis. They are usually not used alone (option A) but are used as an adjunct to DMARDs like methotrexate (option B).

Nonsteroidal anti-inflammatory drugs (NSAIDs) are used as adjunctive therapy for the management of uncontrollable symptoms because of their analgesic and anti-inflammatory properties. Chronic use is not advised due to their side effects namely gastritis and peptic ulcer disease as well as impairment of renal function (option D)

Hence, among the given options methotrexate with anti-TNF is the best therapy in a patient with deformity.

Solution to Question 7:

The given image depicts the ballooning of the intervertebral disc into the adjacent vertebrae leading to biconcave shape of vertebral body and biconvex shape of the disc which is known as the ‘codfish appearance’. This is classically seen in osteoporosis.

Osteoporosis is a metabolic disease of the bone that is caused due to an equivalent decrease in both bone matrix and its mineralization, resulting in a diffuse decrease in both, bone density and bone mass. It is seen in various conditions namely in postmenopausal females, Cushing's disease, and long-term steroid therapy.

The various complications seen in osteoporosis are due to decreased bone mass which contributes to bone pains and increased frequency of pathological fractures such as Colle's fracture and neck of femur fractures. In the vertebrae, the dorso-lumbar spine is the most frequent site to be involved and can present as increased kyphosis due to compression of the anterior part of the vertebral bodies.

Other radiological features of osteoporosis:

• Loss of vertical height of vertebra due to collapse

• Ground glass appearance of bones especially in the bones like pelvis

• Loss of cortical bone (picture frame vertebrae) and trabecular bone (ghost vertebrae)

• Loss of trabeculae in proximal femur area.

DEXA scan is the gold standard for quantifying bone mass in osteoporotic patients. A T-score of less than -2.5 is diagnostic of osteoporosis.

The X-ray given below is of a post-menopausal female with Colle's fracture after trivial trauma, caused secondary to osteoporosis.



Option A: Spondylolisthesis is the anterior translation of one vertebra over another. It commonly occurs between L5-S1 followed by L4-L5. On examination, often there is a visible or palpable 'step'.

A lateral X-ray of the spine shows anterior displacement of one vertebra over another. The oblique view in severe spondylolisthesis shows the 'beheaded Scottish terrier sign' (due to a break in pars interarticularis).

The below-given images depict spondylolisthesis as seen on X-ray:





Option D: Pott's spine also known as TB spine occurs secondary to a pulmonary or gastrointestinal primary and spreads through the Batson's plexus of veins (paravertebral plexus). It most commonly involves the dorso-lumbar region.

On X-ray and CT imaging:

• A reduction in disc space is seen between the vertebrae and it is one of the earlieast signs of TB spine

• Vertebral body destruction can also be seen- based on the number of vertebrae involved they are called by different names. An involvement of 3-4 vertebrae is called as kyphosis

• Cold abscesses such as paravertebral abscess can be visible as a soft tissue shadow

• Advanced destruction and wedging of vertebrae

Solution to Question 8:

The given x-ray showing an expansile, eccentric, multiloculated lytic lesion, at the end of a long bone with a soap-bubble appearance suggests a giant cell tumor (GCT) or osteoclastoma. The treatment of choice for this condition is extended curettage and bone grafting.

Giant cell tumor is a benign but locally aggressive tumor seen involving the metaphysis and /or the epiphysis of long bones, as seen in the image, in young adults of age 20-45 years and rarely teenagers less than 20 years.

Although an aneurysmal bone cyst (ABC) is also a benign, locally destructive, eccentric multiloculated lytic tumor that can present with a similar radiographical picture in any age group (most commonly in adolescents), it is confined within the metaphysis of the long bones, unlike GCT, thus ruling it out in this case. On MRI a fluid sign can be seen.

The following is an x-ray image of the knee showing ABC confined to the metaphysis of the long bones:

An ABC is treated by extended curettage and grafting with a bone graft substitute (option B).

GCT is most commonly seen in the distal femur, proximal tibia, proximal humerus, and distal radius. It presents with complaints of swelling and vague pain and Sometimes, the patient can presents for the first time with a pathological fracture because of the lesion.

Important x-ray features of a GCT:

• Eccentric, solitary lytic lesion

• In epiphysis and metaphysis

• Expansile- expands overlying cortex

• Soap bubble appearance - due to the trabeculae of the remnants of bone traversing the lytic lesion, giving it a loculated appearance.

• No or minimal reactive sclerosis around the tumor

• No calcification

Histologically, GCTs consist of undifferentiated spindle cells profusely interspersed with multi-nucleated giant cells. The tumor stroma is well-vascularized with bands of cellular or collagenous fibrous tissue.

The treatment of choice for GCT is by

• Extended curettage and bone grafting (Option A).

• GCT in the relatively insignificant bone like the distal ulna, clavicle, and proximal fibula is treated by resection without reconstruction.

• Bisphosphonates have been shown to reduce the proliferation of osteoclasts and hence have been shown to stabilize local and metastatic disease.

• Radiotherapy can be used for vertebral tumors

• Additional supportive therapy such as cryotherapy, when used reduce the reccurence rate.

Solution to Question 9:

The following image shows the Jersey finger and baseball finger respectively.

• Jersey finger - is an avulsion injury of the flexor digitorum profundus at the insertion of the base of the distal phalanx with the ring finger most commonly involved. It is caused due to sudden hyperextension of the distal joint.

• Baseball finger (also known as mallet finger)- is a disruption to the extensor tendon at the DIP joint due to collision with a hard object (like a ball) forcing it to flex past its normal range of motion. Clinically, the distal phalanx is in slight flexion in this condition.

Other options:

Boxer's fracture is the break in the metacarpal neck, most commonly of the fifth finger following a blow with the fist.

The x-ray given below shows Boxer's fracture:

March fracture is a stress fracture of the neck of the 2nd or 3rd metatarsal

The X-ray given below shows March fracture:

Orthopedics NEET 2020

Question 1:

Which method of measurement of scoliosis has been depicted in the image given below?



a) Cobb's angle

b) Baumann's angle

c) Ferguson angle

d) Bohler's angle

Question 2:

A 30-year-old woman developed a tumor at the knee joint. Biopsy-image is given below. What is the most likely condition?



a) Osteosarcoma

b) Ewing's sarcoma

c) Giant cell tumour

d) Osteoblastoma

Question 3:

A patient presented with pain in the hand. The joints involved were proximal interphalangeal joint, distal interphalangeal joint and first carpometacarpal joint. The wrist and metacarpophalangeal joints were spared. What is the likely diagnosis?

a) Osteoarthritis

b) Rheumatoid arthritis

c) Psoriatic arthritis

d) Pseudogout

Question 4:

A 55 year old female complains of lower backache. The radiographic image of her lumbosacral spine is given below. What is the probable diagnosis?



a) Osteoporosis

b) Ankylosing spondylitis

c) Diffuse idiopathic skeletal hyperostosis

d) Renal osteodystrophy

Question 5:

A maid is playing with a child by spinning him while holding his hands. A few hours later, the child starts crying, does not use his arm, and does not let anybody touch him. What is the possible diagnosis?

a) Pulled elbow

b) Olecranon fracture

c) Fracture head of radius

d) Elbow dislocation

Question 6:

A patient was admitted to ICU 48 hours after the fracture of the femur. The saturation of oxygen in the rebreathing unit was 100% but his Spo2 remained 60%. The patient was in a state of confusion. Chest radiograph showed lung fields to be clear. What is the most likely diagnosis?

a) Pulmonary embolism

b) Fat embolism

c) ARDS

d) Occult pneumothorax

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	a
5	a
6	b

Detailed Explanations

Solution to Question 1:

The given image depicts the measurement of the curvature of scoliosis using the Cobb's angle.

Scoliosis is characterized by the lateral curvature of the spine in the coronal plane when the patient is upright. Confirmation of the diagnosis is achieved through full-length lateral and posteroanterior X-rays of the spine. A Cobb's angle greater than 10% on the radiograph is diagnostic of scoliosis.

The measurement of Cobb's angle consists of three steps:

• Locating the superior-end vertebra of the curve.

• Locating the inferior-end vertebra of the curve.

• Drawing perpendicular lines from the superior surface of the superior-end vertebra and from the inferior surface of the inferior-end vertebra and making them intersect.

The angle formed by these perpendiculars represents the degree of spinal curvature, referred to as Cobb's angle.



Nash and Moe's method and Perdriolle and Vidal method are tools for assessing the vertebral rotation in scoliosis.

Other options:

Option B: Baumann's angle is the angle formed between a line drawn through the lateral epiphysis and the long axis of the humerus. Normally, it measures less than 80 degrees but increases in cases of supracondylar fractures.



Option C: Ferguson's angle is measured between the upper endplate of the S1 vertebra and a horizontal line. A normal Ferguson angle is typically around 34 degrees. A decreased Ferguson angle can be observed in advanced spondylolisthesis.

Option D: Bohler's angle is the angle formed by two lines tangent to the highest points of the anterior and posterior facets of the calcaneus. This angle is decreased in calcaneal fractures.



Solution to Question 2:

The biopsy image shows multinucleated osteoclast-like giant cells with scattered mononuclear cells, indicating a diagnosis of giant cell tumor (GCT)/osteoclastoma.

GCT usually affects young adults of age 20 to 40 years, with slight female predominance. It is a benign but locally aggressive tumor. It is exclusively seen in adults and affects the ends of long bones, typically involving the epiphysis. The most common location of the tumor is the distal femur followed by the proximal tibia.

Histologically, GCTs show uniform oval mononuclear tumor cells alongside numerous osteoclast-type giant cells containing 100 or more nuclei. Necrosis and mitotic activity may be prominent.



Other options:

Option A: Histologically, osteosarcoma shows pleomorphic tumor cells with hyperchromatic nuclei and abundant mitoses. A definitive diagnosis requires the presence of malignant tumor cells producing unmineralized osteoid or mineralized bone, which is typically fine and lacelike in appearance.



Option B: Histologically, Ewing's sarcoma shows sheets of uniform, small, round cells with scant cytoplasm, which may appear clear because they are rich in glycogen. The presence of Homer–Wright rosettes indicates a greater degree of neuroectodermal differentiation.



Option D: Histologically, osteoblastoma shows randomly interconnecting trabeculae of woven bone, which are prominently rimmed by a single layer of osteoblasts. The stroma surrounding the neoplastic bone is composed of loose connective tissue that contains numerous dilated and congested capillaries.



Solution to Question 3:

The most likely diagnosis is osteoarthritis (OA) of the hand.

In OA of the hand, the joints involved are the distal and proximal interphalangeal joints (DIP & PIP), and the first carpometacarpal joint (base of the thumb), while the metacarpophalangeal (MCP) joint, wrist, elbow, and ankle are usually spared.

The nodules palpable in the DIP are called Heberden's nodes, while those palpable in the PIP are called Bouchard's nodes.



The image below shows OA of the hand:

Other options:

Option B: There is relative sparing of the DIP joint in rheumatoid arthritis.

Option C: Psoriatic arthritis of the hands commonly involves the PIP, DIP, and metacarpophalangeal joints.

Option D: In pseudogout, the knee joint is most commonly affected, followed by the wrist, while the first metatarsophalangeal joint (podagra) is rarely affected.

Solution to Question 4:

The radiographic image of an elderly female's lumbosacral spine shows ballooning of disc spaces, resulting in the characteristic biconcave appearance of vertebral bodies known as the 'codfish' appearance, typically seen in osteoporosis.

Osteoporosis is a common metabolic bone disease characterised by a diffuse reduction in bone density due to a decrease in bone mass.

Risk factors for osteoporosis include:

• Increasing age

• Female sex

• Alcohol

• Smoking

• Previous history of fragility fractures

• Family history of hip fracture

• Low body mass index- less than 18.5 kg/m2

• Excessive usage of glucocorticoids

Osteoporosis presents with fractures that occur without any trauma or as a result of low-impact trauma. The commonly associated fractures are fractures of the distal part of the radius (Colle's fracture), fractures of the hip, and fractures of the vertebrae. In severe cases, the fracture of vertebrae can cause kyphosis.

Radiological findings:

• A decreased vertical height of the vertebra due to collapse

• Ground glass appearance - especially in bones like pelvis

• Codfish appearance - it occurs due to bulging of the disc into the adjacent vertebral bodies so that it becomes biconcave.

DEXA (dual-energy X-ray absorptiometry) is the investigation of choice for the diagnosis of osteoporosis. T-score less than -2.5 in the lumbar spine, femoral neck, or total hip indicates osteoporosis.

Solution to Question 5:

The given scenario of spinning the child by holding his hands, followed by features suggesting pain and inability to use the arm, points to a diagnosis of pulled elbow.

Pulled elbow or nursemaid's elbow refers to the dislocation of the radial head over the annular ligament, commonly seen in children under the age of 6 years. The mechanism of injury is through axial traction on the extended and pronated forearm. This happens when a child is pulled, lifted, or swung by the arm.

Following the dislocation, the child holds the arm relatively immobile, with the elbow extended and the forearm pronated, and resists supination.

The diagnosis is primarily clinical, though X-rays are usually obtained to exclude a fracture.

To treat the condition, simple analgesia is administered, and while the child's attention is diverted, the elbow is extended and pronated to relocate the radial head with a snap. This technique is preferred over the alternative method of supination and flexion, which tends to be more painful and less successful.



Solution to Question 6:

In this clinical scenario, the presence of reduced SpO2 and altered mental status in a patient presenting within 72 hours following a long bone fracture suggests the diagnosis of a fat embolism.

Fat embolism is caused by the fat globules originating from the bone marrow following the fracture of long bones. Clinical features of fat embolism are usually seen within 72 hours. They include tachycardia, fever, breathlessness, petechiae, confusion,and restlessness. Gurd's criteria is used for the diagnosis of fat embolism syndrome.

There is no specific diagnostic test for fat embolism. However, urine analysis to detect fat globules and monitoring of blood pO2 levels are conducted.

The management of fat embolism primarily involves supportive care, including oxygen therapy and prompt stabilization of long-bone fractures.

Option A: Pulmonary embolism most commonly occurs as a result of thrombi from deep vein thrombosis of the leg.

Option C: Acute respiratory distress syndrome is diagnosed using the Berlin's criteria. All four of the following conditions must be met:

• It develops within 1 week of clinical insult or worsening respiratory symptoms.

• Chest radiograph shows bilateral opacities consistent with pulmonary edema. These opacities are not fully explained by effusions, lobar or lung collapse, or nodules.

• The edema is not due to cardiac failure, or fluid overload.

• There is impaired oxygenation. The ratio of arterial oxygen partial pressure to the fraction of inspired oxygen (PaO2/FiO2) is ≤ 300 mmHg as assessed on 5 cm H2O of positive end-expiratory pressure (PEEP).

Option D: Occult pneumothorax is a condition that is not suspected on the basis of the symptoms or plain radiography, but is detected with CT imaging. It usually follows a thoracic injury.

Orthopedics INI-CET 2021

Question 1:

Which of the following is seen in osteoporosis?

a) Normal calcium and increased ALP

b) Normal calcium and normal ALP

c) Decreased calcium and normal ALP

d) Increased calcium and increased ALP

Question 2:

A 15-year-old boy was brought to the ER following a motor vehicle collision complaining of pain over the hip. The x-ray is given below. What is the next step of management?

a) Closed reduction and assessment of hip stability

b) CT pelvis with 3D reconstruction

c) High weight skeletal traction

d) Open reduction and reconstruction of posterior pillar of acetabulum

Question 3:

Which of the following findings appear late in compartment syndrome?

a) Pallor

b) Paralysis

c) Pulselessness

d) Pain on passive strech

Question 4:

Which of the following statements are true about open fractures?

a) 1 and 2

b) 1 and 4

c) 2 and 4

d) 2 and 3

Question 5:

An old lady slipped and fell at home. She was diagnosed with a Colles' fracture and managed with a POP cast. What is the correct sequence of reduction?

a) Traction, POP, palmar flexion, ulnar deviation

b) Traction, palmar flexion, ulnar deviation, POP

c) Traction, ulnar deviation, palmar flexion, POP

d) Palmar flexion, ulnar deviation, traction, POP

Question 6:

Match the following fractures with their sites

a) A-1,B-3,C-2,D-4

b) A-2,B-3,C-1,D-4

c) A-4,B-2,C-4,D-3

d) A-2,B-4,C-1,D-3

Question 7:

Arrange the following in the correct sequence for the Thomas test:

a) 1,2,3,4,5

b) 3,2,1,4,6

c) 3,2,6,5,4

d) 3,6,2,5,4

Question 8:

Following a femoral shaft fracture, your consultant asks you to provide tibial traction. Which of the following will you request from the nurse?

a) 1, 2, 4

b) 3, 5, 6

c) 3, 4, 5

d) 1, 2, 3, 4, 5, 6

Question 9:

The deformity shown in the image below is noted when a patient tries to extend his fingers. Injury to which nerve can lead to this condition?



a) Ulnar nerve

b) Radial nerve

c) Median nerve

d) Posterior interosseous nerve

Question 10:

A 60-year-old woman with a history of Colles' fracture 8 months ago comes to the orthopedics outpatient department. Her T score is -2.5 and she attained menopause at the age of 52 years. She is currently asymptomatic. What treatment will you give this patient?

a) Hormone replacement therapy

b) Alendronate

c) Calcium and vitamin D

d) Repeat DEXA scan after 3 months

Question 11:

Following a road traffic accident, a patient develops type IIIa compound tibial fracture. Arrange the following external fixators in decreasing order of their stability (highest to lowest).

a) 1>2>3>4

b) 2>4>3>1

c) 1>4>3>2

d) 2>3>1>4

Question 12:

A 45-year-old patient with a tibial fracture in a cast presented to the emergency department with complaints of intense pain. The cast was removed and on examination, passive extension was painful. The dorsalis pedis and posterior tibial pulses were present and there was a loss of sensation over the first web space. What is the next step?

a) Duplex imaging

b) Re-apply cast

c) Measure anterior compartment pressure

d) Analgesics

Question 13:

Match the following with the best mode of management:

a) 1-d, 2-c, 3-b, 4-a

b) 1-c, 2-a, 3-d, 4-b

c) 1-a, 2-d, 3-b, 4-c

d) 1-d, 2-a, 3-b, 4-c

Question 14:

In nerve injuries, if the outer sheath and nerve fiber are intact while the inner axons are damaged, what is it known as?

a) Neurapraxia

b) Axonapraxia

c) Axonotmesis

d) Neurotmesis

Answer Key

Question No.	Correct Option
1	b
2	a
3	c
4	b
5	b
6	a
7	c
8	b
9	a
10	b
11	c
12	c
13	c
14	c

Detailed Explanations

Solution to Question 1:

Osteoporosis is associated with normal calcium, normal ALP levels and phosphorous.

Osteoporosis is a common metabolic bone disease characterised by a diffuse reduction in bone density due to a decrease in bone mass.

Risk factors for osteoporosis include:

• Increasing age

• Female sex

• Alcohol

• Smoking

• Previous history of fragility fractures

• Family history of hip fracture

• Low body mass index- less than 18.5 kg/m2

• Excessive usage of glucocorticoids

• They present with fractures that occur without any trauma or as a result of low-impact trauma. The commonly associated fractures are fractures of the distal part of the radius (Colle's fracture), fractures of the hip, and fractures of the vertebrae. In severe cases, the fracture of vertebrae can cause kyphosis.

Radiological findings:

• A decreased vertical height of the vertebra due to collapse

• Ground glass appearance - especially in bones like pelvis

• Codfish appearance - it occurs due to bulging of the disc into the adjacent vertebral bodies so that it becomes biconcave

DEXA (dual-energy X-ray absorptiometry) is the investigation of choice for the diagnosis of osteoporosis. T-score less than -2.5 in the lumbar spine, femoral neck, or total hip indicates osteoporosis.

Treatment - the mainstay is to reduce the risk of fractures. Following medications can be useful:

• Bisphosphonates. They are the first line of medication. They reduce osteoclastic bone resorption and prevent bone loss. Eg. zoledronate

• Denosumab- This is an antibody to the RANK ligand. It prevents the activation of RANK which is required for the proliferation of osteoclasts. Therefore, it decreases bone resorption

• Parathyroid hormone - Preotact and teriparatide are recombinant parathyroid hormone analogues. At low and intermittent doses, they can promote bone formation.

• Selective estrogen receptor modulators (SERMs)- They act on the estrogen receptors and prevent bone resorption. Eg. raloxifene.

• Strontium ranelate- It is thought to increase bone formation and reduce bone resorption.

• Cathepsin K inhibitor- inhibits the activity of collagenases (eg, odanacatib)

• Sclerostin inhibitor- Sclerostin inhibits the activity of osteoblasts. Romosozumab is a humanized antisclerostin monoclonal antibody that increases bone formation.

Operative treatment is required for femoral neck and other long bone fractures.

Solution to Question 2:

The radiograph shows upward and lateral dislocation of the left femoral head with an obscured lesser trochanter which is suggestive of posterior dislocation of the left hip and the next step in the management is closed reduction and assessment of hip stability.

Posterior hip dislocation:

It is commonly seen following dashboard injuries and in this condition, the patient presents with flexion, adduction, and internal rotation deformity of the lower limb.

Radiological findings:

• Upward dislocation of the femoral head

• The femoral head appears smaller than normal

• The lesser trochanter is not visualized as the thigh is internally rotated

• The femoral head and posterior acetabular wall fractures may also be seen

Treatment:

An early reduction should be done as the risk of osteonecrosis and subsequent osteoarthritis increases with time.

Closed reduction under sedation or general anaesthesia is done in most cases. If there is an associated femoral neck fracture, a closed reduction should not be attempted as it can lead to the displacement of the femoral neck and disrupt the blood supply to the femoral head.

In closed reduction the following steps are done:

The pelvis is kept in a steady position and traction is applied in line of the femur with the knee flexed at 90 degrees. The hip is then flexed maintaining the traction along with internal rotation and adduction.

After reduction, the stability of the hip is tested by flexing the hip to 90 degrees, and a longitudinal and posteriorly directed force is applied. While performing this the joint is monitored is an image intensifier to look for any signs of subluxation. If there is evidence of subluxation or any significant fractures then surgical repair should be done. If there is a delay in surgical correction, skeletal traction with a distal femoral traction pin can be done.

In case of damage to the acetabulum, or the presence of intervening structures between the femoral head and the acetabular surface open reduction can be considered.

Early complications:

• Sciatic nerve injury - It can be damaged in 10-20% of the cases

• Vascular injury - Superior gluteal artery tear can occur

• Concurrent femoral shaft fracture - When both occur together hip dislocation may be missed

Late complications:

• Osteonecrosis of the femoral head

• Myositis ossificans

• Secondary osteoarthritis

• Recurrent instability

Solution to Question 3:

Pulselessness is a late sign is compartment syndrome.

Acute compartment syndrome is a surgical emergency caused by increased pressure within a fascial compartment, leading to tissue ischemia. It is characterized by rapid progression of pain and swelling in an extremity, commonly precipitated by traumatic injury.

The classic features of ischemia in acute compartment syndrome include the 5 P's:

• Pain: Pain is the first symptom of acute compartment syndrome. Ischemic muscle groups have an increased sensitivity to touch. On passive stretching of the muscles, severe pain can be elicited.

• Paresthesia: Paresthesia and hypoesthesia are the first signs of ischemia and may be noted due to compression of nerves traversing the affected compartment.

• Pallor: Pallor of the skin may not always be seen as ischemia occurs at the capillary level (late sign).

• Paralysis: Paralysis of muscle groups affected in that compartment (late sign).

• Pulselessness: The absence of distal pulses (extremely late sign).

Treatment:

• Prompt decompression of the affected compartment is done.

• All the dressings, bandages, and cast on the affected limb are removed and the limb is nursed flat.

• Fasciotomy is done if the differential pressure is less than 30 mm Hg

• Fasciotomy wounds are left open and after 2 days if there is necrosis debridement is done. If the wounds are healthy it can be sutured without tension or skin grafting can be done.

The diagnosis of compartment syndrome can be confirmed by measuring the intra-compartmental pressure using a split catheter introduced into the compartment. The difference between diastolic pressure and compartment pressure is known as differential pressure (ΔP). If the differential pressure (ΔP) is less than 30 mm Hg (4.00 kilopascals), immediate compartment decompression is indicated.

Solution to Question 4:

Open fractures commonly involve the tibia and phalanges (statement 1) and necessitate early wound debridement (statement 4).

Open fractures are characterized by a break in the overlying skin and soft tissues, exposing the fracture fragment to the external environment. It is important to note that open fractures can be accompanied by multiple other injuries, and the presence of an open fracture does not exclude the possibility of developing compartment syndrome as a potential complication.

The most common sites involved in open fractures include:

• Phalanges (fingers and toes)

• Tibial diaphysis

• Distal radius

• Ankle

Major components of the management of open fractures are:

1) Antibiotic prophylaxis - It should be given as early as possible. Co-amoxiclav or cefuroxime are the preferred drugs.

2) Early debridement of the wound and fracture - This includes the following steps:

• Excision of the wound margins.

• Extension of the wound if required.

• Delivery of the fracture outside the wound for proper examination.

• All the devitalized tissue should be removed.

• Cleansing of the wound to remove all the foreign materials.

• Nerves and tendons are usually left alone, but if the wound is clean and expertise is available their repair can be considered.

3) Early wound cover:

• In Gustilo type 1 and 2 fractures wound cover can be done at the time of debridement.

• Delayed wound cover is usually practiced in type 3 fractures.

4) Fracture stabilization:

• The method of fracture stabilization depends on the degree of contamination, amount of soft tissue damage, and time from injury to operation.

• External fixation is used as a temporary measure if there is a delay in wound cover.

• If definitive wound cover can be achieved at the time of debridement and there is no significant contamination, then an open fracture can be managed as same as a closed injury.

Solution to Question 5:

The correct order of reduction of Colles' fracture is traction, palmar flexion, ulnar deviation, POP.

Colle's fracture refers to the extra-articular fracture of the distal radius with dorsal angulation of the distal bone fragment.

Displacements that may be seen in Colles' fracture are:

• Impaction of fragments

• Dorsal tilt

• Dorsal displacement

• Lateral tilt (radial tilt)

• Lateral displacement (radial displacement)

• Supination

The patient usually presents with pain, swelling, and deformity of the wrist. There may be a typical 'dinner-fork deformity' where the radial styloid process comes to lie at the same level or a little higher than the ulnar styloid process. A dorsal tilt is characteristic and is seen on a lateral x-ray.

Management of Colles' fracture:

• Undisplaced fractures: immobilization in a below-elbow plaster cast for 6 weeks

• Displaced fractures: closed manipulative reduction and immobilization in Colles' cast. The technique of reduction of Colles' fracture is as follows:

• Dis-impaction of the fracture fragments by longitudinal traction against counter traction provided by the assistant.

• The distal fragment is then pressed into palmar flexion followed by ulnar deviation.

• The patient's hand is drawn into pronation, palmar flexion, and ulnar deviation.

• A Colles' cast is then applied, extending from below the elbow to the metacarpal heads, with the wrist in palmar flexion and ulnar deviation.

• Surgical correction is advisable for comminuted fractures, especially in young adults where the dominant hand is involved. It can be with percutaneous K-wire fixation, external fixators, or locking compression plates.

Complications:

• Stiffness of joints - finger stiffness is commonly seen

• Malunion - can produce dinner-fork deformity

• Subluxation of the inferior radio-ulnar joint

• Carpal tunnel syndrome - uncommon

• Sudek's osteodystrophy - Colles' fracture is the commonest cause of this condition in the upper limb.

• Rupture of extensor pollicis longus tendon - extremely rare



Solution to Question 6:

The correct match is A-1, B-3, C-2, D-4

• Jones fracture - 5th metatarsal

• Bennett's fracture - 1st metacarpal

• March fracture - 2nd metatarsal

• Boxer fracture - 5th metacarpal

Jones fracture - It refers to the fracture of the proximal part of the fifth metatarsal at the metaphyseal-diaphyseal junction. It is usually caused by an adduction injury of the forefoot. Conservative management is done with a short-leg non-weight bearing cast for a period of 6 weeks. Surgical correction by intramedullary screw fixation is advocated for faster healing in highly active individuals.

Bennett's fracture - It occurs at the base of the first metacarpal bone. It is sustained usually by a fall or during punching. The fracture line is oblique and extends into the carpometacarpal(CMC) joints. It is highly unstable due to the pull of abductor pollicis longus on the shaft of the metacarpal.

X-ray shows a triangular fragment in contact with the medial edge of the trapezium and proximal subluxation of the rest of the thumb.

Surgical correction with percutaneous K-wire or open reduction and internal fixation (ORIF) with a lag screw is the preferred treatment.

March fracture - These are stress fractures that usually involve the second metatarsal. It is typically seen in military recruits or runners. On examination, a tender lump may be noted in the midshaft region of the metatarsal. Similar lesions may be also seen in osteoporotic individuals.

The x-ray may be normal initially but later show a hairline crack. After a period of 4-6 weeks, a mass of callus can be identified.

Reduction and splintage are not necessary as it is undisplaced. Normal walking can be done and it's managed by elastic bandage of the forefoot.



Boxer's fracture - It refers to the fracture at the neck of the fifth metacarpal. It is usually produced by striking an object with the hand as seen with a boxer's punch.

It is managed using a gutter splint with flexed MCP joints and straight IP joints until the discomfort settles.

Solution to Question 7:

Among the given options, the correct sequence for the Thomas test is given by option C i.e. 3,2,6,5,4.

Thomas test is used to detect flexion deformity of the hip. Flexion deformity is a common hip deformity, often caused by stronger flexor muscles overpowering the extensor muscles. When there is a spasm of the muscles, the stronger flexors pull the hip in flexion. Soft tissue contractures can convert this into a fixed flexion deformity, making locomotion impossible.

Steps of Thomas test:

• Positioning: The patient is instructed to lie supine on a hard surface with their legs fully extended.

• Observing lumbar lordosis: As the patient assumes the supine position, the curvature of the lower back is visually assessed for any exaggerated lumbar lordosis. It is noted that excessive lumbar lordosis may be a compensatory response to a flexion deformity.

• Gradual flexion of the normal hip: The normal hip is gradually flexed, with care taken not to exceed its normal range of motion.

• Monitoring for overcorrection: During the flexion of the normal hip, signs of overcorrection on the normal side are observed, such as the pelvis starting to lift off the ground. To prevent this, both ischial tuberosities are ensured to remain in contact with the surface.

• Automatic positioning of the affected hip: As a result of the flexion of the normal hip, the affected hip automatically assumes the deformed position characterized by flexion.

• Measuring the degree of flexion: Once the affected hip is in the flexed position, the angle formed between the affected thigh and the bed is measured. This angle represents the actual degree of flexion deformity.

Solution to Question 8:

Among the given options, Steinmann pin, Bohler's stirrup, and Bohler Braun splint are used for tibial traction in femoral shaft fractures.

Steinmann pin: It is a stout, straight steel rod used to provide skeletal traction. Its diameter ranges from 3 to 6 mm. The common sites where it is used include the upper end of the tibia, the

supracondylar region of the femur, and the calcaneum.

Bohler's stirrup: It is used to hold a Steinmann pin and apply traction. The screws on the sides of the stirrup hold the pin. The direction of traction can be changed without moving the pin inside the bone, which prevents the pin from loosening.

Bohler Braun splint: This refers to Bohler's modification of the Braun splint. It consists of a heavy metal frame with four pulleys. It is used for the treatment of fracture shaft femur and supracondylar fractures of the femur. It can also be used for the fractured shaft of the tibia and fibula.

Other options:

Denham pin: This pin is threaded in the center and engages the bony cortex, which reduces the risk of pin sliding. It is used when the bone is cancellous -like calcaneum and osteoporotic bones.

Kirschner wire or K-wire is 1 to 3 mm in diameter. The wire is pointed at both ends for both anterograde and retrograde insertion. The shape of the tips could be trocar, diamond or threaded. It is used in small bone fractures.

Thomas splint: It is used for immobilizing fractures of the femur.

Given below are the images of these instruments






Bohler's stirrup


Denham pin

Thomas Splint
Inner diameter:
25 cm
Length: 102 cm
©MARROW



Solution to Question 9:

The given image shows extension at the metacarpophalangeal joint and flexion at the interphalangeal joints of the ring and little fingers while the patient tries to extend his fingers, suggesting an ulnar nerve injury.

The ulnar nerve supplies the medial two lumbricals in the hand. The lumbricals have a combined action on metacarpophalangeal (MCP) joint flexion and interphalangeal (IP) joint extension (glass holding posture). In ulnar nerve injury, since the medial lumbricals are not functional, the forearm muscles are unopposed. Hence, there is hyperextension at MCP joints and flexion at IP joints. This causes claw hand deformity. There is also numbness of the ulnar one and a half fingers.

Elbow fractures or dislocations lead to high lesions of the ulnar nerve. Here, the hand is not markedly deformed because the ulnar half of the flexor digitorum profundus (which is responsible for flexing the fingers) is paralyzed and the fingers, therefore, appear less clawed. This is called the high ulnar paradox.

The following tests are done to test the integrity of the ulnar nerve:

• Egawa's test - In a positive test, due to the weakness of the dorsal interossei, side-to-side movements of the middle finger is weak.

• Book test or Froment's sign - The patient is asked to hold a book forcefully while the doctor tries to pull it. In a positive test, due to weakness of adductor pollicis muscle, adduction of thumb is lost. The patient tries to hold the book by using flexor pollicis longus muscle. This produces flexion at the inter-phalangeal joint while holding the book.

• Card test - In the case of weak palmar interossei, it is easy for the examiner to pull out the card when held between the fingers. This is due to the loss of adduction by palmar interossei muscles.



Positive Card Test
In the weakness of palmar interossei, there is a weakness of finger adduction. So the patient is unable to hold card firmly between fingers.

Other options

Option B - In radial nerve palsy, the patient presents with 'wrist drop' in case of paralyzed wrist extensors, and 'finger drop' when extensor digitorum is paralyzed. When there is compression in the axilla, radial nerve palsy is called Saturday night palsy. There is paralysis of the triceps, and weakness of the wrist and hand. In high lesions, there is sensory loss on the dorsum of the hand over the anatomical snuff box.

Option C - In median nerve injury, paralysis of opponens pollicis muscle leads to 'ape thumb' deformity. In forearm or elbow injuries, there is paralysis of long flexors of the thumb, index, and middle fingers, radial wrist flexors, and forearm pronators. This leads to the characteristic 'pointing index finger' when the patient tries to make a fist. There is sensory loss over three and a half digits on the radial side.

Option D -Posterior interosseous nerve injury, is seen in posterior interosseous syndrome. The wrist is in radial deviation and there is loss of metacarpophalangeal joint extension. However, wrist extension is preserved.

Solution to Question 10:

The clinical vignette of an elderly post-menopausal woman with a history of Colles' fracture and a T score of -2.5 suggests postmenopausal osteoporosis and the treatment of choice in this condition would be a bisphosphonate such as alendronate.

Options A and C: Hormone replacement therapy, calcium and vitamin D supplements, and calcitonin are not recommended treatments to reduce fracture risk in osteoporosis.

Option D: Repeating the DEXA scan after 3 months will not be required as the patient's T score (-2.5) already suggests osteoporosis. DEXA is done in the following patients to screen for osteoporosis:

• Females aged 65 years or more

• Males aged 70 years or more

• Postmenopausal females under the age of 65 years with risk factors

• Fracture after the age of 50 years

• Males aged 50-69 with risk factors

Osteoporosis is a common metabolic bone disease characterized by a diffuse reduction in bone density due to a decrease in bone mass. It is typically asymptomatic until fractures occur. These fractures are commonly low trauma in nature, defined as a fall from standing height or less. The initial fracture that often occurs is in the distal radius, known as Colles' fracture, followed by fractures in the vertebrae and hip unless treatment is initiated.

A DEXA scan is the investigation of choice in diagnosing osteoporosis and the results are reported in terms of T-score. According to WHO, a T score of

• -1 or higher is considered normal

• -1.01 to -2.49 is defined as osteopenia.

• -2.5 or lower is defined as osteoporosis

Bisphosphonates like alendronate are the drugs of choice for fracture prevention in postmenopausal women. They specifically inhibit bone resorption and thus are very effective in treating osteoporosis. They concentrate at sites of active remodeling by remaining in the matrix until the bone is remodeled. Then, they are released in the acidic environment of the resorption lacunae and induce apoptosis in osteoclasts.

Other treatment options for postmenopausal osteoporosis are:

• Denosumab - This is an antibody to RANKL that promotes osteoclastogenesis. Inhibiting RANKL thus decreases osteoclastic activity.

• Parathyroid hormone - Preotact and teriparatide are anabolic hormones. They promote bone formation when given at intermittent low doses. Aboloparatide is a new drug in this group that has been tried for this condition.

• Raloxifene - It is a selective estrogen receptor modulator (SERM) approved and shown to be effective in preventing vertebral fractures.

• Strontium ranelate - It is thought to promote osteogenesis but is strongly contraindicated in patients prone to thromboembolic events.

• Romosozumab - Sclerostin inhibits osteoblastic activity. Romosozumab shows anti-sclerostin activity, thereby increasing bone formation.

Solution to Question 11:

Among the given options, Ilizarov> Biplanar frame/Ring with cylindrical rod> Uniplanar with double rod> Uniplanar with a single rod (1>4>3>2) represents the correct order of stability of the given external fixators.

External fixation provides rigid fixation of the bones. This is commonly used in severe open types II and III fractures, as the application of a cast, or traction methods and internal fixation is not recommended

The principle of external fixation is the transfixion of bone above and below the fracture with Schanz screws or tensioned wires and these are then connected to each other by rigid bars. There are numerous types of external fixation devices:

• A unilateral frame with a support member and half-pins on one plane constitute a unilateral one-plane configuration. The addition of a second support member and the second plane of half-pins form a unilateral two-plane configuration.

• Transfixation pins connected to support members at either end are used to build the bilateral one-plane configuration. The addition of a second plane of half-pins forms a bilateral two-plane configuration.

• Ring fixators consist of complete or partial rings connected by rods. The rings are fixed to the bone via half pins or wires under high tension.

Circular external fixators confer greater stability and often negate the need to span joints. Unilateral external fixation is commonly used as a temporary method of fixation. Its disadvantages include the need to span joints for sufficient stability. Uniplanar fixation devices are quick and easy to apply but are less sturdy than multiplanar fixation. Bilateral frames are created when the pins are on both sides of the bone and can also add additional stability.

Given below is an Ilizarov ring fixator.

Solution to Question 12:

The clinical presentation of this patient with a history of tibial fracture complaining of intense pain, with painful passive extension and loss of sensation of the first web space upon the removal of the cast, is suggestive of acute compartment syndrome. The next step would be to measure the anterior compartment pressure to confirm the diagnosis.

Acute compartment syndrome refers to the rise in intra-compartmental pressure leading to tissue ischemia and necrosis. The injuries at high risk of developing acute compartment syndrome are fractures of the proximal one-third of the tibia, soft tissue injury, distal radius fracture, and

diaphyseal fracture of the forearm. Other high-risk injuries are crush injuries and circumferential burns.

The classic features of ischemia in acute compartment syndrome include the 5 Ps:

• Pain: is the first symptom of acute compartment syndrome. On passive stretching of the muscles, severe pain can be elicited.

• Paresthesia: paresthesia and hypoesthesia are the first signs of nerve ischemia.

• Paralysis: paralysis of muscle groups affected in that compartment is a late sign.

• Pulselessness: usually, the peripheral pulses and capillary return are intact in the early stages, unless there is a major arterial injury. If pulses are absent, arteriography is indicated.

• Pallor: may not always be seen as ischemia occurs at the capillary level.

The diagnosis can be confirmed by measuring the intra-compartmental pressure using a split catheter introduced into the compartment. The difference between diastolic pressure and compartment pressure is known as differential pressure (ΔP). If the differential pressure (ΔP) is less than 30 mm Hg, immediate compartment decompression is indicated.

Treatment:

• Prompt decompression of the affected compartment is done.

• All the dressings, bandages, and cast on the affected limb are removed and the limb is nursed flat.

• Fasciotomy is done if the differential pressure is less than 30 mm Hg

• Fasciotomy wounds are left open and observed after 2 days. If there is necrosis, debridement is done. If the wounds are healthy, they can be sutured without tension or skin grafting can be done.

Solution to Question 13:

Option C is the correct answer. The given lesions and the correctly matched best modes of management are as follows:

Osteosarcoma: is the most common primary malignant tumor of the bone. It is a high-grade medullary osteoid-producing tumor. The preferred method of treatment of osteosarcoma is wide resection and reconstruction. This is mainly owing to the possibility of limb salvage. Amputation is considered only when tumor-safe surgical margins are compromised.

Multi-agent neoadjuvant chemotherapy with agents like doxorubicin, cisplatin, ifosfamide, and methotrexate is started immediately after diagnosis for 8-12 weeks. The tumor is then re-staged followed by wide resection and limb reconstruction.

Radiographic findings include - a periosteal reaction is the typical finding and it may take the form of a Codman's triangle or it may have a sunburst (sunray) appearance on X-ray. It is shown in the image below:

Bone tumor	Management
1. Osteosarcoma	a. Wide excision

Bone tumor	Management
2. Simple bone cyst	d. Intralesional steroid
3. Giant cell tumor	b. Curettage
4. Osteoid osteoma	c. Radiofrequency ablation

Simple bone cysts: these are cystic bone cavities that are often solitary and unilocular filled with serous or serosanguinous material. The lesions usually regress with bone maturity. In cases where the lesions are active, an injection of steroids is done. Only when there is a high risk of fracture, curettage or fixation and stabilization may be required.

It is seen as a well-outlined lytic centrally placed lesion with cortical thinning on X-ray. When there is a fracture, a small fragment may be visualized within the cavity, known as the fallen leaf sign. The image below shows a bone cyst in the humerus along with a pathological fracture and the fallen leaf sign:

Giant cell tumor: it is a benign, locally aggressive tumor of the bone due to the proliferation of mononuclear cells with scattered macrophages.

In stages 1 and 2, intralesional curettage with debridement of the lesional wall is effective. Stage 3 of the disease can be treated through curettage but some cases require en-bloc resection. Antibodies like denosumab (anti-RANKL) have been effective in controlling osteoclastic activity.

Radiographically, they are observed as an eccentrically located solitary lytic lesion in the metaphyses. They show a characteristic soap bubble appearance and periosteal new bone formation is absent.

Below image shows the soap bubble appearance seen in a giant cell tumor:

Osteoid osteoma: it is a benign tumor formed of osteoid and woven bone surrounded by a halo of reactive bone. The preferred method of treatment in this condition is CT-guided radiofrequency ablation to eradicate the lesion and improve symptoms.

Radiography shows a sclerotic peripheral rim with central radiolucent nidus:

Solution to Question 14:

In nerve injuries, an intact outer sheath and nerve fiber with damaged inner axons is referred to as axonotmesis.

According to Seddon's classification, peripheral nerve injuries can be classified as follows:

• Neurapraxia: This refers to the temporary failure of nerve function without physical axon disruption. There is no degeneration, and normal axonal function typically returns within hours to months, often within the 2-4 week range.

• Axonotmesis: This involves the disruption of axons and myelin while the surrounding connective tissue (including endoneurium and perineurium) remains intact. Axonal degeneration occurs both proximally and distally from the site of injury. Distal degeneration is known as Wallerian degeneration. Recovery may occur after 18 months.

• Neurotmesis: This refers to the disruption of axons and endoneurial tubes, with the degeneration of both proximal and distal axons.

Axonapraxia is a disruption to the axoplasmic flow of neurochemicals temporarily.

Orthopedics NEET 2021

Question 1:

A 20-year-old male patient presented with a history of lower backache and early morning stiffness for two years. He also gave a history of bilateral heel pain since 6 months. Which of the following is the most likely diagnosis?

a) Ankylosing spondylitis

b) Tuberculosis of spine

c) Disc prolapse

d) Mechanical pain

Question 2:

A 5-year-old child was brought to the pediatrician with complaints of bilateral knee joint pain. The x-ray image of the joints is given below. What is the most likely diagnosis?

a) Scurvy

b) Rickets

c) Metaphyseal dysplasia

d) Pycnodysostosis

Question 3:

The following test is done to assess

a) Posterior interosseous nerve

b) Median nerve

c) Ulnar nerve

d) Musculocutaneous nerve

Question 4:

A 30-year-old male patient presented with complaints of a gradually progressive swelling around his wrist joint for 3 months. Given below is the image of the swelling and the X-ray film. What is the most likely diagnosis?

a) Ewing's sarcoma

b) Osteosarcoma

c) Osteoclastoma

d) Osteochondroma

Question 5:

A patient was brought to the hospital with complaints of pain around the left hip joint following a road traffic accident. On examination, the affected limb was flexed, adducted, and medially rotated with obvious shortening. What is the most likely diagnosis?

a) Anterior hip dislocation

b) Posterior hip dislocation

c) Transcervical fracture

d) Intertrochanteric fracture

Question 6:

A 28-year-old man presented with complaints of backache, morning stiffness, and redness of the eyes. An X-ray image of the spine is given below. Which of the following is the most likely diagnosis?



a) Rheumatoid arthritis

b) Paget's disease

c) Osteopetrosis

d) Ankylosing spondylitis

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	c
5	b
6	d

Detailed Explanations

Solution to Question 1:

The given clinical scenario of a young male presenting with lower backache, heel pain, and early morning stiffness is suggestive of ankylosing spondylitis.

Ankylosing spondylitis is commonly seen in young males and has a strong association with HLAB27. The patient usually presents with complaints of recurring backache and early morning stiffness. Later the patient can develop other symptoms like generalized fatigue, pain, swelling of the joints, foot strain, tenderness at insertion of the achilles tendon, and intercostal pain.

Extra-articular manifestations include iritis, pericarditis, aortic incompetence, apical lobe fibrosis, generalized osteoporosis, and atlantoaxial joint subluxation.

On examination, there is a loss of normal lumbar lordosis, increased thoracic kyphosis, diminished spinal movements, and decreased chest expansion.

Other findings are:

• Positive Fleche's test - It is done to assess cervical spine involvement. Here the patient is asked to touch the wall with the back of the head without raising the chin.

• Positive tests for sacroiliac joint involvement such as the pump handle test, pelvic compression test, and Fabre's test.

ASAS diagnostic criteria for ankylosing spondylitis:

Sacroiliitis on imaging plus ≥ 1 SpA feature:	or	HLA-B27 +ve plus ≥2 other SpA features
Sacroiliitis on imaging:Active (acute) inflammation on MRI highly suggestive of SpA-associated sacroiliitisand/orDefinite radiographic sacroiliitis according to modified New York criteria (grade >2 bilateral or grade 3 - 4 unilateral)		SpA features:Inflammatory back painArthritisEnthesitis (heel)Anterior uveitisDactylitis PsoriasisCrohn's disease or ulcerativecolitisGood response to NSAIDsFamily history of SpAHLA-B27Elevated CRP



X-ray findings in ankylosing spondylitis:

• Sacro-iliac joints:

• Cardinal sign/earliest sign– erosion and fuzziness of the sacroiliac joints

• Later- peri-articular sclerosis and bony ankylosis of the joint

• Vertebral changes:

• Earliest- flattening of the normal anterior concavity of the vertebral body ('squaring')

• Later- ossification of the ligaments around the intervertebral discs produces syndesmophytes (bridges) between adjacent vertebrae

• Bamboo spine appearance- due to bridging at several vertebral levels

• Long-standing cases are associated with osteoporosis. Wedging of the vertebral bodies produces hyperkyphosis of the spine.

• Progressive bony ankylosis or erosive arthritis may be seen in peripheral joints.

Bamboo-spine appearance is seen in the Xray of the lumbar spine below:



Other options:

Option B: TB spine (Pott's disease) is the commonest form of bone and joint tuberculosis. The clinical presentation can be variable with pain being the common symptom. The pain can be diffuse initially but later on, it gets localized to the affected segment. Stiffness is also seen as an early symptom. The patient can also present with swelling due to the formation of a cold abscess. Paraplegia and spinal deformities can occur at later stages.

Option C: Disc prolapse refers to protrusion or extrusion of the nucleus pulposus through a rent in the annulus fibrosis. The patient presents with low backache which can be acute or chronic. The patient can also present with pain radiating to the gluteal region, back of the thigh, and leg. This is referred to as sciatic pain. There can be paraesthesia in the dermatomes of the affected nerve roots.

Option D: Mechanical back pain is a broader term that includes all the causes that cause alternation in the spine mechanics. It includes conditions like osteoarthritis, lumbar spondylosis, osteoporosis, etc.

Solution to Question 2:

The clinical presentation and the x-ray showing Pelkan spurs, Trummerfeld zone, and Frenkel's line are suggestive of scurvy.

Rickets can also present with bilateral knee pain, but normal bone mineral density, lack of radiographical signs such as cupping and fraying of the metaphysis and widening of the growth plate makes it an unlikely diagnosis.

Radiological findings in scurvy:

• Frenkel's line - The zone of provisional calcification at the growing metaphysis is seen as a dense, white line called the Frankel's line.

• Pelkan spurs - These are metaphyseal spurs formed due to fractures manifesting at the cortical margins.

• Wimberger's ring sign - The sclerotic rim around the epiphysis forms this appearance.

• Trummerfeld zone. - Lucent metaphyseal band underlying Frankel line.



Scurvy results from vitamin C deficiency, which is required for the hydroxylation of lysine and proline in procollagen. There is a defective formation of connective tissue and collagen in the skin, bone, cartilage, dentine, and blood vessels leading to their fragility.

Early manifestations include irritability, loss of appetite, musculoskeletal pain, and tenderness in the legs. Later on, the child can have swelling of the knees, ankles, and pseudo paralysis. Hemorrhagic manifestations of scurvy include petechiae, purpura, ecchymoses at pressure points, epistaxis, gum bleeding, and characteristic perifollicular hemorrhages.

Other manifestations are poor wound and fracture healing, hyperkeratosis of hair follicles, arthralgia, and muscle weakness. It also results in defective osteoid formation. Costo-chondral separation leads to the formation of tender nodules, which are sharply angulated also known as the scorbutic rosary.

Other options:

Option B: The findings in rickets are given in the pearl below.

Option C: Metaphyseal dysplasia (Pyle's disease) is an autosomal recessive condition caused due to failure of remodeling of the metaphysis. Erlenmeyer flask deformities of the distal femur and proximal tibia are seen in this condition. The patient can also develop genu valgum.

Option D: Pycnodysostosis is an autosomal recessive condition due to a mutation in the cathepsin K (CTSK) gene. It causes osteosclerosis and the affected individual has distinctive facial features and skeletal malformations. The patient has short stature and is prone to recurrent fractures due to their fragile bones.

Solution to Question 3:

The given image shows the pen test which is used to assess the median nerve.

It tests the abductor pollicis brevis muscle which is supplied by the median nerve. The hand must remain flat with the palm facing upwards; a pen is held above the palm and the patient is asked to touch the pen with his thumb - positive sign is unable to touch the pen.

Median nerve lesions can be high lesions or low lesions depending on the site of injury.

Low lesions - It occurs due to injuries at the distal third of the forearm such as carpal dislocation or cut wound of the wrist. There is paralysis of the hand muscles; the patient is unable to abduct the thumb and has a loss of sensation over the radial three-and-a-half digits. The forearm muscles supplied by the median nerve are spared in these lesions. Wasting of thenar eminence is seen in long-standing cases.

High lesions - It occurs due to injuries above the elbow such as forearm fractures or elbow dislocation. There is paralysis of the forearm muscles along with all the features of low lesions.

Muscles supplied by the median nerve are given in the table:

Other tests for median nerve:

Pointing index sign or Oschner's clasp test: The patient is asked to clasp both hands together. When this is done the index and the middle finger fail to flex due to the loss of action of long finger flexors which are supplied by the median nerve.

Benediction test: Due to loss of activity of the long finger flexors the patient is unable to flex the index and middle finger on lifting the hand. As this is the position a clergyman uses to bless the couple during the marriage, it is called the benediction test.

Level	Motor supply
In the arm	No muscles
Proximal one-third of the for earm	All the flexor muscles of the proximal forearm, except the flexor carpi ulnaris and the medial half of the flexor digit orum profundus

Level	Motor supply
Distal one-third of the forear m	No muscles
In the hand	Abductor pollicis brevisOppo nens pollicisFlexor pollicis br evisFirst two lumbricals

Solution to Question 4:

The clinical presentation and the x-ray showing an eccentric expansile lesion with a soap bubble appearance in a 30-year-old man are suggestive of osteoclastoma or giant cell tumor (GCT).

Osteoclastoma usually affects young adults in the age group of 20-40 years. The most common location of the tumor is the distal femur followed by the proximal tibia. Involvement of the distal end of the radius is also seen and the tumor is more aggressive in this location.

Radiographic findings in GCT:

• A solitary, possibly loculated, lytic lesion in the epiphysis

• Eccentric location, most often subchondral

• Expansion of the overlying cortex

• Soap bubble appearance - The tumor is homogeneously lytic with trabeculae of the remnants of bone traversing it, giving it a loculated appearance called soap bubble appearance

• No calcification within the tumor

• No or minimal reactive sclerosis around the tumor.

Treatment :

• Extended curettage and bone grafting.

• En-bloc wide resection and reconstruction are done for advanced stages, local recurrence, and tumors not responding to other methods.

• Resection without reconstruction is done for lesions in expandable bones like distal ulna, clavicle, etc.

• Irradiation and/or embolization are used for inoperable lesions in the spine and pelvis.

• Bisphosphonates have been shown to reduce the proliferation of osteoclasts and hence have been shown to stabilize local and metastatic disease.

• Denosumab is approved for use in unresectable tumors in skeletally mature patients.

Other options:

Option A: Ewing's sarcoma is a primary malignant tumor of the bone commonly involving the age group of 4-25 years involving the diaphysis of the long bones like the femur, tibia, fibula, and humerus.

X-ray shows onion-peel appearance due to deposition of periosteal reaction in layers is the typical finding.

Option B: Osteosarcoma is the most common non-hematologic primary malignancy of bone. It occurs most commonly in the second decade of life and is typically metaphyseal. The distal femur, proximal tibia, and proximal humerus are the common sites.

X-ray shows periosteal reaction and it may take the form of a Codman's triangle or have a sunburst (sunray) appearance.

Option D: Osteochondromas are common benign bone tumors. The lesions consist of a bony mass, often in the form of a stalk, produced by endochondral ossification of a growing cartilaginous cap. The common sites of involvement are the distal femur, proximal tibia, and proximal humerus.



Solution to Question 5:

The given clinical vignette showing a flexed, adducted, and medially rotated limb with apparent limb shortening following trauma is suggestive of posterior hip dislocation.

Posterior hip dislocation is the most common type of hip dislocation commonly seen following dashboard injuries.

X-ray findings:

• Upward dislocation of the femoral head

• Femoral head appears smaller than normal

• Lesser trochanter is not visualized as the thigh is internally rotated

• Femoral head and posterior acetabular wall fractures may also be seen.

Treatment: Early reduction should be done as the risk of osteonecrosis and subsequent osteoarthritis increases with time.

Closed reduction under sedation or general anesthesia is done in most cases. Closed reduction should not be attempted in associated femoral neck fractures as it can lead to the displacement of the femoral neck and disrupt the blood supply to the femoral head.

If there is a failure of closed reduction or any significant fractures then surgical repair should be done. If there is a delay in surgical correction skeletal traction with a distal femoral traction pin can be done.

Early complications include sciatic nerve injury, vascular injury (superior gluteal artery tear can occur), and concurrent femoral shaft fracture (when both occur together hip dislocation may be missed).

Late complications include osteonecrosis of the femoral head, myositis ossificans, secondary osteoarthritis, and recurrent instability.

The image given below shows posterior hip dislocation on the left side.

Other options:

Option A: Anterior hip dislocations are less common than posterior hip dislocations. This injury is usually sustained following a high-energy accident. The affected limb will be abducted, externally rotated, and slightly flexed.



Option C: Transcervical fracture is a type of fracture neck of the femur where the fracture line is in the middle of the neck. In this case, there can be shortening and external rotation of the affected limb.

Option D: Intertrochanteric fracture is usually sustained following a fall or road traffic accident. The leg is short and externally rotated. Clinical findings are more marked than the neck of femur fractures.

Solution to Question 6:

The clinical presentation of a young male with backache, morning stiffness, redness of the eye, and the x-ray showing the classical bamboo-spine appearance are suggestive of ankylosing spondylitis.

Ankylosing spondylitis commonly involves young male adults. It has a strong association with HLAB27.

Radiographic findings in ankylosing spondylitis:

• Sacro-iliac joints:

• Cardinal sign/earliest sign– erosion and fuzziness of the sacroiliac joints

• Later- peri-articular sclerosis and bony ankylosis of the joint

• Vertebral changes:

• Earliest- flattening of the normal anterior concavity of the vertebral body ('squaring')

• Later- ossification of the ligaments around the intervertebral discs produces syndesmophytes (bridges) between adjacent vertebrae

• Bamboo spine appearance- due to bridging at several vertebral levels

• Long-standing cases are associated with osteoporosis. Wedging of the vertebral bodies produces hyperkyphosis of the spine.

• Progressive bony ankylosis or erosive arthritis may be seen in peripheral joints.

ASAS diagnostic criteria for ankylosing spondylitis:

Sacroiliitis on imaging plus ≥ 1 SpA feature:	or	HLA-B27 +ve plus ≥2 other SpA features
Sacroiliitis on imaging:Active (acute) inflammation on MRI highly suggestive of SpA-associated sacroiliitisand/orDefinite radiographic sacroiliitis according to modified New York criteria (grade >2 bilateral or grade 3-4 unilateral).		SpA features:Inflammatory back painArthritisEnthesitis (heel)Anterior uveitisDactylitis PsoriasisCrohn's disease or ulcerativecolitisGood response to NSAIDsFamily history of SpAHLA-B27Elevated CRP



Other options:

Option A: Rheumatoid arthritis is a chronic inflammatory disease with an increased incidence between 25 and 55 years of age. Clinical features include swelling and pain in the small joints of the hands. Metacarpophalangeal (MCP) and proximal interphalangeal (PIP) joints are commonly

involved. The patient also has morning stiffness in the affected joints.

Option B: Paget's disease, also known as osteitis deformans is characterized by an increased bone turnover due to a disorganized, alternating, and increased rate of osteoblastic and osteoclastic activity. It is seen more commonly in people over the age of 50 and most of the patients are asymptomatic. Symptomatic patients present with pain or deformity, with the pelvis and tibia being the commonly involved sites.

Option C: Osteopetrosis/Albers–Schönberg disease occurs due to a defect in the genes that control the osteoclastic activity. It can be of two types:

• Infantile form(autosomal recessive)- The infant presents with growth retardation, abnormal skull development, failure to thrive, and genu valgum. Pancytopenia and hepatosplenomegaly occur as a result of obliteration of the medullary canals and the subsequent extra-medullary hemopoiesis

• Adult form (autosomal dominant) - The patient can be asymptomatic or can present with recurrent fractures and compressive neuropathies. Hemopoiesis is unaffected.

Orthopedics INI-CET 2022

Question 1:

The most common pattern of involvement in Pott's spine is

a) Paradiscal

b) Central

c) Anterior

d) Posterior

Question 2:

Anterior dislocation of the shoulder is likely to result in the damage of which of the following nerves?

a) Axillary nerve

b) Median nerve

c) Radial nerve

d) Suprascapular nerve

Question 3:

A 60 year old patient presents with limping gait and left hip pain. What is the likely diagnosis?

a) Tuberculosis of the hip

b) Rheumatoid arthritis

c) Ankylosing spondylitis

d) Avascular necrosis

Question 4:

A 72-year-old man came with complaints of unilateral knee pain. X-ray examination showed reduced joint space. Pathological findings shown below are of:



a) Osteoarthritis

b) Tuberculosis of the knee

c) Rheumatoid arthritis

d) Lyme disease

Question 5:

Which of the following is false regarding hip dislocation?

a) Posterior dislocation of hip has flexion and adduction

b) Vessel injury more common in anterior

c) Nerve injury more common in anterior

d) Posterior dislocation is more common than anterior

Question 6:

A 35-year-old man presented with knee pain. The x-ray of his knee joint is given below. What is the probable diagnosis?

a) Chondroblastoma

b) Osteochondroma

c) Osteoclastoma

d) Fibrous dysplasia

Question 7:

Book test is used to assess the function of adductor pollicis. Which of the following nerves supplies this muscle?

a) Posterior interosseous nerve

b) Ulnar nerve

c) Radial nerve

d) Median nerve

Question 8:

A 6-year-old girl sustained a fracture of the lateral condyle of the humerus which led to the development of elbow deformity. Which of the following findings can be seen in this patient?

a) Ape thumb deformity

b) Pointing index sign

c) Inability to adduct the fingers

d) Kiloh-Nevin sign

Question 9:

A 65-year-old postmenopausal female patient comes with persistent backache not responding to conservative treatment. She has a history of lifting heavy weights 4 months back and progressively increasing pain that worsens on walking. The pain starts radiating down her legs after walking for a distance of around 100 m. However, she feels no pain on walking uphill. What is the most probable diagnosis?

a) Osteoporotic compression fracture

b) Lumbar stenosis

c) Thromboangiitis obliterans

d) Atherosclerosis

Question 10:

Disruption of the axon with an intact endoneurium is seen in which of the following?

a) Neurapraxia

b) Axonapraxia

c) Axonotmesis

d) Neurotmesis

Question 11:

A patient presents to the OPD with Trendelenburg sign positive on one side. Which of the following muscles are involved?

a) 2 only

b) 1,3

c) 2,4

d) 1,4

Question 12:

You visit a newborn near your house and notice that the baby has CTEV, as shown below. When would you advise the baby's parents for a consultation to put on a cast?

a) In 1 month

b) In 6 months

c) In a few days

d) When the child starts walking

Question 13:

Combination of appearance in CTEV

a) Equinus, eversion, forefoot adduction, cavus

b) Equinus, inversion, forefoot adduction, cavus

c) Equinus, eversion, forefoot abduction, cavus

d) Equinus, inversion, forefoot adduction, planus

Question 14:

A child presents with poor growth and swelling at joints. A radiograph of his wrist is given below. Lab investigations reveal serum ALP levels of >1500. What is the possible diagnosis?



a) Scurvy

b) Rickets

c) Osteogenesis imperfecta

d) Paget's disease

Question 15:

What is not required on an emergency basis in an open fracture?

a) 1, 2, 3

b) 2 only

c) 2 and 4

d) 2 and 3

Question 16:

Which of the following structures are involved in the unhappy triad?



a) A, D, E

b) D, E, F

c) A, B, C

d) C, D, E

Question 17:

A radiograph of the wrist is given below. Identify the true statements regarding this condition.



a) 1, 3

b) 2, 3

c) 3 only

d) 2, 4

Answer Key

Question No.	Correct Option
1	a
2	a
3	d
4	a
5	c
6	c
7	b
8	c
9	b
10	c
11	a
12	c
13	b
14	b

15	b
16	c
17	c

Detailed Explanations

Solution to Question 1:

The most common pattern of involvement in Pott's spine is the paradiscal type.

Pott's spine occurs secondary to a pulmonary or gastrointestinal primary and spreads through the Batson's plexus of veins (paravertebral plexus) and most commonly involves the dorso-lumbar region.

Types of vertebral tuberculosis:

• Paradiscal - contiguous areas of two adjacent vertebrae and the middle intervening disc are affected. This is due to the high oxygen requirements of tubercular bacteria and the corresponding high blood supply in this region by the segmental vessels.

• Central - the body of the vertebra is involved leading to collapse (causing wedging or concertina collapse) of the affected vertebra. This leads to spinal deformities such as kyphosis and gibbus.

• Anterior - the anterior part of the body is affected and spreads through the anterior longitudinal ligament.

• Posterior - the posterior complex (pedicle, lamina, spinous process, and transverse process) is affected.

the image below shows various parts of the vertebra and patterns of involvement in Pott's spine:





Clinical features of spinal tuberculosis:

• Back pain, stiffness and reduced movements are the usual early complaints

• Cautious gait - a person with spinal tuberculosis will walk carefully with short steps to avoid pain resulting from movement

• Pott's paraplegia

• Cold abscess

• Concertina collapse - a type of compression fracture without involvement of the interverterbral disc

• Deformity - kyphus and gibbus (gradually increasing prominence).

Images showing concertina collapse and the deformities seen TB spine.





Solution to Question 2:

Anterior shoulder dislocation is most likely to damage the axillary nerve.

Axillary nerve injury presents with an anaesthetic patch over the deltoid muscle, and deltoid muscle weakness (inability to abduct at the shoulder joint). It is usually a neuropraxia type of injury and resolves spontaneously. Other nerves that can be injured include the radial nerve, musculocutaneous nerve, median nerve or ulnar nerve.

Other complications of anterior dislocation of the shoulder joint:

• Early complications

• Rotator cuff injury is seen in older people and presents with difficult abduction (palpable contraction of deltoid muscle rules out nerve injury).

• Vascular injury: Axillary artery may be injured, especially in older individuals, either at the time of injury or during the reduction.

• Fracture-dislocation: There may be an associated fracture of the humerus (e.g. greater tuberosity) due to the traumatic injury. This may require open reduction and internal fixation.

• Late complications

• Recurrent dislocations: Most common sequelae of anterior dislocation. Usually seen when there is associated injury to the glenoid labrum or stripping of the capsule is present.

• Shoulder stiffness: Due to prolonged immobilization.

• Untreated dislocation: Sometimes it may go unnoticed, especially in older and unconscious individuals. The reduction can be tried up to 6 weeks after the dislocation, after which 'active neglect' (no treatment) is followed.

Radiograph showing anterior dislocation with humeral
head lying below and medial to the glenoid socket.

Solution to Question 3:

The given scenario of limping gait and left hip pain with radiographic findings of the radiolucency and distorted articular surface is suggestive of avascular necrosis of the head of femur.

Avascular necrosis (AVN) or osteonecrosis is a painful bone condition as a result of diminished blood supply mainly due to trauma or non-traumatic etiologies.

The typical features are pain with movements in or near the joint, click in the joint, joint stiffness, limping gait, and deformity. Restriction of movements is also a feature of this condition.

An x-ray shows reactive new bone formation, increased radiolucency in the subchondral bone, and a distorted articular surface.

MRI is the investigation of choice. It allows the identification of AVN of the femoral head and also determines the exact stage and extent of the pathologic process.

Management is based on Ficat and Arlet Classification:

Other options:

Option A: Tuberculosis of the hip is characterized by painful and restricted movements of the hip. The radiographic findings are not appreciable in the early stages. But late stages show periarticular osteoporosis, hazy, and irregular joint margins with a reduction in the joint space, and wandering acetabulum.

Option B: Rheumatoid arthritis (RA) is usually seen in females in the 4th to 5th decade. Patients present with symmetrical polyarthritis of the proximal finger joints, tenosynovitis, early morning stiffness and pain. RA of the hip can be bilateral. Radiographic findings such as joint space narrowing due to hyaline cartilage destruction, cyst formation, and acetabular protrusion-type deformity can be seen.

Option C: Ankylosing spondylitis is a chronic inflammatory disease of the spine and sacroiliac joint. It typically occurs in the age group 15-25 years with a male preponderance. Features include backache, stiffness, sciatica, and fixed flexion contracture. Radiographic findings show erosion of

sacroiliac joints, squaring, and bridging of the vertebral body leading to a bamboo spine appearance.

Stage	Findings	Management	Method
1	X-ray normal, MRI +ve	Conservative	Bisphosphonates
2a	X-ray: sclerosis, cysts, no collapse	Conservative	Bisphosphonates
2b	X-ray: crescent sign, no collapse	Surgical decompression	Core decompression
3	Collapse+loss of sphericity	Surgical	Osteotomy
4	Advanced arthritis	Reconstructive	Total hip replacement

Solution to Question 4:

The scenario of unilateral knee pain, the radiograph findings of reduced joint space, and the histopathology findings of loss of articular cartilage suggest osteoarthritis.

Osteoarthritis is a type of joint disease characterized by degeneration of cartilage leading to synovial joint destruction. Individuals with age > 50 years and overweight are usually affected.

The disease typically involves the knee and initially affects a single joint; eventually, both knees are affected. Pain is the predominant symptom and worsens with use. Other symptoms such as joint stiffness, swelling, giving away or locking can be present. Fixed flexion deformity can be seen in these patients.

A radiograph of the knee joint, the anteroposterior and lateral view is useful to establish a diagnosis. The characteristic findings are the diminution of joint space, subchondral sclerosis, subchondral cysts, osteophytes, and soft tissue calcifications.

Histopathology shows characteristic loss of the articular cartilage, congestion, fibrosis and inflammatory cells.

Management is as follows

• Conservative measures such as weight loss, exercise, physiotherapy for strengthening of the quadriceps muscle, analgesics for pain relief and usage of walking sticks for reducing the load of joints.

• Intraarticular injections of corticosteroids and hyaluronidase are useful.

• Operative measures such as osteotomy and total knee replacement surgery.

Other options:

Option B: Tuberculosis of the knee is more common in childhood with pain and limping being the common symptoms. X-rays show an enlargement of the epiphysis. The characteristic histopathologic findings are caseating and noncaseating granulomas with surrounding epithelioid and multinucleate giant cells.

The below image shows caseating granuloma with Langhans giant cells:



Option C: Rheumatoid arthritis (RA) of the knee is less common. It is usually seen in females in the 4th to 5th decades. It presents as symmetrical polyarthritis of the proximal finger joints, tenosynovitis, early morning stiffness and pain. Histopathological features include synovial cell hyperplasia and proliferation with inflammatory infiltrates and rheumatoid nodules. These are necrotising granulomas surrounded by fibrinoid necrosis.

The below image shows rheumatoid nodules:



Option D: Lyme disease is a tick-borne infection with skin lesions and flu-like symptoms. It presents with asymmetric inflammatory polyarthritis of the large joints when left untreated.

Solution to Question 5:

Statement C is false. Nerve injury is more common in posterior hip dislocation. The sciatic nerve, lying behind the posterior wall of the acetabulum is most commonly involved here.

Sciatic nerve injury is seen in 10-20 % of cases with posterior hip dislocation. It is usually a neuropraxia type of injury and resolves spontaneously. But, if it is suspected after a reduction maneuver, immediate exploration must be carried out.

Posterior hip dislocation is the commonest type of hip dislocation (80%), commonly seen following dashboard injuries. The patient presents with flexion, adduction, and internal rotation deformities.

X-ray findings include:

• Femoral head lying above and behind the acetabulum and appears smaller than normal

• Lesser trochanter is not visualized as the thigh is internally rotated

• Femoral head and posterior acetabular wall fractures may also be seen

Treatment includes an early reduction to reduce the risk of osteonecrosis and osteoarthritis. Closed reduction under sedation or general anaesthesia is done in most cases. If there is an associated femoral neck fracture, the closed reduction should not be attempted as it can lead to the displacement of the femoral neck and disrupt the blood supply to the femoral head.

Anterior hip dislocation is less common than posterior hip dislocations and accounts for about 10-20% of all hip dislocations. This injury is usually sustained following a high-energy accident. The affected limb in this condition will be abducted, externally rotated, and slightly flexed.

Note: Although vascular injuries are seen in posterior dislocations as well, they are most likely with anterior dislocation. This is because of the femoral artery lying anterior to the head. (statement B)

Solution to Question 6:

The given clinical scenario and the x-ray showing a lytic lesion at the end of a long bone with a soap-bubble appearance suggest a giant cell tumor (GCT)/ osteoclastoma.

Osteoclastoma is a benign but locally aggressive tumor that usually affects young adults of age 20 to 40 years with slight female predominance. It affects the ends of long bones, involving metaphysis/epiphysis of long bones. GCT is most commonly seen in the distal femur, proximal tibia, proximal humerus, and distal radius. Common presenting complaints are swelling and vague pain. Sometimes, the patient presents for the first time with a pathological fracture through the lesion.

Important X-ray features of a GCT:

• A solitary, possibly loculated, lytic lesion in the epiphysis.

• Eccentric location, most often subchondral.

• Expansion of the overlying cortex.

• Soap bubble appearance - The tumor is homogeneously lytic with trabeculae of the remnants of bone traversing it, giving it a loculated appearance called soap bubble appearance.

• No calcification within the tumor.

• No or minimal reactive sclerosis around the tumor.

Histologically, GCTs consist of undifferentiated spindle cells profusely interspersed with multi-nucleated giant cells. The tumor stroma is well-vascularized with bands of cellular or collagenous fibrous tissue.

The image below shows the histopathology of osteoclastoma:



The treatment of choice for GCT is extended curettage and bone grafting.

• GCT in the relatively insignificant bone like the distal ulna, clavicle, and proximal fibula is treated by resection without reconstruction.

• Bisphosphonates have been shown to reduce the proliferation of osteoclasts and hence have been shown to stabilize local and metastatic disease.

• Radiation can be used for spinal or sacral tumors.

Other options:

Option A: Chondroblastoma is a benign tumor that usually appears as well-circumscribed round or oval lytic lesions in the center of the epiphysis of a long bone. It presents as a painful mass that causes effusion or stiffness. It is most commonly seen in ages 10-20 years.

The following is an x-ray image of the knee showing chondroblastoma:

Option B: Osteochondromas are common benign bone tumors commonly present in adolescents. The lesions consist of a bony mass, often in the form of a stalk, produced by endochondral ossification of a growing cartilaginous cap. They are usually painless. The common sites of involvement are the distal femur, proximal tibia, and proximal humerus.

The following is an x-ray image of the knee showing osteochondroma of the distal knee:



Option D: Fibrous dysplasia is a condition that affects bone development and is characterized by the replacement of the bone with fibrous tissue. The patient presents with a progressive limb deformity or a fracture through a previously asymptomatic lesion which often occurs in the second decade. Other symptoms include nonspecific bone pain, tenderness, and swelling in the affected area. The commonly involved sites are the proximal femur, tibia, pelvis, and foot.

Polyostotic fibrous dysplasia is associated with McCune-Albright syndrome.

Radiological findings include:

• Well-defined, expansile intramedullary lesions

• Hazy ground glass appearance is seen in the majority of cases

• Shepherd's crook deformity - Expansion with marked varus in the proximal femur

The given image shows the common locations of various bone tumors:



Solution to Question 7:

The deep terminal branch of the ulnar nerve supplies the adductor pollicis muscle.

A positive book test or Froment's sign indicates weakness of the adductor pollicis muscle, due to ulnar nerve weakness.

In this test, the patient is asked to hold a book between the thumb and index finger against resistance, and the sign is positive if there is flexion at the interphalangeal joint. This is because if the weakness of adductor pollicis is present, thumb adduction is weak; the flexor pollicis longus muscle compensates for this weakness by inducing flexion of the interphalangeal joint.

Adduction paralysis occurs as a result of low lesions of the ulnar nerve, which are commonly caused by cuts on shattered glass. The patient has numbness in the medial one-and-a-half fingers. The hand assumes the claw-hand deformity, with hyperextension of the metacarpophalangeal joints of the ring and little fingers, due to weakness of the intrinsic muscles. Hypothenar and interosseous wasting may be seen in the affected hand when compared to the normal hand.

The image below shows the test being demonstrated:



Solution to Question 8:

The finding that can be seen in this patient with a fracture of the lateral condyle of the humerus is the inability to adduct the fingers. This is due to tardy ulnar nerve palsy.

In this condition, there is stretching of the ulnar nerve due to cubitus valgus deformity that occurs as a result of long-standing non-union of a fractured lateral condyle of the humerus.

The normal carrying angle of the elbow is 13 degrees. A cubitus valgus deformity is an angle more than this or an asymmetry between the two sides.

The image below shows cubitus valgus deformity.



The ulnar nerve is most commonly injured in flexion type of supracondylar fracture. In ulnar nerve injury at the wrist, there
is hyperextension at metacarpophalangeal joints and flexion at interphalangeal joints. This causes claw hand deformity. There is numbness in the medial one-third of the palm and the medial one-and-a-half fingers.

The patient is unable to abduct the fingers due to paralysis of the dorsal interossei. The loss of adduction is due to paralysis of the palmar interossei muscles.

A card test is done to test the function of palmar interossei muscles.



Other options:

Option A: Ape thumb deformity is characterized by the inability to abduct the thumb due to paralysis of the thenar muscles supplied by the median nerve. The thumb remains in the same plane as the wrist.

Option B: In median nerve injury of the forearm or elbow, there is paralysis of long flexors of the digits. This leads to the characteristic pointing index finger.

Option C: The inability to perform the 'OK' sign is called as Kiloh Nevin sign. It is seen in the involvement of the anterior interosseous nerve. The patient is unable to bend the terminal phalanx of the thumb and index finger joints due to weakness of the flexor pollicis longus and lateral half flexor digitorum profundus.

Solution to Question 9:

The given clinical scenario of an old woman with back pain that worsens on walking and relieves walking uphill points to the diagnosis of lumbar stenosis.

Walking uphill may be preferred as it flexes the spine and maximizes the spinal canal capacity, relieving the pain.

Spinal stenosis is the narrowing of the spinal canal, causing nerve compression. It can be congenital or acquired, with degeneration being the most common cause. Other factors include conditions like spondylolysis, trauma, infection, and metabolic disorders.

Degenerative spinal stenosis involves disc degeneration, facet joint hypertrophy, and ligamentum flavum calcification. These changes lead to reduced spinal canal dimensions and compression of the nerves resulting in neurogenic claudication.

They present with symptoms such as back pain (most common), sciatica, sensory disturbances in the legs, motor weakness, and sometimes urinary disturbances. Pain typically occurs after standing or walking for a few minutes and is relieved by sitting, squatting, or flexing the spine.

MRI is the imaging method of choice for the investigation of spinal stenosis. Conservative treatment is the first line for spinal stenosis, including reassurance, pain relief, exercise, and activity modification.

Surgery is considered if conservative measures fail and there is significant functional impairment and severe neurological symptoms. Posterior decompression is the surgical method. Laminectomy is performed for central stenosis.

Solution to Question 10:

Disruption of the axon with an intact endoneurium is seen in axonotmesis. There is an axonal disruption (inner axon is damaged) and loss of conductivity, with an intact outer sheath.

Axonotmesis (axonal interruption): There is a loss of conduction, but the continuity of the nerve is maintained and the neural tubes are intact. There is disruption of the axon i.e., the continuity of the axon is broken. But the endoneurium and perineurium are intact, maintaining the continuity of the nerve. It occurs following trauma such as closed fractures and dislocations.

Distal to the lesion, Wallerian degeneration of the disrupted axon occurs, where axons disintegrate and are resorbed by phagocytes. It is then followed by axonal regeneration where numerous fine unmyelinated fibres grow from the proximal stumps to the endoneurial tubes.

Other options:

Option A- Neurapraxia is the reversible physiological nerve conduction block causing some loss of sensation and muscle power. There is spontaneous recovery after a few days or weeks. It occurs following prolonged compression, as in Saturday night palsy, and crutch palsy. Here, there is segmental demyelination but the axon is intact. Nerve conduction is slowed because of the loss of myelin. It is followed by the regeneration of the myelin.

Option B- Neurotmesis is the complete transection of the nerve trunk. It is usually associated with open wounds. The endoneurial tubes are destroyed and the scar tissue prevents the regenerating axons from entering the distal segment. The regenerating fibers mingle with proliferating Schwann cells and fibroblasts to form a neuroma at the site of injury.

Option D- Axonapraxia is a disruption to the axoplasmic flow of neurochemicals temporarily.

Solution to Question 11:

The muscles involved in a positive Trendelenburg sign are the hip abductors—the gluteus medius and gluteus minimus.

The gluteus medius and minimus are powerful abductors of the hip supplied by the superior gluteal nerve. Stabilizing the pelvis on the femur during walking is done by these muscles. It is demonstrated by asking the subject to stand on one limb. If the ipsilateral abductors are normal, the pelvis remains in a horizontal position. If the ipsilateral abductors are weak, then the pelvis tilts towards the opposite (unsupported) side. This is called positive Trendelenburg's sign.

Other causes of positive Trendelenburg signs include shortening of the femoral neck, abductor tendon rupture, and dislocation or subluxation of the hip joint.

Patients are known to have a characteristic gait, which is called Trendelenburg or waddling gait. The trunk is thrown from side to side with each step, and the patient lurches toward the affected limb while walking.

The image below depicts Trendelenburg's gait.



Solution to Question 12:

Treatment of CTEV (congenital talipes equinovarus) should begin as early as the first 2 weeks after birth. In CTEV, or idiopathic clubfoot, the deformity is equinus, inversion, forefoot adduction, and cavus.

Ponseti's method is a successful method of manipulation in CTEV. The order of correction according to Ponseti's method is cavus-adduction-varus-equinus (CAVE). Weekly manipulation of the foot is done with the help of plaster casts.

The first metatarsal is elevated initially to correct the cavus. This creates a flat longitudinal medial border of the foot. Then, the forefoot is abducted in the plane of the metatarsals. This everts the hindfoot and corrects the forefoot out of supination. Finally, the equinus is corrected by dorsiflexing the foot at the ankle joint. In most cases, a percutaneous Achilles' tenotomy is required later for complete correction of the equinus deformity.

After correction, a foot abduction brace (FAB) is used to maintain the corrected position, initially full-time (for 3 months) and then night-time and nap-time for 4 years to prevent relapse of the deformity. CTEV shoes are worn during the day once the child starts walking for up to 5 years. They have a straight inner border (prevents adduction), outer shoe raise (prevents foot inversion), and no heel to prevent equinus.

The image below shows the foot abduction brace:

The image below shows the CTEV shoes:

Solution to Question 13:

In congenital talipes equinovarus (CTEV), or idiopathic clubfoot, the deformity is equinus, inversion, forefoot adduction, and cavus. CTEV is a congenital malformation of the foot. It is predominately seen in male children. Calf and medial foot tissues are underdeveloped.

Components of deformity:

• Cavus - This deformity occurs due to exaggerated arching of the foot at the mid-tarsal joint.

• Adduction - This deformity occurs at the talonavicular joint.

• Inversion - This deformity occurs at the subtalar joint. The calcaneum is inverted, and along with it, the whole foot is inverted. This leads to a medially facing sole.

• Equinus - This deformity occurs at the ankle joint and tarsal joints.

Equinovarus deformity causes inward twisting of the foot (shown below), with the sole facing posteromedially in severe cases. Associated disorders like congenital hip dislocation and spina bifida should be evaluated.

Ponseti's method is a successful method of manipulation in CTEV. It should begin as early as the first 2 weeks after birth. Weekly manipulation of the foot is done with the help of plaster casts. The order of correction according to Ponseti's method is cavus-adduction-varus-equinus (CAVE).

The first metatarsal is elevated initially to correct the cavus. This creates a flat longitudinal medial border of the foot. Then, the forefoot is abducted in the plane of the metatarsals. This everts the hindfoot and corrects the forefoot out of supination. Finally, the equinus is corrected by dorsiflexing the foot at the ankle joint. In most cases, a percutaneous Achilles' tenotomy is required later for complete correction of the equinus deformity.

Typically, five casts are required for correction, followed by a cast for 3 weeks. After correction, a foot abduction brace (FAB) is used to maintain the corrected position, initially full-time (for 3 months) and then night-time and nap-time for 4 years. This helps prevent relapse of the deformity. CTEV shoes are worn during the day once the child starts walking for up to 5 years.

Solution to Question 14:

The clinical vignette describing a child with poor growth, the wrist radiograph findings showing cupping and fraying, and the raised ALP levels are suggestive of rickets.

Inadequate mineralization of bone in children is known as rickets. There is inadequate calcification of the growth plate, leading to irregular chondrocyte accumulation, widened growth plate, poor mineralization in the zone of calcification, and sparse bone formation.

Vitamin D disorders, such as vitamin D deficiency or resistance, calcium deficiency, phosphorus deficiency, and renal loss are among the common causes of rickets. Decreased vitamin D levels reduce calcium and phosphate absorption, leading to low blood calcium. This stimulates increased production of parathyroid hormone (PTH) and elevates alkaline phosphatase (ALP) levels involved in bone mineralization.

Infants with rickets may exhibit symptoms such as tetany, convulsions, failure to thrive, listlessness, and muscle weakness. Softening of the skull bones called craniotabes is seen. Delayed closure of fontanels, frontal bossing, and delayed dentition are also seen.

Widening of the costochondral junctions, forming a rachitic rosary along the ribs is seen in the chest. Harrison groove, a horizontal depression along the lower chest, can occur due to rib softening. Vertebral column abnormalities like scoliosis, kyphosis, and lordosis can also appear. In the distal extremities widening of the wrists and ankles, varus/ valgus deformities, windswept deformity (valgus on the side and varus on the contralateral side), and anterior tibial bowing are seen.



The laboratory findings on vitamin D deficiency rickets would be normal/decreased serum calcium, decreased phosphorous, elevated PTH, and increased ALP levels. Thickening and widening of the growth plate, cupping of the metaphysis, and potential bowing of the diaphysis are seen on X-ray.

Children with nutritional vitamin D deficiency should receive vitamin D supplementation and ensure adequate intake of calcium and phosphorus. Two strategies for vitamin D administration are stoss therapy (300,000-600,000 IU over 1 day) and daily supplementation (2,000 IU/day for at least 3 months).

Other options:

Option A: Scurvy is caused by vitamin C deficiency, resulting in bleeding tendencies and various manifestations such as petechiae, gum bleeding, and scorbutic rosary. Radiographically, it shows characteristic findings like the white line of Frankel, Pelkan spurs, and periosteal elevation.

Option C: Osteogenesis imperfecta (Brittle bone disease) is caused by deficiencies in the synthesis of type I collagen. It presents with the hallmark features of osteoporosis with increased fragility of bones along with blue sclera, dental fragility as well as hearing loss.

Option D: Paget's disease of the bone is a condition where there is an increased bone turnover followed by enlargement and thickening of the bone. In this condition, calcium levels are normal with raised ALP.

Solution to Question 15:

Emergency management of open fractures is mainly focused on the prevention of infection by providing antibiotic coverage and splinting (statements 3 and 4). Reduction of the fracture (statement 1) can be done in the emergency room. Internal fixation is not done on an emergency basis for an open fracture due to the risk of infection (statement 2). It is a part of definitive management and is done after the patient is stabilized.

Open fractures are characterized by a break in the overlying skin and soft tissues. Hence, the fracture fragment is exposed to the external environment. There are more prone to be infected.

The major components of the management of open fractures:

• Antibiotic prophylaxis - It should be given as early as possible (within 3 hours of injury). Amoxiclav is commonly used.

• Early debridement of the wound and fracture - This includes the following steps:

• Excision of the wound margins

• Extension of the wound if required

• Delivery of the fracture outside the wound for proper examination

• Removal of the devitalized tissue

• Cleansing of the wound to remove all the foreign materials

• Nerves and tendons are usually left alone, but if the wound is clean, a repair can be considered

• Wound cover:

• In Gustilo Type I and II fractures, wound cover can be done at the time of debridement (early wound cover)

• Delayed wound cover is usually practiced in Type III fractures

• Fracture stabilization: External fixators are used to stabilize the fractures in the emergency room. Definitive management is later done by open reduction and internal fixation (ORIF).

Solution to Question 16:

The unhappy triad of O'Donoghue consists of injury to

• Anterior cruciate ligament (A)

• Medial collateral ligament (B)

• Medial meniscus (C)





Solution to Question 17:

The given x-ray showing a lytic lesion at the end of a long bone is suggestive of a giant cell tumor (GCT)/osteoclastoma. It affects the ends of long bones, involving metaphysis/epiphysis (statement 3). GCT is most commonly seen in the distal femur, proximal tibia, proximal humerus, and distal radius.

GCTs usually affect young adults of age 20 to 40 years (statement 1), with slight female predominance. It is a benign but locally aggressive tumor. They present with swelling, vague pain, or pathological fracture through the lesion.

A solitary eccentric, expansile, lobulated lytic lesion in the epiphysis/metaphysis is seen on a plain X-ray, which is diagnostic (statement 2). The tumor is homogeneously lytic with trabeculae of the remnants of bone traversing it, giving it a loculated appearance called a soap bubble

appearance (shown below). Soft-tissue extension with a mass can also be observed. They usually lack calcification, new bone formation, or periosteal reaction.

Histologically, GCTs display a mixture of meaty, reddish-purple tissue with soft yellow areas. They are characterized by numerous osteoclast-like giant cells dispersed among round or spindle-shaped mononuclear cells (shown below). The tumor stroma is well-vascularized with bands of cellular or collagenous fibrous tissue.



Malignancy transformation is rare (<1%) in GCTs, though possible (statement 4).

The treatment of choice for GCT is extended curettage and bone grafting. GCT in the relatively insignificant bone like the distal ulna, clavicle, and proximal fibula is treated by resection without reconstruction. Bisphosphonates have been shown to reduce the proliferation of osteoclasts and hence have been shown to stabilize local and metastatic disease. Radiation can be used for spinal or sacral tumors.

Denosumab is an anti-RANKL antibody that can be used to stop the osteolytic process. It can be used in advanced stage 3 lesions or lesions in inoperable locations.

The given image shows the common locations of various bone tumors:



Orthopedics NEET 2022

Question 1:

What could be the diagnosis of a woman with this appearance on x-ray?

a) Multiple brown tumors

b) Fibrous dysplasia

c) Multiple enchondromas

d) Multiple exostoses

Question 2:

Which of the following is true about the type of fixation shown in the image?



a) Fracture tibia, Illizarow fixator

b) Fracture tibia, spanning fixator

c) Fracture femur, spanning fixator

d) Periarticular fracture of knee, spanning fixator

Question 3:

A male patient presented with a bone fracture following a road traffic accident. After 2 days he developed dyspnea, petechiae involving the whole body, and a fall in oxygen saturation. What is the likely diagnosis?

a) Fat embolism

b) Air embolism

c) Venous thromboembolism

d) Pulmonary hypertension

Question 4:

An intrauterine scan at the 13th week of pregnancy showed a fetus with multiple long bone fractures. What is commonly associated with this finding?

a) Achondroplasia

b) Osteogenesis imperfecta

c) Cretinism

d) Marfan syndrome

Question 5:

What is the most common complication of this condition, if left untreated?

a) Malunion and stiffness

b) Non union and cubitus varus

c) Cubitus valgus

d) Myositis ossificans

Question 6:

A child with a history of recurrent respiratory infections presented with complaints of pain around the knee and a high fever. X-ray shows lytic and sclerotic bone. His peripheral blood smear is shown in the image below. Aspirate from the joint will most likely show:



a) Staphylococcus aureus

b) Escherichia coli

c) Salmonella

d) Streptococcus

Question 7:

A boy falls on the left shoulder joint and presents to the emergency department with shoulder pain. His left elbow is flexed and is supported by the right hand. Which bone might be most likely fractured?



a) 1

b) 2

c) 3

d) 4

Answer Key

Question No.	Correct Option
1	c
2	d
3	a
4	b
5	c
6	c
7	a

Detailed Explanations

Solution to Question 1:

The diagnosis of the woman is multiple enchondromas. The X-ray shows multiple calcified lesions in the fingers.

Multiple enchondromatosis is also known as Ollier disease. It is a tumor in the metaphysis of long bones. There is a failure of bone formation and broadening of unossified cartilage within the bone. This results in shortening and angular deformity of the affected bones with an increased risk of pathological fractures. Enchondromas are also known as Maffucci syndrome if it is associated with overlying soft tissue hemangiomas. 25% of the patients with Ollier's disease will develop sarcomas by 40 years of age.

X-ray shows intralesional calcifications with a popcorn appearance. Histologically, enchondromas consist of hyaline cartilage with cytologically benign chondrocytes. Treatment of multiple enchondromas includes correction by osteotomy if the deformities are obvious.

Given below is an X-ray showing calcifications:

Enchondroma



Maffucci syndrome

Hemangiomas of the hand

Other options:

Option A: Multiple brown tumors are caused secondary to hyperparathyroidism and resemble a giant-cell tumor. It can be diagnosed by estimating the levels of calcium, alkaline phosphatase, and parathyroid hormone levels. The bone surrounding the lesion shows peritrabecular fibrosis.

The image below shows a brown tumor on the left hand:



Option B: Fibrous dysplasia is a condition characterized by the replacement of bone with fibrous tissue. The X-ray shows a ground-glass appearance with a well-defined sclerotic rim. It is seen in association with McCune-Albright syndrome. Shepherd's crook deformity in the proximal femur is characteristic of fibrous dysplasia.

The image below shows a child with McCune-Albright syndrome:

Option D: Multiple exostoses is an autosomal dominant type of osteochondroma that affects the phalanx, scaphoid, and metacarpals. The X-ray shows bony erosions and is excised.

The image below shows exostoses at the base of the thumb:

Solution to Question 2:

The image shows a periarticular fracture of the knee and a spanning external fixator used to stabilize the fracture.

External fixators are used if the fracture is associated with severe soft tissue damage. It is also used if internal fixation is risky due to wound contamination. The spanning external fixator will stabilize the fracture until the condition of soft tissue improves.

The technique of an external fixator is that the bone will be fixed above and below the fractures with screws and wires. These screws and wires are then connected to rigid bars. This method is used in fractures of the tibia, femur, pelvis, bones of the hand, etc.

In the image, the bones above and below the knee joint are fixed using a spanning external fixator and hence the fracture is likely to be a periarticular fracture of the knee.

The complications of external fixation include over-distraction, damage to soft tissue structures, and pin-track infections.

Another external fixator that is commonly used in an open type of fracture is the Ilizarov fixator. It is also used for the reconstruction of the limbs and for emergency stabilization in multiple long bone fractures.

Solution to Question 3:

The likely diagnosis is fat embolism.

Fat embolism is caused by the fat globules originating from the bone marrow following fracture of long bones. Clinical features of fat embolism are usually seen within 72 hours. They include tachycardia, fever, breathlessness, petechiae, confusion, and restlessness.

There is no specific test for diagnosis. However, urine analysis to look for fat globules and monitoring of blood pO2 are done. Management of fat embolism is mostly supportive, with oxygen therapy.

Other options:

Option B: Air embolism occurs when air bubbles entering the pulmonary circulation get absorbed in large quantities. Severe hypotension and hypoxemia are seen in this condition. The patient is placed in Trendelenburg's left lateral decubitus position.

Option C: Venous thromboembolism consists of pulmonary embolism (PE) and deep vein thrombosis (DVT). Bedridden patients and patients with chronic systemic diseases are more prone to thromboembolism. Clinical features of PE include dyspnea, syncope, cyanosis,

hypotension etc. DVT is presented as leg cramps, erythema, and tenderness. Heparin, LMWH, or fondaparinux are used for the treatment.

Option D: Pulmonary hypertension is a disease that involves remodeling of the pulmonary vasculature. It results in increased pulmonary artery pressure and vascular resistance. It mainly occurs as a complication of left heart disease, other primary lung diseases, and pulmonary embolism. The clinical features include dyspnea, edema, chest pain, syncope, etc.



Solution to Question 4:

Osteogenesis imperfecta is commonly associated with this multiple long bone fractures.

Osteogenesis imperfecta is a connective tissue disorder that occurs due to abnormal type 1 collagen production. The main features are joints that are vulnerable to fracture, scleral alterations, dentinogenesis imperfecta, and deafness. Multiple bone fractures are a consequence of bone fragility. It is classified into four types based on scleral involvement, natural history, and mode of inheritance.

Protrusio acetabuli, wormian bones, and proximal femoral ‘shepherd’s crook’ deformities of the femurs are the common radiological findings that are observed. Management comprises the use of bisphosphonates, along with physiotherapy and walking aids.

The images below show blue sclera and wormian bones in osteogenesis imperfecta:



Other options:

Option A: Achondroplasia is an autosomal dominant skeletal dysplasia with disproportionately short limbs, sleep apnea, and kyphoscoliosis. Facial features include button nose, frontal bossing, macrocephaly, etc.

The images below show the features of achondroplasia:

(a) Prominent forehead and body asymmetry, (b) a low nasal bridge, (c) trident hands, (d) scoliosis

Option C: Cretinism occurs due to iodine deficiency. Its clinical features are deaf-mutism, squint, mental retardation, and spastic or rigid neuromotor disorder.

Option D: Marfan syndrome is an autosomal dominant disorder. The affected persons will be tall with long limbs and digits. Spontaneous pneumothorax, aortic dissection, and lens dislocation are also seen. An early diastolic murmur is heard on auscultation.

Solution to Question 5:

Cubitus valgus is the most common complication, as the given condition is untreated lateral condyle of humerus fracture.

5–15 degrees of valgus is the normal carrying angle of the elbow. An angle more than this or an asymmetry between the two sides is called a valgus deformity. Long-standing non-union of a fractured lateral condyle is considered the most common cause.

Features of ulnar nerve palsy, such as the weakness of the hand, numbness or tingling of the ulnar fingers are also seen with cubitus valgus

Lateral condyle of humerus fracture is seen mostly in children up to 16 years of age. Falling on the hand with an extended elbow is the mechanism of injury. It is classified into types 1 and 2: the fracture is lateral to the trochlea in type 1 and through the middle, in type 2.

A back slab with the elbow flexed 90 degrees is used if there is no displacement. Reduction and internal fixation are needed for a displaced fracture.

Recurrent dislocations and malunion are the other complications. Since it is a Salter-Harris type IV injury, an imperfect reduction can result in growth arrest.

The image below shows cubitus valgus deformity:



Other options:

Option B: Cubitus varus deformity is also called gunstock deformity. It happens due to the malunion of supracondylar fracture in childhood. Injury to the brachial artery is an important complication.

Option D: Myositis ossificans is a tender swelling seen in the area of injury. It is often confused as a malignant tumor. Calcified lesions are seen in radiological studies. However, the swelling soon becomes less painful.

The X-ray image below shows myositis ossificans:



Solution to Question 6:

The given clinical scenario is suggestive of acute osteomyelitis. Aspirate from the joint will most likely show Salmonella.

The peripheral smear picture is indicative of sickle cell anemia. Patients with sickle cell anemia are prone to acute osteomyelitis caused by Salmonella.

Sickle cell anemia is a hematological disease caused by a mixture of genotypes. Its complication includes acute painful episodes, priapism, stroke, leg ulcers, acute chest syndrome (pneumonia-like illness with fever, cough, and hypoxia), etc.

Acute osteomyelitis refers to infection of the bone. It is mainly seen in children. Staphylococcus aureus is the most common organism causing osteomyelitis. However, in patients with sickle cell anemia, it is usually caused by Salmonella. Hemophilus influenza, E. coli, Streptococcus, and Pseudomonas aeruginosa can also cause the disease. The spread can be through hematogenous route or from exogenous routes such as open fractures, iatrogenic intervention, or spread from local infection.



Cardinal features of acute osteomyelitis in children include:

• Fever

• Pain

• Difficulty in weight-bearing

• Elevated WBC count

• Elevated CRP

• Elevated ESR

X-ray shows periosteal reaction and bony destruction after 10-12 days of infection. Management includes blood culture and sensitivity tests, followed by immediate empirical antibiotic therapy. Surgical drainage is done if there is an abscess or when the patient does not respond to antibiotics.

The complications of inadequately treated acute osteomyelitis include:

• Septicemia

• Epiphyseal damage and altered bone growth

• Chronic osteomyelitis

• Suppurative arthritis

• Pathological fracture

Solution to Question 7:

The clavicle marked as 1 might be most likely fractured.

One of the most frequently fractured bones in the body is the clavicle. Mostly, it occurs due to a direct blow or a fall on an outstretched arm. The midshaft of the clavicle is most commonly fractured. The lateral fragment is pulled down by the weight of the arm and the medial half is

pulled up by the sternomastoid muscle.



The patient will be presented with the flexed elbow as the arm will be held to the chest to prevent movement. It is divided into three groups, based on the location of the fractures.

• Group I – middle-third fractures

• Group II – lateral-third fractures

• Group III – medial-third fractures

Undisplaced fractures will be mostly treated non-operatively, with a simple sling for comfort. It can be used for 2-3 weeks and discarded once the pain subsides. Displaced fractures are treated with internal fixation either by contoured locking plates or intramedullary nails.

Complications include non-union, malunion, and stiffness of the shoulder.

Other options:

Option B: Scapula is marked as '2'. Fracture of the scapula occurs as a result of blunt trauma, falls, seizures, etc. The arm will be held immobile and bruising can be seen over the scapula and chest wall.

Option C: The shaft of the humerus is marked as '3'. Fracture of the shaft of the humerus occurs as a result of a fall on the hand, elbow, direct blow to the arm, or metastasis. The arm will be painful, bruised and swollen.

Option D: The 4th rib is marked as '4'. Rib fracture occurs due to the impact on the chest wall. Multiple rib fractures will result in the loss of structural integrity of the chest wall and cause flail chest. The lung cannot inflate as the flail segment is sucked in during inspiration.

Orthopedics NEET 2023

Question 1:

A child was brought with bony deformities as shown in the radiograph below. It could be due to the deficiency of which of the following?

a) Vitamin A

b) Vitamin D

c) Vitamin K

d) Vitamin C

Question 2:

In which of the following nerve injuries is the instrument shown below used?

a) Radial nerve

b) Median nerve

c) Ulnar nerve

d) Volkmann's ischemic contracture

Question 3:

A patient presented with chronic knee pain but has no history of trauma. Identify the condition shown in the radiograph below and the appropriate management.

a) Patellar avulsion fracture, TBW

b) Bipartite fracture patella, X-ray of other knee

c) Fracture of upper pole of patella, Cylindrical cast

d) Avulsion fracture , interfragmentary screw fixation

Question 4:

A patient at the orthopedics OPD complains of troubled sleep at night due to numbness and tingling sensation involving his lateral 3 digits. His symptoms are relieved as he lays his arms hanging from the bed. Which among the following options correctly describes his condition along with the test used to assess it?

a) Guyon's canal syndrome, Froment's test

b) Carpal tunnel syndrome, Froment's test

c) Guyon's canal syndrome, Durkan's test

d) Carpal tunnel syndrome, Durkan's test

Question 5:

A child is brought to the orthopedics OPD with a deformity in the lower limb and hyperpigmented skin lesions. The x-ray of her thigh is shown below. What is the most likely diagnosis?

a) Non-ossifying fibroma

b) Fibrous dysplasia

c) Paget's disease

d) Osteogenesis imperfecta

Question 6:

The given image is an x-ray of a 22-year-old female. What is the probable diagnosis?

a) Chondroblastoma

b) Osteochondroma

c) Giant cell tumor

d) Aneurysmal bone cyst

Answer Key

Question No.	Correct Option
1	b
2	a
3	b
4	d
5	b
6	c

Detailed Explanations

Solution to Question 1:

The given pediatric radiograph depicts the bowing of legs, which occurs due to rickets that results from the deficiency of vitamin D. Deficiency of vitamin D and calcium results in poor mineralization of bone, leading to rickets in children and osteomalacia in adults.

Rickets presents with skeletal changes like bowing of legs and bone pain once the child starts to walk. Radiological features show cupping at metaphyses, fraying, and widening of the growth plate. Prominent costochondral joints (rachitic rosary), craniotabes, frontal bossing, scoliosis, fractures, and dental anomalies may also be seen. The only cutaneous manifestation of vitamin D deficiency is alopecia.

Other options:

Option A: Vitamin A deficiency results in night blindness, xerophthalmia, and keratinization of the skin.

Option C: Vitamin K deficiency leads to hemorrhagic disease in newborns and elevated prothrombin time and bleeding. Vitamin K acts as a cofactor in the synthesis of coagulation proteins. Toxicity due to vitamin K results in hemolysis, hyperbilirubinemia, and kernicterus in the newborn.

Option D: Vitamin C deficiency leads to scurvy, which occurs due to defective collagen formation resulting in generalized bleeding tendencies and poor wound healing. Radiographic features of scurvy include the white line of Frankel, Trummerfeld zone, Pelkan spurs, and periosteal elevation. Tender nodules at the costochondral junctions, known as the scorbutic rosary, are characteristic.



Solution to Question 2:

The image shown above is that of a cock-up splint. It is used to manage radial nerve palsy.

Radial nerve palsy manifests as an inability to dorsiflex the wrist and digits (wrist drop and finger drop). Numbness occurs on the dorsoradial aspect of the hand and the dorsal aspect of the radial 3 1/2 digits.

Other options:

Option B: Median nerve injury happens in carpel tunnel syndrome where a nocturnal wrist splinting that helps to maintain the hand in a neutral position could be of help during the initial therapy to alleviate symptoms. It prevents prolonged flexion or extension of the wrist.

Option C: Ulnar nerve injury / ulnar claw hand could be managed with a knuckle bender splint if nerve recovery is anticipated. The splint can be used to correct and prevent metacarpophalangeal hyperextension.

Option D: Volkmann's ischaemic contracture (VIN) is most commonly seen following supracondylar humeral fractures. Mild cases of VIN are treated with a Volkmann's splint (Turnbuckle splint).

The first image below shows a knuckle-bender splint and the second image shows a Volkmann's splint

Solution to Question 3:

The given X-Ray shows a patella with a smooth discontinuity suggestive of an unfused accessory ossification center. Since this should arouse the suspicion of a bipartite patella, appropriate management would be to take an X-Ray of the other knee.

A bipartite patella (two-part patella) is a patella with an unfused accessory ossification centre, typically at the superolateral aspect. It usually is asymptomatic and is noted incidentally, however pain, when present, can be caused by overuse (such as in a cyclist or climber).

There are three types of bipartite patella

• Type I (5%) occurs at the inferior pole

• Type II (20%) occurs along the entire lateral border of the patella

• Type III (75%) the most common type, occurs as an elliptical area in the superolateral portion of the patella

Depending on the number of fragments they may be classified as bipartite, tripartite or multipartite. For diagnosis, a conventional skyline view is followed by a skyline view with the patient in a squatting weight-bearing position to assess if the pain is caused by a nonunion at the bipartite site. If the gap is higher in the squatting weight-bearing posture than in the standard skyline view, the test result is considered positive. MRI is not required to diagnose the condition.

If pain occur with a bipartite patella, limiting and restricting activity, correcting the activity that is causing an overuse syndrome, and prescribing nonsteroidal antiinflammatory agents can result in improvement in 3 weeks. Operative treatment with the Ogata technique is rarely necessary.

Solution to Question 4:

The given clinical vignette is suggestive of carpal tunnel syndrome and it can be assessed by Durkan's test.

Durkan's test is the most specific test for the diagnosis of carpal tunnel syndrome. In Durkan's test, direct compression of the median nerve is done for about 30 seconds, which elicits paresthesias and pain in cases of carpal tunnel syndrome.

Carpal tunnel syndrome is the most common entrapment mononeuropathy caused by compression of the median nerve as it passes through the fibro-osseous carpal tunnel beneath the flexor retinaculum. Patients present with pain and paresthesia over the radial 3 and a half digits. There may be a weakness of thumb opposition. In chronic cases, the thenar eminence is wasted. Carpal tunnel syndrome can be associated with rheumatoid arthritis, pregnancy, chronic renal failure, acromegaly, hypothyroidism, and gout. The diagnosis is usually made on clinical grounds but nerve conduction studies can confirm the diagnosis.

Other tests for CTS -

• Phalen's test - Elicited by holding the wrist flexed fully. The appearance of sensory symptoms within 60 seconds implies a positive test.

• Tinel's sign - Sensory symptoms reproduced by percussing over the median nerve imply a positive test.

Guyon’s canal is a fibro-osseous canal, approximately 4 cm long in the adult, on the anteromedial aspect of the wrist under the superficial part of the flexor retinaculum. The ulnar nerve passes in Guyon’s canal along with the ulnar artery and can be compressed here. Froment's test is used in a suspected ulnar nerve injury.

In Froment's test or Book Test, the patient is asked to grasp a book or a sheet of paper between the thumb and index finger. Due to weakness of the adductor pollicis muscle, adduction of the thumb is lost. When asked to hold a book, the patient tries to hold the book by using the flexor pollicis longus muscle instead of the adductor, this produces flexion at the interphalangeal joint.

The picture below shows Froment's test





Non-surgical Management of Carpal Tunnel Syndrome: In mild to moderate cases of carpal tunnel syndrome, non-surgical management options can be explored to alleviate symptoms and improve function. These conservative treatments include:

•

Wrist splinting: Wearing a wrist splint at night can help maintain the wrist in a neutral position, reducing pressure on the median nerve and providing relief from nocturnal symptoms.

•

Activity modification: Identifying and modifying activities that exacerbate symptoms can help in managing carpal tunnel syndrome. For example, taking frequent breaks during repetitive tasks or adjusting the workstation ergonomics can be beneficial.

•

Nonsteroidal anti-inflammatory drugs (NSAIDs): Over-the-counter NSAIDs, such as ibuprofen, can help manage pain and inflammation associated with carpal tunnel syndrome.

•

Corticosteroid injections: In some cases, corticosteroid injections into the carpal tunnel may provide temporary relief from symptoms by reducing inflammation and pressure on the median nerve.

•

Physical and occupational therapy: Therapists can help patients learn exercises to stretch and strengthen the hand and wrist muscles, improving symptoms and function.

If conservative treatments do not provide sufficient relief or the condition worsens, surgical intervention, such as carpal tunnel release surgery, may be considered.

Solution to Question 5:

The above X-ray shows the bending of the femur known as Shepherd's crook deformity. It is seen in fibrous dysplasia of proximal femur.

Fibrous dysplasia is a condition that affects bone development and is characterized by the replacement of the bone with fibrous tissue. The patient presents with a progressive limb deformity or a fracture through a previously asymptomatic lesion. The presentation often occurs in the second decade. Other symptoms include nonspecific bone pain, tenderness, and swelling in the affected area. The commonly involved sites are the proximal femur, tibia, pelvis, and foot. Polyostotic fibrous dysplasia is associated with Albright-McCune syndrome.

Radiological findings include:

• Well-defined, expansile intramedullary lesions

• Hazy ground glass appearance is seen in the majority of cases

• Shepherd's crook deformity - Expansion with marked varus in the proximal femur

Other options:

Option A: Non-ossifying fibroma is the commonest variant of giant cell bone tumors. It is a benign, non-neoplastic lesion most commonly seen in the metaphyses of the long bones in children. Non-ossifying fibromas are usually asymptomatic, and often discovered incidentally on X-rays performed for other reasons. They commonly occur in the metaphyses of long bones and are characterized by well-defined, eccentric, cortically-based lytic lesions. The X-ray findings in the question do not match this description.

Option C: Paget's disease of the bone is a condition where there is an increased bone turnover followed by enlargement and thickening of the bone. In this condition, calcium levels are normal with raised ALP. Paget's disease typically presents in older adults and affects the axial skeleton, including the pelvis, spine, skull, and femur. It is characterized by increased bone turnover, leading to disorganized bone remodeling, resulting in enlarged and deformed bones. The X-ray findings and the patient's age do not match the characteristics of Paget's disease.

Option D: Osteogenesis imperfecta is a genetic disorder characterized by increased bone fragility and susceptibility to fractures. It is associated with blue sclera, dental fragility, and hearing loss. It presents with the hallmark features of osteoporosis with increased fragility of bones along with blue sclera, dental fragility as well as hearing loss. The X-ray findings in the question do not show the characteristic features of osteogenesis imperfecta, such as thin cortices, multiple fractures, or wormian bones.

Solution to Question 6:

The given x-ray showing a lytic lesion at the end of a long bone with a soap-bubble appearance suggests a giant cell tumor (GCT) or osteoclastoma.

Osteoclastoma usually affects young adults of age 20 to 40 years with a slight female predominance. It is a benign but locally aggressive tumor. It is exclusively seen in adults and affects the ends of long bones, involving metaphysis/ epiphysis of long bones. GCT is most commonly seen in the distal femur, proximal tibia, proximal humerus, and distal radius. Common presenting complaints are swelling and vague pain. Sometimes, the patient presents for the first time with a pathological fracture through the lesion.

Important x-ray features of a GCT:

• A solitary, possibly loculated, lytic lesion in the epiphysis

• Eccentric location, most often subchondral

• Expansion of the overlying cortex

• Soap bubble appearance - The tumor is homogeneously lytic with trabeculae of the remnants of bone traversing it, giving it a loculated appearance called soap-bubble appearance.

• No calcification within the tumor

• No or minimal reactive sclerosis around the tumor

Histologically, GCTs consist of undifferentiated spindle cells profusely interspersed with multi-nucleated giant cells. The tumor stroma is well-vascularized with bands of cellular or collagenous fibrous tissue. The image below shows the histopathology of osteoclastoma.



The treatment of choice for GCT is extended curettage and bone grafting.

• GCT in the relatively insignificant bone like the distal ulna, clavicle, and proximal fibula is treated by resection without reconstruction.

• Bisphosphonates have been shown to reduce the proliferation of osteoclasts and hence have been shown to stabilize local and metastatic disease.

• Radiation can be used for spinal or sacral tumors.

Other options:

Option A: Chondroblastoma is a benign tumor that usually appears as well-circumscribed round or oval lytic lesions in the center of the epiphysis of a long bone. It presents as a painful mass that causes effusion or stiffness. It is most commonly seen in ages 10-20 years.

The following is an x-ray image of the knee showing chondroblastoma:

Option B: Osteochondromas are common benign bone tumors commonly present in adolescents. The lesions consist of a bony mass, often in the form of a stalk, produced by endochondral ossification of a growing cartilaginous cap. They are usually painless. The common sites of involvement are the distal femur, proximal tibia, and proximal humerus.

The following is an x-ray image of the knee showing osteochondroma of the distal knee:



Option D: Aneurysmal bone cyst is a tumor that affects all age groups but generally occurs more commonly in adolescents with an equal male: female distribution. It usually occurs in the metaphysis of long bones (like femur and tibia) and the posterior elements of vertebral bodies. Patients usually present with pain and swelling. Radiographically, ABC is seen as an eccentric, expansile, lytic lesion with well-defined margins.

The following is an x-ray image of the knee showing ABC:

The image below shows common locations of various bone tumors:


Location of Bone Tumors within the Bone
Diaphysis
Adamantinoma
Osteofibrous Dysplasia
Osteoid Osteoma
Stress Fracture
Chronic Osteomyelitis
Metaphysis
Fibrous Cortical defect
Non ossifying /fibroma
Osteochondroma
Osteosarcoma
Epiphysis
Articular
Osteochondroma
(Trevor's Disease)
Round Cell Tumors
Langerhans
Cell Histiocytosis
Fibrous Dysplasia
Fibrosarcoma
Chondromyxoid Fibroma
Aneurysmal Bone Cyst
Enchondroma/
Chondrosarcoma
Simple Bone Cyst
Fibrous Dysplasia
Pyogenic Osteomyelitis
Chondroblastoma(child)
Giant Cell tumor(adult)
Fungal/ Tubercular
Osteomyelitis
Aneurysmal Bone Cyst
©MARROW

Orthopedics INI-CET 2023

Question 1:

A 25-year-old male who sustained a shoulder injury presented with swelling around the shoulder joint. An X-ray of the shoulder joint is shown below. Which is/are the ligament(s) likely to have been injured?

a) Both acromioclavicular and coracoclavicular ligament.

b) Coracohumeral ligament

c) Coracoacromial ligament

d) Acromioclavicular ligament

Question 2:

A 14-year-old male presented with a mushroom-like tumor in the distal femur for the past 2 years. Which of the following features will help the clinician to identify a malignant transformation?



a) Initially growth of the tumor was away from the joint, now the growth of tumor is towards the joint.

b) Tumor does not have malignant transformation potential.

c) Tumor is in continuation with bone marrow cavity of the normal bone.

d) Hyaline cap thickness on MRI is 2cms.

Question 3:

A 16-year-old girl presents with multiple swellings and multiple brown rashes. She has increased uptake on bone scans over the femur, skull, and ribs. Biochemical parameters are abnormal. She also has a history of hypothyroidism. An X-ray of her femur is given below. Which of the following is the likely diagnosis?

a) Langerhans cell histiocytosis

b) McCune-Albright syndrome

c) Neuroma with bone involvement

d) Papillary carcinoma of thyroid

Question 4:

A patient came to the emergency department with a history of road traffic accident and a shaft of femur fracture. The patient was stabilized. After 12 hours, he developed sudden onset breathlessness, confusion, and petechial skin rashes over the chest and axilla. What is the probable diagnosis?

a) Fat embolism

b) Head injury

c) Blunt trauma to the chest

d) Hemorrhagic shock

Question 5:

Choose the true statements:

a) ii, iii and iv only

b) i, iii only

c) iii, iv only

d) i, ii only

Question 6:

What type of gait will be seen following a fracture in this region?



a) Waddling

b) High stepping

c) Wide-stepping gait

d) Stamping

Question 7:

True statements about bankart lesion?

a) Avulsion of the anteroinferior glenoid labrum

b) Avulsion of the anterosuperior glenoid labrum

c) Depression in posteroinferior part of the head of the humerus

d) Depression in posterosuperior part of the head of the humerus

Question 8:

Hawkin's classification is used for which fractures?

a) Talus

b) Navicular

c) Tibia

d) Calcaneum

Question 9:

A female patient presented with swollen and tender joints. There is symmetric involvement of multiple joints such as wrists, hands, and feet, which is bilateral, along with morning stiffness lasting more than 1 hour. Erythema is absent. What is the most probable diagnosis?

a) Gout

b) Rheumatoid arthritis

c) Osteoarthritis

d) Psoriatic arthritis

Question 10:

A 45-year-old patient with a tibial fracture in a cast presented to the emergency department with complaints of intense pain. The cast was removed and on examination, passive extension was painful. The dorsalis pedis and posterior tibial pulses were present and there was a loss of sensation over the first web space. What is the next step?

a) Duplex imaging

b) Re-apply cast

c) Measure anterior compartment pressure

d) Analgesics

Question 11:

Three-point bony relationship of the elbow will not be damaged in?

a) Supracondlyar fracture humerus

b) Intercondylar fracture humerus

c) Elbow dislocation

d) Lateral epicondyle fracture

Answer Key

Question No.	Correct Option
1	a

2	d
3	b
4	a
5	a
6	b
7	a
8	a
9	b
10	c
11	a

Detailed Explanations

Solution to Question 1:

The given image shows that the clavicle has moved up and there is a gap between the clavicle and the acromion suggesting injury to both acromioclavicular and coracoclavicular ligament.

The acromioclavicular joint is a synovial plane type between the acromial end of the clavicle and the medial acromial margin. The two ligaments providing support are acromioclavicular(intrinsic) and coracoclavicular(extrinsic). The acromioclavicular ligament has a superior part that extends between the upper aspect of the clavicle's lateral end and the adjacent acromion. Its inferior part, extends between the lower surface of the clavicle's lateral end and the adjoining acromion. Additionally, it serves as an attachment site for the intra-articular disc.

The coracoclavicular ligament connects the clavicle with the coracoid process of the scapula. Although it is separate from the acromioclavicular joint, it plays a crucial role in maintaining the alignment of the acromion and the clavicle. Therefore, it contributes significantly to the suspension mechanism of the scapula. The ligament has two parts - trapezoid and conoid parts.

The primary cause of injury is direct impact to the lateral aspect of the shoulder or the acromion process while the arm is in an adducted position. It can also be caused by falling onto an outstretched hand or elbow.

Standard X-rays are sufficient for diagnosing acromioclavicular joint injuries and should be utilized to assess other potential sources of shoulder pain resulting from trauma. By comparing the coracoclavicular interspace distance with the contralateral view, the injury can be identified.



Solution to Question 2:

The given X-ray shows a well-defined exostosis pointing to the diagnosis of osteochondroma. 2cms thickness of hyaline cap thickness on MRI is suggestive of malignant transformation.

Osteochondroma can turn into low-grade chondrosarcoma but the malignant transformation is extremely rare(Option B). It should be suspected in the following cases:

• Growth after skeletal maturity

• Lucency in the bone

• Additional scintigraphic activity

• Cortical destruction

• Pain after puberty

• Soft tissue mass

• Thickened cartilage cap >2 cm

Osteochondromas are common benign bone tumors mostly arising from the metaphysis of long bones. The cortical and medullary continuity of the tumor with the underlying bone is a pathognomonic feature that establishes the diagnosis. (Option C). The lesions consist of a bony mass, often in the form of a stalk, produced by endochondral ossification of a growing cartilaginous cap. Most of these lesions are asymptomatic and are discovered incidentally.

These are of two types: Pedunculated and sessile, pedunculated types are more common. The definite stalk is directed away from the physis(Option A). The lesions are usually covered with a cartilaginous cap which cannot be seen radiologically and hence the lesions are larger clinically than what their X-Ray appearance suggests. The important feature is that the host bone flares from the cortex into the osteochondroma.

Most patients are asymptomatic and never seek medical attention. However, symptomatic people may undergo resection.

Solution to Question 3:

The above X-ray shows the bending of the femur known as Shepherd's crook deformity. It is seen in fibrous dysplasia of the proximal femur. The diagnosis based on the x-ray and clinical scenario points to McCune-Albright syndrome.

McCune Albright syndrome presents as a triad of cafe-au-lait spots (cutaneous pigmentation), polyostotic fibrous dysplasia, and precocious puberty. It is also associated with multiple endocrinopathies like hyperthyroidism, acromegaly, etc. The locus of involvement is the GNAS1 gene.

Cutaneous pigmentation is more prominent on the side with severe bone involvement. Fibrous dysplasia is a condition that affects bone development and is characterized by the replacement of the bone with fibrous tissue. The patient presents with a progressive limb deformity or a fracture through a previously asymptomatic lesion. The presentation often occurs in the second decade. Other symptoms include nonspecific bone pain, tenderness, and swelling in the affected area. The commonly involved sites are the proximal femur, tibia, pelvis, and foot.

Radiological findings include:

• Well-defined, expansile intramedullary lesions

• Hazy ground glass appearance is seen in the majority of cases

• Shepherd's crook deformity - expansion with marked varus in the proximal femur.

Other options:

Option A: Langerhans cell histiocytosis (LCH) occurs due to the proliferation of a special type of immature dendritic cells, called Langerhans cells. Their normal function is to present antigens. The most common mutation is the BRAFV600E gain-of-function mutation.

It is characterized by single or multiple osteolytic bone lesions and can also infiltrate nearly every organ, like the skin (of the scalp), lungs, liver, spleen, bone marrow, or the central nervous system. It presents with pain and swelling in the affected area. Spinal involvement results in compression fractures and neurological involvement. Vertebral involvement results in a rapid flattening of the vertebral body (vertebra plana).

LCH can present as the following clinical entities:

• Multifocal multisystem Langerhans cell histiocytosis (Letterer-Siwe disease) which is mostly seen in infants

• Unifocal and multifocal system Langerhans cell histiocytosis (eosinophilic granuloma) - The combination of bone defects, diabetes insipidus, and exophthalmos is called the Hand-Schüller-Christian triad that is mostly seen in younger children

• Pulmonary LCH is mostly seen in adults.

Option D: Papillary carcinoma of the thyroid (PTC) presents as a painless nodule or mass in the neck or enlargement of cervical lymph nodes. It usually occurs in women between 25 to 50 years of age. PTC is the most common thyroid cancer associated with previous exposure to ionizing radiation. Conventional PTCs have translocations involving RET or NTRK, and point mutations involving the BRAF gene.

Patients present with painless swelling. Initial presentation may be with cervical lymphadenopathy due to lymph node metastases.

Solution to Question 4:

The most likely diagnosis based on the given clinical scenario is fat embolism.

Fat embolism is caused by the fat globules originating from the bone marrow following the fracture of long bones. Clinical features of fat embolism are usually seen within 72 hours. They include tachycardia, fever, breathlessness, petechiae, confusion, and restlessness. Gurd's criteria is used for the diagnosis of fat embolism syndrome:

There is no specific test for diagnosis. However, urine analysis to look for fat globules and monitoring of blood pO2 are done. Management of fat embolism is mostly supportive, with oxygen therapy.

Other options:

Option C: A majority of chest trauma is caused by blunt injuries. Six life-threatening conditions such as airway obstruction, tension pneumothorax, open pneumothorax, massive hemothorax, flail chest, and pericardial tamponade should be immediately investigated and treated. Only 10% or less of blunt chest trauma patients require surgical treatment, and the remaining patients can be treated conservatively.

Option D: Hemorrhagic shock is the most common cause of shock in surgical or trauma patients. Here the patient presents with agitation, cold clammy extremities, hypotension, tachycardia, and weak peripheral pulses. Acidosis and elevated lactate levels may be seen due to hypoperfusion of tissues.

Solution to Question 5:

Statements ii, iii, and iv are true, and statement i is incorrect.

Statement i: The most commonly fractured carpal bone is the lunate is a false statement. The most common fractured bone is the scaphoid.

Statement ii: The scaphoid is most commonly fractured at the waist is a true statement. A scaphoid fracture is more common in young adults. Commonly, the fracture occurs through the waist of the scaphoid.

Statement iii: The cast for scaphoid fracture is set in a glass holding position is a true statement. The affected hand is immobilized in a scaphoid cast for 3-4 months. The scaphoid cast extends from below the elbow to the metacarpal heads including the thumb up to the inter-phalangeal joint. The wrist is maintained in a little dorsiflexion and radial deviation (glass holding position).

Statement iv: Kienböck's disease is due to osteonecrosis of the lunate is a true statement. Kienböck's disease is usually seen in young adults, who present with pain and stiffness in the wrist. A history of trauma may be present. On examination, localized tenderness can be appreciated over the lunate, and a reduction in grip strength can be noted. Limited and painful wrist movements are noted in the later stages of the disease.

Solution to Question 6:

In the above scenario, the structure marked in the given image is the neck of the fibula. Injury at this site causes failure of dorsiflexion (foot drop) due to injury to the common fibular nerve (also known as common peroneal nerve). Foot drop leads to high stepping gait.

The common peroneal nerve supplies the extensors and evertors of the foot. Injury to the common peroneal nerve results in foot drop. So the patient attempts to lift the leg high enough during walking so that the foot does not drag on the floor. This is known as a high-stepping gait.

The common fibular is derived from the dorsal branches of the L4 - L5 and S1 - S2 ventral rami. It innervates the short head of the biceps femoris in the posterior compartment of the thigh and then continues into the lateral and anterior compartments of the leg and onto the foot.

It gives the following branches:

• Sural communicating nerve (cutaneous branch)

• Lateral sural cutaneous nerve (cutaneous branch)

• Superficial fibular nerve

• Deep fibular nerve.

Injury to the common peroneal nerve will lead to:

• Weakness of dorsiflexion of foot or foot drop

• Loss of dorsiflexion

• Loss of sensation over the lateral aspect of the leg and dorsum of the foot.

• Wasting of fibular and anterior tibial muscles.

Other Options:

Option A: Waddling gait is seen in developmental dysplasia of the hip and in proximal muscle weakness.

Option C: Wide-stepping gait is a broad-based unstable gait and is indicative of cerebellar dysfunction.

Option D: Stamping of the foot is seen in tarsal tunnel syndrome. Tarsal tunnel syndrome is a condition with pain and sensory disturbance occurring in the medial part of the forefoot, unrelated to weight-bearing. This is due to compression of the posterior tibial nerve behind and below the medial malleolus. Typically the pain worsens at night and gets relieved on walking or stamping the foot.

Solution to Question 7:

Bankart lesion is an avulsion in the anteroinferior part of the glenoid labrum caused due to the recurrent anterior dislocation of the humeral head.

Hill-Sachs lesion is a compression fracture on the posterolateral aspect of the humeral head. It is caused by the pressure of the anterior glenoid rim on the humeral head in recurrent dislocations.

Anterior dislocation is the commonest shoulder dislocation and can be caused due to trauma from behind the shoulder pushing the head out anteriorly. It is also caused due to falling on an

outstretched hand in an abducted and externally rotated shoulder. It can be diagnosed on clinical examination with flattening of the shoulder contour and ability to place a rule (Hamilton ruler test) and the inability to touch his/her opposite shoulder (Duga's test). X-ray confirms the diagnosis and may show associated Hill Sachs or bony bankart lesion. The most common complication is recurrent shoulder dislocation.

The following X-ray shows the Bankart lesion and Hill-Sachs lesion:



Solution to Question 8:

Hawkin's classification is used for the fracture of the neck of the talus.

The fracture of the neck of the talus most commonly occur following a high-energy injury causing forced dorsiflexion with axial load. They can be further classified using Hawkin's classification:

• Type I - Undisplaced

• Type II - Subtalar dislocation

• Type III - Subtalar and tibiotalar dislocation

• Type IV - Subtalar, tibiotalar, and talonavicular dislocation

Undisplaced fractures are managed conservatively in a short leg cast for 8-12 weeks and a non-weight bearing cast for 6 weeks. Hawkin's type II-IV are managed by open reduction with internal fixation (ORIF).

The below image shows the Hawkins classification of talar neck fractures.

Solution to Question 9:

In the above clinical scenario, a female patient presenting with swollen and tender joints, symmetric involvement of multiple joints such as wrists, hands, and feet, bilateral in nature, along with morning stiffness lasting for more than 1 hour, and the absence of erythema is rheumatoid arthritis.

Rheumatoid arthritis (RA) is a chronic inflammatory disease of symmetric polyarthritis, mainly affecting the small joints of the hands and feet. Its incidence increases between 25 and 55 years of age. Clinically, patients present with early morning joint pain and stiffness lasting >1 hour that relieves with activity.

Signs and symptoms -

• Swelling and pain of the small joints of the hands initially- metacarpophalangeal (MCP) and proximal interphalangeal (PIP) joints are commonly involved. Distal interphalangeal (DIP) joint involvement is usually due to coexistent osteoarthritis.

• Morning stiffness

• Swan-neck deformity - Hyperextension of the PIP joint with flexion of the DIP joint

• Boutonnière deformity - Flexion of the PIP joint with hyperextension of the DIP joint,

• Z-line deformity - Subluxation of the first MCP joint with hyperextension of the first interphalangeal (IP) joint

• Ulnar deviation of fingers

• Constitutional findings such as fever, weight loss, etc

• Monoarthritis of large joints can occur later

• Tenosynovitis of palm(trigger finger)

• Atlantoaxial joint involvement

Joints commonly affected in Rheumatoid Arthritis: (bilateral and symmetrically involved)

• MCP joints

• Wrists

• PIP joints

• MTP joints

• Atlantoaxial joint

Laboratory investigations - Detection of Rheumatoid factor and anti-CCP antibodies help in differentiating this condition from other forms of arthritis.

Diagnosis is made with clinical signs and symptoms with corroboration from lab and imaging findings.

Other options-

Option (A): Gout is characterized by episodic acute arthritis or chronic arthritis due to the deposition of monosodium urate(MSU) crystals in the joints and connective tissue. It commonly affects postmenopausal women and middle-aged to elderly men. The metatarsophalangeal joint (MTP) of the first toe is commonly involved. Other joints that are commonly affected include the tarsal joints, ankle, and knee.

Option (C): Osteoarthritis is the most common form of arthritis with a high prevalence in the elderly population. Commonly involved joints are the knee, hip, first MTP, and the cervical and lumbosacral spine. In the hands, the DIP and PIP joints and the base of the thumb are commonly involved. The wrist, elbow, and ankle are usually spared.

Option (D): Psoriatic arthritis- In most cases psoriasis precedes arthritis but in some cases, both of them present in a span of 1 year, or arthritis precedes psoriasis. There can be involvement of DIP and PIP joints and the joint involvement tends to follow "ray "distribution. Ray distribution refers to the involvement of all the joints of one finger while the adjacent finger is totally spared. Nail changes in the affected fingers are a differentiating feature of this condition.

Solution to Question 10:

The clinical presentation of this patient with a history of tibial fracture complaining of intense pain, with painful passive extension and loss of sensation of the first web space upon the removal of the cast, is suggestive of acute compartment syndrome. The next step would be to measure the anterior compartment pressure to confirm the diagnosis.

Acute compartment syndrome refers to the rise in intra-compartmental pressure leading to tissue ischemia and necrosis. The injuries at high risk of developing acute compartment syndrome are fractures of the proximal one-third of the tibia, soft tissue injury, distal radius fracture, and diaphyseal fracture of the forearm. Other high-risk injuries are crush injuries and circumferential burns.

The classic features of ischemia in acute compartment syndrome include the 5 Ps:

• Pain: is the first symptom of acute compartment syndrome. On passive stretching of the muscles, severe pain can be elicited.

• Paresthesia: paresthesia and hypoesthesia are the first signs of nerve ischemia.

• Paralysis: paralysis of muscle groups affected in that compartment is a late sign.

• Pulselessness: usually, the peripheral pulses and capillary return are intact in the early stages, unless there is a major arterial injury. If pulses are absent, arteriography is indicated.

• Pallor: may not always be seen as ischemia occurs at the capillary level.

The diagnosis can be confirmed by measuring the intra-compartmental pressure using a split catheter introduced into the compartment. The difference between diastolic pressure and compartment pressure is known as differential pressure (ΔP). If the differential pressure (ΔP) is less than 30 mm Hg, immediate compartment decompression is indicated.

Treatment:

• Prompt decompression of the affected compartment is done.

• All the dressings, bandages, and cast on the affected limb are removed and the limb is nursed flat.

• Fasciotomy is done if the differential pressure is less than 30 mm Hg

• Fasciotomy wounds are left open and observed after 2 days. If there is necrosis, debridement is done. If the wounds are healthy, they can be sutured without tension or skin grafting can be done.

Solution to Question 11:

The three-point bony relationship will not be damaged in supracondylar fracture.

Supracondylar fracture of the humerus is most commonly seen in children of 5-7 years after a fall on an outstretched hand with the elbow in full extension. It is most commonly of extension type followed by flexion type. Gartland classification is used for supracondylar fractures. The three-point bony relationship is maintained in supracondylar fracture (option B). Three-point bony landmarks in the elbow are the tips of medial and lateral epicondyles and the olecranon.

Nerve and vessel injury:

• Most common nerve to be injured in extension type - anterior interosseous nerve > Median nerve > Radial nerve > Ulnar nerve. (AMRU)

• Most common nerve to be injured in flexion type - Ulnar nerve.

• Most common vessel to be injured - brachial artery.

Radiological findings:

• An undisplaced fracture may show a triangular lucency like a sail of a yacht (sail sign), also called a fat pad sign. This is due to the fat pad around the elbow being pushed forward by the hematoma.

• Baumann's angle (humeral-capitellar angle) can help in assessing the degree of medial angulation before and after reduction.

The most common complication of supracondylar fracture is malunion. This leads to cubitus varus or gunstock deformity. Malunion occurs due to improper reduction and not due to physeal damage, as the fracture is extraphyseal (option D). Other complications include myositis

ossificans, compartment syndrome, and Volkmann ischemic contracture.

Undisplaced supracondylar humerus fractures are treated by an above-elbow plaster slab with the elbow in 90° flexion. Displaced fractures are treated by closed reduction and K-wire fixation since they tend to slip after reduction. Open fractures or those associated with brachial artery injury are treated by open reduction and internal fixation.

Cubitus varus deformity is corrected after skeletal maturation and lateral closed-wedge osteotomy (French osteotomy) is the method employed.

Orthopedics INI-CET 2024

Question 1:

Which of the following layers is affected in Salter Harris type 1 fracture?

a) Through the whole of epiphysis

b) Resting zone of the growth plate

c) Calcific layer of the growth plate

d) Mineralisation zone

Question 2:

A 75 year old man presents with fracture of intracapsular neck of femur. What is the "standard of care" management done for this patient?

a) Boot and bar cast

b) Russell's traction

c) Intramedullary nailing

d) Hemiarthroplasty

Question 3:

What is the diagnosis of the following X Ray?

a) Tibial condylar fracture

b) Rheumatoid arthritis

c) Patellar fracture

d) Hemarthrosis

Question 4:

A 22 year old came with thigh pain. X ray was done as shown below. What is the diagnosis?

a) Osteosarcoma

b) Osteomalacia

c) Chronic osteomyelitis

d) Ewings sarcoma

Question 5:

Out of the following, which is the most common site of osteoporotic fracture?

a) Vertebral fracture

b) Glenoid fracture

c) Subtrochanteric fracture

d) Mid shaft of humerus

Question 6:

A 70 yr old female presents with extracapsular neck of femur fracture. What is the best step in management?

a) Russell's traction

b) Interlocking plate

c) Dynamic hip screw

d) Boot and bar cast

Question 7:

A 26 year old female presented with proximal one-third fracture of shaft of femur. What is the treatment of choice ?

a) Intramedullary nailing

b) Above knee slab

c) Hip spica

d) Above knee cast

Question 8:

All of the following statements regarding ankylosing spondylitis are true except?

a) Most commonly seen between the age of 20 to 40.

b) Affects males more than females

c) In 85-95% patients, HLA-B27 is positive

d) Non Erosive arthritis of the joint with sclerosis is seen

Question 9:

A 60-year-old female comes with complaint of unable to bear weight following trivial trauma. X ray of hip and leg is given. What is the diagnosis?

a) Brown tumor

b) Disseminated TB

c) Eosinophilic granulomas

d) Metastasis

Question 10:

A 40-year-old male comes with an intracapsular fracture of the femur. What is the surgery of choice?

a) Open reduction internal fixation

b) Closed reduction internal fixation

c) Hemiarthroplasty

d) Total arthroplasty

Question 11:

FRAX is related to?

a) Fracture risk assessment

b) Fracture rehabilitation assessment

c) Fracture repair assessment

d) Fracture reduction assessment

Answer Key

Question No.	Correct Option
1	c
2	d
3	c
4	c
5	a
6	c
7	a
8	d
9	d
10	b
11	a

Detailed Explanations

Solution to Question 1:

Salter Harris type 1 fracture is a transverse fracture through the hypertrophic or calci■ed zone of the growth plate. The growing zone of the physis is usually not injured and growth disturbance is uncommon.

Solution to Question 2:

In patients over 65 years old with an intracapsular neck of femur fracture and a normal hip otherwise, hemiarthroplasty is the standard management approach.

Reduction and ■xation of fractures in the elderly patient is not recommended due to the associated high rates of subsequent reoperation , non-union and osteonecrosis of the femoral head. In young patients presenting with fracture within 3 weeks, closed reduction internal fixation with cannulated cancellous screws is performed.

Bipolar hip prosthesis is preferred as it has 2 articulating sites. Austin Moore and Thompson prosthesis are also used. If the patient has pre exixting hip conditions like osteoarthritis, total hip replacement is preferred.

Solution to Question 3:

The X-ray shows a transverse fracture of the patella.

The patient presents with pain and swelling over the affected knee. Patellar tenderness and bruise over the front of the joint can be present. Crepitus may be felt in comminuted fractures. The patient cannot lift his leg in full extension(extension lag) due to the disruption of the extensor apparatus. Hemarthrosis can be associated.

Patellar fractures can be transverse, longitudinal, polar, or comminuted.

The radiological examination includes AP and lateral views. A skyline view of the patella is required in some cases of undisplaced fractures.

Management of patellar fractures:

• Undisplaced fractures: It is treated with a plaster cast extending from the groin to just above the malleoli, with the knee in full extension. This is called a cylinder cast and is usually left in place for a period of at least 3 weeks

• A transverse fracture is treated by a surgical procedure consisting of the reduction of the fragments known as tension-band wiring and K- wire fixation. This is followed by the repair of the extensor retinacula and physiotherapy.

• Comminuted fractures are treated by open reduction and fixation. Patellectomy is the preferred option in this case as restoration of a smooth articular surface is difficult in this condition. Patella-saving operations are recently more performed with advancements in fixation techniques.

• Longitudinal fractures usually have intact extensor apparatus. They can be treated with a cylinder cast or extension brace initially, followed by a range of motion.

Solution to Question 4:

The given clinical scenario of a patient with thigh pain and an X-ray of the lower limb revealing involucrum and cloaca in the femur, strongly suggests a diagnosis of chronic osteomyelitis.

Chronic osteomyelitis may occur by direct inoculation of the infective organism during trauma (like road traffic accidents). The infection continues to thrive within the bone leading to alternating quiescent and flare-up stages of chronic osteomyelitis.

Sequestrum is a piece of dead bone. It is formed when a fragment of bone gets separated from the healthy bone due to necrosis. It varies in size from a spicule to a large fragment and acts as a nidus for infection. It leads to abscess and persistent pus discharge in the bone. Sinus tracts are formed when the abscess from the infected bone drains out through the skin.

The involucrum is the dense new sclerotic bone surrounding the sequestrum. It is formed from deep layers of stripped periosteum. Cloacae are the perforations in the involucrum. Pus and tiny spicules of bone may be discharged through them.

The radiograph below shows sequestrum, involucrum and cloacae:

Solution to Question 5:

Out of the following options, vertebral fractures are most commonly seen in osteoporosis

The fractures are classically low trauma de■ned as a fall from standing height or less. Fracture ofthe distal radius (Colles' fracture) is usually the ■rst fracture to occur, followed by vertebrae and hip unless treatment is initiated. Osteoporotic vertebral fractures are particularly dif■cult to diagnose as they may be clinically silent. In severe cases, signi■cant height loss (often exceeding 4 cm)and thoracic kyphosis can occur due to multiple vertebral fractures.

Osteoporosis is a common metabolic bone disease characterized by a diffuse reduction in bone density due to a decrease in bone mass. A DEXA (dual-energy X-ray absorptiometry) scan is the investigation of choice for the diagnosis and assessment of severity.

The mainstay of treatment is to reduce the risk of fractures. The following medications can be useful in the treatment:

• Bisphosphonates - They are the first line of medication. They reduce osteoclastic bone resorption and prevent bone loss. Eg. Zoledronate.

• Denosumab - This is an antibody to the RANK ligand. It prevents the activation of RANK which is required for the proliferation of osteoclasts. Therefore, it decreases bone resorption.

• Parathyroid hormone - Teriparatide is a recombinant parathyroid hormone analog. At low and intermittent doses, they can promote bone formation.

• Selective estrogen receptor modulators (SERMs) - They act on the estrogen receptors and prevent bone resorption. Eg. raloxifene.

• Strontium ranelate - It is thought to increase bone formation and reduce bone resorption.

Solution to Question 6:

Intertrochanteric fractures are the most common type of extracapsular fracture. Dynamic hip screw is the managament for undisplaced intertrochanteric fractures.

In elderly patients, internal fixation is preferred because inthem prolonged bed rest in traction may cause complications related to recumbency i.e., bed sores, pneumonia etc.

Intertrochanteric fractures of the femur can be treated by:

• Extramedullary implants - dynamic hip screw

• Intramedullary implants - proximal femoral nail and gamma nail.

Dynamic hip screws are used in undisplaced intertrochanteric fractures, whereas proximal femoral nails are used in displaced intertrochanteric fractures.





Conservative methods like Russels traction or skeleton traction in Thomas Splint are nowadays not preferred due to the success rate of operative methods.

Solution to Question 7:

Closed intramedullary nail with interlocking screws is the preferred treatment of most femoral shaft fractures.

Newer technique uses Gross-Kempf nails which have perforations at the proximal and distal ends to be locked in place, using transverse screws. Thus, it ensures stability, prevents rotation of the fractured fragments and telescoping of the implant.

In modern practice, conservative treatments like traction, bracing, and spica casts are rarely used. Traction can reduce and maintain alignment in most fractures, except for those in the upper third of the femur. Joint mobility can be preserved through active exercises. However, the major drawback is the extended bed rest required (10–14 weeks for adults), which makes it difficult to keep the femur aligned long enough for callus formation and can lead to increased patient discomfort and frustration.

After surgery, full weight-bearing should be commenced as soon as the ■xation allows. The fracture usually heals within 20 weeks and the complication rate is low; sometimes malunion or delayed union occur.

The image below shows an intramedullary nail in place for a fracture of the shaft of the femur.

Solution to Question 8:

The cardinal and the earliest sign of ankylosing spondylitis is erosion of the sacroiliac joints. Later there may be periarticular sclerosis, especially on the iliac side of the joint and ■nally bony ankylosis

Ankylosing spondylitis is a chronic inflammatory disease affecting the spine and sacroiliac joint. It is more common in young males and has a strong association with HLAB27. Males are affected more frequently than females and the usual age at onset is between 15 and 25 years. Clinically, patients present with recurring backache which worsens in the early morning or with inactivity and improves with exercise. Later they develop fatigue, pain, joint swellings, and heel pain.

Extra-articular manifestations include anterior uveitis, aortic valve insufficiency, heart blocks, and pulmonary fibrosis. Examination findings include loss of lumbar lordosis, increased thoracic kyphosis, diminished spinal movements, and decreased chest expansion.

The ESR and CRP are usually elevated during active phases of the disease. HLA-B27 is present in 95% of cases. Serological tests for rheumatoid factor are usually negative

X-ray findings:

• Erosion of sacroiliac joints- earliest sign

• Periarticular sclerosis in later stages

• Squaring of vertebrae due to the flattening of normal anterior concavity of the vertebral body is the earliest vertebral change

• Bamboo spine appearance due to bridging of adjacent vertebrae

• Wedging of the vertebra leading to hyperkyphosis

Treatment consists of maintaining posture and preserving mobility. Pain and stiffness can be controlled with anti-inflammatory drugs. In severe cases, TNF inhibitors are used. Surgery can be considered to correct the deformity.

Option D: Non erosive arthritis is a feature of systemic lupus erythematosus (SLE).

Solution to Question 9:

The given scenario of a 60-year-old female with trivial trauma unable to bear weight and X ray findings suggestive of destruction of the hip joint and periarticular bone and shaft of the tibia fracture is most likely due to metastasis.

Its more likely this patient has metastasis because:

• Age: 60-year (old age favors cancer)

• Overall bone denisty is normal

Pathological fractures are fractures that occur in a weakened bone that has structurally changed due to some condition. They occur either spontaneously or due to minor trauma. The features suggestive of a pathological fracture include:

• Pain at the site before the fracture

• Multiple recent fractures

• Elderly patient

• History of primary malignancy

• Unusual fracture pattern (Banana fracture - transverse fracture through an abnormal area of bone, similar to breaking a segment from a banana)

Other Options:

Option A: Brown tumor: Seen in hyperparathyroidism. X-rays show lytic lesions, but this is rare and typically involves long-standing disease in younger patients. Focal osteoblastic activity. Overall bone denisty is low.

Option B: Disseminated TB: Pott's disease or monoarticular destruction (e.g., hip or spine). Hip joint is usually destroyed and articular surface erosion is present. Disseminated TB rarely causes widespread fractures. Here there is no joint involvement.

Option C: Eosinophilic granulomas: Typically seen in younger individuals (children/young adults). X-rays reveal punched-out lesions, not diffuse destruction.

Solution to Question 10:

Closed reduction internal fixation is the surgery of choice for a 40-year-old patient with an intracapsular fracture of the femur.

A displaced fracture neck of the femur in an adult who is physiologically less than 65 years of age is best treated by closed reduction and internal fixation with multiple cannulated cancellous screws. Open reduction is only done if closed reduction fails, as the complication rate is higher.

Proximal femur fractures are classified based on their location near the hip joint. The main types include:

• Intracapsular Fractures:: It occurs within the joint capsule and involves the femoral neck and head.

Subtypes:

• Femoral head fracture

• Femoral neck fracture (common site for avascular necrosis and non-union).

2. Extracapsular Fractures: Occur outside the joint capsule.

Subtypes:

• Intertrochanteric fracture: Between the greater and lesser trochanters.

• Subtrochanteric fracture: Below the lesser trochanter.

• Trochanteric fracture: Isolated fracture of the greater or lesser trochanter.

Management of Intracapsular Fractures depends on the age of the patient:

1. > 65 years: Replacement - Hemi/Total Replacement (Option C and D)

2. < 65 years: is based on duration of fracture

• <3 weeks- CRIF with cannulated cancellous screws (avoid hematoma)

• >3 weeks- Check the viability of head- further management depends on the viability of head

Solution to Question 11:

FRAX is related to Fracture risk assessment.

It is used to predict the risk of fracture in a person with osteoporosis within the next 10 years.

It is used to recommend treatment or suggest ways of preventing osteoporosis.

The FRAX tool can be used to guide treatment decisions in people who meet the following three conditions:

• Postmenopausal women or men age 50 and older.

• People with low bone density (osteopenia).

• People who have not taken an osteoporosis medicine.

Orthopedics NEET 2024

Question 1:

A patient met with road traffic accident and complained of severe back ache. The X-ray is shown below. What fracture does the patient have?

a) Chance fracture

b) Burst fracture

c) Compression fracture

d) Fracture of spinous process

Question 2:

An 18-year-old boy complains of fever and pain in the lower limb. There is a localised rise of temperature and tenderness in the region shown in the radiograph. ESR is raised. What is the most probable diagnosis?



a) Osteosarcoma

b) Ewing's sarcoma

c) Giant cell tumour

d) Osteomyelitis

Question 3:

A child sustained an injury as shown in the below radiograph. What is the classification used and the grade used for this type of injury?

a) Salter harris classification-type III

b) Gartland's classification-type IV

c) Gartland's classification-type III

d) Salter harris classification-type IV

Question 4:

A patient sustained a fracture of the base of the fifth metatarsal in the right foot. He was advised below knee pop cast. How long should the cast be put?

a) 6-8 weeks

b) 2-3 weeks

c) 3-5 weeks

d) 16-20 weeks

Question 5:

Following a road traffic accident, a patient with an intertrochanteric fracture of the femur was treated with the implant shown below. Identify the implant.



a) Proximal femoral nail

b) Dynamic hip screw

c) Austin Moore prosthesis

d) Herbert screw

Question 6:

A young woman, following an intense workout involving heavy weight lifting, presents with lower back pain radiating to her right hip and extending down to her right ankle. MRI shows the following finding. Based on the location and radiation of her pain, which of the following nerve roots is most likely involved?



a) L_4

b) L_5

c) S_1

d) L_3

Question 7:

A child with rickets not responding to treatment is being evaluated. Investigations show a serum calcium level of 9 mg/dL, low phosphate level of 2 mg/dL, elevated alkaline phosphatase, and normal parathyroid hormone levels with normal arterial blood gas levels. What is the diagnosis?

a) Vitamin D resistant rickets type 2

b) Hypophosphatemic rickets

c) Vitamin D resistant rickets type 1

d) Nutritional rickets

Question 8:

What is this type of amputation called?



a) Above knee amputation

b) Below knee amputation

c) Knee disarticulation

d) Syme's amputation

Question 9:

A patient complains of decreased sensation in the lateral three-and-a-half fingers on the palmar side of the right hand, along with difficulty moving the upper phalanx of the thumb. Examination reveals thenar atrophy. Which of the following structures, when affected, lead to this presentation?

a) High radial nerve

b) High median nerve

c) High ulnar nerve

d) Carpal tunnel

Question 10:

Patient presented with a twisting injury of the knee. Identify the test being performed.

a) McMurray's test

b) Lachman test

c) Anterior drawer test

d) Posterior drawer test

Question 11:

A child sustained a fracture after a fall. An X-ray taken after fracture management is shown below. By which of the following mechanisms will fracture healing occur?



a) Secondary healing

b) Primary healing

c) Creeping osteogenesis

d) Intermembranous ossification

Question 12:

A 58-year-old female patient presents with a pathological fracture of the hip and femur. Her serum calcium and phosphate are increased. DEXA scan confirms decreased bone density. Which of the following parameters will increase in this condition?

a) Lactate dehydrogenase

b) Aspartate aminotransferase

c) Alanine transaminase

d) Tartrate resistant acid phosphatase

Question 13:

A fracture line passing through which of these will impair longitudinal growth of the bone?

a) Epiphyseal plate

b) Epiphysis

c) Metaphysis

d) Diaphysis

Question 14:

A 56-year-old woman presented with chronic lower back pain. MRI of the lower back shows the following findings. Identify the diagnosis.

a) Renal osteodystrophy

b) Spinal Tuberculosis

c) Spinal metastasis

d) Spondylolisthesis

Question 15:

A 21-year-old patient with a history of a fracture had a cast applied. On removal of the cast, the following deformity was seen. What is the most likely cause of this deformity?



a) Suprachondylar humerus fracture

b) Radial fracure

c) Mid shaft fracture of humerus

d) Lateral condylar fracture

Question 16:

A 10-year-old child fell down the stairs. An X-ray was done. Identify the structure injured as marked in the X-ray.



a) Medial tubercle

b) Tibial tuberosity

c) Gerdy's tubercle

d) Anterior intercondylar area

Answer Key

Question No.	Correct Option
1	a
2	b
3	c
4	a
5	b
6	b
7	b
8	b
9	d
10	b
11	b
12	d
13	a
14	d
15	a
16	b

Detailed Explanations

Solution to Question 1:

The X-ray shows a vertebral fracture with a horizontal fracture line passing through the body of the vertebra. This indicates Chance fracture.

Chance fracture is also known as seat-belt injury or the jack-knife injury. This injury is typically located in the upper lumbar spine in people involved in motor vehicle collisions while wearing a lap seat belt without a shoulder harness. The sudden deceleration causes flexion and posterior distraction of the spine. This causes the mid-lumbar spine to bend like a jack-knife around an axis anterior to the vertebral column.

There is a tear in the vertebra that passes transversely through the bones (bony Chance fracture), or the ligamentous structures (soft Chance fracture), or both. There is little or no crushing of the vertebral body. However, it is an unstable injury as the posterior and middle columns are distracted. Neurological damage is uncommon. Associated intra-abdominal injuries are seen in 50% of these injuries.



X-rays may show a horizontal fracture with the split running through the vertebra. It may involve the spinous process, the transverse processes, pedicles and the vertebral body. There may be opening up of the disc space posteriorly.

Most Chance fractures can be managed conservatively with bed rest and bracing. If posterior fixation is done, it will allow early mobilization. Ligamentous Chance fractures require posterior instrumented fusion.

Other options:

Option B: Burst fracture results from severe axial compression that explodes the vertebral body. The lateral view CT image showing the L4 vertebra burst fracture is given below.

Option C: Spinal compression fractures occur as a result of injury, common fall onto the buttock or pressure from normal activities, to the weakened vertebrae due to osteoporosis. Compression causes a short oblique fracture. The image showing the L4 vertebra compression fracture is given below:

Option D: Fracture of the spinous process of C7(Clay Shoveller's Fracture) is an example of an avulsion injury which may occur with severe voluntary contraction of the muscles at the back of the neck.

Solution to Question 2:

The clinical features and radiograph showing onion peel type of periosteal reaction point towards a diagnosis of Ewing's sarcoma.

Ewing's sarcoma is a malignant tumour that occurs more frequently in males, typically between 10 and 20 years of age and has a diaphyseal long-bone location. The femur is the most common long-bone site, followed by the tibia, humerus and fibula. Ewing's sarcoma is believed to arise from mesenchymal stem cells in the bone marrow.

Pain is the earliest symptom, followed by swelling and a low-grade fever. Radiographically, an aggressive, permeative, poorly defined osteolytic lesion with cortical destruction, periosteal reaction and large, radiolucent soft-tissue mass may be found. Periosteal reaction is common in young patients with the lamellar 'onion-skin' appearance causing fusiform bone enlargement. Ewing's sarcoma is composed of primitive, undifferentiated, small, round blue cells with large nuclei and scant cytoplasm. The characteristic translocation between chromosomes 11 and 22 results in a fusion gene, EWSR1-FL11, which is present in 90% of cases.

Preoperative chemotherapy frequently results in extensive tumour necrosis and shrinkage. Agents used in Ewing's sarcoma include vincristine, ifosfamide, doxorubicin and etoposide. Wide excision may be combined with adjuvant radiotherapy if surgical margins are poor. A more favourable outcome is expected with younger age, distal appendicular sites, small tumour volume, normal

ESR and greater chemotherapy necrosis response.



Other options:

Option A: Osteosarcoma has a periosteal reaction characterised as the Codman triangle. It can also be seen in Ewing's sarcoma.



Option C: Giant cell tumor has a characteristic soap bubble appearance.

Option D: Acute osteomyelitis in children presents with fever, pain, refusal to bear weight, elevated WBC count, elevated CRP, and elevated ESR. During the first week, the X-ray appears to be normal and by the second week, a faint extracortical line may be visible due to periosteal new bone formation. This is a classic sign of early pyogenic osteomyelitis. A combination of regional osteoporosis with a localized segment of apparently increased density is an important late finding in this condition.

Chronic osteomyelitis presents with erythema, chronic pain, swelling, tenderness, and discharging sinus tract. X-ray shows sequestrum is a piece of devitalized bone that has been separated from its surrounding bone during the process of necrosis, involucrum is the thick sheath of periosteal new bone that surrounds the sequestrum and cloacae are openings in the involucrum that allow drainage of purulent and necrotic material out of the dead bone.

Solution to Question 3:

The given clinical features and X-ray image are suggestive of a supracondylar fracture of the humerus (extension type) and Gartland classification is used in this case. The extension type fracture which is displaced with no intact posterior cortex is type 3 as per this classification.

Gartland classification (in children):

• Type I: Undisplaced

• Type II: Displaced but posterior cortex is intact

• Type III: Displaced but no intact posterior cortex and distal fragment could be either displaced:

• Posteromedial

• Posterolateral

Supracondylar fracture of the humerus is most commonly seen in children of 5-7 years after a fall on an outstretched hand with the elbow in full extension. It is most commonly of extension type followed by flexion type. Gartland classification is used for supracondylar fractures. The three-point bony relationship is maintained in supracondylar fracture (option B). Three-point bony landmarks in the elbow are the tips of medial and lateral epicondyles and the olecranon.

Nerve and vessel injury:

• The most common nerve to be injured in extension type - anterior interosseous nerve > Median nerve > Radial nerve > Ulnar nerve. (AMRU)

• The most common nerve to be injured in the flexion type the ulnar nerve.

• The most common vessel to be injured - the brachial artery.

Radiological findings: An undisplaced fracture may show a triangular lucency like a sail of a yacht (sail sign), also called a fat pad sign. This is due to the fat pad around the elbow being pushed forward by the hematoma. Baumann's angle (humeral-capitellar angle) can help in assessing the degree of medial angulation before and after reduction.

The most common complication of supracondylar fracture is malunion. This leads to cubitus varus or gunstock deformity. Malunion occurs due to improper reduction and not due to physeal damage, as the fracture is extraphyseal Other complications include myositis ossificans, compartment syndrome, and Volkmann ischemic contracture.

Undisplaced supracondylar humerus fractures are treated by an above-elbow plaster slab with the elbow in 90° flexion. Displaced fractures are treated by closed reduction and K-wire fixation since they tend to slip after reduction. Open fractures or those associated with brachial artery injury are treated by open reduction and internal fixation.

Other options:

Options A and D: Salter and Harris created a classification of physeal injuries for risk stratification.

• Type 3: Here, the fracture splits the epiphysis and then goes transversely in either direction into the hypertrophic layer of the physis. This leads to damage of the growth plate leading to growth disturbance.

• Type 4: The fracture splits the epiphysis and extends into the metaphysis. Displacement can occur in these types leading to asymmetrical growth.

Solution to Question 4:

The given X-ray indicates a Jones fracture. Conservative management involves a short-leg non-weight bearing cast for 6-8 weeks.

A Jones fracture is typically caused by forced inversion of the foot, such as during a pothole injury, leading to the avulsion of the base of the fifth metatarsal. This injury is due to the pull of the peroneus brevis tendon or the lateral band of the plantar fascia, which inserts onto the base of the fifth metatarsal.

Surgical correction, such as intramedullary screw fixation, is recommended for faster healing in highly active individuals. Proximal avulsion fractures can generally be managed symptomatically with rest and support, followed by early mobilization and return to function.

Solution to Question 5:

The above radiograph shows an extramedullary implant called a dynamic hip screw.

Intertrochanteric fractures of the femur can be treated by:

• Extramedullary implants - dynamic hip screw

• Intramedullary implants - proximal femoral nail and gamma nail.

Dynamic hip screws are used in undisplaced intertrochanteric fractures, whereas proximal femoral nails are used in displaced intertrochanteric fractures.

The below image shows a dynamic hip screw.





Advantages of using a proximal femoral nail over a dynamic hip screw include:

• Smaller incisions

• Reduced blood loss

• Decreased femoral neck shortening

• Decreased risk of infection

Other options:

Option A: The below image shows a proximal femoral nail with locking and stabilization screws.





Option C: Austin Moore's implant is an older implant used in unipolar hemiarthroplasty. It can be identified by the fenestrations on the shaft of the prosthesis, as shown below.

Option D: Herbert screw is cannulated and threaded at both ends. It is designed for use in fractures of small articular bones such as carpals and scaphoid.

Solution to Question 6:

Based on the clinical presentation of lower back pain radiating to the right hip and ankle, along with an MRI showing L4-L5 disc prolapse, the likely nerve root involved is L5. In a typical L4-L5 disc herniation, the L5 nerve root is most commonly compressed.

Other options:

Option A: L4: Involvement typically causes symptoms around the knee and anterior lower leg, but not the ankle.

Option C: S1: Compression here causes symptoms in the posterior leg, calf, and heel, differing from this case's distribution of pain.

Option D: L3: Affects the anterior thigh, with no radiating pain below the knee.





Solution to Question 7:

The given scenario indicates the diagnosis of hypophosphatemic rickets because of the following features:

• normal calcium levels (Normal: 9-11 mg/dL)

• normal parathyroid hormone levels

• decreased phosphate (Normal: 2.5-4.3 mg/dL)

• elevated levels of alkaline phosphatase (Normal: 33-96 U/L)

• normal arterial blood gas levels

Hypophosphatemic rickets:

X-linked dominant hypophosphatemic rickets are the most common genetic cause of rickets while the most common, overall is nutritional vitamin D
deficiency. PHEX gene (phosphate-regulating endopeptidase on x chromosome) products are responsible for the inactivation of phosphatonin. Mutation of this PHEX leads to increased phosphatonin (FGF 23) levels, and hence, the excretion of phosphorous is increased. This is also associated with impaired renal tubular reabsorption of phosphate leading to its excess excretion.

The chart attached below summarizes the evaluation of hypophosphatemic rickets.



Other options:

Options A and C: In vitamin D-resistant rickets, the renal tubule's inability to retain phosphate results in the loss of excessive phosphate through urine. Therefore, even normal or higher supplementations of vitamin D are not sufficient for bone mineralization.

• Vitamin D-dependent Type 1 rickets is due to decreased activation of vitamin D to its active form in the kidney.

• Vitamin D-dependent Type 2 rickets is due to resistance to vitamin D in peripheral tissues. This resistance to vitamin D can be overcome with high levels of this vitamin by supplementation.

Option D: Nutritional rickets occur due to a deficiency of vitamin D. This leads to low calcium levels in the blood. To raise the serum calcium, the secretion of the parathyroid hormone rises. The parathyroid hormone tries to raise the calcium levels by reabsorbing calcium in the kidneys and secreting phosphate in exchange leading to low phosphate levels and increasing bone resorption.

Solution to Question 8:

The given image shows below-knee amputation.

By retaining the knee joint, a below-knee amputation allows for better mobility and function compared to an above-knee amputation. After surgery, patients can usually be fitted with a prosthetic limb, enabling them to walk with a relatively normal gait.

Other options:

Option A: Above-knee amputation (AKA): Surgical removal of the lower limb above the knee joint, typically at the femur, resulting in the loss of the knee and requiring a more complex prosthetic for ambulation.

Option C: Knee disarticulation: Amputation through the knee joint, preserving the femur while removing the tibia and fibula, allowing for better weight-bearing but with a longer residual limb

than AKA.

Option D: Syme's amputation: Removal of the foot at the level of the ankle joint, preserving the heel pad for weight-bearing and creating a durable limb for ambulation without complex prosthetics.

Solution to Question 9:

The most likely nerve injured here is the median nerve, which occurs in Carpal tunnel syndrome.

Carpal tunnel syndrome is the most common entrapment mononeuropathy caused by compression of the median nerve as it passes through the fibro-osseous carpal tunnel beneath the flexor retinaculum. Patients present with pain and paresthesia over the lateral 3 and a half digits. There may be a weakness of thumb opposition. In chronic cases, the thenar eminence is lost due to thenar atrophy. Carpal tunnel syndrome can be associated with rheumatoid arthritis, pregnancy, chronic renal failure, acromegaly, hypothyroidism, and gout. The diagnosis is usually made on clinical grounds but nerve conduction studies can confirm the diagnosis.

Other tests for CTS -

• Phalen's test - Elicited by holding the wrist flexed fully. The appearance of sensory symptoms within 60 seconds implies a positive test.

• Tinel's sign - Sensory symptoms reproduced by percussing over the median nerve imply a positive test.

The given image shows carpal tunnel syndrome:



Non-surgical Management of Carpal Tunnel Syndrome: In mild to moderate cases of carpal tunnel syndrome, non-surgical management options can be explored to alleviate symptoms and improve function. These conservative treatments include:

•

Wrist splinting: Wearing a wrist splint at night can help maintain the wrist in a neutral position, reducing pressure on the median nerve and providing relief from nocturnal symptoms.

•

Activity modification: Identifying and modifying activities that exacerbate symptoms can help in managing carpal tunnel syndrome. For example, taking frequent breaks during repetitive tasks or adjusting the workstation ergonomics can be beneficial.

•

Nonsteroidal anti-inflammatory drugs (NSAIDs): Over-the-counter NSAIDs, such as ibuprofen, can help manage pain and inflammation associated with carpal tunnel syndrome.

•

Corticosteroid injections: In some cases, corticosteroid injections into the carpal tunnel may provide temporary relief from symptoms by reducing inflammation and pressure on the median nerve.

•

Physical and occupational therapy: Therapists can help patients learn exercises to stretch and strengthen the hand and wrist muscles, improving symptoms and function.

If conservative treatments do not provide sufficient relief or the condition worsens, surgical intervention, such as carpal tunnel release surgery, may be considered.

Other options:

Option A: High radial nerve injuries typically cause wrist drop due to weakness in wrist and finger extensors, with sensory loss in the dorsal aspect of the hand, not the lateral three and a half fingers on the palmar side.

Option B: High median nerve injuries can affect sensation in the lateral three-and-a-half fingers on the palmar side and impair thumb flexion and opposition, particularly at the proximal phalanx. There is a loss of pronation due to the paralysis of the pronator teres and pronator quadratus. Paralysis of flexor digitorum superficialis and lateral half flexor digitorum profundus results in loss of wrist flexion. Ape-like hand and pointing index are due to this injury.

Option C: High ulnar nerve injuries result in sensory loss in the medial one-and-a-half fingers (pinky and ring finger) and hand intrinsic muscle weakness, unrelated to thenar muscle involvement.

Solution to Question 10:

The image shows an examination being performed with the knee flexed to about 20 degrees which indicates that it is the Lachman test. The presence of free gliding indicates that it is a positive Lachman test.

Lachman test: The patient's knee is flexed to 20 degrees. The lower part of the thigh is grasped with one hand and the upper part of the leg with the other hand. The joint surfaces are shifted backward and forward upon each other. If the knee is stable, there should be no gliding. The presence of gliding indicates the laxity of the anterior cruciate ligament.

Other options:

Anterior and posterior drawer test: The test is performed with the patient's knees flexed to 90 degrees, and the feet resting on the examining bed. The examiner then grasps the upper part of the leg with both hands and rocks it forward and backward. Excessive anterior movement denotes anterior cruciate laxity (anterior drawer test) Excessive posterior movement denotes posterior cruciate laxity (posterior drawer test).

McMurray's test: The test is used to identify torn meniscus. The knee is flexed as far as possible and the joint is stabilized with one hand. With the other hand, the examiner rotates the leg medially and laterally. If there is a clicking sound while rotating the leg, the test is positive. The sound is produced due to the trapping of the loose tag of the torn meniscus between the articular surfaces.

Solution to Question 11:

The clinical vignette describes a child sustaining a fracture, and the X-rays show compression plating. In this case, the fracture heals through primary healing without callus formation.

The stages of fracture healing are as follows:

• Hematoma Stage (up to 7 days): The periosteum and soft tissues detach from the fracture ends, leading to ischemic necrosis. Some osteocytes die from lack of blood flow, while others transform into daughter cells that aid in healing.

• Granulation Tissue Stage (2-3 weeks): Sensitized daughter cells differentiate into blood vessels, fibroblasts, and osteoblasts, forming soft granulation tissue that fills the fracture gap.

• Callus Stage (4-12 weeks): Granulation tissue differentiates into osteoblasts that produce an intercellular matrix infused with calcium salts, forming a callus (woven bone), which is visible on X-rays around 3 weeks after the fracture.

• Remodeling Stage (1-2 years): The woven bone is replaced by mature lamellar bone, known as the secondary callus, through a multicellular process of substitution.

• Modeling Stage: The bone undergoes gradual strengthening and reshaping of its periosteal and endosteal surfaces, driven by local strains from weight-bearing stresses and muscle forces during normal activities.

Other options:

Option A: Secondary healing is more typical for fractures that have not been perfectly aligned or stabilized, where a callus is formed (soft callus forms first, followed by a hard callus as healing progresses).

Option C: Creeping osteogenesis refers to the process where new bone forms along the surface of existing bone, typically seen in the context of distraction osteogenesis or in cases where there is a gap that needs to be filled.

Option D: Intramembranous ossification is a type of bone formation that occurs primarily in flat bones and does not typically apply to the healing of fractures.

Solution to Question 12:

In this patient, the pathological fracture and elevated serum calcium and phosphate suggest a possible metabolic bone disease such as Paget's disease of bone. The parameter likely to be increased is TRAP (tartrate-resistant acid phosphatase), which is a marker of osteoclast activity.

TRAP is an enzyme produced by osteoclasts, which are cells responsible for bone resorption. Elevated levels of TRAP are often seen in conditions with increased bone turnover, such as Paget's disease, hyperparathyroidism, and certain bone malignancies.

Solution to Question 13:

A fracture line passing through the epiphyseal plate (also known as the growth plate) will impair the longitudinal growth of the bone.

The epiphyseal plate is the area of growing tissue near the ends of the long bones in children and adolescents. It is responsible for bone lengthening (longitudinal growth). If the fracture involves the epiphyseal plate, it can disrupt this process and lead to growth disturbances or deformities.

Other options:

Option B: The epiphysis is the end of a long bone. While fractures involving the epiphysis can affect joint function, they don't directly impair longitudinal growth unless the epiphyseal plate is involved.

Option C: The metaphysis is the region of bone between the diaphysis and the epiphysis, where bone growth in width occurs. A fracture here doesn't directly impact longitudinal growth.

Option D: Diaphysis describes the shaft of the bone. A diaphyseal fracture typically affects the strength and structural integrity of the bone but doesn't impair longitudinal growth.

Solution to Question 14:

The above clinical scenario, the history and MRI scan are suggestive of spondylolisthesis.

Spondylolisthesis is the anterior displacement of a vertebra over another, usually at the lower lumbar spine. Occasionally, the displacement is backward (retrolisthesis). It is characterized by chronic back pain, lumbar lordosis, and step sign. Normally, spinous processes of adjacent vertebrae can be palpated on the back smoothly. In spondylolisthesis, a step-off or step sign can be felt over the spinous process at the affected level.

The Beheaded Scottish terrier sign (shown in the image below) on the x-ray is diagnostic. It is due to a break in the pars interarticularis of the vertebra.



It is usually managed conservatively based on symptoms with analgesics, temporary activity restriction, muscle relaxants, physical therapy, and strengthening exercises.

Other options:

Option A: Renal osteodystrophy is a complex condition that results from the kidneys' inability to maintain the proper levels of calcium and phosphate in the blood, leading to secondary hyperparathyroidism. The below image represents Rugger Jersey spine which is characteristic of hyperparathyroidism seen in renal osteodystrophy.

Option B: TB spine occurs secondary to a pulmonary or gastrointestinal primary and spreads through the Batson's plexus of veins (paravertebral plexus). It most commonly involves the dorso-lumbar region.



Option C: Spinal metastasis: Represents the spread of malignancy to the spine, causing bone destruction. MRI would demonstrate vertebral lesions, bone marrow involvement, or spinal cord compression, without evidence of vertebral slippage. The following image shows a radiograph of spinal metastasis.

Solution to Question 15:

The given image shows cubitus varus deformity, where the forearm has deviated medially in relation to the upper arm, which is a common complication of supracondylar humeral fracture.

Malunion is the commonest complication of a supracondylar fracture and results in a cubitus varus deformity, also known as gunstock deformity. This occurs when the fracture unites with the distal fragment tilted medially and in internal rotation.

The image below shows gunstock deformity:

Cubitus varus is primarily a cosmetic concern with minimal functional impairment. Mild cases may not necessitate treatment, while severe deformities should be corrected through a lateral closed wedge osteotomy (French osteotomy).

Solution to Question 16:

The given X-ray points towards the tibial tuberosity. This image shows an avulsion fracture of the tibial tuberosity.

The tibial tuberosity is a bony prominence located on the anterior part of the tibia, just below the knee. It is the attachment point for the patellar tendon. Injury to the tibial tuberosity is common in adolescents, especially those involved in sports or after trauma like a fall.

Other options:

Option A: The medial tubercle refers to part of the medial condyle of the tibia.

Option C: Gerdy's tubercle is located on the lateral aspect of the tibia, where the iliotibial band (IT band) inserts. It is less commonly injured in falls but can be affected in conditions related to the IT band, like IT band syndrome.

Option D: Anterior intercondylar area is an area is located between the medial and lateral condyles of the tibia and serves as an attachment point for the anterior cruciate ligament (ACL). Injuries in this region are more commonly associated with ACL tears.

Paediatrics AIIMS 2017

Question 1:

A 4-year-old child was brought to the hospital with a history of loose stools. No history of fever or blood in stools. The mother says he is irritable and eagerly drinks water if given. On examination, eyes are sunken and in the skin pinch test, the skin retracted within two seconds but not immediately. What treatment is advised for the child?

a) Give oral fluids and zinc supplementation only and ask mother to come back if danger sign develops.

b) Give oral fluids and ask mother to continue the same and visit again the next day.

c) Consider severe dehydration start IV fluids, IV antibiotics, and refer to higher centre.

d) Wait and observe.

Question 2:

A baby in the NICU is being evaluated for his features of microcephaly and hepatomegaly. CT scan revealed nodular periventricular calcifications. Congenital CMV infection is suspected. Which of the following samples is the best to detect the organism using PCR?

a) Liver biopsy

b) CSF

c) Blood

d) Urine

Question 3:

Identify the disease shown in the karyotype:



a) Cri-du–chat syndrome

b) Bloom syndrome

c) Angelman syndrome

d) Fragile X syndrome

Question 4:

Which of the following milestones would you generally expect a 3 year old child to display?

a) Hopping on one foot

b) Spoon handling

c) Copying a triangle

d) Repeats sentence of 10 syllables

Question 5:

Dried blood spot (DBS) testing is commonly done in infants for:

a) Inborn errors of metabolism

b) Blood grouping

c) Bilirubin testing

d) Coombs Test

Question 6:

A 4 day old baby is active and feeding well. He has a bilirubin level of 8 mg/dl. What is the next step in the management of the baby?

a) Exchange transfusion

b) Stop breastfeeding and start phototherapy

c) IV fluids and phototherapy

d) Continue breastfeeding

Question 7:

The most appropriate expansion of Apgar is:

a) Alert, pulse, grimace, activity, respiration

b) Assessment, pulse, grimace, alive, respiration

c) Appearance, pulse, grimace, activity, respiration

d) Appearance, pressure, grimace, activity, respiration

Question 8:

A 15-day-old infant had conjunctivitis, which later developed into respiratory distress and pneumonia. Chest Xray examination revealed bilateral lung infiltrates. Peripheral smear count showed 14,300/µL (N 43, L 54, E3). Which of the following is the most likely pathogenic organism?

a) Chlamydia trachomatis

b) Streptococcus pneumonia

c) Neisseria gonorrhoeae

d) Mycoplasma pneumonia

Question 9:

Of the following, all are findings of the condition in the below chest x-ray except:

a) Increased pulmonary blood flow

b) Narrow vascular pedicle

c) Pulmonary venous hypertension

d) Right atrial enlargement

Question 10:

In fetal alcohol syndrome, which of the following is not seen?

a) Thinning of corpus callosum

b) Microcephaly

c) Holoprosencephaly

d) Macrocephaly

Question 11:

A child presented with complaints of itching and discharges on flexure aspect of elbow and the lesions were red as shown below. What is your diagnosis?

a) Pemphigus vulgaris

b) Psoriasis

c) Post traumatic pigmentation

d) Atopic dermatitis

Question 12:

Which of the following fits into the criteria of severe variable deceleration, variations less than:

a) 90 beats per minute lasting for 60 seconds

b) 80 beats per minute lasting for 60 seconds

c) 100 beats per minute lasting for 60 seconds

d) 70 beats per minute lasting for 60 seconds

Question 13:

Which of the following constitutes Hutchinson's triad?

a) Interstitial keratitis, 8th nerve deafness, mulberry molars

b) Interstitial keratitis, 8th nerve deafness, Hutchinson molars

c) Interstitial keratitis, 8th nerve deafness, mulberry incisors

d) Interstitial keratitis, 8th nerve deafness, Hutchinson incisors

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	b
5	a
6	d
7	c
8	a
9	d
10	d
11	d
12	d
13	d

Detailed Explanations

Solution to Question 1:

In a child with diarrhea, thirst (eagerly drinks water), sunken eyes, restless and irritable behavior, and skin pinch showing slow retraction (<2 seconds) point towards some dehydration. Treatment includes oral rehydration therapy, zinc supplementation, and continued breastfeeding according to WHO integrated management of neonatal and childhood diseases (IMNCI) plan B.

The degree of dehydration is assessed and managed using the WHO IMNCI PLAN given below:

SIGNS	CLASSIFY AS	IDENTIFY TREATMENT (Urgent pre-referral treatments are in bold print)
Two of the following signs: • Lethargic or unconscious • Sunken eyes • Not able to drink or drinking poorly • Skin pinch goes back very slowly	SEVERE DEHYDRATION	► If child has no other severe classification - Give fluid for severe dehydration (Plan C). OR If child also has another severe classification: Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way. Advise the mother to continue breastfeeding. ► If child is 2 years or older, and there is cholera in your area, give antibiotic for cholera.
Two of the following signs: • Restless, irritable • Sunken eyes • Drinks eagerly, thirsty • Skin pinch goes back slowly.	SOME DEHYDRATION	► Give fluid, Zinc supplements and food for some dehydration (Plan B) ► ***If Child also has a severe classification:*** ***Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way. Advise the mother to continue breastfeeding.*** ► Advise mother when to return immediately. ► Follow-up in 5 days if not improving. ► If confirmed/symptomatic HIV, follow-up in 2 days if not improving.
• Not enough signs to classify as some or severe dehydration	NO DEHYDRATION	► Give fluid, Zinc supplements and food to treat diarrhoea at home (Plan A) ► Advise mother when to return immediately. ► *Follow-up in 5 days if not improving.* ► If confirmed/symptomatic HIV, follow-up in 2 days if not improving.



Solution to Question 2:

Among the following options, urine is the best sample to be sent for the diagnosis of congenital cytomegalovirus (CMV) infection.

In congenital CMV infections, the investigation of choice is PCR or viral culture. It should be done within the first three weeks of life (after this period, it is difficult to distinguish between a congenital and an acquired infection). Since not all affected infants are viremic at birth, urine is a more reliable sample than blood.

Most CMV transmission occurs in the first trimester. The risk of transmission is high in the third trimester. 90% of the infected babies are asymptomatic and have lesser chances of having late sequelae.

The clinical manifestations include
hepatosplenomegaly, petechial rashes, jaundice, chorioretinitis, and microcephaly. Neuroimaging shows intracranial periventricular calcifications. One of the long-term sequelae is sensorineural hearing loss.

There is no definitive treatment for congenital CMV infection. Treatment
with ganciclovir is indicated only in patients with progressive neurologic disease and deafness.

The image below shows periventricular calcifications:

Solution to Question 3:

In the karyotype, there is a partial deletion of the short arm of chromosome 5, which leads to Cri-du-chat syndrome. Partial aneuploidy occurs as a result of the breakdown of the chromosomes.

Cri-du-chat syndrome is a rare genetic disorder. Features of the condition are hypotonia, short stature, characteristic cry called a cat's cry in the first few weeks of life, microcephaly with protruding metopic sutures, hypertelorism, bilateral epicanthic folds, high arched palate, flat nasal bridge, and intellectual disability.

Diagnosis is based on clinical findings and chromosomal analysis. Fluorescence in situ hybridization (FISH) is the specific test to confirm the diagnosis.

Given below is a picture of the facies in cri-du-chat syndrome:

Other options:

Option B: Bloom syndrome is a rare autosomal recessive disorder characterized by short stature, sun sensitivity, predeposition of development of cancer, and genomic instability.

Option C: Angelman syndrome is caused due to deletion in the maternal chromosome 15 or by paternal uniparental disomy. It primarily affects the nervous system. Characteristic features are delayed development, intellectual disability, severe speech impairment, and ataxia.

Option D: Fragile X syndrome is a genetic disorder also termed Martin Bell syndrome. It is the most common cause of inherited mental retardation (among the X-linked disorders), intellectual disability, and autism.

Solution to Question 4:

The ability to handle a spoon well and feed self is usually developed by 24 months (2 years) and would be expected in a 3 year old child.

Other options:

Option A: The ability to hop on one foot usually develops by 48 months (4 years).

Options C and D: The ability to draw a triangle from copy and repeating a sentence of 10 syllables usually develops at around 60 months (5 years) of age.

Some key milestones of children from 15 months of age are listed below:

Age	Motor milestones	Adaptive milestones	Language milestones	Social milestones
15 months	Walks alone, crawls up stairs.	Makes tower of 3 blocks, draws a line with crayon, inserts raisin or coin into bottle.	Jargon, follows simple commands, may name familiar objects, responds to own name.	Hugs parents, indicates some desires or needs by pointing.

Age	Motor milestones	Adaptive milestones	Language milestones	Social milestones
18 months	Runs stiffly, sits on a small chair, walks up stairs with 1 hand held, explores drawers.	Makes tower of 4 blocks, imitates scribbling.	10 words (average), names pictures, identifies 1 or more body parts.	Feeds self, seeks help when in trouble, may complain when wet or soiled.
24 months	Runs well, walks up and down stairs – 1 step at a time, opens doors, jumps.	Makes tower of 7 blocks, scribbles in circular pattern, fold paper once in imitation.	Puts 3 words together.	Handles spoon well,helps to undress, tells immediate stories, listen to stories when pictures are involved.
30 months	Goes up stairs alternating feet.	Makes tower of 9 blocks, makes vertical and horizontal strokes but not a cross.	Refers to self with pronoun 'I', knows full name.	Pretends during play, helps put things away.
36 months	Rides tricycle, stands momentarily on 1 foot.	Makes tower of 10 blocks, imitates construction of bridge with 3 blocks, copies circle, imitates cross.	Knows age and sex, counts 3 objects correctly, repeats 3 numbers or a sentence of 6 syllables.	Helps in dressing, washes hands, plays games (in parallel with other children).
48 months	Hops on 1 foot, throws ball overhand, uses scissors to cut pictures, climbs well.	Copies bridge from model, imitates construction of gate with 5 blocks, copies cross and square, identifies longer of 2 lines.	Counts 4 coins correctly, tells story.	Goes to toilet alone, plays with several children.
60 months	Skips	Draws triangle from copy, names the heavier of two weights.	Names 4 colors,speaks sentence of 10 syllables, counts 10 coins correctly.	Dresses and undresses, asks questions about meaning of words, engages in domestic role-playing.

Solution to Question 5:

Dried blood spots (DBS) taken from the heel are used in neonatal screening for inborn errors of metabolism (IEM) using tandem mass spectroscopy (TMS).

TMS employs two mass spectrometers in series to screen blood samples for their presence and to determine concentrations of various substances like amino acids, fatty acids, and other metabolites. A single sample can be used to screen multiple IEMs such as phenylketonuria, ethylmalonic encephalopathy, and glutaric acidemia.

Newborn screening using DBS is recommended for congenital hypothyroidism, G6PD deficiency, congenital adrenal hyperplasia, and galactosemia.

Universal screening is recommended for hearing loss, cyanotic congenital heart disease (pulse oximetry), hyperbilirubinemia, and congenital hip dysplasia (Ortolani and Barlow tests).

Hearing loss is screened with otoacoustic emission (OAE). Auditory Brainstem Response (ABR) is indicated in infants who fail OAE.

Screening for hypoglycemia should be performed in infants who are small or large for gestational age, born to mothers with diabetes, preterm, or symptomatic.

Solution to Question 6:

Since this child has a bilirubin level of only 8 mg/dl on the 4th day, it points to physiologic jaundice. The child is active and feeding well, the mother should be encouraged to continue breastfeeding frequently and exclusively.

All the other options would be possible treatment choices if the child had pathological jaundice.

Physiological jaundice/Icterus neonatorum is due to increased bilirubin production from the breakdown of fetal red blood cells combined with transient limitation in the conjugation of bilirubin by the immature neonatal liver. It first appears on the 2nd or 3rd days usually peaks on the 4th day at 5-6 mg/dL and decreases to <2 mg/dL between the 5th and 7th days after birth.

On the other hand, in pathological jaundice, appearance is within the first 24 hours after birth and may be present at or beyond the age of 2 weeks in term babies and 3 weeks in preterms babies. There may be an elevated direct (or conjugated) bilirubin level. Jaundice visible on the palms and soles, unresponsiveness to phototherapy, and a rapid rise in bilirubin levels during phototherapy indicate pathological jaundice.

Phototherapy is initiated in babies with pathological jaundice when the bilirubin levels exceed the 95th percentile on the total serum bilirubin (TSB) normogram. Babies are kept at a distance of 30-45 cm from the UV lamp (460-490 nm) with eyes and gonads covered. It is avoided in babies with conjugated hyperbilirubinemia due to the risk of bronze-baby syndrome. Dehydration and hypocalcemia are potential complications and breastfeeding is encouraged.

Exchange transfusion (ET) is done in cases of bilirubin encephalopathy, Rh incompatibility with significant hemolysis, and bilirubin values exceeding TSB cutoffs for ET.

Solution to Question 7:

The parameters to consider while calculating APGAR score in newborn babies are:

• Appearance

• Pulse rate

• Grimace

• Activity(muscle tone)

• Respiratory effort

The Apgar score is a systematic way of assessing the infant after birth at 1 and 5 minutes of life. It gives a maximum score of 10 which indicates the newborn is in the best possible condition.

Changes in the Apgar scores denote the response of the infant to resuscitation. Apgar scores should not be used to determine the need for resuscitation or to guide the steps of resuscitation. Apgar score alone should not be interpreted as evidence of asphyxia.

A score from 7-10 is considered normal. Scores between 0-3 require immediate resuscitation. If the score at 5 minutes remains less than 7, the infant should be assessed again every 5 minutes for the next 20 minutes.

Causes for false positive APGAR score are:

• Prematurity

• Drugs like analgesics, narcotics, sedatives, magnesium sulfate

• Acute cerebral trauma

• Precipitous delivery

• Airway obstruction (choanal atresia)

• Congenital pneumonia or sepsis

• Hemorrhage- hypovolemia

False-negative Apgar score is seen in maternal acidosis and high fetal catecholamine levels.

Solution to Question 8:

The organism most likely to cause conjunctivitis, progressing to respiratory distress, and then pneumonia is Chlamydia trachomatis.

It is not possible to differentiate between chlamydial pneumonia and other causes of atypical pneumonia, like Mycoplasma (option D) based solely on the clinical presentation. Here, the history of conjunctivitis makes it more likely to be a chlamydial infection.

Pneumonia caused by C. trachomatis can develop in infants born to women with an untreated chlamydial infection in 10-20% of cases. It has a very characteristic presentation in infancy, disease onset usually occurs around 1 and 3 months of age and is insidious, with a persistent cough, tachypnea, and absence of fever (atypical pneumonia). Auscultation reveals rales; wheezing is uncommon.

The absence of fever and wheezing helps to distinguish C. trachomatis pneumonia from respiratory syncytial virus pneumonia.

A distinctive laboratory finding is the presence of peripheral eosinophilia (>400 cells/μL). The most consistent finding on chest radiographs is hyperinflation accompanied by minimal interstitial or alveolar infiltrates.

Other options:

Option B: Pneumococcal pneumonia generally presents with features of lobar pneumonia. Fever, dyspnea, and productive cough are the primary symptoms.

Option C: Although Neisseria gonorrhoeae is an important cause of ophthalmia neonatorum, it does not cause pneumonia-like features.

Chlamydiae are obligate intracellular organisms that produce basophilic inclusion bodies. They are called energy parasites as they lack enzymes for ATP synthesis. They cannot grow on artificial media but can grow on cell lines (McCoy, HeLa, or the yolk sac of the chick embryo)

Chlamydiae occur in two forms, the elementary body (extracellular, infective form) and the reticulate body (intracellular, non-infective, and replicative form) which are found inside inclusion bodies.

Chlamydia trachomatis is the most common cause of ophthalmia neonatorum in developed countries. Neonatal inclusion conjunctivitis is caused by serotypes D to K. Chlamydia trachomatis inclusion conjunctivitis manifests relatively late, usually over 1 week after birth.

It is usually a venereal infection derived from the cervix or urethra of the mother during delivery. On examination, the conjunctiva is swollen and edematous. The discharge is initially watery, but later becomes mucopurulent or bloody.

Newborns with suspected chlamydia infection should have samples drawn from both the conjunctivae and oropharynx. Since it is an obligate intracellular organism, swabs should also include conjunctival epithelial cells if possible. Giemsa staining shows characteristic intracellular inclusion bodies. Diagnosis is usually established by PCR.

The treatment includes oral erythromycin for two weeks. Topical erythromycin to tetracycline ointment may also be used.

If left untreated, associated systemic diseases such as chlamydial otitis and pneumonia may develop later. In all cases, both parents must receive appropriate treatment for genital infection.

Given below is an image of ophthalmia neonatorum:

Solution to Question 9:

The given chest x-ray shows an egg-on-string appearance seen in the transposition of great arteries (TGA). Enlargement of the left atrium is seen in TGA, not the right atrium.

The reason for left atrial enlargement is due to increased pulmonary blood flow.

TGA is characterized by the aorta arising from the right ventricle and the pulmonary artery from the left ventricle. This abnormal configuration leads to a narrow vascular pedicle.

As pulmonary vascular resistance drops during the first few days of life, blood flow through the pulmonary vasculature is increased. There is also elevated pulmonary venous pressure seen in TGA, especially in infants with an intact interventricular septum.

The globular (egg) appearance of the heart due to abnormal convexity of the right atrial border and left atrial enlargement, and the narrow superior mediastinum (string) result in the classical egg-on-string appearance.

TGA is the most specific heart disease in infants of diabetic mothers.

Early management includes an immediate infusion of prostaglandin E1 which keeps the ductus arteriosus patent for improving oxygenation. Rashkind balloon atrial septostomy is a palliative catheter-based intervention that can be done to improve the hypoxic state before definitive surgery. The procedure creates an opening in the atrial septum which allows better inter-circulatory mixing of the blood. Jatene arterial switch procedure is the definitive surgery.

Most infants with TGA do not survive till adulthood without corrective surgery.

Solution to Question 10:

Fetal alcohol syndrome (FAS) is associated with microcephaly rather than macrocephaly.

Alcohol exposure during pregnancy results in reduced brain volume in the frontal lobe, striatum, caudate nucleus, thalamus, and cerebellum. It also causes thinning of the corpus callosum and abnormal functioning of the amygdala.

Fetal Alcohol Spectrum Disorders (FASDs) are the leading preventable causes of developmental delay and intellectual disability. FASDs encompass different conditions such as FAS, partial FAS, alcohol-related neurodevelopmental disorder, alcohol-related birth defect, and neurobehavioral disorder associated with prenatal alcohol exposure.

Diagnostic criteria for FASDs include characteristic facial features, prenatal/postnatal growth deficiency, neurobehavioral impairment, and maternal alcohol consumption during pregnancy.

The facial features of FAS include short palpebral fissures, a thin upper lip vermilion border, a smooth philtrum, epicanthic folds, a flattened nasal bridge, lowset and unparallel ears, and retarded midfacial development.

CNS involvement leads to microcephaly, incomplete holoprosencephaly, and agenesis of the corpus callosum and the cerebellar vermis. The neurocognitive features include developmental delays, inattention, impulsivity, social impairments, and behavioral difficulties. Common comorbidities include attention-deficit/hyperactivity disorder (ADHD), self-regulation problems, and impulse control issues. Cardiac anomalies are also seen, with VSD being more common than ASD and tetralogy of Fallot.

Solution to Question 11:

The presence of diffuse, erythematous, itchy plaques on the flexural aspects of the elbow of a child is suggestive of atopic dermatitis.

Atopic dermatitis is the cutaneous manifestation of atopy, characterized by a family history of asthma, allergic rhinitis, or eczema.

There is a clear genetic predisposition. A characteristic defect is an impaired epidermal barrier. In many patients a mutation in the gene encoding filaggrin (a structural protein in the stratum granulosum) is responsible.

Patients may display many immunoregulatory abnormalities, including increased IgE synthesis, increased serum IgE levels, and delayed-type hypersensitivity reactions.

The disease progresses in three phases: Infantile, childhood, and adult.

In infants, the lesions of atopic eczema most frequently start on the face but may occur anywhere on the skin surface. Often, the napkin area is relatively spared. When the child begins to crawl, the exposed surfaces, especially the extensor aspect of the knees and elbows are most involved. The lesions consist of erythema and discrete or confluent oedematous papules. The papules are intensely itchy and may become exudative and crusted as a result of rubbing. Secondary infection and lymphadenopathy are common.

From 18 to 24 months onwards (childhood phase), the sites most characteristically involved are the elbow and knee flexures, a mixture of erythema, crusting, excoriation, hyper and hypopigmentation, and warty lichenification may be seen depending on the skin type on the sides of the neck, wrists, and ankles. The sides of the neck may show reticulate pigmentation, sometimes referred to as an atopic dirty neck. The erythematous and oedematous papules tend to be replaced by lichenification.

The picture in adults is essentially similar to that in later childhood, with lichenification, especially of the flexures and hands.

Solution to Question 12:

Severe variable decelerations are defined as: Less than 70 beats/minute lasting at least 60 seconds.

Variable deceleration is the abrupt decrease in FHR below the baseline. The decrease is 15 beats/minute or more, with a duration of 15 seconds or more but less than 2 minutes.

Mechanism: Umbilical cord compression, leads to occlusion of the umbilical vein, hence there is decreased venous return to maintain Cardiac Output.

Classification:

• Mild: more than 80 bpm irrespective of duration/ less than 30 seconds irrespective of HR/ 70-80 bpm lasting 60 seconds

• Moderate: more than 70 bpm lasting 30-60 seconds / 70-80 bpm lasting more than 60 seconds

• Severe: less than 70 bpm lasting at least 60 seconds

Solution to Question 13:

Hutchinson's triad refers to interstitial keratitis, 8th nerve deafness, and Hutchinson incisors. It is seen in late congenital syphilis.

The image below shows Hutchinson incisors:

Congenital syphilis is due to Treponema pallidum. Early features include hepato-splenomegaly, jaundice, diffuse lymphadenopathy, maculopapular or vesiculobullous rash involving hands and feet, persistent rhinitis (snuffles), and condylomatous lesions. Skeletal changes like saber shins, dental abnormalities like Hutchinson's teeth, and saddle nose (depression of the nasal root) are some late findings.

Other clinical features of congenital syphilis:

• Clutton joint - Due to synovitis, with sterile synovial fluid

• Juvenile tabes and paresis

• Rhagades - Linear scars radiating from mucocutaneous fissures of the mouth, anus, and genitalia

• Mulberry molars - Abnormal 1st lower molars with excessive cusps

• Sabre shin - Anterior bowing of the mid-portion of the tibia

• Olympian brow - Bony prominence of the forehead

Paediatrics AIIMS 2018

Question 1:

Which of the following is not an indicator of fetal lung maturity?

a) Lecithin-sphingomyelin ratio > 2.5

b) Positive shake test

c) Increased phosphatidyl glycerol

d) Nile blue test showing >50% of blue cells

Question 2:

How long should a child be isolated after being diagnosed with bacterial meningitis to prevent further transmission?

a) Till 24 hours after starting antibiotics

b) Till cultures become negative

c) Till antibiotic course is complete

d) Till 12 hours after admission

Question 3:

Choose the correct order of suctioning during neonatal resuscitation.

a) Mouth-nose

b) Nose-mouth

c) Mouth-nose-oesophagus

d) Trachea-nose-mouth

Question 4:

An adolescent girl complaints of dropping objects from hands precipitated during morning hours and exams. There is no history of loss of consciousness. One of her cousins has been diagnosed with epilepsy. EEG was suggestive of epileptic spikes. What is the most probable diagnosis?

a) Juvenile myoclonic epilepsy

b) Atypical absence

c) Choreo-athetoid epilepsy

d) Centrotemporal spikes

Question 5:

A newborn female presents with continuous dribbling of urine. The image below shows the pathology from which the patient is suffering. The most likely diagnosis is:

a) Gastroschisis

b) Bladder exstrophy

c) Superior vesical fissure

d) Anterior ectopic anus

Question 6:

The absence of which of the following milestones in a 3-year-old child is called as developmental delay?

a) Hopping on one leg

b) Drawing a square

c) Feeding by spoon

d) Passing a ball to someone

Question 7:

A 2-year-old child with a history of fall 1 year back with parietal bone fracture, has now presented with a painful and growing parietal swelling. What is the likely diagnosis?

a) Growing scalp hematoma

b) Growing fracture

c) Subdural hygroma

d) Chronic abscess

Question 8:

After the delivery of an infant of a diabetic mother, the blood glucose of the Infant was 60 mg/dl. Which other investigation would you do?

a) Serum potassium

b) Serum sodium

c) Serum chloride

d) Serum calcium

Question 9:

You have been called to declare the brain death of a 12-year-old child in the ICU. Which of the following is not seen in brain death?

a) Normal BP without pharmacological support

b) Positive spinal reflexes on stimulation

c) Sweating and tachycardia

d) Decorticate and decerebrate posturing

Question 10:

Craniopagus is defined as the fusion of _____.

a) Head and spine

b) Head only

c) Thorax and spine

d) Thorax only

Question 11:

Which of the following hormones is the major product of the fetal adrenal glands?

a) DHEA

b) Cortisol

c) Corticosterone

d) Progesterone

Question 12:

A 2-day-old neonate was admitted to the NICU after he was observed to have movements as shown in the clip. His APGAR scores were 8 and 9 at 1 and 5 minutes, respectively. His blood sugar was within the normal range in the first 24 hours. What is the likely diagnosis?



©MARROW

a) Focal tonic seizure

b) Focal clonic seizure

c) Spasms

d) Subtle seizure

Question 13:

A male child with coarse facies, macroglossia, thick lips with hepatosplenomegaly presents with copious mucous discharge from the nose. The most probable underlying diagnosis is _____

a) Hurler disease

b) Beckwith-Wiedemann syndrome

c) Proteus Syndrome

d) Hypothyroidism

Question 14:

Which pattern of inheritance does the pedigree chart denote?



a) Autosomal-dominant disorder

b) X- linked dominant disorder

c) Autosomal-recessive disorder

d) X- linked recessive disorder

Question 15:

A preterm infant in the NICU was observed to have marked nasal flare, distinct xiphoid retractions, minimal intercostal retractions, and inspiratory lag with an audible grunt. What would be the Silverman Anderson score?

a) 10

b) 7

c) 8

d) 9

Question 16:

True regarding the difference between adult and pediatric resuscitation.

a) Ventricular dysrhythmias are common in children

b) The no flow phase precedes the period of cardiac arrest in children.

c) More ventilation to be given compared to chest compressions in children

d) Dysrhythmias are most often due to respiratory insufficiency in children

Answer Key

Question No.	Correct Option
1	d
2	a
3	a
4	a
5	b
6	c
7	b
8	d
9	d
10	b
11	a
12	b
13	a
14	c
15	c
16	d

Detailed Explanations

Solution to Question 1:

Nile blue test showing >50% orange cells is a sign of fetal lung maturity, not blue cells.

Tests for the assessment of fetal lung maturity include:

The Nile blue test involves staining desquamated fetal cells from amniotic fluid with 0.1% Nile blue sulfate. The presence of orange cells > 50% suggests lung maturity. Lecithin/sphingomyelin ratio at 31 to 32 weeks is 1, and at 35 weeks ratio is 2. The ratio ≥2 indicates lung maturity (option A).

The shake test or bubble test (Clement's test) (option B) involves mixing amniotic fluid with 95% ethanol, shaking it for 15 seconds, and observing it for 15 minutes. A complete ring of bubbles at the meniscus is a positive test and indicates lung maturity. This is a useful bedside test.

The presence of phosphatidylglycerol, (option C) saturated phosphatidylcholine and lamellar body count (>50,000/mcL) indicates lung maturity.

Glucocorticoids accelerate lung maturation and lower the incidence of respiratory distress and neonatal mortality in preterms if the birth was delayed for at least 24 hours after initiation of betamethasone. A single course of corticosteroids is recommended for women between 24 and 36 weeks who are at risk of delivery within 7 days. Intramuscular injections of either 2 doses of 12 mg betamethasone are given 24 hourly or 4 doses of 6 mg dexamethasone are given 12 hourly.

Solution to Question 2:

Isolation of the child is recommended until 24 hrs after the initiation of effective therapy after being diagnosed with acute bacterial meningitis to prevent further transmission.

Droplet precautions are recommended for patients with Neisseria meningitides and Haemophilus influenzae type b (Hib) meningitis until they have received 24 hours of effective therapy. Patients should be in private rooms, and hospital personnel should wear a face mask when they are within 3 feet (1 meter) of the patient.

The child with meningitis shows nonspecific symptoms like fever, anorexia, poor feeding, headache, myalgia, hypotension, petechia, purpura, or a rash. Meningial irritation is indicated by nuchal rigidity, back pain, and Kernig, and Brudzinski signs (usually not seen in infants). Seizures and altered mental status may be present.

The recommended empirical antibiotic regimen in a suspected case of meningitis outside the neonatal period is vancomycin combined with a third-generation cephalosporin. Patients allergic to penicillin or cephalosporin can be treated with meropenem, fluoroquinolones, or chloramphenicol.

Chemoprophylaxis with rifampicin/ceftriaxone/ciprofloxacin is recommended for all close contacts with meningococcal meningitis regardless of age or immunization status. Rifampicin prophylaxis should be given to all household contacts of patients with invasive disease caused by Haemophilus influenzae type b, if an immunocompromised child of any age resides in the household or close family member younger than 48 months has not been fully immunized.

Solution to Question 3:

During neonatal resuscitation, the mouth is suctioned first and then the nares, in order to decrease the risk of aspiration. A 12 or 14 French suction catheter is used, with the suction pressure kept between 80-100mmHg.

Routine suctioning of the trachea during neonatal resuscitation is not recommended. The suctioning of the esophagus and stomach is avoided if not indicated specifically, as it can produce a vagal response due to stimulation in the posterior pharynx, resulting in apnea and/or bradycardia.

The steps in neonatal resuscitation follow the ABCs, Airway, Breathing, and Circulation.

The neonatal resuscitation algorithm is given below:



Chest compression should be initiated over the lower third of the sternum at a rate of 90/minutes. The ratio of chest compression to ventilation is 3: 1 (90 compressions: 30 breaths). The thumb technique (preferred) and 2 finger technique are the two different techniques used for performing chest compression. Umbilical vein cannulation is the preferred method for the administration of medications.

Solution to Question 4:

The given clinical scenario describes an adolescent girl with myoclonic jerks (a history of dropping objects) with preserved consciousness. This, along with abnormal EEG findings strongly suggests juvenile myoclonic epilepsy.

Juvenile myoclonic epilepsy (Janz syndrome) is the most common generalized epilepsy in young adults. It starts in early adolescence with one or more of the following features: bilateral myoclonic jerks in the morning, causing patients to drop things, generalized tonic-clonic or

clonic-tonic-clonic seizures upon awakening, and juvenile absence seizure.

Consciousness is usually preserved. Precipitating factors include sleep deprivation, alcohol, and photic stimulation. EEG shows 4 to 5Hz polyspike and slow wave discharges. A family history of epilepsy is often present.

The treatment options for JME include sodium valproate, levetiracetam, and topiramate. Benzodiazepine, lamotrigine, perampanel, phenobarbital, primidone, and zonisamide can be considered as add-on drugs. Complete remission is uncommon and has a high risk of seizure recurrence, hence antiepileptics are continued lifelong in a majority of the patients.

Other options:

Option B: Atypical absence seizures have been associated with myoclonic components and head drop. They are precipitated by drowsiness and EEG shows 1 to 2 Hz spike and slow wave discharges.

Option D: In benign childhood epilepsy with centrotemporal spikes, the child typically wakes at night due to a focal seizure with preserved awareness. Buccal and throat tingling and tonic or clonic contractions on one side of the face are seen, with drooling and preserved consciousness. EEG shows typical wide-based centrotemporal spikes.

Solution to Question 5:

Neonate with continuous urine dribbling a protruding bladder with exposed mucosa suggests bladder exstrophy.

In bladder exstrophy, the bladder protrudes from the abdominal wall, and its mucosa is exposed. The umbilicus is displaced downwards, the pubic rami are widely separated and the rectus muscles are separated. Anomalies of the bladder occur when the mesoderm fails to invade the cephalad extension of the cloacal membrane, the extent of this failure determines the degree of the anomaly.

Males commonly have complete epispadias with dorsal chordee, undescended testes, short penis, and inguinal hernia. Epispadias with separation of the two halves of the clitoris and wide separation of labia is seen in females. Anus is displaced anteriorly in both sexes. Rectal prolapse may be seen.

Management involves a staged surgical reconstruction of the bladder and genitalia. Untreated patients may develop total urinary incontinence and an increased incidence of bladder adenocarcinoma.

Other exstrophy anomalies are cloacal exstrophy and epispadias. Cloacal exstrophy (ileocaecal exstrophy) is a rare condition, in which there is a small exomphalos with an everted caecum and ileum separating halves of the bladder and in males, a split penis. The hindgut is truncated (giving tusks on the face of the elephant appearance).



Other options:

Option A: Gastroschisis is a condition where the intestinal loops herniate through the defect in the abdominal wall (shown below). The defect usually lies on one side of the umbilicus (right side commonly). The loops of the bowel are not covered by amnion as they herniate through the abdominal wall directly into the amniotic cavity. It is associated with elevated alpha-fetoprotein levels.



Option D: Mild forms of the imperforate anus are often called anal stenosis or an anterior ectopic anus.

Solution to Question 6:

Feeding by spoon is a social milestone that typically appears by 18 months. If this is absent, then the child should be termed as having a developmental delay. Hopping on one leg, drawing a square, and passing a ball to someone appear by 48 months of age.

Red flag signs in developmental screening and surveillance are indicators that suggest a serious disorder in a child's development, requiring referral to a pediatrician. These are given below:

Positive indicators: Presence of any of the following

• Loss of developmental skills at any age

• Concerns about vision or confirmed visual impairment

• Hearing loss at any age

• Persistently low muscle tone

• Lack of speech by 18 months without other means of communication

• Asymmetry of movements or features suggestive of cerebral palsy

• Persistent toe walking

• Complex disabilities

• Unusual head circumference compared to norms

• Clinician uncertainty about the developmental disorder.

Negative Indicators:

• Inability to sit unsupported by 12 months

• Inability to walk by 18 months in boys or 2 years in girls.

• Inability to walk normally

• Inability to run by 2.5 years

• Inability to hold objects in hand by 5 months or reach for them by 6 months.

• Inability to point at objects to share interest by 2 years

Solution to Question 7:

The clinical presentation of a child with a painful and growing parietal swelling, coupled with a history of a prior parietal bone fracture one year ago, strongly suggests the diagnosis of a growing fracture, also known as traumatic encephalocele or leptomeningeal cysts.

Skull fractures associated with a dural tear may fail to heal properly. The herniated brain, dilated ventricles, or a leptomeningeal cyst may cause an enlargement of this tear. These fractures are called growing skull fractures. They are usually seen in children < 3 years of age.

They present as a localized swelling or palpable skull defect that increases in size, most commonly seen in the parietal region. However, they may go undetected for years and present with delayed onset of neurologic symptoms (headache, seizure, or neurologic deficit).

Other options:

Option A: The extended duration of one year makes scalp hematoma unlikely, as it usually resolves within a few months.

Option C: Subdural hygromas are characterized by fluid accumulation in the subdural space. Most patients with subdural hygromas are asymptomatic, but uncommon symptoms may include headaches, changes in mental status, nausea and vomiting, focal neurological deficits, and seizures. They don't present with scalp swelling.

Option D: Chronic abscess may show inflammatory changes like erythema, pus, sinus, and systemic features like fever.

Solution to Question 8:

In an infant of a diabetic mother with normal blood glucose levels, the next investigation to consider is serum calcium due to the high risk of developing hypocalcemia. Hypocalcemia in these infants is attributed to the delayed response of the parathyroid hormone system.

Hypocalcemia is defined as a total serum calcium concentration of less than 8 mg/dL for a term baby (ionized calcium below 4mg/dL) and less than 7 mg/dL for a preterm baby. Asymptomatic neonates can be managed conservatively with early nutrition. Symptomatic neonates should receive iv or oral calcium replacement.

Maternal hyperglycemia causes fetal hyperglycemia, and the fetal pancreatic response leads to fetal hyperinsulinemia, which causes neonatal hypoglycemia. Neonatal hypoglycemia is defined as blood glucose <40 mg/dL (plasma glucose <45 mg/dL). The infant tends to be jittery, tremulous, hyperexcitable, hypotonic, lethargic, and poor sucking during the 1st three days after birth. These symptoms can also be related to hypocalcemia and hypomagnesemia which occur in the 1st 24 to 72 hours of life.

Respiratory distress syndrome, macrosomia, hyperbilirubinemia, polycythemia, renal vein thrombosis, visceromegaly, heart failure, and persistent pulmonary hypertension are other features seen in infants of diabetic mothers.

Congenital anomalies are seen in infants of mothers with pregestational diabetes. Neural tube defects (Encephalocele, meningomyelocele, anencephaly), transposition of great vessels (TGA), ventricular septal defect (VSD), atrial septal defect (ASD), hypoplastic left heart, aortic stenosis and coarctation of the aorta are commonly seen.

Caudal regression syndrome is rare, but a specific anomaly seen in infants of diabetic mothers. Small left colon syndrome is a rare anomaly that develops in the second and third trimesters.

Solution to Question 9:

Decerebrate or decorticate posturing is not seen in brain death as they are motor responses to noxious stimuli that originate from the midbrain and cortex.

Brain death, also known as death by neurologic criteria refers to the irreversible and total cessation of all functions of the entire brain, including the brainstem.

The three components for diagnosing brain death are a demonstration of coexisting irreversible coma with a known cause, absence of brain stem reflexes, and apnea. Coma requires the patient to be unresponsive, even to noxious stimuli. The diagnosis of brain death can still be made despite positive spinal reflexes. (option B)

Brainstem reflexes (pupillary responses, oculocephalic reflex, oculovestibular responses, corneal reflex, jaw reflex, facial grimacing, pharyngeal gag reflex, coughing) are absent. Apnea is the absence of respiratory drive at $PaCO_2$ of ≥60mmHg and > 20mmHg above baseline, which is consistent with brain death.

The following manifestations are occasionally seen and should not be misinterpreted as evidence of brainstem function:

• Normal systolic blood pressure - SBP >100 mmHg (option A)

• Movements originating from the spinal cord or peripheral nerve may occur

• Sweating, flushing, tachycardia (option C)

• Deep tendon reflexes; superficial abdominal reflexes; triple flexion response

• Babinski reflex

• Spontaneous movements of limbs other than pathologic flexion or extension response

• Respiratory-like movements (shoulder elevation and adduction, back-arching, intercostal expansion without significant tidal volumes)

Solution to Question 10:

Craniopagus is a type of conjoint twin where there is a fusion of the head or cranium.

Foramen magnum or the skull base never fuses in the case of craniopagus. They usually will have fused skulls and meninges. A shared cerebral cortex may occasionally be found.

Craniopagus has an incidence of 5%, and separation of the twins has a poor prognosis.

Dizygotic/fraternal twins result from the fertilization of two separate ova. Monozygotic/identical twins arise from the division of a single fertilized ovum. The outcome of the monozygotic twinning process depends on when division occurs.

Types of conjoined twins:

• Thoracopagus: united at the thorax

• Craniopagus: united at the head

• Cephalothoracopagus: fusion of both head and thorax

• Omphalopagous or xiphipagous: united anteriorly from the xiphoid to the umbilicus

• Parapagus: lateral fusion with two heads, sharing a single thorax, abdomen, and pelvis

• Ischiopagus: united at the pelvis

• Pygopagus: united in the sacral region

• Heteropagous: asymmetric forms

The image below shows the types of conjoined twins:

Monozygotic types	Time of zygote division
Dichorionic diamniotic	<72 hours
Monochorionic diamniotic(more common)	4 to 8 days
Monochorionic monoamniotic	8 to 12 days
Conjoined twins	>12 days



Common forms of conjoined twins: (A) craniopagus, (B) thoracopagus, (C) omphalopagus, (D) parapagus, (E) ischiopagus, and (F) pygopagus.

Solution to Question 11:

The major steroid products of the fetal adrenal gland are DHEA (dehydroepiandrosterone) and DHEA-S.

They serve as important substrates for estrogen synthesis in the placenta.

The fetal-adrenal-placental axis is vital for normal childbirth. Placental corticotropin-releasing hormone (CRH) stimulates the fetal adrenal gland to produce DHEA-S and cortisol. Cortisol further stimulates the production of placental CRH, creating a positive feedback loop that enhances adrenal steroid hormone production.

The steroid products released by the fetal adrenal gland have effects on the placenta and membranes. These effects contribute to transforming the myometrium from a quiescent to a contractile state.

Congenital adrenal hyperplasia is an autosomal recessive condition in which cortisol synthesis in the adrenal gland is affected. Cortisol deficiency increases the secretion of ACTH, which leads to adrenocortical hyperplasia and overproduction of intermediate metabolites.



Solution to Question 12:

The video clip of a 2-day-old neonate exhibiting repetitive, rhythmic movement in unilateral upper and lower limbs suggests a neonatal seizure of focal clonic type.

A focal clonic seizure is characterized by multiple, repetitive, rhythmic contractions of muscle groups of the limbs, face, or trunk, maybe unifocal or multifocal. It may occur synchronously or asynchronously in muscle groups on one side of the body. It may occur simultaneously but asynchronously on both sides. It cannot be suppressed by restraint.

Other options:

Option A: Focal tonic seizures are characterized by sustained posturing of a single limb, sustained asymmetrical posturing of the trunk, and sustained eye deviation. They cannot be provoked by stimulation or suppressed by restraint.

Option C: Spasms are sudden generalized jerks lasting 1-2 sec and are usually associated with a single, very brief, generalized discharge. They may be flexor, extensor, or mixed extensor/flexor and can occur in clusters. They too cannot be provoked by stimulation or suppressed by restraint.

Option D: Subtle seizures include transient eye deviations, nystagmus, blinking, mouthing, abnormal extremity movements (rowing, swimming, cycling, pedalling), fluctuations in heart rate, hypertension episodes, and apnea. They are more common in preterm than in term babies.

Myoclonic seizures are random, single, rapid muscle contractions of the limbs, face, or trunk. They are not repetitive or may recur at a slow rate. They can be generalized, focal, or fragmentary. They can be provoked by stimulation.

Clinically apparent seizures in a neonate should be treated if they last >3 minutes or are brief serial seizures. In all neonates with seizures, hypoglycemia, CNS infection, and hypocalcemia should be ruled out before starting anti-epileptic drug treatment.

Causes of neonatal seizure according to the age of presentation:

1 to 4 days	4 to 14 days	2 to 8 weeks
Hypoxic ischemic encephalopathyMaternal use of narcotics or barbituratesDrug toxicity: lidocaine, penicillinIntraventricular hemorrhageHypocalcemia, sepsis, hypoglycemia, hypomagnesemia, hypo or hypernatremiaGalactosemia, urea cycle disorders	Meningitis (bacterial)Encephalitis (enteroviral, herpes simplex)Hypocalcemia, hypoglycemiaGalactosemia, fructosemia,Hyperinsulinism, hyperammonemia syndrome	Herpes simplex or enteroviral encephalitis, bacterial meningitisSubdural hematoma, child abuseAminoaciduria, urea cycle defectsMalformations of cortical developmentTuberous sclerosisSturge-Weber syndrome

Solution to Question 13:

The given clinical scenario of a child with coarse facies, macroglossia, thick lips, hepatosplenomegaly, and copious discharge from the nose is suggestive of Hurler's disease.

It is the severe form of mucopolysaccharidoses type 1 (MPS 1), an autosomal recessive disorder caused by a defect in α-L iduronidase, leading to the accumulation of heparan sulfate and dermatan sulfate. The child usually presents between 6 and 24 months with hepatosplenomegaly, coarse facial features (shown below), corneal clouding, large tongue, prominent forehead, joint stiffness, short stature, and skeletal dysplasia.

Most patients have recurrent upper respiratory tract and ear infections, noisy breathing, and persistent copious nasal discharge. Radiograph shows skeletal dysplasia, called dysostosis multiplex - thick ribs, ovoid vertebral bodies, enlarged and trabeculated diaphyses of the long bones with irregular metaphysis and epiphysis.

Macrocephaly, enlarged J-shaped sella, premature closure of lambdoid sagittal sutures, and shallow orbits are seen with disease progression. Obstructive airway disease, cardiac complications, and respiratory infections are the common causes of death.

The image below shows the radiograph shows enlarged diaphyses of the metacarpal and phalanges.



Other options:

Option B: Beckwith-Wiedemann syndrome is an overgrowth syndrome characterized by macrosomia, macroglossia, hemihypertrophy, omphalocele, and renal anomalies. It is associated with an increased risk of Wilms tumor, hepatoblastoma, rhabdomyosarcoma, neuroblastoma, and adrenal cortical carcinoma.

Option C: Proteus syndrome is a disturbance of cellular growth involving ectodermal and mesodermal tissues, caused by mutation in the AKT1 gene. They present with hemimegalencephaly, asymmetric overgrowth of the extremities, verrucous cutaneous lesions, thickening of the bone, and excessive growth of muscles.

Option D: A child with untreated congenital hypothyroidism may have stunted growth, short extremities, narrow palpebral fissures, swollen eyelids, depressed nasal bridge, thick broad protruded tongue, and short and thick neck, dry and scaly skin.

Solution to Question 14:

The above pedigree chart denotes an autosomal recessive pattern. In the above pedigree chart, both males and females are carriers (equally affected) which indicates an autosomal pattern. Parents are unaffected (both are carriers/heterozygotes), and no affected members in other generations. 25% of the children are normal, 50% are carriers, and only 25% are affected. This indicates a a recessive pattern.

Pedigree charts in various modes of inheritance:

Solution to Question 15:

The Silverman-Anderson score of the child would be 8.

Calculation of the Silverman-Anderson score of this child:

Total score: 8

A score of more than 7 indicates respiratory failure.

The Silverman-Anderson score is a commonly used score to assess respiratory distress. It helps to assess the progression of distress over time and initiate treatment. It is more suited for preterms with respiratory distress syndrome. Scoring is to be done every half an hour and charting is carried out to monitor progress.

The image below shows Silverman Anderson retraction score:

Criteria	Finding	Score
Expiratory grunt	Audible with the naked ear	2
Nasal flaring	Marked	2

Criteria	Finding	Score
Xiphoid retraction	Marked	2
Intercostal retraction (lower chest)	Minimal	1
Chest movement (upper chest)	Lag	1



Downes' Score is also used for assessing the severity of respiratory distress. It is more comprehensive and can be applied to any gestational age and condition. Downe's score of >7 means impending respiratory failure.

Score	0	1	2
Respiratory rate	<60/min	60–80/min	>80/min
Air entry	Normal	Mild decrease	Marked decrease
Cyanosis	Nil	In-room air	On >40% O2
Grunting	None	Only on auscultation	Audible with the naked ear
Retraction	Nil	Mild	Moderate

Solution to Question 16:

Dysrhythmias in the pediatric age group are most often the result of respiratory insufficiency or hypoxia and not of primary cardiac causes, as in adults.

Hypoxia, hypercapnia, and acidosis lead to bradycardia, hypotension, and secondary cardiac arrest in children.

Other options:

Option A: Supraventricular tachycardia is the most common tachydysrhythmia that necessitates treatment in pediatric patients. Ventricular dysrhythmia is an uncommon presenting rhythm in infants and children.

Option B: The pre-arrest phase is the period that precedes the period of cardiac arrest.

Four phases of cardiac arrest have been described as follows:

• The prearrest phase- a period that precedes the period of cardiac arrest.

• The no-flow phase- reflects untreated cardiac arrest before it is recognized by a bystander in the community or by a medical provider in the hospital

• The low-flow phase- begins with the onset of cardiopulmonary resuscitation (CPR)

• The post-resuscitation phase - The post-resuscitation phase begins with the return of spontaneous circulation (ROSC).

Option C: The chest compression to ventilation ratio should be 30:2 or 15:2. Excessive ventilation results in decreased venous return into the chest, decreased coronary and cerebral blood flow, diminished cardiac output, and ultimately a decreased likelihood of return of spontaneous circulation.

Guidelines for pediatric basic life support:

Maneuver	Neonate	Infant <1 year	Child 1 year to puberty
Rescue breath	30-60/minute (every 1 to 2 seconds)	12-20/minute (every 3 seconds)	12-20/minute (every 3 seconds)
Pulse check site	Umbilical	Brachial	Carotid/femoral
Chest compression site	one finger below the intermammary line	One finger below the intermammary line or lower half of the sternum	The lower half of the sternum
Compression method	Two fingers or two thumb	Two fingers or two thumb	The heel of one hand or two hands
Compression depth	One-third of the chest	One-third to one-half of the chest	One-third to one-half of the chest
Compression rate	120/minute	100/minute	100/minute
Compression to ventilation ratio	3: 1	15: 2 (single rescuer - 30: 2)	15: 2 (single rescuer - 30: 2)


Chest compressions in pediatric patients:
Two-thumb technique
Two-finger technique
1 Chest compression in infants
2 Chest compression in children from 1–8 years of age
3 Chest compression in children > 8 years of age
MARROW

Paediatrics AIIMS 2019 (May & Nov)

Question 1:

Which of the following is a component of the neuromuscular criteria for maturity according to the expanded New Ballard score?

a) Heel-knee test

b) Finger-nose test

c) Scarf sign

d) Cubital angle

Question 2:

An 18-month-old baby was brought to the hospital with complaints of poor feeding and abnormal twitching of his lower limb. On examination, he has fever with fontanelle bulging and the child was lethargic. The most probable cause is:

a) Febrile seizures

b) Meningitis

c) Pseudotumor cerebri

d) Intracranial haemorrhage

Question 3:

A child presents with brown-colored urine and oliguria for the last 3 days. He has mild facial and pedal edema. His blood pressure is 126/90. He has 2+ proteinuria with >100 red cells/HPF and a few granular casts. His creatinine is 0.9, urea is 56. What is the likely diagnosis?

a) PSGN

b) Nephrolithiasis

c) Posterior Urethral Valve

d) Minimal change disease

Question 4:

Sign of effective neonatal resuscitation is:

a) Color change

b) Bilateral Air entry

c) Increased heart rate

d) None of the above

Question 5:

Gene involved in juvenile myoclonic epilepsy:

a) CHYNN-1

b) GABRA-1

c) FMR-1

d) All of the above

Question 6:

A 3-week neonate presents with ambiguous genitalia with Na+ - 127; K+ - 6 meq/L with a BP of 50/22 mm Hg. Along with IV fluids, what specific management is done?

a) Spironolactone

b) Potassium binding resins

c) Hydrocortisone

d) Broad spectrum antibiotics

Question 7:

A 6 years old child with developmental delay, can ride a tricycle, can climb upstairs with alternate feet, can say their name, and knows own sex, but cannot narrate a story. What is the developmental age?

a) 3 years

b) 4 years

c) 5 years

d) 2 years

Question 8:

The procedure displayed below is:

a) Intubation

b) Nasogastric tube insertion

c) Orogastric tube Insertion

d) Oropharyngeal airway insertion

Question 9:

An infant in the NICU was observed to have the features as seen in the video below. In addition to these features, there is an audible grunt. What would the Silverman Anderson score be?

a) 8

b) 6

c) 5

d) 4

Question 10:

Which of the following denotes an "infant at risk"?

a) Infant of a working mother

b) Infant of mother who has not taken folic acid during pregnancy

c) Infant with history of premature birth at <32 weeks gestation

d) Infant of mother with pre-eclampsia

Question 11:

A 7-year-old child presents with leg pain. She had fever and cough 3 days prior. Currently, there is pain on palpation of the calf muscles. Reflexes and other clinical examinations are normal. Lab test reveals CK levels of 2000 IU. What is the likely diagnosis?

a) Viral Myositis

b) Duchenne

c) GBS

d) Dermatomyositis

Question 12:

An 8-year-old boy with sensorineural hearing loss and eye defects presents to the OPD with hematuria. His renal biopsy on light microscopy was normal. Family history reveals the maternal uncle had Gr 5 CKD an is undergoing dialysis. Which of the following would be the probable diagnosis?

a) Alport syndrome

b) Goodpasture syndrome

c) IgA nephropathy

d) PSGN

Question 13:

The following device shown attached to the baby is used in all except :

a) In CPR

b) To know about apnea

c) In neonatal resuscitation

d) Measure central cyanosis

Answer Key

Question No.	Correct Option
1	c
2	b
3	a
4	c
5	b
6	c
7	a
8	c
9	a
10	c
11	a
12	a
13	d

Detailed Explanations

Solution to Question 1:

Scarf sign is a component of the neuromuscular criteria for maturity according to the expanded new Ballard score.

Scarf sign: If the infant's arm is pulled medially across the chest (as shown in the image below),

• In a hypotonic infant, the elbow will cross the midline.

• In a term infant with a normal tone, the elbow will not reach the midline.

Neuromuscular criteria for maturity (according to the expanded New Ballard score) include:

• Posture

• Square window (wrist)

• Arm recoil

• Popliteal angle

• Scarf sign

• Heel to ear

Physical criteria for maturity (according to the expanded New Ballard score) include:

• Skin

• Lanugo

• Eye/ear

• Plantar crease

• Breast

• Genitals

The expanded new Ballard score is used to assess the physical and neuromuscular maturity of the newborn by scoring six criteria and thereby calculating the gestational age. It is accurate between 20 to 44 weeks.

The lowest score on the expanded New Ballard Score is -10, which corresponds to 20 weeks gestational age, and the highest score is 50, which corresponds to 44 weeks gestational age. An increase in the score by 5 points corresponds to an increase of 2 weeks gestational age

Given in the image below is the scoring system for the physical and neuromuscular criteria:



Neuromuscular Maturity

Score	−1	0	1	2	3	4	5
Posture							
Square window (wrist)	> 90°	90°	60°	45°	30°	0°	
Arm recoil		180°	140–180°	110–140°	90–110°	< 90°	
Popliteal angle	180°	160°	140°	120°	100°	90°	< 90°
Scarf sign							
Heel to ear							

Physical Maturity

Skin	Sticky, friable, transparent	Gelatinous, red, translucent	Smooth, pink, visible veins	Superficial peeling and/or rash; few veins	Cracking, pale areas; rare veins	Parchment, deep cracking; no vessels	Leathery, cracked, wrinkled
Lanugo	None	Sparse	Abundant	Thinning	Bald areas	Mostly bald	
Plantar surface	Heel-toe 40–50 mm: −1 < 40 mm: −2	> 50 mm, no crease	Faint red marks	Anterior transverse crease only	Creases anterior 2/3	Creases over entire sole	
Breast	Imperceptible	Barely perceptible	Flat areola, no bud	Stippled areola, 1–2 mm bud	Raised areola, 3–4 mm bud	Full areola, 5–10 mm bud	
Eye/Ear	Lids fused loosely: −1 tightly: −2	Lids open; pinna flat, stays folded	Slightly curved pinna, soft, slow recoil	Well curved pinna, soft but ready recoil	Formed and firm, instant recoil	Thick cartilage, ear stiff	
Genitals (male)	Scrotum flat, smooth	Scrotum empty, faint rugae	Testes in upper canal, rare rugae	Testes descending, few rugae	Testes down, good rugae	Testes pendulous, deep rugae	
Genitals (female)	Clitoris prominent, labia flat	Clitoris prominent, small labia minora	Clitoris prominent, enlarging minora	Majora and minora equally prominent	Majora large, minora small	Majora cover clitoris and minora	

Maturity Rating

Score	Weeks
−10	20
−5	22
0	24
5	26
10	28
15	30
20	32
25	34
30	36
35	38
40	40
45	42
50	44

Solution to Question 2:

The clinical features of poor feeding, fever, seizures (abnormal lower limb twitching), bulging fontanelle (due to raised ICP), and lethargy are all suggestive of meningitis.

Febrile seizures (Option A) are diagnosed in the absence of any other features suggestive of CNS infection.

Pseudotumor cerebri (Option C) is rarely seen under the age of 10 years. There is an association with obesity, female gender, pregnancy, and drugs such as tetracycline, vitamin A, and retinoid derivatives. It has a chronic course of illness, with papilledema being present in a majority of cases.

Intracranial hemorrhage (Option D) will present with seizures, focal neurological deficits, altered tone, and altered consciousness.

Bacterial meningitis may be caused due to Neisseria meningitidis, Streptococcus pneumoniae, or Haemophilus influenzae type b.

Non-specific symptoms and signs include fever, rash, lethargy, refusal to feed, vomiting, etc. Specific symptoms and signs include photophobia, nuchal rigidity, bulging of the fontanelle, and clinical signs of meningeal irritation (Kernig and Brudzinski signs), but this is not usually seen in infants. Seizures and focal neurological signs may be present.

The diagnosis of meningitis is confirmed by CSF analysis via lumbar puncture - which typically shows micro-organisms on gram stain and culture, neutrophilic pleocytosis, increased protein, and decreased glucose concentration. Blood cultures should be done in all cases of suspected meningitis.

Acute CNS complications include seizures, increased ICP, cranial nerve palsies, stroke, cerebral or cerebellar herniation, and thrombosis of the dural venous sinuses. Subdural effusions and non-communicating hydrocephalus may develop.

Solution to Question 3:

A child presenting with acute onset (3 days) gross hematuria, hypertension, edema, oliguria, and mild to moderate proteinuria is suggestive of a nephritic syndrome picture. The only disease which causes this out of the given options is post-streptococcal glomerulonephritis (PSGN).

PSGN is primarily caused by Group A β-hemolytic streptococcal infections in children aged 5 to 12 years. It is caused by immune complexes containing streptococcal antigens (M protein of cell wall) and specific antibodies. PSGN typically manifests as an acute nephritic syndrome, appearing either 1 to 2 weeks after streptococcal pharyngitis or 3 to 6 weeks after streptococcal pyoderma. The severity of kidney involvement can vary, ranging from asymptomatic microscopic hematuria with normal renal function to gross hematuria with acute renal failure. Edema, hypertension, and oliguria may be present depending on the extent of renal involvement.

Laboratory findings commonly include a significant decrease in serum C3 levels during the acute phase, with values returning to normal within 6 to 8 weeks. To confirm the diagnosis, evidence of a preceding streptococcal infection is necessary. The antistreptolysin O (ASO) titer often increases following pharyngitis, while the anti-deoxyribonuclease B level is a reliable indicator of cutaneous streptococcal infection.

Light microscopic examination reveals enlarged, hypercellular glomeruli with diffuse mesangial and endothelial proliferation, as well as leukocyte infiltration. Immunofluorescence analysis demonstrates granular deposits of IgG and complement C3 along the glomerular basement membrane (GBM) and in the mesangium. Electron microscopy primarily shows subepithelial humps, with subendothelial deposits observed in early stages of the disease.

Managing the acute effects of renal insufficiency (sodium restriction and diuretics) and hypertension is the mainstay. Antibiotic therapy with a 10-day course of penicillin is recommended to limit the spread of nephritogenic pathogens. But antibiotics do not alter the natural course of PSGN. Complete resolution of the hematuria and proteinuria in the majority of children occurs within 3–6 weeks of the onset of nephritis but 3–10% of children may have persistent microscopic hematuria, non-nephrotic proteinuria, or hypertension.

Below is the image showing light microscopy findings of post-streptococcal glomerulonephritis:



Solution to Question 4:

An increase in the heart rate is the best sign of effective neonatal resuscitation. The response to neonatal resuscitation at various steps is assessed based on the heart rate.

As soon as a baby is born, the tone, cry, breathing efforts, and respiratory efforts are assessed. If the tone is poor, or if the baby is not breathing, the following measures are done immediately - warming to maintain the normal temperature, proper positioning of the baby's airway, i.e. sniffing position with the head slightly extended, clearing of airway secretions, and tactile stimulation.

After the initial steps of neonatal resuscitation, the heart rate of the baby must be assessed. If the heart rate is < 100 beats per minute (bpm), positive pressure ventilation is provided along with SPO2 monitoring to ensure adequate ventilation.

If the heart rate remains at < 100 bpm, ventilation corrective steps and/or endotracheal intubation have to be considered. The ventilation corrective steps are as follows and can be remembered with the mnemonic MRSOPA.

• Mask has to be reapplied

• Reposition head

• Suction the secretions

• Open mouth slightly and ventilate

• Pressure is increased slightly

• Airway may be changed

If the heart rate is < 60 bpm, chest compressions are initiated over the lower third of the sternum at a rate of 90 compressions/min. The ratio of compressions to ventilation is 3: 1 (90 compressions: 30 breaths).

If the heart rate still remains <60 bpm, despite effective compressions and ventilation, then epinephrine (1:10,000 solution at 0.1–0.3 mL/kg intravenously or 0.5–1 mL/kg intratracheally) should be administered. The dosage can be repeated every 3–5 minutes. Persistent bradycardia may signify inadequate ventilation or severe cardiac compromise.

The neonatal resuscitation algorithm is given below:



Solution to Question 5:

Mutation in GABRA-1 is associated with juvenile myoclonic epilepsy.

Juvenile myoclonic epilepsy (JME), also known as Janz syndrome or impulsive petit mal, is the most common generalized epilepsy in young adults. It may be caused due to mutation in many genes including CACNB4, CLNC2, GABRD, GABRA1, and Myoclonin1/ EFHC1

It typically begins between the ages of 12 and 18 (adolescence). The defining seizure type in JME is generalized myoclonic seizures, which occur in all patients. These seizures are typically mild and primarily affect the upper extremities in the morning causing the patient to drop things. However, they often go unnoticed and are only brought to medical attention after a generalized tonic-clonic seizure, which tends to occur after sleep deprivation or alcohol consumption. Some cases of childhood absence epilepsy may evolve into JME.

Diagnosis of JME relies on the patient's clinical history and electroencephalogram (EEG) findings. EEG typically shows bursts of generalized irregular 4- to 6-Hz spike-and-wave activity, particularly after awakening. JME is generally a lifelong condition.

Valproate is considered the most effective medication. Lamotrigine, levetiracetam, and topiramate are also approved for JME.

The characteristic EEG pattern is shown in the image below:



Solution to Question 6:

The given clinical case of a neonate presenting with ambiguous genitalia, hyponatremia, hyperkalemia, and hypotension points towards a diagnosis of congenital adrenal hyperplasia (CAH). Hydrocortisone must be administered along with IV fluids for the management of this baby.

The most common enzyme deficiency in CAH is 21-hydroxylase deficiency. Babies with 21-hydroxylase deficiency typically present around 10-14 days of age with the subset of symptoms due to aldosterone deficiency, cortisol deficiency, and androgen excess.

Aldosterone deficiency leads to salt-wasting crises marked by hyponatremia, hyperkalemia, hypotension, and metabolic acidosis.

Cortisol deficiency causes manifestations like hypoglycemia, hyperpigmentation, weight loss, weakness, anorexia, vomiting, and shock.

Androgen excess in 21-hydroxylase deficiency results in virilization of females, leading to female pseudohermaphroditism, and precocious puberty in males. Virilization is usually apparent at birth in females and within 2 to 3 years of life in males.

Without treatment, shock, cardiac arrhythmias, and death may occur within days or weeks.

Diagnosis is confirmed by elevated 17-hydroxyprogesterone levels or by genetic testing.

Glucocorticoid replacement therapy is the primary treatment for cortisol deficiency, aiming to suppress androgen production and prevent complications associated with androgen excess. Fludrocortisone is given to control mineralocorticoid deficiency.

If the condition is diagnosed prenatally, dexamethasone may be administered to the mother.

The image below summarizes the pathway involved in steroid hormonal production:



Solution to Question 7:

The developmental age of a child who can ride a tricycle, can climb upstairs with alternate feet, can say their name, and knows own sex is 3 years. Telling stories is a language milestone expected to be attained at 4 years of age.

The developmental milestones at 3 years of age are as follows:

Domain	Milestones
Motor	Rides tricycleStands momentarily on one footWalks upstairs with alternate foot
Fine Motor	Builds tower of 9 blocksCopies circle
Language	Knows name and sexAsks questions
Social	Shares toysKnows full name and gender

Solution to Question 8:

The procedure depicted is the insertion of an orogastric tube.

It is indicated in conditions like choanal atresia and cleft palate - where a nasogastric tube cannot be passed, or may cause inadvertant injury.

An orogastric tube is used to facilitate enteral rehydration and administer medications in children with dehydration or who are critically ill. It can also be used for gastric decompression in cases of

distention due to small intestinal atresia or stenosis.

Solution to Question 9:

In the given case, the baby shows a see-saw chest movement (2), minimal intercostal retraction (1), visible xiphoid retraction (1), marked nasal flaring (2) and there is an audible respiratory grunt (2). The Silverman-Anderson score of the baby would be 8 (2+1+1+2+2).

Given below is the Silverman-Anderson scoring system (as per 2022 IAP guidelines) to assess the severity of respiratory distress in a child:

Interpretation:

Score	Upper chest retraction(Chest movement)	Lower chest retraction(Intercostal retraction)	Xiphoid retraction	Nasal dilatation	Grunt
0	Synchronized	none	none	none	none
1	Lag on inspiration	Just visible	Just visible	Minimal	Stethoscope
2	See-saw	Marked	marked	marked	naked ear

Score	Inference
0	No distress
1-4	Mild respiratory distress
5-7	Moderate respiratory failure
>7	Severe or impending respiratory distress



Silverman Anderson retraction score is more suited for preterms with hyaline membrane disease.

Downes' Score is a more comprehensive scoring system and can be applied to any gestational age and condition. Scoring is to be done every half an hour and charting is carried out to monitor progress.

Downe's score of >6 means impending respiratory failure.

Score	0	1	2
Respiratory rate	<60/min	60–80/min	>80/min
Air entry	Normal	Mild decrease	Marked decrease
Cyanosis	Nil	In-room air	On >40% O2
Grunting	None	Only on auscultation	Audible with the naked ear
Retraction	Nil	Mild	Moderate

Solution to Question 10:

An infant with a history of premature birth at <32 weeks gestation will be considered a high-risk infant according to AIIMS protocols on Neonatology.

The inclusion criteria for high-risk neonates according to AIIMS protocol are:

1. Birth weight <1500 grams

2. Gestation age <32 weeks

3. Infants with a birth weight of ≥ 1500gm or gestation age ≥ 32 weeks with any of the following risk factors:

• Intrauterine growth centile <3rd centile

• Meningitis

• Received mechanical ventilation for 48 hours or more

• Hypoxic ischemic encephalopathy stage 2 or higher

• Major malformation

• Inborn error of metabolism/ chromosomal or genetic disorders/ intrauterine infections

• Symptomatic hypoglycemia

• Symptomatic polycythemia

• Retrovirus positive mother

• Hyperbilirubinemia requiring exchange transfusion OR Rh isoimmunization/cholestasis

• Abnormal neurological examination at discharge/ seizures

• Major morbidities such as chronic lung diseases, IVH grade III or more and periventricular leukomalacia

Note: Although Park's Textbook of Preventive and Social Medicine mentions infants of working mothers (Option A) as high-risk infants, premature infant with < 32 weeks gestation is chosen over it, as AIIMS protocols are used as the preferred reference for AIIMS exams.

Solution to Question 11:

The clinical picture showing pain in the calves occuring within a week of an upper respiratory tract infection (URI), with elevated creatine kinase levels suggests the diagnosis of viral myositis. The specific history of URI suggests influenza myositis.

Infectious myositis has a male predominance and is typically seen in young adults. Childhood form presents with fever, malaise, and rhinorrhea and is usually followed by severe pain, especially in the calves. The muscle pain is usually worse with movement and the symptoms last for about a week. In children, toe walking and wide-based gait may be seen because of the involvement of the gastrocnemius-soleus muscles.

Other viral causes of myositis include HIV-1 (one of the most common causes of myositis), group B coxsackievirus (epidemic myalgia), HTLV-1 (Human T-lymphotropic virus), and cytomegalovirus (CMV).

Lab features of influenza myositis include elevated CK which may be as high as 500 times normal the value. Urine myoglobin is usually positive. Detection of the virus with PCR testing of nasopharyngeal specimens can be carried out but is not of much clinical significance. The histopathological examination would reveal muscle fiber necrosis without much inflammatory change.

Treatment includes bed rest, IV fluids, and symptomatic management with analgesics and antipyretics. Amantadine and other antiviral agents may be considered in adults. Complications include myocarditis and respiratory dysfunction. Rhabdomyolysis is an uncommon complication.

Solution to Question 12:

The clinical presentation of a child with hematuria, sensorineural hearing loss, and eye defects with normal renal biopsy is consistent with Alport Syndrome. It is a hereditary nephritis, which explains the family history of chronic kidney disease.

Alport syndrome is a genetic disease caused by mutations in the COL4A5 gene, which codes for the alpha chain of type IV collagen. In approximately 85% of cases, the syndrome is X-linked.

In the early years (around 5 years), patients are often asymptomatic but may have persistent microscopic hematuria. Between 5 and 10 years of age, proteinuria, and hypertension may develop, which can be managed with angiotensin-converting enzyme (ACE) inhibitors. By 10 years of age, serum creatinine levels start to rise. Around 15 years, patients may experience ear, nose, and throat (ENT) manifestations, with high-frequency sensory neural hearing loss being a common feature in males. Ocular manifestations, such as anterior lenticonus (pathognomonic) and dot and fleck retinopathy (most common), typically appear around the age of 20. Anterior

lenticonus is particularly correlated with the development of end-stage renal disease (ESRD). By 25 to 30 years of age, most individuals with Alport's syndrome progress to ESRD.

Gross hematuria following upper respiratory tract infections is not typically seen in Alport's syndrome and should prompt consideration of other conditions, such as IgA nephropathy.

Skin immunofluorescence to demonstrate the lack of the α5(IV) collagen chain and genetic testing can be used for the diagnosis. Light microscopy and immunofluorescence (IF) on renal biopsy may be normal. Electron microscopy reveals irregular areas of thinning and thickening of the glomerular basement membrane (GBM), along with splitting of the lamina densa, giving it a basket-weave appearance.

The mainstay of management is the control of hypertension and ACE inhibitors to slow down the progression of renal damage. Although transplant recipients develop anti-GBM antibodies, graft survival is good.

Thin GBM disease is an autosomal dominant disease characterized by persistent microscopic hematuria but no progression to ESRD or extrarenal manifestations. EM can be used to confirm the diagnosis, showing GBM thickness of less than 200 nm.

Other options:

Option B: Goodpasture syndrome is an immune-mediated disease with antibodies against the alpha-3 chain of GBM type IV collagen; it is not hereditary and the light microscopy would reveal extra capillary proliferation with crescents & necrosis.

Option C: In IgA Nephropathy, light microscopy would reveal focal mesangial proliferative glomerulonephritis & mesangial widening.

Option D: In post-streptococcal glomerulonephritis (PSGN), there will be diffuse endocapillary proliferation and accompanying leukocytic infiltration seen in light microscopy.

Solution to Question 13:

The image shows a pulse oximeter. It cannot be reliably used to measure central cyanosis.

A pulse oximeter employs spectrophotometry to measure the percentage saturation of oxygen at a peripheral site. In cases of markedly deranged hemoglobin (Hb) levels and abnormal Hb, pulse oximetry doesn't correlate with central cyanosis, which depends on the absolute value of reduced Hb.

A pulse oximeter can be used for assessing response to resuscitation, titrating oxygen therapy in newborns, monitoring apnea (bradycardia and desaturation), confirming endotracheal tube placement, evaluating the effectiveness of bag and mask ventilation, and monitoring for hypoxia during suction and laryngoscopy.

Cyanosis is characterized by a bluish discoloration of the skin and mucous membranes due to elevated levels of reduced Hb (more than 4 g/dL) or Hb derivatives. It is examined in the nail beds and mucous membranes, earlobes, lips, and fingers. It can be classified into central and peripheral types.

Central cyanosis can be due to reduced oxygen saturation or the presence of abnormal Hb derivatives. Causes include high altitudes, impaired pulmonary function, anatomic shunts, and

abnormal Hb. Both the mucous membranes and skin will be affected.

The absolute amount of reduced Hb rather than its relative quantity is significant in cyanosis. In severe anemia, even with marked arterial desaturation, cyanosis may not be present. Conversely, polycythemia patients are prone to cyanosis even at higher oxygen saturation levels.

The presence of abnormal Hb species may lead to errors in SpO2 readings. In a patient with methemoglobinemia, the SpO2 will be 80% to 85%, irrespective of arterial saturation (SaO2). A patient with carbon monoxide poisoning will have a falsely elevated SpO2 level. A co-oximeter is a better device to measure abnormal hemoglobins.

Carbon monoxide poisoning does not lead to true cyanosis. It causes cherry red discoloration.

Peripheral cyanosis occurs due to decreased blood flow and excessive oxygen extraction from normally saturated arterial blood. The mucous membranes of the oral cavity and sublingual mucosa are spared. The causes include reduced cardiac output, cold exposure, and arterial and venous obstruction. Here, the degree of cyanosis will generally correspond to SpO2 readings. However, the SpO2 levels may be falsely low despite normal SaO2.

Paediatrics AIIMS 2020 (May and Nov INI-CET)

Question 1:

Which of the following is least likely regarding Rh incompatibility in a newborn?

a) Increased unconjugated bilirubin

b) Increased fecal urobilinogen

c) Increased urine bilirubin

d) Increased stercobilinogen

Question 2:

A 3-year-old toxic child presents with tachypnea, high-grade fever, inspiratory stridor, and developed difficulty in swallowing with drooling of saliva within 4-6 hours of fever. Along with airway management which of the following needs to be given?

a) Nebulisation with racemic adrenaline

b) IV ceftriaxone

c) Diphtheria antitoxin

d) Start steroids

Question 3:

A child transfers an object from one hand to another. What does it signify?

a) Visual motor co-ordination

b) Ability to compare objects

c) Ability to handle small objects

d) Voluntary release

Question 4:

Surgical correction of VSD is performed in an infant. This will help in relieving which of the following features?

a) Respiratory alkalosis

b) Failure to thrive

c) Arrythmia

d) Heart block

Question 5:

All the following PFT findings in a child are suggestive of asthma except?

a) Increase in FEV1> 12% after salbutamol inhalation

b) FEV1/FVC< 80%

c) Day- night variation of FEV1 > 15%

d) Decrease in FEV1>15% after exercise

Question 6:

Which of the following is not a symptom of severe dehydration?

a) Drowsy child

b) Skin pinch retracts slowly

c) Thirsty child

d) Sunken fontanelle

Question 7:

Which of the following is not a criteria for severe acute malnutrition?

a) Weight for height less than -3 SD

b) MUAC < 115mm

c) B/L Pedal Edema

d) Weight for age less than -3 SD

Question 8:

Phototherapy converts bilirubin to _____

a) Stercobilin

b) Lumirubin

c) Biliverdin

d) Urobilin

Question 9:

Identify the vitamin which when deficient can lead to neonatal seizures.

a) Pyridoxine

b) Thiamine

c) Riboflavin

d) Pantothenic acid

Question 10:

A baby assessed 5 minutes after birth is found to be pink, with blue extremities, and irregular respiration. Heart rate is 80 beats per minute. There is grimace and some flexion. What is the Apgar score for this baby?

a) 2

b) 3

c) 4

d) 5

Question 11:

A child suffering from steroid-dependent nephrotic syndrome is brought to the OPD for vaccination. Which of the following is true about vaccination in this child?

a) All live vaccines under NIS can be given as per schedule

b) All killed vaccines under NIS can be given

c) Only pneumococcal vaccine should be given to the child in acute phase

d) Live vaccines are contraindicated in the child's sibling

Question 12:

A young girl presents with the complaints of continuous dribbling of urine. However, she also voids normally. Her parents tell you that she had never been totally dry since her birth. What

is the most likely diagnosis?

a) Ectopic ureter

b) Ureterocele

c) Congenital megaureter

d) Vesicoureteric reflux

Question 13:

A 14-month-old child with Hb 6.3 was undergoing repeated blood transfusions. The last transfusion was given in the 9th month. Which of the following investigation is used for a definitive diagnosis?



a) High performance liquid chromatography

b) Ferritin levels estimation

c) Bone marrow aspiration

d) Globin gene sequencing

Answer Key

Question No.	Correct Option
1	c

2	b
3	b
4	b
5	c
6	c
7	d
8	b
9	a
10	d
11	b
12	a
13	d

Detailed Explanations

Solution to Question 1:

Amongst the given options, increased urine bilirubin is least likely with Rh incompatibility in newborns.

Rh incompatibility leads to hemolysis in the newborn, causing the release of large amounts of hemoglobin. This hemoglobin is converted to biliverdin and then to unconjugated bilirubin.

The newborn liver has a limited capacity to convert this large amount of unconjugated bilirubin to its soluble form through glucuronide conjugation. As a result, levels of unconjugated bilirubin increase (option A). Unconjugated bilirubin, being water-insoluble and bound to albumin, is not excreted in urine.

Intestinal bacteria convert unconjugated bilirubin to urobilinogen and stercobilinogen, which are excreted in urine and stool (options B and D).

Solution to Question 2:

The clinical presentation of a toxic-looking child with high-grade fever, tachypnea, inspiratory stridor, dysphagia, and salivary drooling strongly suggests acute epiglottitis. Once the airway is secured, empirical antimicrobial therapy with intravenous ceftriaxone is recommended in the management of acute epiglottitis.

The most common causative agents of epiglottitis in vaccinated children are Streptococcus pyogenes, Streptococcus pneumoniae, non-typeable Haemophilus influenzae, and Staphylococcus aureus. The incidence of Haemophilus influenzae type B has decreased significantly due to vaccination.

The child presents with a high fever, sore throat, dyspnea, dysphagia, and progressive respiratory obstruction. The child may exhibit a toxic appearance, with salivary drooling and assuming a tripod position. The presence of stridor indicates near-complete obstruction of the airway.

Laryngoscopy should be promptly conducted in a controlled setting, such as an operating room or intensive care unit when the diagnosis of epiglottitis is certain or likely based on clinical evaluation.

Acute epiglottitis is a medical emergency due to unpredictable clinical course and sudden airway obstruction. Immediate airway management through endotracheal or nasotracheal intubation or tracheostomy is crucial, regardless of the severity of respiratory distress. Empiric treatment with antibiotics like ceftriaxone, cefepime, or meropenem is recommended until culture and susceptibility results are available. Dexamethasone may also be administered to reduce pharyngeal edema.

Other options:

Option A: Nebulisation with racemic adrenaline is given in moderate to severe croup.

The image below shows the thumb sign in epiglottitis:



Solution to Question 3:

The milestone of transfer of objects from one hand to another in a child signifies the ability of the child to compare objects. The transfer of objects is a motor milestone occurring at 6 months of age.

Other options:

Option A: When the child reaches for objects it implies the onset of visuomotor coordination. It is attained at 4 months

Option C: Thumb finger grasp implies the ability to handle small objects. It is attained at 8 months.

Option D: The voluntary release/ Release phenomenon is used to denote the disappearance of the palmar grasp, which is seen at 4 months.

Solution to Question 4:

Surgery for ventricular septal defect (VSD) mainly helps to relieve congestive cardiac failure which manifests as a failure to thrive in an infant. The main aim of performing surgery in VSD is to get the symptoms of heart failure in control and to prevent the development of pulmonary vascular disease.

Ventricular septal defect (VSD) is the most common congenital heart disease. The most common subtype is the membranous type.

Small VSDs with trivial left-to-right shunts and normal pulmonary artery pressure (PAP) are the most common. These patients are asymptomatic. A loud, harsh, or blowing holosystolic murmur is present and heard best over the lower left sternal border, and it is accompanied by a thrill.

Small VSDs close spontaneously, especially within the first two years of life. Muscular VSDs have a higher closure rate (up to 80%) compared to membranous VSDs.

Large VSDs with excessive pulmonary blood flow and pulmonary hypertension can lead to signs of congestive heart failure in early infancy. These signs may include dyspnea, feeding difficulties, poor weight gain, profuse perspiration, irritability, weak crying, and recurrent pulmonary infections. The holosystolic murmur associated with a large VSD is less harsh and more blowing compared to that of a small VSD. This is due to the absence of a significant pressure gradient across the defect.

The indications of surgery in VSD are:

• Large defects in patients of any age with uncontrolled symptoms or failure to thrive.

• Moderate to large defects associated with pulmonary hypertension in infants between 6 and 12 months, even if symptoms are controlled by medication.

• Patients older than 24 months with a Qp: Qs ratio greater than 2:1 (ratio of total pulmonary blood flow-Qp to total systemic blood flow-Qs).

• Supracristal VSDs of any size, due to the higher risk of developing aortic valve regurgitation.

Severe pulmonary vascular disease nonresponsive to pulmonary vasodilators is a contraindication to VSD closure.

Other options:

Options A and C: Respiratory alkalosis and arrhythmia are other rare complications of VSD.

Options D: Heart block may occur perioperatively in VSD.

Solution to Question 5:

Day-night variation of FEV1 > 15% is not suggestive of asthma, variation ≥20% is suggestive of asthma.

Lung Function Abnormalities in Childhood Asthma

• Spirometry showing airflow limitation: Low FEV1 (relative to the percentage of predicted norms) and FEV1:FVC ratio <0.80 (option B)

• Bronchodilator response (to inhaled β-agonist): Improvement in FEV1 ≥12% and ≥200 mL after inhalation of a short-acting beta agonist (option A).

• Exercise challenge: Worsening in FEV1 ≥15% (option D)

• Daily peak expiratory flow (PEF) or FEV1 monitoring: day to day and/or A.M.-to-P.M. variation ≥20%

• Exhaled nitric oxide (FeNO): A value of >20ppb (parts per billion) supports the clinical diagnosis of asthma in children.

FEV1: Forced expiratory volume in 1 sec; FVC: Forced vital capacity.

Childhood asthma can be classified into two types: recurrent wheezing in early childhood, often triggered by respiratory viral infections and resolves during preschool/lower school years, and chronic asthma associated with allergies that persist into later childhood and adulthood.

Based on the severity, acute asthma is classified as mild, moderate, and severe.

Clinical parameter	Mild	Moderate	Severe
Color	Normal	Normal	Pale
Sensorium	Normal	Anxious	Agitated
Respiratory rate	Increased	Increased	Increased
Dyspnea	Absent	Moderate	Severe
Use of accessory muscles	Nil or minimal	Chest indrawing	Indrawing, nasal flare
Rhonchi	Expiratory and/or inspiratory	Expiratory and/or inspiratory	Expiratory or absent.
Oxygen saturation	>95%	90-95%	<90%
Speech	Sentence	Phrases	Difficulty in speech

Solution to Question 6:

A thirsty child signifies some dehydration, not severe dehydration.

Clinical features of severe dehydration in a child include:

• Drowsy, lethargic, or unconscious, with deeply sunken eyes and absent tears

• Sunken fontanelle

• Drinks poorly or not able to drink

• Deep breathing

• Skin pinch goes back very slowly (recoil in >2 seconds)

• Tachycardia, with bradycardia in most severe cases

• Weak, thready, or impalpable pulse

• Prolonged capillary refill

• Cold, mottled, and cyanotic extremities

• Minimal urine output

Solution to Question 7:

Severe acute malnutrition (SAM) according to the WHO criteria is the weight for height (not age) less than minus -3 SD.

Severe acute malnutrition (SAM) in children of ages 6 to 59 months is defined by the WHO as any of the following:

• Weight-for-length (or height) below –3 SD of the WHO child growth standards

• Presence of bipedal edema

• Mid-upper arm circumference (MUAC) <11.5 cm.

Management of SAM is given below:



Solution to Question 8:

Phototherapy converts bilirubin to lumirubin. It is used to manage clinical jaundice and indirect hyperbilirubinemia.

It involves exposing the patient to high-intensity light in the visible spectrum. Bilirubin absorbs light most effectively in the blue range (420 to 470nm).

Phototherapy converts toxic bilirubin into two forms: lumirubin, formed by irreversible structural isomerization that can be excreted by the kidneys, and a reversible isomer (4Z,15E-bilirubin), formed by photoisomerization that can be excreted in bile without conjugation.

The effectiveness of phototherapy depends on factors such as the appropriate wavelength range, distance to the infant, skin surface area exposure, and the infant's bilirubin metabolism. The infant's eyes should be covered to prevent corneal damage. Continuous conventional phototherapy is typically used, with frequent repositioning of the infant.

Phototherapy can be discontinued once two TSB (total serum bilirubin) values 12 hours apart fall below current age-specific cut-offs.

Complications of phototherapy may include loose stools, erythematous macular rash, purpuric rash (linked to transient porphyrinemia), overheating, dehydration (due to increased insensible water loss and diarrhea), and hypothermia from exposure.

Bronze baby syndrome is a dark grayish-brown skin discoloration seen in infants undergoing phototherapy in the presence of direct hyperbilirubinemia. Porphyria is a contraindication for phototherapy because photosensitivity and blistering can be severe in these infants.

Solution to Question 9:

Pyridoxine deficiency can cause refractory seizures in the neonatal period and infancy.

Pyridoxal 5'-phosphate (P5P) is the active form of pyridoxine and functions as a cofactor for enzymes involved in neurotransmitter synthesis, amino acid metabolism, and glycogen metabolism. Intracellular P5P deficiency in the brain can lead to a seizure disorder that does not respond to typical anticonvulsant medications but shows a response to pyridoxine.

Antiquitin deficiency is the most common cause of pyridoxine-dependent epilepsy (due to mutations in the ALDH7A1 alleles)

They primarily present with generalized seizures in the early days of life. The seizures can be of various types, recurrent focal motor seizures, generalized tonic seizures, and myoclonus. Additional manifestations may include dystonia, respiratory distress, abdominal distention with vomiting, hepatomegaly, hypoglycemia, and hypothermia.

Increased α-aminoadipic semialdehyde and pipecolic acid levels are observed in the CSF, plasma, and urine. Treatment with vitamin B6 (50-100 mg/day) shows significant improvement in seizures. Life-long therapy is needed.

Other options:

Option B: Thiamine (vitamin B1) deficiency causes beriberi.

Option C: Riboflavin (vitamin B2) deficiency causes glossitis, cheilosis, keratitis, and corneal vascularization.

Option D: Pantothenic acid (vitamin B5) deficiency causes burning feet syndrome.

Solution to Question 10:

The Apgar score for this baby will be 5.

• Color; body pink, extremities blue: score-1

• Heart rate; 80 beats per minute: score 1

• Grimace: Score-1

• Some flexion: Score-1

• Irregular respiration: Score-1

The Apgar score is a systematic way of assessing the infant after birth at 1 and 5 minutes of life. It gives a maximum score of 10 which indicates the newborn is in the best possible condition. Scores between 0-3 require immediate resuscitation.

Changes in the Apgar scores denote the response of the infant to resuscitation. If the score at 5 minutes remains less than 7, the infant should be assessed again every 5 minutes for the next 20 minutes.

Apgar Scoring System

Indicator		0 Points	1 Point	2 Points
A	Appearance (skin colour)	Blue/Pale	Pink body, blue extremeties	Pink
P	Pulse	Absent	< 100 BPM	> 100 BPM
G	Grimace (reflex irritability)	No response	Minimal response to stimulation	Cough/sneeze when stimulated
A	Activity (muscle tone)	Floppy	Some flexion	Well flexed
R	Respiratory efforts	Absent	Slow and irregular	Strong cry

©MARROW

Causes for false positive Apgar score are:

• Prematurity

• Drugs like analgesics, narcotics, sedatives, magnesium sulfate

• Acute cerebral trauma

• Precipitous delivery

• Airway obstruction (choanal atresia)

• Congenital pneumonia or sepsis

• Hemorrhage- hypovolemia

False-negative Apgar score is seen in maternal acidosis and high fetal catecholamine levels.

Solution to Question 11:

In a child with steroid-dependent nephrotic syndrome, all killed vaccines can be given.

Option A: Live virus vaccines are administered with caution in children receiving corticosteroid therapy. They can be given to patients in whom the prednisone dose is below 1mg/kg daily or 2mg/kg on alternate days. If the patient is receiving corticosteroid-sparing agents (cyclophosphamide, cyclosporine), live virus vaccines are contraindicated.

Option C: Only live vaccines are contraindicated in this child. Full pneumococcal vaccination (with the 13-valent conjugate vaccine and 23-valent polysaccharide vaccine) and influenza vaccination can be given annually to the child and their household contacts.

Option D: Immunization of siblings/healthy household contacts with live vaccines is necessary to reduce the risk of infection transmission to the immunosuppressed child. Direct exposure to the gastrointestinal/respiratory secretions of the household contacts is avoided for about 3-6 weeks after vaccination.

Minimal change nephrotic syndrome is the most common cause of nephrotic syndrome. It is characterized by heavy proteinuria, edema, hypoalbuminemia, and hyperlipidemia. The diagnosis is confirmed by performing a urinalysis and urine protein: creatinine ratio. Nephrotic range proteinuria is proteinuria greater than 3.5 g per 24 hours (3+ or 4+) or a urine protein : creatinine ratio exceeding 2.

The recommended initial treatment for nephrotic syndrome is a daily dose of 2mg/kg (max 60mg) of prednisolone for 6 weeks followed by an alternate dose of 1.5mg/kg (max 40mg) for the next 6-8 weeks.

Definitions regarding the course of nephrotic syndrome:

• Frequently relapsing nephrotic syndrome: ≥2 relapses in 6 months or 3 or more relapses in 12 months.

• Steroid-dependent nephrotic syndrome: Relapse during steroid tapering or within 2 weeks of discontinuation.

• Steroid resistance: Failure to achieve remission despite therapy with daily prednisolone for 6 weeks

Alternative drugs used in steroid-dependent patients, frequent relapsers, steroid-resistant patients, and those with severe corticosteroid toxicity are cyclophosphamide, calcineurin inhibitors: cyclosporine and Tacrolimus, mycophenolate, levamisole, and rituximab (a chimeric monoclonal antibody against CD20).

Solution to Question 12:

The clinical presentation of a young girl with continuous dribbling of urine, despite normal voiding, strongly suggests the diagnosis of an ectopic ureter.

An ectopic ureter refers to a condition where the ureter drains outside the bladder. It is more common in females and opens either into the urethra below the sphincter or into the vagina. Hence, they present with constant urinary dripping all day, even though the child voids regularly (paradoxical incontinence).

The terminal part of the ureter is frequently narrowed, leading to hydroureteronephrosis. Urinary tract infections (UTIs) are common due to urinary stasis.

In males, ectopic ureters usually enter the posterior urethra above the external sphincter, hence they are usually continent and most patients present with UTI or epididymitis.

Investigations include renal ultrasound (USG), voiding cystourethrogram (VCUG), and renal scan to assess function. USG shows a hydronephrotic kidney or dilated upper pole and ureter. If there is a satisfactory function, the recommended treatment options are ureteral reimplantation into the bladder or ureteroureterostomy. The poor function may require partial or total nephrectomy.

Other options:

Option B: A ureterocele is a cystic dilation of the terminal ureter, typically obstructive due to a small ureteral opening. In childhood, they commonly present with infections. Larger ureteroceles can cause obstruction at the bladder neck.

Option C: Congenital megaureter (dilated ureter) presents UTI, urinary stones, hematuria, and flank pain because of urinary stasis.

Option D: Vesicoureteral reflux (VUR) is characterized by the retrograde flow of urine from the bladder to the ureter and kidney. They usually present with recurrent attacks of UTI.

Solution to Question 13:

The given clinical scenario of repeated transfusion, low hemoglobin level, and target cells in the smear are suggestive of thalassemia. Globin gene sequencing is used for its definitive diagnosis.

High-performance liquid chromatography (HPLC) is used as a screening test for thalassemia (option A).

Thalassemia is a hemoglobinopathy that arises due to deficient synthesis of the globin chains. Depending on the chain affected, it is classified into α-thalassemia and β-thalassemia.

β-thalassemia is caused by point mutations of the β-thalassemia gene on chromosome 11, which diminish the synthesis of β-globin chains. Splicing mutations are the most common cause of (β+) thalassemia.

There is a deficit in the synthesis of HbA with an elevation of HbA2 and HbF levels in β-thalassemia. Children with severe thalassemia present with distinctive facial features such as maxilla hyperplasia, a flat nasal bridge, and frontal bossing. Other manifestations include pathologic bone fractures, significant hepatosplenomegaly, and cachexia.

Blood smears show marked anisocytosis, poikilocytosis, and microcytic hypochromic anemia. Target cells, basophilic stippling, and fragmented red cells also are common. A radiographic image of the skull shows the classic 'crew cut' appearance.

Hb Electrophoresis and high-performance liquid chromatography (HPLC) are used as screening tests for thalassemia. DNA diagnosis to detect β-thalassemia mutations is used for definitive diagnosis. The Mentzer index is the ratio of MCV to RBC count. It is used to differentiate between iron-deficiency anemia and β-thalassemia trait. The Mentzer index is <13 in thalassemia minor syndrome.

Transfusions are administered every 3-4 weeks to maintain a pretransfusion hemoglobin level of 9.5-10.5 g/dL. Transfusion-induced hemosiderosis is a significant complication in transfusion-dependent thalassemia. Iron deposition occurs in the liver, endocrine organs, and

heart, resulting in complications including hypothyroidism, hypogonadotropic hypogonadism, growth hormone deficiency, hypoparathyroidism, diabetes mellitus, heart failure, and arrhythmias.

The image below shows thalassemic facies and crew cut appearance on X-ray:

Paediatrics NEET 2018

Question 1:

A fetus with intrauterine growth restriction was born prematurely with jaundice, hepatosplenomegaly, microcephaly, and diffuse petechiae at birth. A brain CT was done which revealed periventricular calcifications. What is the best method for the diagnosis of the etiological agent?

a) Urine examination

b) Liver biopsy

c) Blood examinaton

d) CSF examination

Question 2:

Which of the following is false regarding the given image?



a) It represents melanocytic hamartoma

b) It causes no visual disturbance

c) It is associated with facial angiofibroma

d) They are diagnostic of a neurocutaneous syndrome

Question 3:

Congenital adrenal hyperplasia most commonly presents as _____.

a) Male pseudohermaphroditism

b) Female pseudohermaphroditism

c) True hermaphroditism

d) 46, XY intersex

Question 4:

A newborn presented with a tuft of hair over the lumbosacral region. What is the likely diagnosis?

a) Spina bifida occulta

b) Spinal lipoma

c) Dermal sinus

d) Any of the above

Question 5:

What is the most common cause of ventriculomegaly in newborns?

a) Arnold-Chiari malformation

b) Dandy-Walker syndrome

c) Arachnoid villi malformation

d) Aqueductal stenosis

Question 6:

When do you consider administering epinephrine in a neonate during resuscitation?

a) Heart rate remains at < 60 beats/minute despite effective compressions and ventilations.

b) Heart rate remains at < 100 beats/minute despite effective compressions and ventilations.

c) Heart rate does not improve after 30 seconds with bag and mask ventilation.

d) Infants with severe respiratory depression fail to respond to positive-pressure ventilation via bag and mask.

Question 7:

Which of the following is not true regarding ataxia-telangiectasia?

a) Mutations in 11q gene is implicated

b) Follows autosomal dominant mode of inheritance

c) Humoral and cellular immunodeficiency

d) Linked to adenocarcinomas

Question 8:

Warthin Finkeldey cells are seen in:

a) Rubella

b) Rubeola

c) Rabies

d) Typhoid

Question 9:

Which of the following is an X-linked disorder?

a) Color blindness

b) Thalassemia

c) Sickle cell anemia

d) Cystic fibrosis

Question 10:

The following parameter in ALL indicate a poor prognosis _____.

a) Age >10 years

b) Leukocyte count <50,000/mm3

c) Hyperdiploidy

d) Trisomy of chromosomes 4, 10, and 17

Question 11:

Viral infection causing transient aplastic anemia is _____

a) HIV

b) Polio

c) Parvovirus B-19

d) HHV-8

Question 12:

What is true about ovotesticular disorders of sex development?

a) Karyotype is 46 XY.

b) Ovotestis is seen in the gonadal biopsy

c) The response of testosterone to hCG is increased.

d) Uterus is usually absent.

Question 13:

Which one of the following is not included in the criteria for neurofibromatosis 1?

a) Two or more iris hamartomas

b) Acoustic neuromas

c) Dysplasia of the sphenoidal and tibial bone

d) Cafe-au-lait spots

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	d
5	d
6	a
7	b

8	b
9	a
10	a
11	c
12	b
13	b

Detailed Explanations

Solution to Question 1:

Periventricular calcifications and microcephaly in a baby with suspected congenital infection point toward cytomegalovirus (CMV). The best diagnostic method for CMV is PCR or viral culture on urine.

Since not all affected infants are viremic at birth, urine is a more reliable sample than blood.

Intrauterine growth restriction (IUGR), prematurity, and hepatosplenomegaly raise the suspicion of congenital infection. Hepatosplenomegaly, jaundice, pneumonitis, petechiae, and meningoencephalitis are common to toxoplasmosis, rubella, CMV, herpes simplex virus (HSV), enteroviruses and syphilis. Hydrocephalus is seen in CMV, toxoplasmosis, HSV, and rubella.

The characteristic clinical features of various congenital infections are described in the table below:

The triad of congenital CMV infection includes intracranial calcifications (periventricular), microcephaly, and chorioretinitis.

Most CMV transmission occurs in the first trimester. The risk of transmission is high in the third trimester. 90% of the infected babies are asymptomatic and have lesser chances of having late sequelae.

In symptomatic individuals, ganciclovir can limit hearing loss and improve the developmental outcome, while there is no definitive management for congenital CMV infection.

The image below shows periventricular calcification in the head CT seen in CMV infection:

Congenital infections	Characteristicfeatures
Cytomegalovirus (CMV)	Periventricular calcifications Microcephaly
Toxoplasmosis	ChorioretinitisDiffuse intracranial calcificationsHydrocephalus
Rubella	Sensorineural hearing lossSalt-and-pepperretinopathyCataractGlaucomaCongenital defects likepatent ductus arteriosus

Congenital infections	Characteristicfeatures
Herpes simplex virus (HSV)	VesiclesMicrocephaly
Syphilis	Bone lesions (osteochondritis, periostitis)Maculopapular rashes
Enterovirus (polio, rhinovirus, hepatitis A)	MyocarditisParalysis
Varicella zoster virus	Limb hypoplasiaCicatricial skin lesionsCortical atrophy
Zika virus	Subcortical calcificationsMicrocephalyCongenital contractures (arthrogryposis)
HIV	ThrushFailure to thriveRecurrent bacterial infectionsCalcification of basal ganglia

The image below shows a blueberry muffin rash, seen in TORCH complex infections and neuroblastoma.

Solution to Question 2:

The given image shows discrete tan-colored nodules over the iris called Lisch nodules (pigmented melanocytic hamartomas). They are seen in neurofibromatosis-1 (NF-1). Facial angiofibroma (adenoma sebaceum) is a feature of tuberous sclerosis (TS) and not neurofibromatosis.

The image below shows facial angiofibroma seen in tuberous sclerosis (Bourneville disease).

Lisch nodules are specifically associated with NF-1 and are usually multiple and bilateral. They are best visualized with a slit lamp examination and don't cause any visual disturbances.

Other important ocular manifestations in NF-1 include eyelid plexiform neurofibroma, which causes S-shaped deformity of the upper lid, with a 'bag of worms' like texture, optic nerve glioma, and spheno-orbital encephalocele due to dysplasia of greater wing of the sphenoid.

NF-1 (von Recklinghausen's disease) is an autosomal dominant disorder caused by a loss-of-function mutation of the NF-1 gene located on chromosome 17q11.2. NF-1 produces neurofibromin, which acts as a tumor suppressor and a negative regulator of RAS-MAP kinase signaling.

Clinical features of NF-1 include café-au-lait macules, axillary freckling, Lisch nodules, neurofibromas, and optic gliomas.

NF-1 is more common than NF2 and typically diagnosed in childhood. NF-2 is rarer but often presents with bilateral acoustic neuroma, juvenile cataract and schwannoma.

The various neurocutaneous syndromes and their ocular manifestations are as follows:

Neurocutaneous syndrome	Ocularmanifestations
Neurofibromatosistype 1	Eyelid plexiform neurofibromaOptic nerve gliomaBilateral Lisch nodulesCongenital ectropion uveaeProminent corneal nervesCongenital glaucomaSpheno-orbital encephalocele
Neurofibromatosistype 2	Posterior subcapsular cataractHamartomas of theretinal pigment epithelium (RPE) and retina
Sturge-Weber syndrome	Choroidal hemangiomaIris heterochromiaIpsilateral glaucoma
Tuberous sclerosis	Atypical iris colobomaIris hypopigmentationRetinal astrocytoma
von Hippel-Lindausyndrome	Retinal hemangiomas

Solution to Question 3:

Congenital adrenal hyperplasia (CAH) most commonly presents as female pseudohermaphroditism as the most common enzyme deficiency is 21-hydroxylase deficiency.

A female pseudohermaphrodite is a genotypically normal female (46, XX), whose external genitalia resembles a male's.

The image below shows ambiguous genitalia in a female with CAH:

The mechanism underlying the endocrine defects in 21-hydroxylase deficiency is depicted in the image below:



Babies with 21-hydroxylase deficiency typically present around 10-14 days of age with the subset of symptoms due to aldosterone deficiency, cortisol deficiency and androgen excess.

Aldosterone deficiency leads to salt-wasting crises marked by hyponatremia, hyperkalemia, hypotension, and metabolic acidosis.

Cortisol deficiency causes manifestations like hypoglycemia, hyperpigmentation, weight loss, weakness, anorexia, vomiting, and shock.

Androgen excess in 21-hydroxylase deficiency results in virilization of females, leading to female pseudohermaphroditism, and precocious puberty in males. Virilization is usually apparent at birth in females and within 2 to 3 years of life in males.

Without treatment, shock, cardiac arrhythmias, and death may occur within days or weeks.

Other options:

Option A: Male pseudohermaphroditism is seen in CAH due to 17α-hydroxylase deficiency resulting in incomplete virilization. 17α-hydroxylase deficiency is 2nd most common enzyme affected in CAH.

Option C: True hermaphroditism is presently referred to as an ovotesticular disorder of sex development(DSD). Here, both ovarian and testicular tissue is present. It is not due to CAH.

Option D: 46 XY intersex (now called 46 XY DSD) has under virilization of genitalia. Adrenal causes include lipoid adrenal hyperplasia, 3β-hydroxysteroid dehydrogenase deficiency, 17-hydroxylase/17,20-lyase deficiency, and not 21 hydroxylase deficiency.

Solution to Question 4:

A tuft of hair over the lumbosacral region in a newborn is suggestive of occult spinal dysraphism or neural tube defect. It comprises a range of conditions, including spina bifida occulta, intradural lipoma, and dorsal dermal sinus.

Spinal dysraphism is due to the defective fusion of the midline embryonic neural, vertebral, and mesenchymal structures. Associated clinical features are high-arched feet, muscle size and strength differences between legs, and abnormal walking patterns. In newborns and young infants, the neurological examination is usually normal, but older children may experience a lack of sensation in the perineal area and back pain.

Most of the patients will have a cutaneous abnormality overlying the lower spine, including a small dimple, a tuft of hair, or a subcutaneous lipoma.

The image shows a tuft of hair in a case of spinal dysraphism:

Spinal dysraphism also includes meningocele, myelomeningocele, lipomeningocele, diastematomyelia (split spinal cord), thickened filum terminale, and cauda equina tumour.

The initial investigation is usually ultrasound in the neonate and the best investigation is MRI.

Spina bifida is the most common type of spinal dysraphism, in which the spinal column does not close completely, leaving the spinal cord and its protective covering exposed.



Myelomeningocele is the most severe form of spina bifida, where the spinal cord and nerve roots are exposed and may be damaged. A sac-like protrusion, containing spinal cord tissue, meninges, and cerebrospinal fluid, is visible outside the body, often leading to significant neurological impairments and disabilities.

Solution to Question 5:

Aqueductal stenosis is the most common cause of ventriculomegaly and hydrocephalus in newborns.

Ventriculomegaly is the dilation of the ventricles. When ventriculomegaly occurs in association with raised cerebrospinal fluid (CSF) pressure, it is called hydrocephalus.

Pathologic etiologies can be congenital or acquired. Congenital causes include structural abnormalities such as aqueductal stenosis, agenesis of the corpus callosum, Dandy-Walker malformations, Chiari malformations, and neural tube defects. Acquired causes are intracranial infections(cytomegalovirus, toxoplasmosis), intracranial hemorrhage, ischemic injury, and brain parenchymal tumors.

Aqueductal stenosis is the narrowing or obstruction of the cerebral aqueduct, a narrow channel that connects the third and fourth ventricles in the brain. As a result, the fluid accumulates within the ventricles, causing them to enlarge.

Treatment for aqueductal stenosis includes ventriculostomy, diuretics, serial lumbar punctures in specific cases, and CSF diversion. In severe cases, CSF diversion procedures (shunting), may be necessary to redirect the flow of CSF and relieve pressure on the brain.



Solution to Question 6:

As per the newborn resuscitation protocol (NRP), a heart rate < 60 beats/minute despite effective compressions and ventilation is the indication to administer epinephrine.

The NRP algorithm is shown below:



The response to neonatal resuscitation at various steps is assessed based on the heart rate, as a rise in heart rate is the most effective indicator of successful ventilatory effort.

If the heart rate is less than 100, positive pressure ventilation (PPV) is provided with a bag and mask.

In case of no response even after doing ventilation corrective steps (Options C and D), endotracheal intubation is done.

The ventilation corrective steps are as follows and can be remembered with the mnemonic MRSOPA.

• Mask has to be reapplied

• Reposition head

• Suction the secretions

• Open mouth slightly and ventilate

• Pressure is increased slightly

• Airway may be changed

If the heart rate is below 60, then chest compressions are initiated over the lower third of the sternum at a rate of 90 compressions/min. The ratio of compressions to ventilation is 3:1.

If the heart rate continues to remain <60 despite effective compressions and ventilation, then IV epinephrine is given.

Solution to Question 7:

Ataxia telangiectasia (AT) or Louis–Bar syndrome is an autosomal recessive disorder.

The defect in AT is in the ATM gene localized to chromosome 11q22-23. The ATM gene is involved in the repair of double-strand DNA breaks. This defect in DNA repair leads to combined immunodeficiency (impaired immunoglobin isotype switching and association with thymic hypoplasia) and predisposition to an increased risk of lymphoreticular tumors such as lymphoma or leukemia.

AT is characterized by progressive ataxia (beginning during infancy), telangiectasia (initially on bulbar conjunctiva), sinopulmonary infections, hypogonadism, sensitivity to ionizing radiation, insulin resistance, and raised alpha-fetoprotein levels.

Solution to Question 8:

Warthin-Finkeldey cells are multinucleated giant cells that are present in the prodromal stage of measles (rubeola), HIV lymphadenitis, and various benign and malignant lymphoid conditions.

The cytopathic effect of measles consists of multinucleate syncytium formation with numerous acidophilic nuclear and cytoplasmic inclusions.

The image given below shows the Warthin-Finkeldey giant cells:



Measles is an infection caused by Paramyxovirus which is a single-stranded enveloped RNA virus. It spreads 4 days before and 5 days after the appearance of the rash through body secretions.

Clinical features include mild fever, conjunctivitis, upper respiratory tract infection, and pathognomic Koplik spots. The Koplik spots are red lesions with bluish-white spots seen in the inner cheeks at the level of premolars. The red maculopapular rashes begin on the forehead, behind the ears, and on the upper neck on the 4th day. It then spreads towards the torso, and extremities and reaches the palms and soles.

The diagnosis is mainly made clinically but a serological demonstration of IgM antibodies and IgG antibodies is also used. Complications include pneumonia, croup, and tracheitis. Subacute sclerosing panencephalitis is a fatal complication of chronic measles diagnosed by elevated IgG levels in cerebrospinal fluid.

Solution to Question 9:

Color blindness is primarily inherited as an X-linked recessive trait. The genes for color vision are located on the X chromosome.

Males have a single X chromosome inherited from mothers. If the X chromosome inherited carries a mutation, males exhibit the disease. However, females become carriers of the color blindness gene and have a 50% chance of passing it on to their children.

Female carriers usually are protected because of the random inactivation of one X chromosome. But there is a possibility that the mutant X chromosome remains active if the normal alleles are inactivated in some of the cells so the heterozygous female can express the disorder partially.

The image shows the pattern of X-linked inheritance:



In color blindness, the mechanism to appreciate one or more primary colors is either defective (anomalous) or absent (anopia). It may be congenital or acquired. Congenital color blindness can be dyschromatopsia (defective color vision of any one of the RGB colors) or achromatopsia (complete color blindness of all three RGB colors). RGB colors are red, green, and blue.

Tests for color blindness include the Ishihara test, Hardy–Rand–Rittler test, and Farnsworth–Munsell 100-hue test.

Solution to Question 10:

Age >10 years indicates a poor prognosis in ALL.

Leukocyte count <50,000/mm3, hyperdiploidy, and trisomy of chromosomes 4,10,17 indicate a good prognosis.

Solution to Question 11:

Parvovirus B19

• Common, most prevalent are the rash illness of erythema infectiosum and transient aplastic crisis.

• Transmission by respiratory route but also transmissible in blood and blood products.

• Shows erythrotropism – it specifically targets the erythroid cell line, especially the pronormoblast stage. Viral infection produces cell lysis, leading to depletion of erythroid precursors and a transient arrest of erythropoiesis. This may lead to a precipitous fall in haemoglobin which may even require blood transfusion.

Clinical manifestations of Parvovirus B19 infection

• Erythema infectiosum/Fifth disease (slapped cheek appearance)

• Transient aplastic anemia

• Petechial/ purpuric skin eruptions: Papular purpuric “gloves and socks” syndrome

• Arthritis and arthralgia

• Chronic anemia in immunocompromised persons

• Primary maternal infection is associated with non-immune fetal hydrops and intrauterine fetal demise.

Solution to Question 12:

Ovotesticular disorder of sex development (DSD) is characterized by the presence of both ovarian and testicular tissue in the same individual. An ovotestis is present in approximately 2/3 of affected individuals.

Most affected individuals have a 46, XX chromosomal karyotype, which normally results in female sexual development. In about 10% of patients, testicular tissue in an individual with a 46, XX karyotype is present as a result of a translocation of the SRY gene on the Y chromosome to the X chromosome or another chromosome. There is a defect in R-spondin1 protein encoded by the RSPO1 gene, described in 46, XX ovotesticular DSD.

An abnormal vagina and an underdeveloped uterus are usually present. The testes are usually undescended, and the penis may present with hypospadias. Upon reaching puberty, breast development, feminization, and menstruation may occur. Most affected individuals are infertile, but ovulation or spermatogenesis is possible. The response of testosterone to hCG is normal or reduced.

Solution to Question 13:

In neurofibromatosis 1 (NF1), two or more of the following clinical manifestations are present:

• Six or more café-au-lait macules

• Axillary or inguinal freckling

• Two or more iris Lisch nodules (hamartomas of iris)

• Two or more neurofibromas or 1 plexiform neurofibroma

• A distinctive osseous lesion such as sphenoidal and tibial bone dysplasia

• Optic gliomas

• A 1st-degree relative with NF-1

Paediatrics NEET 2019

Question 1:

Unconjugated hyperbilirubinemia, which did not subside even after 3 weeks of birth, was observed in a neonate. On investigating, liver enzymes, PT/INR and albumin levels were normal. No hemolysis was seen on a peripheral blood smear. A drop in bilirubin level was observed within a week after treating with phenobarbital. What is the most likely diagnosis?

a) Rotor syndrome

b) Crigler Najjar type 2

c) Dubin Johnson syndrome

d) Crigler Najjar type 1

Question 2:

Which one of the following is not a feature in cerebral palsy?

a) Hypotonia

b) Microcephaly

c) Ataxia

d) Flaccid paralysis

Question 3:

Which of the following is not seen in a child with cystic fibrosis?

a) Sweat chloride test chloride conc of 70 mEq/L

b) Increase immunoreactive trypsinogen level

c) Hyperkalemia

d) Contraction alkalosis

Question 4:

A 9-year-old boy has swelling of knee joints and non-blanching rashes as seen in the picture. Urine exam reveals haematuria +++ and proteinuria +. Platelet levels are normal. What is the

probable diagnosis?

a) Systemic lupus erythematosus (SLE)

b) Henoch Schonlein purpura (HSP)

c) Immune thrombocytopenic purpura (ITP)

d) Polyarteritis Nodosa (PAN)

Question 5:

A child in your unit is diagnosed with severe acute malnutrition (SAM). According to the WHO criteria, SAM is defined best by:

a) Weight for age less than - 2SD

b) Weight for height less than + 2SD

c) Weight for age less than + 3SD

d) Weight for height less than - 3SD

Question 6:

The most common manifestation of congenital toxoplasmosis is _____

a) Hydrocephalus

b) Chorioretinitis

c) Hepatosplenomegaly

d) Thrombocytopenia

Question 7:

Molecular defect causing liver disease in Dubin-Johnson syndrome is_____

a) ATP7A

b) ATP7B

c) ABCC2

d) SERPINA 1

Question 8:

Where would you place the pulse oximeter to measure pre-ductal oxygen saturation in an infant who was born 3 minutes ago?

a) Left upper limb

b) Left lower limb

c) Right upper limb

d) Right lower limb

Question 9:

All among the following are clinical features of necrotizing enterocolitis except _____

a) Vomiting

b) Abdominal mass

c) Erythema of the abdominal wall

d) Metabolic alkalosis

Question 10:

The most common cause of HIV infection in a newborn is _____

a) Perinatal transmission

b) Breast milk

c) Transplacental

d) Exchange transfusion with infected blood

Question 11:

Which of the following is the main factor for the ductal closure postnatally?

a) Increase in partial pressure of oxygen (paO2)

b) Increase in systemic vascular resistance

c) Increase in circulating prostaglandin levels

d) Decrease in pulmonary venous resistance

Question 12:

Cardiac edema and neuropathy are seen in which of the following vitamin deficiencies?

a) Biotin

b) Thiamine

c) Pyridoxine

d) Riboflavin

Question 13:

The commonest indication for liver transplantation in children is _____

a) Alagille syndrome

b) Biliary atresia

c) Caroli disease

d) Hepatcellular carcinoma

Question 14:

While you are evaluating a baby, you show him a bright pink teddy bear, that he reaches out to with both hands. What is the earliest age by which this milestone is typically achieved?

a) 4 months

b) 5 months

c) 6 months

d) 7 months

Question 15:

A baby assessed 5 minutes after birth, is found to be cyanosed with irregular gasping respiration. Heart rate is 60 beats/min, and shows minimal response to stimulation with some flexion of extremities. The Apgar score for this newborn is _____

a) 2

b) 5

c) 3

d) 4

Question 16:

Most common heart defect in congenital rubella syndrome is:

a) ASD

b) VSD

c) PDA

d) PS

Question 17:

Which among the following options describes the chest x-ray image?



a) Egg on string appearance

b) Coeur en sabot

c) Scimitar sign

d) Snow man sign

Answer Key

Question No.	Correct Option
1	b
2	d
3	c
4	b
5	d
6	b
7	c
8	c
9	d
10	a
11	a
12	b
13	b
14	a
15	d
16	c
17	b

Detailed Explanations

Solution to Question 1:

Jaundice persisiting 3 weeks after birth is suggestive of pathological jaundice. Among the given options, Crigler Najjar type 2 presents with unconjugated hyperbilirubinemia which responds to phenobarbital therapy.

Dubin Johnson (option C) and Rotor (option A) syndromes lead to conjugated hyperbilirubinemia.

Crigler Najjar type 2 is an autosomal recessive disease with non-hemolytic, unconjugated hyperbilirubinemia caused by homozygous missense mutations in the enzyme UDP UGT1A1 resulting in reduced (partial) enzymatic activity.

The differences between Crigler-Najjar type 1 and type 2 are described in the table below:

Crigler Najjar Type 1		Crigler Najjar Type 2
Autosomal recessive		Autosomal recessive
UDPGT isabsent		UDPGT isreduced
Severe unconjugated hyperbil irubinemia		Less severe
Pale stools		Normal-coloured stools
May proceed tokernicterusan ddoes notrespond to phenob arbitone		Does not proceed to kernicter us andis responsive to pheno barbitone

Solution to Question 2:

Flaccid paralysis is not a feature of cerebral palsy (CP).

CP is an upper motor neuron disease characterized by rigidity and scissoring gait. Flaccid paralysis would be seen in lower motor neuron disease.

Other options:

Option A: Hypotonia is a characteristic feature of athetoid CP. These infants generally have poor head control.

Option B: Microcephaly can be seen in CP due to perinatal insults to the developing brain.

Option C: Ataxia is a common feature of extrapyramidal CP.

The subtypes of cerebral palsy are given in the table below:

UL - upper limb; LL - lower limb; GDD - global developmental delay

Types	Etiology	Presentation	MRI findings	
Spastic(m/c type)	Diplegia(second m/c in India)	PrematurityIschemiaInfection	Bilateral spasticity (LL>UL)Scissoring gaitCommando crawlLearning disabilities	Periventricular leukomalacia
Hemiplegia	Vascular insultInfectionGenetic	Early hand preference(UL>LL)Walking delayed up to 18-24mEquinovarusdeformity of footSeizures (33%)	Porencephalic cystFocal infarct changes	

Types	Etiology	Presentation	MRI findings	
Quadriplegia(most severe)(m/c type in India)	IschemiaInfection	Marked motor impairmentFlexioncontracturesStrong association withseizures Severeintellectual impairment andGDDDifficulty swallowing (supranuclear palsy)	Multi-cystic encephalomalaciaSevere periventricular leukomalacia	
Dyskinetic(15-20%)	ExtrapyramidalAthetoidChorioathetoid	AsphyxiaKernicterusMitochondrialGenetic	Rigidity/hypotonia, dystonia, athetosis, choreaCognitive disabilityHip displacementSeizures	Basal ganglia scarringThalamic scarring

Solution to Question 3:

Patients with cystic fibrosis (CF) can present with hypokalemia and not hyperkalemia.

Excess production of sweat in CF leads to a significant loss of chloride ions from the extracellular fluid (ECF). The depletion of chloride from the ECF causes an intracellular shift of potassium and activation of renin-aldosterone system, leading to hypokalemia.

The excessive depletion of sodium chloride in CF also affects the acid-base balance. It leads to increased reabsorption of bicarbonate by the kidneys through the chloride-bicarbonate exchanger on the intercalated cells. This process is known as contraction alkalosis (option D) and results in hyponatremic hypochloremic alkalosis.

Cystic fibrosis, also known as mucoviscidosis, is an autosomal recessive disorder caused by a mutation in the cystic fibrosis transmembrane regulator (CFTR) gene on chromosome 7q. This gene is responsible for encoding ATP-gated chloride channels in the apical membranes of

epithelial cells, which regulate the flow of ions across various epithelial membranes in the body, including sweat glands, respiratory airways, and pancreatic ducts. The defect in CFTR leads to elevated salt concentrations in sweat and the production of thick, viscous secretions in the respiratory and gastrointestinal tracts.

Diagnostic tests for CF include the sweat chloride test (option A), where an increased chloride concentration (>60 mEq/L) is considered diagnostic. Immunoreactive trypsinogen levels (option B) are also measured for newborn screening, as they tend to be elevated in CF. Genetic testing for CFTR gene mutations is considered the gold standard and should be performed for all suspected cases. In equivocal cases, nasal transepithelial potential difference can be estimated as an additional diagnostic test.

The criteria for diagnosis is the presence of clinical symptoms affecting at least one organ system (respiratory, gastrointestinal, or genitourinary) or a positive newborn screening or a sibling with cystic fibrosis, along with evidence of dysfunction of the CFTR gene (two positive sweat chloride test results obtained on separate days or identification of two CFTR mutations by genetic testing or an abnormal measurement of nasal potential difference).

The image below shows the organs involved in cystic fibrosis:



Solution to Question 4:

The given clinical scenario showing swelling of knee joints (arthritis), non-blanching rash in the lower limbs (palpable purpura), along with hematuria and proteinuria (renal involvement) in a child is suggestive of Henoch Schonlein purpura (HSP).

HSP (IgA vasculitis) is a small vessel vasculitis most commonly seen in the age group of 3-8 years. It generally follows an upper respiratory tract infection and there is a history of atopy in one-third of the cases.

The pathology of HSP involves leukocytoclastic vasculitis and the deposition of IgA in the affected blood vessels. IgA is deposited in the glomerular mesangium in a distribution similar to that of IgA nephropathy.

Clinical manifestations include palpable purpura, the hallmark feature of HSP, which characteristically involves the extensor surfaces of arms and legs. Musculoskeletal symptoms such as arthritis and arthralgia are common. Gastrointestinal symptoms can include abdominal pain, vomiting, diarrhea, and bleeding. Renal involvement may lead to gross or microscopic hematuria, nephritic syndrome, or nephrotic syndrome. Renal manifestations are more severe in adults. A rapidly progressive form of glomerulonephritis is seen in some cases. Nephritis usually follows the appearance of the rash.

Diagnosis of HSP is primarily based on clinical findings, including the palpable purpura, along with the presence of other symptoms. Laboratory tests are not specific to HSP but may show nonspecific markers of inflammation and sometimes elevated levels of IgA. Imaging studies and biopsies of affected tissues can provide additional diagnostic information

Treatment of HSP is generally supportive, focusing on hydration, pain relief, and nutrition. Corticosteroids may be used in severe cases or for significant gastrointestinal involvement. Intravenous immune globulin (IVIG) and plasma exchange are occasionally employed for severe disease. Nephritis due to IgA vasculitis should be treated with immunosuppressants like prednisolone and azathioprine. Complications of HSP can include serious gastrointestinal problems, kidney disease, and recurrences, but overall, the prognosis for childhood HSP is favorable. Regular monitoring of blood pressure and urine is recommended to detect and manage potential complications.

Solution to Question 5:

Severe acute malnutrition (SAM) is best defined as weight for height less than median minus -3SD

Severe acute malnutrition (SAM) in children of ages 6 to 59 months is defined by the WHO as any of the following:

• Weight-for-length (or height) below –3 SD of the median WHO child growth standards.

• Presence of bipedal edema

• Mid-upper arm circumference <11.5 cm

Once the child is diagnosed with severe acute malnutrition, the doctor must assess for severe edema (+++); low appetite (failed appetite test); medical complications (severe anemia, pneumonia, diarrhea, dehydration, cerebral palsy, tuberculosis, HIV, heart disease, etc.) and one or more danger signs as per IMNCI

If any of the complications are present, the child is diagnosed with complicated SAM and is managed inpatient in the facility.

If the above-mentioned signs are absent, the child is diagnosed with uncomplicated SAM and supervised management at home is given.

Solution to Question 6:

Chorioretinitis is the most common manifestation of congenital toxoplasmosis.

The characteristic triad of congenital toxoplasmosis is chorioretinitis, hydrocephalus, and diffuse intracranial calcifications. Clinical manifestations apart from the triad include hydrops fetalis, intrauterine growth retardation, prematurity, peripheral retinal scars, persistent jaundice, and mild thrombocytopenia.

Congenital toxoplasmosis occurs when a pregnant woman acquires the infection during gestation. The organism may cross to the fetus transplacentally or at the time of vaginal delivery. The severity of the disease in the neonate is greater if the woman gets infected early in her pregnancy. The causative agent is Toxoplasma gondii, an obligate intracellular protozoan.

The diagnosis is made by the detection of IgM anti-toxoplasma antibodies. IgG can cross the placenta is hence, cannot confirm infection in the infant. IgM immunosorbent agglutination assay (ISAGA) and IgA ELISA are the most useful diagnostic tests.

Spiramycin can be administered for cases of acute infection in early pregnancy to reduce vertical transmission. It does not cross the placenta and therefore cannot be used to treat an infected fetus. Pyrimethamine-sulfonamide (to be avoided in the first trimester) is used when a fetal infection is confirmed or in maternal infections after 18 weeks.

Solution to Question 7:

The molecular defect causing liver disease in Dubin–Johnson syndrome is ABCC2.

Dubin-Johnson syndrome is an autosomal recessive disorder due to a defect in ABCC2. ABCC2 codes for canalicular protein with ATP-binding cassette. A defect in this gene leads to the mutation of the multiple drug-resistant protein 2 (MRP2) receptor. This in turn leads to a defective transfer of conjugated bilirubin from the liver to bile.

It presents with abdominal pain, mild icterus, and conjugated hyperbilirubinemia. Biopsy shows black liver due to an accumulation of epinephrine metabolite.

Conditions causing conjugated hyperbilirubinemia are summarised in the table below:

Important diseases and their molecular defects:

Dubin-Johnson syndrome	Rotor syndrome
Autosomal recessive	Autosomal recessive
The absent function ofmultip le drug-resistant proteins (M RP-2)	Defects in anion transport
Black liver on biopsy (due to accumulation of epinephrine metabolite)	No abnormal liver pigmentati on.
Normal levels of coproporph yrin in urine	Increased levels of coproporp hyrin in urine
The gall bladder is not visuali zed on oral cholecystography	The gall bladder is visualized on oral cholecystography

ATP7A	Menke's disease
ATP7B	Wilson disease
CFTR	Cystic fibrosis
SERPINA 1	α-1 antitrypsin deficiency
ABCC2 (MRP2 protein defect)	Dubin Johnson syndrome

Solution to Question 8:

The right upper limb is the best area to check pre-ductal oxygen saturation in a newborn infant as it receives blood supply primarily from the right brachial artery, which arises from the aorta before the ductus arteriosus.

In routine neonatal screening for critical congenital heart diseases (CCHD), saturation probes are placed on the right upper limb (pre-ductal) and the right lower limb (post-ductal) respectively.

If the difference between pre-ductal and post-ductal saturation is more than five percent, it confirms the presence of hypoxemia. Any reading showing an SpO2 <90% or greater than a three percent difference between the right hand and right foot for three successive readings separated by an hour each would suggest the presence of a congenital cardiac disease.

Screening is done to identify critical conditions before they present symptomatically, and before the neonate leaves the hospital post-delivery. This would allow for early and effective treatment of an underlying disease.

Critical congenital heart diseases include coarctation of the aorta, double outlet right ventricle, Ebstein anomaly, hypoplastic left heart syndrome, interrupted aortic arch, pulmonary atresia, single ventricle, tetralogy of Fallot (TOF), total anomalous pulmonary venous return (TAPVC), d-transposition of the great arteries (TGA), tricuspid atresia, and truncus arteriosus.

Solution to Question 9:

Necrotizing enterocolitis (NEC) can cause metabolic or respiratory acidosis (not alkalosis).

Acidosis can be attributed to sepsis, lactic acidosis due to decreased perfusion of the abdominal wall, decreased reabsorption of bicarbonate from the intestine, and reduced respiratory compensation due to apnea.

NEC is the most common life-threatening gastrointestinal disorder in the newborn period. It is characterized by various degrees of mucosal or transmural inflammation and necrosis of the intestinal tract. It most commonly affects the distal ileum and proximal colon.

Three major risk factors implicated are prematurity (greatest risk factor), formula feeding, and bacterial colonization of the intestine. Incidence and mortality increase with decreasing birth weight and gestational age.

It usually presents within the 2nd or 3rd week of life with abdominal distension and tenderness, feed intolerance, and occult/gross blood in the stool. Systemic features include hypotension, bradycardia, lethargy, and temperature instability. Other signs and symptoms include vomiting, changes in stool pattern, abdominal mass, apnea, respiratory distress, hyponatremia, shock, and disseminated intravascular coagulation.

Diagnosis can be made with a plain abdominal X-ray. Pneumatosis intestinalis (air in the bowel wall) is diagnostic. Portal vein gas is a sign of severe disease. Pneumoperitoneum indicates perforation of the intestinal wall.

Staging is done using the modified Bell's staging criteria:

Management of NEC includes supportive measures - nil per oral, nasogastric decompression, intravenous fluids, broad-spectrum antibiotics, and inotropic support if indicated. Perforation is an absolute indication for surgery. The most effective preventive strategy is the use of human milk. Prebiotics and probiotics can be used with variable efficacy.

The images below show pneumatosis intestinalis (A), pneumoperitoneum (B), gasless abdomen (C), and distention of the abdomen with periumbilical erythema (D):

Stage	Features	Radiological signs	Management	Treatment
ISuspected NEC	Mildabdominal distention, vomitingTemperature instability, apnea, bradycardiaI A:Occult blood in stoolsI B:Gross bloodin stools	Normal or intestinal dilatation, mild ileus	MedicalNil per oral Total parenteral nutrition - IV fluids containing dextroseIV antibiotics -ampicillin, gentamycin, metronidazole	Antibiotics for 3 days
II AMildly ill	Absentbowel sounds	Pneumatosis intestinalis	Antibiotics for 14 days	
II BModerately ill	II A + abdominal wall edemaThrombocytopeniaHyponatremiaMetabolic acidosis	II A +portal venous gas, with or without ascites		
III ASeverely ill, bowel intact	PeritonitisDefinite ascites	II B +definite ascites		
III BSeverely ill, bowelperforated	Intestinal perforation	Pneumoperitoneum	Surgical(due to bowel perforation)	Stable- Laparotomy (resection of necrosed part, followed by anastomoses)Unstable-Primary peritoneal drainage, and once stabilized definitive surgery



Solution to Question 10:

The most common cause of HIV infection in a newborn is perinatal transmission. Around 70 to 80% of affected children acquire the virus intrapartum.

The mechanism of perinatal transmission has been postulated to be mucosal exposure to infected blood and cervicovaginal secretions in the birth canal. The intrauterine contractions during active labor/delivery could also increase the risk of late micro transfusions.

Vertical transmission of HIV can occur in utero (during pregnancy), during delivery (perinatal), and after birth by breastfeeding (least common).

The risk factors for vertical transmission are high maternal viremia, preterm delivery (<34 weeks), vitamin A deficiency, prolonged rupture of membranes (>4 hours) low birth weight, and low maternal CD4 count.

Solution to Question 11:

Oxygen is the most important factor controlling ductal closure.



Gestational age also appears to play an important role; the ductus of a premature infant is less responsive to oxygen, even though its musculature is developed. Hence, patent ductus arteriosus (PDA) is more common in premature infants.

The fetal circulatory adjustments after birth are given below:

The patency of ductus arteriosus may either provide an alternative route for blood to bypass congenital defects such as pulmonary atresia or coarctation of the aorta.

Therapeutic medications can be used to either maintain the patency (prostaglandin E1) or promote closure (indomethacin). Drugs like indomethacin are not recommended during the third trimester of pregnancy.

	Functional closure	Anatomical closu re
Ductus venosus	After removal of the placenta	7 days
Foramen ovale	At birth - after breathing is in itiated	Months to years
Ductus arterios us	10–15 hours	10–21 days

Solution to Question 12:

Cardiac edema and neuropathy are associated with the deficiency of thiamine (vitamin B1).

Early symptoms of thiamine deficiency include peripheral neuropathy, irritability, loss of deep tendon reflexes, and proprioception. As the disease progresses, increased intracranial pressure, meningismus, and coma develop.

Wet beriberi (cardiac) is associated with tachycardia, cardiomegaly, cardiac failure, and edema. It is commonly seen in the refugee camps where people consume polished rice based monotonous

diets. Diagnosis of thiamine deficiency can be established by erythrocyte transketolase activity assessment.

Dry beriberi is associated with neurological deficits such as optic atrophy, ataxia, and irritability.

Complications include lactic acidosis and Wernicke encephalopathy.

Solution to Question 13:

Biliary atresia is the most common indication for liver transplantation in children.

Extrahepatic biliary atresia (EHBA) is defined as a partial or total absence of permeable bile duct between the porta hepatis and the duodenum. It is due to inflammatory fibrosis of the biliary tree. It leads to obstruction to bile flow thereby causing the leaking of conjugated bilirubin into the circulation.

Infants have symptoms of jaundice, pale stools, and dark urine right after birth. In advanced disease, infants present with failure to thrive, hepatomegaly, and ascites due to liver cirrhosis.

HIDA scan is the investigation of choice for extrahepatic biliary atresia. An intraoperative cholangiogram is a gold standard investigation for diagnosis.

Children with biliary atresia, who do not achieve successful drainage after hepatoportoenterostomy (Kasai's procedure), require liver transplantation within the first year of life.

Management of extrahepatic biliary atresia		
Type I	Atresia restricted to the common bile duct	Roux-en-Y hepaticojejunostomy
Type II	Atresia of the common hepatic duct	Kasai procedure(Roux-en-Y hepatoportoenterostomy)
Type III	Atresia of the right and left hepatic duct	

Solution to Question 14:

The earliest age of appearance of bidextrous reach is 4 months.

Fine motor reflexes begin to appear only after the palmar grasp reflex disappears at 3-4 months. The development of unidextrous reach occurs by 6 months of age.

Solution to Question 15:

The Apgar score in this baby is 4. It is calculated as:

Appearance: blue - score 0

Pulse: <100 beats/min - score 1

Grimace: with minimal response to stimulation - score 1

Activity: some flexion of the extremities - score 1

Respiratory effort: irregular - score 1

The total score is 4.

Apgar score is not used to guide resuscitation, and resuscitation must be initiated before the 1-minute score. The change of score at sequential time points following birth reflects how well the baby is responding to resuscitative efforts.

Causes for low Apgar score are:

• Fetal distress

• Prematurity

• Drugs like analgesics, narcotics, sedatives, magnesium sulfate

• Acute cerebral trauma

• Precipitous delivery

• Airway obstruction (choanal atresia)

• Congenital pneumonia or sepsis

• Haemorrhage- hypovolemia

False-negative Apgar score is seen in maternal acidosis and high fetal catecholamine levels.

A total score of 10 indicates that the infant is in the best possible condition. An infant with a score of 0-3 requires immediate resuscitation. If the score at 5 minutes remains less than 7, the infant should be assessed again every 5 minutes for the next 20 minutes.

Solution to Question 16:

Patent ductus arteriosus (PDA) is the most common congenital heart disease associated with rubella, followed by pulmonary artery stenosis (PS).

Cardiac anomalies have been observed in 50% of children infected during the first 8 weeks of gestation. The incidence of heart disease in congenital rubella syndrome is: PDA > PS (right >left) > ventricular septal defect > atrial septal defect.

Solution to Question 17:

The given chest x-ray shows 'coeur-en-sabot' or boot-shaped heart, which is seen in tetralogy of Fallot (TOF).

The characteristic silhouette is due to cardiac apex elevation as a result of right ventricular hypertrophy. The hilar areas and lung fields are relatively clear because of diminished pulmonary blood flow or the small size of the pulmonary arteries and the aorta is usually enlarged.

The image below describes the likeness of the cardiac silhouette to a boot:



TOF is cyanotic congenital heart disease with the following four components:

• Obstruction to right ventricular outflow (pulmonary stenosis)

• Malalignment type of ventricular septal defect (VSD)

• Dextroposition of the aorta so that it overrides the ventricular septum (overriding of the aorta)

• Right ventricular hypertrophy (RVH)

The severity of symptoms depends on the degree of the right ventricular outflow obstruction. In mild degrees, the symptoms become pronounced as the baby grows due to RV infundibular hypertrophy and may present with heart failure. In severe degrees, pulmonary blood flow predominantly depends on the ductus arteriosus. As soon as the duct closes, the baby presents with cyanosis and circulatory collapse in the neonatal period.

Clinical features include significant hypercyanotic spells in infants which are characterised by hyperpnea, cyanosis, and gasping followed by syncope. The spells occur most frequently in the morning on initially awakening or after episodes of vigorous crying. These spells are due to a prolonged reduction of pulmonary blood flow resulting in severe systemic hypoxia and metabolic acidosis.

Older children present with exertional dyspnea and assume a squatting position to relieve the dyspneic episode after which the child is able to resume the activity.

Paediatrics NEET 2020

Question 1:

Two girls in the same class are diagnosed with meningococcal meningitis. Their 12-year old friend from the same school is afraid of contracting the disease. What advice should be given to the exposed students?

a) Two doses of polysaccharide vaccine

b) Antibiotic prophylaxis

c) Two doses of conjugate vaccine

d) Single dose of meningococcal vaccine

Question 2:

A laborer's second child is brought to the OPD with a swollen belly and dull face. He has been fed rice water (rice milk) in his diet mostly. On investigations, the child is found to have low serum protein and low albumin. What is the probable diagnosis?

a) Kwashiorkor

b) Marasmus

c) Indian childhood cirrhosis

d) Kawasaki disease

Question 3:

A 3-year-old male has the following buccal mucosa finding. After three days, he presents to your OPD with a rash. What is your diagnosis?



a) Measles

b) Leukoplakia

c) Scarlet fever

d) Kawasaki disease

Question 4:

The given chest X-ray is suggestive of:



a) Tetralogy of Fallot

b) Atrial septal defect

c) Ventricular septal defect

d) Ebstein's anomaly

Question 5:

The greenish-black color of the first stool in the newborn is due to :

a) Bilirubin

b) Biliverdin

c) Meconium

d) Gut flora of neonate

Question 6:

Maternal antibodies that do not provide significant immunity in a neonate:

a) Tetanus

b) Polio

c) Diptheria

d) Measles

Question 7:

Anuj has been■ diagnosed with diphtheria and is being treated in the ward. His 3-year-old brother, Ajay who lives in the same house has received vaccination against diphtheria about 16 months back. What is the next best step for the management of the younger sibling, Ajay?

a) One booster dose

b) Nothing as the child is already exposed

c) Erythromycin + diphtheria toxoid

d) Erythromycin only

Question 8:

According to current recommended dietary guidelines for children, energy from saturated fats should be _____ of total energy intake.

a) < 10%

b) < 5%

c) < 15%

d) < 30%

Question 9:

A 3-year-old girl was brought by her mother with complaints of enlarged clitoris. Karyotype revealed 46XX. What is the likely cause?

a) 21 hydroxylase deficiency

b) 11 hydroxylase deficiency

c) 3 beta dehydrogenase deficiency

d) 2, 3 lyase deficiency

Question 10:

A newborn is brought with evisceration of bowel loops from a defect in the abdominal wall to the right of the umbilicus. The most probable diagnosis is

a) Omphalocele

b) Umbilical hernia

c) Gastroschisis

d) Ectopia vesicae

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	a
5	b
6	b
7	d
8	a

9	a
10	c

Detailed Explanations

Solution to Question 1:

Antibiotic prophylaxis is provided to the close contacts of persons with confirmed meningococcal disease.

Close contacts of individuals with confirmed meningococcal disease are at an increased risk of developing the illness. Prophylaxis is also offered to individuals with direct and prolonged contact with a case of meningococcal disease. Antibiotics are effective in preventing additional cases by eradicating the carriage of the invasive strain.

Treatment is provided to the contacts within 24 hours of identification of the index case. The antibiotics effective include rifampicin, ciprofloxacin, ceftriaxone, and azithromycin.

Prophylaxis is not generally recommended for medical personnel, except in cases where there is exposure to aerosols of respiratory secretions, such as during mouth-to-mouth resuscitation, intubation, or suctioning. This recommendation applies within 24 hours before or after the initiation of antibiotic therapy in the index case.

For the treatment of cases, penicillin is the preferred drug, but ceftriaxone or other third-generation cephalosporins are recommended for patients with penicillin allergy. Droplet infection control precautions should be followed for hospitalized patients for 24 hours after effective therapy has been initiated.

Offering quadrivalent meningococcal A, C, W, Y conjugate vaccines in addition to antimicrobial prophylaxis can further reduce the risk of disease among close contacts of cases caused by vaccine capsular groups.

Meningococcal vaccines are available in two forms: polysaccharide vaccines and polysaccharide-protein conjugate vaccines. Polysaccharide vaccines can be given as a single dose to individuals aged 2 years and older. They are offered in bivalent (A, C), trivalent (A, C, W135), and quadrivalent (A, C, W135, Y) formulations.

Meningococcal conjugate vaccines are available as monovalent (A or C) or quadrivalent (A, C, W135, Y) formulations. Conjugate vaccines, which offer increased immunogenicity and potential for herd protection, are preferred over polysaccharide vaccines, especially in children under 2 years of age. Both types of vaccines are safe to use during pregnancy.

Solution to Question 2:

The given clinical scenario of a child with a swollen belly, dull face with hypoproteinemia along with poor diet history suggests the diagnosis of kwashiorkor.

Kwashiorkor is a form of severe acute malnutrition (SAM) characterized by edema and apathy. In edematous malnutrition, the initial appearance of edema is commonly observed in the feet and lower legs, progressing into generalized edema affecting the hands, arms, and face. The skin over the swollen areas may show dark, crackled, peeling patches (flaky paint dermatosis) with pale skin underneath, making it susceptible to infections.

There may be sparse and easily pluckable hair. Hepatomegaly with fatty infiltration is a common finding. Associated symptoms of misery, apathy, and loss of appetite are seen in children with edema.

Other options:

Option B: Marasmus is a severe form of wasting characterized by significant loss of fat and muscle tissue as the body utilizes these resources for energy. It is often triggered by acute starvation or illness in individuals with borderline nutritional status.

Option C: Indian childhood cirrhosis (ICC) is a chronic liver disease that affects infants and young children. Common symptoms include jaundice, pruritus, lethargy, and hepatosplenomegaly. The disease progresses rapidly, leading to the development of cirrhosis.

Option D: Kawasaki disease (KD) is a systemic inflammatory disorder with arteritis of medium to large-sized vessels. It presents with fever, rash, conjunctivitis, and cervical lymph node enlargement.

Solution to Question 3:

The given image shows small red lesions with bluish-white spots in the center on the inner aspects of the cheek opposite to the lower premolar, indicating the presence of Koplik spots. The appearance of these spots, along with a rash appearing three days later, strongly suggests a diagnosis of measles.

The measles virus is a single-stranded RNA virus. Measles infection has 4 phases: the incubation period, prodromal illness, exanthematous phase (characterized by the rash), and recovery phase.

After an incubation period of 8-12 days, the prodromal phase of measles starts with mild fever and is followed by conjunctivitis, photophobia, coryza, cough, and increasing fever. Koplik spots, appearing 1-4 days prior to the rash, are a characteristic sign of measles.

Symptoms gradually intensify over a period of 2-4 days until the first day of the rash. The rash typically begins on the forehead (around the hairline), behind the ears, and on the upper neck as a red maculopapular eruption. It then spreads downward to the torso and extremities, often becoming confluent on the face and upper trunk.

Complications include acute otitis media (most common), pneumonia (the most common cause of death), croup, bronchiolitis, reactivation of tuberculosis, diarrhea, appendicitis, febrile seizures, and encephalitis. Subacute sclerosing panencephalitis is a rare chronic complication of measles.

Other options:

Option B: Leukoplakia appears as a white patch in the oral cavity that cannot be wiped off.

Option C: Scarlet fever is Group A Streptococcus (GAS) pharyngitis and is characterized by a distinctive sandpaper rash. The rash appears as a diffuse, finely papular, erythematous eruption

that turns the skin bright red and blanches on pressure. It starts around the neck and spreads to the trunk and extremities.

Option D: Kawasaki disease (KD) is a systemic inflammatory disorder with arteritis of medium to large-sized vessels. It presents with fever, rash (maculopapular, erythema multiforme urticarial, or micropustular), conjunctivitis, and cervical lymph node enlargement.

Solution to Question 4:

The given chest X-ray shows a boot-shaped cardiac silhouette which is suggestive of the Tetralogy of Fallot.

The boot-shaped (coeur en sabot) cardiac silhouette is seen due to the right ventricular hypertrophy and upward shift of the cardiac apex. The hilar areas and lung fields appear relatively clear due to reduced pulmonary blood flow or the small size of the pulmonary arteries, or both.

In Ebsteins's anomaly (Option D) box-shaped heart is seen.

Solution to Question 5:

The greenish-black color of the first stool in the newborn is due to biliverdin.

Meconium is a dark, viscous material that is normally present in the intestine of the fetus and passed within 1st 48hour of life. Meconium appears after the 20th week and at term, it is uniformly distributed throughout the gut up to the rectum indicating the presence of intestinal peristalsis.

It is composed of various products of secretion, such as desquamated fetal cells, glycerophospholipids from the lungs, lanugo, scalp hair, vernix, and undigested products from swallowed amniotic fluid. The greenish-black color of meconium is due to the presence of bile pigments excreted by the biliary tract, especially biliverdin (pigment from the oxidation of bilirubin).

During intrauterine hypoxia, the relaxation of the anal sphincter due to vagal stimulation can lead to the passage of meconium into the amniotic fluid. Additionally, the release of arginine vasopressin (AVP) from the fetal pituitary gland due to intrauterine hypoxia can stimulate the colon muscles to contract, resulting in intraamniotic defecation. Meconium aspiration syndrome can occur when meconium is inhaled, as it is toxic to the lungs.

Other options:

Option A: Bilirubin is responsible for the yellow color of the stools.

Option D: Neonates do not harbor gut flora at birth.

Solution to Question 6:

Maternal antibodies do not provide significant immunity to polio infection in a neonate.

Administering an oral polio vaccine (OPV) dose at birth can provide early mucosal protection, safeguarding against enteric pathogens before they can hinder the intestinal mucosal immune response. Furthermore, administering OPV during the period when infants are protected by maternal antibodies may help prevent vaccine-associated paralytic polio (VAPP).

All pregnant women should receive a tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis (Tdap) vaccine during each pregnancy, as early in the 27–36-week-of-gestation window as possible. The maternal antibodies formed from Tdap will provide immunity against tetanus, diphtheria, and pertussis in the newborn. Administering the Tdap vaccine earlier, between the 27th and 36th week of gestation, maximizes the transfer of passive antibodies to the infant (options A and C).

Infants less than 6 months are generally protected from measles by maternal antibodies (option D).

Solution to Question 7:

In the given clinical scenario, a 3-year-old child is likely to have received the 4th dose (1st booster) of the diphtheria vaccine between 16–24 months of age according to National Immunization Schedule (i.e., 16 months ago). Since he has received a booster dose within 5 years, only erythromycin is indicated in case of close contact with the diphtheria patient.

All household contacts with diphtheria patients should be closely monitored for 7 days. Culture should be taken from both the throat and nose. All contacts should receive antibiotic prophylaxis (benzathine penicillin G or erythromycin), regardless of immunization status.

Children who have not received a booster dose of the diphtheria toxoid vaccine within 5 years should be given the vaccine in an age-appropriate form. Children who have not yet received their 4th dose should be vaccinated accordingly.

If fewer than three doses of diphtheria toxoid have been given, or vaccination history is unknown, an immediate dose of diphtheria toxoid should be given and the primary series completed according to the current schedule.

Asymptomatic carriers of diphtheria are treated with antimicrobial prophylaxis for a period of 10-14 days. If an individual has not received a booster dose of diphtheria toxoid within 1 year, an age-appropriate preparation of the vaccine should be administered promptly.

Droplet precautions (respiratory colonization) or contact precautions (cutaneous colonization) of cases and carriers are maintained until two consecutive negative cultures (collected 24 hours apart) are obtained after cessation of therapy.

Solution to Question 8:

According to current dietary guidelines for children, energy from saturated fats should be <10% of total energy intake. Diets that are low in saturated fat and cholesterol (<300mg/day) without transfat are preferred.

Acceptable macronutrient distribution ranges (AMDA % of energy) are given below:

Macronutrient	Age1 to 3years	Age4 to 18years
Fats	30 to 40	25 to 35
Protein	5 to 20	10 to 30
Carbohydrate	45 to 65	45 to 65

Solution to Question 9:

The given clinical scenario of a girl (46XX) with an enlarged clitoris is highly suggestive of congenital adrenal hyperplasia (CAH) due to 21-hydroxylase deficiency.

Congenital adrenal hyperplasia/adrenogenital syndrome is a family of autosomal-recessive disorders caused by the deficiency of enzymes involved in the synthesis of steroid hormones in the adrenal cortex. The majority of cases of CAH are caused by 21-hydroxylase deficiency. This deficiency leads to a lack of synthesis of cortisol and aldosterone, resulting in the salt-wasting form of the disease, which is the most severe form.

Severe 21-hydroxylase deficiency presents with symptoms including weight loss, dehydration, weakness, hypotension, and electrolyte imbalances. Accumulated steroid precursors like 17-hydroxyprogesterone, are shunted into the pathway of androgen synthesis leading to elevated levels of androstenedione and testosterone. This can result in abnormal genital development in affected female fetuses during early gestation.

Females may have masculinized external genitalia. This can include enlargement of the clitoris, partial or complete labial fusion, and a shared opening of the vagina and urethra (urogenital sinus).

Untreated individuals develop signs of androgen excess after birth. These signs include accelerated skeletal maturation, premature epiphyseal closure leading to stunted adult stature, excessive muscular development, the appearance of pubic and axillary hair, acne, and deepening of the voice. In males, enlargement of the penis, scrotum, and prostate. In females, the clitoris may become further enlarged.

Diagnosis of confirmed by elevated 17-hydroxyprogesterone levels. Glucocorticoid replacement with hydrocortisone is done. In a salt-wasting crisis, fludrocortisone supplementation is given. Vaginoplasty and clitoral recession are done for females with ambiguous genitalia.

Solution to Question 10:

The most probable diagnosis for the evisceration of bowel loops, coming out of a defect in the abdominal wall to the right of the umbilicus is gastroschisis. It occurs when there is a failure in the closure of the body wall in the abdominal region.

Gastroschisis is an abdominal wall defect located on one side of the umbilicus (right side commonly) allowing protrusion of the intestines, stomach, bladder, and sometimes ovaries or

undescended testes. The loops of the bowel are not covered by amnion as they herniate through the abdominal wall directly into the amniotic cavity. The affected bowel loops can be damaged by exposure to amniotic fluid.

Risk factors include teenage pregnancy, recreational drug use, smoking, and genitourinary infection during pregnancy. After birth, careful management is required to prevent twisting or kinking of the blood supply to the intestines. The abdomen is wrapped with transparent plastic wrap, a nasogastric tube is inserted, and fluid resuscitation is initiated.

Depending on the severity, primary closure under general anesthesia or gradual reduction using a preformed silo may be performed. Parenteral nutrition is often needed for several weeks to support intestinal recovery.



Other options:

Option A: In omphalocele bowel loops fail to return from the physiologic umbilical hernia. Since the umbilicus is covered by a reflection of the amnion, the omphalocele is also covered by the amnion.



Option B: Congenital umbilical hernia occurs due to the incomplete closure or weakness of the muscular umbilical ring. It presents as a soft swelling covered by skin that protrudes during activities like crying, coughing, or straining, and it can be easily pushed back through the fibrous ring at the umbilicus. The hernia contains the omentum or segments of the small intestine.

The image below shows an umbilical hernia:

Option D: Ectopia vesicae/bladder exstrophy is a congenital abnormality that occurs when the lower abdominal wall does not form properly and the bladder protrudes from the abdominal wall, with its mucosa exposed.

The image below shows ectopia vesicae/bladder exstrophy

Paediatrics INI-CET 2021

Question 1:

Which of the following is a true statement about congenital CMV infection?

a) About 20-30% of the infections are symptomatic

b) Triad of SNHL, periventricular calcification and enamel hypoplasia

c) Most children asymptomatic at birth can develop conductive hearing loss later in life

d) If mother is IgG positive for CMV, the child is unlikely to develop infection

Question 2:

Which of the following are components of kangaroo mother care?

a) Early discharge, skin to skin contact, prevent infection

b) Early discharge, skin to skin contact, exclusive breastfeeding

c) Skin to skin contact, exclusive breastfeeding, prevent infection

d) Early discharge, exclusive breastfeeding and prevent infection

Question 3:

Which of the following is given as a part of the therapy for a child with severe COVID?

a) Corticosteroids

b) Ivermectin

c) Remdesivir

d) All of the above

Question 4:

Which of the following is seen in physiological jaundice?

a) Increased indirect bilirubin, increased urobilinogen, absent bilirubin in urine

b) Increased indirect bilirubin, absent urobilinogen, absent bilirubin in urine

c) Increased direct bilirubin, increased urobilinogen, absent bilirubin in urine

d) Increased direct bilirubin, absent urobilinogen, absent bilirubin in urine

Question 5:

A child presents with fever and vesicular lesions on the upper limb and the lower limb. Neck stiffness was present. Similar lesions were present on the palms, soles, and oral cavity. CSF analysis revealed normal glucose levels and elevated lymphocytes and protein. What is the most likely diagnosis?

a) Bacterial meningitis with sepsis

b) Coxsackie viral meningitis and Hand Foot Mouth disease

c) Herpes simplex gingivostomatitis and meningoencephalitis

d) Tuberculous meningitis

Question 6:

Match the following developmental milestones:

a) 1-a, 2-b, 3-c, 4-d

b) 1-b, 2-a, 3-d, 4-c

c) 1-c, 2-b, 3-a, 4-d

d) 1-d, 2-c, 3-b, 4-a

Question 7:

Which of the following congenital heart disease has equal saturation in all the chambers of the heart?

a) Tetralogy of Fallot

b) Total anomalous pulmonary venous circulation

c) Tricuspid atresia

d) Transposition of great arteries

Question 8:

Which of the following marked structure is damaged in respiratory distress syndrome?



a) 1

b) 2

c) 3

d) 4

Question 9:

Which of the following are causes of neonatal seizures ?

a) 1, 2, 3 only

b) 1, 2, 3, and 4

c) 1, 3, 4 only

d) 2, 4 only

Question 10:

A neonate is found to have increased irritability, poor feeding, and 2 episodes of seizures. Lumbar puncture was done and the findings were suggestive of meningitis. What is the most likely causative organism?

a) Escherichia coli

b) Streptococcus agalactiae

c) Neisseria meningitidis

d) Listeria monocytogenes

Question 11:

At what age does a child attain half-height of the adult height?

a) 12 - 18 months

b) 20 - 24 months

c) 32 - 36 months

d) 40 - 48 months

Answer Key

Question No.	Correct Option
1	d
2	b
3	a
4	a
5	b
6	b
7	b
8	b
9	b
10	a
11	b

Detailed Explanations

Solution to Question 1:

If the mother is IgG positive for CMV, then the child is unlikely to develop the infection.

In one-third of mothers who develop a primary infection during pregnancy, the virus gets transmitted to the fetus. In previously infected mothers with IgG antibodies, only 1-3% would transmit the virus to the fetus.

Most CMV transmission occurs in the first trimester. The risk of transmission is high in the third trimester. Only 10% of infants with congenital CMV infection are symptomatic (option A).

Hepatosplenomegaly, jaundice, and petechiae constitute the clinical triad (option B) in congenital CMV. Neuroimaging shows intracranial periventricular calcifications. Other manifestations include IUGR, hydrops, microcephaly, pneumonitis, chorioretinitis, seizures, sensorineural hearing loss, abnormal dentition, and hypocalcified enamel.

About half of the symptomatic infants develop long-term sequelae which include sensorineural hearing loss (SNHL), vision loss, seizure disorders, mental retardation, and developmental delay. But in asymptomatic infants, only about 13.5% develop long-term sequelae which is commonly SNHL (option C).

The investigation of choice is PCR or viral culture. It should be done within the first two to three weeks of life. After this period, it is difficult to distinguish between a congenital and an acquired infection. Since not all affected infants are viremic at birth, urine and/or saliva are more reliable samples than blood.

There is no definitive treatment for congenital CMV infection. Treatment with ganciclovir is indicated only in patients with progressive neurologic disease and deafness.

Solution to Question 2:

The four components of kangaroo mother care (KMC) are skin-to-skin positioning of the baby on the mother's (or caretaker's) chest, exclusive breastfeeding, early discharge and ambulatory care, and support for the mother and her family for taking care of the baby.

Eligibility criteria for KMC include all hemodynamically stable low birthweight (LBW) babies. All mothers without any serious illness can provide KMC. Short sessions of KMC can be initiated in babies on medical treatment like IV fluids or oxygen therapy.

The initiation of KMC can differ depending on the weight of the newborn. Babies weighing less than 1200 g at birth may require several days to weeks before KMC can begin. For babies weighing between 1200 g and 1800 g, the initiation of KMC may only take a few days. KMC can be started immediately after birth for babies weighing over 1800 g.

The baby should be placed between the mother's breasts in an upright position, with the face turned to one side, The hips are abducted and flexed in a 'frog position' and the arms are also flexed. The abdomen of the baby should be at the level of the mother's epigastrium. In this position, the mother's breathing would stimulate the baby and prevent the development of apnea.

KMC is not necessary once the baby attains 2500 g of weight or 37 weeks of gestation.

The image below shows proper KMC positioning:

The duration of KMC should be gradually increased from short sessions of more than one hour to 24 hours a day.

KMC is comparable to incubators in terms of safety and thermal protection as evidenced by mortality rates. Additionally, it has been shown to reduce severe morbidity by facilitating breastfeeding.

KMC promotes the bonding between mother and child, fostering a strong emotional connection during the early stages of life.

Note: Incubatory care, feeding via nasal catheter, and prevention of infection are part of intensive care for low birthweight babies.

Solution to Question 3:

Among the given options, only corticosteroids are given for children with severe COVID-19.

Severe COVID-19 infection is defined as SpO2 <90% on room air with any of the following findings: acute respiratory distress syndrome (ARDS), signs of severe pneumonia, septic shock, multi-organ dysfunction syndrome (MODS) or pneumonia with cyanosis, grunting, severe chest retraction, lethargy, somnolence, seizures.

It is important to note that CT chest is not indicated in the diagnosis or management of COVID-19 in children and is only considered when there is no improvement in the respiratory status.

Regarding the management of severe COVID-19 infection, the Ministry of Health and Family Welfare (MoHFW) guidelines mention that, in mild or asymptomatic COVID-19 infection, steroids are avoided and are found to be harmful. However, in rapidly progressive moderate or severe cases, corticosteroids are used in the treatment.

This can be administered in two ways: Dexamethasone at a dose of 0.15 mg/kg, with a maximum dose of 6 mg, once a day, or Methylprednisolone at a dose of 0.75 mg/kg, with a maximum dose of 30 mg, once a day.

Steroids should be given for 5-7 days, with a gradual tapering off over 10-14 days, and are avoided within the first 3-5 days of symptom onset, as they prolong viral shedding.

The use of antivirals (option C) or monoclonal antibodies is not recommended for children less than 18 years of age, irrespective of the severity of the infection.

Antimicrobials should not be prescribed unless there is clinical suspicion of a superadded infection or the patient is in septic shock.

Anticoagulants are not indicated routinely, except in patients having a strong personal or family history of thromboembolism or more than four high-risk factors for developing thrombosis.

Solution to Question 4:

In physiological jaundice, there is increased indirect bilirubin, increased urobilinogen, and absent bilirubin in urine.

Physiological jaundice (icterus neonatorum) is due to increased bilirubin production from the breakdown of fetal red blood cells combined with transient limitation in the conjugation of bilirubin by the immature neonatal liver. It first appears on the 2nd or 3rd days usually peaks on the 4th day at 5-6 mg/dL and decreases to <2 mg/dL between the 5th and 7th days after birth.

Physiological jaundice would lead to an increase in indirect (unconjugated) bilirubin. Increased levels of unconjugated bilirubin in the blood lead to increased production of urobilinogen which would be present in the urine. Urine bilirubin is not usually found in this condition.

Risk factors for physiological jaundice include maternal age, maternal diabetes, prematurity, drugs (vitamin K3, novobiocin), polycythemia, male sex, trisomy 21, cutaneous bruising, blood extravasation as in cephalhematoma, breastfeeding, weight loss, and a family history of physiological jaundice.

On the other hand, in pathological jaundice, the appearance of jaundice is within the first 24 hours after birth and may be present at or beyond the age of 2 weeks in term babies and 3 weeks in preterms babies. Other signs of pathological jaundice include an increase in serum bilirubin exceeding 5mg/dL within a 24-hour period, a serum bilirubin level greater than 12mg/dL in a full-term infant or between 10-14mg/dL in a preterm infant, and a direct bilirubin level more than 2mg/dL at any given time. Jaundice visible on the palms and soles, unresponsiveness to phototherapy, and a rapid rise in bilirubin levels during phototherapy indicate pathological jaundice.

When the concentration of unconjugated bilirubin exceeds 20-25mg/dL, bilirubin would penetrate the blood-brain barrier and lead to kernicterus.

Phototherapy is initiated in babies with pathological jaundice when the bilirubin levels exceed the 95th percentile on the total serum bilirubin (TSB) normogram. Babies are kept at a distance of 30-45 cm from the UV lamp (460-490 nm) with eyes and gonads covered. It is avoided in babies with conjugated hyperbilirubinemia due to the risk of bronze-baby syndrome. Dehydration and hypocalcemia are potential complications and breastfeeding is encouraged.

Exchange transfusion (ET) is done in cases of bilirubin encephalopathy, Rh incompatibility with significant hemolysis, and bilirubin values exceeding TSB cutoffs for ET.

Solution to Question 5:

The presence of characteristic vesicular lesions on the palms, soles, and oral cavity is suggestive of Hand-foot-and-mouth disease (HFMD) caused by the coxsackie virus. The presence of neck stiffness and CSF showing elevated lymphocytes with normal glucose and elevated protein levels are suggestive of viral meningitis. Coxsackie A viruses can lead to aseptic meningitis.

HFMD is a viral infection commonly caused by coxsackievirus A16, which belongs to the subgroup of enteroviruses. In larger outbreaks, other viruses such as enterovirus 71, coxsackie A viruses 5, 6, 7, 9, 10, and coxsackie B viruses 2 and 5 can also cause the disease.

HFMD typically presents as a mild illness with vesicular lesions on the tongue, posterior pharynx, buccal mucosa, palate, and lips. Lesions may also appear on the hands, fingers, buttocks, and groin, with a higher prevalence on the hands. These vesicles usually resolve within a week.

Infections with enterovirus 71 can be more severe with neurological and cardiopulmonary manifestations. Coxsackievirus A16 infections occasionally lead to complications such as encephalitis, acute flaccid paralysis, myocarditis, pericarditis, and shock. Coxsackievirus A6 can cause a severe and atypical form of HFMD.

Diagnosing HFMD typically relies on clinical examination of the characteristic lesions. Confirmatory testing can involve viral culture using samples such as blood, cerebrospinal fluid (CSF), urine, and mucosal swabs. However, RT-PCR testing is more sensitive and specific for detecting the viral genome even when cultures are negative.

Treating HFMD involves mainly supportive care as there is no specific proven antiviral agent for enteroviruses. In cases of enterovirus 71-related cardiopulmonary disease, milrinone is suggested as an effective treatment. Immunoglobulins have been utilized to manage severe cases, although outcomes vary.

Apart from HFMD, coxsackie A viruses can cause herpangina characterized by febrile vesicular pharyngitis with ulcerating small vesicles on the oral fauces, palate, uvula, and posterior pharyngeal wall.

On the other hand, coxsackie B viruses can cause pleurodynia or Bornholm disease, myocarditis, and pericarditis in newborns, as well as orchitis and juvenile diabetes.

Other options:

Option C: Herpes simplex virus (HSV) presents similarly with vesicular lesions but lacks typical herpetic signs. The absence of atypical symptoms like mouth pain, drooling, and characteristic herpetic lesions rule out HSV gingivostomatitis. HSV-induced encephalitis causes anosmia, focal

seizures and behavioural changes due to cortex or limbic system damage.

The image shows a typical HSV vesicular lesion below:

Option A: Bacterial meningitis with sepsis is unlikely given the vesicular lesions' distribution pattern (bacteria are less likely to cause vesicles in the oral cavity). Moreover, CSF findings in bacterial meningitis usually show decreased glucose, elevated neutrophils, and elevated protein levels.

Option D: Tuberculous meningitis is less likely as there is no history of TB exposure provided. In most cases, CSF findings in tuberculous meningitis demonstrate elevated lymphocytes and protein levels, along with decreased glucose.

Solution to Question 6:

The correctly matched developmental milestones are as follows:

1.Transfers objects	b. 6-7 months
2.Social smile	a. 1-2 months
3. Pincer grasp	d. 9-12 months
4. Walks 1-2 steps	c. 12-15 months

Solution to Question 7:

In total anomalous pulmonary venous circulation (TAPVC), there is equal saturation in all chambers of the heart.

In TAPVC, the oxygenated blood from the pulmonary veins reaches the right atrium (having deoxygenated blood) instead of the left atrium. It occurs due to the draining of the pulmonary veins to the superior vena cava or to the inferior vena cava or to both. This is a total mixing lesion. The blood from this right atrium (RA) passes to the right ventricle and pulmonary artery. This blood also reaches the left atrium through the atrial septal defect or the patent foramen ovale, and subsequently, to the left ventricle The left atrium is not receiving any blood because all of the pulmonary veins are draining to the right atrium. Therefore, this shunt will be the only source of blood to it. Since blood from right atrium reaches all the other chambers, the oxygen saturation of the blood in all four chambers is the same.



In (TAPVC), there are three main types: supra-cardiac, cardiac, and infra-cardiac. In supra-cardiac TAPVC, the pulmonary veins merge to form a single vein that drains into either the superior vena cava or the left innominate vein. In cardiac TAPVC, the pulmonary veins directly drain into the right atrium or coronary sinus. The most dangerous type is infra-cardiac TAPVC, which is always obstructive. In this type, the pulmonary veins drain into either the portal vein or the inferior vena cava.

Diagnosis of TAPVC is the demonstration of any vein with doppler flow away from the heart because the normal venous flow is usually toward the heart.

Surgical correction of TAPVR is done in infancy. Pulmonary venous confluence is anastomosed directly to the left atrium, the ASD is closed, and any connection to the systemic venous circuit is interrupted.

Other options

Option A - Tetralogy of Fallot (TOF) comprises right ventricular outflow tract obstruction, ventricular septal defect (VSD), overriding of the aorta over the ventricular septum, and right ventricular hypertrophy. Right ventricular outflow tract obstruction increases the pressure in the right ventricle. This causes right to left shunt through the ventricular septal defect. The deoxygenated blood from the right ventricle mixes with the oxygenated blood in the left ventricle. Therefore, in this condition, the left atrium has a higher oxygen saturation compared to the left ventricle due to mixing. The right atria and ventricles have the least oxygen saturation.



Option C - Tricuspid atresia is the congenital absence of the tricuspid valve. Therefore, the blood in the right atrium is shunted to the left atrium through the patent foramen ovale or the atrial septal defect. In the left atrium, the blood from the right atrium mixes with the oxygenated blood and the oxygen saturation decreases. However, the oxygen saturation in the left atrium and left ventricle is still greater compared to that of the hypoplastic right ventricle.



Option D - Transposition of great arteries (TGA) is characterized by the aorta arising from the right ventricle and the pulmonary artery from the left ventricle. Because of this, the pulmonary and systemic circulations remain separate. The right side of the heart has deoxygenated blood and the left side of the heart has oxygenated blood. The pulmonary artery saturation is always higher than the aortic saturation.



Solution to Question 8:

Type II pneumocytes, which are marked as 2 in the above image, are damaged in respiratory distress syndrome.

The cells forming the lining epithelium of alveoli (also called pneumocytes) are mainly of the following types:

• Type I pneumocytes - large flattened cells which present a very thin diffusion barrier for gases. These cells have large cytoplasmic extensions and form the primary lining cells of the alveoli. They are connected to each other by tight junctions.

• Type II pneumocytes (granular pneumocytes) - These are rounded secretory cells bearing microvilli on their free surfaces. These cells secrete the surfactant which decreases the surface tension between the thin alveolar walls and thus prevents alveoli from collapsing during respiration. Type II cells are important in alveolar repair as well.

The below image shows the pneumocytes lining the alveoli:



©MARROW

The alveoli are surrounded by pulmonary capillaries. The alveoli also contain other specialized cells, including pulmonary alveolar macrophages, lymphocytes, plasma cells, neuroendocrine cells, and mast cells.

Respiratory distress syndrome (RDS) is seen in premature infants, due to the decreased production and secretion of the lung surfactant. The surfactant reduces the alveolar surface tension. This phospholipid is synthesized in type II pneumocytes (also called granular pneumocytes) in the endoplasmic reticulum.

The surfactant appears in the amniotic fluid between 28 and 32 weeks of gestation. Mature levels of the pulmonary surfactant are present usually after 35 weeks of gestation. In a premature infant, there is a decrease in both the quantity and quality of surfactant resulting in respiratory distress syndrome.

Administration of antenatal glucocorticoids to mothers who deliver at 24 to 34 weeks can increase surfactant production and reduce the risk of RDS. It includes either giving 2 doses of betamethasone 12mg/dose with 24 hours between each dose, or 4 doses of dexamethasone 6mg/dose, with 12 hours between each dose.

Mild to moderate RDS can be managed with continuous positive airway pressure (CPAP). CPAP is a noninvasive modality of support where a continuous distending pressure (5-7 cm H20) is applied at the nostril level to keep the alveoli open in a spontaneously breathing baby. Severe cases will require mechanical ventilation.

Surfactant administration is reserved for severe cases and moderate cases not improving even with fiO2> 40% in CPAP.

With treatment, it rarely leads to respiratory failure and has a good prognosis.

Solution to Question 9:

All of the above-given options, hyponatremia, hypomagnesemia, hypocalcemia, and hypernatremia are causes of neonatal seizures.

The causes of neonatal seizure according to the age of presentation are summarised in the table below:

Hypoxic-ischemic encephalopathy is the most common cause. Metabolic disorders cause neonatal seizures commonly. Hypoglycemia is the most common metabolic cause of neonatal seizures.

Neonatal seizures may be of different types - subtle, tonic, clonic, spasms, and myoclonic.

Subtle seizures are the most common type of neonatal seizures. They involve movements like transient eye deviations, nystagmus, blinking, mouthing, cycling, and apnea, and are more common in premature infants.

Tonic seizures are characterized by increased muscle tension, either focal (sustained posturing of a single limb) or generalized (bilateral tonic limb extensions or flexions of the upper limbs with tonic extension of the lower limbs and trunk). They can not be provoked by stimulation or suppressed by restraint.

Clonic seizures involve repetitive contractions of muscle groups. Spasms are sudden generalized jerks associated with brief discharges.

Myoclonic seizures are the least common type of neonatal seizures and are rapid jerks (<50 msec) without rhythmicity.

Electroencephalogram (EEG) is the preferred diagnostic investigation for seizures, while MRI is the recommended neuroimaging modality for etiological evaluation. Phenobarbital is the drug of choice for treatment. In term neonates, hypoxic-ischemic encephalopathy (HIE) is the most common cause of seizures, while intraventricular hemorrhage (IVH) is more prevalent in preterm neonates.

1 to 4 days	4 to 14 days	2 to 8 weeks
Hypoxic ischemic encephalopathy	Bacterial meningitisEncephalitis (enteroviral, herpes simplex)	Encephalitis (enteroviral, herpes simplex)Bacterial meningitis
Drug withdrawal, maternal drug use	HypocalcemiaHypoglycemia	Subdural hematomaChild abuse
Drug toxicity: lidocaine, penicillin	GalactosemiaFructosemia	AminoaciduriaUrea cycle defects
Intraventricular hemorrhage		Malformations of cortical development
HypocalcemiaSepsisHypoglycemiaHypomagnesemiaHypo or hypernatremia		Tuberous sclerosis
GalactosemiaUrea cycle disorders		Sturge-Weber syndrome

Solution to Question 10:

The given clinical scenario describes neonatal meningitis. E. coli > Streptococcus agalactiae, part of group B streptococci, are the most likely causative organism of neonatal meningitis.

Early features of meningitis include temperature instability (hypo or hyperthermia), lethargy, bradycardia, and feed intolerance. Late features include seizures, bulging fontanelle, and nuchal rigidity.

The most common cause of bacterial meningitis in a 0 to 2-month baby is group B streptococci (Option B) worldwide and gram-ve bacteria like E.coli and Klebsiella (option A) in India. In babies aged 2 to 24 months, pneumococci are the primary causative pathogens, followed by H. influenzae. For children older than 2 years, pneumococci remain a prominent cause of meningitis, with Neisseria meningitidis being the second most common pathogen.

The most common etiological agents of neonatal meningitis and sepsis are listed in the table below:

Neonates have immature cellular and humoral immunity. This makes them prone to meningitis and sepsis. Thus, an infant with a fever who is 28 days or younger, should be evaluated for sepsis.

Option C: Meningococcal infections are usually asymptomatic. If invasive disease occurs, then the clinical spectrum varies widely from meningococcal meningitis (highest proportion) to bacteremia without sepsis, meningococcal septicemia with or without meningitis, pneumonia, chronic meningococcemia, and occult bacteremia.

Option D: Meningitis caused by Listeria monocytogenes is rare but clinically significant. Infants can acquire the infection in utero or through an infected birth canal. Neonatal listeriosis can manifest as early-onset disease, typically within 1-2 days of birth, primarily as a septicemic form. Alternatively, it can present as a late-onset disease in infants older than 5 days, predominantly as a meningitic form.

	India	Worldwide
Neonatal meningitis	E. coli, Klebsiella	Group BStreptococci
Neonatal sepsis	Klebsiella,S.aures,E. coli	Group BStreptococci

Solution to Question 11:

A child attains half of the adult height by 20 -24 months.

The normal length at birth is 50 cm, increasing to around 75 cm at 1 year of age. By the end of the second year, the length ranges from 85 to 90 cm.

From age 2 to around 12 years, there is a steady growth of about 6 cm per year. During pubertal growth spurts after age 12, females grow an average of 8 cm per year, while males grow around 10 cm per year.

By around 2 years of age, a normal child reaches half of their adult height (around 80-85 cm). Height doubles by age 4 and triples by age 12 compared to the birth length.

Between 18-24 months, a child experiences significant growth and development.

Physically, they improve their balance, learn to run and climb stairs, and reach about half of their adult height. Head growth slows slightly, with 85% of adult head circumference achieved by the

age of 2 years.

Cognitive development includes the establishment of object permanence and problem-solving skills (e.g., using a stick to obtain a toy that is out of reach).

Emotionally, they display self-recognition and recognize when toys are broken. When toddlers see anything abnormal in their noses while gazing into a mirror for the first time, they tend to touch their own faces rather than reaching out toward the reflected image.

Linguistically, their vocabulary expands from 10-15 words to 50-100 words, and they start combining words to form simple sentences while understanding two-step commands (eg., give me the ball and then get your shoes).

Paediatrics NEET 2021

Question 1:

A child with infantile hypertrophic pyloric stenosis has multiple episodes of non-bilious vomiting. Which of the following metabolic abnormality is likely to be seen in this condition?

a) Hypochloremic, hypokalemic acidosis

b) Hypochloremic, hypokalemic alkalosis

c) Hypokalemic hyperchloremic alkalosis

d) Hyperchloremic hypokalemic acidosis

Question 2:

A 6-year-old girl born out of nonconsanguineous marriage presented with rachitic changes in limbs since 1½ years of age for which she has received multiple courses of vitamin D. Investigations showed the following. What is the most likely diagnosis?

a) Vitamin D dependent rickets

b) Hypoparathyroidism

c) Chronic renal failure

d) Hypophosphatemic rickets

Question 3:

During the anthropometric evaluation, the height for age of a child is < - 2SD. This is most likely due to?

a) Chronic malnutrition

b) Acute malnutrition

c) Recent infection

d) No malnutrition

Question 4:

A child presents with generalised edema and was found to have hypercholesterolemia. Urine analysis shows protein 3+ and fat bodies. What is the likely diagnosis?

a) Nephrotic syndrome

b) Nephritic syndrome

c) Goodpasture's disease

d) IgA nephropathy

Question 5:

A young child with congenital hydrocephalus was treated with a successful ventriculoperitoneal (VP) shunt. Four months post surgical repair, the child presented with fever, nuchal rigidity, and irritability. You suspect meningitis. What is the next best step?

a) Wait and watch

b) Check shunt patency by nuclear study

c) Blood culture and take CSF sample from shunt tap

d) Blood culture and take CSF sample by lumbar puncture

Question 6:

In fetal circulation, which vessel carries deoxygenated blood back to the placenta?

a) Umbilical Vein

b) Umbilical artery

c) Descending aorta

d) Pulmonary Artery

Question 7:

A 10-year-old boy presented with seizures. His past history is significant for an episode of fever with rash at 1 year of age which resolved spontaneously. What is the most helpful investigation to diagnose his condition?

a) IgG measles in CSF

b) MRI mesial temporal lobe sclerosis

c) IgM measles in CSF

d) C1Q4 antibodies in the CSF

Question 8:

A 16-month-old boy is found to weigh 8kg. A WHO growth chart(weight-for-age) is provided below. Plot the growth chart for this child. What is the most appropriate advice based on the plotted graph?



a) No malnutrition - Assure the mother

b) Mild malnutrition - home treatment

c) Severe malnutrition - refer to nutrition rehabilitation center

d) Moderate malnutrition - teach the mother on how to feed

Answer Key

Question No.	Correct Option
1	b
2	d
3	a
4	a
5	c
6	b
7	a

8	d

Detailed Explanations

Solution to Question 1:

Multiple episodes of non-bilious vomiting in infantile hypertrophic pyloric stenosis (IHPS) lead to hypochloremic hypokalemic alkalosis.

Emesis causes loss of gastric fluid containing H+ ions, leading to the release of bicarbonate into the bloodstream. Without normal neutralization in the small intestine, metabolic alkalosis develops. This is the generation phase of alkalosis.

Volume depletion during vomiting hinders the urinary excretion of bicarbonate. This is due to reduced glomerular filtration, increased proximal tubule resorption of sodium and bicarbonate, and elevated aldosterone levels promoting bicarbonate resorption and hydrogen ion secretion in the collecting duct. This is the maintenance phase.

Loss of chloride ions in the gastric acid leads to hypochloremia.

Hydrogen ions exit from the cells in exchange for potassium leading to hypokalemia. Loss of fluid due to excessive vomiting activates the RAAS pathway. This worsens hypokalemia as the aldosterone causes active reabsorption of sodium in exchange for the secretion of K+ and H+ ions into the urine. The loss of H+ ions in urine lowers the urinary pH, resulting in paradoxical aciduria.

Infantile hypertrophic pyloric stenosis (IHPS) is associated with hypertrophy of the circular muscle of the pylorus resulting in constriction and obstruction of the gastric outlet. It is common between 3 and 6 weeks of age. First-born male infants are at the highest risk.

Infants usually present with projectile, non-bilious vomiting. Clinically, an olive-shaped mass is palpable in the right upper quadrant or epigastrium. When the infant is given a test feed, visible peristalsis is usually observed moving from left to right (not from right to left), followed by projectile vomiting. It is associated with hypokalemic hypochloremic metabolic alkalosis with paradoxical aciduria.

Ultrasound examination is the investigation of choice for this condition. The diagnostic criteria for pyloric stenosis are a muscle thickness of ≥4 mm and a pyloric length of ≥16 mm. The nipple sign is produced by the double-layered hypertrophic mucosa that protrudes into the stomach. The sonographic appearance of the hypertrophied pylorus resembles that of the uterine cervix and is therefore called the cervix sign.

The image below shows the cervix and nipple sign seen in IHPS:



Management includes the correction of metabolic abnormality prior to surgery. Administering 5% dextrose and 0.45% saline with added potassium at a rate of 150-175 mL/kg per 24 hours can correct the fluid deficit in infants. Monitoring urine output (>2 cc/kg/h) is important to ensure successful rehydration.

The hypertrophic pylorus is surgically corrected by Ramstedt's pyloromyotomy (open or laparoscopic approach). It involves splitting the pyloric muscle while preserving the underlying submucosa.

Solution to Question 2:

A history of resistant rickets with normal levels of calcium (9 to 11 mg/dL), parathyroid hormone (PTH), arterial blood gas (ABG), with decreased phosphate (normal - 2.5 to 4.3 mg/dL), and elevated levels of alkaline phosphatase (ALP) (normal, 33 to 96 U/L) in a child is suggestive of hypophosphatemic rickets.

Hypophosphatemic rickets:

X-linked dominant hypophosphatemic rickets (XLH) is the most common genetic cause of rickets while the most common, overall is nutritional vitamin D deficiency. Mutations in the PHEX gene (phosphate-regulating endopeptidase on x chromosome) leads to increased FGF-23 (phosphatonin), which inhibits phosphate reabsorption in the proximal tubule levels and hence, causes increased excretion of phosphate. It also inhibits renal 1-alpha hydroxylase leading to reduced 1,25 dihydroxy vitamin D.

Causes of hypophosphatemic rickets include familial vitamin D resistance, autosomal dominant hypophosphatemic rickets, autosomal recessive hypophosphatemic rickets, tumor-induced rickets, McCune Albright syndrome, epidermal nevus syndrome, and neurofibromatosis.

In X-linked dominant hypophosphatemic rickets lower extremity abnormalities, short stature, poor growth, delayed dentition, and tooth abscesses are common.

The chart attached below summarizes the evaluation of hypophosphatemic rickets.



Patients with hypophosphatemic rickets respond well to oral phosphorus and 1,25-D (calcitriol). Phosphorus supplementation is usually given in 1-3 g elemental phosphorus divided into 4–5 doses. Frequent dosing reduces diarrhea, which is a complication of high-dose oral phosphorus.

Calcitriol is administered 30–70 ng/kg/day divided into 2 doses. Burosumab, a monoclonal antibody to FGF-23 can be used in children > 1 year of age with XLH rickets.

Further, Joulie's solution is also useful in the treatment of familial hypophosphatemic rickets. Joulie's solution includes contains 30.4 mg of phosphate/mL. The most common side effect is diarrhea.

Other options:

Option A: Vitamin D-dependent rickets type 1A is due to mutation in the gene coding for 1α-hydroxylase leading to reduced calcitriol levels (1,25-D). Type 1B is due to a mutation in the gene for 25-hydroxylase leading to reduced calcidiol (25-D). Type 2A have mutations in the gene coding for vitamin D receptor. Type 2B is due to the over-expression of a hormone response element-binding protein that interferes with the actions of 1,25-D. All of them have normal to low calcium levels, low phosphorus, elevated PTH and ALP levels.

Option B: Hypoparathyroidism will present with features of hypocalcemia. There will be a decrease in serum PTH levels and calcium levels.

Option C: In chronic renal failure, 1α-hydroxylase activity is reduced leading to reduced 1,25-D. They have normal to low calcium levels and elevated PTH and ALP levels. They have hyperphosphatemia due to decreased renal excretion of phosphate.

The image below summarizes the common features of rickets in children.



Solution to Question 3:

Low height for the age (height for age < -2SD) in children indicates chronic malnutrition and stunting.

Classification of nutritional status depending on the anthropometry (height/length for age):

Nutritional status is often assessed in terms of anthropometry:

• Low weight for height, or wasting, indicates acute malnutrition.

• High weight-for-height indicates overweight.

• Low height-for-age, or stunting, indicates chronic malnutrition. This is the best indicator of long-term nutritional status in the pediatric population.

• Low weight-for-age indicates acute or chronic malnutrition. It does not differentiate between wasting and stunting. Hence, it has limited clinical significance.

Mid-upper arm circumference (MUAC) is an age-independent parameter used for the screening of wasting in children. In children aged 6 months to 5 years, MUAC <11.5cm is classified as severe malnutrition.

Classification	Criteria
Moderatelystunted (or) mode rate chronic malnutrition	Height/length-for-age< (–2 S D)and> (–3 SD)
Severelystunted (or) severe c hronic malnutrition	Height/length-for-age< (–3 S D)



Solution to Question 4:

The clinical scenario showing a child with generalized edema, hyperlipidemia, massive proteinuria, and fat bodies in urine is highly suggestive of nephrotic syndrome.

Nephrotic syndrome is characterized by nephrotic range proteinuria/heavy proteinuria >3.5 g/24 hr or urine protein: creatinine ratio > 2 and is associated with a triad of hypoalbuminemia (≤2.5 g/dL), edema, and hyperlipidemia (cholesterol > 200 mg/dL).

Nephrotic syndrome in children is primarily idiopathic, with minimal change disease being the most common glomerular lesion. Podocyte injury or genetic mutations in podocyte proteins can lead to nephrotic-range proteinuria.

Edema is the most common presenting symptom. They have elevated levels of cholesterol, triglycerides, low-density lipoproteins (LDL), and very low-density lipoproteins (VLDL). It is caused by both increased lipid synthesis and decreased lipid catabolism.

They are prone to infections especially by encapsulated bacteria due to factors such as urinary losses of immunoglobulin G (IgG) and complement factors (C3, C5, factors B and D). Nephrotic syndrome is a hypercoagulable state due to vascular stasis, increased platelet, increased hepatic production of fibrinogen, and urinary losses of antithrombotic factors such as antithrombin III and protein S.

The diagnosis is confirmed by performing a urinalysis and urine protein: creatinine ratio. The recommended initial treatment for nephrotic syndrome is a daily dose of 2mg/kg (max 60mg) of prednisolone for 6 weeks followed by an alternate dose of 1.5mg/kg (max 40mg) for the next 6-8 weeks.

Definitions regarding the course of nephrotic syndrome:

• Remission is characterized by a urine protein: creatinine ratio of <0.2 or <1+ protein on urine dipstick testing for 3 consecutive days

• Response in nephrotic syndrome is defined as achieving remission within the initial 4 weeks of corticosteroid therapy

• Relapse is indicated by an increase in urine protein: creatinine ratio >0.2 or a reading of 2+ and higher for 3 consecutive days.

• Frequently relapsing: ≥2 relapses in 6 months or 4 relapses in 12 months.

• Steroid-dependent nephrotic syndrome: Relapse during steroid tapering or within 2 weeks of discontinuation.

• Steroid resistance: Failure to achieve remission despite therapy with daily prednisolone for 4 weeks and alternate days for the next 4 weeks.

Other options :

Option B: Nephritic syndrome classically presents with tea- or cola-colored urine (hematuria), edema, hypertension, and oliguria.

Option C: Goodpasture disease is an autoimmune condition characterized by pulmonary hemorrhage, rapidly progressive glomerulonephritis, and elevated anti-glomerular basement membrane antibody levels.

Option D: IgA nephropathy is an immune complex-mediated disease, with a predominance of IgA within glomerular mesangial deposits. They present with gross hematuria and proteinuria.

Solution to Question 5:

In a child with suspected shunt-associated meningitis, the next best step useful in confirming the diagnosis is by taking a blood culture and a CSF sample from a shunt tap is preferable to a lumbar puncture (LP) (option D).

Shunt aspiration allows for better detection of microbes and avoids the need for a separate procedure, reducing the risk of contamination. VP shunt fluid is obtained from a subcutaneous reservoir using a 25-gauge needle.

Bacterial meningitis may be caused due to meningococcus, streptococcus pneumoniae, or Haemophilus influenzae type b. The clinical presentation of meningitis in children with VP shunts is similar to acute bacterial meningitis in other children. These include fever, rash, vomiting, and refusal to feed. Clinical signs of meningeal irritation (Kernig and Brudzinski signs) are not usually seen in infants.

Blood cultures should be done in all cases of suspected meningitis. Treatment is aggressive with systemic antibiotics. Third-generation cephalosporins along with vancomycin can be used. Wait and watch policy is not followed (option A).

A ventriculoperitoneal (VP) shunt is used to drain excess cerebrospinal fluid (CSF) due to an obstruction in the normal outflow or there is a decreased absorption of the fluid. A silicone rubber device with a proximal portion inserted into the ventricle, a unidirectional valve, and a distal segment is commonly used for shunt placement.

Complications of the shunting procedure include infection due to skin flora entering the shunt (Staphylococcus epidermidis), intracerebral or intraventricular hemorrhage, shunt over drainage, shunt nephritis, shunt disconnection, obstruction, inguinal hernia, and seizures.

Shunt patency studies are performed to determine whether shunt revision surgery is needed in the malfunctioned ventriculoperitoneal shunt. This is done using radionucleotides (option B)

Hydrocephalus represents a group of conditions that result from impaired circulation or absorption of CSF.

Causes of communicating hydrocephalus (increased production or reduced absorption) includes choroid plexus papilloma, meningeal malignancy, and meningitis. Causes of noncommunicating (obstructive) hydrocephalus include aqueductal stenosis, infectious (toxoplasmosis, mumps), Chiari malformation, Dandy-Walker malformation, and mass lesions.

In infants, signs include rapid head enlargement, bulging fontanel, and downward eye deviation. Long-tract signs and dilated scalp veins may also be observed. Older children may experience irritability, lethargy, poor appetite, vomiting, and headaches. Diagnosis can be done using MRI, ultrasound, and measurement of ICP.

Medical management with medications like acetazolamide and furosemide can provide temporary relief, but long-term outcomes are often unsatisfactory. Most cases require surgical intervention, such as ventriculoperitoneal shunting or endoscopic third ventriculostomy.

Solution to Question 6:

Umbilical arteries carry deoxygenated blood back to the placenta during the fetal period. The umbilical vein carries oxygenated blood away from the placenta.

Arteries carry oxygenated blood away from the heart, except for
the pulmonary and umbilical arteries which carry deoxygenated blood.

The blood gets oxygenated in the placenta and this passes to the umbilical vein (pO_2 30 to 35mmHg). From the umbilical vein, 50% of the blood enters the hepatic circulation and 50% bypasses the liver to enter the inferior vena cava (IVC) (pO_2: 26 - 28mmHg) through ductus venosus.

The right atrium receives the oxygenated blood from the IVC and superior vena cava (less oxygenated). The right atrial pressure is more than the left atrial pressure resulting in the major flow of blood from the right atrium to the left atrium via the foramen ovale. The blood from the left atrium flows to the left ventricle. The upper part of the fetal body (coronaries, cerebral arteries, and upper extremities) is exclusively perfused by the left ventricle via the ascending aorta.

The major portion of this blood bypasses the lung and flows from the pulmonary artery into the descending aorta via the ductus arteriosus. Finally, some of this blood from the aorta and the lower parts of the body is pumped back to the placenta via umbilical arteries.

A schematic diagram of fetal circulation is given below:





The concept of fetal circulation and the various embryological remnants of the vascular entities of fetal circulation is a high-yield frequently tested topic in NEET-PG examination. The concepts are summarized in the pearl attached below.

Solution to Question 7:

The clinical scenario of a child with a previous history of fever with rash at 1 year, now presenting with seizures, is highly indicative of subacute sclerosing panencephalitis (SSPE), a rare chronic complication of measles. Anti-measles antibodies are elevated in cerebrospinal fluid (CSF) (IgG >IgM); hence, it is the most helpful investigation to diagnose this condition.

SSPE is a fatal progressive demyelinating disease caused by chronic CNS infection with the genetically altered variant of the measles virus. It occurs several years after primary measles infection and is more common in males.

Clinical manifestations progress through four stages:

• Stage I: involves mild symptoms like irritability and poor school performance.

• Stage II: is characterized by myoclonic seizures.

• Stage III:choreoathetosis, immobility, dystonia, lead-pipe rigidity, and cognitive decline are seen.

• Stage IV: leads to coma and death.

Diagnosis of SSPE is established by documentation of a compatible clinical course along with one of the following supporting findings: elevated measles antibody (IgG >IgM) in CSF analysis or characteristic electroencephalographic (EEG) patterns, suppression-burst episodes (high voltage sharp slow waves) or typical histological findings or isolation of virus from brain tissue.

CT and MRI show evidence of multifocal white matter lesions, cortical atrophy, and ventricular enlargement. Increased T2 signaling in the white matter of the brain is seen as the disease progresses (shown below):



Treatment is mainly supportive. Isoprinosine along with interferons is said to have additional effects. Carbamazepine is found to be useful in controlling myoclonic jerks in the early stages of the illness.

In measles, after an incubation period of 8-12 days, the prodromal phase starts with mild fever and is followed by conjunctivitis, photophobia, coryza, cough, and increasing fever. Koplik spots, appearing 1-4 days prior to the rash, are a characteristic sign of measles. The rash (shown below) appears on the 4th day of fever and typically begins on the forehead (around the hairline), and behind the ears, as a red maculopapular eruption.



The most common complication of measles is acute otitis media. The most common cause of death in measles is pneumonia. Low serum retinol levels in children with measles are associated with higher morbidity and mortality and hence Vitamin A therapy is indicated for all patients with measles.

Other options:

Option B: A common cause of temporal lobe epilepsy is mesial (medial) temporal sclerosis. It is the most common cause of surgically remediable partial epilepsy in adolescents and adults. It is often preceded by febrile seizures. Pathologically, these patients have atrophy and gliosis of the hippocampus. MRI shows hippocampal sclerosis (shown below).



Option D: C1Q4 antibodies in the CSF are seen in neuromyelitis optica. Neuromyelitis optica, also known as Devic's disease, is an aggressive inflammatory demyelinating disorder characterized by recurrent attacks of optic neuritis and myelitis.

Solution to Question 8:

The weight for age of the child falls in between -2SD and -3SD as shown in the growth chart below, which is suggestive of moderate malnutrition and the mother should be taught to feed the baby properly.

The primary treatment in mild and moderate malnutrition involves providing sufficient protein and energy intake, with a recommended minimum of 150 kcal/kg/day. Nutritious home food is advised, and frequent feeding (up to 7 times a day) may be required to achieve the necessary energy intake. High-fat foods are often used to increase energy in therapeutic diets.

Animal-source foods, such as milk and eggs, are recommended by WHO as they provide necessary amino acids and nutrients. Plant-source foods like legumes and cereals can also be good protein sources. A slight increase in protein intake (3g/kg/day) is sufficient for catch-up growth with high energy intake. Milk and vegetable protein mixtures can be used as a protein source and adequate minerals and vitamins are also added.

Weight gain is the best measure of treatment efficacy for mild and moderate malnutrition.

Severe acute malnutrition (SAM) should be referred to a hospital for proper management. Uncomplicated SAM can be managed at home under supervision. Complicated SAM should be treated as an inpatient.

Management of SAM is summarized in the image below.



The World Health Organization(WHO) has established international standards for normal child growth from birth to 5 years of age. The normalization of the anthropometric measures is done in terms of z-scores (standard deviation scores). The anthropometric indicators are marked in a growth chart which is compared with a reference curve.

The anthropometric measures used are weight for age, height for age, weight for height, BMI, and mid-upper arm circumference.

• Low weight for height, or wasting, indicates acute malnutrition.

• High weight-for-height indicates overweight.

• Low height-for-age, or stunting, indicates chronic malnutrition. This is the best indicator of long-term nutritional status in the pediatric population.

• Low weight for the age indicates acute or chronic malnutrition. It does not differentiate between wasting and stunting. Hence, it has limited clinical significance.

Mid-upper arm circumference (MUAC) is used for the screening of wasting in children between 6 months to 5 years. In children aged 6 months to 5 years, MUAC <11.5cm is considered as severe malnutrition.

Anthropometric indicators of different types of malnutrition in Children



©MARROW

Paediatrics INI-CET 2022

Question 1:

In neonatal resuscitation, which of the following is the most effective indicator of successful ventilatory effort?

a) Colour change

b) Rise in heart rate

c) Air entry

d) Chest rise

Question 2:

Which can be used as an alternative to mother's milk in a low birth weight (LBW) baby?

a) Cow's milk

b) Preterm formula

c) Term formula

d) Human donor milk

Question 3:

A 4-year-old male child with a body weight of 15kg and height of 100cm is admitted with renal failure. His blood urea was 100 mg/dL and serum creatinine was 1 mg/dL. What is the closest calculated eGFR in the patient?

a) 33 ml/min/1.73 m2 BSA

b) 40 ml/min/1.73 m2 BSA

c) 55 ml/min/1.73 m2 BSA

d) 80 ml/min/1.73 m2 BSA

Question 4:

Adrenocorticotropic hormone is the drug of choice in which of the following conditions?

a) Juvenile myoclonic epilepsy

b) Rolandic epilepsy

c) West syndrome

d) Lennox Gastaut syndrome

Question 5:

Regarding urinary tract infection (UTI) in children, which of the following is true?

a) 1,3,4

b) 2,4

c) 2,3,4

d) 1,4

Question 6:

Handedness is seen at the age of _____.

a) 1 year

b) 2 years

c) 3 years

d) 4 years

Question 7:

Which of the following cyanotic heart diseases causes increased pulmonary blood flow?

a) 1,4

b) 1,2

c) 3,4

d) 2,4

Question 8:

An 11-year-old boy was brought to the outpatient clinic with intentional tremors, and poor scholastic performance. His sister has similar complaints. On examination, hepatomegaly is seen. The eye finding is shown in the image. What is the probable diagnosis?

a) Wilson's disease

b) Huntington's chorea

c) Glutaric aciduria

d) Hepatitis A

Question 9:

Which alternative method can be used for the procedure shown in the image below?

a) Head circumference

b) Arm span

c) Crown-rump length

d) Knee height

Question 10:

A child presents with acute exacerbation of asthma. Which of the following would you do?

a) 3,4

b) 1,4

c) 1,3,4

d) 2,3,4

Question 11:

A young man is brought to the hospital with high-grade fever and altered consciousness. On examination, the patient had neck rigidity and pain when bending the neck. A lumbar puncture was performed and it showed a WBC count of 1500 cells/µL, majority being neutrophils, elevated pressure with protein of 120 mg/dL, and a glucose level of 70 mg/dL (blood glucose level - 300 mg/dL). How will you manage the index case?

a) Piperacillin+ tazobactum

b) Amphotericin + flucytosine

c) Vancomycin+ ceftriaxone

d) Antitubercular therapy

Question 12:

Which of the following increase the risk of recurrence of febrile seizures?

a) 1,2,4

b) 1,4

c) 1,2,3

d) 1,3,4

Question 13:

Choose the most appropriate answer.

a) Both assertion and reason are independently false statements.

b) The assertion is independently a true statement, but the reason is independently a false statement.

c) Both assertion and reason are independently true statements, but the reason is not the correct explanation for the assertion.

d) Both assertion and reason are independently true statements, and the reason is the correct explanation for the assertion.

Question 14:

A child is brought to the casualty with a history of choking. On examination, there is decreased air entry in the left lung. What would be the correct management?

a) 1, 2, and 4

b) 1 and 2

c) 3 and 4

d) 1 and 4

Question 15:

A 9-month-old boy presents with progressive pallor and abdominal distension. The Mentzer index was found to be less than 13. What are the investigations to be done in this patient?

a) 1, and 2

b) 1, 3, and 4

c) 1 and 4

d) 1, 2, 3, and 4

Question 16:

A child presents with complaints of fever, rash, body ache, and throat ache. He has a history of thorn prick injury a week back. What antibiotics would you give empirically to this child?

a) Ceftriaxone

b) Amoxicillin+ clavulanate

c) Vancomycin

d) Meropenem

Question 17:

A child with thalassemia is undergoing multiple blood transfusions. What is the best method to detect iron overload?

a) Liver iron concentration

b) MRI T2 myocardium

c) Serum ferritin

d) NTBI

Question 18:

How would you manage a child with status epilepticus?

a) 1, 2, 3, 4

b) 2, 1, 4, 3

c) 1, 3, 4, 2

d) 1, 2, 4, 3

Question 19:

A child presents with cold feet and a history of wearing socks even in the summers. On examination, lower limb pulses are diminished as compared to the radial pulse. Prominent radio femoral delay is present. What is the underlying condition?

a) There is hypertrophy of ductus arteriosus.

b) Coarctation distal to the origin of the left subclavian artery

c) Supply is from posterior to anterior thoracic artery.

d) Collaterals develop to supply upper limb.

Question 20:

An unconsious child is brought to the casualty. What is the correct sequence of the management?

a) 3-1-2-4-5

b) 1-2-3-4-5

c) 3-1-2-5-4

d) 1-2-4-3-5

Question 21:

Which of the following statements are true regarding cytomegalovirus (CMV) infection?

a) 3, 4

b) 1, 2

c) 2, 4

d) 1, 3

Question 22:

The hormone that does not play a role in the growth of the fetus is:

a) Insulin

b) Growth hormone

c) Thyroxine

d) Glucocorticoids

Question 23:

Which of the following statements is true regarding the tape shown in the image?

a) 1, 3

b) 2, 4

c) 2, 3

d) 1, 4

Answer Key

Question No.	Correct Option
1	b
2	d
3	b
4	c
5	b
6	c
7	c
8	a
9	b
10	d
11	c
12	c
13	c
14	d
15	c
16	b
17	a
18	d
19	b
20	c
21	d
22	b
23	b

Detailed Explanations

Solution to Question 1:

An increase in the heart rate is the most effective indicator of successful ventilatory effort. The response to neonatal resuscitation at various steps is assessed based on the heart rate.

As soon as a baby is born, the tone, cry, breathing efforts, and respiratory efforts are assessed. If the tone is poor, or if the baby is not breathing, the following measures are done immediately - warming to maintain the normal temperature, proper positioning of the baby's airway, i.e. sniffing position with the head slightly extended, clearing of airway secretions, and tactile stimulation.

After the initial steps of neonatal resuscitation, the heart rate of the baby must be assessed. If the heart rate is < 100 beats per minute (bpm), positive pressure ventilation is provided along with SPO2 monitoring to ensure adequate ventilation.

If the heart rate remains at < 100 bpm, ventilation corrective steps and/or endotracheal intubation have to be considered. The ventilation corrective steps are as follows and can be remembered with the mnemonic MRSOPA.

• Mask has to be reapplied

• Reposition head

• Suction the secretions

• Open mouth slightly and ventilate

• Pressure is increased slightly

• Airway may be changed

If the heart rate is < 60 bpm, chest compressions are initiated over the lower third of the sternum at a rate of 90 compressions/min. The ratio of compressions to ventilation is 3: 1 (90 compressions: 30 breaths).

If the heart rate still remains <60 bpm, despite effective compressions and ventilation, then epinephrine (1:10,000 solution at 0.1–0.3 mL/kg intravenously or 0.5–1 mL/kg intratracheally) should be administered. The dosage can be repeated every 3–5 minutes. Persistent bradycardia may signify inadequate ventilation or severe cardiac compromise.

The neonatal resuscitation algorithm is given below:



Solution to Question 2:

Human donor milk is the best alternative to mother's milk for low birth weight (LBW) babies.

It is recommended that all LBW babies are exclusively fed breast milk regardless of the feeding method. This can be achieved through the use of expressed breast milk (EBM) from the mother or human donor milk.

Donor milk offers advantages over preterm formula, such as reduced risk of necrotizing enterocolitis (NEC) and feeding intolerance. However, due to lower protein and micronutrient levels in donor milk, human milk fortifiers should be added to meet the nutritional needs of infants.

In cases where breastfeeding is not feasible or the mother is ill, formula feeds can be utilized. Term formula feed (option C) is suitable for infants weighing over 1500 grams at birth, while preterm formula feed (option B) is recommended for very low birth weight infants.

Cow milk (option A) is not preferred as it lacks essential nutrients such as vitamin C, iron, and essential fatty acids. Additionally, its high protein and lactose content can lead to digestive issues and allergic reactions in infants.

Once the mother's condition stabilizes or breastfeeding contraindications are resolved, exclusive breastfeeding should be resumed.

The mode of nutrition of LBW depends on their gestational age. Babies born before 28 weeks of gestation receive total parenteral nutrition (TPN). Enteral feeding is not initiated at this stage as the gastrointestinal tract is not fully developed.

Babies born between 28 and 32 weeks of gestation may exhibit suckling bursts but their coordination of suckling, swallowing, and breathing is not fully developed. Hence, EBM given orally may result in aspiration. EBM is administered through a nasogastric tube or orogastric tube.

As gestational age advances to 32-34 weeks, the coordination of feeding improves, and EBM can be given with a paladai or Katori spoon.

For babies born after 34 weeks of gestation, with fully developed coordination, direct breastfeeding is possible.

Solution to Question 3:

eGFR in the 4-year-old child calculated by the modified Schwartz formula is 40 ml/min/1.73 m2 body surface area (BSA).

The modified Schwartz formula is the most commonly used formula in the pediatric age group to estimate creatinine clearance in acute settings. It is based on serum creatinine, patient height, and an empirical constant k. The k value in the modified Schwartz formula is 0.413 for ages 1-17 years. This formula has higher accuracy and precision compared to the original Schwartz formula.

eGFR = 0.413 x height (cm) / serum creatinine (mg/dL)

In the given scenario, the height of the patient is 100 cm and serum creatinine is 1mg/dL. Therefore, the closest eGFR is 40 mL/min/1.73 m2 BSA.

eGFR in adults is calculated using the Modification of Diet in Renal Disease (MDRD) equation or the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation. The calculation is based on age and serum creatinine with specific modifications in females and African Americans.

Glomerular filtration rate (GFR) is a measure of kidney function. It can be estimated accurately by quantitating the clearance of a substance filtered across the capillaries but the substance should not be absorbed nor secreted by tubules.

GFR can be estimated by intravenous injection of radioisotope technetium diethylene triamine pentaacetate (Tc-99m DTPA), but it is expensive and time-consuming. Therefore eGFR is estimated by the clearance of creatinine.

Solution to Question 4:

Adrenocorticotropic hormone (ACTH) is the drug of choice in West syndrome.

West syndrome is a severe generalized epilepsy that begins between the ages of 2 and 12 months. It is characterized by a triad of infantile spasms that occur in clusters, developmental regression, and hypsarrhythmia on electroencephalogram (EEG). Hypsarrhythmia refers to a high voltage, slow, chaotic background with multifocal spikes (shown below).



Cryptogenic cases (unknown etiology) have normal development prior to onset, while symptomatic West syndrome is associated with preceding developmental delay owing to perinatal encephalopathies, metabolic disorders, or malformations.

Hormonal therapy, such as ACTH injections or oral steroids, is the preferred treatment for West syndrome. The patient's response is monitored through awake and asleep EEGs conducted at 1, 2, and 4th weeks of starting hormonal therapy. Side effects may include hypertension, electrolyte imbalance, infections, hyperglycemia and/or glycosuria, and gastric ulcers.

Vigabatrin is a first-line treatment option for infantile spasms in patients with tuberous sclerosis and is also considered a second-line choice if hormonal therapy has been unsuccessful in other cases of infantile spasms.

The ketogenic diet is considered a third-line therapy for infantile spasms. Other alternative treatment options for spasms include valproate, benzodiazepines (such as nitrazepam and clonazepam), topiramate, lamotrigine, zonisamide, pyridoxine, and intravenous gamma globulin (IVIG).

Other options:

Option A: Juvenile myoclonic epilepsy (JME)/ Janz syndrome starts in early adolescence and is characterized by myoclonic jerks in the morning. EEG shows generalized irregular 4- to 6-Hz spike-and-wave activity, particularly after awakening. Valproate is considered the most effective medication. Lamotrigine, levetiracetam, and topiramate are also approved for JME.

Option B: Rolandic epilepsy is benign epilepsy seen in children. They present at night or while awakening. They are partial seizures with no loss of consciousness. They resolve in adulthood. Treatment with carbamazepine is useful.

Option D: Lennox-Gastaut syndrome is seen in 2-10 years and is characterized by a triad of developmental delay, multiple seizures (atypical absences, myoclonic, astatic, and tonic seizures), and specific EEG changes (1 to 2 Hz spike and slow waves, polyspike bursts). Treatment with topiramate, clobazam, felbamate, lamotrigine, and valproate is useful.

Solution to Question 5:

The true statements regarding UTI in children are bowel and bladder dysfunction increases the risk of recurrence, and micturating cystourethrogram (MCU) is done in children with recurrence of UTI.

The chief cause of UTI in children is Escherichia coli, others include Klebsiella, Enterobacter, and Staphylococcus saprophyticus. Risk factors for recurrent UTI are age <6 months, obstructive uropathy, female sex, vesicoureteric reflux (VUR), bladder dysfunction, constipation, and repeated catheterization.

Clinical features in neonates include fever, vomiting, diarrhea, poor weight gain, and lethargy. If the child cries during voiding or has a weak urine stream with a palpable bladder, it implies urinary obstruction. High fever (>39°C), persistent vomiting, dehydration, renal angle tenderness, or raised creatinine is considered a complicated UTI.

A colony count of >105/mL of a clean catch sample is considered significant bacteriuria. Any colonies on suprapubic aspiration and >50,000/mL on urethral catheterization are considered significant. Patients with recurrent UTIs need to be evaluated with ultrasonography, DMSA scan (dimercaptosuccinic acid), and MCU.

Evaluation following the first episode of UTI is given below:

Third-generation cephalosporins or aminoglycosides are preferred empirical antibiotics for urinary tract infections (UTIs). Treatment duration is 7-10 days for simple UTIs and 10-14 days for complicated UTIs. Asymptomatic bacteriuria doesn't require treatment. Infants <3 months and those with complicated UTIs should receive parenteral antibiotics.

Cotrimoxazole, nitrofurantoin, cephalexin, and cefadroxil are given as prophylaxis to prevent recurrent UTIs. Cotrimoxazole and nitrofurantoin can be given in infants >3 months of age.

Age	Evaluation
<1 year	USG, MCU, DMSA renal scan
1 to 5 years	USG, DMSA scan. MCU is done if USG is abnormal
>5 years	USG. If USG is abnormal, MCU and DMSA are done.

Solution to Question 6:

Handedness is seen at the age of 3 years. The child may become frustrated if attempts are made in changing the hand preference.

Solution to Question 7:

The cyanotic congenital heart diseases with increased pulmonary blood flow are transposition of the great arteries (TGA) and total anomalous pulmonary venous communication (TAPVC).

Classification of cyanotic congenital heart diseases based on pulmonary blood flow is as follows:

Atrial and ventricular septal defects are also associated with increased pulmonary blood flow. However, they are acyanotic.

Decreased pulmonary blood flow	Increased pulmonary blood flow
Tetralogy of FallotPulmonary atresia with an intact ventricular septumEbstein anomalyIsolated right ventricular hypoplasia.	Transposition of the great vessels.Total anomalous pulmonary venous return (TAPVR)Truncus arteriosus

Solution to Question 8:

The probable diagnosis in the given case, based on the given clinical scenario and the image which shows icterus in the eyes, is Wilson's disease.

Wilson's disease is an autosomal-recessive condition caused by mutations in the ATP7B gene, which encodes for the ATPase that removes copper, preventing its accumulation in the liver and brain.

Clinical features are jaundice, ascites, hepatomegaly, dysarthria, dystonia, wing-beating tremor, a decline in school performance, and personality change. Deposition of copper in the lens results in sunflower cataracts and Kayser-Fleischer (KF) ring is seen due to the deposition of copper in Descemet's membrane of the cornea.

Diagnosis is made by the presence of a KF ring and decreased levels of serum copper and ceruloplasmin. Increased 24-hour urine copper excretion is seen. Measurement of hepatic copper levels in liver biopsy is the gold standard.

Treatment is the use of intramuscular British anti-lewisite or oral D-penicillamine. These drugs increase the excretion of copper in the urine. Pyridoxine is given concomitantly. Trientine hydrochloride is a suitable alternative to those intolerant to penicillamine.

Zinc acetate helps in copper absorption. It is used in presymptomatic patients, pregnant women, and as maintenance therapy. Liver transplantation is considered in advanced stages.

Other options:

Option B: Huntington's chorea is a trinucleotide (CAG) repeat disorder with an autosomal dominant mode of inheritance. It is characterized by involuntary movements known as chorea.

Option C: Glutaric aciduria is an autosomal recessive disease presenting during the first 6 months of life. Clinical features are hypotonia and jitteriness at birth, progressive hyperkinetic movements (dystonia, choreoathetosis), and may have seizures

Option D: Hepatitis A is a viral infection transmitted by the fecal-oral route, which is the commonest, parenteral route, and also by sexual transmission. The onset of jaundice is preceded by symptoms like nausea, vomiting, anorexia, and mild fever. Hepatitis A resolves completely but a relapse may occur in a few cases.

Solution to Question 9:

The procedure shown in the image is the measurement of the height of a child using a stadiometer. The alternative method that can be used for the procedure is arm span. The proportion of arm span changes with age but is close to height.

For children older than 2 years of age, a stadiometer is used to measure their height. It is measured with the head in the Frankfurt plane, and the back of the head, thorax, buttocks, and heels all on the same axis as that of the stadiometer.

Arm span provides an assessment of proportionality. It is measured by asking the child to stand, the back against the wall with arms outstretched, and the distance between the tip of the middle fingers are measured.

The body proportion in a child is assessed by the ratio of the upper segment (US) and lower segment (LS) and by comparing the arm span with height.

Upper segment: Lower segment ratio

A higher US:LS ratio is associated with bone disorders and conditions involving short-limb dwarfism, such as Turner syndrome, whereas a lower ratio is associated with hypogonadism and Marfan syndrome.

Other options:

Option-A: Head circumference is measured as the maximal circumference along the supraorbital ridge in the front and the occipital prominence at the back of the head. It is measured using a flexible tape. It reflects the brain growth in a child.

Option-C: Crown-rump length is used for the assessment of fetal growth as early as 6–8 weeks of gestation. This length is used to determine the gestational age in the first trimester.

Option-D: Knee height is the distance from the sole of the foot to the anterior surface of the femoral condyle, with the ankle and knee flexed to right angles. It is used when standing height cannot be measured.

Age	US: LS ratio
Birth	1.7
3 years	1.3
7 years	1

Solution to Question 10:

A child coming with acute exacerbation of asthma will be administered oxygen, salbutamol nebulization three times in 60 minutes, and oral corticosteroids.

Exacerbation of asthma is known as an increase in symptoms such as cough, wheezing, and/or breathlessness. If the child has cyanosis, silent chest, poor respiratory effort, altered sensorium, and oxygen saturation of <90%, the exacerbation is considered life-threatening.

Based on the severity, acute asthma is classified as mild, moderate, and severe.

Home treatment for acute exacerbation involves parents administering short-acting, beta-2-agonists by metered-dose inhaler with/without a spacer and facemask. One puff for every 30–60 seconds up to a maximum of 10 puffs with monitoring. If symptoms are relieved after inhalation, the child can continue salbutamol inhalation every 4–6 hours. If there is no improvement or partial improvement, the child should be transferred to a hospital.

In a child presenting to the hospital with severe symptoms, start inhalation of oxygen, short-acting beta-2-agonists such as salbutamol or albuterol that can be used every 15–20 minutes for the first hour and administer oral prednisolone (1–2 mg/kg). Oxygen inhalation is stopped if an oxygen saturation of >95% cannot be maintained.

Clinical parameter	Mild	Moderate	Severe
Color	Normal	Normal	Pale
Sensorium	Normal	Anxious	Agitated
Respiratory rate	Increased	Increased	Increased
Dyspnea	Absent	Moderate	Severe
Use of accessory muscles	Nil or minimal	Chest indrawing	Indrawing, nasal flare

Clinical parameter	Mild	Moderate	Severe
Rhonchi	Expiratory and/or inspiratory	Expiratory and/or inspiratory	Expiratory or absent.
Oxygen saturation	>95%	90-95%	<90%

Solution to Question 11:

The given clinical scenario with high fever, altered consciousness, neck rigidity, and cerebrospinal fluid (CSF) showing neutrophilic pleocytosis with low sugar point toward a diagnosis of bacterial meningitis. The initial treatment involves the administration of 3rd-generation cephalosporin like ceftriaxone along with vancomycin for 10–14 days.

Hyperglycemia due to the pathological stress from the infection can mask the reduced CSF glucose. It can be correctly inferred from CSF/blood glucose ratio. A ratio <0.4 is highly significant for bacterial meningitis.

The most common cause of meningitis in adults over 20 years of age is Streptococcus pneumoniae.

The classic clinical triad of meningitis includes fever, headache, and nuchal rigidity. Other common symptoms include decreased consciousness, nausea, vomiting, photophobia, and seizures. Cutaneous signs such as petechiae, purpura, or an erythematous macular rash may be present.

Kernig's sign and Brudzinski's sign can indicate meningeal irritation. Kernig's sign is positive when there is resistance and pain in extending the knee with the hip flexed at 90 degrees. Brudzinski's sign is positive when there is involuntary flexion of the hips and knees in response to passive neck flexion. However, these signs may not be reliable in certain patient groups, such as infants, elderly individuals, or those with altered mental status.

When bacterial meningitis is suspected, blood cultures should be obtained, and empirical antimicrobial therapy should be initiated without delay. The diagnosis is confirmed by examining cerebrospinal fluid (CSF). CSF analysis shows increased polymorphonuclear leukocytes, decreased glucose concentration, increased protein concentration, and increased opening pressure. CSF bacterial cultures and Gram stain are used to identify the causative organism.

Prompt initiation of empirical antimicrobial therapy is crucial in suspected cases of bacterial meningitis. Commonly used antibiotics include third- or fourth-generation cephalosporins (such as ceftriaxone, cefotaxime, or cefepime) that cover Streptococcus pneumoniae, Neisseria meningitidis, and Haemophilus influenzae and Vancomycin if there is a high suspicion of methicillin-resistant Staphylococcus aureus (MRSA) infection. Ampicillin with metronidazole is added when Listeria monocytogenes is suspected, particularly in neonates, pregnant women, individuals over 50 years, and immunocompromised individuals.

Solution to Question 12:

Age <1 year, temperature of 38–39°C (100.4-102.2°F), and fever duration of <24 h (statement 1,2,3) are the major risk factors for the recurrence of febrile seizures.

Minor risk factors include a family history of epilepsy, male gender, daycare, complex febrile seizure, family history of febrile seizures, and lower serum sodium at the time of presentation.

Febrile seizures occur in children of the age of 6–60 months (peak at 12 to 18 months) with a temperature of 38ºC (100.4°F) or higher, without underlying intracranial infections or prior unprovoked seizures.

Types of febrile seizures:

Lumbar puncture should be performed in the first episode of seizures with fever, for all infants <6 months of age (to exclude meningitis). Electroencephalogram and neuroimaging are advised for children with suspected epilepsy or neurological abnormalities.

Treatment with lorazepam, diazepam, and midazolam for seizures lasting more than 5 minutes. Rectal diazepam is prescribed to use at home as rescue medication if febrile seizures last more than 5 minutes. Buccal or intranasal midazolam may be used alternatively.

For frequently recurring febrile seizures, intermittent oral clonazepam or diazepam can be given during febrile illnesses to reduce the risk of recurrence. Continuous prophylaxis is no longer recommended.

Antipyretics can be given but it does not decrease the risk of recurrence. Chronic antiepileptic therapy may be considered for children at high risk of developing epilepsy. Screening for and treating iron deficiency is recommended as it is associated with an increased risk of febrile seizures.

Simple febrile seizures	Complex (atypical) febrile seizures
Occurs within 24 hours of the onset of fever	Can occur at any time during a fever
Generalized seizures	Focal seizures
Duration is <15 min	Prolonged
No recurrence within 24 hrs of seizures	Recurrent
The chances of going into epilepsy are 1%	The chances of going into epilepsy are 6%

Solution to Question 13:

The correct answer is option C.

The mousy or musty odor of urine in phenylketonuria (PKU) is due to the accumulation of phenylalanine metabolites including phenylpyruvate, phenylacetate, and phenyl lactate.

PKU can lead to low levels of tyrosine, which is synthesized from phenylalanine. Tyrosine gives rise to catecholamines like dopamine, norepinephrine, and epinephrine. Thus, low levels of tyrosine may lead to a deficiency of dopamine, norepinephrine, and epinephrine. However, this is not the reason for the mousy odor of urine.

PKU is caused by a deficiency of the enzyme phenylalanine hydroxylase (classic PKU) or defects in the enzymes involved in producing tetrahydrobiopterin. These deficiencies impair the conversion of phenylalanine to tyrosine, leading to elevated levels of phenylalanine and low levels of tyrosine in the body. The excess phenylalanine in the blood is metabolized to phenyl ketones (phenylpyruvate and phenylacetate) which are excreted in the urine.

Affected infants initially appear normal at birth. If left untreated, they gradually develop profound intellectual disability. Older untreated children may exhibit hyperactivity, autistic behaviors, and neurologic signs such as seizures, spasticity, and tremors. Abnormalities like microcephaly, prominent maxillae, and growth retardation are seen. They are also prone to low bone mineral density and osteopenia.

Untreated PKU patients may exhibit hypopigmentation (fair hair, light skin color, and blue eyes) due to low levels of tyrosine, which is necessary for the formation of melanin pigment.

If a woman with PKU is not following a low-phenylalanine diet during pregnancy, it can lead to maternal PKU syndrome in the offspring. This condition is associated with microcephaly and congenital heart disease.

The preferred screening method is tandem mass spectrometry. If screening is positive, the diagnosis is confirmed by quantitatively measuring phenylalanine levels.

The primary treatment for PKU is a low-phenylalanine diet, which should be initiated as soon as the diagnosis is established. Lifelong restriction of dietary phenylalanine is recommended. Tyrosine becomes an essential amino acid in PKU and it needs to be obtained from the diet.

The series of reactions involving the production of catecholamines is shown in the image below:



Solution to Question 14:

A history of choking and reduced breath sounds over the left lung in a child are suggestive of a foreign body lodged in the left bronchus. The next step in the management of this child is to obtain a chest x-ray, followed by bronchoscopic removal of the foreign body.

Food items (nuts, seeds hard candy, gum, bones raw fruits, and vegetables) account for the majority of choking cases in children. The most common site of foreign body lodging is the bronchus. Foreign body aspiration into the airway has three stages: immediate symptoms of coughing and choking, an asymptomatic interval, and later complications such as fever, atelectasis, and pneumonia.

When a child presents with a history of choking, detailed history taking and physical examination of the nose, oral cavity, pharynx, neck, and lungs are necessary.

The next step for suspected foreign body aspiration is to obtain a chest X-ray, including the abdomen, despite most foreign bodies being radiolucent. Secondary findings on the X-ray can indicate foreign body involvement, such as air trapping, atelectasis, obstructive emphysema, asymmetric hyperinflation, mediastinal shift, or consolidation. While a CT scan has high accuracy, it is not typically performed due to radiation risks.

The treatment of choice for a foreign body in the airway is its endoscopic removal with a rigid instrument, usually a bronchoscope, by a specialist otolaryngologist or pulmonologist. Bronchoscopy must be performed even if there is a high suspicion of an airway foreign body in spite of negative or inconclusive imaging.

Solution to Question 15:

The investigations to be done in this patient are HPLC (high-performance liquid chromatography) and peripheral smear

The clinical history of a child with pallor, Mentzer's index <13, and abdominal distention (suggestive of hepatosplenomegaly) is suggestive of thalassemia.

Peripheral smear shows hypochromic microcytic anemia, target cells, basophilic stippling, fragmented red cells, and nucleated RBCs. HPLC showing decreased levels of HbA and an increase in HbA2 and HbF levels is sufficient to make a diagnosis.

The Mentzer index is the ratio of MCV (mean corpuscular volume) to RBC count. It is used to differentiate between iron-deficiency anemia and β-thalassemia trait. MCV and MCH (mean corpuscular hemoglobin) are low in both IDA and thalassemia. RBC count may be normal in thalassemia, whereas it is reduced in IDA. The Mentzer index is <13 in thalassemia minor syndrome.



Bone marrow examination (option 3) and coagulation studies (option 2) have no role in the diagnosis of thalassemia.

Thalassemia is a hemoglobinopathy that arises due to deficient synthesis of the globin chains. Depending on the chain affected, it is classified into α-thalassemia and β-thalassemia.

There is a deficit in the synthesis of HbA with an elevation of HbA2 and HbF levels in β-thalassemia. Children with severe thalassemia present with facial features (shown below) such as maxilla hyperplasia, a flat nasal bridge, and frontal bossing. Other manifestations include pathologic bone fractures, significant hepatosplenomegaly, and cachexia.

Hb Electrophoresis and HPLC are used as screening tests for thalassemia. DNA diagnosis to detect β-thalassemia mutations is used for definitive diagnosis. A radiographic image of the skull shows the classic 'crew cut' appearance.

Blood transfusions are administered every 3-4 weeks to maintain a pretransfusion hemoglobin level of 9.5-10.5 g/dL. Oral iron chelation to prevent iron toxicity is also recommended.

Hematopoietic stem cell transplantation is the only definitive therapy.

Antenatal counseling and heterozygote screening can be used in the prevention of the condition.

α-thalassemia is caused by deficient synthesis of α-chains. Deletion or malfunction of three α-globin chains can result in HbH disease. Hb Bart's hydrops fetalis is due to the absence of any normally functioning α-globin genes.

Solution to Question 16:

A child with a history of fever with rash and throat ache (sore throat) following a thorn prick injury could be having scarlet fever caused by group A Streptococcus (GAS). This child could be treated empirically with amoxicillin-clavulanate.

GAS commonly presents in children with pharyngitis and rash. It commonly spreads through droplets but can also be inoculated by a wound (thorn prick in this case) or burn infected with GAS.

The rash typically begins as diffuse, erythematous, papular eruptions. Sandpaper rash and the strawberry tongue are characteristic lesions that develop in the course of the disease. Pyrogenic exotoxin (erythrogenic exotoxin) produced by Group A Streptococcus is responsible for scarlet fever.

Streptococcal rapid antigen detection test or throat culture should be done before starting treatment with antibiotics. First-line treatment is by penicillin or amoxicillin for 10 days. Clavulanate can be added to combat any beta-lactamases produced by the bacteria.

Solution to Question 17:

The best method to assess iron overload is quantitative liver iron using T2 MRI. It eliminates the need for invasive procedures like liver biopsy.

Excessive iron stores resulting from transfusions are a major concern in the management of β-thalassemia major. Accurate assessment of iron overload is crucial for optimal treatment.

Quantitative measurement of cardiac iron by T2 MRI (option B) is initiated at the age of 10. However, there can be a discrepancy between liver iron and cardiac iron levels due to variations in tissue loading and unloading rates.

While serial serum ferritin levels (option C) can be used as a screening tool, they may not accurately reflect quantitative iron stores. Relying solely on serum ferritin levels can lead to undertreatment or overtreatment of iron overload.

Normally, iron in the plasma is bound to transferrin. But in iron-overload conditions, iron exceeds the iron binding capacity leading to the appearance of non-transferrin-bound iron (option D). It can act as an indicator of iron overload, but it is influenced by plasma factors like citrate, uric acid, and albumin. Hence, it cannot be used as a tool to measure quantitative iron overload that can help guide in planning chelation therapies.

Other tests that help calculate iron overload include echocardiography, bone mass densitometry, and testing for endocrinopathy. These tests are done at the time of diagnosis and done yearly to monitor iron overload. Serum ferritin tests can be done every 3 months.

Hypothyroidism, hypogonadotropic hypogonadism, growth hormone deficiency, hypoparathyroidism, and diabetes mellitus are seen due to iron deposition in endocrine organs. Iron deposition in the heart can lead to heart failure, arrhythmias, and cardiac-related mortality.

Iron chelation therapy is recommended when serum ferritin levels exceed 1,000 ng/mL and/or liver iron concentration is above 5,000 μg/g dry weight. It is not approved for children under 2 years of age. Three iron chelators are available: deferoxamine, deferasirox, and deferiprone. Deferoxamine is administered subcutaneously or intravenously due to its poor oral bioavailability. Deferasirox is a once-daily oral iron chelator. Deferiprone is an oral iron chelator approved as a second-line agent.

Solution to Question 18:

The order of management of a child with status epilepticus is airway and breathing→ iv lorazepam→IV Fosphenytoin→Midazolam, Propofol, Thiopentone (1, 2, 4, 3)

For generalized tonic-clonic seizures (GTCS), status epilepticus (SE) is defined as continuous convulsive activity or recurrent generalized convulsive seizure activity without regaining consciousness.

t1 is the time at which treatment should be initiated. t2 is the time at which continuous seizure activity leads to long-term sequelae such as neuronal injury.

The image given below depicts the management of status epilepticus in a flowchart:

Type of seizure	t1	t2
GTCS	5 min	30 min
Focal seizures with impaired awareness	10 min	30 min
Absence seizures	10-15 min	unknown



Refractory status epilepticus (SE) refers to SE that does not respond to therapy involving at least two medications (such as benzodiazepine and another medication). Super refractory status epilepticus is SE that does not resolve or recur within 24 hours or more despite treatment that involves continuous infusions of medications like midazolam or pentobarbital.

Solution to Question 19:

The given clinical scenario of a child with cold feet with diminished lower limb pulses and radio femoral delay is suggestive of coarctation of the aorta. It is caused by coarctation distal to the origin of the left subclavian artery.

Coarctation of the aorta is a congenital heart disease, which involves constriction of the aorta at any point from the transverse arch to the iliac bifurcation. In 98% of cases, there is a narrowing of the descending aorta, usually distal to the origin of the left subclavian artery at the origin of the ductus arteriosus.

Coarctation of the aorta can be classified into two types based on the location of constriction: preductal (infantile type) and postductal (adult type). Preductal coarctation occurs proximal to the ductus arteriosus, while postductal coarctation occurs distal to it.

Clinical features include differences in blood pressure and pulse in the upper and lower extremities, radio femoral delay, and prominent pulsations in the upper limb. Cardiac enlargement and pulmonary congestion are noted in infants with severe coarctation. During childhood, the findings are present after the 1st decade. The heart is mildly or moderately enlarged because of left ventricular prominence. The enlarged left subclavian artery typically produces a prominent shadow in the left superior mediastinum. A continuous murmur over the infra-scapular region or the scapula can be auscultated.

The condition is associated with the bicuspid aortic valve, systemic hypertension, an aortic aneurysm, left ventricular hypertrophy, left heart failure, and premature atherosclerosis.

The figure of 3 sign refers to the indentation of the aorta at the coarctation site. There is pre and post-stenotic dilatation along with the left para-mediastinal shadow. Chest radiograph shows notching of ribs and the figure of 3 sign:



Unless surgically corrected in infancy, coarctation of the aorta results in the development of collateral circulation, mainly from branches of the subclavian, superior intercostal, and internal mammary arteries, to create channels for arterial blood to bypass the area of coarctation.

Hypertrophy of the ductus arteriosus (Option A) is not typically associated with coarctation of the aorta as it closes shortly after birth and is not directly involved in coarctation. The blood supply to the upper limb is higher as the coarctation is distal to the origin of the subclavian artery, therefore collaterals do not develop to supply the upper limb (Option D).

Solution to Question 20:

The correct sequence to provide basic life support is to assess responsiveness, check for breathing and pulse, start bag and mask ventilation commence and then cardiopulmonary resuscitation (CPR).

In an unresponsive patient, ensure scene safety first, then check for responsiveness, and activate the emergency response. In the presence of two or more rescuers, one remains with the victim while the other activates the emergency response system. Evaluate breathing and pulse within 10 seconds.

If the person does not have a pulse and is not breathing and there is only one rescuer, start CPR immediately with cycles of 30 compressions and 2 breaths if it is not a witnessed collapse. If it is a witnessed collapse with one rescuer, activate the emergency response system (ERS) if not done already, and retrieve the defibrillator.

After about 2 minutes, if still alone, activate the emergency response system and retrieve the automated external defibrillator (AED). If there are 2 or more rescuers, then one should start CPR while the other retrieves AED. When the second rescuer arrives change the cycle to 2 breaths for

every 15 compressions. Breaths can be provided with the help of a bag and mask.

If the patient has a pulse but no normal breathing or only gasping, provide 1 breath every 3 to 5 seconds or about 12 to 20 breaths/min. Add compressions if pulse remains ≤60/min with signs of perfusion. Activate the emergency response system after 2 minutes. Continue rescue breathing, and check pulse every 2 minutes. If no pulse, start CPR.

When AED arrives, analyze the rhythm. If it is a shockable rhythm, give 1 shock. Resume CPR immediately for about 2 minutes. If it is a non-shockable rhythm, resume CPR immediately for about 2 minutes.

The image below shows chest compressions in pediatric patients.



Solution to Question 21:

The true statements regarding CMV infection are statements 1 and 3.

Most CMV transmission occurs in the first trimester. The risk of transmission is high in the third trimester.

Only 10% (statement 2) of the infected babies are symptomatic. Asymptomatic neonates have lesser chances of having late sequelae (statement 1).

Congenital CMV infection can result from either primary or reactivation infection of the mother. The rate of transmission of infection to the fetus in a mother who has a primary infection is about 30% (statement 3) while in a mother with a non-primary infection, it is 1-2%. However, the total number of infants who have acquired an infection from a mother with a non-primary infection is 3-4 times more than infants who have acquired an infection from mothers with a primary infection.

The clinical manifestations include
hepatosplenomegaly, petechial rashes, jaundice, chorioretinitis, and microcephaly. Neuroimaging shows intracranial periventricular calcifications. One of the long-term sequelae is sensorineural hearing loss.

The investigation of choice is PCR or viral culture. It should be done within the first three weeks of life (statement 4). After this period, it is difficult to distinguish between a congenital and an acquired infection. Since not all affected infants are viremic at birth, urine and/or saliva are more reliable samples than blood.

There is no definitive treatment for congenital CMV infection. Treatment with ganciclovir is indicated only in patients with progressive neurologic disease and deafness.

Below is the image showing the blueberry muffin rash in CMV affected baby:

Solution to Question 22:

Growth hormone, despite being present at higher levels in the fetus, does not play a significant role in fetal growth.

The human fetus produces thyroxine from the 12th week of gestation. Thyroxine and insulin have an important role in regulating tissue growth and differentiation of the fetus, particularly during late gestation. Glucocorticoids play an important role towards the end of gestation in the prepartum maturation of organs such as the liver, lungs, and gastrointestinal tract.

Insulin-like growth factor (IGF)-1 and IGF-II also play an important role in fetal growth.

Solution to Question 23:

The tape shown in the image is Shakir's tape. It is used to assess severe acute malnutrition (SAM). It is used by frontline field workers for nutritional screening of malnourished children and those at risk, aged 6-59 months.

Shakir's tape is used for the measurement of mid-upper arm circumference (MUAC). To locate the midpoint of the left arm, the arm is flexed at the right angle, and the mid-point is marked halfway between the acromion and olecranon process. Color coding is as follows:

• Red is less than 11.5 cm, which indicates severe acute malnutrition

• Yellow is from 11.5 to 12.5 cm, which indicates moderate acute malnutrition

• Green is from 12.5 to 26.5 cm, indicating no acute malnutrition

The presence of any one of the following confers severe acute malnutrition in children 6 to 59 months of age:

• Weight-for-height (length) below -3 standard deviation (<-3SD) on the WHO growth standard

• Presence of bipedal edema

• Mid-upper arm circumference (MUAC) below 11.5 cm

Paediatrics NEET 2022

Question 1:

A 1-day-old neonate has not passed urine since birth. What is the next step in management?

a) Continue breast feeding and observe

b) Admit to NICU

c) Start artificial feeding

d) Start intravenous fluids

Question 2:

A 7-year-old boy presented with abdominal pain, vomiting, oliguria, and periorbital puffiness following chemotherapy. Investigations reveal hyperuricemia, raised creatinine levels, and hyperkalemia. What is the next best step in the management of this condition?

a) Hydration

b) Probenecid

c) Allopurinol

d) Rasburicase

Question 3:

A baby presented with abdominal pain. On examination, a mass is palpated in the right lumbar region. A barium enema is done, and the image is given below. What is the likely diagnosis?



a) Intussusception

b) Volvulus

c) Duodenal atresia

d) Intestinal obstruction

Question 4:

An 8-day-old newborn was found to have thyroid-stimulating hormone level of more than 100 mIU/L. Which of the following will be the next best investigation?

a) Urine iodine excretion

b) Serum thyroid receptor antibody

c) Radiotracer uptake with technetium

d) Perchlorate secretion test

Question 5:

A 10-month-old infant was brought with complaints of jerking movement of limbs towards the body. On examination, there is a regression of developmental milestones. Electroencephalogram shows hypsarrythmia. Which of the following is the drug of choice in this condition?

a) Phenytoin

b) Adrenocorticotropic hormone (ACTH)

c) Levetiracetam

d) Phenobarbitone

Question 6:

A 2-month-old infant born to an HIV-positive mother presents with recurrent diarrhea. What is the next best step?

a) Test stool for giardia and give antibiotics

b) Dried spot sample for HIV DNA PCR

c) Antibody tests for HIV

d) Aerobic culture

Question 7:

An 8-year-old child has difficulty walking and getting up from a squatting position. A muscle biopsy was done and is as shown in the image. Which of the following is true about this condition?



a) Death occurs in 3rd decade

b) Previous history of viral prodrome

c) It is a mitochondrial storage disorder

d) Early treatment has excellent prognosis

Question 8:

A child presented with a history of loose stools with an increase in frequency for 4 days. On examination, he is drowsy, unable to feed, and skin on pinching goes back very slowly. According to the integrated management of neonatal and childhood illness (IMNCI), this child will be classified as having:

a) Mild dehydration

b) Some dehydration

c) Severe dehydration

d) Moderate dehydration

Question 9:

Which of the following children are considered at-risk babies?

a) 2,3

b) 1,2,3,4

c) 4,5

d) 1,4

Question 10:

Identify the condition.

a) Bladder exstrophy

b) Omphalocele

c) Persistent vitellointestinal duct

d) Gastroschisis

Question 11:

A previously healthy child presented with acute-onset dyspnea. A chest X-ray shows unilateral hyperinflation of the lungs. What is true for this patient?

a) Focal area of decreased air entry will be suggestive of foreign body

b) Flexible bronchoscopy used for removal

c) In complete obstruction, ball and valve mechanism causes hyperinflation

d) The child has developed acute laryngotracheobronchitis

Question 12:

A male child presented with arthralgia and abdominal pain. On examination, there was palpable purpura over the lower limbs. There is past history of upper respiratory tract infection prior to the onset of presenting symptoms. Which of the following is the treatment for this condition?

a) Azathioprine

b) Methotrexate

c) Cyclosporine

d) Glucocorticoids

Question 13:

A 10-year-old presents with edema and anasarca. A diagnosis of minimal change disease is made. Which of the following is true about this condition?

a) Light microscopy shows effacement of podocytes

b) Good response to steroids

c) Most common in adults

d) Non-selective proteinuria

Question 14:

A 10-year-old child presents with diarrhea and weight loss. On examination, the height and weight are lesser than expected. Laboratory investigations were positive for class II HLA-DQ2. Which of the following will you advise the child?

a) Fat-free diet

b) Lactose-free diet

c) Low carbohydrate diet

d) Gluten-free diet

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	c
5	b
6	b
7	a
8	c
9	a
10	a
11	a
12	d
13	b
14	d

Detailed Explanations

Solution to Question 1:

In this clinical scenario, a 1 day-old neonate has not passed urine since birth. Most healthy neonates void within 24 hours of life. Delay beyond the first 24 hours of life is generally benign but may result in parental anxiety. Continuing breastfeeding and observing is the next best step in management.

The first urination should occur by 12-24 hours of age and the first passage of meconium should occur by 48 hours of age.

Delay in the passage of urine is seen in conditions causing blockage to urine outflow or birth defects that affect the spinal cord. Some commonly seen conditions are vesicoureteric reflux, ureteropelvic junction obstruction, posterior urethral valve, ureterocele, and spina bifida.

Delay in the passage of the first stool in term neonates may be associated with conditions causing a lower intestinal obstruction, which include meconium plug syndrome, Hirschsprung disease, and imperforate anus. More generalized problems, such as sepsis or hypothyroidism, should be considered.

Solution to Question 2:

The clinical scenario and the laboratory investigations are suggestive of tumor lysis syndrome. Aggressive hydration is the next best step in the management of this condition.

Tumor lysis syndrome (TLS) is a condition that can occur spontaneously after the initiation of chemotherapy.

During chemotherapy, the treatment kills the malignant cells, which leads to increased serum uric acid levels from the turnover of nucleic acids causing hyperuricemia. Tumor lysis results in the release of intracellular potassium, and phosphate pools, which causes hyperkalemia, and hyperphosphatemia. Increased phosphate levels lead to reciprocal depression of serum calcium/hypocalcemia. Deposition of uric acid crystals, calcium phosphate, and hyperphosphatemia leads to renal failure.

Signs and symptoms associated with TLS can develop within 12-48 hours after the initiation of chemotherapy and include:

• Abdominal pain and distension

• Dysuria, oliguria, flank pain and hematuria

• Anorexia, vomiting, cramps, seizures, spasms, altered mental status, and tetany due to hypocalcemia

• Weakness and paralysis due to hyperkalemia

• Hemodialysis in patients refractory to medical treatment

The Cairo-Bishop criteria are used to assess the severity of TLS.

Management of TLS includes:

• Before starting therapy, the electrolytes should be measured and aggressive hydration should be done.

• Allopurinol is a xanthine oxidase inhibitor that decreases the synthesis of uric acid. It should be started in TLS to prevent further accumulation of uric acid.

• Febuxostat is a selective non-purine xanthine oxidase inhibitor that can be used to treat hyperuricemia. It does not require dose calculations and can be used in patients with mildly decreased renal function.

• Rasburicase is an enzyme that degrades uric acid. It is preferred in TLS over allopurinol to decrease the accumulation of uric acid.

Option B: Probenecid inhibits the organic anion transporter in the nephron. It thereby inhibits the reabsorption of uric acid. It is contraindicated in TLS as it increases the risk of uric acid stones due to increased uric acid levels.

Solution to Question 3:

The clinical scenario of pain abdomen with a palpable mass in the right lumbar region along with the barium enema showing claw sign is suggestive of intussusception.

Intussusception occurs when a portion of the bowel is telescoped into an adjacent segment. Intussuscipiens is the one that receives and intussusceptum are the tubes that advance. The apex of the intussusceptum is more prone to ischemia which leads to gangrene.

The most common location of intussusception is ileocolic.

It can be seen in infants during the weaning period following the introduction of new food, vaccination, or upper respiratory tract infection, and the currently approved rotavirus vaccines are associated with very minimal risk of intussusception. An area of enlarged submucosal Payer's patch acts as a lead point for intussusception.

Some lead points in children are Meckel's diverticulum, intestinal polyp, inflamed appendix, submucosal hemorrhage seen in Henoch-Schonlein purpura, and intestinal duplication cysts.

Patients usually present with cramping abdominal pain, palpable sausage-shaped abdominal mass, vomiting, and passage of red currant jelly stools.

Diagnosis is confirmed by abdominal ultrasound which shows target sign/bull's eye sign (shown below) and pseudokidney sign (seen longitudinally).



Barium enema shows a typical claw sign (shown below). This appearance is due to the convex filling defect caused by the apex of the intussusceptum.



The image below shows the coiled spring appearance seen in intussusception. This is due to the barium in the lumen of the intussusceptum and in the intraluminal space.



Treatment of intussusception includes hydrostatic reduction by enema using contrast material or air. Surgical reduction is only done in the presence of refractory shock, suspected bowel necrosis or perforation, peritonitis, and multiple recurrences after hydrostatic reduction. An appendicectomy is an essential component of surgical management.

Option B: Sigmoid volvulus presents with abdominal pain and distension with constipation. A plain x-ray of the abdomen shows an omega sign, coffee-bean sign (shown below), or bent inner tube sign.



Option C: Duodenal atresia is commonly seen in newborns. Babies usually present with bilious/non-bilious vomiting immediately after birth. A plain X-ray abdomen reveals two foci of gas, one in the stomach and the other in the duodenum called the double bubble sign (shown below).



Option D: Intestinal obstruction presents with abdominal pain and distension with constipation. A plain x-ray of the abdomen shows multiple air-fluid levels (shown below). Normally, 3 air-fluid levels can be seen in the plain x-ray at the fundus of the stomach, at the duodenum, and at the cecum. A barium meal is usually contraindicated in acute intestinal obstruction.

Solution to Question 4:

The given laboratory finding of increased thyroid-stimulating hormone level in an 8-day-old baby is suggestive of congenital hypothyroidism. Radiotracer uptake with technetium will be the next investigation in this condition to confirm the presence of the thyroid gland.

Congenital hypothyroidism is the most common preventable cause of mental retardation. Causes of congenital hypothyroidism can be classified based on the presence of goiter as:

Clinical features of congenital hypothyroidism:

• Prolongation of physiologic hyperbilirubinemia- early sign (it is due to delayed maturation of glucuronide conjugation)

• Constipation- usually does not respond to treatment

• Wide open anterior and posterior fontanelles (>0.5 cm) and cranial sutures

• Developmental delay, delayed bone growth, delayed dentition

• Cardiovascular- slow pulse rate, cardiomegaly, asymptomatic pericardial effusion, edema

• Respiratory- macroglossia, feeding difficulties, choking spells, respiratory difficulties

• Hematological- macrocytic anemia refractory to treatment with hematinics

• Nervous system- hypothermia, hypotonia, slow relaxation of deep tendon reflexes, sluggishness, and somnolence.

The classical presentation is described as the 6 P's of congenital hypothyroidism: pot-bellied, pale, puffy-faced child with protruding umbilicus, protuberant tongue, and poor brain development.

The image given below shows a pot belly with an umbilical hernia seen in congenital hypothyroidism.

Goitrous(Palpable goiter)	Non-goitrous(No palpable goiter)
Thyroid dyshormonogenesis (30%)Iodine deficiencyPlacental passage of antithyroid drugs	Thyroid dysgenesis(65%)Thyroid receptor blocking antibodies (5%)Central hypothyroidism

Laboratory investigations in congenital hypothyroidism would demonstrate low T4 levels and increased TSH levels. Screening is usually done on days 1 and 5 days of life, dried blood samples obtained from a heel prick are used for this. Initial investigations in a child with high TSH levels include evaluation of radiotracer uptake study with radioactive iodine or technetium and ultrasound of the thyroid to confirm the presence of a thyroid gland.

An X-ray shows a delay in bone development. The distal femoral and proximal tibial epiphyses normally present at birth are often absent.

In the image below, the left side shows a normal neonate with both knee epiphyses present and the right side shows a neonate with congenital hypothyroidism in whom both epiphyses are small.



Treatment with levothyroxine should be started immediately after diagnosis. Thyroxine (T4) of dose 10-15 μg/kg/day should be started and adjusted to achieve the normal free T4 range. In central hypothyroidism, cortisol replacement should be started before thyroid replacement.

Neonatal hypothyroidism is a sensitive indicator of environmental iodine deficiency and can also be an effective indicator for monitoring the impact of a program.

Other options:

Option A: Urinary iodine excretion is used for surveillance and epidemiological studies for classifying countries according to the level of iodine deficiency by the WHO, and is an effective indicator of the program against iodine deficiency.

Option B: Serum thyroid receptor antibody is used to diagnose transient congenital hypothyroidism caused due to transplacental passage of TSH receptor blocking antibodies. It is usually seen in babies born to mothers suffering from autoimmune hypothyroidism taking thyroxine.

Option D: A perchlorate secretion test can be used to diagnose thyroid peroxidase deficiency.

Solution to Question 5:

The clinical scenario along with the electroencephalogram showing hypsarrythmia is suggestive of the West syndrome. Adreno-corticotropic hormone (ACTH) is the drug of choice for this condition.

West syndrome is severe generalized epilepsy that begins between the ages of 2 and 12 months. It is characterized by a triad of:

- Infantile epileptic spasms
- Developmental regression
- Hypsarrhythmia on electroencephalogram (EEG)

Hypsarrhythmia refers to a high voltage, slow, chaotic background with multifocal spikes as shown below.



West syndrome has been known by many names like infantile spasms, salaam attacks, salaam convulsions infantile spastic epilepsy, and jackknife convulsion syndrome. During an attack babies typically attain complete heaving of the head forward towards their knees, and then immediately relax into the upright position.

The image given below illustrates infantile spasms in a child.

Diagnosis is confirmed by EEG which typically shows hypsarrythmia.

The treatment of choice in West syndrome is either adreno-corticotropic hormone (ACTH) or corticosteroids. An antiseizure medication, mainly vigabatrin can be used when associated with tuberous sclerosis.

Note: The table given below shows the EEG features seen in various seizure disorders.

Options A and C: Phenytoin and levetiracetam are used in focal seizures and status epilepticus.

Option D: Phenobarbitone is used as an antiepileptic in neonatal seizures and as an enzyme inducer in Crigler-Najjar syndrome type II.

Disease	Typical EEG feature
Lennox-Gastaut syndrome	1.5-2.5 Hz spike and polyspike bursts in sleep
Absence seizures	3 Hz spike and slow-wave (spike and dome pattern)
Juvenile myoclonic epilepsy	4-5 Hz spike
West syndrome	Hypsarrhythmia (high-voltage waves with chaotic background)

Solution to Question 6:

In the above scenario, the diagnosis of HIV infection should be established/ruled out before evaluating for secondary infections. Hence, a dried spot sample for HIV DNA PCR is the next best step.

AIDS enteropathy, a syndrome of malabsorption with partial villous atrophy not associated with a specific pathogen, has been postulated to be a

result of direct HIV infection of the gut.

The diagnosis of HIV infection in babies born to HIV-infected mothers is done by HIV proviral DNA or RNA PCR, HIV culture, or HIV p24 antigen. The national program (Early Infant Diagnosis) now uses HIV total nucleic acid (RNA + proviral DNA) PCR test on dried blood spot samples. All positive tests need to be confirmed with an HIV PCR test on another sample.

An infant born to an HIV-positive mother should be breastfed exclusively or by replacement feeds exclusively for 6 months of age. Infants should receive ARV prophylaxis with syrup nevirapine or zidovudine.

Involvement of the gastrointestinal tract is common in HIV-infected babies. Common pathogens identified are bacteria (Salmonella, Campylobacter, Shigella, Mycobacterium avium complex), protozoa (Giardia, Cryptosporidium, Isospora, microsporidia), viruses (CMV, HSV, rotavirus), and fungi (Candida). MAC and protozoal infections are most severe and protracted in children.

Other options:

Option A: Test stool for giardiasis and give antibiotics is not preferred as the immediate next step. First, the diagnosis of HIV infection should be established before treating secondary infections.

Option C: Antibody tests for HIV are not useful in establishing the diagnosis of HIV infection. This is because all infants born to HIV-infected mothers test antibody positive at birth because of the passive transfer of maternal HIV antibodies across the placenta. It can be used as a diagnostic tool after 18 months of age.

Option D: Aerobic culture is not used for the diagnosis of HIV infection.

Solution to Question 7:

The clinical scenario and the muscle biopsy showing fatty replacement of muscle fibers are suggestive of Duchenne muscular dystrophy (DMD). Death in DMD occurs in the late teens and 20s (3rd decade).

DMD is the most common hereditary neuromuscular disease. It is inherited in an X-linked recessive pattern. It is caused by a mutation in the dystrophin gene on chromosome Xp21.

Normally, the dystrophin gene codes for the large protein dystrophin. Dystrophin protein forms a rod that connects the actin filaments to the transmembrane protein dystroglycan in the muscle cell membrane. This dystrophin-glycoprotein complex thereby provides strength to the muscle and connects them to the extracellular environment.

The image given below shows the dystrophin-glycoprotein complex.



Clinical features of DMD:

• Progressive weakness, developmental delay, and intellectual impairment

• In toddlers- delayed walking, falling, toe walking, and trouble running or walking upstairs

• Calf muscular pseudohypertrophy

The image given below shows pseudohypertrophy of the calf muscles.

• Gower's sign: Patient has to use their hands and arms to walk up their own body from squatting position due to the lack of hip and thigh muscle strength due to the weakness of proximal muscles, especially of the lower limbs.

• Trendelenburg gait/hip waddle

• The patient is unable to walk by the age of 12 years and death occurs by late 20 years of age, mostly due to respiratory failure during sleep.

The complications of DMD include contractures, scoliosis, dilated cardiomyopathy, and malignant hyperthermia after anesthesia.

Diagnosis of DMD:

• Elevated serum creatine kinase levels

• Polymerase chain reaction (PCR) analysis for the dystrophin gene

• If PCR is normal, but clinical suspicion is high, muscle biopsy (fatty replacement of muscle fibers) with dystrophin immunocytochemistry (absence of the normal sarcolemmal staining) is done.

Treatment of DMD includes glucocorticoids (prednisone or deflazacort), and multidisciplinary management. Eteplirsen and casimersen are the two new FDA-approved drugs for the treatment of DMD.

Note- The FDA has approved Vamorolone, a novel corticosteroid for the treatment of DMD. The drug offers enhanced anti-inflammatory properties and mineralocorticoid receptor antagonism as well. It is potentially a safer and more effective alternative to glucocorticoids.

Solution to Question 8:

A clinical scenario of a dehydrated child who is drowsy and not able to feed with skin going back very slowly on pinching is suggestive of severe dehydration (as per IMNCI).

Diarrhea is defined as a change in consistency and frequency of stools that occurs more than 3 times a day. Diarrhea, if associated with blood in stools, is called dysentery. If the episodes subside within 7 days, it is called acute diarrhea. If it persists for more than 2 weeks, it is called persistent diarrhea.

Intestinal infections are the most common cause of acute diarrhea. Rotavirus and enterotoxigenic E. coli are the major infectious causes of acute diarrhea. Other causes are certain drugs, food allergies, and systemic infections (e.g. urinary tract infection and otitis media).

The most important assessment in the evaluation of diarrhea is to look for the degree of dehydration. It is assessed as follows:

Based on the degree of dehydration after history and examination, the

estimated fluid loss is calculated as follows:

The treatment plans A,B, and C are explained in the pearl below.

Parameters	No dehydration	Some dehydration	Severe dehydration
Condition	Well alert	Restless, irritable	Lethargic orunconscious,flop py
Eyes	Normal	Sunken	Very sunken and dry
Tears	Present	Absent	Absent
Mouth and ton gue	Moist	Dry	Very dry
Thirst	Drinks normallynot thirst y	Thirsty, drinks eagerly	Drinks poorly or isnot able to drink
Skin pinch	Goes back quickly	Goes back slowly	Goes back very slowly
Treatment	Plan A	Weigh the patient, if possible, and use treatment Plan B	Weigh the patient and use tre atmentPlan C urgently

Degree of dehydration	Assessment of fluid loss
No dehydration	less than 50 mL/kg
Some dehydration	50-100 mL/kg
Severe dehydration	more than 100 mL/kg

Solution to Question 9:

Among the given options, babies on artificial feeds and babies of working/single parents as considered at-risk babies.

The following are considered at-risk infants:

• Birth weight <2.5 kg

• Twins

• Birth order of 5 or more

• Artificial feeding

• Weight <70% of the expected weight (i.e. II and III degrees of malnutrition)

• Failure to gain weight during 3 successive months

• Children with protein-energy malnutrition (PEM) and diarrhea

• Working mother or single parent

It is important to identify children at risk and give special intensive care. This is done because these children contribute largely to perinatal, neonatal, and infant mortality.

Solution to Question 10:

The above image is suggestive of ectopia vesicae or bladder exstrophy.

Ectopia vesicae/bladder exstrophy is a congenital abnormality that occurs when the lower abdominal wall does not form properly and the bladder is open and exposed.

The clinical features of bladder exstrophy include:

• Separation of the pubic bones with non-united symphysis pubis

• Epispadias

• In males- broad and short penis and bilateral inguinal hernia

• In females- bifid clitoris

Management involves a staged surgical reconstruction of the bladder and genitalia. Closure of the abdominal and bladder wall is done in the first year of life. This often requires an iliac osteotomy to relax the tension on the bladder and repair the abdominal wall during wound-healing.

Later, reconstruction of the bladder neck, sphincters, epispadias, and other genital defects is done.

Less satisfactorily, urinary diversion can be carried out by means of ureterosigmoid anastomosis, an ileal or colonic conduit, or continent urinary diversion.

Long-term complications include:

• Stricture at the site of the anastomosis with bilateral hydronephrosis and infection

• Hyperchloremic acidosis

• An increased (20-fold) risk of tumor formation (adenoma and adenocarcinoma) at the site of a ureteroscopic anastomosis.

Other options:

Option B: Omphalocele is a midline anterior abdominal wall defect. It occurs due to the failure of the physiological umbilical hernia to return to the abdominal cavity. The extruded abdominal contents are covered in a sac. It is associated with elevated alpha-fetoprotein levels and chromosomal abnormalities.

The image given below shows omphalocele.



Option C: In a persistent vitellointestinal duct, the vitelline duct remains patent over its entire length. This results in the formation of direct communication between the umbilicus and the intestinal tract. A fecal discharge may then be found at the umbilicus.

Option D: Gastroschisis is a condition where intestinal loops herniate through the defect in the abdominal wall. The defect usually lies to one side of the umbilicus (right side commonly). The loops of the bowel are not covered by amnion as they herniate through the abdominal wall directly into the amniotic cavity. It is associated with elevated alpha-fetoprotein levels.



Solution to Question 11:

The above scenario is suggestive of a foreign body in the airway, most likely in the bronchi. In this condition, auscultation is likely to reveal a focal area of decreased air entry due to incomplete airway obstruction

It is more commonly seen in children below 4 years. The most foreign bodies in the airway are food items. The foreign bodies can be non-irritating types (plastic, glass, or metallic foreign bodies) and irritating types (vegetables, peanuts, beans, and seeds).

The clinical presentation depends upon the site of the obstruction.

Foreign bodies in bronchi are more likely to enter the right bronchus because it is wider and more in line with the tracheal lumen.

The initial evaluation is done with a plain X-ray chest in posteroanterior and lateral views (expiratory films are useful). It may reveal:

• The radio-opaque foreign body- size, shape, and location

• In complete airway obstruction- lobar or segmental atelectasis

• In incomplete airway obstruction- the ball valve mechanism is produced when a foreign body incompletely obstructs a bronchus allowing air to enter inside but not leave. This leads to the appearance of hyperinflation or obstructive emphysema of the affected area when an expiratory X-ray film is taken as the air is not allowed to escape.

Management of foreign body airway is as follows:

• Laryngeal foreign body- Heimlich maneuver should be attempted. If it fails, cricothyrotomy or emergency tracheostomy should be done

• Foreign bodies in the trachea and bronchi are removed through a rigid bronchoscope after adequate hydration and emptying of the stomach.

• Appropriate antibiotics are given to prevent secondary infection.

Option D: Laryngotracheobronchitis or croup is caused by the parainfluenza virus. It commonly affects children of age 6 months-3 years. The clinical features include barking cough, biphasic stridor, and low-grade fever. The chest X-ray shows a narrowing of the subglottic region known as the steeple sign.

The image given below shows the steeple sign seen in croup.

Site of foreign bodies	Signs and symptoms
Larynx	Complete obstruction causes asphyxiaIncomplete obstruction causes croup, hoarseness, cough, stridor, and dyspnea.
Trachea	Dysphonia, dysphagia, dry cough, and biphasic stridor.
Bronchi	May be asymptomatic.Asymmetric breath sound/air entry, coughing, and breathing.



Solution to Question 12:

The clinical scenario is suggestive of Henoch Schonlein's purpura (HSP). Glucocorticoids are used to treat HSP.

Henoch-Schönlein purpura (HSP), also known as IgA vasculitis, is a common small vessel vasculitis in childhood. It is a type III hypersensitivity reaction characterized by the deposition of IgA in the small vessels of the skin, joints, gastrointestinal tract, and kidney, resulting in vasculitis.

The exact etiology of HSP remains unknown. But, it usually occurs following an upper respiratory tract infection caused by group A Streptococcus, Staphylococcus aureus, Mycoplasma, and Adenovirus.

Children usually present with the following features:

• Palpable purpura is symmetric and occurs in the lower extremities, extensor aspect of the upper limbs, and buttocks.

• Arthralgia and arthritis- commonly affecting hip, knee, or ankle joints.

• Gastrointestinal manifestations like abdominal pain, vomiting, diarrhea, and intussusception.

• About 30% of the patients with HSP develop renal manifestations, presenting as microscopic hematuria, hypertension, and progressive glomerulonephritis.

Glucocorticoids such as prednisolone are used for the treatment of gastrointestinal manifestations in HSP. Nephritis due to IgA vasculitis should be treated with immunosuppressants like prednisolone and azathioprine.

The image given below shows palpable purpura in the lower limbs seen in HSP.

The criteria for diagnosis of childhood HSP are:

Palpable purpura with at least one of the following-

• Diffuse abdominal pain

• Any biopsy showing predominant IgA deposition

• Arthritis or arthralgia

• Renal involvement (proteinuria >3 g/24 hr), hematuria, or red cell casts

Lab findings seen in HSP are nonspecific and include leukocytosis, mild anemia, and elevated erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP). The platelet count is normal in HSP. A skin biopsy specimen can be useful in confirming leukocytoclastic vasculitis with IgA and C3 deposition by immunofluorescence.

The image given below shows IgA and C3 deposition in vessels seen on immunofluorescence of skin biopsy in HSP.



Other options:

Immunosuppressants like azathioprine, methotrexate and cyclosporine are used in a variety of rheumatologic disorders like vasculitis, arthritis, inflammatory bowel diseases like Crohn's disease and ulcerative colitis, autoimmune hepatitis, and lupus nephritis.

Solution to Question 13:

The true statement about minimal change disease is that it shows a good response to steroids.

Minimal change disease is the most common cause of nephrotic syndrome in children. It is more common in males and appears between 2 to 6 years of age.

The causes of MCD are mostly idiopathic. It is usually associated with Hodgkin's lymphoma (due to the increased production of cytokines), respiratory infections, prophylactic immunizations, eczema, and rhinitis.

Clinically, they mostly present with proteinuria. The child presents with a history of edema that is initially noted around the eyes and gradually progresses into generalized edema. Hypertension and hematuria are uncommon.

Levels of the following proteins are increased in nephrotic syndrome:

• Fibrinogen - leads to hypercoagulability.

• Lipoproteins - due to increased synthesis.

Levels of the following proteins are decreased in nephrotic syndrome as a consequence of selective proteinuria:

• Albumin—the protein with the largest proportion of loss in nephrotic syndrome

• Transferrin

• Cholecalciferol-binding protein

• Thyroxine-binding globulin (TBG)

• Ferritin

• Coagulation factors

• Antithrombin III



Diagnosis can be made by urinalysis showing proteinuria, ranging from 3+ to 4+ protein on urine dipstick and a spot urine protein/creatinine ratio (PCR) >2. The proteinuria principally contains albumin, hence known as selective proteinuria. Serum albumin levels are <2.5 g/dL and there is an increase in serum cholesterol and triglyceride levels.

Biopsy findings include the following:

• Light microscopy- glomeruli appear normal (hence called "minimal change")

• Immunofluorescence studies- no Ig or complement deposits

• Electron microscopy- diffuse effacement of foot processes of visceral epithelial cells/podocytes

The image given below shows biopsy findings seen in minimal change disease on light microscopy.



The illustration given below shows the effacement of foot processes of podocytes as seen in MCD.



The majority of children will respond to corticosteroid (prednisone/prednisolone) treatment and will achieve complete remission (urine PCR<0.2 or <1+ protein on urine dipstick).

Solution to Question 14:

The clinical scenario of diarrhea, steatorrhea, weight loss, and growth retardation, along with positive class II HLA-DQ2, is consistent with a diagnosis of celiac disease. Patients with celiac disease are advised to avoid gluten.

The dietary recommendation for celiac disease is as follows:

Celiac disease/celiac sprue/gluten-sensitive enteropathy is an immune-mediated disorder triggered by gluten-containing foods. The presence of one of the two locus antigens HLA-DQ2 or

HLA-DQ8 is necessary, but not sufficient for the development of this condition. Detection of these is not useful for the diagnosis, however, their absence helps to exclude celiac disease.

The pathogenesis involves an inflammatory response to gluten and gliadin (gluten-derived peptide). In the small intestine, gliadin is deamidated by the tissue transglutaminase (tTG), an intestinal enzyme. The deamidated gliadin is more immunogenic and is presented to helper T cells, which mediate mucosal inflammation and cause tissue damage.

Children usually present with diarrhea, weight loss, and growth failure. Other features include malabsorption, bloating, irregular bowel habits, iron deficiency anemia, osteoporosis, ataxia, and abnormal liver enzymes.

Screening for celiac disease is done by testing for serum antibodies, including tissue transglutaminase IgA, anti-endomysial, and deaminated anti-gliadin antibodies. The diagnosis in patients with positive antibodies is confirmed by biopsy of the second part of the duodenum or proximal jejunum. Microscopic findings of biopsy show intraepithelial CD8+ lymphocytes, villous atrophy, and crypt hyperplasia.

The image given below shows microscopic findings seen in celiac disease.

Diet avoided (gluten)“BROW”	Diet Allowed“So MR P ”
Barley	Soya bean
Rye	Maize
Oats	Rice
Wheat	Potato



Other options:

Option A: Fat-free diet is advised for various disorders causing steatorrhea. These include pancreatic insufficiency, cystic fibrosis, and familial hypercholesterolemia.

Option B: Lactose-free diet is advised for lactose intolerance. It occurs due to lactase deficiency. Patients usually present with diarrhea, abdominal pain, gassiness, and bloating. A lactose tolerance test or breath hydrogen test may be used for the diagnosis.

Option C: Low-carbohydrate diet is advised for diabetes mellitus.

Paediatrics NEET 2023

Question 1:

A newborn presented with chest retractions, dyspnea, and lethargy. The pediatrician diagnosed the baby with respiratory distress syndrome. This occurs due to the deficiency of:

a) Dipalmitoyl inositol

b) Lecithin

c) Sphingomyelin

d) Dipalmitoylphosphatidylethanolamine

Question 2:

A 10-year-old child weighing 30 kg presents with a history of loose stools for 2 days. On examination, there is severe dehydration. Laboratory investigations are as follows. What is the initial management as per ISPAD guidelines?

a) Manage ABC, NS 20 mL/kg and start insulin after 1 hour

b) Manage ABC, NS 20 mL/kg along with insulin 0.1 IU/kg/hr

c) Manage ABC, NS 10 mL/kg along with insulin 0.1 IU/kg/hr

d) Manage ABC, NS 10 mL/kg and start insulin after 1 hour

Question 3:

Which of the following is the best sign to indicate adequate growth in an infant with a birth weight of 2.8 kg?

a) Increase in length of 25 centimeters in the first year

b) Weight gain of 300 grams per month till 1 year

c) Anterior fontanelle closure by 6 months of age

d) Weight under 75th percentile and height under 25th percentile

Question 4:

A child is brought to the hospital with respiratory distress and biphasic stridor. The radiograph is shown below. What is the diagnosis?



a) Acute epiglottitis

b) Acute laryngotracheobronchitis

c) Foreign body aspiration

d) Laryngomalacia

Question 5:

A 3-month-old baby comes with complaints of deafness, cataract, and patent ductus arteriosus. Which of the following is the most likely diagnosis?

a) Congenital herpes simplex virus infection

b) Congenital toxoplasmosis

c) Congenital cytomegalovirus infection

d) Congenital rubella syndrome

Question 6:

Chloride level in sweat is used in the diagnosis of which disease?

a) Phenylketonuria

b) Cystic fibrosis

c) Gaucher's disease

d) Osteogenesis imperfecta

Question 7:

A 3-month-old baby presents with jaundice and clay-colored stools. Lab investigation reveals that the baby has conjugated hyperbilirubinemia. The liver biopsy shows periductal proliferation. What is the most likely diagnosis?

a) Crigler-Najjar syndrome

b) Rotor syndrome

c) Dubin-Johnson syndrome

d) Biliary atresia

Question 8:

A child presents with fever and a rash spreading from the face, behind cheeks, and buccal mucosa to other body parts. On examination, Koplik's spot is present. What is the likely diagnosis?

a) Measles

b) Rubella

c) Varicella

d) Mumps

Question 9:

An infant presents with hepatosplenomegaly and thrombocytopenia. Neuroimaging with CT shows periventricular calcifications. What is the most likely diagnosis?

a) Congenital rubella syndrome

b) Congenital herpes simplex virus infection

c) Congenital toxoplasmosis

d) Congenital cytomegalovirus infection

Question 10:

A 3-week-old infant presents with a cough and sore throat. The mother reports that the infant develops a paroxysm of cough followed by apnea. The total leucocyte count is >50,000 cells/µL. Which of the following drugs will you prescribe for this patient?

a) Azithromycin

b) Amoxicillin

c) Cotrimoxazole

d) Clarithromycin

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	d
6	b
7	d
8	a
9	d
10	a

Detailed Explanations

Solution to Question 1:

Respiratory distress syndrome in a neonate is likely due to the deficiency of lecithin.

Respiratory distress syndrome (RDS), also known as hyaline membrane disease, is due to the decreased production and secretion of the lung surfactant. Surfactant is a lipoprotein primarily made of phospholipids such as lecithin (otherwise known as dipalmitoylphosphatidylcholine). The other minor components are phosphatidylglycerol, apoproteins, and cholesterol. It is synthesized in type II pneumocytes/granular pneumocytes in the endoplasmic reticulum (shown below).



©MARROW

RDS is commonly seen in preterm babies less than 34 weeks of gestation. Surfactant deficiency leads to an increase in the surface tension over the alveoli, causing the alveoli to collapse easily during expiration. Therefore, the baby has to put more effort to expand the lungs. Tachypnea, cyanosis, nasal flaring, prominent expiratory grunting, and intercostal and subcostal retractions are some of the common findings seen in this condition.

Chest x-ray shows features such as reticulogranular pattern, ground-glass opacity, and air bronchogram. White-out lungs can be seen on chest x-ray in severe diseases.

Management in mild to moderate RDS is with continuous positive airway pressure (CPAP), where the positive pressure keeps the alveoli open in a spontaneously breathing baby. Severe cases will require mechanical ventilation. Steroids increase the production of the surfactant, and therefore, steroids are administered antenatally to mothers with preterm labor.

Other options -

Option A: Dipalmitoyl inositol, or phosphatidylinositol, acts as a precursor for the secondary messenger system. It also helps in cell signaling and membrane trafficking.

Option C: Sphingomyelin is a dominant sphingolipid in the plasma membrane. The lecithin/sphingomyelin ratio in amniotic fluid helps in determining fetal lung maturity.

Option D: Dipalmitoyl ethanolamine is present in the cell membrane.

Solution to Question 2:

The given clinical scenario with severe dehydration, hyperglycemia, glucosuria, and metabolic acidosis is suggestive of diabetic ketoacidosis. Based on the ISPAD guidelines, the initial management includes management of the airway, breathing, and circulation (ABC) and fluid replacement with an IV bolus of 20 mL/kg NS (Normal saline). This is followed by an IV insulin infusion of 0.05 - 0.1 U/kg/hr, at least 1 hour after starting fluid replacement therapy.

The International Society for Pediatric and Adolescent Diabetes (ISPAD) guidelines provide evidence-based recommendations for the management of DKA (diabetic ketoacidosis) in children

and adolescents.

Diabetic ketoacidosis is the most serious complication of type 1 diabetes mellitus. Clinical manifestations of diabetic ketoacidosis include:

• Dehydration, tachypnea with deep, sighing respiration (Kussmaul breathing)

• Nausea, vomiting, and abdominal pain that may mimic an acute abdominal condition

• Confusion, drowsiness, and coma in severe cases.

The diagnosis of DKA is based on the triad of hyperglycemia, ketosis, and metabolic acidosis. All three biochemical criteria are required to diagnose DKA

• Hyperglycemia (blood glucose >200 mg/dl)

• Venous pH < 7.3 or serum bicarbonate <18 mmol/L

• Ketonemia or ketonuria

Initial management includes management of the airway, breathing, and circulation (ABC) according to Pediatric Advanced Life Support (PALS) guidelines. Following that emergency assessment is done, which includes:

• Immediate measurement of blood glucose, blood or urine ketones, serum electrolytes and blood gases

• Assessment of level of consciousness.

• Two peripheral intravenous (IV) lines should be inserted.

The goal of therapy is to correct dehydration, correct acidosis, reverse ketosis, and gradually blood glucose concentration to near normal. Monitor for acute complications, and identify and treat any precipitating event.

Fluid replacement should always begin before starting insulin therapy. One or more boluses of 0.9% saline at 20 mL/kg are given to restore peripheral circulation. Insulin therapy with 0.05–0.1 U/kg/hr must begin at least 1 hour after starting fluid replacement therapy. Insulin of 0.05 U/kg/hr can be considered if pH is > 7.15.

Potassium replacement is given only when potassium is <5.5 mmol/L. In rare cases of hypokalemia with potassium <3.0 mmol/L, defer insulin treatment.

Bicarbonate administration is not recommended except for treatment of severe acidosis (pH < 6.9) or in cases of life-threatening hyperkalemia.

Solution to Question 3:

The best sign to indicate adequate growth in an infant is an increase in length of 25 centimeters in the first year. The growth velocity is the rate of increase in height in one year and varies with age.

The average growth velocity is 25 cm/year in the first year (birth-12 months) and an infant will be 75 cm tall at 1 year. The height velocity is 10 cm/year from 12-24 months and at 2 years, the height is 90 cm. The height velocity is 8 cm/year from 24-36 months. At 4 years, the child is 100 cm tall. After this, the child gains about 6 cm every year until 12 years.

If growth velocity is lower than expected for age, the child is likely to be suffering from a pathological cause of short stature.

Other options:

Option B: During the first 3 months of age, the weight gain is about 20-40 grams per day. From 4 months of age till 1 year of age, the weight gain is about 400 grams per month. The weight of an infant doubles his birth weight at 5 months of age and triples at 1 year of age and quadruples at 2 years of age.

Option C: The anterior fontanelle closes by 18 months of age.

Option D: According to WHO growth charts, weight under the 75th percentile and height under the 25th percentile do not indicate adequate growth.

Solution to Question 4:

The diagnosis in this child with respiratory distress and biphasic stridor is acute laryngotracheobronchitis, which is caused by the parainfluenza virus. The x-ray finding is the 'steeple sign' due to subglottic narrowing.

The age of presentation of laryngotracheobronchitis ranges from 6 months to 6 years. The clinical features include barking cough, inspiratory stridor, hoarseness, and low-grade fever.

Management:

• Racemic epinephrine nebulization leads to the rapid improvement of upper airway obstruction. Epinephrine constricts pre-capillary arterioles in the upper airway mucosa and decreases capillary hydrostatic pressure, leading to fluid resorption and improvement in airway edema.

• Dexamethasone is administered by the least invasive route (Oral > IV > IM) for all the grades of croup.

• Oxygen should be administered to children who are hypoxemic (SpO2 <92 %)

Antibiotics are not indicated in acute laryngotracheobronchitis.

Other options:

Option A: Acute epiglottitis is a medical emergency presenting in children with a rapidly progressive course of fever, sore throat, dyspnea, and respiratory obstruction, followed by a toxic appearance and drooling. The neck is hyperextended and stridor is a late finding. Radiography shows a classic thumb sign.

Option C: Foreign body aspiration initially presents with a paroxysm of cough, choking, and gagging, accompanied by new-onset wheezing and asymmetric breath sounds.

Option D: Laryngomalacia is a congenital laryngeal anomaly presenting in children with low-pitched inspiratory stridor exacerbated by exertion.

Solution to Question 5:

The most likely diagnosis is congenital rubella syndrome based on the clinical findings.

Triad of Gregg is seen in babies with congenital rubella syndrome. The features include:

• Patent ductus arteriosus (PDA)

• Cataract

• Sensorineural Hearing loss (SNHL)

Rubella or German measles is a serious infection in pregnancy that causes abortions or severe fetal defects if infection occurs in the first trimester. Transmission of the virus occurs through nasopharyngeal secretions (droplet transmission). The mother usually has a fever and generalized body rash. The rash is maculopapular in nature that begins from the face and spreads to the trunk and extremities later. The incidence and severity reduce as the pregnancy advances and is rare after 20 weeks of gestation.

There is no specific treatment for rubella virus infection in mothers. Post-exposure passive immunization can be beneficial if given within 5 days of exposure to droplets. The incidence of this disease has reduced due to the availability of MMR vaccines. MMR vaccines can be administered only to non-pregnant females as it is a live, attenuated virus. Pregnancy should be avoided for at least 1 month after vaccination.

Other options:

Option A: Herpes simplex infection can be acquired in utero, intrapartum, or during the neonatal period. The intrauterine infection causes skin vesicles, eye findings (chorioretinitis, keratoconjunctivitis), and microcephaly/hydranencephaly. Intrapartum infection presents with either:

• Skin, eye, and mouth disease

• Encephalitis

• Disseminated disease with sepsis-like presentation

Option B: Congenital toxoplasmosis occurs due to primary infection in immunologically normal pregnant women. First-trimester infection results in severe disease and third-trimester disease result in milder disease. It presents with a triad of chorioretinitis, hydrocephalus, and intracerebral calcifications.

Option C: Pregnant females infected with cytomegalovirus (CMV) present with infectious mononucleosis-like symptoms, that is fever, pharyngitis, lymphadenopathy, and polyarthritis. Transmission to the fetus increases as pregnancy advances and is maximum in the third trimester. Congenital infection includes growth restriction, microcephaly, intracranial (periventricular) calcifications, chorioretinitis, mental and motor deficits, hepatosplenomegaly, hemolytic anemia, thrombocytopenic purpura. Most infected neonates are asymptomatic at birth and later present with these symptoms. Sensorineural hearing loss is a common sequela of congenital CMV infection.

Other infections which cause congenital malformations are:

Organism	Defects
Toxoplasma	Hydro/microcephaly, convulsions, chorioretinitis, diffuse cerebral calcifications

Organism	Defects
Parvovirus	Non-immune hydrops
Syphilis	Hutchinson's triad:Interstitial keratitisHutchinson's teeth Eighth nerve deafness

Solution to Question 6:

The chloride level of sweat, known as the sweat test, is used to diagnose cystic fibrosis (CF). A sweat chloride level of >60 mmol/L is diagnostic of CF.

CF is an autosomal recessive disorder with a mutation in the cystic fibrosis transmembrane regulator (CFTR) gene on chromosome 7q.

The below image depicts the organs affected by CF:



Diagnostic tests include:

1. Immunoreactive trypsinogen for newborn screening of cystic fibrosis.

2. Sweat chloride test: Increased chloride concentration >60 mmol/L is diagnostic.

3. Genetic testing for CFTR gene mutations is the gold standard and should be done for all cases.

4. Nasal transepithelial potential difference can be estimated in equivocal cases.

Diagnostic criteria:

Presence of

• Clinical symptoms affecting at least one organ system (respiratory, gastrointestinal, or genitourinary) or

• Positive newborn screening or

• A sibling with cystic fibrosis.

Evidence of dysfunction of the CFTR gene by

• Two positive sweat chloride test results obtained on separate days or

• Identification of two CFTR mutations by genetic testing or

• An abnormal measurement of nasal potential difference.

Other options:

Option A: Phenylketonuria occurs due to the deficiency of phenylalanine hydroxylase, leading to elevated phenylalanine levels. Diagnosis is confirmed by measuring plasma phenylalanine concentration.

Option C: Gaucher disease is a lysosomal storage disorder caused by a deficiency of β-glucosidase. Diagnosis is confirmed by fibroblast/leukocyte β-glucosidase activity and gene mutations.

Option D: Osteogenesis imperfecta is a genetic connective tissue disorder due to defective type-1 collagen. DNA sequencing is the first diagnostic test.

Solution to Question 7:

The most likely diagnosis in this infant with jaundice, clay-colored stools, and periductal proliferation is biliary atresia.

Biliary atresia presents with progressive jaundice, persistent acholic stools, and palpable liver with abnormal consistency. Laboratory investigations show conjugated hyperbilirubinemia. Ultrasonography reveals a triangular-cord sign (cone-shaped fibrotic mass cranial to the bifurcation of the portal vein).

HIDA test is used for screening and liver biopsy is diagnostic for biliary atresia. Biopsy shows proliferation of the bile ductule, the presence of bile plugs, and portal or perilobular edema and fibrosis, with the basic hepatic lobular architecture intact.

Other options:

Option A: Crigler-Najjar type 2 is an autosomal recessive disease with non-hemolytic unconjugated hyperbilirubinemia caused by homozygous missense mutations in UGT1A1, resulting in reduced (partial) enzymatic activity.

Option B: Rotor syndrome is an autosomal recessive disorder presenting with asymptomatic mild and fluctuating conjugated hyperbilirubinemia with normal liver histology.

Option C: Dubin-Johnson syndrome is an autosomal recessive disorder due to the absent function of multiple drug-resistant protein-2. It presents with abdominal pain, mild icterus, and conjugated hyperbilirubinemia. Biopsy shows black liver due to an accumulation of epinephrine metabolite.

Solution to Question 8:

The likely diagnosis is measles based on the clinical symptoms and the presence of Koplik spots.

Koplik spots are discrete red lesions with central bluish-white spots seen on the buccal mucosa at the level of premolars.

The measles virus has an incubation period of 8-12 days, after which the prodrome phase begins with mild fever, conjunctivitis, upper respiratory tract infection, and pathognomic Koplik spots. This follows the appearance of maculopapular rash after 4th day of fever, which begins at the face and spreads to the torso and extremities. Lymphadenopathy may be present. Treatment is supportive. The most common complication of measles is acute otitis media.

The image below shows Koplik spots.



Other options:

Option B: Rubella presents with low-grade fever, centrifugal macular rash, and prominent lymphadenopathy (suboccipital, postauricular, and anterior cervical lymph nodes). Forchheimer spots (tiny rose-colored lesions on the soft palate) may be present.

The image below shows Forchheimer spots.

Option C: Varicella presents with fever, malaise, and anorexia, followed by a pruritic maculopapular rash, which forms vesicles with a characteristic "cropping" distribution (simultaneous presence of lesions in various stages of evolution).

Option D: Mumps presents with fever, headache, parotitis, and ipsilateral auricular pain. A morbilliform rash may occur.

Solution to Question 9:

The most likely diagnosis in this infant is congenital cytomegalovirus (CMV) infection.

The clinical manifestations are hepatosplenomegaly, petechial rashes, jaundice, chorioretinitis, and microcephaly. Head neuroimaging shows intracranial periventricular calcifications. One of the long-term sequelae is sensorineural hearing loss. Diagnosis is made by isolating the virus from the urine or blood by polymerase chain reaction (PCR).

There is no definitive treatment for congenital CMV infection. Treatment with ganciclovir is indicated only in patients with progressive neurologic disease and deafness. The images below show periventricular calcifications and a blueberry muffin baby.

Other options:

Option A - Congenital rubella syndrome is due to the rubella virus. The classical triad of congenital rubella syndrome is deafness, cataract, and congenital heart disease. Patent ductus arteriosus is a common cardiac defect. Eye findings also include salt and pepper retinopathy as shown below.



Option B - Herpes simplex infection in neonates presents with skin vesicles, eye findings (chorioretinitis, keratoconjunctivitis), and microcephaly/hydranencephaly. It can also present as localized skin, eye, and mouth disease, encephalitis, or disseminated disease with a sepsis-like presentation.

Option C - Congenital toxoplasmosis is due to Toxoplasma gondii. A classic triad of chorioretinitis, hydrocephalus, and intracerebral calcifications is seen. Blueberry muffin baby appearance is also seen in this. Calcifications occur randomly throughout the brain in congenital toxoplasmosis while periventricular calcification is characteristic of congenital CMV infection.

Solution to Question 10:

The given clinical scenario of paroxysm of cough, apnea, and leukocytosis (>50,000 cells/µL) in a neonate is suggestive of pertussis. Azithromycin is the recommended drug of choice for the treatment of pertussis in neonates (< 1 month of age).

Macrolides are recommended for the treatment of pertussis. Azithromycin is the drug of choice in all age groups, for treatment or postexposure prophylaxis. Post-exposure prophylaxis is given to household close contacts and high-risk contacts (infants, immunocompromised, pregnancy).

Other options:

Option B: Amoxicillin is not used for the treatment of pertussis due to its inefficiency in the nasopharynx.

Option C: Trimethoprim-sulfamethoxazole (cotrimoxazole) is recommended as an alternative for infants >2 months old and children who are allergic to macrolides.

Option D: Clarithromycin is an alternative drug that can be used in infants aged >2 months. In children less than 2 months of age, it is contraindicated due to the risk of kernicterus.

Paediatrics INI-CET 2023

Question 1:

In a baby born with meconium-stained liquor, which of the following are recommended as per Neonatal Resuscitation guidelines?

a) 1, 2, 3, 4

b) 2, 4

c) 3, 4

d) 1, 3, 4

Question 2:

Which of the following has the least risk of perinatal transmission?

a) HSV

b) CMV

c) Rubella

d) Hepatitis B

Question 3:

Which of the following is a new drug approved by the FDA for the treatment of Rett syndrome?

a) Olcegepant

b) Lasmiditan

c) Trofinetide

d) Erenumab

Question 4:

A 6-year-old boy was brought with a history of chronic cough, and foul-smelling stools. His sibling who had similar complaints died a few years back. Sweat chloride test and CFTR gene studies were done. CFTR gene codes for proteins involved in the transport of:

a) Na+

b) HCO_3^-

c) Cl^-

d) K^+

Question 5:

The perinatal transmission of infection shows the spectrum of anemia, possibly myocarditis, high output cardiac failure, and ascites. Identify the causative agent.



a) Treponema pallidum

b) CMV

c) Parvovirus B19

d) Toxoplasma gondii

Question 6:

In which of the following conditions is live vaccination not contraindicated?

a) 4 only

b) 1 only

c) 1 and 3

d) 2 and 3

Question 7:

Nada's Minor Criteria includes:

a) 3 and 4

b) 1, 2 and 3

c) 1, 2, 3 and 4

d) 2, 3 and 4

Question 8:

An 18-month-old child with a 3-day history of watery diarrhea and vomiting presented with altered sensorium. Which of the following are differential diagnoses?

a) 1, 2, 3, 4

b) 1, 2, 3

c) 1, 3, 4

d) 2, 3, 4

Question 9:

The causative agent of this condition is associated with which of the following conditions?

a) Gingivostomatitis

b) Pure red cell aplasia

c) Molluscum contagiosum

d) Kaposi sarcoma

Question 10:

A 4-year-old boy presented with a history of fever, conjunctival congestion, and erythematous desquamation of the skin as shown in the images below. A 2D echo shows a coronary aneurysm. What is the first line of treatment for his condition?

a) Aspirin

b) IVIG

c) Clopidogrel

d) LMWH

Question 11:

Which of the following is not associated with Down's syndrome?

a) Conductive hearing loss

b) Short stature

c) Hypothyroidism

d) Caudal regression

Question 12:

Which of the following is correct about congenital CMV?

a) 40-50 % of the cases are symptomatic at birth

b) Optimum sample for PCR is blood/ urine/ saliva sample within 2-3 weeks of birth

c) If asymptomatic at birth, then there is a higher risk of developing SNHL later

d) Most common congenital infection after congenital syphilis

Question 13:

A 3.5 kg term neonate born to a mother with eclampsia, did not cry at birth. Which of the following is the criteria for perinatal asphyxia?

a) APGAR persists at 4-7 for > 5 minutes

b) Cord pH < 7.2

c) Hypocalcemia

d) Hypotonia

Question 14:

Choose the correct statements for cystic fibrosis:

a) i and iii

b) ii and iii

c) i and iv

d) ii and iv

Question 15:

One-day-old term neonate presented with respiratory distress, tachypnea, SpO2-80%, and scaphoid abdomen. Which of the following cannot be used in resuscitation?

a) Positive pressure ventilation

b) Endotracheal intubation

c) Bag and mask ventilation

d) Oxygen suplementation

Question 16:

A child with type 1 DM presents with acidosis, and blood sugar 400 mg/dl. What is the further management?

a) IV insulin stat

b) Normal saline with SC insulin stat

c) Normal saline followed by insulin infusion after 1 hour

d) Normal saline and insulin infusion immediately

Question 17:

A 9-month-old child is brought to the PHC with a fever and cough. She is feeding well and has normal urine output. On examination, a respiratory rate of 48 breaths/min is found and no chest indrawing is seen. What is the next step of management?

a) Explain danger signs and send home

b) No pneumonia - Give amoxyclav + paracetamol and send home

c) Start IV ceftriaxone

d) Refer as the child has severe pneumonia

Answer Key

Question No.	Correct Option
1	c
2	c
3	c
4	c
5	c
6	a
7	a
8	c
9	b
10	b
11	d
12	b
13	d
14	c

15	c
16	c
17	a

Detailed Explanations

Solution to Question 1:

In a baby born with meconium-stained liquor, gentle suction of the nose and mouth is done in the case of a vigorous baby (Statement 3), and positive pressure ventilation (PPV) is done in the case of a non-vigorous baby after early steps (Statement 4).

A baby born through meconium-stained liquor is classified as either a vigorous baby or a non-vigorous baby. A vigorous baby has good breathing and normal tone, whereas, a non-vigorous baby has poor or no breathing with decreased tone. According to neonatal resuscitation guidelines, in a vigorous baby, gentle suction is done and the baby is kept with the mother. In a non-vigorous baby, the initial steps of resuscitation are done and is followed by PPV if the baby is not responding. Routine tracheal suctioning is not recommended (Statement 2).

The response to neonatal resuscitation at various steps is assessed based on the heart rate.

• If the heart rate is less than 100, PPV (positive pressure ventilation) is provided with bag and mask and SPO2 monitoring is done and adequate ventilation is ensured.

• If the heart rate is below 60, then chest compressions are initiated over the lower third of the sternum at a rate of 90 compressions/min.

• Recommended rate of ventilation is 30 inflations/min. (90 compressions + 30 inflation = 120 events/ min). The ratio of compressions to ventilation is 3: 1.

• If the heart rate still remains <60, despite effective compressions and ventilation, then IV epinephrine is given.

Note: For any baby, intrapartum suction of nose and mouth before delivery of shoulder is not recommended. (Statement 1)



Solution to Question 2:

Rubella has the least risk of perinatal transmission.

Rubella or German measles has an intrauterine transmission that causes serious infection in pregnancy which results in abortions or severe fetal defects if infection occurs in the first trimester. The classical triad of congenital rubella syndrome is deafness, cataract, and congenital heart disease.

Other options-

Option A: Herpes simplex infection can be acquired in utero, intrapartum/perinatally, or during the neonatal period. The intrauterine infection causes skin vesicles, eye findings (chorioretinitis, keratoconjunctivitis), and microcephaly/hydranencephaly. Intrapartum infection presents with either:

• Skin, eye, and mouth disease

• Encephalitis

• Disseminated disease with sepsis-like presentation

Option B: Cytomegalovirus, the most common congenital infection with perinatal transmission presents with hepatosplenomegaly, petechial rashes, jaundice, chorioretinitis, and microcephaly. One of the long-term sequelae is hearing loss.

Option D: Hepatitis B virus is associated with significant perinatal transmission. The most important risk factor that determines perinatal transmission is HBeAg.

Solution to Question 3:

Trofinetide is the newly FDA-approved drug for the treatment of Rett syndrome.

The mechanism by which trofinetide exerts therapeutic effects in patients with Rett syndrome is unknown. It is a synthetic analog of the amino■terminal tripeptide of insulin-like growth factor (IGF-1) designed to treat the core symptoms of Rett syndrome by potentially reducing neuroinflammation and supporting synaptic function of the protein fragment glypromate (GPE). IGF-1 is produced by both major types of brain cells – neurons and glia; IGF-1 in the brain is critical for both normal development and response to injury and disease. Trofinetide has been shown to inhibit the production of inflammatory cytokines, inhibit overactivation of microglia and astrocytes, and increase the amount of available IGF-1 that can bind to IGF-1 receptors.

It is Indicated for the treatment of Rett syndrome.

Adverse effects include diarrhea, vomiting, fever, seizure, anxiety, decreased appetite, fatigue, nasopharyngitis

Other options:

Option A: Olcegepant is a calcitonin gene-related peptide (CGRP) antagonist which was considered for the treatment of migraine, but discontinued.

Option B: Lasmiditan is a nontriptan 5HT1F receptor–selective agonist and is used to treat migraine.

Option D: Erenumab is a CGRP (calcitonin gene-related peptide) blocking antibody used in the treatment migraine.

Note: The spelling of the drug in the options has been corrected from Trofinitide to Trofinetide.

Solution to Question 4:

The given clinical scenario is suggestive of cystic fibrosis. In this condition, CFTR gene codes for proteins involved in the transport of Cl- ions.

Cystic fibrosis or mucoviscidosis is an autosomal recessive disorder with a mutation in the cystic fibrosis transmembrane regulator (CFTR) gene on chromosome 7q. The most common CFTR mutation is F508 deletion.

CFTR gene encodes for ATP-gated chloride channels in the apical membranes of epithelial cells. It regulates the flow of ions across epithelial membranes of sweat glands, respiratory airways, pancreatic ducts, etc. The defect leads to high salt concentrations in sweat and viscous luminal secretions in the respiratory and gastrointestinal tracts. Excessive loss of salt in the sweat predisposes young children to salt-depletion episodes. Therefore, these children present with hypochloremic alkalosis, resulting in increased serum bicarbonate levels.

Clinical manifestations/ complications of cystic fibrosis include:

• Sinusitis and nasal polyps

• Respiratory - copious hyper-viscous and adherent pulmonary secretions with abnormal mucociliary clearance obstruct small and medium-sized airways leading to respiratory compromise. The disease can be further complicated by bronchiectasis and allergic bronchopulmonary aspergillosis (ABPA).

• Gastrointestinal -

• Thick and adherent exocrine secretions obstruct pancreatic ducts and impair the production and flow of digestive enzymes

• Tissue destruction of the exocrine pancreas, with fibrotic scarring, cyst proliferation, and loss of acinar tissue - fat malabsorption, steatorrhea, fat-soluble vitamin deficiencies

• Obstruction of intrahepatic bile ducts and parenchymal fibrosis leading to hepatic insufficiency and focal biliary cirrhosis

• Distal intestinal obstructive syndrome in adults and meconium ileus in neonates

• Reproductive - males have infertility due to the congenital absence of bilateral vas deferens and defective sperm transport, resulting in obstructive azoospermia. Females have infertility due to abnormalities of the reproductive tract secretions.

The diagnosis is confirmed if both of the following criteria are met:

• Presence of

• Clinical symptoms affecting at least one organ system (respiratory, gastrointestinal, or genitourinary) or

• Positive newborn screening or

• A sibling with cystic fibrosis

• Evidence of dysfunction of the CFTR gene by

• Two positive sweat chloride test results were obtained on separate days or

• Identification of two CFTR mutations by genetic testing or

• An abnormal measurement of nasal potential difference

Cystic fibrosis requires a multidisciplinary approach to management. The correction of the defect can be via:

• Lumacaftor, an FDA-approved drug that corrects the misfolding and reverts the Phe508del mutation defect

• Ivacaftor potentiates CFTR channel opening and stimulates ion transport

End-stage pulmonary failure requires lung transplantation.

Solution to Question 5:

The given clinical vignette with anemia, myocarditis, high output cardiac failure, ascites and the image showing hydrops fetalis point to a diagnosis of erythema infectiosum (Fifth disease) caused by parvovirus B19.

Parvovirus B19 infects fetal erythroid progenitor cells and causes anemia. Severe anemia leads to high-output cardiac failure and hydrops fetalis. In addition to bone marrow, the liver is also affected causing ascites, because it is involved in erythropoiesis in the early fetal life.

Erythema infectiosum presents as a prodrome of low-grade fever, headache, and a mild upper respiratory tract infection after an incubation period of 4-28 days. A characteristic rash occurs in 3 stages:

• Erythematous facial flushing causing a slapped cheek appearance in the 1st stage

• Diffuse macular erythema spreads to the trunk and proximal extremities in the 2nd stage.

• Central clearing of the macular lesions, giving the rash a lacy, reticulated appearance in the 3rd stage.

The rash is more prominent on extensor surfaces and spares the palms and soles. At this stage, affected children are afebrile and do not appear ill. Children with erythema infectiosum are not likely to be infectious at presentation.

Complications of the disease include:

• Arthralgias or arthritis

• Transient Aplastic Crisis: A transient arrest of erythropoiesis and absolute reticulocytopenia in serum hemoglobin in individuals with chronic hemolytic conditions like sickle cell anemia

• Thrombocytopenic purpura

• Myocarditis

• A papular-purpuric “gloves-and-socks” syndrome

• Neurological: Aseptic meningitis, encephalitis, and peripheral neuropathy, stroke in children with sickle cell disease

• Infection-associated hemophagocytic syndrome in immunocompromised patients

• Maternal infection is associated with nonimmune fetal hydrops and intrauterine fetal demise

There is no data to support the use of IVIG for postexposure prophylaxis, however, it has been used with some success to treat B19-related episodes of anemia and bone marrow failure in immunocompromised children. There is no specific antiviral therapy for B19 infection.

Solution to Question 6:

Live vaccination is not contraindicated in complement deficiency. In DiGeorge syndrome, Wiskott-Aldrich Syndrome, and Ataxia telangiectasia, live vaccines are contraindicated.

In patients with DiGeorge syndrome, Wiskott-Aldrich Syndrome, and Ataxia telangiectasia, there is a risk of severe or fatal infections due to live vaccines because of their impaired T-cell immunity. In contrast, a patient with a complement deficiency has an intact T-cell and B-cell immunity, which are crucial for building an immune response against live vaccines. These patients have an increased susceptibility to certain bacterial infections, especially those caused by Neisseria species, due to defects in the complement system, which is part of the innate immune system. But their adaptive immune system, which is primarily responsible for responding to vaccinations, is generally intact. Therefore, live vaccinations are not typically contraindicated in these patients.

DiGeorge syndrome is due to a developmental defect involving the endodermal derivatives of the 3rd and 4th pharyngeal pouches. Cell-mediated immunity is defective due to decreased T-cells. Affected individuals also have depressed graft rejection, hypocalcemia, and cardiac defects. For severe T cell deficiencies (e.g. severe combined immunodeficiency disease [SCID], complete DiGeorge syndrome), no live viral or live bacterial vaccines should be given.

Wiskott-Aldrich syndrome(WAS) is an X-linked recessive immunodeficiency disorder. It is caused by mutations in the Wiskott-Aldrich syndrome protein (WASp) that primarily affects the T lymphocytes. In patients with partial T cell deficiencies (e.g., partial DiGeorge syndrome, Wiskott-Aldrich syndrome), all live viral vaccines are avoided.

Ataxia telangiectasia is caused due to mutation of the ATM gene. This gene is involved in the repair of double-strand DNA breaks. This defect in DNA repair leads to impaired Ig isotype switching and the formation of antigen receptors. As a result, the patient is immunodeficient with partial T cell deficiency, and hence live vaccines are avoided.

Solution to Question 7:

Nada's Minor Criteria includes abnormal second heart sound and abnormal BP.

Nada's criteria is used to diagnose congenital heart disease in children. The presence of 1 major or 2 minor criteria indicates congenital heart disease.

Major criteria	Minor criteria
Systolic murmur grade 3 or more	Systolic murmur < grade 3
Any diastolic murmur	Abnormal second sound (S2)
Cyanosis	Abnormal ECG
Congestive cardiac failure	Abnormal chest X-ray findings
	Abnormal BP

Solution to Question 8:

The differential diagnoses in a child with a history of watery diarrhea and altered sensorium include severe dehydration, cerebral venous thrombosis, and hyponatremia.

The lack of dysentery(bloody diarrhoea) and the short period of interval between diarrhoeal episodes and the symptom complex, help to rule out Haemolytic Uremic Syndrome (HUS). HUS is a form of microangiopathic hemolytic anemia usually occurring in children less than 5 years of age. There are 2 forms of HUS:

• Typical HUS which is usually followed by a gastrointestinal infection most commonly by E.coli. The infection in turn affects the gastric mucosa and alters the endothelial cells which promotes platelet activation and aggravation.

• Atypical HUS which occurs due to deficiency in certain proteins like complement factor H, factor I, or CD46.

Clinical features of the condition include bloody diarrhea which occurs 5-10 days before the onset of hemolytic uremic syndrome, abdominal pain, fatigue, cramping, and vomiting. The most dangerous complication is renal failure requiring dialysis or renal transplantation. Treatment is

mainly supportive and the nature of the illness is usually self-limiting.

Differential diagnoses in a child presenting with altered sensorium/ seizures with diarrhea:

• Hypo- or hypernatremia

• Hypoglycemia

• Encephalitis or meningitis

• Febrile seizures

• Cerebral or saggital sinus thrombosis

Solution to Question 9:

The given image showing the classical slapped cheek appearance is strongly suggestive of erythema infectiosum. It is caused by Parvovirus. Parvovirus B19 is associated with pure red cell aplasia.

Clinically, parvovirus B-19 infection may have several presentations-

• Erythema infectiosum- Also known as the fifth disease. The incubation period after infection is typically around 1-2 weeks. The hallmark of erythema infectiosum is the characteristic rash, which appears initially as erythematous facial flushing, often described as a “slapped cheek” appearance. Later the rash spreads rapidly or concurrently to the trunk and proximal extremities as diffuse macular erythema. The child may be febrile.

• Transient aplastic crisis- It may be seen in patients with underlying chronic hemolytic anemia like sickle-cell disease, thalassemia, hereditary spherocytosis, and RBC enzyme deficiencies like pyruvate kinase deficiency. Patients with aplastic crisis appear ill with fever, malaise, and pale with tachycardia and tachypnea due to profound anemia. The rash may or may not be present.

• Pure red cell aplasia- Erythrocyte P blood group antigen is the primary cell receptor for the virus. Persistent Parvovirus B19 infection in immunocompromised patients can cause chronic suppression of bone marrow and chronic anemia leading to pure red cell aplasia.

• Maternal infection is associated with non-immune fetal hydrops and intrauterine fetal demise.

• Only about 7% of children with erythema infectiosum have, whereas 80% of adults have joint involvement.

• Infection-associated hemophagocytic syndrome in immunocompromised patients.

The diagnosis of erythema infectiosum is made clinically and does not require investigations routinely. Detection of viral DNA in the blood by PCR or nucleic acid hybridization may be done for immunocompromised cases or for prenatal diagnosis of infection.

It is self-limited and resolves on its own. IVIG may be given to immunocompromised children.

Solution to Question 10:

This child with a history of fever, conjunctival congestion, strawberry tongue, and erythematous desquamation of the skin is likely to have Kawasaki's disease. The first line of treatment in Kawasaki disease is IVIG (intravenous immunoglobulin).

Kawasaki's disease is also known as mucocutaneous lymph node syndrome. It is a vasculitis condition of unknown etiology. Clinical features include a high spiking fever that is unresponsive to antipyretics. The fever can last between 5 days to 3-4 weeks. The five principal clinical criteria for Kawasaki's disease are:

• Bilateral nonexudative conjunctival infection with limbal sparing

• Erythema of the oral and pharyngeal mucosa, strawberry tongue, and red, cracked lips

• Edema and erythema of the hands and feet

• Various forms of rash- maculopapular, erythema multiforme, scarlatiniform

• Nonsuppurative cervical lymphadenopathy

Diagnosis requires fever and at least four of the five principal criteria. Myocarditis develops frequently in acute Kawasaki disease. It presents as tachycardia and reduced ventricular systolic function. The tachycardia will be disproportionate to the fever. Patients may develop coronary artery aneurysms in the 2nd to 3rd week of illness. They are generally asymptomatic and diagnosed on echocardiography. However, if large it can lead to death. Thus, echocardiography should be performed at diagnosis and after 1-2 weeks.

Other symptoms include periungual desquamation, gastrointestinal symptoms (vomiting, abdominal pain, diarrhea), cough, rhinorrhea, and arthritis. Laboratory evaluation will reveal elevated leukocytes, platelets, erythrocyte sedimentation rate, and C-reactive protein levels.

Treatment involves 2g/kg of intravenous immunoglobulin as an infusion over 10-12 hours within 10 days of onset. Moderate to high-dose aspirin should be administered until the patient is afebrile.

<!-- -->

Solution to Question 11:

Among the choices provided, Caudal regression is not associated with Down's syndrome. It occurs more frequently in infants of diabetic mothers (both preconceptionally and early in pregnancy) than in other infants although the exact mechanism is unclear.

Down's syndrome is an autosomal disorder. It was described by Langdon Down in 1866. It is also called trisomy 21 or mongolism. The clinical features of Down's syndrome seen in older children are:

• Short stature

• Small round head

• Narrow tilted eye slits

• Mal-formed ears

• Short broad hands

• Lax limbs

• Mental retardation

Conditions most commonly associated with Down's syndrome:

• Congenital heart disease (in about 35% of cases)

• Gastrointestinal anomalies (in about 10%)

• Chronic serous otitis media (in >50%)

• Conductive hearing loss

• Hypothyroidism

• Alzheimer's disease (in the 30's and 40's)

• Epilepsy (in about 10%)

• Ocular disorders

• Reduced fertility

• Reduced life span (Often due to complications like infections)

The frequency of Down's syndrome increases with rising maternal age. The risk of a woman who is 20 is 1 in 3000 but that of a woman who is 45 is 1 in 50.

It seems to be unaffected by the age of the father.

Note: Genetic counseling: It can either be prospective or retrospective. Prospective genetic counseling refers to the prevention of disease by the prevention or reduction of heterozygous marriage like in sickle cell anemia or thalassemia. Retrospective genetic counseling means that the hereditary disorder has already occurred in the family and methods like contraception, pregnancy termination or sterilization are to be used to avoid any serious health disorder in the unborn baby.

Methods of early diagnosis and treatment are early detection of genetic carriers, prenatal diagnosis, screening of newborn infants, and recognizing pre-clinical cases.

Solution to Question 12:

The correct statement regarding CMV infection is optimum sample for PCR is blood/ urine/ saliva sample within 2-3 weeks of birth

CMV is the most common congenital infection (Option D). The clinical manifestations are hepatosplenomegaly, petechial rashes, jaundice, chorioretinitis, and microcephaly. One of the long-term sequelae is sensorineural hearing loss. Neuroimaging shows intracranial periventricular calcifications. Other manifestations include IUGR, hydrops, microcephaly, pneumonitis, chorioretinitis, seizures, abnormal dentition, and hypocalcified enamel.

Only 10% of the infected babies are symptomatic (Option A). Asymptomatic neonates have lesser chances of having late sequelae i.e. sensorineural hearing loss. About half of the symptomatic infants develop long-term sequelae which include sensorineural hearing loss (SNHL), vision loss, seizure disorders, mental retardation, and developmental delay. However, in asymptomatic infants, only about 13.5% develop long-term sequelae which is commonly SNHL (option C).

Diagnosis can be made by the presence of IgM antibodies in the fetal blood. Isolation of the virus from the urine or blood by PCR is more sensitive. The investigation of choice is PCR or viral culture. It should be done within the first three weeks of life. After this period, it is difficult to distinguish between a congenital and an acquired infection. Since not all affected infants are viremic at birth, urine and/or saliva are more reliable samples than blood (Option B).

There is no definitive treatment for congenital CMV infection. Treatment with ganciclovir is indicated only in patients with progressive neurologic disease and deafness. Below is the image showing the blueberry muffin baby.

Solution to Question 13:

Among the given options, the essential criterion for perinatal asphyxia is hypotonia.

Perinatal asphyxia is an insult to the fetus or newborn due to a lack of oxygen (hypoxia) and/or a lack of perfusion (ischemia) to the various organs. It is often associated with tissue lactic acidosis and hypercarbia.

Essential criteria for perinatal asphyxia:

• Umbilical arterial blood sample indicating prolonged metabolic or mixed acidemia (pH < 7.O).

• Apgar score of 0-3 persisting for >5 minutes.

• Neurological manifestations appearing in the immediate neonatal period Ex:- seizures, coma, hypotonia, or hypoxic ischemic encephalopathy (HIE).

• Neonatal multiorgan dysfunction.

Solution to Question 14:

In cystic fibrosis, there are increased chloride levels in sweat and decreased Chloride levels in pancreatic juice. The chloride level of sweat, known as the sweat test, is used to diagnose cystic fibrosis (CF). A sweat chloride level of >60 mmol/L is diagnostic of CF.

Cystic fibrosis or mucoviscidosis is an autosomal recessive disorder with a mutation in the cystic fibrosis transmembrane regulator (CFTR) gene on chromosome 7q. The most common CFTR mutation is F508 deletion.

CFTR gene encodes for ATP-gated chloride channels in the apical membranes of epithelial cells. The function of the CFTR gene on chloride channels is organ-specific. In sweat glands, the normal function of the CFTR gene is to reabsorb chloride ions. But in the pancreatic ducts and respiratory airways, the normal function of the CFTR gene is to secrete chloride ions. So, in cystic fibrosis, due to defect in the chloride channels, chloride levels in the sweat increase, and chloride levels in the pancreatic juices decreases. Excessive loss of salt in the sweat predisposes young children to salt-depletion episodes. Therefore, these children present with hypochloremic alkalosis, resulting in increased serum bicarbonate levels.

Clinical manifestations/ complications of cystic fibrosis include:

• Sinusitis and nasal polyps

• Respiratory - copious hyper-viscous and adherent pulmonary secretions with abnormal mucociliary clearance obstruct small and medium-sized airways leading to respiratory compromise. The disease can be further complicated by bronchiectasis and allergic bronchopulmonary aspergillosis (ABPA).

• Gastrointestinal -

• Thick and adherent exocrine secretions obstruct pancreatic ducts and impair the production and flow of digestive enzymes

• Tissue destruction of the exocrine pancreas, with fibrotic scarring, cyst proliferation, and loss of acinar tissue - fat malabsorption, steatorrhea, fat-soluble vitamin deficiencies

• Obstruction of intrahepatic bile ducts and parenchymal fibrosis leading to hepatic insufficiency and focal biliary cirrhosis

• Distal intestinal obstructive syndrome in adults and meconium ileus in neonates

• Reproductive - males have infertility due to the congenital absence of bilateral vas deferens and defective sperm transport, resulting in obstructive azoospermia. Females have infertility due to abnormalities of the reproductive tract secretions.

The diagnosis is confirmed if both of the following criteria are met:

• Presence of

• Clinical symptoms affecting at least one organ system (respiratory, gastrointestinal, or genitourinary) or

• Positive newborn screening or

• A sibling with cystic fibrosis

• Evidence of dysfunction of the CFTR gene by

• Two positive sweat chloride test results were obtained on separate days or

• Identification of two CFTR mutations by genetic testing or

• An abnormal measurement of nasal potential difference

Cystic fibrosis requires a multidisciplinary approach to management. The correction of the defect can be via:

• Lumacaftor, an FDA-approved drug that corrects the misfolding and reverts the Phe508del mutation defect

• Ivacaftor potentiates CFTR channel opening and stimulates ion transport

End-stage pulmonary failure requires lung transplantation.

Solution to Question 15:

The given clinical scenario of a one-day term-old neonate presented with respiratory distress, tachypnea, SpO2-80%, and scaphoid abdomen is suggestive of congenital diaphragmatic hernia. Bag and mask ventilation is to be avoided in the management of this baby. Bag and mask ventilation may increase abdominal distension and limit the expansion of the hypoplastic lungs.

Congenital diaphragmatic hernia (CDH) is a birth defect of the diaphragm. Malformation of the diaphragm allows the abdominal organs to push into the chest cavity, hindering proper lung formation and resulting in pulmonary hypoplasia. The most common type of CDH is a Bochdalek hernia and in most cases involves the left side. CDH is a life-threatening pathology in infants and a major cause of morbidity and mortality due to two complications: pulmonary hypoplasia and pulmonary hypertension.

Babies with CDH present with respiratory distress characterized by tachypnea, grunting, use of accessory muscles, and cyanosis. They may also have a scaphoid abdomen and increased chest wall diameter. Auscultation of the chest may reveal bowel sounds with reduced breath sounds.

Diagnosis: A chest radiograph and nasogastric tube passage can confirm the diagnosis of CDH. Prenatal diagnosis can be made using ultrasonography between 16 to 24 weeks of gestation. High-speed fetal magnetic resonance imaging (MRI) can further define the lesion.

Management of CDH:

The immediate step in the management is the naso- or orogastric tube placement for decompression. The airway is to be secured with endotracheal intubation and respiratory distress is stabilised.

Inhaled nitric oxide is used widely for its pulmonary vasodilatory effect. Conventional mechanical ventilation, high-frequency oscillatory ventilation (HFOV), and extracorporeal membrane oxygenation (ECMO) are the 3 main strategies to support respiratory failure in newborn with CDH.

Surgical repair of the defect is to be considered at least 48 hours after stabilization and resolution of pulmonary hypertension. During surgery, the viscera are reduced into the abdominal cavity, and the posterolateral defect in the diaphragm is closed using interrupted, nonabsorbable sutures.

Solution to Question 16:

A given clinical scenario of a child with type 1 diabetes mellitus (DM) presenting with acidosis and a high blood sugar level (hyperglycemia) describes a scenario suggestive of diabetic ketoacidosis (DKA), a serious complication of uncontrolled diabetes. the further management would be Normal saline followed by insulin infusion after 1 hour.

The International Society for Pediatric and Adolescent Diabetes (ISPAD) guidelines provide evidence-based recommendations for the management of DKA (diabetic ketoacidosis) in children and adolescents.

Diabetic ketoacidosis is the most serious complication of type 1 diabetes mellitus. Clinical manifestations of diabetic ketoacidosis include:

• Dehydration, tachypnea with deep, sighing respiration (Kussmaul breathing)

• Nausea, vomiting, and abdominal pain that may mimic an acute abdominal condition

• Confusion, drowsiness, and coma in severe cases.

The diagnosis of DKA is based on the triad of hyperglycemia, ketosis, and metabolic acidosis. All three biochemical criteria are required to diagnose DKA

• Hyperglycemia (blood glucose >200 mg/dl)

• Venous pH < 7.3 or serum bicarbonate <18 mmol/L

• Ketonemia or ketonuria

Initial management includes management of the airway, breathing, and circulation (ABC) according to Pediatric Advanced Life Support (PALS) guidelines. Following that emergency assessment is done, which includes:

• Immediate measurement of blood glucose, blood or urine ketones, serum electrolytes and blood gases

• Assessment of level of consciousness.

• Two peripheral intravenous (IV) lines should be inserted.

The goal of therapy is to correct dehydration, correct acidosis, reverse ketosis, and gradually blood glucose concentration to near normal. Monitor for acute complications, and identify and treat any precipitating event.

Fluid replacement should always begin before starting insulin therapy. One or more boluses of 0.9% saline at 10-20 mL/kg are given to restore peripheral circulation. Insulin therapy with 0.05–0.1 U/kg/hr must begin at least 1 hour after starting fluid replacement therapy. Insulin of 0.05 U/kg/hr can be considered if pH is > 7.15.

Potassium replacement is given only when potassium is <5.5 mmol/L. In rare cases of hypokalemia with potassium <3.0 mmol/L, defer insulin treatment.

Bicarbonate administration is not recommended except for treatment of severe acidosis (pH < 6.9) or in cases of life-threatening hyperkalemia.



Solution to Question 17:

In the given clinical scenario, a 9-month-old child with normal breathing (> 50/min RR for infants 2 months to 12 months is considered as fast breathing) and no chest indrawing is seen. So, the next step in management would be to explain danger signs and send them home.

According to the IMNCI, three key clinical signs are used to assess a sick child with suspected pneumonia:

• Respiratory rate (RR): distinguishes children who have pneumonia from those who do not. Children are classified under pneumonia if:

• > 60 breaths per minute if age is < 2 months

• > 50/min RR for infants 2 months to 12 months

• > 40/min RR for children 12 months to 5 years of age

• Lower chest wall indrawing refers to the indrawing of the lower chest wall with respect to the child's breathing and indicates pneumonia

• Stridor indicates those with severe pneumonia requiring hospital admission.

The revised classification of acute respiratory infections is simplified to include only two categories of pneumonia:

• Cough and cold, no pneumonia - requires home remedy

• Pneumonia - with fast breathing and/or chest indrawing, which requires home therapy with oral amoxicillin.

• Severe pneumonia - pneumonia with any general danger sign,
which requires a referral and injectable therapy.

Danger signs of severe pneumonia are:

• Not able to drink

- Persistent vomiting
- Convulsions
- Lethargic or unconscious
- Stridor in a calm child
- Severe malnutrition



† Not able to drink, persistent vomiting, convulsions, lethargic or unconscious, stridor in a calm child or severe malnutrition.

Paediatrics INI-CET 2024

Question 1:

Which of the following is not seen in Kawasaki disease?

a) Purulent conjunctivitis

b) Strawberry tongue

c) Rash

d) Cervical lymphadenopathy

Question 2:

What is the ideal maintenance fluid of choice in children?

a) 5% dextrose and N/2 saline

b) 5% dextrose and N saline

c) 10% dextrose and N/2 saline

d) 5% dextrose and N/4 saline

Question 3:

Which of the following is not seen in nutritional rickets?

a) Low phosphorus

b) Normal calcium

c) Urine calcium: creatinine ratio > 15

d) High PTH

Question 4:

Which of the following is CORRECT regarding febrile seizure

a) A,B,C

b) A,C,D

c) A,B,C,D

d) A,B,D

Question 5:

What is the rate of increase in head circumference in the first 3 months of life?

a) 1 cm/month

b) 2 cm/month

c) 4 cm/month

d) 0.5 cm/month

Question 6:

Which of the following holds true for "some dehydration" in a child:

a) 1, 3, 5 only

b) 1, 2, 4 only

c) 1, 3, 4, 5 only

d) 1, 2, 3, 4

Question 7:

Which of the following is not a suitable method for free flow oxygen delivery to a newborn baby?

a) Oxygen tube kept close to the baby's nose and mouth

b) Oxygen mask placed firmly over the baby's mouth and nose

c) Self-inflating bag attached to an oxygen mask placed over baby's mouth and nose

d) A reservoir bag attached to oxygen mask kept over the baby's nose and mouth

Question 8:

All are true about renal tubular acidosis in children except

a) Type 4 is due to defective ammoniagenesis and aldosterone resistance

b) Type 2 is Fanconi and hypophosphatemic rickets

c) Type 4 with paradoxical aciduria and hypokalemia

d) Type 2 - abnormal loss of HCO_3

Question 9:

Which of the following is a reliable sign of "some dehydration" in severe acute malnutrition?

a) Increased thirst

b) Sunken eyes

c) Skin pinch that goes back slowly

d) Altered sensorium

Question 10:

A 5-year-old has the following anthropometry values. According to WHO criteria for malnutrition, what is the diagnosis?

a) Severe acute malnutrition

b) Chronic malnutrition

c) Moderate acute malnutrition

d) Acute on chronic malnutrition

Question 11:

Downe's score includes which all of the following?

a) 1,2,3

b) 2,3

c) 1,2,4

d) 2, 4

Question 12:

A 15 year old boy presents with short stature. On examination, his upper segment/lower segment ratio is 0.8:1. Which of the following is the likely diagnosis?

a) Spondyloepiphyseal dysplasia

b) Achondroplasia

c) Growth hormone deficiency

d) Hypothyroidism

Question 13:

Which of the following MRI findings will be seen in a child with spastic quadriplegia?

a) Periventricular leukomalacia

b) Multicystic encephalomalacia

c) Cerebellar degeneration

d) Basal ganglia calcification

Question 14:

A 10 year boy presents to you with sexual maturity rating 5, height for age > chronological age, hyperpigmentation and short stature. His blood pressure measured 180/110mmHg. Which of the following enzymes deficiency is responsible for this condition?

a) 11 beta hydroxylase

b) 21 alpha hydroxylase

c) 3 beta dehydrogenase

d) Cholesterol side chain cleavage enzyme

Question 15:

INSURE is associated with which of the following conditions?

a) Respiratory distress in new born

b) Birth Asphyxia

c) Neonatal jaundice

d) Transient tachypnea of newborn

Question 16:

All of the following are included under diagnostic criteria of severe acute malnutrition except

a) Height for age

b) Weight for height

c) Pedal edema

d) MUAC

Question 17:

In children diagnosed with congenital heart disease, NADA's criteria include all the following, except?

a) Diastolic murmur

b) Increased troponin-T

c) Abnormal X-ray

d) Congestive cardiac failure

Question 18:

Which of the following is false regarding neonatal cholestasis?

a) Kasai procedure should be done within 90 days

b) HIDA scan is an investigation of choice

c) Biliary atresia is the most common cause

d) Liver biopsy rules out other causes

Question 19:

What causes the closure of ductus arteriosus?

a) Increase in Oxygen

b) Decrease in CO_2

c) Increase in pulmonary vascular resistance

d) Fall in concentration of catecholamines

Question 20:

A 4-year-old child was diagnosed with croup. Which of the following is the most common causative organism?

a) Hemophilus influenza

b) Streptococcus pneumonia

c) Influenza virus

d) Diphtheria

Question 21:

What would be the age of a child who copies crosses, tell stories, and goes to the toilet alone?

a) 3 years

b) 4 years

c) 5 years

d) 2 years

Question 22:

Which of the following is not true about breastmilk jaundice?

a) Breastfeeding need not be stopped for the treatment

b) Most of the children require phototherapy

c) Exclusive breastfeeding with a total bilirubin of more than 10mg/dl after 3-4 weeks of birth

d) Unconjugated hyperbilirubinemia

Question 23:

Which of the following is not a feature of cystic fibrosis?

a) Nasal polyposis

b) Biliary atresia

c) Meconium ileus

d) Bleeding diathesis

Question 24:

A 2-week-old infant presents with a history of epistaxis and recurrent rectal bleeding. The parents also reported a history of continuous bleeding from the umbilicus since birth. Laboratory investigations revealed normal prothrombin time, normal thrombin time, normal

partial thromboplastin time, normal platelet count, and normal function. The urea clot solubility test was positive. What is the most likely deficient factor in this patient?

a) Factor XIII

b) Factor XI

c) Factor X

d) Factor XII

Question 25:

Which of the following indicates the primary failure of recovery in an admitted patient with severe acute malnutrition?

a) i,ii,iii

b) i,ii,iv

c) i,iii,iv

d) ii,iii,iv

Answer Key

Question No.	Correct Option
1	a
2	b
3	c
4	b
5	b
6	c
7	c
8	c
9	a
10	b
11	c
12	a
13	b
14	a

15	a
16	a
17	b
18	a
19	a
20	c
21	b
22	b
23	b
24	a
25	a

Detailed Explanations

Solution to Question 1:

Purulent conjunctivitis is not seen in Kawasaki's disease. Non-purulent conjunctivitis is a feature seen.

Kawasaki disease is also known as mucocutaneous lymph node syndrome. It is a vasculitis condition of unknown etiology. Clinical features include a high spiking fever that is unresponsive to antipyretics. The fever can last between 5 days to 3-4 weeks. The five principal clinical criteria for Kawasaki's disease are:

• Bilateral non-purulent conjunctival infection with limbal sparing

• Erythema of the oral and pharyngeal mucosa, strawberry tongue (option B), and red, cracked lips

• Edema and erythema of the hands and feet

• Various forms of rash- maculopapular, erythema multiforme, scarlatiniform (option C)

• Nonsuppurative cervical lymphadenopathy (option D)

The image below shows bilateral non-purulent conjunctival redness, strawberry tongue, and edema and erythema of the hands:

Diagnosis requires fever and at least four of the five principal criteria. Myocarditis develops frequently in acute Kawasaki disease. It presents as tachycardia and reduced ventricular systolic function. The tachycardia will be disproportionate to the fever. Patients may develop coronary artery aneurysms in the 2nd to 3rd week of illness. They are generally asymptomatic and diagnosed on echocardiography. However, if large it can lead to death. Thus, echocardiography should be performed at diagnosis and after 1-2 weeks.

Treatment involves 2g/kg of intravenous immunoglobulin as an infusion over 10-12 hours within 10 days of onset. Moderate to high-dose aspirin should be administered until the patient is afebrile.

Solution to Question 2:

5% dextrose and N saline is the ideal maintenance fluid in children.

5% dextrose provides 17 calories/100 mL and nearly 20% of the daily caloric needs. This level is enough to prevent ketone production and helps minimise protein degradation. However, the child will lose weight on this regimen and therefore it is necessary to start enteral feeds sooner or if not possible, total parenteral nutrition is recommended.

Other options:

Hypotonic fluids like N/2 or N/4 saline can increase the risk of hyponatraemia and free water movement inside RBCs, leading to haemolysis. Hence, isotonic fluids with 5% dextrose are recommended as standard maintenance fluids in children.

Solution to Question 3:

Urine calcium: creatinine ratio > 15 is not seen in a case of nutritional rickets. Instead low urine calcium and high urinary phosphate excretion are seen due to stimulation of PTH(parathormone)

in vitamin D deficiency(nutritional rickets).(Option C)

Initially, low calcium levels in the blood stimulate increased PTH secretion from the parathyroid glands that results in bone resorption and increased reabsorption of calcium and increased excretion of phosphorus in the kidneys.

Hence blood calcium levels can be low to normal with low phosphorus levels and high ALP due to increased bone resorption activity.

Note - High urinary calcium excretion is associated with hereditary hypophosphatemic rickets with hypercalciuria.

Nutritional rickets is a skeletal disorder primarily caused by a deficiency in vitamin D, calcium, or phosphorus, or a combination of these nutrients. Rickets typically affect children during periods of rapid growth and bone development.

25-hydroxy vitamin D level should be measured to diagnose vitamin D deficiency rickets. The initial lab tests in this child should also include serum calcium, phosphorous, alkaline phosphatase (ALP), and parathyroid hormone (PTH). Lab findings reveal decreased calcium and phosphate levels, increased PTH, and ALP.

Treatment with oral vitamin D is preferred. Patients who show no response should be evaluated for other disorders, such as hypophosphatemic rickets, vitamin D-dependent rickets, renal tubular acidosis-associated rickets, and chronic kidney disease-associated rickets.

Solution to Question 4:

Chance of recurrence after one episode of febrile seizure is not 54%

Febrile seizures are seizures that-

• Occur between the age of 6 months and 5 years

• Occur at a temperature of 38°C (100.4°F) or higher

• Are not due to any CNS infection or any metabolic imbalance

• Occur without a history of prior afebrile seizures

• Occur with no post-ictal neurological deficit.

Types of febrile seizures:

Investigations:

• Lumbar puncture should be performed in the first episode of seizures with fever, for all infants <6 months age (to exclude meningitis).

• EEG and neuroimaging are advised for children with atypical febrile seizures.

Treatment:

Antipyretics and proper hydration. Midazolam or diazepam (rectal) can be used if seizures present.

Prophylaxis:

• Intermittent prophylaxis: If the febrile seizures are 3 or more in 6 months, or 4 or more seizures in 1 year, oral clobazam (1mg/kg/day) for 3 days during the fever episode.

• Continuous prophylaxis: If intermittent prophylaxis fails or in recurrent atypical febrile seizures, sodium valproate and phenobarbitone can be used.

Simple febrile seizures	Complex (atypical) febrile seizures
Occurs within 24 hours of the onset of fever	Can occur at any time during a fever
Generalized seizures	Focal seizures
Duration is <15 min	Prolonged
No recurrence within 24 hrs of seizures	Recurrent
Chances of going into epilepsy 1%	Chances of going into epilepsy 6%

Solution to Question 5:

Rate of increase of head circumference in the first 3 months of life is 2 cm/month.

The maximum circumference of the head from the occipital protuberance to the supraorbital ridges on the forehead is recorded using the cross-tape method.

Head circumference increases at the following rates:

• 2 cm/month in first 3 months

• 1 cm/month in the next 3 months

• 0.5 cm/month in the next 6 months

Head circumference is equal to chest circumference at 1 year of age.

Head circumference is more than chest circumference by 2.5 cm at birth.

• At birth: Head circumference > Chest circumference

• At 9-12 months: Head circumference = Chest circumference

• After 1 year: Head circumference < Chest circumference

Age	Head circumference
At birth	33 to 35cm
3 m	40 cm
6 m	43 cm
1 year	46 cm

Age	Head circumference
2 years	48 cm
3 years	49 cm
4 years	50 cm
12 years	52 cm (adult size)

Solution to Question 6:

Increased thirst, delayed skin pinch, sunken eyes, and reduced urine output indicate some dehydration. Lethargy indicates severe dehydration.

The degree of dehydration is assessed as follows:

Based on the degree of dehydration after history and examination, the estimated fluid loss is calculated as follows:

Parameters	No dehydration	Some dehydration	Severe dehydration
Condition	Well alert	Restless, irritable	Lethargic or unconscious,floppy
Eyes	Normal	Sunken	Very sunken and dry
Tears	Present	Absent	Absent
Mouth and tongue	Moist	Dry	Very dry
Thirst	Drinks normallynot thirsty	Thirsty, drinks eagerly	Drinks poorly or isnot able to drink
Skin pinch	Goes back quickly	Goes back slowly	Goes back very slowly
Urine output	Normal	Decreased	Oliguria, anuria
Capillary refill	2-3 seconds	3-4 seconds	>4 seconds
Treatment	Plan A	Weigh the patient, if possible, and use treatment Plan B	Weigh the patient and use treatmentPlan C urgently

Degree of dehydration	Assessment of fluid loss
No dehydration	less than 50 mL/kg
Some dehydration	50-100 mL/kg
Severe dehydration	more than 100 mL/kg

Solution to Question 7:

Self-inflating bag is not used for free flow oxygen delivery. It is used to provide positive pressure ventilation. Self-inflating bag and mask is used in neonatal resuscitation.

Bag and mask ventilation is used as the first step in airway management to provide positive pressure ventilation. It serves as a rescue technique to provide oxygenation and ventilation in patients with difficult airways until tracheal intubation is achieved.

The bag in this apparatus is a self-inflating bag. This allows for it to be used without a pressurized gas supply and is delivered to the patient when the bag is squeezed (compressed). A patent airway and a good seal are essential to facilitate bag and mask ventilation.

The rescuer should be positioned at the patient's head. A one-person technique involves the "E-C seal" in which your first and second digits form a "C" over the mask with your thumb pressing down by the nasal bridge, your second digit over the bottom of the mask by the mouth, and your third through fifth digits forming an "E" and applying pressure to the mandible to hold the mask tight

Effective ventilation can be confirmed by:

• Chest rise on squeezing the bag

• Condensation in the clear mask

• Pulse oximetry shows improved saturation

• Capnography shows exhaled carbon dioxide

Mask ventilation is usually contraindicated in patients with an increased risk of regurgitation. It should be used with caution in patients with severe facial trauma.

The following image shows the parts of a self-inflating bag and mask apparatus with a reservoir bag.

Solution to Question 8:

Hyperkalemia and metabolic acidosis are features of type 4 renal tubular acidosis

Type IV (hyperkalemic) renal tubular acidosis: Hyperkalemia with distal RTA occurring due to aldosterone resistance or deficiency is termed type 4 RT A. Aldosterone directly stimulates the proton pump, increases Na+ absorption resulting in negative intrahibular potential and increases urinary K+ losses and stimulates basolateral Na+ /K+ ATPase. Hence, aldosterone deficiency or resistance is expected to cause hyperkalemia and acidosis.

The clinical features seen in Type IV RTA are as follows:

• Hyperkalemia: Hypoaldosteronism results in an increase in potassium concentration.

• Metabolic acidosis: The important buffers in urine are ammonia and phosphate. Hyperkalemia impairs ammonia production in the proximal tubule, which reduces the buffering of hydrogen ions. This results in decreased hydrogen ion excretion.

• Urinary pH<5.5: Due to the decreased production of ammonia in the proximal tubule, excess protons are excreted in the renal tubules, thus decreasing the urinary pH.

RTA type I (distal RTA) is seen in autoimmune diseases like Sjogren's syndrome, systemic lupus erythematosus, systemic sclerosis, medullary sponge kidney disease, cirrhosis, drugs like amphotericin B, lithium, etc. The patients develop hypokalemia and metabolic acidosis with alkaline urine (urinary pH >5.5).

RTA type II (proximal RTA) is seen in Fanconi syndrome, Wilson's disease, heavy metals ingestion, etc., The patients develop hypokalemia and metabolic acidosis with acidic urine (urinary pH <5.5). It occurs due to generalized proximal tubule dysfunction. Defective HCO_3^- resorption is the primary dysfunctional mechanism causing metabolic acidosis and is also associated with glycosuria, aminoaciduria, and phosphaturia. Rickets is rare in isolated proximal RTA but common in Fanconi syndrome.

RTA type III (mixed RTA) is seen in inherited mutations of carbonic anhydrase II. A combined pathology of type I and type II RTA is seen in this type. It results in hypokalemia and metabolic acidosis.

Solution to Question 9:

Increased thirst is the best indicator of 'some dehydration' in severe acute malnutrition.

Increased thirst is the most reliable indicator because impaired skin pinch and sunken eyes can also be seen due to loss of subcutaneous fat in a child with acute malnutrition. Altered sensorium is a feature of severe dehydration.

The degree of dehydration is assessed as follows:

Based on the degree of dehydration after history and examination, the estimated fluid loss is calculated as follows:

Parameters	No dehydration	Some dehydration	Severe dehydration
Condition	Well alert	Restless, irritable	Lethargic orunconscious,flop py
Eyes	Normal	Sunken	Very sunken and dry
Tears	Present	Absent	Absent
Mouth and ton gue	Moist	Dry	Very dry
Thirst	Drinks normallynot thirst y	Thirsty, drinks eagerly	Drinks poorly or isnot able to drink
Skin pinch	Goes back quickly	Goes back slowly	Goes back very slowly
Urine output	Normal	Decreased	Oliguria, anuria
Capillary refill	2-3 seconds	3-4 seconds	>4 seconds
Treatment	Plan A	Weigh the patient, if possible, and use treatment Plan B	Weigh the patient and use tre atmentPlan C urgently

Degree of dehydrati on	Assessment of fluid loss
No dehydration	less than 50 mL/kg
Some dehydration	50-100 mL/kg
Severe dehydration	more than 100 mL/kg

Solution to Question 10:

According to WHO criteria for malnutrition, the diagnosis of this child is chronic malnutrition. (Option B)

Height for age less than -2 SD (-2.5 in this case) indicates moderate stunting, pointing towards a diagnosis of chronic malnutrition.

Weight for age less than -2 SD (-3.2 in this case) indicates underweight but it cannot differentiate between wasting and stunting.

Weight for height:

• less than - 3SD suggests severe acute malnutrition (SAM)

• between -2 to -3 SD suggests moderate acute malnutrition

Weight for height in this case is -1.7 SD. Hence it is not a case of SAM or moderate acute malnutrition.

Solution to Question 11:

Respiratory rate, cyanosis and grunting are a part of Downe's score. Feeding is not a part of it.

Downes' Score is a comprehensive scoring system and can be applied to any gestational age and condition. Scoring is to be done every half an hour and charting is carried out to monitor progress.

Downe's score of >6 means impending respiratory failure.

Score	0	1	2
Respiratory rate	<60/min	60–80/min	>80/min
Air entry	Normal	Mild decrease	Marked decrease
Cyanosis	Nil	In-room air	On >40% O2
Grunting	None	Only on auscultation	Audible with the naked ear
Retraction	Nil	Mild	Moderate

Solution to Question 12:

Upper segment to lower segment ratio (U/L) less than 1 indicates spondyloepiphyseal dysplasia.

The upper segment to lower segment (U/L) ratio in this child is 0.8. In normal development, the U/L ratio is 1 in children more than 7 years.

The lower segment(L) is defined as the length from the symphysis pubis to the floor. The upper segment(U) is measured by subtracting the lower segment from the height.

The U/L ratio is

• 1.7 at birth

• 1.3 at 3 years

• 1 after 7 years

• 0.9 in adults



High U/L ratio:

• Rickets

• Achondroplasia (short limb dwarfism)

• Turner syndrome

• Untreated congenital hypothyroidism

Low U/L ratio:

• Spondyloepiphyseal dysplasia

• Marfan syndrome

• Morquio syndrome

• Vertebral anomalies

Solution to Question 13:

Multicystic encephalomalacia is the MRI finding seen in children with spastic quadriplegia.

Cerebral palsy is classified into several types based on its clinical features.

Spastic quadriplegia is the most common subtype seen in India. It is often caused by perinatal asphyxia and neonatal illness where all four limbs are affected. MRI would reveal multicystic encephalomalacia. It is associated with a poor outcome.

Spastic diplegia is commonly seen in premature infants with a history of birth asphyxia. It is second most common type of cerebral palsy in India after spastic quadriplegia. Both the lower limbs are involved, with spasticity in the hip adductors. This leads to a characteristic commando crawling in infants and a scissoring gait in older children. Brain MRI shows scarring and shrinkage in the periventricular white matter with compensatory enlargement of the cerebral ventricles, called periventricular leukomalacia (option A). IQ of these children is usually normal

and has a good prognosis.

Spastic hemiplegia affects one-half of the body. It is seen due to a vascular insult leading to a perinatal stroke event. Early hand preference is seen (before 1 year), where one side of the body is always preferred as the other side is weak. MRI shows focal changes and porencephalic cysts due to infarcts.

Dyskinetic/ Extrapyramidal palsy is associated with birth asphyxia and kernicterus in the early neonatal period. Unlike spastic diplegia, the upper limbs are affected more than the lower limb. Brain MRI reveals lesions in the basal ganglia and thalamus (option D).

Cerebellar degenerations are associated with ataxic cerebral palsy (option C).

Note: Periventricular leukomalacia can also be seen in spastic quadriplegia but multicystic encephalomalacia is more common.

Solution to Question 14:

11 beta hydroxylase deficiency is the enzyme deficiency responsible for this condition. It is one of the variants of congenital adrenal hyperplasia(CAH) where there is increased aldosterone and testosterone activity with reduced cortisol levels.

The given clinical scenario of precocious puberty(SMR grading of 5 in a 10 year old boy) with height age> chronological age with short stature and increased blood pressure 180/110 suggests 11 beta hydroxylase deficient variant of CAH.

Solution to Question 15:

INSURE is associated with respiratory distress in newborns.

INSURE is abbreviated as intubate-intratracheal surfactant administration-extubation. It is a treatment modality performed in neonates with respiratory distress especially in preterm where surfactant production is reduced. It is indicated in neonates who are not maintaining oxygen saturation >90% while breathing 40-70% FiO2 in nasal CPAP(continuous positive airway pressure).

Extubation back to nasal CPAP is done immediately once the infant is stable, usually within minutes to <1 hr of surfactant administration. The amount of nasal CPAP required usually decreases after approximately 72 hr of age and most infants can be weaned from nasal CPAP.

Variations of the INSURE method are MIST (minimally invasive surfactant therapy) and LISA (less invasive surfactant administration), in which a small feeding tube, rather than an endotracheal tube (ETT), is used to deliver intratracheal surfactant to a spontaneously breathing infant on nasal CPAP.

The combination of early rescue surfactant by these methods along with nasal CPAP has been associated with the reduced need for mechanical ventilation and prevention of bronchopulmonary dysplasia.

Solution to Question 16:

Height for age is not a diagnostic criterion of severe acute malnutrition.

Criteria for severe acute malnutrition (SAM) in a 6-month-old child are mid-upper arm circumference, symmetrical pedal edema , and weight for height. Height for age is not the criteria for the diagnosis of severe acute malnutrition.

The presence of any of the following confers severe acute malnutrition in children 6 to 59 months of age:

• Weight-for-height below -3 standard deviation (<-3SD) on the WHO Growth Standard (Option B)

• Presence of bipedal edema (Option C)

• Mid-upper arm circumference (MUAC) below 11.5 cm (Option D)

Once the child is diagnosed with severe acute malnutrition, the doctor must assess for the following complications:

• Severe edema (+++)

• Low appetite (failed appetite test)

• Medical complications (severe anemia, pneumonia, diarrhea, dehydration, cerebral palsy, tuberculosis, HIV, heart disease, etc.)

One or more danger signs as per IMNCI

If any of the complications are present, the child is diagnosed with complicated SAM and is managed inpatient in the facility.

If the above-mentioned signs are absent, the child is diagnosed with uncomplicated SAM and supervised management at home is given.

Solution to Question 17:

Increased troponin-T is not included in NADA's criteria.

Nada's criteria is used to diagnose congenital heart disease in children. The presence of 1 major or 2 minor criteria indicates congenital heart disease.

Major criteria	Minor criteria
Systolic murmur grade 3 or moreAny diastolic murmur (Option A)CyanosisCongestive cardiac failure (Option D)	Systolic murmur < grade 3Ab normal second sound (S2)Ab normal ECGAbnormal chest X-ray findings (Option C)Abn ormal BP

Solution to Question 18:

Kasai procedure should be done within 60 days of age, and not 90 days. Delay beyond 60 days significantly reduces success rates and increases the risk of requiring liver transplantation.

Neonatal cholestasis refers to prolonged jaundice in newborns caused by impaired bile flow.

Option B: HIDA scan is the investigation of choice. The hepatobiliary iminodiacetic acid (HIDA) scan is used to evaluate bile flow from the liver into the intestine and can help differentiate biliary atresia from other causes of neonatal cholestasis.

Option C: Biliary atresia is the most common cause of neonatal cholestasis, this condition requires surgical intervention.

Option D: Liver biopsy is a key diagnostic tool in neonatal cholestasis as it can confirm biliary atresia and rule out other causes like metabolic diseases or neonatal hepatitis.

Solution to Question 19:

An increase in oxygen causes the closure of ductus arteriosus.

Mechanism of closure of the ductus arteriosus:

1. Functional Closure (Minutes to Hours): Increased oxygen tension (PaO■) after birth causes smooth muscle contraction in the ductus. Decreased prostaglandin E2 (PGE2) levels (due to loss of placental production and increased pulmonary clearance) also contribute.

2. Anatomical Closure (Days to Weeks): Smooth muscle proliferation and fibrosis lead to permanent closure, forming the ligamentum arteriosum.

Thus, oxygen, reduced PGE2, and smooth muscle constriction drive ductus closure.

Changes in vascular resistance at birth:

Pulmonary vascular resistance (PVR): Before birth, the lungs are fluid-filled, and PVR is high. At birth, with the first breath, the lungs expand, oxygen tension rises, and pulmonary blood flow increases. This causes a decrease in PVR.

Systemic vascular resistance (SVR): At birth, clamping of the umbilical cord eliminates the low-resistance placental circulation. This leads to an increase in SVR.

Solution to Question 20:

Among the options, the Influenza virus is the most common causative organism of croup.

Croup, or laryngotracheobronchitis, is a viral infection affecting the upper airway, primarily in children aged 6 months to 6 years. It causes inflammation of the larynx, trachea, and bronchi, leading to characteristic symptoms such as:

• Barking cough (seal-like cough)

• Inspiratory stridor

• Hoarseness

• Respiratory distress in severe cases

The most common causative organism for croup is the parainfluenza virus. While other viruses like influenza, respiratory syncytial virus (RSV), and adenovirus can also cause croup, parainfluenza is the predominant cause.

Management:

• Racemic epinephrine nebulization leads to the rapid improvement of upper airway obstruction. Epinephrine constricts pre-capillary arterioles in the upper airway mucosa and decreases capillary hydrostatic pressure, leading to fluid resorption and improvement in airway edema.

• Dexamethasone-by the least invasive route (Oral>IV>IM). It is indicated in all the grades of croup.

• Oxygen should be administered to children who are hypoxemic (SpO2 <92 %)

Antibiotics are not indicated in laryngotracheobronchitis.

Other options:

Option A: Hemophilus influenza is associated with epiglottitis.

Option B: Streptococcus pneumonia is associated with pneumonia, otitis media, or meningitis, not croup.

Option D: Diphtheria is associated with pseudomembrane in the throat and airway obstruction

Solution to Question 21:

The combination of copying a cross, telling stories, and independent toileting aligns with the developmental milestones of a 4-year-old child.

Developmental milestones

At 1 year

• Motor: Walks with support

• Adaptive: Mature pincer grasp

• Language: Speaks a few words

• Social: Plays simple ball game

• Domestic mimicry at 18 months

At 2 years (Option D)

• Motor: Walks up and downstairs, one step at a time

• Adaptive: Makes a tower of 6 blocks

• Language: Simple sentences, a vocabulary of 50-100 words, uses pronouns at 2.5 yrs

• Social: Asks for food and starts toilet training, listens to stories

At 3 years (Option A)

• Motor: Rides tricycle

• Adaptive: Makes a tower of 9 blocks, handedness is established

• Language: Knows full name, age, gender

• Social: Plays simple games and engages in parallel play with other children

At 4 years

• Motor: Hops

• Adaptive: Draws square and constructs a bridge with blocks

• Language: Tells stories

• Social: Goes to the toilet alone

At 5 years (Option C)

• Motor: Skips

• Adaptive: Draws a triangle

• Language: Asks the meaning of words, names 4 colors

• Social: Dresses and undresses

Solution to Question 22:

Most of the children suffering from breastmilk jaundice do not require phototherapy.

Breastmilk jaundice is due to significant unconjugated hyperbilirubinemia (Option D), after the 7th day and may last for 2–3 weeks. This may be caused by the breast milk that might contain β-glucuronidase, which deconjugates intestinal bilirubin and promotes its absorption and other inhibitors of glucuronyl transferase. Persistently high bilirubin levels may require changing from

breast milk to infant formula for 24–48 h and/or treatment with phototherapy, without cessation of breastfeeding. (Option A)

Most cases of breast milk jaundice are mild and self-limiting, and phototherapy is required only in rare severe cases where bilirubin levels are significantly elevated. (Option B)

Breast milk jaundice is typically observed in exclusively breastfed infants, often with elevated bilirubin (>10mg/dl) levels persisting beyond the neonatal period (after 3-4 weeks of birth). (Option C)

Solution to Question 23:

Biliary atresia is not a feature of cystic fibrosis.

Cystic Fibrosis (CF) is a genetic disorder that primarily affects the lungs, digestive system, and other organs. It is caused by mutations in the CFTR gene, which leads to the production of thick and sticky mucus. The CFTR gene is on chromosome 7q. The most common CFTR mutation is F508 deletion. This mucus blocks ducts and passages in various organs, causing problems.

Features of CF:

• Respiratory Symptoms: Chronic cough, recurrent lung infections, bronchiectasis, and nasal polyps. (Option A)

• Digestive Issues: Pancreatic insufficiency, malabsorption, meconium ileus (Option C), and fat-soluble vitamin deficiencies which causes bleeding diathesis (Option D).

• Sweat Gland Abnormalities: Elevated sweat chloride levels.

• Reproductive: Infertility in males (due to absent vas deferens) and reduced fertility in females.

• Other Complications: CF-related diabetes (CFRD), osteoporosis, and liver disease.



The diagnosis is confirmed if both of the following criteria are met:

1. Presence of

• Clinical symptoms affecting at least one organ system (respiratory, gastrointestinal, or genitourinary) or

• Positive newborn screening or

• A sibling with cystic fibrosis.

2. Evidence of dysfunction of the CFTR gene by

• Two positive sweat chloride test results were obtained on separate days or

• Identification of two CFTR mutations by genetic testing or

• An abnormal measurement of nasal potential difference.

Cystic fibrosis requires a multidisciplinary approach to management. The correction of the defect can be done via:

• Lumacaftor, corrects the misfolding and reverts the mutation defect.

• Ivacaftor potentiates CFTR channel opening and stimulates ion transport.

End-stage pulmonary failure requires lung transplantation.

Solution to Question 24:

The infant's history of umbilical bleeding, recurrent bleeding episodes, normal coagulation studies, and positive urea clot solubility test strongly suggest Factor XIII deficiency.

FXIII cross-links fibrin polymers and strengthens the clot, making it resistant to premature breakdown. Its deficiency leads to delayed wound healing, spontaneous bleeding, and in severe cases, life-threatening hemorrhages.

Presents with recurrent umbilical bleeding since birth and Epistaxis and rectal bleeding

Laboratory Findings: Normal PT, PTT, TT, platelet count and function: These indicate that the coagulation cascade and platelets are functioning normally up to fibrin formation. Positive urea clot solubility test: This is a classic diagnostic test for Factor XIII deficiency. Clots deficient in Factor XIII dissolve in 5M urea because they lack cross-linked fibrin

The preferred treatment is FXIII concentrates. Fresh frozen plasma (FFP) and cryoprecipitate are alternatives. Antifibrinolytics like tranexamic acid can be adjunctive for minor bleeding. Regular monitoring and prophylaxis are crucial for long-term management and preventing complications.

Other Options:

Option B: Factor XI: Deficiency typically causes mild to moderate bleeding, mainly in mucosal sites, and does not present with umbilical bleeding.

PT and PTT may be prolonged, which is not the case here.

Option C: Factor X: Deficiency causes severe bleeding and affects PT and PTT (both would be prolonged), unlike in this case where all tests are normal.

Option D: Factor XII: Deficiency does not cause bleeding; it is often asymptomatic. It can lead to prolonged PTT, which is not observed here.

Solution to Question 25:

Failure to regain appetite by day 4, failure to start to loose edema by day 4 and presence of edema by day 10 indicate primary failure of recovery in admitted patient with severe acute malnutrition.

For assessing primary failure of recovery in a patient with severe acute malnutrition (SAM), key clinical benchmarks guide the evaluation.

Difference between primary and secondary failure of malnutrition treatment:

Inpatient cases

Day of admission is considered as day 0.

Primary Failure	Secondary failure
Failure to regain appetite by day 4 (i)Failure to start losing edema by day 4 (ii)Presence of edema on day 10 (iii)Failure to gain at least 5 g/kg/day by day 10	Failure to gain 5 g/kg/day for 3 consecutive days during rehabilitationReappearance of edema after 4 weeks

Paediatrics NEET 2024

Question 1:

Three female children graphs were plotted on an IAP growth chart, which of them signals growth failure?



a) Only 2 and 3

b) Only 3

c) 1, 2 and 3

d) Only 1

Question 2:

A neonate has bilateral chorioretinitis with intraparenchymal calcifications. She also has hydrocephalus. What is the causative organism?

a) Trypanosoma

b) Toxoplasma gondi

c) Taenia solium

d) Cytomegalovirus

Question 3:

What is represented by each of the following growth curves?

a) 1 - Liver, 2 - Spleen, 3- Ovary, 4 - Testis

b) 1 - Brain, 2- Tonsils, 3 - BMI, 4 - Genital Organs

c) 1 - Brain, 2 - somatic growth, 3- Ovary, 4 - Height

d) 1- Brain, 2- Waldeyer's ring, 3- somatic growth, 4- Testicular growth

Question 4:

A mother brought her 2-year-old child to the OPD with complaints of lack of weight gain, abdominal pain on eating, and poor eating habits. The child was advised gluten-free diet after which there was improvement in the general condition and weight gain. What is the probable diagnosis?



a) Topical sprue

b) Celiac disease

c) Whipple's disease

d) Intestinal malabsorption

Question 5:

Identify the condition shown in the image:



a) Chiari type 1

b) Dandy-Walker malformation

c) Vein of Galen malformation

d) Corpus callosum agenesis

Question 6:

A newborn baby presented with a difficulty in breathing. Auscultation of the chest revealed gurgling sounds and heart sounds shifted to the right. The chest radiograph confirmed a congenital diaphragmatic hernia. Which among the following is the most common location for this condition?

a) Esophageal opening

b) Left posterolateral

c) Between sternum and coastal cartlage

d) Central tendon

Question 7:

A child presented with failure to thrive and recurrent infections. On examination, tonsils are absent. Further evaluation confirms the deficiency of adenosine deaminase. What is the likely diagnosis?

a) Lesch-Nyhan syndrome

b) Ataxia telengeictasia

c) Severe Combined Immuno Deficiency

d) Friedrich's ataxia

Question 8:

A child presented with diarrhea and dehydration. There was no blood in the stools. A stool examination revealed a spoke wheel pattern. Identify the causative organism.



a) Adenovirus

b) E.coli

c) Norwalk virus

d) Rota virus

Answer Key

Question No.	Correct Option
1	b
2	b
3	d
4	b
5	a
6	b
7	c
8	d

Detailed Explanations

Solution to Question 1:

Out of the growth curves (weight-for-age), only growth curve 3 shows growth failure. This pattern shows fluctuation in growth percentiles, which might indicate a history of growth issues or inconsistent growth. The drop to below -2SD is significant and suggests growth failure as it indicates the child is significantly below the normal range.

The nutritional status of a child is assessed in terms of anthropometry:

The World Health Organization (WHO) has established international standards for normal child growth from birth to 5 years of age. The normalization of the anthropometric measures is done in terms of z-scores (standard deviation scores). The anthropometric indicators are marked in a growth chart which is compared with a reference curve.

The anthropometric measures used are weight for age, height for age, weight for height, BMI, and mid-upper arm circumference.

• Low weight for height, or wasting, indicates acute malnutrition. It is also mentioned as wasting.

• High weight-for-height indicates overweight.

• Low height-for-age, or stunting, indicates chronic malnutrition. This is the best indicator of long-term nutritional status in the pediatric population.

• Low weight-for-age indicates acute or chronic malnutrition. It does not differentiate between wasting and stunting. Hence, it has limited clinical significance.



The below image shows the WHO growth chart for girls (6 months-2years). Boys have similar charting in blue color.



Solution to Question 2:

The presence of bilateral chorioretinitis with intraparenchymal calcifications and hydrocephalus in a neonate is suggestive of congenital toxoplasmosis.

The characteristic triad of congenital toxoplasmosis is chorioretinitis, hydrocephalus, and diffuse intracranial calcifications. Clinical manifestations apart from the triad include hydrops fetalis, intrauterine growth retardation, prematurity, peripheral retinal scars, persistent jaundice, and mild thrombocytopenia.

Congenital toxoplasmosis occurs when a pregnant woman acquires the infection during gestation. The organism may cross to the fetus transplacentally or at the time of vaginal delivery. The severity of the disease in the neonate is greater if the woman gets infected early in her pregnancy. The causative agent is Toxoplasma gondii, an obligate intracellular protozoan.

The diagnosis is made by the detection of IgM anti-toxoplasma antibodies. IgG can cross the placenta is hence, cannot confirm infection in the infant. IgM immunosorbent agglutination assay (ISAGA) and IgA ELISA are the most useful diagnostic tests.

Spiramycin can be administered for cases of acute infection in early pregnancy to reduce vertical transmission. It does not cross the placenta and therefore cannot be used to treat an infected fetus. Pyrimethamine-sulfonamide (to be avoided in the first trimester) is used when a fetal infection is confirmed or in maternal infections after 18 weeks.

Other options:

Option A: Trypanosoma is a genus of protozoan parasites responsible for diseases like African trypanosomiasis (sleeping sickness) and American trypanosomiasis (Chagas disease). These diseases present with different clinical features, such as fever, swollen lymph nodes, and neurological symptoms in African trypanosomiasis, and fever, swelling at the infection site, and cardiomyopathy in Chagas disease.

Option C: Taenia solium is a parasitic tapeworm transmitted through undercooked pork, causing intestinal infection (taeniasis) or more severe cyst formation in tissues, including the brain (cysticercosis). It's prevalent in regions with poor sanitation and where pork is consumed.

Option D: Congenital cytomegalovirus presents with hepatosplenomegaly, petechial rashes, jaundice, chorioretinitis, and microcephaly. Head neuroimaging shows intracranial periventricular calcifications. One of the long-term sequelae is sensorineural hearing loss. Diagnosis is made by isolating the virus from the urine or blood by polymerase chain reaction (PCR)

Solution to Question 3:

The correct answer is 1- Brain, 2- Waldeyer's ring, 3- somatic growth, 4- Testicular growth.

Curve 1: The neural growth(brain) attains 90% of the growth by the age of 2 years.

Curve 2: The tonsils are collections of lymphatic tissue located within the pharynx. They collectively form a ringed arrangement, known as Waldeyer's ring. Tonsils/lymphatic growth in children occur between 4-8 years of age and then shrink down in adult age.

Curve 3: Somatic growth shows a sigmoid curve, rapid growth in the initial years, and then during puberty.

Curve 4: Testicular growth/Gonadal growth starts during puberty, before that the curve is flat.



Solution to Question 4:

The clinical presentation and the histopathological findings are suggestive of celiac disease.

Celiac disease is an autoimmune gluten-induced enteropathy where HLA DQ2 and DQ8 are expressed. In the small intestine, peptides of gluten are presented to helper T cells which mediate mucosal inflammation. Antiendomysial and anti-tTG antibodies are found in serological studies.

Clinical Features:

• Gastrointestinal Symptoms: Chronic diarrhoea, abdominal pain, vomiting, weight loss, and failure to thrive.

• Extraintestinal Symptoms: Anemia, fatigue, elevated liver enzymes, neurologic disorders, short stature, dental enamel defects, arthralgia, aphthous stomatitis.

Investigation:

• Serologic Tests: Initial testing for anti-TG2 IgA antibodies and total IgA. If anti-TG2 is positive and IgA levels are normal, CD is likely.

• Genetic Testing: For HLA-DQ2 or DQ8 haplotypes.

• Endoscopy and Biopsy: Multiple biopsies from the small intestine to check for villous atrophy.

• Biopsy Findings: villous atrophy: flattening of the villi in the small intestine, crypt hyperplasia: enlargement of the crypts, intraepithelial lymphocytosis: Increased number of lymphocytes in the epithelium.

Treatment:

Gluten-Free Diet: To manage celiac disease, patients must strictly avoid all foods containing gluten. Safe alternatives include rice, corn, maize, buckwheat, millet, amaranth, quinoa, sorghum, potato, soybean, tapioca, and teff, along with bean and nut flours.

Biopsy findings can be seen below:



Other options:

Option A: Tropical sprue is another malabsorption disorder seen as diarrhoea, steatorrhea, nutritional deficiencies, and weight loss. It is prevalent in some tropical areas hence the name. It is mostly attributed to infectious agents considering its response to antibiotics. It is diagnosed by a small intestinal biopsy which shows a close resemblance to that seen in celiac disease but it's not specifically triggered by gluten and is more common in adults.

Option C: Whipple Disease is a multisystemic disease caused by T.whipplei. Small intestinal biopsy shows PAS-positive macrophages.

Option D: Intestinal Malabsorption is a general term used for conditions that cause poor nutrient absorption, which could include celiac disease but is not specific enough to explain the clear response to a gluten-free diet.

Solution to Question 5:

The MRI image in the question shows a downward displacement of the cerebellar tonsils through the foramen magnum, which is characteristic of Chiari type 1 malformation.

This condition involves the herniation of the cerebellar tonsils into the spinal canal and is often associated with headaches, neck pain, and sometimes neurological deficits. It may be identified on imaging, such as MRI, where the cerebellar tonsillar descent is clearly visualized.

Arnold Chiari malformation is the most common malformation of the posterior fossa and hindbrain. There are two types of Chiari malformations-

• Chiari I malformation- It is the more common type. There is downward displacement of the cerebellar tonsils through the foramen magnum. It may be associated with syringomyelia.

• Chiari II malformation: Displacement of the medulla, the fourth ventricle, and cerebellar vermis through the foramen magnum. It is almost always associated with a myelomeningocele.

Other options:

Option B: Dandy walker malformation is a developmental anomaly characterized by a cystic dilation of the fourth ventricle, hypoplasia of the cerebellar vermis, an enlarged posterior fossa, and an upward displacement of the cerebellum. The atresia of the foramina of Magendie and Luschka leads to obstructed CSF flow, resulting in non-communicating hydrocephalus, causing symptoms such as vomiting, seizures, bulging fontanelles, and developmental delays. Diagnostic imaging includes ultrasound, CT, or MRI.

Given below is an image of Dandy walker malformation-



Option C: The MR angiography in the vein of Galen malformation shows an enlarged midline venous structure, with multiple arteriovenous communications leading to aneurysmal dilatation. The image below shows the malformation:



Option D: Agenesis of the corpus callosum is one of the most common congenital cerebral malformations which is morphologically the complete or partial absence of corpus callosum and not defined by functional or behavioral abnormalities. The given image shows the MRI findings:



Solution to Question 6:

Congenital diaphragmatic hernia with the features given in the clinical vignette occurs most commonly in a left posterolateral defect.

Bochdalek hernia is the most common congenital diaphragmatic hernia which occurs on the left posterolateral side due to a defect in the pleuroperitoneal membrane. The abdominal contents in the thorax compress the developing lungs causing pulmonary hypoplasia which manifests as respiratory distress in most infants. Chest X-ray shows an indistinct diaphragm with radiolucent para cardiac shadow and absent bowel loops in the abdomen. Opacities can be seen in the thorax if solid organs or omentum also herniate. Surgical repair is indicated in all cases either by a thoracic or abdominal approach.

Given below is an X-ray showing Bochdalek's hernia

Other options:

Option A- Sliding and rolling hernias occur via the oesophagal opening(hiatal hernia). Sliding hernia (more common) occurs when the gastroesophageal junction(GEJ) and a part of the stomach slide up into the thorax via the opening. A rolling hernia occurs when the GEJ stays below the diaphragm, but the stomach goes up, lying parallel to the oesophagus in the thorax. Hiatal hernias occur in adults.

Option C- The space between the sternum and costal cartilage is called the space of Larry or space of Morgagni. A defect in this space causes Morgagni hernia, which is the second most common hernia in children. It occurs on the right anteromedial aspect and is often less symptomatic.

The following is an X-ray showing Morgagni's Hernia :



Option D- A defect in the central tendon causes a central congenital diaphragmatic hernia.



Solution to Question 7:

The given clinical scenario is suggestive of severe combined immunodeficiency (SCID) due to a deficiency of adenosine deaminase.

Affected children present with prominent thrush (oral candidiasis), extensive diaper rash, and failure to thrive. Some patients develop a morbilliform rash shortly after birth because maternal T cells are transferred across the placenta and attack the fetus, causing graft-versus-host disease (GVHD).

Persons with SCID are extremely susceptible to recurrent, severe infections by a wide range of pathogens, including:

• Candida albicans

• Pneumocystis jiroveci

• Pseudomonas

• Cytomegalovirus

• Varicella

The thymus biopsy shows remnants of Hassall's corpuscles, indicating severe combined immunodeficiency (SCID) due to adenosine deaminase (ADA) deficiency. SCID affects both humoral and cell-mediated immune responses, primarily impacting T cells and resulting in the absence or severe underdevelopment of lymphoid tissues like lymph nodes and tonsils.

There are two main types of SCID:

• X-linked SCID: The most common form, caused by a mutation in the common γ-chain (γc) subunit of cytokine receptors. This leads to reduced T-cell numbers and impaired antibody synthesis despite normal B-cell counts. The thymus shows undifferentiated epithelial cells.

• Autosomal recessive SCID: Often due to ADA deficiency, causing toxic metabolite accumulation that damages T cells, resulting in greater T-cell reduction.

In both forms, the thymus is small and lacks lymphoid cells, but ADA deficiency uniquely features remnants of Hassall's corpuscles.

Note: X-linked SCID is the first human disease in which gene therapy has been successful.

Other options:

Option A: Lesch-Nyhan syndrome is due to a deficiency of the enzyme Hypoxanthine-guanine phosphoribosyltransferase(HGPRT). It is an X-linked recessive disease. The clinical features include intellectual disability, self-mutilation, aggression, hyperuricemia, and gout.

Option B: Ataxia Telangiectasia is a progressive neurodegenerative genetic disease causing poor coordination and small dilated blood vessels. It is autosomal recessive in inheritance due to ATM gene mutation on Ch 11q22-23.

Option D: Friedrich's ataxia involves the posterior column, pyramidal, and spinocerebellar tracts. Clinical features include cardiomyopathy, optic atrophy, and diabetes mellitus. It should be differentiated from other forms of ataxia as Tabes dorsalis and SACD(Subacute Combined Degeneration).

Solution to Question 8:

The given clinical scenario along with the electron micrograph image of the stool sample showing a spoke wheel pattern with complaints of non-bloody diarrhea and dehydration in a child is suggestive of rotavirus.

Rotavirus is a nonenveloped, double-stranded (ds) RNA virus. Rotaviruses appear as cartwheel-shaped structures with short spokes radiating from them. It has 11 segments of double-stranded RNA. They exhibit genetic reassortment due to the presence of this segmented RNA.



Rotavirus is the most common identifiable viral cause of diarrhea in children below the age of 5 years and is most frequent between 6-24 months of age. The incubation period is 1-3 days. Most cases are asymptomatic or with mild symptoms like watery diarrhea, vomiting, low-grade fever, and temporary lactose intolerance. Diagnosis can be made by rapid antigen detection of rotavirus in stool specimens.

Rotavirus vaccine is associated with a potential risk of intussusception, especially when the first dose of these vaccines is given to infants aged >12 weeks. Because of this risk, the rotavirus vaccine is not a part of catch-up immunization programs.

Symptoms associated with different levels of dehydration are classified into 3 types no/minimal dehydration, mild to moderate dehydration, and severe dehydration.

Other options:

Option A: Adenovirus contains double-stranded DNA. They have an icosahedral symmetry with structures known as fibers extruding from the vertices. They have a space vehicle-shaped appearance.



Option B: E coli are motile, facultative anaerobes in GIT. It is a gram-negative, flagellated rod. It is the most frequent cause of acute complicated urinary tract infections. It is a lactose fermenter and hence, grows on MacConkey medium to produce pink colonies.



Option C: Norwalk virus is a small, round, non-enveloped virion with a single-stranded positive-sense RNA genome with icosahedral symmetry. When viewed under an electron microscope, it has a cup-shaped morphology. Clinical features include nausea, vomiting, and diarrhoea.



Obstetrics and Gynecology AIIMS 2017

Question 1:

A pregnant lady with PIH was being monitored with bed rest in supine position. Which of the following is a complication of the below representation?



a) Ascending aorta syndrome

b) Superior vena caval syndrome

c) Abdominal aorta syndrome

d) Supine venacava syndrome

Question 2:

Which of the following is part of the WHO guidelines for the management of second stage of labor?

a) Do a routine episiotomy

b) Manual support to perineum to maintain flexion of head

c) Never give warm compress to the perineum

d) Perform labor in lithotomy position

Question 3:

Identify the best parameter to assess in case of male infertility

a) The quantity of sperms

b) The motility of sperms

c) The morphology of sperms

d) The number of sperm

Question 4:

A 26-year-old woman underwent a workup for primary infertility. On hysterosalpingography, there was a bilateral cornual block. What is the next step in management?

a) Tuboplasty

b) Hysteroscopy and laparoscopy-guided surgery

c) Hydrotubation

d) In-vitro fertilization

Question 5:

A 40-year-old woman comes with a history of postcoital bleeding and a visible mass seen in the cervix per speculum. The next step to be done:

a) Conization

b) Cryotherapy

c) Targeted biopsy

d) Pap smear

Question 6:

For emergency contraception, a single tab of 0.75 mg of LNG was taken, it should be followed by which of the following?

a) Single dose of 0.75 after 12 hrs

b) Two doses 12 hrs apart

c) Three doses 12 hrs apart

d) None of the above

Question 7:

A primigravida came to the labor room at 40 weeks + 5 days gestation for induction of labor. On per vaginal examination, the cervix is 1 cm dilated and is 30% effaced. The vertex is at –1 station and the cervix is soft and posterior. What will be the bishop score for this lady?

a) 0

b) 3

c) 5

d) 8

Question 8:

A woman whose obstetric score is G2 P1 L1 had an LMP of Jan 9th 2017. Calculate the EDD according to Naegele's rule.

a) September 19

b) October 19

c) October 16

d) September 16

Question 9:

Identify the grip in the image given below:



a) Second Pelvic grip

b) Abdominal grip

c) Pawlik grip

d) Fundal grip

Question 10:

A primigravida came with 6 cm cervical dilatation and a contraction rate of 3/10 minutes. Which stage of labor is she in?

a) 1st stage

b) 2nd stage

c) 3rd stage

d) 4th stage

Question 11:

Identify the following instrument commonly used in obstetrics.

a) Mayo scissors

b) Curved scissors

c) Suture remover

d) Episiotomy scissors

Question 12:

A 53-year-old postmenopausal woman is found to have endometrial hyperplasia with atypia. What is the next step in management?

a) Progesterone

b) Hysterectomy

c) Mirena

d) Dilatation and curettage

Question 13:

A 16-year-old female patient presents with an ovarian mass. On further evaluation, her serum biomarkers are found to be normal except for LDH, which is found to be elevated. The most likely diagnosis is:

a) Dysgerminoma

b) Endodermal sinus tumor

c) Malignant teratoma

d) Mucinous cystadeno carcinom

Question 14:

A pregnant woman has a fetus with Down's syndrome. Triple test done at 15-16 weeks will have _____.

a) Low MSAFP, high hCG, low UE3

b) High MSAFP, low hCG, high UE3

c) Low MSAFP, high hCG, high UE3

d) High MSAFP, low hCG, low UE3

Question 15:

In a partogram, where do you start plotting with 1st contractions?

a) Before the action line

b) After the action line

c) Before the alert line

d) After the alert line

Question 16:

Which of the following drug must always be available for emergency use in labor ward if a patient is on opioid analgesic?

a) Fentanyl

b) Naloxone

c) Morphine

d) Bupivacaine

Question 17:

A pregnant lady of 49 kg is in her first trimester. What are her additional calorie requirements in this trimester?

a) 500

b) 300

c) Continue same diet

d) 600

Question 18:

A 18 year old girl with primary amenorrhea, no breast development and Prader 3 genitalia was found to have raised FSH. Her karyotype revealed 45 XO/46 XY and USG revealed Mullerian structures with poorly visualized gonads. What would be the next line of management?

a) Vaginoplasty

b) Bilateral gonadectomy

c) HRT to induce puberty

d) Clitoroplasty

Question 19:

Fixative for PAP smear:

a) 10% buffered formalin

b) 95% ethyl alcohol

c) Air dried

d) None of the above

Question 20:

The deepest part of perineal body injury, which causes cystocoele, enterocoele and urethral descent is ____

a) Pubococcygeous

b) Bulbospongiosus

c) Ischiocavernous

d) Superficial transverse perineal

Question 21:

Which is not seen in pregnancy induced hypertension:

a) Systemic hypertension within 10 weeks of gestation

b) Diabetic nephropathy

c) Diabetic retinopathy

d) Thrombocytopenia

Question 22:

A couple with 2 children who are not able to conceive for over an year now present to the OPD. The male partner is diagnosed to be having hypogonadotrophic hypogonadism. Which of the following is not true ?

a) Low LH AND FSH

b) Low testosterone

c) Oligospermia

d) Low prolactin levels

Question 23:

HPV nonavalent vaccine is active against which of the following serotypes?

a) 6, 11, 16, 18, 20, 23, 35, 42, 52

b) 6, 11, 16, 19, 23, 33, 45, 58

c) 6, 11, 16, 18, 32, 34, 45, 52, 58

d) 6, 11, 16, 18, 31, 33, 45, 52, 58

Question 24:

A 30 year old woman came to OPD with a twin pregnancy at 10 weeks of gestation. She has a 3 year old female child who was born at term and had 2 first trimester abortions previously. Which of the following is the correct representation of her obstetric score/formula. G=gravid, P= para:

a) G4P2L1A2

b) G4P1 A2 L1

c) G5P0 L1 A 2

d) G5P1 A2 L1

Question 25:

The partogram is plotted in regular intervals after _ cm dilatation:

a) 5 cm

b) 8 cm

c) 4 cm

d) 6 cm

Question 26:

In fetal alcohol syndrome, which of the following is not seen?

a) Thinning of corpus callosum

b) Microcephaly

c) Holoprosencephaly

d) Macrocephaly

Question 27:

A 32 year old female nearing her due date has been admitted following regular contractions. The resident doctor did an examination and told the intern that the head is at +1 station. Where is the exact position of head?

a) At the perineum

b) Just below ischial spine

c) Just above ischial spine

d) High up in the false pelvis

Question 28:

Ureteric injury is most likely associated with which of the following?

a) Wertheim's hysterectomy

b) Abdominal hysterectomy

c) Vaginal hysterectomy

d) Anterior colporraphy

Question 29:

A lady was discharged after a normal vaginal delivery. On third day she came back with fever, tachycardia and seizures. Fundus showed papilledema with no focal deficits. Diagnosis is:

a) Subarachnoid haemorrhage

b) Meningitis

c) Acute migraine

d) Cortical vein thrombosis

Question 30:

Rh negative mother, ICT negative was given Anti D during 28 week of pregnancy. Which of the following is the ideal one

a) No need additional dose since she is ICT negative

b) Give another dose of Anti D 72 hour postpartum irrespective of baby blood group

c) Give another dose of Anti D 72 hour postpartum depending on the baby blood group

d) All of the above

Question 31:

A 25 year old female came with complaints of 4 months amenorrhea. Her FSH and LH were elevated above the normal value, thyroid function tests were normal. Which is the next step in management ?

a) Estimate serum estradiol values

b) Give progesterone and stop after 10 days for withdrawal bleeding

c) Check USG for polycystic ovaries

d) Do a urine pregnancy test

Question 32:

A 25-year-old female came with complaints of greenish discharge per vaginum and on examination she had a strawberry cervix. Which of the following is the treatment for her

a) Doxycycline

b) Amoxicillin

c) Fluconazole

d) Metronidazole

Question 33:

A 30 year old female came for routine PAP smear testing. The report came as carcinoma in situ. What is the next step?

a) Conisation

b) Hysterectomy

c) Colposcopy and biopsy

d) HPV DNA testing

Question 34:

A 62-year-old female with history of ovarian carcinoma in the mother achieved menopause with smooth transition. And she is on hormone replacement therapy. She developed

abdominal pain and on examination the right adnexal mass was palpable, non tender. On USG examination she has a 3 X 3 cm smooth cyst in her ovary. Which is the next step

a) Reassure and advice regular follow up once in 6 months

b) Confirm USG findings by ordering CT scan

c) Do an exploratory laparoscopy to visualise the nature of cyst

d) Do CA-125 levels, if normal advise regular check up

Question 35:

In endometriotic lesion, histology represents its:

a) High cholesterol

b) High estrogen

c) High levels of prolactin

d) High progesterone

Answer Key

Question No.	Correct Option
1	d
2	b
3	c
4	b
5	c
6	a
7	c
8	c
9	c
10	a
11	d
12	b
13	a
14	a
15	c

16	b
17	c
18	b
19	b
20	a
21	a
22	d
23	d
24	b
25	c
26	d
27	b
28	a
29	d
30	c
31	a
32	d
33	c
34	d
35	b

Detailed Explanations

Solution to Question 1:

Supine vena cava syndrome is the complication of the given representation. The image shows compression of the inferior vena cava by the gravid uterus which can sometimes lead to supine hypotension.

During late pregnancy, the gravid uterus compresses the great vessels against the hard spine, when the patient is in the supine position. Of the two vessels, the vena cava is compressed, but the aorta, owing to its structure is relatively resistant.

Normally, the vena cava compression results in opening up collateral circulation by means of paravertebral and azygous veins but in some cases, when the collateral circulation fails to open up, the venous return of the heart may be seriously curtailed resulting in hypotension, tachycardia, and syncope and is called supine vena cava syndrome.

The normal blood pressure is quickly restored by turning the patient to the lateral position.

Solution to Question 2:

The practice of guarding the perineum during delivery of the fetal head is part of the WHO management guidelines for the second stage of labor.

The fingers of one hand support the perineum, while the second hand applies pressure to the fetal head to control the speed of crowning. The advantage of this is to prevent perineal tears.

The second stage of labor begins with full dilatation of the cervix and ends with the delivery of the fetus.

The main events include the descent of the fetus (bearing down by the mother, uterine contractions), cardinal movements of labor, and delivery of the fetus.

Management of the second stage of labor:

• The most common position used is the dorsal position with the hips flexed, knees flexed and abduction of the thighs. WHO recommends any position of choice including the upright position.

• WHO recommendations for the prevention of perineal tears include warm compresses to the perineum, perineal massage, guarding the perineum, and modified Ritgen's maneuver.

• Modified Ritgen's maneuver: A sterile towel or pad is held in one hand and placed over the perineum; the other hand applies gentle downward pressure on the occiput to promote flexion and controlled delivery of the head (ensuring the smallest diameter of the head passes through the outlet). Once the occiput is born–with gentle pressure, the head is extended–forehead, nose, and mouth are then delivered.

• WHO doesn't recommend fundal pressure and routine episiotomy.

Other options:

Option A: A liberal or routine episiotomy is discouraged. Episiotomy is indicated in cases of:

• Shoulder dystocia

• Breech delivery

• Fetal macrosomia

• Operative vaginal deliveries

• Persistent occiput posterior positions

Option C: Manual techniques, such as massage or application of warm compresses, are recommended for healthy pregnant women requesting pain relief during labor.

Option D: The position that is most comfortable to the mother i.e. upright/squatting position is encouraged. This position gives:

• less discomfort and difficulty in bearing down

• less labor pain

• less perineal/vaginal trauma and wound infections

Second stage	Nulliparous	Multiparous
Mean/average duration	50 minutes	30 minutes
Maximum duration	2 hours (+1 with epidural)	1 hour (+1 with epidural)

Second stage	Nulliparous	Multiparous
WHO maximum duration of 2nd stage/diagnosis of arrest	3 hours	2 hours

Solution to Question 3:

The morphology of sperms is the best parameter to assess male infertility.

Characterization of normal sperm morphology shows improved correlation with fertilization rates during IVF cycles.

The criteria requires careful analysis of the following:

• The shape and size of the sperm head,

• The relative size of the acrosome in proportion to the head,

• And characteristics of the tail, including length, coiling, or presence of two tails.

Significantly decreased fertilization rates are observed when normal morphology of the sample falls below 4 percent.

Abnormal sperm morphology is termed teratospermia or teratozoospermia. Whereas sperm concentration and progressive motility have value for distinguishing fertile from infertile men, strict sperm morphology (as determined by an individual having extensive training and experience) was the one most discriminating value.

Parameters In Semen Analysis	Reference Limits
Volume	>1.5mL
Count	>15 million/mL
Total motility	>40%
Morphology	>4% normal forms
WBCs	< 1 million/mL
Round cells	<5 million/mL

Solution to Question 4:

In the given clinical scenario, the next step in management will be hysteroscopy and laparoscopy-guided surgery.

Tubal contractions, particularly cornual spasms, can give an incorrect impression of proximal fallopian tube obstruction (a false-positive result) on hysterosalpingography. Laparoscopy with chromotubation is therefore considered to confirm the diagnosis of cornual spasm. Laparoscopy is both diagnostic and therapeutic in this condition.

Hysteroscopy is an operative procedure that provides an endoscopic view of the endometrial cavity and tubal ostia via the cervix, performed in the preovulatory phase. It can be diagnostic or therapeutic. It is the gold standard investigation for visualizing the uterine cavity and tubal ostia.

Contraindications to hysteroscopy are as follows:

• Acute genital tract infection (Eg: genital herpes infection)

• Viable intrauterine pregnancy

• During menstruation, the view is obscured, and the infection rate is high.

• Scarred uterus and enlarged uterus of size ≥ 12 weeks are relative contraindications.

• Cervical stenosis can cause cervical tears and uterine perforation.

• Cardiopulmonary disorder due to increased risk of fluid overload and pulmonary edema.

• Known cases of cervical cancer

Laparoscopy is the gold standard for the evaluation of tubal factors in infertility, commonly done in the proliferative phase. It can be diagnostic or therapeutic.

Benefits of laparoscopic evaluation:

• Tubes: For detection of tubal patency, block (site and side), motility, hydrosalpinx change, peritubal adhesions, fimbrial agglutination

• Ovaries: PCOS changes, endometriosis, PID, adhesions

• Uterus: Anomalies, fibroids

• Peritoneal factors: Adhesions, PID, endometriosis, tuberculosis

• Therapy at the same sitting (as appropriate)

Therapeuticindications of hysteroscopy	Diagnosticindications of hysteroscopy
Uterine septumAsherman syndromeRetained IUCDPolypectomySubmucous fibroidAbnormal uterine bleedingTubal sterilization using sclerosing agents	To study the endocervical mucosal liningCongenital malformations of the uterusEndometrial tuberculosisMisplaced IUCDEndometrial lesionsEndometrial polypAsherman syndromeCornual tubal blockage

Diagnostic	Therapeutic (Operative)
Age >35 yearsHistory of pelvic pain, dyspareunia, pelvic surgery, adhesionsAbnormal HSGFailure to conceive after a reasonable period (6 months) with normal HSGWomen with comorbid pelvic pathology (PID, endometriosis)	Unexplained infertilityReconstructive tubal surgeryAdhesiolysisFulguration of endometriotic implantsOvarian drilling in PCOSGIFT and ZIFT procedures

Solution to Question 5:

A targeted biopsy is the next line of management in the patient with a visible cervical mass and postcoital bleeding. A pap smear is done when there are symptoms with no identifiable lesion.

Other options:

Option A: Conization is an invasive procedure where the cervix is cut in a cone-shaped manner. It is both diagnostic and therapeutic. It is indicated for diagnosis in women with HSIL or adenocarcinoma in situ and may be considered under the following conditions:

• The limits of the lesion cannot be visualized with colposcopy.

• SCJ (squamocolumnar junction) is not seen in colposcopy.

• Colposcopy is positive for CIN 2/CIN 3.

• Lack of correlation between cytology, biopsy, and colposcopy.

• Suspected micro invasion in cytology, colposcopy, or biopsy.

• Colposcopist is unable to rule out invasive cancer.

Hence it is very clear that conization is not indicated for all symptomatic women and it should be done following either a PAP, colposcopy or a biopsy with an abnormal or inconclusive result.

Option B: Cryotherapy is ideal only for small superficial lesions. It is indicated for biopsy-proven lesions. Cryotherapy is appropriate for CIN 1 which has persisted for more than 24 months and is confined to the ectocervix.

Option D: The false negative rate of pap smear in the presence of invasive cancer is 50% and a negative pap test should never be relied on in a symptomatic patient, especially one with a visible lesion.

CIN: Cervical intraepithelial neoplasia, HSIL: High squamous intraepithelial lesion, LSIL: Low squamous intraepithelial lesion

Solution to Question 6:

In this case, the 2nd dose of LNG 0.75 mg (levonorgestrel) is taken after 12 hours.

Levonorgestrel can be used in emergency contraception without major side effects. One tablet of LNG 0.75 mg should be taken within 72 hours of unprotected intercourse, and another after 12 hours. Alternatively, two tablets can be as a single dose (1.5 mg).

The tablets can be offered up to 120 hours but the efficacy decreases with the longer coital-drug interval.

Advantages of LNG tablets:

• It has no estrogen and its associated side effects.

• It can be offered to hypertensive, cardiac and diabetic women.

• It can be offered to a lactating woman.

• It can be given as late as 120 hours after unprotected intercourse (but efficacy decreases).

• Single-dose therapy is an advantage.

Contraindications to LNG tablets include liver disease, a history of thrombophlebitis and migraine.

Indications for emergency contraception:

• Unprotected intercourse

• Condom rupture

• Missed pill

• Delay in taking progesterone-only pill for >3 hours

• Rape/Sexual assault

Drugs used in emergency contraception:

• Mifepristone (RU486) is a progesterone antagonist, which blocks the hormonal action on the endometrium, causing the sloughing and shedding of decidua, preventing implantation. A single dose of 25-50 mg is effective in pregnancy in 99.1% of cases. It causes delayed menstruation and expensive than LNG.

• Ulipristal is a synthetic progesterone hormone receptor modulator. It binds to progesterone receptors and prevents/delays ovulation and suppresses endometrium, preventing implantation. A 30 mg tablet should be taken within 5 days. A two per cent pregnancy rate has been reported. Side effects are headaches and mood changes.

• Centchroman: 2 tablets (60 mg) taken twice in 24 hours, within 24 hours of intercourse can prevent implantation in 99% of women.

• Copper T IUCD: inserted within 5 days of unprotected intercourse can prevent implantation of a fertilized ovum. The advantages of copper T IUCD are- it is cheap, has a failure rate of 0.1%, inserted as late as 5 days after unprotected intercourse and remains an ongoing contraceptive method for 3-5 years.

• Yuzpe method: ethinyl estradiol 50 μg + norgestrel 0.25 mg. It is given as 2 tab stat and 2 after 12 hours.

Solution to Question 7:

The Bishop's score, in this case, is 5.

Bishop's pre-induction Cervical Scoring System.

Total score = 13; Favorable score = 6-13; Unfavorable score = 0-5

In the modified Bishop score (1991) the effacement is replaced by cervical length in cm.

From the data given in the question:

• Cervical dilatation is 1 cm, hence score = 1.

• Effacement is 30%, hence score = 0

• Station of the head is –1, hence score = 2

• Consistency of cervix is soft, hence score = 2

• Position of the cervix is posterior, hence score = 0

Hence, Bishop's score is 1+0+2+2+0 = 5.

ScoreCervix	0	1	2	3
Dilatation (cm)	Closed	1-2	3-4	>5
Effacement (%)	0-30	40-50	60-70	≥80
Consistency	Firm	Medium	Soft	-
Position	Posterior	Midline	Anterior	-
Head station	-3	-2	-1,0	+1, +2

Score	0	1	2	3
Cervical length in (cm)	More than 4	2-4	1-2	Less than 1

Solution to Question 8:

As per Naegele's rule, the EDD (expected date of delivery) of this woman is October 16, 2017.

Naegele's rule states that EDD is obtained by adding 7 days and subtracting 3 months from the LMP (the 1st day of the last menstrual period). Another method is adding 9 months and 7 days to LMP. Less than 5% of births follow Naegele's rule.

It is based on a 28-day regular cycle. If the cycle is shorter/longer than 28 days, corrected EDD is calculated as follows:

• For a 35-day cycle, add another 7 (35-28=7) days to the calculated EDD to get the corrected EDD.

• For a 25-day cycle, subtract 3 (28-25=3) days from the calculated EDD to get the corrected EDD.

Naegele's rule cannot be followed in IVF pregnancies. Considering the human embryo takes 266 days to develop, EDD is calculated in IVF pregnancies by the following method:

• Same day as fertilization (day 0): EDD = date of transfer + 266 days

• 3 days after fertilization (day 3): EDD = date of transfer + 263 days

• 5 days after fertilization (day 5): EDD = date of transfer + 261 days

Solution to Question 9:

Leopold's third maneuver (also called the Pawlik's grip) is performed by grasping with the thumb and fingers, the lower portion of the maternal abdomen just above the symphysis pubis. If the presenting part is not engaged, a movable mass will be felt.

Abdominal palpation- Leopold maneuvers:

In the abdominal examination, the mother is made to lie supine and comfortably positioned with her abdomen barred. Leopold maneuvers can be performed throughout the latter months of pregnancy and during and between the contractions of labor.

Leopold's maneuvers may be difficult if not impossible to perform and interpret if the patient is obese, if the amnionic fluid volume is excessive, or if the placenta is anteriorly implanted.

With experience, it is possible to estimate the size of the fetus.

• 1st Leopold (fundal grip): identifies which fetal pole (cephalic or podalic) occupies the uterine fundus. The fetal head is hard and round and is more mobile and ballotable whereas the breech is felt as a large, soft and irregular mass.

• 2nd Leopold (lateral/umbilical grip): palms are placed on either side of the abdomen and gentle, but deep pressure is exerted. The back is felt as a hard and resistant structure. The fetal limbs are felt as numerous, small, irregular parts. By noting whether the back is directed anteriorly, transversely or posteriorly, the fetal orientation can be determined.

• 3rd Leopold (pawlik's grip): it aids in confirmation of fetal presentation. In transverse lie, pawlik's grip is empty.

• 4th Leopold (pelvic grip): it helps to determine the degree of descent.



Solution to Question 10:

In this case, cervical dilatation of 6 cm and contraction rate of 3/10 minutes is indicative of first stage of labor.

The first stage of labor is characterized by dilatation and effacement of cervix and can be divided into latent phase and active labor.

• The onset of latent labor is the point at which the mother perceives regular contractions.

• Cervical dilatation of ≥6 cm in the presence of uterine contractions can be taken to reliably represent the threshold for active labor.

The second stage of labor begins with complete cervical dilatation and ends with fetal delivery.

The third stage of labor is characterized by the delivery of the placenta.

The fourth stage is a stage of observation for at least an hour after the full expulsion of the placenta.

Solution to Question 11:

The instrument given in the picture is episiotomy scissors. The angulation in the scissor is to prevent the extension of pelvic tears into the anal margins and obstetric anal sphincter injuries / complete perineal tears.

Complications of episiotomy:

They can be immediate or remote.

Immediate complications include an extension of the incision into the rectum, vulval hematoma, wound dehiscence and injury to anal sphincter.

Remote complications include dyspareunia, chances of perineal lacerations in subsequent labor and scar endometriosis (rare).

Mayo scissors are long and stout scissors. They can be blunt-tipped/pointed tip/straight or curved. Straight mayo scissors are designed for cutting body tissues near the surface of a wound. They are also used to cut sutures and are also known as suture-cutting scissors.

Curved mayo scissors allow much deeper penetration into the wound and are used to cut thick tissues such as those found in the uterus, muscles, breast, and foot. Mayo scissors used for dissection are placed in tissue with the tips closed. The scissors are then opened so that the tips open and spread out the tissue during the dissection process.

Episiotomy scissor

©MARROW

Normal Mayo scissors

©MARROW

Solution to Question 12:

Among the given options, for post-menopausal women with atypical endometrial hyperplasia, the next step in management would be a hysterectomy because the risk of progression to cancer over time approximates 29 per cent.

Due to a high rate of finding concurrent invasive malignancy coexistent with atypical hyperplasia, the ideal management is to perform a fractional curettage with hysteroscopy before proceeding with a hysterectomy (with removal of both tubes and ovaries).

Endometrial hyperplasia is endometrial thickening with a proliferation of irregularly sized and shaped glands along with an increased gland-to-stroma ratio.

Classification of endometrial hyperplasia:

• Simple hyperplasia is characterised by dilated or cystic glands with round to slightly irregular shapes, an increased glandular-to-stromal ratio without glandular crowding, and no cytologic atypia.

• Complex hyperplasia has architecturally complex (budding and infolding) crowded glands, with less intervening stroma without atypia.

• Atypical hyperplasia refers to cytologic atypia and can be categorised as simple or complex, depending on the corresponding glandular architecture. Criteria for cytologic atypia include

•

• large nuclei of variable size and shape with lost polarity

• increased nuclear-to-cytoplasmic ratios

• prominent nucleoli

• irregularly clumped chromatin with parachromatin clearing

Type of hyperplasia	Progression to cancer (%)
Without atypia	
Simple cystic	1
Complex adenomatous	3
With atypia	
Simple cystic	8
Complex adenomatous	29

Solution to Question 13:

The given clinical history of a 16-year-old female presenting with an ovarian mass and raised LDH, points towards the diagnosis of dysgerminoma.

Dysgerminoma corresponds to the seminoma of the testis and accounts for 3-5% of all ovarian tumors. It usually arises in young girls or in children, with an average age of 20 years. It is the most common malignant tumor of the ovary. It is usually unilateral, but they are the only germ cell malignancy with a significant rate of bilateral ovarian involvement (15-20%) and can be found at gonadal or extragonadal sites.

Genetics:

• Express OCT-3, OCT-4 and NANOG

• Activating mutations in the KIT gene in one-third of cases

Pathology:

• Solid neoplasm with areas of softening due to degeneration with fleshy consistency

• Mimics the pattern of the primitive gonad, lymphocytic infiltration is seen

It presents with abdominal pain, distension, a palpable mass and ascites. Some tumors occur in patients with gonadal dysgenesis, including pseudohermaphroditism. It secretes placental alkaline phosphatase, lactate dehydrogenase and beta-HCG, but no sex hormones.

Morphology and histology:

• Most dysgerminomas (80% to 90%) are unilateral tumors ranging in size from barely visible nodules to masses that virtually fill the abdomen.

• On cut surfaces, they have a solid yellow-white to gray-pink appearance and are often soft and fleshy.

• Like seminoma, it is composed of large vesicular cells with clear cytoplasm, well-defined cell boundaries, and centrally placed regular nuclei.

• The tumor cells grow in sheets or cords separated by scant fibrous stroma, which is infiltrated by lymphocytes and may contain noncaseating granulomas

The image below shows the histopathology of a dysgerminoma, characterized by uniform cells resembling primordial germ cells separated by fibrous septa with lymphocytes.



Management:

• Though dysgerminomas are radiosensitive, they are contraindicated due to ovarian destruction in young girls. It is employed only for residual and recurrent tumors.

• Treatment includes surgical removal and postoperative chemotherapy.

Chemotherapy comprises of

• Injection bleomycin 15 mg i.v (or) i.m weekly for 12 weeks

• Injection etoposide 100mg/m2 1-5 days every 3 weeks

• Injection cisplatin 20 mg/m2 1-5 days 3 weeks

Solution to Question 14:

The triple test findings in pregnancies with fetal Down's syndrome are characterized by:

• Low MSAFP (maternal serum alpha-fetoprotein) levels, approximately 0.7 MoM (multiples of median).

• Low unconjugated estriol levels, approximately 0.8 MoM.

• High hCG levels, approximately 2.0 MoM.

The triple test is done between 15-18 weeks for the diagnosis of Down's syndrome.

Various antenatal screening tests for aneuploidy are as follows:

First-trimester screening with biochemical analytes (↓) PAPP-A and (↑) hCG and USG measurement of NT (↑) can improve the detection rate (87%) of Down's syndrome. For confirmation, a prenatal genetic study (CVS, amniocentesis or cordo■centesis) has to be performed.

Second-trimester screening (quad screening) at 15–18 weeks: (↓) MSAFP, (↓) uE3, (↑) inhibin A and (↑) hCG can detect trisomy 21 in 85% of cases with a false-positive rate of 0.9%. When it is combined with first-trimester screening, the detection rate is increased to 95%.

Dual test (1sttrimester)	Triple test (2ndtrimester)	Quadruple test (2ndtrimester)
Beta HCG	Beta HCG	Beta HCG
PAPPA	Unconjugated estriol	Unconjugated estriol
This can be combined with n uchal translucency by USG	MSAFP	MSAFP
	Seruminhibin A	

Solution to Question 15:

In a partograph, labor is graphed and analysis includes use of action lines and alert lines.

A partograph was designed by WHO for use in developing countries. Partograph is similar for nulliparas and multiparas.

Active phase starts at 3cm cervical dilatation and progress should be no slower than 1cm/hour.

A 4 hour wait is recommended before intervention when the active phase is slow.

In the modified partograph, there is no plotting of the latent phase. The partograph is plotted once active labor begins.



Solution to Question 16:

Naloxone is a narcotic antagonist capable of reversing respiratory depression induced by opioid narcotics.

It acts by displacing the narcotic from specific receptors in the central nervous system.

Naloxone:

It is a competitive antagonist on all types of opioid receptors. It is devoid of any kind of agonistic activity even at high doses. Injected intravenously (0.4–0.8 mg) it promptly antagonizes all actions of morphine: analgesia is gone, respiration is not only normalized but stimulated—probably due to sudden sensitization of the respiratory centre to the retained CO_2, or it may be a manifestation of acute withdrawal; pupils dilate. However, sedation is less completely reversed.

Naloxone is inactive orally because of high first-pass metabolism in the liver. Injected i.v. it acts in 2–3 min. The primary pathway of metabolism is glucuronidation. Plasma t½ is 1 hour in adults and 3 hours in newborns. Duration of action is 1–3 hrs.

Adverse effects of naloxone are uncommon, and may include a rise in BP and pulmonary edema; Naloxone 0.4 mg i.v. precipitates morphine withdrawal in dependent subjects: the syndrome lasting 2–3 hours.

Uses of naloxone:

• drug of choice for morphine poisoning (0.4–0.8 mg i.v. every 2–3 min: max 10 mg)

• reversing neonatal asphyxia due to opioid use during labour (10 μg/kg in the cord)

• treat the overdose with other opioids and agonist-antagonists (except buprenorphine)

• to reverse respiratory depression due to intraoperative use of opioids: 0.1–0.2 mg i.v. (this dose usually preserves analgesia in the postoperative period)

• tried as an adjunct to intraspinal opioid analgesia: reverses respiratory depression without abolishing pain relief

Naloxone is contraindicated in newborns of narcotic-addicted mothers as it can lead to neonatal abstinence syndrome (NAS).

NAS is a drug-withdrawal syndrome that most commonly follows in-utero exposure to maternal opioids. It also can complicate exposure to alcohol or benzodiazepines. The syndrome shows hypertonia, autonomic instability, irritability, poor sucking reflex, and seizures. Care includes close observation and pharmacotherapy with morphine and methadone. Naloxone is not recommended as it can precipitate withdrawal symptoms.

Solution to Question 17:

The pregnant lady can continue the same diet since there are no additional calorie requirements in the first trimester.

Pregnancy requires an additional 80,000 kcal, mostly during the last 20 weeks. To meet this demand, a caloric increase of 100 to 300 kcal/d is recommended during pregnancy. This greater intake, however, should not be divided equally during the course of pregnancy. The Institute of Medicine (2006) recommends adding 0, 340, and 452 kcal/d to the estimated nonpregnant energy requirements in the first, second, and third trimesters respectively.

Whenever caloric intake is inadequate, protein is metabolized rather than being spared for its vital role in fetal growth and development.

Solution to Question 18:

In Turner's syndrome, most affected individuals have a 45,X karyotype, while others have mosaic karyotypes (45,X/46,XX; 45,X/46,XY).The presence of Y chromosome material is associated with a 12% risk of a gonadoblastoma. If a Y chromosome is identified, laparoscopic prophylactic gonadectomy is recommended at the time of diagnosis to eliminate the risk of malignancy.

• Gonadoblastomas can be precursors to germ cell malignancies, such as dysgerminomas (most commonly), teratomas, embryonal carcinomas, or endodermal sinus tumors.

• The karyotype should be determined in any individual with delayed puberty and increased basal FSH concentrations.

• Regardless of the karyotype, the individual with hypergonadotropic hypogonadism has some form of ovarian “failure”

Note:Prader staging system is used for the degree of virilisation of the external genitalia

Prader 0: Classical / typical female external genitalia

Prader 1: Female external genitalia with clitoromegaly

Prader 2: Clitoromegaly with partial labial fusion forming a funnel-shaped urogenital sinus

Prader 3: Increased phallic enlargement. Complete labioscrotal fusion forming a urogenital sinus with a single opening

Prader 4: Complete scrotal fusion with urogenital opening at the base or on the shaft of the phallus.

Prader 5: Classical / typical male external genitalia

Solution to Question 19:

The most commonly used fixative used for pap smear is 95% ethyl alcohol.

Historically ether was used along with alcohol as a fixative for pap smear.Nowadays a fixative ethanol spray is available and can be used conveniently in an office set-up.

For further reading on the development of Pap smear technique
refer https://www.karger.com/Article/PDF/457827

Solution to Question 20:

Pubococcygeus is the deepest part of the levator ani muscle. Injury to it can cause enterocele, cystocele, and urethral descent.

Levator ani muscles are the most important muscles in the pelvic floor and maintain a constant state of contraction. They provide a stable floor and support the weight of the abdominopelvic contents against intra-abdominal forces. Weakening of levator ani and pelvic fascia from stretching or tearing during childbirth, may decrease support for the vagina, bladder, uterus, or rectum or alter the position of the neck of the bladder and urethra.

The levator ani muscle is a complex unit, which consists of 3 muscle components namely

• Pubococcygeus (anterior fibers)

• Pubovaginalis - elevates the urethra and hence contributes to urinary continence

• Puboperinealis - draws the perineal body towards the pubic symphysis

• Puboanalis - elevates the anus and keeps the urogenital hiatus narrowed

• Puborectalis (intermediate fibers)

• Iliococcygeus (posterior fibers)

Origin: The levator ani muscle has a linear origin from the pelvic surface of the body of the pubis, a tendinous arch of obturator fascia, and the pelvic surface of the ischial spine.

Insertion:

• The anterior fibres form a sling around the prostate (levator prostatae) or vagina (sphincter vaginae) and are inserted into the perineal body (a mass of fibrous tissue) in front of the anal canal.

• The intermediate fibres (puborectalis) form a sling around the anorectal junction to be inserted into the anococcygeal raphe (a fibrous raphe which extends from the anorectal junction to the tip

of the coccyx).

• The posterior fibres (iliococcygeus) are inserted into the anococcygeal raphe and the coccyx.

Nerve supply:

• perineal branch of 4th sacral nerve (S4) from its pelvic surface.

• perineal branch of the pudendal nerve (S2, S3) from its perineal surface.

Actions:

• The levator ani muscles of two sides together support the pelvic viscera and resist the intra-abdominal pressure during straining and expulsive efforts of the anterior abdominal wall muscles.

• They also subserve a sphincteric action on the anorectal junction to maintain the continence of faeces in both sexes and vagina in females.



Pelvic diaphragm (superior view)

The given image below shows the pelvic floor:



Solution to Question 21:

Hypertension induced by pregnancy is gestational hypertension which is characterized by BP more than 140/90 mmHg after 20 weeks of gestation in a previously normotensive woman.

The incidence of gestational hypertension and preeclampsia is remarkably increased in diabetic mothers.

40% of pregnant women with diabetes will develop preeclampsia. Chronic hypertension with overt diabetes increased the risk of preeclampsia to 60%

Hypertension is a risk factor associated with the progression of diabetic retinopathy.

The platelet count is usually found to be decreased in gestational hypertension.

Solution to Question 22:

A male with hypogonadotropic hypogonadism will have high prolactin levels.

Hypogonadotropic hypogonadism:

• Low LH &FSH

• Low testosterone

• Low testicular volume (oligospermia)

• High prolactin levels

Hypergonadotropic hypogonadism:

• High LH and FSH

• Low testosterone

• Low testicular volume (oligospermia)

• High prolactin level

Hypogonadotropic hypogonadism:

It can be congenital or acquired. Congenital hypogonadotropic hypogonadism is divided into anosmic hypogonadotropic hypogonadism (Kallmann syndrome) and congenital normosmic isolated hypogonadotropic hypogonadism (idiopathic hypogonadotropic hypogonadism).

Acquired hypogonadotropic hypogonadism can be caused by drugs, infiltrative or infectious pituitary lesions, hyperprolactinemia, encephalic trauma, pituitary/brain radiation, exhausting exercise, abusive alcohol or illicit drug intake, and systemic diseases such as hemochromatosis, sarcoidosis and histiocytosis X.

The clinical characteristics of hypogonadotropic hypogonadism are androgen deficiency and a lack/delay/stop of pubertal sexual maturation. Low blood testosterone levels and low LH and FSH levels confirm the diagnosis. A prolonged stimulated intravenous GnRH test can be useful. In

Kallmann syndrome, cerebral MRI can show an anomalous morphology or even absence of the olfactory bulb.

Therapy for hypogonadotropic hypogonadism depends on the patient's desire for future fertility. Hormone replacement with testosterone is the classic treatment for hypogonadism. Androgen replacement is indicated for men who already have children or have no desire to induce pregnancy, and testosterone therapy is used to reverse the symptoms and signs of hypogonadism. Conversely, GnRH or gonadotropin therapies are the best options for men wishing to have children.

Solution to Question 23:

HPV nonavalent vaccine (Gardasil 9 or HPV9) is active against HPV 6,11,16,18,31,33,45,52,58.

Currently, HPV nonnavalent vaccine has replaced Gardasil which is a quadrivalent vaccine acting against 6,11,16,18.

Vaccines available for HPV:

Brand name	Cervarix	Gardasil	Gardasil-9	Cervavac
Characteristic and HPV subtypes	Bivalent:16,18	Quadrivalent:6,11,16,18	Nonavalent:6,11,16,18,31,33,45,52,58	Quadrivalent:6,11,16,18
Protects against:	CINCA cervix	Anogenital wartsCINCA cervix	Anogenital wartsCINCA cervixVulvalintraepithelialneoplasiaVaginalintraepithelialneoplasiaAnal cancerOropharyngeal cancerHead and neck cancers	Anogenital wartsCINCA cervix
Dose	0.5 ml,IM.	0.5 ml,IM.	0.5 ml,IM.	0.5 ml,IM.
Ideal Age range:	Ideally at 11-12 years or9-26 years	Ideally at 11-12 years or9-26 years	Ideally at 11-12 years or9-26 years	The targeted age group is 9-26 years

Solution to Question 24:

The obstetric score is G4P1A2L1.

Gravida and para refer to pregnancies and not to babies.

Gravida: Total number of pregnancies including the present one.

Para: Total number of pregnancies crossing the period of viability, which is 28 weeks in India.

As such, a woman who delivers twins in the first pregnancy is still a gravida 1 and para 1. A pregnant woman with a previous history of 2 abortions and 1 term delivery can be expressed as 4th Gravida but primipara.

It is customary in clinical practice to summarise the past obstetric history by 2 digits

First digit: Number of viable births

Second digit: Number of abortions

Both digits connected with a "+" sign affixing the letter P.

Thus, P1+2 denotes patient had 1 viable birth and 2 abortions.

In some centres, it is expressed by 4digits connected by dashes, P(A-B-C-D)

A: Number of term pregnancies (37-42 weeks)

B: Number of preterm pregnancies (28 to less than 37 weeks)

C: Number of miscarriages (less than 28 weeks)

D: Number of children alive at present.

Solution to Question 25:

The partograph is plotted in regular intervals after 4 cm dilatation.

In a partograph, the labor process is divided into

• Latent phase ends when cervix dilated upto 4 cm

• Active phase begins with cervical dilatation of ≥6 cm and ends at 10 cm dilatation at the rate of 1 cm/hour.

The latent phase of labour is slow and extremely variable, while the active phase of labour is fairly predictable in terms of duration. Even though the active phase of labor is from more than 6cm dilatation, partograph is marked from 4 cm dilatation.

Solution to Question 26:

Fetal alcohol syndrome is associated with microcephaly rather than macrocephaly.

• High levels of alcohol ingestion during pregnancy can be damaging to embryonic and fetal development.

• The presence of at least one of the below characteristics is defined as fetal alcohol syndrome:

• Growth restriction before and/or after birth

• Facial anomalies: small palpebral fissures, indistinct or absent philtrum, epicanthic folds, flattened nasal bridge, short length of nose, thin upper lip, lowset and unparallel ears, retarded midfacial development.

• CNS dysfunction: Microcephaly, mental retardation, abnormal neurobehavioral development (attention deficit with hyperactivity)

Other features:

• Incomplete holoprosencephaly

• Agenesis of corpus callosum and the cerebellar vermis.

• Cardiac anomalies :VSD>ASD ,tetralogy of fallot.

Solution to Question 27:

The station is just below the ischial spine.

Station of the head is in relation to ischial spines as spines are the most prominent bony projections felt on internal examination.

The level of ischial spines is the halfway between the pelvic inlet and outlet. This level is known as station zero (0).

The levels above and below the spines are divided into fifths to represent centimeters. The station is said to be "O" if the presenting part is at the level of the spines. The station is stated in minus figures, if it is above the spines (–1 cm, –2 cm, –3 cm, –4 cm and –5 cm) and in plus figures if it is below the spines (+1 cm, +2 cm, +3 cm, +4 cm and +5 cm).



Stations of the presenting part

Solution to Question 28:

The ureteric injury is most commonly associated with Wertheim hysterectomy than other hysterectomies due to the radical nature of the procedure.

Overall incidence is 0.5 -1 percent of all pelvic operation.

Procedure	Incidence of Ureteric injury
Abdominal hysterectomy	0.04-3

Procedure	Incidence of Ureteric injury
Vaginal hysterectomy	0.02-0.47
Subtotal hysterectomy	0.03
Wertheim's hysterectomy	1-30

Solution to Question 29:

The given clinical situation likely suggests Cortical Vein Thrombosis.

Option A - Subarachnoid hemorrhage

There are no focal deficits hence unlikely.

Option B - Meningitis

• It would have presented with neck rigidity / photophobia.

• Moreover , presence of early papilledema goes against acute meningitis.

Option C - Acute Migraine

• Migraine does not present with seizures.

• Hence most likely diagnosis is Cortical Vein Thrombosis.

Cortical Vein thrombosis

Risk factors:

• Pregnancy and puerperium

• Oral contraceptives

• Pro thrombotic conditions

Clinical presentation

• Highly variable

• Head ache is the most common symptom.

• With or without

• Vomiting

• Papilledema

• Focal deficits

Investigation

• Brain MRI with MR venography – most sensitive

• CT – normal in 30 % of cases.

Solution to Question 30:

Anti-D should be given postnatally within 72 hours & all women who deliver Rh D-positive baby should have quantification of fetomaternal hemorrhage to guide appropriate dose of anti D prophylaxis.

RCOG and NICE guidelines for Anti–D prophylaxis:

1. All Rh-negative women should be offered Anti-D

First trimester indications: (50 mcg)

• Chorionic villus sampling

• Miscarriage

• Termination of pregnancy

• Ectopic pregnancy

2nd & 3rd-trimester indications (300 mcg)

• Amniocentesis, Cordocentesis

• Obstetric hemorrhage

• Abdominal trauma

• External cephalic version

2. All Rh D-negative women should be given a single dose of 1500 units (300 mcg) of anti D between 28 - 30 weeks of pregnancy.

3. Postnatally within 72 hours

4. All women who deliver Rh D-positive baby should have quantification of fetomaternal hemorrhage to guide appropriate dose of anti D prophylaxis.

Solution to Question 31:

• Since LH and FSH are elevated above the normal value, the possibility of a pregnancy can be ruled out. Therefore no need to do a urine pregnancy test.

• Hence the next step in management would be to check serum estradiol levels to evaluate amenorrhea.

In pregnancy,

• hCG stimulates Relaxin which inhibits LH.

• Inhibin A suppresses FSH.

• Therefore, LH and FSH are low.

Evaluation of amenorrhea :

Categories of amenorrhea based on Gonadotrophin and Estrogen levels.

• Hypergonadotropic hypogonadism

• Hypogonadotropic hypogonadism

• Eugonadotropic hypogonadism

The given clinical picture fits into hypergonadotropic hypogonadism.

Hypergonadotropic hypogonadism describes any process in which:

(1) Ovarian function is decreased or absent (hypogonadism) and

(2) Due to absent negative sex-steroid feedback, LH and FSH, have increased serum levels.

It is also called Premature Ovarian Failure.

Hormonal levels in POF

• Increased LH, FSH

• Low serum estradiol levels

• High levels of serum prolactin

Hence, the next best step would be to test serum estradiol levels.

Differential Diagnosis of Premature Ovarian Failure
GeneticChromosomalNormal karyotypeGonadal dysgenesisSpecific gene defectsFragile X (FMR1 premutation)Galactosemia IatrogenicOvarian surgeryGonadal radiationSystemic chemotherapyAutoimmune diseaseToxinsVirusesMiscellaneous

Solution to Question 32:

The above case suggests Trichomonas vaginalis infection and the treatment of choice is metronidazole.

The greenish-colored vaginal discharge along with the strawberry cervix is characteristic of infection with Trichomonas vaginalis which can be an STD.

The treatment regimen includes:

• Metronidazole 2 g orally in a single dose (or 400 mg 3 times/day for 5 days) is the standard treatment. It should be taken at night, after meals to avoid vomiting. If this too causes vomiting, or in resistant cases consider the use of alternative drugs.

• The alternative regimen includes tinidazole 2g as a single dose or secnidazole in a single dose of 2g. Single-dose therapy ensures better compliance.

• Both partners need to be given treatment at the same time to prevent the risk of recurrence.

The given image below shows strawberry cervix in trichomonas infection:

Solution to Question 33:

Carcinoma in situ is classified as Cervical Intraepithelial Neoplasia-III and the next step in management for such lesions is Colposcopy and biopsy.



Solution to Question 34:

Doing a CA-125 level measurement is the next best step according to the latest RCOG guidelines for Post-menopausal Ovarian cyst.



Initial steps in management include the following:

• Estimate CA – 125 level

• USG

• Calculating Risk of Malignancy index

• RMI = U x M x Ca-125

• U – Ultrasound score

• M – Menopausal status

• Ca 125 – Serum levels

Option B : CT scan is not recommended in initial evaluation.

Option C : Laparotomy/laparoscopy only indicated if size > 5 cm or RMI > 200.

Solution to Question 35:

In endometriotic lesions, histology represents its high estrogen activity.

Endometriosis is defined by the presence of “ectopic” endometrial tissue at a site outside of the uterus. Endometriosis is a proliferative hormone-dependent disease of the childbearing period. It is extremely rare before menarche and disappears after menopause.

The endometriotic implants show certain differences when compared to normal endometrium. There is increased production of estrogen by endometriotic stromal cells due to high levels of the enzyme aromatase. This is absent in normal endometrial stroma.

Management of endometriosis:


Management of symptomatic endometriosis
Medical
Minimal invasive surgery
Surgery
Drug treatment
G - GnRH analogue
O - OCPs
L - Letrozole
D - Danazol
G - Gestrinone
A - Anti progesterone - Mifepristone
P - Progesterone
Mnemonic: GOLD GAP
Laparoscopy
Destruction by cautery, laser vapourization
Excision of cyst
Adhesiolysis
Presacral neurectomy
LUNA (laparoscopic uterosacral nerve ablation)
Laparotomy
Incision of chocolate cyst, and removal of lining
Salpingo-oophorectomy
Hysterectomy and unilateral or bilateral salpingo-oophorectomy
Excision of scar endometriosis

Obstetrics and Gynecology AIIMS 2018

Question 1:

Identify the instrument.

a) Episiotomy scissors

b) Curved artery

c) Curve-bladed Mayo

d) Mosquito forceps

Question 2:

At term pregnancy, the normal amniotic fluid appearance is _____.

a) Clear with white specks

b) Cloudy without white specks

c) Clear without white specks

d) Cloudy with white specks

Question 3:

Identify this condition.

a) Twin to Twin Transfusion Syndrome

b) Monochorionic diamniotic pregnancy

c) Monochorionic monoamniotic pregnancy

d) Dichorionic diamniotic pregnancy

Question 4:

A lady with 7 weeks amenorrhea presented with slight vaginal spotting. Crown Rump Length was 5 mm with a well-formed gestational sac with a calculated gestational age of 5.6 weeks on TVS. What is the next line of management?

a) Wait and watch for 4 weeks

b) Repeat TVS after one week

c) Serum HCG levels

d) Medical intervention

Question 5:

Which of the following would be considered as an absolute contraindication for insertion of copper T to a patient?

a) Unmotivated patient

b) Menorrhagia

c) Previous ectopic pregnancy

d) Previous history of abortion

Question 6:

A 76 year-old female presented with a non-healing ulcer on the labia majora for a period of 6 months, measuring 2 cm x 3 cm. No lymph nodes were palpable. Biopsy shows squamous cell carcinoma. How do you manage this case?

a) Total vulvectomy with unilateral LN dissection

b) Partial vulvectomy with ipsilateral inguinofemoral LN dissection

c) Simple vulvectomy

d) Chemoradiation with resection

Question 7:

A female comes to Gynecology OPD for pre-conceptual counseling, with 2 prior second trimester abortions. What is the next investigation you will advise?

a) Pelvic sonography

b) Hysterosalpingography

c) Endometrial Biopsy

d) Karyotyping of the parents

Question 8:

A pregnant female with known cardiac disease presents in the first trimester with history of warfarin embryopathy. What should be advised now?

a) Continue warfarin throughout the pregnancy

b) Replace warfarin with heparin in first trimester

c) Give acicoumarin

d) Use LMW heparin

Question 9:

A 25-year-old lady with submucosal fibroid was undergoing myomectomy. The surgeon was using 1.5% glycine as irrigating fluid for the cavity. During the surgery, the nurse informs that

there is a 500ml deficit of the irrigating fluid. What is the next step to be done?

a) Stop the surgery

b) Change the fluid to normal saline

c) Continue the surgery with careful monitoring of fluid status

d) Give furosemide to the patient and continue surgery

Question 10:

The most important marker of male fertility in semen analysis is?

a) Morphology

b) Concentration

c) Motility

d) Sperm Count

Question 11:

A 20 year old girl presents with primary amenorrhea. On examination breasts were found to be poorly developed, vagina was present. Ultrasound revealed normal uterus but gonads weren't detected. Further evaluation done showed 46XY karyotype. The most probable diagnosis?

a) Androgen insensitivity syndrome

b) Turner syndrome

c) True hermaphrodite

d) Swyer's syndrome

Question 12:

The results of semen analysis of a young male have been given below. What is the diagnosis?
5 million/ml concentration, 50% motility, 4% normal morphology

a) Oligospermia

b) Teratozoospermia

c) Asthenozoospermia

d) Oligoasthenoteratozoospermia

Question 13:

A woman presents with prolonged labor in emergency. She is being taken for Cesarean section. Which of the following is the correct position in which the nurse should keep the patient on the OT table?

a) Supine with wedge under right hip

b) Semi fowlers

c) Trendelenburg with legs stirrup

d) Prone position

Question 14:

A woman with 31 weeks of gestation presents with lower abdominal pain. The clinician is concerned if the patient is in preterm labor. Which of the following is the most appropriate test in this scenario?

a) Alpha fetoprotein

b) Beta HCG

c) Fetal fibronectin

d) PAPP-A

Question 15:

Identify the procedure done in the labor room?

a) Amniocentesis

b) MCA doppler

c) Cardiotocography

d) Amniotic fluid index determination

Question 16:

A G3P2L0 presents with 9 weeks of amenorrhea. She has a history of conization being done one year ago and is currently on follow-up, with no recurrence on PAP smear. She also has the history of preterm labor at 30 and 32 weeks. What is the next step of management?

a) USG to see cervical length

b) Cervical Cerclage

c) Complete Bed rest

d) Abdominal cerclage

Question 17:

What is the minimum endometrial thickness on transvaginal ultrasonography, above which you perform a biopsy in a case of postmenopausal bleeding?

a) 3mm

b) 4mm

c) 7mm

d) 9mm

Question 18:

A patient has been on MALA-N for 6 months. She is compliant, but she complained of missed periods. What is the next step?

a) Transvaginal ultrasound

b) Reassure her that it's normal to have delayed period

c) Beta HCG estimation

d) Serum FSH and LH

Question 19:

A pregnant woman delivered a baby with the normal expulsion of an intact placenta. After half an hour she started bleeding per vaginal. On examination, she was hypotensive and a boggy mass was palpated per abdomen. USG showed retained placental tissues. What is the likely diagnosis?

a) Placenta membranacea

b) Placenta succenturiata

c) Circummarginate placenta

d) Circumvallate placenta

Question 20:

HPV nonavalent vaccine is active against which of the following serotypes?

a) 6, 11, 16, 18, 20, 23, 35, 42, 52

b) 6, 11, 16, 19, 23, 33, 45, 58

c) 6, 11, 16, 18, 32, 34, 45, 52, 58

d) 6, 11, 16, 18, 31, 33, 45, 52, 58

Question 21:

Sentinel lymph node biopsy is most useful in which of the following gynecological malignancies?

a) Carcinoma vagina

b) Carcinoma vulva

c) Carcinoma cervix

d) Carcinoma endometrium

Question 22:

Which of the following is not included in the active management of the third stage of labor to prevent postpartum hemorrhage?

a) Uterotonic within 1 minute of delivery

b) Delayed clamping, cutting and ligation of cord

c) Gentle massage of uterus

d) Controlled cord traction

Question 23:

Which of the following changes can be seen in the quadruple screening for Down syndrome?

a) Elevated β-hCG, PAPP-A and lower estriol, AFP

b) Elevated β-hCG, Inhibin and lower estriol, AFP

c) Lower β-hCG, PAPP-A and elevated estriol, AFP

d) Lower β-hCG, Inhibin and elevated estriol, AFP

Question 24:

All of the following can be indicators of choriocarcinoma except:

a) Increased number of theca lutein cysts

b) Persistently enlarged uterus

c) Plateau of high serum hCG

d) Suburethral nodule

Question 25:

Which of the following hormones will be secreted in high amounts by the ovarian granulosa cells during the follicular phase of the menstrual cycle?

a) FSH

b) Estrogen

c) LH

d) Progesterone

Question 26:

Which of the following statements is true regarding the instrument shown below?

a) A and B

b) B and C

c) A and C

d) Only C

Question 27:

A young female has history of a spontaneous abortion. She has had amenorrhea since then. Her FSH now is 6 mIU/ml. Which of the following can be a cause for her amenorrhea?

a) Pituitary dysfunction

b) Ovarian dysfunction

c) First pregnancy

d) Uterine synechiae

Question 28:

Which among the following is an ideal treatment for a 55-year-old female with simple hyperplasia of the endometrium with atypia?

a) Simple hysterectomy

b) Medroxy progesterone acetate

c) Levonorgesterol

d) Radiotherapy

Question 29:

Which of the following is an irreversible method of contraception?

a) Laparoscopic tubal ligation

b) Copper-T

c) Implanon

d) LNG-IUD

Question 30:

What is the most useful test in vesicovaginal fistula?

a) Cystoscopy

b) 3 swab test

c) MRI

d) Cystometry

Question 31:

A 40-year-old woman has cervical intraepithelial neoplasia III. What would be the management?

a) Hysterectomy

b) Trachelectomy

c) Conization

d) Colposcopy with LEEP

Question 32:

A pregnant woman has gestational diabetes mellitus (GDM) and her blood glucose levels are well controlled on insulin. What will be the appropriate gestational age for her delivery?

a) 37 weeks

b) 39 weeks

c) 38 weeks

d) 40 weeks

Question 33:

A female presents with 35 weeks and 3 days of gestation, AFI = 5 and there is no end-diastolic blood flow. What is the management?

a) Immediate termination

b) Give surfactant and wait till 36 weeks

c) Do Doppler everyday to check for reversion

d) Terminate after 37 weeks

Question 34:

Which of the following heart sounds would be normal in pregnancy?

a) S3

b) S4

c) Fixed splitting of S2

d) Pericardial knock

Question 35:

A pregnant woman with cardiac disease is on warfarin therapy. When will the final switch to heparin be advised?

a) At the onset of labour

b) 34 weeks

c) 36 weeks

d) 40 weeks

Answer Key

Question No.	Correct Option
1	a

2	d
3	c
4	b
5	a
6	b
7	a
8	b
9	c
10	a
11	d
12	a
13	a
14	c
15	c
16	a
17	b
18	c
19	b
20	d
21	b
22	c
23	b
24	a
25	b
26	c
27	d
28	a
29	a
30	a
31	d
32	b
33	a
34	a
35	c

Detailed Explanations

Solution to Question 1:

The given instrument is Episiotomy scissors.

The episiotommy scissors is angulated to prevent extensions of the perineal tears into the anal margins and anal spinchters and it also prevents complete perineal tears.

Episiotomy is a planned incision made on the posterior vaginal wall and the perineum during the second stage of labor. It is done mostly to avoid spontaneous perineal injuries to the muscles and fascia at the time of delivery.

The ideal time to perform episiotomy is when the perineum is bulged thin during contraction, just prior to crowning. The most commonly performed episiotomy incision is mediolateral, which prevents any rectal involvement even if the tear extends posteriorly. Other types of episiotomy include: Median,lateral and J shaped.

Curved Mayo (option C) is used to cut thicker tissues.

Mosquito forceps (option D) is used to grasp the delicate tissues & blood vessels.

The table below explains major differences between median and mediolateral type of episiotomy.

Midline Episiotomy	Mediolateral Episiotomy
Surgical repair is easy	Surgical repair is difficult
Blood loss is lesser	Blood loss is more
Healing is superior	Healing is not so good
Dyspareunia is rare	Dyspareunia is comparatively more
Postoperative comfort is max imum	Postoperative discomfort is more
Extension, if occurs, may inv olve the rectum	Relative safety fromrectal inv olvementfrom extension

Solution to Question 2:

At term, the amniotic fluid is cloudy and contains a large number of white flakes of vernix.

Amniotic fluid appearance is determined by the proportion of cloudiness and the number of flakes released from the fetal skin over the period of pregnancy. This is summated by the amniotic fluid macroscore.

At the end of the 32 nd week of pregnancy, the amniotic fluid is clear and does not have any flakes (Option C). By the 36 th week, the amniotic fluid starts to have some flakes (Option A) and gradually we will begin to see traces of cloudiness also. By 42 nd week.i.e. term of pregnancy, the amniotic fluid has a large number of flakes and is very cloudy (Option D).

Solution to Question 3:

The given specimen depicts a monochorionic monoamniotic pregnancy as:

• 2 umbilical cords are arising from 1 placenta (monochorionic)

• No intervening amniotic membrane is attached to the placenta. This means that both fetuses must have been inside a single amniotic cavity (monoamniotic).

Option B: In Monochorionic diamniotic pregnancy, on gross examination, there will be 1 placenta and intervening 2 layers of amnion attached to the placenta. In this type of pregnancy, each fetus will have its own amniotic cavity (diamniotic) with a shared placenta (monochorionic). This is seen if the fertilized zygote divides into 2 between 4-8 days.

The below given gross image of monochorionic diamniotic pregnancy shows two umbilical cords arising separately from the placenta with the intervening amniotic membrane held by the forceps.



Option D: In dichorionic diamniotic pregnancy each fetus has it's own amnion & chorion. The placentas here may be fused or separate. This is seen in dizygotic twins as well as monozygotic twins that divide within 0-4 days.

Option A: Twin to twin transfusion syndrome (TTTS) most commonly complicates monochorionic (single placenta) twin pregnancy. In TTTS blood is transfused from the donor twin to the recipient twin through the abnormal vascular connections within the placenta leaving the donor anemic while the recipient becomes polycythemic.

The box below describes the type of amnion and chorion seen in different types of twins depending on the time of division.

Dizygotic twins	Dichorionic diamniotic
Monozygotic twins- division occurs within0-4 days4-8 days8-12 days>13 days	Dichorionic diamnioticMonochorionic diamnioticMonochorionic monoamnioticMonochorionic monoamniotic conjoined twins



Solution to Question 4:

The given clinical scenario of a patient with amenorrhea presenting with spotting, raises a suspicion of possible abortion and in the given case the next line of management would be to repeat TVS after one week.

In a normal intrauterine pregnancy a gestational sac will be visible on TVS by 5 weeks of LMP and the embryonic poles can be visualized by 5.5 weeks. The cardiac activity of the embryo can be first visualized on TVS around 5.5-6 weeks, when the crown rump length (CRL) is around 5mm.

In the event that, the embryonic poles are not visible on TVS, even when the gestational sac diameter has reached ≥25mm or the cardiac activity is not visualized even at CRL ≥7mm a diagnosis of missed abortion can be made. The values, mean sac (gestational sac) diameter ≥25mm and CRL ≥7mm are known as critical cut off values which represent the upper limit beyond which if a structure is not visualized, it is abnormal.

Hence, Missed Abortion should be suspected if, on USG

• CRL ≥7 mm and no heart beat is recorded (CRL-crown rump length)

• MSD ≥25 mm and no embryo is seen (MSD-mean sac diameter)

In this patient the CRL is still at 5 mm and hence, a diagnosis of missed abortion cannot be made. Thus, the next line of management would be to repeat TVS after one week.

Structure	Critical cut-off to visualize th e structure on USG	Structure visible on TVS at
Gestational sac in uterus(intr auterine pregnancy)	On TVS : 2000 IU of beta HC GOn TAS :6500 IU of beta H CG	4 weeks 3 days of LMP to 5 w eeks
Yolk sac	-	5 weeks

Structure	Critical cut-off to visualize th e structure on USG	Structure visible on TVS at
embryonic/fetal poles	Mean sac diameter >/= 25 m m	5 weeks - 5 1/2 weeks
Cardiac activity	CRL >/= 7mm	5 1/2 weeks - 6 weeks

Solution to Question 5:

An unmotivated patient or someone who does not consent to the procedure should never be taken up for a procedure.

Contraindications for use of copper intrauterine devices (IUDs)

• Uterine abnormality with a distorted uterine cavity

• Active uterine and pelvic infection

• Known or suspected uterine or cervical malignancy

• Genital bleeding of unknown etiology

• Mucopurulent cervicitis

• Wilson disease

• A previously placed IUD that has not been removed

• An inability to place or retain the device

Note:

An unmotivated patient is an absolute contraindication. Out of the given options, this would be the best answer.

Solution to Question 6:

According to the recent FIGO guidelines,the patient's lesion measuring 2 cm x 3 cm with no palpable lymph nodes falls under stage Ib vulval cancer and it's management is partial vulvectomy with ipsilateral inguinofemoral lymph node dissection.

For stage IB cancers, radical partial vulvectomy is recommended for the primary tumor and as the women with stage IB cancer can have diseased lymph nodes even without palpable enlargement of nodes, Sentinel Lymph Node Biopsy (SLNB) and/ inguinofemoral lymphadenectomy is performed.

Ipsilateral dissection of lymph nodes is performed if the vulvar lesion is present laterally, >2cm beyond the midline. In cases where the vulvar lesion lies within 2 cm from the midline, bilateral inguinofemoral Lymph Node dissection will be performed.

In this patient, the best management would be to perform Partial vulvectomy with ipsilateral inguinofemoral LN dissection among the given options.

Solution to Question 7:

The most common cause of second trimester abortions are uterocervical structural anomalies and the next investigaton advised in this patient would be pelvic sonography to look for these anomalies.

Although, Hysterosalpingography (Option B) can be used in the patient during the interconceptional period to diagnose internal anatomical abnormalities of the uterus and cervix, it is an invasive procedure and hence is reserved for use only after nonivasive procedures such as sonography is done. Hysterosalpingography (HSG) has limited use in diagnosing the external uterine contour and it is contraindicated in pregnancy.

Pelvic sonography is a non-invasive investigation that can be used to detect certain anatomical anomalies leading to abortions. A 3D sonography will provide better images of uterine and cervical contour than a 2D sonography. This investigation can also be done in pregnant female.

Karyotyping (Option D) is done to rule out any suspicion of chromosomal abnormalities which, more commonly cause first trimester abortions.

NOTE: If a suspicion of cervical incompetence (H/O 1 painless second trimester loss) is present, a TVS is performed in a pregnant female between 18-24weeks and the cervical length and the cervical shape is determined.

• Normal cervical length-3-4cm ; In cervical incompetence ≤2.5cm

• Normal cervix shape-Tshape; In cervical incompetence U shape

In the presence of ≥ 2 painless second trimester abortions, a history based diagnosis of cervical incompetence is made and the pregnant female can undergo prophylactic cervical cerclage with progesterone supplementation.

Some Causes of Midtrimester (second trimester) abortions
Fetal Anomalies- Structural,ChromosomalMaternal Disorders-Autoimmune, Infections, Metabolic disordersPlacental Causes-Abruption, previa Vasculopathy,ChorioamnionitisUterine Defects-Congenital anomalies, submucosal Leiomyomas, cervical incompetance

Solution to Question 8:

In the first trimester, warfarin has to be replaced with heparin due to its potential teratogenic effects such as miscarriage, stillbirths, achondroplasia and stippled epiphyses.

Low molecular weight heparin (LMWH) must replace warfarin and shouldn't be added along with the warfarin (Option D).

The detailed guideline for anticoagulation in pregnancy is as follows:

• LMWH: low molecular weight heparin

• EDD: expected date of delivery

• UFH: unfractionated heparin

• PPH: postpartum hemorrhage

Period of pregnancy	Anticoagulation of choice
First trimester	If dose of warfarin < 5mg/day – continue warfarin till 36 we eksIf dose of warfarin > 5mg /day or warfarin embryopath y – replace it with heparin / L MWH
From 12 weeks to 36 weeks	warfarin
At 36 weeks	Switch to LMWH
1 week before EDD / 3 days b efore EDD	Switch to UFH (as LMWH ha ve longer T1/2– chances of P PH)
On the day of delivery	Stop all anticoagulantS
6 hrs after vaginal delivery / 6-12 hrs after cesarean	Restart UFH and warfarin.W hen INR is 2.5 to 3 withdraw UFH & continue warfarin

Solution to Question 9:

This is a young patient with no comorbidities, subjected to an electrolyte poor fluid for irrigation. Since the deficit is 500 mL, it is recommended to continue the surgery with careful monitoring of fluid status.

The criteria for terminating the procedure for

• Electrolyte deficient fluid is—- a deficit of 1 L

• Electrolyte rich fluid is—-- a deficit of 2.5 L

Glycine irrigation causes hyponatremia. The earliest symptom of hyponatremia is nausea which typically requires fluid absorption (deficit) of at least 1000 mL in women undergoing surgical procedures.

There are two types of distension medias

Electrolyte deficient fluid	Electrolyte rich media
1.5% glycine3% sorbitol5% mannitol	Ringer lactateNormal saline
Instrumentsthat can be used-monopolar / bipolar instrume nts	Only bipolar instruments

Electrolyte deficient fluid	Electrolyte rich media
Complications:Water intoxication Hyponatremia (except mannitol)	Causes Shock & burns when used with monopolar instruments

Solution to Question 10:

Morphology of sperms is the most important marker of male fertility.

While sperm concentration and progressive motility have value for distinguishing infertile men, strict sperm morphology is the most important discriminating value compared toother parameters.

Solution to Question 11:

The given clinical scenario of a 46 XY individual with female internal & external genitalia and poorly developed breasts, presenting with primary amenorrhea points towards the diagnosis of sweyer's syndrome.

Turner syndrome (option B) is a disorder of 45 XO individuals with short stature. The gonads in these individuals are streak ovaries.

Androgen Insensitivity Syndrome (option A) is a disorder of 46 XY individuals with complete insensitivity to testosterone. These individuals have female external genitalia with an absent uterus. Secondary sexual characteristics like the development of breasts are adequate. However,pubic and axillary hairs are scanty.

True hermaphrodite (option C) can be a 46 XX or 46 XY individual with both ovaries and testis. These individuals have both male and female internal genitalia with ambiguous external genitalia.

Sweyer's syndrome is a type of male pseudohermaphroditism where there is an inactivation of the SRY gene that leads to formation of dysgenetic testis.

Thus,in the absence of hormones; AMH & testosterone the mullerian duct grows and the wolffian duct regresses forming female internal genitalia such as uterus (infantile type).

These patients also lack the DHT hormone and hence present with poorly developed breasts (tanner staging 1 or 2), scanty pubic,axillary hair and female external genitalia. These individuals are of normal stature with increased LH & FSH.

Management includes gonadectomy as soon as diagnosed and estrogen + progesterone for breast development.

AMH: Anti mullerian hormone

DHT: dihydro testosterone

Solution to Question 12:

The given scenario of decreased sperm concentration with normal range of motility and morphology points to the diagnosis of oligospermia.

Oligospermia is defined as sperm concentration of <15 million/ml.

It is further subdivided into mild-moderate oligospermia when the concentration is 10-15 million sperms /ml & severe oligospermia when the concentration is <5 million sperms/ml.

Teratozoospermia or teratospermia (option B) is defined as Increased proportion of morphologically abnormal sperms.

Asthenozoospermia or asthenospermia (option C) is used to define sperms with diminished motility.

Term	definition
azoospermia	Semen with absent sperms
aspermia	Both semen and sperms are a bsent
leukocytospermia	Increased WBC count in the semen
necrospermia	Nonviable sperms

Solution to Question 13:

The position of the patient in the Cesarean section is supine, with a wedge under the right hip.

The wedge beneath the right hip creates a left lateral tilt to aid venous return and avoid hypotension.

Solution to Question 14:

Fibronectin detection in cervicovaginal secretions before membrane rupture is a possible marker for impending preterm labor.

Fetal fibronectin:

• Fetal fibronectin is a glycoprotein present in high concentrations in maternal blood and in amniotic fluid.

• Fetal fibronectin levels are measured using ELISA, and values exceeding 50 ng/mL are considered positive.

Screening tests in pregnancy:

1st Trimester:

Dual test + nuchal translucency

DUAL test	Down's syndrome	Trisomy 18
PAPP-A (pregnancy associated plasma protein A)	↓	↓
β-HCG	↑	↓

2nd Trimester:

Integrated screening:

Dual screening + nuchal translucency + quad test

Quadraple test	Down's syndrome	Trisomy 18
α feto protein	↓	↓
β-HCG	↑	↓
Estriol (unconjugated E3)	↓	↓
Inhibin A	↑	↓

Solution to Question 15:

The given procedure is cardiotocography (CTG; electronic fetal monitor).

CTG, is a type of intrapartum fetal monitoring where the fetal heartbeat and the uterine contractions are recorded simultaneously.

The CTG graph can be differentiated from non stress test (NST) graph by the presence of two graphical representations in CTG:

• FHR represented by the upper graph

• Uterine contractions represented by the lower graph.

NST has a single graphical representation.

A normal CTG graph is defined by the following parameters:

• Baseline FHR-110-160

• Beat to beat variability of 5-25 bpm

• Two or more Acceleration during 20 minute period

• No late deceleration

• No variable deceleration

• Early deceleration +/-

The below given depicts a CTG graph



The below given image shows a NST graph



Solution to Question 16:

The next step of management in this patient, is to do USG to measure the cervical length.

In the given clinical scenario the patient has undergone conization which is a risk factor for cervical incompetence. Hence, it is important to measure the cervical length to diagnose cervical incompetence in this patient. If the cervical length is ≤2.5 cm then a cervical cerclage can be performed.

Cervical cerclage (Option B) is performed as the next step in management when the patient presents with ≥2 second trimester pregnancy losses (History based diagnosis of cervical incompetence). Since, the history in this patient is of preterm labour, the first step would still be

to do a TVS to measure the cervical length for diagnosing. Cerclage only comes after diagnosis.

Cervical incompetence is one of the important causes second trimester (mid trimester) abortions.

Risk factors for the incompetence include:

• Past h/o surgery on cervix such as conization, loop electrosurgical excision procedure (LEEP)

• Cervical trauma during labour, instrumental delivery

Diagnosis of cervical incompetence in a pregnant female:

• H/O ≥2 second trimester abortions (history based diagnosis)→ management is by cervical cerclege+progesterone

• H/O 1 second trimester abortion→ TVS done between 18-24 weeks to measure the cervical length→USG based diagnosis of cervical incompetence is made if there is h/o 1 or more second trimester abortions with TVS showing cervical length ≤2.5 cm.

TVS findings in cervical incompetence include:

• Cervical length ≤2.5 cm (Normal:3-4cm)

• Shape of cervix becomes U shaped

• Diameter of internal os ≥ 2cm

Solution to Question 17:

A finding of endometrial thickness ≥ 4 mm (according to ACOG) in a postmenopausal women requires further evaluation, such as endometrial biopsy.

If the endometrial thickness is ≤4 mm in a post menopausal female, malignancy is rare and hence, can be managed by tranexamic acid. If the bleeding still persists, the patient needs to undergo fractional curettage and hysteroscopy.

The algorithm for next step for abnormal uterine bleeding in different age groups is as follows:

Age group	Endometrial thickness on TVS	Action
Age <40years	>12mm	Endometrial sampling/Endometrial biopsy
Age>40 years or Age >45 years	-	Endometrial sampling/Endometrial biopsy is done irrespective of usg findings
Post menopausal women	≥4mm	Endometrial sampling/Endometrial biopsy
Post menopausal women	<4mm	Reassurance+tranexamic acidIf, bleeding still persists then do endometrial biopsy



Solution to Question 18:

In the event of missed periods while on contraceptives, Beta-HCG estimation should be done as the next step to rule out pregnancy.

Serum beta HCG can detect a pregnancy as early as 5 days before the first missed period. This can be done by either serum beta hcg estimation or by urine pregnancy test which estimates the beta-hcg excreted in urine. If the patient is not pregnant she can continue taking the combined pill.

Mala N is a combined oral contraceptive pill that is distributed for free by the Indian government. It contains 0.15 mg levonorgestrel and 30 µg ethinyl estradiol. It contains 21 contraceptive pills + 7 iron tablets. It is started on the first day of each cycle.

The combined oral pills act by suppressesing pituitary hormones and preventing ovulation. The progestogen also causes atrophic changes in the endometrium and prevents implantation. It also causes thickening of the cervical mucus, making it difficult for the sperms to penetrate.

The menstrual side efects includes breakthrough bleeding, menorrhagia and rarely post-pill amenorrhea when the patient is on drug for more than 6 months. Post pill amenorrhea is seen in <1% of patients and should be considered only after ruling out pregnancy.

Pregnancy rate with the combined oral pill is 0.1 per 100 woman-years.

Solution to Question 19:

The reason for the postpartum hemorrhage in the given patient is likely due to
the succenturiate lobe being torn away and left behind at the time of placental delivery, as seen with cases of Placenta succenturiata.

Placenta succenturiata refers to the abnormal placenta where an additional lobe/lobes of placental tissue are located a few centimeters away. The ancillary lobes are smaller in size when compared with the main placental tissue (note: Bilobed placenta has two lobes of equal sizes).These lobes have vessels that course through the membranes connecting them to the main placental tissue.

Placenta succenturiata can be associated with complications like vasa previa (fetal hemorrhage), postpartum hemorrhage due to retained placental tissues, infection such as endometritis, Subinvolution of uterus and Placental polyp formation.

An image of this variant is given below where L1,2,3 depicts smaller lobes, UC is the umbilical cord, and M is the intervening membrane.



The below given image shows a Bilobed or Bipartite or duplex placenta which is different from placenta succenturiata. The lobes in bilobed placenta are nearly equal in size. The umbilical cord is attached into a connecting chorionic bridge or into the intervening membranes in between the two placental lobes.

Other options:

Option A: Placenta membranacea is an unduly large and thin placenta that develops from both the chorionic frondosum and the chorion leave. It is associated with an increased risk of placenta previa or accreta and retained placenta and its complications.

Circummarginate placenta (option C) and Circumvallate placenta (option D) are a type of extrachorial placentation.

In extrachorial placentation, the chorionic plate is smaller than the basal plate.The chorionic plate does not extend upto the placental margin. The membranes (amnion and chorion) are rolled back upon themselves to form a ring leaving a rim of bare placental tissue (extrachorial portion)

Circummarginate placenta-A thin fibrous ring is present at the margin of the chorionic plate where the fetal vessels appear to terminate.

Circumvallate placenta - The fetal surface is divided into a central depressed zone surrounded by a thickened ring which is composed of a double fold of amnion and chorion. Vessels radiate from the cord insertion as far as the ring and then disappear from the view.

Complications associated with circumvallate placenta are preterm delivery, abortions, growth retardation, antepartum hemorrhage, retained placenta, and hydrorrhea gravidarum.

A specimen of circumvallate placenta is shown below:



Solution to Question 20:

HPV nonavalent vaccine (Gardasil 9 or HPV9) is active against HPV 6,11,16,18,31,33,45,52,58.

Currently, HPV nonnavalent vaccine has replaced Gardasil which is a quadrivalent vaccine acting against 6,11,16,18.

Vaccines available for HPV:

Brand name	Cervarix	Gardasil	Gardasil-9	Cervavac
Characteristic and HPV subtypes	Bivalent:16,18	Quadrivalent:6,11,16,18	Nonavalent:6,11,16,18,31,33,45,52,58	Quadrivalent:6,11,16,18
Protects against:	CINCA cervix	Anogenital wartsCINCA cervix	Anogenital wartsCINCA cervixVulvalintraepithelialneoplasiaVaginalintraepithelialneoplasiaAnal cancerOropharyngeal cancerHead and neck cancers	Anogenital wartsCINCA cervix
Dose	0.5 ml,IM.	0.5 ml,IM.	0.5 ml,IM.	0.5 ml,IM.
Ideal Age range:	Ideally at 11-12 years or9-26 years	Ideally at 11-12 years or9-26 years	Ideally at 11-12 years or9-26 years	The targeted age group is 9-26 years

Solution to Question 21:

Among the given gynecological malignancies, sentinel lymph node biopsy (SLNB) is most useful in vulval carcinomas.

This is because women with earlier stages of vulval cancer can have diseased lymph nodes even without palpable enlargement of nodes. Hence, sentinel lymph node biopsy helps in early diagnosis lymph node spread which can be managed by lymph node dissection.

A sentinel lymph node is a physiologically, the first lymph node to receive tumor lymphatic drainage.

Malignancies where sentinel lymph node biopsy is commonly done are :

• Vulval carcinoma

• Breast carcinoma

• Penile carcinoma

• Malignant melanoma

Lymphatic mapping is similar technique to SLNB that can also be used in vulval cancers.

Solution to Question 22:

As per the latest guidelines, uterine massage is no longer a core component of Active Management of the Third Stage of Labor (AMTSL) that is performed for the prevention of postpartum hemorrhage (PPH).

Uterine massage or fundal massage is not recommended as an intervention to prevent PPH in women who have already received prophylactic uterotonics such as oxytocin (core component of AMTSL). Hence, it is now an optional component of AMTSL that is done in patients with uterine atony as a part of the management of PPH.

AMTSL is recommended by the WHO to prevent postpartum hemorrhage (PPH). Uterotonics are the most important components in AMTSL to do so.

The currently recommended components of AMTSL are as follows:

Note: In the event that uterine atony is diagnosed during the assessment of uterine tone, uterine massage can be performed to control PPH.

1.	Injectinguterotonicswithin1 minuteof delivery of the baby or immediately after delivery of theanterior shoulder	Most importantstepOxytocin 10 IU im/iv infusion is thepre ferred uterotonic.(Not recom mendedby WHO-carboprost)
2	Delayed cord clamping	Done within1-3 minsof delive ry80 mlof additional blood g oes to the fetusPreventsfetal anemia
3	Delivery of placenta by contr olled cord traction: Modified Brandt Andrews technique	The uterus is pushed upward s and backwardThe traction o f cord is downwards and forw ards
4	Intermittent assessment of ut erine tone	Done every 15 mins for 2 hou rs

Solution to Question 23:

Quadruple screening is a second-trimester multiple marker screening test used for the detection of chromosomal anomalies like Down syndrome. It is performed between 15-21 weeks.

Pregnancies with Down syndrome are characterized by elevated β-hCG, elevated Inhibin A, reduced estriol and reduced AFP levels.

First-trimester aneuploidy screening combines two maternal serum analytes (Dual screening test) with a sonographic measurement. It is performed between 11-14 weeks.

An increased nuchal translucency indicates an increased possibility of Down's syndrome.

	Trisomy 21	Trisomy 18	Neural tube defect or abdomi nal wall defect
MSAFP	Low	Low	High
β hCG	High	Low	Normal
Unconjugated estri ol	Low	Low	Normal
Inhibin A	High	Low	Normal

Dual test	Down's syndrome	Trisomy 18 & Trisomy 13
PAPP-A (pregnancy associated plasma protein A)	↓	↓
β-HCG	↑	↓

Solution to Question 24:

Increased number of theca lutein cysts is not an indicator of choriocarcinoma. Increased size or persistence of theca lutein cysts indicates choriocarcinoma.

Plateau of high serum β-hCG level for four measurements for ≥3 weeks is an indicator of choriocarcinoma.

Risk factors for malignant change are:

• Molar pregnancy - complete moles have 15-20% risk, partial moles have 1-5% risk

• Patient's age ≥ 40 years or <20 years

• Parity ≥3

• Serum β hCG >1,00,000 IU/mL

• Uterine size >20 weeks

• Previous history of molar pregnancy

• Large theca lutein cysts >6 cm in diameter

Post-evacuation uterine sonographic surveillance can be used as a predictor of subsequent neoplasia. Findings indicative of malignant transformation are:

• Suburethral nodules

• Myometrial nodules

• Hypervascularity.

Solution to Question 25:

During the follicular phase of the menstrual cycle, estrogen is secreted in high amounts by the ovarian granulosa cell.

Follicular phase (days 0–14)

• A primordial follicle develops to the Graafian stage, with atresia of neighboring follicles.

• LH and FSH receptors are up-regulated in theca and granulosa cells.

• Estradiol levels increase and cause the proliferation of the uterus.

• FSH and LH levels are suppressed by the negative feedback effect of estradiol on the anterior pituitary.

• Progesterone levels are low.

Ovulation (day 14)

• Occurs 14 days before menses, regardless of cycle length.

• A burst of estradiol synthesis at the end of the follicular phase has a positive feedback effect on the secretion of FSH and LH (LH surge).

• Ovulation occurs as a result of the estrogen-induced LH surge.

• Estrogen levels decrease just after ovulation (but rise again during the luteal phase).

• Cervical mucus increases in quantity; it becomes less viscous and more penetrable by sperm.

Luteal phase (days 14–28)

• The corpus luteum begins to develop, and it predominantly synthesizes progesterone. Vascularity and secretory activity of the endometrium increase to prepare for receipt of a fertilized egg.

• Basal body temperature increases because of the effect of progesterone on the hypothalamic thermoregulatory center.

• If fertilization does not occur, the corpus luteum regresses at the end of the luteal phase. As a result, estradiol and progesterone levels decrease abruptly.

Menses (days 0–4)

• The endometrium is sloughed because of the abrupt withdrawal of estradiol and progesterone



Solution to Question 26:

The given image shows a pneumatic compression device. It is used in women who are not on any thromboprophylaxis and planned for cesarean section (statement A) and it is discontinued once the patient starts ambulating post-surgery (statement C).

The device is used because pregnancy and surgery increase the risk of thromboembolism.

According to recent guidelines, patients with no additional risk factors and not placed on any pharmacological thromboprophylaxis are placed on pneumatic compression devices before cesarean delivery. Patients undergoing cesarean delivery with additional risk factors are put on both pneumatic compression devices and pharmacological therapy (UFH or LMWH). UFH-unfractionated heparin, LMWH-lower molecular weight heparin.

Indications of pharmacological thromboprophylaxis:

• Inherited coagulation disorder

• Overweight or obese patient

• Heart failure

• Family history of deep vein thrombosis or pulmonary embolism

Solution to Question 27:

The development of uterine synechiae is the most likely cause of amenorrhoea in the given clinical scenario.

Certain types of spontaneous abortions such as inevitable, incomplete and missed abortions might require medical assistance for complete removal of products of conception including procedures such as dilatation and curettage.

Vigorous uterine curettage in early pregnancy damages the basal layer of endometrium leading to scarring and adhesions in the uterus that interferes with normal endometrial growth and shedding.This causes the clinical feature of secondary amenorrhea as seen in this patient.

This intrauterine adhesions are known as uterine synechiae or Asherman syndrome.

As there is no defect in the hypothalamo-pituitary-ovarian axis in the asherman syndrome, the levels of FSH remain within the normal range of 4 - 25 IU/L as seen in this patient.

In the case of Pituitary dysfunction (option A) the FSH levels are usually decreased.

In ovarian dysfunction (option B) the negative feedback of ovarian hormones (estrogen) on pitutary and hypothalamus is lost. Hence, the levels of FSH is increased (≥ 25 IU).

Asherman syndrome is characterised by a spectrum of intra uterine scarring including thin adhesions, dense bands, or complete obliteration of the uterine cavity. It occurs due to endometrial damage following vigorous curettage, usually in miscarriage, or elective abortion. Damage may also occur from other uterine surgeries such as myomectomy or cesarean delivery or from an infection related to an intrauterine device.

Depending on the degree of scarring, patients may present with amenorrhea, hypomenorrhea or recurrent pregnancy loss due to inadequate placentation.

Asherman syndrome shows multiple filling defects on hysterosalpingography (HSG) or saline-infusion sonography. Definitive diagnosis requires hysteroscopy. Hysteroscopic lysis of adhesions is the preferred surgical treatment for the syndrome.

Solution to Question 28:

Simple hysterectomy is the best treatment in a case of atypical endometrial hyperplasia in a peri-menopausal and post-menopausal female.

Endometrial hyperplasia with atypia carries 29% risk of developing into endometrial carcinoma and hence, must be treated with total abdominal hysterectomy (simple hysterectomy).
The presence of any concomitant endometrial cancer must be ruled out by doing fractional curettage (or dilatation and curettage) before doing a hysterectomy.

If a premenopausal woman does not desire to undergo hysterectomy, progesterone can be given and the patient is followed up every 3-6 months. If the atypia persists during the follow-up hysterectomy should be performed.

The table below summarizes the management of patients with endometrial hyperplasia

Management of Endometrial hyperplasia	
Hyperplasia without atypia	
Pre-menopausal women	Progesterone therapy
Postmenopausal women:Simple hyperplasia without atypiaComplex hyperplasia without atypia	Follow up without therapyProgesterone therapy
Hyperplasia with atypia	
Pre-menopausal women	If patient wishes to preserve uterus,give High-dose progesterone therapyIf atypia persists, then perform Total abdominal hysterectomy
Peri-menopausal women	Total abdominal hysterectomy
Post-menopausal women	Total abdominal hysterectomy +BSO (bilateral salphingo-ophorectomy)

Solution to Question 29:

Among the given options, Laparoscopic tubal ligation is an irreversible method of contraception.

Tubal sterilization can be achieved by three methods which are:

• Electrocoagulation is used for the destruction of a segment of the tube

• Mechanical occlusion of the tube using a silicone rubber band (Falope ring) or a spring-loaded clip (Hulka Clemens clip)

• Suture ligation with or without tubal segment excision such as Pomeroy's method, Uchida's method, Irving's method, etc.

Intrauterine devices such as copper T and LNG-IUDs and implants belong to the category of long acting reversible contraception.

In copper-T (Option B) a copper wire is wrapped around a polypropelene vertical frame. The copper in the device causes an intense local inflammatory response in the uterus that leads to lysosomal activation and activation of other inflammatory processes that are spermicidal. The Cu-T 380A is approved for 10 years of continuous use.

LNG-IUD (Option D) is a T-shaped polyethylene structure enclosed by a cylinder containing levonorgesterl. This hormone is released continuously through the permeable cylinder. The progestin makes the endometrium atrophic and stimulates the production of thick cervical mucus that blocks sperm penetration into the uterus. It also decreases tubal motility, thereby preventing fertilization.

Nexplanon has now replaced implanon (Option C) which was a previously used etonogestrel implant. Nexplanon is a single-rod progestin etonogestrel (68 mg) containing subdermal implant that provides contraception over 3 years. Its mechanism of action is by suppression of ovulation, thickened cervical mucus, and endometrial atrophy. Irregular bleeding is an important side effect of the implant. This method of contraception is reversible in nature.

Image below shows implanon:





Solution to Question 30:

Cystoscopy is the most useful test and also the investigation of choice for vesicovaginal fistula (VVF).

Cystoscopy or cystourethroscopy helps to visualize the size of the fistulas, the exact location of multiple fistulas and their proximity to the ureteral orifice. It can also assess the bladder mucosa for any injuries.

The 3-swab test (Option B) is done to identify the types of genitourinary fistula.

In this test, three sterile pieces of cotton swabs are serially placed in the vaginal cavity, while 50-100ml of methylene blue solution is instilled in the bladder using a catheter. The cotton swabs are placed one at the vault, one at the middle, and one just above the introitus.

Interpretation:

• Uppermost swab gets wet and not stained - the leak is from the ureter, as seen in ureterovaginal fistula

• Middle swab gets stained - the leak is from the bladder, as seen in vesicovaginal fistula

• Lowermost swab gets stained- the leak is from the urethra,as seen in urethrovaginal fistula

The below image shows the 3-swab test.



Option C: MRI is usually not needed for the diagnosis of VVF. They can be used to study vesicouterine fistulas.

Option D: Cystometry is done to measure the capacity of the bladder. It is used to investigate urge incontinence.

Solution to Question 31:

Colposcopy with LEEP would be appropriate management for CIN III.

Loop electrosurgical excision procedure (LEEP) is an opd procedure done under colposcopic view and local anesthesia.Here, a low voltage current is passed through the loop and a cone of tissue is removed from the cervix. This procedure causes minimal bleeding and the tissue removed is sent for histopathological examination (HPE). This is the treatment of choice for CIN 2&3 of any age and parity.

Solution to Question 32:

The most appropriate time for the delivery in this patient would be 39 weeks as she has good glycemic control with insulin.

The timing for delivery in a pregnant woman with GDM would vary depending on the control of glucose levels & mode of treatment.

Appropriate timing for delivery in gestational diabetes	
Well-controlledwith diet	39 weeks to 40 weeks +6 days

Appropriate timing for delive ry in gestational diabetes	
Well-controlledwithinsulin/or al agent	39 weeks to 39 weeks + 6 day s
Co-morbidityor suboptimal c ontrol	Induction before 39 weeks
Withpoor glycemic control	Reasonable to deliver at 37 to 38 weeks + 6 days of gestati onIn extreme cases where all efforts for glycemic control h ave failed induction prior to 37 weeks (no clear cut guideli ne)

Solution to Question 33:

The given case scenario is suggestive of fetal compromise. The management in this patient is immediate termination of pregnancy. Since this pregnancy has crossed 34 weeks, it can be terminated by cesarean section.

Abnormal flow patterns in the umbilical artery:

• Reduced end-diastolic flow indicates uteroplacental insufficiency (S/D ratio increases)

• Absent end-diastolic flow suggests fetal compromise, hence the pregnancy is terminated by C-section if >34 weeks.

• Reverse end-diastolic flow suggests impending doom of death, hence the pregnancy is terminated by C-section if >32 weeks.



Solution to Question 34:

Among the given options, the presence of third heart sound (S3) is normal in pregnancy.

The third heart sound (S3) occurs due to rapid diastolic filling that is seen secondary to the physiological hypervolemia that occurs during the pregnancy.

Other normal auscultatory findings include:

• First heart sound is loud with an exaggerated split

• Second heart is loud but normal

• Third heart sound is loud and easily heard

• Systolic murmur of grade<3/6

• Mammary murumur or continous machinery murmur

• Apex beat shifted to 4th intercostal space

Solution to Question 35:

The final switch to heparin in a pregnant women is advised at 36 weeks of pregnancy and is continued till 7 days postpartum.

This switch is made from warfarin to heparin at 36 weeks, as warfarin is a strong anticoagulant which can cause post partum hemorrhage in the mother at the time of labour and can also cross the placenta into the fetal blood and can increase the risk of intracranial hemorrhage in the fetus

The below given algorithm is followed for administration of anticoagulants in a pregnant female:

LMWH- low molecular weight heparin

UFH- Unfractionated heparin

Period of pregnancy	Anticoagulation of choice
First trimester	If dose of warfarin< 5mg/day –continue warfarin till 36 we eksIf dose ofwarfarin > 5mg/ day or warfarin embryopathy present –replace it with hepa rin / LMWH
From12 weeks to 36 weeks	warfarin
At 36 weeks	Switch toLMWH
1 week before EDD / 3 days b efore EDD	Switch toUFH(as LMWH hav e longer T1/2– increased cha nces of PPH)
On the day of delivery	Stop all anticoagulant
6 hrs after vaginal delivery / 6-12 hrs after cesarean	Restart UFH and Warfarin. When INR is 2.5 to 3 withdra w UFH & continue warfarin

Obstetrics and Gynecology AIIMS 2019 (May & Nov)

Question 1:

A 40-year-old woman comes with a history of postcoital bleeding and a visible mass seen in the cervix per speculum. The next step to be done:

a) Conization

b) Cryotherapy

c) Targeted biopsy

d) Pap smear

Question 2:

In preterm labor, which drug has a neuroprotective role?

a) $MgSO_4$

b) Nifedipine

c) Ritodrine

d) Isoxsuprine

Question 3:

The correct statement regarding the physiological changes in the respiratory system in pregnancy?

a) Minute volume decreases by 50%

b) Residual volume increases

c) FRC decreases

d) Total lung capacity increases

Question 4:

On finding a congenital heart block in a fetus, which of the following conditions would you evaluate for in the mother?

a) APLA syndrome

b) Congenital heart defects

c) SLE

d) Hemolytic anaemia

Question 5:

A 33 Weeks multigravida woman presents with epigastric pain, headache, visual symptoms, and proteinuria - 3+. The next step of management would be?

a) Immediate c - section

b) Dexamethasone for fetal lung maturity

c) Induction - vaginal delivery

d) $MgSO_4$ IV

Question 6:

All of the followings are seen in GDM except:

a) Malformations

b) Polyhydramnios

c) Fetal Macrosomia

d) Neonatal Hypoglycemia

Question 7:

Hormone replacement therapy is not indicated for:

a) Hot flashes

b) CAD Prophylaxis

c) Osteoporosis

d) Premature menopause

Question 8:

The risk of fetal varicella syndrome is maximum when the mother is infected between _____.

a) 6 to 10 weeks

b) 8 to 12 weeks

c) 13 to 20 weeks

d) 20 to 28 weeks

Question 9:

The loading dose of MgSO4 in pre-eclampsia is?

a) 4 ml Mgso4 (50% w/v) in 16 ml NS

b) 8 ml MgSo4 (50% w/v) in 12 ml NS

c) 12 ml MgSo4 (50% w/v) in 8 ml NS

d) 10 ml MgSO4 (50% w/v)

Question 10:

Drug of choice for treatment of pregnant woman with bacterial vaginosis?

a) Metronidazole

b) Flucanazole

c) Clindamycin

d) Erythromycin

Question 11:

A 40-year-old female presented with a history of profuse vaginal discharge. The discharge has a fishy odour. Which of the following best describes the microscopic finding given below?



a) Epithelial cell with bacteria attached on cell surface

b) Bacteria engulfed by macrophages

c) Neutrophil englufed bacteria

d) Bacteria arranged in clusters

Question 12:

The following instrument is used for _____

a) VIA

b) VILI

c) PAP smear

d) Colposcopy

Question 13:

Anti-D vaccine is not given in which of the following conditions?

a) Amniocentesis at 16 weeks

b) Manual removal of placenta

c) Intrauterine transfusion at 28 weeks

d) MTP at 63 days

Question 14:

What does this tocograph show?



a) Deceleration

b) Acceleration

c) Normal

d) Deceleration followed by acceleration

Question 15:

A 16-year-old unmarried female has come to the gynecology OPD for cervical cancer vaccination. Which vaccine is preferred?

a) Gardasil

b) Biovac

c) Rabivac

d) Tvac

Question 16:

Which of the following agents cause hydrops fetalis?

a) CMV

b) Parvovirus

c) Toxoplasma

d) HIV

Question 17:

The given instrument is used in:

a) Vaginal hysterectomy

b) Cesearean section

c) Manchester operation

d) Fothergill's surgery

Question 18:

A case of gestational trophoblastic neoplasm is detected to have metastasis in the lungs. It is considered as stage _____.

a) I

b) II

c) III

d) IV

Answer Key

Question No.	Correct Option
1	c
2	a
3	c
4	c
5	d
6	a
7	b
8	c
9	b
10	a
11	a
12	c
13	c
14	a
15	a
16	b
17	b
18	c

Detailed Explanations

Solution to Question 1:

A targeted biopsy is the next line of management in the patient with a visible cervical mass and postcoital bleeding. A pap smear is done when there are symptoms with no identifiable lesion.

Other options:

Option A: Conization is an invasive procedure where the cervix is cut in a cone-shaped manner. It is both diagnostic and therapeutic. It is indicated for diagnosis in women with HSIL or adenocarcinoma in situ and may be considered under the following conditions:

• The limits of the lesion cannot be visualized with colposcopy.

• SCJ (squamocolumnar junction) is not seen in colposcopy.

• Colposcopy is positive for CIN 2/CIN 3.

• Lack of correlation between cytology, biopsy, and colposcopy.

• Suspected micro invasion in cytology, colposcopy, or biopsy.

• Colposcopist is unable to rule out invasive cancer.

Hence it is very clear that conization is not indicated for all symptomatic women and it should be done following either a PAP, colposcopy or a biopsy with an abnormal or inconclusive result.

Option B: Cryotherapy is ideal only for small superficial lesions. It is indicated for biopsy-proven lesions. Cryotherapy is appropriate for CIN 1 which has persisted for more than 24 months and is confined to the ectocervix.

Option D: The false negative rate of pap smear in the presence of invasive cancer is 50% and a negative pap test should never be relied on in a symptomatic patient, especially one with a visible lesion.

CIN: Cervical intraepithelial neoplasia, HSIL: High squamous intraepithelial lesion, LSIL: Low squamous intraepithelial lesion

Solution to Question 2:

In preterm labor, $MgSO_4$ (magnesium sulfate) is considered to have a neuroprotective role. Studies have demonstrated that magnesium exposure (compared with no exposure) significantly lowers the risks for cerebral palsy.

The combination of corticosteroids and magnesium sulfate has been associated with lower rates of neurodevelopmental impairment and death compared to corticosteroids alone.

The proposed mechanisms of action include:

• reduced presynaptic release of glutamate

• blockade of glutamatergic NMDA receptors

• potentiation of adenosine action

• improved calcium buffering by mitochondria

• blockage of calcium entry via voltage-gated channels

$MgSO_4$ is used as an anticonvulsant in the management of eclampsia and as prophylaxis in severe preeclampsia. The therapeutic level of magnesium in eclampsia is 4–7 mEq/L. Measurement of serum magnesium levels is indicated to monitor magnesium toxicity (if serum creatinine > 1.0 mg/dL).

Loss of patellar reflex is the first sign of $MgSO_4$ toxicity and occurs when the plasma magnesium level reaches 10 mEq/L. Elevated magnesium levels above 10 mEq/L weaken breathing, while 12 mEq/L or higher levels can lead to respiratory paralysis and arrest. Other signs of magnesium toxicity include urine output <30 mL/h, chest pain, and heart block visual changes, flushing, muscle paralysis, and drowsiness.

In suspected toxicity, $MgSO_4$ therapy should be stopped and a slow IV infusion of 1g calcium gluconate 10 mL (10% solution) is given.

Other options:

Options B, C, and D: Nifedipine (calcium channel blocker), and ritodrine, isoxsuprine (beta-adrenergic agonists) are used as tocolytics in preterm labor.

Solution to Question 3:

Functional residual capacity (FRC) decreases by 20 to 30 percent (400 to 700 mL) during pregnancy.

FRC = Expiratory Reserve Volume (ERV) + Residual Volume (RV). ERV decreases by 15 to 20 percent (200 to 300 mL). RV decreases by 20 to 25 percent (200 to 400 mL).

FRC and RV decline due to diaphragm elevation. Significant reductions occur by the sixth month with a progressive decline across pregnancy.

The increased minute ventilation during pregnancy is influenced by various factors. These include the stimulatory action of progesterone, reduced expiratory reserve volume, and compensatory respiratory alkalosis. Total lung capacity (FRC + IC) decreases in pregnancy.

Anatomical changes such as the elevation of the diaphragm, an increase in the subcostal angle, and an expansion of the transverse diameter of the thoracic cage are seen.

Respiratory changes in pregnancy are given below:

(Mnemonic: TIMM increases in pregnancy)

Increase	Decrease	Unaffected
Tidal volume	Functional residual capacity	Respiratory rate
Inspiratory capacity (IC)	Expiratory reserve volume	Vital capacity
Minute O2uptake	Residual volume	Inspiratory reserve volum e
Minute ventilation	Total lung capacity	Lung compliance
Peak expiratory flow rate and airway conductance	Total pulmonary resistance	

Solution to Question 4:

On finding a congenital heart block in the fetus, the mother should be evaluated for systemic lupus erythematosus (SLE).

Fetal and neonatal heart blocks occur due to myocarditis and fibrosis in the region between the AV node and the bundle of His. Congenital heart block primarily develops in fetuses of women with antibodies to the SS-A or SS-B antigens, and up to 50 percent of women with SLE test positive for these antibodies. So in pregnancies with either of these antibodies, fetal cardiac monitoring is performed between 18 and 26 weeks of gestation.

The cardiac lesion is permanent, and in most cases, a pacemaker is required. Maternal hydroxychloroquine therapy is associated with a reduced incidence of fetal heart block, particularly in women who have previously had an affected child.

Neonatal lupus erythematosus is thought to be caused by the transplacental passage of maternal antibodies. It is characterized by skin lesions known as lupus dermatitis in newborns, as well as various hematological and systemic abnormalities. In some cases, congenital heart block may also occur. Thrombocytopenia and hepatic involvement are also seen.

The most common cutaneous finding in neonatal lupus erythematosus is an erythematous, slightly scaly eruption on the face and periorbital skin (raccoon sign/owl eye/eye mask). The eruption can be exacerbated by UV exposure. The rash improves over the first few months of life and usually resolves without scarring by 12 months of age.

Avoidance of sunlight should be advised and low■potency topical steroids may be of benefit.

The pregnancy outcomes are best in women with:

- Dormant lupus activity for at least 6 months before conception
- No lupus nephritis manifestation in the form of proteinuria or renal dysfunction
- Absent anti-phospholipid syndrome or lupus anticoagulant
- Superimposed preeclampsia does not develop

Solution to Question 5:

The given clinical scenario of a 33 weeks pregnant woman with epigastric pain, headache, visual symptoms, and proteinuria 3+ indicates severe preeclampsia. The next step of management would be iv administration of $MgSO_4$ for the prevention of seizures.

Preeclampsia can be classified as severe if any of the following are present:

- BP ≥160/110
- Headache
- Visual disturbance
- Upper abdominal pain

• Elevated creatinine, oliguria

• Thrombocytopenia (<1,00,000 cells per mm3)

• Pulmonary edema

• Marked elevation in serum transaminases

• Fetal growth restriction

Dexamethasone (option B) is also given in severe preeclampsia <34 weeks for fetal lung maturity. But the administration of MgSO4 before corticosteroids would be the next best step for the prevention of seizures.

Immediate delivery (options A and C) is not indicated in this case as the patient doesn't have any contraindication for conservative management. Conservative management is contraindicated in the following cases:

• Persistent symptoms, refractory to treatment or severe hypertension

• Eclampsia, pulmonary edema, HELLP syndrome

• Significant renal dysfunction or coagulopathy

• Placental abruption

• Nonreassuring fetal status

• Fetal demise

• Disseminated intravascular coagulation

• Stroke, myocardial infarction

MgSO4 regimens for management of eclampsia and severe preeclampsia:

Magnesium sulfate is discontinued either 24 hours after delivery or 24 hours after the onset of convulsions (whichever is later).

Regimen	Loading Dose	Maintenance Dose
Intramuscular (Pritchard)	4g (20% solution) IV over 3-5 minutefollowed by 10g (5O %), deep IM(5 gm in each bu ttock)	5g (5O%) IM 4 hourly in the alternate buttocks
Intravenous (Zuspan or Sibal)	4g IV slow over 15-20 minute	1-2g/h IV infusion



Solution to Question 6:

Malformations are not associated with gestational diabetes mellitus (GDM). Malformations occur due to the disruption of organogenesis, usually occurring in the first trimester. The abnormal glycemic status in GDM manifests only in the second trimester.

Overt diabetes (pregestational diabetes) is associated with higher rates of malformations compared to the general population. Neural tube defects (encephalocele, meningomyelocele, anencephaly), transposition of great vessels (TGA), ventricular septal defect (VSD), atrial septal defect (ASD), hypoplastic left heart, aortic stenosis and coarctation of the aorta are commonly seen.

Caudal regression syndrome is rare, but a specific anomaly seen in infants of diabetic mothers. Small left colon syndrome is a rare anomaly that develops in the second and third trimesters.

Other options:

Option B: In GDM, increased blood glucose in the mother leads to fetal hyperglycemia. This can cause polyhydramnios due to increased fetal urine production.

Option C: Fetal hyperinsulinemia causes increased hepatic glucose uptake and glycogen synthesis, accelerated lipogenesis, and augmented protein synthesis resulting in macrosomia

Option D: Fetal hyperglycemia in GDM triggers fetal hyperinsulinemia, resulting in neonatal hypoglycemia

Solution to Question 7:

Hormone replacement therapy is not indicated in coronary artery disease prophylaxis. There is currently insufficient evidence to recommend the initiation or continuation of either estrogen therapy (ET) or estrogen-progestin therapy (EPT) for the prevention of primary or secondary

coronary heart disease (CHD).

In an 18-year reevaluation of WHI (women health initiative) participants, it was found that cardiovascular, cancer, and all-cause mortality rates were essentially the same in both the hormone therapy (HT) and placebo groups.

Currently, hormone therapy (HT) is only indicated for the treatment of bothersome vasomotor symptoms (VMS) and genitourinary symptoms, prevention of osteoporosis, and treatment of hypoestrogenism caused by hypogonadism, surgical menopause, or premature ovarian insufficiency.

Estrogen should not be used in women with the following conditions:

• undiagnosed abnormal genital bleeding

• known, suspected, or history of breast cancer

• known or suspected estrogen-dependent neoplasia

• active or prior venous thromboembolism

• active or prior stroke, myocardial infarction, or transient ischemic attack (TIA)

• known thrombophilia, liver dysfunction, or disease

• untreated hypertension

• known hypersensitivity to the ingredients of the estrogen preparation

• known or suspected pregnancy

Risks of HRT:

• Endometrial cancer

• Breast cancer

• Ovarian cancer

• Thromboembolism

• Lipid profile dysfunction

• Gall stones, liver dysfunction

Solution to Question 8:

The risk of developing fetal varicella syndrome (congenital varicella syndrome) is highest when the mother develops varicella infection between 13 to 20 weeks.

Congenital varicella syndrome occurs in 2% of infants with maternal varicella between 13 and 20 weeks of gestation and in 0.4% of infants before 13 weeks of gestation. It is rare after 20 weeks.

Congenital varicella syndrome is characterized by cicatricial skin scarring in a zoster-like pattern (zig-zag scarring), and limb hypoplasia (shown below). They can also present with neurological abnormalities (microcephaly, cortical atrophy, mental retardation), ophthalmic features (chorioretinitis, microphthalmia, cataract), and renal and autonomic system abnormalities (hydronephrosis, neurogenic bladder). Low birth weight is also common.

Infants born to mothers with varicella in the period from 5 days prior to delivery up to 2 days after delivery are at high risk for severe neonatal varicella. They acquire the infection transplacentally.

Since the mother has not developed a strong antibody response, the infant lacks the protective effect of maternal anti-varicella-zoster virus (VZV) antibodies. Therefore varicella-zoster immune globulin (VZIG) should be administered as soon as possible after birth to newborns whose mothers have varicella 5 days before and up to 2 days after the delivery.

Since perinatally acquired varicella can be potentially life-threatening, the infant should typically be treated with intravenous acyclovir (10 mg/kg every 8 hours) when lesions appear.

For pregnant women with uncomplicated primary varicella or herpes zoster that does not require hospitalization, the recommended treatment is oral acyclovir (800 mg five times daily for seven days).

Solution to Question 9:

The loading dose of $MgSO_4$ is 8ml $MgSO_4$ (50% w/v) in 12ml NS. For loading dose IV infusion of 4g of 20% solution is needed.

One 2mL vial has 1g $MgSO_4$. Therefore, to make a 20% solution containing 4g of $MgSO_4$, we would need to take 8 mL of 50% magnesium sulfate, and dilute it with 12 mL NS. Therefore the final 20mL solution (8+12) would have 4g of $MgSO_4$, making it a 20% solution.

$MgSO_4$ is used as an anticonvulsant in the management of eclampsia and as prophylaxis in severe preeclampsia. The therapeutic level of magnesium in eclampsia is 4–7 mEq/L. Measurement of serum magnesium levels is indicated to monitor magnesium toxicity if serum creatinine >1.0 mg/dL.

$MgSO_4$ regimens for management of eclampsia and severe preeclampsia:

Repeat injections of magnesium sulfate are continued only if the patellar reflex is present, respiratory rate is >12, and urine output is >30 mL/h. Magnesium sulfate is discontinued

either 24 hours after delivery or 24 hours after the onset of convulsions (whichever is later).

Loss of patellar reflex is the first sign of MgSO4 toxicity and occurs when the plasma magnesium level reaches 10 mEq/L. Elevated magnesium levels above 10 mEq/L weaken breathing, while 12 mEq/L or higher levels can lead to respiratory paralysis and arrest.

Regimen	Loading Dose	Maintenance Dose
Intramuscular (Pritchard)	4 g (20% solution) IV over 3-5 minutefollowed by 10 g (5O %), deep IM(5 gm in each bu ttock)	5 g (5O%) IM 4 hourly in the alternate buttock
Intravenous (Zuspan or Sibai)	4g IV slow over 15-20 minute	1-2 g/h IV infusion

Solution to Question 10:

The preferred treatment for bacterial vaginosis in pregnancy is either oral metronidazole 500 mg BD for 7 days (or intravaginal metronidazole for 5 days). Treatment is typically reserved for symptomatic women.

An alternative option is oral clindamycin, 300 mg BD for 7 days (or intravaginal clindamycin ovules for 3 days).

Bacterial vaginosis (BV) is characterized by a decrease in lactobacilli and an overgrowth of anaerobic bacterial species, including gardnerella, prevotella, mobiluncus, and bacteroides. Most women with BV are asymptomatic, but those with symptoms may experience a foul-smelling, thin vaginal discharge.

The clinical diagnosis of BV is based on the presence of three out of four Amsel criteria: elevated vaginal pH (>4.5), thin non inflammatory discharge, >20% clue cells, and a fishy odor (positive whiff test) after adding 10% KOH (potassium hydroxide) to a vaginal sample.

BV is associated with adverse pregnancy outcomes such as preterm birth, preterm premature rupture of membranes, and postpartum endometritis. It also increases susceptibility to sexually transmitted infections, including HIV.

The image below shows a vaginal epithelial cell, with adherent bacteria (clue cell) :



Solution to Question 11:

The given clinical scenario of women with fishy odor vaginal discharge is suggestive of bacterial vaginosis (BV). The microscopic finding is vaginal epithelial cells with bacteria attached to their surface called clue cells, creating a poorly defined cellular border. They are the most reliable indicators of BV.

Bacterial vaginosis (BV) is characterized by a decrease in lactobacilli and an overgrowth of anaerobic bacterial species, including gardnerella, prevotella,mobiluncus, and bacteroides. Multiple or new sexual partners, having female partners, and engaging in oral sex are associated with a higher risk of BV. They may experience a foul-smelling (fishy odor) thin vaginal discharge.

The clinical diagnosis of BV is based on the presence of three out of four Amsel criteria: elevated vaginal pH (>4.5), thin noninflammatory discharge, >20% clue cells, and a fishy odor (positive whiff test) after adding 10% potassium hydroxide to a vaginal sample. creating a poorly defined cellular border.

BV is associated with adverse pregnancy outcomes such as preterm birth, preterm premature rupture of membranes, and postpartum endometritis. It also increases susceptibility to sexually transmitted infections, including HIV.

The recommended treatment options in nonpregnant women include oral metronidazole, metronidazole gel, or clindamycin cream. Alternative options include oral secnidazole, tinidazole, or clindamycin. Partner treatment is not needed.

Solution to Question 12:

The image shown is that of Ayre's spatula which is used for preparing PAP smear. The Papanicolaou test (Pap test) is a screening test for cervical cancer.

The squamocolumnar junction is scraped using Ayre's spatula by rotating it around the cervix. The obtained scrapings are spread evenly onto a glass slide and immediately fixed in a solution of 95% ethyl alcohol and ether. After 30 minutes of fixation, the slide is air-dried and then stained with Pap or a short stain method.

Prerequisites for Pap smear testing:

• Should be done prior to vaginal examination.

• Lubricants should not be used.

• The patient should not be menstruating during the pap test.

• The ideal time is around ovulation.

• The patient should not have had intercourse in the 24hrs prior to the test.

Screening guidelines:

• Time for initiation: 21 years, regardless of the age of first sexual intercourse.

• Women aged 21-29 years: Pap test every 3 years

• Women who are aged 30-65 years: Pap test + HPV testing (co-test) every 5 years or Pap alone every 3 years.

• Women should stop having cervical cancer screening after the age of 65 years. If suspicious PAP, screening till 75 years.

• HIV+ women: annual checkup until three consecutive negative tests followed by testing every 3 years for life.

Instruments used for Pap smear are given below:



Solution to Question 13:

The anti-D vaccine is not given in intrauterine transfusion at 28 weeks. Intrauterine transfusion is done in alloimmunized pregnancies, where the maternal immune system (Rh negative) has already been exposed to the fetal blood (Rh positive).

The purpose of the anti-D vaccine is to prevent the development of maternal antibodies against fetal red cells, which is already occurring in alloimmunized pregnancies.

Anti-D vaccine is administered only in ICT-negative non-isoimmunized pregnancies.

Intrauterine transfusion is done when the peak systolic velocity of the middle cerebral artery (MCA) exceeds 1.5 multiples of the median or if hydrops develop or if the fetal hematocrit is <30%. This indicates that the mother already has antibodies against the fetus's red blood cells, making the administration of anti-D unnecessary.

The indications for administering anti-D are:

• Pregnancy loss: spontaneous or elective abortion (option D) and ectopic pregnancy.

• Fetal death: Regardless of the trimester.

• Procedures: Such as amniocentesis (option A), chorionic villus sampling, fetal blood sampling, and molar pregnancy evacuation.

• Other situations: Antepartum bleeding (including threatened abortion), delivery (vaginal or cesarean), external cephalic version, placental abruption, manual removal of placenta (option B), and trauma to the abdomen during pregnancy.

All Rh D-negative women should be given a single dose of 300 mcg of anti-D at 28 weeks of pregnancy (ICT should be negative).



Solution to Question 14:

The given topograph shows a drop in fetal heart rate below the baseline indicating decelerations. Deceleration is the decrease in fetal heart rate below the baseline by 15 bpm for more than 15s.

Following are the various types of decelerations seen on CTG and their clinical significance:

The CTG given below shows a sinusoidal pattern. It resembles a sine wave. It is often associated with fetal anemia, fetomaternal hemorrhage, and fetal acidosis.

Feature	Early deceleration	Late deceleration	Variable deceleration
The pattern of heart rate fall	The maximum dip of the deceleration corresponds with the peak of the contraction.	Onset, maximum dip, and recovery occur after the onset, peak, and recovery of contraction respectively	Abrupt fall in the heart rate.Decelerations are variable in respect of the size, shape, depth, duration, and timing of uterine contractions.
Time taken to reach the maximum dip of deceleration	>30seconds	>30seconds	<30 seconds
Cause	Fetal head compression	Fetal hypoxia due to maternal hypotension, placental insufficiency, and excessive uterine activity	Umbilical cord compression



Solution to Question 15:

Gardasil is a vaccine administered to provide protection against cervical cancer.

Gardasil is a quadrivalent vaccine providing protection against HPV serotypes 6,11,16 and 18. It can be given to both males and females and provides protection against anogenital warts, CIN (cervical intraepithelial neoplasia), and Ca cervix.

Types of available HPV vaccines:

Cervavac is India's first indigenously prepared quadrivalent HPV (qHPV) vaccine against HPV strains 6, 11, 16, and 18.

As per the December 2022 WHO updated dose schedules on HPV vaccines, the recommendations for different age groups are as follows:

• 1 or 2 doses schedule for the primary target of girls aged 9–14 years.

• 1 or 2 doses schedule for young women aged 15–20 years.

• 2 doses with a 6-month interval for women older than 21 years.

• 3 doses (or at least 2 doses) for immunocompromised individuals/HIV infected.

Brand name	Cervarix	Gardasil	Gardasil-9	Cervavac
Characteristic and HPV subtypes	Bivalent: 16,18	Quadrivalent: 6,11,16, 18	Nonavalent: 6,11,16,18, 31,33,45,52,58	Quadrivalent: 6,11,16, 18
Protects against:	CINCA cervix	Anogenital wartsCINCA cervix	Anogenital wartsCINCA cervixVulval intraepithelial neoplasiaVaginal intraepithelial neoplasiaAnal cancerOropharyngeal cancerHead and neck cancers	Anogenital wartsCINCA cervix
Dose	0.5 ml, IM.	0.5 ml, IM.	0.5 ml, IM.	0.5 ml, IM.

Other options:

Option B: Biovac A is a freeze-dried live attenuated vaccine indicated for the prevention of hepatitis A in persons of age 1 year or older.

Option C: Rabivac is a monovalent inactivated rabies vaccine for cats and dogs.

Option D: Tvac is a Tetanus vaccine.

Solution to Question 16:

Parvovirus B19 is the most common infectious cause of nonimmune hydrops fetalis. Parvovirus B19 infects fetal erythroid progenitor cells and causes anemia. Severe anemia leads to high-output cardiac failure and nonimmune hydrops fetalis.

Hydrops is diagnosed by identifying two or more fetal effusions, such as pleural, pericardial, or ascitic effusions, or the presence of one effusion along with anasarca. Hydrops can be seen in association with red cell auto immunization, which is termed immune hydrops. Otherwise, it is nonimmune.

Nonimmune hydrops is more common than immune hydrops. Causes of nonimmune hydrops include cardiac pathology (Ebstein anomaly), chromosomal (Turner's syndrome), hematological (hemoglobinopathies), lymphatic abnormalities (cystic hygroma), and thoracic abnormalities.

Infectious causes of nonimmune hydrops are parvovirus B19 (most common), syphilis, cytomegalovirus, toxoplasmosis, rubella, enterovirus, varicella, herpes simplex, coxsackievirus, listeriosis, leptospirosis, Chagas disease, and Lyme disease.

The following image shows a stillborn baby with hydrops fetalis:

Solution to Question 17:

The image shown is that of a Doyen's retractor which is used in caesarean sections.

It is used to retract the abdominal wall as well as the bladder for proper exposure of the lower uterine segment during cesarean sections.

Solution to Question 18:

Metastases in the lungs are considered stage III GTN (gestational trophoblastic neoplasia) in FIGO classification.

International Federation of Gynecology and Obstetrics (FIGO) staging for GTN is as follows:

The chest X-ray below shows lung metastasis - stage III GTN.

Stage	Description
I	GTN confined to theuterus
II	GTN extends outside of the uterus but is limited to thegenitalstructures(adnexa, vagina, broad ligament)
III	GTN extends to thelungs, with or without known genital tract involvement

Stage	Description
IV	Allothermetastatic sites i.e. kidney, liver, brain, bowel, etc.



Gestational trophoblastic neoplasia (GTN) refers to the aggressive invasion of the endometrium and myometrium by trophoblast cells. It includes invasive mole, gestational choriocarcinoma, placental site trophoblastic tumors, and epithelioid trophoblastic tumors. GTN is diagnosed based on elevated β-hCG levels. FIGO criteria for diagnosis of GTN:

• Persistence of β-hCG level plateau in 4 measurements for 3 weeks or longer

• Rise in β-hCG levels in 3 or more consecutive measurements done weekly, over a period of 2 weeks or longer

• Elevated β-hCG level for 6 months or more

• Histologic diagnosis of choriocarcinoma

Treatment for low-risk disease is chemotherapy with methotrexate, while those with high-risk disease are treated with etoposide, methotrexate, and dactinomycin alternating with cyclophosphamide and vincristine (EMA/CO) chemotherapy.


FIGO staging of gestational trophoblastic neoplasia
Stage I:
Confined to the uterus
Stage II:
Extends outside the uterus but limited to genital structures
Stage III:
Extends to the lungs, with or without genital tract involvement
Stage IV:
All other metastatic sites
©MARROW

Obstetrics and Gynecology AIIMS 2020 (May and Nov INI-CET)

Question 1:

The following groups of people must be screened for Ca cervix:

a) More than 65 years old

b) Less than 21 years old

c) 21 to 65 year old females

d) Teenagers

Question 2:

Oral contraceptives act by all of the following mechanisms except:

a) Inhibit implantation

b) Cervical mucus thickening

c) Inhibit ovulation

d) Increase GnRH release

Question 3:

A 30-year-old woman is diagnosed with a high-grade squamous intraepithelial lesion (HSIL) of the cervix and no obvious macroscopic growth. What is the next line of management?

a) Hysterectomy

b) Conization

c) Colposcopy and biopsy

d) Follow-up after 6 months

Question 4:

A patient with 6 weeks amenorrhea and a positive pregnancy test came with complaints of lower abdominal pain, moderate bleeding PV. She was hemodynamically stable. On USG, there was no gestational sac in the uterus. Ovaries were normal and no free fluid in the pouch

of Douglas. What is the next line of management?

a) Declare as complete abortion and follow-up in 15 days

b) Admit and observe

c) Diagnostic laparoscopy

d) Follow-up with beta-HCG

Question 5:

A hypertensive pregnant woman with abdominal pain, bleeding per vaginum, and loss of fetal movements. What is the probable diagnosis?

a) Abruptio placenta

b) Placenta previa

c) Hydramnios

d) Preterm labour

Question 6:

A 30-year-old woman visits an infertility clinic. She has a regular 28-day cycle. How should she be evaluated?

a) Serum progesterone on day 14

b) Serum progesterone on day 21

c) Serum LH on day 14

d) Serum LH on day 21

Question 7:

A 50-year-old woman presents with abnormal uterine bleeding for two months. What is the next line of management?

a) LNG- IUCD

b) POP for 3 months

c) Hysterectomy

d) Endometrial sampling

Question 8:

A patient with 12 weeks amenorrhea presents with vaginal bleeding. On examination, the cervical os is open and the uterus is of 14 weeks size. USG shows a snowstorm appearance. The next best step in the management is?

a) Medical management

b) Follow up with beta-hCG

c) Suction and evacuation

d) Hysterectomy

Question 9:

A 45-year-old woman is found to have a fibroid of 14 weeks size. She has no specific symptoms. What is the next step in management?

a) Hysterectomy

b) Myomectomy

c) Observation

d) Medical management

Question 10:

All of the following structures are cut in mediolateral episiotomy except?

a) Transverse perineal muscles

b) Obturator muscle

c) Bulbospongiosus

d) Transverse perineal branches of pudendal nerves

Question 11:

Raise in ESR during pregnancy is due to?

a) Albumin

b) Platelet

c) Fibrinogen

d) Anti-thrombin antibody

Question 12:

Which of the following criteria is used for diagnosing abdominal pregnancy?

a) Spiegelberg's criteria

b) Rubin's criteria

c) Studdiford's criteria

d) Godin's criteria

Question 13:

Which of the following is not a component of fetal biophysical profile?

a) Fetal movement

b) Fetal breathing

c) NST

d) Contraction stress test

Question 14:

All of the following can be used for the treatment of PCOD except?

a) Clomiphine citrate

b) Letrozole

c) Ulipristal

d) Laparoscopic ovarian drilling

Question 15:

Which of the following is not a component of the Bishop score?

a) Cervical length

b) Type of maternal pelvis

c) Cervical position

d) Dilatation of cervix

Question 16:

A nursing assistant posted in the obstetric care unit was asked to monitor the progress of labour. After how much dilatation of the cervix, should she start the documentation?

a) 2cm

b) 5cm

c) 4cm

d) 6cm

Question 17:

Identify the grip shown in the image below:

a) Pawlik's grip

b) Lateral grip

c) Pelvic grip

d) Fundal grip

Question 18:

Identify the fetal position in the given image.



a) Right occipito-posterior

b) Left occipito-anterior

c) Left occipito-posterior

d) Right occipito-anterior

Question 19:

A midwife is examining a pregnant woman in labour when she noticed that the cord is prolapsed. What should be the next immediate step of management?

a) Gently replace the cord inside the vagina

b) Inform the doctor and shift to OT

c) Put her in Trendelenberg position

d) Do vaginal examination and find the fetal presentation

Question 20:

A 45-year-old patient comes with repeated infections of trichomonas vaginitis, her physician advises for a Pap smear. She asks you regarding the purpose of this test. What would be your appropriate response?

a) It is a screening test for cervical cancer

b) It is a diagnostic test for trichomonas infection

c) It helps to diagnose cancer in her genital tract

d) It helps to diagnose other associated genital infections in her

Question 21:

Modified Robson's classification class 1 includes:

a) Nulliparous, singleton, cephalic, induced labour

b) Nulliparous, singleton, cephalic, spontaneous labour

c) Multiparous without previous CS, spontaneous labour

d) Multiparous without previous CS, induced/CS before labour

Question 22:

A pregnant woman has cramps in her legs. What will you suggest her to relieve the pain?

a) Extend knee and dorsiflex foot

b) Flex knee and dorsiflex foot

c) Extend knee and plantar flex foot

d) Flex knee and plantar flex foot

Question 23:

A woman in labour presented to the ER. A vaginal breech delivery was planned for her. What statement regarding the management is false?

a) Breech identified for the first time during delivery is a contraindication for vaginal delivery

b) Breech delivery is done in an hospital where a facility for emergency cesarean delivery is available

c) Continous FHR monitoring is required for breech delivery during labour

d) Should be done by a skilled obstetrician

Answer Key

Question No.	Correct Option
1	c

2	d
3	c
4	d
5	a
6	b
7	d
8	c
9	c
10	b
11	c
12	c
13	d
14	c
15	b
16	b
17	b
18	d
19	c
20	a
21	b
22	a
23	a

Detailed Explanations

Solution to Question 1:

Screening for cervical cancer is done in females aged 21 to 65 years.

*Three consecutive negative cytologic tests or two consecutive negative co-tests within the past 10 years, with the most recent test occurring within the past 5 years

Age Group	Screening Recommendations
<21 years	Do not screen
21-29 years	Cytologic testing (Pap smear) every 3 years

Age Group	Screening Recommendations
30-65 years	Preferred: Cytologic and HPV co-testing every 5 yearsAlternative: Cytologic testing every 3 years
>65 years	Discontinue screening if sufficient negative screening results* present, and if there is no history of HSIL, adenocarcinoma in situ, or cancer.
Women who have undergone total hysterectomy with removal of cervix	Discontinue screening

Solution to Question 2:

Oral contraceptive pills (OCPs) do not cause an increase in GnRH, they inhibit the release of GnRH.

OCPs act via the following mechanisms:

• Inhibiting ovulation - Both estrogen and progesterone act on the hypothalamic-pituitary axis to inhibit the release of GnRH (and thus subsequently LH and FSH). Seen in combined oral pills.

• Preventing implantation - Under the effect of OCPs, endometrial stroma undergoes hypoplasia, with atrophy of glands. This makes it unfavorable for implantation of the embryo. Seen primarily with mifepristone, but is a secondary mechanism of action with progesterone-only pills as well.

• Cervical mucus becomes thick and scanty under the effect of progesterone. This makes it difficult for sperm to enter. Seen in progesterone-only pills.

Solution to Question 3:

In a woman diagnosed with a high-grade squamous intraepithelial lesion (HSIL), the next step in management is to perform a colposcopy and biopsy.

Patients who are found to have HSIL are further subjected to biopsy. This is done by colposcopic study for the visual inspection with acetic acid (VIA) for aceto-white areas and taking biopsies of abnormal areas.



Solution to Question 4:

The patient presents with a history of 6 weeks of amenorrhea, a positive urine pregnancy test, abdominal pain and moderate vaginal bleeding. However, the ultrasound examination does not reveal a gestational sac in the uterus, which could indicate either an ectopic pregnancy or an abortion. In such cases, the next line of management is to follow up with beta-HCG (human chorionic gonadotropin) levels.

Beta-HCG levels above 1500 mIU/mL are considered the discriminatory value, beyond which a gestational sac should be visible on transvaginal ultrasound.

If the gestational sac is not visible on ultrasound and the beta-HCG levels are above 1500 mIU/mL, it suggests an ectopic pregnancy. On the other hand, if the beta-HCG levels are below 1500 mIU/mL, they can be repeated after 48 hours and interpreted as follows:

• Decreasing HCG levels between day 1 and day 3 suggest a possible abortion.

• If the HCG levels rise during this period but does not double, it indicates a likely ectopic pregnancy.

Note: While some institutions set the discriminatory threshold for HCG levels at ≥ 1500 mIU/mL, others use ≥ 2000 mIU/mL.

Solution to Question 5:

The given clinical picture with abdominal pain, bleeding per vaginum, and loss of fetal movements in a hypertensive pregnant woman points to the diagnosis of placental abruption.

Placenta previa (option B) refers to when the placenta is implanted partially or completely over the internal os and it usually presents with painless vaginal bleeding.

Hydramnios (option C) refers to a state where the volume of amniotic fluid exceeds 2000mL and it does not present with bleeding per vaginum.

In preterm labor (option D) fetal movements are not compromised.

Abruptio placenta is the partial/total placental separation from its implantation site before delivery. The bleeding may be:

• Revealed (blood dissects downwards between membranes and uterine wall to escape through the cervix)

• Concealed (blood accumulates between the detached placenta and uterus)

Hypertension in pregnancy is the most important risk factor for abruptio placenta. Patients often present with sudden onset abdominal pain, vaginal bleeding, uterine tenderness, and frequent contractions.

The hemorrhage into decidua basalis causes separation and compression of the placenta. This leads to a compromise of circulation to the fetus, leading to fetal distress.

Clinical classification of placental abruption:

Grade 1	Grade 2	Grade 3
Vaginal bleedingis slight	Vaginal bleeding mild to mod erate	Bleeding is moderate to sever e or concealed.
Uterus irritable, minimal ten derness	Uterine tendernessis always present	Uterine tenderness is marked.
Maternal BP and fibrinogen unaffected	Maternal pulse increased, BP maintained	Shock is pronounced
FHS good	Fetal distress present	Fetal death is the rule
Maternal shock absent	Maternal shock absent	Associated coagulation defect or anuria may complicate

Solution to Question 6:

Serum progesterone on day 21 should be evaluated in this woman to check for anovulatory cycles.

Anovulation is the major cause of female infertility. Therefore, tests are done to rule out anovulatory defects in a woman presenting with infertility.

After ovulation, the plasma concentration of progesterone increases and reaches its peak during the mid-luteal phase. Serum progesterone levels are assessed on day 8 and day 21 of a 28-day cycle. If the progesterone value increases from less than 1 ng/mL to greater than 6 ng/mL, it indicates ovulation. Subsequently, as the corpus luteum degenerates, progesterone levels decline.

If the plasma progesterone level remains low during the mid-luteal phase, it suggests a corpus luteal phase defect and indicates the need for hormonal therapy.

Daily estimation of serum LH (options C and D) around the midcycle can detect the LH surge. Ovulation coincides about 10 to 12 hours after the LH peak.

Solution to Question 7:

The next line of management in a 50-year-old woman with abnormal uterine bleeding is endometrial sampling.

Indications for endometrial sampling:

• Any female > 45 years with abnormal uterine bleeding

• A female of age < 45 years having exposure to excessive estrogen such as in PCOD, severe obesity, tamoxifen treatment

• Women with failed medical management and persistent bleeding

• Transvaginal ultrasound findings >12 mm in premenopausal women and > 4 mm in postmenopausal women.

Endometrial sampling may be done by either curettage or aspiration biopsy by pipelle or vibra aspirator.

Note: Transvaginal ultrasound is a better first-line investigation to assess AUB. Since TVS is not included in the options, endometrial sampling is the next best answer. Assessing the reason for AUB and ruling out malignancy in the postmenopausal age group are the initial steps before prescribing any treatment.

Solution to Question 8:

The clinical scenario of vaginal bleeding, an enlarged uterus exceeding the corresponding period of amenorrhea, and a snowstorm appearance on USG strongly indicates a diagnosis of a hydatidiform mole. Suction and evacuation is the next best step in the management of this patient.

Hydatidiform mole is an abnormal condition of the placenta, characterized by partly degenerative and partly proliferative changes in the chorionic villi. It may be either partial or complete.

The patient presents with a characteristic history of abdominal pain, vaginal bleeding or spotting, and vomiting during the first trimester. Also, there may be a history of passing grape-like vesicles vaginally. On examination, the size of the uterus is larger for gestational age and the fetal heart sounds are absent.

A USG scan shows a snowstorm appearance which is characteristic of a complete mole.

Suction and evacuation is the treatment of choice. Hysterectomy (option D) is done only if there is uncontrolled hemorrhage or perforation during surgical evacuation or if the age of the patient is above 35 years and has completed her family.

Serum hCG levels (option B) are checked within 48 hours of evacuation and are used as a baseline for future comparisons. It is then checked every week till it normalizes, which usually takes around 4 to 8 weeks. After the hCG values return to normal, a regular follow-up is done monthly for 6 months.

Solution to Question 9:

In this 45-year-old woman (>40 years) with an asymptomatic fibroid of 14-week size, the next best step in management would be observation.

Fibroids/leiomyomas are slow-growing benign tumours thus regardless of their size asymptomatic fibroids are managed conservatively with observation and annual checkups.

Indications for the treatment of fibroids are:

• Abnormal uterine bleeding

• A rapid increase in the growth of fibroid

• Infertility

• Chronic pelvic pain/ bulk symptoms

Other options:

Option A: Hysterectomy is usually reserved as the definitive treatment of choice for women who have completed their family and do not wish to preserve their fertility.

Option B: Myomectomy is a surgical procedure that involves removing fibroids while preserving the uterus. It is indicated in an infertile woman or a woman desirous of childbearing and wishing to retain the uterus.

Option D: Medical therapy aims to control menorrhagia in cases of symptomatic fibroids, improve haemoglobin levels before surgery, and shrink the fibroids before procedures such as myomectomy.

Solution to Question 10:

All the mentioned structures are cut in mediolateral episiotomy except for the obturator muscle.

Episiotomy refers to the surgically planned incision on the perineum and the posterior vaginal wall during the second stage of labor. The objective is to enlarge the vaginal introitus so as to facilitate easy and safe delivery of the fetus and to minimize overstretching and rupture of the perineal muscles and fascia, which might end up in complete tears.

In mediolateral episiotomy, the incision is made downwards and outwards from the midpoint of the fourchette either to the right or to the left. The other types of episiotomy include median, lateral, and J shaped episiotomy.

Structures cut in mediolateral episiotomy are:

• Posterior vaginal wall

• Superficial and deep transverse perineal muscles (option A)

• Bulbospongiosus and part of levator ani (option C)

• Fascia covering these muscles

• Transverse perineal branches of pudendal vessels and nerves (option D)

• Subcutaneous tissue and skin

Differences between midline and mediolateral episiotomy include:

Midline Episiotomy	Mediolateral Episiotomy
Surgical repair is easy	Surgical repair is difficult
Blood loss is lesser	Blood loss is more
Healing is superior	Healing is not so good
Dyspareunia is rare	Dyspareunia is comparatively more
Postoperative comfort is max imum	Postoperative discomfort is more
Extension, if occurs, may inv olve the rectum	Relative safety from rectal in volvement from extension

Solution to Question 11:

Raised ESR in pregnancy is due to an increase in fibrinogen and globulin levels.

Pregnancy is a hypercoagulable state. The fibrinogen level is raised by 50% during pregnancy. As a result of rise in fibrinogen and globulin level and diminished blood viscosity, erythrocyte sedimentation rate (ESR) gives a much higher value (fourfold increase) during pregnancy.

Total plasma protein increases during pregnancy. However, due to hemodilution, plasma protein concentration falls. This results in diminished viscosity of the blood and reduced colloid osmotic tension. Because of marked fall in albumin level and only slight rise of globulin (mainly alpha-globulin), the normal albumin: globulin ratio of 1.7:1 is diminished to 1:1.

Solution to Question 12:

Studdiford's criteria is used in the diagnosis of primary abdominal pregnancy. It includes:

• Presence of normal tubes and ovaries with no evidence of recent or past injury/pregnancy.

• Absence of uteroperitoneal fistula.

• Presence of a pregnancy related exclusively to the peritoneal surface and early enough to eliminate the possibility of secondary implantation after primary tubal nidation.

Other options:

Option A: Spiegelberg's criteria is used in the diagnosis of ovarian pregnancy:

• Tube on the affected side must be intact.

• The gestation sac must be in the position of the ovary.

• The gestation sac is connected to the uterus by the ovarian ligament.

• Ovarian tissue must be found on its wall on histological examination.

Option B: Rubin's criteria is used in the diagnosis of cervical pregnancy:

• Soft, enlarged cervix equal to or larger than the fundus.

• Uterine bleeding following amenorrhea, without cramping pain.

• Products of conception are entirely confined within and firmly attached to endocervix.

• A closed internal cervical os and a partially opened external os.

Option D:Godin's sonographic criteria is used for the diagnosis of a cesarean scar pregnancy:

• Empty uterine cavity

• Empty cervical canal

• A gestational sac in the anterior part of the uterine isthmus

• Absence of healthy myometrium between the bladder and gestational sac

Solution to Question 13:

The contraction stress test is not a component of the biophysical profile.

Biophysical profile is a multi-parametric screening test to assess uteroplacental insufficiency. It consists of five components and the observations are made for 30 minutes.

Each parameter has a maximum score of 2 and a minimum score of 0. So the total maximum score will be 10. A score of 4 or less indicates fetal acidemia.

Interpretation and management of BPP scores:

The modified biophysical score has only 2 components i.e NST and amniotic fluid index (AFI). It is considered abnormal (nonreassuring) when the NST is non-reactive and/or the AFI is <5.

Note: The contraction stress test (oxytocin challenge test) is an invasive method to assess fetal well-being during pregnancy. When there is an alteration in FHR in response to uterine contractions, it suggests fetal hypoxia.

Parameters	Minimal Normal Criteria
Non-stress Test (NST)	≥2 accelerations within 20-40 minutes.
Fetal breathing movements	≥ 1 episode of rhythmic breathing lasting > 30 seconds.
Gross body movements	≥ 3 discrete body/limb movements
Fetal muscle tone	≥ 1 episode of active extension(limb or trunk) with return of flexion.
Amniotic fluid volume	A pocket of amniotic fluid that measures at least 2 cm in two perpendicular planes (2 × 2 cm pocket)

BPP score	Interpretation	Management
10	Normal, nonasphyxiated fetus	Repeat test weekly
8with normal AFV	Normal, nonasphyxiated fetus	Repeat test weekly
8 with decreased AFV	Suspicion of chronic fetal asphyxia	If ≥36 weeks, delivery is done
6	Possible fetal asphyxia	If AFV is abnormal-Deliver>36 weeks.If AFV is normal- deliver≥ 37 weeks.
4	Probable fetal asphyxia	If ≥32 weeks, deliver
0-2	Almost certain fetal asphyxia	Delivery

Solution to Question 14:

Ulipristal is not used for the treatment of polycystic ovarian disease (PCOD). It is a synthetic progesterone hormone receptor modulator used as an emergency contraceptive pill.

PCOD is a complex multisystemic endocrine disorder affecting women of reproductive age.

Rotterdam criteria is used to diagnose PCOD. The presence of two of the three components of this criteria establishes the diagnosis.

• Oligo/amenorrhoea, anovulation, infertility

• Hirsutism/acne

• Ultrasound findings:

• Enlarged ovaries with ovarian volume will be more than 10 mL

• 12 or more small follicles each of 2-9 mm in size placed peripherally

Clomiphene (option A) is a selective estrogen receptor modulator that is used as the first line of treatment in managing infertility. Clomiphene with dexamethasone improves the fertility rate. In resistant cases, tamoxifen or letrozole (an aromatase inhibitor) (option B) can be tried.

Surgical management of PCOD includes laparoscopic ovarian drilling (option D). Indications of surgery are:

• Failed medical therapy

• Hyperstimulation occurs

• Infertile women

• Previous pregnancy losses

Solution to Question 15:

The type of maternal pelvis is not a component of the Bishop score.

The Bishop score is a scoring system used to assess labor induction outcomes. It is based on the station of the head, dilation, effacement (or length), position, and consistency of the cervix noted by doing a vaginal examination.

Note: In the modified Bishop score (1991), the effacement is replaced by the cervical length in cm.

Solution to Question 16:

The WHO Labour Care Guide begins the documentation of labour progression at 5 cm of cervical dilatation.

The WHO Labour Care Guide is a tool designed to help health personnel involved in labour care. It aims to

• Guide the monitoring and help in the documentation of the progress of labour.

• Guide the health professionals to offer supportive care throughout labour.

• Assist doctors in promptly identifying and addressing emerging labour complications.

• Prevent the use of unnecessary interventions in labour.

• Support audits and quality improvement.

The differences between the WHO-modified partograph and the latest WHO Labour Care Guide are tabulated below:

Note: Previously the answer to the question was 4 cm (according to the modified WHO patograph). However, this information is outdated, therefore the answer to this question has been revised per the latest WHO guidelines.

WHO modified partograph	WHO Labour Care Guide
Documentation of labour pro gression begins at4 cmof cerv ical dilatation.	Documentation of labour pro gression begins at5 cmof cerv ical dilatation.
Fixed1 cm/hour ‘alert’ and ‘ac tion’lines	Time-based cut-offsforeach c entimetreof cervical dilatatio n to guide the obstetrician to take action. For example, at 5 cm of cervical dilatation, if t here is no progress >6 hours, action needs to be taken.
Monitoring of labour is restri cted to thefirst stageonly.	Monitoring ofthe second stag e of labouris included
No section on supportive car e interventions	Addition of a section onsupp ortive careof the mother- lab our companionship, pain reli ef, oral fluid intake and postu re
Documentation ofstrength, d uration, and frequencyof uter ine contractions	Documentation ofonlythedur ation and frequency of uterin e contractions



Solution to Question 17:

The Leopold manoeuvre shown in the image is the lateral grip. It assesses fetal orientation.

Leopold's manoeuvres help assess the fetal pole, lie, presenting part, engagement, and attitude of the fetal head. They are as follows:

• Fundal grip (first Leopold):

• Assesses the uterine fundus.

• Helps in identification of fetal lie and determination of fetal pole at the fundus.

• The breech is felt as a large, irregular mass.

• The head feels hard and round and is more mobile.

• Lateral grip (second Leopold):

• Assesses the fetal orientation.

• The palms are placed on either side of the maternal abdomen, and gentle but deep pressure is exerted.

• The back is felt on one side as a hard, resistant structure.

• The fetal extremities are felt on the other side as numerous small, irregular, mobile parts are felt.

• Fetal orientation can be determined by noting whether the back is directed anteriorly, transversely, or posteriorly.

• The position of the anterior shoulder is mainly assessed by this grip.

• Pawlik's grip / 1st pelvic grip (third Leopold):

• Assesses the presenting part.

• The thumb and fingers of one hand grasp the lower portion of the maternal abdomen just above the symphysis pubis.

• If the presenting part is not engaged, a movable mass will be felt, usually the head.

• Deep pelvic grip / 2nd pelvic grip (fourth Leopold):

• Assesses the engagement and attitude of the fetal head.

• The examiner faces the mother's feet, and the fingertips of both hands are positioned on either side of the presenting part.

• They exert inward pressure and then slide along the axis of the pelvic inlet.

Leopold	Grip	Importance
First	Fundal	To know which fetal pole occupies the uterine fundus.

Leopold	Grip	Importance
Second	Lateral/Umbilical	To know the fetal back and extremities
Third	Pawlik	To know the presenting fetal part in respect to the true pelvis
Fourth	Pelvic	To determine the degree of descent.If the head has already descended into the pelvis, the anterior shoulder or the space created by neck may be differentiated readily from the head



Solution to Question 18:

In the given vertex presentation, the fetal position is right occipitoanterior.

Fetal position refers to the relation of the denominator (arbitrary bony fixed point on the presenting part), with respect to different quadrants of the maternal pelvis. In vertex presentation, the denominator is taken as the occiput (which is close to the triangular posterior fontanelle).

To determine the fetal position the following two steps are followed:

• Determine the direction in which the occiput is facing:

• Towards the pubic symphysis - occipito-anterior

• Towards the sacral promontory - occipito-posterior

• Midway between pubic symphysis and sacral promontory - occipito-transverse

• Determine if the occiput is facing the left or the right in relation to the materal pelvis



Solution to Question 19:

In this scenario the midwife should put the pregnant women in a Trendelenburg position or in an exaggerated and elevated Sims' position with a pillow or wedge under the hip or thigh.

Caesarean section is the recommended treatment of choice for cord prolapse. Prior to performing the procedure, several measures can be taken to alleviate cord compression. These measures involve:

• Manually elevating the fetal presenting part using two fingers or the entire hand through the vagina

• Positioning the patient in steep Trendelenburg, exaggerated Sim's position, or knee-chest position

• Filling the bladder with 500–700 cc or more of saline (known as the Vagos method)

After the diagnosis of cord prolapse is made, it is crucial not to attempt cord replacement. Such attempts are not only ineffective but also result in the inevitable rupture of the membranes, leading to cord prolapse.



Solution to Question 20:

A pap smear is a screening test for cervical cancer. It is not a diagnostic test (option C). The presence of abnormal cells/ a positive test requires further evaluation by colposcopy, cervical biopsy, or fractional curettage.

If trichomonas is reported from Pap testing, it should be confirmed by saline microscopic preparation (option B).

Cervical cancer screening should start at age 21. Women aged 21-29 should have a Pap smear every 3 years, while women aged 30-65 should have both HPV and Pap co-test every 5 years. Women should stop having cervical cancer screening after the age of 65 years. If suspicious PAP, screening till 75 years. HIV+ women should have annual checkups until three consecutive negative tests followed by testing every 3 years for life.

Pap testing should be done prior to vaginal examination. Lubricants should not be used and the patient should not be menstruating during the pap test. The patient should not have intercourse 24hrs prior to the test. The ideal time is around ovulation.

Procedure:

• The patient is put in a dorsal position, and the labia are parted

• Cusco's self-retaining speculum is introduced

• The squamocolumnar junction is scraped with Ayer's spatula:

• Bifid end - ectocervical sample

• Blunt end - vaginal cytology

• Endocervix is scraped with a cytobrush

• The sample is spread on a glass slide

• The slide is immediately fixed using 95% ethyl alcohol

• Stained with Papanicolau stain or short-duration Papanicolau stain (SPS).



Solution to Question 21:

Class 1 of modified Robson's classification includes nullipara, equal to or >37 weeks, single, cephalic, spontaneous labor.

It is the most widely accepted classification system available for analyzing characteristics of delivering women worldwide. It consists of 10 patient population categories. The basis for these categories include:

• Parity

• Onset of labor and mode of delivery

• Gestational age

• Fetal presentation

• No. of fetuses

The modified Robson system classification is as follows:

Class 1	Nullipara, equal to or >37 weeks, single, cephalic, spontaneous labor
Class 2	Nullipara, equal to or >37 weeks, single, cephalic2a:induced labor2b:CS before labor
Class 3	Multipara, equal to or >37 weeks, single, cephalic, spontaneous labor (excludes previous CS)
Class 4	Multipara, equal to or >37 weeks, single, cephalic4a:induced labor4b:CS before labor
Class 5	Multipara, previous CS, equal to or >37 weeks, single, cephalic
Class 6	Nullipara, single, breech
Class 7	Multipara, single, breech (including previous CS)
Class 8	Multiple pregnancies (with or without previous CS)
Class 9	Singleton pregnancy, oblique/transverse lie (with or without previous CS)
Class 10	Single, cephalic <37 weeks (including previous CS)

Solution to Question 22:

Extension at the knee joint and dorsiflexion at the ankle joint helps in stretching the calf muscles which relieves the pain.

Leg cramps are minor ailments during pregnancy mostly due to a deficiency of diffusible serum calcium or elevation of serum phosphorus. Management includes massaging the leg, stretching exercises, application of local heat, and supplementation of vitamin B1 and calcium.

Solution to Question 23:

Breech identified for the first time during delivery is not a contraindication for normal delivery.

Vaginal breech delivery is considered when:

• Fetal weight is between 1.5 kg and 3.5 kg

• There is an adequate pelvis

• The presentation is frank breech

• The fetal head is flexed

• There are no medical or obstetric complications

• Facilities for emergency cesarean section are available (anesthetists, neonatologists) (option B)

• Facilities for continuous labor monitoring are available (option C)

• A skilled obstetrician is present (option D)

• Detailed informed consent has been obtained, explaining the risks of vaginal breech delivery

Cesarean delivery for breech presentation is considered in the following cases:

• Lack of operator experience

• Patient's request for cesarean delivery

• Prior perinatal death or neonatal birth trauma

• Large fetus: > 3800 to 4000 g

• Severe IUGR, term weight <2.5kg to 2.8kg

• Incomplete breech presentation

• Oligohydramnios

• Fetal anomaly incompatible with vaginal delivery

• Hyperextended head

• Contracted pelvis

• Prior cesarean delivery

• Preterm breech

• Twins, with the first baby in the breech position

Obstetrics and Gynecology NEET 2018

Question 1:

In which trimester does acute fatty liver manifest most commonly during pregnancy?

a) First trimester

b) Second trimester

c) Third trimester

d) Both (a) and (b)

Question 2:

A 55-year-old lady comes with abnormal uterine bleeding. A hysterectomy was performed and the following specimen was retrieved. What is the diagnosis?

a) Leiomyoma

b) Endometrial carcinoma

c) Endometrial polyp

d) Adenomyosis

Question 3:

The ligament that maintains anteversion of the uterus during pregnancy is:

a) Round ligament

b) Cardinal ligament

c) Uterosacral ligament

d) Pubocervical ligament

Question 4:

Quadruple testing is done during:

a) 9-11 weeks

b) 12-14 weeks

c) 15-20 weeks

d) 22-28 weeks

Question 5:

What is the ideal number of antenatal visits?

a) 12-14

b) 6-8

c) 7-9

d) 10-11

Question 6:

A woman is diagnosed to be positive for HIV in her first trimester. What is the appropriate management?

a) Start ART immediately and continue till 6 weeks after delivery

b) Start ART after 1st trimester and continue life-long

c) Start ART immediately and continue life-long

d) Start ART after 1st trimester and continue till 6 weeks after delivery

Question 7:

The fate of the Mullerian duct is determined by the:

a) X chromosome

b) Y chromosome

c) 1st chromosome

d) 2nd chromosome

Question 8:

Which of the following is the first step after fitting the pessary to check if the pessary is adequately retained?

a) Asking the patient to void and see if the pessary is in place

b) Seeing if the patient feels any discomfort, which means that the pessary is retained

c) Asking the patient to perform the Valsalva manouvre and see if the pessary is in place

d) Being able to admit two fingers between the ring and the pessary.

Question 9:

A pregnant woman presents with obstructed labor and is grossly dehydrated. Investigations reveal fetal demise. What will be the management?

a) Craniotomy

b) Decapitation

c) Cesarean section

d) Forceps extraction

Question 10:

The best time to identify the type of twin pregnancy on ultrasound is:

a) 18-22 weeks

b) 24-28 weeks

c) 11-14 weeks

d) 14-18 weeks

Question 11:

Which of the following statements is false about the umbilical cord?

a) It has one artery and two veins

b) It contains a gelatinous substance called Wharton's jelly

c) Presence of a single umbilical artery is associated with congenital anomalies

d) Normal average umbilical cord length is 50-60 cms

Question 12:

Which of the following statements is false about non-invasive prenatal testing?

a) It uses cell-free circulating fetal DNA in the maternal serum

b) It is used for the diagnosis of trisomy 13, 18 and 21

c) It can be performed after 10 weeks of gestation

d) It is done after the routine first trimester ultrasound assessment

Question 13:

Which of the following is not a cause of secondary PPH?

a) Retained cotyledon

b) Endometritis

c) Placental polyp

d) Placenta previa

Question 14:

A 16-year-old girl with primary amenorrhea has the following features. What is the most probable diagnosis?

a) Testicular feminizing syndrome

b) Gonadal dysgenesis

c) Kallmann syndrome

d) Turner syndrome

Question 15:

Fimbriectomy is also known as:

a) Pomeroy technique

b) Uchida's procedure

c) Kroener's procedure

d) Irving's procedure

Question 16:

Which of the following cannot cross the placenta?

a) 1 and 2

b) Only 3

c) 2 and 3

d) 3 and 4

Question 17:

When is dilatation and curettage contraindicated?

a) Pelvic inflammatory disease

b) Ectopic pregnancy

c) Menstrual irregularities

d) Endometriosis

Question 18:

A 45-year-old lady complains of post-coital bleeding. She has a positive pap smear. What is the next line of management?

a) Colposcopy directed biopsy

b) Cone biopsy

c) Repeat pap smear

d) Hysterectomy

Question 19:

In superfecundation, which of the following is seen?

a) Fertilisation of 2 ova released at the same time by sperms released at single intercourse

b) Fertilisation of 2 ova released in the same cycle by sperms released at intercourse on 2 different occasion

c) Both of the above

d) None of the above

Question 20:

All are true regarding peripartum cardiomyopathy except _____

a) Development of cardiac failure within 5 months of delivery

b) Left ventricular diastolic dysfunction

c) Absence of identifiable cause

d) Dilated left ventricle

Question 21:

Which of the following is not the mechanism of action of progesterone only pills?

a) Thicken the cervical mucus

b) Prevents ovulation

c) Kills the sperms in the cervical canal

d) Inhibits penetration of sperm

Question 22:

Which of the following is not considered as a factor for a high-risk pregnancy?

a) Obesity

b) Polyhydramnios

c) Anaemia

d) Previous manual removal of placenta

Question 23:

Long-standing pelvic inflammation may lead to which of the following conditions?

a) Pyometra

b) Uterine polyposis

c) Pseudocyesis

d) Endometrial hyperplasia

Question 24:

Which among the following is the most common presentation of postpartum nerve injury?

a) Facial palsy

b) Wrist drop

c) Foot drop

d) Claw hand

Question 25:

What is altered in thyroid profile throughout pregnancy?

a) Free T3 level

b) Free T4 level

c) Total T3 level

d) TSH level

Answer Key

Question No.	Correct Option
1	c
2	a
3	a
4	c
5	a
6	c
7	b

8	c
9	c
10	c
11	a
12	b
13	d
14	b
15	c
16	c
17	a
18	a
19	b
20	b
21	c
22	d
23	a
24	c
25	c

Detailed Explanations

Solution to Question 1:

Acute fatty liver of pregnancy (AFLP) occurs most commonly in the third trimester or early postpartum period.

Risk factors include Pre-eclampsia/HELLP syndrome, male fetus, previous history of AFLP, and multiple gestations.

AFLP (acute yellow atrophy or acute fatty metamorphosis) is the most common cause of acute liver failure in pregnancy. It usually occurs in the later stages of pregnancy, particularly in the third trimester. It occurs due to fetal deficiency of long-chain acyl-coenzyme A dehydrogenase (LCHAD) or poor placental mitochondrial function.

The patients present with persistent nausea and vomiting, progressive jaundice, and epigastric pain. Features like acute hemolysis, hypoglycemia, hypertension, proteinuria, thrombocytopenia, and edema can also be present. Symptoms may progress to fulminant hepatic failure with encephalopathy, disseminated intravascular coagulation, and death.

Liver function tests such as AST, ALT, and serum bilirubin levels are raised. Hypofibrinogenemia, hypocholesterolemia, and prolonged clotting time can be seen. An elevated d-dimer is seen. Microscopy reveals a characteristic microvesicular fatty infiltration.

The condition worsens till delivery after which the hepatic dysfunction resolves. A trial of labor by induction or delivery by Cesarean is done. Intensive supportive measures including blood transfusion should be provided.

The clinical and laboratory findings of acute liver diseases in pregnancy are as follows:

The condition worsens till delivery after which the hepatic dysfunction resolves. Appropriate management is immediate termination of pregnancy irrespective of gestational age by induction or delivery by Cesarean (for obstetric reasons) is done. Intensive supportive measures including blood transfusion should be provided.

Disorders	HELLP syndrome	Obstetric cholestasis	Acute fatty liver of pregnancy	Viral hepatitis
Onset in pregnancy	Mid to late	Late	Late	Variable
Differentiating features	Epigastric pain, hypertension,elevatedindirect bilirubin	Prurituselevateddirect bilirubin	Signs ofliver failurelike hypoglycemia	Fever
AST (U/L)	≤300	≤200	200-800	2000+
Bilirubin (mg/dL)	1-4	1-5	4-10	5-20
Platelets	↓↓	Normal	↓↓	↓
Fibrinogen	Normal	Normal	↓↓↓	Normal
D-dimer	↑	Normal	↑	Normal
Prothrombin time	Normal	Normal	↑↑	↑
Hemolysis	↑↑	No	↑↑↑	No

Solution to Question 2:

The given cut section image showing multiple nodules, well-circumscribed, firm, and white-greyish with a whorled appearance is Leiomyoma (Fibroid).

Leiomyoma is the most common benign and the most common pelvic tumor in females and is more common in obese females(hyper-estrogenism). They are mostly asymptomatic and the most common symptom is menorrhagia.

Leiomyomas can also present with bleeding areas, cystic degeneration, and calcifications. The uterine cavity is compressed.

The image below different types of uterine fibroids classified according to their locations:



Sub-mucous fibroids are the most symptomatic type and intra-mural fibroids are the most common type of uterine fibroids.

Other options:

Option B: Endometrial cancer presents as a polypoid mass protruding into the uterine cavity or as a diffuse thickening of the endometrium. The cut surface may appear white to yellow and may show areas of necrosis or hemorrhage.

Option C: Endometrial polyps are usually small and may be single or multiple. Grossly, they appear as soft, red to pink, often pedunculated (attached by a stalk) masses protruding into the uterine cavity. It presents with heavy cyclic or intermenstrual bleeding, or without complaints as an incidental finding on imaging. Estrogen and progesterone have been implicated in the growth of endometrial polyps, and higher receptor levels are seen compared with the normal endometrium.

Option D: In Adenomyosis (Endometriosis interna), the uterus may appear bulky and symmetrically enlarged. On the cut section, the myometrium may appear thickened and may show tiny cystic spaces filled with old blood (chocolate cysts). The preferred treatment is hysterectomy. Localised excision or levonorgestrel IUCD is tried in younger women.

The image below shows both the gross and histologic appearances of adenomyosis:

Solution to Question 3:

The round ligament helps maintain the anteversion of the uterus during pregnancy.

However, the anteversion in the non-pregnant uterus is maintained by both the uterosacral ligament and round ligament.

Anteflexion of uterus is at the level of internal os.

(Mnemonic - V for version, V for vagina)

The uterus normally lies in a position of anteversion and anteflexion.

In anteversion, the external os is directed downward and backward, so that on vaginal examination, the fingers find that the lowest part of the cervix is the anterior lip.

When the uterus is retroverted, the cervix is directed downward and forward, and the lowest part of the cervix is either the external os or its posterior lip.

Angle of anteversion	Angle of anteflexion
Angle between thecervix and vagina	Angle betweencervix and uter us
900	1200





Solution to Question 4:

The quadruple screening test is ideally done in the second trimester, during 15-20 weeks of gestation. The most accurate values are seen during 16-18 weeks of gestation.

It is the measurement of maternal serum alpha-fetoprotein (MSAFP), unconjugated estriol, hCG, and inhibin-A.

In Down's syndrome (trisomy 21) or Edward's syndrome (trisomy 18), quadruple-marker screenings do not provide any added benefit over first-trimester screenings. It is generally used as a stand-alone test if screening is done for the first time in the second trimester or if first-trimester screening is not available.

	Trisomy 21	Trisomy 18	Neural tube defect or abdominal wall defect	Trisomy 13/ Patau syndrome
MSAFP	Low	Low	High	Normal
β hCG	High	Low	Normal	Normal
Unconjugated estriol	Low	Low	Normal	Normal
Inhibin A	High	Normal/ Low	Normal	High

Solution to Question 5:

The ideal number of antenatal visits is 12-14.

The antenatal visits should ideally be: monthly once up to 28 weeks, once in two weeks between 28-36 weeks, and once weekly visits from 36 weeks onwards.

This translates to a total of 12-15 visits.

The WHO increased the minimum number of antenatal visits from 4 to 8 visits to reduce perinatal mortality.

In the Indian scenario, the earlier minimum of 4 visits were implemented as follows:

• 1st visit: within 12 weeks (preferably as soon as pregnancy is suspected)

• 2nd visit: between 14-26 weeks

• 3rd visit: between 28-34 weeks

• 4th visit: between 36 weeks and term

A booked case is one that has at least 3 antenatal visits with two visits in the third trimester.

Solution to Question 6:

ART should be initiated immediately and continued lifelong in all pregnant and breastfeeding women with HIV. This is regardless of their WHO clinical stage and CD4 cell count.

*Note: Pearl attached below has been updated according to the NACO ART guidelines 2021. Efavirenz has been replaced by Dolutegravir as per the new guidelines.

Solution to Question 7:

The fate of the Mullerian duct is determined by the presence or absence of the Y chromosome as follows:

• In presence of the Y chromosome, the SRY gene (Sex-determining Region of the Y chromosome) produces the protein called the testis determining factor which leads to the gonads developing as

the testes. The Sertoli cells of the testis will then produce a peptide hormone, the Müllerian inhibiting substance (MIS), also known as Anti-Müllerian hormone (AMH), and cause the Müllerian ducts to regress.

• In absence of the Y chromosome and the SRY gene consequently, the gonads will develop as ovaries and the Mullerian duct will differentiate into the fallopian tubes and the upper part of the vagina. The WNT4 gene on the 1st chromosome promotes female sex development.

Solution to Question 8:

The first step after fitting the pessary to check if the pessary is adequately retained is to ask the patient to perform Valsalva's maneuver or cough and see if the pessary is in place.

There is some level of descent of the pessary with these maneuvers is to be expected, but the pessary should return to its normal position when these efforts are stopped. If the pessary descends to the introitus with these maneuvers, then a larger size pessary should be tried.

The pessary is well-fitted if you are able to slide the ring of the pessary up and down along the vaginal sidewall. It is well-fitted if you are able to insert one finger between the vaginal sidewall and the edge of the pessary. When the labia are separated, a well-supported pessary is not usually visible.

The patient should be comfortable, and able to void urine with the pessary in place before leaving the clinic.



Solution to Question 9:

The best management in the given clinical scenario is cesarean section.

Obstructive labor management involves administering intravenous fluids to correct dehydration, providing antibiotics to prevent infection, and promptly performing a cesarean section to relieve

the obstruction. If the fetus is alive, an immediate cesarean section is typically performed. Management of obstructed labor never involves a wait-and-watch approach or augmenting with oxytocin.

Clinical features of obstructed labor include maternal exhaustion, dehydration, tachycardia, tachypnea, genital sepsis, postpartum hemorrhage, and shock.

On P/A examination the upper uterine segment is tender and tonically contracted. The lower uterine segment is thin and stretched. A groove/depression is felt between the upper uterine segment and the lower uterine segment. This is called a Bandls ring.

Complications of obstructed labor include rupture of the uterus and vesicovaginal fistula.

Solution to Question 10:

The best time to identify the type of twin pregnancy on ultrasound is 11-14 weeks.

Twin pregnancies should ideally be dated when the crown–rump length (CRL) measurement is between 45 and 84 mm (i.e. 11 + 0 to 13 + 6 weeks of gestation).

Chorionicity should be ideally determined between 11 + 0 and 13 + 6 weeks of gestation (the earliest time to identify chorionicity is 7 weeks). This is done using the membrane thickness at the site of insertion of the amniotic membrane into the placenta. The T■sign (in monochorionic twins) or lambda sign (in dichorionic twins) can be seen here. As the determination of chorionicity is most accurate when the amnion and chorion have not yet been fused, the first■trimester scan in twin pregnancy is highly important.

Markers of dichorionicity include 4 layers of membranes between the twins, membrane thickness >2mm, different sex of twins, 2 different placentas, and positive twin peak sign.

Women with an uncomplicated dichorionic twin pregnancy should have a first■trimester scan, a detailed second■trimester scan, and scans every 4 weeks thereafter.

Women with uncomplicated monochorionic twins should have a first■trimester scan and then scans every 2 weeks from 16 weeks onwards. This is done in order to detect TTTS and TAPS in a timely manner.

Solution to Question 11:

The umbilical cord normally has two arteries and one vein. The presence of a single umbilical artery is the most common vascular anomaly of the cord. It is associated with congenital anomalies such as genitourinary and cardiac anomalies.

The umbilical cord is a tube that connects the fetus to the placenta. It contains umbilical vessels and an obliterated allantois duct, all surrounded by Wharton's jelly and contained within an outer layer of amnion. Maternal blood enters the fetus through the umbilical vein, and blood exits the fetus via the two umbilical arteries. Wharton's jelly is a gelatinous mucopolysaccharide that protects the umbilical vessels. Local collections of Wharton's jelly produce false knots in the cord.

The average length of the umbilical cord is between 40-70 cm at term. A long cord can form a loop around the fetus' neck and can cause asphyxia. A short cord can result in cord rupture during labor.

Solution to Question 12:

Non-invasive prenatal testing (NIPT) is used for the screening of trisomy 13, 18 and 21 and Turner's syndrome. It is not a diagnostic test.

It is a novel genetic test that uses cell-free circulating fetal DNA in the maternal serum for screening of chromosomal anomalies.

DNA from the fetus is found circulating in the maternal blood in two forms - either inside the intact fetal cells or as cell-free DNA from the breakdown of placental fetal cells. In NIPT, the cell-free chromosome fragments are measured and the quantitative differences from normal fetuses are noted.

NIPT can be performed after 10 weeks of gestation and is recommended after the routine first-trimester ultrasound assessment in high-risk pregnancies or if either 1st or 2nd-trimester screening becomes positive. Its disadvantage is that it is very expensive and it can be used for screening of only a few chromosomal anomalies such as trisomy 13, 18, and 21 and it doesn't analyse the entire karyotype.

Solution to Question 13:

Placenta previa does not cause secondary postpartum hemorrhage. It causes primary postpartum hemorrhage.

Types of PPH:

Primary/early: Hemorrhage occurring within 24 hours of childbirth.

Secondary/ late: Hemorrhage occurring after 24 hours and up to 12 weeks postpartum.

Causes and risk factors for PPH:

The image below shows the prevention and management of PPH:

Primary PPH	Secondary PPH		
Atonic PPH	Traumatic PPH	Coagulopathy	
Primiparity or grand multiparity	Instrumental delivery	Sepsis	Retained bits of cotyledon or membranes
Placental abnormalities like placenta previa/accreta/increta/percreta/abruptio placenta	Precipitate labour	HELLP syndrome	Secondary haemorrhage from Caesarean section wound

Primary PPH	Secondary PPH		
Overdistended uterus as in multi-gestational pregnancy, hydramnios, macrosomia	Macrosomia	IUD	Puerperal uterine inversion
Induction of labour with prostaglandins or oxytocin	Episiotomy		Abnormal involution of placental site
Prolonged labour			Infected fibroid
Precipitate labour			Endometritis
Previous h/o PPH			Placental polyp
Obesity			Carcinoma of cervix
Age >40 years			Uterine av fistula



Solution to Question 14:

The scenario of a 16-year-old girl with absent secondary sexual characters, normal stature, with XX karyotype and high FSH and LH is suggestive of pure gonadal dysgenesis.

46 XX and 46 XY gonadal dysgenesis without the phenotype of Turner's syndrome (short stature, webbed neck) is described as pure gonadal dysgenesis. It is characterized by the presence of primary amenorrhea with or without normal secondary sexual characteristics.

Option A: Testicular feminizing syndrome occurs in karyotype XY. Secondary sexual characters are present and the uterus is absent.

Option C: Kallmann syndrome is a form of hypogonadotropic hypogonadism. GnRH synthesis is reduced, therefore FSH and LH are low, and secondary sexual characters are absent.

Option D: Turner syndrome is associated with short stature and karyotype Xo. Secondary sexual characters and the uterus is present, but the ovaries are streaked.





Solution to Question 15:

Fimbriectomy is also known as Kroener's procedure. This procedure involves ligation of the distal ampulla of the tube with two permanent sutures. Then, resection of the infundibulum of the tube is done.



Some more techniques of permanent sterilization include:

Pomeroy method - The fallopian tube is identified on each side, brought out through the incision. The middle portion is formed into a loop which is tied at the base with catgut and excised. This is the most commonly used technique.



Uchida's method - The tubal serosa is stripped off the muscular layer in the midsegment of the tube. The muscle layer is excised. The proximal end is ligated and buried in the broad ligament. This minimal excision of the tube preserves the potential for tuboplasty.



Irving's procedure - The mid-portion of the tube is ligated and the intervening portion is excised. The proximal end is buried in the myometrium and the distal end is buried in the broad ligament. It is a reliable method but irreversible and may require a laparotomy incision.



Madlener's procedure - A loop of the fallopian tube is crushed and then ligated with a nonabsorbable suture.



Solution to Question 16:

Albumin and IgM cannot cross the placenta.

High molecular weight substances such as proteins (e.g., albumin, IgM, IgA) cannot cross the placenta, except IgG.

Substances which can cross the placenta are as follows:

Note: Maternal hormones like insulin, steroid hormones, and thyroid hormones cross the placenta very slowly, and only in amounts enough to support fetal development. However, calcitonin and parathormone do not cross the placenta.

Substances	Mechanism
O2CO2WaterElectrolytesAnesthetic agents	Simple diffusion
Immunoglobulin G(molecular weight 160,000 Da)	Endocytosis via Fc receptors in the trophoblast
Glucose	Facilitated diffusion via glucose transport proteins (GLUTs)
Amino acids	Transport-protein-mediated active transport
Lipids:GlycerolEssential fatty acidsLDL	Endocytosis via specific receptors
Minerals:IodineIron (unidirectional from mother to fetus) CalciumPhosphorus	Carrier-mediated active transport
Vitamins:Vit AVit CVit D	Carrier-mediated active transport

Solution to Question 17:

Dilatation and curettage (D&C) is contraindicated in pelvic inflammatory disease (PID) as it increases the risk of ascending genital infection.

D&C is a minor gynecological procedure of dilating the cervix and curetting the endometrial tissue from the uterine cavity. Diagnostic indications of D&C include AUB, secondary amenorrhea to detect tubercular endometritis, postmenopausal bleeding to rule out endometrial cancer or hyperplasia, and infertility due to genital TB.

Therapeutic indications of D&C include missed abortion, incomplete abortion, retained products of conception, and endometrial polypectomy.

Absolute contraindication of D&C includes Suspected/ desired pregnancy and lower genital tract infection. However, D&C is performed in some cases with genital infections like septic abortion (where retained pieces of placenta act as a nidus and need to be removed) and pyometra (as there is cervical obstruction with superinfection of menstrual blood and cellular debris, D&C is appropriate to remove the obstruction as well as the source of infection)

Complications of D&C include ascending infection, cervical tear with bleeding, incompetent os, uterine perforation, and ashermann syndrome.



Solution to Question 18:

The appropriate management of a 45-year-old with a positive pap smear and post-coital bleeding is a colposcopy-directed biopsy.

A per speculum examination must be done whenever a female comes with post-coital bleeding in order to evaluate for carcinoma of the cervix. If growth is visible, a punch biopsy should be done, and else if no growth is visible, then a pap smear is done. A pap smear is a screening test for cervical cancer. The presence of abnormal cells/ a positive test requires further evaluation by

colposcopy, cervical biopsy, or fractional curettage.

Cervical cancer screening should start at age 21. Women aged 21-29 should have a pap smear every 3 years, while women aged 30-65 should have both HPV and pap co-test every 5 years. Women should stop having cervical cancer screening after the age of 65 years. If a pap smear is suspicious, then screening is continued till 75 years. HIV+ve women should have annual checkups until three consecutive negative tests followed by testing every 3 years for life.

Pap testing should be done prior to vaginal examination. Lubricants should not be used and the patient should not be menstruating during the pap test. The patient should not have intercourse 24hrs prior to the test. The ideal time is around ovulation.

Solution to Question 19:

Superfecundation is the fertilization of two different ova released in the same cycle, by two separate acts of coitus within a short period of time.

Superfecundation is the fertilization of two different ova in the same menstrual cycle (as suggested by the same gestational age) by separate acts of coitus. It can occur even after artificial reproductive therapy. Hence, after embryo transfer, intercourse must be avoided. Heteropaternal superfecundation is used to refer to instances of two different males fathering fraternal twins. This leads to the possibility of twins also being half-siblings. This may give rise to the doubt of adultery and infidelity.

Superfetation is the fertilization of two ova released in different menstrual cycles. This is not seen in humans. The nidation and development of one fetus over another fetus is theoretically possible until the decidual space is obliterated by 14-16 weeks of pregnancy.

Solution to Question 20:

Left ventricular systolic dysfunction is a feature of peripartum cardiomyopathy (also known as postpartum cardiomyopathy) and not diastolic dysfunction.

Peripartum cardiomyopathy (PPCM) is a type of dilated cardiomyopathy. It develops during the last month of pregnancy or within the first 5 months following delivery.

The predisposing factors include Cardiac gene mutations involving several genes like TTNC1, TTN, and STAT3, Twin pregnancy/ multifetal gestation, Smoking, and Preeclampsia. Other proposed causes include viral myocarditis, abnormal immune response to pregnancy, malnutrition, inflammation, etc.

PPCM is associated with increased secretion of prolactin by the pituitary and antiangiogenic molecule sFlt-1 (soluble fms-like tyrosine kinase) by the placenta at term gestation.

Pearson's diagnostic criteria for PPCM:

• Development of cardiac failure in the last month of pregnancy or within 5 months after delivery

• Absence of an identifiable cause for the cardiac failure

• Absence of recognizable heart disease prior to the last month of pregnancy, and

• Left ventricular systolic dysfunction demonstrated by echocardiographic criteria, such as ejection fraction <45% or fractional shortening <30% along with a dilated left ventricle.

Management involves treating the features of heart failure. Bromocriptine has been found useful in this condition. Many women recover within 6 months of delivery, while some continue to have residual effects.

Solution to Question 21:

Minipills or progesterone-only pills (POP) do not kill sperm in the cervical canal. Progesterone-only pills act by thickening the cervical mucus (the main mechanism of all POPs except cerazette) that inhibits penetration of sperms, inhibiting tubal motility, and suppressing ovulation.

Other advantages of POPs include a reduction in the incidence of pelvic inflammatory diseases and sexually transmitted diseases (due to thickened cervical mucus).

Due to the inhibition of tubal motility, the absolute risk of all pregnancies including ectopic pregnancy decreases, while the relative risk of ectopic pregnancy increases. But a previous history of ectopic pregnancy is not a contraindication for POP.

POPs do not affect clotting, lipid metabolism, and carbohydrate metabolism. So they can safely be used by women who are obese, breastfeeding, hypertensive, have SLE or a history of thromboembolism, diabetic with vascular disease. Significant adverse effects of POPs are irregular bleeding (mostly scanty), and secondary amenorrhea after longtime usage as it causes endometrial atrophy.

Contraindications for POPs include undiagnosed vaginal bleeding, pregnancy, known or suspected breast cancer, and benign or malignant liver tumors.

Solution to Question 22:

Previous manual removal of the placenta does not indicate a high-risk pregnancy.

The high-risk pregnancies according to the Pregnancy Risk Assessment Monitoring System by the CDC are:

Current pregnancy complications	Obstetric history	Other disorders
Preterm labor (<37 Wks) and Post-term (> 42 wk)	Intrauterine growth restriction	Cardiovascularfactorslike chronic hypertension, severe heart failure, andrenaldisorders
Preterm premature rupture of membranes	Stillbirth	Metabolicdisorderslike Obesity,Insulin-dependent diabetes, Previous endocrine ablation, Thyroid disorders

Current pregnancy complications	Obstetric history	Other disorders
Moderate to severe preeclampsia	Late abortion (16–20 wk)	Hematologicalproblems like sickle cell disease, severe anemiaThrombophilia
Polyhydramnios (hydramnios) or oligohydramnios	Fetal exchange transfusion (Rh incompatibility)	Infections(malaria, HIV) and Autoimmunedisorders
Uterine rupture	Multifetal pregnancy	Epilepsy
Antepartum hemorrhage (Placenta previa, and Abruptio placenta)	Previous brachial plexus injury	Anatomicabnormalitieslike uterine malformationand cervical incompetence
Fetal tachycardia or bradycardia for >30 min	Abnormal fetal position	Exposure toteratogenslike alcohol, drugs, smoking
Fetal weight < 2.5 kg or >4kg	Previous h/o myomectomy, repair of complete perineal tear, vesicovaginal fistula, and stress incontinence	Familyhistory of diabetes, hypertension ormultiple pregnancy and congenital malformation
Chorioamnionitis		Lowsocioeconomic status
Prolapsed cord		Abnormal cervical cytologic findings
Fetal acidosis pH ≤7		

Solution to Question 23:

Long-standing pelvic inflammation may lead to pyometra. This refers to the collection of pus in the uterine cavity.

The initial infection involves the cervix uteri and urethra which may extend to Bartholin glands, skene glands, endometrium, and fallopian tubes. Unilateral pain and swelling over the labia may be caused by the inflammation of the Bartholin glands.

Depending on the extent of tissue involved, PID can be classified into five stages

• Stage I - Acute salpingitis without peritonitis - No adhesions

• Stage II - Acute salpingitis with peritonitis - purulent discharge

• Stage III - Acute salpingitis with superimposed tubal occlusion or tubo ovarian complex

• Stage IV - Ruptured tubo-ovarian abscess

• Stage V - Tubercular salpingitis

Long-standing pelvic inflammation may also lead to:

• Dyspareunia

• Infertility

• Ectopic pregnancy

• Formation of adhesions like hydrosalpinx, pyosalpinx, tubo-ovarian abscess

• Chronic pelvic pain

Note: Chronic pelvic inflammation leads to dense adhesions between the uterus and the appendages. Thus, it becomes difficult to define the uterus from the pelvic masses. They form a hard mass which is called the frozen pelvis.

The most common cause of the spread of PID: Ascending infection along with sperms

The only PID that does not spread with the sexual route is genital TB. It has a hematogenous spread.

The most common PID in virgin females is genital TB.

Investigation of choice in PID: Abdominal USG

The surest sign of salpingitis is pus extruding from the fimbrial end of the tube.

Solution to Question 24:

The most common postpartum nerve injury is an injury to the peroneal nerve causing foot drop.

The lateral femoral cutaneous nerve is the most commonly injured nerve after vaginal delivery followed by femoral neuropathies.

Obstetrical neuropathies are due to the pressure on branches of the lumbosacral nerve plexus during labor. They can manifest as complaints of intense neuralgia or cramp-like pains extending down one or both legs as soon as the head descends into the pelvis. Footdrop can be secondary to injury at the level of the lumbosacral plexus, sciatic nerve, or common fibular (peroneal) nerve. Injury of the iliohypogastric and ilioinguinal nerves may occur with cesarean delivery, which may manifest as an inguinal hernia and loss of sensation at the pubic region respectively.

Note: Since there are no options relating to intense neuralgia in the lower limb, foot drop becomes the best possible answer.

Solution to Question 25:

Total T_3 levels are altered throughout pregnancy.

Total T_3 and total T_4 increase due to stimulation of thyroid tissue by hCG (α-subunit of hCG is similar to TSH) during the early period of gestation. However, there is no change in free T_3 and T_4 due to an increase in thyroid-binding globulin (TBG).

This leads to a transient fall in TSH (but remains in the normal range) during the early weeks of gestation. The level of TSH becomes normal during the second and third trimesters.

The hormonal changes during pregnancy are as follows:

Increased	Decreased	Unchanged
Growth Hormone	LH (Luteinizing hormone)	FreeT_3, T_4

Increased	Decreased	Unchanged
Insulin(due toinsulin resistance)	FSH (Follicle-stimulating hormone)	TSH
Prolactin	DHEAS (Di-hydro epiandrosterone)	ADH (Anti- diuretic hormone)
TotalT3, T4		
Aldosterone		
Testosterone, Androstenedione, and cortisol		

Obstetrics and Gynecology NEET 2019

Question 1:

In which trimester does acute fatty liver manifest most commonly during pregnancy?

a) First trimester

b) Second trimester

c) Third trimester

d) Both (a) and (b)

Question 2:

Which of the following is true about vaginal changes in a normal pregnancy?

a) Decreased number of Lactobacilli

b) Increased glycogen metabolism in the epithelium

c) pH > 6.8

d) Epithelial thinning

Question 3:

You discover that there is evidence of cardiac malformation in a fetus during a routine USG scan at 16 weeks. What is the minimum level of fasting blood sugar at which you can suspect overt diabetes?

a) 106 mg/dL

b) 126 mg/dL

c) 116 mg/dL

d) 130 mg/dL

Question 4:

A 45-year-old woman presented with a lower abdominal mass and lower abdominal discomfort. On examination, the uterus appears uniformly enlarged and adnexa is free. MRI shows the presence of a myometrial cyst. What is the most probable diagnosis?

a) Endometriosis

b) Adenomyosis

c) Leiomyoma

d) Endometrial hyperplasia

Question 5:

Which of the following decreases in pregnancy?

a) Respiratory rate

b) Vital capacity

c) Functional residual capacity

d) Inspiratory capacity

Question 6:

Which of the following is not a contraindication for the induction of labor?

a) Pelvic tumor

b) Herpes infection

c) Heart disease

d) History of lower transverse cesarean section in last three pregnancies

Question 7:

The given image shows what grade of placenta previa?

a) IV

b) III

c) II

d) I

Question 8:

All are normal physiological changes in pregnancy except:

a) Left axis deviation

b) Mild ST changes in inferior leads

c) Atrial and ventricular premature contractions

d) Pansystolic murmur

Question 9:

A pregnant woman with a fibroid uterus develops severe acute pain abdomen at 28 weeks of gestation. There is no history of fever and the WBC count is 8000 cells/mL. The most likely diagnosis is:

a) Red degeneration of fibroid

b) Preterm labour

c) Torsion of fibroid

d) None of the above

Question 10:

What is the first line of treatment of mastitis in a lactating mother?

a) Dicloxacillin

b) Cefazolin

c) Ceftriaxone

d) Ampicillin

Question 11:

Major contributor for amniotic fluid post 20 weeks is?

a) Fetal urine

b) Fetal skin

c) Fetal lung fluid

d) Maternal plasma

Question 12:

What will be the level of the anti-Mullerian hormone in low ovarian reserve?

a) <1

b) 1-4

c) >7

d) >10

Question 13:

The maximum dose of PGF2 alpha in postpartum hemorrhage is ____.

a) 0.25 mg

b) 2 mg

c) 20 mg

d) 200 mg

Question 14:

Gestational trophoblastic neoplasm does not include

a) Partial mole

b) Choriocarcinoma

c) Placental site trophoblastic tumor

d) Invasive mole

Question 15:

A baby was born at 34 weeks of pregnancy, the appearance of the baby is given below. What is the mother suffering from ?

a) Gestational diabetes

b) Rubella

c) Hypothyroidism

d) Gestational hypertension

Question 16:

The hormone acting on post ovulatory endometrium is :



a) Luteinizing hormone

b) Follicular stimulating hormone

c) Progesterone

d) Oestrogen

Question 17:

Which of the following is an absolute contraindication to OCP use?

a) Chronic renal disease

b) Thromboembolism

c) History of amenorrhea

d) Diabetes mellitus

Question 18:

What is the chromosome number of a partial hydatidiform mole?

a) 46 XX

b) 45 XY

c) 69 XY

d) 69 XXY

Question 19:

A 55-year-old lady presents with third degree uterine prolapse. Which of the following is the best management for her?

a) Fothergill repair

b) Sling operation

c) Vaginal hysterectomy with pelvic floor repair

d) Shirodkar procedure

Question 20:

What is the dose of Ulipristal used in emergency contraception?

a) 30 mg

b) 300 mg

c) 60 mg

d) 600 mg

Question 21:

What is the ideal time for nuchal translucency evaluation?

a) 6-8 weeks

b) 11-14 weeks

c) 14-16 weeks

d) 16-20 weeks

Question 22:

Magnesium sulphate has no role in:

a) Prevention of seizures in severe pre-eclampsia

b) Prevention of recurrent seizures in eclampsia

c) Prevention of RDS in premature baby

d) Reducing contractility of uterus

Question 23:

Which of the following is seen in the vertex presentation with complete flexion attitude of head?

a) Suboccipito-bregmatic diameter

b) Suboccipito-frontal diameter

c) Occipito-frontal diameter

d) Occipito-posterior position

Question 24:

A pregnant lady with a ■history of recurrent abortions is now diagnosed to have antiphospholipid syndrome. What would be the best mode of treatment for her?

a) Aspirin only

b) Aspirin+heparin

c) Aspirin+heparin+steroids

d) Aspirin+steroids

Question 25:

A 22-year-old pregnant lady presented with a fever and was started on antibiotics. The fever did not subside. On further investigation, she was diagnosed to have septic pelvic thrombophlebitis. Which among the following is the next course of treatment for her?

a) Stop antibiotics and start heparin

b) Antibiotics+Heparin

c) Surgical embolectomy

d) Hysterectomy

Question 26:

What is the treatment of choice for a woman with atrophic vaginitis with complaints of itching?

a) Steroids

b) Topical estrogen

c) Antihistamines only

d) None of the above

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	b
5	c
6	c
7	b
8	d
9	c
10	a
11	a
12	a
13	b
14	a
15	a
16	c
17	b
18	d
19	c
20	a
21	b
22	c
23	a
24	b
25	b
26	b

Detailed Explanations

Solution to Question 1:

Acute fatty liver of pregnancy (AFLP) occurs most commonly in the third trimester or early postpartum period.

Risk factors include Pre-eclampsia/HELLP syndrome, male fetus, previous history of AFLP, and multiple gestations.

AFLP (acute yellow atrophy or acute fatty metamorphosis) is the most common cause of acute liver failure in pregnancy. It usually occurs in the later stages of pregnancy, particularly in the third trimester. It occurs due to fetal deficiency of long-chain acyl-coenzyme A dehydrogenase (LCHAD) or poor placental mitochondrial function.

The patients present with persistent nausea and vomiting, progressive jaundice, and epigastric pain. Features like acute hemolysis, hypoglycemia, hypertension, proteinuria, thrombocytopenia, and edema can also be present. Symptoms may progress to fulminant hepatic failure with encephalopathy, disseminated intravascular coagulation, and death.

Liver function tests such as AST, ALT, and serum bilirubin levels are raised. Hypofibrinogenemia, hypocholesterolemia, and prolonged clotting time can be seen. An elevated d-dimer is seen. Microscopy reveals a characteristic microvesicular fatty infiltration.

The condition worsens till delivery after which the hepatic dysfunction resolves. A trial of labor by induction or delivery by Cesarean is done. Intensive supportive measures including blood transfusion should be provided.

The clinical and laboratory findings of acute liver diseases in pregnancy are as follows:

The condition worsens till delivery after which the hepatic dysfunction resolves. Appropriate management is immediate termination of pregnancy irrespective of gestational age by induction or delivery by Cesarean (for obstetric reasons) is done. Intensive supportive measures including blood transfusion should be provided.

Disorders	HELLP syndrome	Obstetric cholestasis	Acute fatty liver of pre gnancy	Viral hepa titis
Onset in pregnanc y	Mid to late	Late	Late	Variable
Differentiating fea tures	Epigastric pain, hypert ension,elevatedindirect bilirubin	Prurituselevateddirect bilirubin	Signs ofliver failurelike hypoglycemia	Fever
AST (U/L)	≤300	≤200	200-800	2000+
Bilirubin (mg/dL)	1-4	1-5	4-10	5-20
Platelets	↓↓	Normal	↓↓	↓
Fibrinogen	Normal	Normal	↓↓↓	Normal
D-dimer	↑	Normal	↑	Normal
Prothrombin time	Normal	Normal	↑↑	↑
Hemolysis	↑↑	No	↑↑↑	No

Solution to Question 2:

There is an increase in the metabolism of glycogen in the vaginal epithelium during normal pregnancy.

The changes in the vagina during pregnancy include:

• Increased vascularity and hyperemia resulting in a violet/purplish color characteristic of Chadwick's sign

• Vaginal epithelium undergoes marked proliferation (option D) with the extension of the squamocolumnar junction downwards

• Vaginal pH decreases and becomes more acidic (option C) and varies from 3.5-6.0.

• Increase in the number of Lactobacillus acidophilus (option A). It metabolizes the glycogen stores and causes an increased production of lactic acid. Therefore, the pH becomes acidic.

• Increased volume of cervical secretion during pregnancy results in the production of thick white vaginal discharge.

Solution to Question 3:

Overt diabetes is when fasting blood sugar is ≥ 126 mg/dl. Overt diabetics are women who are known to have diabetes before pregnancy.

A patient is said to have overt diabetes if:

• Fasting blood sugar ≥ 126 mg/dL

• A random blood sugar of ≥ 200 mg/dL

• HbA1c of ≥ 6.5%

The maternal complication of diabetes:

• increased Miscarriage and preterm labor.

• increased incidence of preeclampsia

• polyhydramnios

• prolongation of labor due to shoulder dystocia, operative interference, and cephalo-pelvic disproportion.

Fetal and Neonatal complications of diabetes:

• fetal macrosomia (birth weight > 4kg)

• congenital malformations - Musculoskeletal, urogenital, and CNS anomalies.

Solution to Question 4:

Multiparous women >40yrs, presenting with abdominal mass (free from adnexa) and myometrial cysts on MRI, point towards a diagnosis of adenomyosis.

Adenomyosis refers to the growth of the endometrium (both stromal and glandular components) directly into the myometrium. It is predominantly seen in multiparous women above the age of 40 years. Patients generally present with menorrhagia, dysmenorrhea, and sometimes dyspareunia. On examination, there is a gross symmetrical enlargement of the uterus corresponding to 12-14 weeks size. The uterus is soft and tender on bimanual examination.

Transvaginal ultrasonography (TVS) reveals a salt-and-pepper appearance of the myometrium and the presence of sub-endometrial striations producing the Venetian blind appearance. MRI can also be done for diagnosis. However, the only definitive mode of diagnosis is histopathological examination.

Since most women affected by the condition are past the childbearing age, total hysterectomy is the treatment of choice. Localized excision/Levonorgestrel containing IUCD can be tried in younger women.



Other options:

Option A: Endometriosis refers to the presence of endometrial tissue outside the endometrial lining (uterus). They usually present as dysmenorrhea and lower dull abdominal pain. Vaginal examination reveals a tender fixed retroverted uterus. They can be managed medically (oral contraceptives, aromatase inhibitors, etc) and surgically (if failed medical management). Abdominal hysterectomy with bilateral salpingo-oophorectomy is performed in advanced cases.

Option C: Leiomyoma (fibroid) is the commonest benign tumor arising from the uterus. They present as abnormal uterine bleeding, infertility, and abdominal pain. On bimanual examination, the uterus is enlarged and the mass moves with the cervix. It can be managed medically (mifepristone or newer progesterone) and surgically. A myomectomy is performed for large fibroids with significant symptoms.

Option D: Endometrial hyperplasia can be simple, glandular, or atypical. They are seen in women with estrogen excess states like obesity, PCOD, or on tamoxifen therapy. They usually present as abnormal uterine bleeding (AUB).

Solution to Question 5:

Functional residual capacity decreases in pregnancy.

The changes in the respiratory system during pregnancy are:

(Mnemonic: TIMM increases in pregnancy)

Increase	Decrease	Unaffected
Tidal volumeInspiratory capacityMinute O2uptakeMinute ventilation	Functional residual capacityExpiratory reserve volumeResidual volumeTotal lung capacity	Respiratory rateVital capacity Inspiratory reserve volume

Solution to Question 6:

Heart disease is currently not considered a contraindication for the induction of labor. Vaginal delivery is preferred in such patients.

Solution to Question 7:

The given image shows grade III placenta previa.

Placenta previa refers to the placenta implanted (partially/completely) over the lower uterine segment (over or adjacent to the internal os). Grades of placenta previa are:

• Type—I (Low-lying): The major part of the placenta is attached to the upper segment and only the

lower margin encroaches onto the lower segment but not up to the os.

• Type—II (Marginal): The placenta reaches the margin of the internal os but does not cover it.

• Type—III (Incomplete or partial central): The placenta covers the internal os partially (covers the

internal os when closed but does not entirely do so when fully dilated).

• Type—IV (Central or total): The placenta completely covers the internal os even after it is fully dilated.



They typically present with sudden onset, painless vaginal bleeding. On vulval inspection, bright red blood is seen oozing from the cervical opening. Transvaginal ultrasound (TVS) is used to confirm the diagnosis and position of the placenta.

Management is dependent on the presentation of the patient. If there is a need for fetal maturity, expectant management (advocated by McAfee and Johnson) is carried out without increasing undue maternal stress. Definitive management by an elective cesarean section.

Solution to Question 8:

Physiological changes in pregnancy

Clinical examination

• Jugular venous distension and venous hum

• Mammary soufflé: Continuous murmur at 2nd to 4th intercostal space

• Occassional S3at Mitral area

• Aortic or pulmonary flow murmurs at aortic/pulmonary areas

• S2 P increased; S2 split

• S1 M increased and widely split

Electrocardiography

• T wave inversion, biatrial enlargement

• Dysrhythmias

• As the diaphragm is elevated in advancing pregnancy, there is an average 15-degree left-axis deviation in the electrocardiogram (ECG), and mild ST changes may be seen in the inferior leads.

• Atrial and ventricular premature contractions are relatively frequent

• Pregnancy does not alter voltage

• Tricuspid regurgitation

• Increased left atrial end-diastolic dimension and

• Increased left ventricular mass

Solution to Question 9:

The given scenario of severe acute abdominal pain without fever/leucocytosis in a pregnant woman with a fibroid is suggestive of fibroid torsion.

The incidence of fibroid in pregnancy is about 1 in 1,000. Due to increased vascularity in pregnancy, the fibroid increases in size and becomes softer. The torsion is seen in the subserous pedunculated fibroid where the venous return is blocked at its site of attachment in the uterus. This leads to severe acute abdominal pain. Due to the complication being more local than systemic, fever or leucocytosis are generally not seen.

A transvaginal ultrasonogram is immediately done to detect the position and size of the fibroid, which is followed by laparotomic myomectomy as a part of emergency surgical management.

Other options:

Option A: Red degeneration of fibroid is seen commonly in the second half of pregnancy or puerperium. It is due to a hemorrhagic infarction of the tissue and typically presents as acute focal abdominal pain with fever and leucocytosis. Treatment is usually conservative with bed rest, analgesics, and sedatives.

Option B: Preterm labor is where the labor starts before 37 completed weeks. It is characterized by true labor pain. The abdominal pain is slowly increasing in intensity and duration and is also accompanied by show (expulsion of the cervical mucus plug). There are intervals between episodes of contractions where the patient doesn't complain of abdominal pain.

Solution to Question 10:

Dicloxacillin is the first line of treatment for mastitis in lactating mothers and should be taken for 10-14 days.

Mastitis refers to the infection of the breast parenchymal tissue. The most common organism causing breast infection is Staphylococcus aureus. It is common in new mothers at the beginning of lactation who are inexperienced with breastfeeding or during weaning when nipple trauma occurs due to the infant's growing teeth.

Clinical features of mastitis include:

• Fever and malaise

• Localized pain and tenderness

• Purulent nipple discharge

• Redness and swelling of the affected breast

• Positive fluctuation test

A culture from nipple discharge or test-needle aspiration of the cellulitic area is done to confirm the organism and its antibiotic sensitivity. An ultrasonogram of the involved breast is done to exclude abscess. They are treated with antibiotics and adequate breast support. Choice of antibiotics in lactational mastitis:

• Dicloxacillin 500 mg 4 times a day is to be started empirically.

• Erythromycin is given to women who are penicillin-sensitive.

• If the infection is caused by resistant, penicillinase-producing Staphylococcus species then vancomycin, clindamycin, or trimethoprim-sulfamethoxazole is given for at least 10 days.

Solution to Question 11:

Fetal urine is the major contributor to amniotic fluid post 20 weeks.

Amniotic fluid is present in the amniotic cavity which is the space between the epiblast (ectoderm) and trophoblast. The volume of the amniotic fluid increases with pregnancy and reaches a maximum at term. It is contributed by:

• Early pregnancy - ultrafiltrate of maternal plasma (option D).

• Second trimester - the extracellular fluid that diffuses through the fetal skin (option B) and thus reflects the composition of fetal plasma.

• After 20 weeks - keratinization of fetal skin prevents diffusion across the skin. Hence, fetal urine forms amniotic fluid.

Note: Fetal kidneys start producing urine at 12 weeks (8–11 weeks), and by 18 weeks, they produce 7 to 14 mL per day.

Solution to Question 12:

In low ovarian reserve, the level of anti-Mullerian hormone (AMH) will be <1 ng/ml.

AMH is a glycoprotein secreted by the granulosa cells of the primordial follicle of the ovary. It helps in oocyte maturation and mono-follicular development. AMH is also used in assessing ovarian reserve.

In women, AMH is always absent during embryogenesis and is secreted by the granulosa cells after puberty. It helps in follicular development and oocyte maturation. Estimation of serum AMH is used in the study of ovarian reserve in female infertility and secondary amenorrhoea.

In men, AMH starts to be secreted by the Sertoli cells in the testes by the 7th week of intrauterine life. It continues until puberty. It inhibits the development of the Müllerian system. The absence of AMH results in the development of hermaphroditism.

AMH levels in ng/ml	Interpretation
2–6.8	Normal

AMH levels in ng/ml	Interpretation
<1	Poor ovarian reserve
>10	PCODHyperstimulation syndrome

Solution to Question 13:

The maximum dose of prostaglandin F2 alpha in postpartum hemorrhage is 2 mg.

Prostaglandins are paracrine/autocrine hormones and act locally at the site of production. Main sources for:

• PGF2α - Decidua and promotes myometrial contractility

• PGE2 - fetal membranes (amnion) and induces labor and promotes cervical effacement and dilatation

• PGI2 - Myometrium

Carboprost tromethamine (hemabate) is the 15-methyl derivative of prostaglandin F2 alpha. It is approved for the treatment of uterine atony at a dose of 0.25 mg IM. This dose can be repeated if necessary at 15- to 90-minute intervals up to a maximum of eight doses (2 mg).

It is the best drug to control PPH. ls also used an abortifacient in the second trimester.

Contraindications include asthmatic women and those with suspected amniotic fluid embolism, as they cause pulmonary airway and vascular constriction.

Solution to Question 14:

A partial mole is not a gestational trophoblastic neoplasm (GTN).

Gestational trophoblastic disease (GTD) is the term used to describe a group of tumors due to abnormal trophoblast proliferation. Malignant forms of GTD are termed gestational trophoblastic neoplasia (GTN).

Modified WHO Classification of GTD

Molar Pregnancies	Trophoblastic tumors
Hydatidiform mole (benign) - PartialandCompleteInvasive mole (malignant)	Choriocarcinoma (malignant) Placental site trophoblastic tumor (malignant)Epithelioid trophoblastic tumor (malignant)

Solution to Question 15:

The image provided depicts a baby with macrosomia (large for gestational age), the most probable cause of which is gestational diabetes (GDM) in the mother.

GDM in a mother is defined as carbohydrate intolerance with onset/first recognition during the present pregnancy. Diabetes in Pregnancy Societies of India (DIPSI) criteria is used to diagnose GDM (Venous plasma >140mg % after 2h of ingestion of 75g of glucose).

Maternal Complications of GDM:

• During pregnancy - Recurrent spontaneous abortion, preterm labor, urinary tract infection, polyhydramnios, and increased incidence of preeclampsia

• During labor: Prolongation of labor (shoulder dystocia) and postpartum hemorrhage

Neonatal complications -

• Electrolyte imbalances - hypoglycemia, hypocalcemia, hypokalemia, hypomagnesemia

• Respiratory system - respiratory distress syndrome and delayed lung maturity

• Hematological system - polycythemia and single umbilical artery

• Renal system - hydronephrosis (renal agenesis) and polycystic kidneys

• Cardiovascular system - VSD, ASD, Coarctation of the aorta, and TGA

• Gastrointestinal system - duodenal atresia, tracheoesophageal fistula (TEF), and necrotizing enterocolitis

• Central Nervous system - anencephaly, microcephaly, neural tube defects, caudal regression syndrome, and sacral agenesis

Other options:

Option B: Congenital rubella syndrome presents with the classical triad of deafness, cataract, and congenital heart disease (the most common being - patent ductus arteriosus). Eye findings also include salt and pepper retinopathy as shown below.

Option C: Congenital hypothyroidism is the most common preventable cause of mental retardation and presents as 6 P's - pot-bellied, pale, puffy-faced child with protruding umbilicus, protuberant tongue, and poor brain development

Option D: Gestational hypertension primarily occurs due to the failure of the invasion of extravillous trophoblasts into the spiral artery. It is defined as a rise in blood pressure >140/90 mmHg after 20 weeks of pregnancy (without proteinuria). It usually leads to intrauterine growth restriction (IUGR) and fetal death.

Solution to Question 16:

The hormone acting on the post-ovulatory endometrium is progesterone.

Endometrium undergoes changes (endometrial cycle) during the menstrual cycle due to the action of various hormones. The basal zone (lower 1/3rd of the endometrium in contact with the myometrium) is uninfluenced by hormones and shows no change during the cycle. The rest

(functional zone) shows the following changes:

• Regenerative phase (first 2 to 3 days)- the regeneration of glands and the stromal cells from the basal zone. Endometrial thickness is about 2mm.

• Proliferative phase (5 or 6th day to 14th day) - estrogen (option D) causes the glands to become more tubular and perpendicular to the surface and increases the thickness to 10-12mm on the 14th day (at the time of ovulation)

• Secretory phase(15th day till menstrual bleeding) - post ovulation, estrogen-primed endometrium is acted by progesterone (from corpus lutetium) and causes increased blood flow (spiraling of arteries) and increases tortuosity of glands (saw-toothed/cork-screwed glands). The thickness of the endometrium is maximum (12mm) in this phase.

• Menstruation (last 4 or 5 days) - estrogen and progesterone are withdrawn due to the regression of corpus luteum (lack of fertilization). This results in the shedding of the functional zone (menstrual bleeding).



Other options:

Option A: The luteinizing hormone (LH) is secreted by the anterior pituitary and is crucial for ovulation to occur on the 14th day of the menstrual cycle.

Option B: Follicular stimulating hormone (FSH) is secreted by the anterior pituitary. The ovarian follicle with the most sensitivity to FSH becomes the dominant follicle for the current cycle and the rest undergo autolysis.

Solution to Question 17:

Thromboembolism is an absolute contraindication to OCP use.

Oral contraceptives (OCP) are preparations of estrogen or progesterone or a combination of both. Estrogen in OCPs suppresses gonadotropin-releasing hormone (GnRH) and follicle-stimulating hormone (FSH) whereas progesterone suppresses luteinizing hormone (LH), thereby inhibiting

ovulation.

Though OCPs are generally used to prevent pregnancy, they are also indicated for the treatment of dysmenorrhoea, endometriosis, uterine fibroid, and dysfunctional uterine bleeding (DUB)

Contraindications of OCPs:

• Absolute contraindications (WHO category 4)-

• History of thromboembolism- DVT, stroke

• Heart diseases - Valular or ischemic

• Severe hypertension

• Liver diseases like hepatic adenoma, and hepatic carcinoma

• Breastfeeding

• During pregnancy

• Estrogen dependant neoplasms like - breast cancer

• Relative contraindications -

•

WHO category-2 (benefits outweigh the risks) - cancer cervix, cervical intraepithelial neoplasia (CIN), and in patients with sickle cell anemia, disease of the gallbladder, or diabetes mellitus (option D).

• WHO category-3(risks outweigh benefits) - patients with benign liver tumors, unexplained vaginal bleeding, or history of smoking >20 cigarettes/day.

Other options:

Option A: Minipills (progestin-only pills) could be used safely in renal disease as they reduced clotting risks and is not an absolute contraindication.

Option C: A history of amenorrhea is investigated based on the algorithm for primary or secondary amenorrhoea. OCPs are used to regularise menstruation in polycystic ovarian syndrome (PCOD) which is a cause of secondary amenorrhoea.

Solution to Question 18:

Partial hydatidiform mole has a 69XXY or 69XYY karyotype.

Hydatiform mole belongs to the benign group of gestational trophoblastic diseases (GTD). It is an abnormality in chorionic villi making them partly proliferative and degenerative, thereby resulting in the formation of multiple small cysts. Based on the portion of the chorionic villi involved, they could be complete (diffused) or partial (focal)

Solution to Question 19:

In the clinal scenario of a 55-year-old woman with third-degree uterine prolapse, the ideal management is vaginal hysterectomy with pelvic floor repair.

Uterine prolapse is the descent of the uterus from its normal position which usually occurs due to laxity of ligaments that offer support, seen often due to childbirth, aging, or trauma. The uterine descent is clinically categorized into:

• First degree: the external os remains above the vaginal introitus

• Second degree: The external os protrudes below the vaginal introitus but the uterine body remains inside the vagina

• Third degree (Syn: Complete prolapse): The uterine, cervix, body, and fundus descend to lie outside the introitus

• Procidentia involves prolapse of the uterus with the entire vagina everted.

The patient usually presents with a sense of heaviness in the pelvic region with lower back discomfort. Based on the structures involved the patient can additionally have urinary symptoms (like incontinence and urgency), bowel disturbances (constipation and tenesmus), or sexual discomfort during intercourse.

Symptomatic uterine prolapse is managed surgically with different procedures based on the profile of the patient. Conservative management is done for old asymptomatic women with mild degree prolapse.

A vaginal hysterectomy is recommended for patients who satisfy the following criteria:

• >40 years of age

• Have completed their families

• Do not wish to preserve menstrual and child-bearing function

Other options:

Option A: Fothergill repair or Manchester operation is a procedure done to preserve the reproductive function and vaginal prolapse with supravaginal cervix elongation. It initially involves preliminary curettage with low amputation of the cervix. After Mackenrodt's ligament is plicated, anterior colporrhaphy and colpoperineorrhaphy are performed.

Option B: Sling operation (Purandare's operation) is indicated in congenital or nulliparous prolapse without cystocele. The procedure involves the extraperitoneal passage of rectus sheath strips joined to the anterior surface of the cervix by silk sutures.

Option D: Shirodkar procedure is a type of abdominal sling surgery done in young women to retain menstrual and reproductive function. It involves using a sling to fix the cervix to the lumbosacral fascia.

Solution to Question 20:

The dose of Ulipristal in emergency contraception is 30 mg to be taken within 72 hrs (upto 120hrs).

Methods of Emergency contraception

Drug	Dose
Levonorgestrel	0.75mg stat and after 12 hours
Ethinyl estradiol 50 µg + Norgestrel 0.25 mg	2 tab stat and 2 tab after 12 hours
Conjugated estrogen	15 mg BD x 5 days
Ethinyl Estradiol	2.5 mg BD X 5 days
Copper IUDs	Insertion within 5-7 days after accidental unprotected exposure.Not suitable for rape victims
Centchroman	2 tablets(60 mg) to be taken twice at an interval of 12 hours within 24 hours of intercourse.
Ulipristal	30 mg within 72 hours

Solution to Question 21:

Nuchal translucency evaluation, a component of first-trimester aneuploidy screening, is done between 11-14 weeks.

It represents the maximum thickness of the subcutaneous translucent area between the skin and soft tissue, overlying the fetal spine, at the back of the neck. When the nuchal translucency is increased, the risk of fetal aneuploidy and structural anomalies such as heart defects is significantly elevated.

First trimester aneuploidy screening combines two maternal serum analytes and a sonographic measurement:

• Human chorionic gonadotropin (hCG)

• Pregnancy-associated plasma protein A (PAPP-A)

• Nuchal translucency (NT)

It is performed between 11 and 14 weeks of gestation. In cases of fetal Down syndrome, the first-trimester serum-free β-hCG level is higher and the PAPP-A level is lower. With trisomy 18 and trisomy 13, levels of both analytes are lower.

Note: Nuchal fold is a normal fold of skin seen at the back of the fetal neck during the second trimester of pregnancy. Increased nuchal fold thickness is associated with fetal aneuploidy.



Solution to Question 22:

Magnesium sulfate has no role in the prevention of respiratory distress syndrome (RDS) in a premature baby, where there is a deficit of surfactant (which results in alveolar collapse).

Magnesium in the $MgSO_4$ competes with calcium on voltage-gated calcium channels, thereby altering its uptake and release. Functions of $MgSO_4$ include :

• Membrane protector and neuro stabilizer. It is used in the prevention of cerebral palsy in very low birth weight neonates.

• Reduction of motor endplate sensitivity to acetylcholine, and thus is used as an adjunct in the management of acute severe asthma in children and adults

• Blocking of neuronal calcium influx, and is thus used in the conservative management of pre-term labor (tocolytic, option D)

• Inducing cerebral vasodilatation, dilating the uterine arteries, increasing the production of endothelial prostacyclin, and inhibiting platelet activation. Hence it is used in the management of pre-eclampsia (option A) and to prevent progression to eclampsia (option A).

Solution to Question 23:

Suboccipito-bregmatic diameter is seen in vertex presentation with a complete flexion attitude of the head.

Diameters of the fetal skull:

Diameters	Attitude of the head
Suboccipitobregmatic	Complete flexion
Suboccipito-frontal	Incomplete flexion
Occipitofrontal	Marked deflexion
Mento-vertical	Partial extension
Submentovertical	Incomplete extension
Submentobregmatic	Complete extension

Solution to Question 24:

The best mode of treatment for a pregnant woman with antiphospholipid syndrome(APS) is aspirin and heparin.

APS is a prothrombotic disorder (autoantibody-mediated thrombophilia) affecting both arterial and venous systems.

Patients present with recurrent pregnancy loss due to thrombosis of uteroplacental vessels, poor placental perfusion, and prevention of normal decidual changes after zygote implantation. They may also present with preterm delivery, pre-eclampsia, intrauterine growth restriction (IUGR), and fetal death.

Criteria for APS include:

• at least one otherwise unexplained fetal death at or beyond 10 weeks;

• at least one preterm birth before 34 weeks gestation because of eclampsia, severe preeclampsia, placental insufficiency; or

• at least three unexplained consecutive spontaneous abortions before 10 weeks.

Aspirin (60-80mg daily) with heparin (5000-10,000 U subcutaneously every 12h) is the most effective therapy for APS. Treating APS with only aspirin (option A) is not recommended as heparin is crucial in preventing thrombus formation at the decidual-thrombotic interface, Corticosteroids (option C, D) should not be used in primary APS (without an associated connective tissue disorder) but are indicated in patients with pre-existing systemic lupus erythematosus (SLE) and APS (secondary APS).

Solution to Question 25:

Antibiotics plus heparin is the next course of treatment for septic pelvic thrombophlebitis.

Septic pelvic thrombophlebitis (SPT) refers to the pelvic infection with thrombosis of one/both ovarian veins. The pathogenesis of this disease is due to the spreading uterine infection

(secondary to trauma of delivery or surgery) combined with the hypercoagulable state of the pregnancy (thrombus formation).

The patients usually present with constant lower abdominal pain and occasionally with a palpable tender sausage-shaped mass (most diagnostic finding, but rare).

SPT is usually suspected when the fever fails to respond to antibiotic therapy. It can be confirmed by computed tomography (CT) or magnetic resonance imaging (MRI) of the abdomen and pelvis. Patients are treated with a combination of heparin and broad-spectrum antibiotics. Anticoagulation is done to prevent further thrombosis and reduce the spread of septic emboli.

Note: The use of anticoagulation is controversial as not all clinicians favor it. A small randomized study did not demonstrate a benefit, but the majority accept short-term anticoagulation as a part of the treatment regimen.

Solution to Question 26:

Atrophic vaginitis is treated with topical estrogen vaginal cream.

Atrophic vaginitis (senile vaginitis) is seen in postmenopausal women due to the thinning of vaginal mucosa (lack of estrogen). They present as discomfort (itching) and dryness in the vulva with dyspareunia. Occasionally, they may also complain of yellowish or blood-stained vaginal discharge.

The possibility of a coexisting carcinoma of the body or the cervix has to be excluded before treating it with estrogen supplements. Though topical application is the initial treatment, systemic estrogens (hormone replacement therapy) can be considered based on the patient's profile and the severity of the symptoms.

Other options:

Option A: Steroids are not indicated in the management of atrophic vaginitis as the inflammation is due to a lack of estrogen. Furthermore, they may predispose the patient to vaginal infections.

Option C: Antihistamines antagonize histamine receptors and prevent the allergic response. Since atrophic vaginitis is not an allergy response, they are not indicated in the treatment.

Obstetrics and Gynecology NEET 2020

Question 1:

The remnants of Wolffian ducts in females are located in_____.

a) Pouch of Douglas

b) Leaves of broad ligament

c) Uterovesical pouch

d) Iliac fossa

Question 2:

In which part of the fallopian tube does fertilization occur?

a) Ampulla

b) Isthmus

c) Interstitial

d) Infundibular

Question 3:

A patient came to the gynaecology OPD with complaints of foul-smelling frothy vaginal discharge and intense itching. On examination, the cervix was spotted and had the appearance of a strawberry. Identify the causative organism.

a) Trichomonas vaginalis

b) Gardnerella vaginalis

c) Neisseria gonorrhoea

d) Candida albicans

Question 4:

Identify the contraceptive shown in the given image.

a) Male condom

b) Female condom

c) Diaphragm

d) Cervical cap

Question 5:

A woman presents to the OPD with complaints of amenorrhoea for the past year. She has a history of postpartum hemorrhage and lactational failure. What is the probable diagnosis?

a) Sheehan's syndrome

b) Rotor syndrome

c) Prolactinoma

d) Acromegaly

Question 6:

An 18-year-old girl presents with a history of low-grade fever, weight loss, abdominal pain and amenorrhoea for six months. On examination, a pelvic mass is felt on the left side and signs of ascites are positive. What is your diagnosis?

a) Tuberculosis with tubo-ovarian mass

b) Ovarian malignancy

c) Ectopic pregnancy

d) Submucous fibroid

Question 7:

In modern obstetrics, which of the following investigations is used for monitoring the fetus of an Rh-negative sensitized mother?

a) Biophysical profile

b) Fetal blood sampling

c) Amniospectometry

d) MCA doppler peak systolic velocity

Question 8:

What is the causative agent for granuloma inguinale?

a) Calymmatobacterium granulomatis

b) Chlamydia trachomatis

c) Ureaplasma urealyticum

d) Treponema pallidum

Question 9:

Which of the following is the causative organism of vulval warts?

a) Human papilloma virus

b) Herpes simplex virus

c) Human immunodeficiency virus

d) Cytomegalovirus

Question 10:

A 22-year-old pregnant woman at 13 weeks of gestation informs you about accidental exposure to radiation from a chest X-ray at 7 weeks of gestation. What should be the next step advised to her?

a) Continue pregnancy

b) Terminate pregancy

c) Pre-invasive diagnostic test

d) Chromosome screening

Question 11:

After successful delivery of the baby, the obstetrician witnesses that the episiotomy performed has extended backwards beyond the perineal body. Which among the following structures is likely to be injured in this case?

a) External anal sphicter

b) Ischiocavernosus

c) Bulbospongiosis

d) Urethral sphincter

Question 12:

A 16 year old girl presents to the OPD with a history of primary amenorrhoea and cyclical pain. Perineal examination reveals a bulge protruding from the vagina. Identify the cause of this presentation:

a) Transverse septum in the proximal vagina

b) Complete vaginal atresia

c) Imperforate hymen

d) Cervical agenesis

Question 13:

A pregnant woman presents for a routine antenatal checkup at 18 weeks of gestation. On abdominal examination, the uterine size was found to correspond to 16 weeks of gestation. On USG, oligohydramnios was found. Which of the following is a likely cause?

a) Renal agenesis

b) Fetal anemia

c) Barter syndrome

d) Anencephaly

Question 14:

A 24-year-old woman visits the infertility clinic with a complaint of recurrent abortions. Upon investigations, she is found to have a septate uterus. Which of the following corrective procedures has the best obstetric outcome?

a) Jones metroplasty

b) Strassmann metroplasty

c) Tompkins metroplasty

d) Transcervical resection of septum

Question 15:

A 33-year-old female patient comes with complaints of menorrhagia for the past 3 months. The duration and regularity of her menstrual cycles are normal. Non-hormonal management has been tried without success. What is the most appropriate next step?

a) Uterine artery embolisation

b) Endometrial ablation

c) Hormonal therapy

d) Hysterectomy

Question 16:

A 60-year-old woman presented with post-menopausal bleeding. Ultrasonography revealed a mass with feeding vessel as shown in the image below. What is the most likely diagnosis?



a) Fibroid uterus

b) Endometrial polyp

c) Endometrial carcinoma

d) Leiomyosarcoma

Question 17:

A 35-year-old primigravida who conceived after in-vitro fertilization therapy came for an antenatal checkup at 38 weeks of gestation. Her obstetric history revealed that she was pregnant with DCDA twins. On examination, the first twin was found to be in a breech position and the maternal blood pressure was found to be greater than 140/90 mm Hg on two occasions with grade 1+ proteinuria. What should be done next?

a) Monitor BP and terminate pregnancy if BP rises

b) Immediate Caesarean section

c) Terminate pregnancy at 40 weeks of gestation

d) Induce labour with PGE2 gel

Question 18:

Which of the following effects of puberty are not mediated by estrogen?

a) Hair growth

b) Cervical mucus secretion

c) Vaginal cornification

d) Menstruation

Question 19:

A 35-year-old woman comes with complaints of warty lesions on the vulva. On examination, multiple warty growths on the vulva and around the anal canal are seen. This condition is most likely seen in?

a) Condyloma accuminata

b) Molluscum contagiosum

c) Chancroid

d) Herpes

Question 20:

A patient is planned to undergo hysteroscopic polypectomy using bipolar electrocautery. What is the ideal agent for distention?

a) Glycine

b) Normal saline

c) CO_2

d) 70% dextrose

Question 21:

A 52-year-old woman diagnosed with carcinoma cervix stage 2B was advised radiotherapy. Which of the following statements regarding radiotherapy is false?

a) Point B is 3 cm lateral to point A

b) Point A is 2 cm above the external os and 5 cm lateral to the internal uterine canal

c) Point A receives a dose of 7000 cGy

d) Point B receives a dose of 6000 cGy

Question 22:

A woman with a history of infertility is on treatment with human menopausal gonadotropin. She presented with the following finding in the USG. What is the most likely diagnosis?



a) Ovarian hyperstimulation syndrome

b) Theca lutein cyst

c) Polycystic ovarian syndrome

d) None

Question 23:

A lady who is 38 weeks pregnant comes to the OPD for a routine checkup. She has a history of normal twin delivery at term 4 years ago. What is her gravida and para score?

a) G2P2

b) G2P1

c) G3P2

d) G3P1

Question 24:

Identify the finding seen in this hysterosalpingogram:

a) Bilateral hydrosalpinx

b) Extravasation into venous system

c) Normal HSG

d) Bilateral cornual block

Question 25:

Identify the maternal condition that can lead to the findings in the baby as shown below.

a) Pre-gestational diabetes

b) Gestational diabetes mellitus

c) Effect of valproate

d) Hypertensive on ACE inhibitors

Question 26:

A woman with endometrial carcinoma is undergoing radiotherapy. Which of the following is true?

a) Intensity is inversely proportional to squared distance

b) Small blood vessels are most radioresistant

c) Rapidly proliferating cells are most radioresistant

d) Small bowel is most radioresistant

Question 27:

Identify the use of the instrument given below:

a) For tubal ligation

b) For ligation of uterine artery during hysterectomy

c) Ovarian cystectomy

d) Ectopic pregnancy

Question 28:

All of the following are contraceptive methods used post-coitus except:

a) Danazol

b) CuT200

c) RU486

d) Combined high dose estrogen-progestin pills

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	a

6	a
7	d
8	a
9	a
10	a
11	a
12	c
13	a
14	d
15	c
16	b
17	b
18	a
19	a
20	b
21	b
22	a
23	b
24	a
25	a
26	a
27	a
28	a

Detailed Explanations

Solution to Question 1:

The Gartner's duct is a remnant of the Wolffian ducts in women. These are located in the broad ligament and the vagina.

The Gartner's duct is an embryological remnant of the mesonephric duct/Wolffian duct formed during the development of the urinary and reproductive organs. It runs parallel to, but below, the fallopian tube in the mesosalpinx. The duct passes downwards by the side of the uterus to the level of the internal os where it passes into the tissues of the cervix. It then runs forwards to reach the anterolateral aspect of the vaginal wall and may reach as far down as the hymen.

The duct can sometimes form a cyst in the broad ligament or in the vagina. It is called Gartner's cyst. It is a benign condition and can be treated with marsupialization.



Solution to Question 2:

Fertilization usually occurs in the ampulla of the fallopian tube.

During ovulation, the ovum is released and it enters the fallopian tube. It is propelled towards the uterine cavity by the peristaltic action of the fallopian tubes, prostaglandins in the male seminal fluid and oxytocin. It can only be fertilized within 24 hours of ovulation. If coitus takes place during this period, the spermatozoa swimming up into the tubes meet the ovum in the ampulla.

Solution to Question 3:

In the given clinical scenario, Trichomonas vaginalis is the likely causative agent.

Vaginal itching and foul-smelling frothy vaginal discharge are seen in trichomonas vaginitis.

The given image shows colpitis macularis or strawberry cervix, which is a characteristic finding of trichomoniasis. This is due to microscopic, punctate haemorrhages of the cervix.

Solution to Question 4:

The given image shows a female condom. It consists of a pouch made of polyurethane which is 17 cm in length with a flexible ring at each end. The inner ring at the closed end is smaller than the outer ring.

The inner ring is inserted at the apex of the vagina and the outer ring remains outside. It covers the cervix, the entire vagina, and external genitalia. Hence, it provides protection against HIV and STDs. It has a failure rate of 5–15 per 100 woman-years, which is higher than that of a male condom.

It can be fitted by the patient hours before intercourse, and easily removed soon after. It covers the cervix, vagina, and external genitalia and is protective against sexually transmitted diseases. It can also be worn in the puerperal period, unlike the cervical diaphragm.

The female condom should not be used along with a male condom because if worn together they may slip, tear, or become displaced.

The given image below shows the insertion of a female condom:



Solution to Question 5:

The clinical scenario points to a diagnosis of Sheehan's syndrome.

Sheehan's syndrome (SS) is postpartum hypopituitarism secondary to the necrosis of the pituitary gland. It usually results from severe hypotension or shock caused by massive hemorrhage during or after delivery. Patients with SS have varying degrees of anterior pituitary hormone deficiency. The hyperplastic enlargement of the pituitary, which occurs normally during pregnancy, increases the risk of infarction.

Affected women may have persistent hypotension, tachycardia, hypoglycemia, and lactation failure. The most prominent symptoms are lactation failure (due to a deficiency of prolactin) and postpartum amenorrhea (due to a deficiency of gonadotropins). Because deficiencies of some or all pituitary-responsive hormones may develop after the initial insult, Sheehan syndrome can be heterogeneous and may not be identified for years.

It can present acutely as pituitary apoplexy, an endocrine emergency, characterized by a sudden onset of headache, nausea, visual deficits, and in severe cases,
severe hypoglycemia, hypotension and shock, CNS hemorrhage, and death.

Also, it can present late with the following features:

• Loss of pubic and axillary hair

• Hypothyroidism due to decreased TSH secretion

• Adrenal insufficiency due to low ACTH

Treatment is by hormone replacement - first glucocorticoids, later - thyroid hormone, sex steroids, vasopressin and growth hormones are given. If there is evidence of progressive visual loss, cranial nerve involvement or loss of consciousness, immediate surgical decompression must be carried out. Fertility can be achieved in these women with hMG and hCG. Clomiphene is ineffective.

Option B: Rotor syndrome is a type of hereditary hyperbilirubinemia with autosomal recessive inheritance. Affected individuals have mild jaundice and hyperbilirubinemia.

Option C: Prolactinoma or pituitary adenoids is the condition of hyperprolactinemia diagnosed by a prolactin level >200 μg/L. It usually presents with infertility, amenorrhea, and galactorrhea.

Option D: Acromegaly is characterized by the signs and symptoms pertaining to excessive growth hormone secretion.

Solution to Question 6:

The given clinical scenario with abdominal pain, amenorrhoea, pelvic mass and ascites is suggestive of genital tuberculosis.

Genital tuberculosis:

It is caused by M. tuberculosis and almost always occurs secondary to primary infection elsewhere in the extragenital states. The sites and frequency of involvement are as follows:

• Fallopian tubes (90-100%)

• Endometrium (50-60%)

• Ovaries (20-30%)

• Cervix (5-15%)

• Vulva and vagina (< 1%)

Presentation is variable and it can be asymptomatic in 10-20% of cases, infertility, menstrual irregularities, abdominal pain and mass, vaginal discharge, fistula formation, ectopic pregnancy and ascites. The mass is made up of rolled-up omentum that is adherent to the uterus and adnexa.

Constitutional symptoms such as fever, weakness, anorexia or night sweats may be present.

Investigations:

• CBC, ESR and Mantoux test

• Hysterosalpingography shows a lead-pipe appearance of the fallopian tube, calcification of the tube, beading and variation in the filling, cornual block, jagged fluffiness of tubal outline and tobacco pouch appearance of hydrosalpinx and pyosalpinx.

• Laparoscopy establishes the diagnosis of genital TB. It shows tuberculous lesions and perihepatic adhesions (if present).

Differential diagnosis:

• Ovarian cyst, broad ligament cyst or encysted fluid

• Pelvic inflammatory disease

• Ectopic pregnancy

• Carcinoma cervix

• Elephantiasis of the vulva

• Pregnancy

• Puberty menorrhagia and postmenopausal bleeding

• Fungal infection and sarcoidosis

Management:

It is treated with anti-tubercular therapy for 6 months (DOT strategy):

Indications and contraindications for surgery:

Types of surgery include vulvectomy (if vulval hypertrophy is present), removal of the adnexal mass, drainage of pyometra and fistula repair.

TAH+BSO is rarely required nowadays. Tuboplasty is contraindicated.

Option B: Ovarian tumors usually don't affect menstrual cycles. Though virilizing tumors can cause amenorrhoea, other features of virilization such as hirsutism, flattening of breasts, and enlargement of the clitoris are generally present.

Option C: Though ectopic pregnancy may be suspected here, the tube is likely to rupture with exacerbation of pain, bleeding, and peritonitis before six months of gestation. Additionally, the absence of vaginal bleeding/spotting, and the presence of weight loss makes the diagnosis unlikely.

Option D: Submucous fibroid is associated with menorrhagia.

First 2 months	Next 4 months
Isoniazid 15 mg/ kg body weight Rifampicin 450-600 mg Pyrazinamide 30 mg/body weight Ethambutol 30 mg/ kg body weight 3 times a week.	Isoniazid 15 mg/kg body weight Rifampicin 450-600 mg 3 times a week

Indications	Contraindications
Unresponsiveness of the disease in spite of adequate antitubercular chemotherapy. Tubercular pyosalpinx, ovarian abscess or pyometra. Persistent menorrhagia and/or chronic pelvic pain causing deteriorating health status	Presence of active tuberculosis in extragenital site Favorable response to antimicrobial therapy with the diminishing size of the pelvic mass The accidental discovery of tubercular tubo-ovarian mass on laparotomy in a young patient.

Solution to Question 7:

MCA Doppler peak systolic velocity (PSV) is used for the monitoring of fetuses in Rh-negative sensitized mothers.

The middle cerebral artery is used as other fetal vessels require a larger insonating angle and confers significant measurement error.

MCA PSV is enhanced due to increased cardiac output and decreased viscosity of fetal blood, which permits the reliable, noninvasive detection of fetal anemia in cases of blood-group

alloimmunization.

If the MCA PSV exceeds 1.5 MoM (multiples of the median) and the gestational age is less than 34 or 35 weeks, fetal blood sampling should be considered. MCA Doppler PSV has replaced amniocentesis as the preferred test for the detection of anemia.

Solution to Question 8:

The causative agent for granuloma inguinale (donovanosis) is Calymmatobacterium granulomatis. It is now called Klebsiella granulomatis and is a gram-negative bacillus causing chronic ulcerative infection of the vulva.

It is highly contagious and transmitted through repeated sexual or non-sexual contact. The incubation period is 1-12 weeks.

Clinical features:

It begins as a painless nodule which later ulcerates to form multiple beefy red painless ulcers that tend to coalesce; the vulva is progressively destroyed, and minimal adenopathy may occur.

Diagnosis:

Microscopic examination of smears from the lesion/ biopsy specimens reveals pathognomonic intracytoplasmic Donovan bodies and clusters of bacteria with a bipolar (safety pin) appearance. Blue-black stained organisms are seen in the cytoplasm of mononuclear cells.

Management:

Azithromycin 1 g orally once per week or 500 mg daily for at least 3 weeks and until all lesions have completely healed.

The given image below shows the donovanosis in a female patient with a beefy red ulcer:



Solution to Question 9:

The causative agent of vulval warts is the human papillomavirus (HPV).

Genital warts/venereal warts/condyloma acuminata are verrucous growths that affect the skin of the labia majora, perineum, perianal region and vulva. The lesions may be discrete or coalesce to form cauliflower-like growth. The disease is sexually transmitted. The growth of the warts is favoured by vaginal discharge, the use of oral contraceptives, and pregnancy.

HPV causes structural changes in the epithelial cells of the skin called koilocytes, and their presence is characteristic of HPV infection. The koilocytes have large hyperchromatic nuclei and a perinuclear halo, as shown below.



The diagnosis is made by colposcopic studies. The lesions on the vulva are stained by toluidine blue. The abnormal areas of the skin retain the blue dye. These areas are then biopsied for histopathological examination.

The given image below shows condyloma acuminata:

Solution to Question 10:

In the given clinical case, the patient should be advised to continue the pregnancy.

The radiation dose to the fetus during the declared term of pregnancy should be less than 1 mSv. In chest X-ray PA view, the absorbed dose of radiation by the fetus is less than 0.1 mSv. Hence, pregnancy can be continued.

An X-ray has a very limited role in obstetric imaging. There are very few indications where an X-ray will be useful:

• Fetal: congenital malformation of the fetus and neonates, skeletal malformations, birth injuries like fractures or dislocations.

• Maternal: Patients having cardiopulmonary disease may require an X-ray chest during pregnancy and that should be done beyond 12 weeks.

Radiation can cause fetal morbidity (FGR, genetic mutations, neurologic abnormalities, mental retardation, childhood leukemia) and mortality. The risks are high with radiation after the first 2 weeks postconception and within the first trimester (period of organogenesis).

Radiation during pregnancy can damage the early phase of the embryo but the risk can be minimized by using the '10-day rule'. This rule states that no woman should be exposed to an X-ray for a nonurgent indication outside 10 days from her last period during the reproductive period unless the pregnancy is excluded.

The given table below shows the absorbed radiation by the fetus in different diagnostic radiation procedures:

1 cGy = 10 mSv

Procedure	Dose (c Gy)
Abdominal X-ray	0.1-0.3
Pelvic X-ray	0.5-1.1
Chest X-ray	<0.001
Abdominal CT	0.4-0.8
Chest CT	0.002-0.02
Abdominopelvic CT	2.5-3.5
Ventilation scan	0.007-0.05
PET scan	1-1.5

Solution to Question 11:

External anal sphincter is the structure injured in an episiotomy extended posteriorly beyond the perineal body.

Perineal tear have been divided into four degrees:

• First degree- laceration is restricted to the skin of the fourchette

• Second degree- muscles of the perineal body are torn through

• Third degree- the tear extends partially backward through the external sphincter of the anus

• Fourth degree- the external anal sphincter is torn and anal mucosa is also involved

A central tear of the perineum is when the head penetrates first through the posterior vaginal wall, then through the perineal body and appears through the skin of the perineum. It usually occurs in patients with a contracted outlet.

Solution to Question 12:

A history of cyclical abdominal pain, primary amenorrhea and a bulging vaginal membrane suggest cryptomenorrhoea caused by an imperforate hymen.

Imperforate hymen is a urogenital anomaly which occurs due to failure of canalization of the inferior end of the vaginal plate. It usually remains unrecognized until puberty. It can lead to mucocolpos in early infancy.

Patients present with amenorrhea, cyclic abdominal pain, urinary symptoms, and a hypogastric lump on abdominal examination. A bluish bulging membrane at the vaginal opening is noted on genital examination and the Valsalva maneuver.

Ultrasound of the pelvis reveals normal internal genitalia and hematocolpos. Laparoscopy should be performed along with hymenectomy to exclude endometriosis.

Most common causes of cryptomenorrhea:

• Intact/imperforate hymen - a bluish convex bulge is seen

• Transverse vaginal septum - a pinkish concave covering is seen

Ultrasound pelvis confirms the diagnosis. A simple cruciate incision of the hymen allows the free drainage of the collected menstrual blood and restores the reproductive capacity in most cases.

Solution to Question 13:

Among the given options, renal agenesis is the likely cause. It causes oligohydramnios in the early second trimester.

Fetal urine production starts between 8-11 weeks of gestation but becomes a major component of amniotic fluid only in the second trimester. Thus, oligohydramnios is not seen until after 18 weeks of gestation in fetuses with lethal renal abnormalities.

Oligohydramnios occurs when the amniotic fluid index (AFI) is <5 cm or the single deepest vertical pocket is <2 cm. AFI between 5-8 cm is known as borderline oligohydramnios.

Diagnosis of oligohydramnios:

- Uterine size is much smaller than the period of amenorrhea
- Less fetal movements
- The uterus is 'full of fetus' due to scanty liquor
- Malpresentation (breech) is common
- Evidence of intrauterine growth retardation of the fetus is seen
- Sonographic diagnosis is made when the largest liquor pool is less than 2 cm

Other options:

Fetal anemia, Bartter syndrome and Liddle syndrome usually lead to polyhydramnios and not oligohydramnios.

Solution to Question 14:

Transcervical hysteroscopic resection of the septum has the best obstetric outcome.

The procedure involves the resection of the septum under hysteroscopic guidance. Hence, it doesn't scar the uterus and avoids the necessity of cesarean sections for future pregnancies improving obstetric outcomes.

Metroplasty is the surgical resection of the intrauterine septum. This is done by approaching the uterus transabdominally via 3 methods:

- Jones metroplasty
- Strassmann metroplasty
- Tompkins metroplasty

All of these methods are not preferred nowadays as they cause uterine scarring.

Solution to Question 15:

Hormonal therapy is the next best step in the given scenario of menorrhagia that is unresponsive to non-hormonal therapy.

Menorrhagia/heavy menstrual bleeding is defined as heavy but regular menstrual bleeding, >80mL of blood loss, or >8 days of bleeding per menstrual period. It can be acute (<6 months) or chronic (>6 months).

It is managed as follows:

- Medical - first line of treatment.
- NSAIDs

• Tranexamic acid

• OCPs/progestogens - if non-hormonal management fails

• Mirena IUCD

• GnRH agonists

• Blood transfusion, if required

• Surgical - if medical management fails, depending on the diagnosis:

• Endometrial ablation - hysteroscopic, TCRE, RITEA, microwave, NovaSure, etc (option B)

• Uterine artery embolization (option A)

• Uterine tamponade

• Hysterectomy - if all minimally invasive methods fail (option D)

Solution to Question 16:

Visualization of a mass with a single feeding vessel on a color Doppler is characteristic of an endometrial polyp. In contrast, submucous leiomyomas often receive blood flow from multiple vessels (option A).

Endometrial polyps are soft, fleshy growths within the uterus that consist of endometrial glands, fibrous stroma, and surface epithelium. Risk factors for endometrial polyps include increasing age, obesity, and tamoxifen use.

They are usually asymptomatic. When symptomatic, they present with heavy menstrual bleeding, prolonged bleeding, intermenstrual bleeding, or infertility.

The main diagnostic tools for endometrial polyps include transvaginal ultrasound (TVS) with applied color Doppler, saline infusion sonography, and hysteroscopy.

The risk factors for the malignant transformation include postmenopausal status, age older than 60, and abnormal bleeding. Other factors such as larger polyp size (>1.5 cm), tamoxifen use, diabetes, and obesity may also contribute to a higher risk.

Hysteroscopic polypectomy may be considered most effective for symptomatic women or those with risk factors for malignant transformation.

Solution to Question 17:

Immediate cesarean section is done as the first twin is in breech presentation. Uncomplicated dichorionic diamniotic twin pregnancies are terminated at 38 weeks of gestation.

Timing of delivery depends on the type of twin pregnancy:

The mode of delivery depends on the 1st twin:

After delivery of 1st twin, if the 2nd twin is in breech, assisted breech delivery is done. If the 2nd twin is in a transverse lie, the internal podalic version is done under GA in the OT followed by

breech extraction.

Complications in attempting vaginal delivery in non-vertex presentation (i.e) transverse and breech:

• Interlocking of twins

• Fetal head trapping above the cervix

• Umbilical cord prolapse

According to ACOG criteria (2019), preeclampsia requires proteinuria of grade 2+ or more. In the case of pre-eclampsia, termination of pregnancy must be done at 37 weeks. According to the updated criteria, this patient does not have pre-eclampsia. Hence, a Caesarean section can be done at 38 weeks as for any uncomplicated DCDA twin pregnancy.

Type of twin pregnancy	Timing of delivery
Dichorionic diamniotic	38 weeks
Monochorionic diamniotic	34 - 37 weeks + 6 days
Monochorionic monoamniotic	32 - 34 weeks by Caesarean section

1st twin presentation	Mode of delivery
Vertex	Vaginal
Breech	Cesarean section
Transverse	Cesarean section

Solution to Question 18:

Pubarche or the development of axillary and pubic hair is not mediated by estrogen. It occurs due to testosterone in both sexes. Around 50% of testosterone comes from the ovaries and the rest from the adrenal gland.

Puberty in girls involves breast development (thelarche), an increase in adrenal androgen secretion (adrenarche), and the beginning of menstrual cycles (menarche).

Estrogen has the following actions during puberty:

• Breast development

• Development of the reproductive tract

• Starting of menstruation (option D)

• Increased fat in the hips

• Regulation of pubertal growth spurt

• Vaginal cornification (option C)

• Cervical mucus production (option B)

• Epiphyseal closure

• Maintenance of bone formation - acts against parathyroid hormone.

Solution to Question 19:

Multiple warty growths on the vulva (shown below) and around the anal canal are suggestive of condyloma acuminata (genital warts). HPV 6 and 11 are associated with the development of condyloma acuminatum. They are sexually transmitted.

The verrucous growths may appear discrete or coalesce to form large cauliflower-like growths. Infections are often transient in young women but can persist and progress to cancer in older women and immunocompromised individuals. Histopathology will show koilocytes, perinuclear halo, multinucleation, and acanthosis.

Treatment options include podophyllin, trichloroacetic acid, and surgical removal. The use of podophyllin cream is contraindicated in the first trimester of pregnancy. Larger lesions can be removed with a diathermy loop or laser ablation. Cesarean section is advised for women with genital warts during early labor to prevent the risk of transmitting laryngeal papillomatosis to the newborn.

Option B: Molluscum contagiosum presents as a crop of small domed vesicles with central umbilication (shown below). It is caused by the pox virus and it spreads by close sexual or nonsexual contact and by autoinoculation.

Option C: Chancroid appears as small papules that develop into painful ulcerating pustules. They are associated with unilateral inguinal lymphadenopathy. It is caused by H. ducreyi.

Option D: Herpes appears as multiple crops of vesicles and ulcers.

Solution to Question 20:

Normal saline is the ideal fluid media used for bipolar electrocautery as it can function regardless of the fluid electrolyte content.

Normal saline and lactated Ringer solutions are commonly used isotonic, electrolyte-containing fluids in operating rooms for diagnostic hysteroscopy procedures. These solutions can cause volume overload and, if rapidly absorbed in large volumes, may lead to pulmonary edema.

Isotonic, electrolyte-containing fluids are contraindicated for use with monopolar electrosurgical devices because they conduct electrical current, dissipate instrument energy and render the instrument ineffective. Low-viscosity, electrolyte-poor fluids such as 1.5-percent glycine, 3-percent sorbitol, and 5-percent mannitol are preferred as they are non-conductive and can be safely used with monopolar electrosurgical energy devices.

The primary risk of using hypotonic, hypoosmolar solutions is the intravascular absorption of these fluids, which can lead to acute hyponatremia and hypo osmolality.

The criteria for terminating the procedure for electrolyte-deficient fluid is a deficit of 1 L and electrolyte-rich fluid is a deficit of 2.5 L.

Solution to Question 21:

Point A is 2 cm above the external os and 2 cm lateral to the internal uterine canal.

	Point A	Point B
Location	2cm above and 2cm lateral to the external os	2cm above and 5cm lateral to the external os
Structure present	Paracervical/Parametrial lymph node	Obturator lymph node
Dose of radiation	7000-8000 cGy	6000 cGy



Solution to Question 22:

The image showing enlarged ovaries with multiple large cysts along with the history of treatment for infertility with human menopausal gonadotropin suggests the diagnosis of ovarian hyperstimulation syndrome (OHSS).

Human menopausal gonadotropin (HMG) is a medication that is composed of FSH and LH. It is used to stimulate the release of ova in women who do not ovulate spontaneously. It can lead to ovarian hyperstimulation syndrome.

Ovarian Hyperstimulation Syndrome (OHSS) is a condition associated with ovarian enlargement due to exogenous gonadotropin therapy. Symptoms include abdominal pain, distention, ascites, gastrointestinal problems, respiratory compromise, oliguria, and thromboembolism. It can occur during ovulation induction or in early pregnancies conceived through exogenous ovarian stimulation.

Predisposing factors for OHSS include multi-follicular ovaries (such as in PCOS), young age, high estradiol levels during ovulation induction, and pregnancy.

Vasoactive substances produced by the corpus luteum, such as vascular endothelial growth factor (VEGF) and angiotensin II, are believed to contribute to enhanced capillary permeability, resulting in the loss of fluid, protein, and electrolytes into the peritoneal cavity and hemoconcentration.

Sonographic examination reveals enlarged ovaries with numerous follicular cysts and ascites.

Maintaining fluid balance with isotonic fluids like normal saline is crucial. Strategies to prevent OHSS during exogenous ovulation include reducing follicular stimulation (low FSH dose), coasting (withholding FSH administration one or more days prior to the hCG trigger), prophylactic volume expanders, and substituting hCG for FSH towards the end of ovarian stimulation.

Other options:

Option B: Theca lutein cysts are a rare form of follicular cysts characterized by luteinization and hypertrophy of the theca interna layer. Both ovaries are enlarged and cystic. This condition is commonly associated with gestational trophoblastic disease, multifetal gestation, and placentomegaly.

Option C: Polycystic ovarian syndrome shows 12 or more small follicles each 2-9 mm in size placed peripherally (necklace pattern) (shown below). The patient presents with menstrual irregularities, hirsutism, acne, etc.



Solution to Question 23:

The woman has had two pregnancies in total - the previous twin pregnancy and the current pregnancy, hence the gravida score is 2. The first pregnancy resulted in delivery at term (crossed viable period), hence her para score is 1 (parity doesn't include current pregnancy). Hence, her obstetric score would be G2P1 (Gravida - 2, Para - 1).

Multifetal pregnancies count as one for both gravida and para. A pregnant woman who delivers twins for the first time is expressed as primipara.

The obstetric score is used to summarise the past obstetric history as follows:

• G - Gravidity, denoting the total number of pregnancies, including the present one

• P - Parity, denoting the total number of pregnancies crossing the period of viability. This includes both live births and stillbirths. The current pregnancy is not included.

• A - Abortion is the termination of pregnancy, either spontaneous or induced, before fetal viability.

• L - Living children.

The GTPAL system includes the following:

• Gravida

• Term (>37 weeks)

• Preterm (20w - 36w 6d)

• Abortions (<20 weeks)

• Number of living children

Note: The period of viability is when the fetus is capable of independent existence. The WHO defines it to be 24 weeks while the Indian definition is 28 weeks.

Solution to Question 24:

The image showing bilateral ballooned-out tubes filled with contrast is suggestive of bilateral hydrosalpinx.

Hydrosalpinx is a chronic cystic swelling of the fallopian tube caused by distal tubal obstruction, usually bilateral. It is associated with conditions like PID and endometriosis. Grossly, the fimbriae and tubal ostia are replaced by a smooth, dilated end. The ballooned elongated tubes are filled with clear fluid.

Ultrasound reveals a cystic, elongated structure with thin walls, incomplete septa, and anechoic contents. Hysterosalpingography shows dilated, clubbed fallopian tubes filled with contrast. The interstitial end of the tube is open whereas the fimbrial end is indrawn and closed. This permits the dye to enter but not exit the tube during hysterosalpingography.

Laparoscopic treatment options for those not seeking to preserve fertility include adhesion lysis and salpingectomy. For infertile women, laparoscopic neo-salpingostomy is done for mild tubal disease. In vitro fertilization (IVF) may be a better option for severe tubal disease.

The image given below shows bilateral hydrosalpinx:



Solution to Question 25:

The image showing a poorly developed lower limb suggests the diagnosis of caudal regression sequence/sacral agenesis seen in pregestational diabetes. It is the most specific anomaly seen in diabetes.

The caudal regression sequence is a rare malformation, frequently associated with maternal pre-gestational diabetes. It consists of a spectrum of structural defects of the caudal region, including incomplete development of the sacrum and the lumbar vertebrae.

Elevated maternal HbA1c levels correlate with fetal malformations. It occurs approximately 200 times more frequently in infants of diabetic mothers than in other infants.

Overt diabetes (pregestational diabetes) is associated with higher malformation rates than the general population. Neural tube defects (encephalocele, meningomyelocele, anencephaly), transposition of great vessels, ventricular septal defect, atrial septal defect, hypoplastic left heart, aortic stenosis and coarctation of the aorta are commonly seen.

Small left colon syndrome is a rare anomaly that develops in the second and third trimesters.

Option B: Malformations are not associated with gestational diabetes mellitus (GDM). Malformations occur due to the disruption of organogenesis, usually occurring in the first trimester. The abnormal glycemic status in GDM manifests only in the second trimester.

Option C: Valproate leads to neural tube defects and not sacral agenesis.

Option D: ACE inhibitors can cause oligohydramnios.

Solution to Question 26:

The intensity of radiation is inversely proportional to the squared distance. This is known as the inverse square law in radiotherapy.

Chemoradiation is the mainstay of treatment of stage III and IV endometrial cancers. They are treated with brachytherapy followed by external radiation. Brachytherapy is a form of radiation therapy in which the radiation source is placed close to the tumor.

Other options:

Option B: Larger blood vessels are more radioresistant than microvasculature.

Option C: Permanent non-dividing cells are more radioresistant than rapidly proliferating cells.

Option D: Small intestinal epithelial cells (intestinal crypts), which divide rapidly with no differentiation are the most radiosensitive.

Solution to Question 27:

The given instrument is a fallope ring applicator. It is used in tubal ligation.

Falope ring sterilization is accomplished by applying silicone rubber bands to the fallopian tubes. The major advantage is the elimination of thermocoagulation. The occlusion is only in a short segment and can allow for tubal reconstruction.

Solution to Question 28:

Danazol is not used as a post-coitus contraceptive.

Danazol, a synthetic derivative of ethinyl testosterone, acts by inhibiting pituitary gonadotropins. It possesses mild anabolic, anti-estrogenic, and anti-progestational properties. It is used in abnormal uterine bleeding, endometriosis, and cyclic mastalgia.

Emergency or post-coital contraceptives are used to prevent pregnancy after unprotected intercourse. They are interceptives. Their main action is to make the endometrium unsuitable for implantation. They can also delay or inhibit ovulation, and prevent fertilization of the egg by sperm.

Methods of emergency contraception:

Drug	Dose
Levonorgestrel	0.75mg stat and after 12 hours or a single dose of 1.5mg stat. Taken within 72 hours can extend upto 120 hours
Ethinyl estradiol 50 µg + Norgestrel 0.25 mg (Yuzpe method)	2 tab stat and 2 tabs after 12 hours
Conjugated estrogen	15 mg BD x 5 days
Ethinyl Estradiol	2.5 mg BD X 5 days

Drug	Dose
Copper IUDs	Insertion within 5 days after accidental unprotected exposure.Not suitable for rape victims
Centchroman	2 tablets(60 mg) to be taken twice at an interval of 12 hours within 24 hours of intercourse.
Ulipristal	30 mg up to 5 days
RU486 (Mifepristone)	Single dose of 25 to 50mg

Obstetrics and Gynecology INICET 2021

Question 1:

A pregnant woman at 36 weeks gestation is found to have high blood pressure recording and a urine protein of 3+. Concomitant presence of which of the following symptoms would make you suspect imminent eclampsia?

a) 2 and 4 only

b) 1 and 2 only

c) 1, 2, 3

d) 1, 2, 3, 4

Question 2:

What is the probable diagnosis from the given clinical image?

a) Cervical carcinoma

b) Uterine inversion

c) Uterocervical prolapse

d) Fibroid polyp

Question 3:

Which of the following is not used to treat PCOD?

a) Clomiphene citrate

b) Metformin

c) Tamoxifen

d) OCP

Question 4:

A 26-year-old woman who has had 2 children through normal delivery presents with a dull aching pain over the abdomen, with irregular menses. Per vaginal examination revealed a mass felt through the posterior fornix. A USG showed a clear ovarian cyst of size 5 x 5 cm. What is the next best step?

a) Measure CA 125 levels

b) Follow up after 6 weeks

c) Laparoscopy

d) CT pelvis

Question 5:

Which of the following is not included in the quadruple screening test?

a) Inhibin B

b) Unconjugated Estriol (E3)

c) AFP

d) hCG

Question 6:

The following set of instruments are used for which procedure?



a) Pap smear

b) Hysteroscopy

c) Biopsy

d) Dilatation and curettage

Question 7:

What is the fixative used for a pap smear?

a) Formalin

b) Air dried

c) Normal saline

d) 95% ethanol

Question 8:

A doctor tells a 31-year-old woman, who is 20 weeks pregnant, that her baby has a developmental defect. She has come to you for a second opinion. The following USG finding is seen. Identify the defect.

a) Anencephaly

b) Omphalocele

c) Cystic hygroma

d) Encephalocele

Question 9:

Arrange the following in sequential order with regards to the steps of collection of samples for pap smear testing:

a) 1, 2, 4, 3

b) 2, 1, 3, 4

c) 3, 1, 2, 4

d) 1, 2, 3, 4

Question 10:

Which of the following is a part of AMTSL?

a) Uterotonics after delivery of placenta

b) Controlled cord traction

c) Uterine massage

d) Early cord clamping

Question 11:

Which of the following is not a cause of secondary amenorrhea?

a) Turner's mosaic

b) Kallman syndrome

c) Sheehan's syndrome

d) Asherman's syndrome

Question 12:

What should be the minimum percentage for normal sperm morphology in a seminal fluid analysis?

a) 10%

b) 4%

c) 6%

d) 8%

Question 13:

Which of the following best measures ovarian reserve?

a) Luteal phase LH

b) Post luteal phase LH

c) FSH in the follicular phase

d) Anti-Mullerian hormone (AMH)

Question 14:

An 18-year-old girl presented with amenorrhoea. On examination, she had normal genitalia & normal breast development. Her tests reveal a 5 alpha-reductase deficiency. What could be the probable diagnosis?

a) Concealed sex

b) True hermaphroditism

c) Female pseudohermaphroditism

d) Male pseudohermaphroditism

Question 15:

Match the following syndromes with their respective karyotypes.

a) 1-D, 2-C, 3-B, 4-A

b) 1-B, 2-C, 3-D, 4-A

c) 1-A, 2-B, 3-C, 4-D

d) 1-D, 2-A, 3-C, 4-B

Question 16:

Which of the following complication is specific to Monochorionic twins?

a) Selective fetal growth restriction

b) Polyhydramnios

c) Twin to twin transfusion syndrome

d) Neural tube defects

Question 17:

Match the following disorders causing secondary amenorrhoea with their respective lab findings.

a) 1-C, 2-B, 3-D, 4-A

b) 1-D, 2-B, 3-C, 4-A

c) 1-C, 2-D, 3-B, 4-A

d) 1-A, 2-B, 3-D, 4-C

Question 18:

Which of the following instruments are used in the caesarean section?

a) 1, 2, 5, 6

b) 1, 2, 3, 5

c) 1, 2, 4, 6

d) 1, 2, 3, 6

Question 19:

What is the most likely diagnosis?

a) Vuval hematoma

b) Condyloma accuminata

c) Vulval cancer

d) Bartholin cyst

Question 20:

What is the normal uterocervical length?

a) 3 cm

b) 5 cm

c) 7 cm

d) 10 cm

Question 21:

Which of the following investigations is used to diagnose unicornuate uterus?

a) 2, 4

b) 1, 2

c) 1, 3

d) 2, 3

Question 22:

Which of the following symptoms are seen in endometriosis?

a) 3, 4

b) 2, 3

c) 1, 2

d) 1, 2, 4

Question 23:

A female patient having frothy vaginal discharge was found to have a strawberry cervix. Which of the following is the drug of choice?

a) Metronidazole

b) Fluconazole

c) Tetracycline

d) Cefixime

Question 24:

According to IADPSG, what is the cut-off value of fasting blood glucose for the diagnosis of gestational diabetes mellitus?

a) >92 mg/dL

b) >95 mg/dL

c) >106 mg/dL

d) >90 mg/dL

Question 25:

A primigravida in her 10th week of gestation presents with spotting. On examination, the uterus corresponds to 12 weeks. Transvaginal ultrasound was done and it is given below. What is your diagnosis?

a) Blighted ovum

b) Missed abortion

c) Ectopic pregnancy

d) Hydatidiform mole

Question 26:

A 62-year-old patient with a history of diabetes and hypertension came with post-menopausal bleeding. Which of the following you will do to evaluate the cause?

a) Pap smear

b) Visual inspection with iodine

c) Colposcopy

d) Endometrial sampling

Question 27:

Which of the following is recommended by the WHO for the management of the second stage of labor?

a) Mandatory episiotomy

b) Manual perineal support and warm compresses

c) Maintain head in deflexion and deliver

d) Deliver in lithotomy position only

Question 28:

A primigravida is in labour. Her per-vaginal examination revealed a posterior cervix with 5 cm cervical length, 1 cm dilatation, soft consistency, and head at -1 station. Calculate the modified Bishop score.

a) 8

b) 5

c) 3

d) 0

Question 29:

A couple presents to the infertility clinic with a normal hysterosalpingogram and a semen analysis report with azoospermia. Which of the following is the next best test to differentiate between an obstructive and a testicular cause of azoospermia?

a) Serum FSH levels

b) Serum testosterone levels

c) Testicular biopsy

d) Serum LH level

Question 30:

What is the dose of dexamethasone given for fetal lung maturity?

a) 2 doses of 6mg given 12 hours apart

b) 2 doses of 12mg given 24 hours apart

c) 4 doses of 6 mg given 12 hours apart

d) 4 doses of 12mg given 12 hours apart

Question 31:

What is the dose of carbetocin used in the prevention of postpartum hemorrhage?

a) 50 mcg

b) 150 mcg

c) 200 mcg

d) 100 mcg

Question 32:

Milk production in pregnancy is inhibited by :

a) Low luteinizing hormone

b) High estrogen

c) Human somatomammotropin

d) Low thyroid-stimulating hormone

Question 33:

Which of the following emergency contraceptive is used in India?

a) Misoprostol 400 mcg

b) Ethinyl estradiol 600 mcg

c) Ulipristal 30 mg

d) Levonorgestrel 1.5 mg

Question 34:

Which of the following is not a long-acting reversible contraceptive method?

a) Copper T

b) Implanon

c) Laparoscopic tubal ligation

d) Combined oral contraceptives

Question 35:

Which class of ovarian tumor does clear cell carcinoma of the ovary belong to?

a) Epithelial cell tumor

b) Germ cell tumor

c) Sex cord stromal tumor

d) Granulosa cell tumor

Question 36:

Asherman syndrome is most commonly associated with

a) Oligomenorrhea

b) Hypomenorrhea

c) Dysmenorrhea

d) Menorrhagia

Question 37:

A teenage girl presented with irregular cycles and increased facial hair. Her ovaries showed increased volume. Which of the following are used in the first line treatment?

a) 1,2,3

b) 2,3,4

c) 1,3,4

d) 1,2,4

Question 38:

Identify the instrument given below.

a) Cusco's speculum

b) Sims' speculum

c) Proctoscope

d) Doyen's retractor

Question 39:

A 17-year-old girl was brought by her mother to the outpatient department with primary amenorrhea. She had normal breast development and pubic hair. On examination, the vagina was blind-ending and the uterus was not palpable. What is the most probable diagnosis?

a) Mullerian agenesis

b) Gonadal dysgenesis

c) Turner syndrome

d) Testicular feminization syndrome

Question 40:

A 24-year-old female presents with complaints of foul-smelling vaginal discharge. Wet smear of the discharge is shown below. Drug of choice to treat this condition is_____



a) Clindamycin

b) Metronidazole

c) Cotrimoxazole

d) Azithromycin

Question 41:

Match the following :

a) 1-A, 2-C, 3-B, 4-D

b) 1-C, 2-A, 3-B, 4-D

c) 1-C, 2-A, 3-D, 4-B

d) 1-A, 2-C, 3-D, 4-B

Question 42:

Which of the following modalities are used for the diagnosis of placenta accreta?

a) 1 and 3 only

b) 1 and 2 only

c) 2 and 3 only

d) 1 and 4 only

Answer Key

Question No.	Correct Option
1	d
2	c
3	c
4	b
5	a
6	a
7	d
8	a
9	d
10	b
11	b
12	b

13	d
14	d
15	a
16	c
17	c
18	c
19	c
20	c
21	d
22	d
23	a
24	a
25	d
26	d
27	b
28	b
29	a
30	c
31	d
32	b
33	d
34	c
35	a
36	b
37	b
38	a
39	a
40	b
41	b
42	b

Detailed Explanations

Solution to Question 1:

The clinical case of a pregnant women at 36 weeks of gestation with hypertension and proteinuria points to a diagnosis of pre-eclampsia. Certain symptoms if present might suggest impending

eclampsia. These include 1, 2, 3, 4.

Pre-eclampsia is defined as hypertension with blood pressure ≥140/90 mm Hg along with proteinuria that develops after 20 weeks of gestation in a previously normotensive and non-proteinuric woman. Preeclampsia when complicated with generalized tonicclonic convulsions and/or coma is called eclampsia.

Ominous symptoms of impending eclampsia:

• Headaches - usually over the occipital or frontal region

• Visual disturbances - These could include scotomas. These are due to spasm of retinal vessels following retinal infarction, occipital lobe damage secondary to vasogenic edema, or retinal detachment.

• Epigastric or right upper quadrant pain. This is due to hepatocellular necrosis, ischemia, and edema that stretches the Glisson capsule. This is usually seen along with elevated serum hepatic transaminase levels.

• Sudden and generalized or persistent edema may indicate imminent eclampsia. Visible edema over the ankles despite rest, especially on rising from the bed in the morning hours is pathological.

• Disturbed sleep

• Diminished urinary output - less than 400 mL in 24 hours is considered a poor sign ominous

Other symptoms include:

• Renal or cardiac involvement

• Obvious fetal-growth restriction

• Thrombocytopenia - usually due to microangiopathic hemolysis.

It is essential that the patient should be warned and educated against these symptoms. The presence of these symptoms may warn the obstetrician and possibly avert the progression into eclampsia.

Solution to Question 2:

In the provided image, it is evident that both anterior and posterior vaginal walls protrude completely outside the vaginal orifice, and the entire uterus is positioned below this level, indicating a case of procidentia/ grade IV utero-cervical prolapse.

Utero-cervical prolapse is commonly seen in the elderly age group due to weak pelvic muscles.

Differential diagnoses of genital prolapse include:

• Vulval cyst and Gartner cyst.

• Cyst of the anterior vaginal wall - tense with well-defined margins and cannot be reduced with pressure.

• Urethral diverticula - small and located low in the anterior vaginal wall. Urethroscopy aids in diagnosis.

• Congenital elongation of the cervix - vaginal portion of the cervix is elongated without accompanying vaginal wall prolapse. The fornices are deep.

• Cervical fibroid polyps - cervix is located high up and a rim of the cervix is felt above the pedicle of the cervical polyp.

• Chronic inversion of the uterus- identified by the higher position of the cervix and the inability to define the uterus on palpation. Diagnosis is confirmed by the use of a uterine sound and further examination with ultrasound and laparoscopy, which reveal a depression at the fundus and an absence of the uterine fundus in the pelvis.

• Rectal prolapse.

The image below depicts the various degrees of genital organ prolapse:



Other options:

Option A: Cervical carcinoma

It commonly presents with post-coital bleeding or postmenopausal bleeding. When they present as CIN lesions, they can be identified by colposcopy but mass lesions will present with fungating, cauliflower, or punctate hemorrhagic lesions.

The provided image shows a normal transition zone, in the given setting without VIA or colposcopy it is difficult to diagnose the lesion as CA cervix.

The image below shows characteristic hemorrhagic and cauliflower lesions seen in the CA cervix:



Option B: Uterine inversion:

Acute uterine inversion is a rare and life-threatening post-partum complication. Uterine inversion is associated with primiparity, the use of oxytocin, macrosomia, and fundal insertion of the placenta. Premature traction on the umbilical cord and fundal pressure before placental separation are the usual direct causes. During per abdomen examination fundus of the uterus is felt to be absent. Care for acute uterine inversion involves pain management, resuscitation, and replacement of the inverted uterus before edema sets in by the O'Sullivan method.

The image below image shows uterine inversion which can be identified by a completely inverted uterus.

Option D: Fibroid polyp

They are leiomyomas that present as polyps. A fibroid polyp is generally firm, permanent, paler than a mucus polyp, and is of various sizes. Sonosalpingography is also a very good technique for detecting submucous fibroid polyp.

Note:

Cervical fibroid polyps can be easily identified as the cervix is high up in its normal anatomical position unlike in the given image. The image attached below shows degenerated fibroid polyps.

Solution to Question 3:

Tamoxifen is the best answer as it is not commonly used in the treatment of polycystic ovarian disease (PCOD). Tamoxifen is a selective estrogen receptor modulator (SERM).

Treatment of PCOD:

• Weight reduction alone is beneficial in mild hirsutism.

• Lifestyle changes, such as smoking cessation, are recommended.

• Fourth-generation oral contraceptive pills (OCP) containing ethinyl estradiol and drospirenone are used for treating menstrual abnormalities, reducing acne, and hirsutism.

• OCP with cyproterone acetate is prescribed for PCOS patients with hirsutism.

• Patients with PCOD are prone to developing insulin resistance, which can be controlled with metformin. Acarbose can be used as an alternative if metformin is contraindicated.

• Drugs used in the management of infertility in women with PCOS include letrozole and clomiphene citrate. FOGSI, in 2024, updated letrozole as the first-line drug for induction of ovulation in patients with PCOS. However, the FDA approval for the same is still pending.

Surgical options include laparoscopic ovarian drilling or puncture of not more than four cysts in each ovary. This is done either by laser or by unipolar electrocautery. It restores endocrine levels and improves fertility. Surgery is indicated only if:

• Medical therapy fails

• Hyperstimulation occurs

• Infertile women

• Previous pregnancy losses

Note: In cases where clomiphene and dexamethasone fail to induce ovulation, tamoxifen can be considered as an alternative treatment for infertility associated with PCOD. However, it is not recommended as the initial choice. It's important to note that tamoxifen carries a significant risk of endometrial hyperplasia and can also result in unexpected pregnancy with potential teratogenic effects when used for induction.

Solution to Question 4:

A clear ovarian cyst with size less than 7 cm in a woman of reproductive age group is most likely to be a functional ovarian cyst. The next best step in the management of this case is to follow up after 6–12 weeks and perform a repeat USG.

A simple unilocular cyst under 7 cm is often a functional cyst and should be observed. Most functional cysts resolve spontaneously within 4-6 months. A follow-up ultrasound will detect any persistent cyst, which may necessitate laparoscopic evaluation. To expedite resolution or to treat symptoms like pain and irregular menses, oral combined pills may be prescribed for 3-4 months in women of reproductive age.

Features of a functional cyst:

• Related to temporary hormonal variation

• Benign and rarely become complicated

• Usually 6–8 cm in diameter

• Usually asymptomatic

• Unilocular

• Contains clear fluid, unless hemorrhage occurs

• Lining epithelium corresponds to the functional epithelium of the unit from which they arise

Management of functional cysts in women of child-bearing age group is summarized in the flowchart attached below.



Other options:

Option A: Tumour markers such as CA-125 and CEA are particulary useful in postmenopausal women suspected of having a malignant epithelial cell tumor.

Option C: If the cyst does not resolve spontaneoulsy it can be removed laparoscopically or by laparotomy.

Option D: Imaging modalities such as CT scan and MRI help to detect ovarian neoplasms and assist in staging of ovarian cancers.

Solution to Question 5:

Inhibin B is not a component assayed under the quad-screening test. Dimeric inhibin-A is evaluated in this screening test.

Quadruple testing is done in the second trimester between 15-21 weeks of gestation. It comprises:

• Maternal serum AFP

• Human chorionic gonadotropin

• Unconjugated estriol (E3)

• Dimeric inhibin A

The following table summarizes the findings of the quad-screening test in various conditions.

Various antenatal screening tests for aneuploidy are as follows:

Parameter	Trisomy 21	Trisomy 18	Neural tube defect or abdominal wall defect
MSAFP	Low	Low	High
β hCG	High	Low	Normal

Parameter	Trisomy 21	Trisomy 18	Neural tube defect or abdominal wall defect
Unconjugated estriol	Low	Low	Normal
Inhibin A	High	Normal/ Low	Normal

Dual test (1sttrimester)	Triple test (2ndtrimester)	Quadruple test(2ndtrimester)
1. Beta HCG2. PAPPA (Pregnancy-associated plasma protein A)This can be combined with nuchal translucency by USG	1. Beta HCG2. Unconjugated estriol3. MSAFP	1. Beta HCG2. Unconjugated estriol3. MSAFP4. Serum inhibin

Solution to Question 6:

The given set of instruments are commonly used for the collection of pap smears. The given image includes Cusco's bivalved self-retaining speculum, Koplin's jar, smear slides, Ayer's spatula, and endocervical brush.



Pap/Papanicolaou test is a screening test only. The presence of abnormal cells/ a positive test requires further evaluation by colposcopy, cervical biopsy, or fractional curettage.

Screening guidelines:

• Time for initiation: 21 years, regardless of the age of first sexual intercourse.

• Women aged 21-29 years: Every 3 years(earlier it was done annually)

• Women who are aged 30-65 years: Pap test + HPV testing every 5 years.

• Women should stop having cervical cancer screening after the age of 65 years. If suspicious PAP, screening till 75 years.

• HIV+ women: annual checkup until three consecutive negative tests followed by testing every 3 years for life.

• HPV DNA testing in combination with PAP improves the sensitivity of screening.

Prerequisites:

• Should be done prior to vaginal examination.

• Lubricants should not be used.

• The patient should not be menstruating during the pap test.

• The ideal time is around ovulation.

• The patient should not have had intercourse in the 24hrs prior to the test.

Procedure:

• The patient is put in a dorsal position & the labia are separated.

• Cusco's self-retaining speculum is introduced/posterior vaginal wall retractor is used

• The squamocolumnar junction is identified and scraped with Ayre's spatula.

• Endocervix is scraped with a cytobrush.

• Samples are spread separately on 2 different glass slides.

• The slide is immediately fixed using 95% ethyl alcohol.

• Stained with Pap stain or Short stain.

Other options:

Option B: Hysteroscopy is the process of inspection of the uterine cavity.

The image shows the procedure of hysteroscopy and the common instruments used for the same.



Hysteroscopy

Option C: To take biopsy we commonly use Gellhorn's punch biopsy forceps as shown in the image below. It is used to take biopsies from the cervix.

Option D: Instruments used in dilatation and curettage are shown below.



Solution to Question 7:

The most common fixative used in pap smear is 95% ethanol (ethyl alcohol).

Pap smear is a screening test for cervical cancer. Cervical cancer screening should start at age 21. Women aged 21-29 should have Pap smear every 3 years, while women aged 30-65 should have both HPV and Pap co-test every 5 years.

Before the Pap test:

• The Pap test should not be deferred due to unexplained discharge or unscheduled bleeding as they may be signs of cervical or other genital tract cancers.

• Ideally scheduled to avoid menstruation. It is best performed around the time of ovulation.

• Patients should abstain from vaginal intercourse, douching, tampon use, and intravaginal medications for a minimum of 24 to 48 hours prior to the test.

• Avoid touching the cervix before performing the Pap test, as dysplastic epithelium may get removed.

• Ideally, confirmed cervicitis or vaginitis should be treated prior to the test

Procedure:

• The patient is put in a dorsal position and Cusco's self-retaining speculum is introduced.

• The squamocolumnar junction is scraped with Ayre's spatula completing at least one full rotation. It predominantly samples the ectocervix.

• Endocervix is scraped with a cytobrush, which is rotated only one-quarter to one-half turn after the ectocervix has been sampled.

• A broom simultaneously samples both endo- and ectocervix. Five rotations in the same direction are recommended, since reversing direction may cause loss of cellular material.

• Samples are spread separately on 2 different glass slides and fixed using 95% ethyl alcohol.

• Stained with Pap stain or Short stain.



Solution to Question 8:

The given ultrasound image shows an open defect in the calvaria and skin leading to the exposure of the cranial neural tube. This is consistent with the diagnosis of anencephaly.

Anencephaly results from incomplete closure of the neural tube. It is one of the earliest detectable congenital defects that can be detected as early as 10 to 11 weeks of gestation. Screening can also

be done by serum estimation of maternal serum alpha-fetoprotein (MSAFP). The following are the radiological signs seen in anencephaly.

• Frog eye sign - Absence of cranial calvaria, prominent orbits thereby resembling the face of a frog.

• Shower cap sign - The absence of the cranium could lead to the protrusion of a disorganized mass of brain tissue. This resembles a shower cap giving rise to a characteristic triangular facial appearance.

Many neural-tube defects can be prevented with preconceptional supplementation of folic acid. A dose of 4 mg daily is recommended in subsequent pregnancies.



Other options:

Option B: Omphalocele is a midline anterior abdominal wall defect through which the solid abdominal viscera and/or bowel may herniate. The extruded abdominal contents are covered in a sac. These are due to the persistence of physiological herniation of the gut or as a part of

the OEIS complex (omphalocele, exstrophy, imperforate anus, and spinal abnormality) and cloacal exstrophy. The ‘elephant’s trunk’ sign on ultrasound, due to the prolapsed terminal ileum is said to be pathognomonic for cloacal exstrophy of the OEIS complex.

The first image below shows a patient with cloacal exstrophy showing the elephant trunk sign representing the intussuscepted ileum. The second image shows omphalocele while the third image shows gastroschisis (a close differential diagnosis for omphalocele). The fourth image demonstrates an ultrasound image of the fetus suggestive of an omphalocele (asterix)







Option C: Cystic hygroma is a congenital multiloculated lymphatic lesion that is classically found in the left posterior triangle of the neck. It occurs as a result of sequestration or obstruction of developing lymph vessels. It must be carefully differentiated from nuchal translucency (NT) as both may appear similar. This defect can be picked up in a first-trimester ultrasound.

The image below shows a newborn with cystic Hygroma on the right side of the neck and cystic hygroma of the neck in the fetus.

Option D: Encephalocele is caused due to defects in the skull through which the meninges and/or neural tissue herniate. The squamous part of the occipital bone is the most frequently affected bone. In meningoencephalocele or occult spina bifida with Chiari malformations, the scalloping of the frontal bones is observed in the ultrasound as the lemon sign while anterior curvature of the cerebellum with effacement of the cisterna magna in the posterior fossa forms the banana sign.

The image shows a newborn with encephalocele. The second and third images below show the banana and the lemon sign respectively.



Solution to Question 9:

The correct sequence of the procedure involved in the collection of a sample for a pap smear is 1, 2, 3, 4.

The Pap/Papanicolaou smear is used for screening cervical neoplasia.

Procedure:

• The patient is put in a dorsal position & the labia are separated.

• Cusco's self-retaining speculum is introduced/posterior vaginal wall retractor is used

• The squamocolumnar junction is identified and scraped with Ayre's spatula.

• Endocervix is scraped with a cytobrush.

• Samples are spread separately on 2 different glass slides.

• The slide is immediately fixed using 95% ethyl alcohol.

• Stained with Pap stain or Short stain.

Solution to Question 10:

Controlled cord traction during placental delivery is a component of active managament of third stage labor (AMTSL).

AMTSL is recommended by the WHO to prevent postpartum hemorrhage (PPH).

The recommended components of AMTSL are as follows:

Uterine massage or fundal massage is not recommended as an intervention to prevent PPH in women who have already received prophylactic uterotonics such as oxytocin (core component of AMTSL). Hence, it is now an optional component of AMTSL that is done in patients with uterine atony as a part of the management of PPH.

	Component	Remarks
1.	Injectinguterotonicswithin1 minuteof delivery of the baby or immediately after delivery of the anterior shoulderbefor e the delivery of the placenta	Most importantstepOxytocin 10 IU im/iv infusion is thepre ferred uterotonic.(Not recom mendedby WHO-carboprost)
2	Delayed cord clamping	Done within1-3 minsof delive ry80 mlof additional blood g oes to the fetusPreventsfetal anemia

	Component	Remarks
3	Delivery of placenta bycontro lled cord traction:Modified B randt Andrews technique	Done only in settings where s killed birth attendants are av ailableThe uterus is pushed u pwards and backwardThe tra ction of cord is downwards a nd forwards
4	Intermittent assessment of ut erine tone	Done every 15 mins for 2 hou rs

Solution to Question 11:

Kallman syndrome is a not a cause of secondary amenorrhea. It is a cause of primary amenorrhea.

Primary amenorrhea is defined as the absence of menstruation by 14 years without the development of secondary sexual characteristics or by 16 years regardless of the presence of secondary sexual characteristics. Kallmann syndrome presents with primary amenorrhea due to hypogonadotropic hypogonadism from deficiency of GnRH. It is associated with olfactory dysfunction and anosmia.

Secondary amenorrhoea is the absence of menses for more than 3 months in girls or women who previously had regular menstrual cycles or 6 months in girls or women who had irregular menses. Sheehan's syndrome (postpartum hypopituitarism secondary to the necrosis of the pituitary gland) and Asherman's syndrome (formation of intrauterine synechiae, usually following endometrial destructive procedures like curettage or ablation) are causes of secondary amenorrhea.

Note: Turner's syndrome is a cause of primary amenorrhea. Turner's mosaic may present either with primary or with secondary amenorrhea. So, of the given options, the best answer is Kallman syndrome.

The image attached below summarizes the pathology involved in various types of amenorrhea.



Solution to Question 12:

4% is the minimum percentage for normal sperm morphology in a seminal fluid analysis.

Morphology of sperms is the most important marker of male fertility. Characterization of normal sperm morphology shows improved correlation with fertilization rates during IVF cycles.

The criteria requires careful analysis of the following:

• The shape and size of the sperm head,

• The relative size of the acrosome in proportion to the head,

• And characteristics of the tail, including length, coiling, or presence of two tails.

Significantly decreased fertilization rates are observed when normal morphology of the sample falls below 4 percent.

Abnormal sperm morphology is termed teratospermia or teratozoospermia.

Solution to Question 13:

Antimullerian Hormone (AMH), secreted by the granulosa cells, is the best marker of ovarian reserve as it can be tested on any day of the cycle.

AMH is secreted by the Sertoli cells in the testis and granulosa cells in the ovary. In males, it inhibits the development of the Müllerian system and its absence can result in hermaphroditism. In the female, it helps in follicular development and oocyte maturation. The estimation of serum AMH is used in the study of ovarian reserve in an infertile woman and a woman with secondary amenorrhoea. The levels of AMH vary minimally across the cycle and they strongly correlate with the number of ovarian primordial follicles. This makes AMH is the best predictor for measuring ovarian reserve. AMH levels can be affected by the use of oral contraceptives and tobacco smoking.

Values and interpretation:

• Normal value: 2–6.8 ng/mL

• Poor ovarian reserve : <1 ng/mL

• PCOD and hyperstimulation syndrome: >10 ng/mL

Tests for ovarian reserve: Ovarian reserve means the capacity of the ovary to produce eggs. The other indicators include:

• FSH levels are measured in the early follicular phase (Day 2 or 3 of the menstrual cycle) - High FSH means the ovarian reserve is getting depleted (FSH is highest in postmenopausal women). FSH (values >25mIU/ml) levels are higher than normal in women with a reduced ovarian reserve and ovarian hypofunction.

• Serum oestradiol values on day 3 of the cycle. Serum oestradiol <75 pg/mL indicates ovarian hypofunction.

• LH - values <10 IU/L indicates poor reserve

• Antral follicular count - counts less than 4–5 on days 2–5 denotes poor response to hormones.

• Inhibin B - values <40 pg/mL indicates poor reserve

Solution to Question 14:

A girl presenting with amenorrhea, normal female external genitalia and breasts with 5 alpha-reductase deficiency is suggestive of male pseudohermaphroditism.

In pseudohermaphroditism, the sex glands are of a certain sex while the external genitalia is of the opposite sex.

Male pseudohermaphroditism or male intersex is an association of male gonads (XY genotype) with female external genitalia. The production of testosterone from the testes requires 5 alpha-reductase. This enzyme converts testosterone into DHT (dihydrotestosterone), which is capable of acting on peripheral target tissues to produce the male phenotype. The absence of this enzyme results in failure of the testis to descend to the scrotum and ill development of the penis, thereby resulting in the female phenotype. Many individuals develop their attitudes and psyche according to their sex of rearing (as a female). Rehabilitation to their gender roles is the best line of management.

Other options:

Option A: Concealed sex refers to the deliberate concealment of sex usually done by criminals by changing clothes or by other methods to avoid detection. This can be detected by physical examination.

Option B: In true hermaphroditism the glands of both the sexes are present in the same individual. Such cases are very rare. In most cases, the accessory sex gland is atrophic and lacks functional activity.

Option C: Female pseudohermaphrodites are genetic females (XX genotype) with male/ambiguous external genitalia due to the deficiency of 21 alpha-hydroxylase enzymes resulting in an excess of androgens. This causes clitoromegaly resembling the penis,and the fused labioscrotal folds resemble a rudimentary scrotum. Management includes administration of steroids to control excess ACTH, estrogen to confer the secondary sexual characteristics, and surgical restoration of the external genitalia to a feminine pattern.

Solution to Question 15:

The correct answer is 1-D, 2-C, 3-B, 4-A.

Mayer-Rokitansky-Küster-Hauser (MRKH) syndrome, commonly referred to as Mullerian agenesis, refers to the failure of development of the uterus, cervix, and upper part of the vagina in a female with 46 XX karyotype. These individuals have normally functioning gonads and external genitalia.

Condition	Karyotype
1. Turner's syndrome	D. 45 + XO
2. Swyer's syndrome	C. 46+ XY
3. Androgen insensitivity syndrome	B. 46+ XY
4. MRKH (Mayer-Rokitansky-Kuster-Hauser) syndrome	A. 46 + XX

Complete androgen insensitivity syndrome/testicular feminizing syndrome:

In this syndrome, the karyotype is XY and thus, the patient is a genotypic male. Pathology is due to the defective response of the androgen receptors to androgens. Findings include:

• Primary amenorrhea → Due to the absence of uterus and cervix.

• Normally developed breasts → Due to the conversion of the excessive androgens into estrogen.

• Swelling in the inguinal region → Suggestive of an underdeveloped ectopic testis, caused by loss of action of testosterone on its receptor.

• Absence of uterus and cervix → Due to the production of Mullerian inhibiting substance from the Sertoli cells of the testis.

Note:

There is some discrepancy among authors on this topic. Androgen insensitivity syndrome (AIS) with XY karyotype may have mostly female external sexual characteristics. According to William's Textbook of Obstetrics 26th edition, breast development occurs due to peripheral aromatization of testosterone to estrogen whereas the pubic and axillary hair remains scant or absent. But as per Shaw's Textbook of Gynecology 7th edition, XY females with AIS have normal secondary sexual characteristics.

Swyer syndrome or pure gonadal dysgenesis occurs as a result of a mutation in the SRY gene. This leads to underdeveloped, dysgenetic gonads that fail to produce androgens or anti-Mullerian hormone (AMH). The following features are seen:

• Normal prepubertal female phenotype- Absent or poorly developed secondary sexual characteristics.

• Normal Müllerian system development due to the absence of AMH. (Uterus and cervix present)

• Female external genitalia

• Presence of dysgenetic testes that cannot produce AMH or sex hormones. They have a risk of malignant transformation.

As there is gonadal dysgenesis in a patient with 46 XY karyotype, they are actually determined to be males and are a type of male pseudohermaphrodite. They present with primary amenorrhoea, absence of secondary sex characters, and female external genitalia.

Turner's syndrome is a chromosomal abnormality with partial or complete monosomy of the X chromosome (45+XO). In Turner's syndrome, gonadal dysgenesis causes primary ovarian failure

that is hypogonadism. The clinical picture of amenorrhea, short stature, webbed neck, low hairline at the back of the neck, widely spaced nipples, and lack of secondary sexual characteristics is seen in Turner's syndrome.

Solution to Question 16:

Twin to twin transfusion syndrome (TTTS) is a specific complication seen in monochorionic twins. Selective fetal growth restriction, polyhydramnios and neural tube defects are seen as complications to several conditions and are not specific to monochorionic twins.

In monochorionic diamniotic gestation, twins are separated solely by two amniotic membranes, with no intervening chorion. Monochorionic monoamniotic gestation lacks a dividing membrane.

All monochorionic placentas likely have some anastomotic connections, unique to this type. Chronic feto-fetal transfusion may lead to the following clinical syndromes:

• Twin-twin transfusion syndrome (TTTS)

• Twin anemia–polycythemia sequence (TAPS)

• Twin reversed-arterial-perfusion (TRAP) sequence

Twin to twin transfusion syndrome (TTTS):

TTTS is a fetal syndrome in monozygotic twinning with two amnionic sacs and a shared chorion, allowing for anatomical sharing of fetal circulations through anastomoses of placental arteries and veins. This leads to transfusion from donor twin to recipient twin, resulting in donor anemia and restricted growth, while the recipient may suffer polycythemia, circulatory overload, and hydrops.

Features:

The Quintero staging is as follows:

• Stage I—Discordant amnionic fluid volumes+ urine visible in donor's bladder.

• Stage II—Stage I + no visible urine in donor's bladder.

• Stage III— Stage II + abnormal Doppler studies of the umbilical artery, ductus venosus, or umbilical vein.

• Stage IV— frank ascites or hydrops in either twin.

• Stage V—the demise of either fetus.

Treatment:

• Laser photoablation of the placental vascular anastomoses is the preferred treatment for severe TTTS - Quintero stages 2 to 4.

• Septostomy is seldom done.

• Amnioreduction is a treatment modality that involves needle drainage of excessive amniotic fluid.

• Selective feticide is done only if the growth disturbances develop before 20 weeks.

• Of all, selective fetoscopic laser photocoagulation (SFLP) by Solomon technique is the preferred treatment modality as it is associated with better survival rates and a reduced likelihood of

recurrence.

Donor twin	Recipient twin
Oliguria	Polyuria
Anemia	Polycythemia
Oligohydramnios	Polyhydramnios
Activation of Renin-angiotensin system (RAAS)	Activation of Atrial natriuretic peptide (ANP)
Decreased renal blood flow	Increased renal blood flow
Non-visible bladder	Over distended bladder, circulatory overload, hyperviscosity syndrome, hyperbilirubinemia, hypertension
Renal failure	Congestive cardiac failure (CCF), Hydrops twin (late cases)

Solution to Question 17:

The correct answer is 1-C, 2-D, 3-B, 4-A.

Secondary amenorrhoea is the absence of menses for more than three months in girls or women who previously had regular menstrual cycles or six months in girls or women who had irregular menses.

Premature Ovarian failure or premature menopause is the development of menopause before the age of 40 years. It is diagnosed when at least 2 serum FSH levels are elevated (>40 mIU/l) when measured at least 1 month apart in a woman <40 years. The other findings consistent with POF include low estradiol and an elevated FSH/LH ratio.

Asherman's syndrome is the presence of uterine synechiae, due to the destruction of large areas of the endometrium, most commonly as a result of trauma to the endometrium following pregnancy-related curettages. Oligomenorrhoea, amenorrhoea, dysmenorrhoea, abortion, and infertility are seen with uterine synechiae. The laboratory investigations and hormonal assays are generally normal in isolated cases of uterine synechiae. Diagnosis is generally established using hysteroscopy. Moth-eaten appearance on hysterosalpingography suggests a diagnosis of Asherman syndrome. Treatment is by hysteroscopic adhesiolysis.

PCOS is a multisystem, metabolic, endocrine dysfunction that commonly occurs among women in the reproductive age group. The endocrinological changes reflecting in the assays are summarized in the pearl attached below.

Sheehan's syndrome/ pituitary apoplexy is hemorrhagic infarction of large pituitary adenomas or pituitary infarction due to traumatic major blood loss. If this blood loss occurs during delivery, it's called Sheehan's syndrome. Affected women may present with panhypopituitarism and may have persistent hypotension, tachycardia, hypoglycemia, and lactation failure. The most prominent among symptoms are lactation failure (due to deficiency of

prolactin) and postpartum amenorrhea (due to deficiency of gonadotropins).

Condition	Lab markers
1. Sheehan's syndrome	C. Low LH, Low FSH, Low es tradiol
2. Premature Ovarian failure	D. High LH, High FSH, Low estradiol
3. PCOS	B. High LH, Low FSH, Low e stradiol
4. Asherman's syndrome	A. Normal FSH, Normal LH, Normal estradiol

Note: Estradiol level in PCOS is low in comparison to Estrone (E1).

The common differentials for the presentation for secondary amenorrhea and the lab evaluation is summarized in the flowchart shown below.



Solution to Question 18:

Of the given list/combination of instruments 1, 2, 4, 6 are commonly used in the cesarean section.

Cesarean section is an operative procedure in which the fetuses are delivered through an incision on the abdominal and uterine walls.

The given image summarizes the various instruments used in the lower segment cesarean section.

Bard-Parker blade (scalpel with blade) has a handle with a notch to fit the blade. It is usually held by pencil grip. Commonly for abdominal incisions, we use blade no: 22. Other uses include incision of skin and sub-cutaneous tissues, cutting the clamped pedicles, and performing sharp dissections.

The given image shows various Bard-Parker blades.

Doyen's retractor is used to retract the abdominal wall as well as the bladder to facilitate better exposure of the lower uterine segment during cesarian operation. It is also used to pull back the soft tissues to widen the surgical field during laparotomy.

The image below shows Doyen's retractor.



Alli's forceps: It is identified by the distal blades and a tip that has teeth that do not oppose each other. There is also a lock in the proximal part.

Uses:

• To catch hold of the margins and angles of the uterine flaps in LSCS after the delivery of the baby in settings where the Green-Armytage hemostatic clamp is not available.

• To catch hold the anterior lip of the cervix

• To hold the apex of the episiotomy wound during the repair.

• To catch hold of the margins of the peritoneum, rectus sheath, vaginal mucosa during the repair.

• To catch hold of the torn ends of the sphincter ani externus prior to suture in the repair of complete perineal tear.

The given image shows Alli's forceps.



Green armytage forceps have a broad tip as shown in the image and hence are used to compress a wide area. It is used as a hemostat in caesarian sections. In LSCS (Lower segment Caesarean section), the cut uterine edges can bleed and Green Armytage forceps are applied to the edge of the incision for hemostasis.

Cusco's speculum is a self-retaining bivalve speculum used for visualization of the cervix, vaginal fornices, and for performing pap smears.

The instrument shown in the image is Cusco's speculum.

Shirodkar's uterine holding clamp has blades with curved transverse bars with gaps between them when it is locked. This gap makes it atraumatic and is used to lift, hold and steady the uterus during sling surgeries and tubal procedures.

The image below shows Shirodkar's uterine clamp.



Solution to Question 19:

The given image shows a discrete growth on the left labia majora of the vulva. The most likely diagnosis is vulvar carcinoma.

Vulvar carcinoma:

Squamous cell carcinoma is the most common type of vulvar cancer. The most common site of vulvar cancer is labia majora > Clitoris > Labia Minora. Persistent HPV infection and chronic inflammation such as lichen sclerosis are risk factors. It is generally seen in elderly women and presents with vulvar pruritus, vulvar swelling, lump, or an ulcer. The lump is papular with a raised pigmented area. The ulcers often have everted margins with fissured, cracked, and indurated surrounding skin. They usually bleed and cause offensive vulval discharge.

The most common type of spread in vulval cancer is lymphatic while the vascular route of spread is extremely rare in vulvar cancer. The first lymph node to be involved is the superficial inguinal lymph node and thus, a sentinel lymph node biopsy is helpful for evaluating the spread of vulval cancer. Groin node status is the best prognostic predictor.

Other options:

Option A: Vulval hematoma is a collection of blood in the vulval region. The symptoms include persistent perineal pain and rectal tenesmus. There may be retention of urine. The local examination would commonly reveal a tense swelling at the vulva which becomes dusky and purple in color. It is mostly tender to touch. A small hematoma (<5 cm) may be treated conservatively with a cold compress. Larger hematomas should be explored in the operation theatre under general anesthesia

The image shows vulval hematoma. As it is a vascular leak a blue hue can be appreciated as the blood clots.

Option B: Condyloma acuminata (venereal warts) is caused by infection with Human Papilloma Virus 6 or 11. The verrucous growths may appear discrete or coalesce to form large cauliflower-like growths. They may be asymptomatic, but can also cause discomfort discharge, or bleeding. A biopsy is necessary to rule out malignancy and the specific treatment includes electrocautery, cryosurgery, laser, or diathermy.

The image shows multiple warty growths on the vulva and around the anal canal as seen in condyloma acuminata.

Option D: Bartholin cyst is caused by obstruction of the Bartholin ducts by proteinaceous material or by inflammation from infection. Generally caused by gonorrhea and presents as a painful cystic swelling. Symptomatic or recurrent cysts may require marsupialization or gland excision. Bartholin abscess is a sequela to a cyst and it usually requires surgical drainage.

The image below shows Bartholin cyst.

Solution to Question 20:

The normal length of the uterine cavity including the cervical canal is usually 6–7 cm.

The image of the coronal section showing different parts of uterus is attached below.



Clinical application:

• Normally, the uterocervical length is measured using Sim's uterine sound.

• It is based on this length that, lengths of various intra-uterine systems/ intra-uterine devices is determined.

The image below shows Sim's uterine sound.

Solution to Question 21:

Laparoscopy and Hysterosalpingogram (HSG) can be used to diagnose unicornuate uterus.

In the unicornuate uterus, the formation of the uterus and fallopian tube occurs entirely from one Mullerian duct due to the developmental arrest of other Mullerian duct. Women with a unicornuate uterus have an increased incidence of infertility, endometriosis, and dysmenorrhea.

Evaluation:

• The initial evaluation is through sonography. It can also help pick up ipsilateral renal agenesis, which is seen in 40% of the women with unicornuate uterus.

• Saline infusion sonography (SIS) is another modality to identify Mullerian anomalies. It can be used as a screening tool for cavity evaluation as well.

• On Hysterosalpingography, the endometrial cavity usually assumes a fusiform (banana) shape, tapering at the apex and draining into a single fallopian tube. The uterus is generally shifted off the midline.

• Invasive modalities such as hysteroscopy/laparoscopy are reserved for patients with complicated Mullerian anomalies or in patients presenting with infertility to establish the diagnosis.

• These days, MR imaging is used to characterize and pre-operatively evaluate cases that are complicated.

The image below shows a hysterosalpingogram of a unicornuate uterus

Falloposcopy is the inspection of the fallopian tubes through a micro-endoscope. Hysteroscopic falloscopy reveals the tubal pathology of the cornual and interstitial end of the fallopian tube.

Solution to Question 22:

The symptoms of endometriosis include 1, 2, 4. Endometriosis presents with unexplained infertility, dysmenorrhea, and vaginal bleeding.

Endometriosis is the presence of endometrial tissue (glands and stroma) outside the uterus. It is an estrogen-dependent disease of the childbearing age group. The most common site of occurrence is the ovary followed by the pouch of Douglas.

Key clinical features of endometriosis include:

• Dysmenorrhoea

• Lower abdominal pain

• Infertility

• Pain during urination or defecation,

• Constipation or diarrhea

• Irregular bleeding.

The gold standard investigation for the diagnosis of endometriosis is a visual inspection by laparoscopy, preferably with histological confirmation.

The management of endometriosis is summarized in the flowchart below:



Solution to Question 23:

The given clinical scenario of frothy vaginal discharge and strawberry cervix is suggestive of trichomoniasis and metronidazole is the drug of choice for the treatment.

Trichomonas vaginalis is one of the most common causes of vaginal discharge. It is transmitted by sexual intercourse, with an incubation period of 4–21 days. The clinical features include vaginal discharge which is profuse, thin, slightly green in colour, and frothy. It may be associated with vaginal irritation, pruritus, dysuria, and increased frequency. Cervical inflammation with punctate hemorrhagic spots known as the strawberry cervix is typically seen. Motile trichomonads are seen on wet mount examination.

The below image shows the strawberry cervix seen in trichomoniasis.

Administration of metronidazole 2 g orally in a single dose or 500 mg 2 times/day for 7 days is the standard treatment. It should be taken at night and after meals to avoid vomiting. If this too causes vomiting, or in resistant cases an alternate regimen is to be considered. Both the partners need to be treated simultaneously to avoid recurrence.

The alternative regimen includes tinidazole 500 mg orally twice a day for 7 days or secnidazole in a single dose of 1000 mg daily for 2 days. The use of tinidazole is not recommended during pregnancy.

Solution to Question 24:

According to IADPSG, the cut-off value of fasting blood glucose for the diagnosis of gestational diabetes mellitus (GDM) is >92 mg/dL.

Gestational diabetes mellitus (GDM) refers to any degree of glucose intolerance having its onset or first recognition during pregnancy.

For IADPSG criteria, an oral glucose tolerance test (OGTT) is done in the fasting state using 75 g of glucose, and GDM is diagnosed if any one of the following cut-offs is met i.e.

• Fasting: >92 mg/dL

• 1hour: >180 mg/dL

• 2 hr: >153 mg/dL

Earlier, a two-step approach was being used to diagnose GDM. In this, the first step (screening) is a 50-gram one-hour glucose challenge test (GCT). The patients having a positive result of >140mg/dl, are candidates for the OGTT. A 100-gram, three-hour OGTT is the diagnostic test for GDM. GDM is diagnosed if two or more values are positive, as follows.

This has now been replaced by the one-step approach that omits the screening test and simplifies diagnostic testing by performing only a 75-gram, two-hour oral GTT.

Some complications of GDM:

• Maternal

• Hypertension, preeclampsia

• Hydramnios

• Increased operative interventions

• Future development of diabetes mellitus

• Fetal and neonatal

• Fetal macrosomia

• Congenital anomalies

• Metabolic abnormalities such as hypocalcemia, hypomagnesemia, hyperbilirubinemia, and neonatal hypoglycemia.

• Respiratory distress syndrome

• Subsequent childhood and adolescent obesity

Time	Carpenter and Coustan	NDDG
Fasting	95 mg/dL	105 mg/dL
1 hour	180 mg/dL	190 mg/dL
2 hour	155 mg/dL	165 mg/dL
3 hour	140 mg/dL	145 mg/dL

Solution to Question 25:

The given clinical scenario of a primigravida with spotting and an ultrasound showing a characteristic snowstorm appearance (anechoic cystic spaces) is suggestive of a hydatidiform mole (complete mole). It refers to an echogenic uterine mass that has numerous anechoic cystic spaces.

Hydatiform mole belongs to the benign group of gestational trophoblastic diseases (GTD). It is an abnormality in chorionic villi making them partly proliferative and degenerative, thereby resulting in the formation of multiple small cysts. Based on the portion of the chorionic villi involved, they could be complete (diffused) or partial (focal)

The classical clinical features of a complete mole are:

• Abnormal vaginal bleeding -Most common (90%)

• Lower abdominal pain

• Hyperemesis gravidarum due to increased β-hCG

• Features of early-onset preeclampsia

• Uterus appears large for date

• Absent fetal parts and FHS

• Expulsion of vesicular tissues (grape-like vesicles)

• Theca lutein cyst of ovaries of more than 6 cm

• Hyperthyroidism (rare)

• Serum hCG > 100,000 mIU/ml

• USG - snowstorm appearance

The definitive management after diagnosis is suction and evacuation, after making the cervix favorable. Follow-up of the patients with serial hCG values is crucial.

Prophylactic chemotherapy can be started on high-risk women (WHO prognostic score > 7) with methotrexate 1mg/kg/day intramuscular/intravenous on days 1, 3, 5, and 7. Patients are also advised to not get pregnant for at least a year.

Other options:

Option A: Blighted ovum (silent miscarriage) occurs when a fertilized egg implants in the uterus but doesn't develop into an embryo. It is a sonographic diagnosis where there is an absence of a fetal pole in a gestational sac with a diameter of 3 cm or more.

Option B: Missed abortion is when the fetus is dead and retained inside the uterus for a variable period. USG shows an empty sac early in the pregnancy or the absence of fetal cardiac activity and fetal movements.

Option C: Ectopic pregnancy is one in which the fertilized ovum is implanted and develops outside the normal endometrial cavity. The majority of ectopic pregnancies occur in the fallopian tube, the ampulla being the most common site.

Solution to Question 26:

Endometrial sampling is done to evaluate a patient presenting with postmenopausal bleeding.

Postmenopausal bleeding is defined as any bleeding from the genital tract after 12 months of menopause in a woman not on hormone replacement. Any vaginal bleeding occurring after 6 months of amenorrhea in a menopausal woman should also be considered as postmenopausal bleeding and investigated. All postmenopausal women with unexpected uterine bleeding should be evaluated for endometrial malignancy since age is a significant risk factor for this.

Transvaginal ultrasound (TVS) is a better first-line investigation. Since TVS is not included in the options, endometrial sampling is the next best answer.

Endometrial sampling is indicated if:

• The endometrial thickness is >4 mm

• The endometrium shows diffuse or focal increased echogenicity (heterogeneity)

• The endometrium is not adequately visualized on transvaginal sonography

• The woman has persistent bleeding, as it can be a sign of endometrial cancer even when the endometrial thickness is less than 4mm.

The image below gives a summary of the evaluation of postmenopausal bleeding, as per the FOGSI guidelines.



Other options:

Option A: Pap smear should be offered to all women above the age of 21 years who are sexually active for at least 3 years. Colposcopic examination and selective biopsy are considered in patients with abnormal Pap smears.

Option B: Visual inspection with iodine is done where the facilities for Pap screening do not exist. This involves painting the cervix with Lugol's iodine. It is used to detect any abnormal cytology in the cervix and vagina, such as in cases of carcinoma cervix.

The normal vaginal and cervical epithelium contains glycogen, therefore stains mahogany brown on being painted with Lugol's iodine. The abnormal cells remain unstained. These areas need to be biopsied and sent for histopathological examination.

Option C: The colposcopic study is challenging in postmenopausal women because of narrow vaginas, senile vaginitis, indrawn and poorly visible squamocolumnar junction, and atrophic cervix.

Solution to Question 27:

Among the given options, manual perineal support and giving warm compresses are recommended by WHO in the management of the second stage of labor.

The second stage of labor refers to the time period between full cervical dilatation and the birth of the baby. For women in the second stage of labor, the following techniques to reduce perineal trauma and facilitate spontaneous birth such as perineal massage, warm compresses, and manual perineal support (hands-on guarding of the perineum) are recommended.

Option A: Mandatory episiotomy is discouraged. Episiotomy is indicated in cases of:

- Shoulder dystocia
- Breech delivery
- Fetal macrosomia

• Operative vaginal deliveries

• Persistent occiput posterior positions

Option C: Head is delivered by maintaining flexion to ensure that the small sub-occipitofrontal diameter distends the vulval outlet instead of the larger occipitofrontal diameter.

Option D: The position that is most comfortable to the mother including upright positions is recommended.

Solution to Question 28:

The Modified Bishop score of the given patient is calculated to be 5.

From the data given in the question:

• The position of the cervix is posterior, hence score = 0

• Cervical length is 5 cm, hence score = 0

• Cervical dilatation is 1 cm, hence score = 1

• Consistency of cervix is soft, hence score = 2

• The station of the head is –1, hence score = 2

Hence, the Modified Bishop score is 0+0+1+2+2 = 5.

The Bishop score is a scoring system used to assess labor induction outcomes. It is based on the station of the head, dilation, effacement (or length), position, and consistency of the cervix noted by doing a vaginal examination.

According to the NICE guidelines, a score of >/=6 is considered favorable.

According to Williams Obstetrics 26th edition, a score >/=9 is considered favorable.

Note: In the modified Bishop score (1991) the effacement is replaced by the cervical length in cm.

Solution to Question 29:

Serum FSH level is the next best test to differentiate between an obstructive and a testicular cause of azoospermia. FSH is the main hormone stimulating the production of sperm in the testis and has the highest specificity and sensitivity among hormonal assays while differentiating between obstructive and non-obstructive azoospermia.

The absence of sperm in semen is known as azoospermia. It can be due to:

• Pre-testicular azoospermia due to a hypothalamic or pituitary abnormality (hypogonadotropic-hypogonadism)

• Testicular failure or non-obstructive azoospermia

• Post-testicular obstruction implies normal spermatogenesis but failure to deliver the sperm into the ejaculate because of ductal obstruction.

The algorithm for differentiating the causes of normal semen volume azoospermia is as follows:



• A low FSH level indicates pituitary or hypothalamic failure

• A high FSH level denotes primary gonadal failure or testicular cause

• A normal FSH level could be due to either non-obstructive or obstructive azoospermia. A testicular biopsy is usually required in such people to provide a definitive diagnosis.

Solution to Question 30:

The dosage of dexamethasone given for fetal lung maturity is 4 doses of 6 mg, which is given 12 hours apart.

Maternal administration of glucocorticoids is advised in preterm labor when the period of gestation is less than 34 weeks. Intramuscular injection of either betamethasone or dexamethasone is given.

Dosage:

• Betamethasone (Betnesol): 2 doses of 12 mg each, given 24 hours apart.

• Dexamethasone: 4 doses of 6 mg each, given every 12 hours.

This helps in fetal lung maturation and minimize the incidence of:

• Respiratory distress syndrome

• Intraventricular hemorrhage

• Necrotizing enterocolitis

It is beneficial when the delivery is delayed beyond 48 hours after the first dose.

Solution to Question 31:

100 mcg of carbetocin is used in the prevention of postpartum hemorrhage (PPH).

The use of carbetocin (100 μg, IM/IV) is recommended for the prevention of PPH. It is a long-acting synthetic analog of oxytocin with agonist properties. It binds to oxytocin receptors in the uterine smooth muscle, and results in rhythmic contractions, increased frequency of existing contractions, and increased uterine tone.

To effectively prevent PPH, the WHO recommends the use of only one of the following uterotonic agents:

• Oxytocin - (10 IU - IM/IV)

• Carbetocin - (100 μg - IM/IV)

• Misoprostol - (400 μg or 600 μg, per-oral)

• Ergometrine/methylergometrine - (200 μg - IM/IV)

• Oxytocin and ergometrine fixed-dose combination - (5 IU,500μg - IM)

Of these, oxytocin is used as the first-line agent in preventing postpartum hemorrhage. When oxytocin is not available, the other drugs mentioned above can be used based on the clinical condition and availability of the drug.

Solution to Question 32:

Milk production in pregnancy is inhibited by high estrogen.

The synthesis and secretion of milk from the breast alveoli is known as lactogenesis. Prolactin, an anterior pituitary hormone, is responsible for lactogenesis. In spite of high prolactin levels during pregnancy, there is inhibition of milk secretion. This is because of high levels of estrogen and progesterone circulating during pregnancy, which makes the breast tissues unresponsive to prolactin.

Following delivery, there is a fall in estrogen and progesterone levels. As a result, prolactin begins its milk secretory activity in previously fully developed mammary glands.

Other hormones involved in lactogenesis:

• Insulin

• Growth hormone

• Cortisol

• Thyroxine

• Placental lactogen aka human chorionic somatomammotropin: Has growth promoting and lactogenic effects

• Serotonin

Oxytocin is the major hormone for galactokinesis or the ejection of milk.

Solution to Question 33:

Levonorgestrel (LNG) 1.5 mg is the emergency contraceptive used in India.

Emergency contraception refers to backup methods for contraceptive emergencies which women can use within the first few days after unprotected intercourse to prevent an unwanted pregnancy.

Levonorgestrel 1.5 mg single dose can be used for emergency contraception. Alternatively, levonorgestrel 0.75 mg one tablet can be taken within 72 hours of unprotected intercourse and another tablet 12 hours later.

Note: Under the National Reproductive and Child Health Programme, the Drug Controller of India has only approved levonorgestrel tablets for use as an emergency contraceptive pill.

Other options:

Option A: Misoprostol is a prostaglandin E1 analog used in the medical termination of pregnancy.

Option B: High doses of ethinylestradiol can provide emergency contraception when used in combination with levonorgestrel. This is known as the Yuzpe regime.

Option C: Ulipristal is a synthetic progesterone hormone receptor modulator. It attaches to the progesterone receptor and prevents/ delays ovulation, and suppresses the endometrium, thereby preventing implantation. Ulipristal 30 mg should be taken within 5 days of unprotected sexual intercourse. It is approved by US-FDA and European medicine agencies but at present, it is not available in India as an emergency contraceptive pill.

Solution to Question 34:

Laparoscopic tubal ligation is an irreversible method of contraception.

Sterilization is a permanent method of birth control. It includes surgical procedures like tubectomy in woman and vasectomy in man. These methods are advised for those couples, who have completed their families and do not want to bear the inconvenience or cost of the other methods of contraception, and when the other methods are contraindicated.

• Postpartum sterilization is performed in the first week of the postnatal period. It can be combined with a cesarean section

• Interval sterilization is done in non-pregnant women or at any time after 6 weeks of delivery.

Tubal sterilization can be achieved by three methods which are:

• Electrocoagulation is used for the destruction of a segment of the tube

• Mechanical occlusion of the tube using a silicone rubber band (Falope ring) or a spring-loaded clip (Hulka Clemens clip)

• Suture ligation with or without tubal segment excision such as Pomeroy's method, Uchida's method, Irving's method, etc.



Other options:

Option A: Copper T (CuT) is an effective, reversible, and long-term method of contraception. About 50 mcg of copper is eluted daily in the uterus. It can be left in place for a long time and does not interfere with sexual activity. They have an effective life of about 3–5 years. Copper T 380A, known as Paraguard, has a lifespan of 10 years.

Option B: Implanon is a subdermal implant that contains 67mg of desogestrel. It releases 30 mcg of the desogestrel daily and the effect lasts for up to 3 years. It prevents ovulation and is reversible within 1 month of removal.

Option D: Combined oral contraceptive pills (OCP) contain a mixture of ethinyloestradiol with progestogen. It acts by suppressing pituitary hormones i.e. FSH and LH peak and thereby preventing ovulation. It is a reversible method of contraception as 90% of women will have normal ovulatory cycles within 3 months of stopping OCPs.

Solution to Question 35:

Clear cell carcinoma of the ovary is a type of epithelial cell tumor.

Epithelial ovarian tumors arise from the mesoepithelial cells present on the ovarian surface. 80% of all ovarian cancers are epithelial cancers.

Clear cell carcinoma is also known as mesonephroid tumor. The histopathology of clear cell carcinoma reveals large cuboidal epithelial cells with abundant clear cytoplasm characteristically forming tubules, glands, and small cystic spaces. These cystic spaces are lined by hobnail cells. They are clear cells showing large dark nuclei protruding into the lumen. It is a highly malignant tumor.

Malignant epithelial tumors can spread widely via transcoelomic spread. It can spread through the capsule onto the peritoneal surface, omentum, intestinal viscera, and sub-diaphragmatic space. An ascitic tap will reveal blood-stained fluid with tumor cells. It may also spread to the para-aortic nodes and cause metastasis in the other ovary, liver, lungs, and gastrointestinal tract.

Solution to Question 36:

Among the given options, Asherman syndrome is most commonly associated with hypomenorrhea.

Asherman syndrome is the presence of uterine synechiae, due to the destruction of the endometrium. It can occur as a result of trauma to the endometrium following vigorous curettages such as incomplete abortion, postpartum hemorrhage, and elective abortion. Damage may also result from other uterine surgery such as metroplasty, myomectomy.

They usually present with menstrual abnormalities, infertility, recurrent pregnancy loss due to inadequate placentation. Menstrual abnormalities, including hypomenorrhea and amenorrhea, remain the most common clinical features associated with Asherman syndrome. Dysmenorrhea is occasionally present.

The diagnosis is confirmed with hysterosalpingography.

Management includes hysteroscopy, which enables adhesiolysis. Recurrence of adhesions is prevented by insertion of IUCD for 3 months and giving estrogen cyclically (21 days) for 3 months to develop endometrium.

Note:

Amenorrhea indicates the complete absence of menstruation.

Oligomenorrhoea is the term used to describe infrequent and irregular menses. In this condition, the cycle length is prolonged without affecting the duration and amount of flow.

Hypomenorrhoea refers to the condition in which the cycle length remains unaltered, however the duration of bleeding or the amount of blood loss, or both are substantially reduced.

Menorrhagia denotes excessive blood loss (increase in the duration and /or amount of bleeding) without any change in the cycle length.

Dysmenorrhoea is defined as cramping pain accompanying menstruation.

Solution to Question 37:

This patient's irregular menstrual cycles, increased facial hair (signifying clinical hyperandrogenism), and increased ovarian volume are suggestive of polycystic ovarian syndrome (PCOS). Anti-androgens, lifestyle modifications, and combined oral contraceptives (2, 3, and 4) constitue first-line treatment modalities for this condition. Laparoscopic drilling is a surgical procedure done when medical therapy fails.

PCOS is an endocrinopathy seen in women of reproductive age, affecting multiple systems. Rotterdam criteria is used to diagnose PCOD. The presence of two of the three components of this criteria establishes the diagnosis:

• Oligo/amenorrhoea, anovulation, infertility

• Hirsutism/acne

• Ultrasound findings:

• Enlarged ovaries with ovarian volume will be more than 10 mm3

• 12 or more small follicles each of 2-9 mm in size placed peripherally

Ultrasound image given below shows an enlarged ovary seen in PCOS with multiple follicles in the periphery.



Treatment of PCOD:

• Weight reduction alone is beneficial in mild hirsutism.

• Lifestyle changes, such as smoking cessation, are recommended.

• Fourth-generation oral contraceptive pills (OCP) containing ethinyl estradiol and drospirenone are used for treating menstrual abnormalities, reducing acne, and hirsutism.

• OCP with cyproterone acetate is prescribed for PCOS patients with hirsutism.

• Patients with PCOD are prone to developing insulin resistance, which can be controlled with metformin. Acarbosecan be used as an alternative if metformin is contraindicated.

• Infertility in PCOD patients is treated with clomiphene as the first-line drug for inducing ovulation. Dexamethasone is used in combination to improve fertility rates. Letrozole is used off-label in clomiphene-resistant cases.

Surgical options include laparoscopic ovarian drilling or puncture of not more than four cysts in each ovary. This is done either by laser or by unipolar electrocautery. It restores endocrine levels and improves fertility. Surgery is indicated only if:

• Medical therapy fails

• Hyperstimulation occurs

• Infertile women

• Previous pregnancy losses

Solution to Question 38:

The instrument shown in the image is Cusco's speculum.

Cusco's speculum is a self-retaining bivalve speculum used for visualisation of the cervix, vaginal fornices and for performing pap smear.

Other options:

Option B: Sims' speculum is used for the assessment of rectocele and cystocele. The speculum is applied against the vaginal wall and slowly retracted out until a bulge appears, indicating cystocele or rectocele. It can also be used for visualization of the vaginal walls.

Option C: A surgeon uses a lubricated proctoscope to visualize lesions in the anus, anorectal junction, and lower rectum. It is done as an OPD procedure to detect hemorrhoids, polyps, or ulcers.

Option D: Doyen's retractor is used to retract the abdominal wall as well as the bladder to facilitate better exposure of the lower uterine segment during cesarian operation. It is also used to pull back the soft tissues to widen the surgical field during laparotomy.



Solution to Question 39:

The given clinical scenario of a girl with primary amenorrhea, normal breasts, and pubic hair, with a blind-ending vagina and uterus not palpable, is suggestive of Mullerian agenesis.

Absent uterus and vagina in a person with female genotype and phenotype are features of Mullerian agenesis, also known as Mayer-Rokitansky-Kuster-Hauser syndrome.

Ovaries are normal and secondary sexual characters are well developed with normal external genitalia. This condition is associated with anomalies of the spine/skeleton and the kidneys along

with the urinary tract.

Diagnosis is made with the help of pelvic sonography, MRI, and laparoscopy. Patients may have rudimentary Mullerian bulbs without an endometrial cavity. These should be surgically excised if they are symptomatic.

Other options:

Option B: Gonadal dysgenesis patients present with primary amenorrhoea, absence of secondary sex characters (poorly developed breasts), and female external genitalia.

Option C: In Turner syndrome, short stature with poor development of secondary sexual characters and short stature will be seen. It has a karyotype of 45 XO. Barr body will be absent. Female genital organs (vagina, uterus, fallopian tubes, ovaries) are present. The ovaries, though present, are atrophic.

Option D: Testicular feminization syndrome also known as androgen insensitivity syndrome has a karyotype of 46XY. Secondary sexual characters like the development of axillary hair and pubic hair are absent in these females. However, breast development is adequate due to the conversion of androgen into estrogen during puberty.

Solution to Question 40:

A female patient presenting with foul-smelling vaginal discharge and a wet smear showing clue cells points to a diagnosis of bacterial vaginosis, and the drug of choice is metronidazole.

Bacterial vaginosis is a polymicrobial infection caused by Mobiluncus curtisii, Gardnerella vaginalis, Mycoplasma hominis, and Haemophilus vaginalis. It has a foul-smelling (fishy smell) vaginal discharge with minimal vulval irritation.

Amsel's criteria is helpful in diagnosing bacterial vaginosis when 3 of the following 4 are present:

• Thin, white homogenous vaginal discharge

• A pH of vaginal fluid >4.5

• Fishy odor of discharge accentuated by adding 10% KOH-Whiff test

• Presence of clue cells on microscopy: Clue cells are epithelial cells with a fuzzy border or granular cytoplasm caused by small gram-negative bacilli adhering to their surface

Treatment is oral metronidazole 500 mg BD for 7 days.

The image below shows clue cells:



Solution to Question 41:

Option B represents the correct answer. 1-C, 2-A, 3-B, 4-D

Endodermal sinus tumor/yolk sac tumor is a type of ovarian germ cell tumor that is seen in premenarchal girls. Schiller-Duval bodies are pathognomonic on histopathology. Alpha-fetoprotein (AFP) is commonly produced by this tumor and is used as a tumor marker.

Dysgerminona is a type of germ cell tumor and it corresponds to the seminoma of the testis. It does not secrete either male or female sex hormones. It secretes placental alkaline phosphatase (ALP), lactate dehydrogenase (LDH), and β-hCG. Serum lactate dehydrogenase (LDH) is usually used in monitoring individuals or disease recurrence.

Choriocarcinoma is a type of mixed germ cell tumor. It is highly vascular. Histologically, the tumor shows syncytiotrophoblasts and cytotrophoblasts. It secretes large quantities of human chorionic gonadotropin (hCG) hormone, which is used as a tumor marker in the diagnosis and management of the tumor.

Malignant epithelial cell tumors of the ovary secrete CA-125. It is a glycoprotein and surface cell antigen which is secreted by malignant epithelial tumors of the ovary. A level of more than 35 U/mL suggests a malignant and residual tumor and indicates the need for chemotherapy. CA-125 is also raised in abdominal tuberculosis and endometriosis.

Ovarian tumors	Tumor marker
1. Yolk sac tumor	C. AFP
2. Dysgerminoma	A. LDH
3. Choriocarcinoma	B. Beta-HCG

Ovarian tumors	Tumor marker
4. Epithelial cell tumor	D. Ca-125

Solution to Question 42:

Among the given options, ultrasonography and MRI can be used in the diagnosis of placenta accreta.

The term placenta accreta spectrum (PAS)/morbidly adherent placenta refers to abnormal placental implantation characterized by the abnormally implanted, invasive, or adhered placenta. This is due to the absence of decidua basalis and poor development of the fibrinoid or Nitabuch layer.

Risk factors include prior cesarian delivery, prior uterine surgery, prior curettage, and endometrial ablation, associated placenta previa.

Types of the morbidly adherent placenta:

• Placenta accreta: Villi are attached to the myometrium.

• Placenta increta: Villi invade the myometrium.

• Placenta percreta: Villi penetrate through the myometrium and to or through the serosa.

Ultrasound imaging, color Doppler, and MRI have all been useful in the diagnosis of morbidly adherent placenta during pregnancy.

USG findings that suggest PAS are (1) placental lacunae, (2) thinning of the retroplacental myometrium, (3) disruption of the bladder-uterine serosal interface, (4) bridging vessels from the placenta to the bladder-serosal interface, and (5) a placental bulge that pushes outward and distorts the contour of the uterus.

Color flow Doppler study shows hypervascularity of serosa bladder interface.

MRI reveals uterine bulging and dark intraplacental bands on T2-weighted imaging.

Management:

• Partial placenta accreta: Placental tissue is removed as much as possible. Oxytocin helps in uterine contraction and hemostasis. In cases following cesarean delivery, bleeding areas are oversewed.

• Total placenta accreta:

• Parous women- hysterectomy is done

• If the patient desires to have a child- conservative measures include incising the uterus above the placental attachment, clamping, and cutting the umbilical cord as close to its base as possible. The placenta is left behind, which may get autolyzed eventually. Uterine artery embolization or therapy with methotrexate has also been done for the conservation of the uterus.

• If placenta accreta invades the bladder, placental removal alone is avoided. It requires a hysterectomy and partial cystectomy.

The below image shows the placenta accreta, increta, and percreta.



Obstetrics & Gynecology NEET 2021

Question 1:

A pregnant woman at 36 weeks of gestation with a prosthetic valve replacement for mitral stenosis is on warfarin therapy. Her INR is 3. What is the appropriate next step in management?

a) Stop warfarin and start LMWH

b) Stop warfarin and start heparin

c) Continue warfarin

d) Stop warfarin, start LMWH and asprin

Question 2:

A 27-year-old female patient was found to have uterus didelphys. Which of the following is not likely to be a complication of this uterine anomaly?

a) Abortion

b) Endometriosis

c) Preterm labor

d) Transverse lie

Question 3:

A woman who underwent dilatation and curettage for molar pregnancy 6 months ago presents with persistent symptoms of abdominal pain and distention. On further evaluation, the beta-hCG levels were persistently elevated and cannonball foci were found on the chest X-ray. What is the appropriate management in this case?

a) Methotrexate with folinic acid

b) Methotrexate

c) EMACO regimen

d) Hysterectomy

Question 4:

A female patient has a history of three miscarriages. Identify the investigation that was done from the image shown below.



a) Hysteroscopy

b) Saline infusion sonography

c) Hysterosalpingography

d) Transvaginal ultrasonography

Question 5:

A 56-year-old multipara woman presented with grade II/III uterine prolapse with cystocele. She complains of passing urine on coughing and sneezing. What is the type of urinary incontinence seen in this patient?

a) Overflow incontinence

b) Urge incontinence

c) Stress incontinence

d) Neurogenic bladder

Question 6:

A 32-year-old married woman who came for the evaluation of infertility was found to have a septum in her uterine cavity. What would be the appropriate management of this case?

a) Hysterectomy

b) Laparoscopy and metroplasty

c) Hysteroscopy and septoplasty

d) Laparotomy and metroplasty

Question 7:

A 30-year-old G4A3 woman had a history of miscarriage at 8 weeks, the second miscarriage at 11 weeks with no cardiac activity, and the third pregnancy loss at 24 weeks with preterm delivery due to early-onset preeclampsia. What is the most probable diagnosis?

a) Syphilis

b) Antiphospholipid antibody syndrome

c) TORCH infections

d) Gestational diabetes mellitus

Question 8:

A 26-year-old female patient came with a history of fatiguability and constipation for the past 1 year. She has a history of multiple blood transfusions after delivery 4 years ago. She has had a lactational failure and never regained her periods after the delivery. What is the most probable diagnosis?

a) Sheehan syndrome

b) Hypothyroidism

c) Premature ovarian failure

d) Delayed blood transfusion reaction

Question 9:

The ultrasound of a young woman with infertility is suggestive of a uterine anomaly. Which of the following is used to confirm the diagnosis?

a) Hysterosalpingography

b) Hysteroscopy + laparoscopy

c) Laparoscopy

d) Transvaginal sonography

Question 10:

A 26-year-old pregnant woman, on examination was found to have infra-umbilical flattening. The fetal heart sounds were heard over the lateral side. What could be the fetal presentation/position?

a) Brow

b) Knee

c) Occipitoposterior

d) Right dorso-anterior

Question 11:

A primi at term presented with labor pain and multiple painful vesicular lesions on her vulva. No rashes were found on other parts of the body. What will be the management of this case?

a) Acyclovir and elective Cesarean section

b) Acyclovir and induction of labor

c) Spontaneous delivery

d) Antiviral and normal vaginal delivery

Question 12:

A 30-year-old married woman came for an infertility evaluation. On performing further investigations, intramural fibroids of sizes 7x5cm and 5x3cm were seen at the right and left cornual ends of the uterus. Her husband's semen analysis was found to be normal. What is the appropriate management of this case?

a) Uterine artery embolization

b) Myomectomy

c) Artificial reproductive techniques

d) GnRH analogues

Question 13:

A 28-year-old married woman who is anxious to conceive presents with copious vaginal discharge, without any itching. Her last menstrual period was 13 days ago. What is the cause of her vaginal discharge?

a) Candidiasis

b) Bacterial vaginosis

c) Trichomoniasis

d) Physiological

Question 14:

A patient developed atonic postpartum hemorrhage after a normal vaginal delivery. She was shifted to the OT and the obstetrician planned ligation of vessels to arrest the bleeding. What is the order of ligation of the blood vessels supplying the uterus?

a) Uterine artery - Ovarian artery - Internal iliac artery

b) Uterine artery - Ovarian artery- External iliac artery

c) Uterine artery - Ovarian artery - Vaginal artery

d) Uterine artery - pudendal artery - Vaginal artery

Question 15:

A teenage patient presented with a history of dysmenorrhea and chronic pelvic pain. On ultrasonography, a transverse vaginal septum was seen. What is the most common complication that can be seen in this patient?

a) Endometriosis

b) Tubo-ovarian abscess

c) Dermoid cyst

d) Theca lutein cyst

Question 16:

A 28-year old pregnant woman came for a routine antenatal check-up in her second trimester. Four years back, she delivered 2 healthy full-term twins. What would her obstetric score be?

a) G2P2

b) G3P2

c) G2P1

d) G3P1

Question 17:

A 27-year-old woman who is on oral contraceptive pills for contraception has forgotten to take her pills on 4 different days in the last 2 weeks. She presents to the primary health center, asking for advice. What would you suggest to her?

a) Take 4 pills now, continue the schedule, use barrier method of contraception

b) Take 1 pill now, continue the schedule, and use barrier method of contraception

c) Take 1 pill now, use barrier method, and consider emergency contraception in case of recent intercourse

d) Discard the pack, start a new pack with alternate method of contraception

Question 18:

A 30-year-old female patient, on treatment for infertility, came with 6 weeks of amenorrhea, abdominal pain, and mild vaginal bleeding. Her beta hCG level was 2800 mIU/mL. On ultrasound, a 3cm mass was seen in the left adnexa. It contained a gestational sac with no fetal cardiac activity. What is the most appropriate management for this patient?

a) Expectant management

b) Single dose methotrexate

c) Salpingectomy

d) Milking of the tube

Question 19:

A 30-year-old woman on oral contraceptives for the past 6 months presents with 6 weeks of amenorrhea. What would be the best method of assessing the gestational age of this patient?

a) Crown rump length by USG

b) Adding 280 days to the LMP

c) Fundus examination

d) Adding 256 days to the LMP

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	c
5	c
6	c
7	b
8	a
9	b
10	c
11	a
12	b
13	d
14	a
15	a
16	c
17	d
18	b
19	a

Detailed Explanations

Solution to Question 1:

The next appropriate step in the management of a pregnant woman with a prosthetic valve on warfarin therapy at 36 weeks of gestation is to stop warfarin and start low molecular weight heparin (LMWH).

Warfarin is safe and effective in preventing thromboembolism in the mother during the 2nd trimester. However, it is teratogenic and can lead to miscarriage, stillbirths, fetal anticoagulation, and fetal embryopathy if used in the 1st trimester.

Heparin is safe for the fetus as it does not cross the placenta but is less effective than warfarin in preventing thromboembolism.

LMWH- Low molecular weight heparin, UFH- Unfractionated heparin

For patients taking LMWH, factor Xa levels should be monitored weekly.

Heparin should be discontinued just before (at least 6 hours before) delivery. If delivery occurs before discontinuation and there is extensive bleeding, then iv protamine sulfate can be given.

Warfarin or heparin anticoagulation can be restarted after 6 hours of vaginal delivery, usually without any problem.

Warfarin, LMWH, and UFH do not accumulate in breast milk and do not have an anticoagulant effect in the newborn. They are compatible with breastfeeding.

Other options:

Option B: Unfractionated heparin can also be given instead of LMWH. However, in this case, the better answer is to give LMWH; because, in clinical practice, LMWHs have a long half-life, once-daily dosing, and fewer side effects. If LMWH is unavailable or cost-prohibitive, or if anti-Xa levels cannot be monitored, continuous infusion of UFH can be used as an alternative to LMWH.

Option D: Aspirin 75-100 mg can be given daily during pregnancy. It can be continued in women with mechanical heart valves if needed for other indications.

Period of gestation	Anticoagulant
Till 12 completed weeks	LMWH or UFH
13 to 36 weeks	Warfarin
36 weeks till just before deliv ery	LMWH or UFH
6 hours after vaginal delivery and 24 hours after cesarean section	Warfarin, LMWH or UFH

Solution to Question 2:

Among the given options, transverse lie is not a likely complication of uterus didelphys. A transverse lie is more likely to be seen with an arcuate or sub-septate uterus.

A uterus didelphys is a Mullerian defect characterized by two separate uterine horns, each with an endometrial cavity and uterine cervix. It occurs due to the failure of fusion of the Müllerian ducts.

It is usually asymptomatic but can rarely present with dyspareunia, dysmenorrhea, hematocolpos, and hematometra. Rarely, endometriosis (option B) and genital neoplasms may be associated with uterus didelphys.

Pregnancy is possible with uterus didelphys, however, there is an increased risk of the following complications:

• Spontaneous abortion (option A)

• Preterm labor (option C)

• Breech presentation

• Cesarean delivery

• Intrauterine growth restriction

• Low birth weight (<2.5kg)

• Perinatal death

Uterus didelphys can be diagnosed with ultrasound or hysterosalpingography, but MRI provides the most accurate diagnosis.

Uterus didelphys

Solution to Question 3:

The given clinical scenario of a woman with a past history of D(&)C for molar pregnancy, now presenting with abdominal pain and distension, elevated beta-HCG levels, and cannonball metastasis to the lung is suggestive of gestational trophoblastic neoplasia (GTN).

The next step is to categorize the patient as low/high risk based on the WHO prognostic scoring system to decide the treatment modality.

WHO Prognostic Scoring System, adapted by FIGO:

Based on the available clinical information,

• Antecedent gestation: molar pregnancy, score= 0

• Interval: 6 months, score= 1

• Site of metastasis: lung, score= 0

The total score is 1, hence it is a case of low-risk GTN and the management involves treatment with methotrexate with folinic acid.

Treatment of GTN:

Low-risk GTN (FIGO Stages I-III: score <7): Single-agent chemotherapy with methotrexate (MTX) or dactinomycin (less commonly used).

• 50 mg MTX is given IM on days 1, 3, 5, and 7 with folinic acid on alternate days.

High-risk GTN (FIGO Stages II-III: score >7 and Stage IV): Multiple agent chemotherapies, with or without adjuvant surgery or radiotherapy. The preferred multiple-agent chemotherapy regime is the EMACO regime:

• Etoposide

• Methotrexate

• Dactinomycin (actinomycin D)

• Cyclophosphamide

• Vincristine (Oncovin)

The FIGO Anatomic Staging of GTN is shown in the image below:

Prognostic Factor	Score			
0	1	2	4	
Age	< 40	>40	-	-
Antecedent gestation	Mole	Abortion	Term	-
Interval in months	<4	4-6	7-12	>12
Pretreatment serum hCG (IU/L)	< 103	103to < 104	104to < 105	> 105
Largest tumor size (including the uterus)	<3	3-4	≥ 5	-
Site of metastases	Lung	Spleen, kidney	GI tract	Brain, liver
Number of metastases	-	1-4	5-8	>8
Previously failed chemotherapy	-	-	single drug	≥ 2 drugs
Low-risk GTN: score ≤6High-risk GTN: score ≥7				



GTN typically develops along with or after a molar pregnancy. Rarely, it may develop after live birth, miscarriage, or medical termination of pregnancy.

The given image below shows the classification of gestational trophoblastic diseases:



Solution to Question 4:

The investigation shown in the given image is hysterosalpingography. The given hysterosalpingogram shows a unicornuate uterus.

Hysterosalpingography (HSG) is an imaging modality in which radiopaque contrast is injected through the cervical canal to assess the endometrial cavity, the fallopian tubes, and the endocervical canal.

HSG should be done between days 5 and 10 of the menstrual cycle, in the preovulatory phase, as an outpatient procedure. Leech Wilkinson cannula, Rubin's cannula, or Foley's catheter no.14 is introduced into the endometrial cavity through the cervical os, and the contrast is injected. The uterine cavity and fallopian tubes are then visualized and X-rays are obtained.

HSG can be used for the following:

• To assess tubal patency in infertility

• To detect uterine anomalies, synechiae, and polyps

• To detect incompetence of the cervical os

Contraindications to HSG:

• Acute pelvic infection

• Active uterine bleeding

• Pregnancy/ suspected pregnancy

• Contrast allergy

• Premenstrual phase

HSG is a safe procedure, however, complications such as ascending spread of infection, chemical peritonitis, allergic reaction to the dye, or pelvic endometriosis may occur.

Other options:

Option A: Hysteroscopy is an operative procedure in which the uterine cavity can be visualized by passing a fiberoptic telescope via the cervix. It can be diagnostic or therapeutic. A hysteroscopic image of a uterus with carcinoma is shown below.



Option B: Saline infusion sonography is a procedure where normal saline is injected into the uterine cavity and the uterus is imaged with transvaginal sonography. It is shown in the image below.

Option D: Transvaginal ultrasonography is a noninvasive imaging technique using the transvaginal approach. It provides a more detailed evaluation of the pelvic organs as compared to the transabdominal approach.

Solution to Question 5:

The type of urinary incontinence present in a patient with prolapse and cystocele experiencing urination on coughing and sneezing is stress incontinence.

Stress incontinence refers to the involuntary leakage of urine when there is increased intra-abdominal pressure (eg, coughing, sneezing, laughing, Valsalva maneuver). Absence of detrusor muscle contraction is seen in urodynamic studies.

Causes:

• Age and menopause - due to estrogen deficiency

• Multiparity - due to loss of pelvic muscle tone

• Obesity

• Smoking

• Constipation, chronic cough

• Pelvic organ prolapse

• Pregnancy - due to the effect of progesterone and pressure of the gravid uterus on the bladder neck.

• Puerperium - due to descent of the bladder neck, loss of urethro-vesical angle, and reduced pelvic muscle tone

• Surgical repair of vesicovaginal fistula

Grading of genuine stress incontinence (GSI) :

• Grade I: Incontinence with only severe stress, such as coughing, sneezing, and jogging.

• Grade II: Incontinence with moderate stress, such as fast walk, going up and down the stairs.

• Grade III: Incontinence with mild stress such as standing.

Investigations are only done prior to surgery to confirm the diagnosis. They include:

• Stress test

• Cotton swab or Q-tip test

• Marshall and Bonney's test

• Urethroscopy

• Urodynamic studies.

These are not required prior to conservative management.

Conservative management includes Kegel's pelvic exercises, electric nerve stimulation, devices like pessaries, vaginal cones, and artificial sphincters. Medical treatment with estrogen creams, alpha agonists, venlafaxine, and imipramine may be useful.

Other options:

Option A: Overflow incontinence is an old term that refers to the inability to empty the bladder, with involuntary, continuous urinary dribbling. Currently, it refers to another presentation of urge incontinence.

Option B: Urge incontinence is the involuntary escape of urine when there is an imminent need to urinate. It occurs due to detrusor muscle overactivity or detrusor instability. It is seen more commonly in old women with decreased bladder capacity, decreased sensation, and CNS inhibition.

Option D: Neurogenic bladder occurs in patients with neurogenic dysfunction leading to detrusor hyperreflexia. Conditions such as multiple sclerosis, stroke, Parkinson's disease, and spinal injury give rise to this. They present with urgency or urge incontinence.

Solution to Question 6:

The appropriate management for a 32-year-old woman with a septate uterus is hysteroscopy and septoplasty.

Hysteroscopic septoplasty refers to the resection of the uterine septum under hysteroscopic guidance. The septum can be resected either with a resectoscope or a laser. It does not scar the uterus and avoids the necessity of cesarean sections for future pregnancies. There is a reduced risk of pelvic adhesions, compared with transabdominal approaches.

A septate uterus is characterized by a septum in the endometrial cavity. It is caused by the failure of the medial segments of the Müllerian ducts to regress. This creates a permanent septum within the uterine cavity.

The septum can project partially from the fundus or extend completely to the cervical os. It is the most common congenital uterine malformation in women with infertility.

Other options:

Option A: Hysterectomy is not advisable since the patient has come for infertility evaluation and desires pregnancy.

Options B & D: Laparoscopy/laparotomy and metroplasty are not preferred for the treatment of a septate uterus due to the following:

• Requirement of cesarean section for subsequent pregnancies to prevent uterine rupture.

• Postoperative pelvic adhesions leading to infertility.

Note: Jone's and Tompkin's metroplasties are surgeries for the septate uterus that involve a transabdominal approach. They cause uterine scarring and are hence not preferred.

Solution to Question 7:

The given clinical scenario of a G4A3 woman with miscarriages at 8 and 11 weeks, and preterm delivery due to preeclampsia at 24 weeks is suggestive of antiphospholipid antibody (APLA) syndrome.

APLA is an autoimmune disorder commonly associated with fetal death, preterm delivery, early-onset preeclampsia, and fetal growth restriction due to placental insufficiency and thromboses. Pregnancy loss usually occurs before 10 weeks.

Diagnosis of APLA requires at least 1 clinical criterion and 1 laboratory criterion.

• Obstetric criteria

• One or more unexplained deaths of a morphologically normal fetus at or beyond 10 weeks or

• Severe preeclampsia or placental insufficiency requiring delivery before 34 weeks, or

• Three or more unexplained consecutive spontaneous abortions before 10 weeks

• Laboratory criteria

• Presence of lupus anticoagulant, or

• Anticardiolipin antibodies, or

• Anti-β2 glycoprotein antibody

Recurrent pregnancy loss (RPL), also referred to as a recurrent miscarriage or habitual abortion, is historically defined as 3 consecutive pregnancy losses prior to 20 weeks from the last menstrual period.

Common causes include:

• Parental chromosomal abnormalities

• Untreated hypothyroidism

• Uncontrolled diabetes mellitus

• Uterine anatomic abnormalities,

• Antiphospholipid antibody syndrome

Other options:

Option A: There is no history of STD or skin lesions suggestive of syphilis and according to Kassawitz rule, in women with untreated syphilis having a series of pregnancies, the likelihood of fetal infection in later pregnancies becomes less; usually, only the first pregnancy is an abortion, the following pregnancy losses will be stillbirths in the next trimester rather than recurrent pregnancy loss.

Option C: TORCH infections may play a role in sporadic spontaneous pregnancy loss, rather than recurrent pregnancy losses.

Option D: Gestational diabetes mellitus (GDM) refers to the onset or first recognition of diabetes during pregnancy, usually at the 24-28 weeks screening test. Women with uncontrolled overt or pregestational diabetes with elevated HbA1c or preprandial glucose in the first trimester have an increased risk for early miscarriage, but there is a history or lab findings suggestive of gestational diabetes mellitus.

Solution to Question 8:

The most likely diagnosis in a woman with lactational failure and postpartum amenorrhea, now presenting with fatiguability and constipation is Sheehan syndrome. A history of multiple blood transfusions after delivery indicates major obstetric blood loss, which resulted in pituitary ischemia leading to this condition.

Sheehan syndrome is panhypopituitarism due to pituitary ischemia and necrosis following massive obstetrical hemorrhage.

The most severe form presents with shock due to pituitary apoplexy. It is characterized by a sudden onset headache, nausea, visual deficits, and hormonal dysfunction due to acute pituitary hemorrhage or infarction. Less severe forms present with anovulation and amenorrhea.

Damage to specific pituitary cell types leads to lactational failure, loss of axillary and pubic hair, and symptoms of hypothyroidism or adrenal insufficiency. Prolactin deficiency is most commonly encountered, but adrenal insufficiency is the most life-threatening complication. Hence, the adrenal function should be immediately assessed if there is any suspicion of Sheehan syndrome.

Option B: Hypothyroidism can present with vague symptoms such as fatigue, constipation, and amenorrhea. However, postpartum hypothyroidism usually presents 4-8 months after delivery after a thyrotoxic phase with palpitations and thyroid enlargement.

Option C: Premature ovarian failure presents with menopausal symptoms. It is due to primary ovarian dysfunction. It is diagnosed when serum FSH levels are >30-40 mIU/mL on two occasions, at least one month apart.

Option D: Delayed transfusion reaction presents days to weeks after the transfusion. It may be asymptomatic or with low-grade fever. Coombs test may be positive, unconjugated bilirubin increased, and the rise in hemoglobin levels following the transfusion will be less than expected.

Solution to Question 9:

Among the given options, hysteroscopy with laparoscopy is used to confirm the diagnosis.

For infertility evaluation, hysteroscopy and laparoscopy can detect müllerian anomalies, screen for co-existent endometriosis, and exclude other tubal or uterine cavity pathology.

This approach can differentiate between a septate and a bicornuate uterus. On laparoscopy, the presence of a uterine fundus indicates that the uterus is septate, while its absence indicates it is bicornuate. The capacity of each horn in a bicornuate uterus and the depth and thickness of the uterine septum prior to surgical resection can also be assessed.

MRI is the gold standard investigation for the diagnosis of Mullerian anomalies. It clearly delineates both the internal and external uterine anatomy. Complex anomalies and common associations like renal or skeletal anomalies can also be evaluated. It is also less invasive than hysteroscopy and laparoscopy in diagnosing Mullerian anomalies.

Other options:

Option A: Hysterosalpingography (HSG) provides information about the interior contour of the uterine cavity but cannot accurately visualize the external contour of the uterus.

Option C: Laparoscopy alone as a diagnostic procedure cannot visualize the interior contour of a hollow viscus like the uterus. This can be done only with the help of hysteroscopy or imaging such as sonography, HSG, or MRI.

Option D: Transvaginal sonography is used for the initial detection of uterine anomalies causing infertility or spontaneous abortions. Complete evaluation of these anomalies is done by MRI.

Note:

• Hysteroscopy is the gold standard investigation to visualize the uterine cavity and tubal ostia. It also has therapeutic benefits in cases of endometrial polyps, submucous resection of myoma, hysteroscopic adhesiolysis, and resection of uterine septum.

• Laparoscopy is the gold standard investigation to evaluate tubal factors of infertility.

Solution to Question 10:

The presence of infra-umbilical flattening and fetal heart sounds on the lateral side indicates that the fetus is in the occipito-posterior position.

Occipito-posterior (OP) position of the fetus is when the occiput is situated posteriorly over the sacroiliac joint, or directly over the sacrum. The fetus is in vertex presentation.

Examination findings of OP position:

• On inspection, the abdomen appears flattened below the umbilicus (infra-umbilical flattening).

• Pelvic grips:

• The head is not engaged.

• The sinciput (cephalic prominence) is not prominently felt.

• Auscultation: Fetal heart sounds are heard maximally over the flanks. However, in the direct OP position, they are heard in the midline.

There are three variants of OP position:

• Left OP position (fourth position of the vertex) - When the occiput points towards the left sacroiliac joint.

• Right OP position (third position of the vertex) - When the occiput points towards the right sacroiliac joint. It is more common than the left OP position.

• Direct OP position - When the occiput points directly towards the sacrum.

All three positions may be primary (present before the onset of labor) or secondary (develops after the onset of labor).

Other options:

Option A: Brow presentation is a rare type of cephalic presentation where the position of the head is in between full flexion and full extension. The part of the head between the orbital ridge and the anterior fontanel presents at the pelvic inlet.

Option B: Knee presentation is a type of breech presentation. Here, the thighs are extended but the knees are flexed, presenting at the pelvic inlet. The head will be palpable as a hard globular mass at the fundus.

Option D: Right dorso-anterior position is when the fetal back is directed anteriorly. This occurs in a transverse lie. On examination, the abdomen appears wide, and a hard plane of resistance is felt. The fundus extends just above the umbilicus. The head is felt in one iliac fossa and the breech is felt in the other.





Solution to Question 11:

The given clinical scenario of a primi in labor pain with multiple, painful vesicular lesions on her vulva is suggestive of an active herpes simplex virus (HSV) infection. Management involves the administration of acyclovir and elective cesarean section.

HSV infection in pregnancy can be caused by either HSV-1 or HSV-2 and transmitted by sexual contact. The infection can be primary, nonprimary, first episode, and recurrent.

HSV infection (both HSV 1 and 2) during pregnancy has a risk of vertical transmission of the infection to the fetus. The risk is higher in the primary than in the recurrent infection. The maximum risk of fetal infection is during labor, although it can occur in utero and post-natally from direct contact.

Suspicious lesions should be sent for PCR testing or culture. Antiviral treatment with acyclovir should be given if found positive. Women with no active genital lesions, asymptomatic, or those with active lesions in non-genital areas can undergo normal labor and delivery.

Management of HSV infection during pregnancy:

• Oral acyclovir or valacyclovir for primary or recurrent infections during pregnancy.

• IV acyclovir followed by oral acyclovir for severe or disseminated infection

• Oral analgesics or topical anesthetics for pain relief

• Catheterization for urinary retention

Cesarean delivery reduces the risk of vertical transmission during labor and is indicated in:

• Cases of active genital lesions - both primary and recurrent

• Prodromal symptoms like vulval itching or burning

Other options:

Option B: The presene of active lesions is a contraindication for induction of labor.

Option C: Spontaneous delivery is contraindicated in the presence of active lesions.

Option D: Even after treatment with acyclovir, normal delivery is not indicated owing to the risk of dormant viral transmission to the fetus.

Solution to Question 12:

The most appropriate management for a woman with infertility due to intramural fibroids at both the cornual ends of the uterus is myomectomy. The infertility, in this case, is due to the fibroids blocking the right and left cornua of the uterus.

Myomectomy refers to the removal of fibroids and preserving the uterus. It is indicated in women presenting with infertility, those who desire pregnancy, or those who wish to retain the uterus.

Intracavitary fibroids can be resected via the hysteroscopic approach, while subserosal or intramural fibroids require laparotomy or laparoscopic myomectomy.

Other options:

Option A: Uterine artery embolization is not done in this case, as it may decrease fertility, as a result of pelvic adhesions. It is a technique in which polyvinyl alcohol (PVA), gel foam particles, or metal coils are injected into the uterine arteries. This embolization reduces vascularity and the size of the fibroid. It is best suited to control menorrhagia in multiparous women.

Option C: Artificial reproductive techniques are not indicated in this case as infertility is seen to be due to the fibroids blocking the cornual ends of the uterus. However, it may be considered at a later stage.

Option D: GnRH analogs when used for 6 months, can reduce the tumor size by targeting the growth effects of estrogen and progesterone. It can be used in women with infertility caused by cornual fibroids. They are often used preoperatively to reduce the size of the fibroid and facilitate a hysteroscopic approach to surgery. However, it is expensive and can cause menopausal symptoms if used for >6 months. Fibroids are known to regrow 3-4 months after stopping the

therapy.

Solution to Question 13:

The given clinical scenario of a young woman with copious, non-pruritic (without itching) vaginal discharge occurring 13 days after menstrual period (period of ovulation) is most likely to be leucorrhoea (physiological).

The term leucorrhoea can be used in conditions where normal vaginal secretion is increased in amount. An increase in this normal vaginal secretion develops physiologically:

• At puberty

• At ovulation

• During pregnancy

• During the premenstrual phase, occasionally

It may be cervical or vaginal in origin. A speculum examination of the vagina can distinguish between the two. If the discharge is cervical in origin, an excessive mucoid discharge will be visible at the external os.

Normal discharge appears clear or white, without any odor or itching, and will have a pH of 3.8-4.2.

Other options:

Option A: Candidiasis is caused by infection with Candida albicans and presents with pruritus, burning, and dysuria with vulvar erythema, edema, and excoriations. A curdy white discharge will be present.

Option B: Bacterial vaginosis is caused by an alteration in the normal vaginal flora and presents without itching and a malodorous grey-white discharge. It is diagnosed using Amsel's criteria, where at least three of the four following criteria should be present to confirm the diagnosis:

• Homogenous vaginal discharge

• Vaginal pH > 4.5

• Amine-like odor on adding KOH (Whiff test positive)

• Clue cells (>20% of cells)

Option C: Trichomoniasis is caused by Trichomonas vaginalis and presents with a copious, frothy, offensive yellow-green discharge with vaginal soreness or irritation and urinary frequency. On examination, the vagina looks red and inflamed. The inflamed cervix looks like a strawberry with punctate hemorrhagic spots.

Solution to Question 14:

The order of ligation of vessels to arrest bleeding in atonic postpartum hemorrhage (PPH) is the uterine artery - ovarian artery - internal iliac artery.

The ascending branch of the uterine artery is ligated between the upper and lower segments of the uterus, at the lateral border. If the bleeding continues, the ovarian and uterine artery anastomosis is ligated just below the ovarian ligament. If the bleeding still persists, the anterior division of the internal iliac artery is ligated 5 cm distal to the common iliac bifurcation.

Unilateral or bilateral ligation of the internal iliac artery ligation causes a reduction in the pulse pressure in the arteries distal to the ligation. This increases the chances of hemostasis via pressure and clot formation.

Surgical methods to control PPH are outlined as stepwise uterine devascularization procedures:

• Compression sutures: B-Lynch suture

• Ligation of uterine arteries

• Ligation of ovarian and uterine anastomosis

• Ligation of internal iliac artery

• Angiographic embolization of the bleeding vessel

Pelvic packing (usually done prior to embolization) and hysterectomy are other surgical methods employed to control PPH.



Solution to Question 15:

The most common complication that can be seen in this patient is endometriosis.

Transverse vaginal septum leads to obstructed flow of menstrual blood. This may lead to retrograde menstruation, which increases the likelihood of the development of endometriosis, according to Sampson's theory of endometriosis.

Transverse vaginal septum occurs as a result of the failure of Müllerian duct fusion or failure of vaginal plate canalization. It may be obstructive or non-obstructive. The commonest site of a transverse septum is the junction between the upper and middle third of the vagina.

Patients with an obstructive or imperforate vaginal septum present with symptoms similar to the imperforate hymen. Symptoms include amenorrhoea, cyclical abdominal pain, urinary symptoms, and a palpable pelvic mass. The vagina may appear shortened and the cervix may not be identifiable. Secondary sexual characteristics appear normal. The diagnosis can be confirmed with ultrasound or MRI.

There is an increased risk of endometriosis. Other complications such as pyomucocolpos, pyometra, and pyosalpinges can also develop in infants with obstructive vaginal septums.

Treatment is by manual dilatation of the perforation or surgical excision of the septum.

Other options:

Option B: Tubo-ovarian abscess develops in pelvic inflammatory disease.

Option C: Dermoid cysts are cystic ovarian tumors that are often associated with mucinous cystadenomas.

Option D: Theca lutein cysts are found in association with hydatidiform moles, choriocarcinoma, and gonadotropin or clomiphene citrate therapy. They regress with the management of the mole and choriocarcinoma and discontinuation of gonadotropin therapy.

Solution to Question 16:

The obstetric score of a pregnant woman who previously delivered twins would be G2P1 (gravida 2 para 1).

The term gravida refers to the number of times the woman has been pregnant, irrespective of whether it crossed the period of viability. The current pregnancy is included in this count. Since she has been pregnant once before with twins and is currently pregnant now, gravida will be 2.

The term para refers to the number of previous pregnancies reaching a period of viability (gestation of 28 weeks). It does not increase to a higher number if multiples (twins/triplets) are delivered in a given pregnancy and still count as one birth. A current pregnancy, if any, is not included in this count. Therefore, para here will be 1 since she had one birth of twins and the present pregnancy is not counted.

Both gravidity and parity refer to the pregnancies and not the fetuses.

Note: The period of viability is when the fetus is capable of independent existence. The WHO defines it to be 24 weeks while the Indian definition is 28 weeks.

Solution to Question 17:

A woman who has missed 4 oral contraceptive pills in the last two weeks should take one hormonal pill as soon as possible and finish all hormonal pills in the pack as scheduled, throw away the 7 non-hormonal pills in a 28-pill pack and start a new pack the next day.

If any non-hormonal pills were missed, she can discard the remaining pills. Start the new pack of oral contraceptives and continue taking as usual.

If she vomits within 2 hours of taking a pill, another pill from the pack should be taken as soon as possible. After that, continue taking the pills as scheduled. If the vomiting or diarrhea persists for >2 days, follow the instructions for 1 or 2 missed pills as mentioned above.

The contraceptive method should be discontinued and changed in the following situations:

• Unexplained or prolonged heavy vaginal bleeding that suggests an underlying condition not related to the method

• Starting anticonvulsant medication or treatment with rifampicin

• Migraines

• Non-ambulatory conditions lasting for >1 week like major surgery, leg in a cast, etc. Combined oral contraceptives can be restarted 2 weeks after ambulation.

• Suspected heart or liver disease, hypertension, deep vein thrombosis, stroke, breast cancer, diabetes, or gall bladder disease

Missed pills	Management
Missed 1 or 2 pills/ started new pack 1 or 2 days late?	You have two options: take one hormonal pill right away, or take two pills at your usual scheduled time.The risk of pregnancy is minimal to none.
Missed 3 or more pills in the first or second week/started new pack 3 or more days late?	Take one hormonal pill immediately and then continue taking your pills as scheduled.Make sure to use a backup method of contraception for the next seven days.Additionally, if you've had sexual intercourse within the last 72 hours, consider taking emergency contraceptive pills.
Missed 3 or more pills in the third week?	Take one hormonal pill as soon as you can and complete the entire pack of hormonal pills according to the schedule. Discard the seven non-hormonal pills in a 28-pill pack.Begin a new pack the following day.
Missed any nonhormonal pills? (last 7 pills in 28-pill pack)	Dispose of any missed non-hormonal pills.Continue taking your combination oral contraceptives (COCs), one pill each day.Begin the new pack as you normally would.

Solution to Question 18:

A patient on infertility treatment, with 6 weeks of amenorrhea, abdominal pain, and vaginal bleeding with beta hCG levels of 2800 mIU/mL, and a gestational sac in the left adnexa points to the diagnosis of unruptured ectopic pregnancy and the most appropriate management is single-dose methotrexate.

A pregnancy is considered ectopic if implantation occurs at a site outside the uterine cavity. The majority of ectopic pregnancies occur in the fallopian tube, the ampulla being the most common site.

Management of ectopic pregnancy:

Expectant management: Ectopic pregnancies measuring <3 cm and with declining beta hCG levels initially <1500 mIU/mL may resolve without intervention and can be managed expectantly.

Medical management with single or multiple-dose methotrexate is preferred if:

• Hemodynamic stability

• Serum beta-hCG ≥1500 mIU/ml and ≤5000 mIU/mL

• Mean sac diameter <3.5cm with no fetal cardiac activity detected on transvaginal ultrasound.

• Patients are willing and able to comply with post-treatment follow-up and have access to emergency medical services within a reasonable time frame in case of a deterioration in the clinical status.

Methotrexate(MTX) can be given as a single IM dose of 50 mg/m2 or as a multiple-dose regimen of 1mg/kg on days 1, 3, 5, and 7.

Surgical management with salpingectomy or salpingostomy is done when:

• Criteria for expectant and medical management are not met

• Gestational sac >3.5 cm

• Fetal cardiac activity present

• Beta hCG > 5000 mIU/mL

• Significant pain and hemoperitoneum

• Failure of expectant or medical management

• Clinical status of patient warrants urgent intervention

• Doubtful compliance with medical management

Milking of the tube through the fimbriae carries a high risk of bleeding, damage to the tube, the persistence of trophoblastic tissue, and recurrence of ectopic pregnancy.

Solution to Question 19:

The best method of assessing the gestational age of a patient on oral contraceptives with 6 weeks of amenorrhea is by measuring the crown-rump length (CRL) using ultrasonography.

Gestational age refers to the duration of pregnancy calculated from the first day of the LMP. It is usually expressed in weeks. Following the use of oral contraceptives, ovulation may have been later than 2 weeks after the LMP. Therefore, gestational age calculated with the help of LMP will be inaccurate.

First-trimester ultrasound measurement of CRL is the best method to assess gestational age. It is measured from the vertex to the coccyx.

• CRL (in mm) + 6.5 = gestational age in weeks (or)

• CRL (in mm) + 42 days = gestational age in days.

CRL measurement is shown in the image below:



Ultrasonographic parameters for estimation of the gestational age according to the trimester are as follows:

Other options:

Option B: Adding 280 days to the LMP will provide the expected delivery date (EDD). Adding 9 months + 7 days to the LMP or 280 days to the LMP is Naegele's formula for calculating the EDD.

Option C: Measurement of fundal height in cm (with a variation of 1-2cm) is roughly equal to the gestational age in weeks, after 24 weeks of pregnancy. This is measured from the superior border of the symphysis pubis to the top of the uterine fundus when the bladder is empty.

Trimester	Parameter
1st	Crown-rump length (CRL)
2nd and 3rd	Biparietal diameter (BPD)Abdominal Circumference (AC) Head Circumference (HC)Femur Length. (FL)

Obstetrics and Gynecology INICET 2022

Question 1:

A 25-year-old lady had an uneventful vaginal delivery. On postpartum day 1, she presented with postpartum haemorrhage and complains of visual changes and severe headache. A few hours after the complaint, she is found to be in an unresponsive state and is intubated and put on ventilation. On further examination, she has a BP of 60/40 mmHg and tachycardia, blood sugar is 48 mg/dL. Hematocrit and WBC count was normal. Which of the following drugs would help reverse this state?

a) Inj. dobutamine

b) Steroids

c) Thyroxine

d) Activated protien C

Question 2:

All are true about biophysical profile (BPP) except:

a) Includes breathing, movements, tone, and AFI

b) Can tell us about fetal outcome and fetal pH

c) Score is inconsistent with fetal pH at different gestational ages

d) Score <4: fetal pH <7.2; Score 0 : 100% fetal acidemia

Question 3:

Which of the following is not true about lupus nephritis in pregnancy?

a) Immunosuppression can be continued during pregnancy

b) Pregnancy to be planned once the disease has been quiescent for at least 6 months and there is no evidence of renal dysfunction

c) High dose corticosteroids for lupus flare in pregnancy is safe

d) Ecospirin, methotrexate, cyclophosphamide, corticosteroids, azathioprine are safe in pregnancy

Question 4:

Which of the following is false regarding management of diabetes in pregnancy?

a) Elective C-section has no role in reducing incidence of brachial plexus injury

b) In a patient being planned for induction of labor, night dose of intermediate insulin is given as planned, and the morning dose is withheld

c) In active labor, if RBS <70 mg%, D5 is started at 100-150 ml/hr till the RBS is >100 mg%

d) Capillary blood glucose monitoring levels are kept at fasting- 95 mg%; 1 hr postprandial- 140 mg%; 2 hrs postprandial- 120 mg%

Question 5:

Which of the following statements regarding $MgSO_4$ is false?

a) $MgSO_4$ causes uterine relaxation, but is dose dependant

b) $MgSO_4$ is continued upto 24 hours after delivery or last attack of seizure

c) It has inhibitory effect on cerebral cortex thereby preventing seizures and lowering the BP

d) In renal failure, $MgSO_4$ loading dose can be given, but maintenence dose should be adjusted based on renal function

Question 6:

A couple already has 3 children and has been trying for another one for the last 1 year. However, the husband is diagnosed with hypogonadotropic hypogonadism. Which of the following is/are true:

a) 1, 2, 3, 4 and 5

b) 1, 2 and 5

c) 2, 3, 4 and 5

d) 1, 3 and 5

Question 7:

All of the following are theories of causation of preeclampsia except:

a) Prostaglandins play an important role. Thromboxane A2 decreases; Prostacyclin:Thromboxane ratio increases in preeclampsia

b) Endotheilal cell dysfunction leads to reduced nitric oxide levels and this causes the clinical presentation of preeclampsia

c) Vasoconstriction leads to ischemia, necrosis and hemorrhage and end organ damage seen in preeclampsia

d) Anti-angiogenic factors play an important role in preeclampsia. There is an increase in sFlt-1 and decrease in PIGF.

Question 8:

The following cost-effective investigations are routinely recommended in the screening of antenatal mothers except?

a) Echocardiography for cardiac disease

b) Urine analysis for bacteriuria

c) Blood sugar levels for GDM

d) VDRL for syphilis

Question 9:

Which of the following cerclage procedures would be done in a nonpregnant female through a transabdominal approach?

a) Lash and Lash cerclage

b) Benson and Durfee cerclage

c) Wurms cerclage

d) Laparoscopic cerclage

Question 10:

Urinary tract infection in a pregnant lady in her 2nd trimester is best defined as:

a) Growth of 10^3 coliform in suprapubic sample

b) Growth of 10^5 coliform in voided urine sample

c) Frequency with dysuria with >3 WBC/HPF

d) Frequency with dysuria with positive nitrite and leucoesterase dipstick test

Question 11:

Which of the following is incorrect regarding innervation of the uterus?

a) Sensory level is from T10 to L1

b) Uterine contractility is mediated by innervations from level T7 - T8

c) In the 1st stage of labor, pain is due to the fibres at level of T10 - L1

d) In early labor, pain is usually because of the uterine contraction

Question 12:

Seminal vesicles and vas deferens would be bilaterally absent congenitally in which of the following conditions?

a) Cystic fibrosis

b) Klinefelter syndrome

c) Kallmann syndrome

d) Kartagener syndrome

Question 13:

Which of the following graphs represents normal response to OGTT?



a) Graph A

b) Graph B

c) Graph C

d) Graph D

Question 14:

Kassowitz rule is related to:

a) Primary syphillis

b) Congenital syphillis

c) Secondary syphillis

d) Latent syphillis

Question 15:

Modified biophysical profile includes:

a) NST and fetal tone

b) NST and AFI

c) NST and breathing movements

d) NST and gross fetal movements

Question 16:

All of the following statements about anti-Mullerian hormones are true, except:

a) It is best tested in the fasting state on day 2 of menstrual cycle

b) Lipopeptide secreted from granulosa cell of early antral follicle

c) Correlates with pregnancy rate of IVF cycle

d) It decreases sensitivity of the pre-antral follicles to FSH

Question 17:

Which of the following statements is false regarding post-partum hemorrhage?

a) Start colloids initially, 4 times the blood lost

b) Primary PPH occurs within 24 hours of childbirth

c) PPH is loss of 500 ml of blood in vaginal delivery

d) Causes include retained placenta and atony

Question 18:

Recommended HPV vaccination schedule for 9 to 14 year olds according to WHO SAGE guidelines is:

a) 1 or 2 doses

b) 2 doses

c) 1 dose

d) 3 doses

Question 19:

All of the following coagulation factors increase in pregnancy except:

a) Factor II

b) Factor VII

c) Factor VIII

d) Factor XI

Question 20:

A pregnant lady at 20 weeks of gestation comes with complaints of acute onset fever, and pain in the right lumbar region. It is associated with chills, nausea, and vomiting. Urinalysis reveals the presence of E.coli and a diagnosis of acute pyelonephritis is made. Which of the following is incorrect regarding the management of acute pyelonephritis in pregnancy?

a) Hospitalise, send urine and blood cultures and start on antibiotics.

b) Oral antibiotics once afebrile

c) Maintain urine output >50 ml/hr; Intravenous crystalloids can be given

d) Discharge if afebrile for 72 hrs; Continue antibiotics for 5 days

Question 21:

All of the following are the diagnostic criteria for preterm labor except?

a) 4 contractions in 20 min and 8 contractions in 60 min with changes in cervix

b) Cervical dilatation > 2 cm

c) More than 80 percent effaced cervix

d) Cervix posterior in position

Question 22:

Which statement is false regarding late decelerations?



a) Starts after peak of uterine contraction and ends after uterine contraction ends

b) They are smooth and gradual

c) At least 10 beats, 2 times in a 20 min period above baseline

d) Not associated with accelerations

Question 23:

What is the age group distribution for vulval cancer?

a) Not age specific

b) 35-55 years

c) 55-85 years

d) Both of the above

Question 24:

The following are correct regarding normal placenta, except _____.

a) Placenta weight at term is 470 g

b) Round-oval with 22 cm diameter

c) Extraplacental membranes and 2-vesseled umbilical cord present

d) Central thickness >2.5 cm

Question 25:

Arrange the following steps of dilation and curettage in sequence.

a) 4-3-2-1

b) 4-2-3-1

c) 4-3-1-2

d) 4-1-2-3

Question 26:

All are long-term reversible methods of contraception, except:

a) Progestin injection

b) Progestin implants

c) Female sterilization

d) Levonorgestrel-releasing and copper-containing intrauterine devices

Question 27:

According to new WHO guidelines, what is the minimum number of antenatal visits recommended to have a positive pregnancy outcome?

a) 8

b) 4

c) 6

d) 10

Question 28:

Identify the wrong statement regarding the period of gestation calculation.

a) 1st trimester USG is the best assessment

b) Calculated from the last day of the last menstrual period

c) In an IVF pregnancy, day of embryo transfer/day of fertilization is needed to calculate POG

d) Crown-rump length is the best measure of period of gestation in the first trimester

Question 29:

All of the following can be used in first-trimester medical termination of pregnancy (MTP), except _____.

a) Mifepristone + misoprostol

b) Manual vacuum aspiration

c) Dilatation and curettage

d) Extra-amniotic instillation of ethacridine lactate

Question 30:

A woman with eclampsia is started on magnesium sulfate. What is the first sign of magnesium sulfate toxicity?

a) Loss of knee jerk

b) Hypotension

c) Respiratory depression

d) Reduced muscle tone

Question 31:

Which of the following patients would be classified as having primary amenorrhea?

a) 11-year-old girl with Tanner stage 1 breasts

b) 14-year-old girl with Tanner stage 1 breasts

c) 13-year-old girl with Tanner stage 5 breasts

d) 12-year-old with Tanner stage 4 breasts

Question 32:

The daughter of a woman who received diethylstilbestrol during her pregnancy is at higher risk for:

a) Squamous cell carcinoma

b) Clear cell carcinoma

c) Adenocarcinoma

d) Sarcoma botryoides

Question 33:

Which of the following is not a component of biophysical profile?

a) Amniotic fluid volume

b) Fetal tone

c) Daily fetal movement count

d) Non-stress test

Question 34:

All are causes of uterine dysfunction, except _____.

a) High fetal head station

b) Multiparity

c) Neuraxial block

d) Chorioamnionitis

Question 35:

Non-immune hydrops is seen in all, except _____.

a) Human immunodeficiency virus

b) Herpes simplex virus

c) Syphilis

d) Parvovirus B19

Question 36:

Which of the following statements is true regarding recurrent ovarian cancer?

a) Platinum resistant if recurrence > 12 months

b) Partial platinum resistant if recurrence is within 6-12 months

c) Platinum sensitive if recurrence > 6months

d) All of the above

Question 37:

What can be said about this graph?



a) This is a non-stress test.

b) The baseline shows that the fetus has tachycardia.

c) The beat to beat variability is less/inadequate.

d) This is a uterine contraction graph.

Question 38:

The given image depicts:



a) Succenturiate lobe

b) Velamentous insertion of the cord

c) Both succenturiate lobe and velamentous insertion

d) Normal placenta

Question 39:

Which hormone is secreted by placenta?

a) LH

b) hCG

c) FSH

d) GnRH

Question 40:

Antepartum hemorrhage due to abruption shows all except:

a) Uterine tenderness

b) Frequent contractions

c) Painless vaginal bleeding

d) Mild to moderate vaginal bleeding

Question 41:

A woman presents at 8 weeks of gestation. Which of the following signs can be elicited at this gestation?

a) 1,3

b) 1,2

c) 1,4

d) 3,4

Question 42:

The non-contraceptive uses of oral contraceptives include all of the following except:

a) Reduction in the risk of ovarian cancer

b) Ectopic pregnancy

c) Protection against sexually transmitted diseases including HIV

d) Reduction in the risk of endometrial cancer

Question 43:

A 16-year-old presents with absent menses. Examination shows widely spaced nipples, webbed neck, and Tanner grade 1 breast development. USG shows streak ovaries. Investigation shows raised FSH, raised LH, and reduced estradiol. What is the diagnosis?

a) Turner syndrome

b) Kallmann syndrome

c) Androgen insensitivity syndrome

d) Mayer-Rokitansky-Küster-Hauser (MRKH) syndrome

Question 44:

Consider the following actions under AMTSL. Arrange them in the right order:

a) 3,1,2,4

b) 3,2,4,1

c) 2,4,3,1

d) 2.4.1.3

Question 45:

A surgically resected specimen of an ovarian mass is is shown below. What is the likely diagnosis?

a) Immature teratoma

b) Mature cystic teratoma

c) Mucinous cystadenoma

d) Serous cystadenoma

Question 46:

Which of the following is seen in placenta previa?

a) Warning bleeds

b) Dark painful bleeding

c) Uterine tenderness

d) Increased/large uterine size

Question 47:

A pregnant woman with 35 weeks+5 days gestation complains of epigastric pain, headache, and blurring of vision. Her blood pressure is 170/110 mmHg. She is in labor pain.

Examination of the cervix shows a 3 cm dilatation with a bag of liquor. What is the appropriate management?

a) Conservative management

b) Administer tocolytics and deliver at 37 wks

c) Give steroid cover and wait until steroid cover is complete.

d) Treat hypertension, administer $MgSO_4$, and consider termination of pregnancy.

Question 48:

All of the following are risk factors of vulval cancer except

a) Vulval dystrophy

b) Vulval hamartoma

c) Smoking

d) Vulval warts

Question 49:

Which of the following is true regarding perimortem cesarean section?

a) LSCS

b) Pfannenstiel incision

c) Upper segment cesarean section

d) Done in OT

Question 50:

Arrange the following in sequence.

a) B, A, C, D

b) B, A, D, C

c) B, C, A, D

d) B, D, A, C

Question 51:

Match the following

a) A-4, B-3, C-2, D-1

b) A-2, B-3, C-1, D-4

c) A-4, B-1, C-2, D-3

d) A-3, B-2, C-4, D-1

Question 52:

What is vagitus uterinus?

a) Cry of baby in the womb

b) Infection of vagina

c) Infection of uterus

d) Infection of both vagina and uterus

Answer Key

Question No.	Correct Option
1	b
2	c
3	d
4	a
5	c
6	d
7	a
8	a
9	d
10	b
11	b
12	a
13	d
14	b
15	b
16	b

17	a
18	a
19	d
20	d
21	d
22	c
23	c
24	c
25	c
26	c
27	a
28	b
29	d
30	a
31	b
32	b
33	c
34	b
35	a
36	c
37	a
38	c
39	b
40	c
41	b
42	c
43	a
44	b
45	b
46	a
47	d
48	b
49	c
50	a
51	c
52	a

Detailed Explanations

Solution to Question 1:

The given history of a pregnant having postpartum hemorrhage developing headache, visual changes, and hypotension after a vaginal delivery suggests Sheehan syndrome. Steroids are the drug that can reverse this state.

Volume depletion and low cardiac output leading to critically low BP are common in acute pituitary deficiency and recovery of normal cardiovascular status is rapidly achieved under hormone and volume replacement therapy without the need for inotropes like dobutamine.

Pituitary apoplexy is a hemorrhagic infarction of large pituitary adenomas or pituitary infarction due to traumatic major blood loss. If this blood loss occurs during delivery, it's called Sheehan's syndrome. The hyperplastic enlargement of the pituitary (size of gland increases by about 125%), which occurs normally during pregnancy, increases the risk of hemorrhage and infarction.

Pituitary apoplexy presents acutely as an endocrine emergency, with sudden onset of headache, nausea, visual deficits, persistent hypotension, tachycardia, and hypoglycemia leading to shock. The most prominent symptoms are lactation failure (due to a deficiency of prolactin) and postpartum amenorrhea (due to a deficiency of gonadotropins).

Because deficiencies of some or all pituitary-responsive hormones may develop after the initial insult, Sheehan syndrome can be heterogeneous and not be identified for years. It can present late with the following features like loss of pubic and axillary hair, hypothyroidism due to decreased TSH secretion, adrenal insufficiency due to low ACTH.

Treatment is by hormone replacement - first glucocorticoids, later - thyroid hormone, sex steroids, vasopressin, and growth hormones are given. If there is evidence of progressive visual loss, cranial nerve involvement, or loss of consciousness, immediate surgical decompression must be carried out.

Solution to Question 2:

All are true about BPP (biophysical profile) except option C. BPP score is not inconsistent with fetal pH, but has a direct linear relationship with fetal pH and thus can be used to predict fetal acidemia.

BPP is a multi-parametric test to assess uteroplacental insufficiency. It consists of five components/ variables and the observations are made in a 30-minute time period.

Components of BPP:

(Mnemonic: BATMaN)

B - Breathing movements of the fetus

A - Amniotic fluid volume

T - Tone of the fetus

M - Movements of the fetus

N - Non-stress test (NST)

Each parameter has a maximum score of 2 and a minimum score of 0. So the total maximum score will be 10. A score of 4 or less indicates fetal acidemia.

Interpretation and management of BPP scores:

BPP is directly related to fetal pH and fetal outcome. A lower BPP score is associated with a poor outcome (stillbirth, perinatal mortality) and lower APGAR scores at birth.

A score of <10 is associated with a pH of <7.2 and a score of 0 is associated with 100% fetal acidemia.

Note: The modified biophysical score has only 2 components i.e. NST and amniotic fluid index only. It is considered abnormal (nonreassuring) when the NST is non-reactive and/or the AFI is <5.

Parameters	Minimal Normal Criteria
Fetal breathing movements	≥ 1 episode lasting > 30 seco nds
Amniotic fluid volume (AF V)	≥ 1 pocket measuring 2 cm in two perpendicular planes (2 ×2 cm pocket)
Fetal muscle tone	≥ 1 episode of active limb or t runk extension with the retur n of flexion; Eg. opening and closing of the hand, consider ed normal
Gross body movements	≥ 3 discrete body/limb move ments
Non-stress Test (NST)	Reactive pattern

BPP score	Interpretation	Management
10	Normal, nonasphyxiated fetu s	Repeat test weekly (or biweek ly in diabetic mother or post-term pregnancy)
8with normal AFV	Normal, nonasphyxiated fetu s	Repeat test weekly (or biweek ly in diabetic mother or post-term pregnancy)
8 with decreased A FV	Suspicion of chronic fetal asp hyxia	Delivery
6	Possible fetal asphyxia	Deliver, if AFV is abnormal o r >36weeks or if the repeat te st score is ≤ 6. If the repeat s core >6, observation and rep eat test weekly
4	Probablefetal asphyxia	Repeat testing same dayIf the repeat score ≤6, delivery
0-2	Almostcertainfetal asphyxia	Delivery

Solution to Question 3:

Option D is not true about lupus nephritis in pregnancy. Methotrexate and cyclophosphamide are teratogenic and hence they are not safe during pregnancy.

Systemic lupus erythematosus(SLE) is a chronic autoimmune disorder involving the multisystem. It affects young women of childbearing age. The renal involvement of the disease can manifest in the form of active lupus nephritis or flare-ups during pregnancy.

Clinical features include fever, weight loss, migratory joint pain, and myalgias. It can lead to preterm births, preeclampsia, and pregnancy loss while the infant can suffer from neonatal lupus and heart block.

Antinuclear antibodies are useful for screening the disease. Double-stranded DNA (ds DNA) is highly specific in diagnosing. Other antibodies such as lupus anticoagulant, antiphospholipid antibodies, anti-Ro, and anti-La antibodies are useful in diagnosis.

Management is as follows

• Preconceptional planning plays an important role. Conception during a period of quiescence for at least 6 months determines a successful outcome. Stable remission of renal disease for at least 6 months before conception is recommended.

• Corticosteroids are useful during episodes of lupus flare and can be used safely in pregnancy. For patients on chronic corticosteroid therapy, intravenous hydrocortisone 100 mg, three doses are given 8 hourly during labor or at the time of cesarean delivery.

• Immunosuppressants such as azathioprine can be continued during pregnancy. It may be used in case of severe SLE

• Hydroxychloroquine is found to be the best for maintenance and is safe in pregnancy. It can prevent the need for prolonged corticosteroid therapy.



Solution to Question 4:

False statement regarding management of diabetes in pregnancy is option A. Cesarean section is found to be protective against traumatic injuries such as brachial plexus injury, in macrosomic babies born to mothers with diabetes. An elective C-section is found to be more protective than an emergency C-section.

Almost 50 % of pregnant women with diabetes are delivered by c-section. Indications for c-section in diabetic pregnant women are-

• Fetal macrosomia of ≥4.5kgs to avoid trauma to mother or neonate, and also to avoid brachial plexus injury in the baby

• Unsatisfactory progression of labor

• Complicated diabetes mellitus

• Fetal compromise during delivery

• Elderly primigravida (age >30 years)

• Multigravida with bad obstetric history previously

• Pregnancy that is associated with complications like preeclampsia, polyhydramnios or malpresentation

Other options:

Option B: In a planned delivery, although the bedtime dose of intermediate-acting insulin is given, it is recommended that the morning dose must be withheld. This is due to reduced insulin requirements after delivery. Therefore, regular insulin is used for glycemic control during this time.

Option C: In active labor, if RBS (Random blood glucose) <70 mg%, D5 (5% dextrose) is started at 100-150 ml/hr till the RBS is >100 mg%. Before initiation of labor, normal saline is used for iv infusion.

Option D: Self-monitoring of capillary blood glucose is advised in a pregnant woman diagnosed with diabetes. Subcutaneous continuous glucose monitoring devices can also be used for the same. The glucose goals recommended are:

• Fasting: ≤95 mg/dL

• 1 hour post-prandial: ≤140 mg/dL

• 2 hrs post-prandial: ≤120 mg/dL

Solution to Question 5:

The false statement regarding $MgSO_4$ (Magnesium sulfate) is option C. It has an inhibitory effect on the cerebral cortex, thereby preventing seizures, but this action is not responsible for lowering blood pressure.

$MgSO_4$ is mainly used in preeclampsia due to its specific anticonvulsant action and neuroprotective action on the cortex. $MgSO_4$ acts as a membrane stabilizer. It reduces motor endplate sensitivity to acetylcholine and blocks neuronal calcium influx. It is not used to treat hypertension, although it shows some hypotensive action by decreasing vascular response to vasopressors.

$MgSO_4$ is used for the management of eclampsia and as prophylaxis in severe pre-eclampsia. The therapeutic level of magnesium in eclampsia is 4-7mEq/L. The two regimens commonly used are Pritchard and Sibai regimen.

Other actions:

• Cerebral vasodilatation

• Uterine artery vasodilatation

• Uterine tocolytic i.e. inhibits uterine contractility

• Increased production of endothelial prostacyclin

• Inhibits platelet activation

Other options:

Option A: $MgSO_4$ inhibits uterine contractility but is dose-dependent (only above serum values of 8-10 mEq/L).

Option B: $MgSO_4$ has to be administered up to 24 hours after the last seizure (in postpartum seizure) or delivery, whichever is later. For recurrent seizures, 2 g IV bolus is given in the above regimen.

Option D: Magnesium is cleared almost exclusively by renal excretion. But, in renal failure, the initial 4-g loading dose of magnesium sulfate can be safely administered regardless of renal function. Only the maintenance dose needs to be altered according to the renal function. This is because the loading dose is responsible for reaching therapeutic levels, but the maintenance doses are responsible for maintaining steady states.

Regimen	Loading Dose	Maintenance Dose
Intramuscular (Pritchard)	4g (20% solution) IV over 3-5 minutefollowed by 10g (5O %), deep IM(5 gm in each bu ttock)	5g (5O%) IM 4 hourly in alte rnate buttocks
Intravenous (Zuspan or Sibai)	4g IV slow over 15-20 minute	1-2g/h IV infusion

Solution to Question 6:

The given patient is diagnosed with hypogonadotropic hypogonadism. True statements about the condition are 1, 3, and 5.

Hypogonadotropic hypogonadism in a male is the presence of impaired testicular function (hypogonadism), secondary to a hypothalamic-pituitary dysfunction (hypogonadotropic).

Causes include:

• Congenital

• Anosmic hypogonadotropic hypogonadism (Kallmann syndrome)

• Normosmic isolated hypogonadotropic hypogonadism (idiopathic hypogonadotropic hypogonadism)

• Acquired

• Drugs and alcohol

• Infiltrative or infectious pituitary lesions, brain radiation, trauma to the brain

• Hyperprolactinemia

• Systemic diseases such as hemochromatosis, sarcoidosis

A male with hypogonadotropic hypogonadism will have the following findings:

• Hormone levels

• Low LH (Luteinizing hormone), low FSH (follicle-stimulating hormone). This can be confirmed on GnRH stimulation test showing persistently reduced LH, and FSH levels.

• Low testosterone.

• High prolactin levels and high prolactin: testosterone ratio.

• Low testicular volume - oligospermia: the presence of less than 15 million sperm per milliliter of semen. Sperm motility is not impaired.

• MRI showing anomalous or absent olfactory bulb in Kallmann syndrome.

Treatment:

• Testosterone supplementation: Helps reverse the symptoms and signs of hypogonadism. It is advised only in males not planning to have children.

• Pulsatile GnRH or gonadotropin (FSH and hCG) therapies: In males planning to have children.

• Assisted reproductive techniques: Intrauterine insemination (IUI), Intracytoplasmic sperm injection (ICSI)

Solution to Question 7:

All of the following are theories of causation of preeclampsia except option A. There is reduced or absent prostacyclin(PGI2) production with increased production of thromboxane (TXA2) and reduced ratio of prostacyclin: thromboxane in preeclampsia.

The development of pre-eclampsia may be explained by:

• Failure of 2nd wave of trophoblastic invasion into uterine vessels, leading to decreased placental perfusion.

• Maternal inflammatory response directed against foreign fetal antigens, leading to immunological maladaptation.

• Maternal maladaptation to cardiovascular and inflammatory changes of normal pregnancy.

• Genetic factors and epigenetic influences.

The basic pathology is endothelial dysfunction and vascular spasm that affects vessels supplying all organs including the brain, kidney, uterus, etc. This ischemia leads to necrosis, hemorrhage, and end-organ damage of these organs.

Changes leading to vasospasm and endothelial dysfunction include:

• Imbalance of prostaglandins (Reduced vasodilatory PGI2, increased vasoconstrictor TXA2).

• Reduced angiotensinase activity and increased sensitivity of vessels to angiotensin II, which is a vasopressor.

• Reduced production of nitric oxide due to endothelial dysfunction. It normally acts as a vasodilator and inhibits platelet aggregation and thrombus formation.

• Increased production of endothelin 1 by endothelial cells. It is a potent vasoconstrictor.

• Increased inflammatory mediators such as tumor necrosis factor (TNF) and interleukins (IL-6) leading to endothelial injury.

• Increased lipid peroxides, reactive oxygen species (ROS) and superoxide anion radicals, leading to endothelial damage.

• Increased production of anti-angiogenic factors as compared to angiogenic factors. Two main antiangiogenic factors overproduced by the trophoblasts of women with preeclampsia are soluble fms-like tyrosine kinase I (SFlt-1) and soluble endoglin. Increased sFlt-1 inactivates and reduces the levels of free placental growth factor (PlGF) and VEGF concentrations.

All these processes lead to endothelial dysfunction and causes the clinical presentation of preeclampsia.

Solution to Question 8:

Except for echocardiography for cardiac disease, all other tests (urine analysis for bacteriuria, blood sugar levels for GDM, VDRL for syphilis) are cost-effective investigations routinely recommended in the screening of antenatal mothers.

Echocardiography is an expensive investigation and is not routinely recommended to look for cardiac abnormalities in all pregnant ladies. Indications for fetal echocardiography include:

• Abnormality in routine genetic screening such as USG to look for nuchal translucency (NT). NT is the space between the fetal skin and the underlying soft tissue near the neck of the fetus filled with fluid. NT of >3mm is abnormal and warrants the need for detailed USG and fetal echocardiography

• Gestational diabetes mellitus for early detection of fetal anomalies. Done at 20-22 weeks.

• Heart disease in pregnancy like congenital heart disease or Marfan's syndrome needs routine echocardiography (color doppler) at mid-pregnancy to look for structural abnormalities, valve anatomy, etc.

Routine investigations during pregnancy:

• 1st trimester:

• Hemoglobin, hematocrit, blood grouping, blood glucose, and VDRL.

• Midstream urine to look for protein, sugar, pus cells, and evidence of urinary tract infections or bacteriuria by dipstick test (for nitrites).

• Serology for rubella, hepatitis B virus, and HIV.

• Genetic screening like maternal serum alpha-fetoprotein.

• Ultrasound scan to look for gross fetal abnormalities, number of fetuses, and dating of pregnancy.

• 2nd trimester:

• Ultrasound scan (booking scan at 18-20 weeks) is done to look for cardiac activity, structural abnormalities, amniotic fluid index and placental location.

• Triple test at 11-13 weeks for a high risk mother.

• Oral glucose tolerance test (OGTT) at 24-26 weeks with a 1-hour 50-gram glucose load or a diagnostic 2-hour 75-gram glucose load. This is to look for gestational diabetes mellitus.

• 3rd trimester:

• Repeat hemoglobin, hematocrit, and serology for syphilis and HIV.

• Vaginal examination to look for Streptococcus agalactiae.

Solution to Question 9:

The cerclage procedure performed on nonpregnant females through a transabdominal approach is laparoscopic cerclage. Abdominal cerclage is conducted through either laparotomy or laparoscopy and is recommended in cases where the cervix is hypoplastic or in instances of prior failed vaginal cerclage.

Cerclage is usually done for cervical incompetence, defined as the inability to maintain pregnancy till term due to a functional or structural defect of the cervix. Cervical incompetence is diagnosed by clinical examination and ultrasonography showing the presence of short cervix (<25 mm) with or without funneling of internal os.

Prophylactic cerclage is done around 14 weeks of pregnancy or 2 weeks earlier than the previous fetal loss. Rescue or emergency cerclage is done anytime within 24 weeks of gestation. It should only be done when there is bulging of membranes through the dilated cervix.

Commonly used types of cervical cerclage during pregnancy:

• McDonald's procedure - non-absorbable (mersilene) material placed as a purse-string suture at the level of the internal os

• Modified Shirodkar's procedure - non-absorbable (mersilene) tape passed subcutaneously through a transverse anterior incision and tied posteriorly

Removal of cerclage should be done by 37 weeks of gestation. It can be removed earlier in case of preterm labor or if signs of abortion are seen.

Other options:

Option A: Lash and Lash cerclage is the first described treatment for cervical incompetence, involving the placement of stitches in the cervix through the transvaginal approach. It is now replaced by procedures such as McDonald's and Shirodkar's.

Option B: Benson und Durfee cerclage is the technique of placement of cervical cerclage through a transabdominal approach by a laparotomy. It is preferred in pregnant females when the vaginal approach is either not possible or very difficult.

Option C: Wurms cerclage is a type of cerclage used in pregnancy where monofilament material (Prolene) is used to place two higher circumferential sutures through the cervix without the need for bladder dissection.

Solution to Question 10:

Urinary tract infection (UTI) in a pregnant lady in her 2nd trimester is best defined as the growth of 105 coliforms in a voided urine sample. Urine culture is the standard method to evaluate UTI pregnancy.

UTI in pregnancy is a bacterial infection caused by organisms in the normal perineal flora. It can vary from asymptomatic bacteriuria to symptomatic cystitis or pyelonephritis.

Asymptomatic bacteriuria is characterized by persistent and actively multiplying bacteria in the urinary tract without any symptoms. A positive culture is demonstrated during the antenatal visit. It can complicate into a symptomatic infection when untreated.

Symptomatic infections include

• Cystitis is characterized by dysuria, urgency and frequency along with pyuria and bacteriuria. Microscopic hematuria is also seen

• Pyelonephritis is common in the second trimester, nulliparous and young women. It is usually unilateral with fever, chills, and ache in the lumbar region. Nausea, vomiting, and costovertebral tenderness may be seen. The organisms causing pyelonephritis are E. coli >> Streptococcus, S.aureus > Enterobacter, Klebsiella

UTI is evaluated by blood studies including complete blood count, serum creatinine, blood urea nitrogen, urine studies and imaging. Urine studies mainly include a urine culture, urine analysis, and dipstick.

Urine culture is the standard method of evaluating UTIs in pregnancy. Positive urine culture shows a bacterial growth of >100,000 colony-forming units (CFUs) per milliliter in a voided urine sample. The presence of white blood cell (WBC) casts is suggestive of pyelonephritis. No other screening methods are as reliable as a urine culture.

Urine analysis showing positive nitrites, leukocyte esterase, WBCs (pyuria), red blood cells (RBCs), and protein are suggestive of UTI. Pyuria is seen in cystitis. Dipstick for nitrite and leukocyte esterase is used to evaluate asymptomatic bacteriuria.

Management of UTIs is as follows

• Asymptomatic bacteriuria is mainly treated with oral antibiotics such as amoxicillin, ampicillin, nitrofurantoin, and trimethoprim-sulfamethoxazole.

• Pyelonephritis is managed with intravenous hydration, and antibiotics such as ampicillin+gentamycin, cefazolin or ceftriaxone.

Solution to Question 11:

The incorrect regarding innervation of the uterus is statement B. Uterine contractility is mainly mediated by hormonal mechanisms. Estrogen, prostaglandin, and oxytocin produce contraction, while progesterone causes relaxation.

Uterine sensory supply is by ascending afferent fibers which pass through the inferior hypogastric plexus and enter the spinal cord through T10-T12 and L1 roots.

Labor pain is due to stimulation of nociceptors in the genital tract caused by ischemia.

• First stage labor: Pain is mediated by T10 to L1 spinal segments. It is caused by distension of the cervix and low uterine segments along with isometric uterine contraction.

• Second stage labor: Pain is carried by T12 to L1 and S2 to S4 spinal segments. It is caused by tissue damage in the pelvis and perineum

Analgesia in labor is achieved by intramuscular injection of pethidine 50 mg. Metoclopramide is added to prevent vomiting. Pethidine is avoided within 2 hours of delivery as it is a known respiratory depressant to the fetus. When relief of pain is required throughout labor, regional analgesia is preferred through pudendal nerve block, epidural analgesia, spinal anesthesia or combined epidural-spinal anesthesia.

Solution to Question 12:

Seminal vesicles and vas deferens would be bilaterally absent, congenitally in cystic fibrosis. This is known as the congenital bilateral absence of vas deference (CBAVD).

Cystic fibrosis is an autosomal recessive disorder that affects fluid secretion in the exocrine glands and in the epithelial lining. It is caused primarily due to a mutation in the CFTR gene, which codes for an anion channel in the apical membranes of epithelial cells. This leads to abnormally viscous secretions that obstruct organ passages, resulting in recurrent infections, pancreatic insufficiency, steatorrhea, malnutrition, hepatic cirrhosis, intestinal obstruction, and male infertility.

Infertility is due to azoospermia and CBAVD. In some patients, the bilateral absence of the vas deferens may be the only feature suggesting an underlying CFTR mutation.

On examination, a normal reproductive tract (including testis, and epididymis) is seen, with bilateral absence of vas deferens and atrophic or absent seminal vesicles. Semen analysis shows a low-volume (0.5 mL), acidic ejaculate that is devoid of fructose and seminal vesicle fluid. There may be associated renal agenesis in some patients.

Other options:

Option B: Klinefelter syndrome is a genetic disorder that results from a 47, XXY karyotype in phenotypically male patients. They present with gynecomastia, azoospermia, and infertility due

to hyalinization of the seminiferous tubules and a small testis.

Option C: Kallmann syndrome is associated with a low or absent GnRH that is characterized by anosmia, hypogonadism, infertility, cryptorchidism, delayed pubertal maturation, and neurological abnormalities.

Option D: Kartagener syndrome is an autosomal recessive disorder affecting ciliary motility; it is characterized by situs inversus, chronic sinusitis, and bronchiectasis. Male infertility may be present and is due to asthenospermia (immotile spermatozoa due to impaired sperm flagella function).

Solution to Question 13:

The graph representing normal response to oral glucose tolerance test (OGTT) is graph D.

Oral glucose tolerance test is routinely used to diagnose gestational diabetes mellitus. It is performed by giving a bolus of glucose (75g) to the patient, and measuring blood glucose at intervals of time to assess the tolerance and response of the islets of pancreas to glucose. 5 measurements of glucose is noted in blood and urine at half hour intervals.

Indications:

• Patients with symptoms suggestive of diabetes mellitus, but with fasting blood glucose value being borderline (between 100-126 mg/dl).

• Excessive weight gain during pregnancy, a past history of big baby (more than 4 kg) or a past history of miscarriage.

• To rule out benign renal glucosuria.

Glucose Tolerance Curve (GTC) is the graph plotted by taking plasma glucose measurements on Y-axis and plotting against time, on the X-axis.



Graph D: Normal: This is the normal response seen in a healthy individual. It is characterized by a peak glucose value reached within 1 hour, following which glucose levels reach the base value. The

levels don't cross the renal threshold at any point of time.

Graph A and Graph B: Diabetes mellitus: The graph will peak after 1-1.5 hours and does not return to fasting level within 2.5 hours. The highest value exceeds the renal threshold.

Depending on severity, the graph can show mild, moderate, and severe diabetic curve.

Graph C: Lag curve: The fasting blood glucose is normal here, but the curve rapidly rises in 1/2 to 1 hour and exceeds the renal threshold. There is a rapid and complete return to normal value in a few minutes. This is seen in hyperthyroidism, early diabetes, and sometimes during pregnancy.

Solution to Question 14:

Kassowitz rule is related to congenital syphilis.

According to Kassowitz rule, if a woman with untreated syphilis has a series of pregnancies, the likelihood of infection of the fetus in later pregnancies becomes less. Usually, only the first pregnancy is an abortion, the following pregnancy losses will be stillbirths, and later pregnancies will have alive babies with features of congenital syphilis.

Maternal syphilis can cause fetal infection (congenital syphilis) by several routes, the most common being transplacental transmission, followed by transmission during delivery or across the placental membranes. Early maternal syphilis has a higher transmission as compared to late syphilis.

Up to 70 percent of women with maternal syphilis have an adverse outcome. Most common complications include preterm labour, stillbirth, fetal-growth restriction and fetal death. Newborns can have features of infection such as jaundice, ascites, petechiae, purpura, pneumonia, myocarditis and long bone involvement.

Maternal syphilis in all trimesters should be treated with parenteral penicillin G in order to prevent the spread of infection to fetus. The treatment of choice for congenital syphilis is penicillin.

Solution to Question 15:

The modified biophysical profile includes NST (nonstress test) and AFI (amniotic fluid index).

It is a modification of the biophysical profile (BPP) test, including only 2 parameters so that it can be performed more routinely (twice a week) for surveillance and takes less time (10 minutes). Modified BPP is considered nonreassuring when the NST is non-reactive and/ or the AFI is < 5.

The nonstress test is a screening test for fetal wellness. It describes fetal heart rate acceleration (measured by a Doppler) in response to fetal movements (perceived by mother). An acoustic stimulation nonstress test is a type of NST in which a loud external sound is used to startle the fetus and provoke heart rate accelerations.

• Reactive (Reassuring) NST - Two or more accelerations of 15 seconds duration, 15 beats per minute amplitude, in a 20-minute testing time.

• Non-reactive (Non-reassuring) NST - Absence of any fetal reactivity.

BPP is a multi-parametric, screening test to assess uteroplacental insufficiency. It consists of five components/ variables and the observations are made in a 30-minute time period.

Components of BPP:

(Mnemonic: BATMaN)

B - Breathing movements of the fetus

A - Amniotic fluid volume

T - Tone of the fetus

M - Movements of the fetus

N - Non-stress test (NST)

Each parameter has a maximum score of 2 and a minimum score of 0. So the total maximum score will be 10. A score of 4 or less indicates fetal acidemia.

Interpretation and management of BPP scores:

BPP is directly related to fetal pH and fetal outcome. A lower BPP score is associated with a poor outcome (stillbirth, perinatal mortality) and lower APGAR scores at birth.

A score of <10 is associated with a pH of <7.2 and a score of 0 is associated with 100% fetal acidemia.

Note: The modified biophysical score has only 2 components i.e. NST and amniotic fluid index only. It is considered abnormal (nonreassuring) when the NST is non-reactive and/or the AFI is <5.

Parameters	Minimal Normal Criteria
Fetal breathing movements	≥ 1 episode lasting > 30 seco nds
Amniotic fluid volume (AF V)	≥ 1 pocket measuring 2 cm in two perpendicular planes (2 ×2 cm pocket)
Fetal muscle tone	≥ 1 episode of active limb or t runk extension with the retur n of flexion; Eg. opening and closing of the hand, consider ed normal
Gross body movements	≥ 3 discrete body/limb move ments
Non-stress Test (NST)	Reactive pattern

BPP score	Interpretation	Management
10	Normal, nonasphyxiated fetu s	Repeat test weekly (or biweek ly in diabetic mother or post-term pregnancy)

BPP score	Interpretation	Management
8with normal AFV	Normal, nonasphyxiated fetus	Repeat test weekly (or biweekly in diabetic mother or post-term pregnancy)
8 with decreased AFV	Suspicion of chronic fetal asphyxia	Delivery
6	Possible fetal asphyxia	Deliver, if AFV is abnormal or >36weeks or if the repeat test score is ≤ 6. If the repeat score >6, observation and repeat test weekly
4	Probablefetal asphyxia	Repeat testing same dayIf the repeat score ≤6, delivery
0-2	Almostcertainfetal asphyxia	Delivery

Solution to Question 16:

All of the following statements about anti-Mullerian hormones (AMH) are true, except option B. AMH is a glycopeptide secreted by the granulosa cells of the pre-antral follicles and small antral follicles of the ovary.

In males, AMH inhibits the development of the Mullerian system and its absence can result in hermaphroditism. In the female, it helps in follicular development and oocyte maturation. In the absence of AMH, growing follicles are more responsive to FSH. This is because AMH reduces the sensitivity of pre-antral follicles to FSH (option D) thereby regulating the follicular growth.

Serum AMH predicts ovarian reserve in a woman with infertility or secondary amenorrhea. Also, AMH values decrease with age and can indicate menopause, owing to the reduced ovarian reserve. This makes AMH the best predictor for measuring ovarian reserve. However, recent studies have shown intercycle variations of AMH levels, hence it is recommended to measure AMH along with antral follicle count (on USG) on days 1-3 of the cycle (option A).

It is also a useful predictor of pregnancy rate in infertile women undergoing IVF (In-vitro fertilization) treatment (option C). It helps in assessing the ovarian response to exogenous gonadotropin stimulation, blastocyst formation, and miscarriage rates.

Values and interpretation:

• Normal value: 2–6.8 ng/mL

• Poor ovarian reserve: <1 ng/mL

• PCOD and hyperstimulation syndrome: >10 ng/mL

Solution to Question 17:

The false statement regarding PPH is option A. WHO recommends intravenous isotonic crystalloids over intravenous colloids for fluid replacement in PPH. This is because colloids are

not found to be useful, and some colloids (hydroxyethyl starch) can even be harmful.

Postpartum hemorrhage is defined as a loss of more than 500 ml of blood after completion of the third stage of labor (in vaginal delivery).

PPH is blood loss of more than 1000 mL from the genital tract following cesarean section.

Depending upon the amount of blood loss, PPH can be, minor (<1L) or major (>1L). Major is further subdivided into moderate (1L-2L) and severe (>2L).

Types of PPH:

Primary/early PPH: Hemorrhage occurring within 24 hours of childbirth. Causes include

• Atony: Primi, grand multipara, over distended uterus, hydramnios, precipitate labor, prolonged labor, uterine malformations, fibroid uterus

• Trauma: Instrumental deliveries, episiotomy, precipitate labor, macrosomia

• Retained tissues: retained placenta, blood clots

• Bleeding disorders: Sepsis, Intrauterine death, HELLP syndrome.

Secondary/ late PPH: Hemorrhage occurring after 24 hours and up to 12 weeks postpartum. Causes include

• Placental polyp: Retained products undergo necrosis with fibrin deposition and eventually form a placental polyp, which on detachment from the myometrium causes brisk hemorrhage.

• Abnormal involution of the placental site.

• Retention of placental cotyledon or products of conception.

• Uterine artery pseudoaneurysm.

• Endometritis

• Coagulopathy

Resuscitative measures include

• Airway, breathing, circulation assessment

• A high concentration of oxygen (10–15 l/min) via a facemask

• Blood transfusion if clinically required

• Fluid resuscitation with warmed fluids up to 3.5 liters, starting with up to 2 liters of crystalloids such as normal saline and ringer's lactate. Colloids can be used when there is a need for packed RBCs transfusion, until blood is arranged.

Solution to Question 18:

The recommended HPV vaccination schedule for 9 to 14-year-olds according to WHO SAGE guidelines is 1 or 2 doses.

The recommendations for different age groups are as follows:

• 1 or 2 dose schedule for girls aged 9-14 years.

• 1 or 2 dose schedule for young women aged 15-20 years.

• 2 doses with a 6-month interval for women older than 21 years.

• 3 doses (or atleast 2 doses) for immunocompromised individuals.

As per WHO SAGE (Strategic Advisory Group of Experts on Immunization), a single-dose HPV vaccination schedule can provide comparable efficacy and durability of protection to a two-dose regimen. This can help in wider coverage of vaccination of all girls in low-middle income countries. Therefore, a 1 or 2-dose schedule is recommended for females aged between 9-20 years.

The available vaccinations for HPV are:

• Gardasil 9 (HPV 9) is a nonavalent vaccine active against HPV 6,11,16,18,31,33,45,52,58.

• Gardasil 4 (HPV 4) is a quadrivalent vaccine against 6,11,16,18. This is now replaced by HPV 9.

• Cervarix (HPV 2) is a bivalent vaccine active against 16, 18.

Solution to Question 19:

All the above coagulation factors (Factors ll, Vll, Vlll) are increased in pregnancy except factor Xl.

Pregnancy is a hypercoagulable state. This controls bleeding and helps in hemostasis after placental separation.

Hematological changes in pregnancy:

• All clotting factors are increased, except factors 11 and 13. The levels return to normal within 2 weeks of delivery.

• Anticoagulants protein C, protein S, and antithrombin are decreased during pregnancy.

• Fibrinogen levels are increased by 50%.

• Clotting time and bleeding time remain unaffected.

• The viscosity of blood decreases due to relative hemodilution (relatively higher increase in plasma volume as compared to the increase in RBC mass).

• ESR is raised due to an increase in fibrinogen and globulin levels.

• Because of a marked fall in albumin level and only a slight rise of globulin, the normal albumin:globulin ratio of 1.7:1 is decreased to 1:1.

• Platelet counts remain the same or fall by a small amount (15%).

The most common cause of decreased platelet count in pregnancy is benign gestational thrombocytopenia. This is due to hemodilution and increased platelet consumption and the women are generally asymptomatic.

Solution to Question 20:

The incorrect statement regarding the management of acute pyelonephritis in pregnancy is option D. The patient can be discharged once afebrile for 24 hours, and antibiotics have to be continued

for 7 to 10 days.

Acute Pyelonephritis is one of the frequent and serious medical complications occurring in 2nd-trimester of pregnancy. The most common causative organism is E.coli.

Clinical features include sudden onset fever, chills, and aching pain in the lumbar region (usually unilateral). It is associated with nausea, vomiting, and anorexia. Tenderness in the costovertebral angle can be elicited on examination.

Diagnosis is confirmed by the collection of urine (preferably from catheterization) and analysis to look for leukocytes and bacteria.

Management protocol for pyelonephritis during pregnancy is

• Hospitalization is recommended for all patients. Very rare cases of uncomplicated pyelonephritis can be managed on an outpatient basis

• Urine and blood (if temperature is > 39 degrees C) samples are collected for culture.

• Aggressive intravenous hydration to maintain adequate urine output is the mainstay of treatment. A urinary output ≥50 mL/hr is maintained with intravenous crystalloids.

• Vital signs such as blood pressure, pulse, temperature, and oxygen saturation along with urine output should be monitored for signs of worsening sepsis.

• Empirical intravenous antibiotic therapy is started with ampicillin plus gentamicin, cefazolin or ceftriaxone. It is switched to oral antibiotics once the patient is afebrile.

• In case of persistent infection or lack of clinical improvement by 48 to 72 hours, renal sonography is recommended to look for urinary tract abnormalities (like obstruction or calculi).

• The patient is discharged when afebrile for 24 hours and antimicrobial therapy is continued for 7 to 10 days.

• Urine culture is repeated 1–2 weeks after antimicrobial therapy is completed.

Solution to Question 21:

All of the following options are the diagnostic criteria for preterm labor except an posterior position of cervix (Option D). An anterior cervix favors a higher bishop's score, which means a higher chance of preterm labor as compared to the posterior cervix. The midline position of the cervix has also shown higher incidence of preterm labor as compared to posterior cervix.

Preterm labor (PTL) is defined as labor occuring between 20weeks to 37th completed week of gestation.

Diagnostic criteria of preterm labour include:

• Regular uterine contractions at least one in every 10 minutes, or 4 contractions in 20 minutes or 8 contractions in 60 min with/without pain.

• Cervix dilatation ≥2 cm

• Cervix effacement ≥80%

• Length of cervix (on TVS) ≤2.5 cm and funneling of the internal os >1 cm

• Pelvic pressure, backache, vaginal discharge, or bleeding PV

Predictors of preterm labor:

• Clinical predictors:

• History of prior preterm birth

• Multiple pregnancies

• Active genital tract infection

• Biophysical predictors:

• Uterine contractions ≥4/hour

• Bishop score ≥4

• Cervical length (on TVS) ≤25 mm

• Biochemical predictors:

• Fetal fibronectin (fFN) positive (>50 ng/mL) in cervico-vaginal discharge

• Increased levels of IL-6, IL-8, TNF-alpha

Solution to Question 22:

The false statement about late decelerations among the given options, is option C- at least 10 beats, 2 times in a 20 min period above baseline.

Typically, a late deceleration causes a fall in the fetal heart rate (FHR) by a depth of <10 to 20 bpm below the baseline. It is usually not accompanied by accelerations (Option D).

Late decelerations are smooth, gradual, (Option B) symmetrical decline in the FHR that begins at or after the peak of uterine contraction and ends only after the contraction has ended (Option A).

It is usually indicative of uteroplacental insufficiency and fetal hypoxia. Other causes of late decelerations include hypotension from epidural anesthesia, excessive use of oxytocin, maternal hypertension, diabetes, and placental abruption.

A late deceleration plus decreased variability in FHR is predictive of fetal acidemia and neonatal morbidity.

The given CTG shows late decelerations and it's relation with the uterine contraction.

The below CTG shows early decelerations. In early decelerations, the onset, it's nadir and the recovery of deceleration coincide with the beginning, peak, and ending of uterine contraction respectively.

Early decelerations are seen in fetal head compression. They typically cause a fall of ≤ 20 to 30 bpm below the baseline. These early decelerations are common during active labour and are benign in nature.



A variable deceleraton is a drop in the FHR that begins with the uterine contactions onset and reaches a nadir in <30 seconds.Its depth is ≥15 bpm in and the deceleration lasts between 15 seconds to 2 minutes.

The slope of FHR change is erratic, giving the wave form its jagged shape. A variable deceleration is indicative of umbilical cord occlusion.

The below CTG depicts variable deceleration.

Solution to Question 23:

The age group distribution for vulval cancer is 55 to 85 years (Option C).

Vulval cancer is a malignancy of elderly women, usually in their 6th and 7th decades of life. It can also be seen in younger women who have been diagnosed with sexually transmitted infections such as HIV and HPV infections. The average age of diagnosis is 65 years.

Persistent HPV, HSV infections, smoking, chronic immunosuppression and chronic inflammation such as lichen sclerosis are risk factors. Vulvar intraepithelial neoplasia (VIN) is the preinvasive condition of the vulva.

Squamous cell carcinoma is the most common type of vulvar cancer. The most common site of vulvar cancer is labia majora > Clitoris > Labia Minora.

Vulval cancer generally presents with vulvar pruritus, vulvar swelling, a lump or an ulcer. Pain is usually a late feature of the disease.

The lump is seen as a pigmented papular raised area and the ulcers often have everted margins and usually bleed and cause an offensive vulval discharge.

The most common type of spread in vulval cancer is the lymphatic route. The first lymph node to be involved is the superficial inguinal lymph node and hence, the sentinel lymph node biopsy is helpful for evaluating the spread of vulval cancer and helps attain an adequate surgical procedure with a good prognosis.

Treatment of vulval cancer generally involves simple or radical, partial or total vulvectomy for the management of the primary tumour and inguinofemoral lymph node dissection for the involvement of the lymph nodes. This is decided based on the staging of the disease.

Solution to Question 24:

Among the given options, Extraplacental membranes and 2-vesseled umbilical cord present (Option C) is the wrong statement.

At term, the normal placenta weighs 470 g (Option A), is round to oval with a 22-cm diameter (Option B) and has a central thickness of ≥2.5 cm (Option D). The placenta is composed of a placental disc, extraplacental membranes, and a three-vessel umbilical cord. Hence, option C is the wrong statement.

The normal human placenta is discoid, deciduate, and hemochorial.

The term deciduate means that the placenta is shed at the time of parturition. Hemochorial means that the maternal blood directly comes in contact with the fetal trophoblasts.

The placenta consists of two plates. The chorionic plate lies in the fetal surface. It is lined by the amniotic membrane and the umbilical cord is attached to this plate.

The basal plate is the maternal aspect and lies against the uterine wall. It is divided by clefts into portions termed as cotyledons.

Solution to Question 25:

The correct sequence of dilation and curettage (D&C) is 4-3-1-2.

4 - Assessment of uterine size and direction

3 - Uterine sound

1 - Serial dilation

2 - Curettage

Indications:

• Infertility

• Abnormal uterine bleeding (AUB)

• Amenorrhea

• Endometrial tuberculosis

• Endometrial carcinoma

• Postmenopausal bleeding

• Endometrial polyp

• Removal of IUD

• Incomplete abortion

• Molar pregnancy

Absolute contraindication: Viable intrauterine pregnancy.

Steps:

• The patient's bladder is emptied prior to the procedure.

• The operation is performed under general anesthesia or under diazepam sedation with or without a paracervical block.

• With the patient in a lithotomy position, local antiseptic cleaning and draping are carried out.

• A bimanual examination is conducted.

• A posterior vaginal speculum is introduced.

• The anterior lip of the cervix is grasped with an Allis tissue forceps.

• A uterine sound is introduced to confirm the position and note the length of the uterocervical canal.

• The cervical canal is dilated with graduated dilators.

• When the dilator is introduced, the cervix is made steady by traction of the vulsellum (Allis's tissue forceps).

• After the desired dilatation, the uterine cavity is curetted by a uterine curette either in a clockwise or anticlockwise direction starting from the fundus down to the internal os.

• The vulsellum and the speculum are removed.

• The curetted material is preserved in 10% formol-saline, properly labeled, and sent for histological examination.

Complications: Infection, hemorrhage, cervical lacerations, uterine perforation, and post-operative uterine adhesions.



Solution to Question 26:

Female sterilization is an irreversible method of contraception.

It involves surgery and severing, destruction, or ligation of fallopian tubes through laparotomy or laparoscopy. Hence it is normally irreversible.

All other methods are reversible long-term methods of contraception.

Other options:

Option A: Progestin injections, such as NET-EN, are used both as a form of progestogen-only injectable birth control and as components of combined injectable birth control formulations. They are reversible.

Option C: Progestin implants, such as Nexplanon, are used for contraception; they are highly effective, safe, reversible, and the most convenient choice for many women.

Option D: IUCDs are a type of long-term, reversible contraception. They are T-shaped devices that are inserted into the uterine cavity. These can either be
copper-containing like CuT-380A and CuT-200, or hormone-containing IUCDs like mirena.

Solution to Question 27:

The WHO recommends a minimum of 8 antenatal visits to have a positive pregnancy outcome.

According to the latest WHO guidelines the minimum number of antenatal visits has been increased from 4 to 8. This increased number of visits significantly reduces the risk of perinatal deaths and enhances maternal satisfaction.

The ideal number of antenatal visits is 12-15:

• Monthly visits up to 28 weeks

• Two weekly visits between 28-36 weeks

• Weekly visits from 36 weeks onwards

WHO ANC model	
First trimester	Contact 1: up to 12 weeks
Second trimester	Contact 2: 20 weeksContact 3: 26 weeks
Third trimester	Contact 4: 30 weeksContact 5: 34 weeksContact 6: 36 weeksContact 7: 38 weeksContact 8: 40 weeks
Return for delivery at 41 weeks if not given birth.	

Solution to Question 28:

Gestational age or period of gestation (POG) refers to the duration of pregnancy calculated from the first day of the last menstrual period (LMP). It is usually expressed in weeks.

According to Naegele's rule, the estimated due date (EDD) is obtained by adding 7 days and subtracting 3 months from the first day of the last menstrual period (LMP). Another method

involves adding 9 months and 7 days to the LMP. Less than 5% of births follow Naegele's rule. It is based on a 28-day regular cycle.

If the cycle is shorter/longer than 28 days, corrected EDD is calculated as follows:

• For a 35-day cycle, add another 7 (35-28=7) days to the calculated EDD to get the corrected EDD.

• For a 25-day cycle, subtract 3 (28-25=3) days from the calculated EDD to get the corrected EDD.

Other options:

Option A and D: The best method of assessing the gestational age is by measuring the crown-rump length (CRL) using ultrasonography between 7 to 12 weeks of gestation. It is measured from the vertex to the coccyx. CRL (in mm) + 6.5 = gestational age in weeks.

Option C: In an IVF pregnancy, the day of embryo transfer/day of fertilization is needed to calculate POG.

Solution to Question 29:

Extra-amniotic instillation of ethacridine lactate cannot be used in the first trimester. It is generally safe to use in the second trimester.

Intrauterine instillation of hyperosmotic solutions is usually employed in the second trimester:

• Extra-amniotic ethacridine lactate, prostaglandins (PGE2, PGF2α)

• Extra-amniotic saline infusion (isotonic) with a transcervical catheter balloon

• Intra-amniotic hypertonic urea (30%), saline (20%)

The various methods of termination of pregnancy are:

Timing	Medical methods	Surgical methods
1st trimester	MifepristoneMifepristone and misoprostolMethotrexate and misoprostolTamoxifen and misoprostol	Menstrual regulationVacuum aspiration (MVA/EVA)Suction evacuation and/or curettageDilatation and evacuation
2nd trimester	MisoprostolMifepristone and misoprostolDinoprostoneCarboprost	Dilatation and evacuationIntrauterine instillation of hyperosmotic solutionsOxytocin infusionHysterotomy

Solution to Question 30:

Loss of knee jerk reflex is the first sign of $MgSO_4$ toxicity.

$MgSO_4$ is used as an anticonvulsant in the management of eclampsia and for prophylaxis in severe preeclampsia. The therapeutic level of magnesium in eclampsia is 4–7 mEq/L. Measurement of serum magnesium levels is indicated to monitor magnesium toxicity if

serum creatinine >1.0 mg/dL.

Loss of knee jerk reflex is the first sign of $MgSO_4$ toxicity and occurs when the plasma magnesium level reaches 10 mEq/L.

Other signs suggestive of magnesium toxicity:

• Decreased respiratory rate <16/min

• Urine output <30 mL/h

• Chest pain and heart block

• Visual changes, flushing, muscle paralysis, and drowsiness

At > 10 mEq/L, breathing weakens. At ≥ 12 mEq/L respiratory paralysis and respiratory arrest follow. At > 25 mEq/L cardiac arrest occurs.

Magnesium therapy should be stopped and a slow infusion of IV calcium gluconate 10 mL (10% solution) is given. Fluid loading and forced diuresis with estimation of serum magnesium and creatinine levels are done.

Repeat injections of magnesium sulfate are continued only if the patellar reflex is present, respiratory rate is >12, and urine output is >30 mL/h.

Solution to Question 31:

A 14-year-old girl with Tanner stage 1 breasts would be classified as having primary amenorrhea.

Primary amenorrhea is defined as the absence of menstruation by 14 years without the development of secondary sexual characteristics or by 16 years regardless of the presence of secondary sexual characteristics.

Tanner stage 1 breasts correspond to prepubertal breasts. Breast development beyond Tanner stage 2 in girls indicates the onset of puberty.



Solution to Question 32:

The daughter of a woman who received diethylstilbestrol (DES) during her pregnancy is at a high risk of clear cell carcinoma of the vagina. Clear cell carcinoma is a type of adenocarcinoma and hence is the most specific answer.

If the female fetus is exposed to diethylstilbestrol (DES) taken by the mother during pregnancy, columnar epithelium appears in the upper two-thirds of vaginal mucosa, which can develop into vaginal adenosis and clear cell adenocarcinoma during adolescence.

Other options:

Option A: Squamous cell carcinoma arises from non-keratinized stratified epithelium and is commonly associated with human papilloma virus exposure.

Option C: Adenocarcinoma usually arise from cells of glandular tissue. It can be seen in an endometriosis foci, areas of vaginal adenosis, periurethral glands and remnants of the wolffian ducts. Vaginal adenocarcinomas occur as metastatic cancers in most patients and are rarely primary cancers. Metastases may be spread from the endometrium, breast, pancreas, kidneys and colon.

Option D: Sarcoma botryoides is the most common type of embryonal rhabdomyosarcomas in infants and children.

Solution to Question 33:

Daily fetal movement count is not a component of biophysical profile (BPP). Components of BPP are non-stress test (NST), fetal breathing movements, gross body movements, fetal muscle tone, and amniotic fluid volume.

BPP is a multi-parametric screening test to assess uteroplacental insufficiency. The observations are made in a 30-minute time period.

Each parameter has a maximum score of 2 and a minimum score of 0. So the total maximum score will be 10. A score of 4 or less indicates fetal acidemia.

Interpretation and management of BPP scores:

BPP is directly related to fetal pH and fetal outcome. A lower BPP score is associated with a poor outcome (stillbirth, perinatal mortality) and lower APGAR scores at birth.

The modified biophysical score has only 2 components: NST and amniotic fluid index.

BPP score	Interpretation	Management
10	Normal, nonasphyxiated fetus	Repeat test weekly (or biweekly in diabetic mothers or post-term pregnancy)

BPP score	Interpretation	Management
8with normal AFV	Normal, nonasphyxiated fetus	Repeat test weekly (or biweekly in diabetic mothers or post-term pregnancy)
8 with decreased AFV	Suspicion of chronic fetal asphyxia	Delivery
6	Possible fetal asphyxia	Deliver, if AFV is abnormal or > 36 weeks or if the repeat test score is ≤ 6. If the repeat score >6, observation and repeat test weekly
4	Probablefetal asphyxia	Repeat testing same dayIf the repeat score ≤6, delivery
0-2	Almostcertainfetal asphyxia	Delivery

Solution to Question 34:

Multiparity is not a cause of uterine dysfunction.

Adequate uterine contractions are required to dilate the cervix and expel the fetus. Hypotonic uterine dysfunction is a defect in the propulsive power of the uterus that results in prolonged labor. It is a common indication for cesarean section.

Causes of hypotonic uterine dysfunction:

• Neuraxial analgesia

• Chorioamnionitis

• Nulliparity

•

Mechanical disruption of myometrial function from leiomyoma or pelvic tumor

• Maternal obesity

• Uterine müllerian anomaly

• Increased maternal age

• Malposition and malpresentation of the fetus, with inadequate contact of the presenting part onto the lower uterine segment, as seen in the case of a high fetal head station

Solution to Question 35:

Human immunodeficiency virus is not a cause of nonimmune hydrops.

Hydrops is diagnosed by identifying two or more fetal effusions—pleural, pericardial, or ascites—or one effusion plus anasarca. Hydrops can be seen in association with red cell auto

immunization, which is termed immune hydrops. Otherwise, it is nonimmune hydrops.

Pathogenesis of hydrops includes decreased colloid oncotic pressure, increased hydrostatic or central venous pressure, and enhanced vascular permeability. Sonographically measured skin thickness of >5 mm constitutes edema or anasarca.

Etiology of nonimmune hydrops fetalis:

Cardiovascular

• Structural defects:

• Ebstein anomaly

• Tetralogy of Fallot with absent pulmonary valve

• Hypoplastic left or right heart

• Premature closure of ductus arteriosus

• Arteriovenous malformation (vein of Galen aneurysm)

• Cardiomyopathies

• Tachyarrhythmias

• Bradycardia

Chromosomal

• Turner syndrome (45,X), triploidy

• Trisomies 21, 18, and 13

Hematological

• Hemoglobinopathies, such as α4-thalassemia

• Decreased erythrocyte production (myeloproliferative disorders)

• Fetomaternal hemorrhage

Lymphatic Abnormalities

• Cystic hygroma

• Systemic lymphangiectasis

• Pulmonary lymphangiectasis

Infections

• Parvovirus B19

• Syphilis

• Cytomegalovirus

• Toxoplasmosis

• Rubella

• Enterovirus

• Varicella

• Herpes simplex

• Coxsackievirus

• Listeriosis

• Leptospirosis

• Chagas disease

• Lyme disease

Kidney and Urinary Tract

• Kidney malformations

• Bladder outlet obstructions

• Congenital (Finnish) nephrosis

• Bartter syndrome

Placental, Twin, and Cord Abnormalities

• Placental chorioangioma

• Twin-twin transfusion syndrome

• Twin reversed arterial perfusion sequence

• Twin anemia polycythemia sequence

• Cord vessel thrombosis

Idiopathic

Solution to Question 36:

The correct statement regarding the recurrence of ovarian cancer is option C: It is classified as platinum-sensitive disease if recurrence occurs > 6 months of primary therapy.

Gradual elevation of CA125 is considered the first sign of relapse of ovarian cancer. Tamoxifen is recommended for recurrent ovarian cancer if there is only biochemical evidence of the recurrence of ovarian cancer. Otherwise, conventional cytotoxic chemotherapy is administered.

Recurrence of ovarian cancer can be classified as platinum-refractory, platinum-sensitive, and platinum-resistant disease.

• Platinum-refractory disease includes women who progress during primary chemotherapy. It has a worse prognosis.

• Platinum-resistant disease includes women who relapse within 6 months of primary therapy. Palliative, single, non-platinum chemotherapy is recommended for platinum-refractory and platinum-resistant disease. Bevacizumab has been approved for the treatment of platinum-resistant disease. It can also be used in combination with weekly paclitaxel, doxorubicin, and topotecan.

• Platinum-sensitive disease includes women who relapse more than 6 months after primary therapy. These patients are treated with a platinum-based combination such as carboplatin and

paclitaxel or gemcitabine.

Solution to Question 37:

The given graph showing only the tracing for the fetal heart rate (FHR) is a non-stress test (NST).

The NST is a screening test for fetal wellness. It describes fetal heart rate acceleration (measured by a doppler) in response to fetal movements (perceived by the mother).

• Reactive (reassuring) NST - Two or more accelerations of 15 seconds duration and amplitude of 15 beats/min, in a 20-minute testing time

• Non-reactive (non-reassuring) NST - Absence of any fetal reactivity.

Other options:

Option B: Fetal tachycardia is defined as a baseline FHR above 160 beats/min, whereas this tracing shows FHR below 160 beats/min.

Option C: Beat-to-beat variability is the change in baseline fetal heart rate (excluding the accelerations and decelerations) in response to sympathetic and parasympathetic stimulation. A beat-to-beat variability of 6-25 beats/min is considered normal. This tracing shows normal beat-to-beat variability.

Option D: Cardiotocography (CTG) includes tracings for fetal heart rate and uterine contractions.

Solution to Question 38:

The abnormal placenta shown in the image depicts both succenturiate lobe and velamentous insertion of the cord.



The umbilical cord is usually attached at or near the center of the placenta.

In the velamentous placenta, the cord is attached to the membranes. The cord vessels traverse between the membranes for some distance before reaching the placenta. If the leash of blood vessels traverses through the membranes overlying the internal os, in front of the presenting part, the condition is called vasa previa. The rupture of the membranes involving the overlying vessels leads to vaginal bleeding and fetal exsanguination.

The below image shows the velamentous placenta:

The succenturiate placenta is characterized by the presence of one or more small lobes of placenta placed at varying distances from the main placental margins. A leash of blood vessels connects these accessory cotyledons to the main placenta. If these vessels are absent, then it is called placenta spuria. If the succenturiate lobe is retained postpartum, there can be primary or secondary postpartum hemorrhage, subinvolution of the uterus, uterine sepsis, and polyp formation.

The below image shows a succenturiate placenta. L1, L2 and L3 depict smaller lobes, UC is the umbilical cord, and M is the intervening membrane.



Solution to Question 39:

Human chorionic gonadotropin (hCG) is the hormone secreted by the placenta.

hCG is a glycoprotein, produced by the syncytiotrophoblast of the placenta and secreted into the blood of both mother and fetus. It can be detected in the maternal serum or urine as early as 8–9 days postfertilization, with the help of radioimmunoassay. In early pregnancy, hCG concentrations in plasma double in 1.4–2 days.

Functions of hCG:

• Stimulates the secretion of progesterone by the corpus luteum: Rescue and maintenance of the corpus luteum till 6 weeks of pregnancy is the major biological function of hCG.

• Stimulate Leydig cells of the male fetus to produce testosterone.

• Stimulates both adrenal and placental steroidogenesis.

• Stimulates the maternal thyroid because of its thyrotropic activity.

• Promotes secretion of relaxin from the corpus luteum.

High levels of hCG for that particular gestational age are detected in multiple pregnancies, hydatidiform mole or choriocarcinoma, and in pregnancy with a trisomy 21 fetus (Down's syndrome). It is low in cases of ectopic pregnancy or abortion.

Other options:

Option A: LH is secreted by the gonadotrope cells of the anterior pituitary.

Option C: FSH is also secreted by the gonadotrope cells of the anterior pituitary.

Option D: Gonadotropin-releasing hormone (GnRH) regulates the synthesis and secretion of both the hormones, LH and FSH.

Solution to Question 40:

Painless vaginal bleeding is not a feature of abruption of the placenta.

Patients with placental abruption present with sudden-onset abdominal pain, vaginal bleeding, uterine tenderness, and frequent contractions. Painless vaginal bleeding can occur in placenta previa where the placenta is implanted in the lower uterine segment over or very close to the cervical os.

Abruptio placenta is the partial/total separation of a normally implanted placenta before delivery.

The bleeding may be

• Revealed (blood dissects downwards between membranes and uterine wall to escape through the cervix), or

• Concealed (blood accumulates between detached placenta and uterus), or

• Mixed

Risk factors for abruptio placenta include prior abruption history, preterm premature rupture of membranes, and pregnancy-associated hypertension. Depending upon the degree of placental abruption and its clinical effects, the cases are graded as follows:

Uteroplacental apoplexy/Couvelaire uterus is a complication of severe concealed abruptio placentae. There is massive intravasation of blood into the uterine musculature up to the serous coat. It is only diagnosed on laparotomy, where a port-wine-colored uterus (diffuse or patchy) is seen. The presence of Couvelaire uterus alone is not an indication of hysterectomy since it does not interfere with uterine contractions.

Other complications of abruption of placenta:

• Disseminated intravascular coagulation

• Hypovolemic shock due to torrential bleeding

• Acute kidney injury

Abruptio placenta is mostly managed by immediate delivery and management of complications such as shock with adequate blood products.

Grade 1	Grade 2	Grade 3
Slight vaginal bleeding	Mild to moderate vaginal bleeding	Bleeding is moderate to severe or concealed.
Minimal uterine tenderness	Uterine tenderness is always present	Marked uterine tenderness
Maternal BP and fibrinogen will be unaffected	Increased maternal pulse, but BP is maintained, fibrinogen levels may be decreased	BP decreases, associated coagulation defect or anuria may complicate
FHS good	Fetal distress present	Fetal deathoccurs
Maternal shock absent	Maternal shock absent	Pronounced shock

Solution to Question 41:

The signs that can be elicited at 8 weeks of gestation are Goodell's sign and Osiander's sign.

Goodell's sign is a sign of early pregnancy, which describes the marked softening of the cervix in contrast to the non-pregnant state. It is seen as early as the 6th week.

Osiander's sign describes the pulsations felt in the lateral fornix due to increased vascularity. It is seen around 8 weeks of gestation.

Internal ballottement can be elicited between 16–28th weeks of gestation, before which the fetus is too small and after which it is too large to displace. It may not be elicited in cases with scanty liquor amnii, or when the fetus is transversely placed.

Lightening is a sense of relief of the pressure symptoms due to the engagement of the presenting part. It is felt at about 38 weeks of gestation.

Solution to Question 42:

The non-contraceptive uses of oral contraceptives include all except protection against sexually transmitted diseases (STDs), including HIV.

Oral hormonal contraceptives can be combined oral contraceptive pills (COCPs) or progesterone-only minipills.

COCPs are a combination of ethinylestradiol at a dose of 20–30 mcg and an orally active progestogen. It is taken starting on the first day of the cycle for 21 days. They are taken at a fixed time of the day, preferably after a meal. Mala-N and Mala-D both contain levonorgestrel (0.15 mg) + ethinylestradiol (30 mcg). Mala-N is supplied free of cost in Family Planning Clinics in India. Each strip contains 21 hormonal tablets and 7 non-hormonal (iron) tablets.

The mechanism of action is the prevention of ovulation by suppression of the pituitary hormones FSH and LH. Progesterone also acts on the cervical mucus, making it thick and impenetrable by sperm. The pregnancy rate with the COCP is 0.3 per 100 woman-years.

Benefits of the COCP:

• It is effective in controlling fertility.

• It causes regular cycles and reduces menstrual blood loss and are, therefore, useful in menorrhagia and polymenorrhea.

• It lowers the incidence of benign breast neoplastic diseases, such as fibrocystic disease.

• It reduces the incidence of functional ovarian cysts and ovarian (Option A) and endometrial malignancy (Option D). The incidence of ovarian cancer is reduced by 40% and that of uterine malignancy, by 50%, if taken for 1 year; this effect lasts as long as 10 years after stoppage.

• It reduces the incidence of ectopic pregnancy (Option B) by suppressing ovulation and reducing the risk in pelvic inflammatory disease.

• It is useful in endometriosis, acne, and polycystic ovarian disease (PCOD).

Side effects:

• Higher incidence of breast cancer reported among users

• Suppression of lactation

• Occurrence of pituitary adenoma and adenoma of the liver

• Reduced carbohydrate tolerance (therefore, contraindicated in diabetic women)

• Migraine, depression, weight gain, and lethargy (due to progesterone)

• Increased risk of pulmonary embolism and cerebral thrombosis

Ninety-nine percent of women will regain normal menstrual cycles within 6 months of stopping the pills, while the return of fertility is slightly delayed.

Barrier methods like condoms prevent the transmission of STDs.

Solution to Question 43:

The patient's clinical features are strongly suggestive of Turner syndrome.

Turner's syndrome is the most common sex chromosome abnormality in females. The classical karyotype is 45, XO. It is characterized by hypogonadism in phenotypic females.

The clinical features of Turner's syndrome include short stature, short webbed neck, shield chest with widely spaced nipples, primary amenorrhea, congenital heart defects, and renal anomalies. Two X chromosomes are required for the development of ovaries. The loss of one X chromosome results in streak ovaries with inadequate estrogen production. The loss of negative feedback from reduced estrogen raises the FSH and LH levels causing hypergonadotropic hypogonadism.



Other options:

Option B: Kallmnann syndrome presents with delayed puberty due to hypogonadotropic hypogonadism from deficiency of GnRH. It is associated with olfactory dysfunction and anosmia.

Option C: Androgen insensitivity syndrome occurs due to a defect in the androgen receptor in a genotypic male (XY). These individuals are phenotypic females with no male or female internal genitalia. They have normal female external genitalia with an absent or shallow vagina, well-developed breasts, and scanty body hair (axillary and pubic) and are present with primary amenorrhea.

Option D: Mayer-Rokitansky-Küster-Hauser (MRKH) syndrome, commonly referred to as Mullerian agenesis, has primary amenorrhea due to failure of development of the uterus, cervix, and upper vagina. They have normally functioning gonads and external genitalia.

Solution to Question 44:

The correct sequence in AMTSL is 3,2,4,1.

AMTSL is recommended by the WHO to prevent postpartum hemorrhage (PPH). Uterotonic administration is the most important component in AMTSL. However, even before a uterotonic is administered, the birth attendant will often place their hand over the uterus to ensure that there is no undiagnosed twin.

The currently recommended components of AMTSL are as follows:

Note: Uterine massage or fundal massage is not recommended as an intervention to prevent PPH in women who have already received prophylactic uterotonics such as oxytocin (core component of AMTSL). Hence, it is now an optional component of AMTSL that is done in patients with uterine atony as a part of the management of PPH.

	Component	Remarks
1.	Injectinguterotonicswithin1 minuteof delivery of the baby or immediately after delivery of theanterior shoulder	Most importantstepOxytocin 10 IU im/iv infusion is thepre ferred uterotonic.(Not recom mendedby WHO-carboprost)
2	Delayed cord clamping	Done within1-3 minsof delive ry80 mlof additional blood g oes to the fetusPreventsfetal anemia
3	Delivery of placenta by contr olled cord traction:Modified Brandt Andrews technique	Done only in settings where s killed birth attendants are av ailableThe uterus is pushed u pwards and backwardThe tra ction of cord is downwards a nd forwards
4	Intermittent assessment of ut erine tone	Done every 15 mins for 2 hou rs

Solution to Question 45:

The specimen showing an ovarian mass with a tuft of hair in a cystic space is a mature cystic teratoma.

Teratomas are germ cell tumors. They contain embryonic tissues leading to the differentiation of cells along these pathways. There are three classes of teratomas:

• Mature (benign) teratomas (e.g.: dermoid cyst)

• Immature (essentially malignant) teratomas (e.g.: solid teratoma)

• Monodermal (highly specialized) tumors (e.g.: struma ovarii)

Mature cystic teratomas, also known as dermoid cysts are unilocular, containing sebaceous material and hair. They have a smooth surface and are lined by squamous epithelium. Other structures like teeth, bone, cartilage, thyroid tissue, and neural tissues can also be found.

Most patients are asymptomatic. Common presentations include abdominal mass, acute pain due to torsion of the ovary, and rarely due to rupture of the cyst with spillage of contents into the

abdomen.

Treatment of mature teratomas involves excision in order to prevent complications.

Option A - Immature teratomas of the ovary are solid and trabeculated. They also contain cartilage, bone, hair, and sebaceous material. There is no structural order. They are generally malignant. Grossly, they have a smooth external surface and are solid on sectioning. There may be areas of necrosis or hemorrhage.

The following image shows both mature (left) and immature teratomas (right).



Option C- Mucinous cystadenoma is a benign tumor showing multicystic appearance, delicate septa, and the presence of glistening mucin within the cyst.

Option D- Serous cystadenoma have multicystic lesion in which papillary epithelium is contained within fibrous walled cysts or as a mass projecting from the ovarian surface.

Solution to Question 46:

Among the given options, placenta previa shows warning bleeds.

Dark painful bleeding with a tender uterus that is disproportionately enlarged is seen in placental abruption.

Placenta previa is a condition where the placenta is implanted in the lower uterine segment either over or close to the internal cervical os.

Risk factors for placenta previa

• Multigravida

• Age > 35 years

• Smoking

• Previous cesarean section or myomectomy

• Abnormal placental size and lobes

• Prior curettage

Clinical features of placenta previa:

• Sudden-onset recurrent painless vaginal bleeding: First bleeding usually appears after the end of the second trimester or later and is usually mild and not fatal. This is called 'Sentinel bleed'. In about one third of cases, there is a history of 'warning hemorrhage' which is usually slight.

• The size of the uterus corresponds to gestational age.

• The uterus is relaxed, soft, no tenderness

• Floating head

• Fetal heart sound is present unless there is major separation and the patient is exsanguinated.

• Slowing of fetal heart rate on pressing the head down into the pelvis which recovers promptly as pressure is relieved is suggestive of a low-lying placenta especially of posterior type: Stallworthy sign.

Degree of placenta previa:

There are four types of placenta previa depending upon the degree of extension of the placenta to the lower segment

• Type—I (Low-lying): The major part of the placenta is attached to the upper segment and only the lower margin encroaches onto the lower segment but not up to the internal os.

• Type—II (Marginal): The placenta reaches the margin of the internal os but does not cover it.

• Type—III (Incomplete or partial central): The placenta covers the internal os partially (covers the

internal os when closed but does not entirely do so when fully dilated).

• Type—IV (Central or total): The placenta completely covers the internal os even after it is fully dilated.

Per-Vaginal Examination should not be done in suspected Placenta Previa, whereas it can be only done in the operation theatre with all the arrangements for C-section.

The image below shows the types of placenta previa.



Expectant management for placenta previa was advocated by Macafee and Johnson. The aim is to continue the pregnancy for fetal maturity without compromising maternal health.

Prerequisites

• Availability of blood for transfusion whenever required

• Facilities for cesarean section should be available 24 h.

Suitable cases for expectant management for placenta previa:

• Mother in good health (Hb ≥10 g%; hematocrit ≥30%)

• Duration of pregnancy <37 wk

• No active vaginal bleeding

• Fetal well-being is assured (USG)

Conduct of expectant treatment

• Corticosteroids for fetal lung maturity if <34 wk

• Betamethasone 12mg IM 24 hours apart for 2 doses or

• Dexamethasone 6mg IM 12 hours apart for 4 doses

• Bedrest

• Hb, Blood grouping and urine for protein

• Inspection of vulval pads, fetal surveillance with USG

• Supplementary hematinics

• Rh Immunoglobin should be given to all Rh-negative (unsensitized) women

• Use of tocolytics if there are uterine contractions.

The pregnancy is carried till 37 weeks as part of expectant management. During this time, if there is fetal distress or a bleeding episode, then terminate immediately.

Solution to Question 47:

The pregnant woman exhibiting features of severe pre-eclampsia (BP of 170/110 mmHg, epigastric pain, headache, and blurring of vision) and in active labor should be managed by treating hypertension, administering $MgSO_4$ (magnesium sulfate), and considering termination of pregnancy.

Preeclampsia is defined as hypertension with blood pressure ≥140/90 mm Hg along with proteinuria that develops after 20 weeks of gestation in a previously normotensive and non-proteinuric woman. It can be classified as severe if any of the following are present:

• BP ≥160/110

• Headache

• Visual disturbance

• Upper abdominal pain

• Elevated creatinine

• Thrombocytopenia (<1,00,000 cells per mm3)

• Pulmonary edema

• Marked elevation in serum transaminases

This patient has many features of severe pre-eclampsia and is in active labor. The best management for her is to admit her and bring down her BP using labetalol, nifedipine, or hydralazine and give seizure prophylaxis with magnesium sulfate in the same way as in eclampsia. Since the placenta is the main pathology causing pre-eclampsia, termination of pregnancy and delivery of the placenta is the definitive treatment.

Solution to Question 48:

Vulval hamartoma is not a risk factor for vulval cancer.

Vulval cancer is a rare genital tract cancer that may be divided into preinvasive lesions and invasive carcinoma. 90% of the invasive carcinomas are squamous cell carcinoma and are usually present on the labia majora. The risk factors for both types are:

• Chronic vulval irritation

• Immunosuppressive conditions such as pregnancy, HIV infection

• Smoking

• Condyloma (anogenital warts)

• Sexually transmitted diseases

• Dystrophies

Solution to Question 49:

An upper segment cesarean section is done in case of perimortem cesarean section.

Perimortem cesarean section is an emergency cesarean delivery of a woman who is expected to die within the next few moments or has just died. AHA 2020 Guidelines for CPR recommend that perimortem cesarean should be considered if there is no return of spontaneous circulation in 5 minutes in a pregnant woman who has experienced total cardiopulmonary arrest.

The goal of perimortem cesarean section (PMCS) is to improve maternal and fetal outcomes. PMCS improve maternal survival by decreasing compression of the IVC by the gravid uterus. This improves venous return and diaphragmatic displacement, which in turn improves respiratory dynamics. The infant may survive if PMCS is done within 10 minutes of maternal death. A classical upper-segment cesarean section with a vertical uterine incision is done.

Other options:

Option A: Upper segment cesarean section is done, not lower segment cesarean section (LSCS).

Option B: A vertical incision from the xiphoid to the symphysis pubis is done on the abdomen and not the lower segment Pfannenstiel incision.

Option D: There is no standardized guideline about where perimortem cesarean section must be performed.



Solution to Question 50:

The cardinal movements of labor in order are engagement-crowning-restitution-external rotation.

The specific cardinal movements of labor are:

• Engagement is the mechanism by which the biparietal diameter (in occiput presentation) passes through the pelvic inlet.

• Descent is a continuous process that is completed with the expulsion of the fetus.

• Flexion of the head occurs as soon as the descending head meets resistance from the pelvic floor/pelvic walls/cervix.

• Internal rotation is the mechanism by which the occiput gradually turns away from the transverse axis. Crowning occurs after internal rotation.

• Crowning occurs when the vulval outlet is stretched by the maximum diameter of the head and there is non recession of the head after the contraction is over.

• Extension occurs when the flexed head reaches the vulva.

• Restitution is the passive movement of the head due to untwisting of the neck that occurred in internal rotation.

• External rotation is the visible movement of the delivered head occurring due to internal rotation of the shoulders.

• Expulsion is the process by which the anterior shoulder comes under the pubic symphysis to be delivered and is followed by the posterior shoulder and the rest of the fetal body.



Solution to Question 51:

The correct match is A-4, B-1, C-2, D-3

A table differentiating between the invasive prenatal diagnostic techniques is given below.

Fetoscopy is a type of micro laparoscopic technology that aids in fetal diagnosis and therapeutic interventions.

A. Amniocentesis	Amniotic fluid assessed for d etection of AFP.
B. Chorionic villous sampling	Done at 8-12wks
C. Fetal blood sampling	Done at 19-20 wks

A. Amniocentesis	Amniotic fluid assessed for d etection of AFP.
D. Fetoscopy	Fetal organs visualized

	Chorionic villus sampling	Amniocentesis	Cordocentesis
Time	Transcervical 10–13 weeks, Transabdominal 10 weeks t o term	After 15 weeks (early12–14 weeks)	18–20 weeks
Materials for stu dy	Trophoblast cells	Fetal fibroblasts, fluid for bi ochemical study	Fetal white blood cells (othe rs - infection and biochemic al study)
Karyotype result	Direct preparation: 24–48 h ours.Culture: 10–14 days	Culture:3–4 weeks	Culture:24–48 hours
Fetal loss	0.5–1%	0.5%	1–2%
Accuracy	Accurate; may need amnioc entesis for confirmation	Highly accurate	Highly accurate

Solution to Question 52:

Vagitus uterinus is the cry of the baby in the womb.

Vagitus uterinus is the cry of an in-utero baby after the rupture of membranes, during the process of labor but before it has been delivered. In one such instance, it was described as a clear remote cry, much like a baby crying in the next room but close below their heads.

Obstetrics and Gynecology NEET 2022

Question 1:

A pregnant lady with 34 weeks of amenorrhea has the following findings: LDH - 700 IU/L, platelets - 75,000/mm3, serum bilirubin - 1.5 mg/dL, SGOT - 200 U/L, SGPT - 150 U/L, and BP - 140/96 mm Hg. Her coagulation profile and renal function tests are normal. What is the diagnosis?

a) HELLP syndrome

b) Acute fatty liver of pregnancy

c) Viral hepatitis

d) Intrahepatic cholestasis

Question 2:

A woman at 26 weeks of gestation presents for routine evaluation. On examination, fundal height corresponds to 24 weeks. Ultrasonography revealed decreased amniotic fluid. Which of the following conditions would have led to this presentation?

a) Renal agenesis

b) Tracheoesophageal fistula

c) Cardiac abnormalities

d) Ureteral stricture

Question 3:

A type 1 diabetic mother is on magnesium sulfate infusion post-cesarean section for preeclampsia. She develops delirium and is drowsy. She has a respiratory rate of 10/min, random blood glucose level of 240 mg/dL, oliguria, and bilaterally absent knee reflex. What is the cause of her condition?

a) Magnesium sulfate toxicity

b) Diabetic ketoacidosis

c) Eclampsia

d) Diabetes insipidus

Question 4:

Identify the type of hymen shown in the image.



a) Imperforate hymen

b) Semilunar hymen

c) Septate hymen

d) Annular hymen

Question 5:

While discharging a patient who underwent a vesicovaginal fistula repair, which of the following would you recommend?

a) Sexual abstinence for 3 months and avoid pregnancy for a year

b) Sexual abstinence for 3 weeks and avoid pregnancy for 6 months

c) Sexual abstinence for 6 weeks and avoid pregnancy for a year

d) Sexual abstinence for 6 months and avoid pregnancy for 6 years

Question 6:

Which of the following is an absolute contraindication for the insertion of the device shown in the image below?



a) Menstruation

b) Trophoblastic disease

c) Immediately after delivery

d) Ruptured condom during intercourse

Question 7:

A 20-year-old woman is evaluated for primary infertility. Hysterosalpingography was done and reveals the finding shown below. What is the anomaly seen in the image?

a) Septate uterus

b) Uterine didelphys

c) Bicornuate uterus

d) Unicornuate uterus

Question 8:

A pregnant woman with no other comorbid conditions develops preeclampsia. She enquires about the cause of her condition. The doctor explains that it is due to the failure of the invasion of:

a) Spiral artery by villous trophoblasts

b) Radial artery by cytotrophoblasts

c) Spiral artery by extravillous trophoblasts

d) Arcuate artery by extravillous trophoblasts

Question 9:

A woman with an obstetric score of G2P1 comes to the clinic at 14 weeks of gestation for her antenatal checkup. A uterine artery doppler was suggested by the doctor. What would it detect?

a) Early-onset preeclampsia

b) Late-onset preeclampsia

c) Fetal growth restriction

d) Placenta accreta

Question 10:

While conducting a delivery, you perform the maneuver shown below. Following this, there is incomplete separation of the placenta and massive hemorrhage. What is your next step in management?



a) Start oxytocin infusion and wait for spontaneous delivery of placenta

b) Uterine massage

c) Arrange for blood and use Crede's method for placental delivery

d) Manual removal of placenta

Question 11:

A female patient collapses soon after delivery. There is profuse bleeding and features of disseminated intravascular coagulation. Which of the following is the most likely etiology?

a) Amniotic fluid embolism

b) Postpartum hemorrhage

c) Peripartum cardiomyopathy

d) Rupture of the uterus

Question 12:

You are examining a multigravida in the second stage of labor for the past two hours. On examination, contractions are adequate, the cervix is dilated with the head at station 0 with molding 2+ and caput 2+. The sagittal suture is in the right occipitotransverse position. The fetal heart rate is 140 beats/minute. Which of the following is carried out for the management of this patient?

a) Midpelvic forceps

b) Vacuum-assisted delivery

c) Wait for an hour for spontaneous labor

d) Lower segment cesarean section

Question 13:

A primigravida presents to the labor room at 40 weeks of gestation with lower abdominal pain. She has been in labor for 3 hours. Which of the following will determine if she is in active labor?

a) Fetal head 5/5 palpable on abdominal examination

b) Two contractions lasting for 10 seconds in 10 minutes

c) More than 5 cm cervical dilatation with complete effacement

d) Rupture of membranes

Question 14:

Identify the blue-colored area shown in the image below.



a) Obstetric outlet

b) Anatomical outlet

c) Pelvic inlet

d) Midpelvis

Question 15:

What will be the level of the uterus on the postpartum day 2?

a) Two finger breadths below umbilicus

b) One finger breadth below umbilicus

c) Three finger breadths below umbilicus

d) Four finger breadths below umbilicus

Question 16:

A primigravida woman at 12 weeks of gestation comes to the antenatal clinic for nutritional advice. Which of the following will you recommend?

a) Additional 300kcal in 2nd trimester

b) Additional 300kcal in 1st trimester

c) Additional 400kcal in 3rd trimester

d) Additional 300kcal throughout the pregnancy

Question 17:

A 40-year-old G2P1 woman with 18 weeks of amenorrhea comes with a dilated cervix. The cervical length is 15 mm. In spite of explaining the risks, she insisted on cerclage. Which of the following is a contraindication for cervical cerclage?

a) Ruptured membranes

b) Prolapse of membranes into the vagina

c) Fetal fibronectin positive

d) Advanced maternal age

Question 18:

A primigravida presents to the emergency room in the early stage of labor with adequate uterine contractions. On per vaginal examination, a gynecoid pelvis is felt, the membranes are ruptured, and the vertex is felt in the right occipito-posterior position. How will you manage this patient?

a) Vacuum-assisted delivery

b) Cesarean section

c) Normal vaginal delivery

d) Forceps delivery

Question 19:

A pregnant patient, with a history of classical cesarean section in view of fetal growth retardation in the previous pregnancy, presents to you. She is currently at 35 weeks of gestation with breech presentation. What is the next step in management?

a) Cesarean section at 37 weeks

b) Advice USG and visit after 2 weeks

c) Internal podalic version followed by vaginal delivery

d) External cephalic version at 36 weeks

Question 20:

A primigravida at 22 weeks of gestation presents to you with profuse vaginal bleeding. Her blood pressure and glucose levels are within normal limits. Placental implantation at which of the following sites can cause this?

a) Internal os

b) Fallopian tube

c) Ovarian

d) Abdominal

Question 21:

A female patient presents to you with six weeks of amenorrhea, associated with abdominal pain and vaginal bleeding with normal blood pressure. Investigations revealed beta-hCG to be 1400 mIU/mL. An ultrasound scan was done which showed a trilaminar endometrium with normal adnexa. What is the next best step in the management of this patient?

a) Repeat beta-hCG after 48 hours

b) Repeat ultrasound after 3 days

c) Measurement of progesterone

d) Laparoscopy

Question 22:

A 24-year-old lactating female with an 18-month-old child comes with a history of irregular, heavy bleeding seeking contraceptive advice. Which is the contraceptive of choice?

a) Progestasert

b) Copper-T 380A

c) Mala N

d) Norethisterone enanthate - depot injection

Question 23:

A 25-year-old male patient is evaluated for primary infertility. Semen analysis shows azoospermia. A testicular biopsy is done and the image is shown below. The shown finding is consistent with



a) Sertoli cell only syndrome

b) Testicular atrophy

c) Benign testicular neoplasm

d) Orchitis

Question 24:

A 20-year-old woman presented at 7 weeks of gestation, unwilling to continue the pregnancy. What are the drugs used for medical termination of pregnancy in this patient?

a) Misoprostol and Medroxyprogesterone

b) Misoprostol and Mifepristone

c) Mifepristone and Methotextrate

d) Mifepristone and Medroxyprogesterone

Question 25:

Testosterone helps in the development of various organs in the fetus. Which of the following stimulates its production?

a) LH from maternal pituitary

b) hCG from placenta

c) Inhibin from corpus luteum

d) GnRH from fetal hypothalamus

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	c
5	a
6	b
7	a
8	c
9	a
10	d
11	a
12	d
13	c
14	d
15	a
16	a
17	a
18	c

19	a
20	a
21	a
22	c
23	a
24	b
25	b

Detailed Explanations

Solution to Question 1:

The given clinical scenario with elevated liver enzymes and bilirubin, low platelet count, and hypertension indicates the diagnosis of HELLP syndrome.

HELLP is an acronym for hemolysis (H), elevated liver enzymes (EL), and low platelet count (LP). It usually occurs in patients with severe preeclampsia. The patients typically present with right upper quadrant pain, nausea, and vomiting. Hypertension and proteinuria are seen but may be absent in some cases.

In this condition, there is periportal necrosis of the liver causing an elevation in liver enzymes (AST, ALT, and LDH) and bilirubin. Tennessee criteria used to diagnose HELLP syndrome include:

• Platelet count - <1 lakh/mm3

• AST - >70 IU/L

• LDH - >600 IU/L

• Hemolysis

Prompt delivery is indicated for ≥34 weeks gestational age. Vaginal delivery is the preferred route of delivery if cervix is favorable. Platelet transfusion is required if counts are less than 50,000/mm3. Magnesium sulfate is given as antiseizure prophylaxis, and anti-hypertensives are given for severe hypertension.

Complications of HELLP include eclampsia, placental abruption, hepatic subcapsular hematoma, coagulopathy, stroke, and pulmonary edema.

The clinical and laboratory findings of acute liver diseases in pregnancy are as follows:

Disorders	HELLP syn drome	Obstetric cholestasis	Acute fatty liver of pregna ncy	Viral hepatit is
Onset in pregnancy	Mid to late	Late	Late	Variable
Differentiating featur es	-	Pruritus(elevatedserum bil e acids)	Signs ofliver failure	Fever
AST (U/L)	≤300	≤200	200-800	2000+

Disorders	HELLP syndrome	Obstetric cholestasis	Acute fatty liver of pregnancy	Viral hepatitis
Bilirubin (mg/dL)	1-4	1-5	4-10	5-20
Platelets	↓↓	Normal	↓↓	↓
Fibrinogen	Normal	Normal	↓↓↓	Normal
D-dimer	↑	Normal	↑	Normal
Prothrombin time	Normal	Normal	↑↑	↑
Hemolysis	↑↑	No	↑↑↑	No

Solution to Question 2:

The given clinical scenario and decreased amniotic fluid on ultrasonography are suggestive of oligohydramnios. Among the given conditions, renal agenesis is the most likely cause.

Oligohydramnios is a condition where the amount of amniotic fluid is reduced. The major source of amniotic fluid in the second trimester is fetal urine. When the fetus has renal agenesis, urine production is affected. This leads to decreased amniotic fluid/oligohydramnios.

A woman with oligohydramnios usually presents with decreased fundal height for gestational age and decreased fetal movements. On examination, fetal parts can be clearly felt because of less liquor. Fetal malpresentation and intrauterine growth retardation are seen in this condition. It is diagnosed with the help of ultrasonography when the deepest vertical pocket (DVP) of liquor is <2 cm or when the amniotic fluid index (AFI) is <5 cm at gestational age, from 28 to 40 weeks.

It can lead to abortion, deformities like wry neck, club foot, etc, fetal pulmonary hypoplasia, and cord compression. Adequate hydration and amnioinfusion can be used to manage this condition.

Polyhydramnios or hydramnios is a condition where the amount of amniotic fluid is abnormally increased. Ultrasonographic diagnosis is made when the deepest vertical pocket (DVP) is >8 cm or when the amniotic fluid index (AFI) is ≥25 cm.

Other options:

Option D- Ureteral stricture usually arises secondary to trauma or iatrogenic interventions like surgery.

Solution to Question 3:

The given clinical scenario and examination finding of bilaterally absent knee reflex following magnesium sulfate ($MgSO_4$) administration for preeclampsia suggest the cause of her condition to be magnesium sulfate toxicity.

$MgSO_4$ is used as an anticonvulsant in the management of eclampsia and as prophylaxis in severe preeclampsia. It can also be used as a tocolytic to stop preterm labor. The therapeutic level of magnesium in eclampsia is 4–7 mEq/L.

Patients must be monitored for magnesium toxicity:

• Loss of deep tendon reflexes

• Decreased respiratory rate <12/min

• Urine output <30 mL/h

• Chest pain and heart block

• Visual changes, flushing, muscle paralysis, and drowsiness

Measurement of serum magnesium levels is indicated to monitor magnesium toxicity if serum creatinine >1.0 mg/dL.

Management of magnesium toxicity is as follows:

• Stop magnesium therapy

• Estimation of serum magnesium and creatinine levels

• Intravenous injection of calcium gluconate 10 mL (10% solution)

The two commonly used $MgSO_4$ regimens are described below:

Repeated/maintenance dose IM injections of magnesium sulfate are continued only under the following conditions:

• The patellar reflex is present

• Respiration is not depressed

• Urine output >30 mL/h

Other options:

Option B: Pregnant women with diabetic ketoacidosis present with hyperventilation (Kussmaul breathing) and their blood glucose level is around 380 mg/dL.

Option C: Eclampsia patients present with generalized tonic-clonic convulsions.

Option D: Diabetes insipidus patients present with polyuria and polydipsia.

Regimen	Loading Dose	Maintenance Dose
Intramuscular (Pritchard)	4 g (20% solution) IV over 3–5 minutes, followed by 10 g (50%) deep IM (5 g in each bu ttock)	5 g (50%) IM 4 hourly in the alternate buttock
Intravenous (Zuspan orSibai)	4 g IVslow over 15–20 minut e	1–2 g/h IV infusion

Solution to Question 4:

The given image is suggestive of the septate hymen.

The hymen is the membranous vestige of the junction between the sinovaginal bulbs and the urogenital sinus, which perforates during fetal life to establish a connection between the vaginal

lumen and the perineum. Defective perforation of this structure may result in the following variants of hymen

• Imperforate - without any opening

• Microperforate - tiny opening

• Annular - oval central opening

• Septate - two lateral openings partitioned by a bridge of hymenal tissue

• Naviculate - boatlike opening

Patients with septate hymen will typically complain of menstrual irregularities or difficulty with tampon placement or intercourse. Under topical anesthesia, a rapidly absorbable suture or free tie can be placed at the anterior and posterior aspects of the septal tissue, and the septum is resected between the two sutures.

Some of the variants of the hymen are:

Imperforate hymen results due to failure of canalization of the inferior end of the vaginal plate. Clinical features include primary amenorrhea, cyclical abdominal pain, and bluish bulging membranes at the vaginal opening. A simple cruciate incision in the membrane will restore normal reproductive function.



Solution to Question 5:

While discharging a patient who underwent vesicovaginal fistula repair, sexual abstinence for 3 months and avoiding pregnancy for a year would be recommended.

Vesicovaginal fistula (VVF) is a type of urinary fistula which commonly arises due to obstetric trauma. Small fistulas due to difficult childbirth can heal spontaneously. Most established fistulas need to be repaired vaginally usually after 3 months to allow the inflammation to subside. Post-operatively, continuous bladder drainage for 14 days along with antibiotics is recommended. Vaginal or speculum examination and intercourse are not advised for 3 months. Avoiding

pregnancy for a year is also recommended. A cesarean section is usually done for future pregnancies.

Presenting features of VVF include constant dribbling of urine/ true incontinence. Excoriation of the vagina, vulva, and thighs can occur due to constant contact with urine. Few patients might develop bladder stones. Vaginal urine collection can be seen on retraction of the posterior vaginal wall with a speculum. A bimanual examination can be used to assess the fixity and scarring of the surrounding tissues. A cystourethroscopy can be used for localization and assessing bladder mucosa viability.

Solution to Question 6:

The given image shows a copper-containing IUCD (intrauterine contraceptive device). Trophoblastic disease is an absolute contraindication for the insertion of this device.

In cases of uterine invasion like trophoblastic disease, there is an increased likelihood of complications related to IUCD such as perforation, misplaced device, and bleeding.

IUCDs are a type of long-term, reversible contraception. They are T-shaped devices that are inserted into the uterine cavity. These can either be copper-containing like CuT-380A and CuT-200, or hormone-containing IUCDs like Mirena.

The contraceptive action of IUCDs is because of foreign body reaction which leads to various changes in endometrial tissue that interfere with implantation. These changes include prostaglandin release that provokes uterine contractility and tubal peristalsis and leucocytic infiltration in the endometrium. It also interferes with the migration of spermatozoa.

The contraindications for insertion of IUCD include:

• Trophoblastic disease

• Distorted uterine cavity (fibroids, uterine anomalies)

• Ongoing pelvic infection

• Undiagnosed genital tract bleeding

• Suspected pregnancy

• Severe dysmenorrhea

• STIs — current or recurrent

• Wilson disease and copper allergy, for copper-containing IUCDs

• Breast cancer, for hormone-containing IUCDs

The different types of IUCD are shown below:



The timing of insertion of IUCD is as follows:

• Interval insertion (inter-conceptional period)- preferable to insert 2–3 days after menstruation, but it can be inserted at any time during the cycle. It can also be inserted during menstruation which has certain advantages like an open cervical canal and a distended uterine cavity

• Post-partum insertion is done at least 6 weeks after delivery when the uterus involutes to the normal size

• Postplacental insertion can be done immediately following delivery of the placenta

• Inserted within 5 days of coitus as an effective method of emergency contraception (Eg: incase of ruptured condom during intercourse)

Solution to Question 7:

The anomaly seen in the given hysterosalpingogram is suggestive of a septate uterus.

A septate uterus is characterized by the presence of a septum in the endometrial cavity due to the failure of the medial segments of the Müllerian ducts to regress. This creates a permanent septum within the uterine cavity. The septum can project partially from the fundus or can extend completely to the cervical os. On HSG, the septate uterus is seen as widely diverging horns with an intercornual angle <75 degrees.

HSG provides information about the interior of the uterine cavity and not the exterior. Therefore, it cannot reliably differentiate between a septate uterus and a bicornuate uterus. A septate uterus has a smooth fundal contour whereas a bicornuate uterus has an indented fundus.

Transvaginal 3-D ultrasonography (3D TV USG) helps to differentiate between the bicornuate uterus and a septate uterus and is recommended for the assessment of patients with a suspected uterine anomaly. MRI is an accurate investigation to diagnose the septate uterus. Hysteroscopy combined with laparoscopy can confirm both the internal and the external contour of the uterus.

Hysteroscopic septoplasty is a preferred treatment option for the septate uterus. It refers to the resection of the uterine septum under hysteroscopic guidance. There is a reduced risk of pelvic adhesions, scarring and the necessity of cesarean sections for future pregnancies with the hysteroscopic approach, compared with transabdominal approaches.

Solution to Question 8:

Preeclampsia occurs due to the failure of the invasion of extravillous trophoblasts into the spiral artery.

The spiral artery supplies blood to the endometrium. In normal pregnancy, there is an invasion of the extravillous trophoblasts (specifically, endovascular trophoblasts) into the walls of the spiral arteries. In the first wave of invasion (10–12 weeks), endovascular trophoblasts invade up to decidual segments. In the second wave (16–18 weeks), they invade up to the myometrial segments.

This vascular remodeling process replaces the endothelial lining and the muscular arterial wall with endovascular trophoblasts. This results in the distension of the spiral arterioles, thereby transforming them into a low resistance, low pressure, and high flow system.

In preeclampsia, the failure of the second wave of endovascular trophoblastic invasion into uterine vessels leads to decreased placental perfusion.

The images below show the sequential branching of the uterine artery and the vascular remodeling in preeclampsia respectively,





Other contributory etiological factors include genetic factors and maladaptive responses to fetal antigens and cardiovascular changes in pregnancy.

The basic pathology in preeclampsia is endothelial dysfunction leading to decreased nitric oxide production and vascular spasm. This leads to ischemic changes that result in necrosis, hemorrhage, and end-organ damage.

Changes leading to vasospasm and endothelial dysfunction include:

• Reduced vasodilatory prostaglandin PGI2 and increased vasoconstrictor thromboxane (TXA2)

• Reduced angiotensinase activity and increased sensitivity of vessels to angiotensin II, which is a vasopressor

• Increased production of endothelin 1, a potent vasoconstrictor

• Increased inflammatory mediators (tumor necrosis factor- α, interleukins) and reactive oxygen species (ROS) causing endothelial injury

• Increased antiangiogenic factors like soluble fms-like tyrosine kinase I (SFlt-1) and soluble endoglin (sEng)

The definitive treatment of pre-eclampsia is the termination of pregnancy at ≥37 weeks.

Solution to Question 9:

A uterine artery doppler would detect early-onset preeclampsia.

The development of a diastolic notch and resistance to flow in the uterine artery doppler suggests the development of gestational hypertension, pre-eclampsia, and fetal growth restriction later in the pregnancy. A high resistance index in the second trimester is associated with a sixfold increase in the incidence of preeclampsia. The trophoblastic invasion of maternal uterine vessels in the first half of pregnancy results in a decrease in vascular resistance. This can be detected by Doppler studies.

Other screening tests which can be used for the prediction of pre-eclampsia include:

• Mean arterial pressure ≥90 mm Hg in the second trimester

• Absence of end-diastolic frequencies in umbilical artery doppler

• Roll over test

Preeclampsia is a multi-system disorder marked by blood pressure ≥140/90 mm Hg with proteinuria after the 20th week of gestation in a previously normotensive and non-proteinuric woman. Usually, there is an invasion of the extravillous trophoblasts (specifically, endovascular trophoblasts) into the walls of the spiral arteries. In the first trimester, the trophoblastic invasion up to the decidual segments occurs. The myometrial segments are invaded in the second trimester. This second wave of endovascular trophoblast migration fails to happen in preeclampsia.

The pathogenesis can be explained by an imbalance in the prostaglandins - a deficiency of PGI2 (vasodilator) and increased thromboxane (vasoconstrictor). The deficiency of nitric oxide and elevated antiangiogenic factors are other mechanisms responsible.

Features include headache, visual disturbances, epigastric pain, and thrombocytopenia. Risk factors include young maternal age, nulliparity, metabolic syndrome, and hyperhomocysteinemia.

Other options:

Option B: Late-onset preeclampsia occurs > 34 weeks. It often has a favorable outcome.

Option C: Fetal growth restriction can be detected by abnormal uterine artery Doppler notching at 20-24 weeks of gestation.

Option D: Placenta accreta or morbidly adherent placenta arises due to partial or total absence of decidua basalis. Sonography is recommended for diagnosis. Better results can be obtained with power doppler.

Solution to Question 10:

The maneuver shown is the Brandt-Andrews maneuver used in controlled cord traction for delivery of the placenta. The clinical features point to a diagnosis of primary postpartum hemorrhage due to retained tissues and the next step in management is manual removal of the placenta.

Postpartum hemorrhage (PPH) is defined as blood loss >500 mL following the birth of the baby. It is of two types:

• Primary PPH: It occurs within 24 hours after the birth of the baby. This is again classified into third-stage hemorrhage (bleeding before placental expulsion) and true postpartum hemorrhage (bleeding after placental separation).

• Secondary PPH: It occurs after 24 hours but within the puerperal period.

The four main causes of primary PPH are atonic uterus, trauma, retained tissues, and coagulopathies.

Retained tissues such as placental bits and blood clots lead to imperfect uterine retraction. Causes for delayed placental delivery include simple adherent placenta, lower uterine segment constriction, and morbidly adherent placenta. In the presence of brisk bleeding and partial or total attachment of the placenta, manual removal is indicated. It is carried out under adequate analgesia and aseptic precautions.

If the placenta is not expelled even after 30 minutes following the birth of the baby, it is known as retained placenta. It can lead to hemorrhage, shock, and puerperal sepsis. Unseparated retained placenta is manually removed under general anesthesia.

Solution to Question 11:

The given clinical scenario with a diagnosis of disseminated intravascular coagulation (DIC) points to amniotic fluid embolism as the most likely etiology.

Amniotic fluid embolism presents with a classic triad of hemodynamic and respiratory compromise, along with DIC. Usually, it presents during the late stages of labor or immediate postpartum. Gasping for air, seizures, and cardiorespiratory arrest are other features. Clinical features are variable.

Diagnostic criteria for amniotic fluid embolism include:

• Sudden onset cardiorespiratory arrest, or both hypotension and respiratory compromise

• Overt DIC

• Onset during labor or within 30 minutes of delivery of the placenta

• No fever ≥38°C

Since there are no specific diagnostic tests, a clinical diagnosis has to be made. Other causes for a similar presentation such as myocardial infarction, pulmonary embolism, and anaphylactic shock needs to be ruled out.

Risk factors include rapid labor, meconium-stained liquor, and uterine tears. Post-term pregnancy, maternal age, and eclampsia are other predisposing conditions. The currently accepted mechanism is the entry of fetal material into the maternal circulation which leads to the activation

of proinflammatory mediator systems. This results in pulmonary vasoconstriction and hypertension. The tissue factor present in fetal material also activates factor VII resulting in DIC.

The condition can be fatal and management includes cardiopulmonary resuscitation and advanced cardiac life support.

Other options:

Option B: Post-partum hemorrhage is the loss of ≥500mL of blood after the third stage of labor. Symptoms and signs of hypovolemia are commonly seen.

Option C: Peripartum cardiomyopathy presents with features of heart failure in the last month of pregnancy or within 5 months of delivery. Probable causes include viral myocarditis, malnutrition, and hormonal interactions.

Option D: Rupture of the uterus is a disruption in the continuity of all the uterine layers anytime after 28 weeks of pregnancy. Clinical features depend on the specific cause and include abdominal pain, vaginal bleeding, fainting attacks, and rapid pulse rate.

Solution to Question 12:

The given clinical scenario points to a diagnosis of deep transverse arrest. A lower segment cesarean section is carried out for the management of this patient.

In deep transverse arrest, the sagittal suture is placed in the transverse bispinous diameter. There is no descent of the head for at least 1/2-1 hour after full dilatation of the cervix. The mode of delivery is based on the fetal condition and pelvic assessment. Caput succedaneum indicates the static position of the head for a prolonged duration. Molding occurs due to the passage of the fetal head through a resistant birth passage. Grade 2 molding indicates overlapping parietal bones which can be separated easily. The presence of caput and molding of grade 2+ indicates a cephalopelvic disproportion. In such cases, a cesarean section has to be performed.

If the pelvis is adequate, application of ventouse or manual rotation and forceps application can be tried.

Deep transverse arrest, can be either due to incomplete forward rotation of the oblique occipitoposterior position or due to nonrotation in the occipitotransverse position.

Occipitoposterior position is an abnormal position of the vertex and can undergo normal labor in favorable circumstances. In some cases, it undergoes non-rotation resulting in oblique posterior arrest. If it undergoes malrotation (anterior rotation of the sinciput), it can cause a persistent occipitoposterior position. Spontaneous delivery can occur as face to pubis. If arrest occurs in this position, it is termed as occipitosacral arrest. Incomplete forward rotation, as described above, leads to a deep transverse arrest.

Features of occipitoposterior position include infra-umbilical flattening and fetal heart sounds localized to the flank. The fetal parts are more prominently felt near the midline.

Solution to Question 13:

This patient will have presented in active labor if she had more than 5 cm cervical dilatation with complete effacement

The first stage of labor is characterized by dilatation and effacement of the cervix and can be divided into latent phase and active labor. The onset of latent labor is the point at which the mother perceives regular contractions. Cervical dilatation of ≥5 cm in the presence of uterine contractions can be taken to reliably represent the threshold for active labor.

The first symptom to appear is intermittent painful uterine contractions followed by the expulsion of blood-stained mucus (show) per vagina. In the late first stage, the contraction comes at intervals of 3–5 minutes and lasts 45 seconds.

The second stage of labor begins with complete cervical dilatation and ends with fetal delivery.

The third stage of labor is characterized by the delivery of the placenta.

The fourth stage is a stage of observation for at least an hour after the full expulsion of the placenta.

Fetal head 5/5 on abdominal examination, signifies the degree of descent in relation to the pelvic inlet and denotes the head is floating above the brim.

Rupture of membranes at or near term signifies the labor is likely to begin soon.

More than 12 uterine contractions in 1 hour may signify labor onset. Partograph is charted every half an hour recording the number of contractions in 10 minutes and the duration of each contraction in seconds. Contractions can be classified as mild (contraction duration less than 20 seconds), moderate(between 20 and 40 seconds), and strong (more than 40 seconds).

Solution to Question 14:

The blue colored area in the given image shows midpelvis.

Midpelvis is the region of the pelvis that is bounded by the plane of the greatest pelvic dimensions above and the midpelvic plane below.

The plane of greatest pelvic dimensions (most roomy pelvic plane) extends from the middle of the posterior surface of the symphysis pubis to the junction of the second and third sacral vertebrae.

The midpelvic plane extends from the lower margin of the symphysis pubis to the tip of the sacrum at the level of ischial spines (coinciding with the plane of least pelvic dimensions), or from the lower margin of the symphysis pubis to the junction of S4 and S5, making a wedge posteriorly.



Other options:

Option A: Obstetric outlet - It is the region of the pelvis that is bounded by the plane of the least pelvic dimensions above and the anatomical outlet below.



Option B: Anatomical outlet/ bony outlet - It corresponds to the bony outer margin of the pelvis. It is a diamond-shaped region formed by joining two triangular planes with a common base. The anterior triangle is bounded laterally by the ischiopubic rami and the apex is formed by the inferior border of the pubic arch. The posterior triangle is bounded laterally by the ischial tuberosity and sacrotuberous ligament and the apex is formed by the tip of the coccyx. The common base is formed by the line joining the ischial tuberosities.



Option C: Pelvic inlet - It represents the superior plane of the true pelvis. It is bounded by the horizontal pubic rami and the symphysis pubis anteriorly, the linea terminalis laterally, and the promontory and alae of the sacrum posteriorly.



Solution to Question 15:

On the second day post-delivery, the level of the uterus lies at two finger breadths below the umbilicus.

24 hours post delivery is equal to Day 0 when uterus is just below the umbilicus.

Day 1 - 1 finger breadth below umbilicus

Day 2 - 2 fingers breadth below umbilicus

The rate of involution of the uterus can be clinically assessed by noting the fundal height of the uterus in relation to the symphysis pubis.

Involution of the uterus occurs due to a decrease in the size of muscle fibers. Following delivery, the fundus lies about 5 1/2 inches above the symphysis pubis. During the first 24 hours, the level remains constant, after which there is a steady decrease in height by half an inch in 24 hours, which approximately equals two finger breadth below the umbilicus on second-day post delivery. By the end of 2nd week, the uterus becomes a pelvic organ. The rate of involution slows down after this until 6 weeks. The uterus almost becomes normal in size by 6 weeks.

Involution of the uterus occurs due to a decrease in the size of muscle fibers.

Time following delivery	Features	Weight of the uterus
Immediately following delivery	Uterus corresponds to 20 weeks gestational age. Fundus lies just below the umbilicus	1000 g
48 hours	Myometrial involution starts	-
1 week	Rapid involution of uterus	500 g
2 weeks	The uterus becomes apelvicorgan	300 g
4 weeks	Clinically,involution iscomplete	100 g
6-8 weeks	Uterus returns to normal (pre-pregnant) sizeSonographically,involution is complete at 8 weeks	60 g

Solution to Question 16:

For a pregnant woman, an additional 300kcal in the 2nd trimester is recommended.

According to Williams Obstetrics, in the first trimester, energy requirements are generally the same as non-pregnant women. In the second trimester, it increases by 340 kcal and in the third, it increases by 452 kcal per day. According to ICMR (Indian Council of Medical Research), an additional intake of 300 Kcal during the second and third trimesters of pregnancy is recommended. Therefore, option A would be a better answer in this case.

Pregnancy is a period where a mother will require additional nutritional requirements to support the healthy growth and development of the fetus. According to the National Institute of Nutrition,

• Protein intake should increase by 23g per day compared to non-pregnant women

• Fat intake should be 30g per day

• Calcium intake should be 1200mg per day

• Iron intake should be 35mg per day

During lactation, ICMR recommends increasing 550 Kcal during the first six months of lactation and 400 Kcal during 7- 12 months of lactation.

Solution to Question 17:

The clinical scenario of mid-trimester cervical dilatation and a short cervix suggests cervical incompetence. Ruptured membranes is a contraindication for cervical cerclage.

Cervical incompetence or insufficiency is a functional or anatomical impairment in the retentive power of the cervix (or internal os). It is characterized
by repeated, mid-trimester painless cervical dilatation. It leads to the escape of amniotic fluid followed by painless expulsion of the products of conception.

Cerclage is the procedure done to strengthen the cervix in patients with cervical incompetence. Contraindications of cervical cerclage include:

• Ruptured membranes

• Intrauterine infection

• Uterine contractions

• Vaginal bleeding

• Cervical dilatation >4 cm

• Fetal death or defect

Cervical incompetence is associated with various risk factors such as prior preterm birth, congenital uterine anomalies, iatrogenic (dilatation and curettage, induced abortion by dilatation and evacuation, vaginal operative delivery through an undilated cervix, amputation of the cervix, etc), and multiple gestations.

In the interconceptional period, diagnosis is made with the passage of number 6-8 Hegar's dilator beyond the internal os without resistance or pain. Investigations for cervical incompetence during pregnancy include:

• Per speculum examination - shows cervical dilatation with herniation of membranes

• Transvaginal ultrasound - funneling of internal os >1 cm and cervical length <25 mm

Prophylactic cervical cerclage is done between 12-14 weeks of gestation. Rescue or emergency cerclage is done when there is bulging or prolapse of membranes through a dilated cervix.

Shirodkar and McDonald's methods of cervical cerclage are the commonly performed cerclage procedures during pregnancy. In McDonald's method, a non-absorbable (mersilene) suture material is placed as a purse-string suture at the level of the internal os, whereas in the Shirodkar method, non-absorbable (mersilene) tape is used. Removal of stitch should be done at the 37th week of gestation. It can be removed earlier in case of preterm labor or if signs of abortion are seen. Laparoscopic abdominal cerclage is used for non-pregnant patients.

Other options:

Option C- Fetal fibronectin in cervicovaginal secretions is a marker for impending preterm labor before membrane rupture. Prophylactic cervical cerclage is indicated in women if ultrasonography

confirms cervical incompetence.

Option D- Advanced maternal age is not a contraindication for cervical cerclage.

Solution to Question 18:

The given features of right occipito-posterior presentation with the favorable pelvis (gynecoid) and adequate uterine contractions can be managed by normal vaginal delivery.

Right occipito-posterior position (ROP) is the most common malposition, which occurs when the occiput is situated posteriorly over the sacroiliac joint, or directly over the sacrum. It is associated with the risk of delayed engagement of the head, prolonged labor due to transverse arrest, and the need for cesarean sections.

Features of occipito-posterior position include infra-umbilical flattening and fetal heart sounds localized to the flank. The fetal parts are more prominently felt near the midline. The head engages through the right oblique diameter in ROP and the engaging AP diameter is occipitofrontal or suboccipito-frontal. In 90% of circumstances with a favorable pelvis and strong uterine contractions without any complications, it undergoes the mechanism like that of normal labor. Hence, we can wait and watch for the progress of normal vaginal delivery.

In some cases, it undergoes non-rotation resulting in oblique posterior arrest. If it undergoes malrotation (anterior rotation of the sinciput), it can cause a persistent occipito-posterior position. Spontaneous delivery can occur as face to pubis. If an arrest occurs in this position, it is termed an occipito-sacral arrest. Incomplete forward rotation leads to a deep transverse arrest.

Indications of cesarean section in occipito-posterior position include the arrest of labor, inadequate uterine contractions, fetal distress, and other obstetrical complications.

During the second stage of labor, if the anterior rotation of the occiput is completed, delivery occurs spontaneously or may require the application of ventouse or low forceps.

Solution to Question 19:

The next step in the management of this pregnancy with breech presentation and a history of previous classical cesarean delivery is a cesarean section at 37 weeks.

A classical casarean section is done by a longitudinal uterine incision. A previous classical cesarean section is considered a high-risk factor for failure of VBAC (vaginal birth after previous cesarean). Hence, it is contraindicated. There is increased risk of uterine rupture (2-9%) in such cases. In some women, classical cesarean scar can even rupture several weeks before term.

Other contraindications for VBAC include:

- ≥2 previous lower segment cesarean section
- Contracted pelvis or cephalo-pelvic disproportion
- Other obstetric or medical complications

• Prior uterine rupture

If the fetal buttocks or legs enters the pelvis before the head, the presentation is known as breech. Vaginal delivery of a singleton breech fetus can be considered. It can be spontaneous breech delivery, partial breech extraction, or total breech extraction. However, cesarean delivery is considered in the presence of any associated obstetric (previous cesarean delivery) of medical complications.

Other options:

Option B: An ultrasound will not change the decision regarding mode of delivery.

Option C: Internal podalic version results in a breech presentation. It is done in case of transverse lie of second baby in twin pregnancy. It is not indicated in singleton pregnancy.

Option D: External cephalic version is a manipulation through the abdominal wall that results in cephalic presentation. It is usally done at 37 weeks as it can revert back to breech position if carried out before. However, it is contraindicated when vaginal delivery is not an option and in cases of previous cesarean delivery such as in this case.

Solution to Question 20:

The given clinical scenario points to a diagnosis of placenta previa which is caused due to placental implantation at the internal os.

Placenta previa is used to describe a placenta implanted in the lower uterine segment. Painless bleeding towards the end of the second trimester or later is characteristic of the condition. It usually occurs without warning and resolves spontaneously only to recur.

There are four types of placenta previa according to the degree of placental expansion to the lower uterine segment.

• Type-I: The lower margin of the placenta reaches the lower uterine segment but not the os.

• Type-II: The placenta does not cover the internal os but reaches its margin.

• Type-III: The placenta covers the closed internal os but does not do so when fully dilated.

• Type-IV: The placenta covers the internal os even when fully dilated.



Predisposing factors include maternal age, multiparity, smoking, and leiomyomas. Previous cesarean deliveries, raised maternal alpha-fetoprotein levels, and assisted reproductive technology also increase the risk for placenta previa.

Management is individualized on a case-to-case basis. If there is no active bleeding during early gestation, close observation is recommended. Elective cesarean section is usually carried out at 38 weeks. Complications of placenta previa include primary postpartum hemorrhage, morbidly adherent placenta, and coagulation defects.

Solution to Question 21:

The given clinical vignette points to the possibility of diagnosis of either an abortion or ectopic pregnancy. The next best step in the management of this patient is to repeat beta-hCG after 48 hours for diagnosis.

The causes of bleeding in early pregnancy include abortion, ectopic pregnancy, and hydatidiform mole. The characteristic ultrasound finding of a molar pregnancy is snowstorm appearance and so it can be excluded.

A transvaginal ultrasound is done initially in such presentations to locate the gestational sac. If neither intrauterine nor extrauterine pregnancy can be confirmed, it is termed pregnancy of unknown location till definite diagnosis.

A cut-off of ≥1500 mIU/mL for β-hCG levels, known as the discriminatory level is used to aid in diagnosis. Serial monitoring of hCG can be used to assess fetal well-being in cases of threatened abortion. If the initial level is below this cut-off, the gestational sac location usually cannot be identified. In such situations, serial β-hCG levels are measured, and the pattern is assessed. If the value is above the discriminatory level, and an intrauterine pregnancy or abortion cannot be confirmed, then ectopic pregnancy has to be suspected.

The doubling time of hCG during early pregnancy is 1.4-2 days. Failure to double the hCG value along with an empty uterus at 48 hours is highly suggestive of ectopic pregnancy. A trilaminar

endometrial pattern can be seen in ectopic pregnancy along with an adnexal mass separate from the ovary.

Once the diagnosis of ectopic pregnancy is confirmed, administration of methotrexate in single or multiple doses can be considered. A dilatation and curettage can be used to confirm the diagnosis.

Other options:

Option C: Measurement of progesterone cannot be used as a reliable diagnostic test. In general, values <5ng/mL point to a nonliving intrauterine or ectopic pregnancy.

Option D: Laparoscopy is the surgery of choice in hemodynamically stable patients for ectopic pregnancy.

Solution to Question 22:

Among the given options, the contraceptive of choice in this lactating mother with heavy irregular bleeding is MALA N.

MALA N is a combined oral contraceptive (COC) that contains ethinyloestradiol (EE2) in a dose of 20–30 mcg and 0.15 mg of Levonorgesterol. It acts by suppressing ovulation by suppression of pituitary hormones FSH AND LH.

COC has the lowest failure rates and helps in controlling polymenorrhea, dysmenorrhea, and menorrhagia. The risk of thromboembolism is established during the first 6 weeks postpartum (puerperal period) with COC and they are known to suppress lactation. In this clinical scenario, since the child is 18 months old (complementary feeds are usually started at 6 months and child is not entirely dependant on breastmilk for nutrition) and the woman is way past the puerperal period, it can be safely administered.

Other options:

Option A: Progestasert is a hormone-containing IUCD that acts by forming a thick mucus plug at the cervix and prevents the entry of sperms. Its usage is limited in cases with heavy unexplained bleeding.

Option B: Copper T is not used in this patient as it might worsen the symptoms of menorrhagia and may result in anemia.

Option D: Norethisterone enanthate depot is an intramuscular injection given once in 2 months. Though it is suitable for lactating women, it has an increased risk of menstrual irregularity and osteopenia and so not preferred in this case.

Solution to Question 23:

The given clinical scenario and the biopsy findings are consistent with the diagnosis of Sertoli cell-only syndrome.

Patients with Sertoli cell-only syndrome typically present between the age of 20 to 40 years with complaints of primary infertility. Diagnosis can be made on testicular biopsy findings. The histopathology of the biopsy specimen typically shows the tubules which are populated by only

Sertoli cells, with a thickening of the tubular basement membrane and absence of germ cells which is characteristic of Sertoli cell-only syndrome.



The below image shows normal histology of testis:



The Sertoli cells line the seminiferous tubules and have receptors for follicle-stimulating hormone (FSH) and testosterone. There are four sperms per Sertoli cell. It produces a testosterone-binding protein that maintains a high level of testosterone within the testis which is necessary for continuous spermatogenesis.

In Sertoli cell-only syndrome, only Sertoli cells line the seminiferous tubules of the testis, and the patients have very low or absent spermatogenesis. There is no feedback inhibition due to absent spermatozoa, which results in increased FSH levels, but testosterone levels and LH levels remain unaffected since Leydig cells are not affected.

Physical examination is normal as this condition presents with infertility without sexual abnormality. Most cases are usually multifactorial and idiopathic. Patients can still reproduce

with assisted reproductive technology.

Other options:

Option B: Testicular atrophy may be characterized by marked hyalinization and thickening of the basement membrane of the spermatic tubules with the absence of spermatids or germ cells. It may occur due to cryptorchidism, atherosclerotic changes of the supplying vessels, cachexia, and end stage of inflammatory orchitis.

Option C: Sex cord-stromal cells are benign and include Sertoli cell tumors and Leydig cell tumors. Leydig cells are characterized by the presence of crystalloids of Reinke.

Option D: Orchitis is characterized by the inflammatory infiltration of neutrophils and may progress to the formation of a frank abscess. The end-stage may lead to atrophic changes.

Solution to Question 24:

The drugs used for medical termination of pregnancy (MTP) in this patient presenting at 7 weeks of gestation are misoprostol and mifepristone.

Mifepristone 200mg is administered orally on day 1. The minimum recommended interval between mifepristone & misoprostol is 24 hours.

24- 48 hours later, misoprostol 800 mcg is administered by the vaginal, sublingual, or buccal route. Clinical confirmation of MTP is done on day 14.

The various methods of termination of pregnancy are:

Misoprostol and Methotrexate regimen can also be used but takes a longer time of action.

The Medical Termination of Pregnancy Amendment Bill, 2020 has extended the period of termination of pregnancy up to 24 weeks for special categories of women, including survivors of rape, victims of incest, and other vulnerable women (like differently-abled women, minors). The upper gestation limit is not applicable in cases of substantial fetal abnormalities.

Timing	Medical methods	Surgical methods
1st trimester(up to 12 weeks)	MifepristoneMifepristone and misoprostolMethotrexate and misoprostolTamoxifen and misoprostol	Menstrual regulationVacuum aspiration (MVA/EVA)Suction evacuation and/or curettageDilatation and evacuation - Rapid methodSlow method
2nd trimester(13 - 20 weeks)	MisoprostolMifepristone and misoprostolDinoprostoneCarboprost	Dilatation and evacuationIntrauterine instillation of hyperosmotic solutionsOxytocin infusionHysterotomy

Solution to Question 25:

hCG (human chorionic gonadotropin) from placenta stimulates the production of testosterone in the male fetus.

hCG released from the placenta is the initial stimulus for testosterone production in the male fetus. Later by the eighth week of gestation, fetal pituitary LH (luteinizing hormone) stimulates the Leydig cells of the testes to produce testosterone.

Sex differentiation of the embryo depends on the Y chromosome, which contains the testis-determining gene called the SRY (sex-determining region on Y) gene. In a genetically male embryo (46, XY), the gonad is converted into testis under the influence of this gene. Female development occurs in the absence of this gene.

Testosterone released from the Leydig cells acts on the Wolffian duct to develop vas deferens, epididymis, and seminal vesicles. Testosterone is converted to dihydrotestosterone (DHT) with the help of 5-alpha-reductase which helps in the differentiation of male external genitalia.

The Sertoli cells of the seminiferous tubules of testes secrete antimüllerian hormone (AMH)/Müllerian-inhibiting substance (MIS) by 7 weeks of gestation. It causes regression of the Mullerian duct and prevents the development of the uterus, fallopian tube, and upper vagina.

Other options:

Option A: It is LH from the fetal pituitary, and not the maternal pituitary that stimulates testosterone production in the male fetus.

Option C: Inhibin A is produced by the corpus luteum and inhibin B is produced by the granulosa cells of the ovary. They inhibit FSH secretion.

Obstetrics and Gynecology NEET 2023

Question 1:

A partogram of a woman who is in labor is shown below. Oxytocin infusion is already given. How will you manage the condition?

a) Forceps-assisted delivery

b) Vacuum-assisted delivery

c) Oxytocin infusion

d) Cesarean section

Question 2:

A woman presents to you at 36 weeks of gestation with complaints of breathlessness and excessive abdominal distension. Fetal movements are normal. On examination, fetal parts are not easily felt and fetal heartbeat is heard but it is muffled. Her symphysis fundal height is 41 cm. Her abdomen is tense but not tender. What is the most likely diagnosis?

a) Abruptio placenta

b) Hydrocephalus of fetus

c) Polyhydramnios

d) Fetal-maternal ascites

Question 3:

For which of the following procedures in the given OT list, can you preferably do a hysteroscopy?

a) Tubal ligation

b) Asherman syndrome

c) Endocervical polyp

d) Subserosal fibroid

Question 4:

A primigravida presents to you with anemia early in her pregnancy. She is 7 weeks pregnant as seen on ultrasound. Her hemoglobin level is 9 g/dL. When should the iron supplements be started for her?

a) 10 to 12 weeks

b) 8 to 10 weeks

c) After 14 weeks

d) After 20 weeks

Question 5:

A 50-year-old woman presents with foul-smelling bloody discharge per vagina mixed with mucous. On examination, a necrotizing growth is seen in the cervix with lateral parametrium involvement. What is the management for this patient?

a) Chemotherapy

b) Brachytherapy

c) Chemoradiation

d) Surgery

Question 6:

A woman is set to undergo a complete laparoscopic hysterectomy. She wants to know if there are any disadvantages to this procedure. What are the disadvantages of laparoscopic over open surgery?

a) Prolonged recovery time

b) Not knowing the extent of thermal burns

c) Increased bleeding

d) Increased pain

Question 7:

A female presents to you with 6 weeks of amenorrhea complaining of bleeding per vagina and slight abdominal pain. The urine pregnancy test is positive and hCG level is 2800 IU/L. A mass is seen on the left adnexa measuring 3 × 2.5 cm. She is hemodynamically stable. How will you manage this patient?

a) Oral methotrexate

b) Single-dose methotrexate injection

c) Serial methotrexate + leucovorin rescue

d) Salpingectomy

Question 8:

A woman comes with complaints of pain and swelling in the perineal area. She also has complaints of difficulty in walking and sitting. She gives a history of multiple sexual partners. On examination, a tender swelling is seen with redness on the labia, as shown in the image given below. What is the most likely diagnosis?



a) Chlamydial infection

b) Bartholin abscess

c) Genital Tuberculosis

d) Herpes infection

Question 9:

A 23-year-old woman accompanied by her mother-in-law comes to the infertility clinic. She has been having regular intercourse for 6 months but is not able to conceive. What is the next best step?

a) Semen analysis for husband

b) Reassure and review the couple after 6 months

c) Hysterolaparoscopy

d) Diagnostic hysteroscopy

Question 10:

A 28-year-old woman with a history of multiple sexual partners presents with lower abdominal pain for 1 month. She experiences minimal discharge and complaints of intermenstrual bleeding. What is the most likely cause?

a) Herpes simplex

b) Genital tuberculosis

c) Candida

d) Neisseria gonorrhoea

Question 11:

A 27-year-old pregnant G3P2L2 woman presents to you at 36+6 weeks. Ultrasound is done and shows the fetus is in a transverse lie. The liquor is adequate, the placenta is normal, and she has no risk factors. Both her previous deliveries were normal vaginal deliveries. How will you manage this patient?

a) Cesarean section

b) External cephalic version

c) Expectant management

d) Induction of labor

Question 12:

A 24-year-old woman who is being treated for infertility with human menopausal gonadotropin came with complaints of sudden abdominal pain, nausea, vomiting, and breathlessness. The finding of the ultrasound is shown below. What is the most likely diagnosis?



a) Ovarian hyperstimulation syndrome

b) Polycystic ovarian syndrome

c) Theca lutein cysts

d) Granulosa cell tumor

Question 13:

During a hysteroscopic removal of a submucosal fibroid, a fluid deficit of 2000 mL is estimated in a patient. What is the immediate complication of this?

a) Acute tubular necrosis

b) DIC

c) Pulmonary edema

d) Thromboembolism

Question 14:

A 54-year-old woman was diagnosed with advanced cervical cancer. She has a 14-year-old daughter. What advice would you give her daughter?

a) Advise HPV vaccine

b) Screen for BRCA mutation

c) Screen for PTEN mutation

d) Perform cervical biopsy

Question 15:

A 17-year-old girl is seen for primary amenorrhea. There is no development of breasts or hair in the pubic or axillary region. Her height is 155 cm and her weight is 48 kg. She has bilateral inguinal masses. On ultrasound examination, the uterus, fallopian tube, and ovary are absent. What is the most likely diagnosis?

a) Complete androgen insensitivity syndrome

b) Hypergonadotropic hypogonadism

c) Turner syndrome

d) Polycystic ovary syndrome

Question 16:

A 16-year-old girl presents with cyclical pelvic pain every month. She has not achieved menarche yet. On examination, a suprapubic bulge can be seen in the lower abdomen. PR examination reveals a bulging swelling in the anterior aspect. What is the most likely diagnosis?

a) Transverse vaginal septum above the vagina

b) Vaginal atresia

c) Imperforate hymen

d) Cervical agenesis

Question 17:

A pregnant woman comes to the clinic. She already has twins by normal delivery. Which of the following is the correct representation of her obstetric score?

a) G3P2

b) G3P1

c) G2P1

d) G2P2

Question 18:

A woman presents to you at 36 weeks of gestation with complaints of feeling lightheadedness and dizziness when she lies on her back. She says she feels alright if she lies on her side or when walks. What is the most likely reason behind this?

a) Increased intracranial pressure

b) IVC compression

c) Heavy meals

d) Excessive venous pooling at the feet

Answer Key

Question No.	Correct Option
1	d
2	c
3	b
4	b
5	c
6	b
7	b
8	b
9	b
10	d
11	b
12	a
13	c
14	a
15	b
16	c
17	c
18	b

Detailed Explanations

Solution to Question 1:

Based on the above partogram, there is a cephalopelvic disproportion, and the management, in this case, would be to do an emergency cesarian section.

The above partogram shows that the cervix is fully dilated, molding is 2+, the station is 3+, and oxytocin infusion is already given. The patient has adequate uterine contractions, as shown. The amniotic fluid is stained with meconium, and there is a fall in fetal heart rate, which indicates that the baby is in distress. This shows that there is an arrest of the fetal head with molding, which points to cephalopelvic disproportion.

In case of uterine inertia or fetal distress, the management is an emergency cesarean section, not instrumental delivery or elective cesarean section.

Cephalopelvic disproportion is a state where the normal proportion between the size of the fetal head to the size of the pelvis is disturbed. It can either occur due to an average size baby with a small pelvis or a relatively large baby with a normal pelvis.

CPD can occur at the inlet, mid-pelvis, or at pelvic outlet. It is diagnosed at these levels by the following measurements.

Inlet CPD

• Obstetric conjugate < 10 cm

• Diagonal conjugate < 11 cm

• Transverse diameter < 12 cm

Midpelvis CPD

• Sum of the interspinous and posterior sagittal diameters of the mid pelvis ≤13 cm

• Interspinous diameter < 8 cm

Outlet CPD

• Interischial diameter ≤ 8 cm.

The following events may occur over the course of labor, depending on the degree of pelvic contraction.

• Early rupture of the membranes.

• Cord prolapse

• Cervical dilatation is slowed.

• Obstructed labor with features of exhaustion, dehydration, ketoacidosis, and sepsis.

• Operative interference, shock, postpartum hemorrhage, and sepsis.

The degree of disproportion must be assessed by clinical examination and supplemented by imaging pelvimetry. The moderate and the severe degrees can be managed by one of the

following:

• Induction of labor

• Elective cesarean section at term

• Trial labor

The elective cesarean section at term is commonly done in cases of a major degree of inlet contraction and in a moderate degree of inlet contraction associated with outlet contraction or complicating factors like elderly primigravida, malpresentation, post-cesarean pregnancy, etc. It is planned during the last week of pregnancy.

Solution to Question 2:

The diagnosis in this case in which symphysis fundal height (SFH) is 41 cm at 36 weeks of gestation and the woman presents with breathlessness and excessive abdominal distension is most likely polyhydramnios.

Polyhydramnios can be defined as the presence of excessive amniotic fluid, which exceeds 2,000 mL.

The sonographic diagnosis of polyhydramnios can be given as

• AFI ≥25 cm

• Deepest vertical pocket (DVP) measuring >8 cm

Causes of polyhydramnios can be due to maternal and fetal conditions-

• Maternal causes include chorioangioma of the placenta, multiple pregnancies commonly in monozygotic twins, disorders of the cardiac or the renal system, and diabetes mellitus.

• Fetal causes include anencephaly, spina bifida, atretic esophagus or duodenum, facial clefts, and neck masses that lead to abnormal swallowing or hydrops fetalis.

Patients may present with dyspnea, palpitations, and pedal edema.

On examination, the following findings can be elicited:

• Markedly enlarged abdomen with flank fullness

• Tense, shiny skin with striae

• Fundal height more than the appropriate gestational age: The SFH measured in cm after 24 weeks is equivalent to the number of weeks up to 36 weeks. A variance of 2 cm is regarded as typical. The SFH in 36 weeks of gestation should have been 36±2 cms, which is less than in this patient.

• Fetal parts and the presentation cannot be well-defined

• Fluid thrill will be present

On per vaginal examination, tensed bulging membranes can be felt. Investigations to confirm the diagnosis include an ultrasound of the abdomen and pelvis and an evaluation of possible causes.

Other options:

Option A - Placental abruption classically presents with sudden onset of abdominal pain, vaginal bleeding, and uterine tenderness. Frequent contractions and persistent hypertonus are the other findings. When the placenta is completely detached from the uterine wall, except at the margin, blood may collect behind the placenta. This kind of presentation is known as concealed type. In another type, blood can escape via the cervix by dissecting downward between the membranes and the uterine wall. It is called the revealed type, which is the most common type.

Option B - Hydrocephalus of the fetus: Hydrocephalus is a condition where there is an abnormal accumulation of cerebrospinal fluid within the brain, leading to increased head circumference in the fetus. Although it may lead to an increased symphysis fundal height, the clinical presentation of breathlessness and excessive abdominal distension in the mother is not typical of this condition.

Option D - Fetal-maternal ascites can present as Mirror syndrome. There is a connection between fetal hydrops and the onset of maternal edema, in which the mother exhibits the same symptoms as the fetus. Because the mother, the placenta, and the fetus all develop edema, the condition is also called triple edema. At least 20% of cases of hydrops have been observed to be complicated by mirror syndrome. Women most commonly present with edema. Other symptoms may include proteinuria, hypertension, the elevation of liver enzymes, and headache.

Solution to Question 3:

Of the following procedures given in the OT list, Asherman syndrome is the condition for which hysteroscopy would be most preferable. Hysteroscopy is used for the definitive diagnosis of Asherman syndrome.

Asherman syndrome refers to the presence of adhesion within the endometrial cavity. It is also called uterine synechiae and can be asymptomatic tiny bands or can completely or almost completely obliterate the endometrial cavity. It can occur as a result of trauma to the endometrium following vigorous uterine curettage, tuberculous endometritis, pelvic radiation, or endometrial ablation.

Depending on the degree of scarring, patients may have amenorrhea, hypomenorrhea, or recurrent pregnancy loss because of inadequate placentation. Hysterosalpingography is performed to assess the uterine cavity and tubal patency. Intrauterine adhesions are characteristically seen as irregular angulated filling defects in the uterine cavity with a classic 'moth-eaten' appearance. Definitive diagnosis (gold standard) and management are done by hysteroscopy. Hysteroscopic transection is the surgical treatment of choice.

Hysteroscopy has the advantage of being a minimally invasive procedure with lower risks and faster recovery times than traditional open surgery. It is often performed as an outpatient procedure, and patients can usually return to their normal activities within a day or two.

Other options-

Option A - Tubal ligation is done to attain permanent contraception in females, which is usually accomplished by occlusion or division of the fallopian tubes. It prevents ovum passage and fertilization. Procedures done at the time of delivery are termed puerperal sterilization. For puerperal sterilization, suture ligation with tubal segment removal is more frequently employed. A tubal sterilization procedure done at a time unrelated to recent pregnancy is called non-puerperal

tubal sterilization or interval sterilization. Currently, non-puerperal tubal sterilization is mainly accomplished via laparoscopy.

Option C - Endocercival polyps are mucinous columnar epithelium-covered benign endocervical stroma overgrowths. They often present as a single, elongated mass that is red, smooth, and projecting from the endocervical canal. Ring forceps are used to grab endocervical polyps for excision if the stalk is thin. The stalk base of the polyp is repeatedly twisted. Surgical ligation and excision of a thick pedicle may be necessary if further severe bleeding is expected. Although hysteroscopy can help visualize and assess these polyps, their removal is typically performed using forceps, either during a pelvic examination or during a colposcopy. Hysteroscopy is not the primary method for managing endocervical polyps.

Option D - Subserosal fibroids carry higher risks for incomplete resection, difficult visualization, or uterine perforation with hysteroscopic myomectomy. Hence, a number of parameters are taken into consideration while choosing a surgical strategy for subserosa or intramural myomas. Successful results and low recurrence rates are seen after laparoscopic leiomyoma excision.

Solution to Question 4:

The iron supplements for the 7-week pregnant mother presenting with anemia should be started at 8 to 10 weeks.

The Centers for Disease Control and Prevention defined anemia in pregnant women using a cut-off value. The pregnant mother is said to be anemic when the level of hemoglobin is less than 11 g/dL in the first and third trimesters and less than 10.5 g/dL in the second trimester.

Anemia in the first trimester is treated by oral iron therapy. According to the Anemia Mukt Bharat program, pregnant women with mild and moderate anemia are treated with two iron and folic acid tablets. Each tablet contains 60 mg of elemental iron and 500 mcg of folic acid. These tablets are taken daily and are supplied by the healthcare worker. Hence, in this mother iron supplementation should be started by 8-10 weeks.

Drawbacks of oral therapy are intolerance presenting as epigastric pain, nausea, vomiting, constipation, or diarrhea, and unpredictable absorption rate. The improvement in hemoglobin levels is generally evident within 3 weeks of oral iron therapy. The hemoglobin concentration is expected to rise at about 0.7 gm/100 mL/week.

Contraindications to oral therapy are intolerance, and severe anemia in advanced pregnancy. The best time to start corrective iron therapy in pregnancy depends on the individual patient's needs and the presence of anemia. For most pregnant women without anemia, it is recommended to start iron supplementation between 12-16 weeks of gestation to meet the increased iron demands during pregnancy.

If a pregnant woman is found to be anemic, the iron supplementation should be started as soon as the anemia is diagnosed, regardless of the gestational age. Early diagnosis and treatment of anemia are crucial to reduce the risk of complications for both the mother and the baby.

In the case of severe anemia or intolerance to oral iron supplements, intravenous iron therapy may be considered, depending on the clinical situation and the patient's response to oral therapy.

Solution to Question 5:

This patient with a cervical tumor involving the parametrium has stage IIB cervical carcinoma, which is treated primarily with chemoradiation.

Stage IIB cervical cancer refers to cancer that has extended beyond the cervix and upper vagina to the parametrial tissue. It has not spread to the pelvic sidewall or the lower third of the vagina. The primary management for stage IIB cervical cancer is concurrent chemoradiation therapy. This involves the combination of external beam radiation therapy, brachytherapy, and chemotherapy (usually with cisplatin). This approach has been shown to improve overall survival and reduce the risk of local and distant recurrence compared to radiation therapy alone.

Following chemoradiation, patients should be closely monitored for treatment response and potential side effects or complications. This includes regular physical examinations, Pap tests, and imaging studies as needed. The prognosis for stage IIB cervical cancer depends on various factors, such as the size of the tumor, lymph node involvement, and the patient's overall health. With appropriate treatment, the 5-year survival rate for stage IIB cervical cancer is around 58-63%.

Other options:

Option A: Chemotherapy alone: Chemotherapy alone is not the primary treatment for stage IIB cervical cancer. It may be used as an adjunct to other treatments, such as radiation therapy, but is not sufficient on its own.

Option B - Brachytherapy: Brachytherapy is a form of internal radiation therapy, in which a radioactive source is placed close to the tumor. While it is an important component of the treatment for cervical cancer, it is typically not used as a standalone treatment.

Option D - Surgery: Surgery is not the primary treatment option for stage IIB cervical cancer. In some rare cases, pelvic exenteration may be performed, but this is typically reserved for patients with recurrent or persistent disease after chemoradiation.

Solution to Question 6:

The disadvantage of laparoscopic over open surgery is not knowing the extent of the thermal burns.

Diathermy burns are a particular hazard in laparoscopic surgery due to the following reasons:

• Relative lack of visibility of the entire instrument.

• Faulty insulation of laparoscopic instruments

• Intraperitoneal contact of the diathermy with another metal instrument while activating the pedal.

• Inadvertent activation of the pedal while the diathermy tip is out of the vision of the camera.

• Retained heat in the diathermy tip touching susceptible structures, such as the bowel.

The advantages of laparoscopic over open surgery are as follows:

• Less postoperative pain (Option D)

• Less blood loss (Option C)

• Cosmetically better

• Earlier recovery and shorter postoperative stay at the hospital (Option A)

• Fewer intraoperative adhesions

• Better perception of anatomy intraoperatively

Solution to Question 7:

The clinical scenario of a patient with 6 weeks of amenorrhea, abdominal pain, and vaginal bleeding with beta-hCG levels of 2800 mIU/mL and a gestational sac in the left adnexa measuring 3 × 2.5 cm points to the diagnosis of unruptured ectopic pregnancy. The most appropriate management for this patient is a single dose of methotrexate injection.

Symptoms and signs of ectopic pregnancy are often subtle or absent in unruptured ectopic pregnancy. The classic triad includes amenorrhea, which is followed by pain and vaginal bleeding. Lower abdominal and pelvic pain is usually seen with tubal rupture. The pain is severe and frequently described as sharp, stabbing, or tearing.

Management of ectopic pregnancy:

1. Medical management is preferred in the following cases:

• Low initial serum beta-hCG level (Less than 5,000 IU/L)

• Small ectopic pregnancy size (Less than 3.5 cm)

• Absent fetal cardiac activity

• Hemodynamically stable

The most popular method of treating ectopic pregnancies is intramuscular methotrexate injection, which can be done in single-dose or multidose protocols. The dose for single-dose therapy is 50 mg/m2 body surface area (BSA).

Single-dose therapy is preferred over the multidose protocol as the former offers simplicity, lower costs, and less involved post-therapy monitoring.

2. Expectant management: A low initial serum hCG concentration, a considerable reduction in levels over 48 hours, and a sonographic inhomogeneous mass instead of a tubal halo or other gestational features are predictive variables for success.

3. Surgical management with salpingectomy or salpingostomy is done when:

• Criteria for expectant and medical management are not met

• Gestational sac >3.5 cm

• Fetal cardiac activity present

• Beta-hCG >5000 mIU/mL

• Significant pain and hemoperitoneum

• Failure of expectant or medical management

• Clinical status of the patient warrants urgent intervention

• Doubtful compliance with medical management

Other options:

Option A: Oral methotrexate is not the preferred route for treating ectopic pregnancy, as it may lead to variable absorption and unreliable dosing. Intramuscular injection is the preferred route of administration for methotrexate in the management of ectopic pregnancy.

Option C: Serial methotrexate with leucovorin rescue is a multidose protocol that can be used to treat ectopic pregnancy. However, the single-dose methotrexate injection is preferred in this case due to its simplicity, lower costs, and less involved post-therapy monitoring.

Option D: Salpingectomy, the surgical removal of the affected fallopian tube, is reserved for patients who do not meet the criteria for medical management or expectant management, have significant pain and hemoperitoneum, or have failed medical management. This patient does not fulfill the criteria for surgical management.

Solution to Question 8:

The given clinical vignette suggests the diagnosis of Bartholin's abscess.

Bartholin glands are the homologs of the male bulbourethral or Cowper glands. They open into the vestibule at approximately the 5 and 7 o'clock positions. The function of the Bartholin gland is to secrete mucus for lubrication during coitus. Bartholin cyst is caused by obstruction of the duct due to infection, trauma, mucus changes, and congenitally narrowed ducts. If the cyst gets infected, it leads to the formation of a Bartholin abscess. Escherichia coli is the most common causative agent, however, other gram-positive and gram-negative organisms can also cause the same.

Most Bartholin gland cysts are small and asymptomatic except for minor discomfort during sexual contact. If the cyst is larger or infected, however, patients may complain of severe vulvar pain during walking, sitting, or coitus. On examination, cysts typically are unilateral, round, or ovoid, and fluctuant or tense. Infected cysts are often erythematous and tender. Most of the cysts produce labial asymmetry.

Small, asymptomatic cysts can be managed conservatively. Large, symptomatic cysts can be managed by incision and drainage, or marsupialization, or Bartholin gland excision. Bartholin abscess can be managed by incision and drainage by word catheter placement or marsupialization. In postmenopausal women with recurrent Bartholin cysts, a suspicion of Bartholin gland cancer should arise and the cyst wall should be sent for biopsy.



Other options:

Option A - Chlamydial infections typically present with vaginal discharge, dysuria, or cervicitis rather than localized pain and swelling in the perineal area. Though Chlamydia can cause infections in the Bartholin gland, the symptoms described in the question are more suggestive of a Bartholin abscess.

Option C - Genital tuberculosis primarily affects the upper genital tract, leading to menstrual irregularities, infertility, and abdominal pain. While it can also involve the vulva and perineal area, the clinical presentation described in the question is not typical for genital tuberculosis.

Option D - Genital herpes usually presents as painful ulcers or vesicles in the genital area, rather than a tender, erythematous swelling. The clinical presentation in the question is more consistent with a Bartholin abscess.

Solution to Question 9:

The next best step would be to reassure and review the couple after 6 months.

The inability to conceive after 1 year of unprotected intercourse of reasonable frequency is described as infertility.

Infertility evaluation is offered to any couple that has failed to conceive after 1 year of regular unprotected intercourse. There is a decline in the conception rate and a rise in poor pregnancy outcomes in older women hence, the majority of experts concur that in women older than 35, evaluation may be considered after 6 months. Since the woman is 23 years old and has tried conceiving for only 6 months, it is advisable to review them in another 6 months if they fail to conceive.

Infertility workup includes the following if required:

(1) Ovulation and evaluation for the diminished ovarian reserve-

(2) Evaluation of Female reproductive tract abnormality-

• Hysterosalpingography

• Sonography

• Laparoscopy

• Hysteroscopy- The preferred way to diagnose intrauterine anomalies is an endoscopic assessment of the uterine cavity. Both an operating room and a doctor's office can be used for hysteroscopy. It has the ability to concurrently diagnose and treat the abnormalities.

(3) Semen analysis

Solution to Question 10:

This clinical scenario indicates infection with Neisseria gonorrhoeae, which has led to pelvic inflammatory disease.

Gonorrhea is a sexually transmitted disease caused by N. gonorrhoeae. It uses pili in adhesion to the urethra or other mucosal surfaces. The cocci penetrate the intracellular spaces to reach the subepithelial tissue by the third day of infection. The incubation period is 2-8 days.

The major symptoms include vaginal discharge, dyspareunia, and dysuria, usually without urinary frequency or urgency. The specific symptoms seen in males and females are as follows:

In females:

About half the female patients remain asymptomatic. But, patients typically present with urethritis, cervical discharge, pruritus, dysuria, and intermenstrual bleeding.

The ectocervix and vagina are not usually affected in adults as the stratified squamous epithelium is resistant to infection by the cocci. The acidic pH of vaginal secretions is also protective. The principal sites of invasion for gonococcal infection are the columnar and transitional epithelium.

The initial infection involves the cervix uteri and urethra which may extend to Bartholin glands, skene glands, endometrium, and fallopian tubes. Unilateral pain and swelling over the labia may be caused by the inflammation of the Bartholin glands.

Pelvic inflammatory disease and salpingitis may lead to sterility. Other complications are tubo-ovarian abscesses and ectopic pregnancies. Rarely peritonitis may develop with perihepatic inflammation (Fitz-Hugh-Curtis syndrome). The patient presents with right upper quadrant pain and tenderness with abnormal liver function tests.

In males:

• Acute urethritis: This infection may extend along the urethra to involve the prostate, seminal vesicles, and epididymis.

• Genital discharge: It is mucopurulent that can be described as that resembling the 'flow of seed' (semen).

• Strictures: Chronic urethritis may lead to stricture formation.

• Water-can perineum: The infection may spread to the periurethral tissues causing abscesses and multiple discharging sinuses.

Other options:

Option A: Herpes simplex presents as painful genital ulcers.

Option B: While genital tuberculosis may present as menstrual irregularity, it is usually accompanied by systemic symptoms such as fever, malaise, and weight loss. Often with concurrent pulmonary disease. It may also present as infertility or may remain asymptomatic.

Option C: Vaginal candidiasis presents as intense pruritus and copious cottage cheese vaginal discharge.

Solution to Question 11:

A pregnant woman G3P2L2 presenting at 36+6 weeks with a transverse lie without being in labor is managed with an external cephalic version (ECV), followed by vaginal delivery.

In a transverse lie, the fetus' long axis is perpendicular to the mother's. The shoulder is typically placed above the pelvic inlet when lying transversely. The head occupies one iliac fossa, and the other by the breech. This results in a shoulder presentation.

Management of transverse lie:

1. Before labor/early in labor - If the membranes are intact (like in this case), it is recommended that ECV should be offered and attempted whenever possible.

Prerequisites for ECV - Women who have reached 370/7 weeks of gestation. Multiparity, unengaged presenting part, nonanterior placenta, nonobese patient, and abundant amnionic fluid improves the chances of ECV.

Absolute contraindications - When vaginal delivery is contraindicated, e.g. placenta previa

Relative contraindications - Early labor, oligohydramnios or ruptured membranes, known nuchal cord, structural uterine abnormalities, fetal growth restriction, multifetal gestation, and prior abruption of the placenta

2. Active labor - A cesarean delivery is frequently necessary for such a scenario (Option A). The lower uterine segment is not occupied by the fetal feet or head in a dorsoanterior or back down position. A vertical hysterotomy incision is usually necessary for such a position. In a dorsoposterior or backup position, one or both feet can be gripped through a small transverse incision and delivered by breech extraction.

Expectant management refers to allowing vaginal delivery to happen. It is chosen when the version fails or is contraindicated. Vaginal delivery may be considered in the case of small fetuses with congenital malformations or dead fetuses (Option C).

Transverse lie (malpresentation) is a contraindication to induction of labor (Option D).

Solution to Question 12:

The above clinical scenario and the sonogram showing enlarged ovaries having large cysts are suggestive of ovarian hyperstimulation syndrome.

Ovarian hyperstimulation syndrome (OHSS) is characterized by multiple ovarian follicular cysts accompanied by increased capillary permeability. It happens most often as a complication of ovulation induction therapy for infertility. It occurs due to the stimulation of vascular endothelial growth factor (VEGF) expression in granulosa-lutein cells by hCG.

They often present with shortness of breath, abdominal pain, and gastrointestinal disturbances like nausea, vomiting, and oliguria.

Abdominal pain occurs due to ovarian enlargement and accumulation of peritoneal fluid. Increased vascular permeability and loss of fluid, protein, and electrolytes into the peritoneal cavity cause ascites and result in hemoconcentration. Hemoconcentration causes hyperviscosity and leads to hypercoagulability.

Hyponatremia, hyperkalemia, leukocytosis, pleural effusion, and pericardial effusion can be associated with OHSS.

Some of the risk factors for OHSS:

• Polycystic ovarian syndrome

• Young age

• High estradiol during ovulation induction

• Pregnancy

This condition can be prevented by

• Using lowered FSH dose to decrease follicular stimulation

• Coasting- withholding FSH administration for 1 or more days prior to the hCG trigger injection,

• Prophylactic treatment with volume expanders,

• Substitution of hCG for FSH during the final days of ovarian stimulation.

OHSS is primarily managed with general supportive care. Paracentesis is performed to ameliorate abdominal discomfort and relieve respiratory discomfort.

Other options:

Option B- Polycystic ovarian syndrome (PCOS) is the most common endocrine disorder of the reproductive age group. Patients with PCOS present with menstrual irregularities, hirsutism, acne, infertility, sleep apnea, and depression/anxiety. These are all short-term consequences of PCOS. The long-term consequences of PCOS include diabetes mellitus, endometrial cancer, and cardiovascular disease.

According to the American Society of Reproductive Medicine (ASRM) and the European Society of Human Reproduction and Embryology (ESHRE), sonographic criteria for PCOS include

• >20 follicles and/or

• Increased ovarian volume (>10 mL), or both

The ultrasound examination image shown below is of PCOS showing enlarged ovaries with multiple cysts.

Option C - Theca lutein cysts are a type of follicular cysts. Hyperreactio luteinalis is a condition in which one or both ovaries develop multiple large theca lutein cysts. They typically occur after the first trimester due to luteinization and hypertrophy of theca interna layer in response to exceptionally high hCG or luteinizing hormone.

They are commonly seen in

• Gestational trophoblastic disease

• Multiple gestational pregnancy

• Fetal hydrops

• Placentaomegaly

Most patients are asymptomatic and the above symptoms are not seen in this condition. Surgical management is only required if it is complicated by torsion or hemorrhage.

The below image shows bilaterally enlarged ovaries with multiple theca luteal cysts as in hyperreactio luteinalis.



Option D - Most women are diagnosed with adult granulosa cell tumors after the age of 30 years, and the average age is roughly 55 years. Heavy, irregular menstrual bleeding and postmenopausal bleeding are frequent and are caused by the endometrium's extended exposure to estrogen. Breast soreness and enlargement are also seen in this condition. Instead of hormones, the mass of the ovary may be the cause of the symptoms. Abdominal pain and distension that is brought on by an expanding tumor that could potentially be hemorrhagic.

Solution to Question 13:

The given clinical scenario suggests the diagnosis of fluid overload, and pulmonary edema is the immediate complication anticipated in this scenario.

In the context of hysteroscopic procedures, the term fluid deficit refers to the difference between the volume of distension fluid introduced into the uterine cavity and the volume of fluid that is removed or accounted for during the procedure. A large fluid deficit indicates that a significant amount of the distension fluid has been absorbed into the patient's bloodstream.

When a large volume of fluid is absorbed during a hysteroscopic procedure, it can lead to fluid overload. This excess fluid in the bloodstream can increase the pressure in the blood vessels, particularly in the pulmonary circulation. The elevated pressure can cause fluid to leak from the small blood vessels in the lungs into the air spaces, resulting in pulmonary edema.

Pulmonary edema can lead to shortness of breath, difficulty breathing, and impaired gas exchange, as the fluid-filled air spaces in the lungs make it difficult for oxygen to pass from the air into the bloodstream. In severe cases, it can lead to respiratory failure and can be life-threatening. Prompt recognition and management of fluid overload during hysteroscopic procedures are crucial to minimizing the risk of complications like pulmonary edema.

Hysteroscopy is an operative procedure that provides the endoscopic view of endometrial cavity and tubal ostia via the cervix. It can be diagnostic or therapeutic. A hysteroscopic image of a uterus with carcinoma is shown below.



Complications of hysteroscopy include :

• Early complications:

• Anesthesia complications- air embolism can occur

• Uterine perforation

• Thermal injury with cautery and laser

• Sepsis

• Allergic reaction to dextran and glycine

• Fluid overload occurs and can lead to pulmonary edema if the fluid deficit is more than 1000 mL. It can be managed by diuretics.

• Cerebral edema

• Cardiac failure

• Late complications:

• Hematometra following cervical stenosis

• Pelvic inflammatory disease

• Uterine rupture during pregnancy

• Dysmenorrhea

• Amenorrhea

Other options:

Option A: Acute tubular necrosis (ATN) is a kidney disorder involving damage to the tubule cells of the kidneys, which can lead to acute kidney injury (AKI). While fluid overload can be a risk factor for AKI, it is not the immediate complication of a fluid deficit of 2000 mL in this clinical scenario.

Option B: Disseminated intravascular coagulation (DIC) is a serious disorder where the proteins that control blood clotting become overactive. It is not directly related to a fluid deficit during a hysteroscopic procedure and is therefore not the immediate complication in this case.

Option D: Thromboembolism refers to a blood clot that forms in a blood vessel and then breaks loose, traveling through the bloodstream and potentially causing a blockage in another part of the body. Although any surgical procedure carries a risk of thromboembolism, it is not the immediate complication related to a fluid deficit of 2000 mL in this scenario.

Therapeutic indications of hysteroscopy	Diagnostic indications of hysteroscopy
Uterine septumAsherman syndromeRetained IUCDPolypectomySubmucous fibroidAbnormal uterine bleedingTubal sterilization using sclerosing agents	To study the endocervical mucosal liningCongenital malformations of the uterusEndometrial tuberculosisAsherman syndromeMisplaced IUCDEndometrial lesionsEndometrial polypCornual tubal blockage

Solution to Question 14:

The patient's 14-year-old daughter should be advised to take the HPV vaccine.

The patient's daughter is not at a higher risk for developing cervical cancer because it does not run in families. The etiology of cervical cancer is related to human papilloma virus (HPV) infection, and immunization against the virus is protective.

There are three types of available HPV vaccines, Cervarix, Gardasil, and Gardasil-9.

Option 2: BRCA mutation causes breast cancer, it is not related to cervical cancer.

Option 3: PTEN mutation on chromosome 10 causes Cowden syndrome.

Option 4: Cervical biopsy is not indicated in this scenario.

	Cervarix	Gardasil	Gardasil-9
Characteristic and HPV subtypes	Bivalent16,18	Quadrivalent 6,11,16,18	Nonavalent 6,11,16,18,31,33, 45,52,58
Protects against	CIN, Ca cervix	Anogenital warts, CIN, Ca cervix	Anogenital warts, CIN, CA cervix, Vulval intraepithelial neoplasia, Vaginal intraepithelial neoplasia
Dose	0.5 mL	0.5 mL	0.5 mL
Administration	0, 1, 7 months IM	0, 2, 6 months IM	0, 2, 6 months IM
Administered gender	Female andmale	Female andmale	Female andmale
Ideal age range (Routine recommendation by CDC)	11–12 years	11–12 years	11–12 years9–15 years - FDA approved
ACIP age recommendation in previously unimmunized women	13–26 years	13–26 years	13–26 years

Solution to Question 15:

The diagnosis based on the given clinical scenario is hypergonadotropic hypogonadism.

The clinical scenario mentions a female with primary amenorrhea with bilateral inguinal masses (undescended testis) and absent ovary pointing to the possibility of XY karyotype. Absent uterus and fallopian tubes point to the presence of a Mullerian inhibiting substance.

Hypergonadotropic hypogonadism can occur in disorders such as FSH or LH receptor mutations. Leydig cell hypoplasia is an autosomal recessive 46, XY disorder. It is caused by inactivating mutations in the LH/hCG receptor that prevents normal hormone

binding. The decrease in the number or function of Leydig cells decreases fetal testosterone production. This leads to the failure of normal male sexual differentiation.

In patients presenting with LH receptor defects, the action of the anti-mullerian hormone is normal; hence, the derivatives of the Mullerian duct are absent. Due to decreased testosterone production, wolffian duct development is impaired and the testes fail to descend. They appear female at birth and at puberty present with primary amenorrhea. They lack breast development and pubic hair development.

Other options:

Option A: Androgen insensitivity syndrome is an X-linked recessive disorder in which the androgen receptors are absent, scanty, or fail to respond to testosterone. In complete AIS, the individual is born as a phenotypically normal female but generally presents with primary amenorrhea. The external genitalia appears normal, with absent or scant pubic and axillary hair. The vagina is shortened and blind-ending with an absent uterus and ovaries. These patients may also present with a history of childhood inguinal hernias, due to the presence of undescended testis in the abdomen or in the inguinal canal. Due to the constant conversion of testosterone into estrogen, these individuals have normal breast development.

Option C: Turner's syndrome presents with a structural abnormality of the second X chromosome or has a mosaic karyotype, such as 45X/46 XX. The clinical features of Turner's syndrome include short stature, short and wide neck, broad chest, widely spaced nipples, primary amenorrhea, congenital heart defects, and renal anomalies. Two X chromosomes are required for the development of ovaries. Losing one X Chromosome results in streak ovaries and inadequate estrogen production.

Option D: Polycystic ovarian syndrome (PCOS) is the most common reproductive age group endocrine disorder. Patients with PCOS present with menstrual irregularities, hirsutism, acne, infertility, sleep apnea, and depression/anxiety. These are all short-term consequences of PCOS. The long-term consequences of PCOS include diabetes mellitus, endometrial cancer, and cardiovascular disease.

According to American Society of Reproductive Medicine (ASRM) and European Society of Human Reproduction and Embryology (ESHRE), sonographic criteria for PCOS include

• >20 follicles and/or,

• Increased ovarian volume (>10 mL), or both

Solution to Question 16:

The given case scenario is suggestive of cryptomenorrhea due to an imperforate hymen.

Cryptomenorrhea is a condition where menstruation occurs but is not visible due to an obstruction of the outflow tract. The menstrual fluid gets collected in the reproductive tract.

Imperforate hymen is a urogenital anomaly. Imperforate hymen results due to failure of canalization of the inferior end of the vaginal plate. This anomaly usually remains unrecognized until puberty. It can lead to mucocolpos in early infancy. Patients present with amenorrhea, cyclic abdominal pain, urinary symptoms, and a hypogastric lump on abdominal examination. A bluish bulging membrane at the vaginal opening is noted on genital examination and the Valsalva

maneuver.

Ultrasound of the pelvis reveals normal internal genitalia and hematocolpos. Laparoscopy should be performed along with hymenectomy to exclude endometriosis.

Most common causes of cryptomenorrhea:

• Intact/imperforate hymen - a bluish convex bulge is seen

• Transverse vaginal septum - a pinkish concave covering is seen

Ultrasound pelvis confirms the diagnosis. A simple cruciate incision of the hymen allows the free drainage of the collected menstrual blood and restores the reproductive capacity in most cases.

Other options:

Option A: Transverse vaginal septum occurs due to the failure of Müllerian duct fusion or failure of vaginal plate canalization. On examination, the vagina may appear shortened and the cervix may not be identifiable. Secondary sexual characteristics appear normal. The diagnosis can be confirmed with ultrasound or MRI.

Option B: Vaginal atresia is characterized by the absence of the lower portion of the vagina with well-differentiated superior structures like the uterus. On examination, the hymenal ring is normal, proximal to which only a vaginal dimple or small pouch is found.

Option D: Cervical agenesis is the congenital absence of the cervix typically with a normal uterus. These patients also lack the upper vagina in most cases. They present with primary amenorrhea and cyclic pelvic pain. If a functional endometrium is present, a patient may have a distended uterus, and endometriosis may develop. The absence of a cervix is seen on examination.

Solution to Question 17:

The obstetric score of a pregnant woman who previously delivered twins would be G2P1.

The obstetric score is used to summarize the past obstetric history as follows:

• G - Gravidity, denoting the total number of pregnancies, including the present one. It is irrespective of the period of gestation.

• P - Parity, denoting the total number of previous pregnancies crossing the period of viability, which is 28 weeks in India.

• This includes both live births and stillbirths

• The current pregnancy is not included

• A - Abortus or the number of pregnancies that were lost for any reason, including induced abortions or miscarriages.

• L - Living children

Both Gravida and para are the numbers of pregnancies rather than babies.

Terminology

• Nulligravida: One who is not and has never been pregnant

• Primigravida: One who is pregnant for the first time

- Multigravida: One who has previously been pregnant
- Nullipara: One who has never completed a pregnancy to the stage of viability
- Primipara: One who has delivered only once (fetuses born alive or dead after becoming viable)
- Multipara: One with 2 or more pregnancies to the stage of viability or more
- Parturient: A woman in labor

Solution to Question 18:

The most likely reason a woman at 36 weeks of gestation complains of feeling lightheadedness and dizziness when she lies on her back but feels alright if she lies on her side or walks is inferior vena cava (IVC) compression.

When the patient is supine during late pregnancy, the gravid uterus has a compressive effect on the IVC. This results in opening up collateral circulation through the means of paravertebral and azygos veins. However, in some cases, the collateral circulation is unable to open up, and the heart's venous return may get severely impaired. Syncope, tachycardia, and hypotension are produced as a result. The patient is rapidly turned to a lateral posture to return the blood pressure to normal. This phenomenon is called supine hypotension syndrome (postural hypotension).

Other options-

Option A- Individuals with elevated intracranial pressure may experience papilledema along with neck stiffness and visual problems. Several herniation syndromes can result from brain matter being forced through the fixed cranial apertures as the mass grows.

Option C- Pregnant women are advised to avoid heavy meals and not to go to bed immediately after the meal. They can cause heartburn, which is common in pregnancy due to the relaxation of the esophageal sphincter.

Option D- The pressure in the femoral vein is noticeably increased, especially in the later months. The gravid uterus puts pressure on the common iliac veins, particularly on the right side due to the dextrorotation of the uterus. In a lying down position during pregnancy, the femoral venous pressure increases from 8 to 10 cm of water to around 25 cm of water, and in a standing position, it increases to approximately 80 to 100 cm of water. This excessive pooling of blood in the lower limbs explains physiological edema during pregnancy. Edema, varicose veins, piles, and deep vein thrombosis are all caused by the extensibility of the veins and blood stasis in the venous system.

Obstetrics and Gynecology INI-CET 2023

Question 1:

A lady on oral contraceptive pills presented with white discharge per vaginum. It is not associated with itching or foul smell and stains her underwear yellow in color. Examination of the discharge revealed normal lactobacilli and acidic pH. What is the most probable diagnosis?

a) Vaginal candidiasis

b) Increased normal vaginal discharge

c) Bacterial vaginosis

d) Chlamydial infection

Question 2:

What is the most common cause of puberty menorrhagia?

a) Anovulation

b) Endometriosis

c) Malignancy

d) Bleeding disorder

Question 3:

SRY region is located in:

a) Short arm of Y chromosome

b) Short arm of X chromosome

c) Long arm of Y chromosome

d) Long arm of X chromosome

Question 4:

A 32 year old female presents with grey-white vaginal discharge and on gram stain of the discharge, clue cells are seen. Clue cells are

a) Degranulated neutrophils (pus)

b) Epithelial cells covered by bacteria

c) Gardenella vaginalis with Bacillin rods

d) Doderlein bacili in the macrophage

Question 5:

Which of the following is a true association with HRT?

a) Decreases risk of Colon cancer

b) Decreased risk of breast cancer

c) Decreased risk of endometrial cancer

d) Increases risk of osteoporosis

Question 6:

All the following can result in non-immune hydrops except

a) ABO incompatibility

b) Parvovirus B19

c) Chromosomal defect

d) Alpha thalassemia major

Question 7:

In a 46XY female complete gonadal dysgenesis was noted. Complete gonadal dysgenesis is caused by what in the SRY gene?

a) Point mutation

b) Deletion of gene

c) Translocation

d) Inversion

Question 8:

Which of the following statements about primary dysmennorhea is true?

a) Pain begins after few hours after onset of periods and can last up to 48 hours

b) It is dependent on psychological factors

c) Pain begins 3 days prior to the onset of periods and is present upto the second day of the period

d) GnRH antagonists are the first line of treatment

Question 9:

In which of the following statements will you suspect primary amenorrhea

a) 1, 2, 3

b) 1, 2, 3, 4

c) 1, 2, 4

d) 2, 3, 4

Question 10:

A 38-year-old multi gravida presents with 2 years of infertility and dyspareunia and dysmenorrhea. She has a retroverted uterus and tenderness in the pouch of Douglas. What would be the next step of management?

a) Diagnostic laparoscopy

b) Laparotomy

c) HSG

d) TVS

Question 11:

A 23-year-old female with 6 years of infertility, has a BMI of 33 and facial hair. Investigations reveal she has (PCOS) and spills from bilateral tubes. She and her husband are trying to conceive, what would be the best plan of management for this lady?

a) Weight loss + Metformin + induction of ovulation with urinary gonadotrophins

b) Weight loss + folic acid, Try for spontaneous induction, if failed move to induction of ovulation

c) Metformin + finasteride + Induction of ovulation

d) Weight loss + Metformin + Induction of ovulation with clomiphene

Question 12:

Which of the following drug is not given in endometrosis?

a) Medroxyprogesterone acetate

b) Cabergoline

c) Estradiol

d) Letrazole

Question 13:

Lady with 28 weeks of gestation presents with fever, acute pain abdomen, and mild leucocytosis. What is the most probable diagnosis?

a) Uterine torsion

b) Infection of fibroid

c) Red degeneration of fibroid

d) Preterm labor

Question 14:

Choose the incorrectly matched option.

a) Uterine prolapse - Vaginal hysterectomy

b) Vault prolapse - Sacrospinous ligament suspension

c) Cystocele - Colposuspension

d) Rectocele - Posterior colporrhaphy

Question 15:

Fructose is absent in the seminal fluid, which of the following statements is true regarding this:

a) Congenital absence of seminal vesicles

b) Obstruction in the ejaculatory duct

c) Both

d) None

Question 16:

A 15-year-old female presents with primary amenorrhea and signs of virilization. She has prepubertal development of breasts and pubic hair and is tanner stage 2. She has a blind-ended vagina and clitoromegaly. She has normal male testosterone levels. Her genotype is 46XY. What is the probable diagnosis?

a) 5 alpha reductase deficiency

b) 17 alpha hydroxylase deficiency

c) Androgen insensitivity syndrome

d) Swyer syndrome

Question 17:

Which one of the following is a virilising tumor of the ovary?

a) Arrhenoblastoma

b) Thecoma

c) Brenners tumor

d) Carcinoid tumor

Question 18:

A 28-year-old primigravida with a low-risk pregnancy presents at 39 weeks + 6 days with labor pains. Her vitals are stable. On PV, her cervix is 3 cm dilated and 60% effaced. The fetal head is at -3 station and membranes are felt as a bag of worms beneath the head. What is the appropriate course of management?

a) Start oxytocin

b) Rupture of membranes and confirm cord prolapse

c) Emergency Caesarean section

d) Wait and watch

Question 19:

A 20-year-old woman visits the OPD with complaints of irregular menses since menarche and increased facial hair. Her BMI is 29 kg/m2 and her Ferriman-Gallwey score was 6. Her testosterone level was 0.36 ng/dl. Her mother is a diabetic on oral hypoglycemic agents. She should be advised the following, except?

a) OCPs

b) OCPs + Anti-androgens

c) Lifestyle modification

d) Counselled about risk of metabolic syndrome

Question 20:

A 34-year-old woman at 30 weeks of gestation presents with complaints of bleeding PV and abdominal pain. She has drenched her clothes and has experienced decreased fetal movements. On examination, her cervix was 6cm dilated and 50% effaced. Her BP was 150/100mmHg, PR - 100 beats/min. Her abdomen corresponded to 34 weeks and she had a tense and tender uterus with regular contractions and no fetal heart sounds were heard. Her urine protein is 2+ and there is no active bleeding. What is the ideal management?

a) Antihypertensives + ARM (Artificial Rupture of Membranes)

b) Antihypertensives + emergency caesarean section

c) Antihypertensives + tocolytics

d) Observation

Question 21:

Which of the following vessels will serve as an alternate source of blood supply to prevent uterine ischaemia, in case the primary artery is ligated in the event of PPH?

a) Ovarian artery

b) Uterine artery

c) Arcuate Artery

d) Round Ligament Artery

Question 22:

Which of the following are responsible for increased blood volume during pregnancy?

a) Action of estrogen and aldosterone

b) Increased fluid retention by the kidneys

c) Both A and B

d) Action of ADH and aldosterone

Question 23:

Which of the following is not required to be a part of the consent for surgery in a patient who has an obstetric score of G6L1A3 with a history of two previous LSCS?

a) Risk of bladder injury during surgery due to PAS

b) Risk of bowel injury during surgery due to PAS

c) Possible need for hysterectomy during surgery due to PAS

d) Possible need for blood transfusion

Question 24:

A 30-year-old nulligravida is on DMPA and she comes 14 days past her due date for her next DMPA. What should be advised to her?

a) Administer DMPA in the next cycle

b) Take DMPA with back up for 7 days

c) Ask to do UPT before taking the DMPA

d) Take DMPA and no back up needed

Question 25:

A 67-year-old female is brought to the OPD in a poor general condition. On examination, she was found to have a right ovarian mass. Her CT scan indicated peritoneal caking. Her CA-125 levels were 1200 IU/L. What would be the management in this case?

a) Debulking surgery followed by chemotherapy

b) Chemotherapy followed by debulking surgery

c) Debulking surgery followed by radiotherapy

d) Staging laparotomy

Question 26:

Which of the following leads to an increased risk for borderline ovarian tumors?

a) Polycystic Ovarian Syndrome (PCOS)

b) Low parity

c) Oral Contraceptive Pill (OCP) intake

d) Functional ovarian cysts

Question 27:

A 28-year-old multigravida comes to the emergency at night with an inversion of the uterus following the delivery of the baby when the nurse waited till the delivery of the placenta. Her BP is 110/70 mmHg. Pulse rate 90 beats/min. What should be the management in this case?

a) Manually remove placenta and then reposition the uterus

b) Start oxytocin infusion

c) Reposition the uterus without removing placenta

d) Shift the patient to OT and reposition the uterus under anaesthesia

Question 28:

The recombinant form of vaccine to prevent cervical cancer has which of the following proteins?

a) L1

b) E6 and E7

c) E1 and E2

d) L2

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	a
6	a
7	a
8	b
9	c

10	d
11	b
12	c
13	c
14	c
15	c
16	a
17	a
18	c
19	b
20	a
21	a
22	c
23	b
24	d
25	b
26	b
27	c
28	a

Detailed Explanations

Solution to Question 1:

In the given clinical scenario history of vaginal discharge with no itching and foul smell along with normal lactobacilli count with acidic pH is suggestive of normal vaginal discharge.

Leucorrhea is defined as an excessive normal vaginal discharge.

The term leucorrhea should fulfil the following criteria:

• The excess secretion is evident from persistent vulval moistness or staining of the undergarments (brownish yellow on drying) or need to wear a vulval pad.

• It is nonpurulent and nonoffensive.

• It is nonirritant and never causes pruritus.

The physiologic basis involved in normal vaginal secretion is dependent on the endogenous estrogen level. With the rising estrogen level, there is abundant secretory activity of the endocervical glands and the superficial vaginal epithelium becomes rich in glycogen

The normal secretion is expected to increase in conditions when the estrogen levels become high. Such conditions are :

• During puberty—Increased levels of endogenous estrogen lead to marked overgrowth of the endo- cervical epithelium which may encroach onto the ectocervix producing congenital ectopy (erosion) → increased secretion.

• During menstrual cycle

• Around ovulation-Peak rise of estrogen → increase in secretory activity of the cervical glands.

• Premenstrual pelvic congestion and increased mucus secretion from the hypertrophied endometrial glands.

• Pregnancy-There is hyperestrinism with increased vascularity. This leads to increased vaginal transudate and cervical gland secretion.

• During sexual excitement, when there is an abundant secretion from the Bartholin's glands.

Other options:

Vaginal candidiasis: This condition generally presents with itching and a white, "cottage cheese"-like discharge. The absence of itching and the nature of the discharge described in the scenario rule out this option.

Bacterial vaginosis: This condition is characterized by a thin, gray, foul-smelling discharge. The fact that the discharge in this case does not have a foul smell rules out this option.

Chlamydial infection: Chlamydia infections can cause a range of symptoms, including abnormal vaginal discharge. However, many individuals with Chlamydia are asymptomatic, and the discharge associated with this condition does not typically stain underwear yellow.

Solution to Question 2:

The most common cause of puberty menorrhagia is anovulatory cycles.

Anovulatory cycles → unopposed estrogen → endometrial hyperplasia → prolonged and heavy periods.

Other causes

Endocrine dysfunction:

• Polycystic ovary syndrome (PCOS

• Hypothyroidism or hyperthyroidism.

Hematological:

• Idiopathic thrombocytopenic purpura

• Von-Willebrand's disease leukaemia.

Pelvic tumors:

• Fibroid uterus

• Sarcoma botryroides

• Estrogen producing ovarian tumor

Pregnancy complications (abortion)

Treatment:

Reassurance and psychological support. Rest and correction of anemia is done.In refractory cases, progestogens, such as medroxyprogesterone acetate or norethisterone 5 mg thrice daily is given till bleed- ing stops. Usually, the bleeding is controlled within 3–7 days. Medication is continued for 21 days. The condition usually becomes normal following 2–3 courses and then normal cycles resume. In emergency, conjugated equine estrogen 20–40 mg IV is given every 6–8 hours. Once the bleeding is controlled, combined oral pills are started.

Regular menstrual cycle will be established once the hypothalamo-pituitary-ovarian axis is matured.

Solution to Question 3:

SRY region is present on the short arm of the Y chromosome(Yp11).

SRY stands for the sex-determining region on Y. It contains the SRY gene, which produces the protein called the testis determining factor. This causes the gonads to develop into the testes. If the SRY gene or Y chromosome is absent, the gonads develop as ovaries.

Solution to Question 4:

Clue cells are epithelial cells with a fuzzy border or granular cytoplasm caused by small gram-negative bacilli adhering on their surface is characteristic of bacterial vaginosis.

Bacterial vaginosis is a polymicrobial infection caused by Mobiluncus curtisii, Gardnerella vaginalis, Mycoplasma hominis, and Haemophilus vaginalis. It has a malodorous (fishy smell) vaginal discharge with minimal vulval irritation.

Amsel's Criteria is helpful in diagnosing bacterial vaginosis when 3 of the following 4 are present

• Thin, white homogenous vaginal discharge

• A pH of vaginal fluid >4.5

• Fishy odor of discharge accentuated by adding 10% KOH-Whiff test

• Presence of Clue cells on microscopy.

Treatment is oral metronidazole 500 mg BD for 7 days. Alternatively, Ampicillin 500 mg or cephalosporin 500 mg bid for 7 days is also effective.





©MARROW

Solution to Question 5:

HRT decreases the risk of colon cancer.

Indications of HRT:

• Therapeutic

• Symptomatic women who suffer from estrogen deficiency

• Gonadal dysgenesis in adolescents

• Prophylactic

• High-risk cases for menopausal complications such as cardiovascular disease, osteoporosis(Option D), stroke, Alzheimer's disease, and colonic cancer

• Premature menopause, spontaneous or following surgery

• Menopause caused by radiotherapy and chemotherapy for cancer, especially alkylating agents

• Women demanding HRT as prophylaxis

Contraindications of HRT:

• Undiagnosed abnormal genital bleeding

• Known, suspected, or history of breast cancer

• Known or suspected estrogen-dependent neoplasia

• Active or prior venous thromboembolism

• Active or recent (e.g., within the past year) arterial thromboembolic disease

• Liver dysfunction or gallbladder disease

• Known hypersensitivity to the ingredients of the estrogen preparation

• Known or suspected pregnancy

• Uterine fibroids as the fibroids may enlarge in size.

Uses of HRT:

• Short term-hot flushes, vasomotor symptoms

• Dyspareunia, libido

• Urethral syndrome

• Long term-osteoporosis

• Cardiovascular disease

• Alzheimer's disease

Risks of HRT:

• Endometrial cancer(Option C)

• Breast cancer(Option B)

• Ovarian cancer

• Thromboembolism

• Lipid profile dysfunction

• Gall stones, liver dysfunction

Solution to Question 6:

ABO incompatibility is not a cause of nonimmune hydrops.

Hydrops is diagnosed by identifying two or more fetal effusions—pleural, pericardial, or ascites—or one effusion plus anasarca. Hydrops can be seen in association with red cell auto immunization, which is termed immune hydrops. Otherwise, it is nonimmune. Pathogenesis of hydrops includes decreased colloid oncotic pressure, increased hydrostatic or central venous

pressure, and enhanced vascular permeability.

Etiology of hydrops fetalis

Cardiovascular

• Structural defects: Ebstein anomaly, tetralogy of Fallot with absent pulmonary valve, hypoplastic left or right heart, premature closure of ductus arteriosus, arteriovenous malformation (vein of Galen aneurysm)

• Cardiomyopathies

• Tachyarrhythmias

• Bradycardia, as may occur in heterotaxy syndrome with endocardial cushion defect or with anti-Ro/La antibodies

Chromosomal

• Turner syndrome (45,X), triploidy, trisomies 21, 18, and 13

Hematological

• Hemoglobinopathies, such as α4 thalassemia

• Erythrocyte enzyme and membrane disorders

• Erythrocyte aplasia/dyserythropoiesis

• Decreased erythrocyte production (myeloproliferative disorders)

• Fetomaternal hemorrhage

Lymphatic Abnormalities

• Cystic hygroma, systemic lymphangiectasis, pulmonary lymphangiectasis

Infections

• Parvovirus B19, syphilis, cytomegalovirus, toxoplasmosis, rubella, enterovirus, varicella, herpes simplex, coxsackievirus, listeriosis, leptospirosis, Chagas disease, Lyme disease

Syndromic

• Arthrogryposis multiplex congenita, lethal multiple pterygium, congenital lymphedema, myotonic dystrophy type I, Neu-Laxova, Noonan, and Pena-Shokeir syndromes

Thoracic Abnormalities

• Cystic adenomatoid malformation

• Pulmonary sequestration

• Diaphragmatic hernia

• Hydro/chylothorax

• Congenital high airway obstruction sequence (CHAOS)

• Mediastinal tumors

• Skeletal dysplasia with very small thorax

Gastrointestinal

• Meconium peritonitis, gastrointestinal tract obstruction

Kidney and Urinary Tract

• Kidney malformations

• Bladder outlet obstructions

• Congenital (Finnish) nephrosis, Bartter syndrome, mesoblastic nephroma

Placental, Twin, and Cord Abnormalities

• Placental chorioangioma, twin-twin transfusion syndrome, twin reversed arterial perfusion sequence, twin anemia polycythemia sequence, cord vessel thrombosis

Other Rare Disorders

• Inborn errors of metabolism: Gaucher disease, galactosialidosis, GM1 gangliosidosis, sialidosis, mucopolysaccharidoses, mucolipidoses

• Tumors: sacrococcygeal teratoma, hemangioendothelioma with Kasabach-Merritt syndrome

Idiopathic

Solution to Question 7:

Amongst the given options, a point mutation in the SRY gene is most likely to result in complete gonadal dysgenesis but this answer remains controversial as some other sources mention SRY deletion is more commonly associated. The most common cause attributable to about 85% of cases is still idiopathic.

Pure gonadal dysgenesis results from a point mutation in SRY/ SRY deletion or point mutations in another gene with testis-determining effects (DAXJ, SF-1, CBX2). This leads to underdeveloped dysgenetic gonads that fail to produce androgens or AMH. This is characterized by a normal prepubertal female phenotype and a normal mullerian system due to absent AMH.

The child usually presents with failure in the development of secondary sexual characteristics at puberty and amenorrhea. Investigations reveal the absence of estrogen production and they have elevated levels of FSH (>40 miU/mL) and low oestradiol levels (<25 pg/mL).

Women with streak ovaries are infertile, but they can bear children with oocyte donation. They must be treated with cyclic estrogen and progestogen to promote feminization and secondary sexual characteristics and prevent osteoporosis.

Solution to Question 8:

Primary dysmenorrhea is dependent on psychological factors is the true statement.

Dysmenorrhea is defined as cramping pain that occurs during menstruation. Low social support, alexithymia, neuroticism trait, long menstrual bleeding, family history of dysmenorrhea, and high-caffeine diet are important risk factors for women with primary dysmenorrhea. It can be classified as:

A. Etiological:

• Primary dysmenorrhea is a type of dysmenorrhoea that is not associated with any pelvic pathology. It affects women in the age group of 18 to 25 years, with ovulatory cycles. It begins just before or with the onset of menstruation. (Not a few hours after as in Option A or not 3 days prior as in option C)

• Secondary (Acquired) dysmenorrhea is associated with the presence of pelvic pathologies such as fibroids, adenomyosis, pelvic inflammatory disease (PID), and endometriosis.

B. Clinical:

• Spasmodic dysmenorrhea is the most common and is characterized by cramping pain on the first and second days of menstruation.

• Congestive dysmenorrhea presents as worsening pelvic discomfort and pelvic pain, a few days before the onset of menses followed by relief of the symptoms. This is seen in PID, IUCD users, varicosities of pelvic veins, pelvic endometriosis, and fibroids.

• Membranous dysmenorrhea is rare and is characterized by crampy pain that occurs with the passage of the endometrium as a cast at the time of menstruation.

Primary dysmenorrhea can be managed with non-steroidal anti-inflammatory drugs (NSAIDs), combined oral contraceptive pills,(Option D), and progestin-only pills. GnRH agonists and androgens are also useful. Surgical management includes hysterectomy and is indicated if not responding to medical management. In secondary dysmenorrhea, treating the cause is needed.

Solution to Question 9:

Primary amenorrhea is suspected in patients who have not menstruated by age 15 or within 3 years of thelarche or has not menstruated by age 14 and shows signs of hirsutism, excessive exercise, or eating disorder. The conventional definition of primary amenorrhea is a failure of the onset of menstruation beyond 16 years of age or failure to attain seconday sexual characteristics by 13 years of age.

Secondary amenorrhea is absence of menstruation for 3 months in a female who was previously menstruating normally.

Causes of primary amenorrhea:

• Hypogonadotropic hypogonadism

• Hypothalamic and pituitary dysfunction

• Kallmann's syndrome

• CNS tumors - craniopharyngioma

• Delayed puberty

• Hypergonadotropic hypogonadism

• Primary ovarian failure

• Resistant ovarian failure

• 17-alpha-hydroxylase deficiency

- Chromosomal abnormalities
- Turner's syndrome (45 X)
- Pure gonadal dysgenesis (46 XX or 46 XY)
- Androgen insensitivity syndrome (testicular feminization syndrome) (46 XY)
- Developmental defect of the genital tract
- Imperforate hymen
- Transverse vaginal septum
- Vaginal atresia/cervical atresia
- Absent uterus in Mullerian agenesis (Mayer-Rokitansky-Küster-Hauser syndrome)
- Others:
- Uterine synechiae (congenital)
- Adrenogenital syndrome
- Cretinism
- Juvenile diabetes
- Malnutrition
- Anemia
- Weight loss
- Tuberculosis

Solution to Question 10:

This patient presenting with symptoms of dyspareunia and dysmenorrhea with a retroverted uterus and tenderness in the pouch of Douglas is most likely suffering from endometriosis. The next best step in management is to perform TVS. Laparoscopy is the gold standard for the diagnosis of endometriosis but it is not done routinely for diagnostic purposes.

Endometriosis is a condition that is characterized by the presence of endometrial tissue in ectopic sites such as:

- Ovary (chocolate cysts- left/right lower quadrant pain)
- Pouch of Douglas (pain during defecation)
- Uterosacral ligaments (chronic pelvic pain & infertility)
- Appendix (right lower quadrant pain)
- Lower segment cesarean section (LSCS) scar endometriosis
- Umbilicus
- Lung (cyclical hemoptysis)

These extra-uterine sites of endometrial tissue are influenced by cyclical changes in the hormone levels which give rise to location-specific symptoms as mentioned above.

Management:

Investigations include:

• USG: TVS will show a chocolate cyst.

• MRI

• CA-125 more than 35 U/mL

• Laparoscopy: Gold standard for the diagnosis and treatment of endometriosis. It is used to locate the sites, take a biopsy and surgically remove the deposits.

Drugs used for symptom control include the following:

• Oral contraceptive pills

• Mirena IUD

• GnRH analogs

• Letrozole

Surgical management includes:

Minimally invasive surgery may be used to destroy the tissue by cautery, laser vaporization, and lyse adhesions that have formed as a complication of endometriosis.

Major surgical measures include incision of chocolate cyst and removal of lining, salpingo-oophorectomy, hysterectomy, and unilateral or bilateral salpingo-oophorectomy.

Solution to Question 11:

In the given clinical scenario the most appropriate management would be weight loss + folic acid, try for spontaneous induction, if failed move to induction of ovulation. In obese PCOS patients, even modest weight loss (5% of body weight) can help in the restoration of ovulatory cycles in some cases. Hence spontaneous induction is tried. Folic acid is normally administered as prophylaxis before conception to prevent neural tube defects.

Polycystic ovarian syndrome

Polycystic ovarian syndrome (PCOS)/ Polycystic ovarian disease (PCOD)/ Stein-Leventhal syndrome is a multisystem disorder. It involves metabolic and endocrine dysfunctions that commonly occur among women in the reproductive age group. It has multifactorial causation, such as lifestyle changes, stress, diet, obesity, and hyperinsulinemic states.

Important clinical features of PCOS are Menstrual abnormalities (70%) - oligomenorrhea, amenorrhea, obesity, hirsutism, acne, and alopecia due to hyperandrogenism, acanthosis nigricans due to insulin resistance, and infertility.

Rotterdam criterion is used to diagnose PCOS. The presence of any two of the three components of this criteria establishes the diagnosis of PCOS.

• Oligo/ amenorrhoea, anovulation, infertility

• Hirsutism/ acne

• Ultrasound findings-

•

Enlarged ovaries with ovarian volume will be more than 10 mL

•

12 or more small follicles each of 2-9 mm in size placed peripherally.

Investigations

• Hormonal studies

• USG is diagnostic of PCOS - it shows enlarged ovaries with multiple follicles in the periphery. It also rules out other pathologies like ovarian tumors, endometrial hyperplasia.

• Laparoscopy (therapeutic). It reveals bilateral enlarged ovarian cysts.

The ultrasound image below shows an enlarged ovary with multiple follicles in the periphery.



Management of PCOD is individualized. Weight reduction and lifestyle changes are the first-line therapy for PCOD. Oral contraceptive pills (OCP), preferably fourth-generation pills with 30 mcg E2 and 2-3 mg drospirenone are used to treat menstrual abnormalities. Hirsutism and acne are managed with OCP with cyproterone acetate and topical agents such as epiornithine, clindamycin lotion 1%, erythromycin gel 2%, and isotretinoin.

Patients with PCOD are more prone to developing insulin resistance which can be controlled with metformin(Option A, C and D) and acarbose. Patients presenting with infertility are treated by inducing ovulation using clomiphene as a first-line drug, along with dexamethasone and letrozole.

Surgical management includes laparoscopic ovarian drilling or puncture of not more than four cysts in each ovary.

Note:

PCOS in association with diabetes, hypertension, hyperlipidemia and cardiovascular disease presenting in reproductive age group is known as "X syndrome".

Solution to Question 12:

Estradiol is not used in the treatment of endometriosis as estrogen administration leads to further development of endometriotic lesions.

Treatment of endometriosis:

1. Medical management:

For minimal disease, the drug of choice is NSAIDs. Oral contraceptive pills are given if associated menstrual complaints are present. OCPs inhibit gonadotropin release and decrease the menstrual flow.

For moderate to severe diseases, oral progestogens are the first drugs to be started. They antagonize estrogenic effects on the endometrium, causing endometrial atrophy. The drug of choice is GnRH agonist.

Endometriosis requires angiogenesis.Cabergoline, a vascular endothelial growth factor pathway inhibitor, is an effective therapeutic option for women with chronic pain due to endometriosis. (Option B)

Mnemonic to remember drugs used: GOLD GAP

• G: GnRH analog(Leuprolide)

• O: Oral contraceptive pills

• L: Letrozole(Option D)

• D: Danazol

• G: Gestrinone

• A: Antiprogesterone-Mifepristone

• P: Progesterone(Medroxyprogesterone acetate)(Option A)

2. Surgical management:

Primary surgery is preferred in infertile women because hormonal treatment would delay pregnancy.

• Conservative surgery: Endometrial deposits are destroyed by electrocautery/laser.

• Laparoscopic uterosacral nerve ablation (LUNA)

• Radical surgery: Total hysterectomy with bilateral oophorectomy.

Endometriosis is the presence of endometrial tissue (glands and stroma) outside the uterus.

Risk factors for the development of endometriosis

• Family history

• Nulliparity

• Prolonged exposure to endogenous estrogen (early menarche, late menopause)

• Estrogen therapy

• In utero exposure to diethylstilbesterol (DES)

Key clinical features

• Dysmenorrhea

• Lower abdominal pain

• Infertility

• Pain during urination or defecation

• Constipation or diarrhea

• Irregular bleeding

The gold standard for the diagnosis of endometriosis has been a visual inspection by laparoscopy, preferably with histological confirmation. Laparoscopy helps detect and diagnose pelvic endometriosis, locate the site of endometriosis and staging, take a biopsy, and perform surgical ablation as a part of treatment.

Other investigations:

• CA-125 levels: raised in endometriosis.

• Monocyte chemotactic protein (MCP-1) levels: raised in the peritoneal fluid

• MRI

• Color Doppler

• Anti-endometrial antibodies



Solution to Question 13:

In the given scenario history of acute pain abdomen with fever and mild leukocytosis in a 28-week pregnant woman is suggestive of red degeneration of fibroid.

Red degeneration is the most common complication of fibroids in pregnancy. It is due to the breakdown of the fibroid resulting in ischemia and pain. Key clinical features in this condition are:

• Low-grade fever

• Tender uterus on palpation

• Elevated leukocytes

• Elevated ESR

The diagnosis is confirmed with an ultrasound. It is usually self-limiting and can be managed with non-steroidal anti-inflammatory drugs.

Other options

Option A - Uterine torsion is defined as rotation of more than 45 degrees around the long axis of the uterus. The most usual symptoms of uterine torsion are birth obstruction, abdominal pain, vaginal bleeding, shock, and urinary and intestinal symptoms. The treatment in the earlier months of pregnancy is immediate laparotomy and detorsion of the uterus. Near-term or during labor cesarean section is carried out, and elimination of the possible etiologic factors.

Option B - Infection is common in submucous and myomatous polyps if they project into the cervical canal or the vagina. An infected polyp can cause blood-stained purulent discharge. Infection is more likely in the postpartum and postabortal state.

Option D -Preterm labor is diagnosed with the help of the following criteria

• Regular uterine contractions with or without pain (at least one in every 10 minutes)

• Dilatation (> 2 cm) and effacement (80%) of the cervix

• Length of the cervix (measured by TVS) < 2.5 cm and funneling of the internal os

• Pelvic pressure, backache and/or vaginal discharge or bleeding.

Solution to Question 14:

The incorrectly matched option is Cystocele-colposuspension. The surgery performed for cystocoele is anterior colporrhaphy.

Various surgeries were performed for the management of uterovaginal prolapse:

• Anterior colporrhaphy: This is done for cystocele or cystourethrocele.

• Posterior colporrhaphy/ colpoperineorrhaphy: This is done for posterior defects like rectocele or rectal defects with perineal laxity.

• Le Forte's repair: This is preferred only in elderly women with advanced prolapse, or those unfit for surgery. Women wanting an active sexual life are poor candidates for this procedure as it creates an adhesive area in the midline.

• Abdominal sling surgeries: Young women desirous of retaining menstrual and reproductive functions are ideal candidates. These surgeries use tapes as slings to prevent organ descent. These include:

• Abdominal wall cervicopexy (modified by Purandare and Mathre) - sling attached to cervix posteriorly near uterosacral ligaments.

• Shirodkar's sling surgery - the sling fixes the cervix to the lumbosacral fascia.

• Khanna's sling surgery - sling fixed to the anterior superior iliac spine, strengthening the Mackenrodt's ligament.

• Fothergill surgery (Manchester operation): An anterior colporrhaphy is combined with amputation of the cervix. The Mackenrodt ligaments are then sutured in front of the cervix followed by a colpoperineorrhaphy. It is suitable for women <40 years desiring reproductive function, but fertility may be reduced.

• Ward-Mayo surgery: This is a combination of vaginal hysterectomy + pelvic floor repair. It is done in women >40 years, not desiring reproductive function.

• Moschowitz surgery: This is performed for enterocele repair where the pouch of Douglas is obliterated.

Management of Vault prolapse:

• If the patient is fit for abdominal surgery: Transabdominal sacral colpopexy, i.e., suspending the vault to the sacral promontory extraperitoneally. It is best suited for younger women since coital difficulty following vaginal surgery is avoided

• If the patient is not fit for abdominal surgery: Right Transvaginal sacrospinous colpopexy. It is usually preferred in obese and elderly women, who are not fit for abdominal surgery.

• Colpocleisis is permissible if sexual activity is not desired

Solution to Question 15:

Fructose is absent in seminal fluid in case of congenital absence of seminal vesicle and obstruction of the ejaculatory duct.

The ejaculatory duct is formed by the fusion of the ductus deferens and the duct of the seminal vesicle. A major portion of the semen volume is contributed by the seminal vesicles. Seminal fluid is alkaline and protects sperm from acidity in prostatic secretions and in the vagina. Seminal fluid also provides fructose as an energy source for sperm. If seminal vesicle is absent or if there is an obstruction at the level of the ejaculatory duct then semen analysis will reveal low ejaculate volume, oligospermia or azoospermia, low pH, and absence of fructose.

Solution to Question 16:

The clinical scenario points towards the diagnosis of 5 alpha-reductase deficiency.

The 5 alpha-reductase enzyme converts testosterone into a more potent form of the male sex hormone called dihydrotestosterone (DHT). Testosterone is required for the development of male internal genitalia, whereas DHT has a role in the development of male external genitalia. At puberty, testosterone is responsible for aggressive psychosexual behavior, deepening of the voice, increased muscle mass, and initiating spermatogenesis.

In 5 alpha reductase deficiency, there is a failure of conversion of testosterone to dihydrotestosterone. As testosterone levels are normal there is a normal development of Wolffian ducts and thus male internal genitalia are normal. However, some cases may have a blind-ending vagina.

As levels of DHT are low the child will have a female-looking external genitalia at birth. These children tend to be raised as females till puberty, by which they start exhibiting virilization which is caused by testosterone. Hence clitoromegaly is seen. There is a lack of breast development as testosterone is not converted to estrogen. Pubic hair is scanty as its development depends on both testosterone and dihydrotestosterone.

Investigations reveal an increase in the ratio of testosterone to DHT after hCG administration.

If the child wants to be raised as a female, then the testes have to be removed before the child attains puberty and virilization of the child. Estrogen therapy that is appropriate for inducing and maintaining feminization should be started at the time of usual pubertal maturation or immediately after gonadectomy is performed in an adult. Surgical correction in the form of vaginoplasty is performed. If the child is raised as a male corrective procedure include hypospadias correction, correction of chordee, and reconstruction of the urethra.

Other options

Option B: 17 alpha-hydroxylase deficiency results in decreased production of testosterone by adrenal glands. However, there will be normal testosterone production from the testes. Hence depending on the testosterone and dihydrotestosterone levels the child may have female-looking external genitalia or ambiguous genitalia at birth. 17 alpha-hydroxylase deficiency also results in a decrease in 17-OH progesterone levels and cortisol levels. Hence there is an increase in ACTH due to the loss of negative feedback which increases aldosterone. Due to an increase in aldosterone, the child will have hypertension and hypokalemia.

Option C: Androgen insensitivity syndrome is an X-linked recessive disorder in which the androgen receptors are absent, scanty, or fail to respond to testosterone. Hence there will be no signs of virilization. The serum testosterone levels will be elevated. There will be normal breast development as testosterone is converted to estrogen.

Option D: Swyer syndrome results from a mutation in the SRY gene on the Y chromosome. This results in dysgenetic gonads. There is no production of AMH and testosterone and hence there is development of female internal and external genitalia. There will be no signs of virilization.

Solution to Question 17:

Arrhenoblastoma (Sertoli-Leydig cell tumor) of the ovary is a virilizing tumor of the ovary.

Sertoli-Leydig cell tumors are a type of sex-cord stromal ovarian tumors. It is most commonly seen in women of reproductive age group. These produce excess androgens and are hence known as virilizing tumors. Menstrual irregularities and signs of virilization like hirsutism and deepening of the voice are common presenting complaints. Microscopy reveals Sertoli cells and Leydig cells.

Other options:

Option B: Thecomas typically produce estrogen (female sex hormone). Excess estrogen production can lead to symptoms such as abnormal uterine bleeding, endometrial hyperplasia, or

endometrial cancer.

Option C: Brenner tumor (transitional cell tumor) is a benign, epithelial, unilateral tumor, resembling a fibroma. Histologically, Walthard cell rests and coffee bean nuclei are seen. Brenner tumors represent only 2-3% of all ovarian neoplasms, and <2% of these tumors are malignant.

Option D: In carcinoid tumors, excessive levels of serotonin is produced which causes flushing, diarrhea, wheezing, damage to the tricuspid valve, etc. The levels of 5-OH indole acetic acid (serotonin metabolite) are elevated in the urine.

Solution to Question 18:

In this patient, the membranes are intact and the 'bag of worms' palpaeted refers to the umbilical cord within the membrane, suggestive of cord presentation. The best course of management would be to take the patient up for an emergency caesarean section.

Cord presentation refers to the presence of the umbilical cord below the presenting part over the internal os, within an intact bag of membranes, whereas in the case of a cord prolapse, the membranes are ruptured and the cord lies externally. The goal of management in the event of a cord presentation is to expedite delivery while keeping the membranes intact, and hence the best option would be to deliver the fetus via emergency LSCS. While the patient is being prepared for surgery, she may be positioned in the exagerrated Sim's position (shown below) to avoid cord compression.



In comparison, the management algorithm of cord prolapse is based on three important factors, which are: living or dead baby, maturity of the baby, degree of cervical dilatation, according to which the following steps are taken:



Other options:

Option A: Oxytocin should not be started as the patient should not be allowed to proceed to deliver via normal vaginal delivery due to the risk of cord prolapse and fetal death.

Option B: The membranes should not be ruptured as this would lead to an overt cord prolapse and complicate the situation further.

Option D: Cord presentation is an obstetric emergency requiring immediate action.

Solution to Question 19:

This patient's history of irregular menstrual cycles and increased facial hair and hirsutism as per the Ferriman-Gellway score are suggestive of polycystic ovarian syndrome (PCOS). Anti-androgens are not recommended in the initial therapy of PCOS.

Hormonal therapy in PCOS is initiated with combined OCPs which are useful in treating menstrual abnormalities, reducing acne, and hirsutism. Anti-androgens arerecommended in PCOS only if patients have no improvement in hirsutism after 6 months of therapy with OCPs alone. Further, a serum testosterone level of 0.36 ng/dl is within the normal range (0.2-0.8 ng/dl) there is no current indication to begin anti-androgen therapy in this patient.

PCOS is an endocrinopathy seen in women of reproductive age, affecting multiple systems. Rotterdam criteria is used to diagnose PCOD. The presence of two of the three components of this criteria establishes the diagnosis:

• Oligo/amenorrhoea, anovulation, infertility

• Hirsutism/acne

• Ultrasound findings:

• Enlarged ovaries with ovarian volume will be more than 10 mm3

• 12 or more small follicles each of 2-9 mm in size placed peripherally

Ultrasound image given below shows an enlarged ovary seen in PCOS with multiple follicles in the periphery.



Patients with PCOS are at risk of developing metabolic syndrome (Syndrome X), characterised by abdominal obesity, high blood pressure, impaired fasting glucose, high triglyceride levels, and low HDL cholesterol levels.

Treatment of PCOD:

• Lifestyle changes (Option C) such as weight reduction, smoking cessation, are recommended and most important in preventing metabolic syndrome (Option D).

• Fourth-generation oral contraceptive pills (OCP) (Option A) containing ethinyl estradiol and drospirenone are used for treating menstrual abnormalities, reducing acne, and hirsutism.

• OCP with cyproterone acetate is prescribed for PCOS patients with hirsutism.

• Patients with PCOD are prone to developing insulin resistance, which can be controlled with metformin. Acarbose can be used as an alternative if metformin is contraindicated.

• Infertility in PCOD patients is treated with clomiphene as the first-line drug for inducing ovulation. Dexamethasone is used in combination to improve fertility rates. Letrozole is used off-label in clomiphene-resistant cases.

Surgical options include laparoscopic ovarian drilling or puncture of not more than four cysts in each ovary. This is done either by laser or by unipolar electrocautery. It restores endocrine levels and improves fertility. Surgery is indicated only if:

• Medical therapy fails

• Hyperstimulation occurs

• Infertile women

• Previous pregnancy losses

Solution to Question 20:

The clinical scenario of painful antepartum haemorrhage associated with pre-eclampsia (hypertension + proteinuria) and fetal death is suggestive of Grade 2 abruptio placentae. In this case, as there is fetal death and the patient is in labor, the best approach would be to administer antihypertensives and deliver the fetus vaginally.

Abruptio placenta is the partial/total placental separation from its implantation site before delivery. The bleeding may be:

• Revealed - blood dissects downwards between membranes and uterine wall to escape through the cervix

• Concealed - blood accumulates between the detached placenta and uterus, and the uterus is larger than expected for the period of gestation

• Mixed - features of both co-existing

Hypertension in pregnancy is the most important risk factor for abruptio placenta. Spasm of the decidual spiral artery leads to endothelial damage and extravasation of blood in the placental bed, leading to fetal distress. Patients often present with sudden onset abdominal pain, vaginal bleeding, uterine tenderness, and frequent contractions. Based on the degree of placental separation, patients are clinically graded as follows:

• Grade 0 – Asymptomatic, abruption is diagnosed only after delivery by the presence of retroplacental clots.

• Grade 1 – Vaginal bleeding, mild uterine tetani. No maternal and fetal complications.

• Grade 2 – With fetal distress or death

• Grade 3 – With maternal complications like shock, renal failure, coagulopathy, or death.

The management of abruption is based on whether the patient is in labor and the fetal status, with the goal being to expedite delivery and prevent complications such as shock and coagulopathy. Immediate management begins with the administration of IV crystalloids and arrangement of blood for the patient, along with BP control.

• If the patient is in labor and the cervix is favourable, in case the fetal heart rate is reassuring, or in case the fetal has died - amniotomy and vaginal delivery.

• If the patient is not in labor or fetal heart tracing indicate fetal distress - an emergency caesarean section is indicated. (Option B)

Other Options:

Options C and D: Abruption is considered an obstetric emergency and hence tocolytics or expectant management have no place in this situation.

Solution to Question 21:

In the event of postpartum haemorrhage (PPH), bleeding occurs from the uterine artery which is the primary vessel ligated, and the inflow from the ovarian artery compensates to prevent ischaemia.

The arterial supply to the uterus is by two main arteries: the uterine artery, which is a branch of the anterior division of the internal iliac artery, and the ovarian artery which is a direct branch of the abdominal aorta. Ligating the uterine artery limits the supply from the internal iliac artery and hence the ovarian artery serves as the alternate blood supply.

The uterine artery runs from its origin towards the cervix, crosses the ureter and then ascends with a tortuous course along the lateral aspect of the uterus up to ovary, where it anastomoses with the ovarian artery. It is based on this anatomical relationship that the order of ligation of vessels to arrest bleeding in atonic postpartum haemorrhage is the uterine artery > ovarian artery > internal iliac artery.

Other Options:

Option C: The arcuate artery is a branch of the uterine artery itself.

Option D: The artery to the round ligament artery is a branch from the ovarian artery.





Solution to Question 22:

Increased blood volume during pregnancy is due to estrogen and aldosterone, both of which cause increased fluid retention by the kidneys, hence both options A and B are correct.

Increased estrogen during pregnancy directly causes fluid retention and also activates the RAAS pathway causing increased aldosterone, leading to fluid retention by the kidneys. Although there is a reset in the threshold for response to ADH, the levels of ADH remain unchanged.

Although all components of blood are increased during pregnancy, plasma volume is increased more than the RBC volume, leading to a relative hemodilution, responsible for what is termed as physiological anaemia of pregnancy. The increased plasma volume is beneficial as it reduces blood viscosity, allowing for effective gaseous exchange in placental and fetal circulation, it is protective against the possible adverse effects of change in posture causing vascular compression, and also against the possible blood loss during delivery.

Solution to Question 23:

A patient with a history of two previous caesarean deliveries would be at a high risk of having a morbidly adherent placenta or placenta accreta spectrum (PAS). However, bowel injury is an unlikely complication of placenta accreta spectrum (PAS) and hence may not be required as part of the consent for surgery in such a patient.

The placenta accreta spectrum is characterized by abnormally implanted, invasive, or adhered placenta. This occurs due to partial or complete absence of the decidua basalis and imperfect development of the Nitabuch layer. The major risk factors include associated placenta previa and previous cesarean delivery as seen in this patient.

There are three variants of this condition based on the depth of trophoblastic growth.

• Placenta accreta, the villi are attached to the myometrium.

• Placenta increta, the villi invade into the myometrium.

• Placenta percreta, the villi penetrate the myometrium into the serosa, with possible adherence to the bladder wall



PAS can be diagnosed in the antepartum period by ultrasound which will reveal loss of normal hypoechoic retroplacental zone between the placenta and uterus, placental vascular lacunae, bridging vessels, and placental bulging leading to distortion of the contour of the uterus. MRI can be performed when the ultrasound is inconclusive. However, ultrasound remains the modality of choice for diagnosis.

Patients with PAS have a significant risk of haemorrhage and shock. Management of this condition therefore involves appropriate multidisciplinary efforts with adequate surgical, anesthetic, intensive care, and blood banking capabilities. Patients are recommended scheduled delivery between 34 0/7 to 35 6/7 weeks of gestation and the risk of hemorrhage necessitating paripartum hysterectomy and blood transufsion, or bladder injury in percreta must be properly communicated.

Solution to Question 24:

DMPA can be given as a repeat dose up to 4 weeks later from the scheduled date (within grace period) after the previous dose. But if the patient is presenting later than that, she should be given a urine pregnancy test before the next dose of DMPA.

DMPA stands for Depot Medroxy Progesterone Acetate. It is an injectable progesterone formulation available for contraception. It acts by:

• Bringing changes in the cervical mucus

• Making the endometrium hostile

• Preventing ovulation

Disadvantages include spotting and intermediate per vaginal bleeding.

DMPA guidelines:

• Injections to be provided every 3 months

• It can be given up to 2 weeks later without any other contraception.

• DMPA can be given as a repeat dose up to up to 4 weeks later from the scheduled date (within grace period) after the previous dose.

Solution to Question 25:

In the given scenario, the patient presents with poor general condition and the presence of peritoneal metastases is suggestive of Stage III ovarian cancer. The preferred management would be to initiate neoadjuvant chemotherapy with three cycles of paclitaxel + carboplatin over three weeks, followed by debulking/cytoreductive surgery, and then continue another three cycles of chemotherapy.

Epithelial ovarian cancers are treated with a multimodality approach. The usual approach in Stage I and II disease where the patient is fit for surgery, is to perform cytoreductive surgery, which also helps in determining the extent of disease, followed by adjuvant chemotherapy (Option A).

However, patients with advanced disease and poor general condition are unfit for anaesthesia and surgery at the time of presentation and have a higher operative risk, hence neoadjuvant chemotherapy prior to surgery is preferred and has been shown to reduce morbidity in such patients.

Other options:

Option C: Ovarian tumors are not radiosensitive and radiotherapy does not form part of the protocol for treatment of ovarian cancers.

Option D: Staging laparotomy is the term used to describe the procedure of surgically exploring patients with early-stage disease and would not be the approach in a patient with advanced disease in whom chemotherapy followed by debulking surgery is indicated.

Solution to Question 26:

Low parity is a risk factor for the development of borderline ovarian tumors.

Borderline ovarian tumors are ovarian epithelial tumors of low malignant potential, defined histologically by atypical epithelial proliferation without stromal invasion. They are classified as either serous or mucinous. The following are the risk factors for their development:

• low parity / unopposed estrogen

• infertility

• obesity

• smoking

Clinically patients present with symptoms similar to that of a benign ovarian tumor such as a mass per abdomen, abdominal pain and alteration in menstrual cycles. Patients require similar investigations but the diagnosis of borderline ovarian tumor is based on biopsy. Conservative surgery is usually the preferred treatment approach especially in patients who wish to retain fertility. Appendectomy is usually recommended in the mucinous variety to prevent pseudomyxoma peritonei. Patients do not require chemotherapy or radiotherapy for treatment but these tumours tend to recur and hence regular follow-up is advised.

Solution to Question 27:

This patient with a uterine inversion with stable vitals should undergo manual reposition of the uterus without removing the placenta

Uterine inversion occurs when there is excessive traction leading to the loss of the uterine convexity and protrusion of the uterus outside the vaginal introitus.

Risk factors for uterine inversion:

• Fundal attachment of the placenta

• Uterine atony

• Attempts at cord traction before placental delivery

• Adherent placenta

There are 3 degrees of uterine inversion:

• First-degree - the fundus of the uterus inverts into the uterine cavity

• Second-degree - the uterine fundus protrudes through the cervix and lies in the vagina

• Third-degree - the whole of the uterus is inverted and protrudes through the introitus

Since uterine inversion presents with profuse bleeding and a severe degree of shock, it requires prompt diagnosis and management. It is managed by promptly repositioning the uterus by exerting a firm and constant pressure on the inverted uterine fundus. The placenta should not be removed until after the replacement of the uterus if it is still attached to the uterus.

Solution to Question 28:

The recombinant HPV vaccine use virus-like particles (VLPs) that consists of the HPV L1 major capsid protein. The L1 protein self-assembles into VLPs when expressed in eukaryotic cells.

These VLPs contain the same epitopes as the HPV virion. However they do not contain genetic material and cannot transmit infection. The immunogenicity of the HPV vaccines relies on the development of conformational neutralizing antibodies to epitopes displayed on viral capsids.

The nonavalent HPV vaccine works against strains 6, 11, 16, 18, 31, 33, 45, 52 and 58. It is usually commercialized in the name of Gardasil-9. It has received expanded indication which now includes the prevention of oropharyngeal and other head and neck cancers.

As per the December 2022 WHO updated dose schedules on HPV vaccines, the recommendations for different age groups are as follows:

• 1 or 2 doses schedule for the primary target of girls aged 9–14 years.

• 1 or 2 doses schedule for young women aged 15–20 years.

• 2 doses with a 6-month interval for women older than 21 years.

•

A minimum of 2 doses and when feasible 3 doses remain necessary for those known to be immunocompromised and/or HIV-infected.

Obstetrics and Gynecology INI-CET 2024

Question 1:

Which of the following is contraindicated in PPH in a primigravida with chronic hypertension?

a) Methyl ergometrine

b) Misoprostol

c) Oxytocin

d) Prostaglandin F2 alpha

Question 2:

A 16-year-old female came with complaints of primary amenorrhoea. She had normal secondary sexual characteristics and breast development corresponded to Tanner stage IV. Her FSH - 5.3 and LH - 12.1. Her karyotype is 46XX. What is the probable diagnosis?

a) Kallmann syndrome

b) Androgen insensitivity syndrome

c) Gonadal dysgenesis

d) Mullerian agenesis

Question 3:

Which of the following combinations is not seen in the case of twin pregnancy?

a) Monochorionic monoamniotic (MCMA)

b) Dichorionic diamniotic (DCDA)

c) Monochorionic diamniotic (MCDA)

d) Dichorionic monoamniotic (DCMA)

Question 4:

An 18-year-old girl was brought by her mother with complaints of primary amenorrhea and acute retention of urine. She had cyclical abdominal pain and on examination, a supra pubic

mass and a tense bluish bulge were seen in her vagina. What is the most appropriate cause of her symptoms?

a) Imperforate hymen

b) Transverse vaginal septum

c) Vertical vaginal septum

d) Mullerian agenesis

Question 5:

In the quadruple marker test, a 33-year-old G2P1L1 at 18 weeks of gestation was found to be at a high risk of neural tube defects and low risk of trisomy 21. What is the next investigation to be done?

a) Amniocentesis

b) Chorionic villus sampling

c) Non invasive prenatal testing

d) Anomaly Scan

Question 6:

An 18-year-old came with complaints of pain in the abdomen and bleeding PV for the last 2 days. Her last menstrual period was 6 weeks back. On examination, she has hypotension, tachycardia and cervical motion tenderness. Ultrasound of the abdomen shows free fluid in the pouch of Douglas and paracolic gutters. What is the most likely diagnosis?

a) Placenta previa

b) Abruptio placenta

c) Ruptured ectopic pregnancy

d) Missed abortion

Question 7:

A 32 year old primigravida with chronic hypertension came to the hospital at 36 weeks of gestation with headache and blurring vision. Her BP is 170/110 mm Hg. Urine analysis reveals urine albumin 3+. What is the next step to be done?

a) Injection labetalol and magnesium sulphate and expedite the delivery

b) Injection labetalol , antenatal steroids and immediate termination of pregnancy

c) Injection labetalol and magnesium sulphate and deliver at 37 weeks

d) Injection labetalol to control her blood pressure

Question 8:

A female at 40 weeks of gestation presents with labor pains and is in the second stage of labor for the last 2 hours. On examination of the abdomen, there are 4 uterine contractions per 10 minutes each lasting more than a minute. On PV examination, the fetal station is at +3 and the suture is in AP diameter and caput +. The fetal heart rate is 70 beats/ minute in the last 20 minutes. Which is the appropriate method of delivery?

a) Outlet forceps

b) Ventouse delivery

c) Mid cavity forceps

d) Low cavity forceps

Question 9:

Which of the following hormones is not responsible for the in-utero growth of the baby?

a) Insulin

b) Growth hormone

c) Insulin like growth factor

d) Thyroxine

Question 10:

A G2P1L1 delivered a baby via full-term vaginal delivery. The baby is 3 kgs and is active and plays well. Her blood group was found to be O negative and her husband's blood group was found to be B positive. She was given anti-D at 28 weeks of gestation. Which of the following is the appropriate situation to administer postpartum anti D prophylaxis?

a) Baby is O positive and DCT is negative

b) Baby is O negative and DCT is negative

c) Baby is O negative and DCT positive

d) Baby is O positive and DCT positive

Question 11:

What is the correct dose of administration of antenatal steroids for lung maturity?

a) Betamethasone 12 mg 12 hours apart 4 doses

b) Dexamethasone 6 mg 12 hours apart 4 doses

c) Betamethasone 6mg 12hrly 4 doses

d) Dexamethasone 12mg 24hrly 2 doses

Question 12:

Which of the following is not a criterion to diagnose IUGR?

a) Estimated fetal weight < 3rd percentile

b) Abdominal circumference less than 3rd percentile

c) Absent diastolic flow in umbilical artery

d) Uterine artery PI < 95th percentile

Question 13:

Early onset pre-eclampsia is defined before:

a) 28 weeks

b) 30 weeks

c) 32 weeks

d) 34 weeks

Question 14:

Dr Priya is conducting a talk on the screening of cervical cancer with a Pap smear. She works in a clinic in a low-resource setting and she was planning to screen a woman in her 20s. Which of the following is true regarding cervical cancer screening guidelines in her workplace?

a) Pap smear every 3 years, starting at 21 years

b) Co-testing (pap and Hr HPV) every 5 years starting at 25 years

c) Hr HPV testing every 5 years starting at 30 years

d) Pap smear every 5 years starting at 21 years

Question 15:

26-year P1L1 presents with abdominal pain. She has a past history of rheumatic heart disease for which she has undergone mitral valve replacement surgery and is on the oral tablet warfarin 3mg OD. On examination, an abdominal mass corresponding to 14 weeks was felt. On USG, an 8*6 cm unilocular cyst with a reticular pattern was observed in the left ovary. What is the next line of management?

a) Measure CA-125

b) Repeat USG after 6-12 weeks

c) Surgical intervention

d) MRI pelvis

Question 16:

When is delayed cord clamping done?

a) Term and preterm babies irrespective of birth weights

b) Vigorous term and preterm babies requiring resuscitation

c) Vigorous term babies requiring resuscitation

d) All term babies born at 37 weeks

Question 17:

A couple presents with infertility even after 2 years of unprotected intercourse. The seminal analysis reveals a volume of 2 mL, concentration of 40 million/ml, progressive motility of 55%, and morphology of 45%. The wife's hormone levels were normal with bilateral spillage of dye from both fallopian tubes on HSG. What percentage of couples can be attributed to unexplained infertility?

a) 5%

b) 30%

c) 15%

d) 25%

Question 18:

Which of the following is not a physiological change seen in pregnancy?

a) Increase in systolic and diastolic blood pressure after 6 weeks

b) Peripheral resistance to insulin leading to hyperinsuliemia

c) Fasting hypoglycemia and postprandial hyperglycemia

d) Increased plasma volume and RBC mass

Question 19:

Instrument assisted delivery is not indicated in which of the following?

a) Fetal distress

b) Maternal exhaustion

c) Prolonged second stage of labour

d) Cephalo pelvic disproportion

Question 20:

In which stage of meiosis is the primary oocyte arrested?

a) Diplotene

b) Leptotene

c) Zygotene

d) Pachytene

Question 21:

Restitution is best described as:

a) Passive movement of head due to untwisting of neck

b) Movement of the head by 1/8th of the circle due to torsion

c) Stretching of the vulval outlet by the head

d) External rotation after delivery of shoulder

Question 22:

Which of the following is NOT part of AMTSL?

a) Uterine massage

b) Delivery of placenta by controlled cord traction

c) Bimanual compression

d) Administration of oxytocin after shoulder delivery

Question 23:

A 35-year-old P2L2 woman presents with malodorous vaginal discharge and a normal cervix. The whiff test is positive. What is this condition?

a) Candida

b) Chalmydia

c) Bacterial vaginosis

d) Pelvic inflammatory disease

Question 24:

A 14-year-old came with complaints of acute urinary retention. She has not attained menarche and has cyclic abdominal pain. On genital examination, a bluish bulge is seen. What is the probable diagnosis?

a) Transverse vaginal septum

b) Imperforate hymen

c) MRKH

d) OHVIRA

Question 25:

Which of the following does not constitute a germ cell tumor?

a) Yolk cell tumor

b) Dysgerminoma

c) Juvenille granulosa cell tumor

d) Non gestational choriocarcinoma

Question 26:

Which of the following is the smallest pelvic diameter at the brim?

a) Obstetric conjugate

b) Diagonal conjugate

c) True conjugate

d) Anatomical conjugate

Question 27:

A female with abdominal pain and bleeding per vagina with 3 months of amenorrhea. On examination her BP was 90/60mmHg. Which of the following statements is/are true about ectopic pregnancy?

a) 1, 2

b) 3,4

c) 1,2,3

d) 1,2,3,4

Question 28:

A 32-year-old woman, G2P1L1, comes for a quadruple test at 16 weeks. Which of the following is not included in the quadruple test?

a) Dimeric inhibin B

b) Unconjugated estradiol

c) AFP

d) Beta hcg

Question 29:

An athlete with a BMI of 17.5 kg/m2 presentes with secondary amenorrhea and primary infertility. Her laboratory investigations reveal FSH 2 mIU/ml; LH 0.32 IU/L; TSH 1.2mIU/ml. Prolactin 20ng/ml. What is the underlying cause of this condition?

a) Primary ovarian failure

b) Hypogonadotrophic Hypogonadism

c) Hypergonadotrophic Hypogonadism

d) Normogonadotrophic Normogonadism

Answer Key

Question No.	Correct Option
1	a
2	d
3	d
4	a
5	d
6	c
7	a
8	a
9	b
10	a
11	b
12	d
13	d
14	a
15	b
16	a
17	d
18	a
19	d
20	a
21	a
22	c
23	c
24	b
25	c
26	a
27	c
28	a
29	b

Detailed Explanations

Solution to Question 1:

Methylergometrine is contraindicated in the management of postpartum hemorrhage in a female with chronic hypertension.

Methylergometrine is an ergot derivative, which acts directly on the myometrium. It excites uterine contractions which come so frequently one after the other with increasing intensity that the uterus passes into a state of spasm without any relaxation in between. Hence, it should be used only be used either in the late second stage of labor(after the delivery of the anterior shoulder) or following the delivery of the baby.

Indications :

• Prophylactic - Active management of the third stage of labor as prophylaxis to excess bleeding following delivery

• Therapeutic - To stop the atonic uterine bleeding, following delivery, abortion or expulsion of hydatidiform mole. It is used at a dose of 0.2mg im/iv, repeated every 2-4 hours.

Side effects :

• Nausea and vomiting.

• Because of its vasoconstrictive action, it may precipitate a rise in blood pressure, myocardial infarction, stroke and bronchospasm.

• Repeated use may lead to gangrene of the toes due to its vasoconstrictive effect.

• Repeated use in puerperium may interfere with lactation by lowering prolactin levels.

Contraindications :

• Multiple pregnancy - If given accidentally with the delivery of the first baby, the second baby is compromised by the tetanic contractions of the uterus.

• Heart disease - Results in sudden squeezing of blood from the uterine circulation into the systemic circulation causing overloading of the right heart and failure.

• Hypertensive disorders, pre-eclampsia and eclampsia

• Rh-negative mother - More risk of fetomaternal microtransfusion.

• Vascular disease

Other options :

Option B - Misoprostol is a methyl ester of PGE1, used in the induction of labor and in the management of postpartum hemorrhage. Side effects include fever, tachysystole (hyperstimulation), fetal heart rate changes and meconium passage. It should not be used for cases with previous cesarean birth because the risk of uterine rupture is high.

Option C - Oxytocin is used in the active management of the third stage of labor, augmentation of labor and control of postpartum hemorrhage. Contraindications for its use are hypovolemic states, obstructed labor and cardiac disease.

Option D - Prostaglandin F2 alpha derivative (carboprost) is used in the management of postpartum hemorrhage. Contraindications for its use are asthma, active cardiac, renal, or hepatic

disease.

Solution to Question 2:

The given clinical scenario is that of a female with primary amenorrhea, preservation of secondary characteristics, and normal levels of FSH and LH, with a karyotype of 46XX. These are features of Mullerian agenesis.

Congenital absence of both the uterus and vagina is termed Mullerian agenesis or Mayer-Rokitansky-Küster-Hauser syndrome. Females have a shallow vaginal pouch (1 to 2 inches deep). The uterus, cervix, and upper part of the vagina are absent. Typically, normal ovaries persist, as they originate from the genital ridge. Individual presents with primary amenorrhea, have a normal female appearance, secondary sexual characteristics, and 46 XX karyotype. Most patients with Mullerian agenesis have only small rudimentary Mullerian bulbs without endometrial activity.

With Mullerian agenesis, traditional conception is not possible, but pregnancy may be achieved using oocyte retrieval, fertilization, and gestational surrogacy.

Individuals with Mullerian agenesis or hypoplasia have associated congenital renal or skeletal anomalies. Common renal anomalies include ectopic, horseshoe or pelvic kidney and collecting system anomalies. Skeletal malformations include spina bifida, sacralization, sacral bone lumbarization, and cervical vertebral anomalies.

Other options:

Option A - Kallmann syndrome presents with primary amenorrhea due to hypogonadotropic hypogonadism from deficiency of GnRH. The FSH and LH levels are lower than normal, leading to poor development of secondary sexual characteristics. It is associated with olfactory dysfunction and anosmia.

Option B - Androgen insensitivity syndrome occurs due to a defect in the androgen receptor in a genotypic male (XY). These individuals are phenotypic females with no male or female internal genitalia. They have normal female external genitalia with an absent or shallow vagina, well-developed breasts, and scanty body hair (axillary and pubic) and present with primary amenorrhea.

Option C - Gonadal dysgenesis in the form of 45XO (Turner syndrome) or 46XY (Swyer syndrome) presents with poor development of secondary sexual characteristics. Since the gonads are nonfunctional, there is a loss of negative feedback with increased levels of FSH and LH.





Solution to Question 3:

Dichorionic monoamniotic (DCMA) combination is not seen in twin pregnancies.

The table below describes the types of amnion and chorion seen in different types of twins, depending on the time of division.

In monochorionic diamniotic gestation, the twins are separated only by the two amniotic membranes, with no intervening chorion. So, a thin 'T sign' is formed at the point at which the amnions meet the placenta. It is best seen in the latter part of the first trimester or at the beginning of the second trimester. In monochorionic monoamniotic gestation, there is an absence of a dividing membrane.

Time of division after fertilization	Type of twin
0 - 4 days (morula)	Dichorionic diamniotic (DCDA)
4 - 8 days (blastocyst)	Monochorionic diamniotic (MCDA)
8 - 12 days (implanted blastocyst)	Monochorionic monoamniotic (MCMA)
>13 days (formed embryonic disc)	Conjoined twins



In dichorionic gestation, the twins are separated by amniotic membranes and an intervening chorion. So, a thicker twin peak sign or lambda (λ) sign is formed at the point at which the amnions meet the placenta. It is best seen in the first trimester.

Solution to Question 4:

The features of primary amenorrhea, cyclical abdominal pain, acute retention of urine along with a tense bluish bulge make the diagnosis of imperforate hymen more likely.

Imperforate hymen is a type of Mullerian defect, of urogenital origin. Failure of canalization leads to the formation of mucocolpos; this may be recognized in early infancy. However, the anomaly often continues unrecognized until puberty. It presents with amenorrhoea in the presence of secondary sexual characters, cyclic abdominal discomfort, retention of urine and a suprapubic bulge caused by hematocolpos. Examination of the external genitalia reveals the presence of a telltale bluish bulging membrane in the region of the hymen.

A simple cruciate incision followed by an excision of the hymenal tags allows the drainage of the retained menstrual blood. The operation should be performed under aseptic conditions and under an adequate antibiotic cover to avoid any ascending infection.

Other options :

Option B - Transverse vaginal septum is a much rarer condition that occurs due to faulty fusion or canalization of the urogenital sinus and the Müllerian ducts. The majority of the cases involve the upper portion of the vagina. It can be differentiated from imperforate hymen by the presence of a short vagina and the absence of a bulging membrane.

Option C - Vertical vaginal septum occurs when the distal parts of the Müllerian ducts fail to fuse. On examination, there is a longitudinal anteroposterior septum which may be partial or complete, extending right down to the vaginal outlet. It may be associated with a double uterus and double cervix.

Option D - Mullerian agenesis is characterized by congenital absence of the uterus and vagina (small rudimentary uterine bulbs are usually present). The female presents with primary amenorrhea and preservation of female secondary sexual characteristics.

Solution to Question 5:

For a female with a quadruple test showing a low risk of trisomy 21, the next investigation to be done is an anomaly scan, as part of routine antenatal care.

Prenatal genetic counseling involves screening and diagnostic methods. The various screening methods are as follows

The values of the parameters in trisomy 21 are as follows :

PAPP-A - low, MSAFP - low, unconjugated estriol - low, NT - increased, Beta hCG - high, inhibin A - high.

If the screening tests report a high risk of aneuploidy, then the confirmatory diagnostic test is performed, depending on the trimester.

• Chorionic villus sampling - Transcervical - 10-13 weeks, transabdominal - 10 weeks to term

• Amniocentesis - After 15 weeks

• Cordocentesis - 18-20 weeks

Since the screening test report in the question mentions a low risk of trisomy 21, the female should be followed up with an anomaly scan, which is performed as a part of the antenatal care at 18-20 weeks. It gives a detailed survey of fetal anatomy, placental localization and the integrity of the cervical canal.

Other options :

Option A - Amniocentesis is a confirmatory karyotyping test performed if the initial screening test reveals a high risk of aneuploidy.

Option B - Chorionic villus sampling (CVS) is also a confirmatory test, hence done if there is a high risk of aneuploidy. Moreover, CVS is not performed at 18 weeks.

Option C - Non-invasive prenatal testing utilizes cell-free fetal DNA in the maternal blood. However, it is a screening test. Since the female here has already undergone a screening test, there is no benefit is subjecting her to a second screening test.

Dual test (1st trimester)	Triple test (2nd trimester)	Quadruple test(2nd trimester)
1. Beta HCG2. PAPPA (Pregnancy-associated plasma protein A)This can be combined with nuchal translucency by USG	1. Beta HCG2. Unconjugated estriol3. MSAFP	1. Beta HCG2. Unconjugated estriol3. MSAFP4. Serum inhibin A

Solution to Question 6:

The given clinical scenario with a triad of abdominal pain, amenorrhea and vaginal bleeding, along with the ultrasound findings of free fluid, is suggestive of a ruptured ectopic pregnancy.

An ectopic pregnancy is one in which the fertilized ovum is implanted and develops outside the normal endometrial cavity, with the Fallopian tubes being the most common site. The outcomes of such a pregnancy are varied with tubal rupture and massive intraperitoneal hemorrhage being one of them.

Clinical features :

• The incidence is maximum between the age of 20 and 30 years, the maximum fertility period. The onset is acute. The classic triad of symptoms is - abdominal pain (100%), preceded by amenorrhea of 6-8 weeks(75%), and lastly, the appearance of vaginal bleeding (70%).

• Abdominal pain is the most constant feature. Pain is located in the lower abdomen. Shoulder tip pain (25%) (referred pain due to diaphragmatic irritation from hemoperitoneum) may be present.

• Vomiting, fainting attack. Syncopal attack (10%) is due to reflex vasomotor disturbances following peritoneal irritation from the hemoperitoneum.

• Pallor and features of shock - feeble pulse with tachycardia, hypotension, cold clammy extremities.

• Pelvic examination - Vaginal mucosa is blanched white; uterus seems normal in size or slightly bulky; extreme tenderness on fornix palpation(75%).

Investigations :

Estimation of ß-hCG - The findings suggestive of ectopic are a lower concentration of hCG compared to normal intrauterine pregnancy, and doubling in plasma failing to occur in 2 days.

Sonography: Transvaginal sonography (TVS) is more informative. The diagnostic features are

• Absence of intrauterine pregnancy with a positive pregnancy test.

• Fluid (echogenic) in the pouch of Douglas.

• Adnexal mass clearly separated from the ovary (Blob sign).

Color Doppler sonography can identify the placental shape (ring-of-fire pattern) and enhanced blood flow pattern outside the uterine cavity.

Management of ruptured ectopic :

The principle in the management is resuscitation and laparotomy, and not resuscitation followed by laparotomy. Salpingectomy is the definitive surgery.

Other options :

Option A - Placenta previa is a type of antepartum hemorrhage occurring after 28 weeks of pregnancy. It presents as sudden onset painless, recurrent vaginal bleeding. USG findings reveal placenta occupying the lower segment of uterus.

Option B - Abruptio placenta is another form of antepartum hemorrhage occurring due to premature separation of the normally situated placenta. The majority of cases occur in the third trimester. The clinical features vary depending on whether the hemorrhage is concealed or revealed and include abdominal pain, vaginal bleeding. Ultrasound reveals a collection of blood in the retroplacental, subchorionic or preplacental region.

Option D - Missed abortion refers to when the fetus is dead and retained inside the uterus. Clinical features include the persistence of brownish vaginal discharge and subsidence of pregnancy symptoms. Examination findings are uterus size smaller than the period of gestation and closed cervical os. USG reveals an empty sac or a fetus without cardiac activity.

Solution to Question 7:

The given scenario is suggestive of chronic hypertension with superimposed severe pre-eclampsia. The next step in the management is to give injection labetalol, and magnesium sulfate and expedite the delivery, irrespective of the duration of gestation.

The criteria for diagnosis of superimposed pre-eclampsia are :

• New onset proteinuria

• Aggravation of hypertension

• Development of HELLP syndrome

• Development of headache, scotoma, and epigastric pain.

Pre-eclampsia is further divided into :

• Nonsevere pre-eclampsia: Blood pressure >140/90 mmHg but <160/110 mmHg without significant proteinuria

• Severe pre-eclampsia: Persistent increase of systolic blood pressure above 160 mmHg or diastolic above 110 mmHg, proteinuria, oliguria (<400 mL in 24 hours), platelet count <1 lakh/cu.mm, serum creatinine level of >1.1 mg/dL, persistent severe epigastric pain, marked elevation of liver enzyme levels, convulsions or visual disturbances, intrauterine growth restriction of the fetus, and pulmonary edema.

In a patient with severe pre-eclampsia, ie, systolic BP ≥ 160 mm Hg or diastolic ≥ 110 mm Hg, immediate delivery after single-dose steroids is indicated regardless of the period of gestation. Concurrently, injection labetalol is given to reduce the blood pressure and injection magnesium sulfate is given for seizure prophylaxis.

Patients with pre-eclampsia may be expectantly managed up to 34 weeks of gestation. After 34 weeks pregnancy, may be prolonged up to 37 weeks if there is adequate BP control, and delivery is recommended at 37 weeks of gestation. However, in the following events, pregnancy is terminated regardless of the period of gestation:

• Uncontrolled severe hypertension

• Eclampsia

• Pulmonary edema

• HELLP Syndrome

• Renal dysfunction

• Coagulopathy (disseminated intravascular coagulation)

• Abruption

• Fetal compromise



Solution to Question 8:

In the given scenario, outlet forceps would be the best choice of instrumental delivery as the fetus is at +3 station, with suture in AP diameter and has caput.

The criteria for the application of different types of forceps are as follows :

Outlet forceps :

• Scalp is visible at the introitus without separating the labia

• Fetal skull has reached the level of the pelvic floor

• Sagittal suture is in direct anteroposterior diameter or in the right or left occiput anterior or posterior position, but rotation is <=45 degrees

• Fetal head is at or on the perineum

Low forceps :

• Leading point of the fetal skull (station) is at +2 cm or more but has not yet reached the pelvic floor, and

(a) Rotation is <=45°, or (b) Rotation is >45°

Mid forceps :

• Fetal head is engaged. Head is 1/5 palpable per abdomen but the station is above +2 cm but not above the ischial spines (station 0).

High forceps :

• Head is not engaged. This type of forceps is not used in current obstetrics.

Note : Ventouse is not the best choice for a fetus with caput, as there is a risk of scalp avulsion. Also, the vacuum cannot be adequately generated owing to the presence of soft tissue swelling.



Solution to Question 9:

Growth hormone does not play a role in the in-utero growth of a baby. It is, however, essential for postnatal growth.

Normal fetal growth is characterized by cellular hyperplasia followed by hyperplasia and hypertrophy and lastly by hypertrophy alone. The fetal growth increases linearly until the 37th week. It is controlled by genetic factors in the first half and by environmental factors in the second half of pregnancy. Imprinted genes (maternal or paternally acquired alleles) primarily control fetal growth.

The important physiological factors are race (European babies are heavier than Indians); sex (male baby weighs >female); parental height and weight (tall and heavier mothers have heavier babies); birth order (weight rises from first to second pregnancy) and socioeconomic factors (heavier babies in social class I and II).

Fetal growth is predominantly controlled by insulin-like growth factor -1 (IGF-1), insulin and other growth factors. Maternal and fetal thyroid hormones also play a role in the normal development of virtually all fetal tissues, especially the brain.

Solution to Question 10:

Post-partum anti-D prophylaxis is given if the baby is O positive and DCT (direct Coombs' test) is negative.

To prevent active immunization of Rh-negative yet unimmunized, Rh anti-D immunoglobulin (IgG) is administered intramuscularly to the mother following childbirth. The mode of action is antibody-mediated immune suppression.

Anti-D immunoglobulin should be administered within 72 hours or preferably earlier following delivery or abortion. It should be given provided the baby born is Rh-positive and the direct Coombs' test is negative. This reduces the alloimmunization rate by 90%. Use of anti-D immunoglobulin at 28 weeks gestation additionally reduces third-trimester alloimmunization to 0.1%.

Dose: Anti-D-gammaglobulin is administered intramuscularly to the mother (300 μg) following delivery. All Rh-negative unsensitized women should receive 50 μg of Rh-immunoglobulin IM within 72 hours of induced or spontaneous abortion, ectopic or molar pregnancy or chorionic villus sampling in the first trimester. Women with pregnancy beyond 12 weeks should have a full dose of 300 μg. Standard 300 μg of anti-D immunoglobulin IM dose will protect a mother from fetal hemorrhage of up to 30 mL of fetal whole blood or 15 mL of fetal RBCs.

Other options:

Anti-D prophylaxis is futile if -

• the baby has Rh negative blood group - there is no risk of alloimmunization when the Rh antigen is absent

• if the DCT of the baby's blood is positive - it means that alloimmunization has already occurred. Such babies are managed with other treatment options like exchange transfusion.

Solution to Question 11:

The correct dose of administration of antenatal steroids for lung maturity is dexamethasone 6 mg 12 hours apart 4 doses.

Maternal administration of corticosteroids is advocated where the pregnancy is less than 34 weeks. This helps in fetal lung maturation. This is beneficial when the delivery is delayed beyond 48 hours of the first dose. The benefit persists as long as 18 days. Either betamethasone 12 mg IM 24 hours apart for two doses or dexamethasone 6 mg IM every 12 hours for 4 doses is given. Betamethasone is the steroid of choice (RCOG–2004).

Antenatal steroids also reduce the incidence of respiratory distress syndrome, necrotizing enterocolitis, intraventricular hemorrhage and patent ductus arteriosus.

Risks of antenatal corticosteroid use:

• Premature rupture of the membranes especially with evidence of infection, as the infection may flare-up

• Insulin-dependent diabetes mellitus where patients need insulin dose readjustment

• Transient reduction of fetal breathing and body movements.

Solution to Question 12:

Uterine artery pulsatility index (PI) < 95th percentile is not a criterion used to diagnose intrauterine growth restriction (IUGR).

Fetal growth restriction (FGR)/IUGR is said to be present in those babies whose birth weight is below the 10th centile of the average for the gestational age.

The causes of fetal growth restriction are divided into :

• Maternal - constitutional, nutritional, anemia, hypertension, chronic renal disease, alcohol, smoking

• Fetal - Structural anomaly, chromosomal anomaly, infections, multiple pregnancy

• Placental -placenta previa, abruption, circumvallate placenta

• Unknown

Diagnosis :

Clinical -

• Symphysis fundal height (SFH) measurement in centimeters closely correlates with gestational age after 24 weeks. A lag of 3 cm or more suggests growth restriction.

• Maternal weight gain remains stationary or, at times, falling

• Measurement of the abdominal girth showing stationary or falling values

Ultrasound -

Various criteria and thresholds have been used to define FGR. These have included estimated fetal weights (EFW) below the 3rd, 5th, or 10th percentiles; similar abdominal circumference (AC)

percentiles; a specified decline in the EFW percentile or AC percentile over serial assessments; and various abnormal Doppler findings.

Currently, the ACOG (2021) and the Society for Maternal-Fetal Medicine (2020) recommend defining FGR as either an EFW <10th percentile for gestational age or an AC <10th percentile for gestational age.

AC is the single most sensitive parameter to detect FGR.

Doppler velocimetry - The umbilical artery Doppler study should be the primary surveillance tool in FGR (RCOG).

• In normal pregnancy, the systolic/diastolic (S/D) ratio, pulsatility index (PI) and resistance index (RI) decrease as the gestational age advances. Values greater than 2SD indicate reduced diastolic velocities and increased placental vascular resistance. These are associated with FGR and intra-uterine fetal hypoxia.

• Reduced or absent or reversed end-diastolic velocity (AREDV) in the umbilical artery indicates fetal jeopardy and poor perinatal outcome.

• Umbilical venous pulsations indicate inefficient cardiac output with a rise in central venous pressure → impending cardiac failure.

• Middle cerebral artery (MCA): Increased diastolic velocity (brain-sparing effect) is observed in FGR. This is due to cerebral vasodilatation in response to hypoxemia. Abnormal MCA PI, suggests fetal compromise.

• Doppler cerebroplacental ratio (CPR): It is calculated by dividing MCA PI by UA PI. In FGR, placental resistance (UA PI) increases and fetal brain vascular resistance (MCA PI) decreases resulting in a decrease in CPR index. CPR is more sensitive compared to UA or MCA Doppler alone.

Management is as outlined below :



Solution to Question 13:

Early onset pre-eclampsia is defined before 34 weeks.

Pre-eclampsia is characterized by hypertension and end-organ dysfunction that typically develops after 20 weeks of gestation in a previously normotensive individual.

Features of pre-eclampsia:

• Hypertension:

Blood pressure ≥140/90 mmHg on 2 occasions at least 4 hours apart, or ≥160/110 mmHg in a single measurement.

• Proteinuria:

≥300 mg of protein in a 24-hour urine collection.

Protein/creatinine ratio ≥0.3.

Dipstick reading of ≥1+ (though less specific).

• End-Organ Dysfunction (may occur even without proteinuria):

Renal: Elevated serum creatinine >1.1 mg/dL or doubling of baseline.

Hepatic: Elevated liver transaminases (AST/ALT).

Hematologic: Thrombocytopenia (<100,000/µL).

Neurologic: Severe headache, visual disturbances (e.g., scotoma, blurred vision).

Pulmonary: Pulmonary edema.

Early-onset pre-eclampsia:

• Occurs before 34 weeks of gestation.

• Associated with more severe complications, including fetal growth restriction and placental insufficiency.

• Requires careful monitoring and often early delivery if maternal or fetal conditions worsen.

Late-onset pre-eclampsia:

• Occurs at or after 34 weeks of gestation.

• Typically milder, with fewer complications.

Management

• Mild pre-eclampsia: Close monitoring, antihypertensives (e.g., labetalol, nifedipine), and maternal-fetal surveillance.

• Severe pre-eclampsia: Hospitalization, magnesium sulfate for seizure prophylaxis, and delivery planning (often the definitive treatment).

• Timing of Delivery:

Recommended at ≥37 weeks for mild pre-eclampsia.

Earlier delivery for severe cases.

Solution to Question 14:

In the given scenario, cervical cancer is screened by performing a pap smear every 3 years, starting at 21 years.

Pap smear is a screening test. In case of positive test further investigations should be done.

Pap Smear Schedule

• Time for initiation: 21 years, regardless of the age of first sexual intercourse.

• Women aged 21-29 years: Every 3 years(earlier it was done annually)

• Women aged 30-65 years: Pap test + HPV testing every 5 years.

• Women should stop having cervical cancer screening after the age of 65 years. If suspicious PAP, screening till 75 years.

• HIV+ women: annual checkup until three consecutive negative tests followed by testing every 3 years for life.

• HPV DNA testing in combination with PAP improves the sensitivity of screening.

Ayres spatula and endocervical brush are the instruments used for the pap test.

Ayres spatula:

• One end is blunt and the other end is bifid.

• Blunt end is used to take samples for vaginal cytology.

• Bifid end is used to take Pap smear

Endocervical brush

Solution to Question 15:

The USG findings of a unilocular cyst with reticular pattern indicate a functional hemorrhagic cyst. Hence, the next line of management is to repeat USG after 6-12 weeks.

Although the cyst size is >7 cm, invasive treatments are not suitable immediately because of her current use of warfarin which needs time to be tapered. Also, as most functional cysts resolve spontaneously over time, repeat USG might show a decrease in cyst size.

Management of ovarian cysts in reproductive age and perimenopausal women is mentioned in the table below:

Other Options:

Option A: Measure CA-125: In this case, it might not be the best first step because CA-125 is not specific and its level can be influenced by multiple factors. Additionally, the cyst described here is more likely to be benign (reticular pattern suggesting blood), such as a functional ovarian cyst or follicular cyst.

Option C: Surgical Intervention: Surgical intervention may be required if there are signs of torsion, rupture, or malignancy. However, with a simple, unilocular cyst without these features, surgery is usually not warranted immediately, especially as the patient is on warfarin and needs to be tapered. This approach would be more appropriate if the cyst showed signs of complications or did not resolve after monitoring.

Option D: MRI Pelvis: An MRI can offer more detailed imaging if there is concern about the nature of the cyst, but it is typically reserved for complicated cysts (e.g., solid components, suspicious features). It's not the first-line investigation in this case.

Size	Management
< 5cm	Wait & watch; no additional t reatment required

Size	Management
5cm - 7 cm	Follow up with serial TVS in 6-12 weeks; if cyst persists co ntinue TVS on yearly basis
> 7cm	MRI pelvis; surgical evaluatio n

Solution to Question 16:

Delayed cord clamping (DCC) is done in term and preterm babies irrespective of birth weight.

DCC involves waiting at least 30–60 seconds or more after birth before clamping the umbilical cord. It is recommended for both term and preterm babies as part of standard delivery care.

Indications of DCC:

• Term infants: DCC improves iron stores and reduces the risk of anemia in infancy.

• Preterm infants: DCC increases blood volume and reduces the risks of:

• Intraventricular hemorrhage

• Necrotizing enterocolitis

• Provides stabilization benefits by enhancing transitional circulation

Babies irrespective of birth weights: The benefits of DCC apply regardless of the baby's weight.

Contraindications of DCC:

• Babies requiring immediate resuscitation: In emergencies, immediate neonatal stabilization takes priority, and DCC may not be feasible.

• Maternal or placental complications: Conditions like placental abruption or postpartum hemorrhage may necessitate immediate cord clamping for maternal safety.

Other options:

Option B and C: Vigorous term and preterm babies requiring resuscitation: Resuscitation needs immediate intervention, which may preclude delayed cord clamping.

Option D: All term babies born at 37 weeks: While DCC is recommended for term infants, the guideline includes preterm babies too, so this option is incomplete. It also fails to exclude babies requiring resuscitation.

Solution to Question 17:

25% of couples can be attributed to unexplained infertility.

Unexplained infertility refers to cases where no clear cause of infertility can be identified after a thorough evaluation of both partners. In the scenario provided, the male partner has normal seminal parameters, and the female partner has normal hormonal levels and patent fallopian

tubes, making this a case of unexplained infertility.

Some causes of unexplained infertility include:

• Ovulatory Dysfunction

• Impaired sperm function despite normal semen analysis

• Tubal or Uterine Factors

• Endometriosis

• Antisperm antibodies or autoimmune issues.

• Lifestyle and Environmental Factors: Stress, smoking, diet, or environmental toxins.

Unexplained infertility is diagnosed when standard evaluations yield normal results:

• Semen analysis showing normal volume, concentration, motility, and morphology.

• Normal hormonal tests, basal body temperature, or ovulation predictor kits.

• Hysterosalpingography (HSG) or saline sonography shows no blockages.

• Transvaginal ultrasound or hysteroscopy ruling out structural abnormalities.

If no cause is identified after these tests, the couple is categorized as having unexplained infertility.

Management includes:

1. Expectant Management

• Suitable for younger couples (<35 years) with a short duration of infertility.

• Lifestyle modifications (e.g., weight management, smoking cessation, stress reduction).

2. Medical :

• Ovulation Induction: Clomiphene citrate or Letrozole.

• Gonadotropins: Injectable hormones (FSH, LH) for controlled ovarian stimulation.

3. Intrauterine Insemination (IUI)

4. Assisted Reproductive Technology (ART)

• In Vitro Fertilization (IVF)

• Intracytoplasmic Sperm Injection (ICSI)

Solution to Question 18:

Increase in systolic and diastolic blood pressure after 6 weeks is not a physiological change seen in pregnancy.

During early pregnancy, systemic vascular resistance decreases, leading to a drop in blood pressure (both systolic and diastolic), which typically reaches its lowest point in the second trimester. Blood pressure gradually returns to pre-pregnancy levels in the third trimester.

An increase in blood pressure during early pregnancy is not physiological and could indicate an underlying pathology (e.g., chronic hypertension or gestational hypertension later in pregnancy).

Cardiovascular parameters that increase in pregnancy are:

• Cardiac output

• Stroke volume

• Heart rate

Cardiovascular parameters that decrease in pregnancy are:

• Peripheral vascular resistance (due to smooth muscle relaxation by progesterone)

• Both diastolic BP and systolic BP (decrease in DBP > SBP)

• Mean arterial BP

• Artery venous O2 gradient

Other Options:

Option B: Peripheral resistance to insulin leading to hyperinsuliemia: During pregnancy, placental hormones like human placental lactogen (hPL), cortisol, and progesterone induce insulin resistance to ensure adequate glucose availability for the fetus. This leads to compensatory hyperinsulinemia.

Option C: Fasting hypoglycemia and postprandial hyperglycemia: Fasting hypoglycemia occurs due to increased maternal glucose utilization by the fetus. Postprandial hyperglycemia is a result of insulin resistance and slower glucose uptake.

Option D: Increased plasma volume and RBC mass: Plasma volume increases by about 40-50%, and red blood cell (RBC) mass increases by 20-30%, leading to hemodilution and the physiological anemia of pregnancy.

Solution to Question 19:

Instrument-assisted delivery is not indicated in cephalo pelvic disproportion. Ruling out CPD is a prerequisite before attempting instrument-assisted delivery.

Cephalo-pelvic disproportion (CPD) is a condition in which the baby's head is too large, or the pelvis is too small, for the baby to pass through the birth canal. In such cases, instrument-assisted delivery is not indicated because vaginal birth is not feasible. The appropriate management for CPD is usually a cesarean section to avoid complications for both mother and baby.

Conditions where instrument-assisted delivery may be indicated:

• Fetal distress: If the baby is showing signs of distress (such as decelerations in the heart rate), an instrument-assisted delivery may be needed to expedite delivery and reduce the risk of further harm. (Option A)

• Maternal exhaustion: In cases of maternal fatigue, particularly after a prolonged second stage of labor, instruments may be used to assist in delivery if the mother is unable to continue pushing effectively. (Option B)

• Prolonged second stage of labor: If the second stage of labor (when the cervix is fully dilated and the baby is descending) is prolonged, instruments can be used to assist in the delivery of the baby, especially if there are signs of fetal distress or maternal exhaustion. (Option C)

Prerequisites for instrumental vaginal delivery:

• Ruptured membranes vertex presentation

• Cephalopelvic disproportion not suspected

• No fetal coagulopathy or bone demineralization disorder

• Completely dilated cervix

• Precisely assessed fetal head position

• Engaged head

• Experienced operator

Solution to Question 20:

The primary oocyte is arrested in prophase I of meiosis during a specific substage called diplotene.

Primary oocytes initiate meiosis I, but are arrested in prophase I before birth.

The arrest persists until puberty, when hormonal signals trigger the resumption of meiosis in a selected number of oocytes during each menstrual cycle.



Meiosis I (Reduction Division): Homologous chromosomes separate.

• Prophase I: Chromosomes pair, cross over.

• Metaphase I: Homologous pairs align at the equator.

• Anaphase I: Homologous chromosomes separate.

• Telophase I: Two haploid cells form.

Meiosis II (Equational Division): Sister chromatids separate.

• Prophase II: Chromosomes condense.

• Metaphase II: Chromosomes align at the equator.

• Anaphase II: Sister chromatids separate.

• Telophase II: Four genetically unique haploid cells form.

Prophase I consists of several substages (leptotene, zygotene, pachytene, diplotene, and diakinesis). In primary oocytes:

• Leptotene: Chromosomes condense and become visible. (Option B)

• Zygotene: Homologous chromosomes pair up through synapsis. (Option C)

• Pachytene: Crossing over occurs between non-sister chromatids. (Option D)

• Diplotene stage: The homologous chromosomes are connected at chiasmata (sites of crossing over), and the oocyte enters a prolonged resting phase. At ovulation, the primary oocyte completes meiosis I and becomes a secondary oocyte, which then gets arrested in metaphase II of meiosis until fertilization.

• Diakinesis: Chromosomes fully condense; nuclear envelope breaks down.

Solution to Question 21:

Restitution describes a passive adjustment of the head to untwist the neck after internal rotation.

The most common position of the fetus is the left occipito-lateral position where the vertex occupies the left anterior quadrant of the mother's pelvis.

During internal rotation, there will be anterior rotation of the occiput by 2/8th of a circle, placing the occiput behind the symphysis pubis. There is always an accompanying movement of descent with internal rotation. (Option B)

Torsion of the neck occurs during internal rotation of the head. The neck cannot withstand such a high degree of torsion, so the shoulders will rotate in the same direction to the extent of 1/8th of a circle, placing the shoulders in the oblique diameter with one-eighth of the torsion still left behind. Torsion refers to the twisting forces applied to the fetal head and neck during internal rotation to align the head with the birth canal.

In restitution, these twisting forces are released, allowing the head to return to its anatomical alignment with the shoulders.



Other Options:

Option B: Movement of the head by 1/8th of the circle due to torsion: While the head typically does rotate by about 1/8th of a circle (45°) during restitution, this is a passive as the tension in the neck resolves. The phrase "due to torsion" is misleading because torsion refers to twisting forces exerted during internal rotation, not restitution. Restitution is about the release of torsion, not its application.

Option C: Stretching of the vulval outlet by the head: This describes a process that occurs during the crowning stage of labor, not during restitution. At crowning, the fetal head stretches the vulval outlet as it emerges. Restitution occurs after the head is delivered and involves the passive rotation of the head to align with the shoulders.

Option D: External rotation after delivery of shoulder: Once the shoulders are delivered, the head undergoes a second adjustment to align with the shoulders as they rotate to fit through the maternal pelvis. This is an active movement that happens post-delivery of the shoulders, while restitution occurs before shoulder delivery.

Solution to Question 22:

Bimanual compression is not part of AMTSL.

AMTSL is recommended by the WHO to prevent postpartum hemorrhage (PPH). Uterotonics are the most important components in AMTSL to do so.

Note: Uterine massage or fundal massage is not recommended as an intervention to prevent PPH in women who have already received prophylactic uterotonics such as oxytocin (core component of AMTSL). Hence, it is now an optional component of AMTSL that is done in patients with uterine atony as a part of the management of PPH. (Option A)

1.	Injectinguterotonicswithin 1 minute of delivery of the bab y or immediately after deliver y of the anterior shoulder (O ption D)	Most important stepOxytocin 10 IU im/iv infusion is the pr eferred uterotonic.(Not reco mmended by WHO- carbopr ost)
2.	Delayed cord clamping	Done within 1-3 mins of deliv ery80 ml of additional blood goes to the fetusPrevents feta l anemia
3.	Delivery of placenta by contr olled cord traction: Modified Brandt Andrews technique(O ption B)	The uterus is pushed upward s and backwardThe traction o f cord is downwards and forw ards
4.	Intermittentassessment of ut erine tone	Done every 15 mins for 2 hou rs

Solution to Question 23:

The patient with malodorous vaginal discharge and a positive whiff test, strongly suggests bacterial vaginosis.

Bacterial vaginosis (BV) is caused by an imbalance in the vaginal microbiome, characterized by the overgrowth of anaerobic bacteria (Gardnerella vaginalis) and a decrease in the normal protective lactobacilli. Bacterial vaginosis is a polymicrobial infection characterized by malodorous (fishy smell) vaginal discharge. Causative organisms include:

• Gardnerella,

• Mobiluncus,

• Mycoplasma hominis

Bacterial vaginosis is diagnosed using Amsel's criteria:

Amsel's Criteria: (any 3 should be present)

• Thin, white homogenous vaginal discharge

• pH of Vaginal fluid>4.5

• Fishy odor of discharge accentuated on adding 10% KOH (Whiff test)

• Presence of clue cells on microscopy. Many epithelial cells present with granular cytoplasm caused by small gram-negative bacilli adhering on their surface, these cells are called clue cells.



Treatment is oral metronidazole 500 mg BD for 7 days. Alternatively, Ampicillin 500 mg or cephalosporin 500 mg bid for 7 days is also effective.

Other options:

Option A: Candida: Presents with thick, white "cottage cheese-like" discharge, intense itching, and no fishy odor. Whiff test is negative.

Option B : Chlamydia: Often asymptomatic or causes mucopurulent cervical discharge and friable cervix. Does not cause a fishy odor or a positive whiff test.

Option D : Pelvic Inflammatory Disease (PID): Causes pelvic pain, fever, and cervical motion tenderness. Not typically associated with malodorous discharge or a positive whiff test.

Solution to Question 24:

This patient presenting acute urinary retention, cyclic abdominal pain, bluish bulge on genital examination and primary amenorrhea (no menarche), likely has imperforate hymen.

Imperforate hymen results due to failure of canalization of the inferior end of the vaginal plate. This anomaly usually remains unrecognized until puberty. It can lead to mucocolpos in early infancy.

Key Features of Imperforate Hymen:

• Cyclic abdominal pain: Caused by retained menstrual blood (hematocolpos) due to the obstruction.

• Primary amenorrhea: Menstruation cannot occur due to the blockage.

• Bluish bulge: Hematocolpos causes distension of the vaginal canal, which can present as a bluish bulge at the hymen.

• Urinary retention: The distended vagina (hematocolpos) can compress the bladder or urethra, leading to urinary retention.

Ultrasound pelvis confirms the diagnosis.

Management: A simple cruciate incision of the hymen allows the free drainage of the collected menstrual blood, and leads to normal reproductive function.

Other Options:

Option A: Transverse vaginal septum: Would also cause primary amenorrhea and hematocolpos. Typically presents without a bluish bulge, as the obstruction is deeper in the vagina. Extensive dissection and vaginoplasty is required.

Option C: MRKH: Mayer-Rokitansky-Küster-Hauser (MRKH) syndrome causes primary amenorrhea due to absent uterus and upper vagina. There is no cyclic pain or bluish bulge, as menstruation cannot occur.

Option D: OHVIRA: Obstructed hemivagina and ipsilateral renal anomaly: Rare condition with an obstructed hemivagina. Often associated with renal anomalies and a functional contralateral uterus, but it presents with cyclic pain later after menarche onset, not primary amenorrhea.

Solution to Question 25:

Juvenile granulosa cell tumor does not constitute a germ cell tumor. It is a pure-sex cord tumor.

Germ cell tumors:

- Dysgerminoma
- Endodermal sinus tumor (Yolk sac tumor)
- Embryonal carcinoma
- Polyembryoma
- Non-gestational choriocarcinoma
- Teratoma (Mature/immature)
- Mixed forms

Solution to Question 26:

The obstetric conjugate is the smallest pelvic diameter at the brim.

It is the distance between the midpoint of the sacral promontory to the prominent bony projection in the midline on the inner surface of the symphysis pubis. It is the shortest anteroposterior diameter in the anteroposterior plane of the inlet and measures 10 cm. It represents the narrowest fixed diameter of the pelvic inlet that the fetal head must navigate. It cannot be measured directly during a pelvic exam but is estimated using the diagonal conjugate.



Other Options:

Option B: Diagonal Conjugate: It is the distance between the lower border of the symphysis pubis to the midpoint on the sacral promontory. It measures 12 cm. Obstetric conjugate is derived by subtracting 1.5–2 cm from the diagonal conjugate depending upon the height, thickness and inclination of the symphysis pubis.

Option C and D : True conjugate: Also called anatomical conjugate, is the distance between the midpoint of the sacral promontory to the inner margin of the upper border of the symphysis pubis and measures 11 cm.

Solution to Question 27:

An ectopic pregnancy is one in which the fertilized ovum is implanted and develops outside the normal endometrial cavity.

Clinical features :

The classic triad of symptoms is - abdominal pain (100%), preceded by amenorrhea of 6-8 weeks(75%), and lastly, the appearance of vaginal bleeding (70%).

Abdominal pain is the most constant feature. (Statement 1) Pain is located in the lower abdomen. Shoulder tip pain (25%) (referred pain due to diaphragmatic irritation from hemoperitoneum) may be present.

Pelvic examination - Vaginal mucosa is blanched white; uterus seems normal in size or slightly bulky; extreme tenderness on fornix palpation(75%).

Risk factors of ectopic pregnancy:

• Previous ectopic pregnancy

• Tubal corrective surgery, tubectomy

• Assisted reproductive technology

• Documented tubal pathology

• Pelvic inflammatory disease (Chlamydia infection and salpingitis isthmica nodosa), Genital tuberculosis, sexually transmitted diseases (Statement 2)

• Endometriosis, Pelvic adhesions

• Prior abortion, Prior Caesarean delivery

• Multiple sexual partners

• Progesterone-only pills (POPs)

Diagnosis:

• Serum ■rog■■t■rone – concentrations are higher in viable IUPs than in ectopic pregnancies and IUPs that are destined to abort, and measurement of serum ■rog■■t■r■ne may be useful in a patient with abdominal pain and bleeding who has a serum hCG level below that expected for the gestational age. However, the definition of low ■rog■■tero■■ is unclear as there is no specific value to diagnose ectopic pregnancy. (Statement 3)

• ß-hCG - lower concentration of hCG and doubling in plasma failing to occur in 2 days.

• Transvaginal sonography (TVS) features are absence of intrauterine pregnancy with a positive pregnancy test.

• Color Doppler sonography can identify the placental shape (ring-of-fire pattern).

Management of ectopic pregnancy:

1. Medical management is preferred in the following cases:

• Low initial serum beta-hCG level (Less than 5,000 IU/L)

• Small ectopic pregnancy size (Less than 3.5 cm)

• Absent fetal cardiac activity

• Hemodynamically stable

2. Expectant management: A low initial serum hCG concentration, a considerable reduction in levels over 48 hours, and a sonographic inhomogeneous mass instead of a tubal halo or other gestational features are predictive variables for success.

3. Surgical management with salpingectomy or salpingostomy is done when:

• Gestational sac >3.5 cm

• Fetal cardiac activity present

• Beta-hCG >5000 mIU/mL

• Significant pain and hemoperitoneum

Oral methotrexate is not the preferred route for treating ectopic pregnancy, as it may lead to variable absorption and unreliable dosing. Intramuscular injection is the preferred route of administration for methotrexate in the management of ectopic pregnancy. (Statement 4)

Solution to Question 28:

Dimeric inhibin B is not included in the quadruple test.

The quadruple test, performed during the second trimester (15–22 weeks, ideally around 16–18 weeks), includes the following markers:

• Alpha-fetoprotein (AFP)

• Unconjugated estriol (uE3)

• Human chorionic gonadotropin (hCG)

• Inhibin A

The quadruple test is primarily used for screening chromosomal abnormalities such as Down syndrome (trisomy 21), Edward syndrome (trisomy 18), and neural tube defects.

The following table summarizes the findings of the quad-screening test in various conditions.

Parameter	Trisomy 21	Trisomy 18	Neural tube defect or abdominal wall defect
MSAFP	Low	Low	High
β hCG	High	Low	Normal
Unconjugated estriol	Low	Low	Normal
Inhibin A	High	Normal/ Low	Normal

Solution to Question 29:

The low FSH and LH levels, along with the low BMI, suggest hypogonadotropic hypogonadism.

BMI of 17.5 kg/m² indicates low body weight, which can lead to hypothalamic suppression and a disruption in the normal feedback mechanisms that regulate the menstrual cycle. FSH (2mIU/ml), LH (0.3IU/L): These values are low, indicating that the pathology lies at the level of the hypothalamus or pituitary, rather than the ovaries. In hypogonadotropic hypogonadism, there is insufficient secretion of gonadotropins (FSH and LH), which are needed for ovarian function. TSH (1.2mIU/ml) and Prolactin (20 ng/ml) are normal, which rules out thyroid dysfunction or hyperprolactinemia as the cause of the amenorrhea.

Causes of hypogonadotrophic hypogonadism:

• Kallman's syndrome, Prader-Willi syndrome

• Pituitary tumors, apoplexy

• Anorexia nervosa, excess exercise

• Irradiation, Infection,Trauma

Exercise-induced hypogonadotropic hypogonadism (EIHG): intense and prolonged physical activity leads to a decrease in the function of the hypothalamic-pituitary-gonadal (HPG) axis, resulting in lower levels of sex hormones. This occurs due to reduced secretion of gonadotropin-releasing hormone (GnRH), which subsequently lowers levels of luteinizing

hormone (LH) and follicle-stimulating hormone (FSH).

Causes: Excessive physical training, especially in endurance sports. Energy imbalance (low calorie intake vs. high energy expenditure). Psychological stress and overtraining.

Symptoms:

• Men: Decreased libido, erectile dysfunction, fatigue, and muscle weakness.

• Women: Irregular or absent menstrual cycles (amenorrhea).

Management includes rest and reduced exercise intensity. Adequate caloric intake to correct energy deficits. EIHG is a reversible condition with lifestyle modifications and is most commonly seen in athletes or individuals with excessive physical demands.

Other Options:

Option A: Primary ovarian failure (Premature ovarian insufficiency) is associated with elevated FSH and LH levels due to a lack of negative feedback from the ovaries, which is not the case here (FSH and LH are low).

Option C: Hypergonadotropic Hypogonadism refers to primary ovarian failure, where FSH and LH levels are elevated due to the lack of ovarian response. The low levels of FSH and LH here rule out this condition.

Option D: Normogonadotropic Normogonadism would indicate normal levels of FSH, LH, and estrogen, typically seen in conditions like polycystic ovary syndrome (PCOS). However, in this case, the low FSH and LH levels suggest a hypothalamic or pituitary issue.

Obstetrics and Gynecology NEET 2024

Question 1:

Identify the image shown below :



a) Ovarian torsion

b) Germ cell tumor

c) Ovarian cyst

d) Chocolate cyst of ovary

Question 2:

A 14-year-old rape victim is 22 weeks pregnant and she would like to terminate her pregnancy. Which of the following is done under the MTP Act?

a) Opinion of 1 doctor is enough upto 24 weeks

b) Termination allowed between 20-24 weeks only if detrimental to mother's life

c) Cannot be terminated

d) Can terminate pregnancy upto 24 weeks for sexual assault

Question 3:

A ventouse vacuum was placed at the flexion point. What is the correct statement regarding flexion point?

a) 3 cm from posterior fontanelle

b) Midpoint of anterior and posterior fontanelle

c) 3cm from anterior fontanelle

d) 6 cm from posterior fontanelle

Question 4:

A primigravida who is a known case of bronchial asthma develops PPH. Which of the following needs to be avoided in this patient?

a) Carboprost

b) Oxytocin

c) Mifipristone

d) Methyl ergometrine

Question 5:

A patient with a carcinoma cervix lesion showing cervical growth extending to the pelvic side wall along with the presence of hydronephrosis. What is the stage of the tumour?

a) II a

b) II b

c) III B

d) III a

Question 6:

A 60-year-old female presented with a mass per vagina, as shown in the image. What is the diagnosis?

a) Uterovaginal prolapse grade 1

b) Uterovaginal prolapse grade 2

c) Uterovaginal prolapse grade 4

d) Uterovaginal prolapse grade 3

Question 7:

A woman at 37 weeks of gestation comes to the OPD for an antenatal checkup. She is on warfarin for prosthetic valves. What would be the management of this patient?

a) Change to LMWH

b) Continue Warfarin

c) Immediate LSCS

d) Induction of labour

Question 8:

An 18-year-old girl with short stature has primary amenorrhea and has not attained puberty yet. Identify the chromosomal defect as shown in the image.



a) Turner's syndrome

b) Klinefelter's syndrome

c) Kallman syndrome

d) AIS

Question 9:

The illustration given below is used for which female genital pathology?



a) Prolapse

b) Vulval cancer

c) Carcinoma cervix

d) Endometriosis

Question 10:

A teenage girl presents with primary amenorrhea, normal breast and pubic hair development with normal uterus & ovaries. What is the karyotype?

a) 46 XY

b) 45 XO

c) 47 XXY

d) 46 XX

Question 11:

A female complaints of dysuria since 4 days associated with pelvic pain and abnormal vaginal discharge. Pap smear image is given below. What is the diagnosis?



a) Trichomonas

b) Human papilloma virus

c) Cytomegalovirus

d) Candida

Question 12:

A 28-year-old lady with 27 weeks of gestation has multiple congenital malformations in the fetus on radiological investigations. The parents want to terminate the pregnancy for which they went to the MTP board. Who is not a part of the MTP board?

a) Gynaecologist

b) Radiologist

c) Paediatrician

d) Lawyer

Question 13:

A woman at 30 weeks of gestation presents with pain and redness in the calf region. A diagnosis of DVT is made. Which of the following would you use in the management of the patient?

a) LMWH

b) Warfarin

c) Fondaparinux

d) Rivaroxbaran

Question 14:

A G2P1L1 woman with a previous history of breech delivery underwent an LSCS which was uneventful. Now, she was brought to the ER with an antepartum haemorrhage and was shifted for LSCS during which she lost a lot of blood. An obstetric hysterectomy had to be carried out. An image of the uterus is given below. What is the diagnosis?



a) Placenta previa

b) Abruptio placenta

c) Placenta accreta syndrome

d) Uterine rupture

Question 15:

A patient with 32 weeks of gestation had tachycardia and a blood pressure of 160/110 mm of Hg. On examination, she had pedal edema and lab investigations showed proteinuria. Pritchard regimen with magnesium sulphate was started. What is the indication to stop $MgSO_4$?

a) Urine output is 60 ml in 4 hours

b) Respiratory rate >18/min

c) Knee jerk present

d) Visual abnormalities

Question 16:

A G5P4 female comes for her routine scan. She is already a mother of 4 girl children and wants a boy this time. She is now interested to know the sex of the current child. According to PCPNDT act, what should you do as a radiologist?

a) Check for congenital anomalies of the fetus but not look for the sex

b) Check for congenital anomalies, the sex of the child and tell her if it is a female child

c) Check for congenital anomalies, the sex of the child and do not tell her

d) Only check routine ANC and do not do USG

Question 17:

A P1L1 woman, with a history of uneventful previous delivery, presents with secondary infertility and amenorrhea for 2 years. The laparoscopy performed is shown below. What is the likely diagnosis?



a) Tuberculosis

b) Endometriosis

c) Multiple fibroids

d) Sarcoidosis

Question 18:

A 45-year-old female came with complaints of spotting and foul-smelling discharge. On per speculum examination, cervical erosions were present. Which of the following is the most appropriate investigation required?

a) Liquid based cytology, Cervical biospy, HPV antigen testing

b) Pap smear, Cervical biopsy, HCV testing

c) Pap smear, Cervical biopsy, HBV DNA

d) Pap smear, Electrophoresis, HSV DNA

Question 19:

In the absence of other obstetric complications in twin pregnancies A and B. What is the preferred mode of delivery for A and B respectively?



a) A. Vaginal delivery B. LSCS

b) A. LSCS B. Vaginal delivery

c) Both LSCS

d) Both vaginal delivery

Question 20:

A P1L1 woman had delivered a baby 8 hrs before by vaginal delivery. What is the earliest time to insert an IUCD?

a) After 3 months

b) Within 48 hours

c) After 6 weeks

d) Between 48 hours - 6 weeks

Question 21:

A pregnant woman presents with a history suggestive of hyperthyroidism at 8 weeks of POG. What is the drug of choice?

a) Methimazole

b) Carbimazole

c) Propylthiouracil

d) Propranolol

Question 22:

A 42-year-old woman, P2L2, who had her last pregnancy 20 years ago, presents with heavy bleeding. A smooth, round, globular structure is observed on examination. On bimanual examination, the uterus is described as multiparous and firm, and the cervix admits one finger. What is the likely diagnosis?

a) Hypertrophied Cervix

b) Myomatous Polyp

c) Polypoid adenocarcinoma

d) Placental Polyp

Question 23:

The method shown in the image is:



a) Pinard's Maneuver

b) Mauriceau Smellie Veit Method

c) Burns Marshall Technique

d) Ritgen Maneuver

Question 24:

A woman presents with lower abdominal pain. Transvaginal ultrasound (TVS) performed over two days reveals a left adnexal cystic mass with a cogwheel appearance, incomplete septa, and no vascular flow on Doppler imaging. What is the most likely diagnosis?



a) Hydrosalphinx

b) Malignant ovarian cyst

c) Follicular cyst

d) Hemorrhagic cyst

Question 25:

A woman undergoing infertility treatment is found to have bilateral tubal blockage on hysterosalpingography. What is the most appropriate next step in her management?

a) Intrauterine insemination (IUI)

b) In vitro fertilization (IVF)

c) Laparoscopic chromopertubation

d) Ovulation induction

Question 26:

A primigravida experienced precipitous labor while being transported in an ambulance. Upon arrival at the hospital, the given injury is observed. What type of injury is most likely?



a) 2nd degree perineal tear

b) 4th degree perineal tear

c) 3rd degree perineal tear

d) 1st degree perineal tear

Question 27:

What is the most common site for tubal ligation and the reason for ligation at this site?

a) Ampulla; as it has more circular muscles than longitudinal muscles

b) Ampulla; as it is the widest part

c) Isthmus; as it has uniform muscle thickness

d) Isthmus; as it is the narrowest part

Question 28:

A 30-year-old female, P2L2, presents with a mass per vaginum. On examination, infravaginal elongation of the cervix is noted. How will you treat this condition?

a) McCall culdoplasty

b) Le Forte's surgery

c) Fothergill surgery

d) Latzko procedure

Question 29:

A 28-year-old P2L2 woman presents with complaints of a mass descending through the vagina and wishes to retain her fertility. She underwent an abdominal sling procedure with mesh placement. Which is the correct site of attachment of the mesh?

a) Symphysis Pubis

b) Ischial Tuberosity

c) Ischial Spines

d) Sacral Promontory

Question 30:

A woman who had a mediolateral episiotomy during normal labor presents with pain at the episiotomy site, which worsens during her menstrual periods. The findings noted on episiotomy site exploration are shown in the image below. What is the most likely diagnosis?



a) Retained Suture

b) Melanoma

c) Inclusion cyst

d) Endometriosis

Question 31:

During a lower segment cesarean section (LSCS) for a patient with a breech presentation, following findings were noted as given in the image. On per vaginal examination, two cervices are felt. What is the most likely diagnosis?



a) Unicornuate Uterus with Rudimentary Horn

b) Arcuate Uterus

c) Bicornuate Uterus

d) Uterine Didelphys

Question 32:

A 30-year-old lady, G3P2 with 10 weeks intrauterine gestation comes to OPD with an Intra Uterine Contraceptive Device(IUCD) and the tail of the IUCD is noted to be protruding from the cervical os. The most appropriate course of action is?

a) Keep the IUCD in place and continue prophylactic antibiotics

b) Keep the IUCD in place without any treatment

c) Remove the IUCD immediately and continue pregnancy

d) Terminate the pregnancy because of high risk of infection

Question 33:

A 24-year-old woman at 34 weeks of gestation is admitted for preeclampsia with severe features, and you have administered MgSO■ as prophylaxis. Which of the following is not a part of monitoring?

a) Routine measurement of serum magnesium levels

b) Urine output should be > 30 ml/hr

c) Respiratory rate > 12/min

d) Deep tendon reflexes should be present

Question 34:

An unmarried girl presents with amenorrhea and abdominal pain. On genital examination, she is found to have a softened cervix. What is this sign called?

a) Hegar's Sign

b) Braxton Hicks Sign

c) Oslander's Sign

d) Goodell's Sign

Question 35:

Which of the following is not a contraindication to the procedure shown in the image?



a) Twin pregnancy

b) Primigravida

c) Placenta previa

d) Premature rupture of membranes

Question 36:

A woman presents with chronic lower abdominal pain that worsens during menstruation. Ultrasound reveals a right adnexal mass with homogeneous echoes. MRI shows a 4x5 cm T1 hyperintense lesion without fat suppression, while the T2 sequence demonstrates low signal intensity with characteristic dark shading. What is the most likely diagnosis?

a) Cystic tumor of ovary

b) Dermoid cyst

c) Endometrioma

d) Teratoma

Question 37:

A woman's vaginal sample shows an abnormal Nugent's score. What is the most likely diagnosis?

a) Bacterial Vaginosis

b) Herpes Infection

c) Trichomoniasis

d) Candida Albicans

Question 38:

After ovulation, in which phase is the released egg arrested?

a) Primary oocyte with meiosis I completed

b) Secondary oocyte arrested in meiosis II

c) Primary oocyte arrested in meiosis I

d) Secondary oocyte with meiosis II completed

Question 39:

A 45-year-old woman presented with menorrhagia. A hysterectomy was performed, and the specimen is shown below. What is the most likely diagnosis?

a) Intramural Fibroid

b) Myomatous Polyp

c) Endometrioma

d) Adenomyoma

Question 40:

A 55-year-old obese woman with a history of diabetes and hypertension presents with postmenopausal bleeding. Endometrial sampling was performed and it revealed a positive result for PTEN, CTNNB1, KRAS, and MSI markers. What is the most likely diagnosis?

a) Mixed Müllerian Tumor

b) Leiomyosarcoma

c) Endometrioid Cancer

d) Serous Cystadenoma of Ovary

Question 41:

A 21-year-old woman at 10 weeks of gestation, comes to you for medical termination of pregnancy. Which method would you prefer for this patient?

a) Dinoprostone

b) Mifepristone and Misoprostol

c) Oxytocin infusion

d) Intrauterine instillation of hyper-osmotic saline

Question 42:

What is the time taken for the embryo to travel from point 1 to 8?



©MARROW

a) 2 days

b) 8 days

c) 5 days

d) 14 days

Question 43:

A 28-year-old woman, G4P3L3, with 1 neonatal death (NND) and 2 previous lower segment cesarean sections (LSCS), presents at term with an active genital herpes infection. What is the preferred mode of delivery?

a) Elective LSCS

b) Classical Cesarean section

c) Trial of labor (VBAC)

d) Induction of labor

Question 44:

A woman presents with lower abdominal pain. Ultrasound reveals a right adnexal mass with septations. Which tumor markers would be most appropriate to order in this case?

a) CA125, AFP, HCG, CEA

b) CA125, AFP, HCG, LDH

c) CA125, AFP, CA15-3

d) CA125, AFP, CA19-9

Question 45:

Identify the forceps typically used for delivering the aftercoming head in a breech delivery:



a) Wrigley's forceps

b) Kielland's forceps

c) Das's forceps

d) Piper's forceps

Answer Key

Question No.	Correct Option
1	d
2	d
3	a
4	a
5	c
6	c
7	a
8	a
9	a
10	d
11	b
12	d
13	a
14	c

15	a
16	a
17	a
18	a
19	b
20	b
21	c
22	b
23	c
24	a
25	c
26	b
27	c
28	c
29	d
30	d
31	d
32	c
33	a
34	d
35	b
36	c
37	a
38	b
39	a
40	c
41	b
42	b
43	a
44	b
45	d

Detailed Explanations

Solution to Question 1:

The given image shows a chocolate cyst of the ovary.

Chocolate cysts of the ovaries represent the most important manifestation of endometriosis. Ovaries are frequently involved bilaterally. It is lined by columnar epithelium.

Beneath the epithelium, large cells with brown cytoplasm called pseudoxanthoma cells are present. These are large macrophages or scavenger cells and their brown colouration is due to ingested blood pigments such as haemosiderin. They are polyhedral in shape and resemble lutein cells.

Chocolate cyst of ovary:



Solution to Question 2:

According to the MTP Act with 2021 amendments, pregnancy can be terminated upto 24 weeks of gestation in victims of rape and incest, fetal anomalies, or congenital malformations.

Solution to Question 3:

An important step in vacuum extraction is proper cup placement over the flexion point which is a point on the sagittal suture that lies 3 cm in front of the posterior fontanelle.



Advantages of correct positioning of the cup in vacuum extraction:

• Maximum traction

• Minimum cup detachment

• Twisting of the fetal head is avoided

• The smallest head diameter is delivered through the pelvic outlet

• The success rate is improved

• Fetal scalp injury rates are lowered

• Perineal trauma is reduced because the smallest fetal head diameter distends the vulva

Solution to Question 4:

The drug carboprost is contraindicated in patients with bronchial asthma.

Carboprost is a 15-methyl PGF2 alpha derivative used as a 2nd line drug in the treatment of PPH that is unresponsive to oxytocin. The normal dosage is 0.25mg. The maximum dose is 2mg.

Contraindications of carboprost:

Absolute	Relative
AsthmaPulmonary hypertensionSuspected amniotic fluid embolism	Pre-eclampsiaRenal diseaseLiver diseaseHeart disease

Other options :

Option B: Oxytocin is the first-line drug to prevent and treat PPH according to WHO guidelines. Carbetocin, which is a long-acting analogue of oxytocin, can also be used.

Option C: Mifepristone (RU486) is a progesterone antagonist, which blocks the hormonal action on the endometrium, causing the sloughing and shedding of decidua, preventing implantation.

Option D: Methyl ergometrine is contraindicated in hypertensive patients. It is given as an intramuscular injection or slow intravenous to avoid sudden hypertension.

Solution to Question 5:

Cervical cancer with hydronephrosis is classified as Stage III B according to FIGO staging.

Cervical cancer is the second most common gynaecological cancer. The FIGO staging uses clinical, radiological, or pathological findings to stage cervical cancer from Stage I to IV.

Stage III cervical carcinoma involves the lower 1/3rd of the vagina, with or without pelvic wall extension, hydronephrosis, and lymph node involvement. It is further staged as follows:

• Stage III A: Lower 1/3rd of the vagina involvement, no pelvic wall extension

• Stage III B: Pelvic wall involvement with or without hydronephrosis or a non-functioning kidney

• Stage III C: Involvement of the pelvic and/or paraaortic lymph nodes

• Stage III C1: Metastasis to pelvic lymph node only

• Stage III C2: Metastasis to para-aortic lymph nodes

Note: If the carcinoma had extended beyond the true pelvis, involving adjacent pelvic organs like the bladder or rectum, it would've been classified as Stage IV A.

Solution to Question 6:

In the given image, the uterus is visible outside the introitus suggesting the diagnosis of grade IV utero-vaginal prolapse.

The descent of one or more genital organs below their normal anatomical position is called pelvic organ prolapse (POP). It occurs when the muscles and tissues that support the pelvic organs become weakened or damaged, leading to one or more organs bulging into the vaginal canal.

The image below summarises the various degrees of genital organ prolapse.



Types of vaginal prolapse-

• Anterior vaginal wall: prolapse in the upper two-thirds wall is cystocele and in the lower one-third is urethrocele.

• Posterior vaginal wall: prolapse in the upper two-thirds wall is enterocele and in the lower one-third is rectocele.

Image given below shows cystocele.

Treatment options may include pessary, pelvic floor exercises, or surgery depending on the severity of prolapse.

Solution to Question 7:

The next appropriate step in the management of a pregnant woman with a prosthetic valve on warfarin therapy at 37 weeks of gestation is to stop warfarin and start low molecular weight heparin (LMWH).

Warfarin is safe and effective in preventing thromboembolism in the mother during the 2nd trimester. However, it is teratogenic and can lead to miscarriage, stillbirths, fetal anticoagulation, and fetal embryopathy if used in the 1st trimester.

Heparin is safe for the fetus as it does not cross the placenta but is less effective than warfarin in preventing thromboembolism.

LMWH- Low molecular weight heparin, UFH- Unfractionated heparin

For patients taking LMWH, factor Xa levels should be monitored weekly.

Heparin should be discontinued just before (at least 6 hours before) delivery. If delivery occurs before discontinuation and there is extensive bleeding, then iv protamine sulfate can be given. Warfarin or heparin anticoagulation can be restarted after 6 hours of vaginal delivery, usually without any problem.

Warfarin, LMWH, and UFH do not accumulate in breast milk and do not have an anticoagulant effect in the newborn. They are compatible with breastfeeding.

Period of gestation	Anticoagulant
Till 12 completed weeks	LMWH or UFH
13 to 36 weeks	Warfarin
36 weeks till just before deliv ery	LMWH or UFH
6 hours after vaginal delivery and 24 hours after cesarean section	Warfarin, LMWH or UFH

Solution to Question 8:

The given karyotyping image shows a “45 XO” abnormality, which is seen in Turner's syndrome.

Clinical features of Turner's syndrome include short stature, low posterior hairline, broad chest, widely spaced nipples, coarctation of the aorta, bicuspid aortic valve, cubitus valgus, and pigmented nevi.

Turner's syndrome is the most common sex chromosome abnormality in females. The classical karyotype is 45, XO. Other variants have structural abnormalities of X chromosomes or mosaic patterns. Some patients may present during infancy. They mainly present with oedema of the dorsum of the hand and foot due to lymphatic stasis. The most important cause of increased mortality in children with Turner's syndrome is cardiac abnormalities.

Turner's syndrome is a common cause of primary amenorrhea. During embryogenesis, fetal ovaries develop normally and the absence of a second X chromosome leads to accelerated loss of normally formed oocytes. This gets completed by the age of 2. The ovaries are reduced to atrophic fibrous bands without ova and follicles. These are called streak ovaries.

At puberty, they fail to develop secondary sexual characteristics.



Other options:

Option B: Eunuchoid body habitus with abnormally long legs is usually seen in Klinefelter Syndrome (47, XXY). The image below shows the clinical features and karyotyping of Klinefelter Syndrome respectively. Note the two X and one Y chromosomes.





Option C : Kallman's syndrome is a hypothalamic disorder that occurs due to defective gonadotropin-releasing hormone (GnRH).

It is mainly associated with anosmia or hyposmia or hypoplasia. Other associated features such as colour blindness, optic atrophy, deafness, cleft palate, renal abnormalities, and cryptorchidism may be present.

Option D : Androgen insensitivity syndrome is an X-linked recessive disorder in which the androgen receptors are absent, scanty, or fail to respond to testosterone. In complete AIS, the individual is born as a phenotypically normal female but generally presents with primary amenorrhea.

The external genitalia appears normal, with absent or scant pubic and axillary hair. The vagina is shortened and blind-ending with an absent uterus and ovaries. These patients may also present with a history of childhood inguinal hernias, due to the presence of undescended testis in the abdomen or in the inguinal canal. Due to the constant conversion of testosterone into estrogen, these individuals have normal breast development.

Solution to Question 9:

The given image shows POP-Q (pelvic organ prolapse quantification) system which was described by the International Continence Society for the prolapse of pelvic organs and has the following features:

• The hymen is taken as a fixed point

• Six reference points are measured, using a scaled spatula:

• Two on the anterior vaginal wall

• Two on the apical vagina

• Two on the posterior vaginal wall

• Three measurements: The genital hiatus, perineal body, and total vaginal length.

• The points above the hymen are described as minus and points below as plus.

Staging of POP Q:

Stage	Description
Stage 0	No descent of pelvic organs.
Stage 1	The leading edge of the prolapse is>1 cm abovethe level of thehymen.
Stage 2	The leading edge of the prolapse is<1cmproximal or distal to the hymen.
Stage 3	The leading point of prolapse is>1cm belowthehymenbut protrudes no further than 2 cm less than the total length of the vagina.
Stage 4	Completeeversionof vagina



Solution to Question 10:

The given clinical scenario is that of a female with primary amenorrhea, preservation of secondary characteristics with normal uterus & ovaries and an image showing imperforate hymen. The karyotype is normal, i.e, 46 XX karyotype.

Other options:

Option A & B: Gonadal dysgenesis in the form of 45XO (Turner syndrome) or 46XY (Swyer syndrome) presents with poor development of secondary sexual characteristics. Since the gonads are nonfunctional, there is a loss of negative feedback with increased levels of FSH and LH.

Option C: Klinefelter syndrome, characterized by an XXY genotype, is commonly associated with males having atrophic testes with small penis. Other features include tall stature, gynecomastia, lack of secondary sexual characteristics, and infertility.





Solution to Question 11:

This patient with symptoms of dysuria, pelvic pain, and abnormal vaginal discharge and the above pap smear shows large hyperchromatic nuclei and a perinuclear halo which is suggestive of koilocytes, which is a characteristic feature of infection caused by human papillomavirus (HPV).

Other features that can also be seen in histology are:

• Perinuclear halo

• Multinucleation

• Orangeophilic(Keratinisation) cytoplasm

• Acanthosis

• Chronic inflammatory infiltrate

Other options:

Option A: Trichomonas vaginalis is flagellated protozoa that causes trichomonas vaginitis. The pap smear shows a greyish-blue pear-shaped structure without the flagella as shown in the image below. The hanging drop test will reveal the presence of an actively motile pear-shaped flagellate organism.



Option C: Cytomegalovirus infection may be seen in pap smears rarely. Pap smear shows large, basophilic intranuclear inclusions or intracytoplasmic eosinophilic inclusions in endocervical glandular epithelial cells.

Option D: Candida albicans causes candidal vaginitis. On pap smear, It shows pseudohyphae and yeast cells and reactive squamous epithelial cells appear in the form of "shish kebab" as shown in the image below.



Solution to Question 12:

A lawyer is not a part of the medical board. According to the MTP Act with 2021 amendments, a medical board consisting of an obstetrician, a paediatrician, a radiologist and a person appointed by the state.

Solution to Question 13:

The pregnant woman with DVT at 30 weeks of gestation will be managed by low molecular weight heparin.

LMWH is the drug of choice in the management of DVT in any trimester of pregnancy, as it does not cross the placenta to cause fetal malformations. It has better bioavailability, longer half-life, predictable dose response and reduced adverse effects like osteoporosis and thrombocytopenia.

LMWH is given subcutaneously twice a day for up to 3- 6 months, followed by a prophylactic dose for the remaining pregnancy and post-partum period. After the symptoms subside, graduated compression stockings can be used for two years to reduce post-thrombotic syndromes like chronic leg paresthesia, intractable oedema and leg ulcers.

Other options:

Option B- Warfarin is used in the second trimester in patients with mechanical valves even though it is teratogenic, as LMWH does not provide adequate therapeutic effects and is associated with increased maternal thromboembolic events.

Option C and D- Fondaparinux and rivaroxaban are not typically used in pregnant women due to insufficient data about their safety during pregnancy. Fondaparinux is reserved for cases with heparin-induced thrombocytopenia and for those who do not tolerate heparin.

Solution to Question 14:

The given clinical scenario of this woman with previous LSCS, bleeding per vaginum during pregnancy with hysterectomy image showing placenta adherent to the uterine wall suggests that it is a case of placenta accreta syndrome.

Placenta accreta is described as an abnormally implanted, invasive, or adhered placenta. It is due to the partial or total absence of the decidua basalis and imperfect development of the Nitabuch layer.

Nitabuch layer is a layer of fibrin between the compact endometrium and the cytotrophoblastic shell in the placenta. This layer is absent in the placenta accreta.

There are 3 variants of placenta accreta syndrome. They are :

• Placenta accreta - villi are attached to the myometrium.

• Placenta increta - villi invade the myometrium.

• Placenta percreta - villi penetrate through the myometrium and to or through the serosa.



Solution to Question 15:

A urine output of 60 ml in 4 hours (<30ml/hr) is an indication to stop MgSO4 to prevent magnesium toxicity.

MgSO4 is used as an anticonvulsant in the management of eclampsia and as prophylaxis in severe preeclampsia. It can also be used as a tocolytic to stop preterm labour. The therapeutic level of magnesium in eclampsia is 4–7 mEq/L.

Patients must be monitored for magnesium toxicity:

• Loss of deep tendon reflexes

• Decreased respiratory rate <12/min

• Urine output <30 mL/h

• Chest pain and heart block

• Visual changes, flushing, muscle paralysis, and drowsiness

Measurement of serum magnesium levels is indicated to monitor magnesium toxicity if serum creatinine >1.0 mg/dL.

Management of magnesium toxicity is as follows:

• Stop magnesium therapy

• Estimation of serum magnesium and creatinine levels

• Intravenous injection of calcium gluconate 10 mL (10% solution)

The two commonly used $MgSO_4$ regimens are described below:

Repeated/maintenance dose IM injections of magnesium sulfate are continued only under the following conditions:

• The patellar reflex is present

• Respiration is not depressed

• Urine output >30 mL/h

Regimen	Loading Dose	Maintenance Dose
Intramuscular (Pritchard)	4 g (20% solution) IV over 3–5 minutes, followed by 10 g (50%) deep IM (5 g in each bu ttock)	5 g (50%) IM 4 hourly in the alternate buttock
Intravenous (Zuspan orSibai)	4 g IVslow over 15–20 minut e	1–2 g/h IV infusion

Solution to Question 16:

According to PCPNDT act, the radiologist should check for congenital anomalies of the fetus but not look for the sex.

Pre-Conception and Pre-Natal Diagnostic Techniques Act, 1994 is an act of the Parliament of India enacted to stop female foeticides and arrest the declining sex ratio in India.

The few basic requirements of the Act are:

Registration done under Section (18) of the PC-PNDT Act.

Written consent of the pregnant woman and there should be prohibition of communicating the sex of fetus under Section 5 of the Act.

Maintenance of records should be done as provided under Section 29 of the Act.

Creating public awareness at large by placing the board of prohibition on sex determination.

Initiatives Against Sex-Selective Abortions :

Beti Bachao Beti Padhao: Launched by the Government of India in 2015, it aims to address the issue of declining child, sex ratio and empower girls through awareness generation, education, and protection.

The National Plan of Action for Children, 2016 : It identifies the elimination of gender-biased sex selection as one of the key priority areas for children's rights and well-being.

Beti Bachao, Beti Padhao - "Save the daughter, educate the daughter" is a campaign of the Government of India that aims to generate awareness and improve the efficiency of welfare services intended for girls.

The objectives of BBBP are :

To prevent gender-biased sex-selective elimination

To ensure survival and protection of the girl child

To ensure education and participation of the girl child

Solution to Question 17:

The clinical scenario of secondary infertility with previous uneventful delivery and the laparoscopy image showing multiple granulomatous foci which are tuberculous lesions point towards the diagnosis of genital tuberculosis.

Genital tuberculosis is almost always secondary to a primary focus elsewhere. The most frequent location is the lungs (50%), although it can also come from lymph nodes (40%), kidneys, joints, the gastrointestinal system, or as part of a broad miliary infection. It mostly spreads via the haematogenous route or lymphatics sometimes.

In Genital tuberculosis, the leading presenting complaints include infertility, abdominal pain, vaginal discharge, and suspicion of neoplasm. 10-15% of women are found to be asymptomatic as well.

The fallopian tubes are the most common site to be affected. The disease subsequently spreads to other genital organs and the peritoneum.

Investigations:

1. Complete blood count, erythrocyte sedimentation rate (ESR), enzyme-linked immunoabsorbent assay (ELISA) tests - lgG and IgM, and Mantoux tests are done to evaluate the case of genital tuberculosis. The nucleic acid amplification technique (NAAT) can detect tuberculosis within a few hours compared to culture.

2. Hysterosalpingography (HSG) - If genital tuberculosis is suspected, HSG should not be performed because of the risk of spread of infection.

If performed in an asymptomatic woman, the following patterns may be seen:

• The tobacco-pouch appearance of hydrosalpinx and pyosalpinx with blocked fimbrial end.

• Rigid non-peristaltic pipe-like tube (lead-pipe appearance).

• Beading and variation in filling defect.

• The jagged fluffiness of the tubal outline.

• Golf club appearance with slight or moderate dilatation of the ampullary portion of the fallopian tube.

3. The investigation of choice in genital tuberculosis is an endometrial biopsy. Endometrium tissue is obtained by D&C/hystroscopy-directed biopsy. The endometrial scrapings are used for histopathology, PCR testing, and AFB smear and culture.

Diagnostic laparoscopy is widely used to establish the diagnosis of genital tuberculosis/ abdominal pelvic tuberculosis. Tuberculous lesions can be seen on the parietal peritoneum, intestinal serosa, omentum, surface of the uterus, and fallopian tubes.

Solution to Question 18:

Liquid-based cytology, Cervical biopsy, and HPV antigen testing (Option A) are the most appropriate investigations required in this patient with spotting, foul discharge, and cervical erosion which are the features suggestive of cervical cancer.

In liquid-based cytology, a smeared plastic spatula is placed in the liquid fixative (methanol) which removes blood, mucus, and inflammatory cells from epithelial cells. The epithelial cells are then sucked into a filter and pressed onto a glass slide to form a monolayer. This is done to screen patients for cervical cancer.

Differentiating between cervical erosion/ectropion and cervical intraepithelial neoplasia (CIN) and cervical cancer is difficult on macroscopic visualization, therefore cervical biopsy is required to rule out CIN and cervical cancer.

HPV antigen testing is required as HPV-16 is associated with anogenital warts and cervical carcinoma.

Pap/ Papanicolaou test is a screening test only. A positive test requires further evaluation - colposcopy, cervical biopsy, and fractional curettage should be done.

Solution to Question 19:

Image A shows 1st twin in the breech presentation and Image B shows 1st twin in the vertex presentation, Therefore the mode of delivery for images A and B would be LSCS and Vaginal delivery respectively. (Option B)

Timing of delivery depends on the type of twin pregnancy:

The mode of delivery depends on the 1st twin:

After delivery of 1st twin, if the 2nd twin is in breech, assisted breech delivery is done. If the 2nd twin is in a transverse lie, the internal podalic version is done under GA in the OT followed by breech extraction.

Complications in attempting vaginal delivery in non-vertex presentation (i.e) transverse and breech:

• Interlocking of twins

• Fetal head trapping above the cervix

• Umbilical cord prolapse

Type of twin pregnancy	Timing of delivery
Dichorionic diamniotic	38 weeks
Monochorionic diamniotic	34 - 37 weeks + 6 days
Monochorionic monoamniotic	32 - 34 weeks by Caesarean section

1st twin presentation	Mode of delivery
Vertex	Vaginal
Breech	Cesarean section
Transverse	Cesarean section



Solution to Question 20:

The earliest time to insert an IUCD is within 48 hours.

PPIUCD (postpartum IUCD) can be inserted:

(a) post-placental (within 10 minutes of placental delivery)

(b) within 48 hours postpartum

(c) during cesarean section

(d) 4-6 weeks after postpartum.

With PPIUCD, the chances of perforation are less, and the expulsion rate is slightly high.

The timing of the insertion of IUCD is as follows:

Interval insertion (inter-conceptional period)- preferable to insert 2–3 days after menstruation, but it can be inserted at any time during the cycle. It can also be inserted during menstruation which has certain advantages like an open cervical canal and a distended uterine cavity

It can also be inserted within 5 days of coitus as an effective method of emergency contraception (Eg: incase of ruptured condom during intercourse)

Solution to Question 21:

The drug of choice for the treatment of thyrotoxicosis during the first trimester of pregnancy is propylthiouracil (PTU). It is because it has lower teratogenic effects than carbimazole or methimazole.

Carbimazole and methimazole are associated with teratogenic effects such as:

• Aplasia cutis congenita

• Choanal atresia

• Treacheosopageal fistula

• Fetal goiter

However, propylthiouracil use is limited to the first trimester due to the risk of hepatotoxicity. It is replaced with carbimazole or methimazole in the 2nd and 3rd trimesters of the pregnancy.

The lowest effective antithyroid drug dose should be used throughout gestation. This is to maintain the maternal serum-free T4 level in the upper limit of the non-pregnant normal reference range.

Propranolol is a non-selective beta blocker. It is used in thyroid storm preferentially as it blocks the peripheral conversion of inactive T4 to active form T3.

Solution to Question 22:

In the clinical scenario, the smooth, globular structure shown during examination as well as the presentation of heavy menstrual bleeding suggests the diagnosis of myomatous polyp.

Myomatous Polyp, also known as a fibroid polyp, is a type of benign tumor originating from the smooth muscle cells of the uterus, called leiomyomas. These polyps can protrude from the endometrial cavity or cervix and may appear as a smooth, pinkish, round mass, which can sometimes be seen on physical examination, especially when they extend into the vagina.

Fibroids can cause symptoms like heavy menstrual bleeding (menorrhagia), pelvic pain, and pressure. These polyps may remain attached to the uterine wall via a stalk, which can give them a globular appearance.

Since myomatous polyps are associated with fibroids, they tend to occur in women of reproductive age and may persist into the perimenopausal period, as seen in this 42-year-old patient.

Other options:

Option A: Hypertrophied Cervix: A hypertrophied or enlarged cervix is not typically described as a smooth, globular mass. It usually does not present with a history of heavy bleeding.

Option C: Polypoid adenocarcinoma mainly occurs due to malignant changes of uterine or endometrial polyp in post-menopausal women. Endometrial polyps, another name for uterine polyps, are soft, microscopic growths that develop inside a woman's uterus, or womb. They originate from the endometrium, the tissue that lines the uterus.

Option D: Placental Polyp: Placental polyps are remnants of placental tissue and can cause abnormal bleeding but typically do not present with the smooth, globular structure.

Solution to Question 23:

The image shows Marshall Burns's technique used in the delivery of aftercoming head in breech delivery.

Technique:

• Baby is allowed to hang by its weight till the nape of the neck is visible

• When the nape of the neck is visible under the pubic arch, the baby is grasped by the ankles

• Baby is swung in an arc in an upward and forward direction towards the mother's abdomen



Other Options:

Option A: Pinard's manoeuvre used for delivery of extended legs in case of breech delivery.



Option B: The Mauriceau-Smellie-Veit technique is another method that can be used in the delivery of the aftercoming head in case of breech presentation.



Option D: The Ritgen manoeuvre is used for controlled delivery of the head in normal labour.



Solution to Question 24:

Transvaginal ultrasound findings of a left adnexal cystic mass, with incomplete septa and no vascular flow along with the complaints of lower abdominal pain suggest the diagnosis of hydrosalpinx.

A hydrosalpinx is a chronic cystic swelling of the fallopian tube that forms after a distal tubal obstruction. The cogwheel appearance and incomplete septa described in the ultrasound can correspond to the appearance of a hydrosalpinx. The lack of vascular flow on Doppler imaging supports this diagnosis. The tubes are enormously distended due to filling with clear serous fluid and become retort-shaped. On histology, the tubal plicae appear flattened. The interstitial end of the tube is open whereas the fimbrial end is indrawn and closed. This permits the dye to enter but

not exit the tube during HSG.

It is caused due to:

• Pelvic inflammatory disease

• Endometriosis

• Tuberculosis - tubercular salpingitis may also cause tobacco-pouch appearance, lead pipe appearance and pyosalpinx

• Fallopian tube neoplasms (rare)

It can be an incidental finding during a pelvic exam or ultrasound. The patient may be asymptomatic, or present with infertility or chronic pelvic pain.

On HSG, it appears as terminal sacculation and occlusion of both fallopian tubes with a normal uterine cavity. Usually, a thin-walled, hypoechoic, cystic structure is seen. However, it may appear as multiple hyperechoic mural nodules around the inner circumference of the tube creating a 'beads on a string' appearance.

Treatment:

• Treat the underlying cause

• Laparoscopic salpingostomy - in mild disease

• IVF recommended for women with infertility and severe hydrosalpinx



©MARROW

Other options:

Option B: Malignant ovarian cyst: Malignant ovarian cysts can appear as complex masses with irregular shapes and internal vascularity.

Option C: Follicular cyst: A follicular cyst is a common type of ovarian cyst that occurs when a follicle fails to release an egg. It typically appears as a simple cyst on ultrasound.

Option D: Hemorrhagic cyst: A hemorrhagic cyst results from bleeding into a pre-existing cyst and often shows complex internal echoes and potentially some vascular flow on Doppler imaging.

Solution to Question 25:

The most appropriate next step in management for a patient with bilateral tubal blockage on hysterosalpingography is laparoscopic chromopertubation.

Tubal contractions, particularly cornual spasms, can give an incorrect impression of proximal fallopian tube obstruction (a false-positive result) on hysterosalpingography. Laparoscopy with chromotubation is therefore considered to confirm the diagnosis of cornual spasm. Laparoscopy is both diagnostic and therapeutic in this condition.

Laparoscopic chromopertubation involves the laparoscopic visualization of pelvic structures, including the uterus, fallopian tubes, and ovaries, followed by the injection of methylene blue dye through the cervix to assess tubal patency. In addition to evaluating tubal patency, laparoscopy can help diagnose peritubal adhesions, endometriosis, and other pelvic abnormalities.

Chromotubation may be performed, in which a dilute dye is injected through a Leech-Wilkinson cannula placed against the cervix or through a balloon catheter positioned within the uterine cavity. Tubal spill is evaluated through the laparoscope. Indigo carmine dye, if available, is preferable to methylene blue, as methylene blue rarely may induce acute methemoglobinemia, particularly in patients with glucose-6-phosphate dehydrogenase deficiency. It is commonly done in the proliferative phase.

Other options:

Option A: Intrauterine insemination (IUI): IUI is used for treating infertility but relies on open fallopian tubes for natural fertilization.

Option B: In vitro fertilization (IVF): While IVF is an effective treatment for infertility due to tubal blockage, the immediate next step to confirm and evaluate the extent of tubal blockage is typically laparoscopic chromopertubation.

Option D: Ovulation induction: Ovulation induction aims to stimulate egg release but does not address tubal blockage.

Solution to Question 26:

The given image shows a perineal tear extending up to the rectal mucosa, therefore, it is a grade 4 perineal tear.

The involvement of rectal mucosa as shown in the image differentiates it from grade 3 tear. Grade 3 tear involves the vaginal mucosa, perineal skin, perineal muscles, and the anal sphincter complex (external and/or internal), but not rectal mucosa, while grade 4 tear involves all these layers including the rectal mucosa.

Minor perineal injuries are quite common, especially during the first birth. Gross injures like third or fourth degrees of perineal tears are a result of mismanagement of the second stage of labour. This can be prevented by proper conduct of the second stage labour.

Management is as follows:

All recent tears should be repaired immediately after the delivery of the placenta. If there is a delay of 24 hours, the repair has to be withheld and the patient has to be started on antibiotics to prevent infection. Complete perineal tears that were not repaired in the first 24 hours should be operated on between 3-6 months after delivery.

In 3 months of time, any local infection that might have been present would subside, the general condition of the patient will also improve. However, if the perineal tear is untreated for more than 6 months, dense scar tissue forms and making the surgery difficult.

Solution to Question 27:

The most common site for tubal ligation is the isthmus of the fallopian tube due to its uniform muscle thickness, which facilitates effective occlusion.

The isthmus, being a narrow segment closest to the uterine wall, provides an ideal location for surgical intervention. This site ensures a more reliable and efficient ligation, minimizing the risk of complications and maximizing the success rate of the procedure. The uniformity of the muscle layer at this location allows for a consistent occlusion technique. Common procedures for tubal ligation at this site include the Pomeroy method, where a loop of the tube is tied and excised, and other techniques like the Falope ring or the electrocoagulation method.

While the ampulla, which has a larger-diameter segment lateral to the isthmus, is also a part of the fallopian tube, it is less frequently chosen for tubal ligation. Additionally, ligation at the isthmus allows for the possibility of future reversal, should the patient desire to conceive again, making it a favourable choice for women considering sterilization.

The fallopian tube consists of three layers: the mesosalpinx, an outer mesothelial layer serving as visceral peritoneum; the myosalpinx, which has an inner circular smooth muscle layer for pressure during contractions and an outer longitudinal layer for tube movement; and the endosalpinx, an inner columnar epithelium with ciliated and secretory cells, featuring longitudinal folds. Together, these layers transport the ovum efficiently toward the uterine cavity.

Note: The interstitial portion of the fallopian tube is the narrowest segment, located within the uterine wall, and it forms the tubal ostia that opens into the endometrial cavity.

Solution to Question 28:

The appropriate treatment for a 30-year-old female with infravaginal elongation of the cervix is Fothergill surgery.

Fothergill surgery, or the Manchester operation, involves performing an anterior colporrhaphy combined with the amputation of the cervix. During the procedure, the Mackenrodt ligaments are sutured in front of the cervix, and the operation concludes with a colpoperineorrhaphy. This surgical approach is particularly suitable for women under 40 years old who wish to maintain reproductive function, although it's important to note that fertility may be reduced following the procedure.

Various other surgeries were performed for the management of uterovaginal prolapse:

• Anterior colporrhaphy: This is done for cystocele or cystourethrocele.

• Posterior colporrhaphy/ colpoperineorrhaphy: This is done for posterior defects like rectocele or rectal defects with perineal laxity.

• Le Forte's repair(option B): This is preferred only in elderly women with advanced prolapse, or those unfit for surgery. Women wanting an active sexual life are poor candidates for this procedure as it creates an adhesive area in the midline.

• Abdominal sling surgeries: Young women desirous of retaining menstrual and reproductive functions are ideal candidates. These surgeries use tapes as slings to prevent organ descent. These include:

• Abdominal wall cervicopexy (modified by Purandare and Mathre) - sling attached to cervix posteriorly near uterosacral ligaments.

• Shirodkar's sling surgery - the sling fixes the cervix to the lumbosacral fascia.

• Khanna's sling surgery - sling fixed to the anterior superior iliac spine, strengthening the Mackenrodt's ligament.

• Fothergill surgery (Manchester operation)

• Ward-Mayo surgery: This is a combination of vaginal hysterectomy + pelvic floor repair. It is done in women >40 years, not desiring reproductive function.

• Moschowitz surgery: This is performed for enterocele repair where the pouch of Douglas is obliterated.

Management of Vault prolapse:

• If the patient is fit for abdominal surgery: Transabdominal sacral colpopexy, i.e., suspending the vault to the sacral promontory extraperitoneally. It is best suited for younger women since coital difficulty following vaginal surgery is avoided

• If the patient is not fit for abdominal surgery: Right Transvaginal sacrospinous colpopexy. It is usually preferred in obese and elderly women, who are not fit for abdominal surgery.

• Colpocleisis is permissible if sexual activity is not desired.

Other options:

Option A: McCall culdoplasty is primarily used to prevent enterocele formation and provide support to the posterior vaginal apex, particularly following vaginal hysterectomy.

Option D: The Latzko procedure is a surgical technique used to treat vesicovaginal fistula. It involves removing the vaginal epithelium around the fistula and approximating the edges to promote healing.

Solution to Question 29:

In abdominal sling procedures such as sacral colpopexy, which is used to treat pelvic organ prolapse while preserving fertility, the mesh is typically attached to the sacral promontory. This site provides strong support for the prolapsed vaginal apex and allows for a durable suspension without significant compromise to the patient's fertility.

Various surgeries were performed for the management of uterovaginal prolapse:

• Anterior colporrhaphy: This is done for cystocele or cystourethrocele.

• Posterior colporrhaphy/ colpoperineorrhaphy: This is done for posterior defects like rectocele or rectal defects with perineal laxity.

• Le Forte's repair: This is preferred only in elderly women with advanced prolapse, or those unfit for surgery. Women wanting an active sexual life are poor candidates for this procedure as it creates an adhesive area in the midline.

• Abdominal sling surgeries: Young women desirous of retaining menstrual and reproductive functions are ideal candidates. These surgeries use tapes as slings to prevent organ descent. These include:

• Abdominal wall cervicopexy (modified by Purandare and Mathre) - sling attached to cervix posteriorly near uterosacral ligaments.

• Shirodkar's sling surgery - the sling fixes the cervix to the lumbosacral fascia.

• Khanna's sling surgery - sling fixed to the anterior superior iliac spine, strengthening the Mackenrodt's ligament.

• Fothergill surgery (Manchester operation): An anterior colporrhaphy is combined with amputation of the cervix. The Mackenrodt ligaments are then sutured in front of the cervix followed by a colpoperineorrhaphy. It is suitable for women <40 years desiring reproductive function, but fertility may be reduced.

• Ward-Mayo surgery: This is a combination of vaginal hysterectomy + pelvic floor repair. It is done in women >40 years, not desiring reproductive function.

• Moschowitz surgery: This is performed for enterocele repair where the pouch of Douglas is obliterated.

Management of Vault prolapse:

• If the patient is fit for abdominal surgery: Transabdominal sacral colpopexy, i.e., suspending the vault to the sacral promontory extraperitoneally. It is best suited for younger women since coital difficulty following vaginal surgery is avoided

• If the patient is not fit for abdominal surgery: Right Transvaginal sacrospinous colpopexy. It is usually preferred in obese and elderly women, who are not fit for abdominal surgery.

• Colpocleisis is permissible if sexual activity is not desired

Solution to Question 30:

This patient presenting with symptoms of cyclical pain at the episiotomy site with image showing perineal scar endometriosis has ectopic endometrial tissue. Endometriosis can occur at the site of a surgical scar and typically presents with cyclical pain that corresponds with the menstrual cycle.

The extra-uterine sites of endometrial tissue are influenced by cyclical changes in the hormone levels which give rise to location-specific symptoms as mentioned above.

Management:

Investigations include:

• USG: TVS will show a chocolate cyst.

• MRI

• CA-125 more than 35 U/mL

• Laparoscopy: Gold standard for the diagnosis and treatment of endometriosis. It is used to locate the sites, take a biopsy and surgically remove the deposits.

Drugs used for symptom control include the following:

• Oral contraceptive pills

• Mirena IUD

• GnRH analogs

• Letrozole

Surgical management includes:

Minimally invasive surgery may be used to destroy the tissue by cautery, laser vaporization, and lyse adhesions that have formed as a complication of endometriosis.

Major surgical measures include incision of chocolate cyst and removal of lining, salpingo-oophorectomy, hysterectomy, and unilateral or bilateral salpingo-oophorectomy.

Other options:

Option A: Retained Sutures: Retained sutures can cause localized pain and inflammation, but this is usually accompanied by visible signs of infection or irritation rather than cyclical pain linked to menstruation.

Option B: Melanoma: Melanoma is a type of skin cancer that would usually present as a suspicious, irregularly shaped lesion rather than cyclical pain at the site of a previous surgical scar.

Option C: Inclusion Cyst: Inclusion cysts are benign and can cause localized pain or discomfort. However, they typically do not present with pain that worsens specifically during menstruation.

Solution to Question 31:

The most likely diagnosis is this patient with two cervices on examination and two uterine domes as seen in the image is uterine didelphys. It is a congenital anomaly where there are two separate uterine cavities and two cervices, each with its own set of endometrial lining.

Uterine didelphys is commonly associated with renal anomalies such as unilateral renal agenesis, severe renal hypoplasia, etc, and is often associated with

• Recurrent pregnancy loss

• Preterm labor

• Malpresentations

• Prematurity

Per speculum examination will reveal two cervices in such women.

USG is a non-invasive, easily available, convenient, and inexpensive imaging modality. 3-D USG is sufficient to diagnose uterine didelphys without further imaging. The USG findings seen are:

• Two widely separated uterine horns

• A deep fundal indentation

MRI is used in cases where a definitive diagnosis cannot be made based on ultrasonography. Invasive modalities such as hysteroscopy and laparoscopy are reserved for patients with complicated Mullerian anomalies.

Other options:

Option A: Unicornuate Uterus with Rudimentary Horn: This condition involves a single functioning uterine horn and a non-functional rudimentary horn. It presents with a single cervix and an abnormal horn that may cause complications.

Option B: Arcuate Uterus: An arcuate uterus has a normal single uterine cavity with a small indentation at the top of the uterus, which is a minor congenital anomaly.

Option C: Bicornuate Uterus: A bicornuate uterus has two cavities within a single uterus but typically only one cervix. It may have a partial separation or indentation at the top of the uterus but would not present with two cervices.



Solution to Question 32:

The most appropriate course of action in this patient at 10 weeks intrauterine gestation is to Remove the IUCD immediately and continue pregnancy.

Intrauterine Devices in Pregnancy

• If IUD strings are visible and the pregnancy is less than 13 weeks, the IUD should be removed as soon as possible to prevent later septic abortion, premature rupture of the membranes, and premature birth .

• When the strings of the IUD are not visible, an ultrasound examination should be performed to localize the IUD and determine whether expulsion has occurred. If the IUD is present, there are three options for management:

• Therapeutic abortion .

• Ultrasound-guided intrauterine removal of the IUD.

• Continuation of the pregnancy with the device left in place

Note :

If the patient wishes to continue the pregnancy, ultrasound evaluation of the location of the IUD should be considered .

• If the IUD is not in a fundal location, ultrasound-guided removal using small alligator forceps may be successful.

• If the location is fundal, the IUD should be left in place. When pregnancy continues with an IUD in place, the patient must be warned of the symptoms of intrauterine infection and should be cautioned to seek care promptly for fever or flulike symptoms, abdominal cramping, or bleeding. At the earliest sign of infection, high-dose intravenous antibiotic therapy should be given and the pregnancy evacuated promptly.

Solution to Question 33:

Routine measurement of serum magnesium levels is not typically required unless clinical signs of toxicity (e.g., loss of deep tendon reflexes, respiratory depression, decreased urine output) are present.

$MgSO_4$ is used as an anticonvulsant in the management of eclampsia and as prophylaxis in severe preeclampsia. It can also be used as a tocolytic to stop preterm labor. The therapeutic level of magnesium in eclampsia is 4–7 mEq/L.

Patients must be monitored for magnesium toxicity:

• Loss of deep tendon reflexes (Option D)

• Decreased respiratory rate <12/min (Option C)

• Urine output <30 mL/h (Option B)

• Chest pain and heart block

• Visual changes, flushing, muscle paralysis, and drowsiness

Measurement of serum magnesium levels is indicated to monitor magnesium toxicity if serum creatinine >1.0 mg/dL.

Management of magnesium toxicity is as follows:

• Stop magnesium therapy

• Estimation of serum magnesium and creatinine levels

• Intravenous injection of calcium gluconate 10 mL (10% solution)

The two commonly used $MgSO_4$ regimens are described below:

Repeated/maintenance dose IM injections of magnesium sulfate are continued only under the following conditions:

• The patellar reflex is present

• Respiration is not depressed

• Urine output >30 mL/h

Regimen	Loading Dose	Maintenance Dose
Intramuscular (Pritchard)	4 g (20% solution) IVover 3–5 minutes, followed by10 g (50%) deep IM(5 g in each buttock)	5 g (50%) IM 4 hourlyin the alternate buttock
Intravenous (Zuspan orSibai)	4 g IVslow over 15–20 minute	1–2 g/h IVinfusion

Solution to Question 34:

The sign present in this girl is called Goodell's sign. Goodell's sign is a sign of early pregnancy, which describes the marked softening of the cervix in contrast to the non-pregnant state. It is seen as early as the 6th week.

Other options:

Option A: Hegar's Sign: Refers to the softening of the lower uterine segment during early pregnancy, felt during bimanual examination.

Option B: Braxton Hicks Sign: These are irregular, painless uterine contractions that can occur during pregnancy but are not associated with cervical changes.

Option C: Oslander's Sign: Osiander's sign describes the pulsations felt in the lateral fornix due to increased vascularity. It is seen around 8 weeks of gestation.

Solution to Question 35:

The procedure shown in the image is the external cephalic version (ECV). Being a primigravida (first pregnancy) is not a contraindication to ECV. The procedure can be performed in both primigravida and multigravida patients as long as other contraindications are not present.

ECV is recommended to manage transverse lie when the membranes are intact.

Prerequisites for ECV - Women who have reached 370/7 weeks of gestation. Multiparity, unengaged presenting part, nonanterior placenta, nonobese patient, and abundant amnionic fluid improve the chances of ECV.

Absolute contraindications - When vaginal delivery is contraindicated, e.g. placenta previa

Relative contraindications - Early labor, oligohydramnios or ruptured membranes, known nuchal cord, structural uterine abnormalities, fetal growth restriction, multifetal gestation, and prior

abruption of the placenta

Solution to Question 36:

The most likely diagnosis is endometrioma. The patient's chronic lower abdominal pain, which worsens during menstruation, and the ultrasound finding of a right adnexal mass with homogeneous echoes are highly suggestive of an endometrioma.

Endometriosis is defined by the presence of ectopic endometrial tissue at a site outside of the uterus. It is a benign condition but it is locally invasive. Some common sites of endometriosis are ovaries, pelvic peritoneum, pouch of Douglas, uterosacral ligaments, rectovaginal septum, and sigmoid colon. Endometriosis in the ovaries is called endometrioma.

Chocolate cysts of the ovaries represent the most important manifestation of endometriosis.

Patients with endometriosis mainly present with progressively increasing secondary dysmenorrhea, menorrhagia, infertility, dyspareunia, and chronic pelvic pain.

MRI is the diagnostic tool, it shows the characteristic hyperintensity on T1 weighted images and hypointensity on T2 weighted images. Laparoscopy confirms the diagnosis.

Medical management of endometriosis:

• Gestrinone is an anti-estrogenic, anti-progestational, and mildly anabolic drug that reduces sex hormone-binding globulin to release free testosterone. It also decreases the LH surge. It causes regression of lesions and amenorrhea. It is a 19-nortestosterone derivative similar to danazol but with fewer side effects.

• Combined oral contraceptives are used for symptom control and alleviating disease. These also cause menstrual cycle regulation and delay pregnancy.

• Oral progestogens exert an anti-estrogenic effect and cause decidualization and endometrial atrophy. This causes disease regression. It doesn't prevent ovulation, thus suitable for women trying to conceive.

• Mirena IUCDs reduce dysmenorrhea and menorrhagia. It is a single-sitting treatment with minimal side effects.

• Aromatase inhibitors like letrozole are anti-estrogens given for disease regression.

• Anti-inflammatory drugs like mefenamic acid and naproxen relieve dysmenorrhea.

Therapeutic procedures:

• Aspiration of peritoneal fluid from the cul-de-sac removes PGE2, relieves dysmenorrhea, and pelvic pain, and improves fertility.

• Destruction of endometriotic lesions <3cm with diathermy, Nd-YAG laser, or CO2 vaporization can be attempted.

• Larger lesions and chocolate cysts can be excised.

• Laparoscopic breaking of adhesions relieves dysmenorrhea and chronic pelvic pain. It also restores tubal patency and improves ovulation. Pre-sacral neurectomy can be done simultaneously.

• Laser uterosacral nerve ablation (LUNA) is done for pelvic pain.



Laparoscopic image of chocolate cyst of ovary:



Solution to Question 37:

The most likely diagnosis is bacterial vaginosis.

Nugent's criteria are used to evaluate three types of bacteria in a case of suspected bacterial vaginosis: Mobiluncus, Lactobacillus, and Gardnerella/Bacteroides. Bacteria are graded through the gram stain of vaginal samples.

They are each graded on a scale of 1+ to 4+ where:

• 1+ is < 1 cell per field

• 2+ is 1-5 cells per field

• 3+ is 6-30 cells per field

• 4+ is >30 cells per field

Lactobacillus and Bacteroides/Gardnerella are given scores between 0-4, but Mobiluncus is graded only from 0-2. The total scores are then calculated and interpreted as follows:

• 0-3 - Normal

• 4-6 - Intermediate bacterial count

• 7-10 - Bacterial vaginosis

Solution to Question 38:

After ovulation, the released egg is arrested in meiosis II.

The released egg is in the secondary oocyte stage, having completed meiosis I. It then enters meiosis II, but this process is arrested at metaphase II until fertilization occurs. If fertilization takes place, meiosis II will be completed; otherwise, the secondary oocyte remains arrested in meiosis II.

Oogenesis is the process wherein oogonia differentiate into mature oocytes. It begins before birth. At birth, there are approximately 2 million primordial follicles. Primordial germ cells divide by mitosis to form oogonia.

The majority of the oogonia divide by mitosis, however, some enter meiosis I and get arrested in the diplotene stage of prophase and are now called primary oocytes. These primary oocytes remain arrested in this manner until puberty. Primary oocytes haven't completed meiosis, hence they're diploid. The primary oocyte along with surrounding epithelial cells forms the primordial follicle.

Meiosis I is completed after LH surge during puberty to form the secondary oocyte. The cell then enters meiosis II but arrests in metaphase approximately 3 hours before ovulation.

If fertilization does not occur, the secondary oocyte fails to complete the second meiotic division and degenerates about 24 hours after ovulation. If fertilization occurs, penetration by the sperm signals the secondary oocyte to complete meiosis II. This results in one mature oocyte (ovum) and another smaller cell called the second polar body that is shed off.



Solution to Question 39:

The given clinical scenario as well as the specimen suggests that this is a case of leiomyoma uteri or fibroid uterus. The specimen is of a leiomyoma.

Gross specimen appearance:

• The uterine body/corpus is enlarged and deformed.

• The cut surface shows multiple nodules, well-circumscribed, firm, and white-greyish with a whorled appearance formed by the intersecting bundles of brown muscle fibres and white fibrous tissue.

• Leiomyomas can also present bleeding areas, cystic degeneration, and calcifications. The uterine cavity is compressed.

• A typical myoma is a well-circumscribed tumor with a pseudo capsule and firm inconsistency. The cut surface has a whorled appearance.

• The vessels that supply blood to the tumor lie in the capsule and send radial branches into the tumor.

• Therefore, the central portion of the tumour receives the least blood supply and degeneration is noticeable early in this part of the tumour.

• On the other hand, calcification begins at the periphery and spreads inwards along the vessels.

Risk factors:

Increased risk of leiomyomas is seen in nulliparous women, those with obesity, and individuals in a hyper-estrogenic state, such as in conditions like endometriosis, endometrial hyperplasia, or endometrial carcinoma. Black women are also at higher risk.

Conversely, the risk is decreased in multiparous women. Recent studies have shown no significant correlation between smoking and the development of leiomyomas.

Symptoms of fibroid uterus:

Most women with leiomyomas are asymptomatic, but those affected may experience bleeding, pain, pressure, or infertility. Symptom risk increases with the size and number of myomas. Heavy menstrual bleeding (HMB) is common, linked to dilated endometrial venules and dysregulated vasoactive growth factors. When these engorged venules are disrupted during menstruation, bleeding can overwhelm normal hemostatic mechanisms, causing HMB.

Treatment:

Medical therapy for fibroids includes mifepristone, GnRh analogs, Aromatase inhibitors

Drugs to reduce the size of the fibroid

• Antiprogesterone – Mifepristone/RU-486

• Danazol (Usually not used due to androgenic side effects)

• GnRH analogs

• Agonists – goserelin, leuprorelin, nafarelin, buserelin

• Antagonists – cetrorelix, ganirelix

• Levonorgestrel releasing intrauterine system

• Selective progesterone receptor modulator (SPRM's) – Asoprisnil

Other management options

• Uterine artery embolization

• Myomectomy

• Hysterectomy

Other Options:

Option B: Myomatous Polyp: They are typically found on the endometrial lining and would not present as a fibroid embedded in the wall of the uterus.

Option C: Endometrioma: An endometrioma is an ovarian cyst filled with endometrial tissue. It would not be found within the uterine wall but rather on the ovaries.

Option D: Adenomyoma: Adenomyoma is a localized area of endometrial tissue within the myometrium (muscle layer) of the uterus. While it can cause similar symptoms, it does not typically present as a discrete mass protruding from the uterine wall like an intramural fibroid.

Solution to Question 40:

The most likely diagnosis is endometrioid carcinoma. It is the most common type of endometrial cancer, particularly associated with genetic mutations such as PTEN, KRAS, CTNNB1 (beta-catenin), and MSI (microsatellite instability). These genetic alterations are characteristic of Type I endometrial cancers, which have a better prognosis compared to other types.

The histopathological examination (HPE) shows glandular structures that resemble normal endometrial glands but are more crowded and irregular. The cells show nuclear atypia with mitotic activity.

This cancer is generally hormone-dependent (Type I endometrial cancer) and often carries a better prognosis compared to other forms of endometrial cancer. The presence of PTEN and KRAS mutations are indicators of this type of malignancy, which often grows slowly and is diagnosed early due to the symptoms of postmenopausal bleeding.

Endometrial cancer occurs in a state of estrogen excess and is therefore mostly seen in:

• nulliparous women

• early menarche

• late menopause

• PCOS

• HRT

• estrogen-producing tumors

• Hypertension, diabetes, and obesity (adipose tissue produces estrogen E1) together form the corpus cancer syndrome.

Endometrial biopsy is the investigation of choice and if positive should be followed by hysteroscopy with fractional curettage (gold standard) for assessing grade and histology.

Other options:

Option A: Mixed Müllerian Tumor: Also known as carcinosarcoma, this tumor shows both epithelial and mesenchymal components. It is less commonly associated with the mutations listed (PTEN, KRAS, etc.) and typically presents with more aggressive features than endometrioid carcinoma.

Option B: Leiomyosarcoma: This is a malignant tumor of smooth muscle, typically arising from the myometrium, and is not associated with genetic mutations (PTEN, KRAS, CTNNB1).

Option D: Serous Cystadenoma of Ovary: A benign ovarian tumor, this condition would not typically involve the genetic markers found in endometrial carcinoma and would present differently from endometrial cancer. Serous tumors of the ovary are more often associated with TP53 mutations, not PTEN or KRAS.

Solution to Question 41:

Misoprostol and mifepristone are administered during first trimester to induce medical termination of pregnancy.

200mg of mifepristone is administered orally on day 1. The minimum recommended interval between mifepristone & misoprostol is 24 hours. 24- 48 hours later, misoprostol 800 mcg is administered by the vaginal, sublingual, or buccal route. Clinical confirmation of MTP is done on day 14.

The various methods of termination of pregnancy are:

Misoprostol and Methotrexate regimen can also be used, but take a longer time of action.

The Medical Termination of Pregnancy Amendment Bill, 2020 has extended the period of termination of pregnancy up to 24 weeks for special categories of women, including survivors

of rape, victims of incest, and other vulnerable women (like differently-abled women, minors). The upper gestation limit is not applicable in cases of substantial fetal abnormalities.

Timing	Medical methods	Surgical methods
1st trimester(up to 12 weeks)	MifepristoneMifepristoneand misoprostolMethotrexate and misoprostolTamoxifen and misoprostol	Menstrual regulationVacuum aspiration (MVA/EVA)Suction evacuation and/or curettageDilatation and evacuation - Rapid methodSlow method
2nd trimester(13 - 20 weeks)	MisoprostolMifepristone and misoprostolDinoprostone(option A)Carboprost	Dilatation and evacuationIntrauterine instillationofhyperosmotic solutions(option D)Oxytocin infusion(option C)Hysterotomy

Solution to Question 42:

The embryo takes approximately 8 days to travel from point 1 (fertilization) to point 8 (implantation) (typically 6-7 days).

Timeline of implantation:

• 3-4 days after fertilization: The morula (12-16 cells) enters the uterus.

• 4th day after fertilization: The advanced morula reaches the uterine lumen.

• 4-5 days after fertilization: The early blastocyst stage is seen.

• 6-7 days after fertilization: The blastocyst implants in the uterus. Implantation corresponds to 20-22 days of the menstrual cycle.

The blastocyst normally implants in the functional layer of the uterine endometrium. In the secretory phase, the endometrium comprises three layers: stratum compactum, stratum spongiosum, and stratum basale. The first two layers, collectively known as the stratum functionale or functional layer, are crucial for implantation. If implantation does not occur, this functional layer is shed off during menstruation. However, if implantation occurs, the functional layer is retained as the decidua, marking the beginning of a successful pregnancy.





Solution to Question 43:

For women with active genital herpes at the time of labor, elective cesarean section is the preferred mode of delivery to prevent neonatal herpes transmission. A standard lower segment cesarean section (LSCS) is typically performed unless there are other complications necessitating a different approach.

HSV infection in pregnancy can be caused by either HSV-1 or HSV-2 and transmitted by sexual contact. The infection can be primary, nonprimary, first episode, and recurrent.

HSV infection (both HSV 1 and 2) during pregnancy has a risk of vertical transmission of the infection to the fetus. The risk is higher in the primary than in the recurrent infection. The maximum risk of fetal infection is during labor, although it can occur in utero and post-natally from direct contact.

Suspicious lesions should be sent for PCR testing or culture. Antiviral treatment with acyclovir should be given if found positive. Women with no active genital lesions, asymptomatic, or those with active lesions in non-genital areas can undergo normal labor and delivery.

Management of HSV infection during pregnancy:

• Oral acyclovir or valacyclovir for primary or recurrent infections during pregnancy.

• IV acyclovir followed by oral acyclovir for severe or disseminated infection

• Oral analgesics or topical anesthetics for pain relief

• Catheterization for urinary retention

Cesarean delivery reduces the risk of vertical transmission during labor and is indicated in:

• Cases of active genital lesions - both primary and recurrent

• Prodromal symptoms like vulval itching or burning

Other options:

Option B: Classical Cesarean Section: Classical cesarean section involves a vertical incision on the upper part of the uterus and is generally reserved for specific cases such as abnormal uterine anatomy or placenta previa

Option C and D: Trial of Labor (VBAC) and Induction of Labor: Vaginal birth after cesarean (VBAC) and induction of labor are contraindicated in women with active genital herpes due to the risk of transmitting the virus to the newborn during vaginal delivery.

Solution to Question 44:

In cases of adnexal masses, particularly when considering the possibility of ovarian cancer or germ cell tumors, the following tumor markers are useful:

Other options:

CEA (Carcinoembryonic antigen): Used for gastrointestinal cancer.

CA15-3: Marker associated with breast cancer.

CA19-9: Marker associated with pancreatic and gastrointestinal cancers

Tumor markers	Ovarian tumors
AFP (alpha-fetoprotein)	Yolk sac tumor
LDH (lactate dehydrogenase)	Dysgerminoma
HCG (human chorionic gona dotropin)	Choriocarcinoma
CA125	Epithelial cell tumor

Solution to Question 45:

The image shows Piper's forceps, identified by its distinct perineal curve. It is used to assist in the delivery of the after-coming head of breech.

It has a cephalic curve, reverse pelvic curve, and long parallel shanks that permit the baby's body to rest against it during head delivery. It may be applied electively or when the Mauriceau maneuver cannot be accomplished easily.

Other options:

Option A: Wrigley forceps: These are short forceps generally used for low or outlet deliveries but are not suitable for delivering the after-coming head in breech deliveries.

Option B: Kielland forceps: These are rotational forceps used to assist in malpositioned heads during cephalic deliveries.

Option C: Das forceps: These forceps are used for cephalic deliveries.