Ophthalmology Collection

Question Bank with Answers & Explanations

Generated on April 18, 2026

34 Topics • 749 Questions

Need help or updates?
■ *Click here to contact the creator of this PDF*

Viewer Guide

Navigation Tips:

- Use the outline/bookmarks pane (left side in most PDF viewers) to jump between sections.
- Click links at the bottom of each question to switch between Part A (questions) and Part B (answers).
- Master Index links take you to topic parts.

Viewer Settings:

- Adobe Reader: View > Show/Hide > Navigation Panes > Bookmarks
- Chrome: Install PDF viewer extension if needed.
- Mobile: Use apps like Adobe Acrobat or Xodo for outline support.

■ *For queries & support: Message @Neet_pg_bot on Telegram*

DIRECTORY EXPLORER

Click a folder below to open it.

» **01_Orbit**

» **02_Diseases of Lens**

» **03_Uveitis**

» **04_Glaucoma**

» **05_Retina**

» **06_Neuro Ophthalmology**

» **07_Squint _ Strabismus**

» **08_Cornea**

» **09_Conjunctiva**

» **10_Ocular Adnexa**

» **11_Optics**

» **12_Basics of Ophthalmology**

» **13_Sclera**

» **14_Lacrimal Drainage System**

» **15_Community Ophthalmology**

» **16_Ocular Injuries**

» 01_Orbit

« Back to Parent Directory

Path: 01_Orbit

• 001 Orbit - I	*Questions*	*Answers*
• 002 Previous Year Questions	*Questions*	*Answers*

» 02_Diseases of Lens

« Back to Parent Directory

Path: 02_Diseases of Lens

• 003 Lens And Disorders - I	***Questions***	***Answers***
• 004 Previous Year Questions	***Questions***	***Answers***

» 03_Uveitis

« Back to Parent Directory

Path: 03_Uveitis

• 005 Uvea	***Questions***	***Answers***
• 006 Previous Year Questions	***Questions***	***Answers***

» 04_Glaucoma

« Back to Parent Directory

Path: 04_Glaucoma

• 007 Glaucoma-Pacg Treatment Congenital Glaucoma	***Questions***	***Answers***
• 008 Glaucoma - Poag Tonometry Perimetry	***Questions***	***Answers***
• 009 Previous Year Questions	***Questions***	***Answers***

» 05_Retina

« Back to Parent Directory

Path: 05_Retina

• 010 Retinal Investigations And Vascular Disorders	***Questions***	***Answers***
• 011 Macular And Fundus Disorders Retinal Detachment And Retinoblastoma	***Questions***	***Answers***
• 012 Previous Year Questions	***Questions***	***Answers***

» 06_Neuro Ophthalmology

« Back to Parent Directory

Path: 06_Neuro Ophthalmology

• 013 Neuro Ophthalmology	***Questions***	***Answers***
• 014 Previous Year Questions	***Questions***	***Answers***

» 07_Squint _ Strabismus

« Back to Parent Directory

Path: 07_Squint _ Strabismus

• 015 Squint	***Questions***	***Answers***
• 016 Previous Year Questions	***Questions***	***Answers***

» 08_Cornea

« Back to Parent Directory

Path: 08_Cornea

• 017 Cornea- I	***Questions***	***Answers***
• 018 Previous Year Questions	***Questions***	***Answers***

» 09_Conjunctiva

« Back to Parent Directory

Path: 09_Conjunctiva

• 019 Conjunctiva - I	***Questions***	***Answers***
• 020 Previous Year Questions	***Questions***	***Answers***

» 10_Ocular Adnexa

« Back to Parent Directory

Path: 10_Ocular Adnexa

• 021 Lids And Disorders - I	***Questions***	***Answers***
• 022 Previous Year Questions	***Questions***	***Answers***

» 11_Optics

« Back to Parent Directory

Path: 11_Optics

• 023 Optics-Ii	***Questions***	***Answers***
• 024 Optics-I	***Questions***	***Answers***
• 025 Previous Year Questions	***Questions***	***Answers***

» 12_Basics of Ophthalmology

« Back to Parent Directory

Path: 12_Basics of Ophthalmology

• 026 Introduction To Ophthalmology	***Questions***	***Answers***
• 027 Previous Year Questions	***Questions***	***Answers***

» 13_Sclera

« Back to Parent Directory

Path: 13_Sclera

• 028 Sclera	***Questions***	***Answers***
• 029 Previous Year Questions	***Questions***	***Answers***

» 14_Lacrimal Drainage System

« Back to Parent Directory

Path: 14_Lacrimal Drainage System

• 030 Lacrimal Drainage System	Questions	Answers
• 031 Previous Year Questions	Questions	Answers

» 15_Community Ophthalmology

« Back to Parent Directory

Path: 15_Community Ophthalmology

	Questions	*Answers*
• 032 Previous Year Questions		

» 16_Ocular Injuries

« Back to Parent Directory

Path: 16_Ocular Injuries

• 033 Ocular Injuries	***Questions***	***Answers***
• 034 Previous Year Questions	***Questions***	***Answers***

Part A — All Questions

• [01_Orbit] 001 Orbit - I

« Back to Folder Directory

[01_Orbit] 001 Orbit - I — Quick Jump

Q1 (p. 20)	*Q2 (p. 20)*	*Q3 (p. 21)*	*Q4 (p. 21)*	*Q5 (p. 22)*	*Q6 (p. 22)*
Q7 (p. 22)	*Q8 (p. 23)*	*Q9 (p. 23)*	*Q10 (p. 24)*	*Q11 (p. 25)*	*Q12 (p. 25)*
Q13 (p. 25)	*Q14 (p. 26)*	*Q15 (p. 27)*	*Q16 (p. 27)*	*Q17 (p. 28)*	*Q18 (p. 28)*
Q19 (p. 29)	*Q20 (p. 29)*	*Q21 (p. 29)*	*Q22 (p. 30)*	*Q23 (p. 31)*	*Q24 (p. 31)*
Q25 (p. 31)	*Q26 (p. 32)*	*Q27 (p. 32)*			

Q1: A 9-year-old boy presents with a 2-day history of swelling, redness, and pain around his left eye, worsening with eye movements. He had a recent upper respiratory infection. On examination, there is proptosis, restricted eye movement, and chemosis. Vision is intact. CT scan confirms orbital cellulitis. Which staging system is used to classify this condition? (Q853270)

A. Chandler's classification

B. Van Herick classification

C. ETDRS classification

D. Shaffer grading

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 45-year-old patient presents with a large, painless orbital mass that has been causing progressive proptosis and visual disturbance. After imaging and biopsy, the mass is identified as malignant. What is the choice of treatment in this scenario, and what is its definition? (Q853283)

A. Evisceration, it involves removal of the contents of the eyeball, sclera preserved

B. Enucleation, it involves removal of the contents of the eyeball, sclera preserved

C. Enucleation, it involves removal of the entire eyeball including the sclera.

D. Exenteration, it involves the removal of all orbital contents

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 12-year-old patient presents to the OPD with a typical S-shaped deformity of the eyelid. On examination, the patient is found to have multiple café-au-lait spots and axillary freckling. Which of the following findings is most commonly observed in this condition? (Q853282)

A. Early-onset cataracts

B. Lisch nodules

C. B/L Acoustic neuroma

D. Optic nerve sheath meningiomas

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 6-year-old child presents with gradual, painless unilateral proptosis and vision loss. Fundoscopic examination reveals optic atrophy. Imaging studies show fusiform growth along the optic nerve. What is the most appropriate management for this child, considering the tumor is stationary and the vision is still present? (Q853281)

A. Immediate surgical excision of the tumor

B. Radiotherapy to manage the tumor

C. Observation and regular monitoring

D. Chemotherapy as the first-line treatment

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 52-year-old woman presents with progressive, painless proptosis of the left eye and swelling in the temporal region of the face. On examination, she has boggy eyelid swelling, proptosis. A CT scan reveals hyperostosis of the sphenoid bone. Identify the false statement regarding the same. (Q853280)

A. CT scan shows tubular enlargement of the optic nerve

B. They are more common in middle-aged women.

C. More common in NF II

D. They are slow-growing tumors from astrocytes in the optic nerve.

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Which of the following is not part of the classical triad of Sturge-Weber syndrome? (Q853279)

A. Port-wine stain (nevus flammeus) in the distribution of the trigeminal nerve

B. Secondary congenital glaucoma

C. Bilateral optic nerve glioma

D. Ipsilateral meningeal hemangiomatosis

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 7-year-old male child presents with rapidly progressing proptosis in the right eye, which starts suddenly and mimics an acute inflammatory process. Imaging shows a well-defined orbital mass with adjacent bone destruction. What is the most common

histopathological subtype of this tumor? (Q853278)

A. Alveolar sarcoma

B. Embryonal sarcoma

C. Botryoid sarcoma

D. Spindle cell sarcoma

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 50-year-old woman presents with painless proptosis of the right eye for the past year. She denies any visual changes, pain, or other ocular symptoms. On examination, there is mild proptosis. MRI reveals a well-defined, intraconal mass with homogeneous enhancement, suggestive of a cavernous hemangioma. What is the most appropriate management for this patient? (Q853277)

A. Surgical excision

B. Radiotherapy

C. Steroid injections

D. Observation and regular follow-up

See Answer in Part B → Q8

@Neet_pg_bot

Q9: Which among the following statements is/are true regarding orbital varices? Proptosis worsens with the Valsalva maneuver. It is thin-walled distensible high-flow veins. Venous stasis can lead to the formation of phlebolith. Endovascular embolization is contraindicated. (Q853276)

A. 1 and 3

B. 1, 3, and 4

C. Only 2

D. All of the above

See Answer in Part B → Q9

@Neet_pg_bot


Q10: A 35-year-old man presents with a progressive painless swelling in the superolateral part of his left orbit over the past year. Examination reveals mild proptosis and downward and inward displacement of the eye. CT of the orbit shows a well-circumscribed, smooth, encapsulated mass in the lacrimal gland region. Excisional biopsy is shown below. Which of the following is the most likely diagnosis? (Q853275)

A. Adenoid cystic carcinoma

B. Squamous cell carcinoma

C. Pleomorphic adenoma

D. Lymphoma



See Answer in Part B → Q10

@Neet_pg_bot

Q11: Which is the most common non-epithelial malignant lacrimal gland tumor? (Q853274)

A. Pleomorphic adeno-carcinoma

B. Adenoid cystic carcinoma

C. Lymphoma

D. Mucoepidermoid carcinoma

See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 45-year-old male presents with double vision, drooping of the right eyelid, and difficulty moving his right eye in multiple directions. On examination, the patient has normal visual acuity, ptosis, restricted eye movement, decreased sensation over the forehead, and absent corneal reflex on the same side. What is the most likely diagnosis and which cranial nerve is/are most likely affected in this condition? (Q853273)

A. Superior orbital fissure syndrome; III, IV, V, and VI nerves

B. Orbital apex syndrome; optic nerve

C. Tolosa Hunt syndrome; II, III and IV nerves

D. Rochen-Duvigneaud syndrome; only III nerve

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 45-year-old diabetic male presents with complaints of headache, nasal congestion, and facial pain for the past week. On examination, there is periorbital swelling, proptosis, and black eschar on the nasal mucosa. His blood sugar levels are poorly controlled. CT scan of the paranasal sinuses reveals extensive sinus

involvement, and biopsy confirms the diagnosis of rhino-orbital-cerebral mucormycosis (ROCM). Which of the following is NOT a known complication of rhino-orbital-cerebral mucormycosis? (Q853272)

A. Meningitis

B. Brain abscess

C. Retinal vascular occlusion

D. Subarachnoid hemorrhage

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A 50-year-old male, recently recovered from COVID-19, presents with complaints of severe facial pain, nasal congestion, and black eschar on the left eye with periorbital pain. He has a history of uncontrolled diabetes mellitus. On examination, the following is seen. Which of the following is the most likely causative organism? (Q853271)

A. Aspergillus species

B. Rhizopus species

C. Candida species

D. Cryptococcus neoformans



See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 32-year-old male had blunt trauma to the face during a sports event. He reports difficulty moving his right eye upward and experiences double vision when looking up. On examination, there is tenderness at the right infraorbital rim, and the patient exhibits noticeable swelling and bruising around the right eye. Which wall of the orbit is most likely affected in this case? (Q853256)

A. Medial wall

B. Lateral wall

C. Floor

D. Roof

See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 10-year-old girl presents with a swollen, painful left eye that started after a recent cold. She is febrile and on examination, her left eyelid is erythematous and swollen, with chemosis, proptosis, limited ocular movement but normal visual acuity.

What is the most common primary source of this eye infection in the child? (Q853269)

A. Ethmoid sinusitis

B. Maxillary sinusitis

C. Frontal sinusitis

D. Otitis media

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 25-year-old male presents with intermittent bulging of his right eye, worsening with cough and bending forward. His vision is intact, and there is no history of trauma or infection. What is the most likely cause of his condition? (Q853268)

A. Cavernous sinus thrombosis

B. Orbital varix

C. Thyroid orbitopathy

D. Orbital cellulitis

See Answer in Part B → Q17

@Neet_pg_bot

Q18: A 48-year-old man presents with a complaint of eye protrusion. He describes experiencing a rhythmic, pulsatile sensation in his right eye for the past few weeks. Which of the following is NOT a likely cause ? (Q853267)

A. Carotico-cavernous fistula

B. Saccular aneurysm of the ophthalmic artery

C. Congenital meningocele

D. Orbital cellulitis

See Answer in Part B → Q18

@Neet_pg_bot

Q19: If a patient with thyroid eye disease exhibits stable symptoms with no acute inflammation and has experienced a gradual improvement in symptoms over the past 1-2 years, which stage of Rundle's Curve is most likely being described? (Q853266)

A. Initial Phase

B. Peak Phase

C. Improvement Phase

D. Chronic Phase

See Answer in Part B → Q19

@Neet_pg_bot

Q20: A 48-year-old woman with Graves' disease presents with worsening proptosis, diplopia, diminution of vision and difficulty closing her eyes despite treatment with antithyroid medications and corticosteroids. On examination afferent pupillary defects are observed. CT scan shows orbital congestion and muscle enlargement. What is the most appropriate next step in management? (Q853265)

A. Adalimumab

B. Systemic steroids

C. Radiotherapy

D. Decompression Surgery

See Answer in Part B → Q20

@Neet_pg_bot

Q21: A 52-year-old woman with a history of thyroid disease presents with difficulty closing her eyes, noticeable eye protrusion and prominent bilateral lid retraction giving a staring and frightened appearance of the eyes as shown in the image below. Which of the following best describes the sign observed in this patient? (Q853264)

A. Von Graefe's Sign

B. Kocher’s sign

C. Stellwag's Sign

D. Joffroy's Sign



See Answer in Part B → Q21

@Neet_pg_bot

Q22: A 45-year-old woman with a history of thyroid disease presents with bilateral eye bulging and difficulty closing her eyes. On examination, there is noticeable eye protrusion, eyelid swelling, and restricted extraocular movements. Which of the following best describes the pathogenesis of her eye condition? (Q853263)

A. Extraocular muscles are the primary target of inflammation.

B. The earliest defective ocular movement seen in this patient is depression.

C. T-cells attack the orbital fibroblasts and adipocytes in the orbit.

D. Inflammation is primarily due to neutrophils.

See Answer in Part B → Q22

@Neet_pg_bot

Q23: Which of the following is not a cause of pseudoproptosis? (Q853262)

A. High Myopia

B. Eyelid Retraction

C. Contralateral Enophthalmos

D. Orbital Cellulitis

See Answer in Part B → Q23

@Neet_pg_bot

Q24: A 34-year-old woman with a history of hyperthyroidism presents with bilateral proptosis and difficulty closing her eyes. To assess the degree of eye protrusion, a Hertel's exophthalmometer is used. Which of the following statements regarding the use of the exophthalmometer is correct? (Q853261)

A. Normal values for exophthalmometer readings are typically between 20 and 30 mm.

B. Difference of 2–3 mm or more between the two eyes is suspicious regardless of the absolute values

C. A reading greater than 10 mm in both eyes is indicative of proptosis.

D. Hertel's exophthalmometer can only be used to measure the degree of proptosis in one eye at a time.

See Answer in Part B → Q24

@Neet_pg_bot

Q25: Which condition is most commonly associated with bilateral proptosis in adults? (Q853260)

A. Orbital cellulitis

B. Thyroid ophthalmopathy

C. Rhabdomyosarcoma

D. Neuroblastoma

See Answer in Part B → Q25

@Neet_pg_bot

Q26: Which surgical space is typically used for administering a retrobulbar injection? (Q853259)

A. Sub-Tenon Surgical Space

B. Extraconal Surgical Space

C. Intraconal Surgical Space

D. Subarachnoid Surgical Space

See Answer in Part B → Q26

@Neet_pg_bot

Q27: Which of the following statements is TRUE regarding the orbital foramina and fissures? (Q853258)

A. Superior orbital fissure is between the medial and lateral walls.

B. Inferior orbital fissure is between the lesser wing of the sphenoid and the maxillary bone.

C. Optic foramen is between the lateral wall and the orbital floor.

D. Optic nerve and ophthalmic artery pass through the optic foramen.

See Answer in Part B → Q27

• [01_Orbit] 002 Previous Year Questions

« Back to Folder Directory

[01_Orbit] 002 Previous Year Questions — Quick Jump

Q1 (p. 34) *Q2 (p. 35)* *Q3 (p. 35)* *Q4 (p. 35)* *Q5 (p. 36)* *Q6 (p. 36)*
Q7 (p. 37) *Q8 (p. 37)* *Q9 (p. 38)* *Q10 (p. 38)*


Q1: A 15-year-old female presents with a painless, gradually increasing mass located lateral to the lateral canthus, present for the past 10 years. On examination, the mass is non-tender and slowly progressive. What is the most likely diagnosis? (Q866349)

A. Dermoid cyst

B. Capillary hemangioma

C. Lacrimal gland tumor

D. Epidermoid cyst



See Answer in Part B → Q1

@Neet_pg_bot

Q2: Which of the following is true about orbital cellulitis? (Q866363)

A. It is present anterior to the orbital septum

B. Treated effectively with topical antibiotics

C. Presents with proptosis, orbital swelling, normal pupil, and extraocular movements

D. Ethmoid sinusitis is the most common etiology

See Answer in Part B → Q2

@Neet_pg_bot

Q3: What is the most common cause of proptosis in a middle-aged person? (Q853127)

A. Orbital varix B) Thyroid eye disease C) Retinoblastoma D) Cavernous hemangioma

B. Thyroid eye disease

C. Retinoblastoma

D. Cavernous hemangioma

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A patient presents with proptosis that increases when bending down. What is the most likely diagnosis? (Q852503)

A. Orbital Varices

B. Thyroid Eye Disease

C. Cavernous Sinus Thrombosis

D. Orbital Cellulitis

See Answer in Part B → Q4

@Neet_pg_bot


Q5: Which of the following statements is false regarding the condition shown in the image? (Q844005)

A. NOSPEC classification is used

B. Management corresponds to an improvement in the thyrotoxic state

C. Vision loss may occur

D. It is not seen in 10% of patients with hyperthyroidism



See Answer in Part B → Q5

@Neet_pg_bot

Q6: What is the most common cause of proptosis in a middle-aged person? (Q840073)

A. Orbital varix

B. Thyroid eye disease

C. Retinoblastoma

D. Cavernous hemangioma

See Answer in Part B → Q6

@Neet_pg_bot

Q7: Following instrument is used to measure? (Q854708)

A. Proptosis

B. Squint

C. Ptosis

D. Diplopia



See Answer in Part B → Q7

@Neet_pg_bot

Q8: A middle-aged woman presented with bilateral proptosis, restriction of eye movements, and chemosis. Her thyroid profile showed she was hyperthyroid. What is the most probable cause? (Q858505)

A. Orbital pseudotumor

B. Grave's ophthalmopathy

C. Orbital lymphoma

D. Orbital cellulitis

See Answer in Part B → Q8

@Neet_pg_bot

Q9: A 35-year-old female reports recurring episodes of discomfort, pain, and a sensation of pressure behind her left eye. She exhibits intermittent reducible proptosis, and her symptoms intensify when she bends forward. What is the most likely diagnosis? (Q830803)

A. Orbital hemangioma

B. Orbital varices

C. Orbital metastasis

D. Orbital dermoid cyst

See Answer in Part B → Q9

@Neet_pg_bot

Q10: What is the most common primary intraocular malignancy in adults? (Q830808)

A. Retinoblastoma

B. Malignant Choroidal melanoma

C. Reticulum cell sarcoma

D. Neurofibroma

See Answer in Part B → Q10

@Neet_pg_bot

• [02_Diseases of Lens] 003 Lens And Disorders - I

« Back to Folder Directory

[02_Diseases of Lens] 003 Lens And Disorders - I — Quick Jump

Q1 (p. 39)	*Q2 (p. 39)*	*Q3 (p. 41)*	*Q4 (p. 41)*	*Q5 (p. 41)*	*Q6 (p. 42)*
Q7 (p. 42)	*Q8 (p. 44)*	*Q9 (p. 44)*	*Q10 (p. 45)*	*Q11 (p. 46)*	*Q12 (p. 47)*
Q13 (p. 47)	*Q14 (p. 48)*	*Q15 (p. 48)*	*Q16 (p. 49)*	*Q17 (p. 49)*	*Q18 (p. 50)*
Q19 (p. 50)	*Q20 (p. 50)*	*Q21 (p. 51)*	*Q22 (p. 52)*	*Q23 (p. 52)*	*Q24 (p. 53)*
Q25 (p. 53)	*Q26 (p. 54)*	*Q27 (p. 55)*	*Q28 (p. 56)*	*Q29 (p. 56)*	*Q30 (p. 57)*
Q31 (p. 57)	*Q32 (p. 58)*	*Q33 (p. 58)*	*Q34 (p. 58)*	*Q35 (p. 59)*	*Q36 (p. 59)*
Q37 (p. 60)	*Q38 (p. 60)*	*Q39 (p. 60)*	*Q40 (p. 61)*	*Q41 (p. 61)*	

Q1: What kind of laser can be utilized to treat posterior capsular opacification, which is observed during an examination of a patient who has reported decreased visual acuity following cataract surgery? (Q882185)

A. Femto laser

B. Nd YAG laser

C. Argon fluoride

D. Argon

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A newborn presented with the following findings. There is a positive maternal history for fever during the first trimester of pregnancy. Which among the following is least likely to be seen in this newborn? (Q853341)

A. Patent ductus arteriosus

B. Limb hypoplasia

C. Microphthalmia

D. Sensorineural hearing loss





See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which of the following is not a condition associated with a shallow anterior chamber? (Q853331)

A. Wound leak with hypotony

B. Ciliochoroidal detachment

C. Vitreous bulge

D. Iridodialysis

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 67-year-old man had undergone cataract surgery. He presents with endophthalmitis after 3 days. His infection is severe with vision reduced to hand movements close to the face. What is the most appropriate management? (Q853332)

A. Intravitreal antibiotics

B. Systemic antibiotics

C. Steroid therapy

D. Vitrectomy

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 65-year-old man has undergone cataract surgery. He has now returned after 6 weeks with endophthalmitis. What is the most likely organism that caused the infection? (Q853333)

A. Staphylococcus aureus

B. Streptococcus pyogenes

C. Propionibacterium acnes

D. Staphylococcus epidermidis

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Soemmering's Ring and Elschnig's Pearls are examples of which of the following? (Q853334)

A. After cataracts

B. Senile cataracts

C. Congenital cataracts

D. Toxic cataracts

See Answer in Part B → Q6

@Neet_pg_bot


Q7: Match the images of IOL malpositions with their names: Image Syndrome a. 1. Sunrise syndrome b. 2. Lost lens syndrome c. 3. Sunset syndrome (Q853335)

A. a-1,b-2,c-3


B. a-3,b-1,c-2

C. a-2,b-1,c-3


D. a-1,b-3,c-2



@Neet_pg_bot



@Neet_pg_bot



See Answer in Part B → Q7

@Neet_pg_bot

Q8: Which of the following statements is true regarding Cystoid Macular Edema? (Q853336)

A. Collection of fluid in Henle’s layer

B. Honeycomb appearance in fundoscopy

C. Antiprostaglandin is both treatment and prevention

D. All of the above

See Answer in Part B → Q8

@Neet_pg_bot

Q9: What is the common inheritance pattern of hereditary cataracts? (Q853337)

A. Autosomal dominant

B. Autosomal recessive

C. X linked recessive

D. X linked dominant

See Answer in Part B → Q9

@Neet_pg_bot


Q10: Doctor noticed leukocoria in a 3-month-old baby during a routine examination and the eye appeared as shown below. Choose the least appropriate statement regarding the most likely diagnosis. (Q853338)

A. Most common type of bilateral congenital cataract

B. Commonly inherited as autosomal dominant trait

C. Usually zone of fetal nucleus is involved

D. Associated with vitamin-D deficiency



See Answer in Part B → Q10

@Neet_pg_bot


Q11: An infant with leukocoria presents with recurrent gastrointestinal infections and developmental regression. Bilateral eye examination revealed the following. Choose the most appropriate next step in evaluation. (Q853339)

A. Serum estimation of glucose, calcium, and phosphorous levels

B. Estimation of antibody titers for infections

C. Urine chromatography for amino acids

D. Urine assay for the presence of reducing substances



See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 1-month-old infant is diagnosed with a dense unilateral cataract. Which of the following is the most appropriate management strategy? (Q853340)

A. Delay surgery until the child is 1–2 years old to reduce surgical risks

B. Perform immediate cataract surgery

C. Prescribe glasses and monitor the cataract progression

D. Wait until the child develops strabismus before considering surgery

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 70-year-old woman with a history of hypertension and high myopia undergoes cataract surgery. During the procedure, she suddenly experiences severe ocular pain and a rapid rise in intraocular pressure. Fundoscopic examination reveals a dark red elevation in the choroidal region. Which of the following is the most appropriate immediate management step? (Q853330)

A. Intraoperative posterior sclerostomy

B. Increase the infusion pressure in the anterior chamber

C. Perform immediate vitrectomy to drain the hemorrhage

D. Close the surgical wound and monitor intraocular pressure

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A school-going child is diagnosed with a connective tissue disorder characterized by abnormalities of the eye, short stature, brachydactyly, joint stiffness, and cardiovascular defects. Which among the following is incorrect regarding the likely diagnosis? (Q853342)

A. Follows autosomal recessive mode of inheritance

B. Associated with supero-temporal lens dislocation

C. Molecular genetic analysis is confirmatory

D. Usual age of presentation is 5-10 years

See Answer in Part B → Q14

@Neet_pg_bot

Q15: During early ocular development, the optic vesicle induces the formation of the lens placode from adjacent tissue. What is the embryologic origin of the crystalline lens? (Q885161)

A. Neural crest cells

B. Surface ectoderm

C. Neuroectoderm

D. Mesoderm

See Answer in Part B → Q15

@Neet_pg_bot

Q16: The fibers that are most recently formed in the crystalline lens are located in which region? (Q885162)

A. Lens cortex

B. Embryonic nucleus

C. Fetal nucleus

D. Adult nucleus

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 30-year-old patient is being evaluated during a slit-lamp examination. The examiner explains that the crystalline lens is avascular and depends on surrounding structures for metabolic exchange. What is the primary source of nutrition for the lens? (Q885164)

A. Connective tissue diffusion

B. Zonular fibers

C. Aqueous humor

D. Retinal circulation

See Answer in Part B → Q17

@Neet_pg_bot

Q18: A histology slide of the crystalline lens shows a thin capsule with a single layer of cuboidal cells beneath it. The marked region indicates lens epithelium. At which location is this structure normally absent? (Q885166)

A. Posterior capsule region

B. Central anterior surface

C. Equatorial lens region

D. Anterior pole

See Answer in Part B → Q18

@Neet_pg_bot

Q19: In the crystalline lens, the antioxidants ascorbate and alpha-tocopherol remain in their reduced (active) form primarily due to which intracellular substance? (Q885169)

A. Glucose

B. Glycoprotein

C. Glutathione

D. Fatty acids

See Answer in Part B → Q19

@Neet_pg_bot

Q20: A 9-year-old child is brought in by his parents because teachers report poor classroom performance. The child says he cannot clearly see the blackboard and struggles with reading. There is no history of white reflex noticed at birth. Ophthalmic examination reveals a bilateral, disc-shaped opacity involving a particular layer of the lens with relatively clear cortex and nucleus. Which type of congenital cataract most commonly presents this way? (Q885171)

A. Cerulean (bluedot) cataract

B. Lamellar (zonular) cataract

C. Posterior polar cataract

D. Total congenital cataract

See Answer in Part B → Q20

@Neet_pg_bot


Q21: An 18-year-old girl presents with gradual blurring of vision and glare. Slit-lamp examination shows multiple, fine, polychromatic, needle-shaped refractile crystals in the lens cortex, giving a sparkling appearance. Which systemic association is most likely? (Q885175)

A. Alport syndrome

B. Myotonic dystrophy

C. Type 1 diabetes mellitus

D. Lowe syndrome



See Answer in Part B → Q21

@Neet_pg_bot

Q22: Match the type of cataract with the etiology Type of cataract Etiology a. Posterior subcapsular 1. Glassblowers b. Anterior subcapsular-granular 2. Miotics c. Posterior cortical 3. Steroids (Q853319)

A. a1,b2,c3

B. a3,b2,c1

C. a2,b1,c3

D. a2,b3,c1

See Answer in Part B → Q22

@Neet_pg_bot

Q23: What is the consequence of age-related decline in the Na+-K+ ATPase pump in the lens? (Q853310)

A. Increased intralenticular K+ levels and decreased Na+ levels.

B. Increased intralenticular Na+ levels and decreased K+ levels.

C. Enhanced ion exchange efficiency.

D. Reduced intralenticular Ca2+ levels.

See Answer in Part B → Q23

@Neet_pg_bot

Q24: Which among the following is/are true statements regarding lens proteins? α/β/γ-Crystallins are water-soluble proteins Major intrinsic proteins contribute to lens transparency An increase in crystallins can lead to cataract β/γ-Crystallin family are the largest crystallins (Q853311)

A. Only 1

B. 1 & 2

C. 2, 3 & 4

D. 1, 2, 3 & 4

See Answer in Part B → Q24

@Neet_pg_bot


Q25: A 70-year-old male patient comes to your clinic with complaints of glare and diminution of vision for 3 months. On examination, you notice the following appearance Choose the correct statement regarding the correct diagnosis. (Q853312)

A. The Iris shadow test becomes positive only when the whole cortex is involved.

B. Patients see better in dark surroundings.

C. Characterized by saucer-shaped opacities in the center of the posterior cortex.

D. Vision loss occurs at a late stage.



See Answer in Part B → Q25

@Neet_pg_bot


Q26: An 80-year-old female patient came with complaints of gradual progressive diminution of vision for 2 years in both her eyes. She reports that over the last year, she noticed an improvement in her ability to read without glasses. The following finding is seen on slit lamp examination over the right eye and the left eye has similar opacity. Choose the most appropriate course of management for this patient. (Q853313)

A. Advise lifestyle modification and review after 3 months.

B. Recommend corticosteroids eye drops and reassess in 6 months.

C. Plan for Phacoemulsification.

D. Prescribe her rigid gas-permeable contact lenses.



See Answer in Part B → Q26

@Neet_pg_bot

Q27: An elderly male diabetic patient came with loss of vision in the left eye associated with pain in the left eye. On examination, the left eye appears red, and visual acuity- Perception of light+ and IOP is raised in the same eye. The patient gives history of blurring and diminution of vision for the past 1 year. The slit lamp examination is given below. Choose the correct diagnosis and appropriate pathophysiology for this patient. (Q853314)

A. Phacomorphic glaucoma - Secondary open-angle glaucoma

B. Phacomorphic glaucoma - Secondary closed-angle glaucoma

C. Phacolytic glaucoma - Secondary open-angle glaucoma

D. Phacolytic glaucoma - Secondary closed-angle glaucoma



See Answer in Part B → Q27

@Neet_pg_bot

Q28: A 45-year-old man is having difficulty in near vision. The ophthalmologist has prescribed him a convex lens and informed him that it is an age-related physiological condition. Which of the following is an age-related change that would have caused his condition? (Q853315)

A. Decrease in elasticity of lens capsule

B. Sclerosis of lens substance

C. Reduced ciliary muscle power

D. All of the above

See Answer in Part B → Q28

@Neet_pg_bot

Q29: A 6-month-old girl was diagnosed with galactosemia. An ocular examination has revealed a cataract which developed as a complication. Which of the following

regarding her cataract is false? (Q853316)

A. Bilateral

B. Oil droplet appearance

C. Periphery of the lens

D. May be reversed

See Answer in Part B → Q29

@Neet_pg_bot

Q30: Which of the following is not a feature of chalcosis? (Q853317)

A. Kayser-Fleischer ring

B. Sunflower cataract

C. Retinal golden plaques

D. Oil droplet cataract

See Answer in Part B → Q30

@Neet_pg_bot

Q31: A 30-year-old male with retinitis pigmentosa has developed a cataract as a complication. Which of the following is a feature of his cataract? Posterior subcapsular cortical cataract Irregular in outline Variable density Breadcrumb appearance Polychromatic luster Diffuse yellow haze in the adjoining cortex (Q853318)

A. 1,2,3,6

B. 2,4,5,6

C. 1,2,3,4

D. 1,2,3,4,5,6

See Answer in Part B → Q31

@Neet_pg_bot

Q32: Which of the following is the thinnest part of the lens capsule? (Q853309)

A. Anterior pole

B. Posterior pole

C. Equator

D. Pre-equatorial region

See Answer in Part B → Q32

@Neet_pg_bot

Q33: A 35-year-old man presents with decreased vision in his right eye following a blunt trauma a few weeks ago. On examination, feathery opacities are noted. What is the most likely type of cataract this patient has developed? (Q853320)

A. Subepithelial opacities

B. Diffuse concussion cataract

C. Rosette cataract

D. Morgagnian cataract

See Answer in Part B → Q33

@Neet_pg_bot

Q34: A 38-year-old patient with diabetes presents with rapidly worsening vision. Upon examination, the ophthalmologist notes the presence of bilateral, snowflake-like white opacities in the cortex of the lens. Which of the following is the most likely diagnosis? (Q853321)

A. Senile cataract

B. Christmas tree cataract

C. Diabetic cataract

D. Posterior subcapsular cataract

See Answer in Part B → Q34

@Neet_pg_bot

Q35: Which of the following steps is used to maintain the depth of the anterior chamber and protect the corneal endothelium during Conventional ECCE? (Q853323)

A. Anterior capsulotomy

B. Viscoelastic injection

C. Corneoscleral section completion

D. Hydrodissection

See Answer in Part B → Q35

@Neet_pg_bot

Q36: A 70-year-old woman presents for cataract surgery in her right eye. She has moderate cataract density with some nuclear sclerosis and a clear cornea. She expresses a strong desire for a procedure that minimizes recovery time and provides optimal visual outcomes. Which of the following statements accurately reflects the advantage of Phacoemulsification Over Manual SICS? (Q853324)

A. Manual SICS typically requires a smaller incision compared to Phacoemulsification.

B. Manual SICS allows for faster visual recovery compared to Phacoemulsification.

C. Manual SICS is less suitable for very hard cataracts compared to Phacoemulsification.

D. Manual SICS usually results in a higher level of postoperative astigmatism compared to Phacoemulsification.

See Answer in Part B → Q36

@Neet_pg_bot

Q37: What are the steps involved in FLACS? (Q853325)

A. Capsulorrhexis → Corneal Incision → Lens Fragmentation → IOL implantation

B. Corneal Incision → Capsulorrhexis → Lens Fragmentation → IOL implantation

C. Capsulorrhexis → Lens Fragmentation → Corneal Incision → IOL implantation

D. Corneal Incision → Lens Fragmentation → Capsulorrhexis → IOL implantation

See Answer in Part B → Q37

@Neet_pg_bot

Q38: A key advantage of peribulbar anesthesia over retrobulbar anesthesia is: (Q853326)

A. Faster onset of anesthesia

B. Less risk of globe perforation

C. Better akinesia of the eye

D. Reduced risk of systemic toxicity

See Answer in Part B → Q38

@Neet_pg_bot

Q39: Toric IOLs are specifically designed to correct which type of refractive error? (Q853327)

A. Myopia

B. Hypermetropia

C. Astigmatism

D. Presbyopia

See Answer in Part B → Q39

@Neet_pg_bot

Q40: A 55■year■old man underwent myopic LASIK 15 years ago. He now develops senile cataract and undergoes phacoemulsification with posterior chamber IOL implantation. Postoperatively, he complains of poor distance vision, and refraction shows an unexpected hyperopic error of +2.5 D. Which of the following IOL power calculation formulas would have prevented this postoperative refractive error? (Q853328)

A. SRK

B. Hoffer Q

C. Haigis

D. Holladay I

See Answer in Part B → Q40

@Neet_pg_bot

Q41: In which scenario is a posterior capsule tear more commonly observed during cataract surgery? (Q853329)

A. During phacoemulsification with high energy settings

B. During cortical clean-up with a manual irrigation-aspiration technique

C. During hydrodelineation

D. During anterior chamber maintenance with a viscoelastic substance

See Answer in Part B → Q41

@Neet_pg_bot

• [02_Diseases of Lens] 004 Previous Year Questions

« Back to Folder Directory

[02_Diseases of Lens] 004 Previous Year Questions — Quick Jump

Q1 (p. 62)	*Q2 (p. 63)*	*Q3 (p. 63)*	*Q4 (p. 64)*	*Q5 (p. 65)*	*Q6 (p. 65)*
Q7 (p. 66)	*Q8 (p. 66)*	*Q9 (p. 67)*	*Q10 (p. 67)*	*Q11 (p. 68)*	*Q12 (p. 68)*
Q13 (p. 69)	*Q14 (p. 72)*	*Q15 (p. 72)*	*Q16 (p. 73)*	*Q17 (p. 73)*	*Q18 (p. 74)*
Q19 (p. 74)	*Q20 (p. 74)*	*Q21 (p. 75)*	*Q22 (p. 75)*	*Q23 (p. 76)*	*Q24 (p. 77)*
Q25 (p. 78)	*Q26 (p. 78)*	*Q27 (p. 79)*			


Q1: A 65-year-old diabetic woman presents with painless loss of vision. The ocular findings are shown in the image below. What is the likely diagnosis and management for this patient? (Q855468)

A. Zonular cataract and lensectomy

B. Rosette cataract and sugar control

C. Sunflower cataract and lesion extraction

D. Immature senile cataract and phacoemulsification with IOL



See Answer in Part B → Q1

@Neet_pg_bot

Q2: Riya was found to have a height of more than the 95th percentile for her age and an arm-span to height ratio of more than 1.05. On further examination, she had flat feet, mild scoliosis, and long slender fingers. An image of her showing characteristic signs is provided below. What is the most common ocular complication expected in this child? (Q860007)

A. Hypertensive retinopathy

B. Rosette cataract

C. Retinitis pigmentosa

D. Subluxation of lens



See Answer in Part B → Q2

@Neet_pg_bot

Q3: What is the embryological origin of the corneal endothelium? (Q859756)

A. Neuroectoderm

B. Neural crest cells

C. Mesoderm

D. Surface ectoderm

See Answer in Part B → Q3

@Neet_pg_bot


Q4: Identify the given instrument (Q860184)

A. Universal retractor

B. Doyen’s speculum

C. Mouth clamp

D. Eye speculum



See Answer in Part B → Q4

@Neet_pg_bot

Q5: Which of the following best defines blindness ? (Q865357)

A. Visual acuity less than 6/60 in the better eye

B. Visual acuity less than 3/60 in the better eye after best possible correction

C. Visual field less than 10 degrees from fixation in the better eye

D. Inability to perceive light in both eyes

See Answer in Part B → Q5

@Neet_pg_bot


Q6: What is the diagnosis? (Q856030)

A. Intraocular foreign body

B. Pseudoexfoliation syndrome

C. Ocular trauma

D. Vossius ring



See Answer in Part B → Q6

@Neet_pg_bot

Q7: What is the average degree of hypermetropia in a newborn? (Q859755)

A. +2.5 D

B. +5 D

C. +10 D

D. +1 D

See Answer in Part B → Q7

@Neet_pg_bot

Q8: What could be the most probable cause for the progressive vision loss in both eyes of a 25-year-old male who has been undergoing low-dose oral steroid therapy for the past decade? (Q859416)

A. Cataract

B. Glaucoma

C. Cystoid macular edema

D. Retinal detachment

See Answer in Part B → Q8

@Neet_pg_bot

Q9: True statement regarding KF ring: Seen in most patients with neurological involvement Is pathognomonic for Wilson's disease Resolves with desferroxamine Seen in all patients with hepatic involvement Seen in superior and inferior descemets membrane (Q856412)

A. 1 and 5

B. 1,2 and 5

C. 1,2,3,4 and 5

D. 2,3,4 and 5

See Answer in Part B → Q9

@Neet_pg_bot

Q10: What is the formula utilized to calculate the power of the lens during cataract surgery for a patient who has previously undergone LASIK for myopia? (Q856929)

A. SRK 1

B. SRK 3

C. Hoffer Q

D. Haigis L

See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 60-year-old patient with progressive painless loss of vision presents to the ophthalmology OPD. The fundus image of the patient is given below. What could be the probable finding and the cause of it? (Q856121)

A. Soft exudate, hypertension

B. Hard exudate, diabetes mellitus

C. Flame-shaped hemorrhages, hypertension

D. Soft exudate, central retinal vein occlusion



See Answer in Part B → Q11

@Neet_pg_bot


Q12: Identify the surgical step shown in the image given below: [(Q856716)

A. Hydrodissection

B. Capsulorrhexis

C. Intraocular lens implantation

D. Lens aspiration



See Answer in Part B → Q12

@Neet_pg_bot


Q13: Match the given images to their descriptions: 1. A. Posterior subcapsular cataract 2. B. Sunflower cataract 3. C. Morgagnian cataract 4. D. Snowflake cataract (Q857189)

A. 1A 2D 3B 4C


B. 1D 2B 3A 4C


C. 1B 2C 3D 4A

D. 1C 2B 3D 4A


@Neet_pg_bot


@Neet_pg_bot





See Answer in Part B → Q13

@Neet_pg_bot

Q14: What substance is responsible for the development of a diabetic cataract in a 68-year-old female patient, which has been present for 3 months? (Q858160)

A. Glucose

B. Fructose

C. Lactose + Glucose

D. Sorbitol

See Answer in Part B → Q14

@Neet_pg_bot


Q15: An elderly patient presents with white, dandruff-like deposits on the anterior lens surface, seen during slit-lamp examination. What is the most likely diagnosis? (Q866322)

A. Pseudoexfoliation syndrome

B. Glaucoma fleckens

C. Phacolytic glaucoma

D. Pigment dispersion syndrome



See Answer in Part B → Q15

@Neet_pg_bot

Q16: What dye is used in staining for the anterior capsule of lens ? (Q820469)

A. Trypan blue

B. Brilliant blue

C. Fluorescent green

D. Lissamine green

See Answer in Part B → Q16

@Neet_pg_bot

Q17: In which type of cataract surgery is suturing with 10-0 nylon commonly performed? (Q820459)

A. Phacoemulsification

B. Extracapsular cataract extraction (ECCE)

C. Intracapsular cataract extraction (ICCE)

D. Manual small incision cataract surgery (MSICS)

See Answer in Part B → Q17

@Neet_pg_bot

Q18: Which lens is utilized in direct gonioscopy? (Q857414)

A. Goldmann

B. Koeppe

C. Zeiss

D. Sussman

See Answer in Part B → Q18

@Neet_pg_bot

Q19: Which is the type of cataract seen following a blunt trauma? (Q860165)

A. Rosette cataract

B. Christmas tree cataract

C. Snowflake cataract

D. Sunflower cataract

See Answer in Part B → Q19

@Neet_pg_bot

Q20: Oil drop cataract is seen in? (Q860164)

A. Galactosemia

B. Wilson's disease

C. Gaucher's disease

D. Tay-Sachs disease

See Answer in Part B → Q20

@Neet_pg_bot

Q21: After cataract surgery, a patient develops endophthalmitis with vision limited to hand movements. What is the treatment? (Q860543)

A. Intravitreal antibiotic injections only

B. Pars plana vitrectomy +Intravitreal antibiotic injections

C. Redo surgery

D. Topical antibiotics

See Answer in Part B → Q21

@Neet_pg_bot


Q22: Which of the following cataracts is seen in blunt trauma? (Q860514)

A. Rosette cataract

B. Toxic cataract

C. Metabolic cataract

D. Nuclear cataract



See Answer in Part B → Q22

@Neet_pg_bot

Q23: A 2-month-old child is brought to the ophthalmology OPD with the following presentation. What is the ideal management? (Q844026)

A. Medical management

B. Operate immediately

C. Surgery after 6 months

D. Surgery after 2 years



See Answer in Part B → Q23

@Neet_pg_bot


Q24: A patient presents with decreased vision three years after surgery and exhibits this condition. What is the most likely diagnosis? (Q844003)

A. Pseudo-bullous keratopathy

B. Corneal dystrophy

C. Posterior subcapsular cataract

D. Posterior capsular opacification (PCO)



See Answer in Part B → Q24

@Neet_pg_bot

Q25: A patient with a long-standing history of diabetes presents with cataracts. Which of the following metabolic products is primarily responsible for cataract formation in this condition? (Q852482)

A. Mannitol

B. Galactitol

C. Sorbitol

D. Fructose

See Answer in Part B → Q25

@Neet_pg_bot

Q26: What is Anisometropia? (Q853113)

A. Difference in visual acuity

B. Difference in refractive error between 2 eyes

C. Difference in the axial length of the eyeball

D. Difference in the size of the eyeball

See Answer in Part B → Q26

@Neet_pg_bot

Q27: A 15-year-old tall boy with long limbs presents to the OPD. On ocular examination, bilateral ectopia lentis is noted. Which gene is most likely affected in this inherited disorder? (Q866368)

A. COL5A

B. PAX6

C. TGFβR2

D. FBN1 (Fibrillin-1)

See Answer in Part B → Q27

@Neet_pg_bot

• [03_Uveitis] 005 Uvea

« Back to Folder Directory

[03_Uveitis] 005 Uvea — Quick Jump

Q1 (p. 80)	*Q2 (p. 80)*	*Q3 (p. 81)*	*Q4 (p. 81)*	*Q5 (p. 82)*	*Q6 (p. 83)*
Q7 (p. 84)	*Q8 (p. 84)*	*Q9 (p. 84)*	*Q10 (p. 85)*	*Q11 (p. 86)*	*Q12 (p. 86)*
Q13 (p. 87)	*Q14 (p. 87)*	*Q15 (p. 88)*	*Q16 (p. 88)*	*Q17 (p. 89)*	*Q18 (p. 89)*
Q19 (p. 89)	*Q20 (p. 90)*	*Q21 (p. 90)*	*Q22 (p. 91)*	*Q23 (p. 92)*	

Q1: A 42-year-old African American woman presents with visual disturbances. Eye examination reveals vitreous cells, snowball opacities, and candle wax drippings. Which type of uveitis is seen in the underlying disease? (Q853682)

A. Anterior uveitis

B. Panuveitis

C. Posterior uveitis

D. Peripheral multifocal chorioretinitis

See Answer in Part B → Q1

@Neet_pg_bot


Q2: A patient came with redness and pain, and you dilated the pupil to break the synechiae. Which drug was used, looking at this picture? (Q882305)

A. Atropine

B. Phenylephrine

C. Scopolamine

D. Miotics



See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which of the following conditions can cause panuveitis? (Q853710)

A. VKH syndrome

B. Sympathetic Ophthalmia

C. Sarcoidosis

D. All of the above

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 45-year-old male presents with vision loss in his right eye following a traumatic injury that resulted in a penetrating ocular wound two months ago. Recently, he has

developed redness, photophobia, and decreased vision in his left eye as well. On examination, both eyes show signs of bilateral granulomatous uveitis. Which of the following best explains the pathogenesis of the patient's condition? (Q853709)

A. Type I hypersensitivity reaction to lens proteins

B. Delayed hypersensitivity reaction to melanocytes

C. Direct bacterial infection spreading from the injured eye

D. Immune complex deposition in the retinal vasculature

See Answer in Part B → Q4

@Neet_pg_bot


Q5: A 28-year-old woman diagnosed with Vogt–Koyanagi–Harada (VKH) disease, Fundus examination shows the following findings. What does it indicate? (Q853708)

A. Prodromal phase

B. Uveitic phase

C. Convalescent phase

D. Recurrent phase



See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 35-year-old male presents with progressively worsening blurry vision, eye pain, and redness in the right eye. Examination reveals restricted eye movement, and fundoscopic evaluation identifies a well-defined cystic lesion in the vitreous humor, accompanied by surrounding inflammation. What is the most probable diagnosis in this patient? (Q853707)

A. Ocular toxoplasmosis

B. Endophthalmitis

C. Ocular cysticercosis

D. Posterior uveitis due to tuberculosis

See Answer in Part B → Q6

@Neet_pg_bot

Q7: Which of the following ocular conditions is most commonly associated with a positive Amsler sign? (Q853706)

A. Cataract

B. Ocular histoplasmosis syndrome

C. Fuchs heterochromatic uveitis

D. Acute retinal necrosis

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 35-year-old male with a recent travel history to Africa presents with intense itching, skin nodules, and progressive vision loss. On examination, multiple palpable nodules are found on his skin, and there is early corneal clouding. Microfilariae are observed in a skin snip biopsy. What is the most appropriate treatment for this condition? (Q853705)

A. Clindamycin + sulfadiazine

B. Diethylcarbamazine

C. Ivermectin

D. Self-limiting condition, no treatment needed

See Answer in Part B → Q8

@Neet_pg_bot


Q9: A 32-year-old immunocompromised woman living with cats presents with decreased vision in her left eye. On examination, she has mild anterior uveitis with vitritis and the fundoscopy image is shown below. Which of the following is the most likely causative organism? (Q853687)

A. Cytomegalovirus (CMV)

B. Candida albicans

C. Toxoplasma gondii

D. Histoplasma capsulatum



See Answer in Part B → Q9

@Neet_pg_bot

Q10: Match bacterial uveitis in column A with corresponding clinical manifestation in column B Bacterial uveitis Clinical features 1. Tubercular uveitis a. Iris pearls 2. Leprotic uveitis b. Iris roceola 3. Spirochetal uveitis c. Headlight in fog appearance d. Koeppe's nodules (Q853686)

A. 1-d, 2-a, 3-b

B. 1-b, 2-c, 3-d

C. 1-c, 2-d, 3-a

D. 1-b, 2-a, 3-d

See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 34-year-old woman with a history of Crohn's disease presents with sudden onset of eye pain, redness, and blurred vision. Slit-lamp examination reveals fine, grey keratic precipitates, anterior chamber cells, and flare. Which type of uveitis is most consistent with her presentation? (Q853684)

A. Granulomatous uveitis

B. Non-granulomatous anterior uveitis

C. Posterior uveitis

D. Both A and C are correct

See Answer in Part B → Q11

@Neet_pg_bot


Q12: A 30-year-old patient with HIV presents with vision loss and floaters. On examination, the patient has a CD4+ count of 40 cells/mm^3. Fundoscopy reveals areas of necrosis with the following appearance. What is the most likely diagnosis? (Q853683)

A. Toxoplasmosis

B. Herpes Simplex Virus (HSV) Retinitis

C. CMV retinitis

D. Herpes Zoster Ophthalmicus



See Answer in Part B → Q12

@Neet_pg_bot

Q13: Which of the following statements about the uveal tract is false? (Q853669)

A. The choroid connects anteriorly to the iris via the ciliary body.

B. The ciliary body contains both radial and circular smooth muscle fibers.

C. The posterior layer of the iris is continuous with the ciliary body

D. The constrictor pupillae muscle is controlled by sympathetic nerves.

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A 30-year-old man presents with symptoms of recurrent bilateral eye inflammation, including redness, pain, and vision loss. He also has a history of

frequent oral and genital ulcers and skin rashes. He is HLA-B51 positive. What would be the most appropriate approach for managing his uveitis? (Q853681)

A. Start with systemic corticosteroids and consider long-term immunosuppressive therapy

B. Immediate initiation of antiviral therapy

C. Use of topical antibiotics for ocular symptoms

D. Prescription of oral antihistamines for symptom relief

See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 30-year-old man with a recent history of urethritis and arthritis presents with eye symptoms. He is positive for HLA-B27. What is the most appropriate initial treatment? (Q853680)

A. Oral antibiotics

B. Topical corticosteroids

C. Systemic tetracycline

D. Immunosuppressive agents

See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 45-year-old male presents with unilateral eye pain, photophobia, and redness for the past five days. Slit-lamp examination reveals circumcorneal injection, anterior chamber cells and flare, and posterior synechiae. Despite initial treatment with topical corticosteroids and cycloplegics, his symptoms persist, and intraocular pressure (IOP) is found to be elevated. Which of the following is the most likely complication of the condition he is suffering from? (Q853679)

A. Retinal detachment

B. Secondary glaucoma

C. Vitreous hemorrhage

D. Optic neuritis

See Answer in Part B → Q16

@Neet_pg_bot

Q17: Which of the following is not a component of Posterior uveitis? (Q853678)

A. Retinitis

B. Choroiditis

C. Iridocyclitis

D. Retinal vasculitis

See Answer in Part B → Q17

@Neet_pg_bot

Q18: What is the 1st line of treatment in a 35 year old patient presenting with Intermediate uveitis? (Q853676)

A. Pars plana vitrectomy

B. Systemic steroids

C. Periocular injection of triamcinolone

D. Cryotherapy

See Answer in Part B → Q18

@Neet_pg_bot

Q19: A 25-year-old female presents with floaters and gradual deterioration of vision. On examination, there is anterior vitritis with white snowball-like exudates and mild

peripheral periphlebitis. No other relevant history or clinical findings. Which of the following is the most likely diagnosis? (Q853675)

A. CMV Retinitis

B. Tuberculosis

C. Intermediate uveitis

D. Lyme Uveitis

See Answer in Part B → Q19

@Neet_pg_bot

Q20: A 40-year-old patient presents with eye pain, redness, and a constricted pupil. During the examination, fine, grey deposits on the corneal endothelium are noted. Which of the following is the most appropriate next step in investigating and treating this condition? (Q853674)

A. Perform a chest X-ray and initiate treatment with topical cycloplegics.

B. Order a serum ACE level and start treatment with non-steroidal voclosporin.

C. Conduct an HLA typing and begin treatment with topical prednisolone.

D. Arrange for an OCT scan and administer systemic interferon-alpha.

See Answer in Part B → Q20

@Neet_pg_bot

Q21: A 35-year-old patient presents with a 1-week history of worsening eye pain and redness. On examination, a constricted pupil, a hazy anterior chamber, and deposits on the back of the cornea as shown below. Based on these findings, which of the following is the most likely diagnosis? (Q853673)

A. Herpes Simplex Keratitis

B. Conjunctivitis

C. Anterior Uveitis

D. Acute Angle Closure Glaucoma



See Answer in Part B → Q21

@Neet_pg_bot

Q22: A 50-year-old patient presents with anterior uveitis. After a detailed evaluation, the ophthalmologist considers various systemic associations and possible masquerades for the condition. Which of the following conditions is a known systemic association with anterior uveitis, but not a masquerade? (Q853672)

A. Intraocular Foreign Body

B. Behcet's Disease

C. Retinoblastoma

D. Pigment Dispersion Syndrome

See Answer in Part B → Q22

@Neet_pg_bot

Q23: Which of the following statements about the classification of uveitis based on the time of inflammatory activity is false? (Q853670)

A. Acute uveitis is defined as having a duration of less than 3 months and typically has a sudden onset.

B. Chronic uveitis is classified as lasting more than 6 months without interruption.

C. Recurrent uveitis involves repeated episodes with periods of inactivity between them.

D. Remission in uveitis is defined as no inflammatory activity for 3 months or more.

See Answer in Part B → Q23

@Neet_pg_bot

• [03_Uveitis] 006 Previous Year Questions

« Back to Folder Directory

[03_Uveitis] 006 Previous Year Questions — Quick Jump

Q1 (p. 93)	*Q2 (p. 93)*	*Q3 (p. 94)*	*Q4 (p. 94)*	*Q5 (p. 94)*	*Q6 (p. 95)*
Q7 (p. 96)	*Q8 (p. 96)*	*Q9 (p. 96)*	*Q10 (p. 97)*	*Q11 (p. 97)*	*Q12 (p. 97)*

Q1: A 35-year-old male presents with pain, redness, watering, and photophobia in the right eye for 2 days. Vision is mildly reduced. On examination, there is ciliary congestion, small irregular pupil, and presence of keratic precipitates on the corneal endothelium. The intraocular pressure is normal or slightly low.What is the most likely diagnosis? (Q866836)

A. Acute uveitis

B. Acute conjunctivitis

C. Acute congestive glaucoma

D. Central retinal artery occlusion

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A patient presents with sudden diminution of vision, difficulty looking towards the light, and circumcorneal congestion with a hypopyon. What is the most likely diagnosis? (Q844008)

A. Acute conjunctivitis

B. Acute angle closure glaucoma

C. Episcleritis

D. Acute anterior uveitis

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 32-year-old patient presents with blurred vision, photophobia, and mild ocular pain. Examination reveals aqueous flares and keratic precipitates in the anterior chamber. What is the likely diagnosis? (Q860512)

A. Intermediate uveitis

B. Posterior uveitis

C. Toxoplasma uveitis

D. Iridocyclitis

See Answer in Part B → Q3

@Neet_pg_bot

Q4: Iritis is least associated with all of the following diseases: (Q854715)

A. Systemic Lupus Erythematosus

B. Behcet's Disease

C. Psoriatic Arthropathy

D. Ulcerative colitis

See Answer in Part B → Q4

@Neet_pg_bot


Q5: A farmer presents with pain and discharge in his left eye. He says that he sustained an injury caused by a leaf a few days back. What is the finding seen in the image? (Q855559)

A. Hyphema

B. Aqueous flare

C. Arlt line

D. Hypopyon



@Neet_pg_bot

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Iritis is not seen in: Rheumatoid arthritis SLE Bechet’s disease Psoriatic arthritis Ulcerative colitis (Q856415)

A. 1 and 2

B. 1,2,3 and 4

C. 2,4 and 5

D. 1,3 and 4

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 26-year-old patient came with complaints of blurring of vision, photophobia, redness, pain in the left eye, and joint pain. Upon examination keratic precipitates and aqueous cells were observed. Which of the following major histocompatibility complex is be associated with this condition? (Q859309)

A. HLA B27

B. HLA B51

C. HLA B47

D. HLA B57

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 25-year-old male patient with a penetrating injury to the left eye was treated with corneoscleral suturing. Two months later, he returned with complaints of reduced near vision, pain, redness, and sensitivity to light in his right eye. Direct ophthalmoscopy revealed nodules in the fundus. Which part of the eye is affected? (Q859763)

A. Uvea

B. Cornea

C. Conjunctiva

D. Sclera

See Answer in Part B → Q8

@Neet_pg_bot

Q9: Which medication can be administered to a patient diagnosed with acute anterior uveitis, presenting with symptoms of photophobia and blurred vision? (Q859766)

A. Moxifloxacin

B. Atropine

C. Acetazolamide

D. Methotrexate

See Answer in Part B → Q9

@Neet_pg_bot

Q10: What is the specific sign that is pathognomonic for acute anterior uveitis among the options provided? (Q859771)

A. Aqueous flare

B. Keratic precipitates

C. Festooned pupil

D. Bussaca’s nodules

See Answer in Part B → Q10

@Neet_pg_bot

Q11: What will be observed in a patient diagnosed with sympathetic ophthalmitis, and a history of penetrating eye injury? (Q856036)

A. Acute anterior uveitis

B. Pars planitis

C. Panuveitis

D. Chronic anterior uveitis

See Answer in Part B → Q11

@Neet_pg_bot

Q12: On ophthalmologic examination, the presence of Koeppe's nodules is observed in a patient diagnosed with sarcoidosis. In which area of the eye are these nodules typically observed? (Q792994)

A. Cornea

B. Conjunctiva

C. Iris

D. Retina

See Answer in Part B → Q12

@Neet_pg_bot

• [04_Glaucoma] 007 Glaucoma-Pacg Treatment Congenital Glaucoma

« Back to Folder Directory

[04_Glaucoma] 007 Glaucoma-Pacg Treatment Congenital Glaucoma — Quick Jump

Q1 (p. 99)	*Q2 (p. 99)*	*Q3 (p. 100)*	*Q4 (p. 100)*	*Q5 (p. 101)*	*Q6 (p. 101)*
Q7 (p. 101)	*Q8 (p. 102)*	*Q9 (p. 102)*	*Q10 (p. 102)*	*Q11 (p. 103)*	*Q12 (p. 103)*
Q13 (p. 104)	*Q14 (p. 104)*	*Q15 (p. 105)*	*Q16 (p. 105)*	*Q17 (p. 105)*	*Q18 (p. 106)*
Q19 (p. 106)	*Q20 (p. 107)*	*Q21 (p. 107)*	*Q22 (p. 108)*	*Q23 (p. 108)*	*Q24 (p. 108)*
Q25 (p. 109)					

Q1: A 65-year-old female presents with acute onset pain, redness, and blurring of vision in the right eye. On examination, her intraocular pressure (IOP) is 48 mmHg, and there is a shallow anterior chamber with a hazy cornea. Slit-lamp examination reveals an intumescent, swollen lens with the iris pushed forward. Gonioscopy shows a closed angle. What is the most likely underlying mechanism for her glaucoma? (Q853570)

A. Obstruction of the trabecular meshwork by inflammatory cells

B. Forward displacement of the lens due to an abnormal zonular attachment

C. Disruption of aqueous humor flow due to posterior synechiae

D. Pupillary block caused by the swollen lens

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Which of the following conditions is not typically associated with secondary congenital glaucoma? (Q853593)

A. Aniridia

B. Marfan Syndrome

C. Turner Syndrome

D. Angelman syndrome

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 6-month-old infant with congenital glaucoma has corneal enlargement, clouding, and Haab's striae and the angle is difficult to visualize due to corneal clouding. Which surgical procedure is most appropriate for this infant? (Q853592)

A. Goniotomy

B. Trabeculotomy with Trabeculectomy

C. Trabeculectomy with antimetabolites

D. Glaucoma drainage devices

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 1-month-old child presents with persistent eye watering and light sensitivity. Examination shows enlarged corneas with Haab's striae, cloudiness, a deep anterior chamber, and thin, blue sclera. The child has had ocular enlargement since birth. The ophthalmologist suspects congenital glaucoma. Which type of glaucoma best fits this presentation? (Q853591)

A. Juvenile Glaucoma

B. Infantile Glaucoma

C. True Congenital Glaucoma

D. Secondary Congenital Glaucoma

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 50-year-old glaucoma patient is prescribed a prostaglandin analog as first-line therapy. Which of the following is the most likely mechanism of action and a common side effect of this medication? (Q853590)

A. Reduces aqueous humor production; Lid retraction

B. Increases trabecular outflow; Brow ache

C. Increases uveoscleral outflow; Iris pigmentation

D. Decreases aqueous humor secretion; Allergic blepharoconjunctivitis

See Answer in Part B → Q5

@Neet_pg_bot

Q6: All of the following are penetrating glaucoma surgeries except: (Q853589)

A. Trabeculectomy

B. Deep sclerectomy

C. Minimum incision glaucoma surgery

D. Glaucoma drainage device implantation

See Answer in Part B → Q6

@Neet_pg_bot

Q7: Which of the following laser procedures are used for Open angle glaucoma (OAG) and angle closure glaucoma (ACG) respectively? (Q853577)

A. Argon laser iridotomy for open-angle glaucoma and selective laser trabeculoplasty (SLT) for angle closure glaucoma

B. Selective laser trabeculoplasty for open-angle glaucoma and laser iridotomy for angle closure glaucoma

C. Laser iridoplasty for open-angle glaucoma and argon laser iridotomy for angle closure glaucoma

D. Laser iridotomy for open-angle glaucoma and laser iridoplasty for angle closure glaucoma

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 65-year-old male presents to the ophthalmology clinic for a routine eye examination. During the examination, his intraocular pressure (IOP) is measured at 28 mmHg in both eyes. The ophthalmologist diagnoses him with primary open-angle glaucoma. Which of the following medications is primarily designed to lower intraocular pressure by reducing the production of aqueous humor? (Q853576)

A. Travoprost

B. Pilocarpine

C. Acetazolamide

D. Unoprostone

See Answer in Part B → Q8

@Neet_pg_bot

Q9: Which ICE syndrome is characterized by the presence of pigmented nodular lesions on the iris? (Q853575)

A. Progressive iris atrophy

B. Chandler's syndrome

C. Cogan-Reese syndrome

D. Both A & C

See Answer in Part B → Q9

@Neet_pg_bot

Q10: Which of the following statements is false about traumatic glaucoma? (Q853574)

A. Traumatic glaucoma can develop from both perforating and blunt injuries.

B. Angle recession glaucoma typically presents with elevated IOP shortly after the trauma.

C. Lens-induced glaucoma can occur due to a ruptured, swollen, or dislocated lens following trauma.

D. Peripheral anterior synechiae from corneal injury can lead to angle-closure glaucoma.

See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 30-year-old male presents with unilateral eye discomfort, blurred vision, and seeing halos around lights. On examination, his intraocular pressure (IOP) is measured at 48 mm Hg. The anterior segment shows a few small, flat, nonpigmented keratic precipitates located inferiorly, mild redness, and an aqueous flare. The pupil is dilated, but there are no signs of a posterior synechiae. The patient reports recurrent episodes that resolve within days to weeks. Given these findings, what is the most likely diagnosis? (Q853573)

A. Secondary Angle-Closure Glaucoma with Pupillary Block

B. Secondary Angle-Closure Glaucoma without Pupillary Block

C. Open-Angle Glaucoma Due to Uveitis

D. Glaucomatocyclitic Crisis

See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 70-year-old patient with a history of glaucoma undergoes a trabeculectomy to control intraocular pressure. Postoperatively, the patient presents with severe eye pain, blurred vision, a significantly elevated intraocular pressure (IOP), and a shallow anterior chamber. Seidel's test is negative. What is the most appropriate initial treatment for this condition? (Q853572)

A. Oral corticosteroids

B. Pars plana Vitrectomy

C. Atropine drops and oral acetazolamide

D. Peripheral iridectomy

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 58-y/o diabetic patient presents with decreased vision, eye pain, and redness in the right eye. Examination reveals iris neovascularization, elevated intraocular pressure, and proliferative diabetic retinopathy. What is the most appropriate management strategy? (Q853571)

A. Topical beta-blockers and oral carbonic anhydrase inhibitors

B. Panretinal photocoagulation

C. Systemic corticosteroids

D. VEGF inhibitors

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A 55-year-old Asian woman presents with acute eye pain and blurred vision. Her ophthalmologist suspects primary angle closure glaucoma (PACG). Which of the following statements about the etiopathogenesis of PACG is incorrect? (Q853548)

A. Pupillary block is the most common mechanism in PACG

B. Phacomorphic mechanism involves an abnormally large lens pushing the iris forward

C. Plateau iris configuration is more common in younger patients with PACG

D. Caucasians have a higher prevalence of PACG compared to other ethnicities

See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 72-year-old male presents with diminution of vision in the right eye. On slit-lamp examination, dandruff-like material is noted on the anterior lens capsule and iris. Intraocular pressure is elevated, and the patient is diagnosed with pseudoexfoliative glaucoma. Which of the following is most likely to be observed in this patient's eye? (Q853569)

A. Posterior synechiae

B. Zonular instability and phacodonesis

C. Corneal guttata

D. Iris heterochromia

See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 25-year-old myopic male presents with intermittent blurring of vision and mild ocular pain after exercise. On examination, IOP is 28 mmHg, Slit lamp examination reveals Krukenberg spindle on corneal endothelium, trabecular pigmentation on gonioscopy. Which of the following is the first-line treatment? (Q853568)

A. Nd:YAG laser peripheral iridotomy

B. Selective laser trabeculoplasty

C. Topical prostaglandin analogue

D. Argon laser peripheral iridoplasty

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 45-year-old male presents with unilateral eye pain, redness, and moderately elevated intraocular pressure (IOP) of 28 mmHg. On slit-lamp examination, you observe fine, pigmented deposits on the corneal endothelium in a vertical spindle

pattern. The patient O/E has symmetric transillumination defects in the mid-peripheral iris. Based on these findings, which of the following is the most likely cause of this patient's secondary glaucoma? (Q853557)

A. Pseudoexfoliation syndrome

B. Pigment dispersion syndrome

C. Neovascular glaucoma

D. Lens-particle glaucoma

See Answer in Part B → Q17

@Neet_pg_bot

Q18: A 65-year-old male presents with a painful, blind right eye. On examination, the eye is stony hard with no light perception, 100% cupping of the optic disc, and a fixed, mid-dilated pupil. Which of the following is the correct initial management? (Q853556)

A. Topical timolol and brimonidine

B. Trabeculectomy

C. Cyclodestructive procedure

D. Evisceration

See Answer in Part B → Q18

@Neet_pg_bot

Q19: All of the following components of Vogt's triad are typically observed in a patient who has experienced and resolved an acute primary angle-closure glaucoma (APAC) attack except? (Q853555)

A. Glaucoma fleken

B. Iris atrophy

C. Pigment dispersion

D. Progressive retinal damage

See Answer in Part B → Q19

@Neet_pg_bot

Q20: A 55-year-old female presents for a routine eye examination. During gonioscopy, the examination reveals an occludable anterior chamber angle bilaterally. The patient reports no visual symptoms, and intraocular pressure (IOP) measurements are within normal limits. There is no evidence of glaucomatous damage on optic nerve examination. What is the most appropriate next step in management for this patient? (Q853554)

A. I.V Mannitol/Acetazolamide

B. Prophylactic peripheral iridotomy

C. Filtration surgery

D. Topical beta-blocker therapy

See Answer in Part B → Q20

@Neet_pg_bot

Q21: What aspect of the anterior chamber does the Van Herick grading system evaluate during slit lamp examination? (Q853553)

A. The curvature of the cornea

B. The width of the anterior chamber angle

C. The condition of the retina

D. The intraocular pressure

See Answer in Part B → Q21

@Neet_pg_bot

Q22: In the Shaffer classification system for grading the anterior chamber angle, which grade indicates that the angle is fully open with visible structures such as the scleral spur and trabecular meshwork? (Q853552)

A. Grade 0

B. Grade 1

C. Grade 2

D. Grade 3

See Answer in Part B → Q22

@Neet_pg_bot

Q23: Which of the following best describes the principle of gonioscopy? (Q853551)

A. It involves measuring intraocular pressure.

B. It is a technique for visualizing the anterior chamber angle of the eye using a special lens

C. It is a technique for visualizing the posterior chamber angle of the eye using a special lens

D. It evaluates the retina.

See Answer in Part B → Q23

@Neet_pg_bot

Q24: A 35-year-old woman presents with intermittent eye pain and blurred vision. Gonioscopy reveals narrow angles, and the angle remains narrow after laser peripheral iridotomy. Which of the following statements about plateau iris configuration is incorrect ? (Q853550)

A. It is characterized by an anteriorly positioned ciliary body

B. It is more commonly seen in older patients with primary angle closure glaucoma

C. Plateau iris configuration can persist after iridotomy

D. Ultrasound biomicroscopy (UBM) can be useful in diagnosing this condition

See Answer in Part B → Q24

@Neet_pg_bot

Q25: A 50-year-old patient presents with symptoms suggesting possible angle-closure glaucoma. You want to perform a comprehensive assessment of the anterior chamber angle. Which of the following statements is incorrect regarding anterior chamber angle assessment and anatomy? (Q853549)

A. Gonioscopy is the definitive diagnostic tool for primary angle closure.

B. Normal angle structures from anterior to posterior include: Schwalbe's line, non-pigmented trabecular meshwork, pigmented trabecular meshwork, scleral spur, and ciliary body band.

C. An angle is considered occludable if the pigmented trabecular meshwork is not visible in less than 180 degrees without indentation.

D. Indirect gonioscopes used to measure angle of AC are goldmann, posner-ritch, zeiss

See Answer in Part B → Q25

@Neet_pg_bot

• [04_Glaucoma] 008 Glaucoma - Poag Tonometry Perimetry

« Back to Folder Directory

[04_Glaucoma] 008 Glaucoma - Poag Tonometry Perimetry — Quick Jump

Q1 (p. 110)	*Q2 (p. 112)*	*Q3 (p. 112)*	*Q4 (p. 112)*	*Q5 (p. 113)*	*Q6 (p. 113)*
Q7 (p. 114)	*Q8 (p. 114)*	*Q9 (p. 115)*	*Q10 (p. 115)*	*Q11 (p. 115)*	*Q12 (p. 116)*
Q13 (p. 116)	*Q14 (p. 117)*	*Q15 (p. 117)*	*Q16 (p. 117)*	*Q17 (p. 118)*	*Q18 (p. 118)*
Q19 (p. 119)					

Q1: Match the following: 1) Perkins Tonometer a) {{caption_text}} 2) Tonopen b) {{caption_text}} 3) Pulsair tonometer c) {{caption_text}} 4) Rebound Tonometer d) {{caption_text}} (Q853862)

A. 1-a, 2-c, 3-d, 4-b

B. 1-c, 2-b, 3-d, 4-a

C. 1-d, 2-a, 3-b, 4-c

D. 1-c, 2-d, 3-a, 4-b




@Neet_pg_bot


@Neet_pg_bot



See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 62-year-old man with a history of Glaucoma undergoes visual field testing to assess his peripheral vision. The test involves moving a stimulus of constant size and intensity from a non-seeing area into a seeing area to determine the boundaries of his visual field. Which type of perimetry is being described? (Q853871)

A. Goldmann perimetry

B. Automated perimetry

C. Frequency doubling perimetry

D. Short-wavelength automated perimetry

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 55-year-old patient with primary open-angle glaucoma undergoes visual field testing. Which of the following visual field defects is considered the earliest clinically significant field defect occurring in glaucoma? (Q853870)

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 55-year-old patient undergoes a comprehensive eye examination including visual field testing. Which of the following statements is incorrect? (Q853869)

A. A scotoma is an area of decreased vision surrounded by normal vision

B. An isopter is a line connecting points of equal light sensitivity and size detection in the visual field

C. The blind spot is a type of absolute scotoma

D. The visual field extends maximally in the superior direction

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 58-year-old patient is being evaluated for suspected primary open-angle glaucoma. The ophthalmologist wants to perform a thorough examination of the optic nerve head to detect glaucomatous changes. Which of the following techniques is considered the examination of choice for evaluating fundus changes in this patient? (Q853868)

A. Direct ophthalmoscopy

B. Slit-lamp biomicroscopic examination with non-contact lens

C. Optical coherence tomography (OCT)

D. Indirect ophthalmoscopy

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 62-year-old woman with a family history of glaucoma presents with mild visual disturbances. Her intraocular pressure is measured to be 24 mmHg in both eyes. On dilated fundus examination, the following findings are noted: The cup-to-disc ratio of 0.5 Splinter hemorrhages at the optic disc margin Wedge-shaped retinal nerve fiber layer defects on red-free light examination What stage of glaucomatous optic disc changes is most likely present in this patient? (Q853867)

A. Glaucomatous optic atrophy

B. Advanced glaucomatous changes

C. Early glaucomatous changes

D. Physiological cupping

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 52-year-old male presents with progressive vision loss. Fundoscopic examination reveals a vertical elongation of the optic disc and diffuse thinning of the retinal nerve fiber layer (RNFL). Imaging confirms significant damage to the peripapillary region. Which of the following best describes the characteristic peripapillary change seen in glaucomatous optic neuropathy? (Q853866)

A. Increased retinal pigment epithelium (RPE) density in the peripapillary area

B. Presence of alpha zone peripapillary atrophy (PPA)

C. Thickening of peripapillary RNFL on optical coherence tomography (OCT)

D. Uniform retinal nerve fiber layer thickness without focal loss

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 58-year-old woman with a long history of untreated open-angle glaucoma presents with gradual peripheral vision loss. Fundoscopic examination reveals advanced optic disc cupping. Which of the following vascular changes is most characteristic of progressive glaucomatous optic neuropathy? (Q853865)

A. Inferior displacement of retinal vessels

B. Nasal displacement of retinal vessels

C. Arterial attenuation with increased vein diameter

D. Radial peripapillary capillary obliteration

See Answer in Part B → Q8

@Neet_pg_bot

Q9: A 58-year-old male with a family history of glaucoma presents for a comprehensive eye exam. His visual acuity is 20/20 in both eyes, and his intraocular pressure (IOP) is measured at 23 mmHg in the right eye and 21 mmHg in the left eye. Fundoscopic examination reveals the following optic disc findings in the right eye: Cup-to-disc ratio: 0.7 vertically Neuroretinal rim thinning: More pronounced in the superior and inferior quadrants Peripapillary atrophy present No optic disc hemorrhages noted All of the following statements are correct regarding the optic disc assessment in this patient except? (Q853864)

A. The optic disc appearance follows the ISNT rule, indicating a healthy optic nerve.

B. The cup-to-disc ratio indicates early glaucomatous damage.

C. The absence of optic disc hemorrhages does not rule out glaucoma.

D. Peripapillary atrophy is commonly associated with glaucoma and warrants further monitoring.

See Answer in Part B → Q9

@Neet_pg_bot

Q10: Which device is a non contact tonometer that is based on the principle of corneal hysteresis? (Q853863)

A. Schiotz tonometer

B. Goldmann applanation tonometer

C. Ocular Response Analyzer

D. Pneumatonometer

See Answer in Part B → Q10

@Neet_pg_bot

Q11: Which of the following statements about aqueous humor are incorrect? Aqueous humor is formed from the non-pigmented epithelium of ciliary processes Aqueous

constitutes 99% water The non-pigmented epithelium has a Na/Cl symporter responsible for aqueous humor secretion. The rate of formation of Aqueous humor is 2.3 microlitres/min (Q853853)

A. 1, 2, 4 are incorrect

B. Only 3 is incorrect

C. 1,3 is incorrect

D. Only 1 is incorrect

See Answer in Part B → Q11

@Neet_pg_bot

Q12: Which type of tonometer measures intraocular pressure (IOP) by assessing how much a plunger indents the eye? (Q853861)

A. Schiotz Tonometer

B. Goldmann Tonometer

C. Perkins Tonometer

D. Draeger Tonometer

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 58-year-old man presents for a routine eye examination. His intraocular pressure (IOP) is consistently measured at 23 mm Hg on multiple occasions. He reports no visual symptoms, and his fundus examination shows no abnormalities. Visual field testing reveals no defects. Based on these findings, which of the following diagnoses is most appropriate? (Q853860)

A. Normal tension glaucoma

B. Ocular hypertension

C. Acute angle-closure glaucoma

D. Secondary glaucoma

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A 65-year-old woman with a history of glaucoma reports worsening vision, headaches, and difficulty with reading. Her IOP shows significant fluctuations throughout the day, and fundus examination reveals optic disc cupping with visual field defects. Which of the following about IOP changes in POAG is false? (Q853859)

A. Diurnal variation includes a morning rise in IOP in 20% of cases and a biphasic rise in 55% of cases.

B. In early POAG, IOP is consistently above 21 mm Hg.

C. IOP is higher in winter due to seasonal variations.

D. Strenuous exercise temporarily lowers IOP.

See Answer in Part B → Q14

@Neet_pg_bot

Q15: Which gene mutation is linked to the development of primary open angle glaucoma? (Q853858)

A. MYOC gene

B. LOXL1 gene

C. PAX6 gene

D. RB1 gene

See Answer in Part B → Q15

@Neet_pg_bot

Q16: Which of the following is not a recognized risk factor for Primary Open-Angle Glaucoma? (Q853857)

A. Elevated intraocular pressure

B. Diastolic perfusion pressure of more than 55 mm Hg

C. Family history of glaucoma

D. High myopia

See Answer in Part B → Q16

@Neet_pg_bot

Q17: All the following statements about trabecular meshwork are true except; (Q853856)

A. It regulates the production of aqueous humor

B. Trabecular meshwork secrete Glycosaminoglycans

C. It facilitates the drainage of aqueous humor from the anterior chamber of the eye

D. Trabecular meshwork has 3 parts and among the three, juxtacanalicular has the maximum resistance

See Answer in Part B → Q17

@Neet_pg_bot

Q18: Which of the following best describes the role of uveoscleral outflow in intraocular pressure regulation? (Q853855)

A. It involves the drainage of aqueous humor through the trabecular meshwork and Schlemm's canal.

B. It accounts for the drainage of aqueous humor through the ciliary body and sclera, providing an alternative pathway to the conventional outflow route.

C. It is the process by which aqueous humor exits the eye through the vitreous body.

D. It describes the absorption of aqueous humor by the corneal endothelium.

See Answer in Part B → Q18

@Neet_pg_bot

Q19: All the following constituents are more in aqueous humor compared to that of plasma except (Q853854)

A. Lactate

B. Ascorbate

C. Pyruvate

D. Glucose

See Answer in Part B → Q19

@Neet_pg_bot

• [04_Glaucoma] 009 Previous Year Questions

« Back to Folder Directory

[04_Glaucoma] 009 Previous Year Questions — Quick Jump

Q1 (p. 120)	*Q2 (p. 120)*	*Q3 (p. 121)*	*Q4 (p. 121)*	*Q5 (p. 122)*	*Q6 (p. 122)*
Q7 (p. 123)	*Q8 (p. 123)*	*Q9 (p. 124)*	*Q10 (p. 124)*	*Q11 (p. 125)*	*Q12 (p. 125)*
Q13 (p. 126)	*Q14 (p. 126)*	*Q15 (p. 127)*	*Q16 (p. 128)*	*Q17 (p. 128)*	*Q18 (p. 129)*
Q19 (p. 129)	*Q20 (p. 130)*				

Q1: Which of the following instruments are considered the gold standard for measuring intraocular pressure? (Q857736)

A. Gonioscope

B. Goldmann applanation tonometer

C. Hertel’s exophthalmometer

D. Tonopen

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 60-year-old patient presents with sudden onset of severe ocular pain, headache, nausea, vomiting, and blurred vision. On examination, the cornea appears steamy, the pupil is mid -dilated and fixed, and the IOP is 60 mmHg. What is the most likely diagnosis? (Q879003)

A. Acute angle closure glaucoma

B. Secondary angle closure glaucoma due to uveitis

C. Phacomorphic glaucoma

D. Chronic simple glaucoma

See Answer in Part B → Q2

@Neet_pg_bot

Q3: Krukenberg spindle is seen in which of the following conditions? (Q878908)

A. Phacolytic glaucoma

B. Pigmentary glaucoma

C. Juvenile open -angle glaucoma

D. Primary open – angle glaucoma

See Answer in Part B → Q3

@Neet_pg_bot


Q4: What is the probable diagnosis for a 50-year-old woman who is experiencing sudden eye pain and redness? An image of her eye is provided for reference. (Q859359)

A. Acute retrobulbar neuritis

B. Acute angle-closure glaucoma

C. Acute anterior uveitis

D. Severe keratoconjunctivitis



See Answer in Part B → Q4

@Neet_pg_bot

Q5: The drug increasing uveoscleral flow is? (Q855019)

A. Timolol

B. Latanoprost

C. Pilocarpine

D. Carbonic anhydrase inhibitor

See Answer in Part B → Q5

@Neet_pg_bot

Q6: True regarding POAG: Abnormality of trabecular meshwork is seen on gonioscopy First degree relatives have more chances of developing steroid induced

glaucoma Dilatation of pupil is associated with exacerbation of IOP First degree relatives are at 1% increased risk of POAG Visual field defect can exist with normal C:D ratio (Q856418)

A. 1,2,4 and 5

B. 1,2 and 3

C. 2 and 5 only

D. 2,3 and 4

See Answer in Part B → Q6

@Neet_pg_bot

Q7: In a patient with primary open-angle glaucoma, which of the following pupillary abnormalities is most commonly seen? (Q856934)

A. Marcus Gunn pupil

B. Adie’s tonic pupil

C. Argyll-Robertson pupil

D. Hutchinson’s pupil

See Answer in Part B → Q7

@Neet_pg_bot

Q8: What is the likely diagnosis for a 2-month-old infant who was brought in by their mother due to complaints of light sensitivity, difficulty opening the eyes, and excessive tearing? (Q856125)

A. Ophthalmia neonatorum

B. Congenital glaucoma

C. Congenital Dacryocele

D. Congenital Cataract



See Answer in Part B → Q8

@Neet_pg_bot

Q9: Which medication should be avoided in a 30-year-old female presenting with redness, pain, increased intraocular pressure, keratic precipitates, and aqueous flare? (Q856696)

A. Beta-blockers

B. Mannitol

C. Carbonic anhydrase inhibitors

D. Prostaglandin analogues

See Answer in Part B → Q9

@Neet_pg_bot

Q10: Which of the following medications should not be administered to a patient with a known case of angle closure glaucoma who has also been diagnosed with major depressive disorder? (Q855384)

A. Amitriptyline

B. Sertraline

C. Mirtazapine

D. Fluvoxamine

See Answer in Part B → Q10

@Neet_pg_bot

Q11: A patient presents with a vision problem in the right eye. On examination, the intraocular pressure (IOP) of the right eye is 50 mmHg, while the IOP of the left eye is 16 mmHg. Slit-lamp examination of the right eye reveals flare cells in the anterior chamber and a cataractous lens. What is the diagnosis? (Q853114)

A. Phacolytic glaucoma

B. Malignant glaucoma

C. Fuch's heterochromic iridocyclitis

D. Phacomorphic glaucoma

See Answer in Part B → Q11

@Neet_pg_bot

Q12: What is the probable diagnosis for a neonate who presents with symptoms of photophobia and excessive tearing, along with a clinical examination showing a normal tear passage and a clear, but abnormally large cornea, and the presence of Haab's striae in the eye image below? (Q857662)

A. Buphthalmos

B. Keratoconus

C. Trachoma

D. Keratoglobus



See Answer in Part B → Q12

@Neet_pg_bot

Q13: After an acute episode of angle-closure glaucoma, which triad of symptoms is typically observed? (Q855241)

A. Vogt's triad

B. Virchow triad

C. Hutchinson's triad

D. Gaucher triad

See Answer in Part B → Q13

@Neet_pg_bot

Q14: In which of the following conditions is atropine not recommended for use? (Q855597)

A. Acute congestive glaucoma

B. Acute anterior uveitis

C. Malignant glaucoma

D. Corneal ulcer

See Answer in Part B → Q14

@Neet_pg_bot

Q15: What is the probable diagnosis for an infant brought in with symptoms of excessive tear production, sensitivity to light, and blepharospasm? (Q855495)

A. Congenital glaucoma

B. Congenital cataract

C. Congenital Dacryocystitis

D. Ophthalmia neonatorum



See Answer in Part B → Q15

@Neet_pg_bot

Q16: A patient presents with sudden painful decrease in vision and a fixed, vertically oval pupil. What is the likely diagnosis? (Q860174)

A. Macular degeneration

B. Optic neuritis

C. Acute congestive glaucoma

D. Acute anterior uveitis

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A patient presents with a vision problem in the right eye. On examination, the intraocular pressure (IOP) of the Right eye is 50 mmHg, while the IOP of the Left eye is 16 mmHg. Slit lamp examination of the right eye reveals flare cells in the anterior chamber and a cataractous lens. What is the diagnosis? (Q840060)

A. Phacolytic glaucoma

B. Malignant glaucoma

C. Fuch heterochromic iridocyclitis

D. Phacomorphic glaucoma

See Answer in Part B → Q17

@Neet_pg_bot

Q18: What is the gold standard method for measuring intraocular pressure in patients? (Q860572)

A. Hertel’s exophthalmometer

B. Gonioscope

C. Goldmann applanation tonometer

D. Tonopen

See Answer in Part B → Q18

@Neet_pg_bot

Q19: The first line of treatment for glaucoma in adults is? (Q860527)

A. PG analogues

B. Atropine

C. Acetazolamide

D. Pilocarpine

See Answer in Part B → Q19

@Neet_pg_bot

Q20: A patient complains of eye pain and difficulty seeing after watching a movie. What is the first step in management? (Q844037)

A. Mannitol with moxifloxacin

B. Mannitol with pilocarpine

C. Mannitol with atropine

D. Mannitol with carboxymethylcellulose

See Answer in Part B → Q20

@Neet_pg_bot

• [05_Retina] 010 Retinal Investigations And Vascular Disorders

« Back to Folder Directory

[05_Retina] 010 Retinal Investigations And Vascular Disorders — Quick Jump

Q1 (p. 131)	*Q2 (p. 131)*	*Q3 (p. 132)*	*Q4 (p. 132)*	*Q5 (p. 133)*	*Q6 (p. 133)*
Q7 (p. 134)	*Q8 (p. 135)*	*Q9 (p. 135)*	*Q10 (p. 136)*	*Q11 (p. 136)*	*Q12 (p. 137)*
Q13 (p. 137)	*Q14 (p. 138)*	*Q15 (p. 138)*	*Q16 (p. 138)*	*Q17 (p. 139)*	*Q18 (p. 139)*
Q19 (p. 140)	*Q20 (p. 140)*	*Q21 (p. 141)*	*Q22 (p. 142)*	*Q23 (p. 142)*	*Q24 (p. 143)*
Q25 (p. 143)	*Q26 (p. 144)*	*Q27 (p. 144)*	*Q28 (p. 144)*	*Q29 (p. 145)*	*Q30 (p. 145)*
Q31 (p. 146)	*Q32 (p. 146)*	*Q33 (p. 146)*	*Q34 (p. 147)*	*Q35 (p. 147)*	*Q36 (p. 148)*

Q1: A 23-year-old male presents to the ophthalmologist with pain, redness, photophobia or sensitivity to the light, and decreased vision from the last 3 days. Both the eyes of the patient are affected, and the patient is very irritable. Upon examination, infection of both the cornea and ciliary nerve is involved. The doctor diagnosed the patient with acute anterior uveitis. Which of the following HLA (human leukocyte antigen) is associated with uveitis? (Q882186)

A. HLA B27

B. HLA A29

C. HLA B51

D. HLA DR4

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 29-year-old man presents with sudden, severe visual loss following a recent road traffic accident involving chest trauma. Fundus examination reveals multiple white retinal patches near the optic disc, cotton wool spots, and minimal intraretinal hemorrhage. Based on these findings, what is the most likely diagnosis? (Q853784)

A. Diabetic Retinopathy

B. Retinal Vein Occlusion

C. Purtscher Retinopathy

D. Central Serous Retinopathy

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 30-week premature infant weighing 1.4 kg is diagnosed with stage 3 retinopathy of prematurity (ROP) in zone II without plus disease. What is the most appropriate next step in management? (Q853783)

A. Laser photocoagulation

B. Intravitreal anti-VEGF injection

C. Weekly observation and follow-up

D. Cryotherapy of retina

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 64-year-old male with long-standing hypertension comes for a routine eye checkup. On fundoscopy, the ophthalmologist observes the following changes as shown in the image below. What is the most likely stage of hypertensive retinopathy? (Q853696)

A. Grade I

B. Grade II

C. Grade III

D. Grade IV



See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 58-year-old male with a 20-year history of poorly controlled type 2 diabetes and an HbA1c of 10.5% came for an ophthalmic examination. Fundoscopy reveals microaneurysms, hard exudates, and retinal thickening. Which of the following is the most likely cause of future vision loss in this patient? (Q853695)

A. Vitreous hemorrhage

B. Retinal detachment

C. Macular edema

D. Neovascular glaucoma

See Answer in Part B → Q5

@Neet_pg_bot


Q6: A 45-year-old male presents with complaints of severe headache and blurred vision. His blood pressure is 220/120 mmHg. Fundoscopy image is given below. Which of the following fundus changes is most characteristic of this diagnosis? (Q853697)

A. Arteriovenous nicking

B. Copper-wiring

C. Papilledema

D. Hard exudates



@Neet_pg_bot

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 55-year-old patient with a history of uncontrolled hypertension presents for a routine eye examination. Fundoscopy reveals arteriolar narrowing and changes in the arteriovenous (A-V) ratio. What is the most likely A-V ratio finding in this patient with hypertensive retinopathy? (Q853698)

A. 2:3

B. 1:2

C. 1:3

D. 3:2

See Answer in Part B → Q7

@Neet_pg_bot

Q8: All of the following are considered risk factors for the development of central retinal vein occlusion (CRVO), except? (Q853699)

A. Age over 50 years

B. Open-angle glaucoma

C. Hyperthyroidism

D. Hypertension

See Answer in Part B → Q8

@Neet_pg_bot


Q9: A 65-year-old male presents with sudden, severe vision loss in his left eye. On examination, a relative afferent pupillary defect (RAPD) is present. Fundus examination reveals the following changes: Based on these findings, what is the most likely diagnosis? (Q853700)

A. Non-ischemic CRVO

B. Ischemic CRVO

C. Branch Retinal Vein Occlusion (BRVO)

D. Diabetic Retinopathy



See Answer in Part B → Q9

@Neet_pg_bot

Q10: Which of the following investigations is most useful in distinguishing ischemic CRVO from non-ischemic CRVO? (Q853701)

A. Fundus Autofluorescence (FAF)

B. Optical Coherence Tomography (OCT)

C. Goldmann Perimetry and Electroretinogram (ERG)

D. Slit-lamp Examination

See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 72-year-old man with a history of hypertension presents with sudden, painless vision loss in his right eye. Which of the following findings is most consistent with the diagnosis of central retinal artery occlusion? (Q853703)

A. Retinal hemorrhages and cotton wool spots

B. Optic disc edema and retinal hemorrhages

C. Dilated retinal veins with tortuosity

D. Pale retina with a cherry-red spot at the macula

See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 65-year-old male presents to the emergency department with sudden, painless vision loss in his right eye. On examination, the right eye shows a pale retina with a cherry-red spot at the macula. The patient has a history of hypertension and smoking. Which of the following is the most appropriate initial treatment to improve retinal blood flow and potentially restore vision? (Q853704)

A. Oral corticosteroids

B. Intravitreal anti-VEGF injection

C. Ocular massage and anterior chamber paracentesis

D. Systemic anticoagulation therapy

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A cherry-red spot is seen on the retina due to thickening and loss of transparency of the retina at the posterior pole. Which of the following diseases are least associated with this finding on fundoscopy? (Q853778)

A. Tay-Sachs disease

B. Farber disease

C. Central retinal artery occlusion

D. Central retinal vein occlusion

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A patient is being examined for loss of vision in the ophthalmology clinic. His peripheral visual field is lost, and only his central vision remains. Which of the following would not present with central (tunnel) vision? (Q853779)

A. Advanced glaucoma

B. Retinitis pigmentosa

C. CRAO with cilioretinal artery

D. Tobacco amblyopia

See Answer in Part B → Q14

@Neet_pg_bot

Q15: A premature infant, born at 26 weeks of gestation, is receiving oxygen therapy in the neonatal intensive care unit. Retinal examination at 4 weeks post-birth reveals signs of retinal vascular abnormalities. Which of the following mechanisms is most likely responsible for the pathogenesis of retinopathy of prematurity in this patient? (Q853780)

A. Increased oxygen supply leading to retinal vasodilation

B. Disruption of VEGF production

C. Excessive blood glucose levels

D. Retinal hypoxia leading to abnormal collagen formation

See Answer in Part B → Q15

@Neet_pg_bot

Q16: Which of the following is the most significant risk factor for Retinopathy of Prematurity (ROP)? (Q853781)

A. High birth weight

B. Prematurity and low birth weight

C. Prolonged exposure to blue light

D. Deficiency of Vitamin A

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 32-week premature infant presents with dilated and tortuous retinal vessels in the posterior pole, poor pupil dilation, and vitreous haze, indicating a progression of retinopathy of prematurity (ROP). What is the most likely diagnosis? (Q853782)

A. Rush disease

B. Plus disease

C. Cicatricial ROP

D. Stage 4 ROP

See Answer in Part B → Q17

@Neet_pg_bot

Q18: A 23-year-old male presents with sudden, painless vision loss in his left eye. Fundus examination reveals peripheral retinal vasculitis, vessel sheathing and recurrent vitreous hemorrhages. Fluorescein angiography reveals vessel leakage. What is the most appropriate initial treatment? (Q853785)

A. Intravitreal anti-VEGF therapy

B. Systemic corticosteroids

C. Laser photocoagulation

D. Vitrectomy

See Answer in Part B → Q18


Q19: A 3-year-old child is brought to the clinic with a white reflex (leukocoria) noticed in the right eye. An ocular B-scan ultrasound performed to evaluate the child shows a well-defined, echogenic mass within the globe, with associated calcifications. Which is the most likely diagnosis? (Q853671)

A. Retinal detachment

B. Vitreous hemorrhage

C. Retinoblastoma

D. Choroidal malignant melanoma



See Answer in Part B → Q19

Q20: What is the approximate distance of the fovea from the optic disc in a normal eye? (Q853652)

A. 1 disc diameter temporal to the optic disc

B. 2 disc diameters nasal to the optic disc

C. 2 disc diameters temporal to the optic disc

D. 1 disc diameter nasal to the optic disc

See Answer in Part B → Q20

@Neet_pg_bot


Q21: In which layer of the retina, as labeled in the histology image, are the cell bodies of bipolar cells located? (Q853653)

A. A

B. B

C. C

D. D



@Neet_pg_bot

See Answer in Part B → Q21

@Neet_pg_bot

Q22: A 65-year-old male with a history of hypertension and atherosclerosis presents with sudden, painless loss of vision in his left eye. On fundoscopy, you note a pale retina with a cherry-red spot at the macula. What is the most likely underlying cause of this condition? (Q853654)

A. Diabetic Retinopathy

B. Central Retinal Artery Occlusion

C. Central Retinal Vein Occlusion

D. Optic Neuritis

See Answer in Part B → Q22

@Neet_pg_bot

Q23: Which of the following arteries is primarily responsible for supplying the inner layers of the retina? (Q853655)

A. Central retinal artery

B. Short posterior ciliary arteries

C. Long posterior ciliary arteries

D. Ophthalmic artery

See Answer in Part B → Q23

@Neet_pg_bot

Q24: A 55-year-old male with a history of poorly controlled diabetes and hypertension presents with blurred vision. Fundus examination shows bright yellow, well-defined deposits arranged in a star-like pattern around the macula and cloud-like white patches near the optic disc. What is the most likely cause of the yellow deposits? (Q853656)

A. Soft exudates due to axoplasmic flow blockage

B. Hard exudates due to leakage of lipids and proteins

C. Retinal hemorrhages from ruptured microaneurysms

D. Cotton-wool spots from ischemic swelling of nerve fibers

See Answer in Part B → Q24

@Neet_pg_bot

Q25: The blood-retinal barrier separates the blood from the ocular cavity. Which of the following does not constitute the outer blood-retinal barrier? (Q853661)

A. Bruch’s membrane

B. Retinal pigmented epithelium

C. Zona occludens

D. Retinal vascular endothelial cells

See Answer in Part B → Q25

@Neet_pg_bot

Q26: A 60-year-old patient with a history of diabetes comes in for a routine check-up for diabetic retinopathy. The patient also has a cataract, causing a hazy view of the ocular media. Which method of study would be most appropriate for assessing the retina of this patient? (Q853662)

A. Direct ophthalmoscopy

B. Indirect ophthalmoscopy

C. Slit-lamp biomicroscopy

D. Amsler grid test

See Answer in Part B → Q26

@Neet_pg_bot

Q27: A 60-year-old patient with age-related macular degeneration (ARMD) undergoes Fundus Fluorescein Angiography (FFA), revealing pooling of dye under the retinal pigment epithelium (RPE) and leakage from newly formed vessels beneath the retina. What abnormality is most likely detected on FFA in this patient? (Q853663)

A. Hyperfluorescence due to dye pooling under detached RPE and neovascularization

B. Hypofluorescence due to blockage of background fluorescence by retinal hemorrhage

C. Hypofluorescence due to occlusion of retinal vasculature

D. Hyperfluorescence due to staining of the optic disc

See Answer in Part B → Q27

@Neet_pg_bot

Q28: A 35-year-old male presents with difficulty in vision, especially at night, and he reports bumping into objects frequently. Upon examination, he is found to have tunnel vision, and the fundus reveals no abnormalities in the posterior pole. Which region of the retina is most likely responsible for his symptoms? (Q853651)

A. Macula lutea

B. Fovea centralis

C. Optic disc

D. Peripheral retina

See Answer in Part B → Q28

@Neet_pg_bot

Q29: Which of the following electroretinogram (ERG) findings suggests gross disturbances in retinal circulation? (Q853685)

A. Extinguished ERG response

B. Subnormal ERG with a reduced b-wave

C. Negative ERG response

D. Normal ERG with a diminished c-wave

See Answer in Part B → Q29

@Neet_pg_bot

Q30: A 14-year-old patient presents with progressive visual impairment. Upon examination, there is evidence of vitelliform lesions in the macula. Electro-oculography (EOG) is performed to assess retinal function. What Arden ratio would be expected in a patient with Best disease on electro-oculography (EOG)? (Q853688)

A. Greater than 2.0

B. Less than 1.5

C. Between 1.5 and 2.0

D. Exactly 1.85

See Answer in Part B → Q30

@Neet_pg_bot

Q31: A patient with chronic diabetes mellitus complains of diminished vision. Her fundoscopy showed clinically significant macular edema. Which of the following would not be seen as a progressive change leading to diabetic retinopathy? (Q853689)

A. Macular ischemia

B. Neovascularization

C. Vitreous hemorrhage

D. Retinal vasoconstriction

See Answer in Part B → Q31

@Neet_pg_bot

Q32: Which of the following statements is false about laser treatment used for vascular diseases of the retina? (Q853690)

A. Focal laser therapy for diabetic macular edema targets leaking microaneurysms within 300 µm of the fovea.

B. Grid laser acts on diffuse retinal thickening, at least 500 µm from the fovea and optic disc.

C. Subthreshold lasers use very short pulses with longer intervals minimizing collateral damage.

D. Limitations include central vision risks, visual field defects, and the need for multiple sessions.

See Answer in Part B → Q32

@Neet_pg_bot

Q33: A 60-year-old diabetic patient presents with worsening vision. Fundoscopy reveals retinal edema, microaneurysms, and hard exudates. Which of the following mechanisms is not responsible for the development of diabetic retinopathy? (Q853691)

A. Protein kinase C activation

B. Reduced blood viscosity

C. Increased proangiogenic factors

D. Breakdown of the blood-retinal barrier

See Answer in Part B → Q33

@Neet_pg_bot

Q34: A 55-year-old patient with a long history of diabetes mellitus presents for a routine eye examination. The ophthalmologist noted several findings in the retinal examination. Which of the following features is least likely to be observed in early nonproliferative diabetic retinopathy (NPDR)? (Q853692)

A. Microaneurysm

B. Dot and blot hemorrhages

C. Hard exudates

D. Venous beading

See Answer in Part B → Q34

@Neet_pg_bot

Q35: A 60-year-old patient with chronic diabetes is diagnosed with diabetic retinopathy. What is the most specific finding for confirming proliferative diabetic retinopathy (PDR) in this patient? (Q853693)

A. Cotton-wool spots

B. Retinal hemorrhage

C. Neovascularization of disc

D. Intraretinal microvascular abnormalities

See Answer in Part B → Q35

@Neet_pg_bot

Q36: A 58-year-old male with Type 2 diabetes undergoes a fundus examination, revealing extensive retinal hemorrhages, cotton wool spots, venous beading, and moderate intraretinal microvascular abnormalities (IRMA) in one quadrant. What is the most appropriate follow-up interval for this patient? (Q853694)

A. 2 months

B. 4 months

C. 6 months

D. 12 months

See Answer in Part B → Q36

@Neet_pg_bot

• [05_Retina] 011 Macular And Fundus Disorders Retinal Detachment And Retinoblastoma

« Back to Folder Directory

[05_Retina] 011 Macular And Fundus Disorders Retinal Detachment And Retinoblastoma — Quick Jump

Q1 (p. 149)	*Q2 (p. 149)*	*Q3 (p. 150)*	*Q4 (p. 150)*	*Q5 (p. 150)*	*Q6 (p. 151)*
Q7 (p. 152)	*Q8 (p. 152)*	*Q9 (p. 152)*	*Q10 (p. 153)*	*Q11 (p. 153)*	*Q12 (p. 155)*
Q13 (p. 155)	*Q14 (p. 155)*	*Q15 (p. 156)*	*Q16 (p. 157)*	*Q17 (p. 158)*	*Q18 (p. 158)*
Q19 (p. 158)	*Q20 (p. 159)*				

Q1: Which of the following is a characteristic finding in the fundus examination of a patient with Stargardt disease? (Q853584)

A. Beaten bronze appearance

B. Cotton wool spots and Hard exudates

C. Dark/silent choroid

D. Honeycomb appearance

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 14-month-old child is diagnosed with retinoblastoma in the right eye. The tumor is confined to the retina, located posterior to equator, measures 2.5 mm in greatest dimension, and is located 3.5 mm from the fovea. There is no vitreous or subretinal seeding. According to the International Classification of Retinoblastoma (ICRB), how would this tumor be classified, and what will be the appropriate treatment? (Q853598)

A. Group A; Systemic chemotherapy with focal consolidation

B. Group B; Plaque radiotherapy

C. Group B; Enucleation

D. Group A; Laser photocoagulation

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 2-year-old child is brought to the ophthalmology clinic with a white pupillary reflex in the right eye, noticed by the parents in flash photography. On examination, the child has strabismus. Which of the following is NOT a condition to be considered in the differential diagnosis of this presentation? (Q853597)

A. Coats' disease

B. Persistent hyperplastic primary vitreous

C. Retinopathy of prematurity

D. Optic nerve glioma

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 2-year-old child presents with leukocoria in the right eye. Fundus examination reveals a white, elevated mass in the retina. Histopathological examination of the tumor shows rosettes with central lumen. Which of the following is the most specific histological feature for this condition? (Q853596)

A. Homer-Wright rosettes

B. Flexner-Wintersteiner rosettes

C. Pseudorosettes

D. Undifferentiated tumor cells

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 2-year-old child presents with the following findings as shown in the image. Genetic testing reveals a mutation in the RB1 gene. Which of the following statements best describes the genetic findings associated with this condition? (Q853595)

A. Retinoblastoma can be inherited in an autosomal dominant manner, with affected individuals having a 100% chance of passing on the mutation.

B. Retinoblastoma typically arises from a germline mutation in the RB1 gene, followed by a second somatic mutation.

C. Affected children with retinoblastoma often have mutations in both the RB1 and TP53 genes.

D. A single mutation in the RB1 gene is usually sufficient for the development of retinoblastoma in both familial and sporadic cases.



See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 55-year-old woman with a history of proliferative diabetic retinopathy presents with sudden vision changes in her right eye. She reports seeing floaters and flashes of light. On fundoscopy, there is evidence of fibrous tissue contraction and retinal elevation, but no retinal breaks are seen. What is the most appropriate treatment approach for this patient? (Q853594)

A. Laser photocoagulation

B. Cryotherapy with scleral buckling

C. Pars plana vitrectomy

D. Pneumatic retinopexy

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 60-year-old man presents with a sudden onset of flashes of light and a shadow resembling a curtain in the peripheral vision of his right eye. He denies any recent trauma. On examination, fundus evaluation reveals a retinal detachment with a retinal break in the upper temporal quadrant. What is the most appropriate next step in the management of this patient? (Q853588)

A. Intravitreal anti-VEGF injection

B. Pneumatic retinopexy with gas injection

C. Pars plana vitrectomy with silicone oil tamponade

D. Drainage of subretinal fluid with diathermy to seal the break

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 55-year-old man reports the sudden appearance of flashes of light, followed by a dark cloud or veil in the peripheral vision of his left eye. Examination reveals a greyish reflex in the affected quadrant of the retina. What type of retinal detachment is most likely in this patient? (Q853587)

A. Rhegmatogenous Retinal Detachment

B. Tractional Retinal Detachment

C. Exudative Retinal Detachment

D. Macular Hole

See Answer in Part B → Q8

@Neet_pg_bot

Q9: A 10-year-old child presents with a mild reduction in vision in both eyes. Fundus examination reveals a "sunny-side up egg" appearance in the macular region with yellowish subretinal material. The child's father has a similar history of visual impairment since childhood. What is the most likely diagnosis? (Q853586)

A. Stargardt Disease

B. Best Vitelliform Macular Dystrophy

C. Retinitis Pigmentosa

D. Leber Congenital Amaurosis

See Answer in Part B → Q9

@Neet_pg_bot

Q10: A 16-year-old patient presents with progressive vision loss. On examination, the ophthalmologist notes a central scotoma and yellowish-white flecks in the macular region of the retina. A diagnosis of Stargardt disease is made. All of the following are true statements about Stargardt disease, except: (Q853585)

A. It is also known as juvenile macular dystrophy

B. Fundus examination showing flame-shaped hemorrhages

C. Vitamin A supplementation should be avoided

D. Fundus autofluorescence

See Answer in Part B → Q10

@Neet_pg_bot


Q11: A 35-year-old male presents with sudden distrubunce of vision in his right eye for two weeks. He reports distortion of images. He has been intermittently using corticosteroids for allergies. His Visual acuity in the right eye is 6/18. FFA and OCT findings are given below. What is the most likely diagnosis? (Q853558)

A. Central serous chorioretinopathy

B. Age-related macular degeneration

C. Cystoid macular edema

D. Retinal vein occlusion





See Answer in Part B → Q11

@Neet_pg_bot

Q12: Which of the following is not related to retinitis pigmentosa? (Q853583)

A. Refsum's disease

B. Laurence-Moon-Bardet-Biedl syndrome

C. Kearns-Sayer syndrome

D. Conductive deafness

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 30-year-old woman presents with progressive night blindness and gradual loss of peripheral vision over the past few years. Her family history reveals that her father had similar symptoms. On examination, you observe characteristic retinal changes, including bone spicule pigmentation. To evaluate the functional status of her retina more accurately, which diagnostic test would you choose to confirm the suspicion of retinitis pigmentosa (RP)? (Q853567)

A. Electroretinography (ERG)

B. Optical Coherence Tomography (OCT)

C. Visual Field Testing

D. Electrooculogram (EOG)

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A 32-year-old female presents with difficulty seeing in low light and peripheral vision loss that has gradually worsened over several years. On fundoscopy, the

following findings are noted. Which of the following conditions is most likely responsible for these findings? (Q853566)

A. Retinitis Pigmentosa

B. Chronic Central Serous Chorioretinopathy

C. Age-related Macular Degeneration

D. Toxoplasmic Retinochoroiditis



See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 32-year-old woman presents with gradual central vision loss in both eyes. Fundus examination reveals concentric rings of retinal pigment epithelium (RPE) atrophy and hyperpigmentation centred on the fovea, creating a characteristic appearance. The image showing the appearance is given below: Which of the following is least likely to be associated with this condition? (Q853565)

A. Long-term hydroxychloroquine use

B. Cone dystrophy

C. Chloroquine maculopathy

D. Chronic central serous chorioretinopathy



See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 68-year-old woman presents with gradual vision loss in her right eye over the past 6 months. Her ophthalmologist suspects age-related macular degeneration (AMD) and orders further tests. What is the most sensitive modality to detect exudative AMD? (Q853563)

A. Optical Coherence Tomography (OCT)

B. Slit-lamp biomicroscopy using a +90 or +78 D lens

C. Indocyanine green angiography (ICGA)

D. Fluorescein angiography

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 72-year-old man presents with gradual vision loss in both eyes over the past year. His ophthalmologist diagnoses him with age-related macular degeneration (AMD). Which of the following statements about AMD is correct? (Q853562)

A. AMD is a hereditary degeneration exclusively affecting the macula

B. Wet AMD accounts for 90% of all AMD cases but only 10% of severe vision loss

C. Dry AMD is characterised by choroidal neovascularisation

D. AMD is the most common cause of permanent central visual loss after 6th decade

See Answer in Part B → Q17

@Neet_pg_bot

Q18: Which of the following statements about angioid streaks is incorrect? (Q853561)

A. They represent breaks in Bruch's membrane

B. They are commonly associated with sickle cell anemia

C. Choroidal neovascularization (CNV) frequently develops in these patients

D. Steroids are primarily used for the prevention of CNV

See Answer in Part B → Q18

@Neet_pg_bot

Q19: A 28-year-old woman with a history of high myopia since childhood presents for a routine eye examination. She reports occasional floaters and difficulty seeing at night. On fundus examination, the ophthalmologist notes a tilted optic disc, peripapillary atrophy, and a myopic crescent. Which of the following is NOT typically

associated with the condition? (Q853560)

A. Foster-Fuchs spots

B. Lattice degeneration

C. Posterior staphyloma

D. Optic disc edema

See Answer in Part B → Q19

@Neet_pg_bot


Q20: A 65-year-old presents with visual blurring in the left eye, 20/40 vision, and a history of cataract surgery three months ago. Fundus examination shows retinal thickening and a dull foveal reflex. OCT and FFA findings are given below. What is the most likely diagnosis? (Q853559)

A. Diabetic Retinopathy

B. Central Retinal Vein Occlusion

C. Cystoid Macular Edema

D. Central Serous Retinopathy





See Answer in Part B → Q20

@Neet_pg_bot

• [05_Retina] 012 Previous Year Questions

« Back to Folder Directory

[05_Retina] 012 Previous Year Questions — Quick Jump

Q1 (p. 161)	*Q2 (p. 161)*	*Q3 (p. 162)*	*Q4 (p. 162)*	*Q5 (p. 163)*	*Q6 (p. 164)*
Q7 (p. 164)	*Q8 (p. 165)*	*Q9 (p. 166)*	*Q10 (p. 166)*	*Q11 (p. 167)*	*Q12 (p. 168)*
Q13 (p. 168)	*Q14 (p. 168)*	*Q15 (p. 169)*	*Q16 (p. 169)*	*Q17 (p. 170)*	*Q18 (p. 170)*
Q19 (p. 171)	*Q20 (p. 171)*	*Q21 (p. 172)*	*Q22 (p. 172)*	*Q23 (p. 172)*	*Q24 (p. 173)*
Q25 (p. 173)	*Q26 (p. 174)*	*Q27 (p. 174)*	*Q28 (p. 175)*	*Q29 (p. 175)*	*Q30 (p. 176)*
Q31 (p. 176)	*Q32 (p. 177)*	*Q33 (p. 177)*	*Q34 (p. 178)*	*Q35 (p. 178)*	*Q36 (p. 179)*
Q37 (p. 179)	*Q38 (p. 179)*	*Q39 (p. 180)*	*Q40 (p. 180)*		

Q1: Which of the following is not a possible cause of the salt and pepper appearance of the fundus in a patient? (Q857752)

A. Rubella

B. Syphilis

C. Refsum’s disease

D. Tuberculosis

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A myopic patient presents with complaints of flashes and floaters. On examination, a deep anterior chamber is seen. What is the likely diagnosis? (Q866835)

A. Exudative retinal detachment

B. Rhegmatogenous retinal detachment

C. Tractional retinal detachment

D. Central serous retinopathy

See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which of the following is not seen in very severe NPDR? (Q854705)

A. Venous beading

B. Neovascularization of disc

C. Retinal hemorrhage

D. Microaneurysm

See Answer in Part B → Q3

@Neet_pg_bot


Q4: The image below displays the findings of a fundoscopy. What is the probable diagnosis? (Q858140)

A. CRAO

B. CRVO

C. Diabetic retinopathy

D. Cystoid macular edema



See Answer in Part B → Q4

@Neet_pg_bot

Q5: In a three-year-old boy, his parents have noticed a progressive decline in coordination over the last six months. During a physical examination, a growth resembling a cauliflower is observed in the left eye, along with leukocoria and the absence of a red reflex. Microscopic analysis of the sample reveals the presence of Flexner-Wintersteiner rosettes. Which gene is associated with this condition? (Q858150)

A. Rb gene

B. ATP 7B gene

C. Rhodopsin gene

D. FBN 1 gene

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Which of the listed conditions is associated with the presence of punched-out lesions on the retina? (Q855211)

A. Posterior staphyloma

B. Fundal coloboma

C. Congenital Rubella

D. Toxoplasmosis

See Answer in Part B → Q6

@Neet_pg_bot


Q7: What is the likely cause of sudden loss of vision in the left eye for the past two hours in a 58-year-old man who experienced it after his retirement party? The fundoscopic examination reveals a similar image shown below. (Q857205)

A. BRAO

B. CRAO

C. CRVO

D. BRVO



See Answer in Part B → Q7

@Neet_pg_bot

Q8: A male patient, aged 18, seeks medical attention due to an eye injury sustained while climbing a tree five days ago. The patient reports experiencing pain, sensitivity to light (photophobia), and redness in the affected eye for the past two days. Upon performing a slit-lamp examination, a lesion is observed as depicted below. Which of the following characteristics is associated with this particular condition? (Q857224)

A. Reverse hypopyon

B. Dendritic ulcer

C. Ring abscess

D. Satellite lesion



See Answer in Part B → Q8

@Neet_pg_bot

Q9: What is the likely diagnosis for a 7-year-old boy who showed a white pupillary reflex and underwent enucleation, with histopathological examination showing Flexner Wintersteiner rosettes? (Q857233)

A. Retinoblastoma

B. Medulloblastoma

C. Rhabdomyosarcoma

D. Astrocytoma

See Answer in Part B → Q9

@Neet_pg_bot


Q10: What is the condition characterized by painless vision loss? (Q857731)

A. Central retinal arterial occlusion

B. Branched retinal artery occlusion

C. Central retinal vein occlusion

D. Branched retinal vein occlusion



See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 55-year-old male patient presents to the clinic with a complaint of sudden visual blurring that lasted for a short duration . He reports that during the episode, he completely lost vision in his left eye for a few minutes and then regained his vision . The patient denies any pain, headache, or other associated symptoms during the episode. What is the most likely diagnosis ? (Q820648)

A. Amaurosis fugax

B. Central retinal artery occlusion

C. Central retinal vein occlusion

D. Hypertensive retinopathy

See Answer in Part B → Q11

@Neet_pg_bot

Q12: What is the primary mode of dissemination of retinoblastoma? (Q856706)

A. Optic nerve invasion

B. Hematogenous

C. Extraocular extension through sclera

D. Lymphatic spread

See Answer in Part B → Q12

@Neet_pg_bot

Q13: What is the most probable diagnosis for a 7-year-old patient who presented with a white pupillary reflex and underwent enucleation, with histopathological examination revealing the presence of Flexner Wintersteiner rosettes? (Q856178)

A. Retinoblastoma

B. Medulloblastoma

C. Rhabdomyosarcoma

D. Astrocytoma

See Answer in Part B → Q13

@Neet_pg_bot

Q14: Shifting fluid sign is seen in? (Q855022)

A. Exudative retinal detachment

B. Traction retinal detachment

C. Rhegmatogenous retinal detachment

D. Retinal dialysis

See Answer in Part B → Q14

@Neet_pg_bot

Q15: What is the most probable diagnosis for a 3-year-old boy who has displayed growing clumsiness over the last half a year, has leukocoria, and lacks red reflex in his left eye? After diagnosis, enucleation was performed, and the specimen displayed Flexner-Wintersteiner rosettes under microscopic examination? (Q859361)

A. Congenital glaucoma

B. Retinoblastoma

C. Toxocariasis

D. Neurofibroma

See Answer in Part B → Q15

@Neet_pg_bot


Q16: What is the condition in which the cherry red spot, as shown in the image below, is observed? {{caption_text}} (Q859761)

A. Central retinal artery occlusion (CRAO)

B. Central retinal venous occlusion (CRVO)

C. Branched retinal artery occlusion (BRAO)

D. Branched retinal vein occlusion (BRVO)



See Answer in Part B → Q16

@Neet_pg_bot

Q17: In which diseases are microaneurysms observed? (Q859764)

A. Retinal detachment

B. Diabetic retinopathy

C. Uveitis

D. Glaucoma

See Answer in Part B → Q17

@Neet_pg_bot

Q18: What should be the subsequent course of action in managing a diabetic patient who has a visual acuity of 6/9 in one eye and further investigations reveal preretinal

hemorrhages with neovascularization at the optic disc? (Q856002)

A. Focal laser photocoagulation

B. Pan-retinal photocoagulation

C. Grid laser photocoagulation

D. Scleral buckling

See Answer in Part B → Q18

@Neet_pg_bot

Q19: A patient has sudden loss of vision and there is no fundus glow. There is no history of trauma, what could be the likely cause? (Q860200)

A. Vitreous hemorrhage

B. Acute anterior uveitis

C. Optic neuritis

D. Acute congestive glaucoma

See Answer in Part B → Q19

@Neet_pg_bot

Q20: Which of the following criteria is not included in the definition of severe Non-Proliferative Diabetic Retinopathy (NPDR)? (Q860545)

A. Microaneurysms in all quadrants

B. Retinal hemorrhage in all quadrants

C. Looping in all quadrants

D. Bleeding in two quadrants

See Answer in Part B → Q20

@Neet_pg_bot

Q21: Which investigation is most useful for visualizing the individual layers of the retina? (Q865360)

A. Fundus Fluorescein Angiography

B. Optical Coherence Tomography

C. USG B-scan

D. Slit lamp examination

See Answer in Part B → Q21

@Neet_pg_bot

Q22: A woman with a history of proliferative diabetic retinopathy presents with gradual vision changes in her right eye. She reports seeing floaters and flashes of light. What type of retinal detachment is most likely in this patient? (Q865366)

A. Rhegmatogenous Retinal Detachment

B. Tractional Retinal Detachment

C. Exudative Retinal Detachment

D. Primary Retinal Detachment

See Answer in Part B → Q22

@Neet_pg_bot

Q23: Which of the following statements is incorrect about Retinitis Pigmentosa? (Q853115)

A. ERG shows decreased amplitude of a and b waves

B. Rod dystrophy > Cone dystrophy

C. Late presentation is loss of peripheral vision

D. Night vision is lost late and rare

See Answer in Part B → Q23

@Neet_pg_bot


Q24: Which of the following parts of the retina are correctly marked? (Q853129)

A. 1-Retinal pigment epithelium; 2-Outer nuclear layer; 3-Inner plexiform layer; 4-Ganglion layer

B. 1-Retinal pigment epithelium; 2-Inner nuclear layer; 3-Outer plexiform layer; 4-Ganglion layer

C. 1-Ganglion layer; 2-Inner nuclear layer; 3-Outer plexiform layer; 4-RPE

D. 1-Ganglion layer; 2-Outer plexiform layer; 3-Inner nuclear layer; 4-RPE



@Neet_pg_bot

See Answer in Part B → Q24

@Neet_pg_bot

Q25: Which of the following is a complication of diabetic retinopathy? (Q853132)

A. Posterior staphyloma

B. Retinoschisis

C. Tractional retinal detachment

D. Progressive myopia

See Answer in Part B → Q25

@Neet_pg_bot

Q26: Which of the following is the diagnostic investigation for Best disease? (Q853139)

A. EOG

B. ERG

C. Multifocal ERG

D. Visual evoked potential

See Answer in Part B → Q26

@Neet_pg_bot

Q27: A 3-month-old baby, born preterm at 33 weeks of gestation, is brought by parents with complaints of a white pupillary reflex for 2 weeks. CT revealed calcification of the globe in both eyes. What is the probable diagnosis? (Q853144)

A. ROP (Retinopathy of Prematurity)

B. Retinoblastoma

C. Glaucoma

D. Congenital Cataract

See Answer in Part B → Q27

@Neet_pg_bot


Q28: A patient presents with vision problems and has a history of cataract surgery. OCT finding is shown below. What is the syndrome most likely associated with these findings? (Q852496)

A. Posner-Schlossman Syndrome

B. Irvine-Gass Syndrome

C. Central Serous Retinopathy

D. Elschnig Pearls



See Answer in Part B → Q28

@Neet_pg_bot


Q29: A patient presents with a gradual loss of night vision and peripheral vision. What is the most likely diagnosis based on fundoscopy findings? (Q843989)

A. Retinitis pigmentosa

B. Retinal hemorrhage

C. Diabetic retinopathy

D. Hypertensive retinopathy



See Answer in Part B → Q29

@Neet_pg_bot

Q30: A child presents with leukocoria and an intraocular mass. Microscopic examination reveals rosettes with scanty cytoplasm, and the mass is positive for synaptophysin and chromogranin. What is the diagnosis? (Q844034)

A. Retinoblastoma

B. Medulloblastoma

C. Optic glioma

D. Choroidal melanoma

See Answer in Part B → Q30

@Neet_pg_bot

Q31: A 55-year-old hypertensive patient presented with complaints of blurred vision and headaches. On retinal examination, the following findings were observed: multiple flame-shaped hemorrhages, cotton wool spots, and disc edema. Based on the Keith-Wagner classification, which grade of hypertensive retinopathy is this patient most likely classified as? (Q820501)

A. GRADE 1

B. GRADE 2

C. GRADE 3

D. GRADE 4

See Answer in Part B → Q31

@Neet_pg_bot

Q32: A 72-year-old patient, with a history of cataract surgery and diabetes mellitus, presents with decreased vision one month after surgery. Examination reveals a honeycomb appearance or flower petal pattern of the retina. What is the likely diagnosis? (Q860554)

A. Cystoid macular edema

B. Central serous retinopathy

C. Retinal detachment

D. Diabetic retinopathy

See Answer in Part B → Q32

@Neet_pg_bot

Q33: Which of the following statements is incorrect about Retinitis pigmentosa? (Q840062)

A. ERG show decreased amplitude of a and b wave

B. Rod dystrophy > Cone dystrophy

C. Late presentation is loss of peripheral vision

D. Night vision is lost late and rare

See Answer in Part B → Q33

@Neet_pg_bot

Q34: A patient with long-standing rheumatoid arthritis who has been noncompliant with treatment presents with sudden, painless superior field loss in one eye for the past week. Fundus examination shows retinal whitening in the inferior retina with a corresponding superior visual field defect. What is the most likely cause of this condition? (Q840063)

A. Branched retinal artery occlusion

B. Central retinal artery occlusion

C. Branched retinal vein occlusion

D. Central retinal vein occlusion

See Answer in Part B → Q34

@Neet_pg_bot

Q35: A 22-year-old patient with high myopia and thick glasses presents with floaters in his eyes. What is the likely cause of this condition? (Q840065)

A. Subarachnoid hemorrhage

B. Retinal detachment

C. Vitreous hemorrhage

D. Branched retinal artery occlusion

See Answer in Part B → Q35

@Neet_pg_bot

Q36: What could be the cause of sudden painless loss of vision? (Q860179)

A. Chemical injuries to the eyeball

B. Acute congestive glaucoma

C. Acute iridocyclitis

D. Central serous retinopathy

See Answer in Part B → Q36

@Neet_pg_bot

Q37: Which of the following conditions is associated with the curtain falling sign? (Q860191)

A. Retinal detachment

B. Macular degeneration

C. Optic atrophy

D. Acute anterior uveitis

See Answer in Part B → Q37

@Neet_pg_bot

Q38: A 60-year-old male patient presents to the ophthalmology clinic with complaints of visual disturbances. He reports that he is unable to see on the left side from the left eye and the left side from his right eye. However, he notices that his central vision remains intact and unaffected. The patient denies any recent trauma or other significant medical history. What is the most likely visual deficit that the patient is experiencing? (Q820443)

A. Quadrantanopia

B. Heteronymous hemianopia

C. Homonymous hemianopia

D. Homonymous hemianopia with macular sparing

See Answer in Part B → Q38

@Neet_pg_bot

Q39: What is the most probable diagnosis for a 3-year-old boy who has been displaying increased clumsiness for the past 6 months, and upon examination, a cauliflower-like growth, leukocoria, and an absent red reflex in the left eye? Enucleation was performed, and microscopic examination of the specimen reveals the presence of Flexner-Wintersteiner rosettes. (Q793354)

A. Retinoblastoma

B. Neurofibroma

C. Retinopathy of prematurity

D. Toxcariasis

See Answer in Part B → Q39

@Neet_pg_bot


Q40: A 61-year-old university professor complains of vision loss in his left eye. He compares the sensation to a curtain being pulled over his affected eye. Additionally, he experienced flashing lights and floaters a week ago. The provided image shows a B-scan of his left eye. What could be the possible diagnosis for his condition? (Q793050)

A. Glaucoma

B. Posterior vitreous detachment

C. Retinal detachment

D. Corneal ulcer



See Answer in Part B → Q40

@Neet_pg_bot

• [06_Neuro Ophthalmology] 013 Neuro Ophthalmology

« Back to Folder Directory

[06_Neuro Ophthalmology] 013 Neuro Ophthalmology — Quick Jump

Q1 (p. 182)	*Q2 (p. 182)*	*Q3 (p. 183)*	*Q4 (p. 183)*	*Q5 (p. 183)*	*Q6 (p. 184)*
Q7 (p. 184)	*Q8 (p. 185)*	*Q9 (p. 185)*	*Q10 (p. 186)*	*Q11 (p. 186)*	*Q12 (p. 186)*
Q13 (p. 187)	*Q14 (p. 188)*	*Q15 (p. 188)*	*Q16 (p. 188)*	*Q17 (p. 189)*	*Q18 (p. 189)*
Q19 (p. 190)	*Q20 (p. 190)*	*Q21 (p. 191)*	*Q22 (p. 191)*	*Q23 (p. 192)*	*Q24 (p. 192)*
Q25 (p. 193)	*Q26 (p. 193)*	*Q27 (p. 194)*	*Q28 (p. 194)*	*Q29 (p. 195)*	*Q30 (p. 195)*
Q31 (p. 195)	*Q32 (p. 196)*				

Q1: A 50-year-old patient presents with worsening headaches and visual disturbances. Fundoscopic examination reveals optic disc swelling. Which of the following is least likely to be associated with this condition? (Q853766)

A. Increased cup-to-disc ratio

B. Enlargement of the blind spot

C. Blurring of the optic disc margins

D. Engorged and tortuous retinal veins

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Latent nystagmus is seen in which of the below mentioned condition? (Q853821)

A. Infantile esotropia

B. Arnold-Chiari malformation

C. Internuclear ophthalmoplegia

D. Cerebellopontine angle tumours

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 50-year-old woman presents with impaired adduction of the left eye on rightward gaze, and the abducting eye shows ataxic nystagmus. There is no noticeable ptosis, and the convergence remains intact. MRI shows a lesion in the brainstem affecting the medial longitudinal fasciculus. Which of the following is the most likely diagnosis? (Q853820)

A. Internuclear ophthalmoplegia

B. One-and-a-half syndrome

C. Oculomotor nerve palsy

D. Lateral rectus palsy

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 60-year-old woman presents with difficulty moving her eyes to the right. On examination, she is unable to gaze to the right, but has normal eye movements when gazing to the left, up, and down. There is no weakness in the limbs or facial muscles. Which of the following is the most likely location of the lesion? (Q853819)

A. Left abducens nucleus

B. Paramedian pontine reticular formation

C. Trochlear nerve

D. Right oculomotor nucleus

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 32-year-old woman presents with blurred vision in her right eye, photophobia and notices that her right pupil appears larger than the left. On examination, her right

pupil is mid-dilated and reacts very sluggishly to light and has a sluggish and tonic reaction to accommodation. The left pupil appears normal. Which of the following is the most likely diagnosis? (Q853818)

A. Horner’s syndrome

B. Adie’s tonic pupil

C. Third nerve palsy

D. Acute angle-closure glaucoma

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 30-year-old male has difficulty distinguishing between red and green colours. His occupation requires accurate colour perception. Which of the following is the most appropriate test to quantify and classify the severity of his red-green colour vision deficiency? (Q853817)

A. Ishihara colour plates

B. Farnsworth-Munsell 100-Hue Test

C. Nagel Anomaloscope

D. Lantern Test

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 30-year-old male presents with difficulty in colour vision. On further examination, he has a complete inability to perceive the red end of the spectrum. Which of the following is the most likely diagnosis? (Q853816)

A. Protanopia

B. Deuteranopia

C. Tritanopia

D. Total Color Blindness

See Answer in Part B → Q7

@Neet_pg_bot

Q8: On examination, a 45-year-old male was found to have repetitive, involuntary horizontal eye movements after looking at a moving striped pattern. He reports no dizziness, and his neurological examination is otherwise unremarkable. The phenomenon he is experiencing is most likely due to which of the following? (Q853815)

A. Gaze-evoked nystagmus

B. Vestibular Nystagmus

C. Optokinetic Nystagmus

D. Pendular Nystagmus

See Answer in Part B → Q8

@Neet_pg_bot

Q9: A 45-year-old male presents with difficulty moving his eyes upward and convergence-retraction nystagmus triggered by an upward gaze. His pupils react poorly to light but constrict normally when focusing on near objects. Which of the following syndromes is most likely? (Q853811)

A. Parinaud syndrome

B. Weber syndrome

C. Benedict syndrome

D. Millard-Gubler syndrome

See Answer in Part B → Q9

@Neet_pg_bot

Q10: A 50-year-old female patient with anisocoria had cocaine drops applied to both eyes. After 60 minutes, the anisocoria increased. To determine whether the lesion is preganglionic or postganglionic, what is the next step in the diagnostic process? (Q853810)

A. Reapply cocaine drops

B. 0.5% Apraclonidine

C. Adrenaline

D. 1% hydroxyamphetamine

See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 55-year-old male presents with a severe headache and neck stiffness that began a few days ago. His right pupil is smaller than the left, and he has drooping of the right eyelid and reduced sweating on the right side of his face. What is the most likely cause of these symptoms? (Q853809)

A. Cluster headaches

B. Pancoast tumour

C. Carotid artery dissection

D. Cavernous sinus tumor

See Answer in Part B → Q11

@Neet_pg_bot


Q12: A 45-year-old male reports a gradual decrease in vision in his left eye over the past two months, describing it as a "dimming" rather than a complete loss. He also has difficulty reading and seeing clearly. As per the image below, which test is most likely being conducted? (Q853808)

A. Swinging flashlight test

B. Perimetry

C. Tonometry

D. Visual acuity test



See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 48-year-old male with a history of untreated syphilis from several years ago now presents with memory loss, irritability, difficulty walking, and blurred vision. What type of pupil abnormality could be observed in this patient? (Q853807)

A. Marcus Gunn pupil

B. Amaurotic Pupil

C. Adie’s Pupil

D. Argyll Robertson Pupil

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A 45-year-old man presents with difficulty focusing on near objects. Which of the following is responsible for controlling ciliary muscle contraction and pupilary constriction during accomodation? (Q853806)

A. Lateral Geniculate Nucleus

B. Edinger-Westphal Nucleus

C. Superior Colliculus

D. Visual Cortex

See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 60-year-old patient exhibits the following pupillary responses: Light is shown in the left eye; the right pupil constricts, but the left pupil remains dilated. Light is shown in the right eye; only the right pupil constricts, while the left pupil remains dilated. Which of the following lesions is most likely responsible for this pattern of pupillary response? (Q853805)

A. Left optic nerve

B. Right Edinger-Westphal nucleus

C. Left oculomotor nerve

D. Right oculomotor nerve

See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 62-year-old patient with papilledema has been examined. On fundoscopic examination of the eye obscuration of major blood vessels is found on the disc & vessels leaving the disc, with a significant halo around the disc. According to the Frisen grading system, what is the grade of papilledema? (Q853767)

A. Grade 5

B. Grade 2

C. Grade 3

D. Grade 4

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 62-year-old patient presents with difficulty seeing objects in the right half of the visual field in both eyes. Based on this clinical presentation, which of the following is the most likely location of the lesion affecting the visual pathway? (Q853740)

A. Left optic nerve

B. Optic chiasma

C. Left optic tract

D. Right Lateral Geniculate Body

See Answer in Part B → Q17

@Neet_pg_bot

Q18: A 45-year-old patient presents with headaches and transient blurred vision. Fundoscopy reveals optic disc swelling with indistinct margins and engorged veins. What is the most accurate explanation for the pathogenesis of this condition? (Q853765)

A. Compression of the central retinal vein.

B. Increased CSF pressure causes axoplasmic stasis.

C. Inflammation of the optic nerve head.

D. Increased intraocular pressure.

See Answer in Part B → Q18

@Neet_pg_bot

Q19: A 35-year-old patient with multiple sclerosis (MS) notices that her vision becomes temporarily blurred whenever she exercises. Which of the following phenomena is most likely explained by this temporary worsening of neurological symptoms due to increased body temperature? (Q853754)

A. Riddoch's Phenomenon

B. Uhthoff's Phenomenon

C. Pulfrich's Phenomenon

D. Lhermitte sign

See Answer in Part B → Q19

@Neet_pg_bot

Q20: A 25-year-old man presents with a sudden onset of painless, bilateral vision loss that started in the right eye and later affected the left eye. Genetic testing reveals a mutation in the mitochondrial DNA. Which of the following is the most likely diagnosis? (Q853753)

A. Kjer dominant Optic Neuropathy

B. Leber Hereditary Optic Neuropathy (LHON)

C. Wolfram Syndrome

D. Behr syndrome

See Answer in Part B → Q20

@Neet_pg_bot

Q21: A 45-year-old man presents with sudden, progressive bilateral blurring of vision for two days, accompanied by headache, nausea, and vomiting. He has a history of alcohol dependence but denies taking any new medications. Examination shows bilateral central scotomas and hyperemic optic discs with mild retinal edema. His pupils are sluggishly reactive to light. Arterial blood gas analysis reveals a high anion gap metabolic acidosis.Which of the following toxins is most likely responsible for his condition? (Q853752)

A. Methanol

B. Cyanide

C. Arsenic

D. Ethambutol



See Answer in Part B → Q21

@Neet_pg_bot

Q22: A 72-year-old patient presents with sudden vision loss in one eye, severe headache, scalp tenderness, and jaw claudication. Fundoscopic examination reveals a pale, swollen optic disc with haemorrhages. Laboratory tests show elevated ESR and

CRP levels. What is the diagnosis and the appropriate management? (Q853751)

A. Arteritic AION; Lifestyle modifications

B. Non-arteritic AION; Lifestyle modifications

C. Arteritic AION; High dose corticosteroids

D. Non-arteritic AION; High dose corticosteroids

See Answer in Part B → Q22

@Neet_pg_bot

Q23: A 48-year-old woman presents with a history of decreased vision in her right eye. On examining the right eye, there is RAPD, pain on eye elevation, and the fundus is normal. What is the most appropriate management? (Q853750)

A. Conservative management

B. High-dose Corticosteroids

C. Oral antibiotics

D. Surgical intervention

See Answer in Part B → Q23

@Neet_pg_bot


Q24: A 40-year-old woman presents with a 3-week history of difficulty in vision in her left eye, along with persistent headaches and occasional eye pain. She also reports trouble with colour perception. Fundoscopic examination reveals the findings shown in the image. Which type of optic neuritis is most likely to be seen? (Q853749)

A. Papillitis

B. Retrobulbar neuritis

C. Neuroretinitis

D. Papilledema



See Answer in Part B → Q24

@Neet_pg_bot

Q25: A 34-year-old woman has a 6-month history of worsening vision and difficulty with detailed tasks, along with occasional eye pain in her left eye. Fundoscopy shows a chalky white optic disc with normal vessels in her left eye. What is the most common cause of these findings? (Q853748)

A. Papilledema

B. Multiple sclerosis

C. Retinitis pigmentosa

D. Glaucoma

See Answer in Part B → Q25

@Neet_pg_bot

Q26: A 45-year-old female reports a gradual decline in vision in her right eye over the past three months, accompanied by blurry vision and impaired colour perception in that eye. Upon examination, the ophthalmologist suspected optic nerve damage and advised further tests. Which of the following manifestations does not indicate optic nerve dysfunction? (Q853747)

A. Positive scotoma

B. Dyschromatopsia

C. Central scotoma

D. Decrease in contrast sensitivity

See Answer in Part B → Q26

@Neet_pg_bot

Q27: Ophthalmic artery is located Inferiolateral to which part of the optic nerve?? (Q853746)

A. Intraocular part

B. Intraorbital part

C. Intracanalicular part

D. Intracranial part

See Answer in Part B → Q27

@Neet_pg_bot

Q28: Which of the following arteries contributes to the formation of the Circle of Zinn-Haller? (Q853745)

A. Central retinal artery

B. Ophthalmic artery

C. Short posterior ciliary arteries

D. All of the above

See Answer in Part B → Q28

@Neet_pg_bot

Q29: A 62-year-old patient presents with a visual field defect following a stroke. Neuroimaging reveals a lesion in the occipital lobe related to the territory of the posterior cerebral artery affecting the calcarine cortex. Which of the following visual field characteristics is most likely associated with this type of lesion? (Q853744)

A. Congruous Homonymous Hemianopia with Macular Sparing

B. Congruous Homonymous Hemianopia without Macular Sparing

C. Incongruous homonymous hemianopia with Macular sparing

D. Incongruous homonymous hemianopia without Macular sparing

See Answer in Part B → Q29

@Neet_pg_bot

Q30: A 58-year-old patient presents with visual disturbances. Neuroimaging reveals a lesion in the temporal lobe affecting the inferior optic radiations. Which visual field defects are most likely associated with this finding? (Q853743)

A. Inferior homonymous quadrantanopia

B. Superior homonymous quadrantanopia

C. Central scotoma

D. Homonymous hemianopia

See Answer in Part B → Q30

@Neet_pg_bot

Q31: A 58-year-old patient presents with right-sided visual field loss in both eyes, resulting in a homonymous hemianopic defect. Neuroimaging and clinical examination suggest a lesion affecting the visual pathway. Based on the above findings, which anatomical location is most likely implicated in this patient's condition? (Q853742)

A. Right optic tract

B. Left optic tract

C. Right optic nerve

D. Optic chiasma

See Answer in Part B → Q31

@Neet_pg_bot

Q32: A 64-year-old patient presents with difficulty seeing objects. MRI findings reveal a pituitary adenoma causing compression at the optic chiasma. Which visual field defect is most likely associated with this condition? (Q853741)

See Answer in Part B → Q32

@Neet_pg_bot

• [06_Neuro Ophthalmology] 014 Previous Year Questions

« Back to Folder Directory

[06_Neuro Ophthalmology] 014 Previous Year Questions — Quick Jump

Q1 (p. 197)	*Q2 (p. 198)*	*Q3 (p. 198)*	*Q4 (p. 199)*	*Q5 (p. 199)*	*Q6 (p. 200)*
Q7 (p. 200)	*Q8 (p. 200)*	*Q9 (p. 201)*	*Q10 (p. 201)*	*Q11 (p. 202)*	*Q12 (p. 203)*
Q13 (p. 204)	*Q14 (p. 204)*	*Q15 (p. 205)*	*Q16 (p. 205)*	*Q17 (p. 206)*	*Q18 (p. 206)*
Q19 (p. 206)	*Q20 (p. 207)*	*Q21 (p. 207)*	*Q22 (p. 207)*	*Q23 (p. 208)*	*Q24 (p. 208)*
Q25 (p. 209)	*Q26 (p. 209)*	*Q27 (p. 210)*	*Q28 (p. 210)*	*Q29 (p. 211)*	*Q30 (p. 211)*
Q31 (p. 212)	*Q32 (p. 212)*	*Q33 (p. 213)*	*Q34 (p. 213)*	*Q35 (p. 213)*	*Q36 (p. 214)*
Q37 (p. 214)	*Q38 (p. 215)*	*Q39 (p. 215)*	*Q40 (p. 216)*		


Q1: What is the probable diagnosis for a middle-aged man with diabetes, who has reported a decline in vision and whose fluorescein angiography displays the depicted image? {{caption_text}} (Q857807)

A. Mild non proliferative diabetic retinopathy

B. Moderate non proliferative diabetic retinopathy

C. Severe non proliferative diabetic retinopathy

D. Proliferative diabetic retinopathy



See Answer in Part B → Q1

@Neet_pg_bot

Q2: A patient with pituitary adenoma compressing the optic chiasma now presents with loss of visual field. What visual field defect will be seen in the patient? (Q866833)

A. Bitemporal hemianopia

B. Homonymous anopia

C. Homonymous hemianopia

D. Superior quadrantanopia

See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which muscle serves as the yoke muscle for the right lateral rectus? (Q858157)

A. Right superior oblique

B. Left medial rectus

C. Left lateral rectus

D. Right medial rectus

See Answer in Part B → Q3

@Neet_pg_bot

Q4: Which of the following statements accurately describes optic neuritis? (Q855579)

A. Relative afferent pupillary defect is seen

B. There is no pain on ocular movement

C. Multiple myeloma is a common cause

D. Oral corticosteroids are the mainstay of treatment

See Answer in Part B → Q4

@Neet_pg_bot

Q5: Which structures are affected in a pregnant woman who exhibits fever, retroorbital pain, headache, pulsatile proptosis of the right eye, and tinnitus, while having normal blood pressure and fundus? 1. Abducens nerve 2. Oculomotor nerve 3. Internal carotid artery 4. Cavernous sinus (Q855223)

A. A, B, C

B. B, C ,D

C. A, C ,D

D. A, B, C, D

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Which cranial nerves does the oculocephalic reflex evaluate? A. III B. IV C. VI D. VIII E. VII (Q855237)

A. A and C

B. A and E

C. A and D

D. A, C and D

See Answer in Part B → Q6

@Neet_pg_bot

Q7: What would be the next investigative step for a 27-year-old female patient who presents with sudden vision loss accompanied by a relative afferent pupillary defect in her right eye, while her left eye appears normal on examination? 1. MRI brain & orbit 2. Myelin oligodendrocyte glycoprotein (MOG) and neuromyelitis optica (NMO) antibodies 3. Blood sugar 4. Thyroid profile (Q855244)

A. 1, 2 and 3

B. 1 and 2

C. 2 and 3

D. 3 and 4

See Answer in Part B → Q7

@Neet_pg_bot

Q8: What is the probable diagnosis for a man of 45 years who arrives at the emergency department with an abrupt loss of vision and experiencing painful movements of the eye? Upon examination, a relative afferent pupillary defect is observed in the affected eye. Further visual field testing demonstrates a central scotoma in the same eye. (Q857235)

A. Optic nerve glioma

B. Optic neuritis

C. Eale’s disease

D. Neuroretinitis

See Answer in Part B → Q8

@Neet_pg_bot

Q9: What is the probable diagnosis for a man of 45 years who arrives at the emergency department with an abrupt loss of vision and experiencing painful movements of the eye? Upon examination, a relative afferent pupillary defect is observed in the affected eye. Further visual field testing demonstrates a central scotoma in the same eye. (Q857665)

A. Optic nerve glioma

B. Optic neuritis

C. Eale’s disease

D. Neuroretinitis

See Answer in Part B → Q9

@Neet_pg_bot

Q10: In a patient diagnosed with a stroke, their visual field testing indicated macula sparing homonymous hemianopia. Where is the probable location of the lesion within the optic pathway? (Q857749)

A. Optic tract

B. Occipital lobe

C. Optic radiation

D. Optic chiasma

See Answer in Part B → Q10

@Neet_pg_bot


Q11: A patient presents to the OPD with the given visual field effect. What is the probable site of the lesion based on the image displayed? (Q858134)


A. 3

B. 1

C. 4

D. 6





See Answer in Part B → Q11

@Neet_pg_bot


Q12: Identify the site of the lesion. (Q856720)

A. Right occipital lobe

B. Left occipital lobe

C. Right optic tract

D. Optic chiasma



See Answer in Part B → Q12

@Neet_pg_bot

Q13: In a patient, a 33-year-old female presents with symptoms of gradual vision loss specifically affecting the right halves of both eyes. The question pertains to the location of the lesion in the optic pathway. (Q856124)

A. Left optic tract

B. Right visual cortex

C. Optic chiasma

D. Right optic nerve

See Answer in Part B → Q13

@Neet_pg_bot

Q14: What is the likely reason for a patient showing both unilateral and bilateral abducent palsy? (Q855026)

A. Retinoblastoma

B. Orbital pseudotumor

C. Orbital cellulitis

D. Cavernous sinus thrombosis

See Answer in Part B → Q14

@Neet_pg_bot

Q15: The lesion in which of the following nerve is responsible for right eye squint, as depicted below? (Q855031)

A. Oculomotor

B. Trochlear

C. Trigeminal

D. Abducent



See Answer in Part B → Q15

@Neet_pg_bot

Q16: The purpose of the photostress test is to distinguish between: (Q858507)

A. Macula and optic nerve disease

B. Cataract and glaucoma

C. Lens and cornea

D. Retina and vitreous pathologies

See Answer in Part B → Q16

@Neet_pg_bot

Q17: Where is the location of the lesion in Wernicke's hemianopic pupil? (Q858511)

A. Optic nerve

B. Optic tract

C. Optic radiation

D. Optic chiasma

See Answer in Part B → Q17

@Neet_pg_bot

Q18: What is the probable diagnosis for a patient who exhibits miosis, anhidrosis, mild ptosis, and a persistent small pupil even in low light conditions? (Q859316)

A. Adie’s tonic people

B. Horner syndrome

C. Marcus gunn pupil

D. Argyll Robertson pupil

See Answer in Part B → Q18

@Neet_pg_bot

Q19: What is the clinical presentation of a patient with a tumor located in the anterior pituitary gland causing compression on the optic chiasma? (Q859329)

A. Homonymous hemianopia

B. Inferior quadrantanopia hemianopia

C. Bitemporal hemianopia

D. Superior quadrantanopia hemianopia

See Answer in Part B → Q19

@Neet_pg_bot

Q20: Maximum visual field is seen in? (Q859816)

A. Temporal

B. Nasal

C. Inferior

D. Superior

See Answer in Part B → Q20

@Neet_pg_bot

Q21: Sudden painful loss of vision in one eye is due to the lesion at? (Q860534)

A. Optic nerve

B. Optic chiasma

C. Optic tract

D. Visual cortex

See Answer in Part B → Q21

@Neet_pg_bot

Q22: A 45-year-old patient presents with difficulty reacting to bright light. On examination, the pupillary light reflex is absent but the near reflex is preserved. Where is the lesion likely located? (Q866846)

A. Optic nerve

B. Oculomotor nerve

C. Edinger–Westphal nucleus

D. Pretectal area of the midbrain

See Answer in Part B → Q22

@Neet_pg_bot

Q23: A patient presents with a down and out right eye with ptosis since birth. What is the nerve palsy? (Q866852)

A. 4th nerve palsy

B. 3rd nerve palsy

C. 6th nerve palsy

D. 7th nerve palsy

See Answer in Part B → Q23

@Neet_pg_bot


Q24: Based on the eye movement findings shown in the image below, what is the most likely diagnosis? (Q865341)

A. Horizontal gaze palsy

B. Internuclear ophthalmoplegia (INO)

C. 3rd nerve palsy

D. Myasthenia gravis



@Neet_pg_bot

See Answer in Part B → Q24

@Neet_pg_bot

Q25: Which of the following nerves gives sensory supply to the orbit? (Q865355)

A. Vagus nerve

B. Trigeminal nerve

C. Hypoglossal nerve

D. Oculomotor nerve

See Answer in Part B → Q25

@Neet_pg_bot

Q26: In a patient with dense central (macular) corneal or lens opacity, which of the following statements about the visibility of intraocular structures is correct?

(Q853130)

A. Both the iris and pupil are clearly visible

B. Only the pupillary margin is visible

C. Iris details are visible but not the pupil

D. Neither the iris nor the pupillary margin is visible

See Answer in Part B → Q26

@Neet_pg_bot

Q27: A 5-month-old baby is brought by the mother with complaints of the left upper eyelid moving up and down during breastfeeding or thumb sucking, which disappears when the baby is not being fed. What is the most likely diagnosis? (Q852492)

A. Marcus Gunn Jaw-Winking Syndrome

B. Lagophthalmos

C. Myasthenia Gravis

D. 3rd CN palsy

See Answer in Part B → Q27

@Neet_pg_bot

Q28: A patient presents with breathlessness and is diagnosed with a Pancoast tumor. Which of the following findings is not typically associated with this condition? (Q852488)

A. Anhidrosis

B. Miosis

C. Ptosis

D. Exophthalmos

See Answer in Part B → Q28

@Neet_pg_bot


Q29: Identify the visual field defect: (Q843978)

A. Altitudinal defect

B. Roennes nasal step

C. Blind spot enlargement

D. Arcuate scotoma



See Answer in Part B → Q29

@Neet_pg_bot


Q30: Identify the defect shown in the image given below: (Q844012)

A. Binasal hemianopia

B. Bitemporal hemianopia

C. Macula- sparing homonymous hemianopia

D. Altitudinal defect



See Answer in Part B → Q30

@Neet_pg_bot

Q31: What are the yoke muscles for Levodepression? (Q858121)

A. LSO and RIR

B. RSO and LIR

C. RIO and LIR

D. LIO and RIR

See Answer in Part B → Q31

@Neet_pg_bot

Q32: Optic nerve glioma is seen in? (Q860538)

A. NF-1

B. Tuberous sclerosis

C. NF-2

D. Schwannoma

See Answer in Part B → Q32

@Neet_pg_bot

Q33: Which tuberculosis drug causes impairment of color vision and optic neuritis? (Q860567)

A. Rifampicin

B. Ethambutol

C. Pyrazinamide

D. Streptomycin

See Answer in Part B → Q33

@Neet_pg_bot

Q34: Which nerve defect causes lagophthalmos? (Q860170)

A. 7th nerve

B. 6th nerve

C. 4th nerve

D. 5th nerve

See Answer in Part B → Q34

@Neet_pg_bot

Q35: A patient on antitubercular therapy (ATT) started experiencing visual problems. Which ATT drug is most likely responsible for this adverse effect? (Q860207)

A. Rifampicin

B. Streptomycin

C. Ethambutol

D. Pyrazinamide

See Answer in Part B → Q35

@Neet_pg_bot

Q36: In the provided image, which nerve is subjected to blockage? (Q857420)

A. Anterior ethmoidal nerve

B. Nasociliary nerve

C. Supratrochlear nerve

D. Long ciliary nerve



See Answer in Part B → Q36

@Neet_pg_bot

Q37: In a female patient, there is a complaint of vision loss, specifically affecting the right halves of both eyes. Where is the lesion located? (Q855459)

A. Optic nerve

B. Left optic tract

C. Optic chiasma

D. Right optic tract

See Answer in Part B → Q37

@Neet_pg_bot

Q38: A 40-year-old male patient presents with altered sensation in the extremities, difficulty with coordination, and speech problems. On examination, it is noted that the pupillary light reaction is absent, but the accommodation reaction is present. Which condition is most likely responsible for this presentation? (Q858115)

A. Neurosyphilis

B. Horner's syndrome

C. Papilledema

D. Optic Neuritis

See Answer in Part B → Q38

@Neet_pg_bot

Q39: A 40-year-old male patient presents with altered sensation in the extremities, difficulty with coordination, and speech problems. On examination, it is noted that the pupillary light reaction is absent, but the accommodation reaction is present. Which condition is most likely responsible for this presentation? (Q820662)

A. Neurosyphilis

B. Horner’s syndrome

C. Papilledema

D. Optic Neuritis

See Answer in Part B → Q39

@Neet_pg_bot

Q40: A 45-year-old male arrived at the emergency ward with sudden vision loss and experiencing painful eye movements . Upon examination, the affected eye shows a relative afferent pupillary defect, a normal optic disc , and a central scotoma . What is the likely diagnosis? (Q793017)

A. Optic nerve glioma

B. Optic neuritis

C. Eale’s disease

D. Optic atrophy

See Answer in Part B → Q40

@Neet_pg_bot

• [07_Squint _ Strabismus] 015 Squint

« Back to Folder Directory

[07_Squint _ Strabismus] 015 Squint — Quick Jump

Q1 (p. 217)	*Q2 (p. 217)*	*Q3 (p. 218)*	*Q4 (p. 218)*	*Q5 (p. 218)*	*Q6 (p. 219)*
Q7 (p. 219)	*Q8 (p. 220)*	*Q9 (p. 220)*	*Q10 (p. 220)*	*Q11 (p. 221)*	*Q12 (p. 222)*
Q13 (p. 222)	*Q14 (p. 222)*	*Q15 (p. 223)*	*Q16 (p. 223)*	*Q17 (p. 224)*	*Q18 (p. 224)*
Q19 (p. 225)	*Q20 (p. 226)*	*Q21 (p. 226)*	*Q22 (p. 227)*	*Q23 (p. 228)*	*Q24 (p. 229)*
Q25 (p. 229)	*Q26 (p. 230)*	*Q27 (p. 230)*			

Q1: A 3-year-old girl presents with esotropia, which increases on focusing near objects. This issue has progressively worsened over the past few months, and she has difficulty with close visual tasks. A refractive error of +6.50 D has been identified. What is the most likely diagnosis? (Q853529)

A. Accommodative refractive esotropia

B. Accommodative non-refractive esotropia

C. Accommodative mixed esotropia

D. Infantile esotropia

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 7-year old child is brought to the clinic by her parents due to concerns about her right eyes appearing misaligned. On examination, her squint remains constant in all directions of gaze. She does not report any diplopia, and there are no limitations in eye movement. What is the most likely diagnosis? (Q853543)

A. Concomitant squint

B. Incomitant squint

C. Paralytic squint

D. Pseudo-strabismus

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 65 year old woman presented to the OPD with her head tilted towards the right side. On examination, she was having left hypertropia which increases on looking right or medially. The muscle which is most likely paralyzed is: (Q853542)

A. Right superior oblique

B. Left superior oblique

C. Right inferior oblique

D. Left inferior oblique

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 40-year-old patient presents for binocular vision assessment. The ophthalmologist performs the Worth's Four Dot Test. The patient wears red-green glasses and reports seeing three green dots. What does this finding most likely indicate? (Q853541)

A. Diplopia

B. Suppression of right eye

C. Suppression of left eye

D. Normal binocular vision

See Answer in Part B → Q4

@Neet_pg_bot

Q5: During a Maddox rod test, the patient is asked to view a distant light source. With the Maddox rod placed in front of the right eye, the patient reports seeing a vertical red line to the right of the white light. What type of ocular misalignment does this result suggest? (Q853540)

A. Esophoria

B. Exophoria

C. Right hyperphoria

D. Left hyperphoria

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 45-year-old male presents with eyelid retraction on downgaze. He has a history of ocular nerve injury. Which of the following best describes this condition? (Q853539)

A. Von Graefe’s Sign

B. Hutchinson’s Pupil

C. Pseudo-Von Graefe Sign

D. Marcus Gunn Jaw-Winking Phenomenon

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 7-year-old girl has limited abduction of the left eye, with globe retraction and narrowing of the palpebral fissure on adduction and widening of the fissure on attempted abduction. What is the most likely diagnosis? (Q853538)

A. Duane Retraction Syndrome Type I

B. Duane Retraction Syndrome Type II

C. Duane Retraction Syndrome Type III

D. Sixth Cranial Nerve Palsy

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 60-year-old patient presents with the following symptoms: right-sided facial weakness, right-sided lateral gaze impairment, and weakness on the left side of the body. Imaging reveals a ventral pontine lesion. Which of the following syndromes best describes this patient's condition? (Q853537)

A. Gradenigo Syndrome

B. Millard-Gubler Syndrome

C. Moebius Syndrome

D. Foville Syndrome

See Answer in Part B → Q8

@Neet_pg_bot

Q9: A 45-year-old man has double vision and an inward deviation of his left eye. On examination it is found to be associated with 6th Cranial Nerve Palsy. What finding is typical for sixth cranial nerve palsy? (Q853536)

A. Vertical diplopia

B. Pupils are affected

C. Head turn towards the same side, avoiding diplopia

D. Crossed Diplopia

See Answer in Part B → Q9

@Neet_pg_bot

Q10: A 45-year-old patient reports diplopia and is found to have left trochlear nerve palsy. Which of the following findings is not typically associated with it? (Q853534)

A. Diplopia relieves while going downstairs

B. No diplopia with head tilt away from the left side

C. Hypertropia of the affected eye

D. Increase in hypertropia on opposite gaze

See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 65-year-old patient presents with the following findings. Which of the following is least likely to be associated with this condition? (Q853533)

A. Weber's Syndrome

B. Benedict's Syndrome

C. Nothmagel's Syndrome

D. Brown's Syndrome



See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 60-year-old man presents with sudden onset double vision and difficulty aligning his eyes, which began three weeks ago. He has a history of hypertension and recently experienced a mild stroke. Examination reveals that his eye is positioned downward and outward. Which of the following statements about this condition is incorrect? (Q853532)

A. Ptosis is present

B. Mydriasis is present

C. Pupil is fixed

D. Accommodation increases

See Answer in Part B → Q12

@Neet_pg_bot

Q13: How much is the difference in deviation between upward gaze and downward gaze in A-V pattern strabismus? (Q853531)

A. A-pattern ≥ 10 PD, V-pattern ≥ 15 PD

B. A-pattern ≥ 15 PD, V-pattern ≥ 10 PD

C. A-pattern ≤ 10 PD, V-pattern ≤ 15 PD

D. A-pattern ≤ 15 PD, V-pattern ≤ 10 PD

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A 45-year-old male complains of double vision, describing seeing two overlapping images when looking in a specific direction. He reports that this problem started suddenly about two weeks ago and is most noticeable when looking towards the right side. There is no history of trauma. On examination, he has limited movement of the right eye towards the right. What is the most likely cause of this condition? (Q853530)

A. Blowout fracture

B. Thick pterygium

C. Subluxation of lens

D. Paralytic strabismus

See Answer in Part B → Q14

@Neet_pg_bot

Q15: Which of the following is known for its specific pattern of muscle insertions around the sclera, essential for the alignment and movement of the extraocular muscles? (Q853504)

A. Annulus of Zinn

B. Spiral of Tillaux

C. Scleral spur

D. Tenon’s capsule

See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 10-year-old child is having trouble catching a ball, climbing stairs, and performing fine motor tasks. She has a tendency to avoid activities requiring precise hand skills. No developmental delay was detected. Which of the following tests would be appropriate for diagnosing her condition? (Q853528)

A. HESS chart

B. Cover-uncover test

C. Titmus fly test

D. Maddox rod test

See Answer in Part B → Q16


Q17: The parents of an 8-month-old girl have observed her eyes misaligning, especially when she focuses on objects up close, with one eye occasionally turning inward. They consulted an ophthalmologist, who conducted the test shown below. What is the name of this test? (Q853527)

A. Hirschberg test

B. Bruckner test

C. Prism cover test

D. Krimsky test



See Answer in Part B → Q17


Q18: A 10-year-old boy's parents noticed his left eye turning inward, especially when tired or focusing on near objects. An ophthalmologist diagnosed strabismus and performed the test shown below. Which statements about this test are correct? (Q853526)

A. It indicates the angle of squint

B. In an aligned eye, the reflex is centred

C. Reflex at limbus indicates 45° squint

D. All of the above



See Answer in Part B → Q18

@Neet_pg_bot

Q19: A 32-year-old patient presents with eye strain and headaches. The ophthalmologist plans to perform a series of tests to evaluate the patient's strabismus. All of the following are objective tests used to assess strabismus except? (Q853513)

A. Maddox Rod test

B. Prism Bar Cover test

C. Cover Uncover test

D. Hirschberg test

See Answer in Part B → Q19

@Neet_pg_bot

Q20: A 35-year-old patient presents with increased diplopia on upward gaze and restricted movement of the left eye. The patient also has a history of recent orbital trauma. To determine whether the limitation is due to a mechanical restriction or a nerve palsy, which of the following tests is most appropriate? (Q853512)

A. Hess screen test

B. Forced duction test

C. Cover test

D. Worth's 4-dot test

See Answer in Part B → Q20

@Neet_pg_bot


Q21: A 10-year-old patient presents to the clinic with complaints of occasional double vision and eye strain during schoolwork. On examination, you suspect she may have a squint and perform the following test. What is this test known as? (Q853511)

A. Maddox Rod Test

B. Synoptophore

C. Cover-Uncover Test

D. Maddox Wing Test



See Answer in Part B → Q21

@Neet_pg_bot

Q22: Which of the following options correctly identifies the type of strabismus depicted in each image? 1. 2. 3. (Q853510)

A. 1- Esotropia; 2-Exotropia; 3-Hypotropia

B. 1- Esotropia; 2-Hypertropia; 3- Exotropia

C. 1- Hypertropia; 2-Exotropia; 3-Hypotropia

D. 1- Hypertropia; 2-Esotropia; 3-Exotropia



See Answer in Part B → Q22

@Neet_pg_bot

Q23: Which mechanism is responsible for maintaining coordinated movement of both eyes and achieving binocular vision? (Q853509)

A. Conjugate and Dysconjugate movements

B. Synergistic and Antagonistic muscles

C. Yoke muscles

D. All of the above

See Answer in Part B → Q23

@Neet_pg_bot

Q24: A 32-year-old patient presents with diplopia and difficulty looking upwards and to the right, with limited elevation of the right eye. You suspect a lesion affecting a muscle responsible for vertical movement. Which axis describes the rotational movement, and which muscle is likely affected? (Q853508)

A. Fick's x-axis; Right Inferior Rectus

B. Fick's y-axis; Right Medial Rectus

C. Fick's z-axis; Right Inferior Oblique

D. Fick's x-axis; Right Superior Rectus

See Answer in Part B → Q24

@Neet_pg_bot

Q25: Which physiological principle explains that during eye movement, an increase in neural stimulation to a contracting muscle is accompanied by a proportional decrease in stimulation to its opposing muscle? (Q853507)

A. Hering’s Law

B. Sherrington’s Law

C. Henle’s Law

D. Donder’s Law

See Answer in Part B → Q25

@Neet_pg_bot

Q26: A 25-year-old patient presents with a condition in which, upon examination, the left eye fails to adduct during right gaze. The patient reports diplopia, which is more noticeable when looking to the right. Which extraocular muscle is most likely overacting in this patient's condition? (Q853506)

A. Right Medial Rectus

B. Left Lateral Rectus

C. Right Lateral Rectus

D. Left Medial Rectus

See Answer in Part B → Q26

@Neet_pg_bot

Q27: A 45-year-old patient presents with vertical diplopia that worsens when looking down, especially while walking downstairs. Examination reveals weakness in intorsion, depression, and abduction of the affected eye. Which muscle and cranial nerve are most likely affected? (Q853505)

A. Superior Rectus; CN III

B. Inferior Oblique; CN III

C. Superior Oblique; CN IV

D. Inferior Rectus; CN IV

See Answer in Part B → Q27

@Neet_pg_bot

• [07_Squint _ Strabismus] 016 Previous Year Questions

« Back to Folder Directory

[07_Squint _ Strabismus] 016 Previous Year Questions — Quick Jump

Q1 (p. 231)	*Q2 (p. 231)*	*Q3 (p. 232)*	*Q4 (p. 232)*	*Q5 (p. 233)*	*Q6 (p. 234)*
Q7 (p. 235)	*Q8 (p. 235)*	*Q9 (p. 236)*	*Q10 (p. 236)*	*Q11 (p. 237)*	*Q12 (p. 237)*
Q13 (p. 237)	*Q14 (p. 238)*	*Q15 (p. 238)*	*Q16 (p. 238)*	*Q17 (p. 239)*	*Q18 (p. 239)*

Q1: What is the function of the superior oblique muscle? (Q855209)

A. Intorsion, abduction, depression

B. Intorsion, adduction, elevation

C. Extorsion, abduction, depression

D. Extorsion, adduction, elevation

See Answer in Part B → Q1

@Neet_pg_bot

Q2: In a 40-year-old patient with diabetes, a squint was observed. The secondary deviation was higher than the primary deviation during examination. The forced duction test yielded negative results. Which type of squint is expected to be predominant in this individual? (Q858509)

A. Comitant squint

B. Restrictive squint

C. Paralytic squint

D. Pseudosquint

See Answer in Part B → Q2

@Neet_pg_bot

Q3: The movement lost on left gaze as depicted in the image is seen in? (Q855030)

A. 3rd CN palsy

B. 4th CN palsy

C. 6th CN palsy

D. 7th CN palsy



See Answer in Part B → Q3

@Neet_pg_bot

Q4: Which of the following conditions is associated with the finding observed by a physician during an eye examination of a small child? (Q855023)

A. Myopia

B. Astigmatism

C. Hyperopia

D. Emmetropia



See Answer in Part B → Q4

@Neet_pg_bot


Q5: The chart depicted in the image is _____? (Q855387)

A. Hess chart

B. Snellen chart

C. Pelli-Robson chart

D. ETDRS chart



See Answer in Part B → Q5

@Neet_pg_bot


Q6: Identify the condition (Q855339)

A. Medial rectus palsy

B. Trochlear nerve palsy

C. Abducens nerve palsy

D. Oculomotor nerve palsy



See Answer in Part B → Q6

@Neet_pg_bot

Q7: What is the likely diagnosis for a 5-year-old girl presenting with decreased best corrected visual acuity in the right eye after a school vision screening revealed diminution of acuity in the right eye without any examination findings of the eye and visual pathway? (Q857227)

A. Amblyopia

B. Retinoblastoma

C. Congenital cataract

D. Brain lesion

See Answer in Part B → Q7

@Neet_pg_bot

Q8: What is the recommended treatment for a 3-year-old child with amblyopia in the right eye? (Q857188)

A. Observe for 3 months

B. Occlusion of the right eye

C. Occlusion of the left eye

D. Surgical correction

See Answer in Part B → Q8

@Neet_pg_bot


Q9: A 60-year-old male presents with left eyelid drooping and experiencing double vision. Upon examination, it was observed that the pupil in the affected eye was dilated. The following eye movement examination was conducted. Which nerve is most likely injured in this case? (Q857182)

A. Optic nerve

B. Oculomotor

C. Trochlear

D. Abducens



See Answer in Part B → Q9

@Neet_pg_bot

Q10: The cover-uncover test is performed to diagnose various eye conditions. Which of the following is not detected by this test? (Q852497)

A. Latent Squint

B. Manifest Squint

C. Lateral Nystagmus

D. Amblyopia

See Answer in Part B → Q10

@Neet_pg_bot

Q11: What is the recommended treatment for a 3-year-old child with amblyopia in the right eye? (Q858154)

A. Observation

B. Spectacles or refraction with a near addition

C. Occlusion Therapy

D. Radial keratotomy

See Answer in Part B → Q11

@Neet_pg_bot

Q12: Which drug is commonly used for refraction in a child with accommodative squint? (Q858120)

A. 1% Atropine

B. Cyclopentolate

C. Tropicamide

D. Homatropine

See Answer in Part B → Q12

@Neet_pg_bot

Q13: Which nerve innervates the lateral rectus muscle? (Q860162)

A. Sixth nerve

B. Seventh nerve

C. Third nerve

D. Fourth nerve

See Answer in Part B → Q13

@Neet_pg_bot

Q14: The patient has an infection of the cavernous sinus, involving the nerve located on its medial side. To confirm the involvement of the abducens nerve, you will ask the patient to look in which direction? (Q860509)

A. Superior

B. Inferior

C. Medial

D. Lateral

See Answer in Part B → Q14

@Neet_pg_bot

Q15: What is the probable cause of uniocular diplopia in a young boy who has experienced trauma to his right eye? (Q793357)

A. Paresis of extraocular muscle

B. Displacement of one eyeball

C. Symblepharon

D. Iridodialysis

See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 50-year-old female patient visits the ophthalmology clinic with complaints of experiencing vertical diplopia while walking downstairs. Upon further inquiry, it is noted that the patient tends to tilt her head to one side in order to alleviate the

diplopia. Which cranial nerve palsy should be suspected in this patient? (Q820672)

A. 3rd nerve

B. 6th nerve

C. 7th nerve

D. 4th nerve

See Answer in Part B → Q16

@Neet_pg_bot

Q17: Which drug is commonly used for refraction in a child with accommodative squint? (Q820650)

A. 1% Atropine

B. Cyclopentolate

C. Tropicamide

D. Homatropine

See Answer in Part B → Q17

@Neet_pg_bot

Q18: What are the yoke muscles for Levodepression? (Q820639)

A. LSO and RIR

B. RSO and LIR

C. RIO and LIR

D. LIO and RIR

See Answer in Part B → Q18

@Neet_pg_bot

• [08_Cornea] 017 Cornea- I

« Back to Folder Directory

[08_Cornea] 017 Cornea- I — Quick Jump

Q1 (p. 240)	*Q2 (p. 241)*	*Q3 (p. 245)*	*Q4 (p. 245)*	*Q5 (p. 245)*	*Q6 (p. 246)*
Q7 (p. 246)	*Q8 (p. 247)*	*Q9 (p. 247)*	*Q10 (p. 247)*	*Q11 (p. 248)*	*Q12 (p. 248)*
Q13 (p. 251)	*Q14 (p. 251)*	*Q15 (p. 252)*	*Q16 (p. 252)*	*Q17 (p. 253)*	*Q18 (p. 254)*
Q19 (p. 255)	*Q20 (p. 256)*	*Q21 (p. 256)*	*Q22 (p. 256)*	*Q23 (p. 257)*	*Q24 (p. 257)*
Q25 (p. 257)	*Q26 (p. 258)*	*Q27 (p. 258)*	*Q28 (p. 259)*	*Q29 (p. 259)*	*Q30 (p. 260)*
Q31 (p. 260)	*Q32 (p. 261)*	*Q33 (p. 261)*	*Q34 (p. 262)*	*Q35 (p. 262)*	*Q36 (p. 263)*
Q37 (p. 263)	*Q38 (p. 263)*	*Q39 (p. 264)*	*Q40 (p. 264)*	*Q41 (p. 265)*	*Q42 (p. 265)*
Q43 (p. 265)	*Q44 (p. 266)*				


Q1: A 45 y/o patient presents with worsening vision, mild pain, and redness 6 months after Penetrating Keratoplasty. Examination reveals the following. Which feature is most indicative of endothelial rejection? {{caption_text}} (Q853776)

A. Increased Flare and Cells

B. Kayes dots

C. Khodadoust Line

D. Epithelial Rejection Line



See Answer in Part B → Q1

@Neet_pg_bot


Q2: Match the Following: 1) a) Bacterial Keratitis 2) b) Superficial Punctate Keratitis 3) c) Pseudodendritis 4) d) Dendritic Ulcer 5) e) Nummular Keratitis 6) f) Fungal Keratitis (Q853763)


A. 1-b; 2-e; 3-c; 4-f; 5-d; 6-a


B. 1-c; 2-e; 3-d; 4-f; 5-a; 6-b

C. 1-e; 2-c; 3-b; 4-f; 5-d; 6-a


D. 1-c; 2-e; 3-b; 4-f; 5-d; 6-a



@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot





See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 28-year-old female contact lens wearer with severe eye pain and blurred vision shows a ring-shaped corneal infiltrate, and radial keratoneuritis on examination. Corneal scraping showed the presence of cysts. What is the most appropriate initial treatment? (Q853768)

A. Topical moxifloxacin

B. Oral acyclovir

C. Topical polyhexamethylene biguanide (PHMB)

D. Topical corticosteroids

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 23 y/o male presents with B/L eye redness, pain, and severe blurring of vision. He has a history of congenital syphilis. O/E, there is a characteristic pinkish-red discoloration of the cornea with deep stromal vascularization. Which of the following findings is most likely associated with this patient's condition during the healing stage? (Q853769)

A. Pseudodendrites

B. Ghost vessels

C. Ring-shaped abscess

D. Radial keratoneuritis

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 19 y/o female presents with a history of progressive worsening of vision that is not fully corrected by glasses. After evaluation, she is diagnosed with keratoconus. Which of the following examination findings are seen in keratoconus? Munson's sign Scissor reflex Vogt striae Fleischer ring (Q853770)

A. 1,2,3,4

B. 1,3,4

C. 2,3,4

D. 1,4

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Question: A 22 y/o male with a history of worsening vision despite wearing rigid gas-permeable lenses now complains of increased visual distortion. O/E, his corneal topography shows progressive ectasia, and he is advised to consider treatment options to halt disease progression. Which of the following treatments is most appropriate to stabilize the condition and improve contact lens tolerance in this patient? (Q853771)

A. Spectacle correction

B. Intracorneal ring segments (INTACS)

C. Corneal collagen cross-linking (CXL)

D. Penetrating keratoplasty

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 50 y/o patient presents for a corneal transplant evaluation. The patient has a history of systemic infections, including tuberculosis, and is seeking a corneal graft for a therapeutic indication. Which of the following is an absolute contraindication for cornea donation in this patient? (Q853773)

A. Death due to an unknown cause

B. Previous history of ocular surgery

C. Ocular inflammation

D. Retinoblastoma

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A donor cornea is harvested and needs to be preserved for a planned corneal transplant for 10 days. Which of the following storage methods is most appropriate for preserving the donor cornea? (Q853774)

A. Moist Chamber Method

B. McCarey Kaufman (M-K) Medium

C. Optisol GS

D. Cryopreservation

See Answer in Part B → Q8

@Neet_pg_bot

Q9: Which of the following is not an indication of Penetrating Keratoplasty (PK)? (Q853775)

A. Corneal opacity

B. Cosmetic improvement of the eye

C. Advanced keratoconus

D. Secondary glaucoma

See Answer in Part B → Q9

@Neet_pg_bot

Q10: Which layer of the cornea is primarily responsible for maintaining its transparency by regulating fluid content? (Q853607)

A. Stroma

B. Endothelium

C. Epithelium

D. Bowman's Membrane

See Answer in Part B → Q10

@Neet_pg_bot

Q11: All of the following statements about lamellar keratoplasty are true except: (Q853777)

A. It is indicated in pseudophakic bullous keratopathy

B. It has lower risk of graft rejection compared to Penetrating keratoplasty

C. Deep Anterior Lamellar Keratoplasty (DALK) replaces 80-90% of the anterior stroma and is suitable when the endothelium is normal.

D. Descemet’s Stripping Automated Endothelial Keratoplasty (DSAEK) is performed when the endothelium is defective.

See Answer in Part B → Q11

@Neet_pg_bot


Q12: Match the following: 1) Arcus senilis a) 2) Band shaped Keratopatthy b) {{caption_text}} 3) Vortex keratopathy c) 4) Salzmann nodular degeneration d) {{caption_text}} (Q853791)

A. 1-b, 2-d, 3-a, 4-c


B. 1-d, 2-a, 3-c, 4-b


C. 1-c, 2-d, 3-b, 4-a

D. 1-a, 2-b, 3-c, 4-d


@Neet_pg_bot


@Neet_pg_bot





See Answer in Part B → Q12

@Neet_pg_bot

Q13: Which of the following statements about corneal degeneration is incorrect? (Q853792)

A. Salzmann nodular degeneration affects the superficial stroma and Bowman’s membrane.

B. Vortex keratopathy is generally asymptomatic and associated with indomethacin intake.

C. Lipid keratopathy involves both intracellular and extracellular fat deposits.

D. Vortex keratopathy remains permanent even after discontinuing the offending medication.

See Answer in Part B → Q13

@Neet_pg_bot


Q14: A 45 y/o male presents with progressively worsening vision and evident discomfort in his left eye over the past year. O/E, the following findings were observed, which caused significant blurring of his vision. {{caption_text}} Which of the following corneal opacities is most likely causing the greatest reduction in his vision? (Q853793)

A. Nebular corneal opacity

B. Macular corneal opacity

C. Leucomatous corneal opacity

D. Adherent leucoma



See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 25-year-old patient presents with a history of gradual vision loss. O/E, multiple milky, granular deposits are seen in the anterior corneal stroma, with clear cornea between the deposits. Which type of stromal corneal dystrophy is most likely? (Q853812)

A. Granular Corneal Dystrophy, Type 1 (GCD1)

B. Lattice Corneal Dystrophy, Type 1 (LCD1)

C. Macular Corneal Dystrophy (MCD)

D. Schnyder Corneal Dystrophy (SCD)

See Answer in Part B → Q15

@Neet_pg_bot

Q16: Which of the following conditions is most commonly associated with the development of a "beaten-metal" appearance on the cornea? (Q853813)

A. Posterior Polymorphous Corneal Dystrophy (PPCD)

B. Congenital Hereditary Endothelial Dystrophy (CHED)

C. Keratoconus

D. Fuchs Endothelial Corneal Dystrophy (FECD)

See Answer in Part B → Q16

@Neet_pg_bot


Q17: Match the following. A. {{caption_text}} 1. Rose bengal B. {{caption_text}} 2. Lissamine green C. {{caption_text}} 3. Fluorescein sodium (Q853814)


A. A-2, B-3, C-1


B. A-1, B-2, C-3

C. A-3, B-2, C-1

D. A-1, B-3, C-2







See Answer in Part B → Q17

@Neet_pg_bot

Q18: A young patient presents with this sign. Diagnosis is best made by: (Q882199)

A. Keratometry

B. Corneal topography

C. Pachymetry

D. A-scan



See Answer in Part B → Q18

@Neet_pg_bot

Q19: A 70 y/o woman presents with severe pain in her right eye and forehead, along with a rash that started 5 days ago. She was diagnosed with Herpes Zoster Ophthalmicus (HZO). Which of the following statements is true regarding the management of her condition? (Q853762)

A. Systemic steroids alone are sufficient to manage HZO

B. Antiviral therapy is started within 72 hours of rash onset.

C. Postherpetic neuralgia is less common in patients over 70 years of age.

D. Topical antiviral therapy is not recommended to manage eye involvement in HZO.

See Answer in Part B → Q19

@Neet_pg_bot

Q20: What is the average thickness of the central cornea? (Q853606)

A. 1 to 1.2 mm

B. 0.8 to 0.9 mm

C. 0.5 to 0.6 mm

D. 7.8 mm

See Answer in Part B → Q20

@Neet_pg_bot

Q21: What is the primary metabolism of cornea? (Q853608)

A. Anaerobic

B. Aerobic.

C. Both anaerobic and aerobic

D. None of the above

See Answer in Part B → Q21

@Neet_pg_bot

Q22: Which of the following factors is not responsible for maintaining corneal transparency? (Q853609)

A. Corneal Epithelium and Tear Film

B. Avascularity of the Cornea

C. Corneal nerve supply

D. Endothelial Pumps

See Answer in Part B → Q22

@Neet_pg_bot

Q23: Match the following corneal assessment techniques with their correct descriptions: 1) Keratometry a) Measures both anterior and posterior corneal surfaces. 2) Placido's Disc b) Assesses the smoothness and curvature of the corneal surface. 3) Corneal Topography c) Evaluate the anterior curvature of the cornea and generate colour-coded maps. 4) Corneal Tomography (Pentacam) d) Measures the curvature of the central cornea. (Q853610)

A. 1-a, 2-c, 3-d, 4-b

B. 1-d, 2-b, 3-c, 4-a

C. 1-d, 2-a, 3-b, 4-c

D. 1-b, 2-a, 3-c, 4-d

See Answer in Part B → Q23

@Neet_pg_bot

Q24: Which of the following statements is false? (Q853611)

A. Pachymetry measures corneal thickness and is used for diagnosing glaucoma

B. Specular Microscopy is used for photographing the corneal endothelium

C. Aesthesiometer measures corneal sensation and is used to assess neurotrophic keratitis

D. Aesthesiometer measures corneal sensation by evaluating the second cranial nerve

See Answer in Part B → Q24

@Neet_pg_bot

Q25: A 32 y/o contact lens wearer presents with a red, painful eye, blurred vision, and increased sensitivity to light. Examination reveals lid swelling, conjunctival chemosis,

and a large corneal ulcer with greenish exudate, rapidly expanding with hypopyon. What is the most likely cause? (Q853657)

A. Staphylococcus aureus

B. Streptococcus pneumoniae

C. Pseudomonas aeruginosa

D. Enterobacteriaceae

See Answer in Part B → Q25

@Neet_pg_bot

Q26: A 60 y/o woman presents with a painful red eye, photophobia, and a history of chronic use of corticosteroid eye drops. Examination reveals a corneal ulcer with necrosis extending deep into the corneal stroma, approaching Descemet's membrane. Which type of bacterial keratitis is this patient most at risk for? (Q853658)

A. Localized Ulcer

B. Perforating Ulcer

C. Adherent Leucoma

D. Sloughing Ulcer

See Answer in Part B → Q26

@Neet_pg_bot

Q27: Which of the following statements about the treatment of bacterial keratitis is incorrect? (Q853659)

A. Cycloplegics like atropine can be used to reduce ciliary spasms and pain.

B. Ciprofloxacin (750mg twice daily) is given when the sclera gets involved.

C. Topical steroids should be started immediately to reduce inflammation.

D. Duo therapy with fortified cefazolin and tobramycin provides broad-spectrum coverage.

See Answer in Part B → Q27

@Neet_pg_bot

Q28: A 60 y/o man from a tropical region presents with a painful, red eye and a corneal ulcer following recent exposure to decaying vegetable matter. Which is the most common fungal infection affecting the cornea? (Q853755)

A. Candida albicans

B. Fusarium

C. Mucormycosis

D. Aspergillus fumigatus

See Answer in Part B → Q28

@Neet_pg_bot


Q29: A 30 y/o contact lens wearer presents with very severe eye pain, blurred vision, and discomfort in his right eye. O/E, the following corneal finding is noted. Based on the image and the clinical presentation, which of the following is the most likely diagnosis? (Q853764)

A. Herpes Simplex Keratitis

B. Fungal Keratitis

C. Acanthamoeba Keratitis

D. Bacterial Keratitis



See Answer in Part B → Q29

@Neet_pg_bot

Q30: A 65 y/o woman presents with a painful rash on her forehead and eyelid. She also reports blurred vision and photophobia. O/E, she has conjunctivitis, punctate epithelial keratitis, and iridocyclitis. Which of the following is a chronic ocular complication she is at risk of developing? (Q853761)

A. Episcleritis

B. Corneal Scarring

C. Conjunctivitis

D. Secondary Glaucoma

See Answer in Part B → Q30

@Neet_pg_bot

Q31: A 50 y/o male presents with decreased vision, redness, and a dull sensation in his R/E. He has a history of trigeminal neuralgia for which he underwent a Gasserian ganglion block. O/E, there is decreased corneal sensation and epithelial desquamation. Which of the following is the most likely cause of his symptoms? (Q853760)

A. Keratomalacia

B. Herpes Simplex Virus (HSV) Infection

C. Neurotrophic Keratopathy

D. Neuroparalytic Keratopathy

See Answer in Part B → Q31

@Neet_pg_bot

Q32: A 35 y/o patient presents with pain, redness, photophobia, and decreased vision in the right eye. Slit-lamp examination reveals a dendritic ulcer on the cornea. Which of the following is the most appropriate initial treatment for this patient? (Q853759)

A. Topical steroids and antibiotics

B. Oral acyclovir and lubricants

C. Topical acyclovir and cycloplegics

D. Oral steroids and topical antibiotics

See Answer in Part B → Q32

@Neet_pg_bot

Q33: A 45 y/o patient presents with eye redness, excessive tearing, and sensitivity to light. O/E, you observe dendritic ulcers and decreased corneal sensation. Based on these findings, which is the most likely etiological agent responsible for the patient's condition? (Q853758)

A. Adenovirus

B. Varicella Zoster Virus (VZV)

C. Herpes Simplex Virus (HSV)

D. Chlamydia trachomatis

See Answer in Part B → Q33

@Neet_pg_bot

Q34: A 45 y/o male presents with redness, pain, and decreased vision in his right eye. He reports a history of trauma to the eye with a plant material three days ago. O/E, there is a corneal ulcer with feathery edges and satellite lesions. Hypopyon is also noted. Which of the following is the most appropriate initial treatment for this patient? (Q853757)

A. Topical Moxifloxacin

B. Topical Natamycin

C. Oral Acyclovir

D. Intravitreal Amphotericin B

See Answer in Part B → Q34

@Neet_pg_bot

Q35: A 45 y/o man from a tropical region presents with a painful red eye, significant watering, and sensitivity to light after working in a garden with decaying plant material. The patient is diagnosed with keratomycosis. Which of the following is not an examination finding in keratomycosis? (Q853756)

A. Feathery extensions at the margins of the corneal ulcer

B. Extensive corneal vascularization

C. Sterile immune ring

D. Endothelial plaque

See Answer in Part B → Q35

@Neet_pg_bot

Q36: A 55 y/o male with poorly controlled diabetes presents with a non-healing corneal ulcer in his right eye, despite two weeks of appropriate antibiotic therapy. The ulcer shows signs of impending perforation with severe thinning of the cornea. Which of the following is the most appropriate next step in the management of this patient? (Q853660)

A. Increase the frequency of antibiotic eye drops.

B. Apply a pressure bandage and recommend strict bed rest.

C. Perform mechanical debridement and apply tissue adhesive glue.

D. Start topical steroids to reduce inflammation.

See Answer in Part B → Q36

@Neet_pg_bot

Q37: A 58-year-old patient is evaluated before cataract surgery. Specular microscopy shows normal endothelial morphology and adequate reserve to maintain corneal deturgescence. Which of the following values most closely represents the expected endothelial cell density in such an adult? (Q885150)

A. 800-1200 cells/mm²

B. 300-600 cells/mm²

C. 2400-3000 cells/mm²

D. 3500-4200 cells/mm²

See Answer in Part B → Q37

@Neet_pg_bot

Q38: A child presents for ophthalmic evaluation due to an abnormally small-appearing cornea. On examination, the horizontal corneal diameter is found to

be reduced. Microcornea is diagnosed when the horizontal corneal diameter measures below which of the following values? (Q885151)

A. Less than 12 mm

B. 10 mm or less

C. Less than 9 mm

D. Less than 11.5 mm

See Answer in Part B → Q38

@Neet_pg_bot

Q39: A patient develops rapidly progressive corneal ulceration despite no history of trauma or epithelial defect. Which organism is known to invade an intact corneal epithelium? (Q885152)

A. Streptococcus pneumoniae

B. Staphylococcus aureus

C. Neisseria gonorrhoeae

D. Moraxella lacunata

See Answer in Part B → Q39

@Neet_pg_bot

Q40: A young woman presents with sudden onset severe ocular pain, photophobia, tearing and a corneal epithelial defect with stromal infiltrate on slit-lamp examination. A bacterial corneal ulcer is diagnosed. What concentration of topical cefazolin is recommended in fortified therapy? (Q885153)

A. 2%

B. 5%

C. 10%

D. 1%

See Answer in Part B → Q40

@Neet_pg_bot

Q41: A farmer presents with gradually increasing ocular pain, photophobia and blurred vision after minor trauma with plant matter. Slit lamp shows a dry-looking corneal ulcer with feathery margins, satellite lesions and an endothelial plaque. A hypopyon is present. Which organism group is most likely responsible? (Q885155)

A. Gram-positive cocci

B. Filamentous fungi

C. Acanthamoeba

D. Herpes simplex virus

See Answer in Part B → Q41

@Neet_pg_bot

Q42: A farmer develops a corneal ulcer after injury with plant material. Slit-lamp shows a dry ulcer with feathery margins and stromal infiltrates arranged in a curvilinear pattern. Which type of hypersensitivity reaction is mainly responsible for these stromal infiltrates? (Q885156)

A. Type I (IgE-mediated reaction)

B. Type II (antibody-mediated cytotoxic reaction)

C. Type III (immune complex-mediated reaction)

D. Type IV (delayed cell-mediated reaction)

See Answer in Part B → Q42

@Neet_pg_bot

Q43: A patient with infectious keratitis presents to the OPD. In which of the following ocular conditions are topical steroids strictly contraindicated? (Q885157)

A. Vernal keratoconjunctivitis

B. Stromal keratitis

C. Atopic dermatitis-related ocular inflammation

D. Fungal corneal ulcer

See Answer in Part B → Q43

@Neet_pg_bot

Q44: A 38-year-old agricultural worker presents with gradually worsening eye pain and photophobia following trauma with vegetative matter. Slit-lamp examination reveals a dry corneal ulcer with feathery edges and satellite lesions. Corneal culture grows Fusarium species. Which of the following is the drug of choice for treatment? (Q885158)

A. Amphotericin B

B. Fluconazole

C. Natamycin

D. Nystatin

See Answer in Part B → Q44

@Neet_pg_bot

• [08_Cornea] 018 Previous Year Questions

« Back to Folder Directory

[08_Cornea] 018 Previous Year Questions — Quick Jump

Q1 (p. 267)	*Q2 (p. 267)*	*Q3 (p. 268)*	*Q4 (p. 268)*	*Q5 (p. 269)*	*Q6 (p. 269)*
Q7 (p. 270)	*Q8 (p. 270)*	*Q9 (p. 271)*	*Q10 (p. 271)*	*Q11 (p. 272)*	*Q12 (p. 272)*
Q13 (p. 273)	*Q14 (p. 273)*	*Q15 (p. 274)*	*Q16 (p. 274)*	*Q17 (p. 275)*	*Q18 (p. 276)*
Q19 (p. 277)	*Q20 (p. 277)*	*Q21 (p. 278)*	*Q22 (p. 278)*	*Q23 (p. 279)*	*Q24 (p. 279)*
Q25 (p. 280)	*Q26 (p. 281)*	*Q27 (p. 281)*	*Q28 (p. 281)*	*Q29 (p. 282)*	*Q30 (p. 282)*
Q31 (p. 283)	*Q32 (p. 283)*	*Q33 (p. 284)*	*Q34 (p. 285)*	*Q35 (p. 285)*	*Q36 (p. 286)*
Q37 (p. 287)	*Q38 (p. 287)*				

Q1: What is the probable diagnosis for a 30-year-old man who has a previous history of wearing contact lenses, and is currently experiencing pain, redness, and the presence of pus coming from his right eye? During examination, a greyish-white ulcer in the shape of a snake was observed on the lower portion of the cornea. Additionally, a hypopyon measuring 4 mm was found in the anterior chamber, along with a moderate flare reaction. (Q859337)

A. Viral corneal ulcer

B. Bacterial corneal ulcer

C. Fungal corneal ulcer

D. Parasitic corneal ulcer

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Which of the subsequent options does not exhibit a characteristic of keratoconus? (Q855518)

A. Haab's striae

B. Munson's sign

C. Irregular astigmatism

D. Scissoring reflex

See Answer in Part B → Q2

@Neet_pg_bot

Q3: What is the most probable diagnosis for a 23-year-old male who has been experiencing eye pain, sensitivity to light, and redness for the last two days, following an injury caused by brushing his eyes against leaves and thorns while gardening? (Q857661)

A. Simplex keratitis

B. Bacterial keratitis

C. Fungal keratitis

D. Acanthamoeba keratitis

See Answer in Part B → Q3

@Neet_pg_bot


Q4: In the image provided, which refractive error is observed in the depicted condition? (Q857753)

A. Hypermetropia and astigmatism

B. Myopia and astigmatism

C. Myopia and hypermetropia

D. Hypermetropia



See Answer in Part B → Q4

@Neet_pg_bot

Q5: Which of the following statements is incorrect regarding the corneal epithelium? (Q856916)

A. Regeneration is limited to the limbus

B. Bowman’s membrane regenerates

C. Lined by stratified squamous epithelium

D. Apical cells have microvilli

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Which layer is accountable for preserving the moisture and clarity of the cornea? (Q855015)

A. Descemet’s membrane

B. Endothelium

C. Stroma

D. Corneal epithelium

See Answer in Part B → Q6

@Neet_pg_bot

Q7: What is the distinguishing characteristic of a fungal ulcer? (Q855017)

A. Hypopyon

B. Dendritic ulcer on a fluorescein dye

C. Ring abscess

D. Satellite lesion

See Answer in Part B → Q7

@Neet_pg_bot


Q8: A 64-year-old woman with a history of trauma to the left eye 5 days back now presented with complaints of complete loss of vision. Upon examination the following findings were seen. What should be the subsequent course of action to manage this condition? (Q859303)

A. Keratoplasty

B. Keratotomy

C. LASIK

D. DCR



See Answer in Part B → Q8

@Neet_pg_bot

Q9: Which glycosaminoglycan is responsible for the transparency of the cornea? (Q859334)

A. Keratan sulfate

B. Hyaluronic acid

C. Heparan sulfate

D. Dermatan sulfate

See Answer in Part B → Q9

@Neet_pg_bot

Q10: A girl returns from a picnic and develops severe eye pain. She had been using contact lenses but did not follow proper hygiene practices. What is the most likely

causative organism? (Q866848)

A. Acanthamoeba

B. Plasmodium

C. Pseudomonas aeruginosa

D. Herpes simplex virus

See Answer in Part B → Q10

@Neet_pg_bot

Q11: Which of the following is true about pterygium? (Q859749)

A. Inflammation of cornea

B. Fatty deposition of sclera

C. Conjunctival invasion making a flap over cornea

D. Dead epithelial debris accumulation

See Answer in Part B → Q11

@Neet_pg_bot

Q12: For which of the subsequent diseases is collagen cross-linking therapy utilized? (Q859767)

A. Keratoconus

B. Corneal ulcer

C. Band keratopathy

D. Uveitis

See Answer in Part B → Q12

@Neet_pg_bot


Q13: What's the main risk for a 37-year-old man with blurred vision and severe pain in his right eye? He revealed that recently, he went swimming in a pond. During examination, a ring-like lesion was found, as seen in the image. (Q859769)

A. Diabetes

B. Hypertension

C. Steroids

D. Soft Contact lenses



See Answer in Part B → Q13

@Neet_pg_bot

Q14: What is the diagnosis for a female patient, aged 65, who has a history of chickenpox and presents at the clinic with the following findings? (Q859772)

A. Chicken pox exacerbation

B. Drug allergy to acyclovir

C. Herpes zoster ophthalmicus

D. Herpes zoster oticus



See Answer in Part B → Q14

@Neet_pg_bot

Q15: Arlt's line is seen in? (Q859776)

A. Vernal keratoconjunctivitis

B. Phlyctenular keratoconjunctivitis

C. Trachoma

D. Pterygium

See Answer in Part B → Q15

@Neet_pg_bot

Q16: What is the main underlying factor responsible for regular astigmatism when studying the optics of a patient? (Q860009)

A. Abnormal curvature of the cornea

B. Excessive eye rubbing

C. Prolonged exposure to bright light

D. Vitamin A deficiency

See Answer in Part B → Q16

@Neet_pg_bot


Q17: What does this image indicate? (Q856040)

A. Keratoconus

B. Vogt's limbal girdle

C. Keratoglobus

D. Corneal dystrophy



See Answer in Part B → Q17

@Neet_pg_bot

Q18: A 27-year-old male with myopic astigmatism presents with an acute onset of pain and a reduction in visual acuity. He has a history of frequent changes of glasses and eye rubbing. An image of the eye just after the procedure is shown below. Name the procedure that has been done? (Q879000)

A. DSAEK

B. DALK

C. Intracameral C3F8 injection

D. Corneal collagen crosslinking with riboflavin



See Answer in Part B → Q18

@Neet_pg_bot

Q19: Which of the following layers is associated with the beaten bronze appearance? (Q860188)

A. Endothelial Layer

B. Stroma

C. Sub-Epithelial Layer

D. Epithelial Layer

See Answer in Part B → Q19

@Neet_pg_bot

Q20: A patient presents with a painless ulcer in the eye. On examination, a long, branching ulcer with desquamated epithelium is seen on the cornea. What is the most likely diagnosis? (Q866837)

A. Fungal ulcer

B. Dendritic ulcer

C. Bacterial corneal ulcer

D. Neurotrophic ulcer

See Answer in Part B → Q20

@Neet_pg_bot

Q21: In keratoplasty, the commonly used corneal storage medium and suture material are: (Q865358)

A. MK medium and 10-0 nylon

B. Moist chamber and Vicryl

C. Optisol-GS and silk

D. Eusol-C and 6-0 prolene

See Answer in Part B → Q21

@Neet_pg_bot

Q22: Which condition is treated using an Intacs ring, as shown in the image? (Q866345)

A. Keratoconus

B. Cataract

C. Corneal ulcer

D. Glaucoma



See Answer in Part B → Q22

@Neet_pg_bot

Q23: A patient presents with guttate lesions in one eye and bullous keratopathy in the other eye. What is the most likely diagnosis? (Q866353)

A. Fuchs' endothelial dystrophy

B. Interstitial keratitis

C. Viral corneal ulcer

D. Keratoconjunctivitis

See Answer in Part B → Q23

@Neet_pg_bot

Q24: A corneal wisp test was performed, and the corneal reflex was elicited. Which of the following nerves is responsible for the afferent limb of this reflex? (Q852484)

A. Facial Nerve

B. Trigeminal Nerve

C. Oculomotor Nerve

D. Abducens Nerve

See Answer in Part B → Q24

@Neet_pg_bot

Q25: A patient sustained blunt trauma to the eye 6 months ago and now presents with blurring of vision. What is the most likely condition? (Q852489)

A. Iridodialysis

B. Ankyloblepharaon

C. Vossius Ring

D. Cycloiriditis



See Answer in Part B → Q25

@Neet_pg_bot

Q26: A patient presents with eyelid crusting and a thready sensation between the cornea and lids. What is the most likely diagnosis? (Q852500)

A. Chronic Blepharitis

B. Dry Eye Syndrome

C. Meibomian Gland Dysfunction

D. Conjunctivitis

See Answer in Part B → Q26

@Neet_pg_bot

Q27: A patient presents with redness, extreme eye pain, and diminished vision. The patient frequently wears contact lenses. Which of the following organisms is most likely responsible for her condition? (Q844050)

A. Staphylococcus aureus

B. Streptococcus pyogenes

C. Acanthamoeba

D. Listeria

See Answer in Part B → Q27

@Neet_pg_bot

Q28: Shield ulcer is seen in which of the following? (Q860516)

A. Spring catarrh

B. Phlyctenular conjunctivitis

C. Atopic keratoconjunctivitis

D. Autoimmune Conditions

See Answer in Part B → Q28

@Neet_pg_bot

Q29: A 40-year-old male patient, who works in an iron factory, presents to the clinic with complaints of visual disturbances. He reports that his vision has been gradually worsening and he has noticed a change in the color perception. On examination, the ophthalmologist observes the presence of iron deposits in the eye. What is the most likely ocular condition responsible for the iron deposits in the eye and the patient's visual disturbances? (Q820453)

A. Siderosis bulbi

B. Hemochromatosis

C. Retinitis pigmentosa

D. Age-related macular degeneration

See Answer in Part B → Q29

@Neet_pg_bot

Q30: A 45-year-old male patient presents to the ophthalmology clinic with a history of eye trauma with vegetative material .He complains of severe eye pain, redness, tearing, and photophobia. Microscopic examination of the corneal scraping reveals septate hyphae at an acute angle. What is the most likely causative agent responsible for this condition? (Q820465)

A. Fungal corneal ulcer

B. Bacterial corneal ulcer

C. Viral corneal ulcer

D. Acanthamoeba corneal ulcer

See Answer in Part B → Q30

@Neet_pg_bot


Q31: Which of the following combinations corresponds to the image given below? (Q854714)

A. Cobalt blue- Tryptan blue - Corneal ulcer

B. Cobalt green- Rose bengal stain- Corneal ulcer

C. Cobalt blue- Fluorescein stain- Corneal ulcer

D. Cobalt blue- Lissamine green- Corneal ulcer



See Answer in Part B → Q31

@Neet_pg_bot

Q32: The eye examination of a 20-year-old male shows a golden ring on the cornea. What of the following is the investigation of choice to diagnose the condition?

(Q855482)

A. Alpha-1-antitrypsin

B. Alpha-fetoprotein

C. Serum iron

D. Serum ceruloplasmin

See Answer in Part B → Q32

@Neet_pg_bot


Q33: What dye and filter are used in the image, as shown below? (Q855485)

A. Lissamine dye green filter

B. Lissamine dye cobalt blue filter

C. Fluorescein dye green filter

D. Fluorescein dye cobalt blue filter



See Answer in Part B → Q33

@Neet_pg_bot

Q34: A child with elevated liver enzymes presented to the OPD with the following eye finding on gonioscopy. It is associated with? (Q858142)

A. Copper

B. Zinc

C. Selenium

D. Iron



See Answer in Part B → Q34

@Neet_pg_bot


Q35: A patient presents to you with pain and redness in the right eye following an injury during welding. On examination, the image as seen below is observed. Which of the following is the appropriate next step in managing this patient? (Q858146)

A. Steroids and antibiotics

B. Surgical exploration

C. Keratoplasty

D. Removal by 26G needle



See Answer in Part B → Q35

@Neet_pg_bot


Q36: Please analyze the image provided and determine the specific abnormality or injury present . (Q793032)

A. Viral keratitis

B. Fungal keratitis

C. Bacterial keratitis

D. Interstitial keratitis



See Answer in Part B → Q36

@Neet_pg_bot

Q37: What is the probable diagnosis in a 27-year-old male with eye trauma from vegetative matter presenting with a corneal ulcer, hypopyon, satellite lesions, mild pain, photophobia, and redness? (Q793041)

A. Fungal keratitis

B. Acanthamoeba keratitis

C. Viral keratitis

D. Bacterial keratitis

See Answer in Part B → Q37

@Neet_pg_bot


Q38: What is the likely reason for a 21-year-old student to have ocular redness, feeling of a foreign object in the eye, and excessive tearing after wearing a soft contact lens, as indicated by the corneal scraping interface contrast microscopy image? (Q793034)

A. Acanthamoeba

B. Naegleria fowleri

C. Entamoeba histolytica

D. Balamuthia mandrillaris



See Answer in Part B → Q38

@Neet_pg_bot

• [09_Conjunctiva] 019 Conjunctiva - I

« Back to Folder Directory

[09_Conjunctiva] 019 Conjunctiva - I — Quick Jump

Q1 (p. 289)	*Q2 (p. 290)*	*Q3 (p. 290)*	*Q4 (p. 291)*	*Q5 (p. 291)*	*Q6 (p. 292)*
Q7 (p. 292)	*Q8 (p. 293)*	*Q9 (p. 294)*	*Q10 (p. 294)*	*Q11 (p. 295)*	*Q12 (p. 296)*
Q13 (p. 296)	*Q14 (p. 296)*	*Q15 (p. 297)*	*Q16 (p. 297)*	*Q17 (p. 298)*	*Q18 (p. 298)*
Q19 (p. 299)	*Q20 (p. 299)*	*Q21 (p. 300)*	*Q22 (p. 301)*	*Q23 (p. 302)*	*Q24 (p. 302)*
Q25 (p. 302)	*Q26 (p. 303)*				

Q1: A 6-year-old boy presents with a 2-week history of worsening bilateral eye redness, itching, and frequent lacrimation. On examination, hypertrophied, cobblestone-like papillae is seen on the palpebral conjunctiva. What is the most likely diagnosis? (Q853727)

A. Atopic Keratoconjunctivitis

B. Vernal Keratoconjunctivitis

C. Bacterial Conjunctivitis

D. Chlamydial Conjunctivitis



See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 25-year-old male presents to the ophthalmologic clinic for evaluation of his ocular lesions. After carefully assessing the eyes, the attending doctor diagnoses vernal keratoconjunctivitis. Which of the following statements is most likely associated with vernal keratoconjunctivitis? (Q882187)

A. It is a Follicular hypertrophy

B. Hallmark is the presence of Cobblestone papillae

C. Mild itching is present

D. Type IV hypersensitivity response

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 3-year-old child weighing 14 kg is brought to the pediatric clinic with complaints of night blindness and dryness of the eyes. On examination, the ophthalmologist observes conjunctival xerosis, Bitot's spots, and keratomalacia. What is the appropriate treatment for this child's condition? (Q882122)

A. Vitamin A: 2 lahks IU in oral, 0 th , 1 st , and 10 th day

B. Vitamin A: 2 lakhs IU injectable, 0 th , 1 st , and 14 th day

C. Vitamin A: 2 lakhs IU in oral, 0 th , 1 st , and 14 th day

D. Vitamin A: 2 lakhs IU injectable, 0 th , 1 st , and 10 th day

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 64-year-old patient with dry eyes undergoes several tests. Which finding best supports the diagnosis of dry eye syndrome? (Q853738)

A. Tear film break-up time of 15 seconds

B. Schirmer test showing 12 mm wetting

C. Extensive Rose Bengal staining

D. Normal tear film osmolality

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 60-year-old patient diagnosed with dry eye syndrome asks about the typical features of the condition. Which of the following is not commonly associated with dry eye? (Q853737)

A. Frothy tears along the lid margin

B. Punctate epithelial erosions on the cornea

C. Conjunctival xerosis and keratinization

D. Follicles on the conjunctiva

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 65-year-old woman with rheumatoid arthritis presents with dry eyes and mouth. Rose bengal staining shows conjunctival damage, and tear analysis reveals a tear-to-lysozyme ratio of 0.1. Which tear film layer is primarily deficient in this condition? (Q853736)

A. Mucinous layer

B. Aqueous layer

C. Lipid layer

D. All layers are equally affected

See Answer in Part B → Q6

@Neet_pg_bot


Q7: A 70-year-old man with extensive sun exposure has a yellowish, triangular patch on the nasal conjunctiva, as shown below. The lesion does not extend onto the cornea. What is the most likely diagnosis? (Q853735)

A. Pterygium

B. Pinguecula

C. Dermoid

D. Malignant melanoma



See Answer in Part B → Q7

@Neet_pg_bot


Q8: A 50-year-old male comes to the clinic with complaints of a small fleshy triangular mass in the eye. There is no history of pain or discharge. On examination, it is as in the image. What is true about this lesion? (Q853733)

A. This is a non-progressive lesion

B. It can be present in any quadrant of the eye

C. It is caused by thermochemical burn, trauma, and chronic inflammation

D. Histology shows collagenous degenerative changes in the vascularized subepithelial stroma



See Answer in Part B → Q8

@Neet_pg_bot

Q9: A 25-year-old male presents with difficulty seeing at night. Which of the following is not a cause of nyctalopia (night blindness)? (Q853732)

A. Gyrate atrophy

B. Pathological myopia

C. Cone dystrophy

D. Choroideremia

See Answer in Part B → Q9

@Neet_pg_bot

Q10: A 3-year-old child presents with symptoms of night blindness and dry eyes. On examination, there are Bitot's spots on the temporal side of both eyes. The child’s guardian reports that the child has had a history of frequent respiratory infections.

What is the most appropriate treatment regimen to prevent further ocular complications? (Q853731)

A. 100,000 IU of vitamin A orally on diagnosis, next day, and after 2 weeks.

B. Vitamin A eye drops and 10,000 IU orally for 1 month.

C. 200,000 IU of vitamin A orally on diagnosis, next day, and after 2 weeks.

D. Oral multivitamins with increased vitamin A-rich foods for 3 months.

See Answer in Part B → Q10

@Neet_pg_bot


Q11: A 5-year-old boy from a rural area has difficulty seeing at night. Upon examination, you notice the presence of triangular, dry-white spots on the bulbar conjunctiva in both eyes. Which of the following represents the correct diagnosis and grade of xerophthalmia based on the clinical findings? (Q853730)

A. Night Blindness (XN)

B. Conjunctival Xerosis (X1A)

C. Bitot's Spots (X1B)

D. Corneal Xerosis (X2)



See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 23-year-old male presents with a history of atopic dermatitis and asthma. He complains of tearing, burning, itching sensation and blurry vision for the past week. On examination, his eyelids are thickened and swollen, with signs of meibomitis. Slit-lamp examination reveals anterior subcapsular cataracts. What is the most appropriate first-line treatment for this patient's ocular symptoms? (Q853729)

A. Topical corticosteroids

B. Topical dual-acting antihistamine/mast cell stabilizer

C. Topical cyclosporine

D. Oral antihistamines

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 10-year-old child presents with itching, watering, and redness in both eyes. On examination, small round yellow nodules are observed near the limbus, and conjunctival congestion is noted. The child has a past h/o TB. Which of the following best explains the pathophysiology of this condition? (Q853728)

A. Type I hypersensitivity reaction caused by environmental allergens

B. Type IV hypersensitivity reaction to endogenous bacterial antigens

C. Type II hypersensitivity reaction triggered by viral infections

D. Autoimmune reaction against corneal epithelial cells

See Answer in Part B → Q13

@Neet_pg_bot

Q14: Which of the following statements is true regarding the epithelium of the conjunctiva? (Q853712)

A. The epithelium of the limbal conjunctiva is thinner than that of the fornices.

B. The conjunctival epithelium is lined by stratified non-keratinized epithelium.

C. The adenoid layer of the conjunctiva is present at birth and plays a role in neonatal immunity.

D. The dense fibrous layer is superficial to the adenoid layer and contains goblet cells.

See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 2-day-old infant is evaluated for discharge from the eyes. The discharge was initially serous, but it quickly turned into mucopurulent and then purulent. On examination, the eyes are bright red and swollen. What is the most likely causative agent here? (Q853726)

A. Chlamydia trachomatis

B. Silver nitrate

C. Treponema pallidum

D. Neisseria gonorrhoeae

See Answer in Part B → Q15

@Neet_pg_bot

Q16: A patient with characteristic features of trachoma is being evaluated. Which of the following would be the most appropriate systemic antibiotic to be used for the treatment of trachoma? (Q853725)

A. Erythromycin

B. Doxycycline

C. Amoxicillin

D. Azithromycin

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A patient presents with irritation of the eyes and foreign body sensation in the eyes. He is examined by the ophthalmologist, who observes the following in his upper eyelid. According to the WHO classification, which grade is the disease here? (Q853724)

A. TF

B. TI

C. CO

D. TS



See Answer in Part B → Q17

@Neet_pg_bot

Q18: An 8-year-old female is being evaluated for conjunctivitis. She gives a history of itching and mild foreign body sensation in the eyes, stickiness of the lids, and mucoid discharge from the eyes. On examination, there is conjunctival congestion. Which of the following signs would be seen in this stage of the disease? (Q853723)

A. Pannus siccus

B. Arlt's line

C. Herbert's follicle

D. Tylosis

See Answer in Part B → Q18

@Neet_pg_bot

Q19: A 28-year-old man develops sudden redness, Foreign body sensation and subconjunctival haemorrhage in both eyes after returning from a tropical region. His visual acuity is normal. What is the most likely causative agent? (Q853721)

A. Epstein-Barr virus

B. Coxsackievirus

C. Herpes simplex virus

D. Molluscum Contagiosum

See Answer in Part B → Q19

@Neet_pg_bot


Q20: A 30-year-old male presents with irritation and redness at both eye corners, foamy discharge, and skin maceration as shown below. He also has chronic nasal discharge. Which organism is most likely causing his symptoms? (Q853720)

A. Staphylococcus aureus

B. Moraxella axenfeld

C. Neisseria gonorrhoeae

D. Haemophilus influenzae



See Answer in Part B → Q20

@Neet_pg_bot

Q21: A 7-year-old child presents with the eye condition shown in the image. The child reports severe eye pain, redness, and eyelid swelling. Examination reveals an inflamed conjunctiva with thick, adherent membranes that bleed when removed. Which of the following organisms is most likely responsible for this presentation? (Q853719)

A. Adenovirus

B. Neisseria gonorrhoeae

C. Corynebacterium diphtheria

D. Beta-hemolytic streptococci



See Answer in Part B → Q21

@Neet_pg_bot

Q22: A 7-year-old presents with severe eye pain, intense redness, and copious purulent discharge from his right eye. He also has significant lid swelling and tender preauricular lymphadenopathy. Gram stain of the purulent discharge reveals Gram-negative intracellular diplococci. What is the next best step in management? (Q853718)

A. Intravenous ampicillin

B. Eye irrigation with saline

C. Intramuscular Ceftriaxone

D. Topical antibiotic drops

See Answer in Part B → Q22

@Neet_pg_bot

Q23: A 25-year-old woman with a history of soft contact lens use for the past 8 months presents with bilateral itching, redness, and watery discharge. Examination shows large, polygonal papillae on the upper tarsal conjunctiva. What is the most likely diagnosis? (Q853717)

A. Vernal Keratoconjunctivitis

B. Giant Papillary Conjunctivitis

C. Phlyctenular Conjunctivitis

D. Papillary Conjunctivitis

See Answer in Part B → Q23

@Neet_pg_bot

Q24: A 28-year-old woman reports bilateral itching, redness, and watery discharge that worsens in spring. Examination revealed mild conjunctival congestion with small papillae on the upper tarsal conjunctiva. What is the most likely diagnosis? (Q853716)

A. Viral Conjunctivitis

B. Allergic Conjunctivitis

C. Bacterial Conjunctivitis

D. Chlamydial Conjunctivitis

See Answer in Part B → Q24

@Neet_pg_bot

Q25: A 27-year-old male presents with redness, mild swelling, and foreign body sensation in the right eye for 2 days. On examination, no visual impairment is noted, and a diagnosis of acute conjunctivitis is made. Which of the following is not a common sign of conjunctivitis? (Q853715)

See Answer in Part B → Q25

@Neet_pg_bot

Q26: Which of the following glands are primarily responsible for secreting mucus to maintain ocular surface moisture? (Q853713)

A. Glands of Krause

B. Goblet cells

C. Glands of Wolfring

D. Glands of Moll

See Answer in Part B → Q26

@Neet_pg_bot

• [09_Conjunctiva] 020 Previous Year Questions

« Back to Folder Directory

[09_Conjunctiva] 020 Previous Year Questions — Quick Jump

Q1 (p. 304)	*Q2 (p. 304)*	*Q3 (p. 305)*	*Q4 (p. 305)*	*Q5 (p. 306)*	*Q6 (p. 307)*
Q7 (p. 308)	*Q8 (p. 308)*	*Q9 (p. 309)*	*Q10 (p. 310)*	*Q11 (p. 310)*	*Q12 (p. 311)*
Q13 (p. 312)	*Q14 (p. 313)*	*Q15 (p. 313)*	*Q16 (p. 314)*	*Q17 (p. 314)*	*Q18 (p. 315)*
Q19 (p. 315)	*Q20 (p. 316)*	*Q21 (p. 317)*	*Q22 (p. 317)*	*Q23 (p. 318)*	*Q24 (p. 318)*

Q1: What is the most frequently observed eye symptom associated with COVID-19? (Q855594)

A. Uveitis

B. Mucormycosis

C. Follicular conjunctivitis

D. Papillary conjunctivitis

See Answer in Part B → Q1

@Neet_pg_bot


Q2: A 30-year-old contact lens wearer presents with severe pain, redness and watery discharge from the eye. Which of the following stains is most useful for diagnosis? (Q878999)

A. Gram stain

B. Giemsa stain

C. Calcofluor white stain

D. Ziehl-Neelsen stain



See Answer in Part B → Q2

@Neet_pg_bot

Q3: In a patient who has had contact with a COVID-19 patient, presenting with symptoms such as redness, sensitivity to light, excessive tearing, and overall discomfort, a thorough examination was conducted with no eye abnormality. Based on this information, which specific part of the eye is most likely to be affected in this patient? (Q859774)

A. Conjunctiva

B. Uvea

C. Choroid

D. Retina

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 20-year-old male presents with fever and swelling of the left eyelid for the past two days. He has no history of trauma but experienced a mild flu-like illness one week ago. On examination, there is no chemosis or protrusion of the eyeball. Based on these observations, what is the most likely diagnosis? (Q859360)

A. Orbital cellulitis

B. Preseptal cellulitis

C. Panophthalmitis

D. Lid tumour

See Answer in Part B → Q4

@Neet_pg_bot


Q5: Which statement below does not accurately describe the clinical condition mentioned? (Q859319)

A. Abnormal activity of p53 is seen in the pterygium tissue

B. Decreased activity of tissue inhibitors of metalloproteinases is seen

C. Destruction of Bowman’s layer occurs when the pterygium invades the cornea

D. Elastotic degeneration and hyalinization of the connective tissue of the conjunctivaoccur.



See Answer in Part B → Q5

@Neet_pg_bot

Q6: A contact lens user presents with the following clinical picture. He has had watering, redness, and foreign body sensation in the eye for the past 2 months. What is the most probable diagnosis? (Q856131)

A. Follicular conjunctivitis

B. Spring catarrh

C. Giant papillary conjunctivitis

D. Trachoma



See Answer in Part B → Q6

@Neet_pg_bot

Q7: What is the most frequently observed eye symptom of COVID-19? (Q855382)

A. Retinal vein thrombosis

B. Viral keratouveitis

C. Follicular conjunctivitis

D. Pseudomembranous conjunctivitis

See Answer in Part B → Q7

@Neet_pg_bot

Q8: Which of the following treatment options is most frequently used for recurrent pterygium? (Q857717)

A. Conjunctival autograft

B. Beta-radiation

C. Observation

D. Bare sclera technique

See Answer in Part B → Q8

@Neet_pg_bot

Q9: What could be the most likely diagnosis for a 5-year-old boy presenting with severe itching, sensitivity to light, and sticky discharge from his eyes. Upon examination the following findings were seen? (Q857676)

A. Trachoma

B. Vernal keratoconjunctivitis

C. Phlyctenular conjunctivitis

D. Xerophthalmia



See Answer in Part B → Q9

@Neet_pg_bot

Q10: What is the likely diagnosis for a patient who wears contact lenses and has been experiencing eye watering, redness, and a sensation of having a foreign object in the eye for the last two months? (Q857236)

A. Follicular conjunctivitis

B. Spring catarrh

C. Giant papillary conjunctivitis

D. Acute trachoma



See Answer in Part B → Q10

@Neet_pg_bot

Q11: What could be the probable diagnosis for the eye lesion observed in a 21-year-old male patient? (Q857183)

A. Bitot’s spots

B. Pterygium

C. Phlyctenular conjunctivitis

D. Pinguecula



See Answer in Part B → Q11

@Neet_pg_bot


Q12: What is the causative agent of the sago-grain appearance of the conjunctiva? (Q857179)

A. Chlamydia trachomatis

B. Neisseria gonorrhoeae

C. Haemophilus influenzae

D. Vitamin A deficiency



See Answer in Part B → Q12

@Neet_pg_bot


Q13: The finding seen in the image is? (Q866832)

A. Horner-Trantas spots

B. Herbert’s pits

C. Pannus

D. Arlt’s line



See Answer in Part B → Q13

@Neet_pg_bot

Q14: Which of the following is not characteristic of pterygium? (Q855494)

A. Rate of recurrence is 95%

B. It is a fibrovascular growth

C. The parts of a pterygium are the head, neck and body

D. Has progressive and regressive types

See Answer in Part B → Q14

@Neet_pg_bot

Q15: What is the causative agent of the ocular condition as shown in the image? (Q857423)

A. Chlamydia

B. Mycoplasma

C. Pseudomonas

D. Staphylococcus



See Answer in Part B → Q15

@Neet_pg_bot

Q16: Which of the following is the normal conjunctival flora? (Q860198)

A. Staph aureus

B. Staph Epidermidis

C. Strep. Pneumonia

D. H. influenzae

See Answer in Part B → Q16

@Neet_pg_bot

Q17: Which of the following is associated with Vitamin A deficiency? (Q860158)

A. Pinguecula

B. Bitot’s spots

C. Pterygium

D. Stocker’s line

See Answer in Part B → Q17

@Neet_pg_bot

Q18: A young boy from a poor socioeconomic background presents with hyperemia and yellowish nodules at the limbus. What could be the likely diagnosis? (Q860510)

A. Bitot’s spots

B. Pterygium

C. Pinguecula

D. Phlyctenular conjunctivitis



See Answer in Part B → Q18

@Neet_pg_bot


Q19: Which of the following is seen in the given image? (Q844016)

A. Horner trantas spots

B. Herbert pits

C. Pannus

D. Corneal dystrophy



See Answer in Part B → Q19

@Neet_pg_bot


Q20: A patient presents to the outpatient department with the following findings. What is the most likely diagnosis? (Q843999)

A. Pterygium

B. Limbal dermoid

C. Pyogenic granuloma

D. Pinguecula



See Answer in Part B → Q20

@Neet_pg_bot

Q21: What is a common ocular finding in pterygium? (Q853136)

A. Eye pain

B. Macular edema

C. Purulent discharge

D. Astigmatism

See Answer in Part B → Q21

@Neet_pg_bot


Q22: What is the likely diagnosis for a 5-year-old boy who is experiencing severe itching, and thick discharge from his eyes . Upon examination the following finding as shown in the image was observed. (Q793046)

A. Trachoma

B. Vernal keratoconjunctivitis

C. Phylctenular conjunctivitis

D. Xerophthalmia



See Answer in Part B → Q22

@Neet_pg_bot

Q23: What is the most effective treatment for a patient who experiences considerable discomfort when reading text from paper or a screen, despite having normal visual acuity and other eye examination findings, except for a Schirmer's test result of 5mm? (Q793044)

A. Steroids

B. Anti-histamine

C. Lubricating eye drops

D. Prescribed spectacles

See Answer in Part B → Q23

@Neet_pg_bot

Q24: What is the recommended approach for managing a middle-aged woman with a progressive pterygium that is extending towards the pupillary area? (Q793037)

A. Keratoplasty

B. Excision with conjunctival autograft

C. Observation

D. Beta-radiation

See Answer in Part B → Q24

@Neet_pg_bot

• [10_Ocular Adnexa] 021 Lids And Disorders - I

« Back to Folder Directory

[10_Ocular Adnexa] 021 Lids And Disorders - I — Quick Jump

Q1 (p. 320)	*Q2 (p. 321)*	*Q3 (p. 322)*	*Q4 (p. 323)*	*Q5 (p. 324)*	*Q6 (p. 324)*
Q7 (p. 325)	*Q8 (p. 325)*	*Q9 (p. 326)*	*Q10 (p. 327)*	*Q11 (p. 327)*	*Q12 (p. 328)*
Q13 (p. 328)	*Q14 (p. 328)*	*Q15 (p. 329)*	*Q16 (p. 329)*	*Q17 (p. 330)*	*Q18 (p. 330)*

Q1: A 6-month old baby was brought to the clinic with the following finding. Which among the following statements is incorrect regarding the likely diagnosis? (Q853470)

A. More common in boys

B. Undergoes spontaneous regression in majority

C. Oral propranolol can be used for treating this condition

D. Most common orbital tumor of children.



See Answer in Part B → Q1


Q2: Match the image with the condition: Image Condition a. Chronic painless swelling 1. Stye b. Acute painful swelling 2. Chalazion c. Acute painful Swelling 3. Hordeolum Internum (Q853478)

A. a-2,b-3,c-1


B. a-1,b-2,c-3

C. a-3,b-2,c-1

D. a-1,b-3,c-2







See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which is the most common organism causing bacterial blepharitis? (Q853477)

A. Staphylococci

B. Streptococci

C. Propionibacterium acnes

D. Moraxella

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 3-year-old girl has been having difficulty opening her left eye since birth. An image is given below. What is the condition? (Q853476)

A. Symblepharon

B. Ankyloblepharon

C. Blepharitis

D. Blepharophimosis



See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 45-year-old woman with hypothyroidism is now complaining of loss of eyelashes. What is the diagnosis? (Q853475)

A. Madarosis

B. Poliosis

C. Tylosis

D. Trichiasis

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A neonatologist has noted the following finding in a newborn baby. What is the condition called? (Q853474)

A. Trichiasis

B. Pseudotrichiasis

C. Distichiasis

D. None of the above



See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 70-year-old patient presents with complaints of epiphora. On examination, there is an eversion of the punctum noted. Which among the following is incorrect regarding the type of ectropion seen in this patient? (Q853473)

A. Most common type of ectropion

B. Can cause lagophthalmos

C. Age-related changes are implicated in etiopathogenesis

D. Starts from lateral eyelid and progresses medially

See Answer in Part B → Q7

@Neet_pg_bot

Q8: Match the following types of entropion with their correct pair 1) Mechanical entropion A) Trachoma 2) Cicatricial entropion B) Phthisis bulbi 3) Congenital entropion C) Senile entropion 4) Involutional entropion D) Most commonly involves lower eyelid (Q853472)

A. 1-A, 2-B, 3-C, 4-D

B. 1-C, 2-A, 3-B, 4-D

C. 1-B, 2-A, 3-D, 4-C

D. 1-B, 2-A, 3-C, 4-D

See Answer in Part B → Q8

@Neet_pg_bot

Q9: A 75-year-old male patient presents with the following ocular findings and complaints of watering and photophobia. On examination, there are signs of minimal horizontal lid laxity. Which of the following surgical procedures would be the most appropriate for a long-term correction of his condition? (Q853471)

A. Transverse sutures

B. Weis procedure

C. Jones operation

D. Quickert procedure



See Answer in Part B → Q9

@Neet_pg_bot

Q10: Which among the following structures is part of the anterior lamella of the eyelid? (Q853461)

A. Orbicularis oculi

B. Orbital septum

C. Tarsal plate

D. Conjunctiva

See Answer in Part B → Q10

@Neet_pg_bot

Q11: All are causes of ipsilateral pseudoptosis except (Q853469)

A. Enophthalmos

B. Lid retraction

C. Hypotropia

D. Phthisis bulbi

See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 50 year male known smoker presented with hemoptysis and cough. A lung cancer was suspected. On examination, the patient had mild ptosis with good levator function. What would be the preferred management? (LPS-Levator palpebrae superioris) (Q853468)

A. Mullerectomy

B. Minimal resection of LPS

C. Maximal resection of LPS

D. Frontalis sling surgery

See Answer in Part B → Q12

@Neet_pg_bot

Q13: Which of the following measurements is considered indicative of moderate ptosis? (Q853467)

A. 1 mm droop

B. 2 mm droop

C. 3 mm droop

D. 4 mm droop

See Answer in Part B → Q13

@Neet_pg_bot

Q14: Which muscle is most commonly affected in congenital ptosis? (Q853466)

A. Orbicularis oculi

B. Levator palpebrae superioris

C. Frontalis

D. Superior oblique

See Answer in Part B → Q14

@Neet_pg_bot

Q15: Which of the following is a distinguishing feature of aponeurotic ptosis compared to other forms of ptosis? (Q853465)

A. Accompanied by ocular motor dysfunction

B. Presents with normal function of the levator muscle

C. Results from a neurological condition affecting the third cranial nerve

D. Usually involves a sudden onset of symptoms

See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 45-year-old female presents with bilateral ptosis that worsens throughout the day, particularly in the evening, and improves with rest. She also complains of double vision and generalized fatigue. Which of the following is the most likely cause of her symptoms? (Q853464)

A. Horner's Syndrome

B. Myasthenia Gravis

C. Oculopharyngeal Muscular Dystrophy

D. Botulism

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 60-year old male, heavy smoker, presents with persistent shoulder pain radiating to arm for several months. He presents with the following ocular findings. A chest x-ray reveals mass at the apex of the left lung. Which of the following symptoms is most likely to accompany this patient? (Q853463)

A. Mydriasis

B. Contralateral ptosis

C. Anhydrosis and miosis

D. Ophthalmoplegia



See Answer in Part B → Q17

@Neet_pg_bot


Q18: Identify the structure marked ‘X’ in the given image. (Q853462)

A. Orbicularis oculi muscle

B. Tarsal plate

C. Palpebral conjunctiva

D. Levator aponeurosis



See Answer in Part B → Q18

@Neet_pg_bot

• [10_Ocular Adnexa] 022 Previous Year Questions

« Back to Folder Directory

[10_Ocular Adnexa] 022 Previous Year Questions — Quick Jump

Q1 (p. 332) *Q2 (p. 332)* *Q3 (p. 333)* *Q4 (p. 333)* *Q5 (p. 333)* *Q6 (p. 334)*
Q7 (p. 334) *Q8 (p. 335)* *Q9 (p. 335)* *Q10 (p. 335)* *Q11 (p. 336)* *Q12 (p. 336)*
Q13 (p. 337) *Q14 (p. 337)* *Q15 (p. 338)*

Q1: A patient presents with painful lid swelling. On examination, a pus point is observed at the base of an eyelash. Which of the following glands is involved? (Q866843)

A. Zeis gland

B. Meibomian gland

C. Lacrimal gland

D. Moll gland

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 3-month-old child was brought to the outpatient department (OPD) with symptoms of photophobia and watering of the eyes. Which of the following is not a cause of this condition? (Q853128)

A. Congenital nasolacrimal duct obstruction

B. Congenital entropion

C. Congenital glaucoma

D. Congenital distichiasis

See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which muscle is implicated in ptosis in Horner syndrome? (Q843986)

A. Müller muscle

B. Levator aponeurosis

C. Horner muscle

D. Orbicularis oculi

See Answer in Part B → Q3

@Neet_pg_bot

Q4: What is the order of insertion of muscles on Annulus of Zinn? A. Medial rectus B. Inferior rectus C. Lateral rectus D. Superior rectus (Q860517)

A. A, B, C, D

B. B, C, A, D

C. D, C, A, B

D. C, A, B, D

See Answer in Part B → Q4

@Neet_pg_bot

Q5: All of the following are functions of superior oblique muscle except? (Q840068)

A. Adduction

B. Abduction

C. Depression

D. Intorsion

See Answer in Part B → Q5

@Neet_pg_bot

Q6: All of the following are features of oculomotor nerve palsy except? (Q840070)

A. Exotropia

B. Dilated pupil

C. Limited Abduction

D. Ptosis

See Answer in Part B → Q6

@Neet_pg_bot


Q7: Please identify the medical condition depicted in the provided image. (Q857239)

A. Entropion

B. Ectropion

C. Trichiasis

D. Ptosis



See Answer in Part B → Q7

@Neet_pg_bot

Q8: In the context of medical treatment, the V-Y procedure is employed for which of the subsequent conditions? (Q857681)

A. Entropion

B. Ectropion

C. Blepharophimosis

D. Ptosis

See Answer in Part B → Q8

@Neet_pg_bot

Q9: An 11-month-old baby was brought by his mother with complaints of the left upper eyelid moving up and down while breastfeeding that disappears when the baby is not breastfeeding. She gave a history of the child being diagnosed with left eye ptosis. What is this condition called? (Q857685)

A. Symblepharon

B. Blepharophimosis syndrome

C. Lagophthalmos

D. Marcus -Gunn jaw-winking

See Answer in Part B → Q9

@Neet_pg_bot

Q10: What is the most frequently occurring tumor found in the lacrimal gland? (Q855378)

A. Adenoid cystic carcinoma

B. Lymphoma

C. Pleomorphic adenoma

D. Mucoepidermoid carcinoma

See Answer in Part B → Q10

@Neet_pg_bot

Q11: What is the term for the additional row of eyelashes located behind the grey line? (Q858503)

A. Trichiasis

B. Distichiasis

C. Tylosis

D. Madarosis

See Answer in Part B → Q11

@Neet_pg_bot


Q12: Identify the instrument. (Q859327)

A. Evisceration spatula

B. Chalazion clamp

C. Enucleation spoon

D. Bone punch



See Answer in Part B → Q12

@Neet_pg_bot

Q13: What is the recommended treatment for a patient with Bell's palsy, who is experiencing symptoms of dry eye, redness, and irritation, along with the presence of multiple punctate epithelial erosions in the cornea as observed through a slit lamp examination? (Q859358)

A. Tarsorrhaphy

B. Contact lens

C. Corneal replacement

D. LASIK

See Answer in Part B → Q13

@Neet_pg_bot

Q14: In a patient presenting with recurrent chalazion, which cancer should be suspected? (Q820680)

A. Sebaceous cell carcinoma

B. Squamous cell carcinoma

C. Basal cell carcinoma

D. Kaposi’s sarcoma

See Answer in Part B → Q14

@Neet_pg_bot

Q15: What is the likely diagnosis for a patient who has a painless, firm nodule on the upper eyelid that is growing away from the lid margin? (Q793040)

A. Stye

B. Entropion

C. Chalazion

D. Blepharitis

See Answer in Part B → Q15

@Neet_pg_bot

• [11_Optics] 023 Optics-Ii

« Back to Folder Directory

[11_Optics] 023 Optics-Ii — Quick Jump

Q1 (p. 339)	*Q2 (p. 340)*	*Q3 (p. 340)*	*Q4 (p. 340)*	*Q5 (p. 341)*	*Q6 (p. 341)*
Q7 (p. 342)	*Q8 (p. 342)*	*Q9 (p. 342)*	*Q10 (p. 345)*	*Q11 (p. 347)*	*Q12 (p. 348)*
Q13 (p. 348)	*Q14 (p. 349)*	*Q15 (p. 349)*	*Q16 (p. 349)*	*Q17 (p. 350)*	*Q18 (p. 350)*


Q1: Worth's Four Dot Test for a 6 y/o child reveals that the child sees two red and three green lights as shown below in the image. Based on the test result, what is the most likely diagnosis in this child? (Q853632)

A. Binocular Single Vision

B. Left Eye Suppression

C. Right Eye Suppression

D. Diplopia



@Neet_pg_bot

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A 25-year-old patient presents with blurred vision and headaches. On examination, the patient is found to have regular astigmatism with the following prescription: Sphere: +2.00 D, Cylinder: -1.50 D, Axis: 90°.What is the refractive power of the eye along the 90° meridian? (Q853640)

A. +2.00 D

B. +0.50 D

C. +3.50 D

D. +1.25 D

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 45 y/o patient presents with difficulty in reading and blurred vision. An eye examination reveals astigmatism. Based on the classification of astigmatism, which of the following statements is accurate? (Q853639)

A. In With-the-Rule (WTR) astigmatism, the horizontal meridian has greater curvature than the vertical meridian.

B. Compound myopic astigmatism is characterized by both of the eye's two principalmeridians falling in front of the retina

C. Irregular astigmatism involves a predictable pattern of refractive power changes along the meridians.

D. Bi-oblique astigmatism has principal meridians that are perpendicular to each other.

See Answer in Part B → Q3

@Neet_pg_bot

Q4: Which of the following statements about common treatments for presbyopia is correct? (Q853638)

A. Bifocal lenses are designed to provide only distance vision and are not effective for near vision.

B. Multifocal IOLs (intraocular lenses) provide multiple focal points to help with both near and distant vision.

C. Conductive Keratoplasty (CK) offers permanent benefits for presbyopia.

D. Corneal multifocality treatment involves the replacement of the Natural lens with an IOL.

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 45 y/o patient presents with significant astigmatism and is seeking surgical correction. The patient is not a candidate for LASIK. Which of the following procedures is most suitable for correcting high degrees of astigmatism in this case? (Q853637)

A. Limbal Relaxing Incisions (LRI)

B. Photorefractive Keratectomy (PRK)

C. Toric Intraocular Lens (IOL) Implantation

D. Conductive Keratoplasty (CK)

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Which of the following surgical procedures is most suitable for correcting hypermetropia up to +4D and allows for faster visual recovery? (Q853636)

A. Photorefractive Keratectomy (PRK)

B. Conductive Keratoplasty (CK)

C. Laser Thermal Keratoplasty (LTK)

D. Laser in situ Keratomileusis (LASIK)

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 30-year-old patient with a history of myopia is considering LASIK surgery. Which of the following is NOT an indication of the procedure? (Q853635)

A. The patient's refraction has been stable for the past 14 months

B. The patient is over 20 years old

C. Presence of corneal ectasia

D. Myopia of -7 D

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 32 y/o man with -14 D myopia presents for a consultation regarding refractive surgery. He is not a candidate for laser-based procedures due to thin corneas. Which of the following is the most appropriate surgical option for his condition? (Q853634)

A. Phakic intraocular lens (IOL) implantation

B. Radial keratotomy (RK)

C. Clear lens extraction

D. Intracorneal ring (ICR) implantation

See Answer in Part B → Q8

@Neet_pg_bot

Q9: Match the following 1) Titmus fly test a) 2) TNO test b) 3) Frisby test c) 4) Random Dot E test d) (Q853633)

A. 1 - c,2 - b,3 - a,4 - d

B. 1 - b,2 - c,3 - a,4 - d

C. 1 - b,2 -c,3 - d,4 - a

D. 1 - c,2 - b,3 - d,4 - a



@Neet_pg_bot



@Neet_pg_bot



TNO Test

@Neet_pg_bot



@Neet_pg_bot

See Answer in Part B → Q9

@Neet_pg_bot

Q10: Match the following images appropriately: 1.Streak Retinoscope a) 2. Priestly Smith Retinoscope b) 3.Trial Frame c) 4.Refraction Box d) (Q853616)


A. 1- a, 2- d, 3- c, 4- b

B. 1- b, 2- c, 3- a, 4- d


C. 1- c, 2- a, 3- d, 4- b


D. 1- d, 2- c, 3- a, 4- b


MEDIK
@Neet_pg_bot


Occluder, Pin hole, Slit, Red-Green glass, Maddox rod etc
Convex cylindrical lenses
Concave cylindrical lenses
Prisms
Convex spherical lenses
Concave spherical lenses
Trial frame

@Neet_pg_bot





See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 7-year-old child is evaluated for binocular vision using a synoptophore. The child successfully perceives two different images simultaneously and can superimpose

incomplete but similar images into a single complete image. Which grade of binocular single vision is demonstrated by this child? (Q853631)

A. Grade I - Simultaneous Perception

B. Grade II - Fusion

C. Grade III - Stereopsis

D. None of the above

See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 7-year-old child with amblyopia in the right eye is prescribed occlusion therapy. The optometrist recommends patching the sound left eye for a certain number of hours each day. What is the typical recommended duration of patching for effective treatment in children with amblyopia? (Q853630)

A. 1-2 hours per day

B. 3-4 hours per day

C. 4-6 hours per day

D. 12 hours per day

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 7-year-old girl who had a H/0 congenital cataract is presenting with decreased vision in her right eye. Which type of amblyopia is most likely in this case? (Q853629)

A. Strabismic amblyopia

B. Toxic amblyopia

C. Anisometropic amblyopia

D. Stimulus deprivation amblyopia

See Answer in Part B → Q13

@Neet_pg_bot

Q14: Which of the following is an absolute optical indication for contact lens use? (Q853621)

A. Corneal ulcer

B. Bullous keratopathy

C. Unilateral aphakia

D. Aniridia

See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 25-year-old female with myopia is looking for contact lenses that will be comfortable for long hours of wear and require minimal maintenance. She is involved in outdoor activities and is seeking a lens that provides comfort, ease of use, and reduced risk of irritation over extended periods. Which type of contact lens would be most suitable for her? (Q853620)

A. Hard contact lenses

B. Rigid gas permeable lenses

C. Soft contact lenses

D. PMMA lenses

See Answer in Part B → Q15

@Neet_pg_bot

Q16: During subjective refraction, an optometrist uses a Jackson Cross Cylinder (JCC) lens as part of the assessment. Which of the following is the primary purpose of

this test? (Q853619)

A. To determine the spherical refractive error

B. To measure the visual acuity at distance

C. To verify the axis and power of the cylindrical correction

D. To test for binocular vision and eye alignment

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 38-year-old patient presents for an eye examination. During the duochrome test, the patient reports that they can see the red letters more clearly than the green letters. What is the most likely refractive error for this patient? (Q853618)

A. Hyperopia

B. Myopia

C. Astigmatism

D. Presbyopia

See Answer in Part B → Q17

@Neet_pg_bot

Q18: A patient presents with a spectacle prescription of -4.00 D sphere and +1.00 D cylinder at 90°. What is the transposed prescription? (Q853617)

A. -3.00 D sphere -1.00 D cylinder at 180°

B. -5.00 D sphere -1.00 D cylinder at 90°

C. +4.00 D sphere -1.00 D cylinder at 90°

D. -1.00 D sphere +4.00 D cylinder at 180°

See Answer in Part B → Q18

• [11_Optics] 024 Optics-I

« Back to Folder Directory

[11_Optics] 024 Optics-I — Quick Jump

Q1 (p. 352)	*Q2 (p. 352)*	*Q3 (p. 353)*	*Q4 (p. 353)*	*Q5 (p. 354)*	*Q6 (p. 354)*
Q7 (p. 355)	*Q8 (p. 355)*	*Q9 (p. 355)*	*Q10 (p. 356)*	*Q11 (p. 356)*	*Q12 (p. 357)*
Q13 (p. 357)	*Q14 (p. 358)*	*Q15 (p. 358)*	*Q16 (p. 359)*	*Q17 (p. 359)*	*Q18 (p. 360)*
Q19 (p. 360)	*Q20 (p. 360)*	*Q21 (p. 361)*	*Q22 (p. 361)*	*Q23 (p. 362)*	*Q24 (p. 365)*
Q25 (p. 367)					

Q1: Which of the following are features of direct ophthalmoscopy? Erect, virtual image is formed 2.5x magnification Large field of view Peripheral retinal lesions can be examined Routine fundus examination Semi-dark room required (Q853802)

A. 1,5,6

B. 1,2,3,4,5,6

C. 1,3,6

D. 3,4

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Match the following refractive prescriptions with their corresponding types of astigmatism: Prescription Astigmatism Type i) +2D Cyl 1. Simple hypermetropic ii) +1D Sph & +2D Cyl 2. Compound hypermetropic iii) -2D Sph & +3D Cyl @180° 3. Mixed with the rule iv) -2D Sph & -1D Cyl @90° 4. Compound myopic v) -1D Sph & +2D Cyl @90° 5. Mixed against the rule (Q853841)

A. i–1, ii–2, iii–3, iv–4, v–5

B. i–1, ii–2, iii–5, iv–4, v–3

C. i–2, ii–1, iii–5, iv–4, v–3

D. i–1, ii–3, iii–5, iv–2, v–4

See Answer in Part B → Q2

@Neet_pg_bot

Q3: All of the following are true regarding the procedure shown below except (Q853840)

A. It's an objective method for refraction testing

B. Red reflex moving with the movement of this instrument implies Hypermetropia

C. Results are unreliable if the patient’s accommodation is active

D. Cycloplegics are only indicated for children below 5 years of age



See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 50-year-old male presented to the ophthalmology clinic with complaints of difficulty in reading newspapers and books. Which among the following statements is inappropriate regarding the management of this patient? (Q853839)

A. Roughly +1D convex lenses can be prescribed

B. Coexisting refractive error (myopia/hypermetropia) should be corrected first

C. 1.25% Pilocarpine can be beneficial in treatment

D. Conductive keratoplasty can be performed

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 55-year-old patient presents with difficulty reading up close and notices that their vision becomes blurry when trying to focus on objects at near distances. On examination, the near point is found to be farther away than normal. Which of the following best explains the patient's condition? A) The patient has a reduced amplitude of accommodation B) The patient's far point is at infinity C) The patient has a myopic refractive error D) The patient’s near point has moved closer (Q853838)

A. The patient has a reduced amplitude of accommodation B) The patient's far point is at infinity C) The patient has a myopic refractive error D) The patient’s near point has moved closer

B. The patient's far point is at infinity

C. The patient has a myopic refractive error

D. The patient’s near point has moved closer

See Answer in Part B → Q5

@Neet_pg_bot

Q6: All of the following are changes not related to accommodation except (Q853837)

A. Anterior lens curvature almost remains the same

B. Posterior lens curvature is increased

C. Increased zonular tension

D. Ciliary muscle contraction

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A patient post cataract lens extraction refused IOL placement and hence was prescribed spectacles. Later, he presented with complaints of visual discomfort. Choose the right combination of effects the patient would have likely felt. 1) Pincushion effect of distortion 2) Barrel effect of distortion 3) There is magnification of the image 4) There is a minification of the image (Q853836)

A. 1,3

B. 1,4

C. 2,3

D. 2,4

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 45-year-old patient presents with blurry vision and difficulty focusing after undergoing cataract surgery 3 weeks ago. On examination, the patient has no lens in the eye and no evidence of intraocular lens (IOL) implantation. Which of the following is the most appropriate management for this patient? (Q853835)

A. Prescribe corrective glasses with a high refractive index

B. Fit the patient with an intraocular lens

C. Perform a keratoplasty to correct vision

D. Initiate topical corticosteroids for inflammation

See Answer in Part B → Q8

@Neet_pg_bot

Q9: In the case of astigmatism, the vertical meridian is more curved than the horizontal. What kind of lens should be given? (Q853834)

A. Concave cylindrical lens at 180°

B. Convex cylindrical lens at 180

C. Concave cylindrical lens at 90

D. None of the above

See Answer in Part B → Q9

@Neet_pg_bot

Q10: A few rays of light are focused on the retina in one meridian and a few others are focused in front of the retina in another meridian. What is the diagnosis? (Q853833)

A. Simple myopic astigmatism

B. Simple hypermetropic astigmatism

C. Mixed astigmatism

D. Compound myopic astigmatism

See Answer in Part B → Q10

@Neet_pg_bot

Q11: A 35-year-old patient presents with difficulty focusing on both near and distant objects. Examination reveals high hypermetropia. What is the expected fundoscopy finding (Q853832)

A. Smaller optic disc

B. Shotsilk retina

C. Pseudopapillitis

D. All of the above

See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 65-year-old patient presents with gradually worsening blurry vision over the past year with recent improvement in his near vision . On examination, the lens shows yellowish discolouration with increased opacity, consistent with nuclear sclerosis. What kind of myopia has he developed secondary to this? (Q853804)

A. Axial myopia

B. Curvatural myopia

C. Index myopia

D. Positional myopia

See Answer in Part B → Q12

@Neet_pg_bot


Q13: Identify X: (Q853803)

A. Angle alpha

B. Angle beta

C. Angle gamma

D. Angle kappa



@Neet_pg_bot

See Answer in Part B → Q13

@Neet_pg_bot

Q14: Snellen chart is used to assess which of the following? (Q853787)

A. Colour vision

B. Contrast sensitivity

C. Peripheral vision

D. Visual acuity

See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 65-year-old patient with a mature cataract presents for an eye examination. During the assessment, the ophthalmologist observes the absence of a specific Purkinje

image. Which Purkinje image is most likely to be absent in this case? (Q853801)

A. First Purkinje image

B. Second Purkinje image

C. Third Purkinje image

D. Fourth Purkinje image

See Answer in Part B → Q15

@Neet_pg_bot

Q16: Which of the following is the most sensitive test for both congenital and acquired colour vision defects? (Q853800)

A. Ishihara Test

B. Farnsworth-Munsell 100 Hue Test

C. Hardy-Rand-Rittler (HRR) Test

D. Holmgren Wool Test

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A 40-year-old woman presents with a 3-day history of painful vision loss in her right eye. She has a history of multiple sclerosis. On examination, her visual acuity is 20/30 in the right eye, and a RAPD is noted. She reports difficulty distinguishing red and green colors in the right eye, but her left eye is normal on color vision testing. Which of the following is the most likely cause of her acquired color blindness? (Q853799)

A. Retinal vein occlusion

B. Optic neuritis

C. Diabetic retinopathy

D. Macular degeneration

See Answer in Part B → Q17

@Neet_pg_bot

Q18: Which of the following statements regarding congenital colour blindness is TRUE? (Q853798)

A. It is more common in females than males.

B. It is an autosomal recessive condition.

C. It primarily affects the perception of red and green colours and is the most common type.

D. It always results in total loss of colour vision.

See Answer in Part B → Q18

@Neet_pg_bot

Q19: Which type of colour blindness is caused by the absence of green cones in the retina? (Q853797)

A. Protanopia

B. Deuteranopia

C. Tritanopia

D. Achromatopsia

See Answer in Part B → Q19

@Neet_pg_bot

Q20: At what age do infants typically begin to perceive and distinguish between different colors? (Q853796)

A. At birth

B. 1-2 months

C. 3-4 months

D. 6-9 months

See Answer in Part B → Q20

@Neet_pg_bot

Q21: A 55-year-old male presents with complaints of difficulty driving at night and reading in dim light, despite having normal visual acuity during his eye exam. He underwent cataract surgery six months ago but still experiences these visual issues. Which of the following tests is most appropriate to assess his current visual problem? (Q853795)

A. Snellen Chart

B. Ishihara Test

C. Letter Contrast Sensitivity Test

D. Visual Field Test

See Answer in Part B → Q21

@Neet_pg_bot

Q22: A patient’s visual acuity improves significantly when looking through a small, opaque disc with a tiny hole, as shown in the image below. What is the likely cause of this improvement in visual acuity? (Q853794)

A. Cataracts

B. Refractive error

C. Optic nerve damage

D. Macular degeneration



See Answer in Part B → Q22

@Neet_pg_bot

Q23: Match the following visual acuity charts used to assess children of preschool age. 1) Allen test cards a) 2) HOTV test b) 3) Stycar Test c) 4) Cardiff Acuity test d) (Q853790)

A. 1-a; 2-b; 3-c; 4-d


B. 1-a; 2-b; 4-c; 3-d

C. 4-a; 2-b; 3-c; 1-d


D. 1-a; 3-b; 2-c; 4-d




@Neet_pg_bot



@Neet_pg_bot





See Answer in Part B → Q23

@Neet_pg_bot


Q24: Which among the following is used to check visual acuity in children less than 1 year? 1) 2) 3) 4) (Q853789)

A. 1 and 4


B. 2 and 3

C. All except 4


D. All of the above




@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot



See Answer in Part B → Q24

@Neet_pg_bot


Q25: Which among the following is used to check visual acuity in illiterate adults? 1) 2) 3) 4) (Q853788)

A. 1 and 3


B. 2 and 4


C. All except 1

D. None of the above


E
1 20/200
F P
2 20/100
T O Z
3 20/70
L P E D
4 20/50
P E C F D
5 20/40
E D F C Z P
6 20/30
F E L O P Z D
7 20/25
D E F P O T E C
8 20/20
L E F O D P C T
9
F D P L T C E O
10
F E Z O L C F T D
11
@Neet_pg_bot


E
D·36
D·24
D·18
D·12
D·9
D·6





See Answer in Part B → Q25

@Neet_pg_bot

• [11_Optics] 025 Previous Year Questions

« Back to Folder Directory

[11_Optics] 025 Previous Year Questions — Quick Jump

Q1 (p. 371)	*Q2 (p. 371)*	*Q3 (p. 372)*	*Q4 (p. 372)*	*Q5 (p. 373)*	*Q6 (p. 373)*
Q7 (p. 374)	*Q8 (p. 374)*	*Q9 (p. 375)*	*Q10 (p. 375)*	*Q11 (p. 375)*	*Q12 (p. 376)*
Q13 (p. 377)	*Q14 (p. 377)*	*Q15 (p. 378)*	*Q16 (p. 379)*	*Q17 (p. 379)*	*Q18 (p. 380)*
Q19 (p. 380)	*Q20 (p. 380)*	*Q21 (p. 381)*	*Q22 (p. 381)*	*Q23 (p. 382)*	*Q24 (p. 382)*
Q25 (p. 383)	*Q26 (p. 383)*				

Q1: A patient was prescribed with -1D spherical and -2D cylindrical at 180 degrees. What type of refractive error does this patient have? (Q857422)

A. With the rule astigmatism

B. Against the rule astigmatism

C. Simple astigmatism

D. Mixed astigmatism

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Given below are some tests for visual acuity. Which of these is used for a pre-verbal child? ETDRS chart Optokinetic nystagmus Teller’s visual acuity chart Landolt’s C chart (Q878998)

A. 2,3

B. 1,2

C. 2,3,4

D. 2,4

See Answer in Part B → Q2

@Neet_pg_bot

Q3: What would be the suitable course of action for managing a 15-year-old girl who is non-adherent to wearing spectacles for her myopic astigmatism? (Q856123)

A. LASIK

B. Spherical equivalent spectacles

C. Femto LASIK

D. Implantable collamer lens

See Answer in Part B → Q3

@Neet_pg_bot

Q4: What is the statement correct about the instrument given below? (Q857755)

A. Condensing lens is not required

B. It is used as close to the patient’s eye as possible. A real, inverted image is seen, Magnification is about 15 times

C. A real, inverted image is seen

D. Magnification is about 15 times



See Answer in Part B → Q4

@Neet_pg_bot

Q5: Which of the following lenses would be recommended to correct myopia? (Q857245)

A. Convex

B. Concave

C. Cylindrical

D. Abraham lens

See Answer in Part B → Q5

@Neet_pg_bot

Q6: In a 3-year-old child who has recently developed 15-degree accommodative esotropia, what is the most effective treatment approach? (Q857242)

A. Orthoptic exercise

B. Surgery

C. Refractive correction

D. INTACS

See Answer in Part B → Q6

@Neet_pg_bot

Q7: The probable cause for a patient with a corneal scar having a visual acuity of 6/36, which improves to 6/18 with a pinhole, is most likely due to... (Q857209)

A. Glaucoma

B. Malingering

C. Irregular astigmatism

D. Cataract

See Answer in Part B → Q7

@Neet_pg_bot

Q8: During a routine eye examination, a patient's visual acuity was evaluated using Snellen's chart, and it was determined to be 6/6. What is the smallest angle at which this patient can differentiate two separate points or objects? (Q855228)

A. 15 minutes of arc

B. 5 minutes of arc

C. 10 minutes of arc

D. 20 minutes of arc

See Answer in Part B → Q8

@Neet_pg_bot

Q9: When selecting the strength of an intraocular lens (IOL) for a child under 1 year old, what factors should be considered? (Q793030)

A. Myopic shift

B. Hypermetropic shift

C. Length of the lens capsule

D. Growth of the lens capsule

See Answer in Part B → Q9

@Neet_pg_bot

Q10: What is the probable medical condition for a 70-year-old individual who has relied on presbyopia glasses but can now comfortably read the newspaper without them? (Q858148)

A. Index myopia

B. Index hypermetropia

C. Presbyopia

D. None of the above

See Answer in Part B → Q10

@Neet_pg_bot


Q11: What is the term used to describe the refractive error seen in the ocular examination image of a 60-year-old patient who has complained of reduced distant vision, but no longer requires spectacles for near vision? (Q855499)

A. Index myopia

B. Axial myopia

C. Curvatural myopia

D. Positional myopia



See Answer in Part B → Q11

@Neet_pg_bot

Q12: Which of the following refractive errors is associated with the image given below? (Q855491)

A. Presbyopia

B. Astigmatism

C. Hypermetropia

D. Myopia



See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 21 year old boxer wanted to improve the vision . Which of the following surgery is contraindicated? (Q854701)

A. RK

B. PRK

C. LASIK

D. SMILE

See Answer in Part B → Q13

@Neet_pg_bot


Q14: A patient presents with complaints of eye strain. The given image shows the focus of light rays in different meridians in the eye. Which refractive error is present in this patient? (Q866834)

A. Amblyopia

B. Hypermetropia

C. Astigmatism

D. Presbyopia



See Answer in Part B → Q14

@Neet_pg_bot


Q15: Which of the following do not represent the corrected refractive error? (Q860210)

A. Myopia

B. Hyperopia

C. Presbyopia

D. Astigmatism



See Answer in Part B → Q15

@Neet_pg_bot

Q16: What diagnosis is suggested by having different refractive powers (+5D in one eye and +1D in the other eye)? (Q860204)

A. Refractive errors

B. Amblyopia

C. Anisometropic amblyopia

D. Malingering

See Answer in Part B → Q16

@Neet_pg_bot

Q17: What is Anisometropia ? (Q840053)

A. Difference in visual acuity

B. Difference in refractive error between 2 eyes

C. Difference in the axial length of the eyeball

D. Difference in size of eyeball

See Answer in Part B → Q17

@Neet_pg_bot

Q18: During the Worth 4-Dot Test, a patient reports seeing only three green dots . What is the most likely diagnosis? (Q843991)

A. Right eye suppression

B. Left eye suppression

C. Crossed diplopia with esodeviation

D. Crossed diplopia with exodeviation

See Answer in Part B → Q18

@Neet_pg_bot

Q19: Which of the following is a cause of bilateral sudden painless loss of vision? (Q853148)

A. Acute congestive glaucoma

B. Diabetic retinopathy

C. Keratoconus D) Iridocyclitis

D. Iridocyclitis

See Answer in Part B → Q19

@Neet_pg_bot

Q20: What is the normal artery-to-vein ratio at the optic disc margin? (Q853134)

A. 1:3

B. 2:3

C. 2:5

D. 1:5

See Answer in Part B → Q20

@Neet_pg_bot

Q21: A 22-year-old patient with high myopia and thick glasses presents with floaters in his eyes. What is the likely cause of this condition? (Q853121)

A. Subarachnoid hemorrhage

B. Posterior Vitreous Detachment

C. Vitreous hemorrhage

D. Branched retinal artery occlusion

See Answer in Part B → Q21

@Neet_pg_bot


Q22: Binocular single vision consists of the following components: (Q866361)

A. a. SMP b. Fusion

B. a. Stereopsis b. Fusion

C. a. Fusion b. SMP

D. a. SMP b. Stereopsis



See Answer in Part B → Q22

@Neet_pg_bot

Q23: Which of the following is not an effective treatment for amblyopia? (Q865351)

A. Correction of refractive errors

B. Occlusion therapy (patching)

C. Strabismus surgery

D. Use of video games for visual stimulation

See Answer in Part B → Q23

@Neet_pg_bot

Q24: A posterior staphyloma was observed on indirect ophthalmoscopy. What is the likely diagnosis? (Q866854)

A. High myopia

B. Hypermetropia

C. Primary open-angle glaucoma

D. Retinitis pigmentosa

See Answer in Part B → Q24

@Neet_pg_bot

Q25: The following light reflex is seen in a patient. How much is the deviation? (Q866850)

A. 30 PD

B. 15°

C. 90 PD

D. 30°



See Answer in Part B → Q25

@Neet_pg_bot

Q26: A 45-year-old patient with eye examination findings of a deep anterior chamber and jet-black pupil is prescribed +12D glasses. Likely diagnosis? (Q866838)

A. Pseudophakia

B. Aphakia

C. Myopia

D. Hypermetropia

See Answer in Part B → Q26

@Neet_pg_bot

• [12_Basics of Ophthalmology] 026 Introduction To Ophthalmology

« Back to Folder Directory

[12_Basics of Ophthalmology] 026 Introduction To Ophthalmology — Quick Jump

Q1 (p. 385) *Q2 (p. 385)* *Q3 (p. 386)* *Q4 (p. 386)* *Q5 (p. 386)* *Q6 (p. 387)*
Q7 (p. 387) *Q8 (p. 388)* *Q9 (p. 388)* *Q10 (p. 389)*

Q1: A 25-year-old patient presents with redness and discomfort in the eyes. On examination, the physician notes inflammation confined to the conjunctiva attached to the posterior tarsal plate. Which part of the conjunctiva is primarily affected in this patient? (Q853247)

A. Bulbar conjunctiva

B. Forniceal conjunctiva

C. Palpebral conjunctiva

D. Limbal conjunctiva

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Which of the following structures do not contain any ganglion cells? (Q853248)

A. Macula

B. Fovea

C. Foveola

D. Periphery of the retina

See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which of the following is the anteroposterior diameter of the adult human eyeball? (Q853249)

A. 23 mm

B. 23.5 mm

C. 24 mm

D. 75 mm

See Answer in Part B → Q3

@Neet_pg_bot

Q4: Which structure primarily offers resistance to aqueous humor outflow within the iridocorneal angle? (Q853250)

A. Iris

B. Ciliary Body

C. Descemet's Membrane

D. Trabecular Meshwork

See Answer in Part B → Q4

@Neet_pg_bot

Q5: Which of the following statements accurately describes the role of retinal ganglion cells in the visual process? (Q853251)

A. They are the primary photoreceptors responsible for converting light into electrical signals.

B. They are concentrated in the fovea and are responsible for colour vision and high visual acuity.

C. Their axons form the optic nerve, which transmits visual information to the brain.

D. They contain pigments that undergo structural changes when exposed to light, triggering signal transduction.

See Answer in Part B → Q5

@Neet_pg_bot

Q6: What is the primary use of the instrument shown in the image given below? (Q853252)

A. To measure intraocular pressure.

B. To examine the anterior segment of the eye.

C. To provide a wide-field view of the retina.

D. To capture high-resolution images of the retina.



See Answer in Part B → Q6

@Neet_pg_bot

Q7: Which of the following arteries is not a branch of the ophthalmic artery? (Q853253)

A. Supratrochlear Artery

B. Dorsal Nasal Artery

C. Transverse Facial Artery

D. Short Posterior Ciliary Artery

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 32-year-old patient presents with an orbital infection characterized by severe pain, swelling, and redness around the eye. Despite treatment, the patient develops a headache, fever, and neurological deficits, leading to a diagnosis of a brain abscess. What is the most likely route of spread of the infection? (Q853254)

A. Ethmoidal veins to the cavernous sinus

B. Facial vein to the cavernous sinus

C. Superior ophthalmic vein to the cavernous sinus

D. Angular vein to the internal jugular vein

See Answer in Part B → Q8

@Neet_pg_bot

Q9: A patient presents to the ophthalmology OPD with the following features: Which of the following conditions is most likely responsible for these symptoms? (Q853255)

A. Abducens nerve palsy

B. Optic neuritis

C. Oculomotor nerve palsy

D. Trochlear nerve palsy



See Answer in Part B → Q9

@Neet_pg_bot

Q10: During an anatomy lecture, a student learns about the innervation of the eye. The pupil dilates in response to low light conditions. Which nerve fibers are primarily responsible for the dilation of the pupil? (Q853257)

A. Oculomotor nerve fibers

B. Trigeminal nerve fibers

C. Sympathetic nerve fibers

D. Facial nerve fibers

See Answer in Part B → Q10

@Neet_pg_bot

• [12_Basics of Ophthalmology] 027 Previous Year Questions

« Back to Folder Directory

[12_Basics of Ophthalmology] 027 Previous Year Questions — Quick Jump

Q1 (p. 390)	*Q2 (p. 390)*	*Q3 (p. 391)*	*Q4 (p. 391)*	*Q5 (p. 392)*	*Q6 (p. 392)*
Q7 (p. 392)	*Q8 (p. 393)*	*Q9 (p. 393)*	*Q10 (p. 393)*	*Q11 (p. 394)*	*Q12 (p. 395)*
Q13 (p. 395)	*Q14 (p. 395)*	*Q15 (p. 396)*	*Q16 (p. 396)*	*Q17 (p. 397)*	*Q18 (p. 397)*
Q19 (p. 398)	*Q20 (p. 398)*	*Q21 (p. 399)*	*Q22 (p. 399)*	*Q23 (p. 400)*	


Q1: Identify the instrument used for handling delicate tissues during eye surgery. (Q860570)

A. Mcpherson forceps

B. Lim's forceps

C. Artery forceps

D. SR holding forceps



See Answer in Part B → Q1

@Neet_pg_bot

Q2: Which laser is considered the most appropriate for cataract surgery? (Q859777)

A. Nd-YAG

B. Diode

C. Excimer

D. Femtosecond laser

See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which laser is most commonly used for performing a laser iridotomy? (Q859748)

A. Argon laser

B. Nd:YAG laser

C. Excimer laser

D. CO■ laser

See Answer in Part B → Q3

@Neet_pg_bot

Q4: What kind of laser can be utilized to treat posterior capsular opacification, which is observed during an examination of a patient who has reported decreased visual acuity following cataract surgery? (Q858163)

A. Femto laser

B. Nd YAG laser

C. Argon fluoride

D. Argon

See Answer in Part B → Q4

@Neet_pg_bot

Q5: Which of the following is not included in the SAFE strategy for trachoma? (Q856713)

A. Antibiotics

B. Surgery for trichiasis

C. Evaluation of control program

D. Facial cleanliness

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Which organism is most likely causing the infection in a 14-day-old neonate with bilateral conjunctivitis and lung infiltrates, as found in a chest X-ray performed at the outpatient department? (Q855386)

A. Streptococcus

B. Hemophilus influenzae

C. Chlamydia trachomatis

D. Mycoplasma

See Answer in Part B → Q6

@Neet_pg_bot

Q7: What type of visual impairment does a person have if their vision is less than 3/60 but greater than 1/60 in their better eye, and their field of vision is limited to less than 10 degrees from the center of fixation? (Q857712)

A. Curable blindness

B. Economic blindness

C. Social blindness

D. Manifest blindness

See Answer in Part B → Q7

@Neet_pg_bot

Q8: Which is the most common cause of preventable childhood blindness in India? (Q857231)

A. Refractive error

B. Vitamin A deficiency

C. Retinopathy of prematurity

D. Cataract

See Answer in Part B → Q8

@Neet_pg_bot

Q9: According to the National Blindness and Visual Impairment Survey conducted from 2015 to 2019, what is the leading cause of moderate to severe visual impairment? (Q855573)

A. Refractive error

B. Untreated cataract

C. Glaucoma

D. Diabetic retinopathy

See Answer in Part B → Q9

@Neet_pg_bot

Q10: How does the World Health Organization (WHO) define blindness? (Q860171)

A. VA < 3/60

B. VA < 6/60

C. VA > 3/60

D. VA > 6/60

See Answer in Part B → Q10

@Neet_pg_bot

Q11: Which of the following statements about the instrument shown in the image given below is false? {{caption_text}} (Q860010)

A. Used to measure the curvature of the cornea

B. Assess corneal astigmatism.

C. Depends on Purkinje image I.

D. Measures intraocular pressure.



See Answer in Part B → Q11

@Neet_pg_bot

Q12: What is considered the most reliable tonometer in the medical field? (Q860008)

A. Schiotz

B. Goldmann applanation tonometer

C. Tonopen

D. Pascal dynamic contour

See Answer in Part B → Q12

@Neet_pg_bot

Q13: All of the following are functions of the superior oblique muscle except? (Q853122)

A. Adduction

B. Abduction

C. Depression

D. Intorsion

See Answer in Part B → Q13

@Neet_pg_bot

Q14: Which of the following statements is correct regarding indirect ophthalmoscopy? (Q852502)

A. The image is erect and real.

B. The condenser lens needs to be kept close to the examiner’s eye

C. It provides 15x magnification.

D. It provides 3-5x magnification.

See Answer in Part B → Q14

@Neet_pg_bot


Q15: What does the picture depict in terms of the investigation? (Q859780)

A. Fundus fluorescein angiography

B. Optical coherence tomography

C. Retinal CT

D. Retinal MRI



See Answer in Part B → Q15

@Neet_pg_bot

Q16: A 6-year-old child presents with a refractive error of -2D in the right eye and +1D in the left eye. Fundus examination reveals normal visual acuity and retinal findings. What is the most likely diagnosis? (Q852481)

A. Myopia

B. Hyperopia

C. Anisometropia

D. Amblyopia

See Answer in Part B → Q16

@Neet_pg_bot

Q17: A patient presents with right-sided field defects in both eyes, but central vision remains unaffected. What is the most likely diagnosis? (Q852486)

A. Homonymous Hemianopia with Macular Sparing

B. Heteronymous Hemianopia with Central Sparing

C. Optic Chiasm Lesion

D. Optic Tract Lesion

See Answer in Part B → Q17

@Neet_pg_bot

Q18: The person in the image is at the risk of developing? (Q856126)

A. Exposure keratopathy

B. Restricted eye movements

C. Diplopia

D. Amblyopia



See Answer in Part B → Q18

@Neet_pg_bot

Q19: Which of the subsequent options aids in distinguishing a fixed dilated pupil? Phenylephrine 1% Pilocarpine 0.1% Pilocarpine Cocaine 0.5% Epinephrine (Q856408)

A. 1,2,3,4 and 5

B. 2 and 3

C. 1,4and 5

D. 1,3 and 4

See Answer in Part B → Q19

@Neet_pg_bot

Q20: Which of the following combinations of cylindrical lenses can be employed for the treatment of astigmatism in the with-the-rule orientation? (Q793355)

A. +5.00 DC x 180, -3.00 DC x 90

B. +7.00 DC x 180, +5.00 DC x 90

C. -4.00 DC x 180, -5.00 DC x 90

D. -3.00 DC x 180, +1.00 DC x 90

See Answer in Part B → Q20

@Neet_pg_bot

Q21: Which of the following is true about viscoelastic substances? (Q853141)

A. Cause mechanical injury to corneal endothelium B) Prevents Anterior Chamber from collapsing during crucial steps like capsulorrhexis and IOL placement C) It is inert and can be left behind after surgery D) Silicone oil, perfluoropropane, and sodium hexafluoride are most commonly used substances

B. Prevents Anterior Chamber from collapsing during crucial steps like capsulorrhexis and IOL placement

C. It is inert and can be left behind after surgery

D. Silicone oil, perfluoropropane, and sodium hexafluoride are most commonly used substances

See Answer in Part B → Q21

@Neet_pg_bot

Q22: In Fundus Fluorescein Angiography (FFA), how do microaneurysms appear? (Q853133)

A. Hypofluorescent dots

B. Hyperfluorescent dots

C. Hypofluorescent line

D. Hyperfluorescent branching-like structure in the vessel

See Answer in Part B → Q22

@Neet_pg_bot

Q23: All of the following are features of oculomotor nerve palsy except? (Q853124)

A. Exotropia

B. Dilated pupil

C. Limited abduction

D. Ptosis

See Answer in Part B → Q23

@Neet_pg_bot

• [13_Sclera] 028 Sclera

« Back to Folder Directory

[13_Sclera] 028 Sclera — Quick Jump

Q1 (p. 401) *Q2 (p. 401)* *Q3 (p. 402)* *Q4 (p. 403)* *Q5 (p. 403)* *Q6 (p. 404)*
Q7 (p. 404) *Q8 (p. 405)* *Q9 (p. 406)* *Q10 (p. 406)* *Q11 (p. 406)* *Q12 (p. 407)*
Q13 (p. 407) *Q14 (p. 408)* *Q15 (p. 408)*

Q1: Which of the following correctly lists the layers of the sclera from outermost to innermost? (Q853641)

A. Lamina fusca > Sclera proper > Episclera

B. Episclera > Sclera proper > Lamina fusca

C. Sclera proper > Episclera > Lamina fusca

D. Episclera > Lamina fusca > Sclera proper

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Match the scleral apertures with the structures they transmit Scleral Apertures Structures Transmitted A. Anterior scleral apertures 1. Short and long ciliary arteries B. Middle scleral apertures 2. Central retinal artery and vein C. Posterior scleral apertures 3. Anterior ciliary arteries and veins D. Lamina cribrosa 4. Vortex veins (Q853642)

A. A-3, B-4, C-1, D-2

B. A-2, B-1, C-4, D-3

C. A-3, B-2, C-1, D-4

D. A-3, B-1, C-4, D-2

See Answer in Part B → Q2

@Neet_pg_bot


Q3: A 28-year-old patient presents with painless redness of the eye. The ophthalmologist performed a diagnostic test using 2.5% phenylephrine, and no change was noticed. When 10% phenylephrine is used, the following change occurs. What is the most likely diagnosis? Before application of 10% phenylephrine: After application of 10% phenylephrine: (Q853643)


A. Conjunctivitis

B. Scleritis

C. Episcleritis

D. Uveitis





See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 40-year-old patient presents with recurrent episodes of episcleritis. After taking a detailed history and performing a thorough systemic examination, which of the following investigations would you consider ordering? (Q853644)

A. Full blood count and rheumatoid factor (RF)

B. Mantoux test and antinuclear antibody (ANA)

C. VDRL and X-rays of the chest and sacroiliac joints

D. All of the above

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 45-year-old woman with a 15-year history of rheumatoid arthritis (RA) presents with severe ocular pain, redness, and progressively worsening vision in her left eye for the past 2 weeks. Examination reveals a localized area of scleral thinning

with adjacent congestion. There is no posterior involvement on B-scan ultrasound. Which of the following is the most likely diagnosis? (Q853645)

A. Anterior staphyloma

B. Necrotizing scleritis with inflammation

C. Nodular scleritis

D. Scleromalacia perforans

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 38-year-old male presents with persistent eye discomfort, redness, and watering. On examination, the left eye is diffusely tender, sclera appears uniformly reddened and swollen without localized thinning areas. There are no systemic signs of vasculitis or autoimmune disease on further investigation. Which type of anterior scleritis is most consistent with this presentation? (Q853646)

A. Necrotizing anterior scleritis with inflammation

B. Non-necrotizing anterior diffuse scleritis

C. Non-necrotizing anterior nodular scleritis

D. Scleromalacia perforans

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 45-year-old woman presents with severe pain in her right eye, worsened by eye movements, and reduced vision. On examination, proptosis and restricted ocular movements were observed. Ophthalmoscopy shows disc, macular edema, and choroidal folds. Which of the following B-scan ultrasonography findings is most indicative of this disease? (Q853647)

A. T sign

B. Moth-Eaten Appearance

C. Sunray Sign

D. Kiss Sign

See Answer in Part B → Q7

@Neet_pg_bot


Q8: A 60-year-old woman with a 20-year history of rheumatoid arthritis has a painless, progressively thinning area in the sclera of her left eye. The area appears as shown in the image below. Which of the following best describes the pathophysiological mechanism responsible for this condition? (Q853648)

A. Type III hypersensitivity

B. Type II hypersensitivity

C. Type IV hypersensitivity

D. Infective etiology



See Answer in Part B → Q8

@Neet_pg_bot

Q9: Depending on the location of the staphyloma in the eye which among the following options is matched correctly with its cause? (Q853649)

A. Anterior staphyloma - Peripheral corneal ulcer

B. Intercalary staphyloma - Degenerative myopia

C. Equatorial staphyloma - Scleritis

D. Ciliary staphyloma - Pseudocornea

See Answer in Part B → Q9

@Neet_pg_bot

Q10: Blue sclera is commonly associated with all of the following conditions except: (Q853650)

A. Osteogenesis Imperfecta

B. Ehlers-Danlos Syndrome

C. Alkaptonuria

D. Marfan Syndrome

See Answer in Part B → Q10

@Neet_pg_bot

Q11: Clinical question: A 60 y/o man from a tropical region presents with a painful, red eye and a corneal ulcer following recent exposure to decaying vegetable matter. Which is the most common fungal infection affecting the cornea? (Q853664)

A. Candida albicans

B. Fusarium

C. Mucormycosis

D. Aspergillus fumigatus

See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 45 y/o man from a tropical region presents with a painful red eye, significant watering, and sensitivity to light after working in a garden with decaying plant material. The patient is diagnosed with keratomycosis. Which of the following is not an examination finding in keratomycosis? (Q853665)

A. Feathery extensions at the margins of the corneal ulcer

B. Extensive corneal vascularization

C. Sterile immune ring

D. Endothelial plaque

See Answer in Part B → Q12

@Neet_pg_bot

Q13: A 45 y/o male presents with redness, pain, and decreased vision in his right eye. He reports a history of trauma to the eye with a plant material three days ago. O/E, there is a corneal ulcer with feathery edges and satellite lesions. Hypopyon is also noted. Which of the following is the most appropriate initial treatment for this patient? (Q853666)

A. Topical Moxifloxacin

B. Topical Natamycin

C. Oral Acyclovir

D. Intravitreal Amphotericin B

See Answer in Part B → Q13

@Neet_pg_bot

Q14: A 45 y/o patient presents with eye redness, excessive tearing, and sensitivity to light. O/E you observe dendritic ulcers, and decreased corneal sensation. Based on these findings, which virus is most likely responsible for the patient's condition? (Q853667)

A. Adenovirus

B. Varicella Zoster Virus (VZV)

C. Herpes Simplex Virus (HSV)

D. Chlamydia trachomatis

See Answer in Part B → Q14

@Neet_pg_bot

Q15: A 35 y/o patient presents with pain, redness, photophobia, and decreased vision in the right eye. Slit-lamp examination reveals a dendritic ulcer on the cornea. Which of the following is the most appropriate initial treatment for this patient? (Q853668)

A. Topical steroids and antibiotics

B. Oral acyclovir and lubricants

C. Topical acyclovir and cycloplegics

D. Oral steroids and topical antibiotics

See Answer in Part B → Q15

@Neet_pg_bot

• [13_Sclera] 029 Previous Year Questions

« Back to Folder Directory

[13_Sclera] 029 Previous Year Questions — Quick Jump

Q1 (p. 409) *Q2 (p. 409)* *Q3 (p. 410)* *Q4 (p. 412)* *Q5 (p. 412)* *Q6 (p. 413)*

Q1: A patient with long-standing, poorly controlled rheumatoid arthritis presents with a sudden, painless blurring of vision in the superior visual field of one eye. Fundoscopic examination shows retinal hemorrhages and dilated, tortuous veins in the inferior retina. What is the most likely diagnosis? (Q853116)

A. Branch Retinal Artery Occlusion

B. Central Retinal Artery Occlusion

C. Branch Retinal Vein Occlusion

D. Central Retinal Vein Occlusion

See Answer in Part B → Q1

@Neet_pg_bot


Q2: A patient presents with a nodular swelling near the limbus, which blanches after administering phenylephrine and then recurs. What is the most probable diagnosis? (Q844042)

A. Episcleritis

B. Scleritis

C. Bitot spots

D. Pterygium



See Answer in Part B → Q2

@Neet_pg_bot


Q3: What is the probable diagnosis based on the observed blanching of vascularity following the administration of 10 % phenylephrine, as depicted in the provided image? (Q854717)


A. Nodular episcleritis

B. Nodular scleritis

C. Nodular phlycten

D. Scleral ectasia





See Answer in Part B → Q3

Q4: A 5-year-old boy presents with a small nodule on the sclera as shown in the image below. What is the likely diagnosis? (Q855476)

A. Pterygium

B. Pinguecula

C. Dermolipoma

D. Dermoid cyst



See Answer in Part B → Q4

@Neet_pg_bot


Q5: What is the probable diagnosis of a 35-year-old male with a history of prolonged corneal ulcer who now presents with a deformed eye as shown in the image? (Q857704)

A. Megalocornea

B. Conjunctivitis

C. Buphthalmos

D. Anterior staphyloma



See Answer in Part B → Q5

@Neet_pg_bot


Q6: In the image, what are the associated complications observed in a 35-year-old woman who has been diagnosed with rheumatoid arthritis? (Q856012)

A. Scleromalacia perforans

B. Ciliary staphyloma

C. Coloboma

D. Malignant melanoma



See Answer in Part B → Q6

@Neet_pg_bot

• [14_Lacrimal Drainage System] 030 Lacrimal Drainage System

« Back to Folder Directory

[14_Lacrimal Drainage System] 030 Lacrimal Drainage System — Quick Jump

Q1 (p. 415) *Q2 (p. 415)* *Q3 (p. 415)* *Q4 (p. 416)* *Q5 (p. 416)* *Q6 (p. 417)*
Q7 (p. 417) *Q8 (p. 417)* *Q9 (p. 418)* *Q10 (p. 418)*

Q1: Which structure within the lacrimal passages prevents the backflow of tears from the nasal cavity into the nasolacrimal duct? (Q853822)

A. Valve of Hasner

B. The valve of Rosenmüller

C. Orbital lobe

D. Glands of Krause

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Which of the following is a cause of lacrimal pump failure, leading to excessive tearing (epiphora)? (Q853823)

A. Trauma to the canaliculus

B. Bell’s palsy

C. Obstruction of the nasolacrimal duct

D. Narrowing of the punctum

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 65 y/o male presents with persistent tearing in his right eye that has worsened over the past year. The physician orders an imaging study to pinpoint the exact location of the blockage and assess the condition of the lacrimal sac. Which investigation is most appropriate for this purpose? (Q853824)

A. Dacryocystography

B. Fluorescein Dye Disappearance Test (FDDT)

C. Dacryoscintigraphy

D. Lacrimal Syringing

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 50 y/o man undergoes Jones Test 1 for chronic tearing, which shows no dye on the cotton bud placed in the inferior meatus. Following this, Jones Test 2 is performed, and fluorescein dye is detected after lacrimal syringing. What does this result suggest? (Q853825)

A. Partial obstruction within the lacrimal passages

B. Complete obstruction of the nasolacrimal duct

C. Lacrimal pump failure

D. Primary hypersecretion of tears

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 9-month-old infant is brought to the clinic with ongoing tearing and discharge from the right eye. Despite regular lacrimal sac massage and topical antibiotics, the symptoms persist. What is the next best step in management? (Q853826)

A. Continue lacrimal sac massage and antibiotics

B. Probing of the nasolacrimal duct

C. Dacryocystorhinostomy (DCR)

D. Balloon catheter dilation

See Answer in Part B → Q5

@Neet_pg_bot

Q6: Which of the following types of Dacryocystorhinostomy (DCR) is considered for treating nasolacrimal duct obstruction, offering the highest success rate? (Q853827)

A. External Dacryocystorhinostomy (External DCR)

B. Endonasal Dacryocystorhinostomy (Endonasal DCR)

C. Endocanalicular Laser Dacryocystorhinostomy (Laser DCR)

D. Conjunctivodacryocystorhinostomy (CDCR)

See Answer in Part B → Q6

@Neet_pg_bot

Q7: A 2-week-old infant presents with a bluish swelling in the inner corner of the right eye. The parents report excessive tearing and some slimy discharge from the eye. There is no history of trauma or illness. What is the most likely diagnosis? (Q853828)

A. Congenital Dacryocele

B. Dacryoadenitis

C. Dacryops

D. Chronic Dacryocystitis

See Answer in Part B → Q7

@Neet_pg_bot

Q8: A 55 y/o woman presents with a persistent swelling below the inner corner of her left eye. She also reports a history of watery eyes and recurrent conjunctivitis. On examination, there is fluctuant swelling at the lacrimal sac, and mucoid fluid can be expressed. What is the most likely diagnosis and appropriate management? (Q853829)

A. Chronic Dacryocystitis; Probing and syringing.

B. Chronic Dacryocystitis; Dacryocystorhinostomy (DCR).

C. Acute Dacryocystitis; Systemic antibiotics and abscess drainage.

D. Acute Dacryocystitis; Dacryocystectomy (DCT).

See Answer in Part B → Q8

@Neet_pg_bot

Q9: A 60 y/o man presents with watery eyes, redness, swelling at the inner corner of his right eye, and persistent thick yellow discharge. O/E there is pouting of lacrimal puncta and yellow granules from the punctum upon massage. Which organism is most likely responsible, and what is the definitive treatment? (Q853830)

A. Eikenella corrodens; Oral antibiotics and steroid drops.

B. Staphylococcus aureus; Topical antibiotics and warm compresses

C. Actinomyces israelii; Canaliculotomy

D. Chlamydia trachomatis; Irrigation and syringing of the tear duct.

See Answer in Part B → Q9

@Neet_pg_bot

Q10: A 27 yeal old man presents with a painless, slow-growing, firm mass on the upper eyelid that has been gradually increasing in size over the past year. On examination, the lesion is well-circumscribed, mobile, and non-tender, with no signs of ulceration or inflammation. Imaging reveals a well-defined, lobulated mass without bony invasion. What is the most likely diagnosis? (Q853831)

A. Pleomorphic Adenoma

B. Lymphoma

C. Lacrimal Gland Carcinoma

D. Dacryocystitis

See Answer in Part B → Q10

@Neet_pg_bot

• [14_Lacrimal Drainage System] 031 Previous Year Questions

« Back to Folder Directory

[14_Lacrimal Drainage System] 031 Previous Year Questions — Quick Jump

Q1 (p. 420) *Q2 (p. 420)* *Q3 (p. 420)* *Q4 (p. 421)* *Q5 (p. 422)* *Q6 (p. 422)*
Q7 (p. 423) *Q8 (p. 423)* *Q9 (p. 424)* *Q10 (p. 424)* *Q11 (p. 425)* *Q12 (p. 425)*

Q1: A 3-year-old child presents with epiphora, and probing reveals a soft block at 7 mm. What is the best treatment option for this condition? (Q852499)

A. Endonasal Dacryocystorhinostomy (DCR)

B. Conjunctival Dacryocystorhinostomy (DCR)

C. External Dacryocystorhinostomy (DCR)

D. Dacryocystectomy

See Answer in Part B → Q1

@Neet_pg_bot

Q2: Which bones are cut/removed during external Dacryocystorhinostomy? (Q860569)

A. Zygomatic

B. Maxillary

C. Lacrimal

D. Frontal

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 3-month-old child is brought to the outpatient department with watering of eyes and photophobia . On examination, both corneas appear hazy. Which of the following is the most likely diagnosis ? (Q840075)

A. Congenital nasolacrimal duct obstruction

B. Congenital glaucoma

C. Congenital entropion

D. Congenital distichiasis

See Answer in Part B → Q3

@Neet_pg_bot


Q4: Identify the marked structure: (Q857418)

A. Canal of petit

B. Canal of hannover

C. Hyaloid canal

D. Pars plana



See Answer in Part B → Q4

@Neet_pg_bot

Q5: What is the accurate order of staining in patients with dry eyes? i. Fluorescein stain ii. Lissamine green iii. Rose Bengal stain (Q857419)

A. i, iii, ii

B. i, ii, iii

C. ii, i, ii

D. iii, ii, i

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 60-year-old patient presents with chronic dacryocystitis. She complains of recurrent tearing, discharge, and mild swelling in the inner corner of the eye. The clinical findings and history are consistent with chronic dacryocystitis. What should be the treatment of choice for this patient? (Q858129)

A. Lacrimal sac massage

B. Dacryocystorhinostomy

C. Syringing

D. Probing

See Answer in Part B → Q6

@Neet_pg_bot

Q7: What is the most appropriate treatment for a 55-year-old woman who is experiencing long-term mucopurulent discharge from her left eye and swelling near the inner corner of the eye? (Q857191)

A. Dacryocystorhinostomy

B. Dacryocystectomy

C. Incision and drainage

D. Balloon catheter dilation

See Answer in Part B → Q7

@Neet_pg_bot

Q8: Which of the following surgeries use this instrument? (Q857195)

A. Evisceration

B. Enucleation

C. Cataract surgery

D. Dacryocystorhinostomy



See Answer in Part B → Q8

@Neet_pg_bot

Q9: Which of the following treatments is appropriate for a 3-month-old child with a mucopurulent discharge from the right puncta and a history of increased watering in the right eye, as noticed by the mother? (Q857689)

A. Syringing

B. Probing with syringing

C. Sac massage with topical antibiotics

D. Dacryocystorhinostomy

See Answer in Part B → Q9

@Neet_pg_bot

Q10: A 55-year-old woman is being counseled for surgery for her long-standing dacryocystitis. Which of the following is the surgery of choice? (Q859321)

A. Dacryocystorhinostomy

B. Dacryocystectomy

C. Probing

D. Balloon catheter dilation

See Answer in Part B → Q10

@Neet_pg_bot

Q11: In Dacryocystorhinostomy (DCR), the lacrimal sac is opened into which nasal meatus? (Q859753)

A. Inferior meatus

B. Middle meatus

C. Supreme meatus

D. Superior meatus

See Answer in Part B → Q11

@Neet_pg_bot

Q12: A 60-year-old patient presents with chronic dacryocystitis . She complains of recurrent tearing, discharge, and mild swelling in the inner corner of the eye . The clinical findings and history are consistent with chronic dacryocystitis. What should be the treatment of choice for this patient ? (Q820463)

A. Lacrimal sac massage

B. Dacryocystorhinostomy

C. Syringing

D. Probing

See Answer in Part B → Q12

@Neet_pg_bot

• [15_Community Ophthalmology] 032 Previous Year Questions

« Back to Folder Directory

[15_Community Ophthalmology] 032 Previous Year Questions — Quick Jump

Q1 (p. 427) *Q2 (p. 427)* *Q3 (p. 427)* *Q4 (p. 428)* *Q5 (p. 428)* *Q6 (p. 429)*

Q1: Which organism is most likely causing the infection in a 14-day-old neonate with bilateral conjunctivitis and lung infiltrates, as found in a chest X-ray performed at the outpatient department? (Q855386)

A. Streptococcus

B. Hemophilus influenzae

C. Chlamydia trachomatis

D. Mycoplasma

See Answer in Part B → Q1

@Neet_pg_bot

Q2: According to the National Blindness and Visual Impairment Survey conducted from 2015 to 2019, what is the leading cause of moderate to severe visual impairment? (Q855573)

A. Refractive error

B. Untreated cataract

C. Glaucoma

D. Diabetic retinopathy

See Answer in Part B → Q2

@Neet_pg_bot

Q3: Which of the following is not included in the SAFE strategy for trachoma? (Q856713)

A. Antibiotics

B. Surgery for trichiasis

C. Evaluation of control program

D. Facial cleanliness

See Answer in Part B → Q3

@Neet_pg_bot

Q4: Which is the most common cause of preventable childhood blindness in India? (Q857231)

A. Refractive error

B. Vitamin A deficiency

C. Retinopathy of prematurity

D. Cataract

See Answer in Part B → Q4

@Neet_pg_bot

Q5: What type of visual impairment does a person have if their vision is less than 3/60 but greater than 1/60 in their better eye, and their field of vision is limited to less than 10 degrees from the center of fixation? (Q857712)

A. Curable blindness

B. Economic blindness

C. Social blindness

D. Manifest blindness

See Answer in Part B → Q5

@Neet_pg_bot

Q6: How does the World Health Organization (WHO) define blindness? (Q860171)

A. VA < 3/60

B. VA < 6/60

C. VA > 3/60

D. VA > 6/60

See Answer in Part B → Q6

@Neet_pg_bot

• [16_Ocular Injuries] 033 Ocular Injuries

« Back to Folder Directory

[16_Ocular Injuries] 033 Ocular Injuries — Quick Jump

Q1 (p. 430) *Q2 (p. 432)* *Q3 (p. 432)* *Q4 (p. 433)* *Q5 (p. 433)* *Q6 (p. 434)*


Q1: Which of the following types of ocular injuries can result from blunt trauma to the eye? 1) 2) 3) 4) (Q853600)

A. 2 and 4


B. 1 and 2


C. 3 and 1

D. All of the Above




@Neet_pg_bot


@Neet_pg_bot



See Answer in Part B → Q1

@Neet_pg_bot

Q2: A patient presents with a full-thickness corneal wound caused by a pencil. Which of the following is the most likely complication if the injury is left untreated? (Q853601)

A. Ring abscess of the cornea

B. Retinal detachment

C. Glaucoma

D. Strabismus

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A 40-year-old construction worker presents with eye pain and blurred vision after metal struck his eye. CT scan reveals an intraocular foreign body lodged in the vitreous cavity. What is the most appropriate treatment? (Q853602)

A. Observation and follow-up in one week

B. Corticosteroid injection without IOFB removal

C. Immediate surgery with pars plana vitrectomy and IOFB removal

D. Application of topical antibiotics and patching

See Answer in Part B → Q3

@Neet_pg_bot

Q4: A 28-year-old metalworker presents with gradual vision loss and a history of a metallic foreign body injury 4 months ago. Examination reveals iron deposits on Iris, rusty deposits on the lens and pigmentary changes in the retina. What is the most likely cause of his condition? (Q853603)

A. Chalcosis

B. Retinitis pigmentosa

C. Siderosis bulbi

D. Glaucoma

See Answer in Part B → Q4

@Neet_pg_bot

Q5: A 35-year-old male presents to the emergency department after an accidental chemical exposure to his eyes while working in a lab. He reports immediate burning and decreased vision. Upon examination, there is corneal epithelial loss and conjunctival swelling. What is the most appropriate immediate management step? (Q853604)

A. Necrotic tissue removal and administer topical antibiotics

B. Apply corticosteroids to reduce inflammation

C. Immediate irrigation with water or saline for 20-30 minutes

D. Perform corneal transplant

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A 28-year-old male presents to the emergency department 10 days after sustaining blunt trauma to the face during a basketball game. He complains of diplopia and difficulty moving his left eye upwards. On examination, there is significant periorbital swelling. The X-ray is shown below. What does this X-ray indicate in this patient, and what is the most appropriate next step in management? (Q853605)

A. Orbital fat herniation; apply ice and prescribe analgesics

B. Orbital floor fracture; surgical evaluation

C. Orbital haemorrhage; immediate CT scan

D. Maxillary sinus fracture; follow-up with ENT specialist



See Answer in Part B → Q6

@Neet_pg_bot

• [16_Ocular Injuries] 034 Previous Year Questions

« Back to Folder Directory

[16_Ocular Injuries] 034 Previous Year Questions — Quick Jump

Q1 (p. 436) *Q2 (p. 436)* *Q3 (p. 436)* *Q4 (p. 437)* *Q5 (p. 437)* *Q6 (p. 438)*
Q7 (p. 438) *Q8 (p. 439)* *Q9 (p. 439)* *Q10 (p. 440)*

Q1: A patient presents with a foreign body (chuna) in the eye and complaints of irritability and lacrimation. What is the first step in management for this patient? (Q853118)

A. Antibiotic eye drops B) Ascorbic eye drops C) Irrigation of the eye with normal saline D) Steroids

B. Ascorbic eye drops

C. Irrigation of the eye with normal saline

D. Steroids

See Answer in Part B → Q1

@Neet_pg_bot

Q2: A factory worker presented with a history of chuna (lime) exposure in both eyes a few hours ago. What is the first immediate management? (Q853143)

A. Take history B) Copious irrigation of eyes C) Check vision in both eyes D) Check IOP and plan for surgery

B. Copious irrigation of eyes

C. Check vision in both eyes

D. Check IOP and plan for surgery

See Answer in Part B → Q2

@Neet_pg_bot

Q3: A patient has a history of a flying foreign body injury to the eye. Hemochromatosis is subsequently diagnosed. Which of the following substances is most likely involved in the deposition? (Q852494)

A. Iron

B. Aluminum

C. Wood

D. Glass

See Answer in Part B → Q3

@Neet_pg_bot

Q4: The patient presents with foreign body (chuna) in his eyes and complaints of irritability and lacrimation. What is the first step management in this patient? (Q840064)

A. Antibiotics eye drops

B. Ascorbic eye drops

C. Irrigation of eye with normal saline

D. Steroids

See Answer in Part B → Q4

@Neet_pg_bot

Q5: Which investigation is most suitable for localizing a foreign body in the eye of an industrial worker admitted to the hospital due to an eye injury caused by cutting stone with a hammer and iron chisel? (Q857237)

A. B mode scan

B. CT

C. X-ray

D. MRI

See Answer in Part B → Q5

@Neet_pg_bot

Q6: A patient was brought to the outpatient department with a history of trauma near the eye and now presents with symptoms of fever, severe pain, and swelling in the left eye. On examination, no abnormal findings were observed. What is the likely diagnosis? (Q857714)

A. Orbital cellulitis

B. Endophthalmitis

C. Panophthalmitis

D. Preseptal cellulitis



See Answer in Part B → Q6

@Neet_pg_bot

Q7: A man suffered trauma to the eye from a punch while fighting with another person. What is the most common site of fracture in orbit? (Q857721)

A. Medial wall

B. Lateral wall

C. Roof

D. Floor

See Answer in Part B → Q7

@Neet_pg_bot


Q8: Can you determine the observation indicated by the arrow in the image provided? (Q856717)

A. Vossius ring

B. Weiss ring

C. Fleischer’s ring

D. Kayser Fleischer ring



See Answer in Part B → Q8

@Neet_pg_bot

Q9: An industrial worker was admitted to the hospital after an eye injury sustained while cutting stone with a hammer and iron chisel. A foreign body was suspected to be impacted in his eye. Which investigation can be detrimental to this patient? (Q856179)

A. B mode scan

B. CT

C. X-ray

D. MRI

See Answer in Part B → Q9

@Neet_pg_bot

Q10: What is the probable cause of bleeding in a patient who experienced a hyphema after being struck in the eye by a tennis ball? (Q859357)

A. Circulus arteriosus major

B. Anterior ciliary artery

C. Posterior ciliary artery

D. Circulus arteriosus minor

See Answer in Part B → Q10

@Neet_pg_bot

Part B — All Answers & Explanations

• [01_Orbit] 001 Orbit - I

« Back to Folder Directory

[01_Orbit] 001 Orbit - I — Quick Jump

Q1 (p. 441)	*Q2 (p. 442)*	*Q3 (p. 443)*	*Q4 (p. 444)*	*Q5 (p. 445)*	*Q6 (p. 446)*
Q7 (p. 447)	*Q8 (p. 449)*	*Q9 (p. 450)*	*Q10 (p. 450)*	*Q11 (p. 453)*	*Q12 (p. 454)*
Q13 (p. 455)	*Q14 (p. 456)*	*Q15 (p. 457)*	*Q16 (p. 459)*	*Q17 (p. 460)*	*Q18 (p. 461)*
Q19 (p. 462)	*Q20 (p. 463)*	*Q21 (p. 464)*	*Q22 (p. 465)*	*Q23 (p. 466)*	*Q24 (p. 467)*
Q25 (p. 468)	*Q26 (p. 469)*	*Q27 (p. 470)*			

Q1: A 9-year-old boy presents with a 2-day history of swelling, redness, and pain around his left eye, worsening with eye movements. He had a recent upper respiratory infection. On examination, there is proptosis, restricted eye movement, and chemosis. Vision is intact. CT scan confirms orbital cellulitis. Which staging system is used to classify this condition? (Q853270)

A. Chandler's classification


B. Van Herick classification

C. ETDRS classification

D. Shaffer grading

Correct Answer: A. Chandler's classification

Explanation: The staging system used to classify orbital cellulitis is Modified Chandler staging Modified Chandler staging: Stage I: Preseptal cellulitis/ periorbital cellulitis (limited anterior to orbital septum) Stage II: Inflammatory orbital oedema/ Orbital cellulitis (Posterior to orbital septum) Stage III: Subperiosteal abscess Stage IV: Orbital abscess Stage V: Cavernous sinus thrombosis Van Herick classification (Option B) is used to describe the anterior chamber angle status in glaucoma, using ratio of anterior chamber depth to corneal thickness. ETDRS classification (Option C) is the Early Treatment Diabetic Retinopathy Study used in Diabetic retinopathy staging Shaffer grading (Option D) is used to describe anterior chamber angle status in glaucoma, categorizing the risk of angle closure.



@Neet_pg_bot

References:

Parson’s Diseases of the eye, 23rd edition, Page 137.

Kanski’s Clinical ophthalmology, 9th edition, Page 125.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 45-year-old patient presents with a large, painless orbital mass that has been causing progressive proptosis and visual disturbance. After imaging and biopsy, the mass is identified as malignant. What is the choice of treatment in this scenario, and what is its definition? (Q853283)

A. Evisceration, it involves removal of the contents of the eyeball, sclera preserved

B. Enucleation, it involves removal of the contents of the eyeball, sclera preserved

C. Enucleation, it involves removal of the entire eyeball including the sclera.

D. Exenteration, it involves the removal of all orbital contents

Correct Answer: D. Exenteration, it involves the removal of all orbital contents

Explanation: The patient has a malignant orbital tumor with progressive proptosis and visual disturbance. In such cases, the treatment of choice is orbital exenteration , as globe-sparing procedures are insufficient. Surgical Procedures for Orbital Masses: Procedure Extent of Removal Indications Cosmetic and Functional Outcomes Exenteration All orbital contents (Eyeball + EOM + eyelids) Malignant tumors, extensive disease Significant cosmetic and functional deficits Evisceration (Option A) Contents of the eyeball, sclera preserved Not typically used for malignancy Preserves some eye movement Enucleation (Option B, C) Entire eyeball including sclera Malignant tumours, less radical than exenteration Allows for ocular prosthesis placement

References:

Kanski's Clinical Ophthalmology, 10th Edition, Page no, 152.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 12-year-old patient presents to the OPD with a typical S-shaped deformity of the eyelid. On examination, the patient is found to have multiple café-au-lait spots and axillary freckling. Which of the following findings is most commonly observed in this condition? (Q853282)

A. Early-onset cataracts


B. Lisch nodules

C. B/L Acoustic neuroma

D. Optic nerve sheath meningiomas

Correct Answer: B. Lisch nodules

Explanation: This 12-year-old with an S-shaped eyelid deformity (from plexiform neurofibroma) and café-au-lait spots with axillary freckling is a classic case of Neurofibromatosis type 1 (NF1) . The most common ocular finding in NF1 is the presence of Lisch nodules (iris hamartomas) . Ocular Manifestations in Neurofibromatosis Neurofibromatosis Type 1 (NF1) Lisch Nodules: Iris hamartomas, key diagnostic feature. Glaucoma: Common, especially with orbital-facial involvement; may lead to asymmetric globe enlargement. Optic Nerve Glioma: Can cause strabismus and proptosis. Palpebral Neurofibroma: S-shaped deformity of the lid Choroidal Alterations Neurofibromatosis Type 2 (NF2) Early-Onset Cataracts : Present in up to 80% of cases. (Option A) Optic Nerve Hamartomas : Found in some cases. Optic nerve sheath meningiomas (Option D) Combined Pigment Epithelial and Retinal Hamartomas : Also observed. B/L Acoustic neuroma (Option C)



References:

www.ncbi.nlm.nih.gov/pmc/articles/PMC4874640/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 6-year-old child presents with gradual, painless unilateral proptosis and vision loss. Fundoscopic examination reveals optic atrophy. Imaging studies show fusiform growth along the optic nerve. What is the most appropriate management for this child, considering the tumor is stationary and the vision is still present? (Q853281)

A. Immediate surgical excision of the tumor

B. Radiotherapy to manage the tumor

C. Observation and regular monitoring

D. Chemotherapy as the first-line treatment

Correct Answer: C. Observation and regular monitoring

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This child has an optic nerve glioma , suggested by the gradual painless unilateral proptosis, optic atrophy, and fusiform enlargement of the optic nerve on imaging; since the tumor is stationary and useful vision is still preserved, the most appropriate management is conservative observation with regular follow-up rather than surgery or radiotherapy. Optic Nerve Gliomas A slow-growing, tumor of astrocytes It is generally a unilateral tumor that typically affects children (more common in females). More common in patients with neurofibromatosis type 1 (NF1). Treatment: Observation to preserve the nerve as long as possible. Radiotherapy: For inoperable cases. (Option B) Chemotherapy : Used in specific cases when other treatments are not suitable. (Option D) Surgery (cosmetically unacceptable proptosis in a blind eye). (Option A)

References:

Parson’s Diseases of The Eye, 23rd Edition, Page no, 1155-1158.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 52-year-old woman presents with progressive, painless proptosis of the left eye and swelling in the temporal region of the face. On examination, she has boggy eyelid swelling, proptosis. A CT scan reveals hyperostosis of the sphenoid bone. Identify the false statement regarding the same. (Q853280)

A. CT scan shows tubular enlargement of the optic nerve

B. They are more common in middle-aged women.

C. More common in NF II

D. They are slow-growing tumors from astrocytes in the optic nerve.

Correct Answer: D. They are slow-growing tumors from astrocytes in the optic nerve.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above clinical features and CT findings are suggestive of orbital meningiomas. Orbital meningiomas are not tumors derived from astrocytes in the optic nerve. Instead, they originate from the meninges, the protective coverings of the brain and spinal cord. Astrocytes are the cells from which optic nerve gliomas arise, not meningiomas. Meningioma More common in middle-aged women. (Option B) Predominant feature is early visual loss. Patients present with swollen or atrophic optic discs, restriction of eye movements. More common in NF II (Option C) High-resolution CT scans show tubular enlargement of the optic nerve (Option A), and useful vision is retained; the orbit may be explored and the lesion removed without endangering vision.

References:

Parson’s Diseases of The Eye, 23rd Edition, Page no, 1155-1160.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Which of the following is not part of the classical triad of Sturge-Weber syndrome? (Q853279)

A. Port-wine stain (nevus flammeus) in the distribution of the trigeminal nerve

B. Secondary congenital glaucoma

C. Bilateral optic nerve glioma

D. Ipsilateral meningeal hemangiomatosis

Correct Answer: C. Bilateral optic nerve glioma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Bilateral optic nerve glioma is not part a of the classical triad of Sturge-Weber syndrome . This is a characteristic feature of Neurofibromatosis Type 1 (NF1) Sturge-Weber Syndrome Gene involved: Somatic activating mutation in the GNAQ gene Triad: Facial port-wine stain (in trigeminal nerve distribution) Leptomeningeal angiomas Ocular abnormalities (e.g. glaucoma)



References:

Yanoff and Duker Ophthalmology, 5th edition, page 852, 853, 854.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 7-year-old male child presents with rapidly progressing proptosis in the right eye, which starts suddenly and mimics an acute inflammatory process. Imaging shows a well-defined orbital mass with adjacent bone destruction. What is the most common histopathological subtype of this tumor? (Q853278)

A. Alveolar sarcoma

B. Embryonal sarcoma

C. Botryoid sarcoma

D. Spindle cell sarcoma


Correct Answer: B. Embryonal sarcoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The child presents with rapidly progressive, sudden-onset proptosis that mimics acute inflammation. Imaging shows a well-defined orbital mass with bone destruction, which is suggestive of Rhabdomyosarcoma Embryonal sarcoma is the most common subtype of rhabdomyosarcoma, accounting for 60% of cases, particularly in children. Rhabdomyosarcoma Most common so tissue sarcoma of childhood (90% of cases are seen in children under 16 year of age). It is the most common primary orbital malignancy in children Derived from undifferentiated mesenchymal cells, it is a connective tissue tumor with the potential to differentiate into striated muscle cells. Histopathological Types Embryonal : 60% Alveolar (Option A): 31% Botryoid (Option C): 6% Spindle cells (Option D): 3% Anaplastic: Rare





References:

Kanski's Clinical Ophthalmology, 10th Edition, Page no, 148-149.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 50-year-old woman presents with painless proptosis of the right eye for the past year. She denies any visual changes, pain, or other ocular symptoms. On examination, there is mild proptosis. MRI reveals a well-defined, intraconal mass with homogeneous enhancement, suggestive of a cavernous hemangioma. What is the most appropriate management for this patient? (Q853277)

A. Surgical excision

B. Radiotherapy

C. Steroid injections

D. Observation and regular follow-up

Correct Answer: D. Observation and regular follow-up

Explanation: When orbital cavernous hemangioma is small, asymptomatic , and does not affect vision or cause significant cosmetic concerns, observation with regular follow-up is the preferred management. Cavernous hemangioma of orbit: It is a low-flow, non-distensible orbital venous malformation The primary location is intraconal space Treatment: Asymptomatic cases: Observation and periodic follow-up Indications for surgical excision are: (Option A) Symptomatic CVMs Visual loss due to optic nerve compression Disfiguring proptosis Dystopia Other treatment options are stereotactic radiotherapy, pre-operative embolization percutaneous sclerotherapy, and intralesional injection of pingyangmycin (bleomycin A5) (Options B & C)



References:

Parsons diseases of the eye 23rd edition, page 1154

Orbital Vascular Anomalies - StatPearls - NCBI Bookshelf (nih.gov)

Image; Prepladder notes

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Which among the following statements is/are true regarding orbital varices? Proptosis worsens with the Valsalva maneuver. It is thin-walled distensible high-flow veins. Venous stasis can lead to the formation of phlebolith. Endovascular embolization is contraindicated. (Q853276)

A. 1 and 3

B. 1, 3, and 4

C. Only 2

D. All of the above

Correct Answer: A. 1 and 3

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Orbital varices: It is a plexus of thin-walled dilated low-flow vein (Statement 2 ruled out) Clinical manifestation: Soft-tissue compressible swelling with a ‘bag of worms’ Proptosis, diplopia, visual disturbances, or ecchymosis Distension on dependent position, coughing or Valsalva maneuver (Statement 1) Diagnosis: Dynamic investigations, such as Valsalva-augmented MRI/CT Venous stasis can lead to the formation of a phlebolith (Statement 3) , which is visualized on CT or plain X-ray Treatment: Endovascular embolization (Statement 4 ruled out) CO2 laser ablation Surgical excision is very difficult due to the friable nature with the tendency to bleed

References:

Parsons diseases of the eye 23rd edition page 1154

Orbital Varices - EyeWiki

Orbital Vascular Anomalies - StatPearls - NCBI Bookshelf (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot


Q10: A 35-year-old man presents with a progressive painless swelling in the superolateral part of his left orbit over the past year. Examination reveals mild proptosis and downward and inward displacement of the eye. CT of the orbit shows a well-circumscribed, smooth, encapsulated mass in the lacrimal gland region. Excisional biopsy is shown below. Which of the following is the most likely diagnosis? (Q853275)

A. Adenoid cystic carcinoma

B. Squamous cell carcinoma

C. Pleomorphic adenoma

D. Lymphoma




Correct Answer: C. Pleomorphic adenoma


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: A slow-growing, painless, superolateral orbital mass with proptosis and downward/inward globe displacement is typical of a lacrimal gland tumor. On CT, a well-circumscribed, smooth, encapsulated lesion strongly suggests a benign tumor. The most common benign lacrimal gland tumor is pleomorphic adenoma (benign mixed tumor) , composed of both epithelial and stromal elements. Pleomorphic Lacrimal Gland Adenoma: Histology: The inner layer of cells forms glandular tissue associated with squamous differentiation and keratin production. The outer cells undergo metaplastic change leading to the formation of myxoid tissue Adenoid cystic carcinoma (Option A) : Most common malignant lacrimal gland tumor. Presents with rapidly progressive, painful

proptosis due to perineural invasion. Squamous cell carcinoma (Option B): Rare in lacrimal gland. Aggressive, destructive, not encapsulated. Lymphoma (Option D): Presents as a diffuse, firm orbital mass that molds around the globe without causing bone changes.





References:

Parsons diseases of the eye 23rd edition page 1105-06

Image 1: Parsons diseases of the eye 23rd edition, page 1106 FIG. 24.2 (A)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: Which is the most common non-epithelial malignant lacrimal gland tumor? (Q853274)

A. Pleomorphic adeno-carcinoma

B. Adenoid cystic carcinoma

C. Lymphoma


D. Mucoepidermoid carcinoma


Correct Answer: C. Lymphoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Lymphoma is the most common non-epithelial malignant lacrimal gland tumor. Most common lacrimal gland tumor: Pleomorphic adenoma Most common malignant lacrimal gland tumor: Adenoid cystic carcinoma





References:

Parsons diseases of the eye, 23rd edition, pg no 1105-06

Incidence, clinical features, and survival outcomes of primary malignant lacrimal gland tumors: A population■based analysis - PMC (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 45-year-old male presents with double vision, drooping of the right eyelid, and difficulty moving his right eye in multiple directions. On examination, the patient has normal visual acuity, ptosis, restricted eye movement, decreased sensation over the forehead, and absent corneal reflex on the same side. What is the most likely diagnosis and which cranial nerve is/are most likely affected in this condition? (Q853273)

A. Superior orbital fissure syndrome; III, IV, V, and VI nerves

B. Orbital apex syndrome; optic nerve

C. Tolosa Hunt syndrome; II, III and IV nerves

D. Rochen-Duvigneaud syndrome; only III nerve

Correct Answer: A. Superior orbital fissure syndrome; III, IV, V, and VI nerves

Explanation: Superior orbital fissure syndrome typically involves cranial nerves III (oculomotor), IV (trochlear), V1 (ophthalmic branch of the trigeminal nerve), and VI (abducens), which pass through the superior orbital fissure. Clinical manifestation of ptosis (III), ophthalmoplegia (III, IV, VI), sensory loss over the forehead, and absent corneal reflex (V1) point towards the diagnosis of this syndrome. Orbital apex syndrome; optic nerve (Option B) : It is a symptom complex due to the involvement of structures present at the apex of the orbit (structures passing through the superior orbital fissure and the optic canal) So the cranial nerves involved are II, III, IV, V, and VI Optic nerve (II) involvement causes visual loss and afferent pupillary defect Tolosa Hunt syndrome; II, III, and IV nerves (Option C) : Painful ophthalmoplegia from granulomatous inflammation in cavernous sinus/superior orbital fissure. Typically involves III, IV, VI, and V1, vision loss is not a defining feature. Rochon–Duvigneaud syndrome; only III nerve (Option D): This refers to isolated oculomotor nerve palsy (after orbital trauma or fracture).

References:

www.ncbi.nlm.nih.gov/books/NBK592386/

eyewiki.org/Superior_Orbital_Fissure_Syndrome

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 45-year-old diabetic male presents with complaints of headache, nasal congestion, and facial pain for the past week. On examination, there is periorbital swelling, proptosis, and black eschar on the nasal mucosa. His blood sugar levels are poorly controlled. CT scan of the paranasal sinuses reveals extensive sinus involvement, and biopsy confirms the diagnosis of rhino-orbital-cerebral mucormycosis (ROCM). Which of the following is NOT a known complication of rhino-orbital-cerebral mucormycosis? (Q853272)

A. Meningitis

B. Brain abscess

C. Retinal vascular occlusion

D. Subarachnoid hemorrhage

Correct Answer: D. Subarachnoid hemorrhage

Explanation: Mucormycosis commonly causes meningitis, brain abscess, retinal vascular occlusion, and multiple cranial nerve palsies due to its aggressive spread. However, subarachnoid hemorrhage is not a typical complication of ROCM. Rhino Orbital Cerebral Mucormycosis Etiology Fungal infection (Mucoraceae, primarily Mucor and Rhizopus species) in immunocompromised individuals. Clinical Features Facial and periorbital swelling, pain, proptosis, diplopia, fever. (similar to orbital cellulitis) Tissue necrosis (tissue ischemia). Diagnosis Diagnosis is

clinical , Biopsy : Non-septate broad branching hyphae (Option C) CT or MRI may show mild findings despite severe infection. Treatment Correction of the underlying metabolic defect. Antifungal (intravenous) Daily packing and irrigation of the involved areas with antifungal agent. Wide excision of devitalized and necrotic tissues. Exenteration may be required in unresponsive cases in order to reduce the risk of death. Adjunctive hyperbaric oxygen may be helpful. Prognosis The prognosis is poor due to rapid progression to cerebral involvement. Recurrences are common even with treatment. Complications If untreated, mucormycosis can rapidly progress to Meningitis (Option A ruled out) Brain abscess (Option B ruled out) Retinal vascular occlusion (Option C ruled out) Multiple cranial nerve palsies Death

References:

Kanski's Clinical ophthalmology, 9th edition, page 126.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A 50-year-old male, recently recovered from COVID-19, presents with complaints of severe facial pain, nasal congestion, and black eschar on the left eye with periorbital pain. He has a history of uncontrolled diabetes mellitus. On examination, the following is seen. Which of the following is the most likely causative organism? (Q853271)

A. Aspergillus species

B. Rhizopus species

C. Candida species

D. Cryptococcus neoformans



Correct Answer: B. Rhizopus species

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This scenario describes Rhino-Orbital-Cerebral Mucormycosis , commonly caused by Rhizopus , particularly in immunocompromised patients with uncontrolled diabetes. Rhino Orbital Cerebral Mucormycosis Etiology Fungal infection (Mucoraceae, primarily Mucor and Rhizopus species). High risk among the immunocompromised individuals Clinical Features Facial and periorbital swelling, pain, proptosis, diplopia, fever. (similar to orbital cellulitis) Tissue necrosis (tissue ischemia) Aspergillus species (Option A): Causes aspergillosis – invasive pulmonary aspergillosis, allergic bronchopulmonary aspergillosis (ABPA), aspergilloma. Candida species (Option C): Causes candidiasis – oral thrush, vulvovaginal candidiasis, candidemia, disseminated candidiasis. Cryptococcus neoformans (Option D): Causes cryptococcosis – commonly cryptococcal meningitis and pulmonary cryptococcosis (especially in immunocompromised patients).

References:

Kanski's Clinical ophthalmology, 9th edition, Page 126.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 32-year-old male had blunt trauma to the face during a sports event. He reports difficulty moving his right eye upward and experiences double vision when looking up. On examination, there is tenderness at the right infraorbital rim, and the patient exhibits noticeable swelling and bruising around the right eye. Which wall of the orbit is most likely affected in this case? (Q853256)

A. Medial wall

B. Lateral wall

C. Floor


D. Roof

Correct Answer: C. Floor

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: s: In the presented case, the patient exhibits difficulty moving the right eye upward (limitation of upward gaze) and diplopia (double vision) when looking up. These symptoms are characteristic of a blowout fracture. Difficulty in upward gaze and diplopia occur because: Soft tissue edema/hemorrhage Entrapment of the inferior rectus muscle in the fracture site. Note: Weakest wall of the orbit is the medial wall (due to the lamina papyracea of the ethmoid bone, very thin like paper).



References:

Kanski’s clinical ophthalmology, 9th edition, Page 114

www.ncbi.nlm.nih.gov/books/NBK534825/

↑ *Back to this question in Part A*

Q16: A 10-year-old girl presents with a swollen, painful left eye that started after a recent cold. She is febrile and on examination, her left eyelid is erythematous and swollen, with chemosis, proptosis, limited ocular movement but normal visual acuity. What is the most common primary source of this eye infection in the child? (Q853269)

A. Ethmoid sinusitis

B. Maxillary sinusitis

C. Frontal sinusitis

D. Otitis media

Correct Answer: A. Ethmoid sinusitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical features are suggestive of orbital cellulitis. In children, ethmoid sinusitis is the most common cause of orbital cellulitis. Causes of orbital cellulitis: Sinusitis: Ethmoid sinusitis (most common) carries a high risk of causing orbital cellulitis. Underlying dental problem. Dacryocystitis (inflammation of the lacrimal sac). Endogenous cause: Septicemia Exogenous cause: Fracture or surgery of the orbit. Organisms responsible: Staphylococcus aureus (most common) Haemophilus influenzae (most common in children) Streptococcus pneumoniae Streptococcus pyogenes



References:

Parson’s Diseases of the eye, 23rd edition, Page 137.

Kanski’s Clinical ophthalmology, 9th edition Page 125.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A 25-year-old male presents with intermittent bulging of his right eye, worsening with cough and bending forward. His vision is intact, and there is no history of trauma or infection. What is the most likely cause of his condition? (Q853268)

A. Cavernous sinus thrombosis

B. Orbital varix

C. Thyroid orbitopathy


D. Orbital cellulitis

Correct Answer: B. Orbital varix

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The intermittent proptosis triggered by cough and bending forward, along with the absence of pain or inflammation, suggests a vascular cause like orbital varix. Causes of intermittent proptosis: Orbital varix Periodic orbital edema Recurrent orbital hemorrhage Highly vascular tumors Orbital Varices Dilated venous channels in the orbit (part of the spectrum of developmental venous anomalies) Clinical features Most cases are unilateral and the most frequent site is upper nasal. Slowly progressive intermittent proptosis which worsens with head position changes, bending forward, or Valsalva maneuver. Pathology May involve one large vessel or multiple ectatic veins with fibrotic walls. Thrombosed lumens can contain phleboliths from calcified thrombi. Imaging CT - serpiginous, dense structures that enlarge with Valsalva, enhances with contrast administration. MRI - T1 images show low isointense signal, T2 shows dark signal void from flowing blood. Enhancement with Gadolinium is marked, except in areas of thrombosis. Treatment No treatment needed for mild symptoms; Surgical or embolization options for severe cases. Prognosis Good unless large lesions compress the optic nerve, risking vision loss. Cavernous sinus thrombosis (Option A) typically presents with severe pain, ophthalmoplegia, and rapid onset symptoms, unlike this slow intermittent presentation. Thyroid orbitopathy (Option C) usually causes bilateral, progressive proptosis, not intermittent bulging. Orbital cellulitis (Option D) presents with acute onset, pain, fever, and signs of infection, which are absent here.



References:

Yanoff and Duker Ophthalmology, 5th Edition, Page 1328

Kanski's Clinical Ophthalmology, 9th Edition, Page 135

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A 48-year-old man presents with a complaint of eye protrusion. He describes experiencing a rhythmic, pulsatile sensation in his right eye for the past few weeks. Which of the following is NOT a likely cause ? (Q853267)

A. Carotico-cavernous fistula

B. Saccular aneurysm of the ophthalmic artery

C. Congenital meningocele

D. Orbital cellulitis

Correct Answer: D. Orbital cellulitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This presentation suggests pulsatile proptosis . Pulsatile proptosis, also known as pulsatile exophthalmos , is a condition that causes the eyeball to protrude and pulsate. Orbital cellulitis typically causes proptosis due to inflammation and swelling within the orbit. Presents with pain, redness, swelling, and fever, without the rhythmic pulsation characteristic of pulsatile proptosis. Causes of Pulsatile Proptosis Carotico-cavernous Fistula (Option A) : Abnormal connection between the carotid artery and the cavernous sinus Saccular Aneurysm of the Ophthalmic Artery (Option B) : Aneurysm located on the ophthalmic artery causing pulsatile proptosis Congenital Meningocele or Meningoencephalocele (Option C) : Protrusion of the meninges or brain through a defect in the orbital roof Neurofibromatosis: Genetic disorder causing tumors on nerve tissue, which can affect orbital structures. Traumatic or Operative Hiatus: Defects in the orbital roof resulting from trauma or surgery, allowing transmitted cerebral pulsations

References:

Kanski’s Clinical Ophthalmology, 10th Edition, Page 136, 834

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: If a patient with thyroid eye disease exhibits stable symptoms with no acute inflammation and has experienced a gradual improvement in symptoms over the past 1-2 years, which stage of Rundle’s Curve is most likely being described? (Q853266)

A. Initial Phase

B. Peak Phase

C. Improvement Phase

D. Chronic Phase

Correct Answer: D. Chronic Phase

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Since the patient’s disease has stabilized and no longer exhibits significant progression or acute inflammation, they are in the Chronic or “Burnt-Out” Stage, where further changes are unlikely. In the chronic or "burnt-out" stage of thyroid eye disease, the condition has stabilized after reaching its peak severity. This stage is characterized by stable symptoms that have ceased worsening, the absence of acute inflammation indicating the end of the most active phase, and a gradual improvement in symptoms over the preceding months to years, reflecting the natural progression towards stabilization. Rundle’s Curve and Thyroid Eye Disease: Initial (Option A): Few months; progressive worsening. Peak (Option B): Maximum severity. Improvement (Option C): 1–2 yrs; Gradual spontaneous recovery. Chronic/Burnt-out: Disease stabilizes; no active inflammation or further

progression. Fibrotic changes may persist; surgical correction may be needed. Active Disease: Early, inflamed, medically responsive. Inactive Disease: Stable, non-inflamed, surgery only. Significance: Outlines natural course, guides phase-wise treatment.

References:

https://www.bmj.com/content/bmj/338/7695/Clinical_Review.full.pdf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: A 48-year-old woman with Graves' disease presents with worsening proptosis, diplopia, diminution of vision and difficulty closing her eyes despite treatment with antithyroid medications and corticosteroids. On examination afferent pupillary defects are observed. CT scan shows orbital congestion and muscle enlargement. What is the most appropriate next step in management? (Q853265)

A. Adalimumab

B. Systemic steroids

C. Radiotherapy

D. Decompression Surgery

Correct Answer: D. Decompression Surgery

Explanation: This case is of severe thyroid eye disease . Despite initial treatments, her condition has not improved, indicating a need for more advanced management. Orbital decompression surgery is specifically designed to address proptosis by increasing the volume of the orbit. Decompression surgery is the most appropriate treatment for severe thyroid eye disease. Treatment of Thyroid Eye Disease: Management Details Systemic Steroids (Option B) Help reduce inflammation but do not cure the disease. Immunosuppressors Tocilizumab- Targets IL-6. Infliximab- Targets TNF-α Adalimumab- Targets TNF-α (not specifically for TGF) (Option A) Teprotumumab- An anti-IGF-1 receptor antibody, effective in treating thyroid eye disease Radiotherapy (Option C) Anti-inflammatory radiotherapy helps manage symptoms but is not a cure. Decompression Surgery Performed in severe cases. Orbital Decompression Surgery - Addresses proptosis by increasing the orbital volume. Indications for Surgery - Sight-threatening compressive optic neuropathy, subsided inflammation, failure of initial treatments (Steroids, Immunosuppressors, and Radiotherapy). Decompression Surgery Sequence- Deep lateral wall → Medial wall → Inferior wall → Lateral wall.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 122,123

Bailey and Love’s Short Practice of Surgery, 28th Edition, Page 862

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: A 52-year-old woman with a history of thyroid disease presents with difficulty closing her eyes, noticeable eye protrusion and prominent bilateral lid retraction giving a staring and frightened appearance of the eyes as shown in the image below. Which of the following best describes the sign observed in this patient? (Q853264)

A. Von Graefe's Sign

B. Kocher’s sign

C. Stellwag's Sign

D. Joffroy's Sign



Correct Answer: B. Kocher’s sign

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Kocher sign describes a staring and frightened appearance of the eyes with severe bilateral lid retraction which is particularly marked on attentive fixation. This sign directly correlates with the observed prominent lid retraction in the patient. Lid Signs Dalrymple's Sign Upper lid retraction causing a staring appearance (90% of cases). Von Graefe's Sign Lid lag when the globe moves downward (50% of cases). (Option

A) Enroth's Sign Puffy, oedematous swelling of the eyelids. Gifford's Sign Difficulty in eversion of the upper lid. Stellwag's Sign Infrequent blinking. (Option C) Joffroy's Sign Poor forehead wrinkling on looking up. (Option D) Kocher’s sign Increased lid retraction with visual fixation (staring look)



References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 120

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: A 45-year-old woman with a history of thyroid disease presents with bilateral eye bulging and difficulty closing her eyes. On examination, there is noticeable eye protrusion, eyelid swelling, and restricted extraocular movements. Which of the following best describes the pathogenesis of her eye condition? (Q853263)

A. Extraocular muscles are the primary target of inflammation.

B. The earliest defective ocular movement seen in this patient is depression.

C. T-cells attack the orbital fibroblasts and adipocytes in the orbit.

D. Inflammation is primarily due to neutrophils.

Correct Answer: C. T-cells attack the orbital fibroblasts and adipocytes in the orbit.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This scenario is related to thyroid eye disease (Graves' orbitopathy), where T-cells attack orbital fibroblasts and adipocytes. Pathogenesis of Thyroid Eye Disease (Graves' Orbitopathy) Thyroid eye disease (TED) is an autoimmune inflammatory disorder associated with Graves' disease. Primary autoimmune target: orbital fibroblasts → these fibroblasts differentiate into adipocytes and myofibroblasts under the effect of T-cell–mediated cytokines. This leads to: Pleomorphic inflammatory infiltration Increased secretion of glycosaminoglycans (GAGs) Water retention Enlargement of orbital fat and extraocular muscles These changes raise intraorbital pressure → producing the hallmark features of TED: Proptosis Restricted ocular motility (restrictive myopathy, esp. inferior rectus fibrosis → impaired elevation) Optic neuropathy Lid retraction and lid lag Soft tissue congestion Extraocular muscles are affected secondarily (Option A). The autoimmune attack is primarily directed at orbital fibroblasts, not the muscles themselves. The earliest defective ocular movement is elevation (Option B) due to inferior rectus fibrosis (restrictive myopathy). Infiltration is pleomorphic (mixed inflammatory cells), not predominantly neutrophils (Option D).

References:

Kanski's clinical ophthalmology, 9th edition, Pages 118-119.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Which of the following is not a cause of pseudoproptosis? (Q853262)

A. High Myopia

B. Eyelid Retraction

C. Contralateral Enophthalmos

D. Orbital Cellulitis

Correct Answer: D. Orbital Cellulitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Orbital cellulitis is an infection and inflammation of the orbital tissues that causes actual protrusion of the eye (exophthalmos) due to swelling. Pseudoproptosis: Pseudoproptosis is a condition in which the eyeball appears to be proptosed, but actually, there is no forward displacement. Causes of Pseudoproptosis It is seen in high myopia (due to an increase in the size of the eyeball) (Option A) . Lid retraction (Option B) . Shallow orbits (exorbitism) Contralateral ptosis Unilateral enophthalmos. (Option C) .

References:

Parson's Diseases of The Eye, 23rd Edition, Page no, 1132.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: A 34-year-old woman with a history of hyperthyroidism presents with bilateral proptosis and difficulty closing her eyes. To assess the degree of eye protrusion, a Hertel's exophthalmometer is used. Which of the following statements regarding the use of the exophthalmometer is correct? (Q853261)

A. Normal values for exophthalmometer readings are typically between 20 and 30 mm.

B. Difference of 2–3 mm or more between the two eyes is suspicious regardless of the absolute values

C. A reading greater than 10 mm in both eyes is indicative of proptosis.

D. Hertel's exophthalmometer can only be used to measure the degree of proptosis in one eye at a time.

Correct Answer: B. Difference of 2–3 mm or more between the two eyes is suspicious regardless of the absolute values

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Hertel's exophthalmometer is the most commonly used instrument for measuring proptosis. It is preferred because it measures the forward displacement of both eyes simultaneously (Option D) , providing a more accurate and consistent measurement of proptosis and identifying asymmetry between the eyes. Normal exophthalmometer readings are ≤21 mm (Option A) . Anything >21 mm is abnormal. The true threshold for proptosis is >21 mm (Option C)



References:

Kanski's clinical ophthalmology, 9th edition, Page 115-116

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: Which condition is most commonly associated with bilateral proptosis in adults? (Q853260)

A. Orbital cellulitis

B. Thyroid ophthalmopathy

C. Rhabdomyosarcoma

D. Neuroblastoma

Correct Answer: B. Thyroid ophthalmopathy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Thyroid ophthalmopathy is also known as Graves' orbitopathy or thyroid eye disease is the most common cause of bilateral proptosis in adults. Unilateral proptosis Bilateral proptosis Orbital cellulitis Developmental anomalies Inflammatory conditions Vascular anomalies Neoplasms Metabolic diseases Thyroid eye disease (thyroid ophthalmopathy) Orbital myositis Cavernous sinus thrombosis Metastatic neuroblastoma Most common cause of Unilateral proptosis in children is orbital cellulitis (Option A) Rhabdomyosarcoma (Option C) is a cause of proptosis in children Most common cause of bilateral proptosis in children is neuroblastoma (Option D)

References:

Kanski's clinical ophthalmology, 9th edition, Page, 114-116 Parson's Diseases of the Eye, 23rd Edition, Page no, 1132-1134. https://pmc.ncbi.nlm.nih.gov/articles/PMC7204542/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: Which surgical space is typically used for administering a retrobulbar injection? (Q853259)

A. Sub-Tenon Surgical Space

B. Extraconal Surgical Space

C. Intraconal Surgical Space

D. Subarachnoid Surgical Space

Correct Answer: C. Intraconal Surgical Space

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The surgical space typically used for administering a retrobulbar injection is the Intraconal Surgical Space. This space is located within the muscle cone and is the appropriate site for retrobulbar injections to provide anesthesia or other medications directly behind the globe, within the muscle cone. There is a risk of optic nerve injury. Sub-Tenon Surgical Space (Option A) Location: Between the Tenon capsule and the globe Extraconal Surgical Space (Option B) Location: Between the periorbita and the muscle cone Site for: Peribulbar injection Subarachnoid Surgical Space (Option D) Location: Between the optic nerve and the nerve sheath

References:

www.ijooo.org/html-article/13688#:~:text=There%20are%20five%20surgical%20spaces,the%20muscle%20cone

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: Which of the following statements is TRUE regarding the orbital foramina and fissures? (Q853258)

A. Superior orbital fissure is between the medial and lateral walls.

B. Inferior orbital fissure is between the lesser wing of the sphenoid and the maxillary bone.

C. Optic foramen is between the lateral wall and the orbital floor.


D. Optic nerve and ophthalmic artery pass through the optic foramen.


Correct Answer: D. Optic nerve and ophthalmic artery pass through the optic foramen.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The optic foramen, which is part of the roof of the orbit, allows passage of the optic nerve and the ophthalmic artery Superior Orbital Fissure: A bony cleft found at the orbital apex between the roof and lateral wall (Option A) . Located between the lesser and greater wings of the sphenoid bone. Inferior Orbital Fissure: Defined as a space between the lateral wall and the orbital floor (Option B) . Present between the greater wing of the sphenoid and the maxillary bone . Optic Foramen: Part of the roof of the orbit . It is between the lesser wing and the body of the sphenoid (Option C).





References:

Kanski’s clinical ophthalmology, 9th edition, Page 114 The Textbook of Gray’s Anatomy, 42nd Edition, Page no, 770-771.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [01_Orbit] 002 Previous Year Questions

« Back to Folder Directory

[01_Orbit] 002 Previous Year Questions — Quick Jump

Q1 (p. 472) *Q2 (p. 473)* *Q3 (p. 474)* *Q4 (p. 476)* *Q5 (p. 477)* *Q6 (p. 477)*
Q7 (p. 478) *Q8 (p. 480)* *Q9 (p. 481)* *Q10 (p. 482)*


Q1: A 15-year-old female presents with a painless, gradually increasing mass located lateral to the lateral canthus, present for the past 10 years. On examination, the mass is non-tender and slowly progressive. What is the most likely diagnosis? (Q866349)

A. Dermoid cyst

B. Capillary hemangioma

C. Lacrimal gland tumor

D. Epidermoid cyst



Correct Answer: A. Dermoid cyst

Explanation: Dermoid cyst A painless, slowly progressive, lateral canthal mass present since childhood suggests a congenital lesion at a suture line, which is most characteristic of a dermoid cyst. Dermoid cysts are congenital choristomas caused by ectodermal tissue being trapped along embryonic suture lines, most commonly at the frontozygomatic suture. They contain epidermis and dermal appendages like hair follicles and sebaceous glands. The lesion is firm, non-tender, and slow-growing. Imaging (CT/MRI) shows a well-circumscribed cyst with fat density. Definitive management is surgical excision with intact capsule to prevent recurrence and inflammation. Capillary hemangioma (Option B): Appears in infancy, is vascular, red/bluish, and rapidly proliferates before involution — not a 10-year static mass. Lacrimal gland tumor (Option C): Usually in adults, causes superolateral orbital swelling with globe displacement; may have pain. Epidermoid cyst (Option D): Contains only epidermis without dermal appendages; location and imaging features make dermoid more likely in this case.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Which of the following is true about orbital cellulitis? (Q866363)

A. It is present anterior to the orbital septum

B. Treated effectively with topical antibiotics

C. Presents with proptosis, orbital swelling, normal pupil, and extraocular movements

D. Ethmoid sinusitis is the most common etiology

Correct Answer: D. Ethmoid sinusitis is the most common etiology

Explanation: Ethmoid sinusitis is the most common etiology Orbital cellulitis is an infection of orbital (post-septal) tissues that most often occurs by contiguous spread from adjacent paranasal sinuses. The ethmoid sinuses lie immediately medial to the orbit and separated by the very thin lamina papyracea, making ethmoid sinusitis the single most common source—especially in children. Orbital cellulitis: Infection of orbital tissues posterior to the orbital septum (Option A) (orbital fat, extraocular muscles, occasionally optic nerve). Clinical features: Eyelid swelling, erythema, fever, painful, restricted extraocular movements, proptosis, chemosis, decreased vision, possible RAPD (Option C) if optic nerve involved. Differentiation: Key distinction from preseptal cellulitis is painful ophthalmoplegia, proptosis, and visual compromise in orbital cellulitis. Investigation: Urgent contrast CT or MRI of orbits + sinuses to define extent, look for subperiosteal/orbital abscess, and identify sinus disease. Management: Hospitalize → IV broad-spectrum antibiotics (Option B) covering Staph/Strep/H. influenzae/anaerobes; ENT/ophthalmology consult; surgical drainage if abscess or failure to improve. Complications: Orbital/subperiosteal abscess, cavernous sinus thrombosis, meningitis, intracranial abscess, permanent vision loss—hence early aggressive treatment is mandatory.



@Neet_pg_bot



References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: What is the most common cause of proptosis in a middle-aged person? (Q853127)

A. Orbital varix B) Thyroid eye disease C) Retinoblastoma D) Cavernous hemangioma

B. Thyroid eye disease

C. Retinoblastoma

D. Cavernous hemangioma

Correct Answer: B. Thyroid eye disease

Explanation: Thyroid Eye disease Clinical Features : Clinical Features Details Lid Signs Dalrymple's Sign Upper lid retraction causing a staring appearance (90% of cases). Von Graefe's Sign Lid lag when the globe moves downward (50% of cases). Enroth's Sign Puffy, oedematous swelling of the eyelids. Gifford's Sign Difficulty in eversion of the upper lid. Stellwag's Sign Infrequent blinking. Joffroy's Sign Poor forehead wrinkling on looking up. Kocher’s sign Increased lid retraction with visual fixation (staring look) Conjunctival Signs Deep Injection Pronounced redness. Chemosis Swelling of the conjunctiva. Pupillary Signs Less significant; may show unequal dilatation of pupils. Ocular Motility Defects Mobius's Sign Convergence weakness. Restrictive Myopathy Limited movement in the direction opposite to the affected muscle. Exophthalmos Both eyes are often affected with possible unilateral proptosis. (Option B) Orbital varix (Option A) causes intermittent proptosis. Retinoblastoma (Option C) does not cause proptosis. Cavernous hemangioma (Option D) causes proptosis but it is not the most common cause.



References:

Kanski’s clinical ophthalmology, 9th edition, Page 120

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A patient presents with proptosis that increases when bending down. What is the most likely diagnosis? (Q852503)

A. Orbital Varices

B. Thyroid Eye Disease

C. Cavernous Sinus Thrombosis

D. Orbital Cellulitis

Correct Answer: A. Orbital Varices

Explanation: Orbital Varices are abnormal dilations of veins in the orbit. They often present with proptosis (bulging of the eye) that can worsen with increased venous pressure, such as when the patient bends down or performs actions that increase intra-abdominal or intra-thoracic pressure. Causes of intermittent proptosis: Orbital varix Periodic orbital oedema Recurrent orbital hemorrhage Highly vascular tumors Orbital Varices Dilated venous channels in the orbit (part of the spectrum of developmental venous anomalies) Clinical features Most cases are unilateral and the most frequent site is upper nasal. Slowly progressive intermittent proptosis which worsens with head position changes, bending forward, or Valsalva maneuver. Pathology May involve one large vessel or multiple ectatic veins with fibrotic walls. Thrombosed lumens can contain phleboliths from calcified thrombi. Imaging CT - serpiginous, dense structures that enlarge with Valsalva, enhances with contrast administration. MRI - T1 images show low isointense signal, T2 shows dark signal void from flowing blood. Enhancement with Gadolinium is marked, except in areas of thrombosis. Treatment No treatment needed for mild symptoms; Surgical or embolization options for severe cases. Prognosis Good unless large lesions compress the optic nerve, risking vision loss. Thyroid Eye Disease (Option B): This condition typically causes bilateral proptosis due to inflammation and fibrosis of the extraocular muscles. While proptosis occurs, it does not usually worsen with bending down and is often associated with thyroid dysfunction. It has a more gradual onset compared to conditions like orbital varices. Cavernous Sinus Thrombosis (Option C): This serious condition presents with painful proptosis, usually accompanied by headache, cranial nerve palsies, and vision changes. Proptosis does not typically worsen with bending down, and the condition is associated with acute symptoms and systemic signs of infection. Orbital Cellulitis (Option D): This is an infection of the orbit, causing painful proptosis, redness, and swelling, often accompanied by fever. Pr

References:

Yanoff and Duker Ophthalmology, 5th edition, Page 1328 Kanski’s Clinical ophthalmology, 9th edition, Page 135

↑ *Back to this question in Part A*

@Neet_pg_bot


Q5: Which of the following statements is false regarding the condition shown in the image? (Q844005)

A. NOSPEC classification is used

B. Management corresponds to an improvement in the thyrotoxic state

C. Vision loss may occur

D. It is not seen in 10% of patients with hyperthyroidism



Correct Answer: B. Management corresponds to an improvement in the thyrotoxic state

Explanation: The condition shown is Thyroid Eye Disease (Graves' Ophthalmopathy) , an autoimmune inflammatory disorder associated with Graves' disease. The false statement is that management of thyroid eye disease corresponds directly to improvement in the thyrotoxic state — this is incorrect because the course of ophthalmopathy is independent of thyroid hormone levels. NOSPECS classification (Option A): True – used to assess disease severity. Vision loss (Option C): True – may occur due to compressive optic neuropathy or corneal ulceration. Not seen in 10% (Option D): True – eye signs occur in about 25–50% of Graves' patients, so it's absent in some cases. Thyroid eye disease has two distinct phases: Active (Inflammatory) Phase: Managed with corticosteroids or radiotherapy. Inactive (Burnt-out) Phase: Managed surgically if needed. Improving hyperthyroidism may prevent progression but does not directly cause improvement of existing ophthalmopathy.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: What is the most common cause of proptosis in a middle-aged person? (Q840073)

A. Orbital varix

B. Thyroid eye disease

C. Retinoblastoma

D. Cavernous hemangioma

Correct Answer: B. Thyroid eye disease

Explanation: Correct Option B - Thyroid eye disease: The most common cause of both unilateral and bilateral proptosis in adults is thyroid ophthalmopathy. In children, the most common cause of unilateral proptosis is orbital cellulitis and bilateral proptosis is neuroblastoma. Incorrect Options : Option A - Orbital varix: Orbital varix causes intermittent proptosis. Option C - Retinoblastoma: Retinoblastoma does not cause proptosis. Option D - Cavernous hemangioma: Cavernous hemangioma causes proptosis but it is not the most common cause.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: Following instrument is used to measure? (Q854708)

A. Proptosis

B. Squint

C. Ptosis

D. Diplopia



Correct Answer: A. Proptosis

Explanation: The instrument used to measure proptosis is called an exophthalmometer , such as the Hertel or Luedde exophthalmometer. This tool helps assess the degree of ocular protrusion and is particularly useful in diagnosing conditions like proptosis caused by orbital tumors or other space-occupying lesions. {{caption_text}} Proptosis Proptosis refers to the abnormal protrusion of the eyeball, while exophthalmos specifically describes protrusion of the eyeball. Causes: Retrobulbar lesions (most common cause). Shallow orbit (less common). The optic nerve allows significant forward displacement of the globe (25mm length), causing proptosis without excessive nerve stretching. Types of Proptosis: Axial proptosis: Caused by space-occupying lesions within the muscle cone (e.g., cavernous hemangioma, optic nerve tumors). Combined proptosis and dystopia: Due to extraconal lesions (e.g., lacrimal gland tumor) causing displacement in the coronal plane (dystopia). Dystopia: Displacement of the globe in the coronal plane, either horizontal or vertical. Measured using the midline (nose) for horizontal or a vertical scale for vertical displacement. Measuring Severity: Use a plastic rule, Luedde™ exophthalmometer, or a binocular exophthalmometer (Hertel) for accurate measurements of ocular protrusion. Readings >20 mm or a difference of 2–3 mm between eyes suggests proptosis. Palpebral aperture dimensions and any lagophthalmos should be noted. Pseudoproptosis: A false impression of proptosis due to: Facial asymmetry. Enlarged globe (e.g., high myopia, buphthalmos). Lid retraction or contralateral enophthalmos. Clinical evaluation of proptosis Differentiate proptosis from pseudoproptosis Pseudoproptosis (the false impression of proptosis) may be due to facial asymmetry, enlargement of the globe (high myopia, buphthalmos), lid retraction or contralateral enophthalmos. Asymmetrical proptosis is readily detected by looking down at the patient from above and behind Globe location: If it is misplaced superiorly, inferiorly, medially, laterally, or straight out (axially) by an exophthalmometer. An exophthalmometer is used to quantify the distance from the orbital rim to the corneal surface of each eye in order to assess the symmetry of globe protrusion. Readings >20 mm are indicative of proptosis and a dierence of 2–3 mm or more between the two eyes is suspicious regardless of the absolute values. (Option B) Hertel's exophthalmometer: In adults. Luedde exophthalmometer: In children. Naugle exophthalmometer: Non-axial proptosis Squint (Option B) , also known as strabismus, refers to a misalignment of the eyes, where the eyes do not point in the same direction. The condition is measured using ocular alignment tests (like the Hirschberg test or the Cover test), not an exophthalmometer. Ptosis (Option C) refers to the drooping of the upper eyelid. The severity of ptosis is usually assessed by observing the degree of eyelid droop, often with the help of a marginal reflex distance (MRD) test or by measuring the distance from the upper eyelid to the corneal light reflex. This is not measured using an exophthalmometer. Diplopia (Option D) , or double vision, refers to the perception of two images of a single object. This condition is typically evaluated through visual field tests, ocular motility testing, and cover tests, not with an exophthalmometer.



References:

Kanski's Clinical Ophthalmology 9th ed., Page 114- 116

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A middle-aged woman presented with bilateral proptosis, restriction of eye movements, and chemosis. Her thyroid profile showed she was hyperthyroid. What is the most probable cause? (Q858505)

A. Orbital pseudotumor

B. Grave's ophthalmopathy

C. Orbital lymphoma

D. Orbital cellulitis

Correct Answer: B. Grave's ophthalmopathy

Explanation: The clinical scenario of bilateral proptosis, restriction of eye movements, chemosis and hyperthyroidism is suggestive of Grave’s ophthalmology {{caption_text}} Thyroid Eye Disease Clinical Features : Clinical Features Details Lid Signs Dalrymple's Sign Upper lid retraction causing a staring appearance (90% of cases). Von Graefe's Sign Lid lag when the globe moves downward (50% of cases). Enroth's Sign Puffy, oedematous swelling of the eyelids. Gifford's Sign Difficulty in eversion of the upper lid. Stellwag's Sign Infrequent blinking. Joffroy's Sign Poor forehead wrinkling on looking up. Kocher’s sign Increased lid retraction with visual fixation (staring look) Conjunctival Signs Deep Injection Pronounced redness. Chemosis Swelling of the conjunctiva. Pupillary Signs Less significant; may show unequal dilatation of pupils. Ocular Motility Defects Mobius's Sign Convergence weakness. Restrictive Myopathy Limited movement in the direction opposite to the affected muscle. Exophthalmos Both eyes are often affected with possible unilateral proptosis. Orbital pseudotumor (Option A) It is characterized by congestive proptosis, ophthalmoplegia, and optic nerve dysfunction and generally present as a unilateral condition in adults. Orbital lymphoma (Option C) It is characterized by discomfort, bulging eye, diplopia, and visible mass and can be unilateral or bilateral.. Orbital cellulitis (Option D) It is the most common cause of unilateral proptosis in children and is characterized by unilateral, painful proptosis and fever.



References:

Kanski’s clinical ophthalmology, 9th edition, Page 120

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: A 35-year-old female reports recurring episodes of discomfort, pain, and a sensation of pressure behind her left eye. She exhibits intermittent reducible proptosis,

and her symptoms intensify when she bends forward. What is the most likely diagnosis? (Q830803)

A. Orbital hemangioma

B. Orbital varices

C. Orbital metastasis

D. Orbital dermoid cyst

Correct Answer: B. Orbital varices

Explanation: Correct Option B - Orbital varices: Orbital varices are vascular anomalies characterized by dilated and tortuous veins in the orbit, which are associated with intermittent proptosis. Proptosis may worsen when the patient is in an upright position or during activities that increase intra-abdominal pressure, such as coughing or bending forward. Incorrect Options: Option A - Orbital hemangioma: Orbital hemangiomas are benign vascular tumors of the orbit. They typically present as painless, slow-growing masses without the characteristic features of intermittent proptosis. Option C - Orbital metastasis: Orbital metastasis is less probable because of younger age and the recurrent nature of the symptom. Option D - Orbital dermoid cyst: Orbital dermoid cyst is characterized by a painless nodule or proptosis.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: What is the most common primary intraocular malignancy in adults? (Q830808)

A. Retinoblastoma

B. Malignant Choroidal melanoma

C. Reticulum cell sarcoma

D. Neurofibroma

Correct Answer: B. Malignant Choroidal melanoma

Explanation: Correct Option B - Malignant Choroidal melanoma: Malignant choroidal melanoma is the most common primary intraocular malignancy in adults. It originates from the pigment-producing cells (melanocytes) in the choroid, which is a vascular layer beneath the retina in the eye. Choroidal melanoma is more prevalent in adults, particularly in individuals with fair skin and light-colored eyes. Incorrect Options: Option A – Retinoblastoma: It is the most common primary intraocular malignancy in Children. Option C - Reticulum cell sarcoma: Reticulum cell sarcoma, or primary intraocular lymphoma, is a type of non-Hodgkin lymphoma that can occur in the eye. While it is a primary intraocular malignancy, it is less common than malignant choroidal melanoma. Option D – Neurofibroma: Neurofibromas are tumors that arise from nerves. They are often associated

with neurofibromatosis type 1 (NF1), a genetic condition.

References:

..

↑ *Back to this question in Part A*

@Neet_pg_bot

• [02_Diseases of Lens] 003 Lens And Disorders - I

« Back to Folder Directory

[02_Diseases of Lens] 003 Lens And Disorders - I — Quick Jump

Q1 (p. 484)	*Q2 (p. 485)*	*Q3 (p. 486)*	*Q4 (p. 487)*	*Q5 (p. 488)*	*Q6 (p. 489)*
Q7 (p. 492)	*Q8 (p. 496)*	*Q9 (p. 498)*	*Q10 (p. 499)*	*Q11 (p. 500)*	*Q12 (p. 502)*
Q13 (p. 503)	*Q14 (p. 504)*	*Q15 (p. 505)*	*Q16 (p. 506)*	*Q17 (p. 507)*	*Q18 (p. 508)*
Q19 (p. 509)	*Q20 (p. 510)*	*Q21 (p. 511)*	*Q22 (p. 513)*	*Q23 (p. 514)*	*Q24 (p. 516)*
Q25 (p. 517)	*Q26 (p. 518)*	*Q27 (p. 520)*	*Q28 (p. 521)*	*Q29 (p. 522)*	*Q30 (p. 523)*
Q31 (p. 525)	*Q32 (p. 526)*	*Q33 (p. 527)*	*Q34 (p. 529)*	*Q35 (p. 530)*	*Q36 (p. 532)*
Q37 (p. 532)	*Q38 (p. 534)*	*Q39 (p. 535)*	*Q40 (p. 536)*	*Q41 (p. 537)*	

Q1: What kind of laser can be utilized to treat posterior capsular opacification, which is observed during an examination of a patient who has reported decreased visual acuity following cataract surgery? (Q882185)

A. Femto laser

B. Nd YAG laser

C. Argon fluoride

D. Argon

Correct Answer: B. Nd YAG laser

Explanation: Correct Option B - Nd YAG laser: In the clinical scenario mentioned above decreased visual acuity after the cataract surgery with the presence of posterior capsular opacification is suggestive of After cataract, most common late complication of uncomplicated cataract surgery. It is caused by the proliferation of lens epithelial cells that have remained within the capsular bag following cataract extraction. Nd YAG laser is the preferred technique used to create an opening in the posterior capsule termed as posterior capsulotomy to treat patients with after cataract. Indication for Nd YAG laser posterior capsulotomy are: Reduced visual acuity. Glare. Incorrect Options: Option A - Femto laser: FLACS - Femtosecond laser-assisted cataract surgery is used in refractive surgery. The laser used is a neodymium glass (Nd: glass) laser with a wavelength of 1053 nanometers. Option C - Argon fluoride: It is a deep ultraviolet laser with a wavelength of 193 nanometers. It is an excimer laser that is used in refractive surgeries, trabeculoplasty, and glaucoma. The mechanism of action is photoablation which is the flattening of the cornea. Option D - Argon: Argon lasers are photocoagulative lasers with wavelengths of 488 nm and 514 nm used for trabeculoplasty, lipid keratopathy, and glaucoma.

References:

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A newborn presented with the following findings. There is a positive maternal history for fever during the first trimester of pregnancy. Which among the following is least likely to be seen in this newborn? (Q853341)

A. Patent ductus arteriosus

B. Limb hypoplasia

C. Microphthalmia

D. Sensorineural hearing loss





Correct Answer: B. Limb hypoplasia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Limb hypoplasia is a characteristic feature of Congenital Varicella Syndrome , not Rubella. Pearly white cataract and salt & pepper fundus with antenatal history of fever in early pregnancy is classical of Congenital Rubella syndrome. Congenital Rubella syndrome: Cause: Maternal rubella infection acquired in 1st trimester Ocular Defects: Congenital cataract (2nd most common manifestation) Salt and pepper chorioretinopathy (most common feature) Microphthalmia (Option C) Cloudy cornea Poorly dilating pupil Ear Defects: Sensorineural hearing loss (Option D) Cardiovascular Defects: Patent ductus arteriosus (Option A) Pulmonary stenosis Ventricular septal defects

References:

Kanski's clinical ophthalmology, 9th edition, Page 335, 472

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Which of the following is not a condition associated with a shallow anterior chamber? (Q853331)

A. Wound leak with hypotony

B. Ciliochoroidal detachment

C. Vitreous bulge

D. Iridodialysis

Correct Answer: D. Iridodialysis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Iridodialysis, which is a separation of the iris from its attachment to the ciliary body, typically does not cause a shallow anterior chamber. Flat/Shallow anterior chamber Early Postoperative complications of cataract surgery. It may occur due to wound leak, ciliochoroidal detachment and pupil block. Flat anterior chamber with wound leak: (Option A) Associated with hypotony, diagnosed by Seidel's test.Fluorescein diluted by aqueous at site of leakage. Treatment: Most cases cured within 4 days with pressure bandage and oral acetazolamide.If persistent: injection of air in anterior chamber and suturing of leaking wound. Ciliochoroidal detachment: (Option B) Presents as convex brownish mass in an involved quadrant with shallow anterior chamber. Treatment: Most cases cured within 4 days with pressure bandage and oral acetazolamide.If persistent, suprachoroidal drainage with injection of air in anterior chamber. Pupil block due to vitreous bulge: (Option C) Occurs after ICCE and in ECCE technique due to adhesions with IOL (more common with ACIOLs). Leads to formation of iris bombe and shallowing of anterior chamber. If persistent for 5-7 days, it may form permanent peripheral anterior synechiae (PAS) leading to secondary angle closure glaucoma. Treatment: Initially treated with mydriatic, hyperosmotic agents (e.g. 20% mannitol) and acetazolamide.If not relieved, laser peripheral iridotomy should be performed to bypass the pupillary block■■■■■■■■■.

References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 219,220&221. Kanski’s clinical ophthalmology, 9th edition, Page 900

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 67-year-old man had undergone cataract surgery. He presents with endophthalmitis after 3 days. His infection is severe with vision reduced to hand movements close to the face. What is the most appropriate management? (Q853332)

A. Intravitreal antibiotics

B. Systemic antibiotics

C. Steroid therapy


D. Vitrectomy

Correct Answer: D. Vitrectomy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In a patient with post-operative acute bacterial endophthalmitis: Pars plana vitrectomy (if vision ≤ hand movements). If vision is better than hand movements → Intravitreal antibiotic injections. Intravitreal Antibiotics (for bacterial endophthalmitis) Vancomycin (Gram +ve coverage) Ceftazidime (Gram –ve coverage) ± Dexamethasone (small dose) → to control inflammation Fungal Endophthalmitis Intravitreal Amphotericin B Intravitreal Voriconazole Contraindicated Intravitreal Drugs: due to Maculotoxicity) Gentamicin Amikacin Common Causative Organisms Early onset: Staphylococcus epidermidis (most common) Late onset: Propionibacterium acnes or fungus



References:

www.ncbi.nlm.nih.gov/books/NBK559079/

Kanski’s clinical ophthalmology, 9th edition, Page 327

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 65-year-old man has undergone cataract surgery. He has now returned after 6 weeks with endophthalmitis. What is the most likely organism that caused the infection? (Q853333)

A. Staphylococcus aureus

B. Streptococcus pyogenes

C. Propionibacterium acnes

D. Staphylococcus epidermidis

Correct Answer: C. Propionibacterium acnes

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The scenario is that of late-onset postoperative endophthalmitis which is most commonly caused by Propionibacterium acnes Endophthalmitis is inflammation of the inner structures of the eye (uveal tissue and retina) with exudates in the vitreous, anterior, and posterior chambers Late-onset endophthalmitis is also caused by fungus.

References:

https://www.ncbi.nlm.nih.gov/books/NBK559079/ Kanski's clinical ophthalmology, 9th edition, Page 330

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Soemmering's Ring and Elschnig's Pearls are examples of which of the following? (Q853334)

A. After cataracts

B. Senile cataracts

C. Congenital cataracts


D. Toxic cataracts


Correct Answer: A. After cataracts


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Soemmering's Ring and Elschnig's Pearls are examples of after cataract/Secondary cataract M/C postoperative complication following extracapsular lens extraction. Posterior Capsule Opacification (PCO): Common form of after cataract. Dense Membranous After Cataract: Thick membrane formation. Soemmering's Ring: Thick ring behind the iris, enclosed between capsule layers. Elschnig's Pearls: Clustered, vacuolated subcapsular epithelial cells along the posterior capsule. Mechanism: Elschnig’s Pearls: Residual cells from the lens equator migrate onto the posterior capsule. These cells undergo a process of swelling and vacuolation. To an ophthalmologist looking through a slit lamp, they appear as tiny, translucent, grape-like clusters or "pearls." If they migrate into the visual axis, they scatter light and blur vision. Soemmering’s Ring: This occurs when the residual epithelial cells (and sometimes cortical fibers) become trapped between the anterior and posterior capsule flaps. The cells proliferate and form a thick, doughnut-shaped ring of tissue in the periphery of the capsular bag. The center usually remains clear, but the ring can cause issues if the pupil is dilated or if the ring becomes large enough to displace the intraocular lens. Treatment: ND YAG laser posterior capsulotomy







References:

Kanski’s clinical ophthalmology, 9th edition, Page 331

↑ *Back to this question in Part A*

@Neet_pg_bot


Q7: Match the images of IOL malpositions with their names: Image Syndrome a. 1. Sunrise syndrome b. 2. Lost lens syndrome c. 3. Sunset syndrome (Q853335)

A. a-1,b-2,c-3


B. a-3,b-1,c-2


C. a-2,b-1,c-3

D. a-1,b-3,c-2



@Neet_pg_bot







Correct Answer: B. a-3,b-1,c-2


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: IOL malpositions post cataract surgery: Name Malposition Sunset syndrome Inferior subluxation of IOL Sunrise syndrome Superior subluxation of IOL Lost lens syndrome Complete dislocation of an IOL into the vitreous cavity







References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 221

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: Which of the following statements is true regarding Cystoid Macular Edema? (Q853336)

A. Collection of fluid in Henle’s layer

B. Honeycomb appearance in fundoscopy

C. Antiprostaglandin is both treatment and prevention


D. All of the above


Correct Answer: D. All of the above


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Cystoid macular edema: Definition: Collection of fluid in cystic loculi in the Henle's layer of the macula Defect in OPL layer of retina(inner retinal blood barrier break defect) Frequent complication of cataract surgery (Irvine–Gass Syndrome) Mechanism: Ocular surgery, trauma, or uveitis triggers the release of inflammatory markers, most notably prostaglandins , leukotrienes, and VEGF (Vascular Endothelial Growth Factor). These mediators cause the junctions between the endothelial cells of the retinal capillaries to become "leaky." Plasma fluid leaks out of the vessels and into the extracellular space of the retina. Due to the unique anatomy of the macula, this fluid pools in the Henle’s layer and the Inner Nuclear Layer . The Müller cells (supportive cells) can also swell and eventually liquefy, resulting in the formation of the characteristic cystoid spaces. Diagnosis: Fundoscopy: Honeycomb appearance Fluorescein Angiography: Flower petal pattern Management: Prevention: Preoperative and postoperative use of anti prostaglandin eye drops (indomethacin, flurbiprofen, or ketorolac) Treatment: Anterior vitrectomy Steroids Antiprostaglandins


8/23/2013
2
@Neet_pg_bot


02:22.61
8/23/2013
31
@Neet_pg_bot



References:

Kanski's clinical ophthalmology, 9th edition, Page 602

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: What is the common inheritance pattern of hereditary cataracts? (Q853337)

A. Autosomal dominant

B. Autosomal recessive

C. X linked recessive

D. X linked dominant

Correct Answer: A. Autosomal dominant

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Cataracts can be inherited without systemic disorders in the autosomal dominant pattern Hereditary cataract: Inherited without Systemic Disorders: Usually autosomal dominant. Genes: Cry-G (crystallins). Connexins. MIP (Major intrinsic protein). Inherited with Systemic Disorders: Chromosomal disorders (e.g., trisomy 21) Skeletal disorders (e.g., Stickler syndrome) CNS disorders (e.g., cerebro-oculo-facial syndrome) Renal system disorders (e.g., Lowe's syndrome)

References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 188

↑ *Back to this question in Part A*

@Neet_pg_bot


Q10: Doctor noticed leukocoria in a 3-month-old baby during a routine examination and the eye appeared as shown below. Choose the least appropriate statement regarding the most likely diagnosis. (Q853338)

A. Most common type of bilateral congenital cataract

B. Commonly inherited as autosomal dominant trait

C. Usually zone of fetal nucleus is involved

D. Associated with vitamin-D deficiency



Correct Answer: D. Associated with vitamin-D deficiency

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The image shows a lamellar (zonular) cataract , which is the most common type of bilateral congenital cataract . It presents as a disc-shaped opacity surrounded by clear lens layers and is often symmetrical in both eyes . Not associated with vitamin D deficiency. Key features: Involves the fetal nucleus or adjacent lamellae . Autosomal dominant inheritance is common. Can occur after maternal rubella infection or metabolic disturbances (like hypocalcemia) during gestation

References:

Parson’s Diseases of the Eye - 23rd edition, page-634-640.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q11: An infant with leukocoria presents with recurrent gastrointestinal infections and developmental regression. Bilateral eye examination revealed the following. Choose the most appropriate next step in evaluation. (Q853339)

A. Serum estimation of glucose, calcium, and phosphorous levels

B. Estimation of antibody titers for infections

C. Urine chromatography for amino acids

D. Urine assay for the presence of reducing substances



Correct Answer: D. Urine assay for the presence of reducing substances

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: An infant with leukocoria associated with recurrent gastrointestinal infections, developmental regression, and exam findings of oil droplet cataract is suggestive of Galactosemia. Classic Galactosemia: Enzyme Deficient: Galactose-1-phosphate uridyl transferase (GALT) . Pathophysiology: This leads to the accumulation of Galactose-1-phosphate in tissues like the liver, brain, and kidney. Clinical Picture: This is a severe systemic disease. Infants present with: Cataracts (Oil droplet type). Liver dysfunction (Hepatomegaly, jaundice). Mental Retardation (Developmental regression). Susceptibility to Infection: Specifically E. coli neonatal sepsis. Galactokinase Deficiency (The "Milder" Form) Enzyme Deficient: Galactokinase (GALK) . Differentiating Factor: Unlike GALT deficiency, Galactokinase deficiency only results in the accumulation of Galactitol . Clinical Picture: The only major clinical finding is the Oil Droplet Cataract . There is no liver damage, no mental retardation, and no increased risk of sepsis. The next step in evaluation is a urine assay for the presence of reducing substances Levels of blood glucose, calcium and phosphorus to rule out Hyperglycemia and Hypocalcemia. (Option A) Antibody titers for infections such as toxoplasmosis, rubella, cytomegalovirus and herpes simplex virus (TORCH) and hepatitis B virus. (Option B) Screening for amino acids in the urine - if Lowe

syndrome is suspected. (Option C)

References:

Parsons diseases of the Eye - 23rd edition, page-641

www.ncbi.nlm.nih.gov/books/NBK470419/

Kanski's clinical ophthalmology, 9th edition, Page 335, 337

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 1-month-old infant is diagnosed with a dense unilateral cataract. Which of the following is the most appropriate management strategy? (Q853340)

A. Delay surgery until the child is 1–2 years old to reduce surgical risks

B. Perform immediate cataract surgery

C. Prescribe glasses and monitor the cataract progression

D. Wait until the child develops strabismus before considering surgery

Correct Answer: B. Perform immediate cataract surgery

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In congenital cataract, the surgery should be performed as early as possible (Within 1–2 months (Option A) of age) Timing Considerations Ideal timing: After 1 month of age Not before 1 month because: High risk of trabeculitis due to inflammation/surgery Trabeculitis → obstructs aqueous humor drainage → causes glaucoma Why Early Surgery is Essential Foveal reflex development: Begins at 1.5 months ; completes by 5–6 months If surgery delayed → improper foveal development → amblyopia Prescribe glasses and monitor the cataract progression (Option C) : Glasses are used post cataract surgery and are not the mainstay of treatment. Wait until the child develops strabismus before considering surgery (Option D) : Waiting until complications is not advised in dense unilateral cataracts as it can lead to permanent vision impairment.

References:

Parson's diseases of the Eye - 23rd edition, page-641,642

Comprehensive Ophthalmology by AK Khurana - 7th edition, page 192

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 70-year-old woman with a history of hypertension and high myopia undergoes cataract surgery. During the procedure, she suddenly experiences severe ocular pain and a rapid rise in intraocular pressure. Fundoscopic examination reveals a dark red elevation in the choroidal region. Which of the following is the most appropriate immediate management step? (Q853330)

A. Intraoperative posterior sclerostomy

B. Increase the infusion pressure in the anterior chamber

C. Perform immediate vitrectomy to drain the hemorrhage


D. Close the surgical wound and monitor intraocular pressure

Correct Answer: D. Close the surgical wound and monitor intraocular pressure

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical scenario suggests suprachoroidal hemorrhage . The immediate steps involve closing the wound to prevent further prolapse of intraocular contents and monitoring intraocular pressure. A suprachoroidal haemorrhage involves a bleed into the suprachoroidal space from a ruptured short posterior ciliary artery Intraoperative posterior sclerostomy (Option A) should not be performed during the management of suprachoroidal hemorrhage because it increases the risk of complications such as infection and retinal detachment. The immediate focus should be on stabilizing the eye by controlling intraocular pressure and closing the incision. Once stabilized, further surgical interventions can be evaluated in a safer environment. Increase the infusion pressure in the anterior chamber (Option B): Increasing infusion pressure may worsen the bleeding. Perform immediate vitrectomy to drain the hemorrhage (Option C): Immediate vitrectomy is generally avoided initially as it could exacerbate the hemorrhage.



References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 328

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A school-going child is diagnosed with a connective tissue disorder characterized by abnormalities of the eye, short stature, brachydactyly, joint stiffness, and cardiovascular defects. Which among the following is incorrect regarding the likely diagnosis? (Q853342)

A. Follows autosomal recessive mode of inheritance

B. Associated with supero-temporal lens dislocation

C. Molecular genetic analysis is confirmatory


D. Usual age of presentation is 5-10 years

Correct Answer: B. Associated with supero-temporal lens dislocation

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical scenario is suggestive of Weill-Marchesani syndrome , in which there is inferior subluxation of the lens, not supero-temporal displacement Supero-temporal dislocation is seen in Marfan's syndrome Weill-Marchesani Syndrome Autosomal recessive (Option A) The usual age of presentation is 5-10 years (Option D) Molecular genetic analysis is confirmatory (Option C) : Genetic testing for ADAMTS10/ADAMTS17 (Autosomal Recessive) or FBN1 (Autosomal dominant) variants. Clinical features: Microspherophakia, Inferior or forward subluxation of the lens Note: Infero-nasal dislocation is seen in Homocystinuria



References:

https://www.ncbi.nlm.nih.gov/books/NBK1114/

Kanski’s clinical ophthalmology, 9th edition, Page 339

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: During early ocular development, the optic vesicle induces the formation of the lens placode from adjacent tissue. What is the embryologic origin of the crystalline lens? (Q885161)

A. Neural crest cells


B. Surface ectoderm

C. Neuroectoderm

D. Mesoderm

Correct Answer: B. Surface ectoderm

Explanation: The lens develops from the surface ectoderm , not from neural tissues. Origin: The crystalline lens develops from surface ectoderm , not neural or mesodermal tissue. Induction: Around the 4th week, the optic vesicle (neuroectoderm) contacts the surface ectoderm, inducing it to thicken into the lens placode . Lens vesicle formation: The placode invaginates , forming a hollow lens vesicle that later detaches from the surface ectoderm. Anterior cells: Become lens epithelium Posterior cells: Differentiate into lens fibers Nutrition: Temporarily supplied by the hyaloid artery , which later regresses. Other surface ectoderm derivatives: Corneal epithelium, conjunctiva, lacrimal glands, tarsal glands. Neural crest cells (Option A) contribute to corneal stroma, endothelium, sclera, uveal stroma, and trabecular meshwork. Neuroectoderm (Option C) forms retina, retinal pigment epithelium, optic nerve fibers, iris, and ciliary body epithelium. Mesoderm (Option D) contributes to extraocular muscles and vascular endothelium.



References:

Parsons Diseases of the Eye, 23rd Edition, Page 27.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: The fibers that are most recently formed in the crystalline lens are located in which region? (Q885162)

A. Lens cortex


B. Embryonic nucleus

C. Fetal nucleus

D. Adult nucleus

Correct Answer: A. Lens cortex

Explanation: The youngest fibers of the crystalline lens are located in the outer cortex , continuously added throughout life beneath the lens capsule. Lens epithelial cells at the equator proliferate throughout life, elongate, lose nuclei, and become lens fibers. Fiber arrangement: Old fibers → compressed centrally → form the nucleus New fibers → added externally beneath the capsule → form the cortex Age-wise fiber distribution: Embryonic nucleus: Oldest fibers (6-12 weeks IU) (Option B ruled out) Fetal nucleus: 3-8 months IU (Option C ruled out) Infantile nucleus: Birth to puberty Adult nucleus: Early adulthood fibers (Option D ruled out) Cortex: Youngest fibers, continuously added throughout life (Option A) Central fibers undergo compaction over time (nuclear sclerosis), while the cortex remains soft and metabolically active.



References:

Parsons Diseases of the Eye, 23rd Edition, Page 34.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A 30-year-old patient is being evaluated during a slit-lamp examination. The examiner explains that the crystalline lens is avascular and depends on surrounding structures for metabolic exchange. What is the primary source of nutrition for the lens? (Q885164)

A. Connective tissue diffusion

B. Zonular fibers

C. Aqueous humor

D. Retinal circulation

Correct Answer: C. Aqueous humor

Explanation: The avascular lens relies on the aqueous humor for nutrients like glucose and amino acids, while waste products diffuse back, with the anterior capsule and epithelium mediating this metabolic exchange. Minor contribution: Vitreous humor posteriorly Mechanisms: Nutrient transport occurs via diffusion and active transport (ion pumps, channels, transporters). Adequate aqueous supply is crucial for lens metabolism and prevention of cataract formation. Incorrect Options: Connective tissue diffusion (Option A): The lens is enclosed by its capsule and separated from the surrounding connective tissue, which does not provide nutrients. Zonular fibers (Option B): Zonules only suspend and position the lens; they do not supply metabolic support. Retinal circulation (Option D): The posterior lens is separated from retinal vessels by the vitreous and receives no direct blood supply.

References:

Parson’s Diseases of Eye, 23rd Edition, Page 51.

Yanoff and Duker Ophthalmology, 5th Edition, Page 328.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A histology slide of the crystalline lens shows a thin capsule with a single layer of cuboidal cells beneath it. The marked region indicates lens epithelium. At which location is this structure normally absent? (Q885166)

A. Posterior capsule region

B. Central anterior surface

C. Equatorial lens region

D. Anterior pole

Correct Answer: A. Posterior capsule region

Explanation: The lens is derived from surface ectoderm and retains a single layer of cuboidal epithelial cells only on its anterior surface beneath the capsule. During development, the posterior epithelial cells elongate to form primary lens fibers, and the epithelial layer disappears posteriorly . Thus, the posterior capsule directly overlies lens fibers with no epithelial lining. Incorrect Options: Central anterior surface (Option B): Lens epithelium is present here and is metabolically most active. Equatorial lens region (Option C): Germinative zone is located here; epithelial cells proliferate and elongate into secondary lens fibers. Anterior pole (Option D): Epithelium is present beneath the anterior capsule, though cells are flatter centrally.

References:

Yanoff and Duker Ophthalmology, 5th Edition, Page 325.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: In the crystalline lens, the antioxidants ascorbate and alpha-tocopherol remain in their reduced (active) form primarily due to which intracellular substance? (Q885169)

A. Glucose

B. Glycoprotein

C. Glutathione

D. Fatty acids

Correct Answer: C. Glutathione

Explanation: The lens is continuously exposed to oxidative stress from light and metabolic activity. To maintain transparency, it relies on a robust antioxidant system. Reduced glutathione (GSH) is the principal intracellular reducing agent in lens fibers. Regenerates oxidized ascorbate (vitamin C) and alpha-tocopherol (vitamin E) back to their active reduced forms. Detoxifies hydrogen peroxide via glutathione peroxidase , protecting crystallins and membrane lipids. Prevents oxidative damage, delaying cataract formation. Glucose (Option A) is the main metabolic substrate of the lens, but it does not function as the principal intracellular reducing agent. It participates in glycolysis and the sorbitol pathway, not antioxidant redox cycling. Glycoprotein (Option B) is a structural and membrane-associated component of the lens but has no role in maintaining antioxidants in the reduced state. Fatty acids (Option D): Fatty acids are structural components of lens cell membranes and are actually targets of oxidative damage rather than reducers.

References:

Yanoff and Duker Ophthalmology, 5th Edition, Page 331.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: A 9-year-old child is brought in by his parents because teachers report poor classroom performance. The child says he cannot clearly see the blackboard and struggles with reading. There is no history of white reflex noticed at birth. Ophthalmic examination reveals a bilateral, disc-shaped opacity involving a particular layer of the lens with relatively clear cortex and nucleus. Which type of congenital cataract most commonly presents this way? (Q885171)

A. Cerulean (bluedot) cataract

B. Lamellar (zonular) cataract

C. Posterior polar cataract

D. Total congenital cataract

Correct Answer: B. Lamellar (zonular) cataract

Explanation: This child presents with late detection, scholastic difficulty, and a zonular disc-like opacity, which is classical for Lamellar (Zonular) Cataract , the most common visually significant congenital cataract seen in school-age children. Most common visually significant congenital cataract (~50% of cases). Morphology: Disc-shaped opacity in a specific lens lamella, with clear cortex and nucleus; usually bilateral and symmetrical. Etiology: Mainly genetic (autosomal dominant); can be influenced by intrauterine/infantile malnutrition or vitamin D deficiency. Visual impact: Opacity obstructs vision along the visual axis → axial visual impairment, amblyopia risk. Cerulean cataract (Option A): Multiple small bluish cortical opacities; usually stationary and vision is minimally affected, so children are often asymptomatic. Posterior polar cataract (Option C): Localized posterior capsular opacity. Often small and stationary; may cause glare, but usually not the typical progressive scholastic visual decline unless dense. Total congenital cataract (Option D): Dense white lens present early in infancy, producing leukocoria and severe visual deprivation. Such children present much earlier, often with nystagmus or strabismus, not late school complaints.



References:

Parsons Diseases of the Eye, 23rd Edition, Pages 636-637.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q21: An 18-year-old girl presents with gradual blurring of vision and glare. Slit-lamp examination shows multiple, fine, polychromatic, needle-shaped refractile crystals in the lens cortex, giving a sparkling appearance. Which systemic association is most likely? (Q885175)

A. Alport syndrome

B. Myotonic dystrophy

C. Type 1 diabetes mellitus

D. Lowe syndrome



Correct Answer: B. Myotonic dystrophy

Explanation: Christmas tree cataract is characterized by multicolored, needle-like refractile crystals in the lens cortex, caused by cholesterol and cystine deposition, and is strongly associated with myotonic dystrophy , often appearing in adolescence before systemic neuromuscular symptoms. Disease Characteristic Type of Cataract Myotonic dystrophy Christmas tree cataract (Option B) Wilson disease Sunflower cataract Copper deposition (Chalcosis) Sunflower cataract Penetrating trauma Rosette cataract Diabetes mellitus (Option C ruled out) Snowflake or Snowstorm cataract Down syndrome Stellate or Snowflake or Snowstorm cataract Hypoparathyroidism (Hypocalcemia) Congenital lamellar (zonular) cataract Hypovitaminosis D Congenital lamellar (zonular) cataract Atopic dermatitis Shield cataract (anterior subcapsular) and Posterior subcapsular cataract Congenital rubella syndrome Nuclear cataract Galactosemia Oil drop cataract (reversible) Complicated cataract (Iridocyclitis, high myopia) Posterior subcapsular cataract, Breadcrumb appearance, Polychromatic lustre, Rainbow cataract Blunt trauma Vossius ring and Rosette-shaped cataract Steroid induced Posterior subcapsular cataract Alport syndrome (Option A) is characterised by anterior lenticonus, perimacular dot-and-fleck retinopathy, and renal disease; crystalline polychromatic cataract is not typical. Lowe syndrome (Option D) is associated with congenital cataract, glaucoma, and renal tubular dysfunction; cataracts are dense and developmental, not Christmas-tree type.



References:

Parsons Diseases of the Eye, 23rd Edition, Page 631.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: Match the type of cataract with the etiology Type of cataract Etiology a. Posterior subcapsular 1. Glassblowers b. Anterior subcapsular-granular 2. Miotics c. Posterior cortical 3. Steroids (Q853319)

A. a1,b2,c3

B. a3,b2,c1

C. a2,b1,c3


D. a2,b3,c1

Correct Answer: B. a3,b2,c1

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Type of cataract Etiology Posterior subcapsular Prolonged use of high dose steroids, children are more susceptible Anterior subcapsular-granular Miotics especially long-acting cholinesterase inhibitors such as echothiophate, demecarium bromide, and diisopropyl fluorophosphate (DFP) Posterior cortical Infrared aka glassblower cataract, lens capsule exfoliation with discoid posterior opacities X-ray, gamma rays, neutron exposure: Inadequately protected technicians, patients treated for malignant tumors, and workers of atomic energy plants are at risk. Note: Oral steroids are associated with posterior subcapsular cataracts, while topical steroids commonly elevate intraocular pressure (IOP), potentially leading to steroid-induced glaucoma.



References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 315, 316

Kanski's Clinical Opthalmology, 10th edition, Pg 312, 313

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: What is the consequence of age-related decline in the Na+-K+ ATPase pump in the lens? (Q853310)

A. Increased intralenticular K+ levels and decreased Na+ levels.

B. Increased intralenticular Na+ levels and decreased K+ levels.

C. Enhanced ion exchange efficiency.


D. Reduced intralenticular Ca2+ levels.

Correct Answer: B. Increased intralenticular Na+ levels and decreased K+ levels.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The age-related decline in the Na+-K+ ATPase pump in the lens results in increased intralenticular Na+ levels and decreased K+ levels. Pump leak theory: The mechanism that prevents water from getting into the lens is known as the pump-leak theory. Over the Anterior surface of the lens, Na+-K+ ATPase pumps out Na+ and pumps in K+ into the lens. The leak happens in the posterior surface, i.e., ions (K+) and molecules passively diffuse across a concentration gradient . Enhanced ion exchange efficiency (Option C): Dysfunction of the sodium-potassium ATPase pump leads to decreased ion exchange efficiency, not enhanced efficiency.



@Neet_pg_bot

References:

Yanoff & Duker Ophthalmology 5th edition - page 326-e7, 331.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: Which among the following is/are true statements regarding lens proteins? α/β/γ-Crystallins are water-soluble proteins Major intrinsic proteins contribute to lens transparency An increase in crystallins can lead to cataract β/γ-Crystallin family are the largest crystallins (Q853311)

A. Only 1

B. 1 & 2

C. 2, 3 & 4


D. 1, 2, 3 & 4

Correct Answer: B. 1 & 2

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Lens proteins are broadly divided into soluble fractions and insoluble fractions. The water-soluble fraction consists of crystallins α/β/γ (Statement 1) , which make up about 90% of the lens's total protein content. The α crystallins are the largest crystallins (Statement 4) followed by β and γ respectively. The insoluble fraction of the lens consists of albuminoids - mainly major intrinsic proteins (MIP26/Aquaporin-0) and other cytoskeletal proteins. (Statement 2) A decrease in crystallins and an increase in albuminoids can lead to cataract formation. (Statement 3) Other proteins: HMV-3,4; accumulation of HMV-4 can give rise to nuclear cataract.



References:

Yanoff and Duker Ophthalmology, 6th Edition, Page 307

↑ *Back to this question in Part A*

@Neet_pg_bot


Q25: A 70-year-old male patient comes to your clinic with complaints of glare and diminution of vision for 3 months. On examination, you notice the following appearance Choose the correct statement regarding the correct diagnosis. (Q853312)

A. The Iris shadow test becomes positive only when the whole cortex is involved.

B. Patients see better in dark surroundings.

C. Characterized by saucer-shaped opacities in the center of the posterior cortex.

D. Vision loss occurs at a late stage.



Correct Answer: D. Vision loss occurs at a late stage.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: An elderly patient with clinical features (glare/ diminution of vision/ blurred vision) suggestive of cataract and characteristic wedge-shaped opacities starting from the periphery to the center are noted, which points to a diagnosis of Cuneiform Senile Cortical Cataract. Since it starts peripherally, the visual loss occurs late. Type of Cataract Characteristics Visual Impact Cuneiform Senile Cortical Cataract Wedge-shaped opacities with clear areas in between. Extends from periphery to center . First appears in the lower nasal quadrant. Present in both anterior and posterior cortex. The radial spoke-like pattern of opacities is seen. Visual disturbances are noted at a comparatively late stage (since it starts at the periphery). Vision improves in bright light with contracted pupils. (Option B) Cupuliform Senile Cortical Cataract Saucer-shaped opacities are typically in the central part of the posterior cortex (posterior subcapsular cataract). (Option C) Extends from center to periphery. Causes an early

loss of visual acuity.- Due to the involvement of the nodal point Glares seen in PSC Better vision in dim light with dilated pupils. Iris shadow (Option A) (clear cortex visible) is seen only in immature cataract.

References:

1. Kanski's Clinical Ophthalmology, 10th Edition, Page 310 2. Comprehensive Ophthalmology by AK Khurana, 7th Edition, Page 194-196.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q26: An 80-year-old female patient came with complaints of gradual progressive diminution of vision for 2 years in both her eyes. She reports that over the last year, she noticed an improvement in her ability to read without glasses. The following finding is seen on slit lamp examination over the right eye and the left eye has similar opacity. Choose the most appropriate course of management for this patient. (Q853313)

A. Advise lifestyle modification and review after 3 months.

B. Recommend corticosteroids eye drops and reassess in 6 months.

C. Plan for Phacoemulsification.

D. Prescribe her rigid gas-permeable contact lenses.



Correct Answer: C. Plan for Phacoemulsification.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Gradual, progressive loss of vision along with improvement in near vision (referred to as second sight: due to nuclear sclerosis there is an increase in refractive power of lens which leads to improved near vision) combined with characteristic slit-lamp findings, is suggestive of nuclear cataract , for which phacoemulsification is the treatment of choice. Management: Surgical measures: Mainstay of treatment. Cataract surgeries: ICCE (not preferred now) ECCE SICS (Small Incision Cataract Surgery) Phacoemulsification (most preferred) Followed by IOL implantation: PC-IOL (Posterior chamber-intraocular lens) is preferred. Advise lifestyle modification and review after 3 months (Option A) would not address the cataract directly and would not improve the visual acuity. Recommend corticosteroids eye drops and reassess in 6 months (Option B): The use of topical corticosteroids after cataract surgery reduces postoperative noninfectious inflammation but it is not used as the treatment for nuclear cataract. Prescribe her rigid gas-permeable contact lenses (Option D): Rigid gas-permeable contact lenses are indicated for early-stage keratoconus with astigmatism.

References:

Kanski’s Clinical Ophthalmology, 10th Edition, Page 308

Comprehensive Ophthalmology by AK Khurana, 7th Edition, Pages 195-197, 202

Yanoff & Duker Ophthalmology, 6th Edition, Page 361

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: An elderly male diabetic patient came with loss of vision in the left eye associated with pain in the left eye. On examination, the left eye appears red, and visual acuity- Perception of light+ and IOP is raised in the same eye. The patient gives history of blurring and diminution of vision for the past 1 year. The slit lamp examination is given below. Choose the correct diagnosis and appropriate pathophysiology for this patient. (Q853314)

A. Phacomorphic glaucoma - Secondary open-angle glaucoma

B. Phacomorphic glaucoma - Secondary closed-angle glaucoma

C. Phacolytic glaucoma - Secondary open-angle glaucoma

D. Phacolytic glaucoma - Secondary closed-angle glaucoma



Correct Answer: C. Phacolytic glaucoma - Secondary open-angle glaucoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Given clinical scenario is suggestive of Glaucoma and the slit lamp image of a milky white cortex with a sunken nucleus corresponds to the Morgagnian type of hyper mature cataract The lens protein blocks the trabecular meshwork leading to secondary open-angle glaucoma called Phacolytic glaucoma. Feature Phacomorphic Glaucoma Phacolytic Glaucoma Pathogenesis Intumescent cataract bulges forward, pushing the iris into the angle and causing angle-closure glaucoma. Leakage of lens proteins from a hypermature cataract clogs the

trabecular meshwork. Causes Intumescent cataract (mature cataract) Anterior subluxation of lens Spherophakia Hypermature (Morgagnian) cataract Obstruction of the trabecular meshwork by lens proteins. Clinical Features Similar to acute primary angle closure: shallow AC, ciliary congestion, dilated pupil, swollen lens, and closed angle on gonioscopy. Acute congestive glaucoma: high IOP, deep AC, open angle on gonioscopy.

References:

Kanski's Clinical Opthalmology, 10th edition, Page 393

Comprehensive Ophthalmology by AK Khurana, 7th edition, Pages 199, 255

↑ *Back to this question in Part A*

@Neet_pg_bot

Q28: A 45-year-old man is having difficulty in near vision. The ophthalmologist has prescribed him a convex lens and informed him that it is an age-related physiological condition. Which of the following is an age-related change that would have caused his condition? (Q853315)

A. Decrease in elasticity of lens capsule

B. Sclerosis of lens substance

C. Reduced ciliary muscle power

D. All of the above

Correct Answer: D. All of the above

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical scenario is suggestive of presbyopia, and all the above causes are responsible for development of presbyopia. Presbyopia Physiological insufficiency of accommodation, causing worsening near vision after the age of 40 Causes Decrease in elasticity of lens capsule (Option A) Sclerosis of lens substance (Option B) Reduced ciliary muscle power (Option C) Treatment Convex glasses for near-work A guide for powers required at various ages: 45 years: + l to +l.25D 50 years: +1.5 to +1.75D 55 years: +2 to +2.25D 60 years: +2.5 to +30D

References:

Kanski's Clinical Opthalmology, 10th edition, Pg 393 AK Khurana Comprehensive Ophthalmology 7th edition, pages 46-48

↑ *Back to this question in Part A*

@Neet_pg_bot

Q29: A 6-month-old girl was diagnosed with galactosemia. An ocular examination has revealed a cataract which developed as a complication. Which of the following regarding her cataract is false? (Q853316)

A. Bilateral

B. Oil droplet appearance

C. Periphery of the lens


D. May be reversed

Correct Answer: C. Periphery of the lens

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Galactosemia cataracts are central lens cataracts, not peripheral lens cataracts Galactosemia Inborn error of galactose metabolism Enzyme deficiency Galactose-1-phosphate uridyltransferase (GPUT) Galactokinase (GK) Type of cataract Bilateral (Option A) “Oil droplet” (Option B) Center of the lens Features May be reversed (Option D) May be prevented if milk and milk products are eliminated at an early stage



References:

Yanoff & Duker Ophthalmology, 6th Edition, Page 316, 317, 370

Kanski's Clinical Ophthalmology, 10th Edition, Page 337, 338

↑ *Back to this question in Part A*

@Neet_pg_bot

Q30: Which of the following is not a feature of chalcosis? (Q853317)

A. Kayser-Fleischer ring

B. Sunflower cataract

C. Retinal golden plaques

D. Oil droplet cataract


Correct Answer: D. Oil droplet cataract

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Oil droplet cataract is seen in galactosemia, not chalcosis. Chalcosis: Definition Specific changes produced by the alloy of copper in the eye Mechanism Copper ions from the alloy are dissociated electrolytically and deposited under the membranous structures of the eye Clinical features Kayser-Fleischer ring (Option A) : Golden brown ring due to deposition of copper under the peripheral parts of Descemet's membrane of the cornea Sunflower cataract (Option B) : Deposition of copper under the posterior capsule of the lens Retina: Golden plaques (Option C) at the posterior pole that reflect the light with metallic luster





References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 618 Kanski's Clinical Opthalmology, 10th edition, Pg 927,928

↑ *Back to this question in Part A*

Q31: A 30-year-old male with retinitis pigmentosa has developed a cataract as a complication. Which of the following is a feature of his cataract? Posterior subcapsular cortical cataract Irregular in outline Variable density Breadcrumb appearance Polychromatic luster Diffuse yellow haze in the adjoining cortex (Q853318)

A. 1,2,3,6

B. 2,4,5,6

C. 1,2,3,4


D. 1,2,3,4,5,6

Correct Answer: D. 1,2,3,4,5,6

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Any cataract occurring following an anteroposterior segment disease is a complicated cataract. All of the listed features are those of complicated cataracts. Complicated cataract: Causes: Inflammations (Anterior uveitis- most common) Degenerative conditions such as retinitis pigmentosa and myopic chorioretinal degeneration Long-standing retinal detachment Glaucoma (primary or secondary) Intraocular tumors in late stages (retinoblastoma or melanoma). Features : Starts as a posterior subcapsular cortical cataract (PSC) Irregular outline Variable in density Slit-lamp: Breadcrumb appearance Polychromatic luster, i.e. appearance of iridescent colored particles of reds, greens, and blue Diffuse yellow haze in the adjoining cortex Treatment: Surgical Management: Procedure: Phacoemulsification with Intraocular Lens (IOL) Implantation . This is the gold standard. Capsular Tension Ring (CTR): Because retinitis pigmentosa patients often have zonular weakness , a CTR is frequently inserted into the capsular bag to provide 360-degree support and prevent the lens from shifting or the bag from contracting (phimosis) post-operatively. IOL Choice: Monofocal lenses are usually preferred over multifocal lenses, as the latter can further reduce contrast sensitivity in a retina that is already compromised. Pre-operative Evaluation: Macular Assessment: Before surgery, an OCT (Optical Coherence Tomography) is essential to check for Cystoid Macular Edema (CME) , which is common in RP. If CME is present, it must be treated with topical or systemic carbonic anhydrase inhibitors (like Acetazolamide) before proceeding. Potential Acuity Meter (PAM): Used to estimate how much vision the patient can actually regain, as the "limit" is often the retina, not just the cataract. Post-operative Care: Aggressive Anti-inflammatories: Patients are put on a strict regimen of Topical Steroids and NSAIDs to prevent the development or worsening of macular edema. YAG Laser: These patients have a higher rate of Posterior Capsule Opacification (PCO) , which may require a quick laser procedure months or years later to clear the visual axis.



References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 11, 447, 448

Kanski's Clinical Opthalmology, 10th edition, Pg 631-634

↑ *Back to this question in Part A*

@Neet_pg_bot

Q32: Which of the following is the thinnest part of the lens capsule? (Q853309)

A. Anterior pole

B. Posterior pole

C. Equator


D. Pre-equatorial region

Correct Answer: B. Posterior pole

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Capsule at the posterior pole (4 microns) is the thinnest part. Posterior Subcapsular Cataract (PSC): Site: This cataract develops at the posterior pole , where lens epithelial cells migrate abnormally from the equator to the thin posterior capsule's anterior surface, forming opacities. Patients with PSC complain of glare and disproportionate difficulty reading compared to their distance vision. Risk factors: ■■■■■■■ Steroid Use: This is the most classic association. Both systemic and topical steroids can lead to PSC. Diabetes Mellitus: While diabetics often get early-onset nuclear or cortical cataracts, they are specifically predisposed to PSC. High glucose levels lead to the accumulation of sorbitol, causing osmotic stress. Chronic Intraocular Inflammation: Conditions like Uveitis are a major cause of "complicated" PSC. Ionizing Radiation: The posterior pole is highly sensitive to X-rays or radiotherapy. Anterior pole (Option A): Thicker part of the lens capsule, providing support and maintaining curvature. Equator (Option C): Moderately thick, where zonular fibers attach for lens accommodation. Pre-equatorial region (Option D): The thickest portion of the lens capsule measuring 12-14 micrometers supports zonular attachments.



References:

Yanoff & Duker Ophthalmology, 6th edition, Page 307-e1, e2.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q33: A 35-year-old man presents with decreased vision in his right eye following a blunt trauma a few weeks ago. On examination, feathery opacities are noted. What is

the most likely type of cataract this patient has developed? (Q853320)

A. Subepithelial opacities

B. Diffuse concussion cataract

C. Rosette cataract


D. Morgagnian cataract

Correct Answer: C. Rosette cataract

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Rosette Cataract is characterized by feathery opacities along suture lines, as seen in this patient. A traumatic cataract is an acquired cataract resulting from ocular trauma, such as penetrating or blunt injuries. It forms due to degeneration in previously normal lens fibers. It starts from the posterior cortex Mechanism: Shockwave & Fluid Entry: The force of the blunt impact sends a shockwave through the eye, causing sudden expansion and contraction of the lens. This allows fluid to enter the lens through the capsule. Sutural Separation: The fluid tracks along the lens sutures - the natural anatomical lines where lens fibers meet. Feathery Pattern: As the fibers along these sutures become opacified and swollen, they create a "feathery" or "star-shaped" pattern resembling a flower or a rosette. Other features of blunt trauma:- Cornea: Abrasions, recurrent erosions, partial tears, Descemet's membrane tears, edema, blood staining. Sclera: Partial-thickness wounds. Anterior Chamber: Hyphema, exudates. Iris, Pupil, Ciliary Body: Miosis, mydriasis, margin rupture, iris tears, iridodialysis, aniridia, angle recession, iridocyclitis. Lens: Vossius ring, concussion cataract, absorption, subluxation, dislocation. Vitreous: Liquefaction, opacities, detachment, hemorrhage, herniation. Choroid: Rupture, hemorrhage, detachment, choroiditis. Retina: Commotio retinae, hemorrhages, tears, proliferative retinopathy, detachment, macular changes. Intraocular Pressure: Traumatic glaucoma, hypotony. Refraction: Myopia, hypermetropia, loss of accommodation.



References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 139, 307 e.10 Kanski's Clinical Opthalmology, 10th edition, Pg 914-920

↑ *Back to this question in Part A*

@Neet_pg_bot

Q34: A 38-year-old patient with diabetes presents with rapidly worsening vision. Upon examination, the ophthalmologist notes the presence of bilateral, snowflake-like white opacities in the cortex of the lens. Which of the following is the most likely diagnosis? (Q853321)

A. Senile cataract

B. Christmas tree cataract

C. Diabetic cataract


D. Posterior subcapsular cataract

Correct Answer: C. Diabetic cataract

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Diabetic cataract (also known as presenile cataract) is characterized by bilateral, snowflake-like white opacities in the lens cortex, typically seen in young adults with diabetes. Opacities may either resolve spontaneously or progress to maturity within days. Occurs due to sorbitol accumulation. Commonly seen in Type 1 diabetes. Mechanism: The Polyol Pathway When blood sugar is high, glucose in the lens is converted into Sorbitol by the enzyme Aldose Reductase . Sorbitol cannot escape the lens, creating an osmotic gradient that pulls water in, causing the lens fibers to swell and burst. Senile cataract (Option A) typically affects older individuals, although it can occur earlier in diabetics; however, it does not present with snowflake-like opacities. Christmas tree cataract (Option B) is associated with myotonic dystrophy, not diabetes. Posterior subcapsular cataract (Option D) can be seen in various conditions, including myotonic dystrophy, but does not present with the snowflake-like opacity as seen in true diabetic cataract.



References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 310

Kanski’s Clinical Opthalmology, 10th edition, Pg 503

↑ *Back to this question in Part A*

@Neet_pg_bot

Q35: Which of the following steps is used to maintain the depth of the anterior chamber and protect the corneal endothelium during Conventional ECCE? (Q853323)

A. Anterior capsulotomy

B. Viscoelastic injection

C. Corneoscleral section completion


D. Hydrodissection

Correct Answer: B. Viscoelastic injection

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Injecting a viscoelastic substance into the anterior chamber to maintain its depth and protect the corneal endothelium during the surgery. Protects the corneal endothelium. In addition, irrigation of fluids continuously during the procedure helps maintain the anterior chamber depth. Steps of ECCE: Anterior Capsulotomy (Option A) : Cutting the anterior capsule. Methods include can-opener, linear capsulotomy, envelope, or CCC. Corneoscleral Section Completion (Option C) : Extending the corneal incision. Hydrodissection (Option D) : Separates the lens from the capsule.



Surgical steps of conventional extracapsular cataract extraction with posterior chamber intraocular lens implantation: A) Anterior capsulotomy can-opener's technique; B) Removal of anterior capsule; C) Completion of corneoscleral section; D) Removal of nucleus (pressure and counter- pressure method); E) Aspiration of cortex; F) Insertion of inferior haptic of posterior chamber IOL; G) Insertion of superior haptic of PCIOL; H) Dialing of the IOL; I) Corneoscleral suturing.

@Neet_pg_bot

References:

Yanoff and Duker Ophthalmology, 6th Edition, Pages 336, 337

Kanski's Clinical Ophthalmology, 10th Edition, Pages 321, 322

↑ *Back to this question in Part A*

@Neet_pg_bot

Q36: A 70-year-old woman presents for cataract surgery in her right eye. She has moderate cataract density with some nuclear sclerosis and a clear cornea. She expresses a strong desire for a procedure that minimizes recovery time and provides optimal visual outcomes. Which of the following statements accurately reflects the advantage of Phacoemulsification Over Manual SICS? (Q853324)

A. Manual SICS typically requires a smaller incision compared to Phacoemulsification.

B. Manual SICS allows for faster visual recovery compared to Phacoemulsification.

C. Manual SICS is less suitable for very hard cataracts compared to Phacoemulsification.

D. Manual SICS usually results in a higher level of postoperative astigmatism compared to Phacoemulsification.

Correct Answer: D. Manual SICS usually results in a higher level of postoperative astigmatism compared to Phacoemulsification.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Manual SICS usually results in a higher level of postoperative astigmatism compared to Phacoemulsification as they use larger incisions. Aspect Manual SICS Phacoemulsification Incision (Option A) Longer incision (5-5.5mm) Smaller incision (2.75 -3.2mm) Suitability (Option C) Suitable for all cataract densities, including very hard cataracts Typically requires a smaller incision, less suitable for very hard cataracts Visual Recovery (Option B) Slower visual recovery, delayed return to optimal vision Faster visual rehabilitation due to smaller incision and reduced tissue manipulation Astigmatism (Option D) Higher levels of surgically induced astigmatism, larger incision Less postoperative astigmatism due to smaller incision size

References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 308 e1,2,3 Kanski’s Clinical Opthalmology, 10th edition, Pg 321 AK Khurana Comprehensive Ophthalmology 7th edition, page 205, 208-212

↑ *Back to this question in Part A*

@Neet_pg_bot

Q37: What are the steps involved in FLACS? (Q853325)

A. Capsulorrhexis → Corneal Incision → Lens Fragmentation → IOL implantation

B. Corneal Incision → Capsulorrhexis → Lens Fragmentation → IOL implantation

C. Capsulorrhexis → Lens Fragmentation → Corneal Incision → IOL implantation


D. Corneal Incision → Lens Fragmentation → Capsulorrhexis → IOL implantation


Correct Answer: C. Capsulorrhexis → Lens Fragmentation → Corneal Incision → IOL implantation

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Laser-Assisted Surgical Steps: Post-Laser Surgical Steps : After the laser portion, the patient is moved to an operating microscope where the surgeon completes the remaining steps:





References:

https://journals.lww.com/jcrs/abstract/2024/10000/comparison_of_low_energy_flacs_and_conventional.14.aspx
Yanoff & Duker Ophthalmology, 6th edition, Pg 353 Kanski's Clinical Opthalmology, 10th edition, Pg 324, 326

↑ *Back to this question in Part A*

@Neet_pg_bot

Q38: A key advantage of peribulbar anesthesia over retrobulbar anesthesia is: (Q853326)

A. Faster onset of anesthesia

B. Less risk of globe perforation

C. Better akinesia of the eye


D. Reduced risk of systemic toxicity

Correct Answer: B. Less risk of globe perforation

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Peribulbar anesthesia is associated with a lower risk of globe perforation because the anesthetic is injected outside the muscle cone, reducing the likelihood of accidentally penetrating the eye. Features Retrobulbar block Peribulbar block Injection Site Inside the muscle cone (intraconal) Outside the muscle cone (extraconal) Onset of Anesthesia Faster onset (Option A) Slower onset Akinesia (Eye Movement) Better akinesia due to direct nerve blockade (Option C) Variable akinesia,may require multiple injections Risk of Globe Perforation and Optic nerve injury Higher risk due to proximity to the globe and optic nerve Lower risk Hemorrhage Risk Higher risk of retrobulbar hemorrhage Lower risk of severe hemorrhage Reduced risk of systemic toxicity (Option D) is generally low in both techniques, but peribulbar anesthesia does not significantly reduce this risk compared to retrobulbar anesthesia.



↑ *Back to this question in Part A*

@Neet_pg_bot

Q39: Toric IOLs are specifically designed to correct which type of refractive error? (Q853327)

A. Myopia

B. Hypermetropia

C. Astigmatism

D. Presbyopia

Correct Answer: C. Astigmatism

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Astigmatism is a common refractive error where refraction changes in different meridians of the eye. Toric IOLs have an integral cylindrical refractive component to compensate for pre-existing corneal astigmatism and improve vision clarity for patients with astigmatism. Note: Toric IOL: Designed to correct astigmatism (Option C) Aspheric IOL: Designed to reduce spherical aberrations Aniridia IOL: Designed to cover defects of aniridia or partial iris loss. Myopia (Option A): Not specifically corrected by Toric IOLs. Standard monofocal IOLs are typically used for correcting myopia. Hypermetropia (Option B) : Can be corrected with standard IOLs, but Toric IOLs are not designed specifically for this condition. Presbyopia (Option D) : Age-related condition where the eye loses its ability to focus on near objects. Multifocal or EDOF IOLs are typically used to address presbyopia, not Toric IOLs.

References:

www.ncbi.nlm.nih.gov/books/NBK599976/

www.ncbi.nlm.nih.gov/books/NBK570585/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q40: A 55■year■old man underwent myopic LASIK 15 years ago. He now develops senile cataract and undergoes phacoemulsification with posterior chamber IOL implantation. Postoperatively, he complains of poor distance vision, and refraction shows an unexpected hyperopic error of +2.5 D. Which of the following IOL power calculation formulas would have prevented this postoperative refractive error? (Q853328)

A. SRK

B. Hoffer Q

C. Haigis

D. Holladay I

Correct Answer: C. Haigis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Haigis is a 4th generation formula Calculation is not dependent on corneal parameters In LASIK: Corneal curvature is altered The cornea becomes thinner Hence, formulas relying on anterior chamber depth + axial length (like Haigis) are preferred Patients who have undergone prior myopic LASIK are notorious for "hyperopic surprises" after cataract surgery. This occurs for two primary reasons: Overestimation of Corneal Power (K): Standard keratometry measures the anterior corneal surface and assumes a fixed ratio to the posterior surface. LASIK changes this ratio, leading to a K value that is higher than the actual refractive power. Incorrect Effective Lens Position (ELP): Most formulas use the measured corneal power to predict where the IOL will sit in the eye. Since the post-LASIK cornea is flatter, the formula "assumes" a shallow anterior chamber and predicts the IOL will sit more anteriorly than it actually does. Formulas used: SRK (Option A) Hoffer Q (Option B) Holladay I (Option D) P= A-2.5L-0.9K (L= axial length) (K= constant) Normal axial length (24 mm) Uses K and AL Poor performance Post-LASIK Prone to significant hyperopia. Short axial length (<22mm) Uses K and AL Poor performance Post-LASIK Generally better for very short eyes, not post-LASIK. Long axial length (>24.5 mm) Uses K and AL Poor performance Post-LASIK; Tends to result in hyperopia in flat corneas.

References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 216

Kanski's clinical ophthalmology, 9th edition, Page 316

www.ncbi.nlm.nih.gov/books/NBK589643/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q41: In which scenario is a posterior capsule tear more commonly observed during cataract surgery? (Q853329)

A. During phacoemulsification with high energy settings

B. During cortical clean-up with a manual irrigation-aspiration technique

C. During hydrodelineation

D. During anterior chamber maintenance with a viscoelastic substance

Correct Answer: B. During cortical clean-up with a manual irrigation-aspiration technique

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Posterior capsule tears are more common during the cortical clean-up phase of cataract surgery. This is due to the mechanical stress and manipulation involved in removing residual cortex, which can sometimes lead to inadvertent damage to the posterior capsule. Can occur in following situations During forceful hydrodissection. Direct injury with Sinskey's hook, chopper. During cortex aspiration. Manual irrigation-aspiration technique. During phacoemulsification with high energy settings (Option A) : They are more likely to cause issues like

corneal edema or endothelial damage rather than directly causing a posterior capsule tear. During hydrodelineation (Option C) : This technique generally helps separate the lens from the capsule and is less likely to cause a tear if performed correctly. During anterior chamber maintenance with a viscoelastic substance (Option D) : Maintaining the anterior chamber with viscoelastic substances is designed to protect intraocular structures and reduce complications.

References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 218 & 219. Kanski's clinical ophthalmology, 9th edition, Page 316

↑ *Back to this question in Part A*

@Neet_pg_bot

• [02_Diseases of Lens] 004 Previous Year Questions

« Back to Folder Directory

[02_Diseases of Lens] 004 Previous Year Questions — Quick Jump

Q1 (p. 539)	*Q2 (p. 541)*	*Q3 (p. 542)*	*Q4 (p. 543)*	*Q5 (p. 545)*	*Q6 (p. 546)*
Q7 (p. 547)	*Q8 (p. 548)*	*Q9 (p. 549)*	*Q10 (p. 550)*	*Q11 (p. 550)*	*Q12 (p. 553)*
Q13 (p. 555)	*Q14 (p. 560)*	*Q15 (p. 561)*	*Q16 (p. 562)*	*Q17 (p. 562)*	*Q18 (p. 563)*
Q19 (p. 564)	*Q20 (p. 566)*	*Q21 (p. 567)*	*Q22 (p. 568)*	*Q23 (p. 570)*	*Q24 (p. 571)*
Q25 (p. 572)	*Q26 (p. 574)*	*Q27 (p. 574)*			


Q1: A 65-year-old diabetic woman presents with painless loss of vision. The ocular findings are shown in the image below. What is the likely diagnosis and management for this patient? (Q855468)

A. Zonular cataract and lensectomy

B. Rosette cataract and sugar control


C. Sunflower cataract and lesion extraction

D. Immature senile cataract and phacoemulsification with IOL



Correct Answer: D. Immature senile cataract and phacoemulsification with IOL

Explanation: The patient is a 65-year-old diabetic woman with painless loss of vision , which points towards cataracts , a common cause of visual impairment, especially in older individuals and those with diabetes . Immature senile cataract refers to the early stages of cataract formation, where the lens becomes cloudy but is not yet completely opaque. This typically leads to gradual loss of vision without pain. The standard treatment for cataracts, especially in cases with significant visual impairment, is phacoemulsification , a surgical procedure that uses ultrasound to break up the lens, followed by removal and placement of an intraocular lens (IOL) to restore clear vision. Type of Cataract Characteristics Visual Impact Cuneiform Senile Cortical Cataract Wedge-shaped opacities with clear areas in between. Extends from periphery to center . First appears in the lower nasal quadrant. Present in both anterior and posterior cortex. The radial spoke-like pattern of opacities is seen. Visual disturbances are noted at a comparatively late stage (since it starts at the periphery). Vision improves in bright light with contracted pupils. Cupuliform Senile Cortical Cataract Saucer-shaped opacities are typically in the central part of the posterior cortex (posterior subcapsular cataract). Extends from center to periphery. Causes an early loss of visual acuity.- Due to the involvement of the nodal point Glares seen in PSC Better vision in dim light with dilated pupils. Stages of Cortical cataract: Stage Description Characteristics 1. Stage of Lamellar Separation Earliest stage of senile cataract Formation of vacuoles and water clefts within the cortex Reversible change Can only be demonstrated by slit lamp examination 2. Stage of Incipient Cataract Detectable opacities with clear areas present between. Causes visual disturbances. Cuneiform type: Wedge-shaped opacities, extending from the equator appear as radial spoke-like opacities. Cupuliform type: Saucer-shaped opacity in the central posterior cortex. 3. Immature Senile Cataract Progression of opacification Iris shadow visible (clear cortex visible) -Iris Shadow present here due to opacities + clear areas Can progress to Intumescent Cataract leading to the shallow anterior chamber. 4. Mature Senile Cataract (aka intumescent cataract) Complete opacification of cortex The lens appears pearly white Referred to as a "ripe cataract" No iris shadow as the whole lens is opacified with a chalky white color 5. Hypermature Senile Cataract (HMSC) Advanced stage of cataract beyond maturity Morgagnian Hypermature Cataract: Cortex liquefies Nucleus sinks to the bottom Bag of milky fluid appearance {{caption_text}} Sclerotic Hypermature Cataract: The cortex disintegrates, lens and capsule shrink. Deep anterior chamber Iridodonesis may be present. Incorrect Options: Zonular cataract and lensectomy (Option A) : Zonular cataract is a specific type of cataract involving changes to the lens fibers and zonules (the ligaments that hold the lens in place). While lensectomy (removal of the lens) can be performed, it is typically not the preferred term for routine cataract surgery, and this condition is not as commonly associated with the age and symptoms described here. Rosette cataract and sugar control (Option B) : A rosette cataract is a rare type of cataract that forms as a star-shaped pattern, usually associated with trauma or certain diseases, but it is not typically seen in diabetic patients or with the presentation described here. Sugar control is important for diabetes management, but it doesn't directly address the cataract-related vision loss. Sunflower cataract and lesion extraction (Option C) : A sunflower cataract is typically associated with Wilson's disease , a condition characterized by copper accumulation in tissues, leading to a distinctive appearance of the cataract. The patient does not have signs of Wilson's disease, and lesion extraction would not be appropriate for a cataract.



References:

Kanski's Clinical Ophthalmology, 10th edition, Pg 310

Comprehensive Ophthalmology by AK Khurana, 7th edition, Pg 194-196.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Riya was found to have a height of more than the 95th percentile for her age and an arm-span to height ratio of more than 1.05. On further examination, she had flat feet, mild scoliosis, and long slender fingers. An image of her showing characteristic signs is provided below. What is the most common ocular complication expected in this child? (Q860007)

A. Hypertensive retinopathy

B. Rosette cataract

C. Retinitis pigmentosa

D. Subluxation of lens



Correct Answer: D. Subluxation of lens

Explanation: The clinical scenario of tall stature flat feet, mild scoliosis, and long slender fingers in suggestive of Marfan's syndrome. Marfan syndrome is commonly associated with mutations in the FBN1 gene, which encodes the connective protein fibrillin. Autosomal dominant inheritance. Multisystem disorder with involvement of the connective tissues. Clinical features: Skeletal abnormalities: Tall stature Arachnodactyly Chest wall abnormalities like pectus excavatum/Pectus carinatum Thumb sign(On folding thumb, it will protrude beyond the ulnar border of palm) or wrist sign is positive Increased joint laxity CVS abnormalities: Aortic root dilation Mitral insufficiencies Arrhythmias Eye abnormalities: Ectopia lentis or subluxation of the lens of the eye (Option D) Blindness Pulmonary involvement: Increased risk of pneumothorax Diagnosis: Marfan syndrome is clinically diagnosed using Ghent criteria

References:

GHAI Essential Pediatrics, 10th edition pg no: 706

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: What is the embryological origin of the corneal endothelium? (Q859756)

A. Neuroectoderm

B. Neural crest cells

C. Mesoderm

D. Surface ectoderm

Correct Answer: B. Neural crest cells

Explanation: The corneal endothelium originates from neural crest cells, which migrate during embryogenesis to form the inner layer of the cornea. Neural crest cells also contribute to other anterior segment structures, including the stroma and trabecular meshwork. Corneal Development and Cellular Origins Corneal development involves ectodermal-derived tissues. Neural ectoderm-derived optic vesicles interact with cranial ectoderm → form presumptive corneal epithelium and ocular surface epithelia. (Option A ruled out) Neural tube-ectoderm interactions generate neural crest cells, which are migratory and multipotent. Neural crest cells migrate between the lens and corneal epithelium → form corneal endothelium and stromal keratocytes. Trigeminal ganglion (from neural crest and ectodermal placodes) supplies sensory innervation to the corneal stroma and epithelium. Mesoderm (Option C): The mesoderm is responsible for forming structures like muscles, blood vessels, and connective tissue in the body. It does not contribute to the corneal endothelium. Surface ectoderm (Option D): The surface ectoderm forms the corneal epithelium, the outermost layer of the cornea, but not the corneal endothelium.

References:

Classification of corneal endothelial disorders based on neural crest origin - PubMed

Vision - Cornea Development - Embryology

↑ *Back to this question in Part A*

@Neet_pg_bot


Q4: Identify the given instrument (Q860184)

A. Universal retractor

B. Doyen’s speculum


C. Mouth clamp

D. Eye speculum



Correct Answer: D. Eye speculum

Explanation: An eye speculum is a surgical instrument used to hold the eyelids apart during ophthalmic procedures. It ensures proper exposure of the eye, facilitating examination or surgery, like enucleation. Eye speculum Doyen’s speculum (Option B) used in Gynaecology Universal retractor (Option A) and Mouth clamp (Option C) are used in Dentistry



References:

https://www.ncbi.nlm.nih.gov/books/NBK562144/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: Which of the following best defines blindness ? (Q865357)

A. Visual acuity less than 6/60 in the better eye

B. Visual acuity less than 3/60 in the better eye after best possible correction

C. Visual field less than 10 degrees from fixation in the better eye

D. Inability to perceive light in both eyes

Correct Answer: B. Visual acuity less than 3/60 in the better eye after best possible correction

Explanation: Correct Option B -Visual acuity less than 3/60 in the better eye after best possible correction Visual acuity less than 6/60 in the better eye (Option A ruled out): This level of visual acuity (or 20/200 in feet) is often referred to as legal blindness in many countries, including the United States, and is a criterion for eligibility for certain services. However, the international definition of blindness is typically more severe. Visual acuity less than 3/60 in the better eye after best possible correction (Option B): This is the definition of blindness according to the

World Health Organization (WHO) and is widely adopted for international comparison. 3/60 means a person can see at 3 meters what a person with normal vision can see at 60 meters. This is the best definition of blindness among the choices. Visual field less than 10 degrees from fixation in the better eye (Option C ruled out): A restricted visual field can also define blindness, particularly "tunnel vision." The WHO definition of blindness includes "or corresponding visual field loss," which for blindness is typically a visual field of 10 degrees or less from fixation in the better eye. This is also a valid criterion for blindness. Inability to perceive light in both eyes(Option C ruled out): This is known as "No Light Perception" (NLP) and represents total blindness. While this is indeed a form of blindness, it is the most severe form. The general definition of "blindness" encompasses a broader range of severe visual impairment where some light perception or very limited vision might still be present.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: What is the diagnosis? (Q856030)

A. Intraocular foreign body

B. Pseudoexfoliation syndrome

C. Ocular trauma

D. Vossius ring



Correct Answer: B. Pseudoexfoliation syndrome

Explanation: - Pseudoexfoliation syndrome: Pseudoexfoliation syndrome is a condition where an abnormal protein material (pseudoexfoliative material) accumulates in the anterior segment of the eye, especially on the lens capsule , pupillary margin , and the corneal endothelium . This condition can cause the formation of a characteristic white, flaky material around the lens and pupil , which is the hallmark of pseudoexfoliation syndrome. It can lead to increased risk of glaucoma (due to blocked drainage of fluid), cataracts , and other complications. This can lead to phacodonesis (lens wobbling). Ocular manifestation of Pseudoexfoliation Syndrome (XFS) is characterized by the accumulation of fibrillar, proteinaceous material on various ocular structures, including the lens, iris, and trabecular meshwork. This material can be seen as flaky white deposits on the anterior lens capsule and pupillary margin, often with a distinct clear zone at the center of the lens. It causes secondary open-angle glaucoma, known as pseudoexfoliative glaucoma. The excessive material and pigment clog the trabecular meshwork, increasing intraocular pressure (IOP) and causing damage to the optic nerve, which leads to glaucoma. It also predisposes patients to complications during cataract surgery due to weakened zonules, resulting in lens instability or phacodonesis. Features of Pseudoexfoliation Syndrome: White flaky deposits on the anterior lens capsule and iris Increased trabecular meshwork pigmentation Elevated IOP, leading to secondary glaucoma Zonular instability and risk of lens dislocation during cataract surgery Mutations in the LOXL1 gene have been strongly associated with PXF, although not all individuals with these mutations develop the disease. More prevalent in elderly patients. Intraocular foreign body (Option A) : This typically causes localized symptoms such as pain, vision changes , or infection . An image of an intraocular foreign body would typically show the object within the eye, which is not characteristic of the findings in pseudoexfoliation syndrome. Ocular trauma (Option C) : Ocular trauma may lead to various findings such as bruising, swelling , or damage to the structures of the eye . It is not characterized by the deposition of material on the lens and anterior segment like in pseudoexfoliation syndrome. Vossius ring (Option D) : This is a pigmented ring on the cornea that results from trauma (usually from a blow to the eye). It is a permanent imprint of the iris pigment on the corneal endothelium and is not related to the findings of white, flaky material seen in pseudoexfoliation syndrome.

References:

Yanoff and Duker's Ophthalmology, 5th edition, Pages 1074-1077

Kanski Clinical Ophthalmology, 10th edition, Pages 382, 383, 384

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: What is the average degree of hypermetropia in a newborn? (Q859755)

A. +2.5 D

B. +5 D

C. +10 D

D. +1 D

Correct Answer: A. +2.5 D

Explanation: Newborns are physiologically hypermetropic, with an average refractive error of approximately +2.5 to 3 diopters. This occurs because the axial length of the eye is shorter at birth, and the eye grows over time, reducing hypermetropia. Hypermetropia, or long-sightedness, is a refractive error in which parallel rays of light from distant objects focus behind the retina when the eye's accommodation is relaxed. Type of Hypermetropia Cause Axial Hypermetropia Shortening of the eyeball's axial length. Commonest form. Curvatural Hypermetropia Flatter curvature of the cornea, lens, or both. Index Hypermetropia Decreased refractive index of the lens, usually due to cortical sclerosis in old age or in treated diabetics. Positional Hypermetropia Posterior displacement of the crystalline lens. Aphakia Absence of the crystalline lens, either congenital or acquired (e.g., after surgery), leads to high hypermetropia. Consecutive Hypermetropia Overcorrected myopia after refractive surgery (e.g., LASIK, ICL) or underpowered intraocular lens (IOL) after cataract surgery. Features: Small eyeball in high hypermetropia Smaller cornea Shallow anterior chamber Fundus Examination: Optic disc - Small , ill-defined margins, possibly simulating papillitis (pseudopapillitis). Retina - Shot silk appearance (shining retina). +1 D (Option D): +1 D is below the average hypermetropia seen in newborns. This degree of refractive error is more commonly observed in older children or adults after emmetropization has occurred.

References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 28-31

Hyperopia - StatPearls - NCBI Bookshelf

Refractive errors in full-term newborn babies

https://pmc.ncbi.nlm.nih.gov/articles/PMC2636866/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: What could be the most probable cause for the progressive vision loss in both eyes of a 25-year-old male who has been undergoing low-dose oral steroid therapy for the past decade? (Q859416)

A. Cataract

B. Glaucoma

C. Cystoid macular edema

D. Retinal detachment

Correct Answer: A. Cataract

Explanation: The most likely reason for the progressive loss of vision in a 25-year-old male who has been on low-dose oral steroid therapy for the last 10 years is Cataract. Toxic cataract: Steroids - Prolonged use in high doses may cause posterior subcapsular cataracts. Children are more susceptible Other causes of posterior subcapsular cataract: Galactosemia, Myotonic dystrophy, BLUNT trauma, complicated cataract, chloroquine,

busulfan, NF2, Glaucoma by TOPICAL steroids (Option B) and cataract by SYSTEMIC STEROIDS (mnemonic: GTCS) Miotics - Especially long-acting cholinesterase inhibitors such as echothiophate, demecarium bromide, and diisopropyl fluorophosphate (DFP) may cause anterior subcapsular-granular cataracts Other causes of anterior subcapsular cataract - Amiodarone, chlorpromazine, busulfan, gold, and allopurinol AACG (Mnemonic): Atopic dermatitis- shield cataract, Amiodarone, Chlorpromazine, Gold deposit, AACG(acute angle closure glaucoma) Incorrect Options Glaucoma (Option B) is the most common complication of topical steroids. Cystoid macular edema (Option C) may be associated with steroid use, but it typically presents with symptoms such as blurry vision, central vision loss, and distortion. Retinal detachment (Option D) causes progressive loss of vision which is not consistent with the above described case.

References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 315, 316

Kanski's Clinical Opthalmology, 10th edition, Pg 312, 313

AK Khurana Comprehensive Ophthalmology 7th edition, page 201

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: True statement regarding KF ring: Seen in most patients with neurological involvement Is pathognomonic for Wilson's disease Resolves with desferroxamine Seen in all patients with hepatic involvement Seen in superior and inferior descemets membrane (Q856412)

A. 1 and 5

B. 1,2 and 5

C. 1,2,3,4 and 5

D. 2,3,4 and 5

Correct Answer: A. 1 and 5

Explanation: The Kayser–Fleischer (KF) ring is caused by copper deposition in Descemet's membrane of the cornea in Wilson's disease . It appears as a golden-brown to greenish ring , best seen on slit-lamp examination . (1) Seen in most patients with neurological involvement – True KF rings are present in >95% of patients with neurological Wilson's disease . (2) Is pathognomonic for Wilson's disease – False ■■■■■■■■ While highly suggestive, KF rings are not pathognomonic , as similar deposits can rarely occur in chronic cholestatic liver diseases . (3) Resolves with desferrioxamine – False ■■■■■■■■ Desferrioxamine is an iron chelator , not a copper chelator. KF rings resolve with penicillamine or trientine therapy (copper chelators). (4) Seen in all patients with hepatic involvement – False ■■■■■■■■ KF rings are seen in ~50–60% of hepatic cases, not all. (5) Seen in superior and inferior Descemet's membrane – True ■■■■■■■■■■■■■■■■ The rings initially appear superiorly

and inferiorly , where aqueous humor circulation is slowest , before eventually encircling the cornea.

References:

Nelson textbook of pediatrics, ed.21, page no 2103-04

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: What is the formula utilized to calculate the power of the lens during cataract surgery for a patient who has previously undergone LASIK for myopia? (Q856929)

A. SRK 1

B. SRK 3

C. Hoffer Q

D. Haigis L

Correct Answer: D. Haigis L

Explanation: SIK eyes. It uses two variables: axial length & anterior chamber depth, making it more adaptable to eyes that have undergone refractive surgery. This adaptability generally provides more accurate IOL power calculations in post-LASIK cases. Biometry: Method of calculating IOL power by 2 methods - Keratometry and Axial length Equipments used: A scan ultrasound for axial length Formulas used: SRK (Option A and B) Hoffer Q (Option C) Holladay I P= A-2.5L-0.9K (L= axial length) (K= constant) Normal axial length (24 mm) Short axial length (<22mm) Long axial length (>24.5 mm)

References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 216

Kanski’s clinical ophthalmology, 9th edition, Page 316

Biometry

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A 60-year-old patient with progressive painless loss of vision presents to the ophthalmology OPD. The fundus image of the patient is given below. What could be the probable finding and the cause of it? (Q856121)

A. Soft exudate, hypertension

B. Hard exudate, diabetes mellitus


C. Flame-shaped hemorrhages, hypertension


D. Soft exudate, central retinal vein occlusion



Correct Answer: B. Hard exudate, diabetes mellitus

Explanation: Hard exudates are bright yellow, well-defined deposits caused by leakage of fluid rich in lipids and proteins from damaged retinal capillaries. They often form around microaneurysms and, in the macular region, can create a star-shaped pattern (macular fan/ star), which is seen most commonly in diabetic retinopathy (although it can also be seen in Hypertensive retinopathy). Hard and Soft Exudates: Feature Soft Exudates (Cotton-Wool Spots) Hard Exudates Appearance Cloud-like, small, ill-defined, whitish patches Bright yellow, well-defined lipid deposits Formation Mechanism Caused by ischemia and axonal swelling; blockage of axoplasmic flow Formed by leakage of fluid rich in proteins and lipids Location Superficial retinal layers, often seen around the optic nerve Seen around leaking microaneurysms, especially in the macula Common Shape Indistinct and variable; may change shape quickly Radial, star-shaped figure in the macula (macular fan/star) Main Causes Hypertensive retinopathy (Most common cause) Diabetic retinopathy Toxemia of pregnancy Lupus erythematosus (LE) Polyarteritis nodosa (PAN) Retinopathy associated with AIDS Diabetic retinopathy (Most common cause) (Option B) Hypertensive retinopathy Coats' disease Circinate retinopathy Image {{caption_text}} {{caption_text}}





References:

Parson’s diseases of the Eye 23rd Edition, Pages 287,313,318,319

Kanski’s Clinical Ophthalmology 9th Edition, Pages 502-503

Yanoff and Duker Ophthalmology 5th Edition, Page 545

↑ *Back to this question in Part A*

@Neet_pg_bot


Q12: Identify the surgical step shown in the image given below: [(Q856716)

A. Hydrodissection

B. Capsulorrhexis

C. Intraocular lens implantation


D. Lens aspiration




Correct Answer: B. Capsulorrhexis

Explanation: The surgical procedure where tryphan blue dye is being injected in the image is Continuous curvilinear Capsulorrhexis . It refers to the creation of a circular tear in the anterior capsule of the lens. Surgical Steps of Conventional ECCE: Eye Stabilization: Superior rectus suture for downward gaze(known as bridle suture) Conjunctival Flap Creation: A fornix-based flap will be used to expose the limbus. Limbal Incision: Partial-thickness groove across the limbus. Anterior Chamber Access: Entry with a razor blade knife or keratome. Viscoelastic Injection: Maintains chamber depth and protects the cornea. Anterior Capsulotomy/ Capsulorrhexis: Methods include can-opener, linear capsulotomy, envelope, or CCC. Tryphan blue dye is usually injected. (Option B) {{caption_text}} Capsule Removal: Using forceps remove the cut capsule. Corneoscleral Section Completion: Extending the incision. Hydrodissection: Separates the lens from the capsule. (Option A) Nucleus Removal: Methods include pressure/counter pressure or irrigating wire vectis. Lens Cortex Aspiration: Using a two-way cannula. (I/A) (Option D) IOL Implantation: Posterior chamber IOL into the capsular bag. (Option C) Incision

Closure: With sutures. Viscoelastic Removal: Followed by refilling the anterior chamber with irrigating fluid. Conjunctival Flap Repositioning: Secured in place. Subconjunctival Injections: Dexamethasone and gentamicin. Eye Patching: Final step.





Surgical steps of conventional extracapsular cataract extraction with posterior chamber intraocular lens implantation: A) Anterior capsulotomy can-opener's technique; B) Removal of anterior capsule; C) Completion of corneoscleral section; D) Removal of nucleus (pressure and counter- pressure method); E) Aspiration of cortex; F) Insertion of inferior haptic of posterior chamber IOL; G) Insertion of superior haptic of PCIOL; H) Dialing of the IOL; I) Corneoscleral suturing.

References:

Kanski's Clinical Ophthalmology 10th edition, Pages 321,322.

AK Khurana Comprehensive Ophthalmology 7th edition, Pages 204,206-208.

↑ *Back to this question in Part A*


Q13: Match the given images to their descriptions: 1. A. Posterior subcapsular cataract 2. B. Sunflower cataract 3. C. Morgagnian cataract 4. D. Snowflake cataract (Q857189)

A. 1A 2D 3B 4C


B. 1D 2B 3A 4C


C. 1B 2C 3D 4A

D. 1C 2B 3D 4A




@Neet_pg_bot


@Neet_pg_bot




Correct Answer: C. 1B 2C 3D 4A

Explanation: Correct match: 1B 2C 3D 4A 1. {{caption_text}} B) Sunflower cataract: Deposition of copper under the posterior capsule of the lens. Cause: Wilson's disease 2. {{caption_text}} C) Morgagnian cataract: Seen in hypermature senile cataract Cortex liquefies Nucleus sinks to the bottom Bag of milky fluid appearance 3. {{caption_text}} D) Snowflake cataract: Also known as snowburst cataract Cause: Diabetes mellitus 4. {{caption_text}} A) Posterior subcapsular cataract: Opacity below the capsule in the central part of the posterior cortex. Causes: Complicated cataract Blunt trauma Myotonic dystrophy (Christmas tree cataract) Glactosemia (Oil droplet cataract) Drugs: Steroid , chloroquine, busulfan


@Neet_pg_bot


@Neet_pg_bot





References:

Kanski’s Clinical Ophthalmology 10th edition, page 310

Comprehensive Ophthalmology by AK Khurana 7th edition, pages 193-201.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: What substance is responsible for the development of a diabetic cataract in a 68-year-old female patient, which has been present for 3 months? (Q858160)

A. Glucose

B. Fructose

C. Lactose + Glucose

D. Sorbitol

Correct Answer: D. Sorbitol

Explanation: - Sorbitol In diabetic patients, sorbitol accumulation in the lens plays a critical role in the development of diabetic cataracts. This process is primarily mediated by the polyol pathway , where glucose is converted to sorbitol via the enzyme aldose reductase . Sorbitol accumulates in the lens because it is poorly permeable across the lens membrane, leading to osmotic stress and cellular damage, which promotes cataract formation. Glucose Metabolism in the Lens: The lens uses glucose through several metabolic pathways: Glycolytic pathway : This is the primary pathway where about 80% of glucose is metabolized anaerobically. Pentose Phosphate pathway (HMP shunt) : Responsible for metabolizing 15% of glucose. Kreb's citric acid cycle : A small proportion of glucose is metabolized through this oxidative pathway. Sorbitol pathway : In healthy individuals, this pathway is minimally active. However, in diabetic and galactosemic patients , excess glucose is converted into sorbitol by aldose reductase . The accumulation of sorbitol leads to osmotic imbalance, which causes lens fiber swelling and opacity, resulting in cataracts. Glucose (Option A) : Although glucose is involved in cataract formation, it is not directly responsible for the development of diabetic cataracts . In the lens, glucose is metabolized via several pathways, but it is the conversion of glucose to sorbitol (via the polyol pathway) that leads to osmotic stress and cataract formation in diabetes. Fructose (Option B) : Fructose does not significantly contribute to diabetic cataracts . While fructose is metabolized in the body, it is not the primary substance implicated in lens opacity in diabetic patients. The key factor in diabetic cataracts is sorbitol accumulation. Lactose + Glucose (Option C) : Lactose is not involved in the formation of diabetic cataracts . Although glucose plays a role in cataract formation, lactose is more relevant in galactosemic cataracts and not in diabetes-induced cataracts.

References:

K Khurana Comprehensive Ophthalmology 7th edition, page 186

↑ *Back to this question in Part A*

@Neet_pg_bot


Q15: An elderly patient presents with white, dandruff-like deposits on the anterior lens surface, seen during slit-lamp examination. What is the most likely diagnosis? (Q866322)

A. Pseudoexfoliation syndrome

B. Glaucoma fleckens

C. Phacolytic glaucoma

D. Pigment dispersion syndrome



Correct Answer: A. Pseudoexfoliation syndrome

Explanation: Pseudoexfoliation syndrome Based on the above findings, the patient has pseudoexfoliation syndrome, characterized by the deposition of whitish-gray, dandruff-like fibrillar material on the anterior lens capsule and pupillary margin. This appearance is also called the hoarfrost sign . The material is best seen after pupil dilation and often appears in a concentric ring pattern. It is strongly associated with secondary open-angle glaucoma due to blockage of the trabecular meshwork by exfoliated material and pigment. Pseudoexfoliation syndrome is an age-related systemic condition with ocular manifestations involving abnormal production and accumulation of extracellular fibrillar material in ocular tissues. This is due to abnormal basement membrane metabolism → deposition of pseudoexfoliative material in anterior segment structures (lens capsule, iris, zonules, ciliary body). Management: Regular IOP monitoring, antiglaucoma medications as needed, and careful cataract

surgery planning. Iris cyst (Option B): Localized iris mass; does not cause diffuse frosted deposits on the lens capsule. Persistent pupillary membrane (Option C): Remnant of tunica vasculosa lentis; appears as thin strands across the pupil, not concentric whitish deposits. Pigment dispersion syndrome (Option D): Pigment granules are deposited on corneal endothelium (Krukenberg's spindle) and trabecular meshwork; lacks the characteristic fibrillar white deposits of pseudoexfoliation.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: What dye is used in staining for the anterior capsule of lens ? (Q820469)

A. Trypan blue

B. Brilliant blue

C. Fluorescent green

D. Lissamine green

Correct Answer: A. Trypan blue

Explanation: Correct Option A - Trypan Blue Trypan blue is a dye that can be used to stain the anterior capsule of the lens during certain ophthalmic surgeries, such as cataract surgery. It helps to increase the visibility of the anterior capsule of the lens, making the surgical procedure easier and safer. Incorrect Options: Option B - Brilliant Blue: This dye is sometimes used in eye surgery, specifically for staining the internal limiting membrane (ILM) during vitrectomy procedures. Option C - Fluorescent green (Fluorescein): This dye is used in diagnostic procedures to highlight damage to the cornea. Option D - Lissamine Green: This dye is used in eye exams to detect damage to the eye's surface, particularly the cornea and conjunctiva.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: In which type of cataract surgery is suturing with 10-0 nylon commonly performed? (Q820459)

A. Phacoemulsification

B. Extracapsular cataract extraction (ECCE)

C. Intracapsular cataract extraction (ICCE)

D. Manual small incision cataract surgery (MSICS)

Correct Answer: B. Extracapsular cataract extraction (ECCE)

Explanation: Correct Option B - Extracapsular cataract extraction (ECCE) In extracapsular cataract extraction (ECCE), the surgeon makes a larger incision on the eye's surface, typically around 10-12 mm in size. This incision allows the surgeon to remove the cataractous lens while leaving the posterior lens capsule intact. After removing the lens, the incision is usually closed with sutures, often using 10-0 nylon. ECCE requires suturing because the incision size is relatively large, and proper wound closure is necessary to ensure the eye heals correctly and maintains its structural integrity. Incorrect Options: Option A - Phacoemulsification: Phacoemulsification is a modern and minimally invasive cataract surgery technique. In this procedure, a small incision is made through which an ultrasonic probe (phacoemulsifier) is inserted to break up the cataract into tiny fragments that are then suctioned out. Suturing with 10-0 nylon is not commonly performed in phacoemulsification surgery because the incision size is typically very small and may not require sutures to close. Option C - Intracapsular cataract extraction (ICCE): Intracapsular cataract extraction involves removing the entire lens, including the lens capsule, in one piece. This method is now less commonly used due to advances in surgical techniques. Suturing with 10-0 nylon is not commonly performed in ICCE surgery because the procedure involves removing the entire lens and capsule, making the need for suturing less relevant. Option D - Manual small incision cataract surgery (MSICS): Manual small incision cataract surgery is a technique similar to ECCE but uses a smaller incision (around 6-7 mm) compared to traditional ECCE. While suturing may still be performed in MSICS, the use of 10-0 nylon is more commonly associated with traditional ECCE due to the larger incision size.

References:

..

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: Which lens is utilized in direct gonioscopy? (Q857414)

A. Goldmann

B. Koeppe

C. Zeiss

D. Sussman

Correct Answer: B. Koeppe

Explanation: In direct gonioscopy , the Koeppe lens is commonly used. It is designed with a domed or slanted shape that allows light rays to pass through the contact lens/air interface at a steep angle, enabling direct visualization of the anterior chamber angle. Direct Gonioscopy: Direct gonioscopy uses a lens with a domed or slanted configuration to allow light from the eye's angle to pass through the lens to the observer without reflection. It does not require a slit lamp and is typically performed with the patient in a supine position , often under general anesthesia for procedures like infantile glaucoma evaluation and surgery. Lenses Used: Includes Koeppe, Barkan, Swan-Jacob lenses. Technique: Performed with the patient lying down, often using a hand-held microscope or magnifying loupes. Not compatible with desktop slit lamps, limiting clarity and magnification compared to indirect gonioscopy. Indications: Commonly used for evaluating and treating infantile glaucoma. Goldmann lens : Typically used for indirect gonioscopy , which involves using a slit lamp for better magnification and clarity. Zeiss lens : Also used for indirect gonioscopy with a slit lamp, offering high magnification. Sussman lens : A type of indirect gonioscopy lens, offering wide-angle visualization but not used in direct gonioscopy.

References:

Kanski's Clinical ophthalmology, 9e., page 32-34

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: Which is the type of cataract seen following a blunt trauma? (Q860165)

A. Rosette cataract

B. Christmas tree cataract


C. Snowflake cataract


D. Sunflower cataract


Correct Answer: A. Rosette cataract

Explanation: Traumatic cataract: blunt trauma: Rosette cataract (Option A) It is a cortical cataract that starts from the posterior cortex. {{caption_text}} Toxic cataract: Steroids - Prolonged use in high doses may cause posterior subcapsular cataracts. Children are more susceptible Other causes of posterior subcapsular cataract: Galactosemia, Myotonic dystrophy, BLUNT trauma, complicated cataract, chloroquine, busulfan, NF2, Glaucoma by TOPICAL steroids and cataract by SYSTEMIC STEROIDS (mnemonic: GTCS) Miotics - Especially long-acting cholinesterase inhibitors such as echothiophate, demecarium bromide, and diisopropyl fluorophosphate (DFP) may cause anterior subcapsular-granular cataracts Other causes of anterior subcapsular cataract - Amiodarone, chlorpromazine, busulfan, gold, and allopurinol AACG (Mnemonic): Atopic dermatitis- shield cataract, Amiodarone, Chlorpromazine, Gold deposit, AACG(acute angle closure glaucoma) Radiation cataract: Infrared (heat) cataract - {{caption_text}} Infrared aka glassblower cataract Lens capsule exfoliation with discoid posterior opacities X-ray, gamma rays, neutron exposure: Inadequately protected technicians, patients treated for malignant tumors, and workers of atomic energy plants are at risk. Electric Shock cataract/ Lightening cataract: It leads to Stellate ASC {{caption_text}} Incorrect Options: Christmas tree cataract (Option B) is associated with Myotonic

Dystrophy. Snowflake cataract (Option C) Diabetes mellitus is associated with Snowflake cataract Sunflower cataract (Option D) is associated with Wilson’s disease







References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 315, 316

Kanski's Clinical Ophthalmology, 10th edition, Pg 312, 313

AK Khurana Comprehensive Ophthalmology 7th edition, page 201.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: Oil drop cataract is seen in? (Q860164)

A. Galactosemia

B. Wilson's disease


C. Gaucher's disease

D. Tay-Sachs disease

Correct Answer: A. Galactosemia

Explanation: Galactosemia cataracts cause oil drop cataract. Galactosemia Inborn error of galactose metabolism Enzyme deficiency Galactose-1-phosphate uridyltransferase (GPUT) Galactokinase (GK) Type of cataract

Bilateral “Oil droplet” (Option A) Central lens {{caption_text}} Features May be reversed May be prevented if milk and milk products are eliminated at an early stage Wilson's disease (Option B) primarily affects the liver and brain, and produces sunflower cataracts and Kayser-Fleischer ring in the cornea. Gaucher's disease and Tay-Sachs (Option C and D) disease are lysosomal storage disorders and are not associated with oil drop cataracts.



References:

Yanoff & Duker Ophthalmology, 6th edition, Page 316, 317, 370

Kanski’s Clinical Ophthalmology, 10th edition, Pg 337, 338

AK Khurana Comprehensive Ophthalmology 7th edition, page 200

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: After cataract surgery, a patient develops endophthalmitis with vision limited to hand movements. What is the treatment? (Q860543)

A. Intravitreal antibiotic injections only

B. Pars plana vitrectomy +Intravitreal antibiotic injections

C. Redo surgery

D. Topical antibiotics

Correct Answer: B. Pars plana vitrectomy +Intravitreal antibiotic injections

Explanation:

References:

AK Khurana Comprehensive Ophthalmology 7th edition, pages 176-178

https://www.ncbi.nlm.nih.gov/books/NBK559079/

Kanski’s clinical ophthalmology, 9th edition, Page 327

↑ *Back to this question in Part A*

@Neet_pg_bot


Q22: Which of the following cataracts is seen in blunt trauma? (Q860514)

A. Rosette cataract

B. Toxic cataract


C. Metabolic cataract

D. Nuclear cataract



Correct Answer: A. Rosette cataract

Explanation: Rosette-shaped cataract: it is a cortical cataract that starts from the posterior cortex is seen in blunt trauma. Traumatic cataract: A traumatic cataract is an acquired cataract resulting from ocular trauma, such as penetrating or blunt injuries. It forms due to degeneration in previously normal lens fibers. Morphological Forms Subepithelial Opacities: Most common presentation. Rosette Cataracts {{caption_text}} Early Rosette: Feathery opacities along posterior cortex suture lines; typical of concussion cataracts. (Option C) Late Rosette: Develops 1-2 years post-injury, with shorter, compact sutural extensions. Diffuse Concussion Cataract: Common presentation of total cataract following blunt trauma, which may accelerate senile cataract development. Subepithelial Opacities (Option A) are common but do not typically present with feathery opacities along suture lines. Diffuse Concussion Cataract (Option B) involves widespread lens opacity, not localized feathery patterns.. Toxic cataract (Option B) causes posterior subcapsular cataract. It can be caused by steroids, phenothiazines, mong acting miotics, amiodarone, busulfan, gold, chloroquine Metabolic cataract (Option C): Due to Diabetes mellitus It can occur due to 2 enzyme deficiencies Galactokinase - causes lamellar cataract GPUT - causes oil drop cataract Nuclear cataract (Option D): It is due to nuclear sclerosis Sclerosis can be caused due to denaturation of the protein resulting in an increase in insoluble protein and a decrease in soluble protein, and deposition of pigments such as melanin and urochrome.



References:

Yanoff & Duker Ophthalmology, 6th edition, Pg 315, 316, 139, 307 e.10

Kanski's Clinical Ophthalmology, 10th edition, Pg 312, 313,914-920

AK Khurana Comprehensive Ophthalmology 7th edition, page 201, 445-449

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: A 2-month-old child is brought to the ophthalmology OPD with the following presentation. What is the ideal management? (Q844026)

A. Medical management

B. Operate immediately

C. Surgery after 6 months

D. Surgery after 2 years



Correct Answer: B. Operate immediately

Explanation: Correct Option B - Operate immediately: In this situation, the image reveals bilateral congenital cataracts, meaning both eyes are affected. If not treated promptly, this condition can result in irreversible vision loss due to amblyopia. For bilateral cataracts, surgery is typically performed after the child reaches one month of age, so immediate surgery is necessary in this case. Incorrect Options: Options A, C, and D are incorrect, as explained above.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q24: A patient presents with decreased vision three years after surgery and exhibits this condition. What is the most likely diagnosis? (Q844003)

A. Pseudo-bullous keratopathy

B. Corneal dystrophy

C. Posterior subcapsular cataract

D. Posterior capsular opacification (PCO)



Correct Answer: D. Posterior capsular opacification (PCO)

Explanation: The patient has decreased vision occurring years after cataract surgery, which is the classic timeline for posterior capsular opacification (PCO). PCO occurs when residual lens epithelial cells migrate, proliferate, and undergo metaplasia on the posterior capsule following cataract extraction. The image shows pearl-like opacities and fibrotic changes on the posterior capsule, known as Elschnig pearls, which are characteristic of PCO. Elschnig pearls are clusters of proliferating lens epithelial cells that form round, pearl-like opacities on the posterior capsule, leading to progressive visual disturbance, as seen in the given image. Incorrect Options: Option A) Pseudo-bullous keratopathy : Involves corneal edema with epithelial bullae, not posterior capsular changes. Option B) Corneal dystrophy : Usually bilateral, inherited, and unrelated to prior surgery. Option C) Posterior subcapsular cataract : Occurs in a natural lens, not after cataract surgery with IOL implantation.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: A patient with a long-standing history of diabetes presents with cataracts. Which of the following metabolic products is primarily responsible for cataract formation in

this condition? (Q852482)

A. Mannitol


B. Galactitol

C. Sorbitol

D. Fructose

Correct Answer: C. Sorbitol

Explanation: In individuals with diabetes, high blood glucose levels can lead to increased activity of the enzyme aldose reductase. This enzyme converts glucose to sorbitol. Sorbitol is poorly metabolized and can accumulate within the lens of the eye. The accumulation of sorbitol leads to osmotic changes, drawing water into the lens and causing it to swell. This swelling and disruption of lens fibers leads to cataract formation. Pathophysiology of Diabetic Cataract Formation: Glucose enters the lens (insulin-independent uptake). Aldose reductase converts glucose → sorbitol . Sorbitol accumulates (since its conversion to fructose by sorbitol dehydrogenase is slow in the lens). Sorbitol is osmotically active , causing lens swelling , oxidative stress, and crystallin protein aggregation → cataract formation . Mannitol (Option A): Mannitol is an osmotic diuretic and is not involved in cataract formation. Galactitol (Option B): Galactitol (dulcitol) accumulates in the lens in galactosemia, leading to early-onset cataracts, but not in diabetic cataracts. Fructose (Option D): While sorbitol is eventually converted to fructose by sorbitol dehydrogenase, the rate is slow in the lens, making sorbitol the primary culprit in diabetic cataracts.



References:

https://pmc.ncbi.nlm.nih.gov/articles/PMC2903955/#:~:text

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: What is Anisometropia? (Q853113)

A. Difference in visual acuity

B. Difference in refractive error between 2 eyes

C. Difference in the axial length of the eyeball

D. Difference in the size of the eyeball

Correct Answer: B. Difference in refractive error between 2 eyes

Explanation: Anisometropia is when total refraction of the two eyes is unequal. The difference in power of two eyes is more than 2.5 diopters. Anisometropia up to 2.5D is well tolerated and that between 2.5 and 4D can be tolerated depending upon the individual sensitivity. However, if it is more than 4D, it is not tolerated and is a matter of concern. Difference in visual acuity (Option A) could be caused by conditions like myopia, hypermetropia, astigmatism, and presbyopia. Difference in axial length (Option C) of the eyeball is seen in axial myopia. Difference in size of eyeball (Option D) is seen in conditions like Microphthalmia and Buphthalmos.

References:

https://www.ncbi.nlm.nih.gov/sites/books/NBK582146/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: A 15-year-old tall boy with long limbs presents to the OPD. On ocular examination, bilateral ectopia lentis is noted. Which gene is most likely affected in this inherited disorder? (Q866368)

A. COL5A

B. PAX6


C. TGFβR2

D. FBN1 (Fibrillin-1)

Correct Answer: D. FBN1 (Fibrillin-1)

Explanation: FBN1 (Fibrillin-1) Bilateral ectopia lentis in a tall individual with long limbs is classic for Marfan syndrome , which is caused by mutations in the FBN1 gene on chromosome 15. Marfan syndrome is an autosomal

dominant connective tissue disorder. Pathophysiology: Defective fibrillin leads to abnormal connective tissue elasticity and excess TGF-β activation. Clinical features Ocular features: Ectopia lentis (usually superotemporal lens subluxation), myopia, risk of retinal detachment. Skeletal features : Tall stature, arachnodactyly, long limbs (dolichostenomelia), pectus deformities, scoliosis. Cardiovascular: Aortic root dilation, aneurysm, aortic regurgitation, mitral valve prolapse—major cause of mortality. Management: Regular cardiovascular monitoring, beta-blockers or ARBs to slow aortic dilation, surgical repair when needed. COL5A (Option A) mutations cause Ehlers–Danlos syndrome (classic type), presenting with hyperextensible skin, hypermobile joints, and poor wound healing, not isolated ectopia lentis. PAX6 (Option B) is associated with aniridia and other anterior segment dysgenesis disorders, not Marfan syndrome. TGFβR2 (Option C) mutations cause Loeys–Dietz syndrome, which shares aortic aneurysms with Marfan's but usually has hypertelorism, bifid uvula, and no typical lens dislocation.



References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [03_Uveitis] 005 Uvea

« Back to Folder Directory

[03_Uveitis] 005 Uvea — Quick Jump

Q1 (p. 576)	*Q2 (p. 576)*	*Q3 (p. 578)*	*Q4 (p. 578)*	*Q5 (p. 579)*	*Q6 (p. 580)*
Q7 (p. 582)	*Q8 (p. 583)*	*Q9 (p. 584)*	*Q10 (p. 587)*	*Q11 (p. 590)*	*Q12 (p. 591)*
Q13 (p. 592)	*Q14 (p. 594)*	*Q15 (p. 595)*	*Q16 (p. 596)*	*Q17 (p. 597)*	*Q18 (p. 597)*
Q19 (p. 598)	*Q20 (p. 599)*	*Q21 (p. 600)*	*Q22 (p. 603)*	*Q23 (p. 603)*	

Q1: A 42-year-old African American woman presents with visual disturbances. Eye examination reveals vitreous cells, snowball opacities, and candle wax drippings. Which type of uveitis is seen in the underlying disease? (Q853682)

A. Anterior uveitis

B. Panuveitis

C. Posterior uveitis

D. Peripheral multifocal chorioretinitis

Correct Answer: B. Panuveitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical scenario suggests ocular sarcoidosis (granulomatous panuveitis). Ocular Sarcoidosis Granulomatous, multi-system disease with unknown cause Non-caseating epithelioid cell granulomas Type of Uveitis Signs Anterior (Option A) Granulomatous iridocyclitis, iris nodules, large mutton fat KPs, anterior chamber cells and flare, posterior synechiae Posterior (Option C) Choroidal and retinal granulomas, cystoid macular edema, periphlebitis retinae with candle wax droppings appearance Peripheral Multifocal Chorioretinitis (Option D) Small, punched-out atrophic spots, highly suggestive of sarcoidosis

References:

Kanski’s Clinical Ophthalmology, 9th edition, page 443, 450-51

↑ *Back to this question in Part A*

@Neet_pg_bot


Q2: A patient came with redness and pain, and you dilated the pupil to break the synechiae. Which drug was used, looking at this picture? (Q882305)

A. Atropine

B. Phenylephrine

C. Scopolamine

D. Miotics



Correct Answer: A. Atropine

Explanation: Atropine is a muscarinic antagonist (anticholinergic) that causes strong cycloplegia and mydriasis. It is commonly used to break posterior synechiae in uveitis because it dilates the pupil and keeps it from sticking to the lens. The irregular “flower-shaped” pupil after dilation is typical after synechiae are broken. Phenylephrine (Option B) : It’s an α1-adrenergic agonist, causes pupil dilation without cycloplegia.Not effective for breaking posterior synechiae because it doesn’t relax the iris sphincter enough. Scopolamine (Option C) : Another anticholinergic like atropine but less commonly used nowadays. Miotics (Option D) :They constrict the pupil.Would worsen synechiae instead of breaking them.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Which of the following conditions can cause panuveitis? (Q853710)

A. VKH syndrome

B. Sympathetic Ophthalmia

C. Sarcoidosis

D. All of the above

Correct Answer: D. All of the above

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Panuveitis can be caused by various systemic and ocular conditions, including VKH syndrome, Sympathetic Ophthalmia, and Sarcoidosis . Causes of Panuveitis: Autoimmune Infectious Sarcoidosis. Birdshot retinochoroidopathy. Sympathetic ophthalia. Polyarteritis nodosa. Lens-induced. Infections (syphilis, TB, HSV/VZV, toxoplasmosis, cryptococcosis).

References:

Uveitis- A Quick Guide to Essential Diagnosis by C. Stephen Foster, edition 1, page 3.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 45-year-old male presents with vision loss in his right eye following a traumatic injury that resulted in a penetrating ocular wound two months ago. Recently, he has developed redness, photophobia, and decreased vision in his left eye as well. On examination, both eyes show signs of bilateral granulomatous uveitis. Which of the following best explains the pathogenesis of the patient's condition? (Q853709)

A. Type I hypersensitivity reaction to lens proteins

B. Delayed hypersensitivity reaction to melanocytes

C. Direct bacterial infection spreading from the injured eye

D. Immune complex deposition in the retinal vasculature

Correct Answer: B. Delayed hypersensitivity reaction to melanocytes

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient is likely experiencing sympathetic ophthalmia , a condition characterized by bilateral granulomatous uveitis following trauma to one eye . The pathogenesis involves a delayed hypersensitivity reaction (T-cell mediated) to tyrosinase peptide antigens or melanocytes , leading to non-necrotizing granulomatous inflammation in both eyes. Sympathetic ophthalmitis (SO) Definition Bilateral, granulomatous, panuveitis occurring after penetrating trauma where uveal prolapse may have been a feature of the trauma Etiology and pathogenesis Trauma to one eye (exciting eye) — 2 weeks to 3 months — Uveitis in the uninjured eye (sympathizing eye). T-cell mediated autoimmune hypersensitivity reaction to Retinal S antigen. (Option B) Maximum risk upon injury to the ciliary body.

References:

Kanski’s Clinical Ophthalmology, 9th edition, page 449. Yanoff and Duker Ophthalmology, 5th edition, page 770.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q5: A 28-year-old woman diagnosed with Vogt–Koyanagi–Harada (VKH) disease, Fundus examination shows the following findings. What does it indicate? (Q853708)

A. Prodromal phase

B. Uveitic phase

C. Convalescent phase

D. Recurrent phase



Correct Answer: C. Convalescent phase

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The image shown is of Sunset Glow Fundus in the convalescent Phase of Vogt–Koyanagi–Harada Disease . The fundus has a slightly reddish appearance, mainly due to the disappearance of melanocytes in the choroid. Vogt–Koyanagi–Harada (VKH) Syndrome Clinical features Clinically, the disease is categorized into four phases: 1. Prodromal phase (lasts for a few days): Neurological (meningitis, cranial nerve paresis) and auditory manifestations (tinnitus, vertigo and deafness). 2. Acute uveitic phase: Bilateral granulomatous anterior and multifocal posterior uveitis with diffuse choroidal infiltration, vitritis, papillitis and exudative retinal detachments. Dalen–Fuchs nodules can be seen in this phase. 3. Convalescent phase (follows several weeks later) : Localized alopecia, poliosis and vitiligo, depigmented fundus appearance (‘sunset glow’ fundus) and depigmented limbal lesions (Sugiura sign). (Option C) 4. Chronic recurrent phase: smouldering anterior uveitis with exacerbations

References:

Yanoff and Duker Ophthalmology, 5th Edition, Page 764,765,766

Kanski’s Clinical Ophthalmology, 9th Edition, Page 446, 447

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 35-year-old male presents with progressively worsening blurry vision, eye pain, and redness in the right eye. Examination reveals restricted eye movement, and fundoscopic evaluation identifies a well-defined cystic lesion in the vitreous humor, accompanied by surrounding inflammation. What is the most probable diagnosis in this patient? (Q853707)

A. Ocular toxoplasmosis

B. Endophthalmitis

C. Ocular cysticercosis

D. Posterior uveitis due to tuberculosis

Correct Answer: C. Ocular cysticercosis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The presence of a well-defined cystic lesion in the vitreous humor with surrounding inflammation , along with restricted eye movement , is characteristic of ocular cysticercosis , caused by the larval stage of Taenia solium. Inflammation occurs mainly due to the release of toxins from the dead cysts. Ocular cysticercosis Definition Infestation caused by Cysticercus cellulosae , the larval form of the pork tapeworm Taenia solium . Clinical Features Systemic Involvement: Larvae can infest the CNS (neurocysticercosis), muscles, lungs, and subcutaneous tissues. Neurological manifestations include seizures and headaches. Eye involvement: Orbit and adnexal cysticercosis: The extraocular muscles (superior rectus- most common) form is the most common type of orbital and adnexal cysticercosis. Presents as periocular swelling, proptosis, ptosis, pain, diplopia, restriction of ocular motility, strabismus, decreased vision, lid edema and orbital cellulitis. Ocular cysticercosis: Intraocular cysticercosis can involve either the anterior or the posterior segment (more common). Presents as blurring of vision, uveitis, vitritis, scotoma, and photopsia. Occular Toxoplasmosis (Option A): Fundoscopy shows focal necrotizing retinitis surrounded by a distinctive “headlight in the fog” appearance. Endophthalmitis (Option B): It is usually associated with post-surgical infection, trauma, or endogenous spread . It presents with diffuse inflammation, hypopyon, and vitritis , not a well-defined cystic lesion. Posterior uveitis due to TB (Option D): Tubercular uveitis often presents with granulomas, choroiditis, or vasculitis rather than a single cystic lesion with inflammation.

References:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5554854/

Parson’s Diseases of the Eye, 23rd edition, page 220.

Yanoff and Duker Ophthalmology, 5th edition, page 745

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: Which of the following ocular conditions is most commonly associated with a positive Amsler sign? (Q853706)

A. Cataract

B. Ocular histoplasmosis syndrome

C. Fuchs heterochromatic uveitis


D. Acute retinal necrosis


Correct Answer: C. Fuchs heterochromatic uveitis


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Amsler sign (aka Amsler-Verrey sign) is commonly seen in Fuchs heterochromic uveitis (FHU). It is characterized by hemorrhage in the anterior chamber after anterior chamber paracentesis or entry with a blade during anterior segment surgeries Fuchs heterochromatic uveitis: Non-granulomatous anterior uveitis Clinical manifestation: Typically unilateral Atypical anterior uveitis Posterior uveitis - less common Lightening of the colour of the affected iris - Heterochromia Stellate keratic precipitates Amsler sign (aka Amsler-Verrey sign)- hemorrhage in the anterior chamber after anterior chamber paracentesis or entry with the blade during anterior segment surgeries. Small twig-like vessels in the angle likely give rise to the Amsler sign.







References:

Parsons Diseases of the Eye, 23rd edition, pg no 583

www.ncbi.nlm.nih.gov/books/NBK538138

www.ncbi.nlm.nih.gov/books/NBK559148

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 35-year-old male with a recent travel history to Africa presents with intense itching, skin nodules, and progressive vision loss. On examination, multiple palpable nodules are found on his skin, and there is early corneal clouding. Microfilariae are observed in a skin snip biopsy. What is the most appropriate treatment for this condition? (Q853705)

A. Clindamycin + sulfadiazine

B. Diethylcarbamazine

C. Ivermectin

D. Self-limiting condition, no treatment needed

Correct Answer: C. Ivermectin

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient's travel history, symptoms, and findings are consistent with onchocerciasis , caused by Onchocerca volvulus . The primary treatment for onchocerciasis is ivermectin, which kills the microfilariae and reduces the intensity of symptoms. Treatment: Ivermectin - reduces the occurrence of blindness and skin disease (Option C) Older treatments consisting of suramin and diethylcarbamazine are avoided (Option B) Clindamycin + sulfadiazine (Option A): Used in Toxoplasmosis uveitis in combination or as part of quadruple therapy.

References:

Parsons Diseases of the Eye, 23rd edition, pg no 577-78

Onchocerciasis - StatPearls - NCBI Bookshelf (nih.gov)

Kanski's Clinical Ophthalmology, 9th edition, Page 463-65

↑ *Back to this question in Part A*

@Neet_pg_bot


Q9: A 32-year-old immunocompromised woman living with cats presents with decreased vision in her left eye. On examination, she has mild anterior uveitis with vitritis and the fundoscopy image is shown below. Which of the following is the most likely causative organism? (Q853687)

A. Cytomegalovirus (CMV)

B. Candida albicans

C. Toxoplasma gondii

D. Histoplasma capsulatum





Correct Answer: C. Toxoplasma gondii


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The most probable diagnosis in this scenario is toxoplasmosis uveitis, which is caused by Toxoplasma gondii. Fundoscopy shows focal necrotizing retinitis surrounded by a distinctive “ headlight in the fog ” appearance. An immunocompromised person living with cats is a risk factor for toxoplasmosis. Toxoplasmosis Uveitis: It is a protozoan infestation. Organism: Toxoplasma gondii Definitive host - cats Intermediate hosts - humans Systemic lesions are more common in immunocompromised patients Clinical manifestation: Most common cause of posterior uveitis and focal necrotizing retinitis (90% of cases) Congenital Inactive disease - Healed punched out heavily pigmented chorioretinal scars in the macular area In active inflammation - “headlight in the fog” appearance due to intense vitritis {{caption_text}} Cytomegalovirus (CMV) (Option A): The characteristic picture is areas of necrosis with areas of vasculitis, retinal hemorrhages, and loss of fundal details giving a pizza pie appearance. Candida albicans (Option B): The characteristic picture is multiple small, round, whitish areas, which may be associated with areas of hemorrhages with pale centers (Roth's spots). Histoplasma capsulatum (Option D): The characteristic picture is choroidal neovascular membranes or disciform lesions at the macula with well-circumscribed peripheral chorioretinal scars in the absence of inflammatory signs in the aqueous and vitreous, and associated with peripapillary pigment epithelial atrophy.


@Neet_pg_bot


@Neet_pg_bot



References:

Parsons Diseases of the Eye, 23rd Edition, Page 574-578

Kanski's Clinical Ophthalmology, 9th Edition, Page 458-461

Ocular toxoplasmosis: a review of the current diagnostic and therapeutic approaches - PMC (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Match bacterial uveitis in column A with corresponding clinical manifestation in column B Bacterial uveitis Clinical features 1. Tubercular uveitis a. Iris pearls 2. Leprotic uveitis b. Iris roceola 3. Spirochetal uveitis c. Headlight in fog appearance d. Koeppe's nodules (Q853686)

A. 1-d, 2-a, 3-b

B. 1-b, 2-c, 3-d

C. 1-c, 2-d, 3-a

D. 1-b, 2-a, 3-d

Correct Answer: A. 1-d, 2-a, 3-b


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Bacterial uveitis Clinical features 1. Tubercular uveitis d. Koeppe’s nodules 2. Leprotic uveitis a. Iris pearls 3. Spirochetal uveitis b. Iris roceola Tubercular uveitis: Causes both anterior, intermediate, posterior, and panuveitis G ranulomatous inflammation causes iris nodules in the pupillary margin (Koeppe nodule) or anterior mid-iris (Busacca nodule) Koeppe nodule Spirochetal uveitis: Iris roseola - dilated iris vessels that develop into nodules ‘Plastic’ iritis occurs in congenital syphilis and Jarisch–Herxheimer reaction ■■■■■■■ Leprotic uveitis: Chronic granulomatous iritis occurs due to direct invasion of the organismal and is characterized by 'iris pearls' near the pupillary margin in a necklace form.




Busacca nodules



@Neet_pg_bot





Iris pearls

@Neet_pg_bot

References:

Parsons' Diseases of the Eye, 23rd Edition, Page 566-70

www.sciencedirect.com/science/article/pii/S1350946223000289

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A 34-year-old woman with a history of Crohn's disease presents with sudden onset of eye pain, redness, and blurred vision. Slit-lamp examination reveals fine, grey keratic precipitates, anterior chamber cells, and flare. Which type of uveitis is most consistent with her presentation? (Q853684)

A. Granulomatous uveitis

B. Non-granulomatous anterior uveitis

C. Posterior uveitis

D. Both A and C are correct

Correct Answer: B. Non-granulomatous anterior uveitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The most likely diagnosis is anterior uveitis, and the extraintestinal manifestations of IBD (Crohn's disease here) include ocular complications. It is non-granulomatous type. Uveitis in bowel disease: Inflammatory bowel disease Most common in Crohn's disease Ocular complications - conjunctivitis, episcleritis, scleritis, marginal keratitis, anterior uveitis , retinitis, retinal vascular occlusive disease, optic neuritis, and orbital inflammatory disease. Type: Non-granulomatous Whipple's disease

References:

Parsons diseases of the eye, 23rd edition, pg no 545

Ocular Complications of Inflammatory Bowel Disease - PMC (nih.gov)

Ophthalmic manifestations in patients with inflammatory bowel disease: A review - PMC (nih.gov)

Immune recovery uveitis in Whipple's disease: an unusual ocular presentation - PMC (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot


Q12: A 30-year-old patient with HIV presents with vision loss and floaters. On examination, the patient has a CD4+ count of 40 cells/mm³. Fundoscopy reveals areas of necrosis with the following appearance. What is the most likely diagnosis? (Q853683)

A. Toxoplasmosis

B. Herpes Simplex Virus (HSV) Retinitis

C. CMV retinitis

D. Herpes Zoster Ophthalmicus




Correct Answer: C. CMV retinitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: CMV retinitis is common in immunocompromised patients (e.g., those with AIDS and low CD4+ counts). Occurs with CD4+ counts <50 cells/mm³ Characterised by retinal necrosis with mild vitritis, and the sauce & cheese (AKA pizza pie) appearance in fundoscopic examination. Classical fundus finding: Yellow-white retinal lesions with hemorrhages along the vascular arcades , giving a “ pizza pie ” or “ cottage cheese with ketchup ”

appearance. Toxoplasmosis (Option A) usually causes focal necrotizing retinochoroiditis with overlying vitreous inflammation ("headlight in the fog" appearance). Herpes Simplex Virus (HSV) (Option B) can cause acute retinal necrosis but is less common in HIV and usually more peripheral with rapid progression. Herpes Zoster Ophthalmicus (HZO) (Option D) involves the cornea and anterior segment, not characteristic retinal necrosis.



References:

Kanski's Clinical Ophthalmology, 9th edition, Pages 468-71, 458.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: Which of the following statements about the uveal tract is false? (Q853669)

A. The choroid connects anteriorly to the iris via the ciliary body.

B. The ciliary body contains both radial and circular smooth muscle fibers.

C. The posterior layer of the iris is continuous with the ciliary body


D. The constrictor pupillae muscle is controlled by sympathetic nerves.


Correct Answer: D. The constrictor pupillae muscle is controlled by sympathetic nerves.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The constrictor pupillae muscle is controlled by parasympathetic nerves, not sympathetic. Middle Coat of the Eye (Uveal Tract) Choroid: Characteristics: Lines the inner surface of the sclera. Connected anteriorly to the iris via the ciliary body . (Option A) Pierced posteriorly by the optic nerve. Ciliary Body: Ciliary Muscles : Smooth muscle with radial and circular fibers. (Option B) Iris: Attachment: Attached peripherally to the ciliary body; forms the iridocorneal angle (Option C).





@Neet_pg_bot

References:

American Academy of Ophthalmology (AAO)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A 30-year-old man presents with symptoms of recurrent bilateral eye inflammation, including redness, pain, and vision loss. He also has a history of frequent oral and genital ulcers and skin rashes. He is HLA-B51 positive. What would be the most appropriate approach for managing his uveitis? (Q853681)

A. Start with systemic corticosteroids and consider long-term immunosuppressive therapy

B. Immediate initiation of antiviral therapy

C. Use of topical antibiotics for ocular symptoms

D. Prescription of oral antihistamines for symptom relief

Correct Answer: A. Start with systemic corticosteroids and consider long-term immunosuppressive therapy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Systemic corticosteroids can provide initial relief for uveitis symptoms in Behcet's disease, but for long-term management, immunosuppressive therapy is necessary due to the disease's chronic nature and poor long-term prognosis with corticosteroids alone. Behcet's Disease Obliterative vasculitis, likely due to circulating immune complexes Treatment Corticosteroids: Effective initially but not long-term Immunosuppressive Drugs: Methotrexate, azathioprine, cyclosporine for resistant or severe cases Immediate initiation of antiviral therapy (Option B) : Antiviral therapy is not appropriate as Behcet's disease is not caused by viral infection. Use of topical antibiotics for ocular symptoms (Option C) : Topical antibiotics are not effective since the uveitis is due to an autoimmune process. Prescription of oral antihistamines for symptom relief (Option D) : Oral antihistamines are not suitable for treating the autoimmune-related uveitis in Behcet's disease.

References:

Kanski's Clinical Ophthalmology, 9th edition, Page 454-56

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 30-year-old man with a recent history of urethritis and arthritis presents with eye symptoms. He is positive for HLA-B27. What is the most appropriate initial treatment? (Q853680)

A. Oral antibiotics

B. Topical corticosteroids

C. Systemic tetracycline

D. Immunosuppressive agents

Correct Answer: B. Topical corticosteroids

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Topical corticosteroids are the initial treatment for acute iridocyclitis. Reiter's Syndrome Reiter's syndrome includes urethritis, arthritis, and conjunctivitis, often with iridocyclitis. Ocular Features : Acute mucopurulent conjunctivitis, possibly with superficial punctate keratitis. Acute, non-granulomatous iridocyclitis occurs in 20-30% of cases. Treatment : Iridocyclitis: Managed with topical, periocular, and systemic corticosteroids. Post Venereal form (likely due to Chlamydia): Treated with systemic tetracycline (250 mg, four times daily for 10 days).

References:

Kanski’s Clinical Ophthalmology, 9th edition, Page 435-36

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A 45-year-old male presents with unilateral eye pain, photophobia, and redness for the past five days. Slit-lamp examination reveals circumcorneal injection, anterior chamber cells and flare, and posterior synechiae. Despite initial treatment with topical corticosteroids and cycloplegics, his symptoms persist, and intraocular pressure (IOP) is found to be elevated. Which of the following is the most likely complication of the condition he is suffering from? (Q853679)

A. Retinal detachment

B. Secondary glaucoma

C. Vitreous hemorrhage

D. Optic neuritis

Correct Answer: B. Secondary glaucoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Secondary glaucoma is a known complication of anterior uveitis, often resulting from inflammatory trabeculitis, corticosteroid response, or synechial angle closure. In this case, the patient has posterior synechiae, which can lead to pupillary block and increased intraocular pressure (IOP). Chronic inflammation can also cause damage to the trabecular meshwork, leading to impaired aqueous outflow and secondary glaucoma. Complication Description Complicated Cataract Common in iridocyclitis, characterized by posterior subcapsular opacities with " polychromatic luster " and " bread-crumb " appearance. Secondary Glaucoma Can occur early or late due to iridocyclitis. Cyclitic Membrane Late complication where exudates behind the lens become organized in acute plastic iridocyclitis . Choroiditis Long-standing iridocyclitis can lead to choroiditis due to the continuity of the iris, ciliary body, and choroid. Retinal Complications Includes cystoid macular edema, macular scar/hole, epiretinal membrane, exudative retinal detachment, secondary periphlebitis retinae, retinal scars, sub-retinal fibrosis. Papillitis Inflammation of the optic disc, seen in severe cases of iridocyclitis. Band-shaped Keratopathy Common in children with Still’s disease and chronic uveitis. Phthisis Bulbi Final stage of chronic uveitis where the eye becomes soft, shrunken, and atrophic.

References:

Kanski’s Clinical Ophthalmology, 9th edition, Page 425-27 Yanoff and Duker Ophthalmology, 5th edition, Page 696-97

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: Which of the following is not a component of Posterior uveitis? (Q853678)

A. Retinitis

B. Choroiditis

C. Iridocyclitis

D. Retinal vasculitis

Correct Answer: C. Iridocyclitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Posterior uveitis is defined as the inflammation of the choroid and the retina. The components of posterior uveitis include the following, Choroiditis Retinitis Retinal vasculitis Periphlebitis - Venous sheathing Periarteritis - Hemorrhages Retinitis (Option A): Inflammation of the neural retina. The retina lies in the posterior segment of the globe. In many cases of posterior uveitis, the inflammation spreads from the choroid to the retina (chorioretinitis) or starts in the retina itself. Choroiditis (Option B): Inflammation of the choroid, the vascular layer between the retina and the sclera. The choroid makes up the posterior portion of the uveal tract. It is the defining feature of "true" posterior uveitis. Retinal Vasculitis (Option C): Inflammation of the retinal blood vessels. Periphlebitis - Venous sheathing Periarteritis - Hemorrhages

References:

Kanski’s Clinical Ophthalmology, 9th edition, Page 427, 431

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: What is the 1st line of treatment in a 35 year old patient presenting with Intermediate uveitis? (Q853676)

A. Pars plana vitrectomy

B. Systemic steroids

C. Periocular injection of triamcinolone

D. Cryotherapy

Correct Answer: C. Periocular injection of triamcinolone

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The most appropriate initial treatment for intermediate uveitis, especially when it is idiopathic and presents with significant vitreous inflammation, is the periocular injection of triamcinolone. This approach targets inflammation locally with fewer systemic side effects compared to systemic steroids. Treatment (Kaplan's Modified 4-Step Protocol) Step Treatment Details 1 Periocular & Systemic Steroids Triamcinolone injection : 40 mg, 3 injections spaced 3 weeks apart. Oral corticosteroids: Typically considered if periocular steroids are insufficient. Gradual tapering is done. (Option B) 2 Immunosuppressive Drugs Agents : Cyclosporine, azathioprine, methotrexate, cyclophosphamide. Close monitoring due to potential side effects. 3 Cryotherapy/Indirect Laser Photocoagulation (Option D) Cryotherapy : Controlled cold-induced injury. Used for snowbanking when less invasive treatments fail. Laser photocoagulation: Focused light beam treatment. 4 Pars Plana Vitrectomy (Option A) Surgery: For severe cases or complications (e.g., tractional retinal detachment). Removes vitreous gel, and clears debris.

References:

Kanski’s Clinical Ophthalmology, 9th edition, Page 443-45

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: A 25-year-old female presents with floaters and gradual deterioration of vision. On examination, there is anterior vitritis with white snowball-like exudates and mild peripheral periphlebitis. No other relevant history or clinical findings. Which of the following is the most likely diagnosis? (Q853675)

A. CMV Retinitis

B. Tuberculosis

C. Intermediate uveitis

D. Lyme Uveitis


Correct Answer: C. Intermediate uveitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The presence of floaters, anterior vitritis, and white snowball-like exudates near the ora serrata with mild peripheral periphlebitis are characteristic of intermediate uveitis (pars planitis). Intermediate Uveitis: Causes and Features Definition: Intermediate uveitis, also known as pars planitis, primarily affects the pars plana of the

ciliary body and the peripheral choroid. It mostly occurs in children and young adults, with a higher incidence in females. Features Common Features Floaters Vision deterioration due to anterior vitreous opacities Minimal aqueous flare with occasional keratic precipitates Anterior Vitritis White snowball-like exudates near the ora serrata Mild peripheral periphlebitis CMV Retinitis: Typically presents with focal retinochoroiditis rather than the diffuse features seen in intermediate uveitis. (Option A) Tuberculosis: Typically causes anterior uveitis. (Option B) Lyme Uveitis: May cause uveitis but typically presents with more acute dermatological symptoms, and the ocular findings are not as specific to intermediate uveitis as in this case. More commonly seen in the relapsing-remitting type. (Option D)



References:

Parsons Diseases of the eye, 23rd edition, Pg 553.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: A 40-year-old patient presents with eye pain, redness, and a constricted pupil. During the examination, fine, grey deposits on the corneal endothelium are noted. Which of the following is the most appropriate next step in investigating and treating this condition? (Q853674)

A. Perform a chest X-ray and initiate treatment with topical cycloplegics.

B. Order a serum ACE level and start treatment with non-steroidal voclosporin.

C. Conduct an HLA typing and begin treatment with topical prednisolone.

D. Arrange for an OCT scan and administer systemic interferon-alpha.

Correct Answer: C. Conduct an HLA typing and begin treatment with topical prednisolone.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The findings are consistent with a case of anterior uveitis. HLA typing is important in identifying associated systemic conditions that may cause anterior uveitis, such as Behçet's disease or VKH syndrome. Topical prednisolone is a standard treatment for reducing inflammation in anterior uveitis. Investigations for Anterior Uveitis: Ocular Investigations Systemic Investigations USG-B-Scan : To assess the posterior segment of the eye for any abnormalities. Fundus Fluorescein Angiography (FFA) : To evaluate the retinal and choroidal circulation and detect any leakage or abnormalities. Optical Coherence Tomography (OCT) Syphilis Tests : VDRL TPHA Sarcoidosis : Serum ACE Levels : Chest X-Ray : Mantoux Test / QuantiFERON Test X-Ray Spine HLA Typing : HLA-B27 : Associated with various types of arthritis, including ankylosing spondylitis. HLA-B5 and HLA-B51 : Associated with Behçet's disease. HLA-DR4 : Associated with VKH syndrome and sympathetic ophthalmia. Treatment for Anterior Uveitis: Topical Steroids (Treatment of Choice): Prednisolone : Commonly used for reducing inflammation. Loteprednol : Less potential for increasing intraocular pressure. Cycloplegics (Topical): Rest the ciliary muscles to relieve pain. Prevent and break posterior synechiae (adhesions of the iris to the lens). Improve blood supply, leading to better healing. Non-Steroidal Treatment : Voclosporin : Part of the Luminate program, used as a non-steroidal option. For Chronic or Vision-Threatening Uveitis : Monoclonal Antibodies : Adalimumab : TNF-alpha inhibitor, used for severe or chronic cases. Infliximab : TNF-alpha inhibitor. Interferon Alpha : An option for severe cases, helps modulate the immune response.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 424, 430-433

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: A 35-year-old patient presents with a 1-week history of worsening eye pain and redness. On examination, a constricted pupil, a hazy anterior chamber, and deposits on the back of the cornea as shown below. Based on these findings, which of the following is the most likely diagnosis? (Q853673)

A. Herpes Simplex Keratitis

B. Conjunctivitis

C. Anterior Uveitis


D. Acute Angle Closure Glaucoma



Correct Answer: C. Anterior Uveitis

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Anterior uveitis is characterized by the presence of fine, grey keratic precipitates (KPs) on the corneal endothelium, a hazy anterior chamber (aqueous flare), and a constricted pupil. Anterior Uveitis Examination Findings : Keratic Precipitates (KPs) : Non-Granulomatous : Fine, grey deposits; composed of lymphocytes Granulomatous : Large, greasy-looking deposits known as "mutton fat KPs"; composed of macrophages Aqueous Cells : Indicator of active inflammation Aqueous Flare : Haziness in the anterior chamber due to leakage of proteins from inflamed tissues Muddy Iris : Edema in the iris caused by protein leakage Miosis : Constricted pupil due to toxins released from inflamed tissue Nodules on the Iris : Koeppe's Nodules : At the pupillary border; seen in both granulomatous and non-granulomatous uveitis Busacca Nodules : At the b ase of the iris; seen in granulomatous uveitis Koeppe's Nodules Herpes Simplex Keratitis (Option A): Presents with dendritic corneal ulcers, not fine, grey keratic precipitates or a hazy anterior chamber. It also usually does not involve a constricted pupil. Conjunctivitis (Option B): Does not cause keratic precipitates , hazy anterior chamber and a constricted pupil. Acute Angle Closure Glaucoma (Option D): Presents with a mid-dilated or fixed, mid-dilated pupil, severe pain, redness, and vision changes.





References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 424-427

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: A 50-year-old patient presents with anterior uveitis. After a detailed evaluation, the ophthalmologist considers various systemic associations and possible masquerades for the condition. Which of the following conditions is a known systemic association with anterior uveitis, but not a masquerade? (Q853672)

A. Intraocular Foreign Body

B. Behcet's Disease

C. Retinoblastoma

D. Pigment Dispersion Syndrome

Correct Answer: B. Behcet's Disease

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Behcet's Disease is a known systemic association with anterior uveitis, characterized by recurrent oral and genital ulcers, and uveitis. Anterior uveitis , also known as iritis, is an inflammatory condition of the eye, including the iris and ciliary body. Masquerade Syndromes : Conditions that mimic the symptoms of anterior uveitis but are not primarily inflammatory in nature. Neoplastic Masquerades : Lymphomas Anterior Segment Melanoma Retinoblastoma (Option C) Metastasis Non-Neoplastic Masquerades : Intraocular Foreign Body (Option A) Old Retinal Detachment Pigment Dispersion Syndrome (Option D) Coats' Disease

References:

Kanski’s Clinical Ophthalmology 9th ed., Page 424

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Which of the following statements about the classification of uveitis based on the time of inflammatory activity is false? (Q853670)

A. Acute uveitis is defined as having a duration of less than 3 months and typically has a sudden onset.

B. Chronic uveitis is classified as lasting more than 6 months without interruption.

C. Recurrent uveitis involves repeated episodes with periods of inactivity between them.

D. Remission in uveitis is defined as no inflammatory activity for 3 months or more.

Correct Answer: B. Chronic uveitis is classified as lasting more than 6 months without interruption.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Chronic uveitis is defined as uveitis that lasts more than 3 months, not 6 months. Classification Based on Time of Inflammatory Activity Acute Uveitis (Option A): Duration: < 3 months Onset: Sudden Chronic Uveitis: Duration: > 3 months Recurrent Uveitis (Option C): Characterized by repeated episodes Includes periods of inactivity between episodes Remission (Option D): No inflammatory activity for ≥ 3 months

References:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8935739/

↑ *Back to this question in Part A*

@Neet_pg_bot

• [03_Uveitis] 006 Previous Year Questions

« Back to Folder Directory

[03_Uveitis] 006 Previous Year Questions — Quick Jump

Q1 (p. 605) *Q2 (p. 607)* *Q3 (p. 608)* *Q4 (p. 608)* *Q5 (p. 609)* *Q6 (p. 612)*
Q7 (p. 613) *Q8 (p. 617)* *Q9 (p. 619)* *Q10 (p. 619)* *Q11 (p. 622)* *Q12 (p. 624)*

Q1: A 35-year-old male presents with pain, redness, watering, and photophobia in the right eye for 2 days. Vision is mildly reduced. On examination, there is ciliary congestion, small irregular pupil, and presence of keratic precipitates on the corneal endothelium. The intraocular pressure is normal or slightly low.What is the most likely diagnosis? (Q866836)

A. Acute uveitis

B. Acute conjunctivitis


C. Acute congestive glaucoma


D. Central retinal artery occlusion

Correct Answer: A. Acute uveitis

Explanation: The most likely diagnosis is acute anterior uveitis , characterized by pain, redness, photophobia, ciliary flush, keratic precipitates, and a small irregular or sluggish pupil with normal or slightly low IOP . Corneal haze results from cells and flare , and a shallow anterior chamber may occur from ciliary body congestion pushing the iris forward . Anterior Uveitis: Most common type of uveitis affecting the anterior segment of the eye. Clinical Features : Pain, Redness, Photophobia (sensitivity to light), Blepharospasm (involuntary blinking), Serous discharge, Blurring and diminished vision Examination Findings : Keratic Precipitates (KPs) : Non-Granulomatous : Fine, grey deposits; composed of lymphocytes Granulomatous : Large, greasy-looking deposits known as "mutton fat KPs"; composed of macrophages Location : Typically found in the lower cornea due to convection currents in aqueous humor; this area is known as Arlt's triangle Aqueous Cells : Indicator of active inflammation Aqueous Flare : Haziness in the anterior chamber due to leakage of proteins from inflamed tissues Muddy Iris : Edema in the iris caused by protein leakage Miosis : Constricted pupil due to toxins released from inflamed tissue Nodules on the Iris : Koeppe's Nodules : At the pupillary border; seen in both granulomatous and non-granulomatous uveitis Busacca Nodules : At the b ase of the iris; seen in granulomatous uveitis Complications : (Findings) Posterior Synechiae : Iris adheres to the lens; may lead to festooned-shaped pupil Secclusio Pupil/Ring Synechiae : Adhesion along the entire pupillary border; can lead to Iris Bombe and peripheral anterior synechiae, causing angle closure glaucoma Occlusio Pupil : Covers the whole surface of the lens Cyclitic Membrane Rubeosis Iridis : New vessel formation on the iris Acute conjunctivitis (Option B) causes diffuse redness, mucopurulent discharge, and no photophobia or small pupil. Acute congestive glaucoma (Option C) presents with a fixed mid-dilated pupil, hazy cornea, and markedly raised intraocular pressure, not seen here.

Central retinal artery occlusion (Option D) causes sudden painless loss of vision, with a pale retina and cherry-red spot, not pain or redness.







References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A patient presents with sudden diminution of vision, difficulty looking towards the light, and circumcorneal congestion with a hypopyon. What is the most likely diagnosis? (Q844008)

A. Acute conjunctivitis

B. Acute angle closure glaucoma

C. Episcleritis

D. Acute anterior uveitis

Correct Answer: D. Acute anterior uveitis

Explanation: Correct Option D - Acute anterior uveitis: The clinical presentation of the patient is suggestive of acute anterior uveitis. The presence of a hypopyon is often seen in conditions like Behcet disease, which can typically cause non-granulomatous anterior uveitis. Incorrect Options: Options A, B, and C do not present with a hypopyon.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 32-year-old patient presents with blurred vision, photophobia, and mild ocular pain. Examination reveals aqueous flares and keratic precipitates in the anterior chamber. What is the likely diagnosis? (Q860512)

A. Intermediate uveitis

B. Posterior uveitis

C. Toxoplasma uveitis

D. Iridocyclitis

Correct Answer: D. Iridocyclitis

Explanation: Type of Uveitis Signs Anterior (Option D) Granulomatous iridocyclitis , iris nodules, large mutton fat KPs, anterior chamber cells and flare, posterior synechiae Intermediate (Option A) Vitreous cells, snowball opacities, snowbanking Posterior (Option B) Choroidal and retinal granulomas, cystoid macular edema, periphlebitis retinae with candle wax droppings appearance Peripheral Multifocal Chorioretinitis Small, punched-out atrophic spots, highly suggestive of sarcoidosis Uveoparotid Fever (Heerfordt's Syndrome) Bilateral granulomatous panuveitis, painful parotid gland enlargement, cranial nerve palsies, skin rashes, fever, malaise Toxoplasma uveitis (Option C): Definitive host: Cat; Intermediate host: Human Ocular features: granulomatous or non-granulomatous uveitis Primarily it is posterior uveitis and spreads to anterior

References:

Kanski's Clinical Ophthalmology, 9th edition, page 443, 450-51

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: Iritis is least associated with all of the following diseases: (Q854715)

A. Systemic Lupus Erythematosus

B. Behcet's Disease

C. Psoriatic Arthropathy

D. Ulcerative colitis

Correct Answer: A. Systemic Lupus Erythematosus

Explanation: Iritis , also known as Anterior uveitis , is an inflammatory condition of the eye, including the iris and ciliary body. Although SLE is an autoimmune disease that can affect multiple organs, iritis is less commonly seen in SLE compared to the other conditions listed. It more often causes other ocular manifestations, like dry eyes or retinal changes . Systemic Associations and Masquerades of Iritis (Anterior Uveitis) Infectious Causes : Herpes Infections: Varicella Zoster, Herpes Simplex Virus (HSV) Tuberculosis (TB) Syphilis Leprosy Non-Infectious Causes : With Systemic Association : Sarcoidosis Behçet's Disease (Option B ruled out) HLA-B27 Associated Arthritis Psoriatic Arthropathy (Option C ruled out) Inflammatory Bowel Disease: Ulcerative Colitis (UC) > Crohn's Disease (Option D ruled out) Without Systemic Association : Drug-Induced: Pilocarpine, prostaglandin analogues Masquerade Syndromes : Definition : Conditions that mimic the symptoms of anterior uveitis but are not primarily inflammatory in nature. Neoplastic Masquerades : Lymphomas Anterior Segment Melanoma Retinoblastoma Metastasis Non-Neoplastic Masquerades : Intraocular Foreign Body Old Retinal Detachment Pigment Dispersion Syndrome Coats' Disease

References:

Kanski’s Clinical Ophthalmology 9th ed., Page 424

↑ *Back to this question in Part A*

@Neet_pg_bot


Q5: A farmer presents with pain and discharge in his left eye. He says that he sustained an injury caused by a leaf a few days back. What is the finding seen in the image? (Q855559)

A. Hyphema

B. Aqueous flare

C. Arlt line


D. Hypopyon




Correct Answer: D. Hypopyon

Explanation: Given the farmer's symptoms of pain, discharge, and history of trauma, along with the finding of white blood cells in the anterior chamber , the most likely diagnosis is hypopyon , which often results from infection or severe inflammation following eye trauma. Fungal keratitis is a common cause of corneal ulcers following eye trauma, especially when the injury is caused by vegetative matter (such as plant material or dirt). This type of trauma provides an entry point for fungal spores . The classic signs and symptoms of fungal keratitis include: Corneal ulcer with satellite lesions (smaller ulcers surrounding the main ulcer). Hypopyon (accumulation of white blood cells in the anterior chamber of the eye). Mild pain , photophobia , and redness . The history of vegetative matter as the cause of trauma is a significant clue, as fungi are commonly found in plant material. Keratomycosis {{caption_text}} Clinical Features Pain & Foreign Body Sensation Watering of the eye Photophobia Blurred Vision Redness. Examination Findings Corneal Ulcer : Greyish-white, dry-looking with elevated, rolled-out margins. Pigmented Ulcer : Brownish appearance, possible with dematiaceous fungi. Feathery Extensions : Delicate, finger-like projections into the surrounding stroma. Sterile Immune Ring : Yellow line of demarcation where fungal antigen and host antibodies meet. Satellite Lesions : Multiple small lesions around the main ulcer. Hypopyon : Large, non-sterile, even with a small ulcer. {{caption_text}} Endothelial Plaque : Located under the stromal lesion, made of fibrin and leucocytes. Corneal Vascularization : Typically absent in pure fungal ulcers. Perforation : Rare but possible. Investigations Corneal Scraping : Performed under topical anaesthesia for microbiological examination. Wet KOH Mount : Reveals filamentous fungi (branched, septate hyphae) or non-filamentous yeast-like fungi (Candida). Gram's & Giemsa Stains : To detect fungal hyphae and inflammatory cells. Culture : Sabouraud's agar medium is used for fungal culture. PCR : Recommended for rapid identification of fungal pathogens. Anterior Chamber Paracentesis : Considered if there is an intraocular extension. Corneal Biopsy : Indicated for deep stromal abscess or when cultures are negative. Confocal Microscopy : Used for identifying actual fungi within the cornea . Treatment of Keratomycosis (Fungal Corneal Ulcer): Primary Treatment: Antifungal Medications: Topical Antifungal Agents: Natamycin Voriconazole Amphotericin B

(effective against Aspergillus and Fusarium) Nystatin (effective against Candida) Oral Antifungal Agents (for severe ulcers with specific features): Ketoconazole Voriconazole Indications for Oral Therapy: Hypopyon Perforation Suspicion of endophthalmitis Surgical Management (if unresponsive to medical therapy): Repeated Scraping: Removal of necrotic material Cauterization Options: Pure carbolic acid (100%) Trichloroacetic acid (10%–20%) Povidone-iodine 5% Therapeutic Keratoplasty: Corneal transplantation Hyphema (Option A) : Refers to the presence of blood in the anterior chamber, often caused by trauma. While the patient did sustain trauma, the presence of pus (not blood) in the anterior chamber points to hypopyon rather than hyphema. Aqueous flare (Option B) : Refers to the presence of protein in the aqueous humor, which can be seen in inflammatory conditions, but it is generally seen as a cloudy appearance in the anterior chamber, not a white collection of cells like hypopyon. Arlt line (Option C) : This refers to a line of conjunctival scarring seen in trachoma , which is not related to the acute trauma described here and would not be associated with the findings in the anterior chamber.





@Neet_pg_bot

References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 216, 217.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Iritis is not seen in: Rheumatoid arthritis SLE Bechet's disease Psoriatic arthritis Ulcerative colitis (Q856415)

A. 1 and 2

B. 1,2,3 and 4

C. 2,4 and 5

D. 1,3 and 4

Correct Answer: A. 1 and 2

Explanation: Conditions where iritis is seen: Behcet's disease (Statement 3 ruled out): Characterized by recurrent uveitis, including iritis, genital involvement and oral involvement. Psoriatic arthritis (Statement 4 ruled out): A

seronegative spondyloarthropathy frequently associated with anterior uveitis including iritis, urethritis and arthritis. Ulcerative colitis (Statement 5 ruled out): Iritis is common in inflammatory bowel disease, as it shares a systemic inflammatory pathway. Rheumatoid arthritis: Iritis is rarely seen in adult-onset rheumatoid arthritis. However, it is more commonly associated with juvenile idiopathic arthritis (JIA), especially in ANA-positive cases. SLE: Systemic lupus erythematosus typically affects other ocular tissues, such as the retina and optic nerve, rather than causing iritis.

References:

Harrison's Principles of Internal Medicine, 21st Edition, Page 2791, 2793, 2794

Davidson's Principles and Practice of Medicine, 24th Edition, Page 1028, 1030

Apley and Solomon's System of Orthopaedics and Trauma, Edition 10th- Page 66, 67, 68, 104.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 26-year-old patient came with complaints of blurring of vision, photophobia, redness, pain in the left eye, and joint pain. Upon examination keratic precipitates and aqueous cells were observed. Which of the following major histocompatibility complex is be associated with this condition? (Q859309)

A. HLA B27

B. HLA B51


C. HLA B47

D. HLA B57

Correct Answer: A. HLA B27


Explanation: The clinical presentation of blurring vision, photophobia, redness, pain ,and keratic precipitates are suggestive of anterior uveitis. Patients with acute anterior uveitis with joint pain is suggestive of arthritis associated with uveitis. The disease affects young males who are positive for HLA B27. Among patients with ankylosing spondylitis, 90% of them belong to the HLA-B27 antigenic group. Ankylosing Spondylitis (AS) / Marie Strumpell Disease Definition Chronic inflammatory disease primarily affecting the axial skeleton (spine and sacroiliac joints), leading to pain, stiffness, and spinal fusion (bamboo spine) . Etiology Unknown; genetic (HLA-B27) and environmental factors involved. HLA-B27 is positive in 90% of cases, but not all carriers develop AS. Pathophysiology Inflammation at entheses → Erosive changes → New bone formation → Syndesmophytes → Spinal fusion (ankylosis). {{caption_text}} Clinical Features Musculoskeletal Symptoms: Chronic low back pain (worse after inactivity, better with activity) Limited spinal mobility Enthesitis (Achilles, plantar fascia) Peripheral arthritis (hips, shoulders, knees) Extra-articular Symptoms: Ocular: Anterior uveitis (pain, redness, photophobia) Cardiovascular: Aortic regurgitation, conduction issues Pulmonary: Apical fibrosis Gastrointestinal:

Associated with IBD Diagnosis Based on clinical features, imaging studies, and HLA-B27 gene presence. Modified New York Criteria: Clinical Criteria: Low back pain > 3 months (improves with exercise, not with rest) Limited lumbar spine motion (sagittal/frontal planes) Reduced chest expansion (relative to age/sex) Radiographic Criteria (X-ray): Bilateral sacroiliitis (grade 2-4) or unilateral sacroiliitis (grade 3-4) Radiological Features: Blurring/haziness of sacroiliac joint {{caption_text}} Juxta-articular sclerosis Vertical/bridging syndesmophytes Squaring of vertebrae {{caption_text}} Bamboo spine Trolley track sign Dagger sign {{caption_text}} Shiny corner sign/Romanus lesion (inflammatory lesions at edges of vertebrae) {{caption_text}} MRI: Early detection of inflammation Clinical Tests: For Sacroiliitis: Gaenslen's test Patrick/FABER test Figure of 4 test Pump handle test For Lumbar Spine: Schober/ Modified Schober test {{caption_text}} Laboratory Tests HLA-B27 positive (90% of cases) ESR, CRP: Elevated during active disease, but can be normal Differential Diagnosis Mechanical low back pain Other spondyloarthropathies (e.g., psoriatic arthritis) Degenerative disc disease Fibromyalgia Treatment Non-Pharmacological: Physical therapy (mobility/posture exercises) Patient education on disease and exercise importance Pharmacological: NSAIDs (first-line for pain, stiffness) TNF inhibitors (for NSAID-resistant cases) IL-17 inhibitors (Secukinumab) Sulfasalazine (for peripheral arthritis) Corticosteroids (short-term flare control) Surgical: Hip replacement (for severe hip involvement) Spinal surgery (for severe deformities/fractures) Complications Spinal fractures (minimal trauma) Cardiac issues (aortic regurgitation, conduction defects) Osteoporosis leading to fracture risk HLA B51 (Option B) It is associated with Behcet's syndrome with anterior uveitis. Behcet's syndrome is characterized by bilateral recurrent oral ulcers, genital ulcers, uveitis, and skin lesions. HLA B47 (Option C) Is associated with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. It is characterized by symptoms like diarrhea, vomiting, arrhythmias, and low blood pressure. HLA B57 (Option D) Is associated with the detection of abacavir hypersensitivity and slow progression of HIV.





@Neet_pg_bot


@Neet_pg_bot


@neet_pg_bot


@neet_pg_bot



References:

Apley and Solomon’s System of Orthopedics and Trauma, 10th edition, Page 72-76.

https://www.niams.nih.gov/health-topics/ankylosing-spondylitis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 25-year-old male patient with a penetrating injury to the left eye was treated with corneoscleral suturing. Two months later, he returned with complaints of reduced near vision, pain, redness, and sensitivity to light in his right eye. Direct ophthalmoscopy revealed nodules in the fundus. Which part of the eye is affected? (Q859763)

A. Uvea


B. Cornea

C. Conjunctiva

D. Sclera

Correct Answer: A. Uvea

Explanation: The clinical presentation of penetrating injury to the left eye and subsequent reduced near vision, pain, redness, and photophobia along with nodules on fundus i.e., Dalen-Fuchs nodules are suggestive of Sympathetic ophthalmitis. Sympathetic ophthalmitis (SO) Definition Bilateral, granulomatous, panuveitis (Option A) occurring after penetrating trauma where uveal prolapse may have been a feature of the trauma Etiology and pathogenesis Trauma to one eye (exciting eye) — 2 weeks to 3 months — Uveitis in the uninjured eye (sympathizing eye). T-cell mediated autoimmune hypersensitivity reaction to Retinal S antigen. Maximum risk upon injury to the ciliary body. Histopathology Diffuse lymphocytic infiltration of the choroid. Scattered aggregates of epithelioid cells are seen, many of which contain fine granules of melanin. Dalen–Fuchs nodules - granulomas located between Bruch membrane and the RPE (also seen in VKH syndrome) {{caption_text}} Clinical features The exciting eye is frequently red and irritable. The sympathizing eye develops irritation, blurred vision, photophobia and loss of accommodation causing difficulty in near vision (first symptom) Anterior uveitis develops in both eyes, which may be mild or severe and is usually granulomatous Investigations Fundus exam: Dalen–Fuchs nodules, Exudative retinal detachment, vasculitis and optic disc swelling. As inflammation settles, residual chorioretinal scarring may confer a ‘sunset glow’ appearance similar to VKH. B-scan ultrasonography : Choroidal thickening and retinal detachment Treatment Enucleation of a severely injured eye within 10 to 14 days of the injury is effective in preventing or reducing the severity of SO Steroids are the basis of initial treatment. Intravenous methylprednisolone followed by high-dose oral prednisolone is given for several months and gradually tapered according to response. Supplementary topical steroids and cycloplegics may be given Options B, C, and D are incorrect.



References:

Kanski’s Clinical Ophthalmology, 9th edition, page 449.

Yanoff and Duker Ophthalmology, 5th edition, page 770.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Which medication can be administered to a patient diagnosed with acute anterior uveitis, presenting with symptoms of photophobia and blurred vision? (Q859766)

A. Moxifloxacin

B. Atropine

C. Acetazolamide

D. Methotrexate

Correct Answer: B. Atropine

Explanation: Treatment for Anterior Uveitis: Topical Steroids (Treatment of Choice): Prednisolone : Commonly used for reducing inflammation. Loteprednol : Less potential for increasing intraocular pressure. Cycloplegics (Topical): Rest the ciliary muscles to relieve pain. Prevent and break posterior synechiae (adhesions of the iris to the lens). Improve blood supply, leading to better healing. Eg.: Atropine . Non-Steroidal Treatment : Voclosporin: Part of the Luminate program, used as a non-steroidal option. For Chronic or Vision-Threatening Uveitis : Monoclonal Antibodies : Adalimumab : TNF-alpha inhibitor, used for severe or chronic cases. Infliximab : TNF-alpha inhibitor. Interferon Alpha : An option for severe cases, helps modulate the immune response. Incorrect Options: Moxifloxacin (Option A) is an antibiotic commonly used to treat bacterial infections. Acetazolamide (Option C) is a medication primarily used to treat glaucoma by reducing the production of aqueous humor and lowering intraocular pressure. Methotrexate (Option D) is an immunosuppressive medication commonly used in the treatment of various autoimmune and inflammatory conditions. It may be prescribed for severe or chronic cases of uveitis that do not respond to other treatments.

References:

Kanski’s Clinical Ophthalmology, 9th edition, Page 424, 430-33

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: What is the specific sign that is pathognomonic for acute anterior uveitis among the options provided? (Q859771)

A. Aqueous flare


B. Keratic precipitates

C. Festooned pupil


D. Bussaca’s nodules

Correct Answer: B. Keratic precipitates

Explanation: Anterior Uveitis: Definition : Most common type of uveitis affecting the anterior segment of the eye. Clinical Features : Pain Redness Photophobia (sensitivity to light) Blepharospasm (involuntary blinking) Serous discharge Blurring and diminished vision Examination Findings : Keratic Precipitates (KPs) (Option B) : Non-Granulomatous : Fine, grey deposits; composed of lymphocytes Granulomatous : Large, greasy-looking deposits known as "mutton fat KPs"; composed of macrophages Location : Typically found in the lower cornea due to convection currents in aqueous humor; this area is known as Arlt's triangle {{caption_text}} Aqueous Cells : Indicator of active inflammation Aqueous Flare : Haziness in the anterior chamber due to leakage of proteins from inflamed tissues Muddy Iris : Edema in the iris caused by protein leakage Miosis : Constricted pupil due to toxins released from inflamed tissue Nodules on the Iris : Koeppe's Nodules : At the pupillary border; seen in both granulomatous and non-granulomatous uveitis Busacca Nodules : At the b ase of the iris; seen in granulomatous uveitis {{caption_text}} Complications : (Findings) Posterior Synechiae : Iris adheres to the lens; may lead to festooned-shaped pupil {{caption_text}} Secclusio Pupil/Ring Synechiae : Adhesion along the entire pupillary border; can lead to Iris Bombe and peripheral anterior synechiae, causing angle closure glaucoma Occlusio Pupil : Covers the whole surface of the lens Cyclitic Membrane Rubeosis Iridis : New vessel formation on the iris Incorrect Options: Aqueous flare (Option A) refers to the presence of protein and inflammatory cells in the aqueous humor, which may be seen in anterior uveitis. Festooned pupil (Option C) is a characteristic feature of posterior synechiae in chronic uveitis. Bussaca’s nodules (Option D) are larger, granulomatous nodules that are seen in conditions like chronic anterior uveitis, especially in sarcoidosis or tuberculosis.


@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology 9th ed., page 424-27

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: What will be observed in a patient diagnosed with sympathetic ophthalmitis, and a history of penetrating eye injury? (Q856036)

A. Acute anterior uveitis

B. Pars planitis


C. Panuveitis


D. Chronic anterior uveitis

Correct Answer: C. Panuveitis

Explanation: alen–Fuchs nodules, Exudative retinal detachment, vasculitis and optic disc swelling. As inflammation settles, residual chorioretinal scarring may confer a 'sunset glow' appearance similar to VKH.

B-scan ultrasonography : Choroidal thickening and retinal detachment Treatment Enucleation of a severely injured eye within 10 to 14 days of the injury is effective in preventing or reducing the severity of SO Steroids are the basis of initial treatment. Intravenous methylprednisolone followed by high-dose oral prednisolone is given for several months and gradually tapered according to response. Supplementary topical steroids and cycloplegics may be given Causes of Panuveitis: MCP: Multifocal Choroiditis and Panuveitis, BSRC: Birdshot Retinochoroidopathy, ABD: Acute Bilateral Diffuse Uveal Retinitis, VKH: Vogt-Koyanagi-Harada Disease, SFU: Sarcoid-Associated Fuchs Uveitis, TB: Tuberculosis, HSV: Herpes Simplex Virus, VZV: Varicella-Zoster Virus, ARN: Acute Retinal Necrosis Acute anterior uveitis (Option A) : While acute anterior uveitis can occur in sympathetic ophthalmitis, it is not the most characteristic feature. Inflammatory involvement is not limited to just the anterior uvea in sympathetic ophthalmitis, as it typically progresses to involve all layers (hence panuveitis). Pars planitis (Option B) is a type of intermediate uveitis that primarily affects the pars plana of the eye, but it is not typically associated with sympathetic ophthalmitis. It is a distinct form of uveitis, not related to penetrating trauma or autoimmune responses following injury. Chronic anterior uveitis (Option D) : This is not the typical presentation in sympathetic ophthalmitis. While inflammation can persist, it is more commonly panuveitis (involving both anterior and posterior segments), rather than isolated chronic anterior uveitis.





References:

Kanski's Clinical Ophthalmology, 9th edition, page 449.

Yanoff and Duker Ophthalmology, 5th edition, page 770.

Uveitis- A Quick Guide to Essential Diagnosis by C. Stephen Foster, edition 1, page 3.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: On ophthalmologic examination, the presence of Koeppe's nodules is observed in a patient diagnosed with sarcoidosis. In which area of the eye are these nodules typically observed? (Q792994)

A. Cornea


B. Conjunctiva

C. Iris

D. Retina

Correct Answer: C. Iris

Explanation: Correct Option C: Iris Koeppe's nodules are small, yellow-white nodules that appear on the surface of the iris observed in granulomatous uveitis like sarcoidosis, tuberculosis, Vogt-Kanayagi-Harada’s disease, and sympathetic ophthalmitis . They are typically located at the pupillary margin, which is the border between the iris and the pupil. These nodules are composed of collections of inflammatory cells. Incorrect Options: Options A, B, and D are incorrect. Koeppe’s nodules are present at the pupillary margin of the iris.



References:

Prepladder Notes

↑ *Back to this question in Part A*

@Neet_pg_bot

• [04_Glaucoma] 007 Glaucoma-Pacg Treatment Congenital Glaucoma

« Back to Folder Directory

[04_Glaucoma] 007 Glaucoma-Pacg Treatment Congenital Glaucoma — Quick Jump

Q1 (p. 626)	*Q2 (p. 627)*	*Q3 (p. 627)*	*Q4 (p. 628)*	*Q5 (p. 629)*	*Q6 (p. 630)*
Q7 (p. 631)	*Q8 (p. 633)*	*Q9 (p. 634)*	*Q10 (p. 635)*	*Q11 (p. 636)*	*Q12 (p. 638)*
Q13 (p. 639)	*Q14 (p. 641)*	*Q15 (p. 642)*	*Q16 (p. 643)*	*Q17 (p. 644)*	*Q18 (p. 645)*
Q19 (p. 646)	*Q20 (p. 647)*	*Q21 (p. 648)*	*Q22 (p. 648)*	*Q23 (p. 649)*	*Q24 (p. 651)*
Q25 (p. 652)					

Q1: A 65-year-old female presents with acute onset pain, redness, and blurring of vision in the right eye. On examination, her intraocular pressure (IOP) is 48 mmHg, and there is a shallow anterior chamber with a hazy cornea. Slit-lamp examination reveals an intumescent, swollen lens with the iris pushed forward. Gonioscopy shows a closed angle. What is the most likely underlying mechanism for her glaucoma? (Q853570)

A. Obstruction of the trabecular meshwork by inflammatory cells

B. Forward displacement of the lens due to an abnormal zonular attachment

C. Disruption of aqueous humor flow due to posterior synechiae

D. Pupillary block caused by the swollen lens

Correct Answer: D. Pupillary block caused by the swollen lens

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Lens-induced glaucoma is a type of secondary angle-closure glaucoma associated with a disorder of the lens. In this case, the enlarged lens prevents aqueous humor from flowing through the pupil into the anterior chamber, leading to a pupillary block. The iris is pushed forward, resulting in angle closure and increased intraocular pressure. Other options describe mechanisms for other forms of secondary glaucoma, such as uveitic glaucoma (Option A), phacodonesis (Option B), or chronic iritis (Option C) , but are not characteristic of lens-induced glaucoma.

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 377, 378, 393, 394.

https://www.ncbi.nlm.nih.gov/books/NBK574524/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Which of the following conditions is not typically associated with secondary congenital glaucoma? (Q853593)

A. Aniridia

B. Marfan Syndrome

C. Turner Syndrome

D. Angelman syndrome

Correct Answer: D. Angelman syndrome

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Angelman Syndrome (AS) is a rare neurogenetic disorder caused by loss of function of the UBE3A gene on chromosome 15. Ophthalmic findings that can be seen in Angelman syndrome include Strabismus, Refractive errors, Oculocutaneous albinism, It is not associated with secondary glaucoma. Secondary Developmental/ Congenital Glaucoma Associated Ocular Anomalies Associated Systemic Anomalies Glaucoma Associated with Iridodysgenesis: Aniridia (Option A ruled out) Familial Iris Hypoplasia Congenital Ectropion Uvea Congenital Microcornea Congenital Nanophthalmos Chromosomal Disorders: Trisomy 13 (Patau Syndrome) Trisomy 18 (Edward's Syndrome) Trisomy 21 (Down Syndrome) Turner Syndrome (Option C ruled out) Ectopia Lentis Syndromes: Marfan Syndrome (Option B ruled out) Weill-Marchesani Syndrome Homocystinuria Glaucoma Associated with Iridocorneal Dysgenesis: Posterior Embryotoxon Axenfeld-Rieger Syndrome Peter's Anomaly Phakomatosis : Sturge-Weber Syndrome Von Recklinghausen's Neurofibromatosis Metabolic Syndromes: Lowe's Syndrome Hurler's Syndrome Zellweger Syndrome

References:

emedicine.medscape.com/article/1206524-overview?form=fpf

www.aao.org/education/topic-detail/congenital-glaucoma-europe

Parsons Diseases of the Eye, 23rd Edition , Pages 483- 487

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 6-month-old infant with congenital glaucoma has corneal enlargement, clouding, and Haab's striae and the angle is difficult to visualize due to corneal clouding. Which surgical procedure is most appropriate for this infant? (Q853592)

A. Goniotomy

B. Trabeculotomy with Trabeculectomy

C. Trabeculectomy with antimetabolites

D. Glaucoma drainage devices

Correct Answer: B. Trabeculotomy with Trabeculectomy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Congenital glaucoma is diagnosed in this infant based on corneal enlargement , clouding , Haab's striae , and elevated intraocular pressure. When corneal clouding prevents proper angle visualization, Trabeculotomy with Trabeculectomy becomes the procedure of choice. Trabeculotomy helps expose Schlemm's canal to improve aqueous outflow, and trabeculectomy provides additional filtration, offering better intraocular pressure control. Goniotomy (Option A) : Involves making an incision in the angle of the eye and is best suited when the angle can be visualized clearly. Trabeculectomy with antimetabolites (Option C): While this is a common and effective procedure for many glaucoma cases, it might not be the first choice in this scenario due to the presence of corneal clouding and difficulty in initial angle procedures. Glaucoma Drainage Devices (Option D): Reserved for severe, resistant cases when other surgical methods have failed.

References:

Kanski Clinical Ophthalmology, 10th Edition, Page 401, 402, 403

Parsons Diseases of the Eye, 23rd Edition, Page 277, 278

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 1-month-old child presents with persistent eye watering and light sensitivity. Examination shows enlarged corneas with Haab's striae, cloudiness, a deep anterior chamber, and thin, blue sclera. The child has had ocular enlargement since birth. The ophthalmologist suspects congenital glaucoma. Which type of glaucoma best fits this presentation? (Q853591)

A. Juvenile Glaucoma

B. Infantile Glaucoma

C. True Congenital Glaucoma

D. Secondary Congenital Glaucoma

Correct Answer: C. True Congenital Glaucoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: True Congenital Glaucoma is characterized by elevated intraocular pressure present during intrauterine life, resulting in ocular enlargement at birth . The presence of Haab's striae, early onset, enlarged corneas, and the classic triad of symptoms (watering, photophobia, and blepharospasm) are all consistent with newborn glaucoma. Primary Congenital Glaucoma Types 1. Newborn glaucoma/ True congenital glaucoma : Typically presents within the first month of life. 2. Infantile (Option B ruled out) Affects infants older than 1 month and children up to 3 years. 3. Juvenile (Option A ruled out) Also known as Juvenile primary open angle glaucoma (POAG) usually occurs between 10 to 35 years of age. Secondary Congenital Glaucoma (Option D): Usually associated with other ocular or systemic anomalies, which is not indicated in this case.

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 401, 402, 403

Parsons' Diseases of the Eye, 23rd Edition, PDF Pages 277, 278

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 50-year-old glaucoma patient is prescribed a prostaglandin analog as first-line therapy. Which of the following is the most likely mechanism of action and a common side effect of this medication? (Q853590)

A. Reduces aqueous humor production; Lid retraction

B. Increases trabecular outflow; Brow ache

C. Increases uveoscleral outflow; Iris pigmentation

D. Decreases aqueous humor secretion; Allergic blepharoconjunctivitis

Correct Answer: C. Increases uveoscleral outflow; Iris pigmentation

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Increases uveoscleral outflow; Iris pigmentation Anti-Glaucoma Drugs: 1. Prostaglandin F2α Agonists (Option C) Examples: Latanoprost, Bimatoprost, Travoprost, Tafluprost Mechanism: Increases uveoscleral outflow Uses: First-line for primary open-angle and angle-closure glaucoma Side Effects: Sandy eyes, orbital fat atrophy, anterior uveitis, cystoid macular edema, heterochromia iridis, hypertrichosis Contraindications: Cystoid macular edema, diabetes mellitus 2. Beta Blockers (Option D) Non-Selective: Timolol, Levobunolol, Carteolol, Metipranolol More effective but contraindicated in asthma Selective: Betaxolol (retinoprotective), Levobetaxolol Safer in asthma Mechanism: Reduces aqueous secretion from ciliary epithelium Side Effects: Allergic blepharoconjunctivitis, dry eyes, corneal hypoesthesia 3. Alpha-2 Agonists (Option A) Examples: Brimonidine (long-term use), Apraclonidine (IOP spikes) Mechanism: Reduces aqueous production, increases uveoscleral outflow Side Effects: Lid retraction, respiratory depression (Brimonidine), allergic blepharoconjunctivitis Contraindications: Children <2 years Brimonidine: Safest during pregnancy (Category B) 4. Miotics (Option B) Examples: Pilocarpine, Physostigmine, Echothiophate Mechanism: Stimulates M3 receptors, increasing trabecular outflow Side Effects: Brow pain, miosis, cataract (Echothiophate), retinal tears 5. Carbonic Anhydrase Inhibitors Examples: Acetazolamide (systemic), Brinzolamide, Dorzolamide (topical) Mechanism: Inhibits aqueous secretion from ciliary epithelium Uses: Acute congestive glaucoma (with IV mannitol) Side Effects: Ciliary edema, secondary angle-closure glaucoma (Topiramate, Zonisamide) 6. Osmotic Agents Examples: Mannitol, Glycerol These work by increasing blood osmolarity, "sucking" water out of the vitreous to lower IOP quickly. Intravenous Mannitol: For acute attacks. Side effect: Pulmonary edema, congestive heart failure. Oral Glycerol: Sweeter, but contraindicated in Diabetics (metabolized to glucose). 6. Newer Drugs Netarsudil (Rho kinase inhibitor): Increases trabecular outflow; Side effect - vortex keratopathy Omidenepag (PGE2 agonist): Increases both trabecular and uveoscleral outflow

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 404 - 407

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: All of the following are penetrating glaucoma surgeries except: (Q853589)

A. Trabeculectomy

B. Deep sclerectomy

C. Minimum incision glaucoma surgery

D. Glaucoma drainage device implantation


Correct Answer: B. Deep sclerectomy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In penetrating surgeries an incision is made to create a new drainage pathway, while non-penetrating surgeries aim to enhance the eye’s natural drainage mechanism without incisions. Penetrating surgeries:

Trabeculectomy(Option A) : Glaucoma drainage device/seton surgery (Option D): Minimum incision glaucoma surgery (Option C) Non penetrating surgeries: Deep sclerectomy Viscocanalostomy



References:

Kanski Clinical Ophthalmology, 10th edition, Pages 416-422

Parsons' Diseases of the Eye, 23rd Edition, PDF Pages 725, 726

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: Which of the following laser procedures are used for Open angle glaucoma (OAG) and angle closure glaucoma (ACG) respectively? (Q853577)

A. Argon laser iridotomy for open-angle glaucoma and selective laser trabeculoplasty (SLT) for angle closure glaucoma

B. Selective laser trabeculoplasty for open-angle glaucoma and laser iridotomy for angle closure glaucoma

C. Laser iridoplasty for open-angle glaucoma and argon laser iridotomy for angle closure glaucoma


D. Laser iridotomy for open-angle glaucoma and laser iridoplasty for angle closure glaucoma


Correct Answer: B. Selective laser trabeculoplasty for open-angle glaucoma and laser iridotomy for angle closure glaucoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Selective Laser Trabeculoplasty (SLT) It is commonly used for open-angle glaucoma. SLT improves aqueous humor outflow by targeting the trabecular meshwork, which helps lower intraocular pressure over time. Laser Iridotomy It is used for angle closure glaucoma. This procedure creates a small hole in the peripheral iris to allow better fluid drainage and relieve the blockage of the anterior chamber angle. Argon laser iridotomy (Option A) is used for angle closure glaucoma, not open-angle glaucoma. Also, SLT is not typically used for angle closure glaucoma. Laser iridoplasty (Option C) is used for some cases of angle closure glaucoma, particularly when the angle is very narrow or if iridotomy is not feasible. However, argon laser iridotomy is the more common procedure for angle closure glaucoma. Laser iridotomy (Option D) is not used for open-angle glaucoma; it is used for angle-closure glaucoma. Similarly, laser iridoplasty is a treatment for angle closure glaucoma, not open-angle glaucoma.





References:

Kanski Clinical Ophthalmology, 10th edition, Pages 412-416

Parsons' Diseases of the Eye, 23rd Edition, PDF Pages 725, 726

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 65-year-old male presents to the ophthalmology clinic for a routine eye examination. During the examination, his intraocular pressure (IOP) is measured at 28 mmHg in both eyes. The ophthalmologist diagnoses him with primary open-angle glaucoma. Which of the following medications is primarily designed to lower intraocular pressure by reducing the production of aqueous humor? (Q853576)

A. Travoprost

B. Pilocarpine

C. Acetazolamide

D. Unoprostone

Correct Answer: C. Acetazolamide

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Medical management of Glaucoma Topical drugs Beta-blockers: Decrease the formation of aqueous humour. Eg: Timolol, betaxolol Alpha agonists: Have dual action i.e. they decrease the formation of aqueous and increase the trabecular outflow. Eg: Adrenaline/epinephrine, dipivefrin, brimonidine, and apraclonidine Miotics: These increase the trabecular outflow and open the angle blockage. Eg: Pilocarpine (Option B ruled out) Carbonic

anhydrase inhibitors: These decrease the formation of aqueous. Eg: Dorzolamide and Brinzolamide Newer anti-glaucoma drugs: Netarsudil : Increases the trabecular outflow and also decreases the aqueous production. PGF2 alpha agonist: It increases the uveoscleral outflow For eg- Latanoprost, bimatoprost, travoprost (Option A ruled out) , tafluprost, and unoprostone isopropyl (Option D ruled out) Systemic drugs Carbonic anhydrase inhibitor ■■■■■■■Eg: Acetazolamide Hyperosmotic agents: These draw aqueous out of the eye

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 369, 370, 374, 380

Parsons' Diseases of the Eye, 23rd Edition, PDF Pages 724, 725

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Which ICE syndrome is characterized by the presence of pigmented nodular lesions on the iris? (Q853575)

A. Progressive iris atrophy

B. Chandler's syndrome

C. Cogan-Reese syndrome

D. Both A & C

Correct Answer: C. Cogan-Reese syndrome

■ *Listen to Audio 1*

Explanation: Cogan-Reese Syndrome is known for its distinctive iris lesions. It is characterized by the presence of pigmented nodular lesions on the iris, which can be diffuse or localized. These pigmented lesions are a key feature that helps distinguish it from other ICE syndromes. Glaucomas Associated with Iridocorneal Endothelial (ICE) Syndromes: ICE Syndromes Include : Progressive iris atrophy Chandler's syndrome Cogan-Reese syndrome Pathogenesis : Abnormal corneal endothelial cells proliferate to form a membrane at the angle of the anterior chamber. Glaucoma is caused by secondary synechial angle closure due to contraction of this endothelial membrane. Clinical Features : Typically affects middle-aged women . Raised intraocular pressure (IOP) associated with specific features of the causative condition: Progressive iris atrophy : Iris shows corectopia, atrophy, and hole formation. (Options A & D ruled out) Chandler's syndrome : Mild or no iris changes, corneal edema predominates, even at normal IOP. (Option B ruled out) Cogan-Reese syndrome : Iris has pigmented nodular or diffuse lesions (also called iris nevus syndrome), and may have corneal changes. (Option C) Treatment : Glaucoma drainage device (artificial filtration shunt) may help control IOP. Medical treatment is often ineffective . Trabeculectomy surgery usually fails.

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 397, 398

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Which of the following statements is false about traumatic glaucoma? (Q853574)

A. Traumatic glaucoma can develop from both perforating and blunt injuries.

B. Angle recession glaucoma typically presents with elevated IOP shortly after the trauma.

C. Lens-induced glaucoma can occur due to a ruptured, swollen, or dislocated lens following trauma.


D. Peripheral anterior synechiae from corneal injury can lead to angle-closure glaucoma.

Correct Answer: B. Angle recession glaucoma typically presents with elevated IOP shortly after the trauma.


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Angle recession glaucoma often develops months to years after the initial blunt trauma due to long-term fibrosis of the trabecular meshwork. It is not typically associated with immediate elevated IOP. Traumatic Glaucoma: A secondary glaucoma may complicate perforating as well as blunt injuries. (Option A ruled out) Causes: Inflammatory Glaucoma: Results from iridocyclitis. Intraocular Hemorrhage: Hyphaema inside the eye leading to glaucoma. Hyphema frequently causes acute intraocular pressure elevation by blocking the trabecular meshwork (TM). Ghost Cell Glaucoma: RBCs lose their iron and Hb → becomes ghost cells Occurs due to haemorrhage in vitreous Lens-Induced Glaucoma: Due to a ruptured, swollen, or dislocated lens. (Option C ruled out) Angle-Closure: Caused by peripheral anterior synechiae formation after corneal injury. (Option D ruled out) Angle Recession Glaucoma : Due to angle recession (The recession is due to a tear in the ciliary body) from blunt trauma after years (10yrs), leading to fibrosis of the trabecular meshwork over time. (Option B) {{caption_text}} {{caption_text}} Management : Medical Therapy : Includes topical 0.5% timolol and oral acetazolamide. Treatment of Underlying Causes : For instance, using atropine and steroids to control inflammation. Surgical Intervention : Based on the specific cause and situation.




Subluxation of the lens



Total hyphema

@Neet_pg_bot



@Neet_pg_bot

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 395, 396, 397

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A 30-year-old male presents with unilateral eye discomfort, blurred vision, and seeing halos around lights. On examination, his intraocular pressure (IOP) is measured at 48 mm Hg. The anterior segment shows a few small, flat, nonpigmented keratic precipitates located inferiorly, mild redness, and an aqueous flare. The pupil is dilated, but there are no signs of a posterior synechiae. The patient reports recurrent episodes that resolve within days to weeks. Given these findings, what is the most likely diagnosis? (Q853573)

A. Secondary Angle-Closure Glaucoma with Pupillary Block

B. Secondary Angle-Closure Glaucoma without Pupillary Block

C. Open-Angle Glaucoma Due to Uveitis

D. Glaucomatocyclitic Crisis

Correct Answer: D. Glaucomatocyclitic Crisis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Glaucomatocyclitic Crisis (Posner-Schlossman Syndrome) is characterized by very high IOP (often between 40 and 50 mm Hg), minimal inflammation, a few small keratic precipitates, no posterior synechiae, and recurrent episodes that resolve within days to weeks. It is commonly seen in young males. This can be due to HLA-BW54 association has been documented. CMV or H. Pylori infection. The treatment involves controlling IOP with anti-glaucoma medications, followed by steroid therapy to manage the uveitis. {{caption_text}} Secondary Angle-closure glaucoma with a pupillary block (Option A): Caused by pupillary block due to annular synechiae, leading to iris bombe and subsequent angle closure. Secondary Angle-closure glaucoma without a pupillary block (Option B) : Results from inflammatory debris organizing in the angle, which pulls the iris over the trabecular meshwork, causing progressive angle closure and elevated IOP. Open-angle glaucoma (Option C) : This is caused by the blocking/inflammation of the trabecular meshwork by aqueous cells.



References:

Kanski Clinical Ophthalmology, 10th edition, Pages 379, 380, 392, 393

Parsons' Diseases of the Eye, 23rd Edition, Pages 708, 709

Posner-Schlossman Syndrome - StatPearls - NCBI Bookshelf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 70-year-old patient with a history of glaucoma undergoes a trabeculectomy to control intraocular pressure. Postoperatively, the patient presents with severe eye pain, blurred vision, a significantly elevated intraocular pressure (IOP), and a shallow anterior chamber. Seidel's test is negative. What is the most appropriate initial treatment for this condition? (Q853572)

A. Oral corticosteroids

B. Pars plana Vitrectomy

C. Atropine drops and oral acetazolamide


D. Peripheral iridectomy

Correct Answer: C. Atropine drops and oral acetazolamide

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical scenario describes a 70-year-old patient who, following trabeculectomy , experiences severe eye pain and blurred vision, elevated intraocular pressure (IOP), a shallow anterior chamber, a negative Seidel's test, which are characteristic of ciliary block glaucoma (malignant glaucoma) , also known as Aqueous Misdirection Syndrome . This condition occurs due to a mechanical blockage of aqueous humor flow, often after intraocular surgery. {{caption_text}} Pathogenesis: Aqueous Misdirection: The aqueous humor is diverted posteriorly into or behind the vitreous body rather than flowing into the anterior chamber, as the ciliary processes, lens, and anterior vitreous face form a 'ciliary block' that traps the fluid in the posterior segment. Pressure Build-up: As aqueous accumulates in the vitreous cavity, the volume of the posterior segment increases. Forward Displacement: This increased posterior pressure pushes the lens-iris diaphragm forward . Angle Closure: The forward displacement causes the anterior chamber to become uniformly shallow (both centrally and peripherally), leading to a secondary angle closure and a massive spike in IOP. Management: Medical Therapy (Initial): Atropine Drops/Ointment (1%): To dilate the ciliary ring and relieve ciliary block. Acetazolamide (250 mg QID): To reduce aqueous humor production. Timolol Maleate (0.5%): To decrease aqueous production. Intravenous Mannitol: To reduce vitreous gel volume and decrease IOP. Laser Therapy: YAG laser hyaloidotomy for aphakic and pseudophakic patients. Surgical Therapy: Pars plana vitrectomy with or without lensectomy if medical therapy fails within 4-5 days. (Option B ruled out)



References:

Kanski Clinical Ophthalmology, 10th edition, Pages 381, 382, 415, 419, 420, 421

Parsons’ Diseases of the Eye, 23rd Edition, PDF Page 717

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 58-y/o diabetic patient presents with decreased vision, eye pain, and redness in the right eye. Examination reveals iris neovascularization, elevated intraocular pressure, and proliferative diabetic retinopathy. What is the most appropriate management strategy? (Q853571)

A. Topical beta-blockers and oral carbonic anhydrase inhibitors

B. Panretinal photocoagulation


C. Systemic corticosteroids

D. VEGF inhibitors

Correct Answer: B. Panretinal photocoagulation

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Based on these findings, the diagnosis of neovascular glaucoma is supported by the presence of iris neovascularization and elevated IOP in the context of proliferative diabetic retinopathy. Panretinal photocoagulation is the most appropriate treatment as it targets the underlying cause of neovascular glaucoma. This treats the retinal ischemia, reducing the stimulus for abnormal blood vessel growth and neovascularization. Neovascular Glaucoma (NVG) {{caption_text}} Treatment : Panretinal Photocoagulation or Cryotherapy : Aimed at preventing further neovascularization. Anti-VEGF Agents : Used as an adjunctive treatment. Help reduce neovascularization. (Option D) Trabeculectomy : A surgical procedure to lower intraocular pressure, often combined with antifibroblastic agents. Glaucoma Drainage Device : Artificial filtration shunt (Seton operation) can help manage IOP. Causes : Common : Proliferative diabetic retinopathy (M/C) Central retinal vein occlusion Sickle-cell retinopathy Eales' disease Rare : Chronic intraocular inflammation Intraocular tumors Long-standing retinal detachment Central retinal artery occlusion Topical beta-blockers and oral carbonic anhydrase inhibitors (Option A): These medications can help manage intraocular pressure (IOP) but do not address the underlying cause of neovascular glaucoma, which is the retinal ischemia and resulting neovascularization. Systemic corticosteroids (Option C): While corticosteroids can reduce inflammation, they do not address neovascularization or the elevated IOP associated with neovascular glaucoma.



References:

Kanski Clinical Ophthalmology, 10th Edition, Pages 387, 388, 389

Parsons' Diseases of the Eye, 23rd Edition, Page 710

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A 55-year-old Asian woman presents with acute eye pain and blurred vision. Her ophthalmologist suspects primary angle closure glaucoma (PACG). Which of the following statements about the etiopathogenesis of PACG is incorrect? (Q853548)

A. Pupillary block is the most common mechanism in PACG

B. Phacomorphic mechanism involves an abnormally large lens pushing the iris forward

C. Plateau iris configuration is more common in younger patients with PACG

D. Caucasians have a higher prevalence of PACG compared to other ethnicities

Correct Answer: D. Caucasians have a higher prevalence of PACG compared to other ethnicities

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: PACG is more common in Asians compared to all other ethnicities. Mechanisms of PACG 1. Pupillary block (70% of cases) (Option A ruled out): Most common mechanism (70% of cases) Increased contact between iris and lens 2. Plateau iris configuration: Anteriorly positioned ciliary processes More common in

younger patients (Option C ruled out) 3. Phacomorphic mechanism (Option B ruled out): Swollen lens contributes to angle closure Can cause pupillary block or push peripheral iris forward




Fig. 11.34 Mechanism of angle closure. **(A)** Relative pupil block; **(B)** iris bombé; **(C)** iridocorneal contact @Neet_pg_bot

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 372 - 374

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 72-year-old male presents with diminution of vision in the right eye. On slit-lamp examination, dandruff-like material is noted on the anterior lens capsule and iris. Intraocular pressure is elevated, and the patient is diagnosed with pseudoexfoliative glaucoma. Which of the following is most likely to be observed in this patient's eye? (Q853569)

A. Posterior synechiae

B. Zonular instability and phacodonesis

C. Corneal guttata

D. Iris heterochromia

Correct Answer: B. Zonular instability and phacodonesis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Zonular instability is a hallmark of pseudoexfoliation syndrome. This can lead to phacodonesis (lens wobbling). Ocular manifestation of Pseudoexfoliation Syndrome (XFS) is characterized by the accumulation of fibrillar, proteinaceous material on various ocular structures, including the lens, iris, and trabecular meshwork. This material can be seen as flaky white deposits on the anterior lens capsule and pupillary margin, often with a distinct clear zone at the center of the lens. It causes secondary open-angle glaucoma, known as pseudoexfoliative glaucoma. Pathogenesis of Pseudoexfoliation Syndrome: Caused by abnormal extracellular matrix metabolism leading to production of pseudoexfoliative fibrillar material . Strongly associated with polymorphism of the LOXL1 gene , which is involved in elastin fiber formation . Oxidative stress and impaired degradation lead to accumulation of this material. Deposits occur on the lens capsule, iris, zonules, and trabecular meshwork , which may cause secondary open-angle glaucoma (Pseudoexfoliation Glaucoma) . Posterior synechiae (Option A) is incorrect because while inflammation can lead to synechiae, it is not a primary feature of pseudoexfoliation syndrome. Corneal guttata (Option C) and Iris heterochromia (Option D) are associated with Fuchs' endothelial dystrophy, not pseudoexfoliation syndrome. Iris heterochromia (Option D) is typically seen in Fuchs' heterochromic iridocyclitis, not pseudoexfoliation syndrome.

References:

Yanoff and Duker's Ophthalmology, 5th Edition, Page 1074-1077

Kanski Clinical Ophthalmology, 10th Edition, Page 382, 383, 384

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A 25-year-old myopic male presents with intermittent blurring of vision and mild ocular pain after exercise. On examination, IOP is 28 mmHg, Slit lamp examination reveals Krukenberg spindle on corneal endothelium, trabecular pigmentation on gonioscopy. Which of the following is the first-line treatment? (Q853568)

A. Nd:YAG laser peripheral iridotomy

B. Selective laser trabeculoplasty

C. Topical prostaglandin analogue


D. Argon laser peripheral iridoplasty

Correct Answer: C. Topical prostaglandin analogue

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This patient's presentation is consistent with pigmentary glaucoma . The key features are: Young, myopic male Elevated IOP Krukenberg spindle (vertical line of pigment on corneal endothelium) Heavily pigmented trabecular meshwork Iris transillumination defects Symptoms exacerbated by exercise (due to pupillary movement causing increased pigment release) Treatment is along the lines of POAG management, which initially includes medical treatment, with prostaglandin analogues often being the first choice due to their IOP-lowering efficacy and convenient dosing schedule. (Option C) Nd:YAG laser peripheral iridotomy (Option A) i s generally reserved for cases with pronounced iris concavity or those not responding to medical therapy. Selective laser trabeculoplasty (SLT) (Option B) is typically considered after initial medical therapy or in combination with it. Argon laser peripheral iridoplasty (Option D) is primarily used in angle-closure glaucoma to open the angle.



References:

Kanski Clinical Ophthalmology, 10th edition, Pages 385, 386, 387

pubmed.ncbi.nlm.nih.gov/1495789/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A 45-year-old male presents with unilateral eye pain, redness, and moderately elevated intraocular pressure (IOP) of 28 mmHg. On slit-lamp examination, you observe fine, pigmented deposits on the corneal endothelium in a vertical spindle pattern. The patient O/E has symmetric transillumination defects in the mid-peripheral iris. Based on these findings, which of the following is the most likely cause of this patient's secondary glaucoma? (Q853557)

A. Pseudoexfoliation syndrome

B. Pigment dispersion syndrome

C. Neovascular glaucoma

D. Lens-particle glaucoma

Correct Answer: B. Pigment dispersion syndrome

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Fine pigmented deposits in a vertical spindle pattern on corneal endothelium (Krukenberg spindle), mid-peripheral iris transillumination defects (symmetric), moderately raised IOP point strongly to pigment dispersion syndrome (PDS). Pigment dispersion syndrome occurs when pigment granules from the iris epithelium are released into the anterior chamber and deposit on various structures, including the corneal endothelium and trabecular meshwork. This can lead to trabecular outflow obstruction and subsequent glaucoma. Symmetric Iris transillumination defects in the mid-peripheral iris, is characteristic of Pigment Dispersion Syndrome (PDS) . Pseudoexfoliation syndrome (Option A) presents with whitish, dandruff-like material on the lens capsule and pupillary margin. Neovascular glaucoma (Option C) is characterized by new blood vessel growth over the iris and anterior chamber angle, often secondary to ischemic retinal diseases. Lens-particle glaucoma (Option D) occurs when lens material enters the anterior chamber, usually after cataract surgery or lens injury. Pigmentary Dispersion vs Pseudoexfoliation Syndrome: Feature Pigmentary Dispersion (PDS) Pseudoexfoliation (PEX) Typical Patient Young, myopic (nearsighted) males. Elderly patients (>60 years). Iris Finding Mid-peripheral transillumination defects (spoke-like). Pupillary margin atrophy/transillumination. Lens Finding No abnormal material; pigment on the back of the lens. Three-zone dandruff-like material on the anterior capsule. Cornea Krukenberg spindle (vertical pigment line on endothelium). Diffuse pigment or PEX material on endothelium. Zonules Normal. Weak/Fragile (risk of lens subluxation). Gonioscopy Dense, homogenous pigment (360°). Patchy pigment + Sampaolesi Line .

References:

Kanski Clinical Ophthalmology, 10th Edition, Page 382, 385, 387

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A 65-year-old male presents with a painful, blind right eye. On examination, the eye is stony hard with no light perception, 100% cupping of the optic disc, and a fixed, mid-dilated pupil. Which of the following is the correct initial management? (Q853556)

A. Topical timolol and brimonidine

B. Trabeculectomy

C. Cyclodestructive procedure

D. Evisceration

Correct Answer: C. Cyclodestructive procedure

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Absolute glaucoma is the final phase of Primary Angle Closure Glaucoma. Management of absolute glaucoma In absolute glaucoma, the goal of treatment is to relieve pain and reduce IOP, as vision cannot be restored. Retrobulbar alcohol injection - given to relieve pain. It suppresses the ciliary ganglion. A cyclodestructive procedure , such as cyclocryotherapy or cyclophotocoagulation, is the most appropriate management option in this scenario as it reduces aqueous production by damaging the secretory ciliary processes aiming to reduce IOP. (Option C) If pain persists despite cyclodestruction, enucleation/ evisceration may be considered. (Option D)

References:

pmc.ncbi.nlm.nih.gov/articles/PMC3669490/

Kanski Clinical Ophthalmology, 10th edition, Pages 379, 380, 381.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: All of the following components of Vogt's triad are typically observed in a patient who has experienced and resolved an acute primary angle-closure glaucoma (APAC) attack except? (Q853555)

A. Glaucoma fleken

B. Iris atrophy

C. Pigment dispersion

D. Progressive retinal damage

Correct Answer: D. Progressive retinal damage

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Vogt's triad is a clinical set of findings typically associated with acute primary angle-closure glaucoma (APAC) after an attack has been resolved. The triad helps in identifying damage caused during the acute episode. Glaucoma flecken (Opacities develops d/t necrosis of anterior epithelial cells of the lens) (Option A) Patches of iris atrophy (Option B) Dispersion of pigment on cornea (Option C)

References:

Parson’s Textbook of Ophthalmology, 23rd Edition, Page 704

https://journals.lww.com/ijog/fulltext/2023/03020/vogt_s_triad___do_we_find_the_triad_all_the_time_.118.aspx

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: A 55-year-old female presents for a routine eye examination. During gonioscopy, the examination reveals an occludable anterior chamber angle bilaterally. The patient reports no visual symptoms, and intraocular pressure (IOP) measurements are within normal limits. There is no evidence of glaucomatous damage on optic nerve examination. What is the most appropriate next step in management for this patient? (Q853554)

A. I.V Mannitol/Acetazolamide

B. Prophylactic peripheral iridotomy

C. Filtration surgery

D. Topical beta-blocker therapy

Correct Answer: B. Prophylactic peripheral iridotomy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Primary Angle Closure Suspect (PACS) is characterized by an occludable anterior chamber angle as seen during gonioscopy but without evidence of increased IOP or glaucomatous damage. This indicates a risk for potential future angle closure but does not yet present as acute angle-closure glaucoma. PACS: Primary Angle Closure Suspect Iridotrabecular contact or occludable angle of more than 3 quadrants on gonioscopy No peripheral anterior synechiae Intraocular pressure is normal Eclipse sign of PACS - convex iris lens diaphragm Treatment :

Prophylactic Peripheral iridotomy (Option B) is done in both eyes with lens extraction using NdYAG laser

References:

Kanski Clinical Ophthalmology, 10th Edition, Page 375, 376, 381, 382

Parson's Textbook of Ophthalmology, 23rd Edition, Page 702, 706

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: What aspect of the anterior chamber does the Van Herick grading system evaluate during slit lamp examination? (Q853553)

A. The curvature of the cornea

B. The width of the anterior chamber angle

C. The condition of the retina

D. The intraocular pressure

Correct Answer: B. The width of the anterior chamber angle

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The VAN Herick technique is an eye examination method used to determine the width of the anterior chamber angle of the eye using the ratio of the anterior chamber depth and corneal section which gives the status of the angle as per the chart given below. Helps screen for narrow angles / risk of angle-closure glaucoma Width of the empty space (LACD) as compared to the corneal thickness VAN Herick Grade Angle status No black space observed 0 Closed <¼ corneal thickness 1 Extremely narrow ¼ of corneal thickness 2 Narrow >¼ to ½ of corneal thickness 3 Open ≥ 1/2 of corneal thickness 4 Wide open LACD = Limbal anterior Chamber depth Curvature of cornea (Option A): Assessed by keratometry/topography Condition of retina (Option C): Fundus examination Intraocular pressure (Option D): Measured by tonometry

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 375, 376

Interobserver reliability when using the Van Herick method to measure anterior chamber depth - PMC (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: In the Shaffer classification system for grading the anterior chamber angle, which grade indicates that the angle is fully open with visible structures such as the scleral spur and trabecular meshwork? (Q853552)

A. Grade 0

B. Grade 1

C. Grade 2

D. Grade 3

Correct Answer: D. Grade 3

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Shaffer system is one of the various methods used to grade the angle during the gonioscopy. The angle in degree between two imaginary lines is tangential to the inner surface of trabecular meshwork and the interior surface of the iris (1/3rd distance from the periphery) ACA in degrees ACA grade Angle status Visible structures 0 0 Closed No structures visible </= 10 1 Extremely narrow Schwalbe's line 11-19 2 Narrow Trabecular meshwork 20-34 3 Open Scleral spur 35-45 4 Wide open Ciliary body ACA= Anterior Chamber Angle

References:

Kanski Clinical Ophthalmology, 10th edition, Page 388

https://www.researchgate.net/figure/ACA-grading-system-as-described-by-Shaffer_tbl2_276494875

https://www.researchgate.net/publication/360211495_Comparison_of_Iridocorneal_Angle_Assessments_in_Open Angle_Glaucoma_and_Ocular_Hypertension_Patients_Anterior_Segment_Optical_Coherence_Tomography_and _Gonioscopy

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Which of the following best describes the principle of gonioscopy? (Q853551)

A. It involves measuring intraocular pressure.

B. It is a technique for visualizing the anterior chamber angle of the eye using a special lens


C. It is a technique for visualizing the posterior chamber angle of the eye using a special lens


D. It evaluates the retina.

Correct Answer: B. It is a technique for visualizing the anterior chamber angle of the eye using a special lens

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Gonioscopy is the main method used to measure the angle of the anterior chamber The angled structure is not visible to the examiner because of the total internal reflection inside the anterior chamber. Principle: It overcomes total internal reflection Critical angle = 46 degree A Gonioscope with the same refractive index as the cornea and mirror is placed over the cornea. Light from the anterior chamber strikes the cornea and travels through and gets reflected by the mirror in the gonioscope, thus reaching our eye. Measuring intraocular pressure is done with a tonometer, not a gonioscopy. (Option A) It is a technique for visualizing the anterior chamber angle of the eye using a special lens and not the posterior chamber. (Option C) The retina is evaluated with a fundus camera or other retinal imaging techniques, not gonioscopy. (Option D)





@Neet_pg_bot

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 31, 32, 33

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: A 35-year-old woman presents with intermittent eye pain and blurred vision. Gonioscopy reveals narrow angles, and the angle remains narrow after laser peripheral iridotomy. Which of the following statements about plateau iris configuration is incorrect ? (Q853550)

A. It is characterized by an anteriorly positioned ciliary body

B. It is more commonly seen in older patients with primary angle closure glaucoma

C. Plateau iris configuration can persist after iridotomy


D. Ultrasound biomicroscopy (UBM) can be useful in diagnosing this condition

Correct Answer: B. It is more commonly seen in older patients with primary angle closure glaucoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Plateau iris configuration is actually more common in younger patients with primary angle closure glaucoma, not older patients. Plateau Iris Configuration Anatomy Plateau iris is characterized by an anteriorly positioned ciliary body, which pushes the peripheral iris forward. (Option A) This creates a "table-top" or plateau-like appearance of the iris. Diagnostic tools Ultrasound biomicroscopy (UBM) is particularly useful in diagnosing plateau iris configuration. (Option D) Management Due to its persistence after iridotomy, patients with plateau iris may require additional treatments such as: (Option C) Argon laser peripheral iridoplasty Long-term use of miotic drops {{caption_text}}



References:

Pubmed

Kanski Clinical Ophthalmology, 10th edition, Pages 375, 376, 377

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: A 50-year-old patient presents with symptoms suggesting possible angle-closure glaucoma. You want to perform a comprehensive assessment of the anterior chamber angle. Which of the following statements is incorrect regarding anterior chamber

angle assessment and anatomy? (Q853549)

A. Gonioscopy is the definitive diagnostic tool for primary angle closure.

B. Normal angle structures from anterior to posterior include: Schwalbe's line, non-pigmented trabecular meshwork, pigmented trabecular meshwork, scleral spur, and ciliary body band.

C. An angle is considered occludable if the pigmented trabecular meshwork is not visible in less than 180 degrees without indentation.


D. Indirect gonioscopes used to measure angle of AC are goldmann, posner-ritch, zeiss

Correct Answer: C. An angle is considered occludable if the pigmented trabecular meshwork is not visible in less than 180 degrees without indentation.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Correct answer: C) An angle is considered occludable if the pigmented trabecular meshwork is not visible in less than 180 degrees without indentation. Explnation: An angle is considered occludable if the posterior trabecular meshwork is not visible in more than 180 degrees , not less than 180 degrees. Normal angle structures from anterior to posterior (Option B ruled out): Schwalbe's line Non-Pigmented trabecular meshwork Pigmented Trabecular meshwork Scleral spur Ciliary body band Iris {{caption_text}} Gonioscopy is the definitive diagnostic tool for primary angle closure. (Option A) There are 2 types of gonioscopy lenses: Indirect (Option D ruled out) Direct Provides a mirror image of the opposite angle Provides a direct view of the angle 1. Goldmann lenses 2. Zeiss 3. Susman 4. Posner – Ritch 1. Koeppe 2. Swan – Jacob



References:

Pubmed

Parson’s Textbook of Ophthalmology, 23rd Edition, Page 702

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7759936/

↑ *Back to this question in Part A*

• [04_Glaucoma] 008 Glaucoma - Poag Tonometry Perimetry

« Back to Folder Directory

[04_Glaucoma] 008 Glaucoma - Poag Tonometry Perimetry — Quick Jump

Q1 (p. 655) *Q2 (p. 660)* *Q3 (p. 661)* *Q4 (p. 663)* *Q5 (p. 664)* *Q6 (p. 664)*
Q7 (p. 668) *Q8 (p. 669)* *Q9 (p. 671)* *Q10 (p. 673)* *Q11 (p. 676)* *Q12 (p. 677)*
Q13 (p. 678) *Q14 (p. 679)* *Q15 (p. 680)* *Q16 (p. 681)* *Q17 (p. 682)* *Q18 (p. 683)*
Q19 (p. 684)

Q1: Match the following: 1) Perkins Tonometer a) {{caption_text}} 2) Tonopen b) {{caption_text}} 3) Pulsair tonometer c) {{caption_text}} 4) Rebound Tonometer d) {{caption_text}} (Q853862)

A. 1-a, 2-c, 3-d, 4-b

B. 1-c, 2-b, 3-d, 4-a

C. 1-d, 2-a, 3-b, 4-c

D. 1-c, 2-d, 3-a, 4-b




@Neet_pg_bot



@Neet_pg_bot



Correct Answer: C. 1-d, 2-a, 3-b, 4-c

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Perkin's applanation tonometer Hand held version of Goldmann. {{caption_text}} Pulsair A noncontact device that measures intraocular pressure. {{caption_text}} Tonopen Used in scarred and irregular cornea It is an applanation tonometer Tonometer of choice in children {{caption_text}} Rebound tonometer Used for self-measurement {{caption_text}} Goldmann applanation tonometer Gold standard for intraocular pressure measurement





Pulsair tonometer

@Neet_pg_bot



Tonopen

@Neet_pg_bot





References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Pages 29, 30

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 62-year-old man with a history of Glaucoma undergoes visual field testing to assess his peripheral vision. The test involves moving a stimulus of constant size and intensity from a non-seeing area into a seeing area to determine the boundaries of his visual field. Which type of perimetry is being described? (Q853871)

A. Goldmann perimetry

B. Automated perimetry

C. Frequency doubling perimetry


D. Short-wavelength automated perimetry

Correct Answer: A. Goldmann perimetry

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Goldmann perimetry is a type of manual kinetic perimetry. In this technique, a stimulus of constant size and intensity is moved from a non-seeing area (usually the far periphery) towards the center of vision until the patient detects it. This method allows for a detailed mapping of the visual field boundary. {{caption_text}} Various methods of kinetic perimetry: Automated perimetry (Option B): Static perimetry where stationary stimuli of varying intensities are presented at fixed locations in the visual field. It doesn't involve moving stimuli. Frequency doubling perimetry (Option C): Static perimetry that uses a specific stimulus (a sinusoidal grating) to test contrast sensitivity and is particularly useful for detecting early glaucomatous damage. SWAP (Option D): Static perimetry technique that uses a blue light stimulus on a yellow background to selectively test the function of blue-yellow ganglion cells, which may be affected early in glaucoma.



References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Pages9,10

Parson’s Diseases of the Eye, 23rd Edition, Pages 243, 244, 245, 246, 247, 248

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 55-year-old patient with primary open-angle glaucoma undergoes visual field testing. Which of the following visual field defects is considered the earliest clinically significant field defect occurring in glaucoma? (Q853870)


Correct Answer: Unknown


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The earliest clinically significant field defect to appear in patients of glaucoma is small of fixation point ring-shaped paracentral scotoma . It may appear either below or above the blind spot in Bjerrum's area (an arcuate area extending above and below the blind spot between 10° and 20° Visual Field Defects in Primary Open Angle Glaucoma 1. Isopter contraction Mild generalized constriction of central as well as peripheral field 2. Baring of blind spot Early glaucomatous change, but nonspecific and of limited diagnostic value 3. Small ring-shaped paracentral scotoma Earliest clinically significant field defect; appears in Bjerrum's area 4. Seidel's scotoma Paracentral scotoma joins the blind spot to form a sickle-shaped defect 5. Arcuate or Bjerrum's scotoma Formed at a later stage by extension of Seidel's scotoma in an area above or below the fixation point 6. Ring or double arcuate scotoma Develops when the two arcuate scotomas join together 7. Roenne's central nasal step An area of depression above or below the horizontal meridian on the nasal side. This is due to the two arcuate scotomas running in different arcs and meet to form a sharp right-angled gap. 8. Peripheral nasal step Appear

sometimes at an early stage; peripheral nasal step results from unequal contraction of the peripheral isopter 9. Advanced glaucomatous field defects Visual field loss gradually spreads peripherally, eventually only the central and temporal island of vision is left behind.





References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Pages 366, 367

Parson’s Diseases of the Eye, 23rd Edition, Pages 686, 687, 688

https://www.ncbi.nlm.nih.gov/books/NBK441887/#:~:text=an%20area%20of%20depression%20above%20or%20below%20the%20horizontal%20meridian%20on%20the%20nasal%20side

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 55-year-old patient undergoes a comprehensive eye examination including visual field testing. Which of the following statements is incorrect? (Q853869)

A. A scotoma is an area of decreased vision surrounded by normal vision

B. An isopter is a line connecting points of equal light sensitivity and size detection in the visual field

C. The blind spot is a type of absolute scotoma

D. The visual field extends maximally in the superior direction

Correct Answer: D. The visual field extends maximally in the superior direction

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Visual field is the peripheral vision. It extends maximally in the temporal direction and minimally in the superior direction . The extent of the visual field: (mnemonic: TINS) Temporally: 90° to 100° Inferiorly: 70° Nasally: 60° Superiorly: 50° Scotoma (Option A): Area of decreased vision surrounded by normal vision. Isopter (Option B): Line connecting points of equal light sensitivity and size detection in the visual field. Physiological blind spot (Option C): Corresponds to the area where the optic nerve exits the eye, which lacks photoreceptors. It is a type of absolute scotoma.

References:

Visual Field Defects- Pubmed

Neuroanatomy- Visual pathway

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 362-364

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 58-year-old patient is being evaluated for suspected primary open-angle glaucoma. The ophthalmologist wants to perform a thorough examination of the optic nerve head to detect glaucomatous changes. Which of the following techniques is considered the examination of choice for evaluating fundus changes in this patient? (Q853868)

A. Direct ophthalmoscopy

B. Slit-lamp biomicroscopic examination with non-contact lens

C. Optical coherence tomography (OCT)

D. Indirect ophthalmoscopy

Correct Answer: B. Slit-lamp biomicroscopic examination with non-contact lens

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The examination of choice for detecting fundal changes in a glaucoma suspect is Slit-lamp biomicroscopic examination with non-contact lens . The non-contact lenses of +78 or +90 dioptres are commonly used. Direct Ophthalmoscopy (Option A) No depth perception as the examiner uses only one eye. Indirect Ophthalmoscopy (Option D) Magnification is less, hence it is not used to examine the disc and macula. OCT (Optical coherence tomography) (Option C) Useful method to measure optic disc diameter, CDR, NRR thickness and optic disc area

References:

US Ophthalmic Review

Kanski's Clinical Ophthalmology: A Systematic Approach, 9th Edition, Pages 21-27

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 62-year-old woman with a family history of glaucoma presents with mild visual disturbances. Her intraocular pressure is measured to be 24 mmHg in both eyes. On dilated fundus examination, the following findings are noted: The cup-to-disc ratio of 0.5 Splinter hemorrhages at the optic disc margin Wedge-shaped retinal nerve fiber layer defects on red-free light examination What stage of glaucomatous optic disc changes is most likely present in this patient? (Q853867)

A. Glaucomatous optic atrophy

B. Advanced glaucomatous changes

C. Early glaucomatous changes

D. Physiological cupping


Correct Answer: C. Early glaucomatous changes


■ *Listen (English)*

■ *Listen (Hindi)*


Explanation: Key findings: IOP = 24 mmHg → mildly elevated Cup–disc ratio = 0.5 → borderline/moderate Splinter (Drance) hemorrhages → early sign of glaucoma Wedge-shaped RNFL defects → early structural damage These are characteristic of early glaucomatous optic nerve damage , before marked cupping or severe vision loss occurs. Early glaucomatous changes are indicated by signs such as: Cup-to-disc ratio of >0.3. Asymmetry of the cups : A cup-to-disc ratio difference greater than 0.2 between two eyes. Splinter hemorrhages near the optic disc margin. Retinal nerve fiber layer (RNFL) defects: Wedge-shaped dark areas seen on the red-free light examination - an early indicator of glaucomatous damage. Advanced glaucomatous changes Cup-to-disc ratio of 0.7–0.9 Significant neuroretinal rim thinning Vessel bayonetting Glaucomatous optic atrophy Complete destruction of the neural tissue Deeply excavated and white optic disc. Cavernous optic atrophy - large caverns or lacunae are formed due to: Raised IOP leading to modulation of lamina cribrosa. Ischaemic atrophy of the nerve fibers without a corresponding increase of supporting glial tissue. Normal eye (no glaucoma) Early glaucoma Moderate glaucoma Advanced glaucoma








RNFL Thickness Map
350
175
0 µm
Extracted Horizontal Tomogram
T N
Extracted Vertical Tomogram
T N
@Neet_pg_bot






30°
30°
@Neet_pg_bot



RNFL Thickness Map
350
175
0 µm
Extracted Horizontal Tomogram
T N
Extracted Vertical Tomogram
T N
@Neet_pg_bot





30°
30°
@Neet_pg_bot



RNFL Thickness Map
350
175
0 µm


Extracted Horizontal Tomogram
T
N


Extracted Vertical Tomogram
T
N
@Neet_pg_bot




30°
30°
@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology: A Systematic Approach, 9th Edition, Pages 353, 354, 355, 369

Parson's Diseases of the Eye, 23rd Edition, Pages 695, 696

https://glaucoma.org.au/what-is-glaucoma/testing-for-glaucoma/stages-of-glaucoma-progression

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 52-year-old male presents with progressive vision loss. Fundoscopic examination reveals a vertical elongation of the optic disc and diffuse thinning of the retinal nerve fiber layer (RNFL). Imaging confirms significant damage to the peripapillary region. Which of the following best describes the characteristic peripapillary change seen in glaucomatous optic neuropathy? (Q853866)

A. Increased retinal pigment epithelium (RPE) density in the peripapillary area


B. Presence of alpha zone peripapillary atrophy (PPA)

C. Thickening of peripapillary RNFL on optical coherence tomography (OCT)

D. Uniform retinal nerve fiber layer thickness without focal loss

Correct Answer: B. Presence of alpha zone peripapillary atrophy (PPA)

Explanation: Peripapillary atrophy (PPA) in the region around the optic nerve head - is indicative of early damage in glaucoma. It has 2 zones: Alpha zone (outer): Characterized by irregularities in the retinal pigment epithelium (RPE). Larger in size and more commonly observed in glaucomatous eyes. Beta zone (inner): Chorioretinal atrophy. A visible white band of exposed sclera adjacent and distinct to the scleral rim. Larger and more frequently seen in glaucoma. Considered a risk factor for glaucoma progression. The location of the beta zone often correlates with patterns of visual field loss in glaucoma patients. {{caption_text}} Increased retinal pigment epithelium (RPE) density in the peripapillary area (Option A) : The peripapillary area often shows signs of degeneration, not hyperpigmentation. Thickening of peripapillary RNFL on optical coherence tomography (OCT) (Option C) : In glaucoma, the RNFL undergoes thinning , especially in the superior and inferior regions, not thickening, due to the loss of retinal ganglion cells. Uniform retinal nerve fiber layer thickness without focal loss (Option D) : Glaucoma is characterized by focal thinning of the RNFL, particularly around the optic nerve head. Uniform thickness is more typical in healthy eyes, whereas focal thinning is a sign of glaucomatous damage.



References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 354, 355.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 58-year-old woman with a long history of untreated open-angle glaucoma presents with gradual peripheral vision loss. Fundoscopic examination reveals

advanced optic disc cupping. Which of the following vascular changes is most characteristic of progressive glaucomatous optic neuropathy? (Q853865)

A. Inferior displacement of retinal vessels

B. Nasal displacement of retinal vessels

C. Arterial attenuation with increased vein diameter


D. Radial peripapillary capillary obliteration

Correct Answer: B. Nasal displacement of retinal vessels

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In advanced glaucoma, there is progressive optic disc cupping. A characteristic feature of this process is the nasal displacement of retinal vessels, as the optic nerve head undergoes excavation. This displacement is caused by the pulling of the retinal vessels toward the nasal side as the cupping advances, changing the shape of lamina cribrosa. Inferior displacement of retinal vessels (Option A) : This is not characteristic of glaucomatous optic neuropathy; vessels typically shift nasally. Arterial attenuation with increased vein diameter (Option C) : While arterial narrowing can be seen in certain conditions, this pattern is more typical of hypertensive retinopathy. Radial peripapillary capillary obliteration (Option D) : This finding is more associated with ischemic optic neuropathy, not typical in glaucoma.



References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 352, 353

https://www.nature.com/articles/s41598-021-83720-0/figures/5

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: A 58-year-old male with a family history of glaucoma presents for a comprehensive eye exam. His visual acuity is 20/20 in both eyes, and his intraocular pressure (IOP) is measured at 23 mmHg in the right eye and 21 mmHg in the left eye. Fundoscopic examination reveals the following optic disc findings in the right eye: Cup-to-disc ratio: 0.7 vertically Neuroretinal rim thinning: More pronounced in the superior and inferior quadrants Peripapillary atrophy present No optic disc hemorrhages noted All of the following statements are correct regarding the optic disc assessment in this patient except? (Q853864)

A. The optic disc appearance follows the ISNT rule, indicating a healthy optic nerve.

B. The cup-to-disc ratio indicates early glaucomatous damage.

C. The absence of optic disc hemorrhages does not rule out glaucoma.


D. Peripapillary atrophy is commonly associated with glaucoma and warrants further monitoring.


Correct Answer: A. The optic disc appearance follows the ISNT rule, indicating a healthy optic nerve.


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: All of the findings are consistent with early glaucomatous changes except that the disc shows superior and inferior neuroretinal rim thinning violating the ISNT rule. The ISNT rule (Option A) states that, in a healthy optic nerve, the neuroretinal rim thickness typically follows a pattern, being the broadest in: Inferior > Superior > Nasal > Temporal . The patient's optic disc shows several characteristic features of glaucomatous optic neuropathy : 1. Increased cup-to-disc ratio (0.7): A ratio greater than 0.6 is suspicious for glaucoma. 2. Superior and inferior neuroretinal rim thinning: This violates the ISNT rule. 3. Peripapillary atrophy present {{caption_text}} {{caption_text}} The cup-to-disc ratio indicates early glaucomatous damage (Option B) is correct. A cup-to-disc ratio of 0.7 is considered enlarged and suspicious for glaucoma. Normal cup-to-disc ratio is ≤0.3. The absence of optic disc hemorrhages does not rule out glaucoma (Option C) is correct. Their presence is a risk factor for the development and progression of glaucoma. They are more common in NTG, but can also occur in healthy individuals as well as patients with systemic vascular disease. Peripapillary atrophy is commonly associated with glaucoma and warrants further monitoring (Option D) is correct. Peripapillary atrophy surrounding the optic nerve head may be of significance in glaucoma and may be a sign of early damage in patients with ocular hypertension.


C
U
P
Neuroretinal rim (area occupied by the nerve fibres)
Non-neuronal area
@Neet_pg_bot


a) Healthy optic nerve
b) Glaucomatous cupping
Optic Cup
Optic Disk
Cup Radius
Disk Radius
@Neet_pg_bot



References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Pages 351-354

Parson’s Diseases of the Eye, 23rd Edition, Pages 568, 569

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Which device is a non contact tonometer that is based on the principle of corneal hysteresis? (Q853863)

A. Schiotz tonometer

B. Goldmann applanation tonometer

C. Ocular Response Analyzer

D. Pneumatonometer

Correct Answer: C. Ocular Response Analyzer

■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: The Ocular Response Analyzer (ORA) is a non-contact tonometer designed to measure intraocular pressure (IOP) and corneal biomechanics. It uses air impulses to deform the cornea, recording the inward and outward movement to assess corneal elasticity or hysteresis. {{caption_text}} Pascal Dynamic Contour Tonometry (DCT) is a contact tonometer designed to measure intraocular pressure (IOP) by directly conforming to the corneal surface. DCT readings are generally considered more independent of corneal properties compared to Goldmann applanation tonometry (GAT). Schiotz tonometer (Option A) is an indentation tonometer. It doesn't measure corneal hysteresis. Goldmann applanation tonometer (Option B) is the gold standard for IOP measurement and is based on Imbert-Fick law. Pneumatonometer (Option D) is a non-contact applanation tonometer, not based on corneal hysteresis.



Ocular response analyzer

@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot

References:

https://www.ncbi.nlm.nih.gov/books/NBK582132/

https://karger.com/ore/article/55/2/91/258592/Review-on-Dynamic-Contour-Tonometry-and-Ocular

Myron Yanoff and Jay S. Duker, Ophthalmology, 5th Edition, Page 1026

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: Which of the following statements about aqueous humor are incorrect? Aqueous humor is formed from the non-pigmented epithelium of ciliary processes Aqueous constitutes 99% water The non-pigmented epithelium has a Na/Cl symporter responsible for aqueous humor secretion. The rate of formation of Aqueous humor is 2.3 microlitres/min (Q853853)

A. 1, 2, 4 are incorrect

B. Only 3 is incorrect

C. 1,3 is incorrect


D. Only 1 is incorrect

Correct Answer: B. Only 3 is incorrect

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The non-pigmented epithelium has a Na/K/ATPase pump responsible for aqueous humor secretion. Aqueous humor : Aqueous humor is formed from the non-pigmented epithelium of ciliary processes (Option A) Part of ciliary body that secretes aqueous humor is pars plicata. Ciliary processes have blood vessels, stroma, and two layers of epithelium. The inner epithelium is pigmented and the outer epithelium is the non-pigmented epithelium The non-pigmented epithelium has a Na/K/ATPase pump and is responsible for aqueous humor secretion. Aqueous constitutes 99.9% water and 0.1% solutes. (Option B) The rate of formation of aqueous humor is 2.3 microlitres/min (Option D) {{caption_text}}



References:

Kanski's Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 346

Parson's Diseases of the Eye, 23rd Edition, Pages 32, 33, 54, 55

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: Which type of tonometer measures intraocular pressure (IOP) by assessing how much a plunger indents the eye? (Q853861)

A. Schiotz Tonometer

B. Goldmann Tonometer

C. Perkins Tonometer


D. Draeger Tonometer

Correct Answer: A. Schiotz Tonometer

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Schiotz tonometer operates on the principle of indentation tonometry , measuring intraocular pressure (IOP) by determining the depth of corneal indentation produced by a plunger of a known weight. Indentation Tonometry: Concept: Measures IOP by how much a plunger indents the eye; softer eyes indent more than harder eyes. Schiotz Tonometer (Option A): Most widely used tonometer in the world. Principle: Depth of indentation Relies on Friedenwald nomogram Amount of indentation depends on scleral rigidity of patient Advantages: Inexpensive, simple, and reliable. Disadvantages: Gives false readings in eyes with abnormal scleral rigidity (e.g., high myopes, post-surgery). {{caption_text}} Goldmann Tonometer (Option B) → Gold standard Perkins tonometer (Option C) → Hand held version of goldmann Draeger tonometer (Option D) → Measures wide range of IOP



References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 29, 30

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 58-year-old man presents for a routine eye examination. His intraocular pressure (IOP) is consistently measured at 23 mm Hg on multiple occasions. He reports no visual symptoms, and his fundus examination shows no abnormalities. Visual field testing reveals no defects. Based on these findings, which of the following

diagnoses is most appropriate? (Q853860)

A. Normal tension glaucoma

B. Ocular hypertension

C. Acute angle-closure glaucoma

D. Secondary glaucoma

Correct Answer: B. Ocular hypertension

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient has elevated IOP consistently above 21 mm Hg without any visual field defects or fundus changes. These characteristics are consistent with ocular hypertension . ASPECT Ocular Hypertension Characteristic Features IOP consistently above 21 mm Hg (No Damage) No Fundus findings No visual field defects Risk Factors Consistent IOP >30 mm Hg. Central corneal thickness <550 µm. Vertical cup-to-disc ratio >0.7. Increased age. Increased pattern standard deviation (PSD) on Humphrey visual field test. Disc hemorrhages (splinter hemorrhages near or over the disc). Management When these risk factors are present, treatment may be initiated similarly to open-angle glaucoma, but with a less aggressive approach.

References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Pages 347, 367, 370, 378

Parson’s Diseases of the Eye, 23rd Edition, Pages 1392, 1393

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A 65-year-old woman with a history of glaucoma reports worsening vision, headaches, and difficulty with reading. Her IOP shows significant fluctuations throughout the day, and fundus examination reveals optic disc cupping with visual field defects. Which of the following about IOP changes in POAG is false? (Q853859)

A. Diurnal variation includes a morning rise in IOP in 20% of cases and a biphasic rise in 55% of cases.

B. In early POAG, IOP is consistently above 21 mm Hg.

C. IOP is higher in winter due to seasonal variations.

D. Strenuous exercise temporarily lowers IOP.

Correct Answer: B. In early POAG, IOP is consistently above 21 mm Hg.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The diagnosis of Primary Open-Angle Glaucoma (POAG) in this case is supported by significant diurnal fluctuations in intraocular pressure (IOP), characteristic optic disc cupping, and specific visual field defects. In the initial stages of POAG, IOP may not be consistently elevated and often shows exaggerated diurnal variation. Intraocular Pressure (IOP) Changes in POAG Initial Stage: IOP may not be consistently high; diurnal variation (fluctuation throughout the day) is exaggerated. Repeated measurements (every 3-4 hours) are needed to detect this. (Option B) Diurnal Variation Patterns: Morning rise in IOP: 20% of cases Afternoon rise in IOP: 25% of cases Biphasic rise in IOP (morning and afternoon): 55% of cases (Option A) Suspicious Variation: A variation in IOP of over 5 mm Hg is suspicious, and over 8 mm Hg is diagnostic of glaucoma. Later Stages: IOP becomes consistently elevated above 21 mm Hg, typically ranging from 30 to 45 mm Hg. Seasonal Variations: Higher IOP is observed in winter months. (Option C) Exercise: Strenuous exercise lowers IOP temporarily. (Option D)

References:

Kanski's Clinical Ophthalmology: A Systematic Approach, 9th Edition, Pages 349, 350, 351

Parson's Diseases of the Eye, 23rd Edition, Page 693

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: Which gene mutation is linked to the development of primary open angle glaucoma? (Q853858)

A. MYOC gene

B. LOXL1 gene

C. PAX6 gene

D. RB1 gene

Correct Answer: A. MYOC gene

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The MYOC gene (myocilin) is most commonly associated with primary open angle glaucoma (POAG). POAG has been associated with at least 20 loci in the human genome, the most important ones being: Myocillin - MYOC gene Optineurin gene - OPTN gene WTR - 36 LOXL1 gene (Option B): LOXL1 mutations are associated with pseudoexfoliation syndrome, which is the commonest cause of secondary open angle glaucoma. PAX6 gene (Option C): PAX6 gene is crucial for eye development and is associated with congenital eye disorders, including aniridia. RB1 gene (Option D): RB1 gene is primarily associated with retinoblastoma. While it plays a

role in tumor suppression, it is not related to the pathogenesis of primary open-angle glaucoma.

References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 350, 379, 402, 860

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: Which of the following is not a recognized risk factor for Primary Open-Angle Glaucoma? (Q853857)

A. Elevated intraocular pressure

B. Diastolic perfusion pressure of more than 55 mm Hg

C. Family history of glaucoma

D. High myopia

Correct Answer: B. Diastolic perfusion pressure of more than 55 mm Hg

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: A diastolic perfusion pressure of more than 55 mm Hg is generally not considered to be a risk factor for POAG. Lower diastolic perfusion pressure (<55 mm Hg) is a significant risk factor for glaucoma. Primary Open-Angle Glaucoma (POAG): Characteristics: Intraocular Pressure: Elevated (>21 mm Hg) on multiple occasions. Anterior Chamber Angle: Open and normal in appearance. Optic Disc: Characteristic cupping. Visual Field: Specific defects present. Risk Factors of Primary Open-Angle Glaucoma (POAG): Elevated IOP: Asymmetry of IOP of ≥4 mmHg is considered significant. (Option A) Family History: Higher risk in siblings (10%) and offspring (4%). Involves polygenic inheritance with genes like MYOC, OPTN, and WDR36 . (Option C) Age: Increased risk with age, especially between the 5th and 7th decades of life. Race: Higher prevalence and severity in Black individuals compared to Whites. Myopia: Increased risk in myopic individuals. (Option D) Thinner Central Corneal Thickness (CCT) (<555 µm) is an independent risk factor. Cigarette Smoking: Increased risk Blood Pressure: Higher prevalence in hypertensives; diastolic perfusion pressure <55 mm Hg is a significant risk factor. (Option B) Thyrotoxicosis: Increased in Graves' ophthalmic disease. Corticosteroid Responsiveness: Greater risk of IOP rise with topical steroid therapy.

References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 350

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: All the following statements about trabecular meshwork are true except; (Q853856)

A. It regulates the production of aqueous humor

B. Trabecular meshwork secrete Glycosaminoglycans

C. It facilitates the drainage of aqueous humor from the anterior chamber of the eye


D. Trabecular meshwork has 3 parts and among the three, juxtacanalicular has the maximum resistance

Correct Answer: A. It regulates the production of aqueous humor

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Trabecular meshwork has no role in the production of aqueous humor. Aqueous humor is produced from the non-pigmented epithelium of the ciliary process. Trabecular meshwork facilitates the drainage of aqueous humor from the anterior chamber of eye. (Option C) It has 3 parts: (Option D) Uveal trabecular meshwork Corneoscleral trabecular meshwork Juxtacanalicular trabecular meshwork {{caption_text}} Trabecular meshwork secrete Glycosaminoglycans (Option B) Maximum resistance in trabecular meshwork is seen in Juxtacanalicular trabecular meshwork (Option D) Contractile properties are present due to the presence of actin and myosin



References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 346

Parson’s Diseases of the Eye, 23rd Edition, Pages 57, 680, 681

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: Which of the following best describes the role of uveoscleral outflow in intraocular pressure regulation? (Q853855)

A. It involves the drainage of aqueous humor through the trabecular meshwork and Schlemm's canal.

B. It accounts for the drainage of aqueous humor through the ciliary body and sclera, providing an alternative pathway to the conventional outflow route.

C. It is the process by which aqueous humor exits the eye through the vitreous body.


D. It describes the absorption of aqueous humor by the corneal endothelium.

Correct Answer: B. It accounts for the drainage of aqueous humor through the ciliary body and sclera, providing an alternative pathway to the conventional outflow route.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Uveoscleral outflow is an alternative route for aqueous humor drainage, distinct from the conventional trabecular meshwork and Schlemm's canal system. In this process, aqueous humor drains through the ciliary body and into the sclera. This route helps regulate intraocular pressure by providing an additional means for aqueous humor to exit the eye. (Options A, C, and D) describe different aspects of ocular physiology but do not accurately reflect the uveoscleral outflow pathway.



References:

Kanski's Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 346

Parson's Diseases of the Eye, 23rd Edition, Pages 56, 57, 680, 681

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: All the following constituents are more in aqueous humor compared to that of plasma except (Q853854)

A. Lactate

B. Ascorbate

C. Pyruvate

D. Glucose

Correct Answer: D. Glucose

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Due to the difficulty of substances entering the eye through the blood-aqueous barrier and the ease of their exit via drainage channels, many constituents of the aqueous humor are found in lower concentrations compared to blood, except for those that are actively secreted. Glucose levels in the aqueous humor are lower than those in plasma. Blood Aqueous Barrier is formed by: Non-pigmented epithelium (Tight junction) Vascular endothelium of the iris Constituents that are more in aqueous than in plasma are: Lactate (due to anaerobic glycolysis) (Option A) Ascorbate (antioxidant) (Option B) Chloride Pyruvate (Option C)

References:

Parsons Disease of the Eye, 23rd Edition, Page 16f

Myron Yanoff and Jay S. Duker, Ophthalmology, 5th Edition, Page 359

↑ *Back to this question in Part A*

@Neet_pg_bot

• [04_Glaucoma] 009 Previous Year Questions

« Back to Folder Directory

[04_Glaucoma] 009 Previous Year Questions — Quick Jump

Q1 (p. 686)	*Q2 (p. 689)*	*Q3 (p. 689)*	*Q4 (p. 690)*	*Q5 (p. 692)*	*Q6 (p. 692)*
Q7 (p. 693)	*Q8 (p. 695)*	*Q9 (p. 696)*	*Q10 (p. 697)*	*Q11 (p. 698)*	*Q12 (p. 699)*
Q13 (p. 703)	*Q14 (p. 704)*	*Q15 (p. 704)*	*Q16 (p. 707)*	*Q17 (p. 708)*	*Q18 (p. 709)*
Q19 (p. 711)	*Q20 (p. 712)*				

Q1: Which of the following instruments are considered the gold standard for measuring intraocular pressure? (Q857736)

A. Gonioscope


B. Goldmann applanation tonometer

C. Hertel’s exophthalmometer

D. Tonopen

Correct Answer: B. Goldmann applanation tonometer


Explanation: The gold standard instrument for measuring intraocular pressure is Goldmann applanation tonometer . Applanation Tonometry: Concept: Measures IOP based on the force needed to flatten the cornea's surface (Imbert-Fick Law: P = W/A). Goldmann Applanation Tonometer: (Option B) Most popular and accurate; uses a double prism on a slit lamp. Technique: Cornea is anesthetized and stained with fluorescein. Cornea and biprisms are illuminated with cobalt blue light. The prism touches the cornea, and the inner edges of two semicircles are adjusted to just touch. IOP is read by multiplying the dial reading by ten. {{caption_text}} [1] Variable force tonometers: Goldmann applanation tonometer → Gold standard Perkins tonometer (hand held version of goldmann) Draeger tonometer (measures wide range of IOP) Draeger tonometer is similar to Perkins It has a different set of prisms It operates with a motor Pulsair (non-contact tonometer) Tonopen (based on both indentation and applanation) (Option D) Rebound tonometer {{caption_text}} Fixed force tonometers: Maklakov tonometer Barraquer tonometer Gonioscope (Option A): Gonioscope is used to visualize the angle of the anterior chamber. {{caption_text}} Hertel’s exophthalmometer (Option C): Hertel’s exophthalmometer is used to measure the protrusion of the apex of the cornea from the outer orbital margin. {{caption_text}}





Tonopen

@Neet_pg_bot





References:

Kanski’s Clinical Ophthalmology 9th Edition, pages 29,30

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 60-year-old patient presents with sudden onset of severe ocular pain, headache, nausea, vomiting, and blurred vision. On examination, the cornea appears steamy, the pupil is mid -dilated and fixed, and the IOP is 60 mmHg. What is the most likely diagnosis? (Q879003)

A. Acute angle closure glaucoma

B. Secondary angle closure glaucoma due to uveitis

C. Phacomorphic glaucoma

D. Chronic simple glaucoma

Correct Answer: A. Acute angle closure glaucoma

Explanation: The clinical scenario suggests Acute Angle-Closure Glaucoma (AACG) . Pathophysiology: Sudden closure of the anterior chamber angle → blockage of aqueous outflow through the trabecular meshwork → rapid intraocular pressure rise . Predisposing factors: Hypermetropic elderly women ; pupil dilation in dim light narrows the angle and triggers attack. Incorrect Options: Secondary angle closure glaucoma due to uveitis (Option B): Due to posterior synechiae or iris bombe, but develops gradually. The eye is usually quiet or inflamed but not with sudden severe pain and vomiting. Phacomorphic glaucoma (Option C): Lens-induced secondary angle closure due to intumescent cataract. Presents in an elderly patient but with visible swollen lens and shallow anterior chamber; usually subacute onset. Chronic simple glaucoma (Option D): Painless, asymptomatic until late. IOP mildly to moderately elevated, no corneal edema, pupil reactive, no acute symptoms.

References:

Parson’s Diseases of the Eye, 23rd Edition, Page 703, 704, 705

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Krukenberg spindle is seen in which of the following conditions? (Q878908)

A. Phacolytic glaucoma

B. Pigmentary glaucoma

C. Juvenile open -angle glaucoma

D. Primary open – angle glaucoma

Correct Answer: B. Pigmentary glaucoma

Explanation: Krukenberg spindle is characteristically seen in pigmentary glaucoma . It is a vertical, spindle-shaped pigment deposition on the corneal endothelium , usually in the central region . It occurs due to pigment dispersion from the iris pigment epithelium , which deposits on anterior segment structures . Incorrect Options: Phacolytic glaucoma (Option A): Caused by leakage of lens proteins from a hypermature cataract. Juvenile open angle glaucoma (Option C): A primary developmental abnormality of the trabecular meshwork → elevated IOP. Primary open angle glaucoma (Option D): Chronic age-related degeneration of the trabecular meshwork. May have mild pigmentation but krukenberg spindle is not a characteristic feature.

References:

Parson’s Diseases of the Eye, 23rd Edition, Page 716

↑ *Back to this question in Part A*

@Neet_pg_bot


Q4: What is the probable diagnosis for a 50-year-old woman who is experiencing sudden eye pain and redness? An image of her eye is provided for reference. (Q859359)

A. Acute retrobulbar neuritis

B. Acute angle-closure glaucoma

C. Acute anterior uveitis

D. Severe keratoconjunctivitis



Correct Answer: B. Acute angle-closure glaucoma

Explanation: The clinical scenario of a 50-year-old female presenting with an acute painful red-eye and an image showing redness and congestion which gives a diagnosis of Acute angle-closure glaucoma. Acute Angle Closure Glaucoma: Acute angle-closure glaucoma is a sudden and severe form of glaucoma characterized by a rapid increase in intraocular pressure. It occurs when the drainage angle of the eye becomes blocked, leading to a buildup of fluid and pressure within the eye. Clinical features of Glaucoma are: Sudden Painful eye. A ciliary flush around the cornea due to dilated ciliary veins, and in long-standing cases, caput medusae with prominent vessels. Cornea - initially clear but insensitive, later becoming hazy, with epithelial bullae (bullous keratopathy) or filaments (filamentary keratitis). Headache and nausea: Some individuals may experience accompanying headache, nausea, and even vomiting. Anterior chamber - very shallow. Iris - atrophic. Pupil - fixed and dilated. Optic disc - glaucomatous optic atrophy. IOP is high and the eyeball becomes stony hard. Management of absolute glaucoma In absolute glaucoma, the goal of treatment is to relieve pain and reduce IOP, as vision cannot be restored. Retrobulbar alcohol injection - given to relieve pain. First, 1 mL of 2% xylocaine is injected followed after about 5-10 minutes by 1 mL of 80% alcohol. It suppresses the ciliary ganglion. A cyclodestructive procedure, such as cyclocryotherapy or cyclophotocoagulation, is the most appropriate management option in this scenario as it reduces aqueous production by damaging the secretory ciliary processes aiming to reduce IOP. If pain persists despite cyclodestruction, enucleation/ evisceration may be considered. Incorrect Options Acute retrobulbar neuritis (Option A) , also known as optic neuritis, typically presents with vision loss, eye pain, and decreased color vision. It is not characterized by redness and congestion of the eye, so it is less likely to be the correct diagnosis in this case. Acute anterior uveitis (Option C) , also known as iritis, is inflammation of the iris and the anterior chamber of the eye. It commonly presents with symptoms of a red, painful eye with photophobia (sensitivity to light) and decreased vision. The image showing redness and congestion supports this diagnosis. Acute anterior uveitis can be associated with underlying systemic conditions such as ankylosing spondylitis or reactive arthritis. It is the most likely diagnosis based on the given scenario. Severe keratoconjunctivitis (Option D) refers to inflammation of both

the cornea and the conjunctiva. While it can cause redness and irritation of the eye, it is not commonly associated with acute onset and severe pain. It is less likely to be the correct diagnosis in this case.

References:

https://pmc.ncbi.nlm.nih.gov/articles/PMC3669490/#:~:text=An%20eye%20that%20has%20lost,other%20cases%2C%20it%20may%20not.

Kanski Clinical Ophthalmology, 10th edition, Pages 379, 380, 381.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: The drug increasing uveoscleral flow is? (Q855019)

A. Timolol

B. Latanoprost

C. Pilocarpine

D. Carbonic anhydrase inhibitor

Correct Answer: B. Latanoprost

Explanation: Latanoprost is a prostaglandin F■α analog that lowers intraocular pressure (IOP) primarily by increasing uveoscleral outflow . Mechanism of Action: Prostaglandin analogs (Latanoprost, Travoprost, Bimatoprost, Tafluprost) act on the FP receptors in the ciliary muscle , causing remodeling of the extracellular matrix , thereby reducing resistance and enhancing aqueous humor drainage through the uveoscleral pathway . Other Drugs and Their Mechanisms: Drug Class Example Mechanism of IOP Reduction β-blockers Timolol ↓ Aqueous humor production Prostaglandin analogs Latanoprost ↑ Uveoscleral outflow Miotics (Cholinergic agonists) Pilocarpine ↑ Trabecular meshwork outflow Carbonic anhydrase inhibitors Acetazolamide, Dorzolamide ↓ Aqueous humor production α■-agonists Brimonidine ↓ Aqueous production and ↑ Uveoscleral outflow (minor effect)

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 369, 370, 374, 380

Parsons’ Diseases of the Eye, 23rd Edition, PDF Pages 724, 725

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: True regarding POAG: Abnormality of trabecular meshwork is seen on gonioscopy First degree relatives have more chances of developing steroid induced glaucoma Dilatation of pupil is associated with exacerbation of IOP First degree relatives are at 1% increased risk of POAG Visual field defect can exist with normal C:D ratio (Q856418)

A. 1,2,4 and 5

B. 1,2 and 3

C. 2 and 5 only

D. 2,3 and 4

Correct Answer: C. 2 and 5 only

Explanation: Primary Open-Angle Glaucoma (POAG) Characteristics: Intraocular Pressure: Elevated (>21 mm Hg) on multiple occasions. Anterior Chamber Angle: Open and normal in appearance (Statement 3) . Optic Disc: Characteristic cupping , normal C:D ratio in early stages (Statement 5) . Visual Field: Specific defects present. Risk Factors: Elevated IOP: Asymmetry of IOP of ≥4 mmHg is considered significant. Family History: Higher risk in siblings (10%) (Statement 4) and offspring (4%). Involves polygenic inheritance with genes like MYOC, OPTN, and WDR36 . Age: Increased risk with age, especially between the 5th and 7th decades of life. Race: Higher prevalence and severity in Black individuals compared to Whites. Myopia: Increased risk in myopic individuals. Thinner Central Corneal Thickness (CCT) (<555 µm) is an independent risk factor. Cigarette Smoking: Increased risk Blood Pressure: Higher prevalence in hypertensives; diastolic perfusion pressure <55 mm Hg is a significant risk factor. Thyrotoxicosis: Increased in Graves' ophthalmic disease. Corticosteroid Responsiveness: Greater risk of IOP rise with topical steroid therapy (Statement 2). Abnormality of trabecular meshwork is seen on gonioscopy (Statement 1) although trabecular meshwork is seen in gonioscopy, abnormalities in the trabecular meshwork cannot be observed during gonioscopy.

References:

Kanski’s Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 350

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: In a patient with primary open-angle glaucoma, which of the following pupillary abnormalities is most commonly seen? (Q856934)

A. Marcus Gunn pupil


B. Adie’s tonic pupil

C. Argyll-Robertson pupil

D. Hutchinson’s pupil

Correct Answer: A. Marcus Gunn pupil

Explanation: : The pupils paradoxically dilate when the flashlight is swung from the unaffected to the affected eye due to the decreased light perception. This is also called the Marcus Gunn Pupil (Option A) (first sign of optic nerve disorder). Causes: Optic neuritis Optic neuropathy Chiasmal compression Intraocular hemorrhage Central serous retinopathy Cystoid macular edema Retinal detachment Optic tract lesion Central or branch arterial occlusion Central or branch vein occlusion Anisometropic amblyopia Cataract Corneal scar Visual pathway damage Tested by Swinging Flashlight Test . A bright flashlight is shone onto one pupil, causing constriction, then quickly moved to the opposite pupil to observe its response. This process is repeated multiple times to assess the pupillary reactions. Normally, both pupils constrict equally, but the pupil to which light is transferred remains tightly constricted. In RAPD, the affected pupil dilates when the flashlight moves from the normal to the abnormal eye. The dilatation that occurs is commonly called the Marcus Gunn pupil or RAPD (Relative afferent pupillary defect). Adie’s Pupil (Option B) shows no reaction to light, and the near reflex is very slow and tonic. It shows pupillary constriction to 0.125% Pilocarpine due to denervation supersensitivity. Seen in Postganglionic parasympathetic pupillomotor damage. Argyll Robertson Pupil (ARP) (Option C) shows reaction to near reflex, but light reflex (both direct and consensual) is absent, i.e. there is light near dissociation. Seen in Neurosyphilis Hutchinson’s pupil (Option D) This refers to a fixed, dilated pupil due to a midbrain lesion or uncal herniation. It is not typically seen in glaucoma.



References:

Parsons Diseases of the eye, 23rd Edition, Pg 278, 279.

Yanoff & Duker Ophthalmology, 5th edition, Pg 963, 964.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: What is the likely diagnosis for a 2-month-old infant who was brought in by their mother due to complaints of light sensitivity, difficulty opening the eyes, and excessive tearing? (Q856125)

A. Ophthalmia neonatorum

B. Congenital glaucoma

C. Congenital Dacryocele

D. Congenital Cataract




Correct Answer: B. Congenital glaucoma

Explanation: ongenital glaucoma: The presence of classic triad of symptoms- watering, photophobia, and blepharospasm in a 2-month old infant are all consistent with Congenital glaucoma . Primary Congenital Glaucoma: Types Newborn glaucoma/ True congenital glaucoma: Elevated intraocular pressure is present during intrauterine life, resulting in ocular enlargement at birth. Typically presents within the first month of life. Infantile: ≤ 3 yrs of age Juvenile: > 3 yrs of age Pathogenesis Trabeculodysgenesis Angle Recess Absence Iris Insertion Types: Flat Insertion: Reveals the ciliary body and scleral spur. Concave Insertion: Obscures the scleral spur and ciliary body. Clinical Features Classic Triad: Lacrimation, photophobia, and blepharospasm; often accompanied by eye rubbing. {{caption_text}} Corneal Signs: Oedema: Early sign, may progress to permanent opacity. Enlargement: Corneal diameter >13 mm confirms enlargement; >16 mm indicates poor prognosis. Haab’s Striae: Tears in Descemet's membrane, visible as opaque lines. {{caption_text}} Additional Features: Deep Anterior Chamber: Indicates elevated IOP. Optic Disc Changes: Variable cupping and atrophy. Raised IOP: Elevated but not acute. Axial Myopia: Due to increased axial length. Management Direct Goniotomy followed by Trabeculotomy - preferred management for Congenital Glaucoma (Buphthalmos: meaning "bull-like eyes," is a

condition characterized by the enlargement of the eyeball. It typically manifests before the age of 3 years and results from the accumulation and retention of aqueous humor.)





References:

Kanski Clinical Ophthalmology, 10th edition, Pages 401, 402, 403

Parsons' Diseases of the Eye, 23rd Edition, PDF Pages 277, 278

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Which medication should be avoided in a 30-year-old female presenting with redness, pain, increased intraocular pressure, keratic precipitates, and aqueous flare? (Q856696)

A. Beta-blockers

B. Mannitol

C. Carbonic anhydrase inhibitors

D. Prostaglandin analogues

Correct Answer: D. Prostaglandin analogues

Explanation: The clinical presentation of redness, pain, keratic precipitates, aqueous flare, and increased intraocular pressure is suggestive of anterior uveitis induced glaucoma . Prostaglandin analogues are contraindicated in cases of Uveitis as it may aggravate the inflammation. Anti-Glaucoma drugs: Prostaglandin Analogues: Latanoprost, Bimatoprost, Travoprost, Tafluprost Mechanism: Increases uveoscleral outflow Adverse effects: Sandy eyes, orbital fat atrophy, anterior uveitis, cystoid macular edema, heterochromia iridis, hypertrichosis Contraindications: Cystoid macular edema, diabetes mellitus, Anterior Uveitis (Option D) Beta Blockers: (Option A) Non-Selective: Timolol, Levobunolol, Carteolol, Metipranolol Selective: Betaxolol, Levobetaxolol Mechanism: Reduces aqueous secretion Adverse effects: Allergic blepharoconjunctivitis, dry eyes, corneal hypoesthesia Alpha-2 Agonists: Brimonidine (long-term use), Apraclonidine (IOP spikes) Mechanism: Reduces aqueous production, increases uveoscleral outflow Adverse effects: Lid retraction, respiratory depression (Brimonidine), allergic blepharoconjunctivitis Contraindications: Children <2 years Additional: Brimonidine: Safest during pregnancy (Category B) Miotics: Pilocarpine, Physostigmine, Echothiophate Mechanism: Stimulates M3 receptors, increasing trabecular outflow Adverse effects: Brow pain, miosis, cataract (Echothiophate), retinal tears Carbonic Anhydrase Inhibitors: (Option C) Acetazolamide (systemic), Brinzolamide, Dorzolamide (topical) Mechanism: Inhibits aqueous secretion Adverse effects: Ciliary edema, secondary angle-closure glaucoma Osmotic agents: (Option B) Mannitol (IV), Glycerol (oral) Mechanism: Lowers IOP by creating an osmotic gradient Adverse effects: Cardiac overload, urinary retention Newer Drugs: Netarsudil (Rho kinase inhibitor): Increases trabecular outflow; Side effect - vortex keratopathy Omidenepag (PGE2 agonist): Increases both trabecular and uveoscleral outflow

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 404 - 407

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Which of the following medications should not be administered to a patient with a known case of angle closure glaucoma who has also been diagnosed with major depressive disorder? (Q855384)

A. Amitriptyline

B. Sertraline

C. Mirtazapine

D. Fluvoxamine

Correct Answer: A. Amitriptyline

Explanation: Amitriptyline, a Tricyclic Antidepressant (TCA) should not be administered to a known case of Angle closure glaucoma due to its antagonistic action on M1 (Muscarinic) receptor. Amitriptyline: MOA: Primarily blocks Serotonin (5HT3) and Norepinephrine (NE) reuptake ; also has additional blocking action on ■1 (adrenergic), H1 (Histamine) and M1 (Muscarinic) receptors. ADRs : Orthostatic hypotension, GI disturbances, seizures, increases risk of suicidal thoughts, sexual dysfunction. Contraindication: Angle closure glaucoma Common drugs contraindicated in Angle closure glaucoma patients: Anti-Muscarinic Ipratropium bromide, Thiotropium bromide (M1) Toletrodine, Oxybutinin (M3) Scopolamine (M1) Anti-Histaminics Diphenhydramine, Fexofenadine (H1) Cimetidine, Ranitidine (H2) Migraine drugs Sumatriptan Anti-depressants SNRI: Duloxetine and Milnacipram TCA: Amitryptiline Anti-Epileptics Topiramate, Zonisamide Antibiotics Trimethoprim-Sulfametoxazole (Cotrimoxazole)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A patient presents with a vision problem in the right eye. On examination, the intraocular pressure (IOP) of the right eye is 50 mmHg, while the IOP of the left eye is 16 mmHg. Slit-lamp examination of the right eye reveals flare cells in the anterior chamber and a cataractous lens. What is the diagnosis? (Q853114)

A. Phacolytic glaucoma

B. Malignant glaucoma

C. Fuch's heterochromic iridocyclitis

D. Phacomorphic glaucoma

Correct Answer: A. Phacolytic glaucoma

Explanation: The clinical scenario is suggestive of Glaucoma and the slit lamp image flare cells in the anterior chamber and a cataractous lens corresponds to secondary open-angle glaucoma called Phacolytic glaucoma. Feature Phacomorphic Glaucoma (Option D) Phacolytic Glaucoma (Option A) Pathogenesis Intumescent cataract blocks the pupil. this pushes the iris forward, blocking the anterior chamber angle, leading to angle-closure glaucoma. Leakage of lens proteins from a hypermature cataract clogs the trabecular meshwork. Causes

Intumescent cataract (mature cataract) Anterior subluxation of lens Spherophakia Hypermature (Morgagnian) cataract Obstruction of the trabecular meshwork by lens proteins. Clinical Features Symptoms are similar to acute primary angle-closure glaucoma. Shallow anterior chamber. Ciliary congestion, dilated pupil, and swollen cataract is visible. Closed anterior chamber angle on gonioscopy. Features of acute congestive glaucoma due to high IOP. Deep anterior chamber. Open anterior chamber angle on gonioscopy. Management Medical Treatment : IV mannitol, systemic acetazolamide, topical beta-blockers. Laser Iridotomy: To relieve angle closure. Surgical Treatment: Cataract extraction with PCIOL implantation. Medical Therapy: To lower IOP. Surgical Treatment: Extraction of the hypermature cataract and PCIOL implantation. Malignant glaucoma (Option B) shallowing of the anterior chamber is noticed and Seidel's test is negative. Fuch heterochromic iridocyclitis (Option C) , low-grade flare and cells, with stellate keratic precipitates and fine filaments may be seen but IOP is not very high as in the above case.



References:

Kanski's Clinical Ophthalmology, 10th edition, Pg 393 Comprehensive Ophthalmology by AK Khurana 7th edition - page 199, 255

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: What is the probable diagnosis for a neonate who presents with symptoms of photophobia and excessive tearing, along with a clinical examination showing a

normal tear passage and a clear, but abnormally large cornea, and the presence of Haab's striae in the eye image below? (Q857662)

A. Buphthalmos

B. Keratoconus

C. Trachoma

D. Keratoglobus




Correct Answer: A. Buphthalmos

Explanation: The presence of classic symptoms- watering, photophobia with abnormally large cornea in a neonate are all consistent with Buphthalmos- Congenital glaucoma . {{caption_text}} Primary Congenital Glaucoma: Types Newborn glaucoma/ True congenital glaucoma: Elevated intraocular pressure is present during intrauterine life, resulting in ocular enlargement at birth. Typically presents within the first month of life. Infantile: ≤ 3 yrs of age Juvenile: > 3 yrs of age (Buphthalmos: meaning "bull-like eyes," is a condition characterized by the enlargement of the eyeball. It typically manifests before the age of 3 years and results from the accumulation and retention of aqueous humor.) Pathogenesis Trabeculodysgenesis Angle Recess Absence Iris Insertion Types Flat Insertion: Reveals the ciliary body and scleral spur. Concave Insertion: Obscures the scleral spur and ciliary body. {{caption_text}} Clinical Features Classic Triad: Lacrimation, photophobia, and blepharospasm; often accompanied by eye rubbing. Corneal Signs: Oedema: Early sign, may progress to permanent opacity. Enlargement: Corneal diameter >13 mm confirms enlargement; >16 mm indicates poor prognosis. Haab’s Striae: Tears in Descemet's membrane, visible as opaque lines. {{caption_text}} Additional Features: Deep Anterior

Chamber: Indicates elevated IOP. Optic Disc Changes: Variable cupping and atrophy. Raised IOP: Elevated but not acute. Axial Myopia: Due to increased axial length. Management Direct Goniotomy followed by Trabeculotomy - preferred management for Congenital Glaucoma Keratoconus (Option B): is frequently due to congenital weakness of cornea, though it manifests only after puberty. In keratoconus, the cornea thins near the center and progressively bulges forwards, with the apex of the cone always being slightly below the center of the cornea. Important signs include: Munson sign (acute bulge given to the lower lid when the patient looks down), acute hydrops (due to ruptures in the Descemet's membrane), Vogt striae and Fleischer ring may be seen in the later stages. Trachoma (Option C): is the chronic inflammatory disease affecting primarily the conjunctiva, but later secondarily affecting the lids and the cornea. The organism responsible is Chlamydia trachomatis. From the clinical point of view, the diagnostic features of trachoma depend on the following characteristics: Follicles in the upper palpebral conjunctiva (Sago-grain appearance) , pannus , limbal follicles or their sequelae as Herbert pits , in later stages, typical stellate trachomatous scarring in the conjunctiva. Keratoglobus (Option D): is a congenital anomaly in which there is hemispherical protrusion of the whole cornea , occurring bilaterally. It is familial and hereditary. It differs from buphthalmos in that the intraocular pressure is normal, cornea is clear, refractive errors are with-the-rule astigmatism , angle of the anterior chamber normal and there is no cupping of the optical disc.
{{caption_text}}








Haab's Striae

(Double coloured opacities in congenital glaucoma)



References:

Kanski's Clinical Ophthalmology 9th Edition, Pages 217, 218.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: After an acute episode of angle-closure glaucoma, which triad of symptoms is typically observed? (Q855241)

A. Vogt's triad


B. Virchow triad

C. Hutchinson's triad

D. Gaucher triad

Correct Answer: A. Vogt's triad

Explanation: Vogt's triad is a clinical set of findings typically associated with acute primary angle-closure glaucoma (APAC) after an attack has been resolved. The triad helps in identifying damage caused during the acute episode. Features of APAC: Descemet membrane folds Iris atrophy Irregular pupil Glaucomflecken Cataract Cupping +/- Vogt’s triad: It may be seen in both treated and non-treated cases after an attack of acute PAC. It is characterized by: Slightly dilated non-reacting pupil (due to sphincter atrophy) Glaucomflecken (anterior subcapsular lenticular opacity) Patches of iris atrophy {{caption_text}} Virchow's triad (Option B) : This refers to the three factors that predispose to thrombosis: stasis of blood flow , vascular injury , and hypercoagulability . It is used in the context of venous thrombosis, not glaucoma. Hutchinson's triad (Option C) : This refers to a set of findings associated with congenital syphilis , including notched incisors , interstitial keratitis , and sensorineural hearing loss . Gaucher's triad (Option D) : This refers to symptoms seen in Gaucher's disease , a genetic disorder involving lipid storage. The triad includes hepatosplenomegaly , bone pain , and anemia , unrelated to glaucoma.



References:

Parson’s Textbook of Ophthalmology, 23rd Edition, Page 704

Vogt's triad – Do we find the triad all the time?

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: In which of the following conditions is atropine not recommended for use? (Q855597)

A. Acute congestive glaucoma

B. Acute anterior uveitis

C. Malignant glaucoma

D. Corneal ulcer

Correct Answer: A. Acute congestive glaucoma

Explanation: Atropine is not recommended for use in acute congestive glaucoma . This is because atropine is an anticholinergic that causes pupil dilation, which can exacerbate the already elevated intraocular pressure in acute congestive (or angle-closure) glaucoma by further obstructing the drainage of aqueous humor from the eye. This can worsen the condition. Over 60 drugs have been linked to acute angle-closure (AAC) glaucoma, particularly in individuals with ocular predispositions. These include topical anticholinergic pupil dilators (e.g., atropine, cyclopentolate) and systemic medications such as topiramate, sulfonamides, duloxetine, and phenothiazines. Patients on these medications, especially those with AAC risk factors, should have regular ophthalmic examinations. Acute anterior uveitis (Option B): Atropine can be beneficial in this condition as it helps to relieve ciliary muscle spasms, reduces pain, prevents the formation of posterior synechiae, and keeps the pupil mobile. Malignant glaucoma (Option C): Atropine is the first line of treatment for management of malignant glaucoma. It dilates the ciliary ring and relaxes the ciliary muscles which causes stretching of zonules that will pull back the lens, thus opening the block. Corneal ulcer (Option D): Atropine can be used in this case to help relieve pain associated with ciliary muscle spasms, reduces pain, prevents the formation of posterior synechiae, and improve vascularity.

References:

Acute Angle-Closure Glaucoma - StatPearls - NCBI Bookshelf

(PDF) Ophthalmic Uses of Atropine: A Review

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: What is the probable diagnosis for an infant brought in with symptoms of excessive tear production, sensitivity to light, and blepharospasm? (Q855495)

A. Congenital glaucoma

B. Congenital cataract

C. Congenital Dacryocystitis

D. Ophthalmia neonatorum




Correct Answer: A. Congenital glaucoma

Explanation: True Congenital Glaucoma is characterized by elevated intraocular pressure present during intrauterine life, resulting in ocular enlargement at birth . The presence of Haab's striae, early onset, enlarged corneas, and the classic triad of symptoms (watering, photophobia, and blepharospasm) are all consistent with newborn glaucoma. Primary Congenital Glaucoma Types 1. Newborn glaucoma/ True congenital glaucoma : Elevated intraocular pressure is present during intrauterine life, resulting in ocular enlargement at birth . (40% cases) Typically presents within the first month of life. Often associated with developmental anomalies of the anterior segment, such as primary congenital glaucoma, which can result from genetic mutations or other congenital conditions. 2. Infantile: ≤ 3 yrs of age - Infantile glaucoma is labelled when the disease manifests prior to the child's third birthday Affects infants older than 1 month and children up to 3 years. Can also result from genetic factors or be secondary to other conditions like Sturge-Weber syndrome or other syndromic associations. 3. Juvenile: > 3 yrs of age who develop pressure rise after 3 years but before adulthood. Also known as Juvenile primary open angle glaucoma(POAG) usually occurs between 10 to 35 years of age. Buphthalmos , meaning

"bull-like eyes," is a condition characterized by the enlargement of the eyeball. It typically manifests before the age of 3 years and results from the accumulation and retention of aqueous humor. Pathogenesis Trabeculodysgenesis: Maldevelopment of the trabeculum and iridotrabecular junction from neural crest cells impairs aqueous outflow, causing elevated IOP. Angle Recess Absence: The angle recess is absent, with the iris inserting either flatly or concavely into the trabeculum. Iris Insertion Types: Flat Insertion: Reveals the ciliary body and scleral spur. Concave Insertion: Obscures the scleral spur and ciliary body. Clinical Features Classic Triad: Lacrimation, photophobia, and blepharospasm; often accompanied by eye rubbing. {{caption_text}} Corneal Signs: Oedema: Early sign, may progress to permanent opacity. Enlargement: Corneal diameter >13 mm confirms enlargement; >16 mm indicates poor prognosis. Haab's Striae: Tears in Descemet's membrane, visible as opaque lines. {{caption_text}} Additional Features: Thin Sclera: Appears blue due to underlying uveal tissue. Deep Anterior Chamber: Indicates elevated IOP. Iris Changes: Iridodonesis and atrophic patches in late stages. Lens Changes: Becomes flat and may subluxate. Optic Disc Changes: Variable cupping and atrophy. Raised IOP: Elevated but not acute. Axial Myopia: Due to increased axial length. Incorrect Options: Congenital Cataract (Option B) : Causes a cloudy lens and poor vision , but doesn't typically cause excessive tearing, light sensitivity, or blepharospasm. Congenital Dacryocystitis (Option C) : Leads to tearing and discharge from a blocked tear duct but doesn't cause light sensitivity or blepharospasm. Ophthalmia Neonatorum (Option D) : Involves eye redness and discharge due to infection at birth, but doesn't cause light sensitivity or spasms.





References:

Kanski Clinical Ophthalmology, 10th edition, Pages 401, 402, 403

Parsons' Diseases of the Eye, 23rd Edition, PDF Pages 277, 278

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A patient presents with sudden painful decrease in vision and a fixed, vertically oval pupil. What is the likely diagnosis? (Q860174)

A. Macular degeneration


B. Optic neuritis

C. Acute congestive glaucoma

D. Acute anterior uveitis

Correct Answer: C. Acute congestive glaucoma

Explanation: Acute congestive glaucoma is characterized by sudden painful decrease in vision and fixed vertically oval pupil Acute primary angle-closure glaucoma (APAC) after an attack has been resolved. The triad helps in identifying damage caused during the acute episode. Features of APAC: Descemet membrane folds Iris atrophy Irregular pupil (Option C) Glaucomflecken Cataract Cupping +/- Vogt's triad: It may be seen in both treated and non-treated cases after an attack of acute PAC. It is characterized by: Slightly dilated non-reacting pupil (due to sphincter atrophy) Glaucomflecken (anterior subcapsular lenticular opacity) Patches of iris atrophy {{caption_text}} Incorrect Options: Macular degeneration (Option A): It is characterized by irreversible loss of vision Fixed oval pupils is not a feature of this condition Optic neuritis (Option B) are normal in Optic neuritis Acute anterior uveitis (Option D) is characterized by a dilated, festooned-shaped pupil



References:

Parson's Textbook of Ophthalmology, 23rd Edition, Page 704

https://journals.lww.com/ijog/fulltext/2023/03020/vogt_s_triad___do_we_find_the_triad_all_the_time_.118.aspx

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A patient presents with a vision problem in the right eye. On examination, the intraocular pressure (IOP) of the Right eye is 50 mmHg, while the IOP of the Left eye is 16 mmHg. Slit lamp examination of the right eye reveals flare cells in the anterior chamber and a cataractous lens. What is the diagnosis? (Q840060)

A. Phacolytic glaucoma

B. Malignant glaucoma

C. Fuch heterochromic iridocyclitis

D. Phacomorphic glaucoma

Correct Answer: A. Phacolytic glaucoma

Explanation: Correct Option A - Phacolytic glaucoma: Flare cells in the anterior chamber of the eye, cataractous lens, and unequal IOP point towards the diagnosis of phacolytic glaucoma. It is a type of secondary open-angle glaucoma in which trabecular meshwork is clogged by the lens proteins, macrophages which have phagocytosed the lens proteins, and inflammatory debris. Leakage of the lens proteins occurs through an intact capsule in the hypermature (Morgagnian) cataractous lens. The anterior chamber may become deep, and the aqueous may contain fine white protein particles, which settle down as pseudohypopyon. Anterior chamber angle is open on gonioscopy, and aqueous flare or cells may be seen in the anterior chamber. Incorrect Options: Option B - Malignant glaucoma: In Malignant glaucoma, shallowing of the anterior chamber is noticed and Seidel's test is negative. Option C - Fuch heterochromic iridocyclitis: In Fuch heterochromic iridocyclitis, low-grade flare and cells, with stellate keratic precipitates and fine filaments may be seen but IOP is not very high as in the above case. Option D - Phacomorphic glaucoma: In Phacomorphic glaucoma, ciliary congestion, dilated pupil, and intumescent senile cataractous lens is noted.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: What is the gold standard method for measuring intraocular pressure in patients? (Q860572)

A. Hertel's exophthalmometer

B. Gonioscope

C. Goldmann applanation tonometer


D. Tonopen

Correct Answer: C. Goldmann applanation tonometer

Explanation: Types of Tonometry: 1. Indentation Tonometry: Concept: Measures IOP by how much a plunger indents the eye; softer eyes indent more than harder eyes. Schiotz Tonometer: Most widely used tonometer in the world. Advantages: Inexpensive, simple, and reliable. Disadvantages: Gives false readings in eyes with abnormal scleral rigidity (e.g., high myopes, post-surgery). {{caption_text}} 2. Applanation Tonometry: Concept: Measures IOP based on the force needed to flatten the cornea's surface (Imbert-Fick Law: P = W/A). Goldmann Tonometer (Option B): Most popular and accurate; uses a double prism on a slit lamp. Technique: Cornea is anesthetized and stained with fluorescein. Cornea and biprisms are illuminated with cobalt blue light. The prism touches the cornea, and the inner edges of two semicircles are adjusted to just touch. IOP is read by multiplying the dial reading by ten. {{caption_text}} Variable force tonometers: Goldmann applanation tonometer → Gold standard Perkins tonometer (hand held version of goldmann) Draeger tonometer (measures wide range of IOP) Draeger tonometer

is similar to Perkins It has a different set of prisms It operates with a motor Pulsair (non-contact tonometer) Airpuff (non-contact tonometer) Grolman (non-contact tonometer) Tonopen (based on both indentation and applanation) (Option D) MacKay Marg (based on both indentation and applanation) Rebound tonometer Transpalpebral Tonometer Fixed force tonometer: Maklakov tonometer Barraquer tonometer Hertel's exophthalmometer (Option A) is used to measure the degree of exophthalmos (protrusion of the eyeball) in conditions such as thyroid eye disease. Gonioscope (Option B) is a specialized lens used to visualize the iridocorneal angle, which is important in assessing angle-closure glaucoma.





References:

Kanski's Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 29, 30

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: The first line of treatment for glaucoma in adults is? (Q860527)

A. PG analogues

B. Atropine

C. Acetazolamide

D. Pilocarpine

Correct Answer: A. PG analogues

Explanation: Treatment of glaucoma: There are 3 types of treatment modalities: Medical management Laser Surgery 1. Medical: The drugs either decrease the formation of aqueous or increase the drainage to decrease IOP. This is of 2 types: Topical (6 types) Systemic (2 types) Topical: PGF2 alpha agonist: It increases the uveoscleral outflow First line in glaucoma These are also contraindicated in uveitis and asthma For eg- Latanoprost,

bimatoprost, travoprost , tafluprost, and unoprostone isopropyl Latanoprost causes heterochromia iridis Bimatoprost increases both uveoscleral and trabecular outflow Side effects: Hypertrichosis Beta-blockers These decrease the formation of aqueous Eg: Timolol, betaxolol C/I in asthma S/E: corneal anesthesia and blepharoconjunctivitis Alpha agonists: They have dual action i.e. they decrease the formation of aqueous and increase the trabecular outflow. Eg: Adrenaline/epinephrine, dipivefrin, brimonidine, and apraclonidine Brimonidine is safe in pregnancy Adrenaline is C/I in aphakic glaucoma Miotics: These increase the trabecular outflow and open the angle blockage Eg: Pilocarpine S/E of Pilocarpine: Pseudo myopia, retinal detachment, shallow anterior chamber, iris cysts Topical carbonic anhydrase inhibitors: These decrease the formation of aqueous Eg: dorzolamide and brinzolamide These are C/I in sulpha allergies Newer anti-glaucoma drugs: Netarsudil: this increases the contractile property of the trabecular meshwork, thus increasing the trabecular outflow and it also decreases the aqueous production. It is a Rho-kinase inhibitor S/E: vortex keratopathy Systemic: Carbonic anhydrase inhibitor For eg: acetazolamide , methazolamide S/E: Tingling and numbness, constipation, renal stones, and a bad metallic taste Hyperosmotic agents: These draw aqueous out of the eye For eg: I/V mannitol (C/I in heart disease), glycerol (C/I in diabetes), isosorbide, and urea 2. Laser : Laser trabeculoplasty can be done using argon laser (ALT), or Micropulse diode (MLT) and selective laser trabeculoplasty (SLT). 3. Surgery : Surgical treatment of POAG primarily consists of a fistulizing (filtration) surgery which provides a new channel for aqueous outflow and successfully controls the IOP . Trabeculectomy is the most frequently performed filtration surgery nowadays.

References:

Kanski Clinical Ophthalmology, 10th edition, Pages 369, 370, 374, 380

Parsons' Diseases of the Eye, 23rd Edition, PDF Pages 724, 725

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: A patient complains of eye pain and difficulty seeing after watching a movie. What is the first step in management? (Q844037)

A. Mannitol with moxifloxacin

B. Mannitol with pilocarpine

C. Mannitol with atropine

D. Mannitol with carboxymethylcellulose

Correct Answer: B. Mannitol with pilocarpine

Explanation: Correct Option B - Mannitol with pilocarpine: The scenario describes a patient presenting with symptoms after watching a movie, which is suggestive of acute primary angle-closure glaucoma (APACG). Mannitol helps to reduce intraocular pressure rapidly, and pilocarpine is a miotic agent that constricts the pupil and pulls the iris away from the trabecular meshwork, thereby opening the drainage angle. Incorrect Options: Options

A and D: Moxifloxacin and carboxymethylcellulose have no effect on intraocular pressure. Option C: Atropine is a mydriatic agent that dilates the pupil, which can worsen angle closure by increasing the pupillary block.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [05_Retina] 010 Retinal Investigations And Vascular Disorders

« Back to Folder Directory

[05_Retina] 010 Retinal Investigations And Vascular Disorders — Quick Jump

Q1 (p. 714)	*Q2 (p. 715)*	*Q3 (p. 716)*	*Q4 (p. 717)*	*Q5 (p. 721)*	*Q6 (p. 723)*
Q7 (p. 725)	*Q8 (p. 726)*	*Q9 (p. 727)*	*Q10 (p. 728)*	*Q11 (p. 730)*	*Q12 (p. 731)*
Q13 (p. 732)	*Q14 (p. 733)*	*Q15 (p. 734)*	*Q16 (p. 735)*	*Q17 (p. 736)*	*Q18 (p. 738)*
Q19 (p. 739)	*Q20 (p. 743)*	*Q21 (p. 744)*	*Q22 (p. 746)*	*Q23 (p. 747)*	*Q24 (p. 749)*
Q25 (p. 751)	*Q26 (p. 752)*	*Q27 (p. 754)*	*Q28 (p. 755)*	*Q29 (p. 757)*	*Q30 (p. 758)*
Q31 (p. 759)	*Q32 (p. 759)*	*Q33 (p. 760)*	*Q34 (p. 761)*	*Q35 (p. 763)*	*Q36 (p. 763)*

Q1: A 23-year-old male presents to the ophthalmologist with pain, redness, photophobia or sensitivity to the light, and decreased vision from the last 3 days. Both the eyes of the patient are affected, and the patient is very irritable. Upon examination, infection of both the cornea and ciliary nerve is involved. The doctor diagnosed the patient with acute anterior uveitis. Which of the following HLA (human leukocyte antigen) is associated with uveitis? (Q882186)

A. HLA B27

B. HLA A29

C. HLA B51

D. HLA DR4

Correct Answer: A. HLA B27

Explanation: Option A: HLA B27 It is a case in which the patient presents with pain, photophobia, redness, and decreased vision from the last 3 days . Both eyes of the patient are affected. Examination shows the involvement of the cornea and ciliary nerve . It is the scenario of acute anterior uveitis . HLA, known as human leukocyte antigen, are proteins or markers in most human cells. HLA is used by immune cells to decide which cells belong to the body and which are not. HLA are strongly associated with diseases of the uvea . About 50% of acute uveitis cases are associated with the human leukocyte antigen B27 . Other diseases are also associated with this, such as ankylosing spondylitis, reactive arthritis, and inflammatory bowel disease . The actual role of B27 in triggering the inflammation response is still unknown. People with positive or increased HLA-B27 are more likely to have autoimmune diseases. Option B: HLA A29 It is a leukocyte antigen of class 1 type . It is associated with birdshot retinochoroiditis (BRC) , which affects the posterior portion of the eye with choroidal inflammation. It is mostly seen in Caucasians as a rare form of uveitis. HLA A29 is a major histocompatibility complex. Option C: HLA B51 HLA B51 is the major genetic factor in Behcet's syndrome. It is a hallmark of this syndrome. It also shows

susceptibility to the virus that can cause German measles . Patients carrying this allele have 4 times greater chances of developing Bechet's disease. Option D: HLA DR4 The presence of this leukocyte antigen is found in the white blood cells . It is a genetic anomaly. HLA DR4 is associated with rheumatoid arthritis . It is a large serogroup and has a moderate frequency of the alleles.

References:

Parsons disease of eye, 22.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 29-year-old man presents with sudden, severe visual loss following a recent road traffic accident involving chest trauma. Fundus examination reveals multiple white retinal patches near the optic disc, cotton wool spots, and minimal intraretinal hemorrhage. Based on these findings, what is the most likely diagnosis? (Q853784)

A. Diabetic Retinopathy

B. Retinal Vein Occlusion

C. Purtscher Retinopathy


D. Central Serous Retinopathy

Correct Answer: C. Purtscher Retinopathy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Purtscher Retinopathy is a condition typically associated with trauma (especially head or chest trauma) and is characterized by multiple retinal white patches around the optic nerve head and fovea, known as Purtscher flecken, cotton wool spots, and minimal intraretinal hemorrhage. The occlusion of peripapillary arterioles and embolization is the key pathophysiological mechanism. In this case, the sudden severe visual loss after trauma, along with the described retinal findings, suggests Purtscher Retinopathy. Diabetic Retinopathy (Option A) is associated with microvascular changes due to diabetes, including dot-blot hemorrhages, microaneurysms, and neovascularization, but not Purtscher flecken or trauma-related visual loss. Retinal Vein Occlusion (Option B) presents with diffuse retinal hemorrhages, cotton wool spots, and optic disc swelling but is not typically linked to trauma. Central Serous Retinopathy (Option D) involves serous retinal detachment usually due to stress, without the hemorrhages or cotton wool spots seen in Purtscher Retinopathy.



References:

https://www.ncbi.nlm.nih.gov/books/NBK542167/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 30-week premature infant weighing 1.4 kg is diagnosed with stage 3 retinopathy of prematurity (ROP) in zone II without plus disease. What is the most appropriate next step in management? (Q853783)

A. Laser photocoagulation

B. Intravitreal anti-VEGF injection

C. Weekly observation and follow-up

D. Cryotherapy of retina

Correct Answer: C. Weekly observation and follow-up

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Stage 3 ROP (retinopathy of prematurity without plus disease in zone II is considered to be Low risk Pre threshold. It typically warrants close observation with follow-up rather than immediate treatment unless there are signs of progression. Anatomical classification of ROP by Zones: Zone I Bounded by a circle, the radius of which is twice the distance between the center of the optic disc and center of the macula. Zone II Extends from the edge of the zone 1 peripherally to a point tangential to the nasal ora serrata. Zone III Consists of residual temporal crescent of retina anterior to zone II. Threshold Disease: Stage 3+ ROP with plus disease. Located in zone I or II. Involves 5 continuous or 8 discontinuous clock hours. Requires laser therapy within 72 hours. Prethreshold Disease: Classified by the Early Treatment for Retinopathy of Prematurity (ET-ROP) group into two types: Type I

Type II High-Risk Prethreshold Disease , requiring laser photocoagulation. Zone I: Plus disease, any stage. Zone I: Non-plus disease, stage 3. Zone II: Plus disease, stages 2 and 3. Low-Risk Prethreshold Disease , requiring weekly follow-up. Zone I: Non-plus disease, stages 1 and 2. Zone II: Non-plus disease, stage 3. Treatment of ROP: Laser treatment: (Option A) Done in the avascular peripheral retina. Use diode laser or frequency doubled Nd:YAG with LIO (laser indirect ophthalmoscope) delivery. Used for high-risk pre-threshold, threshold, and aggressive posterior ROP. Anti-VEGF: (Option B) Bevacizumab: Effective in patients with zone I, stage 3 ROP with plus disease. Zone I disease responds more than zone II. Recurrence can occur following cessation of anti-VEGF treatment. Pars plana vitrectomy: For tractional retinal detachment not involving macula. Cryotherapy of retina (Option D) is not preferred for stage 3 ROP, especially in zone II without plus disease. It has been largely replaced by laser therapy due to its higher risk of causing inflammation and less favorable outcomes.

References:

Parson’s diseases of the eye, Edition 23,PDF, Page 509,510,532-534 Kanski’s Clinical Ophthalmology, Edition 9, page no: 538-542 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9369302/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 64-year-old male with long-standing hypertension comes for a routine eye checkup. On fundoscopy, the ophthalmologist observes the following changes as shown in the image below. What is the most likely stage of hypertensive retinopathy? (Q853696)

A. Grade I

B. Grade II

C. Grade III

D. Grade IV




Correct Answer: C. Grade III


- *Listen (English)*
- *Listen (Hindi)*


Explanation: The fundoscopy image shows Grade III Hypertensive Retinopathy characterized by generalized arteriolar narrowing , hard exudates arranged in a macular star pattern, and cotton wool spots with no papilledema. Staging of Hypertensive Retinopathy (Keith and Wagner Classification): Stage Key Features Grade I (Option A) Mild generalized arteriolar narrowing, broadening of the light reflex. {{caption_text}} Grade II (Option B) Focal arteriolar narrowing, A-V nicking (Salu's sign) {{caption_text}} Grade III (Option C) Flame-shaped hemorrhages, cotton wool spots, and more pronounced A-V nicking (Bonnet and Gunn signs), hemorrhages, and exudates. {{caption_text}} Grade IV (Option D) All features of Grade III + Papilledema, Silver-wiring of arterioles. {{caption_text}}


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot



References:

Parson's diseases of the eye, Edition 23,PDF, Page 530-534 Kanski's Clinical Ophthalmology,Edition 9, Page 532-535

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 58-year-old male with a 20-year history of poorly controlled type 2 diabetes and an HbA1c of 10.5% came for an ophthalmic examination. Fundoscopy reveals microaneurysms, hard exudates, and retinal thickening. Which of the following is the most likely cause of future vision loss in this patient? (Q853695)

A. Vitreous hemorrhage

B. Retinal detachment

C. Macular edema


D. Neovascular glaucoma


Correct Answer: C. Macular edema

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The most likely cause of future vision loss in this patient is macular edema. Complications of Diabetic Retinopathy: Macular Edema: The most common cause of vision loss in diabetic retinopathy is leakage from dilated capillaries. Vitreous Hemorrhage (VH) (Option A): Seen as blood floating in the vitreous, occluding the view of the retina. Secondary to proliferative diabetic retinopathy (PDR) is a cause of severe vision loss in diabetic patients Retinal Detachment (Option B): PDR causes scars to form in the back of the eye. When the scars pull the retina away from the back of the eye, it's called tractional retinal detachment. Neovascular Glaucoma (Option D): PDR causes abnormal blood vessels to grow out of the retina and block fluid from draining out of the eye, leading to neovascular glaucoma. Can cause vision loss and blindness. Rubeosis Iridis: Neovascularization of the iris is when blood vessels develop on the anterior surface of the iris in response to retinal ischemia. Prone to spontaneous hyphema as the blood vessels are fragile





References:

Parsons Diseases of the Eye, 23rd Edition, Page 289,290 Yanoff and Duker Ophthalmology, 5th Edition, Page 544-551 Kanski's Clinical Ophthalmology, 9th Edition, Page 507,508
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5712956/ https://www.ncbi.nlm.nih.gov/books/NBK559131/
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3980167/
https://www.nei.nih.gov/learn-about-eye-health/eye-conditions-and-diseases/diabetic-retinopathy
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5989268/

↑ *Back to this question in Part A*

@Neet_pg_bot


Q6: A 45-year-old male presents with complaints of severe headache and blurred vision. His blood pressure is 220/120 mmHg. Fundoscopy image is given below. Which of the following fundus changes is most characteristic of this diagnosis? (Q853697)

A. Arteriovenous nicking

B. Copper-wiring

C. Papilledema

D. Hard exudates




@Neet_pg_bot

Correct Answer: C. Papilledema

- *Listen (English)*
- *Listen (Hindi)*

Explanation: The given fundus photograph demonstrates Blurred optic disc margins → Papilledema, f lame-shaped hemorrhages, retinal exudates, venous dilation, and narrowed arterioles. These findings, combined with very high BP (220/120 mmHg) and neurological symptoms (headache, blurred vision), confirm malignant hypertensive retinopathy , where papilledema is a hallmark and diagnostic. Arteriovenous nicking (Option A): Seen in chronic hypertensive changes, not specific to acute or malignant phases. Copper-wiring (Option B): Characteristic of chronic arteriosclerotic changes, not acute hypertensive crisis. Hard exudates (Option D): Present in both chronic and acute stages, but not diagnostic.



References:

Parson’s diseases of the eye, Edition 23,PDF, Page 530-532 Kanski’s Clinical Ophthalmology,Edition 9, Page 532-535 .

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 55-year-old patient with a history of uncontrolled hypertension presents for a routine eye examination. Fundoscopy reveals arteriolar narrowing and changes in the arteriovenous (A-V) ratio. What is the most likely A-V ratio finding in this patient with hypertensive retinopathy? (Q853698)

A. 2:3

B. 1:2

C. 1:3

D. 3:2

Correct Answer: C. 1:3

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In hypertensive retinopathy , arteriolar narrowing occurs as a result of vasoconstriction and thickening of the vessel walls. This causes a change in the normal A-V ratio to 1:3 . In normal fundoscopy, the ratio of the diameters of first- and second-order retinal arterioles to the diameters of venules is approximately 2:3 . Hypertensive Retinopathy: Any long-standing HTN causes changes in retina. Keith-Wagener-Barker Grading: Grade Severity Features Grade 1 Mild Diffuse arterial attenuation Grade 2 Moderate Diffuse arterial attenuation Focal arteriolar constriction AV Nicking Salus sign Grade 3 Severe Grade II features Hemorrhage Exudates Gunn sign Bonnet sign Copper wiring Grade 4 Malignant Grade III features Malignant hypertension Papilledema Silver wiring Atherosclerotic changes: Salus sign S-shaped deflection of the vein at arteriovenous crossing due to atherosclerotic plaque Gunn sign Narrowing of the veins at both ends of arteriovenous crossing Bonnet sign Dilatation (Banking) of vein on one side at arteriovenous crossing Copper-wiring Thining of blood column due to atherosclerotic plaque Silver wiring Narrowing of blood column due to increased atherosclerotic plaque Pregnancy-induced HTN: Presents with hypertensive retinopathy Grade III hypertensive retinopathy → Indication for termination of pregnancy

References:

www.ncbi.nlm.nih.gov/books/NBK525980/

iovs.arvojournals.org/article.aspx?articleid=2126986

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: All of the following are considered risk factors for the development of central retinal vein occlusion (CRVO), except? (Q853699)

A. Age over 50 years

B. Open-angle glaucoma

C. Hyperthyroidism

D. Hypertension

Correct Answer: C. Hyperthyroidism

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Hyperthyroidism is not associated with an increased risk of central retinal vein occlusion (CRVO). It primarily affects metabolism and does not directly influence the vascular system, predisposing to CRVO. Risk Factors of CRVO: Age > 50 years (primary risk factor) (Option A) , Hypertension (Option D) , Open-angle glaucoma (Option B) , Diabetes Hyperlipidemia. Additional Risk Factors: Smoking, optic disc edema, oral

contraceptives, hypercoagulable states (e.g., polycythemia, multiple myeloma, antiphospholipid syndrome, hyperhomocysteinemia, etc.), syphilis, HIV, sickle cell disease.

References:

Parson’s diseases of the eye, Edition 23,PDF, Page 531-533 Kanski’s Clinical Ophthalmology,Edition 9, Page 515-524 https://www.ncbi.nlm.nih.gov/books/NBK525985/

↑ *Back to this question in Part A*

@Neet_pg_bot


Q9: A 65-year-old male presents with sudden, severe vision loss in his left eye. On examination, a relative afferent pupillary defect (RAPD) is present. Fundus examination reveals the following changes: Based on these findings, what is the most likely diagnosis? (Q853700)

A. Non-ischemic CRVO

B. Ischemic CRVO

C. Branch Retinal Vein Occlusion (BRVO)

D. Diabetic Retinopathy




Correct Answer: B. Ischemic CRVO

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above clinical features and fundus examination shows multiple flame-shaped hemorrhages around the disc and multiple cotton wool spots. This is known as splashed-tomato appearance. These findings are seen in Ischemic CRVO. Another characteristic appearance seen in ischemic CRVO is 100 day glaucoma (Neovascular glaucoma) Non-ischemic CRVO (Option A) is more common (75% of cases), but it typically presents with milder symptoms, including gradual vision blurring and the absence or mild presence of RAPD. Fundus findings are less severe, with mild venous congestion and flame-shaped hemorrhages in the periphery. Branch Retinal Vein Occlusion (BRVO) (Option C) typically affects a portion of the retina rather than the entire retina, and the hemorrhages are more localized to the area of the occluded vein. Vision loss is generally less severe compared to ischemic CRVO. Diabetic Retinopathy (Option D) can present with retinal hemorrhages and macular edema, but it is not associated with sudden, severe vision loss and RAPD as in this case.



References:

Parson’s diseases of the eye, Edition 23,PDF, Page 531-535

Kanski’s Clinical Ophthalmology,Edition 9, Page 515-524

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Which of the following investigations is most useful in distinguishing ischemic CRVO from non-ischemic CRVO? (Q853701)

A. Fundus Autofluorescence (FAF)

B. Optical Coherence Tomography (OCT)

C. Goldmann Perimetry and Electroretinogram (ERG)


D. Slit-lamp Examination

Correct Answer: C. Goldmann Perimetry and Electroretinogram (ERG)

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Goldmann Perimetry and ERG are crucial for differentiating between ischemic and non-ischemic CRVO. In ischemic CRVO, there is significant retinal ischemia leading to visual field defects and a reduction in the amplitude of the b-wave on ERG . These findings help to confirm the diagnosis of ischemic CRVO. Fundus Autofluorescence (FAF) (Option A) demonstrates fern-like perivenular hypoautofluorescence due to retinal edema in both non-ischemic and ischemic CRVO. Optical Coherence Tomography (OCT) (Option B) is valuable in evaluating macular edema , which may be present in both non-ischemic and ischemic CRVO. Goldmann perimetry and ERG are most useful in differentiating ischemic from non-ischemic CRVO. Slit-lamp examination (Option D) is essential for detecting anterior segment changes such as neovascularization of the iris (NVI). It is mainly used to monitor for complications like neovascular glaucoma.



References:

Parson’s diseases of the eye, Edition 23,PDF, Page 479-484 Kanski’s Clinical Ophthalmology,Edition 9, Page 515-523

↑ *Back to this question in Part A*

Q11: A 72-year-old man with a history of hypertension presents with sudden, painless vision loss in his right eye. Which of the following findings is most consistent with the diagnosis of central retinal artery occlusion? (Q853703)

A. Retinal hemorrhages and cotton wool spots

B. Optic disc edema and retinal hemorrhages

C. Dilated retinal veins with tortuosity


D. Pale retina with a cherry-red spot at the macula

Correct Answer: D. Pale retina with a cherry-red spot at the macula

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In central retinal artery occlusion, the cherry-red spot is seen at the center of the macula due to the vascular choroid shining through the thin foveal retina, contrasting with the surrounding pale retina caused by ischemia. Central Retinal Artery Occlusion (CRAO) - Key Features: Presentation: Sudden painless vision loss Fundus findings: Retinal edema Cherry red spot at macula Marked attenuation (narrowing) of arteries Retinal hemorrhages and cotton wool spots (Option A): Seen in diabetic retinopathy or hypertensive retinopathy Optic disc edema and retina changes (Option B): Seen in optic disc edema / papilledema Dilated retinal veins with tortuosity (Option C): Seen in papilledema (first sign = dilated veins)



References:

Kanski's Clinical Ophthalmology, 9th edition, pg: 525, 527, 528

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 65-year-old male presents to the emergency department with sudden, painless vision loss in his right eye. On examination, the right eye shows a pale retina with a cherry-red spot at the macula. The patient has a history of hypertension and smoking. Which of the following is the most appropriate initial treatment to improve retinal blood flow and potentially restore vision? (Q853704)

A. Oral corticosteroids

B. Intravitreal anti-VEGF injection

C. Ocular massage and anterior chamber paracentesis

D. Systemic anticoagulation therapy

Correct Answer: C. Ocular massage and anterior chamber paracentesis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical features and fundus examination are suggestive of central retinal artery occlusion (CRAO). The goal of the initial treatment is to restore retinal blood flow. Ocular massage and anterior chamber paracentesis are commonly used methods to dislodge emboli and improve perfusion. Critical time to prevent irreversible damage: 90 minutes Treatment Options Intravenous Acetazolamide : Provides sustained IOP reduction Sublingual Isosorbide Dinitrate : Dilates blood vessels to improve blood flow past the blockage Carbogen Breathing Oral corticosteroids (Option A): Used for Giant Cell Arteritis (GCA), which causes an ischemic optic neuropathy, not a cherry-red spot. Intravitreal anti-VEGF injection (Option B): Used for "wet" Macular Degeneration, Diabetic Retinopathy, or Retinal Vein Occlusion (CRVO) to reduce edema, not for acute arterial blockages. Systemic anticoagulation therapy (Option D): Retinal tissue dies within ~90 minutes; systemic anticoagulants are too slow for acute vision rescue.

References:

Kanski’s Clinical Ophthalmology, 9th edition, pg: 529, 530, 531.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A cherry-red spot is seen on the retina due to thickening and loss of transparency of the retina at the posterior pole. Which of the following diseases are least associated with this finding on fundoscopy? (Q853778)

A. Tay-Sachs disease

B. Farber disease

C. Central retinal artery occlusion


D. Central retinal vein occlusion

Correct Answer: D. Central retinal vein occlusion

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Central retinal vein occlusion causes mild blurring of vision and, on fundoscopy, shows superficial flame-shaped hemorrhages, papilledema, mild or no macular edema in acute cases and sheathing of main veins, chronic cystoid edema in chronic cases. A cherry-red spot is not seen in central retinal vein occlusion. Conditions with Cherry-red spot: 1. Storage Disorders Tay–Sachs disease Niemann–Pick disease GM1 gangliosidosis Metachromatic leukodystrophy Sandhoff disease 2. Vascular Central retinal artery occlusion (CRAO) Branch retinal artery occlusion (rare, if macula involved) 3. Others Commotio retinae (Berlin’s edema) Sialidosis Farber’s disease Mucolipidosis



References:

Kanski's Clinical Ophthalmology, 9th edition, pg: 652. https://www.ncbi.nlm.nih.gov/books/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A patient is being examined for loss of vision in the ophthalmology clinic. His peripheral visual field is lost, and only his central vision remains. Which of the following would not present with central (tunnel) vision? (Q853779)

A. Advanced glaucoma

B. Retinitis pigmentosa

C. CRAO with cilioretinal artery

D. Tobacco amblyopia

Correct Answer: D. Tobacco amblyopia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Tobacco amblyopia presents with loss of central vision and not with its preservation. It is caused by excessive use of tobacco which leads to degeneration of ganglion cells, especially in the macula . Causes centrocecal scotoma: Scotoma involving both the blind spot and the macula It characteristically presents with impairment of central vision, which is bilateral and progressive, with patients complaining of difficulty in doing near work and fogginess. Central vision, tubular vision, or tunnel vision refers to the loss of peripheral vision while preserving central vision. Causes of tubular vision: Terminal stage of advanced glaucomatous field defect (Option A) : The visual field loss is gradual, both centrally and peripherally. The temporal island is more resistant and hence is lost only in the end, leading to loss of even light perception. Advanced stage of retinitis pigmentosa (Option B) : Central vision occurs in advanced cases and is lost ultimately after a prolonged duration. Central retinal artery occlusion (CRAO) with cilioretinal artery (Option C): It provides a second arterial supply to the fovea and hence in the case of CRAO, the central vision is maintained whereas the peripheral is lost. Vigabatrin: It is a drug used in the treatment of epilepsy. It can cause vision loss in 30% of infants and causes bilateral tunnel vision in adults, and causes damage to the central retina and leads to visual acuity

References:

Kanski’s Clinical Ophthalmology, 9th edition, pg: 529 Comprehensive Ophthalmology by A.K. Khurana, 7th edition, pg: 241, 296, 297, 332, 546.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A premature infant, born at 26 weeks of gestation, is receiving oxygen therapy in the neonatal intensive care unit. Retinal examination at 4 weeks post-birth reveals signs of retinal vascular abnormalities. Which of the following mechanisms is most likely responsible for the pathogenesis of retinopathy of prematurity in this patient? (Q853780)

A. Increased oxygen supply leading to retinal vasodilation

B. Disruption of VEGF production

C. Excessive blood glucose levels

D. Retinal hypoxia leading to abnormal collagen formation

Correct Answer: B. Disruption of VEGF production

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In retinopathy of prematurity , the premature infant's retina is vulnerable to abnormal vascular growth. The disruption of VEGF production , often due to fluctuations in oxygen levels (e.g., excess oxygen

therapy), impairs the normal retinal vascular development. This can lead to abnormal blood vessel formation and progression of ROP. Increased oxygen supply leading to retinal vasodilation (Option A): Excessive oxygen therapy causes retinal vessel suppression, not vasodilation. Excessive blood glucose levels (Option C): ROP is not directly caused by blood glucose levels, but by oxygen fluctuations. Retinal hypoxia leading to abnormal collagen formation (Option D): Hypoxia triggers VEGF production, causing abnormal vessel growth, not collagen deposition.

References:

Parsons Diseases of the Eye, 23rd Edition, Page 294,531-534

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: Which of the following is the most significant risk factor for Retinopathy of Prematurity (ROP)? (Q853781)

A. High birth weight

B. Prematurity and low birth weight

C. Prolonged exposure to blue light

D. Deficiency of Vitamin A

Correct Answer: B. Prematurity and low birth weight

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Prematurity and Low birth weight is the most significant risk factor for ROP. In retinopathy of prematurity, the premature infant's retina is vulnerable to abnormal vascular growth. The disruption of VEGF production, often due to fluctuations in oxygen levels (e.g., excess oxygen therapy), impairs the normal retinal vascular development. This can lead to abnormal blood vessel formation and progression of ROP. Retinopathy of prematurity (ROP) is a bilateral proliferative retinopathy in premature infants with low birth weight, commonly due to high oxygen exposure. Prematurity > Low birth weight High birth weight (Option A): Babies with higher birth weights are at a lower risk for ROP. Prolonged exposure to blue light (Option C): No scientific evidence links blue light exposure to ROP. Deficiency of Vitamin A (Option D): Vitamin A deficiency can cause xerophthalmia and night blindness, but it is not a primary cause of ROP.

References:

Parsons Diseases of the Eye, Edition 23, page no: 294,295 Yanoff and Duker Ophthalmology Edition 5 page no:539,540 Kanski’s Clinical Ophthalmology, Edition 9, page no: 540

↑ *Back to this question in Part A*

Q17: A 32-week premature infant presents with dilated and tortuous retinal vessels in the posterior pole, poor pupil dilation, and vitreous haze, indicating a progression of retinopathy of prematurity (ROP). What is the most likely diagnosis? (Q853782)

A. Rush disease

B. Plus disease

C. Cicatricial ROP


D. Stage 4 ROP

Correct Answer: B. Plus disease

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: 32-week premature infant with dilated and tortuous retinal blood vessels in the posterior pole, poor pupillary dilatation, vitreous haze indicates progression of ROP (Plus disease) Significance: Indicates progression of ROP. No hope of regression. Poor prognostic sign. Stages of ROP: Stage 1 – Demarcation line Stage 2 – Ridge formation Stage 3 – Neovascularization Stage 4 (Stage D) – Sub-total retinal detachment (RD) Stage 5 – Total retinal detachment (retinal fibrosis) Rush Disease (Option A): Rapid progression of ROP. Cicatricial Stage of ROP (Option C) : Vitreoretinal fibrosis, Scar formation





References:

Parsons Diseases of the Eye, Edition 23, page no: 294,531-534

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A 23-year-old male presents with sudden, painless vision loss in his left eye. Fundus examination reveals peripheral retinal vasculitis, vessel sheathing and recurrent vitreous hemorrhages. Fluorescein angiography reveals vessel leakage. What is the most appropriate initial treatment? (Q853785)

A. Intravitreal anti-VEGF therapy

B. Systemic corticosteroids

C. Laser photocoagulation


D. Vitrectomy

Correct Answer: B. Systemic corticosteroids

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Young male with sudden painless loss of vision in left eye with fundus showing peripheral retinal vasculitis, venous sheathing, and recurrent vitreous hemorrhage confirms the diagnosis of Eales' Disease (Periphlebitis Retina) Etiology & Pathogenesis Cause: Hypersensitivity to tubercular antigen Type of hypersensitivity: Type IV hypersensitivity (cell-mediated) Type III hypersensitivity (immune complex-mediated) Cause of recurrent vitreous hemorrhage in a young male = Eales' Disease Management: Initial management: Systemic corticosteroids (because of inflammatory vasculitis). Further treatment: Laser photocoagulation (Option C) if required. For vitreous hemorrhage: Vitrectomy (Option D) . Prognosis: In Eales' disease, prognosis after vitrectomy is very good, and patient may regain vision up to 6/6. Intravitreal anti-VEGF therapy (Option A) is used in conditions involving neovascularization, such as diabetic retinopathy, they are not the primary treatment for Eales disease, which is mainly driven by inflammation rather than VEGF-mediated neovascularization.



References:

Parsons Diseases of the Eye, Edition 23, Page 318

↑ *Back to this question in Part A*

@Neet_pg_bot


Q19: A 3-year-old child is brought to the clinic with a white reflex (leukocoria) noticed in the right eye. An ocular B-scan ultrasound performed to evaluate the child shows a well-defined, echogenic mass within the globe, with associated calcifications. Which is the most likely diagnosis? (Q853671)

A. Retinal detachment

B. Vitreous hemorrhage

C. Retinoblastoma


D. Choroidal malignant melanoma



Correct Answer: C. Retinoblastoma


- *Listen (English)*
- *Listen (Hindi)*

Explanation: Retinoblastoma is characterized on B-scan ultrasound by a highly reflective intraocular mass, often accompanied by calcifications. B-scan: Two-dimensional dotted section of the eye. The vitreous cavity appears anechoic (black) with the posterior lens capsule visible anteriorly. The retina, choroid, and posterior lens capsule are hyperechoic structures. Applications: Useful for intraocular structure visualization in opaque media. Identifies vitreoretinal and orbital mass lesions. Detects vascular abnormalities and retinal detachments. Localization of intraocular and intraorbital foreign bodies . Retinal detachment Option A: Opaque and corrugated with retinal folds. Vitreous hemorrhage (Option B): Echogenic dotted and linear pattern. Choroidal malignant melanoma (Option C):




Retinoblastoma

@Neet_pg_bot




@Neet_pg_bot



Retinal detachment

@Neet_pg_bot



Vitreous Hemorrhage

@Neet_pg_bot



References:

Parson’s diseases of the eye, 23rd edition, Pg 320 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2765186/ Kanski’s Clinical Ophthalmology,Edition 9, Page 858-860 Yanoff and Duker Ophthalmology,Edition 5, Page 441,798,799

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: What is the approximate distance of the fovea from the optic disc in a normal eye? (Q853652)

A. 1 disc diameter temporal to the optic disc

B. 2 disc diameters nasal to the optic disc

C. 2 disc diameters temporal to the optic disc


D. 1 disc diameter nasal to the optic disc

Correct Answer: C. 2 disc diameters temporal to the optic disc

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The fovea is situated approximately 2 disc diameters (DD) away from the optic disc on the temporal side . 1 disc diameter is equal to about 1.5 mm, so the fovea is located approximately 3 mm from the optic disc.



@Neet_pg_bot

References:

Parson’s diseases of the eye, Edition 23, Page 286.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q21: In which layer of the retina, as labeled in the histology image, are the cell bodies of bipolar cells located? (Q853653)

A. A

B. B

C. C

D. D




@Neet_pg_bot

Correct Answer: B. B

- *Listen (English)*
- *Listen (Hindi)*

Explanation: The Inner Nuclear Layer is made up of the cell bodies of bipolar cells , horizontal cells, and amacrine cells . It is labeled as B in the Histology image given. Layer Description 1. Pigment Epithelium Outermost layer with a single pigmented cell layer; adheres to Bruch's membrane; supports neurosensory retina, and reduces reflection. Subretinal Space: This is the layer between the Pigment epithelium and the layer of Photoreceptors. 2. Rods and Cones Layer Contains the outer segments of photoreceptors (rods and cones, which are the first-order neurons) ; rods detect low-light vision, cones manage color and detailed vision. 3. External Limiting Membrane A porous membrane allows processes from rods and cones to pass through. 4. Outer Nuclear Layer (Henle’s layer) (Option D) Contains the nuclei of the photoreceptor cells (rods and cones). 5. Outer Plexiform Layer (Option C) Includes connections between rods, cones, bipolar cell dendrites, and horizontal cells. 6. Inner Nuclear Layer (Option B) Made up of the cell bodies of bipolar cells (which are the second-order neurons) , horizontal cells, amacrine cells, and Müller cells; and also contains retinal capillaries. 7. Inner Plexiform Layer (Option A) Houses synaptic connections between bipolar cell axons, ganglion cell dendrites, and amacrine cell processes. 8. Ganglion Cell Layer Contains ganglion cell bodies, which serve as third-order neurons ; midget cells synapse with a single bipolar cell, and polysynaptic cells synapse with multiple. 9. Nerve Fiber Layer Composed of ganglion cell axons that form the optic nerve after passing through the lamina cribrosa. 10. Internal Limiting Membrane The innermost layer separating the retina from the vitreous; is formed by the terminal expansions of Müller cells, serving as a basement membrane.



References:

Parson’s diseases of the eye, Edition 23,PDF, Page 38-41

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: A 65-year-old male with a history of hypertension and atherosclerosis presents with sudden, painless loss of vision in his left eye. On fundoscopy, you note a pale retina with a cherry-red spot at the macula. What is the most likely underlying cause of this condition? (Q853654)

A. Diabetic Retinopathy

B. Central Retinal Artery Occlusion

C. Central Retinal Vein Occlusion

D. Optic Neuritis

Correct Answer: B. Central Retinal Artery Occlusion

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The presentation of sudden, painless vision loss with a cherry-red spot at the macula is highly suggestive of central retinal artery occlusion (CRAO) . Most common cause is embolism from the atherosclerotic plaques. Central Retinal Artery Occlusion (CRAO) Etiology: Obstruction at lamina cribrosa (most often embolism from atherosclerotic plaque; also mucormycosis). Symptoms: Sudden, painless, severe vision loss (seconds). May have history of amaurosis fugax . Critical time: Irreversible damage after 90 min . Fundus finding: Cherry-red spot at macula . Causes of Cherry-Red Spot : 1. Storage Disorders (Metabolic) Tay–Sachs disease Niemann–Pick disease GM1 gangliosidosis Metachromatic leukodystrophy Sandhoff disease 2. Vascular Causes Central retinal artery occlusion (CRAO) 3. Others Trauma-related commotio retinae (Berlin's edema) Farber's disease



References:

Parson's diseases of the eye, Edition 23, Pg 3,4,11,250-252

eyewiki.org/Retinal_Vascularization_and_OCT-Angiography_Interpretation

www.ncbi.nlm.nih.gov/books/NBK525985/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Which of the following arteries is primarily responsible for supplying the inner layers of the retina? (Q853655)

A. Central retinal artery

B. Short posterior ciliary arteries

C. Long posterior ciliary arteries


D. Ophthalmic artery

Correct Answer: A. Central retinal artery

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The central retinal artery is the main vessel responsible for supplying the inner layers of the retina, including the nerve fiber layer, ganglion cells, and inner retinal layers. The short posterior ciliary arteries primarily supply the choroid and the outer retinal layers. Blood Supply of Retina Retinal Blood Supply Outer 4 layers of retina → Supplied by short posterior ciliary arteries The same arteries also supply the choroid. Inner 6 layers of retina → Supplied by central retinal artery (CRA). Course of Central Retinal Artery The CRA runs along the end of the optic nerve. This happens before it enters the eye wall. Capillary Systems in Retina Two capillary systems are present: Between the outer plexiform (OP) and the inner nuclear (IN) layer Capillaries are oriented perpendicular to the retina. Bleeding here → Dot and blot hemorrhage. Along the nerve fiber layer Capillaries are oriented parallel to the retina. Bleeding here → Flame-shaped hemorrhage. Ciliary Arteries (Branches of Ophthalmic Artery) Anterior ciliary arteries Posterior ciliary arteries Short posterior ciliary arteries → 10–12 in number. Long posterior ciliary arteries → Only 2 in number. Run anteriorly. Anastomose with anterior ciliary arteries → Major arterial circle. Branches from this circle → Anastomose around the pupillary border → Minor arterial circle.



References:

Parson’s diseases of the eye, Edition 23, Pg 3,4,11,250-252
https://eyewiki.org/Retinal_Vascularization_and_OCT-Angiography_Interpretation
https://www.ncbi.nlm.nih.gov/books/NBK525985/

↑ *Back to this question in Part A*

Q24: A 55-year-old male with a history of poorly controlled diabetes and hypertension presents with blurred vision. Fundus examination shows bright yellow, well-defined deposits arranged in a star-like pattern around the macula and cloud-like white patches near the optic disc. What is the most likely cause of the yellow deposits? (Q853656)

A. Soft exudates due to axoplasmic flow blockage

B. Hard exudates due to leakage of lipids and proteins

C. Retinal hemorrhages from ruptured microaneurysms


D. Cotton-wool spots from ischemic swelling of nerve fibers


Correct Answer: B. Hard exudates due to leakage of lipids and proteins

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: History of a Diabetic and hypertensive male, complaining of blurred vision. Examination Findings of Bright yellow, well-defined deposits around the macula in a star-like pattern are suggestive of Hard exudates Cause: Increased capillary permeability → leakage of lipids Note: Star-like pattern is not always seen; more typical in neuroretinitis → called macular star. Cloud-like white patches near optic disc → Cotton wool spots (soft exudates) Cause: Axonal degeneration. Hard and Soft Exudates Feature Soft Exudates (Cotton-Wool Spots) Hard Exudates Formation Mechanism Caused by ischemia and axonal swelling; blockage of axoplasmic flow (Option A and D) Formed by leakage of fluid rich in proteins and lipids Location Nerve fibre layer Seen around leaking microaneurysms (Option C) , especially in the macula Main Causes Hypertensive retinopathy Diabetic retinopathy Toxemia of pregnancy Anemic retinopathy Lupus erythematosus (LE) Polyarteritis nodosa (PAN) Scleroderma Leukemic retinopathy Retinopathy associated with AIDS Diabetic retinopathy Hypertensive retinopathy Coats' disease Circinate retinopathy Image





References:

Parson’s diseases of the eye, Edition 23, Pg 287,313,318,319

Kanski’s Clinical Ophthalmology,Edition 9, Page 502-503

Yanoff and Duker Ophthalmology,Edition 5, Page 545

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: The blood-retinal barrier separates the blood from the ocular cavity. Which of the following does not constitute the outer blood-retinal barrier? (Q853661)

A. Bruch’s membrane

B. Retinal pigmented epithelium

C. Zona occludens


D. Retinal vascular endothelial cells

Correct Answer: D. Retinal vascular endothelial cells

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The blood-retinal barrier (BRB) consists of an inner and outer barrier that separates the blood from the ocular cavity, where the retinal vascular endothelial cells form the inner blood-retinal barrier. Features Outer blood-retinal barrier Inner blood-retinal barrier Components Retinal pigment epithelium(RPE) (Option B). Zonula occludens (Option C). Bruch’s membrane (Option A) Formed by the tight junctions between the endothelial cells of the retinal blood vessels. Function It prevents the leakage of extracellular fluid and nutrients from the choroid to the outer retina. Controls permeability of the retinal blood vessels.



References:

Yanoff and Duker Ophthalmology, 5th edition, pg: 427, 428, 626, 627, 628.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: A 60-year-old patient with a history of diabetes comes in for a routine check-up for diabetic retinopathy. The patient also has a cataract, causing a hazy view of the ocular media. Which method of study would be most appropriate for assessing the retina of this patient? (Q853662)

A. Direct ophthalmoscopy

B. Indirect ophthalmoscopy

C. Slit-lamp biomicroscopy

D. Amsler grid test

Correct Answer: B. Indirect ophthalmoscopy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: When the media is hazy, the best method for assessing the retina of this patient is indirect ophthalmoscopy. Direct vs. Indirect Ophthalmoscopy: Feature Direct Ophthalmoscopy Indirect Ophthalmoscopy Instrument used Image Erect, virtual Inverted, real Magnification High magnification (~15x in emmetropes, more in myopes, less in hypermetropes) Moderate magnification (~2.5x with a +20D lens) Field of View Small (~10° or 1-2 disc diameters) Large (~35° or 8 disc diameters) Visualization Central retina only Central and peripheral retina (up to the ora serrata) Refractive Media Poor visualization in hazy or cloudy media (Option A) Can visualize through hazy media due to stronger illumination Examination Distance Close to the eye (~15 mm from the cornea) Performed at a greater distance (~50 cm or an arm's length) Patient Position Performed in a seated/standing position in a semi-dark room Performed with the patient in a supine or sitting position Lens Requirement No external lens is required; correcting lenses for refractive errors are built into the ophthalmoscope Requires a convex lens (commonly +20D) to be placed in front of the patient's eye Peripheral Lesion Detection Limited ability to detect peripheral lesions Excellent for detecting peripheral retinal lesions such as retinal detachment or tears. Slit-Lamp Biomicroscopy (Option C): Although useful for anterior segment examination and some posterior pole assessments, it is ineffective in hazy media due to scattering of light. Amsler grid test (Option D) is a widely used tool for monitoring vision changes, particularly in individuals at risk for age-related macular degeneration (AMD) and in other macular lesions. However, it is less effective in hazy media, such as cataracts or other opacities in the eye.





References:

Parson’s diseases of the eye, 23rdedition, Pg 1145-1146 Kanski’s Clinical Ophthalmology,Edition 9, Page 21-23 Yanoff and Duker Ophthalmology,Edition 5, Page 426

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: A 60-year-old patient with age-related macular degeneration (ARMD) undergoes Fundus Fluorescein Angiography (FFA), revealing pooling of dye under the retinal pigment epithelium (RPE) and leakage from newly formed vessels beneath the retina. What abnormality is most likely detected on FFA in this patient? (Q853663)

A. Hyperfluorescence due to dye pooling under detached RPE and neovascularization

B. Hypofluorescence due to blockage of background fluorescence by retinal hemorrhage

C. Hypofluorescence due to occlusion of retinal vasculature


D. Hyperfluorescence due to staining of the optic disc

Correct Answer: A. Hyperfluorescence due to dye pooling under detached RPE and neovascularization

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient has ARMD , a condition associated with the pooling of dye under the RPE and subretinal neovascularization, both of which lead to hyperfluorescence on FFA. Abnormalities Detected by Fundus Fluorescein Angiography (FFA) 1. Hyperfluorescence Window defect (RPE atrophy): Background choroidal fluorescence. Pooling of dye under RPE: Age-related macular degeneration (Option A). Pooling under sensory retina: Central serous retinopathy (CSR). Leakage into neurosensory retina (inner BRB breakdown): Cystoid macular edema (CME). Leakage from choroidal/retinal neovascularization: Proliferative diabetic retinopathy, subretinal neovascular membrane in ARMD. Staining (dye retention in tissues): Drusen, normal optic disc (Option D). Leakage from optic nerve head: Papilloedema. 2. Hypofluorescence Blockage of background fluorescence: Retinal hemorrhage (Option B), hard exudates, pigmented clumps. Occlusion of retinal/choroidal vasculature: Central retinal artery occlusion (Option C), diabetic retinopathy with capillary occlusion. Loss of vasculature: Choroideremia, myopic degeneration. Normal Study



References:

Parson’s Diseases of the Eye, 23rd edition, Pg 315-318 Kanski’s Clinical Ophthalmology,Edition 9, Page 442-447 Yanoff and Duker Ophthalmology,Edition 5, Page 558-563

↑ *Back to this question in Part A*

@Neet_pg_bot

Q28: A 35-year-old male presents with difficulty in vision, especially at night, and he reports bumping into objects frequently. Upon examination, he is found to have tunnel vision, and the fundus reveals no abnormalities in the posterior pole. Which region of the retina is most likely responsible for his symptoms? (Q853651)

A. Macula lutea

B. Fovea centralis

C. Optic disc


D. Peripheral retina

Correct Answer: D. Peripheral retina

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The presentation of diminished night vision and loss of peripheral vision is suggestive of a defect in the peripheral retina , which is crucial for peripheral and night vision. Additionally, no abnormality was found in the posterior pole, which includes the macula, fovea, and optic disc (Options A, B, C) . Parts of the Retina: Part Description Posterior Pole The region behind the retinal equator includes the optic disc and macula lutea ; best examined using slit-lamp biomicroscopy or by direct ophthalmoscopy. Optic Disc (Option C) Known as the optic nerve head, it is a pink, vertically oval structure, marking the start of the optic nerve. The physiological Cup is a depression in the optic disc through which the central retinal artery and vein pass; the absence of photoreceptors results in the physiological blind spot. Macula Lutea (Option A) A dark, yellow spot at the posterior pole, measuring 5.5 mm in diameter; is responsible for central, high-acuity vision and color detection. Fovea Centralis (Option B) Central area of the macula (1.5 mm in diameter); contains only cones and is the most sensitive part of the retina for visual acuity. Foveola A 0.35 mm diameter shining pit within the fovea, represents the area of highest visual acuity, with the foveal avascular zone (FAZ) surrounding it. Peripheral Retina (Option D) Extends from the retinal equator to the ora serrata, examined via indirect ophthalmoscopy or Goldmann three-mirror lens . It contains rods, hence responsible for night vision. Ora Serrata The serrated peripheral margin is where the retina ends. Here the retina is firmly attached to both the vitreous and choroid. Pars plana extends anteriorly from ora serrata.



References:

Parson’s diseases of the eye, Edition 23,PDF, Page 37,38,40,41,739-741

https://www.aao.org/eye-health/anatomy/peripheral-retina

↑ *Back to this question in Part A*

@Neet_pg_bot

Q29: Which of the following electroretinogram (ERG) findings suggests gross disturbances in retinal circulation? (Q853685)

A. Extinguished ERG response

B. Subnormal ERG with a reduced b-wave

C. Negative ERG response


D. Normal ERG with a diminished c-wave

Correct Answer: C. Negative ERG response

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Negative ERG response suggests gross disturbances in retinal circulation. Electroretinography (ERG) records changes in the eye's resting potential. It is induced by a flash of light. Parts of ERG: A-wave: Negative wave, likely from rods and cones (photoreceptors). B-wave: Large positive wave, generated by Müller cells (inner retinal layers) but reflects bipolar cell activity . C-wave: Positive wave, related to the metabolic activity of the pigment epithelium (RPE), observed in dark-adapted eyes. Foveal ERG provides information about the macula used in diagnosing cataracts. Interpretation Response Inference Extinguished Complete failure in rod and cone function (e.g., pigmentary retinal dystrophy, complete retinal artery occlusion, old retinal detachment, advanced siderosis). Subnormal Large areas of the retina not functioning (e.g., early retinitis pigmentosa, before ophthalmoscopic signs). Negative Gross disturbances in the retinal circulation.



References:

Parson’s diseases of the eye, 23rd edition, Pg 323-327 Yanoff and Duker Ophthalmology, 5th edition, Pg 464 Kanski’s Clinical Ophthalmology,Edition 9, Page 618-630

↑ *Back to this question in Part A*

@Neet_pg_bot

Q30: A 14-year-old patient presents with progressive visual impairment. Upon examination, there is evidence of vitelliform lesions in the macula. Electro-oculography (EOG) is performed to assess retinal function. What Arden ratio would be expected in a patient with Best disease on electro-oculography (EOG)? (Q853688)

A. Greater than 2.0

B. Less than 1.5

C. Between 1.5 and 2.0

D. Exactly 1.85

Correct Answer: B. Less than 1.5

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In Best disease, the Arden ratio on EOG is typically less than 1.5, indicating abnormal retinal pigment epithelium function, which is characteristic of this condition. Electro-oculography (EOG): Measures changes in resting ocular potential via electrodes placed at inner and outer canthi when eyes move side to side. Changes in potential reflect the activity of retinal pigment epithelium (regulated by the bestrophin gene) and outer segments of visual receptors. Clinical Use: Useful in detecting retinal pigment epithelium dystrophies and degenerations before visual symptoms manifest. Helpful in diagnosing Best disease (Arden index <1.5) and differentiating it from adult-onset vitelliform macular dystrophy and pattern dystrophy. (Option B) EOG apparatus can also be used to record eye movements in nystagmus (electronystagmography (ENG)). Arden Index: The result is calculated by dividing the light peak (maximal potential in light) by the dark trough (minimal potential in dark). Arden Ratio Interpretation >1.85 Normal 1.85-1.5 Borderline <1.5 Abnormal

References:

Parson’s Diseases of the Eye, 23rd edition, Pg 325-327 Kanski’s Clinical Ophthalmology,Edition 9, Page 618-630 Yanoff and Duker Ophthalmology, 5th edition, Pg 464 https://www.ncbi.nlm.nih.gov/books/NBK537290/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q31: A patient with chronic diabetes mellitus complains of diminished vision. Her fundoscopy showed clinically significant macular edema. Which of the following would not be seen as a progressive change leading to diabetic retinopathy? (Q853689)

A. Macular ischemia

B. Neovascularization

C. Vitreous hemorrhage

D. Retinal vasoconstriction

Correct Answer: D. Retinal vasoconstriction

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Retinal vasoconstriction is seen in the pathogenesis of hypertensive retinopathy, whereas microvascular adhesion is seen in diabetic retinopathy. CSMO (Clinically Significant Macular Edema) is defined as: Any edema within 500 microns from the center of the fovea Any hard exudates within 500 microns from the center of the fovea Pathogenesis of Diabetic retinopathy (DR): Diabetic macular edema → Macular ischemia (Option A) → Retinal new vessels (Option B) → Vitreous hemorrhage (Option C) → Tractional retinal detachment. Hypertensive retinopathy: Systemic hypertension causes vasoconstriction of retinal arterioles (Option D) Arteriosclerosis, increased vascular permeability, and retinal hemorrhages lead to retinopathy.

References:

Kanski's Clinical Ophthalmology, 10th edition, pg: 496, 532, 533, 536.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4651818/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q32: Which of the following statements is false about laser treatment used for vascular diseases of the retina? (Q853690)

A. Focal laser therapy for diabetic macular edema targets leaking microaneurysms within 300 µm of the fovea.

B. Grid laser acts on diffuse retinal thickening, at least 500 µm from the fovea and optic disc.

C. Subthreshold lasers use very short pulses with longer intervals minimizing collateral damage.

D. Limitations include central vision risks, visual field defects, and the need for multiple sessions.

Correct Answer: A. Focal laser therapy for diabetic macular edema targets leaking microaneurysms within 300 µm of the fovea.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Focal laser therapy for diabetic macular edema targets leaking microaneurysms within 500 µm of the fovea. Type of Laser used in retinal vascular diseases : Photocoagulative lasers Grid laser (Option B): Grid laser is applied for diffuse retinal thickening at least 500 microns away from the fovea and optic disc, used when multiple points leak, while focal therapy is reserved for fewer leaks. Sub-threshold lasers (Option C): These are diode lasers given in very short bursts with longer intervals, minimizing collateral damage and used for sub-fovea or juxtafovea pathologies. Limitations (Option D): Photocoagulative lasers carry a risk of central vision loss if the fovea is hit, can cause visual field defects, and usually require multiple sessions.

References:

Kanski's Clinical Ophthalmology, 5th edition, pg: 507, 508, 510, 511, 516, 523, 536.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q33: A 60-year-old diabetic patient presents with worsening vision. Fundoscopy reveals retinal edema, microaneurysms, and hard exudates. Which of the following mechanisms is not responsible for the development of diabetic retinopathy? (Q853691)

A. Protein kinase C activation

B. Reduced blood viscosity

C. Increased proangiogenic factors

D. Breakdown of the blood-retinal barrier

Correct Answer: B. Reduced blood viscosity

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Reduced blood viscosity is not a contributing factor to diabetic retinopathy; instead, increased blood viscosity is observed. Diabetic Retinopathy develops due to: Cellular damage: From sorbitol, AGEs, Protein kinase C activation, oxidative stress (Option A) Hematological-biochemical changes: Increased blood viscosity (Option B) , platelet adhesiveness, and altered lipids Microangiopathy: Breakdown of the inner blood-retinal barrier (Option D) , capillary damage Neovascularization: ■■■■■■■■ ■■■■■■■■ Driven by increased proangiogenic factors■■■■■■■■ (Option C) like VEGF and PDGF, with reduced anti-angiogenic protection.

References:

Parson’s diseases of the eye, Edition 23,PDF, Page 531-535 Kanski’s Clinical Ophthalmology,Edition 9, Page 543-545,547 Yanoff and Duker Ophthalmology,Edition 5, Page 459-463

↑ *Back to this question in Part A*

@Neet_pg_bot

Q34: A 55-year-old patient with a long history of diabetes mellitus presents for a routine eye examination. The ophthalmologist noted several findings in the retinal examination. Which of the following features is least likely to be observed in early nonproliferative diabetic retinopathy (NPDR)? (Q853692)

A. Microaneurysm

B. Dot and blot hemorrhages

C. Hard exudates


D. Venous beading


Correct Answer: D. Venous beading

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Venous beading is typically associated with moderate non-progressive diabetic retinopathy (NPDR) or preproliferative diabetic retinopathy, and indicates sluggish retinal circulation. Early NPDR or Background Diabetic Retinopathy Moderate NPDR or Preproliferative Diabetic Retinopathy Affects small retinal vessels. Earliest sign: Microaneurysm (Option A) Retinal hemorrhages (Option B) Hard exudates (Option C) Edema is not marked. Increased inner retinal hypoxia Cotton-Wool Spots Venous Beading (Option D) Intraretinal Microvascular Abnormalities (IRMA)


Non-proliferative Diabetic retinopathy
Aneurysm
Hart Exudate
Hemorrhage
@Neet_pg_bot


Hemorrhage
Neovascularization
Hard Exudate
Cotton Wool Spots
@Neet_pg_bot

References:

Parsons Diseases of the Eye, Edition 23, page no: 290-292 Yanoff and Duker Ophthalmology, Edition 5, page no: 544,545 Kanski's Clinical Ophthalmology, Edition 9, page no: 498,499,503

↑ *Back to this question in Part A*

@Neet_pg_bot

Q35: A 60-year-old patient with chronic diabetes is diagnosed with diabetic retinopathy. What is the most specific finding for confirming proliferative diabetic retinopathy (PDR) in this patient? (Q853693)

A. Cotton-wool spots

B. Retinal hemorrhage

C. Neovascularization of disc

D. Intraretinal microvascular abnormalities

Correct Answer: C. Neovascularization of disc

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Neovascularization of the Disc (NVD) is the most specific finding for confirming proliferative diabetic retinopathy (PDR) . Proliferative DR = presence of neovascularization NVD (Neovascularization of Disc) NVE (Neovascularization Elsewhere) Other Findings (Not Specific for PDR) Cotton wool spots (Option A) Seen in early changes Numerous → Pre-proliferative DR (Moderate NPDR) Retinal hemorrhages (Option B) Present at all stages of DR Intraretinal microvascular abnormalities (IRMA) (Option D) Represent capillary shunt vessels Feature of Pre-proliferative DR (Moderate NPDR)

References:

Parsons Diseases of the Eye, Edition 23, page no: 291,292 Yanoff and Duker Ophthalmology, Edition 5, page no: 546 Kanski's Clinical Ophthalmology, Edition 9, page no: 505

↑ *Back to this question in Part A*

@Neet_pg_bot

Q36: A 58-year-old male with Type 2 diabetes undergoes a fundus examination, revealing extensive retinal hemorrhages, cotton wool spots, venous beading, and moderate intraretinal microvascular abnormalities (IRMA) in one quadrant. What is the most appropriate follow-up interval for this patient? (Q853694)

A. 2 months

B. 4 months

C. 6 months


D. 12 months

Correct Answer: B. 4 months

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The presence of extensive retinal hemorrhages, venous beading, and moderate IRMA in one quadrant indicates severe non-proliferative diabetic retinopathy (NPDR). For severe NPDR, the follow-up interval is every 4 months. Non-Proliferative Diabetic Retinopathy (NPDR) No Diabetic Retinopathy Review in 12 months Very Mild NPDR Microaneurysms only Review most patients in 12 months (Option D) Mild NPDR Any or all of the: microaneurysms, retinal hemorrhages, exudates, and cotton-wool spots, up to the level of moderate NPDR. No intraretinal microvascular anomalies (IRMA) or significant beading. Review range 6-12 months, depending on the severity of signs, stability, systemic factors, and patient's circumstances Moderate NPDR Severe retinal hemorrhages in 1-3 quadrants or mild IRMA Significant venous beading can be present in no more than 1 quadrant Cotton wool spots commonly present Review in approximately 6 months (Option C) Severe NPDR The 4-2-1 rule; is one or more of the: Severe hemorrhages in all 4 quadrants Significant venous beading in 2 or more quadrants Moderate IRMA in 1 or more quadrants Review in 4 months (Option B) Very severe NPDR Two or more of the criteria for severe NPDR Review in 2-3 months (Option A)



References:

Parsons Diseases of the Eye, 23rd Edition, Page 292,293

Yanoff and Duker Ophthalmology, 5th Edition, Page 548-551

Kanski’s Clinical Ophthalmology, 9th Edition, Page 497,498

↑ *Back to this question in Part A*

@Neet_pg_bot

• [05_Retina] 011 Macular And Fundus Disorders Retinal Detachment And Retinoblastoma

« Back to Folder Directory

[05_Retina] 011 Macular And Fundus Disorders Retinal Detachment And Retinoblastoma — Quick Jump

Q1 (p. 766)	*Q2 (p. 768)*	*Q3 (p. 769)*	*Q4 (p. 770)*	*Q5 (p. 772)*	*Q6 (p. 773)*
Q7 (p. 774)	*Q8 (p. 776)*	*Q9 (p. 778)*	*Q10 (p. 780)*	*Q11 (p. 780)*	*Q12 (p. 784)*
Q13 (p. 784)	*Q14 (p. 786)*	*Q15 (p. 787)*	*Q16 (p. 788)*	*Q17 (p. 790)*	*Q18 (p. 792)*
Q19 (p. 793)	*Q20 (p. 796)*				

Q1: Which of the following is a characteristic finding in the fundus examination of a patient with Stargardt disease? (Q853584)

A. Beaten bronze appearance

B. Cotton wool spots and Hard exudates

C. Dark/silent choroid


D. Honeycomb appearance


Correct Answer: A. Beaten bronze appearance

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Among the given options Beaten bronze appearance is seen in the fundus examination of a patient with Stargardt disease. Stargardt’s disease The most common macular dystrophy. Presents early, typically in the first decade of life. Mainly a dystrophy of retinal pigment epithelium (RPE). Fundus examination findings: Beaten bronze appearance Snail-like appearance On FFA (Fundus fluorescein angiography) : Dark choroid/ silent choroid sign. (Option C) Not a fundoscopy finding. Due to dense lipofuscin deposition in the RPE, which blocks background choroidal fluorescence. Cotton wool spots and hard exudates (Option B): Seen in vascular retinal diseases such as diabetic retinopathy, hypertensive retinopathy, CRVO, etc. Honeycomb appearance (Option D): Seen in cystoid macular edema (CME).


@Neet_pg_bot


@Neet_pg_bot



References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg.159,287, 301,304

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 14-month-old child is diagnosed with retinoblastoma in the right eye. The tumor is confined to the retina, located posterior to equator, measures 2.5 mm in greatest dimension, and is located 3.5 mm from the fovea. There is no vitreous or subretinal seeding. According to the International Classification of Retinoblastoma (ICRB), how would this tumor be classified, and what will be the appropriate treatment? (Q853598)

A. Group A; Systemic chemotherapy with focal consolidation

B. Group B; Plaque radiotherapy

C. Group B; Enucleation

D. Group A; Laser photocoagulation

Correct Answer: D. Group A; Laser photocoagulation

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The tumor in this case fits Group A criteria of the International Classification of Retinoblastoma: Tumor confined to the retina Located posterior to the equator <3 mm in size >3 mm from the fovea and >1.5 mm from the optic disc No subretinal or vitreous seeding Laser photocoagulation is an appropriate focal treatment for small tumors like this in Group A . International Classification of Retinoblastoma (ICRB) Group Description Treatment Group A Small intraretinal tumors (<3 mm), away from foveola/optic disc, no seeding Focal therapy (laser photocoagulation, cryotherapy) Group B All remaining discrete tumors confined to retina, any size or location; no seeding Systemic chemotherapy ± focal therapy Group C Local subretinal/vitreous seeding, discrete Chemotherapy ± focal Group D Diffuse seeding Chemoreduction ± external beam radiation Group E Extensive tumor, neovascular glaucoma, opaque media, invasion Enucleation

References:

pmc.ncbi.nlm.nih.gov/articles/PMC5998397/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 2-year-old child is brought to the ophthalmology clinic with a white pupillary reflex in the right eye, noticed by the parents in flash photography. On examination, the child has strabismus. Which of the following is NOT a condition to be considered in the differential diagnosis of this presentation? (Q853597)

A. Coats' disease

B. Persistent hyperplastic primary vitreous

C. Retinopathy of prematurity

D. Optic nerve glioma

Correct Answer: D. Optic nerve glioma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The case presented describes a classic presentation of retinoblastoma , for which optic nerve glioma is not a differential diagnosis . Optic nerve gliomas usually present with proptosis, visual loss, and optic nerve swelling , but not typically with leukocoria or an intraocular mass with calcification. Various conditions other than retinoblastoma, which present as leukocoria are collectively called pseudoglioma. Causes of Leukocoria: Congenital cataract Inflammatory deposits in vitreous following a plastic cyclitis or choroiditis, Coloboma of the

choroid, Retinopathy of prematurity (Option C) Persistent hyperplastic primary vitreous (Option B) Toxocara endophthalmitis Exudative retinopathy of Coats (Option A)

References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 864, 865

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 2-year-old child presents with leukocoria in the right eye. Fundus examination reveals a white, elevated mass in the retina. Histopathological examination of the tumor shows rosettes with central lumen. Which of the following is the most specific histological feature for this condition? (Q853596)

A. Homer-Wright rosettes

B. Flexner-Wintersteiner rosettes

C. Pseudorosettes


D. Undifferentiated tumor cells

Correct Answer: B. Flexner-Wintersteiner rosettes

■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: The question describes a classic presentation of retinoblastoma , which is the most common primary intraocular malignancy in children. The specific histological features in retinoblastoma is Flexner-Wintersteiner rosettes Flexner-Wintersteiner rosettes : Rosettes with a central lumen. Homer-Wright rosettes (Option A), Pseudorosettes (Option C), Undifferentiated tumor cells (Option D) are histological features seen in retinoblastoma, but are not highly specific. Homer Wright rosettes → pseudorosettes without lumen.



Flexner-Wintersteiner rosettes

@Neet_pg_bot



Homer Wright rosettes

@Neet_pg_bot

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 860-864

AK Khurana Textbook of Ophthalmology, 7th Edition, Page 314, 315

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 2-year-old child presents with the following findings as shown in the image. Genetic testing reveals a mutation in the RB1 gene. Which of the following statements best describes the genetic findings associated with this condition? (Q853595)

A. Retinoblastoma can be inherited in an autosomal dominant manner, with affected individuals having a 100% chance of passing on the mutation.

B. Retinoblastoma typically arises from a germline mutation in the RB1 gene, followed by a second somatic mutation.

C. Affected children with retinoblastoma often have mutations in both the RB1 and TP53 genes.

D. A single mutation in the RB1 gene is usually sufficient for the development of retinoblastoma in both familial and sporadic cases.



Correct Answer: B. Retinoblastoma typically arises from a germline mutation in the RB1 gene, followed by a second somatic mutation.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical image shows leukocoria in the left eye, suggestive of retinoblastoma. Retinoblastoma is typically caused by a germline mutation in the RB1 gene (inherited or de novo) followed by a second somatic mutation or "second hit" in the retina, leading to the development of the tumor. Knudson's Two-Hit Hypothesis: Development of retinoblastoma requires two hits (mutations) in the RB1 tumor suppressor gene (chromosome 13q14). Hereditary cases: First hit = germline mutation (inherited). Second hit = somatic mutation. These children often develop bilateral and multifocal retinoblastomas. Sporadic cases: Both hits occur as somatic mutations. Tumor usually unilateral and unifocal. Retinoblastoma can be inherited in an autosomal dominant manner, with affected individuals having a 100% chance of passing on the mutation (Option A): While retinoblastoma is

inherited in an autosomal dominant pattern, this option does not explain the second-hit mechanism. Affected children with retinoblastoma often have mutations in both the RB1 and TP53 genes (Option C): TP53 mutations are seen in Li-Fraumeni syndrome, not in retinoblastoma, which is primarily associated with RB1 mutations. A single mutation in the RB1 gene is usually sufficient for the development of retinoblastoma in both familial and sporadic cases (Option D): A single mutation in the RB1 gene is insufficient to cause retinoblastoma; a second somatic mutation is required.

References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Page 314

Kanski's Clinical Ophthalmology, 9th Edition, Page 861

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 55-year-old woman with a history of proliferative diabetic retinopathy presents with sudden vision changes in her right eye. She reports seeing floaters and flashes of light. On fundoscopy, there is evidence of fibrous tissue contraction and retinal elevation, but no retinal breaks are seen. What is the most appropriate treatment approach for this patient? (Q853594)

A. Laser photocoagulation

B. Cryotherapy with scleral buckling

C. Pars plana vitrectomy

D. Pneumatic retinopexy


Correct Answer: C. Pars plana vitrectomy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient's symptoms and fundoscopy findings, including fibrous tissue contraction and retinal elevation without retinal breaks, are consistent with tractional retinal detachment . In this condition, the primary treatment approach is pars plana vitrectomy , which removes the vitreous and addresses the traction caused by the fibrous tissue, thereby relieving the retinal detachment. Treatment for tractional retinal detachment: Pars plana vitrectomy (PPV) is the standard approach for tractional retinal detachment. It involves removing cutting the vitreoretinal tractional bands and internal tamponade with either a long-acting gas or silicon oil. Laser photocoagulation (Option A) is used in proliferative retinopathies to prevent further neovascularization but does not address existing tractional detachment. Cryotherapy with scleral buckling (Option B) is effective in rhegmatogenous retinal detachment, not for tractional detachment. Pneumatic retinopexy (Option D) is not suitable for tractional or exudative detachments, as it primarily addresses rhegmatogenous detachment with retinal

breaks.



References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 681, 683-689

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 60-year-old man presents with a sudden onset of flashes of light and a shadow resembling a curtain in the peripheral vision of his right eye. He denies any recent trauma. On examination, fundus evaluation reveals a retinal detachment with a retinal break in the upper temporal quadrant. What is the most appropriate next step in the management of this patient? (Q853588)

A. Intravitreal anti-VEGF injection

B. Pneumatic retinopexy with gas injection

C. Pars plana vitrectomy with silicone oil tamponade


D. Drainage of subretinal fluid with diathermy to seal the break

Correct Answer: B. Pneumatic retinopexy with gas injection

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This patient has a fresh retinal detachment with a small break in the upper quadrant. Pneumatic retinopexy is an appropriate choice for treating retinal detachments with small breaks in the superior retina. Pneumatic retinopexy: A minimally invasive quick and OP based procedure to reattach the retina. Uses an intravitreal gas bubble with cryotherapy or laser to seal retinal breaks. Common gases: Sulfur hexafluoride (SF6) and perfluoropropane (C3F8). Used for uncomplicated retinal detachments. Ideal for small breaks or multiple breaks covering less than two clock hours in the upper peripheral retina. Intravitreal anti-VEGF injection (Option A): Primarily used for neovascular diseases, such as wet AMD, but not for retinal detachment. Pars plana vitrectomy with silicone oil tamponade (Option C): Usually reserved for complicated cases like extensive vitreoretinal traction or multiple breaks. In this case, the detachment is relatively simple, and pneumatic retinopexy is more suitable. Drainage of subretinal fluid with diathermy to seal the break (Option D): While it can be part of the management for retinal detachment, it is not the primary treatment for simple rhegmatogenous detachments with small breaks.



References:

AK Khurana Comprehensive Ophthalmology, 7th Ed, Pg. 311, 312 Kanski’s Clinical Ophthalmology, 9th Edition, Pg. 681

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 55-year-old man reports the sudden appearance of flashes of light, followed by a dark cloud or veil in the peripheral vision of his left eye. Examination reveals a greyish reflex in the affected quadrant of the retina. What type of retinal detachment is most likely in this patient? (Q853587)

A. Rhegmatogenous Retinal Detachment

B. Tractional Retinal Detachment

C. Exudative Retinal Detachment


D. Macular Hole

Correct Answer: A. Rhegmatogenous Retinal Detachment

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Sudden appearance of flashes of light, followed by a dark cloud or veil in the peripheral, examination reveals a greyish reflex in the affected quadrant of the retina are the features of Rhegmatogenous Retinal Detachment Retinal detachment refers to the separation of the neurosensory retina from the retinal pigment epithelium. Rhegmatogenous retinal detachment (RRD): “Regma” = break in the retina. The break can be either: Hole (degeneration, e.g., lattice degeneration in pathological myopia) Tear (traction-induced, usually horseshoe-shaped) Symptoms : Initial flashes of light (photopsia) from traction on rods and cones. Followed by curtain/wheel falling over peripheral vision. Examination: gray reflex and detached retina. Tractional Retinal Detachment (Option B): Often seen in proliferative diabetic retinopathy or trauma, presenting with gradual vision loss, not acute symptoms. Exudative Retinal Detachment (Option C): Associated with systemic or ocular inflammatory diseases and presents without the classic flashes or floaters. Macular Hole (Option D): Causes central vision loss but does not present with peripheral shadows or flashes.





References:

AK Khurana Comprehensive Ophthalmology, 7th Ed, Pg. 309, 310 Kanski's Clinical Ophthalmology, 9th Edition, Pg. 681

↑ *Back to this question in Part A*

Q9: A 10-year-old child presents with a mild reduction in vision in both eyes. Fundus examination reveals a "sunny-side up egg" appearance in the macular region with yellowish subretinal material. The child’s father has a similar history of visual impairment since childhood. What is the most likely diagnosis? (Q853586)

A. Stargardt Disease

B. Best Vitelliform Macular Dystrophy

C. Retinitis Pigmentosa


D. Leber Congenital Amaurosis

Correct Answer: B. Best Vitelliform Macular Dystrophy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical features and examination findings are suggestive of Best Vitelliform Macular Dystrophy. Best’s vitelliform macular dystrophy Fundoscopy shows the classic “sunny-side up” or “egg yolk” lesion in the macula. Caused by lipofuscin deposition in the macula. Inheritance: autosomal dominant. Disease of the retinal pigment epithelium (RPE). Initially, it causes a mild reduction of vision, but vision drops significantly in the later vitelliruptive stage. Vitelliruptive : Lesion fragments (scrambled egg appearance) Stargardt’s disease (Option A): Presents with yellowish flecks in the macula and beaten bronze/snail-track appearance. Retinitis pigmentosa (Option C): A dystrophy of rods and cones, starting in the mid-periphery. Classic triad: waxy disc pallor, attenuated vessels, bone-spicule pigmentation. Leber’s congenital amaurosis (Option D): Severe form of retinal dystrophy presenting with early-onset blindness.





References:

AK Khurana Comprehensive Ophthalmology, 7th Ed, Pg. 301 Kanski's Clinical Ophthalmology, 9th Edition, Pg. 631-634

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: A 16-year-old patient presents with progressive vision loss. On examination, the ophthalmologist notes a central scotoma and yellowish-white flecks in the macular region of the retina. A diagnosis of Stargardt disease is made. All of the following are true statements about Stargardt disease, except: (Q853585)

A. It is also known as juvenile macular dystrophy

B. Fundus examination showing flame-shaped hemorrhages

C. Vitamin A supplementation should be avoided

D. Fundus autofluorescence

Correct Answer: B. Fundus examination showing flame-shaped hemorrhages

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical case and examination findings are suggestive of Stargardt disease. Fundus examination showing flame-shaped hemorrhages is seen in hypertensive retinopathy. Stargardt disease Also known as juvenile macular dystrophy (Option A) , is the most common macular dystrophy. Presents in first or second decade. Investigations: FFA: Dark choroid syndrome or silent choroid syndrome. Fundus autofluorescence (Option D) : Lipofuscin deposits have inherent autofluorescence, and can be detected without a dye. Treatment Avoid bright sunlight. Use sunglasses Avoid vitamin A (Option C) , as it promotes lipofuscinosis.

References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg.283, 301

↑ *Back to this question in Part A*

@Neet_pg_bot


Q11: A 35-year-old male presents with sudden distrubunce of vision in his right eye for two weeks. He reports distortion of images. He has been intermittently using corticosteroids for allergies. His Visual acuity in the right eye is 6/18. FFA and OCT findings are given below. What is the most likely diagnosis? (Q853558)

A. Central serous chorioretinopathy

B. Age-related macular degeneration

C. Cystoid macular edema

D. Retinal vein occlusion






Correct Answer: A. Central serous chorioretinopathy


■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: This patient’s distortion of vision (metamorphopsia), decreased visual acuity to 6/18, history of steroid intake, with OCT showing foveal detachment , and Fluorescein angiography showing smokestack appearance, is suggestive of Central Serous Chorioretinopathy (CSR) Central Serous Chorioretinopathy (CSCR) Demographics: More common in young/middle-aged men Risk factors: Stress, Type A personality, steroids Symptoms: Blurred central vision (decreased visual acuity, typically 6/6 → 6/18) Metamorphopsia (distortion of images) OCT: Serous detachment of neurosensory retina at the fovea Fluorescein angiography: Smokestack / Inkblot appearance Course: Often self-limiting in the acute stage Age-related macular degeneration (Option B): ARMD typically occurs in older patients (>50) and is associated with drusen and neovascularization, which are absent here. Cystoid macular edema (Option C): CME usually follows ocular surgeries or inflammation and presents with cystic spaces on OCT, not the subretinal detachment seen here. Retinal vein occlusion (Option D): RVO presents with retinal hemorrhages and cotton wool spots on examination, not the isolated macular elevation and leakage seen in CSCR.



@Neet_pg_bot





References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg. 302, 303

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: Which of the following is not related to retinitis pigmentosa? (Q853583)

A. Refsum's disease

B. Laurence-Moon-Bardet-Biedl syndrome

C. Kearns-Sayer syndrome

D. Conductive deafness

Correct Answer: D. Conductive deafness

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Retinitis pigmentosa associated with sensory neural deafness- Usher syndrome Systemic associations in Retinitis Pigmentosa Refsum's disease (Option A) : It is a defect in the phytanic acid metabolism. RP+ Cerebellar ataxia+ Polyneuropathy. It is managed with a low phytanic acid diet. Laurence-Moon-Bardet-Biedl syndrome (Option B) : involves RP, mental retardation, obesity, polydactyly, and paraplegia. Usher's syndrome: It is retinitis pigmentosa (RP) with Sensory neural deafness. Kearns-Sayer syndrome (Option C) : involves RP, heart block, and chronic progressive external ophthalmoplegia (CPEO). Bassen-Kornzweig syndrome involves RP, acanthocytosis, abetalipoproteinemia, and ataxia. It is managed with a low-fat diet and vitamin supplements.

References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg.297, 298

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 30-year-old woman presents with progressive night blindness and gradual loss of peripheral vision over the past few years. Her family history reveals that her father had similar symptoms. On examination, you observe characteristic retinal changes, including bone spicule pigmentation. To evaluate the functional status of her retina more accurately, which diagnostic test would you choose to confirm the suspicion of retinitis pigmentosa (RP)? (Q853567)

A. Electroretinography (ERG)

B. Optical Coherence Tomography (OCT)

C. Visual Field Testing


D. Electrooculogram (EOG)

Correct Answer: A. Electroretinography (ERG)

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: ERG is a sensitive diagnostic test in retinitis pigmentosa that shows reduced scotopic, subnormal responses in early disease and becomes completely extinguished in advanced disease. Electroretinogram (ERG) Investigation of choice for retinitis pigmentosa (RP). ERG has an A wave, a B wave, and a small C wave (corresponding to retinal pigment epithelial activity, usually not emphasized). A wave = activity of rods and cones (negative wave). B wave = activity of bipolar cells. In RP, both A and B waves are diminished. This makes ERG the most important diagnostic tool. Optical Coherence Tomography (OCT) (Option B) Not significant in RP. OCT is more useful in macular diseases, not in diagnosing RP. Visual Field Testing (Option C) Done in RP, but not the investigation of choice. Important finding: Early RP → Ring scotoma. Late RP → Tunnel vision.
Electrooculogram (EOG) (Option D) More significant in retinal pigment epithelial dystrophies, such as Stargardt's disease and Best's disease. Not the main investigation for RP.



References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg.297, 298

Kanski's Clinical Ophthalmology, 9th Edition, Pg. 620, 621

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A 32-year-old female presents with difficulty seeing in low light and peripheral vision loss that has gradually worsened over several years. On fundoscopy, the following findings are noted. Which of the following conditions is most likely responsible for these findings? (Q853566)

A. Retinitis Pigmentosa

B. Chronic Central Serous Chorioretinopathy

C. Age-related Macular Degeneration

D. Toxoplasmic Retinochoroiditis



Correct Answer: A. Retinitis Pigmentosa

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The image given above is retinitis pigmentosa fundoscopy findings showing a pale waxy disc, bony spicules, and attenuation of arteries. The Triad of retinitis pigmentosa: Waxy disc pallor Attenuated arteries Bone-spicule pigmentation Retinitis pigmentosa: Inheritance: Most commonly sporadic, but AD, AR, X-linked are all possible. Most common systemic association: Usher's syndrome (RP + deafness). Treatment: No definitive

cure. Trials with docosahexaenoic acid (DHA) showed possible slowing of progression. Chronic Central Serous Chorioretinopathy (Option B): Typically shows a serous detachment at the macula with a ring reflex. Age-Related Macular Degeneration (Option C): Presents with drusen (dry) or choroidal neovascular membranes (wet) at the macula. Toxoplasmic Retinochoroiditis (Option D): Characterized by the classic “headlight in the fog” appearance.

References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg.296, 297, 298, 303

Kanski’s Clinical Ophthalmology, 9th Edition, Pg. 623

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 32-year-old woman presents with gradual central vision loss in both eyes. Fundus examination reveals concentric rings of retinal pigment epithelium (RPE) atrophy and hyperpigmentation centred on the fovea, creating a characteristic appearance. The image showing the appearance is given below: Which of the following is least likely to be associated with this condition? (Q853565)

A. Long-term hydroxychloroquine use

B. Cone dystrophy

C. Chloroquine maculopathy

D. Chronic central serous chorioretinopathy



Correct Answer: D. Chronic central serous chorioretinopathy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The image shows the characteristic appearance of Bull's eye maculopathy , a retinal disorder characterised by concentric rings of RPE atrophy and hyperpigmentation centred on the fovea , creating an appearance resembling a bull's eye target. While chronic central serous chorioretinopathy can cause retinal pigment epithelium changes, it typically does not present with the characteristic bull's eye maculopathy appearance. Etiological causes of bull's eye maculopathy include: Chloroquine maculopathy (Option C) Hydroxychloroquine (Option A) Cone dystrophy (Option B) Battens disease

References:

Parson's Textbook of Ophthalmology, 23rd Edition, Pages 811, 879

Kanski's Clinical Ophthalmology, 9th Edition, Page 627

journals.lww.com/retinalcases/fulltext/2021/07000/bull_s_eye_maculopathy_possibly_due_to_iron.31.aspx

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A 68-year-old woman presents with gradual vision loss in her right eye over the past 6 months. Her ophthalmologist suspects age-related macular degeneration (AMD) and orders further tests. What is the most sensitive modality to detect exudative AMD? (Q853563)

A. Optical Coherence Tomography (OCT)

B. Slit-lamp biomicroscopy using a +90 or +78 D lens

C. Indocyanine green angiography (ICGA)


D. Fluorescein angiography

Correct Answer: A. Optical Coherence Tomography (OCT)

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: OCT is currently considered the most sensitive modality for detecting exudative (wet) AMD as it picks up RPE detachments and early CNV changes non-invasively. Wet (exudative) AMD → hallmark features: Retinal pigment epithelium (RPE) detachment Choroidal neovascularization (CNV) Note: Dry AMD → OCT shows outer retinal tubulations. Slit-lamp biomicroscopy (Option B): basic clinical exam, not sensitive. Indocyanine green angiography (ICG) (Option C) is better for choroidal pathologies because it binds 98% to plasma proteins and persists longer, so occult CNV is better seen. Fluorescein angiography (Option D) can detect CNV, but limited (misses occult CNV).



References:

Parson’s Textbook of Ophthalmology Edition 23rd Pg. 597, 598, 599

AK Khurana Textbook of Ophthalmology Edition 7th Pg. 306, 307

www.ncbi.nlm.nih.gov/books/NBK560778/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A 72-year-old man presents with gradual vision loss in both eyes over the past year. His ophthalmologist diagnoses him with age-related macular degeneration (AMD). Which of the following statements about AMD is correct? (Q853562)

A. AMD is a hereditary degeneration exclusively affecting the macula

B. Wet AMD accounts for 90% of all AMD cases but only 10% of severe vision loss

C. Dry AMD is characterised by choroidal neovascularisation


D. AMD is the most common cause of permanent central visual loss after 6th decade

Correct Answer: D. AMD is the most common cause of permanent central visual loss after 6th decade

■ *Listen to Audio 1*

Explanation: Age-related Macular Degeneration (AMD) is a nonhereditary degeneration involving the choriocapillaris, Bruch's membrane, retinal pigment epithelium, and photoreceptors. It is the most common cause of permanent central visual loss after 6th decade. AMD is a nonhereditary degeneration (Option A) primarily affecting the macula involving the choriocapillaris, Bruch's membrane, retinal pigment epithelium, and photoreceptors. Wet AMD accounts for only 10% of all AMD cases but is responsible for 90% of severe vision loss in AMD patients (Option B). Choroidal neovascularization is a characteristic of wet AMD, not dry AMD. Dry AMD is characterized by drusen, RPE abnormalities, and geographic atrophy (Option C).





References:

Parson’s Textbook of Ophthalmology Edition 23rd Pg. 597, 598, 599

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: Which of the following statements about angioid streaks is incorrect? (Q853561)

A. They represent breaks in Bruch's membrane

B. They are commonly associated with sickle cell anemia

C. Choroidal neovascularization (CNV) frequently develops in these patients

D. Steroids are primarily used for the prevention of CNV

Correct Answer: D. Steroids are primarily used for the prevention of CNV

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Angioid streaks are crack-like dehiscences in Bruch's membrane, appearing as irregular reddish-brown or gray lines radiating from the optic disc. Asymptomatic angioid streaks do not require treatment. However, anti-VEGF therapy can be used to prevent choroidal neovascularization. Angioid Streaks: Associations: Pseudoxanthoma elasticum (PXE) - Most common Ehlers-Danlos syndrome Paget's disease Sickle cell disease (Option B) Complication: Choroidal neovascularization (CNV) frequently develops in affected patients (Option C) Treatment: Anti-VEGF therapy for choroidal neovascularization



References:

Yanoff and Duker Ophthalmology Edition 5th Pg. 603, 604

www.ncbi.nlm.nih.gov/books/NBK538151/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: A 28-year-old woman with a history of high myopia since childhood presents for a routine eye examination. She reports occasional floaters and difficulty seeing at night. On fundus examination, the ophthalmologist notes a tilted optic disc, peripapillary atrophy, and a myopic crescent. Which of the following is NOT typically associated with the condition? (Q853560)

A. Foster-Fuchs spots

B. Lattice degeneration


C. Posterior staphyloma


D. Optic disc edema


Correct Answer: D. Optic disc edema

- *Listen (English)*
- *Listen (Hindi)*

Explanation: The patient's history and fundus findings are consistent with pathological myopia . Pathological myopia is a progressive form of myopia characterized by excessive axial elongation of the eyeball and degenerative changes in the retina and choroid. Optic disc edema is not a feature of pathological myopia; the disc may appear tilted, but edema is absent unless there is another pathology (e.g., papilledema, optic neuritis). Pathological Myopia Pathological/degenerative/progressive myopia is a rapidly progressive error that starts in childhood at 5-10 years of age and results in high myopia (>-6D) during early adult life. Features: Foster-Fuchs spots (Option A) Lattice degeneration (Option B) Posterior staphyloma (Option C)







References:

AK Khurana Textbook of Ophthalmology Edition 7th Pg. 39, 40, 41

↑ *Back to this question in Part A*

@Neet_pg_bot


Q20: A 65-year-old presents with visual blurring in the left eye, 20/40 vision, and a history of cataract surgery three months ago. Fundus examination shows retinal thickening and a dull foveal reflex. OCT and FFA findings are given below. What is the most likely diagnosis? (Q853559)

A. Diabetic Retinopathy

B. Central Retinal Vein Occlusion

C. Cystoid Macular Edema

D. Central Serous Retinopathy






Correct Answer: C. Cystoid Macular Edema

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Cystoid Macular Edema (CME) is associated with fluid accumulation in the outer plexiform and inner nuclear layers of the retina, leading to cystoid spaces. The OCT findings of multiple cystic spaces centered around the foveola. Fundus Fluorescein Angiography shows leakage and dye accumulation in the macular region, with a "flower petal appearance" in late frames. The history of recent cataract surgery further supports this diagnosis, as CME is a known postoperative complication. (also known as Irvine-Gass syndrome) Causes of Cystoid Macular Edema (CME) Post cataract surgery- Irvine-Gass syndrome Diabetic retinopathy Retinal vein occlusion Uveitis. Retinitis pigmentosa. Drugs: Niacin, epinephrine Fluorescein Angiography: Shows leakage and dye accumulation in the macular region, with a "flower petal appearance" in late frames. Optical Coherence Tomography (OCT): Demonstrates loss of foveal depression, cystoid spaces, and overall retinal thickening. Diabetic Retinopathy (Option A): Typically shows diffuse retinal thickening and other diabetic changes, not the cystoid spaces seen in CME. Central Retinal Vein Occlusion (Option B): Often associated with retinal hemorrhages and diffuse edema, but CME's cystoid spaces are more specific. Central Serous Retinopathy (Option D) is characterized by "ink-blot" or "smoke-stack" leakage patterns on FFA, and neurosensory retinal and RPE detachment on OCT. CSR typically does not present with the cystoid pattern seen in CME.





References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pages 303, 304, 305

↑ *Back to this question in Part A*

• [05_Retina] 012 Previous Year Questions

« Back to Folder Directory

[05_Retina] 012 Previous Year Questions — Quick Jump

Q1 (p. 800)	*Q2 (p. 800)*	*Q3 (p. 804)*	*Q4 (p. 806)*	*Q5 (p. 809)*	*Q6 (p. 811)*
Q7 (p. 813)	*Q8 (p. 815)*	*Q9 (p. 817)*	*Q10 (p. 820)*	*Q11 (p. 823)*	*Q12 (p. 823)*
Q13 (p. 824)	*Q14 (p. 826)*	*Q15 (p. 829)*	*Q16 (p. 830)*	*Q17 (p. 834)*	*Q18 (p. 836)*
Q19 (p. 837)	*Q20 (p. 838)*	*Q21 (p. 839)*	*Q22 (p. 840)*	*Q23 (p. 841)*	*Q24 (p. 844)*
Q25 (p. 847)	*Q26 (p. 849)*	*Q27 (p. 851)*	*Q28 (p. 853)*	*Q29 (p. 856)*	*Q30 (p. 857)*
Q31 (p. 858)	*Q32 (p. 858)*	*Q33 (p. 861)*	*Q34 (p. 861)*	*Q35 (p. 862)*	*Q36 (p. 862)*
Q37 (p. 866)	*Q38 (p. 869)*	*Q39 (p. 870)*	*Q40 (p. 871)*		

Q1: Which of the following is not a possible cause of the salt and pepper appearance of the fundus in a patient? (Q857752)

A. Rubella

B. Syphilis

C. Refsum’s disease

D. Tuberculosis

Correct Answer: D. Tuberculosis

Explanation: Tuberculosis is NOT associated with ‘salt and pepper’ appearance of fundus among the given options. Differential diagnoses of Salt & pepper fundus: Prenatal Rubella (Option A) Congenital Syphilis (Option B) Varicella Mumps Retinitis pigmentosa Leber congenital amaurosis Drugs: Thioridazine (anti-psychotic) Syndromes associated with Retinitis pigmentosa: Usher syndrome Kearns-Sayre syndrome Refsum’s disease (Option C) Bardet-Biedl syndrome Bassen-Kornzweig syndrome

References:

Kanski’s Clinical Ophthalmology 9th Edition, pages 230,472,477,478,622-624.

AK Khurana Comprehensive Ophthalmology 7th edition, pages 546.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A myopic patient presents with complaints of flashes and floaters. On examination, a deep anterior chamber is seen. What is the likely diagnosis? (Q866835)

A. Exudative retinal detachment

B. Rhegmatogenous retinal detachment


C. Tractional retinal detachment


D. Central serous retinopathy

Correct Answer: B. Rhegmatogenous retinal detachment

Explanation: A myopic patient with flashes (photopsia) , floaters , and a deep anterior chamber suggests posterior vitreous detachment with retinal tear. This allows liquefied vitreous to pass into the subretinal space, leading to rhegmatogenous retinal detachment . Key mechanism: Myopia → long axial length → retinal thinning + lattice degeneration → Retinal hole formation→Fluid entry under retina→Rhegmatogenous RD Deep anterior chamber in this context is due to elongation of the globe and posterior shift of the lens–iris diaphragm in high myopia. Type of Retinal Detachment Clinical Features Rhegmatogenous Retinal Detachment 1. Floaters (dark spots) due to rapid vitreous degeneration. 2. Photopsia (flashes of light) caused by retinal irritation from vitreous movements. 3. Gradual visual field loss, initially localized, progressing to total vision loss as detachment spreads. 4. Dark veil/cloud in front of the eye , particularly when detachment reaches the posterior region. 5. Sudden, painless vision loss in large, central detachment. 6. Retinal breaks may be round, horseshoe-shaped, slit-like, or large anterior dialysis. 7. Vitreous pigments (tobacco dusting or Shaffer sign) are seen in the anterior vitreous with posterior vitreous detachment. 8. Old retinal detachment shows retinal thinning, subretinal demarcation lines (high water marks), and secondary intraretinal cysts. Tractional Retinal Detachment (Option C ruled out) 1. Gradual, progressive visual loss 2. Typically painless 3. No associated flashes or floaters 4. Vitreoretinal bands present 5. No retinal breaks 6. Detached area - concave and localized 7. No shifting fluid Exudative Retinal Detachment (Option A ruled out) 1. Painless and sudden or gradual visual loss 2. No photopsia, holes/ tears, folds and undulations. 3. Smooth and convex with pigmentary disturbances 4. Rounded and fixed with neovascularization on the tumor summit. 5. Shifting fluid - changing the position of the detached area with gravity - hallmark feature. Central serous retinopathy (Option D): Localized serous detachment of the neurosensory retina at the macula, seen in young stressed males; presents with central vision blurring and metamorphopsia, not flashes/floaters.


ENJOY NOW
@Neet_pg_bot


Horseshoe shaped
Grey reflex
@Neet_pg_bot



Tractional retinal detachment associated with anteroposterior and bridging traction

@Neet_pg_bot



Area of intraretinal exudates anterior to bullous retinal detachment

@Neet_pg_bot

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Which of the following is not seen in very severe NPDR? (Q854705)

A. Venous beading


B. Neovascularization of disc


C. Retinal hemorrhage

D. Microaneurysm

Correct Answer: B. Neovascularization of disc

Explanation: Neovascularization (such as on the disc) appears only in proliferative diabetic retinopathy (PDR) , which is a more advanced stage of diabetic retinopathy, involving the growth of new blood vessels. Therefore, neovascularization of the disc is not part of very severe NPDR . Classification of Non-Progressive Diabetic Retinopathy (NPDR) Severity Definition Very mild nonproliferative diabetic retinopathy Microaneurysms only Mild nonproliferative diabetic retinopathy Microaneurysms plus hard exudates, cotton-wool spots, and/or mild retinal hemorrhages Moderate nonproliferative diabetic retinopathy Microaneurysms plus mild intraretinal microvascular abnormalities or moderate retinal hemorrhages Severe nonproliferative diabetic retinopathy Any one of the following (4-2-1 Rule) Severe retinal hemorrhages in four quadrants Venous beading in at least two Moderately severe intraretinal microvascular abnormalities (IRMA) in at least one quadrant Very severe nonproliferative diabetic retinopathy Any two of the following (4-2-1 Rule) Severe retinal hemorrhages in four quadrants (Option C ruled out) Venous beading in at least two (Option A ruled out) Moderately severe intraretinal microvascular abnormalities (IRMA) in at least one quadrant (Option D ruled out) Clinical Features: Early NPDR or Background Diabetic Retinopathy Severe NPDR or Preproliferative Diabetic Retinopathy Affects small retinal vessels. Earliest sign: Microaneurysm Fluorescein angiography (FA): Tiny hyperfluorescent dots. Retinal hemorrhages: Both deep, i.e., in the inner nuclear or outer plexiform layer (dot and blot hemorrhage) and superficial, i.e., in nerve fiber layer (splinter or flame-shaped) hemorrhages Hard exudates: White or yellow waxy patches with well-defined margins, often at the posterior pole, in the outer plexiform layer. FA: Hypofluorescence with dense exudates. Edema is not marked. Increased inner retinal hypoxia: Leads to multiple retinal hemorrhages, cotton-wool spots, venous beading, vascular loops, intraretinal microvascular abnormalities (IRMAs), and large areas of capillary nonperfusion. Cotton-Wool Spots: Also called soft exudates or nerve fiber infarcts. Result from ischemia, not exudation. Caused by obstruction of axoplasmic flow, leading to swelling and a white, fluffy appearance. FA: Focal hypofluorescence due to local ischemia. Venous Beading: Indicates sluggish retinal circulation. Typically adjacent to large areas of capillary nonperfusion. (Option D) Intraretinal Microvascular Abnormalities (IRMA): Dilated capillaries act as collateral channels. Bypass capillary bed and is often seen adjacent to capillary nonperfusion. FA: Focal hypofluorescence {{caption_text}}





References:

Parsons Diseases of the Eye, Edition 23, page no: 290-292

Yanoff and Duker Ophthalmology, Edition 5, page no: 544,545

Kanski's Clinical Ophthalmology, Edition 9, page no: 498,499,503

↑ *Back to this question in Part A*

@Neet_pg_bot


Q4: The image below displays the findings of a fundoscopy. What is the probable diagnosis? (Q858140)

A. CRAO

B. CRVO

C. Diabetic retinopathy


D. Cystoid macular edema




Correct Answer: A. CRAO

Explanation: therosclerosis Age Hypertension Cardiac Diseases Atrial fibrillation : promotes blood clot formation, increasing CRAO risk Mitral valve prolapse: this can lead to blood flow abnormalities and clot formation Bacterial endocarditis: causes vegetation that can travel and occlude the retinal artery Thrombophilic disorders

Hyperhomocysteinemia: elevated homocysteine levels increase clot formation risk Antiphospholipid syndrome: an autoimmune disorder causing increased clot formation risk Inherited anticoagulant defects: genetic conditions disrupting natural blood clotting regulation Giant cell arteritis (GCA) Inflammatory condition affecting blood vessels, often with severe visual loss Other factors Smoking Diabetes mellitus Oral contraceptive pills Clinical features: Symptoms: Sudden, painless, profound vision loss Painless, except in GCA: GCA-associated CRAO may present with headache, jaw claudication, scalp tenderness, and/or pain around the temple Clinical Signs: Visual Acuity (VA): severely reduced, often to counting fingers or worse Relative Afferent Pupillary Defect (RAPD): Profound and sometimes total (amaurotic pupil) Fundus Examination: Retinal Whitening: Pale and opaque retina , particularly in the posterior pole Cherry-Red Spot: Bright red area at the fovea due to the contrast between pale retina and underlying choroidal circulation {{caption_text}} Cattle truck appearance: Due to segmentation of blood column in retinal veins {{caption_text}} Attenuated Arteries: Narrowed and segmented retinal arterioles Optic Disc Changes: Pale optic disc over time due to optic atrophy from prolonged ischemia is seen in chronic cases. Fluorescein Angiography (FA) Arterial Filling : Delayed Masking of Background Fluorescence : Edematous retina blocks normal fluorescence from the choroid Patent Cilioretinal Artery : Area of normal perfusion if a cilioretinal artery supplies the fovea, potentially preserving central vision {{caption_text}} Other Diagnostic Tests Reveals retinal edema, structural changes, and possibly an embolic plaque within the retinal artery and other systemic causes. Cranial MRI or CT Carotid duplex scanning Echocardiography Thrombophilia screening CRVO (Option B) (Central Retinal Vein Occlusion): Fundoscopic findings- Diffuse retinal hemorrhages (often in all quadrants), venous dilation, and cotton wool spots. The optic disc may be swollen, and there may be blood and thunder appearance. This is different from CRAO, where the retina is pale and a cherry-red spot is seen. Diabetic Retinopathy (Option C): Fundoscopic findings- Microaneurysms, retinal hemorrhages, cotton wool spots, and hard exudates. These findings reflect chronic vascular changes in diabetes and are distinct from the acute ischemic signs seen in CRAO, which involve retinal pallor and cherry-red spots. Cystoid Macular Edema (Option D) (CME): Fundoscopic findings- Cystic spaces in the macula, often with focal macular edema. This causes central vision blurring, but unlike CRAO, it doesn’t show retinal pallor or cherry-red spots. CME is more related to macular dysfunction rather than ischemia.


@Neet_pg_bot


Cattle trucking of vessel
Cherry red spot
@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology, 9th edition, pg: 525, 527, 528

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: In a three-year-old boy, his parents have noticed a progressive decline in coordination over the last six months. During a physical examination, a growth resembling a cauliflower is observed in the left eye, along with leukocoria and the absence of a red reflex. Microscopic analysis of the sample reveals the presence of Flexner-Wintersteiner rosettes. Which gene is associated with this condition? (Q858150)

A. Rb gene


B. ATP 7B gene

C. Rhodopsin gene

D. FBN 1 gene

Correct Answer: A. Rb gene

Explanation: ge of presentation: Retinoblastoma typically occurs in young children, usually before 5 years of age. Leukocoria: This is the most common presenting sign of retinoblastoma, often described as a white pupillary reflex. Fundus examination: A white, elevated mass in the retina is characteristic of retinoblastoma. Most specific histopathological feature of retinoblastoma is Flexner-Wintersteiner rosettes , which are rosettes with a central lumen as shown in the diagram below. {{caption_text}} Histopathology and Diagnostic modalities for Retinoblastoma Histopathology Flexner-Wintersteiner rosettes (highly specific)- These are composed of an 'empty' lumen surrounded by columnar cells Homer-Wright rosettes - These are pseudo-rosettes and consist of cells surrounding a central lumen made up of their processes Undifferentiated tumor cells Areas of necrosis and calcification Clinical Presentation Leukocoria (60%) Strabismus (20%) Painful red eye (7%) Poor vision (5%) Asymptomatic (3%) Fundus Examination White, elevated mass in the retina Can be endophytic, exophytic, or mixed growth Imaging CT scan: Calcification in 90% of cases MRI: Better for evaluating optic nerve and extraocular extension Ultrasound: Useful for initial screening Genetic Testing RB1 gene mutation on chromosome 13q14 ATP 7B gene (Option B) : Mutations in the ATP7B gene are associated with Wilson's disease , a genetic disorder affecting copper metabolism, leading to copper buildup in various organs, including the liver and brain. It does not cause retinoblastoma. Rhodopsin gene (Option C) : The rhodopsin gene is associated with retinitis pigmentosa , a genetic condition causing progressive vision loss, but not with retinoblastoma. FBN1 gene (Option D) : Mutations in the FBN1 gene are responsible for Marfan syndrome , a connective tissue disorder affecting the heart, eyes, and bones, but not retinoblastoma.



References:

AK Khurana Textbook of Ophthalmology Edition 7th Pg. 314, 315

↑ *Back to this question in Part A*

Q6: Which of the listed conditions is associated with the presence of punched-out lesions on the retina? (Q855211)

A. Posterior staphyloma


B. Fundal coloboma

C. Congenital Rubella

D. Toxoplasmosis

Correct Answer: D. Toxoplasmosis

Explanation: The most probable diagnosis in this scenario is toxoplasmosis uveitis, which is caused by Toxoplasma gondii Toxoplasmosis is caused by the parasite Toxoplasma gondii and is a common cause of posterior uveitis. It can lead to punched-out lesions in the retina, which are characteristic of the disease. These lesions typically appear as well-defined, round areas of retinal scarring. Fundoscopy shows focal necrotizing retinitis surrounded by a distinctive “headlight in the fog” appearance An immunocompromised person living with cats is a risk factor for toxoplasmosis Toxoplasmosis uveitis: It is a protozoan infestation. Organism: Toxoplasma gondii Definitive host - cats Intermediate hosts - humans Systemic lesions are more common in immunocompromised patients Clinical manifestation: Most common cause of posterior uveitis and focal necrotizing retinitis (90% of cases) Congenital Inactive disease - Healed punched out heavily pigmented chorioretinal scars in the macular area {{caption_text}} In active inflammation - “headlight in the fog” appearance due to intense vitritis {{caption_text}} Acquired Asymptomatic Same as congenital Punctate outer retinal toxoplasmosis Treatment: Quadruple therapy: Pyrimethamine + Sulfadiazine + Clindamycin + Oral steroid Steroid eye drops for anterior uveitis Posterior staphyloma (Option A) : This refers to an outpouching of the sclera in the posterior part of the eye, often seen in high myopia (nearsightedness). It typically does not cause punched-out lesions on the retina. Fundal coloboma (Option B) : A coloboma is a congenital defect that can affect various parts of the eye, including the retina. It usually results in a hole or defect in the eye structure but does not typically present as punched-out lesions. Congenital Rubella (Option C) : Rubella can cause retinal changes in infants born with the infection, but it is more associated with "salt-and-pepper" retinopathy rather than punched-out lesions.





References:

Parsons Diseases of the Eye, 23rd edition, PDF pg no 574-78

Ocular toxoplasmosis: a review of the current diagnostic and therapeutic approaches - PMC (nih.gov)

Kanski’s Clinical Ophthalmology, 9th edition, Page 458-61

↑ *Back to this question in Part A*

@Neet_pg_bot


Q7: What is the likely cause of sudden loss of vision in the left eye for the past two hours in a 58-year-old man who experienced it after his retirement party? The fundoscopic examination reveals a similar image shown below. (Q857205)

A. BRAO

B. CRAO

C. CRVO


D. BRVO




Correct Answer: B. CRAO

Explanation: The given image shows a cherry-red spot. The orange reflex from the intact choroid stands out at the thin foveola, in contrast to the surrounding pale retina, giving rise to a ‘ cherry-red spot ’ appearance. Sudden loss

of vision and cherry red spots are characteristic of Central Retinal Artery Occlusion (CRAO) . Central Retinal Artery Occlusion (CRAO): Risk Factors Atherosclerosis Age Hypertension/Cardiac Diseases Thrombophilic Disorders Smoking Diabetes Mellitus Oral Contraceptive Pills Clinical Features Symptoms: Sudden, painless, profound vision loss . Painful only in GCA: May present with headache, jaw claudication, scalp tenderness, and/or pain around the temple. Clinical Signs: Severely reduced Visual Acuity (VA), often to counting fingers or worse. Profound and sometimes total Relative Afferent Pupillary Defect (RAPD), also called an amaurotic pupil. Examination Findings Fundus Examination: Retinal Whitening: Pale and opaque retina, particularly in the posterior pole. Cherry-Red Spot: Bright red area at the fovea due to the contrast between pale retina and underlying choroidal circulation. {{caption_text}} Cattle Truck Appearance: Segmentation of the blood column in retinal veins. {{caption_text}} Attenuated Arteries: Narrowed and segmented retinal arterioles. Optic Disc Changes: Pale optic disc over time due to optic atrophy from prolonged ischemia (chronic cases). Investigations Fluorescein Angiography (FFA): Delayed arterial filling. Masking of background fluorescence due to edematous retina. Patent Cilioretinal Artery: Area of normal perfusion of a cilioretinal artery supplies the fovea, potentially preserving central vision. Other Diagnostic Tests: Retinal edema, structural changes, and possibly an embolic plaque within the retinal artery. Cranial MRI or CT. Carotid duplex scanning. Echocardiography. Management Smoking cessation Ocular massage IOP control agents: Acetazolamide Hyperosmotics: Mannitol Carbogen (95% O2 + 5% CO2) inhalation Nd-YAG laser embolectomy





References:

Kanski's Clinical Ophthalmology 9th edition, pages 525-531

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A male patient, aged 18, seeks medical attention due to an eye injury sustained while climbing a tree five days ago. The patient reports experiencing pain, sensitivity to light (photophobia), and redness in the affected eye for the past two days. Upon performing a slit-lamp examination, a lesion is observed as depicted below. Which of the following characteristics is associated with this particular condition? (Q857224)

A. Reverse hypopyon

B. Dendritic ulcer


C. Ring abscess

D. Satellite lesion



Correct Answer: D. Satellite lesion

Explanation: The clinical scenario of pain, redness, photophobia with presence of thick hypopyon following trauma with vegetative matter is suggestive of Fungal Keratitis . Satellite lesions are characteristic of Fungal keratitis. Fungal Keratitis: Etiology Causative organism: Aspergillus fumigatus >> Fusarium Predisposing factor: Vegetative matter injury (trauma) Clinical Features SIGNS >> symptoms Pain & Foreign Body Sensation Watering/ Photophobia Blurred Vision, Redness Examination Findings Corneal Ulcer: Greyish-white , dry, elevated, rolled-out margins Pigmented Ulcer: Brownish, dematiaceous fungi Feathery Extensions: Finger-like projections in stroma Sterile Immune Ring: Yellow line of demarcation Satellite Lesions (Option D): Small lesions around the main ulcer Hypopyon: Large, immobile, non-sterile , even in small ulcer {{caption_text}} Investigations Corneal Scraping: For microbiological examination Wet KOH Mount: Filamentous fungi or yeast-like fungi (Candida) Culture: Sabouraud's agar for fungal culture PCR: For rapid identification of pathogens Corneal Biopsy: For deep stromal abscess or negative cultures Treatment Topical Antifungal Agents: Natamycin 5% , Voriconazole, Amphotericin-B, Nystatin Oral Antifungal Agents: Ketoconazole, Voriconazole (for severe ulcers) Surgical Management: Repeated Scraping: Removal of necrotic material Cauterization Options: Carbolic acid (100%), Trichloroacetic acid (10%–20%), Povidone-iodine (5%) Therapeutic Keratoplasty: Corneal transplantation Reverse hypopyon (Option A): Collection of silicon oil in the anterior chamber is reverse hypopyon, usually seen as a complication of Vitreoretinal surgeries. Dendritic ulcer (Option B): This is not dendritic ulcer as dendritic ulcer is a feature of epithelial keratitis and not fungal corneal ulcer. Ring abscess (Option C): Ring abscess is seen in helminthic keratitis. OP136



References:

Kanski's Clinical Ophthalmology 9th Edition, Pages 217, 218

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: What is the likely diagnosis for a 7-year-old boy who showed a white pupillary reflex and underwent enucleation, with histopathological examination showing Flexner Wintersteiner rosettes? (Q857233)

A. Retinoblastoma


B. Medulloblastoma


C. Rhabdomyosarcoma

D. Astrocytoma

Correct Answer: A. Retinoblastoma


Explanation: A paediatric patient with leukocoria (white pupillary reflex) and Histiopathology revealing Flexner-Wintersteiner Rosettes is diagnostic of Retinoblastoma . Flexner-Wintersteiner rosette is an example of True rosette and is a highly specific feature of Retinoblastoma. {{caption_text}} {{caption_text}} {{caption_text}} Clinical features and Diagnostic modalities for Retinoblastoma: Clinical Presentation Leukocoria (60%) Strabismus (20%) Painful red eye (7%) Poor vision (5%) Asymptomatic (3%) Fundus Examination White, elevated mass in the retina Can be endophytic, exophytic, or mixed growth Imaging CT scan: Calcification in 90% of cases MRI: Better for evaluating optic nerve and extraocular extension Ultrasound: Useful for initial screening Histopathology Flexner-Wintersteiner rosettes (highly specific): These are composed of an ‘empty’ lumen surrounded by columnar cells Homer-Wright rosettes: These are pseudo-rosettes and consist of cells surrounding a central lumen made up of their processes Undifferentiated tumor cells Areas of necrosis and calcification Genetic Testing RB1 gene mutation on chromosome 13q14 Medulloblastoma (Option B): is a rapidly growing tumor of the brain, is unlikely to present with eye symptoms initially. Histopathology may reveal small round blue cells with Homer-wright pseudorosettes . IHC shows positivity for Synaptophysin+, CD56+, GFAP+ . Rhabdomyosarcoma (Option C): presents with acute painful proptosis. Usually seen in boys in the 1st decade of life. Histopathology may reveal small round blue cells with Homer-wright pseudorosettes . IHC shows positivity for Desmin+ . Astrocytoma (Option D): Histopathology would reveal Fried-egg appearance of tumor cells, Eye manifestations are unlikely to be seen in Astrocytoma. {{caption_text}}






Homer Wright rosettes

@Neet_pg_bot



@Neet_pg_bot



References:

AK Khurana Textbook of Ophthalmology 7th Edition, Pages 314,315.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q10: What is the condition characterized by painless vision loss? (Q857731)

A. Central retinal arterial occlusion

B. Branched retinal artery occlusion

C. Central retinal vein occlusion


D. Branched retinal vein occlusion



Correct Answer: C. Central retinal vein occlusion

Explanation: The above fundoscopy image given is that of ‘splashed tomato’ appearance and sudden painless loss of vision are classical features of ischaemic central retinal vein occlusion (CRVO) . Splashed sauce appearance is characterized by severe tortuosity and engorgement of all branches of the central retinal vein involving the peripheral and posterior retina. CRVO: Risk Factors Age > 50 years (primary risk factor), Hypertension , Diabetes Mellitus, Hyperlipidemia, Glaucoma (increased IOP), Smoking, Hypercoagulable states Pathogenesis Thrombosis driven by venous stasis, endothelial damage, and hypercoagulability Types Non-ischemic CRVO (Venous stasis retinopathy) (75% of cases) Mild to moderate, unilateral blurring of vision Fundus: Mild venous congestion and tortuosity, flame-shaped hemorrhages, mild macular edema Ischemic CRVO (Hemorrhagic retinopathy) Severe, sudden visual loss , massive venous engorgement, "splashed-tomato" hemorrhages, cotton wool spots, macular edema Marked vein sheathing, collaterals, neovascularization, chronic cystoid edema {{caption_text}} Investigations Ocular Exam: Visual acuity, IOP, Slit-lamp exam (NVI), Gonioscopy (NVA) Goldmann Perimetry and ERG: Differentiating ischemic vs non-ischemic CRVO Fundus Fluorescein Angiography (FFA): Assess retinal perfusion Optical Coherence Tomography (OCT): Assess macular edema, sub-retinal fluid Treatment Systemic Management: Control of systemic conditions (hypertension, diabetes, hyperlipidemia, hyperhomocysteinemia) Medical Treatment: Intravitreal anti-VEGF (e.g., Bevacizumab) , intravitreal steroids (Triamcinolone, Dexamethasone implant) Laser Therapy: Panretinal photocoagulation (PRP) for neovascularization Surgical: Pars plana vitrectomy for complications Complications Neovascular glaucoma, Neovascularization of the iris (NVI), Vitreous hemorrhage, Proliferative retinopathy, Chronic macular edema Central retinal artery occlusion (CRAO) (Option A): Cherry red spot seen on fundoscopy. {{caption_text}} Branched retinal artery occlusion (BRAO) (Option B): Focal area of ischaemia due to vascular occlusion of branch of retinal artery. Branched retinal vein occlusion (BRAO) (Option B): Sectorial superficial retinal hemorrhages are seen on fundoscopy.





References:

Kanski's Clinical Ophthalmology 9thEdition pages 383-385,672-674,515-523

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A 55-year-old male patient presents to the clinic with a complaint of sudden visual blurring that lasted for a short duration . He reports that during the episode, he completely lost vision in his left eye for a few minutes and then regained his vision . The patient denies any pain, headache, or other associated symptoms during the episode. What is the most likely diagnosis ? (Q820648)

A. Amaurosis fugax

B. Central retinal artery occlusion

C. Central retinal vein occlusion

D. Hypertensive retinopathy

Correct Answer: A. Amaurosis fugax

Explanation: Correct Option A - Amaurosis Fugax Given the patient's clinical presentation of sudden, transient loss of vision in one eye, without pain, and the subsequent restoration of vision, along with the description of a curtain-like visual disturbance, is suggestive of amaurosis fugax Incorrect Options: Option B - Central Retinal Artery Occlusion: Central retinal artery occlusion involves the sudden blockage of the central retinal artery, leading to sudden and painless vision loss in one eye. However, in this condition, the vision loss is usually more severe and sustained, often leaving the patient with very poor vision or even complete blindness in the affected eye. The sudden and brief nature of the episode described by the patient makes central retinal artery occlusion less likely. Option C - Central Retinal Vein Occlusion: Central retinal vein occlusion results from a blockage in the central retinal vein, causing a sudden painless loss of vision. However, the presentation of central retinal vein occlusion typically involves a more gradual and painless decrease in vision, without the sudden and transient nature of the episode described by the patient. Option D - Hypertensive Retinopathy: Hypertensive retinopathy is associated with long-standing high blood pressure that leads to changes in the retinal blood vessels. It can cause symptoms such as blurred vision, but the sudden and transient nature of the episode, along with the complete loss and restoration of vision, is not consistent with hypertensive retinopathy.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: What is the primary mode of dissemination of retinoblastoma? (Q856706)

A. Optic nerve invasion

B. Hematogenous

C. Extraocular extension through sclera

D. Lymphatic spread

Correct Answer: A. Optic nerve invasion

Explanation: The primary route of spread in retinoblastoma is via direct invasion along the optic nerve . The tumor extends posteriorly through the lamina cribrosa , reaching the optic nerve , and may spread to the brain and meninges . This route is most strongly associated with poor prognosis . Other routes of spread: Hematogenous spread: Secondary route, usually to the bone marrow, brain, and liver . Extraocular (scleral) extension: May occur via invasion through the sclera into the orbit, but it is not the main dissemination route . Lymphatic spread: Extremely rare, as the eye lacks lymphatic channels .

References:

Parson’s Diseases of the Eye 23rd Edition, Pages 394-398

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: What is the most probable diagnosis for a 7-year-old patient who presented with a white pupillary reflex and underwent enucleation, with histopathological examination revealing the presence of Flexner Wintersteiner rosettes? (Q856178)

A. Retinoblastoma

B. Medulloblastoma


C. Rhabdomyosarcoma


D. Astrocytoma

Correct Answer: A. Retinoblastoma

Explanation: linical Presentation Leukocoria (60%) Strabismus (20%) Painful red eye (7%) Poor vision (5%) Asymptomatic (3%) Fundus Examination White, elevated mass in the retina Can be endophytic, exophytic, or mixed growth Imaging CT scan: Calcification in 90% of cases MRI: Better for evaluating optic nerve and extraocular extension Ultrasound: Useful for initial screening Histopathology Flexner-Wintersteiner rosettes (highly specific): These are composed of an ‘empty’ lumen surrounded by columnar cells Homer-Wright rosettes: These are pseudo-rosettes and consist of cells surrounding a central lumen made up of their processes Undifferentiated tumor cells Areas of necrosis and calcification Genetic Testing RB1 gene mutation on chromosome 13q14 Incorrect Options: Medulloblastoma (Option B): is a rapidly growing tumor of the brain, is unlikely to present with eye symptoms initially. Histopathology may reveal small round blue cells with Homer-wright pseudorosettes . IHC shows positivity for Synaptophysin+, CD56+, GFAP+ . Rhabdomyosarcoma

(Option C): presents with acute painful proptosis. Usually seen in boys in the 1st decade of life. Histopathology may reveal small round blue cells with Homer-wright pseudorosettes . IHC shows positivity for Desmin+ . Astrocytoma (Option D): Histopathology would reveal Fried-egg appearance of tumor cells, Eye manifestations are unlikely to be seen in Astrocytoma. {{caption_text}}





References:

AK Khurana Textbook of Ophthalmology 7th Edition, Pages 314,315.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: Shifting fluid sign is seen in? (Q855022)

A. Exudative retinal detachment

B. Traction retinal detachment


C. Rhegmatogenous retinal detachment


D. Retinal dialysis


Correct Answer: A. Exudative retinal detachment

Explanation: Shifting Fluid sign is seen in Exudative retinal detachment Retinal detachment refers to the separation of the neurosensory retina from the retinal pigment epithelium. Etiology: Age: most common in 40-60 years. Sex. More common in males (M:F - 3:2). Myopia. About 40% cases of rhegmatogenous retinal detachment are myopic. Aphakia and pseudophakia. Trauma. It may also act as a predisposing factor. Senile posterior vitreous detachment (PVD): It is associated with retinal detachment in many cases. Retinal degenerations predisposed to retinal detachment: Lattice degeneration, Snail track degeneration, Acquired or degenerative retinoschisis, and Focal pigment clumps. Type of Retinal Detachment Clinical Features Rhegmatogenous Retinal Detachment (Option C) 1. Floaters (dark spots) due to rapid vitreous degeneration. 2. Photopsia (flashes of light) caused by retinal irritation from vitreous movements. 3. Gradual visual field loss, initially localized, progressing to total vision loss as detachment spreads. 4. Dark veil/cloud in front of the eye , particularly when detachment reaches the posterior region. {{caption_text}} 5. Sudden, painless vision loss in large, central detachment. 6. Retinal breaks may be round, horseshoe-shaped, slit-like, or large anterior dialysis. {{caption_text}} 7. Vitreous pigments (tobacco dusting or Shaffer sign) are seen in the anterior vitreous with posterior vitreous detachment. 8. Old retinal detachment shows retinal thinning, subretinal demarcation lines (high water marks), and secondary intraretinal cysts. Tractional Retinal Detachment (Option B) 1. Gradual, progressive visual loss 2. Typically painless 3. No associated flashes or floaters 4. Vitreoretinal bands present 5. No retinal breaks 6. Detached area - concave and localized 7. No shifting fluid Exudative Retinal Detachment (Option A) 1. Painless and sudden or gradual visual loss 2. No photopsia, holes/ tears, folds and undulations. 3. Smooth and convex with pigmentary disturbances 4. Rounded and fixed with neovascularization on the tumor summit. 5. Shifting fluid - changing the position of the detached area with gravity - hallmark feature. {{caption_text}} Retinal dialysis (Option D) involves a circumferential tear at the ora serrata, often due to trauma, but it does not exhibit the shifting fluid sign as seen in exudative detachments.


@Neet_pg_bot


Horseshoe shaped
Grey reflex
@Neet_pg_bot





References:

AK Khurana Comprehensive Ophthalmology, 7th Ed, Pg. 309, 310

Kanski’s Clinical Ophthalmology, 9th Edition, Pg. 681

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: What is the most probable diagnosis for a 3-year-old boy who has displayed growing clumsiness over the last half a year, has leukocoria, and lacks red reflex in his left eye? After diagnosis, enucleation was performed, and the specimen displayed Flexner-Wintersteiner rosettes under microscopic examination? (Q859361)

A. Congenital glaucoma

B. Retinoblastoma


C. Toxocariasis

D. Neurofibroma

Correct Answer: B. Retinoblastoma

Explanation: The clinical scenario of a 3 year old boy having leukocoria, and lacking red reflex in his left eye is consistent with Retinoblastoma. Retinoblastoma {{caption_text}} Retinoblastoma is the most common intraocular cancer in children. Symptoms include leukocoria (white pupillary reflex) and strabismus (misaligned eyes). Diagnosis is primarily clinical, supported by indirect ophthalmoscopy, orbital ultrasonography, and MRI to assess tumour extent. Histopathology: Flexner-Wintersteiner rosettes is a differentiated type of tumor characterized by columnar cells with a central lumen. {{caption_text}} Biopsies are not recommended due to the risk of spreading the tumour. Treatment depends on tumour size, location, and whether the disease is unilateral or bilateral. Options include enucleation, chemotherapy, and focal therapies like laser photocoagulation. Incorrect Options Congenital glaucoma (Option A) is characterized by epiphora, photophobia, blepharospasm, and buphthalmos. Toxocariasis (Option C) It is a parasitic infection caused by roundworm Toxocara canis that can be acquired through ingestion of contaminated soil or food animals. It can present with leukocoria, strabismus, floaters, eosinophilia, and hypergammaglobulinemia. Neurofibroma (Option D) are tumors that arise from nerve sheath cells and are often associated with the genetic disorder neurofibromatosis. In plexiform neurofibroma S-shaped deformity of the lid is observed.





Flexner-Wintersteiner rosettes

@Neet_pg_bot

References:

Nelson Textbook of Pediatrics, 21st edition, Pg.2697

Ghai Essential Pediatrics, 10th Edition, Pg.634

↑ *Back to this question in Part A*

@Neet_pg_bot


Q16: What is the condition in which the cherry red spot, as shown in the image below, is observed? {{caption_text}} (Q859761)

A. Central retinal artery occlusion (CRAO)

B. Central retinal venous occlusion (CRVO)

C. Branched retinal artery occlusion (BRAO)


D. Branched retinal vein occlusion (BRVO)





Correct Answer: A. Central retinal artery occlusion (CRAO)

Explanation: The presentation of sudden, painless vision loss with a cherry-red spot at the macula is suggestive of central retinal artery occlusion (CRAO). Central retinal artery occlusion (CRAO) The central retinal artery is blocked, causing widespread retinal ischemia. Sometimes, if a cilioretinal artery is present, it can continue to supply the macula, preserving central vision even when the central retinal artery is occluded. Clinical features: Symptoms: Sudden, painless, profound vision loss Painless, except in GCA: GCA-associated CRAO may present with headache, jaw claudication, scalp tenderness, and/or pain around the temple Clinical Signs: Visual Acuity (VA): severely reduced, often to counting fingers or worse Relative Afferent Pupillary Defect (RAPD): Profound and sometimes total (amaurotic pupil) Fundus Examination: Retinal Whitening: Pale and opaque retina , particularly in the posterior pole Cherry-Red Spot): Bright red area at the fovea due to the contrast between pale retina and underlying choroidal circulation {{caption_text}} Cattle truck appearance: Due to segmentation of blood column in retinal veins {{caption_text}} Attenuated Arteries: Narrowed and segmented retinal arterioles Optic Disc Changes: Pale optic disc over time due to optic atrophy from prolonged ischemia is seen in chronic

cases. Fluorescein Angiography (FA) Arterial Filling : Delayed Masking of Background Fluorescence : Edematous retina blocks normal fluorescence from the choroid Patent Cilioretinal Artery : Area of normal perfusion if a cilioretinal artery supplies the fovea, potentially preserving central vision {{caption_text}} Other Diagnostic Tests Reveals retinal edema, structural changes, and possibly an embolic plaque within the retinal artery and other systemic causes. Cranial MRI or CT Carotid duplex scanning Echocardiography Thrombophilia screening Incorrect Options: Central retinal venous occlusion (CRVO) (Option B) occurs when the central retinal vein, becomes blocked. It typically presents with sudden, painless vision loss and may be associated with retinal hemorrhages, cotton wool spots, and retinal edema. Branched retinal artery occlusion (BRAO) (Option C) occurs when one of the branches of the central retinal artery becomes blocked, leading to a localized area of retinal ischemia. The affected area of the retina may appear pale, with associated retinal hemorrhages. Branched retinal vein occlusion (BRVO) (Option D) occurs when one of the branches of the central retinal vein becomes blocked, resulting in impaired venous drainage from a particular area of the retina. It can lead to retinal hemorrhages, cotton wool spots, and macular edema.







References:

Parson’s diseases of the eye, Edition 23, Pg 3,4,11,250-252

Kanski’s Clinical Ophthalmology, 9th edition, pg: 525, 527, 528

https://eyewiki.org/Retinal_Vascularization_and_OCT-Angiography_Interpretation

https://www.ncbi.nlm.nih.gov/books/NBK525985/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: In which diseases are microaneurysms observed? (Q859764)

A. Retinal detachment


B. Diabetic retinopathy


C. Uveitis

D. Glaucoma

Correct Answer: B. Diabetic retinopathy

Explanation: Microaneurysms are localized outpouchings of the capillary wall that may form by focal dilatation of the capillary wall where pericytes are absent or by fusion of two arms of a capillary loop. Diabetic retinopathy Clinical Features: Early NPDR or Background Diabetic Retinopathy Severe NPDR or Preproliferative Diabetic Retinopathy Affects small retinal vessels. Earliest sign: Microaneurysm (Option B) Fluorescein angiography (FA): Tiny hyperfluorescent dots. Retinal hemorrhages: Both deep, i.e., in the inner nuclear or outer plexiform layer (dot and blot hemorrhage) and superficial, i.e., in nerve fiber layer (splinter or flame-shaped) hemorrhages Hard exudates: White or yellow waxy patches with well-defined margins, often at the posterior pole, in the outer plexiform layer. FA: Hypofluorescence with dense exudates. Edema is not marked. Increased inner retinal hypoxia: Leads to multiple retinal hemorrhages, cotton-wool spots, venous beading, vascular loops, intraretinal microvascular abnormalities (IRMAs), and large areas of capillary nonperfusion. Cotton-Wool Spots: Also called soft exudates or nerve fiber infarcts. Result from ischemia, not exudation. Caused by obstruction of axoplasmic flow, leading to swelling and a white, fluffy appearance. FA: Focal hypofluorescence due to local ischemia. Venous Beading: Indicates sluggish retinal circulation. Typically adjacent to large areas of capillary nonperfusion. Intraretinal Microvascular Abnormalities (IRMA): Dilated capillaries act as collateral channels. Bypass capillary bed and is often seen adjacent to capillary nonperfusion. FA: Focal hypofluorescence {{caption_text}} Incorrect Options: Retinal detachment (Option A) involves the separation of the sensory retina from the underlying retinal pigment epithelium (RPE). Uveitis (Option C) is inflammation of the uveal tract (iris, ciliary body, and choroid) and can cause symptoms such as pain, photophobia, and redness. Glaucoma(Option D) primarily involves increased intraocular pressure, leading to optic nerve damage .





References:

Parsons Diseases of the Eye, Edition 23, page no: 290-292

Yanoff and Duker Ophthalmology, Edition 5, page no: 544,545

Kanski's Clinical Ophthalmology, Edition 9, page no: 498,499,503

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: What should be the subsequent course of action in managing a diabetic patient who has a visual acuity of 6/9 in one eye and further investigations reveal preretinal hemorrhages with neovascularization at the optic disc? (Q856002)

A. Focal laser photocoagulation

B. Pan-retinal photocoagulation


C. Grid laser photocoagulation

D. Scleral buckling

Correct Answer: B. Pan-retinal photocoagulation

Explanation: R): No Diabetic Retinopathy Review in 12 months Very Mild NPDR Microaneurysms only Review most patients in 12 months Mild NPDR Any or all of the: microaneurysms, retinal hemorrhages, exudates, and cotton-wool spots, up to the level of moderate NPDR. No intraretinal microvascular anomalies (IRMA) or significant beading. Review range 6-12 months, depending on the severity of signs, stability, systemic factors, and patient's circumstances Moderate NPDR Severe retinal hemorrhages in 1-3 quadrants or mild IRMA Significant venous beading can be present in no more than 1 quadrant Cotton wool spots commonly present Review in approximately 6 months Severe NPDR The 4-2-1 rule; is one or more of the: Severe hemorrhages in all 4 quadrants Significant venous beading in 2 or more quadrants Moderate IRMA in 1 or more quadrants Review in 4 months Very severe NPDR Two or more of the criteria for severe NPDR Review in 2-3 months Proliferative Diabetic Retinopathy (PDR): Mild-moderate PDR New vessels on the disc (NVD) or new vessels elsewhere (NVE), but extent insufficient to meet the high-risk criteria Treatment is considered according to the severity of signs, stability, systemic factors, and the patient's circumstances such as reliability of attendance for review. If not treated, review in up to 2 months High-risk PDR NVD greater than ■ disc area Any NVD with vitreous hemorrhage NVE greater than ½ disc area with vitreous hemorrhage Treatment should be performed immediately when possible and certainly on the same day if symptomatic presentation with a good retinal view Advanced diabetic eye disease Is characterized by tractional retinal detachment. Significant persistent vitreous hemorrhage and neovascular glaucoma. Vitreoretinal surgery with photocoagulation Proliferative Diabetic Retinopathy (PDR): Panretinal Photocoagulation (PRP): PRP of ischemic area reduces metabolic demand and prevents the release of vasoproliferative factors by conversion of hypoxic areas into anoxic areas and creates inert scars from leaking vascular anomalies. Laser used: Argon, frequency-doubled YAG or diode laser In PRP, the inferior quadrant of the retina is first coagulated. Untreated Areas: A circular area with a radius of 2.5 DD around the fovea is left untouched during treatment. Each treatment requires 2000-3000 burns, administered over two to four sessions. Anti-VEGF Therapy: Bevacizumab, ranibizumab or aflibercept are used to control neovascularization. Vitreoretinal Surgery: Advanced cases with tractional retinal detachment or unresolved vitreous hemorrhage

require vitrectomy. Focal laser photocoagulation (Option A) : This is typically used for treating macular edema or localized areas of retinal leakage, not for the widespread ischemic retina seen in proliferative diabetic retinopathy. Grid laser photocoagulation (Option C) : This is also used to treat macular edema but is not the treatment of choice for proliferative diabetic retinopathy . Scleral buckling (Option D) : This procedure is used for retinal detachment , often caused by traction from proliferative diabetic retinopathy, but it's not the initial treatment for PDR. Pan-retinal photocoagulation is used before considering surgery in the event of a detachment.



References:

Parsons Diseases of the Eye, Edition 23, page no: 292,293

Yanoff and Duker Ophthalmology,Edition 5, Page 548-551

Kanski’s Clinical Ophthalmology, Edition 9, page no: 497,498

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: A patient has sudden loss of vision and there is no fundus glow. There is no history of trauma, what could be the likely cause? (Q860200)

A. Vitreous hemorrhage

B. Acute anterior uveitis

C. Optic neuritis

D. Acute congestive glaucoma

Correct Answer: A. Vitreous hemorrhage

Explanation: Vitreous Hemorrhage (VH): Seen as blood floating in the vitreous, occluding the view of the retina. Secondary to proliferative diabetic retinopathy (PDR) is a cause of severe vision loss in diabetic patients. Absence of fundus glow is a feature of Vitreous hemorrhage In PDR, fibrovascular tissue lies flat on the retina or attaches to the posterior vitreous, leading to vitreous traction, retinal separation, and vessel rupture. This is the most common cause of VH in adults. Options B, C & D: Acute anterior uveitis, Optic neuritis and Acute congestive glaucoma can also cause sudden loss of vision but there is no loss of fundus glow.

References:

Parsons Diseases of the Eye, Edition 23, page no: 289,290

Yanoff and Duker Ophthalmology,Edition 5, Page 544-551

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: Which of the following criteria is not included in the definition of severe Non-Proliferative Diabetic Retinopathy (NPDR)? (Q860545)

A. Microaneurysms in all quadrants

B. Retinal hemorrhage in all quadrants

C. Looping in all quadrants

D. Bleeding in two quadrants

Correct Answer: C. Looping in all quadrants

Explanation: Looping in all quadrants is not a criterion in severe NPDR. Abbreviated Early Treatment Diabetic Retinopathy Study (ETDRS) classification of diabetic retinopathy Category Management Non-Proliferative Diabetic Retinopathy (NPDR): No Diabetic Retinopathy Review in 12 months Very Mild NPDR Microaneurysms only Review most patients in 12 months Mild NPDR Any or all of the: microaneurysms, retinal hemorrhages, exudates, and cotton-wool spots, up to the level of moderate NPDR. No intraretinal microvascular anomalies (IRMA) or significant beading. Review range 6-12 months, depending on the severity of signs, stability, systemic factors, and patient's circumstances Moderate NPDR Severe retinal hemorrhages in 1-3 quadrants or mild IRMA Significant venous beading can be present in no more than 1 quadrant Cotton wool spots commonly present Review in approximately 6 months Severe NPDR The 4-2-1 rule; is one or more of the: Severe hemorrhages in all

4 quadrants (Option B ruled out) Significant venous beading in 2 or more quadrants (Option D ruled out) Moderate IRMA in 1 or more quadrants (Option A ruled out) Review in 4 months Very severe NPDR Two or more of the criteria for severe NPDR Review in 2-3 months

References:

Parsons Diseases of the Eye, Edition 23, page no: 292,293

Yanoff and Duker Ophthalmology,Edition 5, Page 548-551

Kanski's Clinical Ophthalmology, Edition 9, page no: 497,498

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: Which investigation is most useful for visualizing the individual layers of the retina? (Q865360)

A. Fundus Fluorescein Angiography

B. Optical Coherence Tomography

C. USG B-scan

D. Slit lamp examination

Correct Answer: B. Optical Coherence Tomography

Explanation: Optical Coherence Tomography (OCT) (Option B): Most useful. Provides high-resolution, cross-sectional images. Allows detailed visualization and measurement of individual retinal layers (e.g., nerve fiber layer, photoreceptor layers). Essential for diagnosing and monitoring retinal diseases. Fundus Fluorescein Angiography (FFA) (Option A ruled out): Visualizes retinal and choroidal blood vessels. Highlights leakage, non-perfusion, or abnormal vasculature. Does not provide layer-by-layer retinal images. USG B-scan (Ultrasound B-scan)(Option C ruled out): Uses sound waves; useful when view is obscured (e.g., hemorrhage). Detects gross abnormalities (e.g., detachment, tumors). Resolution not sufficient to see individual retinal layers. Slit lamp examination(Option D ruled out): Clinical tool for magnified, stereoscopic view of the retina. Good for overall integrity and surface observation. Does not provide cross-sectional, layer-by-layer resolution like OCT.

References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 58

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: A woman with a history of proliferative diabetic retinopathy presents with gradual vision changes in her right eye. She reports seeing floaters and flashes of light. What type of retinal detachment is most likely in this patient? (Q865366)

A. Rhegmatogenous Retinal Detachment


B. Tractional Retinal Detachment

C. Exudative Retinal Detachment

D. Primary Retinal Detachment

Correct Answer: B. Tractional Retinal Detachment

Explanation: A diabetic woman with proliferative retinopathy with gradual visual loss is most likely to be due to Tractional Retinal Detachment. Retinal detachment refers to the separation of the neurosensory retina from the retinal pigment epithelium. Proliferative diabetic retinopathy (PDR) leads to neovascularization on the retinal surface. These new vessels grow along with fibrous scar tissue . Over time, this fibrovascular tissue contracts and pulls the neurosensory retina away from the underlying retinal pigment epithelium, resulting in a tractional retinal detachment . Key clinical clues: Gradual vision loss Seen in PDR and other fibrovascular conditions Caused by mechanical traction - no retinal break required Retinal detachment can be classified into three types: Rhegmatogenous or primary retinal detachment Tractional retinal detachment Exudative retinal detachment Tractional Retinal Detachment : Results from the pulling of the retina by fibrous or vascular tissue, commonly seen in fibrovascular proliferative conditions like diabetic retinopathy, without a retinal break. Primary or Rhegmatogenous Retinal Detachment (Option D and A) : This is the most common type and occurs due to a retinal tear or break, allowing subretinal fluid to enter and separate the sensory retina from the pigmentary epithelium. Exudative Retinal Detachment (Option C) : Caused due to choroidal pathologies or by tumors pushing the retina or fluid accumulation under the retina due to inflammation or vascular abnormalities, without any associated retinal tear or traction.



References:

AK Khurana Comprehensive Ophthalmology, 7th Ed, Pg. 309, 310 Kanski's Clinical Ophthalmology, 9th Edition, Pg. 681

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Which of the following statements is incorrect about Retinitis Pigmentosa? (Q853115)

A. ERG shows decreased amplitude of a and b waves


B. Rod dystrophy > Cone dystrophy


C. Late presentation is loss of peripheral vision


D. Night vision is lost late and rare


Correct Answer: D. Night vision is lost late and rare

Explanation: Night Blindness is the characteristic and earliest feature of retinitis pigmentosa and may present several years before the visible changes in the retina appear. Retinitis Pigmentosa: Retinitis pigmentosa is a dystrophy of rods and cones, initially, rods are affected. (Option B ruled out) Most common hereditary fundus dystrophy. It can be sporadic (40-50%) or hereditary. If hereditary, autosomal dominant is the most common mode of inheritance followed by autosomal recessive (AR). X-linked recessive inheritance is the least severe form. Clinical features: Symptoms: Nyctalopia (night blindness) (Option D) Peripheral visual field loss Mid-peripheral superior scotoma leads to ring/ doughnut scotoma resulting in tubular/ tunnel vision. Central vision is affected in later stages (Option C ruled out) Signs: Triad of a p ale waxy disc due to consecutive optic atrophy, attenuation of arteries , and pigmentation disturbance in the form of bony spicules. Tapetal reflex: shinning reflex at macula - seen in female carriers of XLR inheritance. Atypical Retinitis Pigmentosa Atypical retinitis pigmentosa includes heterogeneous disorders that share clinical features with typical pigmentary retinal dystrophy. Conditions included are: RP sine pigmento - absence or paucity of pigment accumulation Retinitis punctata albescens - scattered whitish-yellow spots with arteriolar attenuation, usually sparing the macula. Sectoral RP - only inferior quadrants are involved. Investigation of Retinitis Pigmentosa Electroretinography (ERG) is the investigation of choice - reduced scotopic rod in early disease and reduced photopic with progression. EOG - subnormal- absence of light rise. Perimetry: initially small mid-peripheral scotomas which gradually merge and may progress, leaving only a small island of residual central vision. Electroretinography (ERG): There's a decreased amplitude of both "a" and "b" waves. (Option A ruled out) OCT- to identify cystoid macular edema (CME). Dark adaptometry: Dark adaptation time is prolonged. Decreased Arden ratio may be present.




B
@Neet_pg_bot


C
@Neet_pg_bot



References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg.296, 297, 298, 303

Kanski's Clinical Ophthalmology, 9th Edition, Pg. 623

↑ *Back to this question in Part A*

@Neet_pg_bot


Q24: Which of the following parts of the retina are correctly marked? (Q853129)

A. 1-Retinal pigment epithelium; 2-Outer nuclear layer; 3-Inner plexiform layer; 4-Ganglion layer

B. 1-Retinal pigment epithelium; 2-Inner nuclear layer; 3-Outer plexiform layer; 4-Ganglion layer


C. 1-Ganglion layer; 2-Inner nuclear layer; 3-Outer plexiform layer; 4-RPE


D. 1-Ganglion layer; 2-Outer plexiform layer; 3-Inner nuclear layer; 4-RPE



Correct Answer: A. 1-Retinal pigment epithelium; 2-Outer nuclear layer; 3-Inner plexiform layer; 4-Ganglion layer

Explanation: Histology of Retina: Layers of the Retina: Layer Description 1. Pigment Epithelium Outermost layer with a single pigmented cell layer; adheres to Bruch's membrane; supports neurosensory retina, and reduces reflection. Subretinal Space: This is the layer between the Pigment epithelium and the layer of Photoreceptors. 2. Rods and Cones Layer Contains the outer segments of photoreceptors (rods and cones, which are the first-order neurons) ; rods detect low-light vision, cones manage color and detailed vision. 3. External Limiting Membrane A porous membrane allows processes from rods and cones to pass through. 4. Outer Nuclear Layer (Henle’s layer) Contains the nuclei of the photoreceptor cells (rods and cones). 5. Outer Plexiform Layer Includes connections between rods, cones, bipolar cell dendrites, and horizontal cells. 6. Inner Nuclear Layer Made up of the cell bodies of bipolar cells (which are the second-order neurons) , horizontal cells, amacrine cells, and Müller cells; and also contains retinal capillaries. 7. Inner Plexiform Layer Houses synaptic connections between bipolar cell axons, ganglion cell dendrites, and amacrine cell processes. 8. Ganglion Cell Layer Contains ganglion cell bodies, which serve as third-order neurons ; midget cells synapse with a single bipolar cell, and polysynaptic cells synapse with multiple. 9. Nerve Fiber Layer Composed of ganglion cell axons that form the optic nerve after passing through the lamina cribrosa. 10. Internal Limiting Membrane The innermost layer separating the retina from the vitreous; is formed by the terminal expansions of Müller cells, serving as a basement membrane. The Neuroglial cells, particularly the vertical Müller cells, bind the retina's neural elements. These cells provide structural support and nourish the retinal tissue.





References:

Parson’s diseases of the eye, Edition 23,PDF, Page 38-41

Ophthalmology by Yanoff and Duker, Edition 5, Page 421

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3258978/

↑ *Back to this question in Part A*

Q25: Which of the following is a complication of diabetic retinopathy? (Q853132)

A. Posterior staphyloma

B. Retinoschisis

C. Tractional retinal detachment

D. Progressive myopia

Correct Answer: C. Tractional retinal detachment

Explanation: Diabetic retinopathy is a progressive microvascular complication of diabetes, which can lead to proliferative changes, causing fibrovascular proliferation and subsequent tractional retinal detachment (TRD). This occurs due to contraction of fibrovascular membranes over the retina, leading to vitreoretinal traction and retinal elevation. Complications of Diabetic Retinopathy: 1. Macular Edema: The most common cause of vision loss in diabetic retinopathy is leakage from dilated capillaries. Clinically significant macular edema (CSME): Retinal thickening at the center of the macula. Retinal thickening and/or adjacent hard exudates at or within 500 µ of the center of the macula. An area of retinal thickening greater than or equal to one disc area, any part of which is within 1 disc diameter of the center of the macula. Types: Focal or Diffuse macular edema Diagnosis: Slit-lamp Biomicroscopy: Loss of foveal reflex and macular thickening Fluorescein Angiography Identifies leakage and also diagnoses ischemic maculopathy. Treatment: Anti-VEGF Therapy: Bevacizumab, ranibizumab or aflibercept. Laser treatment should be deferred for 6 months after commencement of anti-VEGF injection. Intravitreal Triamcinolone acetonide: Reduce macular edema and improve visual acuity. Photocoagulation: Focal Photocoagulation: Treatment of choice for focal diabetic macular edema (DME) not involving the center of the fovea using diode or argon burns. Grid Photocoagulation: Considered in recalcitrant cases unresponsive to anti-VEGF agents or intravitreal steroids. Burns are applied to macular areas of diffuse retinal thickening, treating no closer than 500 µ from the foveola and 500 µ from the optic disc. 2. Vitreous Hemorrhage (VH): Seen as blood floating in the vitreous, occluding the view of the retina. Secondary to proliferative diabetic retinopathy (PDR) is a cause of severe vision loss in diabetic patients. In PDR, fibrovascular tissue lies flat on the retina or attaches to the posterior vitreous, leading to vitreous traction, retinal separation, and vessel rupture. This is the most common cause of VH in adults. 3. Retinal Detachment: PDR causes scars to form in the back of the eye. When the scars pull the retina away from the back of the eye , it's called tractional retinal detachment. 4. Neovascular Glaucoma: PDR causes abnormal blood vessels to grow out of the retina and block fluid from draining out of the eye, leading to neovascular glaucoma. Can cause vision loss and blindness. 5. Rubeosis Iridis: Neovascularization of iris, is when blood vessels develop on the anterior surface of the iris in response to retinal ischemia. Prone to spontaneous hyphema as the blood vessels are fragile. Treatment: Panretinal photocoagulation with or without intravitreal injection of anti-VEGF. Posterior staphyloma (Option A): Posterior staphyloma is associated with pathologic myopia, not diabetic retinopathy. It involves progressive thinning and outward bulging of the sclera, primarily in high myopia. Retinoschisis (Option B): Retinoschisis refers to the splitting of the retinal layers, commonly seen in degenerative retinoschisis or X-linked retinoschisis, rather than as a complication of diabetic retinopathy. Progressive myopia (Option D): Progressive myopia is associated with axial elongation of the eyeball, typically seen in high myopia, rather than diabetic retinopathy.




Clinically significant macular edema (CSME)

@Neet_pg_bot



@Neet_pg_bot



References:

Macular Edema

Vitreous Hemorrhage

Retinal Detachment

Neovascular Glaucoma

Rubeosis Iridis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: Which of the following is the diagnostic investigation for Best disease? (Q853139)

A. EOG


B. ERG

C. Multifocal ERG

D. Visual evoked potential

Correct Answer: A. EOG

Explanation: Electro-oculography (EOG) is the gold standard diagnostic test for Best disease. Best disease/ Best Vitelliform Macular Dystrophy (BVMD) , also known as juvenile-onset vitelliform macular dystrophy, is the second most common macular dystrophy after Stargardt disease. Caused by mutations in the BEST1 gene located on chromosome 11q13. Bestrophin, the protein produced by this gene, functions as a transmembrane ion channel in the retinal pigment epithelium (RPE). Inherited in an autosomal dominant (AD) pattern. Usually begins in childhood Clinical Features : Stages of the disease include: Pre-vitelliform : abnormal EOG, normal fundus in asymptomatic patients. Vitelliform : a characteristic "egg yolk or sunny-side up" macular lesion. The lesions can vary in size and developmental stage between the two eyes. Pseudohypopyon: partially regressed lesion. Vitelliruptive : Lesion fragments, visual acuity declines - “scrambled egg appearance”. Atrophic : Pigment disappears, leading to RPE atrophy. Visual acuity - stable early on but declines in adulthood due to complications like choroidal neovascularization (CNV) , scarring, or geographic atrophy. Diagnosis : Fundus Autofluorescence (FAF) : hyper autofluorescence yellowish material. Fluorescein Angiography (FA) : hypofluorescence due to lesion masking. Optical Coherence Tomography (OCT) : hyperreflective material beneath and within the RPE. Electro-oculography (EOG) : Severely subnormal across all stages (Arden index <1.5). Hallmark - normal electroretinogram (ERG) but abnormal electrooculogram (EOG). (Option B ruled out) (Option A) Best disease progresses through five stages, starting with the "egg-yolk" vitelliform lesion and potentially leading to choroidal neovascularization (CNV). Asymptomatic carriers may show a reduced Arden ratio on EOG, making it useful for family screening. Multifocal ERG (Option C): Multifocal ERG assesses macular function and can show localized dysfunction, but it is not the definitive test for Best disease. Visual evoked potential (Option D): Visual evoked potential (VEP) evaluates optic nerve conduction and is mainly used for optic neuritis, demyelinating disorders, and optic neuropathies. It does not help in diagnosing Best disease.





References:

AK Khurana Comprehensive Ophthalmology, 7th Ed, Pg. 301

Kanski's Clinical Ophthalmology, 9th Edition, Pg. 631-634

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: A 3-month-old baby, born preterm at 33 weeks of gestation, is brought by parents with complaints of a white pupillary reflex for 2 weeks. CT revealed calcification of the globe in both eyes. What is the probable diagnosis? (Q853144)

A. ROP (Retinopathy of Prematurity)


B. Retinoblastoma

C. Glaucoma

D. Congenital Cataract

Correct Answer: B. Retinoblastoma

Explanation: The presentation of a white pupillary reflex (leukocoria) in a preterm infant with calcification of the globe on CT scan is most likely indicative of retinoblastoma, most common primary intraocular malignancy in children. The key features pointing to this diagnosis are: Age of presentation: Retinoblastoma typically occurs in young children, usually before 5 years of age. Leukocoria: This is the most common presenting sign of retinoblastoma, often described as a white pupillary reflex. Fundus examination: A white, elevated mass in the retina is characteristic of retinoblastoma. The most specific histopathological feature of retinoblastoma is Flexner-Wintersteiner rosettes , which are rosettes with a central lumen as shown in the diagram below. Histopathology and Diagnostic modalities for Retinoblastoma Histopathology Flexner-Wintersteiner rosettes (highly specific)- These are composed of an ‘empty’ lumen surrounded by columnar cells Homer-Wright rosettes - These are pseudo-rosettes and consist of cells surrounding a central lumen made up of their processes Undifferentiated tumor cells Areas of necrosis and calcification Clinical Presentation Leukocoria (60%) Strabismus (20%) Painful red eye (7%) Poor vision (5%) Asymptomatic (3%) Fundus Examination White, elevated mass in the retina Can be endophytic, exophytic, or mixed growth Imaging CT scan: Calcification in 90% of cases MRI: Better for evaluating optic nerve and extraocular extension Ultrasound: Useful for initial screening Genetic Testing RB1 gene mutation on chromosome 13q14 Retinopathy of Prematurity (Option A): While ROP is common in preterm infants, especially those born before 32 weeks gestation, it typically presents with abnormal blood vessel development in the retina. Leukocoria in ROP is usually due to retinal detachment or scarring, not calcification. ROP is stage-based and does not typically show calcifications on CT unless complicated by retinal detachment. Glaucoma (Option C): Congenital glaucoma may cause enlarged corneas and optic nerve damage, but it does not usually present with white pupillary reflex or calcification. Congenital Cataract (Option D): Congenital cataracts can also cause leukocoria, but they typically do not show calcification on CT. Cataracts are characterized by lens opacity and not calcification within the globe.



References:

AK Khurana Textbook of Ophthalmology Edition 7th Pg. 314, 315, 317 and 318

https://www.ncbi.nlm.nih.gov/books/NBK562319/

https://www.ncbi.nlm.nih.gov/books/NBK572080/

https://www.ncbi.nlm.nih.gov/books/NBK574553/

↑ *Back to this question in Part A*

@Neet_pg_bot


Q28: A patient presents with vision problems and has a history of cataract surgery. OCT finding is shown below. What is the syndrome most likely associated with these findings? (Q852496)

A. Posner-Schlossman Syndrome

B. Irvine-Gass Syndrome

C. Central Serous Retinopathy


D. Elschnig Pearls





Correct Answer: B. Irvine-Gass Syndrome

Explanation: Irvine-Gass Syndrome, also known as cystoid macular edema (CME), is a condition commonly seen after cataract surgery. The OCT finding typically shows loss of foveal depression, cystoid spaces, and overall retinal thickening, which is indicative of this syndrome. Cystoid Macular Edema (CME) Etiology: Ocular Treatment Complications: Post-surgery (e.g., cataract extraction, retinal detachment surgery), ocular laser therapy,

and topical ocular medications. Retinal Vascular Disorders: Diabetic retinopathy, central and branch retinal vein occlusion, retinal telangiectasia, and hypertensive retinopathy. Intraocular Inflammations: Pars planitis, uveitis. Retinitis pigmentosa. Vitreomacular Traction: Associated with macular epiretinal membrane (ERM) and vitreomacular traction (VMT) syndrome. Systemic Diseases: Includes leukemia, chronic renal failure, and multiple myeloma. Miscellaneous Causes: niacin toxicity , PG analogues (risk factor for development of CME) Clinical Features: Visual Loss: Initial minimal to moderate vision loss, potentially leading to permanent vision decrease if persistent. Ophthalmoscopy: Reveals loss of foveal contour, retinal thickening, a yellow spot at the fovea, and a "honeycomb appearance" of the macula . Fluorescein Angiography: Shows leakage and dye accumulation in the macular region, with a "flower petal appearance" in late frames. Optical Coherence Tomography (OCT): Demonstrates loss of foveal depression, cystoid spaces, and overall retinal thickening. Complications: Long-standing CME may result in a lamellar macular hole . Treatment: Addressing the Cause: Intravitreal anti-VEGF or photocoagulation for diabetic CME; discontinuing causative topical medications. Topical NSAIDs: Such as ketorolac, diclofenac, or profenac, are effective in preventing and treating CME. Topical/Systemic Steroids: Useful in established cases. Systemic Carbonic Anhydrase Inhibitors (CAIs): Oral acetazolamide may be beneficial, particularly in cases associated with retinitis pigmentosa. Posner-Schlossman Syndrome (Option A) : This involves high IOP and mild uveitis, but is not associated with macular edema as seen in Irvine-Gass Syndrome. Central Serous Retinopathy (Option C): is characterized by "ink-blot" or "smoke-stack" leakage patterns on FFA, and neurosensory retinal and RPE detachment on OCT. CSR typically does not present with the cystoid pattern seen in CME. Elschnig's Pearls (Option D): Clustered, vacuolated subcapsular epithelial cells along the posterior capsule.







References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg. 303, 304, 305

↑ *Back to this question in Part A*

@Neet_pg_bot


Q29: A patient presents with a gradual loss of night vision and peripheral vision. What is the most likely diagnosis based on fundoscopy findings? (Q843989)

A. Retinitis pigmentosa

B. Retinal hemorrhage


C. Diabetic retinopathy

D. Hypertensive retinopathy



Correct Answer: A. Retinitis pigmentosa

Explanation: Correct Option A - Retinitis pigmentosa: The patient’s presentation and presence of bony spicules, pale waxy disc, and attenuated arteries are suggestive of retinitis pigmentosa. Incorrect Options: Options B, C, and D do not exhibit the fundoscopy findings given in the question.



↑ *Back to this question in Part A*

@Neet_pg_bot

Q30: A child presents with leukocoria and an intraocular mass. Microscopic examination reveals rosettes with scanty cytoplasm, and the mass is positive for synaptophysin and chromogranin. What is the diagnosis? (Q844034)

A. Retinoblastoma

B. Medulloblastoma

C. Optic glioma

D. Choroidal melanoma

Correct Answer: A. Retinoblastoma

Explanation: The patient is a child with leukocoria, which is the classic and most common presentation of retinoblastoma. The tumor is an intraocular mass, consistent with origin from retinal neuroectoderm. Microscopy shows rosettes with scanty cytoplasm, specifically Flexner–Wintersteiner rosettes, characteristic of retinoblastoma. Positivity for synaptophysin and chromogranin indicates neuronal differentiation, seen in retinoblastoma. Incorrect Options: Option B) Medulloblastoma : Occurs in the cerebellum. Presents with raised ICP, ataxia. Not an intraocular tumor Option C) Optic glioma : Associated with NF-1. Causes visual loss and optic nerve enlargement. Does not show rosettes or neuroendocrine markers Option D) Choroidal melanoma : Seen in adults, not children. Pigmented tumor. Negative for synaptophysin/chromogranin

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q31: A 55-year-old hypertensive patient presented with complaints of blurred vision and headaches. On retinal examination, the following findings were observed: multiple flame-shaped hemorrhages, cotton wool spots, and disc edema. Based on the Keith-Wagner classification, which grade of hypertensive retinopathy is this patient most likely classified as? (Q820501)

A. GRADE 1

B. GRADE 2

C. GRADE 3

D. GRADE 4

Correct Answer: D. GRADE 4

Explanation: Grade 4 Grade 4 hypertensive retinopathy is characterized by severe retinal changes including: Flame-shaped hemorrhages Cotton wool spots Papilledema (disc edema) Presence of disc edema distinguishes Grade 4 from Grade 3, which has hemorrhages and cotton wool spots but no disc edema . Symptoms such as blurred vision and headache indicate malignant hypertension often associated with Grade 4. Keith-Wagner Classification of Hypertensive Retinopathy Grade Fundus Findings Clinical Features Grade 1 Mild arteriolar narrowing Usually asymptomatic Grade 2 Moderate arteriolar narrowing + AV nicking Often asymptomatic or mild symptoms Grade 3 Grade 2 + Flame-shaped hemorrhages + Cotton wool spots + hard exudates Visual symptoms possible Grade 4 Grade 3 + Papilledema (disc edema) Malignant hypertension symptoms like headache, blurred vision Grade 1 (Option A): Incorrect, only mild arteriolar narrowing without hemorrhages or edema. Grade 2 (Option B): Incorrect, involves AV nicking but no hemorrhages or disc edema. Grade 3 (Option C): Incorrect, has hemorrhages and cotton wool spots but no disc edema .

References:

.https://pmc.ncbi.nlm.nih.gov/articles/PMC8300046/ https://pmc.ncbi.nlm.nih.gov/articles/PMC6181956/ https://pubmed.ncbi.nlm.nih.gov/7155523/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q32: A 72-year-old patient, with a history of cataract surgery and diabetes mellitus, presents with decreased vision one month after surgery. Examination reveals a honeycomb appearance or flower petal pattern of the retina. What is the likely diagnosis? (Q860554)

A. Cystoid macular edema


B. Central serous retinopathy


C. Retinal detachment

D. Diabetic retinopathy

Correct Answer: A. Cystoid macular edema

Explanation: The clinical scenario of 72 year old, h/o of cataract surgery, flower petal pattern of retina in examination leads to the diagnosis of CME (Cystoid macular edema) Cystoid Macular Edema (CME) Etiology: Ocular Treatment Complications: Post-surgery (e.g., cataract extraction, retinal detachment surgery), ocular laser therapy, and topical ocular medications. Retinal Vascular Disorders: Diabetic retinopathy, central and branch retinal vein occlusion, retinal telangiectasia, and hypertensive retinopathy. Intraocular Inflammations: Pars planitis, uveitis. Retinitis pigmentosa. Vitreomacular Traction: Associated with macular epiretinal membrane (ERM) and vitreomacular traction (VMT) syndrome. Systemic Diseases: Includes leukemia, chronic renal failure, and multiple myeloma. Miscellaneous Causes: niacin toxicity , PG analogues (risk factor for development of CME) Clinical Features: Visual Loss: Initial minimal to moderate vision loss, potentially leading to permanent vision decrease if persistent. Ophthalmoscopy: Reveals loss of foveal contour, retinal thickening, a yellow spot at the fovea, and a "honeycomb appearance" of the macula . Fluorescein Angiography: Shows leakage and dye accumulation in the macular region, with a "flower petal appearance" in late frames. {{caption_text}} Optical Coherence Tomography (OCT): Demonstrates loss of foveal depression, cystoid spaces, and overall retinal thickening. {{caption_text}} Complications: Long-standing CME may result in a lamellar macular hole . Treatment: Addressing the Cause: Intravitreal anti-VEGF or photocoagulation for diabetic CME; discontinuing causative topical medications. Topical NSAIDs: Such as ketorolac, diclofenac, or profenac, are effective in preventing and treating CME. Topical/Systemic Steroids: Useful in established cases. Systemic Carbonic Anhydrase Inhibitors (CAIs): Oral acetazolamide may be beneficial, particularly in cases associated with retinitis pigmentosa. Central Serous Retinopathy (Option B) is characterized by "ink-blot" or "smoke-stack" leakage patterns on FFA, and neurosensory retinal and RPE detachment on OCT. CSR typically does not present with the cystoid pattern seen in CME. Retinal detachment (Option C) Defined as the separation of the retinal pigment epithelium from neurosensory retina. Diabetic Retinopathy (Option D): Typically shows diffuse retinal thickening and other diabetic changes, not the cystoid spaces seen in CME.





References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg. 303, 304, 305

↑ *Back to this question in Part A*

@Neet_pg_bot

Q33: Which of the following statements is incorrect about Retinitis pigmentosa? (Q840062)

A. ERG show decreased amplitude of a and b wave

B. Rod dystrophy > Cone dystrophy

C. Late presentation is loss of peripheral vision

D. Night vision is lost late and rare

Correct Answer: D. Night vision is lost late and rare

Explanation: Correct Option D - Night vision is lost late and rare: Night Blindness is the characteristic and earliest feature of retinitis pigmentosa and may present several years before the visible changes in the retina appear. It occurs due to degeneration of the rods. Incorrect Options: Option A, B, and C: All these statements are true.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q34: A patient with long-standing rheumatoid arthritis who has been noncompliant with treatment presents with sudden, painless superior field loss in one eye for the past week. Fundus examination shows retinal whitening in the inferior retina with a corresponding superior visual field defect. What is the most likely cause of this condition? (Q840063)

A. Branched retinal artery occlusion

B. Central retinal artery occlusion

C. Branched retinal vein occlusion

D. Central retinal vein occlusion

Correct Answer: A. Branched retinal artery occlusion

Explanation: Branched retinal artery occlusion Rheumatoid arthritis , especially when poorly controlled , can cause systemic vasculitis , including involvement of the retinal arterioles . Branch retinal artery occlusion (BRAO) typically presents as a sectoral visual field defect opposite to the involved retinal segment — here, inferior retinal ischemia causes superior field loss . The process is acute, painless , and results from focal vasculitis or embolic occlusion of a branch retinal artery. Incorrect Options B) Central retinal artery occlusion: Causes complete, sudden vision loss , not sectoral (field) loss. C) Branch retinal vein occlusion: Causes sectoral hemorrhages with gradual vision loss and tortuous veins, not acute ischemia or field defect. D) Central retinal vein occlusion: Produces diffuse retinal hemorrhages (“blood and thunder” appearance) , not localized field loss.

References:

.Parsons Diseases of the Eye 23rd edition Pg.no:773,776

↑ *Back to this question in Part A*

@Neet_pg_bot

Q35: A 22-year-old patient with high myopia and thick glasses presents with floaters in his eyes. What is the likely cause of this condition? (Q840065)

A. Subarachnoid hemorrhage

B. Retinal detachment

C. Vitreous hemorrhage

D. Branched retinal artery occlusion

Correct Answer: B. Retinal detachment

Explanation: Correct Option B - Retinal detachment: Muscae volitantes, i.e. floating black opacities in front of the eyes is one of the symptoms in patients with pathological myopia. This is likely due to Rhegmatogenous retinal detachment due to the traction in posterior vitreous detachment. This causes the pigment to disperse in the vitreous cavity. This is called schafer’s sign or Tobacco dust. Incorrect Options: Option A - Subarachnoid hemorrhage: Subarachnoid hemorrhage causes subhyaloid hemorrhage. Option C - Vitreous hemorrhage: Vitreous hemorrhage is caused by blunt force trauma. Option D - Branched retinal artery occlusion: Branched retinal artery occlusion presents with cherry red spot in the retina.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q36: What could be the cause of sudden painless loss of vision? (Q860179)

A. Chemical injuries to the eyeball

B. Acute congestive glaucoma

C. Acute iridocyclitis

D. Central serous retinopathy

Correct Answer: D. Central serous retinopathy


Explanation: Central serous retinopathy is characterized by sudden painless loss of vision. The other causes of sudden painless loss of vision are as follows: Optic neuritis Methyl alcohol amblyopia Nonischaemic central retinal vein occlusion. Central Serous Chorioretinopathy (CSCR) (Option D) CSCR is characterized by the spontaneous detachment of the neurosensory retina in the macular region, with or without retinal pigment epithelium (RPE) detachment. Pathogenesis: Increased sympathetic drive and corticosteroids affect choroidal vascular permeability, raising hydrostatic pressure, and leading to RPE defects and serous detachment of the neurosensory retina. Risk Factors: Males aged 20-50 Type A personality, Emotional stress, Steroid use Systemic lupus erythematosus, Cushing's disease Pregnancy (3rd trimester) Clinical Manifestations: Sudden painless vision loss (6/9 - 6/24) Relative positive scotoma, micropsia, and metamorphopsia Recurrence in 30-50% of cases Examination Findings: Mild macular elevation with circular ring reflex RPE detachment (yellow-grey elevations) Absence or distortion of the foveal reflex Subretinal deposits Investigations: Fundus fluorescein angiography (FFA): Shows "ink-blot" or "smoke-stack" leakage patterns. {{caption_text}} {{caption_text}} Indocyanine green (ICG): Shows hyperfluorescence due to choroidal hyperpermeability. OCT: Shows neurosensory retinal and RPE detachment. {{caption_text}} {{caption_text}} Treatment: Conservative: Reassurance; spontaneous resolution in 80-90% of cases within 1-6 months. Laser Photocoagulation: For cases lasting more than 3 months, recurrent cases, and those with permanent vision loss in the other eye due to CSCR. Photodynamic Therapy (PDT): For severe, sub-foveal leaks, or chronic cases. Anti-VEGF: Considered if choroidal neovascularization (CNV) develops. Incorrect Options: Chemical injuries to the eyeball and acute congestive glaucoma (Option A and B) typically present with pain. Acute iridocyclitis (Option C) is associated with pain, photophobia, and redness, rather than painless vision loss.




Enlarging ink-blot pattern @Neet_pg_bot



Smoke-stack appearance (Mushroom shaped pattern) @Neet_pg_bot




@Neet_pg_bot



Circular ring reflex

@Neet_pg_bot



References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pg. 302, 303

↑ *Back to this question in Part A*

@Neet_pg_bot

Q37: Which of the following conditions is associated with the curtain falling sign? (Q860191)

A. Retinal detachment

B. Macular degeneration

C. Optic atrophy


D. Acute anterior uveitis


Correct Answer: A. Retinal detachment

Explanation: Curtain falling in front of the eye is seen in case of rhegmatogenous retinal detachment. Retinal detachment refers to the separation of the neurosensory retina from the retinal pigment epithelium. Etiology: Age: most common in 40-60 years. Sex. More common in males (M:F - 3:2). Myopia. About 40% cases of rhegmatogenous retinal detachment are myopic. Aphakia and pseudophakia. Trauma. It may also act as a predisposing factor. Senile posterior vitreous detachment (PVD): It is associated with retinal detachment in many cases. Retinal degenerations predisposed to retinal detachment: Lattice degeneration, Snail track degeneration,

Acquired or degenerative retinoschisis, and Focal pigment clumps. Retinal detachment can be classified into three types: Rhegmatogenous or primary retinal detachment Tractional retinal detachment Exudative retinal detachment Type of Retinal Detachment Clinical Features Rhegmatogenous Retinal Detachment Floaters (dark spots) due to rapid vitreous degeneration. Photopsia (flashes of light) caused by retinal irritation from vitreous movements. Gradual visual field loss, initially localized, progressing to total vision loss as detachment spreads. Dark veil/Curtain falling sign/cloud in front of the eye , particularly when detachment reaches the posterior region. {{caption_text}} Sudden, painless vision loss in large, central detachment. Retinal breaks may be round, horseshoe-shaped, slit-like, or large anterior dialysis. {{caption_text}} Vitreous pigments (tobacco dusting or Shaffer sign) are seen in the anterior vitreous with posterior vitreous detachment. Old retinal detachment shows retinal thinning, subretinal demarcation lines (high water marks), and secondary intraretinal cysts. Tractional Retinal Detachment Gradual, progressive visual loss Typically painless No associated flashes or floaters Vitreoretinal bands present No retinal breaks Detached area - concave and localized No shifting fluid Exudative Retinal Detachment Painless and sudden or gradual visual loss No photopsia, holes/ tears, folds and undulations. Smooth and convex with pigmentary disturbances Rounded and fixed with neovascularization on the tumor summit. Shifting fluid - changing the position of the detached area with gravity - hallmark feature. {{caption_text}}








Tractional retinal detachment associated with anteroposterior and bridging traction

@Neet_pg_bot



References:

AK Khurana Comprehensive Ophthalmology, 7th Ed, Pg. 309, 310

Kanski's Clinical Ophthalmology, 9th Edition, Pg. 681

↑ *Back to this question in Part A*

@Neet_pg_bot

Q38: A 60-year-old male patient presents to the ophthalmology clinic with complaints of visual disturbances. He reports that he is unable to see on the left side from the left eye and the left side from his right eye. However, he notices that his central vision remains intact and unaffected. The patient denies any recent trauma or other significant medical history. What is the most likely visual deficit that the patient is experiencing? (Q820443)

A. Quadrantanopia

B. Heteronymous hemianopia

C. Homonymous hemianopia

D. Homonymous hemianopia with macular sparing

Correct Answer: D. Homonymous hemianopia with macular sparing

Explanation: Correct Option D - Homonymous hemianopia with macular sparing This is the correct answer. This condition involves a loss of vision on the same side of both eyes, but with the central field of vision intact. This condition is usually associated with a stroke that affects the occipital lobe of the brain, which processes visual information. The patient's symptoms - inability to see on the left side from both eyes, but with central vision intact - align with this condition. Incorrect Options: Option A - Quadrantanopia: This is a type of visual field loss that happens in one of the four quarters of the visual field. But, the patient's symptoms indicate a loss of vision on the left side completely in both eyes, not just a quarter of the visual field. Option B - Heteronymous hemianopia: This condition involves loss of half the field of view on different sides in each eye. For example, the right half in one eye and the left half in the other. But, the patient's symptoms suggest the same side (left) of both eyes is affected. Option C - Homonymous hemianopia: This is a condition in which a patient loses half of the field of view on the same side in both eyes. This can occur because of stroke or brain injury. However, the patient's central vision remains intact, which suggests macular sparing.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q39: What is the most probable diagnosis for a 3-year-old boy who has been displaying increased clumsiness for the past 6 months, and upon examination, a cauliflower-like growth, leukocoria, and an absent red reflex in the left eye? Enucleation was performed, and microscopic examination of the specimen reveals the presence of Flexner-Wintersteiner rosettes. (Q793354)

A. Retinoblastoma

B. Neurofibroma


C. Retinopathy of prematurity

D. Toxcariasis

Correct Answer: A. Retinoblastoma

Explanation: Correct Option A - Retinoblastoma: Retinoblastoma is a malignant tumor that arises from immature retinal cells in young children, typically before the age of 5. The tumor can present with leukocoria (most commonest presentation), strabismus , or an absent red reflex on examination. Flexner-Wintersteiner rosettes is a differentiated type of tumor characterized by columnar cells with a central lumen. Incorrect Options: Option B - Neurofibroma: Neurofibromas are tumors that arise from nerve sheath cells and are often associated with the genetic disorder neurofibromatosis. In plexiform neurofibroma S-shaped deformity of the lid is observed. Option C - Retinopathy of prematurity: It may cause retinal detachment and leukocoria but there is no history of prematurity and low birth weight. Option D - Toxcariasis: It is a parasitic infection caused by roundworm

Toxocara canis that can be acquired through ingestion of contaminated soil or food. It can present with leukocoria, strabismus, floaters, eosinophilia, and hypergammaglobulinemia.



↑ *Back to this question in Part A*

@Neet_pg_bot


Q40: A 61-year-old university professor complains of vision loss in his left eye. He compares the sensation to a curtain being pulled over his affected eye. Additionally, he experienced flashing lights and floaters a week ago. The provided image shows a B-scan of his left eye. What could be the possible diagnosis for his condition? (Q793050)

A. Glaucoma

B. Posterior vitreous detachment

C. Retinal detachment

D. Corneal ulcer



Correct Answer: C. Retinal detachment

Explanation: Correct Option C - Retinal detachment: The clinical presentation of flashes of light, floaters, and curtain like effect is suggestive of rhegmatogenous retinal detachment. B scan reveals the cyst representing vitreous fluid between the neurosensory layer and retinal pigmented epithelium. Incorrect Options: Option A - Glaucoma: It is a disease that damages the optic nerve, often caused by high pressure inside the eye. It typically causes a gradual loss or sudden loss of vision but curtain like effect is not seen.. Option B - Posterior vitreous detachment Retinal : It is characterized by floaters, flashes of light, and blurred vision. Option D - Corneal ulcers: They are painful sores on the surface of the eye that can cause redness, tearing, and sensitivity to light.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [06_Neuro Ophthalmology] 013 Neuro Ophthalmology

« Back to Folder Directory

[06_Neuro Ophthalmology] 013 Neuro Ophthalmology — Quick Jump

Q1 (p. 873)	*Q2 (p. 874)*	*Q3 (p. 874)*	*Q4 (p. 875)*	*Q5 (p. 876)*	*Q6 (p. 879)*
Q7 (p. 882)	*Q8 (p. 883)*	*Q9 (p. 885)*	*Q10 (p. 886)*	*Q11 (p. 887)*	*Q12 (p. 888)*
Q13 (p. 890)	*Q14 (p. 890)*	*Q15 (p. 892)*	*Q16 (p. 894)*	*Q17 (p. 897)*	*Q18 (p. 898)*
Q19 (p. 899)	*Q20 (p. 900)*	*Q21 (p. 901)*	*Q22 (p. 902)*	*Q23 (p. 903)*	*Q24 (p. 904)*
Q25 (p. 907)	*Q26 (p. 910)*	*Q27 (p. 910)*	*Q28 (p. 911)*	*Q29 (p. 913)*	*Q30 (p. 915)*
Q31 (p. 918)	*Q32 (p. 919)*				

Q1: A 50-year-old patient presents with worsening headaches and visual disturbances. Fundoscopic examination reveals optic disc swelling. Which of the following is least likely to be associated with this condition? (Q853766)

A. Increased cup-to-disc ratio

B. Enlargement of the blind spot

C. Blurring of the optic disc margins

D. Engorged and tortuous retinal veins

Correct Answer: A. Increased cup-to-disc ratio

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Papilledema typically presents with blurring of the optic disc margins, enlargement of the blind spot, and engorged and tortuous retinal veins. As the disc swells, the physiological cup size usually decreases . An increased cup-to-disc ratio is more indicative of glaucomatous changes rather than papilledema . Papilloedema: Swelling of the optic disc due to increased intracranial pressure Characteristic Features: Passive oedema without inflammation. Swelling primarily in the optic nerve head in front of the lamina cribrosa. Engorged, tortuous small blood vessels (Option D) and swollen nerve fibres. Axoplasmic stasis with swollen, disrupted mitochondria. Retinal changes include macular star and cotton-wool spots . Signs: Initial blurring of optic disc margins. (Option C) Enlargement of the blind spot due to the separation of the retina. (Option B) Hyperaemia, swelling, and engorgement of disc vessels. Potential for haemorrhages, exudates, and macular star formation. Chronic cases may lead to optic atrophy and visual loss.

References:

Parsons Diseases of the eye, 23rd Edition, Pg - 842,843 Yanoff & Duker Ophthalmology, 5th edition, Pg - 883

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Latent nystagmus is seen in which of the below mentioned condition? (Q853821)

A. Infantile esotropia

B. Arnold-Chiari malformation

C. Internuclear ophthalmoplegia

D. Cerebellopontine angle tumours

Correct Answer: A. Infantile esotropia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Latent nystagmus is associated with infantile esotropia. With both eyes open, there is no nystagmus, but horizontal nystagmus becomes apparent on covering one eye. The fast phase is in the direction of the uncovered fixating eye. Occasionally, an element of latency may be superimposed on a manifest nystagmus so that when one eye is covered, the amplitude of nystagmus increases (manifest-latent nystagmus). Dissociated vertical deviation is another feature of Infantile esotropia. Arnold-Chiari malformation (Option B) is associated with Downbeat nystagmus (Vertical nystagmus with the fast phase beating downwards, more easily elicited in lateral gaze and downgaze). Internuclear ophthalmoplegia (Option C) is associated with Ataxic nystagmus (Horizontal jerk nystagmus that occurs in the abducting eye of a patient with an INO). Cerebellopontine angle tumours (Option D) are associated with Bruns nystagmus (Coarse cerebellar horizontal jerk nystagmus in one eye and fine high-frequency vestibular nystagmus in the other)

References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 812, 813.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 50-year-old woman presents with impaired adduction of the left eye on rightward gaze, and the abducting eye shows ataxic nystagmus. There is no noticeable ptosis, and the convergence remains intact. MRI shows a lesion in the brainstem affecting the medial longitudinal fasciculus. Which of the following is the most likely diagnosis? (Q853820)

A. Internuclear ophthalmoplegia

B. One-and-a-half syndrome

C. Oculomotor nerve palsy

D. Lateral rectus palsy

Correct Answer: A. Internuclear ophthalmoplegia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This patient's presentation, with impaired adduction of the left eye and the abducting eye shows ataxic nystagmus, along with the intact convergence and MRI findings, is consistent with internuclear ophthalmoplegia (INO). Internuclear Ophthalmoplegia (INO) Injury to the MLF between the abducens nucleus and the contralateral oculomotor nerve's medial rectus subnucleus. Impaired adduction of the ipsilateral eye, with slow or absent adduction during lateral gaze. The abducting eye overshoots the target (dysmetria), resulting in disconjugate nystagmus. Convergence is typically preserved. Bilateral INO Features Abduction saccades may be slow due to impaired inhibition of medial rectus resting tone. Associated with upward beating and torsional nystagmus. One-and-a-Half Syndrome (Option B): This syndrome results from damage to the caudal pons , which involves both the MLF and either the PPRF or abducens nucleus . It causes gaze palsy (inability to look to one side) in addition to INO . . Oculomotor nerve palsy (Option C): Would typically present with ptosis, impaired eye movements (including both vertical and horizontal), and possibly dilated pupil. Lateral rectus palsy (Option D): Lateral rectus palsy results in impaired eye abduction, horizontal diplopia, and is most often due to abducens nerve palsy.

References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 927

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 60-year-old woman presents with difficulty moving her eyes to the right. On examination, she is unable to gaze to the right, but has normal eye movements when gazing to the left, up, and down. There is no weakness in the limbs or facial muscles. Which of the following is the most likely location of the lesion? (Q853819)

A. Left abducens nucleus

B. Paramedian pontine reticular formation

C. Trochlear nerve

D. Right oculomotor nucleus

Correct Answer: B. Paramedian pontine reticular formation

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The paramedian pontine reticular formation (PPRF) is responsible for coordinating ipsilateral horizontal gaze. A lesion in the right PPRF would impair rightward horizontal gaze, while preserving other directions of eye movement, including vertical gaze. The brainstem contains gaze centers, which are the final common pathways for eye movements. The paramedian pontine reticular formation (PPRF) controls ipsilateral horizontal gaze. PPRF is managed by the frontal eye field (FEF). Left abducens nucleus (Option A): The abducens nucleus controls the lateral rectus muscle of the eye, which is responsible for moving the eye laterally. A lesion in the left abducens nucleus would impair the movement of the left eye to the left, but it wouldn't cause an issue with moving both eyes to the right. Trochlear nerve (Option C): A trochlear nerve lesion results in superior oblique palsy, causing vertical diplopia, especially on downward gaze, and leads to compensatory contralateral head tilt. Right oculomotor nucleus (Option D): The oculomotor nucleus controls the medial rectus (responsible for convergence) and the other muscles that move the eye. A lesion in the right oculomotor nucleus would result in pupil dilation (due to the parasympathetic fibers being affected) and ptosis, as well as problems with eye movement, but not isolated difficulty with horizontal gaze.

References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 923, 924

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 32-year-old woman presents with blurred vision in her right eye, photophobia and notices that her right pupil appears larger than the left. On examination, her right pupil is mid-dilated and reacts very sluggishly to light and has a sluggish and tonic reaction to accommodation. The left pupil appears normal. Which of the following is the most likely diagnosis? (Q853818)

A. Horner’s syndrome

B. Adie’s tonic pupil

C. Third nerve palsy


D. Acute angle-closure glaucoma


Correct Answer: B. Adie’s tonic pupil

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In Adie’s tonic pupil, the affected pupil is abnormally dilated (resulting in isocoria). In this condition, the pupil's response to light is absent, and its reaction to near stimuli is slow and tonic. It is caused by damage to the post-ganglionic parasympathetic pupillomotor fibers and is usually unilateral. The affected pupil constricts when treated with weak pilocarpine (0.125%) drops, while the normal pupil does not, due to the denervated iris sphincter becoming supersensitive to parasympathomimetic agents. Horner’s syndrome (Option A): This presents with a small pupil (miosis), not a large one. It is caused by a sympathetic nerve lesion and is associated with ptosis (drooping lid) and anhidrosis (loss of sweating). Third nerve palsy (Option C): While the pupil is dilated and non-reactive, a 3rd nerve palsy almost always presents with significant ptosis and an eye that is deviated "down and out" due to paralysis of the extraocular muscles. Acute angle-closure glaucoma (Option D): This is an ocular emergency. While the pupil is mid-dilated and fixed, the patient would experience excruciating pain , a rock-hard globe, and a cloudy cornea, rather than just "blurred vision and photophobia."




Ptosis
Miosis
@Neet_pg_bot

@Neet_pg_bot



References:

Yanoff & Duker Ophthalmology, 5th edition, Pg 965, 966, 968

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 30-year-old male has difficulty distinguishing between red and green colours. His occupation requires accurate colour perception. Which of the following is the most appropriate test to quantify and classify the severity of his red-green colour vision deficiency? (Q853817)

A. Ishihara colour plates

B. Farnsworth-Munsell 100-Hue Test

C. Nagel Anomaloscope

D. Lantern Test

Correct Answer: C. Nagel Anomaloscope


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Nagel Anomaloscope is the gold standard for diagnosing and quantifying red-green colour vision deficiencies . Test Procedure Evaluation Lantern Tests (Option D) Individuals identify various coloured lights emitted from a lantern. The accuracy of colour identification is evaluated. Farnsworth-Munsell 100-Hue Test (Option B) Measures ability to differentiate between subtle colour differences using hue arrangements. A higher error score indicates poorer colour vision. Ishihara Color Plates (Option A) Coloured dots form numbers or patterns designed to challenge individuals with specific deficiencies. Based on the ability to correctly identify numbers or patterns formed by dots of different colours. Anomaloscope (Nagel, Pickford-Nicolson) Split visual field: adjust red and green lights to match a yellow field. Determines severity and type of red-green colour vision deficiency through light adjustments.




@Neet_pg_bot


@Neet_pg_bot



References:

Parsons Diseases of the eye, 23rd Edition, pg - 255,256. https://www.ncbi.nlm.nih.gov/books/NBK217823/#:~:text=Other%20Tests-,Holmgren%20Wool%20Test,-Wool%20skeins

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 30-year-old male presents with difficulty in colour vision. On further examination, he has a complete inability to perceive the red end of the spectrum. Which of the following is the most likely diagnosis? (Q853816)

A. Protanopia


B. Deuteranopia

C. Tritanopia

D. Total Color Blindness

Correct Answer: A. Protanopia

Explanation: The patient has difficulty in colour vision. Specifically, the absence of perception of the red end of the spectrum is consistent with protanopia. Red-green colour blindness is the most common type of partial colour blindness and is usually congenital. It can be divided into two main types: Protanopia: Characterized by complete inability to perceive red light due to a deficiency or absence of the red cone photopigment. Deuteranopia (Option B): Involves inability to perceive green light due to a deficiency of the green cone photopigment. Protanomalous and Deuteranomalous (milder forms) Blue blindness : Occurs with ageing due to increased lens pigment. Black Cataract : Blue vision decline in individuals with sclerosing lenses.



References:

Parsons Diseases of the eye, 23rd Edition, pg - 215,216

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: On examination, a 45-year-old male was found to have repetitive, involuntary horizontal eye movements after looking at a moving striped pattern. He reports no dizziness, and his neurological examination is otherwise unremarkable. The phenomenon he is experiencing is most likely due to which of the following? (Q853815)

A. Gaze-evoked nystagmus

B. Vestibular Nystagmus

C. Optokinetic Nystagmus

D. Pendular Nystagmus

Correct Answer: C. Optokinetic Nystagmus

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Optokinetic nystagmus is a physiological response that occurs when the eyes follow a moving object, like a striped pattern, and then rapidly return to their original position. It is a normal and non-pathological condition. Optokinetic Nystagmus (OKN): Assessing Visual Acuity: Optokinetic nystagmus (OKN) is an objective method for assessing visual acuity, especially in uncooperative patients. A rotating drum with black stripes is shown to the patient, and their eyes follow the stripes with slow, smooth movements (slow phase), snapping back when the stripe disappears (fast phase) . This involuntary reflex indicates proper visual function, and adjustments in stripe width or distance can assess varying acuity levels. Diagnosing Neurological Conditions: Individuals with homonymous hemianopia (loss of vision in half of the visual field) due to parietal lobe lesions do not exhibit a normal optokinetic nystagmus response. Gaze-evoked nystagmus (Option A) occurs when the eye is held in an eccentric position, and it is typically seen in cerebellar dysfunction. Vestibular Nystagmus(Option B) is an involuntary movement of the eyes due to vestibular system disturbances, often associated with dizziness or vertigo. Pendular Nystagmus (Option D) is characterised by smooth, sinusoidal oscillations without a slow or fast phase and is usually associated with congenital conditions or visual loss.



References:

Parsons Diseases of the eye, 23rd Edition, pg - 241,943-945.

↑ *Back to this question in Part A*

Q9: A 45-year-old male presents with difficulty moving his eyes upward and convergence-retraction nystagmus triggered by an upward gaze. His pupils react poorly to light but constrict normally when focusing on near objects. Which of the following syndromes is most likely? (Q853811)

A. Parinaud syndrome

B. Weber syndrome

C. Benedict syndrome

D. Millard-Gubler syndrome

Correct Answer: A. Parinaud syndrome

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Parinaud syndrome , also known as dorsal midbrain syndrome , is a neurological condition caused by damage to the dorsal midbrain , specifically affecting the pretectal area and superior colliculus . It results in a distinct set of eye movement abnormalities and is commonly linked to lesions, tumours, or strokes affecting this area. Clinical features and signs: Upgaze Palsy (Vertical Gaze Palsy): Difficulty or inability to move their eyes upward (vertical gaze palsy). Convergence-Retraction Nystagmus: When the patient attempts to look upward, the eyes may instead converge (move inward toward each other) and retract into the orbits Pupillary Light-Near Dissociation: The pupils do not constrict to light but react normally when focusing on a near object (accommodation). Eyelid Retraction (Collier's sign): Abnormally high upper eyelids due to damage in the midbrain. Weber syndrome (Option B) is caused by a midbrain lesion involving the oculomotor nerve and corticospinal tract, leading to ipsilateral oculomotor palsy and contralateral hemiparesis. Benedict syndrome (Option C) involves a lesion in the midbrain affecting the oculomotor nerve and red nucleus, resulting in ipsilateral oculomotor nerve palsy, contralateral tremor, and ataxia. Millard-Gubler syndrome (Option D) occurs due to a lesion in the ventral pons, causing ipsilateral facial palsy and contralateral hemiparesis.



References:

Harrison’s Internal Medicine textbook 21st Ed pg- 231

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: A 50-year-old female patient with anisocoria had cocaine drops applied to both eyes. After 60 minutes, the anisocoria increased. To determine whether the lesion is preganglionic or postganglionic, what is the next step in the diagnostic process? (Q853810)

A. Reapply cocaine drops

B. 0.5% Apraclonidine

C. Adrenaline

D. 1% hydroxyamphetamine

Correct Answer: D. 1% hydroxyamphetamine

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given scenario is most likely Horner’s syndrome , and to determine the lesion's location, the next best step is to apply 1% hydroxyamphetamine drops. Algorithm for determining the site of lesion in Horner’s syndrome: Cocaine (4%): Block the reuptake of norepinephrine (noradrenaline) normally released from the nerve

endings. Hence it dilates the normal pupil (Option A) . Apraclonidine (Option B) : α2-adrenergic agonist: Miosis in the normal eye. α1-adrenergic agonist: Mydriasis in Horner's syndrome (presence of adrenergic denervation supersensitivity) Hydroxyamphetamine (1%) : releases norepinephrine from storage vesicles in the postganglionic sympathetic nerve endings at the iris dilator muscle (Option D) . Postganglionic: No dilation (3rd order nerve is dead). Preganglionic lesion: Dilation Adrenaline (Option C) or Adrenaline 1 in 1000 if used in postganglionic cases then pupils dilate due to denervation supersensitivity.

References:

Yanoff & Duker Ophthalmology, 5th edition, Pg 966, 967.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A 55-year-old male presents with a severe headache and neck stiffness that began a few days ago. His right pupil is smaller than the left, and he has drooping of the right eyelid and reduced sweating on the right side of his face. What is the most likely cause of these symptoms? (Q853809)

A. Cluster headaches

B. Pancoast tumour

C. Carotid artery dissection

D. Cavernous sinus tumor

Correct Answer: C. Carotid artery dissection

Explanation: The above case suggests features of Horner's syndrome, which is most likely due to carotid artery dissection since there are complaints of neck stiffness and severe headache and no other significant history. Horner's Syndrome Horner's syndrome is caused by a lesion of the sympathetic chain, which clinically manifests as ipsilateral miosis, facial anhidrosis, and ptosis (drooping of the eyelid). Causes: Central lesions Preganglionic Postganglionic Brainstem pathology (tumours, vascular diseases, demyelinating disorders) Spinal cord tumours Pancoast tumour Carotid and Aortic aneurysms Neck lesions Atherosclerosis of ICA (internal carotid artery)) Nasopharyngeal tumours Cavernous sinus pathology Cluster headaches Clinical manifestations: Miosis, ptosis, enophthalmos. Anhydrosis (loss of sweating): It is seen in preganglionic lesions. Absence of cilio-spinal reflex (involuntary response where the pupil of the eye dilates in response to painful stimuli applied to the skin of the neck or upper back) Acute Horner's syndrome with associated ipsilateral neck or facial pain requires urgent imaging of the neck for evaluation of a carotid dissection (Option C) . Cluster headache (Option A)) causes severe, unilateral periorbital pain with autonomic features, such as ipsilateral tearing, nasal congestion and conjunctival injection along with ptosis, miosis and other features of Horner's syndrome. Pancoast tumour (Option B) may compress the brachial plexus, leading to symptoms like localised pain around the shoulder and upper back, which might be associated with weakness or atrophy of the muscles in the arm on the affected side. Cavernous sinus tumour (Option D) causes ocular pain and diplopia along with other symptoms of Horner's syndrome, and it

is gradual in onset, unlike the above case.

References:

Parsons Diseases of the eye, 23rd Edition, Pg 276, 277. Yanoff & Duker Ophthalmology, 5th edition, Pg 965-968.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q12: A 45-year-old male reports a gradual decrease in vision in his left eye over the past two months, describing it as a "dimming" rather than a complete loss. He also has difficulty reading and seeing clearly. As per the image below, which test is most likely being conducted? (Q853808)

A. Swinging flashlight test

B. Perimetry


C. Tonometry

D. Visual acuity test



Correct Answer: A. Swinging flashlight test

Explanation: The above given scenario is suspective of optic nerve dysfunction and the image depicted above is describing RAPD (Relative Afferent Pupillary Defect) which is tested by swinging flashlight test. Swinging flashlight test: The swinging flashlight test is used to detect a relative afferent pupillary defect (RAPD) , often seen with unilateral optic nerve lesions, even when visual acuity is normal. Procedure: Shine a bright flashlight on one pupil and observe its constriction. Quickly move the light to the contralateral pupil and observe the response. Repeat the process several times, alternating between the two eyes. Normal response: Both pupils constrict equally, and the pupil receiving the light remains tightly constricted. Abnormal response (RAPD): If there is a relative afferent pathway defect, the affected pupil will dilate when the light is moved from the normal eye to the abnormal eye. Significance: The presence of Marcus Gunn pupil (RAPD) is an early indication of optic nerve disease, even when visual acuity appears normal. Causes: Optic neuritis Optic neuropathy Chiasmal compression Intraocular hemorrhage Central serous retinopathy Cystoid macular edema Retinal detachment Optic tract lesion Central or branch arterial occlusion Central or branch vein occlusion Anisometropic amblyopia Cataract Corneal scar Visual pathway damage Perimetry (Option B) measures and maps the visual field, assessing peripheral vision to detect defects or abnormalities. It aids in diagnosing eye conditions like glaucoma, retinal diseases, and neurological disorders affecting visual pathways. Tonometry (Option C) is a diagnostic test used in Glaucoma to measure intraocular pressure (IOP). The visual acuity test (Option D) measures the sharpness or clarity of a person's vision. It evaluates how well a person can see details at a specific distance, typically assessing the ability to read letters or symbols on Snellen’s chart.



References:

Parsons Diseases of the eye, 23rd Edition, Pages 278, 279.

Yanoff & Duker Ophthalmology, 5th edition, Pages 963, 964.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 48-year-old male with a history of untreated syphilis from several years ago now presents with memory loss, irritability, difficulty walking, and blurred vision. What type of pupil abnormality could be observed in this patient? (Q853807)

A. Marcus Gunn pupil

B. Amaurotic Pupil

C. Adie’s Pupil

D. Argyll Robertson Pupil

Correct Answer: D. Argyll Robertson Pupil

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given scenario of cognitive and psychotic symptoms with a history of untreated syphilis suggests Neurosyphilis, where Argyll Robertson's pupil is a clinical finding. Abnormal Pupillary Reactions Reaction Description Causes Amaurotic light reflex (Option B) Absence of direct light reflex on the affected side and an absence of consensual light reflex on the normal side. Optic and retinal lesions on the affected side Marcus Gunn pupil or RAPD (Option A) The pupils paradoxically dilate when the flashlight is swung from the unaffected to the affected eye due to the decreased light perception. It is the first sign of optic nerve disorder. Tested by a swinging flashlight test. Incomplete optic nerve lesions (Optic neuritis) and severe retinal diseases (Retinopathy, CRVO) Adie’s Pupil (Option C) Reaction to light is absent, and the near reflex is very slow and tonic. It is usually unilateral, associated with absent knee jerk and occurs more often in young women. It shows pupillary constriction to 0.125% Pilocarpine due to denervation supersensitivity. Postganglionic parasympathetic pupillomotor damage. Argyll Robertson Pupil (ARP) (Option D) Reaction to near reflex is present, but light reflex (both direct and consensual) is absent, i.e. there is light near dissociation. Neurosyphilis

References:

Parsons Diseases of the eye, 23rd Edition, Pg 280 Yanoff & Duker Ophthalmology, 5th edition, Pg 711 Kanski's Clinical Ophthalmology, 10th Edition, Page 792,795,796

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A 45-year-old man presents with difficulty focusing on near objects. Which of the following is responsible for controlling ciliary muscle contraction and pupilary constriction during accomodation? (Q853806)

A. Lateral Geniculate Nucleus

B. Edinger-Westphal Nucleus


C. Superior Colliculus

D. Visual Cortex

Correct Answer: B. Edinger-Westphal Nucleus

Listen to Audio 1

Explanation: The Edinger-Westphal Nucleus is responsible for the accommodation reflex, controlling the ciliary muscles to adjust the lens and sphincter pupillae constriction for near vision. Accommodation Reflex (Adjustments made by the eye to focus on near objects): It integrates three key responses: Accommodation: Adjusting the lens curvature for clear near vision. Convergence: Both eyes turn inward to align with the near object. Miosis: Pupils constrict to improve depth of field. The Edinger-Westphal Nucleus is the central structure crucial for the near reflex. Convergence: Signals from the medial rectus muscles travel to the Edinger-Westphal nucleus, which then influences the eye muscles to converge. Accommodation: Visual input from the cortex is relayed to the Edinger-Westphal nucleus, which adjusts the ciliary muscles for proper lens focus. Miosis: The Edinger-Westphal nucleus sends parasympathetic signals via the oculomotor nerve to the iris sphincter muscle, causing pupil constriction. Lateral Geniculate Nucleus (LGB) (Option A) is involved in visual processing but not directly in the accommodation reflex. Superior Colliculus (Option C) involves visual reflexes and eye movements but does not control lens adjustment. Visual Cortex (Option D) processes visual information but does not directly manage the accommodation reflex.



@Neet_pg_bot

References:

Parsons Diseases of the eye, 23rd Edition, Page 100

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 60-year-old patient exhibits the following pupillary responses: Light is shown in the left eye; the right pupil constricts, but the left pupil remains dilated. Light is shown in the right eye; only the right pupil constricts, while the left pupil remains dilated. Which of the following lesions is most likely responsible for this pattern of pupillary response? (Q853805)

A. Left optic nerve

B. Right Edinger-Westphal nucleus

C. Left oculomotor nerve


D. Right oculomotor nerve


Correct Answer: C. Left oculomotor nerve

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: A lesion in the left oculomotor nerve would disrupt parasympathetic signals to the left iris muscles, preventing the left pupil from constricting. This results in dilating the left pupil regardless of light stimuli , while the right eye's responses remain unaffected. Light Reflex: Direct Light Reflex: Constriction of the pupil in the same eye that is directly exposed to light. Indirect (Consensual) Light Reflex: Constriction of the pupil in the opposite eye when light is shone into one eye. Optic Nerve (Option A) Direct and consensual reactions are lost on the affected side Both are retained on the opposite side (Option D ruled out) Near reflex and lid reflex remain intact. Nuclear / Supranuclear Lesion (Option B) Ipsilateral absolute pupillary paralysis.





@Neet_pg_bot

References:

Parsons Diseases of the eye, 23rd Edition, Pg - 98

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A 62-year-old patient with papilledema has been examined. On fundoscopic examination of the eye obscuration of major blood vessels is found on the disc & vessels leaving the disc, with a significant halo around the disc. According to the Frisen grading system, what is the grade of papilledema? (Q853767)

A. Grade 5

B. Grade 2

C. Grade 3


D. Grade 4


Correct Answer: D. Grade 4


■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: In the Frisen grading system, Grade 4 is characterised by the obscuration of major vessels on the disc. Although vessels that leave the disc are also obscured, which corresponds to Grade 3, the overall presentation falls under Grade 4 . Grading of Papilledema (Modified Frisen Scale) Grade 1 : C-shaped halo nasally. Grade 2 : Circumferential halo around the optic disc. Grade 3 : Obscuration of one or more major vessels leaving the disc. Grade 4 : Obscuration of major vessels on the disc. Grade 5 : Complete obscuration of all vessels on and leaving the disc




Circumferential halo
@Neet_pg_bot


Circumferential halo + Disc Elevation
Atleast one vessel is not seen while leaving the disc
@Neet_pg_bot



References:

Parsons Diseases of the eye, 23rd Edition, Pg - 846 Yanoff & Duker Ophthalmology, 5th edition, Pg - 883, 884

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A 62-year-old patient presents with difficulty seeing objects in the right half of the visual field in both eyes. Based on this clinical presentation, which of the following is the most likely location of the lesion affecting the visual pathway? (Q853740)

A. Left optic nerve

B. Optic chiasma

C. Left optic tract


D. Right Lateral Geniculate Body

Correct Answer: C. Left optic tract

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical presentation of difficulty seeing objects in the right half of the visual field (nasal from the left field and temporal from the right field) in both eyes indicates a visual field defect known as Right Homonymous Hemianopia, which occurs due to a defect in the Left Optic Tract (lesion posterior to the optic chiasma) affecting fibers from the right visual field of both eyes. The left optic tract carries these fibers (left temporal + right nasal retina). Visual Pathway: Temporal fibers carry impulses from the Nasal field, and Nasal fibres carry the Temporal field. Left optic nerve (Option A): Optic nerve lesion → monocular vision loss. Optic chiasma (Option B) → bitemporal hemianopia. Right Lateral Geniculate Body (Option D): The lateral geniculate body is rarely isolated.



References:

Parsons Diseases of the eye, 23rd Edition, Pg - 85,87,88, 1184

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A 45-year-old patient presents with headaches and transient blurred vision. Fundoscopy reveals optic disc swelling with indistinct margins and engorged veins. What is the most accurate explanation for the pathogenesis of this condition? (Q853765)

A. Compression of the central retinal vein.

B. Increased CSF pressure causes axoplasmic stasis.

C. Inflammation of the optic nerve head.

D. Increased intraocular pressure.

Correct Answer: B. Increased CSF pressure causes axoplasmic stasis.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical presentation suggests Papilledema, a hydrostatic, non-inflammatory condition characterised by swelling of the optic disc due to increased intracranial pressure. Typically bilateral, it occurs because the optic nerve, encased in the meningeal sheaths, is affected by changes in cerebrospinal fluid (CSF) pressure. Pathogenesis: Initially thought to be caused by compression of the central retinal vein, leading to its collapse while the artery remains intact. (Option A ruled out) Current Understanding: Blockage of axoplasmic transport in the optic nerve head due to elevated CSF pressure results in optic disc swelling and vascular changes on the disc surface. Aetiology: Intracranial tumors Brain abscesses, cavernous sinus thrombosis, aneurysms, subarachnoid haemorrhage. Pseudotumour cerebri Malignant hypertension is typically accompanied by increased intracranial pressure. Decrease in intraocular pressure (Option D ruled out)

References:

Parsons Diseases of the Eye, 23rd Edition, Page 839,841,842

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: A 35-year-old patient with multiple sclerosis (MS) notices that her vision becomes temporarily blurred whenever she exercises. Which of the following phenomena is most likely explained by this temporary worsening of neurological symptoms due to increased body temperature? (Q853754)

A. Riddoch's Phenomenon

B. Uhthoff's Phenomenon

C. Pulfrich's Phenomenon

D. Lhermitte sign

Correct Answer: B. Uhthoff's Phenomenon

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Uhthoff's Phenomenon refers to the temporary worsening of neurological symptoms in individuals with multiple sclerosis (MS) due to increased body temperature. Symptoms: Blurring of vision during activities like hot showers or exercise. Increased muscle weakness during a fever. Worsening of balance or coordination after exposure to heat. Riddoch's Phenomenon (Option A) refers to a visual disturbance where a patient can perceive movement but not static objects, typically resulting from damage to the occipital lobe. Pulfrich's Phenomenon (Option C) is a visual anomaly where objects moving in a straight line appear to follow a curved trajectory due to delayed visual processing. It is often associated with optic neuritis but is not related to heat or temperature changes. Lhermitte sign (Option D) is a transient electric shock-like sensation that travels down the spine and often into the limbs when the neck is flexed forward . It is seen in MS.

References:

Harrison’s Internal Medicine textbook, 21st Edition, Page 3463

https://www.ncbi.nlm.nih.gov/books/NBK470244/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: A 25-year-old man presents with a sudden onset of painless, bilateral vision loss that started in the right eye and later affected the left eye. Genetic testing reveals a mutation in the mitochondrial DNA. Which of the following is the most likely diagnosis? (Q853753)

A. Kjer dominant Optic Neuropathy

B. Leber Hereditary Optic Neuropathy (LHON)

C. Wolfram Syndrome

D. Behr syndrome

Correct Answer: B. Leber Hereditary Optic Neuropathy (LHON)

Explanation: Leber Hereditary Optic Neuropathy (LHON) is a mitochondrial genetic disorder that typically presents with subacute, bilateral vision loss, often starting in one eye and affecting the other eye within a few weeks, and it is associated with mutations in mitochondrial DNA. Leber Hereditary Optic Neuropathy Inheritance Mitochondrial DNA disorder; Maternally inherited. Variable Penetrance 50%–70% of males and 10%–15% of females Age of Onset Typically between 15-30 years, later onset is possible, especially in females (40s-50s). Vision Loss Sudden, painless, bilateral, central vision loss, central scotoma common. Kjer dominant Optic Neuropathy (Option A) is an autosomal dominant condition with gradual vision loss beginning in childhood, not sudden onset in young adulthood. It typically does not present with sudden, painless vision loss or significant optic

disc pallor, and visual impairment usually does not progress to severe loss. Wolfram syndrome (Option C): A rare genetic disorder that includes optic atrophy along with diabetes insipidus, diabetes mellitus, and hearing loss . It is inherited in an autosomal recessive pattern. Behr Syndrome (Option D) is a rare autosomal recessive type of spinocerebellar ataxia that presents with Optic Atrophy, Cerebellar Ataxia, and Spastic Paraplegia . Symptoms usually begin in childhood or early adulthood.

References:

Parsons Diseases of the eye, 23rd Edition, Pg 882,883

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: A 45-year-old man presents with sudden, progressive bilateral blurring of vision for two days, accompanied by headache, nausea, and vomiting. He has a history of alcohol dependence but denies taking any new medications. Examination shows bilateral central scotomas and hyperemic optic discs with mild retinal edema. His pupils are sluggishly reactive to light. Arterial blood gas analysis reveals a high anion gap metabolic acidosis.Which of the following toxins is most likely responsible for his condition? (Q853752)

A. Methanol

B. Cyanide

C. Arsenic

D. Ethambutol



Correct Answer: A. Methanol

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The acute onset , bilateral central scotoma , optic disc hyperemia → pallor , and metabolic acidosis strongly indicate methanol poisoning . Methanol is metabolized by alcohol dehydrogenase to formic acid , which accumulates and causes retinal and optic nerve toxicity . Methanol causes a form of toxic optic neuritis, specifically affecting the papillomacular bundle and resulting in bilateral optic disc swelling in the acute stage, followed by optic atrophy later on. The classic triad of systemic symptoms occurring in the course of methanol poisoning includes CNS depression, metabolic disorders manifested mainly as systemic metabolic acidosis, and visual disturbances. Option Reasoning Cyanide (Option B) Causes chronic tobacco–alcohol amblyopia; Chronic: optic disc atrophy, retrobulbar visual tract lesions Arsenic (Option C) Cataracts, Bilateral symmetric optic neuropathy, Retinal pathologies like ARMD (due to chronic exposure and accumulation) Ethambutol (Option D) Causes optic neuritis resulting in loss of red-green colour discrimination, decreased visual acuity, and visual field defects (cecocentral scotoma)

References:

Parsons Diseases of the Eye, 23rd Edition, Page 874

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: A 72-year-old patient presents with sudden vision loss in one eye, severe headache, scalp tenderness, and jaw claudication. Fundoscopic examination reveals a pale, swollen optic disc with haemorrhages. Laboratory tests show elevated ESR and CRP levels. What is the diagnosis and the appropriate management? (Q853751)

A. Arteritic AION; Lifestyle modifications

B. Non-arteritic AION; Lifestyle modifications

C. Arteritic AION; High dose corticosteroids


D. Non-arteritic AION; High dose corticosteroids

Correct Answer: C. Arteritic AION; High dose corticosteroids

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Arteritic AION is often related to Giant Cell Arteritis (GCA) and presents with systemic symptoms such as severe headache, scalp tenderness, and jaw claudication . It requires prompt treatment with high-dose corticosteroids to manage inflammation and prevent further vision loss. Maintenance with long-term low-dose corticosteroids



References:

Parsons Diseases of the eye, 23rd Edition, pg 851-855

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: A 48-year-old woman presents with a history of decreased vision in her right eye. On examining the right eye, there is RAPD, pain on eye elevation, and the fundus is normal. What is the most appropriate management? (Q853750)

A. Conservative management

B. High-dose Corticosteroids

C. Oral antibiotics

D. Surgical intervention

Correct Answer: B. High-dose Corticosteroids

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient's symptoms and clinical findings are consistent with acute retrobulbar optic neuritis , commonly associated with demyelination in multiple sclerosis. While spontaneous recovery can occur , IV methylprednisolone is recommended to reduce inflammation, shorten symptom duration, and decrease the risk of future attacks, especially in high-risk patients. Management of Optic neuritis Treatment Indications Details Conservative management (Observation and Monitoring) Mild symptoms Known history of similar episodes Many cases improve spontaneously. Suitable for mild symptoms but not recommended for severe cases or first episodes. High-Dose Corticosteroids Severe vision loss No history of previous episodes MRI evidence of demyelination Intravenous methylprednisolone (1 g daily for 3 days) followed by oral prednisolone taper as established by the ONTT trial (Optic Neuritis Treatment Trial). It accelerates visual recovery but does not prevent future multiple sclerosis episodes. Patients with optic neuritis when related to MS can also be given Glatiramer acetate and Beta Interferon 1a.

References:

Parsons Diseases of the eye, 23rd Edition, Pg 865

↑ *Back to this question in Part A*

@Neet_pg_bot


Q24: A 40-year-old woman presents with a 3-week history of difficulty in vision in her left eye, along with persistent headaches and occasional eye pain. She also reports trouble with colour perception. Fundoscopic examination reveals the findings shown in the image. Which type of optic neuritis is most likely to be seen? (Q853749)

A. Papillitis

B. Retrobulbar neuritis

C. Neuroretinitis

D. Papilledema




Correct Answer: C. Neuroretinitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical scenario with a macular star in fundoscopy suggests Neuroretinitis , a subtype of optic neuritis. Neuroretinitis: A form of optic neuropathy characterized by optic disc edema + macular star (hard exudates in radial pattern). Etiology: Commonly infectious → especially Bartonella henselae (cat scratch disease); also syphilis, TB, viral causes. Pathophysiology: Leakage from disc capillaries → fluid tracks along outer plexiform layer → forms macular star . Clinical features: Subacute vision loss Mild eye pain Defective color vision Relative afferent pupillary defect (RAPD) may be present Fundus findings: Optic disc swelling (early) Macular star appears later (1-2 weeks) Prognosis & treatment: Generally good visual recovery Treat underlying cause (e.g., antibiotics for Bartonella) Optic neuritis is an inflammatory condition of the optic nerve that can cause vision loss, with symptoms including decreased vision, pain in eye movement, and visual field defects. Classification of Optic neuritis: Papillitis (Option A) Retrobulbar Neuritis (Option B) Neuroretinitis (Option C) Site Optic disc Optic nerve behind the eyeball. Optic disc and surrounding retina in the macular area. Characteristic features Fundoscopy reveals a blurred disc margin due to leakage and increased capillary permeability, edematous and hyperaemic disc, and physiological cup is obliterated. Pain on eye elevation RAPD (relative afferent pupillary defect) is observed. Fundoscopic examination typically reveals normal optic disc , at least initially, which is why the classic phrase, “ the patient sees nothing, and the doctor sees nothing ” is often used to describe retrobulbar neuritis. A macular star is seen in fundoscopy (accumulation of lipids).





References:

Parsons Diseases of the eye, 23rd Edition, Pages 856-864.

Kanski’s Clinical Ophthalmology, 9th edition, Pages 754, 770

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: A 34-year-old woman has a 6-month history of worsening vision and difficulty with detailed tasks, along with occasional eye pain in her left eye. Fundoscopy shows a chalky white optic disc with normal vessels in her left eye. What is the most common cause of these findings? (Q853748)

A. Papilledema

B. Multiple sclerosis

C. Retinitis pigmentosa

D. Glaucoma

Correct Answer: B. Multiple sclerosis


■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: The given scenario suggests Primary optic atrophy, with multiple sclerosis being the most common cause. Optic Atrophy: It is the degeneration of the optic nerve, which occurs as an end result of any pathologic process that damages axons in the anterior visual system. Classification of Optic atrophy: Primary Secondary Consecutive Glaucomatous Etiology Multiple sclerosis (Option B) Retrobulbar neuritis Leber's hereditary optic atrophies Intracranial tumours (e.g. pituitary tumours) Traumatic severance or avulsion of the optic nerve Toxic amblyopia (chronic retrobulbar neuritis) Tabes dorsalis. Long-standing optic nerve disease. Papilledema (Option A) Diffuse chorioretinitis Retinal pigmentary dystrophies (e.g., retinitis pigmentosa (Option C) , pathological myopia and occlusion of the central retinal artery) Glaucoma (Option D) Fundoscopy Chalky white Clear disc margin Lamina cribrosa well seen Normal vessels. Dirty grey Clear disc margin Lamina cribrosa obscured Peripapillary sheathing over vessels Hyaline bodies on the retina. Waxy pallor Blurred disc margin Lamina cribrosa well seen Attenuated vessels Characterized by deep and wide cupping of the optic disc and nasal shift of the blood vessels.


@Neet_pg_bot


@Neet_pg_bot





References:

Parsons Diseases of the eye, 23rd Edition, Page 867-870 Myron Yanoff & Duker Ophthalmology, 6th Edition, Page 808,809

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: A 45-year-old female reports a gradual decline in vision in her right eye over the past three months, accompanied by blurry vision and impaired colour perception in that eye. Upon examination, the ophthalmologist suspected optic nerve damage and advised further tests. Which of the following manifestations does not indicate optic nerve dysfunction? (Q853747)

A. Positive scotoma

B. Dyschromatopsia

C. Central scotoma

D. Decrease in contrast sensitivity

Correct Answer: A. Positive scotoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: A positive scotoma is a visual field defect that is perceived by the patient as something seen abnormally like a dark patch and is commonly associated with macular lesions. Optic nerve dysfunction is usually associated with negative scotoma. Optic nerve damage presents with dyschromatopsia, central scotoma, and contrast sensitivity decreases.

References:

Parsons Diseases of the eye, 23rd Edition, Pg 836, 1336, Yanoff & Duker Ophthalmology, 5th edition, Pg 963, 964.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: Ophthalmic artery is located Inferiolateral to which part of the optic nerve?? (Q853746)

A. Intraocular part


B. Intraorbital part

C. Intracanalicular part

D. Intracranial part

Correct Answer: C. Intracanalicular part

Explanation: The intracanalicular part of the optic nerve is the segment located within the optic canal. The ophthalmic artery enters the optic canal alongside the optic nerve and lies inferiolateral to the optic nerve. Parts of the optic nerve: Part of the optic nerve Description Intraocular (Option A) It starts from the optic disc, pierces the choroid and sclera and becomes continuous with the intraorbital part at the back of the eyeball. Intra-orbital (Option B) Extends from the back of the eyeball to the optic foramina and runs parallel to the ophthalmic artery. Intra-canalicular (Option C) It is closely related to the ophthalmic artery, which lies inferolateral to optic nerve. It crosses obliquely over ophthalmic artery as it enters the orbit to lie on its medial side. Intracranial (Option D) It lies above the cavernous sinus and converges with its fellow to form the optic chiasma. optic nerve" data-author="NA" data-hash="4201" data-license="NA" data-source="NA" data-tags="March2025" height="500" src="" width="500" />



References:

Parsons Diseases of the eye, 23rd Edition, Pg 42, 43. Kanski's Clinical Ophthalmology, 10th Edition, Page 764

↑ *Back to this question in Part A*

@Neet_pg_bot

Q28: Which of the following arteries contributes to the formation of the Circle of Zinn-Haller? (Q853745)

A. Central retinal artery

B. Ophthalmic artery

C. Short posterior ciliary arteries


D. All of the above

Correct Answer: C. Short posterior ciliary arteries

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Circle of Zinn or Circle of Zinn-Haller is primarily formed by the short posterior ciliary arteries, which supply the optic nerve head. Blood supply of optic nerve: Circle of Zinn (also called Circle of Zinn-Haller) : Formed by anastomosis of the short posterior ciliary artery. Central retinal artery (branch of ophthalmic artery)



References:

Parsons Diseases of the eye, 23rd Edition, Page 42,43 Kanski's Clinical Ophthalmology, 10th Edition, Page 763

↑ *Back to this question in Part A*

Q29: A 62-year-old patient presents with a visual field defect following a stroke. Neuroimaging reveals a lesion in the occipital lobe related to the territory of the posterior cerebral artery affecting the calcarine cortex. Which of the following visual field characteristics is most likely associated with this type of lesion? (Q853744)

A. Congruous Homonymous Hemianopia with Macular Sparing

B. Congruous Homonymous Hemianopia without Macular Sparing

C. Incongruous homonymous hemianopia with Macular sparing


D. Incongruous homonymous hemianopia without Macular sparing

Correct Answer: A. Congruous Homonymous Hemianopia with Macular Sparing


■ *Listen (English)*

■ *Listen (Hindi)*


Explanation: A lesion in the occipital lobe, specifically in the Posterior Cerebral Artery (PCA) territory affecting the calcarine cortex, typically causes congruous homonymous hemianopia with macular sparing. Lesions in Visual Cortex: Typically present as congruous homonymous hemianopia (visual field loss is consistent in both eyes). Blood supply of Visual Cortex: The primary visual cortex, mainly supplied by the posterior cerebral artery (PCA), processes visual information, and a PCA stroke can lead to homonymous hemianopia. Macular Area : Located at the posterior tip of the visual cortex and is crucial for detailed central vision. Its dual blood supply (from both PCA and MCA) helps to protect it from complete loss even when there is an isolated PCA lesion. (Macular Sparing) . Injury to the tip of the visual cortex leads to Macular Homonymous Hemianopia .



@Neet_pg_bot



@Neet_pg_bot



References:

Parsons Diseases of the eye, 23rd Edition, Page 1213

Yanoff & Duker Ophthalmology, 5th Edition, Page 919

↑ *Back to this question in Part A*

@Neet_pg_bot

Q30: A 58-year-old patient presents with visual disturbances. Neuroimaging reveals a lesion in the temporal lobe affecting the inferior optic radiations. Which visual field defects are most likely associated with this finding? (Q853743)

A. Inferior homonymous quadrantanopia

B. Superior homonymous quadrantanopia

C. Central scotoma

D. Homonymous hemianopia


Correct Answer: B. Superior homonymous quadrantanopia


■ *Listen (English)*


■ *Listen (Hindi)*


Explanation: A lesion affecting the inferior optic radiations, particularly in Meyer's loop within the temporal lobe, typically results in superior homonymous quadrantanopia. Aspect Inferior Optic Radiations (Meyer's Loop) Superior Optic Radiations (Baum's Loop) Location of Lesion Temporal lobe Parietal lobe Affected Visual Field Superior quadrants of the contralateral visual field of both eyes Inferior quadrants of the contralateral visual field of both eyes. Visual Defect Superior homonymous quadrantanopia Inferior homonymous quadrantanopia (Option A) Central scotoma (Option C) is a defect in central vision that is typically associated with lesions in the retina, optic nerve, and macula rather than the specific pathways in the temporal lobe. Homonymous hemianopia (Option


D) occurs in lesions involving the optic tract, lateral geniculate body (LGB), or both superior and inferior optic radiations.






Visual Field
Left
Right
@Neet_pg_bot


Blind spot
Central Scotoma
@Neet_pg_bot



References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 919

↑ *Back to this question in Part A*

@Neet_pg_bot

Q31: A 58-year-old patient presents with right-sided visual field loss in both eyes, resulting in a homonymous hemianopic defect. Neuroimaging and clinical examination suggest a lesion affecting the visual pathway. Based on the above findings, which anatomical location is most likely implicated in this patient's condition? (Q853742)

A. Right optic tract

B. Left optic tract

C. Right optic nerve


D. Optic chiasma

Correct Answer: B. Left optic tract

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: A lesion in the left optic tract would cause right (contralateral) homonymous hemianopia, where the right visual field is lost in both eyes. Location of the lesion Characteristic features Optic nerve Marked loss of vision or complete blindness on the affected side associated. Near (accommodation) reflex is present. Optic chiasma Bitemporal hemianopia Optic tract Contralateral homonymous hemianopia Contralateral hemianopic pupillary responses, i.e. Wernicke's reaction. Optic radiations Superior quadrantic hemianopia (pie in the sky) Produced when the inferior fibres (Meyer's loop) are involved in temporal lobe lesions) Inferior quadrantic hemianopia (pie on the floor) Produced when superior fibres (Baum's loop) are involved in parietal lobe lesions. Visual cortex Homonymous hemianopia, often with macular sparing



References:

Parsons Diseases of the eye, 23rd Edition, Page 1187

Yanoff & Duker Ophthalmology, 5th edition, Pages 918, 919

↑ *Back to this question in Part A*

@Neet_pg_bot


Q32: A 64-year-old patient presents with difficulty seeing objects. MRI findings reveal a pituitary adenoma causing compression at the optic chiasma. Which visual field


defect is most likely associated with this condition? (Q853741)


Correct Answer: Unknown

- *Listen (English)*
- *Listen (Hindi)*

Explanation: A pituitary adenoma compressing the optic chiasm typically results in Bitemporal hemianopia due to the involvement of crossing nasal fibres of the optic nerves, causing loss of vision in the outer portions of both visual fields. Visual Pathway Lesions and Defects Site of Lesion Fibers Affected Field Defect Optic Tract Lateral Geniculate Body (LGB Lesion) Optic Radiation Ipsilateral Temporal Fibers + Contralateral Nasal Fibers Homonymous Hemianopia (Option C) Visual Cortex without the involvement of tip (Post. Cerebral Artery) Ipsilateral Temporal Fibers + Contralateral Nasal Fibers except macular representation Homonymous Hemianopia with Macular Sparing (Option D) Pie in the Sky (Option B) (Superior Homonymous Quadrantanopia) occurs in Lesion of Inferior Optic Radiation.







References:

Parsons Diseases of the eye, 23rd Edition, Pg - 1184 Yanoff & Duker Ophthalmology, 5th edition, Pg - 913,919

↑ *Back to this question in Part A*

@Neet_pg_bot

• [06_Neuro Ophthalmology] 014 Previous Year Questions

« Back to Folder Directory

[06_Neuro Ophthalmology] 014 Previous Year Questions — Quick Jump

Q1 (p. 923)	*Q2 (p. 925)*	*Q3 (p. 931)*	*Q4 (p. 933)*	*Q5 (p. 935)*	*Q6 (p. 936)*
Q7 (p. 937)	*Q8 (p. 938)*	*Q9 (p. 941)*	*Q10 (p. 943)*	*Q11 (p. 944)*	*Q12 (p. 950)*
Q13 (p. 952)	*Q14 (p. 956)*	*Q15 (p. 958)*	*Q16 (p. 959)*	*Q17 (p. 960)*	*Q18 (p. 961)*
Q19 (p. 962)	*Q20 (p. 964)*	*Q21 (p. 965)*	*Q22 (p. 967)*	*Q23 (p. 968)*	*Q24 (p. 968)*
Q25 (p. 970)	*Q26 (p. 971)*	*Q27 (p. 972)*	*Q28 (p. 974)*	*Q29 (p. 974)*	*Q30 (p. 976)*
Q31 (p. 977)	*Q32 (p. 980)*	*Q33 (p. 980)*	*Q34 (p. 981)*	*Q35 (p. 982)*	*Q36 (p. 982)*
Q37 (p. 984)	*Q38 (p. 989)*	*Q39 (p. 991)*	*Q40 (p. 991)*		


Q1: What is the probable diagnosis for a middle-aged man with diabetes, who has reported a decline in vision and whose fluorescein angiography displays the depicted image? {{caption_text}} (Q857807)

A. Mild non proliferative diabetic retinopathy

B. Moderate non proliferative diabetic retinopathy


C. Severe non proliferative diabetic retinopathy

D. Proliferative diabetic retinopathy



Correct Answer: D. Proliferative diabetic retinopathy

Explanation: In the above given image of FFA (fundus fluorescein angiography), neovascularization is noted which is classical of Proliferative Diabetic Retinopathy (PDR) . {{caption_text}} Abbreviated Early Treatment Diabetic Retinopathy Study (ETDRS) classification of diabetic retinopathy: Category Management Non-Proliferative Diabetic Retinopathy (NPDR): No Diabetic Retinopathy Review in 12 months Very Mild NPDR: Microaneurysms only Review most patients in 12 months (Option D) Mild NPDR: (Option A) Any or all of the: microaneurysms, retinal hemorrhages, exudates, and cotton-wool spots, up to the level of moderate NPDR. No intraretinal microvascular anomalies (IRMA) or significant beading. Review range 6-12 months, depending on the severity of signs, stability, systemic factors, and patient's circumstances Moderate NPDR: (Option B) Severe retinal hemorrhages in 1-3 quadrants or mild IRMA Significant venous beading can be present in no more than 1 quadrant Cotton wool spots commonly present Review in approximately 6 months Severe NPDR: (Option C) The 4-2-1 rule; is one or more of the: Severe hemorrhages in all 4 quadrants Significant venous beading in 2 or more quadrants Moderate IRMA in 1 or more quadrants Review in 4 months Very severe NPDR: Two or more of the criteria for severe NPDR Review in 2-3 months Proliferative Diabetic Retinopathy (PDR): (Option D) Mild-moderate PDR: New vessels on the disc (NVD) or new vessels elsewhere (NVE) , but extent insufficient to meet the high-risk criteria Treatment is considered according to the severity of signs, stability, systemic factors, and the patient's circumstances such as reliability of attendance for review. If not treated, review in up to 2 months High-risk PDR: NVD greater than ■ disc area Any NVD with vitreous hemorrhage NVE greater than ½ disc area with vitreous hemorrhage Treatment should be performed immediately when possible and certainly on the same day if symptomatic presentation with a good retinal view Advanced diabetic eye disease: It is characterized by tractional retinal detachment. Significant persistent vitreous hemorrhage and neovascular glaucoma. Vitreoretinal surgery with photocoagulation



References:

Parsons Diseases of the Eye 23rdEdition, pages 292,293.

Yanoff and Duker Ophthalmology 5thEdition, pages 548-551.

Kanski's Clinical Ophthalmology 9th Edition, pages 497,498.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A patient with pituitary adenoma compressing the optic chiasma now presents with loss of visual field. What visual field defect will be seen in the patient? (Q866833)

A. Bitemporal hemianopia


B. Homonymous anopia

C. Homonymous hemianopia


D. Superior quadrantanopia


Correct Answer: A. Bitemporal hemianopia

Explanation: The optic chiasm contains crossing fibers from the nasal retina of both eyes. Pituitary adenoma commonly compresses the central chiasm from below → affects crossing nasal fibers → loss of temporal visual fields in both eyes → Bitemporal hemianopia Clinical scenario: Gradual, progressive peripheral vision loss, more noticeable while driving or walking. Mechanism: Nasal retinal fibers detect temporal fields → damage leads to bitemporal hemianopia. Visual Pathway Lesions and Defects Site of Lesion Fibers Affected Field Defect Optic Nerve Ipsilateral Temporal + Nasal Fibers Optic Chiasma Bilateral Nasal Fibers Optic Tract Lateral Geniculate Body (LGB Lesion) Optic Radiation Ipsilateral Temporal Fibers + Contralateral Nasal Fibers Visual Cortex without the involvement of tip (Post. Cerebral Artery) Ipsilateral Temporal Fibers + Contralateral Nasal Fibers except macular representation Tip of Visual Cortex (Middle Cerebral Artery) Macular Representation of Ipsilateral Temporal and Contralateral Nasal Fibers. Aspect Inferior Optic Radiations (Meyer's Loop) Superior Optic Radiations (Baum's Loop) Location of Lesion Temporal lobe Parietal lobe Affected Visual Field Superior quadrants of the contralateral visual field of both eyes Inferior quadrants of the contralateral visual field of both eyes. Visual Defect Superior homonymous quadrantanopia (Option D ruled out) Inferior homonymous quadrantanopia Associated Symptoms Memory loss Hearing loss Auditory hallucinations Contralateral hemiplegia Hemianesthesia Optokinetic nystagmus Homonymous anopia (Option B): Complete blindness in corresponding halves of the visual fields of both eyes (e.g., entire right field loss) - occurs with complete optic tract lesion , not chiasmal compression.




Visual Field Defects






@Neet_pg_bot



Anopia


@Neet_pg_bot



Bitemporal hemianopia
(Heteronymous hemianopia)

@Neet_pg_bot

Left eye visual field Right eye visual field



Homonymous hemianopia
(Option C ruled out)

@Neet_pg_bot


Visual Field



Left Right


Homonymous hemianopia with Macular Sparing


@Neet_pg_bot



Macular homonymous hemianopia

@Neet_pg_bot





References:

.

↑ *Back to this question in Part A*

Q3: Which muscle serves as the yoke muscle for the right lateral rectus? (Q858157)

A. Right superior oblique


B. Left medial rectus


C. Left lateral rectus

D. Right medial rectus

Correct Answer: B. Left medial rectus

Explanation: qual Innervation would mean that the right lateral rectus (yoke muscle) should be balanced with the left medial rectus (vs Sherrington's law: This law about reciprocal innervation states that increased innervation and contraction of a muscle is automatically associated with reciprocal decrease in innervation and contraction of its antagonist). The yoke muscle pairs facilitate various eye movements: Horizontal Gaze (Left or Right) : Leftward Gaze (Levoversion) : Left Lateral Rectus (abducts the left eye) Right Medial Rectus (adducts the right eye) Rightward Gaze (Dextroversion) : Right Lateral Rectus (abducts the right eye) Left Medial Rectus (adducts the left eye) Vertical Gaze (Up or Down) : Upward Gaze (Elevation) : Right and Left Superior Rectus (elevate the eyes) Right and Left Inferior Oblique (assist in elevation) Downward Gaze (Depression) : Right and Left Inferior Rectus (depress the eyes) Right and Left Superior Oblique (assist in depression) Inward Gaze (Convergence) : Both Medial Rectus muscles contract to bring the eyes inward (toward the nose) during near-vision tasks, such as reading. Right superior oblique (Option A) : Helps with vertical and torsional eye movement, not horizontal. Left lateral rectus (Option C) : Moves the left eye outward, opposite to what's needed for the right lateral rectus. Right medial rectus (Option D) : Moves the right eye inward, opposite of what is needed for the right lateral rectus.





References:

Yanoff & Duker Ophthalmology, 5th edition, Pg- 1193,1194

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: Which of the following statements accurately describes optic neuritis? (Q855579)

A. Relative afferent pupillary defect is seen

B. There is no pain on ocular movement


C. Multiple myeloma is a common cause

D. Oral corticosteroids are the mainstay of treatment

Correct Answer: A. Relative afferent pupillary defect is seen

Explanation: Optic neuritis is an inflammation of the optic nerve, and one of its characteristic features is the relative afferent pupillary defect (RAPD) . RAPD is observed when a light is shone in the affected eye, and the pupil does not constrict as much as it should. This is due to the impaired function of the optic nerve in transmitting the light reflex to the brain. Classification of Optic neuritis Aspect Papillitis Retrobulbar Neuritis Neuroretinitis Site Optic disc Optic nerve behind the eyeball. Optic disc and surrounding retina in the macular area. Characteristic features Fundoscopy reveals a blurred disc margin due to leakage and increased capillary permeability, edematous and hyperaemic disc, and physiological cup is obliterated. {{caption_text}} Pain on eye elevation RAPD (relative afferent pupillary defect) is observed. Fundoscopic examination typically reveals normal optic disc , at least initially, which is why the classic phrase, “ the patient sees nothing, and the doctor sees nothing ” is often used to describe retrobulbar neuritis. A macular star is seen in fundoscopy (accumulation of lipids). {{caption_text}} Clinical features, It may be asymptomatic or associated with, Visual loss: Monocular sudden, progressive and profound visual loss is the hallmark of acute optic neuritis. Dark adaptation may be lowered. Visual obscuration in bright light is a typical symptom of acute optic neuritis. Impairment of color vision is always present in optic neuritis. Uhthoff's symptom: Episodic transient obscuration of vision on exertion and on exposure to heat, which recovers on resting or moving away from the heat, occurs in patients with isolated optic neuritis. Pulfrich's phenomenon: Depth perception may be impaired, particularly for the moving object. Visual acuity is usually reduced markedly. Visual field changes: The most common field defect in optic neuritis is a relative central or centrocecal scotoma. Contrast sensitivity is impaired. There is no pain on ocular movement (Option B) : Pain on ocular movement is one of the hallmark symptoms of optic neuritis . It often precedes or accompanies vision loss, making it a distinguishing feature of the condition. Multiple myeloma is a common cause (Option C) : Multiple myeloma is a rare cause of optic neuritis. Most cases of optic neuritis are idiopathic or related to multiple sclerosis (MS) . Other systemic conditions like infectious or autoimmune diseases may also be involved, but multiple myeloma is not typically associated with optic neuritis. Oral corticosteroids are the mainstay of treatment (Option D) : Oral corticosteroids are not the first-line treatment for optic neuritis. Intravenous corticosteroids (IV methylprednisolone) are more commonly used in acute optic neuritis, particularly to speed up recovery and reduce the risk of recurrence in conditions like multiple sclerosis





References:

Parsons Diseases of the eye, 23rd Edition, Page 856-864.

Kanski’s Clinical Ophthalmology, 9th Edition, Page 754, 770

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: Which structures are affected in a pregnant woman who exhibits fever, retroorbital pain, headache, pulsatile proptosis of the right eye, and tinnitus, while having normal blood pressure and fundus? 1. Abducens nerve 2. Oculomotor nerve 3. Internal carotid artery 4. Cavernous sinus (Q855223)

A. A, B, C

B. B, C ,D

C. A, C ,D

D. A, B, C, D

Correct Answer: D. A, B, C, D

Explanation: The symptoms described in this pregnant woman, such as fever, retroorbital pain, headache, pulsatile proptosis, and tinnitus, suggest a possible case of cavernous sinus thrombosis. Abducens nerve : The abducens nerve (cranial nerve VI) runs through the cavernous sinus. It is responsible for controlling the lateral rectus muscle, which abducts the eye. Compression or dysfunction can cause pulsatile proptosis and eye movement abnormalities . Oculomotor nerve : The oculomotor nerve (cranial nerve III) also passes through the cavernous sinus and controls most of the eye's movements, including the muscles responsible for pupil constriction. Dysfunction here can lead to ptosis and pupil abnormalities . Internal carotid artery : The internal carotid artery runs through the cavernous sinus and can be affected in cases of sinus thrombosis or vascular pathology. Pulsatile proptosis and tinnitus are often signs of vascular involvement (e.g., arteriovenous fistula or carotid-cavernous fistula). Cavernous sinus : The cavernous sinus is a critical structure that houses several cranial nerves (III, IV, V1, V2, and VI) and the internal carotid artery. Thrombosis or infection here can lead to a constellation of symptoms involving nerve dysfunction and vascular compromise. Given the symptoms and the structures involved, the correct answer is Option D: A, B, C, D , since all these structures (abducens nerve, oculomotor nerve, internal carotid artery, and cavernous sinus) are likely affected in this scenario. Cavernous sinus Thrombosis: Typically results from infections such as sinusitis, orbital cellulitis, or otitis. Symptoms: Severe headache Malaise Nausea and vomiting Unilateral or bilateral proptosis Chemosis Congestion of facial, conjunctival, and retinal veins Reduced vision Cranial nerve dysfunction (third to sixth cranial nerves) Diagnosis: Imaging, especially MRI and MRI venography Systemic infection investigations, including lumbar puncture Treatment: Intravenous antibiotics Possible surgical drainage Prognosis: High mortality rate (20% with treatment, up to 100% untreated) Options A, B, & C are incorrect.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 129, 130

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Which cranial nerves does the oculocephalic reflex evaluate? A. III B. IV C. VI D. VIII E. VII (Q855237)

A. A and C


B. A and E

C. A and D

D. A, C and D

Correct Answer: D. A, C and D

Explanation: The oculocephalic reflex (doll's eyes reflex) evaluates the following cranial nerves: III (Oculomotor nerve) : Controls most of the eye’s movements, including the medial rectus (for adduction of the eye). VI (Abducens nerve) : Controls the lateral rectus muscle, responsible for moving the eye laterally. VIII (Vestibulocochlear nerve) : Involved in the vestibular system, which helps detect head movement and contributes to the reflex response. Thus, the correct answer is Option D: A, C, and D . Oculocephalic Reflex (Doll's Eyes Reflex) Assesses the vestibular-ocular reflex (VOR) to evaluate cranial nerves 3, 6, and 8, brainstem function, and overall neurologic status. It is commonly used in neurologic critical care, for neonates, anesthetized patients, and those with dizziness. Procedure: The examiner holds the patient’s eyelids open and rotates their head from side to side. Positive reflex: Eyes move in the opposite direction of head rotation, indicating intact brainstem function. Negative reflex: Eyes stay midline, indicating possible severe brainstem dysfunction. A similar test can be performed for vertical eye movements. Clinical Significance: Vestibular Lesions: Lesions to the vestibular system can impair the reflex. Dizziness: Abnormal responses indicate possible vestibular dysfunction. Psychiatric Patients: Schizophrenia and bipolar patients show abnormal smooth pursuit eye movements but typically maintain an intact doll's eyes reflex. Intensive Care: The reflex can predict altered mental status (delirium, coma) after sedation cessation. Neonates: The reflex is active in neonates before 11.5 weeks and serves as a developmental milestone. Anesthesia & Ophthalmologic Surgery: The reflex is useful for assessing brainstem function in these contexts. Caution: Do not perform on patients with head or neck injuries that could worsen from the reflex technique.



@Neet_pg_bot

References:

Doll's Eyes - StatPearls - NCBI Bookshelf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: What would be the next investigative step for a 27-year-old female patient who presents with sudden vision loss accompanied by a relative afferent pupillary defect in her right eye, while her left eye appears normal on examination? 1. MRI brain & orbit 2. Myelin oligodendrocyte glycoprotein (MOG) and neuromyelitis optica (NMO) antibodies 3. Blood sugar 4. Thyroid profile (Q855244)

A. 1, 2 and 3

B. 1 and 2

C. 2 and 3

D. 3 and 4

Correct Answer: A. 1, 2 and 3

Explanation: The sudden decrease in vision and relative afferent pupillary defect (RAPD) suggests a pathology with the optic nerve in the right eye, most likely due to optic neuritis, which can be associated with demyelinating diseases like Multiple sclerosis (MS) or Neuromyelitis optica (Devic's disease). MRI scan of the brain and orbit should preferably be performed in each case presenting with acute optic neuritis to check for nerve swelling, optic nerve glioma, cerebral, and brainstem lesion. MOG and NMO antibodies: These antibodies are seen in autoimmune conditions affecting the central nervous system, like Multiple sclerosis and neuromyelitis optica (NMO). Detection of these antibodies can help confirm the diagnosis. Blood sugar levels have to be checked. Because Optic neuritis is Primarily treated with Steroids as Steroids may precipitate hyperglycemia in individuals with DM. A thyroid profile could be relevant in cases of thyroid eye disease, but this typically presents different symptoms, such as eye protrusion, redness, and double vision. Given the scenario, thyroid dysfunction is less likely. Thus Option A: 1, 2 and 3 is the correct answer Options B, C, and D are incorrect.

References:

Kanski’s Clinical Ophthalmology, 9th edition, page 754- 757

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: What is the probable diagnosis for a man of 45 years who arrives at the emergency department with an abrupt loss of vision and experiencing painful movements of the eye? Upon examination, a relative afferent pupillary defect is observed in the affected eye. Further visual field testing demonstrates a central scotoma in the same eye. (Q857235)

A. Optic nerve glioma

B. Optic neuritis


C. Eale’s disease


D. Neuroretinitis

Correct Answer: B. Optic neuritis

Explanation: Clinical scenario of sudden visual loss associated with Relative afferent pupillary defect (RAPD) is suggestive of Optic Neuritis . Optic neuritis: Classification of Optic neuritis: Papillitis Retrobulbar Neuritis Neuroretinitis (Option D) Site Optic disc Optic nerve behind the eyeball. Optic disc and surrounding retina in the macular area. Characteristic features Fundoscopy reveals a blurred disc margin due to leakage and increased capillary permeability, edematous and hyperaemic disc, and physiological cup is obliterated. {{caption_text}} Pain on eye elevation RAPD (relative afferent pupillary defect) is observed. Fundoscopic examination typically reveals normal optic disc, at least initially, which is why the classic phrase, “the patient sees nothing, and the doctor sees nothing” is often used to describe retrobulbar neuritis. A macular star is seen in fundoscopy

(accumulation of lipids). Optic atrophy present {{caption_text}} Clinical features Visual loss: Monocular sudden, progressive and profound visual loss is the hallmark of acute optic neuritis. Dark adaptation may be lowered. Visual obscuration in bright light is a typical symptom of acute optic neuritis. Impairment of color vision is always present in optic neuritis. Uhthoff's symptom: Episodic transient obscuration of vision on exertion and on exposure to heat, which recovers on resting or moving away from the heat, occurs in patients with isolated optic neuritis. Pulfrich's phenomenon: Depth perception may be impaired, particularly for the moving object. Visual acuity is usually reduced markedly. Visual field changes: The most common field defect in optic neuritis is a relative central or centrocecal scotoma. Contrast sensitivity is impaired. Management of Optic neuritis: Treatment Indications Details Conservative management (Observation and Monitoring) Mild symptoms Known history of similar episodes Many cases improve spontaneously. Suitable for mild symptoms but not recommended for severe cases or first episodes. High-Dose Corticosteroids Severe vision loss No history of previous episodes MRI evidence of demyelination Intravenous methylprednisolone (1 g daily for 3 days) followed by oral prednisolone taper as established by the ONTT trial (Optic Neuritis Treatment Trial). Optic nerve glioma (Option A): is a tumor that can affect the optic nerve, but it is more commonly seen in children and young adults . MRI shows fusiform thickening of the intra-orbital portion of the optic nerve. {{caption_text}} Eale's disease (Option C): Also known as retinal vasculitis, typically presents with recurrent episodes of retinal hemorrhages , floaters, and vision loss in young males .





Macular star



Optic nerve glioma (Arrow)

References:

Parsons Diseases of the eye, 23rd Edition, pages 856-865.

Kanski’s Clinical Ophthalmology, 9th edition, pages 754,770

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: What is the probable diagnosis for a man of 45 years who arrives at the emergency department with an abrupt loss of vision and experiencing painful movements of the eye? Upon examination, a relative afferent pupillary defect is observed in the affected eye. Further visual field testing demonstrates a central scotoma in the same eye. (Q857665)

A. Optic nerve glioma

B. Optic neuritis


C. Eale’s disease

D. Neuroretinitis

Correct Answer: B. Optic neuritis

Explanation: Clinical scenario of sudden visual loss associated with Relative afferent pupillary defect (RAPD) is suggestive of Optic Neuritis . Optic neuritis: Classification of Optic neuritis: Papillitis Retrobulbar Neuritis Neuroretinitis (Option D) Site Optic disc Optic nerve behind the eyeball. Optic disc and surrounding retina in the macular area. Characteristic features Fundoscopy reveals a blurred disc margin due to leakage and increased capillary permeability, edematous and hyperaemic disc, and physiological cup is obliterated. {{caption_text}} Pain on eye elevation RAPD (relative afferent pupillary defect) is observed. Fundoscopic examination typically reveals normal optic disc, at least initially, which is why the classic phrase, “the patient sees nothing, and the doctor sees nothing” is often used to describe retrobulbar neuritis. A macular star is seen in fundoscopy (accumulation of lipids). Optic atrophy present {{caption_text}} Clinical features Visual loss: Monocular sudden, progressive and profound visual loss is the hallmark of acute optic neuritis. Dark adaptation may be lowered. Visual obscuration in bright light is a typical symptom of acute optic neuritis. Impairment of color vision is always present in optic neuritis. Uhthoff's symptom: Episodic transient obscuration of vision on exertion and on exposure to heat, which recovers on resting or moving away from the heat, occurs in patients with isolated optic neuritis. Pulfrich's phenomenon: Depth perception may be impaired, particularly for the moving object. Visual acuity is usually reduced markedly. Visual field changes: The most common field defect in optic neuritis is a relative central or centrocecal scotoma. Contrast sensitivity is impaired. Management of Optic neuritis: Treatment Indications Details Conservative management (Observation and Monitoring) Mild symptoms Known history of similar episodes Many cases improve spontaneously. Suitable for mild symptoms but not recommended for severe cases or first episodes. High-Dose Corticosteroids Severe vision loss No history of previous episodes MRI evidence of demyelination Intravenous methylprednisolone (1 g daily for 3 days) followed by oral prednisolone taper as established by the ONTT trial (Optic Neuritis Treatment Trial). Optic nerve glioma (Option A): is a tumor that can affect the optic nerve, but it is more commonly seen in children and young adults . MRI shows fusiform thickening of the intra-orbital portion of the optic nerve. {{caption_text}} Eale's disease (Option C): Also known

as retinal vasculitis, typically presents with recurrent episodes of retinal hemorrhages , floaters, and vision loss in young males .







References:

Parsons Diseases of the eye, 23rd Edition, page 856-865.

Kanski’s Clinical Ophthalmology, 9th edition, page 754,770

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: In a patient diagnosed with a stroke, their visual field testing indicated macula sparing homonymous hemianopia. Where is the probable location of the lesion within the optic pathway? (Q857749)

A. Optic tract

B. Occipital lobe

C. Optic radiation

D. Optic chiasma

Correct Answer: B. Occipital lobe

Explanation: Macula sparing homonymous hemianopia is most commonly caused by a lesion in the occipital lobe, usually due to a PCA stroke. The macula is spared because it receives dual blood supply: Posterior cerebral artery (PCA) Middle cerebral artery (MCA) Therefore, even if PCA territory is damaged, the central (macular) vision remains intact, producing macula-sparing homonymous hemianopia. Incorrect Options: Option A) Optic tract : Produces contralateral homonymous hemianopia. Macular sparing is NOT characteristic. May show Wernicke's hemianopic pupil. Option C) Optic radiation Temporal lobe (Meyer's loop) → contralateral superior quadrantanopia ("pie in the sky") Parietal lobe → contralateral inferior quadrantanopia ("pie on the floor") Complete homonymous hemianopia can occur, and rare cases may show macular sparing, but this is less typical than in occipital lobe lesions. Option D) Optic chiasma: Causes bitemporal hemianopia , NOT homonymous hemianopia. No macular sparing pattern.

References:

Parsons Diseases of the Eye 23rd Edition, Pages 85,87,88,1184

↑ *Back to this question in Part A*

@Neet_pg_bot


Q11: A patient presents to the OPD with the given visual field effect. What is the probable site of the lesion based on the image displayed? (Q858134)


A. 3

B. 1

C. 4


D. 6







Correct Answer: A. 3

Explanation: - 3 Site of Lesion : Optic Chiasma. Effect on Visual Field : Bitemporal Hemianopia (Heteronymous Hemianopia). This means the patient will have loss of vision in the outer (temporal) fields of both eyes, which is a classic sign of a lesion at the optic chiasma where the nasal fibers from both eyes cross. Visual Pathway: Temporal fibers carry impulses from the Nasal field, and Nasal fibres carry the Temporal field. Visual Pathway Lesions and Defects Site of Lesion Fibers Affected Field Defect Optic Nerve (Option B ruled out) Ipsilateral Temporal + Nasal Fibers Anopia Optic Chiasma (Option A) Bilateral Nasal Fibers Bitemporal Hemianopia (Heteronymous Hemianopia) Optic Tract (Option C ruled out) Lateral Geniculate Body (LGB Lesion) Optic Radiation Ipsilateral Temporal Fibers + Contralateral Nasal Fibers Right Homonymous Hemianopia Visual Cortex without the involvement of tip (Post. Cerebral Artery) Ipsilateral Temporal Fibers + Contralateral Nasal Fibers except macular representation Homonymous Hemianopia with Macular Sparing Tip of Visual Cortex (Middle Cerebral Artery) Macular Representation of Ipsilateral Temporal and Contralateral Nasal Fibers. Macular Homonymous Hemianopia Aspect Inferior Optic Radiations (Meyer's Loop) Superior Optic Radiations (Baum's Loop) Location of Lesion Temporal lobe Parietal lobe (Option D ruled out) Affected Visual Field Superior quadrants of the contralateral visual field Inferior quadrants of the contralateral visual field Visual Defect Superior homonymous quadrantanopia Inferior homonymous quadrantanopia Associated Symptoms Memory loss Hearing loss Auditory hallucinations Contralateral hemiplegia Hemianesthesia Optokinetic nystagmus 1 (Option B)- Optic Nerve : Causes complete blindness in one eye, not a bitemporal hemianopia. 4 (Option C)- Optic Tract : Causes homonymous hemianopia, but not bitemporal hemianopia. 6 (Option D)- Visual Cortex : Typically causes homonymous hemianopia with possible macular sparing, not bitemporal hemianopia.







@Neet_pg_bot




@Neet_pg_bot


30
30
Left
Right
@Neet_pg_bot



Visual Field
Left
Right
@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot



References:

Parsons Diseases of the eye, 23rd Edition, Pg - 1184

Yanoff & Duker Ophthalmology, 5th edition, Pg - 913,919

↑ *Back to this question in Part A*

@Neet_pg_bot


Q12: Identify the site of the lesion. (Q856720)

A. Right occipital lobe

B. Left occipital lobe


C. Right optic tract


D. Optic chiasma



Correct Answer: B. Left occipital lobe

Explanation: The visual field chart shows a right homonymous hemianopia — loss of the right visual field in both eyes , indicating a post-chiasmal lesion on the contralateral (left) side , i.e., the left occipital lobe . Right field loss in both eyes → Homonymous hemianopia Same side in both eyes → Post-chiasmal Opposite to defect → Left occipital lobe Macular Sparing: Occipital lobe lesions (e.g., PCA infarcts) often show macular sparing due to dual blood supply from the PCA and MCA . Here, the absence of macular sparing suggests a larger or complete occipital lesion , but it does not exclude an occipital site — it simply indicates involvement of both vascular territories . Right occipital lobe (Option A) and Right optic tract (Option C): Lesion at right occipital lobe/right optic tract will result in Left sided Homonymous hemianopia. ■■■■■■■ Optic Chiasma(Option D): Lesion at optic chiasma will result in Bitemporal hemianopia





References:

Parsons Diseases of the Eye 23rd Edition, Pages 85,87,88,1184

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: In a patient, a 33-year-old female presents with symptoms of gradual vision loss specifically affecting the right halves of both eyes. The question pertains to the location of the lesion in the optic pathway. (Q856124)

A. Left optic tract


B. Right visual cortex


C. Optic chiasma


D. Right optic nerve


Correct Answer: A. Left optic tract

Explanation: - Left optic tract The clinical presentation of difficulty seeing objects in the right half of the visual field in both eyes indicates a visual field defect known as Right Homonymous Hemianopia , which occurs due to a defect in the Left Optic Tract . Visual Pathway: Temporal fibers carry impulses from the Nasal field, and Nasal fibres carry the Temporal field. Components of Visual Pathway Retina First-order Neurons: Bipolar cells in the inner nuclear layer of the retina. Second-order Neurons: Ganglion cells in the retina; their axons form the optic nerve . Optic Nerve Carries visual information from the retina to the optic chiasm. Temporal and Nasal fibres of the same eye. Optic Chiasma Partial decussation occurs Nasal fibres cross to the opposite side, while temporal fibres remain ipsilateral. Optic Tract Carries visual information from the optic chiasma to the Lateral Geniculate body (LGB). Temporal fibres of the same eye and Nasal fibres of the opposite eye. Lateral Geniculate Body (LGB) Third Order Neurons: Relay visual information to the visual cortex. Optic Radiations Meyer's Loop: Carries information from lower retinal quadrants to the inferior part of the visual cortex. Baum’s Loop (Dorsal): Carries information from upper retinal quadrants to the superior part of the visual cortex. Occipital Cortex The primary visual cortex (V1), corresponding to Brodmann area 17 , is situated on the medial side of the occipital lobe, particularly along the borders of the calcarine fissure. Visual Pathway Lesions and Defects Site of Lesion Fibers Affected Field Defect Optic Nerve Ipsilateral Temporal + Nasal Fibers Anopia (Option D) Optic Chiasma Bilateral Nasal Fibers Bitemporal Hemianopia (Heteronymous Hemianopia) (Option C) Optic Tract/ Lateral Geniculate Body (LGB)/ Optic Radiation Ipsilateral Temporal Fibers + Contralateral Nasal Fibers Homonymous Hemianopia Visual Cortex without the involvement of tip (Posterior Cerebral Artery) Ipsilateral Temporal Fibers + Contralateral Nasal Fibers except macular representation Homonymous Hemianopia with Macular Sparing (Option B) Tip of Visual Cortex (Middle Cerebral Artery) Macular Representation of Ipsilateral Temporal and Contralateral Nasal Fibers. Macular Homonymous Hemianopia





@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot



References:

Parsons Diseases of the Eye 23rd Edition, Pages 85,87,88,1184.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: What is the likely reason for a patient showing both unilateral and bilateral abducent palsy? (Q855026)

A. Retinoblastoma

B. Orbital pseudotumor


C. Orbital cellulitis


D. Cavernous sinus thrombosis

Correct Answer: D. Cavernous sinus thrombosis

Explanation: Abducens Nerve (cranial nerve VI) Type Motor Origin Pons (lower border, opposite the pyramid of the medulla). Course Functional component General somatic efferent fibres : Arise from the abducent nucleu s in

the pons and supply the lateral rectus muscle of the eyeball. General somatic afferent fibres carry proprioceptive sensations from the lateral rectus and terminate in the mesencephalic nucleus of the trigeminal nerve. Supply Lateral rectus muscle of the eyeball Clinical aspect Cavernous sinus thrombosis leads to abducens nerve palsy Lesion or injury (elevated intracranial pressure) of the abducens nerve results in paralysis of the lateral rectus muscle, leading to: Inability to abduct the eye Diplopia (double vision) Convergent squint (due to the unopposed action of medial rectus) Incorrect Options: Retinoblastoma (Option A) is a type of eye cancer that primarily affects young children. It originates in the retina. It can occur unilaterally and also be bilateral. Orbital pseudotumor (Option B) , also known as idiopathic orbital inflammatory syndrome, is a condition characterized by inflammation and swelling of the tissues within the orbit. Orbital cellulitis (Option C) is an infection characterized by inflammation and infection of the soft tissues within the orbit. It is usually caused by the spread of an infection from the adjacent structures, such as the sinuses or eyelids.





@Neet_pg_bot

References:

Grey's anatomy 42nd edition, pg 782.

BD Chaurasia’s Human Anatomy, 8th edition, pg 73, 74.

Vishram Singh Textbook of Anatomy, 2nd edition, pg 337, 338.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: The lesion in which of the following nerve is responsible for right eye squint, as depicted below? (Q855031)

A. Oculomotor

B. Trochlear

C. Trigeminal

D. Abducent



Correct Answer: D. Abducent

Explanation: A lesion in the abducent nerve causes paralysis of the lateral rectus muscle, which is responsible for abducting the eye. In its absence, the unopposed pull of the medial rectus causes the eye to deviate inward , resulting in esotropia (convergent squint). This condition is commonly seen in abducent nerve palsy. Paralytic squint : Paralytic squint is a type of incomitant squint in which ocular deviation results from complete or incomplete paralysis of one or more extraocular muscles. History of diplopia and confusion Secondary deviation is greater than primary deviation. Ocular movements are restricted towards the action of paralyzed muscle. Head is turned towards the action of paralyzed muscle. Incorrect Options: Oculomotor (Option A) lesion causes ptosis, mydriasis (pupil dilation), and difficulty moving the eye upward, downward, or inward. The eye often rests in a "down and out" position . Trochlear (Option B) nerve lesion affects the superior oblique muscle, leading to vertical diplopia and difficulty looking downward, particularly in adduction Trigeminal (Option C) nerve is responsible for facial sensation and mastication; it does not innervate any extraocular muscles.

References:

AK Khurana 7th edition page 583, 584

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: The purpose of the photostress test is to distinguish between: (Q858507)

A. Macula and optic nerve disease

B. Cataract and glaucoma

C. Lens and cornea

D. Retina and vitreous pathologies

Correct Answer: A. Macula and optic nerve disease

Explanation: The photostress test is used to differentiate between macular and optic nerve disease. In this test, a bright light is shone into the eye to bleach the photoreceptors, and the recovery time for visual acuity is measured. A prolonged recovery time indicates macular dysfunction, while optic nerve diseases typically do not affect the photostress recovery time. Cataract and glaucoma (Option B) These conditions affect different structures (cataract affects the lens, and glaucoma affects the optic nerve). The photostress test does not assess the lens or intraocular pressure-related damage. Lens and cornea (Option C) The photostress test is unrelated to refractive or structural abnormalities of the lens and cornea. Retina and vitreous pathologies (Option D) While the photostress test involves the retina, it is specific for evaluating macular function, not general retinal issues or vitreous pathologies (e.g., retinal detachment or vitreous hemorrhage).

References:

https://pubmed.ncbi.nlm.nih.gov/836667/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: Where is the location of the lesion in Wernicke's hemianopic pupil? (Q858511)

A. Optic nerve

B. Optic tract

C. Optic radiation

D. Optic chiasma

Correct Answer: B. Optic tract

Explanation: Abnormal Pupillary Reactions Reaction Description Causes Amaurotic light reflex Absence of direct light reflex on the affected side and an absence of consensual light reflex on the normal side. Optic and retinal lesions on the affected side Efferent pathway defect Absence of both direct and consensual light reflex on the affected side and presence of both direct and consensual light reflex on the normal side. Near reflex is also absent on the affected side. Parasympatholytic drugs (e.g. atropine, homatropine) Internal ophthalmoplegia Third-nerve paralysis. Marcus Gunn pupil or RAPD The pupils paradoxically dilate when the flashlight is swung from the unaffected to the affected eye due to the decreased light perception. It is the first sign of optic nerve disorder. Tested by a swinging flashlight test. Incomplete optic nerve lesions (Optic neuritis) (Option A) and severe retinal diseases (Retinopathy, CRVO) Adie’s Pupil Reaction to light is absent, and the near reflex is very slow and tonic. It is usually unilateral, associated with absent knee jerk and occurs more often in young women. It shows pupillary constriction to 0.125% Pilocarpine due to denervation supersensitivity. Postganglionic parasympathetic pupillomotor damage. Argyll Robertson Pupil (ARP) Reaction to near reflex is present, but light reflex (both direct and consensual) is absent, i.e. there is light near dissociation. Neurosyphilis Wernicke's hemianopic pupil Light reflex is absent when light is thrown on the temporal half of the retina of the affected side and the nasal half of the opposite side, while it is present when the light is thrown on the nasal half of the affected side and the temporal half of the opposite side. Optic tract lesion (Option B) Optic radiation (Option C) Lesions in the optic radiation

primarily cause visual field defects but do not typically result in pupillary abnormalities because these fibers are past the pupillary reflex arc. Optic chiasma (Option D) Lesions in the optic chiasma result in bitemporal hemianopia and do not produce Wernicke's hemianopic pupil.

References:

Parsons Diseases of the eye, 23rd Edition, Pg 280

Yanoff & Duker Ophthalmology, 5th edition, Pg 711

Kanski's Clinical Ophthalmology, 10th Edition, Page 792,795,796

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: What is the probable diagnosis for a patient who exhibits miosis, anhidrosis, mild ptosis, and a persistent small pupil even in low light conditions? (Q859316)

A. Adie’s tonic people

B. Horner syndrome

C. Marcus gunn pupil

D. Argyll Robertson pupil

Correct Answer: B. Horner syndrome

Explanation: The clinical presentation of miosis, anhidrosis, mild ptosis, and a persistent small pupil even in low light is suggestive of horner syndrome. Ptosis is a result of the weakness of Müller muscle and miosis due to the unopposed action of the sphincter pupillae with resultant anisocoria. Pancoast tumor can compress the sympathetic nerves, leading to the classic triad of symptoms seen in Horner syndrome: ptosis, miosis (constricted pupil), and ipsilateral facial anhidrosis (lack of sweating). Acquired Neurogenic Ptosis: Acquired ptosis with drooping of the upper eyelid caused by a disruption in the nerve supply to the muscles responsible for elevating the eyelid. Causes: Third Cranial (Oculomotor) Nerve Palsy: The third nerve can be damaged by trauma, ischemia, infection, tumor, or compression, causing ptosis and diplopia.Oculomotor nerve palsy causes ptosis as it supplies the levator palpebrae superioris muscle. The ptotic eyelid can be elevated with a frontalis sling or levator aponeurosis advancement. Horner's Syndrome: Involves a lesion in the sympathetic pathway , affecting the Müller’s muscle, leading to mild ptosis accompanied by miosis (pupil constriction) and anhidrosis (loss of sweating). Ophthalmoplegic migraine Multiple Sclerosis Diagnosis: Topical apraclonidine test is now considered the test of choice. Apraclonidine is an alpha-2 adrenergic agonist that also acts as a very weak alpha 1-adrenergic agonist; when topically applied in Horner’s syndrome, it causes dilation of the affected pupil and possibly lid elevation, without change to the normal pupil. Incorrect Options Adie’s tonic people (Option A) It is a unilateral condition and is characterized by anisocoria (different size of pupil) and direct light reflex is sluggish or absent. Marcus gunn pupil (Option C) Aka relative afferent pupillary defect is characterized by paradoxical dilatation of both the pupil when light is flashed on the diseased eye. Argyll Robertson pupil (Option D) It is a specific sign of late-stage

syphilis. The bilaterally small pupils do not constrict when exposed to bright light.

References:

Yanoff and Duker Ophthalmology 5th edition 1275, 1276

https://www.ncbi.nlm.nih.gov/books/NBK500000/#:~:text=is%20considered%20the-,test%20of%20choice,-due%20to%20its

https://www.reviewofoptometry.com/article/horners-syndrome-a-positive-apraclonidine-testnow-what#:~:text=Apraclonidine%20is%20an%20alpha%2D2%20adrenergic%20agonist%20that%20also%20acts%20as%20a%20very%20weak%20alpha%201%2Dadrenergic%20agonist%3B%20when%20topically%20applied%20in%20Horner%E2%80%99s%20syndrome%2C%20it%20causes%20dilation%20of%20the%20affected%20pupil%20and%20possibly%20lid%20elevation%2C%20without%20change%20to%20the%20normal%20pupil

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: What is the clinical presentation of a patient with a tumor located in the anterior pituitary gland causing compression on the optic chiasma? (Q859329)

A. Homonymous hemianopia

B. Inferior quadrantanopia hemianopia


C. Bitemporal hemianopia

D. Superior quadrantanopia hemianopia


Correct Answer: C. Bitemporal hemianopia

Explanation: Correct Option C - Bitemporal hemianopia: Bitemporal hemianopia is usually caused by tumors in the region of the sella turcica, pressure by a suprasellar aneurysm, or chronic arachnoiditis; these press upon the chiasma, so that the fibers going to the nasal halves of each retina are destroyed. So a tumor in the anterior pituitary pressing on the optic chiasma will present as a case of Bitemporal hemianopia. Pituitary adenoma: The most common visual defect associated with Pituitary Adenoma is Bitemporal hemianopia, which results from the compression of Binasal fibres that cross at the Optic Chiasma. It is the most common Suprasellar mass in adults. Classification: Microadenomas : <10 mm; often incidental or present with endocrine symptoms. Macroadenomas : >10 mm; may cause visual field deficits due to mass effect. Common Types : Prolactinomas : 30% of cases present with amenorrhea and galactorrhea. Non-functioning Adenomas : 20%; no hormone secretion. Growth Hormone-Secreting Adenomas : 15%; causes acromegaly. ACTH-Secreting Adenomas : 10% leads to Cushing's disease. Clinical features: Visual Defect : Compression of the optic chiasm often causes bitemporal hemianopia . Endocrine Symptoms : Amenorrhea, galactorrhea, and other hormonal imbalances. {{caption_text}} Imaging and diagnosis involve MRI as the primary modality, along with serum prolactin and thyroid function tests and visual field charting. Treatment : Medical management Prolactinomas : Dopamine agonists (e.g., bromocriptine, cabergoline). Growth Hormone-Secreting Tumors : Somatostatin analogues (e.g., octreotide). Surgical

management Transsphenoidal Approach : Preferred for most cases; craniotomy may be needed for large tumours. Sublabial Approach Indications : Acute visual deterioration, GH-secreting tumours, Cushing's disease, or large non-functioning adenomas causing symptoms. {{caption_text}} Complications : Pituitary Apoplexy : Sudden haemorrhage or infarction in the tumour, presenting with headache, visual loss, and ophthalmoplegia. Postoperative Risks : CSF leaks, pituitary insufficiency, and diabetes insipidus. Incorrect Options: Homonymous hemianopia (Option A) is caused by post chiasmal lesions like optic tract, optic radiations, and visual cortex lesions. Inferior quadrantanopia hemianopia (Option B) It is caused by a lesion of the fibers of the optic radiations in the anterior parietal lobe. Superior quadrantanopia hemianopia (Option D) It is caused by a lesion of the fibers of the optic radiations in the temporal lobe.



@Neet_pg_bot



@Neet_pg_bot

References:

Bailey & Love's Short Practice of Surgery, 28th edition: Pg - 716,717.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: Maximum visual field is seen in? (Q859816)

A. Temporal

B. Nasal

C. Inferior

D. Superior

Correct Answer: A. Temporal

Explanation: The maximum visual field is observed temporally, extending to about 100-110 degrees. Nasal: ~60 degrees. Superior: ~60 degrees. Inferior: ~70-75 degrees. This distribution is due to anatomical constraints, such as the nose and brows.

References:

https://www.ncbi.nlm.nih.gov/books/NBK220/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: Sudden painful loss of vision in one eye is due to the lesion at? (Q860534)

A. Optic nerve


B. Optic chiasma

C. Optic tract


D. Visual cortex

Correct Answer: A. Optic nerve

Explanation: Sudden painful loss of vision in one eye is typically caused by conditions affecting the optic nerve, such as optic neuritis. It often presents with pain on eye movement and is associated with demyelinating diseases like multiple sclerosis. Classification of Optic neuritis: Papillitis Retrobulbar Neuritis Neuroretinitis Site Optic disc Optic nerve behind the eyeball. Optic disc and surrounding retina in the macular area. Characteristic features Fundoscopy reveals a blurred disc margin due to leakage and increased capillary permeability, an edematous and hyperaemic disc, and an obliterated physiological cup. {{caption_text}} Pain on eye elevation RAPD (relative afferent pupillary defect) is observed. Fundoscopic examination typically reveals normal optic disc , at least initially, which is why the classic phrase, “ the patient sees nothing, and the doctor sees nothing ” is often used to describe retrobulbar neuritis. A macular star is seen in fundoscopy (accumulation of lipids). {{caption_text}} Clinical features: Visual loss: Monocular sudden, progressive and profound visual loss is the hallmark of acute optic neuritis. Dark adaptation may be lowered. Visual obscuration in bright light is a typical symptom of acute optic neuritis. Impairment of color vision is always present in optic neuritis. Uhthoff's symptom: Episodic transient obscuration of vision on exertion and on exposure to heat, which recovers on resting or moving away from the heat, occurs in patients with isolated optic neuritis. Pulfrich's phenomenon: Depth perception may be impaired, particularly for the moving object. Visual acuity is usually reduced markedly. Visual field changes: The most common field defect in optic neuritis is a relative central or centrocecal scotoma. Contrast sensitivity is impaired. {{caption_text}} Optic chiasma (Option B): Lesions here cause bitemporal hemianopia but do not lead to monocular vision loss. Optic tract (Option C): Lesions result in homonymous hemianopia but do not cause sudden vision loss in one eye. Visual cortex (Option D): Lesions here typically lead to cortical blindness or homonymous visual field defects but are not associated with painful monocular vision loss.


@Neet_pg_bot


@Neet_pg_bot



References:

Parsons Diseases of the eye, 23rd Edition, page 856-864.

Kanski’s Clinical Ophthalmology, 9th edition, page 754,770

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: A 45-year-old patient presents with difficulty reacting to bright light. On examination, the pupillary light reflex is absent but the near reflex is preserved. Where is the lesion likely located? (Q866846)

A. Optic nerve

B. Oculomotor nerve

C. Edinger–Westphal nucleus

D. Pretectal area of the midbrain

Correct Answer: D. Pretectal area of the midbrain

Explanation: Absence of the light reflex with preservation of the near reflex is known as light-near dissociation . This occurs when the lesion interrupts the pathway from the optic tract to the Edinger–Westphal nucleus before the convergence–accommodation pathway branches — classically in the pretectal area of the midbrain . Common causes: Argyll Robertson pupil (neurosyphilis), Parinaud's syndrome (pineal tumors). Mechanism: Light reflex pathway: Retina → optic nerve → optic chiasma → optic tract → pretectal area → bilateral Edinger–Westphal nuclei → oculomotor nerve → sphincter pupillae. Near reflex pathway bypasses the pretectal area lesion and reaches Edinger–Westphal nuclei via cortical connections, so it remains intact. Incorrect Options: Optic nerve (Option A): Lesion causes loss of both light and near reflex on the affected side (afferent defect), not selective loss of light reflex. Oculomotor nerve (Option B): Lesion causes loss of both light and near reflex (efferent defect), along with ptosis and eye deviation down and out. Edinger–Westphal nucleus (Option C): Damage here would abolish both light and near reflexes, as it is the parasympathetic motor nucleus for pupillary constriction.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: A patient presents with a down and out right eye with ptosis since birth. What is the nerve palsy? (Q866852)

A. 4th nerve palsy

B. 3rd nerve palsy

C. 6th nerve palsy

D. 7th nerve palsy

Correct Answer: B. 3rd nerve palsy

Explanation: The down and out position of the eye indicates unopposed action of the lateral rectus (CN VI) and superior oblique (CN IV) muscles, suggesting paralysis of the other extraocular muscles supplied by oculomotor nerve (CN III) . Ptosis occurs due to paralysis of the levator palpebrae superioris (also supplied by CN III). Since it is present since birth , this is likely a congenital oculomotor nerve palsy . Key features of 3rd nerve palsy: Eye position: abducted and depressed ("down and out"). Ptosis due to levator palpebrae weakness. May also have pupil dilation if parasympathetic fibers are involved. Causes: congenital, trauma, aneurysm (PComm), diabetes. Incorrect Options: 4th nerve palsy (Option A): Affects superior oblique → vertical diplopia, head tilt to opposite side; eye is slightly up and in, not down and out with ptosis. 6th nerve palsy (Option C): Affects lateral rectus → inability to abduct eye; resting position is inward (esotropia), not down and out. 7th nerve palsy (Option D): Affects facial muscles, not extraocular muscles; presents with facial asymmetry, inability to close eye, no change in eye position.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q24: Based on the eye movement findings shown in the image below, what is the most likely diagnosis? (Q865341)

A. Horizontal gaze palsy

B. Internuclear ophthalmoplegia (INO)

C. 3rd nerve palsy

D. Myasthenia gravis



@Neet_pg_bot

Correct Answer: B. Internuclear ophthalmoplegia (INO)

Explanation: The eye on the side of the MLF lesion cannot turn inward (adduct) fully when attempting to look horizontally away from the lesion. The contralateral eye abducts with possible nystagmus. Horizontal Gaze Palsy (Option A ruled out) Definition: Inability to voluntarily move both eyes beyond the midline in one horizontal direction. Eye Movements: Affected Side: Both eyes cannot look towards the involved side. Unaffected Side: Gaze to the opposite side is typically normal. Convergence: Usually preserved. Oculocephalic Reflex: May overcome the palsy (if supranuclear lesion), indicating brainstem integrity. Internuclear Ophthalmoplegia (INO) (Option B) Definition: Lesion of the medial longitudinal fasciculus (MLF), disrupting horizontal gaze coordination. Eye Movements: Ipsilateral Eye (side of lesion): Impaired or slowed adduction (inward movement) when looking away from the lesion. Contralateral Eye (opposite side): Abducting nystagmus (jerking outward movements). Convergence: Crucially preserved. Vertical Gaze: May be affected in bilateral cases. 3rd Nerve Palsy (Oculomotor Nerve Palsy) (Option C ruled out) Definition: Dysfunction of the oculomotor nerve, controlling most eye movements and eyelid elevation. Eye Movements: Resting Position: Eye is "down and out" (abducted and

depressed). Impaired Movements: Cannot adduct, elevate, or depress past the midline. Pupil: Often dilated (if parasympathetic fibers involved). Ptosis: Drooping of the eyelid present (partial or complete). Myasthenia Gravis (Option D ruled out) Definition: Autoimmune disorder causing fluctuating weakness of voluntary muscles, including ocular muscles. Eye Movements (highly variable and fatiguing): Ptosis: Common, often asymmetrical, fluctuates (worsens with fatigue, improves with rest). Diplopia: Common, variable, affecting any direction of gaze, often changing. Ophthalmoplegia: Weakness in any eye movement, non-localizing (doesn't fit a specific nerve palsy). Fatigability: Eye movements worsen with sustained effort. Pupils: Always normal. Cogan's Lid Twitch: Brief overshoot of upper eyelid after downward gaze, followed by slow drop.

References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 927

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: Which of the following nerves gives sensory supply to the orbit? (Q865355)

A. Vagus nerve

B. Trigeminal nerve

C. Hypoglossal nerve

D. Oculomotor nerve

Correct Answer: B. Trigeminal nerve

Explanation: Trigeminal nerve (Cranial Nerve V) (Option B) : This nerve is the primary sensory nerve for the face, including the orbit and its contents. Its ophthalmic division (V1) specifically provides sensory innervation to: The eye itself (cornea, conjunctiva, iris, ciliary body) The lacrimal gland The skin of the upper eyelid, forehead, and bridge of the nose Parts of the nasal cavity and sinuses. Let's look at why the other options are incorrect: Vagus nerve (Cranial Nerve X) (Option A ruled out) : Primarily involved in parasympathetic control of internal organs (heart, lungs, digestive system), and provides some sensory input from the ear and pharynx, but not the orbit. Hypoglossal nerve (Cranial Nerve XII) (Option C ruled out) : Purely a motor nerve that controls the muscles of the tongue. Oculomotor nerve (Cranial Nerve III) (Option D ruled out) : Primarily a motor nerve that controls most of the extraocular muscles (eye movements) and provides parasympathetic innervation to the pupil and ciliary muscle (for accommodation). It has very limited, if any, direct sensory supply to the orbit itself.

References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 937

Inderbir Singh, 6th edition, head and neck, Page 192

↑ *Back to this question in Part A*

Q26: In a patient with dense central (macular) corneal or lens opacity, which of the following statements about the visibility of intraocular structures is correct? (Q853130)

A. Both the iris and pupil are clearly visible


B. Only the pupillary margin is visible

C. Iris details are visible but not the pupil

D. Neither the iris nor the pupillary margin is visible

Correct Answer: D. Neither the iris nor the pupillary margin is visible

Explanation: A macular opacity refers to a dense central opacity of the cornea or lens involving the visual axis , while the peripheral areas remain clear . Because of this, light cannot pass through the central part , obscuring the view of the central iris and pupil , but the peripheral pupillary margin can still be seen through the clear surrounding area. In other words, the central opacity blocks visualization of intraocular structures behind it, while the edges of the pupil remain faintly visible through the unaffected peripheral region. Clinical Correlation: Seen in cases of healed keratitis , corneal scarring , or early central cataract . The patient typically has reduced central vision but some peripheral visibility remains. In total (leucomatous) opacities or mature cataracts , neither the iris nor the pupil is visible.



References:

Parsons Diseases of the Eye, Edition 23, page no: 441, 442

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: A 5-month-old baby is brought by the mother with complaints of the left upper eyelid moving up and down during breastfeeding or thumb sucking, which disappears when the baby is not being fed. What is the most likely diagnosis? (Q852492)

A. Marcus Gunn Jaw-Winking Syndrome

B. Lagophthalmos

C. Myasthenia Gravis

D. 3rd CN palsy

Correct Answer: A. Marcus Gunn Jaw-Winking Syndrome

Explanation: Marcus Gunn Jaw-Winking Syndrome (also known as the jaw-winking reflex) is a congenital condition where there is an abnormal connection between the trigeminal nerve (supplying Lateral pterygoid) and the oculomotor nerve (supplying LPS). This results in the upper eyelid (ptosis) lifting when the jaw moves, such as during chewing, sucking, or other movements involving the jaw. The eye movement (or "winking") is typically unilateral and can occur during activities like breastfeeding or thumb-sucking, as in this case. This reflex is not present when the baby is not feeding, which is a key feature of this condition. Congenital Ptosis: Associated with congenital weakness or maldevelopment of the levator palpebrae superioris (LPS) muscle. Congenital ptosis probably results from a failure of neuronal migration or development with muscular sequelae secondary to this. A minority of patients have a family history. Characteristic features: Drooping of one or both upper lids. Lid crease is either diminished or absent. Lid lag on downgaze due to tethering effect of abnormal LPS muscle. Types of congenital ptosis: 1. Simple congenital ptosis: Unilateral or bilateral Non-progressive No fluctuation: Unlike myogenic ptosis, congenital ptosis does not worsen throughout the day or improve with rest. 2. Congenital ptosis with associated weakness of superior rectus muscle. 3. Blepharophimosis syndrome: Congenital ptosis Blepharophimosis Telecanthus Epicanthus inversus. 4. Congenital synkinetic ptosis (Marcus Gunn jaw-winking ptosis) Retraction of the ptotic lid with jaw movements due to stimulation of ipsilateral pterygoid muscle. It occurs due to trigemino-oculomotor synkinesis. The pathophysiology of Marcus Gunn Jaw winking phenomenon is a congenital misdirection of mandibular branch of trigeminal nerve, i.e., this synkinetic movement results from a congenital aberrant connection between motor branches of the trigeminal nerve controlling muscles of mastication and superior division of the oculomotor nerve controlling the levator palpebrae superioris. Lagophthalmos (Option B): This condition is characterized by an inability to completely close the eyelids, usually due to facial nerve (VII) dysfunction or other causes of eyelid weakness. Myasthenia Gravis (Option C) : Myasthenia gravis typically presents with fatigable weakness of the muscles, including the eyelids (ptosis), which worsens with activity. 3rd CN Palsy (Option D) : A third cranial nerve palsy typically causes ptosis (drooping of the eyelid) and ocular motility issues (like strabismus or limited eye movement).



References:

Kanski’s Clinical Ophthalmology 9th edition page 77-78 Congenital ptosis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q28: A patient presents with breathlessness and is diagnosed with a Pancoast tumor. Which of the following findings is not typically associated with this condition? (Q852488)

A. Anhidrosis

B. Miosis

C. Ptosis

D. Exophthalmos

Correct Answer: D. Exophthalmos

Explanation: A Pancoast tumor is a type of lung cancer that occurs at the apex of the lung and can invade nearby structures, including the brachial plexus, sympathetic chain, and adjacent ribs. The tumor is commonly associated with a condition known as Horner's syndrome due to its involvement with the sympathetic nervous system. Horner's Syndrome Horner's syndrome is caused by a lesion of the sympathetic chain, which clinically manifests as ipsilateral miosis, facial anhidrosis, and ptosis (drooping of the eyelid). Causes: Central lesions Preganglionic Postganglionic Brainstem pathology (tumours, vascular diseases, demyelinating disorders) Spinal cord tumours Pancoast tumour Carotid and Aortic aneurysms Neck lesions Atherosclerosis of ICA (internal carotid artery)) Nasopharyngeal tumours Cavernous sinus pathology Cluster headaches Clinical manifestations: Miosis, ptosis, enophthalmos. Anhydrosis (loss of sweating): It is seen in preganglionic lesions. Absence of cilio-spinal reflex (involuntary response where the pupil of the eye dilates in response to painful stimuli applied to the skin of the neck or upper back) Acute Horner's syndrome with associated ipsilateral neck or facial pain requires urgent imaging of the neck for evaluation of a carotid dissection

References:

Parsons Diseases of the eye, 23rd Edition, Pg 276, 277. Yanoff & Duker Ophthalmology, 5th edition, Pg 965-968.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q29: Identify the visual field defect: (Q843978)

A. Altitudinal defect

B. Roennes nasal step

C. Blind spot enlargement


D. Arcuate scotoma



Correct Answer: C. Blind spot enlargement

Explanation: Correct Option C - Blind spot enlargement: The given image shows blind spot enlargement. Incorrect Options: Option A - Altitudinal defect: Option B - Roennes nasal step: Option D - Arcuate scotoma:






F

Roenne's
nasal step

@Neet_pg_bot



D

Bjerrum's (Arcuate)
Scotoma

@Neet_pg_bot

↑ *Back to this question in Part A*

@Neet_pg_bot


Q30: Identify the defect shown in the image given below: (Q844012)

A. Binasal hemianopia

B. Bitemporal hemianopia

C. Macula- sparing homonymous hemianopia

D. Altitudinal defect



Correct Answer: C. Macula- sparing homonymous hemianopia

Explanation: The visual field shows loss of the same side in both eyes → a homonymous hemianopia. The central vision (macular area) is preserved → macular sparing. This occurs when the occipital lobe (PCA territory) is affected but the macula is protected by: Dual blood supply from PCA + MCA. Therefore, the pattern fits macular-sparing homonymous hemianopia, most commonly due to occipital lobe stroke. Incorrect Options: Option A)Binasal Hemianopia : Shows loss of nasal fields in both eyes. Rare; associated with bilateral glaucoma or carotid artery calcification. The provided field does not show binasal loss. Option B)Bitemporal Hemianopia : Would show loss of both temporal fields. Seen in optic chiasm compression (e.g., pituitary adenoma). The given defect is homonymous, not bitemporal. Option D) Altitudinal Field Defect : Would show loss of upper or lower half of the visual field horizontally. Seen in ischemic optic neuropathy (AION/NAION). The defect here is vertical, not altitudinal.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q31: What are the yoke muscles for Levodepression? (Q858121)

A. LSO and RIR


B. RSO and LIR

C. RIO and LIR


D. LIO and RIR

Correct Answer: B. RSO and LIR

Explanation: - RSO and LIR The Right Superior Oblique (RSO) and the Left Inferior Rectus (LIR) are the yoke muscles involved in levodepression (the downward movement of both eyes when the gaze is directed to the left). Right Superior Oblique (RSO) : The superior oblique muscle in the right eye helps depress the eye when it is abducted (pushed outward). Left Inferior Rectus (LIR) : The inferior rectus muscle in the left eye helps depress the eye. These two muscles coordinate to cause the downward movement of the eyes when looking to the left. Yoke muscles Yoke muscles are pairs of extraocular muscles, one in each eye, that work together to ensure coordinated eye movements in the same direction. These muscles function according to Hering's Law of Equal Innervation , which ensures that corresponding muscles in both eyes receive equal neural impulses for synchronised movement. Hering’s Law of Equal Innervation would mean that the right lateral rectus (yoke muscle) should be balanced with the left medial rectus (vs Sherrington’s law: This law about reciprocal innervation states that increased innervation and contraction of a muscle is automatically associated with reciprocal decrease in innervation and contraction of its antagonist). The yoke muscle pairs facilitate various eye movements: Horizontal Gaze (Left or Right) : Leftward Gaze (Levoversion) : Left Lateral Rectus (abducts the left eye) Right Medial Rectus (adducts the right eye) Rightward Gaze (Dextroversion) : Right Lateral Rectus (abducts the right eye) Left Medial Rectus (adducts the left eye) Vertical Gaze (Up or Down) : Upward Gaze (Elevation) : Right and Left Superior Rectus (elevate the eyes) Right and Left Inferior Oblique (assist in elevation) Downward Gaze (Depression) : Right and Left Inferior Rectus (depress the eyes) Right and Left Superior Oblique (assist in depression) Inward Gaze (Convergence) : Both Medial Rectus muscles contract to bring the eyes inward (toward the nose) during near-vision tasks, such as reading. LSO and RIR (Option A) : Incorrect because LSO (Left Superior Oblique) helps in depression but not for levodepression. RIR (Right Inferior Rectus) is also involved in depression but doesn't pair with LSO for this movement. RIO and LIR (Option C) : RIO (Right Inferior Oblique) assists with elevation, not depression. LIR does help in depression, but they don't work together for levodepression. LIO and RIR (Option D) : LIO (Left Inferior Oblique) assists with elevation, not depression. RIR does help in depression, but this pair is not correct for levodepression.



@Neet_pg_bot



@Neet_pg_bot

References:

Yanoff & Duker Ophthalmology, 5th edition, Pg- 1193,1194

↑ *Back to this question in Part A*

@Neet_pg_bot

Q32: Optic nerve glioma is seen in? (Q860538)

A. NF-1

B. Tuberous sclerosis

C. NF-2

D. Schwannoma

Correct Answer: A. NF-1

Explanation: Optic nerve gliomas usually present with proptosis, visual loss, and optic nerve swelling , but not typically with leukocoria or an intraocular mass with calcification. Various conditions other than retinoblastoma, which present as leukocoria are collectively called pseudoglioma. Differential diagnosis of Retinoblastoma Leukocoria Congenital cataract Inflammatory deposits in vitreous following a plastic cyclitis or choroiditis, Coloboma of the choroid, Retinopathy of prematurity Persistent hyperplastic primary vitreous Toxocara endophthalmitis Exudative retinopathy of Coats Endophytic retinoblastoma Tuberous sclerosis Neurofibromatosis-1 (Option A) Astrocytoma Exophytic Retinoblastoma Exudative retinopathy of Coats Options B, C, and D are incorrect. Optic nerve glioma is not associated with these conditions.

References:

AK Khurana Textbook of Ophthalmology Edition 7th Pg. 315, 316 and 317

↑ *Back to this question in Part A*

@Neet_pg_bot

Q33: Which tuberculosis drug causes impairment of color vision and optic neuritis? (Q860567)

A. Rifampicin

B. Ethambutol

C. Pyrazinamide

D. Streptomycin

Correct Answer: B. Ethambutol

Explanation: Ethambutol is a first-line anti-tubercular drug that causes retrobulbar neuritis, decreased visual acuity and colour blindness. Ethambutol MOA Inhibits arabinosyl transferases involved in arabinogalactan synthesis, thereby interfering with mycolic acid synthesis in cell wall Mechanism of resistance Mutation of embAB genes, which encodes the enzyme arabinosyl transferase, reducing its affinity for ethambutol Uses In the treatment of Mycobacterium tuberculosis Mycobacterium avium complex (MAC) Mycobacterium kansasii Side effects Painless loss of vision Field defects due to retrobulbar neuritis (Option C) Loss of visual acuity Red-green colour blindness Nausea, rashes, fever Option A, C and D does not cause visual impairment.

References:

KDT 8ed Pg 819-820

↑ *Back to this question in Part A*

@Neet_pg_bot

Q34: Which nerve defect causes lagophthalmos? (Q860170)

A. 7th nerve

B. 6th nerve

C. 4th nerve

D. 5th nerve

Correct Answer: A. 7th nerve

Explanation: Lagophthalmos is the inability to close the eye. It is a feature of 7 th nerve palsy. Neuroparalytic keratitis can occur due to seventh nerve palsy, resulting in lagophthalmos from affected orbicularis oculi and leading to exposure keratitis Bell’s palsy: Description Most common cause of facial palsy Characterized by idiopathic, LMN facial paralysis of acute onset Most commonly the Labyrinthine segment of the facial nerve is involved. Males and females are equally affected. Aetiology Idiopathic: Most common cause Viral Infection: HSV >> Herpes zoster Vascular Ischemia Hereditary: Positive family history in 6-8% Autoimmune Risk Factors More common in diabetics (due to angiopathy) and pregnant women (due to fluid retention). Clinical Features Sudden onset of facial paralysis with inability to close the eye (Bell's phenomenon) (Option A) , asymmetry of the face, epiphora, drooping mouth, and dribbling of saliva. Associated symptoms may include ear pain, noise intolerance (stapedial paralysis), or loss of taste (chorda tympani involvement). Diagnosis Diagnosis is made by excluding other causes of peripheral facial paralysis. Treatment General: Reassurance, pain relief, eye protection and physiotherapy Medical: Prednisolone 60 mg per day for 5 days then reduced by 10 mg per day (for a total treatment time of 10 days) and 50 mg per day (in two divided doses) for 10 days, Acyclovir Surgical: Nerve decompression Prognosis 85%-90% of patients fully recover Incomplete resolution are mainly seen in recurrent/late recovery cases Incorrect Options: 6th nerve (Option B) leads to lateral rectus muscle paralysis, causing impaired abduction. 4th nerve (Option C) affects the superior oblique muscle, resulting in vertical diplopia. 5th nerve (Option D) is responsible for facial sensation and corneal reflex, but not eyelid closure.

References:

Diseases of ENT by Dhingra 8th edition, page- 108, 112-114.

Scott Brown's Otorhinolaryngology Vol-2 8th edition, page- 1400.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q35: A patient on antitubercular therapy (ATT) started experiencing visual problems. Which ATT drug is most likely responsible for this adverse effect? (Q860207)

A. Rifampicin

B. Streptomycin

C. Ethambutol

D. Pyrazinamide

Correct Answer: C. Ethambutol

Explanation: Ethambutol is a first-line anti-tubercular drug that causes retrobulbar neuritis, decreased visual acuity and colour blindness. Ethambutol MOA Inhibits arabinosyl transferases involved in arabinogalactan synthesis, thereby interfering with mycolic acid synthesis in cell wall Mechanism of resistance Mutation of embAB genes, which encodes the enzyme arabinosyl transferase, reducing its affinity for ethambutol Uses In the treatment of Mycobacterium tuberculosis Mycobacterium avium complex (MAC) Mycobacterium kansasii Side effects Painless loss of vision Field defects due to retrobulbar neuritis (Option C) Loss of visual acuity Red-green colour blindness Nausea, rashes, fever Option A, B and D does not cause visual impairment.

References:

KDT 8ed Pg 819-820

↑ *Back to this question in Part A*

@Neet_pg_bot

Q36: In the provided image, which nerve is subjected to blockage? (Q857420)

A. Anterior ethmoidal nerve

B. Nasociliary nerve

C. Supratrochlear nerve

D. Long ciliary nerve



Correct Answer: B. Nasociliary nerve

Explanation: The given image represents the nasociliary nerve block. The nasociliary nerve is typically targeted for blockage in procedures involving the ophthalmic region, especially in the context of blocking sensations to the eye and adjacent structures. The nasociliary nerve , as a branch of the ophthalmic division of the trigeminal nerve (V1), plays a central role in sensory innervation to the eye, cornea, and adjacent regions, which is why it is the primary target in many ophthalmic nerve blocks. Nasociliary nerve block The nasociliary nerve is a branch of the ophthalmic division (V1) of the trigeminal nerve (CN V) . It provides sensory innervation to structures such as the cornea , conjunctiva , sclera , ciliary body , and parts of the nose . Indications for Block: Ocular surgery (e.g., cataract surgery, corneal procedures). Pain management in conditions like corneal ulcers , ocular trauma , or sinusitis . Techniques: Injection Site : The needle is typically inserted near the medial canthus of the eye, in the area where the nasociliary nerve is located close to the lacrimal sac . Injection Approach : The needle is advanced medially, just lateral to the nasal bone , and injected near the nerve's location. Anatomy : The nasociliary nerve travels within the orbit , giving off branches like the long ciliary nerves , posterior ethmoidal , and anterior ethmoidal nerves . Procedure: Patient Positioning : The patient is usually seated or supine with the eye to be blocked exposed. Needle Insertion : A 25-30 gauge needle is used, inserted near the medial canthus , directed medially and slightly superiorly. Local Anesthetic : A small amount (0.5-1 mL) of local anesthetic is injected near the nasociliary nerve for effective blockage. Benefits: Provides anesthesia to the cornea , conjunctiva , and other associated structures in the orbit. Helps in treating ocular pain or managing intraocular procedures with minimal discomfort. Complications: Hematoma Inadvertent block of other branches, leading to side effects like lid drooping or ptosis. Anterior ethmoidal nerve (Option A) is a branch of the nasociliary nerve. It primarily provides sensation to the anterior part of the nasal cavity and the skin of the nose. Although it can be blocked in certain

procedures, it is not the primary nerve blocked in typical ophthalmic procedures. Supratrochlear nerve (Option C) is responsible for sensation in the forehead and scalp, specifically in the area above the eye. This nerve is not typically involved in the same procedures as the nasociliary nerve, which innervates the eye and surrounding structures. Long ciliary nerves (Option D) are branches of the nasociliary nerve and are responsible for innervating the cornea, iris, and the ciliary body. They may be involved in certain procedures but are not usually blocked on their own without affecting the nasociliary nerve.



References:

Parson’s Diseases of the Eye, 23rd Edition, Page 1470

Consensus recommendations for anaesthetic peripheral nerve block - ScienceDirect

↑ *Back to this question in Part A*

@Neet_pg_bot

Q37: In a female patient, there is a complaint of vision loss, specifically affecting the right halves of both eyes. Where is the lesion located? (Q855459)

A. Optic nerve


B. Left optic tract


C. Optic chiasma


D. Right optic tract


Correct Answer: B. Left optic tract

Explanation: The lesion described in the question is located in the left optic tract , which explains the vision loss affecting the right halves of both eyes (right homonymous hemianopia). Visual Pathway: Temporal fibers carry impulses from the Nasal field, and Nasal fibres carry the Temporal field. {{caption_text}} Visual Pathway Lesions and Defects Site of Lesion Fibers Affected Field Defect Optic Nerve Ipsilateral Temporal + Nasal Fibers (Option A ruled out) {{caption_text}} Anopia Optic Chiasma Bilateral Nasal Fibers (Option C ruled out) {{caption_text}} Bitemporal Hemianopia (Heteronymous Hemianopia) Optic Tract Lateral Geniculate Body (LGB Lesion) Optic Radiation Ipsilateral Temporal Fibers + Contralateral Nasal Fibers {{caption_text}} Homonymous Hemianopia Visual Cortex without the involvement of tip (Post. Cerebral Artery) Ipsilateral Temporal Fibers + Contralateral Nasal Fibers except macular representation {{caption_text}} Homonymous Hemianopia with Macular Sparing Tip of Visual Cortex (Middle Cerebral Artery) Macular Representation of Ipsilateral Temporal and Contralateral Nasal Fibers. {{caption_text}} Macular Homonymous Hemianopia Aspect Inferior Optic Radiations (Meyer's Loop) Superior Optic Radiations (Baum's Loop) Location of Lesion Temporal lobe Parietal lobe Affected Visual Field Superior quadrants of the contralateral visual field Inferior quadrants of the contralateral visual field Visual Defect Superior homonymous quadrantanopia Inferior homonymous quadrantanopia Associated Symptoms Memory loss Hearing loss Auditory hallucinations Contralateral hemiplegia Hemianesthesia Optokinetic nystagmus {{caption_text}} {{caption_text}} Right Optic Tract (Option D) Effect: Left homonymous hemianopia (left halves of both fields). Vision loss given here is in the right halves, not the left.





@Neet_pg_bot


@Neet_pg_bot




@Neet_pg_bot




@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot



References:

Parsons Diseases of the eye, 23rd Edition, Pg - 1184

Yanoff & Duker Ophthalmology, 5th edition, Pg - 913,919

↑ *Back to this question in Part A*

@Neet_pg_bot

Q38: A 40-year-old male patient presents with altered sensation in the extremities, difficulty with coordination, and speech problems. On examination, it is noted that the pupillary light reaction is absent, but the accommodation reaction is present. Which condition is most likely responsible for this presentation? (Q858115)

A. Neurosyphilis

B. Horner's syndrome


C. Papilledema

D. Optic Neuritis

Correct Answer: A. Neurosyphilis

Explanation: Neurosyphilis is a complication of syphilis that affects the nervous system. It occurs in the tertiary stage of syphilis, typically years after the initial infection. The condition can present with a variety of neurological symptoms, including altered sensation, coordination difficulties, and speech problems, as seen in the patient. Specific features of this case, including absent pupillary light reaction with intact accommodation, are highly

suggestive of Argyll Robertson pupils , which is a classic sign of neurosyphilis . Neurosyphilis and Argyll Robertson Pupils: Pupillary Light Reflex Absent : The pupillary light reflex is mediated by the optic nerve (CN II) and oculomotor nerve (CN III). In neurosyphilis, the optic nerve is often involved, leading to loss of the pupillary light reflex. Accommodation Reflex Present : Despite the absence of the light reflex, the accommodation reflex (the ability of the pupil to constrict when focusing on a near object) remains intact, which is typical in Argyll Robertson pupils . This occurs because the pathways responsible for accommodation (afferent via the optic nerve and efferent via the oculomotor nerve) are not usually affected in neurosyphilis. Other Symptoms: The patient may also present with neurological symptoms such as sensory disturbances, motor coordination problems (ataxia), and speech difficulties, all of which can be part of neurosyphilis. Spirochetal uveitis: Organism: Treponema pallidum Anterior uveitis: It can present as granulomatous or non-granulomatous iritis typically in the secondary syphilis stage Iris roseola - dilated iris vessels that develop into nodules ‘Plastic’ iritis occurs in congenital syphilis and Jarisch–Herxheimer reaction {{caption_text}} Posterior uveitis: It occurs as disseminated, peripheral, or diffuse choroiditis. Treatment: Systemic penicillin Horner's Syndrome (Option B) : This condition is caused by disruption of the sympathetic pathway to the eye and presents with ptosis, miosis, and anhidrosis. It does not cause absent pupillary light reflex or preserved accommodation reflex. Papilledema (Option C) : Papilledema is swelling of the optic disc due to increased intracranial pressure, which can affect vision but would not explain the absence of the light reflex with preserved accommodation. It also doesn't account for the coordination and speech problems. Optic Neuritis (Option D) : Optic neuritis involves inflammation of the optic nerve, which can cause vision changes, but it does not typically cause the combination of absent light reflex with intact accommodation. Also, optic neuritis usually doesn't present with the full range of neurological symptoms seen here.



References:

Parsons diseases of the eye 23 e, PDF pg no 566-70

Kanski’s Clinical ophthalmology, 9th edition, Page 479, 784, 785

The immune response in tubercular uveitis and its implications for treatment: From anti-tubercular treatment to host-directed therapies - ScienceDirect(3. Clinical perspective of tubercular uveitis: From diagnosis to treatment initiation)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q39: A 40-year-old male patient presents with altered sensation in the extremities, difficulty with coordination, and speech problems. On examination, it is noted that the pupillary light reaction is absent, but the accommodation reaction is present. Which condition is most likely responsible for this presentation? (Q820662)

A. Neurosyphilis

B. Horner’s syndrome

C. Papilledema

D. Optic Neuritis

Correct Answer: A. Neurosyphilis

Explanation: Correct Option A - Neurosyphilis: Neurosyphilis is a bacterial infection of the brain or spinal cord. It occurs in people who have had untreated syphilis for many years. One of the signs of neurosyphilis can be Argyll Robertson pupil, where the pupils do not react to light (absent pupillary light reaction) but do react to accommodation (preserved accommodation reaction). Incorrect Options: Option B - Horner’s syndrome: It is charcaterized by miosis, ptosis, and decreased sweating anhidrosis on the affected side of your face. Option C - Papilledema: This is a condition in which increased pressure in or around the brain causes the part of the optic nerve inside the eye to swell. It can cause visual disturbances but doesn't typically result in an absent pupillary light reaction with preserved accommodation reaction. Option D - Optic Neuritis: It presents with Marcus gunn pupil, i.e., paradoxical s=dilatation of both the pupil when light is flashed to the disease eye.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q40: A 45-year-old male arrived at the emergency ward with sudden vision loss and experiencing painful eye movements . Upon examination, the affected eye shows a relative afferent pupillary defect, a normal optic disc , and a central scotoma . What is the likely diagnosis? (Q793017)

A. Optic nerve glioma

B. Optic neuritis

C. Eale’s disease

D. Optic atrophy

Correct Answer: B. Optic neuritis

Explanation: Correct Option B - Optic neuritis: The sudden loss of vision, painful eye movements, relative afferent pupillary defects, and central scotoma are characteristic of optic neuritis. It typically presents with marked sudden vision loss, dark adaptation may be lowered, and decreased contrast sensitivity. On examination RAPD (relative afferent pupillary defect), central scotoma, and centrocecal scotoma can be seen. Incorrect Options: Option A - Optic nerve glioma: Optic nerve glioma is a type of brain tumor that can affect the optic nerve, but it is relatively rare in adults and typically presents with symptoms of optic nerve with proptosis. Option C - Eale’s disease: Eale's disease is a condition that causes inflammation and occlusion of the retinal blood vessels, leading to vision loss and floaters. Option D - Optic atrophy: It is characterized by atrophy of the all damaged nerve fibers with total afferent pupillary defect and loss of perception of light.

References:

Prepladder Notes

↑ *Back to this question in Part A*

@Neet_pg_bot

• [07_Squint _ Strabismus] 015 Squint

« Back to Folder Directory

[07_Squint _ Strabismus] 015 Squint — Quick Jump

Q1 (p. 993)	*Q2 (p. 994)*	*Q3 (p. 995)*	*Q4 (p. 996)*	*Q5 (p. 997)*	*Q6 (p. 998)*
Q7 (p. 1000)	*Q8 (p. 1001)*	*Q9 (p. 1002)*	*Q10 (p. 1002)*	*Q11 (p. 1003)*	*Q12 (p. 1004)*
Q13 (p. 1006)	*Q14 (p. 1007)*	*Q15 (p. 1008)*	*Q16 (p. 1010)*	*Q17 (p. 1013)*	*Q18 (p. 1017)*
Q19 (p. 1019)	*Q20 (p. 1020)*	*Q21 (p. 1023)*	*Q22 (p. 1024)*	*Q23 (p. 1027)*	*Q24 (p. 1027)*
Q25 (p. 1029)	*Q26 (p. 1029)*	*Q27 (p. 1030)*			

Q1: A 3-year-old girl presents with esotropia, which increases on focusing near objects. This issue has progressively worsened over the past few months, and she has difficulty with close visual tasks. A refractive error of +6.50 D has been identified. What is the most likely diagnosis? (Q853529)

A. Accommodative refractive esotropia

B. Accommodative non-refractive esotropia

C. Accommodative mixed esotropia

D. Infantile esotropia

Correct Answer: A. Accommodative refractive esotropia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above scenario is a case of accommodative reflex esotropia, which has a convergent squint with high hypermetropia. Caused by uncorrected hypermetropia . Esotropia worsens on near fixation due to excessive accommodative effort. Improves/corrects with proper hypermetropic glasses. Accommodative non-refractive esotropia (Option B) Caused by high AC/A ratio (accommodative convergence vs accommodation). Esotropia more at near, but not due to refractive error . Accommodative mixed esotropia (Option C) Combination of hypermetropia + high AC/A ratio . Both refractive and non-refractive components present. Infantile esotropia (Option D) Large-angle, constant esotropia, appearing before 6 months of age . Not related to accommodation or refractive error.

References:

Parsons Diseases of the eye, 23rd Edition, Page 979, 980, 982.

Yanoff & Duker Ophthalmology, 5th edition, Page 1212, 1213.

Comprehensive Ophthalmology, A K Khurana, 7th edition, Page 363.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 7-year old child is brought to the clinic by her parents due to concerns about her right eyes appearing misaligned. On examination, her squint remains constant in all directions of gaze. She does not report any diplopia, and there are no limitations in eye movement. What is the most likely diagnosis? (Q853543)

A. Concomitant squint

B. Incomitant squint

C. Paralytic squint


D. Pseudo-strabismus

Correct Answer: A. Concomitant squint

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The squint/strabismus remains constant in all directions of gaze with no associated diplopia, suggesting that it is a case of concomitant squint. Concomitant Squint/Strabismus: Angle of deviation equal in all 9 gazes. No limitation of eye movements, no diplopia initially. Primary deviation = secondary deviation. Common in early childhood/congenital. Adaptations: amblyopia (negative), abnormal retinal correspondence (positive). Incomitant Squint/Strabismus (Option B): Deviation varies in different gazes. Ocular movements restricted, diplopia present. Late presentation; secondary deviation > primary deviation. Paralytic Squint/Strabismus (Option C) Type of incomitant squint due to paralysis of one/more EOMs. History of diplopia, restricted movement toward paralyzed muscle. Head often turned toward action of paralyzed muscle. Secondary deviation > primary deviation. Pseudo-strabismus (Option D) Eyes actually aligned, only appear misaligned. Causes: Pseudo-esotropia → epicanthal folds, high myopia (negative angle kappa) Pseudo-exotropia → telecanthus, hypertelorism, high hypermetropia (positive angle kappa)



References:

Yanoff & Duker Ophthalmology, 5th Edition, Pages 1211, 1217

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 65 year old woman presented to the OPD with her head tilted towards the right side. On examination, she was having left hypertropia which increases on looking right or medially. The muscle which is most likely paralyzed is: (Q853542)

A. Right superior oblique

B. Left superior oblique

C. Right inferior oblique

D. Left inferior oblique

Correct Answer: B. Left superior oblique

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Woman with head tilt to right, left hypertropia and hypertropia increases on right gaze or on left gaze in adduction. These are classic features of left superior oblique (CN IV) palsy. Superior oblique: depresses, intorts, abducts the eye. In paralysis, hypertropia worsens: On gaze towards opposite side (right gaze) → depression by SO needed. On adduction → SO acts as primary depressor; palsy → hypertropia. Head tilt toward opposite shoulder is compensatory to reduce diplopia (Bielschowsky head tilt test). Right superior oblique (Option A) : Palsy would affect right eye, not left hypertropia. Right inferior oblique (Option C): Palsy would reduce elevation in adduction of right eye, not hypertropia in left eye. Left inferior oblique (Option D) : Palsy would reduce elevation in adduction of left eye, but SO palsy presents with hypertropia, not limitation of elevation.

References:

www.researchgate.net/publication/345627720_Keys_to_Solve_Park-3_Step_Test

eyewiki.org/Three_Step_Test_for_Cyclovertical_Muscle_Palsy

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 40-year-old patient presents for binocular vision assessment. The ophthalmologist performs the Worth's Four Dot Test. The patient wears red-green glasses and reports seeing three green dots. What does this finding most likely indicate? (Q853541)

A. Diplopia

B. Suppression of right eye

C. Suppression of left eye


D. Normal binocular vision

Correct Answer: B. Suppression of right eye

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: If the patient only sees three green dots, it suggests suppression of the right eye, as the red lens over the right eye would normally allow them to see the red dot. The fact that only green dots are seen indicates that the right eye is not contributing to vision. WORTH'S FOUR DOT TEST Setup: Red lens over the right eye, green lens over the left eye; patient views 4 lights (1 red, 2 green, 1 white). Interpretation: Normal binocular vision (Option D): Sees all 4 lights; no squint. Abnormal retinal correspondence: Sees all 4 lights despite manifest squint. Right eye suppression: Sees only 3 green lights; right eye not contributing. Left eye suppression (Option C): Sees only 2 red lights; left eye not contributing. Alternating suppression: Sees 3 green and 2 red lights alternately. Diplopia (Option A): Sees 5 lights (2 red + 3 green); double vision present.



References:

Yanoff & Duker Ophthalmology, 5th edition, Pages 1206, 1207

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: During a Maddox rod test, the patient is asked to view a distant light source. With the Maddox rod placed in front of the right eye, the patient reports seeing a vertical red line to the right of the white light. What type of ocular misalignment does this result suggest? (Q853540)

A. Esophoria

B. Exophoria

C. Right hyperphoria


D. Left hyperphoria

Correct Answer: A. Esophoria

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In the Maddox rod test, if the vertical red line (seen by the right eye) appears to the right of the white light, it suggests that the eyes are converging, which is indicative of esophoria . MADDOX ROD TEST Patient is asked to fixate on a point light in the centre of a Maddox tangent scale at a distance of 6 metres. A Maddox rod is placed in front of one eye with the axis of rod parallel to the axis of deviation. The Maddox rod converts the point source into a line. Thus, the patient will see a point light with one eye and a red line with the other. Due to dissimilar images of the two eyes, fusion is broken and heterophoria becomes manifest . Exophoria (Option B): Eye deviates outward → line appears to the left of the light. Right hyperphoria (Option C): Vertical deviation → Maddox rod placed horizontally (line vertical) detects vertical deviations. Not relevant here. Left hyperphoria (Option D): Vertical deviation of left eye → line placement differs; not indicated in this case.



@Neet_pg_bot

References:

smartoptometryacademy.com/maddox-rod-test-all-you-need-to-know/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 45-year-old male presents with eyelid retraction on downgaze. He has a history of ocular nerve injury. Which of the following best describes this condition? (Q853539)

A. Von Graefe’s Sign

B. Hutchinson’s Pupil

C. Pseudo-Von Graefe Sign

D. Marcus Gunn Jaw-Winking Phenomenon

Correct Answer: C. Pseudo-Von Graefe Sign

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Pseudo-Von Graefe Sign is due to aberrant regeneration of the third cranial nerve following a palsy. The co-contraction of the levator palpebrae superioris and inferior rectus muscles leads to eyelid retraction on downgaze , distinguishing it from other conditions. This Pseudo-Von Graefe Sign is not associated with pupil abnormalities, and the pupil typically remains unaffected. Von Graefe’s Sign (Option A) is seen in Graves’ disease, characterized by delayed descent of the upper eyelid during downgaze caused by restrictive orbitopathy affecting the eyelid. Hutchinson’s Pupil (Option B): Typically seen in uncal herniation, characterized by a unilateral dilated pupil that is unresponsive to light due to compression of the third cranial nerve, often associated with raised intracranial pressure. Marcus Gunn Jaw-Winking Phenomenon (Option D) involves ptosis that improves with jaw movement due to aberrant innervation between the trigeminal nerve (motor division for the jaw) and the levator palpebrae superioris.





References:

Parsons Diseases of the eye, 23rd Edition, Page 1032

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 7-year-old girl has limited abduction of the left eye, with globe retraction and narrowing of the palpebral fissure on adduction and widening of the fissure on attempted abduction. What is the most likely diagnosis? (Q853538)

A. Duane Retraction Syndrome Type I

B. Duane Retraction Syndrome Type II

C. Duane Retraction Syndrome Type III

D. Sixth Cranial Nerve Palsy

Correct Answer: A. Duane Retraction Syndrome Type I

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The child presents with: Limited abduction of the left eye Globe retraction and narrowing of the palpebral fissure on adduction Widening of the palpebral fissure on attempted abduction This clinical picture is characteristic of Duane Retraction Syndrome (DRS) Type I , the most common subtype. Duane Retraction Syndrome Type II (Option B): Incorrect because it shows limited adduction with normal abduction, which is the reverse of this case. Duane Retraction Syndrome Type III (Option C): Incorrect as both abduction and adduction are limited, not just abduction. Sixth Cranial Nerve Palsy (Option D): Incorrect because, although there is limited abduction, there is no globe retraction or palpebral fissure change on adduction.

References:

Yanoff & Duker Ophthalmology, 5th edition, Page 1215

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 60-year-old patient presents with the following symptoms: right-sided facial weakness, right-sided lateral gaze impairment, and weakness on the left side of the body. Imaging reveals a ventral pontine lesion. Which of the following syndromes best describes this patient's condition? (Q853537)

A. Gradenigo Syndrome

B. Millard-Gubler Syndrome

C. Moebius Syndrome

D. Foville Syndrome

Correct Answer: B. Millard-Gubler Syndrome

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Given Clinical scenario with facial weakness, lateral gaze impairment & weakness on the left side of the body along with ventral pontine lesion points toward Millard-Gubler Syndrome Syndromes Associated with Abducens Nerve: Gradenigo Syndrome (Option A): Mastoiditis → petrous apex inflammation → VI nerve palsy + facial pain (V nerve) + deafness. Moebius Syndrome (Option C): Congenital → VI & VII palsy, esotropia/exotropia, orofacial dysmorphism, feeding/speech difficulty. Foville Syndrome (Option D): Dorsal pons lesion → VI palsy + VII palsy + contralateral hemiparesis + central Horner.

References:

Yanoff & Duker Ophthalmology, 5th edition, Page 1215

Harrison's Principles of Internal Medicine, 21st Edition, Page 230

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: A 45-year-old man has double vision and an inward deviation of his left eye. On examination it is found to be associated with 6th Cranial Nerve Palsy. What finding is typical for sixth cranial nerve palsy? (Q853536)

A. Vertical diplopia

B. Pupils are affected

C. Head turn towards the same side, avoiding diplopia

D. Crossed Diplopia

Correct Answer: C. Head turn towards the same side, avoiding diplopia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In sixth cranial nerve palsy, the affected eye cannot abduct properly, leading to diplopia that often improves with a head turn towards the affected side. This compensatory mechanism helps align the eyes to avoid double vision. CN VI supplies the lateral rectus → abduction deficit → esotropia (inward eye). Typical clinical features: Horizontal diplopia (not vertical). Diplopia worse on gaze toward the affected side. Compensatory head turn towards the affected side to reduce diplopia. Pupils are not affected (parasympathetics travel with CN III, not VI). Vertical diplopia (Option A) : Seen in CN IV palsy (superior oblique), not CN VI. Pupils are affected (Option B) : Pupillary involvement is a feature of CN III palsy, not CN VI. Crossed Diplopia (Option D) : Crossed diplopia = seen in exotropia (outward deviation), but CN VI palsy causes esotropia → uncrossed diplopia.

References:

Yanoff & Duker Ophthalmology, 5th edition, Pages 933, 934

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: A 45-year-old patient reports diplopia and is found to have left trochlear nerve palsy. Which of the following findings is not typically associated with it? (Q853534)

A. Diplopia relieves while going downstairs

B. No diplopia with head tilt away from the left side

C. Hypertropia of the affected eye


D. Increase in hypertropia on opposite gaze

Correct Answer: A. Diplopia relieves while going downstairs

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In trochlear nerve palsy, there is vertical diplopia which worsens while going downstairs Trochlear Nerve (CN IV) Palsy: Clinical Features: Hypertropia (Option C) Increase in hypertropia on opposite gaze (Option D) Excyclotorsion (outward rotational movement of the eye) Vertical Diplopia - Increases on Downgaze Depressed Chin with face towards the opposite side Head tilt towards the opposite side (Option B) Poor depression in adduction is typical. Crossed Diplopia



References:

Yanoff & Duker Ophthalmology, 5th edition, Pages 932, 934

Parsons Diseases of the eye, 23rd Edition, Page 992

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A 65-year-old patient presents with the following findings. Which of the following is least likely to be associated with this condition? (Q853533)

A. Weber's Syndrome

B. Benedict's Syndrome

C. Nothmagel's Syndrome

D. Brown's Syndrome



Correct Answer: D. Brown's Syndrome

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Brown's Syndrome is a congenital ocular motility disorder caused by the fibrous tightening of the superior oblique tendon, causing a restriction in eye movement due to issues, and is not associated with third cranial nerve (CN III) palsy. Syndromes associated with Third Cranial Nerve Palsy (Oculomotor Nerve): Weber's Syndrome (Option A) Lesion: Midbrain, involving CN III fascicle + corticospinal tract. Features: Ipsilateral CN III palsy + contralateral hemiplegia. Benedikt's Syndrome (Option B) Lesion: Midbrain, involving CN III fascicle + red nucleus. Features: Ipsilateral CN III palsy + contralateral tremor/involuntary movements. Nothnagel's Syndrome (Option C) Lesion: Midbrain, involving CN III fascicle + superior cerebellar peduncle. Features: Ipsilateral CN III palsy + ipsilateral cerebellar ataxia.

References:

Yanoff & Duker Ophthalmology, 5th edition, Page 931, 932

Harrison's Principles of Internal Medicine, 21st Edition, Page 231

AK Khurana Comprehensive Ophthalmology, 7th Edition, Page 372

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 60-year-old man presents with sudden onset double vision and difficulty aligning his eyes, which began three weeks ago. He has a history of hypertension and recently experienced a mild stroke. Examination reveals that his eye is positioned

downward and outward. Which of the following statements about this condition is incorrect? (Q853532)

A. Ptosis is present

B. Mydriasis is present

C. Pupil is fixed

D. Accommodation increases

Correct Answer: D. Accommodation increases

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given scenario is diagnosed as paralytic strabismus, caused by 3rd nerve (oculomotor) palsy in which accommodation is completely lost due to paralysis of the ciliary muscles. Ptosis is present (Option A): Levator palpebrae superioris (CN III) is paralysed → drooping of the eyelid. Mydriasis is present (Option B): If the parasympathetic fibers are involved (especially in compressive lesions), the pupil is dilated. Pupil is fixed (Option C): When parasympathetic fibers are affected the pupil is dilated and poorly reactive (appears fixed).



References:

Parsons Diseases of the eye, 23rd Edition, Pages 1000, 1024, 1027.

Yanoff & Duker Ophthalmology, 5th edition, Pages 938, 968.

Comprehensive Ophthalmology, A K Khurana, 7th edition, Page 369.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: How much is the difference in deviation between upward gaze and downward gaze in A-V pattern strabismus? (Q853531)

A. A-pattern ≥ 10 PD, V-pattern ≥ 15 PD

B. A-pattern ≥ 15 PD, V-pattern ≥ 10 PD

C. A-pattern ≤ 10 PD, V-pattern ≤ 15 PD


D. A-pattern ≤ 15 PD, V-pattern ≤ 10 PD

Correct Answer: A. A-pattern ≥ 10 PD, V-pattern ≥ 15 PD

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: A–V pattern strabismus : Difference in horizontal deviation between upgaze and downgaze. Diagnostic criteria : A-pattern : Difference ≥ 10 prism dioptres (PD) . V-pattern : Difference ≥ 15 PD .



References:

Yanoff & Duker Ophthalmology, 5th edition, Page 1221, 1222.

Comprehensive Ophthalmology, A K Khurana, 7th edition, Pages 371, 372.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A 45-year-old male complains of double vision, describing seeing two overlapping images when looking in a specific direction. He reports that this problem started suddenly about two weeks ago and is most noticeable when looking towards the right side. There is no history of trauma. On examination, he has limited movement of the right eye towards the right. What is the most likely cause of this condition? (Q853530)

A. Blowout fracture

B. Thick pterygium

C. Subluxation of lens


D. Paralytic strabismus

Correct Answer: D. Paralytic strabismus

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given scenario is diagnosed as binocular diplopia most commonly caused by paralysis of extra ocular muscles (paralytic strabismus). Diplopia refers to the simultaneous perception of two images of a single object. Blowout fracture (Option A): Causes mechanical restriction (commonly inferior rectus entrapment → limitation of upgaze). History of trauma would be present. Thick pterygium (Option B) : Can cause restricted ocular motility but is progressive and gradual, not sudden diplopia. Subluxation of lens (Option C) : Produces monocular diplopia (persists even when one eye is closed). Not gaze-dependent, unlike this case.



References:

Parsons Diseases of the eye, 23rd Edition, Page 211.

Yanoff & Duker Ophthalmology, 5th edition, Page 935.

Comprehensive Ophthalmology, A K Khurana, 7th edition, Page 357.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: Which of the following is known for its specific pattern of muscle insertions around the sclera, essential for the alignment and movement of the extraocular muscles? (Q853504)

A. Annulus of Zinn

B. Spiral of Tillaux

C. Scleral spur


D. Tenon’s capsule


Correct Answer: B. Spiral of Tillaux

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Spiral of Tillaux refers to the spiral arrangement of the insertion points of the extraocular muscles on the sclera, which is essential for coordinating eye movements and maintaining proper ocular alignment. Order of Insertion (Spiral of Tillaux): Medial Rectus Inserts closest to the limbus, about 5.5 mm from the limbus. Inferior Rectus Inserts 6.5 mm from the limbus. Lateral Rectus Inserts 6.9 mm from the limbus Superior Rectus Inserts the farthest, about 7.7 mm from the limbus. Annulus of Zinn (Option A) is a fibrous ring at the back of the orbit around the optic nerve. It serves as the origin for the four rectus muscles (lateral, medial, superior, and inferior), providing a stable attachment point that allows these muscles to control eye movement. Scleral spur (Option C) is the part of the sclera in the anterior chamber angle and plays a role in the drainage of aqueous humour. Tenon’s capsule (Option D) is the fascial sheath that surrounds the eye and helps provide structural support.





References:

Yanoff & Duker Ophthalmology, 5th Edition, Pages 1190,1191

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A 10-year-old child is having trouble catching a ball, climbing stairs, and performing fine motor tasks. She has a tendency to avoid activities requiring precise hand skills. No developmental delay was detected. Which of the following tests would be appropriate for diagnosing her condition? (Q853528)

A. HESS chart

B. Cover-uncover test

C. Titmus fly test

D. Maddox rod test


Correct Answer: C. Titmus fly test

■ *Listen (English)*


■ *Listen (Hindi)*


Explanation: The scenario describes a case of stereopsis, and the appropriate test is the Titmus fly test. Stereopsis is the ability to perceive the 3rd dimension (depth perception). Titmus fly test: The patient has to wear polarized glass, and the fly will be seen from different angles. Evaluate how well the child can perceive the relative depth of the images. HESS chart (Option A) is used to measure the degree of deviation, especially if torsional, and particularly to measure any progressive increase or decrease by Hess screen test. Cover-uncover test (Option B) : This test is used to detect manifest (tropia) and latent (phoria) strabismus by covering one eye and then uncovering it to observe any corrective movement. It evaluates ocular misalignment, not depth perception. Maddox rod test (Option D) is used for detection of latent squint.





@Neet_pg_bot





@Neet_pg_bot

References:

Parsons Diseases of the eye, 23rd Edition, Page 257.

Yanoff & Duker Ophthalmology, 5th edition, Page 1208.

Comprehensive Ophthalmology, A K Khurana, 7th edition, Page 355.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q17: The parents of an 8-month-old girl have observed her eyes misaligning, especially when she focuses on objects up close, with one eye occasionally turning inward. They consulted an ophthalmologist, who conducted the test shown below. What is the name of this test? (Q853527)

A. Hirschberg test

B. Bruckner test

C. Prism cover test


D. Krimsky test




Correct Answer: B. Bruckner test


- *Listen (English)*
- *Listen (Hindi)*

Explanation: The above image depicts the Bruckner test , showing a brighter reflex in a squinted eye . Bruckner Test: The examiner sits at arm's length in front of the patient and shines the beam of a direct ophthalmoscope onto the patient's eyes, wide enough to illuminate both eyes simultaneously. Fundus reflex is depicted as, Squinted eye: Brighter reflex Fixing eye (normal eye) : Dark or dull reflex Hirschberg test (Option A) involves throwing light into the eye from a distance and the patient is asked to look at the light (an infant does this reflexively). A rough indication of the angle of the squint can be obtained from the position of the corneal reflex. Prism cover test (Option C): In the prism bar cover test , prisms of increasing strength (apex towards the deviation) are placed before one eye while the patient fixates with the other; the cover–uncover test is repeated until no corrective movement occurs, and the deviation is then measured in prism dioptres Krimsky test (Option D): In this test, the patient fixates on a point light while prisms of increasing power (apex towards the squint) are placed before the normal eye until the corneal light reflex is centered in the squinting eye; the prism power needed corresponds to the squint angle in prism dioptres.


A
B
@Neet_pg_bot


Normal
15° ET
30° ET
45° ET
@Neet_pg_bot





References:

Parsons Diseases of the Eye, 23rd Edition, Page 960, 961

↑ *Back to this question in Part A*

@Neet_pg_bot


Q18: A 10-year-old boy's parents noticed his left eye turning inward, especially when tired or focusing on near objects. An ophthalmologist diagnosed strabismus and performed the test shown below. Which statements about this test are correct? (Q853526)

A. It indicates the angle of squint

B. In an aligned eye, the reflex is centred

C. Reflex at limbus indicates 45° squint

D. All of the above




Correct Answer: D. All of the above


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above image depicts Hirschberg's test, which is used to indicate the angle of squint by measuring the light reflex on the cornea. Hirschberg's Test The angle of the squint can be roughly estimated (Option A) by observing the position of the corneal reflex when light is directed into the eye. Procedure Shine light into both eyes (ophthalmoscope/torch). Patient fixates on distant light source. Findings Fixing eye : Reflex in centre of pupil (Option B). Squinting eye : Reflex decentred. At pupillary border → ~18–20° deviation Between pupillary border

& limbus → ~35° At limbus → ~45° (Option C)





References:

Parsons Diseases of the eye, 23rd Edition, Page 965, 966

Yanoff & Duker Ophthalmology, 5th Edition, Page 1199

A K Khurana Comprehensive Ophthalmology, 7th Edition, Page 365

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: A 32-year-old patient presents with eye strain and headaches. The ophthalmologist plans to perform a series of tests to evaluate the patient's strabismus. All of the following are objective tests used to assess strabismus except? (Q853513)

A. Maddox Rod test

B. Prism Bar Cover test

C. Cover Uncover test


D. Hirschberg test

Correct Answer: A. Maddox Rod test

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Maddox Rod Test is primarily subjective . It requires the patient to interpret the position of lines projected onto their retina, and their perception is used to measure the angle of deviation. This contrasts with purely objective tests, which rely on direct measurements or observations without relying on the patient's subjective interpretation of visual stimuli. Objective Tests Subjective Tests Corneal Light Reflex Test (Hirschberg Test (Option D) , Krimsky Test) Bruckner Test Cover-Uncover Test (Option C) Prism Bar Test (Option B) Synoptophore Maddox Rod Test Maddox Wing Test Lancaster Red-Green Test Hess Screen Haploscopic Devices (Major Amblyoscope)



@Neet_pg_bot

References:

Parsons Diseases of the eye, 23rd Edition, Pages 973, 974

Yanoff & Duker Ophthalmology, 5th edition, Pages 1198, 1199, 1200

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: A 35-year-old patient presents with increased diplopia on upward gaze and restricted movement of the left eye. The patient also has a history of recent orbital trauma. To determine whether the limitation is due to a mechanical restriction or a nerve palsy, which of the following tests is most appropriate? (Q853512)

A. Hess screen test

B. Forced duction test

C. Cover test


D. Worth’s 4-dot test

Correct Answer: B. Forced duction test

■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: Distinguishes between mechanical restriction and neurogenic palsy . Procedure: Examiner passively moves the eyeball with forceps. If resistance → mechanical restriction (e.g., muscle entrapment, dysthyroid orbitopathy). If free movement → nerve palsy . Hess Screen Test (Option A) measures the degree of misalignment but does not differentiate between mechanical restriction and nerve palsy directly. Cover Test (Option C) is useful for detecting and measuring deviations but not for determining the cause of limited movement. Worth's 4-Dot Test (Option D) assesses binocular vision and suppression but does not provide specific information about mechanical restrictions or nerve palsy.



@Neet_pg_bot

HESS SCREEN



Green in front of left eye


Green in front of right eye

Name _____


Date _____

Notes _____

@Neet_pg_bot



@Neet_pg_bot



References:

Parsons Diseases of the eye, 23rd Edition, Pages 964, 965

Yanoff & Duker Ophthalmology, 5th edition, Page 1201

↑ *Back to this question in Part A*

@Neet_pg_bot


Q21: A 10-year-old patient presents to the clinic with complaints of occasional double vision and eye strain during schoolwork. On examination, you suspect she may have a squint and perform the following test. What is this test known as? (Q853511)

A. Maddox Rod Test

B. Synoptophore

C. Cover-Uncover Test

D. Maddox Wing Test



Correct Answer: C. Cover-Uncover Test

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given image is of Cover-Uncover Test which is used to differentiate between latent squint (heterophoria) and manifest squint (heterotopia). Cover–Uncover Test Purpose : Differentiate latent squint (heterophoria) from manifest squint (heterotropia) . Covers horizontal & vertical deviations (not torsional). Cover Test Detects tropias (manifest squint) . Procedure: Cover one eye → observe uncovered eye for movement. Nasal movement: Exotropia Temporal movement: Esotropia Upward movement: Hypotropia Downward movement: Hypertropia Deviation quantified with prism cover test . Uncover Test Detects phorias (latent squint) . Cover one eye → remove → if eye moves to refixate → phoria/intermittent tropia. Alternate cover test maximizes dissociation; prisms quantify. Other Tests Maddox Rod (Option A) : Detects/quantifies heterophoria ; not for latent vs manifest distinction. Synoptophore (Option B) : Measures squint angle, binocular vision, fusion; not for phoria vs tropia. Maddox Wing (Option D) : Measures eso/exophoria ; mainly for latent deviations.

References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 1199,1200

↑ *Back to this question in Part A*

@Neet_pg_bot


Q22: Which of the following options correctly identifies the type of strabismus depicted in each image? 1. 2. 3. (Q853510)


A. 1- Esotropia; 2-Exotropia; 3-Hypotropia

B. 1- Esotropia; 2-Hypertropia; 3- Exotropia

C. 1- Hypertropia; 2-Exotropia; 3-Hypotropia

D. 1- Hypertropia; 2-Esotropia; 3-Exotropia






Correct Answer: D. 1- Hypertropia; 2-Esotropia; 3-Exotropia

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Squint / Strabismus: A condition where the visual axes of both eyes do not meet at a point (Misalignment of Visual axes) Esotropia : Inward misalignment of the eyes. (Common in Hypermetropes) Exotropia : Outward misalignment of the eyes. (Common in Myopes) Hypertropia : Upward misalignment of one eye relative to the other. Hypotropia: Downward misalignment of one eye relative to the other.



References:

Parsons Diseases of the eye, 23rd Edition, Pages 955, 957, 960, 963, 971

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Which mechanism is responsible for maintaining coordinated movement of both eyes and achieving binocular vision? (Q853509)

A. Conjugate and Dysconjugate movements

B. Synergistic and Antagonistic muscles

C. Yoke muscles

D. All of the above

Correct Answer: D. All of the above

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: All of the above mechanisms are responsible for maintaining coordinated movement of both eyes and achieving binocular vision Mechanism of Binocular Vision Formation: Binocular vision relies on the precise coordination of extraocular muscles in both eyes, ensuring single, clear vision through complex neural mechanisms. Conjugate & Dysconjugate Movements (Option A) Conjugate : Both eyes move in the same direction (e.g., right gaze). Dysconjugate : Eyes move in opposite directions (e.g., convergence/divergence). Synergistic & Antagonistic Muscles (Option B) Synergists : Work together for a movement (e.g., right LR + left MR in right gaze). Antagonists : Oppose the movement (e.g., right MR + left LR in right gaze). Yoke Muscles (Option C) Paired synergists in opposite eyes for conjugate gaze. Example: Right LR + Left MR in rightward gaze.

References:

Parsons Diseases of the eye, 23rd Edition, Pages 931, 933

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: A 32-year-old patient presents with diplopia and difficulty looking upwards and to the right, with limited elevation of the right eye. You suspect a lesion affecting a muscle responsible for vertical movement. Which axis describes the rotational movement, and which muscle is likely affected? (Q853508)

A. Fick's x-axis; Right Inferior Rectus

B. Fick's y-axis; Right Medial Rectus

C. Fick's z-axis; Right Inferior Oblique


D. Fick's x-axis; Right Superior Rectus

Correct Answer: D. Fick's x-axis; Right Superior Rectus

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The vertical movements of the eye occur with the x axis as the center of rotation in primary position. The right superior rectus elevates the eye, particularly in the direction of upward gaze. A lesion in the right superior rectus explains the patient's difficulty with upward movement. Fick's x-axis; Right Inferior Rectus (Option A) Inferior rectus causes depression , not elevation. Wrong muscle for this presentation. Fick's y-axis; Right Medial Rectus (Option B) Fick’s y-axis = torsional movements (intorsion/extorsion). Medial rectus does adduction only , no vertical action. Fick's z-axis; Right Inferior Oblique (Option C) Fick’s z-axis = horizontal movements (adduction/abduction). Inferior oblique elevates in adduction , not abduction. Wrong axis.



References:

Yanoff & Duker Ophthalmology, 5th edition, Pages 927,928

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: Which physiological principle explains that during eye movement, an increase in neural stimulation to a contracting muscle is accompanied by a proportional decrease in stimulation to its opposing muscle? (Q853507)

A. Hering’s Law

B. Sherrington’s Law

C. Henle’s Law

D. Donder’s Law

Correct Answer: B. Sherrington’s Law

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Sherrington’s Law of Reciprocal Innervation of the same eye, states that when one muscle contracts (agonist), its opposing muscle (antagonist) in the same eye must relax to allow coordinated movement. Hering’s Law (Option A) refers to the equal innervation of corresponding eye muscles to ensure simultaneous and symmetrical movements. Henle’s law of accommodation (Option C) refers to anatomical observations related to the length-tension relationship in muscles, particularly in the context of accommodation in the eye. Donder’s Law (Option D) pertains to the torsional movements of the eye.

References:

Yanoff & Duker Ophthalmology, 5th edition, Page 1194

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: A 25-year-old patient presents with a condition in which, upon examination, the left eye fails to adduct during right gaze. The patient reports diplopia, which is more noticeable when looking to the right. Which extraocular muscle is most likely overacting in this patient's condition? (Q853506)

A. Right Medial Rectus

B. Left Lateral Rectus

C. Right Lateral Rectus


D. Left Medial Rectus

Correct Answer: C. Right Lateral Rectus

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Given that the left eye fails to adduct and diplopia is more noticeable during right gaze, it suggests that the left medial rectus is not functioning properly. In INO, the ipsilateral medial rectus fails to adduct due to MLF lesion. The contralateral lateral rectus becomes overacting , often with nystagmus, as it tries to abduct unopposed. Here, left medial rectus fails → right lateral rectus shows overaction. Right Medial Rectus (Option A) Responsible for adduction of right eye (important in left gaze). Not overacting in right gaze. Left Lateral Rectus (Option B) Abducts the left eye in left gaze . Not involved in right gaze → not overacting here. Left Medial Rectus (Option D) This is the underacting muscle in INO, not the overacting one.



References:

Yanoff & Duker Ophthalmology, 5th Edition, Pages 1193, 1194

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: A 45-year-old patient presents with vertical diplopia that worsens when looking down, especially while walking downstairs. Examination reveals weakness in intorsion, depression, and abduction of the affected eye. Which muscle and cranial nerve are most likely affected? (Q853505)

A. Superior Rectus; CN III

B. Inferior Oblique; CN III

C. Superior Oblique; CN IV

D. Inferior Rectus; CN IV

Correct Answer: C. Superior Oblique; CN IV

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The superior oblique muscle , innervated by the trochlear nerve (CN IV) , is responsible for intorsion, depression, and abduction of the eye. Dysfunction of this muscle causes vertical diplopia, particularly noticeable when looking down, such as when walking downstairs. Extraocular muscles and actions Muscle Primary Action Secondary Actions Innervation Medial Rectus Adduction (moves eye inward) None Oculomotor nerve (CN III) Lateral Rectus Abduction (moves eye outward) None Abducens nerve (CN VI) Superior Rectus Elevation (moves eye upward) Intorsion (inward rotation), Adduction Oculomotor nerve (CN III) Inferior Rectus Depression (moves eye downward) Extorsion (outward rotation), Adduction Oculomotor nerve (CN III) Superior Oblique Intorsion (inward rotation) Depression, Abduction Trochlear nerve (CN IV) Inferior Oblique Extorsion (outward rotation) Elevation, Abduction Oculomotor nerve (CN III)

References:

Yanoff & Duker Ophthalmology, 5th edition, Pages 1191, 1192

↑ *Back to this question in Part A*

@Neet_pg_bot

• [07_Squint _ Strabismus] 016 Previous Year Questions

« Back to Folder Directory

[07_Squint _ Strabismus] 016 Previous Year Questions — Quick Jump

Q1 (p. 1032)	*Q2 (p. 1032)*	*Q3 (p. 1033)*	*Q4 (p. 1035)*	*Q5 (p. 1036)*	*Q6 (p. 1039)*
Q7 (p. 1040)	*Q8 (p. 1041)*	*Q9 (p. 1042)*	*Q10 (p. 1043)*	*Q11 (p. 1045)*	*Q12 (p. 1046)*
Q13 (p. 1047)	*Q14 (p. 1047)*	*Q15 (p. 1049)*	*Q16 (p. 1050)*	*Q17 (p. 1050)*	*Q18 (p. 1051)*

Q1: What is the function of the superior oblique muscle? (Q855209)

A. Intorsion, abduction, depression

B. Intorsion, adduction, elevation

C. Extorsion, abduction, depression

D. Extorsion, adduction, elevation

Correct Answer: A. Intorsion, abduction, depression

Explanation: The superior oblique muscle , innervated by the trochlear nerve (CN IV) , is responsible for intorsion, depression, and abduction of the eye. Dysfunction of this muscle causes vertical diplopia, particularly noticeable when looking down, such as when walking downstairs. Extraocular muscles and actions Muscle Primary Action Secondary Actions Innervation Medial Rectus Adduction (moves eye inward) None Oculomotor nerve (CN III) Lateral Rectus Abduction (moves eye outward) None Abducens nerve (CN VI) Superior Rectus Elevation (moves eye upward) (Option B ruled out) Intorsion (inward rotation), Adduction Oculomotor nerve (CN III) Inferior Rectus Depression (moves eye downward) Extorsion (outward rotation), Adduction Oculomotor nerve (CN III) Superior Oblique Intorsion (inward rotation) Depression, Abduction Trochlear nerve (CN IV) Inferior Oblique Extorsion (outward rotation) (Option C & D ruled out) Elevation, Abduction Oculomotor nerve (CN III)

References:

Yanoff & Duker Ophthalmology, 5th edition, Pg- 1191,1192

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: In a 40-year-old patient with diabetes, a squint was observed. The secondary deviation was higher than the primary deviation during examination. The forced duction test yielded negative results. Which type of squint is expected to be

predominant in this individual? (Q858509)

A. Comitant squint

B. Restrictive squint

C. Paralytic squint

D. Pseudosquint

Correct Answer: C. Paralytic squint

Explanation: The clinical scenario of secondary deviation was higher than the primary deviation during examination and forced duction test yielding negative results is suggestive of paralytic squint Squint/Strabismus: Misalignment of the visual axis. This condition is also called Heterotropia. Classification: It is divided into latent squint (phorias) and manifest (tropia). Latent squint - classification Hyperphoria Exophoria Esophoria Manifest squint - classification Esotropia/Exotropia/Hypertropia Concomitant/Incomitant squint Concomitant: Esotropia/Exotropia Incomitant: Paralytic (paralysis of the muscles) Restrictive squint (fibrosis of the muscles) A-V pattern or pattern squint Ocular motility defects Concomitant squint (Option A): It's a type of manifest squint in which the angle of deviation remains equal in all the 9 gazes ; there is no associated limitation of ocular movements . Primary deviation is equal to secondary deviation . Primary deviation is the amount of deviation in a squinted eye. Secondary deviation: It is the amount of deviation in the normal eye (behind the cover). Concomitant squint commonly occurs during early childhood or can be congenital (congenital squint). Eye movements are normal and no double vision (diplopia). Amblyopia develops later (negative adaptation). Abnormal retinal correspondence also plays a role. It is a positive adaptation. Incomitant squint (Option B): This condition is just the opposite of concomitant squints, where the degree of squint varies in different aspects. Secondary deviation is higher than primary deviation in paralytic squint. In contrast to concomitant squints, there is a late presentation. There is palsy of other muscles, which causes limited ocular movements (ocular movements are restricted). Diplopia is also a very important feature of inconcomitant squints. Paralytic squint (Option C): Paralytic squint is a type of incomitant squint in which ocular deviation results from complete or incomplete paralysis of one or more extraocular muscles. History of diplopia and confusion Secondary deviation is greater than primary deviation. Ocular movements are restricted towards the action of paralyzed muscle. Head is turned towards the action of paralyzed muscle. Pseudo-strabismus (Option D): Pseudostrabismus is a condition when the eyes are well aligned but appear to be misaligned. Causes of Pseudo-esotropia: Epicanthal folds High myopia (high negative angle kappa) Causes of Pseudo-Exotropia: Telecanthus or hypertelorism High hypermetropia (high positive angle kappa) Restrictive squint (Option B) occurs when there is a limited range of movement of the affected eye due to restrictions in extraocular muscle function or mechanical factors (e.g., orbital trauma or thyroid eye disease) and would yield a positive forced duction test.

References:

AK Khurana 7th edition page 583, 584

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: The movement lost on left gaze as depicted in the image is seen in? (Q855030)

A. 3rd CN palsy

B. 4th CN palsy

C. 6th CN palsy

D. 7th CN palsy



Correct Answer: A. 3rd CN palsy

Explanation: The image depicts limited adduction of the right eye due to weakness of medial rectus muscle seen in 3rd nerve palsy. The oculomotor nerve is responsible for innervating several extraocular muscles, including the medial rectus, superior rectus, inferior rectus, and inferior oblique muscles. Nerves of the eye: Optic nerve (CN II) Purely sensory Senses the incoming light and image displayed on retina Oculomotor nerve (CN III) Innervates majority of the extraocular muscles Motor innervation to the superior rectus muscle, medial rectus muscle, inferior rectus muscle, inferior oblique muscle, levator palpebrae superioris muscle, ciliary muscle, and the sphincter muscle. Trochlear nerve (CN IV) Motor innervation to the superior oblique muscle Trigeminal Nerve (CN V) Ophthalmic branch of the trigeminal nerve provides sensory innervation to the eye Afferent part of corneal and lacrimation reflex Abducens Nerve (CN VI) Motor innervation to the lateral rectus muscle Facial Nerve (CN VII) Motor innervation of the orbicularis oculi muscle Efferent part of corneal and lacrimation reflex Sympathetic Nervous System Long ciliary nerves: Innervates the pupillary dilator muscles causing mydriasis Parasympathetic nervous system Short ciliary nerves: Innervates the sphincter pupillae muscles causing miosis Incorrect Options: 6 th CN palsy (Option C) This affects lateral rectus muscle and inability to abduct the eye towards the side of gaze. 4 th CN palsy (Option B) This affects the superior oblique muscle, leading to difficulty in looking downward, particularly in an adducted position (e.g., difficulty descending stairs). It does not impact horizontal gaze. 7 th CN palsy (Option D) This nerve innervates facial muscles, not extraocular muscles.

References:

Anatomy, Head and Neck: Eye Nerves - StatPearls - NCBI Bookshelf (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: Which of the following conditions is associated with the finding observed by a physician during an eye examination of a small child? (Q855023)

A. Myopia

B. Astigmatism

C. Hyperopia

D. Emmetropia



Correct Answer: C. Hyperopia

Explanation: The image shows convergent squint (esotropia) — the eyes are deviated inward , a common finding in children with uncorrected hyperopia (farsightedness) . Mechanism: In hyperopia , light focuses behind the retina , so the child must use accommodation (focusing effort) to bring objects into clear focus. Accommodation is linked with convergence (the inward turning of the eyes). Excessive accommodation therefore triggers excessive convergence , leading to accommodative esotropia . This condition is especially common in young children and is corrected with convex (+) lenses , which reduce the need for accommodation. Incorrect Options A) Myopia: Causes divergence (exotropia) rather than convergence. B) Astigmatism: Causes blurred vision due to irregular corneal curvature, not eye deviation D) Emmetropia: Normal refractive state; no squint develops.

References:

Parsons Diseases of the eye, 23rd Edition, Pg - 955,957,960,963,971

↑ *Back to this question in Part A*

@Neet_pg_bot


Q5: The chart depicted in the image is _____? (Q855387)

A. Hess chart

B. Snellen chart

C. Pelli-Robson chart


D. ETDRS chart




@Neet_pg_bot


Correct Answer: A. Hess chart


Explanation: The image in the question shows Hess chart, used for measuring degree of Malalignment in Squint/Starbismus , but it cannot be used to differentiate between Paralytic and Restrictive squint. {{caption_text}} Incorrect options: Snellen Chart (Option B): This is the traditional eye examination tool used worldwide for assessing Visual acuity . {{caption_text}} Pelli-Robson Chart (Option C): This chart is a specific tool used to measure Contrast sensitivity . {{caption_text}} ETDRS Chart (Option D): Standing for Early Treatment Diabetic Retinopathy Study , (also known as the LogMAR chart) this chart is a standardized tool used for testing visual acuity more precisely than the Snellen chart, particularly useful in research settings and monitoring progression of diseases like diabetic retinopathy. {{caption_text}}



@Neet_pg_bot


E
F P
T O Z
L P E D
P E C F D
E D F C Z P
F E L O P Z D
D E F P O T E C
L E F O D P C T
F D P L T C E O
P E Z O L C F T D
1 20/200
2 20/100
3 20/70
4 20/50
5 20/40
6 20/30
7 20/25
8 20/20
9
10
11
@Neet_pg_bot





References:

Kanski’s Clinical Ophthalmology, 9th edition, Pages 2-4,721.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q6: Identify the condition (Q855339)

A. Medial rectus palsy

B. Trochlear nerve palsy


C. Abducens nerve palsy

D. Oculomotor nerve palsy



Correct Answer: D. Oculomotor nerve palsy

Explanation: Given image is suggestive of Paralytic Strabismus of a patient with down and out left eye with ptosis on primary gaze with inability to elevate, adduct completely is suggestive of Oculomotor (Third) nerve palsy. {{caption_text}} Features of third nerve palsy Ptosis: Due to paralysis of the LPS (levator palpebrae superioris) muscle The eyeball is deviated down and outwards (due to actions of the lateral rectus and superior oblique muscles). The pupil is fixed and dilated (due to paralysis of the sphincter pupillae muscle) Ocular movements are restricted Accommodation is completely lost (due to paralysis of the ciliary muscle) Crossed diplopia is elicited by raising the eyelid. Absence of Both direct and consensual reflexes in the affected eye These compensatory head postures are motor adaptations used to minimise the effects of double vision Horizontal Diplopia: The patient often tilts the head downward Vertical Diplopia: The head is positioned with the chin either raised or lowered Torsional Diplopia: The head is tilted sideways Incorrect Options: Medial Rectus Palsy (Option A): This condition is characterized by weakness or paralysis of the medial rectus muscle, which controls the eye's inward movement, often leading to an inability to ADDUCT the eye. Trochlear Nerve Palsy (Option B): This pertains to a neurological disorder where damage to the fourth cranial nerve, or trochlear nerve, results in vertical diplopia when the patient looks down and inability to DEPRESS the eye completely . Abducens Nerve Palsy (Option C): This is a disorder marked by the impairment of the sixth cranial nerve, resulting in lateral gaze palsy.



References:

Parsons Diseases of the Eye 23rd Edition, Pages 1000,1024,1027.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: What is the likely diagnosis for a 5-year-old girl presenting with decreased best corrected visual acuity in the right eye after a school vision screening revealed diminution of acuity in the right eye without any examination findings of the eye and visual pathway? (Q857227)

A. Amblyopia

B. Retinoblastoma

C. Congenital cataract

D. Brain lesion

Correct Answer: A. Amblyopia

Explanation: Decrease in best corrected visual acuity in one or both eyes with no apparent pathology of the eye and visual pathway is suggestive of Amblyopia. Amblyopia: Definition, Types, and Clinical Characteristics: Definition: Partial, reversible vision loss in one or both eyes. Absence of organic diseases of l ocular media, retina, visual pathway The eye appears normal, but vision is impaired. Types of Amblyopia: Type Cause Key Features Strabismic Amblyopia The brain suppresses input from one eye due to constant squint (strabismus). Common cause; underdevelopment of vision in the deviated eye. Stimulus Deprivation Amblyopia Hindered vision due to obstruction from early life (e.g., congenital cataracts, ptosis, dense central corneal opacity). Lack of visual stimulation affects normal development. Anisometropic Amblyopia A significant difference in refractive error between eyes. The brain favours the clearer eye, leading to amblyopia in the other. Ametropic Amblyopia High uncorrected refractive errors in both eyes. Bilateral; is less severe compared to other types. Meridional Amblyopia Uncorrected astigmatism affecting specific visual meridians. Affects one or both eyes depending on the

astigmatism degree. Clinical Characteristics: Reduced Visual Acuity: Affects recognition acuity more than resolution acuity. Graded by severity: Mild: 20/25 to 20/60 Moderate: 20/70 to 20/100 Severe: 20/200 or worse Neutral Density Filter Test: Slight improvement in visual acuity through a filter (unlike organic eye diseases). Crowding Phenomenon: Difficulty identifying letters on crowded charts (e.g., Snellen chart). Fixation Pattern: Eccentric fixation (focusing away from the fovea) indicates severity. Color Vision: Usually normal but affected in severe cases (vision below 6/36). Other ocular examination: usually NO abnormalities are found (Option A) Brain lesion (Option D): A lesion in the visual pathway can result in loss of vision but no such examination findings is mentioned. Retinoblastoma (Option B): Retinoblastoma shows leukocoria which is absent here. Congenital cataract (Option C): Loss of red reflex is seen in congenital cataract.

References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 718-719

Amblyopia - PMC

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: What is the recommended treatment for a 3-year-old child with amblyopia in the right eye? (Q857188)

A. Observe for 3 months

B. Occlusion of the right eye

C. Occlusion of the left eye

D. Surgical correction

Correct Answer: C. Occlusion of the left eye

Explanation: For treating amblyopia, it is typically recommended to occlude the stronger eye (in this case, the left eye) for 4-6 hours per day. Treatment of amblyopia: Amblyopia treatment should be started as early as possible (younger the child, better the prognosis). Amblyopia therapy works best when initiated in young children under 3 years of age and comprise: Occlusion therapy (Option C): Occlusion of the sound (normal) eye to force use of amblyopic eye is the main stay in the treatment of amblyopia. Following methods are deployed, Full-time occlusion: Up to 2 years, the occlusion should be done in 2:1, i.e. 2 days in sound eye and one day in amblyopic eye; at the age of 3 years, 3:1; at the age of 4 years, 4:1; At the age of 5 years, 5:1; and after the age of 6 years 6:1. Part-time occlusion: 4-6 hours/day, as shown by Amblyopia Treatment Study (ATS), and is as effective as full-time occlusion. Maintenance occlusion: After vision equalization, part-time occlusion continues until at least 8 years of age. Ensure no media opacity or refractive errors before starting occlusion therapy. Other Methods: Penalization: Blurring vision in the sound eye with atropine or over-plus lenses. Computerized Home Vision Therapy: Uses specially designed software as an adjunct to occlusion therapy. Pharmacologic Manipulation Levodopa/Carbidopa: Studied as an adjunct to occlusion therapy. Other agents: Citicoline and omega fatty acids

Pleoptics: Pleoptic exercises are done to improve foveal fixation, though less commonly used now.

References:

Kanski’s clinical ophthalmology 10th edition pages 718,719.

Amblyopia - PMC

↑ *Back to this question in Part A*

@Neet_pg_bot


Q9: A 60-year-old male presents with left eyelid drooping and experiencing double vision. Upon examination, it was observed that the pupil in the affected eye was dilated. The following eye movement examination was conducted. Which nerve is most likely injured in this case? (Q857182)

A. Optic nerve

B. Oculomotor

C. Trochlear


D. Abducens



Correct Answer: B. Oculomotor

Explanation: Given image is suggestive of Paralytic Strabismus of a patient with down and out left eye with ptosis on primary gaze with diplopia, inability to elevate, adduct completely is suggestive of Oculomotor (Third) nerve palsy . Features of third nerve palsy Ptosis: Due to paralysis of the LPS (levator palpebrae superioris) muscle The eyeball is deviated down and outwards (due to actions of the lateral rectus and superior oblique muscles). The pupil is fixed and dilated (due to paralysis of the sphincter pupillae muscle) Ocular movements are restricted Accommodation is completely lost (due to paralysis of the ciliary muscle) Crossed diplopia is elicited by raising the eyelid. Absence of Both direct and consensual reflexes in the affected eye These compensatory head postures are motor adaptations used to minimise the effects of double vision Horizontal Diplopia: The patient often tilts the

head downward Vertical Diplopia: The head is positioned with the chin either raised or lowered Torsional Diplopia: The head is tilted sideways Optic nerve injury (Option A): Injury to the optic nerve would cause visual disturbances. Trochlear Nerve injury (Option C): This pertains to a neurological disorder where damage to the fourth cranial nerve, or trochlear nerve, results in vertical diplopia when the patient looks down and inability to DEPRESS the eye completely . Abducens Nerve injury (Option D): This is a disorder marked by the impairment of the sixth cranial nerve, resulting in lateral gaze palsy.



References:

Parsons Diseases of the Eye 23rd Edition, Pages 1000,1024,1027.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: The cover-uncover test is performed to diagnose various eye conditions. Which of the following is not detected by this test? (Q852497)

A. Latent Squint

B. Manifest Squint

C. Lateral Nystagmus


D. Amblyopia

Correct Answer: D. Amblyopia

Explanation: Amblyopia, often called "lazy eye," is a condition where the vision in one eye is reduced because the brain favors the other eye. It can be caused by strabismus, but the cover-uncover test itself does not diagnose amblyopia Cover-Uncover Test is used to differentiate between latent squint (heterophoria) and manifest squint (heterotopia). Cover test: Detects Heterotropia (Manifest Squint) (Option B) Uncover test: Detects Heterophoria (Latent Squint) (Option A) Cover-Uncover Test: The cover and uncover tests assess horizontal and vertical strabismus, but not torsional deviations. They also evaluate the ability of each eye to fixate on a target. Cover Test: Detects Tropias: Measures constant deviations of the visual axis. Procedure: Cover one eye with a paddle, thumb,

or occluder. Observe the uncovered eye for movement: Nasal movement: Esotropia Temporal movement: Exotropia Upward movement: Hypertropia Downward movement: Hypotropia For young children, use small toys instead of bright lights. Quantify the deviation with the simultaneous prism and cover test by introducing a prism before one eye until movement stops. Repeat with the prism before the other eye. Uncover Test: Identifies latent deviations or tropias. Procedure: Cover one eye and then observe the covered eye when the cover is removed. If the eye reverts to fixation, it indicates a phoria or intermittent tropia . If the eye remains deviated, it suggests a tropia with fixation preference for the fellow eye. Alternate Cover Test: Detects phorias by dissociating the visual axes. Alternate covering each eye to maximize visual axis dissociation. Care must be taken to permit time for each eye to reside behind the cover (the cover must not be “fanned” before the eyes). Use prisms to quantify phorias or tropias. If measurements are unclear, estimate the midpoint between defined movements. Lateral Nystagmus (Option C): Nystagmus is involuntary, rhythmic eye movements. The cover-uncover test might exacerbate nystagmus, though it is not the primary test for detecting it.





@Neet_pg_bot

References:

Yanoff & Duker Ophthalmology, 5th edition, Pg - 1199,1200

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: What is the recommended treatment for a 3-year-old child with amblyopia in the right eye? (Q858154)

A. Observation

B. Spectacles or refraction with a near addition

C. Occlusion Therapy

D. Radial keratotomy

Correct Answer: C. Occlusion Therapy

Explanation: ll waking hours or all but one hour. Schedule by Age: Up to 2 years: 2:1 (2 days occluded, 1 day not) 3 years: 3:1 4 years: 4:1 5 years: 5:1 6 years and older: 6:1 Part-time occlusion: 4-6 hours/day , as shown by

Amblyopia Treatment Study (ATS), and is as effective as full-time occlusion. Maintenance occlusion: After vision equalization, part-time occlusion continues until at least 8 years of age. Ensure no media opacity or refractive errors before starting occlusion therapy. Other Methods Penalization : Blurring vision in the sound eye with atropine or over-plus lenses. Computerized Home Vision Therapy : Uses specially designed software as an adjunct to occlusion therapy. Pharmacologic Manipulation Levodopa/Carbidopa : Studied as an adjunct to occlusion therapy. Other agents : Citicoline and omega fatty acids Pleoptics: Pleoptic exercises are done to improve foveal fixation, though less commonly used now. Observation (Option A) : Passive observation is not effective for amblyopia, as early intervention is key to improving visual outcomes. Spectacles or refraction with a near addition (Option B) : These are useful for refractive errors (e.g., hyperopia) but do not directly address amblyopia unless coupled with occlusion therapy. Radial keratotomy (Option D) : This is an outdated surgical procedure for correcting refractive errors, not for treating amblyopia.

References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 718-719

Amblyopia - PMC

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: Which drug is commonly used for refraction in a child with accommodative squint? (Q858120)

A. 1% Atropine

B. Cyclopentolate

C. Tropicamide

D. Homatropine

Correct Answer: A. 1% Atropine

Explanation: ycloplegics can be used in any age group if active accommodation is present. Commonly used cycloplegics: Atropine: used in < 5 years age Homatropine: 5-25 years (Option D ruled out) Phenylephrine (Mydriatic) in elderly Cyclopentolate (Option B) : Often used in children for cycloplegia but is not as commonly used as atropine for long-term treatment or in younger children. Tropicamide (Option C) : A shorter-acting agent, used primarily for pupil dilation rather than cycloplegia.

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no:42

Objective Refraction Technique: Retinoscopy - PubMed

Retinoscopes: Past and present - PMC

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: Which nerve innervates the lateral rectus muscle? (Q860162)

A. Sixth nerve

B. Seventh nerve

C. Third nerve

D. Fourth nerve

Correct Answer: A. Sixth nerve

Explanation: Nerves of the eye: Optic nerve (CN II) Purely sensory Senses the incoming light and image displayed on retina Oculomotor nerve (CN III) (Option C) Innervates majority of the extraocular muscles Motor innervation to the superior rectus muscle, medial rectus muscle, inferior rectus muscle, inferior oblique muscle, levator palpebrae superioris muscle, ciliary muscle, and the sphincter muscle. Trochlear nerve (CN IV) (Option D) Motor innervation to the superior oblique muscle Trigeminal Nerve (CN V) Ophthalmic branch of the trigeminal nerve provides sensory innervation to the eye Afferent part of corneal and lacrimation reflex Abducens Nerve (CN VI) (Option A) Motor innervation to the lateral rectus muscle Facial Nerve (CN VII) Motor innervation of the orbicularis oculi muscle Efferent part of corneal and lacrimation reflex Sympathetic Nervous System Long ciliary nerves: Innervates the pupillary dilator muscles causing mydriasis Parasympathetic nervous system Short ciliary nerves: Innervates the sphincter pupillae muscles causing miosis 7 th CN palsy (Option D) This nerve innervates facial muscles, not extraocular muscles.

References:

Anatomy, Head and Neck: Eye Nerves - StatPearls - NCBI Bookshelf (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: The patient has an infection of the cavernous sinus, involving the nerve located on its medial side. To confirm the involvement of the abducens nerve, you will ask the patient to look in which direction? (Q860509)

A. Superior


B. Inferior


C. Medial

D. Lateral

Correct Answer: D. Lateral

Explanation: Abducens nerve supplied lateral rectus muscle which is involved in abduction movement. Therefore patient is asked to look in lateral direction Abducens Nerve (cranial nerve VI) Type Motor Origin Pons (lower border, opposite the pyramid of the medulla). Course Functional component General somatic efferent fibres : Arise from the abducent nucleu s in the pons and supply the lateral rectus muscle of the eyeball. General somatic afferent fibres carry proprioceptive sensations from the lateral rectus and terminate in the mesencephalic nucleus of the trigeminal nerve. Supply Lateral rectus muscle of the eyeball Clinical aspect Lesion or injury (elevated intracranial pressure) of the abducens nerve results in paralysis of the lateral rectus muscle, leading to: Inability to abduct the eye (Option B) Diplopia (double vision) Convergent squint (due to the unopposed action of medial rectus) Option A, B and C are incorrect.





References:

Grey's anatomy 42nd edition, pg 782.

BD Chaurasia’s Human Anatomy, 8th edition, pg 73, 74.

Vishram Singh Textbook of Anatomy, 2nd edition, pg 337, 338.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: What is the probable cause of uniocular diplopia in a young boy who has experienced trauma to his right eye? (Q793357)

A. Paresis of extraocular muscle

B. Displacement of one eyeball

C. Symblepharon

D. Iridodialysis

Correct Answer: D. Iridodialysis

Explanation: Correct Option D - Iridodialysis: The most likely cause of uniocular diplopia (persistence of diplopia even when the normal eye is closed) in a young boy with trauma to the right eye is iridodialysis. Iridodialysis refers to a separation or tearing of the iris from its ciliary attachment, which can cause the pupil to become distorted and result in double vision. Other causes of uniocular diplopia may include Subluxated clear lens or intraocular lens, incipient cataract, and keratoconus. Incorrect Options: Options A, B, and C are incorrect. These conditions causes binocular diplopia.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A 50-year-old female patient visits the ophthalmology clinic with complaints of experiencing vertical diplopia while walking downstairs. Upon further inquiry, it is noted that the patient tends to tilt her head to one side in order to alleviate the diplopia. Which cranial nerve palsy should be suspected in this patient? (Q820672)

A. 3rd nerve

B. 6th nerve

C. 7th nerve

D. 4th nerve

Correct Answer: D. 4th nerve

Explanation: Correct Option D - 4th nerve: The vertical diplopia in the absence of ptosis, combined with a characteristic head posture, suggests fourth nerve disease. The fourth cranial nerve controls the superior oblique muscle. The main actions of the superior oblique are intorsion and depression in adduction. A compensatory head posture avoids diplopia. To compensate for the weakness of intorsion, there is a contralateral head tilt. To alleviate the weakened depression of the eye, the chin is slightly depressed Peripheral lesions cause ipsilateral and nuclear lesions contralateral superior oblique weakness. Incorrect Options: Option A - Third nerve: Third nerve cranial palsy presents with ptosis , abduction and depression in the primary position,, dilated pupil, and limited adduction, elevation and depression . Option B - 6 CN Palsy (Abducens Nerve Palsy): Sixth nerve palsy presents with inability to move the eye laterally and could cause horizontal diplopia. Option C - 7 CN Palsy (Facial Nerve Palsy): The seventh cranial nerve controls the muscles of facial expression and is responsible for taste sensation on the anterior two-thirds of the tongue.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: Which drug is commonly used for refraction in a child with accommodative squint? (Q820650)

A. 1% Atropine

B. Cyclopentolate

C. Tropicamide

D. Homatropine

Correct Answer: A. 1% Atropine

Explanation: Correct Option A - 1% Atropine: High hypermetropia of around +4 to +7 D causes accommodative squint due to excessive accommodation. Correction of hypermetropia results in full correction of convergent squint. Atropine is often used in pediatric ophthalmology to measure hypermetropia. It works by dilating the pupil and temporarily paralyzing accommodation, or the eye's ability to adjust its focus to see nearby objects clearly. Incorrect Options: Options B, C, and D are incorrect. 1% Atropine is used to measure hypermetropia accurately.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: What are the yoke muscles for Levodepression? (Q820639)

A. LSO and RIR


B. RSO and LIR

C. RIO and LIR

D. LIO and RIR

Correct Answer: B. RSO and LIR

Explanation: .



References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [08_Cornea] 017 Cornea- I

« Back to Folder Directory

[08_Cornea] 017 Cornea- I — Quick Jump

Q1 (p. 1053)	*Q2 (p. 1055)*	*Q3 (p. 1063)*	*Q4 (p. 1063)*	*Q5 (p. 1065)*	*Q6 (p. 1068)*
Q7 (p. 1068)	*Q8 (p. 1069)*	*Q9 (p. 1070)*	*Q10 (p. 1073)*	*Q11 (p. 1075)*	*Q12 (p. 1076)*
Q13 (p. 1081)	*Q14 (p. 1084)*	*Q15 (p. 1087)*	*Q16 (p. 1089)*	*Q17 (p. 1090)*	*Q18 (p. 1094)*
Q19 (p. 1095)	*Q20 (p. 1096)*	*Q21 (p. 1097)*	*Q22 (p. 1097)*	*Q23 (p. 1098)*	*Q24 (p. 1101)*
Q25 (p. 1104)	*Q26 (p. 1105)*	*Q27 (p. 1106)*	*Q28 (p. 1107)*	*Q29 (p. 1108)*	*Q30 (p. 1110)*
Q31 (p. 1111)	*Q32 (p. 1112)*	*Q33 (p. 1113)*	*Q34 (p. 1115)*	*Q35 (p. 1116)*	*Q36 (p. 1118)*
Q37 (p. 1118)	*Q38 (p. 1119)*	*Q39 (p. 1120)*	*Q40 (p. 1121)*	*Q41 (p. 1121)*	*Q42 (p. 1123)*
Q43 (p. 1124)	*Q44 (p. 1125)*				


Q1: A 45 y/o patient presents with worsening vision, mild pain, and redness 6 months after Penetrating Keratoplasty. Examination reveals the following. Which feature is most indicative of endothelial rejection? {{caption_text}} (Q853776)

A. Increased Flare and Cells

B. Kayes dots

C. Khodadoust Line


D. Epithelial Rejection Line



Correct Answer: C. Khodadoust Line

- *Listen (English)*
- *Listen (Hindi)*

Explanation: The most specific sign of endothelial rejection after keratoplasty is Khodadoust’s line. It is a line of immune cells seen on the endothelium. Represents: Demarcation between healthy and damaged endothelium. Types of corneal graft rejection: Epithelial Rejection Stromal Rejection Endothelial Rejection (most common and most likely to lead to graft failure) {{caption_text}} Increased flare and cells (Option A): Not specific to endothelial rejection. Kaye’s dots (Option B) : Specific for epithelial rejection. Appear as small epithelial opacities. Epithelial rejection line (Option D): Indicates epithelial graft rejection, not endothelial. Note: Subepithelial rejection : Presents as Krachmer spots (subepithelial infiltrates).



References:

Kanski's Clinical Ophthalmology, 9th Edition, Pages 278, 279.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q2: Match the Following: 1) a) Bacterial Keratitis 2) b) Superficial Punctate Keratitis 3) c) Pseudodendritis 4) d) Dendritic Ulcer 5) e) Nummular Keratitis 6) f) Fungal Keratitis (Q853763)


A. 1-b; 2-e; 3-c; 4-f; 5-d; 6-a


B. 1-c; 2-e; 3-d; 4-f; 5-a; 6-b

C. 1-e; 2-c; 3-b; 4-f; 5-d; 6-a


D. 1-c; 2-e; 3-b; 4-f; 5-d; 6-a



@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot






Correct Answer: D. 1-c; 2-e; 3-b; 4-f; 5-d; 6-a


■ *Listen (English)*


■ *Listen (Hindi)*


Explanation: Image Description Bacterial Keratitis: {{caption_text}} M/C worldwide: Staph aureus M/C organism in India: Staph epidermidis Neovascularization of cornea- Pannus with keratitis A corneal epithelial defect with circumcorneal injection, often with a hypopyon. Clinical features: Pain, redness, photophobia Blepharospasm, discharge, diminution of vision Fungal Keratitis: {{caption_text}} Grey or yellow-white stromal infiltrate with indistinct fluffy margins Unsterile hypopyon Superficial Punctate Keratitis: {{caption_text}} Punctate lesions Seen in herpes simplex and herpes zoster Nummular Keratitis: {{caption_text}} Characterized by fine granular subepithelial deposits surrounded by a halo of stromal haze. Seen in zoster and adenoviral infections. Dendritic Ulcer: Also known as Viral keratitis Linear branching ulcer with terminal bulbs Pseudodendritis: Corneal epithelial lesions that are smaller and finer than herpes simplex dendrites, have tapered ends without terminal bulbs. Seen in Herpes zoster Acanthamoeba keratitis Rare infection Occurs in soft contact lens users, washing lenses with tap water Swimming while wearing contact lenses without goggles M/C infection after contact lens use: Pseudomonas Clinical features: Pain (Disproportionately more) Due to perineural invasion Lead to radial perineuritis - Pathognomic feature Can lead to thickening of corneal nerves Redness, photophobia, blepharospasm Discharge, diminution of vision On examination: Ring lesion: Can be a/w abscess formation Pseudo-dendrites or micro-dendrites Limbitis with stromal infiltration





@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot





References:

Kanski's Clinical Ophthalmology, 9th Edition, Pages 205, 211, 216, 221, 227.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 28-year-old female contact lens wearer with severe eye pain and blurred vision shows a ring-shaped corneal infiltrate, and radial keratoneuritis on examination. Corneal scraping showed the presence of cysts. What is the most appropriate initial treatment? (Q853768)

A. Topical moxifloxacin

B. Oral acyclovir

C. Topical polyhexamethylene biguanide (PHMB)

D. Topical corticosteroids

Correct Answer: C. Topical polyhexamethylene biguanide (PHMB)

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical features and findings point to Acanthamoeba keratitis. PHMB is an amoebicidal agent effective against both the trophozoites and cysts of Acanthamoeba. It is the treatment of choice for Acanthamoeba keratitis, especially when cysts are identified. Treatment of Acanthamoeba Keratitis: Debridement : Removal of infected epithelium to enhance topical medication penetration. Topical Amoebicides : PHMB (polyhexamethylene biguanide) 0.02% and Chlorhexidine 0.02% : Effective against both trophozoites and cysts. Hexamidine : More effective than Propamidine (Brolene) . Topical Steroids : Considered for persistent inflammation after two weeks of anti-amoebic treatment. Caution against use if corneal thinning is present. Pain Management : Oral NSAIDs are recommended for pain relief. Therapeutic Keratoplasty : Required for cases resistant to medical therapy or when corneal perforation occurs. Topical moxifloxacin (Option A): Moxifloxacin is an antibiotic used to treat bacterial keratitis. Oral acyclovir (Option B): Acyclovir is an antiviral agent used primarily for herpes simplex keratitis. Topical corticosteroids (Option D): While corticosteroids may be used cautiously for persistent inflammation after initiating anti-amoebic treatment, they are not first-line therapy and should be avoided initially due to the risk of worsening the infection if used prematurely.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 232, 233, 234

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 23 y/o male presents with B/L eye redness, pain, and severe blurring of vision. He has a history of congenital syphilis. O/E, there is a characteristic pinkish-red

discoloration of the cornea with deep stromal vascularization. Which of the following findings is most likely associated with this patient's condition during the healing stage? (Q853769)

A. Pseudodendrites

B. Ghost vessels

C. Ring-shaped abscess


D. Radial keratoneuritis

Correct Answer: B. Ghost vessels

- *Listen (English)*
- *Listen (Hindi)*

Explanation: s: The given features with characteristic pinkish-red discoloration of the cornea with deep stromal vascularization describe a salmon patch of Syphilitic Interstitial Keratitis Ghost vessels are remnants of previously filled blood vessels that become empty after the active inflammation of syphilitic interstitial keratitis resolves. They are a characteristic finding during the healing stage of the condition. {{caption_text}} {{caption_text}} Pseudodendrites (Option A) are typically associated with herpetic keratitis, not with interstitial keratitis caused by syphilis. Ring-shaped abscess (Option C): Ring-shaped abscesses are seen in advanced Acanthamoeba keratitis, not in syphilitic interstitial keratitis. Radial keratoneuritis (Option D) is a hallmark feature of Acanthamoeba keratitis and is not seen in syphilitic interstitial keratitis.





References:

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 229, 230, 232

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 19 y/o female presents with a history of progressive worsening of vision that is not fully corrected by glasses. After evaluation, she is diagnosed with keratoconus. Which of the following examination findings are seen in keratoconus? Munson's sign Scissor reflex Vogt striae Fleischer ring (Q853770)

A. 1,2,3,4

B. 1,3,4

C. 2,3,4


D. 1,4

Correct Answer: A. 1,2,3,4


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: All of the above signs are features of keratoconus. Munson’s Sign Definition: V-shaped deformity of lower lid on down gaze. Mechanism: Conical cornea pushes the lower lid forward in a conical manner. Seen in advanced keratoconus. {{caption_text}} Scissor’s Reflex Seen on retinoscopy. Reflex appears as two blades of a scissor moving against each other. Due to irregular astigmatism in keratoconus. Vogt’s Striae Vertical, parallel lines in deep stroma/Descemet’s membrane. Orientation: Parallel to steep axis of the cornea. They disappear on applying pressure to the globe. {{caption_text}} Fleischer’s Ring Iron deposition at the base of the cone. Layer involved: Epithelium. Appears as a yellow-brown to greenish ring on slit-lamp. {{caption_text}} Note Fleischer’s Ring vs Kayser-Fleischer Ring: Fleischer’s = iron deposition in epithelium (keratoconus). Kayser-Fleischer’s = copper deposition in Descemet’s membrane (Wilson’s disease). Treatment of Keratoconus: Spectacles / Toric soft contact lens: For cylindrical correction of astigmatism Choice of contact lens: Rigid gas permeable / Semi soft contact lens - for irregular cornea TOC: In progressive case: C3R (Corneal collagen cross linking) with riboflavin Dresden protocol Instill riboflavin eye drops every 5 minutes, continuing for 30 min Expose to Ultraviolet A rays for another 30 min (continuing riboflavin application) Keratoplasty: Penetrating keratoplasty: best prognosis in non-inflammatory conditions DALK (Deep anterior lamellar keratoplasty) For displaced / eccentric cones and non-progressive cases: INTACS (Intra corneal segments)







References:

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 248, 249.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Question: A 22 y/o male with a history of worsening vision despite wearing rigid gas-permeable lenses now complains of increased visual distortion. O/E, his corneal topography shows progressive ectasia, and he is advised to consider treatment options to halt disease progression. Which of the following treatments is most appropriate to stabilize the condition and improve contact lens tolerance in this patient? (Q853771)

A. Spectacle correction

B. Intracorneal ring segments (INTACS)

C. Corneal collagen cross-linking (CXL)

D. Penetrating keratoplasty

Correct Answer: C. Corneal collagen cross-linking (CXL)

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient has keratoconus showing progressive ectasia. Corneal collagen cross-linking (CXL) with riboflavin is the most appropriate treatment for stabilizing keratoconus progression and improving contact lens tolerance in this patient. It works by strengthening the corneal stroma through the cross-linking of collagen fibers, which helps to halt the progression of ectasia, thereby preserving vision and enhancing the effectiveness of contact lenses. Treatment of Keratoconus Spectacle Correction (Option A) : Effective in the early stages, but contact lenses are more beneficial. Less effective in later stages due to irregular astigmatism. Contact Lenses : Rigid gas permeable lenses improve vision in early cases. Specially designed scleral lenses (e.g., Rose-K) are useful in early to moderate cases. Corneal Collagen Cross-Linking (CXL) : New treatment modality to slow disease progression. Involves riboflavin (0.1%) eye drops and UVA radiation. Keratoplasty (Option D) Intracorneal Ring Segments (INTACS) (Option B) : Effective in early to moderate cases by flattening the cornea. Less effective in halting progression.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 249, 250.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 50 y/o patient presents for a corneal transplant evaluation. The patient has a history of systemic infections, including tuberculosis, and is seeking a corneal graft for a therapeutic indication. Which of the following is an absolute contraindication for cornea donation in this patient? (Q853773)

A. Death due to an unknown cause

B. Previous history of ocular surgery

C. Ocular inflammation

D. Retinoblastoma

Correct Answer: A. Death due to an unknown cause

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Death due to an unknown cause : This is an absolute contraindication as it could signal serious undiagnosed conditions. Contraindications for cornea donation Absolute contraindications Relative contraindications 1) Death due to an unknown cause 2) Systemic infections HIV Hepatitis Congenital rubella TB Syphilis 3) CNS infection or diseases Multiple sclerosis Rabies Creutzfeldt-Jakob disease 4) Most haematological malignancies Leukaemia Lymphoma Myeloma and other myeloproliferative disorders and syndromes Ocular inflammation (Option C) Any congenital or acquired disorders of the Eye or previous ocular surgery (including corneal laser surgery) (Option B) Retinoblastoma (Option D) Malignant tumours or ocular metastases in the anterior segment of the eye.

References:

Contraindications

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 276, 277

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A donor cornea is harvested and needs to be preserved for a planned corneal transplant for 10 days. Which of the following storage methods is most appropriate for preserving the donor cornea? (Q853774)

A. Moist Chamber Method

B. McCarey Kaufman (M-K) Medium

C. Optisol GS

D. Cryopreservation

Correct Answer: C. Optisol GS

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Optisol GS is designed for intermediate-term storage, effective for up to 2 weeks, making it suitable for preserving the cornea for up to 10 days. Storage of Cornea: Short-term Storage (24 to 96 hours) Intermediate-term Storage (up to 2 weeks) Long-term Preservation (months to years) : Moist Chamber Method (Option A) : Preserve the whole globe at 4°C in a refrigerator for up to 48 hours. McCarey Kaufman (M-K) Medium (Option B) : Pink-colored medium containing TC-199, Dextran-40, HEPES buffer, and gentamicin. Optisol GS : Contains chondroitin sulfate, dextran, ascorbic acid, vitamin B12, ATP precursors, gentamicin, and streptomycin. Other Media : Cornisol, Eusol, and Life-GE medium also available for up to 2 weeks. Glycerine Preservation : Effective for 3-4 weeks. Organ Culture Method : Preserves cornea at 30°C to 37°C for up to 35 days. Cryopreservation (Option D) : Preserves corneal button at -70°C for up to 1 year. Note: Enucleation (removal of the eyeball) for corneal donation should ideally be done within 6 hours of death to ensure the best quality tissue for transplantation.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 276.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Which of the following is not an indication of Penetrating Keratoplasty (PK)? (Q853775)

A. Corneal opacity

B. Cosmetic improvement of the eye

C. Advanced keratoconus

D. Secondary glaucoma


Correct Answer: D. Secondary glaucoma

■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: Secondary glaucoma is not an indication of PK but rather an early complication of the procedure. PK addresses corneal diseases and issues directly, not intraocular pressure problems or secondary glaucoma. Penetrating Keratoplasty (PK): It is a type of Allografting (Allo-Keratoplasty) which is the replacement of a

patient’s diseased cornea with a healthy donor cornea. Full-thickness grafting. Indications for Penetrating Keratoplasty: Optical Corneal opacity (Option A) Bullous keratopathy Corneal dystrophies Advanced keratoconus (Option C) Therapeutic To replace an inflamed cornea that does not respond to conventional therapy Tectonic Graft To restore the integrity of the eyeball, e.g., after corneal perforation or in cases of significant corneal thinning. Cosmetic To improve the appearance of the eye (Option B) {{caption_text}} {{caption_text}}







References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 277.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Which layer of the cornea is primarily responsible for maintaining its transparency by regulating fluid content? (Q853607)

A. Stroma

B. Endothelium

C. Epithelium

D. Bowman's Membrane


Correct Answer: B. Endothelium


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The innermost layer of the cornea is the endothelium which consists of a single layer of hexagonal cells. These cells are crucial for maintaining corneal transparency by regulating fluid balance within the cornea through an active pump mechanism. Corneal transparency results from multiple structural and physiological factors acting together. These include stromal collagen architecture, precise hydration control by the endothelium, an intact epithelium and tear film, avascularity, and absence of myelin in corneal nerves. Layers of Cornea Epithelium (Option C ruled out) The outermost layer comprises 5-6 layers of non-keratinized, stratified squamous epithelial cells. Acts as a barrier, protecting the cornea and preventing tear fluid from entering the underlying stroma. The basal cells at the limbus are stem cells responsible for regenerating the corneal epithelium. Bowman’s Membrane (Option D ruled out) Located just beneath the epithelium, this acellular layer is 12 µm thick and composed of densely packed collagen fibrils. It is a protective barrier to the stroma but cannot regenerate if damaged and heals by scarring. Stroma (Option A ruled out) Making up about 90% of the cornea's thickness (approximately 0.5 mm thick), the stroma consists of collagen fibrils arranged in parallel layers. Nearly uniform interfibrillar spacing and ordered fibril organization minimize light scatter, allowing destructive interference of scattered light and thus maintaining clarity. These fibrils are embedded in a hydrated matrix of proteoglycans - Chondroitin and Keratan Sulphate (keratan sulphate maintains corneal transparency). Pre-Descemet's Membrane (Dua's Layer) This layer, discovered in 2013, is about 15 µm thick, acellular, and lies just above Descemet's membrane. It is strong and resistant to air penetration. Descemet's Membrane A robust basement membrane of the corneal endothelium is made of collagen and glycoproteins. It is tense, but when torn, curls inwards on itself. It can regenerate if damaged and is resistant to trauma and pathological processes. Endothelium (Option B) The innermost layer consists of a single layer of flat, hexagonal cells. These cells are crucial for maintaining corneal transparency by regulating fluid balance within the cornea through an active pump mechanism. Using ion transporters (notably Na■/K■-ATPase and associated bicarbonate systems), it pumps fluid from the stroma into the aqueous, maintaining a relatively dehydrated stroma and preventing edema and loss of transparency . {{caption_text}} {{caption_text}} The pre-corneal tear film creates a smooth, optically regular anterior refracting surface over the corneal epithelium. This smooth, well-distributed layer fills in minor epithelial irregularities, reduces surface scatter, and provides an optically polished interface for incoming light. The cornea is normally

avascular, with nutrition coming from tears, aqueous humor, and perilimbal vessels. Absence of blood vessels avoids hemoglobin, cellular elements, and vessel walls in the optical path, reducing scatter and helping keep the tissue clear . Corneal nerves lose their myelin sheaths as they enter the cornea, remaining non-myelinated within the transparent zone. Lack of myelin (which is opaque) and fine caliber of these fibers reduce light scatter, helping preserve transparency while still allowing high corneal sensitivity.





References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 204.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: All of the following statements about lamellar keratoplasty are true except: (Q853777)

A. It is indicated in pseudophakic bullous keratopathy

B. It has lower risk of graft rejection compared to Penetrating keratoplasty

C. Deep Anterior Lamellar Keratoplasty (DALK) replaces 80-90% of the anterior stroma and is suitable when the endothelium is normal.


D. Descemet's Stripping Automated Endothelial Keratoplasty (DSAEK) is performed when the endothelium is defective.

Correct Answer: A. It is indicated in pseudophakic bullous keratopathy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In pseudophakic bullous keratopathy, the primary pathology is in the endothelium. Management: Penetrating keratoplasty (PK) or endothelial keratoplasty (like DSAEK/DMEK) is indicated. Lamellar keratoplasty is NOT useful in this condition. It has a lower risk of graft rejection compared to penetrating keratoplasty (Option B): Especially in Deep Anterior Lamellar Keratoplasty (DALK) → because the recipient's own endothelium is preserved. Endothelial graft rejection (the most serious type) is avoided. Deep anterior lamellar keratoplasty replaces 80–90% of the anterior stroma in situ when the endothelium is normal (Option C): Indicated in stromal diseases where the endothelium remains healthy. Descemet's Stripping Automated Endothelial Keratoplasty (DSAEK) is performed when the endothelium is defective (Option D): Involves removal of endothelium + Descemet's membrane, sometimes with a thin layer of posterior stroma.



References:

Kanski's Clinical Ophthalmology, 9th Edition, Pages 279, 280, 283.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q12: Match the following: 1) Arcus senilis a) 2) Band shaped Keratopatthy b) {{caption_text}} 3) Vortex keratopathy c) 4) Salzmann nodular degeneration d) {{caption_text}} (Q853791)

A. 1-b, 2-d, 3-a, 4-c


B. 1-d, 2-a, 3-c, 4-b


C. 1-c, 2-d, 3-b, 4-a

D. 1-a, 2-b, 3-c, 4-d


@Neet_pg_bot


@Neet_pg_bot






Correct Answer: A. 1-b, 2-d, 3-a, 4-c


■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: CORNEAL DEGENERATION IMAGE Arcus Senilis Cause: Lipoid infiltration, common in elderly Appearance: Grey/white crescent → complete ring; thicker sup/inf; sharp periphery, fades centrally; ≤1 mm wide Effect: No impact on vision/vitality Variants: Arcus juvenilis in young → linked to dyslipidemia {{caption_text}} Band Shaped Keratopathy Common in chronic uveitis, aphakic eyes with prolonged silicone oil, and occasionally linked to hyperparathyroidism or vitamin D poisoning. Characterized by a whitish band across the cornea with round ‘holes’ or cleaves. Caused by hyaline infiltration and deposition of calcareous salts in the superficial stroma. Treatment options include scraping off the opacity, dissolving it with sodium edetate, or using an excimer laser. {{caption_text}} Vortex keratopathy Vortex keratopathy , also known as corneal verticillata. It is a distinctive pattern of corneal epithelial deposits that appear as whorl-like (vortex) lines swirling outward toward the periphery of the cornea. Etiology: Drugs: Chloroquine, Amiodarone (dose related), Tamoxifen, Indomethacin, Netarsudil. Fabry disease Salzmann nodular degeneration A degenerative condition of the cornea. Bluish-white, avascular nodules. Located in the superficial stroma and Bowman’s membrane. Typically occurs in individuals with a history of previous corneal disease.




@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology, 9th Edition, Pages 261, 262, 263, 264, 265.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: Which of the following statements about corneal degeneration is incorrect? (Q853792)

A. Salzmann nodular degeneration affects the superficial stroma and Bowman's membrane.

B. Vortex keratopathy is generally asymptomatic and associated with indomethacin intake.

C. Lipid keratopathy involves both intracellular and extracellular fat deposits.


D. Vortex keratopathy remains permanent even after discontinuing the offending medication.


Correct Answer: D. Vortex keratopathy remains permanent even after discontinuing the offending medication.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The condition of vortex keratopathy is reversible upon stopping the offending medication. CORNEAL DEGENERATION IMAGE Vortex Keratopathy / Cornea verticillata Appearance: Whorl-like corneal epithelial opacity Causes: Drugs (amiodarone, chloroquine, phenothiazines, netarsudil, indomethacin) or Fabry disease (Option B) Symptoms: Usually asymptomatic; may cause glare/surface discomfort Reversibility: Reversible after stopping drug (Option D) Treatment: Lubricants for surface discomfort {{caption_text}} Lipid Keratopathy Deposits: Whitish/yellowish cholesterol & fatty acids in cornea Primary: Rare, normal serum lipids Secondary: With corneal vascularization (infections, interstitial keratitis, trauma, glaucoma, iridocyclitis) Fat deposits: Intracellular → extracellular with stromal necrosis (Option C) {{caption_text}} Salzmann nodular degeneration Type: Degenerative corneal condition Nodules: Bluish-white, avascular Location: Superficial stroma + Bowman's membrane (Option A) Associations: Prior corneal disease Course: Slowly progressive Treatment: Lamellar keratoplasty {{caption_text}}







References:

Kanski's Clinical Ophthalmology, 9th Edition, Pages 264, 265.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q14: A 45 y/o male presents with progressively worsening vision and evident discomfort in his left eye over the past year. O/E, the following findings were observed, which caused significant blurring of his vision. {{caption_text}} Which of the following corneal opacities is most likely causing the greatest reduction in his vision? (Q853793)

A. Nebular corneal opacity

B. Macular corneal opacity

C. Leucomatous corneal opacity

D. Adherent leucoma




Correct Answer: A. Nebular corneal opacity


■ *Listen (English)*


■ *Listen (Hindi)*


Explanation: The image shows a faint opacity covering the central cornea, which suggests a Nebular opacity. Although faint, a nebular opacity can severely impact vision, while also causing maximum discomfort (by inducing irregular astigmatism) The irregular refraction caused by the nebula distorts the image more than a denser opacity like leucoma, leading to greater visual discomfort and blurring. {{caption_text}} Macular corneal opacity

(Option B): This semi-dense opacity affects the cornea less than a nebula regarding irregular refraction, but it still causes significant visual impairment. {{caption_text}} Leucomatous corneal opacity (Option C): While this dense white opacity blocks light more completely, it causes less irregular refraction than a nebula. If it doesn't cover the entire pupillary area, it may not affect vision as much as a nebula. {{caption_text}} Adherent leucoma (Option D): This is a dense opacity associated with iris incarceration and is typically more significant in light blockage, but its effect on vision varies depending on the extent of iris involvement. {{caption_text}} Management of corneal opacity : Optical Iridectomy: This procedure can be used for central macular or leucomatous opacities when vision improves with pupil dilation. Phototherapeutic Keratectomy (PTK): PTK uses an excimer laser, and is effective for treating superficial corneal opacities, such as nebulae. Keratoplasty Cosmetic-Coloured Contact Lenses: These lenses offer a cosmetic solution for unsightly scars in the eyes without visual potential. Corneal Tattooing: It is mainly reserved for firm scars in eyes with no useful vision.




@Neet_pg_bot


@Neet_pg_bot



References:

https://courseware.cutm.ac.in/wp-content/uploads/2020/06/CORNEAL-OPACITIES.pdf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 25-year-old patient presents with a history of gradual vision loss. O/E, multiple milky, granular deposits are seen in the anterior corneal stroma, with clear cornea between the deposits. Which type of stromal corneal dystrophy is most likely? (Q853812)

A. Granular Corneal Dystrophy, Type 1 (GCD1)

B. Lattice Corneal Dystrophy, Type 1 (LCD1)

C. Macular Corneal Dystrophy (MCD)


D. Schnyder Corneal Dystrophy (SCD)


Correct Answer: A. Granular Corneal Dystrophy, Type 1 (GCD1)

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Multiple milky, granular deposits in the anterior corneal stroma, with a clear cornea between the deposits, indicate Granular corneal dystrophy Type 1. Corneal dystrophies are bilateral, non-inflammatory, inherited disorders grouped by primary layer affected:■ Anterior (Epithelial/Bowman's) : Epithelial basement membrane, Meesmann, Reis-Bucklers. Stromal : Granular, lattice, macular, Avellino, Schnyder, congenital stromal. Posterior (Endothelial) : Fuchs, CHED, PPCD. TGFB1 Corneal Dystrophies These dystrophies are associated with mutations in the TGFB1 gene Lattice Corneal Dystrophy (LCD)(Option B) LCD – Corneal dystrophy with amyloid deposits forming lattice lines in stroma. LCD1: Early onset, central lattice lines, recurrent erosions → visual loss. LCD2 (Gelsolin type): Later onset, peripheral lattice lines spreading centrally. {{caption_text}} Granular Corneal Dystrophy (GCD) GCD – Hyaline deposits in corneal stroma. GCD1: Childhood onset, milky granular anterior stromal deposits. GCD2: Early onset, mixed granular + lattice deposits. {{caption_text}} Other Stromal Dystrophies Macular Corneal Dystrophy (MCD) (Option C) Inheritance: Autosomal recessive Features: Diffuse stromal haze + macular opacities Onset: Childhood / early adulthood Effect: Severe visual impairment Schnyder Corneal Dystrophy (SCD) (Option D) Inheritance: Autosomal dominant Deposits: Cholesterol crystals in stroma Features: Ring-shaped corneal opacities Effect: Progressive vision loss





References:

Kanski’s Clinical Ophthalmology, Edition 9, Page- 255, 256, 257, 258, 259.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: Which of the following conditions is most commonly associated with the development of a "beaten-metal" appearance on the cornea? (Q853813)

A. Posterior Polymorphous Corneal Dystrophy (PPCD)

B. Congenital Hereditary Endothelial Dystrophy (CHED)

C. Keratoconus


D. Fuchs Endothelial Corneal Dystrophy (FECD)

Correct Answer: D. Fuchs Endothelial Corneal Dystrophy (FECD)

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Fuchs Endothelial Corneal Dystrophy is strongly associated with the "beaten-metal" appearance on Descemet's membrane. It is a common type of endothelial dystrophy, particularly affecting older adults. Other characteristic feature seen in Fuchs corneal dystrophy is corneal guttata. {{caption_text}} Posterior Polymorphous Corneal Dystrophy (PPCD) (Option A): PPCD is a corneal dystrophy that often presents with unique lesions on the endothelium, such as nodules, vesicles, or band-like patterns that resemble railroad tracks. Congenital Hereditary Endothelial Dystrophy (CHED) (Option B): CHED is a genetic disorder where the endothelium is abnormal from birth or early childhood with diffuse corneal clouding, thickening, and decreased vision. Keratoconus (Option C): Keratoconus is a condition where the cornea progressively thins and bulges into a cone-like shape, causing irregular astigmatism and vision distortion. It does not involve Descemet's membrane or produce the "beaten-metal" appearance.



References:

Kanski’s Clinical Ophthalmology, 9th Edition 9, Pages 259, 260, 261.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q17: Match the following. A. {{caption_text}} 1. Rose bengal B. {{caption_text}} 2. Lissamine green C. {{caption_text}} 3. Fluorescein sodium (Q853814)


A. A-2, B-3, C-1


B. A-1, B-2, C-3

C. A-3, B-2, C-1

D. A-1, B-3, C-2










Correct Answer: C. A-3, B-2, C-1

- *Listen (English)*
- *Listen (Hindi)*

Explanation: {{caption_text}} Fluorescein sodium Fluorescein penetrates poorly into the lipid layer of the corneal epithelium, and therefore, it does not stain normal cornea. Used to stain the ulcer area {{caption_text}} Lissamine green Green colored stain Used to stain devitalized areas- dead cells and mucus. Used in dry eye {{caption_text}} Rose bengal Red colored stain Stains dead and devitalized tissue Used in dry eye. Note: In Dry eye Lissamine green stain is preferred as the rose bengal stains stings when in contact with mucous membranes.


@Neet_pg_bot


@Neet_pg_bot



References:

www.ncbi.nlm.nih.gov/pmc/articles/PMC10276680/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A young patient presents with this sign. Diagnosis is best made by: (Q882199)

A. Keratometry

B. Corneal topography

C. Pachymetry

D. A-scan



Correct Answer: B. Corneal topography

Explanation: The image demonstrates Munson sign, which is suggestive of advanced keratoconus, caused by conical protrusion of the cornea. Corneal topography is the gold standard investigation for diagnosing keratoconus, as it provides a detailed map of corneal curvature, detects early ectatic changes, and identifies asymmetric inferior steepening, even before clinical signs become evident. Keratometry (Option A): Measures central corneal curvature only and may miss early or irregular astigmatism seen in keratoconus. Pachymetry (Option C): Measures corneal thickness and is useful for monitoring progression, but cannot diagnose keratoconus alone. A-scan (Option D): Used for axial length measurement and intraocular lens power calculation, not for corneal ectatic disorders.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: A 70 y/o woman presents with severe pain in her right eye and forehead, along with a rash that started 5 days ago. She was diagnosed with Herpes Zoster Ophthalmicus (HZO). Which of the following statements is true regarding the management of her condition? (Q853762)

A. Systemic steroids alone are sufficient to manage HZO

B. Antiviral therapy is started within 72 hours of rash onset.

C. Postherpetic neuralgia is less common in patients over 70 years of age.

D. Topical antiviral therapy is not recommended to manage eye involvement in HZO.

Correct Answer: B. Antiviral therapy is started within 72 hours of rash onset.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Antiviral therapy is most effective when started within 72 hours of rash onset, its effectiveness diminishes significantly after this period. However, treatment may still be considered beyond 72 hours, especially in severe cases, but the benefits are reduced. Systemic steroids alone are sufficient to manage HZO (Option A): Systemic steroids should not be used as a monotherapy for HZO. They must be combined with antiviral therapy to manage inflammation and pain effectively, as steroids alone do not reduce viral replication or prevent complications. Post-Herpetic Neuralgia: Up to 75% in patients over 70 (Option C) Topical antiviral therapy is not recommended to manage eye involvement in HZO (Option D): Topical antiviral therapy in combination with systemic antiviral therapy is the standard of care for managing the viral infection.

References:

Kanski's Clinical Ophthalmology, 9th edition, Page 226, 227, 229.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: What is the average thickness of the central cornea? (Q853606)

A. 1 to 1.2 mm

B. 0.8 to 0.9 mm

C. 0.5 to 0.6 mm

D. 7.8 mm

Correct Answer: C. 0.5 to 0.6 mm

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: 0.5 to 0.6 mm is the correct average thickness of the cornea at its central region, where it is thinnest . Central corneal thickness (CCT) is measured via pachymetry, typically using ultrasound or optical methods. In procedures like LASIK, sufficient CCT (usually >480-500 µm pre-surgery) ensures a safe residual stromal bed (>250-300 µm post-ablation) to prevent ectasia. Thin corneas increase complication risks, often requiring alternatives like PRK. CCT affects Goldmann applanation tonometry (GAT) accuracy: thick corneas (>550 µm) overestimate IOP, while thin corneas (<500 µm) underestimate it, potentially masking glaucoma risk. 10 microns change in CCT →0.7mmHg change in IOP Diameter 11-11.7 mm Refractive Index 1.376 Refractive Power 45 Dioptre Radius of curvature Anterior - 7.8mm (Option D) Posterior - 6.5mm Thickness at the centre 0.5-0.6 mm (Option C) Thickness at the periphery 1-1.2 mm (Option A) 0.8 to 0.9 mm (Option B): This thickness is greater

than the actual average central thickness, suggesting a thicker cornea.

References:

Kanski's Clinical Ophthalmology, 9th Edition , Page 204.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: What is the primary metabolism of cornea? (Q853608)

A. Anaerobic

B. Aerobic.

C. Both anaerobic and aerobic

D. None of the above

Correct Answer: B. Aerobic.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The cornea uses both aerobic and anaerobic metabolism. But the primary pathway = aerobic metabolism. Oxygen & Nutrition Oxygen comes mainly from the atmosphere (tear film). Nutrients diffuse from the aqueous humor and limbal vessels. Lens In the lens, the primary metabolism is anaerobic. ~80% of glucose in the lens is metabolized via anaerobic glycolysis. Anaerobic Glycolysis in Cornea If anaerobic glycolysis is activated in cornea → corneal edema. Mechanism: Lactic acid accumulates → metabolic acidosis. This inhibits the Na■/K■ pump of endothelial cells. Pump failure → hydration of cornea → corneal edema.

References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 204.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: Which of the following factors is not responsible for maintaining corneal transparency? (Q853609)

A. Corneal Epithelium and Tear Film

B. Avascularity of the Cornea

C. Corneal nerve supply

D. Endothelial Pumps

Correct Answer: C. Corneal nerve supply

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Corneal Nerve Supply is not directly involved in maintaining transparency. It provides sensation to the cornea but does not influence the clarity of the corneal tissue. Factors Responsible for Corneal Transparency: Corneal Epithelium and Tear Film (Option A) Arrangement of Corneal Lamellae Avascularity (Option B) Barrier Function Endothelial Pumps (Option D) Evaporation Normal Intraocular Pressure (IOP) Swelling Pressure (SP) Corneal Crystals/Proteins

References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 204.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Match the following corneal assessment techniques with their correct descriptions: 1) Keratometry a) Measures both anterior and posterior corneal surfaces. 2) Placido's Disc b) Assesses the smoothness and curvature of the corneal surface. 3) Corneal Topography c) Evaluate the anterior curvature of the cornea and generate colour-coded maps. 4) Corneal Tomography (Pentacam) d) Measures the curvature of the central cornea. (Q853610)

A. 1-a, 2-c, 3-d, 4-b

B. 1-d, 2-b, 3-c, 4-a

C. 1-d, 2-a, 3-b, 4-c

D. 1-b, 2-a, 3-c, 4-d


Correct Answer: B. 1-d, 2-b, 3-c, 4-a


■ *Listen (English)*

■ *Listen (Hindi)*


Explanation: Instrument Image Keratometry Definition: Objective measurement of central corneal curvature to assess astigmatism Principle: Image size analysis of anterior cornea using visible doubling Normal Values:

Average ~45 D; >45 D in keratoconus Types: Javal-Schiotz, Bausch & Lomb {{caption_text}} Auto refractive Keratometer Keratoscopy / Placido’s Disc Disc with alternating black & white rings Assesses corneal surface smoothness & curvature Normal: Uniform, sharp ring reflections Abnormal: Distorted rings → corneal irregularities Corneal Topography Evaluates anterior corneal curvature Produces colour-coded maps (shape, curvature, elevation) Uses: Diagnose keratoconus, plan refractive surgery Limitation: No information on posterior corneal surface {{caption_text}} Corneal Tomography (Pentacam) Measures anterior & posterior corneal surfaces Gives 3D corneal structure Detects subclinical keratoconus Assesses refractive surgery candidates Uses Scheimpflug imaging for accurate measurements Helpful in pre-op evaluation & disease monitoring {{caption_text}}






@Neet_pg_bot



@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 209

www.ncbi.nlm.nih.gov/books/NBK562157/#:~:text=Corneal%20topography%20evaluates%20the%20anterior%20surface%20of%20the%20cornea%20and,indications%2C%20and%20interpret%20its%20findings

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: Which of the following statements is false? (Q853611)

A. Pachymetry measures corneal thickness and is used for diagnosing glaucoma

B. Specular Microscopy is used for photographing the corneal endothelium

C. Aesthesiometer measures corneal sensation and is used to assess neurotrophic keratitis

D. Aesthesiometer measures corneal sensation by evaluating the second cranial nerve

Correct Answer: D. Aesthesiometer measures corneal sensation by evaluating the second cranial nerve


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Aesthesiometer (Esthesiometer) measures corneal sensation by evaluating the ophthalmic branch of the fifth cranial nerve (trigeminal nerve), not the second cranial nerve (optic nerve). The fifth cranial nerve is responsible for sensory input from the cornea, which includes detecting tactile sensations. The second cranial nerve (optic nerve) is primarily involved in vision, not sensation. INSTRUMENT IMAGE Pachymetry Measures corneal thickness. Used for diagnosing and treating glaucoma . (Option A) In glaucoma workup: Thick cornea → IOP overestimated by Goldmann tonometer. Thin cornea → IOP underestimated. {{caption_text}} Specular Microscopy: Noninvasive technique for photographing and assessing the corneal endothelium. (Option B) Evaluates endothelial cell number + morphology. Detects polymegathism (variation in size) & pleomorphism (variation in shape). {{caption_text}} {{caption_text}} Esthesiometer (Aesthesiometer): Measures corneal sensation (trigeminal V1) Methods: Cotton tip (qualitative), Cochet-Bonnet esthesiometer (nylon filament, 0–6 cm, 11–200 mm/gm) Uses: Diagnose neurotrophic keratitis (Option C) , monitor corneal health in long-term contact lens wearers {{caption_text}}




@Neet_pg_bot


@Neet_pg_bot



References:

Corneal Pachymetry Test: What It Is, Normal Range & FAQs

Specular Microscopy - StatPearls - NCBI Bookshelf

Corneal Esthesiometry - EyeWiki

Kanski's Clinical Ophthalmology, 9th Edition, Page 209.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: A 32 y/o contact lens wearer presents with a red, painful eye, blurred vision, and increased sensitivity to light. Examination reveals lid swelling, conjunctival chemosis, and a large corneal ulcer with greenish exudate, rapidly expanding with hypopyon. What is the most likely cause? (Q853657)

A. Staphylococcus aureus

B. Streptococcus pneumoniae

C. Pseudomonas aeruginosa

D. Enterobacteriaceae


Correct Answer: C. Pseudomonas aeruginosa

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient's clinical presentation, including the irregular corneal ulcer with greenish exudate and hypopyon, along with a history of contact lens use, suggests a bacterial corneal ulcer. Pseudomonas aeruginosa is known for causing irregular corneal ulcers with greenish mucopurulent exudate, rapid progression, and hypopyon. It is also common in contact lens-related infections. Etiology of Purulent Corneal Ulcer: Corneal epithelial damage: Abrasion, xerosis, exposure keratitis, keratomalacia, bullous keratopathy, neuroparalytic keratitis Pathogens invading intact epithelium: N. gonorrhoeae, N. meningitidis, C. diphtheriae, H. aegyptius, Listeria Sources of infection: Exogenous: conjunctival/lacrimal sac, foreign bodies, vegetative material, airborne/waterborne Common bacteria: General: Staph, Strep pneumoniae, Pseudomonas, Enterobacteriaceae

Contact lens wearers: Pseudomonas (MC), Acanthamoeba (differential) Causative organisms: Gram + cocci: Staph aureus/epidermidis, Strep pneumoniae Gram – cocci: Neisseria spp. Gram + bacilli: Corynebacterium, Bacillus cereus, Propionibacterium, Listeria, Clostridium Gram – bacilli: Pseudomonas, Klebsiella, Proteus, E. coli, Serratia, Moraxella, H. influenzae Filamentous: Actinomyces, Nocardia Mycobacteria: NTM, M. tuberculosis



References:

Kanski's Clinical Ophthalmology, Edition 9, Page- 209, 210, 211.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: A 60 y/o woman presents with a painful red eye, photophobia, and a history of chronic use of corticosteroid eye drops. Examination reveals a corneal ulcer with necrosis extending deep into the corneal stroma, approaching Descemet's membrane. Which type of bacterial keratitis is this patient most at risk for? (Q853658)

A. Localized Ulcer

B. Perforating Ulcer

C. Adherent Leucoma


D. Sloughing Ulcer

Correct Answer: B. Perforating Ulcer

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The description of chronic use of corticosteroid eye drops, the ulcer penetrating deeply into the corneal stroma, approaching Descemet's membrane, suggests a Perforating Ulcer . Most common cause leading to perforation: Pseudomona Localized Ulcer (Option A): This term is too general and does not reflect the severe, deep nature of the ulcer described, which extends near Descemet's membrane, indicating a high risk of progression beyond just being "localized." Adherent Leucoma (Option C): An adherent leucoma is a sequela or complication after a corneal perforation has occurred and healed with iris incarceration, not the type of bacterial keratitis the patient is currently at risk for. Sloughing Ulcer (Option D): While necrosis is present and sloughing might occur, the most immediate and critical risk associated with an ulcer extending deep into the stroma and approaching Descemet's membrane is full-thickness penetration, not merely the shedding of necrotic tissue.



References:

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 211, 212, 213.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: Which of the following statements about the treatment of bacterial keratitis is incorrect? (Q853659)

A. Cycloplegics like atropine can be used to reduce ciliary spasms and pain.

B. Ciprofloxacin (750mg twice daily) is given when the sclera gets involved.

C. Topical steroids should be started immediately to reduce inflammation.

D. Duo therapy with fortified cefazolin and tobramycin provides broad-spectrum coverage.

Correct Answer: C. Topical steroids should be started immediately to reduce inflammation.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Topical steroids should not be used in bacterial keratitis as they can exacerbate the infection and delay healing. Treatment of Bacterial Keratitis Empirical Monotherapy : A single fluoroquinolone (e.g., ofloxacin) is recommended but not standard practice due to high resistance rates in some regions. Perforation is more likely in ulcers treated with ofloxacin . Duo Therapy (Option D) : Use a combination of fortified cefazoline 5% and fortified tobramycin 1.3% to cover both gram-positive and gram-negative bacteria. Ointments : Ocular ointments can be beneficial at bedtime for milder cases and may serve as useful adjunctive therapy. Cycloplegics (Option A) : Administer 1% atropine eye drops to reduce ciliary spasm pain, prevent posterior synechiae, increase blood supply, and decrease exudation. 2% homatropine is an alternative. Vitamin Supplements : Vitamins A, B-complex, and C are recommended to promote early healing of the corneal ulcer. Scleral involvement : In Scleral or systemic involvement cases, systemic antibiotics like oral ciprofloxacin (750 mg twice daily) (Option B) or a combination of cephalosporin and aminoglycoside may be necessary. Avoid Sponge/Pad and steroids. Avoid using an eye sponge or pad to prevent infection from worsening by trapping bacteria. Steroids should be avoided as they may exacerbate the infection and delay the healing process.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 214, 215, 216.

Ocular ointments

↑ *Back to this question in Part A*

@Neet_pg_bot

Q28: A 60 y/o man from a tropical region presents with a painful, red eye and a corneal ulcer following recent exposure to decaying vegetable matter. Which is the most common fungal infection affecting the cornea? (Q853755)

A. Candida albicans

B. Fusarium

C. Mucormycosis

D. Aspergillus fumigatus

Correct Answer: D. Aspergillus fumigatus

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Aspergillus fumigatus is a common causative organism of keratomycosis in tropical and subtropical regions and is often associated with exposure to decaying organic matter. Keratomycosis Etiology Most Common Causative Organisms : Aspergillus fumigatus: Most common in India and tropical regions. Fusarium (Option B): Common in warm climates like Brazil. Candida albicans (Option A): More frequent in temperate climates. Mode of Infection : Injury with Vegetative Material: Typically occurs after trauma with objects like thorns, wooden sticks, crop leaves, or decaying vegetable matter. Common in Field Workers: Especially during harvesting season in rural and tropical areas. Immunocompromised Individuals: Higher risk for developing fungal keratitis. Special Considerations : Candida albicans: Leading cause of endogenous endophthalmitis. Mucormycosis (Option C) & Invasive Aspergillosis: Considered as the cause of orbital cellulitis in immunocompromised patients.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 216.

Endophthalmitis

Tropical

Orbital cellulitis

↑ *Back to this question in Part A*

@Neet_pg_bot


Q29: A 30 y/o contact lens wearer presents with very severe eye pain, blurred vision, and discomfort in his right eye. O/E, the following corneal finding is noted. Based on the image and the clinical presentation, which of the following is the most likely diagnosis? (Q853764)

A. Herpes Simplex Keratitis

B. Fungal Keratitis

C. Acanthamoeba Keratitis


D. Bacterial Keratitis



Correct Answer: C. Acanthamoeba Keratitis

- *Listen (English)*
- *Listen (Hindi)*

Explanation: s: The above clinical features and examination findings are suggestive of Acanthamoeba keratitis. Acanthamoeba keratitis is characterized by severe pain that is often out of proportion to the clinical findings, along with features such as perineural infiltrates and, in advanced cases, ring-shaped abscesses. Symptoms: Blurred Vision Discomfort Severe Pain Ring lesion indicates advanced stage of the disease. Herpes Simplex Keratitis (Option A) typically presents with dendritic ulcers and less severe pain relative to Acanthamoeba keratitis. Fungal Keratitis (Option B) usually presents with feathery-edged infiltrates and satellite lesions. Bacterial Keratitis (Option D) typically presents with a more prominent purulent discharge and a dense, localized infiltrate.



References:

Kanski's Clinical Ophthalmology, 9th Edition, Pages 211, 216, 219, 232

↑ *Back to this question in Part A*

@Neet_pg_bot

Q30: A 65 y/o woman presents with a painful rash on her forehead and eyelid. She also reports blurred vision and photophobia. O/E, she has conjunctivitis, punctate epithelial keratitis, and iridocyclitis. Which of the following is a chronic ocular complication she is at risk of developing? (Q853761)

A. Episcleritis

B. Corneal Scarring

C. Conjunctivitis

D. Secondary Glaucoma

Correct Answer: B. Corneal Scarring

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: s: The above features describe herpes zoster ophthalmicus. A chronic ocular complication she is at risk of developing is corneal scarring. Acute Ocular Manifestations: Conjunctivitis (Option C) : Redness and inflammation of the conjunctiva. Keratitis: Punctate Epithelial Keratitis: Coarse and irregular lesions. Nummular Keratitis: Larger, disc-shaped lesions. Disciform keratitis. Iridocyclitis: Inflammation of the iris and ciliary body, causing pain, photophobia, and blurred vision. Keratouveitis and endothelitis: Leads to acute endothelial cell loss Episcleritis and Scleritis (Option A) : Inflammation of the episclera or sclera, resulting in pain and redness. Chronic Ocular Manifestations: Post-herpetic neuralgia Corneal lesions: Neuroparalytic ulceration. Neurotrophic Keratopathy Corneal scarring Exposure keratitis Mucous plaque keratitis (Sudden development of elevated mucous plaque which stains on rose bengal) Lid lesions: D/t scarring. Include ptosis, trichiasis & entropion. Secondary Glaucoma (Option D): Increased intraocular pressure (IOP) leading to optic nerve damage and vision loss. Hutchinson's rule: The eye will be involved if the nose tip is involved. As it indicates nasociliary nerve involvement.

References:

Kanski's Clinical Ophthalmology, 9th Edition, Pages 224, 225.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q31: A 50 y/o male presents with decreased vision, redness, and a dull sensation in his R/E. He has a history of trigeminal neuralgia for which he underwent a Gasserian ganglion block. O/E, there is decreased corneal sensation and epithelial desquamation. Which of the following is the most likely cause of his symptoms? (Q853760)

A. Keratomalacia

B. Herpes Simplex Virus (HSV) Infection

C. Neurotrophic Keratopathy

D. Neuroparalytic Keratopathy

Correct Answer: C. Neurotrophic Keratopathy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Neurotrophic keratopathy is characterized by decreased corneal sensation and epithelial desquamation due to trigeminal nerve damage or dysfunction. The patient's history of Gasserian ganglion block, a procedure known to impair the trigeminal nerve, strongly supports this diagnosis. Causes of Decreased Corneal Sensation: Herpes Simplex Virus (HSV) Infections (Option B) : HSV keratitis typically presents with dendritic ulcers on the cornea and is associated with pain, photophobia, and possibly vesicular skin lesions. However, the history of a Gasserian ganglion block and the description of decreased corneal sensation point more toward neurotrophic keratopathy. Herpes Zoster Ophthalmicus: Caused by varicella zoster virus. Leads to decreased

corneal sensation. Keratomalacia (Option A) Result of vitamin A deficiency. The cornea becomes dull and insensitive. Signs include haziness and yellow infiltrates (not seen in this patient) Can progress to corneal necrosis. Neurotrophic Keratopathy Neuroparalytic Keratopathy (Option D): This condition occurs due to facial nerve paralysis , leading to lagophthalmos (inability to close the eye) and subsequent corneal exposure. The patient's history does not suggest facial nerve paralysis, making this less likely.

References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 223.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q32: A 35 y/o patient presents with pain, redness, photophobia, and decreased vision in the right eye. Slit-lamp examination reveals a dendritic ulcer on the cornea. Which of the following is the most appropriate initial treatment for this patient? (Q853759)

A. Topical steroids and antibiotics

B. Oral acyclovir and lubricants

C. Topical acyclovir and cycloplegics

D. Oral steroids and topical antibiotics

Correct Answer: C. Topical acyclovir and cycloplegics

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient presents with pain, redness, photophobia, and decreased vision in the right eye, along with a dendritic ulcer on the cornea. This clinical picture is classic for Herpes Simplex Keratitis (HSK), an infection of the cornea caused by the herpes simplex virus (HSV). Treatment of Herpes Simplex Virus (HSV) Keratitis: Topical Therapy: Antivirals: Acyclovir 3% ointment: 5 times daily. Other Topical Medications: Antibiotics: Prevent secondary bacterial infections, especially with epithelial defects. Cycloplegics: (e.g., Homatropine 1%) Relieve pain and photophobia by paralyzing ciliary muscles and dilating the pupil. Topical steroids and antibiotics (Option A): Topical steroids are contraindicated in the treatment of active epithelial HSV keratitis, as they can suppress the immune response and worsen the viral infection, potentially leading to complications like corneal perforation. Antibiotics are not needed unless there is a secondary bacterial infection, which is not indicated here. Oral acyclovir and lubricants (Option B): While oral acyclovir is used in more severe or recurrent cases of HSV keratitis, topical acyclovir is preferred as the initial treatment. Lubricants can be supportive but are not the primary treatment. Oral steroids and topical antibiotics (Option D): Oral steroids are contraindicated in HSV keratitis, similar to topical steroids, as they can exacerbate the infection. Topical antibiotics are unnecessary unless there's evidence of bacterial superinfection.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 221.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q33: A 45 y/o patient presents with eye redness, excessive tearing, and sensitivity to light. O/E, you observe dendritic ulcers and decreased corneal sensation. Based on these findings, which is the most likely etiological agent responsible for the patient's condition? (Q853758)

A. Adenovirus

B. Varicella Zoster Virus (VZV)

C. Herpes Simplex Virus (HSV)

D. Chlamydia trachomatis

Correct Answer: C. Herpes Simplex Virus (HSV)

■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: s: HSV keratitis is characterized by dendritic ulcers on the cornea, which are branching, tree-like lesions. HSV also commonly causes decreased corneal sensation. {{caption_text}} Herpes Simplex Virus (HSV) Keratitis: Dendritic Ulcers: Branching, tree-like ulcers. Geographic Ulcers: Larger, amoeboid-shaped ulcers. Stromal Keratitis: Inflammation of the corneal stroma, leading to oedema, infiltrates, and scarring. Disciform Keratitis: Central, disc-shaped inflammation in the stroma. {{caption_text}} Adenovirus (Option A): Adenoviruses are known for causing epidemic keratoconjunctivitis (EKC) and may present with conjunctivitis and keratitis. They are less likely to cause dendritic ulcers, which are specific to HSV. Herpes Zoster Ophthalmicus (HZO): (Option B) Vesicles and Scars: Painful rash and blisters following the involvement of ophthalmic division of the trigeminal nerve. Nummular Keratitis: Large, coin-shaped, subepithelial corneal infiltrates. {{caption_text}} Chlamydia trachomatis (Option D): Chlamydia typically causes trachoma or inclusion conjunctivitis, which may involve conjunctival scarring and corneal pannus, but not dendritic ulcers.




Dendritic ulcer

@Neet_pg_bot



Geographic (amoebic) ulcer

@Neet_pg_bot



Nummular Keratitis

@Neet_pg_bot

References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 219, 220

↑ *Back to this question in Part A*

@Neet_pg_bot

Q34: A 45 y/o male presents with redness, pain, and decreased vision in his right eye. He reports a history of trauma to the eye with a plant material three days ago. O/E, there is a corneal ulcer with feathery edges and satellite lesions. Hypopyon is also noted. Which of the following is the most appropriate initial treatment for this patient? (Q853757)

A. Topical Moxifloxacin

B. Topical Natamycin

C. Oral Acyclovir

D. Intravitreal Amphotericin B

Correct Answer: B. Topical Natamycin

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The history of trauma with plant material, along with the presence of a corneal ulcer with feathery edges and satellite lesions, is highly suggestive of fungal keratitis . Natamycin (5%) is the first-line treatment for fungal keratitis, especially caused by filamentous fungi like Fusarium and Aspergillus . It is effective in treating the type of fungal infection that is suggested by the clinical presentation in this patient. Topical Moxifloxacin (Option A): Moxifloxacin is an antibiotic effective against bacterial infections, not fungal infections. Given the history of trauma with plant material and the appearance of a corneal ulcer with feathery edges and satellite lesions, the most likely diagnosis is fungal keratitis. Therefore, moxifloxacin would not be appropriate. Oral Acyclovir (Option C): Acyclovir is an antiviral medication used primarily for herpes simplex virus infections. It would not be effective for a fungal corneal ulcer, making it an inappropriate choice in this scenario. Intravitreal Amphotericin B (Option D): Intravitreal Amphotericin B is used in cases of fungal endophthalmitis, not as an initial treatment for keratitis. Although hypopyon suggests severe infection, the primary treatment should still be topical antifungal agents unless there is confirmed or high suspicion of endophthalmitis requiring intravitreal therapy.

References:

Kanski’s Clinical Ophthalmology, 9th Edition 9, Pages 217, 218.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q35: A 45 y/o man from a tropical region presents with a painful red eye, significant watering, and sensitivity to light after working in a garden with decaying plant material. The patient is diagnosed with keratomycosis. Which of the following is not an examination finding in keratomycosis? (Q853756)

A. Feathery extensions at the margins of the corneal ulcer

B. Extensive corneal vascularization

C. Sterile immune ring

D. Endothelial plaque


Correct Answer: B. Extensive corneal vascularization


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Extensive corneal vascularization is not seen in keratomycosis (fungal keratitis). In fact, corneal neovascularization is typically absent , especially in the acute stages. {{caption_text}} Feathery extensions at the

margins of the corneal ulcer (Option A): These are characteristic finger-like projections at the ulcer margins caused by filamentous fungi. Sterile immune ring (Option C): Also known as Wessely ring, formed at the interface of fungal antigens and host antibodies. Endothelial plaque (Option D): Dense, whitish deposits of fibrin and leukocytes under the stromal lesion.





References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 216, 217

↑ *Back to this question in Part A*

@Neet_pg_bot

Q36: A 55 y/o male with poorly controlled diabetes presents with a non-healing corneal ulcer in his right eye, despite two weeks of appropriate antibiotic therapy. The ulcer shows signs of impending perforation with severe thinning of the cornea. Which of the following is the most appropriate next step in the management of this patient? (Q853660)

A. Increase the frequency of antibiotic eye drops.

B. Apply a pressure bandage and recommend strict bed rest.

C. Perform mechanical debridement and apply tissue adhesive glue.

D. Start topical steroids to reduce inflammation.

Correct Answer: C. Perform mechanical debridement and apply tissue adhesive glue.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: For impending corneal perforation, necrotic tissue is first removed (mechanical debridement), then tissue adhesive (e.g., cyanoacrylate glue) is applied to seal the thin area and prevent rupture. Mechanical debridement is necessary to remove necrotic material and promote healing in a non-healing ulcer. Tissue adhesive glue is indicated to prevent perforation, especially when the perforation is less than 2mm. If the perforation is large, then penetrating keratoplasty can be done. Increasing the frequency of antibiotic eye drops (Option A): At this stage, just increasing frequency won’t be useful. Applying a pressure bandage and bed rest (Option B): Pressure can worsen thinning and precipitate perforation. Topical steroids (Option D) are contraindicated as they can worsen the infection.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 214, 215, 216

↑ *Back to this question in Part A*

@Neet_pg_bot

Q37: A 58-year-old patient is evaluated before cataract surgery. Specular microscopy shows normal endothelial morphology and adequate reserve to maintain corneal

deturgescence. Which of the following values most closely represents the expected endothelial cell density in such an adult? (Q885150)

A. 800-1200 cells/mm²

B. 300-600 cells/mm²

C. 2400-3000 cells/mm²

D. 3500-4200 cells/mm²

Correct Answer: C. 2400-3000 cells/mm²

Explanation: In adults, endothelial density averages about 2500-3000 cells/mm². Cell count declines with age because endothelial cells do not regenerate; remaining cells enlarge to cover defects. Age Group Endothelial Cell Density (cells/mm²) At birth 3500-4000 Young adults 2500-3000 Old age (around 80 years) ~2000 Clinical Significance: Endothelial cell loss may occur due to surgical trauma, inflammation, or corneal dystrophies. When the cell density drops below approximately 500 cells/mm² , the endothelium can no longer maintain corneal clarity, leading to corneal edema and decompensation. 800-1200 cells/mm² (Option A): Significantly reduced reserve; increased risk of postoperative corneal edema. 300-600 cells/mm² (Option B): Critically low density; corneal decompensation likely. 3500-4200 cells/mm² (Option D): Typical of neonates and infants, not adults.

References:

Parsons Diseases of the Eye, 23rd Edition, Page 312.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q38: A child presents for ophthalmic evaluation due to an abnormally small-appearing cornea. On examination, the horizontal corneal diameter is found to be reduced. Microcornea is diagnosed when the horizontal corneal diameter measures below which of the following values? (Q885151)

A. Less than 12 mm

B. 10 mm or less

C. Less than 9 mm

D. Less than 11.5 mm

Correct Answer: B. 10 mm or less

Explanation: The given clinical scenario refers to Microcornea , a developmental anomaly characterized by a reduced horizontal corneal diameter. Developmental Anomalies of Cornea Microcornea Horizontal corneal diameter ≤ 10 mm in children/adults (Option B) < 9 mm in newborns (Option C ruled out) Globe size usually normal Associated with hypermetropia and shallow anterior chamber Megalocornea Diameter ≥ 12 mm (≥ 13 mm after 2 years) Associated with high myopia, astigmatism Often X-linked recessive Cornea Plana Abnormally flat cornea Causes high hypermetropia Keratoglobus / Keratectasia Severe corneal thinning with generalized protrusion May be linked to raised IOP or intrauterine keratitis Microphthalmos Entire eye is small Axial length ≥ 2 SD below normal Nanophthalmos Small but structurally normal eye Associated with hypermetropia, strabismus Anophthalmos Complete absence of the globe

References:

Kanski's Clinical Ophthalmology, 9th edition, Page 270.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q39: A patient develops rapidly progressive corneal ulceration despite no history of trauma or epithelial defect. Which organism is known to invade an intact corneal epithelium? (Q885152)

A. Streptococcus pneumoniae

B. Staphylococcus aureus

C. Neisseria gonorrhoeae

D. Moraxella lacunata

Correct Answer: C. Neisseria gonorrhoeae

Explanation: Among the given options, Neisseria gonorrhoeae can invade an intact corneal epithelium, causing rapidly progressive corneal ulceration even in the absence of trauma or a pre-existing epithelial defect. Organisms that can penetrate intact corneal epithelium: Neisseria gonorrhoeae Neisseria meningitidis Corynebacterium diphtheriae Haemophilus aegyptius Listeria Shigella Corneal defenses: Intact epithelium, tear film, and antimicrobial factors prevent infection; most bacteria need an epithelial defect to invade. Clinical significance: Gonococcal keratitis causes hyperacute purulent conjunctivitis with rapid corneal involvement and a high risk of perforation, making it an ophthalmic emergency. Incorrect Options: Streptococcus pneumoniae (Option A) requires epithelial defect; causes serpiginous ulcer. Staphylococcus aureus (Option B): Opportunistic; needs epithelial compromise. Moraxella lacunata (Option D) causes angular conjunctivitis; corneal infection needs a defect.

References:

Kanski's Clinical Ophthalmology, 9th edition, Page 173.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q40: A young woman presents with sudden onset severe ocular pain, photophobia, tearing and a corneal epithelial defect with stromal infiltrate on slit-lamp examination. A bacterial corneal ulcer is diagnosed. What concentration of topical cefazolin is recommended in fortified therapy? (Q885153)

A. 2%

B. 5%

C. 10%

D. 1%

Correct Answer: B. 5%

Explanation: Fortified cefazolin eye drops are used at 5% concentration in combination with an aminoglycoside (e.g., tobramycin) for broad-spectrum coverage. Rationale: Poor corneal vascularity requires high local antibiotic concentrations; fortified drops are prepared from injectable forms. Cefazolin action: First-generation cephalosporin effective against gram-positive organisms (Staphylococcus, Streptococcus). Initial management: Start intensive topical broad-spectrum antibiotics; escalate to fortified drugs if needed. Common fortified regimen: Cefazolin 5%, Tobramycin 1.3%, Vancomycin 5%. Adjuncts: Cycloplegics (e.g., atropine 1%), vitamins A & C, analgesics. Non-healing Corneal Ulcers: Debridement, chemical cautery, or conjunctival flap. Threatened perforation: Lower IOP, cyanoacrylate glue, bandage contact lens.

References:

Kanski’s Clinical Ophthalmology, 9th edition, Page 217.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q41: A farmer presents with gradually increasing ocular pain, photophobia and blurred vision after minor trauma with plant matter. Slit lamp shows a dry-looking corneal ulcer with feathery margins, satellite lesions and an endothelial plaque. A hypopyon is present. Which organism group is most likely responsible? (Q885155)

A. Gram-positive cocci


B. Filamentous fungi


C. Acanthamoeba

D. Herpes simplex virus

Correct Answer: B. Filamentous fungi

Explanation: The clinical scenario is classical for fungal keratitis , particularly infection by filamentous fungi such as Fusarium and Aspergillus , which are the most common cause of fungal keratitis. Fungal Keratitis Etiopathogenesis Filamentous fungi (Fusarium , Aspergillus), hyphal organisms. Usually, after plant matter trauma ; yeast (Candida) affects immunocompromised eyes. Predisposing factors Trauma, chronic ocular disease, steroids, contact lenses, diabetes, and immunosuppression. Clinical features Gradual onset of pain, photophobia, blurred vision, discharge Filamentous keratitis signs: Grey-white corneal infiltrate with feathery margins Satellite lesions Ring infiltrates or branching extensions Stromal necrosis and thinning Endothelial plaque and hypopyon Possible anterior uveitis, raised IOP, or endophthalmitis Investigations Corneal scraping, KOH mount , Gram/Giemsa, culture on Sabouraud agar Management Topical antifungals: Natamycin 5% (first choice), econazole, and amphotericin B. Adjuncts: Cycloplegics, cautious antibiotics. Systemic: Voriconazole/itraconazole for deep lesions. Procedures: Debridement, keratoplasty, tissue adhesive if needed. Gram-positive cocci (Option A) produce suppurative ulcer with dense infiltrate, not feathery margins. Acanthamoeba (Option C): Severe pain out of proportion, ring infiltrate but history of water exposure or contact lenses. Herpes simplex virus (Option D): Dendritic epithelial lesions, reduced corneal sensation.





References:

Kanski's Clinical Ophthalmology, 9th edition, Pages 218-220.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q42: A farmer develops a corneal ulcer after injury with plant material. Slit-lamp shows a dry ulcer with feathery margins and stromal infiltrates arranged in a curvilinear pattern. Which type of hypersensitivity reaction is mainly responsible for these stromal infiltrates? (Q885156)

A. Type I (IgE-mediated reaction)


B. Type II (antibody-mediated cytotoxic reaction)

C. Type III (immune complex-mediated reaction)

D. Type IV (delayed cell-mediated reaction)

Correct Answer: C. Type III (immune complex-mediated reaction)

Explanation: The stromal infiltrates in fungal keratitis result from a Type III immune complex-mediated reaction , where fungal antigens form complexes with antibodies, activating complement and neutrophils, causing enzymatic stromal damage and curvilinear infiltrates. Mechanism: Immune complex deposition in stroma Complement activation Neutrophil chemotaxis Enzymatic stromal destruction Formation of curvilinear/ring-shaped infiltrates. Type I (IgE-mediated reaction) (Option A): Immediate allergy, e.g., vernal keratoconjunctivitis. Type II (antibody-mediated cytotoxic reaction) (Option B): Targets cell surface antigens; not seen in corneal ulcers. Type IV (delayed, cell-mediated reaction) (Option D): Seen in granulomatous infections; not typical for microbial keratitis infiltrates.



References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 218-220.

Robbins & Cotran Pathologic Basis of Diseases,10th Edition, Page 210-212.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q43: A patient with infectious keratitis presents to the OPD. In which of the following ocular conditions are topical steroids strictly contraindicated? (Q885157)

A. Vernal keratoconjunctivitis

B. Stromal keratitis

C. Atopic dermatitis-related ocular inflammation

D. Fungal corneal ulcer

Correct Answer: D. Fungal corneal ulcer

Explanation: Topical steroids are strictly contraindicated in fungal corneal ulcers because they suppress the immune response and worsen fungal infections, increasing the risk of corneal melting and perforation. Topical steroids are strictly contraindicated because they: Promote fungal growth. Suppress host cell-mediated immunity, reducing the body's ability to fight infection. Increase the risk of stromal melting and corneal perforation. Mask clinical signs, potentially delaying diagnosis. Increase the chances of endophthalmitis. Vernal keratoconjunctivitis (Option A): Steroids are used in severe disease to control allergic inflammation. Stromal keratitis (Option B): Often immune-mediated (e.g., herpetic stromal keratitis); steroids are indicated with antiviral cover. Atopic dermatitis ocular involvement (Option C): Steroids may be required to control inflammation.

References:

Kanski's Clinical Ophthalmology, 9th edition, Page 216.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q44: A 38-year-old agricultural worker presents with gradually worsening eye pain and photophobia following trauma with vegetative matter. Slit-lamp examination reveals a dry corneal ulcer with feathery edges and satellite lesions. Corneal culture grows Fusarium species. Which of the following is the drug of choice for treatment? (Q885158)

A. Amphotericin B

B. Fluconazole

C. Natamycin

D. Nystatin

Correct Answer: C. Natamycin

Explanation: Eye trauma with vegetative matter causing a dry corneal ulcer with feathery edges and satellite lesions indicates Fusarium fungal keratitis , for which the drug of choice is topical Natamycin due to its strong activity against filamentous fungi. Risk factor: Eye trauma with vegetative matter (common in agricultural workers) Symptoms: Gradually progressive eye pain and photophobia Slit-lamp findings: Dry corneal ulcer Feathery margins Satellite lesions Diagnosis: Fungal keratitis Causative organism: Fusarium (filamentous fungus)

Drug of choice: Topical Natamycin 5% → Most effective against filamentous fungi, including Fusarium and Aspergillus. Amphotericin B (Option A): More effective against yeasts (e.g., Candida) and has limited efficacy against filamentous fungi like Fusarium. Fluconazole (Option B): Has poor activity against filamentous fungi and is ineffective for Fusarium keratitis. Nystatin (Option D): Primarily active against Candida species and is not used for filamentous fungal keratitis.

References:

Kanski's Clinical Ophthalmology, 9th edition, Pages 219-220.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [08_Cornea] 018 Previous Year Questions

« Back to Folder Directory

[08_Cornea] 018 Previous Year Questions — Quick Jump

Q1 (p. 1127)	*Q2 (p. 1128)*	*Q3 (p. 1130)*	*Q4 (p. 1131)*	*Q5 (p. 1133)*	*Q6 (p. 1134)*
Q7 (p. 1136)	*Q8 (p. 1138)*	*Q9 (p. 1141)*	*Q10 (p. 1141)*	*Q11 (p. 1142)*	*Q12 (p. 1145)*
Q13 (p. 1146)	*Q14 (p. 1147)*	*Q15 (p. 1149)*	*Q16 (p. 1154)*	*Q17 (p. 1157)*	*Q18 (p. 1160)*
Q19 (p. 1163)	*Q20 (p. 1164)*	*Q21 (p. 1165)*	*Q22 (p. 1166)*	*Q23 (p. 1167)*	*Q24 (p. 1168)*
Q25 (p. 1169)	*Q26 (p. 1175)*	*Q27 (p. 1175)*	*Q28 (p. 1176)*	*Q29 (p. 1179)*	*Q30 (p. 1180)*
Q31 (p. 1181)	*Q32 (p. 1183)*	*Q33 (p. 1184)*	*Q34 (p. 1186)*	*Q35 (p. 1188)*	*Q36 (p. 1189)*
Q37 (p. 1190)	*Q38 (p. 1191)*				

Q1: What is the probable diagnosis for a 30-year-old man who has a previous history of wearing contact lenses, and is currently experiencing pain, redness, and the presence of pus coming from his right eye? During examination, a greyish-white ulcer in the shape of a snake was observed on the lower portion of the cornea. Additionally, a hypopyon measuring 4 mm was found in the anterior chamber, along with a moderate flare reaction. (Q859337)

A. Viral corneal ulcer

B. Bacterial corneal ulcer

C. Fungal corneal ulcer

D. Parasitic corneal ulcer

Correct Answer: B. Bacterial corneal ulcer

Explanation: Bacterial corneal ulcers present with redness, pain, and discharge. Signs include a greyish-white serpiginous ulcer seen over the lower third of the cornea, indicative of the bacteria Aetiology of Purulent Corneal Ulcer: Corneal Epithelial Damage: Due to Corneal abrasion, epithelial drying (xerosis, exposure keratitis), necrosis (keratomalacia), desquamation (bullous keratopathy), and trophic changes (neuroparalytic keratitis). Pathogens Invading Intact Epithelium: Neisseria gonorrhoeae, Corynebacterium diphtheriae, Neisseria meningitidis, Haemophilus aegyptius, Listeria species. Sources of Infection: Exogenous Sources: Conjunctival sac, lacrimal sac (dacryocystitis), infected foreign bodies, vegetative material, waterborne or airborne infections. Common Bacteria Associated with Corneal Ulceration: General: Staphylococci , Pseudomonas, Streptococcus pneumoniae, Enterobacteriaceae, Neisseria. In Contact Lens Wearers: Pseudomonas is the most common cause. Acanthamoeba is another differential in contact lens misuse. All causative organisms: Gram-positive Cocci: Staphylococcus aureus, Staphylococcus epidermidis, Streptococcus pneumoniae. Gram-negative Cocci: Neisseria

gonorrhoeae, Neisseria meningitidis. Gram-positive Bacilli: Corynebacterium diphtheriae, C. xerosis, Bacillus cereus, Propionibacterium acne, Listeria, Clostridium. Gram-negative Bacilli: Pseudomonas aeruginosa, Enterobacteriaceae (Klebsiella, Proteus, E. coli, Serratia), Moraxella lacunata, Haemophilus influenzae. Gram-positive Filamentous Bacteria: Actinomyces, Nocardia. Mycobacteria: Non-tuberculous mycobacteria, M. tuberculosis. Clinical Features of Bacterial Corneal Ulcers: Symptoms: Pain and foreign body sensation. Watery eye (reflex hyper lacrimation). Photophobia (light intolerance). Blurred vision. Redness of the eye. Signs: Lid swelling. Blepharospasm (lid twitching). Conjunctival chemosis and hyperemia. Corneal ulcer characteristics: Early stage: Epithelial defect with greyish-white infiltrate . Established: Yellowish-white ulcer, swollen margins, necrotic floor, stromal oedema. Specific bacterial features: Staphylococcus aureus and Streptococcus pneumoniae: Oval, yellowish-white ulcer surrounded by clear cornea. Pseudomonas species: Irregular ulcer with greenish exudate, rapid spread, often with hypopyon. Enterobacteriaceae: Shallow ulcer with greyish-white suppuration, ring-shaped infiltrate. The anterior chamber may contain pus (Hypopyon). Muddy iris colour. Small pupil due to iritis. Possible raised intraocular pressure (inflammatory glaucoma). Incorrect Options Viral corneal ulcer (Option A) It is caused due to low immunity. A red, watery, and bloodshot eye is a prominent feature. There is severe eye pain and pus or other eye discharge. This can lead to vision loss and blindness. Fungal corneal ulcer (Option C) Trauma is one of the causes of Fungal corneal ulcer. There is less pain and Hypopyon if present is thick and immobile. Parasitic corneal ulcer (Option D) Is characterized by disproportionately severe pain and is caused by either contact lens wear or exposure to contaminated water.

References:

Kanski’s Clinical Ophthalmology, Edition 9, Page- 209, 210, 211.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Which of the subsequent options does not exhibit a characteristic of keratoconus? (Q855518)

A. Haab's striae

B. Munson's sign


C. Irregular astigmatism


D. Scissoring reflex

Correct Answer: A. Haab's striae

Explanation: Haab's striae are horizontal breaks or scars in Descemet's membrane and are typically associated with congenital glaucoma (especially buphthalmos) rather than keratoconus . These striae occur as a result of increased intraocular pressure and are not a feature of keratoconus. Keratoconus {{caption_text}} Definition : Progressive, non-inflammatory bilateral ectatic corneal condition affecting the axial part. Pathology : Involves progressive thinning and ectasia of the cornea due to defective synthesis of mucopolysaccharides and collagen. Clinical Features Symptoms : Defective vision due to progressive myopia and irregular astigmatism, uncorrectable by

glasses. Blurred and distorted vision, glare, halos around lights. Light sensitivity and ocular irritation. Visual loss mainly from irregular astigmatism and myopia; secondary loss due to corneal scarring. Examination Findings Corneal Thinning & Bulging : The Cornea thins near the centre and bulges forward, with the cone apex slightly below the corneal centre. Specific Signs : Distorted Window Reflex : Visual irregularity on corneal reflection. Placido Disc Irregularity : Circles appear irregular. Slit-Lamp Examination : Thinning and ectasia of the central cornea. Opacity at the apex, Fleischer Ring : Brownish ring around the cone, possibly due to haemosiderin in the epithelium. Folds in Descemet's and Bowman's membranes. Vogt Striae : Fine, parallel striae at the apex, best seen with a slit lamp, disappearing with external pressure (vs Haab striae: They are curvilinear breaks in Descemet's membrane, resulting acutely from stretching of the cornea in primary congenital glaucoma). {{caption_text}} Oil droplet reflex Retinoscopy : Yawning reflex (scissor reflex) with high oblique/irregular astigmatism. (Option D ruled out) Distant Direct Ophthalmoscopy : Annular dark shadow (oil droplet reflex) separating central and peripheral cornea. A ring of shadow concentric with the margin in the red reflex, resembling an oil droplet. Ophthalmoscope Findings : Irregular Astigmatism : Progressive myopia and irregular astigmatism in later stages, poorly corrected by glasses. (Option C ruled out) Munson's Sign : Localized bulging of the lower lid when looking down (seen in late stages). (Option B ruled out) Keratometer Findings : Malformed, malaligned, and malfocused mires. Corneal Topography : The most sensitive method for early detection (Forme Fruste Keratoconus) shows the shape of the corneal surface.





References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 248, 249.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: What is the most probable diagnosis for a 23-year-old male who has been experiencing eye pain, sensitivity to light, and redness for the last two days, following an injury caused by brushing his eyes against leaves and thorns while gardening? (Q857661)

A. Simplex keratitis

B. Bacterial keratitis

C. Fungal keratitis

D. Acanthamoeba keratitis

Correct Answer: C. Fungal keratitis

Explanation: The clinical scenario of pain, redness in eyes following trauma with vegetative matter is suggestive of Fungal Keratitis . Fungal Keratitis: Etiology Causative organism: Aspergillus fumigatus >> Fusarium Predisposing factor: Vegetative matter injury (trauma) (Option C) Clinical Features SIGNS >> symptoms Pain & Foreign Body Sensation Watering/ Photophobia Blurred Vision, Redness Examination Findings Corneal Ulcer: Greyish-white , dry, elevated, rolled-out margins Pigmented Ulcer: Brownish, dematiaceous fungi Feathery Extensions: Finger-like projections in stroma Sterile Immune Ring: Yellow line of demarcation Satellite Lesions: Small lesions around the main ulcer Hypopyon: Large, immobile, non-sterile , even in small ulcer Perforation: Rare but possible Investigations Corneal Scraping: For microbiological examination Wet KOH Mount: Filamentous fungi or yeast-like fungi (Candida) Culture: Sabouraud's agar for fungal culture PCR: For rapid identification of pathogens Corneal Biopsy: For deep stromal abscess or negative cultures Treatment Topical Antifungal Agents: Natamycin 5% , Voriconazole, Amphotericin-B, Nystatin Oral Antifungal Agents: Ketoconazole, Voriconazole (for severe ulcers) Surgical Management: Repeated Scraping: Removal of necrotic material Cauterization Options: Carbolic acid (100%), Trichloroacetic acid (10%–20%), Povidone-iodine (5%) Therapeutic Keratoplasty: Corneal transplantation Bacterial and Fungal Keratitis: BACTERIAL KERATITIS (Option B) FUNGAL KERATITIS Etiology S. aureus (worldwide) S. epidermidis (India) Aspergillus Examination Findings Fluorescein stain positive ulcer Hypopyon (sterile) Stromal edema Descemet’s membrane folds Keratic precipitates due to anterior uveitis Signs >>> symptoms Dry/rough ulcer Has projecting ends Thick immobile unsterile hypopyon Satellite nodules are infective and have immune reaction- Wessley ring IOP is high Treatment Empirical monotherapy: Moxifloxacin, gatifloxacin, besifloxacin Duo therapy: Gram positive + gram negative. Combination of Cefazolin with Gentamicin or Cefuroxime with Gentamicin. Cycloplegics and mydriatics Antifungals: Natamycin 5% , amphotericin B Steroids are contraindicated Simplex Keratitis (Option A): Caused due to infection by Herpes simplex virus (HSV-1) . Unlike Fungal keratitis, Simplex keratitis is acquired by kissing or in close contact with an infected person. Dendritic ulcer with budding pattern is suggestive of Simplex keratitis. Acanthameoba keratitis (option D): Usually seen in contact lens users , pseudodendritic pattern with tapering ends is suggestive of Acanathamoeba Keratitis.

References:

Kanski’s Clinical Ophthalmology 9th Edition, Pages 217, 218.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q4: In the image provided, which refractive error is observed in the depicted condition? (Q857753)

A. Hypermetropia and astigmatism

B. Myopia and astigmatism

C. Myopia and hypermetropia

D. Hypermetropia



Correct Answer: B. Myopia and astigmatism

Explanation: The image given above is that of Munson’s sign which is the V-shaped conformation of the lower lid produced by the ectatic cornea in downgaze seen in Keratoconus . The refractive error seen in keratoconus is myopia and irregular astigmatism leading to defective vision. Keratoconus: Description Progressive, non-inflammatory bilateral ectatic corneal condition affecting the axial part. Pathology Involves progressive thinning and ectasia of the cornea due to defective synthesis of mucopolysaccharides and collagen. Clinical Features Defective vision due to progressive myopia and irregular astigmatism (uncorrectable by glasses) (Option B) Blurred and distorted vision, glare, and halos around lights Light sensitivity and ocular irritation Visual loss primarily from irregular astigmatism and myopia, secondary loss from corneal scarring Examination Findings Distorted Window Reflex: Visual irregularity on corneal reflection. Placido Disc Irregularity: Irregular circles. Slit-Lamp Examination: Thinning and ectasia of the central cornea. Opacity at the apex, Fleischer Ring (brownish ring around the cone due to haemosiderin in the epithelium). Folds in Descemet's and Bowman's membranes. Vogt Striae: Fine, parallel striae at the apex, disappearing with external pressure. Oil Droplet Reflex: Annular dark shadow separating central and peripheral cornea in distant direct ophthalmoscopy. Retinoscopy: Yawning reflex (scissor reflex) with high oblique/irregular astigmatism. Ophthalmoscope Findings: Irregular astigmatism and progressive myopia, poorly corrected by glasses in later stages. Munson's Sign: Localized bulging of the lower lid when looking down (seen in late stages). Keratometer Findings: Malformed, malaligned, and malfocused mires. Corneal Topography: Sensitive method for early detection, shows corneal surface shape. Management Spectacle Correction: Effective in the early stages, but contact lenses are more beneficial. Less effective in later stages due to irregular astigmatism. Contact Lenses: Rigid gas permeable lenses improve vision in early cases. Specially designed scleral lenses (e.g., Rose-K) are useful in early to moderate cases. Corneal Collagen Cross-Linking (CXL): New treatment modality to slow disease progression. Involves Riboflavin (0.1%) eye drops and UVA radiation . Riboflavin induces cross-linking of corneal collagen to stabilize the stroma and improve contact lens tolerance. Keratoplasty : Recommended if disease progresses or in cases of acute hydrops (Descemet's membrane tear). Reserved for more advanced diseases or when other treatments fail. Penetrating keratoplasty is common, but

Deep Anterior Lamellar Keratoplasty (DALK) is preferred if Descemet's membrane and endothelium are intact, reducing rejection risk. Intracorneal Ring Segments (INTACS) : Effective in early to moderate cases by flattening the cornea. Less effective in halting progression.

References:

Kanski’s Clinical Ophthalmology 9th Edition 9, pages 248-250.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: Which of the following statements is incorrect regarding the corneal epithelium? (Q856916)

A. Regeneration is limited to the limbus


B. Bowman’s membrane regenerates


C. Lined by stratified squamous epithelium

D. Apical cells have microvilli

Correct Answer: B. Bowman’s membrane regenerates

Explanation: ndothelium The innermost layer consists of a single layer of flat, hexagonal cells. These cells are crucial for maintaining corneal transparency by regulating fluid balance within the cornea through an active pump mechanism. {{caption_text}} {{caption_text}}





References:

Kanski's Clinical Ophthalmology, Edition 9, Page- 204.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Which layer is accountable for preserving the moisture and clarity of the cornea? (Q855015)

A. Descemet’s membrane


B. Endothelium


C. Stroma

D. Corneal epithelium

Correct Answer: B. Endothelium

Explanation: Layers of Cornea Epithelium (Option D ruled out) The outermost layer comprises 5-6 layers of non-keratinized, stratified squamous epithelial cells. Acts as a barrier , protecting the cornea and preventing tear fluid from entering the underlying stroma. The basal cells at the limbus are stem cells responsible for regenerating the corneal epithelium. Bowman’s Membrane Located just beneath the epithelium, this acellular layer is 12 µm thick and composed of densely packed collagen fibrils. It is a protective barrier to the stroma but cannot regenerate if damaged and heals by scarring. Stroma (Option C ruled out) Making up about 90% of the cornea's thickness (approximately 0.5 mm thick), the stroma consists of collagen fibrils arranged in parallel layers. These fibrils are embedded in a hydrated matrix of proteoglycans - Chondroitin and Keratan Sulphate (keratan sulphate maintains corneal transparency). Pre-Descemet's Membrane (Dua's Layer) This layer, discovered in 2013, is about 15 µm thick, acellular , and lies just above Descemet's membrane. It is strong and resistant to air penetration. Descemet's Membrane (Option A ruled out) A robust basement membrane of the corneal endothelium is made of collagen and glycoproteins. It is tense, but when torn, curls inwards on itself. It can regenerate if damaged and is resistant to trauma and pathological processes. Endothelium (Option B) The innermost layer consists of a single layer of flat, hexagonal cells. These cells are crucial for maintaining corneal transparency by regulating fluid balance within the cornea through an active pump mechanism. {{caption_text}} {{caption_text}}





References:

Kanski's Clinical Ophthalmology, Edition 9, Page- 204.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: What is the distinguishing characteristic of a fungal ulcer? (Q855017)

A. Hypopyon


B. Dendritic ulcer on a fluorescein dye

C. Ring abscess

D. Satellite lesion

Correct Answer: D. Satellite lesion

Explanation: Keratomycosis {{caption_text}} Clinical Features Pain & Foreign Body Sensation Watering of the eye Photophobia Blurred Vision Redness. Examination Findings Corneal Ulcer : Greyish-white, dry-looking with elevated, rolled-out margins. Pigmented Ulcer : Brownish appearance, possible with dematiaceous fungi. Feathery Extensions : Delicate, finger-like projections into the surrounding stroma. Sterile Immune Ring : Yellow line of demarcation where fungal antigen and host antibodies meet. Satellite Lesions (Option D) : Multiple small lesions around the main ulcer. Hypopyon (Option A ruled out) : Large, non-sterile, even with a small ulcer. Endothelial Plaque : Located under the stromal lesion, made of fibrin and leucocytes. Corneal Vascularization : Typically absent in pure fungal ulcers. Perforation : Rare but possible. Investigations Corneal Scraping : Performed under topical anaesthesia for microbiological examination. Wet KOH Mount : Reveals filamentous fungi (branched, septate hyphae) or non-filamentous yeast-like fungi (Candida). Gram's & Giemsa Stains : To detect fungal hyphae and inflammatory cells. Culture : Sabouraud's agar medium is used for fungal culture. PCR : Recommended for rapid identification of fungal pathogens. Anterior Chamber Paracentesis : Considered if there is an intraocular extension. Corneal Biopsy : Indicated for deep stromal abscess or when cultures are negative. Confocal Microscopy : Used for identifying actual fungi within the cornea . Incorrect Options: Dendritic ulcer on a fluorescein dye (Option B) is a characteristic finding of herpetic keratitis caused by the herpes simplex virus. Ring abscess (Option C) is seen in Acanthamoeba keratitis commonly seen in patients using contact lens and using tap water to clean it.



References:

Kanski's Clinical Ophthalmology, Edition 9, Page- 216, 217.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q8: A 64-year-old woman with a history of trauma to the left eye 5 days back now presented with complaints of complete loss of vision. Upon examination the following findings were seen. What should be the subsequent course of action to manage this condition? (Q859303)

A. Keratoplasty

B. Keratotomy


C. LASIK

D. DCR




Correct Answer: A. Keratoplasty

Explanation: The history of trauma to the eye and complete loss of vision and examination finding revealing congestion and corneal perforation is suggestive of uncontrolled keratitis commonly seen with bacterial keratitis. The choice of treatment for corneal perforation with uncontrolled infection penetrating is keratoplasty. Penetrating Keratoplasty (PK): It is a type of Allografting (Allo-Keratoplasty) which is the replacement of a patient's diseased cornea with a healthy donor cornea. Full-thickness grafting. {{caption_text}} {{caption_text}} {{caption_text}} Indications for Penetrating Keratoplasty: Optical Corneal opacity Bullous keratopathy Corneal dystrophies Advanced keratoconus Therapeutic To replace an inflamed cornea that does not respond to conventional therapy Tectonic Graft To restore the integrity of the eyeball, e.g., after corneal perforation (Option A) or in cases of significant corneal thinning. Cosmetic To improve the appearance of the eye Complications: (ak142) Early Complications: Flat anterior chamber Iris prolapse Infection Secondary glaucoma Epithelial defects Primary graft failure Late Complications: Graft rejection Recurrence of disease Astigmatism Incorrect Options Keratotomy (Option B) is used for correcting refractive error causing flattening of the cornea and reducing its focusing power. LASIK (Option C) surgery is commonly used for keratomileusis, which is altering of the curvature of the cornea, done for refractive correction. . Dacryocystorhinostomy (DCR) (Option D) is a surgical procedure done for dacryocystitis.







References:

Kanski's Clinical Ophthalmology, Edition 9, Page- 277

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Which glycosaminoglycan is responsible for the transparency of the cornea? (Q859334)

A. Keratan sulfate

B. Hyaluronic acid

C. Heparan sulfate

D. Dermatan sulfate

Correct Answer: A. Keratan sulfate

Explanation: Keratan sulfate and chondroitin sulfate are the major glycosaminoglycan present in the corneal stroma. They exist in the proteoglycan form and are responsible for maintaining the space between the regularly oriented layers of collagen fibrils in the stroma. Locations of Various Mucopolysaccharides Mucopolysaccharide Primary Locations Functions Hyaluronic Acid (Option B) Connective tissues, tendons, synovial fluid, vitreous humor Serves as a lubricant in joint cavities and contributes to the structure of connective tissues Heparin (Option C) Liver, lungs, spleen, monocytes Acts as an anticoagulant by activating antithrombin III, which inactivates thrombin and other clotting factors Chondroitin Sulfate Cartilage, bone, tendons, cornea, skin Provides structural support to connective tissues and contributes to their resilience and elasticity Keratan Sulfate (Option A) Cornea, cartilage Dermatan Sulfate (Option D) Skin, blood vessels, heart valves Contributes to the structure of the skin, blood vessels, and heart valves

References:

Harper’s Biochemistry Edition 32; Page 610.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: A girl returns from a picnic and develops severe eye pain. She had been using contact lenses but did not follow proper hygiene practices. What is the most likely causative organism? (Q866848)

A. Acanthamoeba

B. Plasmodium

C. Pseudomonas aeruginosa

D. Herpes simplex virus

Correct Answer: A. Acanthamoeba

Explanation: Improper contact lens hygiene, especially with exposure to contaminated water (picnic scenario, possible swimming), is a major risk factor for Acanthamoeba keratitis . Acanthamoeba is a free-living protozoan found in water, soil, and air; it can adhere to contact lenses and invade the corneal epithelium, causing severe pain (often disproportionate to clinical signs), redness, tearing, and photophobia. Classic sign: ring-shaped stromal infiltrate in advanced cases. Acanthamoeba keratitis: Severe pain out of proportion to clinical findings. History of contact lens wear, especially poor hygiene or water exposure. Resistant to routine antibacterial therapy; requires specific anti-amoebic agents (e.g., polyhexamethylene biguanide, chlorhexidine). Delayed diagnosis can lead to corneal perforation and vision loss. Examination Findings in Acanthamoeba Keratitis: Symptoms: Blurred Vision: Common in corneal conditions due to the disruption of the cornea's transparent surface. Discomfort: Significant discomfort in the affected eye. Severe Pain: Pain is often out of proportion to clinical signs, a key clinical clue. Signs: Early Disease: Irregular and Greyish Epithelium : Early sign indicating amoebic invasion. Pseudodendrites : Lesions resembling herpetic keratitis Limbitis and Stromal Infiltrates : Inflammation at the corneal-scleral junction with diffuse or focal infiltrates in the anterior stroma. Perineural Infiltrates (Radial Keratoneuritis) : Hallmark sign, radial streaks extending towards the cornea's centre, pathognomonic for Acanthamoeba keratitis. Advanced Disease: Ring Abscess : Characteristic of advanced infection, where infiltrates form a ring-shaped abscess. Stromal Opacification and Vascularisation : Corneal opacity and new blood vessel growth, leading to significant vision impairment. Corneal Melting : Serious complication where corneal tissue breaks down, often at the infiltrate edges. Scleritis : Reactive inflammation of the sclera, not a direct infection extension. Investigations and Treatment of Acanthamoeba Keratitis: Investigations: Corneal Scraping : The primary method for obtaining tissue for examination. Scrape the margins and base of the lesion after applying a non-preserved topical anaesthetic. Staining : PAS (Periodic acid–Schiff) and Calcofluor white : Highly effective for visualizing Acanthamoeba cysts. Gram and Giemsa stains : Less sensitive but can also demonstrate cysts. Culture : Definitive diagnosis by culturing on non-nutrient agar with dead E. coli. Additional Investigations : Immunohistochemistry : Uses antibodies to detect specific Acanthamoeba antigens. PCR : Useful for detecting Acanthamoeba DNA. In Vivo Confocal Microscopy : Provides detailed images of Acanthamoeba cysts within the cornea. Corneal Biopsy : Considered if other diagnostic methods are inconclusive. Treatment Modalities: Debridement : Removal of infected epithelium to enhance topical medication penetration. Topical Amoebicides : PHMB (polyhexamethylene biguanide) 0.02% and Chlorhexidine 0.02% : Effective against both trophozoites and cysts. Hexamidine : More effective than Propamidine (Brolene) . Topical Steroids : Considered for persistent inflammation after two weeks of anti-amoebic treatment. Caution against use if corneal thinning is present. Pain Management : Oral NSAIDs are recommended for pain relief. Therapeutic Keratoplasty : Required for cases resistant to medical therapy or when corneal perforation occurs. Plasmodium (Option B): Causes malaria; does not cause keratitis or infection related to contact lenses. Pseudomonas aeruginosa (Option C): Most common bacterial cause of contact lens–related keratitis; produces rapid corneal ulceration, but here the focus is on protozoal infection linked to contaminated water exposure. Herpes simplex virus (Option D): Causes dendritic keratitis; unrelated to contact lens hygiene and not typically triggered by recent outdoor water exposure.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: Which of the following is true about pterygium? (Q859749)

A. Inflammation of cornea


B. Fatty deposition of sclera


C. Conjunctival invasion making a flap over cornea


D. Dead epithelial debris accumulation


Correct Answer: C. Conjunctival invasion making a flap over cornea

Explanation: Pterygium/Surfer’s eye Definition Pterygium is a triangular encroachment of vascularized granulation tissue covered by conjunctiva in the interpalpebral area . {{caption_text}} Parts: Cap, head, neck, body Degenerative process Aetiology Common in dry, sunny climates with sandy soils Ultraviolet light is a probable factor Usually elderly persons Pathology {{caption_text}} Histology shows collagenous degenerative changes in the vascularized subepithelial stroma. Pathogenesis Degenerative subconjunctival proliferation Granulation tissue proliferates as avascularized tissue encroaches on the cornea. Destroys the superficial layers of the stroma and Bowman’s membrane Symptoms Frequently follows a pinguecula Often bilateral Usually on the nasal side Impairment of vision due to: Progression into the pupillary area of the cornea Induced astigmatism by corneal flattening in the axis of pterygium Progressive pterygium: thick and vascular Regressive pterygium: thin, pale, and atrophic Ceases to grow but never disappears A probe can be passed under the lesion Can present as inflammed pterygium {{caption_text}} Signs A probe cannot be passed under the lesion Fusch’s spots/Islets of Vogt: small discrete whitish flecks made of pterygial epithelial cells seen in progressive pterygium and are also called the cap of the pterygium. Stocker line: linear epithelial iron deposition, seen in regressive pterygium. {{caption_text}} Management of pterygium: Management Best left alone Medical treatment: Tear substitutes for small regressive pterygium with dry eye Topical steroids for associated inflammation Protection from UV rays with sunglasses Surgical excision for: Progression of lesion towards the pupillary area causing excessive astigmatism and visual impairment Restricted ocular movements Disfiguring Diplopia Procedures Bare sclera technique: The pterygium is excised from the cornea and conjunctiva, leaving a bare sclera Associated with a high recurrence rate Autoconjunctival graft A piece of limbal conjunctival tissue from the same or the other eye or An amniotic membrane graft is used to cover the bare sclera Sutured at the limbus to cover the defect Surgical excision with lamellar keratectomy and lamellar keratoplasty for deeply infiltrating recurrent recalcitrant pterygia Old methods (not used now): McRaynold’s operation Postoperative use of beta irradiations Post-operative therapy and care Prevent recurrence: Mitomycin C 0.02% drops Complications like scleral necrosis, cataracts, and iritis


@Neet_pg_bot


@Neet_pg_bot





References:

Kanski's Clinical Ophthalmology, 9th edition, Page- 198, 199.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: For which of the subsequent diseases is collagen cross-linking therapy utilized? (Q859767)

A. Keratoconus

B. Corneal ulcer

C. Band keratopathy

D. Uveitis

Correct Answer: A. Keratoconus

Explanation: Treatment of Keratoconus: Spectacle Correction : Effective in the early stages, but contact lenses are more beneficial. Less effective in later stages due to irregular astigmatism. Contact Lenses : Rigid gas permeable lenses improve vision in early cases. Specially designed scleral lenses (e.g., Rose-K) are useful in early to moderate cases. Corneal Collagen Cross-Linking (CXL) (Option A) : New treatment modality to slow disease progression. Involves riboflavin (0.1%) eye drops and UVA radiation. Riboflavin induces cross-linking of corneal collagen to stabilize the stroma and improve contact lens tolerance. Keratoplasty : Recommended if disease progresses or in cases of acute hydrops (Descemet's membrane tear). Reserved for more advanced diseases or when other treatments fail. Penetrating keratoplasty is common, but Deep Anterior Lamellar Keratoplasty (DALK) is preferred if Descemet's membrane and endothelium are intact, reducing rejection risk. Intracorneal Ring Segments (INTACS) : Effective in early to moderate cases by flattening the cornea. Less effective in halting progression. Incorrect Options: Corneal ulcer (Option B) are treated with antibiotics, antivirals, or antifungals, depending on the cause. Band keratopathy (Option C) involves calcium deposition in the superficial cornea and is treated with chelation therapy or surgical interventions like superficial keratectomy. Uveitis (Option B) is inflammation of the uveal tract and is typically managed with corticosteroids, immunosuppressive agents, or biologics.

References:

Kanski’s Clinical Ophthalmology, Edition 9, Page- 249, 250.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q13: What's the main risk for a 37-year-old man with blurred vision and severe pain in his right eye? He revealed that recently, he went swimming in a pond. During examination, a ring-like lesion was found, as seen in the image. (Q859769)

A. Diabetes

B. Hypertension

C. Steroids

D. Soft Contact lenses



Correct Answer: D. Soft Contact lenses

Explanation: The clinical presentation of blurred vision, severe pain after swimming in a pond with ring like lesion on examination is suggestive of Acanthamoeba keratitis. Acanthamoeba keratitis: It is caused by Acanthamoeba , a free-living amoeba found in water, causes severe corneal infections Risk Factors: Contact Lens Use (Primary Risk Factor) Exposure to Contaminated Water Trauma Immunosuppression Use of Homemade Saline Solutions. Poor Lens Storage Conditions Examination Findings in Acanthamoeba Keratitis: Symptoms: Blurred Vision: Common in corneal conditions due to the disruption of the cornea's transparent surface. Discomfort: Significant discomfort in the affected eye. Severe Pain: Pain is often out of proportion to clinical signs, a key clinical clue. Signs: Early Disease: Irregular and Greyish Epithelium : Early sign indicating amoebic invasion. Pseudodendrites : Lesions resembling herpetic keratitis Limbitis and Stromal Infiltrates : Inflammation at the corneal-scleral junction with diffuse or focal infiltrates in the anterior stroma. Perineural Infiltrates (Radial Keratoneuritis) : Hallmark sign, radial streaks extending towards the cornea's centre, pathognomonic for Acanthamoeba keratitis. Advanced Disease: Ring Abscess : Characteristic of advanced infection, where infiltrates form a ring-shaped abscess. Stromal Opacification and Vascularisation : Corneal opacity and new blood vessel growth, leading to significant vision impairment. Corneal Melting : Serious complication where corneal tissue breaks down, often at the infiltrate edges. Scleritis : Reactive inflammation of the sclera, not a direct infection extension. Options A, B, and C are incorrect.

References:

Kanski’s Clinical Ophthalmology, Ed 9, Pg 211,216,219,232

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: What is the diagnosis for a female patient, aged 65, who has a history of chickenpox and presents at the clinic with the following findings? (Q859772)

A. Chicken pox exacerbation

B. Drug allergy to acyclovir

C. Herpes zoster ophthalmicus

D. Herpes zoster oticus



Correct Answer: C. Herpes zoster ophthalmicus

Explanation: The clinical scenario of a patient with history of chickenpox and cutaneous lesions seen along the course of trigeminal nerve is suggestive of Herpes zoster ophthalmicus. Herpes Zoster Ophthalmicus (HZO) Affects the ophthalmic branch of the trigeminal nerve. Anterior uveitis occurs in 40-50% of cases, usually within 2 weeks of skin rash onset. Symptoms: mild iritis, vesicular rash on the nose, small KPs, mild aqueous flare, possible hemorrhagic hypopyon. Complications: iris atrophy, secondary glaucoma, complicated cataracts. Incorrect Options: Chicken pox exacerbation (Option A) , caused by the varicella-zoster virus , usually occurs in children and presents with a generalized vesicular rash . Reactivation of the virus in adults leads to herpes zoster rather than an exacerbation of chickenpox. Drug allergy to acyclovir (Option B) , a common antiviral medication used to treat herpes zoster , would typically present with skin rashes, itching, or swelling after drug exposure. Herpes zoster oticus (Option D) (Ramsay Hunt syndrome) involves the facial nerve and affects the ear, causing pain, vesicular rash in the ear canal , and facial paralysis . It does not primarily affect the eye or lead to keratitis.

References:

Kanski's Clinical Ophthalmology, 9th edition, Page 468-71

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: Arlt's line is seen in? (Q859776)

A. Vernal keratoconjunctivitis


B. Phlyctenular keratoconjunctivitis

C. Trachoma


D. Pterygium

Correct Answer: C. Trachoma

Explanation: Signs in Trachoma: Signs of active trachoma Signs of cicatricial trachoma Conjunctival signs: Conjunctival congestion Follicles in the upper tarsal conjunctiva: Look like sago-grains Papillary enlargement: red, velvety Conjunctival signs: Scarring It may be irregular, star-shaped, or linear Linear scar in the sulcus subtarsalis is called Arlt's line (Option C) accumulation of dead epithelial cells and inspissated mucus in the glands of Henle Pseudocyst Xerosis Symblepharon Corneal signs: Superficial keratitis on slit lamp and fluorescein Herbert's follicles Progressive pannus/ Trachomatous pannus : The upper part of the cornea is cloudy Corneal ulcers are common at the advancing edge of the pannus Shallow, little infiltrated Cause lacrimation and photophobia Corneal signs: Regressive pannus or pannus siccus : Receding infiltration Vessels extend beyond the area of infiltration and haze Herbert pits : oval or circular pitted scars formed after the healing of Herbert's follicles in the limbal area Vascularized and opaque cornea. Corneal xerosis Corneal ectasia Total corneal pannus (blinding sequelae) Lid signs: Distortion of the lids Tylosis : Thickening of the tarsal plate/lid margin Trichiasis due to rubbing of lashes on the cornea Trachomatous ptosis (sleepy appearance) Entropion Madarosis Ankyloblepharon Lacrimal apparatus sequelae: Chronic dacryocystitis Chronic dacryoadenitis Images: Active Trachoma {{caption_text}} Follicles in upper palpebral conjunctiva {{caption_text}} Trachomatous pannus Progressive pannus Cicatricial Trachoma {{caption_text}} ■■■■■■ {{caption_text}} Regressive pannus {{caption_text}} Conjunctival concretions {{caption_text}} Trichiasis {{caption_text}} Tylosis (thickening of tarsal border of eyelid) Incorrect Options: Vernal keratoconjunctivitis (Option A) is characterized by intense itching, redness, watery discharge, and cobblestone appearance of upper palpebral conjunctiva. Phlyctenular keratoconjunctivitis (Option B) is characterized by the formation of small nodules called as phlycten near limbus or ulcers on the cornea. Pterygium (Option D) is conjunctival degeneration characterized by triangular fibrovascular subepithelial ingrowth of bulbar conjunctiva over the limbus.



@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot



References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 173, 174, 175.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: What is the main underlying factor responsible for regular astigmatism when studying the optics of a patient? (Q860009)

A. Abnormal curvature of the cornea

B. Excessive eye rubbing


C. Prolonged exposure to bright light


D. Vitamin A deficiency

Correct Answer: A. Abnormal curvature of the cornea


Explanation: The primary cause of regular astigmatism is an abnormal curvature of the cornea , which results in different refractive powers along different meridians of the eye. This irregularity in corneal shape prevents light rays from focusing on a single point on the retina, causing blurred or distorted vision. Astigmatism: A refractive

error where different meridia of the eye have different refractive errors . Divided into 2 types: Regular and irregular I. Regular Astigmatism: When refraction error changes uniformly from one meridian to the other. (2 principle meridians present) Type Description With-the-Rule (WTR) The vertical meridian has greater curvature than the horizontal meridian. Against-the-Rule (ATR) The horizontal meridian is more curved than the vertical meridian. Oblique Astigmatism Principal meridians are at oblique angles (e.g., 45° and 135°), still perpendicular to each other. Bi-oblique Astigmatism Principal meridians are not perpendicular to each other (e.g., 30° and 100°). Regular Astigmatism Based on Focal Lines: Type Description Simple Astigmatism One meridian focuses on the retina; the other focuses in front (simple myopic) or behind (simple hypermetropic). Compound Astigmatism Both meridians focus light either in front (compound myopic) or behind (compound hypermetropic) the retina. Mixed Astigmatism One meridian focuses in front of the retina (myopic), and the other behind (hypermetropic). II. Irregular Astigmatism Involves multiple, changing meridians of refractive power, making it geometrically complex.Usually due to Corneal scars. Excessive eye rubbing (Option B) can irritate the eyes and potentially lead to infections like keratoconjunctivitis. Prolonged exposure to bright light (Option C) can cause eye strain and discomfort. Vitamin A deficiency (Option D) can lead to xerophthalmia characterized by night blindness and dryness of the eyes.






@Neet_pg_bot



@Neet_pg_bot



References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no:31-33

Astigmatism

Sturm’s Conoid

↑ *Back to this question in Part A*

@Neet_pg_bot


Q17: What does this image indicate? (Q856040)

A. Keratoconus

B. Vogt's limbal girdle

C. Keratoglobus

D. Corneal dystrophy




Correct Answer: A. Keratoconus

Explanation: - Keratoconus: The image most likely represents keratoconus , characterized by the thinning and cone-shaped bulging of the cornea, leading to visual distortion and irregular astigmatism . {{caption_text}} {{caption_text}} Keratoconus Definition : Progressive, non-inflammatory bilateral ectatic corneal condition affecting the axial part. Pathology : Involves progressive thinning and ectasia of the cornea due to defective synthesis of mucopolysaccharides and collagen. Clinical Features Symptoms : Defective vision due to progressive myopia and irregular astigmatism, uncorrectable by glasses. Blurred and distorted vision, glare, halos around lights. Light sensitivity and ocular irritation. Visual loss mainly from irregular astigmatism and myopia; secondary loss due to corneal scarring. Examination Findings Corneal Thinning & Bulging : The Cornea thins near the centre and bulges forward, with the cone apex slightly below the corneal centre. Specific Signs : Distorted Window Reflex : Visual irregularity on corneal reflection. Placido Disc Irregularity : Circles appear irregular. Slit-Lamp Examination : Thinning and ectasia of the central cornea. Opacity at the apex, Fleischer Ring : Brownish ring around the cone, possibly due to haemosiderin in the epithelium. Folds in Descemet's and Bowman's membranes. Vogt Striae : Fine, parallel striae at the apex, best seen with a slit lamp, disappearing with external pressure (vs Haab striae: They are curvilinear breaks in Descemet's membrane, resulting acutely from stretching of the cornea in primary congenital glaucoma). {{caption_text}} Oil droplet reflex Retinoscopy : Yawning reflex (scissor reflex) with high oblique/irregular astigmatism. Distant Direct Ophthalmoscopy : Annular dark shadow (oil droplet reflex) separating central and peripheral cornea. A ring of shadow concentric with the margin in the red reflex, resembling an oil droplet. Ophthalmoscope Findings : Irregular Astigmatism : Progressive myopia and irregular astigmatism in later stages, poorly corrected by glasses. Munson's Sign : Localized bulging of the lower lid when looking down (seen in late stages). Keratometer Findings : Malformed, malaligned, and malfocused mires. Corneal Topography : The most sensitive method for early detection (Forme Fruste Keratoconus) shows the shape of the corneal surface. Vogt's limbal girdle (Option B) : This refers to a white or grayish band at the corneal margin, but it is not associated with corneal thinning or bulging, as seen in keratoconus. Keratoglobus (Option C) :

This is a rare condition where the cornea is uniformly thin and globular in shape, but it is not associated with the same irregular astigmatism and conical shape as keratoconus. Corneal dystrophy (Option D) : This is a group of inherited disorders that cause abnormal deposits in the cornea, but the presentation is different from keratoconus. It does not typically involve the corneal thinning and cone shape seen in keratoconus.







References:

Kanski's Clinical Ophthalmology, Edition 9, Page- 248, 249.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A 27-year-old male with myopic astigmatism presents with an acute onset of pain and a reduction in visual acuity. He has a history of frequent changes of glasses and eye rubbing. An image of the eye just after the procedure is shown below. Name the procedure that has been done? (Q879000)

A. DSAEK

B. DALK


C. Intracameral C3F8 injection

D. Corneal collagen crosslinking with riboflavin




Correct Answer: C. Intracameral C3F8 injection

Explanation: A young man with myopic astigmatism, frequent spectacle changes, and eye rubbing develops sudden painful visual loss—classic for acute hydrops in keratoconus . The image shows a gas bubble in the anterior chamber , confirming intracameral C3F8 injection , which tamponades the Descemet's membrane tear and speeds resolution of corneal edema Incorrect Options: DSAEK (Descemet Stripping Automated Endothelial Keratoplasty) (Option A): A type of endothelial keratoplasty done in endothelial disorders such as Fuchs endothelial dystrophy and pseudophakic bullous keratopathy. Replaces diseased endothelium and Descemet's membrane with donor endothelial layer. DALK (Deep Anterior Lamellar Keratoplasty) (Option B): For advanced keratoconus with healthy endothelium; replaces corneal stroma while preserving Descemet's membrane. Avoided in acute hydrops; Postop images show corneal sutures, not gas bubbles. ■■■■■ Corneal collagen crosslinking (CXL) with riboflavin (Option D): Used to halt progressive keratoconus by UV-A–riboflavin crosslinking. Contraindicated in acute hydrops due to epithelial breaks and marked stromal edema.




@Neet_pg_bot



@Neet_pg_bot





References:

Parson’s Diseases of the Eye, 23rd Edition, Page 499

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: Which of the following layers is associated with the beaten bronze appearance? (Q860188)

A. Endothelial Layer


B. Stroma

C. Sub-Epithelial Layer

D. Epithelial Layer

Correct Answer: A. Endothelial Layer

Explanation: Fuchs Endothelial Corneal Dystrophy is strongly associated with the "beaten-metal" appearance on Descemet's membrane. It is a common type of endothelial dystrophy (Option A) , particularly affecting older adults. {{caption_text}} Option B,C and D are not involved Fusch’s endothelial dystrophy



References:

Kanski’s Clinical Ophthalmology, Edition 9, Page- 259, 260, 261.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: A patient presents with a painless ulcer in the eye. On examination, a long, branching ulcer with desquamated epithelium is seen on the cornea. What is the most likely diagnosis? (Q866837)

A. Fungal ulcer

B. Dendritic ulcer

C. Bacterial corneal ulcer

D. Neurotrophic ulcer

Correct Answer: B. Dendritic ulcer

Explanation: A patient presenting with painless, branching ulcer and desquamated epithelium, points to the diagnosis of Herpes simplex keratitis is caused by Herpes simplex virus type 1 (HSV-1) . The characteristic feature is a dendritic (branching) ulcer with terminal bulbs , best seen with fluorescein staining under cobalt blue light. The ulcer is painless because the virus causes corneal anesthesia due to involvement of trigeminal nerve endings. It has d esquamated epithelium and irregular margins. Fungal ulcer (Option A): Usually follows trauma with vegetative matter; presents with pain, feathery margins, satellite lesions, and hypopyon. Neurotrophic ulcer (Option D): It is an oval or circular, punched-out ulcer , with smooth, rolled edges, due to loss of corneal sensation (often post-HSV, diabetes, trauma, trigeminal nerve lesion) and there is loss of epithelium without infiltration or branching. Bacterial corneal ulcer (Option C): Acute, painful, purulent ulcer with stromal infiltration and hypopyon; rapid progression.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: In keratoplasty, the commonly used corneal storage medium and suture material are: (Q865358)

A. MK medium and 10-0 nylon

B. Moist chamber and Vicryl

C. Optisol-GS and silk

D. Eusol-C and 6-0 prolene

Correct Answer: A. MK medium and 10-0 nylon

Explanation: MK medium and 10-0 nylon (Option A) : While MK medium is less common now for long-term storage, 10-0 nylon is definitively the most commonly used suture material. This makes the suture part of this answer very strong. Moist chamber and Vicryl (Option B ruled out): Neither of these is commonly used for

corneal grafts. Optisol-GS and silk (Option C ruled out): Optisol-GS is the leading modern storage medium, but silk is not the commonly used suture for corneal grafts. Eusol-C and 6-0 prolene (Option D ruled out): Eusol-C is a valid medium, but 6-0 prolene is too thick for corneal suturing.

References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 135

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: Which condition is treated using an Intacs ring, as shown in the image? (Q866345)

A. Keratoconus

B. Cataract

C. Corneal ulcer

D. Glaucoma



Correct Answer: A. Keratoconus

Explanation: Keratoconus. In the above image, the device implanted is an Intacs corneal ring segment, which is used to flatten and reshape the cornea in patients with keratoconus. This helps improve vision by reducing corneal irregularities and correcting refractive error. The transparent ring segments are inserted into the corneal stroma to provide structural support and reduce cone steepness. Keratoconus is a progressive, non-inflammatory disorder of the cornea characterized by thinning and conical protrusion. Symptoms: Progressive blurring, ghost images, glare, frequent change in glasses prescription. Management options: Early stage: Spectacles or rigid gas permeable contact lenses Intermediate stage: Intacs corneal ring segments to reshape the cornea and delay corneal transplantation Advanced stage: Corneal transplantation (DALK or PKP) Intacs: Made of PMMA, placed in mid-peripheral corneal stroma; they reduce myopia and irregular astigmatism, improving best corrected visual acuity. Cataract (Option B): Lens opacity; treated with lens extraction + IOL, unrelated to corneal rings. Corneal ulcer (Option C): Infective/inflammatory; managed medically, not with implants. Glaucoma (Option D): Optic nerve damage from high IOP; managed with drugs, laser, or filtering surgery.



References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: A patient presents with guttate lesions in one eye and bullous keratopathy in the other eye. What is the most likely diagnosis? (Q866353)

A. Fuchs' endothelial dystrophy

B. Interstitial keratitis

C. Viral corneal ulcer

D. Keratoconjunctivitis

Correct Answer: A. Fuchs' endothelial dystrophy

Explanation: Fuchs' endothelial dystrophy The patient has guttate lesions in one eye and bullous keratopathy in the other. Guttate lesions indicate endothelial cell loss , while bullous keratopathy results from corneal edema due to endothelial dysfunction. These are classical features of Fuchs' endothelial dystrophy. Fuchs' endothelial dystrophy is a bilateral, progressive corneal endothelial disorder. Early sign: Guttae (drop-like excrescences on Descemet membrane) seen on slit lamp. Late stage: Stromal edema → epithelial edema → bullae formation (bullous keratopathy). Symptoms: Gradual, painless decrease in vision; glare; haloes; worse in the morning due to overnight edema. Treatment: Early: Hypertonic saline drops, hair dryer to reduce morning edema Advanced: Keratoplasty (DSEK, DMEK, or PK) Interstitial keratitis (Option B) – Chronic stromal inflammation, usually post-infectious (e.g., syphilis, HSV); presents with stromal haze and vascularization, not guttate lesions or bullous keratopathy. Viral corneal ulcer (Option C) – Painful red eye with epithelial defect and stromal infiltrate; acute onset, not a chronic, painless, bilateral process. Keratoconjunctivitis (Option D) – Inflammatory condition involving cornea and conjunctiva; presents with redness, pain, photophobia, discharge; not consistent with long-standing endothelial degeneration.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: A corneal wisp test was performed, and the corneal reflex was elicited. Which of the following nerves is responsible for the afferent limb of this reflex? (Q852484)

A. Facial Nerve

B. Trigeminal Nerve

C. Oculomotor Nerve

D. Abducens Nerve

Correct Answer: B. Trigeminal Nerve

Explanation: The corneal reflex (blink reflex) is a protective response that involves both sensory (afferent) and motor (efferent) pathways: Afferent limb (sensory input): Trigeminal nerve (CN V1 - Ophthalmic division) The cornea is innervated by the nasociliary branch of the ophthalmic division (V1) of the trigeminal nerve. A light touch to the cornea (e.g., with a cotton wisp) stimulates sensory fibers, sending signals to the trigeminal sensory nucleus in the brainstem. Efferent limb (motor response): Facial nerve (CN VII) The facial nerve (CN VII) innervates the orbicularis oculi muscle, which causes eyelid closure in response to the stimulus. Reflex Afferent Nerve Efferent Nerve Accommodation II (Optic nerve) III (Oculomotor nerve) Corneal V■ (Ophthalmic,

nasociliary branch) Bilateral VII (Facial nerve, temporal branch - orbicularis oculi) Cough X (Vagus nerve) X (Vagus nerve, also phrenic and spinal nerves) Gag IX (Glossopharyngeal nerve) X (Vagus nerve) Jaw Jerk V■ (Mandibular branch, sensory from masseter) V■ (Mandibular branch, motor to masseter) Lacrimation V■ (Ophthalmic branch) (loss of reflex does not preclude emotional tears) VII (Facial nerve) Pupillary II (Optic nerve) III (Oculomotor nerve) Facial Nerve (CN VII) (Option A): The facial nerve is responsible for the efferent limb (motor response), not the afferent limb. It causes eyelid closure by contracting the orbicularis oculi muscle. Oculomotor Nerve (CN III) (Option C): CN III controls extraocular muscles and the levator palpebrae superioris (eyelid elevation), but it does not mediate the corneal reflex. Abducens Nerve (CN VI) (Option D): CN VI controls the lateral rectus muscle for eye abduction but is not involved in the corneal reflex.

References:

https://www.ncbi.nlm.nih.gov/books/NBK534247/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: A patient sustained blunt trauma to the eye 6 months ago and now presents with blurring of vision. What is the most likely condition? (Q852489)

A. Iridodialysis

B. Ankyloblepharaon

C. Vossius Ring

D. Cycloiriditis





Correct Answer: A. Iridodialysis


Explanation: Iridodialysis is detachment of iris from its root at the ciliary body. It results in D shaped pupil and a lack biconvex area seen at the periphery. Blunt trauma (also known as non-penetrating or blunt force trauma) Cause: A strong impact, a fall, or a physical assault with a blunt object, is most commonly the result of sports injuries and assaults. Features: Conjunctiva: Subconjunctival hemorrhage Cornea: Abrasions, corneal edema, Descemet's membrane tears. Hyphaema: Blood in the anterior chamber leading to corneal staining. Iris: Iridodialysis (dehiscence of the iris from the ciliary body at its root), iris sphincter tear, Ciliary body: Ciliary body tear leading to angle recession. Lens: Vossious Ring (in the shape of the pupil) (Option C) Rosette-Shaped Cataract (flower-shaped opacity), subluxation of lens. Globe rupture (caused by severe blunt trauma, indicated by iris prolapse) Vitreous haemorrhage : Pigment cells (‘tobacco dust’) can be seen floating in the anterior vitreous. Choroidal rupture Retina: Commotio Retinae (also known as Berlin's edema) , retinal dialysis (disinsertion from ora serrata), giant retinal Tear (horseshoe shaped) , macular hole. Injury to optic nerve Traumatic neuropathy with chorioretinal scarring (decreased visual acuity, visual field defect, afferent pupillary defect, color, and brightness). Ankyloblepharon (Option B): This refers to the abnormal fusion of the eyelids, typically a congenital issue or sometimes a result of trauma leading to scarring of the eyelids. Cycloiriditis (Option D): This is inflammation of the ciliary body and iris. While trauma can trigger this, it usually presents with pain, redness, and photophobia (sensitivity to light) in addition to blurred vision.



@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot





References:

Kanski’s Clinical Ophthalmology, 10th Edition, Pages 914-921.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: A patient presents with eyelid crusting and a thready sensation between the cornea and lids. What is the most likely diagnosis? (Q852500)

A. Chronic Blepharitis

B. Dry Eye Syndrome

C. Meibomian Gland Dysfunction

D. Conjunctivitis

Correct Answer: A. Chronic Blepharitis

Explanation: Chronic Blepharitis is an inflammation of the eyelid margins that can cause symptoms such as eyelid crusting and a thready sensation between the cornea and eyelids, often due to the accumulation of secretions along the lash line. The eyelids may feel gritty or have a foreign body sensation. Bacterial blepharitis: Definition Chronic infection of the anterior part of the lid margin Synonyms Chronic anterior blepharitis Staphylococcal blepharitis Causative Organisms Most commonly Staphylococci (coagulase positive) Rarely Streptococci, Propionibacterium acnes, and Moraxella Clinical Features Chronic irritation Itching Mild lacrimation Gluing of cilia Mild photophobia Yellow crusts at the root of cilia Small ulcers that bleed easily when crusts are removed Red, thickened lid margins with dilated blood vessels Mild papillary conjunctivitis and conjunctival hyperemia Treatment Lid Hygiene: Warm compresses Cotton buds dipped in dilute baby shampoo or 3% sodium bicarbonate solution Avoid touching Antibiotics Topical Steroids Ocular Lubricants Dry Eye Syndrome (Option B): While dry eyes can cause a foreign body sensation, it typically does not cause eyelid crusting. Dry eye symptoms are more related to discomfort from insufficient tear production, but the crusting is not a hallmark feature. Meibomian Gland Dysfunction (Option C): This condition can lead to dry eyes and eyelid discomfort due to insufficient oil production, but it doesn't usually cause eyelid crusting in the way that blepharitis does. Conjunctivitis (Option D): Conjunctivitis (inflammation of the conjunctiva) causes symptoms such as redness, discharge, and itching

References:

Kanski's Clinical Ophthalmology 9th edition page 70-72

↑ *Back to this question in Part A*

@Neet_pg_bot

Q27: A patient presents with redness, extreme eye pain, and diminished vision. The patient frequently wears contact lenses. Which of the following organisms is most likely responsible for her condition? (Q844050)

A. Staphylococcus aureus

B. Streptococcus pyogenes

C. Acanthamoeba

D. Listeria

Correct Answer: C. Acanthamoeba

Explanation: Acanthamoeba is most likely responsible for this patient's condition. Contact lens use (especially with tap water/poor hygiene) is a strong risk factor. Causes Acanthamoeba keratitis. Hallmark features: Severe pain out of proportion to clinical signs (Perineural invasion) Red eye Decreased vision Classic finding: Ring-shaped corneal infiltrate Incorrect Options: Staphylococcus aureus (Option A) → Common cause of bacterial conjunctivitis or keratitis, but pain usually not extreme Streptococcus pyogenes (Option B) → Can cause severe infections, but not classically associated with contact lenses Listeria (Option D) → Rare ocular involvement; more associated with meningitis, neonatal infections

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q28: Shield ulcer is seen in which of the following? (Q860516)

A. Spring catarrh

B. Phlyctenular conjunctivitis


C. Atopic keratoconjunctivitis


D. Autoimmune Conditions


Correct Answer: A. Spring catarrh

Explanation: Vernal keratoconjunctivitis or Spring catarrh It is a recurrent bilateral conjunctivitis that tends to occur with the onset of warm weather. It primarily affects young children and adolescents, particularly boys. Course The condition typically starts in childhood and may resolve on its own over 5–10 years. Symptoms Burning, Itching, Photophobia (sensitivity to light), and lacrimation (excessive tearing). Signs Palpebral Form : Hypertrophied palpebral conjunctiva with a cobblestone appearance; bluish-white color; dense fibrous tissue with thickened epithelium. Limbal or Bulbar Form : Opacification at the limbus, gelatinous thickening, and Horner–Trantas dots {{caption_text}} Pseudogerontoxon/cupid’s bow (which resembles arcus senilis) is a grayish-white ring at the limbus seen in severe limbal vernal keratoconjunctivitis (VKC). {{caption_text}} Shield

ulcers are a significant complication of VKC, particularly in severe cases. These microerosions typically form on the upper third of the cornea. {{caption_text}} Mixed form: Exhibits characteristics of both the tarsal (palpebral) and limbal subtypes, typically presenting in only one eye. Maxwell Lyon's sign is a pseudomembrane formed due to excessive mucus deposition on the papilla resulting in cobblestone papillae. {{caption_text}} Histology In the palpebral form, hypertrophied papillae are observed, rich in eosinophils and other inflammatory cells. Treatment Topical therapy: Antihistamines - Levocabastine & emedastine Mast cell stabilizers - Sodium cromoglicate Mild steroids - Fluorometholone and loteprednol etabonate, Acetylcysteine can control excess mucus. Local therapy : Subtarsal steroid injections and cryotherapy for severe cases. Surgical treatment: Excision of giant papillae and treatment of shield ulcers with debridement and amniotic membrane or bandage contact lens. Cold compresses and tinted glasses for comfort. Avoid eye rubbing Phlyctenular conjunctivitis (Option B): Allergy caused by endogenous antigens → Staph. aureus and TB infection It presents with watering and itching in the eyes O/E: nodule of phlycten near the limbus and conjunctival congestion Atopic keratoconjunctivitis (Option C): Common in temperate regions and winters Dennie-morgan folds present Presence of Madarosis and Hertoghe’s sign Autoimmune Conditions (Option D): Not typically associated with Shield Ulcer : Various autoimmune diseases can affect the eyes, such as rheumatoid arthritis, systemic lupus erythematosus, and Sjögren's syndrome, leading to conditions like dry eye, scleritis, and uveitis. However, shield ulcers are not commonly seen as a direct manifestation of these autoimmune conditions.






@Neet_pg_bot



References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 181, 182, 183, 184, 185, 186.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9376221/

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4108143/

https://www.nature.com/articles/eye198963.pdf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q29: A 40-year-old male patient, who works in an iron factory, presents to the clinic with complaints of visual disturbances. He reports that his vision has been gradually worsening and he has noticed a change in the color perception. On examination, the ophthalmologist observes the presence of iron deposits in the eye. What is the most likely ocular condition responsible for the iron deposits in the eye and the patient's visual disturbances? (Q820453)

A. Siderosis bulbi

B. Hemochromatosis

C. Retinitis pigmentosa

D. Age-related macular degeneration

Correct Answer: A. Siderosis bulbi

Explanation: Correct Option A - Siderosis bulbi This is the correct answer. Siderosis bulbi refers to the condition in which iron particles, usually from a previous trauma, are deposited in the eye's intraocular tissues leading to visual disturbances and discoloration. In this case, the patient's line of work in an iron factory could have exposed him to iron particles. Incorrect Options: Option B - Hemochromatosis: This is a genetic disorder that leads to excessive iron absorption from the diet. It affects various organs in the body, including the liver, heart, and pancreas. While it can cause a number of health problems, it is not typically associated with iron deposits in the eye or the visual disturbances described by the patient. Option C - Retinitis pigmentosa: This is a group of rare, genetic disorders that involve a breakdown and loss of cells in the retina—which is the light sensitive tissue that lines the back of the eye. It is not typically associated with iron deposits in the eye. Option D - Age-related macular degeneration: This is a common eye condition among people over 50. It causes blurred or reduced central vision, due to thinning of the macula, the part of the retina responsible for clear vision in your direct line of sight. It is not associated with iron deposits in the eye.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q30: A 45-year-old male patient presents to the ophthalmology clinic with a history of eye trauma with vegetative material .He complains of severe eye pain, redness, tearing, and photophobia. Microscopic examination of the corneal scraping reveals septate hyphae at an acute angle. What is the most likely causative agent responsible for this condition? (Q820465)

A. Fungal corneal ulcer

B. Bacterial corneal ulcer

C. Viral corneal ulcer

D. Acanthamoeba corneal ulcer

Correct Answer: A. Fungal corneal ulcer

Explanation: Correct Option A - Fungal Corneal Ulcer Fungal corneal ulcers are typically caused by filamentous fungi, and they present with symptoms such as severe eye pain, redness, tearing, and photophobia. Microscopic examination of corneal scrapings often reveals septate hyphae at an acute angle. This presentation is consistent with the information provided in the question. Incorrect Options: Option B - Bacterial Corneal Ulcer: Bacterial

corneal ulcers can also cause similar symptoms, including eye pain, redness, and tearing. However, bacterial ulcers are usually associated with a purulent discharge and the appearance of neutrophils on microscopic examination of corneal scrapings. Option C - Viral Corneal Ulcer: Viral corneal ulcers, such as those caused by herpes simplex virus (HSV), typically present with a dendritic ulcer pattern on the cornea. They may cause symptoms like eye pain and photophobia, but they usually lack the septate hyphae seen in fungal infections. Option D - Acanthamoeba Corneal Ulcer: Acanthamoeba corneal ulcers are caused by a protozoan parasite. They often occur in contact lens wearers or individuals exposed to contaminated water. These ulcers can cause eye pain, redness, and photophobia, but they are not characterized by septate hyphae

↑ *Back to this question in Part A*

@Neet_pg_bot


Q31: Which of the following combinations corresponds to the image given below? (Q854714)

A. Cobalt blue- Tryptan blue - Corneal ulcer

B. Cobalt green- Rose bengal stain- Corneal ulcer


C. Cobalt blue- Fluorescein stain- Corneal ulcer


D. Cobalt blue- Lissamine green- Corneal ulcer




Correct Answer: C. Cobalt blue- Fluorescein stain- Corneal ulcer

Explanation: The combination of cobalt blue light and fluorescein stain is the gold standard for evaluating corneal ulcers , making Option C the correct answer. This setup allows for optimal visualization of corneal defects and provides clear diagnostic information. Fluorescein sodium {{caption_text}} Fluorescein penetrates poorly into the lipid layer of the corneal epithelium, and therefore, it does not stain normal cornea. Although fluorescein is a very effective stain for the diseased cornea, it is more difficult to detect fluorescein staining of the conjunctiva because of the poor scleral contrast. Fluorescein absorbs light in the blue spectrum (490 nm) and emits yellow-green light of a higher wavelength (530 nm), therefore staining can be more readily viewed if a cobalt blue or yellow (blue-free) filter is used. This combination is particularly useful for detecting corneal defects and ulcers. Cobalt blue- Tryptan blue - Corneal ulcer (Option A): Cobalt blue is a filter that is commonly used in conjunction with fluorescein stain, as it enhances the fluorescence of fluorescein, making it easier to detect corneal defects. It absorbs light in the blue spectrum (around 490 nm) and helps the fluorescein stain emit a bright yellow-green fluorescence when illuminated. Tryptan blue is not the preferred or standard stain for detecting corneal ulcers. While it can stain the corneal tissue, it does not provide the clear fluorescence that fluorescein does when combined with a cobalt blue filter. While cobalt blue is useful in conjunction with certain dyes, tryptan blue is not an ideal choice for assessing corneal ulcers. Cobalt green- Rose bengal stain- Corneal ulcer (Option B): Rose bengal {{caption_text}} Rose bengal is a derivative of fluorescein and is used to detect ocular surface damage. It was initially theorized that rose bengal stains dead and devitalized tissue, but it is now thought to stain any part of the ocular surface which is not protected by an overlying tear film, specifically in areas lacking membrane-associated mucins. Cobalt blue- Lissamine green- Corneal ulcer (Option D): Lissamine green {{caption_text}} It has a high affinity for staining dead and degenerated cells, including mucous strands. The dye appears to stain areas where there are disrupted intercellular junctions. Use of a green (red-free) filter shows the stained areas to appear black in color. As it is better tolerated than rose bengal, it is the preferred dye for staining the bulbar conjunctiva.







References:

Ocular surface staining: Current concepts and techniques - PMC (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q32: The eye examination of a 20-year-old male shows a golden ring on the cornea. What of the following is the investigation of choice to diagnose the condition? (Q855482)

A. Alpha-1-antitrypsin

B. Alpha-fetoprotein

C. Serum iron

D. Serum ceruloplasmin

Correct Answer: D. Serum ceruloplasmin

Explanation: The golden ring on the cornea seen in this 20-year-old male is characteristic of Kayser-Fleischer rings , which are typically associated with Wilson's disease . Wilson's disease is a genetic disorder of copper metabolism that leads to copper accumulation in various tissues, including the liver, brain, and cornea. Kayser-Fleischer rings are copper deposits that form in the Descemet membrane of the cornea , appearing as a golden or greenish ring on slit-lamp examination. Wilson's disease Deficiency of ceruloplasmin, leading to abnormal copper deposition in tissues. Symptoms : Liver disease Basal ganglia dysfunction (movement disorders) Psychiatric disturbances Ocular Signs : Kayser-Fleischer ring : Brownish-yellow golden copper deposits in the corneal Descemet membrane, visible on gonioscopy. Sunflower cataract : Anterior capsular cataract in some patients. Diagnosis : Kayser-Fleischer rings detected on gonioscopy. Serum Ceruloplasmin levels Treatment : Penicillamine therapy can reduce copper deposits and improve symptoms. Incorrect Options: Alpha-1-antitrypsin (Option A) : This is a protein associated with lung and liver diseases (like emphysema and cirrhosis), but it is not related to copper metabolism or Wilson's disease. Alpha-fetoprotein (Option B) : This is a tumor marker, often elevated in certain cancers or during pregnancy, but not associated with Wilson's disease or corneal findings. Serum iron (Option C) : Elevated iron levels are associated with hemochromatosis or iron overload disorders, not with Wilson's disease.

References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 267

Wilson Disease - StatPearls - NCBI Bookshelf

↑ *Back to this question in Part A*

@Neet_pg_bot


Q33: What dye and filter are used in the image, as shown below? (Q855485)

A. Lissamine dye green filter

B. Lissamine dye cobalt blue filter


C. Fluorescein dye green filter


D. Fluorescein dye cobalt blue filter



Correct Answer: D. Fluorescein dye cobalt blue filter

Explanation: Fluorescein sodium {{caption_text}} Fluorescein penetrates poorly into the lipid layer of the corneal epithelium, and therefore, it does not stain normal cornea. Although fluorescein is a very effective stain for the diseased cornea, it is more difficult to detect fluorescein staining of the conjunctiva because of the poor scleral contrast. Fluorescein absorbs light in the blue spectrum (490 nm) and emits yellow-green light of a higher wavelength (530 nm), therefore staining can be more readily viewed if a cobalt blue or yellow (blue-free) filter is used. This combination is particularly useful for detecting corneal defects and ulcers. Cobalt blue filter: Cobalt blue is a filter that is commonly used in conjunction with fluorescein stain, as it enhances the fluorescence of fluorescein, making it easier to detect corneal defects. It absorbs light in the blue spectrum (around 490 nm) and helps the fluorescein stain emit a bright yellow-green fluorescence when illuminated. Lissamine green dye {{caption_text}} It has a high affinity for staining dead and degenerated cells, including mucous strands. The dye appears to stain areas where there are disrupted intercellular junctions. Use of a green (red-free) filter shows the stained areas to appear black in color. As it is better tolerated than rose bengal, it is the preferred dye for staining the bulbar conjunctiva. Incorrect Options: Lissamine green dye green filter (Option A): Lissamine green stains dead or degenerated cells and mucous strands on the conjunctiva, not corneal defects. A green filter is used with lissamine green because it enhances contrast, making stained areas appear black. It’s not effective for fluorescein. Lissamine green dye cobalt blue filter (Option B): Lissamine green doesn’t fluoresce, so a cobalt blue filter (which works with fluorescein) won’t enhance its staining. Cobalt blue is useful for fluorescein fluorescence, not for lissamine green, which requires a green filter. Fluorescein dye green filter (Option C): Fluorescein works best with a cobalt blue filter for fluorescence, not a green filter. A green filter doesn’t enhance fluorescein’s fluorescence, making it less effective for detecting corneal defects.





References:

Ocular surface staining: Current concepts and techniques - PMC (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q34: A child with elevated liver enzymes presented to the OPD with the following eye finding on gonioscopy. It is associated with? (Q858142)

A. Copper

B. Zinc

C. Selenium

D. Iron



Correct Answer: A. Copper

Explanation: ye Finding : The child has a Kayser-Fleischer ring , a brownish-yellow copper deposit in the Descemet membrane of the cornea, which is visible on gonioscopy. This is a hallmark sign of Wilson disease. Wilson's disease Deficiency of caeruloplasmin, leading to abnormal copper deposition in tissues. Symptoms : Liver disease Basal ganglia dysfunction (movement disorders) Psychiatric disturbances Ocular Signs : Kayser-Fleischer ring : Brownish-yellow copper deposits in the corneal Descemet membrane, visible on gonioscopy. Sunflower cataract : Anterior capsular cataract in some patients. Diagnosis : Kayser-Fleischer rings detected on gonioscopy. Treatment : Penicillamine therapy can reduce copper deposits and improve symptoms. Zinc (Option B), selenium (Option C), and iron (Option D) are not associated with the Kayser-Fleischer ring or the specific symptoms of Wilson disease.

References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 267

Wilson Disease - StatPearls - NCBI Bookshelf

↑ *Back to this question in Part A*

@Neet_pg_bot


Q35: A patient presents to you with pain and redness in the right eye following an injury during welding. On examination, the image as seen below is observed. Which of the following is the appropriate next step in managing this patient? (Q858146)

A. Steroids and antibiotics

B. Surgical exploration

C. Keratoplasty

D. Removal by 26G needle



Correct Answer: D. Removal by 26G needle

Explanation: : While these medications can be useful for treating inflammation and preventing infection, they are not the first-line treatment for immediate foreign body removal. Surgical exploration (Option B): This option may be considered for more complex injuries or if there are complications, but it is not the first step in straightforward foreign body cases. Keratoplasty (Option C): This surgical procedure is reserved for severe corneal damage or scarring and is not appropriate immediately after an injury

References:

Eyes - flash burns - Better Health Channel

Corneal Flash Burns - Welder's Eye - Photokeratosis

Eye Emergency Manual: An illustrated guide - Third edition, December 2023

↑ *Back to this question in Part A*

@Neet_pg_bot


Q36: Please analyze the image provided and determine the specific abnormality or injury present . (Q793032)

A. Viral keratitis

B. Fungal keratitis

C. Bacterial keratitis

D. Interstitial keratitis



Correct Answer: A. Viral keratitis

Explanation: Correct Option A - Viral keratitis : The lesion in viral keratitis is typically characterized by dendritic or branching patterns on the cornea, as seen in the image. Viral keratitis is a type of corneal infection caused by a virus, such as herpes simplex virus (HSV) or varicella-zoster virus (VZV). Incorrect Options: Option B - Fungal keratitis: The lesion in fungal keratitis often presents as an infiltrate or ulcer with feathery edges (hyphae) and satellite lesions. Option C - Bacterial keratitis: The lesion in bacterial keratitis often presents with ulcer, stromal edema, inflammation, and keratic precipitates due to anterior uveitis. Option D - Interstitial keratitis: This is a condition in which there is inflammation of the cornea that affects the middle layer (stroma) of the cornea. The lesion in interstitial keratitis is typically characterized by a hazy appearance of the cornea, with no obvious ulcer or infiltrate.

References:

Prepladder Notes

↑ *Back to this question in Part A*

@Neet_pg_bot

Q37: What is the probable diagnosis in a 27-year-old male with eye trauma from vegetative matter presenting with a corneal ulcer, hypopyon, satellite lesions, mild pain, photophobia, and redness? (Q793041)

A. Fungal keratitis

B. Acanthamoeba keratitis

C. Viral keratitis

D. Bacterial keratitis

Correct Answer: A. Fungal keratitis

Explanation: Fungal keratitis : Explantion: History of trauma with vegetative matter is the single most important clue → strongly points toward fungal keratitis Key clinical features of fungal keratitis: Corneal ulcer with dry, feathery, irregular margins Satellite lesions around the main ulcer (classic finding) Hypopyon (often thick and immobile) Mild to moderate pain (pain is less than expected for severity) Photophobia and redness Acanthamoeba keratitis (Option B): Typically associated with contact lens use , especially with tap water Causes severe pain out of proportion to signs Classically shows a ring-shaped stromal infiltrate Satellite lesions and hypopyon are uncommon Viral keratitis (Option C): Usually due to HSV Characteristic dendritic or geographic ulcers Hypopyon and satellite lesions are not typical Bacterial keratitis (Option D): Often has severe pain , purulent discharge Ulcer margins are more well-defined Satellite lesions strongly favor fungal rather than bacterial etiology

References:

Prepladder Notes

↑ *Back to this question in Part A*


Q38: What is the likely reason for a 21-year-old student to have ocular redness, feeling of a foreign object in the eye, and excessive tearing after wearing a soft contact lens, as indicated by the corneal scraping interface contrast microscopy image? (Q793034)

A. Acanthamoeba

B. Naegleria fowleri

C. Entamoeba histolytica

D. Balamuthia mandrillaris



Correct Answer: A. Acanthamoeba

Explanation: Correct Option A - Acanthamoeba: The clinical presentation of ocular redness, foreign body sensation, epiphora, history of using a soft contact lens, and image showing a cyst of acanthamoeba characterized by outer wrinkled wall is suggestive of acanthamoeba keratitis caused by acanthamoeba a free living protozoan. It is prevalent in individuals using soft contact lens and using tap water for cleaning or swimming with the contact lens in a swimming pool. Incorrect Options: Option B - Naegleria fowleri: Naegleria fowleri is a free-living amoeba that can cause a rare and often fatal brain infection known as primary amoebic meningoencephalitis

(PAM). Option C - Entamoeba histolytica: Entamoeba histolytica is a parasitic amoeba that can cause amoebiasis, a gastrointestinal infection. Option D - Balamuthia mandrillaris: Balamuthia mandrillaris is a free-living amoeba that can cause a rare and often fatal brain infection known as granulomatous amoebic encephalitis (GAE).

References:

↑ *Back to this question in Part A*

@Neet_pg_bot

• [09_Conjunctiva] 019 Conjunctiva - I

« Back to Folder Directory

[09_Conjunctiva] 019 Conjunctiva - I — Quick Jump

Q1 (p. 1193)	*Q2 (p. 1196)*	*Q3 (p. 1198)*	*Q4 (p. 1199)*	*Q5 (p. 1202)*	*Q6 (p. 1203)*
Q7 (p. 1205)	*Q8 (p. 1208)*	*Q9 (p. 1210)*	*Q10 (p. 1211)*	*Q11 (p. 1212)*	*Q12 (p. 1217)*
Q13 (p. 1217)	*Q14 (p. 1218)*	*Q15 (p. 1219)*	*Q16 (p. 1221)*	*Q17 (p. 1221)*	*Q18 (p. 1225)*
Q19 (p. 1230)	*Q20 (p. 1232)*	*Q21 (p. 1234)*	*Q22 (p. 1236)*	*Q23 (p. 1238)*	*Q24 (p. 1239)*
Q25 (p. 1240)	*Q26 (p. 1241)*				

Q1: A 6-year-old boy presents with a 2-week history of worsening bilateral eye redness, itching, and frequent lacrimation. On examination, hypertrophied, cobblestone-like papillae is seen on the palpebral conjunctiva. What is the most likely diagnosis? (Q853727)

A. Atopic Keratoconjunctivitis

B. Vernal Keratoconjunctivitis

C. Bacterial Conjunctivitis

D. Chlamydial Conjunctivitis





Correct Answer: B. Vernal Keratoconjunctivitis

■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: Vernal Keratoconjunctivitis is the most likely diagnosis given the patient’s age, the characteristic findings of cobblestone-like papillae on the palpebral conjunctiva It commonly affects young children and adolescents, presenting with symptoms such as itching, burning, and lacrimation, and often worsens during warm weather. Vernal keratoconjunctivitis or Spring catarrh Symptoms Burning, Itching, Photophobia (sensitivity to light), and lacrimation (excessive tearing). Signs Palpebral Form : Hypertrophied palpebral conjunctiva with a cobblestone appearance;dense fibrous tissue with thickened epithelium. Limbal or Bulbar Form : Opacification at the limbus, gelatinous thickening, and Horner–Trantas dots Pseudogerontoxon/cupid’s bow (which resembles arcus senilis) is a grayish-white ring at the limbus seen in severe limbal vernal keratoconjunctivitis (VKC). Shield ulcers are a significant complication of VKC, particularly in severe cases. These microerosions typically form on the upper third of the cornea. Mixed form: Exhibits characteristics of both the tarsal (palpebral) and limbal subtypes, typically presenting in only one eye. Maxwell Lyon's sign is a pseudomembrane formed due to excessive mucus deposition on the papilla resulting in cobblestone papillae. Atopic keratoconjunctivitis (Option A): Seen in older patients (teens/adults) with atopy More chronic, less dramatic cobblestone papillae Bacterial conjunctivitis (Option C): Purulent discharge , not intense itching No cobblestone papillae Chlamydial conjunctivitis (Option D): Follicles , not papillae Usually chronic with mucopurulent discharge


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 181, 182, 183, 184, 185, 186.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9376221/

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4108143/

https://www.nature.com/articles/eye198963.pdf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 25-year-old male presents to the ophthalmologic clinic for evaluation of his ocular lesions. After carefully assessing the eyes, the attending doctor diagnoses vernal keratoconjunctivitis. Which of the following statements is most likely associated with vernal keratoconjunctivitis? (Q882187)

A. It is a Follicular hypertrophy

B. Hallmark is the presence of Cobblestone papillae


C. Mild itching is present


D. Type IV hypersensitivity response

Correct Answer: B. Hallmark is the presence of Cobblestone papillae

Explanation: Correct Option B - Hallmark is the presence of Cobblestone papillae: Vernal Keratoconjunctivitis Bilateral, Recurrent, Self-limiting conjunctivitis. Young boys in 1st decade of life are affected. More common in the summer. More common in the tropical climate. Cobblestone papillae are seen: hallmark Horner trantas spots Shield ulcer Cobblestone papillae Incorrect Options: Option A - It is a Follicular hypertrophy: The hypertrophy which occurs in the VKC is papillary, not follicular. Option C - Mild itching is present: The VKC is characterized by severe itching, not mild. Option D - Type IV hypersensitivity response: The type of hypersensitivity reaction is Type I, which is IgE mediated.







References:

Comprehensive Ophthalmology 6th edition A K Khurana Pg 79-80

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 3-year-old child weighing 14 kg is brought to the pediatric clinic with complaints of night blindness and dryness of the eyes. On examination, the ophthalmologist observes conjunctival xerosis, Bitot's spots, and keratomalacia. What is the appropriate treatment for this child's condition? (Q882122)

A. Vitamin A: 2 lahks IU in oral, 0 th , 1 st , and 10 th day

B. Vitamin A: 2 lakhs IU injectable, 0 th , 1 st , and 14 th day

C. Vitamin A: 2 lakhs IU in oral, 0 th , 1 st , and 14 th day

D. Vitamin A: 2 lakhs IU injectable, 0 th , 1 st , and 10 th day

Correct Answer: C. Vitamin A: 2 lakhs IU in oral, 0 th , 1 st , and 14 th day

Explanation: The clinical picture of night blindness, conjunctival xerosis, bitot's spot, and keratomalacia suggest the diagnosis of Xerophthalmia and the treatment is Vitamin A supplementation either orally or injectable. The rule is to inject 1 lakh IU of Vitamin A on the 0 th ,1 st , and 14 th day if given IM for a child above 1 year, or if given orally double the amount of Vitamin A given injectable. If, the child is less than 1 year half the dose. Vitamin A: 2 lakh IU in oral, 0 th , 1 st , and 10 th day (Option A): This option is incorrect, the last dose is given on the 14 th day, not on the 10 th day. Vitamin A: 2 lakhs IU injectable, 0 th , 1 st , and 14 th day (Option B): This option is incorrect, the dose for injectable Vitamin A above 1 year is 1Lakh, not 2 lakhs. Vitamin A: 2 lakhs IU injectable, 0 th , 1 st , and 10 th day (Option D): This option is incorrect, the last dose is given on the 14 th day, not on the 10 th day and moreover the dose given is 1 Lakh if given injectable.

References:

Kanski's Clinical Ophthalmology 9th edition, Page 247, 248

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 64-year-old patient with dry eyes undergoes several tests. Which finding best supports the diagnosis of dry eye syndrome? (Q853738)

A. Tear film break-up time of 15 seconds

B. Schirmer test showing 12 mm wetting

C. Extensive Rose Bengal staining

D. Normal tear film osmolality


Correct Answer: C. Extensive Rose Bengal staining

■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: Extensive Rose Bengal staining indicates significant damage to the corneal and conjunctival epithelium, which is a feature of dry eye. Schirmer’s Test: Schirmer’s test is used to diagnose aqueous deficiency dry eye disease. Schirmer's Test 1 Measures both basal and reflex tear secretions Uses Whatman-41 filter paper, kept in the lower fornix at the junction of lateral one-third and medial two-thirds. After 5 minutes, wetting of the filter strip from the bent end is measured. Normal value: >15 mm 5-10 mm: Mild to moderate keratoconjunctivitis sicca (KCS). < 5 mm: Severe KCS (Option B) Schirmer's Test 2 Measures only basal secretion under anesthesia Tear Film Break-Up Time (TBUT) Tear film stained with sodium fluorescein 1% is observed under a slit lamp, and time is noted after the patient blinks. The interval between the complete blink and the first appearance of a ‘dark’ dry spot on the cornea is recorded as TBUT. Normal: 15-35 seconds < 10 seconds indicates unstable tear film. (Option A) Rose Bengal or Lissamine Green Stain devitalized, desiccated corneal and conjunctival epithelium red or green. (Option C) Rose Bengal stain Lissamine Green Tear film Osmolality Increased in dry eye (Option D)




@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology, Edition 9, Page- 160, 161, 162.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3964175/
https://www.sciencedirect.com/topics/medicine-and-dentistry/schirmer-test
https://www.sciencedirect.com/topics/nursing-and-health-professions/schirmer-test
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3657952/#

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 60-year-old patient diagnosed with dry eye syndrome asks about the typical features of the condition. Which of the following is not commonly associated with dry eye? (Q853737)

A. Frothy tears along the lid margin

B. Punctate epithelial erosions on the cornea

C. Conjunctival xerosis and keratinization

D. Follicles on the conjunctiva

Correct Answer: D. Follicles on the conjunctiva

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Follicles on the conjunctiva are typically seen in viral or chlamydial infections , not in dry eye. Signs Dry eye: Tear Film Signs: Presence of stingy mucus and particulate matter Reduced or absent marginal tear strip Frothy tears along the lid margin indicate meibomian gland dysfunction (Option A) Posterior Blepharitis (Meibomitis): Inflammation of meibomian glands. Conjunctival Signs: Lustreless and mildly congested conjunctiva Conjunctival xerosis and keratinization may occur (Option C) Positive Rose Bengal or Lissamine green staining Corneal Signs: Punctate epithelial erosions (Option B) Filaments and mucus plaques Loss of corneal luster Lesions delineated by fluorescein, Rose Bengal, or Lissamine green staining



References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 157

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 65-year-old woman with rheumatoid arthritis presents with dry eyes and mouth. Rose bengal staining shows conjunctival damage, and tear analysis reveals a

tear-to-lysozyme ratio of 0.1. Which tear film layer is primarily deficient in this condition? (Q853736)

A. Mucinous layer

B. Aqueous layer

C. Lipid layer


D. All layers are equally affected

Correct Answer: B. Aqueous layer

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Keratoconjunctivitis sicca (Sjögren syndrome) is the most likely diagnosis due patient's history of RA and the tear-to-lysozyme ratio of 0.1, dry eyes and mouth, and conjunctival damage shown by Rose Bengal staining, which indicates a deficiency in the aqueous layer of the tear film. Tear Film Layer Secreted By Function Defective in Innermost Mucinous Layer (Option A) Goblet cells, crypts of Henle, Manz's glands Convert hydrophobic corneal surface into hydrophilic one Vitamin A deficiency Middle Aqueous Layer (thickest layer) (Option B) Lacrimal and accessory conjunctival glands (Krause, Wolfring/Ciaccio) Contain tears Also contains antibacterial substances like lysozyme, beta-lysin, and lactoferrin. Sjogren's syndrome Outermost Lipid Layer (thinnest layer) (Option C) Meibomian glands Prevent overflow of tears. Retard evaporation. Lubricates the eyelids. Contact lens wear Meibomian gland defect



References:

Kanski’s Clinical Ophthalmology, 9th edition, Page- 158, 159.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q7: A 70-year-old man with extensive sun exposure has a yellowish, triangular patch on the nasal conjunctiva, as shown below. The lesion does not extend onto the cornea. What is the most likely diagnosis? (Q853735)

A. Pterygium

B. Pinguecula

C. Dermoid

D. Malignant melanoma




Correct Answer: B. Pinguecula


■ *Listen (English)*


■ *Listen (Hindi)*


Explanation: A yellowish, triangular patch on the conjunctiva near the limbus, typically in the elderly, associated with sun exposure and not extending onto the cornea, points towards the diagnosis of Pinguecula. Pinguecula Definition A triangular patch on the conjunctiva appears yellow and resembles fat (pinguis). Pathology Caused by

hyaline infiltration and elastotic degeneration of submucous tissue. Etiology Common in elderly people, particularly those exposed to strong sunlight, dust, and wind. Clinical Feature Bilateral Usually stationary Yellowish triangular patch near the limbus Apex of the triangle is away from the cornea Affects the nasal side first, followed by the temporal side Image Complication Inflammation of pinguecula Intraepithelial abscess formation Rarely calcification Treatment Generally, it requires no treatment. When inflamed, treated with topical steroids. May be excised when cosmetically unaccepted. Pterygium (Option A) is a triangular, vascularized growth extending onto the cornea. Dermoid (Option C) appears as a lenticular yellow tumor near the corneal margin and consists of epidermoid epithelium with sebaceous glands and hair. Malignant melanoma (Option D) typically occurs at the limbus, is usually pigmented, and can spread or metastasize.









References:

Kanski's Clinical Ophthalmology, 9th edition, Page- 197, 198.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q8: A 50-year-old male comes to the clinic with complaints of a small fleshy triangular mass in the eye. There is no history of pain or discharge. On examination, it is as in the image. What is true about this lesion? (Q853733)

A. This is a non-progressive lesion

B. It can be present in any quadrant of the eye

C. It is caused by thermochemical burn, trauma, and chronic inflammation

D. Histology shows collagenous degenerative changes in the vascularized subepithelial stroma




Correct Answer: D. Histology shows collagenous degenerative changes in the vascularized subepithelial stroma


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above clinical scenario points to the diagnosis of pterygium. Pterygium/Surfer's eye Pathology Histology shows collagenous degenerative changes in the vascularized subepithelial stroma. Symptoms Frequently follows a pinguecula Often bilateral Usually on the nasal side (Option B) Impairment of vision due to: Progression into the pupillary area of the cornea Induced astigmatism by corneal flattening in the axis of pterygium Progressive

pterygium: thick and vascular (Option A) Regressive pterygium: thin, pale, and atrophic Ceases to grow but never disappears A probe can be passed under the lesion Can present as inflammed pterygium Pseudopterygium (Option C) is caused by: Trauma Chronic conjunctivitis Thermochemical burn





References:

Kanski’s Clinical Ophthalmology, 9th edition, Page- 198, 199.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: A 25-year-old male presents with difficulty seeing at night. Which of the following is not a cause of nyctalopia (night blindness)? (Q853732)

A. Gyrate atrophy

B. Pathological myopia

C. Cone dystrophy


D. Choroideremia

Correct Answer: C. Cone dystrophy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Cone dystrophy is primarily associated with day blindness (hemeralopia) , which is the inability to see clearly in bright light due to poor light adaptation. Nyctalopia/ Night Blindness is the inability to see in low-light conditions. It is attributed to interference with the functions of the retinal rods. Causes of Nyctalopia: Retinitis pigmentosa Xerophthalmia Pathological myopia (Option B) Advanced stage of primary open-angle glaucoma Peripheral cortical cataract Diffuse choroidal dystrophy Choroideremia (Option D) Gyrate atrophy (Option A): Due to a deficiency of the enzyme ornithine aminotransferase, plasma ornithine levels are elevated. Management: Restriction of arginine (precursor amino acid for ornithine) in diet. Vitamin B6 supplementation lowers plasma ornithine levels.



References:

https://www.sciencedirect.com/topics/neuroscience/choroidal-dystrophy
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7347087/ https://www.sciencedirect.com/topics/pharmacology-toxicology-and-pharmaceutical-science/retina-gyrate-atrophy https://www.ncbi.nlm.nih.gov/books/NBK557759/
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9740538/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: A 3-year-old child presents with symptoms of night blindness and dry eyes. On examination, there are Bitot's spots on the temporal side of both eyes. The child's guardian reports that the child has had a history of frequent respiratory infections. What is the most appropriate treatment regimen to prevent further ocular complications? (Q853731)

A. 100,000 IU of vitamin A orally on diagnosis, next day, and after 2 weeks.

B. Vitamin A eye drops and 10,000 IU orally for 1 month.

C. 200,000 IU of vitamin A orally on diagnosis, next day, and after 2 weeks.

D. Oral multivitamins with increased vitamin A-rich foods for 3 months.

Correct Answer: C. 200,000 IU of vitamin A orally on diagnosis, next day, and after 2 weeks.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This is the recommended treatment for children aged 1 year and older with xerophthalmia. The WHO advises administering 200,000 IU of vitamin A immediately, then repeating the dose on the following day and again 2 weeks later. Population Vitamin A Dose Treatment regimen Children (≥1 year) 200,000 IU orally on diagnosis Day1, Day 2, and 2 weeks later (Option C) Children (Infants under 12 months) 100,000 IU orally Day1, Day 2, and 2 weeks later (Options A and B)

References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 248.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q11: A 5-year-old boy from a rural area has difficulty seeing at night. Upon examination, you notice the presence of triangular, dry-white spots on the bulbar conjunctiva in both eyes. Which of the following represents the correct diagnosis and grade of xerophthalmia based on the clinical findings? (Q853730)

A. Night Blindness (XN)

B. Conjunctival Xerosis (X1A)

C. Bitot's Spots (X1B)


D. Corneal Xerosis (X2)



Correct Answer: C. Bitot’s Spots (X1B)

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Bitot's spots are characteristic of xerophthalmia and appear as triangular, dry-white lesions on the bulbar conjunctiva, a hallmark of grade X1B xerophthalmia. More common on the temporal side Grade Image XN (Night Blindness) (The earliest sign of xerophthalmia) X1A (Conjunctival Xerosis) X1B (Bitot’s Spots) X2 (Corneal Xerosis) X3A (Corneal ulceration/ Keratomalacia involving < 1/3rd of cornea) X3B (Corneal ulceration/ Keratomalacia involving > 1/3rd of cornea) XS (Corneal scarring) XF (Xerophthalmic Fundus)





@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot





References:

KANSKI'S Clinical Ophthalmology, 9th Edition, Page 247.

www.nature.com/articles/s41433-021-01604-z

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 23-year-old male presents with a history of atopic dermatitis and asthma. He complains of tearing, burning, itching sensation and blurry vision for the past week. On examination, his eyelids are thickened and swollen, with signs of meibomitis. Slit-lamp examination reveals anterior subcapsular cataracts. What is the most appropriate first-line treatment for this patient's ocular symptoms? (Q853729)

A. Topical corticosteroids

B. Topical dual-acting antihistamine/mast cell stabilizer

C. Topical cyclosporine

D. Oral antihistamines

Correct Answer: B. Topical dual-acting antihistamine/mast cell stabilizer

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Topical dual-acting antihistamine/mast cell stabilizer is the first-line treatment for atopic keratoconjunctivitis. It addresses both histamine release and mast cell stabilization, helping control inflammation and allergic responses. Atopic Keratoconjunctivitis Treatment First-line therapy: Topical dual-acting antihistamine/mast cell stabilizer If there is no improvement in 2-3 weeks: A brief course of topical corticosteroids. (Option A) For moderate-to-severe disease requiring frequent corticosteroids - Use topical cyclosporine (2%, compounded). (Option C) Eyelid Dermatitis: Topical tacrolimus is preferred over corticosteroids. Oral antihistamines (Option D) may help with systemic symptoms, but they are not as effective for treating ocular manifestations directly.

References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 184, 185, 186.

www.uptodate.com/contents/atopic-keratoconjunctivitis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 10-year-old child presents with itching, watering, and redness in both eyes. On examination, small round yellow nodules are observed near the limbus, and

conjunctival congestion is noted. The child has a past h/o TB. Which of the following best explains the pathophysiology of this condition? (Q853728)

A. Type I hypersensitivity reaction caused by environmental allergens

B. Type IV hypersensitivity reaction to endogenous bacterial antigens

C. Type II hypersensitivity reaction triggered by viral infections

D. Autoimmune reaction against corneal epithelial cells

Correct Answer: B. Type IV hypersensitivity reaction to endogenous bacterial antigens

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above presentation is a case of phlyctenular keratoconjunctivitis primarily mediated by a delayed-type (Type IV) hypersensitivity reaction triggered by bacterial proteins of Staphylococcus aureus or tuberculosis. The child's history of TB is significant, as TB antigens can trigger this hypersensitivity response, leading to the characteristic ocular findings. The immune system reacts to these antigens, causing the formation of nodules (phlyctens) and subsequent inflammation around the limbus. This explains the localized inflammatory response seen in this condition.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: Which of the following statements is true regarding the epithelium of the conjunctiva? (Q853712)

A. The epithelium of the limbal conjunctiva is thinner than that of the fornices.

B. The conjunctival epithelium is lined by stratified non-keratinized epithelium.

C. The adenoid layer of the conjunctiva is present at birth and plays a role in neonatal immunity.


D. The dense fibrous layer is superficial to the adenoid layer and contains goblet cells.

Correct Answer: B. The conjunctival epithelium is lined by stratified non-keratinized epithelium.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The conjunctival epithelium is composed of stratified non-keratinized epithelium . Layers of Conjunctiva Outer Epithelial Layer Lined by stratified squamous epithelium. The epithelium gradually thickens from the fornices to the limbus (Option A) , becoming stratified and nonkeratinized near the corneal margin , while the palpebral conjunctiva lining the eyelids remains thinner and nonkeratinized . It contains: Goblet cells Melanocytes Langerhans cells Adenoid Layer (or lymphoid layer) It is located below the epithelium. This layer develops lymphoid tissue only after 3-4 months of age, which explains why newborns cannot produce a follicular conjunctival reaction. (Option C) Dense Fibrous Layer It is present deep in the adenoid layer, merging into the underlying tissues (lid or sclera). This layer contains the vessels and nerves of the conjunctiva. (Option D)



References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 168.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 2-day-old infant is evaluated for discharge from the eyes. The discharge was initially serous, but it quickly turned into mucopurulent and then purulent. On examination, the eyes are bright red and swollen. What is the most likely causative agent here? (Q853726)

A. Chlamydia trachomatis

B. Silver nitrate

C. Treponema pallidum


D. Neisseria gonorrhoeae

Correct Answer: D. Neisseria gonorrhoeae

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical scenario points to the diagnosis of ophthalmia neonatorum caused by Neisseria gonorrhea and it presents as an inflamed, red, and swollen conjunctiva with discharge that changes fast. Ophthalmia neonatorum or neonatal conjunctivitis is a mucoid, mucopurulent, or purulent discharge from one or both eyes in the first month of life. Source of infection: Neisseria gonorrhoeae: Transmitted at the time of delivery Presents between 2-5 days of birth The discharge becomes mucopurulent and purulent quickly Inflamed, bright red, swollen conjunctiva Thick yellow pus and marked chemosis Cornea seen at the bottom of a crater-like pit in severe cases Bulbar conjunctiva & lids: dense infiltration, swollen and tense Free flow of pus, serum, and often blood later Chlamydia trachomatis: (Option A) {{caption_text}} Other causes: Staphylococci Streptococcus pneumoniae Gram-negative coliforms Herpes simplex virus Chemical: Silver nitrate solution previously used for prophylaxis immediately after birth (Option B)



References:

KANSKI’S Clinical Ophthalmology, 9th edition, Page- 176, 177.
https://www.ncbi.nlm.nih.gov/books/NBK551572/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A patient with characteristic features of trachoma is being evaluated. Which of the following would be the most appropriate systemic antibiotic to be used for the treatment of trachoma? (Q853725)

A. Erythromycin

B. Doxycycline

C. Amoxicillin

D. Azithromycin

Correct Answer: D. Azithromycin

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Azithromycin stands out as the preferred systemic antibiotic or treatment of choice, given its effectiveness and convenient dosing regimen, particularly when taken orally either as a single dose or as part of a mass drug administration approach. Management SAFE is a WHO-recommended combination of interventions to treat trachoma and prevent it from leading to blindness S urgery for cicatricial entropion and trichiasis (Tertiary prevention) A ntibiotics to eliminate the organism (Secondary prevention) F acial cleanliness to avoid infection (Primary prevention) E nvironmental improvement with better hygiene and sanitation (Primordial prevention) Erythromycin (Option A) or doxycycline (Option B) may be considered, although caution is warranted with tetracyclines in pregnancy, breastfeeding, and among children, and hence, it is not preferred over azithromycin.

References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 176.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6760986/ https://www.ncbi.nlm.nih.gov/books/NBK559035/ https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2396018/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A patient presents with irritation of the eyes and foreign body sensation in the eyes. He is examined by the ophthalmologist, who observes the following in his upper eyelid. According to the WHO classification, which grade is the disease here? (Q853724)

A. TF

B. TI

C. CO

D. TS




Correct Answer: D. TS

■ *Listen (English)*


■ *Listen (Hindi)*


Explanation: The above scenario is of trachoma wherein the picture represents a linear scar in the sulcus subtarsalis called Arlt's line , seen in trachomatous scarring stage. WHO Grading System of Trachoma: WHO Classification of Trachoma (FISTO) Stage Description Image TF : Trachomatous Inflammation- Follicular (Option A) The presence of five or more follicles in the upper tarsal conjunctiva Active stage of trachoma TI : Trachamatous Inflammation- Intense (Option B) Pronounced inflammatory thickening of the tarsal conjunctiva that obscures more than half of the normal deep tarsal vessels. TS: Trachamatous Scarring Presence of scarring in tarsal conjunctiva White, bands or sheets of fibrosis in tarsal conjunctiva TT: Trachamatous Trichiasis At least one eyelash rub on eyeball or evidence of recent removal of inturned eyelashes. CO: Corneal Opacity (Option C) Easily visible corneal opacity over the pupil Part of the pupil margin is blurred when seen through the opacity as it is very dense corneal scarring


@Neet_pg_bot


@Neet_pg_bot







References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 174

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: An 8-year-old female is being evaluated for conjunctivitis. She gives a history of itching and mild foreign body sensation in the eyes, stickiness of the lids, and mucoid discharge from the eyes. On examination, there is conjunctival congestion. Which of the following signs would be seen in this stage of the disease? (Q853723)

A. Pannus siccus

B. Arlt’s line

C. Herbert’s follicle

D. Tylosis


Correct Answer: C. Herbert’s follicle


■ *Listen (English)*


■ *Listen (Hindi)*

Explanation: The above clinical scenario points to an active trachomatous infection, and Herbert’s follicles are present in the limbal area of the eye in the active stage of the disease. Signs of active trachoma Signs of cicatricial trachoma Conjunctival signs: Conjunctival congestion Follicles in the upper tarsal conjunctiva: Look like sago-grains Papillary enlargement: red, velvety Conjunctival signs: Scarring It may be irregular, star-shaped, or linear Linear scar in the sulcus subtarsalis is called Arlt’s line (Option B) Accumulation of dead epithelial cells and inspissated mucus in the glands of Henle Pseudocyst Xerosis Symblepharon Corneal signs: Superficial keratitis on slit lamp and fluorescein Herbert’s follicles Progressive pannus/ Trachomatous pannus : The upper part of the cornea is cloudy Trachomatous pannus Progressive pannus Corneal ulcers are common at the advancing edge of the pannus Shallow, little infiltrated Cause lacrimation and photophobia Corneal signs: Regressive pannus or pannus siccus (Option A) : Receding infiltration Vessels extend beyond the area of infiltration and haze Herbert pits : oval or circular pitted scars formed after the healing of Herbert’s follicles in the limbal area Vascularized and opaque cornea. Corneal xerosis Corneal ectasia Total corneal pannus (blinding sequelae) Lid signs: Distortion of the lids Tylosis (Option D) : Thickening of the tarsal plate/lid margin Trichiasis (Option C) due to rubbing of lashes on the cornea Trachomatous ptosis (sleepy appearance) Entropion Madarosis Ankyloblepharon Lacrimal apparatus sequelae: Chronic dacryocystitis Chronic dacryoadenitis





@Neet_pg_bot



@Neet_pg_bot




@Neet_pg_bot



@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot



References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 173, 174, 175.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: A 28-year-old man develops sudden redness, Foreign body sensation and subconjunctival haemorrhage in both eyes after returning from a tropical region. His visual acuity is normal. What is the most likely causative agent? (Q853721)

A. Epstein-Barr virus

B. Coxsackievirus

C. Herpes simplex virus

D. Molluscum Contagiosum


Correct Answer: B. Coxsackievirus

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Sudden redness, foreign body sensation and subconjunctival haemorrhage, especially after travel to a tropical region, are characteristic of acute hemorrhagic conjunctivitis , caused by Coxsackievirus. Causes of Hemorrhagic Conjunctivitis: Viral: Coxsackievirus Enterovirus 70 Echovirus 34 Adenovirus Bacterial: Pneumococcus Haemophilus Neisseria Chlamydia Epstein-Barr virus and Herpes simplex virus (Option A and C): Causes mild conjunctivitis but not acute hemorrhagic conjunctivitis. Herpetic Conjunctivitis - Corneal vesicles merge to form dendritic figures. Molluscum Contagiosum (Option D): Small, white, umbilicated swelling, generally multiple.





Dendritic corneal ulcer due to recurrent keratitis by Herpes Simplex Virus

@Neet_pg_bot



References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 178, 179, 180
https://www.sciencedirect.com/topics/medicine-and-dentistry/acute-hemorrhagic-conjunctivitis

↑ *Back to this question in Part A*

@Neet_pg_bot


Q20: A 30-year-old male presents with irritation and redness at both eye corners, foamy discharge, and skin maceration as shown below. He also has chronic nasal discharge. Which organism is most likely causing his symptoms? (Q853720)

A. Staphylococcus aureus

B. Moraxella axenfeld

C. Neisseria gonorrhoeae

D. Haemophilus influenzae




Correct Answer: B. Moraxella axenfeld

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The presence of localized inflammation, redness, and foamy discharge confined to the angles of the eyes, along with skin maceration around these areas, points towards the diagnosis of angular conjunctivitis. The most common cause is Moraxella axenfeld. Infection is transmitted from the nasal cavity to the eyes via contaminated fingers or handkerchiefs. Treatment: Oxytetracycline (1%) eye ointment applied 2-3 times a day for 9-14 days. Zinc lotion is instilled during the day, and zinc oxide ointment at bedtime to inhibit the proteolytic enzyme and reduce maceration.



References:

AK Khurana Comprehensive Ophthalmology, Edition 7, Pages 66,67

Angular conjunctivitis

Moraxella

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: A 7-year-old child presents with the eye condition shown in the image. The child reports severe eye pain, redness, and eyelid swelling. Examination reveals an inflamed conjunctiva with thick, adherent membranes that bleed when removed. Which of the following organisms is most likely responsible for this presentation? (Q853719)

A. Adenovirus

B. Neisseria gonorrhoeae

C. Corynebacterium diphtheria

D. Beta-hemolytic streptococci



Correct Answer: C. Corynebacterium diphtheria

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Child symptoms like thick, adherent membranes that bleed upon removal, along with the image showing inflamed conjunctiva with adherent membrane, suggest the diagnosis of membranous conjunctivitis. Corynebact erium diphtheria is the most common cause of membranous conjunctivitis. Aspect Membranous Conjunctivitis Pseudomembranous Conjunctivitis Characteristics True membranes grow into the conjunctival epithelium and are more adherent. Removal of these membranes often results in significant bleeding. More superficial, with no growth into conjunctival epithelium, and are less adherent than true membranes. These can be removed with minimal bleeding. Causes Bacterial: Corynebacterium diphtheria. Drug Reactions : Hypersensitivity reactions, Stevens-Johnson syndrome (SJS). Chemical or thermal injury . Infectious : Mild Corynebacterium diphtheria infection, Neisseria gonorrhoeae (Option B ruled out) , Streptococcus pyogenes (Option D ruled out) , Staphylococcus aureus, adenovirus (Option A ruled out) , COVID-19. Inflammatory: Hypersensitivity reactions, foreign body reactions. Ligneous Conjunctivitis Image





References:

https://eyewiki.org/Membranous_Conjunctivitis_and_Pseudomembranous_Conjunctivitis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: A 7-year-old presents with severe eye pain, intense redness, and copious purulent discharge from his right eye. He also has significant lid swelling and tender preauricular lymphadenopathy. Gram stain of the purulent discharge reveals Gram-negative intracellular diplococci. What is the next best step in management? (Q853718)

A. Intravenous ampicillin

B. Eye irrigation with saline

C. Intramuscular Ceftriaxone

D. Topical antibiotic drops


Correct Answer: C. Intramuscular Ceftriaxone

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The presence of copious purulent discharge, tender preauricular lymphadenopathy, and Gram-negative intracellular diplococci strongly suggests gonococcal conjunctivitis, caused by Neisseria gonorrhoeae also termed as acute blennorrhea/ acute purulent conjunctivitis/ hyperacute conjunctivitis, treated with single dose Ceftriaxone IM and oral Azithromycin or doxycycline (twice a day for 7 days). Acute Mucopurulent Conjunctivitis: M/c cause: Staphylococcus aureus Symptoms: Redness Mucus discharge Eyelids stick together, especially in the morning Treatment: Topical Antibiotics: Broad-spectrum antibiotics (e.g., chloramphenicol, lomefloxacin, ofloxacin, ciprofloxacin) applied 4-6 times daily. Antibiotic ointment for lid application to prevent sticking and manage discharge. Acute Purulent Conjunctivitis/ Hyperacute conjunctivitis/ Acute blennorrhoea: M/c cause: Neisseria gonorrhoeae Symptoms: Fiery red conjunctiva (‘pink eye’) Severe pain Lid swelling Copious purulent discharge Pus seen in the fornices and margin of the lid Matting of lashes with dirty yellow crust Coloured halos Tender preauricular lymphadenopathy Treatment: Systemic Antibiotics: Single dose Ceftriaxone IM and oral Azithromycin or doxycycline (twice a day for 7 days) Intravenous ampicillin (Option A) is not the first-line treatment for hyperacute conjunctivitis and is not commonly used. Eye irrigation with saline (Option B) might provide temporary relief but does not address the underlying infection. Systemic antibiotics are needed to prevent complications. Topical antibiotic drops (Option D) are generally insufficient for treating hyperacute conjunctivitis caused by Neisseria gonorrhoeae; systemic antibiotics are required.





References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 171, 172.
https://www.sciencedirect.com/topics/medicine-and-dentistry/gonococcal-conjunctivitis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: A 25-year-old woman with a history of soft contact lens use for the past 8 months presents with bilateral itching, redness, and watery discharge. Examination shows large, polygonal papillae on the upper tarsal conjunctiva. What is the most likely diagnosis? (Q853717)

A. Vernal Keratoconjunctivitis

B. Giant Papillary Conjunctivitis

C. Phlyctenular Conjunctivitis

D. Papillary Conjunctivitis

Correct Answer: B. Giant Papillary Conjunctivitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: A history of prolonged soft contact lens use, along with symptoms like bilateral itching, redness, watery discharge, and the presence of large, polygonal papillae on the upper tarsal conjunctiva, points towards a diagnosis of giant papillary conjunctivitis . Giant Papillary Conjunctivitis Definition Specific allergic inflammation affecting the superior tarsal conjunctiva. Etiology Soft hydrophilic contact lenses Protruding suture ends Ocular prostheses Underlying Mechanism Hypersensitivity reactions (types I and IV) Image Vernal Keratoconjunctivitis (Option A) is characterized by seasonal symptoms, cobblestone-like papillae, limbal thickening with Horner-Trantas dots, and potential corneal involvement, unlike giant papillary conjunctivitis, which typically presents with large papillae associated with contact lens use. Phlyctenular Conjunctivitis (Option C) shows small nodules near the limbus with localized congestion, not large papillae on the tarsal conjunctiva, as seen in giant papillary conjunctivitis. Papillary Conjunctivitis (Option D) is a broad term and lacks the specificity of the large, polygonal papillae and pronounced symptoms seen in Giant Papillary Conjunctivitis, which is linked to contact lens use.



References:

KANSKI'S Clinical Ophthalmology, 9th Edition, Pages 186, 187.

https://www.sciencedirect.com/science/article/pii/S1323893022000831

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: A 28-year-old woman reports bilateral itching, redness, and watery discharge that worsens in spring. Examination revealed mild conjunctival congestion with small papillae on the upper tarsal conjunctiva. What is the most likely diagnosis? (Q853716)

A. Viral Conjunctivitis

B. Allergic Conjunctivitis

C. Bacterial Conjunctivitis

D. Chlamydial Conjunctivitis

Correct Answer: B. Allergic Conjunctivitis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Given the seasonal pattern of symptoms, bilateral itching, watery discharge, mild conjunctival congestion, and the presence of small papillae point toward the diagnosis of allergic conjunctivitis. Viral Conjunctivitis (Option A) typically presents with watery discharge and moderate conjunctival congestion. It is often associated with follicles rather than papillae, and a seasonal pattern is less likely than that of allergic conjunctivitis. Bacterial Conjunctivitis (Option C) is characterized by purulent discharge and marked conjunctival congestion. While bacterial conjunctivitis can sometimes present with papillae, these are not usually as prominent or marked as those in allergic conjunctivitis. Chlamydial Conjunctivitis (Option D) typically presents with mucopurulent discharge and moderate conjunctival congestion. It is often associated with follicles, and papillae may or may not be present. The lack of prominent papillae and specific seasonal patterns make it less likely than allergic conjunctivitis.

References:

Kanski's Clinical Ophthalmology, 9th edition, Pages 170, 181.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: A 27-year-old male presents with redness, mild swelling, and foreign body sensation in the right eye for 2 days. On examination, no visual impairment is noted, and a diagnosis of acute conjunctivitis is made. Which of the following is not a common sign of conjunctivitis? (Q853715)

Correct Answer: Unknown

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Ciliary congestion , also known as ciliary injection, is not a sign of conjunctivitis . It refers to deep, radial vessel congestion around the limbus, usually associated with intraocular conditions like iritis, keratitis, or acute glaucoma. The vessels here are part of the deeper structure of the eye, unlike the superficial vessels involved in conjunctivitis. Symptoms and Signs of Conjunctivitis: Hyperemia or redness of the conjunctiva (Option B) Pain Photophobia (rare, associated corneal involvement or iritis can be present) Blepharospasm Discharge (prominent feature) Foreign body sensation Subconjunctival hemorrhage (Option C) Chemosis or conjunctival edema (Option A) Inflammatory reactions: Follicles (localized aggregation of lymphocytes), papillae (hyperplasia of the normal vascular system), granulomas Pseudomembranes/True membranes



References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 168, 169, 170.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: Which of the following glands are primarily responsible for secreting mucus to maintain ocular surface moisture? (Q853713)

A. Glands of Krause

B. Goblet cells


C. Glands of Wolfring

D. Glands of Moll

Correct Answer: B. Goblet cells

■ *Listen to Audio 1*

Explanation: Goblet cells are located in conjunctival epithelium; secrete mucin , which forms the mucous layer of the tear film, essential for ocular surface moisture and stability. Goblet Cells are unicellular mucous-secreting glands located within the epithelium of the conjunctiva. They produce mucus and maintain the ocular surface's moisture. Glands of Krause (Option A): Accessory lacrimal glands → secrete aqueous component of tears, not mucus. Glands of Wolfring (Option C) : Another type of accessory lacrimal gland → also produce aqueous secretions. Glands of Moll (Option D): Modified sweat glands at lid margin → not mucus-secreting.



References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 168.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [09_Conjunctiva] 020 Previous Year Questions

« Back to Folder Directory

[09_Conjunctiva] 020 Previous Year Questions — Quick Jump

Q1 (p. 1243)	*Q2 (p. 1244)*	*Q3 (p. 1245)*	*Q4 (p. 1246)*	*Q5 (p. 1247)*	*Q6 (p. 1249)*
Q7 (p. 1252)	*Q8 (p. 1252)*	*Q9 (p. 1253)*	*Q10 (p. 1258)*	*Q11 (p. 1261)*	*Q12 (p. 1264)*
Q13 (p. 1266)	*Q14 (p. 1267)*	*Q15 (p. 1270)*	*Q16 (p. 1279)*	*Q17 (p. 1280)*	*Q18 (p. 1281)*
Q19 (p. 1283)	*Q20 (p. 1287)*	*Q21 (p. 1288)*	*Q22 (p. 1290)*	*Q23 (p. 1292)*	*Q24 (p. 1293)*

Q1: What is the most frequently observed eye symptom associated with COVID-19? (Q855594)

A. Uveitis

B. Mucormycosis

C. Follicular conjunctivitis

D. Papillary conjunctivitis

Correct Answer: C. Follicular conjunctivitis

Explanation: The most frequently observed eye symptom associated with COVID-19 is follicular conjunctivitis . Follicular conjunctivitis typically presents with redness, swelling, and irritation of the conjunctiva, along with a follicular reaction on the palpebral conjunctiva. This condition has been reported in patients with COVID-19, and it is often associated with other symptoms of viral infection. Ophthalmic conditions associated with COVID-19 Category Conditions Conjunctiva Acute conjunctivitis Unremitting conjunctivitis Mild follicular conjunctivitis Severe hemorrhagic conjunctivitis Pseudomembrane formation Keratoconjunctivitis Kawasaki-like illness (MIS-C) Sclera/Episclera Episcleritis Nodular episcleritis Necrotizing anterior scleritis Sectoral anterior scleritis Anterior Chamber Acute anterior uveitis Reactivated idiopathic anterior uveitis Retina and Choroid Central retinal vein occlusion Central retinal artery occlusion Acute macular neuroretinopathy (AMN) Paracentral acute middle maculopathy (PAMM) Purtscher-like retinopathy Retinal hemorrhages Cotton wool spots Dilated veins Tortuous vessels Multifocal choroiditis (associated with vaccine) Serpiginous and ampiginous choroiditis Acute retinal necrosis (due to herpes virus reactivation) Retinal vasculitis Retinal degeneration Breakdown of blood-retinal barrier Optic Nerve Optic neuritis Neuromyelitis optica spectrum disorder Anti-myelin oligodendrocyte glycoprotein (anti-MOG) antibodies Papilledema Elevated intracranial pressure (pseudotumor cerebri syndrome) Extraocular Motility, Cranial Nerves Cranial nerve III, IV, VI palsies Binocular diplopia Ocular cranial neuropathies Postinfectious demyelinating conditions (e.g., Miller-Fisher syndrome, Guillain-Barré syndrome) Ocular myasthenia gravis Pupils Mydriasis Cholinergic super-sensitivity Tonic pupils Nystagmus Visual disturbances (oscillopsia) Viral-induced vestibular neuritis Central vestibular nystagmus (associated with rhombencephalitis) Visual Cortex Acute stroke affecting posterior visual pathways Homonymous visual field deficits Posterior cerebral artery ischemic strokes Orbit and Ocular Adnexa Sinusitis Orbital cellulitis Bacterial

orbital abscess Invasive fungal rhino-orbital mucormycosis Orbital myositis Lacrimal System Epiphora (tear overproduction) Manifestations in Newborn Infants Periorbital edema Chemosis Hemorrhagic conjunctivitis Ciliary injection Corneal edema Rubeosis Posterior synechiae Retinopathy of prematurity Vitreous hemorrhage Cotton wool spots Uveitis (Option A) : While uveitis can occur in some cases of COVID-19, it is not the most frequent eye symptom. Uveitis refers to inflammation of the uveal tract (the iris, ciliary body, and choroid), and while it has been reported in some COVID-19 patients, it is less common than conjunctivitis. Mucormycosis (Option B) : This is a rare fungal infection that can occur in COVID-19 patients, particularly in those with diabetes or who are immunocompromised. It can cause serious orbital and sinus involvement, but it is not the most frequently observed eye symptom in COVID-19. Papillary conjunctivitis (Option D) : This refers to inflammation of the conjunctiva with prominent papillae (small, raised bumps) on the inner surface of the eyelids. While it can occur in various infections, it is not as commonly reported as follicular conjunctivitis in COVID-19.

References:

Ophthalmic Manifestations of Coronavirus (COVID-19) COVID-19 and Eye: A Review of Ophthalmic Manifestations of

↑ *Back to this question in Part A*

@Neet_pg_bot


Q2: A 30-year-old contact lens wearer presents with severe pain, redness and watery discharge from the eye. Which of the following stains is most useful for diagnosis? (Q878999)

A. Gram stain

B. Giemsa stain

C. Calcofluor white stain

D. Ziehl-Neelsen stain



Correct Answer: C. Calcofluor white stain

Explanation: This presentation suggests Acanthamoeba keratitis , a severe corneal infection seen mainly in contact lens wearers with poor hygiene or exposure to tap water . Calcofluor white stain is the diagnostic test of choice , as it stains Acanthamoeba cyst walls , which fluoresce under UV light . Incorrect Options: Gram stain (Option A): Detects bacteria/fungi but not Acanthamoeba. Giemsa stain (Option B): Not sensitive or specific for Acanthamoeba. Ziehl-Neelsen stain (Option D): Used for Mycobacterium, Nocardia, etc.

References:

Bailey & Scott's Diagnostic Microbiology, 15th Edition, Page 711

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: In a patient who has had contact with a COVID-19 patient, presenting with symptoms such as redness, sensitivity to light, excessive tearing, and overall discomfort, a thorough examination was conducted with no eye abnormality. Based on this information, which specific part of the eye is most likely to be affected in this patient? (Q859774)

A. Conjunctiva

B. Uvea

C. Choroid

D. Retina

Correct Answer: A. Conjunctiva

Explanation: Correct Option A - Conjunctiva: The symptoms of redness, sensitivity to light (photophobia), excessive tearing, and discomfort are characteristic of conjunctivitis, commonly known as pink eye. Conjunctivitis has been observed as a common ocular manifestation of COVID-19, with the conjunctiva being the most likely part of the eye to be affected. This is usually a mild viral conjunctivitis that resolves with symptomatic treatment. Incorrect Options: Uvea (Option B) is typically affected in uveitis. While uveitis can cause redness, sensitivity to light, and discomfort, it is less common to occur solely as a result of COVID-19. Also, uveitis typically presents with more severe signs upon examination, such as cells in the anterior chamber or posterior synechiae, which were not noted in this case. Choroid (Option C) is a deeper structure within the eye and is not typically affected in viral conjunctivitis or mild ocular manifestations of COVID-19. Choroiditis would present with more severe symptoms and would be visible on specialized imaging such as fluorescein angiography. Retina (Option D) would typically not be directly affected by COVID-19 or the symptoms presented in this patient. Retinal conditions, such as retinitis, often present with more severe visual changes, such as visual field defects or floaters, which were not reported here.

References:

https://www.ncbi.nlm.nih.gov/books/NBK556093/#:~:text=Ocular%20manifestations%20of%20COVID-19%20often%20present%20with%20conjunctivitis%2C%20which,Bacterial%20conjunctivitis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 20-year-old male presents with fever and swelling of the left eyelid for the past two days. He has no history of trauma but experienced a mild flu-like illness one week ago. On examination, there is no chemosis or protrusion of the eyeball. Based on these observations, what is the most likely diagnosis? (Q859360)

A. Orbital cellulitis

B. Preseptal cellulitis


C. Panophthalmitis

D. Lid tumour

Correct Answer: B. Preseptal cellulitis

Explanation: Preseptal cellulitis is an infection of the eyelid and surrounding skin anterior to the orbital septum. It is characterized by eyelid swelling, erythema, fever, and tenderness, often following a minor trauma or spreading

from a nearby infection, or spreading from a remote infection of middle ear and upper respiratory tract infection. In this case, the absence of significant ocular findings, such as chemosis or proptosis, along with the mild flu-like illness history, is more consistent with preseptal cellulitis {{caption_text}} Chandler's Classification of Orbital Cellulitis Stage Description Key Features I - Preseptal cellulitis Infection confined to the eyelid Eyelid swelling, no ocular involvement II - Orbital cellulitis (Option B) Infection spreads to the orbit Pain with eye movement, restricted mobility, no abscess formation, chemosis or proptosis III - Subperiosteal abscess Pus collection beneath periosteum Localized pus, proptosis, pain on movement, swelling IV - Orbital abscess Pus collection within the orbital tissues Severe proptosis, potential vision loss, risk of optic nerve damage V - Cavernous sinus thrombosis Spread of infection to the cavernous sinus Bilateral eye involvement, headache, cranial nerve palsies Panophthalmitis (Option A) It is an intense purulent inflammation of the whole eyeball, including the tenon's capsule, characterized by severe pain,fever, malaise complete loss of vision, and purulent discharge. Lid tumor (Option D) refer to abnormal growths or masses in the eyelid.



References:

Diseases of Ear, Nose, Throat - PL Dhingra, 8th edition, Pg 233.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q5: Which statement below does not accurately describe the clinical condition mentioned? (Q859319)

A. Abnormal activity of p53 is seen in the pterygium tissue

B. Decreased activity of tissue inhibitors of metalloproteinases is seen

C. Destruction of Bowman’s layer occurs when the pterygium invades the cornea

D. Elastotic degeneration and hyalinization of the connective tissue of the conjunctivaoccur.



Correct Answer: D. Elastotic degeneration and hyalinization of the connective tissue of the conjunctivaoccur.

Explanation: The clinical image shows a pterygium , which is a subconjunctival fibrovascular growth extending onto the cornea , most commonly nasally, and associated with chronic UV exposure . Although pterygium and pinguecula share some pathological similarities, the statement in option D is incorrect because it describes histological changes seen in pinguecula , not pterygium. In a pinguecula , there is: Elastotic degeneration of collagen Hyalinization of conjunctival connective tissue No corneal invasion In a pterygium , the pathology instead involves: Fibrovascular proliferation Abnormal expression of p53 due to UV-induced damage Increased matrix metalloproteinases (MMPs) and decreased TIMPs Thus, option D does not correctly describe the condition shown (pterygium), making it the best answer. Abnormal activity of p53 is seen in the pterygium tissue (Option A): This statement is correct. Pterygium is associated with abnormal p53 expression, which is a tumor suppressor gene that plays a role in regulating cell growth and preventing the formation of tumors. Decreased activity of tissue inhibitors of metalloproteinases is seen (Option B): This statement is correct. Pterygium is associated with an imbalance in the activity of matrix metalloproteinases (MMPs) and tissue inhibitors of metalloproteinases (TIMPs), leading to excessive tissue remodeling. Destruction of Bowman’s layer occurs when the pterygium invades the cornea (Option C) : This statement is also correct as the lesion grows over the cornea and disrupts Bowman’s layer.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A contact lens user presents with the following clinical picture. He has had watering, redness, and foreign body sensation in the eye for the past 2 months. What is the most probable diagnosis? (Q856131)

A. Follicular conjunctivitis

B. Spring catarrh

C. Giant papillary conjunctivitis


D. Trachoma



Correct Answer: C. Giant papillary conjunctivitis


Explanation: efinition Specific allergic inflammation affecting the superior tarsal conjunctiva. Etiology Soft hydrophilic contact lenses Protruding suture ends Ocular prostheses Underlying Mechanism Hypersensitivity reactions (types I and IV) Symptoms Itching/ Watering Foreign body sensation Occasionally blurred vision Signs Conjunctival congestion predominantly in the upper palpebral region Large polygonal papillae on superior tarsal conjunctiva Macropapillae (0.3-1.0 mm) and giant papillae (1-2 mm) {{caption_text}} Management Discontinue soft contact lenses, remove offending sutures, clean and polish ocular prosthesis; replace with biocompatible

material (e.g., Biocoat). Ancillary therapy: Mast cell stabilizers (e.g., lodoxamide, cromolyn sodium, olopatadine) along with artificial tears, antihistaminic drops, and decongestants. Follicular Conjunctivitis (Option A): is the most common clinical form of Viral Conjunctivitis (Adenocirus , Molluscum contagiosum, Herpes simplex , COVID-19). Presents as a milder form of acute conjunctivitis (less symptoms). Follicular appearance of conjunctiva is seen on examination. {{caption_text}} Spring Catarrh (Option B): Other name of Vernal Keratoconjunctivitis (type I & IV > type I hypersensitivity reaction) , is characterized by seasonal symptoms, cobblestone-like papillae , limbal thickening with Horner-Trantas dots , and potential corneal involvement, unlike giant papillary conjunctivitis, which typically presents with large papillae associated with contact lens use. {{caption_text}} Trachoma (Option D): Sago-grain appearance of conjunctiva is seen in Trachoma. {{caption_text}}









↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: What is the most frequently observed eye symptom of COVID-19? (Q855382)

A. Retinal vein thrombosis

B. Viral keratouveitis

C. Follicular conjunctivitis

D. Pseudomembranous conjunctivitis

Correct Answer: C. Follicular conjunctivitis

Explanation: Follicular conjunctivitis is the most common ocular manifestation of COVID-19. Ocular manifestations of COVID-19: Conjunctiva Follicular conjunctivitis (Most common) Hemorrhagic conjunctivitis Pseudomembrane conjunctivitis Sclera Scleritis/ Episcleritis Anterior chamber Anterior Uveitis Posterior segment Acute macular neuroretinopathy Purtscher-like retinopathy Retinal arterial/venous occlusion (central/branched) Optic nerve Optic neuritis, optic disc edema Orbit Sinusitis, Preseptal/Orbital cellulitis Lacrimal Epiphora Newborns Periorbital edema (Most common manifestation in newborns)

References:

https://www.ncbi.nlm.nih.gov/books/NBK556093/#:~:text=Examination%20findings%20may%20include%20features,discharge%2C%20and%20mild%20eyelid%20edema .

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: Which of the following treatment options is most frequently used for recurrent pterygium? (Q857717)

A. Conjunctival autograft

B. Beta-radiation

C. Observation

D. Bare sclera technique

Correct Answer: A. Conjunctival autograft

Explanation: A conjunctival autograft (limbal conjunctival tissue from the same or other eye) is the most frequent preferred management for recurrent pterygium. Management of pterygium: Management Reassurance/Observation: if not causing any visual problems (Option C) Medical treatment: Tear substitutes for small regressive pterygium with dry eye Topical steroids for associated inflammation Protection from UV rays

with sunglasses Surgical excision for: Progression of lesion towards the pupillary area causing excessive astigmatism and visual impairment Restricted ocular movements Disfiguring Diplopia Procedures Bare sclera technique: (Option D) The pterygium is excised from the cornea and conjunctiva, leaving a bare sclera Associated with a high recurrence rate Autoconjunctival graft: (Option A) A piece of limbal conjunctival tissue from the same or the other eye or An amniotic membrane graft is used to cover the bare sclera Sutured at the limbus to cover the defect Less recurrence rate Surgical excision with lamellar keratectomy and lamellar keratoplasty for deeply infiltrating recurrent recalcitrant pterygia Old methods (not used now): McRaynold's operation Postoperative use of beta irradiations Post-operative therapy and care Prevent recurrence: Mitomycin C 0.02% drops Complications like scleral necrosis, cataracts, and iritis

References:

Kanski's Clinical Ophthalmology, 9th edition, Page- 199.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q9: What could be the most likely diagnosis for a 5-year-old boy presenting with severe itching, sensitivity to light, and sticky discharge from his eyes. Upon examination the following findings were seen? (Q857676)

A. Trachoma

B. Vernal keratoconjunctivitis

C. Phlyctenular conjunctivitis

D. Xerophthalmia




Correct Answer: B. Vernal keratoconjunctivitis

Explanation: The patient's symptoms of intense itching and sticky (ropy) discharge, along with the presence of hard, flat-topped papillae on the upper tarsal conjunctiva with a cobblestone appearance as shown in the image are characteristic of Vernal Keratoconjunctivitis (VKC). Vernal keratoconjunctivitis (Spring catarrh): It is a recurrent bilateral conjunctivitis that tends to occur with the onset of warm weather. It primarily affects young children and adolescents, particularly boys. Type I & IV HSR reaction > Type IV HSR Course The condition typically starts in childhood and may resolve on its own over 5–10 years. Symptoms Burning, Itching, Photophobia (sensitivity to light), and lacrimation (excessive tearing). Signs Palpebral Form: Hypertrophied palpebral conjunctiva with a cobblestone appearance ; bluish-white color; dense fibrous tissue with thickened epithelium. {{caption_text}} Limbal or Bulbar Form: Opacification at the limbus, gelatinous thickening, and Horner–Trantas dots {{caption_text}} Pseudogerontoxon/cupid’s bow (which resembles arcus senilis) is a grayish-white ring at the limbus seen in severe limbal vernal keratoconjunctivitis (VKC). {{caption_text}} Shield ulcers are a significant complication of VKC, particularly in severe cases. These microerosions typically form on the upper third of the cornea. {{caption_text}} Mixed form: Exhibits characteristics of both the tarsal (palpebral) and limbal subtypes, typically presenting in only one eye. Maxwell Lyon's sign is a pseudomembrane formed due to excessive mucus deposition on the papilla resulting in cobblestone papillae. {{caption_text}} Histology In the palpebral form, hypertrophied papillae are observed, rich in eosinophils and other inflammatory cells. Treatment Topical therapy: Antihistamines - Levocabastine & emedastine Mast cell stabilizers - Sodium cromoglicate Mild steroids - Fluorometholone and loteprednol etabonate, Acetylcysteine can control excess mucus. Local therapy: Subtarsal steroid injections and cryotherapy for severe cases. Surgical treatment: Excision of giant papillae and treatment of shield ulcers with debridement and amniotic membrane or bandage contact lens. Cold compresses and tinted glasses for comfort. Avoid eye rubbing Trachoma (Option A): Sago-grain appearance of conjunctiva is seen in Trachoma. {{caption_text}} Phlyctenular conjunctivitis (Option C): Type of hypersensitivity reaction to endogenous antigens like bacterial antigens, staphylococcal & tuberculosis antigens. It presents as a small, white, or yellowish nodule (phlyctenule) near the limbus and conjunctival congestion. Xerophthalmia (Option D):


Xerophthalmia is a spectrum of ocular disease caused due to vitamin A deficiency . It is characterized by night blindness (earliest feature), conjunctival xerosis, Bitot’s spot, corneal xerosis, ulcer, keratomalacia.











References:

Kanski’s Clinical Ophthalmology 9th edition, pages 181–186

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: What is the likely diagnosis for a patient who wears contact lenses and has been experiencing eye watering, redness, and a sensation of having a foreign object in the eye for the last two months? (Q857236)

A. Follicular conjunctivitis

B. Spring catarrh

C. Giant papillary conjunctivitis

D. Acute trachoma




Correct Answer: C. Giant papillary conjunctivitis


Explanation: A history of prolonged soft contact lens use , along with symptoms like bilateral itching, redness, watery discharge, and the presence of large, polygonal papillae on the upper tarsal conjunctiva, points towards a diagnosis of Giant Papillary Conjunctivitis . Giant Papillary Conjunctivitis: Definition Specific allergic inflammation affecting the superior tarsal conjunctiva. Etiology Soft hydrophilic contact lenses Protruding suture ends Ocular prostheses Underlying Mechanism Hypersensitivity reactions (types I and IV) Symptoms Itching/ Watering Foreign body sensation Occasionally blurred vision Signs Conjunctival congestion predominantly in the

upper palpebral region Large polygonal papillae on superior tarsal conjunctiva Macropapillae (0.3-1.0 mm) and giant papillae (1-2 mm) {{caption_text}} Management Discontinue soft contact lenses, remove offending sutures, clean and polish ocular prosthesis; replace with biocompatible material (e.g., Biocoat). Ancillary therapy: Mast cell stabilizers (e.g., lodoxamide, cromolyn sodium, olopatadine) along with artificial tears, antihistaminic drops, and decongestants. Follicular Conjunctivitis (Option A): is the most common clinical form of Viral Conjunctivitis (Adenocirus , Molluscum contagiosum, Herpes simplex , COVID-19). Presents as a milder form of acute conjunctivitis (less symptoms). Follicular appearance of conjunctiva is seen on examination. {{caption_text}} Spring Catarrh (Option B): Other name of Vernal Keratoconjunctivitis (type I & IV > type I hypersensitivity reaction) , is characterized by seasonal symptoms, cobblestone-like papillae , limbal thickening with Horner-Trantas dots , and potential corneal involvement, unlike giant papillary conjunctivitis, which typically presents with large papillae associated with contact lens use. {{caption_text}} Trachoma (Option D): Sago-grain appearance of conjunctiva is seen in Trachoma. {{caption_text}}




@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology 9th edition, Pages 186,187.

https://www.sciencedirect.com/science/article/pii/S1323893022000831

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: What could be the probable diagnosis for the eye lesion observed in a 21-year-old male patient? (Q857183)

A. Bitot's spots

B. Pterygium


C. Phlyctenular conjunctivitis


D. Pinguecula




Correct Answer: A. Bitot’s spots

Explanation: The provided image illustrates Bitot spots belonging to the X1B classification of xerophthalmia. Classification of Xerophthalmia: Stage Description XN Night blindness: Difficulty seeing in low light or at night. X1A Conjunctival xerosis: Dryness of the conjunctiva without foamy patches (Bitot spots). X1B Bitot spots: Presence of foamy, whitish accumulations on the conjunctiva (Option A) {{caption_text}} X2 Corneal xerosis: The cornea's dryness makes it appear dull and hazy. X3A Corneal ulceration/keratomalacia (<1/3rd of the cornea) X3B Corneal ulceration/keratomalacia (>1/3rd of the cornea) XS Corneal scar: Scarring of the cornea resulting from the healing of ulcers or keratomalacia. XF Xerophthalmic fundus: Retinal changes associated with chronic Vitamin A deficiency. The stages from X1A to X2 are reversible. Pterygium (Option B): Pterygium is a degenerative, subconjunctival fibrovascular tissue encroaching the cornea. May cause visual disturbances. {{caption_text}} Phlyctenular conjunctivitis (Option C): Type of hypersensitivity reaction to endogenous antigens like bacterial antigens, staphylococcal & tuberculosis antigens. It presents as a small, white, or yellowish nodule (phlyctenule) near the limbus and conjunctival congestion. Pinguecula (Option D): Pinguecula is a yellowish-white mound near the limbus formed by elastotic degeneration of conjunctival stroma. Only a cosmetic defect, does not cause any vision problems. {{caption_text}}


@Neet_pg_bot


Head - on Cornea
Cap - Semilunar whitish infiltrate just infront of head
Body - extended between limbus and canthus
Neck - in the limbal area
@Neet_pg_bot



References:

Park’s Textbook of Preventive and Social Medicine, 26th edition, Pg 705,706.

Kanski’s Clinical Ophthalmology 9th edition, pages 197,198,235,236.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q12: What is the causative agent of the sago-grain appearance of the conjunctiva? (Q857179)

A. Chlamydia trachomatis

B. Neisseria gonorrhoeae


C. Haemophilus influenzae

D. Vitamin A deficiency



Correct Answer: A. Chlamydia trachomatis

Explanation: The sago grain appearance of the follicles in conjunctiva is seen in the active stage of trachoma and the organism responsible is Chlamydia trachomatis . Clinical features of Trachoma: Signs of active trachoma Signs of cicatricial trachoma Conjunctival signs: Conjunctival congestion Follicles in the upper tarsal conjunctiva: Look like sago-grains Papillary enlargement: red, velvety {{caption_text}} Conjunctival signs: Scarring It may be irregular, star-shaped, or linear Linear scar in the sulcus subtarsalis is called Arlt's line accumulation of dead epithelial cells and inspissated mucus in the glands of Henle Pseudocyst Xerosis Symblepharon Corneal signs: Superficial keratitis on slit lamp and fluorescein Herbert's follicles Progressive pannus/ Trachomatous pannus : The upper part of the cornea is cloudy Corneal ulcers are common at the advancing edge of the pannus Shallow, little infiltrated Cause lacrimation and photophobia Corneal signs: Regressive pannus or pannus siccus : Receding infiltration Vessels extend beyond the area of infiltration and haze Herbert pits : oval or circular pitted scars formed after the healing of Herbert's follicles in the limbal area Vascularized and opaque cornea. Corneal xerosis Corneal ectasia Total corneal pannus (blinding sequelae) Lid signs: Distortion of the lids Tylosis : Thickening of the tarsal plate/lid margin Trichiasis: due to rubbing of lashes on the cornea Trachomatous ptosis (sleepy appearance) Entropion Madarosis Ankyloblepharon



References:

Kanski's Clinical Ophthalmology 9th edition, pages 173,174,175

↑ *Back to this question in Part A*

@Neet_pg_bot


Q13: The finding seen in the image is? (Q866832)

A. Horner-Trantas spots

B. Herbert's pits

C. Pannus

D. Arlt's line



Correct Answer: B. Herbert’s pits

Explanation: Herbert’s pits are: Small, shallow depressions at the limbus Caused by healing of limbal follicles in trachoma Considered pathognomonic of healed/previous trachoma Pathophysiology Chronic infection with Chlamydia trachomatis Formation of limbal follicles → Follicles ulcerate → Heal by fibrosis → Leave behind depressed pits at the limbus = Herbert’s pits Horner-Trantas spots (Option A): White, chalky dots at limbus composed of eosinophils in vernal keratoconjunctivitis , not trachoma. Pannus (Option C): Superficial vascularization of cornea extending from the limbus, seen in active trachoma and other chronic keratitis. Arlt’s line (Option D): Linear scar on the upper tarsal conjunctiva, seen in late scarring stage of trachoma, not at the limbus.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: Which of the following is not characteristic of pterygium? (Q855494)

A. Rate of recurrence is 95%


B. It is a fibrovascular growth


C. The parts of a pterygium are the head, neck and body


D. Has progressive and regressive types


Correct Answer: A. Rate of recurrence is 95%


Explanation: Pterygium is a fibrovascular growth that arises from the conjunctiva and extends onto the cornea, typically affecting the nasal side of the eye. (Option B ruled out) The parts of a pterygium include: (Option C ruled out) Head : The part that grows onto the cornea. Neck : The transition zone between the conjunctiva and the cornea. Body : The part on the conjunctiva. Pterygium has progressive and regressive types , meaning it can either grow over time (progressive) or shrink back (regressive) in some cases. (Option D ruled out) The rate of recurrence after surgical removal of pterygium is typically around 30-40% , not 95% . This high rate of recurrence is more common in certain settings (e.g., incomplete excision or lack of adjunctive treatments like conjunctival autografting or mitomycin C). A 95% recurrence rate is uncharacteristically high and is not the typical outcome. (Option A) Pterygium/Surfer's eye Pseudopterygium Definition Pterygium is a triangular encroachment of vascularized granulation tissue covered by conjunctiva in the interpalpebral area. Parts: Cap, head, neck, body Degenerative process Pterygium like lesion Due to cicatrizing conjunctival inflammatory overgrowth {{caption_text}} Aetiology Common in dry, sunny climates with sandy soils Ultraviolet light is a probable factor Usually elderly persons Seen at any age Caused by Trauma Chronic conjunctivitis Thermochemical burn Pathology {{caption_text}} Histology shows collagenous degenerative changes in the vascularized subepithelial stroma. Pathogenesis Degenerative subconjunctival proliferation Granulation tissue proliferates as avascularized tissue encroaches on the cornea. Destroys the superficial layers of the stroma and Bowman's membrane Symptoms Frequently follows a pinguecula Often bilateral Usually on the nasal side Impairment of vision due to: Progression into the pupillary area of the cornea Induced astigmatism by corneal flattening in the axis of pterygium Progressive pterygium: thick and vascular Regressive pterygium: thin, pale, and atrophic Ceases to grow but never disappears A probe can be passed under the lesion Can present as inflamed pterygium {{caption_text}} Non-progressive Can be at any quadrant of the corneal limbus Located away from the horizontal Attached to the cornea due to scar tissue Adhered to the adjacent lid Always stationary Signs A probe cannot be passed under the lesion Fusch's spots/Islets of Vogt: small discrete whitish flecks made of pterygial epithelial cells seen in progressive pterygium and are also called the cap of the pterygium. Stocker line: linear epithelial iron deposition, seen in regressive pterygium. {{caption_text}} A probe can be passed under the neck.




@Neet_pg_bot


@Neet_pg_bot





References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 198, 199.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: What is the causative agent of the ocular condition as shown in the image? (Q857423)

A. Chlamydia

B. Mycoplasma

C. Pseudomonas

D. Staphylococcus



Correct Answer: A. Chlamydia

Explanation: The ocular condition shown in the image is active trachoma , which is caused by the bacterium Chlamydia trachomatis . Trachoma is a leading cause of preventable blindness worldwide, particularly in areas with poor sanitation. Signs of active trachoma Signs of cicatricial trachoma Conjunctival signs: Conjunctival congestion Follicles in the upper tarsal conjunctiva: Look like sago-grains Papillary enlargement: red, velvety Conjunctival signs: Scarring It may be irregular, star-shaped, or linear Linear scar in the sulcus subtarsalis is called Arlt's line accumulation of dead epithelial cells and inspissated mucus in the glands of Henle Pseudocyst Xerosis Symblepharon Corneal signs: Superficial keratitis on slit lamp and fluorescein Herbert's follicles Progressive pannus/ Trachomatous pannus : The upper part of the cornea is cloudy Corneal ulcers are common at the advancing edge of the pannus Shallow, little infiltrated Cause lacrimation and photophobia Corneal signs: Regressive pannus or pannus siccus : Receding infiltration Vessels extend beyond the area of infiltration and haze Herbert pits : oval or circular pitted scars formed after the healing of Herbert's follicles in the limbal area Vascularized and opaque cornea. Corneal xerosis Corneal ectasia Total corneal pannus (blinding sequelae) Lid signs: Distortion of the lids Tylosis : Thickening of the tarsal plate/lid margin Trichiasis due to rubbing of lashes on the cornea Trachomatous ptosis (sleepy appearance) Entropion Madarosis Ankyloblepharon Lacrimal apparatus sequelae: Chronic dacryocystitis Chronic dacryoadenitis Images: Active Trachoma {{caption_text}} Follicles in upper palpebral conjunctiva {{caption_text}} Progressive pannus Cicatricial Trachoma {{caption_text}}


{{caption_text}} Regressive pannus {{caption_text}} Conjunctival concretions {{caption_text}} Trichiasis {{caption_text}} Tylosis (thickening of tarsal border of eyelid) WHO Classification of Trachoma (FISTO) Stage Description Image Management TF : Trachomatous Inflammation- Follicular The presence of five or more follicles in the upper tarsal conjunctiva Active stage of trachoma {{caption_text}} Antibiotics to eliminate the organism Topical: Tetracycline 1% or Erythromycin 1% Sulfacetamide Systemic: Tetracycline or erythromycin orally Doxycycline orally Azithromycin single oral dose Combined systemic and topical therapy: For TI or when an associated genital infection is present Tetracycline 1% or erythromycin 1% eye ointment Azithromycin orally TI : Trachamatous Inflammation- Intense Pronounced inflammatory thickening of the tarsal conjunctiva that obscures more than half of the normal deep tarsal vessels. {{caption_text}} TS: Trachamatous Scarring Presence of scarring in tarsal conjunctiva White, bands or sheets of fibrosis in tarsal conjunctiva {{caption_text}} Concretions removed with a hypodermic needle Conjunctival xerosis with artificial tears (lubricating drops) TT: Trachamatous Trichiasis At least one eyelash rub on eyeball or evidence of recent removal of inturned eyelashes. {{caption_text}} Permanent lash removal : Electrolysis Cryolysis Radiofrequency epilation Bilamellar tarsal resection Cicatricial entropion surgery CO: Corneal Opacity Easily visible corneal opacity over the pupil Part of the pupil margin is blurred when seen through the opacity as it is very dense corneal scarring {{caption_text}} After treatment of other trachoma sequelae Penetrating keratoplasty for significant corneal scarring Keratoprosthesis for bilateral blind cases with extensive scarring and surface problems Punctal occlusion and lateral tarsorrhaphy for coexistent eye surface problems Mycoplasma (Option B) : Although Mycoplasma can cause some ocular infections, it is not the causative agent of trachoma. Pseudomonas (Option C) : This is a common pathogen in corneal ulcers or infections, but it is not responsible for trachoma . Staphylococcus (Option D : Staphylococcus infections can cause conjunctivitis and other eye conditions, but it is not the causative agent of trachoma .





Pannus


@Neet_pg_bot





@Neet_pg_bot


@Neet_pg_bot



@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot



@Reel_pg_bot



@Reel_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot



References:

KANSKI'S Clinical Ophthalmology, 9th edition, Page- 173, 174, 175.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: Which of the following is the normal conjunctival flora? (Q860198)

A. Staph aureus

B. Staph Epidermidis


C. Strep. Pneumonia

D. H. influenzae

Correct Answer: B. Staph Epidermidis

Explanation: The normal conjunctival flora is Staph Epidermidis Staph Epidermidis is the most common organism which is responsible for Acute Onset Endophthalmitis. Acute endophthalmitis : {{caption_text}} Definition Inflammation of inner structures of the eye (uveal tissue and retina) with exudates in the vitreous, anterior, and posterior chambers. It spares sclera. Acute bacterial endophthalmitis happens 1-7 days post intraocular surgery. M/C causative organism Staphylococcus epidermidis (Option B) Presentation Severe ocular pain Redness Lacrimation Photophobia Loss of vision Red and swollen lids Conjunctival chemosis Circumcorneal congestion

Cloudy cornea Ring infiltration Yellow necrotic wound edges Hypopyon Oedematous and muddy iris Yellow pupil reflex due to purulent exudation Vitreous exudation which causes amaurotic cat's-eye reflex Raised intraocular pressure initially Treatment Intravitreal antibiotics: First choice: Vancomycin 1 mg + Ceftazidime 2.25 mg Second choice: Vancomycin 1 mg + Amikacin 0.4 mg Topical antibiotics Systemic antibiotics Steroid Therapy Supportive Therapy Vitrectomy - if no improvement in 48-72 hours or if severe infection with vision reduced to hand movements close to the face If blind - no perception of light - Evisceration. Options A, C & D: Staph aureus, Strep. Pneumonia and H. influenzae are disease causing bacteria and do not form the normal conjunctival flora



References:

AK Khurana Comprehensive Ophthalmology 7th edition, pages 176-178

https://www.ncbi.nlm.nih.gov/books/NBK559079/

Kanski's clinical ophthalmology, 9th edition, Page 327

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: Which of the following is associated with Vitamin A deficiency? (Q860158)

A. Pinguecula

B. Bitot’s spots

C. Pterygium

D. Stocker’s line

Correct Answer: B. Bitot’s spots

Explanation: Causes for Vitamin A Deficiency : Decreased intake. Obstructive jaundice: Defective absorption. Liver cirrhosis: Reduce synthesis of RBP (retinol binding protein). Severe malnutrition: Lack of amino acids for RBP synthesis. Chronic nephrosis: RBP excretion through urine. Vitamin A Deficiency Manifestations : 1. Ocular manifestations : Vitamin A deficiency is the most significant preventable cause of blindness, especially in children below 5 in developing countries. Early Signs: Loss of sensitivity to green light. Difficulty adapting to dim light. Night blindness. Prolonged Deficiency: Xerophthalmia: Dry, thick, keratinized conjunctiva and cornea. If untreated, it leads to corneal ulceration and ultimately results in blindness due to the formation of opaque scar tissue. WHO grading of xerophthalmia: Bitot's Spots: Grayish-white triangular plaques firmly adherent to the conjunctiva. Keratomalacia: When xerophthalmia persists for a longer time, it progresses to keratomalacia (corneal softening). XN Night blindness. X1 X1A: Conjunctival xerosis. X1B: Bitot spots. (Option B) X2 Corneal xerosis. X3 X3A: Corneal ulceration, less than one-third. X3B: Corneal ulceration, more than one-third. XS Corneal scar. XF Xerophthalmic fundus. 2. Immune Function: Deficiency increases susceptibility to infections. Infection reduces retinol-binding protein synthesis, lowering vitamin A levels and impairing immune responses. 3. Skin and Mucous Membrane Lesions: Follicular Hyperkeratosis (Phrynoderma): Causes rough skin due to hyperkeratinization of the follicular epithelium. Keratinizing Metaplasia: Affects respiratory, gastrointestinal, and genitourinary tracts. Epithelium is atrophied. 4. Other General Manifestation : Growth retardation due to defective synthesis of chondroitin sulfate. Decreased protein synthesis. Incorrect Options: Pinguecula (Option A) is a degenerative conjunctival condition unrelated to Vitamin A deficiency. Pterygium (Option C) is a fibrovascular growth extending from the conjunctiva onto the cornea, often caused by UV exposure rather than nutritional deficiency. Stocker’s line (Option D) is an iron line seen in front of the advancing edge of a pterygium, also unrelated to Vitamin A deficiency.

References:

Harper Edition 32, Page no: 538.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A young boy from a poor socioeconomic background presents with hyperemia and yellowish nodules at the limbus. What could be the likely diagnosis? (Q860510)

A. Bitot’s spots

B. Pterygium

C. Pinguecula

D. Phlyctenular conjunctivitis



Correct Answer: D. Phlyctenular conjunctivitis

Explanation: Poor socioeconomic status suggests that the person is prone to develop infections. The above image is of Phlyctenular conjunctivitis Phlyctenular keratoconjunctivitis Cause Allergic reactions to endogenous antigens are often linked to bacterial infections such as Staphylococcus aureus (M/C/C in Western countries) or tuberculosis (M/C/C in India). {{caption_text}} Symptoms & signs Watering, itching Reflex blepharospasm (involuntary blinking) Conjunctival congestion Phlycten - Small, round, grayish, or yellowish nodules on the bulbar conjunctiva near the limbus . Corneal involvement can lead to fascicular ulcers or ring ulcers. Pathophysiology Type IV hypersensitivity reaction. Common Complications Corneal involvement: Ulcers form on the cornea, leading to potential scarring. Associated Conditions Long-standing mild infections like tonsillitis or adenoiditis , and in populations with a high prevalence of tuberculosis . Treatment Local therapy: Corticosteroids: Drops/ointment to reduce inflammation. Address underlying cause (e.g., tuberculosis). Antibiotics Cycloplegics: Paralyze ciliary muscle, relieve pain, and prevent complications. Dark Glasses/Eye Shade: For photophobia relief. General therapy: Oral tetracycline or doxycycline is used in recurrent cases linked to Staphylococcal blepharitis. Supportive: Dark glasses or eye shade.



References:

Parsons Diseases of the Eye 23rd Edition Pg.no.: 405-406

↑ *Back to this question in Part A*

@Neet_pg_bot


Q19: Which of the following is seen in the given image? (Q844016)

A. Horner trantas spots

B. Herbert pits


C. Pannus

D. Corneal dystrophy





Correct Answer: B. Herbert pits


Explanation: Correct Option B - Herbert pits: The image shows scarring in the upper limbus, which is referred to as Herbert's pits. It is seen in the chronic cicatricial stage of trachoma. Incorrect Options: Option A - Horner trantas spots: Option C – Pannus: Option D - Corneal dystrophy: Microcystic/Map dot/ Fingerprint Dystrophy Meesmann dystrophy Reis-Bucklers corneal dystrophy




@Neet_pg_bot


@Neet_pg_bot





References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q20: A patient presents to the outpatient department with the following findings. What is the most likely diagnosis? (Q843999)

A. Pterygium

B. Limbal dermoid

C. Pyogenic granuloma

D. Pinguecula



Correct Answer: B. Limbal dermoid

Explanation: Correct Option B - Limbal dermoid: Limbal dermoids are congenital benign tumors that cause astigmatism, encroachment on the visual axis, and fatty component infiltration into the cornea. They consist of overgrowths of epidermal appendages, connective tissues, skin, fat, sweat glands, lacrimal glands, and neurologic tissues located partly over the cornea and partly over the sclera. Incorrect Options: Option A - Pterygium: A pterygium is a fibrovascular growth of the conjunctiva that extends onto the cornea. Option C - Pyogenic granuloma: A pyogenic granuloma is a red, fleshy, vascular lesion that usually arises after trauma or surgery. Option D - Pinguecula: A pinguecula is a yellowish, slightly raised lesion on the conjunctiva, usually near the limbus.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: What is a common ocular finding in pterygium? (Q853136)

A. Eye pain

B. Macular edema


C. Purulent discharge


D. Astigmatism


Correct Answer: D. Astigmatism


Explanation: A pterygium is a fibrovascular growth extending from the conjunctiva onto the cornea, often associated with UV exposure and chronic irritation. The most common visual disturbance caused by a pterygium is astigmatism, due to its pulling effect on the cornea, leading to corneal distortion and irregular refraction. Eye pain (Option A): Pterygium can cause irritation, foreign body sensation, and redness, but severe pain is not a primary feature. Pain occurs only if there is inflammation or ulceration. Macular edema (Option B): Pterygium affects the cornea and conjunctiva, but it does not involve the retina or macula. Macular edema is seen in conditions like diabetic retinopathy or retinal vein occlusion. Purulent discharge (Option C): Pterygium is a non-infectious degenerative condition, so it does not cause purulent discharge. Discharge may be seen in bacterial conjunctivitis or dacryocystitis. Pterygium/Surfer's eye Pseudopterygium Definition Pterygium is a triangular encroachment of vascularized granulation tissue covered by conjunctiva in the interpalpebral area. Parts: Cap, head, neck, body Degenerative process Pterygium like lesion Due to cicatrizing conjunctival inflammatory overgrowth Aetiology Common in dry, sunny climates with sandy soils Ultraviolet light is a probable factor Usually elderly persons Seen at any age Caused by Trauma Chronic conjunctivitis Thermochemical burn Pathology Histology shows collagenous degenerative changes in the vascularized subepithelial stroma. Pathogenesis Degenerative subconjunctival proliferation Granulation tissue proliferates as avascularized tissue encroaches on the cornea. Destroys the superficial layers of the stroma and Bowman's membrane Symptoms Frequently follows a pinguecula Often bilateral Usually on the nasal side Impairment of vision due to: Progression into the pupillary area of the cornea Induced astigmatism by corneal flattening in the axis of pterygium Progressive pterygium: thick and vascular Regressive pterygium: thin, pale, and atrophic Ceases to grow but never disappears A probe can be passed under the lesion Can present as inflammed pterygium Non-progressive Can be at any quadrant of the corneal limbus Located away from the horizontal Attached to the cornea due to scar tissue Adhered to the adjacent lid Always stationary Signs A probe cannot be passed under the lesion Fusch's spots/Islets of Vogt: small discrete whitish flecks made of pterygial epithelial cells seen in progressive pterygium and are also called the cap of the pterygium. Stocker line: linear epithelial iron deposition, seen in regressive pterygium. A probe can be passed under the neck. Management of pterygium: Management Best left alone Medical treatment: Tear substitutes for small regressive pterygium with dry eye Topical steroids for associated inflammation Protection from UV rays with sunglasses Surgical excision for: Progression of lesion towards the pupillary area causing excessive astigmatism and visual impairment Restricted ocular movements Disfiguring Diplopia Procedures Bare sclera technique: The pterygium is excised from the cornea and conjunctiva, leaving a bare sclera Associated with a high recurrence rate Autoconjunctival graft A piece of limbal conjunctival tissue from the same or the other eye or An amniotic membrane graft is used to cover the bare sclera Sutured at the limbus to cover the defect Surgical excision with lamellar keratectomy and lamellar keratoplasty for deeply infiltrating recurrent recalcitrant pterygia

Old methods (not used now): McRaynold’s operation Postoperative use of beta irradiations Post-operative therapy and care Prevent recurrence: Mitomycin C 0.02% drops Complications like scleral necrosis, cataracts, and iritis









References:

Kanski's Clinical Ophthalmology, 9th edition, Page- 198, 199.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q22: What is the likely diagnosis for a 5-year-old boy who is experiencing severe itching, and thick discharge from his eyes . Upon examination the following finding as shown in the image was observed. (Q793046)

A. Trachoma

B. Vernal keratoconjunctivitis


C. Phylctenular conjunctivitis

D. Xerophthalmia



Correct Answer: B. Vernal keratoconjunctivitis

Explanation: Correct Option B - Vernal keratoconjunctivitis: The patient's symptoms of intense itching and thick ropy discharge, along with the presence of hard, flat-topped papillae on the upper tarsal conjunctiva with a cobblestone appearance, are characteristic of vernal keratoconjunctivitis (VKC). Incorrect Options: Option A - Trachoma: Trachoma is a chronic, infectious form of conjunctivitis caused by the bacterium Chlamydia trachomatis. It is characterized by conjunctival scarring, trichiasis, and corneal opacity, which can lead to blindness if left untreated. Option C - Phlyctenular conjunctivitis: Phlyctenular conjunctivitis is a type of hypersensitivity reaction to endogenous antigens like staphylococcal & tuberculosis antigen, typically seen in children. It presents as a small, white, or yellowish nodule (phlyctenule) near limbus and conjunctival congestion.. Option D - Xerophthalmia: Xerophthalmia is a condition caused by severe vitamin A deficiency. It is characterized by dryness and thickening of the conjunctiva and cornea, which can lead to corneal ulceration and blindness.



References:

Prepladder Notes

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: What is the most effective treatment for a patient who experiences considerable discomfort when reading text from paper or a screen, despite having normal visual acuity and other eye examination findings, except for a Schirmer's test result of 5mm? (Q793044)

A. Steroids

B. Anti-histamine

C. Lubricating eye drops

D. Prescribed spectacles

Correct Answer: C. Lubricating eye drops

Explanation: Correct Option C - Lubricating eye drops: A low Schirmer's test value of 5 mm (15 mm normal value)indicates reduced tear production and dry eye syndrome. Lubricants such as artificial tears like cellulose derivatives alleviate dry eye symptoms and improve visual comfort. They work by supplementing the natural tears, improving tear film stability, and reducing ocular surface inflammation. Incorrect Options: Options A, B, and D are incorrect. They are not used for the management of Scirmer’s test.

References:

Prepladder Notes

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: What is the recommended approach for managing a middle-aged woman with a progressive pterygium that is extending towards the pupillary area? (Q793037)

A. Keratoplasty

B. Excision with conjunctival autograft

C. Observation

D. Beta-radiation

Correct Answer: B. Excision with conjunctival autograft

Explanation: Correct Option B - Excision with conjunctival autograft: The most appropriate management for pterygium encroaching the pupillary area is surgical excision with conjunctival autograft. This involves removing the pterygium tissue and replacing it with a graft of healthy conjunctival tissue taken from another area of the patient's own eye. This procedure has been shown to have a low recurrence rate and a high success rate in preventing visual disturbance caused by the pterygium. Incorrect Options: Option A - Keratoplasty: Keratoplasty involves the surgical replacement of the cornea with a healthy donor cornea. Indicated for corneal opacities, bullous keratoplasty, corneal dystrophies, and keratoconus. Option C - Observation: If the pterygium is not causing any visual disturbance, observation may be appropriate. However, if it is steadily progressing towards the pupillary area, it is likely to cause visual disturbances and should be managed surgically. Option D - Beta-radiation: Beta-radiation therapy involves using low doses of radiation to prevent recurrence of the pterygium, not recommended.

References:

Prepladder Notes

↑ *Back to this question in Part A*

@Neet_pg_bot

• [10_Ocular Adnexa] 021 Lids And Disorders - I

« Back to Folder Directory

[10_Ocular Adnexa] 021 Lids And Disorders - I — Quick Jump

Q1 (p. 1294)	*Q2 (p. 1295)*	*Q3 (p. 1299)*	*Q4 (p. 1300)*	*Q5 (p. 1301)*	*Q6 (p. 1303)*
Q7 (p. 1306)	*Q8 (p. 1307)*	*Q9 (p. 1307)*	*Q10 (p. 1309)*	*Q11 (p. 1310)*	*Q12 (p. 1311)*
Q13 (p. 1312)	*Q14 (p. 1313)*	*Q15 (p. 1314)*	*Q16 (p. 1314)*	*Q17 (p. 1315)*	*Q18 (p. 1317)*

Q1: A 6-month old baby was brought to the clinic with the following finding. Which among the following statements is incorrect regarding the likely diagnosis? (Q853470)

A. More common in boys

B. Undergoes spontaneous regression in majority

C. Oral propranolol can be used for treating this condition

D. Most common orbital tumor of children.




Correct Answer: A. More common in boys

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical scenario is suggestive of Capillary Hemangioma . Description Most common orbital tumor in children (Option D) More common in girls than boys (3:2 ratio) (Option A) Onset & Growth Visible at birth in about one-third of cases Rapid growth phase in the first 6 months ~75% regress by 7 years old spontaneously (Option B) Appearance Red, raised nodular mass ("strawberry nevus") Deep orbital mass: proptosis and globe displacement Histopathology Lobules of plump endothelial cells separated by fibrous septa Management Only done for complicated cases (affecting vision loss, systemic,etc.) and for cosmetic indications. Oral Propranolol (Option C) Intralesional corticosteroid Radiotherapy/Cryotherapy Surgery



References:

Yanoff & Duker Ophthalmology, 5th Edition, Page 1297, 1298, 1304, 1307

↑ *Back to this question in Part A*

@Neet_pg_bot


Q2: Match the image with the condition: Image Condition a. Chronic painless swelling 1. Stye b. Acute painful swelling 2. Chalazion c. Acute painful Swelling 3. Hordeolum Internum (Q853478)

A. a-2,b-3,c-1


B. a-1,b-2,c-3

C. a-3,b-2,c-1

D. a-1,b-3,c-2









Correct Answer: A. a-2,b-3,c-1

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Condition Definition & Features Chalazion Chronic non-infective lipogranulomatous inflammation of the meibomian gland. Painless swelling of the eyelid, mild heaviness, blurred vision. Hordeolum Internum Acute Suppurative painful inflammation of the meibomian gland with duct blockage. On tarsal plate Hordeolum externum (Stye) Acute suppurative painful inflammation involving glands of Zeis or Moll. Most common causative organism - Staphylococcus aureus. Acute pain with swelling on the anterior lid margin, watering and photophobia.


@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 39, 41, 67

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Which is the most common organism causing bacterial blepharitis? (Q853477)

A. Staphylococci

B. Streptococci

C. Propionibacterium acnes

D. Moraxella

Correct Answer: A. Staphylococci

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Staphylococci is the most common organism causing bacterial blepharitis Blepharitis: Inflammation of the lid margin Anterior blepharitis: Squamous type → due to dandruff on lashes Ulcerative type → infective cause → Staphylococcus aureus (most common) Posterior blepharitis → also called meibomitis Ulcerative blepharitis findings: Small ulcers at lash roots Sticky scales Management: Lid hygiene Antibiotic ointments

References:

Kanski's Clinical Ophthalmology 9th edition page 70-72

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 3-year-old girl has been having difficulty opening her left eye since birth. An image is given below. What is the condition? (Q853476)

A. Symblepharon

B. Ankyloblepharon

C. Blepharitis

D. Blepharophimosis



Correct Answer: B. Ankyloblepharon

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The image shows adhesion of the upper and lower lid, seen in Ankyloblepharon Symblepharon (Option A): Adhesion between bulbar and palpebral conjunctiva (post-burns/trauma/scarring) Blepharitis (Option C): Inflammation of the lid margin (not the whole lid) Blepharophimosis (Option D): Horizontal shortening of the palpebral fissure (seen in blepharophimosis syndrome)



References:

Kanski's Clinical Ophthalmology 9th edition page 188,189

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 45-year-old woman with hypothyroidism is now complaining of loss of eyelashes. What is the diagnosis? (Q853475)

A. Madarosis

B. Poliosis

C. Tylosis

D. Trichiasis


Correct Answer: A. Madarosis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The scenario is of loss of eyelashes (Madarosis) secondary to hypothyroidism. Madarosis: Partial or complete loss of eyelashes or eyebrows. Causes: Hypothyroidism, Leprosy, Chronic blepharitis Hertoghe's sign: Loss of lateral 1/3rd of eyebrows Poliosis (Option B): Greying of eyelashes and eyebrows. Seen in Vogt-Koyanagi-Harada syndrome, Waardenburg syndrome Tylosis (Option C) - thickening and scarring of the lid margin, a complication of long-standing blepharitis Trichiasis (Option D): Misdirection of eyelashes





References:

Kanski's Clinical Ophthalmology 9th edition page 63-66

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A neonatologist has noted the following finding in a newborn baby. What is the condition called? (Q853474)

A. Trichiasis

B. Pseudotrichiasis

C. Distichiasis

D. None of the above



Correct Answer: C. Distichiasis

- *Listen (English)*
- *Listen (Hindi)*

Explanation: The image shows an extra row of eyelashes, suggestive of distichiasis Trichiasis (Option A): Inward misdirection of lashes with the normal position of the lid margin Pseudotrichiasis (Option B) - The inward turning of lashes along with the lid margin (seen in entropion)


@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology, 9th edition, Pages 62-63

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 70-year-old patient presents with complaints of epiphora. On examination, there is an eversion of the punctum noted. Which among the following is incorrect regarding the type of ectropion seen in this patient? (Q853473)

A. Most common type of ectropion

B. Can cause lagophthalmos

C. Age-related changes are implicated in etiopathogenesis

D. Starts from lateral eyelid and progresses medially

Correct Answer: D. Starts from lateral eyelid and progresses medially

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical scenario is indicative of involutional ectropion . Involutional ectropion: Most common form of ectropion (Option A) Caused by age-related laxity of lower eyelid structures. (Option C) It usually begins medially and extends laterally . (Option D) Progression leads to: Conjunctival inflammation, Keratinization, Lagophthalmos (Option B) Other types of ectropion: Cicatricial Ectropion Paralytic Ectropion Mechanical Ectropion Congenital Ectropion

References:

Yanoff & Duker Ophthalmology. 5th Edition, Page 1287,1288

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: Match the following types of entropion with their correct pair 1) Mechanical entropion A) Trachoma 2) Cicatricial entropion B) Phthisis bulbi 3) Congenital entropion C) Senile entropion 4) Involutional entropion D) Most commonly involves lower eyelid (Q853472)

A. 1-A, 2-B, 3-C, 4-D

B. 1-C, 2-A, 3-B, 4-D

C. 1-B, 2-A, 3-D, 4-C

D. 1-B, 2-A, 3-C, 4-D

Correct Answer: C. 1-B, 2-A, 3-D, 4-C

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Mechanical entropion Due to lack of support from the globe to the lids. Eg, Enophthalmos Phthisis bulbi Cicatricial entropion Due to scarring. Eg, Chemical burns, pemphigus, SJS, symblepharon. Trachoma Congenital entropion Most commonly involves the lower eyelid Involutional entropion Affects the lower eyelid in elderly individuals. Senile entropion

References:

Yanoff & Duker - Ophthalmology 5th edition, page 1281 Kanski’s Clinical Ophthalmology 9th edition page 85-87

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: A 75-year-old male patient presents with the following ocular findings and complaints of watering and photophobia. On examination, there are signs of minimal horizontal lid laxity. Which of the following surgical procedures would be the most appropriate for a long-term correction of his condition? (Q853471)

A. Transverse sutures

B. Weis procedure

C. Jones operation

D. Quickert procedure



Correct Answer: B. Weis procedure

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient's symptoms of watering and photophobia, along with the image of an inward-turned lower lid margin, suggest entropion . The most appropriate surgical procedure for correcting entropion with minimal horizontal laxity is the Weis procedure . Weis procedure: Lid splitting with transverse sutures. Other surgical Techniques Transverse Suture and Everting Suture (Option A): Temporary cure (up to 18 months), suitable for very old patients Jones Operation (Option C): Plication of Lower Lid Retractors, used when the retractor weakness is the problem Quickert Procedure (Option D): For cases with marked horizontal lid laxity, combines horizontal lid shortening with Weis procedure

References:

Parson’s Diseases of the Eye, 23rd Edition, Pages 1054-1057

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Which among the following structures is part of the anterior lamella of the eyelid? (Q853461)

A. Orbicularis oculi

B. Orbital septum

C. Tarsal plate


D. Conjunctiva

Correct Answer: A. Orbicularis oculi

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Orbicularis oculi is part of the anterior lamella of the eyelid. The lid is made up of mainly thick connective tissue, which is called tarsal plate. The eyelid is divided by a grey line into anterior and posterior lamella: Anterior lamella structures include skin, subcutaneous tissue and orbicularis oculi (Option A) . Posterior lamella structures include tarsal plate (Option C ruled out) and conjunctiva (Option D ruled out) . The space between supraorbital margin and infraorbital margin is filled by a thin connective tissue known as orbital septum (Option B ruled out) .



References:

www.ncbi.nlm.nih.gov/pmc/articles/PMC6486383/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: All are causes of ipsilateral pseudoptosis except (Q853469)

A. Enophthalmos

B. Lid retraction

C. Hypotropia

D. Phthisis bulbi

Correct Answer: B. Lid retraction

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Lid retraction is a cause of contralateral pseudoptosis. Pseudoptosis: Apparent drooping of eyelid, unrelated to neurological or muscular abnormalities. Causes of pseudoptosis: Ipsilateral pseudoptosis Contralateral pseudoptosis Microphthalmos Enophthalmos (Option A) Phthisis bulbi (Option D) Hypotropia (Option C) Dermatochalasis Eyelid retraction (Option B) High myopia Proptosis

References:

Kanski's Clinical Ophthalmology 9th edition page 75

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 50 year male known smoker presented with hemoptysis and cough. A lung cancer was suspected. On examination, the patient had mild ptosis with good levator function. What would be the preferred management? (LPS-Levator palpebrae superioris) (Q853468)

A. Mullerectomy

B. Minimal resection of LPS

C. Maximal resection of LPS

D. Frontalis sling surgery

Correct Answer: A. Mullerectomy

Explanation: The given clinical scenario is suggestive of Small cell carcinoma of lung, which is associated with Horner syndrome. Preferred management of horner syndrome is Mullerectomy (aka) Fasanella-Servat operation. Management of ptosis: Tarso-conjunctivo-Mullerectomy (Fasanella-Servat Operation) Mild ptosis (1.5-2 mm) with good levator function Horner syndrome (Option A) LPS resection Conjunctival approach (Blaskowic's operation) Skin approach (Everbusch's operation) Mild ptosis - Minimal resection of LPS (Option B) Moderate ptosis - Moderate resection of LPS Severe ptosis - Maximal resection of LPS (Option C) Frontalis Sling Operation (Brow Suspension) Severe ptosis with zero LPS function (Option D) Marcus gunn jaw winking ptosis - LPS excision with frontalis sling surgery Neurogenic ptosis. Timing of surgery in Congenital ptosis: Mild-moderate ptosis: after 4 years of age Severe ptosis: as early as possible

References:

Kanski's Clinical Ophthalmology 9th edition page 79-80

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: Which of the following measurements is considered indicative of moderate ptosis? (Q853467)

A. 1 mm droop

B. 2 mm droop

C. 3 mm droop


D. 4 mm droop

Correct Answer: C. 3 mm droop

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Ptosis grading Mild Ptosis 2mm droop Moderate Ptosis 3mm droop Severe Ptosis 4mm droop Note: Normal lid position: Upper lid: 2mm below superior limbus Lower lid: Just touches inferior limbus



References:

Kanski's Clinical Ophthalmology 9th edition page 75 MRD 1&2

↑ *Back to this question in Part A*

Q14: Which muscle is most commonly affected in congenital ptosis? (Q853466)

A. Orbicularis oculi

B. Levator palpebrae superioris

C. Frontalis

D. Superior oblique

Correct Answer: B. Levator palpebrae superioris

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Muscle most commonly affected in congenital ptosis is Levator palpebrae superioris Levator palpebrae superioris (LPS) is the main elevator of the eyelid, while Müller's muscle contributes only ~2 mm of lift. Congenital Ptosis Defective development of LPS → congenital ptosis. Absent lid crease → due to failure of LPS to attach to skin. Lid lag on the ptotic side Types of Congenital Ptosis Simple congenital ptosis – most common, isolated LPS defect. Blepharophimosis syndrome – ptosis with reduced horizontal and vertical palpebral fissure dimensions. Marcus Gunn jaw-winking phenomenon Synkinetic ptosis. Abnormal neural connection between LPS and lateral pterygoid muscle. When the jaw moves, the eyelid flicks upward.



Mild right eye ptosis

@Neet_pg_bot

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 77,78

Congenital ptosis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: Which of the following is a distinguishing feature of aponeurotic ptosis compared to other forms of ptosis? (Q853465)

A. Accompanied by ocular motor dysfunction

B. Presents with normal function of the levator muscle

C. Results from a neurological condition affecting the third cranial nerve

D. Usually involves a sudden onset of symptoms

Correct Answer: B. Presents with normal function of the levator muscle

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Acquired Aponeurotic Ptosis: Develops due to defect of levator aponeurosis in the presence of a normal functioning muscle. Also known as involutional ptosis. Lid crease is at a higher level. Accompanied by ocular motor dysfunction (Option A): Aponeurotic ptosis is not associated with ocular motor dysfunction, which is more common in neurological forms of ptosis. Results from a neurological condition affecting the third cranial nerve (Option C): This describes ptosis due to cranial nerve III palsy, not aponeurotic ptosis. Usually involves a sudden onset of symptoms (Option D): Aponeurotic ptosis generally has a gradual onset rather than sudden, distinguishing it from other types of ptosis that may present more abruptly.

References:

Kanski’s Clinical Ophthalmology 9th edition page 74

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A 45-year-old female presents with bilateral ptosis that worsens throughout the day, particularly in the evening, and improves with rest. She also complains of double vision and generalized fatigue. Which of the following is the most likely cause of her symptoms? (Q853464)

A. Horner's Syndrome

B. Myasthenia Gravis

C. Oculopharyngeal Muscular Dystrophy

D. Botulism

Correct Answer: B. Myasthenia Gravis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The clinical scenario is suggestive of Myasthenia gravis It is a neuromuscular disorder characterized by weakness in voluntary muscles, which worsens with activity and improves with rest. It often presents with ptosis, diplopia, and generalized muscle weakness. Horner's Syndrome (Option A): Mild, constant ptosis which does not worsen with activity or improve with rest and no generalized fatigue or double vision. Oculopharyngeal Muscular Dystrophy (OPMD) (Option C): Progressive, constant ptosis but no fluctuation or improvement with rest and no daily variation in symptoms. Botulism (Option D): Sudden onset of ptosis.Symptoms do not improve with rest and not associated with daily fluctuation

References:

Kanski’s Clinical Ophthalmology 9th edition page 74

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A 60-year old male, heavy smoker, presents with persistent shoulder pain radiating to arm for several months. He presents with the following ocular findings. A chest x-ray reveals mass at the apex of the left lung. Which of the following symptoms is most likely to accompany this patient? (Q853463)

A. Mydriasis

B. Contralateral ptosis

C. Anhydrosis and miosis

D. Ophthalmoplegia



Correct Answer: C. Anhydrosis and miosis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The given clinical vignette suggests that the patient most likely has Horner’s syndrome , which can result from a Pancoast tumor at the apex of the lung. This tumor can compress the sympathetic chain. Features of Horner's syndrome: Classic triad of symptoms: Ptosis: Typically unilateral and not contralateral (Option B) Miosis (Option A): constricted pupil Ipsilateral facial anhidrosis (lack of sweating) Loss of ciliospinal reflex Enophthalmos Ophthalmoplegia (Option D): Not typically associated with Horner's syndrome. The syndrome primarily affects the sympathetic pathway and does not directly involve the muscles that move the eye.

References:

Yanoff and Duker Ophthalmology 5th edition 1275, 1276 https://www.ncbi.nlm.nih.gov/books/NBK500000/#:~:text=is%20considered%20the-,test%20of%20choice,-due%20to%20its https://www.reviewofoptometry.com/article/horners-syndrome-a-positive-apraclonidine-testnow-what#:~:text=Apraclonidine%20is%20an%20alpha%2D2%20adrenergic%20agonist%20that%20also%20acts%20as%20a%20very%20weak%20alpha%201%2Dadrenergic%20agonist%3B%20when%20topically%20applied%20in%20Horner%E2%80%99s%20syndrome%2C%20it%20causes%20dilation%20of%20the%20affected%20pupil%20and%20possibly%20lid%20elevation%2C%20without%20change%20to%20the%20normal%20pupil .

↑ *Back to this question in Part A*

@Neet_pg_bot


Q18: Identify the structure marked ‘X’ in the given image. (Q853462)

A. Orbicularis oculi muscle

B. Tarsal plate

C. Palpebral conjunctiva


D. Levator aponeurosis



Correct Answer: B. Tarsal plate

- *Listen (English)*
- *Listen (Hindi)*

Explanation: The structure ‘X’ is the tarsal plate, which belongs to the posterior lamella of the eyelid. The lid is made up of mainly thick connective tissue, which is called tarsal plate. The eyelid is divided by a grey line into anterior and posterior lamella: Anterior lamella structures include skin, subcutaneous tissue and orbicularis oculi. Posterior lamella structures include tarsal plate and conjunctiva. The space between supraorbital margin and infraorbital margin is filled by a thin connective tissue known as orbital septum .



References:

www.ncbi.nlm.nih.gov/books/NBK470320/#:~:text=Understanding%20the%20anterior%20and%20posterior,includes%20the%20conjunctiva%20and%20tarsus.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [10_Ocular Adnexa] 022 Previous Year Questions

« Back to Folder Directory

[10_Ocular Adnexa] 022 Previous Year Questions — Quick Jump

Q1 (p. 1319) *Q2 (p. 1320)* *Q3 (p. 1321)* *Q4 (p. 1322)* *Q5 (p. 1323)* *Q6 (p. 1324)*
Q7 (p. 1325) *Q8 (p. 1326)* *Q9 (p. 1327)* *Q10 (p. 1328)* *Q11 (p. 1329)* *Q12 (p. 1331)*
Q13 (p. 1332) *Q14 (p. 1333)* *Q15 (p. 1333)*

Q1: A patient presents with painful lid swelling. On examination, a pus point is observed at the base of an eyelash. Which of the following glands is involved? (Q866843)

A. Zeis gland

B. Meibomian gland

C. Lacrimal gland

D. Moll gland

Correct Answer: A. Zeis gland

Explanation: Painful lid swelling with a visible opus point (localized yellowish spot of pus) at the base of an eyelash is characteristic of external hordeolum (stye) . An external hordeolum is an acute bacterial infection (usually Staphylococcus aureus) of the sebaceous glands of Zeis located at the eyelash follicle. The pain is due to acute inflammation, and the pus may discharge spontaneously, relieving symptoms. Condition Definition & Features Treatment External Hordeolum (Stye) Acute suppurative inflammation of a lash follicle and associated glands of Zeis or Moll. Most common causative organism - Staphylococcus aureus. Symptoms: Acute pain with swelling on the anterior lid margin, watering and photophobia. Hot compresses 2-3 times daily. Evacuate pus by epilating involved cilia when pus point forms. Rarely, surgical incision for large abscess. Antibiotic eye drops and ointment. Systemic anti-inflammatory and analgesics. Systemic antibiotics for early infection control. Identify and treat predisposing conditions in recurrent cases. Incorrect Options: Meibomian gland (Option B): Sebaceous glands in the tarsal plate; acute infection → internal hordeolum (pus points to conjunctival side), chronic blockage → chalazion; not at eyelash base. Lacrimal gland (Option C): Located in the superolateral orbit; inflammation (dacryoadenitis) causes painful swelling in outer upper eyelid — not localized to eyelash follicle. Moll gland (Option D): Modified sweat glands at eyelash follicles; infection causes less common external hordeolum (Moll type) , but opus point at eyelash base is more classically due to Zeis gland infection.



References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 3-month-old child was brought to the outpatient department (OPD) with symptoms of photophobia and watering of the eyes. Which of the following is not a cause of this condition? (Q853128)

A. Congenital nasolacrimal duct obstruction

B. Congenital entropion

C. Congenital glaucoma

D. Congenital distichiasis

Correct Answer: D. Congenital distichiasis

Explanation: The hallmark of congenital distichiasis is an extra row of eyelashes that typically emerge from the openings of the meibomian glands along the posterior margin of the eyelid. It doesnot necessarily touch the cornea like entropion therefore its less likely to cause lacrimation and photophobia among the given options Distichiasis - Extra row of eyelash Clinical features Foreign body sensation Photophobia, in cases of corneal irritation Irritation

Pain Lacrimation, in cases of corneal irritation Reflex blepharospasm and photophobia (if the cornea is abraded) Conjunctival congestion Complications Recurrent corneal abrasions Superficial corneal opacities Corneal vascularization Non-healing corneal ulcer Treatment Epilation Electrolysis Cryoepilation Surgical correction Options A, B, and C: All of these conditions present with photophobia and watering of the eye.



References:

Kanski's Clinical Ophthalmology 9th edition page 62-63

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Which muscle is implicated in ptosis in Horner syndrome? (Q843986)

A. Müller muscle

B. Levator aponeurosis

C. Horner muscle

D. Orbicularis oculi

Correct Answer: A. Müller muscle

Explanation: Correct Option A - Müller muscle: Ptosis in Horner syndrome is due to the paralysis of the Müller's muscle, a smooth muscle responsible for a minor portion of upper eyelid retraction. Incorrect Options: Options B, C, and D are not responsible for causing ptosis in Horner's syndrome.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: What is the order of insertion of muscles on Annulus of Zinn? A. Medial rectus B. Inferior rectus C. Lateral rectus D. Superior rectus (Q860517)

A. A, B, C, D


B. B, C, A, D

C. D, C, A, B

D. C, A, B, D

Correct Answer: A. A, B, C, D

Explanation: Medial rectus > inferior rectus > lateral rectus > Spiral of tillaux Origin Insertion Nerve supply Recti Common annular tendon or tendinous ring of Zinn. (all recti have the same origin) Sclera little posterior to the limbus (corneoscleral junction) The average distance from Limbus is: Superior: 7.7 mm Inferior: 6.5 mm Medial: 5.5 mm Lateral: 6.9 mm Superior oblique: Trochlear nerve Lateral rectus: Abducens nerve Rest all other recti: Oculomotor nerve Options B, C, and D are incorrect.



References:

Grey's Anatomy 42nd edition, pg 773-777.

BD Chaurasia’s Human Anatomy, 8th edition, pg 232-237.

Vishram Singh Textbook of Anatomy, 2nd edition, pg 285-288.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: All of the following are functions of superior oblique muscle except? (Q840068)

A. Adduction


B. Abduction

C. Depression

D. Intorsion

Correct Answer: A. Adduction

Explanation: The superior oblique muscle primarily causes: Abduction (moves the eye outward, away from the midline) Depression (moves the eye downward when it is adducted) Intorsion (rotates the top of the eye toward the nose) It does not cause adduction , which is moving the eye inward toward the nose; that is the function of the medial rectus . Actions of Extraocular Muscles Muscle Primary Actions Superior oblique Abduction, depression (in adducted eye), intorsion Inferior oblique Abduction, elevation (in adducted eye), extorsion Medial rectus Adduction Lateral rectus Abduction Superior rectus Elevation (in abducted eye), intorsion, slight adduction Inferior rectus Depression (in abducted eye), extorsion, slight adduction Abduction (Option B): Correct function of superior oblique. Depression (Option C): Correct function of superior oblique when eye is adducted. Intorsion (Option D): Correct function of superior oblique.



References:

.Grey's Anatomy 42nd edition, pg 773-777. BD Chaurasia's Human Anatomy, 8th edition, pg 232-237. Vishram Singh Textbook of Anatomy, 2nd edition, pg 285-288.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: All of the following are features of oculomotor nerve palsy except? (Q840070)

A. Exotropia

B. Dilated pupil

C. Limited Abduction

D. Ptosis

Correct Answer: C. Limited Abduction

Explanation: Correct Option C - Limited abduction: Limited abduction (inability to move the eye outward) is not a feature of oculomotor nerve palsy. This movement is controlled by the lateral rectus muscle, which is innervated by the abducens nerve (cranial nerve VI). Dysfunction of the abducens nerve leads to issues with abduction, not

the oculomotor nerve. Incorrect Options : Option A, B, and D: . Oculomotor nerve palsy, or third nerve palsy, typically presents with several distinct features due to the dysfunction of the muscles and structures innervated by the oculomotor nerve (cranial nerve III). These features include: Ptosis : Drooping of the upper eyelid due to paralysis of the levator palpebrae superioris muscle. Exotropia : Outward deviation of the eye because the medial rectus muscle (which normally pulls the eye inward) is paralyzed. Dilated pupil : Also known as mydriasis, this occurs because the oculomotor nerve controls the constrictor pupillae muscle, and its dysfunction leads to unopposed action of the dilator pupillae muscle.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q7: Please identify the medical condition depicted in the provided image. (Q857239)

A. Entropion

B. Ectropion

C. Trichiasis

D. Ptosis



Correct Answer: A. Entropion

Explanation: The above image shows inward rolling of the lower lid margin towards the globe, which is suggestive of Entropion. {{caption_text}} Grading of entropion: Grade-I Only the posterior lid border is in-rolled Grade-II includes in-rolling up to the intermarginal strip Grade-III Whole lid margin including the anterior border is in-rolled Types of Entropion: Type Description Etiology 1) Congenital Entropion Present since birth, more common in the lower eyelid Lower Eyelid: Improper development of lower lid retractors. Upper Eyelid: Mechanical effects due to microphthalmos. 2) Cicatricial Entropion Commonly involves the upper eyelid. Common causes: Trachoma , chemical burns, pemphigus, Stevens-Johnson syndrome, symblepharon. 3) Senile (Involutional) Entropion Affects the lower eyelid in elderly individuals . Horizontal lid laxity Vertical lid instability Overriding of pretarsal orbicularis. Laxity of orbital septum 4) Mechanical Entropion Lack of support from the globe to the lids. Occurs in conditions like phthisis bulbi , enophthalmos and after enucleation or evisceration operations. Ectropion (Option B): Outward rolling of the eyelid away from the globe. Trichiasis (Option C): Misdirected eyelashes rubbing on the cornea but eyelid position is normal. Ptosis (Option D): Drooping of the upper eyelid → not related to lid inversion.



References:

Yanoff & Duker Ophthalmology 5th edition, page 1281

Kanski's Clinical Ophthalmology 9th edition pages 85-87

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: In the context of medical treatment, the V-Y procedure is employed for which of the subsequent conditions? (Q857681)

A. Entropion

B. Ectropion

C. Blepharophimosis

D. Ptosis

Correct Answer: B. Ectropion

Explanation: The V-Y procedure is used in the management of mild cases of Cicatricial Ectropion . Ectropion: Type of Ectropion Description Causes Clinical Features Treatment Involutional Ectropion Most common; age-related Horizontal laxity Canthal tendon laxity Epiphora Tarsal thinning Lagophthalmos Medial conjunctivoplasty for mild cases. Horizontal lid shortening for moderate cases. Byron Smith's modified Kuhnt-Szymanowski operation for severe cases. Lateral tarsal strip technique for generalized ectropion. Cicatricial Ectropion Scarring of anterior lamella Trauma/surgery Scarring conditions (e.g., blepharitis) Scarring Skin fibrosis Mild cases: V-Y operation. (Option B) Mild to moderate cases: Z-plasty (Elschnig's operation). Severe cases: Excision of scar tissue and full-thickness skin grafting. Paralytic Ectropion Facial nerve palsy (Bell's) Bell's palsy Trauma/surgery Exposure keratopathy Temporary measures: Topical lubricants, eyelid taping, suture tarsorrhaphy. Permanent measures (if needed): Horizontal lid tightening with/without middle lamellar buttress, palpebral sling operation. Mechanical Ectropion Mass or lesion pushing eyelid Tumors Conjunctival growths Eyelid displacement Conjunctival chemosis Treat the underlying mechanical causes. Congenital Ectropion Present at birth Tarsal plate atrophy Down syndrome Associated eyelid malformation Mild cases may not require treatment. Moderate or severe cases: Horizontal lid tightening and full-thickness skin graft. V-Y procedures have NO benefit in treatment of Entropion (Option A) , Blepharophimosis (Option C) and Ptosis (Option D) .

References:

Yanoff & Duker - Ophthalmology 5th edition, page 1281

Kanski's Clinical Ophthalmology 9th edition page 85-87

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: An 11-month-old baby was brought by his mother with complaints of the left upper eyelid moving up and down while breastfeeding that disappears when the baby is not breastfeeding. She gave a history of the child being diagnosed with left eye ptosis. What is this condition called? (Q857685)

A. Symblepharon

B. Blepharophimosis syndrome

C. Lagophthalmos

D. Marcus -Gunn jaw-winking

Correct Answer: D. Marcus -Gunn jaw-winking

Explanation: The clinical presentation is suggestive of Marcus-Gunn jaw winking ptosis , characterised by retraction of the ptotic lid with jaw movements due to stimulation of ipsilateral pterygoid muscle. Congenital synkinetic ptosis (Marcus Gunn jaw-winking ptosis): Retraction of the ptotic lid with jaw movements due to stimulation of ipsilateral pterygoid muscle. It occurs due to trigemino-oculomotor synkinesis. The pathophysiology of Marcus Gunn Jaw winking phenomenon is a congenital misdirection of mandibular branch of trigeminal nerve , i.e., this synkinetic movement results from a congenital aberrant connection between motor branches of the trigeminal nerve controlling muscles of mastication and superior division of the oculomotor nerve controlling the levator palpebrae superioris. {{caption_text}} Symblepharon (Option A): Symblepharon is the adhesion between the palpebral and bulbar conjunctiva due to burns, chemical injuries, surgery, and ulcer. Blepharophimosis syndrome (Option B): Blepharophimosis syndrome is a congenital anomaly characterized by blepharophimosis (decreased horizontal length), ptosis , congenital ectropion , and telecanthus . Lagophthalmos (Option C): Lagophthalmos is the inability to close the eyelids voluntarily .



References:

Kanski’s Clinical Ophthalmology 9th edition page 77-78

https://www.ncbi.nlm.nih.gov/books/NBK568688/#:~:text=Congenital%20ptosis%20is%20the%20presence,can%20be%20unilateral%20or%20bilateral

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: What is the most frequently occurring tumor found in the lacrimal gland? (Q855378)

A. Adenoid cystic carcinoma

B. Lymphoma

C. Pleomorphic adenoma

D. Mucoepidermoid carcinoma

Correct Answer: C. Pleomorphic adenoma

Explanation: Pleomorphic adenoma (benign mixed tumor) is the most frequently occurring tumor of the lacrimal gland . It is typically benign , slow-growing, and presents as a painless, firm mass in the superotemporal orbit. Complete surgical excision is essential to prevent recurrence or malignant transformation. Most Common Tumors : Lacrimal Gland Overall / Benign: Pleomorphic adenoma Malignant: Adenoid cystic carcinoma > Mucoepidermoid carcinoma Intraocular Tumors Benign (Adults): Cavernous hemangioma Malignant (Adults): Lymphoma Benign (Children): Dermoid Malignant (Children): Rhabdomyosarcoma

References:

Parsons Diseases of the Eye 23rd Edition, Page 424-427.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: What is the term for the additional row of eyelashes located behind the grey line? (Q858503)

A. Trichiasis

B. Distichiasis


C. Tylosis

D. Madarosis

Correct Answer: B. Distichiasis

Explanation: The case is of distichiasis - extra eyelash row Eyelash rubbing the cornea can happen in: Trichiasis - Inward misdirection of lashes with the normal position of the lid margin (Option A ruled out) {{caption_text}} Pseudotrichiasis - Inward turning of lashes along with the lid margin Distichiasis - Extra row of eyelash (Option B) {{caption_text}} Clinical features Foreign body sensation Photophobia Irritation Pain Lacrimation Reflex blepharospasm and photophobia (if the cornea is abraded) Conjunctival congestion Complications Recurrent corneal abrasions Superficial corneal opacities Corneal vascularization Non-healing corneal ulcer Treatment Epilation Electrolysis Cryoepilation Surgical correction Tylosis (Option C) is a genetic condition characterized by thickening of the lid margin. Madarosis (Option D) refers to the loss or absence of eyelashes and/or eyebrows





References:

Kanski's Clinical Ophthalmology 9th edition page 62-63

↑ *Back to this question in Part A*


Q12: Identify the instrument. (Q859327)

A. Evisceration spatula

B. Chalazion clamp

C. Enucleation spoon

D. Bone punch



Correct Answer: B. Chalazion clamp

Explanation: The above instrument is a chalazion clamp because there is a solid blade (to be kept externally on the skin). There is a ring-shaped blade that needs to be placed on the inner conjunctival surface. The handle has a thumbscrew, which can be tightened and released to achieve the required pressure. There is a forceps-like handle. After aligning the position to enclose the chalazion in the ring, the clamp is tightened and fixed. Evisceration spatula (Option A) Used for removing only intraocular contents leaving the sclera and muscles of the eye intact. {{caption_text}} Enucleation spoon (Option C) Used to remove entire intraocular contents from the eye socket leaving the lining and muscle of the eye intact.



References:

https://pmc.ncbi.nlm.nih.gov/articles/PMC10941851/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: What is the recommended treatment for a patient with Bell's palsy, who is experiencing symptoms of dry eye, redness, and irritation, along with the presence of multiple punctate epithelial erosions in the cornea as observed through a slit lamp examination? (Q859358)

A. Tarsorrhaphy

B. Contact lens

C. Corneal replacement

D. LASIK

Correct Answer: A. Tarsorrhaphy

Explanation: In the case of an elderly patient with Bell's palsy who presents with dry eye, redness, and irritation, along with multiple punctate epithelial erosions in the cornea, the most appropriate management option is Tarsorrhaphy. It is a surgical procedure in which the eyelids are partially sewn together to reduce the exposure of the cornea. Paralytic Ectropion: Cause: Facial nerve palsy (Bell's), Trauma/surgery Clinical Feature: Exposure keratopathy Treatment: Temporary measures: Topical lubricants, eyelid taping, suture tarsorrhaphy . Permanent measures (if needed): Horizontal lid tightening with/without middle lamellar buttress, palpebral sling operation Incorrect Options: Contact lens (Option B) As the cornea is already eroded contact lenses should be avoided Corneal replacement (Option C) , such as corneal transplantation or corneal grafting is not recommended because Bell's palsy will result in exposure keratopathy. LASIK (Option D) (laser-assisted in situ keratomileusis) is a refractive surgery procedure used to correct myopia, hyperopia, and astigmatism.

References:

Yanoff & Duker - Ophthalmology 5th edition, page- 1287,1288

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: In a patient presenting with recurrent chalazion, which cancer should be suspected? (Q820680)

A. Sebaceous cell carcinoma

B. Squamous cell carcinoma

C. Basal cell carcinoma

D. Kaposi's sarcoma

Correct Answer: A. Sebaceous cell carcinoma

Explanation: Sebaceous cell carcinoma arises from sebaceous glands, commonly the meibomian glands of the eyelid. It often mimics a recurrent chalazion, especially in older adults. Any chalazion that recurs or fails to resolve after standard treatment should raise suspicion for sebaceous carcinoma. Early diagnosis is important because it is locally aggressive and can metastasize. Incorrect Options: Option B) Squamous cell carcinoma: Usually presents as a painless, slowly enlarging nodule or ulcer, not as a recurrent chalazion. Option C) Basal cell carcinoma: Common eyelid malignancy; presents as pearly nodules with telangiectasia, rarely mimicking chalazion. Option D) Kaposi's sarcoma: Typically occurs in immunocompromised patients (HIV/AIDS) as violaceous plaques or nodules, not chalazion-like lesions.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: What is the likely diagnosis for a patient who has a painless, firm nodule on the upper eyelid that is growing away from the lid margin? (Q793040)

A. Stye

B. Entropion

C. Chalazion

D. Blepharitis

Correct Answer: C. Chalazion

Explanation: Correct Option C - Chalazion: A hard, painless nodule that is growing outwards, away from the lid margin is a classic presentation of a chalazion. It is typically painless but can cause discomfort or cosmetic concerns if it becomes large. Incorrect Options: Option A - Stye: A stye, also known as a hordeolum externum, is a painful, red, tender lump in the lid margin pointing anteriorly through the lid margin commonly caused by staphylococcus aureus... Option B - Entropion: Entropion is a condition in which the eyelid turns inward, causing the lashes to rub against the eye's surface. It can cause irritation, redness, and tearing. Option D - Blepharitis: Blepharitis is a chronic inflammation of the eyelids that can cause redness, itching, and flaking of the eyelid skin.



References:

,

↑ *Back to this question in Part A*

• [11_Optics] 023 Optics-Ii

« Back to Folder Directory

[11_Optics] 023 Optics-Ii — Quick Jump

Q1 (p. 1336)	*Q2 (p. 1337)*	*Q3 (p. 1338)*	*Q4 (p. 1341)*	*Q5 (p. 1342)*	*Q6 (p. 1343)*
Q7 (p. 1344)	*Q8 (p. 1345)*	*Q9 (p. 1345)*	*Q10 (p. 1352)*	*Q11 (p. 1356)*	*Q12 (p. 1359)*
Q13 (p. 1359)	*Q14 (p. 1360)*	*Q15 (p. 1361)*	*Q16 (p. 1362)*	*Q17 (p. 1363)*	*Q18 (p. 1364)*


Q1: Worth's Four Dot Test for a 6 y/o child reveals that the child sees two red and three green lights as shown below in the image. Based on the test result, what is the most likely diagnosis in this child? (Q853632)

A. Binocular Single Vision

B. Left Eye Suppression

C. Right Eye Suppression

D. Diplopia




@Neet_pg_bot

Correct Answer: D. Diplopia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Seeing two red lights and three green lights on Worth's Four Dot Test indicates the presence of diplopia (double vision) , where the patient perceives two distinct images due to misalignment. Worth’s Four Dot test Procedure: The patient wears a red lens on the right eye and a green lens on the left eye. They are shown a box with four lights: one red, two green, and one white. The patient is asked how many lights they see. Results: All four lights: BSV (binocular single vision). (Option A ruled out) All four lights with manifest squint: Harmonious ARC (anomalous retinal correspondence). Two red lights: Right eye suppression. (Option C ruled out) Three green lights: Left eye suppression. (Option B ruled out) Alternating red and green lights: Alternating suppression. Two red and three green lights: Diplopia (double vision).



References:

Kanski’s clinical ophthalmology, John F. Salmon, 10th edition, Pages 722-725

uomus.edu.iq/img/lectures21/MUCLecture_2024_31858861.pdf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A 25-year-old patient presents with blurred vision and headaches. On examination, the patient is found to have regular astigmatism with the following prescription: Sphere: +2.00 D, Cylinder: -1.50 D, Axis: 90°.What is the refractive power of the eye along the 90° meridian? (Q853640)

A. +2.00 D

B. +0.50 D

C. +3.50 D

D. +1.25 D

Correct Answer: A. +2.00 D

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: To calculate the refractive power along the 90° meridian: Sphere (+2.00 D): This is the baseline power applied equally in all meridians. Cylinder (–1.50 D, Axis 90°): The axis tells us the meridian with no cylinder effect. At 90° (axis meridian) → only the sphere acts. At 180° (perpendicular meridian) → sphere + cylinder = +2.00 + (–1.50) = +0.50 D . So: Power along 90° meridian = +2.00 D (Option A) Power along 180° meridian = +0.50 D

References:

Myron Yanoff and Duker's Textbook of Ophthalmology, 6th Edition, Page 31-33

www.ncbi.nlm.nih.gov/books/NBK582142/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 45 y/o patient presents with difficulty in reading and blurred vision. An eye examination reveals astigmatism. Based on the classification of astigmatism, which of the following statements is accurate? (Q853639)

A. In With-the-Rule (WTR) astigmatism, the horizontal meridian has greater curvature than the vertical meridian.

B. Compound myopic astigmatism is characterized by both of the eye's two principalmeridians falling in front of the retina

C. Irregular astigmatism involves a predictable pattern of refractive power changes along the meridians.


D. Bi-oblique astigmatism has principal meridians that are perpendicular to each other.


Correct Answer: B. Compound myopic astigmatism is characterized by both of the eye's two principalmeridians falling in front of the retina


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In compound myopic astigmatism, the two principal meridians create two separate focal lines, and both focal lines fall in front of the retina. Regular Astigmatism: When refraction error changes uniformly from one meridian to the other (2 principle meridians present) Type Description With-the-Rule (WTR) The vertical meridian has greater curvature than the horizontal meridian. (Option A ruled out) Against-the-Rule (ATR) The horizontal meridian is more curved than the vertical meridian. Oblique Astigmatism Principal meridians are at oblique angles (e.g., 45° and 135°), still perpendicular to each other. Bi-oblique Astigmatism Principal meridians are not perpendicular to each other (e.g., 30° and 100°). (Option D ruled out) Irregular Astigmatism: Involves multiple, changing meridians of refractive power, making it geometrically complex. (Option C ruled out). Usually due to Corneal scars.







References:

Yanoff and Duker's Textbook of Ophthalmology, 6th Edition, Pages 31-33

www.ncbi.nlm.nih.gov/books/NBK582142/

www.ncbi.nlm.nih.gov/books/NBK587355/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: Which of the following statements about common treatments for presbyopia is correct? (Q853638)

A. Bifocal lenses are designed to provide only distance vision and are not effective for near vision.

B. Multifocal IOLs (intraocular lenses) provide multiple focal points to help with both near and distant vision.

C. Conductive Keratoplasty (CK) offers permanent benefits for presbyopia.

D. Corneal multifocality treatment involves the replacement of the Natural lens with an IOL.

Correct Answer: B. Multifocal IOLs (intraocular lenses) provide multiple focal points to help with both near and distant vision.

Explanation: Multifocal IOLs are designed to offer multiple focal points, allowing for improved near and distance vision. I. Surgical and Lens-Based Treatments for Presbyopia: Treatment Description Advantages Limitations Lens Extraction and IOL Implantation Removal of the natural lens and replacement with an artificial IOL. Types: Clear Lens Exchange (CLE) Refractive Lens Exchange (RLE) Presbyopic Lens Exchange (PreLEx) Permanent solution; maintains focus at varying distances. Potential issues with glare and contrast sensitivity. Multifocal IOLs (Option B) Lenses with multiple focal points for different distances. Reduces or eliminates the need for glasses. Potential for glare and reduced contrast sensitivity; ~10% may need IOL exchange. Bifocal IOLs Lenses with two focal points for distance and near vision. (Option A ruled out) Provides clear vision at two distances. May not fully address all visual needs. Conductive Keratoplasty (CK) Uses radiofrequency energy to reshape the cornea, potentially providing multifocal functionality. Can offer benefits for presbyopia. Primarily used for hyperopia ; results may not be permanent. (Option C ruled out) II. Emerging and Experimental Treatments for Presbyopia: Treatment Description Advantages Limitations Laser-Induced monovision Uses LASIK or PRK to reshape one eye for near vision while keeping the other for distance, reducing the need for glasses or contacts. Less need for glasses or contacts for both near and distant vision. Vision correction tailored to individual needs. Rapid visual recovery post-procedure. Touch-ups can maintain optimal vision if needed. Difficulty adjusting, leading to imbalance or discomfort. Impaired spatial judgment, affecting activities like driving. Less clarity in distance or near vision. Further near vision deterioration with presbyopia, requires additional correction. Corneal Multifocality Laser procedures to reshape the cornea to create a multifocal effect. Potential to address presbyopia without lens replacement. (Option D ruled out) Experimental; still in development. Scleral Expansion Surgery Alters the sclera to restore accommodation. Aims to restore accommodation. Inconsistent results; less popular. Intracorneal Inlays Small devices are inserted into the cornea to correct presbyopia. Can significantly benefit presbyopia. Issues with biocompatibility; potential complications. Laser Modification of the Natural Lens Femtosecond lasers modify the natural lens's elasticity to restore accommodation. An experimental approach aiming to restore natural focus.

Remains experimental; not widely available. III. Low Vision Aids for Presbyopia: Type Description Functional Benefits Optical Devices Magnifiers: Handheld, stand, and electronic magnifiers to enlarge text and images. Bifocal and Multifocal Lenses: Provide different optical powers for distance and near vision. (Option A ruled out) Improves ability to see close-up objects; helps with reading and other tasks. Non-Optical Devices Lighting: Enhanced lighting solutions, such as task lamps. Large Print Materials: Books and labels in larger fonts. Improves visibility; aids in reading and daily tasks.

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 286

www.ncbi.nlm.nih.gov/books/NBK585124/

www.visioncenter.org/lasik/monovision/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 45 y/o patient presents with significant astigmatism and is seeking surgical correction. The patient is not a candidate for LASIK. Which of the following procedures is most suitable for correcting high degrees of astigmatism in this case? (Q853637)

A. Limbal Relaxing Incisions (LRI)

B. Photorefractive Keratectomy (PRK)

C. Toric Intraocular Lens (IOL) Implantation

D. Conductive Keratoplasty (CK)

Correct Answer: C. Toric Intraocular Lens (IOL) Implantation

Explanation: Toric Intraocular Lens (IOL) Implantation is appropriate for correcting high degrees of astigmatism, as it uses a specially designed lens with varying refractive powers to address significant astigmatism. Astigmatism Correction: I. Incisional Techniques: Procedure Description Suitability Limitations Limbal Relaxing Incisions (LRI) / Arcuate Keratotomy (Option A ruled out) Paired arcuate incisions were made on opposite sides of the steeper corneal meridian to flatten it, thereby reducing astigmatism. Compression sutures may be added for significant cases. Suitable for reducing large corneal astigmatism, often post-keratoplasty. (unlike this patient) Correction level depends on incision parameters. II. Laser-Based Procedures: Procedure Description Suitability Limitations LASIK Involves creating a corneal flap and using an excimer laser to reshape the underlying stroma, correcting both spherical and cylindrical errors. Contraindicated in patients with corneal thickness less than 450µm. Astigmatism up to 5D Faster recovery, but flap complications are possible. PRK / LASEK (Option B ruled out) Surface ablation techniques using an excimer laser to reshape the cornea after removing the epithelium. Corrects up to 3D of astigmatism. Contraindicated in patients unsuitable for LASIK. Low to moderate astigmatism

Longer healing time, higher discomfort. III. Lens-Based Surgery Procedure Description Suitability Limitations Toric Intraocular Lens (IOL) Implantation (Option C) Replacement of the natural lens with a toric IOL designed to correct astigmatism by having different refractive powers in different meridians. Suitable for astigmatism correction during cataract surgery or high degrees of astigmatism. Implant rotation postoperatively can reduce effectiveness. IV. Conductive Keratoplasty (CK) Procedure Description Suitability Limitations Conductive Keratoplasty (CK) (Option D ruled out) Radiofrequency energy heats and shrinks collagen fibres in the peripheral cornea, increasing corneal curvature. Best suited for low to moderate hypermetropic astigmatism. Results may not be permanent and could regress over time.

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 285

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Which of the following surgical procedures is most suitable for correcting hypermetropia up to +4D and allows for faster visual recovery? (Q853636)

A. Photorefractive Keratectomy (PRK)

B. Conductive Keratoplasty (CK)

C. Laser Thermal Keratoplasty (LTK)

D. Laser in situ Keratomileusis (LASIK)

Correct Answer: D. Laser in situ Keratomileusis (LASIK)

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: LASIK is ideal for correcting hypermetropia up to +4D and offers faster visual recovery and greater postoperative comfort compared to surface ablation procedures like PRK. Treatment Modalities for Hypermetropia: I. Laser Refractive Procedures: Procedure Suitability Limitations LASIK Moderate hypermetropia (up to +4D) Faster recovery, more comfortable than PRK. Surface Ablation - Photorefractive Keratectomy (PRK) (Option A ruled out) Low to moderate hypermetropia Longer healing time, possible regression. Laser Thermal Keratoplasty (LTK) (Option C ruled out) Low degrees of hypermetropia Effectiveness may diminish over time. II. Conductive Keratoplasty (CK): Procedure Description Suitability Limitations Conductive Keratoplasty (Option B ruled out) Radiofrequency energy heats and shrinks collagen fibres in the peripheral cornea, increasing corneal curvature. Low to moderate hyperopia May require repeat procedures; non-permanent.

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 285

↑ *Back to this question in Part A*

Q7: A 30-year-old patient with a history of myopia is considering LASIK surgery. Which of the following is NOT an indication of the procedure? (Q853635)

A. The patient’s refraction has been stable for the past 14 months

B. The patient is over 20 years old

C. Presence of corneal ectasia


D. Myopia of -7 D

Correct Answer: C. Presence of corneal ectasia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: LASIK is contraindicated if there is a presence of ectasia or any other corneal pathology. LASIK (Laser-Assisted In Situ Keratomileusis): Indications: Myopia up to -8 D. (Option D ruled out) Patients must be over 20 years old, have stable refraction for at least 12 months, be motivated, and have no corneal pathology. (Options A & B ruled out) Contraindicated in cases of corneal ectasia or any other corneal pathology.



References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 287-288

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 32 y/o man with -14 D myopia presents for a consultation regarding refractive surgery. He is not a candidate for laser-based procedures due to thin corneas. Which of the following is the most appropriate surgical option for his condition? (Q853634)

A. Phakic intraocular lens (IOL) implantation

B. Radial keratotomy (RK)

C. Clear lens extraction

D. Intracorneal ring (ICR) implantation

Correct Answer: A. Phakic intraocular lens (IOL) implantation

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Phakic IOL implantation is a suitable choice for patients with high myopia (generally greater than -8 D) who are not candidates for laser-based refractive surgery, such as LASIK, due to thin corneas. In this procedure, a lens is implanted in front of the natural lens to correct vision without removing the natural lens, preserving accommodation. Refractive surgeries for myopia Radial Keratotomy (RK) (Option B ruled out) Involves radial cuts in the cornea to flatten it, correcting myopia. Used for low to moderate myopia (2-6 D). No longer recommended due to the advent of safer procedures like LASIK. Clear Lens Extraction (Option C ruled out) Removal of the natural lens (without cloudiness) to correct very high myopia (-16 D to -18 D). Less common now due to risks like retinal detachment. Intracorneal Rings (ICR) : (Option D ruled out) Rings are inserted into the cornea to flatten it and reduce refractive power. Results can be unpredictable, with risks such as corneal inflammation, so is not recommended

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 284-286

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Match the following 1) Titmus fly test a) 2) TNO test b) 3) Frisby test c) 4) Random Dot E test d) (Q853633)

A. 1 - c,2 - b,3 - a,4 - d

B. 1 - b,2 - c,3 - a,4 - d

C. 1 - b,2 -c,3 - d,4 - a

D. 1 - c,2 - b,3 - d,4 - a



@Neet_pg_bot



@Neet_pg_bot



TNO Test

@Neet_pg_bot



@Neet_pg_bot

Correct Answer: B. 1 - b,2 - c,3 - a,4 - d

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Tests for stereopsis Titmus fly test Random Dot E test Lang test Random dot test Frisby test TNO Test



@Neet_pg_bot



@Neet_pg_bot



Lang Test

@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot

References:

Kanski’s clinical ophthalmology, John F. Salmon, 10th edition, Page no: 720-722

www.seevividly.com/info/Binocular_Vision/Vision_Tests/Tests_of_Stereopsis#:~:text=Common%20Stereopsis%20Tests%20Used%20in%20the%20Eye%20Clinic&text=The%20most%20common%20targets%20are,to%20400%20seconds%20of%20arc.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Match the following images appropriately: 1.Streak Retinoscope a) 2. Priestly Smith Retinoscope b) 3.Trial Frame c) 4.Refraction Box d) (Q853616)


A. 1- a, 2- d, 3- c, 4- b

B. 1- b, 2- c, 3- a, 4- d

C. 1- c, 2- a, 3- d, 4- b

D. 1- d, 2- c, 3- a, 4- b






@Neet_pg_bot




@Neet_pg_bot



Correct Answer: D. 1- d, 2- c, 3- a, 4- b

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Aspect Image Streak Retinoscope: Self-illuminated retinoscope. More sensitive than spot retinoscopy More commonly used due to its enhanced ability to detect astigmatism. Priestly Smith Retinoscope: A specific type of mirror retinoscope. Combines plane and concave mirrors. Trial Frame: Pre-requisite for retinoscopy. Adjustable type, suitable for use in both children and adults. Refraction Box: Contains a variety of spherical and cylindrical lenses with plus and minus powers. Includes a pinhole, occluder, and prisms.






@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th Edition, Page 625, 811, 834, 842, 934, 942

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A 7-year-old child is evaluated for binocular vision using a synoptophore. The child successfully perceives two different images simultaneously and can superimpose incomplete but similar images into a single complete image. Which grade of binocular single vision is demonstrated by this child? (Q853631)

A. Grade I - Simultaneous Perception

B. Grade II - Fusion

C. Grade III - Stereopsis


D. None of the above


Correct Answer: B. Grade II - Fusion


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The ability to superimpose two incomplete but similar images into one complete image is characteristic of Grade II - Fusion. Grades of Binocular Single Vision Grade I - Simultaneous Perception Grade II - Fusion Grade III - Stereopsis Ability to see two dissimilar objects simultaneously. (Option A ruled out) Tested by projecting different images to each eye (e.g., bird in one eye, cage in the other). Example: Seeing a bird inside a cage. Power to superimpose two incomplete but similar images into one complete image. Tested by manipulating eye convergence/ divergence for positive and negative fusion ranges. Ability to perceive depth (third dimension). (Option C ruled out) Tested with stereopsis slides using the synoptophore.







References:

Kanski’s clinical ophthalmology, 10th edition, Page no: 716, 724-725

webeye.ophth.uiowa.edu/eyeforum/tutorials/bhola-binocularvision.htm#gsc.tab=0

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 7-year-old child with amblyopia in the right eye is prescribed occlusion therapy. The optometrist recommends patching the sound left eye for a certain number of hours each day. What is the typical recommended duration of patching for effective treatment in children with amblyopia? (Q853630)

A. 1-2 hours per day

B. 3-4 hours per day

C. 4-6 hours per day

D. 12 hours per day

Correct Answer: C. 4-6 hours per day

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: For effective occlusion therapy in treating amblyopia, it is typically recommended to patch the stronger eye for 4-6 hours per day as per ‘Amblyopia Treatment Study’. This duration helps stimulate the weaker eye (amblyopic eye) and encourages improvement in its visual acuity. Occlusion therapy: Occlusion of the sound (normal) eye to force use of amblyopic eye is the main stay in the treatment of amblyopia Full-time occlusion: Up to 2 years, the occlusion should be done in 2:1 , i.e. 2 days in sound eye and one day in amblyopic eye; at the age of 3 years, 3:1 ; at the age of 4 years, 4:1 ; At the age of 5 years, 5:1 ; and after the age of 6 years 6:1 . Part-time occlusion: 4-6 hours/day , as shown by Amblyopia Treatment Study (ATS), and is as effective as full-time occlusion. Maintenance occlusion: After vision equalization, part-time occlusion continues until at least 8 years of age. Ensure no media opacity or refractive errors before starting occlusion therapy.

References:

Kanski’s clinical ophthalmology, 10th edition, Page no: 718-719

pmc.ncbi.nlm.nih.gov/articles/PMC2907781/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 7-year-old girl who had a H/0 congenital cataract is presenting with decreased vision in her right eye. Which type of amblyopia is most likely in this case? (Q853629)

A. Strabismic amblyopia

B. Toxic amblyopia

C. Anisometropic amblyopia

D. Stimulus deprivation amblyopia

Correct Answer: D. Stimulus deprivation amblyopia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Stimulus deprivation amblyopia occurs when there is a physical obstruction preventing adequate visual input from reaching the brain, such as a congenital cataract. In this case, the Congenital cataract is blocking the vision in the right eye, leading to amblyopia, also known as "lazy eye." Types of Amblyopia: Type Cause Key Features Strabismic Amblyopia (Option A ruled out) The brain suppresses input from one eye due to constant squint (strabismus). Common cause; underdevelopment of vision in the deviated eye. Stimulus Deprivation Amblyopia (Option D) Hindered vision due to obstruction from early life (e.g., congenital cataracts, ptosis) . Lack of visual stimulation affects normal development. Anisometropic Amblyopia (Option C ruled out) A significant difference in refractive error between eyes. The brain favours the clearer eye, leading to amblyopia in the other. Toxic amblyopia (Option B): Toxic amblyopia, also known as toxic/nutritional optic neuropathy, is caused by damage to the optic nerve fibres due to exposure to toxins like tobacco, alcohol, and certain medications.

References:

Kanski’s clinical ophthalmology, 10th edition, Page no: 718-719

pmc.ncbi.nlm.nih.gov/articles/PMC2907781/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: Which of the following is an absolute optical indication for contact lens use? (Q853621)

A. Corneal ulcer

B. Bullous keratopathy

C. Unilateral aphakia

D. Aniridia

Correct Answer: C. Unilateral aphakia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Unilateral aphakia (the absence of the lens in one eye) is an absolute optical indication for contact lens use. A contact lens can provide the necessary optical correction for the missing lens, which is crucial for restoring proper vision in the affected eye. Indications for Contact Lens Use: Optical Indications: Cosmetic: For patients with refractive errors seeking aesthetic benefits. Absolute Indications: Anisometropia Unilateral aphakia Keratoconus Irregular astigmatism Therapeutic Indications: Corneal diseases : Corneal ulcers, bullous keratopathy. (Options A & B) Iris diseases : Aniridia, coloboma, albinism. (Option D) Glaucoma Amblyopia Post-surgery: Bandage lenses promote healing after keratoplasty and micro corneal perforations. Preventive Indications: Chemical burns : Prevents symblepharon and preserves fornix depth. Exposure keratitis : Protects the cornea in cases of incomplete eyelid closure. Trichiasis : Shields cornea from irritation due to misdirected eyelashes. Other miscellaneous causes: Used in procedures like gonioscopy, electroretinography, and fundus photography. Utilized during surgeries such as goniotomy, vitrectomy, and endocular photocoagulation. Masks corneal scars, corrects ptosis, and improves appearance in phthisis bulbi with cosmetic scleral lenses. Ideal for athletes, pilots, and actors for a wider field of view and stable vision.

References:

Kanski’s clinical ophthalmology, John F. Salmon, 10th edition, Page no: 268-269

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 25-year-old female with myopia is looking for contact lenses that will be comfortable for long hours of wear and require minimal maintenance. She is involved in outdoor activities and is seeking a lens that provides comfort, ease of use, and reduced risk of irritation over extended periods. Which type of contact lens would be most suitable for her? (Q853620)

A. Hard contact lenses

B. Rigid gas permeable lenses

C. Soft contact lenses

D. PMMA lenses

Correct Answer: C. Soft contact lenses

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Soft contact lenses are the most suitable for long wear due to their comfort, flexibility, and lower risk of causing irritation to the eye. Type of Contact Lens Material Key Properties Hard Lenses (Option A) PMMA (polymethylmethacrylate) (Option D) High optical quality, durable, cheap. RGP Lenses (Option B) Silicone acrylate, Cellulose acetate butyrate (CAB) Oxygen permeable, hard. Soft Lenses (Option C) HEMA (hydroxyethyl-methacrylate), Elastomer Soft, pliable, hydrophilic, oxygen permeable, comfortable and well tolerated.

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 47-48

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: During subjective refraction, an optometrist uses a Jackson Cross Cylinder (JCC) lens as part of the assessment. Which of the following is the primary purpose of this test? (Q853619)

A. To determine the spherical refractive error

B. To measure the visual acuity at distance

C. To verify the axis and power of the cylindrical correction

D. To test for binocular vision and eye alignment

Correct Answer: C. To verify the axis and power of the cylindrical correction

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The Jackson Cross Cylinder (JCC) is used during the refraction process to verify and fine-tune both the axis and power of the cylindrical correction for astigmatism. JCC consists of two cylinders with equal strength but opposite signs, placed at right angles to each other. It is mounted in a handle and commonly used with strengths of ±0.25D and ±0.50D. The test helps to verify the power and axis of a cylindrical lens correction. Power check: When placed along the cylinder axis, the JCC helps determine whether more or less cylindrical power is needed. Axis check: When placed at 45° to the cylinder axis, it helps refine the correct orientation of the astigmatism correction.



References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 37-40

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A 38-year-old patient presents for an eye examination. During the duochrome test, the patient reports that they can see the red letters more clearly than the green letters. What is the most likely refractive error for this patient? (Q853618)

A. Hyperopia

B. Myopia

C. Astigmatism

D. Presbyopia

Correct Answer: B. Myopia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Duochrome Test Refines the spherical component of the prescription The patient views red and green letters to check for chromatic aberration. Individuals with myopia (nearsightedness) tend to see the red letters (longer wavelength) more clearly. Conversely, individuals with hyperopia (farsightedness) would generally see the green letters (shorter wavelength) more clearly. Both clear : Emmetropia . The longer wavelength (red) is refracted less than the shorter (green). Hyperopia (Option A): These patients would see the Green letters more clearly because their eye is "too weak," pushing the green focus point closer to the retina. Astigmatism (Option C): While astigmatism affects clarity, the duochrome test is specifically used to fine-tune the spherical component of a prescription, not the cylinder (astigmatism). Presbyopia (Option D): This is an age-related loss of accommodation (near-focusing power). While it affects vision, the duochrome test is used to measure the refractive state of the eye

at a distance.




Fig. 25.24 *A, Optical principle of duochrome test; B, Duochrome test box consisting of Snellen's 6/18 to 6/5 visual acuity letters with red and green background*

@Neet_pg_bot

References:

Kanski's Clinical Ophthalmology, John F. Salmon, 10th Edition, Page 2

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: A patient presents with a spectacle prescription of -4.00 D sphere and +1.00 D cylinder at 90°. What is the transposed prescription? (Q853617)

A. -3.00 D sphere -1.00 D cylinder at 180°

B. -5.00 D sphere -1.00 D cylinder at 90°

C. +4.00 D sphere -1.00 D cylinder at 90°

D. -1.00 D sphere +4.00 D cylinder at 180°

Correct Answer: A. -3.00 D sphere -1.00 D cylinder at 180°

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: -3.00 D sphere -1.00 D cylinder at 180° is the transposed spectacle prescription. Transposition Process: Calculate the New Spherical Value: Formula: New Spherical = Original Spherical + Cylinder Calculation: -4.00 D sphere + 1.00 D cylinder Result: -4.00 + 1.00 = -3.00 D sphere Change the Sign of the Cylinder: Original Cylinder Value: +1.00 D Transposed Cylinder Value: -1.00 D Adjust the Axis of the Cylinder: Original Axis: 90° Adjustment: Add 90° to the original axis Calculation: 90° + 90° = 180° Transposed Prescription: New Spherical Value: -3.00 D New Cylinder Value: -1.00 D New Axis: 180° Correct Transposed Prescription: -3.00 D sphere -1.00 D cylinder at 180°

References:

www.ncbi.nlm.nih.gov/books/NBK597365/

↑ *Back to this question in Part A*

@Neet_pg_bot

• [11_Optics] 024 Optics-I

« Back to Folder Directory

[11_Optics] 024 Optics-I — Quick Jump

Q1 (p. 1366)	*Q2 (p. 1368)*	*Q3 (p. 1368)*	*Q4 (p. 1370)*	*Q5 (p. 1371)*	*Q6 (p. 1372)*
Q7 (p. 1372)	*Q8 (p. 1374)*	*Q9 (p. 1375)*	*Q10 (p. 1376)*	*Q11 (p. 1377)*	*Q12 (p. 1378)*
Q13 (p. 1379)	*Q14 (p. 1382)*	*Q15 (p. 1383)*	*Q16 (p. 1384)*	*Q17 (p. 1387)*	*Q18 (p. 1388)*
Q19 (p. 1389)	*Q20 (p. 1389)*	*Q21 (p. 1390)*	*Q22 (p. 1392)*	*Q23 (p. 1394)*	*Q24 (p. 1399)*
Q25 (p. 1404)					

Q1: Which of the following are features of direct ophthalmoscopy? Erect, virtual image is formed 2.5x magnification Large field of view Peripheral retinal lesions can be examined Routine fundus examination Semi-dark room required (Q853802)

A. 1,5,6

B. 1,2,3,4,5,6

C. 1,3,6

D. 3,4

Correct Answer: A. 1,5,6

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Characteristic Direct Ophthalmoscopy Indirect Ophthalmoscopy Image of ophthalmoscope Image Characteristics Erect, virtual (Statement 1) Real, inverted Magnification About 15x (Statement 2) About 3–5× with a +20D lens Field of View Small (~2 DD) (Statement 3) Larger field of view (up to 8 DD) Uses Central retina (Statement 4); Routine fundus examination (Statement 5) Essential for assessing retinal detachment and peripheral retinal lesions Prerequisites Semi-darkroom (Statement 6) Darkroom





References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 52-53

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 21-23

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Match the following refractive prescriptions with their corresponding types of astigmatism: Prescription Astigmatism Type i) +2D Cyl 1. Simple hypermetropic ii) +1D Sph & +2D Cyl 2. Compound hypermetropic iii) -2D Sph & +3D Cyl @180° 3. Mixed with the rule iv) -2D Sph & -1D Cyl @90° 4. Compound myopic v) -1D Sph & +2D Cyl @90° 5. Mixed against the rule (Q853841)

A. i–1, ii–2, iii–3, iv–4, v–5

B. i–1, ii–2, iii–5, iv–4, v–3

C. i–2, ii–1, iii–5, iv–4, v–3

D. i–1, ii–3, iii–5, iv–2, v–4

Correct Answer: A. i–1, ii–2, iii–3, iv–4, v–5

Explanation: +2D Cyl (i) : Only one meridian is ametropic (hypermetropic), other is emmetropic → Simple hypermetropic astigmatism +1D Sph & +2D Cyl (ii) : Both meridians are hypermetropic but of different powers → Compound hypermetropic astigmatism -2D Sph & +3D Cyl @180° (iii) : One meridian is myopic, other hypermetropic → Mixed astigmatism . Since plus cylinder is @180°, steepest curve is vertical → With-the-rule -2D Sph & -1D Cyl @90° (iv) : Both meridians are myopic → Compound myopic astigmatism -1D Sph & +2D Cyl @90° (v) : One meridian is myopic, other hypermetropic → Mixed astigmatism , and since plus cylinder is @90°, steepest curve is horizontal → Against-the-rule Classification of Astigmatism Astigmatism Type Definition Example Simple Myopic One meridian is emmetropic; the other is myopic Plano / -2.00D Cyl Simple Hypermetropic One meridian is emmetropic; the other is hypermetropic Plano / +2.00D Cyl Compound Myopic Both meridians are myopic -2.00 Sph / -1.00 Cyl Compound Hypermetropic Both meridians are hypermetropic +1.00 Sph / +2.00 Cyl Mixed One meridian is myopic, other is hypermetropic -1.00 Sph / +2.00 Cyl With-the-rule Steeper meridian is vertical (plus Cyl @180°) Cyl @180° Against-the-rule Steeper meridian is horizontal (plus Cyl @90°) Cyl @90°

References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 31

Astigmatism

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: All of the following are true regarding the procedure shown below except (Q853840)

A. It's an objective method for refraction testing

B. Red reflex moving with the movement of this instrument implies Hypermetropia

C. Results are unreliable if the patient’s accommodation is active


D. Cycloplegics are only indicated for children below 5 years of age



Correct Answer: D. Cycloplegics are only indicated for children below 5 years of age

- *Listen (English)*
- *Listen (Hindi)*

Explanation: The image shows the retinoscopy (Prisky-Smith plane mirror retinoscope) procedure. Cycloplegics are not limited to children <5 years. Used whenever accommodation is active. For 5–7 years: atropine preferred Older children/adolescents: homatropine or cyclopentolate Can be used even till 20–25 years if needed Retinoscopy (skiascopy o r shadow test): Objective method to assess refractive errors (Option A) Procedure: The patient sits 1 meter from the examiner and light is reflected onto the eye. The examiner observes the red reflex and notes its movement in horizontal and vertical meridians. Observations With movement: Emmetropia or hypermetropia (Option B) Against movement: Myopia >1D No movement: Myopia 1D Difficulties in Retinoscopy: Poor red reflex (small pupil, hazy media, high ametropia) Active accommodation (eliminated by cycloplegia) (Option C) Scissor or conflicting shadows



References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no:42

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 50-year-old male presented to the ophthalmology clinic with complaints of difficulty in reading newspapers and books. Which among the following statements is inappropriate regarding the management of this patient? (Q853839)

A. Roughly +1D convex lenses can be prescribed

B. Coexisting refractive error (myopia/hypermetropia) should be corrected first

C. 1.25% Pilocarpine can be beneficial in treatment

D. Conductive keratoplasty can be performed

Correct Answer: A. Roughly +1D convex lenses can be prescribed

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Given clinical scenario is suggestive of Presbyopia . At ~50 years, the average requirement is ≈ +2 D. +1 D is for ~40 years. Approximate Convex glasses power by age (based on Donders table) 40 yrs → +1 D 45 yrs → +1.5 D 50 yrs → +2 D 55 yrs → +2.5 D 60 yrs → +3 D Correct refractive error first, then add near correction (Option B) 1.25% Pilocarpine (Option C) : FDA-approved presbyopia therapy (induces miosis, increases depth of focus). Conductive keratoplasty (Option D) : though limited, used as a monocular procedure for mild presbyopia.

References:

Yanoff and Duker's Textbook of Ophthalmology, 6th Edition, Pages 33-34, 48-49, 118-119

pmc.ncbi.nlm.nih.gov/articles/PMC1705655

www.ncbi.nlm.nih.gov/books/NBK423827/

journals.lww.com/ijo/fulltext/1984/32020/amplitude_of_accommodation_in_different_age_groups.7.aspx

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 55-year-old patient presents with difficulty reading up close and notices that their vision becomes blurry when trying to focus on objects at near distances. On examination, the near point is found to be farther away than normal. Which of the following best explains the patient's condition? A) The patient has a reduced amplitude of accommodation B) The patient's far point is at infinity C) The patient has a myopic refractive error D) The patient’s near point has moved closer (Q853838)

A. The patient has a reduced amplitude of accommodation B) The patient's far point is at infinity C) The patient has a myopic refractive error D) The patient’s near point has moved closer

B. The patient's far point is at infinity

C. The patient has a myopic refractive error

D. The patient’s near point has moved closer

Correct Answer: A. The patient has a reduced amplitude of accommodation B) The patient's far point is at infinity C) The patient has a myopic refractive error D) The patient’s near point has moved closer

Explanation: The amplitude of accommodation refers to the ability of the eye to focus on objects at different distances, specifically the difference between the near point and far point. As individuals age, the amplitude of accommodation naturally decreases, resulting in difficulty focusing on nearby objects (presbyopia). This is likely the cause of the patient's difficulty reading up close. Changes related to accommodation: Far Point (Punctum Remotum) Emmetropic eye - the far point is at infinity (Option B) Hypermetropic eye - behind the eye Myopic eye

- in front of the eye Near Point (Punctum Proximum) Increases with age (Option D) Range of Accommodation The distance between the far point and the near point Amplitude of Accommodation The difference in dioptric power needed to focus between the near point (P) and the far point (R). Expressed as A = P - R The amplitude of accommodation decreases with age (Option A) Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 26-27, 307.e3-e4, e8

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 26-27, 307.e3-e4, e8

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: All of the following are changes not related to accommodation except (Q853837)

A. Anterior lens curvature almost remains the same

B. Posterior lens curvature is increased

C. Increased zonular tension

D. Ciliary muscle contraction

Correct Answer: D. Ciliary muscle contraction

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Accommodation: The mechanism by which the eye increases the power of the crystalline lens to focus on near objects, allowing diverging rays to be focused on the retina. Changes occurring: The curvature of the anterior surface of the lens increases (Option A) Posterior surface curvature almost remains the same (Option B) Contraction of the ciliary muscle Relaxation of zonules (Option C) secondary to ciliary muscle contraction (Helmholtz's capsular theory)

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 26-27, 307.e3-e4, e8

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A patient post cataract lens extraction refused IOL placement and hence was prescribed spectacles. Later, he presented with complaints of visual discomfort.

Choose the right combination of effects the patient would have likely felt. 1) Pincushion effect of distortion 2) Barrel effect of distortion 3) There is magnification of the image 4) There is a minification of the image (Q853836)

A. 1,3

B. 1,4

C. 2,3


D. 2,4


Correct Answer: A. 1,3

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Post-lens extraction, the patient will have hypermetropia . Convex spectacles are prescribed for the correction of hypermetropia, which could cause the Pincushion effect . Pincushion distortion (Statement-1) Barrel distortion (Statement-2) There is magnification of the image. (Statement-3) There is a minification of the image. (Statement-4) Seen with convex lenses . Seen with concave lenses .





References:

Myron Yanoff and Duker's Textbook of Ophthalmology, 6th Edition, Page 45, 66.e3

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 45-year-old patient presents with blurry vision and difficulty focusing after undergoing cataract surgery 3 weeks ago. On examination, the patient has no lens in the eye and no evidence of intraocular lens (IOL) implantation. Which of the following is the most appropriate management for this patient? (Q853835)

A. Prescribe corrective glasses with a high refractive index

B. Fit the patient with an intraocular lens

C. Perform a keratoplasty to correct vision

D. Initiate topical corticosteroids for inflammation

Correct Answer: B. Fit the patient with an intraocular lens

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In this scenario, the patient is presenting with aphakia , which is the absence of a lens in the eye, a condition that can occur after cataract surgery if an intraocular lens (IOL) was not implanted or if it was removed. IOL implantation is the recommended corrective procedure. Prescribe corrective glasses with a high refractive index (Option A): Incorrect as definitive treatment in a 45-year-old. Glasses are bulky, give limited field of vision, and are usually reserved for elderly or unfit patients. Perform a keratoplasty to correct vision (Option C): Incorrect because the cornea is not the issue; aphakia is a refractive error, not a corneal pathology. Initiate topical corticosteroids for inflammation (Option D): Incorrect unless signs of postoperative inflammation are present, which are not described in this case.

References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 318, 370-372 Kanski’s clinical ophthalmology, John F. Salmon, 10th edition, Page no: 340 Secondary congenital aphakia - PMC

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: In the case of astigmatism, the vertical meridian is more curved than the horizontal. What kind of lens should be given? (Q853834)

A. Concave cylindrical lens at 180°

B. Convex cylindrical lens at 180

C. Concave cylindrical lens at 90

D. None of the above

Correct Answer: A. Concave cylindrical lens at 180°

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: This is a case of with-the-rule astigmatism, in which a concave cylindrical lens at 180° or a convex cylindrical lens at 90° is used for correction With-the-Rule Astigmatism (WTR) Against-the-Rule Astigmatism (ATR) The vertical meridian is more curved than the horizontal. Correction requires a concave cylindrical lens at 180° or a convex cylindrical lens at 90°. It is called "with-the-rule" because the vertical meridian is normally more convex than the horizontal due to eyelid pressure. The horizontal meridian is more curved than the vertical meridian. Correction requires a convex cylindrical lens at 180° or a concave cylindrical lens at 90°.

References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 39

Astigmatism

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: A few rays of light are focused on the retina in one meridian and a few others are focused in front of the retina in another meridian. What is the diagnosis? (Q853833)

A. Simple myopic astigmatism

B. Simple hypermetropic astigmatism

C. Mixed astigmatism


D. Compound myopic astigmatism

Correct Answer: A. Simple myopic astigmatism

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In this case, one meridian is on retina, other meridian is in front (myopic), suggestive of s imple myopic astigmatism Astigmatism is classified based on where the two principal meridians focus: Simple Myopic Astigmatism One meridian → on retina (0) Other meridian → in front of retina (–) Simple Hypermetropic Astigmatism (Option B) One meridian → on retina (0) Other meridian → behind retina (+) Compound Myopic Astigmatism (Option D) Both meridians → in front of retina (–, –) Compound Hypermetropic Astigmatism Both meridians → behind retina (+, +) Mixed Astigmatism (Option C) One meridian → in front (–) Other meridian → behind (+)



Types of Astigmatism: A) Simple myopic B) Simple hypermetropic
C) Compound myopic D) Compound hypermetropic E) Mixed

@Neet_pg_bot

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 31

Astigmatism

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: A 35-year-old patient presents with difficulty focusing on both near and distant objects. Examination reveals high hypermetropia. What is the expected fundoscopy finding (Q853832)

A. Smaller optic disc

B. Shotsilk retina

C. Pseudopapillitis


D. All of the above

Correct Answer: D. All of the above

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: All are findings in a hypermetropic eye. Hypermetropia, or long-sightedness, is a refractive error in which parallel rays of light from distant objects focus behind the retina when the eye's accommodation is relaxed. Features: Small eyeball in high hypermetropia Smaller cornea Shallow anterior chamber Fundus Examination: Optic disc - Small (Option A) , ill-defined margins, possibly simulating papillitis (pseudopapillitis) (Option C) . Retina - Shot silk appearance (shining retina) (Option B) .



References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 28-31

www.ncbi.nlm.nih.gov/books/NBK560716/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 65-year-old patient presents with gradually worsening blurry vision over the past year with recent improvement in his near vision . On examination, the lens shows yellowish discolouration with increased opacity, consistent with nuclear sclerosis. What kind of myopia has he developed secondary to this? (Q853804)

A. Axial myopia

B. Curvatural myopia

C. Index myopia

D. Positional myopia

Correct Answer: C. Index myopia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In nuclear cataracts , the lens nucleus becomes sclerotic and denser . This leads to an increase in the refractive index of the lens nucleus. As a result, the eye becomes more myopic. This is known as index myopia . AKA second sight of old age, as there will be improvement in near vision. Axial Myopia (Option A): Increase in the anteroposterior length of the eyeball (most common form). Curvatural Myopia (Option B): Increased curvature of the cornea, lens, or both. Positional Myopia (Option D): Anterior placement of the crystalline lens in the eye.

References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 28-30

Defining and Classifying Myopia

↑ *Back to this question in Part A*

@Neet_pg_bot


Q13: Identify X: (Q853803)

A. Angle alpha

B. Angle beta

C. Angle gamma

D. Angle kappa




@Neet_pg_bot


Correct Answer: D. Angle kappa

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Angle kappa is the angle between the visual axis and the pupillary axis of the eye. The visual axis passes from the fixated object through the nodal point of the eye to the fovea . The pupillary axis is a line perpendicular to the cornea passing through the center of the pupil . The angle formed between these two lines (marked X in the diagram) is called angle kappa . Normally, this angle is slightly positive , meaning the visual axis lies nasal to the pupillary axis — causing a slight temporal displacement of the corneal light reflex. Comparison Angle Definition Key Relation Angle alpha Between optical axis and visual axis At the nodal point Angle beta Between optical axis and line of fixation At the center of rotation Angle gamma Between optical axis and visual line From center of rotation Angle kappa Between visual axis and pupillary axis Clinically most relevant



@Neet_pg_bot



@Neet_pg_bot

References:

Myron Yanoff and Duker's Textbook of Ophthalmology, 6th Edition, Page 20, 1125

www.ncbi.nlm.nih.gov/books/NBK554036/figure/ch14.Fig11/

pmc.ncbi.nlm.nih.gov/articles/PMC3872563/#:~:text=Angle%20kappa%20is%20defined%20as,of%20curvature%20of%20the%20cornea.

ijdmsrjournal.com/issue_dcp/Understanding%20the%20importance%20of%20axes%20and%20angles%20in%20modern%20era%20of%20Ophthalmology%20by%20the%20case%20series%20of%20pseudo%20squint.%20The%20case%20series%20of%20pseudo%20squint.pdf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: Snellen chart is used to assess which of the following? (Q853787)

A. Colour vision

B. Contrast sensitivity

C. Peripheral vision


D. Visual acuity

Correct Answer: D. Visual acuity

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Snellen’s chart is used to measure visual acuity by assessing the eye's ability to distinguish details or separate points. Standard test distance → 6 meters (20 feet in feet-based charts). Fraction (e.g., 6/18, 20/40): Numerator → distance at which the patient reads the line. Denominator → distance at which a normal eye should read that line. Example: 6/18 → A normal person can read the line at 18 m, but the patient reads it only at 6 m. Whole letter subtends an angle of 5 minutes of arc at the nodal point. Stroke/margin of the letter subtends 1 minute of arc. Color vision (Option A): tested by the Ishihara chart Contrast sensitivity (Option B): tested by Pelli–Robson chart Peripheral vision (Option C): assessed by perimetry/visual field testing



References:

Kanski's Clinical Ophthalmology, John F. Salmon, 10th Edition, Page 2-3 Myron Yanoff and Duker's Textbook of Ophthalmology, 6th Edition, Page 35

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 65-year-old patient with a mature cataract presents for an eye examination. During the assessment, the ophthalmologist observes the absence of a specific Purkinje image. Which Purkinje image is most likely to be absent in this case? (Q853801)

A. First Purkinje image

B. Second Purkinje image

C. Third Purkinje image


D. Fourth Purkinje image

Correct Answer: D. Fourth Purkinje image

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In a mature cataract, the posterior lens surface is obscured; therefore, the 4th image is absent. Purkinje images : Reflections of light from ocular surfaces: First (Option A): Anterior corneal surface Brightest and largest image Second (Option B): Posterior corneal surface Third (Option C): Anterior lens surface Absent in Aphakia Fourth: Posterior lens surface Absent in Aphakia Only inverted image



References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 60

Purkinje images

Mature cataract -4th image absent

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: Which of the following is the most sensitive test for both congenital and acquired colour vision defects? (Q853800)

A. Ishihara Test

B. Farnsworth-Munsell 100 Hue Test

C. Hardy-Rand-Rittler (HRR) Test

D. Holmgren Wool Test


Correct Answer: B. Farnsworth-Munsell 100 Hue Test


■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Farnsworth-Munsell 100 Hue test is the most sensitive test for both congenital and acquired color vision defects. It consists of 85 hue caps and color vision is judged by error score. It evaluates a person's ability to discern small differences in hues. Ishihara Test (Option A): Primarily used to detect red-green color blindness. It is not very effective for diagnosing blue-yellow defects or for detecting acquired color vision defects. Hardy-Rand-Rittler (HRR) Test (Option C): This test can detect both red-green and blue-yellow color vision defects and is useful for both congenital and acquired conditions, but it is less sensitive than the Farnsworth-Munsell 100 Hue Test. Holmgren Wool Test (Option D): Older test using colored wool yarns to assess color vision. It is less sensitive than modern tests and is rarely used for clinical evaluation today.




@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 7-8

www.ncbi.nlm.nih.gov/books/NBK217826/#:~:text=The%20most%20effective%20test%20for,(Table%204%2D1).

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A 40-year-old woman presents with a 3-day history of painful vision loss in her right eye. She has a history of multiple sclerosis. On examination, her visual acuity is 20/30 in the right eye, and a RAPD is noted. She reports difficulty distinguishing red and green colors in the right eye, but her left eye is normal on color vision testing. Which of the following is the most likely cause of her acquired color blindness? (Q853799)

A. Retinal vein occlusion

B. Optic neuritis

C. Diabetic retinopathy

D. Macular degeneration

Correct Answer: B. Optic neuritis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Optic neuritis , often associated with multiple sclerosis , commonly affects the optic nerve, leading to disturbances in vision, including red-green color vision defect. Kollner's Rule (for acquired color defects) Optic nerve lesions → Red-Green defect (except Glaucoma → Blue-Yellow) Retinal lesions → Blue-Yellow defect (except Stargardt's disease → Red-Green) Retinal vein occlusion (Option A) typically causes sudden vision loss, often associated with retinal hemorrhages, but does not cause a specific color vision defect in isolation. Diabetic retinopathy (Option C) causes gradual vision loss due to retinal changes (like microaneurysms, hemorrhages, and edema) but does not typically present with acute color vision defects like those seen in optic neuritis. Macular degeneration (Option D) primarily affects central vision and may cause difficulty with reading or recognizing faces but does not typically cause red-green color vision defects.

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 625, 811, 834, 842, 934, 942

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 764, 771, 902

Acquired Color Vision Defects

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: Which of the following statements regarding congenital colour blindness is TRUE? (Q853798)

A. It is more common in females than males.

B. It is an autosomal recessive condition.

C. It primarily affects the perception of red and green colours and is the most common type.

D. It always results in total loss of colour vision.

Correct Answer: C. It primarily affects the perception of red and green colours and is the most common type.

Explanation: Group of inherited conditions that affect an individual's ability to perceive colours correctly. It typically occurs due to defects in the photoreceptor cells (cones) of the retina that are responsible for colour vision. This condition is usually present from birth and remains constant throughout life. Red-Green Color Blindness: The

most common type (Option C). Typically inherited through an X-linked recessive pattern, which is why it is more common in males (Option A,B). Congenital colour blindness usually involves partial colour vision deficiency, not total colour blindness (Option D). Congenital colour blindness is classified into: Dyschromatopsia : Anomalous trichromacy Dichromacy Blue cone monochromatism Achromatopsia/Rod monochromatism

References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 645

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: Which type of colour blindness is caused by the absence of green cones in the retina? (Q853797)

A. Protanopia

B. Deuteranopia

C. Tritanopia

D. Achromatopsia

Correct Answer: B. Deuteranopia

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Deuteranopia is a type of red-green color blindness where green cone cells are absent or nonfunctional, leading to difficulty distinguishing between red and green hues. It is the most common type of color blindness Protanopia (Option A) : Complete red color defect . Tritanopia (Option C) : Absence of blue color appreciation . Achromatopsia (Option D): Total color blindness with day blindness.

References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 6, 645

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 454, 859,

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: At what age do infants typically begin to perceive and distinguish between different colors? (Q853796)

A. At birth

B. 1-2 months

C. 3-4 months

D. 6-9 months

Correct Answer: C. 3-4 months

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: While newborns see mostly in shades of grey, their color perception begins to develop around 3-4 months of age, when they can start distinguishing between primary colors like red and blue. Fovea development begins at 6 weeks. Foveal reflex formation: 5-6 months Color perception begins at 12 weeks (3 months).

References:

www.ncbi.nlm.nih.gov/books/NBK11538/

www.ncbi.nlm.nih.gov/pmc/articles/PMC9314692/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: A 55-year-old male presents with complaints of difficulty driving at night and reading in dim light, despite having normal visual acuity during his eye exam. He underwent cataract surgery six months ago but still experiences these visual issues. Which of the following tests is most appropriate to assess his current visual problem? (Q853795)

A. Snellen Chart

B. Ishihara Test

C. Letter Contrast Sensitivity Test


D. Visual Field Test


Correct Answer: C. Letter Contrast Sensitivity Test

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient's difficulty with night driving and reading in dim light, despite normal visual acuity, suggests a problem with contrast sensitivity. The Letter Contrast Sensitivity Test is specifically designed to evaluate how well the patient can distinguish between varying contrasts, which is often impacted in conditions like post-cataract surgery or early-stage optic nerve diseases. Pelli-Robson chart is the standard test. Contrast Sensitivity Grating Tests Snellen Chart (Option A) measures visual acuity but does not assess the ability to see in low-contrast situations. The Ishihara Test (Option B) evaluates color vision, not contrast sensitivity. The Visual Field Test (Option D) checks for peripheral vision defects, which are unrelated to contrast sensitivity.





(A, B and C) Sinusoidal grating patterns; D) Sqaure grating pattern. (A) and (B) differ in spatial frequency, (B) and (C) in contrast, defined as (Lmax-Lmin) / (Lmax+Lmin) where L is the light intensity recorded by a photocell scanning across the gratings.

@Neet_pg_bot

References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 4

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 35-37

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: A patient's visual acuity improves significantly when looking through a small, opaque disc with a tiny hole, as shown in the image below. What is the likely cause of this improvement in visual acuity? (Q853794)

A. Cataracts

B. Refractive error

C. Optic nerve damage

D. Macular degeneration



Correct Answer: B. Refractive error

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The pinhole test reduces the amount of light scattering in the eye and sharpens vision by narrowing the beam of light entering the pupil. An improvement in vision suggests a refractive error , such as myopia or astigmatism, that can be corrected with lenses. Standard pinhole size: 1mm to 1.25mm Cataracts (Option A): They cloud the lens, but using a pinhole does not typically improve vision, as the issue is related to lens opacification. Optic nerve damage (Option C): This condition, such as optic neuropathy, does not improve with the pinhole, as the issue is with the nerve, not light entering the eye. Macular degeneration (Option D): The visual acuity is unlikely to improve with a pinhole and may even decrease since the underlying problem is retinal damage, not a refractive error.

References:

Kanski’s Clinical Ophthalmology, 10th Edition, Page 2

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9249474/

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3057328/

https://www.ncbi.nlm.nih.gov/books/NBK594235/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Match the following visual acuity charts used to assess children of preschool age. 1) Allen test cards a) 2) HOTV test b) 3) Stycar Test c) 4) Cardiff Acuity test d) (Q853790)


A. 1-a; 2-b; 3-c; 4-d

B. 1-a; 2-b; 4-c; 3-d

C. 4-a; 2-b; 3-c; 1-d

D. 1-a; 3-b; 2-c; 4-d






@Neet_pg_bot



@Neet_pg_bot




Correct Answer: A. 1-a; 2-b; 3-c; 4-d

- *Listen (English)*
- *Listen (Hindi)*

Explanation: Visual acuity assessment- preschool 1) Allen test cards Designed for children aged 2 to 4 years. 4 flashcards are used to test containing 7 schematic figures. a) 2) HOTV test Designed for children older than 4 years. Test involves identifying the letters “H,” “O,” “T,” and “V.” The letters decrease in size from the top to the bottom of the chart. b) 3) Stycar Test (Screening Test for Young Children and Retarded) c) 4) Cardiff Acuity card test Used in preschoolers; but in < 1 year it works on preferential looking principle. d)




@Neet_pg_bot



@Neet_pg_bot





References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 1120-1122

https://www.ncbi.nlm.nih.gov/books/NBK52711/table/ch1.t3/

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1311632/

↑ *Back to this question in Part A*

@Neet_pg_bot


Q24: Which among the following is used to check visual acuity in children less than 1 year? 1) 2) 3) 4) (Q853789)

A. 1 and 4


B. 2 and 3

C. All except 4


D. All of the above




@Neet_pg_bot



@Neet_pg_bot




@Neet_pg_bot




Correct Answer: D. All of the above

- *Listen (English)*
- *Listen (Hindi)*

Explanation: All of the following tests are used to check visual acuity in children less than 1 year old. 1) Cardiff Acuity Test Used in preschoolers; but in < 1 year it works on preferential looking principle. 2) Teller's Acuity Cards (grating test) : Preferential looking test; child fixates on striped vs plain card. 3) Optokinetic Drum: Assesses presence of optokinetic nystagmus → proves intact visual pathway. 4) Visual Evoked Potential (VEP) : Electrophysiological test; measures cortical response to visual stimuli, especially useful in newborns.




@Neet_pg_bot



@Neet_pg_bot





References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 1120-1122

https://www.ncbi.nlm.nih.gov/books/NBK564307/

↑ *Back to this question in Part A*

@Neet_pg_bot


Q25: Which among the following is used to check visual acuity in illiterate adults? 1) 2) 3) 4) (Q853788)

A. 1 and 3


B. 2 and 4


C. All except 1

D. None of the above




D-36
D-24
D-18
D-12
D-9
D-6



Meters (Feet)
Log VAR
H V Z D S
N C V K D
C Z S H N
O N V S R
K D N R O
Z K C S V
D V O H C
O H V C K
H Z C K O
N C K H D
Z H C S R
S Z R D N
H C D R O
R D O S N
@Neet_pg_bot




Correct Answer: B. 2 and 4


- *Listen (English)*
- *Listen (Hindi)*

Explanation: 1) Snellen’s chart Measures visual acuity by assessing the eye's ability to distinguish details or separate points. 2) Tumbling "E" Visual Acuity Chart Involves matching the orientation of the letter "E" in different sizes. Used to check visual acuity in illiterates. 3) LOGMAR chart Modified chart with equal number of letters per line; tests crowding phenomenon and is more accurate (tested at 4m) 4) Landolt's broken ring test Patient identifies the gap in the ring (up, down, left, right)


E
1 20/200
F P
2 20/100
T O Z
3 20/70
L P E D
4 20/50
P E C F D
5 20/40
E D F C Z P
6 20/30
F E L O P Z D
7 20/25
D E F P O T E C
8 20/20
L E F O D P C T
9
F D P L T C E O
10
F E Z O L C F T D
11
@Neet_pg_bot


E
D·36
D·24
D·18
D·12
D·9
D·6





References:

Kanski’s Clinical Ophthalmology, John F. Salmon, 10th Edition, Page 2,3

Myron Yanoff and Duker’s Textbook of Ophthalmology, 6th Edition, Page 1122

https://www.ncbi.nlm.nih.gov/books/NBK564307/

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1298236/

https://pubmed.ncbi.nlm.nih.gov/302151/

↑ *Back to this question in Part A*

@Neet_pg_bot

• [11_Optics] 025 Previous Year Questions

« Back to Folder Directory

[11_Optics] 025 Previous Year Questions — Quick Jump

Q1 (p. 1410)	*Q2 (p. 1411)*	*Q3 (p. 1414)*	*Q4 (p. 1414)*	*Q5 (p. 1417)*	*Q6 (p. 1419)*
Q7 (p. 1420)	*Q8 (p. 1422)*	*Q9 (p. 1424)*	*Q10 (p. 1425)*	*Q11 (p. 1425)*	*Q12 (p. 1427)*
Q13 (p. 1429)	*Q14 (p. 1430)*	*Q15 (p. 1431)*	*Q16 (p. 1433)*	*Q17 (p. 1434)*	*Q18 (p. 1434)*
Q19 (p. 1435)	*Q20 (p. 1437)*	*Q21 (p. 1437)*	*Q22 (p. 1438)*	*Q23 (p. 1439)*	*Q24 (p. 1440)*
Q25 (p. 1441)	*Q26 (p. 1442)*				

Q1: A patient was prescribed with -1D spherical and -2D cylindrical at 180 degrees. What type of refractive error does this patient have? (Q857422)

A. With the rule astigmatism

B. Against the rule astigmatism

C. Simple astigmatism

D. Mixed astigmatism

Correct Answer: A. With the rule astigmatism

Explanation: The prescription given is -1D spherical and -2D cylindrical at 180 degrees . The spherical component (-1D) indicates a myopic refractive error. The cylindrical component (-2D) at 180 degrees indicates astigmatism with the steepest curvature along the 180-degree axis. In with-the-rule astigmatism , the steepest curvature (and therefore the highest cylinder value) is along the vertical axis (90 degrees), and the flatter axis is horizontal (0 or 180 degrees). Since the cylinder is placed at 180 degrees and the patient has a more negative cylindrical value at this axis, it indicates with-the-rule astigmatism . With-the-Rule Astigmatism (WTR) Against-the-Rule Astigmatism (ATR) The vertical meridian is more curved than the horizontal. Correction requires a concave cylindrical lens at 180° ± 20° or a convex cylindrical lens at 90° ± 20°. It is called "with-the-rule" because the vertical meridian is normally more convex than the horizontal due to eyelid pressure. The horizontal meridian is more curved than the vertical meridian. Correction requires a convex cylindrical lens at 180° ± 20° or a concave cylindrical lens at 90° ± 20°. 180 degrees 90 Degrees With the rule [-] [+] Against the rule [+] [-] Against the rule astigmatism (Option B) : This occurs when the steepest curvature is along the horizontal axis (0 or 180 degrees) and the flatter axis is vertical (90 degrees), which is not the case here. Simple astigmatism (Option C) : This refers to astigmatism where one of the meridians has no refractive error (i.e., it's either emmetropic or has no cylinder correction), but this patient has both spherical and cylindrical components. Mixed astigmatism (Option D) : This occurs when one axis has a myopic refractive error and the other has a hypermetropic refractive error. In this case, the prescription is purely myopic in both axes.

References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 39

Astigmatism

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Given below are some tests for visual acuity. Which of these is used for a pre-verbal child? ETDRS chart Optokinetic nystagmus Teller’s visual acuity chart Landolt’s C chart (Q878998)

A. 2,3

B. 1,2

C. 2,3,4

D. 2,4

Correct Answer: A. 2,3


Explanation: The visual-acuity tests which are suitable for a pre-verbal child aka a child who cannot speak or identify letters/symbols must rely on reflexive or preferential-looking responses. Optokinetic nystagmus (OKN): Used for infants and pre-verbal children. Involves moving black-and-white stripes (OKN drum). Presence of optokinetic reflexive eye movements indicates the child can detect the stripes → visual acuity estimation. Teller’s visual acuity chart: Specifically designed for non-verbal infants and toddlers. Based on preferential looking, the child looks toward the panel with stripes instead of a blank panel. Gives a quantitative estimate of visual acuity without requiring cooperation or verbal recognition. ETDRS chart (Option 1): Modified Snellen chart for research-grade visual acuity. Contains letters , requires recognition and verbal response . Used in older children and adults , not useful for pre-verbal kids . ■■■■■■■ ■■■■■■■ Landolt’s C chart (Option 4): Uses Landolt C rings with a gap oriented in different directions. The patient must identify where the gap is → requires cognitive ability and speech/pointing cooperation. Used in literate children and adults , not pre-verbal infants .




@Neet_pg_bot



@Neet_pg_bot





References:

Parson’s Diseases of the Eye, 23rd Edition, Page 239, 240, 241

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: What would be the suitable course of action for managing a 15-year-old girl who is non-adherent to wearing spectacles for her myopic astigmatism? (Q856123)

A. LASIK

B. Spherical equivalent spectacles

C. Femto LASIK

D. Implantable collamer lens

Correct Answer: B. Spherical equivalent spectacles

Explanation: ollamer Lens) are contraindicated because the refractive error is not yet stable . Key principle: Refractive surgeries are generally considered only after 18–21 years of age , when the refractive power has remained stable for at least 1 year . For adolescents, spectacles or contact lenses remain the only safe and effective means of correction. In a non-adherent patient, contact lenses (especially toric lenses for astigmatism) can be considered, but surgery should be deferred until ocular maturity. Management options for Astigmatism: Optical Spectacles: with cylindrical power correction/ spherical equivalent spectacles is the initial treatment option especially for paediatric age group (Option B) Rigid contact lenses: for up to 3D Toric contact lenses: for > 3D astigmatism cases Refractive Procedures Phototherapeutic keratectomy: Using excimer lens, for irregular astigmatism, can correct up to 3D LASIK (Laser-Assisted In Situ Keratomileusis): can correct up to 5D IOL implantation: Toric IOLs are preferred using Collamer lenses Surgical Penetrating or Deep Lamellar Anterior Keratoplasty (DLAK): in cases of corneal scarring/ no improvement with other treatment modalities Incorrect Options: A) LASIK: Contraindicated below 18 years; risk of regression and corneal ectasia. C) Femto LASIK: Similar to LASIK; not indicated before adulthood. D) Implantable collamer lens (ICL): Used for high refractive errors when LASIK is unsuitable, but only in adults with stable refraction.

References:

AK Khurana Textbook of Ophthalmology 7th Edition, Page 44,216.

Kanski Clinical Ophthalmology, 10th edition, Pages 284,285.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: What is the statement correct about the instrument given below? (Q857755)

A. Condensing lens is not required

B. It is used as close to the patient’s eye as possible. A real, inverted image is seen, Magnification is about 15 times

C. A real, inverted image is seen

D. Magnification is about 15 times



Correct Answer: D. Magnification is about 15 times

Explanation: Instrument shown in the image is that of a Direct ophthalmoscope , where a virtual, erect image is produced with magnification of about 15 times . Direct and Indirect Ophthalmoscope: Direct Ophthalmoscope Indirect Ophthalmoscope Image Virtual image, erect (Option B & C) Real image, inverted Magnification 15x magnification (Option D) 2-3x magnification Condensing lens Used (Option A) Not required Visualization Uniocular Binocular Field of view 2 DD 8 DD Advantages Undilated pupil Stereopsis present Entire retina can be seen Visualization through hazy media is possible {{caption_text}} {{caption_text}}





References:

Kanski's Clinical Ophthalmology 9th Edition, Page 25

Yanoff and Duker Ophthalmology 5th Edition, Page 56, 57

Opthalmoscope

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: Which of the following lenses would be recommended to correct myopia? (Q857245)

A. Convex


B. Concave


C. Cylindrical

D. Abraham lens

Correct Answer: B. Concave

Explanation: Concave lenses (diverging lenses) are used to correct Myopia. Myopia - management: General measures Balanced diet Visual hygiene: Posture correction, adequate illumination, reducing usage of video display units (VDUs) Optical Concave lenses are prescribed: Spectacles/ contact lenses (Option B) Surgical Radial Keratotomy (RK): Used for low to moderate myopia (2-6 D). No longer recommended due to the advent of safer procedures like LASIK. Photorefractive Keratectomy (PRK): It uses an excimer laser to reshape the cornea by removing a thin corneal layer. Suitable for correcting myopia up to -6 D. Refractive Lenticule Extraction (SMILE): A minimally invasive laser procedure where a small lenticule is removed to reshape the cornea. Effective for myopia up to -10 D, with or without astigmatism. Phakic Intraocular Lens (IOL) Implantation: Implantable lens placed without removing the natural lens. Best for myopia exceeding -8 D, especially in patients unsuitable for laser surgery. Convex lens (Option A): Used in the case of presbyopia and in the case of hypermetropia . Cylindrical lens (Option C): Used for correction of astigmatism . Abraham lens (Option D): A 66 D lens used for peripheral iridotomy with Nd:Yag laser . {{caption_text}}


Focus in front of retina
Retina
Myopia eye
Light ray from far object
Focus exactly on retina
Light ray from far object
concave lens
Corrected myopia eye
@Neet_pg_bot


Hypermetropia eye
Correction (Convex lens)
for Hypermetropic eye
Focusing Behind
the Retina
Convex Lens
@Neet_pg_bot



↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: In a 3-year-old child who has recently developed 15-degree accommodative esotropia, what is the most effective treatment approach? (Q857242)

A. Orthoptic exercise

B. Surgery


C. Refractive correction

D. INTACS

Correct Answer: C. Refractive correction

Explanation: Refraction correction would be the most effective treatment approach in a child with Accommodative esotropia . Accommodative esotropia: Accommodative Refractive Esotropia Accommodative Non-refractive Esotropia Accommodative Mixed Esotropia Cause High hypermetropia (+4 to +7 D) Abnormal large AC/A (accommodative convergence/accommodation) ratio Combination of hypermetropia and high AC/ A ratio Treatment Correction of hypermetropia (spectacles) {{caption_text}} Corrected by bifocal glasses with +3 DS added for near vision Miotics (Echothiophate) Surgery: Bilateral medial rectus recession Surgery: Bilateral medial rectus recession Orthoptic exercise (Option A): Orthoptic exercises are used for the improvement in the binocular vision to treat paralytic squint . Surgery (Option B): Surgeries are generally performed to treat non-refractive/

non-accommodative esotropia . INTACS (Option D): INTACS (intracorneal ring segments) are used for the management of keratoconus.



References:

Parsons Diseases of the eye 23rd Edition, pages 979- 982.

Yanoff & Duker Ophthalmology 5th edition, pages 1212,1213.

A K KhuranaComprehensive Ophthalmology 7th edition, page 363.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: The probable cause for a patient with a corneal scar having a visual acuity of 6/36, which improves to 6/18 with a pinhole, is most likely due to... (Q857209)

A. Glaucoma

B. Malingering


C. Irregular astigmatism

D. Cataract

Correct Answer: C. Irregular astigmatism

Explanation: Improvement in visual acuity with a pinhole is suggestive of a refractive error . Among the given options, Irregular astigmatism is the only refractive error. The pinhole test reduces the amount of light scattering in the eye and sharpens vision by narrowing the beam of light entering the pupil. Pinhole and Stenopeic slit tests: Pinhole Test Stenopeic Slit Test Mechanism Uses a small opaque disc with holes to reduce retinal blur and improve visual acuity in refractive errors up to 4D. {{caption_text}} Uses a narrow slit to reduce peripheral aberrations and enhance depth of focus. Clinical Applications Determines if reduced vision is due to refractive errors; vision improvement suggests a refractive issue. (Option C) Quick assessment of visual acuity, often in under-resourced settings. WHO recommended blindness surveys. To assess astigmatism, especially irregular astigmatism (as it allows for a more detailed analysis of the patient's visual field by focusing light along a single axis) or complex visual issues. A stenopeic slit is used in cases of corneal opacities to help identify the best location for an optical iridectomy. Limitations Can underestimate post-refractive visual acuity. Reduces light reaching the retina, potentially causing diffraction blur. Limited mainly to complex refractive issues.





References:

Kanski’s clinical ophthalmology 10th edition, page 2.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9249474/

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3057328/

https://www.ncbi.nlm.nih.gov/books/NBK594235/#:~:text=The%20stenopeic%20slit%20is%20often,irregularities%2C%20or%20severe%20dry%20eye.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: During a routine eye examination, a patient's visual acuity was evaluated using Snellen's chart, and it was determined to be 6/6. What is the smallest angle at which this patient can differentiate two separate points or objects? (Q855228)

A. 15 minutes of arc


B. 5 minutes of arc


C. 10 minutes of arc

D. 20 minutes of arc

Correct Answer: B. 5 minutes of arc

Explanation: Visual acuity of 6/6 means the patient has normal visual acuity , which is the ability to see clearly at a distance of 6 meters what should normally be seen at that distance. The smallest angle at which the eye can differentiate two points is referred to as the angular resolution of the eye. For a person with normal vision (6/6), this is approximately 1 minute of arc per line on the Snellen chart. However, the smallest angular separation that the human eye can typically distinguish under optimal conditions is 5 minutes of arc . This corresponds to the smallest angle at which a person with normal vision can resolve two separate points as distinct. Normal visual acuity (6/6) corresponds to a subtended angle of 5 minutes of arc. Snellen's chart Measures visual acuity by assessing the eye's ability to distinguish details or separate points. {{caption_text}} Rows of decreasing-sized letters fit within squares. Each letter subtends an angle of 5 minutes, and the lines forming letters subtend 1 minute of arc. Visual Acuity Notation Snellen Fraction: Denoted as a fraction, with the numerator being 6 meters and the denominator indicating the distance at which normal vision reads the same letter (e.g., 6/18 means the patient reads at 6 meters what normal vision reads at 18 meters). Metric/Imperial: 6/6 = 20/20 vision. Decimal Acuity: 6/6 equals 1.0 decimal acuity. Testing Procedure The patient is positioned 6 meters from the chart, and each eye is tested individually, starting with the right eye. The patient reads the smallest visible letter. Limitations Refractive Errors: Primarily measures uncorrected vision. Pinhole tests help detect refractive errors that can be corrected with lenses. Subjectivity: Dependent on patient literacy, language, and cognitive ability. Scope: Assesses only visual acuity, not contrast sensitivity, color vision, or peripheral vision. 15 minutes of arc (Option A) : This is too large an angle for normal visual acuity. It corresponds to a less sharp visual resolution. 10 minutes of arc (Option C) : This is also larger than the typical 5 minutes of arc resolution of normal vision. 20 minutes of arc (Option D) : This is even larger and would correspond to a more impaired level of visual acuity.





Widening effect of the visual angle with distance

@Neet_pg_bot

References:

Kanski's Clinical Ophthalmology, John F. Salmon, 10th Edition, Page 2-3

Myron Yanoff and Duker's Textbook of Ophthalmology, 6th Edition, Page 35

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: When selecting the strength of an intraocular lens (IOL) for a child under 1 year old, what factors should be considered? (Q793030)

A. Myopic shift

B. Hypermetropic shift

C. Length of the lens capsule

D. Growth of the lens capsule

Correct Answer: A. Myopic shift

Explanation: Correct Option A: When choosing the power of an intraocular lens (IOL) in a child under 1 year of age, the myopic shift should be taken into account. A myopic shift is a phenomenon where the eye becomes more myopic as it grows, and this can lead to a child becoming increasingly nearsighted over time. To compensate for this, a lower power IOL should be chosen to allow for the myopic shift that will occur as the child's eye grows. The length of the lens capsule and growth of the lens capsule are important factors to consider when selecting an IOL, but they are not the primary factors in choosing the power of the IOL in a young child. Incorrect Options: Option B - Hypermetropic shift: Hypermetropic shift is not a relevant factor in choosing the power of an IOL in a child under 1 year of age. The hypermetropic shift is the phenomenon where the eye becomes more hyperopic as it grows, and this is not typically observed in young children. Option C - Length of the lens capsule: The length of the lens capsule is an important factor to consider when selecting an IOL, but it is not the primary factor in choosing the power of the IOL in a young child. Option D - Growth of the lens capsule: The growth of the lens capsule is an important factor to consider when selecting an IOL, but it is not the primary factor in choosing the power of the IOL in a young child.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: What is the probable medical condition for a 70-year-old individual who has relied on presbyopia glasses but can now comfortably read the newspaper without them? (Q858148)

A. Index myopia

B. Index hypermetropia

C. Presbyopia

D. None of the above

Correct Answer: A. Index myopia

Explanation: for the patient's improvement in near vision.

References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 28-30

Defining and Classifying Myopia

↑ *Back to this question in Part A*

@Neet_pg_bot


Q11: What is the term used to describe the refractive error seen in the ocular examination image of a 60-year-old patient who has complained of reduced distant vision, but no longer requires spectacles for near vision? (Q855499)

A. Index myopia

B. Axial myopia

C. Curvatural myopia

D. Positional myopia



Correct Answer: A. Index myopia

Explanation: Myopia = shortsightedness- refractive error where parallel rays of light from distant objects focus in front of the retina when the eye's accommodation is relaxed. Index myopia is a condition that often occurs in older individuals, typically associated with age-related changes in the lens . As we age, the lens of the eye becomes denser and less flexible, which can lead to myopia (nearsightedness) in some cases. The patient has reduced distant vision (a hallmark of myopia) but no longer needs glasses for near vision , which suggests that the lens has become more myopic (thicker or more curved) and is compensating for the near vision loss that typically occurs with presbyopia. Index myopia is usually caused by the increase in refractive index of the lens with age, making it more myopic, particularly for distance vision. Type of Myopia Cause Axial Myopia Increase in the anteroposterior length of the eyeball (most common form). Curvatural Myopia Increased curvature of the cornea, lens, or both. Positional Myopia Anterior placement of the crystalline lens in the eye. Index Myopia Increase in the refractive index of the crystalline lens, often due to nuclear sclerosis. Myopia due to Excessive Accommodation Caused by spasm of accommodation, leading to excessive focusing power. Clinical features Asthenopic symptoms

Half-shutting of eyes for better clarity Prominent eyeball. Prominent, large, and somewhat sluggishly reacting pupils. Slightly deeper anterior chamber. Normal fundus; rarely a temporal myopic crescent may be seen. Treatment Prescription of appropriate concave lenses Surgical Treatment Incorrect Options: Axial myopia (Option B) : Caused by an elongated eyeball . This typically leads to myopia from birth or early in life and doesn't relate to the lens changes seen with aging. Curvatural myopia (Option C) : Caused by a change in the curvature of the cornea , which affects the focusing power of the eye. It's not related to the age-related changes in the lens. Positional myopia (Option D) : Refers to a refractive error caused by the position of the eye or the viewing distance (e.g., looking through different parts of the lens or an unusual gaze position). It's not linked to aging lens changes like index myopia.

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 28-30

Defining and Classifying Myopia

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: Which of the following refractive errors is associated with the image given below? (Q855491)

A. Presbyopia

B. Astigmatism


C. Hypermetropia

D. Myopia



Correct Answer: B. Astigmatism

Explanation: Astigmatism is a refractive error caused by an irregular shape of the cornea or lens, leading to blurred or distorted vision . This results in multiple focal points instead of a single point of focus on the retina. The image shows irregular, blurry distorted vision or varying focal points and is typically associated with astigmatism , where light entering the eye is refracted unevenly. Astigmatism A refractive error where the eye's refraction varies across different meridians. As a result, light rays entering the eye do not converge to a single point, instead form focal lines. {{caption_text}} Regular Astigmatism Refractive power changes uniformly from one meridian to another. There are 2 principal meridians. Oblique Astigmatism- two principal meridians are not horizontal and vertical but are at right angles to each other (e.g., 45° and 135°). Bioblique Astigmatism- two principal meridians are not at right angles to each other (e.g., one at 30° and the other at 100°). Irregular Astigmatism Irregular change in refractive power in different meridians, with multiple meridians that do not admit geometrical analysis Incorrect Options: Presbyopia (Option A) : This is an age-related condition where the eye's ability to focus on near objects decreases, typically around age 40. It causes difficulty in reading or seeing close objects, but it's not associated with distorted vision like astigmatism. Hypermetropia (Option C) (Farsightedness) : In hypermetropia, distant objects are seen clearly, but near objects appear blurry. This is due to a shorter eyeball or less curved cornea , not irregular refraction. Myopia (Option D) (Nearsightedness) : In myopia, nearby objects are seen clearly, but distant objects appear blurry, typically due to a longer eyeball or steeper cornea . It does not cause distorted vision as seen in astigmatism.



References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 31

Astigmatism

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 21 year old boxer wanted to improve the vision . Which of the following surgery is contraindicated? (Q854701)

A. RK

B. PRK

C. LASIK

D. SMILE

Correct Answer: A. RK

Explanation: Radial Keratotomy (RK) involves making radial incisions in the cornea to flatten it and correct myopia. It is no longer recommended due to its unstable results and the risk of complications , especially in young, active individuals. For a 21-year-old boxer , RK is contraindicated because it compromises the structural integrity of the cornea, increasing the risk of corneal weakening and potential rupture due to trauma, which is a significant concern in contact sports. Refractive surgeries for myopia Incisional Method Radial Keratotomy (RK) Involves radial cuts in the cornea to flatten it, correcting myopia. Used for low to moderate myopia (2-6 D). No longer recommended due to the advent of safer procedures like LASIK. Newer Methods Photorefractive Keratectomy (PRK) (Option B ruled out) It uses an excimer laser to reshape the cornea by removing a thin corneal layer. Suitable for correcting myopia up to -6 D. Refractive Lenticule Extraction (SMILE) : (Option D ruled out) A minimally invasive laser procedure where a small lenticule is removed to reshape the cornea. Effective for myopia up to -10 D, with or without astigmatism. Intercorneal Rings (ICR) : Rings are inserted into the cornea to flatten it and reduce refractive power. Results can be unpredictable, with risks such as corneal inflammation, so is not recommended Orthokeratology A non-surgical and reversible method involving rigid contact lenses worn overnight to reshape the cornea. Corrects myopia up to -5 D temporarily. High Myopia Solutions Clear Lens Extraction Removal of the natural lens (without cloudiness) to correct very high myopia (-16 D to -18 D). Less common now due to risks like retinal detachment. Phakic Intraocular Lens (IOL) Implantation Implantable lens placed without removing the natural lens. Best for myopia exceeding -8 D, especially in patients unsuitable for laser surgery. PRK , LASIK , and SMILE are safer alternatives for myopia correction and are well-suited for active individuals like athletes. These methods involve laser reshaping of the cornea with less risk of trauma-related complications than RK.

References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 284-286

↑ *Back to this question in Part A*

@Neet_pg_bot


Q14: A patient presents with complaints of eye strain. The given image shows the focus of light rays in different meridians in the eye. Which refractive error is present in this patient? (Q866834)

A. Amblyopia

B. Hypermetropia

C. Astigmatism


D. Presbyopia



Correct Answer: C. Astigmatism

Explanation: The image of Sturm's conoid demonstrates that parallel light rays do not converge to a single focal point, but rather form two focal lines at different distances due to unequal refractive power in different meridians of the cornea or lens. This optical phenomenon is characteristic of astigmatism , where light rays along one meridian focus in front of the retina while rays along the perpendicular meridian focus either behind or at a different point. Astigmatism causes blurred or distorted vision and eye strain (asthenopia), especially with prolonged near tasks. Sturm's conoid in Astigmatism: Light rays from an object pass through two principal meridians with different curvatures. Each meridian has its own focal length → creates two focal lines (not a single point). The area between these two focal lines is called the Sturm's conoid. The best vision is achieved at the circle of least confusion , located between the two focal lines. Incorrect Options: Amblyopia (Option A): Reduced vision in one or both eyes due to abnormal visual development in childhood; unrelated to optical focusing errors like those shown in Sturm's conoid. Hypermetropia (Option B): Farsightedness due to the eye being too short or the cornea too flat, causing light to focus behind the retina; it produces a single focal point (not two) and does not

show Sturm’s conoid. Presbyopia (Option D): Age-related loss of accommodation due to decreased lens elasticity; affects near vision but does not produce two focal lines in different meridians.



References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q15: Which of the following do not represent the corrected refractive error? (Q860210)

A. Myopia

B. Hyperopia


C. Presbyopia

D. Astigmatism



Correct Answer: C. Presbyopia

Explanation: Presbyopia does not represent in the above given image Presbyopia happens around the age of 40 years for an emmetropic person, less than 40 years in hypermetropic patients, and more than 40 years for myopic patients. At 40 years, around +1 D spherical lens is used to correct it. With every year that passes, around 0.5 D increases happen. The first image is of Myopia correction -It is done by concave lens Second image is of Hyperopia correction -It is done by convex lens Third image is of Astigmatism correction -It is done by cylindrical lens Definition Physiological insufficiency of accommodation is typically associated with aging. Etio - Pathogenesis Age-related factors: Decreased elasticity of the lens capsule. Sclerosis of the lens. Possible decline in ciliary muscle power. Premature causes: Uncorrected hypermetropia Chronic glaucoma Symptoms Difficulty in near vision Asthenopic symptoms (eye fatigue) Intermittent diplopia Treatment Options Optical Treatment: Convex glasses based on age Correct refractive error for distance first Types of Glasses: Unifocal, bifocal, or progressive (varifocal). Pharmacological therapy:1.25% Pilocarpine (FDA-approved) Surgical options: Conductive keratoplasty LASIK (Monovision/presbyopic multifocal-LASIK) Corneal inlays Multifocal IOL placement following lens extraction



References:

Myron Yanoff and Duker’s textbook of ophthalmology, 6th edition, Page no: 33-34, 48-49, 118-119

How to prescribe spectacles for presbyopia - PMC.

Age-related farsightedness (presbyopia)

Amplitude of accommodation in different age groups and age

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: What diagnosis is suggested by having different refractive powers (+5D in one eye and +1D in the other eye)? (Q860204)

A. Refractive errors

B. Amblyopia

C. Anisometropic amblyopia

D. Malingering

Correct Answer: C. Anisometropic amblyopia

Explanation: In this scenario, the child has two different refractive powers in both eyes; +5D in one eye and +1D in the other eye. Anisometropia is the difference between the refractive power of both eyes when > 2.5 D. Anisometropia can lead to Amblyopia Amblyopia: Definition, Types, and Clinical Characteristics Definition: Partial, reversible vision loss in one or both eyes. Absence of organic diseases of ocular media, retina, visual pathway The eye appears normal, but vision is impaired. Types of Amblyopia: Type Cause Key Features Strabismic Amblyopia The brain suppresses input from one eye due to constant squint (strabismus). Common cause; underdevelopment of vision in the deviated eye. Stimulus Deprivation Amblyopia Hindered vision due to obstruction from early life (e.g., congenital cataracts, ptosis, dense central corneal opacity) . Lack of visual stimulation affects normal development. Anisometropic Amblyopia (Option C ruled out) A significant difference in refractive error between eyes. The brain favours the clearer eye, leading to amblyopia in the other. Ametropic Amblyopia High uncorrected refractive errors in both eyes. Bilateral; is less severe compared to other types. Meridional Amblyopia Uncorrected astigmatism affecting specific visual meridians. Affects one or both eyes depending on the astigmatism degree. Refractive errors (Option A) While the condition involves refractive errors, the key point is the significant difference in refractive powers between the two eyes. This disparity leads to anisometropia, which can cause amblyopia. General refractive errors alone do not necessarily lead to amblyopia. Malingering (Option D) refers to the deliberate faking of symptoms.

References:

Kanski’s clinical ophthalmology, John F. Salmon, 10th edition, Page no: 718-719

Amblyopia - PMC

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: What is Anisometropia ? (Q840053)

A. Difference in visual acuity

B. Difference in refractive error between 2 eyes

C. Difference in the axial length of the eyeball

D. Difference in size of eyeball

Correct Answer: B. Difference in refractive error between 2 eyes

Explanation: Correct Option B - Difference in refractive error between 2 eyes: Anisometropia is when total refraction of the two eyes is unequal. The difference in power of two eyes is more than 2.5 diopters. Anisometropia up to 2.5D is well tolerated and that between 2.5 and 4D can be tolerated depending upon the individual sensitivity. However, if it is more than 4D, it is not tolerated and is a matter of concern. Incorrect Options: Option A - Difference in visual acuity: Differences in visual acuity could be caused by conditions like myopia, hypermetropia, astigmatism, and presbyopia. Option C - Difference in axial length: The difference in the axial length of the eyeball is seen in axial myopia. Option D - Difference in size of eyeball: The difference in the size of the eyeball is seen in conditions like Microphthalmia and Buphthalmos.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: During the Worth 4-Dot Test, a patient reports seeing only three green dots . What is the most likely diagnosis? (Q843991)

A. Right eye suppression

B. Left eye suppression

C. Crossed diplopia with esodeviation

D. Crossed diplopia with exodeviation

Correct Answer: A. Right eye suppression

Explanation: Correct Option A - Right eye suppression: The Right Eye (Red Filter): Can only see the Red dot and the White dot (which appears red). It is "blind" to the green light. The Left Eye (Green Filter): Can only see the two Green dots and the White dot (which appears green). It is "blind" to the red light. If the patient reports seeing only three green dots, they are using only their Left eye . The Left eye sees the two green dots plus the white dot (as a 3rd green dot). Because the patient sees no red dots , the brain is actively ignoring (suppressing) the signal from the Right eye . Incorrect Options: Option B (Left Eye Suppression): If the Left eye were suppressed, the patient would see only with their Right eye. They would report seeing two red dots (the one red dot plus the white dot appearing red). Option C & D (Diplopia): In diplopia, neither eye is suppressed, but the brain fails to fuse the images. The patient will see the dots from both eyes simultaneously, totaling five dots (2 red and 3 green). Esodeviation (Uncrossed) results in the red dots appearing to the right of the green dots. Exodeviation (Crossed) results in the red dots appearing to the left of the green dots.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: Which of the following is a cause of bilateral sudden painless loss of vision? (Q853148)

A. Acute congestive glaucoma

B. Diabetic retinopathy


C. Keratoconus D) Iridocyclitis

D. Iridocyclitis

Correct Answer: B. Diabetic retinopathy

Explanation: Diabetic retinopathy often develops in stages, with initial mild symptoms that may progress to sudden vision loss due to complications. Bilateral sudden painless loss of vision can be caused by diabetic retinopathy, particularly in its proliferative form, where there is a sudden onset of vision loss due macular edema, or retinal detachment. Diabetic retinopathy affects both eyes and can lead to significant vision impairment if left untreated. Clinical Features of Proliferative Diabetic Retinopathy (PDR): Includes all features of preproliferative diabetic retinopathy or severe NPDR: Multiple retinal hemorrhages, cotton-wool spots, venous beading, vascular loops, intraretinal microvascular abnormalities (IRMAs), and large areas of capillary nonperfusion along with neovascularization. Neovascularization (2 types): Neovascularization of the Disc (NVD): Arises within 1 disc diameter of the optic nerve from the optic nerve head. Neovascularization Elsewhere (NVE): Arises more than 1 disc diameter from the optic nerve, along the large vessels. Appears adjacent to areas of capillary closure and the accompanying fibrous tissue varies in extent. Fluorescein angiography (FA): Irregular expanding

hyperfluorescence. Severe Vision Loss: Result in vision as poor as hand movements or worse due to severe vitreous hemorrhage or retinal detachment. Fibrovascular Tissue Formation: Formed due to condensation of connective tissue around the new vessels. Lie flat on the retina or attach to the posterior vitreous face. Can lead to vitreous traction, retinal separation, and blood vessel tearing. The most common cause of vitreous hemorrhage in adults. Rubeosis Iridis: Neovascularization of iris. Neovascular glaucoma: Occurs due neovascularization of anterior segment angle. Acute congestive glaucoma (Option A): Acute congestive glaucoma is typically associated with painful loss of vision and is more often unilateral. The condition involves a sudden increase in intraocular pressure, causing symptoms such as severe eye pain, nausea, and vomiting. Keratoconus (Option C): Keratoconus can lead to progressive vision loss due to the distortion of the cornea, but it usually presents gradually, not suddenly, and typically affects one eye more than the other initially. Iridocyclitis (Option D): Iridocyclitis (anterior uveitis) is usually associated with pain , photophobia, and redness in the affected eye. It does not cause sudden, painless loss of vision in both eyes.



References:

Parsons Diseases of the Eye, Edition 23, page no: 291,292

Yanoff and Duker Ophthalmology, Edition 5, page no: 546

Kanski’s Clinical Ophthalmology, Edition 9, page no: 505

Diabetic Retinopathy - StatPearls - NCBI Bookshelf

https://www.ncbi.nlm.nih.gov/books/NBK430857/

https://www.ncbi.nlm.nih.gov/books/NBK470435/

https://www.ncbi.nlm.nih.gov/books/NBK430909/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: What is the normal artery-to-vein ratio at the optic disc margin? (Q853134)

A. 1:3

B. 2:3

C. 2:5

D. 1:5

Correct Answer: B. 2:3

Explanation: The normal artery-to-vein (A:V) ratio at the optic disc margin is approximately 2:3 in a healthy eye. This ratio is clinically important in assessing retinal vascular changes. A significantly reduced A:V ratio suggests worsening microvascular disease and increased risk of retinal complications. Hypertension can reduce the A:V ratio (e.g., 1:3 or 1:4) due to arteriolar narrowing. Diabetes mellitus may also cause changes, but typically with venous dilatation and beading.

References:

https://pmc.ncbi.nlm.nih.gov/articles/PMC9333006/ https://pmc.ncbi.nlm.nih.gov/articles/PMC5437764/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: A 22-year-old patient with high myopia and thick glasses presents with floaters in his eyes. What is the likely cause of this condition? (Q853121)

A. Subarachnoid hemorrhage

B. Posterior Vitreous Detachment

C. Vitreous hemorrhage

D. Branched retinal artery occlusion

Correct Answer: B. Posterior Vitreous Detachment

Explanation: Posterior Vitreous Detachment The likely cause of floaters in a 22-year-old patient with high myopia and thick glasses is Posterior Vitreous Detachment (PVD). In high myopia, the elongated eyeball causes early liquefaction and shrinkage of the vitreous , leading to its detachment from the retina . This results in floaters

(moving black spots) while vision remains normal . A fundus exam is needed to rule out associated retinal tear or detachment . Subarachnoid hemorrhage (Option A): This is a life-threatening brain bleed. The main symptom would be a severe, sudden headache ("worst headache of life"), not isolated floaters. Vitreous hemorrhage (Option C): Causes sudden, dense floaters with decreased vision and a hazy fundus, not just mild floaters with normal vision. Branched retinal artery occlusion (Option D): This is a vascular blockage causing sudden, localized vision loss and visual field defects. It does not typically present with floaters.

References:

AK Khurana Textbook of Ophthalmology Edition 7th Pg. 39, 40, 41.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q22: Binocular single vision consists of the following components: (Q866361)

A. a. SMP b. Fusion

B. a. Stereopsis b. Fusion

C. a. Fusion b. SMP

D. a. SMP b. Stereopsis



Correct Answer: A. a. SMP b. Fusion

Explanation: a. SMP b. Fusion The first image shows two dissimilar pictures meant to be seen simultaneously (e.g., dog and its kennel parts) — this is a test of Simultaneous Macular Perception (SMP). The second image

shows two complementary/similar pictures intended to be merged into one percept (one image completes the other) — that is a test for Fusion. Binocular single vision: SMP (Simultaneous Macular Perception): the baseline ability to perceive two images (one from each fovea) at the same time; absence suggests suppression. Fusion: the cortex's process of aligning and merging the two similar images into one percept — requires both motor alignment and sensory correspondence. Functional relationship: SMP permits inputs; fusion integrates them; without SMP fusion is hard to establish. Clinical tests : SMP and fusion are tested with synoptophore slides and Worth's four-dot before testing stereopsis (Titmus/Randot). Grade Name Description Grade 1 SMP Perception of two dissimilar images simultaneously Grade 2 Fusion Merging of two similar images into one Grade 3 Stereopsis Depth perception from binocular vision a. Stereopsis b. Fusion (Option B) is incorrect because the first image is not a stereopsis (depth) test. Stereopsis tests use binocular disparity/random-dot or contour-based depth targets, not two dissimilar objects; you can't get depth from that first slide. a. Fusion b. SMP (Option C) - Wrong mapping: the first image is not designed for fusion (it shows dissimilar items), and the second is clearly intended for merging — so the pair is reversed. a. SMP b. Stereopsis (Option D) is incorrect because the second image is not a stereopsis/depth stimulus; it's a fusion stimulus. Stereopsis requires disparity-based targets (Titmus/Randot), not complementary pictorial halves.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: Which of the following is not an effective treatment for amblyopia? (Q865351)

A. Correction of refractive errors

B. Occlusion therapy (patching)

C. Strabismus surgery

D. Use of video games for visual stimulation

Correct Answer: C. Strabismus surgery

Explanation: Correction of refractive errors (Option A ruled out) : This is a fundamental and often the first step in amblyopia treatment. Uncorrected significant refractive errors (like high farsightedness, nearsightedness, or astigmatism) can prevent clear image formation and lead to amblyopia. Correcting these errors with glasses or contact lenses allows the brain to receive a clear image, which is essential for visual development. This is an effective treatment. Occlusion therapy (patching) (Option B rule out) : This is a cornerstone of amblyopia treatment. It involves covering the stronger, non-amblyopic eye for specific periods each day, forcing the brain to use and develop the weaker, amblyopic eye. This is a highly effective and widely used treatment. Strabismus surgery (Option C) : Strabismus (misalignment of the eyes) is a common cause of amblyopia. While strabismus surgery can align the eyes, it primarily addresses the cosmetic and functional alignment issue. It does not directly treat the amblyopia itself, which is a brain-based vision deficit. However, aligning the eyes before or during amblyopia treatment can sometimes improve the chances of success, especially in preventing suppression. It is not a direct or primary treatment for the amblyopia deficit itself, but for its cause. Use of video games for visual stimulation (Option D rule out) : There has been research into using specially designed video games (often dichoptic, where each eye sees different parts of the game) to stimulate both eyes simultaneously and improve

visual acuity in amblyopia. While this is a newer and emerging area of research, it has shown promise as an adjunctive or alternative therapy, particularly in older children or those who don't tolerate patching well. This is considered an emerging and potentially effective treatment.

References:

Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 718-719

pmc.ncbi.nlm.nih.gov/articles/PMC2907781/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q24: A posterior staphyloma was observed on indirect ophthalmoscopy. What is the likely diagnosis? (Q866854)

A. High myopia


B. Hypermetropia

C. Primary open-angle glaucoma

D. Retinitis pigmentosa

Correct Answer: A. High myopia

Explanation: A posterior staphyloma is an outpouching of the weakened posterior sclera, visible as an ectatic area on fundus examination. It occurs most commonly in pathological (high) myopia due to progressive elongation of the eyeball, leading to stretching and thinning of the sclera, choroid, and retina at the posterior pole. This structural change can cause progressive visual decline and predispose to retinal degenerations/detachments. Axial length usually >26 mm. Fundus signs: temporal crescent, chorioretinal atrophy, lattice degeneration, and posterior staphyloma. Etiology: mechanical stretching from axial elongation. Clinical impact: distorted vision, higher retinal detachment risk. Staphyloma: It is ectasia of the outer coats of the eye (cornea, sclera, or both) with an incarceration of the uveal tissue. Based on the site affected, staphyloma can be classified as Types of staphyloma Causes Anterior Pseudocornea Intercalary Perforating injuries of the peripheral cornea, marginal corneal ulcer, anterior scleritis, and scleromalacia perforans Ciliary Glaucoma, scleritis, and trauma to the ciliary region Equatorial (approx 14 mm behind the limbus) Scleritis, degenerative myopia, and chronic uncontrolled glaucoma Posterior Degenerative high axial myopia Treatment: Treat underlying condition Local excision and repair with a corneal and scleral patch graft Large staphyloma: Staphylectomy and keratoplasty, or enucleated and replaced with an implant Incorrect Options: Hypermetropia (Option B): Shorter axial length; no scleral ectasia; may cause narrow anterior chamber and angle-closure glaucoma but not posterior staphyloma. Primary open-angle glaucoma (Option C): Optic nerve cupping and visual field loss without scleral ectasia; unrelated to posterior staphyloma formation. Retinitis pigmentosa (Option D): Inherited retinal dystrophy; shows bone-spicule pigmentation, attenuated vessels, and waxy disc pallor — not posterior scleral outpouching.



References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q25: The following light reflex is seen in a patient. How much is the deviation? (Q866850)

A. 30 PD

B. 15°

C. 90 PD

D. 30°



Correct Answer: C. 90 PD

Explanation: The image shows the Hirschberg test result — the corneal light reflex is displaced toward the limbus in one eye, which will correspond to 90PD deviation. In the Hirschberg test: 1 mm corneal reflex shift ≈ 7° deviation 1° deviation ≈ 2 PD (quick approximation) Reflex Displacement (from centre) Approx. Deviation (Degrees) Approx. Prism Diopters (PD) Centered 0° 0 PD 1 mm 7° 14 PD 2 mm 15° 30 PD Midway between centre & limbus 30° 60 PD At limbus 45° 90 PD

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q26: A 45-year-old patient with eye examination findings of a deep anterior chamber and jet-black pupil is prescribed +12D glasses. Likely diagnosis? (Q866838)

A. Pseudophakia

B. Aphakia

C. Myopia

D. Hypermetropia

Correct Answer: B. Aphakia

Explanation: Clinical reasoning: Deep anterior chamber → suggests absence of crystalline lens. Jet-black pupil → no lens reflections seen. +12D prescription → large convex lens to replace lost refractive power of natural lens. Aphakia = complete absence of the natural lens (usually post-cataract surgery without IOL implantation). Without the lens, refractive power decreases drastically (~+15D needed), so strong convex lenses are prescribed. Causes: Post-cataract extraction without IOL, trauma, congenital. Features: Deep anterior chamber, iridodonesis, jet-black

pupil, high hypermetropia (+10 to +15D). Aphakia: Definition Absence of crystalline lens from the eye. Aetiology Congenital absence (rare) Surgical removal (common) Traumatic extrusion of lens (rare) Posterior lens dislocation (optical aphakia) Optical Effects High hypermetropia Reduced eye power to +44 D (from +60 D) Symptoms Defective vision (far and near) Erythropsia and cyanopsia (abnormal red and blue color defect) Signs Limbal/corneal scar (in surgical cases) Deep anterior chamber Iridodonesis & Jet-black pupil Only 2 Purkinje images (normal: 4) Hypermetropic small optic disc Treatment Correct refractive error with: Spectacles: the problem of spherical aberration and roving ring scotoma (jack-in-the-box phenomenon) Contact lenses Intraocular lenses (IOL) - best method Refractive LASIK surgery Incorrect Options: Pseudophakia (Option A): Eye has an artificial intraocular lens post-cataract surgery; patient usually needs minimal refractive correction (not +12D). Myopia (Option C): Nearsightedness; requires concave (minus) lenses; opposite of the strong convex lenses seen here. Hypermetropia (Option D): Farsightedness; needs convex lenses, but typically much lower power (+1D to +4D), not as high as +12D.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [12_Basics of Ophthalmology] 026 Introduction To Ophthalmology

« Back to Folder Directory

[12_Basics of Ophthalmology] 026 Introduction To Ophthalmology — Quick Jump

Q1 (p. 1444) *Q2 (p. 1445)* *Q3 (p. 1447)* *Q4 (p. 1448)* *Q5 (p. 1450)* *Q6 (p. 1451)*
Q7 (p. 1453) *Q8 (p. 1454)* *Q9 (p. 1455)* *Q10 (p. 1457)*

Q1: A 25-year-old patient presents with redness and discomfort in the eyes. On examination, the physician notes inflammation confined to the conjunctiva attached to the posterior tarsal plate. Which part of the conjunctiva is primarily affected in this patient? (Q853247)

A. Bulbar conjunctiva

B. Forniceal conjunctiva

C. Palpebral conjunctiva


D. Limbal conjunctiva


Correct Answer: C. Palpebral conjunctiva

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The palpebral conjunctiva lines the inner surface of the eyelids, which start at lid margins and are attached to posterior tarsal plates; while the bulbar conjunctiva covers the anterior sclera, which is continuous with corneal epithelium at limbus. Anatomical parts: Palpebral conjunctiva: Firmly attached to posterior tarsal plate; vertical vessels. Forniceal conjunctiva (Option B): Loose, redundant fold allowing eye movement. This forms the junction between the palpebral and bulbar conjunctiva in the fornices (superior and inferior). It’s not directly attached to the tarsal plate. Bulbar conjunctiva (Option A): Covers anterior sclera, merges with corneal epithelium at limbus (with palisades of Vogt as corneal stem cell niches). Limbal conjunctiva(Option D) : This is the conjunctiva near the limbus, which is the border between the cornea and sclera — again, not attached to the tarsal plate.





References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 168

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Which of the following structures do not contain any ganglion cells? (Q853248)

A. Macula

B. Fovea

C. Foveola


D. Periphery of the retina

Correct Answer: C. Foveola

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The foveola is the central part of the fovea and is unique in that it lacks ganglion cells, which allows for optimal light entry, contributing to high visual acuity. It contains only cones. Macula lutea Circular, 5–5.5 mm in diameter Covers central 15–20° of visual field Contains yellow pigments lutein and zeaxanthin Fovea Central depression of macula, ~1.5 mm diameter (same as optic disc) Foveola Central floor of fovea, ~0.35 mm diameter Thinnest part of retina Contains only cone photoreceptors & Müller cells (no ganglion cells) Umbo Central pit of foveola Associated with foveolar light reflex (loss = early damage) Foveal avascular zone (FAZ) Vessel-free zone within fovea, ~0.6 mm diameter Surrounded by capillaries Best visualized by fluorescein angiography



References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 556

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Which of the following is the anteroposterior diameter of the adult human eyeball? (Q853249)

A. 23 mm

B. 23.5 mm

C. 24 mm


D. 75 mm

Correct Answer: C. 24 mm

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Anteroposterior diameter (AKA Axial length) of the adult human eyeball is 24 mm. A-scan is used for measurement of axial length. Note: B-scan is used to see the posterior segment of the eye. Depth of anterior chamber: 2.4mm Axes & Angle Kappa Visual axis : Fixation point → nodal point → fovea Anatomical axis : Posterior pole → corneal center Angle α : Visual vs optic axis Angle κ : Visual vs anatomical axis Positive κ (fovea temporal) → nasal corneal reflex, pseudoexotropia Negative κ (fovea nasal, e.g. high myopia) → temporal corneal reflex, pseudoesotropia



References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 698, 726

AK Khurana Comprehensive Ophthalmology, 7th Edition, Page 3

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: Which structure primarily offers resistance to aqueous humor outflow within the iridocorneal angle? (Q853250)

A. Iris

B. Ciliary Body

C. Descemet's Membrane


D. Trabecular Meshwork


Correct Answer: D. Trabecular Meshwork

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Trabecular meshwork (especially the juxtacanalicular portion) is main site of resistance in aqueous humor drainage Three parts of trabecular meshwork: Uveal meshwork – innermost, least resistance Corneoscleral meshwork – intermediate resistance Juxtacanalicular (cribriform) meshwork – maximum resistance to outflow → site of major regulation Iris (Option A) does not play a significant role in regulating the aqueous humour outflow or offering resistance to its drainage. Ciliary body (Option B) produces aqueous humour but does not resist its outflow. Descemet's membrane (Option C) contributes to structural support, not resistance.





References:

Parsons Diseases of the Eye, 23rd Edition, Pages 64-75

AK Khurana Comprehensive Ophthalmology, 7th Edition, Pages 14-19

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: Which of the following statements accurately describes the role of retinal ganglion cells in the visual process? (Q853251)

A. They are the primary photoreceptors responsible for converting light into electrical signals.

B. They are concentrated in the fovea and are responsible for colour vision and high visual acuity.

C. Their axons form the optic nerve, which transmits visual information to the brain.


D. They contain pigments that undergo structural changes when exposed to light, triggering signal transduction.

Correct Answer: C. Their axons form the optic nerve, which transmits visual information to the brain.

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Ganglion cell axons → converge at the optic disc → form the optic nerve. They carry processed visual signals from inner retina to the lateral geniculate body (LGB) and then to the visual cortex. Primary photoreceptors are responsible for converting light into electric signals (Option A): This is the function of rods and cones (photoreceptors). Concentrated in the fovea, responsible for color vision and high acuity (Option B): This describes cones, not ganglion cells. Contain pigments that undergo structural change in response to light (Option D): This refers to rhodopsin in rods (visual pigment in phototransduction). Types of Ganglion Cells: X ganglion cells → responsible for color vision Y ganglion cells → motion detection, peripheral vision W ganglion cells → miscellaneous, including pupillary reflex pathways



References:

Parsons Diseases of the Eye, 23rd Edition, Page 32

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: What is the primary use of the instrument shown in the image given below? (Q853252)

A. To measure intraocular pressure.

B. To examine the anterior segment of the eye.

C. To provide a wide-field view of the retina.

D. To capture high-resolution images of the retina.



Correct Answer: C. To provide a wide-field view of the retina.

- *Listen (English)*
- *Listen (Hindi)*

Explanation: The given image is an Indirect Ophthalmoscope, device used to examine the retina and provide wide-field view of the retina. Clinical use: Best for peripheral retinal examination, e.g., retinal detachment, tears, degenerations. DIRECT OPHTHALMOSCOPE INDIRECT OPHTHALMOSCOPE 15x magnification Virtual image, erect Field of view: 2 DD Monocular Undilated pupil Stereopsis absent 2-3x magnification Real image, inverted Field of view: 8 DD Binocular Dilated pupil Stereopsis present Tonometer: Measures intraocular pressure (Option A) Slit lamp: Examines the anterior segment of the eye (Option B) Fundus camera: Captures high-resolution images of the retina (Option C) , but not wide-field like indirect ophthalmoscopy



References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 25

Myron Yanoff and Jay S. Duker, Ophthalmology, 5th Edition, Page 56, 57

www.ncbi.nlm.nih.gov/pmc/articles/PMC5340107/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: Which of the following arteries is not a branch of the ophthalmic artery? (Q853253)

A. Supratrochlear Artery

B. Dorsal Nasal Artery

C. Transverse Facial Artery


D. Short Posterior Ciliary Artery

Correct Answer: C. Transverse Facial Artery

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Transverse Facial Artery: This artery is a branch of the superficial temporal artery and not ophthalmic artery, which is a branch of the external carotid artery. It supplies the skin of the face. Arterial Supply of eyeball and orbital structures Ophthalmic artery: Branch of internal carotid artery constitutes the main source of blood supply for the eyeball and orbital structures. Ocular Branches Orbital Branches Central retinal artery (supplies inner 6 layers of retina) Long posterior ciliary arteries → Only 2 in number. Short posterior ciliary arteries (Option D) → 10–12 in number. Anterior ciliary arteries - Muscular branch Lacrimal artery - Lateral palpebral artery Recurrent meningeal artery Supraorbital artery Medial palpebral artery Posterior ethmoidal artery Anterior ethmoidal artery Dorsal nasal artery (terminal branch) (Option B) Supratrochlear artery (terminal branch) (Option A) Branches of Ophthalmic artery:



References:

AK Khurana Comprehensive Ophthalmology, 7th Edition, Page 5

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 32-year-old patient presents with an orbital infection characterized by severe pain, swelling, and redness around the eye. Despite treatment, the patient develops a headache, fever, and neurological deficits, leading to a diagnosis of a brain abscess. What is the most likely route of spread of the infection? (Q853254)

A. Ethmoidal veins to the cavernous sinus

B. Facial vein to the cavernous sinus

C. Superior ophthalmic vein to the cavernous sinus


D. Angular vein to the internal jugular vein

Correct Answer: C. Superior ophthalmic vein to the cavernous sinus

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above case is suggestive of orbital cellulitis. The superior ophthalmic vein directly drains blood from the orbit into the cavernous sinus, providing a pathway for infections to enter the cranial cavity. This route is significant in cases of orbital infections, as it can lead to serious complications such as brain abscesses and cavernous sinus thrombosis. The venous drainage of the eye primarily occurs through: Superior ophthalmic vein: It receives drainage from the angular, supratrochlear, and supraorbital veins, superior vortex veins, anterior ethmoidal vein, and receives collateral drainage from the inferior ophthalmic vein. It continues posteriorly to drain into the cavernous sinus. Inferior ophthalmic vein: It originates from a plexus in the inferior orbit and collects blood from the inferior vortex veins. It eventually drains into the cavernous sinus.



References:

Myron Yanoff and Jay S. Duker, Ophthalmology, 5th Edition, Page 1268

↑ *Back to this question in Part A*

@Neet_pg_bot


Q9: A patient presents to the ophthalmology OPD with the following features: Which of the following conditions is most likely responsible for these symptoms? (Q853255)

A. Abducens nerve palsy

B. Optic neuritis

C. Oculomotor nerve palsy

D. Trochlear nerve palsy



Correct Answer: C. Oculomotor nerve palsy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The left eye appears downward and abducted, commonly known as down and out pupil due to paralysis of the inferior oblique, and medial rectus muscles. There is ptosis of left eye (upper eyelid droop), and the pupil is dilated in the left eye (as the constrictor pupillae is denervated). These signs are suggestive of oculomotor nerve palsy (Left side). Oculomotor nerve palsy: Causes ptosis due to paralysis of levator palpebrae superioris . Eye deviates “down and out” because only lateral rectus (VI) and superior oblique (IV) are functioning. Diplopia (double vision) is common due to misalignment. Pupil may be dilated (mydriasis) if parasympathetic fibers are involved. Loss of accommodation and light reflex can occur. Abducens nerve palsy (VI) (Option A): Causes inability to abduct the eye (lateral movement). Eye is pulled medially → medial deviation , not “down and out”. Optic neuritis (II) (Option B): Affects vision (visual acuity, color vision) , not eye movements. No ptosis or ocular deviation. Trochlear nerve palsy (IV) (Option D): Affects superior oblique muscle → difficulty in looking down and in . Causes vertical diplopia , often with head tilt , not full 3rd nerve features. Nerves of the eye: Optic nerve (CN II) Purely sensory Senses the incoming light and image displayed on retina Oculomotor nerve (CN III) Innervates majority of the extraocular muscles Motor innervation to the superior rectus muscle, medial rectus muscle, inferior rectus muscle, inferior oblique muscle, levator palpebrae superioris muscle, ciliary muscle, and the sphincter muscle. Trochlear nerve (CN IV) Motor innervation to the superior oblique muscle Trigeminal Nerve (CN V) Ophthalmic branch of the trigeminal nerve provides sensory innervation to the

eye Afferent part of corneal and lacrimation reflex Abducens Nerve (CN VI) Motor innervation to the lateral rectus muscle Facial Nerve (CN VII) Motor innervation of the orbicularis oculi muscle Efferent part of corneal and lacrimation reflex Sympathetic Nervous System Long ciliary nerves: Innervates the pupillary dilator muscles causing mydriasis Parasympathetic nervous system Short ciliary nerves: Innervates the sphincter pupillae muscles causing miosis

References:

Anatomy, Head and Neck: Eye Nerves - StatPearls - NCBI Bookshelf (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: During an anatomy lecture, a student learns about the innervation of the eye. The pupil dilates in response to low light conditions. Which nerve fibers are primarily responsible for the dilation of the pupil? (Q853257)

A. Oculomotor nerve fibers

B. Trigeminal nerve fibers

C. Sympathetic nerve fibers

D. Facial nerve fibers

Correct Answer: C. Sympathetic nerve fibers

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Sympathetic nerve fibers are responsible for the dilation of the pupil (mydriasis). These fibers originate from the sympathetic chain and supply the dilator pupillae muscle, allowing the pupil to expand in response to low light conditions, thus enabling more light to enter the eye. Autonomic supply of the eye: Sympathetic Nervous System Long ciliary nerves: Innervates the pupillary dilator muscles causing mydriasis Parasympathetic nervous system Short ciliary nerves: Innervates the sphincter pupillae muscles causing miosis

References:

Anatomy, Head and Neck: Eye Nerves - StatPearls - NCBI Bookshelf (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

• [12_Basics of Ophthalmology] 027 Previous Year Questions

« Back to Folder Directory

[12_Basics of Ophthalmology] 027 Previous Year Questions — Quick Jump

Q1 (p. 1458)	*Q2 (p. 1459)*	*Q3 (p. 1460)*	*Q4 (p. 1461)*	*Q5 (p. 1464)*	*Q6 (p. 1464)*
Q7 (p. 1466)	*Q8 (p. 1466)*	*Q9 (p. 1468)*	*Q10 (p. 1468)*	*Q11 (p. 1469)*	*Q12 (p. 1471)*
Q13 (p. 1473)	*Q14 (p. 1474)*	*Q15 (p. 1475)*	*Q16 (p. 1477)*	*Q17 (p. 1478)*	*Q18 (p. 1482)*
Q19 (p. 1486)	*Q20 (p. 1487)*	*Q21 (p. 1488)*	*Q22 (p. 1489)*	*Q23 (p. 1490)*	


Q1: Identify the instrument used for handling delicate tissues during eye surgery. (Q860570)

A. Mcpherson forceps

B. Lim's forceps

C. Artery forceps


D. SR holding forceps



Correct Answer: A. Mcpherson forceps

Explanation: The image shows McPherson forceps , a fine-toothed, spring-handle microsurgical forceps designed for delicate tissue handling during ophthalmic procedures such as cataract surgery, keratoplasty, and intraocular lens implantation. They allow precise grip and minimal trauma to ocular structures, especially the cornea and lens capsule. Incorrect options B) Lim’s forceps: Used in trabeculectomy to handle conjunctival and scleral tissues —

not primarily for fine intraocular manipulation. C) Artery forceps: Used for hemostasis by clamping blood vessels; too coarse for delicate eye tissues. D) SR holding forceps (Superior Rectus holding): Used to grasp the superior rectus muscle during extraocular muscle surgery, not for intraocular work.



References:

Parsons Diseases of the Eye 23rd edition Pg no.: 1046

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Which laser is considered the most appropriate for cataract surgery? (Q859777)

A. Nd-YAG

B. Diode

C. Excimer

D. Femtosecond laser

Correct Answer: D. Femtosecond laser

Explanation: The Femtosecond laser is the most suitable laser for cataract surgery presently. It is used in a procedure called Femtosecond Laser-Assisted Cataract Surgery (FLACS). Femtosecond lasers: Mechanism: They use ultra-short pulses of laser light to ionize tissue, creating a plasma bubble that expands and allows for precise, controlled cutting of the tissue. Advantages: It provides high precision, reduces thermal damage, and improves safety compared to traditional surgical methods. Limitations: Femtosecond lasers are costly and require specialised

surgeon training, which can limit their accessibility and increase the overall cost of surgery. Applications: Femtosecond lasers are commonly used in Femtosecond Laser-Assisted Cataract Surgery (FLACS) and Laser-Assisted In Situ Keratomileusis (LASIK) for precise lens fragmentation, corneal flap creation, and refractive correction. PRK (photorefractive keratectomy) SMILE (small incision lenticule extraction) for correction of myopia Creation of tunnel for intracorneal rings Arcuate incisions to correct corneal astigmatism Keratoplasty incisions Lamellar and full-thickness corneal transplants Incorrect Options: Nd-YAG laser (Option A) is primarily used for posterior capsulotomy for the management of posterior capsular opacification.. Nd-YAG laser is not the primary laser used for cataract removal. Diode laser (Option B) are used in iridotomy for the management of glaucoma. Excimer laser (Option C) are primarily used in refractive surgeries such as LASIK (laser-assisted in situ keratomileusis) to reshape the cornea and correct refractive errors

References:

Parsons Diseases of the eye, 23rd Edition, Pg 357.

Yanoff & Duker Ophthalmology, 5th edition, Pg 98

Kanski's Clinical Ophthalmology, 10th Edition, Page 381

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Which laser is most commonly used for performing a laser iridotomy? (Q859748)

A. Argon laser

B. Nd:YAG laser


C. Excimer laser

D. CO■ laser

Correct Answer: B. Nd:YAG laser

Explanation: Laser Peripheral Iridotomy (LPI), used for treating angle-closure glaucoma , uses an argon or Nd: YAG laser. It operates via a photodisruptive mechanism, which vaporizes the iris tissue to create a precise and clean opening without causing significant thermal damage to surrounding structures. This procedure involves creating a small hole in the peripheral iris to re-establish communication between the anterior and posterior chambers, bypassing the pupillary block and relieving angle crowding in glaucoma. {{caption_text}} Argon laser (Option A): The Argon laser is used for laser iridotomy but is typically employed to pre-treat the iris by thinning the tissue before creating an opening. It uses a photocoagulative mechanism, which involves thermal energy. However, it is less effective than the Nd:YAG laser for standalone iridotomy due to higher chances of tissue damage and incomplete penetration. Excimer laser (Option C): Excimer lasers are not used for iridotomy. These lasers emit ultraviolet light and are designed for corneal refractive procedures like LASIK and PRK. They ablate corneal tissue rather than creating openings in the iris. CO■ laser (Option D): CO■ lasers are suitable in ophthalmology to vaporize lymphangiomas and capillary hemangiomas and to provide hemostasis in patients with

bleeding disorders.



References:

Parsons Diseases of the eye, 23rd Edition, Pg 357,673,675, 676.

Myron Yanoff & Duker Ophthalmology, 6th Edition, Page 98, 429

Kanski's Clinical Ophthalmology, 10th Edition, Page 332, 381

https://pmc.ncbi.nlm.nih.gov/articles/PMC7004285/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: What kind of laser can be utilized to treat posterior capsular opacification, which is observed during an examination of a patient who has reported decreased visual acuity following cataract surgery? (Q858163)

A. Femto laser


B. Nd YAG laser


C. Argon fluoride

D. Argon

Correct Answer: B. Nd YAG laser

Explanation: efinition M/C postoperative complication following extracapsular lens extraction. Clinical Types Posterior Capsule Opacification (PCO) : Common form of after cataract. Dense Membranous After Cataract: Thick membrane formation. {{caption_text}} Soemmering's Ring: Thick ring behind the iris, enclosed between capsule layers. {{caption_text}} Elschnig's Pearls: Clustered, vacuolated subcapsular epithelial cells along the posterior capsule. {{caption_text}} Treatment ND YAG laser posterior capsulotomy Femto laser (Option A) : The femtosecond laser is used in cataract surgery for making precise incisions and can be used for corneal surgery, but it is not used for treating posterior capsular opacification. Argon fluoride (Option C) : This laser is primarily used for retinal treatments , such as for retinal photocoagulation, and is not suitable for posterior capsular opacification. Argon laser (Option D) : The argon laser is used for retinal procedures (like retinal photocoagulation) and does not have the specific characteristics needed to treat PCO effectively.







References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 221

Kanski’s clinical ophthalmology, 9th edition, Page 331

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: Which of the following is not included in the SAFE strategy for trachoma? (Q856713)

A. Antibiotics

B. Surgery for trichiasis

C. Evaluation of control program

D. Facial cleanliness

Correct Answer: C. Evaluation of control program

Explanation: Evaluation of control program is NOT included in the SAFE strategy for Trachoma. Diagnosis and Management of Trachoma: Investigations Micro-immunofluorescence (micro-IF): Using pooled antigens; For routine diagnostic purposes Culture of C. trachomatis: In irradiated McCoy cells; Expensive Monoclonal antibody direct tests: IgA-IPA light microscopy tests are the best combination for diagnosis Management (SAFE strategy) SAFE is a WHO-recommended combination of interventions to treat trachoma and prevent it from leading to blindness S urgery for cicatricial entropion and trichiasis (Tertiary prevention) A ntibiotics to eliminate the organism (Secondary prevention) F acial cleanliness to avoid infection (Primary prevention) E nvironmental improvement with better hygiene and sanitation (Primordial prevention) Surgery (Tertiary prevention) (Option B) Epilation/ Bilamellar tarsal rotation Antibiotics (Secondary prevention) (Option A) For areas with 10% or more prevalence of active trachoma (TF in children 1-9 yrs): Azithromycin single oral dose 20 mg/kg to all community members 5% to 10% prevalence: Antibiotic therapy targeted in the family and close contact only Less than 5% prevalence: Only patients are treated Facial cleanliness (Primary prevention) (Option D) Frequent face wash with clean water, Avoid common towels, handkerchiefs, Surma rods, etc Environmental improvement (Primordial prevention) (Option C) Water latrines and good water supply to reduce flies, Sprays to control flies, Health education

References:

Kanski’s Clinical Ophthalmology, 9th edition, Page- 176

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Which organism is most likely causing the infection in a 14-day-old neonate with bilateral conjunctivitis and lung infiltrates, as found in a chest X-ray performed at the outpatient department? (Q855386)

A. Streptococcus

B. Hemophilus influenzae

C. Chlamydia trachomatis

D. Mycoplasma

Correct Answer: C. Chlamydia trachomatis

Explanation: A 14-day-old neonate with Bilateral conjunctivitis and pneumonia-like presentation is suggestive of Chlamydia trachomatis infection. Neonatal conjunctivitis is also known as Ophthalmia neonatorum. Ophthalmia neonatorum: Definition Ophthalmia neonatorum or neonatal conjunctivitis is a mucoid, mucopurulent, or purulent discharge from one or both eyes in the first month of life. Any discharge, even watery, from a baby's eyes in the first week of life must be viewed under suspicion as tears are not secreted so early in life. Aetiology of ophthalmia neonatorum Mode of infection: Before birth: Very rare through infected liquor amnii During birth: From an infected birth canal After birth: During the first bath of a newborn Soiled clothes Fingers with infected lochia Source of infection: Neisseria gonorrhoeae: Transmitted at the time of delivery Presents between 2-5 days of birth (Option B) The discharge becomes mucopurulent and purulent quickly Thick yellow pus and marked chemosis Cornea seen at the bottom of a crater-like pit in severe cases Chlamydia trachomatis: Transmitted at the time of delivery Onset at 5-14 days after birth (Option C) Serotypes D to K are the most common cause of ophthalmia neonatorum in developed countries Characteristic intracellular inclusion bodies are seen Less severe inflammation Purulent discharge + Other causes: Staphylococci/ Streptococcus pneumoniae (Option A) Gram-negative coliforms Herpes simplex virus Chemical: silver nitrate solution previously used for prophylaxis immediately after birth : erythema, chemosis, lid edema, and discharge within the first 24 hours after birth. Prophylaxis Antenatal measures: Treatment of genital infections and care of mother Surveillance of pregnant women for STDs Natal measures: Aseptic, sterile precautions to be followed during deliveries Clean the infant's closed lids with sterile cotton soaked in normal saline and dried Post-natal measures: Povidone-iodine 2.5% solution If the mother is under suspicion, 1% tetracycline or 0.5% erythromycin eye ointment is used in the baby immediately after birth and watched for a week. Ceftriaxone 50mg/kg IM or IV single dose for infants born to mothers with untreated gonococcal infection. Crede's method: Old method using 1% silver nitrate applied to infants' eyes right after birth It is NOT used now as it can cause chemical neonatal conjunctivitis. Diagnosis Treatment Neisseria gonorrhoea Ceftriaxone IM, gentamicin drops, bacitracin eye ointment Chemical Wash eyes, erythromycin ointment, observe Other bacteria Neomycin-bacitracin ointment, gentamicin, or tobramycin drops Herpes simplex virus Acyclovir 3% eye ointment, systemic acyclovir (systemic involvement) Chlamydia trachomatis Erythromycin or chlortetracycline eye ointment, oral erythromycin (systemic infection) Incorrect options: Streptococci (Option A) and Mycoplasma (Option D): Though they can cause conjunctivitis and pneumonia, it is very rare in the case of a newborn.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: What type of visual impairment does a person have if their vision is less than 3/60 but greater than 1/60 in their better eye, and their field of vision is limited to less than 10 degrees from the center of fixation? (Q857712)

A. Curable blindness

B. Economic blindness

C. Social blindness

D. Manifest blindness

Correct Answer: C. Social blindness

Explanation: Vision < 3/60 but > 1/60 with field of vision < 10 degrees is suggestive of Social Blindness . NPCB Classification of Visual Impairment The National Programme for Control of Blindness (NPCB) in India defines blindness as " visual acuity of <3/60 in a better eye with best possible correction " and central visual field <10 degrees . The NPCB further categorizes visual impairment as: Classification Visual Acuity Visual Impairment Visual acuity of less than 6/18 in the better eye with available correction Low Vision Visual acuity of less than 6/18 to 6/60 Economic Blindness (Option B) Visual acuity of less than 6/60 to 3/60 Social Blindness (Option C) Visual acuity of less than 3/60 to 1/60 Manifest Blindness (Option D) Visual acuity of less than 1/60 to the perception of light Absolute Blindness No perception of light WHO Classification of Visual Impairment The World Health Organization (WHO) defines blindness as " visual acuity of less than 3/60 (Snellen) or its equivalent ". The International Classification of Disease 11 (2018) classifies vision impairment into distance and near-presenting vision impairment Classification Visual Acuity Mild Distance Vision Impairment Presenting visual acuity worse than 6/12 Moderate Distance Vision Impairment Presenting visual acuity worse than 6/18 Severe Distance Vision Impairment Presenting visual acuity worse than 6/60 Blindness Presenting visual acuity worse than 3/60 Near Vision Impairment Presenting near vision acuity worse than N6 or M0.8 at 40 cm with existing correction

References:

Park’s Textbook of Preventive and Social Medicine 27th edition, pages 456,457

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: Which is the most common cause of preventable childhood blindness in India? (Q857231)

A. Refractive error


B. Vitamin A deficiency

C. Retinopathy of prematurity

D. Cataract

Correct Answer: B. Vitamin A deficiency

Explanation: The most common cause of preventable childhood blindness is vitamin A deficiency leading to Xerophthalmia. It can be prevented by timely vitamin A prophylaxis. Blindness in India: Overall most common cause of blindness/visual impairment (VI): Cataract Most common preventable cause of blindness: Vitamin A deficiency Most common cause of blindness in children: Vitamin A deficiency Most common cause of VI in children: Refractive error Other causes of blindness in children: Trachoma, Retinopathy of prematurity; Congenital cataract; Ophthalmia neonatorum. Proposed structure for Vision 2020, NPCB: Tier Number of Units Population Norms Components Centres of Excellence 20 1/50 million Professional Leadership Strategy development Continued Medical Education (CME) Laying of standards and quality assurance Research Training Centres 200 1/5 million Tertiary eye care including retinal surgery, corneal transplantation, glaucoma surgery, etc. Training and CME Service Centres 2000 1/500000 Cataract Surgery Other common eye surgeries Facilities for refraction Referral services Vision Centres 20,000 1/50000 Refraction and prescription of glasses Primary eye care School eye screening programme Screening and referral services



References:

Park’s Textbook of Preventive and Social Medicine 27th edition, page 505.

AK Khurana Comprehensive Ophthalmology 7th edition, pages 499-501

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: According to the National Blindness and Visual Impairment Survey conducted from 2015 to 2019, what is the leading cause of moderate to severe visual impairment? (Q855573)

A. Refractive error

B. Untreated cataract

C. Glaucoma

D. Diabetic retinopathy

Correct Answer: B. Untreated cataract

Explanation: According to the National Blindness and Visual Impairment Survey (2015-2019) , untreated cataract remains the leading cause of moderate to severe visual impairment . Cataracts, which cause the lens of the eye to become cloudy and impair vision, are particularly prevalent in older adults. When left untreated, cataracts can lead to significant visual impairment and blindness, making it a major contributor to visual disability worldwide. Major Causes of Blindness in population aged ≥ 50 years Cataract - 66.2% Corneal opacity (including trachomatous) Cataract surgical complications (including PCO) Posterior segment disease (excluding DR & ARMD) (Option D ruled out) Glaucoma (Option C ruled out) Major Causes of Visual Impairment in population aged ≥ 50 years Cataract - 71.2% (Option B) Refractive error (Option A ruled out) Cataract surgical complications (including PCO) Major Causes of Blindness in population aged 0-49 years Corneal opacity - 37.5 All globe/ CNS abnormality (Amblyopia) Phthisis Other/undetermined Major Causes of Visual Impairment in population aged 0-49 years Refractive error - 29.6 Cataract All globe/ CNS abnormality (Amblyopia) Corneal opacity

References:

National Blindness and Visual Impairment Survey

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: How does the World Health Organization (WHO) define blindness? (Q860171)

A. VA < 3/60

B. VA < 6/60

C. VA > 3/60

D. VA > 6/60

Correct Answer: A. VA < 3/60

Explanation: NPCB Classification of Visual Impairment The National Programme for Control of Blindness (NPCB) in India defines blindness as "visual acuity of <3/60 in a better eye with best possible correction" and central visual field <10 degrees. The NPCB further categorizes visual impairment as: Classification Visual Acuity Visual Impairment Visual acuity of less than 6/18 in the better eye with available correction Low Vision Visual acuity of less than 6/18 to 6/60 Economic Blindness Visual acuity of less than 6/60 to 3/60 Social Blindness Visual acuity of less than 3/60 to 1/60 Manifest Blindness Visual acuity of less than 1/60 to the perception of light Absolute Blindness No perception of light WHO Classification of Visual Impairment The World Health Organization (WHO) defines blindness as "visual acuity of less than 3/60 (Snellen) or its equivalent". (Option A) The International Classification of Disease 11 (2018) classifies vision impairment into distance and near-presenting vision impairment Classification Visual Acuity Mild Distance Vision Impairment Presenting visual acuity worse than 6/12 Moderate Distance Vision Impairment Presenting visual acuity worse than 6/18 Severe Distance Vision Impairment Presenting visual acuity worse than 6/60 Blindness Presenting visual acuity worse than 3/60 Near Vision Impairment Presenting near vision acuity worse than N6 or M0.8 at 40 cm with existing correction Option B,C and D are incorrect.

References:

Park’s Textbook of Preventive and Social Medicine, 27th edition, pg no-456-457

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: Which of the following statements about the instrument shown in the image given below is false? {{caption_text}} (Q860010)

A. Used to measure the curvature of the cornea

B. Assess corneal astigmatism.

C. Depends on Purkinje image I.

D. Measures intraocular pressure.



Correct Answer: D. Measures intraocular pressure.

Explanation: The instrument shown in the image is a Keratometer used to assess the corneal curvature responsible for causing astigmatism. It is based on the size of the image formed by the anterior surface of the cornea i.e., purkinje image I. Intraocular pressure is measured by using a tonometer. Keratometry An objective method for measuring the curvature of the central cornea (Option A ruled out) to estimate corneal astigmatism (Option B ruled out) . Principle of Keratometer: Measures corneal curvature by analyzing the size of the image formed by the anterior corneal surface (Option C ruled out), utilizing visible doubling. Normal Keratometric Values: Average: 45 D Keratoconus >45 D Types of Keratometers: Javal-Schiotz Model Bausch & Lomb Model {{caption_text}} Auto refractive Keratometer



References:

Corneal Imaging - StatPearls - NCBI Bookshelf.

Kanski’s Clinical Ophthalmology, Edition 9, Page- 209.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: What is considered the most reliable tonometer in the medical field? (Q860008)

A. Schiotz

B. Goldmann applanation tonometer


C. Tonopen


D. Pascal dynamic contour

Correct Answer: B. Goldmann applanation tonometer

Explanation: Types of Tonometry: 1. Indentation Tonometry: Concept: Measures IOP by how much a plunger indents the eye; softer eyes indent more than harder eyes. Schiotz Tonometer (Option A): Most widely used

tonometer in the world. Advantages: Inexpensive, simple, and reliable. Disadvantages: Gives false readings in eyes with abnormal scleral rigidity (e.g., high myopes, post-surgery). {{caption_text}} 2. Applanation Tonometry: Concept: Measures IOP based on the force needed to flatten the cornea's surface (Imbert-Fick Law: P = W/A). Goldmann Tonometer (Option B): Most popular and accurate; uses a double prism on a slit lamp. Technique: Cornea is anesthetized and stained with fluorescein. Cornea and biprisms are illuminated with cobalt blue light. The prism touches the cornea, and the inner edges of two semicircles are adjusted to just touch. IOP is read by multiplying the dial reading by ten. {{caption_text}} Variable force tonometers: Goldmann applanation tonometer → Gold standard Perkins tonometer (hand held version of goldmann) Draeger tonometer (measures wide range of IOP) Draeger tonometer is similar to Perkins It has a different set of prisms It operates with a motor Pulsair (non-contact tonometer) Airpuff (non-contact tonometer) Grolman (non-contact tonometer) Tonopen (based on both indentation and applanation) MacKay Marg (based on both indentation and applanation) Rebound tonometer Transpalpebral Tonometer Fixed force tonometer: Maklakov tonometer Barraquer tonometer Tonopen (Option C) is a handheld electronic tonometer that measures IOP by using both applanation and indentation method.. Pascal dynamic contour (Option D) tonometer is a newer technology that measures IOP by analyzing the contour of the cornea during gentle deformation.





References:

Kanski's Clinical Ophthalmology: A Systematic Approach, 9th Edition, Page 29, 30

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: All of the following are functions of the superior oblique muscle except? (Q853122)

A. Adduction

B. Abduction


C. Depression

D. Intorsion

Correct Answer: A. Adduction

Explanation: Superior oblique muscle actions: Primary: Intorsion (internal rotation) (Option D ruled out) Secondary: Depression (mainly when eye is adducted) (Option C ruled out) Tertiary: Abduction (outward

movement) (Option B ruled out) Superior rectus (SR) Medial rectus (MR) Inferior rectus (IR) Lateral rectus (LR) Superior oblique (SO) Inferior oblique (IO) Elevation Adduction Intorsion Adduction Depression Adduction Extorsion Abduction Depression (Option C) Abduction (Option B) Intorsion (Option D) Elevation Abduction Extorsion



References:

Grey's Anatomy 42nd edition, pg 773-777. BD Chaurasia’s Human Anatomy, 8th edition, pg 232-237. Vishram Singh Textbook of Anatomy, 2nd edition, pg 285-288.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: Which of the following statements is correct regarding indirect ophthalmoscopy? (Q852502)

A. The image is erect and real.

B. The condenser lens needs to be kept close to the examiner’s eye

C. It provides 15x magnification.

D. It provides 3-5x magnification.

Correct Answer: D. It provides 3-5x magnification.

Explanation: Indirect ophthalmoscopy is a binocular technique that provides a real, inverted, and reversed image of the retina with moderate magnification (3–5×) and a wide field of view (about 6–8 disc diameters). The condenser lens (commonly +20D) should be held close to the patient's eye , not near the examiner's eye. Holding it too far away reduces image clarity and field of view Key Features of Indirect Ophthalmoscopy: Feature Description Image Real, inverted, and laterally reversed Magnification 3–5× Field of view Wide (6–8 disc diameters) Lens position Held close to patient's eye (~1–2 cm) Depth perception Present (binocular view) Best for Detecting retinal detachment, tears, and peripheral lesions

References:

Myron Yanoff and Duker's textbook of ophthalmology, 6th edition, Page no: 52-53 Kanski's clinical ophthalmology, John F. Salmon, 10th edition, Page no: 21-23

↑ *Back to this question in Part A*

@Neet_pg_bot


Q15: What does the picture depict in terms of the investigation? (Q859780)

A. Fundus fluorescein angiography

B. Optical coherence tomography

C. Retinal CT

D. Retinal MRI



Correct Answer: A. Fundus fluorescein angiography

Explanation: Fundus fluorescein angiography: Principle: Based on the detection of fluorescent light emitted by fluorescein dye in circulation. Absorption & Emission Wavelengths: Irradiating light: 420–490 nm. Emitted fluorescence: 510–530 nm. Retinal pigments and red blood cells can absorb fluorescent light, masking deeper structures. Pigment migration allows visibility of deeper layers. Diagnostic uses: Diabetic Retinopathy: Reveals neovascularization, and assesses vascular tree integrity. Fundus Disorders: Identifies depth of pathological involvement. Optic Nerve Pathologies: Useful in detecting optic nerve issues such as papilloedema. Neovascularization of Iris: Detects leakage from iris vessels, an early sign of rubeosis. {{caption_text}} Normal Study Abnormalities Detected by Fundus Fluorescein Angiography (FFA) Fluorescence Type Cause Examples Hyperfluorescence Window defect in retinal pigment epithelium (RPE) Atrophy showing background choroidal fluorescence Pooling of dye under detached RPE Age-related macular degeneration (ARMD) Pooling of dye under sensory retina Central serous retinopathy (CSR) Leakage into the neurosensory retina due to the breakdown of the inner blood-retinal barrier Cystoid macular edema (CME) Leakage from choroidal or retinal neovascularization Proliferative diabetic retinopathy, subretinal neovascular membrane in ARMD Staining (long retention of dye by tissues) Drusen, normal optic disc Leakage from optic nerve head Papilloedema Hypofluorescence Blockage of background fluorescence Retinal hemorrhage, hard exudates, pigmented clumps Occlusion of retinal/choroidal vasculature Central retinal artery occlusion , diabetic retinopathy capillary occlusion Loss of vasculature Choroideremia, myopic degeneration Incorrect Options: Optical Coherence Tomography (OCT) (Option B) is a non-invasive imaging technique that uses light waves to produce cross-sectional images of the retina.It is particularly useful in assessing conditions like macular edema, macular holes, and retinal detachment. Retinal CT (Option C) is not a recognized imaging modality in ophthalmology, as computed tomography (CT) is generally not suitable for the detailed visualization of retinal layers. Retinal MR (Option D) can be used to assess orbital or optic nerve conditions, it does not provide the level of detail required for retinal imaging



References:

Parson’s Diseases of the Eye, 23rd edition, Pg 315-318

Kanski’s Clinical Ophthalmology,Edition 9, Page 442-447

Yanoff and Duker Ophthalmology,Edition 5, Page 558-563

↑ *Back to this question in Part A*

@Neet_pg_bot

Q16: A 6-year-old child presents with a refractive error of -2D in the right eye and +1D in the left eye. Fundus examination reveals normal visual acuity and retinal findings. What is the most likely diagnosis? (Q852481)

A. Myopia

B. Hyperopia

C. Anisometropia

D. Amblyopia

Correct Answer: C. Anisometropia

Explanation: Anisometropia is a condition where there is a significant difference in the refractive power between the two eyes. In this case: Right Eye (RE): -2D (Total refractive power: 62D → 60 + 2D) Left Eye (LE): +1D (Total refractive power: 59D → 60 - 1D) Difference in refractive power = 62D - 59D = 3D, which is greater than 2.5D, making it clinically significant anisometropia. Anisometropia: is a term used when the dioptric power of both eyes is not the same, and is significant if > 2.5D difference. Myopia (Option A): Myopia (nearsightedness) is present in the right eye (-2D), but the left eye has hyperopia (+1D). The diagnosis should reflect the difference in refractive power rather than just the presence of myopia. Hyperopia (Option B): Hyperopia (farsightedness) is present in the left eye (+1D), but since the right eye has myopia, this case is best classified as anisometropia rather than simple hyperopia. Amblyopia (Option D): Amblyopia (lazy eye) can develop due to anisometropia if left untreated, but the question does not indicate decreased visual acuity. Therefore, anisometropia is the more accurate primary diagnosis at this stage.

References:

Parsons Diseases of the eye, 23rd Edition, Pg 78

↑ *Back to this question in Part A*

@Neet_pg_bot

Q17: A patient presents with right-sided field defects in both eyes, but central vision remains unaffected. What is the most likely diagnosis? (Q852486)

A. Homonymous Hemianopia with Macular Sparing

B. Heteronymous Hemianopia with Central Sparing


C. Optic Chiasm Lesion

D. Optic Tract Lesion

Correct Answer: A. Homonymous Hemianopia with Macular Sparing


Explanation: Homonymous Hemianopia: T his refers to the loss of the same side of the visual field in both eyes. In this case, the patient has a right-sided field defect in both eyes, which suggests a lesion in the contralateral visual pathway, most likely in the right occipital lobe or along the visual tract. Macular Sparing: The macula (central vision) is often spared in cases of cortical lesions, such as those caused by a stroke in the Posterior Cerebral Artery (PCA) territory. The macula receives a dual blood supply from both the PCA and the Middle Cerebral Artery (MCA). If the PCA is affected (for example, in a PCA stroke), but collateral circulation from the MCA is maintained, the macular representation in the visual cortex can still receive enough blood supply to preserve central vision. This phenomenon explains macular sparing in cortical lesions Fig: Homonymous hemianopia with macular sparing. Visual Pathway Lesions and Defects Site of Lesion Fibers Affected Field Defect Optic Nerve Ipsilateral Temporal + Nasal Fibers Anopia Optic Chiasma Bilateral Nasal Fibers Bitemporal Hemianopia (Heteronymous Hemianopia) Optic Tract Lateral Geniculate Body (LGB Lesion) Optic Radiation Ipsilateral

Temporal Fibers + Contralateral Nasal Fibers Homonymous Hemianopia Visual Cortex without the involvement of tip (Post. Cerebral Artery) Ipsilateral Temporal Fibers + Contralateral Nasal Fibers except macular representation Homonymous Hemianopia with Macular Sparing Tip of Visual Cortex (Middle Cerebral Artery) Macular Representation of Ipsilateral Temporal and Contralateral Nasal Fibers. Macular Homonymous Hemianopia







@Neet_pg_bot




@Neet_pg_bot



@Neet_pg_bot



@Neet_pg_bot



References:

Parsons Diseases of the eye, 23rd Edition, Pg - 1184 Yanoff & Duker Ophthalmology, 5th edition, Pg - 913,919

↑ *Back to this question in Part A*

@Neet_pg_bot

Q18: The person in the image is at the risk of developing? (Q856126)

A. Exposure keratopathy

B. Restricted eye movements

C. Diplopia

D. Amblyopia



Correct Answer: A. Exposure keratopathy

Explanation: ongenital eye malformations caused by incomplete closure of the embryonic cleft , leading to underdeveloped or missing tissues in the uvea, retina, or extending onto the back of the iris. Types of Coloboma: Typical: Found along the inferior aspect of the eye structures, including the iris, lens, choroid, retina, and optic nerve. Atypical: Occurs in the non-inferior location of the eye, like the upper iris or peripheral retina, which are not along the path of the embryonic fissure. Types of Coloboma and their characteristics: Iris Coloboma Pear-shaped defect extending from the pupil but not always reaching the ciliary body. It may occur alone or be associated with colobomata of the lens, choroid, or retina. {{caption_text}} Coloboma of lens Notch-shaped defect typically found along the inferior lens margin due to incomplete development of the suspensory ligament. {{caption_text}} Fundus Coloboma Oval or comet-shaped defect with a rounded apex toward the optic disc, often with an irregularly depressed surface. {{caption_text}} Coloboma of Optic Disc It appears as an inferior crescent at the lower edge of the disc due to incomplete closure of the embryonic fissure. Found in Hypermetropic and Astigmatic eyes. {{caption_text}} Coloboma of Lid Notch or gap in the eyelid edge, most commonly on the upper lid, often associated with iris coloboma or accessory auricles. The corneal surface cannot stay lubricated and hence can lead to exposure keratopathy. {{caption_text}} Neurotrophic and Neuroparalytic Keratopathy: Neurotrophic Keratopathy: Caused by trigeminal nerve paralysis. Decreased corneal sensation and epithelial desquamation. May follow procedures like Gasserian ganglion removal or blockage, often used in treating trigeminal neuralgia. Neuroparalytic Keratopathy: This condition occurs due to facial nerve paralysis, leading to lagophthalmos (inability to close the eye) and subsequent corneal exposure. The patient's history does not suggest facial nerve paralysis, making this less likely.


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot



References:

Parsons Diseases of the eye 23rd Edition, Pages 604,605,643,1094

↑ *Back to this question in Part A*

@Neet_pg_bot

Q19: Which of the subsequent options aids in distinguishing a fixed dilated pupil? Phenylephrine 1% Pilocarpine 0.1% Pilocarpine Cocaine 0.5% Epinephrine (Q856408)

A. 1,2,3,4 and 5


B. 2 and 3

C. 1,4and 5

D. 1,3 and 4

Correct Answer: B. 2 and 3

Explanation: To differentiate a fixed dilated pupil, the use of specific pharmacological agents can be helpful. Pilocarpine is a parasympathomimetic drug that stimulates the pupillary sphincter muscle, causing miosis (constriction of the pupil). If the dilated pupil responds to pilocarpine and constricts, it indicates that the dilated pupil is likely due to parasympathetic denervation or dysfunction. Normal pupil react to 1% pilocarpine below this concentration, normal pupil don't react.



References:

Yanoff & Duker Ophthalmology, 5th edition, Pg 965,966,968

↑ *Back to this question in Part A*

@Neet_pg_bot

Q20: Which of the following combinations of cylindrical lenses can be employed for the treatment of astigmatism in the with-the-rule orientation? (Q793355)

A. +5.00 DC x 180, -3.00 DC x 90

B. +7.00 DC x 180, +5.00 DC x 90

C. -4.00 DC x 180, -5.00 DC x 90

D. -3.00 DC x 180, +1.00 DC x 90

Correct Answer: D. -3.00 DC x 180, +1.00 DC x 90

Explanation: Correct Option D: In with-the-rule (WTR) astigmatism, more refractive power (and curvature) along the vertical meridian than the horizontal. A concave cylinder at 180 ± 20 degrees or a convex cylinder at 90 ± 20 is required for this kind of astigmatism. -3.00 DC x 180, +1.00 DC x 90 - This combination of lenses corrects with-the-rule astigmatism. The powers are reasonable for a typical correction. Incorrect Options: Option A. +5.00 DC x 180, -3.00 DC x 90: This combination of lenses corrects against-the-rule astigmatism. It cannot be used to correct with-the-rule astigmatism. Option B. +7.00 DC x 180, +5.00 DC x 90: This combination of lenses corrects against-the-rule astigmatism. It cannot be used to correct with-the-rule astigmatism. Option C. -4.00 DC x 180, -5.00 DC x 90: This combination of lenses corrects with-the-rule astigmatism. However, the powers are too high for a typical correction.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q21: Which of the following is true about viscoelastic substances? (Q853141)

A. Cause mechanical injury to corneal endothelium B) Prevents Anterior Chamber from collapsing during crucial steps like capsulorrhexis and IOL placement C) It is inert and can be left behind after surgery D) Silicone oil, perfluoropropane, and sodium hexafluoride are most commonly used substances

B. Prevents Anterior Chamber from collapsing during crucial steps like capsulorrhexis and IOL placement

C. It is inert and can be left behind after surgery

D. Silicone oil, perfluoropropane, and sodium hexafluoride are most commonly used substances

Correct Answer: B. Prevents Anterior Chamber from collapsing during crucial steps like capsulorrhexis and IOL placement

Explanation: Viscoelastic substances are ophthalmic agents used in anterior segment surgeries like cataract surgery (phacoemulsification) to maintain anterior chamber stability, protect intraocular tissues, and facilitate intraocular lens (IOL) implantation. Cause mechanical injury to corneal endothelium (Option A): Viscoelastic substances are protective, not damaging, to the corneal endothelium. Some high-viscosity viscoelastics (e.g., Healon GV) can increase intraocular pressure if not completely removed, but they do not cause direct mechanical injury. It is inert and can be left behind after surgery (Option C): Viscoelastic substances are not inert and must be removed at the end of surgery to prevent postoperative intraocular pressure (IOP) elevation. Residual viscoelastics can obstruct aqueous outflow, leading to IOP spikes. Silicone oil, perfluoropropane, and sodium hexafluoride are most commonly used substances (Option D): Silicone oil, perfluoropropane (C3F8), and sulfur hexafluoride (SF6) are not viscoelastic substances; they are tamponade agents used in retinal surgeries (e.g., retinal detachment repair). Commonly used ophthalmic viscoelastic devices (OVDs) include: Sodium hyaluronate (Healon, ProVisc) Hydroxypropyl methylcellulose (HPMC) Chondroitin sulfate-based OVDs (Viscoat)

References:

Viscoelastics - StatPearls - NCBI Bookshelf

Agents for Vitreous Tamponade - StatPearls - NCBI Bookshelf

↑ *Back to this question in Part A*

@Neet_pg_bot

Q22: In Fundus Fluorescein Angiography (FFA), how do microaneurysms appear? (Q853133)

A. Hypofluorescent dots


B. Hyperfluorescent dots

C. Hypofluorescent line

D. Hyperfluorescent branching-like structure in the vessel

Correct Answer: B. Hyperfluorescent dots

Explanation: Microaneurysms are the earliest clinically detectable lesions in diabetic retinopathy and appear as small, round dilations of retinal capillaries. In Fundus Fluorescein Angiography (FFA), they exhibit hyperfluorescence due to leakage of fluorescein dye from these abnormal capillaries. Microaneurysms initially appear as hyperfluorescent dots in early-phase FFA due to accumulation of fluorescein dye. In later phases, leakage may lead to diffuse hyperfluorescence in severe cases. Fundus Fluorescein Angiography Principle: Based on the detection of fluorescent light emitted by fluorescein dye in circulation. Absorption & Emission Wavelengths: Irradiating light: 420–490 nm. Emitted fluorescence: 510–530 nm. Retinal pigments and red blood cells can absorb fluorescent light, masking deeper structures. Pigment migration allows visibility of deeper layers. Diagnostic uses: Diabetic Retinopathy: Reveals neovascularization, and assesses vascular tree integrity. Fundus Disorders: Identifies depth of pathological involvement. Optic Nerve Pathologies: Useful in detecting optic nerve issues such as papilloedema. Neovascularization of Iris: Detects leakage from iris vessels, an early sign of rubeosis. Normal Study Hypofluorescent dots (Option A): Microaneurysms do not appear hypofluorescent. Hypofluorescent lesions are usually blockages (e.g., hemorrhages, pigment deposits) or vascular occlusions. Hypofluorescent line (Option C): Hypofluorescent lines are seen in conditions like vascular occlusions (e.g., branch retinal artery occlusion). Hyperfluorescent branching-like structure in the vessel (Option D): This pattern is seen in neovascularization, not microaneurysms.



References:

Parson’s Diseases of the Eye, 23rd edition, Pg 315-318

Kanski’s Clinical Ophthalmology,Edition 9, Page 442-447

Yanoff and Duker Ophthalmology,Edition 5, Page 558-563

↑ *Back to this question in Part A*

@Neet_pg_bot

Q23: All of the following are features of oculomotor nerve palsy except? (Q853124)

A. Exotropia


B. Dilated pupil

C. Limited abduction

D. Ptosis

Correct Answer: C. Limited abduction

Explanation: Limited abduction (inability to move the eye outward) is a feature of abducens nerve palsy. This movement is controlled by the lateral rectus muscle, which is innervated by the abducens nerve (cranial nerve VI). Functions of muscle supplied by oculomotor nerve: Superior rectus (SR) Medial rectus (MR) Inferior rectus (IR) Levator palpebrae superioris (LPS) Inferior oblique (IO) Elevation Adduction Intorsion Adduction (Option A ruled out) Depression Adduction Extorsion Elevation of superior eyelid (Option D ruled out) Elevation Abduction Extorsion Oculomotor nerve (cranial nerve III) Type Motor Origin Midbrain (Oculomotor sulcus on the medial aspect of the cerebral peduncle). Course Components 1. Oculomotor nucleus (somatic component): General somatic efferent (GSE) fibres emerge from this site and supply - Superior division : Levator palpebrae superioris Superior rectus Inferior division : Medial rectus Inferior rectus Inferior oblique 2.Edinger-Westphal nucleus (parasympathetic component) (Option C ruled out) : General visceral efferent (GVE) fibres arise from this site and supply: Sphincter pupillae- causes pupil constriction Ciliaris Note: The nerve fibers innervating the pupillary muscles are located superficially in the nerve trunk and pial blood vessels supply these fibers. The main trunk of the fibers is supplied by the vasa vasorum. Diabetic and hypertensive microangiopathy will affect the vasa vasorum and thus spare the pupillary fibers. Hence, pupil-sparing 3rd nerve lesions are seen with ischemic lesions such as diabetes, hypertension, or arteriosclerotic disease.



Arises from the midbrain
(oculomotor sulcus on the medial aspect of the cerebral peduncle)

Inter-peduncular fossa

Runs forward and laterally between the **posterior cerebral and superior cerebellar arteries,** lateral to the posterior communicating artery.

Tentorial notch of tentorium cerebelli

Middle cranial fossa

Pierces the dura mater in the **oculomotor triangle**


Cavernous sinus
(cavernous sinus, lying above the trochlear, ophthalmic, and maxillary nerves, and lateral to the internal carotid artery)

Superior orbital fissure
- Small upper branch
- Large lower branch


Orbit
Small upper branch supplies :
- Levator palpebrae superioris (half)
- Superior rectus

Large lower branch supplies :
- Medical rectus
- Inferior rectus
- Inferior oblique

@Neet_pg_bot



@Neet_pg_bot



References:

Grey's Anatomy 42nd edition, pg 458, 781. BD Chaurasia’s Human Anatomy, 8th edition, pg 69, 70, 240, 241. Vishram Singh Textbook of Anatomy, 2nd edition, pg 335, 336.
https://www.ncbi.nlm.nih.gov/books/NBK526112/

↑ *Back to this question in Part A*

@Neet_pg_bot

• [13_Sclera] 028 Sclera

« Back to Folder Directory

[13_Sclera] 028 Sclera — Quick Jump

Q1 (p. 1494) *Q2 (p. 1495)* *Q3 (p. 1496)* *Q4 (p. 1498)* *Q5 (p. 1499)* *Q6 (p. 1500)*
Q7 (p. 1503) *Q8 (p. 1505)* *Q9 (p. 1506)* *Q10 (p. 1508)* *Q11 (p. 1509)* *Q12 (p. 1509)*
Q13 (p. 1511) *Q14 (p. 1512)* *Q15 (p. 1515)*

Q1: Which of the following correctly lists the layers of the sclera from outermost to innermost? (Q853641)

A. Lamina fusca > Sclera proper > Episclera

B. Episclera > Sclera proper > Lamina fusca

C. Sclera proper > Episclera > Lamina fusca


D. Episclera > Lamina fusca > Sclera proper

Correct Answer: B. Episclera > Sclera proper > Lamina fusca

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The sclera consists of three ill-defined layers, namely the sclera proper with the episclera on the outside and the lamina fusca interiorly . Layers of the Sclera: Layer Description Composition Remarks Episclera Thin vascularized. connective tissue. Collagen fibers Fibroblasts Blood vessels Nerves Located just outside the sclera proper Sclera proper Relatively avascular. Dense tissue. Dense irregular bundles of collagen (compared to the orderly arrangement in the cornea) Elastic fibers Fibroblasts, Proteoglycans The thickest layer of the sclera Provides structural support and integrity Lamina fusca Innermost layer. Loose connective tissue layer. Melanocytes (migrated from choroid) It contains pigmented cells derived from the choroid



@Neet_pg_bot

References:

Parson’s Diseases of the eye, 23rd Edition, Page 198.

Kanski’s Clinical Ophthalmology, 9th Edition, Page 292

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Match the scleral apertures with the structures they transmit Scleral Apertures Structures Transmitted A. Anterior scleral apertures 1. Short and long ciliary arteries B. Middle scleral apertures 2. Central retinal artery and vein C. Posterior scleral apertures 3. Anterior ciliary arteries and veins D. Lamina cribrosa 4. Vortex veins (Q853642)

A. A-3, B-4, C-1, D-2

B. A-2, B-1, C-4, D-3

C. A-3, B-2, C-1, D-4


D. A-3, B-1, C-4, D-2

Correct Answer: A. A-3, B-4, C-1, D-2

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Apertures of the sclera: Apertures Structures transmitted Anterior scleral apertures Anterior ciliary vessels, nerves, and lymphatics Middle scleral apertures 4 vortex veins (Vena verticose) Posterior scleral apertures 8 to 20 short ciliary vessels (circle of Zinn) and nerves 2 long posterior ciliary arteries anterior to short ciliary Lamina cribrosa Optic nerve fibers Central retinal artery and veins.



References:

Parson’s Diseases of the Eye, 23rd Edition, Page 198

↑ *Back to this question in Part A*

@Neet_pg_bot


Q3: A 28-year-old patient presents with painless redness of the eye. The ophthalmologist performed a diagnostic test using 2.5% phenylephrine, and no change was noticed. When 10% phenylephrine is used, the following change occurs. What is the most likely diagnosis? Before application of 10% phenylephrine: After application of 10% phenylephrine: (Q853643)

A. Conjunctivitis

B. Scleritis

C. Episcleritis

D. Uveitis





Correct Answer: C. Episcleritis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The patient has painless redness of the eye which blanches (as shown in the image) following the application of 10% phenylephrine, suggesting episcleritis . Principle of Phenylephrine Test Phenylephrine causes vasoconstriction of superficial conjunctival vessels. Deeper episcleral or scleral vessels do not respond to phenylephrine. Vascular plexuses Locations of Vessels Clinical correlation. Conjunctival plexus Most superficial Congested in conjunctivitis . Blanched with 2.5% phenylephrine Superficial episcleral plexus Lies at the level of Tenon's capsule. Radial configuration. Congested in episcleritis . Blanching does not occur with 2.5%, but occurs with 10% Deep vascular plexus Lies deep to the Tenon's capsule and over the sclera. Criss-cross pattern. Congested in scleritis . Does not blanch even with 10% phenylephrine. Uveitis (Option D) is associated with pain, redness, photophobia, and excess lacrimation, with associated findings in slit lamp examination, which are not seen here.

References:

Parson’s Diseases of the eye, 23rd Edition, Page 198.

Yanoff and Duker Ophthalmology, 5th Edition, Page 212

Kanski’s Clinical Ophthalmology, 9th Edition, Page 292

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 40-year-old patient presents with recurrent episodes of episcleritis. After taking a detailed history and performing a thorough systemic examination, which of the following investigations would you consider ordering? (Q853644)

A. Full blood count and rheumatoid factor (RF)

B. Mantoux test and antinuclear antibody (ANA)

C. VDRL and X-rays of the chest and sacroiliac joints

D. All of the above

Correct Answer: D. All of the above

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Episcleritis can be associated with systemic conditions such as rheumatoid arthritis (RF), systemic lupus erythematosus (ANA), and other systemic infections or inflammatory diseases that may be assessed with the Mantoux test (Tuberculosis) and VDRL (Syphilis). X-rays of the chest and sacroiliac joints can help rule out conditions like ankylosing spondylitis, which can also present with episcleritis. Therefore, a comprehensive

approach including all listed investigations is appropriate. Conditions associated with Episcleritis: Idiopathic : The exact etiology is unknown in many cases. Systemic Diseases : Gout Rosacea Psoriasis Connective tissue diseases (e.g., rheumatoid arthritis, lupus) Hypersensitivity Reactions : Endogenous tubercular toxins Streptococcal toxins Infectious Episcleritis : Herpes zoster virus Syphilis Lyme disease Tuberculosis

References:

Parson's Diseases of the eye, 23rd Edition, Page 200.

Kanski's Clinical Ophthalmology, 9th Edition, Page 292

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 45-year-old woman with a 15-year history of rheumatoid arthritis (RA) presents with severe ocular pain, redness, and progressively worsening vision in her left eye for the past 2 weeks. Examination reveals a localized area of scleral thinning with adjacent congestion. There is no posterior involvement on B-scan ultrasound. Which of the following is the most likely diagnosis? (Q853645)

A. Anterior staphyloma

B. Necrotizing scleritis with inflammation

C. Nodular scleritis


D. Scleromalacia perforans

Correct Answer: B. Necrotizing scleritis with inflammation

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Severe pain, scleral thinning with adjacent congestion are hallmark signs of necrotizing scleritis , the most severe form of RA-associated scleritis. RA leads to immune-mediated vasculitis , causing ischemic necrosis of the sclera . Classification of Scleritis: Anterior Staphyloma (Option A): Anterior staphyloma is a late-stage complication, where uveal tissue bulges through a thinned, scarred sclera. This patient has active inflammation, whereas staphyloma occurs after chronic necrosis and healing. Nodular Scleritis (Option C): Nodular scleritis presents with discrete, firm nodules on the sclera, without necrosis. This patient has scleral thinning and avascular necrosis, which rules out nodular scleritis. Scleromalacia Perforans (Option D): Scleromalacia perforans is a form of necrotizing scleritis WITHOUT inflammation (painless, non-red eye). This patient has severe pain and congestion, so active necrotizing scleritis is the correct choice.



References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 294

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 38-year-old male presents with persistent eye discomfort, redness, and watering. On examination, the left eye is diffusely tender, sclera appears uniformly reddened and swollen without localized thinning areas. There are no systemic signs of vasculitis or autoimmune disease on further investigation. Which type of anterior scleritis is most consistent with this presentation? (Q853646)

A. Necrotizing anterior scleritis with inflammation

B. Non-necrotizing anterior diffuse scleritis

C. Non-necrotizing anterior nodular scleritis


D. Scleromalacia perforans


Correct Answer: B. Non-necrotizing anterior diffuse scleritis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Clinical presentation of persistent eye discomfort, watering, diffuse redness, and tenderness without localized necrosis or thinning is characteristic of non-necrotizing anterior diffuse scleritis. Clinical classification: Non-necrotizing anterior scleritis: Pain, redness, watering, discharge, and bluish or brownish discoloration. Diffuse Nodular (Option C) : One or more dark red or bluish nodules, which later become purple and semitransparent like porcelain. Necrotizing anterior scleritis With inflammation Without inflammation {{caption_text}} {{caption_text}} Necrotizing anterior scleritis with inflammation■■■■■■■: (Option A) Severe form of scleritis Red, painful eye, and progressive worsening of symptoms {{caption_text}} Scleromalacia perforans (Option D) : Necrotizing, painless form seen in rheumatoid arthritis patients with scleral thinning {{caption_text}}








Necrotizing anterior scleritis
with inflammation

@Neet_pg_bot



References:

Parson’s Diseases of the eye, 23rd Edition, Pages 524-28

Scleritis - StatPearls - NCBI Bookshelf (nih.gov)

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 294, 295, 297

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 45-year-old woman presents with severe pain in her right eye, worsened by eye movements, and reduced vision. On examination, proptosis and restricted ocular movements were observed. Ophthalmoscopy shows disc, macular edema, and choroidal folds. Which of the following B-scan ultrasonography findings is most indicative of this disease? (Q853647)

A. T sign

B. Moth-Eaten Appearance

C. Sunray Sign


D. Kiss Sign


Correct Answer: A. T sign

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Given the clinical presentation of severe pain in the eye, proptosis, restricted ocular movements, reduced vision, disc, and macular edema, and the potential findings of choroidal folds, the diagnosis most consistent with these symptoms is posterior scleritis . Thickening of the posterior sclera, retrobulbar thickening, and fluid surrounding the optic nerve creates a characteristic “T” sign in B-scan ultrasound. In posterior scleritis, fluid accumulates in the sub-Tenon’s space due to inflammation. On B-scan, this fluid tracks along the optic nerve sheath. The horizontal line = scleral thickening + fluid The vertical line = optic nerve Together → form a “T-shaped” sign {{caption_text}} {{caption_text}} Moth-Eaten Appearance (Option B) : Seen in Retinoblastoma , characterized by intraocular calcifications , typically in children , not adults. Sunray Sign (Option C) : Seen in Choroidal Melanoma , due to radial vascularity , but this patient lacks an intraocular mass . Kiss Sign (Option D) : Seen in Total Retinal Detachment , where the detached retina folds inward , but this case has no retinal involvement .





References:

Parson’s Diseases of the eye, 23rd Edition, Page 528

www.ncbi.nlm.nih.gov/books/NBK499944/

www.researchgate.net/figure/B-scan-of-left-eye-showing-thickening-posteriorly-in-the-classic-T-sign-The-T-sign-is_fig2_26254046#:~:text=Download%20Scientific%20Diagram-,B%2Dscan%20of%20left%20eye%20showing%20thickening%20posteriorly%20in%20the,stem%20as%20the%20optic%20nerve.

www.ncbi.nlm.nih.gov/pmc/articles/PMC9128424/

↑ *Back to this question in Part A*

@Neet_pg_bot


Q8: A 60-year-old woman with a 20-year history of rheumatoid arthritis has a painless, progressively thinning area in the sclera of her left eye. The area appears as shown in the image below. Which of the following best describes the pathophysiological mechanism responsible for this condition? (Q853648)

A. Type III hypersensitivity

B. Type II hypersensitivity

C. Type IV hypersensitivity

D. Infective etiology




Correct Answer: A. Type III hypersensitivity

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The diagnosis in this scenario is Scleromalacia Perforans. It is usually associated with long-standing rheumatoid arthritis and presents with scleral thinning, appearing bluish and without signs of inflammation. The underlying cause of scleral thinning is due to Type III hypersensitivity caused by immune-mediated chronic granulomatous inflammation and chronic vaso-occlusion of episcleral vessels, leading to collagen necrosis. {{caption_text}} Type II hypersensitivity (Option B) involves antibody-mediated cellular dysfunction or destruction. Type IV hypersensitivity (Option C): Scleromalacia perforans is caused by immune complex–mediated vasculitis, not delayed-type hypersensitivity. Infective etiology (Option D): Scleromalacia perforans is a sterile, autoimmune complication of systemic diseases.



References:

Parson’s Diseases of the Eye, 23rd Edition, Page 527

eyewiki.org/Scleromalacia_Perforans

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Depending on the location of the staphyloma in the eye which among the following options is matched correctly with its cause? (Q853649)

A. Anterior staphyloma - Peripheral corneal ulcer

B. Intercalary staphyloma - Degenerative myopia

C. Equatorial staphyloma - Scleritis


D. Ciliary staphyloma - Pseudocornea

Correct Answer: C. Equatorial staphyloma - Scleritis

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The causes of equatorial staphyloma are scleritis , degenerative myopia, and chronic uncontrolled glaucoma. Staphyloma: It is ectasia of the outer coats of the eye (cornea, sclera, or both) with an incarceration of the uveal tissue. Types of staphyloma Causes Anterior (Option A) Due to corneal pathology (e.g., perforated corneal ulcer with iris prolapse forming pseudocornea) Intercalary (Option B) Perforating injuries of the peripheral cornea, marginal corneal ulcer, anterior scleritis, and scleromalacia perforans Ciliary (Option D) Glaucoma, scleritis, and trauma to the ciliary region Equatorial (approx 14 mm behind the limbus) Scleritis, degenerative myopia, and chronic uncontrolled glaucoma Posterior (Most common) Degenerative high axial myopia



References:

Parson’s Diseases of the eye, 23rd Edition, Page 531-533

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: Blue sclera is commonly associated with all of the following conditions except: (Q853650)

A. Osteogenesis Imperfecta

B. Ehlers-Danlos Syndrome

C. Alkaptonuria


D. Marfan Syndrome

Correct Answer: C. Alkaptonuria

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Yellow-brown or blue-black pigment deposition occurs in different parts of the eye in alkaptonuria. Classic scleral pigment deposits are called Osler's sign. Blue sclera: Uveal pigment shines through the thin sclera and produces the blue color Causes: Osteogenesis imperfecta (Option A) Ehlers–Danlos syndrome (Option B) Marfan syndrome (Option D) Pseudoxanthoma elasticum Scleromalacia perforans {{caption_text}}



References:

Parson's Diseases of the eye, 23rd Edition, Pages 534-35

Ocular Manifestations of Alkaptonuria - StatPearls - NCBI Bookshelf (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: Clinical question: A 60 y/o man from a tropical region presents with a painful, red eye and a corneal ulcer following recent exposure to decaying vegetable matter. Which is the most common fungal infection affecting the cornea? (Q853664)

A. Candida albicans

B. Fusarium

C. Mucormycosis

D. Aspergillus fumigatus

Correct Answer: D. Aspergillus fumigatus

Explanation: Aspergillus fumigatus is a common causative organism of keratomycosis in tropical and subtropical regions and is often associated with exposure to decaying organic matter. Keratomycosis Etiology Summary Most Common Causative Organisms : Aspergillus fumigatus : Most common in India and tropical regions. Fusarium (Option B ruled out) : Common in warm climates like Brazil. Candida albicans (Option A ruled out) : More frequent in temperate climates. Mode of Infection : Injury with Vegetative Material : Typically occurs after trauma with objects like thorns, wooden sticks, crop leaves, or decaying vegetable matter. Common in Field Workers : Especially during harvesting season in rural and tropical areas. Immunocompromised Individuals : Higher risk for developing fungal keratitis. Special Considerations : Candida albicans : Leading cause of endogenous endophthalmitis. Mucormycosis (Option C ruled out) & Invasive Aspergillosis : Considered as the cause of orbital cellulitis in immunocompromised patients.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 216.

Endophthalmitis

Tropical

Orbital cellulitis

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 45 y/o man from a tropical region presents with a painful red eye, significant watering, and sensitivity to light after working in a garden with decaying plant material. The patient is diagnosed with keratomycosis. Which of the following is not an examination finding in keratomycosis? (Q853665)

A. Feathery extensions at the margins of the corneal ulcer


B. Extensive corneal vascularization

C. Sterile immune ring

D. Endothelial plaque

Correct Answer: B. Extensive corneal vascularization

Explanation: Extensive corneal vascularization is generally absent in pure fungal keratitis, which distinguishes it from bacterial infections where vascularization is more common. Keratomycosis {{caption_text}} Clinical Features Pain & Foreign Body Sensation Watering of the eye Photophobia Blurred Vision Redness. Examination Findings Corneal Ulcer : Greyish-white, dry-looking with elevated, rolled-out margins. Pigmented Ulcer : Brownish appearance, possible with dematiaceous fungi. Feathery Extensions (Option A ruled out) : Delicate, finger-like projections into the surrounding stroma. Sterile Immune Ring (Option C ruled out) : Yellow line of demarcation where fungal antigen and host antibodies meet. Satellite Lesions : Multiple small lesions around the main ulcer. Hypopyon : Large, non-sterile, even with a small ulcer. Endothelial Plaque (Option D ruled out) : Located under the stromal lesion, made of fibrin and leucocytes. Corneal Vascularization : Typically absent in pure fungal ulcers. Perforation : Rare but possible. Investigations Corneal Scraping : Performed under topical anaesthesia for microbiological examination. Wet KOH Mount : Reveals filamentous fungi (branched, septate hyphae) or non-filamentous yeast-like fungi (Candida). Gram's & Giemsa Stains : To detect fungal hyphae and inflammatory cells. Culture : Sabouraud's agar medium is used for fungal culture. PCR : Recommended for rapid identification of fungal pathogens. Anterior Chamber Paracentesis : Considered if there is an intraocular extension. Corneal Biopsy : Indicated for deep stromal abscess or when cultures are negative. Confocal Microscopy : Used for identifying actual fungi within the cornea .



References:

Kanski’s Clinical Ophthalmology, 9th Edition, Page 216, 217.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q13: A 45 y/o male presents with redness, pain, and decreased vision in his right eye. He reports a history of trauma to the eye with a plant material three days ago. O/E, there is a corneal ulcer with feathery edges and satellite lesions. Hypopyon is also noted. Which of the following is the most appropriate initial treatment for this patient? (Q853666)

A. Topical Moxifloxacin

B. Topical Natamycin

C. Oral Acyclovir

D. Intravitreal Amphotericin B

Correct Answer: B. Topical Natamycin

Explanation: The history of trauma with plant material, along with the presence of a corneal ulcer with feathery edges and satellite lesions, is highly suggestive of fungal keratitis . Natamycin is the first-line treatment for fungal keratitis, especially caused by filamentous fungi like Fusarium and Aspergillus . It is effective in treating the type of fungal infection that is suggested by the clinical presentation in this patient. Treatment of Keratomycosis (Fungal Corneal Ulcer): Primary Treatment: Antifungal Medications: Topical Antifungal Agents: Natamycin Voriconazole Amphotericin B (effective against Aspergillus and Fusarium) Nystatin (effective against Candida) Oral Antifungal Agents (for severe ulcers with specific features): Ketoconazole Voriconazole Indications for Oral Therapy: Hypopyon Perforation Suspicion of endophthalmitis Surgical Management (if unresponsive to medical therapy): Repeated Scraping: Removal of necrotic material Cauterization Options: Pure carbolic acid (100%) Trichloroacetic acid (10%–20%) Povidone-iodine 5% Therapeutic Keratoplasty: Corneal transplantation Management of Suspected Endophthalmitis: Ultrasound : To check for exudates in the vitreous Vitreous Tap & intravitreal injections of antibiotics and antifungals (amphotericin B) if confirmed Surgical Options (if medical therapy fails): Corneal graft Conjunctival flap Topical Moxifloxacin (Option A): Moxifloxacin is an antibiotic effective against bacterial infections, not fungal infections. Given the history of trauma with plant material and the appearance of a corneal ulcer with feathery edges and satellite lesions, the most likely diagnosis is fungal keratitis. Therefore, moxifloxacin would not be appropriate. Oral Acyclovir (Option C): Acyclovir is an antiviral medication used primarily for herpes simplex virus infections. It would not be effective for a fungal corneal ulcer, making it an inappropriate choice in this scenario. Intravitreal Amphotericin B (Option D): Intravitreal Amphotericin B is used in cases of fungal endophthalmitis, not as an initial treatment for keratitis. Although hypopyon suggests severe infection, the primary treatment should still be topical antifungal agents unless there is confirmed or high suspicion of endophthalmitis requiring intravitreal therapy.

References:

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 217, 218.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q14: A 45 y/o patient presents with eye redness, excessive tearing, and sensitivity to light. O/E you observe dendritic ulcers, and decreased corneal sensation. Based on these findings, which virus is most likely responsible for the patient's condition? (Q853667)

A. Adenovirus

B. Varicella Zoster Virus (VZV)


C. Herpes Simplex Virus (HSV)

D. Chlamydia trachomatis

Correct Answer: C. Herpes Simplex Virus (HSV)

Explanation: s: HSV keratitis is characterized by dendritic ulcers on the cornea, which are branching, tree-like lesions. HSV also commonly causes decreased corneal sensation. Examination Findings in Viral Keratitis: Signs: Swollen Epithelial Cells: Early signs presenting as swollen, opaque cells in a coarse punctate or stellate pattern. Dendritic Ulcer: Characteristic linear, branching ulcer with terminal bulbs; stains with fluorescein. Geographic Ulcer: Enlarged amoeboid-shaped ulcer, often due to inappropriate steroid use. ulcer labelled" data-author="Alouan A, Cherkaoui O" data-hash="2333" data-license="Open Access" data-source="Ulcère de cornée en carte géographique au test à la flu | Open-i (nih.gov)" data-tags="March2025" src="" /> {{caption_text}} Rose Bengal Staining: Stains virus-laden cells at the ulcer edges, aiding in differentiation from other conditions. Reduced Corneal Sensation: Result of corneal nerve damage. Subepithelial Haze: Mild haziness was observed in the subepithelial layer. Anterior Chamber Activity: Mild inflammatory reaction in the anterior chamber. Elevated IOP: Increased intraocular pressure, requiring monitoring. Persistent Punctate Erosions (PEE): Residual erosions and epithelial irregularities after healing, resolving spontaneously without indicating recurrence. Specific Findings Based on Virus Type: Herpes Simplex Virus (HSV) Keratitis: Dendritic Ulcers: Branching, tree-like ulcers. Geographic Ulcers: Larger, amoeboid-shaped ulcers. Stromal Keratitis: Inflammation of the corneal stroma, leading to oedema, infiltrates, and scarring. Disciform Keratitis: Central, disc-shaped inflammation in the stroma. Ulcer labelled" data-author="Farooq, Asim & Arpeet, Shah & Shukla, Deepak. " data-hash="2336" data-license="CC BY-NC 4.0" data-source="illustration of a dendritic ulcer caused by HSV. Notes: The dendritic... | Download Scientific Diagram (researchgate.net)" data-tags="March2025" src="" /> {{caption_text}} Herpes Zoster Ophthalmicus (HZO): (Option B) Vesicles and Scars: Painful rash and blisters following the involvement of ophthalmic division of the trigeminal nerve. Nummular Keratitis: Large, coin-shaped, subepithelial corneal infiltrates. Keratitis labelled" data-author="Jkokavec" data-hash="2334" data-license="CC BY-SA 3.0" data-source="Nummular keratitis 1 - Nummular keratitis - Wikipedia" data-tags="March2025" src="" /> {{caption_text}} Scleritis: Inflammation of the sclera, causing pain and redness. Iritis: Inflammation of the iris leading to pain, photophobia, and blurred vision. Adenovirus (Option A):

Adenoviruses are known for causing epidemic keratoconjunctivitis (EKC) and may present with conjunctivitis and keratitis. They are less likely to cause dendritic ulcers, which are specific to HSV. Chlamydia trachomatis (Option D): Chlamydia typically causes trachoma or inclusion conjunctivitis, which may involve conjunctival scarring and corneal pannus, but not dendritic ulcers.




Geographic (amoebic) ulcer

@Neet_pg_bot



Dendritic ulcer

@Neet_pg_bot



Nummular Keratitis

@Neet_pg_bot

References:

Kanski’s Clinical Ophthalmology, 9th Edition 9, Pages 219, 220.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q15: A 35 y/o patient presents with pain, redness, photophobia, and decreased vision in the right eye. Slit-lamp examination reveals a dendritic ulcer on the cornea. Which of the following is the most appropriate initial treatment for this patient? (Q853668)

A. Topical steroids and antibiotics

B. Oral acyclovir and lubricants

C. Topical acyclovir and cycloplegics

D. Oral steroids and topical antibiotics

Correct Answer: C. Topical acyclovir and cycloplegics

Explanation: The patient presents with pain, redness, photophobia, and decreased vision in the right eye, along with a dendritic ulcer on the cornea. This clinical picture is classic for Herpes Simplex Keratitis (HSK), an infection of the cornea caused by the herpes simplex virus (HSV). Treatment: Topical Acyclovir: This antiviral medication is the first-line treatment for HSV keratitis. It directly targets the herpes simplex virus, reducing viral replication and promoting healing of the corneal ulcer. Cycloplegics: These medications help to relieve pain caused by ciliary muscle spasms and prevent the formation of synechiae (adhesions) in the iris. Treatment of Herpes Simplex Virus (HSV) Keratitis: Topical Therapy: Antivirals: Common Agents: Aciclovir 3% ointment: 5 times daily. Ganciclovir 0.15% gel: 5 times daily. Trifluridine: Up to 9 times daily. Older Agents: Idoxuridine and vidarabine (less common due to lower efficacy/higher toxicity). Other Topical Medications: Antibiotics: Prevent secondary bacterial infections, especially with epithelial defects. Cycloplegics: (e.g., Homatropine 1%) Relieve pain and photophobia by paralyzing ciliary muscles and dilating the pupil. Steroids: Epithelial Keratitis: Contraindicated due to the risk of worsening infection and corneal perforation. Stromal Keratitis: Used cautiously with antivirals to control inflammation and reduce scarring. Oral Therapy: Antivirals: Indications: immunocompromised patients, children, severe ocular surface disease, intolerance/resistance to topical treatment. Prophylaxis: Low-dose oral acyclovir (400 mg twice daily) reduces recurrence. Alternatives include valaciclovir (500 mg once daily) and famciclovir. Steroids: Generally avoided but may be used with caution in severe stromal inflammation as an adjunct to topical therapy. Surgical Management: Debridement: Removal of infected epithelial cells, helpful in resistant epithelial keratitis. Keratoplasty: Indication: Severe corneal scarring after at least a year of infection-free status. Risk of Recurrence: HSV keratitis may recur post-keratoplasty. Eyes with extensive corneal vascularization are at higher risk of graft rejection and recurrence. Prophylactic oral acyclovir is recommended. Topical steroids and antibiotics (Option A): Topical steroids are contraindicated in the treatment of active epithelial HSV keratitis, as they can suppress the immune response and worsen the viral infection, potentially leading to complications like corneal perforation. Antibiotics are not needed unless there is a secondary bacterial infection, which is not indicated here. Oral acyclovir and lubricants (Option B): While oral acyclovir is used in more severe or recurrent cases of HSV keratitis, topical acyclovir is preferred as the initial treatment. Lubricants can be supportive but are not the primary treatment. Oral steroids and topical antibiotics (Option D): Oral steroids are contraindicated in HSV keratitis, similar to topical steroids, as they can exacerbate the infection.

Topical antibiotics are unnecessary unless there's evidence of bacterial superinfection.

References:

Kanski’s Clinical Ophthalmology, 9th Edition 9, Page 221.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [13_Sclera] 029 Previous Year Questions

« Back to Folder Directory

[13_Sclera] 029 Previous Year Questions — Quick Jump

Q1 (p. 1517) *Q2 (p. 1518)* *Q3 (p. 1519)* *Q4 (p. 1522)* *Q5 (p. 1525)* *Q6 (p. 1526)*

Q1: A patient with long-standing, poorly controlled rheumatoid arthritis presents with a sudden, painless blurring of vision in the superior visual field of one eye. Fundoscopic examination shows retinal hemorrhages and dilated, tortuous veins in the inferior retina. What is the most likely diagnosis? (Q853116)

A. Branch Retinal Artery Occlusion

B. Central Retinal Artery Occlusion

C. Branch Retinal Vein Occlusion

D. Central Retinal Vein Occlusion

Correct Answer: C. Branch Retinal Vein Occlusion

Explanation: The most likely diagnosis for a patient with long-standing, poorly controlled rheumatoid arthritis presenting with sudden, painless blurring of vision in the superior visual field of one eye, and a fundoscopic examination showing retinal hemorrhages and dilated, tortuous veins in the inferior retina, is Branch Retinal Vein Occlusion (BRVO). It occurs when a single branch of the central retinal vein is occluded, causing retinal edema and hemorrhages localized to the area supplied by the blocked vein. Visual Defects: Visual loss is partial, superior or inferior field loss, depending on the location of the occlusion. Prognosis for central vision is better unless the macula is involved, often seen in superior temporal vein occlusion. Complications: Neovascularization and secondary glaucoma are rare but can occur in cases of widespread capillary occlusion. Diagnostic Features: Fluorescein Angiography: Shows areas of hypofluorescence, incomplete perifoveal arcades, and capillary closure in poorly perfused areas. Widespread capillary occlusion is associated with cotton-wool spots, delayed arteriovenous transit time, vessel leakage, and retinal edema. Treatment: No definitive treatment is effective once complete venous blockage occurs. Panphotocoagulation or Cryotherapy: Used to prevent complications like neovascular glaucoma and rubeosis iridis in cases of widespread capillary occlusion. Photocoagulation: Should only be performed after intraretinal blood absorption to address areas of poor perfusion, reducing edema and neovascularization risk. Anti-VEGF Therapy: Intravitreal injections are now standard for improving visual recovery and preventing late complications. Branched Retinal Artery Occlusion (Option A) a cherry red spot in the retina is noticed due to the surrounding edema. Central Retinal Artery Occlusion (Option B) obstruction occurs at the level of the lamina cribrosa, causing sudden painless loss of vision within seconds. Central Retinal Vein Occlusion (Option D) there is a mild blurring of vision, which improves over the day.

References:

Parsons Diseases of the Eye 23rd edition Pg.no:773,776

↑ *Back to this question in Part A*

@Neet_pg_bot


Q2: A patient presents with a nodular swelling near the limbus, which blanches after administering phenylephrine and then recurs. What is the most probable diagnosis? (Q844042)

A. Episcleritis

B. Scleritis


C. Bitot spots

D. Pterygium



Correct Answer: A. Episcleritis

Explanation: Correct Option A – Episcleritis: The patient’s clinical presentation is suggestive of nodular episcleritis. Its diagnosis is confirmed by a phenylephrine test; it blanches upon administering phenylephrine. Incorrect Options: Option B - Scleritis: Scleritis does not respond to phenylephrine. Option C - Bitot spots: Bitot's spot consists of keratinized epithelium and is seen in vitamin A deficiency. Option D - Pterygium: Pterygium is conjunctival degeneration characterized by triangular fibrovascular subepithelial ingrowth of bulbar conjunctiva over the limbus.





References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: What is the probable diagnosis based on the observed blanching of vascularity following the administration of 10 % phenylephrine, as depicted in the provided


image? (Q854717)


A. Nodular episcleritis

B. Nodular scleritis

C. Nodular phlycten

D. Scleral ectasia





Correct Answer: A. Nodular episcleritis

Explanation: The patient has painless redness of the eye which blanches (as shown in the image) following the application of 10% phenylephrine, suggesting episcleritis- Nodular episcleritis Episcleritis Definition Episcleritis refers to inflammation of the loose connective tissue between the sclera and the conjunctiva Etiology Idiopathic Systemic diseases - gout, rosacea, psoriasis, and connective tissue diseases. Hypersensitivity reaction to endogenous tubercular or streptococcal toxins. Infectious episcleritis - herpes zoster virus, syphilis, Lyme disease, and tuberculosis. Clinical features Common in young females Acute onset redness Discomfort - gritty/foreign body sensation Mild to no pain Resolves spontaneously Recurrence is common Clinical types Simple episcleritis: Sectorial (occasionally diffuse) inflammation of episclera. The engorged episcleral vessels are large and run in a radial direction beneath the conjunctiva Nodular episcleritis: Pink or purple flat nodule surrounded by injection, usually situated 2-3 mm away from the limbus. The nodule is firm, tender can be moved separately from the sclera and the overlying conjunctiva also moves freely. Diagnosis 10% Topical phenylephrine blanches overlying conjunctival vessels and inflamed episcleral vessels. Differential diagnosis Conjunctivitis , which is more superficial; blanching occurs with 2.5% phenylephrine Phlyctenulosis (Option C ruled out) Inflamed pinguecula Scleritis , which is deeper and more painful; does not blanch with 10% phenylephrine (Option B ruled out) Treatment Topical NSAIDs Topical mild corticosteroid eye drops Topical artificial tears, Cold compresses applied to the closed lids may offer symptomatic relief from ocular discomfort. Systemic NSAIDs Scleral ectasia (Option D) refers to a thinning and bulging of the sclera, often associated with conditions like keratoconus . It is not an inflammatory condition, and it would not cause the blanching of vascularity seen with phenylephrine. The clinical presentation of scleral ectasia typically involves corneal changes rather than inflammation of the episclera.

References:

Parson’s Diseases of the eye, 23rd ed., page 198.

Yanoff and Duker Ophthalmology, 5e., page 212

Kanski's Clinical ophthalmology, 9e., page 292

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 5-year-old boy presents with a small nodule on the sclera as shown in the image below. What is the likely diagnosis? (Q855476)

A. Pterygium

B. Pinguecula

C. Dermolipoma

D. Dermoid cyst





Correct Answer: D. Dermoid cyst

Explanation: Dermoid cyst appears as a lenticular yellow tumor near the corneal margin and consists of epidermoid epithelium with sebaceous glands and hair. Dermoid cyst is a congenital , often benign growth that contains skin-like structures , such as hair follicles, sweat glands, and sebaceous glands. It can present as a small,

firm, yellowish nodule on the sclera or conjunctiva . These cysts are often located near the limbus (the junction between the cornea and sclera) and may be noticed in childhood, especially if they grow or become more noticeable over time. A dermoid cyst may be diagnosed based on its characteristic appearance and location, and sometimes imaging or excision is needed for confirmation. {{caption_text}} Pterygium (Option A) : A pterygium is a fibrovascular growth that extends from the conjunctiva onto the cornea, typically seen on the nasal side of the eye. It is usually triangular in shape and may cause visual disturbance. It is not typically a nodule confined to the sclera. {{caption_text}} Pinguecula (Option B): A yellowish, triangular patch on the conjunctiva near the limbus, typically in the elderly, associated with sun exposure and not extending onto the cornea. {{caption_text}} Dermolipoma (Option C) : A dermolipoma is a benign tumor consisting of fat and skin-like tissue . While they may be located in the eye area, they typically present as larger, softer masses , and this does not match the presentation of a small nodule on the sclera.







References:

Kanski's Clinical Ophthalmology, 9th edition, Page- 197, 198.

↑ *Back to this question in Part A*


Q5: What is the probable diagnosis of a 35-year-old male with a history of prolonged corneal ulcer who now presents with a deformed eye as shown in the image? (Q857704)

A. Megalocornea

B. Conjunctivitis

C. Buphthalmos


D. Anterior staphyloma



Correct Answer: D. Anterior staphyloma

Explanation: The history of a prolonged corneal ulcer is suggestive of low body immune response or infection by a virulent strain of bacteria leading to sloughing of the whole cornea, leaving a narrow rim behind along with iris prolapse resulting in the false cornea that cannot withstand raised intraocular pressure and results in forward bulging of the cornea with iris tissue. As a result an ectatic cicatrix forms called Anterior Staphyloma as seen in the image. Staphyloma: It is ectasia of the outer coats of the eye (cornea, sclera, or both) with an incarceration of the uveal tissue. Based on the site affected, staphyloma can be classified as {{caption_text}} Types of staphyloma Causes Anterior Pseudocornea (following corneal ulcer) (Option D) Intercalary Perforating injuries of the peripheral cornea, marginal corneal ulcer, anterior scleritis, and scleromalacia perforans Ciliary Glaucoma, scleritis, and trauma to the ciliary region Equatorial (approx 14 mm behind the limbus) Scleritis, degenerative myopia, and chronic uncontrolled glaucoma Posterior Degenerative high axial myopia Treatment: Treat underlying condition Local excision and repair with a corneal and scleral patch graft Large staphyloma:

Staphylectomy and keratoplasty, or enucleated and replaced with an implant Megalocornea (Option A): Rare bilateral nonprogressive congenital defect that is characterized by an increased corneal diameter greater than 12.5 to 13 mm at birth and a deep anterior chamber with normal intraocular pressures and clear cornea. Conjunctivitis (Option B): Is the inflammation or infection of the transparent membrane that lines your eyelid and eyeball. Typical symptoms include redness and a gritty sensation in your eye, along with itching. Buphthalmos (Option C): Meaning "bull-like eyes," is a condition characterized by the enlargement of the eyeball secondary to congenital Glaucoma . It typically manifests before the age of 3 years and results from the accumulation and retention of aqueous humor. OP185



References:

Parsons Diseases of the Eye 23rd Edition, pages 531-533

↑ *Back to this question in Part A*

@Neet_pg_bot


Q6: In the image, what are the associated complications observed in a 35-year-old woman who has been diagnosed with rheumatoid arthritis? (Q856012)

A. Scleromalacia perforans

B. Ciliary staphyloma


C. Coloboma

D. Malignant melanoma



Correct Answer: A. Scleromalacia perforans

Explanation: - Scleromalacia perforans The diagnosis in this scenario is scleromalacia perforans. It is usually associated with long-standing rheumatoid arthritis and presents with scleral thinning appearing bluish and without signs of inflammation. The underlying cause of scleral thinning is probably ischemia because of vessel damage (type III hypersensitivity). Scleromalacia Perforans: Most common in elderly females Pathophysiology: Associated with long-standing rheumatoid arthritis Melting of the sclera is due to obliteration of arterial supply causing ischemia Vessel damage is mediated by type III hypersensitivity (Option A) Clinical manifestations: Painless scleral thinning Underlying dark uveal tissue is visible Rarely perforation of the globe {{caption_text}} Scleromalacia perforans Treatment: No specific therapy Protection of the eye from trauma is very crucial Plenty of topical lubrication Oral NSAIDs Oral steroids Immunomodulatory agents Surgery-scleral patch grafting Ciliary staphyloma (Option B) : This condition involves outpouching of the eye wall near the ciliary body, typically associated with trauma or inflammatory conditions other than rheumatoid arthritis, but it is not a direct complication of RA. Coloboma (Option C) : A coloboma is a congenital defect in the eye, where part of the eye structure (such as the iris or retina) is missing. It is not associated with rheumatoid arthritis . Malignant melanoma (Option D) : This is a type of eye cancer that arises from pigment-producing cells. It is unrelated to rheumatoid arthritis and is not an autoimmune complication.



References:

Parsons diseases of the eye 23e pg 527

Scleromalacia Perforans - EyeWiki

Image :https://eyewiki.org/Scleromalacia_Perforans

↑ *Back to this question in Part A*

@Neet_pg_bot

• [14_Lacrimal Drainage System] 030 Lacrimal Drainage System

« Back to Folder Directory

[14_Lacrimal Drainage System] 030 Lacrimal Drainage System — Quick Jump

Q1 (p. 1529) *Q2 (p. 1531)* *Q3 (p. 1531)* *Q4 (p. 1533)* *Q5 (p. 1536)* *Q6 (p. 1536)*
Q7 (p. 1539) *Q8 (p. 1540)* *Q9 (p. 1541)* *Q10 (p. 1542)*

Q1: Which structure within the lacrimal passages prevents the backflow of tears from the nasal cavity into the nasolacrimal duct? (Q853822)

A. Valve of Hasner

B. The valve of Rosenmüller

C. Orbital lobe


D. Glands of Krause


Correct Answer: A. Valve of Hasner

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The valve of Hasner is a crucial structure located at the lower end of the nasolacrimal duct, where it opens into the inferior meatus of the nose. Its primary function is to prevent the retrograde flow of fluid from the nasal cavity into the nasolacrimal duct, which is particularly important during actions like nose blowing. This valve ensures that tears drain properly from the eye into the nasal cavity without backflow. The valve of Rosenmüller (Option B) is a mucosal fold located at the junction where the common canaliculus (formed by the superior and inferior canaliculi) opens into the lacrimal sac. Orbital Lobe (Option C) is one of the two lobes of the main lacrimal gland, which is involved in the production of reflex tears. Glands of Krause (Option D) are accessory lacrimal glands found in the conjunctival fornices.





References:

Parsons' Diseases of the Eye, 23rd Edition, Pg.1102,1103 Kanski's Clinical Ophthalmology 9th edition page 100-101

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Which of the following is a cause of lacrimal pump failure, leading to excessive tearing (epiphora)? (Q853823)

A. Trauma to the canaliculus

B. Bell’s palsy

C. Obstruction of the nasolacrimal duct

D. Narrowing of the punctum

Correct Answer: B. Bell’s palsy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The lacrimal pump mechanism relies on the proper function of the orbicularis oculi muscle and the correct positioning of the punctum. In Bell’s palsy, weakness of the orbicularis oculi muscle prevents proper eyelid closure and tear drainage, as the orbicularis oculi muscle is involved in both the tear suction and drainage phase of the lacrimal pump mechanism. This leads to epiphora (excessive tearing). Trauma to the canaliculus (Option A): While an injury to the canaliculus can block the tear drainage system, it does not directly disrupt the lacrimal pump mechanism. Obstruction of the nasolacrimal duct (Option C): This condition involves a blockage in the nasolacrimal duct, preventing proper tear drainage. Narrowing of the punctum (Option D): Punctal stenosis refers to a narrowing or closure of the punctum (the opening at the corner of the eyelid that drains tears).

References:

Parsons Diseases of the Eye, 23rd Edition, Page 1114

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 65 y/o male presents with persistent tearing in his right eye that has worsened over the past year. The physician orders an imaging study to pinpoint the exact location of the blockage and assess the condition of the lacrimal sac. Which investigation is most appropriate for this purpose? (Q853824)

A. Dacryocystography

B. Fluorescein Dye Disappearance Test (FDDT)

C. Dacryoscintigraphy

D. Lacrimal Syringing


Correct Answer: A. Dacryocystography


- *Listen (English)*
- *Listen (Hindi)*

Explanation: Dacryocystography (DCG) provides highly detailed anatomical information about the entire lacrimal drainage system. By injecting contrast dye into the lacrimal sac and taking X-rays, DCG can precisely identify the site, extent, and nature of the blockage, making it the investigation of choice when anatomical clarity is required. Fluorescein Dye Disappearance Test (FDDT) (Option B): It is mainly used to assess the overall efficiency of tear drainage and does not provide information on the exact location or nature of any obstruction within the lacrimal system. Dacryoscintigraphy (Option C) or lacrimal scintillography is a functional test that evaluates the rate at which tears flow through the lacrimal drainage system by tracking a radioactive tracer. While it is useful for identifying functional delays or blockages, it lacks the anatomical precision needed to determine the exact location or cause of the obstruction. Lacrimal Syringing (Option D): Normal saline is injected into the lacrimal sac using a syringe. Although it can identify the degree of blockage & approximate blockage site, it does not offer detailed anatomical information.



Normal Dacryocystography





References:

Parsons' Diseases of the Eye, 23rd Edition, Pages 1114, 1115, 1116

Kanski's Clinical Ophthalmology, 9th Edition, Page 104

Yanoff & Duker Ophthalmology, 5th Edition, Pages 1339, 1340

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 50 y/o man undergoes Jones Test 1 for chronic tearing, which shows no dye on the cotton bud placed in the inferior meatus. Following this, Jones Test 2 is performed, and fluorescein dye is detected after lacrimal syringing. What does this result suggest? (Q853825)

A. Partial obstruction within the lacrimal passages

B. Complete obstruction of the nasolacrimal duct

C. Lacrimal pump failure


D. Primary hypersecretion of tears


Correct Answer: A. Partial obstruction within the lacrimal passages

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: A positive Jones Test 2 indicates that the fluorescein dye was able to enter the lacrimal sac but could not reach the nasal cavity without syringing. This finding is typical of a partial blockage somewhere along the lacrimal drainage system, allowing some tear flow but not enough for normal drainage. Jones Test 1: Purpose : To distinguish between partial obstruction and excessive tear production. Procedure : Fluorescein dye is placed in the conjunctival sac, and after 5 minutes, a cotton bud soaked in anaesthetic is placed in the inferior meatus inside the nose. The cotton bud is checked for dye. Interpretation: Positive: Dye on the cotton bud indicates patent tear drainage and suggests hypersecretion. Negative: No dye indicates potential drainage issues, requiring further testing with Jones Test. Jones Test 2: Purpose: Performed if Jones Test 1 is negative to differentiate between partial obstruction and pump failure. Procedure: Lacrimal syringing is done while the cotton bud remains in the nose. Interpretation: Positive: Dye flushed onto the cotton bud indicates partial obstruction. Negative: No dye suggests lacrimal pump failure.





References:

Parsons' Diseases of the Eye, 23rd Edition, Page 1115

Kanski’s Clinical Ophthalmology, 9th Edition, Pages 103, 104

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 9-month-old infant is brought to the clinic with ongoing tearing and discharge from the right eye. Despite regular lacrimal sac massage and topical antibiotics, the symptoms persist. What is the next best step in management? (Q853826)

A. Continue lacrimal sac massage and antibiotics

B. Probing of the nasolacrimal duct

C. Dacryocystorhinostomy (DCR)

D. Balloon catheter dilation

Correct Answer: B. Probing of the nasolacrimal duct

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: In congenital nasolacrimal duct obstruction (CNLDO), persistent symptoms beyond 6–8 months of age, despite conservative management (e.g., Crigler massage and antibiotics), warrant nasolacrimal duct probing . Since this child is 9 months old with no response to massage and antibiotics , the next step is probing , which is often successful when done before 12 months. Congenital Nasolacrimal Duct Obstruction (CNLDO): Aspect Details Definition Blockage of the nasolacrimal duct, usually at the valve of Hasner First-Line Management • Crigler massage to open the duct • Topical antibiotics for infection Indications for Probing • Persistence beyond 6–10 months • Repeated infections or failure of conservative therapy Continue massage and antibiotics (Option A): Incorrect because symptoms have already persisted beyond the expected resolution time despite these measures. Dacryocystorhinostomy (DCR) (Option C): Incorrect as DCR is reserved for older children or adults when probing and intubation fail. Balloon catheter dilation (Option D): Incorrect because balloon dacryoplasty is typically considered for recurrent or failed probing , not first-line.

References:

Parson's Diseases of the Eye, 23rd Edition, Page 1111

Yanoff & Duker Ophthalmology, 5th Edition, Page 1341

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: Which of the following types of Dacryocystorhinostomy (DCR) is considered for treating nasolacrimal duct obstruction, offering the highest success rate? (Q853827)

A. External Dacryocystorhinostomy (External DCR)

B. Endonasal Dacryocystorhinostomy (Endonasal DCR)


C. Endocanalicular Laser Dacryocystorhinostomy (Laser DCR)


D. Conjunctivodacryocystorhinostomy (CDCR)

Correct Answer: A. External Dacryocystorhinostomy (External DCR)

- *Listen (English)*
- *Listen (Hindi)*

Explanation: External Dacryocystorhinostomy (External DCR) procedure is performed through an incision near the inner corner of the eye, providing direct access to the lacrimal sac and nasal mucosa. It is widely done due to its high success rate , which exceeds 90% . However, it leaves a visible scar and may result in greater blood loss compared to other methods. Endonasal Dacryocystorhinostomy (Endonasal DCR) (Option B) involves using an endoscope inserted through the nasal cavity to create a drainage opening, thus avoiding any external incision. While it has the advantage of being scarless and typically involves less bleeding, its success rate is slightly lower, ranging between 70-90%. Endocanalicular Laser Dacryocystorhinostomy (Laser DCR) (Option C): This method utilizes a laser passed through the canaliculli to create a drainage opening. It is minimally invasive and can be performed quickly under local anaesthesia. However, its success rate is lower than external or endonasal DCR Conjunctivodacryocystorhinostomy (CDCR) (Option D): CDCR is specifically used when both the canaliculi and the nasolacrimal duct are blocked. It is a more specialized procedure compared to standard DCR and is not typically the first choice for isolated nasolacrimal duct obstruction. CDCR involves creating a passage directly from the conjunctival sac to the nasal cavity, often using a Jones tube and is reserved for more complex cases where the canaliculi are also involved.





References:

https://pmc.ncbi.nlm.nih.gov/articles/PMC5678323/

Parsons' Diseases of the Eye, 23rd Edition, Pg.1123,1124

Kanski's Clinical Ophthalmology 9th edition page 107, 108

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A 2-week-old infant presents with a bluish swelling in the inner corner of the right eye. The parents report excessive tearing and some slimy discharge from the eye. There is no history of trauma or illness. What is the most likely diagnosis? (Q853828)

A. Congenital Dacryocele

B. Dacryoadenitis

C. Dacryops

D. Chronic Dacryocystitis

Correct Answer: A. Congenital Dacryocele

■ *Listen to Audio 1*

Explanation: The bluish swelling in the inner corner of the eye, along with excessive tearing and mucous discharge, is characteristic of congenital dacryocele. Congenital Dacryocele: A condition where obstruction of the nasolacrimal duct causes amniotic fluid accumulation in the lacrimal sac, leading to a cystic swelling below the inner corner of the eye. Dacryoadenitis (Option B): Presents with a painful, red swelling of the outer third of the upper eyelid, often associated with systemic infections. Dacryops (Option C): Presents as a cystic swelling in the upper fornix due to blocked lacrimal ducts and would typically not be described as bluish or associated with mucous discharge in the medial canthal area. Chronic Dacryocystitis (Option D): Occurs when a blockage leads to recurrent infections of the lacrimal sac over time. It is less likely in a 2-week-old infant presenting for the first time with these symptoms. Additionally, chronic dacryocystitis would typically involve more prolonged symptoms and recurrent infections, which are not indicated in this scenario.



References:

Parsons' Diseases of the Eye, 23rd Edition, Page 1104, 1105

Kanski's Clinical Ophthalmology, 9th Edition, Page 109

Congenital dacryocystocele: diagnosis and treatment - PMC (nih.gov)

Prevalence and clinical characteristics of congenital dacryocystocele - PMC (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: A 55 y/o woman presents with a persistent swelling below the inner corner of her left eye. She also reports a history of watery eyes and recurrent conjunctivitis. On examination, there is fluctuant swelling at the lacrimal sac, and mucoid fluid can be expressed. What is the most likely diagnosis and appropriate management? (Q853829)

A. Chronic Dacryocystitis; Probing and syringing.

B. Chronic Dacryocystitis; Dacryocystorhinostomy (DCR).

C. Acute Dacryocystitis; Systemic antibiotics and abscess drainage.

D. Acute Dacryocystitis; Dacryocystectomy (DCT).

Correct Answer: B. Chronic Dacryocystitis; Dacryocystorhinostomy (DCR).

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The presence of a mucocele suggests chronic dacryocystitis where the lacrimal sac has been obstructed for a prolonged period, leading to the accumulation of mucoid fluid. Milky or gelatinous mucoid fluid expressed from the lower punctum on pressing the swelling (positive Regurgitation test) indicates lacrimal mucocele. Dacryocystorhinostomy (DCR) is the definitive surgical procedure designed to create a new drainage pathway between the lacrimal sac and the nasal cavity, thus resolving the obstruction and restoring proper tear drainage. This approach addresses the underlying blockage and prevents recurrence.

References:

Parsons' Diseases of the Eye, 23rd Edition, Pg.1110-1119 Kanski's Clinical Ophthalmology 9th edition page 110

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: A 60 y/o man presents with watery eyes, redness, swelling at the inner corner of his right eye, and persistent thick yellow discharge. O/E there is pouting of lacrimal puncta and yellow granules from the punctum upon massage. Which organism is most likely responsible, and what is the definitive treatment? (Q853830)

A. Eikenella corrodens; Oral antibiotics and steroid drops.

B. Staphylococcus aureus; Topical antibiotics and warm compresses

C. Actinomyces israelii; Canaliculotomy

D. Chlamydia trachomatis; Irrigation and syringing of the tear duct.

Correct Answer: C. Actinomyces israelii; Canaliculotomy

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Canaliculitis is diagnosed based on symptoms and clinical findings. Actinomyces israelii is the most common cause of canaliculitis with yellow granules. Canaliculotomy is the definitive treatment to remove concretions and resolve the infection. It involves accessing and cleaning out the canaliculus to remove concretions and ensure thorough treatment of the infection. Canaliculitis: Canaliculitis is an inflammation or infection of the lacrimal canaliculus. Causes: Bacterial infection- most commonly Actinomyces israelii Other bacteria: Staphylococcus, Streptococcus, Eikenella, Lactococcus, Nocardia Fungal infections. Viral infections. Treatment:

Initial: Topical and systemic antibiotics, warm compresses, local massage, irrigation and syringing of the tear duct Definitive: Surgery often required for effective treatment Canaliculotomy for direct access and curettage. Safe and effective procedure. Punctal dilation or punctoplasty with curettage Canaliculoplasty with intubation to avoid scarring Follow-up with antibiotic treatment and addressing underlying causes, such as a loose punctal plug



Pouting of puncta

References:

Kanski's Clinical Ophthalmology, 9th Edition, Page 109

Canaliculitis - StatPearls - NCBI Bookshelf (nih.gov)

Canaliculitis: What It Is, Causes & Treatment (clevelandclinic.org)

Lacrimal canaliculitis - PMC (nih.gov)

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: A 27 yeal old man presents with a painless, slow-growing, firm mass on the upper eyelid that has been gradually increasing in size over the past year. On examination, the lesion is well-circumscribed, mobile, and non-tender, with no signs of ulceration or inflammation. Imaging reveals a well-defined, lobulated mass without bony invasion. What is the most likely diagnosis? (Q853831)

A. Pleomorphic Adenoma

B. Lymphoma

C. Lacrimal Gland Carcinoma

D. Dacryocystitis

Correct Answer: A. Pleomorphic Adenoma

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Pleomorphic Adenoma: Pleomorphic adenoma is the most common tumor of the lacrimal gland. It typically present's as a benign, painless , firm, nontender and slow-growing mass. Location: Upper eyelid or superotemporal orbit. Occurs predominantly in young adult males. Lacrimal gland carcinoma (Option C): Lacrimal gland carcinoma is a rare but aggressive malignancy with high morbidity and mortality. The most common type is adenoid cystic carcinoma (50%), followed by pleomorphic adenocarcinoma, mucoepidermoid carcinoma, and squamous cell carcinoma. Lymphoma (Option B): Lymphoma can present as a mass in the lacrimal gland but often has associated systemic symptoms and can cause more diffuse swelling rather than a distinct mass. Dacryocystitis (Option D) : It refers to inflammation of the nasolacrimal sac and would typically present with pain, redness, and possibly discharge, along with swelling in the lower eyelid.

References:

Kanski's Clinical Ophthalmology 9th edition page 141, 142

↑ *Back to this question in Part A*

@Neet_pg_bot

• [14_Lacrimal Drainage System] 031 Previous Year Questions

« Back to Folder Directory

[14_Lacrimal Drainage System] 031 Previous Year Questions — Quick Jump

Q1 (p. 1544) *Q2 (p. 1545)* *Q3 (p. 1546)* *Q4 (p. 1547)* *Q5 (p. 1548)* *Q6 (p. 1552)*
Q7 (p. 1555) *Q8 (p. 1557)* *Q9 (p. 1560)* *Q10 (p. 1560)* *Q11 (p. 1563)* *Q12 (p. 1563)*

Q1: A 3-year-old child presents with epiphora, and probing reveals a soft block at 7 mm. What is the best treatment option for this condition? (Q852499)

A. Endonasal Dacryocystorhinostomy (DCR)


B. Conjunctival Dacryocystorhinostomy (DCR)

C. External Dacryocystorhinostomy (DCR)

D. Dacryocystectomy

Correct Answer: B. Conjunctival Dacryocystorhinostomy (DCR)

Explanation: Epiphora (excessive tearing) and a soft block at 7 mm on probing indicate that the level of obstruction is canalicular (in the canaliculus, not the nasolacrimal duct). For canalicular obstruction, the preferred treatment is Conjunctival DCR (also known as Lester Jones tube placement). This procedure involves creating a bypass from the conjunctival sac to the nasal cavity using a glass tube (Lester Jones tube) to restore tear drainage. Conjunctival DCR is specifically designed for proximal obstructions (e.g., canalicular blocks), whereas Endonasal DCR or External DCR is used for distal obstructions (e.g., nasolacrimal duct blocks). Endonasal Dacryocystorhinostomy (Option A): T his is used for nasolacrimal duct obstructions, not for canalicular blockages. External Dacryocystorhinostomy (Option C): This is more suitable for distal nasolacrimal duct obstructions and is not the first choice for canalicular obstructions. Dacryocystectomy (Option D): This is a last-resort procedure used for chronic infections or tumors and is not indicated for a functional lacrimal system with a canalicular block.



@Neet_pg_bot

References:

Parsons' Diseases of the Eye, 23rd Edition, Pg.1123,1124 Kanski's Clinical Ophthalmology 9th edition page 107, 108

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: Which bones are cut/removed during external Dacryocystorhinostomy? (Q860569)

A. Zygomatic

B. Maxillary

C. Lacrimal

D. Frontal

Correct Answer: C. Lacrimal

Explanation: Dacryocystorhinostomy (DCR) is a surgical procedure performed to treat nasolacrimal duct obstruction, which causes epiphora due to impaired tear drainage. The surgery involves creating a new drainage pathway for tears from the lacrimal sac into the nasal cavity. Indications for DCR: Chronic dacryocystitis (infection of the lacrimal sac) Obstruction of the nasolacrimal duct (either congenital or acquired, e.g., from trauma or aging) Recurrent epiphora (despite conservative treatments) Types of DCR: External DCR (Traditional): Involves making an incision on the skin near the medial canthus of the eye . The lacrimal bone (Mostly) and a portion of the maxillary bone are removed to create an opening between the lacrimal sac and the nasal cavity. A stent is often placed to keep the passage open until healing is complete. Endoscopic DCR (Minimally invasive): No external incision is made. An endoscope is inserted through the nasal cavity to create a passage between the lacrimal sac and nasal cavity. This approach is preferred for cosmetic reasons and for patients who may have anatomical difficulties with external incisions. Postoperative Care: Antibiotics are prescribed to prevent infection. Stent removal (if placed) occurs after 6–8 weeks. Nasal saline irrigation may be recommended to keep the new drainage pathway clear. Complications: Infection Stenosis (narrowing) of the new drainage passage Persistent epiphora or failure of the procedure

References:

https://www.ncbi.nlm.nih.gov/books/NBK557851/

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 3-month-old child is brought to the outpatient department with watering of eyes and photophobia . On examination, both corneas appear hazy. Which of the following is the most likely diagnosis ? (Q840075)

A. Congenital nasolacrimal duct obstruction

B. Congenital glaucoma

C. Congenital entropion

D. Congenital distichiasis

Correct Answer: B. Congenital glaucoma

Explanation: The triad of watering (epiphora), photophobia, and blepharospasm is pathognomonic for congenital glaucoma . Pathophysiology: Due to developmental defect in the trabecular meshwork or angle of anterior chamber , leading to raised intraocular pressure (IOP) . The elevated IOP causes corneal edema , which leads to photophobia and watering . As the cornea enlarges, the eye may appear buphthalmic (large cornea) . Incorrect options A) Congenital nasolacrimal duct obstruction: Causes watering only (no photophobia). C) Congenital entropion: Causes watering from lash irritation but not photophobia. D) Congenital distichiasis: Causes mild watering and irritation, but not corneal haze or photophobia.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q4: Identify the marked structure: (Q857418)

A. Canal of petit

B. Canal of hannover

C. Hyaloid canal


D. Pars plana



Correct Answer: A. Canal of petit

Explanation: The Canal of Petit (also known as Spatia Zonularia) is a sacculated canal that encircles the equator of the lens, formed by the zonular fibers (the zonule of Zinn) that anchor the lens to the ciliary body. This canal is located behind the suspensory ligament of the lens and contains aqueous humor in the interstices between the zonules. It is an important anatomical feature in the structure of the eye. Zonular Spaces (Spatia Zonularia): Zonular fibers, also known as the zonule of Zinn , anchor the lens to the ciliary body and the ciliary part of the retina . These fibers are separated by spaces filled with aqueous humor , referred to as zonular spaces . Anatomy:

The zonules form a fibrous ring around the equator of the lens, connecting it to the ciliary body . The spaces between the zonules form a sacculated canal known as Hannover's Canal or Petit's Canal or Petit's spaces (spatia zonularia), which encircles the lens' equator. Function: These spaces are involved in the structure and function of the ciliary zonule , providing mechanical support for the lens and contributing to accommodation. The canal can be inflated using a fine blowpipe inserted under the suspensory ligament. Canal of Hannover (Option B) : Another term related to zonular anatomy, but it refers to a slightly different structure and is often used interchangeably with the Canal of Petit in some contexts. However, in this case, the correct answer is Canal of Petit . Hyaloid Canal (Option C) : This is a different structure in the eye, located within the vitreous body, and is the remnant of the fetal hyaloid artery. It is not related to the zonules or lens attachment. Pars Plana (Option D) : A region of the retina, part of the ciliary body, involved in ocular surgeries (like vitrectomy) but not related to the canal structures surrounding the lens.



References:

Zonular spaces - e-Anatomy - IMAIOS

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: What is the accurate order of staining in patients with dry eyes? i. Fluorescein stain ii. Lissamine green iii. Rose Bengal stain (Q857419)

A. i, iii, ii


B. i, ii, iii


C. ii, i, ii


D. iii, ii, i


Correct Answer: B. i, ii, iii


Explanation: In the management of dry eye syndrome , different stains are used to assess the health of the ocular surface. The typical order of staining is as follows: Fluorescein Stain : First step : Fluorescein is typically used first to highlight epithelial defects and tear film stability . It is a bright yellow-green dye that fluoresces under cobalt blue light, marking areas of corneal epithelial damage. It can also help assess the tear break-up time (TBUT). Lissamine Green : Second step : Lissamine green is a vital dye that stains dead or degenerated cells in the conjunctiva and cornea, often used to evaluate the conjunctival surface in dry eyes. It is less irritating than Rose Bengal, making it ideal for dry eye patients, and it stains areas with poor tear film or disrupted epithelial integrity . Rose Bengal Stain : Third step : Rose Bengal stains damaged epithelial cells in the conjunctiva and cornea, particularly in cases of severe dryness . It can be more irritating to the eye than the other two stains, so it is typically used last in the staining sequence. It is helpful in identifying severe damage, such as conjunctival filaments or corneal ulcers . Investigations of dry eye 1, Slit-lamp examination Superficial corneal erosions Inadequate tear lake volume Early tear film break-up time Conjunctival hyperemia Conjunctival surface irregularities Meibomian gland dysfunction 2. Schirmer’s Test: Schirmer’s test is used to diagnose aqueous deficiency dry eye disease. Schirmer's Test 1 Measures both basal and reflex tear secretions Uses Whatman-4 filter paper, kept in the lower fornix at the junction of lateral one-third and medial two-thirds. After 5 minutes, wetting of the filter strip from the bent end is measured. Normal value: >15 mm 5-10 mm: Mild to moderate keratoconjunctivitis sicca (KCS). < 5 mm: Severe KCS {{caption_text}} [1] Schirmer's Test 2 Measures only basal secretion under anesthesia 3. Phenol Red Test A modified Schirmer’s test using a cotton thread impregnated with phenol red dye. The thread is placed in the lower fornix for 15 seconds. Phenol dye changes from yellow to red depending on the pH of the tears. Wet length: >20 mm: Normal. <10 mm: Severe dry eye. ≤19 mm as borderline dry eye. {{caption_text}} 4. Tear Film Break-Up Time (TBUT) Tear film stained with sodium fluorescein 1% is observed under a slit lamp, and time is noted after the patient blinks. The interval between the complete blink and the first appearance of a ‘dark’ dry spot on the cornea is recorded as TBUT. It is an indicator of the adequacy of the mucin component of tears. Normal: 15-35 seconds < 10 seconds indicates unstable tear film. {{caption_text}} 5. Rose Bengal or Lissamine Green Stain devitalized, desiccated corneal and conjunctival epithelium red or green. {{caption_text}} {{caption_text}} 6. Alcian Blue Stains particulate matter in the tear film due to mucus 7. Tear Osmolarity Increased in dry eye 8. Tear to Lysozyme ratio Between 0.9 and 0.6 indicates dry eye. Options A, C, & D are incorrect.


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot


@Neet_pg_bot



References:

Kanski's Clinical Ophthalmology, Edition 9, Page- 160, 161, 162.

Diagnosis of dry eye disease and emerging technologies - PMC

Schirmer Test - an overview | ScienceDirect Topics

Evaluation of Phenol Red Thread test versus Schirmer test in dry eyes: A comparative study - PMC

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 60-year-old patient presents with chronic dacryocystitis. She complains of recurrent tearing, discharge, and mild swelling in the inner corner of the eye. The clinical findings and history are consistent with chronic dacryocystitis. What should be the treatment of choice for this patient? (Q858129)

A. Lacrimal sac massage

B. Dacryocystorhinostomy

C. Syringing

D. Probing

Correct Answer: B. Dacryocystorhinostomy

Explanation: ausative organisms: Streptococcus haemolyticus, Pneumococcus, Staphylococcus Stasis and mild infection of long duration Common in women aged 40-60 Causes: anatomical factors (narrow bony canal), nasal diseases, foreign bodies, infection. Clinical Features/ Sequelae Stage of cellulitis: Painful swelling at the lacrimal sac, epiphora, fever, redness, and oedema. Lacrimal abscess: Pus-filled sac, fluctuant swelling, may rupture forming an abscess. Fistula formation: Untreated abscess leads to spontaneous discharge, forming a fistula. Stage of chronic catarrhal dacryocystitis: Watering eye, mild redness, NLD block Lacrimal mucocele: Swelling below inner canthus, mucoid fluid regurgitation. Chronic suppurative dacryocystitis: Purulent discharge (d/t infection), recurrent conjunctivitis. Chronic fibrotic sac: Thickened mucosa, persistent epiphora, irregular folds. Image {{caption_text}} {{caption_text}} Complications Conjunctivitis, corneal ulcer, lid abscess, orbital cellulitis, facial cellulitis, rarely cavernous sinus thrombosis. Conjunctivitis, ectropion of lower lid, risk of corneal ulcers. Treatment Cellulitis: Systemic antibiotics, anti-inflammatory drugs, hot fomentation. Lacrimal abscess: Drainage with incision, antibiotics, possible DCT or DCR later. External fistula: Fistulectomy with DCT or DCR after controlling infection. Conservative: Probing and syringing (early stages). Dacryocystorhinostomy (DCR): Preferred surgery to re-establish lacrimal drainage. Dacryocystectomy (DCT): Indicated if DCR is contraindicated or in cases of fibrosis, infection, or old age. Lacrimal sac massage (Option A) : This is a conservative approach often recommended for mild or early cases of nasolacrimal duct obstruction, especially in infants. It helps express the accumulated tears and debris in the lacrimal sac. However, in chronic dacryocystitis in adults, this approach is typically insufficient and would not resolve the underlying obstruction. Syringing (Option C) (irrigation of the lacrimal system) can help clear the obstruction in some cases and is often used diagnostically to confirm the presence of a blockage. However, it does not provide a long-term solution and is typically not effective for treating chronic dacryocystitis in adults, particularly when there is significant structural obstruction. Probing (Option D) is a procedure used primarily in young children with congenital nasolacrimal duct obstruction. It is typically not appropriate for adults with chronic dacryocystitis, especially in cases where there is long-standing obstruction or scarring of the nasolacrimal duct.





References:

Parsons' Diseases of the Eye, 23rd Edition, Pg.1110-1119

Kanski’s Clinical Ophthalmology 9th edition page 110

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: What is the most appropriate treatment for a 55-year-old woman who is experiencing long-term mucopurulent discharge from her left eye and swelling near the inner corner of the eye? (Q857191)

A. Dacryocystorhinostomy

B. Dacryocystectomy

C. Incision and drainage

D. Balloon catheter dilation

Correct Answer: A. Dacryocystorhinostomy

Explanation: The clinical presentation of mucopurulent discharge from her left eye along with presence of a swelling , i.e., mucocele at inner canthus is suggestive of chronic dacryocystitis . The treatment of choice for chronic dacryocystitis is dacryocystorhinostomy (DCR) which helps in lacrimal drainage. Acute and Chronic Dacryocystitis: Aspect Acute Dacryocystitis Chronic Dacryocystitis Aetiology Acute exacerbation of chronic dacryocystitis Spread from neighbouring infected structures (e.g., sinuses, bones, dental abscess) Causative organisms: Streptococcus haemolyticus, Pneumococcus, Staphylococcus Stasis and mild infection of long duration Common in women aged 40-60 Causes: anatomical factors (narrow bony canal), nasal diseases, foreign bodies, infection. Clinical Features/ Sequelae Stage of cellulitis: Painful swelling at the lacrimal sac, epiphora, fever, redness, and oedema. Lacrimal abscess: Pus-filled sac, fluctuant swelling, may rupture forming an abscess. Fistula formation: Untreated abscess leads to spontaneous discharge, forming a fistula. Stage of chronic catarrhal dacryocystitis: Watering eye, mild redness, NLD block Lacrimal mucocele: Swelling below inner canthus, mucoid fluid regurgitation. Chronic suppurative dacryocystitis: Purulent discharge (d/t infection), recurrent conjunctivitis. Chronic fibrotic sac: Thickened mucosa, persistent epiphora, irregular folds. Image {{caption_text}} {{caption_text}} Complications Conjunctivitis, corneal ulcer, lid abscess, orbital cellulitis, facial cellulitis, rarely cavernous sinus thrombosis. Conjunctivitis, ectropion of lower lid, risk of corneal ulcers. Treatment Cellulitis: Systemic antibiotics, anti-inflammatory drugs, hot fomentation. Lacrimal abscess: Drainage with incision, antibiotics, possible DCT or DCR later. (Option C) External fistula: Fistulectomy with DCT or DCR after controlling infection. Conservative: Probing and syringing (early stages). Dacryocystorhinostomy (DCR): Preferred surgery to re-establish lacrimal drainage. (Option A) Dacryocystectomy (DCT): Indicated if DCR is contraindicated or in cases of fibrosis, infection, or old age (>60y). (Option B)



Lacrimal fistula

@Neet_pg_bot



References:

Parsons’ Diseases of the Eye 23rd Edition, pages 1110-1119.

Kanski’s Clinical Ophthalmology 9th edition, page 110.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q8: Which of the following surgeries use this instrument? (Q857195)

A. Evisceration

B. Enucleation

C. Cataract surgery

D. Dacryocystorhinostomy




Correct Answer: D. Dacryocystorhinostomy

Explanation: The instrument shown is Bone Punch, composed of a sturdy spring handle and two blades that are joined at a right angle. The upper blade features a sharp cutting edge and a small hole, while the lower blade has a cup-shaped indentation. This tool is utilized in DCR (dacryocystorhinostomy) surgery to expand the bony opening by punching the bone from the edges of the opening. Types of DCR: External Dacryocystorhinostomy (External DCR) procedure is performed through an incision near the inner corner of the eye, providing direct access to the lacrimal sac and nasal mucosa. The lacrimal sac is accessed via the Bone punch used to expand the opening

(Option D) . It is widely regarded as the gold standard due to its high success rate , which exceeds 90% . However, it leaves a visible scar and may result in greater blood loss compared to other methods. {{caption_text}} External DCR {{caption_text}} Endonasal Dacryocystorhinostomy (Endonasal DCR): involves using an endoscope inserted through the nasal cavity to create a drainage opening, thus avoiding any external incision. While it has the advantage of being scarless and typically involves less bleeding, its success rate is slightly lower, ranging between 70-90%. Endocanalicular Laser Dacryocystorhinostomy (Laser DCR): This method utilizes a laser passed through the tear duct to create a drainage opening. It is minimally invasive and can be performed quickly under local anaesthesia. However, its success rate is lower than external or endonasal DCR, around 70%, and patients often require additional surgery if the initial procedure fails. Conjunctivodacryocystorhinostomy (CDCR): CDCR is specifically used when both the canaliculi and the nasolacrimal duct are blocked. It is a more specialized procedure compared to standard DCR and is not typically the first choice for isolated nasolacrimal duct obstruction. CDCR involves creating a passage directly from the conjunctival sac to the nasal cavity, often using a Jones tube and is reserved for more complex cases where the canaliculi are also involved.







References:

Parsons' Diseases of the Eye 23rd Edition, pages 1123,1124.

Kanski’s Clinical Ophthalmology 9th edition, pages 107,108.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: Which of the following treatments is appropriate for a 3-month-old child with a mucopurulent discharge from the right puncta and a history of increased watering in the right eye, as noticed by the mother? (Q857689)

A. Syringing

B. Probing with syringing

C. Sac massage with topical antibiotics

D. Dacryocystorhinostomy

Correct Answer: C. Sac massage with topical antibiotics

Explanation: The clinical presentation is suggestive of Congenital Dacryocystitis. As the infant’s age is less than 6 months, Lacrimal sac massage with topical antibiotics would be the preferred management. Treatment of Dacryocystitis: < 6-9 months Lacrimal sac massage with topical antibiotics (Option C) 9 months - 4 years Bowman’s probing with syringing (Option A) (Option B) > 4 years Dacryocystorhinostomy (Option D)

References:

Parsons’ Diseases of the Eye, 23rd Edition, Pg.1111

Yanoff & Duker Ophthalmology 5th edition page 1341

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: A 55-year-old woman is being counseled for surgery for her long-standing dacryocystitis. Which of the following is the surgery of choice? (Q859321)

A. Dacryocystorhinostomy

B. Dacryocystectomy


C. Probing

D. Balloon catheter dilation

Correct Answer: A. Dacryocystorhinostomy

Explanation: Correct Option A - Dacryocystorhinostomy (DCR) The treatment of choice for chronic dacryocystitis is dacryocystorhinostomy (DCR) which helps in lacrimal drainage. Dacryocystorhinostomy (DCR) is the definitive surgical procedure designed to create a new drainage pathway between the lacrimal sac and the nasal cavity, thus resolving the obstruction and restoring proper tear drainage. This approach addresses the underlying blockage and prevents recurrence. Aspect Acute Dacryocystitis Chronic Dacryocystitis Aetiology Acute exacerbation of chronic dacryocystitis Spread from neighbouring infected structures (e.g., sinuses, bones, dental abscess) Causative organisms: Streptococcus haemolyticus, Pneumococcus, Staphylococcus Stasis and mild infection of long duration Common in women aged 40-60 Causes: anatomical factors (narrow bony canal), nasal diseases, foreign bodies, infection. Clinical Features/ Sequelae Stage of cellulitis: Painful swelling at the lacrimal sac, epiphora, fever, redness, and oedema. Lacrimal abscess: Pus-filled sac, fluctuant swelling, may rupture forming an abscess. Fistula formation: Untreated abscess leads to spontaneous discharge, forming a fistula. Stage of chronic catarrhal dacryocystitis: Watering eye, mild redness, NLD block Lacrimal mucocele: Swelling below inner canthus, mucoid fluid regurgitation. Chronic suppurative dacryocystitis: Purulent discharge (d/t infection), recurrent conjunctivitis. Chronic fibrotic sac: Thickened mucosa, persistent epiphora, irregular folds. Image {{caption_text}} {{caption_text}} Complications Conjunctivitis, corneal ulcer, lid abscess, orbital cellulitis, facial cellulitis, rarely cavernous sinus thrombosis. Conjunctivitis, ectropion of lower lid, risk of corneal ulcers. Treatment Cellulitis: Systemic antibiotics, anti-inflammatory drugs, hot fomentation. Lacrimal abscess: Drainage with incision, antibiotics, possible DCT or DCR later. External fistula: Fistulectomy with DCT or DCR after controlling infection. Conservative: Probing and syringing (early stages). Dacryocystorhinostomy (DCR): Preferred surgery to re-establish lacrimal drainage. Dacryocystectomy (DCT): Indicated if DCR is contraindicated or in cases of fibrosis, infection, or old age. Dacryocystectomy (Option B) This procedure involves the removal of the lacrimal sac and is usually reserved for lacrimal sac fibrosis.. Probing (Option C) It is a common treatment approach for congenital nasolacrimal duct obstruction in infants. Balloon catheter dilation (Option D) This procedure is used in some cases to dilate and open a partially blocked nasolacrimal duct but may not be the primary choice for chronic dacryocystitis.





References:

Parsons’ Diseases of the Eye, 23rd Edition, Pg.1110-1119

Kanski’s Clinical Ophthalmology 9th edition page 110

↑ *Back to this question in Part A*

@Neet_pg_bot

Q11: In Dacryocystorhinostomy (DCR), the lacrimal sac is opened into which nasal meatus? (Q859753)

A. Inferior meatus

B. Middle meatus

C. Supreme meatus

D. Superior meatus

Correct Answer: B. Middle meatus

Explanation: DCR creates a new drainage pathway between the lacrimal sac and the nasal cavity , bypassing an obstructed nasolacrimal duct. Surgically, an osteotomy is made through the lacrimal bone and frontal process of maxilla , opening the sac into the middle meatus . This allows direct tear drainage and prevents recurrence of epiphora. Inferior meatus (Option A): This is where the nasolacrimal duct normally opens , but DCR bypasses the duct. Superior/Supreme meatus (Option C & D): Not involved in lacrimal drainage.

References:

Parsons’ Diseases of the Eye, 23rd Edition, Pg.1123,1124

Kanski’s Clinical Ophthalmology 9th edition page 107, 108

↑ *Back to this question in Part A*

@Neet_pg_bot

Q12: A 60-year-old patient presents with chronic dacryocystitis . She complains of recurrent tearing, discharge, and mild swelling in the inner corner of the eye . The clinical findings and history are consistent with chronic dacryocystitis. What should be the treatment of choice for this patient ? (Q820463)

A. Lacrimal sac massage

B. Dacryocystorhinostomy

C. Syringing

D. Probing

Correct Answer: B. Dacryocystorhinostomy

Explanation: Correct Option B - Dacryocystorhinostomy: The treatment of choice for chronic dacryocystitis is dacryocystorhinostomy (DCR) which helps in lacrimal drainage. The procedure involves the removal of the bone near the nasolacrimal sac and the creation of an opening of the lacrimal sac into the mucosa of the middle meatus in the nose. Incorrect Options: Option A - Lacrimal sac massage: Lacrimal sac massage involves gently massaging the area over the lacrimal sac to try and open the blocked nasolacrimal duct in an infant of less than 9 months with congenital dacryocystitis. Option C - Syringing: Syringing, or irrigation, involves flushing the nasolacrimal system with fluid to clear any obstructions. While it can be useful for diagnosing the location of the obstruction, it is not a definitive treatment for chronic dacryocystitis. Option D - Probing: Probing is a procedure in which a thin probe is inserted into the nasolacrimal duct to try and open the blockage. It is a common treatment approach for congenital nasolacrimal duct obstruction in infants of age more than 9 months, success rate is till 2 years.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [15_Community Ophthalmology] 032 Previous Year Questions

« Back to Folder Directory

[15_Community Ophthalmology] 032 Previous Year Questions — Quick Jump

Q1 (p. 1565) *Q2 (p. 1566)* *Q3 (p. 1566)* *Q4 (p. 1567)* *Q5 (p. 1569)* *Q6 (p. 1569)*

Q1: Which organism is most likely causing the infection in a 14-day-old neonate with bilateral conjunctivitis and lung infiltrates, as found in a chest X-ray performed at the outpatient department? (Q855386)

A. Streptococcus

B. Hemophilus influenzae

C. Chlamydia trachomatis

D. Mycoplasma

Correct Answer: C. Chlamydia trachomatis

Explanation: A 14-day-old neonate with Bilateral conjunctivitis and pneumonia-like presentation is suggestive of Chlamydia trachomatis infection. Neonatal conjunctivitis is also known as Ophthalmia neonatorum. Ophthalmia neonatorum: Definition Ophthalmia neonatorum or neonatal conjunctivitis is a mucoid, mucopurulent, or purulent discharge from one or both eyes in the first month of life. Any discharge, even watery, from a baby's eyes in the first week of life must be viewed under suspicion as tears are not secreted so early in life. Aetiology of ophthalmia neonatorum Mode of infection: Before birth: Very rare through infected liquor amnii During birth: From an infected birth canal After birth: During the first bath of a newborn Soiled clothes Fingers with infected lochia Source of infection: Neisseria gonorrhoeae: Transmitted at the time of delivery Presents between 2-5 days of birth (Option B) The discharge becomes mucopurulent and purulent quickly Thick yellow pus and marked chemosis Cornea seen at the bottom of a crater-like pit in severe cases Chlamydia trachomatis: Transmitted at the time of delivery Onset at 5-14 days after birth (Option C) Serotypes D to K are the most common cause of ophthalmia neonatorum in developed countries Characteristic intracellular inclusion bodies are seen Less severe inflammation Purulent discharge + Other causes: Staphylococci/ Streptococcus pneumoniae (Option A) Gram-negative coliforms Herpes simplex virus Chemical: silver nitrate solution previously used for prophylaxis immediately after birth : erythema, chemosis, lid edema, and discharge within the first 24 hours after birth. Prophylaxis Antenatal measures: Treatment of genital infections and care of mother Surveillance of pregnant women for STDs Natal measures: Aseptic, sterile precautions to be followed during deliveries Clean the infant's closed lids with sterile cotton soaked in normal saline and dried Post-natal measures: Povidone-iodine 2.5% solution If the mother is under suspicion, 1% tetracycline or 0.5% erythromycin eye ointment is used in the baby immediately after birth and watched for a week. Ceftriaxone 50mg/kg IM or IV single dose for infants born to mothers with untreated gonococcal infection. Crede's method: Old method using 1% silver nitrate applied to infants' eyes right after birth It is NOT used now as it can cause chemical neonatal conjunctivitis. Diagnosis Treatment Neisseria gonorrhoea Ceftriaxone IM, gentamicin drops, bacitracin eye ointment Chemical Wash eyes, erythromycin ointment, observe

Other bacteria Neomycin-bacitracin ointment, gentamicin, or tobramycin drops Herpes simplex virus Acyclovir 3% eye ointment, systemic acyclovir (systemic involvement) Chlamydia trachomatis Erythromycin or chlortetracycline eye ointment, oral erythromycin (systemic infection) Incorrect options: Streptococci (Option A) and Mycoplasma (Option D): Though they can cause conjunctivitis and pneumonia, it is very rare in the case of a newborn.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: According to the National Blindness and Visual Impairment Survey conducted from 2015 to 2019, what is the leading cause of moderate to severe visual impairment? (Q855573)

A. Refractive error

B. Untreated cataract

C. Glaucoma

D. Diabetic retinopathy

Correct Answer: B. Untreated cataract

Explanation: According to the National Blindness and Visual Impairment Survey (2015-2019) , untreated cataract remains the leading cause of moderate to severe visual impairment . Cataracts, which cause the lens of the eye to become cloudy and impair vision, are particularly prevalent in older adults. When left untreated, cataracts can lead to significant visual impairment and blindness, making it a major contributor to visual disability worldwide. Major Causes of Blindness in population aged ≥ 50 years Cataract - 66.2% Corneal opacity (including trachomatous) Cataract surgical complications (including PCO) Posterior segment disease (excluding DR & ARMD) (Option D ruled out) Glaucoma (Option C ruled out) Major Causes of Visual Impairment in population aged ≥ 50 years Cataract - 71.2% (Option B) Refractive error (Option A ruled out) Cataract surgical complications (including PCO) Major Causes of Blindness in population aged 0-49 years Corneal opacity - 37.5 All globe/ CNS abnormality (Amblyopia) Phthisis Other/undetermined Major Causes of Visual Impairment in population aged 0-49 years Refractive error - 29.6 Cataract All globe/ CNS abnormality (Amblyopia) Corneal opacity

References:

National Blindness and Visual Impairment Survey

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: Which of the following is not included in the SAFE strategy for trachoma? (Q856713)

A. Antibiotics

B. Surgery for trichiasis

C. Evaluation of control program

D. Facial cleanliness

Correct Answer: C. Evaluation of control program

Explanation: Evaluation of control program is NOT included in the SAFE strategy for Trachoma. Diagnosis and Management of Trachoma: Investigations Micro-immunofluorescence (micro-IF): Using pooled antigens; For routine diagnostic purposes Culture of C. trachomatis: In irradiated McCoy cells; Expensive Monoclonal antibody direct tests: IgA-IPA light microscopy tests are the best combination for diagnosis Management (SAFE strategy) SAFE is a WHO-recommended combination of interventions to treat trachoma and prevent it from leading to blindness S urgery for cicatricial entropion and trichiasis (Tertiary prevention) A ntibiotics to eliminate the organism (Secondary prevention) F acial cleanliness to avoid infection (Primary prevention) E nvironmental improvement with better hygiene and sanitation (Primordial prevention) Surgery (Tertiary prevention) (Option B) Epilation/ Bilamellar tarsal rotation Antibiotics (Secondary prevention) (Option A) For areas with 10% or more prevalence of active trachoma (TF in children 1-9 yrs): Azithromycin single oral dose 20 mg/kg to all community members 5% to 10% prevalence: Antibiotic therapy targeted in the family and close contact only Less than 5% prevalence: Only patients are treated Facial cleanliness (Primary prevention) (Option D) Frequent face wash with clean water, Avoid common towels, handkerchiefs, Surma rods, etc Environmental improvement (Primordial prevention) (Option C) Water latrines and good water supply to reduce flies, Sprays to control flies, Health education

References:

Kanski’s Clinical Ophthalmology, 9th edition, Page- 176

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: Which is the most common cause of preventable childhood blindness in India? (Q857231)

A. Refractive error


B. Vitamin A deficiency

C. Retinopathy of prematurity

D. Cataract

Correct Answer: B. Vitamin A deficiency

Explanation: The most common cause of preventable childhood blindness is vitamin A deficiency leading to Xerophthalmia. It can be prevented by timely vitamin A prophylaxis. Blindness in India: Overall most common cause of blindness/visual impairment (VI): Cataract Most common preventable cause of blindness: Vitamin A deficiency Most common cause of blindness in children: Vitamin A deficiency Most common cause of VI in children: Refractive error Other causes of blindness in children: Trachoma, Retinopathy of prematurity; Congenital cataract; Ophthalmia neonatorum. Proposed structure for Vision 2020, NPCB: Tier Number of Units Population Norms Components Centres of Excellence 20 1/50 million Professional Leadership Strategy development Continued Medical Education (CME) Laying of standards and quality assurance Research Training Centres 200 1/5 million Tertiary eye care including retinal surgery, corneal transplantation, glaucoma surgery, etc. Training and CME Service Centres 2000 1/500000 Cataract Surgery Other common eye surgeries Facilities for refraction Referral services Vision Centres 20,000 1/50000 Refraction and prescription of glasses Primary eye care School eye screening programme Screening and referral services



References:

Park’s Textbook of Preventive and Social Medicine 27th edition, page 505.

AK Khurana Comprehensive Ophthalmology 7th edition, pages 499-501

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: What type of visual impairment does a person have if their vision is less than 3/60 but greater than 1/60 in their better eye, and their field of vision is limited to less than 10 degrees from the center of fixation? (Q857712)

A. Curable blindness

B. Economic blindness

C. Social blindness

D. Manifest blindness

Correct Answer: C. Social blindness

Explanation: Vision < 3/60 but > 1/60 with field of vision < 10 degrees is suggestive of Social Blindness . NPCB Classification of Visual Impairment The National Programme for Control of Blindness (NPCB) in India defines blindness as " visual acuity of <3/60 in a better eye with best possible correction " and central visual field <10 degrees . The NPCB further categorizes visual impairment as: Classification Visual Acuity Visual Impairment Visual acuity of less than 6/18 in the better eye with available correction Low Vision Visual acuity of less than 6/18 to 6/60 Economic Blindness (Option B) Visual acuity of less than 6/60 to 3/60 Social Blindness (Option C) Visual acuity of less than 3/60 to 1/60 Manifest Blindness (Option D) Visual acuity of less than 1/60 to the perception of light Absolute Blindness No perception of light WHO Classification of Visual Impairment The World Health Organization (WHO) defines blindness as " visual acuity of less than 3/60 (Snellen) or its equivalent ". The International Classification of Disease 11 (2018) classifies vision impairment into distance and near-presenting vision impairment Classification Visual Acuity Mild Distance Vision Impairment Presenting visual acuity worse than 6/12 Moderate Distance Vision Impairment Presenting visual acuity worse than 6/18 Severe Distance Vision Impairment Presenting visual acuity worse than 6/60 Blindness Presenting visual acuity worse than 3/60 Near Vision Impairment Presenting near vision acuity worse than N6 or M0.8 at 40 cm with existing correction

References:

Park’s Textbook of Preventive and Social Medicine 27th edition, pages 456,457

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: How does the World Health Organization (WHO) define blindness? (Q860171)

A. VA < 3/60

B. VA < 6/60

C. VA > 3/60

D. VA > 6/60

Correct Answer: A. VA < 3/60

Explanation: NPCB Classification of Visual Impairment The National Programme for Control of Blindness (NPCB) in India defines blindness as "visual acuity of <3/60 in a better eye with best possible correction" and central visual field <10 degrees. The NPCB further categorizes visual impairment as: Classification Visual Acuity Visual Impairment Visual acuity of less than 6/18 in the better eye with available correction Low Vision Visual acuity of less than 6/18 to 6/60 Economic Blindness Visual acuity of less than 6/60 to 3/60 Social Blindness Visual acuity of less than 3/60 to 1/60 Manifest Blindness Visual acuity of less than 1/60 to the perception of light Absolute Blindness No perception of light WHO Classification of Visual Impairment The World Health Organization (WHO) defines blindness as "visual acuity of less than 3/60 (Snellen) or its equivalent". (Option A) The International Classification of Disease 11 (2018) classifies vision impairment into distance and near-presenting vision impairment Classification Visual Acuity Mild Distance Vision Impairment Presenting visual acuity worse than 6/12 Moderate Distance Vision Impairment Presenting visual acuity worse than 6/18 Severe Distance Vision Impairment Presenting visual acuity worse than 6/60 Blindness Presenting visual acuity worse than 3/60 Near Vision Impairment Presenting near vision acuity worse than N6 or M0.8 at 40 cm with existing correction Option B,C and D are incorrect.

References:

Park’s Textbook of Preventive and Social Medicine, 27th edition, pg no-456-457

↑ *Back to this question in Part A*

@Neet_pg_bot

• [16_Ocular Injuries] 033 Ocular Injuries

« Back to Folder Directory

[16_Ocular Injuries] 033 Ocular Injuries — Quick Jump

Q1 (p. 1571) *Q2 (p. 1575)* *Q3 (p. 1577)* *Q4 (p. 1577)* *Q5 (p. 1579)* *Q6 (p. 1580)*


Q1: Which of the following types of ocular injuries can result from blunt trauma to the eye? 1) 2) 3) 4) (Q853600)

A. 2 and 4


B. 1 and 2


C. 3 and 1

D. All of the Above




@Neet_pg_bot



@Neet_pg_bot




Correct Answer: D. All of the Above


- *Listen (English)*
- *Listen (Hindi)*

Explanation: All of the given presentations can result from Blunt trauma to the eye. Hyphema: Blood in the anterior chamber leading to corneal staining. 2. Descemet's membrane tear 3. Vossius ring: In the shape of the pupil 4. Lens subluxation Features of blunt trauma: Conjunctiva: Subconjunctival hemorrhage Iris: Iridodialysis (dehiscence of the iris from the ciliary body at its root), iris sphincter tear, Ciliary body: Ciliary body tear leading to angle recession. Lens: Vossious Ring , Rosette-shaped cataract (flower-shaped opacity), subluxation of lens. Globe rupture (caused by severe blunt trauma, indicated by iris prolapse) Vitreous haemorrhage : Pigment cells (‘tobacco dust’) can be seen floating in the anterior vitreous. Choroidal rupture Retina: Commotio Retinae (also known as Berlin's oedema) , retinal dialysis (disinsertion from ora serrata), giant retinal Tear (horseshoe shaped) , macular hole. Injury to optic nerve Traumatic neuropathy with chorioretinal scarring (decreased visual acuity, visual field defect, afferent pupillary defect, colour, and brightness).


@Neet_pg_bot


@Neet_pg_bot





References:

Kanski's Clinical Ophthalmology, 10th Edition, Page 914-921

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A patient presents with a full-thickness corneal wound caused by a pencil. Which of the following is the most likely complication if the injury is left untreated? (Q853601)

A. Ring abscess of the cornea

B. Retinal detachment

C. Glaucoma


D. Strabismus

Correct Answer: A. Ring abscess of the cornea

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Untreated penetrating injuries can introduce microorganisms, leading to severe infections like a ring abscess of the cornea, a sight-threatening condition. Leads to ring abscesses , corneal sloughing, iridocyclitis, endophthalmitis, panophthalmitis, etc. Retinal detachment (Option B) is more commonly associated with blunt trauma rather than penetrating injuries. Glaucoma (Option C) is a potential late complication but not the immediate concern in this scenario. Strabismus (Option D) refers to eye misalignment, which arises from muscle or nerve dysfunction, not corneal trauma.



References:

Kanski’s Clinical Ophthalmology, 10th Edition, Page 923-925.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A 40-year-old construction worker presents with eye pain and blurred vision after metal struck his eye. CT scan reveals an intraocular foreign body lodged in the vitreous cavity. What is the most appropriate treatment? (Q853602)

A. Observation and follow-up in one week

B. Corticosteroid injection without IOFB removal

C. Immediate surgery with pars plana vitrectomy and IOFB removal

D. Application of topical antibiotics and patching

Correct Answer: C. Immediate surgery with pars plana vitrectomy and IOFB removal

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: Pars plana vitrectomy with IOFB removal is essential to prevent complications and preserve vision, and it is ideally done during the initial closure. Intraocular Foreign Bodies (IOFBs): An IOFB can lodge in any part of the anterior or posterior segments, causing mechanical damage, infection, or toxic effects on intraocular structures. Types: Inert: Glass, wood, plastic, rubber, gold, silver, platinum Reactive: Organic and Inorganic (iron causing siderosis, copper causing chalcosis). Diagnosis CT (investigation of choice) with axial and coronal cuts detects and localizes a metallic IOFB. MRI is contraindicated (metallic IOFB). Treatment Magnetic removal (metallic IOFB) Forceps removal through pars plana vitrectomy Prophylaxis against infection (topical/systemic antibiotics) (Option D) Conservative and supportive management (topical corticosteroids , analgesics, etc). (Option B) Observation and follow-up in a week (Option A) in the presence of a foreign body in the vitreous cavity could lead to infection and further complications.

References:

Yanoff and Duker Ophthalmology, 5th Edition, Page 675,676

Kanski’s Clinical Ophthalmology, 10th Edition, Pages 927,928.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: A 28-year-old metalworker presents with gradual vision loss and a history of a metallic foreign body injury 4 months ago. Examination reveals iron deposits on Iris, rusty deposits on the lens and pigmentary changes in the retina. What is the most likely cause of his condition? (Q853603)

A. Chalcosis

B. Retinitis pigmentosa

C. Siderosis bulbi


D. Glaucoma


Correct Answer: C. Siderosis bulbi

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above case is suggestive of siderosis bulbi. It is a degenerative eye condition caused by an iron-containing foreign body that is not promptly removed from the eye. Mechanism: Oxidation and other chemical reactions with surrounding tissues, leading to toxic effects on cellular enzyme systems and subsequent cell death. Clinical Manifestations: Lens deposits Heterochromia iridis (iris pigmentation) Retinal changes (pigmentary retinopathy) Glaucoma (trabecular damage) Electroretinography (ERG): Progressive attenuation of the b-wave over time. Chalcosis (Option A) is caused by copper, not iron, and shows distinct features like Kayser-Fleischer rings. Retinitis pigmentosa (Option B) is genetic, not due to trauma. Glaucoma (Option D) can be a complication but is not the primary diagnosis here.







References:

Yanoff and Duker Ophthalmology, 5th Edition, Page 675

Kanski's Clinical Ophthalmology, 10th Edition, Page 928

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: A 35-year-old male presents to the emergency department after an accidental chemical exposure to his eyes while working in a lab. He reports immediate burning and decreased vision. Upon examination, there is corneal epithelial loss and conjunctival swelling. What is the most appropriate immediate management step? (Q853604)

A. Necrotic tissue removal and administer topical antibiotics

B. Apply corticosteroids to reduce inflammation

C. Immediate irrigation with water or saline for 20-30 minutes

D. Perform corneal transplant

Correct Answer: C. Immediate irrigation with water or saline for 20-30 minutes

■ *Listen (English)*

■ *Listen (Hindi)*

Explanation: The above clinical case is suggestive of chemical injury. Irrigation with water or saline is the priority to dilute and remove the chemical, minimizing damage. Management of Chemical Injury Immediate irrigation: Copious water/saline for 20–30 min or until pH normal; double eversion of lids. Debridement: Removal of dead conjunctiva/corneal epithelium with cotton swab (Option A). Antibiotics: To prevent secondary infection (Option A). Cycloplegics: For ciliary muscle relaxation and pain relief. Steroids: Only for initial 7 days (reduce inflammation but impair healing) (Option B). Vitamin C (Ascorbic acid): 10% eye drops 2-hourly to promote collagen synthesis. Collagenase inhibitors: Citrate 10% drops, tetracycline, acetylcysteine 10%. Early surgical options: Tenon advancement, limbal cell transplant, amniotic membrane graft. Late surgical options: Symblepharon release with conjunctival/mucous graft, keratoplasty (Option D). if corneal opacity. Complications: Secondary glaucoma, limbal stem cell deficiency, pseudopterygium, corneal opacity

References:

Kanski’s Clinical Ophthalmology, 10th Edition, Pages 929-931.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A 28-year-old male presents to the emergency department 10 days after sustaining blunt trauma to the face during a basketball game. He complains of diplopia and difficulty moving his left eye upwards. On examination, there is significant periorbital swelling. The X-ray is shown below. What does this X-ray indicate in this patient, and what is the most appropriate next step in management? (Q853605)

A. Orbital fat herniation; apply ice and prescribe analgesics

B. Orbital floor fracture; surgical evaluation

C. Orbital haemorrhage; immediate CT scan


D. Maxillary sinus fracture; follow-up with ENT specialist




Correct Answer: B. Orbital floor fracture; surgical evaluation

■ *Listen to Audio 1*

Explanation: The " teardrop sign" or “hanging drop sign ” on the X-ray suggests an orbital floor fracture , often associated with muscle entrapment, which leads to restricted eye movement and diplopia. The appropriate next step in management would be to evaluate for surgical intervention to relieve the entrapment and repair the fracture. Orbital imaging: X-rays: A nose-chin (Water's) view showing ' hanging drop ' opacity of the superior maxillary antrum from orbital contents herniating through the floor. Computerised tomography (CT) scanning and magnetic resonance imaging(MRI): Evaluate the extent of the fracture and herniation of contents into maxillary antrum (hanging drop or tear drop sign). CT scan of coronal section of orbit showing fracture floor with herniation of orbital contents Management: Antibiotics and anti-inflammatory drugs Ice packs and nasal decongestants Surgery (dislodge the muscle and do microplating) Indications for surgery: No improvement in diplopia or enophthalmos even after 10 days (indicates muscle entrapment). Enopthalmos is > 2mm Half of the floor is fractured





References:

Kanski’s Clinical Ophthalmology, 10th Edition, Pages 910-912.

↑ *Back to this question in Part A*

@Neet_pg_bot

• [16_Ocular Injuries] 034 Previous Year Questions

« Back to Folder Directory

[16_Ocular Injuries] 034 Previous Year Questions — Quick Jump

Q1 (p. 1584) *Q2 (p. 1585)* *Q3 (p. 1586)* *Q4 (p. 1587)* *Q5 (p. 1588)* *Q6 (p. 1588)*
Q7 (p. 1591) *Q8 (p. 1593)* *Q9 (p. 1596)* *Q10 (p. 1597)*

Q1: A patient presents with a foreign body (chuna) in the eye and complaints of irritability and lacrimation. What is the first step in management for this patient? (Q853118)

A. Antibiotic eye drops B) Ascorbic eye drops C) Irrigation of the eye with normal saline D) Steroids

B. Ascorbic eye drops

C. Irrigation of the eye with normal saline

D. Steroids

Correct Answer: C. Irrigation of the eye with normal saline

Explanation: Chemical injury to the conjunctiva and cornea is a severe ocular emergency that must be treated immediately. The first and the most crucial step in treating chemical injuries is copious eye irrigation by double eversion of the lid with normal saline. Deliver a minimum of 2 liters of water in 20-30 minutes or until pH is restored. Management of Alkali Burns in the Eye: Initial Evaluation and Emergency Response Immediate Irrigation: Continue irrigation during transport to the hospital until ocular surface pH normalizes to 7.0–7.2. Borate buffer (e.g., Cederroth Eye Wash) may be more effective than saline or water. Monitor pH: Check ocular pH with narrow-range strips (6–8) every 15–30 minutes to confirm stabilization. Ophthalmic Examination: Perform a full exam after pH normalization: Treatment Based on Severity (Grade I–IV): Grade I (Mild Injuries): Treatment: Topical antibiotic ointments (e.g., bacitracin, erythromycin). (Option A ruled out) Preservative-free artificial tears for lubrication. Topical steroids (e.g., prednisolone acetate) for 1 week to reduce inflammation. (Option D ruled out) Cycloplegic agents (e.g., cyclopentolate) for comfort. Grades II–IV (Moderate to Severe Injuries): Anti-inflammatory Measures: Hourly topical prednisolone acetate in the first week, tapered after day 10. Long-acting cycloplegics (e.g., atropine) for pain relief. Oral tetracyclines (e.g., doxycycline) to inhibit matrix metalloproteinases and reduce corneal melting risk. High-dose vitamin C (topical/oral) to prevent corneal ulceration. Antibiotics: Topical broad-spectrum antibiotics (e.g., fluoroquinolones) to prevent superinfection. IOP Management: Use aqueous suppressants to control elevated IOP. Amniotic Membrane: Early placement of amniotic membrane (e.g., Prokera) for grades II–IV injuries to support epithelial healing. Surgical Management Debridement: Remove necrotic corneal and conjunctival epithelium with sponges or surgical instruments. Persistent Epithelial Defects: Use amniotic membranes for healing (sutured or Prokera). Ascorbic eye drops (Option B) promote collagen synthesis and improve wound healing but it is not the initial step.

References:

https://www.aao.org/eyenet/article/treating-acute-chemical-injuries-of-cornea

↑ *Back to this question in Part A*

@Neet_pg_bot

Q2: A factory worker presented with a history of chuna (lime) exposure in both eyes a few hours ago. What is the first immediate management? (Q853143)

A. Take history B) Copious irrigation of eyes C) Check vision in both eyes D) Check IOP and plan for surgery

B. Copious irrigation of eyes

C. Check vision in both eyes

D. Check IOP and plan for surgery

Correct Answer: B. Copious irrigation of eyes

Explanation: Chemical eye injuries, particularly alkali burns (such as chuna, calcium hydroxide, or lime), are ophthalmic emergencies. Immediate and aggressive irrigation is the first and most critical step to minimize ocular damage. Chemical Injuries Eye injuries caused by chemicals range from mild irritation to vision loss. Sources of Injury Domestic: Ammonia, detergents, cosmetics Agriculture: Fertilizers, insecticides Laboratory: Acids, alkalis Assaults & Warfare: Acid attacks, chemical weapons Self-inflicted: Malingering or mental health conditions Alkali vs. Acid Burns Alkali: More severe, deeper penetration due to saponification, hygroscopic nature, and soluble compounds formation. Acid: Less severe, protein coagulation forms a protective barrier. Clinical Stages Acute: Conjunctival swelling, corneal epithelial loss, iris inflammation, possible corneal sloughing and staphyloma. Reparation: Epithelial regeneration, corneal vascularization, iris inflammation subsides. Complications: Symblepharon, recurrent ulcers, cataracts, secondary glaucoma. Emergency Management Immediate irrigation : Use water or saline for at least 20-30 minutes or until the eye's pH is restored. A minimum of 2 liters should be used. (Option B) Mechanical removal : Gently remove any chemical particles with a cotton swab, particularly in lime burns. Necrotic tissue removal : Carefully excise dead conjunctiva and corneal epithelium using a cotton swab. Healing & Complication Prevention Topical antibiotics: Use broad-spectrum antibiotics like moxifloxacin to prevent infection. Topical steroids: Control inflammation and anterior uveitis. Use cautiously after two weeks to avoid hindering collagen production. Cycloplegics: Alleviate pain and dilate the pupil with agents like atropine. Ascorbic acid: Promote wound healing and collagen synthesis using topical and systemic Vitamin C. Lubricants: Use preservative-free artificial tears to keep the eye moist and support healing. Autologous serum: Use serum eye drops to provide essential growth factors for healing. Sodium citrate: Stabilise neutrophils and reduce collagenase activity with topical sodium citrate. Doxycycline: Inhibit collagenase activity with oral doxycycline. Symblepharon Prevention:Use a glass shell or sweep a glass rod in the fornices (spaces between the eyelid and eyeball) twice daily to prevent adhesions. Complication Management Secondary glaucoma: Control intraocular pressure with beta-blockers like timolol or carbonic anhydrase inhibitors such as acetazolamide. Limbal stem cell deficiency: Consider amniotic membrane or limbal stem cell transplantation. Pseudopterygium: Surgically remove abnormal conjunctival growth and consider autografting or amniotic membrane transplantation. Corneal opacity: Consider keratoplasty or keratoprosthesis for severe corneal damage.

Implications Chemical injuries can cause permanent vision loss, physical disfigurement and psychological impact despite treatment. Take history (Option A): Taking history is important but not the priority in an emergency. Every second matters in chemical burns, and delaying irrigation can worsen corneal damage. Check vision in both eyes (Option C): Checking vision can be done after irrigation, but delaying decontamination increases ocular morbidity. Check IOP and plan for surgery (Option D): Checking intraocular pressure (IOP) and planning for surgery is important after initial management. Alkali burns can cause secondary glaucoma, but irrigation takes precedence over any diagnostic steps.

References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 456-459

↑ *Back to this question in Part A*

@Neet_pg_bot

Q3: A patient has a history of a flying foreign body injury to the eye. Hemochromatosis is subsequently diagnosed. Which of the following substances is most likely involved in the deposition? (Q852494)

A. Iron

B. Aluminum

C. Wood

D. Glass

Correct Answer: A. Iron

Explanation: Hemochromatosis is a condition characterized by excessive iron accumulation in the body. In the context of a flying foreign body injury, if the foreign body were made of iron, it could lead to iron deposition in the affected tissues, contributing to the development of hemochromatosis. Intraocular Foreign Bodies (IOFBs) Description An IOFB can lodge in any part of the anterior or posterior segments, causing mechanical damage, infection, or toxic effects on intraocular structures. Types Inert: Glass, wood, plastic, rubber, gold, silver, platinum Reactive: Organic and Inorganic (iron causing siderosis, copper causing chalcosis). Diagnosis History. Examination (wound examination, foreign body location, associated signs such as lid laceration and anterior segment damage should be noted). CT (investigation of choice) with axial and coronal cuts detects and localizes a metallic IOFB. MRI is contraindicated (metallic IOFB). Treatment Magnetic removal (metallic IOFB) Forceps removal through pars plana vitrectomy Prophylaxis against infection (topical/systemic antibiotics) Conservative and supportive management (topical corticosteroids, analgesics, etc). Modes of Damage: Mechanical Effects Depend on the size and velocity of the foreign body (FB). Introduction of Infection Metallic FB are usually sterile due to the heat generated on their emission. Pieces of wood or stone carry a greater chance of infection, often leading to endophthalmitis. Reactions to a Foreign Body (FB) Inert FB: No reaction. (Options C and D) Lead & Aluminium: Cause fibrosis (Option B) . Toxic Non-Metallic FB (Zinc, Copper, Mercury, Aluminium): Incite suppuration. Siderosis (Iron FB - Degenerative Changes Due to Electrolytic Dissociation and Dissemination):

Earliest sign: Deposition in the anterior capsule of the lens, appearing as a rusty ring (pathognomonic). Heterochromia iridis: Hyperchromic. Visual loss: Due to pigmentary retinal dystrophy. Secondary chronic glaucoma. Diagnosis: Prussian blue reaction. Chalcosis (Copper from Alloy Deposited on Membranous Structures): Kayser-Fleischer Ring: Deposition in Descemet's membrane (seen in Wilson's disease). Sunflower cataract. No degenerative changes: Visual prognosis is good. Caterpillar Hair: Produces granulomatous iridocyclitis, known as Ophthalmia Nodosa.

References:

Yanoff and Duker Ophthalmology, 5th Edition, Pages 675-676. Kanski's Clinical Ophthalmology, 10th Edition, Pages 927-928.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q4: The patient presents with foreign body (chuna) in his eyes and complaints of irritability and lacrimation. What is the first step management in this patient? (Q840064)

A. Antibiotics eye drops

B. Ascorbic eye drops

C. Irrigation of eye with normal saline

D. Steroids

Correct Answer: C. Irrigation of eye with normal saline

Explanation: Correct Option C - Irrigation of the eye with normal saline: Chemical injury to the conjunctiva and cornea is a severe ocular emergency that must be treated immediately. The first and the most crucial step in treating chemical injuries is copious eye irrigation by double eversion of the lid with normal saline. Deliver a minimum of 2 liters of water in 20-30 minutes or until pH is restored. Incorrect Options : Option A - Antibiotics drops in the eye: Antibiotics are used to prevent infection, but it is not the initial step. Option B - Ascorbic eye drops: Ascorbic eye drops promote collagen synthesis and improve wound healing but it is not the initial step. Option D - Steroids: Steroids reduce inflammation, neutrophil infiltration, and address anterior uveitis. Cautious use is recommended after 2 weeks, as steroids may interfere with collagen synthesis.

References:

.

↑ *Back to this question in Part A*

@Neet_pg_bot

Q5: Which investigation is most suitable for localizing a foreign body in the eye of an industrial worker admitted to the hospital due to an eye injury caused by cutting stone with a hammer and iron chisel? (Q857237)

A. B mode scan

B. CT

C. X-ray

D. MRI

Correct Answer: B. CT

Explanation: CT scan is the investigation of choice in case of an intraocular foreign body and MRI is contraindicated in cases where there is a risk of metallic foreign bodies. Intraocular Foreign Bodies (IOFBs): Definition An IOFB is a foreign object that penetrates the eye and remains lodged inside. It is a serious injury that can result in permanent vision loss. Common Types of IOFBs Metallic (90% of cases): Iron, steel. Others: Glass, stone, lead pellets, copper, aluminium, plastic, wood. Foreign Body Reactions Inorganic Materials: Glass, plastic are inert; lead, aluminium, copper cause inflammatory responses. Organic Materials: Wood causes giant cell reactions and granulomatous inflammation . Diagnosis Detailed History: Understand injury mechanism for IOFB type. Ocular Examination: Slit-lamp examination Gonioscopy Ophthalmoscopy (back of the eye) Signs: Subconjunctival hemorrhage Corneal scars/ Iris holes Opaque tracks in the lens/ IOFB in vitreous or retina Investigations: X-rays: Less common; CT is preferred. (Option C) Ultrasonography: B-mode USG locates both radiopaque and non-radiopaque IOFBs. (Option A) CT Scan: Investigation of choice/ Gold standard for detailed cross-sectional images. (Option B) MRI: Contra-indicated in case of metal foreign bodies (Option D) Treatment IOFB Removal: Essential except for inert, sterile IOFBs not affecting vision or if removal is too risky. Magnetic IOFBs: Removed using a handheld electromagnet. Non-magnetic IOFBs: Requires specialized procedures. Removal Methods: Anterior Chamber: Corneal incision, avoiding central cornea. Magnetic IOFBs: Handheld magnet. Non-magnetic IOFBs: Specialized forceps. Iris: Sector iridectomy for IOFBs in the iris. Lens: Extracapsular cataract extraction (ECCE) with or without lens removal. Vitreous and Retina: Pars plana vitrectomy Post-Removal Management Topical/systemic antibiotics for infection prevention. Topical corticosteroids to reduce inflammation. Cycloplegic agents (e.g., atropine) for pupil dilation and pain relief.

References:

: AK Khurana Comprehensive Ophthalmology 7th edition, page 450-454

↑ *Back to this question in Part A*

@Neet_pg_bot

Q6: A patient was brought to the outpatient department with a history of trauma near the eye and now presents with symptoms of fever, severe pain, and swelling in the left

eye. On examination, no abnormal findings were observed. What is the likely diagnosis? (Q857714)

A. Orbital cellulitis

B. Endophthalmitis

C. Panophthalmitis

D. Preseptal cellulitis




Correct Answer: D. Preseptal cellulitis

Explanation: Symptoms like fever, pain and swelling limited to the region anterior to the orbital septum with no abnormal ocular findings is suggestive of Preseptal cellulitis . {{caption_text}} {{caption_text}} Orbital cellulitis: Definition Orbital cellulitis is an acute infection of the soft tissues of the orbit located behind the orbital septum. Etiology Sinusitis: Ethmoid sinusitis (most common) carries a high risk of causing orbital cellulitis. (Option A) Underlying dental problem. Dacryocystitis (inflammation of the lacrimal sac). Endogenous cause: Septicemia Exogenous cause: Fracture or surgery of the orbit. Causative Organisms Staphylococcus aureus (most common) Haemophilus influenzae (most common in children) Streptococcus pneumoniae Streptococcus pyogenes Clinical Features Preseptal cellulitis: Fever, pain, swelling, tenderness on the outer eyelid region. (Option D) Orbital cellulitis: (Option A) Painful, unilateral Proptosis (swelling and severe pain worsened by eye movement or pressure) Restriction of eye movements (due to proptosis) Decreased visual acuity Staging Modified Chandler staging system: Stage I: Preseptal cellulitis/ periorbital cellulitis (limited anterior to the orbital septum) (Option D) Stage II: Inflammatory orbital oedema/ Orbital cellulitis (Posterior to orbital septum) Stage III: Subperiosteal abscess Stage IV: Orbital abscess Stage V: Cavernous sinus thrombosis {{caption_text}} Treatment Antibiotics (IV): Empirical therapy treats orbital cellulitis by giving Ceftazidime, Vancomycin, and Metronidazole. Monitoring of optic nerve function Surgery: Drainage of an orbital abscess or infected sinuses should be considered if there is a lack of response to antibiotics.







References:

Parson’s Diseases of the eye 23rd edition, Page 137.

Kanski’s Clinical ophthalmology 9th edition, Page 125

↑ *Back to this question in Part A*

@Neet_pg_bot

Q7: A man suffered trauma to the eye from a punch while fighting with another person. What is the most common site of fracture in orbit? (Q857721)

A. Medial wall


B. Lateral wall


C. Roof

D. Floor

Correct Answer: D. Floor

Explanation: Floor is the weakest wal l of orbit due to the presence of Inferior orbital fissure (although medial wall of the orbit is the thinnest wall). Trauma to the orbit can result in fracture of the floor of the orbit - known as ‘ Blowout fracture ’ Blow-out fracture: Description Blow-out orbital fractures result from trauma causing backward displacement of the eyeball and an increase in intraorbital pressure, leading to a resultant fracture. Classification Pure blow-out fractures: Not associated with involvement of the orbital rim. Impure blow-out fractures: Associated with other fractures in the middle third of the facial skeleton. Clinical Features Periorbital oedema and blood extravasation (e.g., subconjunctival ecchymosis) Emphysema of the eyelids, more common with medial wall fractures Paraesthesia and anaesthesia in the distribution of the infraorbital nerve (lower lid, cheek, side of nose, upper lip, and upper teeth) Ipsilateral epistaxis (bleeding from maxillary sinus) Proptosis, often due to orbital oedema and haemorrhage Enophthalmos (due to orbital fat escape, muscle entrapment, and cavity enlargement) Diplopia Radiography X-ray: Water’s view shows ‘hanging drop’ opacity of the superior maxillary antrum due to herniated orbital contents. {{caption_text}} CT/MRI: Coronal sections show extent of the fracture and herniation-hanging drop or tear drop sign {{caption_text}} Management General Measures and Medical Treatment: Avoid nose blowing Systemic antibiotics to prevent sinus infection Analgesics and anti-inflammatory drugs Cold compresses to decrease swelling Surgical Management: Optimal surgery time: 10-14 days after injury Surgical indications: Persistent diplopia, large herniation, tissue incarceration, enophthalmos >3mm Orbital Surgeries: Orbitotomy: 4 approaches: anterior, lateral, transfrontal, and temporofrontal Exenteration: Removal of all orbital contents and periorbita through an incision along orbital margins.





References:

AK Khurana Comprehensive Ophthalmology 7th edition, pages 440-442.

↑ *Back to this question in Part A*

@Neet_pg_bot


Q8: Can you determine the observation indicated by the arrow in the image provided? (Q856717)

A. Vossius ring

B. Weiss ring


C. Fleischer’s ring


D. Kayser Fleischer ring



Correct Answer: A. Vossius ring


Explanation: The given image is suggestive of Vossius ring , a condition that is typically detected during an ophthalmological examination following a blunt trauma to the eye . The Vossius ring itself is identified by the presence of pigment dispersion in a circular shape on the anterior surface of the lens following blunt trauma to the eye. {{caption_text}} Blunt trauma to the eye: Damage Mechanisms Direct Impact: Greatest damage at impact site. Compression Wave Force: Distant damage ("contrecoup"). Indirect Force: Damage from impact against orbital walls. Specific Injuries Cornea: Descemet's membrane tears Conjunctiva: Subconjunctival haemorrhage Anterior Chamber: Hyphema (most common sign) , exudates. Iris, Pupil, Ciliary Body: Iridodialysis,Iridodonesis. Lens: Rosette shaped cataract , ectopia lentis, Vossius ring Vitreous: Liquefaction, opacities, detachment. Choroid: choroidal rupture(bucket handle type) Retina: Commotio retinae , Rhegmatogenous retinal detachment , Gray white retina due to Berlin's edema. Intraocular Pressure: Traumatic glaucoma , hypotony, ghost cell glaucoma , angle recession glaucoma Traumatic optic neuropathy-RAPD (relative afferent pupillary defect) Incorrect Options: Weiss Ring (Option B): This is an opacified ring-shaped structure that can be seen floating within the vitreous humor of the eye, often signifying a posterior vitreous detachment , a condition in which the vitreous gel separates from the retina. {{caption_text}} Fleischer's Ring (Option C): This refers to a pigmented, circular line found at the edge of the cornea in keratoconus patients due to iron deposition in the corneal epithelial layer . It's composed of iron deposits and signifies corneal thinning and protrusion. {{caption_text}} Kayser-Fleischer Ring (Option D): This is a brown-colored ring found at the periphery of the cornea, most commonly seen in patients with Wilson's disease , a rare genetic disorder leading to excessive copper deposition in the body. {{caption_text}}





Wide-field fundus photography. Peripheral retinal tear (white arrow), Weiss ring- sign of PVD (black arrow)





References:

AK Khurana Comprehensive Ophthalmology 7th edition, pages 445-449

↑ *Back to this question in Part A*

@Neet_pg_bot

Q9: An industrial worker was admitted to the hospital after an eye injury sustained while cutting stone with a hammer and iron chisel. A foreign body was suspected to be impacted in his eye. Which investigation can be detrimental to this patient? (Q856179)

A. B mode scan

B. CT

C. X-ray

D. MRI

Correct Answer: D. MRI

Explanation: MRI is contraindicated in cases where there is a risk of metallic foreign bodies , such as in this industrial worker with a suspected foreign body in the eye. The strong magnetic field of an MRI can cause movement or displacement of the metallic object, potentially leading to further injury or damage to the eye.

Intraocular Foreign Bodies (IOFBs): Definition An IOFB is a foreign object that penetrates the eye and remains lodged inside. It is a serious injury that can result in permanent vision loss. Common Types of IOFBs Metallic (90% of cases): Iron, steel. Others: Glass, stone, lead pellets, copper, aluminium, plastic, wood. Foreign Body Reactions Inorganic Materials: Glass, plastic are inert; lead, aluminium, copper cause inflammatory responses. Organic Materials: Wood causes giant cell reactions and granulomatous inflammation . Diagnosis Detailed History: Understand injury mechanism for IOFB type. Ocular Examination: Slit-lamp examination Gonioscopy Ophthalmoscopy (back of the eye) Signs: Subconjunctival hemorrhage Corneal scars/ Iris holes Opaque tracks in the lens/ IOFB in vitreous or retina Investigations: X-rays: Less common; CT is preferred. Ultrasonography: B-mode USG locates both radiopaque and non-radiopaque IOFBs. CT Scan: Investigation of choice/ Gold standard for detailed cross-sectional images. MRI: Contra-indicated in case of metal foreign bodies (Option D) Treatment IOFB Removal: Essential except for inert, sterile IOFBs not affecting vision or if removal is too risky. Magnetic IOFBs: Removed using a handheld electromagnet. Non-magnetic IOFBs: Requires specialized procedures. Removal Methods: Anterior Chamber: Corneal incision, avoiding central cornea. Magnetic IOFBs: Handheld magnet. Non-magnetic IOFBs: Specialized forceps. Iris: Sector iridectomy for IOFBs in the iris. Lens: Extracapsular cataract extraction (ECCE) with or without lens removal. Vitreous and Retina: Pars plana vitrectomy Post-Removal Management Topical/systemic antibiotics for infection prevention. Topical corticosteroids to reduce inflammation. Cycloplegic agents (e.g., atropine) for pupil dilation and pain relief.

References:

AK Khurana Comprehensive Ophthalmology 7th edition, page 450-454

↑ *Back to this question in Part A*

@Neet_pg_bot

Q10: What is the probable cause of bleeding in a patient who experienced a hyphema after being struck in the eye by a tennis ball? (Q859357)

A. Circulus arteriosus major

B. Anterior ciliary artery

C. Posterior ciliary artery

D. Circulus arteriosus minor

Correct Answer: B. Anterior ciliary artery

Explanation: In the case of a patient who sustains an eye injury from a tennis ball and develops hyphema (bleeding in the anterior chamber of the eye) , the likely source of bleeding is the anterior ciliary artery. Anterior ciliary arteries are branches of the ophthalmic artery that supply blood to the anterior segment of the eye, including the conjunctiva, sclera, and iris. These arteries can be injured in cases of trauma to the eye. Blood Supply of the Eye: Structure Arterial Supply Cornea Avascular Conjunctiva Anterior ciliary arteries, branches of ophthalmic artery Sclera Posterior ciliary arteries (short and long branches) (Option C) Iris Major arterial circle of the iris (circulus arteriosus major) (Option A) formed by anterior ciliary and long posterior ciliary arteries Ciliary Body Anterior

ciliary arteries and long posterior ciliary arteries Choroid Short posterior ciliary arteries Retina Central retinal artery Lens Avascular Extraocular Muscles Muscular branches of the ophthalmic artery (including anterior ciliary arteries) Eyelids Ophthalmic artery (superior and inferior palpebral branches), facial artery Lacrimal Gland Lacrimal artery Circulus arteriosus minor (Option D) There is no anatomical structure known as the circulus arteriosus minor. It is an incorrect option.

References:

https://www.ncbi.nlm.nih.gov/books/NBK53329/

↑ *Back to this question in Part A*

@Neet_pg_bot